

US Patent & Trademark Office

Patent Public Search | Text View

United States Plant Patent Application Publication	20250280747
Kind Code	P1
Publication Date	September 04, 2025
Inventor(s)	Mussche; Erik

Hoya Plant Named 'G4Y0002'

Abstract

A new and distinct *Hoya carnososa* plant named 'G4Y0002' which is characterized by uniquely variegated foliage borne on scandent stems, brown juvenile stems, juvenile foliage that is centered brown and broadly margined with a vibrant pink general coloration, mature foliage that is green and broadly margined with a pale-yellow general coloration, and the stability of these characteristics from generation to generation.

Inventors:	Mussche; Erik (Bogor, ID)
Applicant:	Grow 4 You B.V. (Enter, ID)
Family ID:	96880312
Assignee:	Grow 4 You B.V. (Enter, ID)
Appl. No.:	18/831491
Filed:	February 26, 2025

Foreign Application Priority Data

QZ	20240575	Mar. 01, 2024
----	----------	---------------

Publication Classification

Int. Cl.:	A01H6/08 (20180101); A01H5/12 (20180101)
U.S. Cl.:	
CPC	A01H6/08 (20180501); A01H5/12 (20130101);
USPC	PLT/380

Background/Summary

[0001] Latin name of the genus and species: The Latin name of the genus and species of the novel variety disclosed herein is *Hoya carnosa*.

[0002] Variety denomination: The inventive variety of *Hoya* disclosed herein has been given the variety denomination ‘G4Y0002’.

CROSS REFERENCE TO RELATED APPLICATIONS

[0003] This application claims priority to the Community Plant Variety Rights application number 2024/0575, filed Mar. 1, 2024, which is herein incorporated by reference.

BACKGROUND OF THE INVENTION

[0004] Parentage: The new plant originated as a spontaneous, partial-plant, branch mutation. The mutation was discovered at a greenhouse in Bogor, Indonesia in January of 2022, growing amongst a plurality of cultivated *Hoya carnosa* ‘Tri-Color’ plants (U.S. Plant Pat. No. 2,950). The mutation exhibited conspicuously variegated foliage and a high degree of vibrant pink coloration of the juvenile foliage, relative to the parent. The mutation was isolated and grown to a mature size in order to evaluate stability and uniqueness of the characteristics first observed. Upon confirmation of the stability and uniqueness, the new plant was selected for commercialization and given the name, ‘G4Y0002’.

[0005] Asexual Reproduction: Asexual reproduction of the new cultivar ‘G4Y0002’, by way of softwood stem cuttings, was first initiated in May of 2022 in a commercial production greenhouse in Bogor, Indonesia. After three subsequent generations have shown that the unique features of the instant cultivar are stable and reproduced true to type.

SUMMARY OF THE INVENTION

[0006] The cultivar ‘G4Y0002’ has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature, day length, and light intensity, without, however, any variance in genotype. The following traits have been repeatedly observed and are determined to be the unique characteristics of ‘G4Y0002’. These characteristics in combination distinguish ‘G4Y0002’ as a new and distinct *Hoya* cultivar: [0007] 1. *Hoya* ‘G4Y0002’ exhibits a vining habit with an abundance of variegated ovate to elliptic foliage in a whorled arrangement that grows opposite decussate vertically along the entire length of the scandent stems; and [0008] 2. *Hoya* ‘G4Y0002’ exhibits brown juvenile stems; and [0009] 3. *Hoya* ‘G4Y0002’ exhibits juvenile foliage that is centered brown and broadly margined with red to greyed-purple, generally appearing as a vibrant pink color; and [0010] 4. *Hoya* ‘G4Y0002’ exhibits mature foliage that is green and broadly margined with light greyed-yellow, generally appearing as a pale-yellow coloration.

Description

BRIEF DESCRIPTION OF THE FIGURES

[0011] FIG. 1 illustrates, as nearly true as it is reasonably possible to make the same in color photographs of this type, an exemplary 20-week-old ‘G4Y0002’ plant in an 18 cm nursery pot in a greenhouse in Honselersdijk, the Netherlands.

[0012] FIG. 2 illustrates, as nearly true as it is reasonably possible to make the same in color photographs of this type, the mature foliage of ‘G4Y0002’.

[0013] FIG. 3 illustrates, as nearly true as it is reasonably possible to make the same in color photographs of this type, the juvenile foliage of ‘G4Y0002’.

BOTANICAL DESCRIPTION OF THE PLANT

[0014] The following observations and measurements were made in February of 2024 and describe

a 12-month-old 'G4Y0002' plant grown in a 17 cm nursery pot with 21 cm bamboo stakes grown in a climate-controlled greenhouse in Honselersdijk, the Netherlands. Plants were irrigated regularly and given supplemental lighting to provide a consistent 14 hours of light per day. No fertilizer applications or chemical pest control measures were utilized during production.

[0015] Those skilled in the art will appreciate that certain characteristics will vary with older or, conversely, with younger plants. 'G4Y0002' has not been observed under all possible environmental conditions. Where dimensions, sizes, colors and other characteristics are given, it is to be understood that such characteristics are approximations or averages set forth as accurately as practicable. The phenotype of the variety may differ from the descriptions set forth herein with variations in environmental, climatic and cultural conditions. Color notations are based on *The Royal Horticultural Society Colour Chart*, The Royal Horticultural Society, sixth edition.

[0016] A botanical description of 'G4Y0002' and comparisons with the parent and most similar commercial variety of *Hoya* are provided below. [0017] Plant description: [0018] *Growth habit*.—Evergreen vining perennial. [0019] *Average height*.—Vining plant with indeterminate growth; approximately 78.1 cm as observed. [0020] *Average width*.—Vining plant with indeterminate growth; approximately 17.7 cm as observed. [0021] *Propagation method*.—Softwood stem cuttings; rooting is initiated in approximately 20 weeks, and approximately 2 to 4 weeks are required to produce a rooted cutting when grown in a greenhouse with at least 90% humidity and an ambient average temperature of 22 degrees Celsius. [0022] *Time to produce a finished plant*.—Approximately 5 months are required to produce a fully rooted and marketable plant in a 17 cm nursery container, when grown under greenhouse conditions. [0023] *Disease and pest resistance or susceptibility*.—Plants have not been observed to be susceptible or resistant to pathogens and pests common to *Hoya elliptica*. [0024] *Environmental tolerances*.—Adapt to at least USDA Zones 10 through 12; moderate to high tolerance to rain; moderate tolerance to wind; low tolerance to drought. [0025] Root system: [0026] *General*.—Exhibits a relatively shallow root system with adventitious roots arising along the stems. [0027] *Branching*.—Freely branched. [0028] *Density*.—Moderately dense. [0029] *Distribution*.—Relatively shallow. [0030] *Texture*.—Fleshy; smooth; lacking root hairs. [0031] *Color*.—Nearest to a combination of white and greyed-yellow, nearest to RHS NN155D and 162C. [0032] Stem: [0033] *Branching habit*.—Basally clumping scandent stems, vertically climbing and twining around stakes or in trellises. [0034] *Number of main stems per plant*.—3. [0035] *Number of lateral branches per plant*.—None. [0036] *Aspect*.—Terete. [0037] *Length of main stems*.—25.6 cm. [0038] *Diameter of main stems*.—0.225 cm at the base. [0039] *Internode length on main stems*.—5.0 cm average. [0040] *Attitude*.—Vining; scandent. [0041] *Strength*.—Strong, highly flexible. [0042] *Texture and luster*.—Strigose, moderately to densely covered with short hairs, and glossy. Hairs are on average 0.5 mm in length and colored greyed-brown, difficult to distinguish from surrounding, nearest to RHS 199D. [0043] *Color, juvenile*.—Brown, nearest to RHS 200A. [0044] *Color, mature*.—Green, a blend of RHS 197A and 197B. [0045] *Color at internodes*.—Greyed-green, nearest to a combination of RHS 197A and 197B. [0046] Foliage: [0047] *Arrangement*.—Whorled at the base of the stem, as the vining plant climbs leaves become opposite decussate. [0048] *Division*.—Simple. [0049] *Attachment*.—Petiolate. [0050] *Abundance*.—Approximately 10 leaves per lateral branch at the time of observation. [0051] *Lamina*.—Dimensions — On average, 7.9 cm long and 3.8 cm wide; average thickness is 1.75 mm. Shape — Ovate to elliptic. Aspect — Near-flat. Apex — Aristate to apiculate. Base — Acute to shallow truncate. Margin — Entire; not undulated. Texture and luster — Glabrous, coriaceous, and smooth. Color — Juvenile foliage, adaxial surface — Brown, nearest to RHS 200A; broadly and irregularly margined with a blend of red and greyed-purple, nearest to RHS 53D and 184D; sparsely to moderately covered with small, irregularly shaped blotches that are colored greyed-green, nearest 198D. Juvenile foliage, abaxial surface — Greyed-green, nearest to RHS 197B, and lightly suffused yellow-green, nearest to RHS 148B; broadly and irregularly margined with greyed-red, nearest to RHS 182D; narrowly edged with greyed-purple along the margins of the lamina,

nearest to RHS 185A. Mature foliage, adaxial surface — A blend of green and yellow-green, nearest to RHS NN137A and 147A; broadly and irregularly margined with greyed-yellow, nearest to a combination of RHS 160C and 160D; sparsely covered with small, irregularly shaped blotches that are colored greyed-green, nearest 198D. Mature foliage, abaxial surface — Yellow-green, nearest to a blend of RHS 146C and 148B; broadly and irregularly margined greyed yellow, nearest to a combination of RHS 160C and 160D. Venation — Pattern — Parallel venation with brochidodromous secondary lateral veins. Venation color, adaxial surface — Green to yellow-green, nearest a blend of RHS NN137A and 147A. Venation color, abaxial surface — Yellow-green, nearest to RHS 146C and slightly tinted greyed — red proximally, nearest to RHS 182B. [0052] *Petiole*.—Length — 2.0 cm. Width — 0.3 cm. Aspect — Terete. Texture — Juvenile leaves moderately strigose, covered with short adpressed hairs averaging in length at 0.5 mm; colored greyed-brown, nearest to RHS 199D. Once leaves reach maturation, petioles are smooth and glabrous. Luster — Slightly glossy. Strength — Moderately to highly strong. Color, juvenile foliage — Greyed-purple to brown, nearest to a combination of RHS 185C, 185D and N200A. Color, mature foliage — Greyed-purple to brown, nearest to a combination of RHS 185C. [0053] Inflorescence: No Inflorescence observed to date. [0054] Seed and fruit: No seeds or fruit observed to date. [0055] Comparisons with the parent plant: Plants of the new cultivar ‘G4Y0002’ differ from the parent, *Hoya carnosa* ‘Tri-Color’ by the characteristics described in Table 1.

TABLE-US-00001 TABLE 1 Characteristic ‘G4Y0002’ ‘Tri-Color’ General coloration Brown and broadly The typical juvenile foliage is dark of the adaxial margined with red to green and more narrowly margined surface of the greyed-purple, generally with light pink; occasionally, a juvenile foliage. appearing as a vibrant juvenile leaf will emerge nearly pink color. homogenously pale yellow with pink suffusion or pale yellow with a central irregular blotch. General coloration Green and broadly Irregularly centered with pale of the adaxial margined with light green-yellow with light surface of the greyed-yellow, generally yellow-green blotches, and mature foliage. appearing as a pale- margined dark green. yellow coloration. General coloration Brown. Greyed-red to greyed-purple. of the juvenile stems. [0056] Comparison with the most similar *Hoya* cultivar known to the inventor: Plants of the new cultivar ‘G4Y0002’ differ from *Hoya carnosa* ‘Rubra’ (U.S. Plant Pat. No. 3,105) in the following characteristics described in Table 2 below.

TABLE-US-00002 TABLE 2 Characteristic ‘G4Y0002’ ‘Rubra’ General coloration of Variegated; brown and Irregularly centered with pale the adaxial surface of broadly margined with red green-yellow to greyed-yellow; the juvenile foliage. to greyed-purple, generally central portion irregularly appearing as a vibrant pink bordered with light yellow- color. green blotches; margined mid- green; central portions of the laminar are lightly to moderately suffused with pink. General coloration of Green and broadly Irregularly centered with pale the adaxial surface of margined with light greyed-yellow; central portion the mature foliage. greyed-yellow, generally irregularly bordered with light appearing as a pale- yellow-green blotches; yellow coloration. margined mid-green, General coloration of Brown. Greyed-red. the juvenile stems.

Claims

1. A new and distinct *Hoya carnosa* plant named ‘G4Y0002’, substantially as described and illustrated herein.
