



US012415826B2

(12) **United States Patent**
Yi et al.

(10) **Patent No.:** **US 12,415,826 B2**

(45) **Date of Patent:** **Sep. 16, 2025**

(54) **ORGANOMETALLIC COMPOUND,
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME, AND DIAGNOSTIC
COMPOSITION FOR INCLUDING THE
ORGANOMETALLIC COMPOUND**

(71) Applicant: **Samsung Electronics Co., Ltd.**,
Suwon-si (KR)

(72) Inventors: **Jeoungin Yi**, Seoul (KR); **Seungyeon
Kwak**, Suwon-si (KR); **Yoonhyun
Kwak**, Seoul (KR); **Sangdong Kim**,
Seongnam-si (KR); **Sunyoung Lee**,
Seoul (KR); **Aram Jeon**, Seoul (KR);
Hyeonho Choi, Seoul (KR); **Kyuyoung
Hwang**, Anyang-si (KR)

(73) Assignee: **SAMSUNG ELECTRONICS CO.,
LTD.**, Gyeonggi-Do (KR)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 734 days.

(21) Appl. No.: **17/060,731**

(22) Filed: **Oct. 1, 2020**

(65) **Prior Publication Data**

US 2021/0175444 A1 Jun. 10, 2021

(30) **Foreign Application Priority Data**

Dec. 10, 2019 (KR) 10-2019-0163877

(51) **Int. Cl.**
C07F 15/00 (2006.01)
H10K 50/15 (2023.01)
(Continued)

(52) **U.S. Cl.**
CPC **C07F 15/0086** (2013.01); **H10K 85/00**
(2023.02); **H10K 85/322** (2023.02);
(Continued)

(58) **Field of Classification Search**
CPC H10K 85/346; H10K 85/322; H10K
85/324; H10K 85/326; H10K 85/344;
(Continued)

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,358,634 B1 * 3/2002 Igarashi H10K 85/321
428/917
10,147,893 B2 12/2018 Lee et al.
(Continued)

FOREIGN PATENT DOCUMENTS

CN 106939024 A 7/2017
KR 20150082101 A 7/2015
(Continued)

OTHER PUBLICATIONS

Zhang, Jie, et al. "New phosphorescent platinum (II) Schiff base
complexes for PHOLED applications." Journal of Materials Chem-
istry 22.32 (2012): 16448-16457. (Year: 2012).
(Continued)


Primary Examiner — Jennifer A Boyd

Assistant Examiner — Braelyn R Watson

(74) *Attorney, Agent, or Firm* — CANTOR COLBURN
LLP


(57) **ABSTRACT**

Provided are an organometallic compound of Formula 1, an
organic light-emitting device including the organometallic
(Continued)



compound, and a diagnostic composition including the organometallic compound.

Formula 1



In Formula 1, ring A₁, ring A₃, and ring A₄ are each independently a C₅-C₃₀ carbocyclic group or a C₂-C₆₀ heterocyclic group, and ring A₂ is an N-containing 5-membered heterocyclic group.

17 Claims, 1 Drawing Sheet

(51) Int. Cl.

H10K 50/16 (2023.01)
H10K 50/17 (2023.01)
H10K 50/18 (2023.01)
H10K 85/00 (2023.01)
H10K 85/30 (2023.01)
H10K 85/60 (2023.01)

(52) U.S. Cl.

CPC **H10K 85/324** (2023.02); **H10K 85/326** (2023.02); **H10K 85/344** (2023.02); **H10K 85/346** (2023.02); **H10K 85/371** (2023.02); **H10K 85/381** (2023.02); **H10K 85/6572** (2023.02); **H10K 85/6574** (2023.02); **H10K 85/6576** (2023.02); **H10K 50/15** (2023.02); **H10K 50/16** (2023.02); **H10K 50/171** (2023.02); **H10K 50/18** (2023.02)

(58) Field of Classification Search

CPC H10K 85/371; H10K 85/381; H10K 85/6572; H10K 85/6574; H10K 85/6576; H10K 50/15; H10K 50/16; H10K 50/171;

H10K 50/18; H10K 50/11; H10K 2101/10; H10K 50/12; H10K 50/14; H10K 85/331; H10K 85/341; C07F 15/0086; C07F 1/08; C07F 1/10; C07F 1/12; C07F 3/006; C07F 3/02; C07F 3/04; C07F 3/06; C07F 5/00; C07F 5/06; C07F 7/00; C07F 7/28; C07F 7/30; C07F 13/00; C07F 15/06; C07F 15/0046; C07F 15/0073; C07F 15/006; C09K 11/06; C09K 2211/182-188

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

2005/0123792	A1 *	6/2005	Deaton	H05B 33/14
				313/506
2007/0103060	A1 *	5/2007	Itoh	H10K 85/346
				313/504
2009/0236974	A1 *	9/2009	Tamaru	H10K 50/11
				313/504
2013/0274473	A1	10/2013	Che et al.	
2015/0021585	A1 *	1/2015	Yu	C09K 11/06
				257/40
2015/0194615	A1 *	7/2015	Lin	C07F 15/0093
				546/4
2017/0237023	A1	8/2017	Kim et al.	
2021/0210701	A1	7/2021	Lin et al.	

FOREIGN PATENT DOCUMENTS

KR	1020160121369	A	10/2016
KR	1020170094652	A	8/2017

OTHER PUBLICATIONS

Turner, Eric, Nathan Bakken, and Jian Li. "Cyclometalated platinum complexes with luminescent quantum yields approaching 100%." *Inorganic chemistry* 52.13 (2013): 7344-7351. (Year: 2013).*

Baranoff, Etienne, et al. "Cyclometalated iridium (III) complexes based on phenyl-imidazole ligand." *Inorganic chemistry* 50.2 (2011): 451-462. (Year: 2011).*


English Abstract of CN 106939024.

Shiu-Lun Lai, et al., High Efficiency White Organic Light-Emitting Devices Incorporating Yellow Phosphorescent Platinum(II) Complex and Composite Blue Host, *Advanced Functional Materials* (2013), 23(41), 5168-5176.

English Translation of Office Action dated Jul. 22, 2024, issued in corresponding KR Patent Application No. 10-2019-0163877, 8 pp.

Office Action dated Jul. 22, 2024, issued in corresponding KR Patent Application No. 10-2019-0163877, 9 pp.

* cited by examiner



1

**ORGANOMETALLIC COMPOUND,
ORGANIC LIGHT-EMITTING DEVICE
INCLUDING THE SAME, AND DIAGNOSTIC
COMPOSITION FOR INCLUDING THE
ORGANOMETALLIC COMPOUND**

**CROSS-REFERENCE TO RELATED
APPLICATION**

This application claims priority to and the benefit of Korean Patent Application No. 10-2019-0163877, filed on Dec. 10, 2019, in the Korean Intellectual Property Office, the content of which is incorporated herein in its entirety by reference.

BACKGROUND

1. Field

One or more embodiments relate to organometallic compounds, organic light-emitting devices including the same, and diagnostic compositions including the organometallic compounds.

2. Description of Related Art

Organic light-emitting devices are self-emission devices, which have improved characteristics in terms of viewing angles, response time, brightness, driving voltages, and response speed, and produce full-color images.

In an example, an organic light-emitting device includes an anode, a cathode, and an organic layer between the anode and the cathode, wherein the organic layer includes an emission layer. A hole transport region may be between the anode and the emission layer, and an electron transport region may be between the emission layer and the cathode. Holes provided from the anode may move toward the emission layer through the hole transport region, and electrons provided from the cathode may move toward the emission layer through the electron transport region. The holes and the electrons recombine in the emission layer to produce excitons. These excitons transit from an excited state to a ground state, thereby generating light.

Meanwhile, luminescent compounds, for example, phosphorescent compounds, may be used for monitoring, sensing, and detecting biological materials such as various cells and proteins.

SUMMARY


One or more embodiments relate to organometallic compounds, organic light-emitting devices including the same, and diagnostic compositions including the organometallic compounds.

Additional aspects will be set forth in part in the description which follows and, in part, will be apparent from the description, or may be learned by practice of the presented embodiments of the disclosure.

According to an aspect of an embodiment, an organometallic compound represented by Formula 1 is provided.

2

Formula 1



In Formula 1,
ring A₁, ring A₃, and ring A₄ may each independently be a C₅-C₃₀ carbocyclic group or a C₂-C₆₀ heterocyclic group,

ring A₂ is an N-containing 5-membered heterocyclic group,

X₁ may be N, O, or S,

X₂ may be N or C, X₂₁, X₂₂, X₂₃, and X₂₄ may each independently be N, N(R), C(R), C(R)(R'), O, or S, wherein at least one of X₂ and X₂₁ to X₂₄ may be N or N(R),

X₃ and X₄ may each independently be N or C, and at least one of X₃ and X₄ may be C,

a bond between X₁ and M₁, a bond between X₁ and M₁, a bond between X₃ and M₁, and a bond between X₄ and M₁ may each independently be a covalent bond or a coordination bond,

Y₁ to Y₅ may each independently be N or C,

T₁ to T₃ may each independently be a single bond, *—N[(L₁)_{a1}—(R₁)_{b1}]*, *—B(R₁)*, *—P(R₁)*, *—C(R₁)(R₂)*, *—Si(R₁)(R₂)*, *—Ge(R₁)(R₂)*, *—S*, *—Se*, *—O*, *—C(=O)*, *—S(=O)*, *—S(=O)₂*, *—C(R₁)=*, *—C(R₁)=*, *—C(R₁)=C(R₂)*, *—C(=S)*, or *—C≡C*.

L₁ may be a single bond, a substituted or unsubstituted C₅-C₃₀ carbocyclic group, or a substituted or unsubstituted C₁-C₃₀ heterocyclic group,

a₁ may be an integer from 1 to 3, wherein when a₅ is 2 or more, two or more L₁(s) may be identical to or different from each other,

b₁ may be an integer from 1 to 5, wherein when b₁ is 2 or more, two or more R₁(s) may be identical to or different from each other,

R₁ and R₂ may optionally be linked to each other via a single bond, a double bond, or a first linking group to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₁-C₃₀ heterocyclic group,

R, R', R₁, R₂, R₁₀, R₂₀, R₃₀, and R₄₀ may each independently be hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl

3

group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉), b10, b20, b30, and b40 may each independently be an integer from 1 to 10,

when b20 is 2 or more, two or more R₂₀(s) may optionally be linked to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a},

R_{10a} is the same as explained in connection with R₁₀, at least one substituent of the substituted C₅-C₃₀ carbocyclic group, the substituted C₁-C₃₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is:

deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group,

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₁₀ alkoxy group, each substituted with at least one of deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), —B(Q₁₆)(Q₁₇), —P(=O)(Q₁₈)(Q₁₉), or any combination thereof,

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group,

4

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), —B(Q₂₆)(Q₂₇), —P(=O)(Q₂₈)(Q₂₉), or a combination thereof, or —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), —B(Q₃₆)(Q₃₇), or —P(=O)(Q₃₈)(Q₃₉),

wherein Q₁ to Q₉, Q₁₁ to Q₁₉, Q₂₁ to Q₂₉ and Q₃₁ to Q₃₉ may each independently hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryl group substituted with at least one of a C₁-C₆₀ alkyl group, a C₆-C₆₀ aryl group, or any combination thereof, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group.

Another aspect provides an organic light-emitting device including a first electrode, a second electrode, and an organic layer including an emission layer between the first electrode and the second electrode, wherein the organic layer includes at least one of the organometallic compounds.

Another aspect provides a diagnostic composition including at least one organometallic compound represented by Formula 1.

BRIEF DESCRIPTION OF THE DRAWING

The above and other aspects, features, and advantages of certain embodiments of the disclosure will be more apparent from the following description taken in conjunction with

FIGURE which shows a schematic cross-sectional view of an organic light-emitting device according to an exemplary embodiment.

DETAILED DESCRIPTION

Reference will now be made in detail to embodiments, examples of which are illustrated in the accompanying drawings, wherein like reference numerals refer to like elements throughout. In this regard, the present embodi-

5

ments may have different forms and should not be construed as being limited to the descriptions set forth herein.

Accordingly, the embodiments are merely described below, by referring to the figures, to explain aspects. As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. Expressions such as “at least one of,” when preceding a list of elements, modify the entire list of elements and do not modify the individual elements of the list.

It will be understood that when an element is referred to as being “on” another element, it can be directly on the other element or intervening elements may be present therebetween. In contrast, when an element is referred to as being “directly on” another element, there are no intervening elements present.

It will be understood that, although the terms “first,” “second,” “third” etc. may be used herein to describe various elements, components, regions, layers and/or sections, these elements, components, regions, layers and/or sections should not be limited by these terms. These terms are only used to distinguish one element, component, region, layer or section from another element, component, region, layer or section. Thus, “a first element,” “component,” “region,” “layer” or “section” discussed below could be termed a second element, component, region, layer or section without departing from the teachings herein.

The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used herein, “a,” “an,” “the,” and “at least one” do not denote a limitation of quantity, and are intended to cover both the singular and plural, unless the context clearly indicates otherwise. For example, “an element” has the same meaning as “at least one element,” unless the context clearly indicates otherwise.

“Or” means “and/or.” As used herein, the term “and/or” includes any and all combinations of one or more of the associated listed items. It will be further understood that the terms “comprises” and/or “comprising,” or “includes” and/or “including” when used in this specification, specify the presence of stated features, regions, integers, steps, operations, elements, and/or components, but do not preclude the presence or addition of one or more other features, regions, integers, steps, operations, elements, components, and/or groups thereof.

Furthermore, relative terms, such as “lower” or “bottom” and “upper” or “top,” may be used herein to describe one element’s relationship to another element as illustrated in the Figures. It will be understood that relative terms are intended to encompass different orientations of the device in addition to the orientation depicted in the Figures. For example, if the device in one of the figures is turned over, elements described as being on the “lower” side of other elements would then be oriented on “upper” sides of the other elements. The exemplary term “lower,” can therefore, encompass both an orientation of “lower” and “upper,” depending on the particular orientation of the FIGURE. Similarly, if the device in one of the figures is turned over, elements described as “below” or “beneath” other elements would then be oriented “above” the other elements. The exemplary terms “below” or “beneath” can, therefore, encompass both an orientation of above and below.

“About” or “approximately” as used herein is inclusive of the stated value and means within an acceptable range of deviation for the particular value as determined by one of ordinary skill in the art, considering the measurement in question and the error associated with measurement of the particular quantity (i.e., the limitations of the measurement

6


system). For example, “about” can mean within one or more standard deviations, or within 30%, 20%, 10% or 5% of the stated value.

Unless otherwise defined, all terms (including technical and scientific terms) used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this disclosure belongs. It will be further understood that terms, such as those defined in commonly used dictionaries, should be interpreted as having a meaning that is consistent with their meaning in the context of the relevant art and the present disclosure, and will not be interpreted in an idealized or overly formal sense unless expressly so defined herein.

Exemplary embodiments are described herein with reference to cross section illustrations that are schematic illustrations of idealized embodiments. As such, variations from the shapes of the illustrations as a result, for example, of manufacturing techniques and/or tolerances, are to be expected. Thus, embodiments described herein should not be construed as limited to the particular shapes of regions as illustrated herein but are to include deviations in shapes that result, for example, from manufacturing. For example, a region illustrated or described as flat may, typically, have rough and/or nonlinear features. Moreover, sharp angles that are illustrated may be rounded. Thus, the regions illustrated in the figures are schematic in nature and their shapes are not intended to illustrate the precise shape of a region and are not intended to limit the scope of the present claims.

An aspect of the present disclosure provides an organo-metallic compound represented by Formula 1 below:

Formula 1



M_1 in Formula 1 may be beryllium (Be), magnesium (Mg), aluminum (Al), calcium (Ca), titanium (Ti), manganese (Mn), cobalt (Co), copper (Cu), zinc (Zn), gallium (Ga), germanium (Ge), zirconium (Zr), ruthenium (Ru), rhodium (Rh), palladium (Pd), silver (Ag), rhenium (Re), platinum (Pt), or gold (Au).

In one or more embodiments, M_1 may be Pd, Pt, or Au, but embodiments of the present disclosure are not limited thereto.

Ring A_1 , ring A_3 , and ring A_4 in Formula 1 may each independently be a C_5 - C_{30} carbocyclic group or a C_2 - C_{60} heterocyclic group.

In one or more embodiments, ring A_1 , ring A_3 , and ring A_4 may each independently be a benzene group, a naphthalene group, an anthracene group, a phenanthrene group, a triphenylene group, a pyrene group, a chrysene group, a cyclopentadiene group, a 1,2,3,4-tetrahydronaphthalene group, a furan group, a thiophene group, a silole group, an indene group, a fluorene group, an indole group, a carbazole group, a benzofuran group, a dibenzofuran group, a benzo-

7

thiophene group, a dibenzothiophene group, a benzosilole group, a dibenzosilole group, an azafluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a quinoxaline group, a quinazoline group, a phenanthroline group, a pyrrole group, a pyrazole group, an imidazole group, a triazole group, a tetrazole group, an oxazole group, an isooxazole group, a thiazole group, an isothiazole group, an oxadiazole group, a thiadiazole group, a benzopyrazole group, a benzimidazole group, an indazole group, a benzoxazole group, a benzothiazole group, a benzoxadiazole group, a benzothiadiazole group, a benzotriazole group, a diazaindene group, a triazaindene group, a 5,6,7,8-tetrahydroisoquinoline group, or a 5,6,7,8-tetrahydroquinoline group.

For example, ring A_1 , ring A_3 , and ring A_4 may each independently be a benzene group, a naphthalene group, a 1,2,3,4-tetrahydronaphthalene group, a fluorene group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, or a dibenzosilole group.

Ring A_2 in Formula 1 may be an N-containing 5-membered heterocyclic group.

In one or more embodiments, ring A_2 may be a pyrrole group, a pyrazole group, an imidazole group, a triazole group, a tetrazole group, an oxazole group, an isooxazole group, a thiazole group, an isothiazole group, an oxadiazole group, or a thiadiazole group.

In one or more embodiments, ring A_2 may be a pyrazole group, an imidazole group, a triazole group, or a tetrazole group.

In one or more embodiments, ring A_2 may optionally be condensed with a ring that is formed by linking two or more $R_{20}(s)$ together.

X_1 in Formula 1 may be N, O, or S.

In one or more embodiments, in Formula 1, X_1 may be O or S, and a bond between X_1 and M_1 may be a covalent bond.

For example, X_1 may be O, but embodiments of the present disclosure are not limited thereto.

In Formula 1, X_2 may be N or C, and X_{21} , X_{22} , X_{23} , and X_{24} may each independently be N, N(R), C(R), C(R)(R'), O, or S, wherein at least one of X_2 and X_{21} to X_{24} may be N or N(R).

In one or more embodiments, X_2 may be N, and a bond between X_2 and M_1 may be a coordination bond.

In one or more embodiments, X_{21} to X_{24} may each be C(R) or C(R)(R').

In one or more embodiments, at least one of X_{21} to X_{24} may be N or N(R).

X_3 and X_4 in Formula 1 may each independently be N or C, and at least one of X_3 and X_4 may be C.

For example, X_3 may be C and X_4 may be N; or X_3 may be N and X_4 may be C.

In one or more embodiments, in Formula 1, X_3 may be C and X_4 may be N.

In Formula 1, one of a bond between X_3 and M_1 and a bond between X_4 and M_1 may be a covalent bond, and the other bond may be coordination bond. Thus, the organometallic compound represented by Formula 1 may be electrically neutral.

In one or more embodiments, a bond between X_3 and M_1 may be a covalent bond, and a bond between X_4 and M_1 may be a coordination bond, but embodiments of the present disclosure are not limited thereto.

8


In Formula 1, X_{21} may be N or C(R_{2a}), X_{22} may be N or C(R_{2b}), X_{23} may be N or C(R_{2c}), and at least one of X_{21} to X_{23} may be N.

For example, one or two of X_{21} to X_{23} may be N.

In one or more embodiments, in Formula 1, X_{21} may be N, X_{22} may be C(R_{2b}), and X_{23} may be C(R_{2c}); or X_{21} may be N, X_{22} may be C(R_{2b}), and X_{23} may be N, but embodiments of the present disclosure are not limited thereto.


In one or more embodiments, R_{2a} , R_{2b} , and R_{2c} may each independently be the same as described in connection with R, R', and R_{20} .

In one or more embodiments, a moiety represented by




in Formula 1 may be a group represented by one of Formulae CY1-1 to CY1-16:


CY1-1




CY1-2



CY1-3









CY1-4



9


-continued




10

-continued


CY1-5




CY1-6




CY1-7




CY1-8



CY1-9



CY1-10



In Formulae CY1-1 to CY1-16,

X₁₁ may be N or C(R₁₁), X₁₂ may be N or C(R₁₂), X₁₃ may be N or C(R₁₃), X₁₄ may be N or C(R₁₄), X₁₅ may be N or C(R₁₅), X₁₆ may be N or C(R₁₆), X₁₇ may be N or C(R₁₇), and X₁₈ may be N or C(R₁₈), X₁₉ may be C(R_{19a})(R_{19b}), N(R₁₉), O, S, or Si(R_{19a})(R_{19b}), R₁₁ to R₁₉, R_{19a}, and R_{19b} are each independently the same as described in connection with R₁₀,

CY1-11

CY1-12

CY1-13

CY1-14


CY1-15

CY1-16








11

* and *¹ each indicate a binding site to a neighboring atom.

In one or more embodiments, a moiety represented by




in Formula 1 may be a group represented by one of Formulae CY2-1 to CY2-10:




12

-continued


CY2-8



CY2-9



CY2-10




In Formulae CY2-1 to CY2-10,

X₂₅ may be N or C(R₂₃), X₂₆ may be N or C(R₂₄), X₂₇ may be N or C(R₂₅), and X₂₈ may be N or C(R₂₆).

R₂₁ to R₂₆ are each independently the same as described in connection with R₂₀, and


*, *¹ and *¹¹ may each independently be a binding site to a neighboring atom.

In one or more embodiments, a moiety represented by




in Formula 1 may be a group represented by one of Formulae CY3-1 to CY3-22:

CY3-1









CY3-2



13

-continued




14

-continued

CY3-3


5



CY3-4

10

15




20

CY3-5

25


30



CY3-6

35

40




CY3-7

45

50


55

CY3-8



60

65



CY3-9

CY3-10

CY3-11

CY3-12







CY3-13

CY3-14

CY3-15

15

-continued



16


-continued

CY3-16

CY3-22

5

10



CY3-17

15

20

In Formulae CY3-1 to CY3-22,

X_{31} may be N or C(R_{31}), X_{32} may be N or C(R_{32}), X_{33} may be N or C(R_{33}), X_{34} may be N or C(R_{34}), X_{35} may be N or C(R_{35}), X_{36} may be N or C(R_{36}), and X_{37} may be N or C(R_{37}),

X_{39} may be C(R_{39a})(R_{39b}), N(R_{39}), O, S, or Si(R_{39a})(R_{39b}),

R_{31} to R_{39} , R_{39a} , and R_{39b} are each independently the same as described in connection with R_{30} , and

CY3-18

25


*, *,*, and *'' may each independently be a binding site to a neighboring atom.

In one or more embodiments, a moiety represented by

30

CY3-19

35




40

in Formula 1 may be a group represented by one of Formulae CY4-1 to CY4-8:

CY4-1


45



CY3-20

50

CY4-2




55

CY4-3

CY3-21






60



65

17

-continued



In Formulae CY4-1 to CY4-8,

X_{41} may be N or C(R_{41}), X_{42} may be N or C(R_{42}), X_{43} may be N or C(R_{43}), X_{44} may be N or C(R_{44}), X_{45} may be N or C(R_{45}), X_{46} may be N or C(R_{46}), X_{47} may be N or C(R_{47}), and X_{48} may be N or C(R_{48}),

R_{41} to R_{48} are each independently the same as described in connection with R_{40} , and

* and *' each indicate a binding site to a neighboring atom.

T_1 to T_3 in Formula 1 may each independently be a single bond, *—N[(L_1)_{a1}—(R_1)_{b1}—]*, *—B(R_1)—*, *—P(R_1)—*, *—C(R_1)(R_2)—*, *—Si(R_1)(R_2)—*, *—Ge(R_1)(R_2)—*, *—S—*, *—Se—*, *—O—*, *—(=O)—*, *—S(=O)—*, *—S(=O)₂—*, *—C(R_1)=*, *—C(R_1)—*, *—C(R_1)=C(R_2)—*, *—C(=S)—*, or *—C≡C—*. R_1 and R_2 may be understood by referring to the related description to be presented later.

In one or more embodiments, T_1 and T_2 may each be a single bond.

In one or more embodiments, T_3 may be *—N[(L_1)_{a1}—(R_1)_{b1}—]*, *—C(R_1)(R_2)—*, *—Si(R_1)(R_2)—*, *—O—*, or *—S—*.

L_1 in Formula 1 may be a single bond, a substituted or unsubstituted C₅-C₃₀ carbocyclic group, or a substituted or unsubstituted C₁-C₃₀ heterocyclic group.

18

a1 may be 1 to 3, wherein when a1 is 2 or more, two or more L_1 (s) may be identical to or different from each other. For example, a1 may be 1 or 2.

b1 may be 1 to 5, wherein when b1 is 2 or more, two or more of R_1 (s) may be identical to or different from each other. For example, b1 may be 1, 2, or 3.

In one or more embodiments, L_1 may be: a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, or a pentacenylene group; or

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, or a pentacenylene group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₆₀ heterocycloalkyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, or any combination thereof.

R_1 and R_2 may optionally be linked together, via a single bond, a double bond, or a first linking group, to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a} (for example, a fluorene group, a xanthene group, or an acridine group, each unsubstituted or substituted with at least one R_{10a}). R_{10a} is the same as described in connection with R_1 .

The first linking group may be *—N(R_3)—*, *—B(R_3)—*, *—P(R_3)—*, *—C(R_3)(R_4)—*, *—Si(R_3)(R_4)—*, *—Ge(R_3)(R_4)—*, *—S—*, *—Se—*, *—O—*, *—C(=O)—*, *—S(=O)—*, *—S(=O)₂—*, *—C(R_3)=*, *—C(R_3)=C(R_4)—*, *—C(=S)—*, or *—C≡C—*, R_3 and R_4 are the same as described in connection with R_1 , and * and *' may each be a binding site to a neighboring atom.

R , R' , R_1 , R_2 , R_{10} , R_{20} , R_{30} , and R_{40} in Formula 1 may each independently be hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted

19

C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉).

b10, b20, b30 and b40 in Formula 1 may each independently be an integer from 1 to 10, and

when b20 is 2 or more, two or more R₂₀(s) may optionally be linked together to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a}. R_{10a} is the same as described in connection with R₁₀.

In one or more embodiments, R, R', R₁, R₂, R₁₀, R₂₀, R₃₀, and R₄₀ may each independently be: hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, —SF₅, C₁-C₂₀ alkyl group, or a C₁-C₂₀ alkoxy group;

a C₁-C₂₀ alkyl group or a C₁-C₂₀ alkoxy group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuran group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuran group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group; or

20

tenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuran group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthrenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuran group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuran group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, or any combination thereof; or —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(Q₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉),

wherein Q₁ to Q₉ may each independently be:

—CH₃, —CD₃, —CD₂H, —CDH₂, —CH₂CH₃, —CH₂CD₃, —CH₂CD₂H, —CH₂CDH₂, —CHDCH₃, —CHDCD₂H, —CHDCDH₂, —CHDCD₃, —CD₂CH₃, —CD₂CD₃, —CD₂CD₂H, or —CD₂CDH₂;

an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, or a naphthyl group; or

21

an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, or a naphthyl group, each substituted with at least one deuterium, a C₁-C₁₀ alkyl group, a phenyl group, or any combination thereof.

For example, R₁ and R₂ may each independently be:

a C₁-C₃₀ alkyl group;

a C₁-C₃₀ alkyl group, substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group; or

a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzo-

22

carbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group, each substituted with deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof.

In one or more embodiments, R₁, R₂, R₁₀, R₂₀, R₃₀, and R₄₀ may each independently be:

hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isoheptyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, or a dibenzothiophenyl group;

a methyl group, an ethyl group, an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, an n-hexyl group, an isoheptyl group, a sec-hexyl group, a tert-hexyl group, an n-heptyl group, an isoheptyl group, a sec-heptyl group, a tert-heptyl group, an n-octyl group, an isooctyl group, a sec-octyl group, a tert-octyl group, an n-nonyl group, an isononyl group, a sec-nonyl group, a tert-nonyl group, an n-decyl group, an isodecyl group, a sec-decyl group, a tert-decyl group, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, or a dibenzothiophenyl group, each substituted with deuterium, —F, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a cyano group, a nitro group, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, or a dibenzothiophenyl group,

23

pyridinyl group, a pyrimidinyl group, a carbazolyl group, a dibenzofuranyl group, a dibenzothiophenyl group, or any combination thereof; or
 $-\text{N}(\text{Q}_1)(\text{Q}_2)$, $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$, $-\text{B}(\text{Q}_6)(\text{Q}_7)$, or
 $-\text{P}(=\text{O})(\text{Q}_8)(\text{Q}_9)$,







wherein Q_1 to Q_9 may each independently be:

$-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CH}_2\text{CH}_3$,
 $-\text{CH}_2\text{CD}_3$, $-\text{CH}_2\text{CD}_2\text{H}$, $-\text{CH}_2\text{CDH}_2$, $-\text{CHDCD}_3$,
 $-\text{CHDCD}_2\text{H}$, $-\text{CHDCDH}_2$, $-\text{CHDCD}_3$,
 $-\text{CD}_2\text{CH}_3$, $-\text{CD}_2\text{CD}_3$, $-\text{CD}_2\text{CD}_2\text{H}$, or
 $-\text{CD}_2\text{CDH}_2$;

an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, or a naphthyl group; or

an n-propyl group, an isopropyl group, an n-butyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, an n-pentyl group, an isopentyl group, a sec-pentyl group, a tert-pentyl group, a phenyl group, or a naphthyl group, each substituted with at least one deuterium, a C_1 to C_{10} alkyl group, a phenyl group, or any combination thereof.

For example, R , R' , R_1 , R_2 , R_{10} , R_{20} , R_{30} , and R_{40} may each independently be hydrogen, deuterium, $-\text{F}$, a cyano group, a nitro group, $-\text{SF}_5$, $-\text{CH}_3$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a group represented by one of Formulas 9-1 to 9-19, a group represented by one of Formulae 10-1 to 10-46, or $-\text{Si}(\text{Q}_3)(\text{Q}_4)(\text{Q}_5)$:



5

10

15

20

25

30

9-1

9-2

9-3

40

9-4

9-5

9-6

9-7

55

9-8















60

9-9

65

24

-continued



9-10

9-11

9-12

9-13

9-14

9-15

9-16

9-17

9-18

9-19

10-1













10-2

10-3

10-4

25

-continued


**26**

-continued

10-5


5

10-6



10


10-7



15

10-8


20



10-9

25

10-10



30

10-11



35

10-12

40

10-13

45





10-14

50



10-15

55



10-16

65



10-17

10-18

10-19

10-20

10-21

10-22

10-23









10-24

10-25

10-26

27


-continued

**28**

-continued


10-27

5



10-28 10

15




10-29

20


10-30

30




10-31

35




10-32

45




10-33

50



10-34

60



65

10-35

10-36

10-37

10-38






10-39

10-40

10-41

29

-continued



* in Formulae 9-1 to 9-19 and 10-1 to 10-46 indicates a binding site to a neighboring atom.

In one or more embodiments, at least one R_{10} may be a group represented by one of Formulae 9-1 to 9-19.

In one or more embodiments, at least one R_{20} may be a group represented by one of Formulae 10-1 to 10-46.

In one or more embodiments, in Formula 1, ring A_1 and ring A_3 may each be a benzene group and ring A_4 may be a pyridine group,

X_2 and X_4 may each be N and X_3 may be C,

a bond between X_2 and M_1 and a bond between X_4 and M_1 may each be a coordination bond, and a bond between X_3 and M_1 may be a covalent bond,

Y_1 to Y_5 may each be C, at least one of R, R', R_1 , R_2 , R_{10} , R_{20} , R_{30} , and R_{40} may be:

a C_1 - C_{30} alkyl group; or

a C_1 - C_{30} alkyl group substituted with at least one deuterium, —F, —Cl, —Br, —I, — CD_3 , — CD_2H , — CDH_2 , — CF_3 , — CF_2H , — CFH_2 , a hydroxyl group, a cyano

30

group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C_1 - C_{10} alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group; or

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, — CD_3 , — CD_2H , — CDH_2 , — CF_3 , — CF_2H , — CFH_2 , a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a

31

salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolynyl group, an isoquinolynyl group, a benzoquinolynyl group, a quinoxalynyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, an imidazopyrimidinyl group, or any combination thereof.

b10, b20, b30, and b40 in Formula 1 respectively indicate the number of R₁₀, R₂₀, R₃₀, and R₄₀ and may each independently be 1, 2, 3, or 4. When b10 is 2 or more, two or more of R₁₀(s) may be identical to or different from each other, when b20 is 2 or more, two or more of R₂₀(s) may be identical to or different from each other, when b30 is 2 or more, two or more of R₃₀(s) may be identical to or different from each other, and when b40 is 2 or more, two or more of R₄₀(s) may be identical to or different from each other.

In one or more embodiments, b1, b3, and b4 may each independently be 0, 1, or 2.





When b20 in Formula 1 is 2 or more, two or more R₂₀(s) may optionally be linked together to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a}, and R_{10a} may be the same as described in connection with R₁₀.

For example, when b20 in Formula 1 is 2 or more, two or more R₂₀(s) may optionally be linked together to form a cyclopentane group, a cyclopentadiene group, a furan group, a thiophene group, a pyrrole group, a silole group, an adamantane group, a norbornane group, a norbornene group, a cyclohexane group, a cyclohexene group, a benzene group, a naphthalene group, an indene group, an indole group, a benzofuran group, a benzothiophene group, a benzosilole group, a fluorene group, a carbazole group, a dibenzofuran group, a dibenzothiophene group, or dibenzosilole group, each unsubstituted or substituted with at least one R_{10a}.

In Formula 1, i) at least one of R₁ and R₂ and R₃₀ may optionally be linked together to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a}, ii) at least one of R₁ and R₂ and R₄₀ may optionally be linked together to form a C₁-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a}, and R_{10a} is the same as described in connection with R₁₀.




In one or more embodiments, the organometallic compound may be one of Compounds 1 to 12, but embodiments of the present disclosure are not limited thereto:

32



33

-continued

**34**

-continued

5

8

5

10

15

20

6

25

30

35

9

40

7

45

50





55

60

65


10

11



35

-continued




In the organometallic compound represented by Formula 1, a second ring represented by A_2 may be an N-containing 5-membered heterocyclic group, and at least one of a third ring represented by A_3 and a fourth ring represented by A_4 may be covalently bonded to M_1 via a C atom.

Although it is not intended to limit embodiments by any theory,

- i) Compounds A and B, in which a ring corresponding to the second ring of Formula 1 is a 6-membered ring, each show a longer wavelength than the organometallic compound, and thus, the luminescent characteristics of Compounds A and B are different from those of the organometallic compound, and
- ii) Compounds B and C, in which a ring corresponding to the third ring of Formula 1 and a ring corresponding to the fourth ring of Formula 1 are each linked to M_1 via a C atom, charge-transfer characteristics deteriorate and thus efficiency may be decreased.


Compound A




36

-continued

Compound B



Compound C



In the organometallic compound represented by Formula 1, the second ring represented by A_2 may be an N-containing 5-membered heterocyclic group, and at least one of the third ring represented by A_3 and the fourth ring represented by A_4 may be covalently bonded to M_1 via a C atom. Accordingly, luminescent characteristics and charge-transfer characteristics are excellent. Accordingly, an electronic device using the organometallic compound represented by Formula 1, for example, an organic light-emitting device using the organometallic compound represented by Formula 1 may have excellent luminescent efficiency and/or lifespan characteristics.

For example, the highest occupied molecular orbital (HOMO), lowest unoccupied molecular orbital (LUMO), and singlet (S_1) and triplet (T_1) energy levels of Compounds 1 to 6 were evaluated by using a DFT method of Gaussian program (structurally optimized at a level of B3LYP, 6-31G (d,p)). Evaluation results are shown in Table 1 below.


TABLE 1

Compound No.	HOMO (eV)	LUMO (eV)	Band gap	S_1 energy level (eV)	T_1 energy level (eV)
1	-4.51	-1.30	3.22	2.68	2.53
2	-4.44	-1.62	2.82	2.36	2.28
3	-4.34	-1.07	3.28	2.73	2.55
4	-4.48	-1.36	3.12	2.59	2.48
5	-4.46	-1.40	3.06	2.53	2.41
7	-4.50	-1.62	2.88	2.41	2.33
A	-4.54	-1.49	3.05	2.48	2.22
B	-4.42	-1.61	2.81	2.24	2.04
C	-4.56	-1.83	2.73	2.25	1.99


37

TABLE 1-continued


Compound				S ₁ energy	T ₁ energy
No.	HOMO (eV)	LUMO (eV)	Band gap	level (eV)	level (eV)




1



2



3




4


38

TABLE 1-continued

Compound				S ₁ energy	T ₁ energy
No.	HOMO (eV)	LUMO (eV)	Band gap	level (eV)	level (eV)



5




20

25

30

35



7

40

45


50

A

55


60

65



B

TABLE 1-continued

Compound		S ₁ energy		T ₁ energy
No.	HOMO (eV) LUMO (eV) Band gap	level (eV)	level (eV)	(eV)
				
C				

From Table 1, it is confirmed that the organometallic compound represented by Formula 1 has such electric characteristics that are suitable for use as a material for an emission layer for an electric device, for example, an organic light-emitting device.

Synthesis methods of the organometallic compound represented by Formula 1 may be recognizable by one of ordinary skill in the art by referring to Synthesis Examples provided below.

Accordingly, the organometallic compound represented by Formula 1 is suitable for use as a material for an organic layer of organic light-emitting device, for example, an emission layer. Thus, another aspect provides an organic light-emitting device including: a first electrode; a second electrode; and an organic layer placed between the first electrode and the second electrode and including an emission layer, and the organic layer includes at least one organometallic compound represented by Formula 1.

The organic light-emitting device may have, due to the inclusion of an organic layer including the organometallic compound represented by Formula 1, a low driving voltage, high efficiency, high power, high quantum efficiency, a long lifespan, a low roll-off ratio, and excellent color purity.

In one or more embodiments, in the organic light-emitting device, the first electrode is an anode, and the second electrode is a cathode, and the organic layer further includes a hole transport region between the first electrode and the emission layer and an electron transport region between the emission layer and the second electrode, and the hole transport region includes a hole injection layer, a hole transport layer, an electron blocking layer, or any combination thereof, and the electron transport region includes a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.

In one or more embodiments, the organometallic compound represented by Formula 1 may be included in the emission layer.

The organometallic compound is included in the emission layer may act as an emitter. For example, an emission layer including the organometallic compound represented by Formula 1 may emit phosphorescent light generated by the transfer of the triplet excitons of the organometallic compound into the ground state.

In one or more embodiments, the emission layer including the organometallic compound represented by Formula 1 may further include a host. The host may be any host, and details of the host may be the same as described above. The

amount of the host in the emission layer may be greater than the amount of the organometallic compound represented by Formula 1.

In one or more embodiments, the emission layer may include a host and a dopant, the host may be any host, and the dopant may include the organometallic compound represented by Formula 1. The emission layer may emit phosphorescent light generated by the transfer of triplet excitons of the organometallic compound, which acts as a dopant, to the ground state.

In one or more embodiments, the emission layer may emit green light having a maximum emission wavelength of about 490 nm to about 580 nm, for example, about 500 nm to about 570 nm.

In one or more embodiments, the emission layer may emit green light having a maximum emission wavelength of about 580 nm to about 780 nm, for example, about 590 nm to about 750 nm.

The expression “(an organic layer) includes at least one of organometallic compounds” used herein may include a case in which “(an organic layer) includes identical organometallic compounds represented by Formula 1” and a case in which “(an organic layer) includes two or more different organometallic compounds represented by Formula 1”.

For example, the organic layer may include, as the organometallic compound, only Compound 1. In this regard, Compound 1 may exist in an emission layer of the organic light-emitting device. In one or more embodiments, the organic layer may include, as the organometallic compound, Compound 1 and Compound 2. In this regard, Compound 1 and Compound 2 may exist in an identical layer (for example, Compound 1 and Compound 2 all may exist in an emission layer).

The term “organic layer” used herein refers to a single layer and/or a plurality of layers between the first electrode and the second electrode of the organic light-emitting device. The “organic layer” may include, in addition to an organic compound, an organometallic complex including metal.

FIGURE is a schematic cross-sectional view of an organic light-emitting device 10 according to an embodiment. Hereinafter, the structure of an organic light-emitting device according to an embodiment and a method of manufacturing an organic light-emitting device according to an embodiment will be described in connection with FIGURE. The organic light-emitting device 10 includes a first electrode 11, an organic layer 15, and a second electrode 19, which are sequentially stacked.

A substrate may be additionally located under the first electrode 11 or above the second electrode 19. For use as the substrate, any substrate that is used in organic light-emitting devices available in the art may be used, and the substrate may be a glass substrate or a transparent plastic substrate, each having excellent mechanical strength, thermal stability, transparency, surface smoothness, ease of handling, and water resistance.

In one or more embodiments, the first electrode 11 may be formed by depositing or sputtering a material for forming the first electrode 11 on the substrate. The first electrode 11 may be an anode. The material for forming the first electrode 11 may be materials with a high work function to facilitate hole injection. The first electrode 11 may be a reflective electrode, a semi-transmissive electrode, or a transmissive electrode. The material for forming the first electrode 11 may be indium tin oxide (ITO), indium zinc oxide (IZO), tin

41

oxide (SnO₂), or zinc oxide (ZnO). In one or more embodiments, the material for forming the first electrode **11** may be metal, such as magnesium (Mg), aluminum (Al), aluminum-lithium (Al—Li), calcium (Ca), magnesium-indium (Mg—In), or magnesium-silver (Mg—Ag).

The first electrode **11** may have a single-layered structure or a multi-layered structure including two or more layers. For example, the first electrode **11** may have a three-layered structure of ITO/Ag/ITO, but the structure of the first electrode **11** is not limited thereto.

The organic layer **15** is located on the first electrode **11**.

The organic layer **15** may include a hole transport region, an emission layer, and an electron transport region.

The hole transport region may be between the first electrode **11** and the emission layer.

The hole transport region may include a hole injection layer, a hole transport layer, an electron blocking layer, a buffer layer, or any combination thereof.

The hole transport region may include only either a hole injection layer or a hole transport layer. In one or more embodiments, the hole transport region may have a hole injection layer/hole transport layer structure or a hole injection layer/hole transport layer/electron blocking layer structure, wherein, for each structure, each layer is sequentially stacked in this stated order from the first electrode **11**.

When the hole transport region includes a hole injection layer (HIL), the hole injection layer may be formed on the first electrode **11** by using one or more suitable methods, for example, vacuum deposition, spin coating, casting, and/or Langmuir-Blodgett (LB) deposition.


When a hole injection layer is formed by vacuum deposition, the deposition conditions may vary according to a material that is used to form the hole injection layer, and the structure and thermal characteristics of the hole injection layer. For example, the deposition conditions may include a deposition temperature of about 100° C. to about 500° C., a vacuum pressure of about 10⁻⁸ torr to about 10⁻³ torr, and a deposition rate of about 0.01 Å/sec to about 100 Å/sec. However, the deposition conditions are not limited thereto.

When the hole injection layer is formed using spin coating, coating conditions may vary according to the material used to form the hole injection layer, and the structure and thermal properties of the hole injection layer. For example, a coating speed may be from about 2,000 rpm to about 5,000 rpm, and a temperature at which a heat treatment is performed to remove a solvent after coating may be from about 80° C. to about 200° C. However, the coating conditions are not limited thereto.


Conditions for forming a hole transport layer and an electron blocking layer may be understood by referring to conditions for forming the hole injection layer.

The hole transport region may include at least one m-MT-DATA, TDATA, 2-TNATA, NPB, β-NPB, TPD, Spiro-TPD, Spiro-NPB, methylated-NPB, TAPC, HMTPD, 4,4',4"-tris (N-carbazolyl)triphenylamine (TCTA), polyaniline/dodecylbenzenesulfonic acid (PANI/DBSA), poly(3,4-ethylenedioxythiophene)/poly(4-styrenesulfonate) (PEDOT/PSS), polyaniline/camphor sulfonic acid (PANI/CSA), polyaniline/poly(4-styrenesulfonate) (PANI/PSS), a compound represented by Formula 201 below, a compound represented by Formula 202 below, or any combination thereof.

42




m-MTDATA




TDATA

43



-continued



2-TNATA




NPB

 β -NPB


TPD

44


-continued




Spiro-TPD




Spiro-NPB



methylated NPB



TAPC





HMTPD

45

-continued

Formula 201



Ar₁₀₁ to Ar₁₀₂ in Formula 201 may each independently be:
 a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, or a pentacenylene group; or

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthrenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a naphthacenylene group, a picenylene group, a perylenylene group, or a pentacenylene group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀

46

alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, or any combination thereof.

xa and xb in Formula 201 may each independently be an integer from 0 to 5, or 0, 1, or 2. For example, xa may be 1 and xb may be 0, but xa and xb are not limited thereto.

R₁₀₁ to R₁₀₈, R₁₁₁ to R₁₁₉ and R₁₂₁ to R₁₂₄ in Formulae 201 and 202 may each independently be:

hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group (for example, a methyl group, an ethyl group, a propyl group, a butyl group, a pentyl group, a hexyl group, and so on), or a C₁-C₁₀ alkoxy group (for example, a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentoxy group, and so on);

a C₁-C₁₀ alkyl group or a C₁-C₁₀ alkoxy group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, or any combination thereof;

a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, or a pyrenyl group; and

a phenyl group, a naphthyl group, an anthracenyl group, a fluorenyl group, or a pyrenyl group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, or any combination thereof, but embodiments of the present disclosure are not limited thereto.

R₁₀₉ in Formula 201 may be:


a phenyl group, a naphthyl group, an anthracenyl group, or a pyridinyl group; and

a phenyl group, a naphthyl group, an anthracenyl group, or a pyridinyl group, each substituted with at least one a deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₂₀ alkyl group, a C₁-C₂₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyridinyl group, or any combination thereof.

According to an embodiment, the compound represented by Formula 201 may be represented by Formula 201A below, but embodiments of the present disclosure are not limited thereto:

47


Formula 201A



48

-continued


HT2




R₁₁, R₁₁₁, R₁₁₂, and R₁₀₉ in Formula 201A may be understood by referring to the description provided herein.

For example, the compound represented by Formula 201, and the compound represented by Formula 202 may include a compound of HT1 to HT20 illustrated below, but are not limited thereto:

HT1




HT3



49

-continued

HT4

**50**

-continued

HT6

5

10

15

20


25

30

35

40

HT5



45


50

55

60

65


HT7



51

-continued

HT8



5

10

15

20


25

30

35

40

HT9



45

50

55


60

65


52

-continued

HT10







HT11







53

-continued


**54**

-continued



55


-continued



A thickness of the hole transport region may be from about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å. When the hole transport region includes at least one of a hole injection layer and a hole transport layer, a thickness of the hole injection layer may be in a range of about 100 Å to about 10,000 Å, for example, about 100 Å to about 1,000 Å, and a thickness of the hole transport layer may be in a range of about 50 Å to about 2,000 Å, for example, about 100 Å to about 1,500 Å. When the thicknesses of the hole transport region, the hole injection layer and the hole transport layer are within these ranges, satisfactory hole transporting characteristics may be obtained without a substantial increase in driving voltage.


The hole transport region may further include, in addition to these materials, a charge-generation material for the improvement of conductive properties. The charge-generation material may be homogeneously or non-homogeneously dispersed in the hole transport region.

The charge-generation material may be, for example, a p-dopant. The p-dopant may be a quinone derivative, a metal oxide, a cyano group-containing compound, or any combination thereof, but embodiments of the present disclosure are not limited thereto. Examples of the p-dopant are: a quinone derivative, such as tetracyanoquinonodimethane (TCNQ) or 2,3,5,6-tetrafluoro-tetracyano-1,4-benzoquinonodimethane (F4-TCNQ); a metal oxide, such as a tungsten oxide or a molybdenum oxide; and a cyano group-containing compound, such as Compound HT-D1 below, but are not limited thereto.



56

-continued



The hole transport region may include a buffer layer.


Also, the buffer layer may compensate for an optical resonance distance according to a wavelength of light emitted from the emission layer, and thus, efficiency of a formed organic light-emitting device may be improved.

Then, an emission layer (EML) may be formed on the hole transport region by vacuum deposition, spin coating, casting, LB deposition, or the like. When the emission layer is formed by vacuum deposition or spin coating, the deposition or coating conditions may be similar to those applied in forming the hole injection layer although the deposition or coating conditions may vary according to a material that is used to form the hole transport layer.

Meanwhile, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be materials for the hole transport region described above and materials for a host to be explained later. However, the material for the electron blocking layer is not limited thereto. For example, when the hole transport region includes an electron blocking layer, a material for the electron blocking layer may be mCP, which will be explained later.


The emission layer may include a host and a dopant, and the dopant may include the organometallic compound represented by Formula 1.

The host may include at least one TPBi, TBADN, ADN (also referred to as "DNA"), CBP, CDBP, TCP, mCP, Compound H50, Compound H51, or any combination thereof:




57


-continued




TBADN




ADN



CBP




CDBP




TCP

58


-continued



mCP




H50



H51

In one or more embodiments, the host may further include a compound represented by Formula 301 below.

Formula 301



Ar₁₁₁ and Ar₁₁₂ in Formula 301 may each independently be:

a phenylene group, a naphthylene group, a phenanthrenylene group, or a pyrenylene group; or

a phenylene group, a naphthylene group, a phenanthrenylene group, or a pyrenylene group, each substituted with at least one a phenyl group, a naphthyl group, an anthracenyl group, or any combination thereof.

Ar₁₁₃ to Ar₁₁₆ in Formula 301 may each independently be:

a C₁-C₁₀ alkyl group, a phenyl group, a naphthyl group, a phenanthrenyl group, or a pyrenyl group; or


a phenyl group, a naphthyl group, a phenanthrenyl group, or a pyrenyl group, each substituted with at least one a phenyl group, a naphthyl group, an anthracenyl group, or any combination thereof.

g, h, i, and j in Formula 301 may each independently be an integer from 0 to 4, and may be, for example, 0, 1, or 2.

59


Ar₁₁₃ and Ar₁₁₆ in Formula 301 may each independently be:

- a C₁-C₁₀ alkyl group, substituted with at least one a phenyl group, a naphthyl group, an anthracenyl group, or any combination thereof;
- a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl, a phenanthrenyl group, a fluorenyl group, or any combination thereof;
- a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, or a fluorenyl group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₁₀ alkoxy group, a phenyl group, a naphthyl group, an anthracenyl group, a pyrenyl group, a phenanthrenyl group, a fluorenyl group, or any combination thereof; or



but embodiments of the present disclosure are not limited thereto.

In one or more embodiments, the host may include a compound represented by Formula 302 below:



Formula 302

Ar₁₂₂ to Ar₁₂₅ in Formula 302 are the same as described in detail in connection with Ar₁₁₃ in Formula 301.

Ar₁₂₆ and Ar₁₂₇ in Formula 302 may each independently be a C₁-C₁₀ alkyl group (for example, a methyl group, an ethyl group, or a propyl group).

k and l in Formula 302 may each independently be an integer from 0 to 4. For example, k and l may be 0, 1, or 2.

When the organic light-emitting device is a full-color organic light-emitting device, the emission layer may be patterned into a red emission layer, a green emission layer, and a blue emission layer. In one or more embodiments, due to a stacked structure including a red emission layer, a green emission layer, and/or a blue emission layer, the emission layer may emit white light.

When the emission layer includes a host and a dopant, an amount of the dopant may be in a range of about 0.01 parts by weight to about 15 parts by weight based on 100 parts by weight of the host, but embodiments of the present disclosure are not limited thereto.

60

A thickness of the emission layer may be in a range of about 100 Å to about 1,000 Å, for example, about 200 Å to about 600 Å. When the thickness of the emission layer is within this range, excellent light-emission characteristics may be obtained without a substantial increase in driving voltage.


Then, an electron transport region may be located on the emission layer.

The electron transport region may include a hole blocking layer, an electron transport layer, an electron injection layer, or any combination thereof.


For example, the electron transport region may have a hole blocking layer/electron transport layer/electron injection layer structure or an electron transport layer/electron injection layer structure, and the structure of the electron transport region is not limited thereto. The electron transport layer may have a single-layered structure or a multi-layered structure including two or more different materials.

Conditions for forming the hole blocking layer, the electron transport layer, and the electron injection layer which constitute the electron transport region may be understood by referring to the conditions for forming the hole injection layer.

When the electron transport region includes a hole blocking layer, the hole blocking layer may include, for example, at least one of BCP, Bphen, BAQ, or any combination thereof but embodiments of the present disclosure are not limited thereto.





BCP




Bphen

A thickness of the hole blocking layer may be in a range of about 20 Å to about 1,000 Å, for example, about 30 Å to about 300 Å. When the thickness of the hole blocking layer is within these ranges, the hole blocking layer may have excellent hole blocking characteristics without a substantial increase in driving voltage.


The electron transport layer may further include at least one BCP, Bphen, Alq3, BAQ, TAZ, NTAZ, or any combination thereof.

61Alq₃

BAlq



TAZ



NTAZ

62

In one or more embodiments, the electron transport layer may include at least one of ET1 to ET25, but are not limited thereto:

5

ET1

10

15

20

25

30

ET2

35

40

45




50

ET3

55


60

65



63

-continued

**64**

-continued

ET4

ET7

5

10

15

20

25

ET5

ET8

30

35

40

45

ET6





50

ET9

55

60


65



65

-continued

ET10




5

10

15

20


ET11



25

30


ET12



40

45

ET13



55


60

65


66

-continued


ET14



ET15




ET16



67

-continued

ET17



5

10


15

20


68

-continued

ET20



ET18




25

30


35

40

ET21



ET19



45


50

55


60

65

ET22



-continued



A thickness of the electron transport layer may be in a range of about 100 Å to about 1,000 Å, for example, about

70

ET23

5

Also, the electron transport layer may further include, in addition to the materials described above, a metal-containing material.

10



ET-D1



ET-D2

3.

The electron injection layer may include at least one LiF, NaCl, CsF, Li₂O, BaO, or any combination thereof.

40

ET25

50

60

0

6.

The diagnostic composition may be used in various applications including a diagnosis kit, a diagnosis reagent, a biosensor, and a biomarker.

The term “C₁-C₆₀ alkyl group” as used herein refers to a linear or branched saturated aliphatic hydrocarbon monovalent group having 1 to 60 carbon atoms, and examples thereof include a methyl group, an ethyl group, a propyl group, an isobutyl group, a sec-butyl group, a tert-butyl group, a pentyl group, an isoamyl group, and a hexyl group.

The term “C₁-C₆₀ alkylene group” used herein refers to a divalent group having the same structure as that of the C₁-C₆₀ alkyl group.

The term “C₁-C₆₀ alkoxy group” used herein refers to a monovalent group represented by —OA₁₀₁ (wherein A₁₀₁ is the C₁-C₆₀ alkyl group), and examples thereof include a methoxy group, an ethoxy group, and an isopropoxy group.

The term “C₂-C₆₀ alkenyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon double bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethenyl group, a propenyl group, and a butenyl group. The term “C₂-C₆₀ alkenylene group” used herein refers to a divalent group having the same structure as that of the C₂-C₆₀ alkenyl group.

The term “C₂-C₆₀ alkynyl group” as used herein refers to a hydrocarbon group formed by substituting at least one carbon-carbon triple bond in the middle or at the terminus of the C₂-C₆₀ alkyl group, and examples thereof include an ethynyl group, and a propynyl group. The term “C₂-C₆₀ alkynylene group” as used herein refers to a divalent group having the same structure as that of the C₂-C₆₀ alkynyl group.

The term “C₃-C₁₀ cycloalkyl group” as used herein refers to a monovalent saturated hydrocarbon monocyclic group having 3 to 10 carbon atoms, and examples thereof include a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, and a cycloheptyl group. The term “C₃-C₁₀ cycloalkylene group” as used herein refers to a divalent group having the same structure as that of the C₃-C₁₀ cycloalkyl group.

The term “C₁-C₁₀ heterocycloalkyl group” as used herein refers to a monovalent saturated monocyclic group having at least one heteroatom N, O, P, Si, and S as a ring-forming atom and 1 to 10 carbon atoms, and examples thereof include a tetrahydrofuran group, and a tetrahydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkylene group” as used herein refers to a divalent group having the same structure as the C₁-C₁₀ heterocycloalkyl group.

The term “C₃-C₁₀ cycloalkenyl group” as used herein refers to a monovalent monocyclic group that has 3 to 10 carbon atoms and at least one carbon-carbon double bond in the ring thereof and no aromaticity, and examples thereof include a cyclopentenyl group, a cyclohexenyl group, and a cycloheptenyl group. The term “C₃-C₁₀ cycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₃-C₁₀ cycloalkenyl group.

The term “C₂-C₁₀ heterocycloalkenyl group” as used herein refers to a monovalent monocyclic group that has at least one N, O, P, Si, B, Se, Ge, Te, S, or any combination thereof as a ring-forming atom, 2 to 10 carbon atoms, and at least one carbon-carbon double bond in its ring. Examples of the C₂-C₁₀ heterocycloalkenyl group are a 2,3-dihydrofuran group, and a 2,3-dihydrothiophenyl group. The term “C₁-C₁₀ heterocycloalkenylene group” as used herein refers to a divalent group having the same structure as the C₂-C₁₀ heterocycloalkenyl group.

The term “C₆-C₆₀ aryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms, and the term “C₆-C₆₀ arylene group” as used herein refers to a divalent group having a carbocyclic aromatic system having 6 to 60 carbon atoms. Examples of the C₆-C₆₀ aryl group include a phenyl group, a naphthyl group, an anthracenyl group, a phenanthrenyl group, a pyrenyl group, and a chrysenyl group. When the C₆-C₆₀ aryl group and the C₆-C₆₀ arylene group each include two or more rings, the rings may be fused to each other. The C₇-C₆₀ alkylaryl group refers to a C₆-C₆₀ aryl group substituted with at least one C₁-C₁₀ alkyl group.

The term “C₁-C₆₀ heteroaryl group” as used herein refers to a monovalent group having a carbocyclic aromatic system that has at least one N, O, P, Si, B, Se, Ge, Te, S, or any combination thereof as a ring-forming atom, and 1 to 60 carbon atoms. The term “C₁-C₁₀ heteroarylene group” as used herein refers to a divalent group having a carbocyclic aromatic system that has at least one N, O, P, Si, B, Se, Ge, Te, S, or any combination thereof as a ring-forming atom, and 1 to 60 carbon atoms. Examples of the C₁-C₁₀ heteroaryl group include a pyridinyl group, a pyrimidinyl group, a pyrazinyl group, a pyridazinyl group, a triazinyl group, a quinolinyl group, and an isoquinolinyl group. When the C₆-C₆₀ heteroaryl group and the C₆-C₆₀ heteroarylene group each include two or more rings, the rings may be fused to each other. The C₂-C₆₀ alkylheteroaryl group refers to a C₁-C₁₀ heteroaryl group substituted with at least one C₁-C₁₀ alkyl group.

The term “C₆-C₆₀ aryloxy group” as used herein indicates —OA₁₀₂ (wherein A₁₀₂ is the C₆-C₆₀ aryl group), and the term “C₆-C₆₀ arylthio group” as used herein indicates —SA₁₀₃ (wherein A₁₀₃ is the C₆-C₆₀ aryl group).

The term “monovalent non-aromatic condensed polycyclic group” as used herein refers to a monovalent group (for example, having 8 to 60 carbon atoms) having two or more rings condensed to each other, only carbon atoms as ring-forming atoms, and no aromaticity in its entire molecular structure. Examples of the monovalent non-aromatic condensed polycyclic group include a fluorenyl group. The term “divalent non-aromatic condensed polycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed polycyclic group.

The term “monovalent non-aromatic condensed heteropolycyclic group” as used herein refers to a monovalent group (for example, having 2 to 60 carbon atoms) having two or more rings condensed to each other, a heteroatom N, O, P, Si, and S, other than carbon atoms, as a ring-forming atom, and no aromaticity in its entire molecular structure.

Examples of the monovalent non-aromatic condensed heteropolycyclic group include a carbazolyl group. The term “divalent non-aromatic condensed heteropolycyclic group” as used herein refers to a divalent group having the same structure as the monovalent non-aromatic condensed heteropolycyclic group.

The term “C₅-C₃₀ carbocyclic group” as used herein refers to a saturated or unsaturated cyclic group having, as a ring-forming atom, 5 to 30 carbon atoms only. The C₅-C₃₀ carbocyclic group may be a monocyclic group or a polycyclic group.

The term “C₁-C₃₀ heterocyclic group” as used herein refers to a saturated or unsaturated cyclic group having, as a ring-forming atom, at least one N, O, Si, P, B, Se, Ge, Te, S, or any combination thereof, other than 1 to 30 carbon atoms. The C₁-C₃₀ heterocyclic group may be a monocyclic group or a polycyclic group.

73

At least one substituent of the substituted C₅-C₃₀ carbocyclic group, the substituted C₁-C₃₀ heterocyclic group, the substituted C₁-C₁₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₁₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₇-C₆₀ alkylaryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₁₀ heteroaryl group, the substituted C₂-C₆₀ alkyl heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group may be:

deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₇-C₆₀ alkylaryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a C₂-C₆₀ alkyl heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁₁)(Q₁₂), —Si(Q₁₃)(Q₁₄)(Q₁₅), —B(Q₁₆)(Q₁₇), —P(=O)(Q₁₈)(Q₁₉), or any combination thereof;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₇-C₆₀ alkylaryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a C₂-C₆₀ alkyl heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group;

a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₇-C₆₀ alkylaryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a C₂-C₆₀ alkyl heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ hetero-

74

cycloalkenyl group, a C₆-C₆₀ aryl group, a C₇-C₆₀ alkylaryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a C₂-C₆₀ alkyl heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, —N(Q₂₁)(Q₂₂), —Si(Q₂₃)(Q₂₄)(Q₂₅), —B(Q₂₆)(Q₂₇), —P(=O)(Q₂₈)(Q₂₉), or any combination thereof; or —N(Q₃₁)(Q₃₂), —Si(Q₃₃)(Q₃₄)(Q₃₅), —B(Q₃₆)(Q₃₇), or —P(=O)(Q₃₈)(Q₃₉),


wherein Q₁ to Q₉, Q₁₁ to Q₁₉, Q₂₁ to Q₂₉, and Q₃₁ to Q₃₉ may each independently be hydrogen, deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₁-C₆₀ alkyl group substituted with at least one deuterium, a C₁-C₆₀ alkyl group, and a C₆-C₆₀ aryl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryl group substituted with at least one deuterium, a C₁-C₆₀ alkyl group, and a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a C₂-C₆₀ alkyl heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group.

Hereinafter, a compound and an organic light-emitting device according to embodiments are described in detail with reference to Synthesis Example and Examples.

However, the organic light-emitting device is not limited thereto. The wording “‘B’ was used instead of ‘A’” used in describing Synthesis Examples means that an amount of ‘A’ used was identical to an amount of ‘B’ used, in terms of a molar equivalent.


EXAMPLES

Synthesis Example 1: Synthesis of Compound 7




75


-continued



7(2)



7(3)



7

76


organic layer was concentrated. The crude product was purified by column chromatography (EA/Hex(hexane) with gradient elution from 20% to 35% to complete the production of 2.2 g of Intermediate 7-3 (Yield of 70%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for C₅₂H₄₇N₃O₂: m/z 745.3668, Found: 745.3665.


(2) Synthesis of Compound 7

2.2 g (2.95 mmol) of Intermediate 7(3) and 1.5 g (3.54 mmol, 1.0 eq) of K₂PtCl₄ were mixed with 60 mL of AcOH and 20 mL of H₂O and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the precipitate was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layer was concentrated and the crude product was purified by column chromatography (MC 40%, EA 1%, Hex 59%) to obtain 1.7 g of Compound 6 (Yield of 61%). The obtained material was identified by Mass and HPLC.


Synthesis Example 2: Synthesis of Compound 1



1(1)





1(2)



1(3)

(1) Synthesis of Intermediate 7(3)

2.4 g (0.004 mmol, 1.2 eq) of Intermediate 7(1), 1.5 g (0.004 mol, 1 eq) of Intermediate 7(2), 0.34 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.7 g (0.015 mmol, 3 eq) of potassium carbonate were mixed with 80 mL of tetrahydrofuran (THF):distilled water (H₂O) at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, the precipitate was filtered therefrom to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O and the



(1) Synthesis of Intermediate 1(3)

2.0 g (0.005 mmol, 1.0 eq) of Intermediate 1(1), 1.6 g (0.005 mol, 1.2 eq) of Intermediate 1(2), 0.37 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.9 g (0.015 mmol, 3 eq) of potassium carbonate were mixed with 50 mL THF:H₂O at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, the mixture was filtered therefrom to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O and the organic layer was concentrated. The crude product was purified by column chromatography (EA/Hex(hexane) with gradient elution from 10% to 20% to complete the production of 2.2 g of Intermediate 1(3) (Yield of 67%). The obtained material was identified by Mass and HPLC.



HRMS(MALDI) calcd for C₃₅H₃₇N₃O₂: m/z 531.2886, Found: 531.2884.

(2) Synthesis of Compound 1

2.2 g (4.14 mmol) of Intermediate 1(3) and 2.1 g (4.97 mmol, 1.2 eq) of K₂PtCl₄ were mixed with 60 mL of AcOH and 20 mL of H₂O and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the mixture was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layer was concentrated and the crude product was purified by column chromatography (MC 40%, EA 1%, Hex 59%) was performed to obtain 1.3 g of Compound 6 (Yield of 43%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for C₃₅H₃₅N₃O₂Pt: m/z 724.2377, Found: 724.2378.

Synthesis Example 3: Synthesis of Compound 2



(1) Synthesis of Intermediate 2(3)

2.0 g (0.004 mmol, 1.0 eq) of Intermediate 2(1), 1.7 g (0.004 mol, 1.2 eq) of Intermediate 2(2), 0.31 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.6 g (0.015 mmol, 3 eq) of potassium carbonate were mixed with 80 mL THF:H₂O at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, the reaction mixture was filtered therefrom to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O. The organic layer was concentrated and the crude product was purified by column chromatography (EA/Hex(hexane) with gradient elution

79

from 10% to 30% to complete the production of 1.8 g of Intermediate 2(3) (Yield of 68%). The obtained material was identified by Mass and HPLC.


HRMS(MALDI) calcd for C₄₇H₄₅N₃O₂: m/z 683.3512, Found: 683.3513.

(2) Synthesis of Compound 2


1.8 g (2.63 mmol) of Intermediate 2(3) and 1.3 g (3.16 mmol, 1.2 eq) of K₂PtCl₄ were mixed with 80 mL AcOH: H₂O (3:1) were mixed and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the reaction mixture was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layer was concentrated and purified by column chromatography (MC 40%, EA 1%, Hex 59%) to obtain 1.1 g of Compound 2 (Yield of 48%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for C₄₇H₄₃N₃O₂Pt: m/z 876.3003, Found: 876.3006.


Synthesis Example 4: Synthesis of Compound 3



3(1)




3(2)



3(3)

80

-continued



3

(1) Synthesis of Intermediate 3(3)

2.0 g (0.005 mmol, 1.0 eq) of Intermediate 3(1), 2.0 g (0.005 mol, 1.2 eq) of Intermediate 3(2), 0.37 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.9 g (0.014 mmol, 3 eq) of potassium carbonate were mixed with 80 mL THF:H₂O at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, filtered to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O, and the organic layer was concentrated to provide the crude product. The crude product was purified by column chromatography (EA/Hex(hexane)) with gradient elution from 10% to 30% to complete the production of 2.0 g of Intermediate 3(3) (Yield of 73%). The obtained material was identified by Mass and HPLC.


HRMS(MALDI) calcd for: C₄₁H₄₂N₄O m/z 606.3359, Found: 606.3357.

(2) Synthesis of Compound 3

2.0 g (3.30 mmol) of Intermediate 3(3) and 1.6 g (3.96 mmol, 1.2 eq) of K₂PtCl₄ were mixed with 80 mL AcOH: H₂O (3:1) were mixed and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the reaction mixture was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layer was concentrated to provide the crude product, which was purified by column chromatography (MC 40%, EA 1%, Hex 59%) to obtain 1.3 g of Compound 3 (Yield of 49%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for: C₄₁H₄₀N₄OPt m/z 799.2850, Found: 799.2852.



Synthesis Example 5: Synthesis of Compound 4



4(1)

81

-continued



(1) Synthesis of Intermediate 4(3)

2.0 g (0.005 mmol, 1.0 eq) of Intermediate 4(1), 2.4 g (0.005 mol, 1.2 eq) of Intermediate 4(2), 0.37 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.9 g (0.014 mmol, 3 eq) of potassium carbonate were mixed with 80 mL of a THF:H₂O at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, the reaction mixture was filtered to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O and the organic layer was concentrated to provide the crude product, which was purified by column chromatography (EA/Hex(hexane)) with gradient elution from 10% to 30% to complete the production of 2.2 g of Intermediate 4(3) (Yield of 71%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for: C₄₈H₄₅N₃O m/z 679.3563, Found: 679.3565.

(2) Synthesis of Compound 4


2.2 g (3.24 mmol) of Intermediate 4(3) and 1.6 g (3.88 mmol, 1.2 eq) of K₂PtCl₄ were mixed with 80 mL AcOH: H₂O (3:1) and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the reaction mixture

82

was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layer was concentrated and the resulting crude product was purified by column chromatography (MC 40%, EA 1%, Hex 59%) to obtain 0.8 g of Compound 4 (Yield of 28%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for: C₄₈H₄₃N₃OPt m/z 872.3054, Found: 872.3056.

Synthesis Example 5: Synthesis of Compound 5



83

(1) Synthesis of Intermediate 5(3)

2.0 g (0.005 mmol, 1.0 eq) of Intermediate 4(1), 2.4 g (0.005 mol, 1.2 eq) of Intermediate 4(2), 0.37 g (0.001 mmol, 0.07 eq) of tetrakis(triphenylphosphine)palladium (0), and 1.9 g (0.014 mmol, 3 eq) of potassium carbonate were mixed with 80 mL THF:H₂O at a ratio of 3:1 and refluxed for 12 hours. The reaction mixture was cooled to room temperature, and then, the reaction mixture was filtered to obtain a filtrate. The filtrate was washed with ethyl acetate (EA)/H₂O, the organic layer was concentrated, and the resulting crude product was purified by column chromatography (EA/Hex(hexane)) with gradient elution from 10% to 30% to complete the production of 2.1 g of Intermediate 5(3) (Yield of 68%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for: C₄₈H₄₇N₃O m/z 681.3719, Found: 681.3718.

(2) Synthesis of Compound 5

2.1 g (3.08 mmol) of Intermediate 5(3) and 1.53 g (3.70 mmol, 1.2 eq) of K₂PtCl₄ were mixed with 80 mL AcOH:H₂O (3:1) were mixed and then refluxed for 16 hours. After cooling the reaction mixture to room temperature, the reaction mixture was filtered and the obtained precipitate was dissolved in methylene chloride (MC) and washed with H₂O. The organic layers was concentrated and the resulting crude product was purified by column chromatography (MC 40%, EA 1%, Hex 59%) to obtain 0.5 g of Compound 5 (Yield of 19%). The obtained material was identified by Mass and HPLC.

HRMS(MALDI) calcd for: C₄₈H₄₅N₃OPt m/z 874.3210, Found: 874.3211.

Example 1

As an anode, an ITO-patterned glass substrate was cut to a size of 50 mm×50 mm×0.5 mm, sonicated with isopropyl alcohol and pure water, each for 5 minutes, and then cleaned by exposure to ultraviolet rays and ozone for 30 minutes. Then, the ITO-patterned glass substrate was provided to a vacuum deposition apparatus.


HT3 and F6TCNNQ were vacuum-deposited at the weight ratio of 98:2 on the anode to form a hole injection layer having a thickness of 100 Å, and HT3 was vacuum-deposited on the hole injection layer to form a hole transport layer having a thickness of 1350 Å. H-H1 was deposited on the hole transport layer to form a hole transport layer having a thickness of 300 Å.

Then, H-H1, H-H2, and Compound 1(dopant) were co-deposited at the weight ratio of 47.5:47.5:5 on the hole transport layer to form an emission layer having a thickness of 400 Å.


Then, ET3 and ET-D1 were co-deposited at the volume ratio of 50:50 on the emission layer to form an electron transport layer having a thickness of 350 Å, and ET-D1 was vacuum-deposited on the electron transport layer to form an electron injection layer having a thickness of 10 Å, and Al was vacuum-deposited on the electron injection layer to form a cathode having a thickness of 1000 Å, thereby completing the manufacture of an organic light-emitting device.

84


HT3



F6-TCNNQ






H-H1



85

-continued



Examples 2 to 6 and Comparative Examples 1 and 2

Organic light-emitting devices were manufactured in the same manner as in Example 1, except that Compounds shown in Table 2 were each used instead of Compound 1 as a dopant in forming an emission layer.




Evaluation Example 1: Evaluation of Characteristics of Organic Light-Emitting Devices

The driving voltage, the luminescence quantum efficiency, and the lifespan (T_{95}) of each of the organic light-emitting devices manufactured according to Examples 1 to 6 and Comparative Examples 1 and 2 were evaluated. Results thereof are shown in Table 2. This evaluation was performed using a current-voltage meter (Keithley 2400) and a luminescence meter (Minolta Cs-1,000A), and the lifespan (T_{95}) (at 6000 nit) was evaluated by measuring, as a relative value (%), the amount of time that elapsed until luminance was reduced to 95% of the initial brightness of 100%.

86

TABLE 2

			Luminescence		
5		Driving	quantum		
	Dopant	voltage	efficiency	lifespan (T ₉₅)	
	Compound	(Relative	(Relative	(Relative	
	No.	value, %)	value, %)	value, %)	
10	Example 1	1	76%	78%	40%
	Example 2	2	77%	95%	57%
	Example 3	3	80%	73%	50%
	Example 4	4	79%	69%	28%
15	Example 5	5	79%	67%	35%
	Example 6	7	77%	100%	50%
	Comparative	B	93%	40%	8%
	Example 1				
20	Comparative	C	100%	27%	1%
	Example 2				




3


87

TABLE 2-continued


No.	Dopant Compound No.	Driving voltage (Relative value, %)	Luminescence quantum efficiency (Relative value, %)	lifespan (T ₉₅) (Relative value, %)




4



5



7




B

88

TABLE 2-continued

No.	Dopant Compound No.	Driving voltage (Relative value, %)	Luminescence quantum efficiency (Relative value, %)	lifespan (T ₉₅) (Relative value, %)



C

From Table 2, it was confirmed that the organic light-emitting devices of Example 1 to 6 have a lower driving voltage, higher luminescence quantum efficiency, and a longer lifespan, than the organic light-emitting devices of Comparative Examples 1 and 2.


The organometallic compounds have excellent electrical characteristics and/or thermal stability. Accordingly, an organic light-emitting device using the organometallic compounds has low driving voltage and excellent luminescence quantum efficiency and lifespan characteristics. Such organometallic compounds have excellent phosphorescent luminescent characteristics, and thus, when used, a diagnostic composition having a high diagnostic efficiency may be provided.

It should be understood that embodiments described herein should be considered in a descriptive sense only and not for purposes of limitation. Descriptions of features or aspects within each embodiment should typically be considered as available for other similar features or aspects in other embodiments. While one or more embodiments have been described with reference to the figures, it will be understood by those of ordinary skill in the art that various changes in form and details may be made therein without departing from the spirit and scope as defined by the following claims.

What is claimed is:

1. An organometallic compound represented by Formula 1:

Formula 1



89


wherein, in Formula 1,

M₁ is beryllium (Be), magnesium (Mg), aluminum (Al), calcium (Ca), titanium (Ti), manganese (Mn), cobalt (Co), copper (Cu), zinc (Zn), gallium (Ga), germanium (Ge), zirconium (Zr), ruthenium (Ru), rhodium (Rh), palladium (Pd), silver (Ag), rhenium (Re), platinum (Pt), or gold (Au),







ring A₁ and ring A₃ are each independently a C₅-C₃₀ carbocyclic group or a C₂-C₆₀ heterocyclic group,

ring A₄ is a C₂-C₆₀ heterocyclic group,

wherein a moiety represented by





in Formula 1 is a group represented by one of Formulae CY2-1 to CY2-8:



90

-continued



wherein, in Formulae CY2-1 to CY2-8,

*, *, and * are each independently a binding site to a neighboring atom,

X₁ is N, O, or S,

X₃ is C, and X₄ is N,

a bond between X₁ and M₁, a bond between X₁ and M₁, a bond between X₃ and M₁, and a bond between X₄ and M₁ are each independently a covalent bond or a coordination bond,

Y₁ to Y₅ are each C,

T₁ and T₂ are each independently a single bond, *—N [L₁]_{a1}—(R₁)_{b1}—*, *—B(R₁)—*, *—P(R₁)—*, *—C(R₁)(R₂)—*, *—Si(R₁)(R₂)—*, *—Ge(R₁)(R₂)—*, *—S—*, *—Se—*, *—O—*, *—C(=O)—*, *—S(=O)—*, *—S(=O)₂—*, *—C(R₁)=*, *—C(R₁)—*, *—C(R₁)=C(R₂)—*, *—C(=S)—*, or *—C≡C—*,

T₃ is *—N[L₁]_{a1}—(R₁)_{b1}—*, *—B(R₁)—*, *—P(R₁)—*, *—C(R₁)(R₂)—*, *—Si(R₁)(R₂)—*, *—Ge(R₁)(R₂)—*, *—S—*, *—Se—*, *—O—*, *—C(=O)—*, *—S(=O)—*, *—S(=O)₂—*, *—C(R₁)=*, *—C(R₁)—*, *—C(R₁)=C(R₂)—*, *—C(=S)—*, or *—C≡C—*,

L₁ is a single bond, a substituted or unsubstituted C₅-C₃₀ carbocyclic group, or a substituted or unsubstituted C₁-C₃₀ heterocyclic group,

a₁ is an integer from 1 to 3, wherein when a₁ is 2 or more, two or more L₁ (s) are identical to or different from each other,


b₁ is an integer from 1 to 5, wherein when b₁ is 2 or more, two or more R₁ (s) are identical to or different from each other,

R₁ and R₂ are optionally linked to each other via a single bond, a double bond, or a first linking group to form a substituted or unsubstituted C₅-C₃₀ carbocyclic group or a substituted or unsubstituted C₁-C₃₀ heterocyclic group,

R₁, R₂, R₃₀, and R₄₀ are each independently hydrogen, deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl

91

group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(O₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉), R₂₁ and R₂₂ are each independently deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(O₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉), provided that when a moiety represented by




in Formula 1 is a group represented by Formula CY2-2, then R₂₁ and R₂₂ are different, b10 is an integer from 2 to 10, Each R₁₀ is independently deuterium, —F, —Cl, —Br, —I, —SF₅, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a substituted or unsubstituted C₁-C₆₀ alkyl group, a substituted or unsubstituted C₂-C₆₀ alkenyl group, a substituted or unsubstituted C₂-C₆₀ alkynyl group, a substituted or unsubstituted C₁-C₆₀ alkoxy group, a substituted or unsubstituted C₃-C₁₀ cycloalkyl group, a substituted or unsubstituted C₁-C₁₀ heterocycloalkyl group, a substituted or unsubstituted C₃-C₁₀ cycloalkenyl group, a substituted or unsubstituted C₂-C₁₀ heterocycloalkenyl group, a substituted or unsubstituted C₆-C₆₀ aryl group, a substituted or unsubstituted C₆-C₆₀ aryloxy group, a substituted or unsubstituted C₆-C₆₀ arylthio group, a substituted or unsubstituted C₁-C₆₀ heteroaryl group, a substituted or unsubstituted monovalent non-aromatic condensed polycyclic group, a substituted or unsubstituted monovalent non-aromatic condensed heteropolycyclic group, —N(Q₁)(Q₂), —Si(Q₃)(Q₄)(O₅), —B(Q₆)(Q₇), or —P(=O)(Q₈)(Q₉),

92

b30, and b40 are each independently an integer from 1 to 10,

R₂₁ and R₂₂ are optionally linked together to form a C₅-C₃₀ carbocyclic that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a}, provided that wherein a moiety represented by



in Formula 1 is a group represented by one of Formulae CY2-4 and CY2-8, then R₂₁ and R₂₂ are not optionally linked to form a C₆-carbocyclic group,

- i) R₃₀ and at least one of R₁ and R₂ are optionally linked together to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a},
- ii) R₄₀ and at least one of R₁ and R₂ are optionally linked together to form a C₅-C₃₀ carbocyclic group that is unsubstituted or substituted with at least one R_{10a} or a C₁-C₃₀ heterocyclic group that is unsubstituted or substituted with at least one R_{10a},

R_{10a} is the same as explained in connection with R₁₀, and at least one substituent of the substituted C₅-C₃₀ carbocyclic group, the substituted C₁-C₃₀ heterocyclic group, the substituted C₁-C₆₀ alkyl group, the substituted C₂-C₆₀ alkenyl group, the substituted C₂-C₆₀ alkynyl group, the substituted C₁-C₆₀ alkoxy group, the substituted C₃-C₁₀ cycloalkyl group, the substituted C₁-C₁₀ heterocycloalkyl group, the substituted C₃-C₁₀ cycloalkenyl group, the substituted C₂-C₁₀ heterocycloalkenyl group, the substituted C₆-C₆₀ aryl group, the substituted C₆-C₆₀ aryloxy group, the substituted C₆-C₆₀ arylthio group, the substituted C₁-C₆₀ heteroaryl group, the substituted monovalent non-aromatic condensed polycyclic group, and the substituted monovalent non-aromatic condensed heteropolycyclic group is:

deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group;

a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, or a C₁-C₆₀ alkoxy group, each substituted with at least one of deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₃-C₁₀ cycloalkyl group, a C₁-C₁₀ heterocycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic

93

condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{N}(\text{Q}_{11})$ (Q_{12}), $-\text{Si}(\text{Q}_{13})(\text{Q}_{14})(\text{Q}_{15})$, $-\text{B}(\text{Q}_{16})(\text{Q}_{17})$, $-\text{P}(=\text{O})(\text{Q}_{18})(\text{Q}_{19})$, or any combination thereof;

a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group;

a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group, each substituted with at least one of deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, $-\text{CD}_3$, $-\text{CD}_2\text{H}$, $-\text{CDH}_2$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, $-\text{N}(\text{Q}_{21})(\text{Q}_{22})$, $-\text{Si}(\text{Q}_{23})(\text{Q}_{24})(\text{Q}_{25})$, $-\text{B}(\text{Q}_{26})(\text{Q}_{27})$, $-\text{P}(=\text{O})(\text{Q}_{28})(\text{Q}_{29})$, or any combination thereof; or $-\text{N}(\text{Q}_{31})(\text{Q}_{32})$, $-\text{Si}(\text{Q}_{33})(\text{Q}_{34})(\text{Q}_{35})$, $-\text{B}(\text{Q}_{36})(\text{Q}_{37})$, or $-\text{P}(=\text{O})(\text{Q}_{38})(\text{Q}_{39})$,


wherein Q_1 to Q_9 , Q_{11} to Q_{19} , Q_{21} to Q_{29} and Q_{31} to Q_{39} are each independently hydrogen, deuterium, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{I}$, a hydroxyl group, a cyano group, a nitro group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a $\text{C}_1\text{-C}_{60}$ alkyl group, a $\text{C}_2\text{-C}_{60}$ alkenyl group, a $\text{C}_2\text{-C}_{60}$ alkynyl group, a $\text{C}_1\text{-C}_{60}$ alkoxy group, a $\text{C}_3\text{-C}_{10}$ cycloalkyl group, a $\text{C}_1\text{-C}_{10}$ heterocycloalkyl group, a $\text{C}_3\text{-C}_{10}$ cycloalkenyl group, a $\text{C}_2\text{-C}_{10}$ heterocycloalkenyl group, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryl group substituted with at least one a $\text{C}_1\text{-C}_{60}$ alkyl group, or any combination thereof, a $\text{C}_6\text{-C}_{60}$ aryl group, a $\text{C}_6\text{-C}_{60}$ aryloxy group, a $\text{C}_6\text{-C}_{60}$ arylthio group, a $\text{C}_1\text{-C}_{60}$ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, or a monovalent non-aromatic condensed heteropolycyclic group.

2. The organometallic compound of claim 1, wherein ring A_4 is an azafluorene group, an azacarbazole group, an azadibenzofuran group, an azadibenzothiophene group, an azadibenzosilole group, a pyridine group, a pyrimidine group, a pyrazine group, a pyridazine group, a triazine group, a quinoline group, an isoquinoline group, a quinoxaline group, aquinazoline group, a phenanthroline group, a pyrrole group, a pyrazole group, an imidazole group, a triazole group, a tetrazole group, an oxazole group, an isooxazole group, a thiazole group, an isothiazole group, an oxadiazole group, a thiadiazole group, a benzopyrazole group, a benzimidazole group, an indazole group, a benzoxazole group, a benzothiazole group, a benzoxadiazole

94


group, a benzothiadiazole group, a benzotriazole group, a diazaindene group, a triazaindene group, a 5,6,7,8-tetrahydroisoquinoline group, or a 5,6,7,8-tetrahydroquinoline group.

3. The organometallic compound of claim 1, wherein a moiety represented by




in Formula 1 is a group represented by one of Formulae CY1-1 to CY1-4 and CY1-11 to CY1-16:


CY1-1




CY1-2




CY1-3



CY1-4








CY1-11



95

-continued



wherein, in Formulae CY1-1 to CY1-16,

X₁₁ is N or C(R₁₁), X₁₂ is N or C(R₁₂), X₁₃ is N or C(R₁₃),
X₁₄ is N or C(R₁₄), X₁₅ is N or C(R₁₅), X₁₆ is N or
C(R₁₆), X₁₇ is N or C(R₁₇), and X₁₈ is N or C(R₁₈),

X₁₉ is C(R_{19a})(R_{19b}), N(R₁₉), O, S, or Si(R_{19a})(R_{19b}),

R₁₁ to R₁₉, R_{19a}, and R_{19b} are each independently the
same as described in connection with R₁₀ in claim 1,
and


* and *' each indicate a binding site to a neighboring
atom.

96

4. The organometallic compound of claim 1, wherein a
moiety represented by

CY1-12

5




10

in Formula 1 is a group represented by one of Formulae
CY3-1 to CY3-11:


CY1-13 15

20



CY3-1


25



CY3-2

CY1-14

30




CY3-3

35

CY1-15 40


45



CY3-4


CY1-16

50



CY3-5






55



CY3-6

97

-continued



wherein, in Formulae CY3-1 to CY3-11,

X_{31} is N or C(R_{31}), X_{32} is N or C(R_{32}), X_{33} is N or C(R_{33}), X_{34} is N or C(R_{34}), X_{35} is N or C(R_{35}), X_{36} is N or C(R_{36}), and X_{37} is N or C(R_{37}),

X_{39} is C(R_{39a})(R_{39b}), N(R_{39}), O, S, or Si(R_{39a})(R_{39b}),


R_{31} to R_{39} , R_{39a} , and R_{39b} are each independently the same as described in connection with R_{30} in claim 1, and

*, *', and *'' are each independently a binding site to a neighboring atom.

98

5. The organometallic compound of claim 1, wherein a moiety represented by

CY3-7



in Formula 1 is a group represented by one of Formulae CY4-1 to CY4-4:

CY3-8

15

CY4-1

20

CY4-2

CY3-9

25

CY4-3

30

35

CY3-10

40

CY4-4

CY3-11

45

wherein, in Formulae CY4-1 to CY4-84,

X_{41} is N or C(R_{41}), X_{42} is N or C(R_{42}), X_{43} is N or C(R_{43}), X_{44} is N or C(R_{44}), X_{45} is N or C(R_{45}), X_{46} is N or C(R_{46}), X_{47} is N or C(R_{47}), and X_{48} is N or C(R_{48}),

R_{41} to R_{48} are each independently the same as described in connection with R_{40} in claim 1, and

* and *' each indicate a binding site to a neighboring atom.

6. The organometallic compound of claim 1, wherein T_3 is $^*-\text{N}[(L_1)_{a1}-(R_1)_{b1}]-^*$, $^*-\text{C}(R_1)(R_2)-^*$, $^*-\text{Si}(R_1)(R_2)-^*$, $^*-\text{O}-^*$, or $^*-\text{S}-^*$,

L_1 is:

a phenylene group, a pentalenylene group, an indenylene group, a naphthylene group, an azulenylene group, a heptalenylene group, an acenaphthylene group, a fluorenylene group, a phenalenylene group, a phenanthrenylene group, an anthracenylene group, a fluoranthenylene group, a triphenylenylene group, a pyrenylene

group, a chrysenylenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, or a pentacenylenylene group; or
 a phenylene group, a pentalenylenylene group, an indenylene group, a naphthylene group, an azulenylenylene group, a heptalenylenylene group, an acenaphthylenylene group, a fluorenylene group, a phenalenylenylene group, a phenanthrenylene group, an anthracenylenylene group, a fluoranthenylenylene group, a triphenylenylene group, a pyrenylene group, a chrysenylenylene group, a naphthacenylenylene group, a picenylene group, a perylenylene group, or a pentacenylenylene group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₆₀ alkyl group, a C₂-C₆₀ alkenyl group, a C₂-C₆₀ alkynyl group, a C₁-C₆₀ alkoxy group, a C₃-C₁₀ cycloalkyl group, a C₃-C₁₀ cycloalkenyl group, a C₁-C₁₀ heterocycloalkyl group, a C₂-C₁₀ heterocycloalkenyl group, a C₆-C₆₀ aryl group, a C₆-C₆₀ aryloxy group, a C₆-C₆₀ arylthio group, a C₁-C₆₀ heteroaryl group, a monovalent non-aromatic condensed polycyclic group, a monovalent non-aromatic condensed heteropolycyclic group, or any combination thereof,

R₁ and R₂ are each independently:


a C₁-C₃₀ alkyl group;

a C₁-C₃₀ alkyl group, substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a dibenzosilolyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group; or
















a phenyl group, a naphthyl group, a fluorenyl group, a phenanthrenyl group, an anthracenyl group, a fluoranthenyl group, a triphenylenyl group, a pyrenyl group, a chrysenyl group, a pyrrolyl group, a thiophenyl group, a furanyl group, an imidazolyl group, a pyrazolyl group, a thiazolyl group, an isothiazolyl group, an oxazolyl group, an isoxazolyl group, a pyridinyl group, a pyrazinyl group, a pyrimidinyl group, a pyridazinyl group, an isoindolyl group, an indolyl group, an indazolyl group, a purinyl group, a quinolinyl group, an isoquinolinyl group, a benzoquinolinyl group, a quinoxalinyl group, a quinazolinyl group, a cinnolinyl group, a carbazolyl group, a phenanthrolinyl group, a benzimidazolyl group, a benzofuranyl group, a benzothiophenyl group, an isobenzothiazolyl group, a benzoxazolyl group, an isobenzoxazolyl group, a triazolyl group, a tetrazolyl group, an oxadiazolyl group, a triazinyl group, a dibenzofuranyl group, a dibenzothiophenyl group, a benzocarbazolyl group, a dibenzocarbazolyl group, an imidazopyridinyl group, or an imidazopyrimidinyl group, each substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano group, a nitro group, an amino group, an amidino group, a hydrazine group, a hydrazone group, a carboxylic acid or a salt thereof, a sulfonic acid or a salt thereof, a phosphoric acid or a salt thereof, a C₁-C₁₀ alkyl group, a C₁-C₁₀ alkoxy group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group, an adamantanyl group, a norbornanyl group, a norbornenyl group, a cyclopentenyl group, a cyclohexenyl group, a cycloheptenyl group, a phenyl group, a naphthyl group, a pyridinyl group, a pyrimidinyl group, or any combination thereof.

7. The organometallic compound of claim 1, wherein R₁, R₂, R₃₀, and R₄₀ are each independently hydrogen, deuterium, —F, a cyano group, a nitro group, —SF₅, —CH₃, —CD₃, —CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a group represented by one of Formulae 9-1 to 9-19, a group represented by one of Formulae 10-1 to 10-46, or —Si(Q₃)(Q₄)(Q₅):



101


-continued



102

-continued


9-7



10-3

5


9-8



10-4

10


9-9



10-5

15

9-10




10-6

9-11

20

9-12




10-7

9-13

25


9-14



10-8

30


9-15



10-9

35


9-16



10-10

40


9-17



10-11

45


9-18



10-12

50


9-19



10-13

55


10-1



10-14

60











10-2



65

103

-continued




104

-continued

10-15


5



10-25

10-16


10



10-26

10-17


15



10-27

10-18


20



10-28

10-19


25



10-29

10-20


30



10-30

10-21


35



10-31

10-22


40



10-32

10-23


45



10-33








10-24

50



105


-continued

**106**

-continued

10-34


5



10

10-35


15



20

10-36


25



30

10-37


35



40

10-38


45



50

10-39

55



and

10-40

60

* in Formulae 9-1 to 9-19 and 10-1 to 10-46 indicates a binding site to a neighboring atom.

8. The organometallic compound of claim 7, wherein at least one R₁₀ is of Formulae 9-1 to 9-19.

9. The organometallic compound of claim 7, wherein at least one R₂₁ to R₂₂ is of Formulae 10-1 to 10-46.

10. The organometallic compound of claim 1, wherein ring A₁ and ring A₃ are each a benzene group and ring A₄ is a pyridine group,

107

a bond between X_2 and M_1 and a bond between X_4 and M_1 are each a coordination bond, and a bond between X_3 and M_1 is a covalent bond,

Y_1 to Y_5 are each C,

at least one of R_1 , R_2 , R_{10} , R_{21} to R_{22} , R_{30} , and R_{40} is:

a C_1 - C_{30} alkyl group;

a C_1 - C_{30} alkyl group substituted with at least one deuterium, —F, —Cl, —Br, —I, —CD₃, —CD₂H, —CDH₂,

—CF₃, —CF₂H, —CFH₂, a hydroxyl group, a cyano

group, a nitro group, an amino group, an amidino

group, a hydrazine group, a hydrazone group, a car-

boxylic acid or a salt thereof, a sulfonic acid or a salt

thereof, a phosphoric acid or a salt thereof, a C_1 - C_{10}

alkyl group, a C_1 - C_{10} alkoxy group, a cyclopentyl

group, a cyclohexyl group, a cycloheptyl group, a

cyclooctyl group, an adamantanyl group, a norbornanyl

group, a norbornenyl group, a cyclopentenyl group, a

cyclohexenyl group, a cycloheptenyl group, a phenyl

group, a naphthyl group, a pyridinyl group, a pyrim-

idinyl group, or any combination thereof;

a cyclopentyl group, a cyclohexyl group, a cycloheptyl

group, a cyclooctyl group, an adamantanyl group, a

norbornanyl group, a norbornenyl group, a cyclopen-

tenyl group, a cyclohexenyl group, a cycloheptenyl

group, a phenyl group, a naphthyl group, a fluorenyl

group, a phenanthrenyl group, an anthracenyl group, a

fluoranthenyl group, a triphenylenyl group, a pyrenyl

group, a chrysenyl group, a pyrrolyl group, a thiophe-

nyl group, a furanyl group, an imidazolyl group, a

pyrazolyl group, a thiazolyl group, an isothiazolyl

group, an oxazolyl group, an isoxazolyl group, a pyridi-

nyl group, a pyrazinyl group, a pyrimidinyl group, a

pyridazinyl group, an isoindolyl group, an indolyl

group, an indazolyl group, a purinyl group, a quinolinyl

group, an isoquinolinyl group, a benzoquinolinyl

group, a quinoxalinyl group, a quinazolinyl group, a

cinnolinyl group, a carbazolyl group, a phenanthrolinyl

group, a benzimidazolyl group, a benzofuranyl group,

a benzothiophenyl group, an isobenzothiazolyl group, a

benzoxazolyl group, an isobenzoxazolyl group, a tri-

azolyl group, a tetrazolyl group, an oxadiazolyl group,

a triazinyl group, a dibenzofuranyl group, a dibenzo-

thiophenyl group, a benzocarbazolyl group, a dibenzo-

carbazolyl group, an imidazopyridinyl group, or an

imidazopyrimidinyl group; and

a cyclopentyl group, a cyclohexyl group, a cycloheptyl

group, a cyclooctyl group, an adamantanyl group, a

norbornanyl group, a norbornenyl group, a cyclopen-

tenyl group, a cyclohexenyl group, a cycloheptenyl

group, a phenyl group, a naphthyl group, a fluorenyl

group, a phenanthrenyl group, a triphenylenyl group, a pyrenyl

group, a chrysenyl group, a pyrrolyl group, a thiophe-

nyl group, a furanyl group, an imidazolyl group, a

pyrazolyl group, a thiazolyl group, an isothiazolyl

group, an oxazolyl group, an isoxazolyl group, a pyridi-

nyl group, a pyrazinyl group, a pyrimidinyl group, a

pyridazinyl group, an isoindolyl group, an indolyl

group, an indazolyl group, a purinyl group, a quinolinyl

group, an isoquinolinyl group, a benzoquinolinyl

group, a quinoxalinyl group, a quinazolinyl group, a

cinnolinyl group, a carbazolyl group, a phenanthrolinyl


group, a benzimidazolyl group, a benzofuranyl group,

a benzothiophenyl group, an isobenzothiazolyl group, a

108

benzoxazolyl group, an isobenzoxazolyl group, a tri-
azolyl group, a tetrazolyl group, an oxadiazolyl group,
a triazinyl group, a dibenzofuranyl group, a dibenzo-
thiophenyl group, a benzocarbazolyl group, a dibenzo-
carbazolyl group, an imidazopyridinyl group, and an
imidazopyrimidinyl group, each substituted with at
least one deuterium, —F, —Cl, —Br, —I, —CD₃,
—CD₂H, —CDH₂, —CF₃, —CF₂H, —CFH₂, a
hydroxyl group, a cyano group, a nitro group, an amino
group, an amidino group, a hydrazine group, a hydra-
zone group, a carboxylic acid or a salt thereof, a
sulfonic acid or a salt thereof, a phosphoric acid or a
salt thereof, a C_1 - C_{20} alkyl group, a C_1 - C_{20} alkoxy
group, a cyclopentyl group, a cyclohexyl group, a
cycloheptyl group, a cyclooctyl group, an adamantanyl
group, a norbornanyl group, a norbornenyl group, a
cyclopentenyl group, a cyclohexenyl group, a cyclo-
heptenyl group, a phenyl group, a naphthyl group, a
fluorenyl group, a phenanthrenyl group, an anthracenyl
group, a fluoranthenyl group, a triphenylenyl group, a
pyrenyl group, a chrysenyl group, a pyrrolyl group, a
thiophenyl group, a furanyl group, an imidazolyl group,
a pyrazolyl group, a thiazolyl group, an isothiazolyl
group, an oxazolyl group, an isoxazolyl group, a pyridi-
nyl group, a pyrazinyl group, a pyrimidinyl group, a
pyridazinyl group, an isoindolyl group, an indolyl
group, an indazolyl group, a purinyl group, a quinolinyl
group, an isoquinolinyl group, a benzoquinolinyl
group, a quinoxalinyl group, a quinazolinyl group, a
cinnolinyl group, a carbazolyl group, a phenanthrolinyl
group, a benzimidazolyl group, a benzofuranyl group,
a benzothiophenyl group, an isobenzothiazolyl group, a
benzoxazolyl group, an isobenzoxazolyl group, a tri-
azolyl group, a tetrazolyl group, an oxadiazolyl group,
a triazinyl group, a dibenzofuranyl group, a dibenzo-
thiophenyl group, a benzocarbazolyl group, a dibenzo-
carbazolyl group, an imidazopyridinyl group, or an
imidazopyrimidinyl group.


11. The organometallic compound of claim 1, wherein the organometallic compound is of Compounds 1 to 5, 7 to 9, and 12:




1

109

-continued


**110**

-continued



111

-continued



12. An organic light-emitting device comprising:
 a first electrode;
 a second electrode; and
 an organic layer including an emission layer between the
 first electrode and the second electrode,
 wherein the organic layer comprises at least one organo-
 metallic compound of claim 1.
13. The organic light-emitting device of claim 12, wherein
 the first electrode is an anode,
 the second electrode is a cathode,

112

the organic layer further comprises a hole transport region
 between the first electrode and the emission layer and
 an electron transport region between the emission layer
 and the second electrode,

the hole transport region comprises a hole injection layer,
 a hole transport layer, an electron blocking layer, or any
 combination thereof, and

the electron transport region comprises a hole blocking
 layer, an electron transport layer, an electron injection
 layer, or any combination thereof.

14. The organic light-emitting device of claim 12, wherein
 the organometallic compound is included in the emission
 layer.

15. The organic light-emitting device of claim 14, wherein
 the emission layer further comprises a host and the amount
 by weight of the host is greater than the amount by weight
 of the organometallic compound.

16. The organic light-emitting device of claim 14, wherein
 the emission layer emits green light having a maximum
 emission wavelength of about 490 nm to about 580 nm or
 red light having a maximum emission wavelength of about
 580 nm to about 780 nm.

17. A diagnostic composition comprising at least one
 organometallic compound of claim 1.

* * * * *