



US 20250277059A1

(19) **United States**(12) **Patent Application Publication**
SAITO et al.(10) **Pub. No.: US 2025/0277059 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **ANTIBODY USED TO TREAT
CORONAVIRUS INFECTION**(22) PCT Filed: **Apr. 12, 2023**(86) PCT No.: **PCT/JP2023/014855**

§ 371 (c)(1),

(2) Date: **Oct. 10, 2024**(71) Applicants: **RIKEN**, Wako-shi, Saitama (JP); **The University of Tokyo**, Bunkyo-ku, Tokyo (JP); **National University Corporation Shiga University of Medical Science**, Otsu-shi, Shiga (JP)(30) **Foreign Application Priority Data**

Apr. 12, 2022 (JP) 2022-065547

(72) Inventors: **Takashi SAITO**, Saitama (JP); **Michishige HARADA**, Saitama (JP); **Mikako SHIROUZU**, Saitama (JP); **Takehisa MATSUMOTO**, Saitama (JP); **Akiko IDEI**, Saitama (JP); **Jun-ichiro INOUE**, Tokyo (JP); **Mizuki YAMAMOTO**, Tokyo (JP); **Jin GOHDA**, Tokyo (JP); **Yasushi KAWAGUCHI**, Tokyo (JP); **Yasushi ITO**, Shiga (JP); **Hirohito ISHIGAKI**, Shiga (JP); **Misako NAKAYAMA**, Shiga (JP); **Yoshinori KITAGAWA**, Shiga (JP)**Publication Classification**(51) **Int. Cl.****C07K 16/40** (2006.01)**A61K 39/00** (2006.01)**A61P 31/14** (2006.01)(52) **U.S. Cl.**CPC **C07K 16/40** (2013.01); **A61P 31/14** (2018.01); **A61K 2039/505** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/33** (2013.01); **C07K 2317/34** (2013.01); **C07K 2317/55** (2013.01); **C07K 2317/732** (2013.01); **C07K 2317/76** (2013.01); **C07K 2317/92** (2013.01)(73) Assignees: **RIKEN**, Wako-shi, Saitama (JP); **The University of Tokyo**, Bunkyo-ku, Tokyo (JP); **National University Corporation Shiga University of Medical Science**, Otsu-shi, Shiga (JP)

(57)

ABSTRACT

The present invention provides an antibody that binds to an extracellular domain of TMPRSS2, capable of inhibiting beta coronavirus infection of cells, or an antigen-binding fragment thereof, and a composition including the antibody or an antigen-binding fragment thereof.



Specification includes a Sequence Listing.(21) Appl. No.: **18/855,890**Target cells: Daudi **human TMPRSS2**Target cells: Daudi **Cynomolgus Monkey TMPRSS2**

FIG. 1




FIG. 2




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 4




FIG. 5A

SPD

1: Futhan binding site

Name	Sequence	752_1	2228_15	1864_10	2020_12		
Peptide_181	AKAYRPGVYGNVMVFTDWIY		☐		☐	Futhan	SEQ ID NO: 41
Peptide_182	AYRPGVYGNVMVFTDWIYRQ		☐		☐	Futhan	SEQ ID NO: 42
Peptide_183	RPGVYGNVMVFTDWIYRQMR		☐	☐	☐	Futhan	SEQ ID NO: 43
Peptide_184	GVYGNVMVFTDWIYRQMRAD		☐		☐		SEQ ID NO: 44
Peptide_185	VYGNVMVFTDWIYRQMRADG		☐				SEQ ID NO: 45

SEQ ID NO: 85 : **GVYGNVMVFTDWIY**

2 : Active center of Futhan binding site

Name	Sequence	752_1	2228_15	1864_10	2020_12		
Peptide_168	DSGGPLVTSKNNIWWLIGDT	☐			☐	Active center/Futhan	SEQ ID NO: 46
Peptide_169	GGPLVTSKNNIWWLIGDTSW	☐					SEQ ID NO: 47
Peptide_170	PLVTSKNNIWWLIGDTSWGS	☐					SEQ ID NO: 48
Peptide_171	VTSKNNIWWLIGDTSWGS GC	☐				Futhan	SEQ ID NO: 49
Peptide_172	SKNNIWWLIGDTSWGS GCAK	☐				Futhan	SEQ ID NO: 50
Peptide_173	NNIWWLIGDTSWGS GCAKAY	☐				Futhan	SEQ ID NO: 51

SEQ ID NO: 86 : **PLVTSKNNIWWLIGDTSW**

3

Name	Sequence	752_1	2228_15	1864_10	2020_12		
Peptide_146	VLNAAKVLLIETQRCNSRYV	☐	☐		☐		SEQ ID NO: 52
Peptide_147	NAAKVLLIETQRCNSRYVYD		☐		☐		SEQ ID NO: 53
Peptide_148	AKVLLIETQRCNSRYVYDNL	☐	☐		☐		SEQ ID NO: 54
Peptide_149	VLLIETQRCNSRYVYDNLIT	☐	☐		☐	1258 chimera	SEQ ID NO: 55

SEQ ID NO: 87 : **VLLIETQRCNSRYV**

4

SPD

Name	Sequence	752_1	2228_15	1864_10	2020_12		
Peptide_102	WTAFAgilRQSFMFYgAGYQ				☐	Chimera (1831)	SEQ ID NO: 56
Peptide_103	AFAGILRQSFMFYgAGYQVE				☐	Chimera (1831)	SEQ ID NO: 57
Peptide_104	AGILRQSFMFYgAGYQVEKV				☐	Chimera (1831)	SEQ ID NO: 58
Peptide_105	ILRQSFMFYgAGYQVEKVIS	☐	☐	☐	☐		SEQ ID NO: 59
Peptide_106	RQSFMFYgAGYQVEKVISHP	☐	☐		☐		SEQ ID NO: 60
Peptide_107	SFMFYgAGYQVEKVISHPNY	☐	☐		☐		SEQ ID NO: 61
Peptide_108	MFYgAGYQVEKVISHPNYDS				☐		SEQ ID NO: 62
Peptide_109	YGAGYQVEKVISHPNYDSKT				☐		SEQ ID NO: 63

SEQ ID NO: 88 : **MFYgAGYQVEKV**

5, 6

Name	Sequence	752_1	2228_15	1864_10	2020_12		
Peptide_093	WVTAAHcVEKPLNNPWHWT	☐		☐		Active center/Chimera (1831)	SEQ ID NO: 64
Peptide_094	VTAAHCVEKPLNNPWHWTAF	☐		☐	☐	Active center/Chimera (1831)	SEQ ID NO: 65
Peptide_095	AAHCVEKPLNNPWHWTAFAG	☐				Active center/Chimera (1831)	SEQ ID NO: 66
Peptide_096	HCVEKPLNNPWHWTAFAGIL					Active center/Chimera (1831)	SEQ ID NO: 67
Peptide_097	VEKPLNNPWHWTAFAGILRQ					Chimera (1831)	SEQ ID NO: 68
Peptide_098	KPLNNPWHWTAFAGILRQSF		☐	☐	☐	Chimera (1831)	SEQ ID NO: 69

SEQ ID NO: 89 : **VTAAHCVEKPLNNPWHWT**SEQ ID NO: 90 : **KPLNNPWHWTAFAGILRQSF**

FIG. 5B

7: Self-cleavage site

Name	Sequence	752_1	2228_15	1864_10	2020_12	
Peptide_065	KAVVSLRCIACGVNLNSSRQ	☐				Self-cleavage site SEQ ID NO: 70
Peptide_066	VVSLRCIACGVNLNSSRQSR	☐	☐			Self-cleavage site SEQ ID NO: 71
Peptide_067	SLRCIACGVNLNSSRQSRIV	☐			☐	Self-cleavage site SEQ ID NO: 72
Peptide_068	RCIACGVNLNSSRQSRIVGG	☐			☐	Self-cleavage site SEQ ID NO: 73
Peptide_069	IACGVNLNSSRQSRIVGGES	☐				Self-cleavage site SEQ ID NO: 74

SEQ ID NO: 91 : **IACGVNLNSSRQSR**

8: Common to all TMPRSS2

Name	Sequence	752_1	2228_15	1864_10	2020_12	
Peptide_036	WNENYGRAACRDMGYKNNFY	☐			☐	SEQ ID NO: 75

SEQ ID NO: 92 : **WNENYGRAACRDMGYKNNFY**

9

Name	Sequence	752_1	2228_15	1864_10	2020_12	
Peptide_023	RLYG ENFI LQVYSSQRKSWH	☐	☐		☐	SEQ ID NO: 76
Peptide_024	YCP ENFI LQVYSSQRKSWHPV	☐			☐	SEQ ID NO: 77
Peptide_025	<u>PNFI</u> LQVYSSQRKSWHPVCQ	☐			☐	SEQ ID NO: 78
Peptide_026	<u>FNFI</u> LQVYSSQRKSWHPVCQDD	☐			☐	SEQ ID NO: 79
Peptide_027	LQVYSSQRKSWHPVCQDDWN	☐				SEQ ID NO: 80
Peptide_028	VYSSQRKSWHPVCQDDW NEN	☐				SEQ ID NO: 81
Peptide_029	SSQRKSWHPVCQDDW NENY C	☐				SEQ ID NO: 82
Peptide_030	QKSWHPVCQDDW NENY GRA	☐				SEQ ID NO: 83
Peptide_031	KSWHPVCQDDW NENY GRAAC	☐				SEQ ID NO: 84

SEQ ID NO: 93 : **PNFI**LQVYSSQRKSWHPVCQDD

FIG. 6




FIG. 7A




FIG. 7C




FIG. 7D

Dye antibody

ALL

316 to 321 a.d.

TMPRSS2- wild type

ROSE

TMRSS2 chimera
human/hamster
Hamster

SQLMS

Alanine substitution
F321A

POSTER




FIG. 8




FIG. 9




FIG. 10A

CLUSTAL O(1.2.4) multiple sequence alignment

```
Golden-hamster  MALNSGSPPGIGPYENHGFQSEHIYPPRPVAPDVYNFYPPQNYPPVPQYFPRVTTQA  60
mouse          MALNSGSPPGIGPCYENHGFQSEHICPPRPVAPNGYNLYPAQYYPSPVPQYAPRITTQA  60
human          MALNSGSPPAIGPYENHGFQSEHYPYPAQPTVVPTVYEVHQAQYYPSPVPQYAPRVLTA  60
green-monkey   MALNSGSPPGVGPFYENHGFQSEHYPYPAQPTVAFNVYEVHQAQYYPSPVPQYIPSVLTHA  60
crab-eating-macaque MALNSGSPPGVGPFYENHGFQSEHYPYPAQPTVAFNVYEVHQAQYYPSPVPQYTPRVLTHA  60
Rhesus-macaque MALNSGSPPGVGPFYENHGFQSEHYPYPAQPTVAFNVYEVHQAQYYPSPVPQYTPRVLTHA  60
cat            MALNSGSPPGVGPFYENHGFQSEGLYPPRPVAVVPGAYSVYPAPYPPAVPQYAPRVLTHA  60
Ferret         MALNAGSPPGVGPFYENHGFQSESLYPAPPATVPSVYVAYPAAYPPAVPQYTPRVLTA  60
               ****:* **::** *****: * * * * . * * * : * ** ***** * : * : *

Golden-hamster  STTVTHTQPHSS--GKLCSTSTKTKKSLCFALSLGIVLVGAAVAALLWKFL--PGCSTSEM  118
mouse          STSVINTHPKSS--GALCTS--KSKKSLCLALALGTVLTGAAVAALLWRFDSCNSTSEM  117
human          SNPVVCTQPKSPSGTVCTS--KTKKALCITLTTLGTFVLGAALAAGLLWKFMGSKCSNGGI  118
green-monkey   SNPAVRTQPKSPSGTVCTS--KTKKALCVTMTLGAVLVGAALAAGLLWKFMGSKCSDSGI  118
crab-eating-macaque SNPAVCRQPKSPSGTVCTS--KTKKALCVTMTLGAVLVGAALAAGLLWKFMGSKCSDSGI  118
Rhesus-macaque SNPAVCRQPKSPSGTVCTS--KTKKALCVTMTLGAVLVGAALAAGLLWKFMGSKCSDSGI  118
cat            STPSHTQPKSPSGTVCTS--KAKKVLCTITTLGAVLAGAVLAALLWKFMENKCLVSGI  118
Ferret         STPAVRTQPKSPSGTACTA--KAKKALCITISLGAVLAGAAVTAVLLWKFMENKCSVSGI  118
               * . : * * * * : * * * * * : : * * . * . : * * * * * * * * :

Golden-hamster  ECHSSGTCISSSLWCDGTSCHPCNGEDENRCVRLYGPSFTLQVYSSQRKAWYPVCQDDWND  178
mouse          ECGSSGTCISSSLWCDGVAHCPNGEDENRCVRLYGQSFILQVYSSQRKAWYPVCQDDWSE  177
human          ECDSSGTCINPSNWC DGVSCHPCNGEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWNE  178
green-monkey   ECDSSGTCISSSNWCDGVSCHPCNGEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWNE  178
crab-eating-macaque ECDSSGTCISSSNWCDGVSCHPCNGEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWNE  178
Rhesus-macaque ECDSSGTCISSSNWCDGVSCHPCNGEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWNE  178
cat            ECGSSGTCVSPSHWCDGVLHCPGSEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWSE  178
Ferret         ECGSSGTCISPSHWCDGVLHCPGSEDENRCVRLYGPNFILQVYSSQRKSWHPVCQDDWSD  178
               * * * * * : * * * * * * * * * * * * * * * * * * * * * :

Golden-hamster  SYGRAACKDMGYKNFFYTTQGIPOSSGATSFMKLNISAGNIDLYKKLYHSDSCSSRMVVS  238
mouse          SYGRAACKDMGYKNFFYSSQGIPOSSGATSFMKLNVSSGNVDLYKKLYHSDSCSSRMVVS  237
human          NYGRAACRDMGYKNFFYSSQGIPOSSGATSFMKLNTSAGNVDIYKKLYHSDACSSKAVVS  238
green-monkey   NYARAACRDMGYKNFFYSSQGIADNSGATSFMKLNTSAGNVDIYKKLYHSDACSSKAVVS  238
crab-eating-macaque NYARAACRDMGYKNSFYSSQGIPOSSGATSFMKLNTSAGNVDIYKKLYHSDACSSKAVVS  238
Rhesus-macaque NYARAACRDMGYKNSFYSSQGIPOSSGATSFMKLNTSAGNVDIYKKLYHSDACSSKAVVS  238
cat            GYGRAACQDMGYRNSFYSSHGVDSSGATSFMRVNTSANHMDLYKKLYHSDVCSSETVVS  238
Ferret         SYGRAACQDMGYRNSFYSSQGIPOSSGATSFMKLNTSAGNTDLYKKLYHSDICASKTVVS  238
               . * , * * * * : * * * * * : * * : * * : * * * * * * * * :

Golden-hamster  LRCIQCGVRS--ATRQSRIVGGSSNAPGDWPFQVSLHVQGVHVCSSIIITPEWIVTAAHCV  297
mouse          LRCIECGVRS--VKRQSRIVGGLNASPGDWPWFQVSLHVQGVHVCSSIIITPEWIVTAAHCV  296
human          LRCIACGVNLSNRQSRIVGGESALPGAWPFQVSLHVQNVHVCSSIIITPEWIVTAAHCV  298
green-monkey   LRCIACGVRSNLSNRQSRIVGGQNALPGAWPFQVSLHVQNIHVCSSIIITPEWIVTAAHCV  298
crab-eating-macaque LRCIACGVRSNLSNRQSRIVGGQNALPGAWPFQVSLHVQNIHVCSSIIITPEWIVTAAHCV  298
Rhesus-macaque LRCIACGVRSNLSNRQSRIVGGQNALPGAWPFQVSLHVQNIHVCSSIIITPEWIVTAAHCV  298
cat            LRCIECGVTAKMRQSRIVGGSSAAPGDWPFQVSLHVQGVHVCSSIIISPEWIVTAAHCV  298
Ferret         LRCIECGVAGKTHNRQSRIVGGSSASPGDWPWFQVSLHVQGVHVCSSIIITPEWIVTAAHCV  298
               * * * * * * * * * * * * * * * * * * * * * * * * * * *

Golden-hamster  SEQ ID No: 1
mouse          SEQ ID No: 2
human          SEQ ID No: 3
green-monkey   SEQ ID No: 4
crab-eating-macaque SEQ ID No: 5
Rhesus-macaque SEQ ID No: 6
cat            SEQ ID No: 7
Ferret         SEQ ID No: 8
```

FIG. 10B

Golden-hamster	EEPLNSPRYWTAFAGILSQSLMFYGSRHQVEKVISHPNYNSETKNNNDIALMKLQTPLTFN	357
mouse	EEPLSSPRIWTAFAGILRQSLMFYGSRHQVEKVISHPNYDSKTKNNNDIALMKLQTPLAFN	356
human	EKPLNNPWHWTAFAGILRQSFMFYAGYQVEKVISHPNYDSKTKNNNDIALMKLQKPLTFN	358
green-monkey	EKPLNSPWQWTAFAGILTQSSMFYVKGHRVEKVISHPNYDSKTKNNNDIALMKLHTPLTFN	358
crab-eating-macaque	EKPLNSPWQWTAFVGTTLRQSSMFYKGRHVEKVISHPNYDSKTKNNNDIALMKLHTPLTFN	358
Rhesus-macaque	EKPLNSPWQWTAFVGTTLRQSSMFYKGRHVEKVISHPNYDSKTKNNNDIALMKLHTPLTFN	358
cat	EEPLNNPRHWTAFVGTTLRQSFMFYGHGYRVGKVISHPNYDSKTKNNNDIALMKLQTPLTFD	358
Ferret	EEPLNNPRIYWTVFAGVLRQSFMFYGHGYRVGKVISHPSYDESKTKNNNDIALMKLQTPLTFS	358
	*:***_* **_* * ** ** * :* *****;*:*****;***;*,	
Golden-hamster	DLVKPVCLPNPGMMLDPAQECWISGWGSTYKGTSDMLNAAMVPLIERSKCNKXIYNN	417
mouse	DLVKPVCLPNPGMMLDLQECWISGWGATYKGTSDVLNAAMVPLIEPSKCNKXIYNN	416
human	DLVKPVCLPNPGMMLQPEQLCWISGWGATEKGTSEVLNAAKVLLIETQRCSRYVYDN	418
green-monkey	ELVKPVCLPNPGMMLPEPQHCWISGWGATQEKGTSDMLNAAMVPLIEPRRCNSKRVYDG	418
crab-eating-macaque	EVVKPVCLPNPGMMLPEPQHCWISGWGATQEKGTSDVLNAAMVPLIEPRRCNKKXIYDG	418
Rhesus-macaque	EVVKPVCLPNPGMMLPEPQHCWISGWGATQEKGTSDVLNAAMVPLIEPRRCNKKXIYDG	418
cat	DKVKPVCLPNPGLMLEPQPCWISGWGATHEKGTSDVLNAAMVPLIEPWRCSKXIYNN	418
Ferret	DKVKPVCLPNPGMMLPEPQSCWISGWGATHEKGTSDVLNAAMVPLIEPWRCSKXIYNS	418
	: *****;*: * *****;* *****; ***, * ** * **:; :*:;	
Golden-hamster	LITPAMVCAGFLQGTVDSCQGDSCGGPLVTLKNDIWWLIGDTSWGSACAALRPGVYGNVT	477
mouse	LITPAMICAGFLQGSVDSCQGDSCGGPLVTLKNGIWWLIGDTSWGSACAALRPGVYGNVT	476
human	LITPAMICAGFLQGNVDSCQGDSCGGPLVTSNNIWWLIGDTSWGSACAALRPGVYGNVT	478
green-monkey	LVTPAMICAGFLQGTVDSCQGDSCGGPLVTLKNDVWWLIGDTSWGSACAALRPGVYGNVT	478
crab-eating-macaque	LITPAMICAGFLRGTVDSCQGDSCGGPLVTLKNDVWWLIGDTSWGSACAALRPGVYGNVT	478
Rhesus-macaque	LITPAMICAGFLQGTVDSCQGDSCGGPLVTLKNDVWWLIGDTSWGSACAALRPGVYGNVT	478
cat	LVTPAMICAGYLQGTIDSCQGDSCGGPLVTMKSIIWWLIGDTSWGSACAALRPGVYGNVT	478
Ferret	LVTPAMICAGYLGGTIDSCQGDSCGGPLVTLKRIWWLIGDTSWGSACAALRPGVYGNVT	478
	*:*****;***;*: ***** ***** * _ :*****;*: *****	
Golden-hamster	VFTDWIYQQMMANS	491
mouse	VFTDWIYQQMRANS	490
human	VFTDWIYRQMRADG	492
green-monkey	VFTDWIYRQMRADD	492
crab-eating-macaque	VFTDWIYRQMRADD	492
Rhesus-macaque	VFTDWIYRQMRADD	492
cat	VFTDWIYRQMRANG	492
Ferret	VFTDWIYRQMRANS	492
	*****;** *	

FIG. 10C




FIG. 10D




FIG. 11A




FIG. 11B




FIG. 12




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 13D




FIG. 14




FIG. 15





FIG. 16

Target cells: Daudi human TMPRSS2



Target cells: Daudi Cynomolgus Monkey TMPRSS2




FIG. 17




FIG. 18A

Cynomolgus monkey in vivo experiment 2





FIG. 18B

Body temperature change at night



Body weight change




FIG. 19A

Amount of viruses in airway and bronchus swabs
(titration)





FIG. 19B

Amount of viruses in bronchus swab
(RT-PCR)



Student T-
test, $p < 0.05$

FIG. 20




FIG. 21





FIG. 22



ANTIBODY USED TO TREAT CORONAVIRUS INFECTION

TECHNICAL FIELD

[0001] The present invention relates to an antibody for use in treating coronavirus infection.

BACKGROUND ART

[0002] Nowadays, the coronavirus pandemic has been greatly affecting the global economy. Because of mRNA vaccine against coronavirus being developed and quickly spread out across the world, the threats of the coronavirus infection seem to have come to an end for a while. However, variants of the coronavirus have emerged and rapidly replaced the original strain and variants of the past. For example, the α variant was confirmed in the UK in September 2020. The β variant was confirmed in South America in May 2020. The γ variant was confirmed in Brazil in November 2020. The δ variant was confirmed in India in October 2020. Also, the omicron variant was confirmed in multiple countries in November 2021. Moreover, the λ variant was confirmed in Peru in December 2020, and the μ variant was confirmed in Colombia in January 2021. Besides that, the World Health Organization (WHO) has designated from time to time a large number of variants as variants of concern (VOC) and variants of interest (VOI), and the threats of coronavirus variants seem to continue to happen. As such, a large number of variants emerged within the past short period of 1 to 2 years, which causes threats across the world every time a new variant emerges.

[0003] Prophylactic and therapeutic drugs for coronavirus infection have been developed worldwide. In particular, mRNA vaccines that encode the spike(S) proteins of coronaviruses have been administered by intramuscular injection all over the world, and their extremely high effectiveness has been confirmed. The mRNA vaccines contain, as an active ingredient, lipid nanoparticles (LNPs) that encapsulate mRNA containing pseudo-uridine (Patent Literatures 1 and 2).

[0004] However, the emergence of a large number of variants raises concerns about the effectiveness of the vaccines, and therefore, every time a new variant emerges, the effectiveness of the vaccines needs to be confirmed, and if not effective, vaccines must be redesigned and developed again.

CITATION LIST

Patent Literature

- [0005]** Patent Literature 1: WO2007/024708A
[0006] Patent Literature 2: WO2009/127060A

SUMMARY OF INVENTION

[0007] The present invention provides an antibody against coronaviruses, the antibody being expected to be effective regardless of virus mutations.

[0008] Specifically, the present inventors have found that antibodies that target human protein TMPRSS2 required to activate the S protein used by coronaviruses for infection, can widely prevent coronavirus infection regardless of variant.

[0009] According to the present invention, the following inventions are provided.

[0010] [1] An antibody or an antigen-binding fragment thereof that binds to an extracellular domain of TMPRSS2, capable of inhibiting coronavirus infection of cells.

[0011] [2] The antibody or an antigen-binding fragment thereof according to [1] above, capable of binding to (i) one or more extracellular domains of TMPRSS2 selected from the group consisting of mice, green monkeys, cats, hamsters, and macaque monkeys, and (ii) an extracellular domain of human TMPRSS2.

[0012] [3] The antibody or an antigen-binding fragment thereof according to [1] or [2] above, wherein a binding affinity (KD) to human TMPRSS2 is 10^{-8} M or less.

[0013] [4] The antibody or an antigen-binding fragment thereof according to any of [1] to [3] above, wherein IC₅₀ referring to inhibition of the coronavirus infection of human cells is 10 μ g/mL or less.

[0014] [5] The antibody or an antigen-binding fragment thereof according to any of [1] to [4] above, wherein the antibody binds to a peptide having the amino acid sequence consisting of the amino acid sequence selected from the group consisting of the amino acid sequence set forth in SEQ ID NO: 75 (WNE-NYGRAACRDMGYKNNFY), and the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT).

[0015] [6] The antibody or an antigen-binding fragment thereof according to any of [1] to [4] above, wherein the antibody binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and wherein:

[0016] (α) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 75 (WNE-NYGRAACRDMGYKNNFY) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 94: WSEGYGRAACQDMGYRNSFY);

[0017] (β) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 95: VTAAHCVEEPLNNPRHWT); or

[0018] (γ) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 96 (RQSFMF) of the human TMPRSS2 is substituted for a corresponding sequence of a hamster (i.e., the amino acid sequence set forth in SEQ ID NO: 97: SQSLMF).

[0019] [7] The antibody or an antigen-binding fragment thereof according to any of [1] to [4] above, wherein the antibody binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and wherein:

[0020] (A) the antibody does not significantly bind to a human TMPRSS2 variant having a point mutation at N179A; or

- [0021] (B) the antibody does not significantly bind to a human TMPRSS2 variant having a point mutation at W306A.
- [0022] [8] The antibody or an antigen-binding fragment thereof according to any of [1] to [4] above, wherein the antibody is: (1) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 11,
- [0023] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 12, and
- [0024] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 13,
- [0025] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 14,
- [0026] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 15, and
- [0027] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 16.
- [0028] (2) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 19,
- [0029] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 20, and
- [0030] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 21,
- [0031] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 22,
- [0032] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 23, and
- [0033] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 24.
- [0034] (3) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 27,
- [0035] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 28, and
- [0036] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 29,
- [0037] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 30,
- [0038] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 31, and
- [0039] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 32.
- [0040] (4) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 35,
- [0041] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 36, and
- [0042] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 37,
- [0043] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 38,
- [0044] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 39, and
- [0045] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 40.
- [0046] (5) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 100,
- [0047] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 101, and
- [0048] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 102,
- [0049] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 103,
- [0050] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 104, and
- [0051] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 105.
- [0052] (6) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 108,
- [0053] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 109, and
- [0054] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 110,
- [0055] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 111,
- [0056] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 112, and
- [0057] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 113.
- [0058] (7) an antibody or an antigen-binding fragment thereof, competing with the antibody according to any of (1) to (6) above for binding with TMPRSS2; or
- [0059] (8) an antibody or an antigen-binding fragment thereof, including the amino acid sequence having one or several amino acid mutations in the CDR sequence of the antibody according to any of (1) to (6) above; or
- [0060] (9) a human chimeric antibody or a humanized antibody of the antibody according to any of (1) to (8) above.
- [0061] [9] A composition including the antibody according to any of [1] to [8] above or an antigen-binding fragment thereof.
- [0062] [10] The composition according to [9] above, for use in preventing a subject in need thereof from being infected with coronavirus.
- [0063] [11] A pharmaceutical composition according to [9] above, for use in administering to a subject with coronavirus.
- [0064] [12] The composition according to [9] above, for use in treating coronavirus infection in a subject infected with a beta coronavirus.
- [0065] [21] The antibody according to any of [1] to [8] above, wherein the antibody is a humanized antibody, or an antigen-binding fragment thereof.
- [0066] [22] The antibody according to any of [1] to [8] above, wherein the antibody is a human chimeric monoclonal antibody, or an antigen-binding fragment thereof.
- [0067] [23] A composition, including the antibody according to [21] or [22] above or an antigen-binding fragment thereof.

- [0068] [24] The composition according to [23] above, for use in preventing a subject in need thereof from being infected with coronavirus.
- [0069] [25] A pharmaceutical composition according to [23] above, for use in administering to a subject with coronavirus.
- [0070] [26] The composition according to [23] above, for use in treating coronavirus infection in a subject infected with a beta coronavirus.
- [0071] [101] The antibody or composition according to any of the above, wherein the coronavirus is a beta coronavirus.
- [0072] [102] The antibody or composition according to any of the above, wherein the coronavirus is SARS-COV, MERS, or SARS-COV-2.
- [0073] [103] The antibody or composition according to any of the above, wherein the coronavirus is one or more selected from the alpha variant, the beta variant, the gamma variant, the delta variant, and the omicron variant of SARS-COV-2.
- [0074] [111] The antibody or composition according to any of the above, wherein TMPRSS2 is human TMPRSS2, and the cell is a human cell.
- [0075] [112] The antibody or composition according to [101] above, wherein TMPRSS2 is human TMPRSS2, and the cell is a human cell.
- [0076] [113] The antibody or composition according to [102] above, wherein TMPRSS2 is human TMPRSS2, and the cell is a human cell.
- [0077] [114] The antibody or composition according to [103] above, wherein TMPRSS2 is human TMPRSS2, and the cell is a human cell.
- [0078] [121] The antibody or an antigen-binding fragment thereof according to [1] above, wherein a binding affinity (KD) to the human TMPRSS2 of the antibody or the antigen-binding fragment is 10^{-8} M or less, and IC_{90} referring to inhibition of the coronavirus infection of human cells is 10 μ g/mL or less.
- [0079] [122] The antibody or an antigen-binding fragment thereof according to [1] above, wherein the binding affinity (KD) to the human TMPRSS2 of the antibody or the antigen-binding fragment is 10^{-8} M or less, and the antibody or the antigen-binding fragment binds to the amino acid sequence set forth in SEQ ID NO: 85 (GVYGNVMVFTDWIY).
- [0080] [123] The antibody or an antigen-binding fragment thereof according to [1] above, wherein IC_{90} referring to inhibition of the coronavirus infection of human cells of the antibody or the antigen-binding fragment is 10 μ g/mL or less, and the antibody or the antigen-binding fragment binds to the amino acid sequence set forth in SEQ ID NO: 85 (GVYGNVMVFTDWIY).
- [0081] [124] The antibody or an antigen-binding fragment thereof according to [1] above, wherein the binding affinity (KD) to the human TMPRSS2 of the antibody or the antigen-binding fragment is 10^{-8} M or less, IC_{90} referring to inhibition of the coronavirus infection of human cells is 10 μ g/mL or less, and the antibody or the antigen-binding fragment binds to the amino acid sequence set forth in SEQ ID NO: 85 (GVYGNVMVFTDWIY).
- [0082] [131] The antibody or an antigen-binding fragment thereof according to [121] above, competing with the antibody according to any of (2) to (4) above for binding to TMPRSS2.
- [0083] [132] The antibody or an antigen-binding fragment thereof according to [122] above, competing with the antibody according to any of (2) to (4) above for binding to TMPRSS2.
- [0084] [133] The antibody or an antigen-binding fragment thereof according to [123] above, competing with the antibody according to any of (2) to (4) above for binding to TMPRSS2.
- [0085] [134] The antibody or an antigen-binding fragment thereof according to [124] above, competing with the antibody according to any of (2) to (4) above for binding to TMPRSS2.
- [0086] [135] The antibody or an antigen-binding fragment thereof according to [121] above, competing with the antibody according to (1) above for binding to TMPRSS2.
- [0087] [136] The antibody or an antigen-binding fragment thereof according to [122] above, competing with the antibody according to (1) above for binding to TMPRSS2.
- [0088] [137] The antibody or an antigen-binding fragment thereof according to [124] above, competing with the antibody according to (1) above for binding to TMPRSS2.
- [0089] [138] The antibody or an antigen-binding fragment thereof according to [124] above, competing with the antibody according to (1) above for binding to TMPRSS2.
- [0090] [141] An antibody or an antigen-binding fragment thereof that binds to an extracellular domain of TMPRSS2, capable of inhibiting coronavirus infection of cells and inhibiting an action of TMPRSS2 on an S protein.
- [0091] [142] An antibody or an antigen-binding fragment thereof that binds to an extracellular domain of TMPRSS2, capable of inhibiting coronavirus infection of cells and inhibiting an interaction of TMPRSS2 with an S protein.
- [0092] [151] A composition including the antibody according to any of the above or an antigen-binding fragment thereof.
- [0093] [152] The composition according to [151] above, for use in preventing cells from being infected with coronavirus.
- [0094] [153] The composition according to [151] above, for use in preventing the coronavirus infection.
- [0095] [154] The composition according to [151] above, for use in treating a subject infected with coronavirus.
- [0096] [161] A method for treating a subject, including administering to the subject an effective amount of the antibody according to any of the above or an antigen-binding fragment thereof.
- [0097] [162] The method according to [161] above, wherein a subject carries coronavirus.
- [0098] [163] The method according to [161] or [162] above, wherein further coronavirus infection of cells in the body of a subject is suppressed.
- [0099] [164] A method for treating a subject infected with coronavirus, including administering to the sub-

ject an effective amount of the antibody according to any of the above or an antigen-binding fragment thereof.

[0100] [171] The antibody according to the above or an antigen-binding fragment thereof, or a pharmaceutical composition including the antibody or an antigen-binding fragment thereof, for use in the method according to any of [161] to [164] above.

[0101] [172] Use of the antibody according to the above or an antigen-binding fragment thereof, in manufacture of medicament for use in the method according to any of [161] to [164] above.

BRIEF DESCRIPTION OF DRAWINGS

[0102] FIG. 1 shows that TMPRSS2 which is essential for beta coronavirus infection of cells is expressed on a membrane surface of an infected cell (host cell), and that the antibody of the present invention binds to TMPRSS2, which is expressed on the membrane surface of the infected cell. The antibody of the present invention binds to an antigen that is expressed in a host cell, and it is therefore theoretically unaffected by viral mutation.

[0103] FIG. 2 shows the binding properties to human TMPRSS2 of the monoclonal antibodies that bind to human TMPRSS2 newly obtained in Examples. The human TMPRSS2 was expressed in Daudi cell lines, and the binding between the antibodies and the cells was detected by flow cytometry. Data on the antibodies whose bindings were particularly good were enclosed in square frames.

[0104] FIG. 3A shows results from a test system that tests whether or not the monoclonal antibodies that bind to human TMPRSS2 newly obtained in Examples compete with a 752_1 antibody for binding to the human TMPRSS2.

[0105] FIG. 3B shows results from the test system which tests whether or not the monoclonal antibodies that bind to human TMPRSS2 newly obtained in Examples compete with a 2228_15 antibody for binding to the human TMPRSS2.

[0106] FIG. 3C shows results from the test system that tests whether or not the monoclonal antibodies that bind to human TMPRSS2 newly obtained in Examples compete with an 1831_15 antibody for binding to the human TMPRSS2.

[0107] FIG. 3D shows results from the test system that tests whether or not the monoclonal antibodies that bind to human TMPRSS2 newly obtained in Examples compete with an 1864_10 antibody for binding to the human TMPRSS2.

[0108] FIG. 4 is data showing the binding properties of various monoclonal antibodies to 20 amino acid-long peptides that were designed by shifting 2 amino acids each in a whole area of the human TMPRSS2.

[0109] FIG. 5A shows the binding properties of each monoclonal antibody to each peptide.

[0110] FIG. 5B shows the binding properties of each monoclonal antibody to each peptide.

[0111] FIG. 6 shows epitope regions of the antibody on human TMPRSS2 predicted from the results of FIGS. 5A and 6B.

[0112] FIG. 7A is data showing the binding properties between each monoclonal antibody and the human TMPRSS2 variants.

[0113] FIG. 7B is data showing the binding properties between each monoclonal antibody and the human TMPRSS2 variants.

[0114] FIG. 7C is data showing the binding properties between each monoclonal antibody and the human TMPRSS2 variants.

[0115] FIG. 7D is data showing the binding properties between each monoclonal antibody and the human TMPRSS2 variants.

[0116] FIG. 8 shows three-dimensional structure models of the epitope region obtained by observing complexes of a Fab fragment of the 752_1 antibody, a Fab fragment of the 2228 antibody, and the human TMPRSS2 with a cryo-electron microscope. Graphics in string-shape shows human TMPRSS2, and a surface model drawn in the back shows an antibody.

[0117] FIG. 9 shows the binding properties of the obtained various antibodies to other TMPRSS family members.

[0118] FIG. 10A shows amino acid sequence alignment results of TMPRSS2 from each animal species.

[0119] FIG. 10B shows amino acid sequence alignment results of TMPRSS2 from each animal species.

[0120] FIG. 10C shows the binding properties of each monoclonal antibody to mouse, cynomolgus monkey, or human TMPRSS2.

[0121] FIG. 10D shows the binding properties of each monoclonal antibody to mouse or human TMPRSS2.

[0122] FIG. 11A shows the binding properties of human chimeric antibodies obtained by replacing the Fc regions of the obtained monoclonal antibodies with the Fc region of human IgG4, to human TMPRSS2.

[0123] FIG. 11B shows the binding properties of human chimeric antibodies derived from the 752_1 antibody obtained by replacing the Fc region thereof with the Fc region of human IgG4, to human and cynomolgus monkey TMPRSS2.

[0124] FIG. 12 shows inhibitory effects on viral infection by the obtained various monoclonal antibodies.

[0125] FIG. 13A shows infection inhibiting effects on various virus variants by the obtained various monoclonal antibodies.

[0126] FIG. 13B shows infection inhibiting effects on various virus variants by the obtained various monoclonal antibodies.

[0127] FIG. 13C shows infection inhibiting effects on various virus variants by the obtained various monoclonal antibodies.

[0128] FIG. 13D shows infection inhibiting effects on various virus variants by the obtained various monoclonal antibodies.

[0129] FIG. 14 shows inhibitory effects on cell fusion (fusion between TMPRSS2/ACE-2 expressing 293T cells and spike protein expressing 293T cells) by the obtained various monoclonal antibodies. The fact that the cell fusion was inhibited suggests that virus entry to cells is inhibited.

[0130] FIG. 15 shows measurement results of antibody-dependent cell cytotoxicity (ADCC) of Antibody-Dependent Effect (ADE) substituted (L244A/E245A/P339A) recombinant antibodies in the Fc regions of the obtained monoclonal antibodies.

[0131] FIG. 16 shows measurement results of antibody-dependent cell cytotoxicity (ADCC) of the obtained human chimeric monoclonal antibodies.

[0132] FIG. 17 shows results of SARS-COV-2 infection inhibition experiment on cells expressing human or macaque TMPRSS2.

[0133] FIG. 18A shows an experimental scheme for inhibitory effects of antibody administration on SARS-CoV-2 infection of cynomolgus monkeys.

[0134] FIG. 18B shows changes in body temperature and body weight of individuals after administering antibodies against SARS-COV-2 infection of cynomolgus monkeys.

[0135] FIG. 19A shows measurement results (by titration method) of the number of viruses in the airway and bronchus swabs of the cynomolgus monkeys.

[0136] FIG. 19B shows measurement results (by RT-PCR method) of the number of viruses in the bronchus swab of the cynomolgus monkeys.

[0137] FIG. 20 shows results of lung histopathologic diagnosis score of the cynomolgus monkeys by hematoxylin-eosin (HE) staining and immunohistochemical staining of SARS-COV-2 N protein.

[0138] FIG. 21 shows results of binding tests for the prepared humanized antibodies to TMPRSS2 expressing cells.

[0139] FIG. 22 shows results of an experiment of inhibiting infection of cells with pseudo-typed virus by the prepared humanized antibodies.

DESCRIPTION OF EMBODIMENTS

Definition

[0140] As used herein, the “subject” is a vertebrate, and for example, may be a mammal including a human, such as a mammal to be infected with coronavirus (e.g., SARS-COV-2) (cats, ferrets, bats, and pangolins). The subject may be a subject infected with coronavirus (e.g., SARS-COV-2), may be an asymptomatic carrier infected with coronavirus (e.g., SARS-COV-2), and may be a subject infected with coronavirus (e.g., SARS-COV-2) and developed COVID-19. The subject may be a subject who has a possibility (is at a risk) of having been infected with coronavirus (e.g., SARS-COV-2), or may be a subject who has a possibility (is at a risk) of being infected with coronavirus (e.g., SARS-COV-2). The subject may be a young child (e.g., infant (1- to 6-year-old), a schoolchild (6- to 12-year-old), an adolescent (12-year-old or above), or an adult (20-year-old or above). The adult may be an adult of 30-year-old or above, 40-year-old or above, 50-year-old or above, 60-year-old or above, or 70-year-old or above.

[0141] As used herein, the “coronavirus” is a virus belonging to the subfamily Ortho coronavirinae in the family Coronaviridae and the order Nidovirales, and a single-stranded plus-strand RNA virus. Coronaviruses have spike protein projections (S protein) on the surface of the viruses and was named as coronavirus because the appearance resembles the solar corona. In humans, it causes respiratory infection including common cold. The viruses belonging to the subfamily Ortho coronavirinae are classified into Alphacoronavirus, Beta coronavirus, Gamma coronavirus and Delta coronavirus. SARS-related coronaviruses are classified in Beta coronavirus. The SARS-related coronaviruses include SARS-coronavirus (SARS-COV) and SARS-coronavirus 2 (SARS-COV-2). SARS-CoV-2 has been causing novel coronavirus infection (COVID-19) pandemic since the end of 2019. The SARS-related coronaviruses infect host cells by binding to ACE2 receptors of the host cells through

S proteins. The SARS-related coronaviruses have a common infection mechanism in that they infect cells using the ACE2 receptors. The S proteins of SARS-COV-2 have furin cleavage sites that increase infectivity and pathogenicity therein (Andersen et al., Nature Medicine, 26, 450-452, 2020). Coronaviruses are known to cause respiratory infection such as common cold in humans, and SARS coronavirus (SARS-COV), MERS coronavirus (MERS-COV), and 2019 novel coronavirus (SARS-COV-2) have lethality. As other coronaviruses, lethal viruses are known such as mouse hepatitis virus (MHV) and feline infectious peritonitis virus (FIPV). Coronaviruses infect cells by binding to target cell surfaces through the binding of the spike proteins exposed on the envelope surfaces and angiotensin-converting enzymes 2 (ACE2) in cell surface molecules, and then by being taken into the cells by endocytosis. Examples of coronaviruses include coronaviruses belonging to the subfamily Coronavirinae, i.e., Alphacoronavirus (e.g., canine coronavirus, alphacoronavirus 1, human coronavirus 229E, human coronavirus NL63, and porcine epidemic diarrhea virus), Beta coronavirus (e.g., subgenus Embecovirus, subgenus Sarbecovirus, subgenus Merbecovirus, and subgenus Nobecovirus, such as human enteric coronavirus 4408, human coronavirus OC43, mouse coronavirus, human coronavirus HKU1, SARS-related coronavirus (e.g., SARS coronavirus (SARS-COV), 2019 novel coronavirus (SARS-COV-2), and MERS coronavirus, and equine coronavirus), Gamma coronavirus (e.g., avian coronavirus, and beluga whale coronavirus SW1), and Delta coronavirus (e.g., bulbul coronavirus HKU11, munia coronavirus HKU13, and thrush coronavirus HKU12). Although vaccines against coronavirus have been developed, these are vaccines against antigens that coronavirus has, such as S proteins, and therefore, theoretically it cannot be said that these vaccines maintain their effectiveness against mutation of coronavirus.

[0142] As used herein, “SARS-COV-2” is the coronavirus that caused the pandemic that occurred in 2020. On Jan. 7, 2020, The World Health Organization (WHO) provisionally named this virus 2019-ncov. Also, on February 11 of the same year, The International Committee on Taxonomy of Viruses (ICTV) officially named this virus SARS-COV-2. Coronaviruses can cause illnesses ranging from common cold to serious respiratory diseases such as severe acute respiratory syndrome (SARS) or Middle East respiratory syndrome (MERS). The WHO named the disease caused by this novel coronavirus COVID-19. The International Committee on Taxonomy of Viruses determined that SARS-CoV-2 belongs to the genus Beta coronavirus and is the same species as SARS-COV (or its sister lineage). The complete genomic sequence of SARS-COV-2 is registered to the National Center for Biotechnology Information (NCBI) under GenBank Accession Number: MN908947.3. Virus particles (virions) have particle sizes of about 50 to 200 nm and, similar to common coronaviruses, contain spike proteins, nucleocapsid protein, membrane proteins and envelope proteins, and viral genome RNA. Nucleocapsid proteins form a complex with RNA, around which spike proteins, membrane proteins and envelope proteins that are bound with lipid surround to form a virion envelope. The spike protein located on the outermost surface of the envelope is thought to facilitate the infection of cells by binding to an ACE2 receptor on the cell surface. There are people who do not show symptoms of the disease despite infection with SARS-COV-2, and they are called asymptomatic car-

riers. It is pointed out that asymptomatic carriers have a possibility to infect others with the virus they carry. It is pointed out that because of infection with SARS-CoV-2, senses of smell and/or taste are decreased or lost. SARS-CoV-2 may cause severe acute respiratory syndrome. As the main symptoms of severe acute respiratory syndrome, fever of around 40° C., cough and shortness of breath have been reported. Significant complication is pneumonia. The presence or absence of SARS-CoV-2 infection is mainly determined by PCR testing. The PCR test is to evaluate whether or not the SARS-CoV-2 gene is present in the body, depending on whether or not a band is amplified specific to SARS-CoV-2. Examples of the treatment for SARS-CoV-2 include antiviral drugs against SARS-CoV-2 (e.g., remdesivir), steroidal anti-inflammatory drugs (e.g., dexamethasone), and inhibitors of proinflammatory cytokines (e.g., IL-6 inhibitors such as anti-IL-6 antibody, TNF- α inhibitor such as etanercept).

[0143] As used herein, the “spike protein” can be a protein to be encoded at positions 21563 to 25384 of SARS-CoV-2 genome registered to the National Center for Biotechnology Information (NCBI) under GenBank Accession Number: MN908947.3, or a protein having the amino acid sequence set forth in SEQ ID NO: 1; and the above spike protein of SARS-CoV-2 has the amino acid sequence registered to NCBI under GenBank Accession Number: QHD43416.1. The spike proteins include S1 and S2; S1 is present at positions 13 to 541 of the above amino acid sequence, and S2 is present at positions 543 to 1208 of the above amino acid sequence. S1 further has an N-terminal domain (NTD) and a receptor binding domain (RBD); NTD is present at positions 13 to 304 of the above amino acid sequence, and RBD is present at positions 319 to 541. S1 and S2 are cleaved in the cell, produced as separate peptides, and form a complex during virus particle formation. The spike protein is also referred to as S protein. As the spike protein, any spike protein that natural virus (e.g., including virus variants) has (those having the amino acid sequence corresponding to the amino acid sequence registered to NCBI under GenBank Accession Number: QHD43416.1) can be used.

[0144] As used herein, the “TMPRSS2” is a cell membrane protein called transmembrane protease, serine 2 (type II transmembrane serine protease), and is expressed in the prostate and airway epithelium. TMPRSS2 is expressed as an inactive precursor, and is converted into an active type by self-cleavage at the QSR sequence. Further, the active TMPRSS2 has been confirmed to cleave and activate respiratory viruses such as influenza A virus, influenza B virus, Sendai virus, human parainfluenza 1 to 4, and human metapneumovirus. However, for these viruses, TMPRSS2 is involved in the formation of virus particles produced in the cell and the like but is not involved in the invasion of these viruses into cells. On the other hand, TMPRSS2 does not particularly act against coronaviruses in the cell, but it acts on the S protein of coronaviruses that are entering cells from outside of the cells and cleaves and activates the S protein to facilitate cell entry of coronaviruses. TMPRSS2 knockout mice exhibit a healthy phenotype similar to that of normal mice (Kim T S. et al., *Mol. Cell Bio.*, 26:965-975, 2006; and Sakai K. et al., *J. Virol.*, 88:5608-5616, 2014). Therefore, when inhibiting functions of TMPRSS2 though, there will be little or no adverse effect on the living body. Ferret TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 1 or the amino acid sequence corre-

sponding thereto. Hamster TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 2 or the amino acid sequence corresponding thereto. Mouse TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 3 or the amino acid sequence corresponding thereto. Feline TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 4 or the amino acid sequence corresponding thereto. Human TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 5 or the amino acid sequence corresponding thereto. Green monkey TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 6 or the amino acid sequence corresponding thereto. Cynomolgus monkey TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 7 or the amino acid sequence corresponding thereto. Rhesus macaque TMPRSS2 has, for example, the amino acid sequence set forth in SEQ ID NO: 8 or the amino acid sequence corresponding thereto. Human TMPRSS2 includes a cytoplasmic region on the N-terminal side, a transmembrane region for 85th to 105th amino acids, an extracellular domain for amino acids thereafter, an LDLRA region for 112th to 149th amino acids, an SRCR region for 150th to 242nd amino acids, and a serine protease region for 255th to 492nd amino acids. The serine protease region contains catalytic triad of H296, D345, and S441.

[0145] As used herein, the phrase “amino acid sequence corresponding to the amino acid sequence of SEQ ID NO: n” refers to a sequence corresponding to the amino acid sequence of SEQ ID NO: n {where n is a natural number} when aligning two amino acid sequences.

[0146] As used herein, the term “peptide” refers to an amino acid polymer. Polymers usually have no branches. The term “partial peptide” refers to a part of a particular peptide. Peptides and partial peptides can be produced from nucleic acids that encode the peptides. Peptides and partial peptides can also be chemically synthesized. Peptides and partial peptides are also isolated, concentrated or purified. Isolation means to separate peptides and partial peptides from at least other components, and purification means to separate peptides and partial peptides at least selectively. Concentration means that the concentration of peptides and partial peptides is increased.

[0147] As used herein, the term “antibody” refers to an immunoglobulin and is a protein having a structure wherein two heavy chains (H chains) and two light chains (L chains) both of which are stabilized by a pair of disulfide bonds, are associated with each other. The heavy chain consists of a heavy chain variable region VH, a heavy chain constant regions CH1, CH2 and CH3, and a hinge region located between CH1 and CH2; and the light chain consists of a light chain variable region VL, and a light chain constant region CL. Among these, a variable region fragment (Fv) consisting of VH and VL is a region directly involved in the antigen binding and adding diversity to the antibody. Also, the antigen-binding region consisting of VL, CL, VH and CH1 is referred to as Fab region, and a region consisting of the hinge region, CH2 and CH3 is referred to as Fc region.

[0148] Within the variable region, a region that directly contacts the antigen is particularly variable and is referred to complementarity-determining region (CDR). Portions other than CDR with relatively less mutations are referred to as framework regions (FR). There are 3 CDRs in each of the light and heavy chain variable regions, which are respec-

tively referred to, from the N-terminal side, heavy chain CDR1 to 3 and light chain CDR1 to 3.

[0149] The antibody may be a monoclonal antibody or polyclonal antibody. Also, the antibody may be any isotype among IgG, IgM, IgA, IgD, and IgE. The antibody may be produced by immunizing a non-human animal, such as mouse, rat, hamster, guinea pig, rabbit, or chicken. The antibody may be a recombinant antibody. The antibody may be a chimeric antibody, humanized antibody, fully humanized antibody, human antibody, or the like.

[0150] The chimeric antibody is an antibody wherein fragments of antibodies derived from different species are linked. The chimeric antibody may be a human chimeric antibody wherein at least the constant region is derived from a human antibody.

[0151] The term “humanized antibody” refers to an antibody wherein the corresponding position of a non-human antibody is substituted with the amino acid sequence (e.g., framework region) characteristic of a human-derived antibody, and examples thereof include those having the heavy chain CDR1 to 3 and the light chain CDR1 to 3 of the antibody produced by immunizing a mouse or rat, and wherein all other regions including 4 framework regions (FR) of each of the heavy chain and the light chain are derived from a human antibody. Such an antibody may be referred to CDR-grafted antibody.

[0152] The “human chimeric antibody” is, in a non-human-derived antibody, an antibody wherein the constant region of the non-human-derived antibody is substituted for the constant region of a human antibody. For the human chimeric antibody, from the viewpoint of enhancing ADCC activity, for example, the subtype of the human antibody used for the constant region may be IgG1.

[0153] As used herein, the “composition” is a mixture of one or more components. The composition can contain, for example, a partial peptide and an aqueous solvent (e.g., water). The composition can further contain a pharmaceutically acceptable additive (e.g., excipient and/or carrier). The composition used to treat a subject is referred to as pharmaceutical composition. The antibody is usually present in the form of a composition. The isolated antibody can be contained in the composition together with other pharmaceutically acceptable additives.

[0154] As used herein, the “treatment” includes prophylactic treatment and therapeutic treatment. The therapeutic treatment can be directed against infected viruses, whereas the prophylactic treatment can be conducted in order to prevent future infection, to delay the onset of coronavirus infection (e.g., COVID-19) due to infection, to reduce, delay deterioration of, or cease deterioration of the symptoms of the coronavirus infection (e.g., COVID-19) that has developed, or to prevent the exacerbation. The therapeutic treatment can be administered to symptomatic patients or asymptomatic carriers. The prophylactic treatment can be administered to non-infected individuals. The therapeutic treatment can be administered to infected individuals.

[0155] As used herein, the phrase “effective amount” refers to an amount of active ingredients that achieve expected pharmacological effects.

<Antibody of the Present Invention or Antigen-Binding Fragment Thereof>

[0156] According to the present invention, provided is an antibody that binds to an extracellular domain of TMPRSS2,

the antibody being capable of inhibiting coronavirus infection of cells, or an antigen-binding fragment thereof. In an embodiment, the antibody of the present invention can inhibit an interaction between TMPRSS2 and an S protein, resulting in inhibiting coronavirus infection of cells. In a preferred embodiment, TMPRSS2 can be human TMPRSS2. The inhibition of TMPRSS2 by the antibody of the present invention is considered not to affect the healthy phenotype of an individual and does not cause unacceptable side effects to the living body.

[0157] In an embodiment of the present invention, the antibody of the present invention can bind to (i) one or more extracellular domains of TMPRSS2 selected from the group consisting of mice, green monkeys, cats, hamsters, and macaque monkeys, as well as to (ii) human TMPRSS2. In a preferred embodiment, the antibody of the present invention can bind to mouse TMPRSS2 and human TMPRSS2. In a preferred embodiment, the antibody of the present invention can bind to macaque monkey TMPRSS2 and human TMPRSS2. In a preferred embodiment, the antibody of the present invention can bind to green monkey TMPRSS2 and human TMPRSS2.

[0158] In an embodiment of the present invention, the antibody of the present invention may have a binding affinity (KD) to human TMPRSS2 of 10^{-6} M or less, 10^{-7} M or less, 10^{-8} M or less, 10^{-9} M or less, or 10^{-10} M or less.

[0159] In an embodiment of the present invention, the antibody of the present invention may have IC₅₀ referring to inhibition of coronavirus infection of human cells of 100 µg/mL or less, 90 µg/mL or less, 80 µg/mL or less, 70 µg/mL or less, 60 µg/mL or less, 50 µg/mL or less, 40 µg/mL or less, 30 µg/mL or less, 20 µg/mL or less, 10 µg/mL or less, 9 µg/mL or less, 8 µg/mL or less, 7 µg/mL or less, 6 µg/mL or less, 5 µg/mL or less, 4 µg/mL or less, 3 µg/mL or less, 2 µg/mL or less, or 1 µg/mL or less.

[0160] In an embodiment of the present invention, the antibody of the present invention can bind to a peptide having the amino acid sequence of any of SEQ ID NOs: 41 to 93. In an embodiment of the present invention, the antibody of the present invention can bind to a peptide having the amino acid sequence of any of SEQ ID NOS: 85 to 93. In an embodiment of the present invention, the antibody of the present invention can bind to a peptide having the amino acid sequence of SEQ ID NO: 85.

[0161] In an embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and can bind to a peptide having the amino acid sequence consisting of the amino acid sequence selected from the group consisting of the amino acid sequence set forth in SEQ ID NO: 75 (WNEYGRAACRDMGYKNNFY), the amino acid sequence set forth in SEQ ID NO: 89 (VTAACVCEKPLNNPWHWT), and the amino acid sequence set forth in SEQ ID NO: 96 (RQSFMF).

[0162] In an embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and:

[0163] (α) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 75 (WNEYGRAACRDMGYKNNFY) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 94: WSEGYGRAACQDMGYRNSFY) with

binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto;

[0164] (β) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 95: VTAAHCVEEPLNNPRHWT) with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto; or

[0165] (γ) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 96 (RQSFMF) of the human TMPRSS2 is substituted for a corresponding sequence of a hamster (i.e., the amino acid sequence set forth in SEQ ID NO: 97: SQSLMF) with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto.

[0166] In a preferred embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and

[0167] (α) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 75 (WNE-NYGRAACRDMGYKNNFY) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 94: WSEG YGRAACQDMGYRNSFY) with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto.

[0168] In a preferred embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and

[0169] (β) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 95: VTAAHCVEEPLNNPRHWT) with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto.

[0170] In a preferred embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and

[0171] (γ) binds to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 96 (RQSFMF) of the human TMPRSS2 is substituted for a corresponding sequence

of a hamster (i.e., the amino acid sequence set forth in SEQ ID NO: 97: SQSLMF) with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto.

[0172] In an embodiment, the antibody of the present invention binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and:

[0173] (A) binds to a human TMPRSS2 variant having a point mutation at N179A with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto; or

[0174] (B) binds to a human TMPRSS2 variant having a point mutation at W306A with binding affinity less than that for the human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5 (e.g., KD value of less than 1-fold, $\frac{1}{2}$ or less, $\frac{1}{3}$ or less, $\frac{1}{4}$ or less, $\frac{1}{10}$ or less, $\frac{1}{100}$ or less, or $\frac{1}{1,000}$ or less), or does not significantly bind thereto.

[0175] In an embodiment, the antibody of the present invention may be an antagonist antibody against TMPRSS2.

[0176] In an embodiment of the present invention, the antibody of the present invention or an antigen-binding fragment thereof can be:

[0177] (1) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 11,

[0178] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 12, and

[0179] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 13,

[0180] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 14,

[0181] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 15, and

[0182] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 16.

[0183] (2) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 19,

[0184] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 20, and

[0185] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 21,

[0186] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 22,

[0187] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 23, and

[0188] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 24.

[0189] (3) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 27,

- [0190] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 28, and
- [0191] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 29,
- [0192] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 30,
- [0193] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 31, and
- [0194] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 32.
- [0195] (4) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 35,
- [0196] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 36, and
- [0197] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 37,
- [0198] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 38,
- [0199] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 39, and
- [0200] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 40.
- [0201] (5) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 100,
- [0202] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 101, and
- [0203] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 102,
- [0204] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 103,
- [0205] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 104, and
- [0206] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 105.
- [0207] (6) an antibody or an antigen-binding fragment thereof, including a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 108,
- [0208] HCDR2 having the amino acid sequence set forth in SEQ ID NO: 109, and
- [0209] HCDR3 having the amino acid sequence set forth in SEQ ID NO: 110,
- [0210] a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 111,
- [0211] LCDR2 having the amino acid sequence set forth in SEQ ID NO: 112, and
- [0212] LCDR3 having the amino acid sequence set forth in SEQ ID NO: 113.
- [0213] (7) an antibody or an antigen-binding fragment thereof, competing with the antibody according to any of (1) to (6) above for binding with TMPRSS2; or
- [0214] (8) an antibody or an antigen-binding fragment thereof, including the amino acid sequence having one or several amino acid mutations in the CDR sequence of the antibody according to any of (1) to (6) above; or
- [0215] (9) a human chimeric antibody or a humanized antibody of the antibody according to any of (1) to (8) above.
- [0216] In an embodiment of the present invention, the antibody of the present invention or an antigen-binding fragment thereof may be an antibody or an antigen-binding fragment thereof:
- [0217] (7) having a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 9, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 10 {e.g., having a heavy chain variable region and a light chain variable region of the 752_1 antibody},
- [0218] (8) having a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 17, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 18 {e.g., having a heavy chain variable region and a light chain variable region of the 1864_10 antibody},
- [0219] (9) having a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 25, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 26 {e.g., having a heavy chain variable region and a light chain variable region of the 2020_12 antibody},
- [0220] (10) having a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 33, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 34 {e.g., having a heavy chain variable region and a light chain variable region of the 2228_15 antibody},
- [0221] (11) having a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 98, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 99 {e.g., having a heavy chain variable region and a light chain variable region of the 1831_15 antibody},
- [0222] (12) competing with the antibody according to (7) above for binding with TMPRSS2.
- [0223] (12) competing with the antibody according to (10) above for binding with TMPRSS2.
- [0224] (13) competing with the antibody according to (11) above for binding with TMPRSS2; or (14) competing with the antibody according to any of (7) to (10) above for binding with TMPRSS2.
- [0225] In a preferred embodiment of the present invention, the antibody of the present invention binds to human TMPRSS2 and competes with the antibody according to (7) above for binding with TMPRSS2. Accordingly, it is possible to suppress infection of human cells with beta coronavirus including SARS-COV-2.
- [0226] In a preferred embodiment of the present invention, the antibody of the present invention binds to human TMPRSS2 and competes with the antibody according to (10) above for binding with TMPRSS2. Accordingly, it is possible to suppress infection of human cells with beta coronavirus including SARS-COV-2.
- [0227] In a preferred embodiment of the present invention, the antibody of the present invention binds to human TMPRSS2 and competes with the antibody according to (11) above for binding with TMPRSS2. Accordingly, it is possible to suppress infection of human cells with beta coronavirus including SARS-COV-2.

[0228] In an embodiment of the present invention, the antibody of the present invention may be an antagonist antibody against TMPRSS2. In an embodiment of the present invention, the antibody of the present invention inhibits the activating effect of TMPRSS2 on the S protein of coronavirus. In an embodiment of the present invention, the antibody of the present invention inhibits the interaction between TMPRSS2 and the S protein of coronavirus. In an embodiment of the present invention, the antibody of the present invention does not inhibit the interaction between TMPRSS2 and the S protein of coronavirus.

[0229] In an embodiment, the antibody of the present invention may be at least one of IgG1, IgG2, IgG3, IgG4, IgE, IgA, IgD, and IgM. In an embodiment, the antibody of the present invention contains a human κ light chain constant region and/or a human heavy chain. In an embodiment, the antibody of the present invention contains a human λ light chain constant region and/or a human heavy chain. In an embodiment, the antibody of the present invention contains a human κ light chain constant region and/or a human IgG2 heavy chain constant region. In an embodiment, the antibody of the present invention contains a human λ light chain and/or a human IgG4 heavy chain constant region. In an embodiment, the antibody of the present invention contains a human λ light chain constant region and/or a human IgG2 heavy chain constant region. In an embodiment, the antibody of the present invention contains a human κ light chain and/or a human IgG4 heavy chain constant region.

[0230] In a preferred embodiment of the present invention, the antibody of the present invention does not have significant ADCC activity. In an embodiment, the antibody of the present invention may be a human chimeric antibody, humanized antibody, or human antibody. In an embodiment, the subtypes of the human chimeric antibody, humanized antibody, or human antibody may be IgG4. In an embodiment, IgG4 may have serine at position 108 from the N-terminal of the CH1 region being substituted for proline in the CH1 region. In an embodiment, IgG4 may have serine at position 120 from the N-terminal of the CH1 region being substituted for proline in the CH1 region.

[0231] In an embodiment, the humanized antibody has a light chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 115 or 120, a framework region 2 having the amino acid sequence of SEQ ID NO: 116 or 121, a framework region 3 having the amino acid sequence of SEQ ID NO: 117 or 122, and a framework region 4 having the amino acid sequence of SEQ ID NO: 118 or 123.

[0232] In an embodiment, the humanized antibody has a light chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 115, a framework region 2 having the amino acid sequence of SEQ ID NO: 116, a framework region 3 having the amino acid sequence of SEQ ID NO: 117, and a framework region 4 having the amino acid sequence of SEQ ID NO: 118.

[0233] In an embodiment, the humanized antibody has a light chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 120, a framework region 2 having the amino acid sequence of SEQ ID NO: 121, a framework region 3 having the amino acid sequence of SEQ ID NO: 122, and a framework region 4 having the amino acid sequence of SEQ ID NO: 123.

[0234] In an embodiment, the humanized antibody has a heavy chain variable region containing a framework region

1 having the amino acid sequence of SEQ ID NO: 125 or 130, a framework region 2 having the amino acid sequence of SEQ ID NO: 126 or 131, a framework region 3 having the amino acid sequence of SEQ ID NO: 127 or 132, and a framework region 4 having the amino acid sequence of SEQ ID NO: 128 or 133.

[0235] In an embodiment, the humanized antibody has a heavy chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 125, a framework region 2 having the amino acid sequence of SEQ ID NO: 126, a framework region 3 having the amino acid sequence of SEQ ID NO: 127, and a framework region 4 having the amino acid sequence of SEQ ID NO: 128.

[0236] In an embodiment, the humanized antibody has a heavy chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 130, a framework region 2 having the amino acid sequence of SEQ ID NO: 131, a framework region 3 having the amino acid sequence of SEQ ID NO: 132, and a framework region 4 having the amino acid sequence of SEQ ID NO: 133.

[0237] In an embodiment, the humanized antibody has:

[0238] a heavy chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 125 or 130, a framework region 2 having the amino acid sequence of SEQ ID NO: 126 or 131, a framework region 3 having the amino acid sequence of SEQ ID NO: 127 or 132, and a framework region 4 having the amino acid sequence of SEQ ID NO: 128 or 133; and

[0239] a light chain variable region containing a framework region 1 having the amino acid sequence of SEQ ID NO: 115 or 120, a framework region 2 having the amino acid sequence of SEQ ID NO: 116 or 121, a framework region 3 having the amino acid sequence of SEQ ID NO: 117 or 122, and a framework region 4 having the amino acid sequence of SEQ ID NO: 118 or 123.

[0240] Any one or more of the framework regions 1 to 4 above may have one or several amino acid mutations as long as the binding affinity to the antigen is not significantly lowered. In an embodiment, the amino acid mutation is amino acid substitution.

[0241] In an embodiment, the humanized antibody has:

[0242] (1) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 124, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 114.

[0243] (2) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 129, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 114.

[0244] (3) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 124, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 119.

[0245] (4) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 129, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 119.

[0246] (5) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 144, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 134.

[0247] (6) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 149, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 134.

[0248] (7) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 144, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 139; and

[0249] (8) a heavy chain variable region having the amino acid sequence set forth in SEQ ID NO: 149, and a light chain variable region having the amino acid sequence set forth in SEQ ID NO: 139.

[0250] The antibody is not the specific antibody disclosed in WO2019/147831A (which is incorporated herein by reference in its entirety). More specifically, the antibody does not include any one or more or all of the light chain CDRs contained in the amino acid sequence of SEQ ID NO: 4 or 18 in WO2019/147831A, or any one or more or all of the heavy chain CDRs contained in the amino acid sequence of SEQ ID NO: 2, 17, or 19 in WO2019/147831A. Further, the antibody does not include any or all of the CDRs set forth in SEQ ID NOS: 25 to 27, or any or all of the CDRs set forth in SEQ ID NOS: 29 to 31 in WO2019/147831A. Moreover, the antibody does not include any or all of the CDRs set forth in SEQ ID NOS: 33 to 35, or any or all of the CDRs set forth in SEQ ID NOS: 37 to 39 in WO2019/147831A. The antibody does not include any or all of the CDRs set forth in SEQ ID NOS: 41 to 43, or any or all of the CDRs set forth in SEQ ID NOS: 45 to 47 in WO2019/147831A.

[0251] The antibody is not the specific antibody disclosed in WO2021/163076A (which is incorporated herein by reference in its entirety). Specifically, the antibody does not include any one or more or all of the CDRs contained in the heavy chain variable region set forth in SEQ ID NO: 2, or any one or more or all of the CDRs contained in the light chain variable region set forth in SEQ ID NO: 10 in WO2021/163076A. The antibody does not include any one or more or all of the CDRs contained in the heavy chain variable region set forth in SEQ ID NO: 22, or any one or more or all of the CDRs contained in the light chain variable region set forth in SEQ ID NO: 30 in WO2021/163076A. The antibody does not include any one or more or all of the CDRs contained in the heavy chain variable region set forth in SEQ ID NO: 42, or any one or more or all of the CDRs contained in the light chain variable region set forth in SEQ ID NO: 50 in WO2021/163076A.

[0252] The antibody is not the specific antibody disclosed in WO2021/211406 (which is incorporated herein by reference in its entirety). The antibody is not the specific antibody disclosed in WO2021/211416 (which is incorporated herein by reference in its entirety). The specific antibody refers to, for example, an antibody whose whole or a part (e.g., all or a part of CDRs) is defined by the amino acid sequence.

[0253] In an embodiment, the antibody is monospecific. In an embodiment, the antibody is Mult specific. In an embodiment, the antibody is bispecific.

[0254] The antibody can be obtained by a method well known to those skilled in the art. For example, recombinant proteins or cells that express recombinant proteins are used to immunize a TMPRSS2 KO animals, and then, antibodies that bind to TMPRSS2 can be obtained. Monoclonal antibodies can be obtained as culture supernatants of hybridomas obtained by fusing splenocytes of the immunized animals with myeloma. Monoclonal antibodies can also be

obtained from the ascites or the like of animals into which hybridomas was transplanted. Also, monoclonal antibodies may be obtained from B cells by gene cloning. Alternatively, the antibodies can be obtained by a well-known method such as phage display. The antibodies can be modified to prepare chimeric antibodies (e.g., human chimeric antibodies) or humanized antibodies as necessary. The antibodies can be produced from animals (e.g., mice) having human antibody loci (e.g., variable regions). The obtained antibodies are tested in terms of whether or not they have properties to bind to TMPRSS2, and whether or not they have properties to inhibit coronavirus infection of cells, and antibodies that have such properties can be selected from the obtained antibodies. The obtained antibodies can also be tested depending on whether or not they further have one or more properties selected from the group consisting of the binding affinity (KD) to TMPRSS2, IC₉₀ referring to infection inhibition, properties to bind to specific amino acids, and properties to compete with reference antibodies for binding to TMPRSS2, and antibodies having further useful properties can be selected from antibody pools.

[0255] When useful antibodies are obtained, the antibodies can be industrially produced from antibody-producing cells such as Chinese hamster ovary (CHO) cells, which have genes for encoding antibodies.

[0256] The competition can be determined, for example, by binding labeled antibodies to TMPRSS2 expressing cells and incubating them in the presence of unlabeled antibodies at various concentrations and based on the decreased amount of label bound to cells. For example, when an antibody competes with another antibody, the amount of label bound to cells can be decreased to 50% or less, 40% or less, 30% or less, 20% or less, or 10% or less.

[0257] KD of the antibodies can be appropriately determined by those skilled in the art. KD can be determined from K_{on} and K_{off} by, for example, the surface plasmon resonance (SPR) method. For example, by the following formula:

$$KD = K_{off} / K_{on}$$

[0258] KD can be determined. In other words, antibodies having small KD are considered to have strong binding affinity, and for example, the antibodies tend to easily bind to their antigen and not easily dissociate therefrom. The surface plasmon resonance (SPR) method can be performed by, for example, commercially available apparatus (e.g., Biacore (trademark)).

[0259] IC₉₀ referring to infection inhibition is the concentration of test antibodies that can inhibit infection by 90% in the absence of the test antibodies. It is indicated that the smaller IC₉₀ is, the greater the infection inhibiting effects of the test antibodies. IC₉₀ can be determined by examining the infection inhibiting effects in the presence of the test antibodies having various concentrations. Usually, IC₉₀ can be more precisely determined by conducting experiments using solutions of test antibodies at a lower concentration and a higher concentration than IC₉₀.

[0260] The properties to bind to a specific amino acid can be confirmed, for example, depending on whether or not the test antibodies bind to a peptide having a specific amino acid sequence (e.g., peptide having the amino acid sequence of any of SEQ ID NOS: 41 to 93 and 96). This confirmation method can be appropriately conducted using a well-known technique such as ELISA by those skilled in the art.

[0261] In an embodiment of the present invention, the antibody of the present invention is an antibody that binds to an extracellular domain of TMPRSS2 and is capable of inhibiting coronavirus infection of cells. In an embodiment of the present invention, the antibody of the present invention is an antibody that binds to an extracellular domain of human TMPRSS2 and is capable of inhibiting coronavirus infection of human cells. The coronavirus may be a beta coronavirus, and in a preferred embodiment, may be a beta coronavirus such as SARS-COV, MERS, or SARS-COV-2. In a particularly preferred embodiment, the coronavirus may be the alpha variant, the beta variant, the gamma variant, the delta variant, and the omicron variant of SARS-COV-2. According to the present invention, since the coronavirus infection is inhibited by inhibiting human proteins, it is theoretically assured to provide an antibody that are effective in inhibiting coronavirus infection, regardless of mutations in S proteins of the coronavirus.

[0262] In an embodiment of the present invention, the antibody of the present invention is an antibody that binds to an extracellular domain of human TMPRSS2, whose binding affinity (KD) to the human TMPRSS2 is 10^{-8} M or less, and that is capable of inhibiting coronavirus infection of human cells. In an embodiment of the present invention, the antibody of the present invention may be an antibody that binds to an extracellular domain of human TMPRSS2, whose IC_{90} referring to inhibition of the coronavirus infection of human cells is $10 \mu\text{g/mL}$ or less. In an embodiment of the present invention, the antibody of the present invention may be an antibody that binds to an extracellular domain of human TMPRSS2, whose binding affinity (KD) to the human TMPRSS2 is 10^{-8} M or less, and whose IC_{90} referring to inhibition of the coronavirus infection of human cells is $10 \mu\text{g/mL}$ or less. The coronavirus may be a beta coronavirus, and in a preferred embodiment, may be a beta coronavirus such as SARS-COV, MERS, or SARS-COV-2. In a particularly preferred embodiment, the coronavirus may be the alpha variant, the beta variant, the gamma variant, the delta variant, and the omicron variant of SARS-COV-2.

[0263] The Centers for Disease Control and Prevention (CDC) designated as variants of concern (VOC) and variants of interest (VOI) of SARS-COV-2, Alpha (B.1.1.7, Q.1-Q.8), Beta (B.1.351, B.1.351.2, B.1.351.3), Gamma (P.1, P.1.1, P.1.2), Epsilon (B.1.427 and B.1.429), Eta (B.1.525), Iota (B.1.526), Kappa (B.1.617.1), B.1.617.3, Mu (B.1.621, B.1.621.1), Zeta (P.2). In addition to that, the omicron variant (B.1.1.259, BA lineage) was added as a VOC, in November 2021. The antibody of the present invention can be effective against all of these variants.

[0264] Preferably, the antibody of the present invention also exhibits the binding properties with one or more extracellular domains of TMPRSS2 in laboratory animals, for example, selected from the group consisting of mice, green monkeys, cats, hamsters and macaque monkeys, from the viewpoint of conducting animal testing before human clinical trials.

<Composition Containing Antibody of the Present Invention>

[0265] According to the present invention, a composition containing the antibody of the present invention is provided. According to the present invention, a pharmaceutical composition containing the antibody of the present invention is also provided. The pharmaceutical composition of the pres-

ent invention can further contain a pharmaceutically acceptable additive in addition to the antibody.

[0266] The composition or pharmaceutical composition of the present invention can be administered in a subject for use in inhibiting cell entry of coronavirus in the subject. Therefore, the composition or pharmaceutical composition of the present invention can be used in subjects who are not infected with coronavirus to prevent them from being infected with coronavirus. Also, the composition or pharmaceutical composition of the present invention can be used in subjects infected with coronavirus to prevent coronavirus from being spread from the infected cells. Therefore, the composition or pharmaceutical composition of the present invention can also be used to treat subjects infected with coronavirus. Administering to subjects infected with coronavirus the composition or pharmaceutical composition of the present invention make it possible to prevent the symptoms of the coronavirus infection from deteriorating or becoming severe, or to improve the symptoms in the subjects. Preventing deterioration includes delaying deterioration and inhibiting deterioration. The coronavirus may be a beta coronavirus, may be a coronavirus such as SARS-COV, MERS, or SARS-COV-2, and in particular, may be a coronavirus variant designated as VOC or VOI by The World Health Organization (WHO).

[0267] A subject to whom the composition or pharmaceutical composition of the present invention is administered is preferably a human. However, the coronavirus infection is known to be zoonotic infection. Also, coronaviruses having high pathogenicity in animals are known such as mouse hepatitis virus, feline infectious peritonitis virus, and porcine epidemic diarrhea virus. Accordingly, the composition or pharmaceutical composition of the present invention can be used to prevent infection of these coronaviruses and treat their symptoms. In this case, the composition or pharmaceutical composition of the present invention needs to contain an antibody capable of binding to TMPRSS2 in the target animal species and inhibiting the infection of cells of the animal species with the target coronavirus.

<Other Applications>

[0268] According to the present invention, provided is use of the antibody of the present invention or an antigen-binding fragment thereof, in production of a medicine for inhibiting cell entry of coronavirus in a subject.

[0269] According to the present invention, provided is use of the antibody of the present invention or an antigen-binding fragment thereof, for inhibiting cell entry of coronavirus in a subject.

[0270] According to the present invention, provided is the composition or pharmaceutical composition of the present invention for use in a method for inhibiting cell entry of coronavirus in a subject.

[0271] According to the present invention, provided is use of the antibody of the present invention or an antigen-binding fragment thereof for use in a method for inhibiting cell entry of coronavirus in a subject.

[0272] According to the present invention, provided is a method for preventing coronavirus infection in a subject, including administering to the subject an effective amount of the antibody of the present invention or an antigen-binding fragment thereof.

[0273] According to the present invention, provided is a method for treating a subject infected with coronavirus,

including administering to the subject an effective amount of the antibody of the present invention or an antigen-binding fragment thereof.

[0274] According to the present invention, provided is a method for preventing coronavirus infection from deteriorating and/or becoming severe in a subject infected with coronavirus, including administering to the subject an effective amount of the antibody of the present invention or an antigen-binding fragment thereof.

EXAMPLES

[Materials and Methods]

Gene Synthesis

[0275] All amino acid sequences were obtained from UniProtKB. From the amino acid sequences, DNA that encodes the optimum human codon usage sequence was designed and chemically synthesized to produce a recombinant protein and a transfection cell line (Eurofins Genomics).

Construction of Plasmid Vector

[0276] The synthesized DNA was amplified by PCR using a forward primer and a reverse primer that were added with tag sequences at the C-terminal and/or the N-terminal and the 3' and 5' regions of homology required for In-Fusion (trademark) cloning. The sequences of these extensions must match precisely the 15 bp of the 5' end and 3' end of the recipient vector exposed by linearization of the vector at the position into which the PCR product is inserted.

[0277] For the retrovirus expression vector pMys-IRES-GFP (CELL BIOLABS, INC.), the synthesized DNA was amplified by PCR using a primer set with a restriction digestion site added in place of the 15 bp addition sequence for In-Fusion (trademark) cloning. Using a DNA fragment after being digested with each restriction enzyme, the retrovirus expression vector was inserted. The correct sequence of the cDNA in the expression vector was confirmed by DNA sequencing.

Serine Protease Domain of Human TMPRSS2 against *E. coli*-Derived Immunizing Antigen

[0278] DNA fragments that encode the serine protease domain of human TMPRSS2 (containing a His sequence at the C-terminal) were inserted into pE-SUMOstar Amp (LifeSensors, Inc.) and pMAL-p5X Vector (BioLabs) using In-Fusion (trademark) cloning system (Takara Bio Inc.). In order to produce recombinant proteins, plasmid vectors were introduced into *E. coli* BL21 strain by heat shock, and colonies grown by culturing in a nutrient agar medium containing ampicillin at 37° for 24 hours, were cultured in a nutrient broth medium containing ampicillin for 24 hours, and the cultured bacteria were transferred to an inducing medium to induce the production of the recombinant proteins, and placed in a shaker incubator at 24 rpm. After the number of bacteria reached an appropriate level (absorbance of 0.6), 1 mM IPTG solution was added into bacterial cell suspension. At 2 hours and 4 hours after the addition of the IPTG solution, centrifugation was performed at 4000 rpm for 8 minutes to precipitate bacteria. The bacterial cells were pelleted by centrifugation (6000 g×15 minutes) and washed with PBS. The pellets were suspended in 10 mL lysis buffer (50 mM potassium phosphate buffer, pH 7.8, 0.5 M NaCl, 5 mM MgCl₂, 1 mg/mL lysozyme, 10 µg/mL DNase) and

subjected to sonification in a water bath (Branson 200) for 5 minutes, and then the lysate was incubated on a shaker at room temperature for 30 minutes. Ultracentrifugation (150000×g for 35 minutes) was performed to collect supernatant. Fractions from sucrose gradient were collected using a bent Pasteur pipette.

Production of Recombinant Protein of Human Full-Length TMPRSS2 Against Immunizing Antigen from Insect Cells

[0279] Full-length cDNA of human TMPRSS2 having a His sequence at the C-terminal was inserted into a pFast-Bac1 vector using In-Fusion (trademark) cloning system (Takara Bio Inc.).

[0280] Using Bac-to-Bac Baculovirus Expression System (Thermo Fisher Scientific, Inc.), recombinant proteins were expressed in HighFive insect cells (Thermo Fisher Scientific, Inc.), and the cultured cells were homogenized and disrupted. Membrane fractions were separated and solubilized with 1% dodecyl maltoside (DDM, Anatrace, Maumee, OH).

Harvest of Recombinant Proteins for ELISA and BIAcore Assays from Mammalian Cells.

[0281] A pcDNA3.4 expression vector was purchased from Thermo Fisher, Inc. PCR fragments that encode Strep-tag II (WSHPQFEK) at the C-terminal and His tag at the N-terminal of the extracellular region of the human TMPRSS family were inserted into an expression vector.

[0282] Expi293F cells were cultured in a humidified 8% CO₂ incubator at 37° C., 120 rpm, using Expi293 Expression Medium (Thermo Fisher, Inc., Cat. A1435101). On the day of transfection, the density of Expi293 cells was approximately 4.0×10⁶ cells/mL. The cells were centrifuged at 200×g for 15 minutes at room temperature and the medium was decanted, and then the cells were resuspended in 100% fresh medium to a density of 2.5×10⁶ cells/mL. Expi293 transfection was performed according to the manufacturer's protocol except for 100% replacement of the medium before the transfection described above. For all Expi293 transfections, Expi293 Expression System Kit (Thermo Fisher, Inc., Cat. A14635) containing a transfection enhancer and ExpiFectamine 293 transfection reagent was used. The ExpiFectamine 293 transfection reagent and plasmid DNA that were separately diluted with a complex medium, OptiMem (Thermo Fisher, Inc., Cat. 31985062) were used. After 5-minute incubation, the ExpiFectamine 293 and the DNA mixture were combined and incubated for further 20 minutes. The mixture of ExpiFectamine 293-DNA-OptiMEM was then added into the cells. An enhancer 1 and an enhancer 2 were added to the transfected culture 16 to 18 hours after transfection. The recombinant proteins were purified with Ni-NTA affinity chromatography.

Purification of Recombinant Proteins

[0283] The recombinant proteins were purified with Ni-NTA affinity chromatography. Resin was added thereto to bind to the supernatant at 4° C. The next day, the mixture was moved to a column, and the pass-through fractions were removed, and then the resultant was washed with Tris buffer (pH 7.0) containing 0.05% DDM, 0.002% Cholesteryl Hemisuccinate (CHS Anatrace Cat no CH210) and 5 mM DTT. After that, using wash buffer added with 2.5 mM desthiobiotin, recombinant proteins were eluted from the resin.

Retroviral Vector and Transfection Cell Line

[0284] Into the retrovirus expression vector pMys-IRES-GFP, the full-length TMPRSS2 family gene having an HA-tagged sequence (amino acid sequence: YPYDVPDYA) at the C-terminal was inserted.

[0285] Daudi lymphoma cell line was obtained from JCRB Cell Bank (JCRB9071). A20 and EL4 lymphoma cell lines were obtained from RIKEN-BRC Cell Bank. These cell lines were introduced using a retroviral vector as follows. A retrovirus expression vector and an envelope expression vector, p10A1 or pAmpho, were co-introduced into GP2-293 cells using a Retro-X Universal Packaging system (Takara Bio Inc., #631530). 2 to 3 days after transfection, the culture supernatant was filtered with a 0.45 μ m filter, and then infected with the above cell lines using 10 μ g/ml polybrene (Sigma Aldrich Co. LLC, #P8155). After 1 week from the introduction, the cells expressed the introduced molecules on the surface. Finally, the cells were sorted with a purity of 98% or more using FACS Aria III (BD Biosciences).

Immunization

[0286] TMPRSS2 KO mice were provided by Dr. Takeda (The National Institute of Infectious Diseases), and subcutaneously primed with TMPRSS2-serine protease domain recombinant protein (50 μ g/mouse) using adjuvant TiterMax Gold (TiterMax USA, Inc.), mice were alternately immunized in the abdominal cavity with TMPRSS2 transfectants (2×10^7 cells/mouse) and recombinant proteins (50 μ g/mouse) at 2-week intervals. The TMPRSS2 transfectant was treated with gamma radiation (10 Gy) before immunization. ODN 1826 Vaccine Grade (Nacalai tesque, Inc., Japan) was used for immunization of the TMPRSS2 transfectant cells. These mice were boost-immunized with full-length TMPRSS2 recombinant protein (10 μ g/mouse) by tail vein injection 3 days before the fusion. Spleen B-cells were purified from the immunized mice and used for hybridoma preparation.

Enrichment of IgG+ B Cells from Immunized Mice

[0287] Mature B cells were separated by negative selection using the MASC system (Miltenyi Biotec, Bergisch Gladbach, Germany). Unwanted mononuclear cells (T cells, NK cells, monocytes, dendritic cells, granulocytes, and immature B cells prior to B cell receptor class switching) were targeted via a biotin-labeled antibody cocktail specific to the cell surface receptors of these cell types. Also, mature B cells were specifically enriched from the sample without retention in the LS column in the presence of a magnetic field.

Cell Fusion and Hybridoma Selection

[0288] Mature B cells enriched from the spleens of immunized mice were mixed with P3U1 myeloma cells at a ratio of 1:1, and the mixture was subjected to electrofusion using a Nepagene ECFG21 electroporator (Nepa Gene Co., Ltd., Ichikawa, Chiba, Japan) according to the manufacturer's instruction. The cells were then seeded at 2×10^3 cells/well in 96-well flat-bottom plates and selected in a culture solution containing a HAT supplement (Sigma Aldrich Co. LLC) and a BM Condensed H1 supplement (F. Hoffmann-La Roche Ltd.). When the cells covered half of the bottom of the wells, the culture supernatant was examined by ELISA for reactivity with TMPRSS2 recombinant protein. A portion of the

hybridoma cells obtained from positive culture supernatants was frozen in CELLBANKER 1 plus (Takara Bio Inc.). The remaining cells were passage cultured to confluent, and then each supernatant was used for flow cytometry screening and inhibition screening for cell fusion analysis.

ELISA Screening

[0289] To a Nunc MaxiSorp flat-bottom 96-well microplate, 50 mM carbonate coating buffer containing 0.2 μ g/mL recombinant protein of human TMPRSS2, pH 9.6 (100 μ L) was applied and incubate at 4° C. overnight. The plate was then washed with PBST wash buffer (0.05% Tween 20/PBS) three times, 1% BSA blocking buffer was added thereto at 200 μ L/well to incubate for 2 hours at room temperature. After washing, 100 μ L hybridoma supernatant was added thereto, and the mixture was incubated for 2 hours followed by washing. 100 μ L HRP-labeled anti-mouse IgG antibody (MBL #330) was added to each well, and then the microplate was incubated for 1 hour. After washing the plate, 100 μ L/well of TMB substrate solution (SeraCare Life Sciences, Inc. 5120-0077) was added thereto, and reacted in the dark for 15 minutes. The reaction was stopped by adding 2 M sulfuric acid (50 μ L/well), and then the absorbance was measured at OD450 using a microplate reader (ARVO $\times$ 3, PerkinElmer, Inc.).

Screening by Flow Cytometry

[0290] For screening of the ELISA positive supernatant by flow cytometry, the Daudi cell line expressing TMPRSS2 and GFP and the parent cells of Daudi were mixed at a ratio of 1:1. The cells were then stained with 100 μ L hybridoma supernatant in a 96-well plate, stored on ice for 20 minutes, and then incubated with allophycocyanin (APC)-labeled anti-mouse IgG (BioLegend, Inc., Poly4053). The cells were washed, resuspended in propidium iodide (Nacalai tesque, Inc., Japan), and the staining of TMPRSS2 was analyzed using LSR Fortessa X-20 (BD Biosciences, Franklin Lakes, NJ). Hybridomas whose supernatant reacted with TMPRSS2+Daudi transfectant and did not react with the parent cell line were selected.

Screening of Dual Split Protein (DSP) Reporter Assay to Monitor Membrane Fusion (Cell Fusion Assay)

[0291] In order to establish a stable 293 FT cell line that expresses spike proteins (S proteins) with DSP8-11, and expresses ACE2 and TMPRSS2 with DSP1-7, 293T cells introduced with VSV-G expressing plasmid were used to produce recombinant pseudo-typed retroviruses that express any of these proteins. Reporter cells derived from 293 FT infected with pseudo-typed virus were selected with 1 μ g/mL puromycin, 10 μ g/mL blastidine, and 300 μ g/mL hygromycin for at least 1 week. Fusion assay was performed using these batch-selected cells.

[0292] In DSP assays using 293 FT cells, effector cells that express S proteins using DSP8-11, and target cells that express ACE2 and TMPRSS2 using DSP1-7 were seeded on a 12-well cell culture plate 1 day before the assay (2×10^5 cells/500 μ L). 2 Hours before the DSP assay, the cells were treated with a *Renilla* luciferase substrate, 6 μ M EnduRen (Promega, Madison, WI, USA) to activate EnduRen. 50 μ L Target cells that express ACE2 and TMPRSS2 were seeded on a 384-well plate using a multidrop dispenser (Thermo Scientific, Waltham, MA, USA), and the hybridoma super-

nanatant was added to the wells. After incubation at 37° C. for 1 hour, suspension of the effector cells expressing S proteins was added to the wells using a multidrop dispenser. After incubation at 37° C. for 4 hours, the RL activity was measured using a Centro xS960 luminometer (Berthold Technologies GmbH, Germany). The inhibition rate calculated as relative light units (RLU) of the luciferase activity compared to control was reported as neutralizing antibody sensitivity of the hybridoma supernatant in the DSP reporter assay.

First Cloning of Hybridomas

[0293] Hybridomas of interest among 3,723 candidates were collected from a frozen cell stock, cultured in a HT medium overnight, and then cloned by limiting dilution (0.3 cells/well). Clones generally appeared in 8 to 15 days, and 120 clones obtained by limiting dilution were screened again for antibody reactivity, using flow cytometry analysis and inhibitory functions for DSP assays with 293 FT effector cells that express S proteins and 293 FT target cells that express ACE2 and TMPRSS2 or Calu-3 target cells introduced with DSP1-7.

[0294] Further, mouse IgG concentrations of the 120 hybridoma supernatants were analyzed by ELISA, and mAb concentrations (IC50) that reduces RLU by 50% compared to control were reported as neutralizing antibody titers for the DSP reporter assay with the 293 FT target cells and 293 effector cells. Titers were calculated using a non-linear regression curve fit (GraphPad Prism Software Inc., La Jolla, CA).

Second Cloning of Hybridomas

[0295] 20 Clones of interest from the independent hybridomas selected by the above functional screening targeting 120 clones were collected from the frozen cell stock, cultured in HT and BM-Condensed H1 media overnight, and then cloned again by limiting dilution (0.3 cells/well) to confirm the reactivity to TMPRSS2 by FACS analysis.

[0296] In order to adapt to a passage medium, concentrations of FCS, HT, BM-Condensed H1 were stepwisely reduced in the HT medium, cell growth was repeatedly confirmed, and eventually the cells were adapted to culture in 10% FCS RPMI-1640 medium without adding HT and BM-Condensed H1.

Identification of mAb Isotype

[0297] After subcloning twice, the isotype of mouse mAb was identified with Rapid Mouse Monoclonal Antibody Isotyping Kit according to the manufacturer's instruction (Antigen, ISO-M8ac).

Adaptation of Cells to Serum-Free Medium and Production of mAb

[0298] Hybridomas were passaged in 10% FCS RPMI-1640 medium and subjected to 3 passages in each of hybridoma-SFM media (manufactured by Thermo Fisher Scientific, Inc.) containing 6%, 4%, 2%, or 1% fetal bovine serum (FBS) followed by serial culture in a serum-free medium. Samples of 5×10^7 cells were gathered and transferred to CELLline flask culture (BD Biosciences) using the serum-free medium according to the manufacturer's instruc-

tion. After culturing at 37° C. with 5% CO₂ for 14 days, the cells and media were collected from a cell culture chamber. Purification of mAbs from Hybridoma Supernatant Fluid **[0299]** Monoclonal antibodies in culture supernatant collected in the serum-free medium or 10% Ultra Low IgG FBS medium (Thermo Fisher Science, Inc.) were purified with Protein-A Sepharose 4 Fast Flow according to the manufacturer's instruction. The monoclonal antibodies were purified with (GE healthcare), aseptically filtered, and then stored at 4° C.

Production of SARS-COV-2 Pseudo Virus and Neutralization Assay of Anti-TMPRSS2 mAbs

[0300] In order to establish stable cell lines that express SARS-COV-2 pseudo-typed S proteins, lentiviruses were prepared using HEK293T cells containing lentivirus transfer plasmids, psPAX2 packaging plasmids, and vesicular stomatitis virus (VSV)-G expression plasmids.

[0301] Neutralization assay was performed with Calu-3 cell lines. Pseudo viruses having a luciferase activity titer of about 10⁶ RLU/ml were incubated with an antibody at 37° C. for 1 hour. The mixture of pseudo viruses and the antibody (100 µl) was then seeded on a 96-well plate in which 3.0×10^4 cells/well have been seeded 1 day before infection. The infectivity of the pseudo viruses was scored in terms of the luciferase activity after 48 hours.

Isolation of SARS-COV-2

[0302] VeroE6 (ATCC CRL-1586) cells were maintained in Eagle's minimal essential media (MEM) containing 10% FBS. Respiratory swabs were taken from a patient who tested positive for COVID-19. SARS-COV-2 virus was proliferated in VeroE6 cells at 37° C. using Opti-MEM (Invitrogen) containing 0.3% bovine serum albumin (BSA) and 1 µg L-1-tosylamide-2-phenylethyl chloromethyl ketone (TPCK)-trypsin/ml. All experiments using SARS-COV-2 virus were conducted in an enhanced biosafety level 3 (BSL3).

SARS-COV-2 Infection Assay

[0303] Calu-3 cells were cultured in an environment of 37° C. with 5% CO₂ in MEM containing 10% FBS. In vitro infection experiments using Calu-3 cells as target cells were conducted in 2 groups of "pretreatments" and "without pretreatment". In the "pretreatment" group, cells were pretreated with nafamostat mesylate (10-fold serial dilutions from 100 µM to 1 nM, 4 wells per dilution) or anti-TMPRSS2 mAb 1 hour before infection. SARS-COV-2 was then added thereto at a multiplicity of infection (MOI) of 0.1 in the absence of TPCK-trypsin, and the cells were further incubated for 24 hours to facilitate virus entry.

[0304] All cell samples were thoroughly washed with PBS, and then used for RNA extraction and RT-PCR according to SuprePrep II Cell Lysis & RT Kit for qPCR (TOYOBO Co., Ltd.). In a real-time RT-PCR multiplex used for SARS-CoV-2 detection, the expression levels of SARS-COV-2 N and rpl13a genes were quantified, and the expression level of SARS-COV-2 N was corrected by the expression level of rpl13a.

[0305] Plots of SARS-COV-2 N and rpl13a were each evaluated in the green channel (excitation wavelength of 470 nm, detection wavelength of 510 nm, fluorescent FAM-labeled probe) and the orange channel (excitation wave-

length of 585 nm, detection wavelength of 610 nm, fluorescent ROX-labeled probe). The results were analyzed taking into consideration of cycle threshold (Ct) values. The mAb concentration that was reduced by 90% (IC90) compared to control was reported as the neutralizing antibody titer. Titers were calculated using a non-linear regression curve fit (GraphPad Prism Software Inc., La Jolla, CA).

Competitive Inhibition Between Anti-TMPRSS2 mAbs

[0306] For these studies, analysis using a flow cytometer was employed. Briefly, the staining concentrations of APC-labeled mAbs were determined by the concentrations showing a value of 80% of the maximum fluorescence intensity against Daudi-TMPRSS2 transfectant cell line. Competitors of unlabeled mAb were diluted 3-fold at concentrations ranging from 0.1- to 10-fold of APC-labeled mAb and stained on ice for 30 minutes. Thereafter, a labeled mAb for detection was added to the cell suspension and further stained for 30 minutes. The cells were washed and resuspended in propidium iodide for analysis.

ELISA Binding Assay for Cross-Reactivity of Anti-TMPRSS2 mAb to Extracellular Regions of Human TMPRSS Family Proteins

[0307] The reactivity of anti-TMPRSS2 mAbs to extracellular regions of human TMPRSS family proteins was analyzed by ELISA method. To a streptavidin-coated 96-well plate (Thermo Scientific, 436014), 100 μ l each of 78 nM Strep-tagged recombinant proteins in PBS was captured at room temperature for 2 hours and washed 5 times with PBST (0.1% Tween 20/PBS). Anti-TMPRSS2 mAbs (10 μ g/mL), mouse IgG (Equitech-Bio, SLM66) or Ni-NTA-HRP (KPL, 24-01-01) as controls, diluted to 0.2% BSA in PBST was added thereto to incubate for 2 hours. After washing with PBST, the bound IgG was detected with 100 μ l horseradish peroxidase Goat anti-mouse IgG antibody (BioLegend, Inc.) diluted to 1:15,000 in PBST at room temperature for 1 hour and washed with PBST. 100 μ l TMB substrate solution (SeraCare Life Sciences, Inc. 5120-0077) was added to the well and allowed to color at room temperature for a few minutes. The reaction was stopped with 100 μ l 2M H₂SO₄. The optical density (OD) was read at 450 nm while referring to OD with a microplate reader (ARVO $\times$ 3, PerkinElmer, Inc.).

BIACore Binding Assay

[0308] BIAcore surface plasmon resonance system (BIAcore, Inc., Piscataway, NJ) was used at 25° C. for kinetic binding analysis of extracellular regions of anti-TMPRSS2 mAbs and human TMPRSS2. Sensor chips to which rabbit anti (mouse IgG) was covalently attached by amino coupling were prepared according to the manufacturer's instruction (BIAcore AB, Stevenage, Hertfordshire, UK).

[0309] Human TMPRSS2 recombinant proteins at various concentrations were injected in PBS containing 0.05% Tween 20 at a flow rate of 35 μ l/min for binding assay. The change in refractive index at the time of binding was used for kinetic measurement. The affinity of TMPRSS2 binding to mAb was calculated from on and off rates. Samples whose concentrations were raised to obtain stable measurements were injected therein, and the net increase in response units (RU) with background correction was recorded. Data analysis was performed by a method such as the Scatchard method, where RU is plotted relative to injected VEGF165

concentration, or values of RU divided by concentration (bound/total) are plotted relative to RU (bound) (the affinity as Kd is obtained from $-(1/\text{slope})$).

FACS Analysis for Cross-Reactivity of Anti-TMPRSS2 mAbs to Human TMPRSS Family Proteins and TMPRSS2 Derived from Various Animal Species

[0310] Daudi-TMPRSS family transfected cell line was stained on ice with anti-TMPRSS2 mAbs (1 μ g/mL) for 20 minutes, and then incubated with allophycocyanin (APC)-labeled anti-mouse IgG (BioLegend, Inc., Poly4053). The cells were washed, resuspended in propidium iodide (Nacalai tesque, Inc., Japan), and then the staining of TMPRSS2 was analyzed using LSR Fortessa X-20 (BD Biosciences, Franklin Lages, NJ).

Epitope Mapping by Peptide Array

[0311] Epitopes of mAbs were defined with synthesized peptides purified with a microarray (JPT Peptide Technologies GmbH, Berlin, Germany). A 20-meric peptide library that covers the human TMPRSS2 extracellular region was synthesized, immobilized on a glass slide of a microarray, and then detected with anti-TMPRSS2 mAbs for the sequential antibody profiling experiments (Table 2). The profiling experiments were conducted at JPT Peptide Technologies GmbH (Berlin, Germany).

Mutation Introduction of Human TMPRSS2 and Transfectant Thereof

[0312] For mutagenesis, the inserted human TMPRSS2 CDNA was ligated into the expression vector pIRES2-ZsGreen using its unique XhoI and EcoRI sites (Takara 632478). Mutant human TMPRSS2 cDNA was obtained by KOD-Plus-Mutagenesis Kit (TOYOBO Co., Ltd., #SMK-101) using primers in Table 3. All constructs were confirmed by DNA sequencing. Transfection was performed using COS7 cells. Expression of wild-type and mutant TMPRSS2 was detected by staining with indicated monoclonal antibodies.

Analysis of mAb Sequences

[0313] The cDNAs that encodes variable heavy (VH) and light (VL) genes of mouse mAbs were obtained by PCR using single-stranded cDNA libraries from each of the hybridoma cell lines. Total RNA was extracted from the hybridoma cells using TRIzol Reagent (Invitrogen, 15596026) according to the manufacturer's instruction. SuperScript (trademark) III First-Strand Synthesis System (Thermo Fisher Science, Inc., 18080-051) was used. The amplification of VH-CH1 and VL-CL genes was performed using a mixture of 12 or 10 forward primers designed for complementing each N-terminal sequence of FVH- or FVL-encoding region, and a reverse primer designed for complementing each C-terminal sequence of CH1 or CL region. The PCR products were cloned into pCR2.1-TOPO vector and the sequencing analysis was conducted.

Preparation of Recombinant Human-Mouse Chimeric IgG4

[0314] Gene sequences that encode VH and VL were fused to gene that each encode human IgG4 heavy chain and human κ light chain constant regions. The above genes that encode chimeric antibody chains were cloned into a pcDNA3.4 vector downstream of a leader sequence of the immunoglobulin kappa light chain. Before transfection into

expi293F cells, all the expression vectors were verified by DNA sequencing. Strategies similar to those used to prepare recombinant antigens were used for obtaining chimeric antibodies as well.

[0315] To avoid Fab arm replacement, mouse/human chimeric antibodies were prepared in which the heavy chain constant region was replaced with the human IgG4 S108P variant (Silva et al., J Biol Chem, 2015, vol. 290:5462-5469).

Antibody-Dependent Cell Cytotoxicity Assay (ADCC)

[0316] For these studies, calcein release analysis was used. Daudi that expresses TMPRSS2 was used for target cells, and KHYG-1 that expresses Fc receptor was used for effector cells. Target cells were labeled using Calcein-AM (1 μ g/ml, Nacalai tesque, Inc., Japan) at 37° C. for 1 hour while occasionally agitating, washed, and then plated on a 96-well U bottom plate at a density of 1×10^4 cells/well. The indicated mAbs were added at various concentrations, and the effector cells were added at an effector:target (E:T) ratio of 10:1. After incubation at 37° C. with 5% CO₂ for 3 hours, each supernatant was collected and analyzed for the fluorescence with an excitation filter using ARVO $\times 3$ (PerkinElmer, Inc.). 485 $\pm$ 9 nm; band pass filter. Analysis was performed using 530 $\pm$ 9 nm). Spontaneous release was determined by incubating the target cells in the medium alone, and the maximum release was obtained by suspending the cells in 1% Triton X-100. The ADCC rate of each sample (triplicate) was calculated using the following formula.

$$\% \text{ Lysis} = (\text{experimental release} - \text{spontaneous release}) / (\text{maximum release} - \text{spontaneous release}) \times 100.$$

[Results]

[0317] TMPRSS2 is a human protein expressed in human airway cells. It is required for TMPRSS2 that spike(S) proteins of viruses are cleaved by TMPRSS2 such that coronaviruses such as betacoronaviruses (e.g., SARS-COV-2) infect human airway cells (Tomita et al., Journal of Virology, 95(12), 2021, doi/10.1128/JVI.00434-21).

[0318] In a dual split protein (dsp) reporter assay to monitor membrane fusion (cell fusion assay), ACE2 expressing cells and S protein expressing cells are fused by the interaction between ACE2 and S protein. DSP1-7 is expressed in ACE2 expressing cells and DSP9-11 is expressed in S protein expressing cells; when the two are fused, DSP1-7 and DSP8-11 are associated within cells to become GFP to emit fluorescence (Yamamoto et al, Viruses, 2020). The fact that the above cell fusion is inhibited in the presence of the obtained monoclonal antibodies indicates the ability of the antibody to be capable of inhibiting virus entry to ACE2 expressing cells of the virus. In this example, it was tested whether or not the above cell fusion was inhibited in the presence of a monoclonal antibody. The concentrations of the monoclonal antibodies were set to 0.08 μ g/mL, 0.31 μ g/mL, 1.25 μ g/mL, or 5 μ g/mL. The infection inhibiting effects of the antibodies were evaluated by showing the cell fusion inhibitory activity (%) as the vertical axis, and the antibody concentration as the transverse axis. The results were as shown in FIG. 14. As shown in FIG. 14, all 20 antibodies were able to inhibit the cell fusion between the ACE2 expressing cells and the S protein expressing cells in a concentration-dependent manner.

[0319] Next, in this example, a group of antibodies that bind to human TMPRSS2 were produced, and those having the protection effect against coronavirus infection were obtained (see FIG. 1). The advantage of this strategy is that, as demonstrated in this example, effective antibodies can be obtained regardless of virus mutation, by targeting proteins on the side of hosts such as humans, rather than proteins on the side of viruses where mutation occurs (see FIGS. 13A to D).

[0320] Monoclonal antibodies that bind to human TMPRSS2 were obtained, and the binding properties with TMPRSS2 expressing cells were confirmed under various concentration conditions. Some of the results were as shown in FIG. 2. As shown in FIG. 2, it bound to TMPRSS2 expressing cells. Also, the results that the binding affinity of various antibodies to TMPRSS2 was measured by BIAcore were as shown in Table 1.

TABLE 1

Binding affinity of anti-human TMPRSS2 antibody to TMPRSS2			
Antibody ID	ka (1/Ms)	kd (1/s)	KD (M)
752-1	3.98×10^5	0.003836	9.63×10^{-9}
1109-1	1.54×10^5	0.00178	1.16×10^{-8}
1831-15	5.63×10^5	2.36×10^{-4}	4.18×10^{-10}
1864-10	5.63×10^5	0.002894	5.14×10^{-9}
2020-12	6.54×10^5	0.001759	2.69×10^{-9}
2114-16	5.70×10^5	5.91×10^{-4}	1.04×10^{-9}
2123-13	1.22×10^5	0.001143	9.40×10^{-9}
2228-15	2.61×10^5	0.00219	8.41×10^{-9}

[0321] As shown in FIG. 2, all of the obtained monoclonal antibodies were antibodies having excellent binding affinity to TMPRSS2. Therefore, 752_1 antibody, 2228_15 antibody, 1864_10 antibody, and 2020_12 antibody were selected from the obtained monoclonal antibodies and subjected to the following analysis (see FIG. 2).

Competition Experiment

[0322] A test was performed to determine whether or not the 20 monoclonal antibodies having infection inhibiting effects obtained for the binding to human TMPRSS2 competed with each other. The results were as shown in FIGS. 3A to 3C. FIG. 3A each show the results of whether or not the antibodies compete with 752_1 antibody (FIG. 3A), 2228_15 antibody (FIG. 3B), 1831_15 antibody (FIG. 3C), 1864_10 antibody (FIG. 3D) for binding to human TMPRSS2. As shown in FIG. 3A, 752_1 antibody competed with 1109 antibody, 2123 antibody, 2179 antibody, 1477 antibody, 2156 antibody, 1185 antibody, 377 antibody, 2419 antibody, 2576 antibody, 1749 antibody, and 2355 antibody. Also, as shown in FIG. 3B, 2228_15 antibody competed with 2020_12 antibody, 1864_10 antibody, 617 antibody, 3723 antibody, and 2909 antibody. Further, as shown in FIG. 3C, 1831_15 antibody competed with 2123 antibody, and 2114 antibody. 1831_15 antibody weakly competed with 2114 antibody. 2576 antibody weakly competes with only 752_1 antibody. From these results, the obtained antibodies having infection inhibiting effects were classified into 3 competitive groups. In Table 2 below, the antibodies included in the same competitive groups were classified into bin 1 to bin 3. Note that the results shown in FIG. 3D closely resembled the results shown in FIG. 3B. 2228_15 antibody and 1864_10 antibody are thought to probably bind to nearly the same location.

TABLE 2

Results of epitope binning		
Bin 1	Bin 2	Bin 3
752__1	1864__10	2114
1109	2020__12	1831__15
2123	2228__15	2123
2179	617	
1477	3723	
2156	2909	
1185		
377		
2419		
2576		
1749		
2355		

[0323] Competitive antibodies are likely to bind to the same or overlapping epitopes. The groups of antibodies that can strongly inhibit SARS-COV-2 infection can be classified into the 3 competitive groups, suggesting that the epitopes to which these antibodies bind have significant technical meanings in relation to the infection inhibition. 2576 antibody is an antibody that can strongly inhibit SARS-COV-2 infection but does not strongly compete with either 1864_10 or 2114, and it weakly competed with 752_1. This result indicates that although 2576 antibody has a small epitope overlap with 752_1 antibody, it is still enough to inhibit SARS-COV-2 infection. Also, 2123 antibody strongly competed with 752_1.

Epitope Mapping

[0324] Peptides with a length of 20 amino acids were prepared by shifting 2 amino acids each to the extracellular region (387 amino acids) of TMPRSS2 as an antigen. Total number of obtained peptides were 185. The 20 monoclonal antibodies were each allowed to react with them, information was obtained to which peptides they showed binding properties, and based on the information, epitope mapping was performed. The results were as shown in FIG. 4.

[0325] Epitope candidates for 4 monoclonal antibodies (752_1 antibody, 2228_15 antibody, 1864_10 antibody, and 2020_12 antibody) were as shown in FIGS. 5A to 5B. 2228_15 antibody, 1864_10 antibody, and 2020_12 antibody particularly bound to the amino acid sequence set forth in SEQ ID NO: 85 (GVYGNVMVFTDWIY). The circled regions are epitopes in FIG. 6. In the leftmost panel in FIG. 6, peptide 3 is shown in the lower circle, and peptide 4 is shown in the upper circle. In the second panel from the left in FIG. 6, peptides 1 and 2 are shown in the circle. In the third panel from the left in FIG. 6, peptide 5 is shown in the circle. In the fourth panel from the left in FIG. 6, peptide 6 is shown in the center circle, peptide 7 is shown in the right circle, and peptide 8 is shown in the left circle.

Binding Properties of TMPRSS2 to Amino Acid Substitution Variants

[0326] For the purpose of identification of amino acids critical for binding between TMPRSS2 and antibodies, TMPRSS2 variants having amino acid point mutation were produced to confirm the binding with the antibodies. The TMPRSS2 variants thus produced were linked to GFP via IRES and forcibly expressed in COS7 cells. GFP-positive cells are thought to express TMPRSS2 variants, and the

binding between the cells and each antibody was analyzed by flow cytometry. The results were as shown in FIGS. 7A to 7D. As shown in FIG. 7A, 752_1 antibody lost binding properties to TMPRSS2 chimera (human/176 to 195 cat) in which the 176th to 195th amino acid sequences of the human TMPRSS2 were replaced with a corresponding region of cat TMPRSS2. Therefore, N179 of the human TMPRSS2 was shown to be important for the binding of 752_1 antibody to human TMPRSS2. Also, as shown in FIG. 7A, 1831_15 antibody lost binding properties to TMPRSS2 chimera (human/292 to 309 cat) in which the 292nd to 309th amino acid sequences of the human TMPRSS2 were replaced with a corresponding region of cat TMPRSS2. From this, the 292nd to 309th amino acid sequences of the human TMPRSS2 were shown to be important for the binding of 1831_15 antibody to human TMPRSS2. Next, as shown in FIG. 7B, 752_1 antibody lost binding properties to N179A variant of TMPRSS2. Therefore, N179 of the human TMPRSS2 was shown to be important for the binding of 752_1 antibody to human TMPRSS2. Next, as shown in FIG. 7C, 1831_15 antibody lost binding properties to W306A variant of human TMPRSS2. Therefore, W306 of the human TMPRSS2 was shown to be important for the binding of 1831_15 antibody to human TMPRSS2. Further, as shown in FIG. 7D, 2020_12 antibody, 1864_10 antibody, and 2228_15 antibody lost binding properties to TMPRSS2 chimera (human/316 to 321 hamster) in which the 316th to 321st amino acid sequences of the human TMPRSS2 were replaced with a corresponding region of hamster TMPRSS2. From this, the 316th to 321st amino acid sequences of the human TMPRSS2 were shown to be important for the binding of 2020_12 antibody, 1864_10 antibody, and 2228_15 antibody to human TMPRSS2.

[0327] A Fab fragment of the 752_1 antibody and a Fab fragment of the 2228_15 antibody described above were prepared and formed into a complex with the human TMPRSS2 to observe with a cryo-electron microscope. Electron microscope images were analyzed on a computer, and the structures of the binding sites between TMPRSS2 and respective Fab fragments as shown in FIG. 8 were obtained. As shown in FIG. 8, the 752 Fab fragment was close to N177, N179, R182, R186, and I221 of TMPRSS2 in the complex and predicted to bind to these amino acids. Also, this result coincides with the binding data of TMPRSS2 variants shown in FIG. 7A. In addition to that, as shown in FIG. 8, the 2228 Fab fragment was close to R316, F319, and F321 of TMPRSS2 in the complex and predicted to bind to these amino acids. Further, this result coincides with the binding data of TMPRSS2 variants shown in FIG. 7D. According to the above epitope binning, many antibodies were found to bind to these sites, and therefore, these epitopes are likely to be useful binding sites of the antibodies for inhibiting SARS-COV-2 infection of cells.

[0328] The human TMPRSS2 is a cytoplasmic region on the N-terminal, the 84th to 106th amino acids are a trans-membrane region, the 133rd to 148th amino acids are a LDLRA region, the 149th and 242nd amino acids are a SRCR region, and the 255th to 492nd are a serine protease region. The serine protease region contains catalytic triad of H296, D345, and S441.

Binding Properties to TMPRSS Family

[0329] The binding properties of the 4 monoclonal antibodies (752_1 antibody, 2228_15 antibody, 1864_10 anti-

body, and 2020_12 antibody) to the TMPRSS family were confirmed. The results were as shown in FIG. 9. As shown in FIG. 9, these antibodies bound specifically only to TMPRSS2.

Cross-Reactivity

[0330] Cross-reactivity to TMPRSS2 of non-human animal species was confirmed. Specifically, the cross-reactivity of ferret (SEQ ID NO: 1), hamster (SEQ ID NO: 2), mouse (SEQ ID NO: 3), feline (SEQ ID NO: 4), human (SEQ ID

NO: 5), green monkey (SEQ ID NO: 6), cynomolgus monkey (SEQ ID NO: 7), and rhesus macaque (SEQ ID NO: 8) to TMPRSS2 were confirmed. Note that the alignments of these TMPRSS2 were as shown in FIGS. 10A and 10B. Moreover, the results were as shown in FIGS. 10C and 10D. **[0331]** As a result, 1864_10 and 2020_12 exhibited the cross-reactivity to mouse TMPRSS2. Also, 752_1 exhibited the cross-reactivity to cynomolgus monkey TMPRSS2.

[0332] Characteristics of the above 20 infection-inhibiting monoclonal antibodies were determined in the same manner, and the results are shown in Table 3.

TABLE 3

Characteristics of 20 infection-inhibiting monoclonal antibodies							
mAb	Activity			Binding inhibition			
	Virus IC90	Calu-3 Inhibitor	293 FT [®] AC50 (ug/ml) [®]	between antibodies	Cross-reactivity	KD (M)	Isotype
752_1	0.3	68.4	0.04	752_1	Cynomolgus monkey	9.6E-09	IgG1, κ
1864_10	0.1	67.6	0.03	2228_15	Mouse	5.1E-09	IgG2b, κ
2020_12	0.2	66.0	0.04	2228_15	Mouse	2.7E-09	IgG2c, κ
2228_15	0.5	75.6	0.07	2228_15	—	8.4E-09	IgG2c, κ
1109_1	1.3	56.5	0.53	752_1	Cynomolgus monkey	1.2E-08	IgG2c, κ
2114_16	1.5	47.6	0.54	1831_15	—	1.0E-09	IgG2c, λ
2123_18	1.4	62.3	0.77	752_1	Mouse-Cynomolgus monkey	9.4E-09	IgG2c, κ
1831_15	—	44.8	0.26	1831_15	Cynomolgus monkey	4.2E-10	IgG2b, κ
617_8	—	52.2	0.46	2228_15	Mouse	—	IgG2c, κ
2179_7	—	14.4	0.11	752_1	Cynomolgus monkey	—	IgG2c, κ
3723_11	—	52.8	0.05	2228_15	Mouse	—	IgG2c, κ
2909_2	2.4	61.7	0.13	2228_15	—	—	IgG3, κ
1477_14	2.7	50.1	0.52	752_1	Cynomolgus monkey	—	IgG2c, κ
2156_8	2.9	61.6	0.22	752_1	Cynomolgus monkey	—	IgG2c, κ
1185_10	3.0	53.0	0.10	752_1	Cynomolgus monkey	—	IgG2c, κ
377_7	4.7	58.9	0.36	752_1	Cynomolgus monkey	—	IgG2c, κ
2419_16	—	5.6	0.26	752_1	Cynomolgus monkey	—	IgG2c, κ
2576_2	—	1.1	0.43	752_1	Cynomolgus monkey	—	IgG2c, κ
1749_14	—	43.2	0.53	752_1	Cynomolgus monkey	—	IgG2b, κ
2355_3	—	43.1	0.49	752_1	Cynomolgus monkey	—	IgG2c, κ

* The symbol “—” indicates not determined.

The blank spaces also indicate not determined.

⑦ indicates text missing or illegible when filed

Antibody Sequencing

[0333] The amino acid sequences of the heavy chain and light chain of the 4 antibodies were determined, and the heavy chain CDR1 to 3 and the light chain CDR1 to 3 were presumed (see Table 4).

TABLE 4

Amino acid sequences of each monoclonal antibody					
Antibody	Region	SEQ ID NO.	Antibody	Region	SEQ ID NO.
752_1	Heavy chain variable region	SEQ ID NO: 9	2020_12	Heavy chain variable region	SEQ ID NO: 25
	Light chain variable region	SEQ ID NO: 10		Light chain variable region	SEQ ID NO: 26
	Heavy chain CDR1	SEQ ID NO: 11		Heavy chain CDR1	SEQ ID NO: 27
	Heavy chain CDR2	SEQ ID NO: 12		Heavy chain CDR2	SEQ ID NO: 28
	Heavy chain CDR3	SEQ ID NO: 13		Heavy chain CDR3	SEQ ID NO: 29
	Light chain CDR1	SEQ ID NO: 14		Light chain CDR1	SEQ ID NO: 30
	Light chain CDR2	SEQ ID NO: 15		Light chain CDR2	SEQ ID NO: 31
	Light chain CDR3	SEQ ID NO: 16		Light chain CDR3	SEQ ID NO: 32
1864_10	Heavy chain variable region	SEQ ID NO: 17	2228_15	Heavy chain variable region	SEQ ID NO: 33
	Light chain variable region	SEQ ID NO: 18		Light chain variable region	SEQ ID NO: 34
	Heavy chain CDR1	SEQ ID NO: 19		Heavy chain CDR1	SEQ ID NO: 35
	Heavy chain CDR2	SEQ ID NO: 20		Heavy chain CDR2	SEQ ID NO: 36
	Heavy chain CDR3	SEQ ID NO: 21		Heavy chain CDR3	SEQ ID NO: 37
	Light chain CDR1	SEQ ID NO: 22		Light chain CDR1	SEQ ID NO: 38
	Light chain CDR2	SEQ ID NO: 23		Light chain CDR2	SEQ ID NO: 39
	Light chain CDR3	SEQ ID NO: 24		Light chain CDR3	SEQ ID NO: 40

TABLE 4-continued

Amino acid sequences of each monoclonal antibody					
Antibody Region	SEQ ID NO.	Antibody	Region	SEQ ID NO.	
1831_15 Heavy chain variable region	SEQ ID NO: 98	2114_16	Heavy chain variable region	SEQ ID NO: 106	
Light chain variable region	SEQ ID NO: 99		Light chain variable region	SEQ ID NO: 107	
Heavy chain CDR1	SEQ ID NO: 100		Heavy chain CDR1	SEQ ID NO: 108	
Heavy chain CDR2	SEQ ID NO: 101		Heavy chain CDR2	SEQ ID NO: 109	
Heavy chain CDR3	SEQ ID NO: 102		Heavy chain CDR3	SEQ ID NO: 110	
Light chain CDR1	SEQ ID NO: 103		Light chain CDR1	SEQ ID NO: 111	
Light chain CDR2	SEQ ID NO: 104		Light chain CDR2	SEQ ID NO: 112	
Light chain CDR3	SEQ ID NO: 105		Light chain CDR3	SEQ ID NO: 113	

Production of Human IgG4 Chimeric Antibody

[0334] Human IgG4 chimeric antibodies were produced from the 4 monoclonal antibodies (752_1 antibody, 2228_15 antibody, 1864_10 antibody, and 2020_12 antibody) and human IgG4 antibody. The binding capacity of each of the human IgG4 chimeric antibodies to TMPRSS2 was evaluated by flow cytometry using TMPRSS2 and GFP expression Daudi cells, indicating that all of the human IgG4 chimeric antibodies maintained good binding capacity to TMPRSS2 (see FIG. 11A). Further, S108P (where, 108 means the 108th amino acid as the 1st amino acid in the CH1 region) or S120P mutation was further introduced into the CH1 region of the human IgG4 chimeric antibody obtained from 752_1. From this mutation, Fab arm replacement is expected to be prevented. The binding properties between monkey TMPRSS2 and 752_1 human IgG4 chimeric antibody (S108P) were confirmed in the same manner as above. The results were as shown in FIG. 11. As shown in FIG. 11B, 752_1 human IgG4 chimeric antibody (S108P) exhibited better binding properties to human and monkey TMPRSS2 than to mouse antibody, which was its parent antibody.

Infection Inhibition Experiment

[0335] Next, the inhibition capacity of the obtained antibody to virus infection of human cells was tested. Calu-3 cells were used as the human cells and treated with serially diluted antibodies (0.1, 1 and 10 $\mu\text{g/mL}$). Virus (MOI=0.1) was added 1 hour later. After 24 hours, the cells were washed with PBS to obtain cell lysates, and total RNA was extracted from the cells to synthesize cDNA. Viral genome and GAPDH as internal control were quantified by a quantitative PCR method. The amount of viral genome was estimated as the ratio of the amount of viral genome/the amount of GAPDH mRNA (S/G ratio). Also, for the above S/G ratio, the inhibitory efficiency of the antibody was estimated by determining the ratio to the negative control containing no antibody (i.e., no inhibitory effect). IC_{90} was determined as the antibody concentration achieving 90% inhibition. The results were as shown in FIG. 12 and Table 5.

TABLE 5

Infection inhibiting effects on raw virus by TMPRSS2 antibody					
Clone	IC_{90} ($\mu\text{g/ml}$)		Clone	IC_{90} ($\mu\text{g/ml}$)	
	First run	Second run		First run	Second run
377_7	4.7	n.t.	1477_14	2.7	n.t.
617_8	—	n.t.	1749_14	—	n.t.
752_1	0.29	0.47	1831_15	—	n.t.
1109_1	1.3	2	1864_10	0.11	0.16
1185_10	3	2.5	2020_12	0.23	0.13

TABLE 5-continued

Infection inhibiting effects on raw virus by TMPRSS2 antibody					
Clone	IC_{90} ($\mu\text{g/ml}$)		Clone	IC_{90} ($\mu\text{g/ml}$)	
	First run	Second run		First run	Second run
2114_16	1.5	3.4	2355_3	—	n.t.
2123_13	1.4	1.6	2419_16	—	—
2156_8	2.9	n.t.	2576_2	—	n.t.
2179_7	—	n.t.	2909_2	2.4	2.4
2228_15	0.47	0.24	3723_11	—	n.t.

* The “n.t.” means not tested.

[0336] As shown in FIG. 12, all of the produced antibodies significantly inhibited the infection of the cells with live viruses. IC_{90} was as shown in Table 5.

Infection Inhibiting Effect on Viruses Having S Protein Variants

[0337] Pseudo-typed viruses having various S protein variants were prepared. Specifically, the following 3 variants were prepared: Pseudo-typed viruses having S proteins derived from SARS-COV-2 (WT) derived from Wuhan, SARS COV_2 S VOC202012/01 (B.1.1.7: α variant), and SARS COV_2 S 501Y.V2 (B.1.351: β variant). As test samples, commercially available anti-spike protein RBD antibody (Sino Biological, Inc., 40592-MM57, hereinafter, referred to as “MM57”), nafamostat, and the antibodies of the present example (752_1 antibody, 2228_15 antibody, 1864_10 antibody, and 2020_12 antibody) were used. MM57 antibody is an antibody capable of binding to the spike proteins to neutralize the S proteins of SARS-COV-2. Nafamostat is a serine protease inhibitor known as an active ingredient for anticoagulant drugs. Recently, nafamostat has been shown to have inhibitory effects on SARS-COV-2 infection of cells.

[0338] The results were as shown in FIG. 13A. As shown in FIG. 13A, MM57 exhibited strong inhibitory effects on WT-type virus and α variant-type virus, whereas it hardly exhibited significant inhibitory effects on β variant-type virus. Nafamostat exhibited infection inhibiting effects on all the WT-type, α variant-type, and β variant-type viruses. On the other hand, all of the antibodies of the present example exhibited infection inhibiting effects on all the WT-type, α variant-type, and β variant-type viruses.

[0339] MM57 is an antibody against S proteins, which greatly reduced or eliminated the infection inhibiting effects on β variant-type virus. On the other hand, the antibodies of the present example that bind to TMPRSS2, a protein of human cells, robustly exhibited infection inhibiting effects

even on variants. Coronavirus proteins may induce mutation. Therefore, antibodies targeting the coronavirus S proteins may reduce or eliminate the binding properties to the S protein variants, and thereby reducing or eliminating the infection inhibiting effects. On the other hand, as for antibodies against human proteins, since mutation of human proteins is unlikely to occur, their effectiveness is unlikely to depend on coronavirus mutation. The results of the present example were consistent with this logic.

[0340] Furthermore, the infection inhibiting effects on pseudo-typed viruses having SARS-COV-2 δ variant (B.1.617.2) and SARS-COV-2 κ variant (B.1.617.1) were also examined. As shown in FIGS. 13B and 13C, all of the antibodies of the present example exhibited strong infection inhibiting effects on δ variant-type, and κ variant-type viruses. In addition to that, the infection inhibiting effects on pseudo-typed viruses having the S protein of the omicron variant ($\omicron$ variant) were examined. As shown in FIG. 13D, the antibodies exhibited strong infection suppressing effects on pseudo-typed viruses having the S protein derived from the $\omicron$ variant, as with the α variant, β variant, and δ variant. As such, the effectiveness of the antibodies against all variants was demonstrated regardless of the virus variant from which it is derived. Also, none of the antibodies exhibited the infection inhibiting effect on VSV-G virus, which was used as a negative control. From this, it is clearly indicated that the virus infection inhibiting effects of the antibodies of the present example is exerted via the interaction between TMPRSS2 and S proteins.

[0341] The ADCC activities were tested for 1864_10 antibody (mIgG2b) and 2020_12 antibody (mIgG2c). ADE-substituted recombinant antibodies (1864_10ADE- and 2020_12 antibody ADE-) having mutations of L244A/E245A/P339A and L244A/E245A/P338A were each produced for these antibodies (parent antibodies). Daudi cells (Daudi; TM2-HA-k13-LD10, mouse TM2 2nd LD1, 1×10^4 cells/well) in which human TMPRSS2 or mouse TMPRSS2 was forcibly expressed were prepared. The above Daudi cells were incubated at 37° C. for 1 hour in the presence of Calcein-AM. KHYG-1 mFcRy3 transfectant (hNK cell line) at 1×10^5 /well was used as the effector cells. The ratio of the effector cells and Daudi cells was set to 10:1. As a positive control antibody, rituximab was used as an antibody having ADCC activity. As a negative control, a mouse IgG isotype control antibody was used. As test antibodies, 1864_10 antibody (mIgG2b) and 2020_12 antibody (mIgG2c), and ADE-substituted recombinant antibodies thereof were used. The lysis rate (%) of Daudi cells by ADCC activity was measured in the presence of various antibodies. The results were as shown in FIG. 15. As shown in FIG. 15, the ADE-substituted recombinant antibodies had no detectable ADCC activity.

[0342] In addition to that, the presence or absence of ADCC activity of hIgG4 chimeric antibody was confirmed. In the same manner as above, Daudi cells expressing human TMPRSS2 or cynomolgus monkey TMPRSS2 were used, and KHYG-1 hFcRy3 transfectant (hNK cell line) at 1×10^5 /well was used as the effector cells. The results were as shown in FIG. 16. As a result, antibodies other than rituximab did not exhibit significant cell lysis (%), and no significant ADCC activity was observed.

[0343] For 752hIgG4 (S120P) chimeric antibody, SARS-COV-2 infection inhibition experiment was conducted in vitro. The cells used were X293T-ACE2 expressing human

or macaque TMPRSS2. The results were as shown in FIG. 17. As shown in FIG. 17, the antibodies inhibited the infection of X293T cells with SARS-COV-2 in a concentration-dependent manner.

[0344] In an in vivo test using cynomolgus monkeys, an infection experiment of cynomolgus monkeys with SARS-COV-2 was conducted. As shown in FIG. 18A, the δ variant was administered on Day 0, and 752hIgG4 (S120P) chimeric antibody (100 mg) was administered twice on Day 0 and Day 1. Swabs were collected on Days 3, 5, and 7. The body temperature and weight changes of the cynomolgus monkeys were as shown in FIG. 18B. As shown in FIG. 18B, the body temperature of the antibody administration group was lower than the control group, indicating the therapeutic effect of the antibody administration. The number of viruses in the airway and bronchus swabs was measured by the titration method. As a result, as shown in FIG. 19A, the number of viruses in the airway and bronchus swabs reduced in the antibody administration group. Also, as shown in FIG. 19B, the number of viruses in the airway swabs reduced in the antibody administration group by RT-PCR. The number of viruses was approximately 1/18 of that in the control group.

[0345] The lung tissues were resected from the cynomolgus monkeys on Day 7, and the lung tissues were observed. The tissues were evaluated according to Table 6 below. The average value across all the visual fields was used as the visual lung injury score.

TABLE 6

Lung histopathologic diagnosis score	
Score	Tissue states
0	Healthy lung
1	Mild destruction of epithelial cells in airway and bronchus
2	Mild infiltration of inflammatory cells around periphery of bronchus
3	Moderate infiltration of inflammatory cells around alveolar walls, and resulting alveolar thickening
4	Mild alveolar injury with vascular injury of 10% or less
5	Moderate alveolar and vascular injury (11% to approximately 30%)
6	Severe alveolar injury with hyaline membrane-associated alveolar hemorrhage of 31% to approximately 50%
7	Severe alveolar injury with hyaline membrane-associated alveolar hemorrhage of 51% or more

[0346] As a result, as shown in FIG. 20, the lung histopathologic diagnosis score for the cynomolgus monkeys infected with SARS-COV-2 was improved in the antibody administration group. Also, a viral antigen test was conducted. The percentage of viral antigen positive cells was evaluated using 8G8A, which is mAb against SARS nucleocapsid protein (0: none, 1: hardly observed, 2: moderately observed, 3: highly observed). As a result, as shown in FIG. 20, the amount of viral antigen was significantly reduced in the antibody administration group. As such, the lung histopathologic diagnosis score for the cynomolgus monkeys infected with SARS-COV-2 on Day 7 of infection was improved by antibody administration, and the amount of viral antigen was significantly reduced.

[0347] In the following, 752 antibody and 2228 antibody were humanized.

TABLE 7

Amino acid sequences of humanized light chain variable regions (VL) and heavy chain variable regions (VH)				
subtype	⑦	⑦	amino acid sequence	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	
⑦	⑦	⑦	⑦	

⑦ indicates text missing or illegible when filed

[0348] When humanizing, CDRs were grafted onto 2 different frameworks, and 2 VHs and 2 VLs per 1 antibody clone were obtained (see Table 7). 4 Humanized antibody prototypes were obtained by appropriately combining the above VHs and VLs. The amino acid sequence of human IgG4 (S108P) was used as the constant region. The binding affinity of the obtained humanized antibody was measured by BIACore.

TABLE 8

KD values of humanized antibodies							
	recombinant	H chain subtype	L chain subtype	ka (1/Ms)	kd (1/s)	KD (M)	CI⑦ (RU ²)
recombinant 752-1 mIgG	mIgG	mIgG1	mlg kappa	5.42E+05	0.001409	2.60E-09	14.6
humanized 752-1 IgG4 (S108P)	prototype #1	752_hVH-1	752_hVL-1	4.12E+05	0.009321	2.26E-08	30.2
	prototype #2	752_hVH-1	752_hVL-2	1.13E+06	0.01521	1.34E-08	85.8
	prototype #3	752_hVH-2	752_hVL-1	2.22E+06	0.004691	2.11E-09	6.98
	prototype #4	752_hVH-2	752_hVL-2	2.50E+06	0.006114	2.44E-09	11.7
recombinant 2228-15 mIgG	mIgG	mIgG2C	mlg kappa	3.38E+05	0.001257	3.72E-09	3.63
humanized 2228-15 IgG4 (S108P)	prototype #1	2228_hVH-1	2228_hVL-1	5.22E+06	0.0641	1.23E-10	87.5
	prototype #2	2228_hVH-1	2228_hVL-2	7.31E+05	0.001039	1.42E-08	7.2
	prototype #3	2228_hVH-2	2228_hVL-1	8.25E+05	7.74E-04	9.38E-10	3.03
	prototype #4	2228_hVH-2	2228_hVL-2	1.44E+06	1.00E-04	2⑦E-10	2.63

⑦ indicates text missing or illegible when filed

[0349] As shown in Table 8, all the humanized antibody prototypes exhibited preferable KD values. Each of these antibodies can be used as a humanized antibody. In FIG. 21, concentration-dependent binding between these antibodies (each independently, also referred to as humanized #1 to #4 or hu #1 to #4) and Tmprss2 was confirmed. Infection inhibition experiments were conducted in vitro using the

obtained humanized IgG4 (S120P) antibody and pseudo-typed virus. The cells were human lung epithelial adenocarcinoma (Calu3 cells). As a result, as shown in FIG. 22, all the antibodies exhibited a preferable infection inhibiting effect.

[0350] The documents cited herein are incorporated herein by reference in their entirety.

Sequence Listing Contents	
SEQ ID NO: 1: Example of ferret Tmprss2	
MALNSGSPPG IGPYYENHGF QSEHIYPPRP PVAPDVYNPY PPQNYPPVP QYFPRVTQQA	60
STTVTHTQPH SSGKLCSTST KTKKSLCFAL SLGIVLVGAA VAAVLLWKFL PGCSTSEMEC	120
MSSGTCISS LWCDDGTSCHP NGEDENCAV RLYGPSFTLQ VYSSQRKAWY PVCQDDWNDS	180
YGRAACKDMG YKNNFYTTQG IPDSSGATSF MKLNISAGNI DLYKKLYHSD SCSSRMVVS	240
RCIQCGVRSR TRQSRIVGGS NASPGDWPWQ VSLHVQGVHV CGGSIITPEW IVTAAHCVEE	300
PLNSPRYWT A FAGILSQSLM FYGSRHQVEK VISHPNYNSE TKNNDIALMK LQTPLTFNDL	360
VKPVCLPNPG MMLDPAQECW ISGWGSTYEK GKTSMDMLNAA MVPLIERSKC NNKYIYNLNI	420
TPAMVCAGFL QGTVDSCQGD SGGPLVTLKN DIWWLIGDTS WSGGCAKALR PGVYGNVTVF	480
TDWIYQQMMA NS	492

-continued

Sequence Listing Contents			
SEQ ID NO: 2: Example of hamster Tmprss2			
MALNSGSPPG	IGPCYENHGY	QSEHICPPRP	PVAPNGYNLY PAQYYPSVPV QYAPRITTQA 60
STSVIHTHPK	SSGALCTSKS	KKSLCLALAL	GTVLTGAAVA AVLLWRFWDS NCSTSEMECG 120
SSGTCISSSL	WCDGVAHCPN	GEDENRCVRL	YQGSFILQVY SSQRKAWYPV CQDDWSESYG 180
RAACKDMGYK	NNFYSSQGI	DQSGATSPK	LNVSNGNVDL YKKLYHSDSC SSRMVVSLRC 240
IECGVRSVKR	QSRIVGGLNA	SPGDWPQVVS	LHVQGVHVCV GSIITPEWIV TAAHCVEEPL 300
SSPRYWTAF	GILRQSLMFY	GSRHQVEKVI	SHPNYDSKTK NNDIALMKLQ TPLAFNDLVK 360
PVCLPNPGMM	LDLDQECWIS	GWGATYEKVK	TSDLVNAAMV PLIEPSKCNK KYIYNLITP 420
AMICAGFLQG	SVDSCQDGS	GPLVTLKNGI	WWLIGDTSWG SGCAKALRPG VYGNVTFTD 480
WIYQQMRANS			490
SEQ ID NO: 3: Example of mouse Tmprss2			
MALNSGPPPG	VGPYYENHGY	QPEGLYPPRP	AVVPGAYSIVY PAPYYPPAVP QYAPRVLTQA 60
STPSSHTQPK	SPSGTVCTSK	AKKVLCTFT	LGAVLAGAVL AAVLLWKFMK NKCLVSGIEC 120
GSSGTCVSPS	HWCDGVLHCP	SGEDENRCVR	LYGPNFILQV YSSQRKSWHP VCQDDWSEGY 180
GRAACQDMGY	RNSFYSSHGV	ADDSGATSPK	RVNTSANHMD LYKKLYHSDV CSSKTVVSRL 240
CIECGVTAKM	LDLDQECWIS	SAAPGDWPQV	ASLHVQGVHV CCGSIITPEW IVTAAHCVEE 300
PLNNPRHWT	FVGILRQSEM	FYGHGYRVGK	VISHPNYDSK TKNNDIALMK LQTPFTFDDK 360
VKPVCLPNPG	LMLEPEQPCW	ISGWGATHEK	GKTSDELNAV MVPLIEPWRC NSKYVYNNLV 420
TPAMICAGYL	QGTIDSCQGD	SGGPLVTMKS	HIWWLIGDTS WSGCAKANR PGVYGNVTVF 480
TDWIYRQMRA	NG		492
SEQ ID NO: 4: Example of cat Tmprss2			
MALNSGSPPA	IGPYENHGY	QPENPYPAQP	TVVPTVYEVH PAQYYPSVPV QYAPRVLTQA 60
SNPVVCTQPK	SPSGTVCTSK	TKKALCITLT	LGTFLVGAAL AAGLLWKFMG SKCSNSGIEC 120
DSSGTCINPS	NWCDGVSHCP	GEDENRCVR	LYGPNFILQV YSSQRKSWHP VCQDDWNENY 180
GRAACRDMGY	KNNFYSSQGI	VDDSGSTSPK	KLNTSAGNVD IYKKLYHSDA CSSKAVVSLR 240
CIACGVNLMS	SRQSRIVGGE	SALPGAWPWQ	VSLHVQNVHV CCGSIITPEW IVTAAHCVEK 300
PLNNPWHWT	FAGILRQSEM	FYGAGYQVEK	VISHPNYDSK TKNNDIALMK LQKPLTFNDL 360
VKPVCLPNPG	MMLEPEQPCW	ISGWGATEEK	GKTSEVLNAA KVLLETQRC NSRYVYDNL 420
TPAMICAGFL	QGNVDSQGD	SGGPLVTSKN	NIWWLIGDTS WSGCAKAYR PGVYGNVMVF 480
TDWIYRQMRA	DG		492
SEQ ID NO: 5: Example of human Tmprss2			
MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH PAQYYPSVPV QYIPSVLTQA 60
SNPAVRTQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL AAGLLWKFMG SKCSDSGIEC 120
DSSGTCISS	NWCDGVSHCP	SGEDENRCVR	LYGPNFILQV YSSQRKSWHP VCRDDWNENY 180
ARAACRDMGY	QNNFYSSQGI	ADNSGATSPK	KLNTSAGNVD IYKKLYHSDA CSSKAVVSLR 240
CIACGVRSNL	SRQSRIVGGQ	NALPGAWPWQ	VSLHVQNIHV CCGSIITPEW IVTAAHCVEK 300
PLNSPWQWTA	FAGLLTQSSM	FYVKGHRVEK	VISHPNYDSK TKNNDIALMK LHTPLTFNEV 360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSMDLNAA MVPLIEPRRC NSKRVDGLV 420
TPAMICAGFL	QGTVDSCQGD	SGGPLVTLKN	DVWWLIGDTS WSGCAQANR PGVYGNVTVF 480
TDWIYRQMRA	DD		492
SEQ ID NO: 6: Example of green monkey Tmprss2			
MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH PAQYYPSVPV QYTPRVLTQA 60
SNPAVCRQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL AAGLLWKFMG SKCSDSGIEC 120
DSSGTCISS	NWCDGVSHCP	NGEDENRCVR	LYGPNFILQV YSSQRKSWHP VCRDDWNENY 180
ARAACRDMGY	KNSFYSSQGI	VDNSGATSPK	KLNTSAGNVD IYKKLYHSDA CSSKAVVSLR 240
CIACGVRSNL	NRQSRIVGGQ	NALLGAWPWQ	VSLHVQNIHV CCGSIITPEW IVTAAHCVEK 300
PLNSPWQWTA	FVGTLRQSSM	FYEKGHRVEK	VISHPNYDSK TKNNDIALMK LHTPLTFNEV 360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSVDLNAA MVPLIEPRRC NNKYIYDGLI 420
TPAMICAGFL	RGTVDSQGD	SGGPLVTLKN	DVWWLIGDTS WSGCAQANR PGVYGNVTVF 480
TDWIYRQMRA	DD		492
SEQ ID NO: 7: Example of cynomolgus monkey Tmprss2			
MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH PAQYYPSVPV QYTPRVLTQA 60
SNPAVCRQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL AAGLLWKFMG SKCSDSGIEC 120
DSSGTCISS	NWCDGVSHCP	NGEDENRCVR	LYGPNFILQV YSSQRKSWHP VCRDDWNENY 180
ARAACRDMGY	KNSFYSSQGI	VDNSGATSPK	KLNTSAGNVD IYKKLYHSDA CSSKAVVSLR 240
CIACGVRSNL	SRQSRIVGGQ	NALLGAWPWQ	VSLHVQNIHV CCGSIITPEW IVTAAHCVEK 300
PLNSPWQWTA	FVGTLRQSSM	FYEKGHRVEK	VISHPNYDSK TKNNDIALMK LHTPLTFNEV 360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSVDLNAA MVPLIEPRRC NNKYIYDGLI 420
TPAMICAGFL	QGTVDSCQGD	SGGPLVTLKN	DVWWLIGDTS WSGCAQANR PGVYGNVTVF 480
TDWIYRQMRA	DD		492
SEQ ID NO: 8: Example of rhesus macaque Tmprss2			
MALNAGSPPG	VGPYYENHGY	QPESLYPAPP	ATVPSVYVAY PAAYPPAVP QYTPRVLTQA 60
STPAVRTQPK	SPSGTACTAK	AKKALCITIS	LGAVLAGAAV TAVLLWKFMK NKCSVSGIEC 120
GSSGTCISPS	HWCDGVLHCP	SGEDENRCVR	LYGPNFILQV YSAQRKSWHP VCQDDWSDSY 180
GRAACQDMGY	RNSFYSSQGI	VDDSGASSPM	KLNTSAGNTD LYKKLYHSDI CASKTVVSRL 240
CIECGVAGKT	MRQSRIVGGS	SASPGDWPQV	VSLHVQGTHV CCGSIITPEW IVTAAHCVEE 300

-continued

Sequence Listing Contents		
PLNNPRYWTV FAGVLRQSFM FYGHGYRVGK VISHPSYDSK TKNNIDIALMK LQTPLTFSDK		360
VKPVCLPNPG MMLEPNQSCW ISGWGATHEK GKTSDELNAV MVPLIEPWRC NSKYVYNSLV		420
TPAMICAGYL RGGTDSQCGD SGGPLVTLKS RIWWLIGDTS WSGCAKANR PGVYGNVTVF		480
TDWIYQMRA NS		492
SEQ ID NO: 9: Heavy chain variable region of 752_1 antibody		
QVQLQQSGAE LARPGASVKL SCKASGYTFT SCGISWVKQR TGQGLEWIGE IYPRGGNTYY		60
NEKFKGKATL TADKSSSTAY MELRSLTSED SAVYFCAREN GNYDPLFAYW GQGTLLTVSA		120
SEQ ID NO: 10: Light chain variable region of 752_1 antibody		
DIQMTQSPSS LSASLGKVT ITCKASQDIN KYIAWYQHKP GKGPRLLIHY TSTLQPGIPS		60
RFSGSGSGRD YFSIYNLEP EDIATYYCLQ YYNLWTFGGG TKLEIK		106
SEQ ID NO: 11: Heavy chain CDR1 SCGIS of 752_1 antibody		
SEQ ID NO: 12: Heavy chain CDR2 EIYPRGGNTY YNEKFKG of 752_1 antibody		
SEQ ID NO: 13: Heavy chain CDR3 ENGNVDPLFA Y of 752_1 antibody		
SEQ ID NO: 14: Light chain CDR1 KASQDINKYI A of 752_1 antibody		
SEQ ID NO: 15: Light chain CDR2 YTSTLQP of 752_1 antibody		
SEQ ID NO: 16: Light chain CDR3 LQYYNLWT of 752_1 antibody		
SEQ ID NO: 17: Heavy chain variable region of 1864_10 antibody		
QVQLQQSGPA LVKPGASLKI SCEASGYTFT YYSINWVKQR PGQGLEWIGW IFPGSGSTYY		60
NEKFKGKATL TIDKSSSTAY MLLSSLTSED SAVYFCARSD YGSAYGYFDV WGTGTTVTVS		121
S		
SEQ ID NO: 18: Light chain variable region of 1864_10 antibody		
DIVLTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSYMHY QQKPGQPPKL LIYLASNLES		60
GVPARFSGSG SGTDFTLNIH PVEEEDAATY YCQHSRDLPL TFGAGTKLEL K		111
SEQ ID NO: 19: Heavy chain CDR1 YYSIN of 1864_10 antibody		
SEQ ID NO: 20: Heavy chain CDR2 WIFPGSGSTY YNEKFKG of 1864_10 antibody		
SEQ ID NO: 21: Heavy chain CDR3 SDYGSAYGYF DV of 1864_10 antibody		
SEQ ID NO: 22: Light chain CDR1 RASKSVSTSG YSYMH of 1864_10 antibody		
SEQ ID NO: 23: Light chain CDR2 LASNLES of 1864_10 antibody		
SEQ ID NO: 24: Light chain CDR3 QHSRDLPLT of 1864_10 antibody		
SEQ ID NO: 25: Heavy chain variable region of 2020_12 antibody		
QVQLQQPGAE LAKPGASVKL SCKASGYTFT SYWMHWVKQR PGQGLEWIGL IPPNSGSSNY		60
NEKFKSKATL TVDKSSRTAY QLSSLTSED SAVYICARWD YGSSYGYFDV WGTGTTVTVS		120
S		121
SEQ ID NO: 26: Light chain variable region of 2020_12 antibody		
DIVLTQSPAS LAVSLGQRAT ISCRASESVS TSGYSYMHY QQKPGQPPKL LIYLASNLES		60
GVPVRFSGSG SGTDFTLNIH PVEEEDAATY YCQHSRELPP TFGAGTKLEL K		111
SEQ ID NO: 27: Heavy chain CDR1 SYWMH of 2020_12 antibody		
SEQ ID NO: 28: Heavy chain CDR2 LIPPNSGSSN YNEKFKS of 2020_12 antibody		
SEQ ID NO: 29: Heavy chain CDR3 WDYGSSYGYF DV of 2020_12 antibody		
SEQ ID NO: 30: Light chain CDR1 RASESVSTSG YSYMH of 2020_12 antibody		
SEQ ID NO: 31: Light chain CDR2 LASNLES of 2020_12 antibody		
SEQ ID NO: 32: Light chain CDR3 QHSRELPPPT of 2020_12 antibody		
SEQ ID NO: 33: Heavy chain variable region of 2228_15 antibody		
QVQLQQSGPE LVKPGTSVKI SCKASGYTFT DYYINWVKQR PGQGLEWIGW IFPGSGSSYY		60
NAIFKKGATL TVDTSSNTAH MSLSSLTSD SAVYFCARGD FGNFGGFFTY WQGTLLTVTS		120
A		121

-continued

Sequence Listing Contents	
SEQ ID NO: 34: Light chain variable region of 2228_15 antibody DVVVTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSFMHWY QQKPGQPPKL LIYLASNLES GVPARFSGSG SGTDFILNIH PVEEEDAATY YCQHSRELPL TFGAGTKLEL K	60 111
SEQ ID NO: 35: Heavy chain CDR1 DYYIN of 2228_15 antibody	
SEQ ID NO: 36: Heavy chain CDR2 WIFPGSGSSY YNAIFKG of 2228_15 antibody	
SEQ ID NO: 37: Heavy chain CDR3 GDFGNFGGFF TY of 2228_15 antibody	
SEQ ID NO: 38: Light chain CDR1 RASKSVSTSG YSFMH of 2228_15 antibody	
SEQ ID NO: 39: Light chain CDR2 LASNLES of 2228_15 antibody	
SEQ ID NO: 40: Light chain CDR3 QHSRELPLT of 2228_15 antibody	
SEQ ID NO: 41: TMPRSS2 portion peptide AKAYRPGVYG NVMVFTDWIY	
SEQ ID NO: 42: TMPRSS2 portion peptide AYRPGVYGNV MVFTDWIYRQ	
SEQ ID NO: 43: TMPRSS2 portion peptide RPGVYGNVMV FTDWIYRQMR	
SEQ ID NO: 44: TMPRSS2 portion peptide GVGNNVMVFT DWIYRQMRAD	
SEQ ID NO: 45: TMPRSS2 portion peptide VYGNVMVFTD WIYRQMRADG	
SEQ ID NO: 46: TMPRSS2 portion peptide DSGGPLVTSK NNIWWLIGDT	
SEQ ID NO: 47: TMPRSS2 portion peptide GGPLVTSKNN IWWLIGDTSW	
SEQ ID NO: 48: TMPRSS2 portion peptide PLVTSKNNIW WLIGDTSWG	
SEQ ID NO: 49: TMPRSS2 portion peptide VTSKNNIWWL IGDTSWGSGC	
SEQ ID NO: 50: TMPRSS2 portion peptide SKNNIWWLIG DTSWGSGCAK	
SEQ ID NO: 51: TMPRSS2 portion peptide NNIWWLIGDT SWGSGCAKAY	
SEQ ID NO: 52: TMPRSS2 portion peptide VLNAAKVLLI ETQRCNSRYV	
SEQ ID NO: 53: TMPRSS2 portion peptide NAAKVLLIET QRCNSRYVYD	
SEQ ID NO: 54: TMPRSS2 portion peptide AKVLLIETQR CNSRYVDNL	
SEQ ID NO: 55: TMPRSS2 portion peptide VLLIETQRCN SRYVDNLIT	
SEQ ID NO: 56: TMPRSS2 portion peptide WTAFAGILRQ SEMFYGAGYQ	
SEQ ID NO: 57: TMPRSS2 portion peptide AFAGILRQSE MFYGAGYQVE	
SEQ ID NO: 58: TMPRSS2 portion peptide AGILRQSFMF YGAGYQVEKV	
SEQ ID NO: 59: TMPRSS2 portion peptide ILRQSEMFY GAGYQVEKVIS	
SEQ ID NO: 60: TMPRSS2 portion peptide RQSEMFYAG YQVEKVISHP	
SEQ ID NO: 61: TMPRSS2 portion peptide SEMFYGAGY QVEKVISHPNY	
SEQ ID NO: 62: TMPRSS2 portion peptide MFYGAGYQVE KVISHPNYDS	
SEQ ID NO: 63: TMPRSS2 portion peptide YGAGYQVEKV ISHPNYDSKT	
SEQ ID NO: 64: TMPRSS2 portion peptide WIVTAAHCVE KPLNNPWHWT	
SEQ ID NO: 65: TMPRSS2 portion peptide VTAACHVEKP LNNPWHWTAF	
SEQ ID NO: 66: TMPRSS2 portion peptide AAHCVEKPLN NPWHWTAFAG	
SEQ ID NO: 67: TMPRSS2 portion peptide HCVEKPLNPN WHWTAFAGIL	
SEQ ID NO: 68: TMPRSS2 portion peptide VEKPLNPNWH WTAFAGILRQ	
SEQ ID NO: 69: TMPRSS2 portion peptide KPLNNPWHWT AFAGILRQSF	

-continued

Sequence Listing Contents

SEQ ID NO: 70: TMPRSS2 portion peptide KAVVSLRCIA CGVNLN SSRQ

SEQ ID NO: 71: TMPRSS2 portion peptide VVSLRCIACG VNLN SSRQSR

SEQ ID NO: 72: TMPRSS2 portion peptide SLRCIACGVN LN SSRQSRIV

SEQ ID NO: 73: TMPRSS2 portion peptide RCIACGVNLN SSRQSRIVGG

SEQ ID NO: 74: TMPRSS2 portion peptide IACGVNLN SS RQSRIVGGES

SEQ ID NO: 75: TMPRSS2 portion peptide WNENYGRAAC RDMGYKNNFY

SEQ ID NO: 76: TMPRSS2 portion peptide RLYGPNFILQ VYSSQRKSWH

SEQ ID NO: 77: TMPRSS2 portion peptide YGPNFILQVY SSQRKSWHPV

SEQ ID NO: 78: TMPRSS2 portion peptide PNFILQVYSS QRKSWHPVCQ

SEQ ID NO: 79: TMPRSS2 portion peptide FILQVYSSQR KSWHPVCQDD

SEQ ID NO: 80: TMPRSS2 portion peptide LQVYSSQRKS WHPVCQDDWN

SEQ ID NO: 81: TMPRSS2 portion peptide VYSSQRKSWH PVCQDDWNEN

SEQ ID NO: 82: TMPRSS2 portion peptide SSQRKSWHPV CQDDWNENYG

SEQ ID NO: 83: TMPRSS2 portion peptide QRKSWHPVCQ DDWNENYGRA

SEQ ID NO: 84: TMPRSS2 portion peptide KSWHPVCQDD WNENYGRAAC

SEQ ID NO: 85: TMPRSS2 portion peptide GVGYNMVMFT DWIY

SEQ ID NO: 86: TMPRSS2 portion peptide PLVTSKNNIW WLIGDTSW

SEQ ID NO: 87: TMPRSS2 portion peptide VLLIETQRCN SRYV

SEQ ID NO: 88: TMPRSS2 portion peptide MPYGAGYQVE KV

SEQ ID NO: 89: TMPRSS2 portion peptide VTAACVCEKP LNNPWHWT

SEQ ID NO: 90: TMPRSS2 portion peptide KPLNNPWHWT AFAGILRQSE

SEQ ID NO: 91: TMPRSS2 portion peptide IACGVNLN SS RQSR

SEQ ID NO: 92: TMPRSS2 portion peptide WNENYGRAAC RDMGYKNNFY

SEQ ID NO: 93: TMPRSS2 portion peptide PNFILQVYSS QRKSWHPVCQ DD

SEQ ID NO: 94: TMPRSS2 portion peptide WSEG YGRAAC QDMGYRNSFY

SEQ ID NO: 95: TMPRSS2 portion peptide VTAACVVEEP LNNPRHWT

SEQ ID NO: 96: TMPRSS2 portion peptide RQSEME

SEQ ID NO: 97: TMPRSS2 portion peptide SQSLMF

SEQ ID NO: 98: Heavy chain variable region of 1831_15 antibody
 QVQLQQPGAE LVKPGASVKL SCKASGYSFT TYWMHWVRQR PGQGLEWIGI IHPNSGTTNY
 NEKFKSKASL TVDKSSSTAY MQLSSLTSED SAVYYCARLL WDYFDYWGQG TLTIVSS

SEQ ID NO: 99: Light chain variable region of 1831_15 antibody
 DIVMSQSPSS LAVSAGEKVT MSCKSSQSLL NSRTRKNFLA WYQQKPGQSP KLLIYWASTR
 ESGVPDRFTG SGSGTDFTLT ISSVQAEDLA VYICKQSYNL YTFGGGTKLE IK

SEQ ID NO: 100: Heavy chain CDR1 GYSFTTYW of 1831_15 antibody

SEQ ID NO: 101: Heavy chain CDR2 IHPNSGTT of 1831_15 antibody

SEQ ID NO: 102: Heavy chain CDR3 ARLLDYFDY of 1831_15 antibody

SEQ ID NO: 103: Light chain CDR1 QSLNLSRTRK NF of 1831_15 antibody

-continued

Sequence Listing Contents

SEQ ID NO: 104: Light chain CDR2 WAS of 1831_15 antibody

SEQ ID NO: 105: Light chain CDR3 KQSYNLYT of 1831_15 antibody

SEQ ID NO: 106: Heavy chain variable region of 2113_16 antibody
 EVQLQQSGPE LVKPGASVKI SCKASGYTFT DYYMNWVKRR HGKSLEWIGN INPNNGGTSYQ
 NQKFKGKATL TVDKSSSTAY MELRSLTSED SVVYYCAREG DLFPFAYWGQ GTLVTVSA

SEQ ID NO: 107: Light chain variable region of 2113_16 antibody
 QAVVTQESAL TTSPGETVTI TCRSSIGTVT TSNFANWVQE KPDHLFTGLI GASNNRAPGV
 PARFSGSLIG DKAALTITGA QTEDEAIYFC ALWFSNHWWF GGGTKLTVL

SEQ ID NO: 108: Heavy chain CDR1 GYTFTDYY of 2113_16 antibody

SEQ ID NO: 109: Heavy chain CDR2 INPNNGGT of 2113_16 antibody

SEQ ID NO: 110: Heavy chain CDR3 AREGDLFPFA Y of 2113_16 antibody

SEQ ID NO: 111: Light chain CDR1 IGTVTTSNF of 2113_16 antibody

SEQ ID NO: 112: Light chain CDR2 ASN of 2113_16 antibody

SEQ ID NO: 113: Light chain CDR3 ALWFSNHWW of 2113_16 antibody

SEQ ID NO: 114: 752 hVL-1
 DIQMTQSPSS LSASVGDRVT ITCKASQDIN KYIAWYQQKP GKAPKLLIYY TSTLQPGVPS
 RFSGSGSGTD FTFTISLQP EDIATYYCLQ YYNLWTFGGG TKVEIK

SEQ ID NO: 115: 752 hVL-1 FR1
 DIQMTQSPSS LSASVGDRVT ITC

SEQ ID NO: 116: 752 hVL-1 FR2
 WYQQKPGKAP KLLIY

SEQ ID NO: 117: 752 hVL-1 FR3
 GVPSRFSGSG SGTDFFTIS SLQPEDIATY YC

SEQ ID NO: 118: 752 hVL-1 FR4
 FGGGTKVEIK

SEQ ID NO: 119: 752 hVL-2
 DIQMTQSPSS LSASVGDRVT ITCKASQDIN KYIAWYQQKP GKAPKLLIHY TSTLQPGVPS
 RFSGSGSGTD FTFTISLQP EDIATYYCLQ YYNLWTFGGG TKVEIK

SEQ ID NO: 120: 752 hVL-2 FR1
 DIQMTQSPSS LSASVGDRVT ITC

SEQ ID NO: 121: 752 hVL-2 FR2
 WYQQKPGKAP KLLIH

SEQ ID NO: 122: 752 hVL-2 FR3
 GVPSRFSGSG SGTDFFTIS SLQPEDIATY YC

SEQ ID NO: 123: 752 hVL-2 FR4
 FGGGTKVEIK

SEQ ID NO: 124: 752 hVH-1
 QVQLVQSGAE VKKPGASVKV SCKASGYTFT SCGISWVRQA PGQGLEWMGE IYPRGGNTYY
 NEKFKGRVTM TTDSTSTAY MELRSLRSD TAVYYCAREN GNYDPLPAYW GQGTLVTVSS

SEQ ID NO: 125: 752 hVH-1 FR1
 QVQLVQSGAE VKKPGASVKV SCKASGYTFT

SEQ ID NO: 126: 752 hVH-1 FR2
 WVRQAPGQGL EWMG

SEQ ID NO: 127: 752 hVH-1 FR3
 RVTMTDTST STAYMELRSL RSDDTAVYYC AR

SEQ ID NO: 128: 752 hVH-1 FR4
 WQGTLVTVS S

-continued

Sequence Listing Contents

SEQ ID NO: 129: 752 hVH-2
QVQLQQSGTG LARPGASVKL SCKASGYTFT SCGISWVTQR AGQGLEWIGE IYPRGGNTYY
NEKFKGKATL TADKSSSSAY MELRGLTAED SAVYFCAREN GNYDPLFAYW GQGTTTLTVSS

SEQ ID NO: 130: 752 hVH-2 FR1
QVQLQQSGTG LARPGASVKL SCKASGYTFT

SEQ ID NO: 131: 752 hVH-2 FR2
WVTQRAGQGL EWIG

SEQ ID NO: 132: 752 hVH-2 FR3
KATLTADKSS SSAYMELRGL TAEDSAVYFC AR

SEQ ID NO: 133: 752 hVH-2 FR4
WGQGTTLTVS S

SEQ ID NO: 134: 2228 hVL-1
DVVMTQSPLS LPVTLGQPAS ISCRASKSVS TSGYSFMHWF QQRPGQSPRR LIYLASNLES
GVPDRFSGSG SGTDFTLKIS RVEAEDVGVI YCQHSRELPL TFGGGTKVEI K

SEQ ID NO: 135: 2228 hVL-1 FR1
DVVMTQSPLS LPVTLGQPAS ISC

SEQ ID NO: 136: 2228 hVL-1 FR2
WFQQRPGQSP RRLIY

SEQ ID NO: 137: 2228 hVL-1 FR3
GVPDRFSGSG SGTDFTLKIS RVEAEDVGVI YC

SEQ ID NO: 138: 2228 hVL-1 FR4
FGGGTKVEIK

SEQ ID NO: 139: 2228 hVL-2
DIVMTQSPDS LAVSLGERAT INCRASKSVS TSGYSEMHWY QQKPGQPPKL LIYLASNLES
GVPDRFSGSG SGTDFTLTIR PLQAEDVAVY YCQHSRELPL TFGGQTKLEI K

SEQ ID NO: 140: 2228 hVL-2 FR1
DIVMTQSPDS LAVSLGERAT INC

SEQ ID NO: 141: 2228 hVL-2 FR2
WYQQKPGQPP KLLIY

SEQ ID NO: 142: 2228 hVL-2 FR3
GVPDRFSGSG SGTDFTLTIR PLQAEDVAVY YC

SEQ ID NO: 143: 2228 hVL-2 FR4
FGGQTKLEIK

SEQ ID NO: 144: 2228 hVH-1
QVQLVQSGAE VKKPGASVKV SCKASGYTFT DYYINWVRQA PGQGLEWMGW IFPGSGSSYY
NAIFKGRVTM TRDTSISTAY MELSRRLSDD TAVYYCARGD FGNFGGFITY WGQGLTVTVS
S

SEQ ID NO: 145: 2228 hVH-1 FR1
QVQLVQSGAE VKKPGASVKV SCKASGYTFT

SEQ ID NO: 146: 2228 hVH-1 FR2
WVRQAPGQGL EWMG

SEQ ID NO: 147: 2228 hVH-1 FR3
RVTMTRDTSI STAYMELSRRL RSDDTAVYYC AR

SEQ ID NO: 148: 2228 hVH-1 FR4
WGQGLTVTVS S

SEQ ID NO: 149: 2228 hVH-2
EIQLVQSGAE VKKPGSSVKV SCKASGYTFT DYYINWVRQA PGQGLEWIGW IFPGSGSSYY
NAIFKGRATL TVDTSINTAY MELSSLRSED TAFYFCARGD FGNFGGFITY WGQGLTVTVS
S

-continued

Sequence Listing Contents

SEQ ID NO: 150: 2228 hVH-2 FR1
EIQLVQSGAE VKKPGSSVKV SCKASGYTFT

SEQ ID NO: 151: 2228 hVH-2 FR2
WMRQAPGQGL EWIG

SEQ ID NO: 152: 2228 hVH-2 FR3
RATLTVDST NTAYMELSSL RSEDATFYFC AR

SEQ ID NO: 153: 2228 hVH-2 FR4
WGQGTLLVTVS S

SEQUENCE LISTING

Sequence total quantity: 153

SEQ ID NO: 1 moltype = AA length = 492
FEATURE Location/Qualifiers
source 1..492
mol_type = protein
organism = Mesocricetus auratus

SEQUENCE: 1
MALNSGSPPG IGPYYENHGF QSEHIYPPRP PVAPDVYNPY PPQNYPPFVP QYFPRVTQA 60
STTVTHTQPH SSGKLCSTST KTKKSLCFAL SLGIVLVGAA VAAVLLWKFL PGCSTSEMEC 120
MSSGTCISSS LWCDGTSHCP NGEDENRCAV RLYGPSFTLQ VYSSQKAWY PVCQDDWNDS 180
YGRAACKDMG YKNNFYITQG IPDSSGATSF MKNLNISAGNI DLYKKLYHSD SCSSRMVVS 240
RCIQCGVRSR TRQSRIVGGS NASPGDWPWQ VSLHVQGVHV CGGSIITPEW IVTAAHCVEE 300
PLNSPRYWTA FAGILSQSLM FYGSRHQVEK VISHPNYNSE TKNNIDIALMK LQTPLTFNDL 360
VKPVCLPNPG MMLDPAQECW ISGWGSTYK GKTSDMLNAA MVPLIERSKC NNKYIYNNLI 420
TPAMVCAGFL QGTVDSCQGD SGGPLVTLKN DIWWLIGDTS WSGCAKALR PGVYGNVTVF 480
TDWIYQQMMA NS 492

SEQ ID NO: 2 moltype = AA length = 490
FEATURE Location/Qualifiers
source 1..490
mol_type = protein
organism = Mus musculus

SEQUENCE: 2
MALNSGSPPG IGPCYENHGY QSEHICPPRP PVAPNGYNLY PAQYYPSPVP QYAPRITTQA 60
STSVIHTHPK SSGALCTSKS KKSLLCLALAL GTVLTGAAVA AVLLWRFWDS NCSTSEMECG 120
SSGTCISSSL WCDGVAHCPN GEDENRCVRL YGQSFILQVY SSQKAWYVP CQDDWSESYG 180
RAACKDMGYK NNFYSSQGIP DQSGATSFMK LNVSSGNVDL YKKLYHSDSC SSRMVVSLRC 240
IECGVRSVVR QSRIVGGLNA SPGDWPWQVS LHVQGVHVCG GSIITPEWIV TAAHCVEEPL 300
SSPRYWTAPA GILRQSLMFY GSRHQVEKVI SHPNYDSKTK NNDIALMKLQ TPLAFNDLVK 360
PVCLPNPGMM LDLDQECWIS GWGATYKGGK TSDVLNAAMV PLIEPSKCNS KYIYNNLITP 420
AMICAGFLQG SVDSCQGDGSG GPLVTLKNGI WWLIGDTSWG SGCAKALRPG VYGNVTVFTD 480
WIYQQMRANS 490

SEQ ID NO: 3 moltype = AA length = 492
FEATURE Location/Qualifiers
source 1..492
mol_type = protein
organism = Felis catus

SEQUENCE: 3
MALNSGPPPG VGPYYENHGY QPEGLYPPRP AVVPGAYSIVY PAPYYPPAVP QYAPRVLTHA 60
STPSSHTQPK SPSGTVCSTK AKKVLCTIFT LGAVLAGAVL AAVLLWKFME NKCLVSGIEC 120
GSSGTCVSPS HWCAGVLHCP SGEDENRCVR LYGNPFILQV YSSQKSWHP VCQDDWSEGY 180
GRAACQDMGY RNSFYSSHGVS ADDSGATSPM RVNTSANHMD LYKKLYHSDV CSSKTVVSLR 240
CIECGVTAKM GRQSRIVGGS SAAPGDWPWQ ASLHVQGVHV CGGSIISPEW IVTAAHCVEE 300
PLNNPRHWT A FVGILRQSFY FYGHGYRVGK VISHPNYDSK TKNNIDIALMK LQTPLTFDDK 360
VKPVCLPNPG LMLEPEQPCW ISGWGATHEK GKTSDDELNAV MVPLIEPWR NSKYVYNNLV 420
TPAMICAGYL QGTIDSCQGD SGGPLVTMKS HIWWLIGDTS WSGCAKANR PGVYGNVTVF 480
TDWIYQMRAN NG 492

SEQ ID NO: 4 moltype = AA length = 492
FEATURE Location/Qualifiers
source 1..492
mol_type = protein
organism = Homo sapiens

SEQUENCE: 4
MALNSGSPPA IGPYYENHGY QPENPYPAQP TVVPTVYEVH PAQYYPSPVP QYAPRVLTHA 60

-continued

SNPVVCTQPK	SPSGTVCTSK	TKKALCITLT	LGTFLVGAAL	AAGLLWKFMG	SKCSNSGIEC	120
DSSGTCINPS	NWCDGVSHCP	GGEDENRCVR	LYGPNFILQV	YSSQRKSWHP	VCQDDWNENY	180
GRAACRDMGY	KNNFYSSQGI	VDDSGSTFSM	KLNTSAGNVD	IYKKLYHSDA	CSSKAVVSLR	240
CIACGVNLS	SRQSRIVGGE	SALPGAWPWQ	VSLHVQNVHV	CGGSIITPEW	IVTAAHCVEK	300
PLNNPWHWTA	FAGILRQSF	FYGAGYQVEK	VISHPNYDSK	TKNNDIALMK	LQKPLTFNDL	360
VKPVCLPNPG	MMLQPEQLCW	ISGWGATEEK	GKTSEVLNAA	KVLLIETQRC	NSRYVYDNL	420
TPAMICAGFL	QGNVDSQGD	SGGPLVTSKN	NIWWLIGDTS	WGSACAAYR	PGVYGNVMVF	480
TDWIYRQMRA	DG					492

SEQ ID NO: 5 moltype = AA length = 492
 FEATURE Location/Qualifiers
 REGION 1..492
 note = Chlorocebus sabaeus TMPRESS2
 source 1..492
 mol_type = protein
 organism = unidentified

SEQUENCE: 5

MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH	PAQYYPSFVP	QYIPSVLTHA	60
SNPAVRTQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL	AAGLLWKFMG	SKCSDSGIEC	120
DSSGTCISS	NWCDGVSHCP	SGEDENRCVR	LYGPNFILQV	YSSQRKSWHP	VCRDDWNENY	180
ARAACRDMGY	QNNFYSSQGI	ADNSGATFSM	KLNTSAGNVD	IYKKLYHSDA	CSSKAVVSLR	240
CIACGVRSNL	SRQSRIVGGQ	NALPGAWPWQ	VSLHVQNIHV	CGGSIITPEW	IVTAAHCVEK	300
PLNSPWQWTA	FAGILTQSSM	FYVKGHRVEK	VISHPNYDSK	TKNNDIALMK	LHTPLTFNEL	360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSMDLNAA	MVPLIEPRRC	NSKRVDGLV	420
TPAMICAGFL	QGTVDSCQGD	SGGPLVTLKN	DVWWLIGDTS	WGSACAQANR	PGVYGNVTVF	480
TDWIYRQMRA	DD					492

SEQ ID NO: 6 moltype = AA length = 492
 FEATURE Location/Qualifiers
 source 1..492
 mol_type = protein
 organism = Macaca fascicularis

SEQUENCE: 6

MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH	PAQYYPSFVP	QYTPRVLTHA	60
SNPAVCRQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL	AAGLLWKFMG	SKCSDSGIEC	120
DSSGTCISS	NWCDGVSHCP	NGEDENRCVR	LYGPNFILQV	YSSQRKSWHP	VCRDDWNENY	180
ARAACRDMGY	KNSFYSSQGI	VDNSGATFSM	KLNTSAGNVD	IYKKLYHSDA	CSSKAVVSLR	240
CIACGVRSNL	NRQSRIVGGQ	NALLGAWPWQ	VSLHVQNIHV	CGGSIITPEW	IVTAAHCVEK	300
PLNSPWQWTA	FVGTLRQSSM	FYEKGHRVEK	VISHPNYDSK	TKNNDIALMK	LHTPLTFNEV	360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSVDLNAA	MVPLIEPRRC	NNKYIYDGLI	420
TPAMICAGFL	RGTVDSCQGD	SGGPLVTLKN	DVWWLIGDTS	WGSACAQANR	PGVYGNVTVF	480
TDWIYRQMRA	DD					492

SEQ ID NO: 7 moltype = AA length = 492
 FEATURE Location/Qualifiers
 source 1..492
 mol_type = protein
 organism = Macaca mulatta

SEQUENCE: 7

MALNSGSPPG	VGPYYENHGY	QPENPYPAQP	TVAPNVYEVH	PAQYYPSFVP	QYTPRVLTHA	60
SNPAVCRQPK	SPSGTVCTSK	TKKALCVTMT	LGAVLVGAAL	AAGLLWKFMG	SKCSDSGIEC	120
DSSGTCISS	NWCDGVSHCP	NGEDENRCVR	LYGPNFILQV	YSSQRKSWHP	VCRDDWNENY	180
ARAACRDMGY	KNSFYSSQGI	VDNSGATFSM	KLNTSAGNVD	IYKKLYHSDA	CSSKAVVSLR	240
CIACGVRSNL	SRQSRIVGGQ	NALLGAWPWQ	VSLHVQNIHV	CGGSIITPEW	IVTAAHCVEK	300
PLNSPWQWTA	FVGTLRQSSM	FYEKGHRVEK	VISHPNYDSK	TKNNDIALMK	LHTPLTFNEV	360
VKPVCLPNPG	MMLEPEQHCW	ISGWGATQEK	GKTSVDLNAA	MVPLIEPRRC	NNKYIYDGLI	420
TPAMICAGFL	QGTVDSCQGD	SGGPLVTLKN	DVWWLIGDTS	WGSACAQANR	PGVYGNVTVF	480
TDWIYRQMRA	DD					492

SEQ ID NO: 8 moltype = AA length = 492
 FEATURE Location/Qualifiers
 REGION 1..492
 note = Mustela putorius furo TMPRESS2
 source 1..492
 mol_type = protein
 organism = unidentified

SEQUENCE: 8

MALNAGSPPG	VGPYYENHGY	QPESLYPAPP	ATVPSVYVAY	PAAYYPPAVP	QYTPRVLTAQ	60
STPAVRTQPK	SPSGTACTAK	AKKALCITIS	LGAVLAGAAV	TAVLLWKFME	NKCSVSGIEC	120
GSSGTCISPS	HWCDGVSHCP	SGEDENRCVR	LYGPNFILQV	YSAQRKSWHP	VCQDDWSDSY	180
GRAACQDMGY	RNSFYSSQGI	VDDSGASSFM	KLNTSAGNTD	LYKKLYHSDI	CASKTVVSLR	240
CIECGVAGKT	MRQSRIVGGS	SASPGDWPWQ	VSLHVQGTHV	CGGSIITPEW	IVTAAHCVEE	300
PLNNPRYWTV	FAGVLRQSF	FYGHGYRVGK	VISHPSYDSK	TKNNDIALMK	LQTPLTFSK	360
VKPVCLPNPG	MMLEPNQSCW	ISGWGATHEK	GKTSDELNAV	MVPLIEPWRC	NSKYVYNSLV	420
TPAMICAGYL	RGGTDSQGD	SGGPLVTLKS	RIWWLIGDTS	WGSACAKANR	PGVYGNVTVF	480
TDWIYRQMRA	NS					492

-continued

```

SEQ ID NO: 9          moltype = AA  length = 120
FEATURE              Location/Qualifiers
REGION              1..120
                    note = 752_1 VH
source              1..120
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 9
QVQLQQSGAE LARPGASVKL SCKASGYTFT SCGISWVKQR TQGLEWIGE IYPRGGNTYY 60
NEKFKGKATL TADKSSSTAY MELRSLTSED SAVYFCAREN GNYDPLPAYW GQGLVTVSA 120

SEQ ID NO: 10         moltype = AA  length = 106
FEATURE              Location/Qualifiers
REGION              1..106
                    note = 752_1 VL
source              1..106
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 10
DIQMTQSPSS LSASLGKVT ITCKASQDIN KYIAWYQHQP GKGPRLLIHY TSTLQPGIPS 60
RFGSGSGGRD YFSFIYNLEP EDIATYYCLQ YYNLWTFGGG TKLEIK 106

SEQ ID NO: 11         moltype = AA  length = 5
FEATURE              Location/Qualifiers
REGION              1..5
                    note = 752_1 HCDR1
source              1..5
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 11
SCGIS 5

SEQ ID NO: 12         moltype = AA  length = 17
FEATURE              Location/Qualifiers
REGION              1..17
                    note = EIYPRGGNTYYNEKFKG
source              1..17
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 12
EIYPRGGNTY YNEKFKG 17

SEQ ID NO: 13         moltype = AA  length = 11
FEATURE              Location/Qualifiers
REGION              1..11
                    note = 752_1 HCDR3
source              1..11
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 13
ENGNYDPLFA Y 11

SEQ ID NO: 14         moltype = AA  length = 11
FEATURE              Location/Qualifiers
REGION              1..11
                    note = 752_1 LCDR1
source              1..11
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 14
KASQDINKYI A 11

SEQ ID NO: 15         moltype = AA  length = 7
FEATURE              Location/Qualifiers
REGION              1..7
                    note = 752_1 LCDR2
source              1..7
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 15
YTSTLQP 7

SEQ ID NO: 16         moltype = AA  length = 8
FEATURE              Location/Qualifiers
REGION              1..8

```


-continued

source	note = 752_1 LCDR3 1..8 mol_type = protein organism = synthetic construct	
SEQUENCE: 16 LQYYNLWT		8
SEQ ID NO: 17 FEATURE REGION	moltype = AA length = 121 Location/Qualifiers 1..121 note = 1864_10 VH	
source	1..121 mol_type = protein organism = synthetic construct	
SEQUENCE: 17 QVQLQQSGPA LVKPGASLKI SCEASGYTFT YYSINWVKQR PGQGLEWIGW IFPGSGSTYY NEKFKGKATL TIDKSSSTAY MLLSSLTSED SAVYFCARSD YGSAYGYFDV WGTGTTVTVS S		60 120 121
SEQ ID NO: 18 FEATURE REGION	moltype = AA length = 111 Location/Qualifiers 1..111 note = 1864_10 VL	
source	1..111 mol_type = protein organism = synthetic construct	
SEQUENCE: 18 DIVLTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSYMHY QQKPGQPPKL LIYLASNLES GVPARFSGSG SGTDFTLNIH PVVEEDAATY YCQHSRDLPL TFGAGTKLEL K		60 111
SEQ ID NO: 19 FEATURE REGION	moltype = AA length = 5 Location/Qualifiers 1..5 note = 1864_10 HCDR1	
source	1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 19 YYSIN		5
SEQ ID NO: 20 FEATURE REGION	moltype = AA length = 17 Location/Qualifiers 1..17 note = 1864_10 HCDR2	
source	1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 20 WIFPGSGSTY YNEKFKG		17
SEQ ID NO: 21 FEATURE REGION	moltype = AA length = 12 Location/Qualifiers 1..12 note = 1864_10 HCDR3	
source	1..12 mol_type = protein organism = synthetic construct	
SEQUENCE: 21 SDYGSAYGYF DV		12
SEQ ID NO: 22 FEATURE REGION	moltype = AA length = 15 Location/Qualifiers 1..15 note = 1864_10 LCDR1	
source	1..15 mol_type = protein organism = synthetic construct	
SEQUENCE: 22 RASKSVSTSG YSYMH		15
SEQ ID NO: 23 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = 1864_10 LCDR2	
source	1..7 mol_type = protein	

-continued

```

SEQUENCE: 23
LASNLES
organism = synthetic construct
7

SEQ ID NO: 24
FEATURE
REGION
moltype = AA length = 9
Location/Qualifiers
1..9
note = 1864_10 LCDR3
source
1..9
mol_type = protein
organism = synthetic construct

SEQUENCE: 24
QHSRDLPLT
9

SEQ ID NO: 25
FEATURE
REGION
moltype = AA length = 121
Location/Qualifiers
1..121
note = 2020_12 VH
source
1..121
mol_type = protein
organism = synthetic construct

SEQUENCE: 25
QVQLQQPGAE LAKPGASVKL SCKASGYTFT SYWMHWVKQR PGQGLEWIGL IPPNSGSSNY 60
NEKFKSKATL TVDKSSRTAY MQLSSLTSED SAVYYCARWD YGSSYGYFDV WGTGTTVTVS 120
S 121

SEQ ID NO: 26
FEATURE
REGION
moltype = AA length = 111
Location/Qualifiers
1..111
note = 2020_12 VL
source
1..111
mol_type = protein
organism = synthetic construct

SEQUENCE: 26
DIVLTQSPAS LAVSLGQRAT ISCRASESVS TSGYSYMHY QKPGQPPKL LIYLASNLES 60
GVPVRFSGSG SGTDFTLNIH PVVEEDAATY YCQHSRELPP TFGAGTKLEL K 111

SEQ ID NO: 27
FEATURE
REGION
moltype = AA length = 5
Location/Qualifiers
1..5
note = 2020_12 HCDR1
source
1..5
mol_type = protein
organism = synthetic construct

SEQUENCE: 27
SYWMH
5

SEQ ID NO: 28
FEATURE
REGION
moltype = AA length = 17
Location/Qualifiers
1..17
note = 2020_12 HCDR2
source
1..17
mol_type = protein
organism = synthetic construct

SEQUENCE: 28
LIPPNSGSSN YNEKFKS
17

SEQ ID NO: 29
FEATURE
REGION
moltype = AA length = 12
Location/Qualifiers
1..12
note = 2020_12 HCDR3
source
1..12
mol_type = protein
organism = synthetic construct

SEQUENCE: 29
WDYGSSYGYF DV
12

SEQ ID NO: 30
FEATURE
REGION
moltype = AA length = 15
Location/Qualifiers
1..15
note = 2020_12 LCDR1
source
1..15
mol_type = protein
organism = synthetic construct

SEQUENCE: 30
RASESVSTSG YSYMH
15

```

-continued

```

SEQ ID NO: 31      moltype = AA  length = 7
FEATURE           Location/Qualifiers
REGION           1..7
                  note = 2020_12 LCDR2
source           1..7
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 31
LASNLES                                     7

SEQ ID NO: 32      moltype = AA  length = 9
FEATURE           Location/Qualifiers
REGION           1..9
                  note = 2020_12 LCDR3
source           1..9
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 32
QHSRELPTT                                   9

SEQ ID NO: 33      moltype = AA  length = 121
FEATURE           Location/Qualifiers
REGION           1..121
                  note = 2228_15 VH
source           1..121
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 33
QVQLQQSGPE LVKPGTSVKI SCKASGYTFT DYYINWVKQR PGQGLEWIGW IFPGSGSSYY 60
NAIFKGKATL TVDTSSNTAH MSLSSLTSDS SAVYFCARGD FGNFGGFPTY WQGTLVTVS 120
A                                                    121

SEQ ID NO: 34      moltype = AA  length = 111
FEATURE           Location/Qualifiers
REGION           1..111
                  note = 2228_15 VL
source           1..111
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 34
DVVVTQSPAS LAVSLGQRAT ISCRASKSVS TSGYSFMHWY QQKPGQPPKL LIYLASNLES 60
GVPARFSGSG SGTDFILNIH PVEEEDAATY YCQHSRELPL TFGAGTKLEL K          111

SEQ ID NO: 35      moltype = AA  length = 5
FEATURE           Location/Qualifiers
REGION           1..5
                  note = 2228_15 HCDR1
source           1..5
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 35
DYYIN                                       5

SEQ ID NO: 36      moltype = AA  length = 17
FEATURE           Location/Qualifiers
REGION           1..17
                  note = 2228_15 HCDR2
source           1..17
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 36
WIFPGSGSSY YNAIFKG                        17

SEQ ID NO: 37      moltype = AA  length = 12
FEATURE           Location/Qualifiers
REGION           1..12
                  note = 2228_15 HCDR3
source           1..12
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 37
GDFGNFGGFF TY                             12

SEQ ID NO: 38      moltype = AA  length = 15
FEATURE           Location/Qualifiers

```

-continued

REGION	1..15	
	note = 2228_15 LCDR1	
source	1..15	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 38		
RASKSVSTSG YSFMH		15
SEQ ID NO: 39	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = 2228_15 LCDR2	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 39		
LASNLES		7
SEQ ID NO: 40	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = 2228_15 LCDR3	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 40		
QHSRELPLT		9
SEQ ID NO: 41	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_181	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 41		
AKAYRPGVYG NVMVFTDWIY		20
SEQ ID NO: 42	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_182	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 42		
AYRPGVYGNV MVFTDWIYRQ		20
SEQ ID NO: 43	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_183	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 43		
RPGVYGNVMV FTDWIYRQMR		20
SEQ ID NO: 44	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_184	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 44		
GVYGNVMVFT DWIYRQMRAD		20
SEQ ID NO: 45	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_185	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 45		

-continued

VYGNVMVFTD WIYRQMRADG	20
SEQ ID NO: 46 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_168 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 46 DSGGPLVTSK NNIWWLIGDT	20
SEQ ID NO: 47 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_169 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 47 GGPLVTSKNN IWWLIGDTSW	20
SEQ ID NO: 48 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_170 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 48 PLVTSKNNIW WLIGDTWGS	20
SEQ ID NO: 49 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_171 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 49 VTSKNNIWWL IGDTSWGGC	20
SEQ ID NO: 50 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_172 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 50 SKNNIWWLIG DTSWGGCAK	20
SEQ ID NO: 51 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_173 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 51 NNIWWLIGDT SWGSGCAKAY	20
SEQ ID NO: 52 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_146 source 1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 52 VLNAKVLLI ETQRCNSRYV	20
SEQ ID NO: 53 moltype = AA length = 20 FEATURE Location/Qualifiers REGION 1..20 note = Peptide_147	

-continued

source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 53 NAAKVLLIET QRCNSRYVYD		20
SEQ ID NO: 54 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_148	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 54 AKVLLIETQR CNSRYVDNL		20
SEQ ID NO: 55 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_149	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 55 VLLIETQRCN SRYVYDNLIT		20
SEQ ID NO: 56 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_102	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 56 WTAFAGILRQ SFMFYGAGYQ		20
SEQ ID NO: 57 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_103	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 57 AFAGILRQSF MFYGAGYQVE		20
SEQ ID NO: 58 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_104	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 58 AGILRQSFMF YGAGYQVEKV		20
SEQ ID NO: 59 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_105	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 59 ILRQSFMPYG AGYQVEKVIS		20
SEQ ID NO: 60 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Peptide_106	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 60 RQSFMPYGAG YQVEKVISHP		20

-continued

SEQ ID NO: 61	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_107	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 61		
SFMFYGAGYQ VEKVISHPNY		20
SEQ ID NO: 62	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_108	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 62		
MPYGAGYQVE KVISHPNYDS		20
SEQ ID NO: 63	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_109	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 63		
YGAGYQVEKV ISHPNYDSKT		20
SEQ ID NO: 64	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_093	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 64		
WIVTAAHCVE KPLNNPWHWT		20
SEQ ID NO: 65	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_094	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 65		
VTAACVCVEKP LNNPWHWTAF		20
SEQ ID NO: 66	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_095	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 66		
AAHCVEKPLN NPWHWTAFAG		20
SEQ ID NO: 67	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_096	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 67		
HCVEKPLNPNP WHWTAFAGIL		20
SEQ ID NO: 68	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_097	
source	1..20	
	mol_type = protein	

-continued

SEQUENCE: 68	organism = synthetic construct	
VEKPLNNPWH WTAFAGILRQ		20
SEQ ID NO: 69	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_098	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 69		
KPLNNPWHWT AFAGILRQSF		20
SEQ ID NO: 70	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_065	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 70		
KAVVSLRCIA CGVNLN SSRQ		20
SEQ ID NO: 71	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_066	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 71		
VVSLRCIACG VNLN SSRQSR		20
SEQ ID NO: 72	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_067	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 72		
SLRCIACGVN LN SSRQSRIV		20
SEQ ID NO: 73	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_068	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 73		
RCIACGVNLN SSRQSRIVGG		20
SEQ ID NO: 74	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_069	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 74		
IACGVNLN SS RQSRIVGGES		20
SEQ ID NO: 75	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_036	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 75		
WNENYGRAAC RDMGYKNNFY		20
SEQ ID NO: 76	moltype = AA length = 20	
FEATURE	Location/Qualifiers	

-continued

REGION	1..20	
	note = Peptide_023	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 76		
RLYGPNFILQ VYSSQRKSWH		20
SEQ ID NO: 77	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_024	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 77		
YGNPNFILQVY SSQRKSWHPV		20
SEQ ID NO: 78	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_025	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 78		
PNFILQVYSS QRKSWHPVCQ		20
SEQ ID NO: 79	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_026	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 79		
FILQVYSSQR KSWHPVCQDD		20
SEQ ID NO: 80	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_027	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 80		
LQVYSSQRKS WHPVCQDDWN		20
SEQ ID NO: 81	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_028	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 81		
VYSSQRKSWH PVCQDDWNEN		20
SEQ ID NO: 82	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_029	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 82		
SSQRKSWHPV CQDDWNENYG		20
SEQ ID NO: 83	moltype = AA length = 20	
FEATURE	Location/Qualifiers	
REGION	1..20	
	note = Peptide_030	
source	1..20	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 83		

-continued

QRKSWHPVCQ DDWNNYGRA	20
SEQ ID NO: 84	moltype = AA length = 20
FEATURE	Location/Qualifiers
REGION	1..20
source	note = Peptide_031
	1..20
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 84	
KSWHPVCQDD WNNYGRAAC	20
SEQ ID NO: 85	moltype = AA length = 14
FEATURE	Location/Qualifiers
REGION	1..14
source	note = Fig6A, 1 peptide
	1..14
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 85	
GVYGNVMVFT DWIY	14
SEQ ID NO: 86	moltype = AA length = 18
FEATURE	Location/Qualifiers
REGION	1..18
source	note = Fig6A, 2 peptide
	1..18
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 86	
PLVTSKNNIW WLIGDTSW	18
SEQ ID NO: 87	moltype = AA length = 14
FEATURE	Location/Qualifiers
REGION	1..14
source	note = Fig6A, 3 peptide
	1..14
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 87	
VLLIETQRCN SRYV	14
SEQ ID NO: 88	moltype = AA length = 12
FEATURE	Location/Qualifiers
REGION	1..12
source	note = Fig6A, 4 peptide
	1..12
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 88	
MPYGAGYQVE KV	12
SEQ ID NO: 89	moltype = AA length = 18
FEATURE	Location/Qualifiers
REGION	1..18
source	note = Fig6A, 5 peptide
	1..18
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 89	
VTAHCVEKP LNNPWHWT	18
SEQ ID NO: 90	moltype = AA length = 20
FEATURE	Location/Qualifiers
REGION	1..20
source	note = Fig6A, 6 peptide
	1..20
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 90	
KPLNNPWHWT AFAGILRQSF	20
SEQ ID NO: 91	moltype = AA length = 14
FEATURE	Location/Qualifiers
REGION	1..14
	note = Fig6B, 7 peptide

-continued

source	1..14 mol_type = protein organism = synthetic construct	
SEQUENCE: 91 IACGVNLNSS RQSR		14
SEQ ID NO: 92 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Fig6B, 8 peptide	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 92 WNENYGRAAC RDMGYKNNFY		20
SEQ ID NO: 93 FEATURE REGION	moltype = AA length = 22 Location/Qualifiers 1..22 note = Fig6B, 9 peptide	
source	1..22 mol_type = protein organism = synthetic construct	
SEQUENCE: 93 PNFILQVYSS QRKSWHPVCQ DD		22
SEQ ID NO: 94 FEATURE REGION	moltype = AA length = 20 Location/Qualifiers 1..20 note = Cat TMPRSS2 176-195	
source	1..20 mol_type = protein organism = synthetic construct	
SEQUENCE: 94 WSEGYGRAAC QDMGYRNSFY		20
SEQ ID NO: 95 FEATURE REGION	moltype = AA length = 18 Location/Qualifiers 1..18 note = Cat TMPRSS2 292-306	
source	1..18 mol_type = protein organism = synthetic construct	
SEQUENCE: 95 VTAACVVEEP LNNPRHWT		18
SEQ ID NO: 96 FEATURE REGION	moltype = AA length = 6 Location/Qualifiers 1..6 note = Human TMPRSS2 316-321	
source	1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 96 RQSFMF		6
SEQ ID NO: 97 FEATURE REGION	moltype = AA length = 6 Location/Qualifiers 1..6 note = Hamster TMPRSS2 316-321	
source	1..6 mol_type = protein organism = synthetic construct	
SEQUENCE: 97 SQSLMF		6
SEQ ID NO: 98 FEATURE REGION	moltype = AA length = 117 Location/Qualifiers 1..117 note = 1831_15 VH	
source	1..117 mol_type = protein organism = synthetic construct	
SEQUENCE: 98 QVQLQQPGAE LVKPGASVKL SCKASGYSFT TYWMHWVRQR PGQGLEWIGI IHPNSGTTNY NEKFKSKASL TVDKSSSTAY MQLSSLTSED SAVYYCARLL WDYPDYWGQG TTLTVSS		60 117

-continued

```

SEQ ID NO: 99      moltype = AA  length = 112
FEATURE           Location/Qualifiers
REGION            1..112
                  note = 1831_15 VL
source            1..112
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 99
DIVMSQSPSS LAVSAGEKVT MSCKSSQSLL NSRTRKNFLA WYQOKPGQSP KLLIYWASTR 60
ESGVDPDRFTG SGSGTDFTLT ISSVQAEDLA VYYCKQSYNL YTFGGGKLE IK      112

SEQ ID NO: 100     moltype = AA  length = 8
FEATURE           Location/Qualifiers
REGION            1..8
                  note = 1831_15 HCDR1
source            1..8
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 100
GYSFTTYW                                     8

SEQ ID NO: 101     moltype = AA  length = 8
FEATURE           Location/Qualifiers
REGION            1..8
                  note = 1831_15 HCDR2
source            1..8
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 101
IHPNSGTT                                     8

SEQ ID NO: 102     moltype = AA  length = 10
FEATURE           Location/Qualifiers
REGION            1..10
                  note = 1831_15 HCDR3
source            1..10
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 102
ARLLWDYFDY                                     10

SEQ ID NO: 103     moltype = AA  length = 12
FEATURE           Location/Qualifiers
REGION            1..12
                  note = 1831_15 LCDR1
source            1..12
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 103
QSLLSRTRK NF                                 12

SEQ ID NO: 104     moltype =    length =
SEQUENCE: 104
000

SEQ ID NO: 105     moltype = AA  length = 8
FEATURE           Location/Qualifiers
REGION            1..8
                  note = 1831_15 LCDR3
source            1..8
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 105
KQSYNLYT                                     8

SEQ ID NO: 106     moltype = AA  length = 118
FEATURE           Location/Qualifiers
REGION            1..118
                  note = 2114_16 VH
source            1..118
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 106
EVQLQQSGPE LVKPGASVKI SCKASGYTFT DYYMNWVKRR HGKSLEWIGN INPNNGGTSY 60
NQKFKGKATL TVDKSSSTAY MELRSLTSED SVVYYCAREG DLFPFAYWGQ GTLVTVSA   118

```

-continued

```

SEQ ID NO: 107      moltype = AA  length = 109
FEATURE            Location/Qualifiers
REGION              1..109
                    note = 2114_16 VL
source              1..109
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 107
QAVVTQESAL TTSPGETVTL TCRSSIGT VT TSNFANWVQE KPDHLFTGLI GASNNRAPGV 60
PARFSGSLIG DKAALTITGA QTEDEAIYFC ALWFSNHWVF GGGTKLTVL 109

SEQ ID NO: 108      moltype = AA  length = 8
FEATURE            Location/Qualifiers
REGION              1..8
                    note = 2114_16 HCDR1
source              1..8
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 108
GYTFTDYY 8

SEQ ID NO: 109      moltype = AA  length = 8
FEATURE            Location/Qualifiers
REGION              1..8
                    note = 2114_16 HCDR2
source              1..8
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 109
INPNNGGT 8

SEQ ID NO: 110      moltype = AA  length = 11
FEATURE            Location/Qualifiers
REGION              1..11
                    note = 2114_16 HCDR3
source              1..11
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 110
AREGDLFPFA Y 11

SEQ ID NO: 111      moltype = AA  length = 9
FEATURE            Location/Qualifiers
REGION              1..9
                    note = 2114_16 LCDR1
source              1..9
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 111
IGTVTTSNF 9

SEQ ID NO: 112      moltype =  length =
SEQUENCE: 112
000

SEQ ID NO: 113      moltype = AA  length = 9
FEATURE            Location/Qualifiers
REGION              1..9
                    note = 2114_16 LCDR3
source              1..9
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 113
ALWFSNHWV 9

SEQ ID NO: 114      moltype = AA  length = 106
FEATURE            Location/Qualifiers
source              1..106
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 114
DIQMTQSPSS LSASVGDVRT ITCASQDIN KYIAWYQQKP GKAPKLLIYY TSTLQPGVPS 60
RFSGSGSGTD FTFTISSLPQ EDIATYYCLQ YYNLWTFGGG TKVEIK 106

SEQ ID NO: 115      moltype = AA  length = 23

```

-continued

FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 115		
DIQMTQSPSS LSASVGDRVT ITC		23
SEQ ID NO: 116	moltype = AA length = 15	
FEATURE	Location/Qualifiers	
source	1..15	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 116		
WYQQKPGKAP KLLIY		15
SEQ ID NO: 117	moltype = AA length = 32	
FEATURE	Location/Qualifiers	
source	1..32	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 117		
GVPSRFGSGG SGTDFFTIS SLQPEDIATY YC		32
SEQ ID NO: 118	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 118		
FGGGTKVEIK		10
SEQ ID NO: 119	moltype = AA length = 106	
FEATURE	Location/Qualifiers	
source	1..106	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 119		
DIQMTQSPSS LSASVGDRVT ITCKASQDIN KYIAWYQQKPKAPKLLIHY TSTLQPGVPS		60
RFGSGSGSGTD FTFTISSLQP EDIATYYCLQ YYNLWTFGGG TKVEIK		106
SEQ ID NO: 120	moltype = AA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 120		
DIQMTQSPSS LSASVGDRVT ITC		23
SEQ ID NO: 121	moltype = AA length = 15	
FEATURE	Location/Qualifiers	
source	1..15	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 121		
WYQQKPGKAP KLLIH		15
SEQ ID NO: 122	moltype = AA length = 32	
FEATURE	Location/Qualifiers	
source	1..32	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 122		
GVPSRFGSGG SGTDFFTIS SLQPEDIATY YC		32
SEQ ID NO: 123	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 123		
FGGGTKVEIK		10
SEQ ID NO: 124	moltype = AA length = 120	
FEATURE	Location/Qualifiers	
source	1..120	
	mol_type = protein	

		organism = synthetic construct	
SEQUENCE: 124			
QVQLVQSGAE VKKPGASVKV	SKKASGYTFT	SCGISWVRQA	PGQGLEWMGE IYPRGGNTYY 60
NEKFKGRVTM TTDSTSTAY	MELRSLRSD	TAVYYCAREN	GNYPDLFAYW GQGTLVTVSS 120
SEQ ID NO: 125	moltype = AA length = 30		
FEATURE	Location/Qualifiers		
source	1..30		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 125			
QVQLVQSGAE VKKPGASVKV	SKKASGYTFT		30
SEQ ID NO: 126	moltype = AA length = 14		
FEATURE	Location/Qualifiers		
source	1..14		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 126			
WVRQAPGQGL EWMG			14
SEQ ID NO: 127	moltype = AA length = 32		
FEATURE	Location/Qualifiers		
source	1..32		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 127			
RVTMTTDTST STAYMELRSL	RSDDTAVYYC	AR	32
SEQ ID NO: 128	moltype = AA length = 11		
FEATURE	Location/Qualifiers		
source	1..11		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 128			
WGQGLTVTS S			11
SEQ ID NO: 129	moltype = AA length = 120		
FEATURE	Location/Qualifiers		
source	1..120		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 129			
QVQLQQSGTG LARPGASVKL	SKKASGYTFT	SCGISWVTQR	AGQGLEWIGE IYPRGGNTYY 60
NEKFKGKATL TADKSSSAY	MELRGLTAED	SAVYFCAREN	GNYPDLFAYW GQGTTLVSS 120
SEQ ID NO: 130	moltype = AA length = 30		
FEATURE	Location/Qualifiers		
source	1..30		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 130			
QVQLQQSGTG LARPGASVKL	SKKASGYTFT		30
SEQ ID NO: 131	moltype = AA length = 14		
FEATURE	Location/Qualifiers		
source	1..14		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 131			
WVTQRAGQGL EWIG			14
SEQ ID NO: 132	moltype = AA length = 32		
FEATURE	Location/Qualifiers		
source	1..32		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 132			
KATLTADKSS SSAYMELRGL	TAEDSAVYFC	AR	32
SEQ ID NO: 133	moltype = AA length = 11		
FEATURE	Location/Qualifiers		
source	1..11		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 133			

-continued

WGQGTTLTVS S	11
SEQ ID NO: 134 moltype = AA length = 111	
FEATURE	Location/Qualifiers
source	1..111
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 134	
DVVMTQSPLS LPVTLGQPAS ISCRASKSVS TSGYSFMHWF QQRPGQSPRR LIYLASNLES	60
GVPDRFSGSG SGTDFTLKIS RVEAEDVGVI YCQHSRELPL TFGGGTKVEI K	111
SEQ ID NO: 135 moltype = AA length = 23	
FEATURE	Location/Qualifiers
source	1..23
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 135	
DVVMTQSPLS LPVTLGQPAS ISC	23
SEQ ID NO: 136 moltype = AA length = 15	
FEATURE	Location/Qualifiers
source	1..15
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 136	
WFQQRPGQSP RRLIY	15
SEQ ID NO: 137 moltype = AA length = 32	
FEATURE	Location/Qualifiers
source	1..32
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 137	
GVPDRFSGSG SGTDFTLKIS RVEAEDVGVI YC	32
SEQ ID NO: 138 moltype = AA length = 10	
FEATURE	Location/Qualifiers
source	1..10
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 138	
FGGGTKVEIK	10
SEQ ID NO: 139 moltype = AA length = 111	
FEATURE	Location/Qualifiers
source	1..111
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 139	
DIVMTQSPDS LAVSLGERAT INCRASKSVS TSGYSFMHWY QQKPGQPPKL LIYLASNLES	60
GVPDRFSGSG SGTDFTLTIR PLQAEDVAVY YCQHSRELPL TFGGGTKLEI K	111
SEQ ID NO: 140 moltype = AA length = 23	
FEATURE	Location/Qualifiers
source	1..23
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 140	
DIVMTQSPDS LAVSLGERAT INC	23
SEQ ID NO: 141 moltype = AA length = 15	
FEATURE	Location/Qualifiers
source	1..15
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 141	
WYQQKPGQPP KLLIY	15
SEQ ID NO: 142 moltype = AA length = 32	
FEATURE	Location/Qualifiers
source	1..32
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 142	
GVPDRFSGSG SGTDFTLTIR PLQAEDVAVY YC	32

-continued

SEQ ID NO: 143	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 143		
FGGTKLEIK		10
SEQ ID NO: 144	moltype = AA length = 121	
FEATURE	Location/Qualifiers	
source	1..121	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 144		
QVQLVQSGAE VKKPGASVKV SCKASGYTFT DYYINWVRQA PGQGLEWMGW IFPGSGSSYY		60
NAIFKGRVTM TRDTSISTAY MELSLRSDD TAVYYCARGD FGNFGGFPTY WQGTLVTVS		120
S		121
SEQ ID NO: 145	moltype = AA length = 30	
FEATURE	Location/Qualifiers	
source	1..30	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 145		
QVQLVQSGAE VKKPGASVKV SCKASGYTFT		30
SEQ ID NO: 146	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 146		
WVRQAPGQGL EWMG		14
SEQ ID NO: 147	moltype = AA length = 32	
FEATURE	Location/Qualifiers	
source	1..32	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 147		
RVTMTRDTSI STAYMELSLR RSDDTAVYYC AR		32
SEQ ID NO: 148	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 148		
WQGTLVTVS S		11
SEQ ID NO: 149	moltype = AA length = 121	
FEATURE	Location/Qualifiers	
source	1..121	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 149		
EIQLVQSGAE VKKPGSSVKV SCKASGYTFT DYYINWVRQA PGQGLEWIGW IFPGSGSSYY		60
NAIFKGRATL TVDTSTNTAY MELSSLRSED TAFYFCARGD FGNFGGFPTY WQGTLVTVS		120
S		121
SEQ ID NO: 150	moltype = AA length = 30	
FEATURE	Location/Qualifiers	
source	1..30	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 150		
EIQLVQSGAE VKKPGSSVKV SCKASGYTFT		30
SEQ ID NO: 151	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 151		
WVRQAPGQGL EWIG		14

-continued

SEQ ID NO: 152	moltype = AA	length = 32	
FEATURE	Location/Qualifiers		
source	1..32		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 152			
RATLTVDTST NTAYMELSSL	RSEDTAFYFC	AR	32
SEQ ID NO: 153	moltype = AA	length = 11	
FEATURE	Location/Qualifiers		
source	1..11		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 153			
WGQGTLLTVS	S		11

1. An antibody or an antigen-binding fragment thereof that binds to an extracellular domain of TMPRSS2, capable of inhibiting coronavirus infection of cells.

2. The antibody or an antigen-binding fragment thereof according to claim 1, capable of binding to (i) one or more extracellular domains of TMPRSS2 selected from the group consisting of mice, green monkeys, cats, hamsters, and macaque monkeys, and (ii) an extracellular domain of human TMPRSS2.

3. The antibody or an antigen-binding fragment thereof according to claim 1, wherein a binding affinity (KD) to human TMPRSS2 is 10^{-8} M or less.

4. The antibody or an antigen-binding fragment thereof according to claim 1, wherein IC_{50} for inhibition of the coronavirus infection of human cells is 10 μ g/mL or less.

5. The antibody or an antigen-binding fragment thereof according to claim 1, wherein the antibody binds to a peptide having the amino acid sequence consisting of the amino acid sequence selected from the group consisting of the amino acid sequence set forth in SEQ ID NO: 75 (WNE-NYGRAACRDMGYKNNFY), and the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT).

6. The antibody or an antigen-binding fragment thereof according to claim 1, wherein the antibody binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and wherein:

- (α) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 75 (WNE-NYGRAACRDMGYKNNFY) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 94 WSEGYGRAACQDMGYRNSFY).
- (β) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 89 (VTAAHCVEKPLNNPWHWT) of the human TMPRSS2 is substituted for a corresponding sequence of a cat (i.e., the amino acid sequence set forth in SEQ ID NO: 95: VTAAHCVEEPLNNPRHWT); or
- (γ) the antibody does not significantly bind to chimeric TMPRSS2 having the amino acid sequence wherein the amino acid sequence set forth in SEQ ID NO: 96 (RQSFMF) of the human TMPRSS2 is substituted for a corresponding sequence of a hamster (i.e., the amino acid sequence set forth in SEQ ID NO: 97: SQSLMF).

7. The antibody or an antigen-binding fragment thereof according to claim 1, wherein the antibody binds to human TMPRSS2 having the amino acid sequence set forth in SEQ ID NO: 5, and wherein:

(A) the antibody does not significantly bind to a human TMPRSS2 variant having a point mutation at N179A; or

(B) the antibody does not significantly bind to a human TMPRSS2 variant having a point mutation at W306A.

8. The antibody or an antigen-binding fragment thereof according to claim 1, being:

(1) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 11,

HCDR2 having the amino acid sequence set forth in SEQ ID NO: 12, and

HCDR3 having the amino acid sequence set forth in SEQ ID NO: 13,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 14,

LCDR2 having the amino acid sequence set forth in SEQ ID NO: 15, and

LCDR3 having the amino acid sequence set forth in SEQ ID NO: 16.

(2) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 19,

HCDR2 having the amino acid sequence set forth in SEQ ID NO: 20, and

HCDR3 having the amino acid sequence set forth in SEQ ID NO: 21,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 22,

LCDR2 having the amino acid sequence set forth in SEQ ID NO: 23, and

LCDR3 having the amino acid sequence set forth in SEQ ID NO: 24.

(3) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 27,

HCDR2 having the amino acid sequence set forth in SEQ ID NO: 28, and

HCDR3 having the amino acid sequence set forth in SEQ ID NO: 29,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 30, LCDR2 having the amino acid sequence set forth in SEQ ID NO: 31, and LCDR3 having the amino acid sequence set forth in SEQ ID NO: 32.

(4) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 35, HCDR2 having the amino acid sequence set forth in SEQ ID NO: 36, and HCDR3 having the amino acid sequence set forth in SEQ ID NO: 37,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 38, LCDR2 having the amino acid sequence set forth in SEQ ID NO: 39, and LCDR3 having the amino acid sequence set forth in SEQ ID NO: 40.

(5) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 100, HCDR2 having the amino acid sequence set forth in SEQ ID NO: 101, and HCDR3 having the amino acid sequence set forth in SEQ ID NO: 102,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 103, LCDR2 having the amino acid sequence set forth in SEQ ID NO: 104, and LCDR3 having the amino acid sequence set forth in SEQ ID NO: 105.

(6) an antibody or an antigen-binding fragment thereof, comprising a heavy chain variable region that has HCDR1 having the amino acid sequence set forth in SEQ ID NO: 108, HCDR2 having the amino acid sequence set forth in SEQ ID NO: 109, and HCDR3 having the amino acid sequence set forth in SEQ ID NO: 110,

a light chain variable region that has LCDR1 having the amino acid sequence set forth in SEQ ID NO: 111, LCDR2 having the amino acid sequence set forth in SEQ ID NO: 112, and LCDR3 having the amino acid sequence set forth in SEQ ID NO: 113.

(7) an antibody or an antigen-binding fragment thereof, competing with the antibody according to any of (1) to (6) above for binding with TMPRSS2; or

(8) an antibody or an antigen-binding fragment thereof, comprising the amino acid sequence having one or several amino acid mutations in the CDR sequence of the antibody according to any of (1) to (6) above; or

(9) a human chimeric antibody or a humanized antibody of the antibody according to any of (1) to (8) above.

9. A composition comprising the antibody or an antigen-binding fragment thereof according to claim **1**.

10. The composition according to claim **9**, for use in preventing a subject in need thereof from being infected with coronavirus.

11. A pharmaceutical composition according to claim **9**, for use in administering to a subject with coronavirus.

12. The composition according to claim **9**, for use in treating coronavirus infection in a subject infected with a beta coronavirus.

* * * * *