

(54)	ENHANCED CELL-FREE BACTERIOPHAGE SYNTHESIS BY GENETIC MODULATION OF BACTERIAL TRANSCRIPTION/TRANSLATION MACHINERY (TXTL) MACHINERY		Publication Classification	
(71)	Applicant:	The Administrators of the Tulane Educational Fund, New Orleans, LA (US)	(51)	Int. Cl.
(72)	Inventor:	Rani Brooks, New Orleans, LA (US)		C12N 7/00 (2006.01)
(73)	Assignee:	The Administrators of the Tulane Educational Fund, New Orleans, LA (US)		C12N 1/06 (2006.01)
				C12N 1/20 (2006.01)
				C12N 9/22 (2006.01)
				C12N 15/11 (2006.01)
				C12N 15/70 (2006.01)
				C12R 1/19 (2006.01)
			(52)	U.S. Cl.
				CPC C12N 7/00 (2013.01); C12N 1/066 (2013.01); C12N 1/20 (2013.01); C12N 9/226 (2013.01); C12N 15/111 (2013.01); C12N 15/70 (2013.01); C12N 2310/20 (2017.05); C12N 2795/00051 (2013.01); C12R 2001/19 (2021.05)
(21)	Appl. No.:	18/858,248	(57)	ABSTRACT
(22)	PCT Filed:	Apr. 19, 2023		The present disclosure relates to compositions including or obtained from genetically modified bacterial host cells (e.g., <i>E. coli</i>) and methods for using the same for cell-free bacteriophage synthesis (CFBS). In particular, the present technology relates to genetically modified <i>E. coli</i> that over-
(86)	PCT No.:	PCT/US2023/019145		
	§ 371 (c)(1),			
	(2) Date:	Oct. 18, 2024		


Related U.S. Application Data

(60) Provisional application No. 63/332,901, filed on Apr. 20, 2022.

ABSTRACT

The present disclosure relates to compositions including or obtained from genetically modified bacterial host cells (e.g., *E. coli*) and methods for using the same for cell-free bacteriophage synthesis (CFBS). In particular, the present technology relates to genetically modified *E. coli* that over-express one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or repress RecC subunit exonuclease RecBCD, and methods for using the same to obtain improved CFBS yields.

Specification includes a Sequence Listing.



in vivo bacteriophage replication




FIG. 1A

cell-free bacteriophage synthesis (CFBS)




FIG. 1B




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 4A




FIG. 4B




FIG. 5A




FIG. 5B




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7A




FIG. 7B




FIG. 8




FIG. 9




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B

[illegible]

[illegible]

[illegible]

gaagt (SEQ ID NO: 1)

[illegible]

FIG. 12 (Cont.)

pelB- ClyF gBlock	tttaagaggagataacataTGAAA TACTTATTGCCGACCGCCGCGCTGGTTTGGCTTTAGCAGCCACGCCCTGCAATGGCCATGGCTCTGCCCAAACGG GCAA GCGGACTGCTAAGCAGGTGGTTGATTGGGCTATCAATCTTATCGGCTCTGGCGTAGATGGATGGGACTATGGCGCCAAATGTTGGGATTGG CGAATTACATTTTAAACCGTTATTGGAAATTTAAGACTCCTGGGAATGCTCGTGATATGSCCTGGTATCGCTATCCGGAGGGAATTAAGTTTTCGGTA ACACTTCIGATTTCGTCCCAAAACCGGCSATATGCGGTATGGACGGTGGGAACATAATTGGAA TACTTGGGGGCATACTGGCATTTAGTTGGAC CGTCGACAAAATCCATTTTACTCGGTTGACCAAAACTGGAACAATTCAAATCGTATGTAGGGTCGCCAGCGGCCAAAATTAACATTCCTATTTCGG TGTGACTCATTTGGTCCGTCAGATACAAGCGGAGCCGAGCCCTACACCACCTAGTCCGCCAGCTATCGCGAACAACAGGCACAAATGACTGTCA CAGTGGACGCCCTTAA TGTGCGCGTGCGCCGAACACTAGCGGAGAAATCGTTGCCGTCTACAAGCGCGGGAGTCCCTCGATTACGACACAGTCAT TATCGACGTAAATGGGTACGTGTGGTGTCTGATACATTGGAGGTAGCGGAAAGCGCAATTACGTGGCTACCGGAGCGCACAAAGGACGCGGAAGCGCTT GGAACGCGTGGGGAACITTTAAATAAgtagacggctgcta (SEQ ID NO: 2)
pJL1_ FWD	GTCGACCGGCTGCTAACAAAG (SEQ ID NO: 3)
pJL1_ REV	ATGTATATCTCCTTCTTAAAGTTAAACAAAATTAATTCTAGAG (SEQ ID NO: 4)
Colon y_PC R_F	Tatagtcctgtgggttgcgc (SEQ ID NO: 5)
Colon y_PC R_R	Ctgcctgtgaacaagictgg (SEQ ID NO: 6)
pelB- ClyF T7 primer F	Accgcctgagcaataataatagacacacatagggag (SEQ ID NO: 7)
pelB- ClyF T7 primer R	Aaggggtatgclaggcaaaaaacccctcaagac (SEQ ID NO: 8)
T7_Fr ag A_F	TCTCAGTGTACGGACCTAAAGTTCCGCC (SEQ ID NO: 9)

FIG. 12 (Cont.)

T7_Fr ag A_R	ATTACGGCATGACAGTAGACAACCTTTCCG (SEQ ID NO: 10)
T7_Fr ag B_F	TGCAGCAATACCGGAAGGTGTCTACIGT (SEQ ID NO: 11)
T7_Fr ag B_R	ATATGTCCTCATAGATGTGCCATATGTGG (SEQ ID NO: 12)
T7_Fr ag C_F	ACTTGTGACTCCACATAGGCACATCTATGA (SEQ ID NO: 13)
T7_Fr ag C_R	GAATAACCTGAGGGTCAATACCCCTGCTTGT (SEQ ID NO: 14)
T7_Fr ag D_F	GACATGATGGACAAAGCAGGGTATTGACCCT (SEQ ID NO: 15)
T7_Fr ag D_R	AGGGACACAGAGAGACACTCAAGGTAACAC (SEQ ID NO: 16)
T7_Fr ag E_F	Acttgactccacalaggcacatcica (SEQ ID NO: 17)
T7_Fr ag E_R	Tagtgagtcgtattatttgctcagcgggtggc (SEQ ID NO: 18)
T7_Fr ag_F_... F	Tgaggggtttttgctcagcataacccctggg (SEQ ID NO: 19)
T7_Fr ag_F_... R	Gaataacctgagggtcaataccctgtgt (SEQ ID NO: 20)

FIG. 12 (Cont.)

[illegible]




FIG. 13




FIG. 14A




FIG. 14B




FIG. 14C

hcat FWD	CGCTCGGCTATTTACATACT	SEQ ID NO: 24
hcat REV	GGTTTCTGGCGTTAAACCAATC	SEQ ID NO: 25
cysG FWD	CCGTCTGGTTTCTGACGATATT	SEQ ID NO: 26
cysG REV	CGCAGCAGGATCTGGTTAAT	SEQ ID NO: 27
idnT FWD	GTGCGCCTCTTCTTTGAATTT	SEQ ID NO: 28
idnT REV	TCGATGGTGGCTCCATTAC	SEQ ID NO: 29
trxA FWD	AATATGGCATCCGTGGTATCC	SEQ ID NO: 30
trxA REV	GTTAGCGTCGAGGAACCTCTT	SEQ ID NO: 31
FndCas12a FWD	CCTGTTTCGATGAGAGGAATCTG	SEQ ID NO: 32
FndCas12a REV	CCTTAGCTGGGTGAGTGATTT	SEQ ID NO: 33

FIG. 15

FIG. 16

Name	Sequence
pCRJ 001	GGCTCACCTTCGGGTGGGCTTCCTGCTTATACAGAGACCTCAGCCGCTACAGGGCGCGCTCCCATTCGCCATTTCAGGCTGCGCAACTGTGGGAA GGGCGATCGGTGCGGGCTCTTCGCTATTACGCCAGCTGGCGAAAGGGGATGCTGCGAAGGCGATTAGTTGGTAACGCCAGGGTTTCCCAGT CAAGACGTGTAAACAGCGGCCAGTAGCGCGCTAATACGACTCACTATAGGCGCAATTGGGTACCGGGCCCCCTCGAGGTCCTCCAGCTTT GTTCCCTTAGTGAGGGTTAATTGCGGCTTGGGTAATCATGGTCATAGCTGTTCCTGTGTAAGGCTCAATCTACAAACAGTAGAAATTACGGTGT TTACTAGTATTAACCTAGGACTGAGCTAGCTAGCTCAAGGATATATCCGCTCCCTGAGAGATACCTTAACAGGGAAGTAGAGGGCCGCGCAACCGCTTTTCCATAG GAAATGGCTTACGAACGGGGCGGAGATTTCCTGGAAGATGCCAGGAAGATACCTTAACAGGGAAGTAGAGGGCCGCGCAACCGCTTTTCCATAG GCTCGCCCCCTGACAAGCATCACGAAATCTGACGCTCAATCAGTGGTGGCGAAACCCGACAGGACTATAAGATACCAGGGCTTCCGCCCTGGCG GCTCCCTCGTGGCTCTCCCTGCTCCCTGGCTTACCGGTGTCTCCGCTGTATGGCGGCTTGTCTCATTCACGCTGACACTCAGTTCG GGTAGGCAGTTCGCTCCAAGCTGGACTGTATGCACGAACCCCGCTCAGTCCGACCGCTGGCTTATCCGGTAACCTATCGCTTGAAGTCAACCC GGAAAGACATGCAAAAGCACCACTGGCAGCGACCTGGTAATTGATTAGAGGAGTAGTCTTGAAGTCATGCGCCGCTTAAGGCTAAACTGAAAGG ACAAAGTTTGGTGACTGGCTCTCCAGCCAGTTACCTCGGTCAAGAGTTGGTAGCTCAGAGAACCTTCGAAACCCGCTTCAAGGGCGGTTT TCGTTTTCAGAGCAAGAGATTACGCGCAGAACCAACGATCTCAAGAGATCATCTTAAATCAGATAAAATATTTCTAGATTTCAGTGCAATTTATCTC TTCAAATGTAGCACTGAGTCAAGCCCATACGATATAAGTTGTACTAGTCTGGATTCTCACCATAAAMMGCCCGCGGCAACGAGCGTTCT GAACAAATCCAGATGGAGTTCTGAGGTCAATTACTGGATCTATCAACAGGAGTCCAAAGCGAGCTCGATATCAAAATACGCCCGCCCTGCCACTCATCGC AGTACTGTGTAATTCATTAAAGCATTTGCGGACATGGAAGCATCAAAACGGCATGATGAACCTGAATCGCCAGCGCATCAGCACCTTGTGCGCTT GCGTATAATTGCCCATTGTTGAAACGGGGCGGCAAGATTGTCCAATTGCGCCAGCTTTAAATCAAAACTGGTGAACCTCACCCAGGGATTGGCTG AGACGAAACATAATCTCAATAAACCTTTAGGGAATAGGCGAGTTTTCACCGTAACACGCCACATCTTGGCAATATATGTAGAACTGCGGGA ATCGTCTGGTATTCACCTCAGAGCGATGAAACGTTTCAGTTTGTCTCATGGAACCGGTGTAACAAGGGTGAACACTATCCCATATCACCACTCACCC GTCTTTCAATTGCCATACGAAATTCGGGATGAGCATTCATCAGCGGGGCAAGAAATGTGAATAAGGCGCGGATAAACTTGTGCTTATTTCTTTACGGTC TTTAAACAGGCGTAATATCCAGCTGAACGGCTCGTTATAGGTACATGAGCAACTGACTGAATGCCCTCAAAATGTCTTACGATGCCATTTGGGATA TATCAACGGTGGTATATCCAGTGATTTTCTCCATTTTAGCTTCTTAGCTCCTGAAATCTOGATAACTCAAAATAACGCCCGGTAGTGATCTTATTT CAATTATGGTGAAAGTTGGAACCTCTTAAGTCCCGATCAACGCTCATTTTCGCCAGATACGAGTCCACCAATTCAGCAAAATTCAGCAATCATCAC GTTTCATCTTCCCTGGTTGCCAATGGCCCAATTTTCTGTGATACGAGAAGGTCGCGAATTCAGGCGCTTTTAGACTGGTGGTAAATGAAATCTTTTT AAGAAGGAGATATACATTAAAGAGGAGAAAGGATCTATAGCACTACCAAGGATTCGTCAACAAGTATTCAGTGAAGACACTGCGGTTTCGAGCTG ATCCACAGGGCAACACACTCGAACAATCAAGGCCCGAGGCTGATTCCTGACCATGAGAACCGGCAACACTATAAGAAACCAAGCAATCA TTGATAAATACCACCATCTTTATCGAGGAAATTCAGCTCCGTGTGCTATCAGTGAGGATCTGCTGCAGAAATTTCTCAGAGTGTACTTCAAGCTGAA GAAGCGACGATGACAACCTGCAGAAGGACTTCAAGTCCGCCAAGGACACCATCAAGAAAGTACAGCTCTGAGTAAATTCAGAGATTAGCGAGTACAT CAAGGACTCCGAAAGTTTAAATACTGTTCAACCAAGATCTGATCGATGCTAAGAAAGCCAGGAGTCGACCTGATCCTGTGGCTGAACAGCTCAA GGACAAATGGGATTGAAGTGTCAAGGCTAACTCCGATATCAGTATGATTTGACGAGGCACTGGAATATCATCAAGAGCTTCAAGGATGGACCACTACTT TAAAGGCTTCCACGAGAACCGCAAGACGTTACTCCAGCAACGACATCTCTACCTCCATCATCTACCGAATCTGTCGATGACAACTCGCCAAAGTTCCCT GGAGAACAAAGGCCAAATATGAATCTCTGAAGGACAAAGCTCCGAGGCAATTAATACGAACAGATCAAGAAAGATCTGGCTGAGGAACCTGACATTCGA TATCGACTATAAGACTAGCGAGGTGAACCAAGAGGGTCTTTCCTTGGACGAGGTGTTTGAATCGCCCAATTTCAACAAATTACCTGAAACCAAGTCCGGCAT

FIG. 16 (Cont.)

TACTAAATTCAAATACCATCATTTGGCGGGAAGTTTGTGAACGGGGGAGAAATACCAAGCGCAAGGGAATTAACGAATACATCAATCTGTATAGCCAGCAGAT
CAACGACAAAACCTCTGAAGAAATACAAGATGTCTGTGCTGTTTCAAAACAGATCCTGAGTGATACCGAGTCCAAAGCTTTTGTGCTATTGATAAACTGGAAGAT
GACTCAGACGTTGTCATACCATGCAGAGCTTTTATGAGCAGATGCGCGCTTCAAGACAGTGGAGGAAAAATCTATTAAAGGAACTCTGAGTCTGCTG
TTGATGAACCTGAAAGCCAGAAAGCTGGACCTGAGTAAATCTACTTCAAAACGATAAGAGTCTGACAGACCTGTACAGCAGGTGTTTGTGATGACTAT
TCCGTGATTGGACCGCGTCCGTGGAGTACATTACACAGCAGATCGCTCCAAAGAACTGGATAATCCCTTAAGAAAGAGCAAGAACTGATCGCTAA
GAAAACCGAGAAGGCAAAATATCTGAGTCTGGAACAAATTAAGCTGGCACTGGAGGAGTTCAACAAGCACAGGATATTGACAAACAGTCCCGCTTTGA
GGAAATCCTGGCCAACTTCGCAGCCATCCCATGATTTTGTGATGAGATCGCCAGAACAAAGACAACTCTGGCTCAGATCAGTATTAAATACCAAGAACCA
GGCAAGAAAGACCTGCTGCAGGCTTCAGCAGAAGATGACGTGAAGCCATCAAGGATCTGCTGGACCAACCAATCTGCTGCACAAGCTGAAAA
TCTTCCATATTAGTCAGTCAGAGGATAAGGCTAATACTCTGGATAAAGACGAACACTCTACCTGGTTCGAGGAATGTTACTTCGAGCTGGCAAACAT
TGTCCTCTGTATAACAAGATTAGGAACCTACATCACACAGAGCCCTTACTGTACGAGAAGTTTAACTGAACCTCGAAAAATAGTACCTGGCCAAACGG
GTGGGATAAGAACAAAGGAGCCTGACAACACAGCTATCCTGTTTCATCAAGGATGACAAGTACTATCTGGGAGTGATGAATAAGAAAAACAATAAGATCTT
CGATGACAAAGCCATTAAGGAGAACAAAGGGAAGGATACAAAGAAAATCGTGATTAAGCTGCTGCCCGCGCAATAAGATGCTGCCTAAGGTGTTCT
TCAGCGCCAAAGGATACAAATCTACAACCCATCCGAGGACATCCTGCGGATTAGAAATCACTCAACACATACCTAAAGACGGGAGCCCCAGAGGGAT
ATGAGAAATTTGAGTTCACATCGAGGATTGCGAGGAAGTTTATTGACTCTACAAGCAGAGCATCTCCAAACACCTGAAATGGAAGGATTTTGGCTTCGG
GTTTCCGACACACAGAGATATACTCTATCGACGAGTTTACCGGAGGTGGAATAACAAAGATTTTTCAGCATACAGCAAGGCGAGCAACCACTGCAACACTG
ATCGACAGCGTGGTCAATCAGGGAAGCTGTAACCTGTTCCAGATCTATACAAAGATTTTTCAGCATACAGCAAGGCGAGCAACCACTGCAACACTG
TACTGGAAGGCGCTGTTGATGAGAGGAATCTGCAGGACGTGGTCTATAAACTGAACGAGAGGCGCAACTGTTTACCGGAAGCAGTCTATTCTTAA
GAAATCACTCACCAGCTAAGGAGGCCATCGCTAACAAAGAACAGGACAACTCTAAGAAAGAGAGCGGTTCGAAATCGATCTGATTAAGGACAAGC
GGTCAACCGAAGATAAGTCTTTTCCATTGTCCAAATCACCAATTAACCTCAAGTCAAGCGCGCTAACCAAGTTCAACGACGAGATCAATCTGCTGCTGAA
GGAAAGCAACGATGTGCACATCTTGAGCATTTGCTCGAGGAGAGCGGCACTCTGGCCCTACTATACCTGGTGGATGGCAAGGGAATATCATTAAGC
AGGATACATTCACATCATTTGGCAATGACCGGATGAACCAACTACCAAGATAAACTGGCTGCAATCGAGAAGGATAGAGACTCAGCTAGGAAGGACT
GGAAGAAATCAACAACATTAAGGAGATGAAGGAAGGCTATCTGAGCCAGGTGGTCCATGAGATTGCAAGCTGGTCAATCGAATACAATGCCATTGTGG
TGTTGAGGATCTGAACCTCGGCTTTAAGAGGGGCGCTTTAAGGTGGAATAACAGGCTATCAGAAAGCTGGAGAAATGCTGATCGAAAAAGCTGAATT
ACCTGGTGTTAAGATAACGAGTTCGACAAGACCGGAGCGGTCCTGAGAGCCTACCAAGCTGACAGCTCCCTTTGAAACTTTCAAGAAATGGGAAAAAC
AGACAGGCATCATCTACTATGTGCCAGCGGATTCATCTCCAAGATCTGCCCGGTGACCGGCTTTGTCAACCAAGCTGACCCCTAAATATGAGTCAGTGA
GCAAGTCCAGGAATTTTCAGCAAGTTCGATAAGATCTGTTATAATCTGGACAAGGGTACTTCGAGTTTCCCTCGATTACAAGAACTTCGGCGACAA
GGCCGCTAAGGGGAAATGGACCAATGGCTCTCTGGATCTCGGCTGATCAACCTTCGAAATTCGAAATTCGGAATAAAACCACTTGGGACACIAGGGAGGTGTA
CCCAACCAAGGAGCTGGAAAAAGCTGCTGAAAGACTACTCTATCGAGTATGGACATGGCGAATGCGAATGCAATCAAGCGAGCCATCTGTGGCGAGAGTGAAGA
AATTTTGGCCAAAGCTGACCTCAGTCTGATTAACATCTCGAGATCGGGAACCTCAAGACCGGGACAGAACTGGACTATCTGATTAGCCCCGTGGCTG
ATGTCAACGGAAACTTCTTCGACAGCAGACAGGCAACCAAAATATGGCTCAGGATGCAGACGCCAACCGGGCTACCCACATCGGGCTGAAGGAGCT
GATGCTGCTGGGCGGATCAAGAACAAATCAGGAGGGGAAGAGCTGAACCTGGTCAATTAAGAACGAGGAATACTTCGAGTTTGCCAGAAATAGAAATA
ACAAAGGCGCGCGGCCACGAAAAAGGCGCGCCAGGCAAAAAAGAAAAAGGATCTCTACCCATACGATGTTCCAGATTACGCTTATCCCTACGAGGCTG

FIG. 17

[illegible]

FIG. 18

Name	Sequence	Description	SEQ ID NO
g001 top	agatcgaaagcggtatccggtgaaa	guide RNA Goldengate insert top strand targeting trxA promoter	SEQ ID NO: 46
g001 bottom	atactttcacgggatacgtcttgcg	guide RNA Goldengate insert bottom strand targeting trxA promoter	SEQ ID NO: 47
g002 top	agattatagagcgataaaatttat	guide RNA Goldengate insert top strand targeting trxA CDS	SEQ ID NO: 48
g002 bottom	atacataattttatgcgtcatata	guide RNA Goldengate insert bottom strand targeting trxA CDS	SEQ ID NO: 49
g003 top	agattggttcgcctgtatccttt	guide RNA Goldengate insert top strand targeting nusG promoter	SEQ ID NO: 50
g003 bottom	atacaaaaggataccaggcgaaacca	guide RNA Goldengate insert bottom strand targeting nusG promoter	SEQ ID NO: 51
g004 top	agataagggcggtagcaacgtcgcg	guide RNA Goldengate insert top strand targeting nusG CDS	SEQ ID NO: 52
g004 bottom	ataccgacgttgcctacgcgcctt	guide RNA Goldengate insert bottom strand targeting nusG CDS	SEQ ID NO: 53
g005 top	agatctgcacaaagcagcatcaaggg	guide RNA Goldengate insert top strand targeting pgk promoter	SEQ ID NO: 54
g005 bottom	ataccctgtgctgctttttgcag	guide RNA Goldengate insert bottom strand targeting pgk promoter	SEQ ID NO: 55
g006 top	agatagatgacgcgtcgtgatctt	guide RNA Goldengate insert top strand targeting pgk CDS	SEQ ID NO: 56
g006 bottom	atacaagatccagatcgggtcatct	guide RNA Goldengate insert bottom strand targeting pgk CDS	SEQ ID NO: 57
g007 top	agattcttcttcgtctaaagatt	guide RNA Goldengate insert top strand targeting subH(ssyA) promoter	SEQ ID NO: 58
g007 bottom	atacaatctttagacgaagaaga	guide RNA Goldengate insert bottom strand targeting subH(ssyA) promoter	SEQ ID NO: 59
g008 top	agatgtagcaaaogttagataaagc	guide RNA Goldengate insert top strand targeting subH(ssyA) CDS	SEQ ID NO: 60
g008 bottom	atacgtttatctacgtttgtcac	guide RNA Goldengate insert bottom strand targeting subH(ssyA) CDS	SEQ ID NO: 61
g009 top	agattgcacagactgtaagtaacccg	guide RNA Goldengate insert top strand targeting eno promoter	SEQ ID NO: 62
g009 bottom	ataccgggtactacagictgcca	guide RNA Goldengate insert bottom strand targeting eno promoter	SEQ ID NO: 63
g010 top	agataagcgcgaagtacatctggag	guide RNA Goldengate insert top strand targeting eno CDS	SEQ ID NO: 64
g010 bottom	ataccaccagatgtaacttcgcctt	guide RNA Goldengate insert bottom strand targeting eno CDS	SEQ ID NO: 65
g011 top	agattcggaaactggatagatggt	guide RNA Goldengate insert top strand targeting mukB promoter	SEQ ID NO: 66
g011 bottom	atacaccaatcatatccagttccga	guide RNA Goldengate insert bottom strand targeting mukB promoter	SEQ ID NO: 67
g012 top	agatgctcacitgacgcgtgattaac	guide RNA Goldengate insert top strand targeting mukB CDS	SEQ ID NO: 68
g012 bottom	atacgttaatcacagcgtcagtgagc	guide RNA Goldengate insert bottom strand targeting mukB CDS	SEQ ID NO: 69
g013 top	agataaaatgcacaaacattttata	guide RNA Goldengate insert top strand targeting hemL promoter	SEQ ID NO: 70
g013 bottom	atactataaaagttttgtcattt	guide RNA Goldengate insert bottom strand targeting hemL promoter	SEQ ID NO: 71
g014 top	agattcagcgcagcgcgcgcgcgtga	guide RNA Goldengate insert top strand targeting hemL CDS	SEQ ID NO: 72
g014 bottom	atactcagctcgcgcgcgcgcgcgtg	guide RNA Goldengate insert bottom strand targeting hemL CDS	SEQ ID NO: 73

FIG. 18 (Cont.)

g015 top	agatttaagggtatcatcctgttc	guide RNA Goldengate insert top strand targeting cyaR (yveE) promoter	SEQ ID NO: 74
g015 bottom	atacgaacggatgataaccttaa	guide RNA Goldengate insert bottom strand targeting cyaR (yveE) promoter	SEQ ID NO: 75
g016 top	egatgcctgtgtaactcctttac	guide RNA Goldengate insert top strand targeting cyaR (yveE) CDS	SEQ ID NO: 76
g016 bottom	atacgaaggagattacacagc	guide RNA Goldengate insert bottom strand targeting cyaR (yveE) CDS	SEQ ID NO: 77
g017 top	egatcgtatatactacacagcata	guide RNA Goldengate insert top strand targeting lexA promoter	SEQ ID NO: 78
g017 bottom	atactatgctgtgagtataacag	guide RNA Goldengate insert bottom strand targeting lexA promoter	SEQ ID NO: 79
g018 top	agatactctatcctgtgatcacatc	guide RNA Goldengate insert top strand targeting lexA CDS	SEQ ID NO: 80
g018 bottom	atacgaigtgatcacggatgagat	guide RNA Goldengate insert bottom strand targeting lexA CDS	SEQ ID NO: 81
g019 top	egatgctttaaatacgcgagcac	guide RNA Goldengate insert top strand targeting infC promoter	SEQ ID NO: 82
g019 bottom	atacgtgctcgcggattttaaagc	guide RNA Goldengate insert bottom strand targeting infC promoter	SEQ ID NO: 83
g020 top	egatttcgtataatgattacggca	guide RNA Goldengate insert top strand targeting infC CDS	SEQ ID NO: 84
g020 bottom	atactgcggaatcattataaga	guide RNA Goldengate insert bottom strand targeting infC CDS	SEQ ID NO: 85
g021 top	egatgcgtagtctcccggggc	guide RNA Goldengate insert top strand targeting recC promoter	SEQ ID NO: 86
g021 bottom	atacgcctccgggtgaactagcg	guide RNA Goldengate insert bottom strand targeting recC promoter	SEQ ID NO: 87
g022 top	agatttgcgaacgcgaaaggctg	guide RNA Goldengate insert top strand targeting recC CDS	SEQ ID NO: 88
g022 bottom	ataccagcgtctcgttcgacaa	guide RNA Goldengate insert bottom strand targeting recC CDS	SEQ ID NO: 89
g023 top	agataaggcatagttaccatgog	guide RNA Goldengate insert top strand targeting dgt promoter	SEQ ID NO: 90
g023 bottom	atacgcgaiggtaaactatgctt	guide RNA Goldengate insert bottom strand targeting dgt promoter	SEQ ID NO: 91
g024 top	agatcactggagcgcaatgcggcc	guide RNA Goldengate insert top strand targeting dgt CDS	SEQ ID NO: 92
g024 bottom	atacggcgcatcgtcctccagt	guide RNA Goldengate insert bottom strand targeting dgt CDS	SEQ ID NO: 93
g025 top	agatgactggtatcaggcggaiga	guide RNA Goldengate insert top strand targeting udk promoter	SEQ ID NO: 94
g025 bottom	atactcatcgcgcgtatataccagtc	guide RNA Goldengate insert bottom strand targeting udk promoter	SEQ ID NO: 95
g026 top	agattcgtgaattacgtgagcaag	guide RNA Goldengate insert top strand targeting udk CDS	SEQ ID NO: 96
g026 bottom	atacttgcctacgttaattcacga	guide RNA Goldengate insert bottom strand targeting udk CDS	SEQ ID NO: 97
g027 top	agattgatttaataagatagatc	guide RNA Goldengate insert top strand targeting rna promoter	SEQ ID NO: 98
g027 bottom	atacgaatactatctattaaatca	guide RNA Goldengate insert bottom strand targeting rna promoter	SEQ ID NO: 99
g028 top	agatatcgtctatgtctcgtggccic	guide RNA Goldengate insert top strand targeting rna CDS	SEQ ID NO: 100
g028 bottom	atacgaaggccagacatagcgat	guide RNA Goldengate insert bottom strand targeting rna CDS	SEQ ID NO: 101
g029 top	agattgcctacccgtagagtcctcgg	guide RNA Goldengate insert top strand targeting gp2 CDS	SEQ ID NO: 102

FIG. 18 (Cont.)

g029 bottom	atactgggclacogtagagtcclcg	guide RNA Goldengate insert bottom strand targeting gp2 CDS	SEQ ID NO: 103
g030 top	agatgctctggcctagaggacaag	guide RNA Goldengate insert top strand targeting gp2 CDS	SEQ ID NO: 104
g030 bottom	atacctgtctctagggccagagc	guide RNA Goldengate insert bottom strand targeting gp2 CDS	SEQ ID NO: 105
g031 top	agattccctatcaagcattcigac	guide RNA Goldengate insert top strand targeting oxys	SEQ ID NO: 106
g031 bottom	atacGTCAGAAATGCTTGATAGGGA	guide RNA Goldengate insert bottom strand targeting oxys	SEQ ID NO: 107
g032 top	agatacccttgaagtcacigcccg	guide RNA Goldengate insert top strand targeting oxys	SEQ ID NO: 108
g032 bottom	atacggggcagtgacitcaagggt	guide RNA Goldengate insert bottom strand targeting oxys	SEQ ID NO: 109
g033 top	agatcagtggaataatgagccgt	guide RNA Goldengate insert top strand targeting me promoter	SEQ ID NO: 110
g033 bottom	atacacggccctcattatccacig	guide RNA Goldengate insert bottom strand targeting me promoter	SEQ ID NO: 111
g034 top	agatttgattacggcgcigaacgt	guide RNA Goldengate insert top strand targeting me CDS	SEQ ID NO: 112
g034 bottom	atacacgttcagcgccgtaatacaa	guide RNA Goldengate insert bottom strand targeting me CDS	SEQ ID NO: 113
g035 top	agatcgtctglaaatccclacaa	guide RNA Goldengate insert top strand targeting trxB promoter	SEQ ID NO: 114
g035 bottom	atactgtaggggaattacagacg	guide RNA Goldengate insert bottom strand targeting trxB promoter	SEQ ID NO: 115
g036 top	agataaactgagatcattttgat	guide RNA Goldengate insert top strand targeting trxB CDS	SEQ ID NO: 116
g036 bottom	atacatbaaaaatgatctcagitt	guide RNA Goldengate insert bottom strand targeting trxB CDS	SEQ ID NO: 117
g037 top	agatGCACACTGGGCAATACGAGG	guide RNA Goldengate insert top strand targeting NT-control	SEQ ID NO: 118
g037 bottom	atacCCCTCGTATTGCCCCAGTGTGC	guide RNA Goldengate insert bottom strand targeting NT-control	SEQ ID NO: 119

**ENHANCED CELL-FREE BACTERIOPHAGE
SYNTHESIS BY GENETIC MODULATION OF
BACTERIAL
TRANSCRIPTION/TRANSLATION
MACHINERY (TXTL) MACHINERY**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application claims the benefit of and priority to U.S. Provisional Patent Application No. 63/332,901, filed Apr. 20, 2022, the contents of which are incorporated herein by reference in its entirety.

GOVERNMENT SUPPORT

[0002] This invention was made with government support under W81XWH-20-1-0071 awarded by the Other Agency, and 1144646 awarded by the National Science Foundation. The government has certain rights in the invention.

TECHNICAL FIELD

[0003] The present disclosure relates to compositions including or obtained from genetically modified bacterial host cells (e.g., *E. coli*) and methods for using the same for cell-free bacteriophage synthesis (CFBS). In particular, the present technology relates to genetically modified *E. coli* that overexpress one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or repress RecC subunit of the RecBCD exonuclease, and methods for using the same to obtain improved CFBS yields.

BACKGROUND

[0004] The following description of the background of the present technology is provided simply as an aid in understanding the present technology and is not admitted to describe or constitute prior art to the present technology.

[0005] Multidrug-resistant (MDR) bacteria pose one of the greatest emerging health threats. These pathogens possess intrinsic and acquired mechanisms of resistance to most common antibiotics, leaving few treatment options for the most at-risk patients. The Centers for Disease Control reports 2.8 million MDR infections per year leading to 35,000 deaths with numbers are on the rise². Phage therapy, a resurgent anti-bacterial treatment, has the potential to cure bacterial infections resistant to small molecule antibiotics. Unlike small molecule antibiotics, phage do not harm beneficial bacterial flora that protect against future infections.

[0006] Despite a promising future, phage therapy is encumbered by logistical challenges in the manufacturing process. Once a suitable phage has been identified for a particular application, it needs to be propagated using a working host, which ideally is well characterized, non-pathogenic, and free of genomically encoded prophage to prevent potential contamination of phage preparations with lysogenic phage⁶. Such a host is often not immediately available. Next, crude phage preparations must be purified to remove endotoxin, a component of outer membranes of Gram-negative bacteria that induces septic shock⁷. While several organic solvent extraction or affinity column purification methods exist to remove endotoxin, these approaches have widely varying efficiencies depending on the phage undergoing purification and typically result in significant loss of titer⁸. Additionally, low shelf-stability can result in a

significant loss of titer prior to clinical administration due to storage conditions, shipping conditions, and phage-dependent variability⁹⁻¹¹.

[0007] Accordingly, there is an urgent need for phage production methods that achieve enhanced yields over conventional cell-free bacteriophage synthesis (CFBS) methods.

SUMMARY OF THE PRESENT TECHNOLOGY

[0008] In one aspect, the present disclosure provides genetically modified bacterial host cells (e.g., *E. coli*) that overexpress one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or comprise a vector that includes a nucleic acid sequence encoding a CRISPR enzyme and a vector that includes a nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC.

[0009] In some embodiments, the genetically modified bacterial host cells comprise a non-endogenous expression vector that comprises a translation initiation factor IF-3 (infC) nucleic acid sequence of SEQ ID NO: 39, an OxyS nucleic acid sequence of SEQ ID NO: 41, a CyaR nucleic acid sequence of SEQ ID NO: 42, or any combination thereof.

[0010] Additionally or alternatively, in some embodiments, the non-endogenous expression vector comprising the infC nucleic acid sequence and the non-endogenous expression vector comprising the OxyS nucleic acid sequence and/or the CyaR nucleic acid sequence are the same or distinct. In other embodiments, the non-endogenous expression vector comprising the OxyS nucleic acid sequence and the non-endogenous expression vector comprising the CyaR nucleic acid sequence are the same or distinct.

[0011] In any of the preceding embodiments of the genetically modified bacterial host cells disclosed herein, the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is a plasmid, a cosmid, a bacmid, a bacterial artificial chromosome (BAC), or a viral vector. In certain embodiments, the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is operably linked to an expression control sequence. The expression control sequence may be an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter.

[0012] Additionally or alternatively, in some embodiments, the genetically modified bacterial host cells of the present technology comprise a vector that includes a nucleic acid sequence encoding a CRISPR enzyme and a vector that includes a nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC. In some embodiments, the vector including the nucleic acid sequence encoding the CRISPR enzyme and the vector including the nucleic acid sequence encoding the crRNA are

the same or distinct. In certain embodiments, the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter.

[0013] Additionally or alternatively, in certain embodiments of the genetically modified bacterial host cells of the present technology, the CRISPR enzyme is a nuclease-deficient *Francisella novicida* Cas12a (dFnCas12a) and/or the nucleic acid sequence encoding the crRNA is selected from the group consisting of SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88 and SEQ ID NO: 89.

[0014] Additionally or alternatively, in some embodiments, the genetically modified bacterial host cells of the present technology comprise a deletion or mutation in (a) a promoter region located upstream of a transcription start site (TSS) of *recC*, or (b) a *recC* coding region located downstream of a transcription start site (TSS) of *recC*, wherein the mutation or deletion represses transcription of *recC*. The deletion may be located upstream or downstream of the transcription start site (TSS) of *recC*. In some embodiments, the deletion is about 10-about 100 base pairs (bps) in length. In other embodiments, the mutation is a nonsense mutation, a frameshift mutation, or a missense mutation.

[0015] In any of the foregoing embodiments described herein, the genetically modified bacterial host cells are *E. coli* cells.

[0016] In one aspect, the present disclosure provides a cell lysate obtained from any and all embodiments of the genetically modified bacterial host cells disclosed herein. The cell lysates of the present technology comprise an effective amount of transcription/translation (TXTL) machinery that is configured to synthesize bacteriophage under cell-free conditions. The cell lysates may be prepared using one or more of French-press cell lysis, sonication, runoff reactions, or lysate dialysis.

[0017] In another aspect, the present disclosure provides an in vitro method for synthesizing bacteriophage virions comprising contacting a bacteriophage genome with any and all embodiments of the cell lysates disclosed herein and an energy buffer in vitro to obtain a reaction mixture, and incubating the reaction mixture under conditions to produce viable phage virions, wherein the energy buffer comprises canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes. In some embodiments, the reaction mixture is incubated at 16° C. to 42° C. for about 1-24 hours. The bacteriophage genome may be derived from a naturally occurring bacteriophage, or a genetically engineered bacteriophage. In some embodiments, the bacteriophage genome is obtained from a bacteriophage that specifically infects the naturally occurring counterpart of the genetically modified bacterial host cells disclosed herein, such as *E. coli*. Additionally or alternatively, in some embodiments, the bacteriophage genome is obtained from a bacteriophage selected from among T7 phage, T7-like phage, Lambda, K1E, T3, T5, T4, or PhiX174.

[0018] Additionally or alternatively, in some embodiments, the energy buffer further comprises one or more of a polyethylene glycol (PEG) polymer (such as PEG-8000 or PEG-6000), deoxynucleotide triphosphates (dNTPs), and stabilizers. In certain embodiments, the energy buffer further comprises one or more of Mg-glutamate, K-glutamate, tRNA, cAMP, folinic acid, spermidine, 3-PGA, HEPES,

Nicotinamide adenine dinucleotide (NAD), coenzyme A (CoA), DTT, and maltodextrin.

[0019] In any of the preceding embodiments of the methods disclosed herein, the reaction mixture further comprises a vector including a nucleic acid sequence encoding a reporter gene, wherein the nucleic acid sequence encoding the reporter gene is operably linked to a promoter that is responsive to a RNA polymerase (RNAP) encoded by the bacteriophage genome. In certain embodiments, the RNAP is selected from among SP6 RNA Polymerase, T3 RNA Polymerase and T7 RNA polymerase. The reporter gene may encode a bioluminescent protein, a fluorescent protein, or a chemiluminescent protein. Examples of bioluminescent proteins include, but are not limited to, Aequorin, firefly luciferase, *Renilla* luciferase, red luciferase, luxAB, or nanoluciferase. Examples of chemiluminescent proteins include, but are not limited to, β -galactosidase, horseradish peroxidase (HRP), or alkaline phosphatase. Examples of fluorescent proteins include, but are not limited to, sfGFP, TagBFP, Azurite, EBFP2, mKalamal, Sirius, Sapphire, T-Sapphire, ECFP, Cerulean, SCFP3A, mTurquoise, monomeric Midoriishi-Cyan, TagCFP, mTFP1, EGFP, Emerald, Superfolder GFP, Monomeric Azami Green, TagGFP2, mUKG, mWasabi, EYFP, Citrine, Venus, SYFP2, TagYFP, Monomeric Kusabira-Orange, mKOx, mKO2, mOrange, mOrange2, mRaspberry, mCherry, dsRed, mStrawberry, mTangerine, tdTomato, TagRFP, TagRFP-T, mApple, mRuby, mPlum, HcRed-Tandem, mKate2, mNeptune, NirFP, TagRFP657, IFP1.4, iRFP, mKeima Red, LSS-mKate1, LSS-mKate2, PA-GFP, PAmCherry1, PATagRFP, Kaede (green), Kaede (red), KikGR1 (green), KikGR1 (red), PS-CFP2, PS-CFP2, mEos2 (green), mEos2 (red), PSmOrange, or Dronpa.

[0020] Additionally or alternatively, in some embodiments of the methods disclosed herein, the phage virions show increased efficiency of plating (EOP) relative to phage virions obtained with cell-free bacteriophage synthesis (CFBS) in a control lysate obtained from a wild-type bacterial cell, wherein the wild-type bacterial cell and the genetically modified bacterial host cell are the same species.

[0021] In one aspect, the present disclosure provides a kit comprising one or more expression vectors comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO: 39, SEQ ID NO: 41 and SEQ ID NO: 42, and instructions for using the same to prepare donor bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In certain embodiments, the at least one nucleic acid sequence is operably linked to an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter. Additionally or alternatively, in some embodiments, the kits further comprise vials containing naturally-occurring or non-natural bacterial host cells that can be transformed with the one or more expression vectors comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO: 39, SEQ ID NO: 41 and SEQ ID NO: 42. In some embodiments, the bacterial host cells are *E. coli*.

[0022] In another aspect, the present disclosure provides a kit comprising one or more expression vectors comprising at least one nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of *recC* or a *recC* coding region located downstream of a transcription start site (TSS) of *recC*, and instructions for using the same to prepare donor

bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In some embodiments, the one or more expression vectors further comprise a nucleic acid sequence encoding a CRISPR enzyme. Examples of CRISPR enzymes include, but are not limited to, nuclease-deficient Cas12a from *F. novicida* (dFnCas12a), Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9, Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, and Csf4. Additionally or alternatively, in some embodiments, the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the at least one nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter. In some embodiments, the nucleic acid sequence encoding the crRNA is selected from the group consisting of SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88 and SEQ ID NO: 89. Additionally or alternatively, in some embodiments, the kits further comprise vials containing naturally-occurring or non-natural bacterial host cells that can be transformed with the one or more expression vectors comprising the nucleic acid sequence encoding the CRISPR enzyme and/or the at least one nucleic acid sequence encoding the crRNA. In some embodiments, the bacterial host cells are *E. coli*.

[0023] In yet another aspect, the present disclosure provides a kit comprising any and all embodiments of the genetically modified bacterial host cell described herein, and instructions for using the same to prepare donor bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In some embodiments, the genetically modified bacterial host cells are *E. coli*.

[0024] In any and all embodiments of the kits disclosed herein, the kits further comprise one or more of polyethylene glycol (PEG) polymers (e.g., PEG-8000, PEG-6000), deoxynucleotide triphosphates (dNTPs), stabilizers, and an energy buffer comprising canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] FIGS. 1A-1B: Gene expression background in TXTL source influences CFBS yields. FIG. 1A: The number of progeny resulting from bacteriophage T7 infection of host *E. coli* BL21 is influenced by the genetic background of the host at the time of infection. Knockdown (KD) or overexpression (OX) of certain genes may positive, negative, or neutral impact on T7 progeny yield relative to wild-type BL21 (WT). FIG. 1B: Cell-free bacteriophage synthesis (CFBS) can be modulated by modifying the genetic background of the source of transcription/translation (TXTL) machinery derived from cell lysates.

[0026] FIGS. 2A-2C: Inducible CRISPRi of *trxA* knockdown mRNA and T7 efficiency of plating (EOP). Induced expression of (FIG. 2A) dFnCas12a with crRNA targeting the *trxA* promoter results in repression of (FIG. 2B) *trxA* by $90 \pm 2.4\%$, which is associated with a (FIG. 2C) significant efficiency of plating (EOP) reduction to $12 \pm 7.0\%$. Data represented as mean SD (n=3). Rhamnose induction significantly lowers *trxA* expression vs. the NT-control ($p < 0.0001$, 2-Way ANOVA). Welch's two-tailed t-test was performed indicating significant reduction of EOP ($p < 0.05$).

[0027] FIGS. 3A-3C: CRISPRi-mediated gene repression in *E. coli* BL21 has varied impact on T7 lysis onset time and mean lysis time (T50%). CRISPRi induced by 2% (w/v) L-rhamnose for 4 h changes lysis profiles of T7 infecting log-phase *E. coli* BL21 depending on gene target and whether crRNA targets promoters (odd numbers) or coding sequences (even numbers). A non-targeting control was included in each experiment (g037). Optical densities (OD) were normalized with max OD=1 for clearer comparisons. Shaded regions represent max and min OD of triplicate experiments performed on different days. Representative curve effects (FIG. 3A): neutral (NT-control), shift (*trxA* (g001)), widening (*mukB* (g012)), and shoulder (*infC* (g020)). Lysis onset time (FIG. 3B) and mean lysis time (FIG. 3C) are represented as mean $\pm$ SD (n=3). Welch's two-tailed t-test was performed indicating significant change in lysis and mean lysis timing for as a result of CRISPRi induction ($p < 0.05$).

[0028] FIGS. 4A-4B: Gene overexpression and CRISPRi-mediated gene repression modulates efficiency of plating (EOP). Log-phase *E. coli* BL21 knockdown (FIG. 4A) and overexpression (FIG. 4B) strains were infected with phage T7 at MOI=0.0001. After 30 min (roughly once lysis cycle) at 37° C., further phage replication was stopped by adding chloroform to each culture followed by pipette mixing. Phage yields were enumerated by plaque counting on double-layer agar plates containing *E. coli* BL21 as propagation host. Efficiency of plating (EOP) represents the fraction change in assembled phage between (FIG. 4A) gene-repressed CRISPRi strains and their non-repressed counterparts carrying the same plasmid without CRISPRi induction or (FIG. 4B) fraction change in assembled phage between overexpressing strains and their non-induced counterparts carrying the same plasmid. Data represented as mean $\pm$ SD (n=3). Welch's two-tailed t-test was performed indicating significant change in EOP as a result of CRISPRi induction ($p < 0.05$). $p < 0.5$, *; $p < 0.01$, **; $p < 0.001$, ***; $p < 0.0001$, ****.

[0029] FIGS. 5A-5B: T7 genome pJ11-T7-Pr-sfGFP cascade phage-dependent TXTL mechanism. A cell-free expression system synthesizing sfGFP was used to troubleshoot CFBS reaction. (FIG. 5A) 0.5 nM T7 genome was included in a standard CFBS reaction using sfGFP expression controlled by T7 RNA polymerase (RNAP). (FIG. 5B) Transcription of sfGFP only occurs if T7 RNAP (T7 gene 1) itself is transcribed by endogenous *E. coli* RNAP. sfGFP expression reflects functional endogenous TXTL capacity and T7 RNAP synthesis and transcription activity.

[0030] FIGS. 6A-6C: Cell-free protein and bacteriophage synthesis yields are influenced by genetic background of TXTL donor. Biological duplicates (1 and 2) of cell lysates were prepared from *E. coli* BL21 overexpression (UP) or repression (DOWN) of various effector genes. Protein synthesis (FIG. 6A) was carried out using the T7 genome pJ11-T7-Pr-sfGFP cascade. (FIG. 6B) CFBS yields were measure by plaque count assays using wild-type BL21 as propagation host. Gray bars indicate protein and phage yields at 4 h. Black dotted lines indicate 4 h yields in wild-type BL21 lysates and colored lines indicate 20 h yields. (FIG. 6C) Relative sfGFP fluorescence and T7 titers shown as heatmaps. Data shows mean $\pm$ SD (n=3). Welch's two-tailed t-test was performed indicating significant increase in sfGFP fluoresce or T7 titer relative to the wild-type *E. coli* BL21 baseline ($p < 0.05$).

[0031] FIGS. 7A-7B: Comparison of cell-free systems for protein or phage synthesis. In vitro synthesis of (FIG. 7A) fluorescent reporter protein sfGFP and cell-free bacteriophage synthesis (FIG. 7B) of phage T7. Each system contains transcription/translation (TXTL) machinery derived from *E. coli* BL21 lysates, a DNA template, and energy buffer (20 canonical amino acids, phosphoenol pyruvate (PEP), NTPs, cofactors, and coenzymes). sfGFP is encoded on a circular plasmid (a) and phage T7 expression encoded on purified T7 gDNA. CFBS reactions are supplemented with molecular crowder PEG-8000 and dNTPs to support genome replication.

[0032] FIG. 8: CRISPRi/overexpression induction experimental workflow. Phenotypic readouts for gene repression and overexpression on T7 fitness included impacts on lysis curve profile (e.g. lysis timing, mean lysis timing) and efficiency of plating. Gene repression was validated by RT-qPCR.

[0033] FIG. 9: Single-plasmid CRISPR interference (CRISPRi) vector pCRJ001.

[0034] FIGS. 10A-10B: CRISPRi (KD) and overexpression (OX) inducer titration. CRISPRi knockdowns (KD) were induced in strains carrying pCRJ001 vectors using 0.02, 0.1, and 0.2% (w/v) L-rhamnose (FIG. 10A) and overexpression from pBAD vectors induced using 0.002, 0.01, 0.02, and 0.2% (w/v) L-arabinose (FIG. 10B). Bars reflect efficiency of plating (EOP) from 30 min T7 infections (MOI=0.0001). Data represented as mean $\pm$ SD (n=3). Ordinary One-way ANOVAs were performed indicating significant differences in EOP (p<0.05). A non-targeting control (NT-control) was included for KD strains and pBAD-sfGFP for OX strains.

[0035] FIG. 11A-11B: Rapid T7 enumeration assay. T7 infected log-phase BL21 carrying pJ11-T7-Pr-sfGFP. T7 RNA polymerase expressed during T7 infection drives sfGFP expression from the plasmid. Timing of sfGFP expression was dependent on T7 MOI (FIG. 11A) measured by normalized relative fluorescence units (RFU). A standard curve of log phage titer vs time to maximum sfGFP expression rate (T_{vmax}) (FIG. 11B) was generated and used to rapidly estimate T7 titers in cell-free bacteriophage synthesis experiments to inform dilutions for plaque assays. Dotted lines represent 95% CI for simple linear regression.

[0036] FIG. 12: T7 engineering primers and DNA oligos used in the present disclosure.

[0037] FIG. 13: Gibson Assembly of synthetic phage T7 expression heterologous genes. 0.2 pmol of each of T7 genome fragments A, B, E, F, and D mixed with 0.6 pmol pelB-ClyF (left) or sfGFP (right) expression cassettes into NEBuilder® HiFi DNA Assembly mix from New England Biolabs (Ipswich, MA) to assemble synthetic T7 genomes. 1 μ L of assembly mixture was used in 9 μ L CFBS reaction mixture to reboot into functional phage.

[0038] FIGS. 14A-14C: Spot test of heterologous gene expression using engineered T7. T7_ClyF lysates were prepared by infecting *E. coli* BL21 with phage rebooted via CFBS using synthetic genomes. Lysates were spotted onto LB agar plates overlaid with 100 μ L log-phase bacteria (OD=0.3)+4 mL SM agarose (50 mM TrisCl pH 7.5, 100 mM NaCl, 8 mM MgSO₄, 0.3% (w/v) agarose. 3 μ L of lysates were spotted on (FIG. 14A) Methicillin-resistant *S. aureus* (MRSA), (FIG. 14B) BL21, and (FIG. 14C) a MRSA/BL21 co-culture. Top rows show T7_ClyF lysates clearing bacteria in all 3 cases. Middle row was spotted with

T7_ColE1 lysates (ColE1 is bacteriocin that kills certain strains of Enterobacteriales and not Gram-positive bacteria such as MRSA). T7_ColE1 lysates have no effect on MRSA alone, but clear BL21. T7_ColE1 creates a halo of reduced bacterial growth on MRSA/BL21, but the center of the spot is turbid with MRSA. T7(+) was included as a control on the third row with no effect on MRSA alone and similar effects to those observed in the T7_ColE1 spots.

[0039] FIG. 15: RT-qPCR primers used in the present disclosure.

[0040] FIG. 16: Vectors used in the present disclosure.

[0041] FIG. 17: overexpression sequences used in the present disclosure.

[0042] FIG. 18: sgRNA sequences used in the present disclosure.

DETAILED DESCRIPTION

[0043] It is to be appreciated that certain aspects, modes, embodiments, variations and features of the present methods are described below in various levels of detail in order to provide a substantial understanding of the present technology.

[0044] In practicing the present methods, many conventional techniques in molecular biology, protein biochemistry, cell biology, immunology, microbiology and recombinant DNA are used. See, e.g., Sambrook and Russell eds. (2001) *Molecular Cloning: A Laboratory Manual*, 3rd edition; the series Ausubel et al. eds. (2007) *Current Protocols in Molecular Biology*; the series *Methods in Enzymology* (Academic Press, Inc., N.Y.); MacPherson et al. (1991) *PCR 1: A Practical Approach* (IRL Press at Oxford University Press); MacPherson et al. (1995) *PCR 2: A Practical Approach*; Harlow and Lane eds. (1999) *Antibodies, A Laboratory Manual*; Freshney (2005) *Culture of Animal Cells: A Manual of Basic Technique*, 5th edition; Gait ed. (1984) *Oligonucleotide Synthesis*; U.S. Pat. No. 4,683,195; Hames and Higgins eds. (1984) *Nucleic Acid Hybridization*; Anderson (1999) *Nucleic Acid Hybridization*; Hames and Higgins eds. (1984) *Transcription and Translation; Immobilized Cells and Enzymes* (IRL Press (1986)); Perbal (1984) *A Practical Guide to Molecular Cloning*; Miller and Calos eds. (1987) *Gene Transfer Vectors for Mammalian Cells* (Cold Spring Harbor Laboratory); Makrides ed. (2003) *Gene Transfer and Expression in Mammalian Cells*; Mayer and Walker eds. (1987) *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); and Herzenberg et al. eds (1996) *Weir's Handbook of Experimental Immunology*. Methods to detect and measure levels of polypeptide gene expression products (i.e., gene translation level) are well-known in the art and include the use of polypeptide detection methods such as antibody detection and quantification techniques. (See also, Strachan & Read, *Human Molecular Genetics*, Second Edition. (John Wiley and Sons, Inc., NY, 1999)).

[0045] Bacteriophage research has been boosted by a rising interest in using phage therapy to treat antibiotic-resistant bacterial infections. In addition, there is a desire to use phages and their unique proteins for specific biocontrol applications and diagnostics. Success in the bacteriophage industry requires development of robust scalable manufacturing platforms for upstream production of high phage titres and their downstream purification and concentration whilst achieving processing yields en route. Malik D. *Current Opinion in Biotechnology* 68:262-271 (2021). Phage therapy

is encumbered by significant challenges in the manufacturing process. Malik and Resch, *Front Microbiol.* 11: 584137 (2020).

[0046] The present disclosure demonstrates that lysates obtained from genetically modified bacterial host cells (e.g., *E. coli*) that overexpress one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or repress RecC subunit exonuclease RecBCD achieve enhanced CFBS yields relative to conventional cell-free phage synthesis methodologies.

Definitions

[0047] Unless defined otherwise, all technical and scientific terms used herein generally have the same meaning as commonly understood by one of ordinary skill in the art to which this technology belongs. As used in this specification and the appended claims, the singular forms “a”, “an” and “the” include plural referents unless the content clearly dictates otherwise. For example, reference to “a cell” includes a combination of two or more cells, and the like. Generally, the nomenclature used herein and the laboratory procedures in cell culture, molecular genetics, organic chemistry, analytical chemistry and nucleic acid chemistry and hybridization described below are those well-known and commonly employed in the art.

[0048] As used herein, the term “about” in reference to a number is generally taken to include numbers that fall within a range of 1%, 5%, or 10% in either direction (greater than or less than) of the number unless otherwise stated or otherwise evident from the context (except where such number would be less than 0% or exceed 100% of a possible value).

[0049] The term “amino acid” refers to naturally occurring and non-naturally occurring amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally encoded amino acids are the 20 canonical amino acids (alanine, arginine, asparagine, aspartic acid, cysteine, glutamine, glutamic acid, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, and valine) and pyrrolysine and selenocysteine. Amino acid analogs refer to agents that have the same basic chemical structure as a naturally occurring amino acid, i.e., an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, such as, homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (such as, norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. In some embodiments, amino acids forming a polypeptide are in the D form. In some embodiments, the amino acids forming a polypeptide are in the L form. In some embodiments, a first plurality of amino acids forming a polypeptide is in the D form and a second plurality is in the L form.

[0050] Amino acids are referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, are referred to by their commonly accepted single-letter code.

[0051] As used herein, “bacteriophage” or “phage” refers to a virus that infects bacteria. Bacteriophages are obligate intracellular parasites that multiply inside bacteria by co-opting some or all of the host biosynthetic machinery (i.e.,

viruses that infect bacteria). Though different bacteriophages may contain different materials, they all contain nucleic acid and protein, and can under certain circumstances be encapsulated in a lipid membrane. Depending upon the phage, the nucleic acid can be either DNA or RNA (but not both) and can exist in various forms.

[0052] The term “CRISPR enzyme system” refers to a programmable nuclease system for genetic editing that includes a CRISPR enzyme protein (e.g., Cas12a), or derivative thereof, and one or more non-coding guide RNAs (“gRNAs”) that provide the function of a CRISPR RNA (crRNA) and trans-activating RNA (tracrRNA) for the Cas9. The crRNA and tracrRNA can be separate RNA molecules or can be combined into a single RNA molecule to produce a “single guide RNA” (sgRNA). The crRNA or the tracrRNA portion of the sgRNA provide sequence that is complementary to the genomic target.

[0053] As used herein, the term “effective amount” refers to a quantity sufficient to achieve a desired effect or outcome, e.g., an amount which results in the manufacturing or synthesis of bacteriophage under in vitro cell-free conditions, e.g., cell-free bacteriophage synthesis (CFBS).

[0054] As used herein, “expression” includes one or more of the following: transcription of the gene into precursor mRNA; splicing and other processing of the precursor mRNA to produce mature mRNA; mRNA stability; translation of the mature mRNA into protein (including codon usage and tRNA availability); and glycosylation and/or other modifications of the translation product, if required for proper expression and function.

[0055] As used herein, an “expression control sequence” refers to polynucleotide sequences which are necessary to affect the expression of coding sequences to which they are operably linked. Expression control sequences are sequences which control the transcription, post-transcriptional events and translation of nucleic acid sequences. Expression control sequences include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (e.g., ribosome binding sites); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion. The nature of such control sequences differs depending upon the host organism; in prokaryotes, such control sequences generally include promoter, ribosomal binding site, and transcription termination sequence. The term “control sequences” is intended to encompass, at a minimum, any component whose presence is essential for expression, and can also encompass an additional component whose presence is advantageous, for example, leader sequences.

[0056] As used herein, “heterologous nucleic acid sequence” is any sequence placed at a location in the genome where it does not normally occur. A heterologous nucleic acid sequence may comprise a sequence that does not naturally occur in a bacteriophage, or it may comprise only sequences naturally found in the bacteriophage, but placed at a non-normally occurring location in the genome. In some embodiments, the heterologous nucleic acid sequence is not a natural phage sequence. In certain embodiments, the heterologous nucleic acid sequence is a natural phage sequence that is derived from a different phage. In other embodiments, the heterologous nucleic acid sequence

is a sequence that occurs naturally in the genome of a wild-type phage but is then relocated to another site where it does not naturally occur, rendering it a heterologous sequence at that new site.

[0057] “Homology” or “identity” or “similarity” refers to sequence similarity between two peptides or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same nucleobase or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences. A polynucleotide or polynucleotide region (or a polypeptide or polypeptide region) has a certain percentage (for example, at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 99%) of “sequence identity” to another sequence means that, when aligned, that percentage of bases (or amino acids) are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art. In some embodiments, default parameters are used for alignment. One alignment program is BLAST, using default parameters. In particular, programs are BLASTN and BLASTP, using the following default parameters: Genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+SwissProtein+SPupdate+PIR. Details of these programs can be found at the National Center for Biotechnology Information. Biologically equivalent polynucleotides are those having the specified percent homology and encoding a polypeptide having the same or similar biological activity. Two sequences are deemed “unrelated” or “non-homologous” if they share less than 40% identity, or less than 25% identity, with each other.

[0058] As used herein, a “host cell” is a bacterial cell that can be infected by a phage to yield progeny phage particles. A host cell can form phage particles from a particular type of phage genomic DNA. In some embodiments, the phage genomic DNA is introduced into the host cell by infecting the host cell with a phage. In some embodiments, the phage genomic DNA is introduced into the host cell using transformation, electroporation, or any other suitable technique. In some embodiments, the phage genomic DNA is substantially pure when introduced into the host cell. In some embodiments, the phage genomic DNA is present in a vector when introduced into the host cell. The definition of host cell can vary from one phage to another. For example, *E. coli* may be the natural host cell for a particular type of phage, but *Klebsiella pneumoniae* is not.

[0059] As used herein, the term “isolated” refers to a substance or entity that has been separated from at least some of the components with which it was associated when initially produced (whether in nature or in an experimental setting). Isolated substances and/or entities may be separated from at least about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, or more of the other components with which they were initially associated. In some embodiments, isolated substances and/or entities are more than about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%,

or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components.

[0060] As used herein, “operably linked” means that expression control sequences are positioned relative to the nucleic acid of interest to initiate, regulate or otherwise control transcription of the nucleic acid of interest.

[0061] As used herein, a “phage genome” includes naturally occurring phage genomes and derivatives thereof. Generally, the derivatives possess the ability to propagate in the same hosts as the naturally occurring phage. In some embodiments, the only difference between a naturally occurring phage genome and a derivative phage genome is at least one of a deletion and an addition of nucleotides from at least one end of the phage genome (if the genome is linear) or at least one point in the genome (if the genome is circular).

[0062] As used herein, the term “polynucleotide” or “nucleic acid” means any RNA or DNA, which may be unmodified or modified RNA or DNA. Polynucleotides include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, RNA that is mixture of single- and double-stranded regions, and hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, polynucleotide refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons.

[0063] As used herein, the term “recombinant” when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the material is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all.

[0064] As used herein, an endogenous nucleic acid sequence in the genome of an organism (or the encoded protein product of that sequence) is deemed “recombinant” herein if a heterologous sequence is placed adjacent to the endogenous nucleic acid sequence, such that the expression of this endogenous nucleic acid sequence is altered. In this context, a heterologous sequence is a sequence that is not naturally adjacent to the endogenous nucleic acid sequence, whether or not the heterologous sequence is itself endogenous to the organism (originating from the same organism or progeny thereof) or exogenous (originating from a different organism or progeny thereof). By way of example, a promoter sequence can be substituted (e.g., by homologous recombination) for the native promoter of a gene in the genome of an organism, such that this gene has an altered expression pattern. This gene would be “recombinant” because it is separated from at least some of the sequences that naturally flank it. A nucleic acid is also considered “recombinant” if it contains any modifications that do not naturally occur in the corresponding nucleic acid in a genome. For instance, an endogenous coding sequence is considered “recombinant” if it contains an insertion, deletion or a point mutation introduced artificially, e.g., by

human intervention. A “recombinant nucleic acid” also includes a nucleic acid integrated into a host cell chromosome at a heterologous site and a nucleic acid construct present as an episome.

[0065] As used herein, a “recombinant bacteriophage genome” is a bacteriophage genome that has been genetically modified by the insertion of a heterologous nucleic acid sequence into the bacteriophage genome. A “recombinant bacteriophage” means a bacteriophage that comprises a recombinant bacteriophage genome. In some embodiments, the bacteriophage genome is modified by recombinant DNA technology to introduce a heterologous nucleic acid sequence into the genome at a defined site. In some embodiments, the heterologous nucleic acid sequence is introduced with no corresponding loss of endogenous phage genomic nucleotides. In other words, if bases N1 and N2 are adjacent in the wild-type bacteriophage genome, the heterologous nucleic acid sequence is inserted between N1 and N2. Thus, in the resulting recombinant bacteriophage genome, the heterologous nucleic acid sequence is flanked by nucleotides N1 and N2. In some embodiments, endogenous phage nucleotides are removed or replaced during the insertion of the heterologous nucleic acid sequence. For example, in some embodiments, the heterologous nucleic acid sequence is inserted in place of some or all of the endogenous phage sequence which is removed. In some embodiments, endogenous phage sequences are removed from a position in the phage genome distant from the site(s) of insertion of the heterologous nucleic acid sequences.

[0066] As used herein, the term “sample” refers to clinical samples obtained from a subject or isolated microorganisms. In certain embodiments, a sample is obtained from a biological source (i.e., a “biological sample”), such as tissue, bodily fluid, or microorganisms collected from a subject. Sample sources include, but are not limited to, mucus, sputum, bronchial alveolar lavage (BAL), bronchial wash (BW), whole blood, bodily fluids, cerebrospinal fluid (CSF), urine, plasma, serum, or tissue.

Bacteriophage

[0067] Bacteriophage are obligate intracellular parasites that multiply inside bacteria by co-opting some or all of the host biosynthetic machinery. Phages contain nucleic acid and protein, and may be enveloped by a lipid membrane. Depending upon the phage, the nucleic acid genome can be either DNA or RNA but not both, and can exist in either circular or linear forms. The size of the phage genome varies depending upon the phage. The simplest phages have genomes that are only a few thousand nucleotides in size, while the more complex phages may contain more than 100,000 nucleotides in their genome, and in rare instances no more than 500,000 bp. The number and amount of individual types of protein in phage particles will vary depending upon the phage. The proteins function in infection and to protect the nucleic acid genome from environmental nucleases.

[0068] Phage genomes come in a variety of sizes and shapes (e.g., linear or circular). Most phages range in size from 24-200 nm in diameter. The capsid is composed of many copies of one or more phage proteins, and acts as a protective envelope around the phage genome. Many phages have tails attached to the phage capsid. The tail is a hollow tube through which the phage nucleic acid passes during infection. The size of the tail can vary and some phages do

not even have a tail structure. In the more complex phages, the tail is surrounded by a contractile sheath which contracts during infection of the bacterial host cell. At the end of the tail, phages have a base plate and one or more tail fibers attached to it. The base plate and tail fibers are involved in the binding of the phage to the host cell.

[0069] Lytic or virulent phages are phages which can only multiply in bacteria and lyse the bacterial host cell at the end of the life cycle of the phage. The lifecycle of a lytic phage begins with an eclipse period. During the eclipse phase, no infectious phage particles can be found either inside or outside the host cell. The phage nucleic acid takes over the host biosynthetic machinery and phage specific mRNAs and proteins are produced. Early phage mRNAs code for early proteins that are needed for phage DNA synthesis and for shutting off host DNA, RNA and protein biosynthesis. In some cases, the early proteins actually degrade the host chromosome. After phage DNA is made late mRNAs and late proteins are made. The late proteins are the structural proteins that comprise the phage as well as the proteins needed for lysis of the bacterial cell. In the next phase, the phage nucleic acid and structural proteins are assembled and infectious phage particles accumulate within the cell. The bacteria begin to lyse due to the accumulation of the phage lysis protein, leading to the release of intracellular phage particles. The number of particles released per infected cell can be as high as 1000 or more. Lytic phage may be enumerated by a plaque assay. The assay is performed at a low enough concentration of phage such that each plaque arises from a single infectious phage. The infectious particle that gives rise to a plaque is called a PFU (plaque forming unit).

[0070] Lysogenic phages are those that can either multiply via the lytic cycle or enter a quiescent state in the host cell. In the quiescent state, the phage genome exists as a prophage (i.e., it has the potential to produce phage). In most cases, the phage DNA actually integrates into the host chromosome and is replicated along with the host chromosome and passed on to the daughter cells. The host cell harboring a prophage is not adversely affected by the presence of the prophage and the lysogenic state may persist indefinitely. The lysogenic state can be terminated upon exposure to adverse conditions. Conditions which favor the termination of the lysogenic state include: desiccation, exposure to UV or ionizing radiation, exposure to mutagenic chemicals, etc. Adverse conditions lead to the production of proteases (rec A protein), the expression of the phage genes, reversal of the integration process, and lytic multiplication.

[0071] In some embodiments, a phage genome comprises at least 5 kilobases (kb), at least 10 kb, at least 15 kb, at least 20 kb, at least 25 kb, at least 30 kb, at least 35 kb, at least 40 kb, at least 45 kb, at least 50 kb, at least 55 kb, at least 60 kb, at least 65 kb, at least 70 kb, at least 75 kb, at least 80 kb, at least 85 kb, at least 90 kb, at least 95 kb, at least 100 kb, at least 105 kb, at least 110 kb, at least 115 kb, at least 120 kb, at least 125 kb, at least 130 kb, at least 135 kb, at least 140 kb, at least 145 kb, at least 150 kb, at least 175 kb, at least 200 kb, at least 225 kb, at least 250 kb, at least 275 kb, at least 300 kb, at least 325 kb, at least 350 kb, at least 375 kb, at least 400 kb, at least 425 kb, at least 450 kb, at least 475 kb, or at least 500 kb of nucleic acids.

Cell-Free Bacteriophage Synthesis Compositions and Methods of the Present Technology

[0072] In one aspect, the present disclosure provides genetically modified bacterial host cells (e.g., *E. coli*) that overexpress one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or comprise a vector that includes a nucleic acid sequence encoding a CRISPR enzyme and a vector that includes a nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC. In some embodiments, the vector including the nucleic acid sequence encoding the CRISPR enzyme and the vector including the nucleic acid sequence encoding the crRNA are the same or distinct.

[0073] An exemplary nucleic acid sequence of translation initiation factor IF-3 (infC) is set forth in SEQ ID NO: 39, as provided below:

(SEQ ID NO: 39)

ATTAAAGGCGGAAACGAGTTCAAACGGCGCGCCCTAACCGTATCAATG
 GCGAAATTCGCGCCAGGAAGTTCGCTTAACAGGTCTGGAAGCGGAGCA
 GCTTGGTATTGTGAGTCTGAGAGAAGCTCTGGAGAAAGCAGAAGAAGCC
 GGAGTAGACTTAGTCGAGATCAGCCCTAACGCCGAGCCCGGTTTGTC
 GTATAATGGATTACGGCAAATTCCTCTATGAAAAGAGCAAGTCTTCTAA
 GGAACAGAAGAAAAAGCAAAAGTTATCCAGGTTAAGGAAATTAATTC
 CGTCCTGGTACAGATGAAGGCGACTATCAGGTAAACTCCGAGCCTGA
 TTCGCTTTCTCGAAGAGGGTGATAAGCCAAATCACGCTGCGTTTCCG
 CGGTCGTGAGATGGCGCACCAGCAAATCGGTATGGAAGTGCTTAATCGC
 GTGAAAGACGATTGTGAAGAACTGGCAGTGGTGAATCCTTCCCAACGA
 AGATCGAAGGCCCGCAGATGATCATGGTCTCGCTCCTAAGAAGAAACA
 GTAA

[0074] An exemplary nucleic acid sequence of OxyS is set forth in SEQ ID NO: 41, as provided below:

(SEQ ID NO: 41)

ATTCTGACTGATAATTGCTCACAGAAACGGAGCGCACCTCTTTAAAC
 CTTGAAGTCACTGCCCGTTTCGAGAGTTTCTCAACTCGAATAACTAAAG
 CCAACGTGAACCTTTTGGGATCTCCAGGATCCGCTTTTTTTTGCATAA
 AAA

[0075] An exemplary nucleic acid sequence of CyaR is set forth in SEQ ID NO: 42, as provided below:

(SEQ ID NO: 42)

GCTGAAAACATAACCCATAAAATGCTAGCTGTACCAGGAACCACTCC
 TTAGCCTGTGTAATCTCCCTTACACGGGCTTATTTTTT

[0076] In some embodiments, the genetically modified bacterial host cells comprise a non-endogenous expression vector that comprises a translation initiation factor IF-3 (infC) nucleic acid sequence of SEQ ID NO: 39, an OxyS nucleic acid sequence of SEQ ID NO: 41, a CyaR nucleic

acid sequence of SEQ ID NO: 42, or any combination thereof. In some embodiments, the genetically modified bacterial host cells comprise a heterologous nucleic acid sequence that is at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to SEQ ID NO: 39, SEQ ID NO: 41 or SEQ ID NO: 42. Additionally or alternatively, in some embodiments, the expression levels and/or activity of infC, OxyS or CyaR in the genetically modified bacterial host cell is at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, at least 90, at least 100, at least 200, at least 300, at least 400, at least 500, at least 600, at least 700, at least 800, at least 900, or at least 1000 times higher compared to that observed in a native bacterial host cell, wherein the genetically modified bacterial host cell is the same species as the native bacterial host cell.

[0077] Additionally or alternatively, in some embodiments, the non-endogenous expression vector comprising the infC nucleic acid sequence and the non-endogenous expression vector comprising the OxyS nucleic acid sequence and/or the CyaR nucleic acid sequence are the same or distinct. In other embodiments, the non-endogenous expression vector comprising the OxyS nucleic acid sequence and the non-endogenous expression vector comprising the CyaR nucleic acid sequence are the same or distinct.

[0078] In any of the preceding embodiments of the genetically modified bacterial host cells disclosed herein, the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is a plasmid, a cosmid, a bacmid, a bacterial artificial chromosome (BAC), or a viral vector. In certain embodiments, the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is operably linked to an expression control sequence. The expression control sequence may be an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter.

[0079] Additionally or alternatively, in some embodiments, the genetically modified bacterial host cells of the present technology comprise a vector that includes a nucleic acid sequence encoding a CRISPR enzyme and a vector that includes a nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC. In some embodiments, the vector including the nucleic acid sequence encoding the CRISPR enzyme and the vector including the nucleic acid sequence encoding the crRNA are the same or distinct. In certain embodiments, the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter.

[0080] Additionally or alternatively, in certain embodiments, the CRISPR enzyme is a Type II CRISPR enzyme. In some embodiments, the CRISPR enzyme catalyzes DNA cleavage. In some embodiments, the CRISPR enzyme cata-

lyzes RNA cleavage. In some embodiments, the CRISPR enzyme is any Cas12a or Cas9 protein, for instance any naturally-occurring bacterial Cas12a or Cas9 as well as any variants, homologs or orthologs thereof. Non-limiting examples of Cas proteins include Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn1 and Csx12), Cas10, Cas12a from *F. novicida* (dFnCas12a), Csy1, Csy2, Csy3, Cse1, Cse2, Cse1, Cse2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, Csf4, homologues thereof, or variants thereof. In some embodiments, the CRISPR enzyme cleaves both strands of the target nucleic acid at the Protospacer Adjacent Motif (PAM) site. In certain embodiments of the genetically modified bacterial host cells of the present technology, the CRISPR enzyme is a nuclease-deficient *Francisella novicida* Cas12a (dFnCas12a) and/or the nucleic acid sequence encoding the crRNA is selected from the group consisting of SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88 and SEQ ID NO: 89.

[0081] Additionally or alternatively, in some embodiments, the genetically modified bacterial host cells of the present technology comprise a deletion or mutation in (a) a promoter region located upstream of a transcription start site (TSS) of *recC*, or (b) a *recC* coding region located downstream of a transcription start site (TSS) of *recC*, wherein the mutation or deletion represses transcription of *recC*. The deletion may be located upstream or downstream of the transcription start site (TSS) of *recC*. In some embodiments, the deletion is about 10-about 100 base pairs (bps) in length. In certain embodiments, the deletion is about 10 bps, about 11 bps, about 12 bps, about 13 bps, about 14 bps, about 15 bps, about 16 bps, about 17 bps, about 18 bps, about 19 bps, about 20 bps, about 21 bps, about 22 bps, about 23 bps, about 24 bps, about 25 bps, about 26 bps, about 27 bps, about 28 bps, about 29 bps, about 30 bps, about 31 bps, about 32 bps, about 33 bps, about 34 bps, about 35 bps, about 36 bps, about 37 bps, about 38 bps, about 39 bps, about 40 bps, about 41 bps, about 42 bps, about 43 bps, about 44 bps, about 45 bps, about 46 bps, about 47 bps, about 48 bps, about 49 bps, about 50 bps, about 51 bps, about 52 bps, about 53 bps, about 54 bps, about 55 bps, about 56 bps, about 57 bps, about 58 bps, about 59 bps, about 60 bps, about 61 bps, about 62 bps, about 63 bps, about 64 bps, about 65 bps, about 66 bps, about 67 bps, about 68 bps, about 69 bps, about 70 bps, about 71 bps, about 72 bps, about 73 bps, about 74 bps, about 75 bps, about 76 bps, about 77 bps, about 78 bps, about 79 bps, about 80 bps, about 81 bps, about 82 bps, about 83 bps, about 84 bps, about 85 bps, about 86 bps, about 87 bps, about 88 bps, about 89 bps, about 90 bps, about 91 bps, about 92 bps, about 93 bps, about 94 bps, about 95 bps, about 96 bps, about 97 bps, about 98 bps, about 99 bps, or about 100 bps in length. In other embodiments, the mutation is a nonsense mutation, a frameshift mutation, or a missense mutation.

[0082] In any of the foregoing embodiments described herein, the genetically modified bacterial host cells are *E. coli* cells.

[0083] In one aspect, the present disclosure provides a cell lysate obtained from any and all embodiments of the genetically modified bacterial host cells disclosed herein. The cell lysates of the present technology comprise an effective amount of transcription/translation (TXTL) machinery that

is configured to synthesize bacteriophage under cell-free conditions. The cell lysates may be prepared using one or more of French-press cell lysis, sonication, runoff reactions, or lysate dialysis.

[0084] In another aspect, the present disclosure provides an in vitro method for synthesizing bacteriophage virions comprising contacting a bacteriophage genome with any and all embodiments of the cell lysates disclosed herein and an energy buffer in vitro to obtain a reaction mixture, and incubating the reaction mixture under conditions to produce viable phage virions, wherein the energy buffer comprises canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes. In some embodiments, the reaction mixture is incubated at 16° C. to 42° C. for about 1-24 hours. In certain embodiments, the reaction mixture is incubated at 16° C., 17° C., 18° C., 19° C., 20° C., 21° C., 22° C., 23° C., 24° C., 25° C., 26° C., 27° C., 28° C., 29° C., 30° C., 31° C., 32° C., 33° C., 34° C., 35° C., 36° C., 37° C., 38° C., 39° C., 40° C., 41° C., or 42° C. for about 1 hour, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, or 24 hours. The bacteriophage genome may be derived from a naturally occurring bacteriophage, or a genetically engineered bacteriophage. In some embodiments, the bacteriophage genome is obtained from a bacteriophage that specifically infects the naturally occurring counterpart of the genetically modified bacterial host cells disclosed herein, such as *E. coli*. Additionally or alternatively, in some embodiments, the bacteriophage genome is obtained from a bacteriophage selected from among T7 phage, T7-like phage, Lambda, K1E, T3, T5, T4, or PhiX174.

[0085] Additionally or alternatively, in some embodiments, the energy buffer further comprises one or more of a polyethylene glycol (PEG) polymer (such as PEG-8000 or PEG-6000), deoxynucleotide triphosphates (dNTPs), and stabilizers. In certain embodiments, the energy buffer further comprises one or more of Mg-glutamate, K-glutamate, tRNA, cAMP, folinic acid, spermidine, 3-PGA, HEPES, Nicotinamide adenine dinucleotide (NAD), coenzyme A (CoA), DTT, and maltodextrin.

[0086] In any of the preceding embodiments of the methods disclosed herein, the reaction mixture further comprises a vector including a nucleic acid sequence encoding a reporter gene, wherein the nucleic acid sequence encoding the reporter gene is operably linked to a promoter that is responsive to a RNA polymerase (RNAP) encoded by the bacteriophage genome. In certain embodiments, the RNAP is selected from among SP6 RNA Polymerase, T3 RNA Polymerase and T7 RNA polymerase. The reporter gene may encode a bioluminescent protein, a fluorescent protein, or a chemiluminescent protein. Examples of bioluminescent proteins include, but are not limited to, Aequorin, firefly luciferase, *Renilla* luciferase, red luciferase, luxAB, or nanoluciferase. Examples of chemiluminescent proteins include, but are not limited to, β -galactosidase, horseradish peroxidase (HRP), or alkaline phosphatase. Examples of fluorescent proteins include, but are not limited to, sfGFP, TagBFP, Azurite, EBFP2, mKalamal, Sirius, Sapphire, T-Sapphire, ECFP, Cerulean, SCFP3A, mTurquoise, monomeric Midoriishi-Cyan, TagCFP, mTFP1, EGFP, Emerald, Superfolder GFP, Monomeric Azami Green, TagGFP2, mUKG, mWasabi, EYFP, Citrine, Venus, SYFP2, TagYFP, Mono-

meric Kusabira-Orange, mKOx, mKO2, mOrange, mOrange2, mRaspberry, mCherry, dsRed, mStrawberry, mTangerine, tdTomato, TagRFP, TagRFP-T, mApple, mRuby, mPlum, HcRed-Tandem, mKate2, mNeptune, NirFP, TagRFP657, IFP1.4, iRFP, mKeima Red, LSS-mKate1, LSS-mKate2, PA-GFP, PAmCherry1, PAtagRFP, Kaede (green), Kaede (red), KikGR1 (green), KikGR1 (red), PS-CFP2, PS-CFP2, mEos2 (green), mEos2 (red), PSmOrange, or Dronpa.

[0087] Additionally or alternatively, in some embodiments of the methods disclosed herein, the phage virions show increased efficiency of plating (EOP) relative to phage virions obtained with cell-free bacteriophage synthesis (CFBS) in a control lysate obtained from a wild-type bacterial cell, wherein the wild-type bacterial cell and the genetically modified bacterial host cell are the same species.

Polynucleotides, Polypeptides and Analogs

[0088] Also included in the presently disclosed subject matter are infC, OxyS, CyaR and/or RecC cRNA polynucleotides and their corresponding polypeptides or fragments that may be modified in ways that enhance their functional activity when expressed in a bacterial host cell. The presently disclosed subject matter provides methods for optimizing an amino acid sequence or a nucleic acid sequence by producing an alteration in the sequence. Such alterations can comprise certain mutations, deletions, insertions, or post-translational modifications. The presently disclosed subject matter further comprises analogs of any naturally-occurring polypeptide of the presently disclosed subject matter. Analogs can differ from a naturally-occurring polypeptide of the presently disclosed subject matter by amino acid sequence differences. Analogs of the presently disclosed subject matter can generally exhibit at least about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or more identity or homology with all or part of a naturally-occurring amino acid sequence of the presently disclosed subject matter. The length of sequence comparison is at least about 5, about 10, about 15, about 20, about 25, about 50, about 75, about 100 or more amino acid residues. Again, in an exemplary approach to determining the degree of identity, a BLAST program can be used, with a probability score between e^{-3} and e^{-100} indicating a closely related sequence.

[0089] Analogs can differ from the naturally-occurring polypeptides of the presently disclosed subject matter by alterations in primary sequence. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethanemethyl sulfate or by site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, *Molecular Cloning: A Laboratory Manual* (2nd ed.), CSH Press, 1989, or Ausubel et al., supra). Also included are cyclized peptides, molecules, and analogs which contain residues other than L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g., beta (β) or gamma (γ) amino acids.

[0090] In addition to full-length polypeptides, the presently disclosed subject matter also provides fragments of any one of the polypeptides or peptide domains of the presently disclosed subject matter. A fragment can be at least about 5, about 10, about 13, or about 15 amino acids. In some embodiments, a fragment is at least about 20 contiguous

amino acids, at least about 30 contiguous amino acids, or at least about 50 contiguous amino acids. In some embodiments, a fragment is at least about 60 to about 80, about 100, about 200, about 300 or more contiguous amino acids.

[0091] In accordance with the presently disclosed subject matter, the polynucleotides encoding infC, OxyS, CyaR and/or RecC cRNA can be modified by codon optimization. Codon optimization can alter both naturally occurring and recombinant gene sequences to achieve the highest possible levels of productivity in any given expression system. Factors that are involved in different stages of protein expression include codon adaptability, mRNA structure, and various cis-elements in transcription and translation. Any suitable codon optimization methods or technologies that are known to ones skilled in the art can be used to modify the polynucleotides of the presently disclosed subject matter, including, but not limited to, OptimumGene™, Encor optimization, and Blue Heron.

Vectors

[0092] As used herein, a “vector” is a replicable nucleic acid from which one or more heterologous proteins or RNAs can be expressed when the vector is introduced into an appropriate host cell. The vector is used to introduce the nucleic acid encoding the polypeptide or RNA into the host cell for amplification of the nucleic acid or for expression/display of the polypeptide or RNA encoded by the nucleic acid. As used herein, a vector also includes “virus vectors” or “viral vectors.” Viral vectors are engineered viruses that are operatively linked to exogenous genes to transfer (as vehicles or shuttles) the exogenous genes into cells.

[0093] Many expression vectors are available and known to those of skill in the art and can be used for nonendogenous expression of infC, OxyS, CyaR, and/or a RecC cRNA. The choice of expression vector will be influenced by the choice of host expression system. Such selection is well within the level of skill of the skilled artisan. In general, expression vectors can include transcriptional promoters and optionally enhancers, translational signals, and transcriptional and translational termination signals. Expression vectors that are used for stable transformation typically have a selectable marker which allows selection and maintenance of the transformed cells. In some cases, an origin of replication can be used to amplify the copy number of the vector in the cells. The vectors typically remain episomal, but can be designed to effect integration of a gene or portion thereof into a chromosome of the genome. Also contemplated are vectors that are artificial chromosomes, such as yeast artificial chromosomes and mammalian artificial chromosomes. Selection and use of such vehicles are well known to those of skill in the art.

[0094] Vectors also can contain additional nucleotide sequences operably linked to the ligated nucleic acid molecule, such as, for example, an epitope tag such as for localization, e.g. a hexa-his tag or a myc tag, hemagglutinin tag or a tag for purification, for example, a GST fusion, and a sequence for directing protein secretion and/or membrane association.

[0095] Expression of the infC, OxyS, or CyaR genes and/or a RecC cRNA can be controlled by any promoter/enhancer known in the art. Suitable bacterial promoters are well known in the art and described herein below. Selection of the promoter used to direct expression of a heterologous

nucleic acid depends on the particular application and is within the level of skill of the skilled artisan. Promoters which can be used include but are not limited to prokaryotic expression vectors such as the β -lactamase promoter (Jay et al. (1981) *Proc. Natl. Acad. Sci. USA* 75:5543), the tac promoter (DeBoer et al. (1983) *Proc. Natl. Acad. Sci. USA* 50:21-25); see also "Useful Proteins from Recombinant Bacteria" (1980) in *Scientific American* 242:79-94), rhaB promoter and the like.

[0096] In addition to the promoter, the expression vector typically contains a transcription unit or expression cassette that contains all the additional elements required for the expression of the heterologous nucleic acid, in host cells. A typical expression cassette contains a promoter operably linked to the gene sequence and signals required for efficient polyadenylation of the transcript, ribosome binding sites and translation termination. Additional elements of the cassette can include enhancers. In addition, the cassette typically contains a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region can be obtained from the same gene as the promoter sequence or can be obtained from different genes.

[0097] Any methods known to those of skill in the art for the insertion of DNA fragments into a vector can be used to construct expression vectors containing a nucleic acid encoding any of the genes disclosed herein. These methods can include in vitro recombinant DNA and synthetic techniques and in vivo recombinants (genetic recombination). The insertion into a cloning vector can, for example, be accomplished by ligating the DNA fragment into a cloning vector which has complementary cohesive termini. If the complementary restriction sites used to fragment the DNA are not present in the cloning vector, the ends of the DNA molecules can be enzymatically modified. Alternatively, any site desired can be produced by ligating nucleotide sequences (linkers) onto the DNA termini; these ligated linkers can contain specific chemically synthesized nucleic acids encoding restriction endonuclease recognition sequences.

[0098] Genetic modification of engineered host cells can be accomplished by transducing a substantially homogeneous cell composition with a recombinant DNA or RNA construct. The vector can be a retroviral vector (e.g., gamma retroviral), which is employed for the introduction of the DNA or RNA construct into the host cell genome. For example, a polynucleotide encoding infC, OxyS, CyaR and/or a RecC cRNA can be cloned into a retroviral vector and expression can be driven from its endogenous promoter, from the retroviral long terminal repeat, or from an alternative internal promoter.

[0099] Non-viral vectors or RNA can be used as well. Random chromosomal integration, or targeted integration (e.g., using a nuclease, transcription activator-like effector nucleases (TALENs), Zinc-finger nucleases (ZFNs), and/or clustered regularly interspaced short palindromic repeats (CRISPRs), or transgene expression (e.g., using a natural or chemically modified RNA) can be used.

[0100] For initial genetic modification of the cells to provide cells expressing infC, OxyS, or CyaR genes and/or a RecC cRNA, a retroviral vector can be employed for transduction. However, any other suitable viral vector or non-viral delivery system can be used for genetic modification of cells.

[0101] Possible methods of transduction also include direct co-culture of the cells with producer cells, e.g., by the method of Bregni et al. (1992) *Blood* 80: 1418-1422, or culturing with viral supernatant alone or concentrated vector stocks with or without appropriate growth factors and polycations, e.g., by the method of Xu et al. (1994) *Exp. Hemat.* 22:223-230; and Hughes et al. (1992) *J. Clin. Invest.* 89: 1817.

[0102] In some embodiments, the chosen vector exhibits high efficiency of infection and stable integration and expression (see, e.g., Cayouette et al. (1997) *Human Gene Therapy* 8:423-430; Kido et al. (1996) *Current Eye Research* 15:833-844; Bloomer et al. (1997) *Journal of Virology* 71:6641-6649; Naldini et al. (1996) *Science* 272:263-267; and Miyoshi et al. (1997) *Proc. Natl. Acad. Sci. U.S.A.* 94: 10319). Other viral vectors that can be used include, for example, adenoviral, lentiviral, and adeno-associated viral vectors, vaccinia virus, a bovine papilloma virus, or a herpes virus, such as Epstein-Barr Virus (also see, for example, the vectors of Miller (1990) *Human Gene Therapy* 15-14; Friedman (1989) *Science* 244: 1275-1281; Eglitis et al. (1988) *BioTechniques* 6:608-614; Tolstoshev et al. (1990) *Current Opinion in Biotechnology* 1:55-61; Sharp (1991) *The Lancet* 337: 1277-1278; Cornetta et al. (1987) *Nucleic Acid Research and Molecular Biology* 36:311-322; Anderson (1984) *Science* 226:401-409; Moen (1991) *Blood Cells* 17:407-416; Miller et al. (1989) *Biotechnology* 7:980-990; Le Gal La Salle et al. (1993) *Science* 259:988-990; and Johnson (1995) *Chest* 107:77S-83S). Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg et al. (1990) *N. Engl. J. Med* 323:370; Anderson et al., U.S. Pat. No. 5,399,346).

[0103] In certain non-limiting embodiments, the vector expressing an infC, OxyS, CyaR and/or RecC cRNA nucleic acid sequence is a retroviral vector, e.g., an oncoretroviral vector. In some instances, the retroviral vector is a SFG retroviral vector or murine stem cell virus (MSCV) retroviral vector. In certain non-limiting embodiments, the vector expressing an infC, OxyS, CyaR and/or RecC cRNA nucleic acid sequence is a lentiviral vector. In certain non-limiting embodiments, the vector expressing an infC, OxyS, CyaR and/or RecC cRNA nucleic acid sequence is a transposon vector.

[0104] Non-viral approaches can also be employed for the expression of a protein in cell. For example, a nucleic acid molecule can be introduced into a cell by administering the nucleic acid in the presence of lipofection (Feigner et al. (1987) *Proc. Nat'l. Acad. Sci. U.S.A.* 84:7413; Ono et al. (1990) *Neuroscience Letters* 17:259; Brigham et al. (1989) *Am. J. Med. Sci.* 298:278; Staubinger et al. (1983) *Methods in Enzymology* 101:512), asialoorosomucoid-polylysine conjugation (Wu et al. (1988) *Journal of Biological Chemistry* 263: 14621; Wu et al. (1989) *Journal of Biological Chemistry* 264: 16985), or by micro-injection under surgical conditions (Wolff et al. (1990) *Science* 247: 1465). Other non-viral means for gene transfer include transfection in vitro using calcium phosphate, DEAE dextran, electroporation, and protoplast fusion. Liposomes can also be potentially beneficial for delivery of DNA into a cell. Transient expression can be obtained by RNA electroporation.

[0105] The resulting cells can be grown under conditions similar to those for unmodified cells, whereby the modified cells can be expanded and used for a variety of purposes.

Kits

[0106] The present technology provides kits for preparing donor bacterial host cell lysates for cell-free bacteriophage synthesis (CFBS). In one aspect, the present disclosure provides a kit comprising one or more expression vectors comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO: 39, SEQ ID NO: 41 and SEQ ID NO: 42, and instructions for using the same to prepare donor bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In certain embodiments, the at least one nucleic acid sequence is operably linked to an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter. Additionally or alternatively, in some embodiments, the kits further comprise vials containing naturally-occurring or non-natural bacterial host cells that can be transformed with the one or more expression vectors comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO: 39, SEQ ID NO: 41 and SEQ ID NO: 42. In some embodiments, the bacterial host cells are *E. coli*. In certain embodiments, the bacterial host cells are *E. coli* strain BL21.

[0107] In another aspect, the present disclosure provides a kit comprising one or more expression vectors comprising at least one nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of *recC* or a *recC* coding region located downstream of a transcription start site (TSS) of *recC*, and instructions for using the same to prepare donor bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In some embodiments, the one or more expression vectors further comprise a nucleic acid sequence encoding a CRISPR enzyme. Examples of CRISPR enzymes include, but are not limited to, nuclease-deficient Cas12a from *E. novicida* (dFnCas12a), Cas1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9, Cas10, Csy1, Csy2, Csy3, Cse1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx15, Csf1, Csf2, Csf3, and Csf4. Additionally or alternatively, in some embodiments, the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the at least one nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter. In some embodiments, the nucleic acid sequence encoding the crRNA is selected from the group consisting of SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88 and SEQ ID NO: 89. Additionally or alternatively, in some embodiments, the kits further comprise vials containing naturally-occurring or non-natural bacterial host cells that can be transformed with the one or more expression vectors comprising the nucleic acid sequence encoding the CRISPR enzyme and/or the at least one nucleic acid sequence encoding the crRNA. In some embodiments, the bacterial host cells are *E. coli*. In certain embodiments, the bacterial host cells are *E. coli* strain BL21.

[0108] In yet another aspect, the present disclosure provides a kit comprising any and all embodiments of the genetically modified bacterial host cell described herein, and instructions for using the same to prepare donor bacterial cell lysates for cell-free bacteriophage synthesis (CFBS). In some embodiments, the genetically modified bacterial host cells are *E. coli*. In certain embodiments, the bacterial host cells are *E. coli* strain BL21.

[0109] In any and all embodiments of the kits disclosed herein, the kits further comprise one or more of polyethylene glycol (PEG) polymers (e.g., PEG-8000, PEG-6000), deoxynucleotide triphosphates (dNTPs), stabilizers, and an energy buffer comprising canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes.

[0110] Additionally or alternatively, in some embodiments, the kits further comprise a vector including a nucleic acid sequence encoding a reporter gene, wherein the nucleic acid sequence encoding the reporter gene is operably linked to a promoter that is responsive to a phage RNA polymerase (RNAP), such as SP6 RNA Polymerase, T3 RNA Polymerase and T7 RNA polymerase. In some embodiments, the reporter gene encodes a bioluminescent protein, a fluorescent protein, or a chemiluminescent protein. Examples of bioluminescent proteins include, but are not limited to, Aequorin, firefly luciferase, Renilla luciferase, red luciferase, luxAB, or nanoluciferase. Examples of chemiluminescent proteins include, but are not limited to, β -galactosidase, horseradish peroxidase (HRP), or alkaline phosphatase. Examples of fluorescent proteins include, but are not limited to, sfGFP, TagBFP, Azurite, EBFP2, mKalamal, Sirius, Sapphire, T-Sapphire, ECFP, Cerulean, SCFP3A, mTurquoise, monomeric Midoriishi-Cyan, TagCFP, mTFP1, EGFP, Emerald, Superfolder GFP, Monomeric Azami Green, TagGFP2, mUKG, mWasabi, EYFP, Citrine, Venus, SYFP2, TagYFP, Monomeric Kusabira-Orange, mKOx, mKO2, mOrange, mOrange2, mRaspberry, mCherry, dsRed, mStrawberry, mTangerine, tdTomato, TagRFP, TagRFP-T, mApple, mRuby, mPlum, HcRed-Tandem, mKate2, mNeptune, NirFP, TagRFP657, IFP1.4, iRFP, mKeima Red, LSS-mKate1, LSS-mKate2, PA-GFP, PAmCherry1, PAtagRFP, Kaede (green), Kaede (red), KikGR1 (green), KikGR1 (red), PS-CFP2, PS-CFP2, mEos2 (green), mEos2 (red), PSmOrange, or Dronpa.

[0111] In any of the preceding embodiments, the kits of the present technology comprise one or more coded/labeled vials that contain a plurality of bacteriophage genomes. The plurality of bacteriophage genomes may be obtained from naturally occurring bacteriophage or genetically engineered bacteriophage. In some embodiments, each coded/labeled vial containing a plurality of bacteriophage genomes corresponds to a different bacteriophage type. In other embodiments, each coded/labeled vial containing a plurality of bacteriophage genomes corresponds to the same bacteriophage type. In some embodiments, each phage vial is assigned a unique code that identifies the bacteriophage in the phage vial, or the types of bacteria that the bacteriophage strain infects. The unique code can be encoded by a machine discernible pattern, such as a bar code, a QR code, an alphanumeric string, or any other pattern that can be discerned by a reader. Each unique code may be shown as, for example, a bar code sticker on a vial or container storing a corresponding phage sample. In some embodiments, the kit is stored under conditions that permit the preservation of the bacteriophage genomes for extended periods, such as under bacteriophage-specific, controlled temperature, moisture, and pH conditions.

[0112] The kit may further comprise one or more of: wash buffers and/or reagents, hybridization buffers and/or reagents, labeling buffers and/or reagents, and detection means. The buffers and/or reagents are usually optimized for the particular detection technique for which the kit is

intended. Protocols for using these buffers and reagents for performing different steps of the procedure may also be included in the kit. Further optional components of the kits may include expression media for reporter gene products disclosed herein, such as a medium containing nutrients and cofactors for bioluminescence, devices such as a lamp configured to illuminate at specific wavelengths of light to detect biofluorescence, and devices for measuring the extent of expression of the reporter genes, such as a photometer or photodetector.

[0113] In some embodiments, the kits further comprise positive control heterologous nucleic acid sequences to correct for any variability in the CFBS systems between experimental runs. The kits may also comprise instructions for use, software for automated analysis, containers, packages such as packaging intended for commercial sale and the like.

EXAMPLES

[0114] The present technology is further illustrated by the following Examples, which should not be construed as limiting in any way.

Example 1: Materials and Methods

Reagents

[0115] The following CFES/CFBS components, buffer salts, and media supplements were purchased from Sigma-Aldrich (St. Louis, MO): ATP, CTP, GTP, *E. coli* total tRNA mixture (from strain MRE600), 5-Formyl-5,6,7,8-tetrahydrofolic acid (folinic acid), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES), nicotinamide adenine dinucleotide (NAD), cyclic adenosine monophosphate (cAMP), coenzyme A (CoA), spermidine, 18 canonical amino acids excluding glutamate and leucine, polyethylene glycol 8000 (PEG-8000), maltodextrin, sodium oxalate, 1,4-dithiothreitol (DTT), potassium glutamate (K(glu)), magnesium glutamate (Mg(glu)₂), magnesium sulfate, calcium chloride, DNase I, lysozyme, and proteinase K. Leucine, phosphoenolpyruvate (PEP), and agar were purchased from Fisher Scientific (Hampton, NH) and UTP from Alfa Aesar (Haverhill, MA). RecBCD nuclease inhibitor GamS,

dNTPs, and all plasmid cloning reagents were purchased from New England Biolabs (NEB, Ipswich, MA). Tryptone, yeast extract, sodium chloride, potassium phosphate monobasic, potassium phosphate dibasic, agarose were purchased from VWR (Radnor, PA). TURBO DNA-free™ Kit and SuperScript™ II Reverse Transcriptase were purchased from Invitrogen (Waltham, MA). All statistical analysis was performed using GraphPad Prism 9.

Bacterial Strains and Growth Conditions

[0116] *E. coli* DH5α [genotype F[−]Φ80lacZΔM15Δ (lacZYA-argF) U169 recA1 endA1 hsdR17(r_K[−], m_K⁺)phoA supE44 thi-1 gyrA96 relA1 λ[−]] was used for plasmid cloning (NEB, Ipswich, MA). *E. coli* BL21 ATCC BAA-1025 [genotype F[−]ompT lon hsdS_B(r_B[−] m_B[−])gal dem [malB⁺]_{K-12}(λ^S)] was used for phage propagation, in vivo phage fitness and quantification assays, and as a source of cell extract for cell-free bacteriophage synthesis (CFBS) (American Tissue Culture Collection, Manassas, VA). The *E. coli* cells were grown at 37° C. in Luria-Bertani (LB) media (10 g/L of tryptone, 5 g/L of yeast extract, 10 g/L of sodium chloride in Milli-Q water). For T7 phage experiments, ΦLB (LB+2.5 mM MgSO₄ and 2.5 mM CaCl₂) was used. Solid media incorporated 1.500 agarose. All *E. coli* and phage strains used in this study are listed in Table A.

TABLE A

<i>E. coli</i> and phage strains		
Strain	Relevant genotype	Source
<i>E. coli</i> DH5α	F [−] Φ80lacZΔM15Δ(lacZYA-argF) U169 recA1 endA1 hsdR17(r _K [−] , m _K ⁺)phoA supE44 thi-1 gyrA96 relA1 λ [−]	NEB C2987I
<i>E. coli</i> BL21	F [−] ompT lon hsdS _B (r _B [−] m _B [−])gal dem [malB ⁺] _{K-12} (λ ^S)	ATCC BAA-1025
<i>E. coli</i> bacteriophage T7	N/a	ATCC BAA-1025-B2

[0117] All plasmids used in this study are listed in Table B.

TABLE B

Plasmids		
Plasmid	Relevant features	Reference or source
pJL1-sfGFP	T7 promoter-T7 gene 10 RBS-sfGFP-T7 terminator kanR (kanamycin resistance) ColE1 ori	Addgene # 69496
pBAD-sfGFP	araBAD promoter-T7gene 10 RBS-sfGFP-rmB1 T1 terminator ampR (ampicillin resistance) ColE1 ori	Addgene #54519
pCRJ001	rhaB promoter-RBS-dFnCas12a (aka dFnCpf1)-rmB1 terminator; J23119 promoter-FnCas12a sgRNA scaffold-LacZα cat (chloramphenicol resistance) p15A ori	This study
pCRJ001-g001	pCRJ001 derivative expressing sgRNA targeting trxA promoter	This study
pCRJ001-g002	pCRJ001 derivative expressing sgRNA targeting trxA CDS	This study
pCRJ001-g003	pCRJ001 derivative expressing sgRNA targeting nusG promoter	This study
pCRJ001-g004	pCRJ001 derivative expressing sgRNA targeting nusG CDS	This study

TABLE B-continued

Plasmids		
Plasmid	Relevant features	Reference or source
pCRJ001-g005	pCRJ001 derivative expressing sgRNA targeting pgk promoter	This study
pCRJ001-g006	pCRJ001 derivative expressing sgRNA targeting pgk CDS	This study
pCRJ001-g007	pCRJ001 derivative expressing sgRNA targeting subH promoter	This study
pCRJ001-g008	pCRJ001 derivative expressing sgRNA targeting subH CDS	This study
pCRJ001-g009	pCRJ001 derivative expressing sgRNA targeting eno promoter	This study
pCRJ001-g010	pCRJ001 derivative expressing sgRNA targeting eno CDS	This study
pCRJ001-g011	pCRJ001 derivative expressing sgRNA targeting mukB promoter	This study
pCRJ001-g012	pCRJ001 derivative expressing sgRNA targeting mukB CDS	This study
pCRJ001-g013	pCRJ001 derivative expressing sgRNA targeting hemL promoter	This study
pCRJ001-g014	pCRJ001 derivative expressing sgRNA targeting hemL CDS	This study
pCRJ001-g015	pCRJ001 derivative expressing sgRNA targeting cyaR promoter	This study
pCRJ001-g016	pCRJ001 derivative expressing sgRNA targeting cyaR CDS	This study
pCRJ001-g017	pCRJ001 derivative expressing sgRNA targeting lexA promoter	This study
pCRJ001-g018	pCRJ001 derivative expressing sgRNA targeting lexA CDS	This study
pCRJ001-g019	pCRJ001 derivative expressing sgRNA targeting infC promoter	This study
pCRJ001-g020	pCRJ001 derivative expressing sgRNA targeting infC CDS	This study
pCRJ001-g021	pCRJ001 derivative expressing sgRNA targeting recC promoter	This study
pCRJ001-g022	pCRJ001 derivative expressing sgRNA targeting recC CDS	This study
pCRJ001-g023	pCRJ001 derivative expressing sgRNA targeting dgt promoter	This study
pCRJ001-g024	pCRJ001 derivative expressing sgRNA targeting dgt CDS	This study
pCRJ001-g025	pCRJ001 derivative expressing sgRNA targeting udk promoter	This study
pCRJ001-g026	pCRJ001 derivative expressing sgRNA targeting udk CDS	This study
pCRJ001-g027	pCRJ001 derivative expressing sgRNA targeting rna promoter	This study
pCRJ001-g028	pCRJ001 derivative expressing sgRNA targeting rna CDS	This study
pCRJ001-g030	pCRJ001 derivative expressing sgRNA targeting Φ T7 gene 3 CDS	This study
pCRJ001-g031	pCRJ001 derivative expressing sgRNA targeting oxyS promoter	This study
pCRJ001-g032	pCRJ001 derivative expressing sgRNA targeting oxyS CDS	This study
pCRJ001-g033	pCRJ001 derivative expressing sgRNA targeting rne promoter	This study
pCRJ001-g034	pCRJ001 derivative expressing sgRNA targeting rne CDS	This study
pCRJ001-g035	pCRJ001 derivative expressing sgRNA targeting trxB promoter	This study
pCRJ001-g036	pCRJ001 derivative expressing sgRNA targeting trxB CDS	This study
pCRJ001-g037	pCRJ001 derivative expressing non-targeting control sgRNA	This study
pBAD-trxA	pBAD derivative for trxA overexpression	This study
pBAD-trxB	pBAD derivative for trxB overexpression	This study
pBAD-mukB	pBAD derivative for mukB overexpression	This study
pBAD-infC	pBAD derivative for infC overexpression	This study
pBAD-me	pBAD derivative for rne overexpression	This study
pBAD-oxyS	pBAD derivative for oxyS overexpression	This study

TABLE B-continued

Plasmids		
Plasmid	Relevant features	Reference or source
pBAD-cyaR	pBAD derivative for cyaR overexpression	This study
pBAD-dgt	pBAD derivative for dgt overexpression	This study

sgRNA = single-guide RNA

CDS = coding sequence downstream of transcription start sites

[0118] Primers and DNA oligos were purchased from integrated DNA Technologies (Coralville, IA) and are listed in FIG. 12.

Phage Propagation

[0119] *E. coli* bacteriophage T7 (ATCC BAA-1025-B2) was propagated using *E. coli* BL21 as the host strain. BL21 was grown for overnight (16 h) in 10 mL Φ LB at 37° C. inoculated with a single colony. The overnight culture was transferred to 100 mL prewarmed Φ LB in a 500 mL Erlenmeyer flask and incubated for 1 h shaking at 250 rpm to bring the culture to log phase. This culture was then inoculated with 100 μ L 10^8 PFU/mL T7 lysate and allowed to incubate until complete bacterial lysis was visually observed. The lysate was allowed to shake incubate a further 10 min after addition of 1 mL chloroform to lyse remaining BL21 cells. The lysate was transferred to 50 mL conical tubes and centrifuged at 10,000 $\times$ g for 5 min to pellet cell debris and separate chloroform from the mixture. Supernatants were recovered, 0.22 μ m sterile-filtered, then placed on ice for 1 h. Phage were concentrated by PEG-precipitation (Jakociune, D.; Moodley, A., A Rapid Bacteriophage DNA Extraction Method. *Methods Protoc* 2018, 1 (3)). Briefly, 0.22 μ m sterile-filtered PEG-8000/NaCl solution was mixed with lysate supernatants to final concentrations of 4% (w/v) and 0.5 M respectively followed by overnight incubation at 4° C. to encourage precipitation. Phage were pelleted by centrifugation at 10,000 $\times$ g at 4° C. for 30 min. Supernatants were carefully decanted and discarded. Following a second centrifugation at 6000 $\times$ g at 4° C. for 1 min, residual PEG/NaCl solution was aspirated by pipette. Phage pellets were resuspended and consolidated in minimal SM buffer (50 mM Tris-Cl pH 7.5, 100 mM NaCl, 8 mM MgSO₄). Phage stocks typically achieved titers of 10^{12} - 10^{13} PFU/mL and were stored at 4° C.

Phage Enumeration

[0120] T7 titers were determined using the standard double-layer agar (DLA) plaque assay using BL21 as the host strain (Liu, Y.; Huang, H.; Wang, H.; Zhang, Y., A novel approach for T7 bacteriophage genome integration of exogenous DNA. *J Biol Eng* 2020, 14, 2). Overnight cultures of BL21 in Φ LB were diluted 1:10 in fresh prewarmed Φ LB then incubated for 1 h at 37° C. and 250 rpm to bring cultures to log-phase. 10 μ L of log-phase BL21 was mixed with 3 μ L of appropriate dilutions of T7 sample, incubated for 5 min, then mixed with 1 mL of 50° C. 0.7% SM agarose (SM agarose+0.7% agarose (w/v)) and poured over 37° C. pre-warmed 60 $\times$ 15 mm Φ LB agar plates. The SM agarose was allowed to solidify at room temperature for 15 min, then plates were incubated at 37° C. for 1 h. Plates were then moved to a benchtop to incubate at room temperature overnight. T7 titers were calculated from plaque counts.

Alternatively, titers were determined using DLA spot tests. 100 μ L log-phase BL21 was mixed with 4 mL 50° C. 0.7% SM agarose then poured over 37° C. pre-warmed 100 $\times$ 15 mm Φ LB agar plates and allowed to solidify at room temperature for 15 min. Phage samples diluted in SM buffer were applied to solidified plates spotting with 3 μ L sample. Spots were allowed to dry completely at room temp (~10 min) then plates were incubated at 37° C. for 1 h. Finally, plates were incubated at room temperature overnight. T7 titers were calculated from the average of triplicate spots for each sample. Titers given in plaque forming units/mL (PFU/mL) were calculated from plaque counts using the following formula:

$$\text{titer} = \frac{\# \text{ plaques}}{3 \mu\text{L}} \cdot \frac{1000 \mu\text{L}}{1 \text{ mL}} \cdot \text{dilution factor} \quad (1)$$

Rapid Phage Titer Estimation

[0121] A microtiter plate-based assay was used to determine approximate phage titer to inform appropriate phage sample dilutions for plaque assays using a Spectramax iD5 plate reader. *E. coli* BL21 carrying pJL1-sfGFP (T7 promoter-RBS-sfGFP-T7 terminator) was grown overnight at 37° C. at 250 rpm in LB+35 μ g/mL kanamycin (Φ LB/kan). The overnight culture was sub-cultured 1:10 in fresh pre-warmed Φ LB+kan and allowed to incubate for a further 1 h to bring cell into log-phase. The culture was then diluted to OD₆₀₀=0.3 and transferred to a Corning® black/clear-bottom 3631 non-treated microplate to 200 μ L per well. The bacteria were then inoculated with 2 μ L of phage samples and infection kinetics tracked via OD₆₀₀ and sfGFP fluorescence (485 nm excitation/515 emission) on a Molecular Devices ID5 plate reader. In vivo T7-mediated T7 RNA polymerase expression drives sfGFP expression off of the reporter plasmid. Assay plates were incubated at 37° C. and “medium” speed orbital shaking between reads. Approximate T7 phage titers were calculated using a linear standard curve of phage known phage titers (1-10 log PFU/mL) vs time to V_{max} for relative fluorescence intensity with earlier V_{max} onset correlated with higher phage titers. Triplicate T7 standards, Φ LB, and bacteria without phage were included as controls in each assay plate. Limit-of-detection was 10 PFU/mL detectable within 2 hours of inoculation.

Phage Genome Extraction

[0122] T7 genomes were extracted using Qiagen DNeasy Blood and Tissue Kits (Hilden, Germany) as previously described (Jakociune, D. et al., supra). Briefly, residual bacterial nucleic acids were removed from high titer lysates by 1 h incubation with nucleases at 37° C.: 450 μ L lysate, 50

μL 10× nuclease buffer (100 mM Tris-Cl pH 7.5, 25 mM MgCl₂, 1 mM CaCl₂), 1 μL 1 mg/mL DNase I (Sigma-Aldrich, St. Louis, MO), and 1 μL 10 mg/mL RNase A (ThermoFisher Scientific, Roskilde, Denmark). Nucleases were inactivated by addition of 20 μL 0.5 M ethylenediaminetetraacetic acid (EDTA) pH 8.0 and incubation at 70° C. for 10 min. Then, capsids were degraded by the addition of 1.25 μL 20 mg/mL proteinase K and incubation at 56° C. for 1.5 h with gentle mixing by inversion every 30 min. Liberated T7 genomes were then purified following Qiagen DNeasy Kit Protocols. Typical yields for a 10¹² PFU/mL lysate were 100 μL of ~500-2000 ng/L genomic DNA (gDNA). DNA integrity was confirmed by gel electrophoresis (not shown).

Plasmid Purification and Transformation

[0123] Plasmids were purified using Qiagen kits as described by manufacturer. For cloning, plasmids were purified using Qiaprep® Spin Miniprep Kits. Plasmids used in cell-free experiments were purified using Plasmid Plus Midi Kits. CRISPRi and overexpression constructs were transformed into NEB DH5α by heat-shock. Competent cells (10 μL) were mixed with assembled vectors (1 μL) and left on ice for 30 min followed by a 30s shock at 42° C. the 5 min recovery on ice. Pre-warmed SOC broth (20 g/L tryptone, 5 g/L yeast extract, 10 mM NaCl, 2.5 mM KCl, 10 mM MgCl₂, 10 mM MgSO₄ and 20 mM glucose) was added to each transformation tube which were then incubated at 37° C. at 250 rpm for 1 h then plated on appropriate selective media. BL21 was transformed by electroporation. Electrocompetent BL21 was prepared as follows. BL21 was grown overnight in LB, then sub-cultured 1:100 in 100 mL pre-warmed LB in a 500 mL Erlenmeyer flask and incubated at 37° C. at 250 rpm. When, the culture reached OD₆₀₀=0.4-0.6, cells were harvested by centrifugation at 10,000×g for 5 min at 4° C. Recovered pellets were washed three times with ice-cold sterile 10% glycerol, then resuspended in 10% to achieve 50× concentration. Electrocompetent cells were aliquoted and stored at -80° C. until use. Thawed 50 μL aliquots were mixed with 50-100 ng of desired plasmid (maximum 5 μL) and kept on ice for 10 min. The cell/plasmid mixtures were transferred to chilled 1 mm gap cuvettes and exposed to a 1.8 kV pulse for 5 ms followed by immediate addition of 1 mL pre-warmed SOC. Transformants were incubated at 37° C. at 250 rpm for 1 h, then plated on selective media. Recovered transformants were plated on LB agar supplemented with appropriate antibiotics (35 μg/mL kanamycin, 35 μg/mL chloramphenicol, or 100 μg/mL ampicillin) and incubated overnight at 37° C. Plasmids were validated by Sanger sequencing by Azenta (South Plainfield, NJ). Transformants bearing sequence confirmed plasmids were banked in glycerol stocks at -80° C.

Goldengate Cloning Protocol of CRISPRi Constructs

[0124] Gene targets were selected based only with experimentally confirmed promoter positions and transcription start sites. CRISPRi constructs were all prepared using pCRJ001 as the vector backbone by replacing a LacZα with sgRNA sequences targeting either promoters or coding sequences (CDS) as close to the start codon as possible. Where possible, stringent TTTV PAM-sites were used and less stringent TTV when the alternative was not available (Chen, P.; Zhou, J.; Wan, Y.; Liu, H.; Li, Y.; Liu, Z.; Wang,

H.; Lei, J.; Zhao, K.; Zhang, Y.; Wang, Y.; Zhang, X.; Yin, L., A Cas12a ortholog with stringent PAM recognition followed by low off-target editing rates for genome editing. *Genome Biol* 2020, 21 (1), 78). In 50 μL reactions, pCRJ001 (1.5 μg) was digested by BsaI-HF®v2 (60 U, NEB, Ipswich, MA) @37° C. for 2 h with pipet mixing every 30 min followed by heat inactivation at 80° C. for 30 min then held at 53° C. until the ligation step. sgRNA inserts were prepared by annealing complementary 20 bp DNA oligos with 4 bp overhangs complementary sticky ends on pCRJ001 BsaI-digests. In 10 μL reactions, 10 μM of each oligo were heated to 95° C. for 5 min, then stepped down to 4° C. at 0.1° C./s. Linearized pCRJ001 and annealed inserts were then ligated at 25° C. for 2 h: 1.5 μL linear vector (45 ng, 2.25 nM), 1 μL annealed oligos (300 ng, 2000 nM), and 2.5 μL 2× Instant Sticky-end Ligase Master Mix (NEB). Ligated CRISPRi vectors were transformed into DH5α by heat-shock. Transformants were plated on LB agar+35 μg/mL chloramphenicol (cm) and 40 μg/mL 5-bromo-4-chloro-3-indolyl-β-D-galactopyranoside (X-gal) (Messing, J.; Gronenborn, B.; Müller-Hill, B.; Hofschneider, P. H., Filamentous coliphage M13 as a cloning vehicle: insertion of a HindIII fragment of the lac regulatory region in M13 replicative form in vitro. *Proc Natl Acad Sci USA* 1977, 74 (9), 3642-3646). White colonies were selected for propagation in LB+cm and plasmids were purified. Assembly of desired constructs was confirmed by Sanger sequencing. Sequence confirmed CRISPRi vectors were then transformed into BL21 by electroporation. Validated DH5α and BL21 containing CRISPRi vectors were banked as glycerol stocks at -80° C. Preparation of pBAD Vectors by Gibson Assembly

[0125] Overexpression vectors were prepared by Gibson assembly as previously described (Gibson, D. G.; Young, L.; Chuang, R. Y.; Venter, J. C.; Hutchison, C. A., 3rd; Smith, H. O., Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 2009, 6 (5), 343-5). pBAD-sfGFP (Table B) was used as the backbone for arabinose inducible gene expression. pBAD-sfGFP was linearized by restriction digestion with XhoI and EcoRI (NEB) for 1 h @37° C. The linear vector was PCR amplified by high-fidelity NEB Q5® DNA polymerase to generate blunt-ended linear PCR products. Gene inserts were PCR amplified with Q5 using BL21 genomic DNA template liberated by boiling a colony in 100 μL nuclease-free water. PCR products were designed with 20 bp overhangs complementary to the 5' and 3' ends of the linearized vector to replace sfGFP while retaining regulatory elements (FIG. 12). Overexpression constructs were assembled in 5 μL reactions containing 2.5 μL NEBuilder® HiFi DNA Assembly 2× Master Mix (E2621L), 0.1 pmol gene inserts, and 0.05 pmol linear vectors incubated at 50° C. for 1 h. Assembled constructs were transformed into DH5α by heat-shock and miniprep for sequence confirmation. Correctly assembled overexpression constructs were finally transformed into BL21 by electroporation and banked as glycerol stock at -80° C.

Gene Expression Titration Experiments

[0126] Gene repression and overexpression assay conditions were optimized for inducer concentrations and induction timing. For CRISPRi experiments, trxA repression was evaluated by mRNA level by RT-qPCR and T7 efficiency of plating after rhamnose induction at 0.02, 0.1, 0.2, 2% (w/v) rhamnose at 2, 4, 8 and 24 h post-induction. Overexpression

induction conditions were determined by overexpression of sfGFP after 4 h induction with 0.002, 0.01, 0.02, 0.1, 0.2% (w/v) arabinose. sfGFP levels were measured relative fluorescence intensity.

RNA Extractions and RT-qPCR

[0127] RNA extractions were performed using Qiagen RNeasy Kit with enzymatic lysis and an additional DNase treatment as previously described (Hay, M.; Li, Y. M.; Ma, Y., RNA extraction of *Escherichia coli* grown in Lysogeny Broth for use in RT-qPCR. *JEMI Methods* 2017, 1, 1-6). Log-phase cells (0.5 mL, OD600=0.4-0.6) were pelleted and resuspended in 200 μ L lysis buffer pH 8.5 (15 mg/mL lysozyme, 1 mg/mL proteinase K, 30 mM Tris-Cl, 1 mM EDTA) then incubated at room temperature for 10 min with gentle vortexing every 2 min. The lysate was processed using the standard Qiagen RNeasy Kit protocol including on-column DNase I treatment and eluted using 30 μ L nuclease-free water. A second DNase treatment was performed using a 30 μ L reaction volume containing 26 μ L RNA extract, 3 μ L 10 $\times$ TURBOTM DNase Buffer and 1 μ L TURBOTM DNase incubated at 37 $^{\circ}$ C. for 30 min. DNase activity was stopped using 3 μ L DNase Inactivation reagent. RNA was quantified using a DeNovix DS-11 microvolume spectrophotometer. cDNA was generated using SuperScriptTM II Reverse Transcriptase with RNA normalized to 50 ng per reaction. RT-qPCR was performed in 20 μ L reaction containing 10 μ L iTaq Universal SYBR $^{\circ}$ Green Supermix (2 $\times$) (Bio-Rad, Hercules, CA), 500 nM forward and reverse primers (FIG. 12), and 1 ng cDNA. Relative mRNA expression was calculated using the $2^{-\Delta\Delta C_t}$ method (Livak, K. J.; Schmittgen, T. D., Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001, 25 (4), 402-8). Expression of *cysG* *hcaT*/*idnT* was averaged to be used as reference genes due to their stable expression in the context of induced protein expression in *E. coli* BL21 (Zhou, K.; Zhou, L.; Lim, Q.; Zou, R.; Stephanopoulos, G.; Too, H. P., Novel reference genes for quantifying transcriptional responses of *Escherichia coli* to protein overexpression by quantitative PCR. *BMC Mol Biol* 2011, 12, 18).

Preparation of Gene-Knockdown or -Overexpressing Cultures for Phage Fitness Assays

[0128] *E. coli* BL21 carrying CRISPRi or overexpression plasmids were streaked from glycerol stocks onto Φ LB+50 μ g/mL chloramphenicol (Φ LB/cm) and Φ LB+100 μ g/mL ampicillin (Φ LB/amp) agar respectively to generate isolated colonies. A single colony per construct was picked to inoculate overnight selective broth cultures incubated at 37 $^{\circ}$ C. at 250 rpm. These cultures were then sub-cultured 1:10 into fresh 37 $^{\circ}$ C. broth and incubated at 37 $^{\circ}$ C. at 250 rpm for 1 h to bring cells to log-phase. Gene repression was induced by 1% inoculation of 1.2 mL of 37 $^{\circ}$ C. Φ LB/cm+2% rhamnose with log-phase cultures. Gene overexpression was induced by 1% inoculation of 1.2 mL of 37 $^{\circ}$ C. Φ LB/amp+0.2% arabinose with log-phase cultures. Induced cultures were incubated in Corning Costar $^{\circ}$ 3738 Not Treated 24-well plates at 37 $^{\circ}$ C. at 250 rpm for at least 4 h until OD600 reached 0.4-0.6, then placed on ice for 15 min. Induced and non-induced controls were centrifuged at 6000 $\times$ g at 4 $^{\circ}$ C. for 5 min, then resuspended in fresh media to OD=0.3 and kept on ice until phage fitness assays (lysis

timing and efficiency of plating). Non-targeting (NT)-controls were included in each CRISPRi experiment. Overexpression of sfGFP was included as a negative control in overexpression experiments. Biological replicates (n=3) were performed on different days with cultures started from unique isolated colonies. Antibiotics and inducers were 0.22 μ m filter sterilized and added to broth cultures immediately before use.

Kinetic Assay of Lysis

[0129] Induced and non-induced control cultures were loaded into Corning Costar $^{\circ}$ 3631 black/clear-bottom 96-well plates to 200 μ L and incubated at 37 $^{\circ}$ C. for 15 min to pre-warm. The cultures were then inoculated with 20 μ L T7 diluted in LB to a multiplicity of infection (MOI) of 3 using a multichannel pipette. Φ LB was added to phage-free controls. Lysis/growth kinetics were recorded by plate reader by measuring OD600 very 3 min with orbital shaking between reads. Lysis time was defined as the time post-infection at which a decrease in OD600 was first recorded. Mean lysis time ($T_{50\%}$) was calculated as the midpoint of the lysis curve fit to a cumulative normal distribution (Heineman, R. H.; Bull, J. J., Testing optimality with experimental evolution: lysis time in a bacteriophage. *Evolution* 2007, 61 (7), 1695-709; Heineman, R. Lysis time, optimality, and the genetics of evolution in a T7 phage model system. University of Texas at Austin, 2007). Effect of gene repression or overexpression on lysis times and mean lysis times were evaluated by paired two-tailed t-test on averages from biological triplicates (p<0.05).

Efficiency of Plating (EOP) Assay

[0130] Fifty microliters of knockdown and overexpressing BL21 strains at OD600=0.3 were infected with T7 at an MOI=0.0001 using a multichannel pipette and allowed to incubate in Greiner Bio-One #655201 polypropylene flat-bottom 96-well microplates for 30 min at 37 $^{\circ}$ C. and 250 rpm. After incubation, 100 μ L of ice-cold chloroform was added to each well to halt phage replication and lyse remaining intact cells. Samples were transferred to microcentrifuge tubes containing 500 μ L ice-cold SM buffer and centrifuged at 17,000 $\times$ g for 1 min for phase separation. The supernatants were further diluted in SM buffer as necessary. Phage titers were counted by spot tests on wild-type BL21 overlay plates from the average of three 3 μ L technical replicate spots. Non-induced knockdown and overexpressing strains were included as controls. For each strain, efficiency of plating (EOP) was calculated using the following formula:

$$EOP = \frac{\text{mean PFU}_{\text{induced}}}{\text{mean PFU}_{\text{non-induced}}} \quad (2)$$

Preparation of Cell-Extracts for CFES and CFBS

[0131] Cell-extracts were prepared based on Kwon et al. (2019) with modification. Each knockdown and overexpression strains was grown overnight in 1 mL LB, supplemented with appropriate antibiotics, inoculated by a single isolated colony. Overnight cultures were diluted 1:10 in prewarmed selective LB and incubated for 1 h at 37 $^{\circ}$ C. to bring the cultures to log-phase. Next, cultures were diluted 1:100 in

pre-warmed 2×YTP media (16 g/L of tryptone, 10 g/L of yeast extract, 5 g/L of sodium chloride, 7 g/L of potassium phosphate dibasic, 3 g/L of potassium phosphate monobasic, pH 7.2) plus appropriate antibiotics in 500 mL baffled-flasks and incubated at 37° C. at 250 rpm and growth monitored by OD600. When OD600 reached ~0.2, L-rhamnose was added to knockdown cultures and L-arabinose to overexpression cultures to 2% and 0.2% respectively. When OD600 reached 2.0-3.0, flasks were immediately placed on ice and gently mixed to quickly bring down temperature. Samples were kept cold from this step onward. After 15 min on ice, each culture was centrifuge at 10,000×g for 5 min at 4° C. The supernatants were removed and pellets washed three times with ice-cold S30B buffer (10 mM Tris-Cl, pH 8.2, 14 mM magnesium glutamate, 60 mM potassium glutamate, 2 mM DTT). Pellet wet-mass (g) was recorded followed by flash-freezing in liquid nitrogen and storage at -80° C. overnight. Frozen pellets were thawed slowly in ice-water then resuspended in S30B (0.8 mL per g wet-mass) and transferred to 1.5 mL microcentrifuge tubes with final volumes 500-1000 µL. Cell slurries were lysed in an ice-water bath using the Qsonica Q125 Sonicator with 1/8" probe set to 50% amplitude with 10 s on/off pulses to avoid overheating samples. Target total energy input was calculated using the following formula (Kim, J.; Copeland, C. E.; Seki, K.; Vogeli, B.; Kwon, Y. C., Tuning the Cell-Free Protein Synthesis System for Biomanufacturing of Monomeric Human Filaggrin. *Front Bioeng Biotechnol* 2020, 8, 590341):

$$\text{Energy (J)} = \frac{\text{sample volume (}\mu\text{L)} - 33.6}{1.8} \quad (3)$$

[0132] DTT was added to each lysate for a final 3 mM concentration. Cell-lysates were centrifuged at 12,000×g for 10 min at 4° C. Supernatants from each strain were consolidated, then incubated at 37° C. shaking for 80 min for a runoff reaction followed by another centrifugation at 10,000×g for 10 min at 4° C. Finally, these clarified supernatants were aliquoted and flash-frozen with liquid nitrogen then stored at -80° C. Total protein was measured using Pierce BCA Protein Assay Kit (ThermoFisher) using bovine serum albumin standards. Presence of residual bacteria was checked by spotting clarified lysates on LB agar and incubating overnight at 37° C. Inducers were dissolved in 2×YTP and 0.22 µm filter-sterilized. Biological duplicates were prepared for each selected knockdown and overexpression strain. Extracts based on wild-type BL21 absent plasmids were prepared as baseline references for protein and phage synthesis.

CFBS TXTL Protein Synthesis Reporter Assay (T7 gDNA+ pJL1-sfGFP Cascade)

[0133] Activity of cell extracts for T7 gDNA-dependent transcription and protein synthesis was evaluated using sfGFP synthesis as a reporter. Protein synthesis was performed in 15 µL reactions incubated at 30° C. in 1.5 mL tubes with cell extracts occupying 33% volume and reaction buffer and DNA templates the remaining 67%. Reactions contained 57 mM HEPES, pH 8, 130 mM K(glu), 12 mM Mg(glu)₂, 0.4 mM NAD, 0.27 mM CoA, 0.75 mM cAMP, 2 mM spermidine, 1 mM DTT, 1.5 mM ATP and GTP, 0.9

mM CTP and UTP, 0.2 mg/mL *E. coli* tRNA, 0.068 mM folinic acid, 2 mM of each canonical amino acid except glutamate, 33 mM PEP, 12.66 mg/mL maltodextrin, 0.5 nM T7 gDNA, and 5 nM pJL1-sfGFP. Reaction components were mixed together and with DNA added last then kept on ice for 5 min prior to incubation at 30° C. Five microliter samples were taken at 4 h and 20 h, diluted in 20 µL 50 mM HEPES, pH 8, and sfGFP fluorescence measured as described above in Greiner Bio-One #781209 black flat-bottom 384-well plates. All reactions were run in triplicate.

CFBS for Phage Synthesis

[0134] Bacteriophage synthesis was performed using reaction setup described above with modifications: pJL1-sfGFP was omitted and reaction buffer was supplemented with 3.5% (w/v) PEG-8000 and 0.5 mM dNTPs. At 4 h and 20 h, 3 µL samples were taken and diluted in 30 µL SM buffer. Approximate T7 titers calculated by the rapid titer estimation assay described above. More precise titers were calculated from plaque counts using the DLA method.

Example 2: CRISPRi-Mediated *trxA* Knockdown Influences T7 Fitness

[0135] To demonstrate the efficacy of this system, a CRISPRi vector with a crRNA targeting the *trxA* promoter (pCRJ001_g001, inserted via Golden Gate assembly) was transformed into T7 host *E. coli* BL214². The *trxA* gene encodes thioredoxin 1, a non-essential *E. coli* protein involved with cytoplasmic redox homeostasis and an essential host-factor for T7 genome replication^{33, 43}. In *trxA*-deficient mutants, T7 will still induce host lysis and produce progeny virions, but will not replicate, making *trxA* an ideal target to demonstrate the effects of gene knockdown on T7 fitness⁴⁴. The functionality of the CRISPRi system was evaluated using RT-qPCR and direct plaque assays to enumerate T7 progeny resulting from infection of host *E. coli* BL21 with a *trxA* knockdown (*trxA*-KD) background. Gene repression was induced in log-phase cells with 2% L-rhamnose (w/v) for 4 h at which point cells were harvested for RNA extraction and T7 infection assays. RT-qPCR revealed 90±2.4% (SD, n=3) drop in *trxA* mRNA compared to the non-induced control carrying the same CRISPRi vector (FIG. 2B). Rhamnose induction increased dFnCas12a mRNA by over 70-fold compared to un-induced controls (FIG. 2A). T7 infection in a *trxA*-KD background resulted in an efficiency of plating (EOP) of 12±7% relative to the uninduced control (FIG. 2C). The *trxA*-KD EOP was also significantly lower compared to that of the non-targeting (NT) control strain (vector pCRJ001_g037) (p<0.01, Welch's two-tailed t-test).

Example 3: CRISPRi Screen of Potential T7 Fitness Effector Genes

[0136] Having confirmed our inducible CRISPRi system could be used to influence T7 fitness, additional CRISPRi vectors were prepared to target each of the potential T7 effectors selected in this study (Table C).

TABLE C

CRISPRi/Overexpression gene targets.					
Target Gene	Category	Function	Essential (<i>E. coli</i>) ^a	Essential (Φ T7) ^b	Ref.
infC	protein	translation initiation factor IF-3	Yes	—	33
pgk	energy	phosphoglycerate kinase	Yes	—	33
subH (ssyA)	energy	inositol monophosphatase	Yes	—	33
hemL (gsa)	protein	glutamate-1-semialdehyde aminotransferase	Yes	—	33
eno	energy/nucleic acid	enolase in glycolysis/gluconeogenesis/mRNA degradosome	Yes	—	33
mukB	protein/cell-division	degradosome/chromosomal DNA condensation	Yes	—	33
lexA	transcription factor	SOS response regulator	Yes	—	33
rne	nucleic acid	RNase E	Yes	No	30, 33, 34
nusG	transcription factor	transcription termination factor	Yes	No	33, 35, 36
trx	redox regulator	oxidized thioredoxin	No	Yes	30, 33
T7gp3	nucleic acid	Φ T7 DNA replication	No	Yes	37
dgt	energy	deoxy guanosinetriphosphate (dGTP) triphosphohydrolase	No	No	30, 33
udk	nucleic acid	uridine kinase/cytidine kinase	No	No	30, 33
recC	nucleic acid	Exodeoxyribonuclease V (RecBCD) subunit	No	No	30, 33, 38
cyaR (ryeE)	transcription factor	CyaR small RNA, oxidative stress regulator	No	—	39
rna	nucleic acid	Rnase I, cleaves phosphodiester bond between any two nucleotides	No	—	33
oxyS	transcription factor	OxyS small RNA, oxidative stress regulator	No	No	40
trxB	redox regulator	thioredoxin reductase	No	No	30, 33, 41

^aChromosomal deletion results in non-viable or non-replicative cells.^bHost/phage chromosomal deletion causes non-productive phage infections.

[0137] Two vectors were designed for each gene with one crRNA directed towards promoters and one towards coding sequences (CDS) as near to transcription start sites (TSS) as possible as available PAM sites allowed. The exception was T7 exonuclease gene 3 (T7gp3) for which only the CDS was targeted. Each CRISPRi vector was then transformed into BL21 for experiments evaluating the impact of gene knock-downs on T7 fitness.

[0138] Lysis time courses and EOP assays were used to compare relative T7 fitness. Here, lysis onset time is defined as the time post-T7 infection at which optical density begins to decrease as previously described²⁶. Log-phase BL21 carrying CRISPRi vectors (OD=0.3) were infected with T7 to a multiplicity of infection (MOI)=3 then OD kinetics were tracked (FIGS. 3A and 8). Overall, in cultures without CRISPRi induction, lysis onset was ~25 min, later than the typical T7 lysis onset of ~15 min²⁶. We attribute this to the metabolic burden of plasmid maintenance and chloramphenicol in selective media. Initial screens found delayed lysis onset in induced CRISPRi strains with crRNAs targeting the dgt CDS (g024) (p<0.01) and both the eno promoter (g009) and CDS (g010) (p<0.05) relative to the same strains without CRISPRi induction FIG. 3B.

[0139] While only three CRISPRi targets resulted in significant changes in lysis onset time, examination of lysis curves revealed differences in the lysis kinetic profiles for many targeted genes as indicated by the widened curve profiles when CRISPRi was induced (FIG. 3A). Based on this observation, we also compared mean lysis times, which represents the time at which 50% of cells are lysed⁴⁵. In addition to dgt and eno, repressors trxA promoter (g001), mukB promoter and CDS (g011 and g012 respectively), and infC CDS (g020) were also found to have significant delays

in mean lysis time. This suggests that each of these effector candidates may warrant further investigation.

[0140] While indicative of altered T7 life cycle kinetics, lysis onset and mean lysis timing are not necessarily indicative of effects on number of progeny phage resulting from infection²⁸. Therefore, the efficiency of plating (EOP) in knockdown strains was also evaluated. Log-phase CRISPRi *E. coli* BL21 stains (OD=0.3) were infected at MOI=0.0001 (1 virion per 10,000 cells) for 30 min, then phage replication was halted with chloroform and progeny phage enumerated by plaque assay using wild-type BL21 as host. EOP assays found that most CRISPRi constructs had a significant negative impact on T7 titer with the exceptions of those targeting *lexA*, *trxB*, *rna*, *pgk*, *cyaR*, and *recC* (FIG. 4A). SOS response transcriptional regulator *lexA* expression is tightly limited under normal growth conditions, so its repression was not expected to elicit a strong effect on T7 fitness⁴⁶. Likewise, starvation response regulator *cyaR* repression did not affect T7 lysis nor EOP during growth on rich media⁴⁷. RNase I encoded by *rna* localized in the periplasm and is thus unlikely to interfere with T7 mRNA stability⁴⁸. Phosphoglycerate kinase (*pgk*) did not affect lysis timing or EOP, which is unexpected given the lysis delay and EOP drop induced by the enolase (*eno*) knockdown as both enzymes are part of the canonical glycolysis pathway⁴⁹.

[0141] *eno* was chosen as a CRISPRi target because of its roles in glycolysis, as part of the mRNA degradosome complex, and interactions between enolase and other phage. Notably, enolase associates with RNase E (*rne*), whose major role is degradation of mRNA. The T7 early gene gp0.7 inhibits RNase E activity early during infections to stabilize phage-derived mRNAs³⁴. There is evidence of protein-protein interactions between coliphage T1 and *E.*

coli enolase (significance of interaction unknown) and direct inhibition of enolase by *Bacillus subtilis* phage SPO1, suggesting *eno* may play a role in T7 fitness directly or indirectly^{34, 50, 51}. Because *me* inhibition is part of the T7 lifecycle, we expected *me* knockdown to have a neutral to positive effect on EOP. However, as with *eno*, *me* knockdown lowered EOP. This was perhaps due to impairment of RNase E's role in rRNA maturation, which could lead to reduced translational capacity. In silico modeling of infection kinetics suggests that translation rate is the primary bottleneck in T7 progeny assembly³¹. The hypothesis that translation rates influence T7 fitness is supported by our observations of lysis delay and EOP loss in translation initiation factor IF-3 (*infC*) (FIG. 4A).

[0142] Of the essential *E. coli* genes investigated, *mukB*, *subH*, *hemL*, and *nusG* repression caused significant loss of EOP (FIG. 4A). While each of these knockdown strains were viable under experimental conditions, they also experienced severe growth rate defects, which have been associated with diminished phage bursts³¹.

[0143] Knockdown of known antagonistic *E. coli* genes *udk* (uridine-cytidine kinase; uridine/cytidine+ATP→UMP/CMP+ADP+H⁺)⁵², and *dgt* (dGTPase; dGTP→dG+PPPi)⁵³ unexpectedly resulted in lower EOP (FIG. 4A). T7 encodes Udk and Dgt inhibitors (*gp0.7* and *gp1.2*, respectively), so we hypothesized that repressing these genes could increase T7 fitness. However, the interfering with *udk* only caused a small non-significant drop in EOP. The *dgt* promoter repressor (*dgt023*) caused a significantly lower EOP, perhaps due to perturbed ribo- and deoxyribonucleotide pool homeostasis, thereby preventing normal T7 activities²⁹.

[0144] CRISPRi-based knockdown of the small regulatory RNA *oxyS* also lowered EOP (FIG. 4A), which was consistent with our expectation. *OxyS* acts as a regulator of stress-response sigma factor *RpoS* (σ^S) expression and oxidative stress response⁵⁴. Previous studies demonstrate that $\Delta oxyS$ *E. coli* has higher expression of *RpoS* and *RpoS*-regulated genes relative to wild-type and that inducible *oxyS* expression results in *RpoS* suppression⁵⁴. Early T7 transcription is mediated primarily by *E. coli* RNA polymerase-sigma 70 complex ($E\sigma^{70}$), but can also be carried out using the alternate $E\sigma^S$ complex⁵⁵. During middle and late gene transcription, $E\sigma^{70}$ and $E\sigma^S$ are inhibited by T7 *gp2* and *gp5.7*, respectively, and transcription is taken over by T7 RNAP. T7 *gp2* mutants experience abortive infections associated with “aberrant” transcription (interrupted transcripts and atypical, terminator read-through) caused by competition between $E\sigma^{70}$ and T7 RNAP⁵⁶. We suspect that the loss of EOP in *oxyS*-knockdown strains may be due to incomplete $E\sigma^S$ inactivation due to greater background *RpoS* accumulation and competition between active $E\sigma^S$ and T7 RNAP.

[0145] Another interesting observation was the neutral impact of *trxB* (thioredoxin reductase) knockdown on EOP. In the *TrxA/TrxB* redox system, *TrxA* acts as a recyclable reducing agent with diverse roles in over 80 protein-protein interactions and transcriptional regulation of at least 26 genes⁵⁷. These complex interactions are implicated in most cellular processes including oxidative stress response, translation, energy transduction. *TrxB* recharges oxidized *TrxA* to restore the pool of reduced-form *TrxA*. Under normal conditions, balanced *TrxA* concentrations and *TrxB* activity maintains the majority of *TrxA* in its reduced form^{58, 59}. *TrxA* exists in its reduced state in the functional T7 DNA

polymerase holoenzyme and does not meaningfully interact with T7 DNA polymerase in its oxidized state⁴¹. *trxB* deletion causes cytosolic *TrxA* to exist solely in its oxidized form, which has been shown to lower EOP by 10⁻⁸-fold for coliphage *f1*, which also requires reduced form *TrxA*^{41, 60}. As discussed above, *trxA* knockdown lowered EOP, likely by depletion of free *TrxA* (FIGS. 2, 4A). We anticipated *trxB* knockdown would cause a similar effect by depleting reduced-form *TrxA*. The lack of impact on EOP in the *trxB*-KD strains suggests there is sufficient *TrxB* activity to supply adequate reduced-form *TrxA* to support T7 replication.

[0146] The *recC* CDS (*g022*) was the only target whose knockdown resulted in a significant positive effect on EOP (+21±3.1%). *RecC* was selected as a target because of its role in the double-stranded DNA repair RecBCD holoenzyme, which has exonuclease activity on linear DNA such as the T7 genome⁶¹. Deletion of *recB* or *recC* results in loss of nuclease activity³⁸. Exonuclease inhibitor *GamS* is typically included in CFES to protect linear DNA templates from RecBCD. Improved EOP here was consistent with recent work by Batista et al. (2022) showing that TXTL with *recB* or *recBCD* deletions improved CFES sfGFP yields for linear templates compared to wild-type TXTL¹⁵.

[0147] Inducer titration experiments were conducted on a subset of CRISPRi strains to determine if gene knockdown effects were tunable (FIG. 10A). The *trxA* knockdowns showed concentration dependence where higher L-rhamnose concentrations caused greater loss of EOP. Repression of *rna* had no impact on EOP at any concentration of rhamnose, consistent with our previous results (FIG. 4A). At lower rhamnose concentrations (0.02-0.1% w/v), *recC* repression lowered EOP in contrast to the EOP gains observed at higher concentrations (0.2-2% w/v).

[0148] In brief, we demonstrated a novel dFnCas12a-based single-plasmid CRISPRi system could be used to influence phage fitness in an inducible and tunable manner. The system can be used to knockdown essential and non-essential genes of host and phage as well as genes encoding sRNAs rather than proteins. The strength of interference with phage fitness depends on the strength of gene repression, so care must be taken when designing crRNAs as which provides tighter control, targeting promoters versus CDS, appears to be gene dependent. One approach could be to design additional crRNA constructs for each gene or to multiplex promoter and CDS targeting as our FnCas12a system is readily multiplexed by constructing vectors with consecutive guide RNA scaffold arrays⁶².

Example 4: Overexpression of Potential Phage Fitness Effectors

[0149] To complement the investigation of phage effectors by CRISPRi, we sought to explore modulating T7 fitness by effector overexpression. L-arabinose inducible pBAD vectors were constructed to overexpress a subset of the T7 effector targets. This subset was chosen from among the genes whose repression caused significant loss in EOP (FIG. 4A) under the hypothesis that if repression causes a negative impact on T7 fitness, overexpression may have a positive impact. Among these genes, only overexpression of sRNAs *oxyS* and *cyaR* increased T7 EOP (FDR $q < 0.01$, Welch's two-tailed t-test), *dgt*, *trxB*, and *trxA* decreased EOP, and *infC* and *me* had a neutral effect as did the control of overexpressed sfGFP (FIG. 4B). *mukB* overexpression,

which has also been implicated in acetate tolerance⁶³, also decreased EOP but was not significant. In contrast to the knockdown studies, none of these overexpression impacted lysis timing nor the profile of their lysis curves (data not shown).

[0150] We suspect that CyaR and OxyS may support T7 transcription by interfering with RpoS translation^{39, 54}. OxyS may also act to reduce oxidative stress on T7 by activating oxidative stress-response in an RpoS-independent manner^{64, 65}.

[0151] Huber et. al (1988) found that a Dgt overexpressing optA1 mutant *E. coli* maintained 50× higher dGTPase concentrations, which lowered dGTP pools 5-fold⁶⁶. Wild-type T7 is able to replicate in these dGTP depleted backgrounds, but foundational studies utilizing optA1 strains do not report impact on EOP⁶⁶⁻⁶⁸. Here, we again demonstrate that T7 does propagate in a Dgt overexpressing strain, but EOP is diminished.

[0152] Unexpectedly, TrxA and TrxB overexpression decreased EOP. We expected these genes to have a positive effect on T7 titer in contrast to the near abolition of T7 replication in trxA CRISPRi strain g001. It is unclear why this occurred, but overexpression of these genes may have interfered with redox balance or had other unpredictable effects given TrxA's many protein-protein and protein-DNA interactions^{57, 59}.

[0153] As with the CRISPRi constructs, our overexpression strains showed L-arabinose concentration dependence of EOP effects (FIG. 10B). Most notably, there is a negative relationship between L-arabinose concentration (0.002-0.2% w/v) and EOP in the pBAD-trxA strain. pBAD-oxyS and pBAD-cyaR only reach significantly increased EOP at 0.2% L-arabinose. Likewise, pBAD-dgt only significantly lower EOP at 0.2% L-arabinose. Meanwhile, pBAD-infC showed a steady but small EOP increase up to 0.02% L-arabinose, but dropped back to baseline at 0.2%. Importantly, T7 EOP was insensitive to induction of control pBAD-sfGFP at all L-arabinose concentrations.

Example 5: Effect of Gene Expression Background in TXTL Source on Cell-Free Expression Systems and Bacteriophage Synthesis

[0154] To determine if in vivo effectors of T7 fitness had the same qualitative effect in cell-free systems, CRISPRi and overexpression strains were selected to prepare transcription/translation machinery (TXTL) for cell-free protein express systems (CFES) and cell-free bacteriophage synthesis (CFBS) based on prior research indicating their usefulness in cell-free protein synthesis or positive effects on T7 fitness during in vivo studies. Exogenous IF-3 (infC) supplementation, recC deletion, and rna deletion each have been shown to improve CFES yields of GFP^{15, 48, 69}. oxyS- and cyaR-overexpression strains were chosen as positive in vivo effectors, Dgt-overexpression and trxA-knockdown strains as likely negative effectors, and trxA-overexpression for its strong negative effects. All lysates were harvested 4 h after induction with appropriate inducer and harvested in mid-log-phase with OD₆₀₀~2-3 yielding total protein yields ranging from 15-25 mg/mL. No growth defects were observed, but recC-KD strains had an extended lag-phase (~1 h) before entering log-phase. Lysates were produced from each strain in duplicate grown from different colonies.

[0155] To confirm that each lysate could carry out transcription from the T7 genome and produce proteins, we

developed a simple cascade CFES reaction containing T7 genomes and fluorescent reporter plasmid pJL1-T7-Pr-sfGFP (FIG. 5A). In these reactions endogenous *E. coli* RNA polymerase transcribes T7 early genes including T7 RNAP (gp1) (FIG. 5B). T7 RNAP then drives sfGFP expression via the reporter plasmid. sfGFP is only synthesized in the presence of functional *E. coli* TXTL machinery and T7 transcriptional activity. CFES using each lysate showed sfGFP yields matching (infC_UP, recC_DOWN, trxA_UP) or exceeding that of wild-type BL21 TXTL after 20 hr (oxyS_UP, cyaR_UP, rna_DOWN, dgt_UP) (FIG. 6A). However, these CFES reactions did not result in detectable plaque forming units (PFU).

[0156] Having confirmed all lysates were capable of T7-mediated protein synthesis, CFBS reactions were carried out. infC_UP, oxyS_UP, cyaR_UP, recC_DOWN, and rna_DOWN all matched or exceeded wild-type BL21 lysate T7 yields by 20 hrs for at least one biological replicate with infC_UP lysates resulting in the greatest fold increase in T7 titer (9.7 to 16-fold) (FIG. 6B). Consistent with expectations, dgt_UP and trxA_DOWN eliminated T7 yields in some reactions and lowered yields by 10² to 10⁵-fold in reactions where T7 was detected. trxA_UP reactions also had T7 yields diminished by ~10³-fold, which was consistent with in vivo effector assays.

[0157] It is notable that high or low CFES sfGFP yields did not necessarily predict CFBS T7 yields (FIG. 6C). For instance, infC_UP matched wild-type BL21 TXTL for sfGFP yields but increased T7 yields whereas dgt_UP and trxA_DOWN exceeded wild-type BL21 for sfGFP yields but suffered substantial loss of T7 titer. This suggests that the influence of TXTL background on phage yields is not strictly tied to protein synthesis capacity. Multiple reports describe differential optimization of cell-free synthesis of different proteins usually by varying reaction conditions (e.g. temperature, volume), composition (e.g. [Mg²⁺], linear vs. circular template, redox state), and bacterial source of transcriptional/translation machinery (e.g. *E. coli* A19 vs BL21 vs non-*E. coli* bacteria)^{15, 70-73}. These factors affect the physicochemical environment in which protein synthesis takes place and some degree of optimization may be required for each new protein or phage to be produced using cell-free systems. On the road to engineering of chassis strains for CFBS TXTL donors, further investigation is required to determine if the positive/negative effectors described here are unique to T7 or can be generalized to develop a generalized CFBS platform optimized for more production of more diverse phage.

[0158] The trends for effector impact on T7 fitness were the same in vivo and in CFBS, (e.g. positive effector in vivo and in CFBS) suggesting in vivo screening of effectors is an appropriate approach for determining TXTL backgrounds may support CFBS yield improvements.

[0159] Prior work has demonstrated that CFES supplementation with exogenous TrxA and IF-3 improved synthesis of sfGFP transcribed from plasmids by T7 RNAP⁶⁹. Here, overexpression of IF-3 increased T7 but did not have a significant impact on sfGFP yields. Conversely, TrxA overexpression negatively impacted CFBS T7 titers again without significant impact on sfGFP. These results may be explained by the impact of pleiotropic effects of overexpressed endogenous protein. IF-3 does not have any known transcriptional regulation roles whereas TrxA interacts with numerous proteins and genes with a wide variety of meta-

bolic functions and indirectly as a transcriptional regulator⁵⁷. Likewise, OxyS and CyaR primarily function as transcription and transcription regulators of *E. coli* genes and may significantly alter the proteome of TXTL machinery. To avoid pleiotropic effects, future work supplementing CFBS with exogenous proteins could help elucidate whether T7 effectors influence CFBS yields directly or indirectly. In particular, proteomics of oxyS- and cyaR-OX lysates may reveal more CRISPRi and OX targets to improve CFBS yields while minimizing global transcriptional changes. A combination of proteomics, CRISPRi, OX has the potential to inform the design of mutant *E. coli* lysates optimized for complex phage production rather than simply protein synthesis.

Example 6: Rebooting Synthetic T7 Using CFBS Methods of the Present Technology

[0160] T7 genome fragments A, B, C, D, E, and F were generated by PCR using NEB® Q5 DNA polymerase (100 µL each) and purified phage T7 gDNA as template. PCR primers were designed to have 20 bp overlap with adjacent sequences.

[0161] PCR product size was confirmed by gel electrophoresis. After size confirmation, fragments were purified using Qiagen QiaQuick PCR purification kit.

[0162] Purified PCR products were mixed (0.2 pmol each) with NEBuilder® HiFi DNA Assembly mix from New England Biolabs (Ipswich, MA) to assemble synthetic T7 genomes. Fragments A+B+C+D generated synthetic wild-type T7 genomes. Fragments A+B+E+CASSETTE+F+D were assembled to generate synthetic T7 genomes encoding expression cassettes (“CASSETTE”) for production of heterologous proteins concomitant to T7 replication. Expression cassettes were loaded at 0.6 pmol in assembly reactions. Assembled T7 genomes were rebooted using cell-free bacteriophage synthesis: 1 µL of Gibson assembly mixture was used in 9 µL CFBS reaction mixture to reboot into functional phage.

[0163] As a demonstration of this phage engineering workflow, T7_{ClyF} (expresses anti-MRSA antimicrobial peptide) and T7_{sfGFP} were assembled and rebooted entirely in vitro. See FIGS. 13-14.

[0164] Without wishing to be bound by theory, it is believed that these improved CFBS yields achieved with the methods of the present technology can be replicated in T7-like viruses given their structural and behavioral similarities to T7 phage. See e.g., Volozhantsev et al., *Arch Virol* (2016) 161:499-501; Lavigne et al., *Virology* 312 (2003) 49-59 and Leon-Velarde et al., *Virology Journal* 2014, 11:188.

Example 7: CFBS Methods of the Present Technology with Different Phage Types

[0165] The CFBS methods described in Example 1 will be applied to additional phage types such as one or more of Lambda, K1E, T3, T5, T4, and PhiX174. It is anticipated that lysates obtained from genetically modified bacterial host cells (e.g., *E. coli*) that overexpress one or more of translation initiation factor IF-3 (infC), OxyS and CyaR and/or repress RecC subunit exonuclease RecBCD will achieve enhanced CFBS yields with one or more of these additional phage types relative to conventional CFBS methods.

EQUIVALENTS

[0166] The present technology is not to be limited in terms of the particular embodiments described in this application, which are intended as single illustrations of individual aspects of the present technology. Many modifications and variations of this present technology can be made without departing from its spirit and scope, as will be apparent to those skilled in the art. Functionally equivalent methods and apparatuses within the scope of the present technology, in addition to those enumerated herein, will be apparent to those skilled in the art from the foregoing descriptions. Such modifications and variations are intended to fall within the scope of the present technology. It is to be understood that this present technology is not limited to particular methods, reagents, compounds compositions or biological systems, which can, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

[0167] In addition, where features or aspects of the disclosure are described in terms of Markush groups, those skilled in the art will recognize that the disclosure is also thereby described in terms of any individual member or subgroup of members of the Markush group.

[0168] As will be understood by one skilled in the art, for any and all purposes, particularly in terms of providing a written description, all ranges disclosed herein also encompass any and all possible subranges and combinations of subranges thereof. Any listed range can be easily recognized as sufficiently describing and enabling the same range being broken down into at least equal halves, thirds, quarters, fifths, tenths, etc. As a non-limiting example, each range discussed herein can be readily broken down into a lower third, middle third and upper third, etc. As will also be understood by one skilled in the art all language such as “up to,” “at least,” “greater than,” “less than,” and the like, include the number recited and refer to ranges which can be subsequently broken down into subranges as discussed above. Finally, as will be understood by one skilled in the art, a range includes each individual member. Thus, for example, a group having 1-3 cells refers to groups having 1, 2, or 3 cells. Similarly, a group having 1-5 cells refers to groups having 1, 2, 3, 4, or 5 cells, and so forth.

[0169] All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety, including all figures and tables, to the extent they are not inconsistent with the explicit teachings of this specification.

REFERENCES

- [0170]** 1. ANTIBIOTIC RESISTANCE THREATS IN THE UNITED STATES. Antibiotic Resistance Coordination and Strategy Unit within the Division of Healthcare Quality Promotion, National Center for Emerging and Zoonotic Infectious Diseases, Centers for Disease Control and Prevention: CDC, 2019.
- [0171]** 2. *Antibiotic Resistance Threats in the United States*; US Centers for Disease Control: 2019.
- [0172]** 3. Hyman, P., Phages for Phage Therapy: Isolation, Characterization, and Host Range Breadth. *Pharmaceuticals (Basel)* 2019, 12 (1).
- [0173]** 4. Yehl, K.; Lemire, S.; Yang, A. C.; Ando, H.; Mimeo, M.; Torres, M. T.; de la Fuente-Nunez, C.; Lu, T.

- K., Engineering Phage Host-Range and Suppressing Bacterial Resistance through Phage Tail Fiber Mutagenesis. *Cell* 2019, 179 (2), 459-469 e9.
- [0174] 5. Panwar, R. B.; Sequeira, R. P.; Clarke, T. B., Microbiota-mediated protection against antibiotic-resistant pathogens. *Genes Immun* 2021, 22 (5-6), 255-267.
- [0175] 6. Glonti, T.; Pirnay, J. P., In Vitro Techniques and Measurements of Phage Characteristics That Are Important for Phage Therapy Success. *Viruses* 2022, 14 (7).
- [0176] 7. Wilding, K. M.; Porter Hunt, J.; Wilkerson, J. W.; Funk, P. J.; Swensen, R. L.; Christian, M. L.; Bundy, B. C., Endotoxin-free *E. coli*-based cell-free protein synthesis: Pre-expression endotoxin removal approaches for on-demand cancer therapeutic production. *Biotechnology Journal* 2019, 14 (3).
- [0177] 8. Van Belleghem, J. D.; Merabishvili, M.; Vergaunen, B.; Lavigne, R.; Vanechoutte, M., A comparative study of different strategies for removal of endotoxins from bacteriophage preparations. *J Microbiol Methods* 2017, 132, 153-159.
- [0178] 9. Lynch, S. Phage down under: stability of a travelling phage *Capsid & Tail* [Online], 2022. <https://phage.directory/capsid/phage-shipping-around-australia>.
- [0179] 10. Duyvejonck, H.; Merabishvili, M.; Vanechoutte, M.; de Soir, S.; Wright, R.; Friman, V. P.; Verbeke, G.; De Vos, D.; Pirnay, J. P.; Van Mechelen, E.; Vermeulen, S. J. T., Evaluation of the Stability of Bacteriophages in Different Solutions Suitable for the Production of Magistral Preparations in Belgium. *Viruses* 2021, 13 (5).
- [0180] 11. Jonczyk, E.; Klak, M.; Miedzybrodzki, R.; Gorski, A., The influence of external factors on bacteriophages—review. *Folia Microbiol (Praha)* 2011, 56 (3), 191-200.
- [0181] 12. Silverman, A. D.; Karim, A. S.; Jewett, M. C., Cell-free gene expression: an expanded repertoire of applications. *Nat Rev Genet* 2020, 21 (3), 151-170.
- [0182] 13. Caschera, F.; Noireaux, V., Synthesis of 2.3 mg/ml of protein with an all *Escherichia coli* cell-free transcription-translation system. *Biochimie* 2014, 99, 162-8.
- [0183] 14. Romantseva, E.; Alperovich, N.; Ross, D.; Lund, S. P.; Strychalski, E. A., Effects of DNA template preparation on variability in cell-free protein production. *Synth Biol (Oxf)* 2022, 7 (1), ysac015.
- [0184] 15. Batista, A. C.; Levrier, A.; Soudier, P.; Voyvodic, P. L.; Achmedov, T.; Reif-Trauttmansdorff, T.; DeVisch, A.; Cohen-Gonsaud, M.; Faulon, J. L.; Beisel, C. L.; Bonnet, J.; Kushwaha, M., Differentially Optimized Cell-Free Buffer Enables Robust Expression from Unprotected Linear DNA in Exonuclease-Deficient Extracts. *ACS Synth Biol* 2022, 11 (2), 732-746.
- [0185] 16. Jew, K.; Smith, P. E. J.; So, B.; Kasman, J.; Oza, J. P.; Black, M. W., Characterizing and Improving pET Vectors for Cell-free Expression. *Front Bioeng Biotechnol* 2022, 10, 895069.
- [0186] 17. Guzman-Chavez, F.; Arce, A.; Adhikari, A.; Vadhin, S.; Pedroza-Garcia, J. A.; Gandini, C.; Ajioka, J. W.; Molloy, J.; Sanchez-Nieto, S.; Varner, J. D.; Federici, F.; Haseloff, J., Constructing Cell-Free Expression Systems for Low-Cost Access. *ACS Synth Biol* 2022, 11 (3), 1114-1128.
- [0187] 18. Silverman, A. D.; Kelley-Loughnane, N.; Lucks, J. B.; Jewett, M. C., Deconstructing Cell-Free Extract Preparation for in Vitro Activation of Transcriptional Genetic Circuitry. *ACS Synth Biol* 2019, 8 (2), 403-414.
- [0188] 19. Garenne, D.; Thompson, S.; Brisson, A.; Khakimzhan, A.; Noireaux, V., The all-*E. coli* TXTL toolbox 3.0: new capabilities of a cell-free synthetic biology platform. *Synth Biol (Oxf)* 2021, 6 (1), ysab017.
- [0189] 20. Rustad, M.; Eastlund, A.; Jardine, P.; Noireaux, V., Cell-free TXTL synthesis of infectious bacteriophage T4 in a single test tube reaction. *Synth Biol (Oxf)* 2018, 3 (1), ysy002.
- [0190] 21. Sun, Z. Z.; Hayes, C. A.; Shin, J.; Caschera, F.; Murray, R. M.; Noireaux, V., Protocols for implementing an *Escherichia coli* based TX-TL cell-free expression system for synthetic biology. *J Vis Exp* 2013, (79), e50762.
- [0191] 22. Wilding, K. M.; Zhao, E. L.; Earl, C. C.; Bundy, B. C., Thermostable lyoprotectant-enhanced cell-free protein synthesis for on-demand endotoxin-free therapeutic production. *N Biotechnol* 2019, 53, 73-80.
- [0192] 23. Hunt, J. P.; Yang, S. O.; Wilding, K. M.; Bundy, B. C., The growing impact of lyophilized cell-free protein expression systems. *Bioengineered* 2017, 8 (4), 325-330.
- [0193] 24. Falgenhauer, E.; von Schonberg, S.; Meng, C.; Muckl, A.; Voge, K.; Emslander, Q.; Ludwig, C.; Simmel, F. C., Evaluation of an *E. coli* Cell Extract Prepared by Lysozyme-Assisted Sonication via Gene Expression, Phage Assembly and Proteomics. *Chembiochem* 2021, 22 (18), 2805-2813.
- [0194] 25. Garenne, D.; Bowden, S.; Noireaux, V., Cell-free expression and synthesis of viruses and bacteriophages: applications to medicine and nanotechnology. *Curr Opin Syst Biol* 2021, 28.
- [0195] 26. Xu, H.; Bao, X.; Hong, W.; Wang, A.; Wang, K.; Dong, H.; Hou, J.; Govinden, R.; Deng, B.; Chenia, H. Y., Biological Characterization and Evolution of Bacteriophage T7-big up tri, openholin During the Serial Passage Process. *Front Microbiol* 2021, 12, 705310.
- [0196] 27. Shin, J.; Jardine, P.; Noireaux, V., Genome replication, synthesis, and assembly of the bacteriophage T7 in a single cell-free reaction. *ACS Synth Biol* 2012, 1 (9), 408-13.
- [0197] 28. Heineman, R. H.; Bull, J. J., Testing optimality with experimental evolution: lysis time in a bacteriophage. *Evolution* 2007, 61 (7), 1695-709.
- [0198] 29. Mutalik, V. K.; Adler, B. A.; Rishi, H. S.; Piya, D.; Zhong, C.; Koskella, B.; Kutter, E. M.; Calendar, R.; Novichkov, P. S.; Price, M. N.; Deutschbauer, A. M.; Arkin, A. P., High-throughput mapping of the phage resistance landscape in *E. coli*. *PLoS Biol* 2020, 18 (10), e3000877.
- [0199] 30. Qimron, U.; Marintcheva, B.; Tabor, S.; Richardson, C. C., Genomewide screens for *Escherichia coli* genes affecting growth of T7 bacteriophage. *Proc Natl Acad Sci USA* 2006, 103 (50), 19039-44.
- [0200] 31. You, L.; Suthers, P. F.; Yin, J., Effects of *Escherichia coli* physiology on growth of phage T7 in vivo and in silico. *J Bacteriol* 2002, 184 (7), 1888-94.
- [0201] 32. Joseph, R. C.; Sandoval, N. R., Single and multiplexed gene repression in solventogenic *Clostridium* via Cas12a-based CRISPR interference. *Synth Syst Biotechnol* 2023, 8 (1), 148-156.
- [0202] 33. Keseler, I. M.; Gama-Castro, S.; Mackie, A.; Billington, R.; Bonavides-Martinez, C.; Caspi, R.

- Kothari, A.; Krummenacker, M.; Midford, P. E.; Muniz-Rascado, L.; Ong, W. K.; Paley, S.; Santos-Zavaleta, A.; Subhraveti, P.; Tierrafría, V. H.; Wolfe, A. J.; Collado-Vides, J.; Paulsen, I. T.; Karp, P. D., The EcoCyc Database in 2021. *Front Microbiol* 2021, 12, 711077.
- [0203] 34. Marchand, I.; Nicholson, A. W.; Dreyfus, M., Bacteriophage T7 protein kinase phosphorylates RNase E and stabilizes mRNAs synthesized by T7 RNA polymerase. *Mol Microbiol* 2001, 42 (3), 767-776.
- [0204] 35. Emily C. A. Goodall, a. A. R., a Iain G. Johnston, a Sara Jabbari, a Keith A. Turner, b Adam F. Cunningham, a; Peter A. Lund, a. J. A. C., a Ian R. Henderson, The Essential Genome of *Escherichia coli* K-12. *mBio* 2018, 9 (1).
- [0205] 36. Burova, E.; Hung, S. C.; Sagitov, V.; Stitt, B. L.; Gottesman, M. E., *Escherichia coli* NusG Protein Stimulates Transcription Elongation Rates In Vivo and In Vitro. *J Bacteriol* 1995, 177 (5), 1388-1392.
- [0206] 37. Massy, B. d.; A. Weisberg, R.; Studier, F. W., Gene 3 endonuclease of bacteriophage T7 resolves conformationally branched structures in double-stranded DNA. *J Mol Biol* 1987, 193, 359-376.
- [0207] 38. Seroka, K.; Wackernagel, W., In vivo effects of recBC DNase, exonuclease I, and DNA polymerases of *Escherichia coli* on the infectivity of native and single-stranded DNA of bacteriophage T7. *J Virol* 1977, 21 (3).
- [0208] 39. Kim, W.; Lee, Y., Mechanism for coordinate regulation of rpoS by sRNA-sRNA interaction in *Escherichia coli*. *RNA Biol* 2020, 17 (2), 176-187.
- [0209] 40. Barshishat, S.; Elgrably-Weiss, M.; Edelstein, J.; Georg, J.; Govindarajan, S.; Haviv, M.; Wright, P. R.; Hess, W. R.; Altuvia, S., OxyS small RNA induces cell cycle arrest to allow DNA damage repair. *EMBO J* 2018, 37 (3), 413-426.
- [0210] 41. Russel, M.; Model, P., Direct cloning of the trxB gene that encodes thioredoxin reductase. *J Bacteriol* 1985, 163 (1), 238-242.
- [0211] 42. Chitboonthavisuk, C.; Luo, C. H.; Huss, P.; Fernholz, M.; Raman, S., Engineering a Dynamic Controllable Infectivity Switch in Bacteriophage T7. *ACS Synth Biol* 2022, 11 (1), 286-296.
- [0212] 43. Adler, S.; Modrich, P., T7-induced DNA polymerase. Requirement for thioredoxin sulfhydryl groups. *Journal of Biological Chemistry* 1983, 258 (11), 6956-6962.
- [0213] 44. Himawan, J. S.; Richardson, C. C., Genetic analysis of the interaction between bacteriophage T7 DNA polymerase and *Escherichia coli* thioredoxin. *Proc Natl Acad Sci USA* 1992, 89 (20), 9774-9778.
- [0214] 45. Heineman, R. Lysis time, optimality, and the genetics of evolution in a T7 phage model system. University of Texas at Austin, 2007.
- [0215] 46. Yaguchi, K.; Mikami, T.; Igari, K.; Yoshida, Y.; Yokoyama, K.; Makino, K., Identification of LexA regulated promoters in *Escherichia coli* O157:H7. *J Gen Appl Microbiol* 2011, 57, 219-230.
- [0216] 47. De Lay, N.; Gottesman, S., The Crp-activated small noncoding regulatory RNA CyaR (RyeE) links nutritional status to group behavior. *J Bacteriol* 2009, 191 (2), 461-76.
- [0217] 48. Snappyan, M.; Robin, S.; Yeretssian, G.; Lecocq, M.; Marc, F.; Sakanyan, V., Cell-Free Protein Synthesis by Diversifying Bacterial Transcription Machinery. *Bio Tech* 2021, 10 (4).
- [0218] 49. Morita, T.; Kawamoto, H.; Mizota, T.; Inada, T.; Aiba, H., Enolase in the RNA degradosome plays a crucial role in the rapid decay of glucose transporter mRNA in the response to phosphosugar stress in *Escherichia coli*. *Mol Microbiol* 2004, 54 (4), 1063-75.
- [0219] 50. Gassner, C.; Schneider-Scherzer, E.; Lottspeich, F.; Schweiger, M.; Auer, B., *Escherichia coli* bacteriophage T1 DNA methyltransferase appears to interact with *Escherichia coli* enolase. *Biological Chemistry* 1998, 379 (4-5), 621-623.
- [0220] 51. Zhang, K.; Li, S.; Wang, Y.; Wang, Z.; Mulvenna, N.; Yang, H.; Zhang, P.; Chen, H.; Li, Y.; Wang, H.; Gao, Y.; Wigneshweraraj, S.; Matthews, S.; Zhang, K.; Liu, B., Bacteriophage protein PEIP is a potent *Bacillus subtilis* enolase inhibitor. *Cell Rep* 2022, 40 (1), 111026.
- [0221] 52. Gawel, D.; Hamilton, M. D.; Schaaper, R. M., A novel mutator of *Escherichia coli* carrying a defect in the dgt gene, encoding a dGTP triphosphohydrolase. *J Bacteriol* 2008, 190 (21), 6931-9.
- [0222] 53. Shadrin, A.; Sheppard, C.; Savalia, D.; Severinov, K.; Wigneshweraraj, S., Overexpression of *Escherichia coli* udk mimics the absence of T7 Gp2 function and thereby abrogates successful infection by T7 phage. *Microbiology (Reading)* 2013, 159 (Pt 2), 269-274.
- [0223] 54. Zhang, A.; Altuvia, S.; Tiwari, A.; Argaman, L.; Hengge-Aronis, R.; Storz, G., The OxyS regulatory RNA represses rpoS translation and binds the Hfq (HF-I) protein. *EMBO J* 1998, 17 (20), 6061-6068.
- [0224] 55. Tabib-Salazar, A.; Liu, B.; Barker, D.; Burchell, L.; Qimron, U.; Matthews, S. J.; Wigneshweraraj, S., T7 phage factor required for managing RpoS in *Escherichia coli*. *Proc Natl Acad Sci USA* 2018, 115 (23), E5353-E5362.
- [0225] 56. Savalia, D.; Robins, W.; Nechaev, S.; Molineux, I.; Severinov, K., The role of the T7 Gp2 inhibitor of host RNA polymerase in phage development. *J Mol Biol* 2010, 402 (1), 118-26.
- [0226] 57. Kumar, J. K.; Tabor, S.; Richardson, C. C., Proteomic analysis of thioredoxin-targeted proteins in *Escherichia coli*. *Proc Natl Acad Sci USA* 2004, 101 (11), 3759-64.
- [0227] 58. Jurado, P.; de Lorenzo, V.; Fernandez, L. A., Thioredoxin fusions increase folding of single chain Fv antibodies in the cytoplasm of *Escherichia coli*: evidence that chaperone activity is the prime effect of thioredoxin. *J Mol Biol* 2006, 357 (1), 49-61.
- [0228] 59. Miranda-Vizuet, A.; Rodriguez-Ariza, A.; Toribio, F.; Holmgren, A.; Lopez-Barea, J.; Pueyo, C., The levels of ribonucleotide reductase, thioredoxin, glutaredoxin 1, and GSH are balanced in *Escherichia coli* K12. *J Biol Chem* 1996, 271 (32), 19099-103.
- [0229] 60. Russel, M.; Model, P., Thioredoxin is required for filamentous phage assembly. *Proc Natl Acad Sci USA* 1985, 82.
- [0230] 61. Dillingham, M. S.; Kowalczykowski, S. C., RecBCD Enzyme and the Repair of Double-Stranded DNA Breaks. *Microbiology and Molecular Biology Reviews* 2008, 72 (4), 642-671.
- [0231] 62. Liao, C.; Ttofali, F.; Slotkowski, R. A.; Denny, S. R.; Cecil, T. D.; Leenay, R. T.; Keung, A. J.; Beisel, C. L., Modular one-pot assembly of CRISPR arrays enables library generation and reveals factors influencing crRNA biogenesis. *Nat Commun* 2019, 10 (1), 2948.

- [02332] 63. Sandoval, N. R.; Mills, T. Y.; Zhang, M.; Gill, R. T., Elucidating acetate tolerance in *E. coli* using a genome-wide approach. *Metab Eng* 2011, 13 (2), 214-24.
- [02333] 64. González-Flecha, B.; Demple, B., Role for the oxyS Gene in Regulation of Intracellular Hydrogen Peroxide in *Escherichia coli*. *J Bacteriol* 1999, 181 (12), 3833-3836.
- [0234] 65. Loison, P.; Majou, D.; Gelhaye, E.; Boudaud, N.; Gantzer, C., Impact of reducing and oxidizing agents on the infectivity of Qbeta phage and the overall structure of its capsid. *FEMS Microbiol Ecol* 2016, 92 (11).
- [0235] 66. Myers, J. A.; Beauchamp, B. B.; Richardson, C. C., Gene 1.2 protein of bacteriophage T7. Effect on deoxyribonucleotide pools. *Journal of Biological Chemistry* 1987, 262 (11), 5288-5292.
- [0236] 67. Huber, H. E.; Beauchamp, B. B.; Richardson, C. C., *Escherichia coli* dGTP triphosphohydrolase is inhibited by gene 1.2 protein of bacteriophage T7. *Journal of Biological Chemistry* 1988, 263 (27), 13549-13556.
- [0237] 68. Saito, H.; Richardson, C. C., Genetic analysis of gene 1.2 of bacteriophage T7: isolation of a mutant of *Escherichia coli* unable to support the growth of T7 gene 1.2 mutants. *J Virol* 1981, 37 (1), 343-351.
- [0238] 69. Airen, I. O. Genome-wide Functional Genomic Analysis for Physiological Investigation and Improvement of Cell-Free Protein Synthesis. Stanford University, 2011.
- [0239] 70. Kim, J.; Copeland, C. E.; Seki, K.; Vogeli, B.; Kwon, Y. C., Tuning the Cell-Free Protein Synthesis System for Biomanufacturing of Monomeric Human Filaggrin. *Front Bioeng Biotechnol* 2020, 8, 590341.
- [0240] 71. Noireaux, J. S. V., Efficient cell-free expression with the endogenous *E. Coli* RNA polymerase and sigma factor 70. *Journal of Biological Engineering* 2010, 4 (8).
- [0241] 72. Yim, S. S.; Johns, N. I.; Park, J.; Gomes, A. L.; McBee, R. M.; Richardson, M.; Ronda, C.; Chen, S. P.; Garenne, D.; Noireaux, V.; Wang, H. H., Multiplex transcriptional characterizations across diverse bacterial species using cell-free systems. *Mol Syst Biol* 2019, 15 (8), e8875.
- [0242] 73. Li, J.; Wang, H.; Kwon, Y. C.; Jewett, M. C., Establishing a high yielding *streptomyces*-based cell-free protein synthesis system. *Biotechnol Bioeng* 2017, 114 (6), 1343-1353.
- [0243] 74. Jakociune, D.; Moodley, A., A Rapid Bacteriophage DNA Extraction Method. *Methods Protoc* 2018, 1 (3).
- [0244] 75. Chen, P.; Zhou, J.; Wan, Y.; Liu, H.; Li, Y.; Liu, Z.; Wang, H.; Lei, J.; Zhao, K.; Zhang, Y.; Wang, Y.; Zhang, X.; Yin, L., A Cas12a ortholog with stringent PAM recognition followed by low off-target editing rates for genome editing. *Genome Biol* 2020, 21 (1), 78.
- [0245] 76. Liu, Y.; Huang, H.; Wang, H.; Zhang, Y., A novel approach for T7 bacteriophage genome integration of exogenous DNA. *J Biol Eng* 2020, 14, 2.
- [0246] 77. Hay, M.; Li, Y. M.; Ma, Y., RNA extraction of *Escherichia coli* grown in Lysogeny Broth for use in RT-qPCR. *JEMI Methods* 2017, 1, 1-6.
- [0247] 78. Livak, K. J.; Schmittgen, T. D., Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001, 25 (4), 402-8.
- [0248] 79. Zhou, K.; Zhou, L.; Lim, Q.; Zou, R.; Stephanopoulos, G.; Too, H. P., Novel reference genes for quantifying transcriptional responses of *Escherichia coli* to protein overexpression by quantitative PCR. *BMC Mol Biol* 2011, 12, 18.
- [0249] 80. Kim, J.; Copeland, C. E.; Padumane, S. R.; Kwon, Y. C., A Crude Extract Preparation and Optimization from a Genomically Engineered *Escherichia coli* for the Cell-Free Protein Synthesis System: Practical Laboratory Guideline. *Method Protocol* 2019, 2 (3).

SEQUENCE LISTING

Sequence total quantity: 120
 SEQ ID NO: 1 moltype = DNA length = 32759
 FEATURE Location/Qualifiers
 source 1..32759
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 1

tctcacagtg	tacggacctaa	aagttcccc	atagggggta	cctaaagccc	agccaatcac	60
ctaaagtcaa	ccttcgggtg	accttgaggg	ttccctaagg	gttgggggatg	acccttgggt	120
ttgtcttttg	gtgttacctt	gagtggtctt	ctgtgtccct	atctgttaca	gtctcctaaa	180
gtatctctct	aaagtcacct	cctaactgct	atcctaagc	caacacctaa	agcctacacc	240
taaagaccca	tcaagtoaac	gcttatctta	aagtttaaac	ataaagacca	gacctaaaga	300
ccagacctaa	agacactaca	taaagaccag	acctaagac	gccttgttgt	tagccataaa	360
gtgataacct	ttaatacttg	tctttattaa	tacaactcac	tataaggaga	gacaacttaa	420
agagacttaa	aagatttaatt	taaaatttat	caaaaagagt	attgacttaa	agtctaacct	480
ataggatact	tacagccatc	gagagggaca	cggcggaatg	ccatcccaat	cgacaccggg	540
gtcaaccgga	taagtagaca	gcttgataag	tgcgcagaaa	aacaggattt	gacaacatga	600
agtaacatgc	agtaagatac	aaatcgctag	gtaacactag	cagcgtoaac	cgggcgcaca	660
gtgccttcta	ggtgacttaa	gcgcaccacg	gcacataagg	tgaacacaaa	cgggtgacaa	720
catgaagtaa	acacggtacg	atgtaccaca	tgaacagaca	gtgagtcacc	acactgaaag	780
gtgatgcggt	ctaacgaaac	ctgacctaa	acgctcttta	acaatctggt	aaatagctct	840
tgagtgcagt	actagcggat	aactcaaggg	tatcgcaagg	tgccctttat	gatattcact	900
aataaactgca	cgaggttaaca	caagatggct	atgtctaaca	tgacttacaa	caacgttttc	960
gaccacgctt	acgaaatgct	gaaagaaaac	atccgttatg	atgacatccg	tgacactgat	1020
gacctgcacg	atgctattca	catggctgcc	gataatgcag	ttccgcacta	ctacgctgac	1080
atcttttagcg	taattggcaag	tgagggcatt	gaccttgagt	tcgaagactc	tggtctgatg	1140
cctgacacca	aggacgtaat	cgcgcatcgt	caagcgcgta	tctatgagca	attaacgatt	1200
gacctctggg	aagacgcaga	agacttgctc	aatgaatact	tggagggaag	cgaggagtac	1260
gaggaggatg	aagagtaatg	tctactacca	acgtgcaata	cggctctgacc	gctcaaaactg	1320

-continued

tacttttcta	tagcgacatg	gtgcgctgtg	gctttaactg	gtcactcgca	atggcacagc	1380
tcaaagaact	gtacgaaaac	aacaaggcaa	tagcttttaga	atctgctgag	tgatagactc	1440
aaggctcgctc	ctagcgagtg	gcctttatga	ttatcacttt	acttatgagg	gagtaatgta	1500
tatgcttact	atcggctctac	tcaccgctct	aggctagct	gtaggtgcat	cctttgggaa	1560
ggcttttagt	gtagctgtag	gttccacttt	taccgcttg	atcatcatag	gaatcatcaa	1620
aggggcacta	cgcaaatgat	gaagcactac	gttatgccaa	tcacacagtc	caacggggca	1680
accgtatgta	cacctgatgg	gttcgcaatg	aaacaacgaa	tcgaacgcct	taagcgtgaa	1740
ctccgcatta	accgcaagat	taacaagata	ggttccggct	atgacagaac	gcactgatgg	1800
cttaagaaaa	ggttatatgc	ccaatggcac	actatacgct	gcaaatcggc	gaatagttag	1860
aacttggcga	gagaacaacc	tcgaacgcgc	caaggacaag	agagggcgcc	gtggcataga	1920
cgaaaggaaa	aggttaaaagc	caagaaactc	gccgcacttg	aacaggcact	agccaacaca	1980
ctgaacgcta	tctcataacg	aacataaagg	acacaatgca	atgaacatta	ccgacatcat	2040
gaacgctatc	gacgcaatca	agacactgcc	aatctgtgaa	cttgacaagc	gtcaaggtat	2100
ggcttatcgac	ttactgtctg	agatgggtcaa	cagcgagacg	tgtagtgccg	agctaaccga	2160
actaaatcag	gcacttgagc	atcaagattg	gtggactacc	ttgaagtgtc	tcacggctga	2220
cgcagggttc	aagatgctct	gtaatgggtca	cttctcggtc	gcttatagtc	accgctgct	2280
acctaacaga	gtgattaaag	tgggctttaa	gaaagaggat	tcaggcgcat	cctataccgc	2340
attctgcgcg	atgtatcagg	gtcgtcctgg	tatccctaac	gtctacgatg	tacagcgcca	2400
cgtggatgc	tatacgggtg	tacttgacgc	acttaaggat	tcgagcggtt	tcaacaatga	2460
tgcccatat	aaatacgtcg	agattgcaag	cgacatcatt	gattgcaatt	cggatgagca	2520
tgatgagtta	actggatggg	atgggtgagtt	tgttgaaact	tgtaaacata	tcggcaagtt	2580
ctttgagggc	atcgctctcat	tcgacatgca	tagcgggaac	atcatgttct	caaatggaga	2640
cgtaccatac	atcacccgac	cggtatcatt	ctcgacagaag	aaagacgggt	gcgcattcag	2700
catcgaccct	gaggaaactca	tcaaggaaat	cgaggaaagt	gcacgcagaga	aagaaatgta	2760
ccgcgctaag	gcccgtaaa	aacgtcacga	ggggcgctta	gaggcacgca	gattcaaacc	2820
tcgcaaccgc	aaggcacgta	aagcacacaa	agctaagcgc	gaaagaatgc	ttgctgcgtg	2880
gcgatgggct	gaacgtcaag	aacggcgtaa	ccatgaggtta	gctgtagatg	tactaggaag	2940
aaaccaataac	gctatgtctct	gggtcaacat	gttctctggg	gactttaagg	cgcttgagga	3000
acgaatcgcg	ctgcactggc	gtaatgctga	ccggatggct	atcgctaatg	gtcttacgct	3060
caacatgat	aagcaacttg	acgcaatggt	aatgggctga	tagtcttatt	ttacagggtca	3120
tctgcgggtg	gcctgaatag	gtacgattta	ctaactggaa	gaggcactaa	atgaacacga	3180
ttaaacatcg	ttaagaacgac	ttctctgaca	tcgaactggc	tgctatcccc	ttcaaacactc	3240
tggctgacca	ttacgggtgag	cgtttagctc	gcgaacagtt	ggcccttgag	catgagttct	3300
acgagatggg	tgaagcgacg	ttccgcaaga	tgtttgagcg	tcaacttaaa	gctggtaggg	3360
ttgcggataa	cgctgcgcgc	aagcctctca	tcactaccct	actccctaag	atgatgacac	3420
gcatcaacga	ctgggttgag	gaagtgaag	ctaagcgccg	caagcgcccc	acagccttcc	3480
agttcctgca	agaaatcaag	ccggaagccg	tagcgtacat	caccattaa	accactctgg	3540
cttgccctaac	cagtgctgac	aatacaaccg	ttcaggctgt	agcaagcgca	atcggtcggg	3600
ccattgagga	cgaggctcgc	ttcggctgta	tccgtgacct	tgaagctaa	cacttcaaga	3660
aaaacgttga	ggaacaaactc	aacaagcgcg	tagggcacgt	ctacaagaaa	gcattttatg	3720
aagtgtgcga	ggctgacatg	ctctctaaag	gtctactcgg	tggcgaggcg	tggtcttctg	3780
ggcataagga	agactctatt	catgtaggag	tacgctgcat	cgagatgctc	attgagttca	3840
ccggaatggg	tagctttacac	cgccaaaatg	ctggcgtagt	aggtaaacg	tctgagacta	3900
tcgaactcgc	acctgaatac	gctgagggtta	tcgcaaccgc	tgagggtgag	ctggctggca	3960
tctctccgat	gttccaaact	tcggtagttc	ctcctaagcc	gtggactggc	attactgggt	4020
gtggctattg	ggctaacggg	cgtcgtctct	tggcgctggg	gcgtactcac	agtaagaaag	4080
cactgatgcg	ctacgaagac	gtttacatgc	ctgagggtga	caaagcgatt	aacattgcgc	4140
aaaaaccgcg	atggaaaaatc	aacaagaaag	tcttagcggt	cgccaacgta	atcaccgaag	4200
ggaagcattg	tcgggtcgag	gcattccctg	cgattgagcg	tgaagaactc	ccgatgaaac	4260
gggaagacat	cgacatgaat	cctgagggtc	tcaccgcgtg	gaaacgtgct	gccgctgctg	4320
tgtaccgcaa	ggacaaggct	cgcgaagtct	gccgtatcag	ccttgagttc	atgcttgagc	4380
aagccaataa	ggttgctaac	cataaggcca	ctcgggtccc	ttacaacatg	gactggcgcg	4440
gtcgtgttta	cgctgtgtca	atgttcaacc	cgcaaggtaa	cgatatgacc	aaaggactgc	4500
ttacgctggc	gaaaggtaaa	ccaatcggtta	aggaagggtta	ctactggctg	aaaatccacg	4560
gtgcaaaactg	tgccgggtgc	gataagggtc	cgttccctga	gcgcataca	ttcattgagg	4620
aaaaccacga	gaacatcatg	gcttgcgcta	agtctccact	ggagaacact	tggtgggctg	4680
agcaagattc	tccgttctgc	ttccttgctg	tctgctttga	gtacgctggg	gtacagcacc	4740
acggcctgag	ctataacgc	ttccctccgc	tggcggttga	cgggtcttgc	tctgggcatcc	4800
agcacttctc	cgcgatgctc	cgagatgagg	taggtgggtc	cgcgggttaac	ttgcttctca	4860
gtgaaaccgt	tcaggacatc	tacgggattg	ttgctaagaa	agtcaacgag	attctacaag	4920
cagacgcaat	caatgggacc	gataacgaag	tagttaccgt	gaccgatgag	aacactgggt	4980
aaatctctga	gaaagtcaag	ctgggcacta	aggcactggc	tggtcaatgg	ctggcttacg	5040
gtgttactcg	cagtgtgact	aagcgttcag	tcatgacgtc	ggcttacggg	tccaaagagt	5100
tcggcttccg	tcaacaagtg	ctggaagata	ccattcagcc	agctattgat	tccggcaagg	5160
gctgatgtt	cactcagccg	aatcaggctg	ctggatacat	ggtaagctg	atttgggaat	5220
ctgtgagcgt	gacgggtgta	gctgcgggtg	aagcaatgaa	ctggcttaag	tctgctgcta	5280
agctgctggc	tgctgaggtc	aaagataaga	agactggaga	gattctctgc	aagcgttgcg	5340
ctgtgcattg	ggtaactcct	gatgggttcc	ctgtgtggca	ggaatacaag	aagcctattc	5400
agacgcgctt	gaacctgatg	ttcctcggtc	agttccgctt	acagcctacc	attaacacca	5460
acaaagatag	cgagattgat	gcacacaaac	aggagtctgg	tatcgctcct	aactttgtac	5520
acagccaaga	cggtagccac	ctcgttaaga	ctgtagtgtg	ggcacacgag	aagtacggaa	5580
tcgaatcttt	tgcaactgatt	cacgactcct	tcgggtaccat	tccggctgac	gctgcgaacc	5640
tgttcaaaagc	agtgcgcgaa	actatgggtg	acacatatga	gtcttgtagt	gtactggctg	5700
atthctacga	ccagttcgct	gaccagttgc	acgagtctca	attggacaaa	atgccagcac	5760
tccgggttaa	aggttaacttg	aaactccgtg	acattcttaga	gtcggacttc	gcgttcgctg	5820
aacgccaat	caatacgact	cactatagag	ggacaaaactc	aaggtcattc	gcaagagtgg	5880

-continued

cctttatgat	tgaccttctt	cgggttaata	cgactcacta	taggagaacc	ttaaggttta	5940
actttaagac	ccttaagtgt	taattagaga	tttaaatata	agaattacta	agagaggact	6000
ttaagtatgc	gtaacttcga	aaagatgacc	aaacgttcta	accgtaatgc	tcgtgacttc	6060
gaggcaacca	aaagtcgcaa	gttgaataag	actaagcgtg	accgctctca	caagcgtagc	6120
tgggagggct	agtaagatgg	gacgtttata	tagtggtaat	ctggcagcat	tcaaggcagc	6180
aacaacaacg	ctgttccagt	tagacttagc	ggtcatttat	gatgactggg	atgatgccta	6240
tacaagaaaa	gattgcatac	ggttacgtat	tgaggacagg	agtggaaacc	tgattgatac	6300
tagcaccttc	taccaccacg	acgaggacgt	tctgttcaat	atgtgtactg	attgggtgaa	6360
ccatattgat	gaccagttga	aggactggaa	gtaatacgac	tcagtatagg	gacaatgctt	6420
aaggctcgct	tctaggagtg	gccttagtca	tttaaccaat	aggagataaa	cattatgatg	6480
aacattaaga	ctaaccggtt	taagccgtg	tctttcgtag	agtctgcat	taagaaggct	6540
ctggataacg	ctgggtatct	tatcgctgaa	atcaagtacg	atgggtgacg	cgggaacatc	6600
tgcgtagaca	atactgctaa	cagttactgg	ctctctcggt	tatctaaaac	gattccggca	6660
ctggagcact	taaacgggtt	tgatgttcgc	tggaagcgtc	tactgaacga	tgaccgttgc	6720
ttctacaaag	atggctttat	gcttgatggg	gaactcatgg	tcaagggcgt	agactttaac	6780
acagggtccg	gcctactgcg	taccaaatgg	actgacacga	agaaccaaga	gttccatgaa	6840
gagttattcg	ttgaaccaat	ccgtaagaaa	gataaagttc	cctttaagct	gcacactgga	6900
caccttcaca	taaaactgta	cgctatcctc	ccgctgcaca	tcgtggagtc	tggaagaagc	6960
tgtagtgta	tcacgttgct	catgcaggaa	cacgttaaga	acatgctgcc	tctgctacag	7020
gaataacttc	ctgaaactga	atggcaagcg	gctgaatctt	acgaggtcta	cgatatggta	7080
gaactacagc	aactgtacga	gcagaagcga	gcagaagggc	atgagggctc	cattgtgaaa	7140
gacccgatgt	gtatctataa	gcgcggtaag	aaatctggct	gggtggaagt	gaaacctgag	7200
aaacgaagcg	acggtatcat	tcagggtctg	gtatggggta	caaaaggtct	ggctaataga	7260
ggtaaatgta	ttgggtttga	gggtcttctt	gagagtggct	gtttagttaa	cgccacgaat	7320
atctctcgcg	ccttaatgga	tgagttcact	gagacagtaa	aagaggccac	cctaagtcaa	7380
tggggattct	ttagcccata	cggattttgc	gacaacgatg	cttgtactat	taacccttac	7440
catggctggg	cggtgcaaat	tagctacatg	gaggaacac	ctgatggctc	tttgcggcac	7500
ccatcgcttc	taatgttccg	tgccaccgag	gacaaccctc	aagagaaaat	gtaatcacac	7560
tgggtcacc	tcgggtgggc	ctttctgctg	ttataaggag	acactttatg	tttaagaagg	7620
ttggtaaat	ccttgccggt	ttggcagcta	tcctgacgct	tgcttatatt	cctgcccgtat	7680
accctcaagt	agcactagta	gtagttggcg	cttgttactt	agcggcagtg	tgtgcttgcg	7740
tgtggagtat	agttaaactg	taatacgact	cactaaagga	ggtagacac	atgatgtact	7800
taatgccatt	actcatcgct	attgtaggat	gccttgcgct	ccactgtagc	gatgatgata	7860
tgccagatgg	tcacgcttaa	taacgactcac	taaggagagc	actatatgtt	tcgacttcat	7920
tacaacaaaa	gcgttaagaa	tttcacgggt	cccggtgctg	accgttcaat	cgtatgtgcg	7980
agcgaagcgc	gagctaaagt	acctcttatt	ggtaaacacg	ttcctttggc	accgagcgtc	8040
cacatcatta	tcacccgtgg	tgactttgag	aaagcaatag	acaagaaacg	tccggttcct	8100
agtggtggcg	tgacccgctt	cccggttcgt	cgtctgttac	tcaaacgaat	caaggaggtg	8160
ttctgatggg	actgttagat	gggtgaagcct	gggaaaaaga	aaacccgccca	gtacaagcaa	8220
ctgggtgtat	agcttgctta	gagaaagatg	accgttatcc	acacacctgt	aacaaaggag	8280
ctaacgatat	gacgaacgt	gaacaagaga	tgatcattaa	gttgatagac	aataatgaag	8340
gtcgccacga	tgattttgaat	ggctgcccgt	ttctctgctc	caatgtccct	tgccacctct	8400
gccccgcaaa	taacgactcaa	aagataacct	taggtgaaat	ccgagcgatg	gacccacgta	8460
aaccacatct	gaataaaact	gaggtaactc	ctacagatga	ccagccttcc	gctgagacaa	8520
tcgaagggtg	cactaagcct	tcccactaca	tgtgttttga	cgacattgag	gctatcgaa	8580
tgattgctcg	ttcaatgaac	gttagcagct	tcaagggata	ctgcttcggt	aacactttaa	8640
agtaacagct	acgtgctggg	aagaagtcag	agttagcgt	cttagagaaa	gacctagcga	8700
aagcagactt	ctataaagaa	ctcttttgaga	aacataagga	taaatgttat	gcataacttc	8760
aagtcaaccc	cacctgcoga	cagcctatct	gatgacttca	catcttgctc	agagtgggtg	8820
cgaagagatg	gggaagagac	attcgacgat	gcgtacatca	agctgtatga	actttggaaa	8880
tcgagaggtc	aatgactatg	tcaaacgtaa	atacaggttc	acttagtggt	gacaataaga	8940
agttttgggc	tctcttgtag	tctcggagc	attccttcga	gggtccaatc	tacgctgaga	9000
cctagacga	agctctggag	ttagccgaat	ggcaatacgt	tccggctggc	tttgagggt	9060
ctcgtgtgct	tctctgtgta	gcacgaaggt	aatacgaact	actattaggg	aagactccct	9120
ctgagaaccc	aaacgaaacc	taaggagat	taacattatg	gctaagaaga	ttttcacctc	9180
tgcgctgggt	accgctgaac	cttacgctta	catcgccaag	ccggactacg	gcaacgaaga	9240
gcgtggcttt	gggaacccct	gtgggtgcta	taaagttgac	ctgactattc	ccaacaaaga	9300
cccgcgctgc	cagcgatatg	tcgatgaaat	cgtgaagtgt	cacgaagagg	cctatgctgc	9360
tgccgttgag	gaatacgaag	ctaataccacc	tgcgttagct	cgtggtaaga	aaccgctgaa	9420
accgatagag	ggtagacatg	cgttcttcga	taacgggtgac	ggtagcacta	cctttaagtt	9480
caaatgctac	gcgtcttttc	aagacaagaa	gaccaaagag	accaagcaca	tcaatctggt	9540
tgtggttgac	tcaaaaggta	agaagatgga	agacgttccg	attatcggtg	gtggctctaa	9600
gctgaaagtt	aaatatcttc	tggttccata	caagtggaa	actgctgtag	gtgcgagcgt	9660
taagctgcaa	ctggaatccg	tgatgctggg	cgaactggct	accttgggtg	gcgggtgaaga	9720
cgattgggct	gacgaagtgt	aagagaacgg	ctatgttgcc	tctgggtctg	ccaaagcgag	9780
caaacccagc	gacgaagaaa	gctgggacga	agacgacgaa	gagtcaggag	aagcagacga	9840
agacggagac	ttctaagtgg	aactgcggga	gaaaatcctt	gagcgaatca	aggtgacttc	9900
ctctgggtgt	tgggagtggc	agggcgctac	gaacaataaa	gggtacgggc	aggtgtgggtg	9960
cagcaatacc	ggaaagggtg	tctactgtca	tcgcgtaatg	tctaattgctc	cgaagggttc	10020
taccgtcctg	cactccgtgt	ataatccatt	atgtgtgaac	cctgaacacc	tatccatagg	10080
aactccaaaa	gagaactcca	ctgacatggg	aaataagggt	cgctcacaca	aggggtataa	10140
actttcagac	gaagacgtaa	tggaatcat	ggagtccagc	gagtcacatg	tatccttagc	10200
tcgcacctat	gggtgtctcc	aacagactat	ttgtgatata	cgcaaaggga	ggcgacatgg	10260
caggttacgg	cgctaaagga	atccgaaagg	ttggagcggt	tcgctctggc	ctagaggaca	10320
aggtttcaaa	cgagttggaa	tcaaaaggta	ttaaattcga	gtatgaagag	tggaaggtgc	10380
cttatgtaat	tccggcgagc	aatcacactt	acactccaga	cttcttactt	ccaaacggta	10440

-continued

tattcgttga	gacaaaggggt	ctgtgggaaa	gcgatgatag	aaagaagcac	ttattaatta	10500
gggagcagca	ccccgagcta	gacatccgta	ttgtcttctc	aagctcacgt	actaagttat	10560
acaaaggttc	tccaacgtct	tatggagagt	tctgcgaaaa	gcatgggtatt	aagttcgctg	10620
ataaaactgat	acctgtcgag	tggataaagg	aacccaagaa	ggaggtcccc	ttttagatagat	10680
taaaaagga	aggaggaaag	aaataatggc	tcgtgtacag	tttaacaac	gtgaatctac	10740
tgacgcaatc	tttgttcaact	gctcggctac	caagccaagt	cagaatgttg	gtgtccgtga	10800
gattcgcag	tggcacaaag	agcagggttg	gctcgatgtg	ggataccact	ttatcatcaa	10860
gcgagacggt	actgtggagg	caggacgaga	tgagatggct	gtaggctctc	acgctaagg	10920
ttacaaccac	aactctatcg	gcgtctgcct	tgttgggtgt	atcgacgata	aaggtaagtt	10980
cgacgctaac	tttacgcccag	cccaaatgca	atcccttcgc	tcaactgctt	tcacactgct	11040
ggctaagtac	gaaggcgctg	tgtctcgccg	ccatcatgag	gtggcgccga	aggcttgccc	11100
ttcgttcgac	cttaagcggt	gggtgggagaa	gaacgaactg	gtcacttctg	accgtggata	11160
attaattgaa	ctcactaaag	ggagaccaca	gcggtttccc	tttgttcgca	ttggaggtca	11220
aataatgccc	aagtcttata	aaacaattcta	taaggctccg	aggaggcata	tccaagtgtg	11280
ggaggcagcc	aatgggccta	taccaaaagg	ttattatata	gaccacattg	acggcaatcc	11340
actcaacgac	gccttagaca	cttccgtct	ggctctccca	aaagaaaact	catggaacat	11400
gaagactcca	aagagcaata	cctcaggact	aaagggactg	agtgggagca	aggaaaaggga	11460
gatgtggaga	ggcactgtaa	cagctgagg	taaacagcat	aactttcgta	gtagagatct	11520
attggaagtc	gttgcgtgga	ttatagaac	taggaggga	tgcatggag	aattcgcacg	11580
attccgata	gttatttctt	taccacattc	cttgtgacaa	ctgtgggag	agtgtggga	11640
actcgtgtt	ctctgacgga	cacacgttct	gctacgtatg	cgagaagtgg	actcgtggt	11700
atgaagacac	taaagagagg	gcttcaaaac	ggaaacccctc	aggaggtaaa	ccaatgactt	11760
acaaactgtg	gaacttcggg	gaatccaatg	gacgctactc	cgcgttaact	gcgagaggaa	11820
tctccaagga	aacctgtcag	aaggctggct	actggattgc	caaagtagac	gggtgtgat	11880
accaagtggc	tgactatcgg	gaccagaacg	gcaacattgt	gagtcagaag	gttcagagata	11940
aagataagaa	ctttaagacg	actggtagtc	acaagagtga	cgtctgttgc	gggaaagcact	12000
tgtggaatgg	tggttaagaag	attgtcgtta	cagaagggtga	aatcgacatg	cttaccgtga	12060
tggaaactca	agacttgaag	tatcctgtag	tgtcgttggg	tcacggtgcc	tctgccgcta	12120
agaaagacatg	cgctgcacaa	tacgaatact	ttgaccagtt	cgaacagatt	atctttaatgt	12180
tcgatatgga	cgaagcaggg	cgcacaaagc	tcgaagaggc	tgacacaggt	ctacctgctg	12240
gtaaggtacg	agtggaagtt	cttccgtgta	aggatgcaaa	cgagtgtcac	ctaaatgggtc	12300
acgacccgtga	aatcatggag	caagtgtgga	atgctgggtcc	ttggattcct	gatgggtggg	12360
tatcggctct	ttcgttaact	gaacgaatcc	gtgagcacct	atcgtccgag	gaatcagtag	12420
gtttaacttt	cagtggctgc	actggtatca	acgataagac	cttaggtgcc	cgtgggtggtg	12480
aagtcattat	ggtcacttcc	gggtccggta	tgggtaagtc	aacgttcgtc	cgtcaacaag	12540
ctctacaatg	gggcacagcg	atgggcaaga	aggtaggctt	agcgatgctt	gaggagtcgg	12600
ttgaggagac	cgctgaggag	cttataggtc	tacacaaccg	tgtccgactg	agacaatccg	12660
actcactaaa	gagagagatt	attgagaacg	gtaagttcga	ccaatgggtc	gatgaactgt	12720
tcggcaacga	tacgttccat	ctatatgact	cattcgcgga	ggctgagacg	gatagactgc	12780
tcgctaagct	ggctacagct	cgtcaggct	tgggctgtga	cgtaatcatt	ctagaccaca	12840
tctcaatcgt	cgatcccgct	tctgggtgaat	ccgatgagcg	taagatgatt	gacaaacctga	12900
tgaccaagct	caaaagggttc	gctaaagtcaa	ctgggggtgt	gctggctgta	atttgtcacc	12960
ttaaagaacc	tcgcaaaagt	aaagcacatg	aggaaaggtcg	cccgtttctt	attactgacc	13020
tacgtgggtc	tggcgcaact	cgccaactat	ctgatactat	tattgccctt	gagcgtaatc	13080
agcaaggcga	tatgcctaac	cttgtcctcg	ttcgtattct	caagtgcgcg	tttactgggtg	13140
atactggtat	cgtcggctac	attggaatata	acaaggaaac	cggatggctt	gaacctcaa	13200
gttactcagg	ggaagaagag	tcacactcag	agtcaacaga	ctgggtccaa	gacactgact	13260
cttgacagga	ttcttgatga	cttccagac	gactacgaga	agtttcgctg	gagagtcoca	13320
ttctaatacg	actcactata	ggagacacac	catgtttcaa	ctgattaaga	agttaggcca	13380
actcgtggtt	cgtatgtaca	acgtggaagc	caagcgactg	aacgatgagg	ctcgtaaaaga	13440
ggccacacag	tcacgcgctc	tggcgattcg	ctccaacgaa	ctggctgaca	gtgcattccac	13500
taaaagtacc	gaggctgccc	gtgtggcaaa	ccaagctcaa	cagctttcca	aattctttga	13560
gtaatcaaac	aggagaaacc	attatgtcta	acgtagctga	aactatccgt	ctatccgata	13620
cagctgacca	gtggaaccgt	cgaagtcaca	tcaacgttcg	caacggtaag	gcgactatgg	13680
tttaccgctg	gaaggactct	aagtctctta	agaatcacac	tcagcgtatg	acgttgacag	13740
atgagcaagc	aactcgtctg	gtcaatgcgc	ttaccaaaagc	tgccgtgaca	gcaattcatg	13800
aagctgggtc	cgtcaatgaa	gctatggctc	tcctcgacaa	gattgataac	taagagtgggt	13860
atcctcaagg	tcgcaaaagt	ggcgccctta	atgaataacta	ttcgactcac	tataggagat	13920
attaccatgc	gtgaccttaa	agttatccaa	gcagaaatcg	ctaaaactgga	agctgaactg	13980
gaggacgtta	agtaccatga	agctaagact	cgctccgctg	ttcacatctt	gaagaactta	14040
ggctggactt	ggacaagaca	gactggctgg	aagaaaccag	aagttacca	gctgagtcac	14100
aagggtgttc	ataaggacac	tatgacccac	atcaaggctg	gtgattgggt	taaggttgac	14160
atgggagttg	ttggtggata	cggctacgtc	cgctcagtta	gtggcaata	tgacaagtg	14220
tcatacatct	caggtgttac	tccacgcggg	gcaatcgttg	ccgataaagc	caacatgatt	14280
cacacaggtt	tcttgacagt	gttttcatat	gaagagattg	ttaagtcacg	ataatcaata	14340
ggagaaatca	atatgatcgt	ttctgacatc	gaagctaacg	ccctcttaga	gagcgtcaact	14400
aagttccact	gcgggggttat	ctacgactac	tccaccgctg	agtcagtaag	ctaccgtccg	14460
agtgactctg	gtgcgtatct	ggatgcgctg	gaagccgagg	ttgcacgagg	cggtcttatt	14520
gtgttcacac	acgggtcacaa	gtatgacgtt	cctgcattga	ccaaactggc	aaagttgcaa	14580
ttgaaccgag	agttccacct	tctcgtgtag	aactgtattg	acacccttgt	gttgtcacgt	14640
ttgattcatt	ccaacctcaa	ggacaccgat	atgggtcttc	tgcgttcggg	caagttgccc	14700
ggaaaacgct	ttgggtctca	cgttttggag	gcgtgggggt	atcgcttagg	cgagatgaag	14760
ggtgaataca	aagacgactt	taagcgtatg	cttgaagagc	agggtgaaga	atacgttgac	14820
ggaatggagt	ggtggaactt	caacgaagag	atgatggact	ataacgttca	ggacgttggt	14880
gtactctaa	ctctccttga	gaagctactc	tctgacaaac	attactccc	tctgagatt	14940
gactttacgg	acgtaggata	cactacgttc	tggtcagaat	cccttgaggc	cgttgacatt	15000

-continued

gaacatcgctg	ctgcatggct	gctcgctaaa	caagagcgca	acgggttccc	gtttgacaca	15060
aaagcaatcg	aagagttgta	cgtagagtta	gctgctcgcc	gctctgagtt	gtcccgtaaa	15120
ttgaccgaaa	cggttcggctc	gtggatcacg	cctaagagtg	gcactgagat	gttctgccat	15180
ccgcgaacag	gtaagccact	acctaataac	cctcgcat	agacacctaa	agttggtggt	15240
atctttaaga	agcctaagaa	caaggcacag	cgagaaggcc	gtgagccttg	cgaacttgat	15300
acccgcgagt	acgtttgctgg	tgctccttac	accccagttg	aacatgttgt	gtttaaccct	15360
tcgtctcgctg	accacattca	gaagaaactc	caagaggctg	ggtgggtccc	gaccaagtac	15420
accgataaagg	gtgctcctgt	ggtggacgat	gaggtactcg	aaggagtacg	tgtagatgac	15480
cctgagaagc	aagcgcgtat	cgacctcatt	aaagagtact	tgatgatcca	gaagcgaaatc	15540
ggacagtcgtg	ctgagggaga	caaagcatgg	cttcgttatg	ttgctgagga	tggttaagatt	15600
catggttctg	ttaaacctaa	tgagcagatt	acgggtcgtg	cgacctatgc	gttcccaaac	15660
cttgccgaaa	ttccgggtgt	acgttctcct	tatggagagc	agtgctcgcc	tgcttttgcc	15720
gctgagcacc	attgggatgg	gataaactgg	aagccttggg	ttcaggctgg	catcgacgca	15780
tcgggtctgt	agctacgcgt	cttgggtcac	ttcatggctc	gctttgataa	cggcgagtac	15840
gctcacgaga	ttcttaacgg	cgacatccac	actaagaacc	agatagctgc	tgaactacct	15900
acccgagata	acgctaagac	gttcatctat	gggttctctc	atggtgctgg	tgatgagaag	15960
attggagaca	ttgtttggtg	tggttaagag	cgcgtaagg	aactcaagaa	gaaattcctt	16020
gagaacaccc	ccgcgattgc	agcactccgc	gagttctatc	aacagacact	tgctgagttc	16080
ttcctaatggg	tagctgggtg	gcaacaagtc	aagtggaaac	gccgctggat	taaaggctctg	16140
gatggtcgta	aggtacacgt	tcgtagtctc	cacgctgcct	tgaataacct	actgcaatct	16200
gctggtgctc	tcctctgcaa	actgtggatt	atcaagaccg	aagagatgct	cgtagagaaa	16260
ggcttgaagc	atggctggga	tggggacttt	gcgtacatgg	catgggtaca	tgatgaaatc	16320
caagtaggct	gccgtaccca	agagattgct	caggtgggtca	ttgagacccg	acaagaagcg	16380
atgcgctggg	ttggagacca	ctggaaacttc	cggtgtcttc	tggtataccg	aggttaagatg	16440
ggctcctaatt	ggggcgattt	ccactgatac	aggaggctac	tcatagaaca	aagacactta	16500
acaggtgctg	cttctgaaat	gctagtagcc	tacaaattta	ccaaagctgg	gtactactgtc	16560
tattacccta	tgctgactca	gagtaaaag	gacttgggtg	tatgtaagga	tggttaaaatt	16620
agtaaggttc	agggttaaaac	agccacaacg	gttcaaacca	acacaggaga	tgccaagcag	16680
gttagggctag	gtggatggcg	taggtccgaa	tataaggatg	gagacttga	cattcttgcg	16740
gttgtggttg	acgaagatgt	gcttattttc	acatgggacg	aagtaaaagg	taagacatcc	16800
atgtgtgtcg	gcaagagaaa	caaaggcata	aaactatagg	agaaattatt	atggctatga	16860
caaagaaatt	ttaagtgtcc	ttcgacgtta	ccgcaagat	gtcgtctgac	gttcaggcaa	16920
tttagagaaa	agatagctg	catctatgta	agcaggctcg	ctcagggtcg	attgtcccca	16980
atggttaaca	gaaggaaatg	attgtccagt	ttctgacaca	cggtatggaa	ggattgatga	17040
cattcgtagt	acgtacatca	ttctgtgagg	ccattaagga	catgcacgaa	gagtatgcag	17100
ataaggactc	tttcaaacaa	ttcctgcaa	cagtacggga	gggtgtctga	tgctctgacta	17160
cctgaaagtg	tcgcaagcaa	tcaaaagttg	ccctaagact	ttccagtcca	actatgtacg	17220
gaacaatcg	agcctcgtag	cgaggccgcg	ttcccggtg	cacatctcgt	gctgactac	17280
tagtgagcgt	aacgggtggc	cttgggaaat	cactgcttcc	ggtagctcgt	ttctgaaacg	17340
aatgggagga	gtgtcttaat	gtctcgtgac	cttgtgacta	ttccacgcga	tgtgtggaac	17400
gatatacagg	gctacatcga	ctctctggaa	cgtgagaaac	atagccttaa	gaatcaacta	17460
atggaagctg	acgaatacgt	agcggaaacta	gaggagaaac	ttaatggcac	ttcttgacct	17520
taaaacaattc	gtgaaggtcg	cgacgacaag	ggatcccttg	tgatggacgg		17580
cgactggctg	gtcttccaag	ctatgagtc	tgctgagttt	gatgcctctt	gggagggaaga	17640
gatttggcac	cgatgctgtg	accacgctaa	ggcccgctcag	attcttgagg	attccattaa	17700
gtcctacgag	accgctaaga	aggcttgggc	agggtgctcca	attgtccttg	cggtccaccga	17760
tagtgtaaac	tgggcgtaag	aactgggtga	cccgaactat	aaggctaacc	gtaaggccgt	17820
gaagaaacct	gtagggtact	ttgagttcct	tgatgctctc	tttgagccgc	aagagttcta	17880
ttgcatccgt	gagctatgc	ttgaggggtga	tgacgttatg	ggagtatttg	cttccaatcc	17940
gtctgccttc	gggtcctgta	aggctgtaat	catctcttgc	gataaggact	ttaagacct	18000
ccctaactgt	gacttctcgt	gggtgaccac	tggttaacatc	ctgactcaga	ccgaagagtc	18060
cgctgactgg	tgggacacct	tcacagacct	caagggtgac	atcactgatg	gttactcagg	18120
gattgctgga	tggggtgata	ccgcgagga	cttcttgaat	aaccggttca	taaccgagcc	18180
taaaacgtct	gtgcttaagt	ccggtaagaa	caaaggccaa	gaggttacta	aatgggttaa	18240
acgcgaacct	gagcctcatg	agacgctttg	ggactgcatt	aagtccattg	gcgcgaagcc	18300
tggtatgacc	gaagaggata	ttatcaagca	gggccaagt	gctcgaatcc	tacgggtcaa	18360
cgagtacaac	tttattgaca	aggagattta	cctgtggaga	ccgtagcgta	tattggtctg	18420
ggctcttctg	ttctcggaag	gtgcctcatt	tcgtggggcc	tttgggactt	agccagaata	18480
atcaagtcgt	tacacgacac	taagtataaa	actcaaggct	cctaataata	tacgactcac	18540
tatagggaga	taggggcctt	tacgattatt	actttaagat	ttaactctaa	gaggaaatctt	18600
tattatgtta	acacctatta	accaattact	taagaacct	aacgatattc	cagatgtacc	18660
tcgtgcaacc	gctgagatc	tacagggtcg	attcaactat	gcgtacctcg	aagcgtctgg	18720
tcataatgga	cttatcgctg	ctaattgggtg	tagtgaggcc	cacatcttgg	gtttcatcca	18780
gggctacag	tatgctctca	acgtcattga	cgagattgag	ttacgcaagg	aacaactaag	18840
agatgatggg	gaggtatgac	actatgtgtt	ttcaccgaa	aattaaaact	ccgaagatgg	18900
ataccaatca	gattcgagcc	gttgagccag	cgccctctgac	ccaagaagtg	tcaagcgtgg	18960
agttcggtgg	gtcttctgat	gagacggata	ccgagggcac	cgaagtgtct	ggacgcaaa	19020
gctcaagggt	cgaaacgtgat	gattccgtag	cgaagtctaa	agccagcgcc	aatggctccg	19080
ctcgatgaa	atcttccatc	cgtaagtcgg	catttgagg	taagaagtga	tgtctgagtt	19140
cacatgtgtg	gaggctaaga	gtcgttccg	tgcaatccgg	tggaactgtg	aacaccttgg	19200
gttgccctaaa	ggattccgaag	gaacactttg	gggtctacagc	ctctacgtag	acgaagtgat	19260
ggacatgtct	gggtgcccgtg	aagagtacat	tctggactct	accggaaaac	atgtagcgta	19320
cttcgcgtgg	tgcgtaagct	gtgacattca	ccacaaagga	gacattctgg	atgtaacgtc	19380
cgttgtcatt	aactcctgag	cagactctaa	gggcttacag	cgattctctag	cgaacgcctt	19440
taagtacctt	ccggaactcc	acgattgcga	ttgggtgtct	cggtgtaagc	atgaaggcga	19500
gacaatgcgt	gtatacttta	aggaggtata	agttatgggt	aagaaagtta	agaaggccgt	19560

-continued

gaagaaagtc	accaagtcgc	ttaagaaagt	cgttaaggaa	ggggctcgtc	cggttaaaca	19620
ggttgctggc	ggcttagctg	gtctggctgg	tggtactggg	gaagcacaga	tggtggaagt	19680
accacaagct	gccgcacaga	ttgttgacgt	acctgagaaa	gaggtttcca	ctgaggacga	19740
agcacagaca	gaaagcggag	gcaagaaagc	tcgtgctggc	ggtaagaaat	ccttgagtgt	19800
agcccgtagc	tcgggtggcg	gtatcaacat	ttaatcaggga	ggttatcgtg	gaagactgca	19860
ttgaatggac	cggagggtgc	aactctaagg	gttatggctg	taagtggggt	aatggtaaac	19920
ttgtgactcc	acataggcac	atctatgagg	agacatatgg	tccagttcca	acaggaattg	19980
tggtgatgca	tatctgcgat	aacctaggt	gctataacat	aaagcacctt	acgcttgga	20040
ctocaaaggga	taattccgag	gacatggtta	ccaaaggtag	acaggctaaa	ggagagggaac	20100
taagcaagaa	acttacagag	tcagacgttc	tcgctatacg	ctcttcaacc	ttaagccacc	20160
gctccttagg	agaactgtat	ggagtcagtc	aatcaaccat	aacgcgaata	ctacagcgta	20220
agacatggag	acacatttaa	tggtcgagaa	acgaacaggga	cttgccggagg	atggcgcaaa	20280
gtctgtctat	gagcgtttaa	agaacgaccg	tgctccctat	gagacacgcg	ctcagaattg	20340
gcctcaatat	accatcccat	catgtgtccc	taaggactcc	gataacgcct	ctacagatta	20400
tcaaaactccg	tggaagcccg	tggtgcgtcg	tggtctgaac	aatctagcct	ctaagctcat	20460
gctggctcta	ttccctatgc	agacttggat	gcgacttact	atatctgaat	atgaagcaaa	20520
gcagttactg	agcgactccg	atggactcgc	taaggtcgat	gagggcctct	cgatggtaga	20580
gcgtatcatc	atgaactaca	ttgagtctaa	cagttaccgc	gtgactctct	ttgaggctct	20640
caaacagtta	gtcgtagctg	gtaacgtcct	gctgtacctc	ccggaaccgg	aagggtcaaa	20700
ctataatccc	atgaagctgt	accgattgtc	ttcttatgtg	gtccaacgag	acgcattccg	20760
caacgttctg	caaattgtga	ctcgtgacca	gatagctttt	gggtcctccc	ctgaggacat	20820
ccgtaaggct	gtagaaggta	aaggtgggtga	gaagaaagct	gatgagacaa	tcgacgtgta	20880
cactcacatc	tatctggatg	aggactcagg	tgaatacctc	cgatacgaag	aggtcggagg	20940
tatggaagtc	caaggctccg	atgggactta	tcctaaagag	gcttgcccat	acatcccgat	21000
tcggatggtc	agactagatg	gtgaatccta	cggtcgttcg	tacattgagg	aatacttagg	21060
tgacttacgg	tcctctgaaa	atctccaaga	ggctatcgtc	aagatgtcca	tgattagctc	21120
taagggtatc	ggcttagtga	atcctgctgg	tatcccacga	ccacgcgcag	tgaccaaagc	21180
tcagactggg	gacttcgtta	ctggctcgtc	agaagacatc	tcgttctccc	aactggagaa	21240
gcaagcgagc	tttactgtag	ctaaagccgt	aagtgaacgt	atcgaggctc	gcctttcgtt	21300
tgcccttatg	ttgaactctg	cggttcagcg	tacaggtgaa	cggtgacccg	ccgaagagat	21360
tcggtatgta	gcttctgaac	ttgaagatac	tttaggtggg	gtctactcta	tcctttctca	21420
agaattacaa	ttgcctctgg	tacgagtgtc	cttgaagcaa	ctacaagcca	cgcaacagat	21480
tcctgagtta	cctaaggaag	ccgtagagcc	aaccattagt	acaggctctg	aagcaattgg	21540
tcgaggacaa	gaccttgata	agctggagcg	gtgtgtcact	gcgtgggctg	cactggcacc	21600
tatgcgggac	gaccttgata	ttaaccttgc	gatgattaag	ttacgtattg	ccaacgctat	21660
cggtattgac	actctcggta	ttctactcac	cgaagaacag	aagcaacaga	agatggccca	21720
acagtcctatg	caaatgggta	tggaataatg	tgctgctcgc	ctggctcaag	gtatggctgc	21780
acaagctaca	gcttccactg	aggctatggc	tgctgccgct	gattccgtag	gtttacagcc	21840
gggaatttaa	tacgactcac	tatagggaga	cctcatcttt	gaaatgagcg	atgacaagag	21900
gttgaggtcc	tcggtctccc	tgtagtccaa	ctttaaggag	acaataataa	tggtgaatc	21960
taatgcagac	gtatatgcat	cttttggcgt	gaactccgct	gtgatgtctg	gtgggtccgt	22020
tgaggaaacat	gagcagaaca	tgctggctct	tgatgttgct	gcccgtagtg	gcgatgatgc	22080
aatcgagtta	gcgtcagacg	agaacgtgac	ctgtatgaca	actctgaccc		22140
gttcgggtcaa	gaggtatgac	aaggccgcct	tcagggtcgt	atcggtgatg	gctctgagcc	22200
gaccgatgtg	gacactggag	aagaaggcgt	tgagggcacc	gaagggtccg	aagagtttac	22260
cccactgggc	gagactccag	aagaactggg	agctgcctct	gagcaacttg	gtgagcacga	22320
agaggggcttc	caagagatga	ttaacattgc	tgctgagcgt	ggcatgagtg	tcgagaccat	22380
tgaggctatc	cagcgtgagt	acgaggagaa	cgaagagttg	tcgcgcgagt	cctacgctaa	22440
gctggctgaa	attggctaca	cgaaggcttt	cattgactcg	tatatccgtg	gtcaagaagc	22500
tcgtgtggag	cagtacgtaa	acagtgctcat	tgagtacgct	gggtggctcg	aacgttttga	22560
tgactgtat	aaccaccttg	agacgcacaa	ccctgaggct	gcacagtcgc	tgataaatgc	22620
gttgaccaat	cgtgacttag	cgaccgttaa	ggctatcatc	aacttggctg	gtgagtctcg	22680
cgctaaggcg	ttcggctcgt	agcccaactcg	tagtgtgact	aatcggtgct	ttccggctaa	22740
acctcaggct	accaagcgtg	aaggctttgc	ggaccgtagc	gagatgatta	aagctatgag	22800
tgaccctcgg	tatcgcacag	atgccaaacta	tcgtcgtcaa	gtcgaacaga	aagtaatcga	22860
ttcgaacttc	tgatagactt	cgaataat	acgactcact	atagggagac	cacaacgggt	22920
tcctctctaga	aataattttg	tttaacttta	agaaggagat	atacatatgg	ctagcatgac	22980
tggtggacag	caaatgggta	ctaaccgaag	taaagggtga	gttgctgctg	gagataaaact	23040
ggcgttgttc	ttgaaggtat	ttggcgggtga	agtcctgact	gcgttcgctc	gtacctccgt	23100
gaccacttct	cgccacatgg	tacgttccat	ctccagcggg	aaatccgctc	agttccctgt	23160
tcctgggtcgc	actcaggcag	cgtatctggc	tcggggcgag	aacctcgacg	ataaacgtaa	23220
ggacatcaaa	cacaccgaga	aggtaatcac	cattgacggg	ctcctgacgg	ctgacgttct	23280
gatttatgat	attgaggacg	cgatgaacca	ctacgacggt	cgctctgagt	atacctctca	23340
gttggggtgaa	tctctggcga	tggtgcggga	tggtgcgggt	ctggctgaga	ttgccgggtc	23400
gtgtaacgtg	gaaagcaaat	ataatgagaa	catcgagggc	ttaggtactg	ctaccgtaat	23460
tgagaccact	cagaacaagg	ccgcacttac	cgaccaagtt	gcgtgggta	aggagattat	23520
tcgggctctg	actaaggctc	gtcgggctct	gaccaagaac	tatgttccgg	ctgctgacgg	23580
tggtttctac	tgtagcccg	atagctactc	tgcgattctg	gcagcactga	tgccgaacgc	23640
agcaaaactac	gctgctctga	ttgacctga	gaagggttct	atccgcgaac	ttatgggctt	23700
tgagggttga	gaagttccgc	acctcaccgc	tggtgggtgct	ggtagccgctc	gtgagggcac	23760
tactggctcag	aagcacgtct	tcctcgccaa	taaagggtgag	ggtaagtgtca	aggttgctaa	23820
ggacaacggt	atcggtctgt	tcctgcaccg	ctctgcggta	ggtagctgtta	agctgcgtga	23880
cttggtctctg	gagcgcgctc	gccgtgctaa	cttccaagcg	gaccagatta	tcgctaagta	23940
cgcaatgggc	cagcgtgggc	ttcgcccaga	agctgctggg	gcagtgggtt	tcaaaagtga	24000
gtaatgctgg	gggtggcctc	aacggctcgt	gctagtcccg	aagaggcgag	tgtaacttca	24060
acagaagaaa	ccttaacgcc	agcacaggag	gccgcacgca	cccgcgctgc	taacaaagcc	24120

-continued

cgaaaggaag	ctgagttggc	tgtgtgccacc	gctgagcaat	aactagcata	acccttggg	24180
gcctctaacc	gggtottgag	gggttttttg	ctgaaaggag	gaactatatg	cgctcatcag	24240
atatgaacgt	tgagactgcc	gctgagttat	cagctgtgaa	cgacattctg	gcgtctatcg	24300
gtgaacctcc	ggtatcaacg	ctggaagggtg	acgctaaccg	agatgcagcg	aacgctcggc	24360
gtattctcaa	caagattaac	cgacagattc	aatctcgtgg	atggacgttc	aacattgagg	24420
aaggcataac	gctactacct	gatgtttact	ccaacctgat	tgtatacagt	gacgactatt	24480
tatccctaata	gtctacttcc	ggtcaatcca	tctacgttaa	cggaggtggc	tatgtgtatg	24540
accgaacgag	tcaatcagac	cgctttgact	ctgggtattac	tgtgaacatt	attcgtctcc	24600
gcgactacga	tgagatgcct	gagtgcttcc	gttactggat	tgtcaccacg	gcttcccgtc	24660
agttcaacaa	ccgattcttt	ggggcaccgg	aagtagaggg	tgtactccaa	gaagaggaag	24720
atgagggtag	acgtctctgc	atggagtatg	agatggacta	cgggtgggtac	aatatgctgg	24780
atggagatgc	gttcaactct	ggtctactga	ctcgctaaca	ttaataaata	aggaggtctt	24840
aatggcactc	attagccaat	caatcaagaa	cttgaagggt	ggtatcagcc	aacagcctga	24900
catccttcgt	tatccagacc	aagggtcacg	ccaagttaac	ggttgggtct	cggagaccga	24960
gggctcccaa	aagcgtccac	ctcttgtttt	cttaaatata	cttggagaca	acggtgctgt	25020
aggccaagct	cgctacatcc	acctgattaa	cggagatgag	cacgaacagt	attacgtgtg	25080
gttcaactgg	acgggaatcc	gagtggttca	cctttctggg	aacgagaagc	aagtttaggt	25140
tctaacgggt	tccaactaca	tcaagaccgc	taatccacgt	aacgacctgc	gaatgggtac	25200
tgtagcagac	tatacgttca	tcggttaaccg	taacgtttgt	gcacagaaga	acacaaagtc	25260
tgtcaactta	ccgaattaca	accctaatac	agacggattg	attaacgttc	gtgggtgggtca	25320
gtatggtagg	gaactaatgt	tacacattaa	cggtaaaagac	gttgcgaagt	ataagatacc	25380
agatggtagt	caacctgaac	acttaacaaa	tacggatgcc	caatgggttag	ctgaagagtt	25440
agccaaagcag	attgcgacta	acttgtctga	tgggactgta	aatgtagggc	aaggggttcat	25500
ccatgtgacc	gcacctagtgt	gtcaacagat	tgaactcctc	acgactaaag	atggctacgc	25560
agaccagtgt	attaacctgt	tgaccacta	cgctcagtcg	ttctctaagc	tgccacctaa	25620
tgtccttaac	ggctacatgg	tgaaaatcgt	aggggacgcc	tctaagctcg	ccgaccagta	25680
ttacgttcgg	tatgacgtcg	agcggaaaagt	tggactgag	actttagggt	ggaacactga	25740
ggaccaagtt	ctatgggaaa	ccatgccaca	cgctcttgtg	cgagccgctg	acggtaaatt	25800
cgacttcaag	tggcttgagt	ggctctcctaa	gtcttgggtg	gacgttgaca	ccaacctctg	25860
gcctctcttt	gttgggtcaa	gtattaacga	tgtgttcttc	ttccgtaacc	gcttaggatt	25920
ccttagtggtg	gagaacatca	tattgagtcg	tacagccaaa	tacttcaact	tctacctctg	25980
gtccattgcg	aaccttagtg	atgacgaccc	tatagacgta	gctgtgagta	ccaaccgaat	26040
agcaatcctt	aagtacgcgc	ttcgttcttc	agaagagtta	ctcatctggt	ccgatgaagc	26100
acaattcgtc	ctgactgcct	ccggtactct	cacatctaag	tcggttgagt	tgaacctaac	26160
gaccacagtt	gacgtacagg	accgacgcgag	accttttggg	attgggcgta	atgtctactt	26220
tgtcagtcgc	aggtccagct	tcacgtccat	ccacaggtag	tacgctgtgc	aggatgtcag	26280
ttccgttaag	aatgctgagg	acattacatc	acacgttctc	aactacatcc	ctaattgggtg	26340
gttcagtat	tgccggaagt	gtacggaaaa	cttctgtctg	gtactatctc	acggggagcc	26400
tagtaaaatc	ttcatgtaca	aattctctgt	cctgaacgaa	gagttaaggc	aacagtcgtg	26460
gtctcattgg	gactttgggg	aaaaacgtaca	ggttctagct	tgtcagagta	tcagctcaga	26520
tatgtatgtg	attcttccga	atgagttcaa	tacgttctca	gctagaatct	ctttcactaa	26580
gaacgccatt	gacttacagg	gagaacccta	tcgtgccttt	atggacatga	agattcgata	26640
cacgattcct	agtggaaact	acaacgatga	cacattcact	acctctattc	atattccaac	26700
aatttatggt	gcaaaccttc	ggaggggcga	aatcactgta	tgggagcctg	atggtaagat	26760
aaccgtgttt	gagcaacctc	cggtctgggtg	gaatagcgac	ccttggctga	gactcagcgg	26820
taacttggag	ggacgcctgg	gttcaacttg	gttcaacatt	aacttcgtat	atgagttctc	26880
taagttctct	atcaagcaga	ctgcgcacga	cggtctctac	tcacaggaag	acattgggag	26940
cttacagtta	cgccgagcgt	gggttaacta	cgagaacctc	ggtagctttg	acatttatgt	27000
tgagaaccaa	tcgtctaaact	ggaagtacac	aatggctggg	gcccgattag	gctctaacac	27060
tctgagggtc	gggagactga	acttagggac	cggaacaat	cgattccctg	tggttggtaa	27120
cgccaaagtc	aacactgtat	acattctgtc	agatgagact	accctctcga	acatcattgg	27180
gtgtgggtcg	gaaggtaaact	acttacggag	aagttccggt	atttaattaa	atattctccc	27240
tgtgtgtggt	cgaaattaat	acgactcact	atagggagaa	caatacagct	acgggagggt	27300
tttcttatga	tgactataag	acctactaaa	agtacagact	ttaggttat	cactccggct	27360
caccatgaca	ttcttgaagc	taaggctgct	ggtattgagc	cgagtttccc	tgatgcttcc	27420
gagtggtgca	cgttgagcct	ctatgggttc	cctctagcta	tcggtggtaa	ctgcggggac	27480
cagtgctggg	tcgttacgag	cgaccaagtg	tggcgactta	gtggaaaagg	taagcgaaag	27540
ttccgttaagt	taatcatgat	gtatcgcgat	aagatgcttg	agaagtatga	tactctttgg	27600
aattacgtat	gggtaggcaa	tacgtcccac	attcgtttcc	tcaagactat	cggtgcggta	27660
ttccatgaag	agtacacacg	agatgggtcaa	tttcagttat	ttacaatcac	gaaaggagga	27720
taaccatatg	tgttggggcag	ccgcaataacc	tatcgctata	ctggcgctc	aggctatcag	27780
tggtcagaac	gctcaggcca	aaatgattgc	cgctcagacc	gctgctgggc	gtcgtcaagc	27840
tatggaaatc	atgaggcaga	cgaacatcca	gaatgctgac	ctatcgttgc	aagctcgaag	27900
taactttag	gaagcgtccg	ccgagttgac	ctcacagaac	atgcagaagg	tccaagctat	27960
tgggtctatc	cgagcggcta	tcggagagag	tatgcttgaa	gggttctcaa	tggaccgcat	28020
taagcgagtc	acagaaggac	agttcattcg	ggaagccaat	atggtaactg	agaactatcg	28080
ccgtgactac	caagcaatct	tcgcacagca	acttggtggg	actcaagtg	ctgcaagtca	28140
gattgacgaa	atctataaga	cggaacagaa	acagaagagt	aagctacaga	tggttctgga	28200
cccactggct	atcatggggg	cttccgctgc	gagtgcttac	gcatecgggtg	cgttcgactc	28260
taagtcacac	actaaggcac	ctattgttgc	cgctaaagga	accaagacgg	ggaggtaatg	28320
agctatgagt	aaaattgagt	ctgccttca	agcggcacaa	ccgggactct	ctcggttacg	28380
tgggtgggtc	ggaggtatgg	gctatcgtgc	agcaaccact	caggccgaac	agccaagggtc	28440
aagcctattg	gacaccattg	gtcggttcgc	taaggctggg	gccgatattg	ataccgctaa	28500
ggaacaacga	gcacgagacc	tagctgatga	acgctctaac	gagattatcc	gtaagctgac	28560
ccctgagcaa	cgtcgagaag	ctctcaacaa	cgggaccctt	ctgtatcagg	atgacccata	28620
cgctatggaa	gcactccgag	tcaagactgg	tcgtaacgct	gcgtatcttg	tggacgatga	28680

-continued

cggtatgcag	aagataaaag	agggtgtctt	ccgtactcgc	gaagagatgg	aagagtatcg	28740
ccatagtcgc	cttcaagagg	gcgcataagg	atacgcctgag	cagttcggca	tcgaccctga	28800
ggacgttgat	tatcagcgtg	gtttcaacgg	ggacattacc	gagcgtaca	tctcgctgta	28860
tggtgcgcac	gataacttct	tgagccagca	agctcagaag	ggcgcctatca	tgaacagccg	28920
agtggaaactc	aacggtgtgc	ttcaagaccc	tgatatgctg	cgtcgctccag	actctgctga	28980
cttcttttag	aagtataatcg	acaacggctct	ggttactggc	gcaatcccat	ctgatgctca	29040
agccacacag	cttataagcc	aagcgcttcag	tgacgcttct	agccgtgctg	gtggtgctga	29100
cttctgtag	cgagtcgggtg	acaagaagg	aacacttaac	ggagccacta	cgacttaccg	29160
agagttgatt	ggtgaggaac	agtggaaacgc	tctcatggtc	acagcacaac	gttctcagtt	29220
tgagactgac	gcgaagctga	acgagcagta	tcgcttgaag	attaactctg	cgctgaacca	29280
agaggaccca	aggacagctt	gggagatgct	tcaaggatc	aaggctgaac	tagataaagg	29340
ccaacctgat	gagcagatga	caccacaacg	tgagtggcta	atctccgcac	aggaacaagt	29400
tcagaatcag	atgaacgcac	ggacgaaagc	tcaggccaag	gctctggacg	attccatgaa	29460
gtcaatgaac	aaacttgacg	taatcgacaa	gcaattccag	aagcgaatca	acgggtgagtg	29520
gggtctcaacg	gattttaagg	atatgccagt	caacgagaac	actggtgagt	tcaagcatag	29580
cgatatggtt	aactacgcca	ataagaagct	cgctgagatt	gacagtatgg	acattccaga	29640
cggtgccaaag	gactgctatga	agttgaagta	ccttcaagcg	gactctaagg	acggagcatt	29700
ccgtacacgc	atcggaacca	tggtcactga	cgctgggtcaa	gagtggtctg	ccgctgtgat	29760
taacggtaag	ttaccagaac	gaacccagc	tatggatgct	ctgcgcagaa	tccgcaatgc	29820
tgaccctcag	ttgattgctg	cgctataccc	agaccaagct	gagctattcc	tgacgatgga	29880
catgatggac	aagcagggtg	ttgacctca	ggttattctt	gatgccgacc	gactgactgt	29940
taagcgggtcc	aaagagcaac	gctttgagga	tgataaagca	ttcgagtctg	cactgaatgc	30000
atctaaggct	cctgagattg	cccgatgccc	agcgtaactg	cgcgaaatctg	cacgtaagat	30060
ttatgactcc	gttaagtatc	gctcggggaa	cgaaagcatg	gctatggagc	agatgaccaa	30120
gttctttaag	gaattctacct	acacgttcac	tggtgatgat	gttgacggtg	ataccgttgg	30180
tytgattcct	aaagaataga	tgcaagttaa	ctctgacccg	aaatcatggg	agcaaggctg	30240
ggatattctg	gaggaagcac	gtaagggaat	catttgcgagc	aacccttgga	taaccaataa	30300
gcaactgacc	atgtattctc	aaggtgactc	catttacctt	atggacacca	caggtcaagt	30360
cagagtcgga	tacgacaaag	agttactctc	gaaggtctgg	agtgagaacc	agaagaaact	30420
cgaaagagaaa	gctcgtgaga	aggctctggc	tgatgtgaac	aagcgcagac	ctatagttgc	30480
cgctacgaag	gcccgtgaag	ctgctgctaa	acgagtcgga	gagaaacgta	aacagactcc	30540
taagttcatc	tacggacgta	aggagtaact	aaaggctaca	taaggaggcc	ctaaatggat	30600
aagtagcata	agaacgtacc	aagtgattat	gatggtctgt	tccaaaaggc	tgctgatgcc	30660
aaacgggtctg	tcgccaagga	tttactgtaa	gtcgcttgga	cagaatcacg	atttgtgcct	30720
acagcaaaat	ctaagactgg	accattaggg	atgatgcaat	ttaccaaggc	aaccgctaag	30780
gcccctcggtc	tgcgagttac	cgatggtcca	gacgcagacc	gactgaaccc	tgagttagct	30840
attaatgctg	ccgctgaagc	cttgcaggtt	ctggtaggga	agtttgatgg	cgatgaactc	30900
aaagctgccc	ttgcgtacaa	ccaaggcgag	ggacgcttgg	gtaatccaca	acttgaggcg	30960
tactctaagg	gagactctgc	atcaatctct	gaggaggagc	gtaactacat	gcgtaacctt	31020
ctggatgttg	ctaatgcaac	tatggctgga	cagttggaaa	cttttggtgg	cataacccca	31080
aagggtaaa	gcattccggc	tgaggtagga	ttggctggaa	tggttcacaa	gcagaaagta	31140
acacaggaac	ttcctgagtc	cacaagtttt	gacgttaagg	gtatcgaaac	ggaggctacg	31200
gcgaaaacct	tcgccaagga	cttttgggag	acccacggag	aaacacttga	cgagtacaac	31260
agtcgttcaa	ccttcttcgg	attcaaaaat	gctgcggaag	ctgaactctc	caactcagtc	31320
gctgggtagg	cttccgctg	tggtcgtctc	gataatggtt	ttgatgtgtt	taagacacc	31380
attacgcgga	ctcgcgtgaa	ctctcacatc	tggaactccag	aggagttaga	gaagattcga	31440
acagagggtta	agaacctgac	gtacatcaac	gttgtaactg	gtggtcccc	tgagaacctc	31500
gatgacctca	ttaaattggc	tacagagaac	tttgagaatg	actccgcgcg	tgccgaggct	31560
ggcctagggtg	ccaaactgag	tgctgggtatt	attggtgctg	gtgtggaacc	gcttagctat	31620
gttctctatgg	tcggtgtcac	tggttaagggc	tttaagttaa	tcaataaggc	tcttgtagtt	31680
ggtgccgaaa	gtcgtgctct	gaacgttgca	tccgaaggct	tcggtacctc	cgtagctggt	31740
ggtgacgcag	actatgcggg	tgctgcctta	ggtggctttg	tggttgccgc	aggcatgtct	31800
gcaatcagtg	acgctgtagc	tgctggactg	aaacgcagta	aaccagaagc	tgagttcgac	31860
aatgagttca	tcggtcctat	gatgcgattg	gaagcccgtg	agacagcacg	aaacgccaac	31920
tctgcggacc	tctctcggtg	gaacactgag	aacatgaagt	tgaaagtgta	acataatggt	31980
gtcccttatg	aggacttacc	aacagagaga	ggtgcgctgg	tggttacatga	tggtccggtt	32040
ctaagtgcac	gcaacccaat	caaccctaag	actctaaaag	agttctccga	ggttgacctt	32100
gagaaggctg	cgcgaggaa	caaaactggc	gggttcaccg	agatttgctt	gaagaccttg	32160
gggtctgacg	atgctgacat	ccgtagagtg	gctatcgacc	tcggttcgctc	tctactggtt	32220
atgcagctcg	gtgcctcagg	taagttcggt	gcaacagctt	ctgacatcca	tgagagactt	32280
catggtactg	accagcgtac	ttataatgac	ttgtacaaa	caatgtctga	cgctatgaaa	32340
gacctgagtg	tctctactgg	cgccgctaa	atgtcccgtg	aagaaactcg	atacactatc	32400
taccgtagag	cggcactagc	tattgagcgt	ccagaactac	agaaggcact	cactccgtct	32460
gagagaatcg	ttatggacat	cattaaagcgt	cactttgaca	ccaagcgtga	acttatggaa	32520
aaaccagcaa	tattcggttaa	cacaaaaggct	gtgagtatct	tccctgagag	tcgccacaaa	32580
ggtacttacg	ttcctcacgt	atatgaccgt	catgccaaag	cgctgatgat	tcaacgctac	32640
ggtgccgaag	gtttgcagga	agggattgcc	cgctcatgga	tgaacagcta	cgtctccaga	32700
ctgaggtca	aggccagagt	cgatgagatg	cttaaggaat	tacacggggt	gaaggaagt	32759

SEQ ID NO: 2 moltype = DNA length = 836
 FEATURE Location/Qualifiers
 source 1..836
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 2
 tttaagaagg agatatacat atgaaatact tattgccgac gccgcgcgct gggttgcctt 60

-continued

```

tggttagcagc ccagcctgca atggccatgg ctctgcccac aacgggcaag ccgactgcta 120
agcagggtggg tgattgggct atcaatctta tcggctctgg cgtagatgtg gatgggtact 180
atgggcgccca atgttgggat ttgccgaatt acatttttaa ccgttatgtg aattttaaga 240
ctcctgggaa tgctcgtgat atggcctggg atcgctatcc ggagggattt aaagttttct 300
gtaacacttc tgatttcgtc ccaaaacccg gcgatattgc ggtatggacc ggtgggaact 360
ataattgga tacttggggg cactactggca ttgtagtgtg accgtcgaca aaatcctatt 420
tttactcggg tgacccaaac tggacaacatt caaatcgtg ttaggggtcg ccagcggcaa 480
aaattaaaca ttcttatttc ggtgtgactc atttcgtccg tccagcatac aaggcggagc 540
cgaagcctac accacctaag cgtcccgca gctatcgca aacaggcaca atgactgtca 600
cagtggacgc ccttaatgtg cgcctgctgc cgaacactag cggagaaatc gttgccgtct 660
acaagcgcgg ggagtccttc gattacgaca cagtcattat cgacgtaaat gggtagtgtg 720
gggtgtcgtg cattggaggt agcggaaagc gcaattacgt ggctaccgga gcgacaaagg 780
acgggaagcg ctttggaac gcgtggggaa cttttaata agtcgaccgg ctgcta 836

```

```

SEQ ID NO: 3      moltype = DNA length = 21
FEATURE          Location/Qualifiers
source          1..21
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 3
gtcgaccggc tgctaacaaa g 21

```

```

SEQ ID NO: 4      moltype = DNA length = 43
FEATURE          Location/Qualifiers
source          1..43
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 4
atgtatatct ccttcttaaa gttaaacaaa attatttcta gag 43

```

```

SEQ ID NO: 5      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source          1..22
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 5
tatagtcttg tcgggtttcg cc 22

```

```

SEQ ID NO: 6      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source          1..22
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 6
ctggcctggt gaacaagtct gg 22

```

```

SEQ ID NO: 7      moltype = DNA length = 37
FEATURE          Location/Qualifiers
source          1..37
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 7
accgctgagc aataataata cgactcacta tagggag 37

```

```

SEQ ID NO: 8      moltype = DNA length = 34
FEATURE          Location/Qualifiers
source          1..34
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 8
aaggggttat gctaggcaaa aaaccctca agac 34

```

```

SEQ ID NO: 9      moltype = DNA length = 30
FEATURE          Location/Qualifiers
source          1..30
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 9
tctcacagtg tacggaccta aagttcccc 30

```

```

SEQ ID NO: 10     moltype = DNA length = 30
FEATURE          Location/Qualifiers
source          1..30
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 10

```

-continued

attacgcgat gacagtagac aacctttccg	30
SEQ ID NO: 11	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 11	
tgcagcaata ccgaaaggt tgtctactgt	30
SEQ ID NO: 12	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 12	
atatgtctcc tcatagatgt gcctatgtgg	30
SEQ ID NO: 13	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 13	
acttgtgact ccacataggc acatctatga	30
SEQ ID NO: 14	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 14	
gaataacctg agggtaacata cctgcttgt	30
SEQ ID NO: 15	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 15	
gacatgatgg acaagcaggg tattgacct	30
SEQ ID NO: 16	moltype = DNA length = 30
FEATURE	Location/Qualifiers
source	1..30
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 16	
agggacacag agagacactc aaggaacac	30
SEQ ID NO: 17	moltype = DNA length = 27
FEATURE	Location/Qualifiers
source	1..27
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 17	
acttgtgact ccacataggc acatcta	27
SEQ ID NO: 18	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 18	
tagtgagtcg tattattatt gctcagcggg ggc	33
SEQ ID NO: 19	moltype = DNA length = 33
FEATURE	Location/Qualifiers
source	1..33
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 19	
tgaggggttt tttgcctagc ataaccctt ggg	33
SEQ ID NO: 20	moltype = DNA length = 30
FEATURE	Location/Qualifiers

-continued

source 1..30
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 20
gaataacctg agggtaata cctgcttgt 30

SEQ ID NO: 21 moltype = DNA length = 22
FEATURE Location/Qualifiers
source 1..22
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 21
gaaaccttaa cgccagcaca gg 22

SEQ ID NO: 22 moltype = DNA length = 22
FEATURE Location/Qualifiers
source 1..22
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 22
accgatagac gccagaatgt cg 22

SEQ ID NO: 23 moltype = DNA length = 916
FEATURE Location/Qualifiers
source 1..916
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 23
taatacgact cactataggg agaccacaac ggtttccctc tagaaataat ttgtttaac 60
ttaagaagg agatatacat atgagcaaag gtgaagaact gttaccggc gttgtgccga 120
ttctgggtga actggatggc gatgtgaacg gtcacaaatt cagcgtgcgt ggtgaagggtg 180
aaggcgatgc cagcattggc aaactgacgc tgaaatttat ctgaccacc ggcaaaactgc 240
cggtgccgtg gccgacgctg gtgaccaccc tgacctatgg cggtcagtg tttagtcgct 300
atccggatca catgaaacgt cagcatttct ttaaatctgc aatgccggaa ggctatgtgc 360
aggaacgtac gattagcttt aaagatgatg gcaaatataa aacgcgcgcc gttgtgaaat 420
ttgaaggcga taccctgggt aaccgcattg aactgaaagg caccgatttt aaagaagatg 480
gcaatatcct gggccataaa ctggaataca actttaatag ccataatggt tatattacgg 540
cggataaaca gaaaaatggc atcaaaagcga attttaccgt tcgccataac gttgaagatg 600
gcagtgtgca gctggcagat cattatcagc agaatacccc gattggtgat ggtccgggtgc 660
tgctgccgga taatcattat ctgagcacgc agaccgttct gtctaaagat ccgaacgaaa 720
aaggcacgcg ggaccacatg gttctgcacg aatatgtgaa tgcggcaggt attacgtgga 780
gcatccgcga gttcgaaaaa taagtcgacc ggctgctaac aaagcccga aggaagctga 840
gttggtgct gccaccgtg agcaataact agcatacccc cttggggcct ctaaacgggt 900
cttgaggggt tttttg 916

SEQ ID NO: 24 moltype = DNA length = 21
FEATURE Location/Qualifiers
source 1..21
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 24
cgctcggcta tttcacatac t 21

SEQ ID NO: 25 moltype = DNA length = 22
FEATURE Location/Qualifiers
source 1..22
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 25
ggtttctggc gttaaaccaa tc 22

SEQ ID NO: 26 moltype = DNA length = 22
FEATURE Location/Qualifiers
source 1..22
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 26
ccgtctggtt tctgacgata tt 22

SEQ ID NO: 27 moltype = DNA length = 20
FEATURE Location/Qualifiers
source 1..20
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 27
cgcagcagga tctggttaat 20

SEQ ID NO: 28	moltype = DNA length = 21		
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 28			
gtgcgcctct tctttgaatt t			21
SEQ ID NO: 29	moltype = DNA length = 19		
FEATURE	Location/Qualifiers		
source	1..19		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 29			
tcgatggggtc gtccattac			19
SEQ ID NO: 30	moltype = DNA length = 21		
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 30			
aatatggcat ccgtgggtatc c			21
SEQ ID NO: 31	moltype = DNA length = 21		
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 31			
gttagcgctcg aggaactctt t			21
SEQ ID NO: 32	moltype = DNA length = 22		
FEATURE	Location/Qualifiers		
source	1..22		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 32			
cctgttcgat gagaggaatc tg			22
SEQ ID NO: 33	moltype = DNA length = 21		
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 33			
ccttagctgg gtgagtgatt t			21
SEQ ID NO: 34	moltype = DNA length = 6458		
FEATURE	Location/Qualifiers		
source	1..6458		
	mol_type = other DNA		
	organism = synthetic construct		
SEQUENCE: 34			
gggtcacctt cgggtgggct tttctgcgtt tatacagaga cctcagccgc tacaggcgcc 60			
gtccattcgc ccattcaggc tgcgcgaactg ttgggaagggt cgatcgggtgc gggccctcttc 120			
gctattacgc cagctggcgca aaggggggatg tgcgtcaagg cgattaagtt gggtaacgcc 180			
agggttttcc cagtcacgac gttgtaaaaac gacggccagtt gagcgcgcggt aatacgaactc 240			
actatagggtc gaattgggta ccggggccccc cctcgaggtc ctccagcttt tgttcccttt 300			
agtgagggtt aattgcgcgc ttggcgtaaat catggtcata gctgtttctc gtgtgaaagg 360			
tctcaatcta caacagttag aattacgggtg tttactagta ttatacctag gactgagcta 420			
gctgtcaagg atatatctgc ctctctcgct cactgactcg ctacgctcgg tcggttcgact 480			
gcggcgagcg gaaatggctt acgaacgggg cggagatttc ctggaagatg ccaggaagat 540			
actaacagc gaagtgaagc gcgcggcgcca aagccggttt tccattagcct ccgccccct 600			
gccaagcattc acgaaatctg agcctcaaat cagtggtggc gaacacccgac aggactataa 660			
agataccagg cgtttccccc ttggcggtccc ctcgtgcgct ctctctgttc tgcctttcgg 720			
tttaccgggtg tcatctccgt gttatggccg cgtttgtctc attccacgcg ttacactcag 780			
ttccgggtag tcatctctgc ccaagcttga ctgtatgcac gaaccccccg ttacgttcga 840			
ccgctgcgct tcattccgcta actatcgctt tgagttccaac ccggaagaag atgcaaaagc 900			
accactggca gcagccactg gtaattgatt tagaggagtt agtcttgaag tcatgcgcgcg 960			
gttaaggcta aactgaaagg acaagttttt gtgactgcgc tccttccaage cagttacctc 1020			
ggttcaaga gttggtagct cagagaacctc tcgaaaaaac gccctgcgaag gcggtttttt 1080			
cgttttcaga gcaagagatt acgcgcagac caaacgatc tcaagaagat catcttatta 1140			
atcagataaa atattttctag atttctagtc aatttatctc ttcaaatgta gcacctgaag 1200			
tcaqccccat acgatataaa ttgttactag tcttttqatt ctcaccaata aaaaacqccc 1260			

-continued

ggcggcaacc	gagcgttctg	aacaaatcca	gatggagtcc	tgaggtcatt	actggatcta	1320
tcaacaggag	tccaagcgag	ctcgatatca	aattacgccc	cgccctgcca	ctcatcgag	1380
tactgttgta	attcattaag	cattctgccc	acatggaagc	catcacaaac	ggcatgatga	1440
acctgaatcg	ccagcgccat	cagcaccttg	tcgccttgcc	tataaatatt	gcccattggtg	1500
aaaacggggg	cgaagaagtt	gtccatattg	gccacgttta	aatcaaaact	ggtgaaactc	1560
acccagggat	tggctgagac	gaaaaacata	ttctcaataa	accccttagg	gaaataggcc	1620
agggtttcac	cgtaacacgc	cacatcttgc	gaatatatgt	gtagaaactg	ccggaatcgc	1680
tcgtgggtatt	cactccagag	cgatgaaaac	gtttcagttt	gctcatggaa	aacgggtgaa	1740
caagggtgaa	cactatccca	tatcaccagc	tcaccgtctt	tcattgccat	acgaaattcc	1800
ggatgagcat	tcatcaggcg	ggcaagaatg	tgaataaagg	ccggataaaa	cttgtgctta	1860
tttttcttta	cggtctttaa	aaaggccgta	atatccagct	gaacggctcg	gttataggta	1920
cattgagcaa	ctgactgaaa	tgctctaaaa	tggtctttac	gatgccattg	ggatatatca	1980
acgggtggtat	atccagtgat	ttttttctcc	attttagctt	ccttagctcc	tgaaaatctc	2040
gataactcaa	aaaatacgcc	cggtagtgtg	cttatttcat	tatggtgaaa	gttgaaacct	2100
cttacgtgcc	gatcaacgtc	tcattttcgc	cagatatcga	cgctccaccac	aattcagcaa	2160
attgtgaaca	tcatcacgtt	ctcttttccc	tggttgccaa	tggtccattt	tcctgtcagt	2220
aacgagaagg	tcgcgaattc	aggcgctttt	tagactggtc	gtaatgaaat	tccttttaag	2280
aaggagat	atgataaaga	ggagaagga	tctatgagca	tctaccagga	gttcgtcaac	2340
aagtattcac	tgatgaagac	actgcggttc	gagctgatcc	cacagggcaa	gacactggag	2400
aacatcaagg	cccagggcct	gattctggac	gatgagaagc	gggcaaaaaga	ctataagaaa	2460
gccaagcaga	tcattgtata	ataccaccag	ttctttatcg	aggaatctct	gagctccgtg	2520
tgcatcagtg	aggatctgtg	gcagaattac	tcagacgtgt	acttcaagct	gaagaagagc	2580
gacgatgaca	acttcgagaa	ggacttcaag	tcgcgcaagg	acaccatcaa	gaaaagtaca	2640
cgtctgagta	attccagatt	agcgagtaca	tcaaggagtc	cgaaaagtgt	aaaaatctgt	2700
tcaaccagaa	tctgatcgat	gctaagaaag	gccaggagtc	cgacctgac	ctgtggctga	2760
aacagtcata	ggacaattgg	attgaactgt	tcaaggctaa	ctccgatatc	actgatattg	2820
acgagggcact	ggaaatcatc	aagagcttca	agggatggac	cacatacttt	aaaggcttcc	2880
acgagaaccg	caagaacgtg	tactccagca	acgacattcc	tacctccatc	atctaccgaa	2940
tcgtcgatga	caatctgcca	aagttcctgg	agaacaaggc	caaatatgaa	tcctctgaagg	3000
acaaagctcc	cgagggaatt	aattacgaac	agatcaagaa	agatctggct	gagggaactga	3060
cattcgatat	cgactataag	actagcgagg	tgaaccagag	ggtcttttcc	ctggacgagg	3120
tgtttgaagt	cgccaatttc	aacaattacc	tgaaccagtc	cggcattact	aaattcaata	3180
ccatcattgg	cggggaagtt	gtgaacgggg	agaataccaa	gcgcaaggga	attaacgaat	3240
acatcaatct	gtatagccag	cagatcaacg	acaaaactct	gaagaataac	aagatgtctg	3300
tgctgttcaa	acagatcctg	agtgtaccgc	agtcgaagtc	ttttgtcatt	gataaaactgg	3360
aagatgactc	agacgtggct	actaccatgc	agagctttta	tgagcagatc	gccgctttca	3420
agacagtggg	ggaaaaatct	attaaaggaaa	ctctgagctc	gctgttcgat	gacctgaaag	3480
cccagaagct	ggacctgagt	aagatctact	tcaaaaacga	taagagtcct	acagacctgt	3540
cacagcaggt	tcttgatgac	tattccgtga	ttgggaccgc	cgctcctggg	tacattacac	3600
agcagatcgc	tccaaagaa	ctggataatc	cctctaagaa	agagcaggaa	ctgatcgcta	3660
agaaaaacga	gaaggcaaaa	tatctgagtc	tggaacaact	taagctggca	ctggaggagt	3720
tcaacaagca	cagggatatt	gacaaacagt	gccgctttga	ggaaatcctg	gccaaactcg	3780
cagccatccc	catgatcttt	gatgagatcg	cccagaacaa	agacaatctg	gctcagatca	3840
gtattaagta	ccagaaccag	ggcaagaaag	acctgctgca	ggcttcagca	gaagatgacg	3900
tgaaagccat	acaggatctg	ctggaccaga	ccaacaatct	gctgcacaag	ctgaaaatct	3960
tcctatttag	tcagtcagag	gataaggcta	atatcctgga	taaagacgaa	cacttctacc	4020
tggtgttcga	ggaatgttac	ttcgagctgg	caaacattgt	cccctgttat	aacaagatta	4080
ggaactacat	cacacagaag	ccttactctg	acgagaagtt	taaaactgaac	ttcgaaaata	4140
gtaccctggc	caacgggtgg	gataaagaac	aggagcctga	caacacagct	atcctgttca	4200
tcaaggatga	caagtactat	ctgggagtga	tgaataagaa	aaacaataag	atcttcgatg	4260
acaaagccat	taaggagaac	aaaggggaag	gatacaagaa	aatcggtgat	aagctgctgc	4320
ccggcgcaaaa	taagatgcctg	cctaagggtg	ctctcagcgc	caagagtatc	aaattctaca	4380
acccatccga	ggacatcctg	cggattagaa	atcactcaac	acatactaa	aacggggagcc	4440
cccagaaggg	atatgagaaa	tttgagttca	acatcgagga	ttgcaggaa	tttattgact	4500
tctacaagca	gagcatctcc	aaacaccctg	aatggaaagg	ttttggcttc	cggttttccg	4560
acacacagag	atataactct	atcgacgagt	tctaccgca	gggtggaaaat	caggggtata	4620
agctgacttt	tgagaacatt	tctgaaagtt	acatcgacag	cggtgtcaat	caggggaaagc	4680
tgtaacctgt	ccagatctat	aacaaagatt	tttcagcata	cagcaagggc	agaccaaacc	4740
tgcatacact	gtactggaag	gcctgttctg	atgagaggaa	tctgcaggac	gtggtctata	4800
aactgaacgg	agaggccgaa	ctgttttacc	ggaagcagtc	tattcctaag	aaaatcactc	4860
acccagctaa	ggaggccatc	gctaacaaga	acaaggacaa	tcctaagaaa	gagagcggtg	4920
tcgaatacga	tctgattaag	gacaagcggg	tcaccgaaga	taagttcttt	ttccattgtc	4980
caatcaccat	taacttcaag	tcaagcgccg	ctaacaagtt	caacgacgag	atcaatctgc	5040
tgctgaagga	aaaagcaaac	gatgtgcaca	tcctgagcat	tgctcgagga	gagcggcatc	5100
tggtctacta	taccctgggt	gatggcaaa	ggaatatcat	taagcaggat	acattcaaca	5160
tcattggcaa	tgaccggatg	aaaaccaact	accacgataa	actggctgca	atcgagaagg	5220
atagagactc	agctaggga	gactggaaga	aaatcaacaa	cattaaggag	atgaaggga	5280
gctatctgag	ccagggtgtc	catgagattg	caaagctggt	catcgaaatc	aatgccattg	5340
tggtgttcga	ggatctgaac	ttcggttcta	agagggggcg	ctttaagggtg	gaaaaacagg	5400
tctatcagaa	gctggagaaa	atgctgatcg	aaaagctgaa	ttacctgggtg	tttaagata	5460
acgaggtcga	caagaccgga	ggcgtcctga	gagcctacca	gctgacagct	ccctttgaaa	5520
ctttcaagaa	aatgggaaaa	cagacaggca	tcatctacta	tggtccagcc	ggattcactt	5580
ccaagatctg	cccgtgacc	ggctttgtca	accagctgta	ccctaataat	gagtcagtga	5640
gcaagtccca	ggaatttttc	agcaagttcg	ataagatctg	ttataatctg	gacaaggggt	5700
acttcgagtt	ttccttcgat	tacaagaact	tcggcgacaa	ggcggtcaag	gggaaatgga	5760
ccattgcctc	cttcgagatc	cgctgatca	actttcgaaa	ttccgataaa	aaccacaatt	5820

-continued

gggacactag	ggaggtgtac	ccaaccaagg	agctggaaaa	gctgctgaaa	gactactcta	5880
tcgagtatgg	acatggcgaa	tgcatacaagg	cagccatctg	tggcgagagt	gataagaaat	5940
ttttcgccaa	gctgacctca	gtgctgaata	caatcctgca	gatgcggaac	tcaagaccg	6000
ggacagaact	ggactatctg	attagccccg	tggctgatgt	caacggaaac	ttcttcgaca	6060
gcagacaggc	acccaaaaat	atgcctcagg	atgcagacgc	caacggggcc	taccacatcg	6120
ggctgaagg	actgatgctg	ctgggcccga	tcaagaacaa	tacggagggg	aagaagctga	6180
acctgggtcat	taagaacgag	gaatacttcg	agtttgtcca	gaatagaat	aacaaaaggc	6240
cggcgggccc	gaaaaaggcc	ggccaggcaa	aaaagaaaaa	gggatccctac	ccatacgatg	6300
ttccagatta	cgcttatccc	tacgacgtgc	ctgattatgc	atacccatat	gatgtccccg	6360
actatgccta	acaaataaaa	cgaagggtc	agtcgaaaga	ctgggcccctt	cgttttatct	6420
gttggtttgc	ggtgaacgct	ctctactaga	gtcacact			6458

SEQ ID NO: 35 moltype = DNA length = 2486
 FEATURE Location/Qualifiers
 source 1..2486
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 35

tccaggggga	aacgcctggt	atctttatag	tctgtcggg	tttcgccacc	tctgacttga	60
gcgtcgattt	ttgtgatgct	cgtcaggggg	gcggagccta	tggaaaaacg	ccagcaacgc	120
gatcccgca	aattaatacg	actcactata	gggagaccac	aacggtttcc	ctctagaaat	180
aattttgttt	aactttaaga	aggagatata	catatgagca	aaggtgaaga	actgtttacc	240
ggcggttgct	cgattctggt	ggaactggat	ggcgatgtga	acggtcacaa	attcagcgtg	300
cgtggtgaag	gtgaaggcga	tgccacgatt	ggcaaaactga	cgctgaaatt	tatctgcacc	360
acccggcaaac	tgccggtgct	gtggccgacg	ctggtgacca	ccctgacctg	tggcggtcag	420
tgttttagtc	gtctatccgga	tcacatgaaa	cgctcacgatt	tctttaaatc	tgcaatgccg	480
gaaggctatg	tgcaggaaacg	tacgattagc	tttaaatgatg	atggcaataa	taaaacgcgc	540
gccgttgtga	aatttgaagg	cgataccctg	gtgaaccgca	ttgaactgaa	aggcacggat	600
tttaagaag	atggcaatat	ctcgggccc	aaactgggaat	acaactttaa	tagccataat	660
gtttatatta	cgccgggata	acagaaaaat	ggcatcaaac	cgaattttac	cgttcgccat	720
aacgttgaag	atggcagttg	gcagctggca	gatcattatc	agcagaatac	cccgaattgt	780
gatggtccgg	tgtctgtgcc	ggataatcat	tatctgagca	cgcagaccgt	tctgtctaaa	840
gatccgaacg	aaaaaggcac	gcgggaccac	atggttctcg	acgaatatgt	gaatgcggca	900
ggtattacgt	ggagccatgc	gcagttcgaa	aaataagtcg	accggctgct	aacaaagccc	960
gaaaggaagc	tgagttggct	gctgccaccg	ctgagcaata	actagcataa	ccctctgggg	1020
ccctctaaacg	ggtcttgagg	ggttttttgc	tgaagaccaa	ttctgattag	aaaaactcat	1080
cgagcatcaa	atgaaactgc	aattttattca	tatcaggatt	atcaatacca	tatttttgaa	1140
aaagccggtt	ctgtaatgaa	ggagaaaact	caccgaggca	gttccatagg	atggcaagat	1200
cctggtatcg	gtctgcgatt	cgcactcgtc	caacatcaat	acaacctatt	aatttcccct	1260
cgctcaaaaat	aaggttatca	agtgagaaat	caccatgagt	gacgactgaa	tccggtgaga	1320
atggcaaaaag	cttatgcatt	tctttccaga	cttgttcaac	aggccagcca	ttacgctcgt	1380
catcaaaaatc	actcgcacat	accaaaaccg	tattcattcg	tgattgcgcc	tgagcgagac	1440
gaaatcacgcg	tcaaggtgta	aaaggacaat	tacaaacagg	aatcgaatgc	aaccggcgca	1500
ggaacactgc	cagcgcatca	acaatatatt	cacctgaatc	aggatattct	tctaatacct	1560
ggaatgctgt	tttcccgggg	atcgcagtg	tgagtaacca	tgcatcatca	ggagtacgga	1620
taaaatgctt	gatggtcgga	agaggcataa	attccgtcag	ccagtttagt	ctgacctct	1680
catctgtaac	atcattggca	acgctacctt	tgccatgttt	cagaaacaac	tctggcgcat	1740
cggtcttccc	atacaatcga	tagattgtcg	cacctgattg	cccgacatta	tccgagagcc	1800
atttatatccc	atataaatca	gcacatcatg	tggaaattta	tccggtcttc	gagcaagacg	1860
tttcccggtg	aatatggctc	ataaaccccc	ttgtattact	gtttatgtaa	gcagacagtt	1920
tattgtttca	tgatgatata	tttttatctt	gtgcaatgta	acatcagaga	ttttgagaca	1980
caacgtgaga	tcaaaagatc	ttcttgagat	ccttttttct	tgcgcgtaat	ctgctgcttg	2040
caaacaaaaa	aaccaccgct	accagcggtg	gtttgtttgc	cggatcaaga	gctaccaact	2100
ctttttccga	aggtaactgg	cttcagcaga	gcgcagatac	caaatactgt	tcttctagt	2160
tagccgtagt	taggcccacca	cttcaagaac	tctgtagcac	cgctacata	cctcgctctg	2220
ctaactcctg	taccagtgcc	tgtgcccagt	ggcgataagt	cgtgtcttac	cgggttggtg	2280
tcaagacgat	agttaccgga	taaggcgag	cggtcgggct	gaacgggggg	ttcgtgcaca	2340
cagcccagct	tggagcgaaac	gaactgagat	acctacacgc	tgagctatga		2400
gaaagcgcca	cgcttcccga	agggagaaaag	gcggacaggt	atccggttaag	cggcagggtc	2460
ggaacaggag	agcgcacgag	ggagct				2486

SEQ ID NO: 36 moltype = DNA length = 5429
 FEATURE Location/Qualifiers
 source 1..5429
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 36

ttatgacaac	ttgacggcta	catcattcac	tttttcttca	caaccggcac	ggaactcgct	60
cgggtggtgc	ccggtgcatt	ttttaaatat	ccgcgagaaa	tagagttgat	cgtcaaaacc	120
aacattgcga	ccgacggtgg	cgatagggcat	ccgggtgtgtg	ctcaaaagca	gcttcgcctg	180
gctgatacgt	tggctcctgc	gccagcttaa	gacgctaata	cctaactgct	ggcggaaaag	240
atgtgacaga	cgcgacggcg	acaagcaaac	atgctgtgcg	acgctggcga	tatcaaaatt	300
gctgtctgcc	agggtgatcg	tgatgtactg	acaagcctcg	cgtacccgat	tatccatcgg	360
tggatggagc	gactcgttaa	tgccttccat	gcgcgcgagt	aacaattgct	caagcagatt	420
tatcgccagc	agctccgaat	agcgcccttc	cccttgcccg	gcgttaatga	tttgcccaaa	480
caggtcgctg	aaatgcggct	ggtgcgcttc	atccggggca	aagaaccccc	tattggcaaa	540

-continued

tattgacggc	cagttaagcc	attcatgcc	gtaggcgcgc	ggacgaaagt	aaaccactg	600
gtgatcccat	tcgcgagcct	ccggatgacg	accgtagtga	tgaatctctc	ctggcgggaa	660
cagcaaaata	tcaccccgct	ggcaaaacaa	ttctcgtccc	tgatttttca	ccacccctcg	720
accgcgaatg	gtgagattga	gaatataacc	tttcattccc	agcggctcgt	cgataaaaa	780
atcgagataa	ccgttgccct	caatcgccgt	taaaccgcc	accagatggg	cattaaacga	840
gtatcccgcc	agcaggggat	cattttgccg	ttcagccata	cttttcatac	tcccgcctat	900
cagagaagaa	accaattgtc	catattgcat	cagacattgc	cgctactcgc	tcttttactg	960
gctcttctcg	ctaaccaaac	cggtaacccc	gcttattaaa	agcattctgt	aacaaagcgg	1020
gaccaaagcc	atgacaaaa	cgcgtaacaa	aagtgtctat	aatcacggca	gaaaagtcca	1080
cattgattat	ttgcacggcg	tcacactttg	ctatgccata	gcatttttat	ccataagatt	1140
agcggatact	acctgacgct	ttttatcgca	actctctact	gtttctccat	acccgttttt	1200
ttgggctaga	aataattttg	tttaacttta	agaaggagat	atacatatgc	gggggtctca	1260
tcacatcat	catcatggta	tggtagcat	gactgggtga	cagcaaatgg	gtcgggatct	1320
gtacgagaac	ctgtacttcc	agggctcgag	catgggtgag	aagggcgagg	agctgttcac	1380
cggggtgggt	cccactctgg	tcgagctgga	cggcgacgta	aacggccaca	agttcagcgt	1440
gcgcggcgag	ggcgagggcg	atgccaccaa	cggcaagctg	accctgaagt	tcactctgcac	1500
caaccggcaag	ctgcccgtgc	cctggcccac	cctcgtgacc	accctgaact	acggcgttga	1560
gtgcttcagc	cgctaccccg	accacatgaa	gcgcacacgac	ttcttcaagt	ccgccatgcc	1620
cgaaggctac	gtccaggagc	gcacatcag	cttcaaggac	gacggcacct	acaagaccg	1680
cgccgaggtg	aagttcgagg	gcgacacct	ggtgaaccgc	atcgagctga	agggcattcg	1740
cttcaaggag	gacggcaaca	tcttggggca	caagctggag	tacaacttca	acagccacaa	1800
cgtctatatc	accgccgaca	agcagaagaa	cggcatcaag	gccaaacttca	agatccgcca	1860
caacgtggag	cacggcagcg	tgacgtctgc	cgaccactac	cagcagaaca	ccccactcgg	1920
cgacggcccc	gtgctgtctg	ccgacaacca	ctacctgagc	accagtcg	tgctgagcaa	1980
agaccccaac	gagaagcgcg	atcacatggt	cctgctggag	ttcgtgaccg	ccgccgggat	2040
caactcacggc	atggacgagc	tgtacaagta	agaattcgaa	gcttggctgt	ttggcggtat	2100
gagagaagat	tttcagcctg	atacagatta	aatcagaacg	cagaagcggg	ctgataaaac	2160
agaatttgcc	tgggcgcgagt	agcgcggtgg	tcccacctga	ccccatgccg	aactcagaag	2220
tgaaaacgccc	tagcgcgcat	ggtagtgtgg	ggtctcccca	tgcgagagta	gggaaactgcc	2280
aggcatcaaa	taaaacgaaa	ggctcagtcg	aaagactggg	cctttcgctt	tatctgttgt	2340
ttgtcgggtg	acgctctcct	gagtaggaca	aatccgcg	gagcggattt	gaacgttgcc	2400
aagcaacggc	ccggagggtg	gcgggcagga	cgcccgccat	aaactgccag	gcatacaatt	2460
aagcagaagg	ccatccctgac	ggatggcctt	tttgcgttcc	tacaaactct	tttgtttatt	2520
tttctaata	cattcaaaata	tgtatccgct	catgagacaa	taaccctgat	aaatgcttca	2580
ataatattga	aaaaggaaga	gtatgagtat	tcaacatttc	cgtgtcgc	ttattccctt	2640
ttttgcggca	ttttgccttc	ctgtttttgc	tcaaccagaa	acgtgtgtga	aagtaaaaga	2700
tgctgaagat	cagttgggtg	ccagagtggg	ttacatcgaa	ctggatctca	acagcggtaa	2760
gatccttgag	agttttcgcc	ccgaagaacg	ttttccaatg	atgagcactt	ttaaagtctt	2820
gctatgtggc	gcggtattat	cccggtgtga	cgccggggcaa	gagcaactcg	gtcgcgcgat	2880
acactattct	cagaatgact	tggttgagta	ctcaccagtc	acagaaaagc	atcttacgga	2940
tggtcatgaca	gtaagagaat	tatgcagtgc	tgccataacc	atgagtgaata	acactcggcg	3000
caacttactt	ctgacaacga	tcggaggacc	gaaggagcta	accgcttttt	tgcaacaacat	3060
gggggatcat	aggtacccgc	tggtcgttg	ggaaccggag	ctgaatgaag	ccataccaaa	3120
cgacgagcgt	gacaccacga	tgccctcagc	aatggcaaca	acgttgcgca	aactattaac	3180
tgggcaacta	cttactctag	cttcccggca	acaattaata	gactggatgg	aggcgggata	3240
agttgcaggga	ccactttctg	ctccggccct	tcgggtggcg	tggtttattg	ctgataaatc	3300
tgagcccggt	gagcgtgggt	ctcgcgggtat	cattgcagca	ctggggccag	atggtaagcc	3360
ctcccgatc	gtagtattct	acacgacggg	gagtcaggca	actatggatg	aacgaaatag	3420
acagatcgct	gagatagggt	cctcactgat	taagcattgg	taactgtcag	accaagttta	3480
ctcatatata	cttttagattg	atttacgcgc	cctgtagcgg	cgcattaagc	gcggcgggtg	3540
tggtgggttac	gcgcagcgtg	accgctacac	ttgcagcgc	cctagcgc	gctcctttcg	3600
ctttcttccc	ttcctttctc	gccagtttcg	ccggctttcc	ccgtcaagct	ctaaactcgg	3660
ggctcccttt	aggggttccga	tttagtgctt	tacggcacct	cgaccccaaa	aaacttgatt	3720
tggtgtgagg	ttcacgtagt	ggggccatcg	cctgatatag	gggttttcgc	cctttgacgt	3780
tgaggtccac	gttctttaat	agtggactct	tgttccaaac	tggaacaaca	ctcaacccta	3840
tctcgggcta	ttcttttgat	ttataaggga	ttttgcccgt	ttcggcctat	tggttaaaaa	3900
atgagctgat	ttacaaaaaa	tttaacgcga	attttaacaa	aatattaacg	tttacaattt	3960
aaaaggatct	aggtcaagat	cctttttgat	aatctcatga	ccaaaatccc	ttaacgtgag	4020
ttttcgttcc	actgagcgtc	agaccccgta	gaaaagatca	aaggatcttc	ttgagatcct	4080
ttttttctgc	gcgtaactctg	ctgcttgcaa	acaaaaaaac	caccgctacc	agcgggtggt	4140
tggttgccgg	atcaagagct	accaactctt	tttccgaagg	taactggctt	cagcagagcg	4200
cagataccaa	atactgtcct	tctagtgtag	ccgtagttag	gccaccactt	caagaactct	4260
gtagcaccgc	ctacatacct	cgtctgtcta	atcctgttac	cagtggctgc	tgccagtgcc	4320
gataagtcgt	gtcttaccgg	ggtggactca	agacgatagt	taccggataa	ggcgcagcgg	4380
tcgggctgaa	cggggggttc	gtgcacacag	ccagcttgg	agcgaacgac	ctacacgcaa	4440
ctgagatacc	tacagcgtga	gctatgagaa	agcgcacgc	ttcccgagg	gagaaaaggcg	4500
gacaggtatc	cggtaagcgg	cagggtcggga	acaggagagc	gcacgaggga	gcttccaggg	4560
ggaaacgcct	ggatctctta	tagtctctgc	gggtttcgcc	acctctgact	tgagcgtcga	4620
tttttggtgat	gctcgtcagg	ggggcgaggc	ctatggaaaa	acgccagcaa	cgccgctttt	4680
ttacgggtcc	tggtcttttg	ctggcctttt	gctcacatgt	tctttctcgc	gttatccctt	4740
gattctgtgg	ataacgctat	taccgccttt	gagtgagctg	ataccgctcg	ccgcagccga	4800
acgacgcagc	gcagcgagtc	agtgagcggg	gaagcgggaag	agcgcctgat	gcgggtattt	4860
ctccttacgc	atctgtgcgg	tatttcacac	cgcactctggt	gcactctcag	tacaactctg	4920
tctgatgccg	catagttaag	ccagtatata	ctccgctatc	gctacgtgac	tggtgcatgg	4980
ctgcgcccg	acaccccgca	acacccgctg	acggcccgctg	acgggcttgt	ctgctcccg	5040
catccgctta	cagacaagct	gtgaccgtct	ccgggagctg	catgtgtcag	aggtttttcac	5100

-continued

```

cgctcatcacc gaaacgcgcg aggcagaag agatggcgcc caacagtcce ccggccacgg 5160
ggcctgcccac catacccacg ccgaaacaag cgctcatgag cccgaagtgg cgagcccgat 5220
cttccccato ggtgatgtcg gcgatatagg cggcagcaac cgcacctgtg gcgccggtga 5280
tgccggccac gatgcgtccg gcgtagagga tctgctcatg tttgacagct tatcatcgat 5340
gcataatgtg cctgtcaaat ggacgaagca gggattctgc aaacccctatg ctactccgtc 5400
aagccgtcaa ttgtctgatt cgttaccaa 5429

```

```

SEQ ID NO: 37      moltype = DNA length = 330
FEATURE           Location/Qualifiers
source            1..330
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 37
atgagcgata aaattattca cctgactgac gacagttttg acacggatgt actcaaagcg 60
gacggggcga tcctcgtcga tttctgggca gagtgggtgc gtccgtgcaa aatgatcgcc 120
ccgattctgg atgaatcgcg tgacgaatat cagggcaaac tgaccgttgc aaaactgaac 180
atcgatcaaa accctggcac tgccgcgaaa tatggcatcc gtggtatccc gactctgctg 240
ctggtcaaaa acggtgaagt ggcggcaacc aaagtgggtg cactgtctaa aggtcagttg 300
aaagagttcc tcgacgctaa cctggcgtaa 330

```

```

SEQ ID NO: 38      moltype = DNA length = 966
FEATURE           Location/Qualifiers
source            1..966
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 38
atgggcacga ccaaacacag taaactgctt atcctggggt caggcccgcc ggataacacc 60
gctgctgtct acgcggcgcg cgccaacctg caacctgtgc tgattaccgg catggaaaaa 120
ggcggccaac tgaccaccac caccgaagtg gaaaactggc ctggcgatcc aaacgatctg 180
accggtccgt tattaatgga gcgcatgcac gaacatgcca ccaagtttga aactgagatc 240
atttttgatc atatcaacaa ggtggatctg caaaaccgtc cgttccgtct gaatggcgat 300
aacggcgaat acacttgoga cgcgctgatt attgccaccg gagcttctgc acgctatctc 360
ggcctgcccct ctgaagaagc ctttaaaggc cgtggggttt ctgcttctgc aaactgcgac 420
ggtttcttct atcgcaacca gaaagtgtcg gtcacggcgg cggcaatac cgcgggtgaa 480
gaggcgctgt atctgtctaa catcgcttcg gaagtgcac tgattcaccg ccgtgacggt 540
ttccgcgcgg aaaaaatcct cattaaagcg ctgatggata aagtggagaa cggcaacatc 600
attctgcaca ccaaccgtac gctggaagaa gtgacggcgg atcaaatggg tgtcactggc 660
gttcgtctgc gcgatacgca aaacagcgat aacatcgagt cactcgacgt tgccggtctg 720
ttgtttgcta tcggttcacg ccgaataact gcgattttcg aagggcagct ggaactggaa 780
aacggctaca tcaaatgata gtcgggtatt catggtaatg ccaccagac cagcattcct 840
ggcgtctttg ccgcaggcga cgtgatggat cacatttacc gccaggccat tacttcggcc 900
ggtacaggct gcattggcagc acttgatgcy gaacgctacc tcgatggttt agctgacgca 960
aataaa 966

```

```

SEQ ID NO: 39      moltype = DNA length = 543
FEATURE           Location/Qualifiers
source            1..543
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 39
attaaaggcg gaaaacagat tcaaacggcg cgccctaacc gtatcaatgg cgaaattcgc 60
gccaggaag ttcgcttaac aggtctggaa ggcgagcagc ttggtattgg gagtctgaga 120
gaagctctcg agaaagcaga agaagccgga gtagacttag tcgagatcag ccctaaccgc 180
gagccgcggg tttgtcgat atgggattac ggcaaatccc tctatgaaa gagcaagtct 240
tctaagggaac agaagaaaaa gcaaaaaagt atccaggtta aggaatttaa attccgtcct 300
ggtacagatg aaggcgacta tcaggtaaaa ctccgcagcc tgattcgctt tctcgaagag 360
ggtgataaag ccaaaatcac gctgcgtttc cgcggtcgct agatggcgca ccagcaaatc 420
ggatgggaag tgcttaatcg cgtgaaagac gatttgcaag aactggcagt ggtcgaatcc 480
ttccaacga agatcgaagg ccgccagatg atcatgggtc tcgctcctaa gaagaacag 540
taa 543

```

```

SEQ ID NO: 40      moltype = DNA length = 1518
FEATURE           Location/Qualifiers
source            1..1518
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 40
atggcacaga ttgatttccg aaaaaaata aactggcatc gtcgttaccg ttcaccgcag 60
ggcgtaaaaa ccgaacatga gatectgcgg atcttcgaga gcgatcgcg gcgtatcatc 120
aactctccgg caattcgtcg tctgcaacaa aagacccagg tttttccact ggagcgcaat 180
gcccgcgtgc gcacgcgtct taccactcgt atggaagtcc agcagggtgg acgctacatc 240
gccaagaaaa ttttaagcgc tctgaaagag cttaaattgc tggaagcata cggcctggat 300
gaactgaccg gcccttttga aagcattgtt gagatgtcat gcctgatgca cgatatcgcc 360
aatccgcggt ttggtcatct tggcgaagca gcgataaatg actggtttgc ccaacgtttg 420
caccgcgaag atgcggaag ccagcctctg actgacgata gctgcagcgt ggccgcaata 480
cgtttacggg acggggaaga accgcttaac gagctgcggc gcaagattcg tcaggactta 540

```

-continued

tgctcatttg	aggggaatgc	acaaggcatt	cgtctggtgc	atacattgat	gcggatgaat	600
ctcacctggg	cacagggttg	cggatatttta	aaatataccc	gtccggcggtg	gtggcggtggc	660
gaaacgcctg	agacacatca	ctattttaatg	aaaaagccgg	gttattatct	ttctgaagaa	720
gcctatattg	cgcgggtgcg	taaagaactt	aatttggcgc	tttacagtcg	tttccatta	780
acgtgggatta	tggaaagctgc	cgaacacatc	tcctattgtg	tggcagacct	tgaagatgcg	840
gtagagaaaa	gaatatttac	cgttgagcag	ctttatcatc	atttgacga	agcgtggggc	900
cagcatgaga	aaggttcgct	cttttcgcta	gtggttgaaa	atgcctggga	aaaatcacgc	960
tcaaatagtt	taagccgcag	tacggaagat	cagtttttta	tgtatttacg	agtaaacacc	1020
ctaaataaac	tggtaacccta	cgcggcacaa	cgatttattg	ataatctgcc	tgcgattttc	1080
gccggaacgt	ttaatcatgc	attattggaa	gatgccagcg	aatgcagcga	tcttcttaag	1140
ctatataaaa	atgtcgctgt	aaaacatgtg	tttagccatc	cagatgtcga	gcagcttgaa	1200
ttgcagggct	atcgggtcat	tagcggatta	ttagagattt	atcgctcttt	attaagcctg	1260
tcgttatcag	actttactga	actggttagaa	aaagaacggg	tgaacggttt	ccctattgaa	1320
tcgcgcttat	tccacaaact	ctcgacgcgc	catcggtggg	cctatgtcga	ggctgtcagt	1380
aaattaccgt	cagattctcc	tgaagttccg	ctatggggaat	attattaccg	ttgccgcctg	1440
ctgcaggatt	atatcagcgg	tatgaccgac	ctctatgcgt	gggatgaata	ccgacgtctg	1500
atggccgtag	aacaataa					1518

SEQ ID NO: 41 moltype = DNA length = 150
 FEATURE Location/Qualifiers
 source 1..150
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 41
 attctgactg ataattgtct acagaaacgg agcggcacct cttttaaccc ttgaagtac 60
 tgcccgtttc gagagtttct caactcgaat aactaaagcc aacgtgaact ttgcggtac 120
 tccaggatcc gctttttttt gccataaaaa 150

SEQ ID NO: 42 moltype = DNA length = 87
 FEATURE Location/Qualifiers
 source 1..87
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 42
 gctgaaaaac ataaccata aaatgctagc tgtaccagga accacctctc tagcctgtgt 60
 aatctcccctt acacgggctt attttttt 87

SEQ ID NO: 43 moltype = DNA length = 723
 FEATURE Location/Qualifiers
 source 1..723
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 43
 atgagcaaa gtagaagaact gtttaccggc gttgtgccga ttctgggtga actggatggc 60
 gatgtgaacg gtcacaaaatt cagcgtgcgt ggtgaagggt aaggcgatgc cagcattggc 120
 aaactgacgc tgaattttat ctgcaccacc ggcaaaactgc cgggtccgtg gccgacgctg 180
 gtgaccaccc tgacctatgg cgttcagtgt tttagtctgt atccggatca catgaaacgt 240
 cagcatttct ttaaatctgc aatgccggaa ggctatgtgc aggaacgtac gattagcttt 300
 aaagatgatg gcaaatataa aacgcgcgcc gttgtgaaat ttgaaggcga taccctggtg 360
 aaccgcattg aactgaaagg caccgatttt aaagaagatg gcaataatct gggccataaa 420
 ctggaataca actttaatag ccataatgtt tatattacgg cggataaaca gaaaaatggc 480
 atcaaaagcg attttaccgt tcgccataac gttgaagatg gcagtgtgca gctggcagat 540
 cattatcagc agaatacccc gatttggtgat ggtccggtgc tgctgcccga taatcattat 600
 ctgagcacgc agaccgttct gtctaaagat ccgaacgaaa aaggcacgcg ggaccacatg 660
 gttctgcacg aatatgtgaa tgcggcaggt attacgtgga gccatccgca gttcgaaaaa 720
 taa 723

SEQ ID NO: 44 moltype = DNA length = 4461
 FEATURE Location/Qualifiers
 source 1..4461
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 44
 atgattgaac gcggtaaatt tcgctcactg acgctgatta actggaacgg cttttttgcc 60
 cgaaacttttg accttgacga gctggtcacg acgctttctg gcggtaacgg ggccggtaaa 120
 tccaccacca tggccggcgtt cgttacggcg ctgatccccg acctgacctt gttgcatttc 180
 cgtaacacta cgggaagccgg ggccaccagc ggttcgcgcg ataaaggtct gcacggtaag 240
 ctgaaagcgg gtgtctgtta ttcgatgctc gacaccatta actcgcgcga ccagcgcgtg 300
 gtggtcgggtg tgcgtctgca acaggttgcc ggacgcgacg gtaaaagtga tatcaaacca 360
 tttgccattc agggactgcc gatgtcgggt cagccgacac agctgggtgac cgaaccctgt 420
 aacgaacgcc aggcgcgcgt gctgccgctt aacgagctga aagacaagct cgaggcgatg 480
 gaaggcgtgc agtttaaaac gttcaactcc attacgatt accactcgt gatgttcgat 540
 ctgggcacat tcgcgcgtcg tctgcgctcc gcacttgacc gtagcaaat ctatcgtctg 600
 atcgaagctt cgtgtatggt cgggatctcc agtgccatta cccgttctct gcgcgactac 660
 ctggtgcggg aaaacagcgg cgtgcgtaaa gcgttccagg acatggaagc ggcgtgcgt 720
 gaaaaccgca tgacgctgga agcgattcgt gtcaccacag cggaccgtga cctgtttaag 780

-continued

```

catctgatca gcgaagctac caactacgtg gcggcgagact acatgcgtca cgccaacgag 840
cgccgtgtcc atctogacaa agccctggag ttctgtcgcg agttacatac ttccggtcag 900
caactggcgg ctgagcagta caaacacgtc gatattggcg gtgagctggc agagcacaac 960
ggtgcccgaag gcgatctgga agcggattat caggcgggcca gcgatcacct gaacctggtg 1020
caaacccgac tgcgtcagca ggagaagatc gaacgctacg aagcggatct cgatgagctg 1080
cagatccgtc tggaaagagca aaacgaagtg gtggcagaag ccacgaacg ccagggaagag 1140
aacgaggctc gcgcggaagc tgccgaactg gaagtggacg agctgaaaag ccagcttgct 1200
gactaccagc aggcgctgga cgtccagcaa acgcgcgcga tccagtataa ccaggcgatt 1260
gctgcactta atcgttgcaa agaactgtgc catctgccgg acttaaccgc cgacagcgcc 1320
gccgaatggc tggaaacctt ccaggcgaaa gagctggaag cgactgaaaa aatgctctct 1380
cttgagcaga aactggacat ggcgcaaac ccgacacgac agtttgagca ggcttatcag 1440
ctgggtggtg caatcaacgg cccactggcg cgtaacgagg cgtgggatgt cgctcgcgaa 1500
ctattgcgcg aaggggtcga tcagcgtcac ctggcagagc aggttcagcc gctgaggatg 1560
cgattaagcg aactggacaa gcgtctacgc gagcagcaag aagctgagcg tctgctggca 1620
gatttctgca aacgtcaggg caagaatttt gatatcgacg aactggaagc cctgcatcag 1680
gaactggaag cagcattgct ctctctttcc gatagcgtgt ctaacgcccg tgaagagcgc 1740
atggcactgc gccaggagca ggaacagctg cagtctcgca ttcagagttt gatgcagcgt 1800
gcgcgcgttt ggctggcagc gcaaaacagt ctcaaccagt tgagcgaaca gtgcggcgaa 1860
gaggtttacc ccagccagga cgtcaccgaa tatctgcaac agttgctgga gcgtgagcga 1920
gaggcgattg ttgaacgcga tgaagtgggc gcgcgcaaaa acgcgcgcga tgaagagatc 1980
gaacgtttaa gccagcctgg cggctctgaa gatcagcgtc tgaacgcgct ggccgagcgt 2040
tttggtggtg tgctgtgtc agaaatttat gacgacgtta gcctggaaga tgcgcgctac 2100
ttctcagcgc tgcattggcc gtcaacgccac gccatcgtgg tgcagatct gtcacaggta 2160
actgaacacc tggaaagcgt gaccgattgc ccggaagatc tctatctgat cgaaggagat 2220
cccgagtcac tcatgacagc cgtgttcagc gttgatgagc tggaaaaagc ggtagtgggtg 2280
aaaaatcgcc atcgtcagtg gcgttattca cgtttcccg aagtgcgctg gtttggctgt 2340
gctgcgcgtg aaagccgctat tgaagccctc catgcccagc gtgaagtgtc ttccgaacgc 2400
ttcggccagc tctcctttga ggtacagaaa actcagcgtc tgcacaggc gttcagccgc 2460
tttatcgcca gtcactctggc ggttgcgttt gagtctgacc cggaagcaga aatccgtcaa 2520
ctgaacagcc tgcgcgtcga actggagcgg gcgttaagta atcatgaaa tgataaccag 2580
cagcagcgtc ttacgtttga gcaggcgaaa gaggcgctga cggcgctgaa ccgcattctg 2640
ccgcgtctca acctgttggc tgatgacagc ctggcgagtc gcctcgatga aatccgcgaa 2700
cgtctggatg aagcccagga agccgcgcgt tttgttcagc agtttggcaa tcaactggcg 2760
aaactggaac cgattgtttc ggtattgcag agcgaccccg aacagttcga acagttaaaa 2820
gaagattacg cgtactctca gcagatgcag cgcgatgccc gtcagcagcg gtttgcctcg 2880
acggaagtgg tgcagcgtc tgcgcacttt agctattctg actcggcaga aatgcttagc 2940
ggtaaacagc atctcaacga aaaaactgcgt gaacgtctgg aacaggcgga agcggagcgt 3000
acccgcgcct gcgaagcgtt gcgcggtcac gcagcgcagt tgagtcaagta caaccagggtg 3060
ctggcttcgc tgaaaagtct ttacgacacc aaaaaagagc tactcaacga tctgcaacgt 3120
gaattgcagg atatcgccgt gcgtgctgat agcggggcag aagagcgggc gcgtattcgc 3180
cgtgacgagc tgcattgcga actgagcaat aaccgtttcac gccgcaatca actggaaaaa 3240
gcgcttacct tctgcgaagc ggagatggac aacctgaccc gcaaaactgc caagctggag 3300
gggattacct ttgagatgag cgaagcaggtg gtgacggcca aagcgggctg gttgctgggtg 3360
atgcgcattg tgaagatcaa cggcgcttgc cgcgccttac accgtcgtga gctggcttat 3420
ctctccgctg atgatttgcg ttccatgtcg gataaggcgt taggtgcgct gcgtctggcg 3480
gtggcgggata acgaacatct cgcgcagctg ctgcgcattg cgaagatcc gaaacgtccg 3540
gagcgtaaaa ttacgttctt cgtggcggtt tatcagcacc tgcgtgaacg tattcgtcag 3600
gatattatc tgaccgatga tccggtggaa gctatcgaac agatggagat tgaacttagc 3660
cgtctgaccc agaatattac ctcccgtgaa cagaaactgg cgtacagttc ccgcagcgtg 3720
gcgaacatca ttccgcaaac cattcagcgc gagcagaacc gtatccgtat gctcaaccag 3780
gggttgacga acgtatcgtt tggtcagggt aacagcgtgc gtctcaacgt gaacgtgcgt 3840
gaaacgcagc ccattgctact ggaatgtgct tctgaacagc acgagcagca tcaggatctg 3900
tttaacagca accgtttgac ctctcgggaa gcgctggcga aactgtatca acgtcttaac 3960
ccgcagattg atatggggcg gcgcacgcgc cagaccatcg gtgaagaact gctggattac 4020
cgcaactatc tggaaatgga agttgaggtt aaccgtgggt ccgatggctg gctgcgcgca 4080
gagtcgtggt cattgtcgac cgggtgaggc attggtaccg gtatgtcgat tctgggtgatg 4140
gtggtacaaa gctgggaaga tgaatctcgc cgcctgcgcg gtaagatat ctctccttgc 4200
cgctctgtgt tctcgtatga agcagcgca ctggatgctc gttctatgc cagcgtgttt 4260
gaattgtgtg agcgtttgca aatgcaactc atcatcgag cgccggaaaa tatcagcccg 4320
gagaaaggca ccacctataa actggtgcgt aaagtcttcc agaataccga acacgttcat 4380
gtcgtcggcc tgcgaggatt tgcgcgcgaa ctccctgaaa cgcttccagg aactgacgaa 4440
gcgccttctc aggcgagtta a

```

```

SEQ ID NO: 45      moltype = DNA length = 3186
FEATURE            Location/Qualifiers
source              1..3186
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 45
atgaaaagaa tghtaatcaa cgcaactcag cagggaagagt tgcgcgttgc ccttgtagat 60
gggcagcgtc tgtatgacct ggatatcgaa agtccagggc acgagcagaa aaaggcaaac 120
atctacaaag gtaaaaatcac ccgcattgaa ccgagtcgtg aagctgcttt tgttgattac 180
ggcgtgaac gtcacggttt cctcccacta aaagaaattg ccgcgcaata ttccctgtct 240
aactacagtg ctcatggtcg tcccaacatt aaagatgtgt tgcgtgaagg tcagggaagt 300
attgttcaga tcgataaaga agagcgcgcc cggcattaac cacctttatc 360
agtctggcgg gtatgctatc ggttctgat ccgaacaacc gcgcgcgggg tggcatttct 420

```

-continued

```

cgccgtatcg aaggcgacga cgtaccgaa ttaaaagaag cactggcaag ccttgaactg 480
ccggaaggca tggggcttat cgtgcgcacc gctggcgctg gcaaatctgc tgaggcgctg 540
caatgggatt taagcttccg tctgaaacac tgggaagcca tcaaaaaagc cgctgaaagc 600
cgcccgcccc cgctctctgat tcatcaggag agcaacgtaa tgcgtcgccg attcccgcat 660
tacttacgtc aggacatcgg cgaatacctt atcgataacc cgaaagtgtc cgaactggca 720
cgtcagcata tgcgtcgatt aggtcgcccc gatttcagca gcaaaatcaa actgtacacc 780
ggcgagatcc cgctgtttcag ccaactaccg atcgagtcac agatcgagtc cgccttcacg 840
cgtgaagttc gtctgcgcgtc tgggtgttcc attgttatcg acagcacoga agcgttaacg 900
gccatcgaca tcaactccgc acgcgcgacc cgccggcgcg atatogaaga aaccgcgttt 960
aacactaacc tcgaagctgc cgatgagatt gctcgtcagc tgcgcctgcg tgacctcggc 1020
ggcctgattg ttatcgactt catcgacatg acgcagtagc gccaccagcg tgcggtagaa 1080
aaccgtctgc gtgaagcggt gcgtcaggac cgtgcgcgta ttcaaatcag ccatatttct 1140
cgctttggcc tgctggaatg gtcctgtcag cgctgagcc catcactggg tgaatccagc 1200
catcacgtct gcccgcgctg ctccggtacc ggtacggtgc gtgacaacga atcgctgtcg 1260
ctctctatct tgcgtctgat cgaagaagaa gcgctgaaag agaaccacca ggaagtccac 1320
gccattgttc ctgtgcgaat cgcttcttat ctgctgaatg aaaaacgttc tgcggtgaat 1380
gccattgaaa cgtgtcagga cgtgtgttgc tgcgtgattg tgccaaacga tcagatggaa 1440
accccgctac accacgtgct gcgctgtcgt aaagggggaa aaacgcgcag cttaagctac 1500
atgtgcccga agctgcatag agaagcgatg gcgctgccc ctgaagaaga gttcgctgaa 1560
cgtgaagctc gggaaacaac tgcgtgggca acctttgcca tgcggtgagt gccgcccagc 1620
ccaaccccag ctgaacctgc cgcgcctgtc gtacccccag cacctaaatc tgcaccggca 1680
aacaccagcg tctctgcccc acctgggctg ttgagccgct tcttcggcgc actgaaagcg 1740
ctgttcagcg gtggtgaaga aaccaaacgc tccgagcaac caacaccgaa agcagaagcg 1800
aaaccggaac gtcaacagga tcgtcgcaag cctcgtcaga acaaccgcgc tgaccgtaat 1860
gagcgccgag acacccgtag tgaacgtact gaagcgagcg ataatcgcca agaaaaccgt 1920
cgtaatctgc ccagggcaca gcagcagact gccgagagcg gtgagagcgc tcagcaggtt 1980
gaggtaacgg aaaaagcgcg taccaccgac gagcagcaag cgccgcgctg tgaacgtagc 2040
cgccgccgta atgatgataa acgtcaggcg caacaagaag cgaaggcgct gaatgttga 2100
gagcaatctg ttcaggaaac cgaacaggaa gaacgtgtac gtcgggttca gccgcgtcgt 2160
aaacagcgct agctcaatca gaaagtgcgt tacgagcaaa gcgtagccga agaagcggt 2220
gtcgaccggg tgggtgaaga aactgtcgct gccgaaccga ttgttcagga agcgcagct 2280
ccacgcacag aactggtgaa agtcccgctg ccagtcgtag cgcaaaactg accagaacag 2340
caagaagaga acaatgttga taaccgtgac aacggtggca tgccgcgctg ttctcgccgc 2400
tcgctcgctc acctgcgcgt aagtgttcag cgtcgctcgc gctatcgta cgagcgctat 2460
ccaacccagt cgccaatgcc gttgaccgta gcgtgcgcgt ctccggaact ggccctctggc 2520
aaagtctgga tccgctatcc aattgtacgt ccgcaagatg tacaggttga agagcagcgc 2580
gaacaggaaag aagtacatgt gcagccgatg gtgactgagg tccctgtcgc cgccgctatc 2640
gaaccgggtg ttgagcgccc agttgttgaa gaagtggccg gtgtcgtaga agcccccggt 2700
caggttgccc aaccgcaacc ggaagtgggt gaaacgagcg atcctgaagt gattgtctgc 2760
gcggtaactg aacagccgca ggtgattacc gagtctgatg ttgcccagc ccagggaagt 2820
gcagaacaag cagaaccggt ggttgaaccg caggaagaga cggcagatat tgaagaagt 2880
gtcgaacctg ctgaggttgt agttgtgtaa cctgaagttg ttgctcaacc tgcgcgcgca 2940
gtagtgcgtc aagtcgcagc agaagttgaa acggtagctg cggtcgaacc tgaggtcacc 3000
gttagacata accacgctac cgcgccaatg acgcgcgctc cagcaccgga atatgttccg 3060
gaggcaccgc gtcacagtga ctggcagcgc cctacttttg ccttcgaagg taaaggtgcc 3120
gcaggtggtc ataccggaac acatcatgcc tctgctgctc ctgcccgtcc gcaacctgtt 3186
gagtaa

```

```

SEQ ID NO: 46      moltype = RNA length = 24
FEATURE           Location/Qualifiers
source            1..24
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 46
agatcgaaag cgtatccggt gaaa                                     24

```

```

SEQ ID NO: 47      moltype = RNA length = 24
FEATURE           Location/Qualifiers
source            1..24
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 47
atactttcac cggatacgct ttgc                                     24

```

```

SEQ ID NO: 48      moltype = RNA length = 24
FEATURE           Location/Qualifiers
source            1..24
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 48
agattatatg agcgataaaa ttat                                     24

```

```

SEQ ID NO: 49      moltype = RNA length = 24
FEATURE           Location/Qualifiers
source            1..24
                  mol_type = other RNA

```

-continued

```

                                organism = synthetic construct
SEQUENCE: 49
atacataatt ttatcgctca tata                                     24

SEQ ID NO: 50                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 50
agattggttc gcctggtatc cttt                                     24

SEQ ID NO: 51                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 51
atacaaagga taccaggcga acca                                     24

SEQ ID NO: 52                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 52
agataaggcc gcgtagcaac gtcg                                     24

SEQ ID NO: 53                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 53
ataccgacgt tgctacgagg cctt                                     24

SEQ ID NO: 54                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 54
agatctgcaa aaagcagcac aagg                                     24

SEQ ID NO: 55                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 55
atacccttgt gctgcttttt gcag                                     24

SEQ ID NO: 56                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 56
agatagatga ccgatctgga tctt                                     24

SEQ ID NO: 57                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 57
atacaagatc cagatcggtc atct                                     24

SEQ ID NO: 58                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 58
agattcttctc ttcgtctaaa gatt                                     24

```

-continued

SEQ ID NO: 59	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 59		
atacaatctt tagacgaaga aaga		24
SEQ ID NO: 60	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 60		
agatgtgacc aacgtagata aagc		24
SEQ ID NO: 61	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 61		
atacgcttta tctacgttgg tcac		24
SEQ ID NO: 62	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 62		
agattgccag actgtaagta cccg		24
SEQ ID NO: 63	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 63		
ataccgggta cttacagtct ggca		24
SEQ ID NO: 64	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 64		
agataagccg aagtacatct ggag		24
SEQ ID NO: 65	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 65		
atacctccag atgtacttcg gctt		24
SEQ ID NO: 66	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 66		
agattcggaa ctggatatga tggc		24
SEQ ID NO: 67	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 67		
atacaccatc atatccagtt ccga		24
SEQ ID NO: 68	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	

-continued

SEQUENCE: 68	organism = synthetic construct	
agatgctcac tgacgctgat taac		24
SEQ ID NO: 69	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 69		
atacgttaat cagcgtcagt gagc		24
SEQ ID NO: 70	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 70		
agataaatgc acaaaacttt tata		24
SEQ ID NO: 71	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 71		
atactataaa agttttgtgc attt		24
SEQ ID NO: 72	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 72		
agatcagcgc agcgcgcgag ctga		24
SEQ ID NO: 73	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 73		
atactcagct cgcgcgctgc gctg		24
SEQ ID NO: 74	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 74		
agatttaagg ttatcatccg tttc		24
SEQ ID NO: 75	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 75		
atacgaaaacg gatgataacc ttaa		24
SEQ ID NO: 76	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 76		
agatgcctgt gtaatctccc ttac		24
SEQ ID NO: 77	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 77		
atacgtaagg gagattacac aggc		24

-continued

SEQ ID NO: 78	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 78		
agatctgtat atactcacag cata		24
SEQ ID NO: 79	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 79		
atactatgct gtgagtatat acag		24
SEQ ID NO: 80	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 80		
agatatctca tccgtgatca catc		24
SEQ ID NO: 81	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 81		
atacgatgtg atcacggatg agat		24
SEQ ID NO: 82	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 82		
agatgcttta aaatccgcga gcac		24
SEQ ID NO: 83	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 83		
atacgtgctc gcggatttta aagc		24
SEQ ID NO: 84	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 84		
agattcgtat aatggattac ggca		24
SEQ ID NO: 85	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 85		
atactgccgt aatccattat acga		24
SEQ ID NO: 86	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 86		
agatcgctag tttcacccgg gggc		24
SEQ ID NO: 87	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	

-continued

```

                                organism = synthetic construct
SEQUENCE: 87
atacgccccc gggtgaaact agcg                                     24

SEQ ID NO: 88                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 88
agatttgctg aacgcgaacg gctg                                     24

SEQ ID NO: 89                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 89
ataccagccg ttcgcgttcg acaa                                     24

SEQ ID NO: 90                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 90
agataaggca tagtttacca tgcg                                     24

SEQ ID NO: 91                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 91
ataccgcatg gtaaaactatg cctt                                     24

SEQ ID NO: 92                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 92
agatcactgg agcgcaatgc cgcc                                     24

SEQ ID NO: 93                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 93
atacggcggc attgcgctcc agtg                                     24

SEQ ID NO: 94                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 94
agatgactgg tatcaggcgg atga                                     24

SEQ ID NO: 95                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 95
atactcatcc gcctgatacc agtc                                     24

SEQ ID NO: 96                moltype = RNA length = 24
FEATURE                      Location/Qualifiers
source                      1..24
                                mol_type = other RNA
                                organism = synthetic construct
SEQUENCE: 96
agattcgtga attacgtgag caag                                     24

```

-continued

SEQ ID NO: 97	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 97		
ataccttgct cacgtaattc acga		24
SEQ ID NO: 98	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 98		
agattgattt aatagatagt attc		24
SEQ ID NO: 99	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 99		
atacgaatac tatctattaa atca		24
SEQ ID NO: 100	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 100		
agatatcgct atgtcctggc cctc		24
SEQ ID NO: 101	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 101		
atacgagggc caggacatag cgat		24
SEQ ID NO: 102	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 102		
agatggctac cgtagagtcc tcgg		24
SEQ ID NO: 103	moltype = RNA length = 27	
FEATURE	Location/Qualifiers	
source	1..27	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 103		
atacttgggc taccgtagag tcctcgg		27
SEQ ID NO: 104	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 104		
agatgctctg gcctagagga caag		24
SEQ ID NO: 105	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 105		
ataccttgtc ctctaggcca gage		24
SEQ ID NO: 106	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	

-continued

```

                                organism = synthetic construct
SEQUENCE: 106
agattcccta tcaagcattc tgac                                     24

SEQ ID NO: 107            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 107
atacgtcaga atgcttgata ggga                                     24

SEQ ID NO: 108            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 108
agataccctt gaagtcactg cccg                                     24

SEQ ID NO: 109            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 109
ataccgggca gtgacttcaa gggt                                     24

SEQ ID NO: 110            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 110
agatcagtg aataatgagg ccgt                                     24

SEQ ID NO: 111            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 111
atacacggcc tcattattcc actg                                     24

SEQ ID NO: 112            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 112
agatttgatt acggcgctga acgt                                     24

SEQ ID NO: 113            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 113
atacacgttc agcgccgtaa tcaa                                     24

SEQ ID NO: 114            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 114
agatcgtctg taaattccct acaa                                     24

SEQ ID NO: 115            moltype = RNA length = 24
FEATURE                  Location/Qualifiers
source                  1..24
                        mol_type = other RNA
                        organism = synthetic construct
SEQUENCE: 115
atactttag ggaatttaca gacg                                     24

```

-continued

SEQ ID NO: 116	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 116		
agataaactg agatcatttt tgat		24
SEQ ID NO: 117	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 117		
atacatcaaa aatgatctca gttt		24
SEQ ID NO: 118	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 118		
agatgcacac tgggcaatac gagg		24
SEQ ID NO: 119	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	organism = synthetic construct	
SEQUENCE: 119		
ataccctcgt attgccagt gtgc		24
SEQ ID NO: 120	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
source	1..6	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 120		
HHHHHH		6

1. A genetically modified bacterial host cell comprising
 - a non-endogenous expression vector that comprises a translation initiation factor IF-3 (infC) nucleic acid sequence of SEQ ID NO: 39, an OxyS nucleic acid sequence of SEQ ID NO: 41, and/or a CyaR nucleic acid sequence of SEQ ID NO: 42; and/or
 - a vector that includes a nucleic acid sequence encoding a CRISPR enzyme and a vector that includes a nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC, optionally wherein the vector including the nucleic acid sequence encoding the CRISPR enzyme and the vector including the nucleic acid sequence encoding the crRNA are the same; and/or
 - a deletion or mutation in (a) a promoter region located upstream of a transcription start site (TSS) of recC, or (b) a recC coding region located downstream of a transcription start site (TSS) of recC, wherein the mutation or deletion represses transcription of recC, optionally wherein the mutation is a nonsense mutation, a frameshift mutation, or a missense mutation or the deletion is a 10-100 base pair deletion located upstream or downstream of the transcription start site (TSS) of recC.
2. (canceled)
3. (canceled)
4. (canceled)

5. (canceled)
6. (canceled)
7. (canceled)
8. The genetically modified bacterial host cell of claim 1, wherein the non-endogenous expression vectors comprising the infC nucleic acid sequence, the OxyS nucleic acid sequence and/or the CyaR nucleic acid sequence are the same or distinct.
9. (canceled)
10. The genetically modified bacterial host cell of claim 1, wherein the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is a plasmid, a cosmid, a bacmid, a bacterial artificial chromosome (BAC), or a viral vector.
11. The genetically modified bacterial host cell of claim 1, wherein the non-endogenous expression vector comprising the infC nucleic acid sequence, and/or the non-endogenous expression vector comprising the OxyS nucleic acid sequence, and/or the non-endogenous expression vector comprising the CyaR nucleic acid sequence is operably linked to an expression control sequence, optionally wherein the expression control sequence is an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter.
12. (canceled)
13. (canceled)

14. (canceled)
15. The genetically modified bacterial host cell of claim 1, wherein the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter; or wherein the CRISPR enzyme is a nuclease-deficient *Francisella novicida* Cas12a (dFnCas12a) and/or wherein the nucleic acid sequence encoding the crRNA is selected from the group consisting of SEQ ID NO: 86, SEQ ID NO: 87, SEQ ID NO: 88 and SEQ ID NO: 89.
16. (canceled)
17. (canceled)
18. (canceled)
19. (canceled)
20. (canceled)
21. The genetically modified bacterial host cell of claim 1, wherein the genetically modified bacterial host cell is an *E. coli* cell.
22. A cell lysate obtained from the genetically modified bacterial host cell of claim 1, wherein the cell lysate comprises an effective amount of transcription/translation (TXTL) machinery that is configured to synthesize bacteriophage under cell-free conditions, optionally wherein the cell lysate is prepared using one or more of French-press cell lysis, sonication, runoff reactions, or lysate dialysis.
23. (canceled)
24. An in vitro method for synthesizing bacteriophage virions comprising
contacting a bacteriophage genome with the cell lysate of claim 22, and an energy buffer in vitro to obtain a reaction mixture, and
incubating the reaction mixture under conditions to produce viable phage virions,
wherein the energy buffer comprises canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes.
25. The method of claim 24, wherein the energy buffer further comprises one or more of a polyethylene glycol (PEG) polymer, deoxynucleotide triphosphates (dNTPs), and stabilizers, optionally wherein the PEG polymer comprises PEG-8000 or PEG-6000; or
wherein the energy buffer further comprises one or more of Mg-glutamate, K-glutamate, tRNA, cAMP, folic acid, spermidine, 3-PGA, HEPES, Nicotinamide adenine dinucleotide (NAD), coenzyme A (CoA), DTT, and maltodextrin.
26. (canceled)
27. (canceled)
28. The method of claim 24, wherein the reaction mixture further comprises a vector including a nucleic acid sequence encoding a reporter gene, wherein the nucleic acid sequence encoding the reporter gene is operably linked to a promoter that is responsive to a RNA polymerase (RNAP) encoded by the bacteriophage genome, wherein the RNAP is selected from among SP6 RNA Polymerase, T3 RNA Polymerase and T7 RNA polymerase; or wherein the reporter gene encodes a bioluminescent protein, a fluorescent protein, or a chemiluminescent protein.
29. (canceled)
30. (canceled)
31. The method of claim 28, wherein the bioluminescent protein is Aequorin, firefly luciferase, Renilla luciferase, red luciferase, luxAB, or nanoluciferase; or

- wherein the chemiluminescent protein is β -galactosidase, horseradish peroxidase (HRP), or alkaline phosphatase; or
wherein the fluorescent protein is sfGFP, TagBFP, Azurite, EBFP2, mKalamal, Sirius, Sapphire, T-Sapphire, ECFP, Cerulean, SCFP3A, mTurquoise, monomeric Midoriishi-Cyan, TagCFP, mTFP1, EGFP, Emerald, Superfolder GFP, Monomeric Azami Green, TagGFP2, mUKG, mWasabi, EYFP, Citrine, Venus, SYFP2, TagYFP, Monomeric Kusabira-Orange, mKOx, mKO2, mOrange, mOrange2, mRaspberry, mCherry, dsRed, mStrawberry, mTangerine, tdTomato, TagRFP, TagRFP-T, mApple, mRuby, mPlum, HcRed-Tandem, mKate2, mNeptune, NirFP, TagRFP657, IFP1.4, iRFP, mKeima Red, LSS-mKate1, LSS-mKate2, PA-GFP, PAmCherry1, PAmCherry2, Kaede (green), Kaede (red), KikGR1 (green), KikGR1 (red), PS-CFP2, PS-CFP2, mEos2 (green), mEos2 (red), PSmOrange, or Dronpa.
32. (canceled)
33. (canceled)
34. The method of claim 24, wherein the bacteriophage genome is derived from a naturally occurring bacteriophage, or a genetically engineered bacteriophage and/or wherein the bacteriophage genome is derived from a bacteriophage selected from among T7 phage, T7-like phage, Lambda, K1E, T3, T5, T4, or PhiX174.
35. The method of claim 24, wherein the phage virions show increased efficiency of plating (EOP) relative to phage virions obtained with cell-free bacteriophage synthesis (CFBS) in a control lysate obtained from a wild-type bacterial cell, wherein the wild-type bacterial cell and the genetically modified bacterial host cell are the same species.
36. A kit comprising one or more expression vectors comprising at least one nucleic acid sequence selected from the group consisting of SEQ ID NO: 39, SEQ ID NO: 41 and SEQ ID NO: 42, and instructions for using the same to prepare donor bacterial host cell lysates for cell-free bacteriophage synthesis (CFBS).
37. The kit of claim 36, wherein the at least one nucleic acid sequence is operably linked to an inducible promoter, a constitutive promoter, an endogenous promoter, or a heterologous promoter.
38. A kit comprising one or more expression vectors comprising at least one nucleic acid sequence encoding a crRNA that specifically targets a promoter region located upstream of a transcription start site (TSS) of recC or a recC coding region located downstream of a transcription start site (TSS) of recC, and instructions for using the same to prepare donor bacterial host cell lysates for cell-free bacteriophage synthesis (CFBS), optionally wherein the one or more expression vectors further comprise a nucleic acid sequence encoding a CRISPR enzyme.
39. The kit of claim 38, wherein the nucleic acid sequence encoding the CRISPR enzyme is operably linked to an inducible promoter and/or the at least one nucleic acid sequence encoding the crRNA is operably linked to a constitutive promoter.
40. A kit comprising the genetically modified bacterial host cell of claim 1, and instructions for using the same to prepare donor bacterial host cell lysates for cell-free bacteriophage synthesis (CFBS).
41. The kit of claim 36 further comprising one or more of polyethylene glycol (PEG) polymers, deoxynucleotide triphosphates (dNTPs), stabilizers, and an energy buffer com-

prising canonical amino acids, phosphoenol pyruvate (PEP), nucleoside triphosphates (NTPs), cofactors, and coenzymes.

42. The kit of claim **41**, further comprising a vector including a nucleic acid sequence encoding a reporter gene, wherein the nucleic acid sequence encoding the reporter gene is operably linked to a promoter that is responsive to a phage RNA polymerase (RNAP).

* * * * *