



US 20250283113A1

(19) **United States**(12) **Patent Application Publication**
KIM et al.(10) **Pub. No.: US 2025/0283113 A1**(43) **Pub. Date: Sep. 11, 2025**(54) **COMPOSITIONS AND METHODS FOR
INCREASING DELETION EFFICIENCY OF
NUCLEIC ACID SEGMENT BY
MODULATION OF NHEJ REPAIR PATHWAY**(71) Applicant: **GenKore, Inc.**, Daejeon (KR)(72) Inventors: **Yong Sam KIM**, Seoul (KR); **Do Yon
KIM**, Daejeon (KR); **Hyun Jung
CHIN**, Seoul (KR)(21) Appl. No.: **18/845,663**(22) PCT Filed: **Mar. 10, 2023**(86) PCT No.: **PCT/KR2023/003336**

§ 371 (c)(1),

(2) Date: **May 12, 2025**(30) **Foreign Application Priority Data**

Mar. 10, 2022 (KR) 10-2022-0030000


May 27, 2022 (KR) 10-2022-0065600

Publication Classification(51) **Int. Cl.***C12N 15/90* (2006.01)*C12N 9/22* (2006.01)*C12N 15/11* (2006.01)(52) **U.S. Cl.**CPC *C12N 15/90* (2013.01); *C12N 9/226*(2025.05); *C12N 15/11* (2013.01); *C12N**2310/20* (2017.05); *C12N 2310/531* (2013.01)


(57)

ABSTRACT


The present invention relates to a composition and method for increasing deletion efficiency of nucleic acid segments in a target gene by modulation of non-homologous end joining (NHEJ) repair pathway. Specifically, the present invention relates to a CRISPR/Cas system (for example, CRISPR/Cas12f1 system) for increasing deletion efficiency of nucleic acid segments in a target gene by modulation of non-homologous end joining repair pathway.

Specification includes a Sequence Listing.


[FIG. 1]



[FIG. 2]



[FIG. 3]



[FIG. 4A]

Name	Amino acid sequence	SEQ ID NO
TnpB protein	MGEKSSRRRRNGKSGAWTAAITSCVGGKMAKNTITKTLKLRIVRPYN SAEVEKIVADEKNNREKIALEKNKDKVKEACSKHLKVAAYCTTQVER NACLFCKARKLDDKFYQKLRGQFPDAVFWQEI SEIFRQLQKQAAEIY NQSLIELYYEIFIKGKGIANASSVEHYLSDVCYTRAAELFKNAAIAS GLRSKIKSNFRLKELKNMKSGLPTTKSDNFP IPLVKQKGGQYTGF EI SNHNSDFI IKIPFGRWQVKKEIDKYRPWEKFD FEQVQKSPKPI SLLL STQRRKRNGWSKDEGTEAEIKKVMNGDYQTSYIEVKRGSKIGEKSA WMLNLSIDVPKIDKGVDPSIIGGIDVGKSPFLVCAINNAFSRYSISD NDLFHFNKKMFARRRILLKKNRHKRAGHGAKNKLKPIITILTEKSERF RKKLIERWACEIADFFIKNKVGTVMENLESMKRKEDSYFNIIRLRGF WPYAEMONKIEFKLKQYGI EIRKVAPNNTSKTCSKCGHLNNYFNFEY RKKNKFPHFKCEKCNFKENADYNAALNISNPCLKSTKEEP	1
TnpB-v1 protein	MEKRINKIRKKLSADNATKPVSRSGPMAKNTITKTLKLRIVRPYN SAEVEKIVADEKNNREKIALEKNKDKVKEACSKHLKVAAYCTTQVER NACLFCKARKLDDKFYQKLRGQFPDAVFWQEI SEIFRQLQKQAAEIY NQSLIELYYEIFIKGKGIANASSVEHYLSDVCYTRAAELFKNAAIAS GLRSKIKSNFRLKELKNMKSGLPTTKSDNFP IPLVKQKGGQYTGF EI SNHNSDFI IKIPFGRWQVKKEIDKYRPWEKFD FEQVQKSPKPI SLLL STQRRKRNGWSKDEGTEAEIKKVMNGDYQTSYIEVKRGSKIGEKSA WMLNLSIDVPKIDKGVDPSIIGGIDVGKSPFLVCAINNAFSRYSISD NDLFHFNKKMFARRRILLKKNRHKRAGHGAKNKLKPIITILTEKSERF RKKLIERWACEIADFFIKNKVGTVMENLESMKRKEDSYFNIIRLRGF WPYAEMONKIEFKLKQYGI EIRKVAPNNTSKTCSKCGHLNNYFNFEY RKKNKFPHFKCEKCNFKENADYNAALNISNPCLKSTKEEP	2

[FIG. 4B]

Name	Amino acid sequence	SEQ ID NO
TnpB-v2 protein	MAGGPGAGSAAPVSSTSSLPLAALNMRVMAKNTITKTLKLRIVRPYNSAE VEKIVADEKNNREKIALEKNKDKVKEACSKHLKVAAYCTTQVERNACLFC KARKLDDKFYQKLRGQFPDAVFWQEI SEIFRQLQKQAAEIYNQSLIELYY EIFIKGKGIANASSVEHYLSDVCYTRAAELFKNAAIASGLRSKIKSNFRL KELKNMKSGLPPTTKSDNFP IPLVKQKGGQYTGFESI SNHNSDFI IKIPFGR WQVKKEIDKYRPWEKFD FEQVQKSPKPI SLLLSTQRRKRKNGWSKDEGTE AEIKKVMNGDYQTSYIEVKGSKIGEKS AWMLNLSIDVPKIDKGVDPSSI GGIDVGVSPLVCAINNAFSRYSISDNDLFHFNKKMFARRRILLKKNRHK RAGHGAKNKLKPIITILTEKSERFRKKLIERWACEIADFFIKNKVGTVMEN NLESMKRKEDSYFNIRLRGFWPYAEMQKIEFKLKQYGI EIRKVAPNNTS KTCCKGHLNNYFNFEYRKKNKFP HFKCEKCNFKENADYNAALNISNPKL KSTKEEP	3
TnpB-v3 protein	MAGGPGAGSAAPVSSTSSLPLAALNMMAKNTITKTLKLRIVRPYNSAEVE KIVADEKNNREKIALEKNKDKVKEACSKHLKVAAYCTTQVERNACLFCCKA RKLDDKFYQKLRGQFPDAVFWQEI SEIFRQLQKQAAEIYNQSLIELYYEI FIKGKGIANASSVEHYLSDVCYTRAAELFKNAAIASGLRSKIKSNFRLKE LKNMKSGLPPTTKSDNFP IPLVKQKGGQYTGFESI SNHNSDFI IKIPFGRWQ VKKEIDKYRPWEKFD FEQVQKSPKPI SLLLSTQRRKRKNGWSKDEGTEAE IKKVMNGDYQTSYIEVKGSKIGEKS AWMLNLSIDVPKIDKGVDPSSIIGG IDVGVSPLVCAINNAFSRYSISDNDLFHFNKKMFARRRILLKKNRHKRA GHGAKNKLKPIITILTEKSERFRKKLIERWACEIADFFIKNKVGTVMENL ESMKRKEDSYFNIRLRGFWPYAEMQKIEFKLKQYGI EIRKVAPNNTSKT CCKGHLNNYFNFEYRKKNKFP HFKCEKCNFKENADYNAALNISNPKLKS TKEEP	4

[FIG. 4C]

Name	Amino acid sequence	SEQ ID NO
Cas12f1 protein	MAKNTITKTLKLRIVRPYNSAEVEKIVADEKNNREKIALEKNKDKVKEAC SKHLKVAAYCTTQVERNACLFCKARKLDDKFYQKLRGQFPDAVFWQEISE IFRQLQKQAAEIYNQSLIELYYEIFIKGKGIANASSVEHYLSDVCYTRAA ELFKNAATASGLRSKIKSNFRLKELKNMKSGLPPTKSDNFP IPLVKQKGG QYTGFEISNHNSDFIIKIPFGRWQVKKEIDKYRPWEKFD FEQVQKSPKPI SLLLSTQRRKKNKGWSKDEGTEAEIKKVMNGDYQTSYIEVKRGSKI GEKS AWMLNLSIDVPKIDKGVDPSSIIGGIDVGKSPLVCAINNAFSRYSISDND LFHFNKKMFARRRI LLKKNRHKRAGHGAKNKLKPIITILTEKSERFRKKLI ERWACEIADFFIKNKVGTVMENLESMKRKEDSYFNIRLRGFWPYAEMQN KIEFKLKQYGI EIRKVAPNNTSKTCSKCGHLNNYFNFEYRKKNKFPHFKC EKC�FKENADYNAALNISNPCLKSTKEEP	5

[FIG. 4D]

Name	Nucleotide sequence (5' to 3')	SEQ ID NO
Human codon-optimized nucleic acid encoding TnpB protein	ATGGGGGAGAAAAGTTCCCGCCGCCGACGGAATGGAAAAAGCGGTGCGTGGACTGCTGCTATA ACAAGCTGTGTTGGGGGTAAAGATGGCCAAGAACACAATTACAAAGACACTGAAGCTGAGGATC GTGAGACCATACAACAGCGCTGAGGTGAGAAAGATTGTGGCTGATGAAAAGAACACAGGGAA AAGATCGCCCTCGAGAAGAACAAGGATAAGGTGAAGGAGGCCTGCTCTAAGCACCTGAAAGTG GCCGCCTACTGCACCACACAGGTGGAGAGGAACGCCCTGTCTGTTTTGTAAAGCTCGGAAGCTG GATGATAAGTTTTACCAGAAGCTGCGGGGCCAGTTCCCCGATGCCGTCTTTTGGCAGGAGATT AGCGAGATCTTCAGACAGCTGCAGAAGCAGGCCGCCGAGATCTACAACCAGAGCCTGATCGAG CTCTACTACGAGATCTTCATCAAGGGCAAGGGCATTGCCAACGCCCTCTCCGTGGAGCACTAC CTGAGCGACGTGTGCTACACAAGAGCCGCCGAGCTCTTTAAGAACGCCGCTATCGCTTCCGGG CTGAGGAGCAAGATTAAAGAGTAACCTCCGGCTCAAGGAGCTGAAGAACATGAAGAGCGGCCCTG CCCACTACAAAGAGCGACAACCTCCCAATTCCACTGGTGAAGCAGAAGGGGGGCCAGTACACA GGGTTGAGATTTCCAACCACAACAGCGACTTTATTATTAAGATCCCTTTGGCAGGTGGCAG GTCAAGAAGGAGATTGACAAGTACAGGCCCTGGGAGAAGTTTGATTTCGAGCAGGTGCAGAAG AGCCCCAAGCCTATTTCCCTGCTGCTGTCCACACAGCGCGGAAGAGGAACAAGGGGTGGTCT AAGGATGAGGGGACCGAGGCCGAGATTAAGAAAGTGATGAACGGCGACTACCAGACAAGCTAC ATCGAGGTCAAGCGGGGAGTAAGATTGGCGAGAAGAGCGCCTGGATGCTGAACCTGAGCATT GACGTGCCAAAGATTGATAAGGGCGTGGATCCAGCATCATCGGAGGGATCGATGTGGGGGTC AAGAGCCCCCTCGTGTGCGCCATCAACAAGCCCTTCAGCAGGTACAGCATCTCCGATAACGAC CTGTTCCACTTTAACAAGAAGATGTTGCCCCGGGGAGGATTTTGCTCAAGAAGAACCGGCAC AAGCGGGCCGGACAGGGGCCAAGAACAAGCTCAAGCCCATCACTATCCTGACCGAGAAGAGC GAGAGGTTCAAGGAAGAAGCTCATCGAGAGATGGGCCCTGCGAGATCGCCGATTTCTTTATTAAG AACAAGGTCCGAACAGTGCAGATGGAGAACCTCGAGAGCATGAAGAGGAAGGAGGATTCTTAC TTCAACATTCCGGCTGAGGGGGTTCTGCCCCACGCTGAGATGCAGAACAGATTGAGTTTAAG CTGAAGCAGTACGGGATTGAGATCCGGAAGGTGGCCCCCAACAACACCAGCAAGACCTGCAGC AAGTGCGGGCACCTCAACAACACTACTCAACTTCGAGTACCGGAAGAAGAACAAGTTCCACAC TTCAAGTGCAGAGAAGTGCAACTTTAAGGAGAAGCGCGATTACAACGCCGCCCTGAACATCAGC AACCTTAAGCTGAAGAGCACTAAGGAGGAGCCC	6

[FIG. 4E]

Name	Nucleotide sequence (5' to 3')	SEQ ID NO
Human codon-optimized nucleic acid encoding TnpB-v1 protein	ATGGAAAAGAGAATCAACAAGATCAGGAAGAAGCTGAGCGCCGACAACGCCACCAAGCCTGTG TCTAGGAGTGGCCCCCATGGCCAAGAACAATTACAAAGACACTGAAGCTGAGGATCGTGAG ACCATACAACAGCGCTGAGGTCGAGAAGATTGTGGCTGATGAAAAGAACAAACAGGGAAAAGAT CGCCCTCGAGAAGAACAAGGATAAGGTGAAGGAGGCCTGCTCTAAGCACCTGAAAGTGGCCGC CTACTGCCACCACACAGGTGGAGAGGAACGCCTGTCTGTTTGTAAAGCTCGGAAGCTGGATGA TAAGTTTACCAGAAGCTGCGGGGCCAGTTCCCCGATGCCGTCTTTTGGCAGGAGATTAGCGA GATCTTCAGACAGCTGCAGAAGCAGGCCGCCGAGATCTACAACCAGAGCCTGATCGAGCTCTA CTACGAGATCTTCATCAAGGGCAAGGGCATTGCCAAGCCTCCTCCGTGGAGCACTACCTGAG CGACGTGTGCTACACAAGAGCCGCCGAGCTCTTTAAGAACGCCGCTATCGCTTCGGGGCTGAG GAGCAAGATTAAGAGTAACTTCGGCTCAAGGAGCTGAAGAACATGAAGAGCGGCCTGCCAC TACAAAGAGCGACAACCTTCCCAATTCCACTGGTGAAGCAGAAGGGGGGCCAGTACACAGGGTT CGAGATTTCACACCACAACAGCGACTTTATTATTAAGATCCCTTTGGCAGGTGGCAGGTCAA GAGGAGATTGACAAGTACAGGCCCTGGGAGAAGTTTGATTTCGAGCAGGTGCAGAAGAGCCC CAAGCCTATTTCCCTGCTGCTGTCCACACAGCGCCGAAGAGGAACAAGGGGTGGTCTAAGGA TGAGGGGACCGAGGCCGAGATTAAAGAAAGTGATGAACGGCGACTACCAGACAAGCTACATCGA GGTCAAGCGGGCAGTAAGATTGGCGAGAAGAGCGCCTGGATGCTGAACCTGAGCATTGACGT GCCAAAGATTGATAAGGGCGTGGATCCAGCATCATCGGAGGGATCGATGTGGGGTCAAGAG CCCCCTCGTGTGCGCCATCAACAACGCCTTCAGCAGGTACAGCATCTCCGATAACGACCTGTT CCACTTTAACAAGAAGATGTTGCCCCGCCGAGGATTTTGCTCAAGAAGAACCGCCACAAGCG GGCCGGACACGGGGCCAAGAACAAGCTCAAGCCCATCACTATCTGACCGAGAAGAGCGAGAG GTTCAGGAAGAAGCTCATCGAGAGATGGGCCTGCGAGATCGCCGATTTCTTTATTAAGAACA GGTCGGAACAGTGCAGATGGAGAACCCTCGAGAGCATGAAGAGGAAGGAGGATTCTACTTCAA CATTGGCTGAGGGGGTTCTGGCCCTACGCTGAGATGCAGAACAAAGATTGAGTTTAAGCTGAA GCAGTACGGGATTGAGATCCGGAAGGTGGCCCCCAACAACACCAGCAAGACCTCCAGCAAGTG CGGGCACCTCAACAACCTACTTCAACTTCGAGTACCGGAAGAAGAACAAGTTCCACACTTCAA GTGCGAGAAGTGCAACTTTAAGGAGAACCGGATTACAACGCCGCCCTGAACATCAGCAACCC TAAGCTGAAGAGCACTAAGGAGGAGCCC	7

[FIG. 4F]

Name	Nucleotide sequence (5' to 3')	SEQ ID NO
Human codon-optimized nucleic acid encoding TnpB-v2 protein	ATGGCTGGCGGACCAGGCGCAGGTAGTGCTGCGCCAGTTTCTTCAACTTCCTCCCTGCCCTG GCTGCGCTTAACATGCGCGTGATGGCCAAGAACAACAATTACAAAGACACTGAAGCTGAGGATC GTGAGACCATACAACAGCGCTGAGGTGAGAGATTGTGGCTGATGAAAAGAACAACAGGGAA AAGATCGCCCTCGAGAAGAACAAGGATAAGGTGAAGGAGGCCCTGCTCTAAGCACCTGAAAGTG GCCGCCTACTGCACCACACAGGTGGAGAGGAACGCCGTGTCTGTTTTGTAAAGTCTGGAAGCTG GATGATAAGTTTTACCAGAAGCTGCGGGGCCAGTTCCCCGATGCCGTCTTTTGGCAGGAGATT AGCGAGATCTTCAGACAGCTGCAGAAGCAGGCCGCCGAGATCTACAACCAGAGCCTGATCGAG CTCTACTACGAGATCTTCATCAAGGGCAAGGGCATTGCCAACGCCCTCTCCGTGGAGCACTAC CTGAGCGACGTGTGCTACACAAGAGCGCGCGAGCTCTTTAAGAACGCCGCTATCGCTTCCGGG CTGAGGAGCAAGATTAAGAGTAACCTCCGGCTCAAGGAGCTGAAGAACATGAAGAGCGGCCCTG CCCACTACAAGAGCGACAACCTTCCCAATTCCACTGGTGAAGCAGAAGGGGGGCCAGTACACA GGGTTCGAGATTTCCAACCACAACAGCGACTTTATTATTAAGATCCCCTTTGGCAGGTGGCAG GTCAAGAAGGAGATTGACAAGTACAGGCCCTGGGAGAAGTTTGATTTCGAGCAGGTGCAGAAG AGCCCCAAGCCTATTTCCCTGCTGCTGTCCACACAGCGCGGAAGAGGAACAAGGGGTGGTCT AAGGATGAGGGGACCGAGGCCGAGATTAAGAAAGTGATGAACGCGGACTACCAGACAAGCTAC ATCGAGGTCAAGCGGGGCAGTAAGATTGGCGAGAAGAGCGCCTGGATGCTGAACCTGAGCATT GACGTGCCAAAGATTGATAAGGGCGTGGATCCAGCATCATCGGAGGGATCGATGTGGGGGTC AAGAGCCCCCTCGTGTGCGCCATCAACAACGCCCTTCAGCAGGTACAGCATCTCCGATAACGAC CTGTTCCACTTTAACAAGAAGATGTTGCGCCGGCGGAGGATTTTGCTCAAGAAGAACCGGCAC AAGCGGGCCGGACACGGGGCCAAGAACAAGCTCAAGCCCATCACTATCCTGACCGAGAAGAGC GAGAGGTTCAAGGAAGAAGCTCATCGAGAGATGGGCCCTGCGAGATCGCCGATTTCTTTATTAAG AACAAGGTCCGAACAGTCCAGATGGAGAACCTCGAGAGCATGAAGAGGAAGGAGGATTCTTAC TTCAACATTGCGCTGAGGGGGTTCTGGCCCTACGCTGAGATGCAGAACAAGATTGAGTTTAAG CTGAAGCAGTACGGGATTGAGATCCGGAAGGTGGCCCCCAACAACACCAGCAAGACCTGCAGC AAGTGCGGGCACCTCAACAACACTTCAACTTCGAGTACCGGAAGAAGAACAAGTTCCACAC TTCAAGTGGGAGAAGTGCAACTTTAAGGAGAACGCCGATTACAACGCCGCCCTGAACATCAGC AACCTTAAGCTGAAGAGCACTAAGGAGGAGCCC	8


[FIG. 4G]

Name	Nucleotide sequence (5' to 3')	SEQ ID NO
Human codon-optimized nucleic acid encoding TnpB-v3 protein	ATGGCTGGCGGACCAGGCGCAGGTAGTGCTGCGCCAGTTTCTTCAACTTCCTCCCTGCCCTTG GCTGCGCTTAACATGATGGCCAAGAACAATTACAAGACACTGAAGCTGAGGATCGTGAGA CCATACAACAGCGCTGAGGTCGAGAAGATTGTGGCTGATGAAAAGAACAACAGGGAAAAGATC GCCCTCGAGAAGAACAAGGATAAGGTGAAGGAGGCCCTGCTCTAAGCACCTGAAAGTGGCCGCC TACTGCACCACACAGGTGGAGAGGAACGCCCTGTCGTGTTTGTAAAGCTCGGAAGCTGGATGAT AAGTTTACCAGAAGCTGCGGGGCCAGTTCCCGATGCCGTCTTTTGGCAGGAGATTAGCGAG ATCTTCAGACAGCTGCAGAAGCAGGCCGCCGAGATCTACAACCAGAGCCTGATCGAGCTCTAC TACGAGATCTTCATCAAGGGCAAGGGCATTGCCAACGCCCTCCTCCGTGGAGCACTACCTGAGC GACGTGTGCTACACAAGAGCCGCCGAGCTCTTTAAGAACGCCGCTATCGCTTCGGGGCTGAGG AGCAAGATTAAGAGTAACCTCCGGCTCAAGGAGCTGAAGAACATGAAGAGCGGCCCTGCCACT ACAAGAGCGACAACCTTCCCAATTCCACTGGTGAAGCAGAAGGGGGGCCAGTACACAGGGTTC GAGATTTCCAACCACAACAGCGACTTTATTATTAAGATCCCCCTTTGGCAGGTGGCAGGTCAAG AAGGAGATTGACAAGTACAGGCCCTGGGAGAAGTTTGATTTGAGCAGGTGCAGAAGAGCCCC AAGCCTATTTCCCTGCTGCTGTCCACACAGCGCGGAAGAGGAACAAGGGGTGGTCTAAGGAT GAGGGGACCGAGGCCGAGATTAAGAAAGTGATGAACGGCGACTACCAGACAAGCTACATCGAG GTCAAGCGGGGCAGTAAGATTGGCGAGAAGAGCGCCTGGATGCTGAACCTGAGCATTGACGTG CCAAAGATTGATAAGGGCGTGATCCACGATCATCGGAGGGATCGATGTGGGGGTCAAGAGC CCCCCTGTGTGCCCATCAACAACGCCCTCAGCAGGTACAGCATCTCCGATAACGACCTGTTT CACTTTAACAAGAAGATGTTGCCCCGGCGGAGGATTTTGCTCAAGAAGAACCGGCACAAGCGG GCCGGACACGGGGCCAAGAACAAGCTCAAGCCCATCACTATCCTGACCGAGAAGAGCGAGAGG TTCAGGAAGAAGCTCATCGAGAGATGGGCCTGCGAGATCGCCGATTCTTTATTAAGAACAAG GTGGGAACAGTGCAGATGGAGAACCCTCGAGAGCATGAAGAGGAAGGAGGATTCTACTTCAAC ATTCGGCTGAGGGGGTTCTGGCCCTACGCTGAGATGCAGAACAAGATTGAGTTTAAGCTGAAG CAGTACGGGATTGAGATCCGAAGGTGGCCCCCAACAACACCAGCAAGACCTGCAGCAAGTGC GGGCACCTCAACAACCTACTTCAACTTCGAGTACCGGAAGAAGAACAAGTTCCACACTTCAAG TGCGAGAAGTGCAACTTTAAGGAGAACGCCGATTAC AACGCCGCCCTGAACATCAGCAACCCCT AAGCTGAAGAGCACTAAGGAGGAGCCC	9


[FIG. 4H]

Name	Nucleotide sequence (5' to 3')	SEQ ID NO
Human codon-optimized nucleic acid encoding Cas12f1 protein	ATGGCCAAGAACACAATTACAAAGACACTGAAGCTGAGGATCGTGAGACCATACAACAGCGCT GAGGTCGAGAAGATTGTGGCTGATGAAAAGAACAACAGGGAAAAGATCGCCCTCGAGAAGAAC AAGGATAAGGTGAAGGAGGCCTGCTCTAAGCACCTGAAAGTGGCCGCCCTACTGCACCACACAG GTGGAGAGGAACGCCTGTCTGTTTTGTAAAGCTCGGAAGCTGGATGATAAGTTTTACCAGAAG CTGCGGGGCCAGTTCCCGGATGCCGTCTTTTGGCAGGAGATTAGCGAGATCTTCAGACAGCTG CAGAAGCAGGCCGCCGAGATCTACAACCAGAGCCTGATCGAGCTCTACTACGAGATCTTCATC AAGGGCAAGGGCATTGCCAAGCCCTCTCCGTGGAGCACTACCTGAGCGACGTGTGCTACACA AGAGCCGCCGAGCTCTTTAAGAAGCCGCTATCGCTTCCGGGCTGAGGAGCAAGATTAAGAGT AACTTCCGGCTCAAGGAGCTGAAGAACATGAAGAGCGGCCCTGCCCACTACAAAGAGCGACAAC TTCCCAATTCCACTGGTGAAGCAGAAGGGGGGCCAGTACACAGGGTTCGAGATTTCCAACCAC AACAGCGACTTTATTATTAAAGATCCCCCTTTGGCAGGTGGCAGGTCAAGAAGGAGATTGACAAG TACAGGCCCTGGGAGAAGTTTGATTTTCGAGCAGGTGCAAGAAGAGCCCCAAGCCTATTTCCCTG CTGCTGTCCACACAGCGGCGGAAGAGGAACAAGGGGTGGTCTAAGGATGAGGGGACCGAGGCC GAGATTAAGAAAGTGATGAAGGGCGACTACCAGACAAGCTACATCGAGGTCAAGCGGGGCGAGT AAGATTGGCGAGAAGAGCGCCTGGATGCTGAACCTGAGCATTGACGTGCCAAAGATTGATAAG GGCGTGGATCCAGCATCATCGGAGGGATCGATGTGGGGTCAAGAGCCCCCTCGTGTGCCCC ATCAACAACGCCCTTCAGCAGGTACAGCATCTCCGATAACGACCTGTTCACCTTTAACAAGAAG ATGTTCCGCCCGCGGAGGATTTTGCTCAAGAAGAACCGGCACAAGCGGGCCGACACGGGGCC AAGAACAAGCTCAAGCCCATCACTATCCTGACCGAGAAGAGCGAGAGGTTCAAGGAAGAAGCTC ATCGAGAGATGGGCCTGCGAGATCGCCGATTTCCTTTATTAAGAACAAAGGTCCGAACAGTGCAG ATGGAGAACCTCGAGAGCATGAAGAGGAAGGAGGATTCTTCACTTCAACATTCCGCTGAGGGGG TTCTGGCCCTACGCTGAGATGCAGAACAAAGATTGAGTTTAAGCTGAAGCAGTACGGGATTGAG ATCCGGAAGGTGGCCCCCAACAACACCAGCAAGACCTGCAGCAAGTGGGGCACCTCAACAAC TACTTCAACTTCGAGTACCGGAAGAAGAACAAGTTCCACACTTCAAGTGGGAGAAGTGAAC TTTAAGGAGAACGCCGATTACAACGCCGCCCTGAACATCAGCAACCCCTAAGCTGAAGAGCACT AAGGAGGAGCCC	10


[FIG. 5]




[FIG. 6A]




[FIG. 6B]




[FIG. 6C]




[FIG. 7A]




[FIG. 7B]




[FIG. 7C]




[FIG. 8]




[FIG. 9]




[FIG. 10A]




[FIG. 10B]




[FIG. 11]




[FIG. 12]





[FIG. 13]





[FIG. 14]




[FIG. 15]




[FIG. 16]




[FIG. 17]




[FIG. 18]




[FIG. 19]




[FIG. 20]




[FIG. 21]



[FIG. 22]



[FIG. 23]



**COMPOSITIONS AND METHODS FOR
INCREASING DELETION EFFICIENCY OF
NUCLEIC ACID SEGMENT BY
MODULATION OF NHEJ REPAIR PATHWAY**

TECHNICAL FIELD

[0001] This application claims priorities based on Korean Patent Application No. 2022-0030000, filed on Mar. 10, 2022, and Korean Patent Application No. 2022-0065600, filed on May 27, 2022, the entire disclosures of which are incorporated herein by reference.

[0002] The present disclosure relates to a composition and a method for increasing deletion efficiency for a nucleic acid segment in a target gene through regulation of the non-homologous end joining (NHEJ) repair pathway. Specifically, the present disclosure relates to a CRISPR/Cas system (for example, a CRISPR/Cas12f1 system) for increasing deletion efficiency for a nucleic acid segment in a target gene by regulation of the non-homologous end joining repair pathway.

BACKGROUND ART

[0003] Gene scissors technology, represented by a CRISPR/Cas system, is a new type of gene editing technique that targets a desired gene sequence in mammalian cells and the like. Since discovery of the CRISPR/Cas system, extensive research has been conducted to improve gene-editing efficiency, such as including double-strand breaks, and to develop a CRISPR/Cas system in a form (for example, small), which can be easily delivered into cells, such as AAV.

[0004] Double-strand DNA breaks within cells can be repaired through a DNA repair mechanism called non-homologous end joining (NHEJ) or homology-directed repair (HDR). For non-homologous end joining (NHEJ), random insertion or deletion of bases occurs at double-strand break sites, which results in frameshift mutation or premature mutation in the gene where DNA double-strand breaks have occurred, thereby knocking out the gene. On the other hand, homology-directed repair (HDR) requires a donor DNA (homologous template) to repair double-strand breaks, and a new sequence with the sequence of this donor DNA as a template is introduced into a break site. Gene editing (for example, knock-out of a specific gene or introduction of a new gene caused by indel) is mainly accomplished by utilizing the NHEJ or HDR pathway in cells.

[0005] Meanwhile, it is known that some genetic diseases can be treated by deletion or removal of a specific gene segment (for example, specific exon) which has undergone a genetic mutation or becomes problematic due to the genetic mutation. For example, Duchenne muscular dystrophy, which is caused by a mutation in the dystrophin gene, can be treated by deleting the entire exon 51 (exon skipping), which has undergone a frameshift mutation or premature-termination mutation, thereby allowing for production of a protein having a near-normal function. However, despite the existence of such a definitive therapeutic strategy, effective deletion of a specific gene segment with a relatively long sequence remains challenging.

[0006] Despite recent advances in gene editing techniques, there is still a need for a gene editing technique that can enhance removal (large deletion) of a gene segment or deletion efficiency therefor.

DISCLOSURE OF INVENTION

Technical Problem

[0007] The object of the present disclosure is to solve the above-mentioned problems of the prior art.

[0008] In addition, an object of the present disclosure is to provide a composition and a method for increasing deletion efficiency for a nucleic acid segment.

[0009] In addition, another object of the present disclosure is to provide a gene editing system with increased deletion efficiency of a nucleic acid segment.

[0010] The object of the present disclosure is not limited to the above-mentioned objects. The objects of the present disclosure will become clearer from the following description and may be realized by means and combinations thereof as set forth in the claims.

Solution to Problem

[0011] Representative configurations of the present disclosure to achieve the above purpose are as follows.

[0012] According to an aspect of the present disclosure, there is provided a composition for increasing deletion of a nucleic acid segment, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule, or a use of a composition for increasing deletion of a nucleic acid segment, the composition comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule.

[0013] In an embodiment, the gene involved in non-homologous end joining pathway may comprise at least one selected from the group consisting of ATM1, XRCC4, XLF, XRCC6, LIG4, and DCLRE1C.

[0014] In an embodiment, the gene involved in non-homologous end joining pathway may comprise at least one selected from the group consisting of XRCC6 and DCLRE1C.

[0015] In an embodiment, the inhibitory molecule may be shRNA, dsRNA, siRNA, miRNA, or an antisense oligonucleotide.

[0016] In an embodiment, the shRNA molecule may comprise at least one selected from the group consisting of shXRCC6 and shDCLRE1C.

[0017] In an embodiment, the shRNA molecule may comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS: 360 to 389, SEQ ID NOS: 400 to 430, SEQ ID NOS: 433 to 445, and SEQ ID NO: 473.

[0018] In an embodiment, the shRNA molecule may comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS: 375 to 379, SEQ ID NOS: 385 to 389, SEQ ID NOS: 410 to 414, SEQ ID NOS: 420 to 424, and SEQ ID NOS: 433 to 445.

[0019] In an embodiment, the composition may comprise at least two of the molecules that inhibit expression of a gene involved in non-homologous end joining (NHEJ) pathway or nucleic acid constructs encoding the molecules.

[0020] In an embodiment, the at least two inhibitory molecules may each inhibit expression of the same or different genes.

[0021] In an embodiment, the composition may further comprise (i) a Cas endonuclease and (ii) a first guide RNA and a second guide RNA that hybridize to a first target

sequence and a second target sequence, respectively; the nucleic acid segment may exist between the first target sequence and the second target sequence; and the Cas endonuclease may form a complex with the first guide RNA to induce first cleavage, and the Cas endonuclease may form a complex with the second guide RNA to induce second cleavage.

[0022] In an embodiment, the composition may further comprise at least one nucleic acid construct comprising (i) a nucleic acid encoding a Cas endonuclease, (ii) a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence, and (iii) a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence; the nucleic acid segment may exist between the first target sequence and the second target sequence; and the Cas endonuclease may form a first complex with the first guide RNA to induce first cleavage, and the Cas endonuclease may form a second complex with the second guide RNA to induce second cleavage.

[0023] In an embodiment, the first cleavage and the second cleavage may be each independently a single-strand DNA break or a double-strand DNA break.

[0024] In an embodiment, the composition or nucleic acid construct may comprise an additional guide RNA that hybridizes to an additional target sequence or a nucleic acid encoding the additional guide RNA.

[0025] In an embodiment, the Cas endonuclease may be an endonuclease comprising a Cas12f1 protein.

[0026] In an embodiment, the Cas12f1 protein may have at least 70% sequence identity to the amino acid sequence of SEQ ID NO: 1 or 5.

[0027] In an embodiment, the first or second guide RNA may be an engineered guide RNA.


[0028] In an embodiment, the engineered guide RNA may comprise a U-rich tail sequence linked to the 3'-end of its guide sequence, in which the U-rich tail is represented by $5'-(U_mV)_nU_o-3'$, wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5.

[0029] In an embodiment, the engineered guide RNA may comprise a nucleotide sequence having at least 50% sequence identity to a scaffold region of a wild-type Cas12f1 guide RNA sequence that sequentially comprises, from the 5'-end, a first stem-loop region, a second stem-loop region, a third stem-loop region, a fourth stem-loop region, and a tracrRNA-crRNA complementarity region, and the engineered guide RNA may comprise at least one modification selected from the group consisting of the following (1) to (5) with respect to the wild-type Cas12f1 guide RNA sequence:

- [0030]** (1) deletion of at least a part of the first stem-loop region;
- [0031]** (2) deletion of at least a part of the second stem-loop region;
- [0032]** (3) deletion of at least a part of the tracrRNA-crRNA complementarity region;
- [0033]** (4) replacement of one or more uracil (U) residues with A, G, or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region; and
- [0034]** (5) addition of a U-rich tail to the 3'-end of its crRNA sequence (a sequence of the U-rich tail is represented by $5'-(U_mV)_nU_o-3'$, wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5).

[0035] In an embodiment, the wild-type Cas12f1 guide RNA may comprise tracrRNA comprising the nucleotide sequence of SEQ ID NO: 11 and crRNA comprising the nucleotide sequence of SEQ ID NO: 12.

[0036] In an embodiment, the engineered guide RNA may consist of a sequence represented by Formula (I) or has at least 80% sequence identity thereto:



[0037] in Formula (I),

[0038] X^a , X^{b1} , X^{b2} , X^{c1} , and X^{c2} each independently consist of 0 to 35 (poly)nucleotides,

[0039] X^g is a first or second guide sequence,

[0040] Lk is a polynucleotide linker of 2 to 20 nucleotides or is absent, and

[0041] $(U_mV)_nU_o$ is present as a U-rich tail or absent, and when $(U_mV)_nU_o$ is present, U is uridine, V is each independently A, C, or G, m and o are each independently an integer between 1 and 20, and n is an integer between 0 and 5.

[0042] In an embodiment, X^a may comprise the nucleotide sequence of SEQ ID NO: 14 or a nucleotide sequence having the sequence of SEQ ID NO: 14 from which 1 to 20 nucleotides are deleted.

[0043] In an embodiment, X^{b1} may comprise the nucleotide sequence of SEQ ID NO: 25 or a nucleotide sequence having the sequence of SEQ ID NO: 25 from which 1 to 13 nucleotides are deleted.

[0044] In an embodiment, X^{b2} comprises the nucleotide sequence of SEQ ID NO: 29 or a nucleotide sequence having the sequence of SEQ ID NO: 29 from which 1 to 14 nucleotides are deleted.

[0045] In an embodiment, the sequence $5'-X^{b1}UUAGX^{b2}-3'$ in Formula (I) may be a nucleotide sequence selected from the group consisting of SEQ ID NOS: 34 to 38.

[0046] In an embodiment, X^{c1} comprises the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having the sequence of SEQ ID NO: 39 from which 1 to 28 nucleotides are deleted.

[0047] In an embodiment, in a case where three or more consecutive uracil (U) residues are present in a sequence of X^{c1} , the sequence of X^{c1} comprises a modification in which at least one U residue thereof is replaced with A, G, or C.

[0048] In an embodiment, X^{c2} comprises the nucleotide sequence of SEQ ID NO: 58 or a nucleotide sequence having the sequence of SEQ ID NO: 58 from which 1 to 27 nucleotides are deleted.

[0049] In an embodiment, in a case where the sequence $5'-ACGAA-3'$ is present in X^{c2} , the sequence may be replaced with $5'-NGNNN-3'$, and N may be each independently A, C, G, or U.

[0050] In an embodiment, the sequence 5'-X^{c1}-Lk-X^{c2}-3' in Formula (I) may be a nucleotide sequence selected from the group consisting of SEQ ID NOS: 80 to 86.

[0051] In an embodiment, Lk may comprise a nucleotide sequence selected from the group consisting of 5'-GAAA-3', 5'-UUAG-3', 5'-UGAAAA-3', 5'-UUGAAAA-3', 5'-UUCGAAAGAA-3' (SEQ ID NO: 76), 5'-UUCAGAAUGAA-3' (SEQ ID NO: 77), 5'-UUC AUGAAAAUGAA-3' (SEQ ID NO: 78), and 5'-UUC CAUUGAAAAUGAA-3' (SEQ ID NO: 79).

[0052] In an embodiment, the engineered guide RNA may comprise an engineered tracrRNA having a nucleotide sequence selected from the group consisting of SEQ ID NOS: 87 to 132.

[0053] In an embodiment, the engineered guide RNA may comprise an engineered crRNA sequence having a nucleotide sequence selected from the group consisting of SEQ ID NOS: 133 to 148.

[0054] In an embodiment, the engineered guide RNA may be a dual guide RNA or a single guide RNA.

[0055] In an embodiment, the engineered single guide RNA may comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS: 149 to 186.

[0056] In an embodiment, the nucleic acid construct may be or be contained in an adeno-associated virus vector.

[0057] In an embodiment, the composition may not comprise a donor sequence required for homology-directed repair (HDR) pathway or a nucleic acid construct encoding the donor sequence.

[0058] According to another aspect of the present disclosure, there is provided a method for increasing deletion of a nucleic acid segment in a target gene of a cell, comprising bringing, into contact with the cell, the above-described composition.

[0059] In an embodiment, the nucleic acid segment may comprise a gene fragment that needs to be removed for gene correction.

[0060] In an embodiment, the nucleic acid segment may comprise a gene fragment that needs to be removed for treating a genetic disease.

[0061] In an embodiment, the nucleic acid segment may comprise a mutated sequence.

[0062] In an embodiment, the nucleic acid segment may comprise a mutated sequence that causes a genetic disease.

[0063] According to yet another aspect of the present disclosure, there is provided a kit or system for gene editing, comprising: a composition for increasing deletion of a nucleic acid segment, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule; and a gene editing composition for deletion of a nucleic acid segment, comprising (i) a Cas endonuclease and a first guide RNA and a second guide RNA that hybridize to a first target sequence and a second target sequence, respectively, or (ii) one or more nucleic acid constructs comprising a nucleic acid encoding a Cas endonuclease, a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence, and a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence, wherein the nucleic acid segment exists between the first target sequence and the second target sequence, and the Cas endonuclease forms a complex with

the first guide RNA to induce first cleavage, and the Cas endonuclease forms a complex with the second guide RNA to induce second cleavage.

[0064] Regarding the respective components of the kit and system for gene editing, it is clear that reference may be made to the detailed description disclosed herein, and reference may also be made to all embodiments or combinations thereof described for the compositions or methods disclosed herein.

Advantageous Effects of Invention

[0065] The present inventors have identified that in a case of inducing deletion or removal of a nucleic acid segment with a CRISPR/Cas12f1 system, efficiency of the deletion or removal actually increases when expression or activity of a factor involved in the NHEJ repair pathway is inhibited, thereby completing the present disclosure. According to an embodiment of the present disclosure, it has been shown that inhibition of NHEJ pathway factors by shRNA can increase efficiency of large-scale deletion induced by UnCas12f1. In particular, interference of DCLRE1C gene, which encodes the Artemis protein, and XCCR6 gene, which encodes the Ku70 protein, has been found to most effectively increase deletion efficiency. In addition, in an embodiment, it has been identified that deletion efficiency for a nucleic acid segment is increased in a case of using a gene-editing system, which is more efficient and has broader applications, together with shRNA that inhibits expression of DCLRE1C or XCCR6, wherein the system comprises a Cas12f1 protein (for example, UnCas12f1, CWCas12f1, or a variant protein thereof), which is a hypercompact nucleic acid cleavage protein, and an engineered guide RNA in which certain regions have been modified to exhibit excellent indel efficiency when used with the protein. Accordingly, the present disclosure can be utilized in gene therapy, gene correction, or the like that requires effective deletion of a relatively long nucleic acid segment, such as exon skipping.

BRIEF DESCRIPTION OF DRAWINGS

[0066] FIG. 1 illustrates a schematic diagram showing canonical DSB repair pathways including NHEJ and HDR.

[0067] FIG. 2 illustrates a schematic diagram showing how a patient with Duchenne muscular dystrophy is treated by deletion of a nucleic acid segment comprising exon 51 in the dystrophin gene.

[0068] FIG. 3 illustrates modification sites in the engineered guide RNA.

[0069] FIG. 4 illustrates a human codon-optimized amino acid sequence of the Cas12f1 protein (FIGS. 4A and 4C) and a human codon-optimized nucleotide sequence encoding the protein (FIGS. 4D to 4H).

[0070] FIG. 5 illustrates results obtained by comparing the effects of RNA PolIII-dependent promoters on inhibition of DCLRE1C expression.

[0071] FIGS. 6A to 6C illustrate results obtained by performing qPCR analysis on inhibited expression of respective NHEJ component genes caused by shRNA (mean±standard error, n=2 independent experiments).

[0072] FIGS. 7A to 7C illustrate results obtained by performing Western blot analysis on inhibited expression of respective NHEJ component genes caused by shRNA.

[0073] FIG. 8 illustrates that increased deletion efficiency for E51 is achieved by UnCas12f1 in a case of being treated with shRNA.

[0074] FIG. 9 illustrates deletion efficiency for E51 depending on treatment with various shRNAs (mean±standard error, n=3 independent experiments. * p<0.05, ** p<0.01).

[0075] FIG. 10A illustrates results obtained by identifying changes in deletion caused by shDCLRE1C using qPCR on day 3 after treatment of HEK293T cells according to Example 6.

[0076] FIG. 10B illustrates results obtained by measuring DCLRE1C mRNA levels using qPCR on day 3 after treatment of HEK293T cells according to Example 6.

[0077] FIG. 11 illustrates results obtained by performing whole genome sequencing (WGS) analysis to measure deletion efficiency for E51. The section indicated by 800 bp represents the analyzed 800 bp region between the two gRNAs.

[0078] FIG. 12 illustrates results obtained by identifying the effect of shDCLRE1C on increased deletion efficiency through complementation of DCLRE1C expression. Results from two independent experiments on HEK293T and AC16 cells are presented on agarose gels.

[0079] FIG. 13 illustrates results obtained by identifying deletion efficiency for E51 in DCLRE1C-KO cell line.

[0080] FIG. 14 illustrates results obtained by measuring murine Dclre1c mRNA levels depending on treatment with shRNAs for various mouse Dclre1c genes.

[0081] FIG. 15 illustrates a graph showing mRNA expression levels of ATM1 and XRCC4 measured in cells transfected with shRNAs according to an embodiment.

[0082] FIG. 16 illustrates a graph showing mRNA expression levels of XLF-1 and XRCC6 measured in cells transfected with shRNAs according to an embodiment.

[0083] FIG. 17 illustrates a graph showing mRNA expression levels of LIG4 and DCLRE1C measured in cells transfected with shRNAs according to an embodiment.

[0084] FIG. 18 illustrates a graph obtained by identifying, with qRT-PCR, deletion efficiency for exon 51 achieved by inhibited expression of NHEJ-related genes in HEK293 cells. The experiments using UnCas12f1 protein are indicated by “Cas12f1,” and the experiments using CWCas12f1 protein are indicated by “TaRGET.”

[0085] FIG. 19 illustrates a graph obtained by identifying, with qRT-PCR, deletion efficiency for exon 51 achieved by inhibited expression of NHEJ-related genes in AC16 cells. The experiments using UnCas12f1 protein are indicated by “Cas12f1,” and the experiments using CWCas12f1 protein are indicated by “TaRGET.”

[0086] FIG. 20 illustrates a graph obtained by identifying, with qRT-PCR, deletion efficiency for exon 51 achieved using a single type of shRNA or a combination of two or more types of shRNA in HEK293 cells.

[0087] FIG. 21 illustrates a graph obtained by identifying, with qRT-PCR, deletion efficiency for exon 51 achieved using a single type of shRNA or a combination of two or more types of shRNA in AC16 cells.

[0088] FIG. 22 illustrates a graph obtained by identifying, with qRT-PCR, deletion efficiency for exon 51 depending on the number of days post transfection using systems comprising shRNA according to an embodiment. The results are indicated by SaCas9, Cas12f1 (UnCas12f1), and TaRGET depending on the CRISPR protein used in the system.

[0089] FIG. 23 illustrates results obtained by analyzing deletion efficiency for exon 51 depending on treatment with shRNADCLRE1C and shXRCC6 (mean±standard error, n=2 independent experiments).

MODES FOR CARRYING OUT INVENTION

[0090] The detailed description of the present disclosure set forth below will be described with reference to specific drawings (only if there are drawings) with respect to specific embodiments in which the present disclosure may be practiced; however, the present disclosure is not limited thereto and is limited only by the appended claims, along with the full scope of equivalents to which such claims are entitled. It should be understood that various embodiments/examples of the present disclosure, although different, are not necessarily mutually exclusive. For example, a particular feature, structure, or characteristic described herein may be changed from one embodiment/example to another embodiment/example or implemented in combinations of embodiments/examples without departing from the technical spirit and scope of the present disclosure. Unless defined otherwise, technical and scientific terms used herein have the same meaning as generally used in the art to which the present disclosure belongs. For purposes of interpreting this specification, the following definitions will apply and whenever appropriate, terms used in the singular will also include the plural and vice versa.

[0091] Hereinafter, in order to enable those skilled in the art to easily practice the present disclosure, various preferred embodiments/examples of the present disclosure will be described in detail with reference to the attached drawings (only if there are drawings).

I. Definition

[0092] As used herein, “nucleic acid,” “nucleotide,” “nucleoside,” and “base” have the meanings commonly understood by those skilled in the art. Specifically, “nucleic acid” is a biological molecule composed of nucleotides, and is used interchangeably with polynucleotide. The nucleic acid comprises both DNA and RNA, which is double-stranded or single-stranded. “Nucleotide” is a unit composed of phosphoric acid, a pentose sugar, and a base (or nucleobase). In RNA (ribonucleic acid), the pentose sugar is ribose; and in DNA (deoxyribonucleic acid), the pentose sugar is deoxyribose. The nucleotide has one selected from adenine (A), guanine (G), cytosine (C), thymine (T), and uracil (U) as a nucleobase. Adenine, guanine, and cytosine exist both in RNA and DNA, thymine exists only in DNA, and uracil exists only in RNA. In addition, the pentose sugar and nucleobase constituting the nucleotide may be referred to as “nucleoside.” The nucleoside is classified into adenosine, thymidine, cytidine, guanosine, and uridine depending on the type of nucleobase. The abbreviations for base, nucleoside, and nucleotide may be identical and may be appropriately interpreted depending on the context. For example, the sequence 5'-UUUUU-3' may be a sequence of five consecutive bases (uracil residues), a sequence of five consecutive nucleosides (uridine residues), and/or a sequence of five consecutive nucleotides (uridine monophosphate). In addition, when describing a nucleic acid, RNA, and DNA, nucleotides constituting the same are abbreviated as uridine, adenosine, thymidine, cytidine, and guanosine according to the type of nucleoside. The above

abbreviation may be appropriately interpreted depending on the context. For example, RNA comprising a sequence of four consecutive uridine residues may be interpreted as RNA comprising four consecutive uridine monophosphate nucleotides. In addition, the terms nucleic acid, nucleotide, nucleoside, and base as used herein may include modified nucleic acids, nucleotides, nucleosides, and bases known in the art for improving, for example, safety or immunogenicity thereof.

[0093] As used herein, “target nucleic acid” or “target gene” refers to a nucleic acid or gene that is a subject of gene editing (for example, double-strand break or deletion of a gene segment) or targeted by a gene editing system (for example, Cas12f1 system or TaRGET system). These terms may be used interchangeably and refer to the same subject. Unless otherwise defined, the target gene may be a unique gene or nucleic acid possessed by a target cell (for example, a prokaryotic cell, a eukaryotic cell, an animal cell, a mammalian cell, or a plant cell), a gene or nucleic acid of external origin, or an artificially synthesized nucleic acid or gene, and may mean single-stranded or double-stranded DNA or RNA. The target gene or target nucleic acid may be a mutated gene involved in a genetic disease. In an embodiment, the target gene or target nucleic acid may be a gene having a mutation. In an embodiment, the target gene or target nucleic acid may be a mutated human gene.

[0094] As used herein, “target region” means a region of a target gene to which a guide RNA is designed to bind and in which cleavage occurs. The target region may comprise a target sequence. The target region may comprise a target sequence. In addition, in double-stranded nucleic acids, the target region may refer to a region that comprises a target sequence (included in a target strand) and a sequence complementary thereto (included in a non-target strand).

[0095] As used herein, “target sequence” refers to a sequence located in a target nucleic acid or a target gene, which is recognized by a guide RNA, or a sequence to be modified by a gene editing system such as CRISPR/Cas12f1 system. Specifically, the target sequence refers to a sequence complementary to a guide sequence included in a guide RNA or a sequence that binds complementarily to the guide sequence. The strand including the target sequence is referred to as a “target strand.” When the target nucleic acid or the target gene is single-stranded, the strand may be a target strand. When the target nucleic acid or the target gene is double-stranded, one of the double strands may be a target strand, and a strand complementary to the target strand may exist. The strand complementary to the target strand is referred to as a “non-target strand.” The “non-target strand” comprises a PAM (Protospacer Adjacent Motif) sequence and a protospacer sequence. The PAM sequence is a sequence recognized by Cas endonucleases such as Cas12f1 protein. The protospacer sequence, which is located at the 5'-end or the 3'-end of the PAM sequence, is a sequence having complementarity to a target sequence or a sequence that forms a complementary bond with a target sequence. Correlation between the protospacer sequence and the target sequence is similar to correlation between the target sequence and the guide sequence. Due to these characteristics, in general, a guide sequence may be designed using a protospacer sequence. That is, a guide sequence which binds complementarily to a target sequence may be designed as a nucleotide sequence having the same nucleotide sequence as

the protospacer sequence, and the guide sequence is designed by replacing T in the protospacer sequence with U.

[0096] As used herein, “stem” refers to a nucleic acid region having a secondary structure that comprises a nucleotide region capable of forming a double strand. A configuration in which a double strand is connected primarily by a region of single-stranded nucleotides (a loop region) is referred to as a “stem-loop.” The terms “stem” and “stem-loop” may be used interchangeably and should be interpreted appropriately depending on the context.

[0097] The term “non-homologous end joining (NHEJ)” refers to a mechanism that repairs a double-strand break in a nucleotide sequence by direct ligation of the broken ends without the requirement for a homologous template (as opposed to homology-directed repair, which requires a homologous sequence to induce healing of a double-strand break in a nucleotide sequence). NHEJ often leads to loss (deletion) of a nucleotide sequence near the double-strand break site.

[0098] The term “vector,” unless otherwise specified, refers to any material capable of transporting a genetic material into a cell. For example, a vector may be a nucleic acid, typically a DNA molecule, comprising a genetic material of interest, for example, a nucleic acid encoding an effector protein (Cas protein) of a CRISPR/Cas system, and/or a nucleic acid encoding a guide RNA; however, the vector is not limited thereto.

[0099] The term “operably linked” means a functional linkage between two or more elements arranged in such a way that allows the described element to function in an intended manner. For example, when a promoter sequence is operably linked to a sequence encoding protein A, it means that the promoter is linked to the sequence encoding the protein A so as to transcribe and/or express the sequence encoding the protein A in a cell. In addition, the term includes all other meanings generally recognized by those skilled in the art and may be appropriately interpreted depending on the context.

[0100] The term “engineered” is used to distinguish a substance or molecule from one having a naturally occurring configuration, and means that the substance or molecule is obtained by application of artificial modification. For example, “engineered guide RNA” refers to a guide RNA obtained by applying an artificial modification to the configuration of a naturally occurring guide RNA.

[0101] The term “NLS (nuclear localization sequence or signal)” refers to an amino acid sequence that promotes introduction of a substance from outside the nucleus into the nucleus, for example, by nuclear transport. The term “NES (nuclear export sequence or signal)” refers to an amino acid sequence that promotes transport of a substance from inside the nucleus to the outside of the nucleus, for example, by nuclear transport. The terms NLS or NES are known in the relevant art and may be clearly understood by those skilled in the art.

[0102] The term “about” refers to an amount, level, value, number, frequency, percent, dimension, size, amount, weight or length that varies by approximately 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2, or 1% with respect to a reference amount, level, value, number, frequency, percent, dimension, size, amount, weight or length. For example, the term “about” may mean $x \pm 5\%$ when used in relation to a value x expressed as a number or numerical value.

[0103] The term “subject” is used interchangeably with “patient” and may be a mammal in need of prevention or treatment of a genetic disease, such as primate (for example, human), companion animal (for example, dog and cat), domestic animal (for example, cow, pig, horse, sheep, and goat), and laboratory animal (for example, rat, mouse, and guinea pig). In an embodiment of the present disclosure, the subject is a human.

[0104] The term “treatment” generally means obtaining a desired pharmacological and/or physiological effect. Such an effect has a therapeutic effect in that it partially or completely cures a disease and/or harmful effects caused by the disease. Desirable therapeutic effects include, but are not limited to, prevention of occurrence or recurrence of a disease, improvement of symptoms, reduction of any direct or indirect pathological consequences of a disease, prevention of metastasis, reduction of disease progression rate, improvement or alleviation of disease state, and remission or improved prognosis. Preferably, “treatment” may refer to medical intervention for an already manifested disease or disorder. More preferably, “treatment” may be deletion of a specific segment in a specific gene or restoration of the reading frame of the specific gene resulting therefrom.

[0105] As used herein, the term “target nucleic acid editing system,” “gene editing system,” or “gene restoration system” refers to a system that comprises a nucleic acid degrading enzyme, such as nucleic acid editing protein or endonuclease, and a nucleic acid-targeting molecule corresponding to the nucleic acid degrading enzyme, and this system binds to or interacts with a target nucleic acid or target gene so that a target region of the target nucleic acid or target gene can be cleaved, edited, repaired, and/or restored. Here, the nucleic acid-targeting molecule may be represented by an engineered guide RNA (gRNA), but is not limited thereto. Meanwhile, the target nucleic acid editing system may exist in any form capable of editing the target nucleic acid. For example, the system may be in a form of a composition that comprises a complex comprising a nucleic acid degrading enzyme and a nucleic acid-targeting molecule, may be in a form of a kit in which the nucleic acid degrading enzyme and the nucleic acid-targeting molecule are each included in separate compositions, or may be a vector system or composition comprising at least one vector that comprises a nucleic acid encoding the nucleic acid degrading enzyme and a nucleic acid encoding the nucleic acid-targeting molecule.

[0106] The term “hypercompact TaRGET system” refers to a gene editing system that comprises a nucleic acid degrading enzyme such as hypercompact CRISPR/Cas protein or tiny endonuclease (for example, Cas12f1 or a variant thereof) and a nucleic acid-targeting molecule corresponding to the nucleic acid degrading enzyme, and is used for differentiation from the existing gene editing system. Here, the nucleic acid-targeting molecule may be represented by an engineered guide RNA (gRNA), but is not limited thereto. The system may be any type of gene editing system capable of binding to a target nucleic acid or target gene so that a target region of the target nucleic acid or gene is cleaved, edited, repaired, and/or restored.

[0107] The term “nuclease” or “endonuclease” refers to an enzyme that possesses catalytic activity for DNA cleavage and may be used interchangeably.

[0108] The term “nuclease” or “endonuclease” refers to an enzyme that possesses catalytic activity for nucleic acid

cleavage and may be used interchangeably. These terms may also be used interchangeably with “nucleic acid editing protein,” “gene editing protein,” or “nucleic acid degrading protein.” The molecule referred to as “nucleic acid editing protein,” “gene editing protein,” or “nucleic acid degrading protein” refers to a (endo-) nuclease that recognizes the targeting nucleic acid, DNA or RNA, or a protospacer adjacent motif (PAM) present in a target gene, and then allows double-strand breaks (DSBs) to occur at nucleotide sequences within or outside the target nucleotide sequence. In addition, the endonuclease, the nucleic acid editing protein, or the like is also referred to as an effector protein that constitutes a nucleic acid construct for a nucleic acid editing system. Here, the effector protein may be a nucleic acid degrading protein capable of binding to a guide RNA (gRNA) or engineered gRNA, or may be a peptide fragment capable of binding to a target nucleic acid or target gene.

[0109] The term “guide RNA (gRNA)” refers to RNA that is capable of forming a complex with a molecule referred to as an endonuclease, a gene editing protein, a nucleic acid degrading protein, or the like, and interacting with (for example, hybridizing to, forming a complementary bond(s) with, or forming a hydrogen bond(s) with) a target nucleotide sequence, and comprises a guide sequence having sufficient complementarity with the target nucleotide sequence to cause sequence-specific binding of the complex to the target nucleotide sequence. In the present disclosure, a guide RNA and a guide molecule may be used interchangeably.

[0110] The terms “tracrRNA (trans-activating crRNA)” and “crRNA (CRISPR RNA)” include the meanings commonly understood by those skilled in the art. These terms may be used to refer to respective molecules of a dual guide RNA found in nature, and may also be used to refer to respective portions of a single guide RNA (sgRNA) in which the tracrRNA and the crRNA are connected by a linker. Unless otherwise stated, the description tracrRNA and crRNA means tracrRNA and crRNA that constitute a guide RNA.

[0111] The term “scaffold region” refers collectively to a portion of a guide RNA (gRNA) which can interact with a molecule called endonuclease, gene editing protein, nucleic acid degrading protein, or the like, and may be used to refer to the remaining portion of a guide RNA found in nature, excluding a spacer.

[0112] The terms “guide sequence,” “spacer,” or “spacer sequence” may be used interchangeably, and refer to a polynucleotide within the CRISPR/Cas system which is capable of interacting with (for example, hybridizing to, forming a complementary bond(s) with, or forming a hydrogen bond(s) with) a target sequence portion. For example, the guide sequence or spacer sequence refers to 10 to 50 consecutive nucleotides linked directly or indirectly through a linker or the like to or near the 3-end of crRNA, which constitutes a guide RNA, in a target nucleic acid editing system.

[0113] The term “wild type” has the meaning commonly understood by those skilled in the art and means a typical form of an organism, strain, gene, or characteristic as it occurs in nature to the extent that it is distinguishable from mutant or variant forms.

[0114] The term “variant” should be understood to mean expression of qualities having a pattern that deviates from

what occurs in nature. For example, the variant protein may mean a variant of (wild-type) Cas12f1.

[0115] The term “nucleic acid construct” refers to a nucleic acid molecule that comprises one or more distinct segments and/or elements. The term includes linear nucleic acid constructs (for example, λ -phage constructs, PCR products) as well as circular nucleic acid constructs such as plasmid constructs, viral vector constructs, cosmid vectors, and the like. In an embodiment, the nucleic acid construct is a single-stranded or double-stranded nucleic acid molecule, which may be modified to contain a nucleic acid segment. In an embodiment, the nucleic acid construct may be such that specific nucleic acids are operably linked. In another embodiment, the nucleic acid construct of the present disclosure is an expression vector, which comprises sequences that render it suitable for replication and integration in prokaryotes. In yet another embodiment, the expression vector is suitable for expression in eukaryotes. In still yet another embodiment, the expression vector is suitable for expression in both prokaryotes and eukaryotes (for example, a shuttle vector). The expression vector also comprises an expression signal such as a promoter and/or an enhancer. Nucleotide sequences required for expression in prokaryotes often include promoters, operators (non-essential), and ribosomal binding sites along with other sequences. Eukaryotic cells are known to utilize promoters, enhancers, and termination and polyadenylation signals.

[0116] The terms “protein,” “polypeptide,” and “peptide” may be used interchangeably and refer to a polymeric form of amino acids of any length which may comprise genetically coded and non-genetically coded amino acids, chemically or biochemically modified or derivatized amino acids, and polypeptides having modified peptide backbones. The terms include fusion proteins, including, but not limited to, fusion proteins with a heterologous amino acid sequence, fusions with heterologous and homologous leader sequences, with or without N-terminal methionine residues; immunologically tagged proteins; and the like.

[0117] The term “hybridizable” means, for example, that a guide sequence has at least 60%, for example at least 70%, at least 80%, at least 90%, at least 95%, at least 98%, at least 99%, or 100% sequence complementarity to a target sequence or has less than 5 mismatches and binds complementarily to the target sequence.

[0118] The term “microRNA” or “miRNA” is used interchangeably and has the meaning commonly understood by those skilled in the art. miRNA is a small, non-coding, single-stranded RNA of approximately 22 nucleotides in length (typically between 19 and 25 nucleotides in length). miRNA typically targets more than one gene. Targeting with miRNA allows mismatches and inhibition of mRNA translation is mediated by incomplete complementarity, whereas siRNA and shRNA are specific to their targets due to perfect sequence complementarity.

[0119] The term “siRNA” refers to a small interfering or short interfering RNA and has the meaning commonly understood by those skilled in the art. siRNA refers to an RNA duplex of nucleotides that can target a desired gene and inhibit expression of a gene that shares homology therewith. siRNA is formed from a long double strand RNA (dsRNA) or shRNA.

[0120] The term “shRNA” refers to a small hairpin RNA or a short hairpin RNA, and has the meaning commonly understood by those skilled in the art. shRNA refers to an

RNA duplex in which the siRNA portion is part of a hairpin structure. shRNA may be processed into functional siRNA in cells.

[0121] The term “antisense oligonucleotide” refers to a nucleotide sequence that can interact and/or hybridize with a target sequence in pre-mRNA or mRNA molecule having a complementary nucleotide sequence, thereby modifying gene expression, the meaning of which will be clearly understood by those skilled in the art.

[0122] All technical terms used in the present disclosure, unless otherwise defined, have meanings commonly understood by those skilled in the relevant technical field and may be interpreted appropriately depending on the context.

II. Increased Deletion Efficiency for Nucleic Acid Segment Caused by Inhibited Expression of Genes Involved in NHEJ Pathway

[0123] In mammalian cells, the “canonical” or “classical” NHEJ pathway (C-NHEJ) requires several factors, including DNA-PK, Ku70-80, Artemis, ligase IV (Lig4), XRCC4, CLF, and Pol μ , to repair double-strand breaks (Kasperek & Humphrey Seminars in Cell & Dev. Biol. 22:886-897, 2011). In a case of artificially inducing a double-strand break in a target gene using a gene-editing system such as Cas12f1, random base insertion and deletion (Indel) occur at the double-strand break site through the NHEJ pathway in a cell. The present disclosure is based, in part, on the finding that, when it is desired to induce large-scale deletion of a relatively long nucleic acid segment, that is, a nucleotide sequence, in a target gene, increased deletion efficiency for the nucleic acid segment is achieved by inhibiting expression or activity of factors (in particular, XRCC6 and DCLRE1C) involved in the NHEJ repair pathway.

[0124] According to an aspect of the present disclosure, there is provided a composition for increasing deletion of a nucleic acid segment in a target gene, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining pathway, or a nucleic acid construct encoding the molecule.

[0125] In an embodiment, the gene involved in non-homologous end joining pathway may comprise at least one selected from the group consisting of ATM1, XRCC4, XLF, XRCC6, LIG4, and DCLRE1C. The ATM1 gene encodes ATM1 protein. The XRCC4 gene encodes XRCC4 protein. The XLF gene encodes XLF protein. The XRCC6 gene encodes Ku70 protein. The LIG4 gene encodes DNA ligase IV protein. The DCLRE1C gene encodes ARTEMIS protein. The proteins expressed by these six genes are all known to be involved in NHEJ. Specifically, KU70 is a DNA recognition protein that binds to the ends of double-strand breaks to assist in DNA repair; ARTEMIS participates in end processing during NHEJ repair; and XLF-XRCC4-DNA ligase IV complex provides a scaffold for joining double-strand breaks, ensuring accurate and efficient ligation and repair.

[0126] In another embodiment, the gene involved in non-homologous end joining pathway may comprise at least one selected from the group consisting of XRCC6 and DCLRE1C.

[0127] In an embodiment, the inhibitory molecule may be a small molecule or an inhibitory nucleic acid.

[0128] In an embodiment, the inhibitory molecule may be shRNA, dsRNA, siRNA, miRNA, or an antisense oligonucleotide. Preferably, the inhibitory molecule may be

shRNA, siRNA, or an antisense oligonucleotide. More preferably, the inhibitory molecule may be shRNA or siRNA.

[0129] In another embodiment, the inhibitory molecule may be targeted to enzymes involved in NHEJ, HDR, or upstream regulation thereof, by post translational modification, for example, through phosphorylation, ubiquitination, and/or sumoylation.

[0130] In another embodiment, the composition may comprise an shRNA molecule, a dsRNA molecule, a siRNA molecule, a miRNA molecule, or an antisense oligonucleotide molecule that inhibits expression of an XRCC6 gene; an shRNA molecule, a dsRNA molecule, a siRNA molecule, a miRNA molecule, or an antisense oligonucleotide molecule that inhibits expression of a DCLRE1C gene; or a combination thereof.

[0131] In yet another embodiment, the shRNA molecule may comprise at least one selected from the group consisting of shXRCC6 and shDCLRE1C.

[0132] In still yet another embodiment, the inhibitory molecule or shRNA molecule may comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS: 360 to 389, SEQ ID NOS: 400 to 430, SEQ ID NOS: 433 to 445, and SEQ ID NO: 473.

[0133] In still yet another embodiment, the inhibitory molecule or shRNA molecule may comprise a nucleotide sequence selected from the group consisting of SEQ ID NOS: 375 to 379, SEQ ID NOS: 385 to 389, SEQ ID NOS: 410 to 414, SEQ ID NOS: 420 to 424, and SEQ ID NOS: 433 to 445.

[0134] In an embodiment, the composition may comprise at least one nucleic acid construct encoding a molecule that inhibits expression of a gene involved in non-homologous end joining pathway.

[0135] In an embodiment, the composition may comprise at least two of the above-described nucleic acid constructs encoding molecules that inhibit expression of a gene involved in non-homologous end joining pathway. Specifically, the at least two of the nucleic acid constructs may each encode a molecule that inhibits expression of the same gene, or may encode molecules that inhibit expression of different genes. For example, the composition may comprise at least two of the nucleic acid constructs encoding molecules (for example, shXRCC6) that inhibit XRCC6. For example, the composition may comprise at least two of the nucleic acid constructs encoding molecules (for example, shDCLRE1C) that inhibit expression of DCLRE1C. In addition, the composition may comprise a nucleic acid construct encoding a molecule (for example, shXRCC6) that inhibits XRCC6 and a nucleic acid construct encoding a molecule (for example, shDCLRE1C) that inhibits expression of DCLRE1C.

[0136] In an embodiment, the nucleic acid construct may be or comprise the above-described nucleotide sequence encoding a molecule that inhibits expression of a gene involved in non-homologous end joining pathway.

[0137] In an embodiment, the nucleic acid construct may comprise, in an operably-linked form, the above-described nucleotide sequence encoding a molecule that inhibits expression of a gene involved in non-homologous end joining pathway. For example, the nucleic acid construct may comprise the above-described nucleotide sequence encoding a molecule that inhibits expression of a gene involved in non-homologous end joining pathway, in a form

of being operably linked to another element, such as a promoter, that enables the nucleotide sequence to be expressed in a cell.

[0138] In an embodiment, the nucleic acid construct may be a vector. In another embodiment, at least two of the nucleic acid constructs according to an embodiment of the present disclosure may be contained in a single vector. Specifically, the at least two of the nucleic acid constructs may each encode the above-described molecule that inhibits expression of the same gene involved in non-homologous end joining pathway, and may also encode the above-described molecules that inhibit expression of different genes. For example, at least two of the nucleic acid constructs encoding shXRCC6 may be contained in a single vector. For example, the vector may comprise at least one nucleic acid construct encoding shXRCC6 and at least one nucleic acid construct encoding shDCLRE1C. Details on vectors or vector systems are given in a separate section below.

[0139] In another aspect, the composition may be a gene editing composition for deleting a nucleic acid segment (for example, a CRISPR/Cas system such as a CRISPR/Cas12f1 system). In an embodiment, the composition may be a gene editing composition for deleting a nucleic acid segment in a gene of a cell. In another embodiment, the composition may be a gene editing composition or an adjuvant composition for gene editing to increase deletion efficiency for a nucleic acid segment in a gene of a cell. Specifically, the composition may further comprise (i) a Cas endonuclease and (ii) a first guide RNA and a second guide RNA that hybridize to a first target sequence and a second target sequence, respectively, in a target gene. The nucleic acid segment may exist between the first target sequence and the second target sequence, and the Cas endonuclease may form a complex with the first guide RNA to induce first cleavage, and the Cas endonuclease may form a complex with the second guide RNA to induce second cleavage.

[0140] In yet another aspect, the composition may further comprise at least one nucleic acid construct comprising (i) a nucleic acid encoding a Cas endonuclease, (ii) a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence in a target gene, and (iii) a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence in the target gene. The nucleic acid segment exists between the first target sequence and the second target sequence. Once the nucleic acid constructs are expressed in a cell, the Cas endonuclease may form a first complex with the first guide RNA to induce first cleavage, and the Cas endonuclease may form a second complex with the second guide RNA to induce second cleavage.

[0141] The endonuclease, together with the guide RNAs, may be referred to as a gene editing system, a CRISPR/Cas system, or a CRISPR/Cas12f1 system.

[0142] In an embodiment, the first cleavage and the second cleavage may be each independently a single-strand DNA break or a double-strand DNA break.

[0143] In another embodiment, the composition or nucleic acid construct may comprise an additional guide RNA that hybridizes to an additional target sequence in the target gene or a nucleic acid encoding the same. The target sequences may overlap with each other.

[0144] In another embodiment, the guide RNA may target a region adjacent to a nucleic acid segment to be deleted in

a target gene to generate cleavage (for example, a single-strand break or a double-strand break).

[0145] In yet another embodiment, two guide RNAs may target the upstream and downstream regions, respectively, of a nucleic acid segment to be deleted in a target gene, thereby generating at least one cleavage (for example, two single-strand breaks or two double-strand breaks).

[0146] In still yet another embodiment, at least two guide RNAs may be used to generate at least two sets of cleavage (for example, two double-strand breaks, one double-strand break, and one single-strand break; or two pairs of single-strand breaks).

[0147] Without wishing to be bound by any particular theory, it is thought that the composition of the present disclosure, which comprises a molecule that inhibits expression of a gene involved in non-homologous end joining pathway or a nucleic acid construct encoding the molecule, blocks repair of a double-strand break site by the NHEJ pathway when the double-strand break is formed by a gene editing system, thereby maintaining the double-strand break state for a long time, which increases likelihood that a double break can occur on both sides of a nucleic acid segment to be deleted so that the nucleic acid segment is removed.

[0148] Details on the gene editing system, CRISPR/Cas system, and CRISPR/Cas12f1 system included in the composition of the present disclosure will be given in a separate section below.

[0149] In an embodiment, the endonuclease comprising the Cas protein and the guide RNA may be included in the form of a ribonucleoprotein particle (RNP).

[0150] In an aspect, the composition of the present disclosure may not comprise a donor sequence required for homology-directed repair (HDR) pathway or a nucleic acid construct encoding the donor sequence. Without wishing to be bound by any particular theory, the composition of the present disclosure is intended to remove a specific nucleic acid segment by a double-strand break.

[0151] According to another aspect of the present disclosure, there is provided a method for increasing deletion of a nucleic acid segment in a target gene of a cell, comprising bringing, into contact with the cell, the composition according to the above-described embodiment.

[0152] In an embodiment, the nucleic acid segment may be a gene fragment that needs to be removed or deleted for various purposes, such as gene therapy or improvement of breed. For example, the nucleic acid segment may be a gene fragment that needs to be removed for gene correction. In addition, the nucleic acid segment may comprise a gene fragment that needs to be removed for treatment of a genetic disease.

[0153] In an embodiment, the nucleic acid segment may have a length of 1 bp to 1000 kbp, 100 bp to 1000 kbp, 200 bp to 1000 kbp, 300 bp to 1000 kbp, 400 bp to 1000 kbp, 500 bp to 1000 kbp, 600 bp to 1000 kbp, 700 bp to 1000 kbp, 800 bp to 1000 kbp, 900 bp to 1000 kbp, 1 kbp to 1000 kbp, 100 kbp to 1000 kbp, 200 kbp to 1000 kbp, 300 kbp to 1000 kbp, 400 kbp to 1000 kbp, 500 kbp to 1000 kbp, 600 kbp to 1000 kbp, 700 kbp to 1000 kbp, 800 kbp to 1000 kbp, or 900 kbp to 1000 kbp. In addition, the length of the nucleic acid segment may have a length within any range of the specified numerical ranges, which can be readily determined and understood by those skilled in the art.

[0154] In an embodiment, the nucleic acid segment may comprise a (undesirable) mutated sequence. In another embodiment, the nucleic acid segment may comprise a sequence altered by a genetic mutation. The mutation may include, but is not limited to, frameshift mutations, premature termination mutations, point mutations, mutations caused by expanded repeat sequences, and mutations caused by repeat sequences.

[0155] In an embodiment, the nucleic acid segment may comprise a mutated sequence that causes a genetic disease. For example, the mutated sequence that causes a genetic disease may be, but is not limited to, a frameshift mutation or a premature termination mutation in dystrophin exon 51 which causes Duchenne dystrophy, or a point mutation where adenine in intron 26 of the CEP290 gene is replaced with guanine which causes Leber congenital amaurosis type 10 (LCA10).

[0156] In another embodiment, the cell may be a prokaryotic cell, a eukaryotic cell, an animal cell, a mammalian cell, or a plant cell. Specifically, the cell may be an animal cell, a mammalian cell, or a human cell. In addition, the cell may be a plant cell.

[0157] In an embodiment, the bringing into contact with the cell may be delivery or introduction of a composition comprising a nucleic acid construct into the cell. Specifically, the nucleic acid construct (for example, a vector) may be delivered or introduced into a cell, for example, by in vivo electroporation, liposomes, nanoparticles, or DNA injection or DNA vaccination, with or without a recombinant vector.

[0158] In addition, the composition comprising a nucleic acid construct of the present disclosure may be delivered or introduced by a virus, such as a retrovirus, a lentivirus, an adenovirus, an adeno-associated virus, a vaccinia virus, a poxvirus, a herpes simplex virus or a phage. Specifically, the composition may be contained in a packaging virus and delivered into a cell in the form of a virus produced by the packaging virus.

[0159] Specifically, the bringing-into-contact, delivery, or introduction may be made by electroporation, gene gun, sonoporation, magnetofection, nanoparticles, and/or transient cell compression or squeezing method. When the cell is a eukaryotic cell, cationic liposome method, lithium acetate-DMSO, lipid-mediated transfection, calcium phosphate precipitation, lipofection, polyethylenimine (PEI)-mediated transfection, DEAE-dextran-mediated transfection, and/or nanoparticle-mediated nucleic acid delivery (see Panyam et al., *Adv Drug Deliv Rev.* 2012 Sep. 13. pii: 50169-409X(12)00283-9) may be used.

[0160] In another embodiment, the bringing-into-contact, delivery, or introduction may be performed in vitro, in vivo, or ex vivo.

[0161] According to yet another aspect of the present disclosure, there is provided a kit or system for gene editing, comprising: a composition for increasing deletion of a nucleic acid segment, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule; and a gene editing composition for deletion of a nucleic acid segment, comprising (i) a Cas endonuclease and a first guide RNA and a second guide RNA that hybridize to a first target sequence and a second target sequence, respectively, or (ii) at least one nucleic acid construct comprising a nucleic acid encoding a Cas endo-

nuclease, a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence, and a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence, wherein the nucleic acid segment exists between the first target sequence and the second target sequence, and the Cas endonuclease forms a complex with the first guide RNA to induce first cleavage, and the Cas endonuclease forms a complex with the second guide RNA to induce second cleavage.

[0162] For respective components of the kit and system for gene editing, it is clear that reference may be made to the detailed description disclosed herein, and reference may also be made to all embodiments or combinations thereof described for the compositions or methods disclosed herein.

III. CRISPR/Cas System

[0163] As disclosed herein, the CRISPR/Cas12f1 system is provided as a CRISPR/Cas system for editing or modifying a target gene; however, the CRISPR/Cas system is not limited thereto. The CRISPR/Cas system may be a CRISPR/Cas9 system, a CRISPR/Cas10 system, a CRISPR/Cas12f1 system, or a CRISPR/TnpB system. Recent studies have shown that transposon-associated transposase B (TnpB) can be used as an endonuclease for the CRISPR/Cas system (Altae-Tran H, Kannan S, Demircioglu F E, Oshiro R, Nety S P, McKay U, Dlakic M, Inskeep W P, Makarova K S, Macrae R K, Koonin E V, Zhang F The widespread IS200/IS605 transposon family encodes diverse programmable RNA-guided endonucleases. *Science*. 2021 October; 374(6563):57-65. doi: 10.1126/science.abj6856. Epub 2021 Sep. 9. PMID: 34591643; PMCID: PMC8929163).

[0164] The disclosed system comprises (i) an endonuclease comprising at least one Cas protein or a nucleic acid construct encoding the endonuclease and (ii) at least one (for example, two) guide RNA molecule or a nucleic acid construct encoding the molecule.

[0165] In an embodiment, the Cas protein may be a Cas 9 protein, a Cpf1 protein, a Cas12f1 protein, or a TnpB protein.

[0166] In another embodiment, the Cas protein may be a Cas12f1 protein. The Cas12f1 protein may be in a wild-type, variant, or engineered form.

[0167] The present inventors have confirmed that TnpB (Transposon-associated transposase B) protein derived from *Candidatus Woesearchaeota* archaeon has an amino acid sequence similar to the UnCas12f1 protein (and thus, TnpB having an amino acid sequence similar to the UnCas12f1 protein is also named CWCas12f1; CWCas12f1 may be collectively referred to as Cas12f1 protein together with UnCas12f1, and may belong to a variant of Cas12f1 in its relationship with UnCas12f1), has a molecular weight that is about 1/3 smaller than that of an existing nucleic acid degrading protein including the Cas9 protein, which has been studied the most to date, and has significantly higher nucleic acid cleavage efficiency for a target nucleic acid or target gene. In addition, the present inventors have confirmed that engineered guide RNAs having a small size obtained by modifying the wild-type Cas12f1 guide RNA may induce excellent nucleic acid cleavage efficiency (for example, a double-strand break) together with the Cas12f1 protein such as CwCas12f1 or UnCas12f1. The hypercompact gene editing system comprising an engineered guide RNA and Cas12f1 or a variant thereof, such as CWCas12f1 or UnCas12f1, disclosed herein may be referred to as

“CRISPR/Cas12f1 system” or “TaRGET system,” and these terms may be used interchangeably. (However, for convenience, in the examples, the system using the UnCas12f1 protein is referred to as Cas12f1 system, and the system using the CWCas12f1 protein is referred to as TaRGET system). Hereinafter, respective components of the Cas12f1 gene editing system are described.

1. Endonuclease Comprising Cas12f1 Protein

[0168] The gene editing system based on CRISPR/Cas12f1 comprises an endonuclease comprising Cas12f1 or a nucleic acid encoding the endonuclease. The Cas12f1 protein is a (small) endonuclease characterized by exhibiting excellent activity in cleaving a target site of a target nucleic acid and being significantly smaller in size by about 1/3 compared to the existing CRISPR/Cas9 system.

[0169] Cas12f1 is one of the effector proteins named Cas14 in a previous study (see Harrington et al., *Science*, 362, 839-842, 2018), and is also called Cas14a1 protein. The Cas12f1 protein disclosed herein may be a wild-type Cas12f1 protein existing in nature. In addition, the Cas12f1 protein may be a variant of the wild-type Cas12f1 protein. A variant of Cas12f1 is referred to as a “Cas12f1 variant.” The Cas12f1 variant may be a variant having the same or equivalent function as the wild-type Cas12f1 protein, a variant of which some or all functions are modified, and/or a variant in which additional functions are added.

[0170] It has been reported that the Cas12f1 protein forms a complex with a guide RNA such that two Cas12f1 protein molecules bind to a guide RNA in the form of a dimer, and that all or part of the domain of the Cas12f1 protein recognizes a specific part of the scaffold region of the guide RNA to form a CRISPR/Cas12f1 complex (see Takeda et al., *Structure of the miniature type V-F CRISPR-Cas effector enzyme*, *Molecular Cell* 81, 1-13, 2021; and Xiao et al., *Structural basis for the dimerization-dependent CRISPR-Cas12f1 nuclease*, *bioRxiv*, 2020). Cas12f1 may generate a double-strand or single-strand break in a target nucleic acid or a target gene. Deletion of a desired gene segment may be induced by such a double-strand or single-strand break.

[0171] The Cas12f1 protein may recognize a protospacer adjacent motif (PAM) sequence present in a target nucleic acid or target gene. The PAM sequence is a unique sequence determined depending on the CRISPR protein. The PAM sequence recognized by Cas12f1 may be a T-rich sequence. The PAM sequence recognized by Cas12f1 may be a sequence of 5'-TTTR-3', wherein R may be T, A, C, or G. Preferably, the PAM sequence may be 5'-TTTA-3', 5'-TTTT-3', 5'-TTTC-3' or 5'-TTTG-3'. More preferably, the PAM sequence may be 5'-TTTA-3' or 5'-TTTG-3'.

[0172] In an embodiment, the Cas12f1 protein may be derived from a Cas14 family (see Harrington et al., *Science* 362, 839-842 (2018); US 2020/0172886 A1).

[0173] In another embodiment, the Cas12f1 protein may be a Cas14a1 (UnCas12f1) protein derived from an uncultured archaeon (Harrington et al., *Science* 362, 839-842 (2018); US 2020/0172886 A1). For example, the UnCas12f1 protein may comprise or consist of the amino acid sequence of SEQ ID NO: 5 (see FIG. 4).

[0174] In another embodiment, the Cas12f1 protein may be TnpB (transposon-associated transposase B) protein derived from the *Candidatus Woesearchaeota* archaeon. The TnpB protein is a protein conventionally known as a transposase. To date, the TnpB protein has been known only as

a transposon-encoded nuclease, and it is not known whether the TnpB protein has Cas endonuclease activity. In addition, a guide RNA for the TnpB protein has also not been known. The present inventors have confirmed for the first time that TnpB variant or engineered TnpB, which is based on the TnpB protein sequence, has excellent endonuclease activity of targeting a target nucleic acid or a target gene and cleaving a double-stranded DNA of the target site while having a similar size to a Cas12f1 protein, which belongs to the group with the smallest molecular weight among nucleic acid degrading proteins, and have constructed an engineered guide RNA that exhibits excellent gene editing activity when used together with TnpB or a variant protein thereof. This TnpB protein is named CWCas12f1 protein. For example, the CWCas12f1 protein may comprise or consist of the amino acid sequence of SEQ ID NO: 1 (see FIG. 4).

[0175] In an embodiment, the Cas12f1 protein may be a Cas12f1 variant. The Cas12f1 variant may comprise a modification of at least one amino acid, such as deletion, substitution, insertion, or addition, compared to the amino acid sequence of the wild-type Cas12f1 protein.

[0176] In another embodiment, the Cas12f1 variant may comprise deletion of at least one amino acid or substitution with another amino acid sequence compared to the amino acid sequence of the wild-type Cas12f1 protein (for example, the amino acid sequence of RuvC domain or PAM recognition domain).

[0177] In another embodiment, the Cas12f1 variant may be a variant having at least one amino acid residue added to the N-terminus and/or C-terminus of the amino acid sequence of wild-type Cas12f1 (for example, UnCas12f1 or CWCas12f1) or a variant protein thereof. The present inventors have confirmed that among the variants having amino acids added to the N-terminus and/or C-terminus of the wild-type Cas12f1 protein, there are variants having a function equivalent to the wild-type Cas12f1. For this purpose, reference may be made to Korean Patent Application No. 10-2021-0181875, the entire disclosure of which should be deemed to be incorporated herein. Preferably, the Cas12f1 variant may be such that it has 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 amino acids added to the N-terminus and/or C-terminus of wild-type Cas12f1 or a variant protein thereof. In an embodiment, Cas12f1 (for example, UnCas12f1 or CWCas12f1) or a variant protein thereof may comprise an amino acid sequence having the amino acid sequence of SEQ ID NO: 1 in which 1 to 28 amino acids at the N-terminus are removed or substituted. For example, the Cas12f1 variant may comprise or consist of TnpB-v1 protein (SEQ ID NO: 2), which further comprises 26 amino acids derived from the N-terminus of CasX at the N-terminus of the UnCas12f1 protein, TnpB-v2 protein (SEQ ID NO: 3), which further comprises 28 random amino acid sequences at the N-terminus of the UnCas12f1 protein, or TnpB-v3 protein (SEQ ID NO: 4), which further comprises 26 random amino acid sequences at the N-terminus of the UnCas12f1 protein (see FIG. 4).

[0178] In an embodiment, the Cas12f1 variant may be such that it is engineered to recognize a PAM sequence other than 5'-TTTA-3' or 5'-TTTG-3'. In an embodiment, the Cas12f1 variant may comprise substitution of at least one amino acid residue selected from the group consisting of amino acids at position 170 (serine), position 174 (tyrosine), position 184 (alanine), position 188 (serine), position 191

(arginine), position 225 (glutamine), position 230 (tyrosine), position 271 (valine), and position 272 (glutamine) with respect to the wild-type sequence of CWCas12f1 (TnpB) (for example, amino acid sequence of SEQ ID NO: 1). Preferably, the Cas12f1 variant may comprise substitution of at least one amino acid residue selected from the group consisting of amino acids at position 170 (serine, S), position 188 (serine, S), position 191 (arginine, R), position 225 (glutamine, Q), and position 272 (glutamine, Q). More preferably, the Cas12f1 variant may comprise one or more selected from the following substitutions with respect to the wild-type sequence (for example, the amino acid sequence of SEQ ID NO: 1): S170T, S188Q, S188H, S188K, R191K, Q225T, Q225F, and Q272K (wherein T is threonine, Q is glutamine, H is histidine, K is lysine, and F is phenylalanine). In addition, the Cas12f1 variant may comprise an amino acid sequence selected from the group consisting of SEQ ID NOS: 392 to 399. These Cas12 variants may further recognize 5'-TNTN-3', 5'-TTTN-3', 5'-TGTA-3', 5'-TCTG-3', 5'-TG TG-3', or 5'-TTTC-3' as a PAM sequence, wherein N is A, T, C, or G.

[0179] In another embodiment, the Cas12f1 variant may be a fusion protein. The fusion protein may comprise two or more heterologous polypeptide domains, wherein one polypeptide domain comprises a Cas12f1 protein or a variant protein thereof, and the other domain comprises a (poly)peptide having another function or activity. For example, the (poly)peptide having another function or activity may have methylase activity, demethylase activity, transcription activation activity, transcription repression activity, transcription release factor activity, histone modification activity, RNA cleavage activity, or nucleic acid binding activity. In addition, the (poly)peptide, which has a different function or activity, may be a tag or reporter protein for separation and/or purification. For example, the tag or reporter protein includes, but is not limited to, a tag protein such as a histidine (His) tag, a V5 tag, a FLAG tag, an influenza hemagglutinin (HA) tag, a Myc tag, a VSV-G tag, and a thioredoxin (Trx) tag; a fluorescent protein such as green fluorescent protein (GFP), yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), blue fluorescent protein (BFP), HcRED, and DsRed; and a reporter protein (enzyme) such as glutathione-S-transferase (GST), horseradish peroxidase (HRP), chloramphenicol acetyltransferase (CAT), β -galactosidase, β -glucuronidase, and luciferase.

[0180] In addition, the (poly)peptide having another function or activity may be, but is not limited to, a reverse transcriptase, a deaminase or another proteolytic enzyme.

[0181] In another embodiment, the Cas12f1 protein may comprise an amino acid sequence having at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, or at least 95% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NOS: 1 to 5.

[0182] In an embodiment, the Cas12f1 protein may comprise one selected from the following sequences: (i) the amino acid sequence of SEQ ID NO: 5; (ii) the amino acid sequence of SEQ ID NO: 1; (iii) an amino acid sequence having the amino acid sequence of SEQ ID NO: 1 in which 1 to 28 amino acids at the N-terminus have been removed or substituted; or (iv) an amino acid sequence having the amino acid sequence of SEQ ID NO: 1 in which 1 to 600 amino acids have been added to the N-terminus or C-terminus.

[0183] In another embodiment, the Cas12f1 variant protein may be a protein comprising or consisting of one

selected from the amino acid sequences having the amino acid sequence of SEQ ID NO: 1 in which 1 to 600 amino acids have been added to the N-terminus or C-terminus. Here, there is no limitation on the added sequence of 1 to 600 amino acids. For example, the added 1 to 600 amino acids may be the amino acid sequence of SEQ ID NO: 390 or SEQ ID NO: 391. An NLS or NES sequence may further be included between the added sequence and the Cas12f1 variant protein.

[0184] In an embodiment, since the target nucleic acid editing system cleaves a nucleic acid at a target site of a target nucleic acid or target gene, the target site may be located in the nucleus of a cell. The Cas12f1 protein may comprise one or more nuclear localization signal (NLS) sequences that localize the molecule into the nucleus. For example, one or more nuclear localization signal sequences may have a sufficient amount or activity to induce the Cas12f1 protein to be targeted to the nucleus of a eukaryotic cell (for example, a mammalian cell) in a detectable amount. For example, differences in the strength of activity may result from the number of NLSs included in the Cas12f1 protein, the type of specific NLS(s) used, or a combination of these factors. For example, the NLS may be, but is not limited to, an NLS sequence derived from NLS of SV40 virus large T-antigen, NLS from nucleoplasmin, c-myc NLS; hRNPA1 M9 NLS, the sequence of IBB domain from importin- α , the sequence of myoma T protein, the sequence of human p53, the sequence of mouse c-abl IV, the sequence of influenza virus NS1, the sequence of hepatitis virus delta antigen, the sequence of mouse Mx1 protein, the sequence of human poly(ADP-ribose) polymerase, or the sequence of steroid hormone receptor (human) glucocorticoid.

[0185] In another embodiment, the Cas12f1 protein may comprise a nuclear export sequence (NES).

[0186] In another embodiment, the Cas12f1 protein may be a fusion of various enzymes that may be involved in a gene expression process within cells. Here, the Cas12f1 protein to which the enzymes are fused may cause various quantitative and/or qualitative changes in gene expression in cells. For example, the various enzymes to be additionally bound may be DNMT, TET, KRAB, DHAC, LSD, p300, Moloney Murine Leukemia Virus (M-MLV) reverse transcriptase, or variants thereof. The Cas12f1 protein to which the reverse transcriptase is fused may also function as a prime editor.

[0187] In an embodiment, there is provided a nucleic acid encoding the Cas12f1 protein. The nucleic acid encoding

Cas12f1 (including a variant thereof) may be codon optimized for a subject (for example, a human) to which the Cas12f1 protein is to be introduced. For example, the human codon optimized nucleotide sequence encoding Cas12f1 may be, for example, at least one selected from SEQ ID NOS: 6 to 10.

2. Guide RNA

[0188] As disclosed herein, the CRISPR/Cas12f1 system comprises at least one guide RNA or a nucleic acid construct encoding the guide RNA. Cas12f1 guide RNA provides targeting for CRISPR/Cas12f1. The guide RNA of the CRISPR/Cas12f1 system may be a Cas12f1 guide RNA found in nature or an engineered Cas12f1 guide RNA. The Cas12f1 guide RNA found in nature or engineered Cas12f1 guide RNA comprises a scaffold region and a spacer region. The scaffold region of the Cas12f1 guide RNA is a region that comprises parts of tracrRNA (trans-activating CRISPR RNA) and crRNA (CRISPR RNA) and functions to interact with the Cas12f1 protein. The spacer region of the Cas12f1 guide RNA comprises a guide sequence.

[0189] The wild-type gRNA includes two structures in which apart of tracrRNA (tracrRNA anti-repeat) and a part of crRNA repeat (crRNA repeat) are complementarily bound to form a duplex, which are conveniently referred to as R:AR1 and R:AR2. The wild-type guide RNA may comprise (i) at least one stem region, (ii) a tracrRNA-crRNA complementarity region, and optionally (iii) a region comprising three or more consecutive uracil (U) residues. Specifically, the wild-type guide RNA may sequentially comprise, from the 5'-end, a first stem region, a second stem region, a third stem region, a fourth stem region, and a fifth stem region (tracrRNA-crRNA complementarity region). For example, referring to FIG. 2, the scaffold region of the wild-type guide RNA comprises five stem regions, that is, a first stem region (stem 1), a second stem region (stem 2), a third stem region (stem 3), a fourth stem region (stem 4), and a fifth stem region (stem 5 (R:AR2)), from the 5'-end. The region comprising stem 5 (R:AR2) is also referred to as a tracrRNA-crRNA complementarity region.

[0190] More specifically, the wild-type gRNA may comprise a wild-type tracrRNA having the nucleotide sequence of SEQ ID NO: 11, or a wild-type crRNA having the nucleotide sequence of SEQ ID NO: 12. In addition, the wild-type gRNA may be fused in the form of a single guide RNA to become a single guide RNA (sgRNA) having the nucleotide sequence of SEQ ID NO: 13.

TABLE 1

Name	Nucleotide sequence	SEQ ID NO
Wild-type tracrRNA	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAG UAACCCUCGAAACAAAUUCAUUUUUCUCUCCAAUUCUG CACAA	11
Wild-type crRNA	GUUGCAGAACCCGAAUAGACGAAUGAAGGAUUGCAAC	12
Canonical sgRNA	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAG UAACCCUCGAAACAAAUUCAUUUUUCUCUCCAAUUCUG	13

TABLE 1-continued

Name	Nucleotide sequence	SEQ ID NO
	CACAAgaaaGUUGCAGAACCCGAAUAGacgaaUGAAGGA AUGCAACNNNNNNNNNNNNNNNNNNNNNNNN	

2.1. Guide Sequence

[0191] The guide RNA may comprise at least one guide sequence that hybridizes with a target sequence in a target gene. Since a protospacer sequence complementary to the target sequence is located at the 5'- or 3'-end of the PAM sequence recognized by the Cas12f1 protein, the guide sequence may be designed using the protospacer sequence. A guide sequence that binds complementarily to a target sequence may be designed as a nucleotide sequence having the same nucleotide sequence as the protospacer sequence. When the protospacer sequence is a DNA sequence, the guide sequence may be such that T in the protospacer sequence is replaced with U.

[0192] In an embodiment, the guide sequence may be hybridizable with or complementary to a target sequence of contiguous 15 to 30 bp in length, wherein the target sequence is located in a region 5000 bp, 4000 bp, 3000 bp, 2000 bp, or 1000 bp upstream of a nucleic acid segment to be deleted in a target gene, or a region 5000 bp, 4000 bp, 3000 bp, 2000 bp, or 1000 bp downstream of the nucleic acid segment, and is adjacent to the 5'-end or 3'-end of a PAM sequence recognized by the Cas12f1 protein.

[0193] In an embodiment, the guide sequence of the guide RNA may bind complementarily to the target sequence. Complementary binding between the guide sequence and the target sequence may include at least one mismatch bond. For example, complementary binding between the guide sequence and the target sequence may include 0 to 5 mismatches. The guide sequence may be a sequence having at least 70% sequence complementarity to the target sequence. Unless stated otherwise, "complementary" may mean including 0 to 5 mismatches or having at least 70% complementarity, and should be interpreted appropriately depending on the context. When the target sequence is DNA, for an adenosine (A) present in the target sequence, the guide sequence may comprise a uridine (U) residue that can form a complementary bond with A.

[0194] In an embodiment, the target sequence may be a sequence of 15 to 40 nucleotides. For example, the target sequence may be a sequence of 15 to 20, 15 to 25, 15 to 30, 15 to 35, or 15 to 40 nucleotides. The target sequence may be a sequence of 20 to 25, 20 to 30, 20 to 35, or 20 to 40 nucleotides. In addition, the target sequence may be a sequence of 25 to 30, 25 to 35, or 25 to 40 nucleotides. In addition, the target sequence may be a sequence of 30 to 35 or 30 to 40 nucleotides. In addition, the target sequence may be a sequence of 35 to 40 nucleotides. In addition, the target sequence may be a sequence of 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40 nucleotides.

[0195] In an embodiment, the guide sequence may be a sequence that is at least 70% to 75%, at least 70% to 80%, at least 70% to 85%, at least 70% to 90%, at least 70% to 95%, at least 70% to 100%, at least 75% to 80%, at least 75% to 85%, at least 75% to 90%, at least 75% to 95%, or

at least 75% to 100% complementary to the target sequence. Specifically, the guide sequence may be a sequence that is at least 80% to 85%, at least 80% to 90%, at least 80% to 95%, at least 80% to 100%, at least 85% to 90%, at least 85% to 95% or at least 85% to 100% complementary to the target sequence. More specifically, the guide sequence may be a sequence that is at least 90% to 95%, at least 90% to 100%, or at least 95% to 100% complementary to the target sequence. More specifically, the guide sequence may be a sequence that is at least 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% complementary to the target sequence.

[0196] In another embodiment, the guide sequence may be identical to or similar to the protospacer sequence. The guide sequence may have at least 70% sequence identity to the protospacer sequence. For thymine (T) present in the protospacer sequence, the guide sequence may comprise uracil (U) instead of thymine (T).

[0197] In an embodiment, the guide sequence may have at least 70% to 75%, at least 70% to 80%, at least 70% to 85%, at least 70% to 90%, at least 70% to 95%, at least 70% to 100%, at least 75% to 80%, at least 75% to 85%, at least 75% to 90%, at least 75% to 95%, or at least 75% to 100% sequence identity or similarity to the protospacer sequence. Specifically, the guide sequence may have at least 80% to 85%, at least 80% to 90%, at least 80% to 95%, at least 80% to 100%, at least 85% to 90%, at least 85% to 95%, or at least 85% to 100% sequence identity or similarity to the protospacer sequence. More specifically, the guide sequence may have at least 90% to 95%, at least 90% to 100%, or at least 95% to 100% identity or similarity to the protospacer sequence. Even more specifically, the guide sequence may have at least 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100% identity or similarity to the protospacer sequence.

[0198] In an embodiment, the guide sequence may be a sequence that is hybridizable with or complementary to a nucleotide sequence present in an upstream region and/or a downstream region of a nucleic acid segment to be deleted. For example, a guide sequence of a first guide RNA may target a nucleotide sequence present in an upstream region of a nucleic acid segment to be deleted, and a guide sequence of a second guide RNA may target a nucleotide sequence present in a downstream region of the nucleic acid segment to be deleted.

[0199] In an embodiment, the guide sequence may be present at the 5'-end of the crRNA. In another embodiment, a U-rich tail may be added to the 5'-end of the guide sequence. The U-rich tail will be described later.

2.2. Engineered Guide RNA

[0200] Since no naturally occurring gRNA has been found for CWCas12f1 according to an embodiment of the present disclosure, it was desired to produce an optimal gRNA

exhibiting highly efficient targeting and editing activity not only for the engineered UnCas12f1 protein but also for the engineered CWCas12f1 protein. From this perspective, the gRNA may be a wild-type gRNA found in nature for wild-type UnCas12f1, which is similar in size to the CWCas12f1 protein. That is, in the present disclosure, the “wild-type” gRNA for the engineered Cas12f1 protein was used to mean “basic” or “canonical” gRNA.

[0201] In an embodiment, the guide RNA for the engineered Cas12f1 protein is characterized in that it is an engineered guide RNA in which a new configuration is added to a wild-type guide RNA found in nature, or the existing structure is removed and/or substituted, or whose structure is partially modified.

[0202] In an embodiment, the engineered gRNA is an engineered gRNA comprising a sequence having the wild-type gRNA sequence in which at least one nucleotide has been substituted, deleted, inserted, or added, wherein the sequence excluding the guide sequence has at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to the wild-type Cas12f1 gRNA. In the context of RNA, nucleic acids, or polypeptides, the term “sequence identity” refers to a value determined by comparing two sequences that are optimally aligned over a comparison window, in which a sequence portion of RNA, nucleic acid, and the like within the comparison window may comprise insertions or deletions (that is, gaps) relative to the reference sequence to achieve optimal alignment.

[0203] Hereinafter, the structures of wild-type and engineered gRNAs and modifications thereof will be described in detail for each of the five modification sites. The modification site is abbreviated as “MS” throughout this specification, and the numbers following “modification site” or “MS” are sequentially assigned depending on engineering flow of each modification site according to an embodiment. However, this does not mean that engineering (modification) at a modification site with a later number necessarily includes engineering (modification) at a modification site with an earlier number. FIG. 2 illustrates modification sites MS1 to MS5 included in the engineered guide RNA according to an embodiment of the present disclosure on the wild-type guide RNA sequence.

[0204] The modifications applied to the engineered guide RNA (gRNA) of the present disclosure are ultimately intended to achieve high gene editing efficiency while deriving a gRNA that is shorter in length. That is, the modifications disclosed in the present disclosure are intended to produce an engineered gRNA of a shorter length having equal or improved recognition/cleavage efficiency for a target nucleic acid compared to the wild-type gRNA of a longer length, thereby allowing more space to be allocated to other components (for example, additional guide RNAs, shRNAs for inhibiting specific gene expression) for various purposes or uses within the packaging limit (about 4.7 kb) of a delivery vehicle such as adeno-associated virus (AAV). This provides a highly efficient gene editing effect that could not be achieved with the existing CRISPR/Cas system.


[0205] Therefore, the engineered gRNA provided in the present disclosure basically comprises a sequence having the wild-type Cas12f1 gRNA sequence in which one or more nucleotides are substituted, deleted, inserted, or added. Here, for the engineered gRNA, a portion thereof excluding the guide sequence may have sequence identity of 50% or more,

60% or more, 70% or more, 80% or more, 85% or more, 90% or more, or 95% or more to the wild-type Cas12f1 gRNA.

[0206] In an embodiment, compared to a wild-type Cas12f1 gRNA comprising (i) at least one stem region, (ii) a tracrRNA-crRNA complementarity region and optionally (iii) a region comprising three or more consecutive uracil (U) residues, the engineered gRNA of the present disclosure may comprise at least one modification selected from the group consisting of (a) deletion of at least a part of the at least one stem region; (b) deletion of at least a part of the tracrRNA-crRNA complementarity region; (c) replacement of one or more of uracil (U) residues when three or more consecutive uracil (U) residues are present; and (d) addition of one or more uridine residues to the 3'-end of the crRNA sequence.

[0207] In another embodiment, the engineered guide RNA may comprise at least one modification selected from the group consisting of (a1) deletion of at least a part of the first stem region; (a2) deletion of at least a part of the second stem region; (b) deletion of at least a part of the tracrRNA-crRNA complementarity region; (c) replacement of one or more U residues with A, G or C in three or more consecutive uracil (U) residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region; and (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence, wherein a sequence of the U-rich tail is represented by 5'-(U_mV)_nU_o-3', wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5.

[0208] In another embodiment, the engineered guide RNA may be represented by Formula (I).



[0209] In Formula (I),

[0210] X^a, X^{b1}, X^{b2}, X^{c1}, and X^{c2} each independently consist of 0 to 35 (poly)nucleotides,

[0211] X^g is a guide sequence,

[0212] Lk is a polynucleotide linker of 2 to 20 nucleotides or is absent, and

[0213] (U_mV)_nU_o is present as a U-rich tail or absent, and when (U_mV)_nU_o is present, U is uridine, V is each independently A, C, or G, m and o are each independently an integer between 1 and 20, and n is an integer between 0 and 5.

[0214] [In Formula (I), the black solid line refers to a chemical bond (for example, phosphodiester bond) between nucleotides or specific molecules, and the gray thick line refers to a complementary bond between nucleotides].

[0215] In an embodiment, X^a may be absent or a (poly)nucleotide having a stem-loop conformation.

[0216] In an embodiment, X^{b1} and X^{b2} may be (poly)nucleotides capable of complementary binding to each other.

[0217] In an embodiment, X^{c1} and X^{c2} may be (poly) nucleotides capable of complementary binding to each other.

[0218] In another embodiment, the engineered guide RNA may have at least 70%, at least 80%, at least 85%, at least 90%, at least 95%, or at least 98% sequence identity to the sequence represented by Formula (I). Here, the sequence identity with Formula (I) is based on the sequence excluding the regions indicated by the symbols.

[0219] When referring to the scaffold region of the wild-type guide RNA, the first stem region of the scaffold sequence may be a region corresponding to X^a in Formula (I). The second stem region of the scaffold sequence may be a region corresponding to X^{b1} and X^{b2} in Formula (I). The third stem region of the scaffold sequence may be a region corresponding to the sequence 5'-GGCUGCUUG-CAUCAGCC-3' in Formula (I). The fourth stem region of the scaffold sequence may be a region corresponding to the sequence 5'-UCGAGAAGUGCUUUCUUCGGAA-AGUAACCCUCGA-3' in Formula (I). In addition, the tracrRNA-crRNA complementarity region (the fifth stem region) of the scaffold sequence may be a region corresponding to X^{c1} and X^{c2} in Formula (I).

[0220] Hereinafter, modifications at respective modification sites in the engineered gRNA will be described in detail.

(1) Modification at Modification Site 1 (MS1)

[0221] This section describes a modification at MS1. In an embodiment, wild-type tracrRNA (for example, SEQ ID NO: 11), which may be a guide RNA (gRNA) existing in nature, may have a sequence comprising five consecutive uracil (U) residues therein. This poses a problem in that, in a case of attempting to express the wild-type tracrRNA in a cell using a vector or the like, such a sequence acts as a transcription termination signal under certain conditions, thereby causing unintended early termination of transcription. That is, in a case where the sequence containing five consecutive U residues acts as a transcription termination signal, normal or complete expression of the tracrRNA is inhibited, and formation of normal or complete gRNA is also inhibited, which consequently decreases efficiency of cleavage or homology-directed repair of the target nucleic acid editing system of the present disclosure.

[0222] Therefore, in order to solve the above-mentioned problem, the engineered gRNA may be such that at least one uracil (U) of three or more, four or more, or five or more consecutive U residues, preferably four or five U residues, which are contained in the wild-type tracrRNA (for example, SEQ ID NO: 11), is artificially modified into another nucleotide such as A, C, T, or G.

[0223] In an embodiment, the engineered gRNA is provided which comprises a modification in which at least one of three or more consecutive U residues is substituted with a different type of nucleotide in a region containing three or more consecutive U residues, referred to as MS1. For example, the three or more consecutive U residues may be present in the tracrRNA-crRNA complementarity region of the tracrRNA, wherein a modification may be made by substituting at least one of the three or more U residues with A, G, or C such that no sequence with three or more consecutive U residues exists.

[0224] Here, it is preferable that the sequence within the tracrRNA-crRNA complementarity region of crRNA, which corresponds to the sequence to be modified, is also modified together. In an embodiment, when there is the sequence

5'-ACGAA-3' within the tracrRNA-crRNA complementarity region of crRNA, which forms a partial complementary bond with the sequence 5'-UUUUU-3' within the tracrRNA-crRNA complementarity region of tracrRNA, this sequence may be replaced with 5'-NGNNN-3'. Here, N is each independently A, C, G, or U.

[0225] In another embodiment, MS1 may be present in the polynucleotides indicated by X^{c1} and X^{c2} in Formula (I).

[0226] In an embodiment, in the engineered gRNA of Formula (I), when three or more consecutive uracil (U) residues are present in the X^{c1} sequence, the sequence may comprise a modification in which one or more of U residues thereof are replaced with A, G, or C. For example, when the sequence 5'-UUUUU-3' is present in the X^{c1} sequence, the sequence may be replaced with 5'-NNNCN-3'. Here, N is each independently A, C, G, or U. As a more specific example, the sequence 5'-UUUUU-3' in the X^{c1} sequence may be replaced with any one nucleotide sequence selected from the group consisting of the following sequences; however, the replacing sequence is not limited to the following sequences as long as it prevents appearance of a sequence containing three or more consecutive U residues: 5'-UUUCU-3', 5'-GUUCU-3', 5'-UCUCU-3', 5'-UUGCU-3', 5'-UUUCC-3', 5'-GCUCU-3', 5'-GUUCC-3', 5'-UCGCU-3', 5'-UCUCC-3', 5'-UUGCC-3', 5'-GCGCU-3', 5'-GCUCC-3', 5'-GUGCC-3', 5'-UCGCC-3', 5'-GCGCC-3', and 5'-GUGCU-3'.

[0227] In another embodiment, in the engineered gRNA of Formula (I), the X^{c2} sequence comprises a region in which at least a part of the sequence forms a complementary bond with the X^{c1} sequence (also referred to as a tracrRNA-crRNA complementarity region), wherein a corresponding sequence in the X^{c2} sequence, which forms at least one complementary bond with 3 or more consecutive U residues present in the X^{c1} sequence, may also be modified. For example, when the sequence 5'-ACGAA-3' is present in the X^{c2} sequence of Formula (I), the sequence may be replaced with 5'-NGNNN-3'. Here, N is each independently A, C, G, or U. As a more specific example, the sequence 5'-ACGAA-3' in the X^{c1} sequence of Formula (I) may be replaced with any one nucleotide sequence selected from the group consisting of the following sequences; however, the replacing sequence is not limited to the following sequences: 5'-AGGAA-3', 5'-AGCAA-3', 5'-AGAAA-3', 5'-AGCAU-3', 5'-AGCAG-3', 5'-AGCAC-3', 5'-AGCUA-3', 5'-AGCGA-3', 5'-AGCCA-3', 5'-UGCAA-3', 5'-UGCUA-3', 5'-UGCGA-3', 5'-UGCCA-3', 5'-GGCAA-3', 5'-GGCUA-3', 5'-GGCGA-3', 5'-GGCCA-3', 5'-CGCAA-3', 5'-CGCUA-3', 5'-CGCGA-3', and 5'-CGCCA-3'.

[0228] In another embodiment, when a sequence containing 3 or more consecutive U residues in the X^{c1} sequence of Formula (I) is modified into another sequence, it is preferred that the corresponding nucleotides in the X^{c2} sequence (that is, at least some of which forms a complementary bond therewith) are modified so that they can form a complementary bond with the modified nucleotides. For example, when the sequence 5'-UUUUU-3' in the X^{c1} sequence is modified into 5'-GUGCU-3', it is preferred that the sequence 5'-ACGAA-3' in the X^{c2} sequence is modified into 5'-AGCAA-3'; however, complementary bonding is not necessarily required.

(2) Modification at Modification Site 2 (MS2)

[0229] This section describes a modification at MS2. In an embodiment, the engineered guide RNA (gRNA) may be obtained by adding a new configuration to the gRNA found in nature, and may be such that one or more uridine residues are added to the 3'-end of the crRNA sequence. Here, the 3'-end of the crRNA sequence may be the 3'-end of the guide sequence (spacer). In the present disclosure, the one or more uridine residues added to the 3'-end are also referred to herein as a U-rich tail. The engineered gRNA comprising one or more uridine residues or a U-rich tail added to the 3'-end serves to increase nucleic acid cleavage or indel efficiency of the hypercompact CRISPR/Cas12 system for a target gene or target nucleic acid.

[0230] The term "U-rich tail" as used herein may refer not only to an RNA sequence itself that is rich in uridine (U), but also a DNA sequence encoding the same, and this may be appropriately interpreted depending on the context. The present inventors have experimentally elucidated the structure and effects of the U-rich tail sequence in detail, and the U-rich tail sequence will be described in more detail below with specific embodiments.

[0231] In an embodiment, the U-rich tail sequence may be represented by U_x , wherein x may be 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20. For example, x may be an integer within a range of two numerical values selected from the numerical values listed above. For example, x may be an integer between 1 and 6. As another example, x may be an integer between 1 and 20. In an embodiment, x may be an integer of 20 or higher.

[0232] In another embodiment, the U-rich tail sequence is represented by $5'-(U_mV)_nU_o-3'$, wherein V may be each independently A, C or G, m and o may be integers from 1 to 20, and n may be an integer from 0 to 5. As an example, n may be 0, 1, or 2. As an example, m and o may be each independently 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

[0233] In another embodiment, the engineered gRNA may be gRNA consisting of a sequence represented by Formula (I) or having at least 80%, at least 85%, at least 90%, or at least 95% sequence identity thereto. Here, MS2 is a region corresponding to $(U_mV)_nU_o$ in Formula (I), wherein U is uridine, and V , m , o , and n are as defined above.

[0234] Preferably, in the engineered gRNA represented by Formula (I), $(U_mV)_nU_o$ may be a U-rich tail in which (i) n is 0, o is an integer between 1 and 6, or (ii) V is A or G, m and o are each independently an integer between 3 and 6, and n is an integer between 1 and 3. In a specific example, $(U_mV)_nU_o$ in Formula (I) may be a U-rich tail consisting of any one sequence selected from the group consisting of $5'-U-3'$, $5'-UU-3'$, $5'-UUU-3'$, $5'-UUUU-3'$, $5'-UUUUU-3'$, $5'-UUUUUU-3'$, $5'-UUUUUUU-3'$, $5'-UUUUUUUU-3'$, $5'-UUUUUUUUU-3'$, $5'-UUUUUUUUUU-3'$, $5'-UUUUUUUUUUU-3'$, $5'-UUUUUUUUUUUU-3'$, and $5'-UUUUUUUUUUUUU-3'$, wherein R is A or G.

[0235] In yet another embodiment, the U-rich tail sequence may comprise a modified uridine repeat sequence that contains a non-uridine ribonucleoside (A, C, or G) for every 1 to 5 repetitions of uridine. The modified uridine repeat sequence is particularly useful in a case of designing a vector that expresses an engineered crRNA. In an embodiment, the U-rich tail sequence may comprise a sequence in which UV, UUV, UUUU, UUUUV, and/or UUUUUUV are repeated one or more times. Here, V is one of A, C or G.

[0236] In addition, the U-rich tail sequence may be a combination of the sequence represented by U_x and the sequence represented by $(UaV)_n$. In an embodiment, the U-rich tail sequence may be represented by $(U)n1-V1-(U)n2-V2-U_x$. Here, $V1$ and $V2$ are each one of adenine (A), cytosine (C), and guanine (G). Here, $n1$ and $n2$ may each be an integer between 1 and 4. Here, x may be an integer between 1 and 20. In addition, the U-rich tail sequence may have a length of 1 nt, 2 nts, 3 nts, 4 nts, 5 nts, 6 nts, 7 nts, 8 nts, 9 nts, 10 nts, 11 nts, 12 nts, 13 nts, 14 nts, 15 nts, 16 nts, 17 nts, 18 nts, 19 nts, or 20 nts. In an embodiment, the U-rich tail sequence may have a length of 20 nts or longer.

[0237] In another embodiment, when the engineered gRNA is expressed in a cell, the U-rich tail may exist in a plurality of forms due to premature termination of transcription. For example, according to an embodiment, when a gRNA intended to contain a U-rich tail of the sequence $5'-UUUUUAUUUUUUU-3'$ is transcribed in a cell, four or more or five or more T residues may act as a termination sequence, and thus gRNAs containing a U-rich tail such as $5'-UUUUUAUUUUU-3'$, $5'-UUUUUAUUUUU-3'$, or $5'-UUUUUAUUUUUU-3'$ may be produced simultaneously. Therefore, in the present disclosure, a U-rich tail containing four or more U residues may be understood to also include a U-rich tail sequence having a shorter length than the intended length.

[0238] In yet another embodiment, the U-rich tail sequence may comprise additional nucleotides other than uridine, depending on the environment where the CRISPR/Cas12 system is actually used and expression environment, such as the internal environment of a eukaryotic cell or a prokaryotic cell.

(3) Modification at Modification Site 3 (MS3)

[0239] This section describes a modification at MS3. As described above, MS3 refers to a region (which may be referred to as the first stem region) that comprises at least a part of the nucleotides forming a stem structure within a complex of the gRNA with an effector protein. The MS3 may comprise a region that does not interact with the effector protein when the gRNA and effector protein form a complex. The modification at MS3 involves removal of at least a part of the first stem region near the 5'-end of tracrRNA.

[0240] In an embodiment, the engineered gRNA comprises a modification in which at least a part of the first stem region (for example, the sequence of SEQ ID NO: 14) is deleted.

[0241] In another embodiment, the engineered gRNA comprises a modification in which at least a part of the first stem region on tracrRNA is deleted, wherein at least a part of the first stem region to be deleted may consist of 1 to 20 nucleotides. Specifically, at least a part of the first stem region may consist of 2 to 20, 3 to 20, 4 to 20, 5 to 20, 6 to 20, 7 to 20, 8 to 20, 9 to 20, 10 to 20, 11 to 20, 12 to 20, 13 to 20, 14 to 20, 15 to 20, 16 to 20, 17 to 20, 18 to 20, 19, or 20 nucleotides.

[0242] In yet another embodiment, the MS3 or the first stem region is a portion corresponding to the polynucleotide indicated by X^a of Formula (I), wherein due to a modification in which at least a part of the first stem region is deleted, X^a may consist of 0 to 35 (poly)nucleotides, preferably 0 to 20, 0 to 19, 0 to 18, 0 to 17, 0 to 16, 0 to 15, 0 to 14, 0 to

13, 0 to 12, 0 to 11, 0 to 10, 0 to 9, 0 to 8, 0 to 7, 0 to 6, 0 to 5, 0 to 4, 0 to 3, 0 to 2, 1 or 0 (poly)nucleotides.

[0243] In an embodiment, in the engineered gRNA of Formula (I), X^a may comprise the nucleotide sequence of SEQ ID NO: 14 or may comprise a nucleotide sequence having at least a part thereof, preferably a nucleotide sequence having the sequence of SEQ ID NO: 14 from which 1 to 20 nucleotides are deleted. For example, the nucleotide deletion may involve random deletion of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides from the sequence of SEQ ID NO: 14. As a preferred example, the nucleotide deletion may involve sequential deletion of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 nucleotides from the 5'-end of the sequence of SEQ ID NO: 14. More specifically, X^a may comprise or consist of CUUCACUGAUAAAGUGGAGA-3' (SEQ ID NO: 14), 5'-UUCACUGAUAAAGUGGAGA-3' (SEQ ID NO: 15), 5'-UCACUGAUAAAGUGGAGA-3' (SEQ ID NO: 16), 5'-CACUGAUAAAGUGGAGA-3' (SEQ ID NO: 17), 5'-ACUGAUAAAGUGGAGA-3' (SEQ ID NO: 18), 5'-CUGAUAAAGUGGAGA-3' (SEQ ID NO: 19), 5'-UGAUAAAGUGGAGA-3' (SEQ ID NO: 20), 5'-GAUAAAGUGGAGA-3' (SEQ ID NO: 21), 5'-AUAAAGUGGAGA-3' (SEQ ID NO: 22), 5'-UAAAGUGGAGA-3' (SEQ ID NO: 23), 5'-AAAGUGGAGA-3' (SEQ ID NO: 24), 5'-AAGUGGAGA-3', 5'-AGUGGAGA-3', 5'-GUGGAGA-3', 5'-UGGAGA-3', 5'-GGAGA-3', 5'-GAGA-3', 5'-AGA-3', 5'-GA-3', or 5'-A-3', or X^a may be absent.

(4) Modification at Modification Site 4 (MS4)

[0244] This section describes a modification at MS4. MS4 refers to a region spanning the 3'-end of tracrRNA and the 5'-end of crRNA, or, in a case of a single guide RNA form, a region where the sequence corresponding to tracrRNA and the sequence corresponding to crRNA form at least partial complementary bonding. MS4 may comprise at least a part of the sequence referred to as the tracrRNA-crRNA complementarity region (which may also be referred to as the fifth stem region). In the present disclosure, the tracrRNA-crRNA complementarity region may comprise both modification site 1 (MS1) and modification site 4 (MS4). The modification at MS4 comprises deletion of at least a part of the tracrRNA-crRNA complementarity region. The tracrRNA-crRNA complementarity region may comprise a part of tracrRNA and a part of crRNA. In this regard, the tracrRNA-crRNA complementarity region may comprise nucleotides such that partial nucleotides contained in tracrRNA can form complementary bonds with partial nucleotides contained in crRNA within a complex of gRNA with the nucleic acid degrading protein, and may comprise nucleotides adjacent thereto. The tracrRNA-crRNA complementarity region of tracrRNA may comprise a region that does not interact with the nucleic acid degrading protein within a complex of gRNA with the nucleic acid degrading protein.

[0245] In some embodiments, the engineered gRNA comprises deletion of at least a part of the tracrRNA-crRNA complementarity region in tracrRNA, deletion of at least a part of the tracrRNA-crRNA complementarity region in crRNA, or deletion of at least a part of the tracrRNA-crRNA complementarity region in both the tracrRNA and the crRNA.

[0246] In another embodiment, the engineered gRNA comprises a modification in which a part of the tracrRNA-crRNA complementarity region is deleted, wherein the part of the complementary region to be deleted may consist of 1 to 54 nucleotides.

[0247] In yet another embodiment, the engineered gRNA comprises a modification in which the entire tracrRNA-crRNA complementarity region is deleted, wherein the entire complementary region to be deleted may consist of 55 nucleotides.

[0248] In an embodiment, the tracrRNA-crRNA complementarity region may comprise the nucleotide sequence of SEQ ID NO: 39 and/or the nucleotide sequence of SEQ ID NO: 58.

[0249] In another embodiment, the tracrRNA-crRNA complementarity region may further comprise a linker sequence.

[0250] Specifically, at least a part of the tracrRNA-crRNA complementarity region may consist of 3 to 55, 5 to 55, 7 to 55, 9 to 55, 11 to 55, 13 to 55, 15 to 55, 17 to 55, 19 to 55, 21 to 55, 23 to 55, 25 to 55, 27 to 55, 29 to 55, 31 to 55, 33 to 55, 35 to 55, 37 to 55, 39 to 55, or 41 to 55 nucleotides, preferably 42 to 55, 43 to 55, 44 to 55, 45 to 55, 46 to 55, 47 to 55, 48 to 55, 49 to 55, 50 to 55, 51 to 55, 52 to 55, 53 to 55, or 54, or 55 nucleotides.

[0251] In yet another embodiment, MS4 or the tracrRNA-crRNA complementarity region is a region corresponding to the polynucleotide indicated by X^{c1} and X^{c2} in Formula (I), in which due to the modification where at least a part of the tracrRNA-crRNA complementarity region is deleted, X^{c1} and X^{c2} may each independently consist of 0 to 35 (poly) nucleotides.

[0252] Preferably, X^{c1} may consist of 0 to 28, 0 to 27, 0 to 26, 0 to 25, 0 to 24, 0 to 23, 0 to 22, 0 to 21, 0 to 20, 0 to 19, 0 to 18, 0 to 17, 0 to 16, 0 to 15, 0 to 14, 0 to 13, 0 to 12, 0 to 11, 0 to 10, 0 to 9, 0 to 8, 0 to 7, 0 to 6, 0 to 5, 0 to 4, 0 to 3, 0 to 2, 1, or 0 (poly)nucleotides. In addition, preferably, X^{c2} may consist of 0 to 27, 0 to 26, 0 to 25, 0 to 24, 0 to 23, 0 to 22, 0 to 21, 0 to 20, 0 to 19, 0 to 18, 0 to 17, 0 to 16, 0 to 15, 0 to 14, 0 to 13, 0 to 12, 0 to 11, 0 to 10, 0 to 9, 0 to 8, 0 to 7, 0 to 6, 0 to 5, 0 to 4, 0 to 3, 0 to 2, 1, or 0 (poly)nucleotides.

[0253] In an embodiment, in the engineered gRNA of Formula (I), X^{c1} may comprise the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having the sequence of SEQ ID NO: 39 from which 1 to 28 nucleotides are deleted. Preferably, the nucleotide deletion may involve sequential removal of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, or 28 nucleotides from the 5'-end of the sequence of SEQ ID NO: 39. More specifically, X^{c1} may comprise or consist of 5'-UUCAUUUUUCCUCUCCAAUUCUGCACAA-3' (SEQ ID NO: 39), 5'-UUCAUUUUUCCUCUCCAAUUCUGCAC-3' (SEQ ID NO: 40), 5'-UUCUUUUUCCUCUCCAAUUCUGCAC-3' (SEQ ID NO: 41), 5'-UUCAUUUUUCCUCUCCAAUUCUGCA-3' (SEQ ID NO: 42), 5'-UUCAUUUUUCCUCUCCAAUUCUGC-3' (SEQ ID NO: 43), 5'-UUCAUUUUUCCUCUCCAAUUCUG-3' (SEQ ID NO: 44), 5'-UUCAUUUUUCCUCUCCAAUUCU-3' (SEQ ID NO: 45), 5'-UUCUUUUUCCUCUCCAAUUC-3' (SEQ ID NO: 46), 5'-UUCAUUUUUCCUCUCCAAU-3' (SEQ ID NO: 47), 5'-UUCAUUUUUCCUCUCCAA-3' (SEQ ID NO: 48), 5'-UUCAUUUUUCCUCUCCAA-3' (SEQ ID NO: 49),

5'-UUCAUUUUUCCUCUCCA-3' (SEQ ID NO: 50), 5'-UUCAUUUUUCCUCUCC-3' (SEQ ID NO: 51), 5'-UUC AUUUUUUCCUCUC-3' (SEQ ID NO: 52), 5'-UUC AUUUUUUCCUCU-3' (SEQ ID NO: 53), 5'-UUC AUUUUUUCCUC-3' (SEQ ID NO: 54), 5'-UUCAUUUUUCCU-3' (SEQ ID NO: 55), 5'-UUC AUUUUUUCC-3' (SEQ ID NO: 56), 5'-UUCAUUUUUC-3' (SEQ ID NO: 57), 5'-UUCAUUUUU-3', 5'-UUC AUUUU-3', 5'-UUCAUUU-3', 5'-UUCAUU-3', 5'-UUCAU-3', 5'-UUC-3', 5'-UU-3', or 5'-U-3', or X^{c1} may be absent.

[0254] Here, in a case where there is a region containing 3, 4, or 5 or more uracil (U) residues in the sequence of X^{c1} from which some nucleotides have been removed, the modification at MS1 as described above may also apply. For details about MS1, see the section "(1) Modification at modification site 1 (MS1)."

[0255] In another embodiment, in the engineered gRNA of Formula (I), X^{c2} may comprise the nucleotide sequence of SEQ ID NO: 58 or a nucleotide sequence having the sequence of SEQ ID NO: 58 from which 1 to 27 nucleotides are deleted. Preferably, the nucleotide deletion may involve sequential removal of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or 27 nucleotides from the 5'-end of the sequence of SEQ ID NO: 58. More specifically, X^{c2} may comprise or consist of 5'-GUUGCAGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 58), 5'-UUGCAGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 59), 5'-UGCAGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 60), 5'-GCAGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 61), 5'-CAGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 62), 5'-AGAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 63), 5'-GAACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 64), 5'-AACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 65), 5'-ACCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 66), 5'-CCCGAAUAGACGAAUGAA-3' (SEQ ID NO: 67), 5'-CCGAAUAGACGAAUGAA-3' (SEQ ID NO: 68), 5'-CGAAUAGACGAAUGAA-3' (SEQ ID NO: 69), 5'-GAUAGACGAAUGAA-3' (SEQ ID NO: 70), 5'-AAUAGACGAAUGAA-3' (SEQ ID NO: 71), 5'-UAUAGACGAAUGAA-3' (SEQ ID NO: 72), 5'-UAGACGAAUGAA-3' (SEQ ID NO: 73), 5'-AGACGAAUGAA-3' (SEQ ID NO: 74), 5'-GACGAAUGAA-3' (SEQ ID NO: 75), 5'-ACGAAUGAA-3', 5'-CGAAUGAA-3', 5'-GAAUGAA-3', 5'-AAUGAA-3', 5'-AUGAA-3', 5'-UGAA-3', 5'-GAA-3', 5'-AA-3', or 5'-A-3', or X^{c2} may be absent.

[0256] Here, in a case where there is a sequence corresponding a sequence containing 3 or more, or 3, 4, or 5 or more uracil (U) residues in the sequence of X^{c2} from which some nucleotides have been removed, the modification at MS1 as described above may also apply. For details regarding MS1, see the section "(1) Modification at modification site 1 (MS1)."

[0257] In the engineered gRNA of Formula (I), the regions corresponding to X^{c1} and X^{c2} may each independently undergo the above-described modification. However, MS4 or the tracrRNA-crRNA complementarity region is a region where tracrRNA and crRNA form complementary bonds. For the tracrRNA and the crRNA to function as a dual guide RNA, it is preferable that the position and number of nucleotides to be deleted in each of X^{c1} and X^{c2} be identical

with or similar to each other. That is, in order to preserve complementarity, in a case of sequentially deleting nucleotides from the 3'-end of tracrRNA in MS4 (tracrRNA-crRNA complementarity region), it is preferable to sequentially delete nucleotides from the 5'-end of crRNA.

[0258] In some embodiments, the 3'-end of X^{c1} and the 5'-end of X^{c2} in the engineered gRNA of Formula (I) may be linked by a linker (Lk) so that the gRNA is modified into a single guide RNA (sgRNA) form. The linker Lk is a sequence that physically or chemically connects tracrRNA and crRNA, and may be a polynucleotide sequence having a length of 1 to 30 nucleotides. In an embodiment, Lk may be a sequence of 1 to 5, 5 to 10, 10 to 15, 2 to 20, 15 to 20, 20 to 25, or 25 to 30 nucleotides. For example, Lk may be, but is not limited to, 5'-GAAA-3'. As another example, Lk may be a linker comprising or consisting of 5'-UUAG-3', 5'-UGAAAA-3', 5'-UUGAAAA-3', 5'-UUCGAAAGAA-3' (SEQ ID NO: 76), 5'-UUCAGAAAGAA-3' (SEQ ID NO: 77), 5'-UUCAGAAAAAGAA-3' (SEQ ID NO: 78), or 5'-UUCAUUGAAAAAGAA-3' (SEQ ID NO: 79).

[0259] Meanwhile, while it is possible to use a linker (Lk) to make a single guide RNA (sgRNA), it is also possible to directly connect the 3'-end of tracrRNA, of which a partial sequence has been removed, to the 5'-end of crRNA of which a partial sequence has been removed.

[0260] In another embodiment, a case where X^{c1} and X^{c2} in the engineered gRNA of Formula (I) are linked by a linker may be indicated by 5'- X^{c1} -Lk- X^{c2} -3' as in Formula (I), and the 5'- X^{c1} -Lk- X^{c2} -3' may be any one nucleotide sequence selected from the group consisting of SEQ ID NOS: 80 to 86, but is not limited thereto.

(5) Modification at Modification Site 5 (MS5)

[0261] This section describes a modification at MS5. As described above, MS5 corresponds to a region located toward the 3'-end of tracrRNA, which is referred to as the second stem region. The second stem region may comprise nucleotides that form a stem structure within a complex of the guide RNA (gRNA) with nucleic acid editing protein, and may comprise nucleotides adjacent thereto. Here, the stem structure is distinct from the stem included in the above-described first stem region.

[0262] In an embodiment, the engineered gRNA comprises a modification in which at least a part of the second stem region is deleted.

[0263] In another embodiment, the engineered gRNA comprises deletion of at least a part of the second stem region, wherein at least a part of the second stem region to be deleted may consist of 1 to 27 nucleotides. Specifically, the at least a part of the second stem region may consist of 2 to 27, 3 to 27, 4 to 27, 5 to 27, 6 to 27, 7 to 27, 8 to 27, 9 to 27, 10 to 27, 11 to 27, 12 to 27, 13 to 27, 14 to 27, 15 to 27, 16 to 27, 17 to 27, 18 to 27, 19 to 27, 20 to 27, 21 to 27, 22 to 27, 23 to 27, 24 to 27, 25 to 27, 26, or 27 nucleotides.

[0264] In an embodiment, the second stem region may comprise or consist of the nucleotide sequence of SEQ ID NO: 25 and/or the nucleotide sequence of SEQ ID NO: 29.

[0265] In another embodiment, MS5 or the second stem region is a region comprising a (poly)nucleotide (comprising a loop of 5'-UUAG-3') that is adjacent to the polynucleotide indicated by X^{b1} and X^{b2} in Formula (I), in which due to the modification where at least the part of the second stem

region is deleted, X^{b1} and X^{b2} may each independently consist of 0 to 35 (poly)nucleotides.

[0266] Preferably, X^{b1} in Formula (I) may consist of 0 to 13, 0 to 12, 0 to 11, 0 to 10, 0 to 9, 0 to 8, 0 to 7, 0 to 6, 0 to 5, 0 to 4, 0 to 3, 0 to 2, 1, or 0 (poly)nucleotides. In addition, preferably, X^{b2} may consist of 0 to 14, 0 to 13, 0 to 12, 0 to 11, 0 to 10, 0 to 9, 0 to 8, 0 to 7, 0 to 6, 0 to 5, 0 to 4, 0 to 3, 0 to 2, 1, or 0 (poly)nucleotides.

[0267] In an embodiment, in the engineered gRNA of Formula (I), X^{b1} may comprise the nucleotide sequence of SEQ ID NO: 25 or a nucleotide sequence having the sequence of SEQ ID NO: 25 from which 1 to 13 nucleotides are deleted. Preferably, the nucleotide deletion may involve sequential removal of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, or 13 nucleotide from the 5'-end of the sequence of SEQ ID NO: 25. More specifically, X^{b1} may comprise or consist of 5'-CAAAGCUGUCCC-3' (SEQ ID NO: 25), 5'-CAAAGCUGUCC-3' (SEQ ID NO: 26), 5'-CAAAGCUGUC-3' (SEQ ID NO: 27), 5'-CAAAGCUGU-3' (SEQ ID NO: 28), 5'-CAAAGCUG-3', 5'-CAAAGCU-3', 5'-CAAAGC-3', 5'-CAAAAG-3', 5'-CAAAA-3', 5'-CAAA-3', 5'-CAA-3', 5'-CA-3', or 5'-C-3', or X^{b1} may be absent.

[0268] In another embodiment, in the engineered gRNA of Formula (I), X^{b2} may comprise the nucleotide sequence of SEQ ID NO: 29 or a nucleotide sequence having the sequence of SEQ ID NO: 29 from which 1 to 14 nucleotides are deleted. Preferably, the nucleotide deletion may involve sequential removal of at least 1, 2, 3, 5, 6, 7, 8, 9, 10, 11, 12, 13, or 14 nucleotides from the 5'-end of the sequence of SEQ ID NO: 29. More specifically, X^{b2} may comprise or consist of 5'-GGGAUUAGAACUUG-3' (SEQ ID NO: 29), 5'-GGAUUAGAACUUG-3' (SEQ ID NO: 30), 5'-GAUUA-GAACUUG-3' (SEQ ID NO: 31), 5'-AUUAGAACUUG-3' (SEQ ID NO: 32), 5'-UUAGAACUUG-3' (SEQ ID NO: 33), 5'-UAGAACUUG-3', 5'-AGAACUUG-3', 5'-GAACUUG-3', 5'-AACUUG-3', 5'-ACUUG-3', 5'-CUUG-3', 5'-UUG-3', 5'-UG-3', or 5'-G-3', or X^{b2} may be absent.

[0269] In the engineered gRNA of Formula (I), the regions corresponding to X^{b1} and X^{b2} may be each independently modified. However, for normal preservation of the stem-loop structure, it is preferable that the position and number of nucleotides to be deleted in each of X^{b1} and X^{b2} be identical with or similar to each other. For example, in a case of sequentially deleting nucleotides from the 5'-end direction in X^{b1} , it is preferable to sequentially delete nucleotides from the 3'-end direction in X^{b2} .

[0270] In another embodiment, a sequence of the loop portion connecting X^{b1} and X^{b2} in the engineered gRNA of Formula (I) is indicated by 5'-UUAG-3', and this may be replaced with another sequence such as 5'-NNNN-3' and 5'-NNN-3', if necessary. Here, N is each independently A, C, G, or U. For example, the 5'-NNNN-3' may be 5'-GAAA-3', and the 5'-NNN-3' may be 5'-CGA-3'.

[0271] For example, in the engineered gRNA of Formula (I), a sequence of the loop portion connecting X^{b1} and X^{b2} is 5'-UUAG-3', and the sequence 5'- X^{b1} UUAG X^{b2} -3' in Formula (I) may comprise or consist of any one nucleotide sequence selected from the group consisting of SEQ ID NOS: 34 to 38.

(6) Examples of gRNAs to which Modifications at Modification Sites 1 to 5 have been Applied

[0272] The engineered guide RNA (gRNA) included in the target nucleic acid editing system of the present disclosure

may comprise modifications at two or more of the above-mentioned modification sites 1 (MS1) to 5 (MS5).

[0273] In some embodiments, the engineered guide RNA may comprise one or more modifications selected from the group consisting of (a1) deletion of at least a part of the first stem region; (a2) deletion at least a part of the second stem region; (b) deletion of at least a part of the tracrRNA-crRNA complementarity region; (c) replacement of one or more uracil (U) residues with A, G, or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region; and (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence. The U-rich tail sequence may be represented by 5'-(U_mV)_nU_o-3', wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5.

[0274] For example, the engineered guide RNA may comprise (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence and (c) replacement of one or more uracil (U) residues with A, G, or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region.

[0275] As another example, the engineered guide RNA may comprise (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence, (c) replacement of one or more U residues with A, G or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region, and (a1) deletion of at least a part of the first stem region.

[0276] As yet another example, the engineered guide RNA may comprise (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence, (c) replacement of one or more U residues with A, G or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region, and (a1) deletion of at least a part of the first stem region.

[0277] As still yet another example, the engineered guide RNA may comprise (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence, (a1) deletion of at least a part of the first stem region, and (b) deletion of at least a part of the tracrRNA-crRNA complementarity region, wherein the engineered guide RNA may further comprise replacement of one or more U residues with A, G or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region containing partial deletion.

[0278] As still yet another example, the engineered guide RNA may comprise (d1) addition of a U-rich tail to the 3'-end of the crRNA sequence, (a1) deletion of at least a part of the first stem region, (b) deletion of at least a part of the tracrRNA-crRNA complementarity region, and (a2) deletion of at least a part of the second stem region, wherein the engineered guide RNA may further comprise replacement of one or more U residues with A, G or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region containing partial deletion.

[0279] As an example of tracrRNA to which modifications at the plurality of modification sites (MS) as described above have been applied, there is provided an engineered tracrRNA comprising the nucleotide sequence of any one of SEQ ID NOS: 87 to 132.

[0280] Specifically, the engineered tracrRNA of the present disclosure may comprise or consist of the nucleotide

sequence of SEQ ID NO: 87 (MS1), SEQ ID NO: 88 (MS1/MS3-1), SEQ ID NO: 89 (MS1/MS3-2), SEQ ID NO: 90 (MS1/MS3-3), SEQ ID NO: 91 (MS1/MS4*-1), SEQ ID NO: 92 (MS1/MS4*-2), SEQ ID NO: 93 (MS1/MS4*-3), SEQ ID NO: 94 (MS1/MS5-1), SEQ ID NO: 95 (MS1/MS5-2), SEQ ID NO: 96 (MS1/MS5-3), SEQ ID NO: 97 (MS1/MS3-3/MS4*-1), SEQ ID NO: 98 (MS1/MS3-3/MS4*-2), SEQ ID NO: 99 (MS1/MS3-3/MS4*-3), SEQ ID NO: 100 (MS1/MS4*-2/MS5-1), SEQ ID NO: 101 (MS1/MS4*-2/MS5-2), SEQ ID NO: 102 (MS1/MS4*-2/MS5-3), SEQ ID NO: 103 (MS1/MS3-3/MS5-1), SEQ ID NO: 104 (MS1/MS3-3/MS5-2), SEQ ID NO: 105 (MS1/MS3-3/MS5-3), SEQ ID NO: 106 (MS1/MS3-3/MS4*-2/MS5-3), SEQ ID NO: 107 (mature form, MF), SEQ ID NO: 108 (MF/MS3-1), SEQ ID NO: 109 (MF/MS3-2), SEQ ID NO: 110 (MF/MS3-3), SEQ ID NO: 111 (MF/MS4-1), SEQ ID NO: 112 (MF/MS4-2), SEQ ID NO: 113 (MF/MS4-3), SEQ ID NO:

114 (MF/MS5-1), SEQ ID NO: 115 (MF/MS5-2), SEQ ID NO: 116 (MF/MS5-3), SEQ ID NO: 117 (MF/MS5), SEQ ID NO: 118 (MF/MS3-3/MS4-1), SEQ ID NO: 119 (MF/MS3-3/MS4-2), SEQ ID NO: 120 (MF/MS3-3/MS4-3), SEQ ID NO: 121 (MF/MS4-3/MS5-1), SEQ ID NO: 122 (MF/MS4-3/MS5-2), SEQ ID NO: 123 (MF/MS4-3/MS5-3), SEQ ID NO: 124 (MF/MS4-3/MS5), SEQ ID NO: 125 (MF/MS3-3/MS5-1), SEQ ID NO: 126 (MF/MS3-3/MS5-2), SEQ ID NO: 127 (MF/MS3-3/MS5-3), SEQ ID NO: 128 (MF/MS3-3/MS5), SEQ ID NO: 129 (MF/MS3-3/MS4-3/MS5-3), SEQ ID NO: 130 (MF/MS3-3/MS4-1/MS5), SEQ ID NO: 131 (MF/MS3-3/MS4-2/MS5), or SEQ ID NO: 132 (MF/MS3-3/MS4-3/MS5).

[0281] In some embodiments, exemplary sequences of the engineered tracrRNA, which has one or more modifications at any one or more of the modification sites selected from MS1, MS3, MS4, and MS5, are provided in Table 2.

TABLE 2

tracrRNA	Nucleotide sequence	SEQ ID NO
MS1	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCU CGGAAAGUAACCCUCGAAACAAUUCAGUGCUCUCU CCAAUUCGCACAA	87
MS1/MS3-1	GAUAAAGUGGAGAACCGCUUCACCAAAGCUGUCCU UAGGGGAUAGAACUUGAGUGAAGGUGGGCUGCUUG CAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGAAAG UAACCCUCGAAACAAUUCAGUGCUCUCUCCAAUUC UGCACAA	88
MS1/MS3-2	UGGAGAACCGCUUCACCAAAGCUGUCCCUUAGGGGA UUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAGCC UAAUGUCGAGAAGUGCUUUCUUCGAAAGUAACCCUC GAAACAAUUCAGUGCUCUCUCCAAUUCGCACAA	89
MS1/MS3-3	ACCGCUUCACCAAAGCUGUCCCUUAGGGGAUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUCUUCGAAAGUAACCCUCGAAACA AAUUCAGUGCUCUCUCCAAUUCGCACAA	90
MS1/MS4*-1	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCU CGGAAAGUAACCCUCGAAACAAUUCAGUGCUCUCU CCAAUUC	91
MS1/MS4*-2	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCU CGGAAAGUAACCCUCGAAACAAUUCAGUGCUCUCU C	92
MS1/MS4*-3	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCU CGGAAAGUAACCCUCGAAACAAUUCAGUGCUCU	93
MS1/MS5-1	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UGUUUAGAUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGAAAG GUAACCCUCGAAACAAUUCAGUGCUCUCUCCAAUUC CUGCACAA	94
MS1/MS5-2	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGC UUAGGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAG CCUAAUGUCGAGAAGUGCUUUCUUCGAAAGUAACCC UCGAAACAAUUCAGUGCUCUCUCCAAUUCGCACAA A	95

TABLE 2-continued

tracrRNA	Nucleotide sequence	SEQ ID NO
MS1/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAUAG UUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUC GAGAAGUGCUUUUUCUGGAAAGUAACCCUCGAAACAA AUUCAGUGCUCUCUCCAAUUCUGCACAA	96
MS1/MS3- 3/MS4*-1	ACCGCUUCACCAAAGCUGUCCUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUUUCUGGAAAGUAACCCUCGAAACA AAUUCAGUGCUCUCUCCAAUUC	97
MS1/MS3- 3/MS4*-2	ACCGCUUCACCAAAGCUGUCCUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUUUCUGGAAAGUAACCCUCGAAACA AAUUCAGUGCUCUCUC	98
MS1/MS3- 3/MS4*-3	ACCGCUUCACCAAAGCUGUCCUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUUUCUGGAAAGUAACCCUCGAAACA AAUUCAGUGCU	99
MS1/MS4*- 2/MS5-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUUUAGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUUUCUGGAAA GUAACCCUCGAAACAAAUUCAGUGCUCUCUC	100
MS1/MS4*- 2/MS5-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UUAGGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAG CCUAAUGUCGAGAAGUGCUUUUUCUGGAAAGUAACCC UCGAAACAAAUUCAGUGCUCUCUC	101
MS1/MS4*- 2/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAUAG UUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUC GAGAAGUGCUUUUUCUGGAAAGUAACCCUCGAAACAA AUUCAGUGCUCUCUC	102
MS1/MS3- 3/MS5-1	ACCGCUUCACCAAAGCUGUUUAGAUUAGAACUUGAG UGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGAA GUGCUUUUCUGGAAAGUAACCCUCGAAACAAAUUCA GUGCUCUCUCCAAUUCUGCACAA	103
MS1/MS3- 3/MS5-2	ACCGCUUCACCAAAGCUUAGGAACUUGAGUGAAGGU GGGCUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUU CUUCGAAAGUAACCCUCGAAACAAAUUCAGUGCUCU UCUCCAAUUCUGCACAA	104
MS1/MS3- 3/MS5-3	ACCGCUUCACCAAUAGUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUGGAA AGUAACCCUCGAAACAAAUUCAGUGCUCUCUCCAAU UCUGCACAA	105
MS1/MS3- 3/MS4*-2/ MS5-3	ACCGCUUCACCAAUAGUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUGGAA AGUAACCCUCGAAACAAAUUCAGUGCUCUCUC	106
Mature Form (MF)	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUAGGGGAUUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCU CGAAAGUAACCCUCGAAACAAAUUCAUUU	107
MF/MS3-1	GAUAAAGUGGAGAACCGCUUCACCAAAGCUGUCCU UAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUUG CAUCAGCCUAAUGUCGAGAAGUGCUUUUCUGGAAAG UAACCCUCGAAACAAAUUCAUUU	108
MF/MS3-2	UGGAGAACCGCUUCACCAAAGCUGUCCCUAGGGGA UUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAGCC UAAUGUCGAGAAGUGCUUUUCUGGAAAGUAACCCUC GAAACAAAUUCAUUU	109
MF/MS3-3	ACCGCUUCACCAAAGCUGUCCCUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUUCUGGAAAGUAACCCUCGAAACA AAUUCAUUU	110

TABLE 2-continued

tracrRNA	Nucleotide sequence	SEQ ID NO
MF/MS4-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUU CGGAAAGUAACCCUCGAAACAAAUCAU	111
MF/MS4-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUU CGGAAAGUAACCCUCGAAACAAAUUC	112
MF/MS4-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGG CUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUU CGGAAAGUAACCCUCGAAACAA	113
MF/MS5-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUUUAGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAA GUAACCCUCGAAACAAAUUCAUUU	114
MF/MS5-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UUUAGAGAACUUGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAAGUAAC CCUCGAAACAAAUUCAUUU	115
MF/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAUAG UUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUC GAGAAGUGCUUUCUUCGGAAAGUAACCCUCGAAACAA AUUCAUUU	116
MF/MS5	CUUCACUGAUAAAGUGGAGAACCGCUUCACUAGAGU GAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGAAG UGCUUUCUUCGGAAAGUAACCCUCGAAACAAAUUCAU UU	117
MF/MS3- 3/MS4-1	ACCGCUUCACCAAAGCUGUCCCUUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUCUUCGGAAAGUAACCCUCGAAACA AAUUCAU	118
MF/MS3- 3/MS4-2	ACCGCUUCACCAAAGCUGUCCCUUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUCUUCGGAAAGUAACCCUCGAAACA AAUUC	119
MF/MS3- 3/MS4-3	ACCGCUUCACCAAAGCUGUCCCUUAGGGGAUUAGAA CUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGU CGAGAAGUGCUUUCUUCGGAAAGUAACCCUCGAAACA AA	120
MF/MS4- 3/MS5-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UGUUUAGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAA GUAACCCUCGAAACAA	121
MF/MS4- 3/MS5-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGC UUUAGAGAACUUGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAAGUAAC CCUCGAAACAA	122
MF/MS4- 3/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAUAG UUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUC GAGAAGUGCUUUCUUCGGAAAGUAACCCUCGAAACAA A	123
MF/MS4-3/ MS5	CUUCACUGAUAAAGUGGAGAACCGCUUCACUAGAGU GAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGAAG UGCUUUCUUCGGAAAGUAACCCUCGAAACAA	124
MF/MS3- 3/MS5-1	ACCGCUUCACCAAAGCUGUUUAGAUUAGAACUUGAG UGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGAA GUGCUUUCUUCGGAAAGUAACCCUCGAAACAAAUUCA UUU	125

tracrRNA	Nucleotide sequence	SEQ ID NO
MF/MS3-3/MS5-2	ACCGCUUACACAAAAGCUUUGAGAACUUGAGUGAAG GUGGGCUGCUUGCAUCAGCCUAAUGUCGAGAAGUGCU UUCUUCGGAAGUAACCCUCGAAACAAAUUCAUUU	126
MF/MS3-3/MS5-3	ACCGCUUACCAAUUAGUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGA AGUAACCCUCGAAACAAAUUCAUUU	127
MF/MS3-3/MS5	ACCGCUUACCUUAGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUAAC CCUCGAAACAAAUUCAUUU	128
MF/MS3-3/MS4-3/MS5-3	ACCGCUUACCAAUUAGUUGAGUGAAGGUGGGCUGCU UGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGA AGUAACCCUCGAAACAA	129
MF/MS3-3/MS4-1/MS5	ACCGCUUACCUUAGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUAAC CCUCGAAACAAAUUCAU	130
MF/MS3-3/MS4-2/MS5	ACCGCUUACCUUAGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUAAC CCUCGAAACAAAUUC	131
MF/MS3-3/MS4-3/MS5	ACCGCUUACCUUAGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUAAC CCUCGAAACAA	132

NO: 139 (MF/MS4-2), SEQ ID NO: 140 (MF/M4-3), SEQ ID NO: 141 (MS1/M2), SEQ ID NO: 142 (MS1/MS2/MS4*1), SEQ ID NO: 143 (MS1/MS2/MS4*-2), SEQ ID NO: 144 (MS1/MS2/MS4*3), SEQ ID NO: 145 (MF/M32), SEQ ID NO: 146 (MF/M2/MS4-1), SEQ ID NO: 147 (MF/MU2/MC4-2), or SEQ ID NO: 148 (MF/MS2/MS4-3). In some embodiments, exemplary sequences of the engineered crRNA, which has one or more modifications at any one or more modification sites selected from MS 1, MS2, and MS4 are provided in Table 3.

TABLE 3

crRNA	Nucleotide sequence	SEQ ID NO
MS1	GUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUGCAAC	133
MS1/MS4*-1	GAACCCGAAUAGAGCAAUGAAGGAAUGCAAC	134
MS1/MS4*-2	GAAUAGAGCAAUGAAGGAAUGCAAC	135
MS1/MS4*-3	AGCAAUGAAGGAAUGCAAC	136
MF	GAAUGAAGGAAUGCAAC	137
MF/MS4-1	AUGAAGGAAUGCAAC	138
MF/MS4-2	GAAGGAAUGCAAC	139
MF/MS4-3	GGAAUGCAAC	140
MS1/MS2	GUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUGCAA CNNNNNNNNNNNNNNNNNNNNUUUAUUUUUU	141
MS1/MS2/MS4*- -1	GAACCCGAAUAGAGCAAUGAAGGAAUGCAACNNNNN NNNNNNNNNNNNNNNNUUUAUUUUUU	142
MS1/MS2/MS4*- -2	GAAUAGAGCAAUGAAGGAAUGCAACNNNNNNNNNN NNNNNNNNNNUUUAUUUUUU	143

TABLE 3-continued

crRNA	Nucleotide sequence	SEQ ID NO
MS1/MS2/MS4*-3	AGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNNNN NNNUUUUAUUUUUU	144
MF/MS2	GAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNNNNNN NUUUUAUUUUUU	145
MF/MS2/MS4-1	AUGAAGGAAUGCAACNNNNNNNNNNNNNNNNNNNNNNU UUUAUUUUUU	146
MF/MS2/MS4-2	GAAGGAAUGCAACNNNNNNNNNNNNNNNNNNNNNNNUU UAUUUUUU	147
MF/MS2/MS4-3	GGAAUGCAACNNNNNNNNNNNNNNNNNNNNNNNUUUUAU UUUUU	148

[0283] In Table 3, indication of a guide sequence (spacer) is omitted from all crRNA sequences unless necessary, and the sequence indicated by ‘NNNNNNNNNNNNNNNNNNNNNN’ indicates any guide sequence (spacer) that can hybridize with a target sequence in a target gene. The guide sequence may be appropriately designed by those skilled in the art depending on a desired target gene and/or a target sequence in the target gene as described above, and therefore is not limited to a specific sequence of a particular length. In another embodiment, the engineered gRNA may comprise tracrRNA comprising or consisting of any one nucleotide sequence selected from the group consisting of SEQ ID NOS: 87 to 132; and crRNA comprising or consisting of any one nucleotide sequence selected from the group consisting of SEQ ID NOS: 133 to 148. In another embodiment, when the engineered gRNA of the present disclosure is in the form of a single guide RNA (sgRNA), the engineered sgRNA may sgRNA comprising or consisting of any one nucleotide sequence selected from the group consisting of SEQ ID NOS: 149 to 186.

[0284] Specifically, the engineered sgRNA may be sgRNA of SEQ ID NO: 149 comprising a modification at MS1, sgRNA of SEQ ID NO: 150 comprising modifications at MS1/MS2, sgRNA of SEQ ID NO: 151 comprising modifications at MS1/MS2/MS3, sgRNA of SEQ ID NO: 152 comprising modifications at MS2/MS3/MS4, or sgRNA of SEQ ID NO: 153 comprising modifications at MS2/MS3/MS4/MS5.

[0285] In another specific example, the engineered sgRNA may be sgRNA comprising or consisting of the nucleotide sequence of SEQ ID NO: 154 (MS1/MS3-1), SEQ ID NO: 155 (MS1/MS3-2), SEQ ID NO: 156 (MS1/MS3-3), SEQ ID NO: 157 (MS1/MS4*-1), SEQ ID NO: 158 (MS1/MS4*-2), SEQ ID NO: 159 (MS1/MS4*-3), SEQ ID NO: 160 (MS1/MS5-1), SEQ ID NO: 161 (MS1/MS5-2), SEQ ID NO: 162 (MS1/MS5-3), SEQ ID NO: 163 (MS1/MS2/MS4*-2), SEQ ID NO: 164 (MS1/MS3-3/MS4*-2), SEQ ID NO: 165 (MS1/MS2/MS5-3), SEQ ID NO: 166 (MS1/MS3-3/MS5-3), SEQ ID NO: 167 (MS1/MS4*-2/MS5-3), SEQ ID NO: 168 (MS1/MS2/MS3-3/MS4*-2), SEQ ID NO: 169 (MS1/MS2/MS3-3/MS5-3), SEQ ID NO: 170 (MS1/MS2/MS4*-2/MS5-3), SEQ ID NO: 171 (MS1/MS3-3/MS4*-2/MS5-3), or SEQ ID NO: 172 (MS1/MS2/MS3-3/MS4*-2/MS5-3).

[0286] In addition, the sgRNA may be sgRNA comprising or consisting of the nucleotide sequence of SEQ ID NO: 173, which is a mature form (abbreviated as MF) of sgRNA.

[0287] In another embodiment, there is provided an exemplary sgRNA which comprises partial modification of the nucleotide sequence of the MF sgRNA. Specifically, the MF sgRNA may be an sgRNA comprising or consisting of the nucleotide sequence of SEQ ID NO: 174 (MS3-1), SEQ ID NO: 175 (MS3-2), SEQ ID NO: 176 (MS3-3), SEQ ID NO: 177 (MS4-1), SEQ ID NO: 178 (MS4-2), SEQ ID NO: 179 (MS4-3), SEQ ID NO: 180 (MS5-1), SEQ ID NO: 181 (MS5-2), SEQ ID NO: 182 (MS5-3), SEQ ID NO: 183 (MS3-3/MS4-3), SEQ ID NO: 184 (MS3-3/MS5-3), SEQ ID NO: 185 (MS4-3/MS5-3), or SEQ ID NO: 186 (MS3-3/MS4-3/MS5-3).

[0288] In a preferred embodiment, the engineered sgRNA may consist of the nucleotide sequence of SEQ ID NO: 151 (Cas12f_ge3.0), SEQ ID NO: 152 (Cas12f_ge4.0), or SEQ ID NO: 153 (Cas12f_ge4.1).

(7) Additional Sequence

[0289] The engineered tracrRNA of the present disclosure may optionally further comprise an additional sequence. The additional sequence may be located at the 3'-end of the engineered tracrRNA. In addition, the additional sequence may be located at the 5'-end of the engineered tracrRNA. For example, the additional sequence may be located at the 5'-end of the first stem region.

[0290] The additional sequence may consist of 1 to 40 nucleotides. In an embodiment, the additional sequence may be any nucleotide sequence or a randomly arranged nucleotide sequence. For example, the additional sequence may be 5'-AUAAGGUGA-3' (SEQ ID NO: 187).

[0291] In addition, the additional sequence may be a known nucleotide sequence. For example, the additional sequence may be a hammerhead ribozyme nucleotide sequence. Here, the hammerhead ribozyme nucleotide sequence may be 5'-CUGAUGAGU-CCGUGAGGACGAAACGAGUAAGCUCGUC-3' (SEQ ID NO: 188) or 5'-CUGCUCGAAUGAGCAAAGCAG-GAGUGCCUGAGUAGUC-3' (SEQ ID NO: 189). The sequences listed above are merely examples, and the additional sequence is not limited thereto.

(8) Chemical Modification

[0292] In some embodiments, the engineered tracrRNA or engineered crRNA included in the engineered gRNA may have chemical modification in at least one or more nucleotides, if necessary. Here, the chemical modification may be a modification in various covalent bonds that may occur in a nucleotide base and/or sugar portion.

[0293] For example, the chemical modification may be methylation, halogenation, acetylation, phosphorylation, phosphorothioate (PS) linkage, locked nucleic acid (LNA), 2'-O-methyl 3'phosphorothioate (MS) or 2'-O-methyl 3'thiopropyl (MSP). The above example is a simple example and the modification is not limited thereto.

[0294] In a case of using the hypercompact gene editing system comprising a complex of the engineered gRNA with engineered Cas12f1 (CWCas12f1 or UnCas12f1) of the present disclosure, indel efficiency for a target gene or target nucleic acid in a cell is significantly improved compared to a case of using the guide RNA or Cas12f1 found in nature.

[0295] Above all, the engineered gRNA may involve optimized length for high efficiency and resulting cost reduction in gRNA synthesis, creation of additional space or capacity in a case of being inserted into a viral vector, normal expression of tracrRNA, increased expression of operable gRNA, increased gRNA stability, increased stability of complex of gRNA with nucleic acid editing protein, induction of formation of complex of gRNA with nucleic acid editing protein at high efficiency, increased cleavage efficiency of target nucleic acid by hypercompact nucleic acid editing system comprising complex of gRNA with nucleic acid editing protein, and increased homology-directed repair efficiency for target nucleic acid caused by such a system. Accordingly, in a case of using the above-described engineered gRNA for Cas12f1 or an engineered Cas12f1 protein, it is possible to overcome the limitations of the above-mentioned prior art, thereby cleaving or editing a gene with high efficiency in a cell.

[0296] In addition, the engineered gRNA has a short length compared to gRNA found in nature, and thus has high applicability in the field of gene editing technology. Using the engineered gRNA, the hypercompact gene editing system comprising a complex of the gRNA with nucleic acid editing protein has advantages of being very small in size and having excellent editing efficiency, which allows the system to be utilized in various gene editing technologies.

3. Nucleic Acid Construct Encoding Each Component of Cas12f1 System

[0297] Since each component of the gene editing system provided in the present disclosure is intended to be expressed in a cell, according to an embodiment, there is provided a nucleic acid construct encoding each component of the gene editing system. The nucleic acid construct may comprise a synthetic nucleotide sequence.

[0298] In an embodiment, the nucleic acid may be DNA or RNA (for example, mRNA). The nucleic acid or polynucleotide encoding each component of the gene editing system are disclosed herein as a representative example, or the nucleotide sequence thereof may be readily determined by those skilled in the art by referring to the specific sequence of each component.

[0299] In an embodiment, the nucleic acid construct may comprise a human codon-optimized nucleotide sequence

encoding a Cas12f1 protein. The term “codon optimization” refers to a process of modifying a native nucleic acid sequence for enhanced expression in a cell of interest by replacing at least one codon in the native sequence with a codon, which is used more frequently or most frequently in a gene of the target cell, while maintaining its native amino acid sequence. Different species have specific biases for specific codons for specific amino acids, and codon bias (differences in codon usage between organisms) is often correlated with translation efficiency of an mRNA, which is considered to be dependent on the nature of codons being translated and availability of specific tRNA molecules. Predominance of tRNA selected in a cell generally reflects the most frequently used codon in peptide synthesis. Thus, genes may be tailored for optimal gene expression in a given organism based on codon optimization.

[0300] For example, the nucleic acid encoding the human codon optimized CWCas12f1 protein or a variant thereof may comprise or consist of a sequence selected from SEQ ID NOS: 6 to 9. In addition, the nucleic acid encoding the human codon optimized UnCas12f1 protein may comprise or consist of the sequence of SEQ ID NO: 10.

[0301] In another embodiment, the nucleic acid or polynucleotide may be DNA or RNA that exists in nature, or may be a modified nucleic acid in which a chemical modification has occurred in at least a part of the nucleic acid or polynucleotide. For example, the nucleic acid or polynucleotide may be one in which one or more nucleotides have been chemically modified. Here, the chemical modification may include any modification of nucleic acids known to those skilled in the art.

IV. Vector or Vector System

[0302] As disclosed herein, the nucleic acid construct may be a vector or may be contained in a vector. The components of the composition according to an embodiment of the present disclosure may be one nucleic acid construct or two or more nucleic acid constructs. In addition, the nucleic acid construct(s) may be present in one vector or may be present separately in two or more vectors. In an embodiment, when the composition comprises one or more vectors, the composition may be a vector system. Since the vector or vector system allows each component of the above-described Cas12f1 system to be expressed in a cell, the nucleic acid construct (for example, nucleotide sequence) included in the vector system comprises at least one nucleotide sequence encoding each component of an embodiment of the present disclosure. In addition, since the disclosed vector system allows each component of the present disclosure to be expressed in a cell, all effects and advantages that are achieved by the present disclosure are applied as is.

[0303] In the disclosed vector system, each nucleic acid construct is capable of expressing each component (for example, an inhibitory molecule, an endonuclease, and first and second guide RNAs) in a cell. The vector system enables removal of a nucleic acid segment to be deleted in a cell.

[0304] In the vector system disclosed herein, for the nucleotide sequence of each nucleic acid construct and the components expressed thereby, see the above-described details.

[0305] In order to use the composition disclosed herein for deletion of a nucleic acid segment, a method may be used in which one or more vectors comprising nucleotide sequences encoding respective components are introduced directly or

through an appropriate delivery means or delivered through a vehicle, such as a virus, into a target cell and the respective components of the gene editing system are allowed to be expressed in the target cell. Preferably, the nucleic acid constructs, whose nucleotide sequences encoding respective components are operably linked, may be contained in a single vector.

[0306] In an embodiment, the nucleic acid construct encoding one or more components of the above-described composition may be present in two or more vectors.

[0307] In another embodiment, the nucleic acid constructs encoding one or more components of the above-described composition may be present in a single vector (for example, AAV).

[0308] In addition, the vector system of the present disclosure may comprise, in addition to the above-described components, a nucleotide sequence encoding an additional expression element that is desired to be expressed as needed by those skilled in the art. For example, the additional expression element may be a tag. Specifically, the additional expression element may be a herbicide resistance gene such as glyphosate, glufosinate ammonium, or phosphinothricin, or an antibiotic resistance gene such as ampicillin, kanamycin, G418, bleomycin, hygromycin, or chloramphenicol.

[0309] In another embodiment, the vector or vector system needs to comprise one or more regulatory and/or control components so that it is directly expressed in a cell. Specifically, the regulatory and/or control components may include, but are not limited to, a promoter, an enhancer, an intron, a polyadenylation signal, a Kozak consensus sequence, an internal ribosome entry site (IRES), a splice acceptor, a 2A sequence, and/or a replication origin. The replication origin may be, but is not limited to, an f1 origin of replication, an SV40 origin of replication, a pMB1 origin of replication, an adeno origin of replication, an AAV origin of replication, and/or a BBV origin of replication.

[0310] In another embodiment, in order to express, in a cell, the nucleic acid construct encoding each component in a cell, a promoter sequence may need to be operably linked to the sequence encoding each component so that an RNA transcription factor can be activated in the cell. The promoter sequence may be designed differently depending on the corresponding RNA transcription factor or expression environment, and is not limited as long as it can properly express the components of the composition of the present disclosure in a cell.

[0311] For example, the promoter sequence may be a promoter that promotes transcription of RNA polymerase RNA Pol I, Pol II, or Pol III. Specifically, the promoter may be one of U6 promoter, EFS promoter, EF1- α promoter, H1 promoter, 7SK promoter, CMV promoter, LTR promoter, Ad MLP promoter, HSV promoter, SV40 promoter, CBA promoter, or RSV promoter.

[0312] In another embodiment, when a sequence of the vector comprises the promoter sequence, transcription of a sequence operably linked to the promoter is induced by an RNA transcription factor, and the vector may comprise a termination signal that induces termination of transcription of the RNA transcription factor. The termination signal may vary depending on the type of the promoter sequence. Specifically, when the promoter is a U6 or H1 promoter, the promoter recognizes a TTTTT (T5) or TTTTTT (T6) sequence, which is a thymidine (T) repeat sequence, as a termination signal.

[0313] The sequence of the engineered guide RNA according to an embodiment may comprise a U-rich tail sequence at its 3-end. Accordingly, the sequence encoding the engineered guide RNA comprises a T-rich sequence corresponding to the U-rich tail sequence at its 3-end. As described above, some promoter sequences recognize a thymidine (T) repeat sequence, for example, a sequence consisting of five or more consecutive thymidine (T) residues, as a termination signal, and therefore, in some cases, the T-rich sequence may be recognized as a termination signal. In other words, when the vector sequence provided herein comprises a sequence encoding the engineered guide RNA, a sequence encoding the U-rich tail sequence included in the engineered gRNA sequence may be used as a termination signal.

[0314] In an embodiment, when the vector sequence comprises a U6 or H1 promoter sequence and a sequence encoding the engineered guide RNA operably linked thereto, a sequence portion that encodes the U-rich tail sequence included in the guide RNA sequence may be recognized as a termination signal. Specifically, the U-rich tail sequence may comprise a sequence consisting of five or more consecutive uridine (U) residues.

[0315] In an embodiment, the vector may be a viral vector. Specifically, the viral vector may be at least one selected from the group consisting of a retrovirus vector, a lentivirus vector, an adenovirus vector, an adeno-associated virus vector, a vaccinia virus vector, a poxvirus vector, a herpes simplex virus vector, and a phagemid vector. Preferably, the viral vector may be an adeno-associated viral vector (AAV). In addition, the viral vector includes, but is not limited to, a SIN lentivirus vector, a retrovirus vector, a foamy virus vector, an adenovirus vector, an adeno-associated virus (AAV) vector, a hybrid vector and/or a plasmid transposon (for example, the Sleeping Beauty transposon system), or an integrase-based vector system.

[0316] In another embodiment, the vector may be a non-viral vector. Specifically, the non-viral vector may be at least one selected from the group consisting of, but not limited to, plasmid, naked DNA, DNA complex, mRNA (transcript), and amplicon. For example, the plasmid may be selected from the group consisting of pcDNA series, pSC101, pGV1106, pACYC177, ColE1, pKT230, pME290, pBR322, pUC8/9, pUC6, pBD9, pHc79, pIJ61, pLAFR1, pHV14, pGEX series, pET series, and pUC19.

[0317] The term “naked DNA” refers to DNA (for example, histone-free DNA) that encodes a protein, such as Cas12f1 of the present disclosure, cloned into a suitable expression vector (for example, plasmid) in an appropriate orientation for expression.

[0318] The term “amplicon,” when used with respect to a nucleic acid, means a product of copying the nucleic acid, wherein the product has a nucleotide sequence that is identical with or complementary to at least a portion of the nucleotide sequence of the nucleic acid. For example, an amplicon may be produced by any of a variety of amplification methods that use a nucleic acid or an amplicon thereof as a template, including polymerase extension, polymerase chain reaction (PCR), rolling circle amplification (RCA), multi-displacement amplification (MDA), ligation extension, or ligation chain reaction. The amplicon may be a nucleic acid molecule having a single copy of a particular nucleotide sequence (for example, a PCR product) or multiple copies of the nucleotide sequence (for example, a concatemeric product of RCA).

[0319] The vector disclosed herein may be designed in the form of a linear or circular vector. In a case where the vector is a linear vector, RNA transcription is terminated at the 3-end even if a sequence of the linear vector does not separately comprise a termination signal. However, in a case where the vector is a circular vector, RNA transcription is not terminated unless a sequence of the circular vector separately comprises a termination signal. Therefore, when using a circular vector, a termination signal corresponding to a transcription factor related to each promoter sequence has to be included in order for the vector to express an intended target.

[0320] In an embodiment, the viral vector or non-viral vector may be delivered by a delivery system such as liposomes, polymeric nanoparticles (for example, lipid nanoparticles), oil-in-water nanoemulsions, or combinations thereof, or in the form of a virus.

[0321] In another embodiment, the virus may be selected from the group consisting of a retrovirus, a lentivirus, an adenovirus, an adeno-associated virus, a vaccinia virus, a poxvirus, a herpes simplex virus and a phage.

[0322] In yet another embodiment, the phage may be selected from the group consisting of λ gt4,B, λ -Charon, λ Δz1, and M13.

[0323] In order to efficiently deliver the nucleic acid construct included in the composition of the present disclosure into a target cell or target site via a virus, in particular, an adeno-associated virus (AAV), it is important to design a size of the nucleotide sequence encoding all components of the editing system to be within 4.7 kb that is a packaging limit of AAV. This has an advantage in that in a case where the Cas12f1 system of the present disclosure is used, a very small size of the hypercompact nucleic acid editing protein and two engineered gRNAs included in the system allows sufficient packaging by AAV even if an additional regulatory molecule (for example, a nucleic acid construct that expresses the inhibitory molecule of the present disclosure) is further included.

V. Formulation

[0324] As disclosed herein, the composition may be a pharmaceutical composition.

[0325] In an embodiment, the pharmaceutical composition may be for deletion of a nucleic acid segment. In addition, the pharmaceutical composition may be for treating or delaying onset or progression of a genetic disease caused by an undesirable mutation.

[0326] In an embodiment, the pharmaceutical composition may be formulated according to the mode of administration to be used. For example, in a case where the pharmaceutical composition is an injectable pharmaceutical composition, it may be desirable to use an isotonic agent. An additive for isotonicity may generally include sodium chloride, dextrose, mannitol, sorbitol, and lactose. In an embodiment, isotonic solutions such as phosphate buffered saline are preferred. A stabilizer may include gelatin and albumin. In an embodiment, a vasoconstrictor is added to the formulation.

[0327] In another embodiment, the composition may further comprise a pharmaceutically acceptable excipient. The pharmaceutically acceptable excipient may be a functional molecule that acts as a vehicle, an adjuvant, a carrier, or a diluent. The pharmaceutically acceptable excipient may be a gene transfer enhancer (which may include a surfactant) such as an immune stimulating complex (ISCOMS), Freund's

incomplete adjuvant, a LPS analogue (including monophosphoryl lipid A), a muramyl peptide, a quinone analogue, a vesicle such as squalene and squalane, hyaluronic acid, a lipid, a liposome, a calcium ion, a viral protein, a polyanion, a polycation, or a nanoparticle, or other known gene transfer facilitating agent.

[0328] In another embodiment, the composition may comprise a gene transfer enhancer. The gene transfer enhancer may be a polyanion, a polycation (including poly-L-glutamic acid (LGS)), or a lipid. The gene transfer enhancer is poly-L-glutamic acid, and more preferably, the poly-L-glutamic acid may be present in the composition for genome editing of skeletal muscle or cardiac muscle at a concentration of less than 6 mg/ml. The gene transfer enhancer may also include a surfactant, such as an immune stimulating complex (ISCOMS), Freund's incomplete adjuvant, a LPS analogue (including monophosphoryl lipid A), a muramyl peptide, a quinone analogue and a vesicle, such as squalene and squalane; and hyaluronic acid may also be used.

[0329] In an embodiment, the composition comprising one or more vectors included in the above-described vector system may comprise a gene transfer enhancer, such as a lipid, a liposome (including lecithin liposomes, or other liposomes known in the art), a DNA-liposome mixture, a calcium ion, a viral protein, a polyanion, a polycation, or a nanoparticle, or other known gene transfer enhancer. Preferably, the gene transfer enhancer is a polyanion, a polycation (for example, poly-L-glutamic acid (LGS)), or lipid 17.

[0330] An actual dosage of the (pharmaceutical) composition may vary greatly depending on various factors, such as the choice of vector, the target cell, organism, or tissue, the condition of the subject to be treated, the degree of transformation/modification sought, the route of administration, the method of administration, the form of transformation/modification sought, and the like. The administration may be performed by a route of administration selected from subretinal administration, subcutaneous administration, intradermal administration, intraocular administration, intravitreal administration, intratumoral administration, intranodal administration, intramedullary administration, intramuscular administration, intravenous administration, intralymphatic administration, and intraperitoneal administration. The pharmaceutical composition may further comprise a carrier (water, saline, ethanol, glycerol, lactose, sucrose, calcium phosphate, gelatin, dextran, agar, pectin, peanut oil, sesame oil, and the like), a diluent, a pharmaceutically acceptable carrier (for example, phosphate buffered saline), a pharmaceutically acceptable excipient, and/or other compounds known in the art.

[0331] For example, delivery for treatment of a disease may be via AAV. A therapeutically effective dosage for in vivo delivery of AAV to a human may be a saline solution in a range of about 20 ml to about 50 ml containing about 1×10^{10} to about 1×10^{100} AAV per ml of solution. The dosage may be adjusted to balance the therapeutic benefit against any adverse effects.

Form for Carrying Out the Disclosure

[0332] Hereinafter, the present disclosure will be described in more detail by the following examples. However, these examples are only intended to illustrate the present disclosure, and the scope of the present disclosure is not limited to these examples.

Example 1. Experimental Method

1. Preparation of Genomic DNA (gDNA)

[0333] AC16 or HEK293T cells were transfected and harvested. Genomic DNA was prepared using a gDNA prep kit (Maxwell® RSC Cultured Cells DNA, PROMEGA, AS1620). Samples were prepared according to the manufacturer's instructions.

2. PCR and Gel Purification

[0334] This experiment was performed using the GEL & PCR Purification System (GP104-200, Biofact). To the PCR product was added UB buffer in an amount equivalent to 3 times the volume of the PCR product and thorough mixing was performed. Then, isopropanol was added thereto in an amount equivalent to 2 times the volume of the PCR product and thorough mixing was performed. In a case of the gel, the gel of the corresponding band was cut and weighed. Then, UB buffer was added thereto in an amount equivalent to 3 times the weight of the gel. The gel was dissolved by incubation at 65° C. for 10 minutes, and then isopropanol was added thereto in an amount equivalent to 1 time the gel volume and thorough mixing was performed. The column was prepared, 200 µl of HelpB buffer was added to the column, centrifugation was performed at 13,000 rpm for 30 seconds. Then, the filtered solution was discarded. The reaction solution was added to the column, centrifugation was performed at 7,000 rpm for 1 minute. Then, the filtered solution was discarded. 750 µl of 80% EtOH was added thereto, centrifugation was performed at 13,000 rpm for 30 seconds. Then, the filtered solution was discarded. After repeating the process twice, centrifugation was performed at 13,000 rpm for 3 minutes. The centrifuged column was placed in a 1.5 ml tube, 30 µl of EB buffer was added dropwise to the center, and the reaction was allowed to occur at room temperature for 1 minute. Centrifugation was performed at 13,000 rpm for 1 minute. The DNA collected in the 1.5 ml tube was quantified and stored at 4° C.

3. Collection of Plasmid Vector

[0335] For transfection or Sanger sequencing, the vector-transformed DH5u was used. Plasmid Mini prep kit (PM105-200, Biofact) was used according to the manufacturer's instructions. The culture medium of the vector-transformed DH5u was placed in a 1.5 ml tube, and centrifugation was performed at 13,000 rpm for 5 minutes. After centrifugation, the supernatant was discarded, and the pellet was sufficiently dispersed by vortexing. 350 µl of B1 buffer was added thereto, and the tube was shaken to ensure sufficient reaction. Next, 350 µl of A1 buffer containing RNase A was added thereto, and the tube was inverted until the blue color disappeared. Then, centrifugation was performed at 13,000 rpm for 5 minutes. The column was prepared, 200 µl of HelpB buffer was added thereto, and the solution was centrifuged at 13,000 rpm for 30 seconds. Then, the filtered solution was discarded. 750 µl of the centrifuged supernatant was added to the prepared column, centrifugation was performed at 7,000 rpm for 1 minute, and the filtered solution was discarded. 750 µl of 80% EtOH was added thereto, centrifugation was performed at 13,000 rpm for 30 seconds, and the filtered solution was discarded. This process was repeated twice. After repeating the process twice, centrifugation was performed at 13,000 rpm for 3 minutes. The centrifuged column was placed into a 1.5 ml

tube, 30 µl of EB buffer was added dropwise to the center, and then the reaction was allowed to occur at room temperature for 1 minute. Centrifugation was performed at 13,000 rpm for 1 minute, and the plasmid vectors collected in the 1.5 ml tube were quantified and stored at -20° C.

4. Preparation of DNA Cassette

[0336] To confirm indel efficiency of the spacer sequences of Cas12f1, a cassette containing the U6 promoter, scaffold sequence, guide sequence, and U-rich tail sequence (T4AT6) was amplified by PCR and used. The process was performed as follows.

1) Selection of Spacer and Order of Oligo

[0337] The spacer was selected from the 20mer sequence followed by TTTA or TTTG, which are PAM, and spacers whose sequences end with T were excluded. In addition, to minimize off-target effects, the spacers were designed using CRISPR RGEN TOOL by classifying them with less than 2 mismatches. In addition, the reverse complement sequence comprising a DR (direct repeat) and U-rich sequence was custom-made to be used as an R primer.

2) PCR

[0338] The PCR was performed under the composition and condition shown in Table 4 below.

TABLE 4

Reagent composition		PCR condition	
2x pfu PCR Master mix	205 µl	Pre-denaturation	95° C., 5 min
hU6 F primer (10 P)	2.05 µl	Denaturation (D)	95° C., 30 s
Target oligo (10 P)	2.05 µl	Annealing	60° C., 30 s
Template	1µl (200 ng)	Extension (E)	72° C., 2 min
DW	199.9 µl	D-E Cycle	30 cycles
Total	410 µl	Final extension	72° C., 3 min
Prepared in 8 PCR tubes, each containing 50µl		Storage	4° C., ∞

[0339] 400 µl of the mixture was added to 8 PCR tubes, each containing 50 µl, and each sample was amplified.

3) Gel Analysis

[0340] 1% agarose gel was prepared, and the size marker and PCR products were added to the gel. Electrophoresis was performed to confirm the amplified size.

4) Purification and Quantification

[0341] After confirming the amplified size, the gel was purified according to Experimental Method 2 to quantify the PCR products.

5. Cell Culture

[0342] For use in the experiments, AC16 and HEK 293T cells were used as human cells, and Hepa-1c17 cells were used as mouse cells. At 37° C. in a 5% CO₂ incubator, AC16 and HEK 293T cells were maintained in DMEM medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin, and Hepa-1c17 cells were maintained in Alpha minimal medium supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin.

The attached cells were washed twice with cold PBS and then immediately stored at -80°C . for RNA or protein extraction.

6. Transfection

[0343] The day before transfection, HEK293 and AC16 cells (80% confluency) cultured in 100 mm dishes were treated with trypsin to detach from the bottom of the dish. The detached cells were placed in 50 ml of each pre-warmed medium and slowly dissolved with a pipette. 24-well plates were prepared according to the number of samples and repetitions, and 500 μl of cell suspension medium was added to each well ($1/100$ dilution). Then, incubation was performed overnight in a CO_2 incubator at 37°C . until transfection.

[0344] When the cell confluency reached approximately 70% to 80%, 200 μl of the 500 μl medium per well was removed and the plates were placed in the incubator. 1.5 ml tubes were prepared according to the number of samples, and 200 μl of Opti-MEM was added to each tube. 1.5 μg of Cas12f1 and 0.5 μg of gRNA were added to the tube containing Opti-MEM, and vortexed for 5 seconds (nucleic acid mixture). Then, the nucleic acid mixture and FuGENE HD were added at a ratio of 1:3, and reaction was allowed to occur at room temperature for 20 minutes (that is, in a case where the nucleic acid mixture was 2 μg , 6 μl of FuGENE HD was administered). The 24-well plate was taken out from the incubator, and 200 μl of the solution containing the nucleic acid mixture and FuGENE HD was gently added along the well wall. After shaking the plate sufficiently in an S shape, it was incubated in a CO_2 incubator at 37°C . for 72 hours. After 72 hours, the cells were harvested and gDNA was extracted therefrom according to Experimental Method 1.

7. Construction of Vector

[0345] The following procedure was performed using the Cas12f1 ge4.0 dual gRNA vector (see Korean Patent Application Nos. 10-2021-0051552 and 10-2022-0043768). The restriction enzyme ends of the vector to be cloned were confirmed, and dual gRNA oligos were designed and custom-made. The custom-made oligos were diluted to a concentration of 100 pmol. 4.5 μl each of the diluted forward and reverse primers was taken and placed into a PCR tube, and then 1 μl of $10\times$ annealing buffer was added thereto to adjust the total volume to 10 μl . Then, annealing was performed under the conditions of 95°C . for 5 minutes and $-1^{\circ}\text{C}/\text{min}$ from 95°C . to 4°C . The Cas12f1 ge4.0 dual gRNA vector was prepared and incubated at 500 rpm, 37°C . for 2 hours under the digestion conditions in Table 5 below.

TABLE 5

Reagent	Volume
NEB 10X 2.1 buffer	5 μl
Vector	10 μg
BbsI	10 μl
DW	Amount to make total volume of 50 μl
Total	50 μl

[0346] After digestion, the digested vector was obtained through electrophoresis and gel elution. Ligation was performed using the digested vector and annealed oligo (see Table 6).

TABLE 6

Reagent	Volume
DNA ligation mix (TAKARA)	2 μl
Annealed oligo	1.5 μg
Vector digested with BbsI	0.5 μl
Total	4 μl

[0347] After ligation, transformation was performed on DH5 α . After incubation on an LB plate, positive colonies were confirmed through colony PCR and then incubated in 3 ml LB medium. After miniprep, sequencing was performed to confirm whether the final sequences matched.

8. DH5 α Transformation

[0348] The previously-produced vector was transformed into *E. coli* to produce the vector. DH5 α competent cells were taken out and thawed on ice. The ligated vector was added up to $1/10$ of the amount of DH5 α , and the incubation was performed on ice for 30 minutes. After heat shock at 42°C . for 30 seconds, cooling was performed on ice for 2 minutes. Incubation was performed using 100 μl of LB medium or S.O.C medium at 37°C . for 1 hour. The cells were spread on LB plates warmed to room temperature (containing ampicillin or kanamycin depending on the vector) and incubated at 37°C . for 14 to 16 hours.

9. PCR of NGS Sample

[0349] NGS samples for confirming indel efficiency for a target were prepared by purification of the PCR product after performing the 1st to 3rd PCRs. The preparation of NGS samples for confirming indel efficiency was conducted over a total of 3 PCRs. The first PCR produced a band of approximately 450 to 500 bp, and the second PCR was performed using this PCR product as a template. After the 2nd PCR, the sample was loaded onto a 2% agarose gel to confirm whether the band was properly displayed within 250 bp. If the band was not properly displayed, the cause was determined. Then, the process was restarted from the 1st PCR. If the correct band was confirmed, the 3rd PCR was performed using the 2nd PCR product as a template. Here, if the concentration of the 2nd PCR product was high, DW was added to adjust the concentration. After completing the 3rd PCR, the sample was loaded onto a 2% agarose gel to identify the bands. The completed PCR products were pooled in equal amounts (5 μl each) and then subjected to PCR purification.

[0350] This experiment was performed using the GEL & PCR Purification System (GP104-200, Biofact). UB buffer was added to the PCR product in an amount equivalent to 5 times the volume of the PCR product and thorough mixing was performed. The column was prepared, 200 μl of HelpB buffer was added to the column, centrifugation was performed at 13,000 rpm for 30 seconds, and then the filtered solution was discarded. The reaction solution was added to the column, centrifugation was performed at 7,000 rpm for 1 minute, and then the filtered solution was discarded. 750 μl of 80% EtOH was added thereto, centrifugation was performed at 13,000 rpm for 30 seconds, and then the filtered solution was discarded. After repeating the process twice, centrifugation was performed at 13,000 rpm for 3 minutes. The centrifuged column was placed in a 1.5 ml

tube, 100 µl of EB buffer was added dropwise to the center, and the reaction was allowed to occur at room temperature for 1 minute. Centrifugation was performed at 13,000 rpm for 1 minute. The DNA collected in the 1.5 ml tube was quantified to obtain a concentration of 15 ng/µl and stored at 4° C. until NGS analysis.

TABLE 7

Reagent composition		PCR condition	
KAPA HiFi PCR mix	5 µℓ	Pre-denaturation	95° C., 3 min
Forward primer (10 pmol/ul)	0.5 µℓ	Denaturation (D)	98° C., 20 s
Reverse primer (10 pmol/ul)	0.5 µℓ	Annealing (A)	60° C., 15 s
Template (gDNA)	1 µℓ	Extension (E)	72° C., 2 min
DW	3 µℓ	D-E Cycle	30 cycles
Total	10 µℓ	Final extension	72° C., 3 min
		Storage	4° C., ∞

10. Preparation of Cell Extracts and Western Blot Analysis

[0351] Cell extracts were prepared by scraping cells with 60 to 80 µl of RIPA Lysis and Extraction Buffer (THERMOFISHER, 89900) containing protease inhibitor cocktail (ROCHE, 11836153001) at 4° C. and kept on ice for 10 minutes. The concentration of the cleared lysate was measured by Bradford assay (BIO-RAD), in which 20 to 65 µg of protein lysate, depending on the target protein, was loaded onto Mini-PROTEAN® TGX™ Precast Protein Gels (BIO-RAD), electrophoresed, and transferred to PVDF membranes using the Trans-Blot Turbo Transfer System (BIO-RAD). The membrane was incubated at room temperature for 1 hour in a blocking solution containing 5% non-fat dry milk in TBS-T and then incubated at 4° C. overnight with primary antibodies in TBS-T. After incubation with primary antibodies, the membrane was washed four times with TBS-T and then incubated at room temperature for 1 hour in a blocking solution containing anti-rabbit or anti-mouse immunoglobulin conjugated with HRP. The membrane was additionally washed four times with TBS-T and specific protein complexes were visualized with ECL prime (CYTIVA). Antibodies used were against KU70 (CST, 4588S), DNA ligase IV (CST, 146495), XLF (CST, 2854S), Artemis (CST, 13381), GAPDH (CST, 2118L), and XRCC4 (THERMO, MA5-24383).

11. Quantitative Real-Time PCR (qRT-PCR)

[0352] Purification of total RNA was performed, and reverse transcription of total 1 µg RNA was performed with Superscript IV reverse transcriptase (INVITROGEN) using random hexamers. For real-time PCR, 1/20 of the RT product was amplified with the KAPA SYBR FAST qPCR kit (KAPA BIOSYSTEM, Wilmington, MA, USA) using the real-time cycler Quantstudio (THERMO FISHER SCIEN-TIFIC). The relative amount of mRNA transcripts was measured by the 2ΔΔCT method. GAPDH and 18s rRNA were used as endogenous controls. The primer sequences used for qRT-PCR are shown in Table 8 below.

TABLE 8

Primer name	Sequence (5'-3')	SEQ ID NO
hATM-F	GGACAGTGGAGGCACAAAAT	451
hATM-R	GTGTCGAAGACAGCTGGTGA	452

TABLE 8-continued

Primer name	Sequence (5'-3')	SEQ ID NO
hKU70-F	CGGGAACAAATGAACCACT	453
hKU70-R	TGAAACCCATGAGCATCAAA	454
hLIG4-F	CACCTTGCCTTTTCCACGAA	455
hLIG4-R	CAGATGCCTTCCCCCTAAGT TG	456
hXRCC4-F	CTGATGGTCATTTCAGCATGG	457
hXRCC4-R	TCCTGCTCTGACAACAATG	458
hXLF-F	TCTCTGGCCTCCCCTTCTAT	459
hXLF-R	TAGCTCCCTCACTTGGCACT	460
hARTEMIS-F	AATTCCAAGTCGGGAGGAGT	461
hARTEMIS-R	GGATCTGAGTGTTCGGGTCT	462
GAPDH-F	GGAAGGACTCATGACCACAGT	463
GAPDH-R	CAGTGAGCTTCCCGTTTCAG	464
18s_rRNA-F	TCAACTTTCGATGGTAGTCG CC	465
18s_rRNA-R	GGCCTCGAAAGAGTCTCTGTA TTGT	466
mDclre1c-F	GCTTCGGGTGAGAAGGAAG	467
mDclre1c-R	TGCCAGTCTGAAGTCTCCTG	468

12. shRNA Cloning

[0353] shRNA sequences for the factors involved in the NHEJ repair pathway used in this experiment were shown in Table 9 below. To clone shRNA, annealing was performed for the target oligos for each shRNA. The annealed insert was inserted into a plasmid together with a U6 or H1 promoter.

TABLE 9

Species	Gene	#	5'-3'	SEQ ID NO
hUMAN	XRCC4	1	GUGAUGACACUGGCACAUUA	400
		2	GUGAGAGUACUGAUGAGGA	401
		3	GAAUCCACCUUUGUUUCUGA	402
		4	GCUACAAGUAUCUUGGGAGA	403
		5	GAAAUGCAGCUCAAGAACGA	404
	XLF	1	GCAUGAGUCUGGCAUUAACA	405
		2	GAAAGCCUUUGUCAUGAA	406
		3	AAACAGUCUUCUCCUGCAA	401
		4	GUGAAAGACCUAGAGAUCCA	408
		5	GAUAUGGCAGUCACCACACA	409
	XRCC6	1	GCAGCAUUGUGCAGAUACA	410
		2	GCAGGAACAUCUCCUCCUUA	411
		3	GUCAGUGCUCUGCUCAUCA	412
		4	GGAUCAUGCUGUUCACCAA	413
		5	GGAUCUGACUACUCACUCA	414
	LIG4	1	GCACAAAGAUUGAGAUUA	415
		2	GCCAGACACGUACUGUGUAA	416
		3	GCGAGCAGACUCCUCAAGAA	417
		4	AGAGGAUUCUGAUCUGCAA	418
		5	GCAUGAUCCUUCUGUAGGA	419

TABLE 9--continued

Species	Gene	#	5' - 3'	SEQ ID NO
	DCLRE1C	1	GCGAGACUCCUACCCAGAU	420
		2	GGACAAAGCUGACUACAGA	421
		3	GCGAGCUCUCCUUUCACA	422
		4	GUGACUCUGAUGGAGAAUCA	423
		5	GCCAGAAUUUUCUCCAGUCA	424
	ATM1	1	GGAGCCAGAUAGUUUGUAU	425
		2	GCAAGCAGCUGAAACAAU	426
		3	GGAGCUGAUUGUAGCAACA	427
		4	GCACAGAAGUGCCUCCAAU	428
		5	GGACAUAUGUUUCUGGGAGA	429
		6	GAACUUCAGUGGACCUUCA	430
	Scrambled	1	GUUCCCGAAGCUGUCACGU	431
		2	GCAACAGAUGAAGAGCACC AA	432
		3	CAGAGCUAACUCAGAUAGUA CU	446
Mouse	Scrambled	1	CCUAAGGUUAAGUCGCCUCG	469
	Dclre1c	1	GAAGACCAGUGCUAUAUCUA	433
		2	GGACCUUGAUGUGCUAACA	434
		3	GCACCGCAAGCCUUAUA	435
		4	GAAACUGGCAACAGGUCAA	436
		5	GGAUCAUGAGGGAGUAAGA	437
		6	GAGUGUUAGACUAUUA	438
		7	GGACAGUGCACCUGACUAA	439
		8	GAAUGUAUGGGUACUGUCA	440
		9	GAGGUGUAAGGGUCUGAA	441
		10	GAAUGGAGCUUCUGCACUCU	443
		11	GCCUAUGGCUACGAGUAUUU	443
		12	GAGCUGGGAGUUUCAGGUUCA	444
		13	GCAUCAAGCCAUUACCAU	445

[0354] The DCLRE1C mRNA levels are shown in FIG. 5 when the U1 and H1 promoters were used together with shRNA for DCLRE1C. From the results, it can be seen that each of the U1 and H1 promoters is a suitable promoter for use with the shRNA of the present disclosure.

13. Quantification of Large-Scale Deletion by qPCR (Dystrophin Exon 51, E51)

[0355] Quantitative real-time PCR was used to quantify genomic DNA. Primer sequences were designed for each of the E51 internal regions where large-scale deletion occurs. For qPCR, 20 ng of gDNA was amplified with the KAPA SYBR FAST qPCR kit (KAPA BIOSYSTEM, Wilmington, MA, USA) using the real-time cyclers Quantstudio (THERMO FISHER SCIENTIFIC). The primer sequences used for qPCR are shown in Table 10 below.

TABLE 10

Primer name	Sequence (5'-3')	SEQ ID NO
E51_LD F	TGTCATGAATAAGAGTTGGCTCA	447
E51_LD R	AGGTTGTGTACCAGAGTAACA	448
18s_rRNA-F	TCAACTTTCGATGGTAGTCGCC	449
18s_rRNA-R	GGCCTCGAAGAGTCCTGTATTGT	450

14. Whole Genome Sequencing (WGS) Analysis (E51)

[0356] HEK293T cells were transfected with UnCas12f1, a pair of gRNA (F142/R52), and shRNA, and then harvested 5 days after transfection. Genomic DNA was purified using a gDNA prep kit (Maxwell® RSC Cultured Cells DNA, PROMEGA, AS1620), 400 to 500 ng of genomic DNA was fragmented, and libraries were prepared using the MGIEasy FS DNA Library Prep. Kit (MGI Tech) according to the manufacturer's instructions. The libraries were then subjected to WGS using DNBSEQ-T7 (MGI Tech). Sequence reads were mapped to the human genome reference standard, GRCh38. Large-scale deletion efficiency was analyzed by counting reads mapped to the deleted region between a pair of gRNAs, and counts per million (CPM) was calculated. The percentage of deletion was calculated based on the difference in CPM between the sample and the control.

Example 2. Construction of Nucleic Acid Editing System for Deletion of Target Gene (Dystrophin Exon 51)

Example 2.1. Production of Engineered gRNA

[0357] The most common type among patients with Duchenne muscular dystrophy (DMD) is a type in which a stop codon occurs in dystrophin exon 51. As shown in FIG. 2, deletion of exons 49 and 50 leads to production of a stop codon in exon 51, which serves as a signal to stop protein synthesis, thereby preventing production of dystrophin protein. Here, deletion of exon 51 prevents production of the stop codon, thereby allowing for production of a dystrophin protein that is shorter than normal but has normal function.

[0358] The CRISPR/Cas 12f1 system and the TaRGET system were constructed for deletion of dystrophin exon 51. In the systems, for the gRNAs having a guide sequence that hybridizes with a target sequence for deletion of exon 51, engineered gRNAs having at least one of the five modification sites (MS1, MS2, MS3, MS4, and MS5) as shown in FIG. 3 were produced, and the specific sequences thereof are shown in Table 11.

TABLE 11

gRNA	Sequence (5' to 3')	SEQ ID NO
Canonical sgRNA	CUUCACUGAUAAGUGGAGAACCGCUACCAAAAGCUG UCCCUuagGGGAUUAGAACUUGAGUGAAGGUGGGCUGC GCAUCAGCCUAAUGUCGAGAAUGUCUUUCUUCGGAAAG UAACCCUCGAAACAAAUCAUUUUUCCUCUCCAAUUCUG CACAAGaaaGUUCGAGAACCGCUUAGaGacgaaUGAAGGAA UGCAACNNNNNNNNNNNNNNNNNNNNNNNN	13
MS1	CUUCACUGAUAAGUGGAGAACCGCUACCAAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAUGUCUUUCUUCGGAA	149

TABLE 11-continued

gRNA	Sequence (5' to 3')	SEQ ID NO
	AGUAACCCUCGAAACAAAUUCAGUGCUCUCUCCAAUUC UGCACAAGaaaGUUGCAGAACCCGAAUAGAGCAAUGAAGG AAUGCAACNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	
MS1/MS2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAAGUCUUUUCUCGGAA AGUAACCCUCGAAACAAAUUCAGUGCUCUCUCCAAUUC UGCACAAGaaaGUUGCAGAACCCGAAUAGAGCAAUGAAGG AAUGCAACNNNNNNNNNNNNNNNNNNNNNNUUUUAUUUUU U	150
MS1/MS2/MS (ge3.0)	ACCGCUUCACCAAAGCUGUCCUUAGGGGAUUAGAACU UGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGA GAAGUGCUUUUCUCGGAAAGUAACCCUCGAAACAAAUUC AGUGCUCUCUCCAAUUCUGCACAAGaaaGUUGCAGAACCC CGAAUAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNN NNNNNNNNUUUUAUUUUUU	151
MS2/MS3/MS (ge4.0)	ACCGCUUCACCAAAGCUGUCCUUAGGGGAUUAGAACU UGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGA GAAGUGCUUUUCUCGGAAAGUAACCCUCGAAACAAAGaaa GGAAUGCAACNNNNNNNNNNNNNNNNNNNNNNUUUUAUUU UUU	152
MS2/MS3/MS4 /MS5 (ge4.1)	ACCGCUUCACUUAGAGUGAAGGUGGGCUGCUUGCAUCAG CCUAAUGUCGAGAAGUGCUUUUCUCGGAAAGUAACCCUC GAAACAAAGaaaGGAAUGCAACNNNNNNNNNNNNNNNNNN NNUUUUAUUUUUU	153
MS1/MS3-1	GAUAAAGUGGAGAACCGCUUCACCAAAGCUGUCCCUUA GGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUC AGCCUAAUGUCGAGAAGUGCUUUUCUCGGAAAGUAACCC UCGAAACAAAUUCAGUGCUCUCUCCAAUUCUGCACAAG aaaGUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUGCAAC NNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	154
MS1/MS3-2	UGGAGAACCGCUUCACCAAAGCUGUCCUUAGGGGAUU AGAAUCUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAA UGUCGAGAAGUGCUUUUCUCGGAAAGUAACCCUCGAAAC AAAUUCAGUGCUCUCUCCAAUUCUGCACAAGaaaGUUGC AGAAACCCGAAUAGAGCAAUGAAGGAAUGCAACNNNNNN NNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	155
MS1/MS3-3	ACCGCUUCACCAAAGCUGUCCUUAGGGGAUUAGAACU UGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGA GAAGUGCUUUUCUCGGAAAGUAACCCUCGAAACAAAUUC AGUGCUCUCUCCAAUUCUGCACAAGaaaGUUGCAGAACCC CGAAUAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNN NNNNNNNNNN	156
MS1/MS4*-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUCGGAA AGUAACCCUCGAAACAAAUUCAGUGCUCUCUCCAAUUC gaaaGAACCCGAAUAGAGCAAUGAAGGAAUGCAACNNNNNN NNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	157
MS1/MS4*-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUCGGAA AGUAACCCUCGAAACAAAUUCAGUGCUCUCUCCgaaaGAA UAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNN NNNN	158
MS1/MS4*-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUCGGAA AGUAACCCUCGAAACAAAUUCAGUGCgaaaAGCAAUGAA GGAAUGCAACNNNNNNNNNNNNNNNNNNNNNNNNNNNNNN	159
MS1/MS5-1	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UuuagAUUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCA GCCUAAUGUCGAGAAGUGCUUUUCUCGGAAAGUAACCCU	160

TABLE 11-continued

gRNA	Sequence (5' to 3')	SEQ ID NO
	CGAAACAAAUUCAGUGCUCUCUCCAAUUCGCACAAGaa aGUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUGCAACN NNNNNNNNNNNNNNNNNNNN	
MS1/MS5-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCuua gGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAU GUCGAGAAGUGCUUUCUUCGGAAGUAACCCUCGAAACA AAUUCAGUGCUCUCUCCAAUUCGCACAAGaaGUUGC GAACCCGAAUAGAGCAAUGAAGGAAUGCAACNNNNNNNN NNNNNNNNNNNNNNNNNNNN	161
MS1/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUCAG UGCUCUCUCCAAUUCGCACAAGaaGUUGCAGAACCCG AAUAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNN	162
MS1/MS2/ MS4*-2	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAAGCUG UCCCUUAGGGGAUUAGAACUUGAGUGAAGGUGGGCUGC UUGCAUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGA AGUAACCCUCGAAACAAUUCAGUGCUCUCUcgaagAA UAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNN NNNNUUUUAUUUU	163
MS1/MS3- 3/MS4*-2	ACCGCUUCACCAAAGCUGUCCUAGGGGAUUAGAACU UGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGA GAAGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUC AGUGCUCUCUcgaagAAUAGAGCAAUGAAGGAAUGCAA CNNNNNNNNNNNNNNNNNNNNNN	164
MS1/MS2/ MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUCAG UGCUCUCUCCAAUUCGCACAAGaaGUUGCAGAACCCG AAUAGAGCAAUGAAGGAAUGCAACNNNNNNNNNNNNNN NNNNNNUUUUAUUUU	165
MS1/MS3- 3/MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUA ACCCUCGAAACAAUUCAGUGCUCUCUCCAAUUCUGCA CAAGaaGUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUG CAACNNNNNNNNNNNNNNNNNNNNNN	166
MS1/MS4*- 2/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUCAG UGCUCUCUcgaagAAUAGAGCAAUGAAGGAAUGCAACN NNNNNNNNNNNNNNNNNNNNNN	167
MS1/MS2/MS3 -3/MS4*-2	ACCGCUUCACCAAAGCUGUCCuuagGGGAUUAGAACUU GAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAG AAGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUC GUGCUCUCUcgaagAAUAGAGCAAUGAAGGAAUGCAAC NNNNNNNNNNNNNNNNNNNNUUUUAUUUU	168
MS1/MS2/MS3 -3/MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUA ACCCUCGAAACAAUUCAGUGCUCUCUCCAAUUCUGCA CAAGaaGUUGCAGAACCCGAAUAGAGCAAUGAAGGAAUG CAACNNNNNNNNNNNNNNNNNNNNUUUUAUUUU	169
MS1/MS2/MS4 -2/MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAUUCAG UGCUCUCUcgaagAAUAGAGCAAUGAAGGAAUGCAACN NNNNNNNNNNNNNNNNNNNNUUUUAUUUU	170
MS1/MS3- 3/MS4*- 2/MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUA ACCCUCGAAACAAUUCAGUGCUCUCUCCAAUUCUGA GCAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNNNNNN NNNNNNNNNNNNNNNNNNNNNN	171

TABLE 11-continued

gRNA	Sequence (5' to 3')	SEQ ID NO
MS1/MS2/MS3-3/MS4*-2/MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUUCUUCGGAAAGUA ACCCUCGAAACAAAUUCAGUGCUCUUCGaaaGAAUAGA GCAAUGAAGGAUGCAACNNNNNNNNNNNNNNNNNNNN UUUUUUUUU	172

[0359] In addition, mature form gRNAs were produced by removing the modification site MS1 from the canonical gRNA, and the specific sequences thereof are shown in Table 12.

TABLE 12

gRNA	Sequence (5' to 3')	SEQ ID NO
Mature form gRNA	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUG UCCCuagGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAG UAACCCUCGAAACAAAUUCAUUUgaaaGAAUGAAGGAUG CAACNNNNNNNNNNNNNNNNNNNN	173
MS3-1	GAUAAAGUGGAGAACCGCUUCACCAAAGCUGUCCCuag GGGAUUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCA GCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAGUAACCCU CGAAACAAAUUCAUUUgaaaGAAUGAAGGAUGCAACNNN NNNNNNNNNNNNNNNNNNNN	174
MS3-2	UGGAGAACCGCUUCACCAAAGCUGUCCCuagGGGAUUA GAACUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAU GUCGAGAAGUGCUUUUCUUCGAAAGUAACCCUCGAAACA AAUUCAUUUgaaaGAAUGAAGGAUGCAACNNNNNNNNNN NNNNNNNNNN	175
MS3-3	ACCGCUUCACCAAAGCUGUCCCuagGGGAUUAGAACUU GAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAG AAGUGCUUUUCUUCGAAAGUAACCCUCGAAACAAAUUCA UUUgaaaGAAUGAAGGAUGCAACNNNNNNNNNNNNNN NNNNNN	176
MS4-1	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUG UCCCuagGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAG UAACCCUCGAAACAAAUUCAUgaaaGAAUGAAGGAUGCAAC NNNNNNNNNNNNNNNNNNNN	177
MS4-2	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUG UCCCuagGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAG UAACCCUCGAAACAAAUUCgaaaGAAGGAUGCAACNNNN NNNNNNNNNNNNNNNNNN	178
MS4-3	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUG UCCCuagGGGAUUAGAACUUGAGUGAAGGUGGGCUGCUU GCAUCAGCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAG UAACCCUCGAAACAAAgaaaGGAUGCAACNNNNNNNNNN NNNNNNNNNN	179
MS5-1	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUG UuuagAUUAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCA GCCUAAUGUCGAGAAGUGCUUUUCUUCGAAAGUAACCCU CGAAACAAAUUCAUUUgaaaGAAUGAAGGAUGCAACNNN NNNNNNNNNNNNNNNNNN	180
MS5-2	CUUCACUGAUAAGUGGAGAACCGCUUCACCAAAGCUu uagAGAACUUGAGUGAAGGUGGGCUGCUUGCAUCAGCCU AAUGUCGAGAAGUGCUUUUCUUCGAAAGUAACCCUCGA AACAAAUUCAUUUgaaaGAAUGAAGGAUGCAACNNNNNN NNNNNNNNNNNNNNNNNN	181

TABLE 12-continued

gRNA	Sequence (5' to 3')	SEQ ID NO
MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAAUUCAU UUgaaaGAAUGAAGGAAUGCAACNNNNNNNNNNNNNNNN NNNN	182
MS3-3/ MS4-3	ACCGCUUCACCAAAGCUGUCCuuagGGGAUUAGAACUU GAGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAG AAGUGCUUUCUUCGGAAGUAACCCUCGAAACAAgaaaG GAAUGCAACNNNNNNNNNNNNNNNNNNNNNNNN	183
MS3-3/ MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUA ACCCUCGAAACAAAUUCAUUUgaaaGAAUGAAGGAUGCA ACNNNNNNNNNNNNNNNNNNNNNNNN	184
MS4-3/ MS5-3	CUUCACUGAUAAAGUGGAGAACCGCUUCACCAAuuagUUG AGUGAAGGUGGGCUGCUUGCAUCAGCCUAAUGUCGAGA AGUGCUUUCUUCGGAAGUAACCCUCGAAACAAgaaaGG AAUGCAACNNNNNNNNNNNNNNNNNNNNNNNN	185
MS3-3/MS4- 3/MS5-3	ACCGCUUCACCAAuuagUUGAGUGAAGGUGGGCUGCUUGC AUCAGCCUAAUGUCGAGAAGUGCUUUCUUCGGAAGUA ACCCUCGAAACAAgaaaGGAUGCAACNNNNNNNNNNNNNN NNNNNNNN	186

[0360] The sequence indicated by NNNNNNNNNNNNNNNNNNNNN in Tables 11 and 12 refers to any guide sequence (spacer sequence) that can hybridize with the target sequence. The guide sequence may be appropriately designed by those skilled in the art according to a desired target gene and/or a target sequence, and is not limited to a specific sequence of a particular length.

Example 2.2. Cas12f1 and TaRGET Systems

[0361] UnCas12f1 and CWCas21f1 were used together with the guide RNA of Example 2.1. PCR amplification was performed using the human codon-optimized nucleotide sequence (SEQ ID NOS: 10 and 6) of the protein as a template, and cloning was performed, according to the desired cloning sequence, into a vector having a promoter capable of expression in a eukaryotic system and a poly(A) signal sequence using the Gibson assembly method. After cloning, the sequence of the obtained recombinant plasmid vector was finally identified by the Sanger sequencing method. The nucleic acid construct thus produced was cloned into the pMAL-c2 plasmid vector, and transformed into BL21(DE3) *E. coli* cells. The transformed *E. coli* colonies were grown in LB broth at 37° C. until the optical density reached 0.7. The transformed *E. coli* cells were cultured at 18° C. overnight in the presence of 0.1 mM isopropylthio- α -D-galactoside. Thereafter, the cultured cells were collected by centrifugation at 3,500 g for 30 minutes, and the collected cells were resuspended in a buffer containing 20 mM Tris-HCl (pH 7.6), 500 mM NaCl, 5 mM P-mercaptoethanol, and 5% glycerol. The cells were lysed in a lysis buffer and then disrupted by sonication. The sample containing the disrupted cells was centrifuged at 15,000 g for 30 minutes, and the supernatant obtained was filtered through a 0.45 μ m syringe filter (Millipore). The filtered supernatant was loaded onto a Ni²⁺-affinity column using an FPLC purification system (KTA Purifier, GE Healthcare). The bound fractions were eluted with a gradient of 80-400 mM imidazole, 20 mM Tris-HCl (pH 7.5).

[0362] The eluted proteins were cleaved by treatment with TEV protease for 16 hours. The cleaved proteins were purified on a heparin column with a linear gradient of 0.15-1.6 M NaCl. The recombinant Cas12f1 variant protein purified on the heparin column was dialyzed against a solution of 20 mM Tris (pH 7.6), 150 mM NaCl, 5 mM P-mercaptoethanol, and 5% glycerol. The dialyzed protein was purified by passing it through an MBP column, and then repurified on a monoS column (GE Healthcare) or EnrichS with a linear gradient of 0.5-1.2 M NaCl.

[0363] The repurified proteins were collected and dialyzed against a solution of 20 mM Tris (pH 7.6), 150 mM NaCl, 5 mM β -mercaptoethanol, and 5% glycerol to purify the hypercompact gene editing protein (miniature endonuclease) used in the present disclosure. The concentration of the produced hypercompact gene editing protein was quantified by the Bradford quantitative method using bovine serum albumin (BSA) as a standard and measured electrophoretically on a coomassie blue-stained SDS-PAGE gel.

Comparative Example 1. SaCas9 System

[0364] The guide sequence was cloned into plasmids containing U6 promoter and SaCas9 scaffold, respectively. Then, the sequence from each U6 promoter to the guide RNA was inserted into the plasmid encoding SaCas9 to produce a one-vector module. Information on the target sequences is shown in Table 13.

TABLE 13

Target	PAM	Sequence	SEQ ID NO
F68	CTGAAT	GTGTATTGCTTGCTACTACTCA	471
R84	GAGAGT	GTGTATTACTTGCTACTGCA	472

Example 2.3. Selection of Target Sequences (E51)

[0365] The regions 2000 bp upstream and 2000 bp downstream of exon 51 were set as target regions for deletion of exon 51, and the target regions are referred to as the front region (F region) and the rear region (R region), respectively. Various protospacer sequences were selected from the above regions, guide RNAs were designed based thereon, and analysis of indel and deletion efficiency was performed. As a result, the sequences shown in Table 14 were confirmed to have high indel and deletion efficiency, and were selected as target sequences to perform subsequent experiments.

TABLE 14

Target name	PAM	Target seq	SEQ ID NO
R52	TTTA	CTCTCCTAGACCATTTCCCA	218
F142	TTTG	CTCATTCTCATGCCTGGACA	213

Example 3. qPCR Analysis on shRNA-Mediated Inhibition of Expression of NHEJ Component Gene

[0366] 5 µg of plasmid vector encoding shRNA was prepared using All-in-one vector (Biofact), transfected into AC16 cells, and harvested 3 days after transfection. Then, mRNA expression of each NHEJ component gene was measured by qRT-PCR. As a result, XRCC4-3, XLF-1, XRCC6-1, LIG4-5, DCLRE1C-3, and ATM-3 were selected as the most effective shRNAs. The results are shown in FIG. 6. The sequences of sense strands of the shRNAs used are described in Table 9.

Example 4. Western Blot Analysis on shRNA-Mediated Inhibition of Expression of NHEJ Component Gene

[0367] shRNA-mediated inhibition of gene expression was analyzed by Western blot analysis. The experimental conditions were the same as in Example 3. The results are shown in FIG. 7. From the results, it was possible to select the best shRNAs for XRCC4-3, XLF-1, XRCC6-1, LIG4-5, DCLRE1C-3, and ATM-3 and it can be seen that these shRNAs effectively reduced expression of the target genes. As a result, the Western blot results were consistent with the qRT-PCR results.

Example 5. Confirmation of Increased Deletion Efficiency Caused by UnCas12f1 Following shRNA Treatment (E51)

[0368] The most effective shRNA module for each target identified in Examples 3 and 4 was cloned into the UnCas12f1 expression vector, and the shRNA was designed to be expressed under the U6 promoter. A pair of guide RNAs for the targets F142 and R52 as shown in Table 14 was used to induce E51 deletion. In order to accurately find the optimal shRNA for a large-scale deletion strategy, comparison of the deletion efficiency altered by the selected shRNAs was conducted in HEK 293T cells. The E51 deletion efficiency was measured on an agarose gel in which the PCR-amplified E51 region was decomposed. The results are shown in FIG. 8. From the results, it can be seen that treatment with shRNA increases the E51 deletion efficiency caused by UnCas12f1 .

Example 6. Quantification of Deletion Efficiency (E51)

[0369] To compare the large-scale deletion efficiency between various treatments with shRNA, intensity analysis was conducted on the deletion band (Del) and the non-deletion band (WT) in FIG. 8, and the relative values were calculated using Image J software. The results are shown in FIG. 9. From the results, it can be seen that shRNA for the DCLRE1C gene is most effective in improving the large-scale deletion efficiency.

Example 7. Confirmation of Optimal shRNA for Deletion (E51)

Example 7.1. Analysis of Deletion Efficiency Following Treatment with shRNADCLRE1C

[0370] The efficiency of large-scale deletion was analyzed by qPCR 3 days after treatment of HEK293T cells with the selected shRNA (shDCLRE1C-2, -3, or -5)+UnCas12f1 and a pair of gRNA (F142/R52), and the results are shown in FIG. 10A. In addition, the DCLRE1C mRNA levels were measured under the same conditions, and the results are shown in FIG. 10B.

[0371] As can be seen from FIG. 10A, the sample treated with shDCLRE1C-3 showed the highest deletion level compared to the remaining shRNAs. These results are thought to stem from the fact that shDCLRE1C-3 most effectively downregulates DCLRE1C expression compared to the remaining shRNAs (see FIG. 10B).

Example 7.2. Analysis of Deletion Efficiency Following Treatment with shRNADCLRE1C and shXRCC6

[0372] For shDCLRE1C and shXRCC6 among the shRNAs selected in Example 4, experiments were conducted in the same manner as in Example 7.1 using the two shRNAs alone, in combination, or one shRNA thereof in two copies, to analyze the efficiency of large-scale deletion of E51. The results are shown in FIG. 23.

[0373] As can be seen from FIG. 23, shDCLRE1C and shXRCC6 resulted in high deletion of E51 when used alone, in combination, or in two copies. In particular, the highest deletion was observed when shDCLRE1C was used in two copies, and very high deletion was also observed when shDCLRE1C and shXRCC6 were used in combination.

Example 8. Whole Genome Sequencing (WGS) Analysis for Measurement of Deletion Efficiency (E51)

[0374] HEK293T cells were transfected with UnCas12f1, a pair of gRNA (F142/R52), and shRNA, and then harvested 5 days after transfection. gDNA was prepared therefrom for WGS that is performed to verify large-scale deletion efficiency. To compare the degree of deletion between the two gRNAs which is induced at both sides of E51, reads were mapped to the reference standard sequence and counting was performed within the deleted region. The data were presented as CPM (Count Per Million) along with each read count. The results are shown in FIG. 11. From the results, it can be seen that in a case of being treated with shDCLRE1C-3 in two copies (shDC-3×2), the mapped reads in the deleted region decreased, which indicates increased deletion of E51.

Example 9. Confirmation of Effect of shDCLRE1C
on Increased Deletion Efficiency Through
Complementation of DCLRE1C Expression (E51)

[0375] To confirm the effect of functional Artemis protein on large-scale deletion efficiency caused by UnCas12f1, DCLRE1C cDNA was co-transfected with shDCLRE1C-3 into HEK293T and AC16 cells. The results are shown in FIG. 12. From the results, it can be seen that the increased large-scale deletion caused by shDCLRE1C was nullified by restoration of DCLRE1C expression.

Example 10. Increased Deletion Efficiency in
DCLRE1C-KO Cell Line (E51)

[0376] To confirm the effect of DCLRE1C gene on large-scale deletion caused by UnCas12f1, DCLRE1C-KO cell lines of HEK293T cells were generated using a CRISPR-Cas12f1 system. Then, deletion of E51 induced by UnCas12f1 and a pair of gRNA (F142/R52) was analyzed by qPCR at the deleted locus. The results are shown in FIG. 13. From the results, it can be seen that increased large-scale deletion was observed in all KO cell lines compared to the wild type. Therefore, it is thought that the loss of function of the Artemis protein in KO cells resulted in increased large-scale deletion.

Example 11. Selection of shRNA for Inhibited
Expression of Murine Dclre1c

[0377] shRNAs for the Dclre1c gene encoding mouse Artemis protein were selected in Hepa-1c1c7 cells. The

shRNAs, which are expressed under the U6 promoter, were generated, and qRT-PCR was performed to measure of the Dclre1c mRNA levels. The sequences of the shRNAs used here are listed in Table 9. The results are shown in FIG. 14. From the results, the greatest decrease in the Dclre1c mRNA level was observed in cells treated with shDclre1c-12. The thus selected shRNAs can be used for in vivo studies using mouse models (for example, DMD mouse model with endogenous murine Dclre1c gene).

Example 12. Comparison of Deletion of Exon 51
Caused by Various Gene Editing Systems
Comprising shRNA

Example 12.1. Selection of shRNA

[0378] Candidate shRNAs for six genes known to be involved in the NHEJ repair pathway were transfected into previously prepared AC16 cells at a dose of 5 ug, and incubation was performed for 3 days. Then, the cells were harvested and qRT-PCR was used to measure the mRNA expression levels of the genes. The results are shown in FIG. 15 (ATM1 and XRCC4), FIG. 16 (XLF-1 and XRCC6), and FIG. 17 (LIG4 and DCLRE1C). From the results, the optimal shRNAs were selected. The candidate shRNAs used for selection are shown in Table 15. Scrambled indicates the control shRNA.

TABLE 15

Gene	No.	shRNA	SEQ ID NO
ATM1	1	GGAGCCAGAUAGUUUGUAUUUUAAGAGAAUACA AACUAUCUGGCUCC	360
	2	GCAAGCAGCUGAAACAAUUUUAAGAGAAUUUG UUUCAGCUGCUUGC	361
	3	GGAGCUGAUUGUAGCAACAUUUAAGAGAUUUUG CUACAAUCAGCUCC	362
	4	GCACAGAAGUGCCUCCAAUUUUAAGAGAAUUGGA GGCACUUCUGUGC	363
	5	GGACAUAGUUUCUGGGAGAUUUAAGAGAUUCCCC AGAAACUAUGUCC	364
	6	GAACUUCAGUGGACCUUUAUUUAAGAGAUGAAG GUCCACUGAAGUUC	473
XRCC4	1	GGAUGACACUGGCACAUUAUUAAGAGAUAAUG UGCCAGUGUCAUCC	365
	2	GGAGAGUACUGAUGAGGAAUUAAGAGAUUCCU CAUCAGUACUCUCC	366
	3	GAAUCCACCUUGUUUCUGAUUUAAGAGAUCAAGAA ACAAGGUGGAUUC	367
	4	GUACAAGUAUCUUGGGAGAUUUAAGAGAUUCCCC AAGAUACUUGUAC	368
	5	GAAUGCAGCUCAAGAACGAUUAAGAGAUUGUUC UUGAGCUGCAUUC	369
XLF-1	1	GCAUGAGUCUGGCAUUAUUAAGAGAUUGAA UGCCAGACUCAUGC	370
	2	GAAAGCCCUUUGUCAUGAAUUAAGAGAUUCAU GACAAAGGGCUUUC	371
	3	GAACAGUGCUUCCUGCAAUUUAAGAGAUUGCAG GGAAGCACUGUUC	372
	4	GGAAAGACCUAGAGAUUCAUUAAGAGAUUGGAU CUCUAGGUCUUUCC	373
	5	GUUAGGCAGUACCAACACAUUAAGAGAUUGUGUG GUGACUGCCAUAC	374

TABLE 15-continued

Gene	No.	shRNA	SEQ ID NO
XRCC6	1	GCAGCAUUGGCAGAUACAUAAGAGAUGUAU CUGCACAAUGCUGC	375
	2	GCAGGAACAUCUCCUCCUUAUUAAGAGAUAAAGGA GGGAUGUUCUGC	376
	3	GCAGUGCUCUGCUCACAAUUAAGAGAUGAUG AGCAGAGCACUGC	377
	4	GGAUCAUGCUGUUCACCAAUUAAGAGAUGGU GAACAGCAUGAUCC	378
	5	GGAUCUGACUACUCACUCAUUAAGAGAUGAGUG AGUAGUCAGAUCC	379
LIG4	1	GCACAAAGAUGGAGAUGUAUUAAGAGAUACAU CUCCAUCUUUGUGC	380
	2	GCAGACACGUCUGUGUAAUUAAGAGAUUACAC AGUACGUGUCUGC	381
	3	GGAGCAGACUCCUGAAGAAUUAAGAGAUUCUUC AGGAGUCUGCUC	382
	4	GGAGGAUUCUGAUCUGCAAUUAAGAGAUUGCA GAUCAGAAUCCUCC	383
	5	GCAUGAUCCUUCUGUAGGAUUAAGAGAUCCUAC AGAAGGAUCAUGC	384
DCLRE1C	1	GGAGACUCCUACCCAGAUUAUUAAGAGAUUUCUG GGUAGGAGUCUCC	385
	2	GGACAAAGCUGACUACAGAUUAAGAGAUUCUGU AGUCAGCUUUGUCC	386
	3	GCAGAGCUCUCGUUUCACAUUAAGAGAUUGGAA ACGAGAGCUCUGC	387
	4	GGACUCUGAUGGAGAAUCAUUAAGAGAUUAU CUCCAUCAGAUCC	388
	5	GCAGAAUUCUCCAGUCAUUAAGAGAUAGACUG GGAGAAUUCUGC	389
Scrambled	3	CAGAGCUAACUCAGAUAGUACU	470

Example 12.2. Deletion of Exon 51 by System Comprising shRNA

Measurement of Deletion Efficiency for Exon 51 in Case of Using Single shRNA

[0379] Based on the qRT-PCR results, for each gene, the shRNA that exhibited the highest inhibition efficiency for mRNA expression was selected. Nucleic acids encoding the shRNA, the two guide RNAs, and CWCas12f1 or Cas12f1 were inserted into a single vector, and the vector was transfected into AC16 cells and HEK293 cells. The experiment was independently repeated three times to measure the relative deletion efficiency for exon 51. Here, relative means a relative value under the same conditions, as indel efficiency varies depending on the transfection time, vector type, and concentration. The results are shown in FIG. 18 (HEK293 cells) and FIG. 19 (AC 16 cells).

[0380] Referring to FIGS. 18 and 19, it was confirmed that exon 51 was effectively deleted through inhibition of expression of NHEJ-related genes in all Cas12f1 systems, and in particular, higher deletion level of exon 51 was observed in a case of using shXRCC6 and shDCLRE1C.

Measurement of Deletion Efficiency for Exon 51 in Case of Using Two or More shRNAs

[0381] Deletion of exon 51 was induced using two or more identical or different shRNAs. Nucleotide sequences encoding the two guide RNAs, Cas12f1, and shDCLRE1C (one, two, or three selected from shDCLRE1C2, shDCLRE1C3, and shDCLRE1C5) were inserted into a single vector and the vector was transfected into HEK293 cells and AC16 cells using the same method as described above to measure

the relative deletion efficiency of exon 51. Here, the shDCLRE1Cs used were three that showed the highest mRNA expression inhibition efficiency among the five shRNAs (right one in FIG. 17). The results are shown in FIG. 20 (HEK293 cells) and FIG. 21 (AC16 cells).

[0382] Referring to FIGS. 20 and 21, deletion of exon 51 was successfully achieved in all experimental groups, and the deletion efficiency for exon 51 was particularly excellent in a case where two or more shDCLRE1Cs were introduced.

Example 12.3. Deletion of Exon 51 Over Time after Transfection with System Comprising shRNA

[0383] To determine deletion efficiency for exon 51 over different transfection periods (3 days, 5 days, and 7 days post transfection) using one or more shRNAs, nucleic acids encoding the two guide RNAs, TnpB or Cas12f1, and one or more shRNAs were inserted into a single vector, and then the deletion efficiency for exon 51 was determined at various time points starting from the day of transfection. In all experiments, AC16 cells were used, and the empty vector and the SaCas9 system were prepared as controls for comparison. The results are shown in FIG. 22.

[0384] As shown in FIG. 22, the deletion efficiency of exon 51 increased over time from the start of transfection in all experimental groups. In particular, 7 days after transfection, the deletion efficiency for exon 51 in the Cas12f1 and TaRGET systems that use a combination of two shDCLRE1Cs was similar to that of SaCas9.

CONCLUSION

[0385] As such, it was shown that inhibition of NHEJ pathway factors by shRNA can increase efficiency of large-scale deletion induced by UnCas12f1. It has been found that among the various shRNAs selected, interference of the DCLRE1C gene encoding the Artemis protein most effectively increases the efficiency of large-scale deletion. Large-scale deletion achieved by using Cas nuclease and a pair of

gRNA can be utilized as a gene editing approach for treatment of a genetic disease such as DMD, and low editing efficiency in vivo is likely to be one of the biggest obstacles in determining whether to proceed to the next stage. Therefore, it is expected that increased efficiency of large-scale deletion achieved by shRNA-mediated inhibition of NHEJ component gene expression will be widely applied in development of deletion-based gene editing therapies.

SEQUENCE LISTING

Sequence total quantity: 473

SEQ ID NO: 1 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = TnpB
 organism = unidentified

SEQUENCE: 1
 MGEKSSRRRR NGKSGAWTAA ITSCVGGKMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60
 NREKIALEKN KDKVKEACSK HLKVAAYCTT QVERNACLFC KARKLDDKFY QKLRGQFPDA 120
 VFWEISEIF RQLQKQAAEI YNQSLIELYY EIFIKGKGIA NASSVEHYLS DVCYTRAAEL 180
 FKNAAIASGL RSKIKSNFRL KELKNMKSGL PTTKSDNFPI PLVKQKGGQY TGFEISNHNS 240
 DFIKIPFGR WQVKKEIDKY RPWEKDFEFQ VQKSPKPISL LLSTQRRKRK KGWSKDEGTE 300
 AEIKKVMNGD YQTSYIEVKR GSKIGEKSAM MLNLSIDVPK IDKGVDPSII GGIDVGVKSP 360
 LVCAINNAFS RYSISDNDLF HFNKKMFARR RILKKNRHK RAGHGAKNKL KPITILTEKS 420
 ERFRKKLIER WACEIADFFI KNKVGTVQME NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
 EFKLKQYGIE IRKVAPNNTS KTCCKGHLN NYFNFEYRKK NKFPFKCEK CNFKENADYN 540
 AALNISNPKL KSTKEEP 557

SEQ ID NO: 2 moltype = AA length = 555
 FEATURE Location/Qualifiers
 source 1..555
 mol_type = protein
 note = TnpB-v1
 organism = synthetic construct

SEQUENCE: 2
 MEKRINKIRK KLSADNATKP VSRSGPMAKN TITKTLKLRI VRPYNSAEVE KIVADEKNNR 60
 EKIALEKNKD KVEACSKHL KVAAYCTTQV ERNACLFC KA RLDDKFYQK LRGQFPDAVF 120
 WQEISEIFRQ LQKQAAEIYN QSLIELYYEI FIKGKGIANA SSVHYLSDV CYTRAAELFK 180
 NAAIASGLRS KIKSNFRLKE LKNMKSGLPT TKSDNFPIPL VKQKGGQYTG FEISNHNSDF 240
 IIKIPFGRWQ VKKEIDKYRP WEKDFEFQVQ KSPKPISLLL STQRRKRKNG WSKDEGTEAE 300
 IKKVMNGDYQ TSYIEVKRGS KIGEKSAMWL NLSIDVPKID KGVDPISIGG IDVGVSPLV 360
 CAINNAFSRY SISDNDLFHF NKKMFARRRI LLKKNRHKRA GHGAKNKLKP ITILTEKSER 420
 FRKKLIERWA CEIADFFIKN KVGTVQMENL ESMKRKEDSY FNIRLRGFWP YAEMQNKIEF 480
 KLKQYGEIR KVAPNNTSKT CSKCGHLNLY FNFEYRKNK FPFKCEKCN FKENADYNAA 540
 LNISNPKLKS TKEEP 555

SEQ ID NO: 3 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = TnpB-v2
 organism = synthetic construct

SEQUENCE: 3
 MAGGPGAGSA APVSSSTSLP LAALNMRVMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60
 NREKIALEKN KDKVKEACSK HLKVAAYCTT QVERNACLFC KARKLDDKFY QKLRGQFPDA 120
 VFWEISEIF RQLQKQAAEI YNQSLIELYY EIFIKGKGIA NASSVEHYLS DVCYTRAAEL 180
 FKNAAIASGL RSKIKSNFRL KELKNMKSGL PTTKSDNFPI PLVKQKGGQY TGFEISNHNS 240
 DFIKIPFGR WQVKKEIDKY RPWEKDFEFQ VQKSPKPISL LLSTQRRKRK KGWSKDEGTE 300
 AEIKKVMNGD YQTSYIEVKR GSKIGEKSAM MLNLSIDVPK IDKGVDPSII GGIDVGVKSP 360
 LVCAINNAFS RYSISDNDLF HFNKKMFARR RILKKNRHK RAGHGAKNKL KPITILTEKS 420
 ERFRKKLIER WACEIADFFI KNKVGTVQME NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
 EFKLKQYGIE IRKVAPNNTS KTCCKGHLN NYFNFEYRKK NKFPFKCEK CNFKENADYN 540
 AALNISNPKL KSTKEEP 557

SEQ ID NO: 4 moltype = AA length = 555
 FEATURE Location/Qualifiers
 source 1..555
 mol_type = protein
 note = TnpB-v3
 organism = synthetic construct

SEQUENCE: 4
 MAGGPGAGSA APVSSSTSLP LAALNMAKN TITKTLKLRI VRPYNSAEVE KIVADEKNNR 60

-continued

EKIALEKNKD	KVKEACSKHL	KVAAYCTTQV	ERNACLFCKA	RKLDKIFYQK	LRGQFPDAVF	120
WQEISEIFRQ	LQKQAAEIYN	QSLIELYYEI	FIKKGIGIANA	SSVEHYLSDV	CYTRAAELFK	180
NAAIASGLRS	KIKSNFRLKE	LKNMKSGLPT	TKSDNFPIPL	VKQKGGQYTG	FEISNHNSDF	240
IIKIPFRWRQ	VKKEIDKYRP	WEKFDFEQVQ	KSPKPISLLL	STQRRKRKNG	WSKDEGTAE	300
IKKVMNGDYQ	TSYIEVKRGS	KIGEKSAWML	NLSIDVPKID	KGVDPSIIGG	IDVGVKSPVL	360
CAINNAFSRY	SISDNDLFHF	NKKMFARRRI	LLKKNRHKRA	GHGAKNKLKP	ITILTEKSER	420
FRKKLIERWA	CEIADFFIKN	KVGTVMENL	ESMKRKEDSY	FNIRLRGFWP	YAEMQNKIEF	480
KLKQYGIER	KVAPNNTSKT	CSKCGHLNNY	FNFEYRKKNK	PPHFKCEKCN	FKENADYNAA	540
LNISNPKLKS	TKEEP					555

SEQ ID NO: 5 moltype = AA length = 529
FEATURE Location/Qualifiers
source 1..529
 mol_type = protein
 note = Wild-type Cas12f1
 organism = synthetic construct

SEQUENCE: 5

MAKNTITKTL	KLRIVRPYNS	AEVEKIVADE	KNNREKIALE	KNKDKVKEAC	SKHLKVAAYC	60
TTQVERNACL	FCRKARKLDDK	FYQKLRGQFP	DAVFWQEIIE	IFRQLQKQAA	EIYNQSLIEL	120
YYEIFIKGKG	IANASSVEHY	LSDV CYTRAA	ELFKNAAIAS	GLRSKIKSNF	RLKELKNMKS	180
GLPTTKSDNF	PIPLVKQKGG	QYTGFEISNH	NSDFI IKIPF	GRWQVKKEID	KYRPWEKFDF	240
EQVQKSPKEI	SLLLSTQRRK	RNKGWSKDEG	TEAEIKVMN	GDYQTSYIEV	KRGSKIGESK	300
AWMLNLSIDV	PKIDKGVDP	IIGGIDVGK	SPLVCAINNA	FSRYSISDND	LFHFNKKMFA	360
RRRILLKKNR	HKRAGHGAKN	KLKPITILTE	KSERFRKKLI	ERWACEIADF	FIKNKVGTVQ	420
MENLESMKKR	EDSYFNIRLR	GFWPYAEMQN	KIEFKLKQYG	IEIRKVAPNN	TSKTCCKCGH	480
LNNYFNFEYR	KKNKFPHPKC	EKCNFKENAD	YNAALNISNP	KLKSTKEEP		529

SEQ ID NO: 6 moltype = DNA length = 1671
FEATURE Location/Qualifiers
source 1..1671
 mol_type = other DNA
 note = Human codon-optimized TnpB
 organism = synthetic construct

SEQUENCE: 6

atgggggaga	aaagtcccg	cgcgcgacgg	aatggaaaaa	gcgggtgcgtg	gactgctgct	60
ataacaagct	gtgttgggg	taagatggcc	aagaacacaa	ttacaaagac	actgaagctg	120
aggatcgtga	gaccatacaa	cagcgcgtgag	gtcgagaaga	ttgtggctga	tgaaaagaac	180
aacagggaaa	agatgcgcct	cgagaagaac	aaggataaag	tgaaggaggg	ctgctctaag	240
cacctgaaag	tggccgccta	ctgcaccaca	caggtggaga	ggaacgcctg	tctgttttgt	300
aaagctcgga	agctggatga	taagttttac	cagaagctgc	ggggccagtt	ccccgatgcc	360
gtcttttggc	aggagattag	cgagatcttc	agacagctgc	agaagcaggg	cgcgcgagtc	420
tacaaccaga	gcctgatcga	gctctactac	gagatcttca	tcaagggcaa	gggcattgct	480
aacgcctcct	ccgtggagca	ctacctgagc	gacgtgtgct	acacaagagc	cgccgagctc	540
tttaagaacg	ccgctatcgc	ttccgggctg	aggagcaaga	ttaagagtaa	cttccggctc	600
aaggagctga	agaacatgaa	gagcgcgcctg	cccactacaa	agagcgcacaa	cttcccaatt	660
ccactgggtga	agcagaaggg	gggccaagtac	acagggttcg	agattttcaa	ccacaacagc	720
gactttatta	ttaagatccc	ctttggcagg	tggcaggcca	agaaggagat	tgacaagtac	780
aggccctggg	acaagtttga	tttcgagcag	gtgcagaaga	gccccaaagg	tatttccctg	840
ctgctgtcca	cacagcggcg	gaagagggaac	aagggttggt	ctaaggatga	ggggaccgag	900
gccgagatta	agaaagtgat	gaacggcgac	taccagacaa	gctacatcga	ggtaacggcg	960
ggcagtaaga	tggcgcagaa	gagcgcctgg	atgctgaacc	tgagcattga	cgtgccaaaag	1020
attgataagg	ctgtggatcc	cagcatcatc	ggagggatcg	atgtgggggtg	caagagcccc	1080
ctcgtgtgcg	ccatcaacaa	cgccttcagc	agggtacaga	tctccgataa	cgacctgttc	1140
cactttaaca	agaagatggt	cgcccgggcg	aggattttgc	tcaagaagaa	ccggcacaag	1200
cgggcccggac	acggggccaa	gaacaagctc	aagcccatac	ctatcctgac	cgagaagagc	1260
gagagggttca	ggaagaagct	catcgagaga	tgggcctgcg	agatcgccga	tttctttatt	1320
aagaacaagg	tcggaacagt	gcagatggag	aacctcgaga	gcatgaagag	gaaggaggat	1380
tctactttca	acattcggct	gaggggggttc	tggccctacg	ctgagatgca	gaacaagatt	1440
gagtttaagc	tgaagcagta	cgggatttag	atccggaagg	tggcccccac	caacaccagc	1500
aagacctgca	gcaagtgcgg	gcacctcaac	aactacttca	acttcagtag	ccggaagaag	1560
aaacaagttcc	cacacttcaa	gtgcgagaag	tgcaacttta	aggagaacgc	cgattacaac	1620
gccgccctga	acatcagcaa	ccctaagctg	aagagcacta	aggaggagcc	c	1671

SEQ ID NO: 7 moltype = DNA length = 1666
FEATURE Location/Qualifiers
source 1..1666
 mol_type = other DNA
 note = Human codon-optimized TnpB-V1
 organism = synthetic construct

SEQUENCE: 7

atggaaaaaga	gaatcaacaa	gatcaggaag	aagctgagcg	ccgacaacgc	caccaagcct	60
gtgtctagga	gtggccccc	tggccaagaa	cacaattaca	aagacactga	agctgaggat	120
cgtgagacca	tacaacagcg	ctgaggtcga	gaagattgtg	gctgatgaaa	agaacaacag	180
ggaaaagatc	gccctcgaga	agaacaagga	taaggtgaag	gaggcctgct	ctaagcacct	240
gaaagtggcc	gcctactgca	ccacacaggt	ggagagggaac	gcctgtctgt	tttgtaagc	300
tcggaagctg	gatgataagt	tttaccagaa	gctgcggggc	cagttccccg	atgccgtctt	360

-continued

```

ttggcaggag attagecgaga tcttcagaca gctgcagaag caggccgccc agatctacaa 420
ccagagcctg atcgagctct actacgagat cttcatcaag ggcaaggcca ttgccaacgc 480
ctcctccgtg gagcactacc tgagcgacgt gtgctacaca agagccgccc agctctttaa 540
gaacgccgct atcgcttccg ggctgaggag caagattaag agtaacttcc ggctcaagga 600
gctgaagaac atgaagagcg gectgcccac tacaaagagc gacaacttcc caattccact 660
ggtagaagcag aagggggggc agtacacagg gtctcgagatt tccaaccaca acagcgactt 720
tattattaag atcccccttg gcaggtggca ggccaagaag gagattgaca agtacaggcc 780
ctgggagaag tttgatttcc agcaggtgca gaagagcccc aagcctattt cctgctgct 840
gtccacacag cggcggaaga ggaacaaggg gtggtctaag gatgagggga ccgaggccga 900
gattaagaaa gtgatgaacg gcgactacca gacaagctac atcgagggtc agcggggcag 960
taagattggc gagaagagcg cctggatgct gaacctgagc attgacgtgc caaagattga 1020
taagggcggt gatcccgaca tcacggaggg gatcgatgtg ggggtcaaga gccccctcgt 1080
gtgcgccatc aacaacgcct tcagcaggta cagcatctcc gataacgacc tgttccactt 1140
taacaagaag atgttcgccc ggcgaggat tttgctcaag aagaaccggc acaagcgggc 1200
cggcacaggc gccaaagaca agctcaagcc catcactatc ctgaccgaga agagcgagag 1260
gttcaggaag aagctcatcg agagatgggc ctgcgagatc gccgatttct ttattaagaa 1320
caaggtcgga acagtgcaga tggagaacct cgagagcatg aagagggaag aggatctcta 1380
cttcaacatt cggctgaggg ggttctggcc ctacgctgag atgcagaaca agattgagtt 1440
taagctgaag cagtacggga ttgagatccg gaaggtggcc cccaacaaca ccagcaagac 1500
ctgcagcaag tgcggggcacc tcacaacta cttcaacttc gagtaccgga agaagaacaa 1560
gttccccacac ttcaagtgcg agaagtgcaa ctttaaggag aacgccgatt acaacgccgc 1620
cctgaacatc agcaacccta agctgaagag cactaaggag gagccc 1666

```

```

SEQ ID NO: 8      moltype = DNA length = 1671
FEATURE
source            Location/Qualifiers
                  1..1671
                  mol_type = other DNA
                  note = Human codon-optimized TnpB-V2
                  organism = synthetic construct

```

```

SEQUENCE: 8
atggctggcg gaccaggcgc aggtagtgtc gcgccagttt cttcaacttc ctccctgccc 60
ctggctggcg ttaacatgcg cgtgatggcc aagaacacaa ttacaaagac actgaagctg 120
aggatcgtga gaccatacaa cagcgtgtag gtgcagaaga ttgtggctga tgaagaagac 180
aacagggaag gagatgcctt cgagaagaac aaggataagg tgaaggagcc ctgctctaag 240
cacctgaaag tggccgccta ctgcaccaca caggtggaga ggaacgcctg tctgttttgt 300
aaagctcgga agctggatga taagttttac cagaagctgc ggggccagtt ccccgatgcc 360
gtcttttggc agggagattag cgagatcttc agacagctgc agaagcagcc cgccgagatc 420
tacaaccaga gctgatcgca gctctactac gagatcttca tcaagggcaa gggcattgcc 480
aacgcctcct ccgtggagca ctacctgagc gacgtgtgct acacaagagc cgccgagctc 540
tttaagaacg ccgctatcgc ttccgggctg aggaagcaaa ttaagagtaa cttccggctc 600
aaggagctga agaacatgaa gagcggcctg cccactacaa agagcgacaa cttcccaatt 660
ccactggtga agcagaaggg gggccagtag acaggggttc agatttccaa ccacaacagc 720
gactttatta ttaagatccc ctttggcagg tggcagggtc agaaggagat tgacaagtag 780
agggcctggg agaagtttga tttcgagcag gtgcagaaga gccccaaagc tatttccctg 840
ctgctgtcca cacagcggcg gaagaggaac aaggggtggt ctaaggatga ggggaccgag 900
gccagagtta agaaagtgat gaacggcgac taccagacaa gctacatcga ggtcaagcgg 960
ggcagtaaga ttggcgagaa gagcgctcgt atgctgaacc tgagcattga cgtgccaaa 1020
attgataaag gcgtggatcc cagcatcatc ggagggatcg atgtgggggt caagagcccc 1080
ctcgtgtgcg ccatacaaca cgcttcagc aggtacagca tctccgataa cgacctgttc 1140
cactttaaca agaagatggt cgcccgccgg aggattttgc tcaagaagaa ccggcacaag 1200
cgggcccggc acggggccaa gaacaagctc aagcccatca ctatcctgac cgagaagagc 1260
gagaggttca ggaagaagct gcagagaga tgggcctcgc agatcgccga tttctttatt 1320
aagaacaagg tcggaacagt gcagatggag aacctcgaga gcatgaagag gaaggaggat 1380
tcctacttca acattcggct gaggggggtt tggccctacg ctgagatgca gaacaagatt 1440
gagtttaagc tgaagcagta cgggattgag atccggaagg tggcccccac caacaccagc 1500
aagacctgca gcaagtgcgg gcacctcaac aactacttca acttcagta ccggaagaag 1560
aacaagttcc cacacttcaa gtgcgagaag tgcaacttta aggagaacgc cgattacaac 1620
gccgccttga acatcagcaa ccctaagctg aagagcacta aggaggagcc c 1671

```

```

SEQ ID NO: 9      moltype = DNA length = 1665
FEATURE
source            Location/Qualifiers
                  1..1665
                  mol_type = other DNA
                  note = Human codon-optimized TnpB-V3
                  organism = synthetic construct

```

```

SEQUENCE: 9
atggctggcg gaccaggcgc aggtagtgtc gcgccagttt cttcaacttc ctccctgccc 60
ctggctggcg ttaacatgag ggccaagaac acaattacaa agacactgaa gctgaggatc 120
gtgagaccat acaacagcgc tgaggtcgag aagattgtgg ctgatgaaa gaacaacagg 180
aaaaagatcg cctcagagaa gaacaaggat aaggtgaagg aggcctgtct taagcacctg 240
aaagtggcgg cctactgac cacaagggtg gagaggaacg cctgtctgtt ttgtaagct 300
cggaaagctg atgataagtt ttaccagaag ctgcgggggc agttcccga tgcgtcttt 360
tggcaggaga ttacgagat cttcagacag ctgcagaagc aggcgcgcca gatctacaac 420
cagagcctga tcgagctcta ctacgagatc ttcataaagg gcaagggcat tgccaacgcc 480
tcctccgtgg agcaactacc gagcgacgtg tgctacacaa gagccgcga gctctttaag 540
aacgcgcgta tcgcttccgg gctgaggagc aagattaaga gtaacttccg gctcaaggag 600

```

-continued

```

ctgaagaaca tgaagagcgg cctgcccact acaaagagcg acaacttccc aattccactg 660
gtgaagcaga agggggggcca gtacacaggg ttcgagattt ccaaccacaa cagcgacttt 720
attattaaga tccccttttg caggtggcag gtcaagaagg agattgacaa gtacaggccc 780
tgggagaagt ttgatttcga gcaggtgcag aagagcccca agcctatttc cctgctgctg 840
tccacacagc ggcggaagag gaacaagggg tggctaaagg atgaggggac cgaggccgag 900
attaagaaag tgatgaacgg cgactaccag acaagctaca tcgagggtcaa gcggggcagt 960
aagattggcg agaagagcgc ctggatgctg aacctgagca ttgacgtgcc aaagattgat 1020
aagggcgtgg atcccagcat catcgaggag atcgatgtgg gggccaagag ccccctcgtg 1080
tgcgccatca acaacgcctt cagcagggtac agcatctccg ataacgacct gtccactttt 1140
aacaagaaga tgttcgcccc gcggaggatt ttgctcaaga agaaccggca caagcgggcc 1200
ggacacgggg ccaagaacaa gctcaagccc atcactatcc tgaccgagaa gagcgagagg 1260
ttcaggaaga agctcatcga gagatggggc tgcgagatcg ccgatttctt tattaagaac 1320
aaggtcggaa cagtgcagat ggagaacctc gagagcatga agaggaagga ggattcctac 1380
ttcaacattc ggctgagggg gttctggccc tacgctgaga tgcagaacaa gattgagttt 1440
aagctgaagc agtacgggat tgagatccgg aaggtggccc ccaacaacac cagcaagacc 1500
tgcagcaagt cggggcacct caacaactac ttcaacttcg agtaccggaa gaagaacaag 1560
tcccacact tcaagtgcga gaagtgcac tttaaggaga acgcccatta caacgcgcgc 1620
ctgaacatca gcaaccctaa gctgaagagc actaaggagg agccc 1665

```

```

SEQ ID NO: 10      moltype = DNA length = 1587
FEATURE           Location/Qualifiers
source            1..1587
                  mol_type = other DNA
                  note = Human codon-optimized Cas12f1
                  organism = synthetic construct

```

```

SEQUENCE: 10
atggccaaga acacaattac aaagacactg aagctgagga tcgtgagacc atacaacagc 60
gctgaggtcg agaagattgt ggctgatgaa aagaacaaca gggaaaagat cgccctcgag 120
aagaacaagg ataaggtgaa ggaggcctgc tctaagcacc tgaagtgagg cgctactgac 180
accacacagg tggagaggaa cgctgtctg ttttgtaaag ctcggaagct ggatgataag 240
ttttaccaga agctgcgggg ccagttcccc gatgccgtct tttggcagga gattagcgag 300
atcttcagac agctgcagaa gcaggccgcc gagatctaca accagagcct gatcgagctc 360
tactacgaga tcttcacaa gggcaagggc attgccaacg cctcctccgt ggagcactac 420
ctgagcgagc tgtgctacac aagagccgcc gagctcttta agaaccgccg tatcgcttcc 480
gggctgagga gcaagattaa gactaacctc cggctcaagg agctgaagaa catgaagagc 540
ggcctgcccc ctacaagag cgacaacttc ccaattccac tggtgaaagca gaaggggggc 600
cagtagacag ggttcagat ttccaaccac aacagcgact ttattattaa gatccctttt 660
ggcaggtggc aggtcaagaa ggagattgac aagtacaggg cctgggagaa gtttgatttc 720
gagcaggtgc agaagagccc caagcctatt tccctgctgc tgtccacaca gcggcggaag 780
aggaacaagg ggtggtctaa ggatgagggg accgagggcc agattaagaa agtgatgaac 840
ggcgactacc agacaagcta catcgaggtc aagcggggca gtaagattgg cgagaagagc 900
gcctggatgc tgaacctgag cattgacgtg ccaaagattg ataagggcgt ggatcccgag 960
atcatcggag ggcagatgat ggggggtcaag agccccctcg tgtgcgccat caacaacgcc 1020
ttcagcaggt acagcatctc cgataacgac ctgttccact ttaacaagaa gatgttcgcc 1080
cggcgaggga ttttgctcaa gaagaaccgg cacaagcggg ccggacacgg ggccaagaac 1140
aagctcaagc ccacacatct cctgaccgag aagagcgaga ggttcaggaa gaagctcctc 1200
gagagatggg cctgcgagat cgccgatttc tttattaaga acaaggtcgg aacagtgcag 1260
atggagaacc tcgagagcat gaagaggaag gaggattcct acttcaacat tcggctgagg 1320
gggttctggc cctacgctga gatgcagaa aagattgagt ttaagctgaa gcagtacggg 1380
attgagatcc ggaaggtggc ccccaacaac accagcaaga cctgcagcaa gtgcggggac 1440
ctcaacaact acttcaactt cgagtaccgg aagaagaaca agttcccaca cttcaagtgc 1500
gagaagtgca actttaagga gaaccccgat tacaacggcg ccctgaacat cagcaaccct 1560
aagctgaaga gcactaagga ggagccc 1587

```

```

SEQ ID NO: 11      moltype = RNA length = 161
FEATURE           Location/Qualifiers
source            1..161
                  mol_type = other RNA
                  note = Wild-type tracrRNA for TnpB or Cas12f1 gRNA
                  organism = synthetic construct

```

```

SEQUENCE: 11
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaagataa 120
ccctcgaaac aaattcattt ttctctcca attctgcaca a 161

```

```

SEQ ID NO: 12      moltype = RNA length = 37
FEATURE           Location/Qualifiers
source            1..37
                  mol_type = other RNA
                  note = Wild-type crRNA for TnpB or Cas12f1 gRNA
                  organism = synthetic construct

```

```

SEQUENCE: 12
gttgacgaac ccgaatagac gaatgaagga atgcaac 37

```

```

SEQ ID NO: 13      moltype = RNA length = 222
FEATURE           Location/Qualifiers

```

-continued

```

source          1..222
                 mol_type = other RNA
                 note = Canonical sgRNA for TnpB or Cas12f1
                 organism = synthetic construct
variation       203..222
                 note = n is a, c, g or u

SEQUENCE: 13
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcattt ttctctcca attctgcaca agaaagttgc agaaccgaa 180
tagacgaatg aaggaatgca acnnnnnnnn nnnnnnnnnn nn                      222

SEQ ID NO: 14      moltype = RNA length = 20
FEATURE           Location/Qualifiers
source            1..20
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 14
cttcactgat aaagtggaga                                         20

SEQ ID NO: 15      moltype = RNA length = 19
FEATURE           Location/Qualifiers
source            1..19
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 15
ttcactgata aagtggaga                                         19

SEQ ID NO: 16      moltype = RNA length = 18
FEATURE           Location/Qualifiers
source            1..18
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 16
tcactgataa agtggaga                                         18

SEQ ID NO: 17      moltype = RNA length = 17
FEATURE           Location/Qualifiers
source            1..17
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 17
cactgataaa gtggaga                                         17

SEQ ID NO: 18      moltype = RNA length = 16
FEATURE           Location/Qualifiers
source            1..16
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 18
actgataaag tggaga                                         16

SEQ ID NO: 19      moltype = RNA length = 15
FEATURE           Location/Qualifiers
source            1..15
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 19
ctgataaagt ggaga                                         15

SEQ ID NO: 20      moltype = RNA length = 14
FEATURE           Location/Qualifiers
source            1..14
                 mol_type = other RNA
                 note = Xa of Formula (I)
                 organism = synthetic construct

SEQUENCE: 20
tgataaagtg gaga                                         14

SEQ ID NO: 21      moltype = RNA length = 13

```

-continued

FEATURE	Location/Qualifiers	
source	1..13 mol_type = other RNA note = Xa of Formula (I) organism = synthetic construct	
SEQUENCE: 21		
gataaagtgg aga		13
SEQ ID NO: 22	moltype = RNA length = 12	
FEATURE	Location/Qualifiers	
source	1..12 mol_type = other RNA note = Xa of Formula (I) organism = synthetic construct	
SEQUENCE: 22		
ataaagtgga ga		12
SEQ ID NO: 23	moltype = RNA length = 11	
FEATURE	Location/Qualifiers	
source	1..11 mol_type = other RNA note = Xa of Formula (I) organism = synthetic construct	
SEQUENCE: 23		
taaagtggag a		11
SEQ ID NO: 24	moltype = RNA length = 10	
FEATURE	Location/Qualifiers	
source	1..10 mol_type = other RNA note = Xa of Formula (I) organism = synthetic construct	
SEQUENCE: 24		
aaagtggaga		10
SEQ ID NO: 25	moltype = RNA length = 13	
FEATURE	Location/Qualifiers	
source	1..13 mol_type = other RNA note = Xb1 of Formula (I) organism = synthetic construct	
SEQUENCE: 25		
caaaagctgt ccc		13
SEQ ID NO: 26	moltype = RNA length = 12	
FEATURE	Location/Qualifiers	
source	1..12 mol_type = other RNA note = Xb1 of Formula (I) organism = synthetic construct	
SEQUENCE: 26		
caaaagctgt cc		12
SEQ ID NO: 27	moltype = RNA length = 11	
FEATURE	Location/Qualifiers	
source	1..11 mol_type = other RNA note = Xb1 of Formula (I) organism = synthetic construct	
SEQUENCE: 27		
caaaagctgt c		11
SEQ ID NO: 28	moltype = RNA length = 10	
FEATURE	Location/Qualifiers	
source	1..10 mol_type = other RNA note = Xb1 of Formula (I) organism = synthetic construct	
SEQUENCE: 28		
caaaagctgt		10
SEQ ID NO: 29	moltype = RNA length = 14	
FEATURE	Location/Qualifiers	
source	1..14 mol_type = other RNA note = Xb2 of Formula (I)	

-continued

SEQUENCE: 29	organism = synthetic construct	
gggattagaa cttg		14
SEQ ID NO: 30	moltype = RNA length = 13	
FEATURE	Location/Qualifiers	
source	1..13	
	mol_type = other RNA	
	note = Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 30		
ggattagaac ttg		13
SEQ ID NO: 31	moltype = RNA length = 12	
FEATURE	Location/Qualifiers	
source	1..12	
	mol_type = other RNA	
	note = Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 31		
gattagaact tg		12
SEQ ID NO: 32	moltype = RNA length = 11	
FEATURE	Location/Qualifiers	
source	1..11	
	mol_type = other RNA	
	note = Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 32		
attagaactt g		11
SEQ ID NO: 33	moltype = RNA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = other RNA	
	note = Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 33		
ttagaacttg		10
SEQ ID NO: 34	moltype = RNA length = 31	
FEATURE	Location/Qualifiers	
source	1..31	
	mol_type = other RNA	
	note = Xb1-UUAG-Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 34		
caaaagctgt cccttagggg attagaactt g		31
SEQ ID NO: 35	moltype = RNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other RNA	
	note = Xb1-UUAG-Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 35		
caaaagctgt ttagattaga acttg		25
SEQ ID NO: 36	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = Xb1-UUAG-Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 36		
caaaagcttt agagaacttg		20
SEQ ID NO: 37	moltype = RNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other RNA	
	note = Xb1-UUAG-Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 37		
caaaagctta ggaacttg		18

-continued

SEQ ID NO: 38	moltype = RNA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = other RNA	
	note = Xb1-UUAG-Xb2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 38		
caattagttg		10
SEQ ID NO: 39	moltype = RNA length = 28	
FEATURE	Location/Qualifiers	
source	1..28	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 39		
ttcatttttc ctctccaatt ctgcacaa		28
SEQ ID NO: 40	moltype = RNA length = 27	
FEATURE	Location/Qualifiers	
source	1..27	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 40		
ttcatttttc ctctccaatt ctgcaca		27
SEQ ID NO: 41	moltype = RNA length = 26	
FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 41		
ttcatttttc ctctccaatt ctgcac		26
SEQ ID NO: 42	moltype = RNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 42		
ttcatttttc ctctccaatt ctgca		25
SEQ ID NO: 43	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 43		
ttcatttttc ctctccaatt ctgc		24
SEQ ID NO: 44	moltype = RNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 44		
ttcatttttc ctctccaatt ctg		23
SEQ ID NO: 45	moltype = RNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 45		
ttcatttttc ctctccaatt ct		22
SEQ ID NO: 46	moltype = RNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other RNA	

-continued

	note = Xc1 of Formula (I) organism = synthetic construct	
SEQUENCE: 46		
ttcatttttc ctctccaatt c		21
SEQ ID NO: 47	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 47		
ttcatttttc ctctccaatt		20
SEQ ID NO: 48	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 48		
ttcatttttc ctctccaat		19
SEQ ID NO: 49	moltype = RNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 49		
ttcatttttc ctctccaa		18
SEQ ID NO: 50	moltype = RNA length = 17	
FEATURE	Location/Qualifiers	
source	1..17	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 50		
ttcatttttc ctctcca		17
SEQ ID NO: 51	moltype = RNA length = 16	
FEATURE	Location/Qualifiers	
source	1..16	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 51		
ttcatttttc ctctcc		16
SEQ ID NO: 52	moltype = RNA length = 15	
FEATURE	Location/Qualifiers	
source	1..15	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 52		
ttcatttttc ctctc		15
SEQ ID NO: 53	moltype = RNA length = 14	
FEATURE	Location/Qualifiers	
source	1..14	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 53		
ttcatttttc ctct		14
SEQ ID NO: 54	moltype = RNA length = 13	
FEATURE	Location/Qualifiers	
source	1..13	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 54		
ttcatttttc ctc		13

-continued

SEQ ID NO: 55	moltype = RNA length = 12	
FEATURE	Location/Qualifiers	
source	1..12	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 55		
ttcatttttc ct		12
SEQ ID NO: 56	moltype = RNA length = 11	
FEATURE	Location/Qualifiers	
source	1..11	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 56		
ttcatttttc c		11
SEQ ID NO: 57	moltype = RNA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = other RNA	
	note = Xc1 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 57		
ttcatttttc		10
SEQ ID NO: 58	moltype = RNA length = 27	
FEATURE	Location/Qualifiers	
source	1..27	
	mol_type = other RNA	
	note = Xc2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 58		
gttgcagaac ccgaatagac gaatgaa		27
SEQ ID NO: 59	moltype = RNA length = 26	
FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other RNA	
	note = Xc2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 59		
ttgcagaacc cgaatagacg aatgaa		26
SEQ ID NO: 60	moltype = RNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other RNA	
	note = Xc2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 60		
tcgagaaccc gaatagacga atgaa		25
SEQ ID NO: 61	moltype = RNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other RNA	
	note = Xc2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 61		
gcagaaccg aatagacgaa tgaa		24
SEQ ID NO: 62	moltype = RNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other RNA	
	note = Xc2 of Formula (I)	
	organism = synthetic construct	
SEQUENCE: 62		
cagaaccgga atagacgaat gaa		23
SEQ ID NO: 63	moltype = RNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	

-continued

	mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 63 agaacccgaa tagacgaatg aa		22
SEQ ID NO: 64 FEATURE source	moltype = RNA length = 21 Location/Qualifiers 1..21 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 64 gaacccgaat agacgaatga a		21
SEQ ID NO: 65 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 65 aaccggaata gacgaatgaa		20
SEQ ID NO: 66 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 66 accggaatag acgaatgaa		19
SEQ ID NO: 67 FEATURE source	moltype = RNA length = 18 Location/Qualifiers 1..18 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 67 ccggaataga cgaatgaa		18
SEQ ID NO: 68 FEATURE source	moltype = RNA length = 17 Location/Qualifiers 1..17 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 68 ccgaatagac gaatgaa		17
SEQ ID NO: 69 FEATURE source	moltype = RNA length = 16 Location/Qualifiers 1..16 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 69 cgaatagacg aatgaa		16
SEQ ID NO: 70 FEATURE source	moltype = RNA length = 15 Location/Qualifiers 1..15 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 70 gaatagacga atgaa		15
SEQ ID NO: 71 FEATURE source	moltype = RNA length = 14 Location/Qualifiers 1..14 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct	
SEQUENCE: 71		

-continued

aatagacgaa tgaa	14
SEQ ID NO: 72 FEATURE source	moltype = RNA length = 13 Location/Qualifiers 1..13 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct
SEQUENCE: 72 atagacgaat gaa	13
SEQ ID NO: 73 FEATURE source	moltype = RNA length = 12 Location/Qualifiers 1..12 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct
SEQUENCE: 73 tagacgaatg aa	12
SEQ ID NO: 74 FEATURE source	moltype = RNA length = 11 Location/Qualifiers 1..11 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct
SEQUENCE: 74 agacgaatga a	11
SEQ ID NO: 75 FEATURE source	moltype = RNA length = 10 Location/Qualifiers 1..10 mol_type = other RNA note = Xc2 of Formula (I) organism = synthetic construct
SEQUENCE: 75 gacgaatgaa	10
SEQ ID NO: 76 FEATURE source	moltype = RNA length = 10 Location/Qualifiers 1..10 mol_type = other RNA note = Linker (Lk) organism = synthetic construct
SEQUENCE: 76 ttcgaaagaa	10
SEQ ID NO: 77 FEATURE source	moltype = RNA length = 12 Location/Qualifiers 1..12 mol_type = other RNA note = Linker (Lk) organism = synthetic construct
SEQUENCE: 77 ttcagaaatg aa	12
SEQ ID NO: 78 FEATURE source	moltype = RNA length = 14 Location/Qualifiers 1..14 mol_type = other RNA note = Linker (Lk) organism = synthetic construct
SEQUENCE: 78 ttcatgaaaa tgaa	14
SEQ ID NO: 79 FEATURE source	moltype = RNA length = 16 Location/Qualifiers 1..16 mol_type = other RNA note = Linker (Lk) organism = synthetic construct
SEQUENCE: 79 ttcattgaaa aatgaa	16
SEQ ID NO: 80 FEATURE	moltype = RNA length = 59 Location/Qualifiers

-continued

```

source                1..59
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 80
ttcagtgctc ctctccaatt ctgcacaaga aagttgcaga acccgaatag agcaatgaa 59

SEQ ID NO: 81          moltype = RNA length = 46
FEATURE               Location/Qualifiers
source                1..46
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 81
ttcagtgctc ctctccaatt cgaagaacc cgaatagagc aatgaa 46

SEQ ID NO: 82          moltype = RNA length = 34
FEATURE               Location/Qualifiers
source                1..34
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 82
ttcagtgctc ctctcgaaag aatagagcaa tgaa 34

SEQ ID NO: 83          moltype = RNA length = 22
FEATURE               Location/Qualifiers
source                1..22
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 83
ttcagtgctg aaaagcaatg aa 22

SEQ ID NO: 84          moltype = RNA length = 18
FEATURE               Location/Qualifiers
source                1..18
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 84
ttcatttgaa agaatgaa 18

SEQ ID NO: 85          moltype = RNA length = 14
FEATURE               Location/Qualifiers
source                1..14
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 85
ttcatgaaaa tgaa 14

SEQ ID NO: 86          moltype = RNA length = 10
FEATURE               Location/Qualifiers
source                1..10
                      mol_type = other RNA
                      note = Xc1-Lk-Xc2 of Formula (I)
                      organism = synthetic construct

SEQUENCE: 86
ttcgaaagaa 10

SEQ ID NO: 87          moltype = RNA length = 161
FEATURE               Location/Qualifiers
source                1..161
                      mol_type = other RNA
                      note = Engineered tracrRNA for TnpB or Cas12f1 gRNA (MS1)
                      organism = synthetic construct

SEQUENCE: 87
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctctctcca attctgcaca a 161

SEQ ID NO: 88          moltype = RNA length = 154
FEATURE               Location/Qualifiers
source                1..154
                      mol_type = other RNA

```

-continued

```

                                note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                                (MS1/MS3-1)
                                organism = synthetic construct

SEQUENCE: 88
gataaagtgg agaaccgctt caccaaaagc tgtcccttag gggattagaa cttgagtga 60
ggtagggctgc ttgcatcagc ctaatgtcga gaagtgtttt ctcggaaag taaccctcga 120
aacaattca gtgtcctct ccaattctgc acaa 154

SEQ ID NO: 89      moltype = RNA length = 147
FEATURE           Location/Qualifiers
source           1..147
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS3-2)
                  organism = synthetic construct

SEQUENCE: 89
tggagaaccg cttcaccaaa agctgtccct taggggatta gaacttgagt gaaggtgggc 60
tgcttgcatc agcctaattg cgagaagtgc tttcttcgga aagtaaccct cgaacaaat 120
tcagtgtccc tctccaattc tgcacaa 147

SEQ ID NO: 90      moltype = RNA length = 141
FEATURE           Location/Qualifiers
source           1..141
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS3-3)
                  organism = synthetic construct

SEQUENCE: 90
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
ctcctctcca attctgcaca a 141

SEQ ID NO: 91      moltype = RNA length = 154
FEATURE           Location/Qualifiers
source           1..154
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS4*-1)
                  organism = synthetic construct

SEQUENCE: 91
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctcctctcca attc 154

SEQ ID NO: 92      moltype = RNA length = 148
FEATURE           Location/Qualifiers
source           1..148
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS4*-2)
                  organism = synthetic construct

SEQUENCE: 92
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctcctctc 148

SEQ ID NO: 93      moltype = RNA length = 142
FEATURE           Location/Qualifiers
source           1..142
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS4*-3)
                  organism = synthetic construct

SEQUENCE: 93
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ct 142

SEQ ID NO: 94      moltype = RNA length = 155
FEATURE           Location/Qualifiers
source           1..155
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                  (MS1/MS5-1)
                  organism = synthetic construct

SEQUENCE: 94

```

-continued

```

cttcactgat aaagtggaga accgcttcac caaaagctgt ttagattaga acttgagtga 60
agggtgggctg cttgcatcag cctaattgtcg agaagtgcctt tcttcggaaa gtaaccctcg 120
aaacaaattc agtgctcctc tccaattctg caca 155

SEQ ID NO: 95      moltype = RNA length = 148
FEATURE           Location/Qualifiers
source            1..148
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS5-2)
                  organism = synthetic construct

SEQUENCE: 95
cttcactgat aaagtggaga accgcttcac caaaagctta ggaacttgag tgaaggtggg 60
ctgcttgcat cagcctaatt tcgagaagtg ctttcttcgg aaagtaacct tcgaaacaaa 120
ttcagtgtctc ctctccaatt ctgcacaa 148

SEQ ID NO: 96      moltype = RNA length = 140
FEATURE           Location/Qualifiers
source            1..140
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS5-3)
                  organism = synthetic construct

SEQUENCE: 96
cttcactgat aaagtggaga accgcttcac caattagttg agtgaagggt ggctgcttgc 60
atcagcctaa tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca aaattcagtg 120
tcctctccaa ttctgcacaa 140

SEQ ID NO: 97      moltype = RNA length = 134
FEATURE           Location/Qualifiers
source            1..134
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS3-3/MS4*-1)
                  organism = synthetic construct

SEQUENCE: 97
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaagggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
ctcctctcca attc 134

SEQ ID NO: 98      moltype = RNA length = 128
FEATURE           Location/Qualifiers
source            1..128
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS3-3/MS4*-2)
                  organism = synthetic construct

SEQUENCE: 98
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaagggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
ctcctctc 128

SEQ ID NO: 99      moltype = RNA length = 122
FEATURE           Location/Qualifiers
source            1..122
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS3-3/MS4*-3)
                  organism = synthetic construct

SEQUENCE: 99
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaagggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
ct 122

SEQ ID NO: 100     moltype = RNA length = 142
FEATURE           Location/Qualifiers
source            1..142
                  mol_type = other RNA
                  note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                    (MS1/MS4*-2/MS5-1)
                  organism = synthetic construct

SEQUENCE: 100
cttcactgat aaagtggaga accgcttcac caaaagctgt ttagattaga acttgagtga 60
agggtgggctg cttgcatcag cctaattgtcg agaagtgcctt tcttcggaaa gtaaccctcg 120
aaacaaattc agtgctcctc tc 142

```

-continued

SEQ ID NO: 101 moltype = RNA length = 135
 FEATURE Location/Qualifiers
 source 1..135
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS4*-2/MS5-2)
 organism = synthetic construct

SEQUENCE: 101
 cttcactgat aaagtgagga accgcttcac caaaagctta ggaacttgag tgaaggtggg 60
 ctgcttgcat cagcctaagt tcgagaagtg ctttcttcgg aaagtaacct tcgaaacaaa 120
 ttcagtgtct ctctc 135

SEQ ID NO: 102 moltype = RNA length = 127
 FEATURE Location/Qualifiers
 source 1..127
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS4*-2/MS5-3)
 organism = synthetic construct

SEQUENCE: 102
 cttcactgat aaagtgagga accgcttcac caattagtgt agtgaagggt ggctgcttgc 60
 atcagcctaa tgcgagaagt tgctttcttc ggaaagtaac cctcgaaaca aattcagtgc 120
 tcctctc 127

SEQ ID NO: 103 moltype = RNA length = 135
 FEATURE Location/Qualifiers
 source 1..135
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS3-3/MS5-1)
 organism = synthetic construct

SEQUENCE: 103
 accgcttcac caaaagctgt ttagattaga acttgagtga aggtgggctg cttgcatcag 60
 cctaattgtc agaagtgtct tcttcggaaa gtaaccctcg aaacaaattc agtgcctctc 120
 tccaattctg caca 135

SEQ ID NO: 104 moltype = RNA length = 128
 FEATURE Location/Qualifiers
 source 1..128
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS3-3/MS5-2)
 organism = synthetic construct

SEQUENCE: 104
 accgcttcac caaaagctta ggaacttgag tgaaggtggg ctgcttgcat cagcctaagt 60
 tcgagaagtg ctttcttcgg aaagtaacct tcgaaacaaa ttcagtgtct ctctccaatt 120
 ctgcacaa 128

SEQ ID NO: 105 moltype = RNA length = 120
 FEATURE Location/Qualifiers
 source 1..120
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS3-3/MS5-3)
 organism = synthetic construct

SEQUENCE: 105
 accgcttcac caattagtgt agtgaagggt ggctgcttgc atcagcctaa tgcgagaag 60
 tgctttcttc ggaaagtaac cctcgaaaca aattcagtgc tcctctccaa ttctgcacaa 120

SEQ ID NO: 106 moltype = RNA length = 107
 FEATURE Location/Qualifiers
 source 1..107
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MS1/MS3-3/MS4*-2/MS5-3)
 organism = synthetic construct

SEQUENCE: 106
 accgcttcac caattagtgt agtgaagggt ggctgcttgc atcagcctaa tgcgagaag 60
 tgctttcttc ggaaagtaac cctcgaaaca aattcagtgc tcctctc 107

SEQ ID NO: 107 moltype = RNA length = 140
 FEATURE Location/Qualifiers
 source 1..140
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA (MF)
 organism = synthetic construct

-continued

SEQUENCE: 107
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcattt 140

SEQ ID NO: 108 moltype = RNA length = 133
FEATURE Location/Qualifiers
source 1..133
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS3-1)
 organism = synthetic construct

SEQUENCE: 108
gataaagtgg agaaccgctt caccaaaagc tgtcccttag gggattagaa cttgagttaa 60
ggtaggctgc ttgcatcagc ctaatgtcga gaagtgtttt ctcggaaag taaccctcga 120
aacaattca ttt 133

SEQ ID NO: 109 moltype = RNA length = 126
FEATURE Location/Qualifiers
source 1..126
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS3-2)
 organism = synthetic construct

SEQUENCE: 109
tggagaaccg cttcaccaaa agctgtccct taggggatta gaacttgagt gaaggtgggc 60
tgcttgcatc agcctaattg cgagaagtgc tttcttcgga aagtaaccct cgaacaaat 120
tcattt 126

SEQ ID NO: 110 moltype = RNA length = 120
FEATURE Location/Qualifiers
source 1..120
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS3-3)
 organism = synthetic construct

SEQUENCE: 110
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcattt 120

SEQ ID NO: 111 moltype = RNA length = 138
FEATURE Location/Qualifiers
source 1..138
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS4-1)
 organism = synthetic construct

SEQUENCE: 111
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcat 138

SEQ ID NO: 112 moltype = RNA length = 136
FEATURE Location/Qualifiers
source 1..136
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS4-2)
 organism = synthetic construct

SEQUENCE: 112
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattc 136

SEQ ID NO: 113 moltype = RNA length = 133
FEATURE Location/Qualifiers
source 1..133
 mol_type = other RNA
 note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
 (MF/MS4-3)
 organism = synthetic construct

SEQUENCE: 113
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaa 133

```

SEQ ID NO: 114           multype = RNA length = 134
FEATURE                 Location/Qualifiers
source                  1..134
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS5-1)
                        organism = synthetic construct

SEQUENCE: 114
cttcactgat aaagtggaga accgcttcac caaaagctgt ttagattaga acttgagtga 60
agggtgggctg cttgcatcag cctaattgtcg agaagtgcctt tcttcggaaa gtaaccctcg 120
aaacaaattc attt                                     134

SEQ ID NO: 115           multype = RNA length = 129
FEATURE                 Location/Qualifiers
source                  1..129
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS5-2)
                        organism = synthetic construct

SEQUENCE: 115
cttcactgat aaagtggaga accgcttcac caaaagcttt agagaacttg agtgaaggtg 60
ggctgcttgc atcagcctaa tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca 120
aattcattt                                     129

SEQ ID NO: 116           multype = RNA length = 119
FEATURE                 Location/Qualifiers
source                  1..119
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS5-3)
                        organism = synthetic construct

SEQUENCE: 116
cttcactgat aaagtggaga accgcttcac caattagtgt agtgaaggtg ggctgcttgc 60
atcagcctaa tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca aattcattt 119

SEQ ID NO: 117           multype = RNA length = 113
FEATURE                 Location/Qualifiers
source                  1..113
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA (MF/MS5)
                        organism = synthetic construct

SEQUENCE: 117
cttcactgat aaagtggaga accgcttcac ttagagttaa ggtgggctgc ttgcatcagc 60
ctaattgtcg gaagtgcctt cttcggaag taaccctcga aacaaattca ttt 113

SEQ ID NO: 118           multype = RNA length = 118
FEATURE                 Location/Qualifiers
source                  1..118
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS3-3/MS4-1)
                        organism = synthetic construct

SEQUENCE: 118
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa cctcgaaac aaattcat 118

SEQ ID NO: 119           multype = RNA length = 116
FEATURE                 Location/Qualifiers
source                  1..116
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS3-3/MS4-2)
                        organism = synthetic construct

SEQUENCE: 119
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa cctcgaaac aaattc 116

SEQ ID NO: 120           multype = RNA length = 113
FEATURE                 Location/Qualifiers
source                  1..113
                        mol_type = other RNA
                        note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                          (MF/MS3-3/MS4-3)
                        organism = synthetic construct

SEQUENCE: 120
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60

```

-continued

```

catcagccta atgtcgagaa gtgctttctt cggaaagtaa cctcgaaac aaa      113

SEQ ID NO: 121      moltype = RNA  length = 127
FEATURE            Location/Qualifiers
source             1..127
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS4-3/MS5-1)
                   organism = synthetic construct

SEQUENCE: 121
cttcactgat aaagtggaga accgcttcac caaaagctgt ttagattaga acttgagtga  60
aggtgggctg cttgcatcag cctaattgtcg agaagtgtct tcttcggaaa gtaaccctcg 120
aaacaaa                                127

SEQ ID NO: 122      moltype = RNA  length = 122
FEATURE            Location/Qualifiers
source             1..122
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS4-3/MS5-2)
                   organism = synthetic construct

SEQUENCE: 122
cttcactgat aaagtggaga accgcttcac caaaagcttt agagaacttg agtgaagggtg  60
ggctgcttgc atcagcctaa tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca 120
aa                                122

SEQ ID NO: 123      moltype = RNA  length = 112
FEATURE            Location/Qualifiers
source             1..112
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS4-3/MS5-3)
                   organism = synthetic construct

SEQUENCE: 123
cttcactgat aaagtggaga accgcttcac caattagtgt agtgaagggtg ggctgcttgc  60
atcagcctaa tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca aa      112

SEQ ID NO: 124      moltype = RNA  length = 106
FEATURE            Location/Qualifiers
source             1..106
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS4-3/MS5)
                   organism = synthetic construct

SEQUENCE: 124
cttcactgat aaagtggaga accgcttcac ttagagttaa ggtgggctgc ttgcatcagc  60
ctaattgtcg gaagtgtctt cttcggaaaag taaccctcga aacaaa      106

SEQ ID NO: 125      moltype = RNA  length = 114
FEATURE            Location/Qualifiers
source             1..114
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS3-3/MS5-1)
                   organism = synthetic construct

SEQUENCE: 125
accgcttcac caaaagctgt ttagattaga acttgagtga aggtgggctg cttgcatcag  60
cctaattgtcg agaagtgtct tcttcggaaa gtaaccctcg aaacaaattc attt      114

SEQ ID NO: 126      moltype = RNA  length = 109
FEATURE            Location/Qualifiers
source             1..109
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS3-3/MS5-2)
                   organism = synthetic construct

SEQUENCE: 126
accgcttcac caaaagcttt agagaacttg agtgaagggt ggctgcttgc atcagcctaa  60
tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca aattcattt      109

SEQ ID NO: 127      moltype = RNA  length = 99
FEATURE            Location/Qualifiers
source             1..99
                   mol_type = other RNA
                   note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                     (MF/MS3-3/MS5-3)

```

-continued

```

                organism = synthetic construct
SEQUENCE: 127
accgcttcac caattagttg agtgaaggtg ggctgcttgc atcagcctaa tgcgagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aattcattt 99

SEQ ID NO: 128      moltype = RNA length = 93
FEATURE            Location/Qualifiers
source              1..93
                    mol_type = other RNA
                    note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                      (MF/MS3-3/MS5)
                    organism = synthetic construct
SEQUENCE: 128
accgcttcac ttagagttaa ggtgggctgc ttgcatcagc ctaatgtcga gaagtgcttt 60
cttcggaaaag taaccctcga aacaaattca ttt 93

SEQ ID NO: 129      moltype = RNA length = 92
FEATURE            Location/Qualifiers
source              1..92
                    mol_type = other RNA
                    note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                      (MF/MS3-3/MS4-3/MS5-3)
                    organism = synthetic construct
SEQUENCE: 129
accgcttcac caattagttg agtgaaggtg ggctgcttgc atcagcctaa tgcgagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aa 92

SEQ ID NO: 130      moltype = RNA length = 91
FEATURE            Location/Qualifiers
source              1..91
                    mol_type = other RNA
                    note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                      (MF/MS3-3/MS4-1/MS5)
                    organism = synthetic construct
SEQUENCE: 130
accgcttcac ttagagttaa ggtgggctgc ttgcatcagc ctaatgtcga gaagtgcttt 60
cttcggaaaag taaccctcga aacaaattca t 91

SEQ ID NO: 131      moltype = RNA length = 89
FEATURE            Location/Qualifiers
source              1..89
                    mol_type = other RNA
                    note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                      (MF/MS3-3/MS4-2/MS5)
                    organism = synthetic construct
SEQUENCE: 131
accgcttcac ttagagttaa ggtgggctgc ttgcatcagc ctaatgtcga gaagtgcttt 60
cttcggaaaag taaccctcga aacaaattc 89

SEQ ID NO: 132      moltype = RNA length = 86
FEATURE            Location/Qualifiers
source              1..86
                    mol_type = other RNA
                    note = Engineered tracrRNA for TnpB or Cas12f1 gRNA
                      (MF/MS3-3/MS4-3/MS5)
                    organism = synthetic construct
SEQUENCE: 132
accgcttcac ttagagttaa ggtgggctgc ttgcatcagc ctaatgtcga gaagtgcttt 60
cttcggaaaag taaccctcga aacaaa 86

SEQ ID NO: 133      moltype = RNA length = 37
FEATURE            Location/Qualifiers
source              1..37
                    mol_type = other RNA
                    note = Engineered crRNA for TnpB or Cas12f1 gRNA (MS1)
                    organism = synthetic construct
SEQUENCE: 133
gttcagaaac ccgaatagag caatgaagga atgcaac 37

SEQ ID NO: 134      moltype = RNA length = 31
FEATURE            Location/Qualifiers
source              1..31
                    mol_type = other RNA
                    note = Engineered crRNA for TnpB or Cas12f1 gRNA
                      (MS1/MS4*-1)
                    organism = synthetic construct

```

-continued

SEQUENCE: 134
gaaccggaat agagcaatga aggaatgcaa c 31

SEQ ID NO: 135 moltype = RNA length = 25
FEATURE Location/Qualifiers
source 1..25
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA
(MS1/MS4*-2)
organism = synthetic construct

SEQUENCE: 135
gaatagagca atgaaggaat gcaac 25

SEQ ID NO: 136 moltype = RNA length = 19
FEATURE Location/Qualifiers
source 1..19
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA
(MS1/MS4*-3)
organism = synthetic construct

SEQUENCE: 136
agcaatgaag gaatgcaac 19

SEQ ID NO: 137 moltype = RNA length = 17
FEATURE Location/Qualifiers
source 1..17
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA (MF)
organism = synthetic construct

SEQUENCE: 137
gaatgaagga atgcaac 17

SEQ ID NO: 138 moltype = RNA length = 15
FEATURE Location/Qualifiers
source 1..15
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA (MF/MS4-1)
organism = synthetic construct

SEQUENCE: 138
atgaaggaat gcaac 15

SEQ ID NO: 139 moltype = RNA length = 13
FEATURE Location/Qualifiers
source 1..13
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA (MF/MS4-2)
organism = synthetic construct

SEQUENCE: 139
gaaggaatgc aac 13

SEQ ID NO: 140 moltype = RNA length = 10
FEATURE Location/Qualifiers
source 1..10
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA (MF/MS4-3)
organism = synthetic construct

SEQUENCE: 140
ggaatgcaac 10

SEQ ID NO: 141 moltype = RNA length = 68
FEATURE Location/Qualifiers
source 1..68
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA (MS1/MS2)
organism = synthetic construct

variation 38..57
note = n is a, c, g or u

SEQUENCE: 141
gttcagaaac ccgaatagag caatgaagga atgcaacnnn nnnnnnnnnn nnnnnnnnttt 60
tatttttt 68

SEQ ID NO: 142 moltype = RNA length = 62
FEATURE Location/Qualifiers
source 1..62
mol_type = other RNA
note = Engineered crRNA for TnpB or Cas12f1 gRNA

-continued

```

(MS1/MS2/MS4*-1)
organism = synthetic construct
variation      32..51
               note = n is a, c, g or u
SEQUENCE: 142
gaaccggaat agagcaatga aggaatgcaa cnnnnnnnnn nnnnnnnnnn nttttatttt 60
tt                                                62

SEQ ID NO: 143      moltype = RNA length = 56
FEATURE            Location/Qualifiers
source             1..56
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA
                   (MS1/MS2/MS4*-2)
                   organism = synthetic construct
variation          26..45
                   note = n is a, c, g or u
SEQUENCE: 143
gaatagagca atgaaggaat gcaacnnnnn nnnnnnnnnn nnnnntttta tttttt 56

SEQ ID NO: 144      moltype = RNA length = 50
FEATURE            Location/Qualifiers
source             1..50
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA
                   (MS1/MS2/MS4*-3)
                   organism = synthetic construct
variation          20..39
                   note = n is a, c, g or u
SEQUENCE: 144
agcaatgaag gaatgcaacn nnnnnnnnnn nnnnnnnnnt tttatttttt 50

SEQ ID NO: 145      moltype = RNA length = 48
FEATURE            Location/Qualifiers
source             1..48
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA (MF/MS2)
                   organism = synthetic construct
variation          18..37
                   note = n is a, c, g or u
SEQUENCE: 145
gaatgaagga atgcaacnnn nnnnnnnnnn nnnnnnnntt tatttttt 48

SEQ ID NO: 146      moltype = RNA length = 46
FEATURE            Location/Qualifiers
source             1..46
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA
                   (MF/MS2/MS4-1)
                   organism = synthetic construct
variation          16..35
                   note = n is a, c, g or u
SEQUENCE: 146
atgaaggaat gcaacnnnnn nnnnnnnnnn nnnnntttta tttttt 46

SEQ ID NO: 147      moltype = RNA length = 44
FEATURE            Location/Qualifiers
source             1..44
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA
                   (MF/MS2/MS4-2)
                   organism = synthetic construct
variation          14..33
                   note = n is a, c, g or u
SEQUENCE: 147
gaaggaatgc aacnnnnnnn nnnnnnnnnn nnnttttatt tttt 44

SEQ ID NO: 148      moltype = RNA length = 41
FEATURE            Location/Qualifiers
source             1..41
                   mol_type = other RNA
                   note = Engineered crRNA for TnpB or Cas12f1 gRNA
                   (MF/MS2/MS4-3)
                   organism = synthetic construct
variation          11..30
                   note = n is a, c, g or u

```

-continued

SEQUENCE: 148
ggaatgcaac nnnnnnnnnn nnnnnnnnnn ttttattttt t 41

SEQ ID NO: 149 moltype = RNA length = 222
FEATURE Location/Qualifiers
source 1..222
mol_type = other RNA
note = Engineered sgRNA with modification of MS1 for TnpB
or Cas12f1
organism = synthetic construct
variation 203..222
note = n is a, c, g or u

SEQUENCE: 149
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctctctcca attctgcaca agaaagttgc agaaccgaa 180
tagagcaatg aaggaatgca acnnnnnnnn nnnnnnnnnn nn 222

SEQ ID NO: 150 moltype = RNA length = 233
FEATURE Location/Qualifiers
source 1..233
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS2 for
TnpB or Cas12f1
organism = synthetic construct
variation 203..222
note = n is a, c, g or u

SEQUENCE: 150
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctctctcca attctgcaca agaaagttgc agaaccgaa 180
tagagcaatg aaggaatgca acnnnnnnnn nnnnnnnnnn nntttattt ttt 233

SEQ ID NO: 151 moltype = RNA length = 213
FEATURE Location/Qualifiers
source 1..213
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS2/MS3
for TnpB or Cas12f1
organism = synthetic construct
variation 183..202
note = n is a, c, g or u

SEQUENCE: 151
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
ctctctcca attctgcaca agaaagttgc agaaccgaa tagagcaatg aaggaatgca 180
acnnnnnnnn nnnnnnnnnn nntttattt ttt 213

SEQ ID NO: 152 moltype = RNA length = 158
FEATURE Location/Qualifiers
source 1..158
mol_type = other RNA
note = Engineered sgRNA with modification of MS2/MS3/MS4
for TnpB or Cas12f1
organism = synthetic construct
variation 128..147
note = n is a, c, g or u

SEQUENCE: 152
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaagaaagga 120
atgcaacnnn nnnnnnnnnn nnnnnnnntt tatttttt 158

SEQ ID NO: 153 moltype = RNA length = 129
FEATURE Location/Qualifiers
source 1..129
mol_type = other RNA
note = Engineered sgRNA with modification of
MS2/MS3/MS4/MS5 for TnpB or Cas12f1
organism = synthetic construct
variation 99..118
note = n is a, c, g or u

SEQUENCE: 153
accgcttcac cgagtgaagg tgggctgctt gcacagcct aatgtcgaga agtgccttct 60
tcggaagta accctcgaaa caaagaaagg aatgcaacnn nnnnnnnnnn nnnnnnnntt 120
ttatttttt 129

-continued

SEQ ID NO: 154 moltype = RNA length = 215
 FEATURE Location/Qualifiers
 source 1..215
 mol_type = other RNA
 note = Engineered sgRNA with modification of MS1/MS3-1 for
 TnpB or Cas12f1
 organism = synthetic construct
 variation 196..215
 note = n is a, c, g or u

SEQUENCE: 154
 gataaagtgg agaaccgctt caccaaaagc tgtcccttag gggattagaa cttgagtga 60
 ggtgggctgc ttgcatcagc ctaatgtcga gaagtgttt ctcggaaag taaccctcga 120
 aacaaattca gtgtcctctt ccaattctgc acaagaaagt tgcagaacc gaatagagca 180
 atgaaggaat gcaacnnnnn nnnnnnnnn nnnnn 215

SEQ ID NO: 155 moltype = RNA length = 208
 FEATURE Location/Qualifiers
 source 1..208
 mol_type = other RNA
 note = Engineered sgRNA with modification of MS1/MS3-2 for
 TnpB or Cas12f1
 organism = synthetic construct
 variation 189..208
 note = n is a, c, g or u

SEQUENCE: 155
 tggagaaccg cttcaccaaa agctgtccct taggggatta gaacttgagt gaaggtgggc 60
 tgcttgcatc agcctaattg cgagaagtgc tttcttcgga aagtaaccct cgaacaaaat 120
 tcagtgtctc tctccaattc tgcacaagaa agttgcagaa cccaataga gcaatgaag 180
 aatgcaacnn nnnnnnnnnn nnnnnnnnn 208

SEQ ID NO: 156 moltype = RNA length = 202
 FEATURE Location/Qualifiers
 source 1..202
 mol_type = other RNA
 note = Engineered sgRNA with modification of MS1/MS3-3 for
 TnpB or Cas12f1
 organism = synthetic construct
 variation 183..202
 note = n is a, c, g or u

SEQUENCE: 156
 accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
 catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcagtg 120
 ctctctcca attctgcaca agaaagttgc agaaccgaa tagagcaatg aaggaatga 180
 acnnnnnnnn nnnnnnnnnn nn 202

SEQ ID NO: 157 moltype = RNA length = 209
 FEATURE Location/Qualifiers
 source 1..209
 mol_type = other RNA
 note = Engineered sgRNA with modification of MS1/MS4*-1 for
 TnpB or Cas12f1
 organism = synthetic construct
 variation 190..209
 note = n is a, c, g or u

SEQUENCE: 157
 cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
 gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
 ccctcgaaac aaattcagtg ctctctcca attcgaaaga acccgaaatg agcaatgaag 180
 gaatgcaacn nnnnnnnnnn nnnnnnnnn 209

SEQ ID NO: 158 moltype = RNA length = 197
 FEATURE Location/Qualifiers
 source 1..197
 mol_type = other RNA
 note = Engineered sgRNA with modification of MS1/MS4*-2 for
 TnpB or Cas12f1
 organism = synthetic construct
 variation 178..197
 note = n is a, c, g or u

SEQUENCE: 158
 cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
 gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
 ccctcgaaac aaattcagtg ctctctcga aagaatagag caatgaagga atgcaacnn 180
 nnnnnnnnnn nnnnnnn 197

SEQ ID NO: 159 moltype = RNA length = 185

-continued

FEATURE
source Location/Qualifiers
1..185
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS4*-3 for
TnpB or Cas12f1
organism = synthetic construct
variation 166..185
note = n is a, c, g or u

SEQUENCE: 159
cttcactgat aaagtgagga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctgaaaagca atgaaggaat gcaacnnnnn nnnnnnnnnn 180
nnnnn 185

SEQ ID NO: 160 moltype = RNA length = 216
FEATURE Location/Qualifiers
source 1..216
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS5-1 for
TnpB or Cas12f1
organism = synthetic construct
variation 197..216
note = n is a, c, g or u

SEQUENCE: 160
cttcactgat aaagtgagga accgcttcac caaaagctgt ttagattaga acttgagtga 60
aggtgggctg cttgcatcag cctaattgtcg agaagtgcctt tcttcggaaa gtaaccctcg 120
aaacaaattc agtgctcctc tccaattctg cacaagaaag ttgcagaacc cgaatagagc 180
aatgaaggaa tgcaacnnnn nnnnnnnnnn nnnnnn 216

SEQ ID NO: 161 moltype = RNA length = 209
FEATURE Location/Qualifiers
source 1..209
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS5-2 for
TnpB or Cas12f1
organism = synthetic construct
variation 190..209
note = n is a, c, g or u

SEQUENCE: 161
cttcactgat aaagtgagga accgcttcac caaaagctta ggaacttgag tgaagtgagg 60
ctgcttgcat cagcctaagtg tcgagaagtg ctttcttcgg aaagtaacct tcgaaacaaa 120
ttcagtgtct cttcccaatt ctgcacaaga aagttgcaga acccgaatag agcaatgaag 180
gaatgcaacn nnnnnnnnnn nnnnnnnnn 209

SEQ ID NO: 162 moltype = RNA length = 201
FEATURE Location/Qualifiers
source 1..201
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS5-3 for
TnpB or Cas12f1
organism = synthetic construct
variation 182..201
note = n is a, c, g or u

SEQUENCE: 162
cttcactgat aaagtgagga accgcttcac caattagttg agtgaagggt ggctgcttgc 60
atcagcctaa tgtcgagaag tctttcttc ggaaagtaac cctcgaacaa aattcagtg 120
tcctctccaa ttctgcacaa gaaagttgca gaaccgcaat agagcaatga agaatgcaa 180
cnnnnnnnnn nnnnnnnnnn n n 201

SEQ ID NO: 163 moltype = RNA length = 206
FEATURE Location/Qualifiers
source 1..206
mol_type = other RNA
note = Engineered sgRNA with modification of MS1/MS2/MS4*-2
for TnpB or Cas12f1
organism = synthetic construct
variation 178..197
note = n is a, c, g or u

SEQUENCE: 163
cttcactgat aaagtgagga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcagtg ctcctctcga aagaatagag caatgaagga atgcaacnnn 180
nnnnnnnnnn nnnnnnnntt tatatt 206

SEQ ID NO: 164 moltype = RNA length = 177
FEATURE Location/Qualifiers

-continued

```

source          1..177
                mol_type = other RNA
                note = Engineered sgRNA with modification of
                    MS1/MS3-3/MS4*-2 for TnpB or Cas12f1
                organism = synthetic construct
variation       158..177
                note = n is a, c, g or u

SEQUENCE: 164
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cgaaaagtaa cctcgaaac aaattcagtg 120
ctcctctcga aagaatagag caatgaagga atgcaacnnn nnnnnnnnnn nnnnnnn 177

SEQ ID NO: 165      moltype = RNA length = 210
FEATURE            Location/Qualifiers
source             1..210
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of MS1/MS2/MS5-3
                      for TnpB or Cas12f1
                  organism = synthetic construct
variation          182..201
                  note = n is a, c, g or u

SEQUENCE: 165
cttcactgat aaagtggaga accgcttcac caattagttg agtgaagggt ggctgcttgc 60
atcagcctaa tgctcgagaag tgctttcttc ggaaagtaac cctcgaaaca aaattcagtg 120
tcctctccaa ttctgcacaa gaaagttgca gaaccogaat agagcaatga aggaatgcaa 180
cnnnnnnnnn nnnnnnnnnn nttttatatt 210

SEQ ID NO: 166      moltype = RNA length = 181
FEATURE            Location/Qualifiers
source             1..181
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                      MS1/MS3-3/MS5-3 for TnpB or Cas12f1
                  organism = synthetic construct
variation          162..181
                  note = n is a, c, g or u

SEQUENCE: 166
accgcttcac caattagttg agtgaagggt ggctgcttgc atcagcctaa tgctcgagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aaattcagtg tcctctccaa ttctgcacaa 120
gaaagttgca gaaccogaat agagcaatga aggaatgcaa cnnnnnnnnn nnnnnnnnnn 180
n                                                    181

SEQ ID NO: 167      moltype = RNA length = 176
FEATURE            Location/Qualifiers
source             1..176
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                      MS1/MS4*-2/MS5-3 for TnpB or Cas12f1
                  organism = synthetic construct
variation          157..176
                  note = n is a, c, g or u

SEQUENCE: 167
cttcactgat aaagtggaga accgcttcac caattagttg agtgaagggt ggctgcttgc 60
atcagcctaa tgctcgagaag tgctttcttc ggaaagtaac cctcgaaaca aaattcagtg 120
tcctctcgaa agaatagagc aatgaaggaa tgcaacnnn nnnnnnnnnn nnnnnnn 176

SEQ ID NO: 168      moltype = RNA length = 186
FEATURE            Location/Qualifiers
source             1..186
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                      MS1/MS2/MS3-3/MS4*-2 for TnpB or Cas12f1
                  organism = synthetic construct
variation          158..177
                  note = n is a, c, g or u

SEQUENCE: 168
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cgaaaagtaa cctcgaaac aaattcagtg 120
ctcctctcga aagaatagag caatgaagga atgcaacnnn nnnnnnnnnn nnnnnnttt 180
tattttt                                                    186

SEQ ID NO: 169      moltype = RNA length = 190
FEATURE            Location/Qualifiers
source             1..190
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of

```

-continued

```

MS1/MS2/MS3-3/MS5-3 for TnpB or Cas12f1
organism = synthetic construct
variation      162..181
               note = n is a, c, g or u

SEQUENCE: 169
accgcttcac caattagttg agtgaaggtg ggctgcttgc atcagcctaa tgcgcagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aattcagtcg tcctctccaa ttctgcacaa 120
gaaagttgca gaaccggaat agagcaatga aggaatgcaa cnnnnnnnnn nnnnnnnnnn 180
nttttatttt                                     190

SEQ ID NO: 170      moltype = RNA length = 185
FEATURE            Location/Qualifiers
source             1..185
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                    MS1/MS2/MS4*-2/MS5-3 for TnpB or Cas12f1
                    organism = synthetic construct
variation          157..176
                  note = n is a, c, g or u

SEQUENCE: 170
cttcactgat aaagtggaga accgcttcac caattagttg agtgaaggtg ggctgcttgc 60
atcagcctaa tgcgcagaag tgctttcttc ggaaagtaac cctcgaaaca aattcagtcg 120
tcctctcgaa agaatagagc aatgaaggaa tgcaacnnnn nnnnnnnnnn nnnnnntttt 180
atttt                                             185

SEQ ID NO: 171      moltype = RNA length = 156
FEATURE            Location/Qualifiers
source             1..156
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                    MS1/MS3-3/MS4*-2/MS5-3 for TnpB or Cas12f1
                    organism = synthetic construct
variation          137..156
                  note = n is a, c, g or u

SEQUENCE: 171
accgcttcac caattagttg agtgaaggtg ggctgcttgc atcagcctaa tgcgcagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aattcagtcg tcctctcgaa agaatagagc 120
aatgaaggaa tgcaacnnnn nnnnnnnnnn nnnnnn                                     156

SEQ ID NO: 172      moltype = RNA length = 165
FEATURE            Location/Qualifiers
source             1..165
                  mol_type = other RNA
                  note = Engineered sgRNA with modification of
                    MS1/MS2/MS3-3/MS4*-2/MS5-3 for TnpB or Cas12f1
                    organism = synthetic construct
variation          137..156
                  note = n is a, c, g or u

SEQUENCE: 172
accgcttcac caattagttg agtgaaggtg ggctgcttgc atcagcctaa tgcgcagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aattcagtcg tcctctcgaa agaatagagc 120
aatgaaggaa tgcaacnnnn nnnnnnnnnn nnnnnntttt atttt                                     165

SEQ ID NO: 173      moltype = RNA length = 181
FEATURE            Location/Qualifiers
source             1..181
                  mol_type = other RNA
                  note = Mature form sgRNA for TnpB or Cas12f1
                    organism = synthetic construct
variation          162..181
                  note = n is a, c, g or u

SEQUENCE: 173
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgctgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcatth gaaagaatga aggaatgcaa cnnnnnnnnn nnnnnnnnnn 180
n                                                  181

SEQ ID NO: 174      moltype = RNA length = 174
FEATURE            Location/Qualifiers
source             1..174
                  mol_type = other RNA
                  note = Mature form sgRNA with modification of MS3-1 for
                    TnpB or Cas12f1
                    organism = synthetic construct
variation          155..174
                  note = n is a, c, g or u

```

-continued

SEQUENCE: 174
gataaagtgg agaaccgctt caccaaaagc tgtcccttag gggattagaa cttgagttaa 60
gggtgggctgc ttgcatcagc ctaatgtcga gaagtgcctt cttcggaag taaccctcga 120
aacaattca tttgaaagaa tgaaggaatg caacnnnnnn nnnnnnnnn nnnn 174

SEQ ID NO: 175 moltype = RNA length = 167
FEATURE Location/Qualifiers
source 1..167
 mol_type = other RNA
 note = Mature form sgRNA with modification of MS3-2 for
 TnpB or Cas12f1
 organism = synthetic construct
variation 148..167
 note = n is a, c, g or u

SEQUENCE: 175
tggagaaccg cttcaccaaa agctgtccct taggggatta gaacttgagt gaagggtggc 60
tgcttgcatc agcctaagt ctagaagtcg tttcttcgga aagtaaccct cgaaacaaat 120
tcatttgaaa gaatgaagga atgcaacnnn nnnnnnnnn nnnnnnn 167

SEQ ID NO: 176 moltype = RNA length = 161
FEATURE Location/Qualifiers
source 1..161
 mol_type = other RNA
 note = Mature form sgRNA with modification of MS3-3 for
 TnpB or Cas12f1
 organism = synthetic construct
variation 142..161
 note = n is a, c, g or u

SEQUENCE: 176
accgcttcac caaaagctgt cccttagggg attagaactt gagtgaaggt gggctgcttg 60
catcagccta atgtcgagaa gtgctttctt cggaaagtaa ccctcgaaac aaattcattt 120
gaaagaatga aggaatgcaa cnnnnnnnnn nnnnnnnnn n 161

SEQ ID NO: 177 moltype = RNA length = 177
FEATURE Location/Qualifiers
source 1..177
 mol_type = other RNA
 note = Mature form sgRNA with modification of MS4-1 for
 TnpB or Cas12f1
 organism = synthetic construct
variation 158..177
 note = n is a, c, g or u

SEQUENCE: 177
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcattg aaatgaagga atgcaacnnn nnnnnnnnn nnnnnnn 177

SEQ ID NO: 178 moltype = RNA length = 173
FEATURE Location/Qualifiers
source 1..173
 mol_type = other RNA
 note = Mature form sgRNA with modification of MS4-2 for
 TnpB or Cas12f1
 organism = synthetic construct
variation 154..173
 note = n is a, c, g or u

SEQUENCE: 178
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaattcgaaa gaaggaatgc aacnnnnnnn nnnnnnnnn nnn 173

SEQ ID NO: 179 moltype = RNA length = 167
FEATURE Location/Qualifiers
source 1..167
 mol_type = other RNA
 note = Mature form sgRNA with modification of MS4-3 for
 TnpB or Cas12f1
 organism = synthetic construct
variation 148..167
 note = n is a, c, g or u

SEQUENCE: 179
cttcactgat aaagtggaga accgcttcac caaaagctgt cccttagggg attagaactt 60
gagtgaaggt gggctgcttg catcagccta atgtcgagaa gtgctttctt cggaaagtaa 120
ccctcgaaac aaagaaagga atgcaacnnn nnnnnnnnn nnnnnnn 167

SEQ ID NO: 180 moltype = RNA length = 175

-continued

FEATURE	Location/Qualifiers		
source	1..175		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS5-1 for TnpB or Cas12f1		
	organism = synthetic construct		
variation	156..175		
	note = n is a, c, g or u		
SEQUENCE: 180			
cttcactgat	aaagtggaga	accgcttcac caaaagctgt ttagattaga acttgagtga	60
aggtgggctg	cttgcacacg	cctaagtgtc agaagtgtt tcttcgaaa gtaaccctcg	120
aaacaaattc	atttgaaaga	atgaaggaat gcaacnnnnn nnnnnnnnnn nnnnn	175
SEQ ID NO: 181			
FEATURE	Location/Qualifiers		
source	1..170		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS5-2 for TnpB or Cas12f1		
	organism = synthetic construct		
variation	151..170		
	note = n is a, c, g or u		
SEQUENCE: 181			
cttcactgat	aaagtggaga	accgcttcac caaaagcttt agagaacttg agtgaagggtg	60
ggctgcttgc	atcagcctaa	tgtcgagaag tgctttcttc ggaaagtaac cctcgaaaca	120
aattcatttg	aaagaatgaa	ggaatgcaac nnnnnnnnnn nnnnnnnnnn	170
SEQ ID NO: 182			
FEATURE	Location/Qualifiers		
source	1..160		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS5-3 for TnpB or Cas12f1		
	organism = synthetic construct		
variation	141..160		
	note = n is a, c, g or u		
SEQUENCE: 182			
cttcactgat	aaagtggaga	accgcttcac caattagttg agtgaagggtg ggctgcttgc	60
atcagcctaa	tgtcgagaag	tgctttcttc ggaaagtaac cctcgaaaca aattcatttg	120
aaagaatgaa	ggaatgcaac	nnnnnnnnnn nnnnnnnnnn	160
SEQ ID NO: 183			
FEATURE	Location/Qualifiers		
source	1..147		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS3-3/MS4-3 for TnpB or Cas12f1		
	organism = synthetic construct		
variation	128..147		
	note = n is a, c, g or u		
SEQUENCE: 183			
accgcttcac	caaaagctgt	cccttagggg attagaactt gaggtaagggt ggctgcttgc	60
catcagccta	atgtcgagaa	gtgctttctt cggaagtaa cctcgaaac aaagaaagga	120
atgcaacnnn	nnnnnnnnnn	nnnnnnnn	147
SEQ ID NO: 184			
FEATURE	Location/Qualifiers		
source	1..140		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS3-3/MS5-3 for TnpB or Cas12f1		
	organism = synthetic construct		
variation	121..140		
	note = n is a, c, g or u		
SEQUENCE: 184			
accgcttcac	caattagttg	agtgaagggt ggctgcttgc atcagcctaa tgtcgagaag	60
tgctttcttc	ggaaagtaac	cctcgaaaca aattcatttg aaagaatgaa ggaatgcaac	120
nnnnnnnnnn	nnnnnnnnnn		140
SEQ ID NO: 185			
FEATURE	Location/Qualifiers		
source	1..146		
	mol_type = other RNA		
	note = Mature form sgRNA with modification of MS4-3/MS5-3 for TnpB or Cas12f1		
	organism = synthetic construct		

-continued

```

variation          127..146
                   note = n is a, c, g or u

SEQUENCE: 185
cttcactgat aaagtgagga accgcttcac caattagttg agtgaagggtg ggctgcttgc 60
atcagcctaa tgcgagaag tgctttcttc ggaaagtaac cctcgaaaca aagaaaggaa 120
tgcaacnnnn nnnnnnnnnn nnnnnn 146

SEQ ID NO: 186      moltype = RNA length = 126
FEATURE            Location/Qualifiers
source             1..126
                   mol_type = other RNA
                   note = Mature form sgRNA with modification of
                     MS3-3/MS4-3/MS5-3 for TnpB or Cas12f1
                   organism = synthetic construct

variation          107..126
                   note = n is a, c, g or u

SEQUENCE: 186
accgcttcac caattagttg agtgaagggtg ggctgcttgc atcagcctaa tgcgagaag 60
tgctttcttc ggaaagtaac cctcgaaaca aagaaaggaa tgcaacnnnn nnnnnnnnnn 120
nnnnnn 126

SEQ ID NO: 187      moltype = RNA length = 10
FEATURE            Location/Qualifiers
source             1..10
                   mol_type = other RNA
                   note = The additional sequence for TnpB or Cas12f1 gRNA
                   organism = synthetic construct

SEQUENCE: 187
ataaagggtga 10

SEQ ID NO: 188      moltype = RNA length = 37
FEATURE            Location/Qualifiers
source             1..37
                   mol_type = other RNA
                   note = The additional sequence for TnpB or Cas12f1 gRNA
                   organism = synthetic construct

SEQUENCE: 188
ctgatgagtc cgtgaggagc aaacagagtaa gctcgtc 37

SEQ ID NO: 189      moltype = RNA length = 37
FEATURE            Location/Qualifiers
source             1..37
                   mol_type = other RNA
                   note = The additional sequence for TnpB or Cas12f1 gRNA
                   organism = synthetic construct

SEQUENCE: 189
ctgctcgaat gagcaaagca ggagtgcctg agtagtc 37

SEQ ID NO: 190      moltype = DNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                   mol_type = other DNA
                   note = F102 -protospacer
                   organism = synthetic construct

SEQUENCE: 190
aggttgcttc ctcattagag 20

SEQ ID NO: 191      moltype = DNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                   mol_type = other DNA
                   note = F104 -protospacer
                   organism = synthetic construct

SEQUENCE: 191
atattcttag aatcggtcac 20

SEQ ID NO: 192      moltype = DNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                   mol_type = other DNA
                   note = F108 -protospacer
                   organism = synthetic construct

SEQUENCE: 192
ccaccaatct tctggttata 20

SEQ ID NO: 193      moltype = DNA length = 20

```

-continued

FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F109 -protospacer	
	organism = synthetic construct	
SEQUENCE: 193		
tagacaacag actcaagagc		20
SEQ ID NO: 194	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F110 -protospacer	
	organism = synthetic construct	
SEQUENCE: 194		
actgactact ccccaagtat		20
SEQ ID NO: 195	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F111 -protospacer	
	organism = synthetic construct	
SEQUENCE: 195		
gggccttatc tccagtttct		20
SEQ ID NO: 196	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F114 -protospacer	
	organism = synthetic construct	
SEQUENCE: 196		
taaacgcctg gctagtaaga		20
SEQ ID NO: 197	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F115 -protospacer	
	organism = synthetic construct	
SEQUENCE: 197		
taaaaacagt aatacctaac		20
SEQ ID NO: 198	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F118 -protospacer	
	organism = synthetic construct	
SEQUENCE: 198		
ccttgcttac tgcttattgc		20
SEQ ID NO: 199	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F119 -protospacer	
	organism = synthetic construct	
SEQUENCE: 199		
ttcagtacta gcaataagca		20
SEQ ID NO: 200	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F121 -protospacer	
	organism = synthetic construct	
SEQUENCE: 200		
tttaatgact ttgaaacagt		20
SEQ ID NO: 201	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F123 -protospacer	

-continued

SEQUENCE: 201	organism = synthetic construct	
aaacagtatt tcatgtctaa		20
SEQ ID NO: 202	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F124 -protospacer	
	organism = synthetic construct	
SEQUENCE: 202		
gacatgaaat actgtttcaa		20
SEQ ID NO: 203	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F128 -protospacer	
	organism = synthetic construct	
SEQUENCE: 203		
agaaaatatt gtatcttggg		20
SEQ ID NO: 204	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F130 -protospacer	
	organism = synthetic construct	
SEQUENCE: 204		
tttcagattg ttagtaaact		20
SEQ ID NO: 205	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F131 -protospacer	
	organism = synthetic construct	
SEQUENCE: 205		
tactgggagc agtttatcat		20
SEQ ID NO: 206	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F133 -protospacer	
	organism = synthetic construct	
SEQUENCE: 206		
atttccttag ggtccagctt		20
SEQ ID NO: 207	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F134 -protospacer	
	organism = synthetic construct	
SEQUENCE: 207		
aagctggacc ctagggaaat		20
SEQ ID NO: 208	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F135 -protospacer	
	organism = synthetic construct	
SEQUENCE: 208		
aaattccctt gaataggaag		20
SEQ ID NO: 209	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F136 -protospacer	
	organism = synthetic construct	
SEQUENCE: 209		
aatcagaaag aagatcttat		20

-continued

SEQ ID NO: 210	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F138 -protospacer	
	organism = synthetic construct	
SEQUENCE: 210		
ttcaagaaaa aacaaaggca		20
SEQ ID NO: 211	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F140 -protospacer	
	organism = synthetic construct	
SEQUENCE: 211		
ccacttccac aatgtatatg		20
SEQ ID NO: 212	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F141 -protospacer	
	organism = synthetic construct	
SEQUENCE: 212		
acttaagtta cttgtccagg		20
SEQ ID NO: 213	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F142 -protospacer	
	organism = synthetic construct	
SEQUENCE: 213		
ctcatttcca tgcctggaca		20
SEQ ID NO: 214	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F143 -protospacer	
	organism = synthetic construct	
SEQUENCE: 214		
aaaaattggtt aaatgtatat		20
SEQ ID NO: 215	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F147 -protospacer	
	organism = synthetic construct	
SEQUENCE: 215		
attgaagagt aacaatttga		20
SEQ ID NO: 216	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F148 -protospacer	
	organism = synthetic construct	
SEQUENCE: 216		
acttattggtt attgaaattg		20
SEQ ID NO: 217	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F43 -protospacer	
	organism = synthetic construct	
SEQUENCE: 217		
gtatcaattc acaccagcaa		20
SEQ ID NO: 218	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	

-continued

	note = R52 -protospacer organism = synthetic construct	
SEQUENCE: 218 ctctcctaga ccatttccca		20
SEQ ID NO: 219 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R55 -protospacer organism = synthetic construct	
SEQUENCE: 219 ctgagagaga aacagttgcc		20
SEQ ID NO: 220 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R56 -protospacer organism = synthetic construct	
SEQUENCE: 220 tatccttgat tatacttagg		20
SEQ ID NO: 221 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R57 -protospacer organism = synthetic construct	
SEQUENCE: 221 cattaattta tatccttgat		20
SEQ ID NO: 222 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R58 -protospacer organism = synthetic construct	
SEQUENCE: 222 ttatttgcac taatttatat		20
SEQ ID NO: 223 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R59 -protospacer organism = synthetic construct	
SEQUENCE: 223 actgagacaa ctattcttgt		20
SEQ ID NO: 224 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R60 -protospacer organism = synthetic construct	
SEQUENCE: 224 gtaaatttaa ctgagacaac		20
SEQ ID NO: 225 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R61 -protospacer organism = synthetic construct	
SEQUENCE: 225 aaataccagg ttgtttagta		20
SEQ ID NO: 226 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R62 -protospacer organism = synthetic construct	
SEQUENCE: 226 taccaaataa gtcactcaac		20

-continued

SEQ ID NO: 227	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R63 -protospacer	
	organism = synthetic construct	
SEQUENCE: 227		
aatgtaaata gctcagttga		20
SEQ ID NO: 228	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R64 -protospacer	
	organism = synthetic construct	
SEQUENCE: 228		
gcactacgca gcccaacata		20
SEQ ID NO: 229	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R65 -protospacer	
	organism = synthetic construct	
SEQUENCE: 229		
attctgcaat atgttgggct		20
SEQ ID NO: 230	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R66 -protospacer	
	organism = synthetic construct	
SEQUENCE: 230		
ccttctccaa tgataaggat		20
SEQ ID NO: 231	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R68 -protospacer	
	organism = synthetic construct	
SEQUENCE: 231		
agttatagct ctctttcaat		20
SEQ ID NO: 232	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R69 -protospacer	
	organism = synthetic construct	
SEQUENCE: 232		
aaattaccct agatcttaaa		20
SEQ ID NO: 233	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R70 -protospacer	
	organism = synthetic construct	
SEQUENCE: 233		
tgcaatgccca tgttcaaatg		20
SEQ ID NO: 234	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R71 -protospacer	
	organism = synthetic construct	
SEQUENCE: 234		
aacatggcat tgcataaatg		20
SEQ ID NO: 235	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	

-continued

	mol_type = other DNA note = R73 -protospacer organism = synthetic construct	
SEQUENCE: 235 ccaaatggat tacactgagg		20
SEQ ID NO: 236 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R74 -protospacer organism = synthetic construct	
SEQUENCE: 236 gtaaaattat gacatcaact		20
SEQ ID NO: 237 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R75 -protospacer organism = synthetic construct	
SEQUENCE: 237 cctagctaga tcaaatatac		20
SEQ ID NO: 238 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R76 -protospacer organism = synthetic construct	
SEQUENCE: 238 atctagctag gtaaaccata		20
SEQ ID NO: 239 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R77 -protospacer organism = synthetic construct	
SEQUENCE: 239 tatttgtgaa tgattaagaa		20
SEQ ID NO: 240 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R79 -protospacer organism = synthetic construct	
SEQUENCE: 240 atgtataaca attccaacat		20
SEQ ID NO: 241 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R82 -protospacer organism = synthetic construct	
SEQUENCE: 241 gaaaggcttg aaagctgtta		20
SEQ ID NO: 242 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R84 -protospacer organism = synthetic construct	
SEQUENCE: 242 tatttcttta gaaaggcttg		20
SEQ ID NO: 243 FEATURE source	moltype = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R85 -protospacer organism = synthetic construct	
SEQUENCE: 243		

-continued

atacctaaat accttcagca	20
SEQ ID NO: 244	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R86 -protospacer
	organism = synthetic construct
SEQUENCE: 244	
aaaaaacaag aagtgaggca	20
SEQ ID NO: 245	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R89 -protospacer
	organism = synthetic construct
SEQUENCE: 245	
taaaataaac ttcaccaatt	20
SEQ ID NO: 246	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R90 -protospacer
	organism = synthetic construct
SEQUENCE: 246	
caaagaaatt ccccttactc	20
SEQ ID NO: 247	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R91 -protospacer
	organism = synthetic construct
SEQUENCE: 247	
ttcctatgga attggtgaag	20
SEQ ID NO: 248	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R92 -protospacer
	organism = synthetic construct
SEQUENCE: 248	
aaggccttga gcttgaatac	20
SEQ ID NO: 249	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R93 -protospacer
	organism = synthetic construct
SEQUENCE: 249	
aatgcaggtt gtttactatc	20
SEQ ID NO: 250	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R94 -protospacer
	organism = synthetic construct
SEQUENCE: 250	
caaatgatga ttgtggtcaa	20
SEQ ID NO: 251	moltype = DNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other DNA
	note = R95 -protospacer
	organism = synthetic construct
SEQUENCE: 251	
caaatggcac tgaattcagg	20
SEQ ID NO: 252	moltype = DNA length = 20
FEATURE	Location/Qualifiers

-continued

source	1..20 mol_type = other DNA note = R97 -protospacer organism = synthetic construct	
SEQUENCE: 252		
tatacatcca tatccataac		20
SEQ ID NO: 253	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = R98 -protospacer organism = synthetic construct	
SEQUENCE: 253		
catatattta tatacatcca		20
SEQ ID NO: 254	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = R99 -protospacer organism = synthetic construct	
SEQUENCE: 254		
cttctgcttt aaaaaaagta		20
SEQ ID NO: 255	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F1 -protospacer organism = synthetic construct	
SEQUENCE: 255		
tcatcccaat ggcatattta		20
SEQ ID NO: 256	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F2 -protospacer organism = synthetic construct	
SEQUENCE: 256		
gtcatcatga accattctct		20
SEQ ID NO: 257	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F3 -protospacer organism = synthetic construct	
SEQUENCE: 257		
agtctcaggc cctggccttg		20
SEQ ID NO: 258	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F4 -protospacer organism = synthetic construct	
SEQUENCE: 258		
gaaaaagaca gaaaggaaga		20
SEQ ID NO: 259	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F5 -protospacer organism = synthetic construct	
SEQUENCE: 259		
ttgaaattta gtgtgacatg		20
SEQ ID NO: 260	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20 mol_type = other DNA note = F6 -protospacer organism = synthetic construct	

-continued

SEQUENCE: 260		
gtgtgacatg atcatttcca		20
SEQ ID NO: 261	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F7 -protospacer	
	organism = synthetic construct	
SEQUENCE: 261		
accctcaaat aagagcagag		20
SEQ ID NO: 262	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F8 -protospacer	
	organism = synthetic construct	
SEQUENCE: 262		
agggtcaaat tttcaataag		20
SEQ ID NO: 263	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F9 -protospacer	
	organism = synthetic construct	
SEQUENCE: 263		
aaaagtgact tattgaaat		20
SEQ ID NO: 264	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F10 -protospacer	
	organism = synthetic construct	
SEQUENCE: 264		
aaaaagcatt ttctgacact		20
SEQ ID NO: 265	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F11 -protospacer	
	organism = synthetic construct	
SEQUENCE: 265		
ggaggatttc aagcttatgg		20
SEQ ID NO: 266	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F12 -protospacer	
	organism = synthetic construct	
SEQUENCE: 266		
ggaacactcc atccacattt		20
SEQ ID NO: 267	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F13 -protospacer	
	organism = synthetic construct	
SEQUENCE: 267		
ccagtcagcc tgggtgtctg		20
SEQ ID NO: 268	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F14 -protospacer	
	organism = synthetic construct	
SEQUENCE: 268		
atttagctct cttttcatcc		20
SEQ ID NO: 269	moltype = DNA length = 20	

-continued

FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F15 -protospacer	
	organism = synthetic construct	
SEQUENCE: 269		
gctctctttt catcctcaca		20
SEQ ID NO: 270	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F16 -protospacer	
	organism = synthetic construct	
SEQUENCE: 270		
gctcctttcc tggaatccct		20
SEQ ID NO: 271	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F17 -protospacer	
	organism = synthetic construct	
SEQUENCE: 271		
cttccatgat ttcttctcct		20
SEQ ID NO: 272	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F18 -protospacer	
	organism = synthetic construct	
SEQUENCE: 272		
tgtcacagtc caggagaaga		20
SEQ ID NO: 273	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F19 -protospacer	
	organism = synthetic construct	
SEQUENCE: 273		
cattacattt aattttttct		20
SEQ ID NO: 274	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F20 -protospacer	
	organism = synthetic construct	
SEQUENCE: 274		
attctttcta gaaagagcct		20
SEQ ID NO: 275	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F21 -protospacer	
	organism = synthetic construct	
SEQUENCE: 275		
tttctcaaag ctatctgact		20
SEQ ID NO: 276	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F22 -protospacer	
	organism = synthetic construct	
SEQUENCE: 276		
ttgcctcact gttactgect		20
SEQ ID NO: 277	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F23 -protospacer	

-continued

SEQUENCE: 277	organism = synthetic construct	
tgtcacgaaa caatgattga		20
SEQ ID NO: 278	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F24 -protospacer	
	organism = synthetic construct	
SEQUENCE: 278		
attgtcagag agaataaaaa		20
SEQ ID NO: 279	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F25 -protospacer	
	organism = synthetic construct	
SEQUENCE: 279		
ttctctctga caattaaaac		20
SEQ ID NO: 280	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = F26 -protospacer	
	organism = synthetic construct	
SEQUENCE: 280		
atcaatgcag acagaaaaaa		20
SEQ ID NO: 281	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R1 -protospacer	
	organism = synthetic construct	
SEQUENCE: 281		
acatctatga gcctcagtta		20
SEQ ID NO: 282	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R2 -protospacer	
	organism = synthetic construct	
SEQUENCE: 282		
aatccttggt ctgctactta		20
SEQ ID NO: 283	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R3 -protospacer	
	organism = synthetic construct	
SEQUENCE: 283		
attcctaagc ttgtggttatt		20
SEQ ID NO: 284	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R4 -protospacer	
	organism = synthetic construct	
SEQUENCE: 284		
gtgatttgta tgtagatgta		20
SEQ ID NO: 285	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R5 -protospacer	
	organism = synthetic construct	
SEQUENCE: 285		
tatgtagatg tagatgtagt		20

-continued

SEQ ID NO: 286	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R6 -protospacer	
	organism = synthetic construct	
SEQUENCE: 286		
tggttgctat gtactgatac		20
SEQ ID NO: 287	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R7 -protospacer	
	organism = synthetic construct	
SEQUENCE: 287		
caataccata ttgagttata		20
SEQ ID NO: 288	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R8 -protospacer	
	organism = synthetic construct	
SEQUENCE: 288		
gtaaataaaa gtctctgggag		20
SEQ ID NO: 289	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R9 -protospacer	
	organism = synthetic construct	
SEQUENCE: 289		
tttaccaaaag gaaacaatat		20
SEQ ID NO: 290	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R10 -protospacer	
	organism = synthetic construct	
SEQUENCE: 290		
ccaaaggaaa caatatattta		20
SEQ ID NO: 291	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R11 -protospacer	
	organism = synthetic construct	
SEQUENCE: 291		
aacattataa aatattgttt		20
SEQ ID NO: 292	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R12 -protospacer	
	organism = synthetic construct	
SEQUENCE: 292		
taatgtttaa agcccaggtt		20
SEQ ID NO: 293	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R13 -protospacer	
	organism = synthetic construct	
SEQUENCE: 293		
aagcccaggt tttgaagtta		20
SEQ ID NO: 294	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	

-continued

	note = R14 -protospacer organism = synthetic construct	
SEQUENCE: 294 aaccagaca atgtaacttc		20
SEQ ID NO: 295 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R15 -protospacer organism = synthetic construct	
SEQUENCE: 295 aagttacatt gtctgggttc		20
SEQ ID NO: 296 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R16 -protospacer organism = synthetic construct	
SEQUENCE: 296 tagctagata aacttgggct		20
SEQ ID NO: 297 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R17 -protospacer organism = synthetic construct	
SEQUENCE: 297 tctagctata aaatggggat		20
SEQ ID NO: 298 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R18 -protospacer organism = synthetic construct	
SEQUENCE: 298 tcacaggat taagttgggt		20
SEQ ID NO: 299 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R19 -protospacer organism = synthetic construct	
SEQUENCE: 299 ttcaatacta gcagtaagca		20
SEQ ID NO: 300 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R20 -protospacer organism = synthetic construct	
SEQUENCE: 300 ccttgcttac tgcttactgc		20
SEQ ID NO: 301 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R21 -protospacer organism = synthetic construct	
SEQUENCE: 301 gttccaccac gaactctaga		20
SEQ ID NO: 302 FEATURE source	mol_type = DNA length = 20 Location/Qualifiers 1..20 mol_type = other DNA note = R22 -protospacer organism = synthetic construct	
SEQUENCE: 302 cataatacaa atgccatcat		20

-continued

SEQ ID NO: 303	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R23 -protospacer	
	organism = synthetic construct	
SEQUENCE: 303		
tattatgcaa actgtatatc		20
SEQ ID NO: 304	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R24 -protospacer	
	organism = synthetic construct	
SEQUENCE: 304		
tgagtcacgc atttagggaa		20
SEQ ID NO: 305	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R25 -protospacer	
	organism = synthetic construct	
SEQUENCE: 305		
gggaagccat tgatgtgctc		20
SEQ ID NO: 306	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R26 -protospacer	
	organism = synthetic construct	
SEQUENCE: 306		
gagaactctg gagactactg		20
SEQ ID NO: 307	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R27 -protospacer	
	organism = synthetic construct	
SEQUENCE: 307		
ttagagaac tctggagact		20
SEQ ID NO: 308	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R28 -protospacer	
	organism = synthetic construct	
SEQUENCE: 308		
atgccccctc acagagatcg		20
SEQ ID NO: 309	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R29 -protospacer	
	organism = synthetic construct	
SEQUENCE: 309		
aagacgatct ctgtgagggg		20
SEQ ID NO: 310	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = R30 -protospacer	
	organism = synthetic construct	
SEQUENCE: 310		
acctatattt aaagacgac		20
SEQ ID NO: 311	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	

-continued

	mol_type = other DNA note = R31 -protospacer organism = synthetic construct	
SEQUENCE: 311 aatataggtc aaaaactaat		20
SEQ ID NO: 312 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = Spacer 1 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 312 ctcattctca tgctggac		19
SEQ ID NO: 313 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = Spacer 2 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 313 ctcattctca tgctggaca		20
SEQ ID NO: 314 FEATURE source	moltype = RNA length = 21 Location/Qualifiers 1..21 mol_type = other RNA note = Spacer 3 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 314 ctcattctca tgctggaca a		21
SEQ ID NO: 315 FEATURE source	moltype = RNA length = 22 Location/Qualifiers 1..22 mol_type = other RNA note = Spacer 4 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 315 ctcattctca tgctggaca ag		22
SEQ ID NO: 316 FEATURE source	moltype = RNA length = 24 Location/Qualifiers 1..24 mol_type = other RNA note = Spacer 5 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 316 ctcattctca tgctggaca agta		24
SEQ ID NO: 317 FEATURE source	moltype = RNA length = 25 Location/Qualifiers 1..25 mol_type = other RNA note = Spacer 6 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 317 ctcattctca tgctggaca agtaa		25
SEQ ID NO: 318 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = Spacer 7 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 318 ctcattctca tgctggac		19
SEQ ID NO: 319 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = Spacer 8 for dystrophin(w/o U-tail) organism = synthetic construct	
SEQUENCE: 319		

-continued

ctcattctca tgcctggaca	20
SEQ ID NO: 320	moltype = RNA length = 21
FEATURE	Location/Qualifiers
source	1..21
	mol_type = other RNA
	note = Spacer 9 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 320	
ctcattctca tgcctggaca a	21
SEQ ID NO: 321	moltype = RNA length = 22
FEATURE	Location/Qualifiers
source	1..22
	mol_type = other RNA
	note = Spacer 10 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 321	
ctcattctca tgcctggaca ag	22
SEQ ID NO: 322	moltype = RNA length = 24
FEATURE	Location/Qualifiers
source	1..24
	mol_type = other RNA
	note = Spacer 11 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 322	
ctcattctca tgcctggaca agta	24
SEQ ID NO: 323	moltype = RNA length = 25
FEATURE	Location/Qualifiers
source	1..25
	mol_type = other RNA
	note = Spacer 12 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 323	
ctcattctca tgcctggaca agtaa	25
SEQ ID NO: 324	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = Spacer 13 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 324	
ctctcctaga ccatttccc	19
SEQ ID NO: 325	moltype = RNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other RNA
	note = Spacer 14 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 325	
ctctcctaga ccatttccca	20
SEQ ID NO: 326	moltype = RNA length = 21
FEATURE	Location/Qualifiers
source	1..21
	mol_type = other RNA
	note = Spacer 15 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 326	
ctctcctaga ccatttccca c	21
SEQ ID NO: 327	moltype = RNA length = 22
FEATURE	Location/Qualifiers
source	1..22
	mol_type = other RNA
	note = Spacer 16 for dystrophin(w/o U-tail)
	organism = synthetic construct
SEQUENCE: 327	
ctctcctaga ccatttccca cc	22
SEQ ID NO: 328	moltype = RNA length = 23
FEATURE	Location/Qualifiers

-continued

```

source          1..23
                 mol_type = other RNA
                 note = Spacer 17 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 328
ctctcctaga ccatttccca cca                               23

SEQ ID NO: 329      moltype = RNA  length = 24
FEATURE            Location/Qualifiers
source            1..24
                 mol_type = other RNA
                 note = Spacer 18 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 329
ctctcctaga ccatttccca ccag                               24

SEQ ID NO: 330      moltype = RNA  length = 19
FEATURE            Location/Qualifiers
source            1..19
                 mol_type = other RNA
                 note = Spacer 19 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 330
ctctcctaga ccatttccc                                     19

SEQ ID NO: 331      moltype = RNA  length = 20
FEATURE            Location/Qualifiers
source            1..20
                 mol_type = other RNA
                 note = Spacer 20 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 331
ctctcctaga ccatttccca                                     20

SEQ ID NO: 332      moltype = RNA  length = 21
FEATURE            Location/Qualifiers
source            1..21
                 mol_type = other RNA
                 note = Spacer 21 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 332
ctctcctaga ccatttccca c                                   21

SEQ ID NO: 333      moltype = RNA  length = 22
FEATURE            Location/Qualifiers
source            1..22
                 mol_type = other RNA
                 note = Spacer 22 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 333
ctctcctaga ccatttccca cc                                  22

SEQ ID NO: 334      moltype = RNA  length = 23
FEATURE            Location/Qualifiers
source            1..23
                 mol_type = other RNA
                 note = Spacer 23 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 334
ctctcctaga ccatttccca cca                                23

SEQ ID NO: 335      moltype = RNA  length = 24
FEATURE            Location/Qualifiers
source            1..24
                 mol_type = other RNA
                 note = Spacer 24 for dystrophin(w/o U-tail)
                 organism = synthetic construct

SEQUENCE: 335
ctctcctaga ccatttccca ccag                               24

SEQ ID NO: 336      moltype = RNA  length = 28
FEATURE            Location/Qualifiers
source            1..28
                 mol_type = other RNA
                 note = Spacer 1 for dystrophin(w/ U-tail)
                 organism = synthetic construct

```

-continued

SEQUENCE: 336
ctcattctca tgcctggact tttatttt 28

SEQ ID NO: 337 moltype = RNA length = 29
FEATURE Location/Qualifiers
source 1..29
 mol_type = other RNA
 note = Spacer 2 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 337
ctcattctca tgcctggaca tttattttt 29

SEQ ID NO: 338 moltype = RNA length = 30
FEATURE Location/Qualifiers
source 1..30
 mol_type = other RNA
 note = Spacer 3 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 338
ctcattctca tgcctggaca attttatttt 30

SEQ ID NO: 339 moltype = RNA length = 31
FEATURE Location/Qualifiers
source 1..31
 mol_type = other RNA
 note = Spacer 4 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 339
ctcattctca tgcctggaca agttttattt t 31

SEQ ID NO: 340 moltype = RNA length = 33
FEATURE Location/Qualifiers
source 1..33
 mol_type = other RNA
 note = Spacer 5 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 340
ctcattctca tgcctggaca agtatatttat ttt 33

SEQ ID NO: 341 moltype = RNA length = 34
FEATURE Location/Qualifiers
source 1..34
 mol_type = other RNA
 note = Spacer 6 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 341
ctcattctca tgcctggaca agtaatttta tttt 34

SEQ ID NO: 342 moltype = RNA length = 23
FEATURE Location/Qualifiers
source 1..23
 mol_type = other RNA
 note = Spacer 7 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 342
ctcattctca tgcctggact ttt 23

SEQ ID NO: 343 moltype = RNA length = 24
FEATURE Location/Qualifiers
source 1..24
 mol_type = other RNA
 note = Spacer 8 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 343
ctcattctca tgcctggaca tttt 24

SEQ ID NO: 344 moltype = RNA length = 25
FEATURE Location/Qualifiers
source 1..25
 mol_type = other RNA
 note = Spacer 9 for dystrophin(w/ U-tail)
 organism = synthetic construct

SEQUENCE: 344
ctcattctca tgcctggaca atttt 25

SEQ ID NO: 345 moltype = RNA length = 26

-continued

FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other RNA	
	note = Spacer 10 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 345		
ctcattctca tgcctggaca agtttt		26
SEQ ID NO: 346	moltype = RNA length = 28	
FEATURE	Location/Qualifiers	
source	1..28	
	mol_type = other RNA	
	note = Spacer 11 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 346		
ctcattctca tgcctggaca agtatattt		28
SEQ ID NO: 347	moltype = RNA length = 29	
FEATURE	Location/Qualifiers	
source	1..29	
	mol_type = other RNA	
	note = Spacer 12 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 347		
ctcattctca tgcctggaca agtaatttt		29
SEQ ID NO: 348	moltype = RNA length = 28	
FEATURE	Location/Qualifiers	
source	1..28	
	mol_type = other RNA	
	note = Spacer 13 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 348		
ctctcctaga ccatttcctt tttatttt		28
SEQ ID NO: 349	moltype = RNA length = 29	
FEATURE	Location/Qualifiers	
source	1..29	
	mol_type = other RNA	
	note = Spacer 14 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 349		
ctctcctaga ccatttcctt ttttatttt		29
SEQ ID NO: 350	moltype = RNA length = 30	
FEATURE	Location/Qualifiers	
source	1..30	
	mol_type = other RNA	
	note = Spacer 15 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 350		
ctctcctaga ccatttcctt cttttatttt		30
SEQ ID NO: 351	moltype = RNA length = 31	
FEATURE	Location/Qualifiers	
source	1..31	
	mol_type = other RNA	
	note = Spacer 16 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 351		
ctctcctaga ccatttcctt cttttatttt t		31
SEQ ID NO: 352	moltype = RNA length = 32	
FEATURE	Location/Qualifiers	
source	1..32	
	mol_type = other RNA	
	note = Spacer 17 for dystrophin(w/ U-tail)	
	organism = synthetic construct	
SEQUENCE: 352		
ctctcctaga ccatttcctt ccattttatt tt		32
SEQ ID NO: 353	moltype = RNA length = 33	
FEATURE	Location/Qualifiers	
source	1..33	
	mol_type = other RNA	
	note = Spacer 18 for dystrophin(w/ U-tail)	

-continued

```

                                organism = synthetic construct
SEQUENCE: 353
ctctcctaga ccatttccca ccagttttat ttt                                     33

SEQ ID NO: 354      moltype = RNA  length = 23
FEATURE            Location/Qualifiers
source             1..23
                   mol_type = other RNA
                   note = Spacer 19 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 354
ctctcctaga ccatttcctt ttt                                               23

SEQ ID NO: 355      moltype = RNA  length = 24
FEATURE            Location/Qualifiers
source             1..24
                   mol_type = other RNA
                   note = Spacer 20 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 355
ctctcctaga ccatttccca tttt                                             24

SEQ ID NO: 356      moltype = RNA  length = 25
FEATURE            Location/Qualifiers
source             1..25
                   mol_type = other RNA
                   note = Spacer 21 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 356
ctctcctaga ccatttccca ctttt                                           25

SEQ ID NO: 357      moltype = RNA  length = 26
FEATURE            Location/Qualifiers
source             1..26
                   mol_type = other RNA
                   note = Spacer 22 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 357
ctctcctaga ccatttccca cttttt                                          26

SEQ ID NO: 358      moltype = RNA  length = 27
FEATURE            Location/Qualifiers
source             1..27
                   mol_type = other RNA
                   note = Spacer 23 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 358
ctctcctaga ccatttccca ccatttt                                          27

SEQ ID NO: 359      moltype = RNA  length = 28
FEATURE            Location/Qualifiers
source             1..28
                   mol_type = other RNA
                   note = Spacer 24 for dystrophin(w/ U-tail)
                   organism = synthetic construct
SEQUENCE: 359
ctctcctaga ccatttccca ccagtttt                                          28

SEQ ID NO: 360      moltype = RNA  length = 47
FEATURE            Location/Qualifiers
source             1..47
                   mol_type = other RNA
                   note = shRNA for ATM1-1
                   organism = synthetic construct
SEQUENCE: 360
ggagccagat agtttgtatt tcaagagaat acaaactatc tggctcc                  47

SEQ ID NO: 361      moltype = RNA  length = 47
FEATURE            Location/Qualifiers
source             1..47
                   mol_type = other RNA
                   note = shRNA for ATM1-2
                   organism = synthetic construct
SEQUENCE: 361
gcaagcagct gaaacaaatt tcaagagaat ttgtttcagc tgcttgc                  47

```

-continued

SEQ ID NO: 362	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for ATM1-3	
	organism = synthetic construct	
SEQUENCE: 362		
ggagctgatt gtagcaacat tcaagagatg ttgctacaat cagctcc		47
SEQ ID NO: 363	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for ATM1-4	
	organism = synthetic construct	
SEQUENCE: 363		
gcacagaagt gcctccaatt tcaagagaat tggaggcact tctgtgc		47
SEQ ID NO: 364	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for ATM1-5	
	organism = synthetic construct	
SEQUENCE: 364		
ggacatagtt tctgggagat tcaagagatc tcccagaaac tatgtcc		47
SEQ ID NO: 365	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC4-1	
	organism = synthetic construct	
SEQUENCE: 365		
ggatgacact ggcacattat tcaagagata atgtgccagt gtcattc		47
SEQ ID NO: 366	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC4-2	
	organism = synthetic construct	
SEQUENCE: 366		
ggagagtact gatgaggaaat tcaagagatt cctcatcagt actctcc		47
SEQ ID NO: 367	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC4-3	
	organism = synthetic construct	
SEQUENCE: 367		
gaatccacct tgtttctgat tcaagagatc agaaacaagg tggattc		47
SEQ ID NO: 368	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC4-4	
	organism = synthetic construct	
SEQUENCE: 368		
gtacaagtat cttgggagat tcaagagatc tccaagata cttgtac		47
SEQ ID NO: 369	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC4-5	
	organism = synthetic construct	
SEQUENCE: 369		
gaatgcagct caagaacgat tcaagagatc gttcttgagc tgcattc		47
SEQ ID NO: 370	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	

-continued

	note = shRNA for XLF-1 organism = synthetic construct	
SEQUENCE: 370		
gcatgagtct ggcattacat tcaagagatg taatgccaga ctcatgc		47
SEQ ID NO: 371	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XLF-2	
	organism = synthetic construct	
SEQUENCE: 371		
gaaagccctt tgtcatgaat tcaagagatt catgacaaaag ggctttc		47
SEQ ID NO: 372	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XLF-3	
	organism = synthetic construct	
SEQUENCE: 372		
gaacagtgtc tccttgcaat tcaagagatt gcaggggaagc actgttc		47
SEQ ID NO: 373	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XLF-4	
	organism = synthetic construct	
SEQUENCE: 373		
ggaaagacct agagatccat tcaagagatg gatctctagg tctttcc		47
SEQ ID NO: 374	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XLF-5	
	organism = synthetic construct	
SEQUENCE: 374		
gtatggcagt caccacacat tcaagagatg tgtggtgact gccatac		47
SEQ ID NO: 375	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC6-1	
	organism = synthetic construct	
SEQUENCE: 375		
gcagcattgt gcagatacat tcaagagatg tatctgcaca atgctgc		47
SEQ ID NO: 376	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC6-2	
	organism = synthetic construct	
SEQUENCE: 376		
gcaggaacat ccctccttat tcaagagata aggagggatg ttcctgc		47
SEQ ID NO: 377	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC6-3	
	organism = synthetic construct	
SEQUENCE: 377		
gcagtgtctc gctcatcaat tcaagagatt gatgagcaga gcactgc		47
SEQ ID NO: 378	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC6-4	
	organism = synthetic construct	
SEQUENCE: 378		
ggatcatgct gttcaccaat tcaagagatt ggtgaacagc atgatcc		47

-continued

SEQ ID NO: 379	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for XRCC6-5	
	organism = synthetic construct	
SEQUENCE: 379		
ggatctgact actcactcat tcaagagatg agtgagtagt cagatcc		47
SEQ ID NO: 380	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for LIG4-1	
	organism = synthetic construct	
SEQUENCE: 380		
gcacaaagat ggagatgtat tcaagagata catctccatc tttgtgc		47
SEQ ID NO: 381	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for LIG4-2	
	organism = synthetic construct	
SEQUENCE: 381		
gcagacacgt actgtgtaat tcaagagatt acacagtacg tgtctgc		47
SEQ ID NO: 382	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for LIG4-3	
	organism = synthetic construct	
SEQUENCE: 382		
ggagcagact cctgaagaat tcaagagatt cttcaggagt ctgctcc		47
SEQ ID NO: 383	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for LIG4-4	
	organism = synthetic construct	
SEQUENCE: 383		
ggaggattct gatctgcaat tcaagagatt gcagatcaga atcctcc		47
SEQ ID NO: 384	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for LIG4-5	
	organism = synthetic construct	
SEQUENCE: 384		
gcatgatcct tctgtaggat tcaagagatc ctacagaagg atcatgc		47
SEQ ID NO: 385	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for DCLRE1C-1	
	organism = synthetic construct	
SEQUENCE: 385		
ggagactcct acccagatat tcaagagata tctgggtagg agtctcc		47
SEQ ID NO: 386	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	
	mol_type = other RNA	
	note = shRNA for DCLRE1C-2	
	organism = synthetic construct	
SEQUENCE: 386		
ggacaaagct gactacagat tcaagagatc tgtagtacgc tttgtcc		47
SEQ ID NO: 387	moltype = RNA length = 47	
FEATURE	Location/Qualifiers	
source	1..47	

-continued

```

mol_type = other RNA
note = shRNA for DCLRE1C-3
organism = synthetic construct

SEQUENCE: 387
gcagagctct cgtttcacat tcaagagatg tgaaacgaga gctctgc 47

SEQ ID NO: 388      moltype = RNA length = 47
FEATURE            Location/Qualifiers
source              1..47
                    mol_type = other RNA
                    note = shRNA for DCLRE1C-4
                    organism = synthetic construct

SEQUENCE: 388
ggactctgat ggagaaatcat tcaagagatg attctccatc agagtcc 47

SEQ ID NO: 389      moltype = RNA length = 47
FEATURE            Location/Qualifiers
source              1..47
                    mol_type = other RNA
                    note = shRNA for DCLRE1C-5
                    organism = synthetic construct

SEQUENCE: 389
gcagaattct tcccagtcac tcaagagatg actgggaaga attctgc 47

SEQ ID NO: 390      moltype = AA length = 249
FEATURE            Location/Qualifiers
source              1..249
                    mol_type = protein
                    note = N-term 249aa extension
                    organism = synthetic construct

SEQUENCE: 390
MVSKGEELFT GVVPIVELD GDVNGHKFSV SGEGEDATY GKLTCLKFICT TGKLPVPWPT 60
LVTTLTYGVO CFSRYPDHMK QHDFPKSAMP EGYVQERTIF FKDDGNYKTR AEVKFEGDTL 120
VNRIELKGID FKEDGNILGH KLEYNYNVSHN VYIMADKQKN GIKVNFKIRH NIEDGSVQLA 180
DHYQQNTPIG DGPVLLPDNH YLSTQSALSK DPNEKRDHNV LLEFVTAAGI TLGMDELYKS 240
GLRSRAQAL 249

SEQ ID NO: 391      moltype = AA length = 248
FEATURE            Location/Qualifiers
source              1..248
                    mol_type = protein
                    note = C-term 248aa extension
                    organism = synthetic construct

SEQUENCE: 391
RARDPPVATM VSKGEELFTG VVPIVELDGD DVNGHKFSVS GEGEGDATYG KLTCLKFICTT 60
GKLPVPWPTL VTTLTYGVQC FSRYPDHMKQ HDPFKSAMPE GYVQERTIFF KDDGNYKTRA 120
EVKFEGDTLV NRIELKGIDF KEDGNILGHK LEYNYNVSHN VYIMADKQKNG IKVNFKIRHN 180
IEDGSVQLAD HYQQNTPIGD GPVLLPDNH YLSTQSALSKD PNEKRDHNV LLEFVTAAGIT 240
LGMDELYK 248

SEQ ID NO: 392      moltype = AA length = 557
FEATURE            Location/Qualifiers
source              1..557
                    mol_type = protein
                    note = Engineered CWCas12f1 (S170T)
                    organism = synthetic construct

SEQUENCE: 392
MGEKSSRRRR NGKSGAWTAA ITSCVGGKMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60
NREKIALEKN KDKVKEACSK HLKVAAYCTT QVERNACLFC KARKLDDKFY QKLRGQFPDA 120
VFWQEISEIF RQLQQAQAEI YNQSLIELYY EIFIKGKGIA NASSVEHYLT DVCYTRAAEL 180
FKNAAIASGL RSKIKSNFRL KELKNMKSGL PTTKSDNFPI PLVKQKGGQY TGFEISNHNS 240
DFIIKIPFGR WQVKKEIDKY RPWEKFDPEQ VQKSPKPISL LLSTQRRKRN KGWSKDEGTE 300
AEIKKVMNGD YQTSYIEVFR GSKIGESAW MLNLSIDVPK IDKGVDPSII GGIDVGKSP 360
LVCAINNAFS RYSISDNDLF HFNNKMFARR RILLKKNRHK RAGHGAKNKL KPITILTEKS 420
ERFRKKLIER WACEIADFFI KNKVGTVMQE NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
EPFLKQYQIE IRKVAPNNTS KTCCKCGHLN NYFNFEYRKK NKFPHPKCEK CNFKENADYN 540
AALNISNPKL KSTKEEP 557

SEQ ID NO: 393      moltype = AA length = 557
FEATURE            Location/Qualifiers
source              1..557
                    mol_type = protein
                    note = Engineered CWCas12f1 (S188Q)
                    organism = synthetic construct

SEQUENCE: 393
MGEKSSRRRR NGKSGAWTAA ITSCVGGKMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60

```

-continued

NREKIALEKN	KDKVKEACSK	HLKVAAYCTT	QVERNACLF	KARKLDDKFY	QKLRGQFPDA	120
VFWQEISEIF	RQLQKQAAEI	YNQSLIELYY	EIFIKGKGIA	NASSVEHYLS	DVCYTRAAEL	180
FKNAIAAAGL	RSKIKSNFRL	KELKNMKSGL	PTTKSDNFPI	PLVKQKGGQY	TGFEISNHNS	240
DFIIKIPFGR	WQVKKEIDKY	RPWEKFDPEQ	VQKSPKPISL	LLSTQRRKRN	KGWSKDEGTE	300
AEIKKVMNGD	YQTSYIEVCR	GSKIGEKSAA	MLNLSIDVPK	IDKGVDPDSII	GGIDVGVKSP	360
LVCAINNAFS	RYSISDNDFL	HFNNKMFARR	RILLKKNRHK	RAGHGAKNKL	KPITILTEKS	420
ERFRKKLIER	WACEIADFFI	KNKVGTVQME	NLESMKRKED	SYFNIRLRGF	WPYAEMQNKI	480
EPFLKQYGIE	IRKVAPNNTS	KTCSKCGHLN	NYFNFEYRKK	NKPPHFKCEK	CNFKENADYN	540
AALNISNPKL	KSTKEEP					557

SEQ ID NO: 394 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = Engineered CWCas12f1 (S188H)
 organism = synthetic construct

SEQUENCE: 394

MGEKSSRRRR	NGKSGAWTAA	ITSCVGGKMA	KNTITKTLKL	RIVRPYNsAE	VEKIVADEKN	60
NREKIALEKN	KDKVKEACSK	HLKVAAYCTT	QVERNACLF	KARKLDDKFY	QKLRGQFPDA	120
VFWQEISEIF	RQLQKQAAEI	YNQSLIELYY	EIFIKGKGIA	NASSVEHYLS	DVCYTRAAEL	180
FKNAIAAHGL	RSKIKSNFRL	KELKNMKSGL	PTTKSDNFPI	PLVKQKGGQY	TGFEISNHNS	240
DFIIKIPFGR	WQVKKEIDKY	RPWEKFDPEQ	VQKSPKPISL	LLSTQRRKRN	KGWSKDEGTE	300
AEIKKVMNGD	YQTSYIEVCR	GSKIGEKSAA	MLNLSIDVPK	IDKGVDPDSII	GGIDVGVKSP	360
LVCAINNAFS	RYSISDNDFL	HFNNKMFARR	RILLKKNRHK	RAGHGAKNKL	KPITILTEKS	420
ERFRKKLIER	WACEIADFFI	KNKVGTVQME	NLESMKRKED	SYFNIRLRGF	WPYAEMQNKI	480
EPFLKQYGIE	IRKVAPNNTS	KTCSKCGHLN	NYFNFEYRKK	NKPPHFKCEK	CNFKENADYN	540
AALNISNPKL	KSTKEEP					557

SEQ ID NO: 395 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = Engineered CWCas12f1 (S188K)
 organism = synthetic construct

SEQUENCE: 395

MGEKSSRRRR	NGKSGAWTAA	ITSCVGGKMA	KNTITKTLKL	RIVRPYNsAE	VEKIVADEKN	60
NREKIALEKN	KDKVKEACSK	HLKVAAYCTT	QVERNACLF	KARKLDDKFY	QKLRGQFPDA	120
VFWQEISEIF	RQLQKQAAEI	YNQSLIELYY	EIFIKGKGIA	NASSVEHYLS	DVCYTRAAEL	180
FKNAIAAGL	RSKIKSNFRL	KELKNMKSGL	PTTKSDNFPI	PLVKQKGGQY	TGFEISNHNS	240
DFIIKIPFGR	WQVKKEIDKY	RPWEKFDPEQ	VQKSPKPISL	LLSTQRRKRN	KGWSKDEGTE	300
AEIKKVMNGD	YQTSYIEVCR	GSKIGEKSAA	MLNLSIDVPK	IDKGVDPDSII	GGIDVGVKSP	360
LVCAINNAFS	RYSISDNDFL	HFNNKMFARR	RILLKKNRHK	RAGHGAKNKL	KPITILTEKS	420
ERFRKKLIER	WACEIADFFI	KNKVGTVQME	NLESMKRKED	SYFNIRLRGF	WPYAEMQNKI	480
EPFLKQYGIE	IRKVAPNNTS	KTCSKCGHLN	NYFNFEYRKK	NKPPHFKCEK	CNFKENADYN	540
AALNISNPKL	KSTKEEP					557

SEQ ID NO: 396 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = Engineered CWCas12f1 (R191K)
 organism = synthetic construct

SEQUENCE: 396

MGEKSSRRRR	NGKSGAWTAA	ITSCVGGKMA	KNTITKTLKL	RIVRPYNsAE	VEKIVADEKN	60
NREKIALEKN	KDKVKEACSK	HLKVAAYCTT	QVERNACLF	KARKLDDKFY	QKLRGQFPDA	120
VFWQEISEIF	RQLQKQAAEI	YNQSLIELYY	EIFIKGKGIA	NASSVEHYLS	DVCYTRAAEL	180
FKNAIAASGL	RSKIKSNFRL	KELKNMKSGL	PTTKSDNFPI	PLVKQKGGQY	TGFEISNHNS	240
DFIIKIPFGR	WQVKKEIDKY	RPWEKFDPEQ	VQKSPKPISL	LLSTQRRKRN	KGWSKDEGTE	300
AEIKKVMNGD	YQTSYIEVCR	GSKIGEKSAA	MLNLSIDVPK	IDKGVDPDSII	GGIDVGVKSP	360
LVCAINNAFS	RYSISDNDFL	HFNNKMFARR	RILLKKNRHK	RAGHGAKNKL	KPITILTEKS	420
ERFRKKLIER	WACEIADFFI	KNKVGTVQME	NLESMKRKED	SYFNIRLRGF	WPYAEMQNKI	480
EPFLKQYGIE	IRKVAPNNTS	KTCSKCGHLN	NYFNFEYRKK	NKPPHFKCEK	CNFKENADYN	540
AALNISNPKL	KSTKEEP					557

SEQ ID NO: 397 moltype = AA length = 557
 FEATURE Location/Qualifiers
 source 1..557
 mol_type = protein
 note = Engineered CWCas12f1 (Q225T)
 organism = synthetic construct

SEQUENCE: 397

MGEKSSRRRR	NGKSGAWTAA	ITSCVGGKMA	KNTITKTLKL	RIVRPYNsAE	VEKIVADEKN	60
NREKIALEKN	KDKVKEACSK	HLKVAAYCTT	QVERNACLF	KARKLDDKFY	QKLRGQFPDA	120
VFWQEISEIF	RQLQKQAAEI	YNQSLIELYY	EIFIKGKGIA	NASSVEHYLS	DVCYTRAAEL	180
FKNAIAASGL	RSKIKSNFRL	KELKNMKSGL	PTTKSDNFPI	PLVKTKGGQY	TGFEISNHNS	240
DFIIKIPFGR	WQVKKEIDKY	RPWEKFDPEQ	VQKSPKPISL	LLSTQRRKRN	KGWSKDEGTE	300

-continued

```

AEIKKVMNGD YQTSYIEVKR GSKIGEKS AW MLNLSIDVPK IDKGVDPSII GGIDVGKSP 360
LVCAINNAFS RYSISDNDLF HFNKKMFARR RILLKKNRHK RAGHGAKNKL KPITILTEKS 420
ERFRKKLIER WACEIADFFI KNKVGTVQME NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
EPKLLQYGIE IRKVAPNNTS KTCCKCGHLN NYFNFEYRKK NKPFHPKCEK CNFKENADYN 540
AALNISNPPL KSTKEEP 557

```

```

SEQ ID NO: 398      moltype = AA length = 557
FEATURE            Location/Qualifiers
source             1..557
                  mol_type = protein
                  note = Engineered CWCas12f1 (Q225F)
                  organism = synthetic construct

```

```

SEQUENCE: 398
MGEKSSRRRR NGKSGAWTAA ITSCVGGKMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60
NREKIALEKN KDKVKEACSK HLKVAAYCTT QVERNACLFC KARKLDDKFY QKLRGQFPDA 120
VFWQEISEIF RQLQQA AEI YNQLIELYY EIFIKGKGIA NASSVEHYLS DVCYTRAAEL 180
FKNAAIASGL RSKIKSNFRL KELKNMKSGL PTTKSDNFPI PLVKFKGGQY TGFEISNHNS 240
DFIIKIPFGR WQVKEIDKY RPWEKDFDEQ VQKSPKISL LLSTQRRKRN KGWSKDEGTE 300
AEIKKVMNGD YQTSYIEVKR GSKIGEKS AW MLNLSIDVPK IDKGVDPSII GGIDVGKSP 360
LVCAINNAFS RYSISDNDLF HFNKKMFARR RILLKKNRHK RAGHGAKNKL KPITILTEKS 420
ERFRKKLIER WACEIADFFI KNKVGTVQME NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
EPKLLQYGIE IRKVAPNNTS KTCCKCGHLN NYFNFEYRKK NKPFHPKCEK CNFKENADYN 540
AALNISNPPL KSTKEEP 557

```

```

SEQ ID NO: 399      moltype = AA length = 557
FEATURE            Location/Qualifiers
source             1..557
                  mol_type = protein
                  note = Engineered CWCas12f1 (Q272K)
                  organism = synthetic construct

```

```

SEQUENCE: 399
MGEKSSRRRR NGKSGAWTAA ITSCVGGKMA KNTITKTLKL RIVRPYNSAE VEKIVADEKN 60
NREKIALEKN KDKVKEACSK HLKVAAYCTT QVERNACLFC KARKLDDKFY QKLRGQFPDA 120
VFWQEISEIF RQLQQA AEI YNQLIELYY EIFIKGKGIA NASSVEHYLS DVCYTRAAEL 180
FKNAAIASGL RSKIKSNFRL KELKNMKSGL PTTKSDNFPI PLVKQKGGQY TGFEISNHNS 240
DFIIKIPFGR WQVKEIDKY RPWEKDFDEQ VKKSPKISL LLSTQRRKRN KGWSKDEGTE 300
AEIKKVMNGD YQTSYIEVKR GSKIGEKS AW MLNLSIDVPK IDKGVDPSII GGIDVGKSP 360
LVCAINNAFS RYSISDNDLF HFNKKMFARR RILLKKNRHK RAGHGAKNKL KPITILTEKS 420
ERFRKKLIER WACEIADFFI KNKVGTVQME NLESMKRKED SYFNIRLRGF WPYAEMQNKI 480
EPKLLQYGIE IRKVAPNNTS KTCCKCGHLN NYFNFEYRKK NKPFHPKCEK CNFKENADYN 540
AALNISNPPL KSTKEEP 557

```

```

SEQ ID NO: 400      moltype = RNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                  mol_type = other RNA
                  note = shRNA for human XRCC4-1
                  organism = synthetic construct

```

```

SEQUENCE: 400
gtgatgacac tggcacatta 20

```

```

SEQ ID NO: 401      moltype = RNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                  mol_type = other RNA
                  note = shRNA for human XRCC4-2
                  organism = synthetic construct

```

```

SEQUENCE: 401
gtgagagtac tgatgaggaa 20

```

```

SEQ ID NO: 402      moltype = RNA length = 19
FEATURE            Location/Qualifiers
source             1..19
                  mol_type = other RNA
                  note = shRNA for human XRCC4-3
                  organism = synthetic construct

```

```

SEQUENCE: 402
gaatccacct tgtttctga 19

```

```

SEQ ID NO: 403      moltype = RNA length = 20
FEATURE            Location/Qualifiers
source             1..20
                  mol_type = other RNA
                  note = shRNA for human XRCC4-4
                  organism = synthetic construct

```

```

SEQUENCE: 403

```

-continued

gctacaagta tcttgggaga	20
SEQ ID NO: 404	moltype = RNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other RNA
	note = shRNA for human XRCC4-5
	organism = synthetic construct
SEQUENCE: 404	
gaaatgcagc tcaagaacga	20
SEQ ID NO: 405	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = shRNA for human XLF-1
	organism = synthetic construct
SEQUENCE: 405	
gcatgagtct ggcattaca	19
SEQ ID NO: 406	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = shRNA for human XLF-2
	organism = synthetic construct
SEQUENCE: 406	
gaaagccctt tgtcatgaa	19
SEQ ID NO: 407	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = shRNA for human XLF-3
	organism = synthetic construct
SEQUENCE: 407	
aaacagtgct tccctgcaa	19
SEQ ID NO: 408	moltype = RNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other RNA
	note = shRNA for human XLF-4
	organism = synthetic construct
SEQUENCE: 408	
gtgaaagacc tagagatcca	20
SEQ ID NO: 409	moltype = RNA length = 20
FEATURE	Location/Qualifiers
source	1..20
	mol_type = other RNA
	note = shRNA for human XLF-5
	organism = synthetic construct
SEQUENCE: 409	
gatatggcag tcaccacaca	20
SEQ ID NO: 410	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = shRNA for human XRCC6-1
	organism = synthetic construct
SEQUENCE: 410	
gcagcattgt gcagataca	19
SEQ ID NO: 411	moltype = RNA length = 19
FEATURE	Location/Qualifiers
source	1..19
	mol_type = other RNA
	note = shRNA for human XRCC6-2
	organism = synthetic construct
SEQUENCE: 411	
gcaggaacat ccctcctta	19
SEQ ID NO: 412	moltype = RNA length = 20
FEATURE	Location/Qualifiers

-continued

source	1..20 mol_type = other RNA note = shRNA for human XRCC6-3 organism = synthetic construct	
SEQUENCE: 412 gtcagtgctc tgctcatcaa		20
SEQ ID NO: 413 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = shRNA for human XRCC6-4 organism = synthetic construct	
SEQUENCE: 413 ggatcatgct gttcaccaa		19
SEQ ID NO: 414 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = shRNA for human XRCC6-5 organism = synthetic construct	
SEQUENCE: 414 ggatctgact actcactca		19
SEQ ID NO: 415 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = shRNA for human LIG4-1 organism = synthetic construct	
SEQUENCE: 415 gcacaaagat ggagatgta		19
SEQ ID NO: 416 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = shRNA for human LIG4-2 organism = synthetic construct	
SEQUENCE: 416 gccagacacg tactgtgtaa		20
SEQ ID NO: 417 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = shRNA for human LIG4-3 organism = synthetic construct	
SEQUENCE: 417 gcgagcagac tcctgaagaa		20
SEQ ID NO: 418 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = shRNA for human LIG4-4 organism = synthetic construct	
SEQUENCE: 418 agaggattct gatctgcaa		19
SEQ ID NO: 419 FEATURE source	moltype = RNA length = 19 Location/Qualifiers 1..19 mol_type = other RNA note = shRNA for human LIG4-5 organism = synthetic construct	
SEQUENCE: 419 gcatgacct tctgtagga		19
SEQ ID NO: 420 FEATURE source	moltype = RNA length = 20 Location/Qualifiers 1..20 mol_type = other RNA note = shRNA for human DCLRE1C-1 organism = synthetic construct	

-continued

SEQUENCE: 420		
gcgagactcc taccagata		20
SEQ ID NO: 421	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human DCLRE1C-2	
	organism = synthetic construct	
SEQUENCE: 421		
ggacaaagct gactacaga		19
SEQ ID NO: 422	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human DCLRE1C-3	
	organism = synthetic construct	
SEQUENCE: 422		
gcagagctct cgtttcaca		19
SEQ ID NO: 423	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = shRNA for human DCLRE1C-4	
	organism = synthetic construct	
SEQUENCE: 423		
gtgactctga tggagaatca		20
SEQ ID NO: 424	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = shRNA for human DCLRE1C-5	
	organism = synthetic construct	
SEQUENCE: 424		
gccagaattc ttcccagtca		20
SEQ ID NO: 425	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-1	
	organism = synthetic construct	
SEQUENCE: 425		
ggagccagat agtttgat		19
SEQ ID NO: 426	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-2	
	organism = synthetic construct	
SEQUENCE: 426		
gcaagcagct gaaacaaat		19
SEQ ID NO: 427	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-3	
	organism = synthetic construct	
SEQUENCE: 427		
ggagctgatt gtagcaaca		19
SEQ ID NO: 428	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-4	
	organism = synthetic construct	
SEQUENCE: 428		
gcacagaagt gcctccaat		19
SEQ ID NO: 429	moltype = RNA length = 19	

-continued

FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-5	
	organism = synthetic construct	
SEQUENCE: 429		
ggacatagtt tctgggaga		19
SEQ ID NO: 430	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for human ATM1-6	
	organism = synthetic construct	
SEQUENCE: 430		
gaacttcagt ggaccttca		19
SEQ ID NO: 431	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = scrambled-1 shRNA	
	organism = synthetic construct	
SEQUENCE: 431		
gttctccgaa cgtgtcacgt		20
SEQ ID NO: 432	moltype = RNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other RNA	
	note = scrambled-2 shRNA	
	organism = synthetic construct	
SEQUENCE: 432		
gcaacaagat gaagagcacc aa		22
SEQ ID NO: 433	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-1	
	organism = synthetic construct	
SEQUENCE: 433		
gaagaccagt gctaatacta		19
SEQ ID NO: 434	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-2	
	organism = synthetic construct	
SEQUENCE: 434		
ggaccttgat gtgctaaca		19
SEQ ID NO: 435	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-3	
	organism = synthetic construct	
SEQUENCE: 435		
gcaccgcaag ccttctata		19
SEQ ID NO: 436	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-4	
	organism = synthetic construct	
SEQUENCE: 436		
gaaactggca acaggtcaa		19
SEQ ID NO: 437	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-5	

-continued

SEQUENCE: 437	organism = synthetic construct	
ggatcatgag ggagtaaga		19
SEQ ID NO: 438	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-6	
	organism = synthetic construct	
SEQUENCE: 438		
gaagtgttag actattaca		19
SEQ ID NO: 439	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-7	
	organism = synthetic construct	
SEQUENCE: 439		
ggacagtgca cctgactaa		19
SEQ ID NO: 440	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-8	
	organism = synthetic construct	
SEQUENCE: 440		
gaatgtatgg gtactgtca		19
SEQ ID NO: 441	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-9	
	organism = synthetic construct	
SEQUENCE: 441		
gaaggtgtaa gggctctgaa		19
SEQ ID NO: 442	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-10	
	organism = synthetic construct	
SEQUENCE: 442		
gaatggagct tctgcactct		20
SEQ ID NO: 443	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-11	
	organism = synthetic construct	
SEQUENCE: 443		
gcttatggct acgagtattt		20
SEQ ID NO: 444	moltype = RNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-12	
	organism = synthetic construct	
SEQUENCE: 444		
gagctgggag ttcaggttca		20
SEQ ID NO: 445	moltype = RNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other RNA	
	note = shRNA for mouse DCLRE1C-13	
	organism = synthetic construct	
SEQUENCE: 445		
gcatcaagcc atctaccat		19

-continued

SEQ ID NO: 446	moltype = RNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other RNA	
	note = scrambled-3 shRNA	
	organism = synthetic construct	
SEQUENCE: 446		
cagagctaac tcagatagta ct		22
SEQ ID NO: 447	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	note = primer_E51_LD_F for qPCR	
	organism = synthetic construct	
SEQUENCE: 447		
tgtcatgaat aagagtttgg ctca		24
SEQ ID NO: 448	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	note = primer_E51_LD_R for qPCR	
	organism = synthetic construct	
SEQUENCE: 448		
aggttggtgc accagagtaa ca		22
SEQ ID NO: 449	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	note = primer_18s_rRNA-F for qPCR	
	organism = synthetic construct	
SEQUENCE: 449		
tcaactttcg atggtagtcg cc		22
SEQ ID NO: 450	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	note = primer_18s_rRNA-R for qPCR	
	organism = synthetic construct	
SEQUENCE: 450		
ggcctcgaaa gagtctgta ttgt		24
SEQ ID NO: 451	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hATM-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 451		
ggacagtgga ggcacaaaat		20
SEQ ID NO: 452	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hATM-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 452		
gtgtcgaaga cagctggtga		20
SEQ ID NO: 453	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hXRCC6-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 453		
cgggaaacaa atgaaccagt		20
SEQ ID NO: 454	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	

-continued

SEQUENCE: 454	note = primer_hXRCC6-R for qRT-PCR	
tgaaacccat gagcatcaaa	organism = synthetic construct	20
SEQ ID NO: 455	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hLIG4-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 455		
caccttgctggt tttccacgaa		20
SEQ ID NO: 456	mol_type = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	note = primer_hLIG4-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 456		
cagatgcctt ccccctaagt tg		22
SEQ ID NO: 457	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hXRCC4-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 457		
ctgatgggtca ttcagcatgg		20
SEQ ID NO: 458	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hXRCC4-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 458		
tcctgctcct gacaacaatg		20
SEQ ID NO: 459	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hLIG4-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 459		
tctctggcct ccccttctat		20
SEQ ID NO: 460	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hLIG4-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 460		
tagctccctc acttggcact		20
SEQ ID NO: 461	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hDCLRE1C-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 461		
aattccaagt cgggaggagt		20
SEQ ID NO: 462	mol_type = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_hDCLRE1C-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 462		
ggatctgagt gttgcggtct		20

-continued

SEQ ID NO: 463	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	note = primer_GAPDH-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 463		
ggaaggactc atgaccacag t		21
SEQ ID NO: 464	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	note = primer_GAPDH-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 464		
cagtgagctt cccgttcag		19
SEQ ID NO: 465	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	note = primer_18s_rRNA-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 465		
tcaactttcg atggtagtcg cc		22
SEQ ID NO: 466	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	note = primer_18s_rRNA-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 466		
ggcctcgaaa gagtctcgta ttgt		24
SEQ ID NO: 467	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	note = primer_mDclrelc-F for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 467		
gcttcgggtg agaaggaag		19
SEQ ID NO: 468	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	note = primer_mDclrelc-R for qRT-PCR	
	organism = synthetic construct	
SEQUENCE: 468		
tgccagtcctg aagtcctcctg		20
SEQ ID NO: 469	moltype = RNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other RNA	
	note = scrambled-1 shRNA for mouse	
	organism = synthetic construct	
SEQUENCE: 469		
cctaaggtta agtcgcctc g		21
SEQ ID NO: 470	moltype = RNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other RNA	
	note = scrambled-3 shRNA	
	organism = synthetic construct	
SEQUENCE: 470		
cagagctaac tcagatagta ct		22
SEQ ID NO: 471	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	

-continued

```

mol_type = other DNA
note = F68-protospacer for SaCas9
organism = synthetic construct

SEQUENCE: 471
gtgtattgct tgtactactc a                                21

SEQ ID NO: 472      moltype = DNA length = 21
FEATURE            Location/Qualifiers
source              1..21
                    mol_type = other DNA
                    note = R84-protospacer for SaCas9
                    organism = synthetic construct

SEQUENCE: 472
gtgtattac ttgctactgc a                                21

SEQ ID NO: 473      moltype = RNA length = 47
FEATURE            Location/Qualifiers
source              1..47
                    mol_type = other RNA
                    note = shRNA for ATM1-6
                    organism = synthetic construct

SEQUENCE: 473
gaacttcagt ggaccttcac tcaagagatg aaggtccact gaagttc    47

```

1. A composition for increasing deletion efficiency of a nucleic acid segment, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule.

2. The composition of claim 1, wherein the gene involved in non-homologous end joining pathway comprises at least one selected from the group consisting of ATM1, XRCC4, XLF, XRCC6, LIG4, and DCLRE1C.

3. The composition of claim 2, wherein the gene involved in non-homologous end joining pathway comprises at least one selected from the group consisting of XRCC6 and DCLRE1C.

4. (canceled)

5. The composition of claim 1, wherein the molecule is a shRNA molecule that inhibits expression of at least one selected from the group consisting of shXRCC6 and shDCLRE1C.

6. The composition of claim 1, wherein the molecule is a shRNA molecule that comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS: 360 to 389, SEQ ID NOS: 400 to 430, SEQ ID NOS: 433 to 445, and SEQ ID NO: 473.

7. The composition of claim 5, wherein the shRNA molecule comprises a nucleotide sequence selected from the group consisting of SEQ ID NOS: 375 to 379, SEQ ID NOS: 385 to 389, SEQ ID NOS: 410 to 414, SEQ ID NOS: 420 to 424, and SEQ ID NOS: 433 to 445.

8. (canceled)

9. (canceled)

10. The composition of claim 1, wherein the composition further comprises (i) a Cas endonuclease and (ii) a first guide RNA and a second guide RNA that hybridize to a first target sequence and a second target sequence, respectively,

wherein the nucleic acid segment exists between the first target sequence and the second target sequence; and

wherein the Cas endonuclease forms a first complex with the first guide RNA to induce first cleavage, and the Cas endonuclease forms a second complex with the second guide RNA to induce second cleavage.

11. The composition of claim 1, wherein the composition further comprises at least one nucleic acid construct comprising (i) a nucleic acid encoding a Cas endonuclease, (ii) a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence, and (iii) a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence;

wherein the nucleic acid segment exists between the first target sequence and the second target sequence; and

wherein the Cas endonuclease forms a first complex with the first guide RNA to induce first cleavage, and the Cas endonuclease forms a second complex with the second guide RNA to induce second cleavage.

12. (canceled)

13. (canceled)

14. The composition of claim 10, wherein the Cas endonuclease is an endonuclease comprising a Cas12f1 protein.

15. (canceled)

16. (canceled)

17. The composition of claim 10, wherein the first or second guide RNA comprises a U-rich tail sequence linked to the 3'-end of its guide sequence, in which the U-rich tail is represented by 5'-(U_mV)_nU_o-3', wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5.

18. The composition of claim 10, wherein the first or second guide RNA is an engineered guide RNA comprising a nucleotide sequence having at least 50% sequence identity to a scaffold region of a wild-type Cas12f1 guide RNA sequence that sequentially comprises, from the 5'-end, a first stem-loop region, a second stem-loop region, a third stem-loop region, a fourth stem-loop region, and a tracrRNA-crRNA complementarity region, and


the engineered guide RNA comprises at least one modification selected from the group consisting of the following (1) to (5) with respect to the wild-type Cas12f1 guide RNA sequence:

- (1) deletion of at least a part of the first stem-loop region;
- (2) deletion of at least a part of the second stem-loop region;

- (3) deletion of at least a part of the tracrRNA-crRNA complementarity region;
- (4) replacement of one or more uracil (U) residues with A, G, or C in three or more consecutive U residues when the consecutive U residues are present in the tracrRNA-crRNA complementarity region; and
- (5) addition of a U-rich tail to the 3'-end of its crRNA sequence (a sequence of the U-rich tail is represented by 5'-(U_mV)_nU_o-3', wherein V is each independently A, C, or G, m and o are integers between 1 to 20, and n is an integer between 0 to 5).

19. (canceled)

20. The composition of claim 18, wherein the engineered guide RNA consists of a sequence represented by Formula (I) or has at least 80% sequence identity thereto:



in Formula (I),

X^a, X^{b1}, X^{b2}, X^{c1}, and X^{c2} each independently consist of 0 to 35 (poly)nucleotides,

X^g is a first or second guide sequence,

Lk is a polynucleotide linker of 2 to 20 nucleotides or is absent, and

(U_mV)_nU_o is present as a U-rich tail or absent, and when (U_mV)_nU_o is present, U is uridine, V is each independently A, C, or G, m and o are each independently an integer between 1 and 20, and n is an integer between 0 and 5.

21. The composition of claim 20, wherein X^a comprises the nucleotide sequence of SEQ ID NO: 14 or a nucleotide sequence having the sequence of SEQ ID NO: 14 from which 1 to 20 nucleotides are deleted.

22. The composition of claim 20, wherein X^{b1} comprises the nucleotide sequence of SEQ ID NO: 25 or a nucleotide sequence having the sequence of SEQ ID NO: 25 from which 1 to 13 nucleotides are deleted.

23. The composition of claim 20, wherein X^{b2} comprises the nucleotide sequence of SEQ ID NO: 29 or a nucleotide sequence having the sequence of SEQ ID NO: 29 from which 1 to 14 nucleotides are deleted.

24. The composition of claim 20, wherein the sequence 5'-X^{b1}UUAGX^{b2}-3' in Formula (I) is a nucleotide sequence selected from the group consisting of SEQ ID NOS: 34 to 38.

25. The composition of claim 20, wherein X^{c1} comprises the nucleotide sequence of SEQ ID NO: 39 or a nucleotide sequence having the sequence of SEQ ID NO: 39 from which 1 to 28 nucleotides are deleted.

26. The composition of claim 25, wherein in a case where three or more consecutive uracil (U) residues are present in a sequence of X^{c1}, the sequence of X^{c1} comprises a modification in which at least one U residue thereof is replaced with A, G, or C.

27. The composition of claim 20, wherein X^{c2} comprises the nucleotide sequence of SEQ ID NO: 58 or a nucleotide sequence having the sequence of SEQ ID NO: 58 from which 1 to 27 nucleotides are deleted.

28. The composition of claim 27, wherein in a case where the sequence 5'-ACGAA-3' is present in X^{c2}, the sequence is replaced with 5'-NGNNN-3', and N is each independently A, C, G, or U.

29. The composition of claim 20, wherein the sequence 5'-X^{c1}-Lk-X^{c2}-3' in Formula (I) is a nucleotide sequence selected from the group consisting of SEQ ID NOS: 80 to 86.

30. The composition of claim 20, wherein Lk comprises a nucleotide sequence selected from the group consisting of 5'-GAAA-3', 5'-UUAG-3', 5'-UGAAAA-3', 5'-UUGAAAAA-3', 5'-UUCGAAAGAA-3' (SEQ ID NO: 76), 5'-UUCAGAAAUGAA-3' (SEQ ID NO: 77), 5'-UU-CAUGAAAAUGAA-3' (SEQ ID NO: 78), and 5'-UU-CAUUGAAAAAUGAA-3' (SEQ ID NO: 79).

31. The composition of claim 10, wherein the first or second guide RNA comprises an engineered tracrRNA having a nucleotide sequence selected from the group consisting of SEQ ID NOS: 87 to 132.

32. The composition of claim 10, wherein the first or second guide RNA comprises an engineered crRNA sequence having a nucleotide sequence selected from the group consisting of SEQ ID NOS: 133 to 148.

33. (canceled)

34. The composition of claim 10, wherein the first or second guide RNA is a single guide RNA comprising a nucleotide sequence selected from the group consisting of SEQ ID NOS: 149 to 186.

35. (canceled)

36. (canceled)

37. A method for increasing deletion of a nucleic acid segment in a target gene of a cell, comprising bringing into contact with the cell the composition of claim 1.

38. (canceled)

39. (canceled)

40. A kit or system for gene editing, comprising:

a composition for increasing deletion of a nucleic acid segment, comprising a molecule that inhibits expression of a gene involved in non-homologous end joining (NHEJ) pathway, or a nucleic acid construct encoding the molecule; and

a gene editing composition for deletion of a nucleic acid segment, comprising (i) a Cas endonuclease and a first guide RNA and a second guide RNA that hybridize to a first target sequence and a second target sequence, respectively, or (ii) at least one nucleic acid construct comprising a nucleic acid encoding a Cas endonuclease, a nucleic acid encoding a first guide RNA that hybridizes to a first target sequence, and a nucleic acid encoding a second guide RNA that hybridizes to a second target sequence,

wherein the nucleic acid segment exists between the first target sequence and the second target sequence, and the Cas endonuclease forms a first complex with the first guide RNA to induce first cleavage, and the Cas endonuclease forms a second complex with the second guide RNA to induce second cleavage.

* * * * *