



US012383617B2

(12) **United States Patent**
Hubbell et al.

(10) **Patent No.: US 12,383,617 B2**

(45) **Date of Patent: Aug. 12, 2025**

(54) **COMPOSITIONS AND METHODS
CONCERNING IMMUNE TOLERANCE**

(71) Applicants: **The University of Chicago**, Chicago,
IL (US); **Anokion SA**, Ecublens (CH)

(72) Inventors: **Jeffrey A. Hubbell**, Chicago, IL (US);
David Scott Wilson, Chicago, IL (US);
Kym Brünggel, Chicago, IL (US);
Kristen Marie Lorentz, Ecublens (CH)

(73) Assignees: **The University of Chicago**, Chicago,
IL (US); **Anokion SA**, Ecublens (CH)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 951 days.

(21) Appl. No.: **17/250,015**

(22) PCT Filed: **May 9, 2019**

(86) PCT No.: **PCT/US2019/031440**

§ 371 (c)(1),

(2) Date: **Nov. 3, 2020**

(87) PCT Pub. No.: **WO2019/217628**

PCT Pub. Date: **Nov. 14, 2019**

(65) **Prior Publication Data**

US 2021/0244812 A1 Aug. 12, 2021

Related U.S. Application Data

(60) Provisional application No. 62/669,044, filed on May
9, 2018.

(51) **Int. Cl.**

A61K 39/385 (2006.01)

A61K 39/00 (2006.01)

A61K 47/54 (2017.01)

A61K 47/60 (2017.01)

A61P 37/06 (2006.01)

C07D 249/16 (2006.01)

C08F 220/58 (2006.01)

C08G 65/332 (2006.01)

C08G 65/334 (2006.01)

C08G 81/02 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/385** (2013.01); **A61K 39/0008**
(2013.01); **A61K 39/001154** (2018.08); **A61K**
39/001158 (2018.08); **A61K 39/4611**
(2023.05); **A61K 39/4644** (2023.05); **A61K**
47/549 (2017.08); **A61K 47/60** (2017.08);
A61P 37/06 (2018.01); **C07D 249/16**
(2013.01); **C08F 220/58** (2013.01); **C08G**
65/332 (2013.01); **C08G 65/334** (2013.01);
C08G 81/025 (2013.01); **A61K 2039/6087**
(2013.01); **A61K 2039/6093** (2013.01); **A61K**
2239/38 (2023.05)

(58) **Field of Classification Search**

CPC **A61K 39/385**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,671,958 A	6/1987	Rodwell et al.
4,741,900 A	5/1988	Alvarez et al.
4,859,449 A	8/1989	Mattes
4,867,973 A	9/1989	Goers et al.
4,950,738 A	8/1990	King et al.
5,086,002 A	2/1992	Hillyard et al.
5,140,104 A	8/1992	Coughlin et al.
5,156,840 A	10/1992	Goers et al.
5,162,512 A	11/1992	King et al.
5,227,165 A	7/1993	Domb et al.
5,227,293 A	7/1993	Stengelin et al.
5,346,696 A	9/1994	Kim et al.
5,358,857 A	10/1994	Stengelin et al.
5,470,570 A	11/1995	Taylor et al.
5,487,890 A	1/1996	Taylor et al.
5,681,571 A	10/1997	Holmgren et al.
5,698,679 A	12/1997	Nemazee et al.
5,718,915 A	2/1998	Virtanen et al.
5,879,679 A	3/1999	Taylor et al.
5,885,808 A	3/1999	Spooner et al.
5,886,143 A	3/1999	Theodore et al.
5,948,639 A	9/1999	Gimeno et al.
5,985,826 A	11/1999	Theodore et al.
5,994,104 A	11/1999	Anderson et al.
5,997,861 A	12/1999	Virtanen et al.
6,022,564 A	2/2000	Takechi et al.
6,090,925 A	7/2000	Woiszwilllo et al.
6,120,770 A	9/2000	Adams et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN	1289256	3/2001
CN	1756560	4/2006

(Continued)

OTHER PUBLICATIONS

“EPFL School of Life Sciences—Annual Report SV 2011,” 156
Pages (Dec. 31, 2011).

“Integer”, Meriam-Webster, available online at <https://www.merriam-webster.com/dictionary/integer>, 12 pages (2019) (Year: 2019).

“SubName: Full=Phosphate ABC Transporter, Inner Membrane
Subunit PstC;”, XP002717162, Retrieved From EBI Accession No.
UNIPROT:C7QKI6, Database Accession No. C7QKI6 (Oct. 13,
2009).

“SubName: Full=Putative Integron Gene Cassette Protein; Flags:
Fragment;”, XP002717159, Retrieved From EBI Accession No.
UNIPROT:BOBITO, Database Accession No. BOBITO (Feb. 26,
2008).

(Continued)

Primary Examiner — Paul V Ward

(74) *Attorney, Agent, or Firm* — Norton Rose Fulbright
US LLP

(57) **ABSTRACT**

The present disclosure provides compositions comprising
mannose-fused antigens to target mannose receptors. The
compositions may be used to prevent immunity or reduce an
immune response protein-based drugs that would otherwise
elicit an immune response.

17 Claims, 7 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

6,153,203	A	11/2000	Holmgren et al.	8,961,991	B2	2/2015	Zurawski et al.
6,217,869	B1	4/2001	Meyer et al.	8,992,917	B2	3/2015	Goldenberg et al.
6,224,794	B1	5/2001	Amsden et al.	9,005,903	B2	4/2015	Rubin-Bejerano et al.
6,264,950	B1	7/2001	Staerz	9,066,984	B2	6/2015	Mi et al.
6,322,796	B1	11/2001	Holmgren et al.	9,102,730	B2	8/2015	Zurawski et al.
6,365,163	B1	4/2002	Holmgren et al.	9,102,734	B2	8/2015	Zurawski et al.
6,379,699	B1	4/2002	Virtanen et al.	9,187,561	B2	11/2015	Goldenberg et al.
6,488,927	B2	12/2002	Muzykantov et al.	9,216,156	B2	12/2015	Fleury et al.
6,512,103	B1	1/2003	Dairaghi et al.	9,233,072	B2	1/2016	Alexis et al.
6,562,347	B1	5/2003	Kwak et al.	9,234,040	B2	1/2016	Zurawski et al.
6,703,488	B1	3/2004	Burton et al.	9,260,692	B2	2/2016	Martin et al.
6,737,057	B1	5/2004	Zaghoulani et al.	9,308,280	B2	4/2016	Shi et al.
6,814,964	B2	11/2004	Virtanen et al.	9,326,939	B2	5/2016	Paulson et al.
6,905,688	B2	6/2005	Rosen et al.	9,416,186	B2	8/2016	Zurawski et al.
6,953,675	B2	10/2005	Leung et al.	9,439,859	B2	9/2016	Alexis et al.
7,041,287	B2	5/2006	Muzykantov et al.	9,453,074	B2	9/2016	Oh et al.
7,132,475	B2	11/2006	Hubbel et al.	9,457,047	B2	10/2016	Rubin-Bejerano et al.
7,144,569	B1	12/2006	Anderson et al.	9,474,717	B2	10/2016	von Andrian et al.
7,148,329	B1	12/2006	Figdor et al.	9,517,257	B2	12/2016	Hubbell et al.
7,172,760	B2	2/2007	Muzykantov et al.	9,518,087	B2	12/2016	Hubbell et al.
7,175,988	B2	2/2007	Roschke et al.	9,522,183	B2	12/2016	Paulson et al.
7,192,582	B2	3/2007	Hudson et al.	9,539,210	B2	1/2017	von Andrian et al.
7,285,642	B2	10/2007	Figdor et al.	9,561,272	B2	2/2017	Thomas et al.
7,420,040	B2	9/2008	Young et al.	9,688,991	B2	6/2017	Levy et al.
7,420,041	B2	9/2008	Young et al.	9,751,945	B2	9/2017	Ploegh et al.
7,541,180	B2	6/2009	Valiante et al.	9,814,780	B2	11/2017	Hubbell et al.
7,585,508	B1	9/2009	Prendergast	9,850,296	B2	12/2017	Hubbell et al.
7,612,180	B2	11/2009	Goldenberg et al.	9,878,048	B2	1/2018	Hubbell et al.
7,704,943	B2	4/2010	Griffin et al.	9,901,645	B2	2/2018	Hubbell et al.
7,704,964	B2	4/2010	Delcayre et al.	9,901,646	B2	2/2018	Hubbell et al.
7,786,267	B2	8/2010	Zurawski et al.	10,046,056	B2	8/2018	Hubbell et al.
7,811,809	B2	10/2010	Heyduk et al.	10,800,838	B2	10/2020	Hubbell et al.
7,837,997	B2	11/2010	Muzykantov et al.	10,821,157	B2	11/2020	Hubbell et al.
7,884,109	B2	2/2011	Cohen et al.	10,919,963	B2	2/2021	Hubbell et al.
7,888,460	B2	2/2011	Anderson et al.	10,940,209	B2	3/2021	Hubbell et al.
7,892,743	B2	2/2011	Owen et al.	10,946,079	B2	3/2021	Hubbell et al.
7,932,294	B2	4/2011	Satyam	10,953,101	B2	3/2021	Hubbell et al.
7,994,283	B2	8/2011	Valiante et al.	2002/0004037	A1	1/2002	Koteliansky et al.
8,007,805	B2	8/2011	George et al.	2002/0038002	A1	3/2002	Zaghoulani
8,021,689	B2	9/2011	Reddy et al.	2002/0081298	A1	6/2002	Zaghoulani
8,057,798	B2	11/2011	Zurawski et al.	2002/0103343	A1	8/2002	Taylor et al.
8,058,400	B2	11/2011	Figdor et al.	2002/0187131	A1	12/2002	Hawiger et al.
8,058,406	B2	11/2011	Mi et al.	2002/0193572	A1	12/2002	Leung et al.
8,105,599	B2	1/2012	Figdor et al.	2003/0022826	A1	1/2003	Haynes
8,236,934	B2	8/2012	Banchereau et al.	2003/0082643	A1	5/2003	Hudson et al.
8,252,902	B2	8/2012	Barbas et al.	2003/0103967	A1	5/2003	Zaghoulani
8,273,357	B2	9/2012	Hacohen et al.	2003/0104045	A1	6/2003	Virtanen et al.
8,277,812	B2	10/2012	Iannacone et al.	2003/0175921	A1	9/2003	Barbas et al.
8,318,912	B2	11/2012	Simon	2003/0190676	A1	10/2003	Barbas et al.
8,323,696	B2	12/2012	Hubbel et al.	2003/0211078	A1	11/2003	Heavner
8,329,144	B2	12/2012	Anderson et al.	2004/0052815	A1	3/2004	Lycke
8,333,973	B2	12/2012	Muzykantov et al.	2004/0077843	A1	4/2004	Burton et al.
8,343,497	B2	1/2013	Shi et al.	2004/0146948	A1	7/2004	Britton et al.
8,343,498	B2	1/2013	Alexis et al.	2004/0147721	A1	7/2004	Valiante
8,425,910	B2	4/2013	Mi et al.	2004/0185057	A1	9/2004	Kirkby et al.
8,449,888	B2	5/2013	Zurawski et al.	2004/0197314	A1	10/2004	Delcayre et al.
8,507,237	B2	8/2013	Hermet et al.	2004/0258688	A1	12/2004	Hawiger et al.
8,518,410	B2	8/2013	Zurawski et al.	2005/0031628	A1	2/2005	George et al.
8,551,476	B2	10/2013	Mi et al.	2005/0053579	A1	3/2005	Galipeau et al.
8,562,998	B2	10/2013	Shi et al.	2005/0113297	A1	5/2005	Francois et al.
8,580,253	B2	11/2013	Rubin-Bejerano et al.	2005/0118168	A1	6/2005	Figdor et al.
8,586,052	B2	11/2013	Zurawski et al.	2005/0201973	A1	9/2005	Virtanen et al.
8,591,905	B2	11/2013	Von Andrian et al.	2005/0203022	A1	9/2005	Gotwals et al.
8,592,364	B2	11/2013	Swartz et al.	2005/0220804	A1	10/2005	Figdor et al.
8,613,903	B2	12/2013	Goldenberg et al.	2005/0250936	A1	11/2005	Oppermann et al.
8,617,823	B2	12/2013	Rubin-Bejerano et al.	2006/0034864	A1	2/2006	Zaghoulani
8,637,028	B2	1/2014	Alexis et al.	2006/0127929	A1	6/2006	Swager et al.
8,673,293	B2	3/2014	Martin et al.	2006/0153881	A1	7/2006	Narum et al.
8,685,408	B2	4/2014	Tartour et al.	2006/0173168	A1	8/2006	Carlock et al.
8,722,047	B2	5/2014	Goldenberg et al.	2006/0178299	A1	8/2006	Anderson et al.
8,728,481	B2	5/2014	Banchereau et al.	2006/0257412	A1	11/2006	Bowdish et al.
8,859,629	B2	10/2014	Van Delft et al.	2006/0280679	A1	12/2006	Bowdish et al.
8,889,140	B2	11/2014	Lee et al.	2007/0059794	A1	3/2007	Ideno et al.
8,906,381	B2	12/2014	Lannacone et al.	2007/0111222	A1	5/2007	Chasin et al.
8,932,595	B2	1/2015	Lannacone et al.	2007/0122409	A1	5/2007	Zaghoulani
				2007/0190615	A1	8/2007	Cohen et al.
				2007/0218053	A1	9/2007	Zaghoulani
				2008/0031899	A1	2/2008	Reddy et al.
				2008/0131428	A1	6/2008	Young et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2008/0160041 A1	7/2008	Figdor et al.	2012/0121592 A1	5/2012	Oh et al.
2008/0175971 A1	7/2008	Anderson et al.	2012/0128635 A1	5/2012	Gregory et al.
2008/0178299 A1	7/2008	Merkle et al.	2012/0129210 A1	5/2012	Bourdeaux et al.
2008/0206262 A1	8/2008	Banchereau et al.	2012/0178139 A1	7/2012	Hubbel et al.
2008/0213267 A1	9/2008	Young et al.	2012/0207745 A1	8/2012	Godfrin et al.
2008/0227707 A1	9/2008	Carlock et al.	2012/0237513 A1	9/2012	Zurawski et al.
2008/0233143 A1	9/2008	Jackson et al.	2012/0276095 A1	11/2012	Langermann et al.
2008/0241170 A1	10/2008	Zurawski et al.	2012/0282281 A1	11/2012	Banchereau et al.
2008/0254044 A1	10/2008	Zurawski	2013/0004427 A1	1/2013	El-Sayed et al.
2008/0261262 A1	10/2008	Godfrin	2013/0022634 A1	1/2013	Lycke
2008/0274092 A1	11/2008	Godfrin et al.	2013/0053543 A1	2/2013	Davis et al.
2008/0305104 A1	12/2008	Young et al.	2013/0059299 A1	3/2013	Parr et al.
2008/0318852 A1	12/2008	Anderson et al.	2013/0071413 A1	3/2013	Simon
2009/0004218 A1	1/2009	Hacohen et al.	2013/0078216 A1	3/2013	Dunlevy et al.
2009/0017039 A1	1/2009	Mi et al.	2013/0078267 A1	3/2013	Anderson et al.
2009/0123467 A1	5/2009	Bedi et al.	2013/0101463 A1	4/2013	Mambrini et al.
2009/0130104 A1	5/2009	Muzykantov et al.	2013/0115230 A1	5/2013	Simon
2009/0142263 A1	6/2009	Young et al.	2013/0129790 A1	5/2013	Alexis et al.
2009/0149656 A1	6/2009	Singaram et al.	2013/0164364 A1	6/2013	Paulson et al.
2009/0181011 A1	7/2009	Zaghouani	2013/0171074 A1	7/2013	Barbas et al.
2009/0191118 A1	7/2009	Young et al.	2013/0171233 A1	7/2013	Paulson et al.
2009/0202622 A1	8/2009	Fleury et al.	2013/0236533 A1	9/2013	Von Andrian et al.
2009/0269285 A1	10/2009	Anderson et al.	2013/0287810 A1	10/2013	Mohamadzadeh et al.
2009/0280132 A1	11/2009	Zaghouani	2013/0287857 A1	10/2013	Von Andrian et al.
2009/0317381 A1	12/2009	Plaut et al.	2013/0295120 A1	11/2013	Zurawski et al.
2009/0324538 A1	12/2009	Wong et al.	2013/0318648 A1	11/2013	Anderson et al.
2010/0003266 A1	1/2010	Simon	2013/0323786 A1	12/2013	Mi et al.
2010/0003338 A1	1/2010	Hubbell et al.	2013/0336991 A1	12/2013	Mi et al.
2010/0015131 A1	1/2010	Mi et al.	2014/0037736 A1	2/2014	Shi et al.
2010/0055189 A1	3/2010	Hubbell et al.	2014/0079728 A1	3/2014	Jackson et al.
2010/0092425 A1	4/2010	Von Andrian et al.	2014/0127198 A1	5/2014	Zurawski et al.
2010/0098718 A1	4/2010	Valiante	2014/0127301 A1	5/2014	Alexis et al.
2010/0129392 A1	5/2010	Shi et al.	2014/0134168 A1	5/2014	Zurawski et al.
2010/0129439 A1	5/2010	Alexis et al.	2014/0199315 A1	7/2014	Mi et al.
2010/0129820 A1	5/2010	Kool et al.	2014/0205630 A1	7/2014	Tartour
2010/0222407 A1	9/2010	Segura et al.	2014/0212445 A1	7/2014	Martin et al.
2010/0233251 A1	9/2010	Von Andrian et al.	2014/0227268 A1	8/2014	Banchereau et al.
2010/0239575 A1	9/2010	Banchereau et al.	2014/0234344 A1	8/2014	Banchereau et al.
2010/0285015 A1	11/2010	Muzykantov et al.	2014/0308238 A1	10/2014	Rubin-Bejerano et al.
2010/0291080 A1	11/2010	Lee et al.	2014/0314865 A1	10/2014	Von Andrian et al.
2010/0291082 A1	11/2010	Zurawski	2014/0377291 A1	12/2014	Fischbach et al.
2010/0297114 A1	11/2010	Zurawski	2014/0377838 A1	12/2014	Maynard et al.
2010/0310612 A1	12/2010	DuFour et al.	2015/0104478 A1	4/2015	Lee et al.
2010/0316620 A1	12/2010	Bourdeaux et al.	2015/0166659 A1	6/2015	Goldenberg et al.
2010/0322929 A1	12/2010	Zurawski et al.	2015/0191730 A1	7/2015	Levy et al.
2010/0330115 A1	12/2010	Zurawski et al.	2015/0250862 A1	9/2015	Cantor et al.
2011/0014171 A1	1/2011	Bourdeaux et al.	2015/0299329 A1	10/2015	Zurawski et al.
2011/0033426 A1	2/2011	Martin et al.	2015/0307545 A1	10/2015	Jackson et al.
2011/0044912 A2	2/2011	Anderson et al.	2016/0015821 A1	1/2016	Hubbell et al.
2011/0045049 A1	2/2011	Rubin-Bejerano et al.	2016/0022792 A1	1/2016	Zurawski et al.
2011/0064709 A1	3/2011	Miller et al.	2016/0024212 A1	1/2016	Goldenberg et al.
2011/0064754 A1	3/2011	Taylor et al.	2016/0031988 A1	2/2016	Zurawski et al.
2011/0082075 A1	4/2011	Prendergast	2016/0058792 A1	3/2016	Quintana et al.
2011/0091493 A1	4/2011	Mohamadzadeh et al.	2016/0060324 A1	3/2016	Paulson et al.
2011/0105379 A1*	5/2011	Shulman A61K 47/54 536/18.3	2016/0060358 A1	3/2016	Hay
2011/0123536 A1	5/2011	Chermann et al.	2016/0083468 A1	3/2016	Mi et al.
2011/0143994 A1	6/2011	Lycke	2016/0108096 A1	4/2016	Thompson et al.
2011/0177532 A1	7/2011	Rubin-Bejerano et al.	2016/0243248 A1	8/2016	Hubbell et al.
2011/0200632 A1	8/2011	Jackson et al.	2016/0346384 A1	12/2016	Porcelli et al.
2011/0206759 A1	8/2011	Swartz et al.	2016/0354453 A1	12/2016	Hubbell et al.
2011/0268804 A1	11/2011	Shi et al.	2016/0375126 A1	12/2016	Oh et al.
2011/0268805 A1	11/2011	Alexis et al.	2017/0007708 A1	1/2017	Hubbell et al.
2011/0293644 A1	12/2011	Anderson et al.	2017/0020926 A1	1/2017	Mata-Fink et al.
2011/0311542 A1	12/2011	Mi et al.	2017/0066825 A1	3/2017	Hubbell et al.
2012/0004643 A1	1/2012	Zurawski et al.	2017/0066828 A1	3/2017	Goldenberg et al.
2012/0009140 A1	1/2012	Godfrin et al.	2017/0121379 A1	5/2017	Zhang et al.
2012/0014960 A1	1/2012	Mi et al.	2017/0137513 A1	5/2017	Vallera et al.
2012/0027808 A1	2/2012	Iannacone	2017/0252417 A1	9/2017	Irvine et al.
2012/0039989 A1	2/2012	Hubbel et al.	2017/0296636 A9*	10/2017	Hubbell A61K 47/549
2012/0058180 A1	3/2012	Kren et al.	2017/0320933 A1	11/2017	Mannie
2012/0076831 A1	3/2012	Miller et al.	2017/0326213 A1	11/2017	Jajosky et al.
2012/0087890 A1	4/2012	Iannacone et al.	2018/0000916 A1	1/2018	Zurawski et al.
2012/0107301 A1	5/2012	Bowdish et al.	2018/0094071 A1	4/2018	Zurawski et al.
2012/0121570 A1	5/2012	Godfrin	2018/0100011 A1	4/2018	Hubbell et al.
			2018/0104284 A1	4/2018	Wallecha et al.
			2018/0112034 A1	4/2018	Maynard
			2018/0117171 A1	5/2018	Mooney et al.
			2018/0271986 A1	9/2018	Hubbell et al.
			2018/0303951 A1	10/2018	Hubbell et al.

(56)

References Cited**U.S. PATENT DOCUMENTS**

2019/0382479	A1	12/2019	Hubbell et al.
2020/0101146	A1	4/2020	Hubbell et al.
2020/0101169	A1	4/2020	Hubbell et al.
2020/0121762	A1	4/2020	Hubbell et al.
2020/0129601	A1	4/2020	Hubbell et al.
2020/0129625	A1	4/2020	Hubbell et al.
2020/0129629	A1	4/2020	Hubbell et al.

FOREIGN PATENT DOCUMENTS

CN	101443351	5/2009
CN	101750244	6/2010
CN	102791293	11/2012
CN	103282380	9/2013
CN	103547272	1/2014
CN	106432371	2/2017
EP	0119650	9/1984
EP	0175617	10/1991
EP	0088695	6/1992
EP	0173629	6/1992
EP	0480041	6/1993
EP	0308208	12/1993
EP	0251455	5/1994
EP	0294294	5/1995
EP	0789715	8/1997
EP	0808366	11/1997
EP	0722340	4/1998
EP	0505357	3/1999
EP	0602290	8/1999
EP	0978564	2/2000
EP	1012308	6/2000
EP	630407	8/2000
EP	1046651	10/2000
EP	1093464	4/2001
EP	1301541	4/2003
EP	0743856	7/2003
EP	1370588	12/2003
EP	1409009	4/2004
EP	1292621	9/2006
EP	1838734	10/2007
EP	1853313	11/2007
EP	1028978	1/2008
EP	1086137	6/2008
EP	1938836	7/2008
EP	1440156	8/2008
EP	1619208	10/2008
EP	1996700	12/2008
EP	1996701	12/2008
EP	1045861	3/2009
EP	2125012	12/2009
EP	2178896	4/2010
EP	1516881	6/2010
EP	2238986	10/2010
EP	2315779	5/2011
EP	1417229	6/2011
EP	2344185	7/2011
EP	2344187	7/2011
EP	2394657	12/2011
EP	2394661	12/2011
EP	2406290	1/2012
EP	2428226	3/2012
EP	2478917	7/2012
EP	2066294	10/2012
EP	2527363	11/2012
EP	2598120	6/2013
EP	2618817	7/2013
EP	2620157	7/2013
EP	2630967	8/2013
EP	1904104	9/2013
EP	1991564	9/2013
EP	2115129	11/2013
EP	2684889	1/2014
EP	1443963	5/2014
EP	1664270	5/2014
EP	2115002	8/2014

EP	1605974	11/2014
EP	1850832	12/2014
EP	2114985	12/2014
EP	2283358	4/2015
EP	2213742	1/2016
EP	2982695	2/2016
EP	2983791	2/2016
EP	2989123	3/2016
EP	2346528	4/2016
EP	2406286	5/2016
EP	2205273	9/2016
EP	3091034	11/2016
EP	2406288	12/2016
EP	2406289	2/2017
EP	2217269	4/2017
EP	2344186	4/2017
EP	2630966	4/2017
JP	S5742852	3/1982
JP	S59173762	10/1984
JP	H09-509572	9/1997
JP	2003-519619	6/2003
JP	2004-526452	9/2004
JP	2007-510915	4/2007
JP	2007-312776	12/2007
JP	2009-505049	2/2009
JP	2009-060894	3/2009
JP	2009-149664	7/2009
JP	2013-516967	5/2013
JP	2014-159492	9/2014
JP	2014-532799	12/2014
JP	106432371	2/2017
JP	2017507180	3/2017
JP	2018-072431	8/2018
WO	1991/008770	6/1991
WO	WO 1992/05801	4/1992
WO	WO 1992/22310	12/1992
WO	WO 1995/06737	3/1995
WO	WO 1995/22977	8/1995
WO	WO 1996/023882	8/1996
WO	WO 1996/040245	12/1996
WO	WO 1998/06737	2/1998
WO	WO 1999/036437	7/1999
WO	WO 1999/38536	8/1999
WO	WO 00/01732	1/2000
WO	WO 2000/074717	12/2000
WO	WO 2001/022995	4/2001
WO	WO 2001/025793	4/2001
WO	WO 2002/004522	1/2002
WO	WO 2002/072799	9/2002
WO	WO 2002/083262	10/2002
WO	WO 2002/102407	12/2002
WO	WO 2003/064464	8/2003
WO	WO 2003/066820	8/2003
WO	WO 2003/104273	12/2003
WO	WO 2004/034966	4/2004
WO	WO 2004/035619	4/2004
WO	WO 2004/045520	6/2004
WO	WO 2004/098645	11/2004
WO	WO 2005/045436	5/2005
WO	WO 2005/105129	11/2005
WO	WO 2006/002382	1/2006
WO	WO 2006/016247	2/2006
WO	WO 2006/093524	9/2006
WO	WO 2007/008300	1/2007
WO	WO 2007/017556	2/2007
WO	WO 2007/095748	8/2007
WO	WO 2007/097934	8/2007
WO	WO 2007/098254	8/2007
WO	WO 2007/099387	9/2007
WO	WO 2007/099446	9/2007
WO	WO 2007/101698	9/2007
WO	WO 2007/150020	12/2007
WO	WO 2008/063849	5/2008
WO	WO 2009/018500	2/2009
WO	WO 2009/019317	2/2009
WO	WO 2009/056332	5/2009
WO	WO 2009/078796	6/2009
WO	WO 2009/086552	7/2009
WO	WO 2009/120893	10/2009

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2010/042870	4/2010
WO	WO 2010/045518	4/2010
WO	WO 2010/060155	6/2010
WO	WO 2010/076517	7/2010
WO	WO 2010/085509	7/2010
WO	WO 2011/012715	2/2011
WO	WO 2011/051346	5/2011
WO	WO 2011/086143	7/2011
WO	WO 2011/092715	8/2011
WO	WO 2011/112482	9/2011
WO	WO 2012/021512	2/2012
WO	WO 2012/057671	5/2012
WO	WO 2012/083185	6/2012
WO	WO 2012/112690	8/2012
WO	WO 2012/167088	12/2012
WO	WO 2013/121296	8/2013
WO	WO 2013/160865	10/2013
WO	WO 2014/011465	1/2014
WO	WO 2014/023709	2/2014
WO	WO 2014/052545	4/2014
WO	WO 2014/135528	9/2014
WO	WO 2014/169255	10/2014
WO	WO 2015/140648	9/2015
WO	WO 2015/157595	10/2015
WO	WO 2015/171863	11/2015
WO	WO 2015/175957	11/2015
WO	WO 2016/022971	2/2016
WO	WO 2016/044655	3/2016
WO	WO 2016/044661	3/2016
WO	WO 2016/070050	5/2016
WO	WO 2016/210447	12/2016
WO	WO 2017/015141	1/2017
WO	WO 2017/023779	2/2017
WO	WO 2017/025889	2/2017
WO	WO 2017/041053	3/2017
WO	WO 2017/044308	3/2017
WO	WO 2017/046652	3/2017
WO	WO 2017/058996	4/2017
WO	WO 2017/066484	4/2017
WO	WO 2017/109134	6/2017
WO	WO 2017/112899	6/2017
WO	WO 2017/139498	8/2017
WO	WO 2017/139787	8/2017
WO	WO 2016104647	10/2017
WO	WO 2017/192785	11/2017
WO	WO 2017/192786	11/2017
WO	WO 2018/232176	12/2018
WO	WO 2019/098682	5/2019
WO	WO 2019/191079	10/2019
WO	WO 2019/0215590	11/2019
WO	WO 2021/053589	3/2021

OTHER PUBLICATIONS

"SubName: Full=Putative Transcriptional Regulator, ArsR Family," XP002717163, Retrieved From EBI Accession No. UNIPROT:D2RZT2, Database Accession No. D2RZT2 (Mar. 2, 2010).

"SubName: Full=Putative Uncharacterized Protein," XP002717158, Retrieved From EBI Accession No. UNIPROT:CONJE0, Database Accession No. CONJE0 (May 5, 2009).

"SubName: Full=Putative Uncharacterized Protein," XP002717160, Retrieved From EBI Accession No. UNIPROT:B9PUP0, Database Accession No. B9PUP0 (Mar. 24, 2009).

"SubName: Full=Uncharacterized Protein," XP002717157, Retrieved From EBI Accession No. UNIPROT:B5E9K2 Database Accession No. B5E9K2 (Oct. 14, 2008).

Ahmed et al., "Carbohydrate-based materials for targeted delivery of drugs and genes to the liver." *Nanomedicine (Lond.)* (2015) 10(14), 2263-2288.

Albert et al., "Immature dendritic cells phagocytose apoptotic cells via v135 and CD36, and cross-present antigens to cytotoxic T

lymphocytes," *Journal of Experimental Medicine*, vol. 188(7): 1359-1368 (Oct. 5, 1998).

Amagai, et al., "Desmoglein as a target in skin disease and beyond", *J Invest Dermatol*, Mar. 2012; in 23 pages.

Andre et al., "Determination of modulation of ligand properties of synthetic complex-type biantennary N-glycans by introduction of bisecting GlcNAc in silico, in vitro and in vivo" *Eur. J. Biochem.* 2004;271(1):118-134.

Arnaboldi et al., "Suppression of Th 1 and Th17, but not Th2, responses in a CD8+ T cell-mediated model of oral tolerance," *Mucosal Immunology*, vol. 2(5):427-438 (Sep. 2009).

Bailon et al., "Rational design of a potent, long-lasting form of interferon: A 40 kDa branched polyethylene glycol-conjugated interferon-2a for the treatment of hepatitis C," *Bioconjugate Chemistry*, vol. 12(2):195-202 (2001).

Baker, et al. "Hybrid Insulin Peptides are Autoantigens in Type 1 Diabetes", *Diabetes*, Sep. 2019, vol. 68, pp. 1830-1840.

Benaglia et al. "Searching for More Effective Agents and Conditions for the RAFT Polymerization of MMA: Influence of Dithioester Substituents, Solvent, and Temperature" *Macromolecules* 2005, 38, 3129-3140 (Year: 2005).

Bielekova et al., "Expansion and Functional Relevance of High-Avidity Myelin-Specific CD4 T Cells in Multiple Sclerosis," *J Immunol* 2004; 172:3893-3904.

Bigbee et al., "Binding specificities of eight monoclonal antibodies to human glycophorin A—studies with McM, and MkEn(UK) variant human erythrocytes and M- and MNv-type chimpanzee erythrocytes," Dec. 1, 1984, *J. Immunol.*, 133(6): 3149-3155 (1984).

Blancher et al., "Reactivity of anti-glycophorin monoclonal antibodies (Mabs) in tests with red cells of non-human primates," Jan. 1, 1997, *Transfus Clin Biol* 4, 81-85 (1997).

Boyer et al. "Bioapplications of RAFT Polymerization". *Chem. Rev.* 2009, 109, 5402-5436 (Year: 2009).

Brack et al., "Tumor-targeting properties of novel antibodies specific to the large isoform of tenascin-C," *Clinic Cancer Research*, vol. 12(10):3200-3208 (May 15, 2006).

Bursch et al., "Langerhans cells are not required for the COB T cell response to epidermal self-antigens," *Journal of Immunology*, vol. 182(8):4657-4664 (Apr. 15, 2009).

Caja et al., "Antibodies in celiac disease: implications beyond diagnostics," *Cellular & Molecular Immunology* (2011) 8, 103-109.

Calvaresi and Hergenrother. "Glucose conjugation for the specific targeting and treatment of cancer"; *Chem Sci. Jun.* 2013; 4(6): 2319-2333 (Year: 2013).

Cao et al., "Molecular Beacon Aptamers for Protein Monitoring in Real-Time and in Homogeneous Solutions," *Current Proteomics*, 2:31-401, (2005).

Chasis et al., "Signal Transduction by Glycophorin A: Role of Extracellular and Cytoplasmic Domains in a Modulatable Process", *The Journal of Cell Biology*, 107:1351-1357, (Oct. 1988).

Cheremisinoff. *Condensed Encyclopedia of Polymer Engineering Terms*, Butterworth-Heinemann, 2001, pp. 39-81. (Year: 2001).

Chiarantini et al., "Red Blood Cells as Delivery System for Recombinant HSV-1 Glycoprotein B: Immunogenicity and Protection in Mice," *Vaccine*, 15(3):276-280, (1997).

Ciccocioppo, R. et al., "The immune recognition of gluten in coeliac disease", *British Society for Immunology, Clinical and Experimental Immunology*, Feb. 1, 2005, pp. 408-416.

Clements et al., "The Crystal Structure of Myelin Oligodendrocyte Glycoprotein, a key autoantigen in multiple sclerosis," *Proc. Natl. Acad. Sci. (PNAS)* vol. 100: 11059-11064 (Sep. 2003).

Coulstock et al., "Liver-targeting of interferon-alpha with tissue-specific domain antibodies" *Plos One, Public Library of Science*, US, vol. 8, No. 2, Jan. 1, 2013.

Craig et al., "Processing of C3b-Opsonized Immune Complexes Bound to Non-Complement Receptor 1 (CR1) Sites on Red Cells: Phagocytosis, Transfer and Associations with CR1," *J. Immunol* 2005, 174, 3059-3066.

Crispe et al., "Cellular and molecular mechanisms of liver tolerance," *Immunol Rev.*, 213: 101-118 (2006).

Dane et al., "Isolation of cell specific peptide ligands using fluorescent bacterial display libraries-" *Journal of Immunological Methods*, vol. 309(1-2):120-129, (Jan. 2006).

(56)

References Cited

OTHER PUBLICATIONS

- Darrah et al., "IL-10 production differentially influences the magnitude, quality, and protective capacity of Th1 responses depending on the vaccine platform," *Journal of Experimental Medicine*, vol. 207(7):1421-1433 (2010).
- Deng, et al., "Disulfide-Based Self-Immolative Linkers and Functional Bioconjugates for Biological Applications," *Macromolecular Rapid Communication, Advanced Science News* (2020), 41, 1900531, in 14 pages.
- Dennis et al., "Albumin Binding as a General Strategy for Improving the Pharmacokinetics of Proteins," *Journal of Biological Chemistry*, vol. 277(38):35035-35043 (Sep. 20, 2002).
- Devalapally et al., "Poly(ethylene oxide)-modified Poly(beta-amino ester) Nanoparticles as a pH-sensitive System for Tumor-targeted Delivery of Hydrophobic Drugs: Part 3. Therapeutic Efficacy and Safety Studies in Ovarian Cancer Xenograft Model," *Cancer Chemotherapy Pharmacology*, 59:477-484, (2007).
- Dhalluin et al., "Structural and biophysical characterization of the 40 kDa PEG-interferon-2a and its individual positional isomers," *Bioconjugate Chemistry*, vol. 16(3):504-517 (2005).
- Di Lorenzo et al., "Translational Mini-Review Series on Type 1 Diabetes: Systemic analysis of T cell epitopes in autoimmune diabetes," 2007, *Clin Exp Immunol*, vol. 148: 1-146.
- Dienst et al., "Specific occlusion of murine and human tumor vasculature by VCAM-1-targeted recombinant fusion proteins," *Journal of The National Cancer Institute*, vol. 97(10):733-747, (2005).
- Dieterich et al., "Identification of Tissue Transglutaminase as the Autoantigen of Celiac Disease," *Nature Medicine* vol. 3 p. 797-801 (1997).
- Dominguez-Soto, et al., "The DC-SIGN-related lectin LSECtin mediates antigen capture and pathogen binding by human myeloid cells" www.bloodjournal.org. Immunobiology, Jun. 15, 2007, vol. 109, No. 12, pp. 5337-5345.
- Dornmair Klaus et al: "T-cell-mediated autoimmunity: Novel techniques to characterize autoreactive T-cell receptors", *American Journal of Pathology*, vol. 163, No. 4, Oct. 2003 (Oct. 2003), pp. 1215-1226, ISSN: 0002-9440.
- Ducan, R. Development of HPMA copolymer-anticancer conjugates: Clinical experience and lessons learnt. *Advanced Drug Delivery Reviews* 61 (2009) pp. 1131-1148.
- Ferguson et al., "Armed response: How dying cells influence T-cell functions," *Immunology Review*, vol. 241 (1):77-88 (May 2011).
- Fife et al., "Insulin-induced remission in new-onset NOD mice is maintained by the PD-1-PD-L1 pathway," *The Journal of Experimental Medicine*, vol. 203(12):2737-2747, (Nov. 27, 2006).
- Fishburn, "The pharmacology of PEGylation: balancing PD with PK to generate novel therapeutics," *Journal of Pharmaceutical Sciences* vol. 97(10):4167-4183 (Oct. 10, 2008).
- Folgori A et al: "A general strategy to identify mimotopes of pathological antigens using only random peptide libraries and human sera", *EMBO (European Molecular Biology Organization) Journal*, vol. 13, No. 9, May 1, 1994 (May 1, 1994), pp. 2236-2243, ISSN: 0261-4189.
- Fonsatti et al., "Targeting cancer vasculature via endoglin/CD105: A novel antibody-based diagnostic and therapeutic strategy in solid tumours," *Cardiovascular Research*, vol. 86(1):12-19, (2010).
- Gadaleta et al., "Trans-arterial chemoembolization as a therapy for liver tumours: New clinical developments and suggestions for combination with angiogenesis inhibitors," *Critical Reviews in Oncology/Hematology*, vol. 80:40-53 (2011).
- Gao et al., "In situ growth of a stoichiometric PEG-like conjugate at a protein's N-terminus with significantly improved pharmacokinetics," *Proceedings of the National Academy Sciences* vol. 106(36): 15231-15236 (Sep. 8, 2009).
- Geng et al., "Site-directed conjugation of "clicked" glycopolymers for form glycoprotein mimics: binding to mammalian lectin and induction of immunological function," *J Am Chem Soc*. Dec. 12, 2007;129(49):15156-63.
- Getts et al., "Have We Overestimated the Benefit of Human(ized) Antibodies?" *Landes Bioscience*, 2(6):682-694, (Nov./Dec. 2010).
- Getz et al., "Protease-Resistant Peptide Ligands From a Knottin Scaffold Library," *ACS Chemical Biology*, 8 Pages, (May 26, 2011).
- Godsel et al., "Prevention of autoimmune myocarditis through the induction of antigen-specific peripheral immune tolerance," *Circulation* vol. 103(12):1709-1714 (2001).
- Gorovits et al., "Proposed mechanism of off-target toxicity for antibody-drug conjugates driven by mannose receptor uptake," *Cancer Immunol Immunother* (2013) 62:217-233.
- Gorzalany et al., "Protein replacement therapies for rare diseases: a breeze for regulatory approval?" *Science Translational Medicine* 5, 178fs10 (2013).
- Granoff et al., "A Novel Mimetic Antigen Eliciting Protective Antibody to *Neisseria meningitidis*" *J Immunol* 2001; 167:6487-6496.
- Green et al., "Immunogenic and tolerogenic cell death," *National Review of Immunology* vol. 9(5):353-363, (May 2009).
- Grimm et al., "Memory of tolerance and induction of regulatory T cells by erythrocyte-targeted antigens," *Scientific Reports* 2015, 5:159907, 11 pages.
- Gupta et al., "Expression, purification, and characterization of an anti-RBC Fab-p24 fusion protein for hemagglutination-based rapid detection of antibodies to HIV in whole blood," *Protein Expression and Purification* 26 (2002) 162-170.
- Gurwitz, "Peptide Mimetics: Fast-Forward Look" *Drug Development Research* 78:231-235, Year 2017.
- Hackel et al., "Picomolar affinity fibronectin domains engineered utilizing loop length diversity, recursive mutagenesis, and loop shuffling," *Journal of Molecular Biology*, vol. 381(5):1238-1252, (Sep. 19, 2008).
- Hall et al., "Identification of peptide ligands facilitating nanoparticle attachment to erythrocytes," *Biotechnology Progress*, vol. 23(3):749-754 (2007).
- Hasselberg et al., "ADP-ribosylation controls the outcome of tolerance or enhanced priming following mucosal immunization" *The Journal of Immunology*, Aug. 24, 2016.
- Hirosue et al., "Antigen delivery to dendritic cells by poly(propylene sulfide) Nanoparticles with disulfide conjugated peptides: Cross-presentation and T cell activation," *Vaccine* 28, Elsevier Inc. (2010) p. 7897-7906.
- Holz et al., "CD8+ T cell tolerance following antigen recognition on hepatocytes," *Journal of Autoimmunity*, vol. 34 (1):15-22 (2010). <https://en.wikipedia.org/wiki/Retepase> accessed Apr. 13, 2020, printed Apr. 23, 2020 in 2 pages.
- <https://en.wikipedia.org/wiki/Tenecteplase>, accessed Apr. 13, 2020, printed Apr. 23, 2020 in 4 pages.
- Huang et al., "Characterization of poly(ethylene glycol) and PEGylated products by LC/MS with postcolumn addition of amines," *Analytical Chemistry*, vol. 81(2):567-577 (Jan. 15, 2009).
- Huang et al., "Tumor infarction in mice by antibody-directed targeting of tissue factor to tumor vasculature," *Science*, vol. 275(5299):547-550 (Jan. 24, 1997).
- Ichikawa et al., "Hepatic stellate cells function as regulatory bystanders," *Journal of Immunology*, vol. 186 (10):5549-5555 (May 15, 2011).
- Immunogenic, Definition of Immunogenic by Merriam-Webster, <https://www.merriam-webster.com/dictionary/immunogenic> [May 10, 2019 11:59:27 AM], retrieved on May 10, 2019, in 9 pages.
- International Search Report and Written Opinion from corresponding PCT Application No. PCT/IB2013/000684, 12 pages, dated Jul. 9, 2013.
- International Search Report and Written Opinion from corresponding PCT Application No. PCT/US2011/047078, 4 pages, dated May 1, 2012.
- International Search Report for Application No. PCT/EP2014/054161 issued on May 26, 2014.
- Jain, et al., "Mutual prodrugs containing bio-cleavable and drug releasable disulfide linkers," *Bioorganic Chemistry* 49, Elsevier Inc. (2013) p. 40-48.
- Janeway et al., "The complement system and innate immunity," *Immunology: the Immune System in Health and Disease*, 5th Edition. New York: Garland Science (2001).

(56)

References Cited

OTHER PUBLICATIONS

- Janeway et al., *Immuno Biology*, 8th Edition, Garland Science (2012).
- Jewett et al., "Cu-free click cycloaddition reactions in chemical biology," *Chem Soc Rev.* Apr. 2010; 39(4): 1272-1279.
- Jones et al., "Localization of Pectic Galactan in Tomato Cell Walls Using a Monoclonal Antibody Specific to (1->4)- β -D-Galactan" *Plant Physiol.* 1997; 113:1405-1412.
- Julyan et al. "Preliminary clinical study of the distribution of HPMa copolymers bearing doxorubicin and galactosamine" *Journal of Controlled Release* 57 (1999) pp. 281-290.
- Karim et al. "Hepatic expression and cellular distribution of the glucose transporter family"; *World J Gastroenterol* Dec. 14, 2012; 18(46): 6771-6781 (Year: 2012).
- Keefe et al., "Aptamers as therapeutics," *Nature Reviews Drug Discovery*, vol. 9(7):537-550 (2010).
- Kenrick et al., "Bacterial Display Enables Efficient and Quantitative Peptide Affinity Maturation," *Protein Engineering Design & Selection*, vol. 23(1):9-17 (2010).
- Khandelwal et al., "Assessment of survival of aging erythrocyte in circulation and attendant changes in size and CD147 expression by a novel two step biotinylation method," *Experimental Gerontology*, vol. 41(9):855-861 (Aug. 4, 2006).
- Kim et al. "Imaging and therapy of liver fibrosis using bioreducible polyethylenimine/siRNA complexes conjugated with N-acetylglucosamine as a targeting moiety" *Biomaterials* 34:6504-6514 (2013).
- Kim et al., "Specific Binding of Glucose-derivatized Polymers to the Asialoglycoprotein Receptor of Mouse Primary Hepatocytes," *The Journal of Biological Chemistry*, vol. 276, No. 38, pp. 35312-35319, Sep. 21, 2001.
- Kina et al., "The Monoclonal Antibody TER-119 Recognizes a Molecule Associated with Glycophorin A and Specifically Marks the Late Stages of Murine Erythroid Lineage," *British Journal of Haematology*, vol. 109:280-287 (2000).
- King et al. "Antibody responses to bee melittin (Api m 4) and hornet antigen 5 (Dol m 5) in mice treated with the dominant T-cell Epitope peptides" *Journal of Allergy and Clinical Immunology*, vol. 101, Issue 3, Mar. 1998, pp. 397-403.
- Klebe. "Optimization of Lead Structures", *Drug Design*, Springer-Verlag Berlin Heidelberg 2013, pp. 153-171. (Year: 2013).
- Kontos et al., "Engineering antigens for in situ erythrocyte binding induces T-cell deletion," *Proceeding of the National Academy Sciences*, Dec. 17, 2012, vol. 110, No. 1, p. E60-E68.
- Kontos et al., "Improving Protein Pharmacokinetics by Engineering Erythrocyte Affinity," *Molecular Pharmaceutics*, 2010, vol. 7, No. 6, p. 2141-2147.
- Kontos, "Engineering Erythrocyte Affinity for Improved Pharmacokinetics and Immune Tolerogenesis", Thesis, 106 Pages (Jun. 23, 2011).
- Kontos, et al., "Engineering antigen-specific immunological tolerance", www.sciencedirect.com, *Current Opinion in Immunology*, Jul. 8, 2015, 35:80-88.
- Kopecek et al. "HPMA copolymers: Origins, early developments, present, and future." *Advanced Drug Delivery Reviews* 62, (2010) pp. 122-149.
- Kravtsoff et al., "Tolerance Evaluation of L-asparaginase loaded in red blood cells," 1996, *Eur J Clin Pharmacol*, vol. 51: 221-225.
- Krebber et al., "Reliable Cloning of Functional Antibody Variable domains from Hybridomas and Spleen Cell Repertoires Employing a Reengineered Phage Display System," *Journal of Immunological Methods*, vol. 201:35-55 (1997).
- La Rosa, et al., "The Innate Immune System in Allograft Rejection and Tolerance," *J. Immunol.*, 2007, 178:7503-7509.
- Langer et al., "Optimization of the Preparation Process for Human Serum Albumin (HSA) Nanoparticles," *International Journal of Pharmaceutics*, 257:169-180, (2003).
- Lee et al., "Aptamers as Molecular Recognition Elements for Electrical Nanobiosensors," *Analytical and Bioanalytical Chemistry*, 390:1023-1032, (2008).
- Lee et al., "Signaling pathways downstream of pattern-recognition receptors and their cross talk," *Annual Review of Biochemistry*, vol. 76:447-480 (Feb. 28, 2007).
- Lehrman et al., "The Binding of Fucose-containing Glycoproteins by Hepatic Lectins" *The Journal of Biological Chemistry* Jun. 5, 1986; 261, 7426-7432.
- Lepeniec et al., "Targeting C-type lectin receptors with multivalent carbohydrate ligands," *Adv. Drug Deliv. Rev.* (2013).
- Li et al., "Targeting self- and foreign antigens to dendritic cells via DC-ASGPR generates IL-10-producing suppressive CD4+ T cells," *Jan. 2, 2012, Journal of Experimental Medicine* 209, 109-121 (2012).
- Liu et al. "Hapten may play an important role in food allergen-related intestinal immune inflammation," *North American Journal of Medical Sciences*, vol. 3. No. 3. (Mar. 2011).
- Liu et al., "Immune tolerance after delivery of dying cells to dendritic cells in situ," *Journal of Experimental Medicine*, vol. 196(8): 1091-1097 (Oct. 21, 2002).
- Liu et al., "Functional Nucleic Acid Sensors", *Chemical Reviews*, 109(5):1948-1998, (May 2009).
- Lobst et al., "Selective Sugar Binding to the Carbohydrate Recognition Domains of the Rat Hepatic and Macrophage Asialoglycoprotein Receptors," *The Journal of Biological Chemistry*, vol. 271, No. 12, Issue Mar. 22, 1996, 6686-6693.
- Loma et al., "Multiple Sclerosis: Pathogenesis and Treatment" *Department of Neurology, Curr. Neuroparmacology*, 9:409-416, Year 2011.
- Lorentz et al., "Engineered binding to erythrocytes induces immunological tolerance to *E. coli* asparaginase," *Sci. Adv.* 2015, 2015;1:e1500112, 10 pages.
- Luo et al., "ECDI-fixed allogeneic splenocytes induce donor-specific tolerance for long-term survival of islet transplants via two distinct mechanisms," *Proceedings of National Academy of Science*, vol. 105(38):14527-14532 (Sep. 23, 2008).
- Lutolf et al., "Synthesis and Physicochemical Characterization of End-Linked Poly(ethylene glycol)-co-peptide Hydrogels Formed by Michael-Type Addition," *Biomacromolecules*, 4:713-722, (Feb. 1, 2003).
- Lutolf et al., "Systematic modulation of Michael-type reactivity of thiols through the use of charged amino acids," *Bioconjugate Chemistry* vol. 12(6):1051-1056 (2001).
- Lutterotti, A. et al., "Antigen-Specific Tolerance by Autologous Myelin Peptide-Coupled Cells: A Phase I Trial in Multiple Sclerosis," *Science Translational Medicine* 5, 188ra75-188ra75 (2013).
- Magnani et al., "Red blood cells as an antigen-delivery system," *Biotechnol Appl Biochem.* Oct. 1992; 16(2):188-94.
- Maluccio et al., "Transcatheter arterial embolization with only particles for the treatment of unresectable hepatocellular carcinoma," *Journal of Vascular and Interventional Radiology*, vol. 19(6):862-869 (2008).
- Mamidyala, S. et al., "Glycomimetic ligands for the human asialoglycoprotein receptor" *J. Am Chem. Soc.* Feb. 1, 2012, 134(4), pp. 1978-1981.
- Martini, S., Nielsen, M., Peters, B. et al. *The Immune Epitope Database and Analysis Resource Program 2003-2018: reflections and outlook.* *Immunogenetics* 72, 57-76 (2020).
- Maynard et al., "Antibody engineering," *Annual Review of Bio-medical Engineering*, vol. 2:339-376 (2000).
- Meager et al., "Anti-cytokine autoantibodies in autoimmunity: preponderance of neutralizing autoantibodies against interferon-alpha, interferon-omega and interleukin-12 in patients with thymoma and or myasthenia gravis" *Clinical and Experimental Immunology*, Wiley-Blackwell Publishing Ltd, GB, vol. 132, No. 1, Apr. 1, 2003.
- Medina et al., "Targeting hepatic cancer cells with pegylated dendrimers displaying N-acetylgalactosamine and SP94 peptide ligands" *Advanced Healthcare Materials*, vol. 2, Issue 10, pp. 1337-1350, Oct. 2013.
- Meyer et al. *Metformin and Insulin in Type 1 Diabetes*; *Diabetes Care* 26:1655-1656, Year: 2003.
- Miller et al., "Antigen-specific tolerance strategies for the prevention and treatment of autoimmune disease," *Nature Reviews Immunology* 7(9):665-677, (Sep. 2007).

(56)

References Cited

OTHER PUBLICATIONS

- Mitea, C. et al., "A Universal Approach to Eliminate Antigenic Properties of Alpha-Gliadin Peptides in Celiac Disease," *PLoS One*, vol. 5, Issue 12, e15637, pp. 1-9, Dec. 2010.
- Moad et al., "Mechanism and Kinetics of Dithiobenzoate-Mediated RAFT Polymerization—Status of the Dilemma," *Macromolecular Journals Chem. Phys.* 2014, 215: 9-26.
- Moghim et al., "Stealth liposomes and long circulating nanoparticles: critical issues in pharmacokinetics, opsonization and protein-binding properties," *Progress in Lipid Research*, vol. 42(6):463-478 (2003).
- Mohandas et al., "Red cell membrane: past, present, and future," *Blood*, vol. 112(10):3939-3948 (Nov. 15, 2008).
- Moreau et al., "PEPOP: Computational design of immunogenic peptides" *BioMed Central*, Jan. 30, 2008, 15 pages.
- Moss et al., "Glossary of Class Names of Organic Compounds and Reactive Intermediates Based on Structure", *Pure & Appl. Chem.*, vol. 67, Nos. 819, pp. 1307-1375, 1995 (Year: 1995).
- Mueller, "Mechanisms maintaining peripheral tolerance," *Nature Immunology*, vol. 11(1):21-27 (Jan. 2010).
- Murphy, "Antigen Recognition by B-Cell and T-cell Receptors," 2012, Janeway's Immuno Biology, 8th Edition, Chapter 4, Garland Science Taylor & Francis Group, London and New York.
- Murray et al., "The Mouse Immune Response to Carrier Erythrocyte Entrapped Antigens," *Vaccine*, 24:6129-6139, (2006).
- Muzykantov, "Drug Delivery by Red Blood Cells: Vascular Carriers Designed by Mother Nature", *Expert Opinion Drug Delivery*, 7(4):403-427, (Apr. 2010).
- Nakayama, et al., "Determining Antigen Specificity of Human Islet Infiltrating T Cells in Type 1 Diabetes", *Frontiers in Immunology*, Mar. 8, 2019, vol. 10, pp. 1-7.
- Nardin et al., "How are immune complexes bound to the primate erythrocyte complement receptor transferred to acceptor phagocytic cells," *Mol. Immunol.* 1999, 36, 827-835.
- Nishikawa et al., "Galactosylated proteins are recognized by the liver according to the surface density of galactose moieties" *The American journal of physiology* Jun. 1995; 268(5 Pt 1):G849-56, Abstract.
- O'Neil et al., "Extracellular matrix binding mixed micelles for drug delivery applications," *Journal of Control Release*, vol. 137(2):146-151, (Mar. 27, 2009).
- Parmeggiani et al., "Designed armadillo repeat proteins as general peptide-binding scaffolds: consensus design and computational optimization of the hydrophobic core," *Journal of Molecular Biology*, vol. 376(5):1282-1304 (2008).
- Pasut et al., "PEG conjugates in clinical development or use as anticancer agents: An overview," *Advanced Drug Delivery Reviews*, vol. 61(13):1177-1188 (2009).
- Qin et al., "Galactosylated N-2-Hydroxypropyl Methacrylamide-b-N-3-Guanidinopropyl Methacrylamide Block Copolymers as Hepatocyte-Targeting Gene Carriers," *Bioconjugate Chem.* 22:1503-1512 (2011).
- Qin, et al., Preparation and bioactivity of anti-human red blood cell ScFv and CSFV E₀ bifunctional fusion protein, *Chin J. Biotech* 2010, Jan. 25: 26(1): 28-34, *Chinese Journal of Biotechnology* (2010).
- Rajpal, Arvind, et al. "A general method for greatly improving the affinity of antibodies by using combinatorial libraries", *PNAS*, www.pnas.org/cgi/doi/10.1073/pnas.0503543102, vol. 102 No. 24, pp. 8466-8471, Jun. 14, 2005.
- Reddy et al., "Exploiting lymphatic transport and complement activation in nanoparticle vaccines," *Nature Biotechnology*, vol. 25(10):1159-1164 (Oct. 2007).
- Reddy et al., "In vivo targeting of dendritic cells in lymph nodes with poly(propylene sulfide) nanoparticles," *Journal of Controlled Release*, vol. 112(1):26-34, (Mar. 10, 2006).
- Reinagel et al., "The Primate Erythrocyte Complement Receptor (CR1) as a Privileged Site: Binding of Immunoglobulin G to Erythrocyte CR1 Does Not Target Erythrocytes for Phagocytosis," 1997, *Blood*, vol. 89: p. 1068-1077.
- Rice et al., "Directed evolution of a biterminal bacterial display scaffold enhances the display of diverse peptides," *Protein Engineering, Design & Selection*, vol. 21(7):435-442 (2008).
- Rigopoulou et al., "Asialoglycoprotein receptor (ASGPR) as target autoantigen in liver autoimmunity: Lost and found," *Autoimmunity Reviews*, 12 (2012) 260-269.
- Rockey et al., "Synthesis and radiolabeling of chelator-RNA aptamerbioconjugates with copper-64 for targeted molecular imaging," *Bioorganic & Medicinal Chemistry*, vol. 19(13):4080-4090 (2011).
- Ruuslahti et al., "Targeting of drugs and nanoparticles to tumors," *Journal of Cell Biology*, vol. 188(6):759-768 (2010).
- Rybick et al., "The extra-domain A of fibronectin is a vascular marker of solid tumors and metastases," *Cancer Research*, vol. 67(22):10948-10957 (2007).
- Saibeni et al., "Antibodies to tissue-type plasminogen activator (t-PA) in patients with inflammatory bowel disease: high prevalence, interactions with functional domains of t-PA and possible implications in thrombosis," *J. Thrombosis and Haemostasis*, 4:1510-1516 (2006).
- Saint-Lu, N. et al., "Targeting the allergen to oral dendritic cells with mucoadhesive chitosan particles enhances tolerance induction," *Allergy*, vol. 64(7):1003-1013 (2009).
- Sakaguchi et al., "Regulatory T Cells and Immune Tolerance," *Cell* 133, May 30, 2008, 775-787.
- Sampson, "Aptamers and SELEX: the technology," *World Patent Information*, vol. (25):123-129 (2003).
- Savla et al., "Tumor targeted quantum dot-mucin 1 aptamer-doxorubicin conjugate for imaging and treatment of cancer," *Journal of Controlled Release*, vol. 153(1):16-22, Feb. 20, 2011.
- Schliemann et al., "In vivo biotinylation of the vasculature in B-cell lymphoma identifies BST-2 as a target for antibody-based therapy," *Vascular Blood*, vol. 115(3):736-744 (Jan. 21, 2010).
- Seamons et al. *Immune Tolerance to Myelin Proteins (Immunologic Research 2003; 28/3:201-221).*
- Sehon et al., "Conversion of Antigens to Tolerogenic Derivatives by Conjugation with Monomethoxypolyethylene Glycol", *The Pharmacology and Toxicology of Proteins*, pp. 205-219 (1987).
- Sehon et al., *The Pharmacology and Toxicology of Proteins*, Proceedings of Cetus—UCLA Symposium Held at Lake Tahoe, Ca, Feb. 21-28, 1987, Alan R. Liss, Inc.—New York.
- Seymour et al., "Hepatic Drug Targeting: Phase I evaluation of polymer-bound doxorubicin" *Journal of Clinical Oncology*, vol. 20, No. 6, Mar. 15, 2002, pp. 1668-1676.
- Seymour et al., "N-(2-Hydroxypropyl)methacrylamide copolymers targeted to the hepatocyte galactose-receptor: pharmacokinetics in DBA2 mice." *Br. J. Cancer* (1991) 63, pp. 859-866.
- Shan et al., "Structural Basis for Gluten Intolerance in Celiac Sprue," *Science*, 297, 2275 (2002).
- Shen "A galactosamine-mediated drug delivery carrier for targeted liver cancer therapy" *Pharmacological Research* 64 (2011) 410-419.
- Sheridan "Fresh from the biologic pipeline-2009," *Nature Biotechnology*, vol. 28(4):307-310 (Apr. 2010).
- Sigma-Aldrich, "RAFT Agents," available online at <https://www.sigmaaldrich.com/materials-science/material-science-products.html?TablePage=103936134>, 4 pages (accessed on Sep. 21, 2020) (Year: 2020).
- Silverman et al., "Engineered cystine-knot peptides that bind $\alpha\beta$ integrin with antibody-like affinities," *Journal of Molecular Biology*, vol. 382(4):1064-1075 (Jan. 30, 2009).
- Sørensen et al., "Role of sialic acid for platelet life span: exposure of β -galactose results in the rapid clearance of platelets from the circulation by asialoglycoprotein receptor-expressing liver macrophages and hepatocytes." *Blood*, Aug. 20, 2009. vol. 114, No. 8.
- Spitzer et al., "ScFv-Mediated In Vivo Targeting of DAF to Erythrocytes Inhibits Lysis by Complement," *Molecular Immunology*, vol. 40:911-919 (Oct. 30, 2003).
- St. Clair et al., "New Reagents on the Horizon for Immune Tolerance," Sep. 20, 2006, *Annu. Rev. Med.* 2007. 58:329-46.
- Staud et al., "Liver uptake and hepato-biliary transfer of galactosylated proteins in rats are determined by the extent of galactosylation" *Biochimica et Biophysica Acta* May 1999; 1427(2):183-192, Abstract.

(56)

References Cited**OTHER PUBLICATIONS**

- Steiner et al., "Efficient selection of DARPins with sub-nanomolar affinities using SRP phage display," *Journal of Molecular Biology*, vol. 382(5):1211-1227 (2008).
- Stern et al., "Promoting Tolerance to Proteolipid protein-induced experimental autoimmune encephalomyelitis through targeting dendritic cells," *Proc. Natl. Acad. Sci. (PNAS)* vol. 107: 17280-17285, (Oct. 2010).
- Sun, et al., "Comparison between Ovalbumin and Ovalbumin Peptide 323-339 Responses in Allergic Mice: Humoral and Cellular Aspects," *Scandinavian Journal of Immunology*, vol. 71: 329-335 (Jan. 2010).
- Supplementary European Search Report from corresponding PCT Application No. PCT/US2011047078, 21 Pages, Dated Jan. 22, 2014.
- Taneja et al., "Lessons from animal models for human autoimmune diseases," Sep. 1, 2001, *Nature Immunology*, vol. 2, No. 9, 781-784 (Sep. 2001).
- Taylor et al., "Anti-glycophorin single-chain Fv fusion to low-affinity mutant erythropoietin improves red blood cell-lineage specificity," *Protein Engineering, Design & Selection*, vol. 23, No. 4 pp. 251-260, 2010.
- Teitelbaum et al., Immunomodulation of experimental autoimmune encephalomyelitis by oral administration of copolymer I. *Pro. Natl. Acad. Sci. USA* vol. 96, pp. 3842-3847, Mar. 1999.
- Thijssen et al., "Galectin-1 is essential in tumor angiogenesis and is a target for antiangiogenesis therapy", *Proceeding of the National Academy Sciences*, vol. 103(43):15975-15980 (2006).
- Thomson et al., "Antigen-presenting cell function in the tolerogenic liver environment," *National Reviews Immunology*, vol. 10(11):753-766 (Nov. 2010).
- Tobio et al., "Stealth PLA-PEG Nanoparticles as Protein Carriers for Nasal Administration," *Pharmaceutical Research*, 15(2):270-275, (1998).
- Trahtenherts, A. et al., "An internalizing antibody specific for the human asialoglycoprotein receptor" *Hybridoma*, vol. 28, No. 4, Aug. 1, 2009.
- Turley et al., "Prospects for Antigen-Specific Tolerance Based Therapies for the Treatment of Multiple Sclerosis," *Results and Problems in Cell Differentiation*, 51 :217-235, (2010).
- Tye-Din, et al. "Comprehensive, Quantitative Mapping of T Cell Epitopes in Gluten in Celiac Disease", *www.Science TranslationalMedicine.org*, Jul. 21, 2010, vol. 2 Issue 41, in 14 pages.
- Updike et al., "Infusion of red blood cell-loaded asparaginase in monkey: Immunologic, metabolic, and toxicologic consequences," 1983, *J Lab Clin Med*, vol. 101(5): p. 679-691.
- Van Der Vlies et al., "Synthesis of pyridyl disulfide-functionalized nanoparticles for conjugating thiol-containing small molecules, peptides, and proteins," *Bioconjugate Chemistry*, vol. 21(4):653-662 (2010).
- Velluto et al., "PEG-b-PPS Diblock Copolymer Aggregates for Hydrophobic Drug Solubilization and Release: Cyclosporin A as an Example," *Molecular Pharmaceutics*, 11 Pages, (May 2, 2008).
- Vogl et al., "Review on transarterial chemoembolization in hepatocellular carcinoma: Palliative, combined, neoadjuvant, bridging, and symptomatic indications," *European Journal Radiology*, vol. 72(3):505-516 (2009).
- Walker et al., "Anti-serum albumin domain antibodies in the development of highly potent, efficacious and long-acting interferon," *Protein Engineering Design & Selection*, vol. 23(4):271-278 (2010).
- Wan, "Regulatory T cells: immune suppression and beyond," May 1, 2010, *Cell Mol Immunol*. May 2010; 7(3):204-210.
- Wang et al in "Diagnostic imaging and therapeutic application of nanoparticles targeting the liver"; *J. Mater. Chem. B*, 2015, 3, 939. (Year: 2015).
- Wang et al., "Synthesis and Micellization of Thermoresponsive Galactose-Based Diblock Copolymers," *J Polymer Sci.* 49:3280-3290 (2011).
- Weisser et al., "Applications of single-chain variable fragment antibodies in therapeutics and diagnostics," *Biotechnology Advances*, vol. 27(4):502-520 (2009).
- Wilcock, H. et al. "End Group Removal and Modification of RAFT polymers," *Polymer Chemistry*, vol. 1, Jan. 1, 2010, pp. 149-157.
- Wilson et al., "Rapid Whole Blood Assay for HIV-1 Seropositivity Using an Fab-Peptide Conjugate," *Journal of Immunological Methods*, vol. 138:111-119 (1991).
- Wilson, D.B., "Kent et al. Replying to: D.B. Wilson", *Nature*, 438, 2005.
- Wu, Herren, "Simultaneous Humanization and Affinity Optimization of Monoclonal Antibodies", *Methods in Molecular Biology*, vo. 207: Recombinant Antibodies for Cancer Therapy: Methods and Protocols, Tolowa, NJ, pp. 197-212, Jan. 1, 2003.
- Yamazaki et al., "CD8+ CD205+ splenic dendritic cells are specialized to induce Foxp3+ regulatory T cells," *Journal of Immunology*, vol. 181(10):6923-6933 (2008).
- Yeste Ada et al: "Antigen Microarrays for the Study of Autoimmune Diseases", *Clinical Chemistry*, vol. 59, No. 7, Jul. 2013 (Jul. 2013), pp. 1036-1044, ISSN: 0009-9147(print).
- Yoo et al., "N-Acetylgalactosamine dendrons as clearing agents to enhance liver targeting of model antibody-fusion protein." *Bioconjugate Chemistry*, vol. 24, No. 12, Dec. 18, 2013, pp. 2088-2103.
- Zaitsev et al., "Targeting of a Mutant Plasminogen Activator to Circulating Red Blood Cells for Prophylactic Fibrinolysis", *The Journal of Pharmacology and Experimental Therapeutics*, 332(3):1022-1031 and 976 (Nov. 30, 2009).
- Zhao, X. et al "Construction and characterization of an anti-asialoglycoprotein receptor single-chain variable-fragment-targeted melittin" *Biotechnology and Applied Biochemistry*, Nov.-Dec. 2011; 58(6): pp. 405-411.
- Zhong et al., "Ligand-directed Reduction-Sensitive Shell-Sheddable Biodegradable Micelles Actively Deliver Doxorubicin into the Nuclei of Target Cancer Cells," *Biomacromolecules* 14:3723-3730 (2013).
- English translation of Office Communication issued in Japanese Patent Application No. 2020-563514, dated Feb. 7, 2024.
- Li et al., "Shape Effect of Glyco-Nanoparticles on Macrophage Cellular Uptake and Immune Response," *ACS Macro Letters*, 5(9):1059-1064, 2016.
- Brungel, K. L., *Introducing tolerance to foreign and partly foreign proteins by erythrocyte binding and artificial glycosylations*. 2018. *Ecole Polytechnique Federale de Lausanne*, Thesis 8190.
- Office Action issued in Corresponding Singapore Application No. 11202011078V, dated Jun. 30, 2022.
- International Search Report and Written Opinion issued in Corresponding PCT Application No. PCT/US2019/31440, dated Sep. 13, 2019.
- Office Action issued in corresponding Japanese Application No. 2020563514 dated Jun. 19, 2023.
- English translation of Office Communication issued in Chinese Patent Application No. 201980045533.2, dated Nov. 8, 2024.
- Chungsong et al., "Bioconjugate Strategies for the Induction of Antigen-Specific Tolerance in Autoimmune Diseases," *Bioconjug. Chem.*, 29(3):719-732, 2018.
- Schuetz et al., "Mannose receptor induces T-cell tolerance via inhibition of CD45 and up-regulation of CTLA-4," *PNAS*, 113(38):10649-10654, 2016.

* cited by examiner




FIG. 1A-D




FIG. 1E-G




FIG. 2A-B




FIG. 3A-C




FIG. 3D-E




FIG. 4A-B




FIG. 5




FIG. 6




FIG. 7A-B

1

COMPOSITIONS AND METHODS CONCERNING IMMUNE TOLERANCE

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a national phase application under 35 U.S.C. § 371 of International Application No. PCT/US2019/031440 filed May 9, 2019, which claims priority to U.S. Provisional Patent Application No. 62/669,044, filed May 9, 2018, all of which are incorporated in their entirety by reference.

BACKGROUND OF THE INVENTION

1. Field

Embodiments of the present invention relate generally to the fields of organic chemistry, biochemistry, and immunology.

2. Background

Small molecules, e.g., non-peptide or nucleic acid-based molecules having a molecular weight of less than 500 Daltons, have historically dominated the FDA's new molecular entity (NME) approvals. Since the FDA's approval of insulin as the first drug produced by recombinant DNA technology in 1982, protein-based drug approvals have experienced an upward trend, as demonstrated by the almost yearly increase in FDA biologics license application (BLA) approvals in the last 23 years.

A primary difference between a small molecule drug and a protein-based drug lies in the ability of antigen presenting cells to take up proteins, including protein-based drugs, process them, and present them as peptides to the major histocompatibility complex (MHC) class I and II to the immune system. Protein-based drugs are expressed and purified to be free of any pathogen-associated molecular patterns (PAMPs) and danger associated molecules (DAMPs). Notwithstanding the anti-immunogenic design of protein-based drugs, a significant portion of patients who are treated with these drugs develop antibodies against the drugs (anti-drug antibodies).

These and other examples demonstrate that protein-based drug immunogenicity can revert the effect of therapy for certain patients or even render a new approach to treat a disease inefficient. The negative effects provided by anti-drug antibodies highlight the need to develop a platform that allows clinicians induce tolerance to foreign and partly foreign proteins.

Technologies to induce antigen-specific immunological tolerance are still in their infancy, but are needed to prevent immunity to many protein-based drugs, to reverse immunity to allergens, and to prevent and reverse immunity to autoimmune antigens.

SUMMARY

A useful tolerogenic drug would consist of an antigen and a tolerance inducing component. The tolerance inducing component can be a chemical conjugated entity, part of a fusion protein, nanoparticles, or cells that are pulsed with the antigen. Ideally, a tolerance-inducing drug is based on the drug it tries to tolerize against, and can easily be modified.

Disclosed herein are compositions and methods for inducing tolerance towards therapeutic proteins, e.g., protein-


2

based drugs. In some aspects, the present disclosure provides therapeutic, polymer-tethered antigens that include mannose monomers or derivatives thereof. In some embodiments, derivatives of mannose include optionally substituted mannose substituents. In several embodiments, the mannose derivative comprises a mannose with a phosphate at the C1, C2, C3, C4, C5, and/or C6 position. In several embodiments, the targeting moiety comprises mannose-6-phosphate. In some embodiments, compositions including a mannose or mannose-derived moiety induce tolerance to immunogenic protein-based therapeutics, delete antigen-specific CD4 and CD8 T cells, elevate levels of regulatory T cell responses and IL-10 producing Breg cells, and/or reduce antigen-specific plasma cells and memory B cells. The compositions disclosed herein may be targeted for delivery to antigen-presenting cells including, but not limited to, hepatocytes, LSECs, Kupffer cells, and stellate cells. In some aspects, the compositions disclosed herein exhibit affinity for and specifically bind to mannose-binding receptors.

In several embodiments, there are provided compounds, as well as compositions comprising such compounds, of Formula 1:



where X comprises an antigen, a tolerogenic portion thereof, or a mimetic thereof, Y comprises a linker moiety, Z comprises a moiety that specifically targets a mannose receptor, p is an integer from about 2 to about 250, m is an integer from about 1 to about 100, and R² is any of functional groups I-III:










where Ar is a substituted or unsubstituted aromatic group, where R³ is any carbon-containing linear or heterocyclic moiety, and R¹¹ is hydrogen or an alkyl group.







In several embodiments, the moiety that specifically targets a mannose receptor is selected from the group consisting of α-linked mannose, β-linked mannose, substituted mannose, mannose-6-phosphate, N-acetyl mannosamine, and mannan having β(1-4), α(1-6), α(1-2), and/or α(1-3) linkages. In several embodiments, a plurality of these targeting moieties are used in combination in order to enhance targeting of and/or binding to the mannose receptor.

In several embodiments, Y is a linker resulting from reaction of at least one of a N-hydroxysuccinamidyl linker, maleimide linker, PEG linker, vinylsulfone linker, pyridyl di-thiol-poly(ethylene glycol) linker, pyridyl di-thiol linker, n-nitrophenyl carbonate linker, NHS-ester linker, and nitro-phenoxy poly(ethylene glycol)ester linker. In some embodiments, Y is covalently bound to X.

Depending on the embodiment, the following portion of Formula 1 ($-\text{[Y(Z)p)]-}$) is represented by one of Formula Ya to Yr:


$$Y_a$$

$$Y_c$$







Ye

 Y_g 

5

6

-continued


$$Y_n$$

$$Y_0$$
 Y_p  Y_q 

Yr

where

n is an integer from about 1 to about 100;

q is an integer from about 1 to about 44;

k is an integer from about 1 to about 12;

i is an integer from 0 to about 20;


v is an integer from about 1 to about 4;

p is an integer from about 2 to about 250;


r is an integer from 0 to about 250;

R_1 is $-\text{CH}_2-$, $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)(\text{CN})-$, $-(\text{CH}_2)_2$
 $-\text{C}(\text{CH}_3)(\text{CH}_3)-$, $-(\text{CH}_2)_2-\text{CH}(\text{CH}_3)-$ or
 $-\text{CH}(\text{CH}_3)-$; 55

W^1 and W^2 are as defined below:



-continued



50

R⁹ is a direct bond, $-(CH_2)_2-NH-C(O)-$ (an ethylacetamido group or "EtAcN") or $-(CH_2)_2-(O-CH_2-CH_2)_t-NH-C(O)-$ (a pegylated ethylacetamido group or "Et-PEG_t-AcN")

60 t is an integer from 1 to 5, Z is mannose or a mannose
receptor-targeting moiety; and R¹⁰ is an aliphatic
group, an alcohol, an aliphatic amine-containing group,
or an aliphatic alcohol.

In several embodiments, Y is an antibody, antibody frag-
65 ment, peptide or other ligand that binds to X.

In several embodiments, X is an antigen against which a patient may develop or has developed an unwanted immune

response. For example, depending on the embodiment, the antigen may be a foreign transplant antigen, an alloantigen, an autoimmune antigen, a food antigen, an animal antigen, a plant antigen, an environmental antigen, a therapeutic antigen, a synthetic self-antigen, or a tolerogenic (e.g., immunogenic, or capable of inducing an immune response) portion thereof. In several embodiments, X is an asparaginase antigen or an ovalbumin antigen. In several embodiments, the antigen is comprised in a vesicle, cell fragment, or cell.

In several embodiments, the antigen comprises at least one autoimmune antigen or tolerogenic portion thereof. In several embodiments, the at least one autoimmune antigen comprises at least one of an immunogenic fragment or fragments of myelin basic protein (MPB), an immunogenic fragment or fragments of myelin oligodendrocyte glycoprotein (MOG), an immunogenic fragment or fragments of myelin proteolipid protein (PLP), MBP, MOG, or PLP. In several embodiments, the at least one autoimmune antigen comprises at least one of SEQ ID Nos. 23-47. In several embodiments, the at least one autoimmune antigen comprises at least one of SEQ ID Nos. 24, 25, 27, 28, 31, 32, 33, 34, 35, 36, 43, 44, 45, 46, and 47. In some embodiments, the compound optionally further comprises at least one of SEQ ID NOs: 29, 38, 39, 40, 41, and 42. In several embodiments, such compounds are for use in treatment of or prevention of multiple sclerosis. In some embodiments, such compounds are administered for the use of preventing multiple sclerosis in a subject predicted to have multiple sclerosis. In some aspects, such compounds are administered to a subject presenting one or more symptoms of multiple sclerosis.


In several embodiments, the at least one autoimmune antigen comprises at least one of insulin, proinsulin, pre-proinsulin, glutamic acid decarboxylase-65 (GAD-65 or glutamate decarboxylase 2), GAD-67, glucose-6 phosphatase 2, islet-specific glucose 6 phosphatase catalytic subunit related protein (IGRP), insulinoma-associated protein 2 (IA-2), insulinoma-associated protein 2 β (IA-2 β), ICA69, ICA12 (SOX-13), carboxypeptidase H, Imogen 38, GLIMA 38, chromogranin-A, HSP-60, carboxypeptidase E, peripherin, glucose transporter 2, hepatocarcinoma-intestine-pancreas/pancreatic associated protein, S100 β , glial fibrillary acidic protein, regenerating gene II, pancreatic duodenal homeobox 1, dystrophin myotonia kinase, and SST G-protein coupled receptors 1-5, or an immunogenic fragment of any of these antigens. In several embodiments, the autoimmune antigen comprises an immunogenic fragment of proinsulin. Optionally an immunogenic fragment of IA-2, GAD-65, GAD-67, insulin, and/or IGRP is included. In several embodiments, the at least one autoimmune antigen comprises at least one of SEQ ID NOs: 1-19, or an immunogenic fragment of any of SEQ ID NOs: 1-19. In several embodiments, the at least one autoimmune antigen comprises at least one of SEQ ID NOs: 4-19. In several embodiments, such compounds are for use in the treatment or prevention of Type 1 Diabetes. In some embodiments, such compounds are administered for the use of preventing Type 1 Diabetes in a subject predicted to have Type 1 Diabetes. In some aspects, such compounds are administered to a subject presenting one or more symptoms of Type 1 Diabetes.

In several embodiments, the antigen comprises a food antigen, or a tolerogenic portion thereof. In several embodiments, the antigen comprises at least one of tissue transglutaminase, high molecular weight glutenin, low molecular weight glutenin, gluten, alpha-gliadin, gamma-gliadin, omega-gliadin, hordein, secalin, avenin, and deamidated

forms thereof. In several embodiments, the antigen comprises a tolerogenic portion of at least one of tissue transglutaminase, high molecular weight glutenin, low molecular weight glutenin, gluten, alpha-gliadin, gamma-gliadin, omega-gliadin, hordein, secalin, avenin, and deamidated forms thereof. In several embodiments, the antigen comprises at least one of SEQ ID NOs. 54-61. In several embodiments, such compounds are for use in the treatment or prevention of Celiac Disease. In some embodiments, such compounds are administered for the use of preventing Celiac Disease in a subject predicted to have Celiac Disease. In some aspects, such compounds are administered to a subject presenting one or more symptoms of Celiac Disease.

In several embodiments, the mannose receptor is mannose-6-phosphate receptor. In several embodiments, Y and X are connected through a bond configured to cleave when the compound reaches a target area. Advantageously, this cleavage, in several embodiments, is triggered when the compound is at a target site (e.g., site where the mannose receptor is bound). This allows, in several embodiments, delivery of the free antigen to the target site.


In several embodiments, Ar is selected from:



where each instance of R'', when present, is independently selected from an optionally substituted C1-6-alkyl, optionally substituted C1-6 alkoxy, optionally substituted amino, OH, or halogen and wherein, X'' is a heteroatom. In several embodiments, X'' is N. In several embodiments, R11 is C1-6-alkyl. In several embodiments, R11 is —CH3. In several embodiments, R3 is C1-6-alkyl.




There are also provided herein, compositions comprising the compounds as described above, or elsewhere herein. Likewise, there are provided for herein uses of these compounds for inducing tolerance (or treating an unwanted immune response) to an antigen, a tolerogenic portion (or portions) of one or more antigens, and/or to mimetics of the antigens or portions of antigens. Also provided are uses of such compounds in the preparation of a medicament for inducing tolerance (or treating an unwanted immune response) to an antigen, a tolerogenic portion (or portions) of one or more antigens, and/or to mimetics of the antigens or portions of antigens.

Certain aspects of the disclosure are directed towards compositions comprising a compound of Formula 1:







Formula 1

where X comprises an antigen or a tolerogenic portion thereof, Y comprises a linker moiety, Z comprises a moiety that specifically targets a mannose receptor, p is an integer from about 2 to about 250, m is an integer from about 1 to about 100, R² is any of functional groups I-III:



where Ar is a substituted or unsubstituted aromatic group,

R^3 is any carbon-containing linear or heterocyclic moiety, and R^{11} is hydrogen or an alkyl group. In some embodiments, R^2 comprises an end-capping group. In some embodiments, R^2 when disconnected from the construct, forms a stable or substantially stable free radical. In some embodiments, R^2 is a reversible addition-fragmentation chain transfer (RAFT) agent for a living polymerization. In some embodiments, R^2 can be reversibly added and removed to the construct to lengthen the linker region. In some embodiments, R^2 is a RAFT agent. In some embodiments, R^2 is not a RAFT agent. In some embodiments, R^2 is H or is absent. In some embodiments, R^2 is an optionally substituted dithiobenzoate, a trithiocarbonate, or a xanthate. In some embodiments, R^3 or R^{11} may be hydrogen, optionally substituted alkyl, optionally substituted cycloalkyl, optionally substituted aryl, or optionally substituted heteroaryl. In some embodiments, R^3 is hydrogen, optionally substituted C_6 -aryl, or C_{1-6} -alkyl (optionally substituted with halogen, or hydroxyl). In some embodiments, Ar as provided above is phenyl (optionally substituted with one or more OH groups, NH_2 groups, and/or halogens). In some embodiments, R^2 is one of the functional groups:




where R³ is as defined above. In several embodiments, Ar in any of functional groups I or II is an optionally substituted C₆-C₁₄ aryl or an optionally substituted heteroaryl having 6 to 14 ring members of which 1-4

are heteroatoms. In several embodiments, the optionally substituted C₆-C₁₄ aryl is optionally substituted with one or more functional groups selected from C₁-C₆alkyl, amino, halogen, —OH, or combinations thereof. In several embodiments, Ar is selected from the group consisting of:


I
II

III




where each instance of Rⁿ, when present, is independently selected from an optionally substituted C₁₋₆-alkyl, optionally substituted C₁₋₆ alkoxy, optionally substituted amino, OH, or halogen. In several embodiments, Xⁿ is a heteroatom. In several embodiments, Xⁿ is N. In several embodiments, R¹¹ is C₁₋₆-alkyl. In several embodiments, R¹¹ is —CH₃. In several embodiments, R³ is C₁₋₆-alkyl.

In some embodiments, Y and X are connected via a bond that cleaves or is configured to cleave at a target site for the compound. In several embodiments, Y and X are bonded through a disulfanyl ethyl ester or a disulfide bond. In several embodiments, the bond between Y and X is configured to cleave when the compound of Formula 1 reaches its biological target (e.g., the liver, liver cells, and/or the cytosol of cells in the liver) in a patient. In several embodiments, the Y—X bond cleaves in the presence of a cellular reducing agent (e.g., glutathione). In some embodiments, advantageously, once the bond between X and Y is cleaved, X is left in its native form and/or an active form. In some embodiments, once X is cleaved from Y, it is in a form that is more active than when bound to Y. In some embodiments, dithiol-containing compounds, particularly disulfanylethyl carbamate-containing links between X and Y (named including a free amine of X, otherwise named a “disulfanyl ethyl ester” without including the free amine of X) are advantageous as having the ability to cleave and release an antigen in its original form once inside a cell, for example as illustrated below (where Y' indicates the remaining portion of the linker and X and Z are as defined)



11




In several embodiments, the liver targeting moiety of the compound of Formula 1 is not galactose, galactosamine, N-acetylgalactosamine, glucose, glucoseamine and/or N-acetylglucosamine.

In some embodiments, Y is a linker. In several embodiments, Y is a reaction product resulting from one or more

12


reactions involving at least one of the following: N-hydroxysuccinamidyl (NHS) linker, NHS ester linker, PEG linker, maleimide linker, vinylsulfone linker, pyridyl di-thiol-poly (ethylene glycol) linker, pyridyl di-thiol linker, n-nitrophenyl carbonate linker, or a nitrophenoxy poly(ethylene glycol) ester linker. The linker may have one or more mannose moieties or mannose receptor-targeting moieties bound to it. In several embodiments, Y comprises an antibody, an antibody fragment, a peptide, or a disulfanyl ethyl ester to which one or more mannose moieties or mannose receptor-targeting moieties are bound. In some aspects, $-[Y(Z)_p]-$ is a group represented by one of Formula Ya to Yr:



13

14

-continued




where the left, opening parentheses “(” signifies the location of the bond between X and Y, the right, closing parentheses “)” signifies the location of the bond between Y and R². In several embodiments, n is an integer greater than or equal to about: 1, 10, 20, 40, 50, 75, 100, 150 or ranges including and/or spanning the aforementioned values. In several embodiments, n is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, or ranges including and/or spanning the aforementioned values. In several embodiments, q is an integer greater than or equal to about: 1, 10, 20,

40, 50, 75, 100, 150 or ranges including and/or spanning the aforementioned values. In several embodiments, q is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 75, 100, 150, or ranges including and/or spanning the aforementioned values. In several embodiments, k is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, v is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values.

15

In several embodiments, k is 2. In several embodiments, v is 2. In several embodiments, n is 4. In several embodiments, n is 44. In several embodiments, q is 3. As used herein, variables disclosed as having structure, a value, or a range of values for one embodiment, may also have those values when the variable is used in another embodiment (even where the variable is not defined with respect to that other embodiment). In several embodiments, n is an integer from 1 to 100. In several embodiments, q is an integer from 1 to 44. In several embodiments, k is an integer from 1 to 12. In several embodiments, i is an integer from 0 to 20. In several embodiments, v is an integer from 1 to 4. In several embodiments, R₁ is —CH₂—, —(CH₂)₂—C(CH₃)(CN)—, —(CH₂)₂—C(CH₃)(CH₃)—, —(CH₂)₂—CH(CH₃)— or —CH(CH₃)—. In several embodiments, W¹ and W² are as depicted below:







where Z is mannose or a mannose receptor-targeting moiety, R⁹ is a direct bond, —(CH₂)₂—NH—C(O)— (an ethylacetamido group or “EtAcN”) or —(CH₂)₂—(O—CH₂—CH₂)_t—NH—C(O)— (a pegylated ethylacetamido group or “Et-PEG_t-AcN”), t is an integer from 1 to 5, p is an integer from 2 to 250, R¹⁰ is an aliphatic group, an alcohol, an aliphatic amine-containing group, or an aliphatic alcohol, and r is an integer from 0 to 250. In several embodiments, —W¹_p—W²_r— (e.g., as provided in —[Y(Z)_p]_r— or in linker structures) is a random copolymer or block copolymer of W₁ and W₂. In several embodiments, the number of repeat units of W¹ is denoted as p and wherein p is an integer of at least about 1. In several embodiments, the number of repeat units of W² is denoted as r and wherein r is an integer of at least about 1. In some embodiments, R¹⁰ is a C_falkyl or C_falkylOH_g, where f represents the number of carbons in the alkyl group and is an integer between 0 and 10, and g represents the number of hydroxyl groups present on the alkyl group and is an integer between 0 and 10. In some embodiments, R¹⁰ is 2-hydroxyethyl. In some aspects —W¹_p—W²_q— represents a block copolymer or a random copolymer of W¹ and W² monomers.

In several embodiments, the linker comprises a polymeric chain with pendant liver targeting moieties decorating the polymeric chain. In some embodiments, the polymeric chain (or Y) comprises Y' as disclosed elsewhere herein. In several embodiments, the polymeric chain comprises an acrylate portion (e.g., acrylate-based polymers and/or acrylate-based copolymers). In several embodiments, the acrylate portion comprises one or more acrylate units (e.g., acrylate derivatives, including methacrylates and derivatives thereof) comprising a pendant liver targeting agent. In several embodi-


16

ments, the polymeric chain comprises a hydrophilic portion and/or region. In several embodiments, the hydrophilic portion comprises a length of one or more regions having —(CH₂CH₂O)_s— where s is an integer from about 1 to about 44. In several embodiments, s is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 75, 100, 150, or ranges including and/or spanning the aforementioned values. In some embodiments, the hydrophilic portion comprises one or more polyethylene glycol (PEG) regions. In some embodiments, the PEG may have polydispersity as measured by the weight average molecular weight in g/mol (Mw) of the PEG divided by the number average molecular weight in g/mol (Mn) of the PEG (e.g., Mw/Mn). In some embodiments, the PEG chains have a number average or weight average molecular weight (g/mol) of equal to or at least about: 500, 1000, 2000, 5000, 10000, or ranges including and/or spanning the aforementioned values. In several embodiments, the polymeric chain is optionally substituted. In some embodiments, the polymeric chain comprises pendant hydrophilic groups such as a —OH, —SO(OH)₂, optionally substituted polyether, optionally substituted polyamino, and the like.


In several embodiments, the antigen and liver targeting portion of the compound are joined using click chemistry, for example, by functionalizing the antigen with a first linker arm comprising an alkynyl group (or an azide), functionalizing the liver targeting moiety with a second linker arm comprising an azide (or an alkynyl group), and clicking them together via “click” chemistry. In some embodiments, an alkynyl group that can be clicked in copper-free conditions is used. In some embodiments, —[Y(—Z)_p]_r— is a group represented by one or more of Formulae AI-AIV:



where the left, opening parentheses “(” signifies the location of the bond between X and Y, the right, closing parentheses “)” signifies the location of the bond between Y and R², Y' is a random copolymer or block copolymer of two or more different types of repeat units, wherein at least one type of repeat unit comprises a pendant Z group, (or plurality of pendant Z groups) where Z is mannose and/or a mannose receptor-targeting moiety. In some embodiments, Y' is a random copolymer or block copolymer of W¹ and W², where W¹ and W² are as depicted below:



17



where Z is mannose and/or a mannose receptor-targeting moiety (including, but not limited to, one or more of mannosamine, N-acetylmannosamine, or N-acetylglucosamine), R^9 is a direct bond, optionally substituted $\text{---C(O)---NH---(CH}_2\text{)}_2\text{---}$ (an ethylacetamido group or “EtAcN”) or optionally substituted $\text{---C(O)---NH---(CH}_2\text{)}_2\text{---(O---CH}_2\text{---CH}_2\text{)}_t\text{---}$ (a pegylated ethylacetamido group or “Et-PEG_t-AcN”), t is an integer from 1 to 5. In some embodiments, t is an integer of equal to or at least about: 1, 2, 3, 4, 5, 10, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, R^{10} is an aliphatic group, an alcohol, an aliphatic amine-containing group, or an aliphatic alcohol. In some embodiments, R^9 or R^{10} are independently optionally substituted alkyl, an optionally substituted polyether, or optionally substituted polyamino. In some embodiments, R^{10} is an optionally substituted C_{alkyl}, optionally substituted C_{alkyl}OH_g, or an optionally substituted $\text{---(C}_{f\text{alkyl}}\text{OH}_g\text{)}_e\text{---H}$ where f represents the number of carbons in the alkyl group and is an integer between 0 and 10, g represents the number of hydroxyl groups present on the alkyl group and is an integer between 0 and 10, and e represents the number of alkyl/ether repeat units and is an integer between 0 and 10. In some embodiments, e, f, and g are independently selected integers of equal to or at least about: 0, 1, 2, 3, 4, 5, 10, or ranges including and/or spanning the aforementioned values. In some embodiments, R^{10} is a 2-hydroxyethyl (e.g., $\text{---CH}_2\text{CH}_2\text{OH}$). In some embodiments, R^{10} is an optionally substituted 2-hydroxyethyl. In some embodiments, R^{10} is an optionally substituted polyether.





In some embodiments, Y' is represented as $\text{---W}^1_p\text{---W}^2_r\text{---}$. As noted elsewhere herein, $\text{---W}^1_p\text{---W}^2_r\text{---}$ may represent a block copolymer or a random copolymer of W^1 and W^2 monomers having p repeat units of W^1 and r repeat units of W^2 . In some embodiments, p is an integer equal to or greater than about: 1, 50, 85, 100, 150, 165, 200, 225, 250, 300, 400, or ranges including and/or spanning the aforementioned values. In some embodiments, r is an integer equal to or greater than about: 1, 50, 85, 100, 150, 165, 200, 225, 250, 300, 400, or ranges including and/or spanning the aforementioned values. In some embodiments, Y' is a homopolymer of W^1 or W^2 . In some embodiments, r is 0. In some embodiments, the sum of p and r is an integer equal to or greater than about: 1, 50, 85, 100, 150, 165, 170, 200, 225, 250, 300, 400, 600, 800, or ranges including and/or spanning the aforementioned values.

In some embodiments, polymeric chain^a and polymeric chain^b are present or optionally not present. In some embodiments, where present, polymeric chain^a and polymeric chain^b can independently comprise hydrophilic polymers. In some embodiments, where present, polymeric chain^a and polymeric chain^b can independently comprise one or more optionally substituted $\text{---(CH}_2\text{CH}_2\text{O)}_s\text{---}$, optionally substituted $\text{---(CH}_2\text{)}_u\text{---}$, or optionally substituted alkylene. In several embodiments, u is an integer less than or equal to about: 1, 5, 10, 20, or ranges including and/or spanning the


18

aforementioned values. In some embodiments, polymeric chain^a and polymeric chain^b comprise or consist of one or more of the following structures, or a portion thereof:


5




25




30




35




40




45



50



55




wherein the variables (e.g., i, k, n, q, v, etc.) are as disclosed elsewhere herein. In several embodiments, for example, n is an integer from about 1 to about 100, q is an integer from about 1 to about 100, k is an integer from about 1 to about 20, i is an integer from about 0 to about 20, and v is an integer from about 1 to about 20. In several embodiments, n or q represents the number of repeat units in a PEG chain. In some embodiments, the PEG chain may have some polydispersity. In some embodiments, n and q do not indicate a number of repeat units but instead independently indicate the presence of a PEG polymer chain having a Mn (in g/mol) or Mw (in g/mol) of equal to or at least

19


about 500, 1000, 2000, 5000, 10000, or ranges including and/or spanning the aforementioned values. In some embodiments, k, i, and v can each independently comprise an optionally substituted alkylene.

In several embodiments, n is an integer greater than or equal to about: 1, 10, 20, 40, 50, 75, 100, 150 or ranges including and/or spanning the aforementioned values. In several embodiments, n is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, or ranges including and/or spanning the aforementioned values. In several embodiments, q is an integer greater than or equal to about: 1, 10, 20, 40, 50, 75, 100, 150 or ranges including and/or spanning the aforementioned values. In several embodiments, q is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 75, 100, 150, or ranges including and/or spanning the aforementioned values. In several embodiments, k is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, v is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, k is 2. In several embodiments, v is 2. In several embodiments, n is 4. In several embodiments, n is 44. In several embodiments, q is 3. As used herein, variables disclosed as having structure, a value, or a range of values for one embodiment, may also have those values when the variable is used in another embodiment (even where the variable is not defined with respect to that other embodiment).

In several embodiments, the “CLICK” group and/or $-\text{[Y (-Z)}_p\text{]-}$, more generally, comprises the following functional unit:




In several embodiments, the “CLICK” group and/or $-\text{[Y (-Z)}_p\text{]-}$, more generally, comprises one or more of the following units (each of which may be optionally substituted):




20

-continued

5




10



20


25



30


35

40



45


50




55

60

65




In several embodiments, $-\text{[Y}(-\text{Z})\text{p]}-$ comprises the one or more of the following functional units:



wherein each variable (e.g., i, k, n, q, v, CLICK, R^1 , Y' , etc.) is as disclosed elsewhere herein. In some embodiments, for example, n is an integer from about 1 to about 44, q is an integer from about 1 to about 44, k is an integer from about 1 to about 12, i is an integer from about 0 to about 20, v is an integer from about 1 to


about 4, and R_1 is $-\text{CH}_2-$, $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)(\text{CN})-$, $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)(\text{CH}_3)-$, $-(\text{CH}_2)_2-\text{CH}(\text{CH}_3)-$, $-\text{CH}(\text{CH}_3)-$, or is absent.

In several embodiments, $-\text{[Y}(\text{Z})_p]-$ is a group represented by any one or more of Formula Ya' to Yr':





Ya'

Yb'



Yc'

Yd'





23

24

-continued



Ye'

Ye''





Yf'

Yf''




Yg'


Yh'




Yi'




Yj'




Yk'



Yl'




Ym'




25

26


-continued




Yn'




Yo'



Yp'



Yq'



Yr'

wherein the variables (e.g., i, k, n, q, v, R¹, Y', etc.) are as disclosed elsewhere herein. For example, in several embodiments, n is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, or ranges including and/or spanning the aforementioned values. In several embodiments, q is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, 30, 40, 50, 75, 100, 150, or ranges including and/or spanning the aforementioned values. In several embodiments, k is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, v is an integer greater than or equal to about: 1, 2, 3, 4, 5, 10, 15, 20, or ranges including and/or spanning the aforementioned values. In several embodiments, k is 2. In several embodiments, v is 2. In several embodiments, n is 4. In several embodiments, n is 43 or 44. In several embodiments, q is 3. In several embodiments, R₁ is —CH₂—, —(CH₂)₂—C(CH₃)(CN)—, —(CH₂)₂—C(CH₃)(CH₃)—, —(CH₂)₂—CH(CH₃)— or —CH

(CH₃)—. In some embodiments, Y' is a random copolymer or block copolymer of W¹ and W² having p repeat units of W¹ and r repeat units of W².

In several embodiments, as shown elsewhere herein, the targeting portion comprises one or more pendant liver targeting moieties decorating a portion of the linker. In several embodiments, the portion of the linker is a polymeric chain with pendant targeting agents attached randomly or in blocks along the chain. In some embodiments, the polymeric chain comprises an acrylate portion (e.g., acrylate polymers and/or acrylate copolymers). In several embodiments, the acrylate portion comprises an acrylate unit comprising a pendant liver targeting agent. In several embodiments, the acrylate portion further comprises an acrylate unit not comprising a pendant liver targeting agent.

In some embodiments, Y is a linker resulting from one or more reactions involving at least one of the following: N-hydroxysuccinamidyl (NHS) linker, NHS ester linker, PEG linker, maleimide linker, vinylsulfone linker, pyridyl di-thiol-poly(ethylene glycol) linker, pyridyl di-thiol linker,







27

n-nitrophenyl carbonate linker, or a nitrophenoxy poly(ethylene glycol)ester linker. The linker may have one or more mannose and/or a mannose receptor-targeting moieties (including, but not limited to, one or more of mannosamine, N-acetylmannosamine, or N-acetylglucosamine) bound to it. In embodiments, Y comprises an antibody, an antibody fragment, a peptide, or a disulfanyl ethyl ester to which one

28


or more mannose and/or mannose receptor-targeting moieties (including, but not limited to, one or more of mannosamine, N-acetylmannosamine, or N-acetylglucosamine) are bound.

In some embodiments, $-[Y(-Z)_p]-$ comprises one of the following structures:




-continued


Yk1




YL1




Ym1



Yo1



Yp1



where the variables are as disclosed elsewhere herein.

In some embodiments, other linker structures can be found in U.S. Application Publication Nos. U.S. 2017/0007708A1 and 2016/0243248A1 and International Publication No. WO 2017/046652, each of which is incorporated by reference in its entirety.

In several embodiments, various ratios of W^1 to W^2 are used (e.g., W^1 and W^2 as provided in any of the Formulae disclosed elsewhere herein). In some embodiments, a majority of Y^1 repeat units comprise W^1 . In some embodiments, the ratio of W^1 to W^2 is equal to or greater than about 50:1, about 25:1, about 10:1, about 5:1, about 4:1, about 2:1, about 1:1, about 1:2, about 1:4, about 1:5, about 1:10, about 1:25, about 1:50, and any ratio in between those listed, including

endpoints. In some embodiments, the ratio of p to r is equal to or greater than about 50:1, about 25:1, about 10:1, about 5:1, about 4:1, about 2:1, about 1:1, about 1:2, about 1:4, about 1:5, about 1:10, about 1:25, about 1:50, and any ratio in between those listed, including endpoints. In some embodiments, a homopolymer of W^1 is provided without a W^2 portion.

X may be a foreign transplant antigen, or alloantigen, or autoimmune antigen. In some aspects, X represents an antigen against which a patient may develop or has developed an unwanted immune response. In some embodiments, X is an antigen against which a subject, such as a transplant recipient or autoimmune patient, develops an unwanted immune response. In several embodiments, X is a foreign

31

extracellular vesicle, cell fragment, or cell containing alloantigens against which transplant recipients or autoimmune patients develop and unwanted immune response. In still further embodiments, X is a foreign food, animal, plant or environmental antigen against which patients develop an unwanted immune response. In certain aspects, X is a foreign therapeutic agent against which patients develop an unwanted immune response. In a further aspect, X is a synthetic self-antigen to which patients develop an unwanted immune response. In several embodiments, X is a tolerogenic portion of a larger antigen. In certain embodiments, X, or a portion of X, is, is at least, or is at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, or 500 amino acids in length (or any range derivable therein). In some embodiments, X is an asparaginase antigen or an ovalbumin antigen. In several embodiments, X is an immunogenic fragment of one or more of myelin basic protein, myelin oligodendrocyte glycoprotein, proteolipid protein, insulin, proinsulin, preproinsulin, high molecular weight glutenin, low molecular weight glutenin, alpha- or gamma-gliadin, hordein, secalin, or avenin.

In some embodiments, Z, the moiety that specifically targets a mannose receptor, is selected from the group consisting of α -linked mannose, β -linked mannose, substituted mannose, mannose-6-phosphate, N-acetyl mannosamine, and mannan having β (1-4), α (1-6), α (1-2), and/or α (1-3) linkages. In some aspects, the mannose receptor is the mannose-6-phosphate receptor.


32

Some aspects of the disclosure are directed towards the use of a composition as disclosed herein in any method disclosed herein. Some embodiment provide for the use of any composition disclosed herein for the induction of tolerance to the antigen or a tolerogenic portion thereof. It is specifically contemplated that any step or element of an embodiment may be implemented in the context of any other step(s) or element(s) of a different embodiment disclosed herein.


In some embodiments, a method of inducing immunological tolerance to an antigen target is provided. The method comprises administering a composition of Formula 1:



where the variables are as disclosed elsewhere herein. In some embodiments, for example, X comprises an antigen or a tolerogenic portion thereof, Y comprises a linker moiety, Z comprises a moiety that specifically targets a mannose receptor, p is an integer from 2 to 250, m is an integer from 1 to 100, R^2 is any of functional groups I-III:



where Ar is a substituted or unsubstituted aromatic group, R^3 is any carbon-containing linear or heterocyclic moiety, and R^{11} is hydrogen or an alkyl group. In some embodiments, R^2 is one of the functional groups:



where R^3 is as defined above.


In some embodiments, Y is a linker as described elsewhere herein. In several embodiments, Y is a linker resulting

33

from one or more reactions involving at least one of the following N-hydroxysuccinamidyl (NHS) linker, NHS ester linker, PEG linker, maleimide linker, vinylsulfone linker, pyridyl di-thiol-poly(ethylene glycol) linker, pyridyl di-thiol linker, n-nitrophenyl carbonate linker, or a nitrophenoxy poly(ethylene glycol)ester linker. The linker may have one or more mannose moieties or mannose receptor-targeting

34


moieties bound to it. In embodiments, Y comprises an antibody, an antibody fragment, a peptide, or a disulfanyl ethyl ester to which one or more mannose moieties or mannose receptor-targeting moieties are bound. In several embodiments, $-\text{[Y}(-\text{Z})_p\text{]}-$ is as disclosed elsewhere herein. For example, in some aspects, $-\text{[Y}(\text{Z})_p\text{]}-$ is a group represented by one of sequence Formula Ya to Yr:




35

36


-continued




Ym




Yn




Yo



Yp



Yq




Yr

and

37

where the left, opening parentheses “(” signifies the location of the bond between X and Y, the right, closing parentheses “)” signifies the location of the bond between Y and R², n is an integer from 1 to 100, q is an integer from 1 to 44, k is an integer from 1 to 12, i is an integer from 0 to 20, v is an integer from 1 to 4, R₁ is —CH₂—, —(CH₂)₂—C(CH₃)(CN)—, —(CH₂)₂—C(CH₃)(CH₃)—, —(CH₂)₂—CH(CH₃)— or —CH(CH₃)—, W¹ and W² are as depicted below:



where Z is mannose or a mannose receptor-targeting moiety, R⁹ is a direct bond, —(CH₂)₂—NH—C(O)— (an ethylacetamido group or “EtAcN”) or —(CH₂)₂—(O—CH₂—CH₂)_t—NH—C(O)— (a pegylated ethylacetamido group or “Et-PEG_t-AcN”), t is an integer from 1 to 5, 1 to 3, or 1 or 2, p is an integer from 2 to 250, R¹⁰ is an aliphatic group, an alcohol, an aliphatic amine-containing group, or an aliphatic alcohol, and r is an integer from 0 to 250. In some embodiments, R¹⁰ is a C_{alkyl} or C_{alkyl}OH_g, where f represents the number of carbons in the alkyl group and is an integer between 0 and 10, and g represents the number of hydroxyl groups present on the alkyl group and is an integer between 0 and 10. In some embodiments, R¹⁰ is 2-hydroxyethyl. In some aspects —W¹_p—W²_q— represents a block copolymer or a random copolymer of W¹ and W² monomers.

X may be a foreign transplant antigen, or alloantigen, or autoimmune antigen. In some aspects, X represents an antigen against which a patient may develop or has developed an unwanted immune response. In some embodiments, X can be an antigen against which a subject, such as a transplant recipient or autoimmune patient, develops an unwanted immune response. In several embodiments, X can be a foreign extracellular vesicle, cell fragment, or cell containing alloantigens against which transplant recipients or autoimmune patients develop and unwanted immune response. In still further embodiments, X can be a foreign food, animal, plant or environmental antigen against which patients develop an unwanted immune response. In certain aspects X can be a foreign therapeutic agent against which patients develop an unwanted immune response. In a further aspect X can be a synthetic self-antigen to which patients develop an unwanted immune response. In several embodiments, X can be a tolerogenic portion of a larger antigen. In certain embodiments, X, or a portion of X, is at least, or is at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98,

38

99, 100, 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 251, 252, 253, 254, 255, 256, 257, 258, 259, 260, 261, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 285, 286, 287, 288, 289, 290, 291, 292, 293, 294, 295, 296, 297, 298, 299, 300, 301, 302, 303, 304, 305, 306, 307, 308, 309, 310, 311, 312, 313, 314, 315, 316, 317, 318, 319, 320, 321, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335, 336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350, 351, 352, 353, 354, 355, 356, 357, 358, 359, 360, 361, 362, 363, 364, 365, 366, 367, 368, 369, 370, 371, 372, 373, 374, 375, 376, 377, 378, 379, 380, 381, 382, 383, 384, 385, 386, 387, 388, 389, 390, 391, 392, 393, 394, 395, 396, 397, 398, 399, 400, 401, 402, 403, 404, 405, 406, 407, 408, 409, 410, 411, 412, 413, 414, 415, 416, 417, 418, 419, 420, 421, 422, 423, 424, 425, 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, or 500 amino acids in length (or any range derivable therein). In some embodiments, X is an asparaginase antigen or an ovalbumin antigen. In specific embodiments, X is an asparaginase antigen or an ovalbumin antigen, Z, the moiety that specifically targets a mannose receptor, may be selected from the group consisting of α-linked mannose, β-linked mannose, substituted mannose, mannose-6-phosphate, N-acetyl mannosamine, and mannan having β(1-4), α(1-6), α(1-2), and/or α(1-3) linkages.

In some aspects, a method of treating or preventing an unwanted immune response against an antigen is provided. The method comprises administering to a subject in need of suppression of an immune response to the antigen an effective amount of a composition comprising a compound of Formula 1. The composition can be administered for clearance of a circulating protein or peptide or antibody that specifically binds to antigen moiety X, which circulating protein or peptide or antibody is causatively involved in transplant rejection, immune response against a therapeutic agent, autoimmune disease, hypersensitivity and/or allergy. The composition can be administered in an amount effective to reduce a concentration of the antibodies that are causatively involved in transplant rejection, immune response against a therapeutic agent, autoimmune disease, hypersensitivity and/or allergy in blood of the patient by at least 50% w/w, as measured at a time between about 12 to about 48 hours after the administration. The composition can be administered for tolerization of a patient with respect to antigen moiety X.

Other embodiments are discussed throughout this application. Any embodiment discussed with respect to one aspect applies to other aspects as well and vice versa. Each embodiment described herein is understood to be embodiments that are applicable to all aspects. It is contemplated

that any embodiment discussed herein can be implemented with respect to any method or composition, and vice versa. Furthermore, compositions and kits can be used to achieve methods disclosed herein.

The term “comprising,” which is synonymous with “including,” “containing,” or “characterized by,” is inclusive or open-ended and does not exclude additional, unrecited elements or method steps. The phrase “consisting of” excludes any element, step, or ingredient not specified. The phrase “consisting essentially of” limits the scope of described subject matter to the specified materials or steps and those that do not materially affect its basic and novel characteristics. It is contemplated that embodiments described in the context of the term “comprising” may also be implemented in the context of the term “consisting of” or “consisting essentially of.”

The terms “effective amount” or “therapeutically effective amount” refer to that amount of a composition of the disclosure that is sufficient to effect treatment, as defined herein, when administered to a mammal in need of such treatment. This amount will vary depending upon the subject and disease condition being treated, the weight and age of the subject, the severity of the disease condition, the particular composition of the disclosure chosen, the dosing regimen to be followed, timing of administration, manner of administration and the like, all of which can readily be determined by one of ordinary skill in the art.

The “numerical values” and “ranges” provided for the various substituents are intended to encompass all integers within the recited range. For example, when defining n as an integer representing a mixture including from 1 to 100, where the mixture typically encompasses the integer specified as $n \pm 10\%$ (or for smaller integers from 1 to about 25, ± 3), it should be understood that n can be an integer from 1 to 100 (e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 25, 30, 34, 35, 37, 40, 41, 45, 50, 54, 55, 59, 60, 65, 70, 75, 80, 82, 83, 85, 88, 90, 95, 99, 100, or any between those listed). The terms “ $\pm 10\%$ ” or “ ± 3 ” should be understood to disclose and provide specific support for equivalent ranges wherever used.

The term “optional” or “optionally” means that the subsequently described event or circumstance may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not.

A peptide, protein, or fragment that specifically binds a particular target is referred to as a “ligand” for that target.

A “polypeptide” is a term that refers to a chain of amino acid residues, regardless of post-translational modification (e.g., phosphorylation or glycosylation) and/or complexation with additional polypeptides, and/or synthesis into multisubunit complexes with nucleic acids and/or carbohydrates, or other molecules. Proteoglycans therefore also are referred to herein as polypeptides. A long polypeptide (having over 50 amino acids) is referred to as a “protein.” A short polypeptide (having 50 amino acids or fewer) is referred to as a “peptide.” Depending upon size, amino acid composition and three dimensional structure, certain polypeptides can be referred to as an “antigen-binding molecule,” “antibody,” an “antibody fragment” or a “ligand.” Polypeptides can be produced by a number of methods, many of which are well known in the art. For example, polypeptides can be obtained by extraction (e.g., from isolated cells), by expression of a recombinant nucleic acid encoding the polypeptide, or by chemical synthesis. Polypeptides can be produced by, for example, recombinant technology, and expression vec-

tors encoding the polypeptide introduced into host cells (e.g., by transformation or transfection) for expression of the encoded polypeptide.

As used herein, “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents and the like. In several embodiments, these media and agents can be used in combination with pharmaceutically active substances. Except insofar as any conventional media or agent is incompatible with the active ingredient, its use in the therapeutic compositions is contemplated. Supplementary active ingredients can also be incorporated into the compositions.

The term “purified” as used herein with reference to a polypeptide refers to a polypeptide that has been chemically or biologically synthesized and is thus substantially uncontaminated by other polypeptides, or has been separated or isolated from most other cellular components by which it is naturally accompanied (e.g., other cellular proteins, nucleic acids, or cellular components such as lipid membrane). An example of a purified polypeptide is one that is at least 70%, by dry weight, free from the proteins and naturally occurring organic molecules with which it naturally associates. A preparation of a purified polypeptide therefore can be, for example, at least 80%, at least 90%, or at least 99%, by dry weight, the polypeptide. Polypeptides also can be engineered to contain a tag sequence (e.g., a polyhistidine tag, a myc tag, a FLAG® tag, a SNAP® tag, or other affinity tag) that facilitates purification or marking (e.g., capture onto an affinity matrix, visualization under a microscope). Thus a purified composition that comprises a polypeptide refers to a purified polypeptide unless otherwise indicated. The term “isolated” indicates that the polypeptides or nucleic acids of the disclosure are not in their natural environment. Isolated products of the disclosure can thus be contained in a culture supernatant, partially enriched, produced from heterologous sources, cloned in a vector or formulated with a vehicle, etc.

The term “sequence identity” is used with regard to polypeptide or polynucleotide sequence comparisons. This expression in particular refers to a percentage of sequence identity, for example at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% to the respective reference polypeptide or to the respective reference polynucleotide. Particularly, the polypeptide in question and the reference polypeptide exhibit the indicated sequence identity over a continuous stretch of 20, 30, 40, 45, 50, 60, 70, 80, 90, 100 or more amino acids or over the entire length of the reference polypeptide. In several embodiments, despite differing sequences from a reference nucleotide (or corresponding polypeptide) a polynucleotide exhibits at least some degree of functional equivalence to the reference sequence, and in some embodiments, enhanced function.

The term “treatment” or “treating” means any treatment of a disease or disorder in a mammal, including: preventing or protecting against the disease or disorder, that is, causing the clinical symptoms not to develop; inhibiting the disease or disorder, that is, arresting or suppressing the development of clinical symptoms; and/or relieving the disease or disorder, that is, causing the regression of clinical symptoms.

The term “unwanted immune response” refers to a reaction by the immune system of a subject, which in the given situation is not desirable. The reaction of the immune system is unwanted if such reaction does not lead to the prevention,

reduction, or healing of a disease or disorder but instead causes, enhances or worsens a disorder or disease. Typically, a reaction of the immune system causes, enhances or worsens a disease if it is directed against an inappropriate target. Exemplified, an unwanted immune response includes but is not limited to transplant rejection, immune response against a therapeutic agent, autoimmune disease, and allergy or hypersensitivity.

The term “operatively linked” refers to a situation where two components are combined to form the active complex prior to binding at the target site. For example, a molecule conjugated to one-half of a biotin-streptavidin complex and an antigen complexed to the other one-half of the biotin-streptavidin complex are operatively linked through complexation of the biotin and streptavidin molecules. The term operatively linked is also intended to refer to covalent or chemical linkages that conjugate two molecules together.

Throughout this application, the term “about” is used according to its plain and ordinary meaning in the area of protein chemistry to indicate that a value includes the standard deviation of error for the device or method being employed to determine the value.

As used in this specification and claim(s), the words “comprising” (and any form of comprising, such as “comprise” and “comprises”), “having” (and any form of having, such as “have” and “has”), “including” (and any form of including, such as “includes” and “include”) or “containing” (and any form of containing, such as “contains” and “contain”) are inclusive or open-ended and do not exclude additional, unrecited elements or method steps.

Methods may involve multiple administrations of one or more compounds, compositions, and/or agents. In certain embodiments, cells or a subject are provided with a tolerance inducing agent prior to administering the composition for which a tolerance is being induced. It is contemplated that compounds, compositions, and/or agents may be formulated in a pharmaceutically acceptable formulation in certain embodiments of the invention.

A “disease” is defined as a pathological condition of a body part, an organ, or a system resulting from any cause, such as infection, genetic defect, or environmental stress. A “health-related condition” is defined herein to refer to a condition of a body part, an organ, or a system that may not be pathological, but for which treatment is sought. Examples include conditions for which cosmetic therapy is sought, such as skin wrinkling, skin blemishes, and the like. The disease can be any disease, and non-limiting examples include hyperproliferative diseases such as cancer and premalignant lesions, wounds, and infections.

A subject may be “predicted to have” a disease if the subject exhibits a characteristic, condition, or behavior that increases the likelihood of getting a disease. The characteristic, condition, or behavior that increases the likelihood of getting a disease is known as a risk factor, and may be behavioral, physiological, demographic, environmental, or genetic in nature. Behavioral risk factors usually relate to actions that a subject has chosen to take. Demographic risk factors are those that relate to the overall population, such as age or gender. Environmental risk factors include those that are related to exposure to objects in an environment, such as air pollution and access to clean water. Genetic risk factors are based on an individual’s genetic makeup, and may reflect interaction between the genes of the individual and environmental factors. A subject having a high number of manifestations of risk factors associated with a disease may be at increased risk of developing the disease, and may be “predicted to have” the disease.

“Prevention” and “preventing” are used according to their ordinary and plain meaning to mean “acting before” or such an act. In the context of a particular disease or health-related condition, those terms refer to administration or application of an agent, drug, or remedy to a subject or performance of a procedure or modality on a subject for the purpose of blocking the onset of a disease or health-related condition.


As used herein, an “antigen” is any substance that serves as a target for the receptors of an adaptive immune response, such as the T cell receptor, major histocompatibility complex class I and II, B cell receptor or an antibody. In some embodiments, an antigen may originate from within the body (e.g., “self,” “auto” or “endogenous”). In additional embodiments, an antigen may originate from outside the body (“non-self,” “foreign” or “exogenous”), having entered, for example, by inhalation, ingestion, injection, or transplantation, transdermal, etc. In some embodiments, an exogenous antigen may be biochemically modified in the body. Foreign antigens include, but are not limited to, food antigens, animal antigens, plant antigens, environmental antigens, therapeutic agents, as well as antigens present in an allograft transplant. Non-limiting examples of antigens are provided herein.

An “antigen-binding molecule” as used herein relates to molecules, in particular to proteins such as immunoglobulin molecules, which contain antibody variable regions providing a binding (specific binding in some embodiments) to an epitope. The antibody variable region can be present in, for example, a complete antibody, an antibody fragment, and a recombinant derivative of an antibody or antibody fragment. The term “antigen-binding fragment” of an antibody (or “binding portion”), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind a target sequence. Antigen-binding fragments containing antibody variable regions include (without limitation) “Fv”, “Fab”, and “F(ab’)₂” regions, “single domain antibodies (sdAb)”, “nanobodies”, “single chain Fv (scFv)” fragments, “tandem scFvs” (V_HA-V_LA-V_HB-V_LB), “diabodies”, “tribodies” or “tribodies”, “single-chain diabodies (scDb)”, and “bi-specific T-cell engagers (BiTEs)”.

An “epitope”, also known as antigenic determinant, is the segment of a macromolecule, e.g. a protein, which is recognized by the adaptive immune system, such as by antibodies, B cells, major histocompatibility complex molecules, or T cells. An epitope is that part or segment of a macromolecule capable of binding to an antibody or antigen-binding fragment thereof. In this context, the term “binding” in particular relates to a specific binding. In the context of several embodiments of the present invention it is preferred that the term “epitope” refers to the segment of protein or polypeptide that is recognized by the immune system.

The term mannose or mannosylating moiety refers to a monosaccharide sugar that exists both in open-chain form and in cyclic form, having D- and L-isomers. In the cyclic form, there are two anomers, namely alpha and beta. In the alpha form, the C1 alcohol group is in the axial position, whereas in the beta form, the C1 alcohol group is in the equatorial position. In particular, “mannose” refers to the cyclic six-membered pyranose, more in particular the D-isomer and even more particularly the alpha-D-form (α-D-mannose). The structure and numbering of mannose on non-limiting examples of stereochemical illustration. In the current formulation, the mannose residue is a single sugar that is connected to the backbone of the polymer via a single site such as the primary alcohols that attach to C1-C4 and C6. Thus, one of the alcohols is used to connect the mannose

to the polymer while the other alcohols remain OH groups at neutral pH. In several embodiments, the advantageous polyfunctionality of the approaches disclosed herein comes, at least in part, from the multiple monomers that are used to decorate the side of the polymer.



Whenever a group is described as being “optionally substituted” that group may be unsubstituted or substituted with one or more of the indicated substituents. Likewise, when a group is described as being “unsubstituted or substituted” if substituted, the substituent(s) may be selected from one or more the indicated substituents. If no substituents are indicated, it is meant that the indicated “optionally substituted” or “substituted” group may be substituted with one or more group(s) individually and independently selected from C₁₋₆ alkyl, C₁₋₆ alkenyl, C₁₋₆ alkynyl, C₁₋₆ cycloalkyl, C₁₋₆ aryl, C₁₋₆ heteroaryl, C₁₋₆ heterocyclyl, C₁₋₆ alkoxy, C₁₋₆ acyl, cyano, hydroxyl, an amino, halogen substituted C₁₋₆alkyl, halogen substituted C₁₋₆alkoxy, and halogen.

As used herein, a “chemical modification” refers to a change in the naturally-occurring chemical structure of one or more amino acids of a polypeptide. Such modifications can be made to a side chain or a terminus, e.g., changing the amino-terminus or carboxyl terminus. In some embodiments, the modifications are useful for creating chemical groups that can conveniently be used to link the polypeptides to other materials, or to attach a therapeutic agent.

“Conservative changes” can generally be made to an amino acid sequence without altering activity. These changes are termed “conservative substitutions” or mutations; that is, an amino acid belonging to a grouping of amino acids having a particular size or characteristic can be substituted for another amino acid. Substitutes for an amino acid sequence can be selected from other members of the class to which the amino acid belongs. For example, the nonpolar (hydrophobic) amino acids include alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, methionine, and tyrosine. The polar neutral amino acids include glycine, serine, threonine, cysteine, tyrosine, asparagine and glutamine. The positively charged (basic) amino acids include arginine, lysine and histidine. The negatively charged (acidic) amino acids include aspartic acid and glutamic acid. Such substitutions are not expected to substantially affect apparent molecular weight as determined by polyacrylamide gel electrophoresis or isoelectric point. Conservative substitutions also include substituting optical iso-

mers of the sequences for other optical isomers, specifically D amino acids for L amino acids for one or more residues of a sequence. Moreover, all of the amino acids in a sequence can undergo a D to L isomer substitution. Exemplary conservative substitutions include, but are not limited to, Lys for Arg and vice versa to maintain a positive charge; Glu for Asp and vice versa to maintain a negative charge; Ser for Thr so that a free —OH is maintained; and Gln for Asn to maintain a free —NH₂. Yet another type of conservative substitution constitutes the case where amino acids with desired chemical relativities are introduced to impart reactive sites for chemical conjugation reactions, if the need for chemical derivatization arises. Such amino acids include but are not limited to Cys (to insert a sulfhydryl group), Lys (to insert a primary amine), Asp and Glu (to insert a carboxylic acid group), or specialized noncanonical amino acids containing ketone, azide, alkyne, alkene, and tetrazine side-chains. Conservative substitutions or additions of free —NH₂ or —SH bearing amino acids can be particularly advantageous for chemical conjugation with the linkers and mannosylating moieties of Formula 1. Moreover, point mutations, deletions, and insertions of the polypeptide sequences or corresponding nucleic acid sequences can in some cases be made without a loss of function of the polypeptide or nucleic acid fragment. Substitutions can include, e.g., 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50 or more residues (including any number of substitutions between those listed). A variant usable in embodiments herein may exhibit a total number of up to 200 (e.g., up to 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, or 200, including any number in between those listed) changes in the amino acid sequence (e.g., exchanges, insertions, deletions, N-terminal truncations, and/or C-terminal truncations). In several embodiments, the number of changes is greater than 200. Additionally, in several embodiments, the variants include polypeptide sequences or corresponding nucleic acid sequences that exhibit a degree of functional equivalence with a reference (e.g., unmodified or native sequence). In several embodiments, the variants exhibit about 80%, about 85%, about 90%, about 95%, about 97%, about 98%, about 99% functional equivalence to an unmodified or native reference sequence (and any degree of functional equivalence between those listed). The amino acid residues described herein employ either the single letter amino acid designator or the three-letter abbreviation in keeping with the standard polypeptide nomenclature, *J. Biol. Chem.*, (1969), 243, 3552-3559. All amino acid residue sequences are represented herein by formulae with left and right orientation in the conventional direction of amino-terminus to carboxy-terminus.

The term “liver-targeting moiety” refers to mannose moieties having the ability to direct, e.g., a polypeptide, to the cells of a liver expressing mannose receptors. The liver comprises different cell types, including but not limited to hepatocytes, sinusoidal epithelial cells, Kupffer cells, stellate cells, and/or dendritic cells. Typically, a liver-targeting moiety directs a polypeptide to one or more of these cells. On the surface of the respective liver cells, receptors are present which recognize and specifically bind the liver-targeting moiety. Liver-targeting can be achieved by chemical conjugation of an antigen or ligand to a mannosylating or mannansylating moiety. Naturally occurring desilylated proteins are not encompassed within the scope of certain embodiments of the present disclosure.

The term “random copolymer” refers to the product of simultaneous polymerization of two or more monomers in

admixture, where the probability of finding a given monomeric unit at any given site in a polymer chain is independent of the nature of the neighboring units at that position (Bernoullian distribution). Thus, when the variable group identified as Wp represents a random copolymer, the chain can comprise any sequence from 2 up to about 150 W1 and W2 groups, such as: —W1-W2-W1-W2-; —W2-W1-W2-W1-; —W1-W1-W1-W2-; —W1-W1-W2-W2-; —W1-W2-W2-W1-; —W1-W2-W1-W2-W2-W1-W2-W1-; and W2-W2-W1-W2-W1-W1-W1-W2-W2-W1-W2-W2-W1; ad infinitum, where Z attached to the various W1 groups and the W1 and W2 groups themselves can be the same or different.

The term “sequence identity” is used with regard to polypeptide (or nucleic acid) sequence comparisons. This expression in particular refers to a percentage of sequence identity, for example at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% to the respective reference polypeptide or to the respective reference polynucleotide. Particularly, the polypeptide in question and the reference polypeptide exhibit the indicated sequence identity over a continuous stretch of 20, 30, 40, 45, 50, 60, 70, 80, 90, 100 or more amino acids (or any range derivable therein) or over the entire length of the reference polypeptide.

“Specific binding,” as that term is commonly used in the biological arts, refers to a molecule that binds to a target with a relatively high affinity as compared to non-target tissues, and generally involves a plurality of non-covalent interactions, such as electrostatic interactions, van der Waals interactions, hydrogen bonding, and the like. Specific binding interactions characterize antibody-antigen binding, enzyme-substrate binding, and certain protein-receptor interactions; while such molecules might bind tissues besides their specific targets from time to time, to the extent that such non-target binding is inconsequential, the high-affinity binding pair can still fall within the definition of specific binding.

The term “unwanted immune response” refers to a reaction by the immune system of a subject, which in the given situation is not desirable. The reaction of the immune system is unwanted if such reaction does not lead to the prevention, reduction, or healing of a disease or disorder but instead causes, enhances or worsens, or is otherwise associated with induction or worsening of a disorder or disease. Typically, a reaction of the immune system causes, enhances or worsens a disease if it is directed against an inappropriate target. Exemplified, an unwanted immune response includes but is not limited to transplant rejection, immune response against a therapeutic agent, autoimmune disease, and allergy or hypersensitivity.

The term “variant” is to be understood as a protein (or nucleic acid) which differs in comparison to the protein from which it is derived by one or more changes in its length, sequence, or structure. The polypeptide from which a protein variant is derived is also known as the parent polypeptide or polynucleotide. The term “variant” comprises “fragments” or “derivatives” of the parent molecule. Typically, “fragments” are smaller in length or size than the parent molecule, whilst “derivatives” exhibit one or more differences in their sequence or structure in comparison to the parent molecule. Also encompassed are modified molecules such as but not limited to post-translationally modified proteins (e.g. glycosylated, phosphorylated, ubiquitinated, palmitoylated, or proteolytically cleaved proteins) and

modified nucleic acids such as methylated DNA. Also mixtures of different molecules such as but not limited to RNA-DNA hybrids, are encompassed by the term “variant”. Naturally occurring and artificially constructed variants are to be understood to be encompassed by the term “variant” as used herein. Further, the variants usable in the present invention may also be derived from homologs, orthologs, or paralogs of the parent molecule or from artificially constructed variant, provided that the variant exhibits at least one biological activity of the parent molecule, e.g., is functionally active. A variant can be characterized by a certain degree of sequence identity to the parent polypeptide from which it is derived. More precisely, a protein variant in the context of the present disclosure may exhibit at least 80% sequence identity to its parent polypeptide. Preferably, the sequence identity of protein variants is over a continuous stretch of 20, 30, 40, 45, 50, 60, 70, 80, 90, 100 or more amino acids. As discussed above, in several embodiments variants exhibit about 80%, about 85%, about 90%, about 95%, about 97%, about 98%, about 99% functional equivalence to an unmodified or native reference sequence (and any degree of functional equivalence between those listed).

It is specifically contemplated that any limitation discussed with respect to one embodiment of the invention may apply to any other embodiment of the invention. Furthermore, any composition of the invention may be used in any method of the invention, and any method of the invention may be used to produce or to utilize any composition of the invention.

The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternative are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.”

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE FIGURES

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIGS. 1A-1G. OVA-p(Man) induces T cell deletion. BLK6 mice were treated with saline, or 10 µg of OVA in the form of free OVA or OVA conjugated to p(Man) (OVA-p(Man)) one day and 7 days after an adoptive transfer of 7.0×10^5 OTI and OTII T cells. The mice were challenged with an intradermal injection of LSP and OVA on 14 days after the initial OTI and OTII T cell transfer, then the immune response in the draining lymph nodes (dLNs) was assessed on day 19 via flow cytometry. FIG. 1A Fraction of OTI T cells in the dLNs on day 19. FIG. 1B Fraction of OTI T cells in the dLNs on day 19. FIG. 1C Fraction of OTII T cells in the liver on day 19. FIG. 1D Fraction of OTI T cells in the liver on day 19. FIG. 1E Fraction of Tregs in the dLN on day 19. FIG. 1F Fraction of Tregs in the liver on day 19.

FIG. 1G Fraction of T follicular helper cells (Tfh) as a fraction of total CD4 T cells in the dLNs on day 19.

FIGS. 2A-2B: OVA-p(Man) induces T cells anergy in DLs. BLK6 mice were treated with saline, or 10 µg of OVA in the form of free OVA or OVA conjugated to p(Man) (OVA-p(Man)) one day and 7 days after an adoptive transfer of 7.0×10^5 OTI and OTII T cells. The mice were challenged with an intradermal injection of LPS and OVA on 14 days after the initial OTI and OTII T cell transfer, and immune response in the draining lymph nodes (dLNs) was assessed on day 19. Cells from the dLN were restimulated with OVA (A) or SIINFELK (peptide) for 6 hours, then the percentage of IFN-γ producing cells was determined by flow cytometry. FIG. 2A Percentage of IFN-γ producing CD4+ T cells in the dLNs. FIG. 2B Percentage of IFN-γ producing CD8+ T cells in the dLNs.

FIGS. 3A-3D. Tolerance induction to protein therapeutics. Five BALB/c mice per group were injected with 2.5 µg of asparaginase formulated as free asparaginase (ASNase) or conjugated to p(Man) (ASNase-p(Man)) once a week for 3 weeks and then were switched to 15 µg of ASNase i.v. one a week for 8 weeks. During the initial 3 weeks ASNase-p(Man) was administered via either i.v. or subcutaneous injection. Sera was taken from the mice and monitored weekly for the presence of αASNase. FIG. 3A Pan IgG αASNase titers of mice treated with ASNase and ASNase-p(Man) via i.v. and subcutaneous injection. FIG. 3B αASNase IgG1 titers of treatment groups on day 38. FIG. 3C αASNase IgG2a titers of treatment groups on day 38. FIG. 3D αASNase IgG2b titers of treatment groups on day 38. FIG. 3E. αASNase IgG3 titers of treatment groups on day 38.

FIGS. 4A-4B. Response to p(Man) conjugates. FIG. 4A Bone marrow of animals treated with ASNase-p(Man) had fewer αASNase plasma cells than animals that were treated with ASNase. FIG. 4B Spleens of animals treated with ASNase-p(Man) had a greater percentage of IL-10 producing B regulatory cells.

FIG. 5. Tolerance induction to protein therapeutics. FIG. 5 is a graph depicting tolerance induction to the protein therapeutic asparaginase conjugated to p(Man) linker and mannose. Five BALB/c mice per group were injected with 2.5 µg of asparaginase formulated as free asparaginase (ASNase) or conjugated to p(Man) (ASNase-p(Man)) once a week for 3 weeks and then were switched to 15 µg of ASNase i.v. one a week for 7 weeks. Sera was taken from the mice and monitored weekly for the presence of anti-ASNase. Over the entire time of analysis none of the animals treated with ASNase-p(Man) developed measureable levels of antibodies against ASNase.

FIG. 6 p(Man)-ASNase administration regimen for assessment of p(Man)-protein conjugates on anti-asparaginase (anti-ASNase) humoral immune response.

FIGS. 7A-7B. Assessment of p(Man)-protein conjugates on anti-asparaginase (anti-ASNase) humoral immune response. FIG. 7A is a graph depicting serum asparagine concentration for days 71, 73, and 76. Mice treated with only saline or p(Man)-ASNase have a significantly lower serum asparagine concentration as compared to animals that had been treated with saline and then administered wt ASNase. FIG. 7B is a graph depicting serum asparagine concentration vs the anti-ASNase titer for each animal in the study. A strong correlation ($r=0.8$) between serum asparagine concentration and anti-ASNase titer is evident.

DETAILED DESCRIPTION

Several embodiments disclosed herein overcome the deficiencies of the prior art by providing compositions compris-

ing mannose-fused antigens. The compositions may be used to prevent immunity or reduce an immune response protein-based drugs that would otherwise elicit an immune response.

Multiple mannose binding receptors are expressed by antigen presenting cells (APCs) and serve as gateways for antigen uptake and antigen cross presentation by these cells to T cells. Antigens taken up by APCs and presented to T cells in the absence of co-stimulation leads to T cell deletion, inactivity, and the formation of T regulator cells, that control antigen specific immune responses.

The present disclosure provides, in several embodiments, certain therapeutic compositions that are targeted for delivery to (and for uptake by) antigen presenting cells, particularly hepatocytes, LSECs, Kupffer cells and/or stellate cells, more particularly hepatocytes and/or LSECs, and even more particularly to specifically bind mannose-binding receptors.

Liver targeting facilitates two mechanisms of treatment: tolerization and clearance. Tolerization takes advantage of the liver's role in clearing apoptotic cells and processing their proteins to be recognized by the immune system as "self," as well as the liver's role in sampling peripheral proteins for immune tolerance. Clearance takes advantage of the liver's role in blood purification by rapidly removing and breaking down toxins, polypeptides and the like. Targeting of these compositions to the liver is accomplished by a mannosating moiety. The mannosylating moiety is chemically conjugated. The antigen can be endogenous (a self-antigen) or exogenous (a foreign antigen), including but not limited to: a foreign transplant antigen against which transplant recipients develop an unwanted immune response (e.g., transplant rejection), a foreign food, animal, plant or environmental antigen to which patients develop an unwanted immune (e.g., allergic or hypersensitivity) response, a therapeutic agent to which patients develop an unwanted immune response (e.g., hypersensitivity and/or reduced therapeutic activity), a self-antigen to which patients develop an unwanted immune response (e.g., autoimmune disease), or a tolerogenic portion (e.g., a fragment or an epitope) thereof; these compositions are useful for inducing tolerization to the antigen. Accordingly, the compositions of the present disclosure can be used for treating an unwanted immune response, e.g., transplant rejection, an immune response against a therapeutic agent, an autoimmune disease, and/or an allergy, depending on the embodiment. Also provided are pharmaceutical compositions containing a therapeutically effective amount of a composition of the disclosure admixed with at least one pharmaceutically acceptable excipient. In another aspect, the disclosure provides methods for the treatment of an unwanted immune response, such as transplant rejection, response against a therapeutic agent, autoimmune disease or allergy.

B. CHEMICAL DEFINITIONS

As used herein, a "small molecule" refers to an organic compound that is either synthesized via conventional organic chemistry methods (e.g., in a laboratory) or found in nature. Typically, a small molecule is characterized in that it contains several carbon-carbon bonds, and has a molecular weight of less than about 1500 grams/mole. In certain embodiments, small molecules are less than about 1000 grams/mole. In certain embodiments, small molecules are less than about 550 grams/mole. In certain embodiments, small molecules are between about 200 and about 550 grams/mole. In certain embodiments, small molecules exclude peptides (e.g., compounds comprising 2 or more

amino acids joined by a peptidyl bond). In certain embodiments, small molecules exclude nucleic acids.


As used herein, the term “amino” means —NH_2 ; the term “nitro” means —NO_2 ; the term “halo” or “halogen” designates —F , —Cl , —Br or —I ; the term “mercapto” means —SH ; the term “cyano” means —CN ; the term “azido” means —N_3 ; the term “silyl” means —SiH_3 , and the term “hydroxy” means —OH . In certain embodiments, a halogen may be —Br or —I .

As used herein, a “monovalent anion” refers to anions of a -1 charge. Such anions are well-known to those of skill in the art. Non-limiting examples of monovalent anions include halides (e.g., F— , Cl— , Br— and I—), NO_2^- , NO_3^- , hydroxide (OH—) and azide (N_3^-).

As used herein, the structure indicates that the bond may be a single bond or a double bond. Those of skill in the chemical arts understand that in certain circumstances, a double bond between two particular atoms is chemically feasible and in certain circumstances, a double bond is not. The present invention therefore contemplates that a double bond may be formed only when chemically feasible.

The term “alkyl” includes straight-chain alkyl, branched-chain alkyl, cycloalkyl (alicyclic), cyclic alkyl, heteroatom-unsubstituted alkyl, heteroatom-substituted alkyl, heteroatom-unsubstituted C_n -alkyl, and heteroatom-substituted C_n -alkyl. In certain embodiments, lower alkyls are contemplated. Examples of branched alkyl groups include, but are not limited to, iso-propyl, sec-butyl, t-butyl and the like. Examples of straight chain alkyl groups include, but are not limited to, methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl and the like. The alkyl group may have 1 to 30 carbon atoms (whenever it appears herein, a numerical range such as “1 to 30” refers to each integer in the given range; e.g., “1 to 30 carbon atoms” means that the alkyl group may consist of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The “alkyl” group may also be a medium size alkyl having 1 to 12 carbon atoms. The term “lower alkyl” refers to alkyls of 1-6 carbon atoms (that is, 1, 2, 3, 4, 5 or 6 carbon atoms). The term “heteroatom-unsubstituted C_n -alkyl” refers to a radical, having a linear or branched, cyclic or acyclic structure, further having no carbon-carbon double or triple bonds, further having a total of n carbon atoms, all of which are nonaromatic, 3 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C_1 - C_{10} -alkyl has 1 to 10 carbon atoms. The groups, —CH_3 (Me), $\text{—CH}_2\text{CH}_3$ (Et), $\text{—CH}_2\text{CH}_2\text{CH}_3$ (n-Pr), $\text{—CH}(\text{CH}_3)_2$ (iso-Pr), $\text{—CH}(\text{CH}_2)_2$ (cyclopropyl), $\text{—CH}_2\text{CH}_2\text{CH}_2\text{CH}_3$ (n-Bu), $\text{—CH}(\text{CH}_3)\text{CH}_2\text{CH}_3$ (sec-butyl), $\text{—CH}_2\text{CH}(\text{CH}_3)_2$ (iso-butyl), $\text{—C}(\text{CH}_3)_3$ (tert-butyl), $\text{—CH}_2\text{C}(\text{CH}_3)_3$ (neo-pentyl), cyclobutyl, cyclopentyl, and cyclohexyl, are all non-limiting examples of heteroatom-unsubstituted alkyl groups. The term “heteroatom-substituted C_n -alkyl” refers to a radical, having a single saturated carbon atom as the point of attachment, no carbon-carbon double or triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C_1 - C_{10} -alkyl has 1 to 10 carbon atoms. The following groups are all non-limiting examples of heteroatom-substituted alkyl groups: trifluoromethyl, $\text{—CH}_2\text{F}$, $\text{—CH}_2\text{Cl}$, $\text{—CH}_2\text{Br}$,

$\text{—CH}_2\text{OH}$, $\text{—CH}_2\text{OCH}_3$, $\text{—CH}_2\text{OCH}_2\text{CF}_3$, $\text{—CH}_2\text{OC}(\text{O})\text{CH}_3$, $\text{—CH}_2\text{NH}_2$, $\text{—CH}_2\text{NHCH}_3$, $\text{—CH}_2\text{N}(\text{CH}_3)_2$, $\text{—CH}_2\text{CH}_2\text{Cl}$, $\text{—CH}_2\text{CH}_2\text{OH}$, $\text{—CH}_2\text{CH}_2\text{OC}(\text{O})\text{CH}_3$, $\text{—CH}_2\text{CH}_2\text{NHCO}_2\text{C}(\text{CH}_3)_3$, and $\text{—CH}_2\text{Si}(\text{CH}_3)_3$.

As used herein, the term “alkylene” refers to a bivalent fully saturated straight chain aliphatic hydrocarbon group. Examples of alkylene groups include, but are not limited to, methylene, ethylene, propylene, butylene, pentylene, hexylene, heptylene and octylene. An alkylene group may be represented by , followed by the number of carbon atoms, followed by a “*”. For example,

to represent ethylene. The alkylene group may have 1 to 30 carbon atoms (whenever it appears herein, a numerical range such as “1 to 30” refers to each integer in the given range; e.g., “1 to 30 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, etc., up to and including 30 carbon atoms, although the present definition also covers the occurrence of the term “alkylene” where no numerical range is designated). The alkylene group may also be a medium size alkyl having 1 to 12 carbon atoms. The alkylene group could also be a lower alkyl having 1 to 4 carbon atoms. An alkylene group may be substituted or unsubstituted. For example, a lower alkylene group can be substituted by replacing one or more hydrogen of the lower alkylene group and/or by substituting both hydrogens on the same carbon with a C3-6 monocyclic cycloalkyl group (e.g.,



The term “alkenyl” includes straight-chain alkenyl, branched-chain alkenyl, cycloalkenyl, cyclic alkenyl, heteroatom-unsubstituted alkenyl, heteroatom-substituted alkenyl, heteroatom-unsubstituted C_n -alkenyl, and heteroatom-substituted C_n -alkenyl. In certain embodiments, lower alkenyls are contemplated. The term “lower alkenyl” refers to alkenyls of 1-6 carbon atoms (that is, 1, 2, 3, 4, 5 or 6 carbon atoms). The term “heteroatom-unsubstituted C_n -alkenyl” refers to a radical, having a linear or branched, cyclic or acyclic structure, further having at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, a total of n carbon atoms, three or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C_2 - C_{10} -alkenyl has 2 to 10 carbon atoms. Heteroatom-unsubstituted alkenyl groups include: —CH=CH_2 (vinyl), —CH=CHCH_3 , $\text{—CH=CHCH}_2\text{CH}_3$, $\text{—CH}_2\text{CH=CH}_2$ (allyl), $\text{—CH}_2\text{CH=CHCH}_3$, and $\text{—CH=CH—C}_6\text{H}_5$. The term “heteroatom-substituted C_n -alkenyl” refers to a radical, having a single nonaromatic carbon atom as the point of attachment and at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one heteroatom, wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C_2 - C_{10} -alkenyl has 2 to 10 carbon

51

atoms. The groups, $-\text{CH}=\text{CHF}$, $-\text{CH}=\text{CHCl}$ and $-\text{CH}=\text{CHBr}$, are non-limiting examples of heteroatom-substituted alkenyl groups.

The term “aryl” includes heteroatom-unsubstituted aryl, heteroatom-substituted aryl, heteroatom-unsubstituted C_n-aryl, heteroatom-substituted C_n-aryl, heteroaryl, heterocyclic aryl groups, carbocyclic aryl groups, biaryl groups, and single-valent radicals derived from polycyclic fused hydrocarbons (PAHs). The term “heteroatom-unsubstituted C_n-aryl” refers to a radical, having a single carbon atom as a point of attachment, wherein the carbon atom is part of an aromatic ring structure containing only carbon atoms, further having a total of n carbon atoms, 5 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₆-C₁₀-aryl has 6 to 10 carbon atoms. Non-limiting examples of heteroatom-unsubstituted aryl groups include phenyl (Ph), methylphenyl, (dimethyl)phenyl, $-\text{C}_6\text{H}_4\text{CH}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CH}(\text{CH}_3)_2$, $-\text{C}_6\text{H}_4\text{CH}(\text{CH}_2)_2$, $-\text{C}_6\text{H}_3(\text{CH}_3)\text{CH}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CH}=\text{CH}_2$, $-\text{C}_6\text{H}_4\text{CH}=\text{CHCH}_3$, $-\text{C}_6\text{H}_4\text{C}\equiv\text{CH}$, $-\text{C}_6\text{H}_4\text{C}\equiv\text{CCH}_3$, naphthyl, and the radical derived from biphenyl. The term “heteroatom-substituted C_n-aryl” refers to a radical, having either a single aromatic carbon atom or a single aromatic heteroatom as the point of attachment, further having a total of n carbon atoms, at least one hydrogen atom, and at least one heteroatom, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-unsubstituted C₁-C₁₀-heteroaryl has 1 to 10 carbon atoms. Non-limiting examples of heteroatom-substituted aryl groups include the groups: $-\text{C}_6\text{H}_4\text{F}$, $-\text{C}_6\text{H}_4\text{Cl}$, $-\text{C}_6\text{H}_4\text{Br}$, $-\text{C}_6\text{H}_4\text{I}$, $-\text{C}_6\text{H}_4\text{OH}$, $-\text{C}_6\text{H}_4\text{OCH}_3$, $-\text{C}_6\text{H}_4\text{OCH}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{OC}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{NH}_2$, $-\text{C}_6\text{H}_4\text{NHCH}_3$, $-\text{C}_6\text{H}_4\text{N}(\text{CH}_3)_2$, $-\text{C}_6\text{H}_4\text{CH}_2\text{OH}$, $-\text{C}_6\text{H}_4\text{CH}_2\text{OC}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CH}_2\text{NH}_2$, $-\text{C}_6\text{H}_4\text{CF}_3$, $-\text{C}_6\text{H}_4\text{CN}$, $-\text{C}_6\text{H}_4\text{CHO}$, $-\text{C}_6\text{H}_4\text{CHO}$, $-\text{C}_6\text{H}_4\text{C}(\text{O})\text{CH}_3$, $-\text{C}_6\text{H}_4\text{C}(\text{O})\text{C}_6\text{H}_5$, $-\text{C}_6\text{H}_4\text{CO}_2\text{H}$, $-\text{C}_6\text{H}_4\text{CO}_2\text{CH}_3$, $-\text{C}_6\text{H}_4\text{CONH}_2$, $-\text{C}_6\text{H}_4\text{CONHCH}_3$, $-\text{C}_6\text{H}_4\text{CON}(\text{CH}_3)_2$, furanyl, thienyl, pyridyl, pyrrolyl, pyrimidyl, pyrazinyl, quinolyl, indolyl, and imidazolyl. In certain embodiments, heteroatom-substituted aryl groups are contemplated. In certain embodiments, heteroatom-unsubstituted aryl groups are contemplated. In certain embodiments, an aryl group may be mono-, di-, tri-, tetra- or penta-substituted with one or more heteroatom-containing substituents.

The term “aralkyl” includes heteroatom-unsubstituted aralkyl, heteroatom-substituted aralkyl, heteroatom-unsubstituted C_n-aralkyl, heteroatom-substituted C_n-aralkyl, heteroaralkyl, and heterocyclic aralkyl groups. In certain embodiments, lower aralkyls are contemplated. The term “lower aralkyl” refers to aralkyls of 7-12 carbon atoms (that is, 7, 8, 9, 10, 11 or 12 carbon atoms). The term “heteroatom-unsubstituted C_n-aralkyl” refers to a radical, having a single saturated carbon atom as the point of attachment, further having a total of n carbon atoms, wherein at least 6 of the carbon atoms form an aromatic ring structure containing only carbon atoms, 7 or more hydrogen atoms, and no heteroatoms. For example, a heteroatom-unsubstituted C₇-C₁₀-aralkyl has 7 to 10 carbon atoms. Non-limiting examples of heteroatom-unsubstituted aralkyls are: phenylmethyl (benzyl, Bn) and phenylethyl. The term “heteroatom-substituted C_n-aralkyl” refers to a radical, having a single saturated carbon atom as the point of attachment, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one heteroatom, wherein at least one of the carbon atoms is incorporated an aromatic ring

52

structures, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-heteroaralkyl has 2 to 10 carbon atoms.

The term “acyl” includes straight-chain acyl, branched-chain acyl, cycloacyl, cyclic acyl, heteroatom-unsubstituted acyl, heteroatom-substituted acyl, heteroatom-unsubstituted C_n-acyl, heteroatom-substituted C_n-acyl, alkylcarbonyl, alkoxy carbonyl and aminocarbonyl groups. In certain embodiments, lower acyls are contemplated. The term “lower acyl” refers to acyls of 1-6 carbon atoms (that is, 1, 2, 3, 4, 5 or 6 carbon atoms). The term “heteroatom-unsubstituted C_n-acyl” refers to a radical, having a single carbon atom of a carbonyl group as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 1 or more hydrogen atoms, a total of one oxygen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-acyl has 1 to 10 carbon atoms. The groups, $-\text{CHO}$, $-\text{C}(\text{O})\text{CH}_3$, $-\text{C}(\text{O})\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{C}(\text{O})\text{CH}(\text{CH}_3)_2$, $-\text{C}(\text{O})\text{CH}(\text{CH}_2)_2$, $-\text{C}(\text{O})\text{C}_6\text{H}_5$, $-\text{C}(\text{O})\text{C}_6\text{H}_4\text{CH}_3$, $-\text{C}(\text{O})\text{C}_6\text{H}_4\text{CH}_2\text{CH}_3$, and $-\text{COC}_6\text{H}_3(\text{CH}_3)_2$, are non-limiting examples of heteroatom-unsubstituted acyl groups. The term “heteroatom-substituted C_n-acyl” refers to a radical, having a single carbon atom as the point of attachment, the carbon atom being part of a carbonyl group, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom, in addition to the oxygen of the carbonyl group, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-acyl has 1 to 10 carbon atoms. The groups, $-\text{C}(\text{O})\text{CH}_2\text{CF}_3$, $-\text{CO}_2\text{H}$, $-\text{CO}_2-$, $-\text{CO}_2\text{CH}_3$, $-\text{CO}_2\text{CH}_2\text{CH}_3$, $-\text{CO}_2\text{CH}_2\text{CH}_2\text{CH}_3$, $-\text{CO}_2\text{CH}(\text{CH}_3)_2$, $-\text{CO}_2\text{CH}(\text{CH}_2)_2$, $-\text{C}(\text{O})\text{NH}_2$ (carbamoyl), $-\text{C}(\text{O})\text{NHCH}_3$, $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_3$, $-\text{CONHCH}(\text{CH}_3)_2$, $-\text{CONHCH}(\text{CH}_2)_2$, $-\text{CON}(\text{CH}_3)_2$, and $-\text{CONHCH}_2\text{CF}_3$, are non-limiting examples of heteroatom-substituted acyl groups.

The term “alkoxy” includes straight-chain alkoxy, branched-chain alkoxy, cycloalkoxy, cyclic alkoxy, heteroatom-unsubstituted alkoxy, heteroatom-substituted alkoxy, heteroatom-unsubstituted C_n-alkoxy, and heteroatom-substituted C_n-alkoxy. In certain embodiments, lower alkoxy are contemplated. The term “lower alkoxy” refers to alkoxy of 1-6 carbon atoms (that is, 1, 2, 3, 4, 5 or 6 carbon atoms). The term “heteroatom-unsubstituted C_n-alkoxy” refers to a group, having the structure $-\text{OR}$, in which R is a heteroatom-unsubstituted C_n-alkyl, as that term is defined above. Heteroatom-unsubstituted alkoxy groups include: $-\text{OCH}_3$, $-\text{OCH}_2\text{CH}_3$, $-\text{OCH}_2\text{CH}_2\text{CH}_3$, $-\text{OCH}(\text{CH}_3)_2$, and $-\text{OCH}(\text{CH}_2)_2$. The term “heteroatom-substituted C_n-alkoxy” refers to a group, having the structure $-\text{OR}$, in which R is a heteroatom-substituted C_n-alkyl, as that term is defined above. For example, $-\text{OCH}_2\text{CF}_3$ is a heteroatom-substituted alkoxy group.

The term “alkenyloxy” includes straight-chain alkenyloxy, branched-chain alkenyloxy, cycloalkenyloxy, cyclic alkenyloxy, heteroatom-unsubstituted alkenyloxy, heteroatom-substituted alkenyloxy, heteroatom-unsubstituted C_n-alkenyloxy, and heteroatom-substituted C_n-alkenyloxy. The term “heteroatom-unsubstituted C_n-alkenyloxy” refers to a group, having the structure $-\text{OR}$, in which R is a heteroatom-unsubstituted C_n-alkenyl, as that term is defined above. The term “heteroatom-substituted C_n-alkenyloxy” refers to

a group, having the structure —OR, in which R is a heteroatom-substituted C_n-alkenyl, as that term is defined above.

The term “alkynyloxy” includes straight-chain alkynyloxy, branched-chain alkynyloxy, cycloalkynyloxy, cyclic alkynyloxy, heteroatom-unsubstituted alkynyloxy, heteroatom-substituted alkynyloxy, heteroatom-unsubstituted C_n-alkynyloxy, and heteroatom-substituted C_n-alkynyloxy. The term “heteroatom-unsubstituted C_n-alkynyloxy” refers to a group, having the structure —OR, in which R is a heteroatom-unsubstituted C_n-alkynyl, as that term is defined above. The term “heteroatom-substituted C_n-alkynyloxy” refers to a group, having the structure —OR, in which R is a heteroatom-substituted C_n-alkynyl, as that term is defined above.

The term “aryloxy” includes heteroatom-unsubstituted aryloxy, heteroatom-substituted aryloxy, heteroatom-unsubstituted C_n-aryloxy, heteroatom-substituted C_n-aryloxy, heteroaryloxy, and heterocyclic aryloxy groups. The term “heteroatom-unsubstituted C_n-aryloxy” refers to a group, having the structure —OAr, in which Ar is a heteroatom-unsubstituted C_n-aryl, as that term is defined above. A non-limiting example of a heteroatom-unsubstituted aryloxy group is —OC₆H₅. The term “heteroatom-substituted C_n-aryloxy” refers to a group, having the structure —OAr, in which Ar is a heteroatom-substituted C_n-aryl, as that term is defined above.

The term “aralkyloxy” includes heteroatom-unsubstituted aralkyloxy, heteroatom-substituted aralkyloxy, heteroatom-unsubstituted C_n-aralkyloxy, heteroatom-substituted C_n-aralkyloxy, heteroaralkyloxy, and heterocyclic aralkyloxy groups. The term “heteroatom-unsubstituted C_n-aralkyloxy” refers to a group, having the structure —OAr, in which Ar is a heteroatom-unsubstituted C_n-aralkyl, as that term is defined above. The term “heteroatom-substituted C_n-aralkyloxy” refers to a group, having the structure —OAr, in which Ar is a heteroatom-substituted C_n-aralkyl, as that term is defined above.

The term “acyloxy” includes straight-chain acyloxy, branched-chain acyloxy, cycloacyloxy, cyclic acyloxy, heteroatom-unsubstituted acyloxy, heteroatom-substituted acyloxy, heteroatom-unsubstituted C_n-acyloxy, heteroatom-substituted C_n-acyloxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, and carboxylate groups. The term “heteroatom-unsubstituted C_n-acyloxy” refers to a group, having the structure —OAc, in which Ac is a heteroatom-unsubstituted C_n-acyl, as that term is defined above. For example, —OC(O)CH₃ is a non-limiting example of a heteroatom-unsubstituted acyloxy group. The term “heteroatom-substituted C_n-acyloxy” refers to a group, having the structure —OAc, in which Ac is a heteroatom-substituted C_n-acyl, as that term is defined above. For example, —OC(O)OCH₃ and —OC(O)NHCH₃ are non-limiting examples of heteroatom-unsubstituted acyloxy groups.

The term “alkylamino” includes straight-chain alkylamino, branched-chain alkylamino, cycloalkylamino, cyclic alkylamino, heteroatom-unsubstituted alkylamino, heteroatom-substituted alkylamino, heteroatom-unsubstituted C_n-alkylamino, and heteroatom-substituted C_n-alkylamino. The term “heteroatom-unsubstituted C_n-alkylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing a total of n carbon atoms, all of which are nonaromatic, 4 or more hydrogen atoms, a total of 1 nitrogen atom, and no

additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-alkylamino has 1 to 10 carbon atoms. The term “heteroatom-unsubstituted C_n-alkylamino” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-alkyl, as that term is defined above. A heteroatom-unsubstituted alkylamino group would include —NHCH₃, —NHCH₂CH₃, —NHCH₂CH₂CH₃, —NHCH(CH₃)₂, —NHCH(CH₃)CH₂CH₃, —NHCH₂CH₂CH₂CH₃, —NHCH(CH₃)CH₂CH₂CH₃, —NHCH₂CH(CH₃)CH₂CH₃, —NHC(CH₃)₃, —N(CH₃)₂, —N(CH₃)CH₂CH₃, —N(CH₂CH₃)₂, N-pyrrolidinyl, and N-piperidinyl. The term “heteroatom-substituted C_n-alkylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, no carbon-carbon double or triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-alkylamino has 1 to 10 carbon atoms. The term “heteroatom-substituted C_n-alkylamino” includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-alkyl, as that term is defined above.

The term “alkenylamino” includes straight-chain alkenylamino, branched-chain alkenylamino, cycloalkenylamino, cyclic alkenylamino, heteroatom-unsubstituted alkenylamino, heteroatom-substituted alkenylamino, heteroatom-unsubstituted C_n-alkenylamino, heteroatom-substituted C_n-alkenylamino, dialkenylamino, and alkyl(alkenyl)amino groups. The term “heteroatom-unsubstituted C_n-alkenylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing at least one nonaromatic carbon-carbon double bond, a total of n carbon atoms, 4 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₂-C₁₀-alkenylamino has 2 to 10 carbon atoms. The term “heteroatom-unsubstituted C_n-alkenylamino” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted C_n-alkenyl, as that term is defined above. The term “heteroatom-substituted C_n-alkenylamino” refers to a radical, having a single nitrogen atom as the point of attachment and at least one nonaromatic carbon-carbon double bond, but no carbon-carbon triple bonds, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkenylamino has 2 to 10 carbon atoms. The term “heteroatom-substituted C_n-alkenylamino” includes groups, having the structure —NHR, in which R is a heteroatom-substituted C_n-alkenyl, as that term is defined above.

The term “alkynylamino” includes straight-chain alkynylamino, branched-chain alkynylamino, cycloalkynylamino, cyclic alkynylamino, heteroatom-unsubstituted alkynylamino, heteroatom-substituted alkynylamino, heteroatom-unsubstituted C_n-alkynylamino, and heteroatom-substituted C_n-alkynylamino. The term “heteroatom-unsubstituted C_n-alkynylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing a total of n carbon atoms, all of which are nonaromatic, 4 or more hydrogen atoms, a total of 1 nitrogen atom, and no

stituted Cn-alkynylamino, dialkynylamino, alkyl(alkynyl) amino, and alkenyl(alkynyl)amino groups. The term “heteroatom-unsubstituted Cn-alkynylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, containing at least one carbon-carbon triple bond, a total of n carbon atoms, at least one hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₂-C₁₀-alkynylamino has 2 to 10 carbon atoms. The term “heteroatom-unsubstituted Cn-alkynylamino” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted Cn-alkynyl, as that term is defined above. The term “heteroatom-substituted Cn-alkynylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having one or two carbon atoms attached to the nitrogen atom, further having at least one nonaromatic carbon-carbon triple bond, further having a linear or branched, cyclic or acyclic structure, and further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₂-C₁₀-alkynylamino has 2 to 10 carbon atoms. The term “heteroatom-substituted Cn-alkynylamino” includes groups, having the structure —NHR, in which R is a heteroatom-substituted Cn-alkynyl, as that term is defined above.

The term “arylamino” includes heteroatom-unsubstituted arylamino, heteroatom-substituted arylamino, heteroatom-unsubstituted Cn-arylamino, heteroatom-substituted Cn-arylamino, heteroarylamino, heterocyclic arylamino, and alkyl(aryl)amino groups. The term “heteroatom-unsubstituted Cn-arylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having at least one aromatic ring structure attached to the nitrogen atom, wherein the aromatic ring structure contains only carbon atoms, further having a total of n carbon atoms, 6 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₆-C₁₀-arylamino has 6 to 10 carbon atoms. The term “heteroatom-unsubstituted Cn-arylamino” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted Cn-aryl, as that term is defined above. The term “heteroatom-substituted Cn-arylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having a total of n carbon atoms, at least one hydrogen atom, at least one additional heteroatoms, that is, in addition to the nitrogen atom at the point of attachment, wherein at least one of the carbon atoms is incorporated into one or more aromatic ring structures, further wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₆-C₁₀-arylamino has 6 to 10 carbon atoms. The term “heteroatom-substituted Cn-arylamino” includes groups, having the structure —NHR, in which R is a heteroatom-substituted Cn-aryl, as that term is defined above.

The term “aralkylamino” includes heteroatom-unsubstituted aralkylamino, heteroatom-substituted aralkylamino, heteroatom-unsubstituted Cn-aralkylamino, heteroatom-substituted Cn-aralkylamino, heteroaralkylamino, heterocyclic aralkylamino groups, and diaralkylamino groups. The term “heteroatom-unsubstituted Cn-aralkylamino” refers to a radical, having a single nitrogen atom as the point of

attachment, further having one or two saturated carbon atoms attached to the nitrogen atom, further having a total of n carbon atoms, wherein at least 6 of the carbon atoms form an aromatic ring structure containing only carbon atoms, 8 or more hydrogen atoms, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₇-C₁₀-aralkylamino has 7 to 10 carbon atoms. The term “heteroatom-unsubstituted Cn-aralkylamino” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted Cn-aralkyl, as that term is defined above. The term “heteroatom-substituted Cn-aralkylamino” refers to a radical, having a single nitrogen atom as the point of attachment, further having at least one or two saturated carbon atoms attached to the nitrogen atom, further having a total of n carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom, that is, in addition to the nitrogen atom at the point of attachment, wherein at least one of the carbon atom incorporated into an aromatic ring, further wherein each heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₇-C₁₀-aralkylamino has 7 to 10 carbon atoms. The term “heteroatom-substituted Cn-aralkylamino” includes groups, having the structure —NHR, in which R is a heteroatom-substituted Cn-aralkyl, as that term is defined above.

The term “amido” includes straight-chain amido, branched-chain amido, cycloamido, cyclic amido, heteroatom-unsubstituted amido, heteroatom-substituted amido, heteroatom-unsubstituted Cn-amido, heteroatom-substituted Cn-amido, alkylcarbonylamino, arylcarbonylamino, alkoxy-carbonylamino, aryloxy-carbonylamino, acylamino, alkylaminocarbonylamino, arylaminocarbonylamino, and ureido groups. The term “heteroatom-unsubstituted Cn-amido” refers to a radical, having a single nitrogen atom as the point of attachment, further having a carbonyl group attached via its carbon atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 1 or more hydrogen atoms, a total of one oxygen atom, a total of one nitrogen atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C₁-C₁₀-amido has 1 to 10 carbon atoms. The term “heteroatom-unsubstituted Cn-amido” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted Cn-acyl, as that term is defined above. The group, —NHC(O)CH₃, is a non-limiting example of a heteroatom-unsubstituted amido group. The term “heteroatom-substituted Cn-amido” refers to a radical, having a single nitrogen atom as the point of attachment, further having a carbonyl group attached via its carbon atom to the nitrogen atom, further having a linear or branched, cyclic or acyclic structure, further having a total of n aromatic or nonaromatic carbon atoms, 0, 1, or more than one hydrogen atom, at least one additional heteroatom in addition to the oxygen of the carbonyl group, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-amido has 1 to 10 carbon atoms. The term “heteroatom-substituted Cn-amido” includes groups, having the structure —NHR, in which R is a heteroatom-unsubstituted Cn-acyl, as that term is defined above. The group, —NHCO₂CH₃, is a non-limiting example of a heteroatom-substituted amido group.

The term “alkylthio” includes straight-chain alkylthio, branched-chain alkylthio, cycloalkylthio, cyclic alkylthio, heteroatom-unsubstituted alkylthio, heteroatom-substituted alkylthio, heteroatom-unsubstituted Cn-alkylthio, and het-

eroatom-substituted Cn-alkylthio. The term "heteroatom-unsubstituted Cn-alkylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted Cn-alkyl, as that term is defined above. The group, —SCH₃, is an example of a heteroatom-unsubstituted alkylthio group. The term "heteroatom-substituted Cn-alkylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted Cn-alkyl, as that term is defined above.

The term "alkenylthio" includes straight-chain alkenylthio, branched-chain alkenylthio, cycloalkenylthio, cyclic alkenylthio, heteroatom-unsubstituted alkenylthio, heteroatom-substituted alkenylthio, heteroatom-unsubstituted Cn-alkenylthio, and heteroatom-substituted Cn-alkenylthio. The term "heteroatom-unsubstituted Cn-alkenylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted Cn-alkenyl, as that term is defined above. The term "heteroatom-substituted Cn-alkenylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted Cn-alkenyl, as that term is defined above.

The term "alkynylthio" includes straight-chain alkynylthio, branched-chain alkynylthio, cycloalkynylthio, cyclic alkynylthio, heteroatom-unsubstituted alkynylthio, heteroatom-substituted alkynylthio, heteroatom-unsubstituted Cn-alkynylthio, and heteroatom-substituted Cn-alkynylthio. The term "heteroatom-unsubstituted Cn-alkynylthio" refers to a group, having the structure —SR, in which R is a heteroatom-unsubstituted Cn-alkynyl, as that term is defined above. The term "heteroatom-substituted Cn-alkynylthio" refers to a group, having the structure —SR, in which R is a heteroatom-substituted Cn-alkynyl, as that term is defined above.

The term "arylthio" includes heteroatom-unsubstituted arylthio, heteroatom-substituted arylthio, heteroatom-unsubstituted Cn-arylthio, heteroatom-substituted Cn-arylthio, heteroarylthio, and heterocyclic arylthio groups. The term "heteroatom-unsubstituted Cn-arylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-unsubstituted Cn-aryl, as that term is defined above. The group, —SC₆H₅, is an example of a heteroatom-unsubstituted arylthio group. The term "heteroatom-substituted Cn-arylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-substituted Cn-aryl, as that term is defined above.

The term "aralkylthio" includes heteroatom-unsubstituted aralkylthio, heteroatom-substituted aralkylthio, heteroatom-unsubstituted Cn-aralkylthio, heteroatom-substituted Cn-aralkylthio, heteroaralkylthio, and heterocyclic aralkylthio groups. The term "heteroatom-unsubstituted Cn-aralkylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-unsubstituted Cn-aralkyl, as that term is defined above. The group, —SCH₂C₆H₅, is an example of a heteroatom-unsubstituted aralkyl group. The term "heteroatom-substituted Cn-aralkylthio" refers to a group, having the structure —SAr, in which Ar is a heteroatom-substituted Cn-aralkyl, as that term is defined above.

The term "acylthio" includes straight-chain acylthio, branched-chain acylthio, cycloacylthio, cyclic acylthio, heteroatom-unsubstituted acylthio, heteroatom-substituted acylthio, heteroatom-unsubstituted Cn-acylthio, heteroatom-substituted Cn-acylthio, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, and carboxylate groups. The term "heteroatom-unsubstituted Cn-acylthio" refers to a group, having the structure —SAc, in which Ac is a heteroatom-unsubstituted Cn-acyl, as that term is defined above. The group, —SCOCH₃, is an example of a heteroatom-unsubstituted acylthio group. The term

"heteroatom-substituted Cn-acylthio" refers to a group, having the structure —SAc, in which Ac is a heteroatom-substituted Cn-acyl, as that term is defined above.

The term "alkylsilyl" includes straight-chain alkylsilyl, branched-chain alkylsilyl, cycloalkylsilyl, cyclic alkylsilyl, heteroatom-unsubstituted alkylsilyl, heteroatom-substituted alkylsilyl, heteroatom-unsubstituted Cn-alkylsilyl, and heteroatom-substituted Cn-alkylsilyl. The term "heteroatom-unsubstituted Cn-alkylsilyl" refers to a radical, having a single silicon atom as the point of attachment, further having one, two, or three saturated carbon atoms attached to the silicon atom, further having a linear or branched, cyclic or acyclic structure, containing a total of n carbon atoms, all of which are nonaromatic, 5 or more hydrogen atoms, a total of 1 silicon atom, and no additional heteroatoms. For example, a heteroatom-unsubstituted C1-C10-alkylsilyl has 1 to 10 carbon atoms. An alkylsilyl group includes dialkylamino groups. The groups, —Si(CH₃)₃ and —Si(CH₃)₂C(CH₃)₃, are non-limiting examples of heteroatom-unsubstituted alkylsilyl groups. The term "heteroatom-substituted Cn-alkylsilyl" refers to a radical, having a single silicon atom as the point of attachment, further having at least one, two, or three saturated carbon atoms attached to the silicon atom, no carbon-carbon double or triple bonds, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, all of which are nonaromatic, 0, 1, or more than one hydrogen atom, and at least one additional heteroatom, that is, in addition to the silicon atom at the point of attachment, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-substituted C₁-C₁₀-alkylsilyl has 1 to 10 carbon atoms.

The term "phosphonate" includes straight-chain phosphonate, branched-chain phosphonate, cyclophosphonate, cyclic phosphonate, heteroatom-unsubstituted phosphonate, heteroatom-substituted phosphonate, heteroatom-unsubstituted Cn-phosphonate, and heteroatom-substituted Cn-phosphonate. The term "heteroatom-unsubstituted Cn-phosphonate" refers to a radical, having a single phosphorous atom as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 2 or more hydrogen atoms, a total of three oxygen atoms, and no additional heteroatoms. The three oxygen atoms are directly attached to the phosphorous atom, with one of these oxygen atoms doubly bonded to the phosphorous atom. For example, a heteroatom-unsubstituted C₀-C₁₀-phosphonate has 0 to 10 carbon atoms. The groups, —P(O)(OH)₂, —P(O)(OH)OCH₃, —P(O)(OH)OCH₂CH₃, —P(O)(OCH₃)₂, and —P(O)(OH)(OC₆H₅) are non-limiting examples of heteroatom-unsubstituted phosphonate groups. The term "heteroatom-substituted Cn-phosphonate" refers to a radical, having a single phosphorous atom as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 2 or more hydrogen atoms, three or more oxygen atoms, three of which are directly attached to the phosphorous atom, with one of these three oxygen atoms doubly bonded to the phosphorous atom, and further having at least one additional heteroatom in addition to the three oxygen atoms, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-unsubstituted C₀-C₁₀-phosphonate has 0 to 10 carbon atoms.

The term "phosphinate" includes straight-chain phosphinate, branched-chain phosphinate, cyclophosphinate, cyclic phosphinate, heteroatom-unsubstituted phosphinate, heteroatom-substituted phosphinate, heteroatom-unsubstituted

Cn-phosphinate, and heteroatom-substituted Cn-phosphinate. The term "heteroatom-unsubstituted Cn-phosphinate" refers to a radical, having a single phosphorous atom as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 2 or more hydrogen atoms, a total of two oxygen atoms, and no additional heteroatoms. The two oxygen atoms are directly attached to the phosphorous atom, with one of these oxygen atoms doubly bonded to the phosphorous atom. For example, a heteroatom-unsubstituted C0-C10-phosphinate has 0 to 10 carbon atoms. The groups, —P(O)(OH)H , —P(O)(OH)CH_3 , $\text{—P(O)(OH)CH}_2\text{CH}_3$, $\text{—P(O)(OCH}_3\text{)CH}_3$, and $\text{—P(O)(OC}_6\text{H}_5\text{)H}$ are non-limiting examples of heteroatom-unsubstituted phosphinate groups. The term "heteroatom-substituted Cn-phosphinate" refers to a radical, having a single phosphorous atom as the point of attachment, further having a linear or branched, cyclic or acyclic structure, further having a total of n carbon atoms, 2 or more hydrogen atoms, two or more oxygen atoms, two of which are directly attached to the phosphorous atom, with one of these two oxygen atoms doubly bonded to the phosphorous atom, and further having at least one additional heteroatom in addition to the two oxygen atoms, wherein each additional heteroatom is independently selected from the group consisting of N, O, F, Cl, Br, I, Si, P, and S. For example, a heteroatom-unsubstituted C0-C10-phosphinate has 0 to 10 carbon atoms.

Any apparently unfulfilled valency is to be understood to be properly filled by hydrogen atom(s). For example, a compound with a substituent of —O or —N is to be understood to be —OH or —NH_2 , respectively.

Any genus, subgenus, or specific compound discussed herein is specifically contemplated as being excluded from any embodiment described herein.

Compounds described herein may be prepared synthetically using conventional organic chemistry methods known to those of skill in the art and/or are commercially available (e.g., ChemBridge Co., San Diego, Calif.).

The claimed invention is also intended to encompass salts of any of the compounds of the present invention. The term "salt(s)" as used herein, is understood as being acidic and/or basic salts formed with inorganic and/or organic acids and bases. Zwitterions (internal or inner salts) are understood as being included within the term "salt(s)" as used herein, as are quaternary ammonium salts such as alkylammonium salts. Nontoxic, pharmaceutically acceptable salts are preferred, although other salts may be useful, as for example in isolation or purification steps during synthesis. Salts include, but are not limited to, sodium, lithium, potassium, amines, tartrates, citrates, hydrohalides, phosphates and the like. A salt may be a pharmaceutically acceptable salt, for example. Thus, pharmaceutically acceptable salts of compounds of the present invention are contemplated.

The term "pharmaceutically acceptable salts," as used herein, refers to salts of compounds of this invention that are substantially non-toxic to living organisms. Typical pharmaceutically acceptable salts include those salts prepared by reaction of a compound of this invention with an inorganic or organic acid, or an organic base, depending on the substituents present on the compounds of the invention.

Non-limiting examples of inorganic acids which may be used to prepare pharmaceutically acceptable salts include: hydrochloric acid, phosphoric acid, sulfuric acid, hydrobromic acid, hydroiodic acid, phosphorous acid and the like. Examples of organic acids which may be used to prepare pharmaceutically acceptable salts include: aliphatic mono- and dicarboxylic acids, such as oxalic acid, carbonic acid,

citric acid, succinic acid, phenyl-heteroatom-substituted alkanic acids, aliphatic and aromatic sulfuric acids and the like. Pharmaceutically acceptable salts prepared from inorganic or organic acids thus include hydrochloride, hydrobromide, nitrate, sulfate, pyrosulfate, bisulfate, sulfite, bisulfate, phosphate, monohydrogenphosphate, dihydrogen phosphate, metaphosphate, pyrophosphate, hydroiodide, hydrofluoride, acetate, propionate, formate, oxalate, citrate, lactate, p-toluenesulfonate, methanesulfonate, maleate, and the like.

Suitable pharmaceutically acceptable salts may also be formed by reacting the agents of the invention with an organic base such as methylamine, ethylamine, ethanolamine, lysine, ornithine and the like.

Pharmaceutically acceptable salts include the salts formed between carboxylate or sulfonate groups found on some of the compounds of this invention and inorganic cations, such as sodium, potassium, ammonium, or calcium, or such organic cations as isopropylammonium, trimethylammonium, tetramethylammonium, and imidazolium.

Derivatives of compounds of the present invention are also contemplated. In certain aspects, "derivative" refers to a chemically modified compound that still retains the desired effects of the compound prior to the chemical modification. Such derivatives may have the addition, removal, or substitution of one or more chemical moieties on the parent molecule. Non-limiting examples of the types modifications that can be made to the compounds and structures disclosed herein include the addition or removal of lower alkanes such as methyl, ethyl, propyl, or substituted lower alkanes such as hydroxymethyl or aminomethyl groups; carboxyl groups and carbonyl groups; hydroxyls; nitro, amino, amide, and azo groups; sulfate, sulfonate, sulfono, sulphydryl, sulfonyl, sulfoxido, phosphate, phosphono, phosphoryl groups, and halide substituents. Additional modifications can include an addition or a deletion of one or more atoms of the atomic framework, for example, substitution of an ethyl by a propyl; substitution of a phenyl by a larger or smaller aromatic group. Alternatively, in a cyclic or bicyclic structure, heteroatoms such as N, S, or O can be substituted into the structure instead of a carbon atom.

Compounds employed in methods may contain one or more asymmetrically-substituted carbon or nitrogen atoms, and may be isolated in optically active or racemic form. Thus, all chiral, diastereomeric, racemic form, epimeric form, and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated. Compounds may occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. In some embodiments, a single diastereomer is obtained. The chiral centers of the compounds of the present invention can have the S- or the R-configuration, as defined by the IUPAC 1974 Recommendations. Compounds may be of the D- or L-form, for example. It is well known in the art how to prepare and isolate such optically active forms. For example, mixtures of stereoisomers may be separated by standard techniques including, but not limited to, resolution of racemic form, normal, reverse-phase, and chiral chromatography, preferential salt formation, recrystallization, and the like, or by chiral synthesis either from chiral starting materials or by deliberate synthesis of target chiral centers.

In addition, atoms making up the compounds described herein are intended to include all isotopic forms of such atoms. Isotopes, as used herein, include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of

hydrogen include tritium and deuterium, and isotopes of carbon include ^{13}C and ^{14}C .

As noted above, compounds described herein may exist in prodrug form. As used herein, "prodrug" is intended to include any covalently bonded carriers which release the active parent drug or compounds that are metabolized in vivo to an active drug or other compounds employed in the methods described herein in vivo when such prodrug is administered to a subject. Since prodrugs are known to enhance numerous desirable qualities of pharmaceuticals (e.g., solubility, bioavailability, manufacturing, etc.), the compounds employed in some methods described herein may, if desired, be delivered in prodrug form. Thus, the invention contemplates prodrugs of compounds described herein as well as methods of delivering prodrugs. Prodrugs of the compounds employed in embodiments may be prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or in vivo, to the parent compound.

Accordingly, prodrugs include, for example, compounds described herein in which a hydroxy, amino, or carboxy group is bonded to any group that, when the prodrug is administered to a subject, cleaves to form a free hydroxyl, free amino, or carboxylic acid, respectively. Other examples include, but are not limited to, acetate, formate, and benzoate derivatives of alcohol and amine functional groups; and alkyl, carbocyclic, aryl, and alkylaryl esters such as methyl, ethyl, propyl, iso-propyl, butyl, isobutyl, sec-butyl, tert-butyl, cyclopropyl, phenyl, benzyl, and phenethyl esters, and the like.

It should be recognized that the particular anion or cation forming a part of any salt in any embodiment is not critical, so long as the salt, as a whole, is pharmacologically acceptable. Additional examples of pharmaceutically acceptable salts and their methods of preparation and use are presented in Handbook of Pharmaceutical Salts: Properties, Selection and Use (2002), which is incorporated herein by reference.

C. PHARMACEUTICAL FORMULATIONS AND ADMINISTRATION THEREOF

1. Pharmaceutical Formulations and Routes of Administration

Pharmaceutical compositions disclosed herein comprise an effective amount of one or more candidate substance or additional agent dissolved or dispersed in a pharmaceutically acceptable carrier. The phrases "pharmaceutical or pharmacologically acceptable" refers to molecular entities and compositions that do not produce an adverse, allergic or other untoward reaction when administered to an animal, such as, for example, a human, as appropriate. The preparation of a pharmaceutical composition that contains at least one candidate substance or additional active ingredient will be known to those of skill in the art in light of the present disclosure, as exemplified by Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, incorporated herein by reference. Moreover, for animal (e.g., human) administration, it will be understood that preparations should meet sterility, pyrogenicity, general safety and purity standards as required by FDA Office of Biological Standards.

As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, absorption delaying agents, salts, preservatives, drugs, drug stabilizers, gels,

binders, excipients, disintegration agents, lubricants, sweetening agents, flavoring agents, dyes, such like materials and combinations thereof, as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 18th Ed. Mack Printing Company, 1990, pp. 1289-1329). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated.

The compounds administered according to embodiments disclosed herein may comprise different types of carriers depending on whether it is to be administered in solid, liquid or aerosol form, and whether it need to be sterile for such routes of administration as injection. Embodiments can be administered intravenously, intradermally, intraarterially, intraperitoneally, intrasessionally, intracranially, intraarticularly, intraprostatically, intrapleurally, intratracheally, intranasally, intravitreally, intravaginally, intrarectally, topically, intratumorally, intramuscularly, systemically, subcutaneously, subconjunctival, intravesicularly, mucosally, intrapericardially, intraumbilically, intraocularly, orally, locally, via inhalation (e.g., aerosol inhalation), via injection, via infusion, via continuous infusion, via localized perfusion bathing target cells directly, via a catheter, via a lavage, in cremes, in lipid compositions (e.g., liposomes), or by other method or any combination of the foregoing as would be known to one of ordinary skill in the art (see, for example, Remington's Pharmaceutical Sciences, 1990).

The actual dosage amount of a composition of embodiments disclosed herein that are administered to an animal patient can be determined by physical and physiological factors such as body weight, severity of condition, the type of disease being treated, previous or concurrent therapeutic interventions, idiopathy of the patient and on the route of administration. The practitioner responsible for administration will, in any event, determine the concentration of active ingredient(s) in a composition and appropriate dose(s) for the individual subject.

In certain embodiments, pharmaceutical compositions may comprise, for example, at least about 0.1% of a compound of the present disclosure. In other embodiments, the compound may comprise between about 2% to about 75% of the weight of the unit, or between about 25% to about 60%, for example, and any range derivable therein. In other non-limiting examples, a dose may also comprise from about 1 microgram/kg/body weight, about 5 microgram/kg/body weight, about 10 microgram/kg/body weight, about 50 microgram/kg/body weight, about 100 microgram/kg/body weight, about 200 microgram/kg/body weight, about 350 microgram/kg/body weight, about 500 microgram/kg/body weight, about 1 milligram/kg/body weight, about 5 milligram/kg/body weight, about 10 milligram/kg/body weight, about 50 milligram/kg/body weight, about 100 milligram/kg/body weight, about 200 milligram/kg/body weight, about 350 milligram/kg/body weight, about 500 milligram/kg/body weight, to about 1000 mg/kg/body weight or more per administration, and any range derivable therein. In non-limiting examples of a derivable range from the numbers listed herein, a range of about 5 mg/kg/body weight to about 100 mg/kg/body weight, about 5 microgram/kg/body weight to about 500 milligram/kg/body weight, etc., can be administered, based on the numbers described above.

In any case, the composition may comprise various antioxidants to retard oxidation of one or more component. Additionally, the prevention of the action of microorganisms can be brought about by preservatives such as various antibacterial and antifungal agents, including but not limited

to parabens (e.g., methylparabens, propylparabens), chlorobutanol, phenol, sorbic acid, thimerosal, or combinations thereof.

The candidate substance may be formulated into a composition in a free base, neutral or salt form. Pharmaceutically acceptable salts, include the acid addition salts, e.g., those formed with the free amino groups of a proteinaceous composition, or which are formed with inorganic acids such as for example, hydrochloric or phosphoric acids, or such organic acids as acetic, oxalic, tartaric or mandelic acid. Salts formed with the free carboxyl groups can also be derived from inorganic bases such as for example, sodium, potassium, ammonium, calcium or ferric hydroxides; or such organic bases as isopropylamine, trimethylamine, histidine, or procaine.

In embodiments where the composition is in a liquid form, a carrier can be a solvent or dispersion medium comprising but not limited to, water, ethanol, polyol (e.g., glycerol, propylene glycol, liquid polyethylene glycol, etc.), lipids (e.g., triglycerides, vegetable oils, liposomes) and combinations thereof. The proper fluidity can be maintained, for example, by the use of a coating, such as lecithin; by the maintenance of the required particle size by dispersion in carriers such as, for example liquid polyol or lipids; by the use of surfactants such as, for example hydroxypropylcellulose; or combinations thereof such methods. It may be preferable to include isotonic agents, such as, for example, sugars, sodium chloride or combinations thereof.

In other embodiments, one may use eye drops, nasal solutions or sprays, aerosols or inhalants. Such compositions are generally designed to be compatible with the target tissue type. In a non-limiting example, nasal solutions are usually aqueous solutions designed to be administered to the nasal passages in drops or sprays. Nasal solutions are prepared so that they are similar in many respects to nasal secretions, so that normal ciliary action is maintained. Thus, in certain embodiments the aqueous nasal solutions usually are isotonic or slightly buffered to maintain a pH of about 5.5 to about 6.5. In addition, antimicrobial preservatives, similar to those used in ophthalmic preparations, drugs, or appropriate drug stabilizers, if required, may be included in the formulation. For example, various commercial nasal preparations are known and include drugs such as antibiotics or antihistamines.

In certain embodiments the candidate substance is prepared for administration by such routes as oral ingestion. In these embodiments, the solid composition may comprise, for example, solutions, suspensions, emulsions, tablets, pills, capsules (e.g., hard or soft shelled gelatin capsules), sustained release formulations, buccal compositions, troches, elixirs, suspensions, syrups, wafers, or combinations thereof. Oral compositions may be incorporated directly with the food of the diet. In certain embodiments, carriers for oral administration comprise inert diluents, assimilable edible carriers or combinations thereof. In other aspects, the oral composition may be prepared as a syrup or elixir. A syrup or elixir, and may comprise, for example, at least one active agent, a sweetening agent, a preservative, a flavoring agent, a dye, a preservative, or combinations thereof.

In certain embodiments an oral composition may comprise one or more binders, excipients, disintegration agents, lubricants, flavoring agents, and combinations thereof. In certain embodiments, a composition may comprise one or more of the following: a binder, such as, for example, gum tragacanth, acacia, cornstarch, gelatin or combinations thereof, an excipient, such as, for example, dicalcium phosphate, mannitol, lactose, starch, magnesium stearate, sodium

saccharine, cellulose, magnesium carbonate or combinations thereof; a disintegrating agent, such as, for example, corn starch, potato starch, alginic acid or combinations thereof; a lubricant, such as, for example, magnesium stearate; a sweetening agent, such as, for example, sucrose, lactose, saccharin or combinations thereof; a flavoring agent, such as, for example peppermint, oil of wintergreen, cherry flavoring, orange flavoring, etc.; or combinations thereof the foregoing. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, carriers such as a liquid carrier. Various other materials may be present as coatings or to otherwise modify the physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar, or both.

Additional formulations which are suitable for other modes of administration include suppositories. Suppositories are solid dosage forms of various weights and shapes, usually medicated, for insertion into the rectum, vagina, or urethra. After insertion, suppositories soften, melt or dissolve in the cavity fluids. In general, for suppositories, traditional carriers may include, for example, polyalkylene glycols, triglycerides, or combinations thereof. In certain embodiments, suppositories may be formed from mixtures containing, for example, the active ingredient in the range of about 0.5% to about 10%, and preferably about 1% to about 2%.

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and/or the other ingredients. In the case of sterile powders for the preparation of sterile injectable solutions, suspensions or emulsion, certain methods of preparation may include vacuum-drying or freeze-drying techniques which yield a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered liquid medium thereof. The liquid medium should be suitably buffered if necessary and the liquid diluent first rendered isotonic prior to injection with sufficient saline or glucose. The preparation of highly concentrated compositions for direct injection is also contemplated, where the use of DMSO as solvent is envisioned to result in extremely rapid penetration, delivering high concentrations of the active agents to a small area.

The composition must be stable under the conditions of manufacture and storage, and preserved against the contaminating action of microorganisms, such as bacteria and fungi. It will be appreciated that endotoxin contamination should be kept minimally at a safe level, for example, less than 0.5 ng/mg protein.

In particular embodiments, prolonged absorption of an injectable composition can be brought about by the use in the compositions of agents delaying absorption, such as, for example, aluminum monostearate, gelatin, or combinations thereof.

2. Combination Therapy

In some embodiments, it is contemplated that the tolerance-inducing compositions disclosed herein may be used in conjunction with the compositions for which a tolerance is being induced as part of a treatment regimen. This process may involve contacting the cell(s) with the agents at the same time or within a period of time wherein separate administration of the agents produces a desired therapeutic benefit. This may be achieved by contacting the cell, tissue or organism with a single composition or pharmacological

formulation that includes two or more agents, or by contacting the cell with two or more distinct compositions or formulations, wherein one composition includes one agent and the other includes another.

Compounds discussed herein may precede, be co-current with and/or follow the other agents by intervals ranging from minutes to weeks. In embodiments where the agents are applied separately to a cell, tissue or organism, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agents would still be able to exert an advantageously combined effect on the cell, tissue or organism. For example, in such instances, it is contemplated that one may contact the cell, tissue or organism with two, three, four or more modalities substantially simultaneously (i.e., within less than about a minute) as the candidate substance. In other aspects, one or more tolerance-inducing compositions may be administered or provided within 1 minute, 5 minutes, 10 minutes, 20 minutes, 30 minutes, 45 minutes, 60 minutes, 2 hours, 3 hours, 4 hours, 5 hours, 6 hours, 7 hours, 8 hours, 9 hours, 10 hours, 11 hours, 12 hours, 13 hours, 14 hours, 15 hours, 16 hours, 17 hours, 18 hours, 19 hours, 20 hours, 21 hours, 22 hours, 23 hours, 24 hours, 25 hours, 26 hours, 27 hours, 28 hours, 29 hours, 30 hours, 31 hours, 32 hours, 33 hours, 34 hours, 35 hours, 36 hours, 37 hours, 38 hours, 39 hours, 40 hours, 41 hours, 42 hours, 43 hours, 44 hours, 45 hours, 46 hours, 47 hours, 48 hours, 1 day, 2 days, 3 days, 4 days, 5 days, 6 days, 7 days, 8 days, 9 days, 10 days, 11 days, 12 days, 13 days, 14 days, 15 days, 16 days, 17 days, 18 days, 19 days, 20 days, 21 days, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, or 8 weeks or more, and any range derivable therein, prior to administering the compositions for which a tolerance is being induced. In some embodiments, more than one course of therapy may be employed. It is contemplated that multiple courses may be implemented.

D. ORGANISMS AND CELL SOURCE

Cells that may be used in some methods can be from a variety of sources. Embodiments include the use of mammalian cells, such as cells from monkeys, chimpanzees, rabbits, mice, rats, ferrets, dogs, pigs, humans, and cows. Alternatively, the cells may be from fruit flies, yeast, or *E. coli*, which are all model systems for evaluating homologous recombination.

Methods can involve cells, tissues, or organs involving the heart, lung, kidney, liver, bone marrow, pancreas, skin, bone, vein, artery, cornea, blood, small intestine, large intestine, brain, spinal cord, smooth muscle, skeletal muscle, ovary, testis, uterus, and umbilical cord.

Moreover, methods can be employed in cells of the following type: platelet, myelocyte, erythrocyte, lymphocyte, adipocyte, fibroblast, epithelial cell, endothelial cell, smooth muscle cell, skeletal muscle cell, endocrine cell, glial cell, neuron, secretory cell, barrier function cell, contractile cell, absorptive cell, mucosal cell, limbus cell (from cornea), stem cell (totipotent, pluripotent or multipotent), unfertilized or fertilized oocyte, or sperm.

Moreover, methods can be implemented with or in plants or parts of plants, including fruit, flowers, leaves, stems, seeds, cuttings. Plants can be agricultural, medicinal, or decorative.

E. ANTIGENS

The antigen employed as X in the compositions of Formula 1, or in any of the compositions or methods of the

current disclosure, can be a protein or a peptide, e.g. the antigen may be a complete or partial therapeutic agent, a full-length transplant protein or peptide thereof, a full-length autoantigen or peptide thereof, a full-length allergen or peptide thereof, and/or a nucleic acid, or a mimetic of an aforementioned antigen. Combinations of multiple fragments may also be used, depending on the embodiment. For example, if a longer peptide identified as P has antigenic regions A, B, C, and D, compositions disclosed herein for induction of tolerance to P can comprise any combination of A, B, C, and D, and repeats of any of A, B, C, and D. A listing of any particular antigen in a category or association with any particular disease or reaction does not preclude that antigen from being considered part of another category or associated with another disease or reaction.

In several embodiments, the antigen comprises one or more therapeutic agents that are proteins, peptides, antibodies and antibody-like molecules (including antibody fragments and fusion proteins with antibodies and antibody fragments), and gene therapy vectors. These include human, non-human (such as mouse) and non-natural (e.g., engineered) proteins, antibodies, chimeric antibodies, humanized antibodies, viruses and virus-like particles, and non-antibody binding scaffolds, such as fibronectins, DARPins, knottins, and the like. In several embodiments, human allograft transplantation antigens against which transplant recipients develop an unwanted immune response are used. In several embodiments, the antigen comprises one or more self-antigens that cause an unwanted, autoimmune response. While self-antigens are of an endogenous origin in an autoimmune disease patient, according to several embodiments, the polypeptides employed in the disclosed compositions are, depending on the embodiment, synthesized exogenously (as opposed to being purified and concentrated from a source of origin).

In several embodiments, the antigen to which tolerance is desired comprises one or more foreign antigens, such as food, animal, plant and environmental antigens, against which a patient experiences an unwanted immune response. While a therapeutic protein can also be considered a foreign antigen due to its exogenous origin, for purposes of clarity in the description of the present disclosure such therapeutics are described as a separate group. Similarly, a plant or an animal antigen can be eaten and considered a food antigen, and an environmental antigen may originate from a plant. They are, however, considered foreign antigens. In the interest of simplicity no attempt will be made to describe distinguish and define all of such potentially overlapping groups, as those skilled in the art can appreciate the antigens that can be employed in the compositions of the disclosure, particularly in light of the detailed description and examples.

In several embodiments, X is selected from the group consisting of insulin, proinsulin, preproinsulin, gluten, gliadin, myelin basic protein, myelin oligodendrocyte glycoprotein and proteolipid protein, Factor VIII, Factor IX, asparaginase, uricase and fragments of any of the preceding. In several embodiments, the antigen X is not a full length protein. For example, in some embodiments, the antigen is not full length gliadin, insulin, or proinsulin. In several embodiments, the antigen is not full length myelin basic protein, not full length myelin oligodendrocyte protein, or not full length proteolipid protein. In several embodiments, the antigen X is not a fragment of a protein. As discussed in more detail below, there exist a variety of antigens to which tolerance may be desired. These may include, but are not limited to, exogenous antigens that result in an adverse immune response when a subject is exposed to the antigen.

67

In several embodiments, the adverse immune response could be a result of ingestion of the antigen, e.g., orally or nasally, or via some other mucosal route. These routes could be the case, for example, with food antigens. In some embodiments, the antigen may be purposefully administered to a subject, for example, with the administration of a therapeutic composition to treat a disease or condition that the subject is affected by. In still additional embodiments, the antigen may be produced by the subject, e.g., an autoimmune antigen. For example, in several embodiments, X comprises a foreign transplant antigen against which transplant recipients develop an unwanted immune response or a tolerogenic portion thereof. In several embodiments, X comprises a foreign food, animal, plant or environmental antigen against which patients develop an unwanted immune response or a tolerogenic portion thereof. In several embodiments, X comprises a foreign therapeutic agent against which patients develop an unwanted immune response or a tolerogenic portion thereof. In several embodiments, X comprises a synthetic self-antigen against the endogenous version of which patients develop an unwanted immune response or a tolerogenic portion thereof.

In further detail to the above, there are provided in several embodiments, compounds where X is a food antigen. In some such embodiments, X is one or more of conarachin (Ara h 1), allergen II (Ara h 2), arachis agglutinin, conglutinin (Ara h 6), α -lactalbumin (ALA), lactotransferrin, Pen a 1 allergen (Pen a 1), allergen Pen m 2 (Pen m 2), tropomyosin fast isoform, high molecular weight glutenin, low molecular weight glutenin, alpha-gliadin, gamma-gliadin, omega-gliadin, hordein, seclain, and avenin. Fragment of any of these antigens and/or mimotopes of any of these antigens are also used, in several embodiments. In several embodiments, X is selected from the group consisting of gluten, high molecular weight glutenin, low molecular weight glutenin, alpha-gliadin, gamma-gliadin, omega-gliadin, hordein, seclain, and avenin and fragments thereof. In several embodiments, X is selected from the group consisting of gluten, high molecular weight glutenin, low molecular weight glutenin, alpha-gliadin, gamma-gliadin, and omega-gliadin and fragments thereof. In several embodiments, X is gluten or fragment thereof. In several embodiments, X is gliadin or fragment thereof.

In several embodiments, there are provided compounds where X is a therapeutic agent. In several embodiments, X is selected from the group consisting of Factor VII, Factor IX, asparaginase, and uricase and fragments thereof. In several embodiments, X is a therapeutic agent selected from the group consisting of Factor VII and Factor IX and fragments thereof. In several embodiments, X is a therapeutic agent selected from the group consisting of Factor VIII or fragment thereof. In several embodiments, when X is a therapeutic agent, the compound can be used in the treatment, prevention, reduction, or otherwise amelioration of an immune response developed against a therapeutic agent for hemophilia. As discussed herein, mimotopes of any antigenic portion of the antigens above can be used in several embodiments.

In several embodiments, X comprises asparaginase or a fragment thereof. In several embodiments, X comprises uricase or a fragment thereof. In several such embodiments, the compound can be used in the treatment, prevention, reduction, or otherwise amelioration of an immune response developed against an anti-neoplastic agent. As discussed herein, mimotopes of any antigenic portion of the antigens above can be used in several embodiments.

68

In several embodiments, X is associated with an autoimmune disease. For example, in several embodiments, the associated autoimmune disease is one or more of Type I diabetes, multiple sclerosis, rheumatoid arthritis, vitiligo, uveitis, pemphigus vulgaris and neuromyelitis optica.

In several embodiments, the autoimmune disease is Type I diabetes and X comprises insulin or a fragment thereof. In several embodiments, the autoimmune disease is Type I diabetes and X comprises proinsulin or a fragment thereof. In several embodiments, the autoimmune disease is Type I diabetes and X comprises preproinsulin or a fragment thereof. As discussed herein, mimotopes of any antigenic portion of the antigens above can be used in several embodiments. In several embodiments, combinations of these antigens can be incorporated into the tolerogenic compound which may aid in reducing immune responses to self-antigens at multiple points along the insulin pathway.

In several embodiments, the autoimmune disease is multiple sclerosis and X comprises myelin basic protein or a fragment thereof. In several embodiments, the autoimmune disease is multiple sclerosis and X comprises myelin oligodendrocyte glycoprotein or a fragment thereof. In several embodiments, the autoimmune disease is multiple sclerosis and X comprises proteolipid protein or a fragment thereof. As discussed herein, mimotopes of any antigenic portion of the antigens above can be used in several embodiments. In several embodiments, combinations of these antigens can be incorporated into the tolerogenic compound (e.g., a mixture of antigens or fragments of MOG, MBP and/or PLP) which may aid in reducing immune responses to self-antigens at multiple points along the enzymatic pathways that control myelination or myelin repair.

As discussed herein, mimotopes of any antigenic portion of the self-antigens above (or otherwise disclosed herein) can be used in several embodiments.

In several embodiments, the pharmaceutically acceptable composition consists of, or consists essentially of a compound wherein X is a food antigen, therapeutic agent, a self antigen, or fragment thereof, a linker Y, and a liver targeting moiety Z selected from mannose and/or a mannose receptor-targeting moiety (including, but not limited to, one or more of mannosamine, N-acetylmannosamine, or N-acetylglucosamine).

The tolerogenic antigen can be a complete protein, a portion of a complete protein, a peptide, or the like, and can be derivatized (as discussed above) for attachment to a linker and/or antigen-binding moiety, can be a variant and/or can contain conservative substitutions, particularly maintaining sequence identity, and/or can be desialylated.

In the embodiments where the antigen is a therapeutic protein, peptide, antibody or antibody-like molecule, specific antigens can be selected from: Abatacept, Abciximab, Adalimumab, Adenosine deaminase, Ado-trastuzumab emtansine, Agalsidase alfa, Agalsidase beta, Aldeslakin, Alglucerase, Alglucosidase alfa, α -1-proteinase inhibitor, Anakinra, Anistreplase (anisoyleated plasminogen streptokinase activator complex), Antithrombin III, Antithymocyte globulin, Ateplase, Bevacizumab, Bivalirudin, Botulinum toxin type A, Botulinum toxin type B, C1-esterase inhibitor, Canakinumab, Carboxypeptidase G2 (Glucarpidase and Voraxaze), Certolizumab pegol, Cetuximab, Collagenase, Crotalidae immune Fab, Darbepoetin- α , Denosumab, Digoxin immune Fab, Eculizumab, Etanercept, Factor VIIa, Factor VIII, Factor IX, Factor XI, Factor XIII, Fibrinogen, Filgrastim, Galsulfase, Golimumab, Histrelin acetate, Hyaluronidase, Idursulphase, Imiglucerase, Infliximab, Insulin [including recombinant human insulin

("rHu insulin") and bovine insulin], Interferon- α 2a, Interferon- α 2b, Interferon- β 1a, Interferon- β 1b, Interferon- γ 1b, Ipilimumab, L-arginase, L-asparaginase, L-methionase, Lactase, Laronidase, Lepirudin/hirudin, Mecasermin, Mecasermin rinfabate, Methoxy Natalizumab, Octreotide, Ofatumumab, Oprelvekin, Pancreatic amylase, Pancreatic lipase, Papain, Peg-asparaginase, Peg-doxorubicin HCl, PEG-epoetin- β , Pegfilgrastim, Peg-Interferon- α 2a, Peg-Interferon- α 2b, Pegloticase, Pegvisomant, Phenylalanine ammonia-lyase (PAL), Protein C, Rasburicase (uricase), Sacrosidase, Salmon calcitonin, Sargramostim, Streptokinase, Tenecteplase, Teriparatide, Tocilizumab (atlizumab), Trastuzumab, Type 1 alpha-interferon, Ustekinumab, vW factor. The therapeutic protein can be obtained from natural sources (e.g., concentrated and purified) or synthesized, e.g., recombinantly, and includes antibody therapeutics that are typically IgG monoclonal or fragments or fusions.

Particular therapeutic protein, peptide, antibody or antibody-like molecules include, but are not limited to, Abciximab, Adalimumab, Agalsidase alfa, Agalsidase beta, Aldeslakin, Alglucosidase alfa, Factor VIII, Factor IX, Infliximab, Insulin (including rHu Insulin), L-asparaginase, Laronidase, Natalizumab, Octreotide, Phenylalanine ammonia-lyase (PAL), or Rasburicase (uricase) and generally IgG monoclonal antibodies in their varying formats.

Some embodiments employ hemostatic agents (e.g., Factor VIII and IX), Insulin (including rHu Insulin), and the non-human therapeutics uricase, PAL and asparaginase.

In several embodiments, therapeutic agents are delivered through the use of, e.g., a gene therapy vector. In some such embodiments, an immune response may be developed against a portion of such vectors and/or their cargo (e.g., the therapeutic agent). Thus, in several embodiments, the antigen to which tolerance is desired comprises a gene therapy vector, including, but are not limited to: adenoviruses and adeno-associated virus (and corresponding variants-1, -2, -5, -6, -8, -9, and/or other parvoviruses), lentiviruses, and retroviruses.

Unwanted immune response in hematology and transplant includes autoimmune aplastic anemia, transplant rejection (generally), and Graft vs. Host Disease (bone marrow transplant rejection). In the embodiments where the tolerogenic antigen is a human allograft transplantation antigen, specific sequences can be selected from: subunits of the various MHC class I and MHC class II haplotype proteins (for example, donor/recipient differences identified in tissue cross-matching), and single-amino-acid polymorphisms on minor blood group antigens including RhCE, Kell, Kidd, Duffy and Ss. Such compositions can be prepared individually for a given donor/recipient pair.

In type 1 diabetes mellitus, antigens include, but are not limited to: insulin, proinsulin, preproinsulin, glutamic acid decarboxylase-65 (GAD-65 or glutamate decarboxylase 2), GAD-67, glucose-6 phosphatase 2 (IGRP or islet-specific glucose 6 phosphatase catalytic subunit related protein), insulinoma-associated protein 2 (IA-2), and insulinoma-associated protein 2 β (IA-2 β); other antigens include ICA69, ICA12 (SOX-13), carboxypeptidase H, Imogen 38, GLIMA 38, chromogranin-A, HSP-60, carboxypeptidase E, peripherin, glucose transporter 2, hepatocarcinoma-intestine-pancreas/pancreatic associated protein, S100 β , glial fibrillary acidic protein, regenerating gene II, pancreatic duodenal homeobox 1, dystrophin myotonic kinase, islet-specific glucose-6-phosphatase catalytic subunit-related protein, and SST G-protein coupled receptors 1-5, or immunogenic fragments or portions of any of such antigens. It should be noted that insulin is an example of an antigen that

can be characterized both as a self-antigen and a therapeutic protein antigen. For example, rHu Insulin and bovine insulin are therapeutic protein antigens (that are the subject of unwanted immune attack), whereas endogenous human insulin is a self-antigen (that is the subject of an unwanted immune attack). Because endogenous human insulin is not available to be employed in a pharmaceutical composition, a recombinant form is employed in certain embodiments of the compositions of the disclosure.

Human insulin, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT P01308):

(SEQ ID NO: 1)
MALWMRLRLPLALLALWGPDPAAAFVNQHLGSHLVEALYLVCGERGFF
YTPKTRREAEDLQVGQVELGGGPGAGSLQPLALEGSLQKRGIVEQCCTS
ICSLYQLENYCN

GAD-65, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT Q05329):

(SEQ ID NO: 2)
MASPGSGFWSFGSEDSGSDSENPGTARAWCQVAQKFTGGIGNKLCALLY
GDAEKPAESGGSQPPRAAARKAACACDQKPCSCSKVDVNYAFLHATDLL
PACDGERPTLAFLDVMNILLQYVVKSFDRSTKVIDFHYPNELLQEYNW
ELADQPQNLEELMHQCQTTLKYAIKTGHPRYFNQLSTGLDMVGLAADWL
TSTANTNMFTYEIAPVFVLLLEYVTLKKMRIIGWPGSGDGIFSPGGAI
SNMYAMMIARFKMFPEVKEKGMALPRLIARTSESHFSLLKKGAAALGI
GTDSVILIKCDERGMIPSDLERRILEAKQKGFVPFLVSATAGTTVYGA
FDPLLAVADICKKYIWMHVDAAWGGGLLSRKHKWLKSGVERANSVTW
NPHKMMGVPLQCSALLVREEGLMQNCNMHASLYLQDQKHYDLSYDTGD
KALQCGRHVDVFKLWLMWRAGKTGFEAHVDKCLEAEYLYNIIKNREG
YEMVFDGKPGQHTNVCFWYIIPPSLRTLEDNEERMRLSKVAPVIKARMME
YGTMTVSQPLGDKVNFRRMVISNPAATHQDIDFLIEEIERLGQDL

IGRP, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT QN9QR9):

(SEQ ID NO: 3)
MDFLHRNGVLI IQHLQKDYRAYYTFNLNFMNSVGDPRNIFFIYFPLCFQF
NQTVGTKMIWVAVIGDWLNLIFKWILFGHRPYVWVQETQIYPNHSSPCL
EQFPPTCTETGPGSPSGHAMGASCWVYVMVTAALSHTVCGMDKFSITLHR
LTWSFLWSVFWLIQISVCISRVFIATHEPHQVILGVIGGMLVAEAFEHT
PGIQTASLGTLYLKTNLFLFLFAVGFYLLLRVLNIDLLWSVPIAKKWCAN
PDWIIHIDTTPFAGLVRLNGLVLFGLGFAINSEMFLLSCRGNNYTLFSRL
LCALTSLTILQLYHFLQIPTHEHLFVYVLSFCKSASIPLTVVAFIPYSV
HMLMKQSGKKSQ.

In several embodiments, human proinsulin, including an exogenously obtained form useful in the tolerogenic compositions of the disclosure, has the following sequence:

71

(SEQ ID NO: 4)
FVNQHLGSHLVEALYLVCGERGFFYTPKTRREAEDLQVGQVELGGGPG
AGSLQPLALEGSLQKRGIVEQCCTSIICSLYQLENYCN.

Depending on the embodiment, peptides/epitopes useful
in the tolerogenic compositions of the disclosure for treating
type 1 diabetes include some or all of the following
sequences, individually in a tolerogenic composition or
together in a cocktail of tolerogenic compositions:
Human Proinsulin 1-70:

(SEQ ID NO: 5)
FVNQHLGSHLVEALYLVCGERGFFYTPKTRREAEDLQVGQVELGGGPG
AGSLQPLALEGSLQKRGIVEQ;

Human Proinsulin 9-70:

Human Proinsulin 1-70:
(SEQ ID NO: 6)
SHLVEALYLVCGERGFFYTPKTRREAEDLQVGQVELGGGPGAGSLQPLA
LEGSLQKRGIVEQ;

Human Proinsulin 9-38:

(SEQ ID NO: 7)
SHLVEALYLVCGERGFFYTPKTRREAEDLQ;

Human Proinsulin 1-38:

(SEQ ID NO: 8)
FVNQHLGSHLVEALYLVCGERGFFYTPKTRREAEDLQ;

Human Proinsulin 9-23:

(SEQ ID NO: 9)
SHLVEALYLVCGERG;

Human Proinsulin 45-71 (C13-A6):

(SEQ ID NO: 10)
GGGPGAGSLQPLALEGSLQKRGIVEQC;

Human Proinsulin C24-A1:

(SEQ ID NO: 11)
LALEGSLQKRG;

Human Proinsulin C19-A3:

(SEQ ID NO: 12)
GSLQPLALEGSLQKRGIV;

Human Proinsulin C13-32:

(SEQ ID NO: 13)
GGGPGAGSLQPLALEGSLQK;

Human Proinsulin B9-C4:

(SEQ ID NO: 14)
SHLVEALYLVCGERGFFYTPKTRREAED;

Human Proinsulin C22-A5:

(SEQ ID NO: 15)
QPLALEGSLQKRGIVEQ;

Human IA-2 718-782:

(SEQ ID NO: 16)
AYQAEPTNCATAQEGEKNKRNHPDPLPYDHARIKLKVESPSRSDYIN
ASPIIEHDPMPAYIA;

Human IA-2 785-819:

(SEQ ID NO: 17)
GPLSHTIADFQWQMVESGCTVIVMLTPLVEDGVKQ;

Human IA-2 828-883:

(SEQ ID NO: 18)
GASLYHVYEVNLVSEHIWCEDFLVRSFYLNKVTQETRTLQTFHFLSWP
AEGTPAS;

Human IA-2 943-979:

(SEQ ID NO: 19)
EHVRDQRPGLVRSKQDFEFALTAVAEVNAIKALPQCG.

In autoimmune diseases of the thyroid, including
Hashimoto's thyroiditis and Graves' disease, main antigens

72

include, but are not limited to, thyroglobulin (TG), thyroid
peroxidase (TPO) and thyrotropin receptor (TSHR); other
antigens include sodium iodine symporter (NIS) and mega-
lin. In thyroid-associated ophthalmopathy and dermopathy,
in addition to thyroid autoantigens including TSHR, an
antigen is insulin-like growth factor 1 receptor. In hypopara-
thyroidism, a main antigen is calcium sensitive receptor.

In Addison's Disease, main antigens include, but are not
limited to, 21-hydroxylase, 17 α -hydroxylase, and P450 side
chain cleavage enzyme (P450_{scc}); other antigens include
ACTH receptor, P450_{c21} and P450_{c17}.

In premature ovarian failure, main antigens include, but
are not limited to, FSH receptor and α -enolase.

In autoimmune hypophysitis, or pituitary autoimmune
disease, main antigens include, but are not limited to,
pituitary gland-specific protein factor (PGSF) 1a and 2;
another antigen is type 2 iodothyronine deiodinase.

In multiple sclerosis, main antigens include, but are not
limited to, myelin basic protein ("MBP"), myelin oligoden-
drocyte glycoprotein ("MOG") and myelin proteolipid pro-
tein ("PLP").

MBP, including an exogenously obtained form useful in
the compositions of the disclosure, has the following
sequence (UNIPROT P02686):

(SEQ ID NO: 20)
MGNHAGKRELNAEKASTNSETNRGESEKKRNLGELSRTTSEDNEVFGEA

DANQNNGTSSQDTAVTDSKRTADPKNAWQDAHPADPGSRPHLIRLFSRD

APGREDNTFKDRPSEDELQTIQEDSAATESLDMASQKRPSQRHGSK

YLATASTMDHARHGFLPRHRDTGILDSIGRFFGGDRGAPKRGSGKDSHH

PARTAHYGLSLPQKSHGRTQDENPVVHFFKNIVTPRTPPPSGQKGRGLSL

SRFSWGAEGQRPFGYGGGRASDYKSAHKGFKGVDQAQGTLSKIFKLGGRD

SRSGSPMARR.

MOG, including an exogenously obtained form useful in
the compositions of the disclosure, has the following
sequence (UNIPROT Q16653):

(SEQ ID NO: 21)
MASLSRPSLPSCLCSFLLLLLLQVSSSYAGQFRVIGPRHPHIALVGVDEV

ELPCRISPGKNATGMEVGWYRPPFSRVVHLYRNGKDQDQDAPEYRGRT

ELLKDAIGEGKVTLRIRNVRFSDGGFTCFRDRHSYQEEAAAMELKVEDP

FYWVSPGVLVLLAVLPVLLLQITVGLIFLCLQYRLRGLKRAEIEINLHRT

FDPHFLRVPCWKITLFVIVPVLGPLVALIICYNWLHRRLAGQFLEELRN

PF.

PLP, including an exogenously obtained form useful in
the compositions of the disclosure, has the following
sequence (UNIPROT P60201):

(SEQ ID NO: 22)
MGLLECCARCLVGAPFASLVATGLCFFGVALFCGCGHEALTGTCKLIET

YFSKNYQDYEYLINVIHAFQYVIYGTASFFFLYGALLAEGFYTTGAVR

QIFGDYKTTICGKLSATVTGGQKGRSGRQHQHQSLSLERVCHCLGKWL

HPDKFVGITYALTVVWLLVFACSAVPVYIYFNTWTTCQSIAFPKTSAS

IGSLCADARMYGVLFPWNAPFGKVCNSLLSICKTAEFQMTFHLFIAAFV

GAAATLVSLTFFMIAATYNFAVLKLMGRGTFK.

Peptides/epitopes useful in the compositions of the disclosure for treating multiple sclerosis include some or all of the following sequences, individually in a tolerogenic composition as disclosed herein or together in a combination (e.g., a cocktail) of tolerogenic compositions:

MBP 13-32: (SEQ ID NO: 23)
 KYLATASTMDHARHGFLPRH;
 MBP 83-99: (SEQ ID NO: 24)
 ENPWHEFFKNIVTPRTP;
 MBP 111-129: (SEQ ID NO: 25)
 LSRFSWGAEGQRPFGYGG;
 MBP 146-170: (SEQ ID NO: 26)
 AQGTLSKIFKLGRDSRSGSPMARR;
 MOG 1-20: (SEQ ID NO: 27)
 GQFRVIGPRHPHALVGDEV;
 MOG 35-55: (SEQ ID NO: 28)
 MEVGWYRPPFSRWHLRYNGK;
 PLP 139-154: (SEQ ID NO: 29)
 HCLGKWLGHDPKFVGI;
 MOG 30-60: (SEQ ID NO: 30)
 KNATGMEVGWYRSPFSRVVHLRYNGKDQDAE;
 MBP 83-99: (SEQ ID NO: 31)
 ENPVVHFFKNIVTPRTP;
 MOG 35-55: (SEQ ID NO: 32)
 MEVGWYRPPFSRVVHLRYNGK;
 MBP 82-98: (SEQ ID NO: 33)
 DENPVVHFFKNIVTPRT;
 MBP 82-99: (SEQ ID NO: 34)
 DENPVVHFFKNIVTPRTP;
 MBP 82-106: (SEQ ID NO: 35)
 DENPVVHFFKNIVTPRTPPPSQGKG;
 MBP 87-106: (SEQ ID NO: 36)
 VHFFKNIVTPRTPPPSQGKG;
 MBP 131-155: (SEQ ID NO: 37)
 ASDYKSAHKLKGVDAQGTLSKIFK;
 PLP 41-58: (SEQ ID NO: 38)
 GTEKLIETYFSKNYQDYE;
 PLP 89-106: (SEQ ID NO: 39)
 GFYTTGAVRQIFGDYKTT;
 PLP 95-116: (SEQ ID NO: 40)
 AVRQIFGDYKTTICGKGLSATV;

-continued

PLP 178-197: (SEQ ID NO: 41)
 NTWTTCQSIAPFSKTSASIG;
 5 PLP 190-209: (SEQ ID NO: 42)
 SKTSASIGSLCADARMYGVL;
 10 MOG 11-30: (SEQ ID NO: 43)
 PIRALVGDEVELPCRISPGK;
 MOG 21-40: (SEQ ID NO: 44)
 ELPCRISPGKNATGMEVGWY;
 15 MOG 64-86: (SEQ ID NO: 45)
 EYRGRTTELLKDAIGEGKVTLRIR;
 20 MOG 1-62: (SEQ ID NO: 46)
 GQFRVIGPRHPHALVGDEVELPCRISPGKNATGMEVGWYRPPFSRVVH
 LYRNGKDQGDQA
 MBP 76-136: (SEQ ID NO: 47)
 25 SHGRTQDENPVVHFFKNIVTPRTPPPSQGKGRGLSLSRFSWGAEGQRP
 FGYGGRASDYKSCG
 In rheumatoid arthritis, main antigens include, but are not limited to, collagen II, immunoglobulin binding protein, the fragment crystallizable region of immunoglobulin G, double-stranded DNA, and the natural and cirtullinated forms of proteins implicated in rheumatoid arthritis pathology, including fibrin/fibrinogen, vimentin, collagen I and II, and alpha-enolase.
 30 In autoimmune gastritis, a main antigen is H+, K+-AT-Pase.
 In pernicious angemis, a main antigen is intrinsic factor.
 40 In celiac disease, main antigens include, but are not limited to, tissue transglutaminase and the natural and deamidated forms of gluten or gluten-like proteins, such as alpha-, gamma-, and omega-gliadin, glutenin, hordein, secalin, and avenin. Those skilled in the art will appreciate, for example, that while the main antigen of celiac disease is alpha gliadin, alpha gliadin turns more immunogenic in the body through deamidation by tissue glutaminase converting alpha gliadin's glutamines to glutamic acid. Thus, while alpha gliadin is originally a foreign food antigen, once it has been modified in the body to become more immunogenic it can be characterized as a self-antigen, depending on the embodiment.
 45 In vitiligo, a main antigen is tyrosinase, and tyrosinase related protein 1 and 2.
 50 MART1, Melanoma antigen recognized by T cells 1, Melan-A, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT Q16655):
 55 (SEQ ID NO: 48)
 MPREDAHFYIGYPKKGHGHSYTTAEAAAGIGILTIVILGVLIIICWYCR
 60 RRNGYRALMDKSLHVGTCALTRRCPPQEGFDHRDSKVSLEKNCPEVPV
 65 NAPPAYEKLSEQSPPPYSP.

75

Tyrosinase, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT P14679):

(SEQ ID NO: 49)

```

MLLAVLYCLLWSFQTSAGHFPRACVSSKNLMEKECCPPWSGDRSPCGQL
SGRGSCQNILLSNAPLGPQFPFTGVDDRESWPSVFYNRTCQCSGNFMGF
NCGNCKFGFWGPNCTERRLLVRRNIFDLSAFEKDKFFAYLTAKHTISS
DYVIPIGTYGQMKNGSTPMENDINIYDLFVWMHYVYVSMALLGGSEIWR
DIDFAHEAPAFLPWHRLFLLRWEQEIQKLTGDNFTIPYDWRDAEKCD
ICTDEYMGQGHPTNPNNLSPASFFSSWQIVCSRLEEYNSHQSLCNGTPE
GPLRRNPNGNHDKSRTPLRPSSADVEFCLSLTQYESGSMKKAANFSFRNT
LEGFASPLTGIADASQSSMHNLHIYMGNTMSQVQGSANDPIFLHHAF
VDSIFEQWLRHRRLPEQVEYPEANAPIGHNRESYMPFIPLYRNGDFFIS
SKDLGYDYSYLQSDPDSPFQDIKSYLEQASRIWSWLLGAAMVGAULTA
LLAGLVSLLCRHKRKLPEEKQPLLMKEDYHSLYQSHL
  
```

Melanocyte protein PMEL, gp100, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT P40967):

(SEQ ID NO: 50)

```

MDLVLRCLLHLAVIGALLAVGATKVPNRQDLGVSRLRTKAWNRQLY
PEWTEAQRDLDCWRGGQVSLKVSNDGPTLIGANASFSIALNFPQSQKVL
DQGVIVWNNTIINGSQVWGQPVYPQETDDACIFPDGGPCPSGWSQKR
SFVYVWKTGWQYVQVLGGPVSGLSIGTGRAMLGTHTEMTVYHRRGSR
YVPLAHSSSAFTITDQVPFVSQVSLRALDGGNKHFLRNQPLTFALQLH
DPSGLAEADLSYTWDFDSSGTLISRALVVTHTYLEPGPVTAQVVLQA
AIPLTSCGSSPVP GTTDGHRPTAEAPNTTAGQVPTTEVGTTPGQAPTA
EPSGTTSVQVPTTEVISTAPVQMPTAESTGMTPEKVPVSEVMGTTLAEM
STPEATGMTPAEVSIVVLSGTTAAQVTTTEWVETTARELPPIPEPEGPDA
SSIMSTESITGSLGPLLDGTATLRLVKRQVPLDCVLRYGSPSVTLDIV
QGIESAEILQAVPSGEGDAFELTVSCQGGPKKEACMEISSPGCQPPAQR
LCQPVLPSACQLVLHQLKGGSGTYCLNVSLADTNSLAVVSTQLIMPG
QEAGLGQVPLIVGILLVMAVVLASLIYRRRLMKQDFSPVQLPHSSSHW
LRLPRIFCSCPIGENSPLLSGQOV.
  
```

In myasthenia gravis, a main antigen is acetylcholine receptor.

In pemphigus vulgaris and variants, main antigens include, but are not limited to, desmoglein 3, 1 and 4; other antigens include pemphaxin, desmocollins, plakoglobin, periplakin, desmoplakins, and acetylcholine receptor.

In bullous pemphigoid, main antigens include BP180 and BP230; other antigens include plectin and laminin 5.

In dermatitis herpetiformis Duhring, main antigens include, but are not limited to, endomysium and tissue transglutaminase.

In epidermolysis bullosa acquisita, a main antigen is collagen VII.

In systemic sclerosis, main antigens include, but are not limited to, matrix metalloproteinase 1 and 3, the collagen-

76

specific molecular chaperone heat-shock protein 47, fibrillin-1, and PDGF receptor; other antigens include Scl-70, U1 RNP, Th/To, Ku, Jo1, NAG-2, centromere proteins, topoisomerase I, nucleolar proteins, RNA polymerase I, II and III, PM-Slc, fibrillarin, and B23.

In mixed connective tissue disease, a main antigen is U1snRNP.

In Sjogren's syndrome, the main antigens include, but are not limited to, nuclear antigens SS-A and SS-B; other antigens include fodrin, poly(ADP-ribose) polymerase and topoisomerase, muscarinic receptors, and the Fc-gamma receptor IIIb.

In systemic lupus erythematosus, main antigens include nuclear proteins including the "Smith antigen," SS-A, high mobility group box 1 (HMGB1), nucleosomes, histone proteins and double-stranded DNA (against which auto-antibodies are made in the disease process).

In Goodpasture's syndrome, main antigens include, but are not limited to, glomerular basement membrane proteins including collagen IV.

In rheumatic heart disease, a main antigen is cardiac myosin.

In autoimmune polyendocrine syndrome type 1 antigens include aromatic L-amino acid decarboxylase, histidine decarboxylase, cysteine sulfinic acid decarboxylase, tryptophan hydroxylase, tyrosine hydroxylase, phenylalanine hydroxylase, hepatic P450 cytochromes P4501A2 and 2A6, SOX-9, SOX-10, calcium-sensing receptor protein, and the type 1 interferons interferon alpha, beta and omega.

In neuromyelitis optica, a main antigen is AQP4.

Aquaporin-4, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT P55087):

(SEQ ID NO: 51)

```

MSDRPTARRWGKCGPLCTRENIMVAFKGVWTAQAFKAVTAEFMLLIFV
LLSLGSTINWGGTEKPLPVDMLISLCFGLSIATMVQCFGHISGGHINP
AVTVAMVCTRKISIAKSVFYIAAQCLGAIIGAGILYLVTPPSVVGGLGV
TMVHGNLTAGHGLLVELIITPQLVFTIFASCDSKRTDVTGSIALAIGFS
VAIGHLFAINYTGASMNPARSFGPAVIMGNWENHWIYVWGPPIIGAVLAG
GLYEVVFCPDVEFKRRFKEAFSKAAQQTGKSYMEVEDNRSQVETDDLIL
KPGVVHVIDVDRGEEKKGDQSGEVLSSV.
  
```

In uveitis, main antigens include Retinal S-antigen or "S-arrestin" and interphotoreceptor retinoid binding protein (IRBP) or retinol-binding protein 3.

S-arrestin, including an exogenously obtained form useful in the compositions of the disclosure, has the following sequence (UNIPROT P10523):

(SEQ ID NO: 52)

```

MAASGKTSKSEPNHVIFFKKISRDKSVTIYLGNDYIDHVSQVQVVDGTV
LVDPDLVKGKVVYVTLTCAFRYGQEDIDVIGLTFRRDLYFSRVQVYPPV
GAASTPTKLQESLLKGLSNTYFPLLTFPDYLPSCSVMLQPAPQDSGKSC
GVDFEVKAPATDSTDAEEDKIPKKSSVRLLRKQVHAPLEMGPQPRABE
AWQFMSDKPLHLAVSLNKEIYFHGEPITVTVTNTEKTVKKIKAFV
EQVANVVLVYSSDYVVKPVAMEEAQEKVPPNSTLTTLTLLPLANNRER
RGIALDGKIKHEDTNLASSIIKEGIDRTVLGLVSYQIKVKLTVSGFL
  
```

In the embodiments where the antigen is a foreign antigen against which an unwanted immune response is developed, such as to animal, plant and environmental antigens, specific antigens can, for example, be: cat, mouse, dog, horse, bee, dust, tree and goldenrod, including the following proteins or peptides derived from: weeds, (including ragweed allergens amb a 1, 2, 3, 5, and 6, and Amb t 5; pigweed Che a 2 and 5; and other weed allergens Par j 1, 2, and 3, and Par o 1); grass (including major allergens Cyn d 1, 7, and 12; Dac g 1, 2, and 5; Hol i 1.01203; Lol p 1, 2, 3, 5, and 11; Mer a 1; Pha a 1; Poa p 1 and 5); pollen from ragweed and other weeds (including curly dock, lambs quarters, pigweed, plantain, sheep sorrel, and sagebrush), grass (including Bermuda, Johnson, Kentucky, Orchard, Sweet vernal, and Timothy grass), and trees (including *catalpa*, elm, hickory, olive, pecan, sycamore, and walnut); dust (including major allergens from species *Dermatophagoides pteronyssinus*, such as Der p 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 14, 15, 18, 20, 21, and 23; from species *Dermatophagoides farina*, such as Der f 1, 2, 3, 6, 7, 10, 11, 13, 14, 15, 16, 18, 22, and 24; from species *Blomia tropicalis* such as Blo t 1, 2, 3, 4, 5, 6, 10, 11, 12, 13, 19, and 21; also allergens Eur m 2 from *Euroglyphus maynei*, Tyr p 13 from *Tyrophagus putrescentiae*, and allergens Bla g 1, 2, and 4; Per a 1, 3, and 7 from cockroach); pets (including cats, dogs, rodents, and farm animals; major cat allergens include Fel d 1 through 8, cat IgA, Bla g 2, and cat albumin; major dog allergens include Can f 1 through 6, and dog albumin); bee stings, including major allergens Api m 1 through 12; and fungus, including allergens derived from, species of *Aspergillus* and *Penicillium*, as well as the species *Alternaria alternata*, *Davidiella tassiana*, and *Trichophyton rubrum*.

79

In Parkinson's disease, the main antigen is alpha synuclein. Alpha synuclein, including an exogenously obtained form useful in the tolerogenic compositions of the disclosure, has the following sequence (UNIPROT P37840):

(SEQ ID NO: 62)
MDVFMKGLSKAKEGVVAAAEKTKQGVAAAGKTKEGVLYVGSKTKEGVV
HGVATVAEKTKEQVTNVGGAVVTGVTAVAQKTVEGAGSIAAATGFVKKD
QLGKNEEGAPQEGILEDMPVDPNEAYEMPSEEGYQDYEPKA.

The antigen can be a complete protein, a portion of a complete protein, a peptide, or the like, and can be derivatized (as discussed above) for attachment to a linker and/or mannosylating moiety, can be a variant and/or can contain conservative substitutions, particularly maintaining sequence identity, and/or can be desilylated.

F. EXAMPLES

The following examples are included to demonstrate non-limiting embodiments of the invention disclosed herein. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the embodiments of the invention. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the embodiments of the invention disclosed herein.

Example 1

Polymer and Conjugate Synthesis

All reactions were carried out at room temperature unless specified. Unless otherwise stated, chemicals were reagent grade and purchased from Sigma-Aldrich (Saint Louis, Mo., USA). Size exclusion chromatography was carried out on an ÄKTA protein purification system (General Electric Healthcare Lifesciences), using a Superdex 200 10/300 column (General Electric Healthcare Lifesciences). All NMR spectra were collected on a Bruker Avance-II 400 MHz NMR, unless otherwise noted, and NMR spectra were analyzed with MnovaNMR (Mestrelab). High pressure size exclusion chromatography was performed on a Dionex Ultimate 3000 UHPLC (Thermo Fisher Scientific). Gels were imaged using a Biorad Universal Hood Gel Doc 2000 System (Biorad). Antigens conjugated to polymers as disclosed in the Examples are non-limiting examples of tolerogenic antigens according to embodiments disclosed herein, including embodiments using immunogenic fragments of antigens.


Compound 1: 1-(2-chloroethyl)- α -D-mannose

Acetyl chloride (4.35 mL, 61.05 mmol) was added dropwise to the ice-cold solution of D-mannose (10.0 g, 55.51 mmol) in chloroethanol (40 mL, 413.68 mmol). The mixture was stirred for 15 minutes at 4° C. and then was transferred to the oil bath at 70° C. The reaction was then stirred for 4 h. After cooling to room temperature, a dark brown solution was poured into a 400 mL solution of ethyl acetate and DCM (3:1, v/v) in order to remove excess chloroethanol. The mixture was placed at -20° C. for 30 minutes and then a dark brown sticky precipitate was collected from the supernatant.

80


The precipitate was dissolved in anhydrous ethanol and 3 spoons of activated charcoal was added. The suspension was mixed for 1.5 h and then filtered through Celite and washed with ethanol. In the last step, ethanol was evaporated via rotary evaporation to provide 12.8 g (95.24%) of product: $C_8H_{15}ClO_6$, ESI-MS $[M+Na]^+_{theor}=m/z$ 265.0455, $[M+Na]^+_{found}=m/z$ 265.0458; 1H NMR (400 MHz, D_2O) δ 4.86 (s, C1, 1H), 3.92 (dd, C2, 1H), 3.62-3.85 (multiple signals from 7H: C3, C4, C5, C6, C6', ethoxy group); ^{13}C NMR (100 MHz, D_2O) δ 99.84, 76.32, 72.94, 70.52, 69.94, 69.65, 67.76, 66.75, 60.96, 43.39.

Compound 2: 1-(2-azidoethyl)- α -D-mannose




Compound 1 (12.7 g, 52.48 mmol) was dissolved in 15 mL of N,N-dimethylformamide. To that solution, sodium azide was added (5.0 g, 76.92 mmol) and the suspension was placed in an oil bath and stirred over night at 90° C. After 16 h, the reaction mixture was filtered through Celite and the solvent was then removed via rotary evaporation to provide a oily, brown substance. The residual was adsorbed on silica gel and purified using flash chromatography (DCM:MeOH 92:8, v/v) to yield 5.6 g (42.86%) of pure product: $C_8H_{15}N_3O_6$, ESI-MS $[M+Na]^+_{theor}=m/z$ 272.2578, $[M+Na]^+_{found}=m/z$ 272.0850; 1H NMR (400 MHz, D_2O) δ 4.84 (s, C1, 1H), 3.91 (dd, C2, 1H), 3.39-3.87 (multiple signals from 7H: C3, C4, C5, C6, C6', ethoxy group); ^{13}C NMR (100 MHz, D_2O) δ 99.85, 72.94, 70.44, 69.98, 66.73, 66.34, 50.24.

Compound 3: 1-(2-aminoethyl)- α -D-mannose





81



A suspension of 2 (5.5 g, 22.09 mmol) and 10% palladium on carbon (one spoon, ca. 500 mg) in 10 mL of ethanol was hydrogenated in a Shlenk flask with an initial pressure of 2 bars of hydrogen gas. The reduction process was monitored by TLC. After 3 h reaction was completed and the suspension was filtered through Celite. The solvent was evaporated in vacuo to give 4.9 g (99.48%) of product: C₈H₁₇NO₆, ESI-MS [M+Na]⁺+theor=m/z 246.0954, [M+Na]⁺+found=m/z 246.0955; ¹H NMR (400 MHz, D₂O) δ 4.80 (s, C1, 1H), 3.89 (dd, C2, 1H), 3.83-3.44 (multiple signals from 7H: C3, C4, C5, C6, C6', ethoxy group); ¹³C NMR (100 MHz, D₂O) δ 99.87, 72.74, 70.55, 70.03, 68.82, 66.81, 60.97, 39.94.

Compound 4: N-[2-(α-D-mannose)ethyl] methacrylamide





Compound 3 (4.5 g, 20.17 mmol) was dissolved in 10 mL of N,N-dimethylformamide. To that solution, triethylamine (3 mL, 22.28 mmol) was added and the mixture was cooled down to 4° C. Subsequently, pentafluorophenyl methacrylate (4.38 mL, 24.21 mmol) was added drop-wise with constant stirring. After 30 minutes, ice-bath was removed

82


and the reaction was allowed to stir at room temperature for the next 4 h. Next, the solvent was evaporated and the residual was adsorbed on silica gel. The purification of crude material using flash chromatography (DCM:MeOH 95:5, v/v) provided 3.8 g (64.73%) of mannose monomer: C₁₂H₂₁NO₇, ESI-MS [M+Na]⁺+theor=m/z 314.1216, [M+Na]⁺+found=m/z 314.1208; ¹H NMR (400 MHz, D₂O) δ 5.6 (s, 1H), 5.38 (s, 1H), 4.78 (s, 1H), 3.84 (s, C2, 1H), 3.77-3.34 (multiple signals from 7H: C3, C4, C5, C6, C6', ethoxy group), 1.85 (s, 3H); ¹³C NMR (100 MHz, D₂O) δ 172.06, 139.06, 121.00, 99.63, 72.78, 70.47, 69.99, 66.58, 65.73, 60.78, 39.04, 17.68.

Compound 17: 2-(2-(2-(2-hydroxyethoxy)ethoxy)ethoxy)ethyl 4-methylbenzenesulfonate




Tetraethylene glycol (2.5 g, 12.9 mmol) and pyridine (1.0 g, 12.6 mmol) were added to 50 mL of DCM and stirred for 20 minutes at 0° C. To that solution, p-toluenesulfonyl chloride (2.37 g, 10 mmol) in 15 mL of DCM was added slowly. The reaction mixture was then stirred for 2 h at 0° C. followed by 4 h at room temperature. After that time, the solvent was evaporated and crude product was purified via flash chromatography (ethyl acetate:hexane 6:4, v/v). The final yield was 1.95 g (72.22%): C₁₅H₂₄O₇S, ESI-MS [M+H]⁺+theor=m/z 349.1321, [M+H]⁺+found=m/z 349.1325; ¹H NMR (400 MHz, CDCl₃) δ 7.68 (d, 2H), 7.24 (d, 2H), 4.00-4.12 (m, 2H), 3.42-3.70 (m, 14H), 2.89 (t, 1H), 2.33 (s, 3H); ¹³C NMR (100 MHz, DMSO-d₆) δ 144.68, 132.76, 129.68, 127.73, 72.34, 70.46, 70.41, 70.22, 70.10, 69.17, 68.44, 61.40, 21.41.

83




Sodium azide (1.5 g, 23.1 mmol) was added to a solution of 17 (1.5 g, 4.3 mmol) in N,N-dimethylformamide (75 mL) at room temperature. The reaction mixture was stirred overnight at 90° C. The reaction was then filtered and the solvent was removed in vacuo. The resulting viscous liquid was then purified by flash column chromatography (ethyl acetate:hexane 6:4, v/v) to yield a pure product (1.25 g, 83%): C₈H₁₇N₃O₄, ESI-MS [M+Na]⁺+theor=m/z 242.1117, [M+Na]⁺+found=m/z 242.1171; ¹H NMR (400 MHz, DMSO-d₆) δ 3.49-3.65 (m, 14H), 3.30 (t, 2H), 2.91 (t, 1H); ¹³C NMR (100 MHz, DMSO-d₆) δ 72.43, 70.56, 70.52, 70.46, 70.21, 69.91, 61.51, 50.54.



4-Cyano-4-[(dodecylsulfanylthiocarbonyl)sulfanyl]pentanoic acid (500 mg, 1.24 mmol), compound 18 (226 mg, 1.03 mmol) and DCC (255 mg, 1.24 mmol) were dissolved in 4 mL of DCM. The solution was placed in an ice bath and let stir for 30 minutes. After that time, 4-dimethylaminopyridine (12.6 mg, 0.10 mmol) dissolved in 1 mL of DCM was added drop-wise. The solution was stirred at 0° C. for 2 h and then at room temperature for 1 hour. Once reaction was completed, DCM was evaporated in vacuum. The crude product was purified via column chromatography (DCM: ethyl acetate 97:3, v/v) to yield 300 mg (88.57%) of the pure product: C₂₇H₄₈N₄O₅S₃, ESI-MS [M+H]⁺+theor=m/z 604.2865 [M+H]⁺+found=m/z 604.2862; ¹H NMR (400 MHz, CDCl₃) δ 4.26 (t, 2H), 3.76-3.65 (m, 12H), 3.39 (t, 2H), 3.33 (t, 2H), 2.66 (dd, 2H), 2.53 (m, 1H), 2.38 (m, 1H), 1.88 (s, 3H), 1.69 (dt, 2H), 1.38 (m, 2H), 1.26 (s, 16H), 0.88 (t, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 216.92, 171.46, 119.00, 70.72, 70.68, 70.61, 70.07, 68.96, 64.16, 50.69, 46.35, 37.06, 33.81, 31.91, 29.71, 29.62, 29.54, 29.42, 29.34, 29.07, 28.93, 27.68, 24.84, 22.69, 14.13.



84

Compound 21: p (Man)



The mannose monomer (4) (150.0 mg, 0.51 mmol) and (20) (130.0 mg, 1.03 mmol) were dissolved in 600 µl of anhydrous DMF and added to a schlenk tube. Then, RAFT agent (19) (18.5 mg, 0.03 mmol) in 100 µl of DMF and AIBN (1 mg, 0.006 mmol) in 10 µl of DMF were added to the schlenk tube. The tube was degassed via four freeze-pump-thaw cycles and then immersed in an oil bath preheated at 70° C. to initiate polymerization. The reaction was left stirring for 14 hours. After that time, the polymer was precipitated by transferring its viscous solution into 20 mL of cold acetone. The light-yellow suspension was placed in the freezer for 30 minutes. The precipitate was then centrifuged and re-suspended in fresh acetone. The processed was repeated 3 times. In the final step, the resultant glycopolymers (150 mg, 47.6%) was dried in a vacuum oven at reduced pressure and characterized by means of ¹H NMR and GPC. The p(HPMA-TLR7) used in the biological studies had a number average molecular weight of 15,425 Da, as determined by size exclusion chromatography, using a dextran standard, a degree of polymerization of 82.1, and were composed of a 1:2.25 molar ratio of Mannose:HPMA, as determined by ¹H NMR.



Compound 25: Ethanol disulfanyl polyethylene glycol amine



A solution of thiol polyethylene glycol amine (JenKem Technology, USA) (1.0 g, 0.5 mmol) in DCM (5 ml) was added dropwise to a stirred solution of 2-(2-pyridinyldithio) ethanol (24) (467.5 mg, 2.5 mmol) in MeOH (3 ml). The solution was stirred at room temperature for 10 h then approximately half the solvent was removed via rotary evaporation. The remaining crude product was then decanted into ice cold hexanes (40 ml) and placed at -20° C.

for 4 h. The precipitate and solvent mixture was centrifuged at 2000 g for 3 min. The solvent was then decanted and excess solvent was removed from the pelleted precipitate under reduced pressure. The crude product was then used in the next step without further purification (65% crude yield). The final structure was characterized by ¹H NMR and reverse phase chromatography.

Compound 26: Ethanol disulfanyl polyethylene glycol (1R,8S,9s)-Bicyclo[6.1.0]non-4-yn-9-ylmethyl carbamate




87

A solution of (1R,8S,9s)-Bicyclo[6.1.0]non-4-yn-9-ylmethyl N-succinimidyl carbonate (90 mg, 0.30 mmol) in DCM (0.5 ml) was added dropwise to an ice-cooled stirred solution of ethanol disulfanyl polyethylene glycol amine 25 (0.5 g, 0.24 mmol) and trimethylamine (48 mg, 0.48 mmol) in DCM (5 ml). After the addition of (1R,8S,9s)-Bicyclo[6.1.0]non-4-yn-9-ylmethyl N-succinimidyl carbonate, the reaction was allowed to come to room temperature and stirred for another 6 h. The reaction mixture was then poured

88


into ice-cold hexanes (40 ml) and placed at -20°C . for 4 h. The precipitate and solvent mixture was centrifuged at 2000 g for 3 min. The solvent was then decanted and excess solvent was removed from the pelleted precipitate under reduced pressure. The crude product was then used in the next step without further purification (75% crude yield). The final structure was characterized by ^1H NMR and reverse phase chromatography.

Compound 27: N-succinimidyl carboamate Ethanol disulfanyl polyethylene glycol (1R,8S,9s)-Bicyclo[6.1.0]non-4-yn-9-ylmethyl carbamate (Self-immolative Linker)




A solution of Ethanol disulfanyl polyethylene glycol (1R,8S,9s)-Bicyclo[6.1.0]non-4-yn-9-ylmethyl carbamate (300 mg, 0.13 mmol) in anhydrous acetonitrile (ACN) (1.5 ml) was added dropwise to a stirred solution of N,N'-Disuccinimidyl carbonate 26 (0.5 g, 0.24 mmol) and trimethylamine (48 mg, 0.48 mmol) in anhydrous ACN (5 ml). The reaction mixture was stirred overnight and was then poured into ice-cold hexanes (40 ml) and placed at -20°C . for 4 h. The precipitate and solvent mixture was centrifuged at 2000 g for 3 min. The solvent was then decanted and excess solvent was removed from the pelleted precipitate under reduced pressure. The crude product was purified via silica gel flash chromatography through a thin pad of silica DCM:MeOH (85:15) (yield: 43%, 129 mg). The final structure was characterized by ^1H NMR and reverse phase chromatography.

Compound 28: General procedure for OVA-, asparaginase-self-immolative linker conjugates




EndoGrade® Ovalbumin (OVA) (Hyglos) (10 mg, 222.2 nmol), recombinant asparaginase (7.2 mg, 222.2 nmol) and self-immolative Linker (27) (5 mg) were added to an endotoxin free tube. Phosphate buffer at pH 7.7 (200 μ L) was added to the tube and the tube was stirred at 1 h at room temperature. The reaction mixture was then filtered (0.22 μ M) and the conjugates were purified via Zeba Spin Desalting Columns with a 30 kDa cutoff limit (Thermo Fisher). Chemical conjugation was verified via gel electrophoresis and high pressure size exclusion chromatography.



Compound 28 in PBS, prepared as described above, was added to an endotoxin free tube and p(Man) (21) (30 mg) was added and the reaction was stirred for 30 min at room temperature. The reaction mixture was then filtered (0.22 μ M) and the final product was purified via size exclusion chromatography. Chemical conjugation was verified via gel electrophoresis and high pressure size exclusion chromatography.

91

92



30

Example 2

OTI/OTII Challenge to Tolerance Model

BLK6 mice were treated with saline, or 10 µg of OVA (as a non-limiting example of an immunogenic antigen) in the form of free OVA or OVA conjugated to p(Man) (OVA-p(Man)) one day and 7 days after an adoptive transfer of 7.0×10^5 OTI and OTII T cells. These mice were challenged with an intradermal injection of LPS and OVA 14 days after the initial OTI and OTII T cell transfer and then the immune response in the draining lymph nodes (dLNs) was assessed on day 19 and compared to mice that were treated with saline, but did not receive the challenge of LPS and OVA (No-challenge).

Profound tolerance was induced in the CD4⁺ T cell compartment, as shown in FIGS. 1-2. In terms of total cell frequencies, both dosing regimens of both OVA-p(Man) resulted in equivalent low levels of OTII cells after challenge, statistically lower than by treatment of OVA (* indicates $p < 0.05$, ** indicates $p < 0.01$), as shown in FIG. 1. When the cells that remained were analyzed by flow cytometry for the presence of the transcription factor FoxP3 and the receptor CD25, the numbers of FoxP3⁺CD25⁺ cells (markers of T regulatory cells) was statistically significantly elevated compared to treatment with OVA alone, as shown in FIG. 1. Additionally, the spleens of animals treated with OVA-p(Man) contained a lower percentage of T follicular helper cells (Tfh) as compared to other groups that were challenged with LPS and OVA. When the cells that remained were analyzed by flow cytometry for the expression of IFN-γ after exposure to OVA antigen, the frequency of CD4⁺ T cells expressing this inflammatory cytokine was decreased in the groups receiving OVA-p(Man), as shown in FIG. 2.

Profound tolerance was also induced in the CD8⁺ T cell compartment, as shown in FIG. 1. In terms of total cell

frequencies, OVA-p(Man) resulted in equivalent low levels of OTI cells after challenge, statistically lower than by treatment of OVA (* indicates $p < 0.05$, ** and indicates $p < 0.01$), as shown in FIG. 1. When the cells that remained were analyzed by flow cytometry for the expression of IFN-γ after exposure to SIINFELKE antigen, the frequency of CD8⁺ T cells expressing this inflammatory cytokine was decreased in the groups receiving OVA-p(Man), as shown in FIG. 2.

Example 3

Tolerance Induction to Intravenously Administered Asparaginase

Five BALB/c mice per group were injected with 2.5 µg of asparaginase (as a non-limiting example of an immunogenic antigen) formulated as free asparaginase (ASNase) or conjugated to p(Man) (ASNase-p(Man)) once a week for 3 weeks and then, at week 4, were switched to 15 µg of ASNase i.v. once a week for 8 weeks. During the initial 3 weeks ASNase-p(Man) was administered via either i.v. or subcutaneous injection. Sera was taken from the mice and monitored weekly for the presence of αASNase.

Upon intravenous injection of ASNase at week 4, animals treated with saline and subcutaneously administered ASNase-p(Man) experienced a rapid increase in serum αASNase IgG (FIG. 3), animals treated with i.v. administered ASNase-p(Man) did not incur an increase in serum αASNase IgG for the duration of the experiment. Furthermore, after 38 days of treatment, animals treated with ASNase-p(Man) via intravenous infusion, experienced significantly lower αASNase IgG subclass titers (FIG. 3).

At week 22 (3 weeks of tolerization, 8 weeks of ASNase treatment, and 11 weeks after last dose of Asnase), animals treated with saline and ASNase-p(Man) via intravenous

infusion were sacrificed and the spleens, livers, and bone marrow of these animals was collected and processed into single cell suspensions. The cells from these organs were stimulated *in vitro* with recombinant ASNase lacking activity for 3 days. After three days the cells were analyzed via flow cytometry for the existence of α ASNase producing cells (plasma cells) and IL-10 producing B regulatory cells. The results show that the bone marrow of animals treated with ASNase-p(Man) had a fewer α ASNase plasma cells than animals that were treated with ASNase (FIG. 4). On the contrary, the spleens of animals treated with ASNase-p(Man) had a greater percentage of IL-10 producing B regulatory cells.

Example 4

Effect on Anti-Asparaginase (Anti-ASNase) Humoral Immune Response

To assess the ability of p(Man)-protein conjugates to prevent an anti-asparaginase (anti-ASNase) humoral immune response and thus avoid the loss of efficacy that is the result of rapid antibody-mediated clearance of biological therapeutics from the serum, mice were treated on days -21, -14, and -7 with saline or 2.5 μ g of ASNase conjugated to p(Man) (p(Man)-ASNase) via iv infusion (n=5 animals per group). After 7 days, the mice were then treated with weekly iv infusions of 15 μ g of wt ASNase for 4 weeks. The mice were then treated with 15 μ g of wt ASNase on day 49. On day 70, mice that had been treated with saline on days -21, -14, -7, 0, 14, 21, and 49 were treated with 15 μ g of wt ASNase to assess the efficacy of ASNase in naïve mice (FIG. 6). On day 70, the mice that had received saline or p(Man)-ASNase then treated with wt ASNase were also adminis-

tered 15 μ g of wt ASNase. The serum asparagine concentration of each animal was assessed on days 71, 73, and 76.

The results show that one day after being treated with ASNase, animals in each group have a similar serum concentration of asparagine. However, three days after being administered asparagine on day 7, animals treated with only saline or p(Man)-ASNase have a significantly lower serum asparagine concentration as compared to animals that had been treated with saline and then administered wt ASNase on days 0, 7, 14, 21, and 49 (FIG. 7A). In addition, animals treated with p(Man)-ASNase had significantly lower asparagine serum concentrations on day 76 than the animals in other groups. When serum asparagine concentration is plotted against the anti-ASNase titer for each animal in the study, a strong correlation ($r=0.8$) between serum asparagine concentration and anti-ASNase titer becomes evident (FIG. 7B). These results demonstrate that p(Man)-ASNase inhibits the loss of efficacy associated with an anti-ASNase immune response.

All of the methods and compositions disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the methods and apparatuses and in the steps or in the sequence of steps of the methods described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 62

<210> SEQ ID NO 1

<211> LENGTH: 110

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
Met Ala Leu Trp Met Arg Leu Leu Pro Leu Leu Ala Leu Leu Ala Leu
1           5           10           15
Trp Gly Pro Asp Pro Ala Ala Ala Phe Val Asn Gln His Leu Cys Gly
20          25          30
Ser His Leu Val Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe
35          40          45
Phe Tyr Thr Pro Lys Thr Arg Arg Glu Ala Glu Asp Leu Gln Val Gly
50          55          60
Gln Val Glu Leu Gly Gly Gly Pro Gly Ala Gly Ser Leu Gln Pro Leu
65          70          75          80
Ala Leu Glu Gly Ser Leu Gln Lys Arg Gly Ile Val Glu Gln Cys Cys
85          90          95
Thr Ser Ile Cys Ser Leu Tyr Gln Leu Glu Asn Tyr Cys Asn
100         105         110
```

<210> SEQ ID NO 2

<211> LENGTH: 585

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met Ala Ser Pro Gly Ser Gly Phe Trp Ser Phe Gly Ser Glu Asp Gly
 1          5          10          15
Ser Gly Asp Ser Glu Asn Pro Gly Thr Ala Arg Ala Trp Cys Gln Val
 20          25          30
Ala Gln Lys Phe Thr Gly Gly Ile Gly Asn Lys Leu Cys Ala Leu Leu
 35          40          45
Tyr Gly Asp Ala Glu Lys Pro Ala Glu Ser Gly Gly Ser Gln Pro Pro
 50          55          60
Arg Ala Ala Ala Arg Lys Ala Ala Cys Ala Cys Asp Gln Lys Pro Cys
 65          70          75          80
Ser Cys Ser Lys Val Asp Val Asn Tyr Ala Phe Leu His Ala Thr Asp
 85          90          95
Leu Leu Pro Ala Cys Asp Gly Glu Arg Pro Thr Leu Ala Phe Leu Gln
100          105          110
Asp Val Met Asn Ile Leu Leu Gln Tyr Val Val Lys Ser Phe Asp Arg
115          120          125
Ser Thr Lys Val Ile Asp Phe His Tyr Pro Asn Glu Leu Leu Gln Glu
130          135          140
Tyr Asn Trp Glu Leu Ala Asp Gln Pro Gln Asn Leu Glu Glu Ile Leu
145          150          155          160
Met His Cys Gln Thr Thr Leu Lys Tyr Ala Ile Lys Thr Gly His Pro
165          170          175
Arg Tyr Phe Asn Gln Leu Ser Thr Gly Leu Asp Met Val Gly Leu Ala
180          185          190
Ala Asp Trp Leu Thr Ser Thr Ala Asn Thr Asn Met Phe Thr Tyr Glu
195          200          205
Ile Ala Pro Val Phe Val Leu Leu Glu Tyr Val Thr Leu Lys Lys Met
210          215          220
Arg Glu Ile Ile Gly Trp Pro Gly Gly Ser Gly Asp Gly Ile Phe Ser
225          230          235          240
Pro Gly Gly Ala Ile Ser Asn Met Tyr Ala Met Met Ile Ala Arg Phe
245          250          255
Lys Met Phe Pro Glu Val Lys Glu Lys Gly Met Ala Ala Leu Pro Arg
260          265          270
Leu Ile Ala Phe Thr Ser Glu His Ser His Phe Ser Leu Lys Lys Gly
275          280          285
Ala Ala Ala Leu Gly Ile Gly Thr Asp Ser Val Ile Leu Ile Lys Cys
290          295          300
Asp Glu Arg Gly Lys Met Ile Pro Ser Asp Leu Glu Arg Arg Ile Leu
305          310          315          320
Glu Ala Lys Gln Lys Gly Phe Val Pro Phe Leu Val Ser Ala Thr Ala
325          330          335
Gly Thr Thr Val Tyr Gly Ala Phe Asp Pro Leu Leu Ala Val Ala Asp
340          345          350
Ile Cys Lys Lys Tyr Lys Ile Trp Met His Val Asp Ala Ala Trp Gly
355          360          365
Gly Gly Leu Leu Met Ser Arg Lys His Lys Trp Lys Leu Ser Gly Val
370          375          380
Glu Arg Ala Asn Ser Val Thr Trp Asn Pro His Lys Met Met Gly Val
385          390          395          400

```


-continued

```

Pro Leu Gln Cys Ser Ala Leu Leu Val Arg Glu Glu Gly Leu Met Gln
      405                        410                        415

Asn Cys Asn Gln Met His Ala Ser Tyr Leu Phe Gln Gln Asp Lys His
      420                        425                        430

Tyr Asp Leu Ser Tyr Asp Thr Gly Asp Lys Ala Leu Gln Cys Gly Arg
      435                        440                        445

His Val Asp Val Phe Lys Leu Trp Leu Met Trp Arg Ala Lys Gly Thr
      450                        455                        460

Thr Gly Phe Glu Ala His Val Asp Lys Cys Leu Glu Leu Ala Glu Tyr
      465                        470                        475                        480

Leu Tyr Asn Ile Ile Lys Asn Arg Glu Gly Tyr Glu Met Val Phe Asp
      485                        490                        495

Gly Lys Pro Gln His Thr Asn Val Cys Phe Trp Tyr Ile Pro Pro Ser
      500                        505                        510

Leu Arg Thr Leu Glu Asp Asn Glu Glu Arg Met Ser Arg Leu Ser Lys
      515                        520                        525

Val Ala Pro Val Ile Lys Ala Arg Met Met Glu Tyr Gly Thr Thr Met
      530                        535                        540

Val Ser Tyr Gln Pro Leu Gly Asp Lys Val Asn Phe Phe Arg Met Val
      545                        550                        555                        560

Ile Ser Asn Pro Ala Ala Thr His Gln Asp Ile Asp Phe Leu Ile Glu
      565                        570                        575

Glu Ile Glu Arg Leu Gly Gln Asp Leu
      580                        585

```

<210> SEQ ID NO 3

<211> LENGTH: 355

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

Met Asp Phe Leu His Arg Asn Gly Val Leu Ile Ile Gln His Leu Gln
 1          5          10          15

Lys Asp Tyr Arg Ala Tyr Tyr Thr Phe Leu Asn Phe Met Ser Asn Val
 20          25          30

Gly Asp Pro Arg Asn Ile Phe Phe Ile Tyr Phe Pro Leu Cys Phe Gln
 35          40          45

Phe Asn Gln Thr Val Gly Thr Lys Met Ile Trp Val Ala Val Ile Gly
 50          55          60

Asp Trp Leu Asn Leu Ile Phe Lys Trp Ile Leu Phe Gly His Arg Pro
 65          70          75          80

Tyr Trp Trp Val Gln Glu Thr Gln Ile Tyr Pro Asn His Ser Ser Pro
 85          90          95

Cys Leu Glu Gln Phe Pro Thr Thr Cys Glu Thr Gly Pro Gly Ser Pro
100          105          110

Ser Gly His Ala Met Gly Ala Ser Cys Val Trp Tyr Val Met Val Thr
115          120          125

Ala Ala Leu Ser His Thr Val Cys Gly Met Asp Lys Phe Ser Ile Thr
130          135          140

Leu His Arg Leu Thr Trp Ser Phe Leu Trp Ser Val Phe Trp Leu Ile
145          150          155          160

Gln Ile Ser Val Cys Ile Ser Arg Val Phe Ile Ala Thr His Phe Pro
165          170          175

His Gln Val Ile Leu Gly Val Ile Gly Gly Met Leu Val Ala Glu Ala
180          185          190

```

-continued

Phe Glu His Thr Pro Gly Ile Gln Thr Ala Ser Leu Gly Thr Tyr Leu
 195 200 205
 Lys Thr Asn Leu Phe Leu Phe Leu Phe Ala Val Gly Phe Tyr Leu Leu
 210 215 220
 Leu Arg Val Leu Asn Ile Asp Leu Leu Trp Ser Val Pro Ile Ala Lys
 225 230 235 240
 Lys Trp Cys Ala Asn Pro Asp Trp Ile His Ile Asp Thr Thr Pro Phe
 245 250 255
 Ala Gly Leu Val Arg Asn Leu Gly Val Leu Phe Gly Leu Gly Phe Ala
 260 265 270
 Ile Asn Ser Glu Met Phe Leu Leu Ser Cys Arg Gly Gly Asn Asn Tyr
 275 280 285
 Thr Leu Ser Phe Arg Leu Leu Cys Ala Leu Thr Ser Leu Thr Ile Leu
 290 295 300
 Gln Leu Tyr His Phe Leu Gln Ile Pro Thr His Glu Glu His Leu Phe
 305 310 315 320
 Tyr Val Leu Ser Phe Cys Lys Ser Ala Ser Ile Pro Leu Thr Val Val
 325 330 335
 Ala Phe Ile Pro Tyr Ser Val His Met Leu Met Lys Gln Ser Gly Lys
 340 345 350
 Lys Ser Gln
 355

<210> SEQ ID NO 4
 <211> LENGTH: 86
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val Glu Ala Leu Tyr
 1 5 10 15
 Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro Lys Thr Arg Arg
 20 25 30
 Glu Ala Glu Asp Leu Gln Val Gly Gln Val Glu Leu Gly Gly Gly Pro
 35 40 45
 Gly Ala Gly Ser Leu Gln Pro Leu Ala Leu Glu Gly Ser Leu Gln Lys
 50 55 60
 Arg Gly Ile Val Glu Gln Cys Cys Thr Ser Ile Cys Ser Leu Tyr Gln
 65 70 75 80
 Leu Glu Asn Tyr Cys Asn
 85

<210> SEQ ID NO 5
 <211> LENGTH: 70
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val Glu Ala Leu Tyr
 1 5 10 15
 Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro Lys Thr Arg Arg
 20 25 30
 Glu Ala Glu Asp Leu Gln Val Gly Gln Val Glu Leu Gly Gly Gly Pro
 35 40 45
 Gly Ala Gly Ser Leu Gln Pro Leu Ala Leu Glu Gly Ser Leu Gln Lys
 50 55 60

-continued

Arg Gly Ile Val Glu Gln
65 70

<210> SEQ ID NO 6
<211> LENGTH: 62
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Ser His Leu Val Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe
1 5 10 15
Phe Tyr Thr Pro Lys Thr Arg Arg Glu Ala Glu Asp Leu Gln Val Gly
20 25 30
Gln Val Glu Leu Gly Gly Gly Pro Gly Ala Gly Ser Leu Gln Pro Leu
35 40 45
Ala Leu Glu Gly Ser Leu Gln Lys Arg Gly Ile Val Glu Gln
50 55 60

<210> SEQ ID NO 7
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Ser His Leu Val Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe
1 5 10 15
Phe Tyr Thr Pro Lys Thr Arg Arg Glu Ala Glu Asp Leu Gln
20 25 30

<210> SEQ ID NO 8
<211> LENGTH: 38
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Phe Val Asn Gln His Leu Cys Gly Ser His Leu Val Glu Ala Leu Tyr
1 5 10 15
Leu Val Cys Gly Glu Arg Gly Phe Phe Tyr Thr Pro Lys Thr Arg Arg
20 25 30
Glu Ala Glu Asp Leu Gln
35

<210> SEQ ID NO 9
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Ser His Leu Val Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly
1 5 10 15

<210> SEQ ID NO 10
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Gly Gly Gly Pro Gly Ala Gly Ser Leu Gln Pro Leu Ala Leu Glu Gly
1 5 10 15
Ser Leu Gln Lys Arg Gly Ile Val Glu Gln Cys
20 25

-continued

<210> SEQ ID NO 11
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

Leu Ala Leu Glu Gly Ser Leu Gln Lys Arg Gly
1 5 10

<210> SEQ ID NO 12
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

Gly Ser Leu Gln Pro Leu Ala Leu Glu Gly Ser Leu Gln Lys Arg Gly
1 5 10 15

Ile Val

<210> SEQ ID NO 13
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

Gly Gly Gly Pro Gly Ala Gly Ser Leu Gln Pro Leu Ala Leu Glu Gly
1 5 10 15

Ser Leu Gln Lys
20

<210> SEQ ID NO 14
<211> LENGTH: 28
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

Ser His Leu Val Glu Ala Leu Tyr Leu Val Cys Gly Glu Arg Gly Phe
1 5 10 15

Phe Tyr Thr Pro Lys Thr Arg Arg Glu Ala Glu Asp
20 25

<210> SEQ ID NO 15
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

Gln Pro Leu Ala Leu Glu Gly Ser Leu Gln Lys Arg Gly Ile Val Glu
1 5 10 15

Gln

<210> SEQ ID NO 16
<211> LENGTH: 65
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Ala Tyr Gln Ala Glu Pro Asn Thr Cys Ala Thr Ala Gln Gly Glu Gly
1 5 10 15

Asn Ile Lys Lys Asn Arg His Pro Asp Phe Leu Pro Tyr Asp His Ala
20 25 30

-continued

Arg Ile Lys Leu Lys Val Glu Ser Ser Pro Ser Arg Ser Asp Tyr Ile
 35 40 45

Asn Ala Ser Pro Ile Ile Glu His Asp Pro Arg Met Pro Ala Tyr Ile
 50 55 60

Ala
 65

<210> SEQ ID NO 17
 <211> LENGTH: 35
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

Gly Pro Leu Ser His Thr Ile Ala Asp Phe Trp Gln Met Val Trp Glu
 1 5 10 15

Ser Gly Cys Thr Val Ile Val Met Leu Thr Pro Leu Val Glu Asp Gly
 20 25 30

Val Lys Gln
 35

<210> SEQ ID NO 18
 <211> LENGTH: 56
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

Gly Ala Ser Leu Tyr His Val Tyr Glu Val Asn Leu Val Ser Glu His
 1 5 10 15

Ile Trp Cys Glu Asp Phe Leu Val Arg Ser Phe Tyr Leu Lys Asn Val
 20 25 30

Gln Thr Gln Glu Thr Arg Thr Leu Thr Gln Phe His Phe Leu Ser Trp
 35 40 45

Pro Ala Glu Gly Thr Pro Ala Ser
 50 55

<210> SEQ ID NO 19
 <211> LENGTH: 39
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

Glu His Val Arg Asp Gln Arg Pro Gly Leu Val Arg Ser Lys Asp Gln
 1 5 10 15

Phe Glu Phe Ala Leu Thr Ala Val Ala Glu Glu Val Asn Ala Ile Leu
 20 25 30

Lys Ala Leu Pro Gln Cys Gly
 35

<210> SEQ ID NO 20
 <211> LENGTH: 304
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

Met Gly Asn His Ala Gly Lys Arg Glu Leu Asn Ala Glu Lys Ala Ser
 1 5 10 15

Thr Asn Ser Glu Thr Asn Arg Gly Glu Ser Glu Lys Lys Arg Asn Leu
 20 25 30

Gly Glu Leu Ser Arg Thr Thr Ser Glu Asp Asn Glu Val Phe Gly Glu

-continued

35	40	45
Ala Asp Ala Asn Gln Asn Asn Gly Thr Ser Ser Gln Asp Thr Ala Val		
50	55	60
Thr Asp Ser Lys Arg Thr Ala Asp Pro Lys Asn Ala Trp Gln Asp Ala		
65	70	80
His Pro Ala Asp Pro Gly Ser Arg Pro His Leu Ile Arg Leu Phe Ser		
	85	90
Arg Asp Ala Pro Gly Arg Glu Asp Asn Thr Phe Lys Asp Arg Pro Ser		
	100	105
Glu Ser Asp Glu Leu Gln Thr Ile Gln Glu Asp Ser Ala Ala Thr Ser		
	115	120
Glu Ser Leu Asp Val Met Ala Ser Gln Lys Arg Pro Ser Gln Arg His		
	130	135
Gly Ser Lys Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His		
	145	150
Gly Phe Leu Pro Arg His Arg Asp Thr Gly Ile Leu Asp Ser Ile Gly		
	165	170
Arg Phe Phe Gly Gly Asp Arg Gly Ala Pro Lys Arg Gly Ser Gly Lys		
	180	185
Asp Ser His His Pro Ala Arg Thr Ala His Tyr Gly Ser Leu Pro Gln		
	195	200
Lys Ser His Gly Arg Thr Gln Asp Glu Asn Pro Val Val His Phe Phe		
	210	215
Lys Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys Gly		
	225	230
Arg Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln Arg		
	245	250
Pro Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys Ser Ala His		
	260	265
Lys Gly Phe Lys Gly Val Asp Ala Gln Gly Thr Leu Ser Lys Ile Phe		
	275	280
Lys Leu Gly Gly Arg Asp Ser Arg Ser Gly Ser Pro Met Ala Arg Arg		
	290	295

<210> SEQ ID NO 21

<211> LENGTH: 247

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

Met Ala Ser Leu Ser Arg Pro Ser Leu Pro Ser Cys Leu Cys Ser Phe		
1	5	10
Leu Leu Leu Leu Leu Leu Gln Val Ser Ser Ser Tyr Ala Gly Gln Phe		
	20	25
Arg Val Ile Gly Pro Arg His Pro Ile Arg Ala Leu Val Gly Asp Glu		
	35	40
Val Glu Leu Pro Cys Arg Ile Ser Pro Gly Lys Asn Ala Thr Gly Met		
	50	55
Glu Val Gly Trp Tyr Arg Pro Pro Phe Ser Arg Val Val His Leu Tyr		
	65	70
Arg Asn Gly Lys Asp Gln Asp Gly Asp Gln Ala Pro Glu Tyr Arg Gly		
	85	90
Arg Thr Glu Leu Leu Lys Asp Ala Ile Gly Glu Gly Lys Val Thr Leu		
	100	105

-continued

```

Arg Ile Arg Asn Val Arg Phe Ser Asp Glu Gly Gly Phe Thr Cys Phe
   115                               120                   125

Phe Arg Asp His Ser Tyr Gln Glu Glu Ala Ala Met Glu Leu Lys Val
   130                               135                   140

Glu Asp Pro Phe Tyr Trp Val Ser Pro Gly Val Leu Val Leu Leu Ala
   145                               150                   155                   160

Val Leu Pro Val Leu Leu Leu Gln Ile Thr Val Gly Leu Ile Phe Leu
   165                               170                   175

Cys Leu Gln Tyr Arg Leu Arg Gly Lys Leu Arg Ala Glu Ile Glu Asn
   180                               185                   190

Leu His Arg Thr Phe Asp Pro His Phe Leu Arg Val Pro Cys Trp Lys
   195                               200                   205

Ile Thr Leu Phe Val Ile Val Pro Val Leu Gly Pro Leu Val Ala Leu
   210                               215                   220

Ile Ile Cys Tyr Asn Trp Leu His Arg Arg Leu Ala Gly Gln Phe Leu
   225                               230                   235                   240

Glu Glu Leu Arg Asn Pro Phe
   245

```

<210> SEQ ID NO 22

<211> LENGTH: 277

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 22

```

Met Gly Leu Leu Glu Cys Cys Ala Arg Cys Leu Val Gly Ala Pro Phe
  1      5      10      15

Ala Ser Leu Val Ala Thr Gly Leu Cys Phe Phe Gly Val Ala Leu Phe
  20      25      30

Cys Gly Cys Gly His Glu Ala Leu Thr Gly Thr Glu Lys Leu Ile Glu
  35      40      45

Thr Tyr Phe Ser Lys Asn Tyr Gln Asp Tyr Glu Tyr Leu Ile Asn Val
  50      55      60

Ile His Ala Phe Gln Tyr Val Ile Tyr Gly Thr Ala Ser Phe Phe Phe
  65      70      75      80

Leu Tyr Gly Ala Leu Leu Leu Ala Glu Gly Phe Tyr Thr Thr Gly Ala
  85      90      95

Val Arg Gln Ile Phe Gly Asp Tyr Lys Thr Thr Ile Cys Gly Lys Gly
  100     105     110

Leu Ser Ala Thr Val Thr Gly Gly Gln Lys Gly Arg Gly Ser Arg Gly
  115     120     125

Gln His Gln Ala His Ser Leu Glu Arg Val Cys His Cys Leu Gly Lys
  130     135     140

Trp Leu Gly His Pro Asp Lys Phe Val Gly Ile Thr Tyr Ala Leu Thr
  145     150     155     160

Val Val Trp Leu Leu Val Phe Ala Cys Ser Ala Val Pro Val Tyr Ile
  165     170     175

Tyr Phe Asn Thr Trp Thr Thr Cys Gln Ser Ile Ala Phe Pro Ser Lys
  180     185     190

Thr Ser Ala Ser Ile Gly Ser Leu Cys Ala Asp Ala Arg Met Tyr Gly
  195     200     205

Val Leu Pro Trp Asn Ala Phe Pro Gly Lys Val Cys Gly Ser Asn Leu
  210     215     220

Leu Ser Ile Cys Lys Thr Ala Glu Phe Gln Met Thr Phe His Leu Phe
  225     230     235     240

```

-continued

Ile Ala Ala Phe Val Gly Ala Ala Ala Thr Leu Val Ser Leu Leu Thr
 245 250 255

Phe Met Ile Ala Ala Thr Tyr Asn Phe Ala Val Leu Lys Leu Met Gly
 260 265 270

Arg Gly Thr Lys Phe
 275

<210> SEQ ID NO 23
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

Lys Tyr Leu Ala Thr Ala Ser Thr Met Asp His Ala Arg His Gly Phe
 1 5 10 15

Leu Pro Arg His
 20

<210> SEQ ID NO 24
 <211> LENGTH: 16
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 24

Glu Asn Pro Trp His Phe Phe Lys Asn Ile Val Thr Pro Arg Thr Pro
 1 5 10 15

<210> SEQ ID NO 25
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 25

Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro Gly Phe Gly
 1 5 10 15

Tyr Gly Gly

<210> SEQ ID NO 26
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 26

Ala Gln Gly Thr Leu Ser Lys Ile Phe Lys Leu Gly Gly Arg Asp Ser
 1 5 10 15

Arg Ser Gly Ser Pro Met Ala Arg Arg
 20 25

<210> SEQ ID NO 27
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 27

Gly Gln Phe Arg Val Ile Gly Pro Arg His Pro Ile Arg Ala Leu Val
 1 5 10 15

Gly Asp Glu Val
 20

<210> SEQ ID NO 28
 <211> LENGTH: 20

-continued

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 28
Met Glu Val Gly Trp Tyr Arg Pro Pro Phe Ser Arg Trp His Leu Tyr
1 5 10 15
Arg Asn Gly Lys
20

<210> SEQ ID NO 29
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 29
His Cys Leu Gly Lys Trp Leu Gly His Pro Asp Lys Phe Val Gly Ile
1 5 10 15

<210> SEQ ID NO 30
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 30
Lys Asn Ala Thr Gly Met Glu Val Gly Trp Tyr Arg Ser Pro Phe Ser
1 5 10 15
Arg Val Val His Leu Tyr Arg Asn Gly Lys Asp Gln Asp Ala Glu
20 25 30

<210> SEQ ID NO 31
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 31
Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg Thr
1 5 10 15
Pro

<210> SEQ ID NO 32
<211> LENGTH: 21
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 32
Met Glu Val Gly Trp Tyr Arg Pro Pro Phe Ser Arg Val Val His Leu
1 5 10 15
Tyr Arg Asn Gly Lys
20

<210> SEQ ID NO 33
<211> LENGTH: 17
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 33
Asp Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1 5 10 15
Thr

<210> SEQ ID NO 34
<211> LENGTH: 18

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 34
Asp Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1          5          10          15
Thr Pro

<210> SEQ ID NO 35
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 35
Asp Glu Asn Pro Val Val His Phe Phe Lys Asn Ile Val Thr Pro Arg
1          5          10          15
Thr Pro Pro Pro Ser Gln Gly Lys Gly
          20          25

<210> SEQ ID NO 36
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 36
Val His Phe Phe Lys Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser
1          5          10          15
Gln Gly Lys Gly
          20

<210> SEQ ID NO 37
<211> LENGTH: 25
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 37
Ala Ser Asp Tyr Lys Ser Ala His Lys Gly Leu Lys Gly Val Asp Ala
1          5          10          15
Gln Gly Thr Leu Ser Lys Ile Phe Lys
          20          25

<210> SEQ ID NO 38
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 38
Gly Thr Glu Lys Leu Ile Glu Thr Tyr Phe Ser Lys Asn Tyr Gln Asp
1          5          10          15
Tyr Glu

<210> SEQ ID NO 39
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 39
Gly Phe Tyr Thr Thr Gly Ala Val Arg Gln Ile Phe Gly Asp Tyr Lys
1          5          10          15
Thr Thr

```

-continued

<210> SEQ ID NO 40
<211> LENGTH: 22
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 40

Ala Val Arg Gln Ile Phe Gly Asp Tyr Lys Thr Thr Ile Cys Gly Lys
1 5 10 15

Gly Leu Ser Ala Thr Val
20

<210> SEQ ID NO 41
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Asn Thr Trp Thr Thr Cys Gln Ser Ile Ala Phe Pro Ser Lys Thr Ser
1 5 10 15

Ala Ser Ile Gly
20

<210> SEQ ID NO 42
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 42

Ser Lys Thr Ser Ala Ser Ile Gly Ser Leu Cys Ala Asp Ala Arg Met
1 5 10 15

Tyr Gly Val Leu
20

<210> SEQ ID NO 43
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 43

Pro Ile Arg Ala Leu Val Gly Asp Glu Val Glu Leu Pro Cys Arg Ile
1 5 10 15

Ser Pro Gly Lys
20

<210> SEQ ID NO 44
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 44

Glu Leu Pro Cys Arg Ile Ser Pro Gly Lys Asn Ala Thr Gly Met Glu
1 5 10 15

Val Gly Trp Tyr
20

<210> SEQ ID NO 45
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 45

Glu Tyr Arg Gly Arg Thr Glu Leu Leu Lys Asp Ala Ile Gly Glu Gly
1 5 10 15

-continued

Lys Val Thr Leu Arg Ile Arg
20

<210> SEQ ID NO 46
<211> LENGTH: 62
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 46

Gly Gln Phe Arg Val Ile Gly Pro Arg His Pro Ile Arg Ala Leu Val
1 5 10 15

Gly Asp Glu Val Glu Leu Pro Cys Arg Ile Ser Pro Gly Lys Asn Ala
20 25 30

Thr Gly Met Glu Val Gly Trp Tyr Arg Pro Pro Phe Ser Arg Val Val
35 40 45

His Leu Tyr Arg Asn Gly Lys Asp Gln Asp Gly Asp Gln Ala
50 55 60

<210> SEQ ID NO 47
<211> LENGTH: 63
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 47

Ser His Gly Arg Thr Gln Asp Glu Asn Pro Val Val His Phe Phe Lys
1 5 10 15

Asn Ile Val Thr Pro Arg Thr Pro Pro Pro Ser Gln Gly Lys Gly Arg
20 25 30

Gly Leu Ser Leu Ser Arg Phe Ser Trp Gly Ala Glu Gly Gln Arg Pro
35 40 45

Gly Phe Gly Tyr Gly Gly Arg Ala Ser Asp Tyr Lys Ser Cys Gly
50 55 60

<210> SEQ ID NO 48
<211> LENGTH: 118
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 48

Met Pro Arg Glu Asp Ala His Phe Ile Tyr Gly Tyr Pro Lys Lys Gly
1 5 10 15

His Gly His Ser Tyr Thr Thr Ala Glu Glu Ala Ala Gly Ile Gly Ile
20 25 30

Leu Thr Val Ile Leu Gly Val Leu Leu Leu Ile Gly Cys Trp Tyr Cys
35 40 45

Arg Arg Arg Asn Gly Tyr Arg Ala Leu Met Asp Lys Ser Leu His Val
50 55 60

Gly Thr Gln Cys Ala Leu Thr Arg Arg Cys Pro Gln Glu Gly Phe Asp
65 70 75 80

His Arg Asp Ser Lys Val Ser Leu Gln Glu Lys Asn Cys Glu Pro Val
85 90 95

Val Pro Asn Ala Pro Pro Ala Tyr Glu Lys Leu Ser Ala Glu Gln Ser
100 105 110

Pro Pro Pro Tyr Ser Pro
115

<210> SEQ ID NO 49
<211> LENGTH: 529

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 49

```

Met Leu Leu Ala Val Leu Tyr Cys Leu Leu Trp Ser Phe Gln Thr Ser
 1          5          10          15

Ala Gly His Phe Pro Arg Ala Cys Val Ser Ser Lys Asn Leu Met Glu
 20          25          30

Lys Glu Cys Cys Pro Pro Trp Ser Gly Asp Arg Ser Pro Cys Gly Gln
 35          40          45

Leu Ser Gly Arg Gly Ser Cys Gln Asn Ile Leu Leu Ser Asn Ala Pro
 50          55          60

Leu Gly Pro Gln Phe Pro Phe Thr Gly Val Asp Asp Arg Glu Ser Trp
 65          70          75          80

Pro Ser Val Phe Tyr Asn Arg Thr Cys Gln Cys Ser Gly Asn Phe Met
 85          90          95

Gly Phe Asn Cys Gly Asn Cys Lys Phe Gly Phe Trp Gly Pro Asn Cys
100          105          110

Thr Glu Arg Arg Leu Leu Val Arg Arg Asn Ile Phe Asp Leu Ser Ala
115          120          125

Pro Glu Lys Asp Lys Phe Phe Ala Tyr Leu Thr Leu Ala Lys His Thr
130          135          140

Ile Ser Ser Asp Tyr Val Ile Pro Ile Gly Thr Tyr Gly Gln Met Lys
145          150          155          160

Asn Gly Ser Thr Pro Met Phe Asn Asp Ile Asn Ile Tyr Asp Leu Phe
165          170          175

Val Trp Met His Tyr Tyr Val Ser Met Asp Ala Leu Leu Gly Gly Ser
180          185          190

Glu Ile Trp Arg Asp Ile Asp Phe Ala His Glu Ala Pro Ala Phe Leu
195          200          205

Pro Trp His Arg Leu Phe Leu Leu Arg Trp Glu Gln Glu Ile Gln Lys
210          215          220

Leu Thr Gly Asp Glu Asn Phe Thr Ile Pro Tyr Trp Asp Trp Arg Asp
225          230          235          240

Ala Glu Lys Cys Asp Ile Cys Thr Asp Glu Tyr Met Gly Gly Gln His
245          250          255

Pro Thr Asn Pro Asn Leu Leu Ser Pro Ala Ser Phe Phe Ser Ser Trp
260          265          270

Gln Ile Val Cys Ser Arg Leu Glu Glu Tyr Asn Ser His Gln Ser Leu
275          280          285

Cys Asn Gly Thr Pro Glu Gly Pro Leu Arg Arg Asn Pro Gly Asn His
290          295          300

Asp Lys Ser Arg Thr Pro Arg Leu Pro Ser Ser Ala Asp Val Glu Phe
305          310          315          320

Cys Leu Ser Leu Thr Gln Tyr Glu Ser Gly Ser Met Asp Lys Ala Ala
325          330          335

Asn Phe Ser Phe Arg Asn Thr Leu Glu Gly Phe Ala Ser Pro Leu Thr
340          345          350

Gly Ile Ala Asp Ala Ser Gln Ser Ser Met His Asn Ala Leu His Ile
355          360          365

Tyr Met Asn Gly Thr Met Ser Gln Val Gln Gly Ser Ala Asn Asp Pro
370          375          380

Ile Phe Leu Leu His His Ala Phe Val Asp Ser Ile Phe Glu Gln Trp
385          390          395          400

```

Leu

```
<210> SEQ ID NO 50
<211> LENGTH: 661
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

<400> SEQUENCE: 50

Met	Asp	Leu	Val	Leu	Lys	Arg	Cys	Leu	Leu	His	Leu	Ala	Val	Ile	Gly
1				5					10					15	
Ala	Leu	Leu	Ala	Val	Gly	Ala	Thr	Lys	Val	Pro	Arg	Asn	Gln	Asp	Trp
			20					25					30		
Leu	Gly	Val	Ser	Arg	Gln	Leu	Arg	Thr	Lys	Ala	Trp	Asn	Arg	Gln	Leu
		35					40					45			
Tyr	Pro	Glu	Trp	Thr	Glu	Ala	Gln	Arg	Leu	Asp	Cys	Trp	Arg	Gly	Gly
	50					55					60				
Gln	Val	Ser	Leu	Lys	Val	Ser	Asn	Asp	Gly	Pro	Thr	Leu	Ile	Gly	Ala
65					70					75					80
Asn	Ala	Ser	Phe	Ser	Ile	Ala	Leu	Asn	Phe	Pro	Gly	Ser	Gln	Lys	Val
				85					90					95	
Leu	Pro	Asp	Gly	Gln	Val	Ile	Trp	Val	Asn	Asn	Thr	Ile	Ile	Asn	Gly
			100					105					110		
Ser	Gln	Val	Trp	Gly	Gly	Gln	Pro	Val	Tyr	Pro	Gln	Glu	Thr	Asp	Asp
		115					120					125			
Ala	Cys	Ile	Phe	Pro	Asp	Gly	Gly	Pro	Cys	Pro	Ser	Gly	Ser	Trp	Ser
	130					135					140				
Gln	Lys	Arg	Ser	Phe	Val	Tyr	Val	Trp	Lys	Thr	Trp	Gly	Gln	Tyr	Trp
145					150					155					160
Gln	Val	Leu	Gly	Gly	Pro	Val	Ser	Gly	Leu	Ser	Ile	Gly	Thr	Gly	Arg
			165						170					175	
Ala	Met	Leu	Gly	Thr	His	Thr	Met	Glu	Val	Thr	Val	Tyr	His	Arg	Arg
		180						185					190		
Gly	Ser	Arg	Ser	Tyr	Val	Pro	Leu	Ala	His	Ser	Ser	Ser	Ala	Phe	Thr
		195					200					205			
Ile	Thr	Asp	Gln	Val	Pro	Phe	Ser	Val	Ser	Val	Ser	Gln	Leu	Arg	Ala
	210					215					220				
Leu	Asp	Gly	Gly	Asn	Lys	His	Phe	Leu	Arg	Asn	Gln	Pro	Leu	Thr	Phe
225					230					235					240

Ala	Leu	Gln	Leu	His	Asp	Pro	Ser	Gly	Tyr	Leu	Ala	Glu	Ala	Asp	Leu	
				245					250					255		
Ser	Tyr	Thr	Trp	Asp	Phe	Gly	Asp	Ser	Ser	Gly	Thr	Leu	Ile	Ser	Arg	
				260					265					270		
Ala	Leu	Val	Val	Thr	His	Thr	Tyr	Leu	Glu	Pro	Gly	Pro	Val	Thr	Ala	
				275					280					285		
Gln	Val	Val	Leu	Gln	Ala	Ala	Ile	Pro	Leu	Thr	Ser	Cys	Gly	Ser	Ser	
				290					295					300		
Pro	Val	Pro	Gly	Thr	Thr	Asp	Gly	His	Arg	Pro	Thr	Ala	Glu	Ala	Pro	
				305					310					315		
Asn	Thr	Thr	Ala	Gly	Gln	Val	Pro	Thr	Thr	Glu	Val	Val	Gly	Thr	Thr	
				325					330					335		
Pro	Gly	Gln	Ala	Pro	Thr	Ala	Glu	Pro	Ser	Gly	Thr	Thr	Ser	Val	Gln	
				340					345					350		
Val	Pro	Thr	Thr	Glu	Val	Ile	Ser	Thr	Ala	Pro	Val	Gln	Met	Pro	Thr	
				355					360					365		
Ala	Glu	Ser	Thr	Gly	Met	Thr	Pro	Glu	Lys	Val	Pro	Val	Ser	Glu	Val	
				370					375					380		
Met	Gly	Thr	Thr	Leu	Ala	Glu	Met	Ser	Thr	Pro	Glu	Ala	Thr	Gly	Met	
				385					390					395		
Thr	Pro	Ala	Glu	Val	Ser	Ile	Val	Val	Leu	Ser	Gly	Thr	Thr	Ala	Ala	
				405					410					415		
Gln	Val	Thr	Thr	Thr	Glu	Trp	Val	Glu	Thr	Thr	Ala	Arg	Glu	Leu	Pro	
				420					425					430		
Ile	Pro	Glu	Pro	Glu	Gly	Pro	Asp	Ala	Ser	Ser	Ile	Met	Ser	Thr	Glu	
				435					440					445		
Ser	Ile	Thr	Gly	Ser	Leu	Gly	Pro	Leu	Leu	Asp	Gly	Thr	Ala	Thr	Leu	
				450					455					460		
Arg	Leu	Val	Lys	Arg	Gln	Val	Pro	Leu	Asp	Cys	Val	Leu	Tyr	Arg	Tyr	
				465					470					475		
Gly	Ser	Phe	Ser	Val	Thr	Leu	Asp	Ile	Val	Gln	Gly	Ile	Glu	Ser	Ala	
				485					490					495		
Glu	Ile	Leu	Gln	Ala	Val	Pro	Ser	Gly	Glu	Gly	Asp	Ala	Phe	Glu	Leu	
				500					505					510		
Thr	Val	Ser	Cys	Gln	Gly	Gly	Leu	Pro	Lys	Glu	Ala	Cys	Met	Glu	Ile	
				515					520					525		
Ser	Ser	Pro	Gly	Cys	Gln	Pro	Pro	Ala	Gln	Arg	Leu	Cys	Gln	Pro	Val	
				530					535					540		
Leu	Pro	Ser	Pro	Ala	Cys	Gln	Leu	Val	Leu	His	Gln	Ile	Leu	Lys	Gly	
				545					550					555		
Gly	Ser	Gly	Thr	Tyr	Cys	Leu	Asn	Val	Ser	Leu	Ala	Asp	Thr	Asn	Ser	
				565					570					575		
Leu	Ala	Val	Val	Ser	Thr	Gln	Leu	Ile	Met	Pro	Gly	Gln	Glu	Ala	Gly	
				580					585					590		
Leu	Gly	Gln	Val	Pro	Leu	Ile	Val	Gly	Ile	Leu	Leu	Val	Leu	Met	Ala	
				595					600					605		
Val	Val	Leu	Ala	Ser	Leu	Ile	Tyr	Arg	Arg	Arg	Leu	Met	Lys	Gln	Asp	
				610					615					620		
Phe	Ser	Val	Pro	Gln	Leu	Pro	His	Ser	Ser	Ser	His	Trp	Leu	Arg	Leu	
				625					630					635		
Pro	Arg	Ile	Phe	Cys	Ser	Cys	Pro	Ile	Gly	Glu	Asn	Ser	Pro	Leu	Leu	

-continued

Ser Gly Gln Gln Val
660

<210> SEQ ID NO 51
 <211> LENGTH: 323
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 51

Met Ser Asp Arg Pro Thr Ala Arg Arg Trp Gly Lys Cys Gly Pro Leu
 1 5 10 15
 Cys Thr Arg Glu Asn Ile Met Val Ala Phe Lys Gly Val Trp Thr Gln
 20 25 30
 Ala Phe Trp Lys Ala Val Thr Ala Glu Phe Leu Ala Met Leu Ile Phe
 35 40 45
 Val Leu Leu Ser Leu Gly Ser Thr Ile Asn Trp Gly Gly Thr Glu Lys
 50 55 60
 Pro Leu Pro Val Asp Met Val Leu Ile Ser Leu Cys Phe Gly Leu Ser
 65 70 75 80
 Ile Ala Thr Met Val Gln Cys Phe Gly His Ile Ser Gly Gly His Ile
 85 90 95
 Asn Pro Ala Val Thr Val Ala Met Val Cys Thr Arg Lys Ile Ser Ile
 100 105 110
 Ala Lys Ser Val Phe Tyr Ile Ala Ala Gln Cys Leu Gly Ala Ile Ile
 115 120 125
 Gly Ala Gly Ile Leu Tyr Leu Val Thr Pro Pro Ser Val Val Gly Gly
 130 135 140
 Leu Gly Val Thr Met Val His Gly Asn Leu Thr Ala Gly His Gly Leu
 145 150 155 160
 Leu Val Glu Leu Ile Ile Thr Phe Gln Leu Val Phe Thr Ile Phe Ala
 165 170 175
 Ser Cys Asp Ser Lys Arg Thr Asp Val Thr Gly Ser Ile Ala Leu Ala
 180 185 190
 Ile Gly Phe Ser Val Ala Ile Gly His Leu Phe Ala Ile Asn Tyr Thr
 195 200 205
 Gly Ala Ser Met Asn Pro Ala Arg Ser Phe Gly Pro Ala Val Ile Met
 210 215 220
 Gly Asn Trp Glu Asn His Trp Ile Tyr Trp Val Gly Pro Ile Ile Gly
 225 230 235 240
 Ala Val Leu Ala Gly Gly Leu Tyr Glu Tyr Val Phe Cys Pro Asp Val
 245 250 255
 Glu Phe Lys Arg Arg Phe Lys Glu Ala Phe Ser Lys Ala Ala Gln Gln
 260 265 270
 Thr Lys Gly Ser Tyr Met Glu Val Glu Asp Asn Arg Ser Gln Val Glu
 275 280 285
 Thr Asp Asp Leu Ile Leu Lys Pro Gly Val Val His Val Ile Asp Val
 290 295 300
 Asp Arg Gly Glu Glu Lys Lys Gly Lys Asp Gln Ser Gly Glu Val Leu
 305 310 315 320
 Ser Ser Val

<210> SEQ ID NO 52
 <211> LENGTH: 405
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 52

```

Met Ala Ala Ser Gly Lys Thr Ser Lys Ser Glu Pro Asn His Val Ile
1      5      10      15
Phe Lys Lys Ile Ser Arg Asp Lys Ser Val Thr Ile Tyr Leu Gly Asn
20     25     30
Arg Asp Tyr Ile Asp His Val Ser Gln Val Gln Pro Val Asp Gly Val
35     40     45
Val Leu Val Asp Pro Asp Leu Val Lys Gly Lys Lys Val Tyr Val Thr
50     55     60
Leu Thr Cys Ala Phe Arg Tyr Gly Gln Glu Asp Ile Asp Val Ile Gly
65     70     75     80
Leu Thr Phe Arg Arg Asp Leu Tyr Phe Ser Arg Val Gln Val Tyr Pro
85     90     95
Pro Val Gly Ala Ala Ser Thr Pro Thr Lys Leu Gln Glu Ser Leu Leu
100    105    110
Lys Lys Leu Gly Ser Asn Thr Tyr Pro Phe Leu Leu Thr Phe Pro Asp
115    120    125
Tyr Leu Pro Cys Ser Val Met Leu Gln Pro Ala Pro Gln Asp Ser Gly
130    135    140
Lys Ser Cys Gly Val Asp Phe Glu Val Lys Ala Phe Ala Thr Asp Ser
145    150    155    160
Thr Asp Ala Glu Glu Asp Lys Ile Pro Lys Lys Ser Ser Val Arg Leu
165    170    175
Leu Ile Arg Lys Val Gln His Ala Pro Leu Glu Met Gly Pro Gln Pro
180    185    190
Arg Ala Glu Ala Ala Trp Gln Phe Phe Met Ser Asp Lys Pro Leu His
195    200    205
Leu Ala Val Ser Leu Asn Lys Glu Ile Tyr Phe His Gly Glu Pro Ile
210    215    220
Pro Val Thr Val Thr Val Thr Asn Asn Thr Glu Lys Thr Val Lys Lys
225    230    235    240
Ile Lys Ala Phe Val Glu Gln Val Ala Asn Val Val Leu Tyr Ser Ser
245    250    255
Asp Tyr Tyr Val Lys Pro Val Ala Met Glu Glu Ala Gln Glu Lys Val
260    265    270
Pro Pro Asn Ser Thr Leu Thr Lys Thr Leu Thr Leu Leu Pro Leu Leu
275    280    285
Ala Asn Asn Arg Glu Arg Arg Gly Ile Ala Leu Asp Gly Lys Ile Lys
290    295    300
His Glu Asp Thr Asn Leu Ala Ser Ser Thr Ile Ile Lys Glu Gly Ile
305    310    315    320
Asp Arg Thr Val Leu Gly Ile Leu Val Ser Tyr Gln Ile Lys Val Lys
325    330    335
Leu Thr Val Ser Gly Phe Leu Gly Glu Leu Thr Ser Ser Glu Val Ala
340    345    350
Thr Glu Val Pro Phe Arg Leu Met His Pro Gln Pro Glu Asp Pro Ala
355    360    365
Lys Glu Ser Tyr Gln Asp Ala Asn Leu Val Phe Glu Glu Phe Ala Arg
370    375    380
His Asn Leu Lys Asp Ala Gly Glu Ala Glu Glu Gly Lys Arg Asp Lys
385    390    395    400
Asn Asp Val Asp Glu
405

```

-continued

```

<210> SEQ ID NO 53
<211> LENGTH: 1247
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 53
Met Met Arg Glu Trp Val Leu Leu Met Ser Val Leu Leu Cys Gly Leu
1          5          10          15
Ala Gly Pro Thr His Leu Phe Gln Pro Ser Leu Val Leu Asp Met Ala
20          25          30
Lys Val Leu Leu Asp Asn Tyr Cys Phe Pro Glu Asn Leu Leu Gly Met
35          40          45
Gln Glu Ala Ile Gln Gln Ala Ile Lys Ser His Glu Ile Leu Ser Ile
50          55          60
Ser Asp Pro Gln Thr Leu Ala Ser Val Leu Thr Ala Gly Val Gln Ser
65          70          75          80
Ser Leu Asn Asp Pro Arg Leu Val Ile Ser Tyr Glu Pro Ser Thr Pro
85          90          95
Glu Pro Pro Pro Gln Val Pro Ala Leu Thr Ser Leu Ser Glu Glu Glu
100         105         110
Leu Leu Ala Trp Leu Gln Arg Gly Leu Arg His Glu Val Leu Glu Gly
115         120         125
Asn Val Gly Tyr Leu Arg Val Asp Ser Val Pro Gly Gln Glu Val Leu
130         135         140
Ser Met Met Gly Glu Phe Leu Val Ala His Val Trp Gly Asn Leu Met
145         150         155         160
Gly Thr Ser Ala Leu Val Leu Asp Leu Arg His Cys Thr Gly Gly Gln
165         170         175
Val Ser Gly Ile Pro Tyr Ile Ile Ser Tyr Leu His Pro Gly Asn Thr
180         185         190
Ile Leu His Val Asp Thr Ile Tyr Asn Arg Pro Ser Asn Thr Thr Thr
195         200         205
Glu Ile Trp Thr Leu Pro Gln Val Leu Gly Glu Arg Tyr Gly Ala Asp
210         215         220
Lys Asp Val Val Val Leu Thr Ser Ser Gln Thr Arg Gly Val Ala Glu
225         230         235         240
Asp Ile Ala His Ile Leu Lys Gln Met Arg Arg Ala Ile Val Val Gly
245         250         255
Glu Arg Thr Gly Gly Gly Ala Leu Asp Leu Arg Lys Leu Arg Ile Gly
260         265         270
Glu Ser Asp Phe Phe Phe Thr Val Pro Val Ser Arg Ser Leu Gly Pro
275         280         285
Leu Gly Gly Gly Ser Gln Thr Trp Glu Gly Ser Gly Val Leu Pro Cys
290         295         300
Val Gly Thr Pro Ala Glu Gln Ala Leu Glu Lys Ala Leu Ala Ile Leu
305         310         315         320
Thr Leu Arg Ser Ala Leu Pro Gly Val Val His Cys Leu Gln Glu Val
325         330         335
Leu Lys Asp Tyr Tyr Thr Leu Val Asp Arg Val Pro Thr Leu Leu Gln
340         345         350
His Leu Ala Ser Met Asp Phe Ser Thr Val Val Ser Glu Glu Asp Leu
355         360         365
Val Thr Lys Leu Asn Ala Gly Leu Gln Ala Ala Ser Glu Asp Pro Arg

```

370					375					380					
Leu 385	Leu	Val	Arg	Ala	Ile 390	Gly	Pro	Thr	Glu	Thr 395	Pro	Ser	Trp	Pro	Ala 400
Pro	Asp	Ala	Ala	Ala 405	Glu	Asp	Ser	Pro	Gly 410	Val	Ala	Pro	Glu	Leu 415	Pro
Glu	Asp	Glu	Ala	Ile 420	Arg	Gln	Ala	Leu 425	Val	Asp	Ser	Val	Phe	Gln	Val
Ser	Val	Leu 435	Pro	Gly	Asn	Val	Gly 440	Tyr	Leu	Arg	Phe	Asp 445	Ser	Phe	Ala
Asp	Ala	Ser	Val	Leu	Gly 455	Val	Leu	Ala	Pro	Tyr	Val 460	Leu	Arg	Gln	Val
Trp 465	Glu	Pro	Leu	Gln	Asp 470	Thr	Glu	His	Leu	Ile 475	Met	Asp	Leu	Arg	His 480
Asn	Pro	Gly	Gly	Pro 485	Ser	Ser	Ala	Val	Pro 490	Leu	Leu	Leu	Ser	Tyr	Phe 495
Gln	Gly	Pro	Glu	Ala 500	Gly	Pro	Val	His 505	Leu	Phe	Thr	Thr	Tyr	Asp	Arg 510
Arg	Thr	Asn 515	Ile	Thr	Gln	Glu	His 520	Phe	Ser	His	Met	Glu 525	Leu	Pro	Gly
Pro	Arg	Tyr 530	Ser	Thr	Gln	Arg 535	Gly	Val	Tyr	Leu	Leu 540	Thr	Ser	His	Arg
Thr 545	Ala	Thr	Ala	Ala 550	Glu	Glu	Phe	Ala	Phe	Leu	Met 555	Gln	Ser	Leu	Gly 560
Trp	Ala	Thr	Leu	Val 565	Gly	Glu	Ile	Thr	Ala 570	Gly	Asn	Leu	Leu	His 575	Thr
Arg	Thr	Val 580	Pro	Leu	Leu	Asp	Thr	Pro 585	Glu	Gly	Ser	Leu	Ala	Leu	Thr 590
Val	Pro	Val 595	Leu	Thr	Phe	Ile	Asp 600	Asn	His	Gly	Glu	Ala 605	Trp	Leu	Gly
Gly	Gly	Val 610	Val	Pro	Asp	Ala 615	Ile	Val	Leu	Ala	Glu 620	Glu	Ala	Leu	Asp
Lys 625	Ala	Gln	Glu	Val	Leu 630	Glu	Phe	His	Gln	Ser 635	Leu	Gly	Ala	Leu	Val 640
Glu	Gly	Thr	Gly	His 645	Leu	Leu	Glu	Ala	His 650	Tyr	Ala	Arg	Pro	Glu	Val 655
Val	Gly	Gln	Thr	Ser 660	Ala	Leu	Leu	Arg 665	Ala	Lys	Leu	Ala	Gln	Gly	Ala 670
Tyr	Arg	Thr 675	Ala	Val	Asp	Leu	Glu 680	Ser	Leu	Ala	Ser	Gln 685	Leu	Thr	Ala
Asp	Leu	Gln 690	Glu	Val	Ser	Gly 695	Asp	His	Arg	Leu	Leu 700	Val	Phe	His	Ser
Pro	Gly	Glu 705	Leu	Val	Val 710	Glu	Glu	Ala	Pro	Pro 715	Pro	Pro	Pro	Ala	Val 720
Pro	Ser	Pro	Glu	Glu 725	Leu	Thr	Tyr	Leu	Ile 730	Glu	Ala	Leu	Phe	Lys	Thr 735
Glu	Val	Leu	Pro	Gly 740	Gln	Leu	Gly	Tyr 745	Leu	Arg	Phe	Asp	Ala	Met	Ala 750
Glu	Leu	Glu 755	Thr	Val	Lys	Ala	Val 760	Gly	Pro	Gln	Leu	Val 765	Arg	Leu	Val
Trp	Gln	Gln 770	Leu	Val	Asp 775	Thr	Ala	Ala	Leu	Val	Ile 780	Asp	Leu	Arg	Tyr
Asn 785	Pro	Gly	Ser	Tyr	Ser 790	Thr	Ala	Ile	Pro	Leu 795	Leu	Cys	Ser	Tyr	Phe 800

Phe 805	Glu 805	Ala 805	Glu 805	Pro 805	Arg 805	Gln 805	His 805	Leu 805	Tyr 805	Ser 805	Val 805	Phe 805	Asp 805	Arg 805	Ala 805
Thr 820	Ser 820	Lys 820	Val 820	Thr 820	Glu 820	Val 820	Trp 820	Thr 825	Leu 825	Pro 825	Gln 825	Val 830	Ala 830	Gly 830	Gln 830
Arg 835	Tyr 835	Gly 835	Ser 835	His 835	Lys 835	Asp 840	Leu 840	Tyr 840	Ile 840	Leu 840	Met 845	Ser 845	His 845	Thr 845	Ser 845
Gly 850	Ser 850	Ala 850	Ala 850	Glu 850	Ala 855	Phe 855	Ala 855	His 855	Thr 855	Met 860	Gln 860	Asp 860	Leu 860	Gln 860	Arg 860
Ala 865	Thr 865	Val 865	Ile 865	Gly 870	Glu 870	Pro 870	Thr 870	Ala 870	Gly 875	Gly 875	Ala 875	Leu 875	Ser 875	Val 875	Gly 880
Ile 885	Tyr 885	Gln 885	Val 885	Gly 885	Ser 885	Ser 885	Pro 885	Leu 885	Tyr 890	Ala 890	Ser 890	Met 890	Pro 890	Thr 895	Gln 895
Met 900	Ala 900	Met 900	Ser 900	Ala 900	Thr 900	Thr 900	Gly 905	Lys 905	Ala 905	Trp 905	Asp 910	Leu 910	Ala 910	Gly 910	Val 910
Glu 915	Pro 915	Asp 915	Ile 915	Thr 915	Val 915	Pro 920	Met 920	Ser 920	Glu 920	Ala 920	Leu 925	Ser 925	Ile 925	Ala 925	Gln 925
Asp 930	Ile 930	Val 930	Ala 930	Leu 930	Arg 935	Ala 935	Lys 935	Val 935	Pro 935	Thr 935	Val 940	Leu 940	Gln 940	Thr 940	Ala 940
Gly 945	Lys 945	Leu 945	Val 945	Ala 945	Asp 950	Asn 950	Tyr 950	Ala 950	Ser 955	Ala 955	Glu 955	Leu 955	Gly 955	Ala 960	Lys 960
Met 965	Ala 965	Thr 965	Lys 965	Leu 965	Ser 965	Gly 965	Leu 965	Gln 965	Ser 970	Arg 970	Tyr 970	Ser 970	Arg 970	Val 975	Thr 975
Ser 980	Glu 980	Val 980	Ala 980	Leu 980	Ala 980	Glu 980	Ile 985	Leu 985	Gly 985	Ala 985	Asp 985	Leu 985	Gln 985	Met 985	Leu 985
Ser 995	Gly 995	Asp 995	Pro 995	His 995	Leu 995	Lys 995	Ala 995	Ala 995	His 995	Ile 995	Pro 995	Glu 995	Asn 995	Ala 995	Lys 995
Asp 1010	Arg 1010	Ile 1010	Pro 1010	Gly 1010	Ile 1010	Val 1015	Pro 1015	Met 1015	Gln 1015	Ile 1015	Pro 1020	Ser 1020	Pro 1020	Glu 1020	
Val 1025	Phe 1025	Glu 1025	Glu 1025	Leu 1025	Ile 1025	Lys 1030	Phe 1030	Ser 1030	Phe 1030	His 1030	Thr 1035	Asn 1035	Val 1035	Leu 1035	
Glu 1040	Asp 1040	Asn 1040	Ile 1040	Gly 1040	Tyr 1040	Leu 1045	Arg 1045	Phe 1045	Asp 1045	Met 1045	Phe 1050	Gly 1050	Asp 1050	Gly 1050	
Glu 1055	Leu 1055	Leu 1055	Thr 1055	Gln 1055	Val 1055	Ser 1060	Arg 1060	Leu 1060	Leu 1060	Val 1060	Glu 1065	His 1065	Ile 1065	Trp 1065	
Lys 1070	Lys 1070	Ile 1070	Met 1070	His 1070	Thr 1070	Asp 1075	Ala 1075	Met 1075	Ile 1075	Ile 1075	Asp 1080	Met 1080	Arg 1080	Phe 1080	
Asn 1085	Ile 1085	Gly 1085	Gly 1085	Pro 1085	Thr 1085	Ser 1090	Ser 1090	Ile 1090	Pro 1090	Ile 1090	Leu 1095	Cys 1095	Ser 1095	Tyr 1095	
Phe 1100	Phe 1100	Asp 1100	Glu 1100	Gly 1100	Pro 1100	Pro 1105	Val 1105	Leu 1105	Leu 1105	Asp 1105	Lys 1110	Ile 1110	Tyr 1110	Ser 1110	
Arg 1115	Pro 1115	Asp 1115	Asp 1115	Ser 1115	Val 1115	Ser 1120	Glu 1120	Leu 1120	Trp 1120	Thr 1120	His 1125	Ala 1125	Gln 1125	Val 1125	
Val 1130	Gly 1130	Glu 1130	Arg 1130	Tyr 1130	Gly 1130	Ser 1135	Lys 1135	Lys 1135	Ser 1135	Met 1135	Val 1140	Ile 1140	Leu 1140	Thr 1140	
Ser 1145	Ser 1145	Val 1145	Thr 1145	Ala 1145	Gly 1145	Thr 1150	Ala 1150	Glu 1150	Glu 1150	Phe 1150	Thr 1155	Tyr 1155	Ile 1155	Met 1155	
Lys 1160	Arg 1160	Leu 1160	Gly 1160	Arg 1160	Ala 1160	Leu 1165	Val 1165	Ile 1165	Gly 1165	Glu 1165	Val 1170				

-continued

Ser	Ser	Trp	Glu	Gly	Val	Gly	Val	Thr	Pro	His	Val	Val	Val	Pro
1205						1210					1215			
Ala	Glu	Glu	Ala	Leu	Ala	Arg	Ala	Lys	Glu	Met	Leu	Gln	His	Asn
1220						1225					1230			
Gln	Leu	Arg	Val	Lys	Arg	Ser	Pro	Gly	Leu	Gln	Asp	His	Leu	
1235						1240					1245			

<210> SEQ ID NO 54
 <211> LENGTH: 33
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 54

Leu	Gln	Leu	Gln	Pro	Phe	Pro	Gln	Pro	Gln	Leu	Pro	Tyr	Pro	Gln	Pro
1			5					10						15	
Gln	Leu	Pro	Tyr	Pro	Gln	Pro	Gln	Leu	Pro	Tyr	Pro	Gln	Pro	Gln	Pro
			20					25					30		

Phe

<210> SEQ ID NO 55
 <211> LENGTH: 33
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 55

Leu	Gln	Leu	Gln	Pro	Phe	Pro	Gln	Pro	Glu	Leu	Pro	Tyr	Pro	Gln	Pro
1			5						10					15	
Glu	Leu	Pro	Tyr	Pro	Gln	Pro	Glu	Leu	Pro	Tyr	Pro	Gln	Pro	Gln	Pro
			20					25					30		

Phe

<210> SEQ ID NO 56
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 56

Gln	Gln	Tyr	Pro	Ser	Gly	Gln	Gly	Ser	Phe	Gln	Pro	Ser	Gln	Gln	Asn
1			5					10						15	

Pro Gln

<210> SEQ ID NO 57
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 57

Gln	Pro	Phe	Pro	Gln	Pro	Glu	Gln	Pro	Phe	Pro	Trp
1			5					10			

<210> SEQ ID NO 58
 <211> LENGTH: 15
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 58

Glu	Leu	Gln	Pro	Phe	Pro	Gln	Pro	Glu	Leu	Pro	Tyr	Pro	Gln	Pro
1			5					10					15	

<210> SEQ ID NO 59
 <211> LENGTH: 18

-continued

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 59

Gly Cys Arg Gly Gly Gly Pro Gln Pro Gln Pro Phe Pro Ser Gln Gln
 1 5 10 15

Pro Tyr

<210> SEQ ID NO 60

<211> LENGTH: 51

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 60

Gly Cys Arg Gly Gly Gly Pro Gln Pro Gln Pro Phe Pro Ser Gln Gln
 1 5 10 15

Pro Tyr Leu Gln Leu Gln Pro Phe Pro Gln Pro Gln Leu Pro Tyr Pro
 20 25 30

Gln Pro Gln Leu Pro Tyr Pro Gln Pro Gln Leu Pro Tyr Pro Gln Pro
 35 40 45

Gln Pro Phe
 50

<210> SEQ ID NO 61

<211> LENGTH: 51

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 61

Gly Cys Arg Gly Gly Gly Pro Gln Pro Gln Pro Phe Pro Ser Gln Gln
 1 5 10 15

Pro Tyr Leu Gln Leu Gln Pro Phe Pro Gln Pro Glu Leu Pro Tyr Pro
 20 25 30

Gln Pro Glu Leu Pro Tyr Pro Gln Pro Glu Leu Pro Tyr Pro Gln Pro
 35 40 45

Gln Pro Phe
 50

<210> SEQ ID NO 62

<211> LENGTH: 140

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 62

Met Asp Val Phe Met Lys Gly Leu Ser Lys Ala Lys Glu Gly Val Val
 1 5 10 15

Ala Ala Ala Glu Lys Thr Lys Gln Gly Val Ala Glu Ala Ala Gly Lys
 20 25 30

Thr Lys Glu Gly Val Leu Tyr Val Gly Ser Lys Thr Lys Glu Gly Val
 35 40 45

Val His Gly Val Ala Thr Val Ala Glu Lys Thr Lys Glu Gln Val Thr
 50 55 60

Asn Val Gly Gly Ala Val Val Thr Gly Val Thr Ala Val Ala Gln Lys
 65 70 75 80

Thr Val Glu Gly Ala Gly Ser Ile Ala Ala Ala Thr Gly Phe Val Lys
 85 90 95

Lys Asp Gln Leu Gly Lys Asn Glu Glu Gly Ala Pro Gln Glu Gly Ile
 100 105 110


-continued

Leu Glu Asp Met Pro Val Asp Pro Asp Asn Glu Ala Tyr Glu Met Pro
 115 120 125

Ser Glu Glu Gly Tyr Gln Asp Tyr Glu Pro Glu Ala
 130 135 140

The invention claimed is:

1. A compound of Formula 1:



Formula 1

where:




X comprises an antigen, a tolerogenic portion thereof, or a mimetic thereof;

Y comprises a linker moiety;


Z comprises a moiety that specifically targets a mannose receptor;

p is an integer from 2 to 250;


m is an integer from 1 to 100;

R² is any of functional groups I-III:


Ya




Yb




Yc




Yd




Ye




Yf



Yg




Yh



Yi

-continued

III



where Ar is a substituted or unsubstituted aromatic group and one or more of:

R³ is C₁₋₆-alkyl; orR¹¹ is C₁₋₆-alkyl.

2. The compound of claim 1, wherein the moiety that specifically targets a mannose receptor is selected from the group consisting of α -linked mannose, β -linked mannose, substituted mannose, mannose-6-phosphate, N-acetyl mannosamine, and mannan having β (1-4), α (1-6), α (1-2), and/or α (1-3) linkages.

3. The compound of claim 1, wherein Y is a linker resulting from reaction of at least one of a N-hydroxysuccinamidyl linker, maleimide linker, PEG linker, vinylsulfone linker, pyridyl di-thiol-poly (ethylene glycol) linker, pyridyl di-thiol linker, n-nitrophenyl carbonate linker, NHS-ester linker, and nitrophenoxy poly (ethylene glycol) ester linker.


4. The compound of claim 3, wherein Y is covalently bound to X.

5. The compound of claim 1, wherein $-[Y(Z)_p]-$ is represented by one of Formula Ya to Yr:


143

144


-continued




Yj




Yk




Yl




Ym



Yn



Yo




Yp

145

146

-continued

Y_qY_r

20

where

n is an integer from 1 to 100;

q is an integer from 1 to 44;


k is an integer from 1 to 12;

i is an integer from 0 to 20;

v is an integer from 1 to 4;

p is an integer from 2 to 250;

r is an integer from 0 to 250;

 R_1 is $-\text{CH}_2-$, $-(\text{CH}_2)_2-\text{C}(\text{CH}_3)(\text{CN})-$, $-(\text{CH}_2)_2-$, $-\text{C}(\text{CH}_3)(\text{CH}_3)-$, $-(\text{CH}_2)_2-\text{CH}(\text{CH}_3)-$ or $-\text{CH}(\text{CH}_3)-$;
 W^1 and W^2 are as defined below:
 R^9 is a direct bond, $-(\text{CH}_2)_2-\text{NH}-\text{C}(=\text{O})-$ (an ethylacetamido group or "EtAcN") or $-(\text{CH}_2)_2-(\text{O}-\text{CH}_2-\text{CH}_2)_t-\text{NH}-\text{C}(=\text{O})-$ (a pegylated ethylacetamido group or "Et-PEGt-AcN")

t is an integer from 1 to 5;

Z is mannose or a mannose receptor-targeting moiety; and

 R^{10} is an aliphatic group, an alcohol, an aliphatic amine-containing group, or an aliphatic alcohol.

6. The compound of claim 1, wherein Y is an antibody, antibody fragment, peptide or other ligand that binds to X.

7. The compound of claim 1, wherein X is an antigen against which a patient may develop or has developed an unwanted immune response.


8. The compound of claim 7, wherein the antigen is a foreign transplant antigen, an alloantigen, an autoimmune antigen, a food antigen, an animal antigen, a plant antigen, an environmental antigen, a therapeutic antigen, a synthetic self-antigen, or a tolerogenic portion thereof.

9. The compound of claim 8, wherein X is an asparaginase antigen or an ovalbumin antigen.

10. The compound of claim 1, wherein the mannose receptor is mannose-6-phosphate receptor.


11. The compound of claim 1, wherein Y and X are connected through a bond configured to cleave when the compound reaches a target area.

12. The compound of claim 1, wherein Ar is selected from:

where each instance of R'' , where present, is independently selected from an optionally substituted C_{1-6} -alkyl, optionally substituted C_{1-6} alkoxy, optionally substituted amino, OH, or halogen and wherein, X'' is a heteroatom.13. The compound of claim 12 wherein X'' is N.14. The compound of claim 1, wherein R^{11} is $-\text{CH}_3$.

15. A composition comprising the compound of claim 1.

16. A method of inducing immunological tolerance to an antigen target comprising administering to a subject a composition Formula 1:



Formula 1

where:


X comprises an antigen, a tolerogenic portion thereof, or a mimetic thereof;

Y comprises a linker moiety;

Z comprises a moiety that specifically targets a mannose receptor;


p is an integer from 2 to 250;

m is an integer from 1 to 100;

147R² is any of functional groups I-III:


I

5



II

10



III

15

where Ar is a substituted or unsubstituted aromatic group
and one or more of:

R³ is C₁₋₆-alkyl; or

R¹¹ is C₁₋₆-alkyl.

17. The method of claim 16, wherein the moiety that
specifically targets a mannose receptor is selected from the
group consisting of α -linked mannose, β -linked mannose,
substituted mannose, mannose-6-phosphate, N-acetyl man-
nosamine, and mannan having β (1-4), α (1-6), α (1-2), and/or
 α (1-3) linkages.

* * * * *

148