

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12409125
Kind Code	B2
Date of Patent	September 09, 2025
Inventor(s)	Cochran; Brooke Michele et al.

Shampoo compositions containing a sulfate-free surfactant system and sclerotium gum thickener

Abstract

A stable shampoo composition containing a surfactant system that is substantially free of sulfated surfactants. The shampoo composition also contains an anionic surfactant, an amphoteric surfactant, a cationic polymer and a sclerotium gum thickener. The shampoo composition has a consumer-preferred viscosity of greater than 2500 cP, a pH greater than 5.5 and lacks in-situ coacervate.

Inventors: Cochran; Brooke Michele (Cincinnati, OH), Brown; Mark Anthony (Union, KY), Zhao; Jean Jianqun (Cincinnati, OH), Ballhaus; Lauren Elizabeth (Williamsburg, OH), Mazzei; Nicole Marie (Cincinnati, OH)

Applicant: The Procter & Gamble Company (Cincinnati, OH)

Family ID: 81851689

Assignee: The Procter & Gamble Company (Cincinnati, OH)

Appl. No.: 17/738697

Filed: May 06, 2022

Prior Publication Data

Document Identifier	Publication Date
US 20220378684 A1	Dec. 01, 2022

Related U.S. Application Data

us-provisional-application US 63276108 20211105
us-provisional-application US 63188517 20210514

Publication Classification

Int. Cl.: A61K8/60 (20060101); A61K8/19 (20060101); A61K8/368 (20060101); A61K8/49 (20060101); A61K8/73 (20060101); A61K8/92 (20060101); A61K8/9789 (20170101);

U.S. Cl.:

CPC **A61K8/604** (20130101); **A61K8/19** (20130101); **A61K8/368** (20130101); **A61K8/4926** (20130101); **A61K8/4946** (20130101); **A61K8/4953** (20130101); **A61K8/73** (20130101); **A61K8/737** (20130101); **A61K8/922** (20130101); **A61K8/9789** (20170801); **A61Q5/02** (20130101); **A61Q17/005** (20130101); A61K2800/33 (20130101); A61K2800/5426 (20130101)

Field of Classification Search

CPC: A61K (8/604); A61K (8/19); A61K (8/368); A61K (8/4926); A61K (8/4946); A61K (8/4953); A61K (8/73); A61K (8/737); A61K (8/922); A61K (8/9789); A61K (8/42); A61K (8/44); A61K (8/466); A61K (8/731); A61K (8/732); A61K (8/8158); A61K (2800/30); A61K (2800/596); A61K (2800/33); A61K (2800/5426); A61K (2800/52); A61Q (5/02); A61Q (5/12); A61Q (17/005)

References Cited**U.S. PATENT DOCUMENTS**

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
1489388	12/1923	Glenn	N/A	N/A
1600340	12/1925	Hoffman	N/A	N/A
1612255	12/1925	William	N/A	N/A
2438091	12/1947	Lynch	N/A	N/A
2528378	12/1949	Mannheimer	N/A	N/A
2658072	12/1952	Milton	N/A	N/A
2809971	12/1956	Bernstein et al.	N/A	N/A
2879231	12/1958	Marshall	N/A	N/A
3219656	12/1964	Boettner	N/A	N/A
3236733	12/1965	Karsten et al.	N/A	N/A
3373208	12/1967	Blumenthal	N/A	N/A
3636113	12/1971	Hall	N/A	N/A
3709437	12/1972	Wright	N/A	N/A
3716498	12/1972	Hall	N/A	N/A
3753196	12/1972	Kurtz et al.	N/A	N/A
3761418	12/1972	Parran	N/A	N/A
3792068	12/1973	Luedders et al.	N/A	N/A
3887692	12/1974	Gilman	N/A	N/A
3904741	12/1974	Jones et al.	N/A	N/A
3950532	12/1975	Bouillon et al.	N/A	N/A
3959160	12/1975	Horsler et al.	N/A	N/A
4049792	12/1976	Elsnau	N/A	N/A
4089945	12/1977	Brinkman et al.	N/A	N/A
4120948	12/1977	Shelton	N/A	N/A
4137180	12/1978	Naik	N/A	N/A
4237155	12/1979	Kardouche	N/A	N/A
4309119	12/1981	Wittersheim	N/A	N/A
4323683	12/1981	Bolich, Jr. et al.	N/A	N/A
4329334	12/1981	Su et al.	N/A	N/A
4345080	12/1981	Bolich	N/A	N/A
4359456	12/1981	Gosling et al.	N/A	N/A

4379753	12/1982	Bolich, Jr.	N/A	N/A
4430243	12/1983	Bragg	N/A	N/A
4470982	12/1983	Winkler	N/A	N/A
4726945	12/1987	Patel	N/A	N/A
4732696	12/1987	Urfer	N/A	N/A
4839166	12/1988	Grollier et al.	N/A	N/A
4854333	12/1988	Inman et al.	N/A	N/A
4867971	12/1988	Ryan et al.	N/A	N/A
4931274	12/1989	Barabino et al.	N/A	N/A
4973416	12/1989	Kennedy	N/A	N/A
4985238	12/1990	Tanner et al.	N/A	N/A
4997641	12/1990	Hartnett	N/A	N/A
5019375	12/1990	Tanner et al.	N/A	N/A
5093112	12/1991	Birtwistle et al.	N/A	N/A
5104646	12/1991	Bolich, Jr.	N/A	N/A
5106609	12/1991	Bolich, Jr.	N/A	N/A
5135747	12/1991	Faryniarz et al.	N/A	N/A
5156834	12/1991	Beckmeyer et al.	N/A	N/A
5294644	12/1993	Login et al.	N/A	N/A
5296157	12/1993	Macgilp et al.	N/A	N/A
5296622	12/1993	Uphues	N/A	N/A
5298640	12/1993	Callaghan et al.	N/A	N/A
5332569	12/1993	Wood et al.	N/A	N/A
5364031	12/1993	Taniguchi et al.	N/A	N/A
5374421	12/1993	Tashiro	N/A	N/A
5374614	12/1993	Behan et al.	N/A	N/A
5409695	12/1994	Abrutyn et al.	N/A	N/A
5415810	12/1994	Lee et al.	N/A	N/A
5417965	12/1994	Janchitraponvej et al.	N/A	N/A
5429816	12/1994	Hofrichter et al.	N/A	N/A
5439682	12/1994	Wivell	N/A	N/A
5441659	12/1994	Minor	N/A	N/A
5486303	12/1995	Capeci	N/A	N/A
5489392	12/1995	Capeci	N/A	N/A
5496488	12/1995	Kacher et al.	N/A	N/A
5516448	12/1995	Capeci	N/A	N/A
5536493	12/1995	Dubief	N/A	N/A
5554588	12/1995	Behan et al.	N/A	N/A
5560918	12/1995	Wivell	N/A	N/A
5565422	12/1995	Del Greco	N/A	N/A
5569645	12/1995	Dinniwell	N/A	N/A
5574005	12/1995	Welch	N/A	N/A
5576282	12/1995	Miracle	N/A	N/A
5578298	12/1995	Berthiaume	N/A	N/A
5595967	12/1996	Miracle	N/A	N/A
5597936	12/1996	Perkins	N/A	N/A
5599549	12/1996	Wivell	N/A	N/A
5624666	12/1996	Coffindaffer et al.	N/A	N/A
5635469	12/1996	Fowler et al.	N/A	N/A
5665267	12/1996	Dowell et al.	N/A	N/A
5691297	12/1996	Nassano	N/A	N/A
5714137	12/1997	Trinh	N/A	N/A
5747436	12/1997	Patel	N/A	N/A

5776444	12/1997	Birtwistle et al.	N/A	N/A
5800897	12/1997	Sharma	N/A	N/A
5816446	12/1997	Steindorf et al.	N/A	N/A
5830440	12/1997	Sturla et al.	N/A	N/A
5853618	12/1997	Barker	N/A	N/A
5879584	12/1998	Bianchetti	N/A	N/A
5891424	12/1998	Bretzler et al.	N/A	N/A
5902225	12/1998	Monson	N/A	N/A
5925603	12/1998	D'Angelo	N/A	N/A
5942217	12/1998	Woo et al.	N/A	N/A
5976514	12/1998	Guskey et al.	N/A	N/A
5980877	12/1998	Baravetto	N/A	N/A
5985939	12/1998	Minor	N/A	N/A
6015547	12/1999	Yam	N/A	N/A
6015780	12/1999	Llosas Bigorra et al.	N/A	N/A
6020303	12/1999	Cripe et al.	N/A	N/A
6039933	12/1999	Samain et al.	N/A	N/A
6046152	12/1999	Vinson et al.	N/A	N/A
6060443	12/1999	Cripe et al.	N/A	N/A
6087309	12/1999	Vinson et al.	N/A	N/A
6110451	12/1999	Matz et al.	N/A	N/A
6133222	12/1999	Vinson et al.	N/A	N/A
6139828	12/1999	Mccullough	N/A	N/A
6153567	12/1999	Hughes	N/A	N/A
6153569	12/1999	Halloran	N/A	N/A
6162834	12/1999	Sebillotte-Arnaud et al.	N/A	N/A
6180121	12/2000	Guenin et al.	N/A	N/A
6225464	12/2000	Hiler, II	N/A	N/A
6231844	12/2000	Nambu	N/A	N/A
6232302	12/2000	Alberico et al.	N/A	N/A
6248135	12/2000	Trinh et al.	N/A	N/A
6268431	12/2000	Snyder et al.	N/A	N/A
6284225	12/2000	Bhatt	N/A	N/A
6329331	12/2000	Aronson	N/A	N/A
6335312	12/2001	Coffindaffer et al.	N/A	N/A
6352688	12/2001	Scavone et al.	N/A	N/A
6386392	12/2001	Argentieri	N/A	N/A
6413920	12/2001	Bettiol	N/A	N/A
6423305	12/2001	Cauwet-Martin et al.	N/A	N/A
6436442	12/2001	Woo et al.	N/A	N/A
6451300	12/2001	Dunlop et al.	N/A	N/A
6488943	12/2001	Beerse et al.	N/A	N/A
6511669	12/2002	Garnier et al.	N/A	N/A
6565863	12/2002	Guillou et al.	N/A	N/A
6579907	12/2002	Sebillotte-Arnaud et al.	N/A	N/A
6627585	12/2002	Steer	N/A	N/A
6642194	12/2002	Harrison	N/A	N/A
6649155	12/2002	Dunlop et al.	N/A	N/A
6656923	12/2002	Trinh	N/A	N/A
6660288	12/2002	Behan et al.	N/A	N/A
6679324	12/2003	Den Boer et al.	N/A	N/A
6716455	12/2003	Birkel	N/A	N/A
6716805	12/2003	Sherry	N/A	N/A

6740713	12/2003	Busch et al.	N/A	N/A
6743760	12/2003	Hardy et al.	N/A	N/A
6764986	12/2003	Busch et al.	N/A	N/A
6767507	12/2003	Woo et al.	N/A	N/A
6794356	12/2003	Turner	N/A	N/A
6814088	12/2003	Barnabas et al.	N/A	N/A
6827795	12/2003	Kasturi et al.	N/A	N/A
6869923	12/2004	Cunningham	N/A	N/A
6897253	12/2004	Schmucker-castner	N/A	N/A
6908889	12/2004	Niemiec et al.	N/A	N/A
6930078	12/2004	Wells	N/A	N/A
6992054	12/2005	Lee et al.	N/A	N/A
7018978	12/2005	Miracle et al.	N/A	N/A
7030068	12/2005	Clare et al.	N/A	N/A
7100767	12/2005	Chomik et al.	N/A	N/A
7151079	12/2005	Fack et al.	N/A	N/A
7172099	12/2006	Hoefte	N/A	N/A
7202198	12/2006	Gordon et al.	N/A	N/A
7217752	12/2006	Schmucker-Castner et al.	N/A	N/A
7220408	12/2006	Decoster et al.	N/A	N/A
7223361	12/2006	Kvietok	N/A	N/A
7223385	12/2006	Gawtrex et al.	N/A	N/A
7485289	12/2008	Gawtrex et al.	N/A	N/A
7504094	12/2008	Decoster et al.	N/A	N/A
7531497	12/2008	Midha et al.	N/A	N/A
7541320	12/2008	Dabkowski et al.	N/A	N/A
7598213	12/2008	Geary et al.	N/A	N/A
7659233	12/2009	Conopco	N/A	N/A
7666825	12/2009	Wagner et al.	N/A	N/A
7820609	12/2009	Soffin et al.	N/A	N/A
7829514	12/2009	Paul et al.	N/A	N/A
7841036	12/2009	Smith	N/A	N/A
7867505	12/2010	Elliott et al.	N/A	N/A
7928053	12/2010	Hecht et al.	N/A	N/A
7977288	12/2010	SenGupta	N/A	N/A
8007545	12/2010	Fujii et al.	N/A	N/A
8058500	12/2010	Sojka et al.	N/A	N/A
8084407	12/2010	Soffin et al.	N/A	N/A
8088721	12/2011	Soffin et al.	N/A	N/A
8119168	12/2011	Johnson	N/A	N/A
8124063	12/2011	Harichian et al.	N/A	N/A
8158571	12/2011	Alonso	N/A	N/A
8300949	12/2011	Xu	N/A	N/A
8322631	12/2011	Richardson et al.	N/A	N/A
8343469	12/2012	Bierganns et al.	N/A	N/A
8349300	12/2012	Wells	N/A	N/A
8357359	12/2012	Woo et al.	N/A	N/A
8361450	12/2012	Johnson et al.	N/A	N/A
8388699	12/2012	Wood	N/A	N/A
8401304	12/2012	Cavallaro et al.	N/A	N/A
8435501	12/2012	Peffly et al.	N/A	N/A
8437556	12/2012	Saisan	N/A	N/A
8491877	12/2012	Schwartz et al.	N/A	N/A

8539631	12/2012	Catalfamo et al.	N/A	N/A
8574561	12/2012	Patel et al.	N/A	N/A
8580725	12/2012	Kuhlman et al.	N/A	N/A
8609600	12/2012	Warr et al.	N/A	N/A
8628760	12/2013	Carter et al.	N/A	N/A
8629095	12/2013	Deleersnyder	N/A	N/A
8653014	12/2013	Hilvert	N/A	N/A
8675919	12/2013	Maladen	N/A	N/A
8679316	12/2013	Brunner et al.	N/A	N/A
8680035	12/2013	Kuhlman et al.	N/A	N/A
8699751	12/2013	Maladen	N/A	N/A
8709337	12/2013	Gruenbacher et al.	N/A	N/A
8709385	12/2013	Tamarkin	N/A	N/A
8741275	12/2013	Dente et al.	N/A	N/A
8741363	12/2013	Albrecht et al.	N/A	N/A
8771765	12/2013	Fernandez	N/A	N/A
8772354	12/2013	Williams et al.	N/A	N/A
8795635	12/2013	Tamarkin et al.	N/A	N/A
8877316	12/2013	Hasenoehrl et al.	N/A	N/A
8883698	12/2013	Scheibel et al.	N/A	N/A
8931711	12/2014	Gruenbacher	N/A	N/A
8980239	12/2014	Staudigel et al.	N/A	N/A
8987187	12/2014	Smets et al.	N/A	N/A
9006162	12/2014	Rizk	N/A	N/A
9186642	12/2014	Dihora et al.	N/A	N/A
9187407	12/2014	Koshti et al.	N/A	N/A
9265727	12/2015	Lowenborg	N/A	N/A
9272164	12/2015	Johnson et al.	N/A	N/A
9296550	12/2015	Smith	N/A	N/A
9308398	12/2015	Hutton et al.	N/A	N/A
9393447	12/2015	Zasloff	N/A	N/A
9428616	12/2015	Wagner	N/A	N/A
9511007	12/2015	Frantz et al.	N/A	N/A
9512275	12/2015	Wagner	N/A	N/A
9610239	12/2016	Feng	N/A	N/A
9655821	12/2016	Carter et al.	N/A	N/A
9662291	12/2016	Johnson et al.	N/A	N/A
9682021	12/2016	Tamarkin et al.	N/A	N/A
9776787	12/2016	Nakajima	N/A	N/A
9949901	12/2017	Zhao et al.	N/A	N/A
9949911	12/2017	Cetti	N/A	N/A
9968535	12/2017	Kitko	N/A	N/A
9968537	12/2017	Sharma	N/A	N/A
9993419	12/2017	Glenn, Jr.	N/A	N/A
9993420	12/2017	Glenn, Jr. et al.	N/A	N/A
10039706	12/2017	Meralli et al.	N/A	N/A
10039939	12/2017	Xavier et al.	N/A	N/A
10113140	12/2017	Frankenbach	N/A	N/A
10182976	12/2018	Staudigel	N/A	N/A
10238685	12/2018	Dunn et al.	N/A	N/A
10265261	12/2018	Park et al.	N/A	N/A
10311575	12/2018	Stofel	N/A	N/A
10392625	12/2018	Jin et al.	N/A	N/A

10426713	12/2018	Song	N/A	N/A
10441519	12/2018	Zhao	N/A	N/A
10552557	12/2019	Frankenbach et al.	N/A	N/A
10610473	12/2019	Hertenstein et al.	N/A	N/A
10653590	12/2019	Torres Rivera	N/A	N/A
10799434	12/2019	Torres Rivera	N/A	N/A
10842720	12/2019	Thompson	N/A	N/A
10881597	12/2020	Lane et al.	N/A	N/A
10888505	12/2020	Johnson	N/A	N/A
10912732	12/2020	Gillis	N/A	N/A
11116703	12/2020	Song et al.	N/A	N/A
11116704	12/2020	Song et al.	N/A	N/A
11129775	12/2020	Song et al.	N/A	N/A
11334694	12/2021	Cetti et al.	N/A	N/A
11334695	12/2021	Cetti et al.	N/A	N/A
11904036	12/2023	Song	N/A	N/A
2001/0000467	12/2000	Murray	N/A	N/A
2001/0006088	12/2000	Lyle	N/A	N/A
2001/0006621	12/2000	Coupe et al.	N/A	N/A
2001/0016565	12/2000	Bodet et al.	N/A	N/A
2002/0012646	12/2001	Royce et al.	N/A	N/A
2002/0028182	12/2001	Dawson	N/A	N/A
2002/0037299	12/2001	Turowski-Wanke et al.	N/A	N/A
2002/0172648	12/2001	Hehner et al.	N/A	N/A
2002/0193265	12/2001	Perron et al.	N/A	N/A
2002/0197213	12/2001	Schmenger et al.	N/A	N/A
2003/0003070	12/2002	Eggers et al.	N/A	N/A
2003/0008787	12/2002	Mcgee et al.	N/A	N/A
2003/0022799	12/2002	Alvarado et al.	N/A	N/A
2003/0049292	12/2002	Turowski-Wanke et al.	N/A	N/A
2003/0050150	12/2002	Tanaka	N/A	N/A
2003/0059377	12/2002	Riley	N/A	N/A
2003/0083210	12/2002	Goldberg	N/A	N/A
2003/0108501	12/2002	Hofrichter	N/A	N/A
2003/0147842	12/2002	Restle et al.	N/A	N/A
2003/0154561	12/2002	Patel	N/A	N/A
2003/0161802	12/2002	Flammer	N/A	N/A
2003/0180238	12/2002	Sakurai et al.	N/A	N/A
2003/0180246	12/2002	Frantz et al.	N/A	N/A
2003/0185867	12/2002	Kerschner et al.	N/A	N/A
2003/0192922	12/2002	Ceppaluni et al.	N/A	N/A
2003/0202952	12/2002	Wells et al.	N/A	N/A
2003/0223951	12/2002	Geary et al.	N/A	N/A
2003/0228272	12/2002	Amjad et al.	N/A	N/A
2004/0014879	12/2003	Denzer et al.	N/A	N/A
2004/0064117	12/2003	Hammons	N/A	N/A
2004/0131660	12/2003	Lange et al.	N/A	N/A
2004/0144863	12/2003	Kendrick	N/A	N/A
2004/0151793	12/2003	Paspaleeva-kuhn et al.	N/A	N/A
2004/0157754	12/2003	Geary et al.	N/A	N/A
2004/0229963	12/2003	Stephane	N/A	N/A
2004/0234484	12/2003	Peffly	N/A	N/A
2004/0235689	12/2003	Sakai et al.	N/A	N/A

2005/0003975	12/2004	Browne et al.	N/A	N/A
2005/0003980	12/2004	Baker	N/A	N/A
2005/0020468	12/2004	Frantz et al.	N/A	N/A
2005/0136011	12/2004	Nekludoff	N/A	N/A
2005/0152863	12/2004	Brautigam	N/A	N/A
2005/0192207	12/2004	Morgan, III et al.	N/A	N/A
2005/0201967	12/2004	Albrecht et al.	N/A	N/A
2005/0202984	12/2004	Schwartz et al.	N/A	N/A
2005/0208106	12/2004	Lange et al.	N/A	N/A
2005/0227902	12/2004	Erazo-majewicz et al.	N/A	N/A
2005/0233929	12/2004	Queen	N/A	N/A
2005/0245407	12/2004	Ishihara	N/A	N/A
2005/0276831	12/2004	Dihora	N/A	N/A
2006/0002880	12/2005	Peffly	N/A	N/A
2006/0005333	12/2005	Catalfamo et al.	N/A	N/A
2006/0009337	12/2005	Smith	N/A	N/A
2006/0030509	12/2005	Modi	N/A	N/A
2006/0034778	12/2005	Kitano et al.	N/A	N/A
2006/0057075	12/2005	Arkin et al.	N/A	N/A
2006/0057097	12/2005	Derici	N/A	N/A
2006/0079417	12/2005	Wagner	N/A	N/A
2006/0079418	12/2005	Wagner et al.	N/A	N/A
2006/0079419	12/2005	Wagner et al.	N/A	N/A
2006/0079420	12/2005	Wagner et al.	N/A	N/A
2006/0079421	12/2005	Wagner et al.	N/A	N/A
2006/0084589	12/2005	Vlad et al.	N/A	N/A
2006/0090777	12/2005	Hecht et al.	N/A	N/A
2006/0094610	12/2005	Yamato et al.	N/A	N/A
2006/0110415	12/2005	Gupta	N/A	N/A
2006/0120982	12/2005	Derici et al.	N/A	N/A
2006/0120988	12/2005	Bailey et al.	N/A	N/A
2006/0135397	12/2005	Bissey-beugras	N/A	N/A
2006/0166857	12/2005	Surburg et al.	N/A	N/A
2006/0171911	12/2005	Schwartz et al.	N/A	N/A
2006/0182703	12/2005	Arisz	N/A	N/A
2006/0183662	12/2005	Crotty	N/A	N/A
2006/0210139	12/2005	Carroll	N/A	N/A
2006/0229227	12/2005	Goldman	N/A	N/A
2006/0252662	12/2005	Soffin	N/A	N/A
2006/0263319	12/2005	Fan et al.	N/A	N/A
2006/0269503	12/2005	Yamazaki	N/A	N/A
2006/0276357	12/2005	Smith et al.	N/A	N/A
2006/0292104	12/2005	Guskey	N/A	N/A
2007/0003499	12/2006	Shen et al.	N/A	N/A
2007/0020263	12/2006	Shitara et al.	N/A	N/A
2007/0072781	12/2006	Soffin et al.	N/A	N/A
2007/0110700	12/2006	Wells	N/A	N/A
2007/0154402	12/2006	Trumbore et al.	N/A	N/A
2007/0155637	12/2006	Smith, III et al.	N/A	N/A
2007/0160555	12/2006	Staudigel	N/A	N/A
2007/0179207	12/2006	Fernandez de Castro et al.	N/A	N/A
2007/0225193	12/2006	Kuhlman et al.	N/A	N/A
2007/0269397	12/2006	Terada	N/A	N/A

2007/0275866	12/2006	Dykstra	N/A	N/A
2007/0292380	12/2006	Staudigel	N/A	N/A
2007/0298994	12/2006	Finke et al.	N/A	N/A
2008/0003245	12/2007	Kroepke et al.	N/A	N/A
2008/0008668	12/2007	Harichian et al.	N/A	N/A
2008/0019928	12/2007	Franzke	N/A	N/A
2008/0063618	12/2007	Johnson	N/A	N/A
2008/0138442	12/2007	Johnson	N/A	N/A
2008/0152610	12/2007	Cajan	N/A	N/A
2008/0160093	12/2007	Schwartz et al.	N/A	N/A
2008/0176780	12/2007	Warr	N/A	N/A
2008/0194454	12/2007	Morgan	N/A	N/A
2008/0206179	12/2007	Peffly et al.	N/A	N/A
2008/0229512	12/2007	Syed et al.	N/A	N/A
2008/0260655	12/2007	Tamarkin et al.	N/A	N/A
2008/0260665	12/2007	Guerchet et al.	N/A	N/A
2008/0261844	12/2007	Ruppert et al.	N/A	N/A
2008/0317698	12/2007	Wells et al.	N/A	N/A
2009/0005280	12/2008	Woo et al.	N/A	N/A
2009/0029900	12/2008	Cetti et al.	N/A	N/A
2009/0041702	12/2008	Molenda	N/A	N/A
2009/0062406	12/2008	Loeffler	N/A	N/A
2009/0155383	12/2008	Kitko et al.	N/A	N/A
2009/0178210	12/2008	Bistram	N/A	N/A
2009/0197784	12/2008	Ainger	N/A	N/A
2009/0221463	12/2008	Kitko et al.	N/A	N/A
2009/0240223	12/2008	Warren	N/A	N/A
2009/0246236	12/2008	Kitko	N/A	N/A
2009/0312223	12/2008	Yang et al.	N/A	N/A
2009/0312224	12/2008	Yang et al.	N/A	N/A
2009/0324505	12/2008	Seidling	N/A	N/A
2010/0000116	12/2009	Aouad et al.	N/A	N/A
2010/0001116	12/2009	Johnson	N/A	N/A
2010/0009285	12/2009	Daems et al.	N/A	N/A
2010/0061946	12/2009	Scherner et al.	N/A	N/A
2010/0087357	12/2009	Morgan, III et al.	N/A	N/A
2010/0152083	12/2009	Velazquez	N/A	N/A
2010/0168251	12/2009	Warr et al.	N/A	N/A
2010/0183539	12/2009	Bernhardt	N/A	N/A
2010/0215775	12/2009	Schmaus et al.	N/A	N/A
2010/0287710	12/2009	Denutte et al.	N/A	N/A
2010/0310644	12/2009	Liebmann	N/A	N/A
2010/0322878	12/2009	Stella et al.	N/A	N/A
2011/0008267	12/2010	Arkin et al.	N/A	N/A
2011/0023266	12/2010	Gross et al.	N/A	N/A
2011/0098209	12/2010	Smets et al.	N/A	N/A
2011/0107524	12/2010	Chieffi et al.	N/A	N/A
2011/0118691	12/2010	Nishitani	N/A	N/A
2011/0139170	12/2010	Hippe et al.	N/A	N/A
2011/0150815	12/2010	Woo et al.	N/A	N/A
2011/0165107	12/2010	Derks et al.	N/A	N/A
2011/0171155	12/2010	Federle	N/A	N/A
2011/0177017	12/2010	Coffindaffer et al.	N/A	N/A

2011/0232668	12/2010	Hoffmann et al.	N/A	N/A
2011/0245126	12/2010	Tsaur et al.	N/A	N/A
2011/0245134	12/2010	Smets	N/A	N/A
2011/0245136	12/2010	Smets	N/A	N/A
2011/0268778	12/2010	Dihora et al.	N/A	N/A
2011/0269657	12/2010	Dihora	N/A	N/A
2011/0300095	12/2010	Dente et al.	N/A	N/A
2011/0303766	12/2010	Smith	N/A	N/A
2011/0305739	12/2010	Royce	N/A	N/A
2011/0305778	12/2010	Caggioni et al.	N/A	N/A
2011/0308555	12/2010	Smets et al.	N/A	N/A
2011/0308556	12/2010	Smets et al.	N/A	N/A
2011/0319790	12/2010	Kost et al.	N/A	N/A
2012/0004328	12/2011	Huchel et al.	N/A	N/A
2012/0009285	12/2011	Wei et al.	N/A	N/A
2012/0014901	12/2011	Sunkel et al.	N/A	N/A
2012/0031419	12/2011	Batt	N/A	N/A
2012/0034173	12/2011	Batt	N/A	N/A
2012/0052031	12/2011	Troccaz et al.	N/A	N/A
2012/0100091	12/2011	Hata et al.	N/A	N/A
2012/0100092	12/2011	Murray	N/A	N/A
2012/0129924	12/2011	Park et al.	N/A	N/A
2012/0219610	12/2011	Smith, III et al.	N/A	N/A
2012/0230936	12/2011	Mikkelsen	N/A	N/A
2012/0237469	12/2011	Dente et al.	N/A	N/A
2012/0246851	12/2011	Smith et al.	N/A	N/A
2012/0258150	12/2011	Rauckhorst et al.	N/A	N/A
2012/0291911	12/2011	Smith	N/A	N/A
2012/0309660	12/2011	Kawasoe	N/A	N/A
2012/0316095	12/2011	Wei et al.	N/A	N/A
2013/0029932	12/2012	Kachi et al.	N/A	N/A
2013/0034515	12/2012	Stone et al.	N/A	N/A
2013/0043145	12/2012	Smith, III et al.	N/A	N/A
2013/0043146	12/2012	Smith, III et al.	N/A	N/A
2013/0043147	12/2012	Smith, III et al.	N/A	N/A
2013/0045285	12/2012	Stella et al.	N/A	N/A
2013/0053295	12/2012	Kinoshita et al.	N/A	N/A
2013/0053300	12/2012	Scheibel et al.	N/A	N/A
2013/0089587	12/2012	Staudigel	N/A	N/A
2013/0115173	12/2012	Trumbore et al.	N/A	N/A
2013/0143784	12/2012	Rizk	N/A	N/A
2013/0150338	12/2012	Ananthapadmanabhan	N/A	N/A
2013/0156712	12/2012	Frantz	N/A	N/A
2013/0189212	12/2012	Jawale et al.	N/A	N/A
2013/0211952	12/2012	Sugaya	N/A	N/A
2013/0216491	12/2012	Ogihara et al.	N/A	N/A
2013/0243718	12/2012	Pasquet	N/A	N/A
2013/0244922	12/2012	Bartelt	N/A	N/A
2013/0266642	12/2012	Hollingshead et al.	N/A	N/A
2013/0280192	12/2012	Carter et al.	N/A	N/A
2013/0280202	12/2012	Stella et al.	N/A	N/A
2013/0284195	12/2012	Murdock	N/A	N/A
2013/0296289	12/2012	Hall	N/A	N/A

2013/0319463	12/2012	Policicchio	N/A	N/A
2014/0037703	12/2013	Dihora et al.	N/A	N/A
2014/0039066	12/2013	Grimadell et al.	N/A	N/A
2014/0086893	12/2013	Gutmann et al.	N/A	N/A
2014/0112879	12/2013	Molenda	N/A	N/A
2014/0127149	12/2013	Lepilleur	N/A	N/A
2014/0131395	12/2013	Chang	N/A	N/A
2014/0134125	12/2013	Dahl et al.	N/A	N/A
2014/0162979	12/2013	Palla-venkata et al.	N/A	N/A
2014/0171471	12/2013	Krueger	N/A	N/A
2014/0186864	12/2013	Kato et al.	N/A	N/A
2014/0201927	12/2013	Bianchetti et al.	N/A	N/A
2014/0216495	12/2013	Bureiko	N/A	N/A
2014/0221269	12/2013	Sobel et al.	N/A	N/A
2014/0228268	12/2013	Fahl et al.	N/A	N/A
2014/0237732	12/2013	Zuedel Fernandes et al.	N/A	N/A
2014/0246515	12/2013	Nakajima	N/A	N/A
2014/0308227	12/2013	Mabille	N/A	N/A
2014/0309154	12/2013	Carter et al.	N/A	N/A
2014/0335041	12/2013	Peffly et al.	N/A	N/A
2014/0348884	12/2013	Hilvert et al.	N/A	N/A
2014/0348886	12/2013	Johnson et al.	N/A	N/A
2014/0349902	12/2013	Allef et al.	N/A	N/A
2015/0017152	12/2014	Potechin et al.	N/A	N/A
2015/0021496	12/2014	Shabbir	N/A	N/A
2015/0037273	12/2014	Wagner	N/A	N/A
2015/0050231	12/2014	Murase	N/A	N/A
2015/0057208	12/2014	Frantz et al.	N/A	N/A
2015/0071977	12/2014	Dihora	N/A	N/A
2015/0093420	12/2014	Snyder	N/A	N/A
2015/0093429	12/2014	Carter et al.	N/A	N/A
2015/0098921	12/2014	Franzke et al.	N/A	N/A
2015/0099684	12/2014	Boutique	N/A	N/A
2015/0108163	12/2014	Smith et al.	N/A	N/A
2015/0110728	12/2014	Jayaswal	N/A	N/A
2015/0141310	12/2014	Smets et al.	N/A	N/A
2015/0147286	12/2014	Barrera	N/A	N/A
2015/0157548	12/2014	De Feij et al.	N/A	N/A
2015/0218496	12/2014	Schmiedel et al.	N/A	N/A
2015/0231045	12/2014	Krohn et al.	N/A	N/A
2015/0262354	12/2014	Periaswamy	N/A	N/A
2015/0297489	12/2014	Kleinen et al.	N/A	N/A
2015/0299400	12/2014	Wagner et al.	N/A	N/A
2015/0313818	12/2014	Stagg	N/A	N/A
2015/0352027	12/2014	Thomas et al.	N/A	N/A
2015/0359725	12/2014	Glenn, Jr. et al.	N/A	N/A
2015/0359726	12/2014	Glenn, Jr. et al.	N/A	N/A
2015/0359728	12/2014	Glenn, Jr. et al.	N/A	N/A
2016/0008257	12/2015	Zhou et al.	N/A	N/A
2016/0022566	12/2015	Figura	N/A	N/A
2016/0089317	12/2015	Cetti et al.	N/A	N/A
2016/0089318	12/2015	Cetti et al.	N/A	N/A
2016/0089322	12/2015	Santos Nogueira et al.	N/A	N/A

2016/0089462	12/2015	Frankenbach	N/A	N/A
2016/0089464	12/2015	Frankenbach et al.	N/A	N/A
2016/0089465	12/2015	Frankenbach et al.	N/A	N/A
2016/0090555	12/2015	Frankenbach et al.	N/A	N/A
2016/0090556	12/2015	Frankenbach et al.	N/A	N/A
2016/0090557	12/2015	Frankenbach et al.	N/A	N/A
2016/0090558	12/2015	Frankenbach et al.	N/A	N/A
2016/0092661	12/2015	Hollingshead et al.	N/A	N/A
2016/0095804	12/2015	Xavier et al.	N/A	N/A
2016/0113849	12/2015	Grimadell et al.	N/A	N/A
2016/0128944	12/2015	Chawrai	N/A	N/A
2016/0193125	12/2015	Jones et al.	N/A	N/A
2016/0206522	12/2015	Ribaut et al.	N/A	N/A
2016/0235643	12/2015	Mathonneau et al.	N/A	N/A
2016/0250115	12/2015	Li et al.	N/A	N/A
2016/0279048	12/2015	Jayaswal et al.	N/A	N/A
2016/0287503	12/2015	Schroeder	N/A	N/A
2016/0287509	12/2015	Peffly	N/A	N/A
2016/0296656	12/2015	Scavone et al.	N/A	N/A
2016/0303043	12/2015	Khoury	N/A	N/A
2016/0306909	12/2015	Hollingshead et al.	N/A	N/A
2016/0309871	12/2015	Torres Rivera et al.	N/A	N/A
2016/0310369	12/2015	Thompson	N/A	N/A
2016/0310370	12/2015	Smith	N/A	N/A
2016/0310371	12/2015	Thompson	N/A	N/A
2016/0310375	12/2015	Torres Rivera	N/A	N/A
2016/0310386	12/2015	Smith, III et al.	N/A	N/A
2016/0310388	12/2015	Smith, III et al.	N/A	N/A
2016/0310389	12/2015	Zhao	N/A	N/A
2016/0310390	12/2015	Smith, III et al.	N/A	N/A
2016/0310391	12/2015	Smith, III et al.	N/A	N/A
2016/0310393	12/2015	Chang et al.	N/A	N/A
2016/0310402	12/2015	Zhao	N/A	N/A
2016/0317424	12/2015	Kadir et al.	N/A	N/A
2016/0326458	12/2015	Smets et al.	N/A	N/A
2016/0338929	12/2015	Zasloff	N/A	N/A
2016/0354300	12/2015	Thompson et al.	N/A	N/A
2017/0066579	12/2016	Zillges	N/A	N/A
2017/0071837	12/2016	Schelges et al.	N/A	N/A
2017/0101609	12/2016	Vargas	N/A	N/A
2017/0110690	12/2016	Lamansky et al.	N/A	N/A
2017/0110695	12/2016	Nishikawa et al.	N/A	N/A
2017/0119917	12/2016	Frankenbach et al.	N/A	N/A
2017/0137752	12/2016	Frankenbach et al.	N/A	N/A
2017/0137753	12/2016	Frankenbach et al.	N/A	N/A
2017/0165164	12/2016	Zhao et al.	N/A	N/A
2017/0165165	12/2016	Zhao et al.	N/A	N/A
2017/0209359	12/2016	Zhao et al.	N/A	N/A
2017/0224607	12/2016	Li et al.	N/A	N/A
2017/0239155	12/2016	Hartnett	N/A	N/A
2017/0249407	12/2016	Cetti et al.	N/A	N/A
2017/0249408	12/2016	Cetti et al.	N/A	N/A
2017/0252273	12/2016	Renock et al.	N/A	N/A

2017/0255725	12/2016	Frankenbach et al.	N/A	N/A
2017/0273880	12/2016	Hertenstein	N/A	N/A
2017/0278249	12/2016	Stofel	N/A	N/A
2017/0283959	12/2016	Shellef	N/A	N/A
2017/0304172	12/2016	Chang et al.	N/A	N/A
2017/0304184	12/2016	Glenn, Jr.	N/A	N/A
2017/0304185	12/2016	Glenn, Jr. et al.	N/A	N/A
2017/0304186	12/2016	Glenn, Jr.	N/A	N/A
2017/0333321	12/2016	Carnali	N/A	N/A
2017/0333591	12/2016	Scavone et al.	N/A	N/A
2017/0367963	12/2016	Kadir et al.	N/A	N/A
2018/0004875	12/2017	Cetti et al.	N/A	N/A
2018/0044097	12/2017	Zeik	N/A	N/A
2018/0057451	12/2017	Owens et al.	N/A	N/A
2018/0066210	12/2017	Frankenbach et al.	N/A	N/A
2018/0098923	12/2017	Hutton, III	N/A	N/A
2018/0110594	12/2017	Atkin	N/A	N/A
2018/0110688	12/2017	Torres Rivera et al.	N/A	N/A
2018/0110689	12/2017	Torres Rivera et al.	N/A	N/A
2018/0110690	12/2017	Torres Rivera	N/A	N/A
2018/0110691	12/2017	Torres Rivera et al.	N/A	N/A
2018/0110692	12/2017	Torres Rivera et al.	N/A	N/A
2018/0110693	12/2017	Renock et al.	N/A	N/A
2018/0110694	12/2017	Renock et al.	N/A	N/A
2018/0110695	12/2017	Thompson	N/A	N/A
2018/0110696	12/2017	Johnson et al.	N/A	N/A
2018/0110704	12/2017	Zhao et al.	N/A	N/A
2018/0110707	12/2017	Zhao et al.	N/A	N/A
2018/0110710	12/2017	Zhao et al.	N/A	N/A
2018/0110714	12/2017	Glenn, Jr. et al.	N/A	N/A
2018/0116937	12/2017	L'Oreal	N/A	N/A
2018/0116941	12/2017	L'Oreal	N/A	N/A
2018/0133133	12/2017	Kleinen et al.	N/A	N/A
2018/0177708	12/2017	Lee et al.	N/A	N/A
2018/0221266	12/2017	Zhao et al.	N/A	N/A
2018/0256481	12/2017	Glenn, Jr	N/A	N/A
2018/0280270	12/2017	Rughani et al.	N/A	N/A
2018/0311135	12/2017	Chang	N/A	N/A
2018/0311136	12/2017	Chang	N/A	N/A
2018/0318194	12/2017	Hoffmann et al.	N/A	N/A
2018/0344611	12/2017	Zhao et al.	N/A	N/A
2018/0344612	12/2017	Zhao et al.	N/A	N/A
2018/0344613	12/2017	Zhao et al.	N/A	N/A
2018/0344614	12/2017	Zhao et al.	N/A	N/A
2018/0360713	12/2017	Jouy et al.	N/A	N/A
2019/0105242	12/2018	Song	N/A	N/A
2019/0105243	12/2018	Song	N/A	N/A
2019/0105244	12/2018	Song	N/A	N/A
2019/0105245	12/2018	Song	N/A	N/A
2019/0105246	12/2018	Cochran	N/A	N/A
2019/0105247	12/2018	Song	N/A	N/A
2019/0117543	12/2018	Zhao	N/A	N/A
2019/0117544	12/2018	Zhao	N/A	N/A

2019/0117545	12/2018	Zhao	N/A	N/A
2019/0125650	12/2018	Lee et al.	N/A	N/A
2019/0142711	12/2018	Torres Rivera	N/A	N/A
2019/0142800	12/2018	Ghosh et al.	N/A	N/A
2019/0155975	12/2018	Cetti et al.	N/A	N/A
2019/0167554	12/2018	Wankhade	N/A	N/A
2019/0183777	12/2018	Gillis	N/A	N/A
2019/0183778	12/2018	Glenn, Jr.	N/A	N/A
2019/0192405	12/2018	Zhao	N/A	N/A
2019/0240121	12/2018	Torres Rivera	N/A	N/A
2019/0307298	12/2018	Zhao	N/A	N/A
2019/0328647	12/2018	Chang et al.	N/A	N/A
2019/0365611	12/2018	Brown et al.	N/A	N/A
2019/0365619	12/2018	Ceballos et al.	N/A	N/A
2019/0365633	12/2018	Glenn, Jr.	N/A	N/A
2020/0000690	12/2019	Renock	N/A	N/A
2020/0078284	12/2019	Botto et al.	N/A	N/A
2020/0129402	12/2019	Jamadagni	N/A	N/A
2020/0163846	12/2019	Song	N/A	N/A
2020/0170894	12/2019	Park et al.	N/A	N/A
2020/0197272	12/2019	Hertenstein et al.	N/A	N/A
2020/0206110	12/2019	Hertenstein et al.	N/A	N/A
2020/0237628	12/2019	Torres Rivera	N/A	N/A
2021/0022986	12/2020	Glenn, Jr.	N/A	N/A
2021/0093543	12/2020	Parikh et al.	N/A	N/A
2021/0121385	12/2020	Muller et al.	N/A	N/A
2021/0128444	12/2020	Muller et al.	N/A	N/A
2021/0128447	12/2020	Galpin	N/A	A61K 8/368
2021/0169765	12/2020	Renock	N/A	N/A
2021/0212927	12/2020	Hutton, III et al.	N/A	N/A
2021/0267853	12/2020	Johnson et al.	N/A	N/A
2021/0275410	12/2020	Hutton, III	N/A	N/A
2021/0353518	12/2020	Ballhaus et al.	N/A	N/A
2021/0353522	12/2020	Ballhaus et al.	N/A	N/A
2021/0401716	12/2020	Gogineni et al.	N/A	N/A
2022/0062136	12/2021	Feng	N/A	A61K 8/44
2022/0160606	12/2021	Renock	N/A	N/A
2022/0175640	12/2021	Herteinstein et al.	N/A	N/A
2022/0323337	12/2021	Yamazaki	N/A	N/A
2022/0378680	12/2021	Ballhaus et al.	N/A	N/A
2022/0395444	12/2021	Hutton, III	N/A	N/A
2023/0053056	12/2022	Renock et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
825146	12/1974	BE	N/A
199400875	12/1995	BR	N/A
704195	12/1964	CA	N/A
1248458	12/1988	CA	N/A
2078375	12/1993	CA	N/A
1263455	12/1999	CN	N/A
1286612	12/2000	CN	N/A
1545404	12/2003	CN	N/A

1823929	12/2005	CN	N/A
100534415	12/2008	CN	N/A
101112349	12/2010	CN	N/A
101690697	12/2010	CN	N/A
101559034	12/2012	CN	N/A
102895151	12/2012	CN	N/A
102973437	12/2012	CN	N/A
102697668	12/2012	CN	N/A
103356408	12/2012	CN	N/A
102697670	12/2013	CN	N/A
104107401	12/2013	CN	N/A
102851015	12/2013	CN	N/A
105726393	12/2015	CN	N/A
105769617	12/2015	CN	N/A
106659664	12/2016	CN	N/A
106750361	12/2016	CN	N/A
107595657	12/2017	CN	N/A
107595673	12/2017	CN	N/A
107648096	12/2017	CN	N/A
107737329	12/2017	CN	N/A
107961212	12/2017	CN	N/A
108186385	12/2017	CN	N/A
108283583	12/2017	CN	N/A
108451858	12/2017	CN	N/A
110279591	12/2018	CN	N/A
113041184	12/2020	CN	N/A
2145204	12/1972	DE	N/A
3018456	12/1980	DE	N/A
4315396	12/1993	DE	N/A
102004012009	12/2004	DE	N/A
202005009618	12/2004	DE	N/A
102004023720	12/2004	DE	N/A
102014225083	12/2014	DE	N/A
102014225606	12/2014	DE	N/A
102015204987	12/2015	DE	N/A
0108517	12/1983	EP	N/A
0574086	12/1992	EP	N/A
0666358	12/1994	EP	N/A
0674898	12/1994	EP	N/A
1340485	12/2002	EP	N/A
1346720	12/2002	EP	N/A
067898	12/2005	EP	N/A
1714678	12/2005	EP	N/A
1842572	12/2006	EP	N/A
2005939	12/2007	EP	N/A
1970045	12/2008	EP	N/A
2042216	12/2014	EP	N/A
3121210	12/2016	EP	N/A
3260171	12/2016	EP	N/A
3622946	12/2019	EP	N/A
2052450	12/1993	ES	N/A
2669531	12/1991	FR	N/A
2795955	12/2000	FR	N/A

190110699	12/1900	GB	N/A
191023922	12/1910	GB	N/A
1347950	12/1973	GB	N/A
2048229	12/1979	GB	N/A
2450727	12/2008	GB	N/A
42318	12/1985	HU	N/A
S56011009	12/1980	JP	N/A
S58113300	12/1982	JP	N/A
S58198412	12/1982	JP	N/A
AS60004598	12/1984	JP	N/A
S61236708	12/1985	JP	N/A
S62205200	12/1986	JP	N/A
S63501221	12/1987	JP	N/A
S63165308	12/1987	JP	N/A
H04364114	12/1991	JP	N/A
H06220495	12/1993	JP	N/A
H0753340	12/1994	JP	N/A
07252134	12/1994	JP	N/A
H08310924	12/1995	JP	N/A
09020618	12/1996	JP	N/A
09030938	12/1996	JP	N/A
H09175961	12/1996	JP	N/A
H10017894	12/1997	JP	N/A
H11139944	12/1998	JP	N/A
2964226	12/1998	JP	N/A
2000178586	12/1999	JP	N/A
3069802	12/1999	JP	N/A
2001011492	12/2000	JP	N/A
2001011497	12/2000	JP	N/A
2001254099	12/2000	JP	N/A
2001261529	12/2000	JP	N/A
2003201217	12/2000	JP	N/A
2002179552	12/2001	JP	N/A
2002226889	12/2001	JP	N/A
2002285191	12/2001	JP	N/A
2002336337	12/2001	JP	N/A
2003055699	12/2002	JP	N/A
2003082398	12/2002	JP	N/A
2003171688	12/2002	JP	N/A
2003176497	12/2002	JP	N/A
2003261413	12/2002	JP	N/A
2003268398	12/2002	JP	N/A
3480165	12/2002	JP	N/A
2003342131	12/2002	JP	N/A
3634988	12/2004	JP	N/A
3634991	12/2004	JP	N/A
3634996	12/2004	JP	N/A
2005187359	12/2004	JP	N/A
2005232113	12/2004	JP	N/A
2006063044	12/2005	JP	N/A
2006104149	12/2005	JP	N/A
2006124312	12/2005	JP	N/A
2006183039	12/2005	JP	N/A

2006193549	12/2005	JP	N/A
2006249092	12/2005	JP	N/A
2006282565	12/2005	JP	N/A
2007131687	12/2006	JP	N/A
2007177047	12/2006	JP	N/A
2007223935	12/2006	JP	N/A
2007238607	12/2006	JP	N/A
2007297380	12/2006	JP	N/A
2008001626	12/2007	JP	N/A
2008214292	12/2007	JP	N/A
2009096778	12/2008	JP	N/A
2009120559	12/2008	JP	N/A
2009161866	12/2008	JP	N/A
2010138130	12/2009	JP	N/A
2011153167	12/2010	JP	N/A
2011190221	12/2010	JP	N/A
2011241353	12/2010	JP	N/A
5041113	12/2011	JP	N/A
2013010757	12/2012	JP	N/A
2013091641	12/2012	JP	N/A
2013151434	12/2012	JP	N/A
2013155143	12/2012	JP	N/A
2013193968	12/2012	JP	N/A
2013216639	12/2012	JP	N/A
6046394	12/2013	JP	N/A
2014009177	12/2013	JP	N/A
2014024875	12/2013	JP	N/A
2014037383	12/2013	JP	N/A
2014091723	12/2013	JP	N/A
2014234350	12/2013	JP	N/A
5667790	12/2014	JP	N/A
2015034157	12/2014	JP	N/A
2015101545	12/2014	JP	N/A
2015129099	12/2014	JP	N/A
2016013973	12/2015	JP	N/A
2016030722	12/2015	JP	N/A
2016088910	12/2015	JP	N/A
6184550	12/2016	JP	N/A
2018012673	12/2017	JP	N/A
2018083781	12/2017	JP	N/A
100290589	12/2000	KR	N/A
100821846	12/2007	KR	N/A
1020080111280	12/2007	KR	N/A
20090095359	12/2008	KR	N/A
20100040180	12/2009	KR	N/A
20140060882	12/2013	KR	N/A
101494008	12/2014	KR	N/A
101503922	12/2014	KR	N/A
101532070	12/2014	KR	N/A
50333	12/2009	UA	N/A
8603679	12/1985	WO	N/A
9114759	12/1990	WO	N/A
91017237	12/1990	WO	N/A

9213520	12/1991	WO	N/A
199325650	12/1992	WO	N/A
9417783	12/1993	WO	N/A
9502389	12/1994	WO	N/A
9726854	12/1996	WO	N/A
9823258	12/1997	WO	N/A
9906010	12/1998	WO	N/A
9918928	12/1998	WO	N/A
9924004	12/1998	WO	N/A
9924013	12/1998	WO	N/A
9949837	12/1998	WO	N/A
9957233	12/1998	WO	N/A
0012553	12/1999	WO	N/A
0032601	12/1999	WO	N/A
0119949	12/2000	WO	N/A
0142409	12/2000	WO	N/A
0148021	12/2000	WO	N/A
2001076552	12/2000	WO	N/A
2003051319	12/2002	WO	N/A
03096998	12/2002	WO	N/A
03105793	12/2002	WO	N/A
2004078901	12/2003	WO	N/A
2005023975	12/2004	WO	N/A
2008017540	12/2007	WO	N/A
2008128826	12/2007	WO	N/A
2008145582	12/2007	WO	N/A
2009016555	12/2008	WO	N/A
2009030594	12/2008	WO	N/A
2009053931	12/2008	WO	N/A
2010026009	12/2009	WO	N/A
2010052147	12/2009	WO	N/A
2011124560	12/2010	WO	N/A
2012017091	12/2011	WO	N/A
2012052536	12/2011	WO	N/A
2012055587	12/2011	WO	N/A
2012055812	12/2011	WO	N/A
2012084970	12/2011	WO	N/A
2012127009	12/2011	WO	N/A
2012136651	12/2011	WO	N/A
2013010706	12/2012	WO	N/A
2013018805	12/2012	WO	N/A
2013119908	12/2012	WO	N/A
2014073245	12/2013	WO	N/A
2014073456	12/2013	WO	N/A
2014111667	12/2013	WO	N/A
2014111668	12/2013	WO	N/A
2014148245	12/2013	WO	N/A
2015016037	12/2014	WO	N/A
2015067779	12/2014	WO	N/A
2015085376	12/2014	WO	N/A
2015122371	12/2014	WO	N/A
2015141787	12/2014	WO	N/A
2016049389	12/2015	WO	N/A

2016147196	12/2015	WO	N/A
2017052161	12/2016	WO	N/A
2017140798	12/2016	WO	N/A
2017140802	12/2016	WO	N/A
2017207685	12/2016	WO	N/A
2018023180	12/2017	WO	N/A
2018064511	12/2017	WO	N/A
2018109148	12/2017	WO	N/A
2019030458	12/2018	WO	N/A
2019074990	12/2018	WO	N/A
2019074992	12/2018	WO	N/A
2019200027	12/2018	WO	N/A
2020005309	12/2019	WO	N/A
2020030732	12/2019	WO	N/A
2020076881	12/2019	WO	N/A
2020131836	12/2019	WO	N/A
2021026572	12/2020	WO	N/A
2021099088	12/2020	WO	N/A
2021127318	12/2020	WO	N/A
2021144326	12/2020	WO	N/A
2021231510	12/2020	WO	N/A
2022144161	12/2021	WO	N/A
2024037872	12/2023	WO	N/A

OTHER PUBLICATIONS

“Anti-Dandruff Shampoo”, Mintel Database, Record No. 752198, dated Aug. 2007 ; pp. 1-3. cited by applicant

“Dandruff Control Shampoo” , Mintel Database, Record No. 2300131, dated Jan. 2014; pp. 1-2. cited by applicant

“Foam & chemical contamination in waterways”, Retrieved From

<https://www.epa.nsw.gov.au/-/media/epa/corporate-site/resources/epa/foam-chemical-contamination-in-waterway.pdf>, Dec. 2015, 2 Pages. cited by applicant

“Natural Detangling Shampoo”, Mintel Database, dated Sep. 13, 2017; 2 pages. cited by applicant

“Soda Shampoo”, Mintel Database, dated Apr. 2015; pp. 1-4. cited by applicant

“Treatment Foam for Recurrent Scaling Conditions”, Mintel Database, Aug. 2007; pp. 1-2. cited by applicant

Acne Foaming Cleanser, Database accession No. 4172863, Jul. 29, 2016, 3 pages. cited by applicant

Air Quality of the Iowa Department of Natural Resources. A Review of the Science and Technology of Odor Measurement, 2005, 51 pages (2005). cited by applicant

Anonymous: “MERQUAT Polyquaternium 47 Series, Water Soluble Polymers for Personal Care”, Jul. 30, 2017, URL: https://www.in-cosmetics.com/_novadocuments/2729, retrieved on Dec. 21, 2018 ; 1 page. cited by applicant

Anonymous: “Naturally Derived Body Wash”, Database GNPD [Online] MINTEL; Feb. 15, 2021, 2 pages. cited by applicant

Anonymous: “Peptide Shampoo”, Database GNPD [Online] MINTEL; Dec. 14, 2015, 3 pages. cited by applicant

Anonymous: “Replenishing Moisture Shampoo”, Database GNPD [Online] MINTEL, Mar. 2, 10015br. cited by applicant

Anonymous: “Shampoo”, Database GNPD [Online] MINTEL, Jan. 26, 2021, 3 pages. cited by applicant

Anonymous: “Shampooing au Phytolait d'abricot—Formule N°102-MP06-MI3-AA03”, Internet Citation, Feb. 19, 2005, Retrieved from the Internet: URL:

<http://web.archive.org/web/20050219040350/www.albanmuller.com/francais/catalogue/formules/formul10.asp>, 1 page. cited by applicant

ASTM D3954-94, Reapproved 2010, vol. 15.04, Standard Test Method for Dropping Point of Waxes. cited by applicant

BASF, "Practical Guide to Rheology Modifiers", download from https://insights.basf.com/files/BASF_ED_RheologyModifiers_download.pdf on Nov. 1, 2022. (Year: 2022). cited by applicant

Brattoli et al. Odour Detection Methods: Offactometry and Chemical Sensors. Sensors (Basel), 2011; 11(5); 5290-5322 (2011). cited by applicant

Carbopol Aqua SF-1 Polymer Technical Data Sheet, TDS-294, dated Dec. 2000 ; pp. 1-9. cited by applicant

Chemical Book (Chemical Book, Isolongifolone, available at http://www.chemicalbook.com/ProductChemicalPropertiesCB5318980_EN.htm), no date available. cited by applicant

Christensen et al., "Experimental Determination of Bubble Size Distribution in a Water Column by Interferometric Particle Imaging and Telecentric Direct Image Method", Student Report, Aalborg University; dated Jun. 3, 2014; 123 pages. cited by applicant

Crepaldi, E.L., et al., Chemical, Structural, and Thermal Properties of Zn(II)—Cr(III) Layered Double Hydroxides Intercalated with Sulfated and Sulfonated Surfactants, Journal of Colloid and Interface Science, 2002, pp. 429-442, vol. 248. cited by applicant

D'Souza et al., Shampoo and Conditioners: What a Dermatologist Should Know? Indian J Dermatol, dated May-Jun. 2015; pp. 60(3), 248-254 (2015). cited by applicant

Database GNPD [Online] MINTEL; Mar. 28, 2018 (Mar. 28, 2018), anonymous: Dandruff Control Shampoo 11, XP055787038, Database accession No. 5556267 abstract. cited by applicant

Database GNPD [Online] MINTEL; Apr. 5, 2005 (Apr. 5, 2005), anonymous: "Anticaspa-Graso Anti-Dandruff Shampoo", XPC:155787029, Database accession No. 351776 paragraph [ingredients]. cited by applicant

Database GNPD [Online] MINTEL; Jan. 6, 2020 (Jan. 6, 2020), anonymous: 11 Shampoo 11, 3 pages. cited by applicant

Database WPI; Week 201459; Thomson scientific, London, GB; An 2014-P66521; XP002752638. cited by applicant

Datasheet: Empigen Total Active TC/U, Datasheet, dated Jan. 31, 2017 (Innospec) ; 2 pages. cited by applicant

Dehyquart Guar: Published dated Nov. 2010 ; pp. 1-34. cited by applicant

Fevola, Michael J. "Guar Hydroxypropyltrimonium Chloride." Cosmetics and toiletries; vol. 127.1; Jan. 2012 ; pp. 16-21. cited by applicant

Grillet et al., "Polymer Gel Rheology and Adhesion", Rheology, 2012, pp. 59-80. cited by applicant

Hair Care/Conditioning Polymers Differentiation, Anonymous, Feb. 1, 2017, URL: <http://www.biochim.it/assets/site/media/allegati/cosmetica/hair-care/tab-merquat-hair-care.pdf>, retrieved on Dec. 20, 2018; p. 1. cited by applicant

Happi: "Sulfate-Free Surfactants Conditioning Shampoo", Retrieved from the Internet: URL: https://www.happi.com/contents/view_formulary/2009-10-01/sulfate-free-surfactants-conditioning-shampoo/, XP002804301, Jan. 10, 2019, 1 page. cited by applicant

Inspection certificate for Hostapon® CCG, Clariant Ibérica Production, S.A., May 6, 2019; p. 1-2. cited by applicant

McGinley et al. American Association of Textile Chemists and Colorists, 2017, 17 pages, (2017). cited by applicant

McGinley et al. Performance Verification of Air Freshener Products and Other Odour Control Devices for Indoor Air Quality Malodours. Presented at the 8th Workshop on Odour and Emissions of Plastic Materials Universitat Kassel Institut for Werkstofftechnik Kassel, Germany, Mar. 27-28, 2006, 13 pages. cited by applicant

Medvedev, Diffusion Coefficients in Multicomponent Mixtures, PhD Thesis from Technical University of Denmark, dated 2005, 181 pages. cited by applicant

Mintel GNPD Base, Bright Blonde Shampoo Record No. 3412889 Feb. 29, 2016 ; 2 pages. cited by applicant

Mintel GNPD Base, Mineral Conquer Blonde Silver Shampoo Record No. 3953107 Apr. 30, 2016; 2 pages. cited by applicant

Mintel GNPD Base, Royal Treatment Collection, Record No. 1946223 dated Dec. 31, 2011, 3 pages. cited by applicant

Morioka, H. et al. "Effects of Zinc on the New Preparation Method of Hydroxy Double Salts" Inorg. Chem.

1999, 38, 4211-6. cited by applicant

Musazzi, “Emulsion versus nonoemulsion: how much is the formulative shift critical for a cosmetic product?” (Drug Deliv. and Trans. Res. (2018) 8: pp. 414-421 (Year: 2018). cited by applicant

Natural oils: why specific carbon chains are chosen for certain surfactant properties, Chemlink, URL Link: <https://www.chemlink.co.uk/natural-oils-why-specific-carbon-chains-are-chosen-for-certain-surfactant-properties/a> (Year: 2022), 4 pgs. cited by applicant

Naturally Rich Moisturizing Shampoo, Database accession No. 6421011, Mar. 27, 2019, 3 pages. cited by applicant

Noritomi H. Formation and Solubilization Property of Water-in-Oil Microemulsions of Alkyl Glucosides. Advances in Nanoparticles, 2013, 2, 366-371 (Year: 2013). cited by applicant

Parchem fine & specialty chemicals. MIPA-laureth sulfate supplier distributor—CAS 83016-76-6; dated 2021; pp. 1-7. cited by applicant

PERM Inc, , Diffusion Coefficient: Measurement Techniques, <https://perminc.com/resources/fundamentals-of-fluid-flow-in-porous-media/chapter-3-molecular-diffusion/diffusion-coefficient/measurement-techniques>, dated Oct. 2020; p. 1-4. cited by applicant

Polyquaternium: “Final Report on the Safety Assessment of the Polyquaternium-10”, Journal of the American College of Toxicology, Jan. 1, 1988, URL: <http://www.beauty-review.nl/wp-content/uploads/2015/02/Final-Report-on-theSafety-Assessment-of-Polyquaternium-10.pdf>, retrieved on Dec. 20, 2018; 9 pages. cited by applicant

Practical Modern Hair Science, Published 2012; 43 pages. cited by applicant

Product Bulletin, Amphosol® CG, Cocamidopropyl Betaine, Stepan Company, Jun. 2011; 1-2 pages. cited by applicant

Product Data Sheet for Chemoryl™LS Surfactant, Sodium Lauroyl Sarcosinate, Lubrizol Advanced Materials, Inc., Mar. 24, 2020; 1-2 pages. cited by applicant

Product Data Sheet, Eversoft™ UCS-40S, Disodium Cocoyl Glutamate (Sodium Cocoyl Glutamate*), Sino Lion USA, Jul. 2018; 2 pages. cited by applicant

Product Fact Sheet—Hostapon® CCG, mild anionic surfactant for the cosmetic industry, Clariant International Ltd., Aug. 2014 ; 1-3 pages. cited by applicant

Product Fact Sheet, Hostapon® CGN, Mild anionic surfactant for the cosmetic industry, Clariant International Ltd., Jan. 2016; 1-2 pages. cited by applicant

Rajendran A. et al: “Study on the Analysis of Trace Elements in Aloe vera and Its Biological Importance Study on the Analysis of Trace Elements in Aloe vera and Its Biological Importance”, Journal of Applied Sciences Research, Jan. 1, 2007 (Jan. 1, 2007), XP055799133, pp. 1476-1478. cited by applicant

Robinson et al., Final Report of the Amended Safety Assessment of Sodium Laureth Sulfate and Related Salts of Sulfated Ethoxylated Alcohols, International Journal of Toxicology 29 (Supplement 3); dated 2010; pp. 151S-161S. cited by applicant

S. Herrwerth et al.: “Highly Concentrated Cocamidopropyl Betaine—The Latest Developments for Improved Sustainability and Enhanced Skin Care”, Tenside, Surfactants, Detergents, vol. 45, No. 6, dated Nov. 1, 2008, pp. 304-308, p. 305—left-hand column; 3 pages. cited by applicant

Safety assessment of amino acid alkyl amides used in cosmetics , dated Sep. 20, 2013, 46 pages. cited by applicant

Schaefer, Katie, “Eco-friendly, Non-flammable Liquified Gas Propellant”, <https://www.cosmeticsandtoiletries.com/formulating/function/aids/138418589.html#close-olyticsmodal>. Published Jan. 30, 2012; 1-2 pages. cited by applicant

Sensory., “A Review of The Science and Technology of Odor Measurement”, Prepared for the Air Quality Bureau of the Iowa Department of Natural Resources, Dec. 30, 2005 51 pages. cited by applicant

Shampoo C, Database accession No. 1632217, Sep. 29, 2011, 3 pages. cited by applicant

Softazoline CL-R, Kawaken Singapore PTE Ltd. Website printout from <http://kawaken.com.sg/softazoline-ch-r/a>, accessed on Nov. 30, 2022. cited by applicant

Todd et al., Volatile Silicone Fluids for Cosmetics, Cosmetics and Toiletries, vol. 91, pp. 27-32 (Jan. 1976). cited by applicant

UL Prospector® Product Data Sheet, Plantacare® 818 UP, C8-16 fatty alcohol glucoside, BASF, dated May 21, 2015; 1-3 pages. cited by applicant

Unhale Shrikrushna Subhash et al: Formulation and Development of Sulphate Free Shampoo About an Updates and Guidelines of Corona Virus View project health and beauty science View project Rohit Bhavsar Reliance Industries Limited; International Journal for Research in Applied Science & Engineering Technology, Apr. 1, 2020 (Apr. 1, 2020) XP055842327, DOI: 10.22214, 14 pages. cited by applicant

“Deep Image Matting”, Ning Xu et al, Beckman Institute for Advanced Science and Technology, University of Illinois at Urbana-Champaign, Adobe Research, dated Mar. 10, 2017; 10 pages. cited by applicant

Anonymous: “Anti-Dandruff Scalp Care Shampoo”, MINTEL, Database accession No. 301924, Sep. 16, 2004, 2 pages. cited by applicant

“Personal care solutions Guide”, Solvay, Publication date: May 2018, 84 pages. cited by applicant

Product Data Record Tego® Betain F KB 5, dated Jul. 1, 2015, 4 pages. cited by applicant

16158 PCT Search Report and Written Opinion for PCT/US2022/028256 dated Oct. 7, 2022, 15 pages. cited by applicant

PubChem CID 3033856 for decyl glucoside. downloaded Jun. 22, 2023, 32 pages. (Year: 2023). cited by applicant

“Comparative Study on the Chemical constituents of Aloe Vera and Aloe Kula in China”, Zhang Xiaohua et al., Flavor Cosmetics, No. 63, dated Dec. 31, 2000, pp. 7-11. cited by applicant

Shampoo, ID# 6148479. Mintel GNPD [online], URL: <http://www.gnpd.com>, Nov. 2018, 4 pages. cited by applicant

Anonymous, “Medicated Cleanser”, Nioxin Research Laboratories, Mintel GNPD [online], ID:1060983, URL: <http://www.gnpd.com>, dated Feb. 2009, 4 pages. cited by applicant

P. A. Cornwell, “A review of shampoo surfactant technology: consumer benefits, raw materials and recent developments”, International Journal of Cosmetic Science, 40, Oct. 27, 2017, pp. 16-30. cited by applicant

Sbhatu et al, “Formulation and Physicochemical Evaluation of Lab-Based Aloe adigratana Reynolds Shampoos”, Hindawi, International Journal of Analytical Chemistry, vol. 2020, Article ID 6290617, URL: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7160718/>, Published Apr. 4, 2020, 7 pages. cited by applicant

Shampoo Manufacturing Plant, Lodha International, URL: <https://www.lodhapharma.com/shampoo-manufacturing-plant.php#:~:text=Shampoos%20are%20categorized%20as%20viscous,if%20not%20all%20of%20them>, Published: 2018, 4 pages. cited by applicant

“Jaguar® Optima”, Solvay, site: www.solvay.com, year 2023, 1 page. cited by applicant

Cafasso, “What's the proper order to use shampoo and conditioner while bathing”, Healthline, Retrieved from Internet: <https://www.healthline.com/health/beauty-skin-care/shampoo-or-conditioner-first>, dated Jul. 13, 2020, 2 pages. cited by applicant

Duis et al., “Environmental fate and effects of water-soluble synthetic organic polymers used in cosmetic products”, Environmental Sciences Europe, 33:21, Year 2021; 20 pages. cited by applicant

Matsouka et al., “Vesicle formation of disodium lauryl sulfosuccinate”, Journal of Molecular Liquids, 348, 118422, Nov. 1, 2021, 7 pages. cited by applicant

Ramachandra et al., “Processing of Aloe Vera Leaf Gel: A Review”, American Journal of Agricultural and Biological Sciences 3 (2), year 2008, pp. 502-510. cited by applicant

Yang et al., “Synthesis and foaming performance of Lauramidopropyl Betaine Derivate Surfactants”, Materials Science Forum, vol. 953, Jan. 9, 2019, 1 page. cited by applicant

Anonymous, “Healing + Anti-Breakage Shampoo”, ID# 3383875, Mintel Gnpd, URL: <http://www.gnpd.com>, dated Aug. 2015, 3 pages. cited by applicant

Anonymous, “Shampoo”, ID# 1743027, Mintel GNPD, URL: <http://www.gnpd.com>, dated Mar. 2012, 3 pages. cited by applicant

Anonymous; “Advanced Oil Control Anti Dandruff Shampoo”, XP093149356, Database GNPD Mintel, Database accession No. 9347772, dated Feb. 1, 2022, 03 Pages. cited by applicant

Bryant Taylor, “Here's What You Need To Know About The Latest Korean Beauty Obsession”, Retrieved from Internet: <https://www.refinery29.com/en-us/scalp-ph-facts>, Jun. 5, 2016, 8 pages. cited by applicant

Goh et al., “New cationic conditioning polymers for hair care”, Retrieved from Internet: <https://cosmeticsbusiness.com/new-cationic-conditioning-polymers-for-hair-care-47152>, Dec. 18, 2006, 10 pages. cited by applicant

Harvey Gedeon; Journal of Cosmetic Science, “The Challenges Facing Today's Cosmetic Chemist”, 57, 2005,

73-94 (22 pages). cited by applicant
Hossel et al. "Conditioning polymers in today's shampoo formulations-efficacy, mechanism and test methods", International Journal of Cosmetic Science. 2000, vol. 1, 2000, pp. 1-10. cited by applicant
Manuel, F. et al., "A new postulate on two stages of dandruff: A clinical perspective", International Journal of Trichology, 2011, vol. 3, 3-6 (Year: 2011). cited by applicant
Tarun et al. "Evaluation of pH of Bathing Soaps and Shampoos for Skin and Hair Care", Indian Journal of Dermatology. 2014, vol. 59, No. 5, No Known Date, 6 Pages. cited by applicant
Technical Information; TEGO® Betain F KB 5 / TEGO® Betain F KM 1; Mild amphoteric surfactants: www.evonik.com/personal-care; Jul. 2010; 2 pages; Product Specification; 5 pages. cited by applicant
"Selecting the Right Surfactant for Cosmetics", Retrieved from Internet: <https://cosmetics.specialchem.com/centers/selecting-the-right-surfactantforcosmetics/surfactantsinhaircareapplications#:~:text=Amphoteric%20surfactants%20are%20used%20to,be%20based%20ntirely%20on%20amphoterics>, Aug. 11, 2020, 4 pages. cited by applicant
"UltraPureGel", Retrieved from Internet: <https://www.ingredientstodiefor.com/item/UltraPureGel/1264,pages>, cited by applicant

Primary Examiner: Packard; Benjamin J

Assistant Examiner: Tien; Lucy M

Background/Summary

FIELD OF THE INVENTION

(1) The present invention relates to a shampoo composition, in particular a phase-stable shampoo composition with a surfactant system that is free of or substantially free of sulfate-based surfactants and a thickener comprising sclerotium gum.

BACKGROUND OF THE INVENTION

(2) Historically, most commercial cleansing compositions, such as shampoo compositions, contain sulfate-based surfactant systems because they provide effective cleaning and a good user experience. Sulfate-based surfactant systems generally have acceptable viscosity making it easy to apply and distribute the shampoo composition throughout a user's hair. In addition, sulfate-based surfactant systems can generally be paired with cationic polymers that can form coacervate with the sulfate-based surfactant system during use thereby providing a shampoo with effective conditioning benefits.

(3) Some consumers, especially those with color-treated or otherwise treated hair, may prefer a shampoo composition that is substantially free of sulfate-based surfactant systems. However, it can be difficult to formulate a shampoo with non-sulfate-based surfactants with sufficient viscosity. Many current sulfate-free shampoos have a viscosity that is too low, which makes it difficult to hold in a user's hand and apply across the hair and scalp. These consumers may also want conditioning polymers in their shampoo because higher conditioning shampoos feel less stripping to the hair. However, it can be difficult to formulate a shampoo with a sulfate-free surfactant system that also has cationic polymers that deliver a conditioning benefit because the shampoo can have low viscosity, which makes it difficult for a user to hold the liquid shampoo composition in their palm and distribute across their hair and scalp.

(4) When traditional anionic thickeners, such as xanthan gum, carrageenan, konjac gum, and gellan gum are added to shampoo compositions containing sulfate-free surfactants and cationic polymers, the composition can be unstable and separate into multiple phases. The traditional anionic thickener can complex with the cationic polymer, thereby reducing active deposition, which decreases wet conditioning and causes phase separation.

(5) Therefore, there is a need for a stable shampoo composition with a sufficient viscosity and superior product performance that contains one or more non-sulfated anionic surfactants.

SUMMARY OF THE INVENTION

(6) A stable shampoo composition comprising: (a) a surfactant system comprising: (i) from about 3 to

about 35% of an anionic surfactant; (ii) from about 5 to about 15% of an amphoteric surfactant; wherein the surfactant system is substantially free of sulphated surfactants; (b) from about 0.01% to about 2% of a cationic polymer; (c) from about 0.15% to about 1.5% of a thickener comprising sclerotium gum; wherein the composition comprises a viscosity greater than 2500 cP and a pH greater than 5.5.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) FIG. 1 shows a typical sclerotium gum structure;
- (2) FIG. 2 shows a β -(1,3)- β -(1,6) glucan structure;
- (3) FIG. 3 shows a tridimensional conformation of scleroglucan triplex;
- (4) FIG. 4 is a 20 $\times$ micrograph of a shampoo that contains in situ coacervate;
- (5) FIG. 5 is 10 $\times$ micrograph of the shampoo composition of FIG. 1;
- (6) FIG. 6A is an image of Comparative Example C5; and
- (7) FIG. 6B is an image of Comparative Example C6.

DETAILED DESCRIPTION OF THE INVENTION

(8) Some consumers prefer shampoo compositions that are substantially free of sulfate-based surfactants. These consumers may also prefer shampoos that feel less stripping to the hair and therefore may prefer a shampoo with cationic conditioning polymers. However, it is difficult to formulate these compositions to be stable and have a consumer preferred viscosity. They are often too thin, which makes them difficult to hold in a user's palm and distribute across their hair and scalp.

(9) There are at least three common ways to thicken shampoo compositions: 1. Add an anionic thickener (e.g. xanthan gum, carrageenan, konjac gum, gellan gum) 2. Decrease the pH 3. Add inorganic salts (e.g. NaCl)

(10) However, these methods are not effective in a shampoo composition that contains sulfate-free anionic surfactants and a cationic conditioning polymer. First, when traditional anionic thickeners are added, it was found that the composition becomes unstable, impacting product appearance and performance. Second, when the pH was decreased it was found that many sulfate-free surfactant systems can hydrolyze at low pH resulting in viscosity and performance changes over time and will eventually lead to phase separation. Third, when inorganic salt was added to the liquid shampoo becomes unstable due to formation of a gel-like complex known as coacervate in the composition (referred to herein as “in situ coacervate” or an “in situ coacervate phase,” which is a coacervate that forms in the composition, prior to dilution), as opposed to when it is diluted with water when a user washes their hair.

(11) It was found that sclerotium gum, which has a unique triple helix structure, could be used to thicken shampoo compositions that contain sulfate-free anionic surfactants and cationic polymers while also maintaining phase stability. It was also found that a thickener comprising sclerotium gum could be used to thicken shampoo compositions that contain sulfate-free surfactants and cationic polymers while also maintaining phase stability and pH (e.g. >5.5), and providing superior wet conditioning.

(12) It was also found that in some examples the amount of inorganic salt in the composition can also impact product stability and viscosity. In some examples, product stability and acceptable viscosity can be achieved by (1) maintaining a low inorganic salt concentration in formulas (e.g. from about 0 to about 1 wt %); and/or (2) having a ratio of polymer charge density to salt of greater than 1.2:1.

(13) One way to lower the inorganic salt concentration is to avoid or minimize adding extra inorganic salt to the formula and/or by using low inorganic salt containing raw materials. For example, commercially available sulfate free surfactants such as disodium cocoyl glutamate typically comes with high levels of inorganic salt such as 5% or higher. Amphoteric surfactant such as betaines or sultaines also typically come with high levels of inorganic salt such as sodium chloride. Use of these high salt containing raw materials in sulfate-free surfactant based cleaning formulations in excess of about 1% total sodium chloride in the formulation can cause formation of undesired coacervate in the product. If the inorganic salt level is lowered in the surfactant raw materials so that total salt in the composition is less than about 1% or lower, a stable 1-phase clear product can be formulated. Whereas, if the regular material with high inorganic salt is used, the product is cloudy, 2-phase, and unstable. The cloudy 2-

phase product is likely the result of in situ coacervate formation.

(14) In some examples, sclerotium gum can be the only thickener in the shampoo composition. In other examples, the thickener can include sclerotium gum, inorganic salt, and combinations thereof. The shampoo composition can be free of or substantially free of anionic thickeners. Non-limiting examples of anionic thickeners that may be excluded from the shampoo compositions can include carboxylic acid polymers, crosslinked polyacrylate polymers, polyacrylamide polymers, polysaccharides, and gums. Non-limiting examples of gums can include acacia, agar, algin, alginic acid, ammonium alginate, amylopectin, calcium alginate, calcium carrageenan, carrageenan, dextrin, gelatin, gellan gum, karaya gum, konjac gum, locust bean gum, natto gum, potassium alginate, potassium carrageenan, propylene glycol alginate, sodium carboxymethyl dextran, sodium carrageenan, tragacanth gum, xanthan gum, and mixtures thereof.

(15) The pH can be greater than 5.5, alternatively greater than 5.7, alternatively greater than 5.8. Alternatively, the pH can be from about 5 to about 8, alternatively from about 5.2 to about 7.5, alternatively from about 5.3 to about 7.25, alternatively from about 5.4 to about 7.1, alternatively from about 5.5 to about 7, alternatively from about 5.6 to about 6.9, alternatively from about 5.7 to about 6.8, and alternatively from about 5.8 to about 6.7. The pH can be from about 4 to about 8, alternatively from about 4.5 to about 7.5, alternatively from about 5 to about 7, alternatively from about 5.5 to about 6.5, alternatively from about 5.5 to about 6, and alternatively from about 6 to about 6.5. The pH is determined by the pH Test Method, described herein.

(16) The shampoo composition can have a viscosity greater than 2500 cP, alternatively greater than 4000 cP, alternatively greater than 4500 cP, and alternatively greater than 5000 cP. The shampoo composition can have a viscosity of about 2500 cP to about 20,000 cP, alternatively from about 3000 cps to about 15,000 cps, alternatively from 4000 cP to about 13,000 cP, alternatively from about 5,000 cP to about 11,000 cP, and alternatively from about 7,000 cP to about 10,000 cP, alternatively from about 2500 cP to about 8500 cP, alternatively 2500 cP to about 6500 cP as measured at 26.6° C., as measured by the Cone/Plate Viscosity Measurement Test Method, described herein.

(17) The shampoo composition can have a yield stress, Herschel-Bulkley @ shear rate 10.sup.-2 to 10.sup.-4 s.sup.-1 of from about 0.001 Pa to about 0.5 Pa, alternatively from about 0.002 Pa to about 0.3 Pa, alternatively from about 0.003 Pa to about 0.2 Pa, alternatively from about 0.004 Pa to about 0.18 Pa, alternatively from about 0.007 Pa to about 0.17 Pa, and alternatively from about 0.008 Pa to about 0.15 Pa. The shampoo composition can have a yield stress, Herschel-Bulkley @ shear rate 10.sup.-2 to 10.sup.-4 s.sup.-1 of greater than 0.003 Pa, alternatively greater than 0.005 Pa, alternatively greater than 0.010 Pa, alternatively greater than 0.015 Pa, and alternatively greater than 0.1 Pa. The yield stress is measured at 26.7° C. by flow sweep at a shear rate 100 to 10.sup.-4 s.sup.-1 using Discovery Hybrid Rheometer (DHR-3) available from TA Instruments. To apply the Herschel-Bulkley model, the TA software to fit the model in the log space at a shear rate from 10.sup.-2 to 10.sup.-4 s.sup.-1 is used. The geometry used to measure the yield stress and viscosity of the cleansing phase is a 60 mm 2° aluminum cone (with a Peltier steel plate). The geometry should be run at the gap specified by the manufacturer for the geometry. Trimming the sample during the initial conditioning step in step 1 is recommended to ensure data integrity and reproducibility. Torque map the geometry prior to running the yield stress or shear stress methods when the instrument+geometry is out of calibration. The version of Trios software used to generate the rheology data herein is TRIOS 5.1.1 In some examples, it was found that a thickener that contained a combination of sclerotium gum and citrus peel fiber (commercially available as FiberDesign™ Sensation from Cargill® with an INCI citrus limon peel powder (and) sclerotium gum) the shampoo composition could have a higher yield stress, which can help suspend actives (e.g. cationic polymers; silicones; non-silicone organic conditioning agents such as botanical hydrocarbon oils, waxes, conditioning polymers; anti-dandruff actives) in the formulation, which can lead to improved product performance. In some examples the thickener can be selected from the group consisting of sclerotium gum, citrus limon peel powder, inorganic salts, and combinations thereof. In other examples the thickener can be selected from the group consisting of sclerotium gum, citrus limon peel powder, and combinations thereof. In other examples, the thickener can contain both sclerotium gum and citrus limon peel powder.

(18) The shampoo composition may have less than 1% of inorganic salt, alternatively less than 0.75%

inorganic salt, alternatively less than 0.5% inorganic salt, alternatively less than 0.25% inorganic salt, alternatively less than or equal to 0.2% inorganic salt. The shampoo composition may have from about 0% to about 1% inorganic salt, alternatively from about 0% to about 0.9% inorganic salt, alternatively from about 0% to about 0.8%, alternatively from about 0% to about 0.5% inorganic salt, alternatively from about 0% to about 0.3% inorganic salt; and alternatively from about 0.05% to about 0.2% inorganic salt. The shampoo composition may have greater than 0% and less than 1% inorganic salt, alternatively greater than 0% and less than 0.75%, alternatively greater than 0% and less than 0.5%, and alternatively greater than 0% and less than 0.25%. The shampoo composition can include from about 0.5% to about 5% inorganic salt, alternatively from about 0.55% to about 4%, alternatively about 0.75% to about 3.5%, alternatively 0.6% to about 3.25%, alternatively from about 0.8% to about 3%, alternatively from about 0.8% to about 2.5%, alternatively from about 1% to about 2%, and alternatively from about 1% to about 1.5%. The shampoo composition can include from about 0.75% to about 1.5% inorganic salt, alternatively from about 0.8% to about 1.4%, alternatively from about 0.9% to about 1.4%. The wt. % inorganic chloride salt can be determined by the Argentometry Method to Measure wt % Inorganic Chloride Salt Test Method, described herein. The inorganic salt can be an inorganic chloride salt. The inorganic salt can enter the composition with the surfactants or other ingredients, or it can be added separately as a thickener. If the inorganic salt enters the composition along with the surfactants, it can still be considered a thickener.

(19) The ratio of polymer charge density to total inorganic salt can be from about 0.5:1 to about 3:1, from about 0.7:1 to about 2.5:1, alternatively 0.75:1 to about 2.25:1, alternatively from about 1:1 to about 2:1, alternatively from about 1.2:1 to about 1.5:1, alternatively from about 1.1:1 to about 1.4:1, and alternatively from about 1.2:1 to about 1.4:1. The ratio of polymer charge density to total inorganic salt can be $\geq 1.1:1$, alternatively $\geq 1.2:1$, alternatively $\geq 1.3:1$, and alternatively $\geq 1.5:1$. The ratio of polymer charge density to total inorganic salt can be $< 15:1$, $< 10:1$, $< 7:1$, $< 5:1$, alternatively $< 4:1$, alternatively $< 3:1$, alternatively $< 2:1$, and alternatively $< 1.5:1$. The ratio of polymer charge density to inorganic salt is the charge density of the cationic polymer (meq/gm) to the wt. % of the inorganic salt, disregarding the units. If the composition contains more than one cationic polymer, then the ratio is calculated according to the polymer with the lowest charge density.

(20) The ratio of anionic surfactant to amphoteric surfactant can be from about 0.25:1 to about 3:1, alternatively from about 0.3:1 to about 2.5:1, alternatively from about 0.4:1 to about 2:1, alternatively from about 0.5:1 to about 1.5:1, alternatively from about 0.6:1 to about 1.25:1, and alternatively from about 0.75:1 to about 1:1. The ratio of anionic surfactant to amphoteric surfactant can be from about 0.4:1 to about 1.25:1, alternatively from about 0.5:1 to about 1.1:1, and alternatively from about 0.6:1 to about 1:1. In some examples, the ratio of anionic surfactant to amphoteric surfactant is less than 2:1, alternatively less than 1.75:1, alternatively less than 1.5:1, alternatively less than 1.1:1, and alternatively less than 1:1.

(21) The shampoo composition can be used to clean and/or condition hair. First, the user dispenses the liquid shampoo composition from the bottle into their hand or onto a cleaning implement. Then, they massage the shampoo into their wet hair. While they are massaging the shampoo composition into the hair the shampoo is diluted and a coacervate can form and the shampoo can lather. After massaging into hair, the shampoo composition is rinsed from the user's hair and at least a portion of the cationic polymers can be deposited on the user's hair, which can provide a wet conditioning benefit. Shampooing can be repeated, if desired, and/or a conditioner can be applied. The conditioner can be a rinse-off conditioner or a leave-in conditioner.

(22) It may be consumer desirable to have a shampoo composition with a minimal level of ingredients. The shampoo composition can be formulated without polymeric thickeners or suspending agents such as carbomer, EGDS or thixcin. The shampoo composition may contain 11 or fewer ingredients, 10 or fewer ingredients, 9 or fewer ingredients, 8 or fewer ingredients, 7 or fewer ingredients, 6 or fewer ingredients. The minimal ingredient formula can include water, anionic surfactant, amphoteric surfactant, cationic polymer, inorganic salt, and perfume. It is understood that perfumes can be formed from one or more materials. In some examples, the composition can be free of or substantially free of fragrance. In another example, the composition can be free of or substantially free of polyethylene glycol (PEG).

(23) The composition can be free of alkyl polyglucoside. Non-limiting examples of alkyl polyglucosides

can include decyl glucoside, cocoyl glucoside, lauroyl glucoside and combinations thereof. It has been found that alky polyglucosides can impact lather properties and thickening of shampoo compositions containing sulfate-free anionic surfactants and cationic conditioning polymers.

(24) The composition can be substantially free of or free of fatty esters having at least 10 carbon atoms. The composition can be substantially free of or free of fatty esters. The composition can be substantially free of or free of fatty esters with hydrocarbyl chains derived from fatty acids or alcohols. The composition can be substantially free of or free of oligomeric or polymeric esters, prepared from unsaturated glyceryl esters can also be used as conditioning materials.

(25) While the specification concludes with claims particularly pointing out and distinctly claiming the invention, it is believed that the present disclosure will be better understood from the description.

(26) As used herein, "cleansing composition" includes personal cleansing products such as shampoos, conditioners, conditioning shampoos, shower gels, liquid hand cleansers, facial cleansers, and other surfactant-based liquid compositions.

(27) The shampoo composition can be clear prior to dilution with water. The term "clear" or "transparent" as used herein, means that the compositions have a percent transparency (% T) of at least about 70% transmittance at 600 nm. The % T may be at 600 nm from about 70% to about 100%, from about 75% to about 100%, from about 80% to about 100%, from about 85% to about 100%, from about 90% to about 100%, from about 95% to about 100%. In the present invention, the percent transparency (% T) may be at least about 80% transmittance at 600 nm; percent transparency (% T) may be at least about 90% transmittance at 600 nm.

(28) As used herein, the term "fluid" includes liquids and gels.

(29) As used herein, "molecular weight" or "M.Wt." refers to the weight average molecular weight unless otherwise stated. Molecular weight is measured using industry standard method, gel permeation chromatography ("GPC"). The molecular weight has units of grams/mol.

(30) As used herein, "substantially free" refers to less than 0.5%, alternatively less than 0.25%, alternatively less than 0.1%, alternatively less than 0.05%, alternatively less than 0.02%, and alternatively less than 0.01%.

(31) As used herein, "sulfate free" and "substantially free of sulfates" means essentially free of sulfate-containing compounds except as otherwise incidentally incorporated as minor components.

(32) Sulfate free contains no detectable sulfated surfactants.

(33) As used herein, "sulfated surfactants" or "sulfate-based surfactants" means surfactants which contain a sulfate group. The term "substantially free of sulfated surfactants" or "substantially free of sulfate-based surfactants" means essentially free of surfactants containing a sulfate group except as otherwise incidentally incorporated as minor components.

(34) All percentages, parts and ratios are based upon the total weight of the compositions of the present invention, unless otherwise specified. All such weights as they pertain to listed ingredients are based on the active level and, therefore, do not include carriers or by-products that may be included in commercially available materials.

(35) Unless otherwise noted, all component or composition levels are in reference to the active portion of that component or composition, and are exclusive of impurities, for example, residual solvents or by-products, which may be present in commercially available sources of such components or compositions.

(36) It should be understood that every maximum numerical limitation given throughout this specification includes every lower numerical limitation, as if such lower numerical limitations were expressly written herein. Every minimum numerical limitation given throughout this specification will include every higher numerical limitation, as if such higher numerical limitations were expressly written herein. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

(37) Surfactant

(38) The cleansing compositions described herein can include one or more surfactants in the surfactant system. The one or more surfactants can be substantially free of sulfate-based surfactants. As can be appreciated, surfactants provide a cleaning benefit to soiled articles such as hair, skin, and hair follicles by facilitating the removal of oil and other soils. Surfactants generally facilitate such cleaning due to

their amphiphilic nature which allows for the surfactants to break up, and form micelles around, oil and other soils which can then be rinsed out, thereby removing them from the soiled article. Suitable surfactants for a cleansing composition can include anionic moieties to allow for the formation of a coacervate with a cationic polymer. The surfactant can be selected from anionic surfactants, amphoteric surfactants, zwitterionic surfactants, non-ionic surfactants, and combinations thereof.

(39) Cleansing compositions typically employ sulfate-based surfactant systems (such as, but not limited to, sodium lauryl sulfate) because of their effectiveness in lather production, stability, clarity and cleansing. The cleansing compositions described herein are substantially free of sulfate-based surfactants. "Substantially free" of sulfate based surfactants as used herein means from about 0% to about 3%, alternatively from about 0% to about 2%, alternatively from about 0% to about 1%, alternatively from about 0% to about 0.5%, alternatively from about 0% to about 0.25%, alternatively from about 0% to about 0.1%, alternatively from about 0% to about 0.05%, alternatively from about 0% to about 0.01%, alternatively from about 0% to about 0.001%, and/or alternatively free of sulfates. As used herein, "free of" means 0%.

(40) Additionally, the surfactants can be added to the composition as a solution, instead of the neat material and the solution can include inorganic salts that can be added to the formula. The surfactant formula can have inorganic salt that can be from about 0% to about 2% of inorganic salts of the final composition, alternatively from about 0.1% to about 1.5%, and alternatively from about 0.2% to about 1%.

(41) Suitable surfactants that are substantially free of sulfates can include sodium, ammonium or potassium salts of isethionates; sodium, ammonium or potassium salts of sulfonates; sodium, ammonium or potassium salts of ether sulfonates; sodium, ammonium or potassium salts of sulfosuccinates; sodium, ammonium or potassium salts of sulfoacetates; sodium, ammonium or potassium salts of glycines; sodium, ammonium or potassium salts of sarcosines; sodium, ammonium or potassium salts of glutamates; sodium, ammonium or potassium salts of alanines; sodium, ammonium or potassium salts of carboxylates; sodium, ammonium or potassium salts of taurates; sodium, ammonium or potassium salts of phosphate esters; and combinations thereof.

(42) The concentration of the surfactant in the composition should be sufficient to provide the desired cleaning and lather performance. The cleansing composition can include a total surfactant level of from about 5% to about 50%, alternatively from about 8% to about 40%, alternatively from about 10% to about 30%, alternatively from about 12% to about 25%, alternatively from about 13% to about 23%, alternatively from about 14% to about 21%, alternatively from about 15% to about 20%.

(43) The cleansing composition can include from about 3% to about 30% anionic surfactant, alternatively from about 4% to about 20%, alternatively from about 5% to about 15%, alternatively from about 6% to about 12%, and alternatively from about 7% to about 10%. The cleansing composition can include from about 3% to about 40% amphoteric surfactant, alternatively from about 4% to about 30%, alternatively from about 5% to about 25%, alternatively from about 6% to about 18%, alternatively from about 7% to about 15%, alternatively from about 8% to about 13%, and alternatively from about 9% to about 11%.

(44) The ratio of anionic surfactant to amphoteric surfactant can be from about 0.25:1 to about 3:1, alternatively from about 0.3:1 to about 2.5:1, alternatively from about 0.4:1 to about 2:1, alternatively from about 0.5:1 to about 1.5:1, alternatively from about 0.6:1 to about 1.25:1, and alternatively from about 0.75:1 to about 1:1. The ratio of anionic surfactant to amphoteric surfactant can be from about 0.4:1 to about 1.25:1, alternatively from about 0.5:1 to about 1.1:1, and alternatively from about 0.6:1 to about 1:1. In some examples, the ratio of anionic surfactant to amphoteric surfactant is less than 2:1, alternatively less than 1.75:1, alternatively less than 1.5:1, alternatively less than 1.1:1, and alternatively less than 1:1.

(45) In some examples, inorganic salt is added to the shampoo composition with the surfactant raw materials. In one example, the surfactant raw materials include less than 1.5% inorganic salt, alternatively less than 1.25%, alternatively less than 1%, alternatively less than 0.7%, alternatively less than 0.5%, alternatively less than 0.25%, alternatively less than 0.2%, alternatively less than 0.15%, alternatively less than or equal to 0.1%. In some examples, at least 0.05% inorganic salt is added to the formula via the surfactant raw materials, alternatively at least 0.07%, and alternatively at least 0.1%.

(46) The surfactant system can include one or more amino acid based anionic surfactants. Non-limiting examples of amino acid based anionic surfactants can include sodium, ammonium or potassium salts of acyl glycines; sodium, ammonium or potassium salts of acyl sarcosinates; sodium, ammonium or potassium salts of acyl glutamates; sodium, ammonium or potassium salts of acyl alaninates and combinations thereof.

(47) The amino acid based anionic surfactant can be a glutamate, for instance an acyl glutamate. Non-limiting examples of acyl glutamates can be selected from the group consisting of sodium cocoyl glutamate, disodium cocoyl glutamate, ammonium cocoyl glutamate, diammonium cocoyl glutamate, sodium lauroyl glutamate, disodium lauroyl glutamate, sodium cocoyl hydrolyzed wheat protein glutamate, disodium cocoyl hydrolyzed wheat protein glutamate, potassium cocoyl glutamate, dipotassium cocoyl glutamate, potassium lauroyl glutamate, dipotassium lauroyl glutamate, potassium cocoyl hydrolyzed wheat protein glutamate, dipotassium cocoyl hydrolyzed wheat protein glutamate, sodium capryloyl glutamate, disodium capryloyl glutamate, potassium capryloyl glutamate, dipotassium capryloyl glutamate, sodium undecylenoyl glutamate, disodium undecylenoyl glutamate, potassium undecylenoyl glutamate, dipotassium undecylenoyl glutamate, disodium hydrogenated tallow glutamate, sodium stearyl glutamate, disodium stearyl glutamate, potassium stearyl glutamate, dipotassium stearyl glutamate, sodium myristoyl glutamate, disodium myristoyl glutamate, potassium myristoyl glutamate, dipotassium myristoyl glutamate, sodium cocoyl/hydrogenated tallow glutamate, sodium cocoyl/palmoil/sunfloweroil glutamate, sodium hydrogenated tallowoil Glutamate, sodium olivoyl glutamate, disodium olivoyl glutamate, sodium palmoil glutamate, disodium palmoil Glutamate, TEA-cocoyl glutamate, TEA-hydrogenated tallowoil glutamate, TEA-lauroyl glutamate, and mixtures thereof.

(48) The amino acid based anionic surfactant can be an alaninate, for instance an acyl alaninate. Non-limiting example of acyl alaninates can include sodium cocoyl alaninate, sodium lauroyl alaninate, sodium N-dodecanoyl-L-alaninate and combination thereof.

(49) The amino acid based anionic surfactant can be a sarcosinate, for instance an acyl sarcosinate. Non-limiting examples of sarcosinates can be selected from the group consisting of sodium lauroyl sarcosinate, sodium cocoyl sarcosinate, sodium myristoyl sarcosinate, TEA-cocoyl sarcosinate, ammonium cocoyl sarcosinate, ammonium lauroyl sarcosinate, dimer dilinoleyl bis-lauroylglutamate/lauroylsarcosinate, disodium lauroamphodiacetate lauroyl sarcosinate, isopropyl lauroyl sarcosinate, potassium cocoyl sarcosinate, potassium lauroyl sarcosinate, sodium cocoyl sarcosinate, sodium lauroyl sarcosinate, sodium myristoyl sarcosinate, sodium oleoyl sarcosinate, sodium palmitoyl sarcosinate, TEA-cocoyl sarcosinate, TEA-lauroyl sarcosinate, TEA-oleoyl sarcosinate, TEA-palm kernel sarcosinate, and combinations thereof.

(50) The amino acid based anionic surfactant can be a glycinate for instance an acyl glycinate. Non-limiting example of acyl glycines can include sodium cocoyl glycinate, sodium lauroyl glycinate and combination thereof.

(51) The composition can contain additional anionic surfactants selected from the group consisting of sulfosuccinates, isethionates, sulfonates, sulfoacetates, glucose carboxylates, alkyl ether carboxylates, acyl taurates, and mixture thereof.

(52) Non-limiting examples of sulfosuccinate surfactants can include disodium N-octadecyl sulfosuccinate, disodium lauryl sulfosuccinate, diammonium lauryl sulfosuccinate, sodium lauryl sulfosuccinate, disodium laureth sulfosuccinate, tetrasodium N-(1,2-dicarboxyethyl)-N-octadecyl sulfosuccinate, diamyl ester of sodium sulfosuccinic acid, dihexyl ester of sodium sulfosuccinic acid, dioctyl esters of sodium sulfosuccinic acid, and combinations thereof. The composition can comprise a sulfosuccinate level from about 2% to about 22%, by weight, from about 3% to about 19%, by weight, 4% to about 17%, by weight, and/or from about 5% to about 15%, by weight. Suitable isethionate surfactants can include the reaction product of fatty acids esterified with isethionic acid and neutralized with sodium hydroxide. Suitable fatty acids for isethionate surfactants can be derived from coconut oil or palm kernel oil including amides of methyl tauride. Non-limiting examples of isethionates can be selected from the group consisting of sodium lauroyl methyl isethionate, sodium cocoyl isethionate, ammonium cocoyl isethionate, sodium hydrogenated cocoyl methyl isethionate, sodium lauroyl isethionate, sodium cocoyl methyl isethionate, sodium myristoyl isethionate, sodium oleoyl isethionate, sodium oleyl methyl isethionate, sodium palm kerneloil isethionate, sodium stearyl methyl isethionate,

and mixtures thereof.

(53) Non-limiting examples of sulfonates can include alpha olefin sulfonates, linear alkylbenzene sulfonates, sodium laurylglucosides hydroxypropylsulfonate and combination thereof.

(54) Non-limiting examples of sulfoacetates can include sodium lauryl sulfoacetate, ammonium lauryl sulfoacetate and combination thereof.

(55) Non-limiting example of glucose carboxylates can include sodium lauryl glucoside carboxylate, sodium cocoyl glucoside carboxylate and combinations thereof.

(56) Non-limiting example of alkyl ether carboxylate can include sodium laureth-4 carboxylate, laureth-5 carboxylate, laureth-13 carboxylate, sodium C12-13 pareth-8 carboxylate, sodium C12-15 pareth-8 carboxylate and combination thereof.

(57) Non-limiting example of acyl taurates can include sodium methyl cocoyl taurate, sodium methyl lauroyl taurate, sodium methyl oleoyl taurate, sodium caproyl methyltaurate and combination thereof.

(58) The surfactant system may further comprise one or more amphoteric surfactants and the amphoteric surfactant can be selected from the group consisting of betaines, sultaines, hydroxysultanes, amphohydroxypropyl sulfonates, alkyl amphoacetates, alkyl amphodiacetates, alkyl amphopropionates and combination thereof.

(59) Examples of betaine amphoteric surfactants can include coco dimethyl carboxymethyl betaine, cocoamidopropyl betaine (CAPB), cocobetaine, lauryl amidopropyl betaine (LAPB), oleyl betaine, coco-betaine, cetyl betaine, lauryl dimethyl carboxymethyl betaine, lauryl dimethyl alphacarboxyethyl betaine, cetyl dimethyl carboxymethyl betaine, lauryl bis-(2-hydroxyethyl) carboxymethyl betaine, stearyl bis-(2-hydroxypropyl) carboxymethyl betaine, oleyl dimethyl gamma-carboxypropyl betaine, lauryl bis-(2-hydroxypropyl)alpha-carboxyethyl betaine, and mixtures thereof. Examples of sulfobetaines can include coco dimethyl sulfopropyl betaine, stearyl dimethyl sulfopropyl betaine, lauryl dimethyl sulfoethyl betaine, lauryl bis-(2-hydroxyethyl) sulfopropyl betaine and mixtures thereof.

(60) Non-limiting example of alkylamphoacetates can include sodium cocoyl amphoacetate, sodium lauroyl amphoacetate and combination thereof.

(61) The amphoteric surfactant can comprise cocamidopropyl betaine (CAPB), lauramidopropyl betaine (LAPB), and combinations thereof.

(62) The surfactant system may further comprise one or more non-ionic surfactants and the non-ionic surfactant can be selected from the group consisting alkyl polyglucoside, alkyl glycoside, acyl glucamide and mixture thereof. Non-limiting examples of alkyl polyglucosides can include decyl glucoside, cocoyl glucoside, lauroyl glucoside and combination thereof.

(63) Non-limiting examples of acyl glucamide can include lauroyl/myristoyl methyl glucamide, capryloyl/caproyl methyl glucamide, lauroyl/myristoyl methyl glucamide, cocoyl methyl glucamide and combinations thereof.

(64) The composition can contain a non-ionic deterative surfactants that can include cocamide, cocamide methyl MEA, cocamide DEA, cocamide MEA, cocamide MIPA, lauramide DEA, lauramide MEA, lauramide MIPA, myristamide DEA, myristamide MEA, PEG-20 cocamide MEA, PEG-2 cocamide, PEG-3 cocamide, PEG-4 cocamide, PEG-5 cocamide, PEG-6 cocamide, PEG-7 cocamide, PEG-3 lauramide, PEG-5 lauramide, PEG-3 oleamide, PPG-2 cocamide, PPG-2 hydroxyethyl cocamide, and mixtures thereof.

(65) Sclerotium Gum

(66) The shampoo composition comprises 0.15 wt % to 1.05 wt % sclerotium gum, 0.15 wt % to 1.0 wt % sclerotium gum, 0.2 wt % to 0.8 wt % sclerotium gum, 0.4 wt % to 0.8 wt % sclerotium gum, and/or 0.4 wt % to 0.6 wt % sclerotium gum, and any combination thereof. Sclerotium gum is also called scleroglucan, and it is a branched polysaccharide. In some instances, the primary structure of the scleroglucan consists of glucose molecules linked by β -(1,3) linkage, and every third glucose molecule in the primary structure contains an additional glucose molecule linked by a β -(1,6) linkage. In certain solutions, scleroglucan forms a triple helix shape.

(67) FIG. 1 shows an example of a typical sclerotium gum structure. FIG. 2 shows an example of a β -(1,3)- β -(1,6) glucan structure exhibiting the (3:1) side branching ratio of scleroglucan (Martin et al., 2007). FIG. 3 shows an example of a tridimensional conformation of a scleroglucan triplex (Crescenzi et al., 1988). Specific examples of sclerotium gum include Amigum ER commercially available from Alban

Muller, Actigum CS 11 QD commercially available from Cargill.

(68) Cationic Polymer

(69) A cleansing composition can include a cationic polymer to allow formation of a coacervate. As can be appreciated, the cationic charge of a cationic polymer can interact with an anionic charge of a surfactant to form the coacervate. Suitable cationic polymers can include: (a) a cationic guar polymer, (b) a cationic non-guar galactomannan polymer, (c) a cationic starch polymer, (d) a cationic copolymer of acrylamide monomers and cationic monomers, (e) a synthetic, non-crosslinked, cationic polymer, which may or may not form lyotropic liquid crystals upon combination with the detergent surfactant, and (f) a cationic cellulose polymer. In certain examples, more than one cationic polymer can be included.

(70) A cationic polymer can be included by weight of the cleansing composition at about 0.05% to about 3%, about 0.075% to about 2.0%, or at about 0.1% to about 1.0%. Cationic polymers can have cationic charge densities of about 0.6 meq/g or more, of about 0.9 meq/g or more, about 1.2 meq/g or more, and about 1.5 meq/g or more. However, cationic charge density can also be about 7 meq/g or less and alternatively about 5 meq/g or less. Cationic polymers can have cationic charge densities of from about 0.2 meq/g to about 2.2 meq/g, from about 0.3 meq/g to about 2.0 meq/g, from about 0.4 meq/g to about 1.8 meq/g; from about 0.5 meq/g to about 1.7 meq/g and from about 0.6 meq/g to about 1.3. In some examples the composition can include a cationic polymer with charge density of about 1.7 to about 2.1 meq/g and about 1 to about 1.5% total inorganic salt. The charge densities can be measured at the pH of intended use of the cleansing composition. (e.g., at about pH 3 to about pH 9; or about pH 4 to about pH 8). The average molecular weight of cationic polymers can generally be between about 10,000 and 10 million, between about 50,000 and about 5 million, and between about 100,000 and about 3 million, and between about 300,000 and about 3 million and between about 100,000 and about 2.5 million. Low molecular weight cationic polymers can be used. Low molecular weight cationic polymers can have greater translucency in the liquid carrier of a cleansing composition. The cationic polymer can be a single type, such as the cationic guar polymer guar hydroxypropyltrimonium chloride having a weight average molecular weight of about 2.5 million g/mol or less, and the cleansing composition can have an additional cationic polymer of the same or different types.

(71) Charge density of cationic polymers other than cationic guar polymers can be determined by measuring % Nitrogen. % Nitrogen is measured using USP <461> Method II. % Nitrogen can then be converted to Cationic Polymer Charge Density by calculations known in the art.

(72) The charge density of cationic guar polymers can be calculated as follows: first, calculate the degree of substitution, as disclosed in WO 2019/096601, page 3, lines 4-22, and then cationic charge density can be calculated from the degree of substitution, as described in WO 2013/011122, page 8, lines 8-17, the disclosure of these publications are incorporated by reference.

(73) Cationic Guar Polymer

(74) The cationic polymer can be a cationic guar polymer, which is a cationically substituted galactomannan (guar) gum derivative. Suitable guar gums for guar gum derivatives can be obtained as a naturally occurring material from the seeds of the guar plant. As can be appreciated, the guar molecule is a straight chain mannan which is branched at regular intervals with single membered galactose units on alternative mannose units. The mannose units are linked to each other by means of $\beta(1-4)$ glycosidic linkages. The galactose branching arises by way of an $\alpha(1-6)$ linkage. Cationic derivatives of the guar gums can be obtained through reactions between the hydroxyl groups of the polygalactomannan and reactive quaternary ammonium compounds. The degree of substitution of the cationic groups onto the guar structure can be sufficient to provide the requisite cationic charge density described above.

(75) A cationic guar polymer can have a weight average molecular weight ("M.Wt.") of less than about 3 million g/mol, and can have a charge density from about 0.05 meq/g to about 2.5 meq/g. Alternatively, the cationic guar polymer can have a weight average M.Wt. of less than 1.5 million g/mol, from about 150 thousand g/mol to about 1.5 million g/mol, from about 200 thousand g/mol to about 1.5 million g/mol, from about 300 thousand g/mol to about 1.5 million g/mol, and from about 700,000 thousand g/mol to about 1.5 million g/mol. The cationic guar polymer can have a charge density from about 0.2 meq/g to about 2.2 meq/g, from about 0.3 meq/g to about 2.0 meq/g, from about 0.4 meq/g to about 1.8 meq/g, from about 0.5 meq/g to about 1.7 meq/g, and from about 0.6 meq/g to about 1.3 meq/g.

(76) A cationic guar polymer can have a weight average M.Wt. of less than about 1 million g/mol, and

can have a charge density from about 0.1 meq/g to about 2.5 meq/g. A cationic guar polymer can have a weight average M.Wt. of less than 900 thousand g/mol, from about 150 thousand to about 800 thousand g/mol, from about 200 thousand g/mol to about 700 thousand g/mol, from about 300 thousand to about 700 thousand g/mol, from about 400 thousand to about 600 thousand g/mol, from about 150 thousand g/mol to about 800 thousand g/mol, from about 200 thousand g/mol to about 700 thousand g/mol, from about 300 thousand g/mol to about 700 thousand g/mol, and from about 400 thousand g/mol to about 600 thousand g/mol. A cationic guar polymer has a charge density from about 0.2 meq/g to about 2.2 meq/g, from about 0.3 meq/g to about 2.0 meq/g, from about 0.4 meq/g to about 1.8 meq/g; and from about 0.5 meq/g to about 1.5 meq/g.

(77) A cleansing composition can include from about 0.01% to less than about 0.7%, by weight of the cleansing composition of a cationic guar polymer, from about 0.04% to about 0.55%, by weight, from about 0.08% to about 0.5%, by weight, from about 0.16% to about 0.5%, by weight, from about 0.2% to about 0.5%, by weight, from about 0.3% to about 0.5%, by weight, and from about 0.4% to about 0.5%, by weight.

(78) The cationic guar polymer can be formed from quaternary ammonium compounds which conform to general Formula II:

(79) ##STR00001##

(80) wherein where R^{sup.3}, R^{sup.4} and R^{sup.5} are methyl or ethyl groups; and R^{sup.6} is either an epoxyalkyl group of the general Formula III:

(81) ##STR00002##

or R^{sup.6} is a halohydrin group of the general Formula IV:

(82) ##STR00003##

wherein R^{sup.7} is a C_{sub.1} to C_{sub.3} alkylene; X is chlorine or bromine, and Z is an anion such as Cl[—], Br[—], I[—] or HSO_{sub.4}[—].

Suitable cationic guar polymers can conform to the general formula V:

(83) ##STR00004##

wherein R^{sup.8} is guar gum; and wherein R^{sup.4}, R^{sup.5}, R^{sup.6} and R^{sup.7} are as defined above; and wherein Z is a halogen. Suitable cationic guar polymers can conform to Formula VI:

(84) ##STR00005##

(85) wherein R^{sup.8} is guar gum.

(86) Suitable cationic guar polymers can also include cationic guar gum derivatives, such as guar hydroxypropyltrimonium chloride. Suitable examples of guar hydroxypropyltrimonium chlorides can include the Jaguar series commercially available from Solvay® S.A., Hi-Care Series from Rhodia®, and N-Hance™ and AquaCat™ from Ashland™ Inc. For example, N-Hance™ BF-17 is a borate (boron) free guar polymers. N-Hance™ BF-17 has a charge density of about 1.7 meq/g and M.Wt. of about 800,000. BF-17 has a charge density of about 1.7 meq/g and M.Wt. of about 800,000. BF-17 has a charge density of about 1.7 meq/g and M.Wt. of about 800,000. BF-17 has a charge density of about 1.7 meq/g and M.Wt. of about 800,000.

(87) Cationic Non-Guar Galactomannan Polymer

(88) The cationic polymer can be a galactomannan polymer derivative. Suitable galactomannan polymer can have a mannose to galactose ratio of greater than 2:1 on a monomer to monomer basis and can be a cationic galactomannan polymer derivative or an amphoteric galactomannan polymer derivative having a net positive charge. As used herein, the term “cationic galactomannan” refers to a galactomannan polymer to which a cationic group is added. The term “amphoteric galactomannan” refers to a galactomannan polymer to which a cationic group and an anionic group are added such that the polymer has a net positive charge.

(89) Galactomannan polymers can be present in the endosperm of seeds of the Leguminosae family Galactomannan polymers are made up of a combination of mannose monomers and galactose monomers. The galactomannan molecule is a straight chain mannan branched at regular intervals with single membered galactose units on specific mannose units. The mannose units are linked to each other by means of β (1-4) glycosidic linkages. The galactose branching arises by way of an α (1-6) linkage. The ratio of mannose monomers to galactose monomers varies according to the species of the plant and can be affected by climate. Non Guar Galactomannan polymer derivatives can have a ratio of mannose

to galactose of greater than 2:1 on a monomer to monomer basis. Suitable ratios of mannose to galactose can also be greater than 3:1 or greater than 4:1. Analysis of mannose to galactose ratios is well known in the art and is typically based on the measurement of the galactose content.

(90) The gum for use in preparing the non-guar galactomannan polymer derivatives can be obtained from naturally occurring materials such as seeds or beans from plants. Examples of various non-guar galactomannan polymers include Tara gum (3 parts mannose/1 part galactose), Locust bean or Carob (4 parts mannose/1 part galactose), and Cassia gum (5 parts mannose/1 part galactose).

(91) A non-guar galactomannan polymer derivative can have a M. Wt. from about 1,000 g/mol to about 10,000,000 g/mol, and a M.Wt. from about 5,000 g/mol to about 3,000,000 g/mol.

(92) The cleansing compositions described herein can include galactomannan polymer derivatives which have a cationic charge density from about 0.5 meq/g to about 7 meq/g. The galactomannan polymer derivatives can have a cationic charge density from about 1 meq/g to about 5 meq/g. The degree of substitution of the cationic groups onto the galactomannan structure can be sufficient to provide the requisite cationic charge density.

(93) A galactomannan polymer derivative can be a cationic derivative of the non-guar galactomannan polymer, which is obtained by reaction between the hydroxyl groups of the polygalactomannan polymer and reactive quaternary ammonium compounds. Suitable quaternary ammonium compounds for use in forming the cationic galactomannan polymer derivatives include those conforming to the general Formulas II to VI, as defined above.

(94) Cationic non-guar galactomannan polymer derivatives formed from the reagents described above can be represented by the general Formula VII:

(95) ##STR00006##

(96) wherein R is the gum. The cationic galactomannan derivative can be a gum hydroxypropyltrimethylammonium chloride, which can be more specifically represented by the general Formula VIII:

(97) ##STR00007##

(98) The galactomannan polymer derivative can be an amphoteric galactomannan polymer derivative having a net positive charge, obtained when the cationic galactomannan polymer derivative further comprises an anionic group.

(99) A cationic non-guar galactomannan can have a ratio of mannose to galactose which is greater than 4:1, a M.Wt. of about 100,000 g/mol to about 500,000 g/mol, a M.Wt. of about 50,000 g/mol to about 400,000 g/mol, and a cationic charge density from about 1 meq/g to about 5 meq/g, and from about 2 meq/g to about 4 meq/g.

(100) Cleansing compositions can include at least about 0.05% of a galactomannan polymer derivative by weight of the composition. The cleansing compositions can include from about 0.05% to about 2%, by weight of the composition, of a galactomannan polymer derivative.

(101) Cationic Starch Polymers

(102) Suitable cationic polymers can also be water-soluble cationically modified starch polymers. As used herein, the term "cationically modified starch" refers to a starch to which a cationic group is added prior to degradation of the starch to a smaller molecular weight, or wherein a cationic group is added after modification of the starch to achieve a desired molecular weight. The definition of the term "cationically modified starch" also includes amphoterically modified starch. The term "amphoterically modified starch" refers to a starch hydrolysate to which a cationic group and an anionic group are added.

(103) The cleansing compositions described herein can include cationically modified starch polymers at a range of about 0.01% to about 10%, and/or from about 0.05% to about 5%, by weight of the composition.

(104) The cationically modified starch polymers disclosed herein have a percent of bound nitrogen of from about 0.5% to about 4%.

(105) The cationically modified starch polymers can have a molecular weight from about 850,000 g/mol to about 15,000,000 g/mol and from about 900,000 g/mol to about 5,000,000 g/mol.

(106) Cationically modified starch polymers can have a charge density of from about 0.2 meq/g to about 5 meq/g, and from about 0.2 meq/g to about 2 meq/g. The chemical modification to obtain such a charge density can include the addition of amino and/or ammonium groups into the starch molecules. Non-

limiting examples of such ammonium groups can include substituents such as hydroxypropyl trimmonium chloride, trimethylhydroxypropyl ammonium chloride, dimethylstearylhydroxypropyl ammonium chloride, and dimethyldodecylhydroxypropyl ammonium chloride. Further details are described in Solarek, D. B., Cationic Starches in Modified Starches: Properties and Uses, Wurzburg, O. B., Ed., CRC Press, Inc., Boca Raton, Fla. 1986, pp 113-125 which is hereby incorporated by reference. The cationic groups can be added to the starch prior to degradation to a smaller molecular weight or the cationic groups may be added after such modification.

(107) A cationically modified starch polymer can have a degree of substitution of a cationic group from about 0.2 to about 2.5. As used herein, the “degree of substitution” of the cationically modified starch polymers is an average measure of the number of hydroxyl groups on each anhydroglucose unit which is derivatized by substituent groups. Since each anhydroglucose unit has three potential hydroxyl groups available for substitution, the maximum possible degree of substitution is 3. The degree of substitution is expressed as the number of moles of substituent groups per mole of anhydroglucose unit, on a molar average basis. The degree of substitution can be determined using proton nuclear magnetic resonance spectroscopy (“¹H NMR”) methods well known in the art. Suitable ¹H NMR techniques include those described in “Observation on NMR Spectra of Starches in Dimethyl Sulfoxide, Iodine-Complexing, and Solvating in Water-Dimethyl Sulfoxide”, Qin-Ji Peng and Arthur S. Perlin, Carbohydrate Research, 160 (1987), 57-72; and “An Approach to the Structural Analysis of Oligosaccharides by NMR Spectroscopy”, J. Howard Bradbury and J. Grant Collins, Carbohydrate Research, 71, (1979), 15-25.

(108) The source of starch before chemical modification can be selected from a variety of sources such as tubers, legumes, cereal, and grains. For example, starch sources can include corn starch, wheat starch, rice starch, waxy corn starch, oat starch, cassaya starch, waxy barley, waxy rice starch, glutenous rice starch, sweet rice starch, amioca, potato starch, tapioca starch, oat starch, sago starch, sweet rice, or mixtures thereof. Suitable cationically modified starch polymers can be selected from degraded cationic maize starch, cationic tapioca, cationic potato starch, and mixtures thereof. Cationically modified starch polymers are cationic corn starch and cationic tapioca.

(109) The starch, prior to degradation or after modification to a smaller molecular weight, can include one or more additional modifications. For example, these modifications may include cross-linking, stabilization reactions, phosphorylations, and hydrolyzations. Stabilization reactions can include alkylation and esterification.

(110) Cationically modified starch polymers can be included in a cleansing composition in the form of hydrolyzed starch (e.g., acid, enzyme, or alkaline degradation), oxidized starch (e.g., peroxide, peracid, hypochlorite, alkaline, or any other oxidizing agent), physically/mechanically degraded starch (e.g., via the thermo-mechanical energy input of the processing equipment), or combinations thereof.

(111) The starch can be readily soluble in water and can form a substantially translucent solution in water. The transparency of the composition is measured by Ultra-Violet/Visible (“UV/VIS”) spectrophotometry, which determines the absorption or transmission of UV/VIS light by a sample, using a Gretag Macbeth Colorimeter Color. A light wavelength of 600 nm has been shown to be adequate for characterizing the degree of clarity of cleansing compositions.

(112) Cationic Copolymer of an Acrylamide Monomer and a Cationic Monomer

(113) A cleansing composition can include a cationic copolymer of an acrylamide monomer and a cationic monomer, wherein the copolymer has a charge density of from about 1.0 meq/g to about 3.0 meq/g. The cationic copolymer can be a synthetic cationic copolymer of acrylamide monomers and cationic monomers.

(114) Suitable cationic polymers can include:

(115) (i) an acrylamide monomer of the following Formula IX:

(116) ##STR00008##

where R⁹ is H or C₁₋₄ alkyl; and R¹⁰ and R¹¹ are independently selected from the group consisting of H, C₁₋₄ alkyl, CH₂OCH₃, CH₂OCH₂CH₃, and phenyl, or together are C₃₋₆cycloalkyl; and

(ii) a cationic monomer conforming to Formula X:

(117) ##STR00009##

where $k=1$, $v=v'$, and v'' is independently an integer of from 1 to 6, w is zero or an integer of from 1 to 10, and $X_{sup.-}$ is an anion.

(118) A cationic monomer can conform to Formula X where $k=1$, $v=3$ and $w=0$, $z=1$ and $X_{sup.-}$ is $Cl_{sup.-}$ to form the following structure (Formula XI):

(119) ##STR00010##

(120) As can be appreciated, the above structure can be referred to as diquat.

(121) A cationic monomer can conform to Formula X wherein v and v'' are each 3, $v'=1$, $w=1$, $y=1$ and $X_{sup.-}$ is $Cl_{sup.-}$, to form the following structure of Formula XII:

(122) ##STR00011##

(123) The structure of Formula XII can be referred to as triquat.

(124) The acrylamide monomer can be either acrylamide or methacrylamide.

(125) The cationic copolymer can be AM:TRIQUAT which is a copolymer of acrylamide and 1,3-Propanediaminium,N-[2-[[[dimethyl[3-[(2-methyl-1-oxo-2-propenyl)amino]propyl]ammonio[acetyl]amino]ethyl]2-hydroxy-N,N,N',N'-pentamethyl-, trichloride. AM:TRIQUAT is also known as polyquaternium 76 (PQ76). AM:TRIQUAT can have a charge density of 1.6 meq/g and a M.Wt. of 1.1 million g/mol.

(126) The cationic copolymer can include an acrylamide monomer and a cationic monomer, wherein the cationic monomer is selected from the group consisting of: dimethylaminoethyl (meth)acrylate, dimethylaminopropyl (meth)acrylate, ditertiobutylaminoethyl (meth)acrylate, dimethylaminomethyl (meth)acrylamide, dimethylaminopropyl (meth)acrylamide; ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine; trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride, and mixtures thereof.

(127) The cationic copolymer can include a cationic monomer selected from the group consisting of: trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, and mixtures thereof.

(128) The cationic copolymer can be formed from (1) copolymers of (meth)acrylamide and cationic monomers based on (meth)acrylamide, and/or hydrolysis-stable cationic monomers, (2) terpolymers of (meth)acrylamide, monomers based on cationic (meth)acrylic acid esters, and monomers based on (meth)acrylamide, and/or hydrolysis-stable cationic monomers. Monomers based on cationic (meth)acrylic acid esters can be cationized esters of the (meth)acrylic acid containing a quaternized N atom. Cationized esters of the (meth)acrylic acid containing a quaternized N atom can be quaternized dialkylaminoalkyl (meth)acrylates with C.sub.1 to C.sub.3 in the alkyl and alkylene groups. The cationized esters of the (meth)acrylic acid containing a quaternized N atom can be selected from the group consisting of ammonium salts of dimethylaminomethyl (meth)acrylate, dimethylaminoethyl (meth)acrylate, dimethylaminopropyl (meth)acrylate, diethylaminomethyl (meth)acrylate, diethylaminoethyl (meth)acrylate; and diethylaminopropyl (meth)acrylate quaternized with methyl chloride. The cationized esters of the (meth)acrylic acid containing a quaternized N atom can be dimethylaminoethyl acrylate, which is quaternized with an alkyl halide, or with methyl chloride or benzyl chloride or dimethyl sulfate (ADAME-Quat). The cationic monomer when based on (meth)acrylamides are quaternized dialkylaminoalkyl(meth)acrylamides with C.sub.1 to C.sub.3 in the alkyl and alkylene groups, or dimethylaminopropylacrylamide, which is quaternized with an alkyl halide, or methyl chloride or benzyl chloride or dimethyl sulfate.

(129) The cationic monomer based on a (meth)acrylamide can be a quaternized dialkylaminoalkyl(meth)acrylamide with C.sub.1 to C.sub.3 in the alkyl and alkylene groups. The cationic monomer based on a (meth)acrylamide can be dimethylaminopropylacrylamide, which is quaternized with an alkyl halide, especially methyl chloride or benzyl chloride or dimethyl sulfate. The cationic monomer can be a hydrolysis-stable cationic monomer. Hydrolysis-stable cationic monomers

can be, in addition to a dialkylaminoalkyl(meth)acrylamide, any monomer that can be regarded as stable to the OECD hydrolysis test. The cationic monomer can be hydrolysis-stable, and the hydrolysis-stable cationic monomer can be selected from the group consisting of: diallyldimethylammonium chloride and water-soluble, cationic styrene derivatives.

(130) The cationic copolymer can be a terpolymer of acrylamide, 2-dimethylammoniummethyl(meth)acrylate quaternized with methyl chloride (ADAME-Q) and 3-dimethylammoniumpropyl(meth)acrylamide quaternized with methyl chloride (DIMAPA-Q). The cationic copolymer can be formed from acrylamide and acrylamidopropyltrimethylammonium chloride, wherein the acrylamidopropyltrimethylammonium chloride has a charge density of from about 1.0 meq/g to about 3.0 meq/g.

(131) The cationic copolymer can have a charge density of from about 1.1 meq/g to about 2.5 meq/g, from about 1.1 meq/g to about 2.3 meq/g, from about 1.2 meq/g to about 2.2 meq/g, from about 1.2 meq/g to about 2.1 meq/g, from about 1.3 meq/g to about 2.0 meq/g, and from about 1.3 meq/g to about 1.9 meq/g.

(132) The cationic copolymer can have a M.Wt. from about 100 thousand g/mol to about 2 million g/mol, from about 300 thousand g/mol to about 1.8 million g/mol, from about 500 thousand g/mol to about 1.6 million g/mol, from about 700 thousand g/mol to about 1.4 million g/mol, and from about 900 thousand g/mol to about 1.2 million g/mol.

(133) The cationic copolymer can be a trimethylammoniopropylmethacrylamide chloride-N-Acrylamide copolymer, which is also known as AM:MAPTAC. AM:MAPTAC can have a charge density of about 1.3 meq/g and a M.Wt. of about 1.1 million g/mol. The cationic copolymer can be AM:ATPAC. AM:ATPAC can have a charge density of about 1.8 meq/g and a M.Wt. of about 1.1 million g/mol.

(134) Synthetic Polymers

(135) A cationic polymer can be a synthetic polymer that is formed from: i) one or more cationic monomer units, and optionally ii) one or more monomer units bearing a negative charge, and/or iii) a nonionic monomer,

(136) wherein the subsequent charge of the copolymer is positive. The ratio of the three types of monomers is given by “m”, “p” and “q” where “m” is the number of cationic monomers, “p” is the number of monomers bearing a negative charge and “q” is the number of nonionic monomers

(137) The cationic polymers can be water soluble or dispersible, non-crosslinked, and synthetic cationic polymers which have the structure of Formula XIII:

(138) ##STR00012##

where A, may be one or more of the following cationic moieties:

(139) ##STR00013##

where @=amido, alkylamido, ester, ether, alkyl or alkylaryl;

where Y=C1-C22 alkyl, alkoxy, alkylidene, alkyl or aryloxy;

where ψ =C1-C22 alkyl, alkyloxy, alkyl aryl or alkyl arylox;

where Z=C1-C22 alkyl, alkyloxy, aryl or aryloxy;

where R1=H, C1-C4 linear or branched alkyl;

where s=0 or 1, n=0 or ≥ 1 ;

where T and R7=C1-C22 alkyl; and

where X=halogen, hydroxide, alkoxide, sulfate or alkylsulfate.

(140) Where the monomer bearing a negative charge is defined by R2'=H, C.sub.1-C.sub.4 linear or branched alkyl and R3 is:

(141) ##STR00014##

where D=O, N, or S;

where Q=NH.sub.2 or O;

where u=1-6;

where t=0-1; and

where J=oxygenated functional group containing the following elements P, S, C.

(142) Where the nonionic monomer is defined by R2''=H, C.sub.1-C.sub.4 linear or branched alkyl, R6=linear or branched alkyl, alkyl aryl, aryl oxy, alkyloxy, alkylaryl oxy and β is defined as

(143) ##STR00015##

and

where G' and G'' are, independently of one another, O, S or N—H and L=0 or 1.

(144) Suitable monomers can include aminoalkyl (meth)acrylates, (meth)aminoalkyl (meth)acrylamides; monomers comprising at least one secondary, tertiary or quaternary amine function, or a heterocyclic group containing a nitrogen atom, vinylamine or ethylenimine; diallyldialkyl ammonium salts; their mixtures, their salts, and macromonomers deriving from therefrom.

(145) Further examples of suitable cationic monomers can include dimethylaminoethyl (meth)acrylate, dimethylaminopropyl (meth)acrylate, ditertiobutylaminoethyl (meth)acrylate, dimethylaminomethyl (meth)acrylamide, dimethylaminopropyl (meth)acrylamide, ethylenimine, vinylamine, 2-vinylpyridine, 4-vinylpyridine, trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride, diallyldimethyl ammonium chloride.

(146) Suitable cationic monomers can include quaternary monomers of formula —NR₃⁺, wherein each R can be identical or different, and can be a hydrogen atom, an alkyl group comprising 1 to 10 carbon atoms, or a benzyl group, optionally carrying a hydroxyl group, and including an anion (counter-ion). Examples of suitable anions include halides such as chlorides, bromides, sulphates, hydrosulphates, alkylsulphates (for example comprising 1 to 6 carbon atoms), phosphates, citrates, formates, and acetates.

(147) Suitable cationic monomers can also include trimethylammonium ethyl (meth)acrylate chloride, trimethylammonium ethyl (meth)acrylate methyl sulphate, dimethylammonium ethyl (meth)acrylate benzyl chloride, 4-benzoylbenzyl dimethylammonium ethyl acrylate chloride, trimethyl ammonium ethyl (meth)acrylamido chloride, trimethyl ammonium propyl (meth)acrylamido chloride, vinylbenzyl trimethyl ammonium chloride. Additional suitable cationic monomers can include trimethyl ammonium propyl (meth)acrylamido chloride.

(148) Examples of monomers bearing a negative charge include alpha ethylenically unsaturated monomers including a phosphate or phosphonate group, alpha ethylenically unsaturated monocarboxylic acids, monoalkylesters of alpha ethylenically unsaturated dicarboxylic acids, monoalkylamides of alpha ethylenically unsaturated dicarboxylic acids, alpha ethylenically unsaturated compounds comprising a sulphonic acid group, and salts of alpha ethylenically unsaturated compounds comprising a sulphonic acid group.

(149) Suitable monomers with a negative charge can include acrylic acid, methacrylic acid, vinyl sulphonic acid, salts of vinyl sulfonic acid, vinylbenzene sulphonic acid, salts of vinylbenzene sulphonic acid, alpha-acrylamidomethylpropanesulphonic acid, salts of alpha-acrylamidomethylpropanesulphonic acid, 2-sulphoethyl methacrylate, salts of 2-sulphoethyl methacrylate, acrylamido-2-methylpropanesulphonic acid (AMPS), salts of acrylamido-2-methylpropanesulphonic acid, and styrenesulphonate (SS).

(150) Examples of nonionic monomers can include vinyl acetate, amides of alpha ethylenically unsaturated carboxylic acids, esters of an alpha ethylenically unsaturated monocarboxylic acids with an hydrogenated or fluorinated alcohol, polyethylene oxide (meth)acrylate (i.e. polyethoxylated (meth)acrylic acid), monoalkylesters of alpha ethylenically unsaturated dicarboxylic acids, monoalkylamides of alpha ethylenically unsaturated dicarboxylic acids, vinyl nitriles, vinylamine amides, vinyl alcohol, vinyl pyrrolidone, and vinyl aromatic compounds. Suitable nonionic monomers can also include styrene, acrylamide, methacrylamide, acrylonitrile, methylacrylate, ethylacrylate, n-propylacrylate, n-butylacrylate, methylmethacrylate, ethylmethacrylate, n-propylmethacrylate, n-butylmethacrylate, 2-ethyl-hexyl acrylate, 2-ethyl-hexyl methacrylate, 2-hydroxyethylacrylate and 2-hydroxyethylmethacrylate.

(151) The anionic counterion (X[−]) in association with the synthetic cationic polymers can be any known counterion so long as the polymers remain soluble or dispersible in water, in the cleansing composition, or in a coacervate phase of the cleansing composition, and so long as the counterions are physically and chemically compatible with the essential components of the cleansing composition or do not otherwise unduly impair product performance, stability or aesthetics. Non limiting examples of

suitable counterions can include halides (e.g., chlorine, fluorine, bromine, iodine), sulfate, and methylsulfate.

(152) The cationic polymer described herein can also aid in repairing damaged hair, particularly chemically treated hair by providing a surrogate hydrophobic F-layer. The microscopically thin F-layer provides natural weatherproofing, while helping to seal in moisture and prevent further damage. Chemical treatments damage the hair cuticle and strip away its protective F-layer. As the F-layer is stripped away, the hair becomes increasingly hydrophilic. It has been found that when lyotropic liquid crystals are applied to chemically treated hair, the hair becomes more hydrophobic and more virgin-like, in both look and feel. Without being limited to any theory, it is believed that the lyotropic liquid crystal complex creates a hydrophobic layer or film, which coats the hair fibers and protects the hair, much like the natural F-layer protects the hair. The hydrophobic layer can return the hair to a generally virgin-like, healthier state. Lyotropic liquid crystals are formed by combining the synthetic cationic polymers described herein with the aforementioned anionic detergent surfactant component of the cleansing composition. The synthetic cationic polymer has a relatively high charge density. It should be noted that some synthetic polymers having a relatively high cationic charge density do not form lyotropic liquid crystals, primarily due to their abnormal linear charge densities. Such synthetic cationic polymers are described in PCT Patent App. No. WO 94/06403 which is incorporated by reference. The synthetic polymers described herein can be formulated in a stable cleansing composition that provides improved conditioning performance, with respect to damaged hair.

(153) Cationic synthetic polymers that can form lyotropic liquid crystals have a cationic charge density of from about 2 meq/gm to about 7 meq/gm, and/or from about 3 meq/gm to about 7 meq/gm, and/or from about 4 meq/gm to about 7 meq/gm. The cationic charge density is about 6.2 meq/gm. The polymers also have a M. Wt. of from about 1,000 to about 5,000,000, and/or from about 10,000 to about 2,000,000, and/or from about 100,000 to about 2,000,000.

(154) Cationic synthetic polymers that provide enhanced conditioning and deposition of benefit agents but do not necessarily form lyotropic liquid crystals can have a cationic charge density of from about 0.7 meq/gm to about 7 meq/gm, and/or from about 0.8 meq/gm to about 5 meq/gm, and/or from about 1.0 meq/gm to about 3 meq/gm. The polymers also have a M.Wt. of from about 1,000 g/mol to about 5,000,000 g/mol, from about 10,000 g/mol to about 2,000,000 g/mol, and from about 100,000 g/mol to about 2,000,000 g/mol.

(155) Cationic Cellulose Polymer

(156) Suitable cationic polymers can be cellulose polymers. Suitable cellulose polymers can include salts of hydroxyethyl cellulose reacted with trimethyl ammonium substituted epoxide, referred to in the industry (CTFA) as Polyquaternium 10 and available from Dwo/Amerchol Corp. (Edison, N.J., USA) in their Polymer LR, JR, and KG series of polymers. Other suitable types of cationic cellulose can include the polymeric quaternary ammonium salts of hydroxyethyl cellulose reacted with lauryl dimethyl ammonium-substituted epoxide referred to in the industry (CTFA) as Polyquaternium 24. These materials are available from Dow/Amerchol Corp. under the tradename Polymer LM-200. Other suitable types of cationic cellulose can include the polymeric quaternary ammonium salts of hydroxyethyl cellulose reacted with lauryl dimethyl ammonium-substituted epoxide and trimethyl ammonium substituted epoxide referred to in the industry (CTFA) as Polyquaternium 67. These materials are available from Dow/Amerchol Corp. under the tradename SoftCAT Polymer SL-5, SoftCAT Polymer SL-30, Polymer SL-60, Polymer SL-100, Polymer SK-L, Polymer SK-M, Polymer SK-MH, and Polymer SK-H. Cationic cellulose polymers can have a cationic charge density of from about 0.2 meq/g to about 2.2 meq/g, from about 0.3 meq/g to about 2.0 meq/g, from about 0.4 meq/g to about 1.8 meq/g; from about 0.5 meq/g to about 1.7 meq/g and from about 0.6 meq/g to about 1.3.

(157) Additional cationic polymers are also described in the CTFA Cosmetic Ingredient Dictionary, 3rd edition, edited by Estrin, Crosley, and Haynes, (The Cosmetic, Toiletry, and Fragrance Association, Inc., Washington, D.C. (1982)), which is incorporated herein by reference. Techniques for analysis of formation of complex coacervates are known in the art. For example, microscopic analyses of the compositions, at any chosen stage of dilution, can be utilized to identify whether a coacervate phase has formed. Such coacervate phase can be identifiable as an additional emulsified phase in the composition. The use of dyes can aid in distinguishing the coacervate phase from other insoluble phases dispersed in

the composition. Additional details about the use of cationic polymers and coacervates are disclosed in U.S. Pat. No. 9,272,164 which is incorporated by reference.

(158) Liquid Carrier

(159) As can be appreciated, cleansing compositions can desirably be in the form of pourable liquid under ambient conditions. Inclusion of an appropriate quantity of a liquid carrier can facilitate the formation of a cleansing composition having an appropriate viscosity and rheology. A cleansing composition can include, by weight of the composition, about 20% to about 95%, by weight, of a liquid carrier, and about 60% to about 85%, by weight, of a liquid carrier. The liquid carrier can be an aqueous carrier such as water.

(160) Optional Components

(161) As can be appreciated, cleansing compositions described herein can include a variety of optional components to tailor the properties and characteristics of the composition. As can be appreciated, suitable optional components are well known and can generally include any components which are physically and chemically compatible with the essential components of the cleansing compositions described herein. Optional components should not otherwise unduly impair product stability, aesthetics, or performance. Individual concentrations of optional components can generally range from about 0.001% to about 10%, by weight of a cleansing composition. Optional components can be further limited to components which will not impair the clarity of a translucent cleansing composition.

(162) Suitable optional components which can be included in a cleansing composition can include co-surfactants, deposition aids, conditioning agents (including hydrocarbon oils, fatty esters, silicones), anti-dandruff agents, suspending agents, viscosity modifiers, dyes, nonvolatile solvents or diluents (water soluble and insoluble), pearlescent aids, foam boosters, pediculocides, pH adjusting agents, perfumes, preservatives, chelants, proteins, skin active agents, sunscreens, UV absorbers, and vitamins. The CTFA Cosmetic Ingredient Handbook, Tenth Edition (published by the Cosmetic, Toiletry, and Fragrance Association, Inc., Washington, D.C.) (2004) (hereinafter "CTFA"), describes a wide variety of non-limiting materials that can be added to the composition herein.

(163) Conditioning Agents

(164) A cleansing composition can include a silicone conditioning agent. Suitable silicone conditioning agents can include volatile silicone, non-volatile silicone, or combinations thereof. If including a silicone conditioning agent, the agent can be included from about 0.01% to about 10%, by weight of the composition, from about 0.1% to about 8%, from about 0.1% to about 5%, and/or from about 0.2% to about 3%. Examples of suitable silicone conditioning agents, and optional suspending agents for the silicone, are described in U.S. Reissue Pat. No. 34,584, U.S. Pat. Nos. 5,104,646, and 5,106,609, each of which is incorporated by reference herein. Suitable silicone conditioning agents can have a viscosity, as measured at 25° C., from about 20 centistokes ("csk") to about 2,000,000 csk, from about 1,000 csk to about 1,800,000 csk, from about 50,000 csk to about 1,500,000 csk, and from about 100,000 csk to about 1,500,000 csk.

(165) The dispersed silicone conditioning agent particles can have a volume average particle diameter ranging from about 0.01 micrometer to about 50 micrometer. For small particle application to hair, the volume average particle diameters can range from about 0.01 micrometer to about 4 micrometer, from about 0.01 micrometer to about 2 micrometer, from about 0.01 micrometer to about 0.5 micrometer. For larger particle application to hair, the volume average particle diameters typically range from about 5 micrometer to about 125 micrometer, from about 10 micrometer to about 90 micrometer, from about 15 micrometer to about 70 micrometer, and/or from about 20 micrometer to about 50 micrometer.

(166) Additional material on silicones including sections discussing silicone fluids, gums, and resins, as well as manufacture of silicones, are found in *Encyclopedia of Polymer Science and Engineering*, vol. 15, 2d ed., pp 204-308, John Wiley & Sons, Inc. (1989), which is incorporated herein by reference.

(167) Silicone emulsions suitable for the cleansing compositions described herein can include emulsions of insoluble polysiloxanes prepared in accordance with the descriptions provided in U.S. Pat. No. 4,476,282 and U.S. Patent Application Publication No. 2007/0276087 each of which is incorporated herein by reference. Suitable insoluble polysiloxanes include polysiloxanes such as alpha, omega hydroxy-terminated polysiloxanes or alpha, omega alkoxy-terminated polysiloxanes having a molecular weight within the range from about 50,000 to about 500,000 g/mol. The insoluble polysiloxane can have

an average molecular weight within the range from about 50,000 to about 500,000 g/mol. For example, the insoluble polysiloxane may have an average molecular weight within the range from about 60,000 to about 400,000; from about 75,000 to about 300,000; from about 100,000 to about 200,000; or the average molecular weight may be about 150,000 g/mol. The insoluble polysiloxane can have an average particle size within the range from about 30 nm to about 10 micron. The average particle size may be within the range from about 40 nm to about 5 micron, from about 50 nm to about 1 micron, from about 75 nm to about 500 nm, or about 100 nm, for example.

(168) Other classes of silicones suitable for the cleansing compositions described herein can include i) silicone fluids, including silicone oils, which are flowable materials having viscosity less than about 1,000,000 csk as measured at 25° C.; ii) aminosilicones, which contain at least one primary, secondary or tertiary amine; iii) cationic silicones, which contain at least one quaternary ammonium functional group; iv) silicone gums; which include materials having viscosity greater or equal to 1,000,000 csk as measured at 25° C.; v) silicone resins, which include highly cross-linked polymeric siloxane systems; vi) high refractive index silicones, having refractive index of at least 1.46, and vii) mixtures thereof.

(169) Alternatively, the cleansing composition can be substantially free or free of silicones.

(170) Organic Conditioning Materials

(171) The conditioning agent of the cleansing compositions described herein can also include at least one organic conditioning material such as oil or wax, either alone or in combination with other conditioning agents, such as the silicones described above. The organic material can be non-polymeric, oligomeric or polymeric. The organic material can be in the form of an oil or wax and can be added in the cleansing formulation neat or in a pre-emulsified form. Suitable examples of organic conditioning materials can include: i) hydrocarbon oils; ii) polyolefins, iii) fatty esters, iv) fluorinated conditioning compounds, v) fatty alcohols, vi) alkyl glucosides and alkyl glucoside derivatives; vii) quaternary ammonium compounds; viii) polyethylene glycols and polypropylene glycols having a molecular weight of up to about 2,000,000 including those with CTFA names PEG-200, PEG-400, PEG-600, PEG-1000, PEG-2M, PEG-7M, PEG-14M, PEG-45M and mixtures thereof.

(172) Emulsifiers

(173) A variety of anionic and nonionic emulsifiers can be used in the cleansing composition of the present invention. The anionic and nonionic emulsifiers can be either monomeric or polymeric in nature. Monomeric examples include, by way of illustrating and not limitation, alkyl ethoxylates, alkyl sulfates, soaps, and fatty esters and their derivatives. Polymeric examples include, by way of illustrating and not limitation, polyacrylates, polyethylene glycols, and block copolymers and their derivatives. Naturally occurring emulsifiers such as lanolins, lecithin and lignin and their derivatives are also non-limiting examples of useful emulsifiers.

(174) Chelating Agents

(175) The cleansing composition can also comprise a chelant. Suitable chelants include those listed in A E Martell & R M Smith, Critical Stability Constants, Vol. 1, Plenum Press, New York & London (1974) and A E Martell & R D Hancock, Metal Complexes in Aqueous Solution, Plenum Press, New York & London (1996) both incorporated herein by reference. When related to chelants, the term “salts and derivatives thereof” means the salts and derivatives comprising the same functional structure (e.g., same chemical backbone) as the chelant they are referring to and that have similar or better chelating properties. This term includes alkali metal, alkaline earth, ammonium, substituted ammonium (i.e. monoethanolammonium, diethanolammonium, triethanolammonium) salts, esters of chelants having an acidic moiety and mixtures thereof, in particular all sodium, potassium or ammonium salts. The term “derivatives” also includes “chelating surfactant” compounds, such as those exemplified in U.S. Pat. No. 5,284,972, and large molecules comprising one or more chelating groups having the same functional structure as the parent chelants, such as polymeric EDDS (ethylenediaminedisuccinic acid) disclosed in U.S. Pat. Nos. 5,747,440, and 5,747,440 are each incorporated by reference herein. Suitable chelants can further include histidine.

(176) Levels of an EDDS chelant or histidine chelant in the cleansing compositions can be low. For example, an EDDS chelant or histidine chelant can be included at about 0.01%, by weight. Above about 10% by weight, formulation and/or human safety concerns can arise. The level of an EDDS chelant or histidine chelant can be at least about 0.01%, by weight, at least about 0.05%, by weight, at least about

0.1%, by weight, at least about 0.25%, by weight, at least about 0.5%, by weight, at least about 1%, by weight, or at least about 2%, by weight, by weight of the cleansing composition.

(177) Gel Network

(178) A cleansing composition can also include a fatty alcohol gel network. Gel networks are formed by combining fatty alcohols and surfactants in the ratio of from about 1:1 to about 40:1, from about 2:1 to about 20:1, and/or from about 3:1 to about 10:1. The formation of a gel network involves heating a dispersion of the fatty alcohol in water with the surfactant to a temperature above the melting point of the fatty alcohol. During the mixing process, the fatty alcohol melts, allowing the surfactant to partition into the fatty alcohol droplets. The surfactant brings water along with it into the fatty alcohol. This changes the isotropic fatty alcohol drops into liquid crystalline phase drops. When the mixture is cooled below the chain melt temperature, the liquid crystal phase is converted into a solid crystalline gel network. Gel networks can provide a number of benefits to cleansing compositions. For example, a gel network can provide a stabilizing benefit to cosmetic creams and hair conditioners. In addition, gel networks can provide conditioned feel benefits to hair conditioners and shampoos.

(179) A fatty alcohol can be included in the gel network at a level by weight of from about 0.05%, by weight, to about 14%, by weight. For example, the fatty alcohol can be included in an amount ranging from about 1%, by weight, to about 10%, by weight, and/or from about 6%, by weight, to about 8%, by weight.

(180) Suitable fatty alcohols include those having from about 10 to about 40 carbon atoms, from about 12 to about 22 carbon atoms, from about 16 to about 22 carbon atoms, and/or about 16 to about 18 carbon atoms. These fatty alcohols can be straight or branched chain alcohols and can be saturated or unsaturated. Nonlimiting examples of fatty alcohols include cetyl alcohol, stearyl alcohol, behenyl alcohol, and mixtures thereof. Mixtures of cetyl and stearyl alcohol in a ratio of from about 20:80 to about 80:20 are suitable.

(181) A gel network can be prepared by charging a vessel with water. The water can then be heated to about 74° C. Cetyl alcohol, stearyl alcohol, and surfactant can then be added to the heated water. After incorporation, the resulting mixture can pass through a heat exchanger where the mixture is cooled to about 35° C. Upon cooling, the fatty alcohols and surfactant crystallized can form crystalline gel network. Table 1 provides the components and their respective amounts for an example gel network composition.

(182) To prepare the gel network pre-mix of Table 1, water is heated to about 74° C. and the fatty alcohol and gel network surfactant are added to it in the quantities depicted in Table 1. After incorporation, this mixture is passed through a mill and heat exchanger where it is cooled to about 32° C. As a result of this cooling step, the fatty alcohol, the gel network surfactant, and the water form a crystalline gel network.

(183) TABLE-US-00001 TABLE 1 Premix % Gel Network Surfactant.sup.1 11.00 Stearyl Alcohol 8% Cetyl Alcohol 4% Water QS .sup.1 For anionic gel networks, suitable gel network surfactants above include surfactants with a net negative charge including sulfonates, carboxylates and phosphates among others and mixtures thereof.

(184) For cationic gel networks, suitable gel network surfactants above include surfactants with a net positive charge including quaternary ammonium surfactants and mixtures thereof.

(185) For Amphoteric or Zwitterionic gel networks, suitable gel network surfactants above include surfactants with both a positive and negative charge at product usage pH including betaines, amine oxides, sultaines, amino acids among others and mixtures thereof.

(186) Benefit Agents

(187) A cleansing composition can further include one or more benefit agents. Exemplary benefit agents include, but are not limited to, particles, colorants, perfume microcapsules, gel networks, and other insoluble skin or hair conditioning agents such as skin silicones, natural oils such as sunflower oil or castor oil. The benefit agent can be selected from the group consisting of particles; colorants; perfume microcapsules; gel networks; other insoluble skin or hair conditioning agents such as skin silicones, natural oils such as sunflower oil or castor oil; and mixtures thereof.

(188) Suspending Agent

(189) A cleansing composition can include a suspending agent at concentrations effective for suspending water-insoluble material in dispersed form in the compositions or for modifying the viscosity of the

composition. Such concentrations range from about 0.1% to about 10%, and from about 0.3% to about 5.0%, by weight of the compositions. As can be appreciated however, suspending agents may not be necessary when certain glyceride ester crystals are included as certain glyceride ester crystals can act as suitable suspending or structuring agents.

(190) Suitable suspending agents can include anionic polymers and nonionic polymers. Useful herein are vinyl polymers such as cross linked acrylic acid polymers with the CTFA name Carbomer, cellulose derivatives and modified cellulose polymers such as methyl cellulose, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl methyl cellulose, nitro cellulose, sodium cellulose sulfate, sodium carboxymethyl cellulose, crystalline cellulose, cellulose powder, polyvinylpyrrolidone, polyvinyl alcohol, guar gum, hydroxypropyl guar gum, xanthan gum, arabia gum, tragacanth, galactan, carob gum, guar gum, karaya gum, carrageenin, pectin, agar, quince seed (*Cydonia oblonga* Mill), starch (rice, corn, potato, wheat), algae colloids (algae extract), microbiological polymers such as dextran, succinoglucan, pulleran, starch-based polymers such as carboxymethyl starch, methylhydroxypropyl starch, alginic acid-based polymers such as sodium alginate, alginic acid propylene glycol esters, acrylate polymers such as sodium polyacrylate, polyethylacrylate, polyacrylamide, polyethyleneimine, and inorganic water soluble material such as bentonite, aluminum magnesium silicate, laponite, hectonite, and anhydrous silicic acid.

(191) Other suitable suspending agents can include crystalline suspending agents which can be categorized as acyl derivatives, long chain amine oxides, and mixtures thereof. Examples of such suspending agents are described in U.S. Pat. No. 4,741,855, which is incorporated herein by reference. Suitable suspending agents include ethylene glycol esters of fatty acids having from 16 to 22 carbon atoms. The suspending agent can be an ethylene glycol stearates, both mono and distearate, but particularly the distearate containing less than about 7% of the mono stearate. Other suitable suspending agents include alkanol amides of fatty acids, having from about 16 to about 22 carbon atoms, alternatively from about 16 to about 18 carbon atoms, suitable examples of which include stearic monoethanolamide, stearic diethanolamide, stearic monoisopropanolamide and stearic monoethanolamide stearate. Other long chain acyl derivatives include long chain esters of long chain fatty acids (e.g., stearyl stearate, cetyl palmitate, etc.); long chain esters of long chain alkanol amides (e.g., stearamide diethanolamide distearate, stearamide monoethanolamide stearate); and glyceryl esters as previously described. Long chain acyl derivatives, ethylene glycol esters of long chain carboxylic acids, long chain amine oxides, and alkanol amides of long chain carboxylic acids can also be used as suspending agents.

(192) Other long chain acyl derivatives suitable for use as suspending agents include N,N-dihydrocarbyl amido benzoic acid and soluble salts thereof (e.g., Na, K), particularly N,N-di(hydrogenated) C.sub.16, C.sub.18 and tallow amido benzoic acid species of this family, which are commercially available from Stepan® Company (Northfield, Ill., USA).

(193) Examples of suitable long chain amine oxides for use as suspending agents include alkyl dimethyl amine oxides, e.g., stearyl dimethyl amine oxide.

(194) Other suitable suspending agents include primary amines having a fatty alkyl moiety having at least about 16 carbon atoms, examples of which include palmitamine or stearamine, and secondary amines having two fatty alkyl moieties each having at least about 12 carbon atoms, examples of which include dipalmitoylamine or di(hydrogenated tallow)amine. Still other suitable suspending agents include di(hydrogenated tallow)phthalic acid amide, and crosslinked maleic anhydride-methyl vinyl ether copolymer.

(195) Viscosity Modifiers

(196) The shampoo composition can be free of or substantially free of viscosity modifiers other than organic salt.

(197) In some examples, the composition can contain a viscosity modifier instead of or in addition to organic salt. Viscosity modifiers can be used to modify the rheology of a cleansing composition. Suitable viscosity modifiers can include Carbomers with tradenames Carbopol 934, Carbopol 940, Carbopol 950, Carbopol 980, and Carbopol 981, all available from B. F. Goodrich Company, acrylates/stearate-20 methacrylate copolymer with tradename ACRY SOL 22 available from Rohm and Hass, nonoxynyl hydroxyethylcellulose with tradename AMERCELL POLYMER HM-1500 available from Amerchol,

methylethylcellulose with tradename BENECEL, hydroxyethyl cellulose with tradename NATROSOL, hydroxypropyl cellulose with tradename KLUCCEL, cetyl hydroxyethyl cellulose with tradename POLYSURF 67, all supplied by Hercules, ethylene oxide and/or propylene oxide based polymers with tradenames CARBOWAX PEGs, POLYOX WASRs, and UCON FLUIDS, all supplied by Amerchol. Other suitable rheology modifiers can include cross-linked acrylates, cross-linked maleic anhydride co-methylvinylethers, hydrophobically modified associative polymers, and mixtures thereof.

(198) Dispersed Particles

(199) Dispersed particles as known in the art can be included in a cleansing composition. If including such dispersed particles, the particles can be incorporated, by weight of the composition, at levels of about 0.025% or more, about 0.05% or more, about 0.1% or more, about 0.25% or more, and about 0.5% or more. However, the cleansing compositions can also contain, by weight of the composition, about 20% or fewer dispersed particles, about 10% or fewer dispersed particles, about 5% or fewer dispersed particles, about 3% or fewer dispersed particles, and about 2% or fewer dispersed particles.

(200) As can be appreciated, a cleansing composition can include still further optional components. For example, amino acids can be included. Suitable amino acids can include water soluble vitamins such as vitamins B1, B2, B6, B12, C, pantothenic acid, pantothenyl ethyl ether, panthenol, biotin, and their derivatives, water soluble amino acids such as asparagine, alanin, indole, glutamic acid and their salts, water insoluble vitamins such as vitamin A, D, E, and their derivatives, water insoluble amino acids such as tyrosine, tryptamine, and their salts.

(201) Anti-dandruff agents can be included. As can be appreciated, the formation of a coacervate can facilitate deposition of the anti-dandruff agent to the scalp.

(202) Suitable anti-dandruff agents can include pyridinethione salts, azoles, selenium sulfide, particulate sulfur, and mixtures thereof. Such anti-dandruff particulate should be physically and chemically compatible with the essential components of the composition, and should not otherwise unduly impair product stability, aesthetics or performance. A shampoo composition can include a cationic polymer to enhance deposition of an anti-dandruff active.

(203) An anti-dandruff agent can be a pyridinethione particulate such as a 1-hydroxy-2-pyridinethione salt. The concentration of pyridinethione anti-dandruff particulates can range from about 0.1% to about 4%, about 0.1% to about 3%, and from about 0.3% to about 2% by weight of the composition. Suitable pyridinethione salts include those formed from heavy metals such as zinc, tin, cadmium, magnesium, aluminum and zirconium, particularly suitable is the zinc salt of 1-hydroxy-2-pyridinethione (known as "zinc pyridinethione" or "ZPT"), 1-hydroxy-2-pyridinethione salts in platelet particle form, wherein the particles have an average size of up to about 20 μm , up to about 5 μm , up to about 2.5 μm . Salts formed from other cations, such as sodium, can also be suitable. Pyridinethione anti-dandruff agents are further described in U.S. Pat. Nos. 2,809,971; 3,236,733; 3,753,196; 3,761,418; 4,345,080; 4,323,683; 4,379,753; and 4,470,982, each of which are incorporated herein by reference. It is contemplated that when ZPT is used as the anti-dandruff particulate, that the growth or re-growth of hair may be stimulated or regulated, or both, or that hair loss may be reduced or inhibited, or that hair may appear thicker or fuller.

(204) In addition to the anti-dandruff active selected from polyvalent metal salts of pyrrithione, a shampoo composition can further include one or more anti-fungal or anti-microbial actives in addition to the metal pyrrithione salt actives. Suitable anti-microbial actives include coal tar, sulfur, whitfield's ointment, castellani's paint, aluminum chloride, gentian violet, octopirox (piroctone olamine), ciclopirox olamine, undecylenic acid and its metal salts, potassium permanganate, selenium sulphide, sodium thiosulfate, propylene glycol, oil of bitter orange, urea preparations, griseofulvin, 8-hydroxyquinoline ciloquinol, thiobendazole, thiocarbamates, haloprogin, polyenes, hydroxypyridone, morpholine, benzylamine, allylamines (such as terbinafine), tea tree oil, clove leaf oil, coriander, palmarosa, berberine, thyme red, cinnamon oil, cinnamic aldehyde, citronellic acid, hinokitol, ichthyol pale, Sensiva SC-50, Elestab HP-100, azelaic acid, lyticase, iodopropynyl butylcarbamate (IPBC), isothiazalinones such as octyl isothiazalinone and azoles, and combinations thereof. Suitable anti-microbials can include itraconazole, ketoconazole, selenium sulphide and coal tar.

(205) A suitable anti-microbial agent can be one material or a mixture selected from: azoles, such as climbazole, ketoconazole, itraconazole, econazole, and elubiol; hydroxy pyridones, such as piroctone

oleamine, ciclopirox, rilopirox, and MEA-Hydroxyoctyloxypyridinone; kerolytic agents, such as salicylic acid and other hydroxy acids; strobilurins such as azoxystrobin and metal chelators such as 1,10-phenanthroline. Examples of azole anti-microbials can include imidazoles such as benzimidazole, benzothiazole, bifonazole, butaconazole nitrate, climbazole, clotrimazole, croconazole, eberconazole, econazole, elubiol, fenticonazole, fluconazole, flutimazole, isoconazole, ketoconazole, lanoconazole, metronidazole, miconazole, neticonazole, omoconazole, oxiconazole nitrate, sertaconazole, sulconazole nitrate, tioconazole, thiazole, and triazoles such as terconazole and itraconazole, and combinations thereof. When present in a shampoo composition, the soluble anti-microbial active can be included in an amount from about 0.01% to about 5%, from about 0.5% to about 6%, from about 0.1% to about 3%, from about 0.1% to about 9%, from about 0.1% to about 1.5%, from about 0.1% to about 2%, and more from about 0.3% to about 2%, by weight of the composition.

(206) Selenium sulfide is a particulate anti-dandruff agent suitable for use as an anti-microbial compositions when included at concentrations of about 0.1% to about 4%, by weight of the composition, from about 0.3% to about 2.5% by weight, and from about 0.5% to about 1.5% by weight. Selenium sulfide is generally regarded as a compound having one mole of selenium and two moles of sulfur, although it may also be a cyclic structure that conforms to the general formula $\text{Se}_{x-1}\text{S}_y$, wherein $x+y=8$. Average particle diameters for the selenium sulfide are typically less than 15 μm , as measured by forward laser light scattering device (e.g. Malvern 3600 instrument), less than 10 μm . Selenium sulfide compounds are described, for example, in U.S. Pat. Nos. 2,694,668; 3,152,046; 4,089,945; and 4,885,107, each of which are incorporated herein by reference.

(207) Sulfur can also be used as a particulate anti-microbial/anti-dandruff agent. Effective concentrations of the particulate sulfur are typically from about 1% to about 4%, by weight of the composition, alternatively from about 2% to about 4%.

(208) Keratolytic agents such as salicylic acid can also be included in a shampoo composition described herein.

(209) A cleansing composition can optionally include pigment materials such as inorganic, nitroso, monoazo, disazo, carotenoid, triphenyl methane, triaryl methane, xanthene, quinoline, oxazine, azine, anthraquinone, indigoid, thionindigoid, quinacridone, phthalocyanine, botanical, natural colors, including water soluble components such as those having C. I. Names.

(210) One or more stabilizers and preservatives can be included. For example, one or more of trihydroxystearin, ethylene glycol distearate, citric acid, sodium citrate dihydrate, a preservative such as kathon, sodium chloride, sodium benzoate, sodium salicylate and ethylenediaminetetraacetic acid ("EDTA") can be included to improve the lifespan of a personal care composition. The stabilizer and/or preservative can be used at a level of from about 0.10% to about 2%. Particularly suitable is sodium benzoate at a level of from about 0.10% to about 0.45%. The personal care composition may also include citric acid at a level of from about 0.5% to about 2%. The sodium benzoate and the citric acid can be added to the personal care composition alone or in combination.

(211) Method of Making a Cleansing Composition

(212) A cleansing composition described herein can be formed similarly to known cleansing compositions. For example, the process of making a cleansing composition can include the step of mixing the surfactant, cationic polymer, and liquid carrier together to form a cleansing composition. The sclerotium gum can be incorporated into the composition by dispersing it first in water, at about a 1:80 polymer-to-water ratio respectively, using a high shear milling device (e.g. IKA T25 DS1 Digital Ultra Turrax homogenizer), adjusting the mill speed and mill time to achieve the desired batch viscosity and rheology. The remaining ingredients were then added to complete the batch.

(213) Additional information on sulfate-free surfactants and other ingredients that are suitable for shampoo compositions is found at U.S. Pub. Nos. 2019/0105247 and 2019/0105246, incorporated by reference.

Test Methods

(214) Argentometry Method to Measure Wt % Inorganic Chloride Salts

(215) The weight % of inorganic chloride salt in the composition can be measured using a potentiometric method where the chloride ions in the composition are titrated with silver nitrate. The silver ions react with the chloride ions from the composition to form an insoluble precipitate, silver chloride. The method

an electrode (Mettler Toledo DM141) that is designed for potentiometric titrations of anions that precipitate with silver. The largest change in the signal occurs at the equivalence point when the amount of added silver ions is equal to the amount of chloride ions in solution. The concentration of silver nitrate solution used should be calibrated using a chloride solution known to one of skill in the art, such as a sodium chloride solution that contains a standard and known amount of sodium chloride to confirm that the results match the known concentration. This type of titration involving a silver ion is known as argentometry and is commonly used to determine the amount of chloride present in a sample.

(216) Methods to Determine Lack of In Situ Coacervate in Composition Prior to Dilution

(217) 1. Microscopy Method to Determine Lack of In Situ Coacervate

(218) Lack of in situ coacervate can be determined using a microscope. The composition is mixed to homogenize, if needed. Then, the composition is sampled onto a microscope slide and mounted on a microscope, per typical microscopy practices. The sample is viewed at, for example, a 10× or 20× objective. If in situ coacervate is present in the sample, an amorphous, gel-like phase with about 20 nm to about 200 nm particle size can be seen throughout the sample. This amorphous, gel-like phases can be described as gel chunks or globs. In this method, the in situ coacervate is separate from other ingredients that were intentionally added to the formula that form flocks or otherwise appear as particles under microscopy.

(219) FIG. 4 is an example microscopy photograph at 20× objective of a marketed sulfate-free shampoo composition that contains a cationic polymer and also has in situ coacervate. FIG. 4 at reference numeral 1 shows an amorphous, gel-like phase that is about 130 nm long that is the in situ coacervate. FIG. 5 is an example microscopy photograph at 10× objective the same marketed shampoo composition that was used in FIG. 4 at 20× objective. FIG. 5 shows many of these amorphous, gel-like phases present with a length from about 20 nm to about 200 nm.

(220) 2. Clarity Assessment—Measurement of % Transmittance (% T)

(221) Lack of in situ coacervate can be determined by composition clarity. A composition that does not contain in situ coacervate will be clear, if it does not contain any ingredients that would otherwise give it a hazy appearance.

(222) Composition clarity can be measured by % Transmittance. For this assessment to determine if the composition lacks coacervate, the composition should be made without ingredients that would give the composition a hazy appearance such as silicones, opacifiers, non-silicone oils, micas, and gums or anionic rheology modifiers. It is believed that adding these ingredients would not cause in situ coacervate to form prior to use, however these ingredients will obscure measurement of clarity by % Transmittance.

(223) Clarity can be measured by % Transmittance (% T) using Ultra-Violet/Visible (UV/VIS) spectrophotometry which determines the transmission of UV/VIS light through a sample. A light wavelength of 600 nm has been shown to be adequate for characterizing the degree of light transmittance through a sample. Typically, it is best to follow the specific instructions relating to the specific spectrophotometer being used. In general, the procedure for measuring percent transmittance starts by setting the spectrophotometer to 600 nm. Then a calibration “blank” is run to calibrate the readout to 100 percent transmittance. A single test sample is then placed in a cuvette designed to fit the specific spectrophotometer and care is taken to ensure no air bubbles are within the sample before the % T is measured by the spectrophotometer at 600 nm. Alternatively, multiple samples can be measured simultaneously by using a spectrophotometer such as the SpectraMax M-5 available from Molecular Devices. Multiple samples are transferred into a 96 well visible flat bottom plate (Greiner part #655-001), ensuring that no air bubbles are within the samples. The flat bottom plate is placed within the SpectraMax M-5 and % T measured using the Software Pro v.5™ software available from Molecular Devices.

(224) 3. Lasentec FBRM Method

(225) Lack of in situ coacervate can also be measured using Lasentec FBRM Method with no dilution. A Lasentec Focused Beam Reflectance Method (FBRM) [model S400A available from Mettler Toledo Corp] may be used to determine floc size and amount as measured by chord length and particle counts/sec (counts per sec).

(226) 4. In Situ Coacervate Centrifuge Method

(227) Lack of in situ coacervate can also be measured by centrifuging a composition and measuring in situ coacervate gravimetrically. For this method, the composition should be made without a suspending agent to allow for separation of an in situ coacervate phase. The composition is centrifuged for 20 minutes at 9200 rpm using a Beckman Coulter TJ25 centrifuge. Several time/rpm combinations can be used. The supernatant is then removed and the remaining settled in situ coacervate assessed gravimetrically. % In Situ Coacervate is calculated as the weight of settled in situ coacervate as a percentage of the weight of composition added to the centrifuge tube using the equation below. This quantifies the percentage of the composition that participates in the in situ coacervate phase.

$$(228) \% \text{InSituCoacervate} = \frac{\text{Weight of settled in situ coacervate}}{\text{Weight of composition added to centrifuge tube}} \times 100$$

Measures of Improved Performance Due to No In Situ Coacervate Prior to Dilution

(229) The composition does not contain in situ coacervate prior to dilution. Because of this, coacervate quantity and quality upon dilution is better than a composition that does contain in situ coacervate prior to dilution. This provides better wet conditioning and deposition of actives from a composition that does not contain coacervate prior to dilution compared to a composition that does contain coacervate prior to dilution.

(230) 1. Measurement of % Transmittance (% T) During Dilution

(231) Coacervate formation upon dilution for a transparent or translucent composition can be assessed using a spectrophotometer to measure the percentage of light transmitted through the diluted sample (% T). As percent light transmittance (% T) values measured of the dilution decrease, typically higher levels of coacervate are formed. Dilutions samples at various weight ratios of water to composition can be prepared, for example 2 parts of water to 1 part composition (2:1), or 7.5 parts of water to 1 part composition (7.5:1), or 16 parts of water to 1 part composition (16:1), or 34 parts of water to 1 part composition (34:1), and the % T measured for each dilution ratio sample. Examples of possible dilution ratios may include 2:1, 3:1, 5:1, 7.5:1, 11:1, 16:1, 24:1, or 34:1. By averaging the % T values for samples that span a range of dilution ratios, it is possible to simulate and ascertain how much coacervate a composition on average would form as a consumer applies the composition to wet hair, lathers, and then rinses it out. Average % T can be calculated by taking the numerical average of individual % T measurements for the following dilution ratios: 2:1, 3:1, 5:1, 7.5:1, 11:1, 16:1, 24:1, and 34:1. Lower average % T indicates more coacervate is formed on average as a consumer applies the composition to wet hair, lathers and then rinses it out.

(232) % T can be measured using Ultra-Violet/Visible (UV/VIS) spectrophotometry which determines the transmission of UV/VIS light through a sample. A light wavelength of 600 nm has been shown to be adequate for characterizing the degree of light transmittance through a sample. Typically, it is best to follow the specific instructions relating to the specific spectrophotometer being used. In general, the procedure for measuring percent transmittance starts by setting the spectrophotometer to 600 nm. Then a calibration "blank" is run to calibrate the readout to 100 percent transmittance. A single test sample is then placed in a cuvette designed to fit the specific spectrophotometer and care is taken to insure no air bubbles are within the sample before the % T is measured by the spectrophotometer at 600 nm.

Alternatively, multiple samples can be measured simultaneously by using a spectrophotometer such as the SpectraMax M-5 available from Molecular Devices. Multiple dilution samples can be prepared within a 96 well plate (VWR catalog #82006-448) and then transferred to a 96 well visible flat bottom plate (Greiner part #655-001), ensuring that no air bubbles are within the sample. The flat bottom plate is placed within the SpectraMax M-5 and % T measured using the Software Pro v.5™ software available from Molecular Devices.

(233) 2. Assessment of Coacervate Floc Size Upon Dilution

(234) Coacervate floc size upon dilution can be assessed visually. Dilutions samples at various weight ratios of water to composition can be prepared, for example 2 parts of water to 1 part composition (2:1), or 7.5 parts of water to 1 part composition (7.5:1), or 16 parts of water to 1 part composition (16:1), or 34 parts of water to 1 part composition (34:1), and the % T measured for each dilution ratio sample. Examples of possible dilution ratios may include 2:1, 3:1, 5:1, 7.5:1, 11:1, 16:1, 24:1, or 34:1.

(235) 3. Wet Combing Force Method

(236) Hair switches of 4 grams general population hair at 8 inches length are used for the measurement.

Each hair switch is treated with 4 cycles (1 lather/rinse steps per cycle, 0.1 gm cleansing composition/gm hair on each lather/rinse step, drying between each cycle) with the cleansing composition. Four switches are treated with each shampoo. The hair is not dried after the last treatment cycle. While the hair is wet, the hair is pulled through the fine tooth half of two Beautician 3000 combs. Force to pull the hair switch through the combs is measured by a friction analyzer (such as Instron or MTS tensile measurement) with a load cell and outputted in gram-force (gf). The pull is repeated for a total of five pulls per switch. Average wet combing force is calculated by averaging the force measurement from the five pulls across the four hair switches treated with each cleansing composition. Data can be shown as average wet combing force through one or both of the two combs.

(237) 4. Deposition Method

(238) Deposition of actives can be measured in vitro on hair tresses or in vivo on panelist's heads. The composition is dosed on a hair tress or panelist head at a controlled amount and washed according to a conventional washing protocol. For a hair tress, the tress can be sampled and tested by an appropriate analytical measure to determine quantity deposited of a given active. To measure deposition on a panelist's scalp, the hair is then parted on an area of the scalp to allow an open-ended glass cylinder to be held on the surface while an aliquot of an extraction solution is added and agitated prior to recovery and analytical determination of a given active. To measure deposition on a panelist's hair, a given amount of hair is sampled and then tested by an appropriate analytical measure to determine quantity deposited of a given active.

(239) Cone/Plate Viscosity Measurement

(240) The viscosities of the examples are measured by a Cone/Plate Controlled Stress Brookfield Rheometer R/S Plus, by Brookfield Engineering Laboratories, Stoughton, MA The cone used (Spindle C-75-1) has a diameter of 75 mm and 1° angle. The liquid viscosity is determined using a steady state flow experiment at constant shear rate of 2 s.sup.-1 and at temperature of 26.7° C. The sample size is about 2.5 ml to about 3 ml and the total measurement reading time is 3 minutes.

(241) Lather Characterization—Kruss DFA100 Lather Characterization

(242) A cleansing composition dilution of 10 parts by weight water to 1 part by weight cleanser is prepared. The shampoo dilution is dispensed into the Kruss DFA100 which generates the lather and measures lather properties.

(243) pH Method

(244) First, calibrate the Mettler Toledo Seven Compact pH meter. Do this by turning on the pH meter and waiting for 30 seconds. Then take the electrode out of the storage solution, rinse the electrode with distilled water, and carefully wipe the electrode with a scientific cleaning wipe, such as a Kimwipe®. Submerge the electrode in the pH 4 buffer and press the calibrate button. Wait until the pH icon stops flashing and press the calibrate button a second time. Rinse the electrode with distilled water and carefully wipe the electrode with a scientific cleaning wipe. Then submerge the electrode into the pH 7 buffer and press the calibrate button a second time. Wait until the pH icon stops flashing and press the calibrate button a third time. Rinse the electrode with distilled water and carefully wipe the electrode with a scientific cleaning wipe. Then submerge the electrode into the pH 10 buffer and press the calibrate button a third time. Wait until the pH icon stops flashing and press the measure button. Rinse the electrode with distilled water and carefully wipe with a scientific cleaning wipe. Submerge the electrode into the testing sample and press the read button. Wait until the pH icon stops flashing and record the value.

EXAMPLES

(245) The following examples further describe and demonstrate embodiments within the scope of the present invention. The examples are given solely for the purpose of illustration and are not to be construed as limitations of the present invention, as many variations thereof are possible without departing from the spirit and scope of the invention.

(246) The following Examples illustrate various shampoo compositions. The Examples in Table 2 and Table 3 were prepared by conventional formulation and mixing techniques. The sclerotium gum was incorporated using high shear milling, as described herein.

(247) The total sodium chloride in the examples in Table 2 and Table 3 was 0.07%. The total sodium chloride in the tables below was calculated based on the product specifications from the suppliers. The

ratio of anionic to amphoteric surfactant in Table 2 and (248) Table 3 was 0.9:1. The ratio of anionic surfactant to amphoteric surfactant is calculated by wt. %. The ratio of polymer charge density to inorganic salt in Table 2 and Table 3 was 18:1.

(249) The ratio of polymer charge density to inorganic salt is the charge density of the polymer (meq/gm) to the wt. % of the inorganic salt, disregarding the units. If the composition contains more than one cationic polymer, then the ratio is calculated according to the polymer with the lowest charge density.

(250) For the examples and comparative examples Table 2 and Table 3, the phase stability was determined as follows. The examples were prepared and immediately put in a clear, glass jar of at least 1 inch width. The cap was screwed on the jar, finger tight. The example was stored at ambient temperatures (20-25° C.), away from direct sunlight, for 5 days. Then the composition was inspected to see if either haze and/or precipitate was visually detectable. The haze and/or precipitate can be suspended across the liquid shampoo composition or a portion thereof and/or at or near the bottom of the container. If either haze or precipitate were present, it was determined that the composition had multiple phases and was not phase stable. FIGS. 6A and 6B show Comparative Example C5 (containing 0.5% konjac gum thickener) and C6 (containing 0.5% gellan gum thickener), respectively. A precipitate can be seen towards the bottom of the container in both FIGS. 6A and 6B. If neither haze nor precipitate were present, it was determined that there was no in situ coacervate and the composition was a stable, single phase. It is believed that the stable shampoo examples would have improved product performance as compared to the examples that were not phase stable.

(251) As used herein, “visually detect” or “visually detectable” means that a human viewer can visually discern the quality of the example with the unaided eye (excepting standard corrective lenses adapted to compensate for near-sightedness, farsightedness, or stigmatism, or other corrected vision) in lighting at least equal to the illumination of a standard 100 watt incandescent white light bulb at a distance of 1 meter.

(252) The examples in Table 2 and Table 3, could also be formulated with silicones, opacifiers (e.g. glycol distearate, glycol stearate), non-silicone oils, micas, gums or anionic rheology modifiers and other ingredients that would cause the shampoo to have a hazy or hazier appearance. However, it is believed that adding these ingredients would not cause in situ coacervate to form or other phase instability prior to use.

(253) TABLE-US-00002 TABLE 2 Comparative Shampoo Composition Examples

	C1	C2	C3	C4	C5	C6
One	One	Multiple	Multiple	Multiple	Multiple	Phase
Stability	phase	phase	phase	phases	phases	phases
pH	5.62	5.54	5.75	5.81	5.79	5.87
Viscosity @ 2 s ⁻¹ (cps)	2236	893	4846	8588	3896	3053
Lauramidopropyl Betaine .sup.1	9.75	9.75	9.75	9.75	9.75	9.75
Sodium Cocoyl Isethionate .sup.2	6.00	6.00	6.00	6.00	6.00	6.00
Sodium Lauryl Sarcosinate .sup.3	2.50	2.50	2.50	2.50	2.50	2.50
Polyquaternium 10 .sup.4	0.55	0.25	0.55	0.55	0.55	0.55
(JR-30M, CD 1.25 meq/g)						
Xanthan Gum .sup.8	0.20	0.20	0.20	0.20	0.20	0.20
Carrageenan .sup.9	0.20	0.20	0.20	0.20	0.20	0.20
Konjac Gum .sup.10	0.50	0.50	0.50	0.50	0.50	0.50
Gellan Gum .sup.11	0.50	0.50	0.50	0.50	0.50	0.50
Citric Acid .sup.12						
To pH 5.5-6.1						
Water, Preservatives, Q.S.						
to 100						
Perfume, and Optional Components						

(254) C1 and C2 in Table 2 were phase stable. However, neither example contains a thickener, and the viscosity is too low to be consumer preferred. It is suspected that these examples will be difficult to hold in a user's palm and apply across the hair.

(255) C3-C6 contained an anionic thickener (e.g. xanthan gum, carrageenan, konjac gum, and gellan gum) separated into multiple phases, which is believed to not only impact the appearance of the product, but will also impact product performance. Thus, C3-C6 will likely not be consumer preferred.

(256) Yield stress was not measured for the examples in Table 2 because the examples were not consumer acceptable.

(257) TABLE-US-00003 TABLE 3 Shampoo Composition Examples

	Ex. 1	Ex. 2	Ex. 3	Ex. 4
Stability	One	One	One	One
phase	phase	phase	phase	phase
pH	5.86	5.84	5.82	5.50
Viscosity @ 2 s ⁻¹ (cps)	4192	2681	4889	10615
Yield Stress, Herschel-0.004	0.008	0.017	0.16	0.16
Bulkley @ shear rate 10.sup.-2 to 10.sup.-4 s.sup.-1 (Pa)				
Lauramidopropyl Betaine .sup.1	9.75	9.75	9.75	9.75
Sodium Cocoyl Isethionate .sup.2	6.00	6.00	6.00	6.00
Sodium Lauryl Sarcosinate .sup.3	2.50	2.50	2.50	2.50
Polyquaternium 10 .sup.4	0.55	0.55	0.55	0.55
(JR-30M, CD 1.25 meq/g)				
Sclerotium Gum .sup.5,6	0.50	1.00	0.50	1.00
Sclerotium Gum and Citrus 0.50	1.00	1.00	1.00	1.00
Limon Peel Powder .sup.7				
Citric Acid .sup.12				
To pH 5.5-6.1				
Water,				

Preservatives, Q.S. to Q.S. to Q.S. to Q.S. to Perfume, and Optional 100 100 100 100 Components

The Examples in

(258) Table 3 were phase stable, have a pH that prevents the surfactant from hydrolyzing (e.g. greater than or equal to 5.5), and have a consumer acceptable viscosity (e.g. greater than or equal to 2700 Pa). It is believed that these examples would be consumer preferred, as compared to the comparative examples in Table 2.

(259) The Examples in Table 4 and

(260) Table 5 could be made and it is believed that these examples would be stable and have consumer acceptable viscosity. The Examples in Table 4 would have a total sodium chloride of 0.7% based on the specifications from suppliers and a ratio of polymer charge density to inorganic salt of 18:1.

(261) TABLE-US-00004 TABLE 4 Ex. 5 Ex. 6 Ex. 7 Lauramidopropyl Betaine .sup.1 9.75 9.75 9.75 Sodium Cocoyl Isethionate .sup.2 6.00 6.00 6.00 Sodium Lauryl Sarcosinate .sup.3 4.00 Polyquaternium 10 .sup.4 0.55 0.55 0.55 (JR-30M, CD 1.25 meq/g) Sclerotium Gum .sup.5,6 0.50 0.50 0.50 Sodium Benzoate .sup.13 0.75 0.75 0.45 Sodium Salicylate .sup.14 0.45 0.45 Perfume 1.00 1.00 1.00 Citric Acid .sup.12 To pH 5.5-6.5 Water QS to 100 QS to 100 QS to 100 Number of Ingredients 8 9 10 Ratio of Anionic to 0.6:1 0.6:1 1.0:1 Amphoteric Surfactant

(262) TABLE-US-00005 TABLE 5 Ex. 8 Ex. 9 Ex. 10 Ex. 11 Ex. 12 Ex. 13 Ex. 14 Lauramidopropyl Betaine .sup.1 9.75 9.75 5.36 2.44 2.44 Low Salt Cocamidopropyl 9.75 Betaine .sup.15 Cocamidopropyl Betaine .sup.16 4.39 7.31 7.31 7.50 Sodium Cocoyl Isethionate .sup.2 6.00 6.00 6.00 6.00 6.00 6.00 4.50 Sodium Lauryl Sarcosinate .sup.3 2.50 2.50 2.50 2.50 2.50 2.50 Polyquaternium 10 .sup.4 0.55 0.55 0.55 0.55 (JR-30M, CD 1.25 meq/g) Polyquaternium 10 .sup.19 0.10 0.40 0.25 (KG-30M, CD 1.9 meq/g) Sclerotium Gum .sup.5, 6 0.50 1.50 0.50 0.50 0.50 0.50 0.50 Piroctone Olamine .sup.17 0.50 Zinc Pyrithione .sup.18 1.00 Water, Preservatives, pH QS to QS to QS to QS to QS to QS to QS to adjusters, Perfume, and 100 100 100 100 100 100 100 100 Optional Components Ratio of Anionic to 0.9:1 0.9:1 0.9:1 0.9:1 0.9:1 0.9:1 0.6:1 Amphoteric Surfactant Total Sodium Chloride 0.07 0.07 0.06 0.81 1.3 1.3 1.3 (including from surfactant) Ratio of Polymer Charge 18:1 18:1 21:1 1.5:1 1.5:1 1.5:1 1.5:1 Density to Inorganic Salt

Suppliers for Examples in Table 2 to Table 5 1. Mackam® DAB-ULS available from Solvay®.

Specification Range: Solids=34-36%, Sodium Chloride=0-0.5%. Average values are used for

calculations: Actives=35%, Sodium Chloride=0.25%. 2. Hostapon® SCI-85 C available from Clariant®

3. SP Crodasinic™ LS30/NP MBAL available from Croda® 4. UCARE™ Polymer JR-30M available

from Dow® 5. Sclerotium Gum, Amigum ER available from Alban Muller® 6. Sclerotium Gum,

Actigum® CS 11 QD available from Cargill® 7. Sclerotium Gum and Citrus Limon Peel Powder, Fiber

Design Sensation® available from Cargill® 8. Xanthan Gum, Keltrol® CG-SFT available from CP

Kelco® 9. Carrageenan, GENUVISCO® Carrageenan CG-131 available from CP Kelco® 10. Konjac

Gum, Nutricol® XP 3464 available from FMC Corporation 11. Gellan Gum, Kelcogel® CG-LA

available from CP Kelco® 12. Citric Acid USP Anhydrous Fine Granular available from ADM® 13.

Sodium Benzoate available from Kalama® 14. Sodium Salicylate available from JQC (Huayin)

Pharmaceutical Co., Ltd 15. Dehyton® PK 45 from BASF® with Sodium Chloride removed, resulting in

33.05% Dry Residue, 0.21% Sodium Chloride, 32.84% Active used for calculations. 16. TEGO® Betain

CK PH 12 available from Evonik®. Specification Range: Actives=28-32%, Sodium Chloride=4.5-6%.

Average values are used for calculations: Actives=30%, Sodium Chloride=5.25%. 17. Octopirox®

available from Clariant® 18. Zinc Pyrithione available from Lonza® 19. UCARE™ Polymer KG-30M

available from Dow®

(263) The dimensions and values disclosed herein are not to be understood as being strictly limited to the exact numerical values recited. Instead, unless otherwise specified, each such dimension is intended to mean both the recited value and a functionally equivalent range surrounding that value. For example, a dimension disclosed as “40 mm” is intended to mean “about 40 mm.”

(264) Every document cited herein, including any cross referenced or related patent or application and any patent application or patent to which this application claims priority or benefit thereof, is hereby incorporated herein by reference in its entirety unless expressly excluded or otherwise limited. The citation of any document is not an admission that it is prior art with respect to any invention disclosed or claimed herein or that it alone, or in any combination with any other reference or references, teaches,

suggests or discloses any such invention. Further, to the extent that any meaning or definition of a term in this document conflicts with any meaning or definition of the same term in a document incorporated by reference, the meaning or definition assigned to that term in this document shall govern.

(265) While particular embodiments of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

Claims

1. A phase-stable shampoo composition comprising: a) a surfactant system comprising: (i) about 3% to about 35% of an anionic surfactant comprising sodium, ammonium or potassium salts of isethionates, sodium, ammonium or potassium salts of sarcosinates, or a combination thereof; (ii) about 5% to about 15% of an amphoteric surfactant comprising a betaine, wherein the surfactant system is substantially free of sulfated surfactants; b) about 0.01% to about 2% of a cationic polymer comprising polyquaternium-10; and c) about 0.15% to about 1.5% of a thickener comprising sclerotium gum, wherein the composition has a viscosity of greater than 2500 cP and a pH of greater than 5.5.
 2. The phase-stable shampoo composition of claim 1, further comprising less than about 1% inorganic salt.
 3. The phase-stable shampoo composition of claim 2, wherein the inorganic salt is selected from sodium chloride, potassium chloride, sodium sulfate, ammonium chloride, sodium bromide, and combinations thereof.
 4. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition has a viscosity of greater than 4000 cP.
 5. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition has a pH greater than 5.7.
 6. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition has a yield stress of greater than 0.003 Pa at a shear rate 10^{-2} to 10^{-4} s $^{-1}$ according to the Herschel-Bulkley model.
 7. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition has a ratio of polymer charge density to total inorganic salt of greater than or equal to 1.2:1.
 8. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition is substantially free of alkyl polyglucoside.
 9. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition is substantially free of fatty esters.
 10. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition has a percent transparency % T at 600 nm of greater than 70%.
 11. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition lacks an in situ coacervate, as determined by the Microscopy Method to Determine Lack of In Situ Coacervate.
 12. The phase-stable shampoo composition of claim 1, further comprising an antidandruff agent.
 13. The phase-stable shampoo composition of claim 12, wherein the antidandruff agent is selected from piroctone olamine, zinc pyrithione, and combinations thereof.
 14. The phase-stable shampoo composition of claim 1, wherein the phase-stable shampoo composition is substantially free of silicones.
 15. The phase-stable shampoo composition of claim 1, wherein the composition has 9 or fewer ingredients.
 16. A method of making the phase-stable shampoo composition of claim 1, wherein the sclerotium gum is added using high shear milling.
-