



US012410455B2

(12) **United States Patent**
Lee et al.

(10) **Patent No.:** **US 12,410,455 B2**
(45) **Date of Patent:** **Sep. 9, 2025**

(54) **PHOSPHOENOLPYRUVATE CARBOXYLASE
VARIANT AND METHOD FOR PRODUCING
5'-INOSINIC ACID USING SAME**

(71) Applicant: **DAESANG CORPORATION**, Seoul
(KR)

(72) Inventors: **Sun Hee Lee**, Seoul (KR); **Tae Yeol
Choi**, Seoul (KR); **Hyun Ho Kim**,
Seoul (KR); **Dong Hyun Kim**, Seoul
(KR); **Hyun Sook Kim**, Seoul (KR);
Jong Hwan Shin, Seoul (KR)

(73) Assignee: **DAESANG CORPORATION**, Seoul
(KR)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **18/879,948**

(22) PCT Filed: **Aug. 31, 2023**

(86) PCT No.: **PCT/KR2023/012968**
§ 371 (c)(1),
(2) Date: **Dec. 30, 2024**

(87) PCT Pub. No.: **WO2024/210280**
PCT Pub. Date: **Oct. 10, 2024**

(65) **Prior Publication Data**
US 2025/0257379 A1 Aug. 14, 2025

(30) **Foreign Application Priority Data**
Apr. 6, 2023 (KR) 10-2023-0045160

(51) **Int. Cl.**
C12P 19/32 (2006.01)
C12N 9/88 (2006.01)

(52) **U.S. Cl.**
CPC **C12P 19/32** (2013.01); **C12N 9/88**
(2013.01); **C12Y 401/01031** (2013.01)

(58) **Field of Classification Search**
CPC **C12P 19/32**; **C12N 9/88**; **C12Y 401/01031**
See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

KR	10-1166027	7/2012
KR	10-2016-0078694	7/2016
KR	10-2016-0125170	10/2016
KR	10-1904675	10/2018
KR	10-1916622	11/2018

OTHER PUBLICATIONS

International Search Report issued Jan. 19, 2024 in International
(PCT) Application No. PCT/KR2023/012968.


Primary Examiner — Iqbal H Chowdhury

(74) *Attorney, Agent, or Firm* — Wenderoth, Lind &
Ponack, L.L.P.

(57) **ABSTRACT**

The present invention relates to a novel phosphoenolpyruvate carboxylase variant and a method of producing 5'-inosinic acid using the same. The phosphoenolpyruvate carboxylase variant is obtained by substituting one or more amino acids in the amino acid sequence constituting phosphoenolpyruvate carboxylase to change the activity of the protein, and a recombinant microorganism comprising the phosphoenolpyruvate carboxylase variant is capable of efficiently producing 5'-inosinic acid.

8 Claims, 1 Drawing Sheet
Specification includes a Sequence Listing.



1

PHOSPHOENOLPYRUVATE CARBOXYLASE VARIANT AND METHOD FOR PRODUCING 5' ϵ -A, γ -A, ϵ -INOSINIC ACID USING SAME

SEQUENCE LISTING

A sequence listing in electronic (XML file) format is filed with this application and incorporated herein by reference. The name of the XML file is "Sequence Listing-1947A.xml"; the file was created on Dec. 29, 2024; the size of the file is 20,189 bytes.

TECHNICAL FIELD

The present invention relates to a novel phosphoenolpyruvate carboxylase variant and a method of producing 5'-inosinic acid using the same.

BACKGROUND ART

5'-inosinic acid (or inosine monophosphate (IMP)) is an intermediate in the metabolic system of nucleic acid biosynthesis, and not only plays an important physiological role in the bodies of plants and animals, but is also used in various applications, including food, medicine, and various medical applications. In particular, 5'-inosinic acid is a nucleic acid-based seasoning, which has drawn much attention as a savory seasoning, because it has significant synergistic effects on taste when used together with monosodium glutamate (MSG).

Methods for producing 5'-inosinic acid include a method of enzymatically degrading ribonucleic acid extracted from yeast cells, a method of chemically phosphorylating inosine produced by fermentation, etc. Recently, a method of culturing a 5'-inosinic acid-microorganism and recovering 5'-inosinic acid producing accumulated in the medium has been mainly used.

For the production of 5'-inosinic acid using microorganisms, in order to improve the efficiency of production of 5'-inosinic acid, there has been development of a variety of recombinant strains or mutant strains having excellent 5'-inosinic acid productivity by applying genetic recombination technology to microorganisms such as *Escherichia coli* and *Corynebacterium*, which are widely used in the production of useful substances such as nucleic acids or L-amino acids, and methods of producing 5'-inosinic acid using the same. In particular, there have been attempts to increase the production of 5'-inosinic acid by targeting genes such as enzymes, transcription factors and transport proteins, which are involved in the biosynthetic pathway of 5'-inosinic acid, or by inducing mutations in promoters that regulate the expression of these genes. However, there are dozens to hundreds of types of proteins such as enzymes, transcription factors and transport proteins, which involved directly or are indirectly in the production of 5'-inosinic acid, and thus much research is still needed on the increase in 5'-inosinic acid productivity by changes in the activity of these proteins.

PRIOR ART DOCUMENTS

Patent Documents

Korean Patent No. 10-116602

DISCLOSURE

Technical Problem

An object of present invention is to provide a novel phosphoenolpyruvate carboxylase variant.

2

Another object of the present invention is to provide a polynucleotide encoding the variant.

Still another object of the present invention is to provide a transformant comprising the variant Or polynucleotide.

Yet another object of the present invention is to provide a method of producing 5'-inosinic acid using the transformant.

Technical Solution

One aspect of the present invention provides a phosphoenolpyruvate carboxylase variant in which one or more of the amino acids at positions 472 and 673 in the amino acid sequence of SEQ ID NO: 4 is/are substituted with other amino acid(s).

As used in the present invention, the term "phosphoenolpyruvate carboxylase" catalyzes a reaction of removing carbon dioxide from phosphoenolpyruvate, and it may be a polypeptide or protein consisting of the amino acid sequence of SEQ ID NO: 4 and having phosphoenolpyruvate carboxylase activity.

Information on the nucleic acid and protein sequences of the phosphoenolpyruvate carboxylase is available from known sequence databases (e.g., GenBank, UniProt).

According to one embodiment of the present invention, the phosphoenolpyruvate carboxylase may be encoded by the nucleotide sequence of SEQ ID NO: 3.

The amino acid sequence of the phosphoenolpyruvate carboxylase according to the present invention or the nucleotide sequence encoding the same may include a nucleotide sequence or amino acid sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% homology or identity to the amino acid sequence of SEQ ID NO: 4 or the nucleotide sequence of SEQ ID NO: 3. As used herein, the term "homology" or "identity" means the percentage rate of identity between two sequences, which is determined by aligning the reference nucleotide sequence or amino acid sequence and any other nucleotide sequence of amino acid sequence to correspond to each other as much as possible and analyzing the aligned sequences.

According to one embodiment of the present invention, the phosphoenolpyruvate carboxylase may be derived from wild-type *Corynebacterium stationis*.

As used in the present invention, the term "variant" refers to a protein that has an amino acid sequence different from the amino acid sequence before mutation by the conservative substitution and/or modification of one or more amino acids at the N-terminus, C-terminus of and/or within the amino acid sequence, which result(s) from mutation in the nucleotide sequence of the gene encoding the protein, but retains the functions or properties of the protein before mutation. As used herein, the term "conservative substitution" means substituting one amino acid with another amino acid having similar structural and/or chemical properties.

The conservative substitution may have little or no impact on the activity of the protein or polypeptide. In addition, the term "modification" refers to substitution, insertion, deletion, or the like of one or more amino acids. The amino acid is selected from among alanine (Ala, A), isoleucine (Ile, I), valine (Val, V), leucine (Leu, L), methionine (Met, M), asparagine (Asn, N), cysteine (Cys, C), glutamine (Gln, Q), serine (Ser, S), threonine (Thr, T), phenylalanine (Phe, F), tryptophan (Trp, W), tyrosine (Tyr, Y), aspartic acid (Asp, D), glutamic acid (Glu, E), arginine (Arg, R), histidine (His, H), lysine (Lys, K), glycine (Gly, G), and proline (Pro, P).

In addition, some variants include those in which one or more portions, such as an N-terminal leader sequence of

transmembrane domain, have been removed, or those in which a portion has been removed from the N- and/or C-terminus of a mature protein.

The variant may have increased (enhanced), unchanged, or decreased (weakened) ability compared to that of the protein before mutation. Here, the term “increased or enhanced” includes: a case in which the activity of the protein itself has increased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is higher than that in the wild-type strain or the strain expressing the protein before mutation due to increased expression or translation of the gene encoding the protein; and a combination thereof. In addition, the term “decreased or weakened” includes: a case in which the activity of the protein itself has decreased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is lower than that in the wild-type strain or the strain expressing the protein before mutation due to reduced expression or translation of the gene encoding the protein; and a combination thereof. In the present invention, the term “variant” may be used interchangeably with terms such as variant type, modification, variant polypeptide, mutated protein, mutation, and the like.

According to one embodiment of the present invention, the variant may consist of the amino acid sequence of SEQ ID NO: 2 in which proline at position 472 in the amino acid sequence of SEQ ID NO: 4 is substituted with serine and glycine at position 673 is substituted with aspartic acid.

More specifically, the phosphoenolpyruvate carboxylase variant may comprise an amino acid sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100% homology or identity to the amino acid sequence of SEQ ID NO: 2.

Another aspect of the present invention provides a polynucleotide encoding the phosphoenolpyruvate carboxylase variant.

As used in the present invention, the term “polynucleotide” refers to a DNA or RNA strand having a certain length or more, which is a long-chain polymer of nucleotides formed by linking nucleotide monomers via covalent bonds. More specifically, the term “polynucleotide” refers to a polynucleotide fragment encoding the variant.

According to one embodiment of the present invention, the polynucleotide may comprise nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2.

More specifically, the polynucleotide may comprise the nucleotide sequence of SEQ ID NO: 1 in which nucleotide “c” at position 1414 in the nucleotide sequence of SEQ ID NO: 3 encoding phosphoenolpyruvate carboxylase is substituted with nucleotide “t” and nucleotide “g” at position 2018 is substituted with nucleotide “a”.

Still another aspect of the present invention provides a vector comprising a polynucleotide encoding the phosphoenolpyruvate carboxylase variant.

Yet another aspect of the present invention provides a transformant comprising the phosphoenolpyruvate carboxylase variant or the polynucleotide.

As used in the present invention, the term “vector” refers to any type of nucleic acid sequence transfer structure that is used as a means for transferring and expressing a gene of interest in a host cell. Unless otherwise specified, the term “vector” may mean one allowing the nucleic acid sequence contained therein to be expressed after insertion into the host cell genome and/or one allowing the nucleic acid sequence to be expressed independently. This vector comprises essential regulatory elements operably linked so that an inserted

gene can be expressed. As used herein, the term “operably linked” means that a gene of interest and regulatory sequences thereof are functionally linked together in a manner enabling gene expression, and the “regulatory elements” include a promoter for initiating transcription, any operator sequence for regulating transcription, a sequence encoding suitable mRNA ribosome-binding sites, and a sequence for regulating termination of transcription and translation.

The vector in the present invention is not particularly limited as long as it may replicate in a host cell, and any vector known in the art may be used. Examples of the vector include a natural or recombinant plasmid, cosmid, virus and bacteriophage. Examples of a phage vector or cosmid vector include, but are not limited to, pWE15, M13, λ MBL3, λ MBL4, λ IXII, λ ASHII, λ APII, λ t10, λ t11, Charon4A, and Charon21A, and examples of a plasmid vector include, but are not limited to, pBR series, pUC series, pBluescriptII series, pGEM series, pTZ series, pCL series, and pET series.

The vector may typically be constructed as a vector for cloning or as a vector for expression. The vector for expression may be a conventional vector that is used in the art to express a foreign gene or protein in a plant, animal, or microorganism, and may be constructed through various methods known in the art.

As used in the present invention, the term “recombinant vector” may be transformed into a suitable host cell, and then may replicate regardless of the genome of the host cell or may be integrated into the genome itself. In this case, the “suitable host cell” may contain a replication origin, which is a particular nucleotide sequence which enables the vector to replicate in the suitable host cell and from which replication starts. For example, when the vector used is an expression vector and uses a prokaryotic cell as a host, the vector generally comprises a strong promoter capable of promoting transcription (e.g., PL λ promoter, CMV promoter, trp promoter, lac promoter, tac promoter, T7 promoter, etc.), a ribosome binding site for initiation of translation, and a transcription/translation termination sequence. When a eukaryotic cell is used as a host, the vector comprises a replication origin operating in the eukaryotic cell, and examples of the replication origin include, but are not limited to, an f1 replication origin, an SV40 replication origin, a pMB1 replication origin, an adeno replication origin, an AAV replication origin, and a BBV replication origin. In addition, the recombinant vector may comprise a promoter derived from the genome of a mammalian cell (e.g., metallothionein promoter) or a promoter derived from a mammalian virus (e.g., adenovirus late promoter, vaccinia virus 7.5K promoter, SV40 promoter, cytomegalovirus promoter, HSV-tk promoter, etc.), and generally has polyadenylation sequence transcription termination sequence.

The recombinant vector may comprise a selection marker. The selection marker serves to select a transformant (host cell) transformed with the vector, and since only cells expressing the selection marker can survive in the medium treated with the selection marker, it is possible to select transformed cells. Representative examples of the selection marker include, but are not limited to, kanamycin, streptomycin, and chloramphenicol.

The transformant may be produced by inserting the recombinant vector into a host cell, and the transformant may be obtained by introducing the recombinant vector into an appropriate host cell. The host cell is a cell capable of stably and continuously cloning or expressing the expression vector, and any host cell known in the art may be used.

Where the vector is transformed into prokaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, *E. coli* sp. strains such as *E. coli* DH5 α , *E. coli* JM109, *E. coli* BL21, *E. coli* RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776, *E. coli* W3110, and *E. coli* XL1-Blue, *Bacillus* sp. strains such as *Bacillus subtilis* and *Bacillus thuringiensis*, *Corynebacterium* sp. strains such as *Corynebacterium glutamicum* and *Corynebacterium stationis*, and various Enterobacteriaceae strains such as *Salmonella typhimurium*, *Serratia marcescens*, and *Pseudomonas* species.

Where the vector is transformed into eukaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, yeast (e.g., *Saccharomyces cerevisiae*), insect cells, plant cells and animal cells, such as Sp2/0, CHO K1, CHO DG44, PER.C6, W138, BHK, COS7, 293, HepG2, Huh7, 3T3, RIN, and MDCK cell lines.

As used in the present invention, the term “transformation” refers to a phenomenon in which external DNA is introduced into a host cell, thereby artificially causing genetic changes, and the term “transformant” refers to a host cell into which external DNA has been introduced and in which the expression of the gene of interest is stably maintained.

The transformation may be performed using a suitable vector introduction technique selected depending on the host cell, so that the gene of interest or a recombinant vector comprising the same may be expressed in the host cell. For example, introduction of the vector may be performed by electroporation, heat-shock, calcium phosphate (CaPO₄) precipitation, calcium chloride (CaCl₂) precipitation, microinjection, polyethylene glycol (PEG) method, DEAE-dextran method, cationic liposome method, lithium acetate-DMSO method, or any combination thereof, without being limited thereto. As long as the transformed gene may be expressed in the host cell, it may be inserted into the chromosome of the host cell, or exist extrachromosomally, without being limited thereto.

The transformant may include a cell transfected, transformed, or infected with the recombinant vector of the present invention in vivo or in vitro, and may be used in the same sense as a recombinant host cell, a recombinant cell, or a recombinant microorganism.

Genes inserted into the recombinant vector of the present invention may be introduced into a host cell such as a *Corynebacterium* sp. strain by homologous recombination crossover.

According to one embodiment of the present invention, the transformant may be a *Corynebacterium* sp. microorganism.

The *Corynebacterium* sp. microorganism may be, but is not limited to, *Corynebacterium glutamicum*, *Corynebacterium crudilactis*, *Corynebacterium deserti*, *Corynebacterium callunae*, *Corynebacterium suranareae*, *Corynebacterium lubricantis*, *Corynebacterium doosanense*, *Corynebacterium efficiens*, *Corynebacterium uterequi*, *Corynebacterium stationis*, *Corynebacterium pacaense*, *Corynebacterium singulare*, *Corynebacterium humireducens*, *Corynebacterium marinum*, *Corynebacterium halotolerans*, *Corynebacterium spheniscorum*, *Corynebacterium freiburgense*, *Corynebacterium striatum*, *Corynebacterium canis*, *Corynebacterium ammoniagenes*, *Corynebacterium renale*, *Corynebacterium pollutisoli*, *Corynebacterium imitans*, *Corynebacterium caspium*, *Corynebacterium testudinis*, *Corynebacterium pseudopelargi*, or *Corynebacterium flavesens*.

The transformant in the present invention may be a strain either comprising the above-described phosphoenolpyruvate carboxylase variant or a polynucleotide encoding the same or comprising the vector comprising the same, a strain expressing the phosphoenolpyruvate carboxylase variant or the polynucleotide, or a strain having activity for the phosphoenolpyruvate carboxylase variant, without being limited thereto.

The transformant of the present invention may comprise other protein variants or genetic mutations, in addition to the phosphoenolpyruvate carboxylase variant.

According to one embodiment of the present invention, the transformant may have the ability to produce 5'-inosinic acid.

The 5'-inosinic acid is a nucleic acid-based compound that gives flavor to food, especially umami (savory) taste, and is used with the same meaning as inosine monophosphate (IMP).

The transformant may naturally have the ability to produce 5'-inosinic acid or may be one artificially endowed with the ability to produce 5'-inosinic acid.

According to one embodiment of the present invention, the transformant may have an increased ability to produce 5'-inosinic acid, due to a change in phosphoenolpyruvate carboxylase activity.

As used in the present invention, the term “increased ability to produce” means that 5'-inosinic acid productivity has increased compared to that of the parent strain. As used herein, the term “parent strain” refers to a wild-type strain or mutant strain to be mutated, and includes a strain that is to be mutated directly or to be transformed with a recombinant vector or the like. In the present invention, the parent strain may be a wild-type *Corynebacterium* sp. strain or a *Corynebacterium* sp. strain mutated from the wild-type microorganism.

The transformant according to the present invention exhibits an increased ability to produce 5'-inosinic acid compared to the parent strain, due to the change in phosphoenolpyruvate carboxylase activity caused by introduction of the phosphoenolpyruvate carboxylase variant thereinto. More specifically, the amount of 5'-inosinic acid produced by the transformant may be at least 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% higher than that produced by the parent strain, or may be 1.1-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 3.5-fold, 4-fold, 4.5-fold, 5-fold, 5.5-fold, 6-fold, 6.5-fold, 7-fold, 7.5-fold, 8-fold, 8.5-fold, 9-fold, 9.5-fold, or 10-fold higher than that produced by the parent strain, without being limited thereto. For example, the amount of 5'-inosinic acid produced by the transformant comprising the phosphoenolpyruvate carboxylase variant may be at least 5%, specifically 5 to 50% (preferably 10 to 40%) higher than that produced by the parent strain.

Still yet another aspect of the present invention provides a method for producing 5'-inosinic acid, comprising steps of: culturing the transformant in a medium; and recovering 5'-inosinic acid from the transformant or the medium in which the transformant has been cultured.

The culturing may be performed using a suitable medium and culture conditions known in the art, and any person skilled in the art may easily adjust and use the medium and the culture conditions. Specifically, the medium may be a liquid medium, without being limited thereto. Examples of the culturing method include, but are not limited to, batch culture, continuous culture, fed-batch culture, or a combination thereof.

According to one embodiment of the present invention, the medium should meet the requirements of a specific strain in a proper manner, and may be appropriately modified by a person skilled in the art. For culture media for *Escherichia* sp. strains, reference may be made to, but not limited to, a known document (Manual of Methods for General Bacteriology, American Society for Bacteriology, Washington D.C., USA, 1981).

According to one embodiment of the present invention, the medium may contain various carbon sources, nitrogen sources, and trace element components. Examples of carbon sources that may be used include: sugars and carbohydrates such as glucose, sucrose, lactose, fructose, maltose, starch, and cellulose; oils and fats such as soybean oil, sunflower oil, castor oil, and coconut oil; fatty acids such as palmitic acid, stearic acid, and linoleic acid; alcohols such as glycerol and ethanol; and organic acids such as acetic acid. These substances may be used individually or as a mixture, without being limited thereto. Examples of nitrogen sources that may be used include peptone, yeast extract, meat extract, malt extract, corn steep liquor, soybean meal, urea, or inorganic compounds such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, and ammonium nitrate. The nitrogen sources may also be used individually or as a mixture, without being limited thereto. Examples of phosphorus sources that may be used include, but are not limited to, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts. In addition, the culture medium may contain, but is not limited to, metal salts such as magnesium sulfate or iron sulfate, which are required for growth. In addition, the culture medium may contain essential growth substances such as amino acids and vitamins. Moreover, suitable precursors may be used in the culture medium. The medium or individual components may be added to the culture medium batchwise or in a continuous manner by a suitable method during culturing, without being limited thereto.

According to one embodiment of the present invention, the pH of the culture medium may be adjusted by adding compounds such as ammonium hydroxide, potassium hydroxide, ammonia, phosphoric acid and sulfuric acid to the microorganism culture medium in an appropriate manner during the culturing. In addition, during the culturing, foaming may be suppressed using an anti-foaming agent such as a fatty acid polyglycol ester. Additionally, to keep the culture medium in an aerobic condition, oxygen or an oxygen-containing gas (for example, air) may be injected into the culture medium. The temperature of the culture medium may be generally 20° C. to 45° C., for example, 25° C. to 40° C. The culturing may be continued until a desired amount of a useful substance is produced. For example, the culturing time may be 10 hours to 160 hours.

According to one embodiment of the present invention, in the step of recovering 5'-inosinic acid from the cultured transformant the medium in which the transformant has been cultured, the produced 5'-inosinic acid may be collected or recovered from the medium using a suitable method known in the art depending on the culture method. Examples of a method that may be used to recover the produced 5'-inosinic acid include, but are not limited to, centrifugation, filtration, extraction, spraying, drying, evaporation, precipitation, crystallization, electrophoresis, fractional dissolution (e.g., ammonium sulfate precipitation), chromatography (e.g., ion exchange, affinity, hydrophobicity and size exclusion), and the like.

According to one embodiment of the present invention, the step of recovering 5'-inosinic acid may be performed by centrifuging the culture medium at a low speed to remove biomass and separating the obtained supernatant through ion-exchange chromatography.

According to one embodiment of the present invention, the step of recovering 5'-inosinic acid may include a process of purifying the 5'-inosinic acid.

Advantageous Effects

The phosphoenolpyruvate carboxylase variant according to the present invention is obtained by substituting one or more amino acids in the amino acid sequence constituting phosphoenolpyruvate carboxylase to change the activity of the protein, and a recombinant microorganism comprising the phosphoenolpyruvate carboxylase variant is capable of efficiently producing 5'-inosinic acid.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 shows the structure of a pk19msb plasmid according to one embodiment of the present invention.

MODE FOR INVENTION

Hereinafter, the present invention will be described in more detail. However, this description is merely presented by way of example to facilitate the understanding of the present invention, and the scope of the present invention is not limited by this exemplary description.

Example 1. Construction of Strain Expressing Phosphoenolpyruvate Carboxylase Variant

To evaluate the effect of a variant (SEQ ID NO: 2) having a substitution of serine(S) for proline (P) at position 472 and a substitution of aspartic acid (D) for glycine (G) at position 673 in the amino acid sequence of phosphoenolpyruvate carboxylase (SEQ 4) on the production of 5'-inosinic acid, the present inventors constructed a vector for expressing the phosphoenolpyruvate carboxylase variant and a strain into which the vector has been introduced.

1-1. Construction of Vector for Expression of Phosphoenolpyruvate Carboxylase Variant

Using the genomic DNA of wild-type *Corynebacterium stationis* ATCC6872 as a template, PCR was performed using a primer pair of primers 1 and 2. The PCR fragment and a pK19msb plasmid (SEQ ID NO: 5) were treated with the restriction enzyme *smal* (NEB) and ligated together using T4 ligase. The resulting plasmid was named pK_PC.

The PCR amplification was performed using Pfu PreMix (Bioneer) under the following conditions: denaturation at 95° C. for 5 min, and then 30 cycles, each consisting of 95° C. for 30 sec, 58° C. for 30 sec, and 72° C. for 1 min and 30 sec, followed by reaction at 72° C. for 5 min.

The primer sequences used for plasmid construction are shown in Table 1 below.

TABLE 1

Primer name	SEQ ID NO.	Primer sequence (5'-3')
Primer 1	6	CTCTGCTCGCTCACTGGTCT
Primer 2	7	TACTCGCTGTGTCGTAGCGGG

1-2. Construction of Mutant Strain into which Phosphoenolpyruvate Carboxylase Variant has been Introduced

An electrocompetent cell preparation method, a modification of the method of van der Rest et al., was used as a method for transformation of *Corynebacterium stationis* KCCM13339P.

First, *Corynebacterium stationis* KCCM13339P was primarily cultured in 10 mL of 2YT medium (containing 16 g/l of tryptone, 10 g/l of yeast extract, and 5 g/l of sodium chloride) supplemented with 2% glucose, thus preparing a seed culture. Isonicotinic acid hydrazine at a concentration of 1 mg/ml and 2.5% glycine were added to 100 ml of 2YT medium free of glucose. Next, the seed culture was inoculated into the 2YT medium to reach an OD₆₁₀ value of 0.3, and then cultured at 30° C. and 180 rpm for 5 to 8 hours so that the OD₆₁₀ value reached 0.6 to 0.7. The culture was kept on ice for 30 minutes, and then centrifuged at 3,500 rpm at 4° C. for 10 minutes. Thereafter, the supernatant was discarded and the precipitated *Corynebacterium stationis* KCCM13339P was washed 4 times with a 10% glycerol solution and finally re-suspended in 0.5 ml of a 10% glycerol solution, thereby preparing competent cells. Electroporation was performed using a Bio-Rad electroporator. The prepared competent cells and the constructed pK_PC vector were placed in an electroporation cuvette (0.2 mm), and then subjected to electroporation under conditions of 2.5 kV, 200Ω and 12.5 μF. Immediately after completion of the electroporation, 1 ml of regeneration (RG) medium (containing 18.5 g/l brain heart infusion and 0.5 M sorbitol) was added to the cells which were then heat-treated at 46° C. for 6 minutes. Next, the cells were cooled at room temperature, transferred into a 15-ml cap tube, incubated at 30° C. for 2 hours, and plated on a selection medium (containing 5 g/l tryptone, 5 g/l NaCl, 2.5 g/l yeast extract, 18.5 g/l brain heart infusion powder, 15 g/l agar, 91 g/l sorbitol, and 20 μg/l kanamycin). The cells were cultured at 30° C. for 72 hours, and the generated colonies were cultured in medium until the stationary phase to induce secondary recombination. Then, the cells were diluted to 10⁻⁵ to 10⁻⁷, and plated on an antibiotic-free plate medium (containing 10% sucrose), and a strain having no kanamycin resistance and grown on the medium containing 10% sucrose was selected and named IPC-1.

Experimental Example 1. Evaluation of 5'-Inosinic Acid Productivity of Strain Expressing Phosphoenolpyruvate Carboxylase Variant

5'-Inosinic acid productivity was compared between the parent strain KCCM13339P and the mutant strain IPC-1 into which the phosphoenolpyruvate carboxylase variant has been introduced.

Each strain (parent strain or mutant strain) was inoculated at 1% by volume into a 100-mL flask containing 10 mL of the medium for 5'-inosinic acid production shown in Table 2 below, and cultured with shaking at 200 rpm at 34° C. for 45 hours. After completion of the culturing, the concentra-

tion of 5'-inosinic acid in the medium was measured using HPLC (Agilent), and the results are shown in Table 3 below.

TABLE 2

Component	Content
Glucose	70 g/L
(NH ₄) ₂ SO ₄	2 g/L
MgSO ₄	1 g/L
Urea	2 g/L
Yeast extract	20 g/L
KH ₂ PO ₄	2 g/L
FeSO ₄	10 mg/L
MnSO ₄	10 mg/L
Thiamine_HCl	5 mg/L
biotin	20 μg/L
Cystein	20 mg/L
Bata-alanine	20 mg/L
Adenine	30 mg/L

TABLE 3

Strain	5'-inosinic acid production (g/L)
KCCM13339P	20.0
IPC-1	23.2

As shown in Table 3 above, it was confirmed that the amount of 5'-inosinic acid produced by the mutant strain into which the phosphoenolpyruvate carboxylase variant has been introduced was increased by about 16% compared to that produced by the parent strain, due to substitution of the amino acids at positions 472 and 673 with other amino acids. These results suggest that introduction of point mutations into phosphoenolpyruvate carboxylase provides a significant effect on 5'-inosinic acid productivity.

So far, the present invention has been described with reference to the preferred embodiments. Those of ordinary skill in the art to which the present invention pertains will appreciate that the present invention may be embodied modified in forms without departing from the essential characteristics of the present invention. Therefore, the disclosed embodiments should be considered from an illustrative point of view, not from a restrictive point of view. The scope of the present invention is defined by the claims rather than the foregoing description, and all differences within the scope equivalent thereto should be construed as being included in the present invention

[Accession Number]

Depository Authority: Korean Culture Center of Microorganisms (KCCM)

Accession Number: KCCM13339P

Deposit Date: Mar. 29, 2023

SEQUENCE LISTING

Sequence total quantity: 7
 SEQ ID NO: 1 moltype = DNA length = 2805
 FEATURE Location/Qualifiers
 source 1..2805
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 1
 atgttcaata gcactgacgt ttcgctggaa aggaacccca ctgcaatgag tgaacaggtg 60
 cgcatgata tccgactgct aggacgcac ctcggccgag taattgccga gcaagaaggc 120

-continued

```

gaagatgtat ttgagttggt ggaagcact cgcgcttg ccttcggggg ggcgcgcggt 180
gaggaaaaaa ctgaggcttt gctggcgacc ttccggacta tcgatgaaaa ccgcataaac 240
ctcgttgcgc gttctttcag ccactttgcg ctgatggcga atatcgaga ggacctcgac 300
gatgaaacgg cgttgacagc ccgcgaggat gctggtgcac aggcctccga tgettccctg 360
cacggagttt tgggcaagct caaggcggcg aaagacatca aaacaagtga tgcacccgt 420
atcttgtagt ccgcgcaagt ctccccgttt ttaccgcac accccacgga gacacggcgt 480
cgtacggtat ttgatgtcca ggagcgcatc gtggcgctat tgcgtgagcg ccacagcatc 540
ttggcccaac cgcaaacagc aaggcttgaa gcccgctttg ctgagattga acgcgaggct 600
cacctgcgca tgaccatcct gtggcaaac ccggttaatt gcattgcacg gccacagatc 660
gaagatgaag taaacgtagg tctgcgctac ttcaagcgca gcctcttaga gcaggttaccg 720
gcgattaaac ggcacaccat cgcgggctt cgcgatgtct tcggcgcatc agtgcccaac 780
cggcaagtgt tgccgacagg gtcattggatc ggccgcgacc acgatggcaa tccttatgtc 840
accggcgata ctttgaccta cgcgacgcca caggccgagg ataccatcct ggaatactac 900
gttgatgagc ctcgagcact agaaaaggaa ctgagcttat cagaccgcta ttcagaaagt 960
tcgcggaat tacaagaact ggcacgcgcg gggagaaaac atattccaag tcgcgtcgat 1020
gagccttacc gcagggcaat ccacggagtg caccgacgca tgatagctac ccgcgcagcc 1080
gttgatgagc aagccgagc tgagcttggt gaattttcac cttatgcctc ccgcgaccag 1140
tttcgcacgg atttgacagt tattgaccgg tctttgcggc aatttaata tgcgatcatc 1200
gccgacgata ggctgtgcg aattctttct gccgtggata cttttggctt ccacctcaat 1260
gctttggatc tgcgtcagaa ctcggagctt ttccgagccg tcttggaaga acttttcgct 1320
gccgcggcgt ttactgcacc agatgccgga tacgccagcc tggatgagag cgcgaagcgt 1380
gaactactag tagctgagct cacctctgct cgtcactgg tcttcccggt gtctcaagac 1440
ttcagtgcag acacggaacg cgaactaggg atcttccggg cagcagccca ggccgtcgag 1500
ttactggggc caggtgttgt ccacactgc attgtttcaa tgaccggtag agtcagcgat 1560
attttggaac cgatggtact gctcaaagaa gtaggcatca taagctttaa ccacgcccag 1620
caacgcctcg ttggttcagt ggacatcgca ccgctttttg aaaccattga agacttaaa 1680
gctggcgcgag agatccttca agaactatgg gaagttgacc tttaccgagg atacttgccg 1740
gggcgcgaag agctccaaga agtcgtgttg gggatttccg actccaataa agacggcgagg 1800
tattgtctg cgaattgggc ttatatgat gcgcagctag ccacgtcaa ggctgcgat 1860
acgcacggca tcgggttgcg cttttcacat ggccgcggcg gtgctgtggg tcgcggtggc 1920
gggcgcagct acgatgccat cttggcccag ccgaagggtg cggctccggg cagcgtgcgt 1980
attaccgagc aaggtgaggt catctctgcc cgtacgaca cagcagtagc cgcgcgccg 2040
cacctggaag cttttgtcgc gggcacgcta gaagcctctc tactggatac tgaagagcta 2100
aaagagcctg aacgcgccta tgccatcatg cgggatatcg cctcgctggc cggtagagaa 2160
tatgcgcaat tgggtcgaga tgactccggg ttatcgact acttcacgca atccacgcca 2220
ttgcacgaaa ttggcgacct caatatggga tcacgtccaa ctgctcgcaa gcagaccgaa 2280
tccatcgggt acctgccgcg gattccatgg gtccgtgcat ggtcgcagtc gcgcgttaac 2340
ctgcgggctt ggttcggagt gggcacaggc atcatgcgct gggctggcga agacgagacc 2400
cgctggcaag atttgcccaa gctttatcag gcatggcgtt tctttcgctt cgtgctcgac 2460
aacatggcgc aggtcattgg gaaagcgtcg atgaatctcg caaagatcta ctccacgctt 2520
gtcgatgacc ccacgggtct ccagcgcgtg tttaacacca tctggacga atacgaatta 2580
accaaagaag tattcttccg catcactggt catgattcgc tcatggctgg taatgagcgt 2640
ttggagcgtt cagtacagcg acgtatcca tacttactgc cgctcaacgc catccagatt 2700
gagttgctgc gccgctaccg tgcggcgag gatacattct tggctcttaa gaccatccag 2760
gtgaccatga acggacttgc gacaggacta cgcacctcag gctag 2805

```

```

SEQ ID NO: 2      moltype = AA length = 934
FEATURE          Location/Qualifiers
source           1..934
                 mol_type = protein
                 organism = synthetic construct

```

```

SEQUENCE: 2
MFNSTDVSLE RNPTAMSEQV RDDIRLLGRI LGRVIAEQEG EDVFELVEST RRLAPGVARG 60
EEKAEALLAT FRITDENRIN LVARSFSHFA LMANIAEDLD DETALAAED AGAQAPDASL 120
HGVLGKLLKAA KDIKTSVTR ILDAAQVSPV FTAHPTEERR RTVFDVQERI VALLRERHSI 180
LAQPQTARLE ARLAEIEREA HLRMTILWQT ALIRIARQPI EDEVNVGLRY FKRSLLQVQP 240
AINRDTIAGL RDVFGASVPN RQVVRTGSKI GGDHNGNPNV TGDTLTYATR QAADTILEY 300
VDELALEKE LSLSDRYSES SAELQELATR GRNDIPSRVD EPYRRAIHGV HGRMIATRAA 360
VAGEAAAESG EFSPYASPTQ FRDQLQVIDR SLRQFNDAII ADDRLLRILS AVDTFGFHLN 420
ALDLRQNSSE FEAVLDLFLA AAGVTAPDAG YASLDESAGR ELLVAELTS RSLVFPWSQD 480
FSADTERELG IFRAAAQAVE LLGPGVPHC IVSMTGTVSD ILEPMVLLKE VGIISFNPAQ 540
QLVGSVDIA PLFETIEDLK AGAEILQELW EVDLYRGYLR GREDEVQEVVL GYSDSNKDDG 600
YLSANWALYD AQLAIVKACD THGIGLRFSH GRGGAVGRGG GPTYDAILAQ PEGAVRGSVR 660
ITEQGEVISA RYDTASTARR HLEAFVAGTL EASLLDTEKL KEPERAYGIM RDIASLAGEK 720
YAQLVRDDSG FIDYFTQSTP LHEIGDLNMG SRPTARKQTE SIGDLRAIPW VLSWSQSRVN 780
LPGWFGVGTG IMRWAGEDET RWQDLRLKLYQ AWPFFRSVLD NMAQVMGKAS MNLAKIYSTL 840
VDDPTVSQRV PNTILDEYEL TKEVFFRITG HDLSLMAGNER LERSVQRYP YLLPLNAIQI 900
ELLRRYRAGE DFLVSKTIQ VTMNGLATGL RTSG 934

```

```

SEQ ID NO: 3      moltype = DNA length = 2805
FEATURE          Location/Qualifiers
source           1..2805
                 mol_type = genomic DNA
                 organism = Corynebacterium stationis

```

```

SEQUENCE: 3
atgttcaata gcactgacgt ttcgctggaa aggaacccca ctgcaatgag tgaacagggt 60
cgcgatgata tccgactgct aggacgcac ctcggccgag taattgccga gcaagaaggc 120
gaagatgtat ttgagttggt ggaagcact cgcgcttg ccttcggggg ggcgcgcggt 180
gaggaaaaaa ctgaggcttt gctggcgacc ttccggacta tcgatgaaaa ccgcataaac 240

```


-continued

```

ctcgttgccg gttctttcag ccactttgcg ctgatggcga atategcaga ggacctcgac 300
gatgaaacgg cgcttgacgc ccgcgaggat gctgggtgcac aggcctccga tgcttccctg 360
cacggagttt tgggcaagct caaggcgggc aaagacatca aaacaagtga tgtcaccctg 420
atcttgatg cgcgcgaagt ctccccgtt ttaccgcac accccacgga gacacggcgt 480
cgtacgggat ttgatgtcca ggagcgcac gtggcgctat tgcgtgagcg ccacagcatc 540
ttggcccaac cgaacaacgc aaggtctgaa gcccgctctg ctgagattga acgcgaggct 600
cacctgcgca tgaccatctc gtggcaaac gcgttaattc gcatgacac gccacagatc 660
gaagatgaag taaacgtagg tctgcgtac ttcaagcgca gcctcttaga gcaggtagccg 720
gcgattaacc gcgacacccat cgcggggtct cgcgatgtct tcggcgcatc agtgcccaac 780
cggcaagtgt tgcgcacagg gtcattggatc ggccggcgacc acgatggcaa tccttatgtc 840
accggcgata ctttgaccta cgcgacgcga caggcgcggg ataccatcct ggaatactac 900
gttgatgagc tcgcagcact agaaaaggaa ctgagcttat cagaccgcta ttcagaaagt 960
tcgcgggaat tacaagaact ggccacgcgc gggagaaacg atattccaag tcgcgtcgat 1020
gagccttacc gcagggaact ccacggagtg caccgacgca tgatagctac ccgcgacgcc 1080
gttgaggag aagccgcgc tgagctctgg gaattttcac cttatgcctc ccgcagccag 1140
tttcgcacgg atttgcagg tattgaccgg tctttgcggc aatttaatga tgcgatcacc 1200
gcccagcata ggttgcgtgc aattctttct gcctgggata cttttggctt ccacctcaat 1260
gctttggatc tgcgtcagaa ctccgagctc ttccgagccg tcttggacga acttttcgct 1320
gccgcggggc ttaactgcac agatgccgga taacgcagcc tggatgagag gcccaagcgt 1380
gaactactag tagctgacct cactctgct cgcctcactg tcttcccggt gtctcaagac 1440
ttcagtgacg acacgggaac cgaactaggg atcttccggg cagcagccca gccgctcgag 1500
ttaactgggc caggtgttgt cccacactgc attgtttcaa tgaccggtac agtcagcgat 1560
attttggaac cgatggtact gctcaaaaga gttagcatca taagctttaa cccagccag 1620
caacgcctcg ttggttcagt ggacatcgca ccgctttttg aaaccattga agacttaag 1680
gctggcgccg agatccttca agaactatgg gaagttagc tttaccgagg atacttgcgc 1740
ggggcgcaag acgtccaaga agtcgtgttg gggatttccg actccaataa agacggcgccg 1800
tatttgcctc cgaattgggc tttatatgat gcgcagctag ccacgtcaa gccctgcgat 1860
acgcacggca tcgggttgcg cttttcacat ggccgcggcg gtgctgtggg tcgcggtggc 1920
ggggcgacgt acgatgacct cttggcccag cccgaagggt cggtccgggg cagcgtgcgt 1980
attaccgagc aaggtgaggt catctctgcc cgtacgcgca cagcagatc cgcgcgccgc 2040
cacctggaag cttttgtcgc gggcacgcta gaagcctctc tactggatac tgaaggcta 2100
aaagagcctg aacgcgcata tggcatcatg cgggatatcg cctcgtggc cggtgagaaa 2160
tatgcgcaat tgggtgcgca tgactccggg tttatcgact acttcaacga atccacgcca 2220
ttgcacgaaa ttggcgacct caatatggga tcacgtccaa ctgctcgcaa gcagaccgaa 2280
tccatcggtg acttgcgcgc gattccatgg gtctgtcat ggtcgcagtc gcgcgttaac 2340
ctgcgcggct ggttcggagt gggcacaggg atcatgcgct gggctggcga agacgagacc 2400
cgctggcaag atttgcgcaa gctttatcag gcatggcgt tctttcgctt cgtgctgac 2460
aacatggcgc aggtcatggg gaaagcgtcg atgaatctcg caaagatcta ctccacgctt 2520
gtcgatgacc ccacgggtct ccagcgcgtg tttaacacca tcttggacga atacgaatta 2580
accaaaagag tattcttcgc catcactggt catgattcgc tcatggctgg taatgagcgt 2640
ttggagcgtt cagtagacgc acgctatcca tacttactgc cgctcaacgc catccagatt 2700
gagttgctgc gccgctaccg tgcgggcgag gatacattct tggtctctaa gaccatccag 2760
gtgaccatga acggacttgc gacaggacta cgcacctcag gctag 2805

```

```

SEQ ID NO: 4      moltype = AA length = 934
FEATURE          Location/Qualifiers
source           1..934
                 mol_type = protein
                 organism = Corynebacterium stationis

```

```

SEQUENCE: 4
MFNSTDVSLE RNPAMSEQV RDDIRLLGRI LGRVIAEQEG EDVFELVEST RRLAFGVARG 60
EEKAEALLAT FRITDENRIN LVARSFSHFA LMANIAEDLD DETALAARED AGAQAPDASL 120
HGVLGKLLAA KDIKTSVTR ILDAAQVSPV FTAHPTETRR RTVFDVQERI VALLRERHSI 180
LAQPQTARLE ARLAEIEREA HLRMTILWQT ALIRIARPOI EDEVNVLGRY FKRSLLSEQV 240
AINRDTIAGL RDVFGASVPN RQVVRTGSWI GGDHGNPYV TGDLTLYATR QAADTILEYV 300
VDELAALKE LSLSDRYSES SAELOELATR GRNDIPSRVD EPYRRAIHGV HGRMIATRAA 360
VAGEAAAESG EFSFYASPTQ FRITDLQVIDR SLRQFNDAII ADDRLLRILS AVDTFGFHLN 420
ALDLRQNSSE FEAVLDELFA AAGVTAPDAG YASLDESAGR ELLVAELTSA RPLVFPWSQD 480
FSADTERELG IFRAAAQAVE LLGPGVVPHC IVSMTGTVSD ILEPMVLLKE VGIISFNPAQ 540
QRLVGSVDIA PLFETIEDLK AGAEILQELW EVDLYRGYLR GREDEVQEVV L GYSDSNKDG 600
YLSANWALYD AQLAIVKACD THGIGLRFSH GRGGAVGRGG GPTYDAILAQ PEGAVRGSVR 660
ITEQGEVISA RYGTASTARR HLEAFVAGTL EASLLDTEKL KEPERAYGIM RDIASLAGEK 720
YAQLVRDDSG FIDYFTQSTP LHEIGDLNMG SRPTARKQTE SIGDLRAIPW VLSWSQSRVN 780
LPQWFGVGTG IMRWAGEDET RWQDLRKLQY AWPFFRSVLD NMAQVMGKAS MNLAKIYSTL 840
VDDPTVSQRV FNTILDEYEL TKEVFFRITG HDLSLMAGNER LERSVQRRYP YLLPLNAIQI 900
ELLRRYRAGE DTFLVSKTIQ VTMNGLATGL RTSG 934

```

```

SEQ ID NO: 5      moltype = DNA length = 5719
FEATURE          Location/Qualifiers
source           1..5719
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 5
tgccgcaagc actcaggcgc caaggcgtgc taaaggaagc ggaacacgta gaaagccagt 60
ccgcagaaac ggtgctgacc ccggatgaat gtcagctact gggctatctg gacaagggaa 120
aacgcaagcg caaagagaaa gcaggtagct tgcagtgggc ttacatggcg atagctagac 180
tgggcggttt tatggacagc aagcgaaccg gaattgccag ctggggcgcc ctctggttaag 240
gttgggaagc cctgcaagc aaactggatg gcttcttcgc cgccaaggat ctgatggcgc 300
aggggatcaa gatctgatca agagacagga tgaggatcgt ttcgcatgat tgaacaagat 360

```

-continued

ggattgcacg	caggttctcc	ggcgcgttgg	gtggagagggc	tattcggtcta	tgactgggca	420
caacagacaa	tcggctgctc	tgatgcgcgc	gtgttccggc	tgtcagcgca	ggggcgcccg	480
gtttctttttg	tcaagaccga	cctgtccgggt	gccctgaatg	aactccaaga	cgaggcagcg	540
cggtatcgt	ggctggccac	gacgggcgtt	ccttgccgag	ctgtgctcga	cgttgteact	600
gaagcgggaa	gggactggct	gctattgggc	gaagtgccgg	ggcaggatct	cctgtcatct	660
caccttgctc	ctgcgcagaa	agtatccatc	atggctgatg	caatgcggcg	gctgcatacg	720
cttgatccgg	ctacctgccc	attcgaccac	caagcgaaac	atcgcatcga	gcgagcacgt	780
actcggatgg	aagccggtct	tgatgatcag	gatgatctgg	acgaagagca	tcaggggctc	840
gcgcgagccg	aactgttcgc	caggctcaag	gcgcgggatgc	ccgacggcga	ggatctcgtc	900
gtgacccatg	gcgatgacct	cctggccgaat	atcatggtgg	aaaatggccg	cttttctgga	960
ttcatcgact	gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttggctacc	1020
cgtgatattg	ctgaagagct	tggcggcgaa	tgggctgacc	gcttcctcgt	gctttacggt	1080
atcgccgctc	ccgattcgca	gcgcacgcgc	ttctatcgcc	ttcttgacga	gttctcttga	1140
gcgggactct	ggggttcgct	agaggatcga	tcctttttta	cccatcacat	atactggcgc	1200
ttcactatta	tttagtgaaa	tgagatatta	tgatattttc	tgaattgtga	ttaaaaaggc	1260
aactttatgc	ccatgcaaca	gaaactataa	aaaatacaga	gaatgaaaaa	aaacagatag	1320
atttttttagt	ctctttaggc	ctagtctgc	aaatcctttt	atgattttct	atcaaacaaa	1380
agaggaaaaa	agaccagttg	caatccaaac	gagagtctaa	tagaatgagg	tcgaaaaagta	1440
aatcgccgcg	gtttgttact	gataaagcag	gcaagaccta	aaatgtgtaa	agggcaaaagt	1500
gtataactttg	gcgtcacccc	ttacatat	taggtctttt	tttattgtgc	gttaactaact	1560
tgccatcttc	aaacaggagg	gctggaagaa	gcgacccgct	aacacagtag	ataaaaaagg	1620
agacatgaac	gatgaacatc	aaaaagtgtg	caaaacaagc	aacagtatta	acctttacta	1680
ccgcactgct	ggcaggaggc	gcaactcaag	cgtttgcgaa	agaaacgaac	caaaagccat	1740
ataagggaaa	atacggcatt	ttccatatta	cacgccatga	tatgctgcga	atccctgaac	1800
agcaaaaaaa	tgaaaaaatat	caagtttctg	aatttgattc	gtccacaatt	aaaaatatct	1860
cttctgcaaa	aggcctggac	gtttgggaca	gctggccatt	acaaaacgct	gacggcactg	1920
tcgcaaaacta	tcacggctac	cacatcgtct	ttgcattagc	cggagatcct	aaaaatgcgg	1980
atgacacatc	gatttacaat	ttctatcaaa	aagtcggcga	aacttctatt	gacagctgga	2040
aaaacgctgg	ccgcgtcttt	aaagacagcg	acaaattcga	tgcaaatgat	tctatcctaa	2100
aagacccaaac	acaagaatgg	tcaggttcag	ccacatttac	atctgacgga	aaaatccggt	2160
tattctcac	tgatttctcc	ggttaaacatt	acggcaaaac	aacactgaca	actgcacaag	2220
ttaacgtatc	agcatcagac	agctctttga	acatcaacgg	tgtagaggat	tataaatcaa	2280
cttttgacgg	tgacggaaaa	acgtatcaaa	atgtacagca	gttcatcgat	gaaggcaact	2340
acagctcagg	cgacaaccat	acgtgagag	atcctcacta	cgtagaagat	aaaggccaca	2400
aatacttagt	atttgaagca	aacactggaa	ctgaagatgg	ctaccaaggc	gaagaatctt	2460
tatttaacaa	agcatactat	ggcaaaagca	catcattctt	ccgtcaagaa	agtcaaaaac	2520
ttctgcacaa	cgataaaaaa	cgcacggctg	agttagcaaa	cggcgctctc	ggtatgattg	2580
agctaaacga	tgattacaca	ctgaaaaaaa	tgatgaaacc	gctgattgca	tctaacaacag	2640
taacagatga	aattgaacgc	gcgaacgtct	ttaaaaatga	cggcaaatgg	tacctgttca	2700
ctgactcccg	cggatcaaaa	attgacgattg	acggcattac	gtctaacgat	atttcatatg	2760
ttgggtatgt	ttctaattct	ttaaactggc	catcaaacgc	gctgaacaaa	actggccttg	2820
tggttaaaat	ggatcttgat	cctaacgatg	taacctttac	ttactcacac	ttcgctgtac	2880
ctcaagcgaa	aggaaaacaat	gtcgtgatta	caagctatat	gacaaacaga	ggattctacg	2940
cagacaaaac	atcaacgttt	gcgcgcgagct	tcctgctgaa	catcaaaagg	aagaaaaacat	3000
ctgttgtcaa	agacagcatc	cctgaacaag	gacaattaac	agttaacaaa	taaaaacgca	3060
aaagaaaaatg	ccgatgggta	ccgagcgaaa	tgaccgacca	agcgacgccc	aacctgccat	3120
cacgagattt	cgattccacc	gcgcgcctct	atgaaagggt	gggcttcgga	atcgttttcc	3180
gggacgcctc	cgcggaacgtg	ctcatagctc	acgacgcccg	tgattttgta	gccctggccg	3240
acggccagca	ggttagccga	caggctcatg	ccggccgccc	ccgccttttc	ctcaatcgct	3300
cttctgtctg	ctggaaggca	gtacaccttg	ataggtgggc	tgcccttctc	ggttggtctg	3360
gtttcatcat	ccatccgctt	gcctcatct	gttacgcggg	cggtagccgg	ccagcctcgc	3420
agagcaggat	tcctgttgag	caccgccagg	tgcaataaag	ggacagtga	gaaggaaacac	3480
ccgctcgcgg	gtgggctac	ttcacctatc	ctgcccggct	gacgcggttg	gatacaccaa	3540
ggaaagtcta	cacgaaccct	ttggcaaaat	cctgtatata	gtgcgaaaaa	ggatggatat	3600
accgaaaaaa	tcgctataat	gacccggaag	cagggttatg	cagcgaaaaa	gcgctgcttc	3660
cctgtgcttt	tgtggaatat	ctaccgactg	gaaaacaggca	aatgcaggaa	attactgaac	3720
tgaggggaca	ggcgagagac	gatgccaaaag	agctcctgaa	aatctcgata	actcaaaaaa	3780
tacgcccgtg	agtgatctta	ttctattatg	gtgaaagtgg	gaacctctta	cgtgccgatc	3840
aacgtctcat	tttcgcaaaa	agttggccca	gggcttcccg	gtatcaacag	ggacaccagg	3900
atttatttat	tctgcgaagt	gatcttccgt	cacaggattt	tattcgccgc	aaagtgcgtc	3960
gggtgatgct	gccaacttac	tgatttagtg	tatgatggtg	tttttgaggt	gctccagtgg	4020
cttctgtttc	tatcagctcc	tgaataatctc	gataaactcaa	aaaatacgcc	cggtagtgat	4080
cttatttcat	tatggtgaaa	gttggaacct	cttactgtcc	gatcaacgtc	tcattttctc	4140
caaaagtgtg	cccaggcctt	cccggtatca	acagggaacac	caggatttat	ttattctgcg	4200
aagtgatctt	ccgtcacagg	tatttattcg	gcgcaaaagt	cgtcggttga	tgctgccaac	4260
ttactgattt	agtgatgat	ggtgtttttg	aggtgctcca	gtggcttctg	ttctatcag	4320
ggctggatga	tcctccagcg	cggggatctc	atgctggagt	tcctcgccca	ccccaaaagg	4380
atctaggtga	agatcctttt	tgataatctc	atgacaaaaa	tccttaacg	tgagttttct	4440
ttccactgag	cgtcagaccc	cgtagaaaag	atcaaaaggat	cttcttgaga	tccttttttt	4500
ctgcgcgtaa	tctgctgctt	gcaaacaaaa	aaaccaccgc	taccagcggt	gggttgtttg	4560
ccggatcaag	agctaccaac	tccttttccg	aaggtaactg	gcttcagcag	agcgagata	4620
ccaaataactg	ttcttctagt	gtagccgtag	ttaggccacc	acttcaagaa	ctctgtagca	4680
ccgcctacat	acctcgctct	gctaactcctg	ttaccagtgg	ctgctgccag	tggcgataag	4740
tcgtgtctta	ccgggttgga	ctcaagacga	tagttaccgg	ataaggcgca	gcggtcgggc	4800
tgaacggggg	gttcgtgcac	acagcccgac	ttggagcgaa	cgacctacac	cgaactgaga	4860
tacctacagc	gtgagctatg	agaaagcgcc	acgcttcccg	aaggagagaa	ggcggacagg	4920
tatccggtaa	gcggcagggt	cggaaacagga	gagcgacaga	gggagcttcc	agggggaaac	4980
gcctgggtac	tttatagctc	tgctgggttt	cgccacctct	gaacttgagc	tcgatttttg	5040
tgatgctcgt	cagggggggc	gagcctatgg	aaaaacgcca	gcaacgcggc	ctttttacgg	5100

-continued

```

ttcttgacct tttgtggcc tttgtctac atgttcttct ctgcgttacc cctgattct 5160
gtggataaac gtattaccgc ctttgagtga gctgataccg ctgcgcgcag ccgaacgacc 5220
gagcgcagcg agtcagtgag cgaggaagcg gaagagcgcc caatacgcaa accgcctctc 5280
cccgcgcggt ggccgattca ttaatgcagc tggcagcaca gggttcccg ctggaaagcg 5340
ggcagtgagc gcaacgcaat taatgtgagt tagctcactc attaggcacc ccaggcttta 5400
cactttatgc ttccggctcg tatgttgtgt ggaattgtga gcggataaca atttcacaca 5460
ggaaacagct atgacatgat tacgccaagc ttgcatgcct gcaggctcgac tctagaggat 5520
ccccgggtac cgagctcgaa ttcaactggc gtcgttttac aacgtcgtga ctgggaaaac 5580
cctggcggtta cccaacttaa tcgccttgca gcacatcccc ctttcgcccag ctggcgtaat 5640
agcgaagagg cccgcaccga tcgccttcc caacagttgc gcagcctgaa tggcgaatgg 5700
cgataagcta gcttcacgc 5719

```

```

SEQ ID NO: 6      moltype = DNA length = 20
FEATURE          Location/Qualifiers
source           1..20
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 6
ctctgctcgc tcaactggtct 20

```

```

SEQ ID NO: 7      moltype = DNA length = 21
FEATURE          Location/Qualifiers
source           1..21
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 7
tactcgctgt gtcgtagcgg g 21

```

The invention claimed is:

1. A phosphoenolpyruvate carboxylase variant in which one or more of the amino acids at positions 472 and 673 in the amino acid sequence of SEQ ID NO: 4 is/are substituted with other amino acid(s).
2. The phosphoenolpyruvate carboxylase variant of claim 1, which consists of the amino acid sequence of SEQ ID NO: 2 in which proline at position 472 in the amino acid sequence of SEQ ID NO: 4 is substituted with serine and glycine at position 673 is substituted with aspartic acid.
3. A polynucleotide encoding the variant of claim 1.
4. A transformant host cell comprising the variant of claim 1 or the polynucleotide of claim 3.

5. The transformant host cell of claim 4, which is a *Corynebacterium* sp. microorganism.

6. The transformant host cell of claim 4, which has ability to produce 5'-inosinic acid.

7. A method for producing 5'-inosinic acid, comprising steps of: culturing the transformant host cell of claim 4 in a medium; and recovering 5'-inosinic acid from the transformant host cell or the medium in which the transformant host cell has been cultured.

8. A transformant host cell comprising the polynucleotide of claim 3.

* * * * *