

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12414551
Kind Code	B2
Date of Patent	September 16, 2025
Inventor(s)	Macdonald; Lynn et al.

Genetically modified major histocompatibility complex mice

Abstract

The invention provides genetically modified non-human animals that express chimeric human/non-human MHC I and MHC II polypeptides and/or human or humanized $\beta 2$ microglobulin polypeptide, as well as embryos, cells, and tissues comprising the same. Also provided are constructs for making said genetically modified animals and methods of making the same. Methods of using the genetically modified animals to study various aspects of human immune system are provided.

Inventors: Macdonald; Lynn (Harrison, NY), Murphy; Andrew J. (Croton-on-Hudson, NY), Voronina; Vera (North Bethesda, MD), Gurer; Cagan (Chappaqua, NY)

Applicant: Regeneron Pharmaceuticals, Inc. (Tarrytown, NY)

Family ID: 1000008821038

Assignee: Regeneron Pharmaceuticals, Inc. (Tarrytown, NY)

Appl. No.: 17/550945

Filed: December 14, 2021

Prior Publication Data

Document Identifier	Publication Date
US 20220201994 A1	Jun. 30, 2022

Related U.S. Application Data

continuation parent-doc US 16184473 20181108 US 11224208 child-doc US 17550945
continuation parent-doc US 14185316 20140220 US 10154658 20181218 child-doc US 16184473
us-provisional-application US 61767811 20130222

Publication Classification

Int. Cl.: A01K67/0278 (20240101); C07K14/74 (20060101); C07K14/705 (20060101); C07K14/73 (20060101); C12N15/85 (20060101)

U.S. Cl.:

CPC A01K67/0278 (20130101); C07K14/70539 (20130101); A01K2207/15 (20130101); A01K2217/072 (20130101); A01K2267/03 (20130101); C07K14/70514 (20130101); C07K14/70517 (20130101); C12N15/8509 (20130101)

Field of Classification Search

CPC: A01K (67/0278)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5413923	12/1994	Kucherlapati et al.	N/A	N/A
5416260	12/1994	Koller et al.	N/A	N/A
5574205	12/1995	Kucherlapati et al.	N/A	N/A
5644065	12/1996	Benoist et al.	N/A	N/A
5859312	12/1998	Littman et al.	N/A	N/A
5942435	12/1998	Wheeler	N/A	N/A
5965787	12/1998	Luthra et al.	N/A	N/A
6002066	12/1998	Leung et al.	N/A	N/A
6139835	12/1999	Kucherlapati et al.	N/A	N/A
6150584	12/1999	Kucherlapati et al.	N/A	N/A
6270772	12/2000	Burrows et al.	N/A	N/A
6372955	12/2001	Karlsson et al.	N/A	N/A
6514752	12/2002	Kucherlapati et al.	N/A	N/A
6586251	12/2002	Economides et al.	N/A	N/A
6596541	12/2002	Murphy et al.	N/A	N/A
6815171	12/2003	Burrows et al.	N/A	N/A
7105348	12/2005	Murphy et al.	N/A	N/A
7265218	12/2006	Burrows et al.	N/A	N/A
7294754	12/2006	Poueymirou et al.	N/A	N/A
7339089	12/2007	Gotch	N/A	N/A
7663017	12/2009	Lone et al.	N/A	N/A
7745690	12/2009	Kanazawa et al.	N/A	N/A
8178653	12/2011	Tureci et al.	N/A	N/A
8847005	12/2013	Macdonald et al.	N/A	N/A
9043996	12/2014	Macdonald et al.	N/A	N/A
9585373	12/2016	Macdonald et al.	N/A	N/A
9591835	12/2016	Macdonald et al.	N/A	N/A
9615550	12/2016	Macdonald et al.	N/A	N/A
9657311	12/2016	Bauche et al.	N/A	N/A
10154658	12/2017	Macdonald et al.	N/A	N/A

10278373	12/2018	Harada et al.	N/A	N/A
2002/0164721	12/2001	Firat et al.	N/A	N/A
2003/0093818	12/2002	Belmont et al.	N/A	N/A
2003/0124524	12/2002	Kornman et al.	N/A	N/A
2004/0137537	12/2003	Montero-Julian et al.	N/A	N/A
2005/0050580	12/2004	Gotch et al.	N/A	N/A
2005/0066375	12/2004	Thiam et al.	N/A	N/A
2005/0114910	12/2004	Lone et al.	N/A	N/A
2006/0107339	12/2005	Gotch et al.	N/A	N/A
2007/0209083	12/2006	Thiam et al.	N/A	N/A
2008/0152131	12/2007	Sendo et al.	N/A	N/A
2008/0181899	12/2007	Papadopoulos et al.	N/A	N/A
2008/0219947	12/2007	Linette et al.	N/A	N/A
2009/0328240	12/2008	Sing et al.	N/A	N/A
2010/0011450	12/2009	Garcia et al.	N/A	N/A
2010/0111993	12/2009	Tureci et al.	N/A	N/A
2010/0138938	12/2009	Garcia et al.	N/A	N/A
2011/0067121	12/2010	Lone et al.	N/A	N/A
2013/0111617	12/2012	Macdonald et al.	N/A	N/A
2013/0117873	12/2012	Wang et al.	N/A	N/A
2013/0185819	12/2012	Macdonald et al.	N/A	N/A
2014/0134662	12/2013	Flavell et al.	N/A	N/A
2014/0245466	12/2013	Macdonald et al.	N/A	N/A
2015/0040253	12/2014	Macdonald et al.	N/A	N/A
2015/0245598	12/2014	Macdonald et al.	N/A	N/A
2015/0342163	12/2014	Voronina et al.	N/A	N/A
2017/0142944	12/2016	Macdonald et al.	N/A	N/A
2017/0164590	12/2016	Macdonald et al.	N/A	N/A
2017/0273286	12/2016	Macdonald et al.	N/A	N/A
2018/0139940	12/2017	Macdonald et al.	N/A	N/A
2018/0288986	12/2017	Macdonald et al.	N/A	N/A
2019/0223418	12/2018	Murphy et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
0437576	12/2001	EP	N/A
1878342	12/2007	EP	N/A
1878798	12/2007	EP	N/A
0950707	12/2008	EP	N/A
1017721	12/2008	EP	N/A
1409646	12/2011	EP	N/A
1991001140	12/1990	WO	N/A
1992011753	12/1991	WO	N/A
1993005817	12/1992	WO	N/A
1995003331	12/1994	WO	N/A
1997032603	12/1996	WO	N/A
1998024893	12/1997	WO	N/A
2001027291	12/2000	WO	N/A
2002059263	12/2001	WO	N/A
2003006639	12/2002	WO	N/A

2004/083244	12/2003	WO	N/A
2005004592	12/2004	WO	N/A
2007131092	12/2006	WO	N/A
2008010099	12/2007	WO	N/A
2008010100	12/2007	WO	N/A
2009114400	12/2008	WO	N/A
2011004192	12/2010	WO	N/A
2011123708	12/2010	WO	N/A
2012007951	12/2011	WO	N/A
2012039779	12/2011	WO	N/A
2012018610	12/2011	WO	N/A
2013063340	12/2012	WO	N/A
2013063346	12/2012	WO	N/A
2014130671	12/2013	WO	N/A
2014164638	12/2013	WO	N/A
2014164640	12/2013	WO	N/A
2016164492	12/2015	WO	N/A
2017210586	12/2016	WO	N/A

OTHER PUBLICATIONS

Abarrategui and Krangel (2006) "Regulation of T cell receptor alpha gene recombination by transcription," *Nat. Immunol.*, 7:1109-1115 and Corrigendum. cited by applicant

Anderson (2008) "Post-transcriptional control of cytokine production," *Nature Immunology*, 9(4):353-359. cited by applicant

Allen et al. (1986) "B2-Microglobulin is not required for Cell surface expression of the murine class I histocompatibility antigen H-2Db or of a truncated H-2Db," *PNAS*, 83:7447-7451. cited by applicant

Altmann et al. (1995) "The T Cell Response to HLA-DR Transgenic Mice to Human Myelin Basic Protein and other Antigens in the Presence and Absence of Human CD4," *J. Exp. Med.*, 181:867-875. cited by applicant

Alvarez et al. (1995) "V(D)J Recombination and Allelic Exclusion of a TCR B-Chain Minilocus Occurs in the Absence of a Functional Promoter," *J. Immunol.*, 155:1191-1202. cited by applicant

Arnold and Hammerling (1991) "MHC Class-1 Transgenic Mice," *Annu. Rev. Immunol.*, 9:297-322. cited by applicant

Auerbach et al. (2000) "Establishment and Chimera Analysis of 129/SvEV- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines," *BioTechniques*, 29:1024-1032. cited by applicant

Baker et al. (1996) "Adaptation of TCR Expression Vectors for the Construction of Mouse-Human Chimeric MBP-Specific TCR Transgenes," *J. Neurosci. Research*, 45:487-491. cited by applicant

Barber and Lechler (1991) "Interactions between the amino-terminal domains of MCH class II molecules have a profound effect on serologic and T cell recognition: an analysis using recombinant HLA-DR/H-2E molecules," *J. Immunol.*, 147:2346-2353. cited by applicant

Barthold (2004) "Genetically altered mice: phenotypes, no phenotypes, and Faux phenotypes," *Genetica*, 122:75-88. cited by applicant

Basha et al. (2008) "MHC Class I Endosomal and Lysosomal Trafficking Coincides with Exogenous Antigen Loading in Dendritic cells," *PLoS ONE*, 3:e3247, 11 pages. cited by applicant

Bassing et al. (2000) Recombination signal sequences restrict chromosomal V(D)J recombination beyond the 12/23 rule, *Nature*, 405:583-586. cited by applicant

Benmohamed et al. (2000) "Induction of CTL Response by a Minimal Epitope Vaccine in HLA-A*0201/DR1 Transgenic Mice: Dependence on HLA Class II Restricted TH Response," *Hum. Immunol.*, 61:764-779. cited by applicant

Bernabeu et al. (1984) "B2-Microglobulin from serum associates with MHC class I antigens on the surface of cultured cells," *Nature*, 308:642-645. cited by applicant

Beta-2 microglobulin https://en.wikipedia.org/w/index.php?title=Beta-2_microglobulin&oldid=452509865. cited by applicant

Betser-Cohen et al. (2010) "The Association of MHC Class I Proteins with the 2B4 Receptor Inhibits Self-Killing of Human NK cells," *J. Immunol.*, 184:2761-2768. cited by applicant

Bonnet, et al. (2009) "Molecular Genetics at the T-Cell Receptor β Locus: Insights into the Regulation of V(D)J Recombination," *V(D)J Recombination*, 650:116-132. cited by applicant

Bouffard et al. (1997) "A Physical Map of Human Chromosome 7: An Integrated YAC Contig Map with Average STS Spacing of 79 kb," *Genome Research*, 7:673-692. cited by applicant

Boyd and Wood (2009) "Variation in MHC expression between undifferentiated mouse ES cells and ES cell derived insulin-producing cell clusters," *Transplantation*, 87(9):1300-1304. cited by applicant

Brevini et al. (2010) "Embryonic Stem Cells in Domestic Animals; No shortcuts to pig embryonic stem cells," *Theriogenology*, 74:544-550. cited by applicant

Brouwers et al. (2015) "Unexpected Phenotypes in Mouse Models Carrying the Human Growth Hormone Minigene to Enhance Transgene Expression," *Journal of Steroids & Hormonal Science*, 6(2):1000.e115 (2 pages). cited by applicant

Buta et al. (2013) "Reconsidering pluripotency tests: Do we still need teratoma assays?," *Stem Cell Res.*, 11:552-562. cited by applicant

Cameron, E. (1997) "Recent Advances in Transgenic Technology," *Molecular Biotechnology*, 7:253-265. cited by applicant

Cao et al., (2009) "Isolation and Culture of Primary Bovine Embryonic Stem Cell Colonies by a Novel Method," *Journal of Experimental Zoology*, 311A:368-376. cited by applicant

Carpenter, et al. (2014) "Post-transcriptional regulation of gene expression in innate immunity," *Nature Reviews Immunology*, 14:361-376. cited by applicant

Carstea et al. (2009) "Germline competence of mouse ES and iPS Cell lines: Chimera technologies and genetic background," *World Journals of Stem Cells*, 1(1):22-29. cited by applicant

Chamberlain et al. (1988) "Tissue-specific and Cell surface expression of human major histocompatibility complex class I heavy (HLA-B7) and light (B2-microglobulin) chain genes in transgenic mice," *PNAS*, 86:7690-7694. cited by applicant

Cheuk et al. "Human MHC Class I Transgenic Mice Deficient for H2 Class I Expression Facilitate Identification and Characterization of New HLA Class I-Restricted Viral T Cell Epitopes," *Journal of Immunology*, 2002, 169:5571-5580. cited by applicant

Choi et al., (2011) "Expression of the metabotropic glutamate receptor 5 (mGluR5) induces melanoma in transgenic mice," *PNAS*, 108(37):1519-15224. cited by applicant

Chung et al. (1994) "Functional three-domain single-chain T-Cell receptors," *PNAS*, 91:12654-12658. cited by applicant

Clark et al. (1987) "Peptide and nucleotide sequences of rat CD4 (W3/25) antigen: evidence for derivation from a structure with four immunoglobulin-related domains," *PNAS*, 84(6):1649-1653. cited by applicant

Connolly et al. (1988) "The Lyt-2 Molecule Recognizes Residues in the Class I alpha3 Domain in Allogeneic Cytotoxic T Cell Responses," *J. Exp. Med.*, 168:325-341. cited by applicant

Cooper et al. (2007) "An Impaired Breeding Phenotype in Mice with a Genetic Deletion of Beta-2 Microglobulin and Diminished MHC Class I Expression: Role in Reproductive Fitness," *Biol. Reprod.*, 77:274-279. cited by applicant

Corbeil et al. (1996) "HIV-induced Apoptosis Requires the CD4 Receptor Cytoplasmic Tail and Is Accelerated by Interaction of CD4 with p56lck," *J. Exp. Med.*, 183:39-48. cited by applicant

Cosson and Bonifacio (1992) "Role of Transmembrane Domain Interactions in the Assembly of Class II MHC Molecules," *Science*, 258:659-662. cited by applicant

Crusio et al. (2004) "Flanking Gene and Genetic Background Problems in Genetically Manipulated Mice," *Biol. Psychiatry*, 56:381-385. cited by applicant

Danner et al. (2011) "Expression of HLA Class II Molecules in Humanized NOD.Rag1KO.IL2RgKO Mice is Critical for Development and Function of Human T and B cells," *PLoS ONE*, 6:e19826, 12 pages. cited by applicant

Database entry for NCB I Reference Sequence: NG 001333.2 (Sep. 19, 2006) "*Homo sapiens* T cell receptor beta locus (TRB) on chromosome 7". cited by applicant

De Bakker et al. (2006) "A high-resolution HLA and SNP haplotype map for disease association studies in the extended human MHC," *Nature Genet. Online Supplement*, 33 pages. cited by applicant

De Bakker et al. (2006) "A high-resolution HLA and SNP haplotype map for disease association studies in the extended human MHC," *Nature Genet.*, 38:1166-1172. cited by applicant

De Gassart et al. (2008) "MHC class II stabilization at the surface of human dendritic cells is the result of maturation-dependent MARCH I down-regulation," *PNAS*, 105:3491-3496. cited by applicant

Dennis (2002) "Welfare issues of genetically modified animals," *ILAR Journal*, 43(2):100-109. cited by applicant

Doetschman (2009) "Influence of Genetic Background on Genetically Engineered Mouse Phenotypes," *Methods Mol. Biol.*, 530:423-433. cited by applicant

Dolan et al. (2004) "Invariant Chain and the MHC Class II Cytoplasmic Domains Regulate Localization of MHC Class II Molecules to Lipid Rafts in Tumor Cell-Based Vaccines," *J. Immunol.*, 172:907-914. cited by applicant

Due (1989) "Self-Recognition by T Cell," *J. Exp. Med.* 170:59-71. cited by applicant

El Fakhry et al. (2004) "Delineation of the HLA-DR Region and the Residues Involved in the Association with the Cytoskeleton," *J. Biol. Chem.*, 279:18472-18480. cited by applicant

Ellmeier et al. (1998) "Multiple developmental stage-specific enhancers regulate CD8 expression in developing thymocytes and in thymus-independent T cells," *Immunity*, 9(4):485-496. cited by applicant

Epstein et al. "Expression and function of HLA-A2.1 in transgenic mice," *Eur. J. Immunol.*, 1989, 19:1575-1583. cited by applicant

Festing et al. (1999) "Revised nomenclature for strain 129 mice," *Mamm. Genome*, 10:836. cited by applicant

Firat et al. (2002) "Comparative analysis of the CO8+ T Cell repertoires of H-2 class I wild-type/HLA-2.1 and H-2 class I knockout/HLA-A2.1 transgenic mice," *Internat. Immunol.*, 14:925-934. cited by applicant

Fleischer et al. (1996) Reactivity of Mouse T-Cell Hybridomas Expressing Human V β Gene Segments with Staphylococcal and Streptococcal Superantigens, *Infection and Immunity*, 64(3):997-994. cited by applicant

Fooksman et al. (2009) "Cutting Edge: Phospholidylinositol 4, 5-Bisphosphate Concentration at the APC Side of the Immunological Synapse Is Required for Effector T Cell Function," *J. Immunol.*, 182:5179-5182. cited by applicant

Friese et al., (2008) "Opposing effects of HLA class I molecules in tuning autoreactive CD8+ T cells in multiple sclerosis," *Nature Med.*, 14(11): 1227-1235. cited by applicant

Fugger et al. (1994) "Expression of HLA-DR4 and human CD4 transgenes in mice determines the variable region beta-chain T-Cell repertoire and mediates an HLA-DR-restricted immune response," *PNAS*, 91:6151-6155. cited by applicant

Fukui et al. (1993) "T-cell repertoire in a strain of transgenic C57BL/6 mice with the HLA-DRA gene on the X-chromosome," *Immunogenetics*, 37(3):204-211. cited by applicant

Fukui et al. (1997) "Differential requirement of MHC class II molecules expressed on hematopoietic cells for positive selection of CD4+ thymocytes in TCR α β and TCR β transgenic

mice,” Internal. Immunol., 9(9):1385-1391. cited by applicant

Gao et al. (1997) “Crystal structure of the complex between human CD88alpha-alpha and HLA-A2,” Nature, 387:630-634. cited by applicant

Gao et al. (2000) “Molecular interactions of coreceptor CD8 and MHC class 1: the molecular basis for functional coordination with the T-Cell receptor,” Immunol. Today, 21 (12):630-636. cited by applicant

Germain et al., (2002) “T-Cell Development and The CD4-CD8 Lineage Decision, Nature Reviews, Immunol.,” 2:309-322. cited by applicant

Giraldo and Montoliu (2001) “Size matters: use of YACs, BACs and PACs in transgenic animals,” Transgenic Research, 10:83-103. cited by applicant

Glick and Pasternak Dzh. Moleculyarnaya biotekhnologiya. Printsipy i primeneniye. Moscow: Mir, 2002 (with English translation). cited by applicant

Goldman et al. (2004) “Transgenic animals in medicine: Integration and expression of foreign genes, theoretical and applied aspects,” Med. Sci. Monit., 10(11): RA274-285. cited by applicant

Gomez et al. (2010) “Derivation of cat embryonic stem-like cells from in vitro-produced blastocysts on homologous and heterologous feeder cells,” Theriogenology, 74:498-515. cited by applicant

Gruda et al. (2007) “Intracellular Cysteine Residues in the Tail of MHC Class I Proteins Are Crucial for Extracellular Recognition by Leukocyte Ig-Like Receptor 1,” J. Immunol., 179:3655-3661. cited by applicant

Günther and Walter (2001) “The major histocompatibility complex of the rat (*Rattus norvegicus*),” Immunogenetics, 53:520-542. cited by applicant

Gur et al. (1997) “Structural Analysis of Class I MHC Molecules: The Cytoplasmic Domain Is Not Required for Cytoskeletal Association, Aggregation and Internalization,” Mol. Immunol., 34:125-132. cited by applicant

Güssow et al. (1987) “The Human B2-Microglobulin Gene. Primary Structure and Definition of the Transcriptional Unit,” J. Immunol., 139:3132-3138. cited by applicant

Haks et al. (1999) “Cell-fate decisions in early T cell development: regulation by cytokine receptors and the pre-TCR,” Immunol., 11:23-37. cited by applicant

Hall (2008) “Porcine Embryonic Stem Cells: A Possible Source for Cell Replacement Therapy,” Stem Cell Rev., 4:275-282. cited by applicant

Hanna et al. (1994) “Specific expression of the human CD4 gene in mature CD4 + CD8- and immature CD4+ and CD8+ T cells and in macrophages of transgenic mice,” Mol. Cellular Biol., 14(2):1084-1094. cited by applicant

Harton et al. (2016) “Immunological Functions of the Membrane Proximal Region of MHC Class II Molecules,” F1000Research, 5:1-12, doi: 10.12688/f1000research.7610.1. cited by applicant

Holdsworth et al. (2009) “The HLA dictionary 2008: a summary of Hla-A, -B, -C-DRB1/3/4/5, and -DQB1 alleles and their association with serologically defined Hla-A, -B, -C, -DR, and -DO antigens,” Tissue Antigens, 73:95-170. cited by applicant

Hong et al. (2012) “Derivation and Characterization of Embryonic Stem Cells Lines Derived from Transgenic Fischer 344 and Dark Agouti Rats,” Stem Cells and Development, 21(9):1571-1586. cited by applicant

Hostert et al. (1997) “A CD8 genomic fragment that directs subset-specific expression of CD8 in transgenic mice,” J. Immunol., 158(9):4270-4281. cited by applicant

Hou et al. (2016) “Derivation of Porcine Embryonic Stem-Like Cells from In Vitro-Produced Blastocyst-Stage Embryos,” Scientific Research, 6:1-13. cited by applicant

Houdebine, (1994) “Production of pharmaceutical proteins from transgenic animals,” J. Biotechnology, 34:269-287. cited by applicant

Houdebine, et al (2002) “The methods to generate transgenic animals and to control transgene expression,” J. of Biotech., 98:145-160. cited by applicant

Houdebine, et al (2007) "Transgenic Animal Models in Biomedical Research," *Methods in Molecular Biology*, 360:163-202. cited by applicant

Houdebine (2009) "Methods to Generate Transgenic Animals," *Genetic Engineering in Livestock, New Applications and Interdisciplinary Perspectives*, Engelhard M, et al., 2009, XVI, 1 46 p. 8 illus., pp. 31-48, see p. 36. cited by applicant

Huang et al. (1997) "Analysis of the contact sites on the CD4 Molecule with Class II MHC Molecule," *J. Immunol.*, 158:215-225. cited by applicant

Intarapat and Stern (2013) "Chick stem cells: Current progress and future prospects," *Stem Cell Res.*, 11:1378-1392. cited by applicant

International MHC and Autoimmunity Genetics Network (IMAGEN) (2009) "Mapping of multiple 40 susceptibility variants within the MHC region for 7 immune-mediated diseases," *PNAS*, 106: 18680-18685. cited by applicant

Irie et al. (1998) "The cytoplasmic domain of CD8 beta regulates Lck kinase activation and CD8 T cell development," *J. Immunol.*, 161(1):183-191. cited by applicant

Irwin et al. (1989) "Species-restricted interactions between CD8 and the alpha3 domain of class I influence the magnitude of the xenogeneic response," *J. Exp. Med.*, 170:1091-1101. cited by applicant

Ishimoto et al. (1997) "In vitro and in vivo evidence for high frequency of I-Ab-reactive CD4+ t cells in HLA-DQ or HLA-DRA transgenic mice lacking endogenous MHC class I and/or class II expression," *J. Immunol.*, 159(8):3717-3722. cited by applicant

Itano et al. (1996) "The Cytoplasmic Domain of CD4 Promotes the Development of CD4 Lineage T Cells," *J. Exp. Med.*, 183(3):731-741. cited by applicant

Ito et al. (1996) "HLA-DR4-IE Chimeric Class II Transgenic, Murine Class II-Deficient Mice Are Susceptible to Experimental Allergic Encephalomyelitis," *J. Exp. Med.*, 183:2635-2644. cited by applicant

Jakobovits (1994) "Humanizing the mouse genome," *Cur. Biol.*, 4(8):761-763. cited by applicant

Jean et al. (2013) "Pluripotent genes in avian stem cells," *Develop. Growth Differ.*, 55:41-51. cited by applicant

Johansson et al. (2005) "Natural killer Cell education in mice with single or multiple major histocompatibility complex class I molecules," *J. Exp. Med.*, 201:1145-1155. cited by applicant

Josson et al. (2011) "B2 microglobulin induces epithelial to mesenchymal transition and confers cancer lethality and bone metastasis in human cancer cells," *Cancer Res.*, 71:1-11. cited by applicant

Kalinke et al. (1990) "Strong Xenogeneic HLA Response in Transgenic Mice after Introducing an Alpha3 Domain into HLA B27," *Nature*, 348:642-644. cited by applicant

Kaplan et al. (2005) "A new murine tumor model for studying HLA-A2-restricted anti-tumor immunity," *Cancer Letters*, 224:153-166. cited by applicant

Kappel et al. (1992) "Regulating gene expression in transgenic animals," *Current Opinion in Biotechnology*, 3:548-553. cited by applicant

Kawamata and Ochiya (2010) "Generation of genetically modified rats from embryonic stem cells," *PNAS*, 107(32):14223-14228. cited by applicant

Khor et al. (2002) "Allelic exclusion at the TCR β locus," *Curr. Opin. Immunol.*, 14:230-234. cited by applicant

Kievits et al. (1987) "HLA-restricted recognition of viral antigens in HLA transgenic mice," *Nature* 329:447-449. cited by applicant

Killeen et al. (1993) "Regulated expression of human CD4 rescues helper T cell development in mice lacking expression of endogenous CD4," *EMBO J.*, 12(4): 1547-1553. cited by applicant

Kioussis et al. (2002) "Chromatin and CD4, CD8A and CD8B gene expression during thymic differentiation," *Nature Rev. Immunol.*, 2(12):909-919. cited by applicant

Kirwan et al. (2005) "Killer Cell Ig-Like Receptor-Dependent Signaling by Ig-Like Transcript 2

(ILT2/CD85j/LILRB1/LIR-1)," J. Immunol., 175:5006-5015. cited by applicant

Koller and Orr (1985) "Cloning and complete sequence of an HLA-A2 gene: Analysis of two HLA-A Alleles at the nucleotide Level" J. Immunol. 134(4):2727-2733. cited by applicant

Koller et al. (1990) "Normal Development of Mice Deficient in B2M, MHC Class I Proteins, and CD8+ T cells," Science, 248:1227-1230. cited by applicant

Koop et al., (1994) "The human T-cell receptor TCRAC/TCRDC (C alpha/C delta) region: organization, sequence, and evolution of 97.6 kb of DNA," Genomics 19(3):478-493. cited by applicant

Krangel et al. (1998) "Development regulation of V(D)J recombination at the TCR α/δ locus," Immunol. Rev., 165:131-147. cited by applicant

Kruisbeek et al., (2000) "Branching out to gain control: how the pre-TCR is linked to multiple functions," Rev. Immunol. Today, 21 (12):637-644. cited by applicant

Kumanovics et al. (2003) "Genomic Organization of the Mammalian MHC," Annu. Rev. Immunol., 21:629-657. cited by applicant

Laface et al. (1995) "Human CD8 Transgene Regulation of HLA Recognition by Murine T cells," J. Exp. Med., 182:1315-1325. cited by applicant

Landau et al. (1988) "The envelope glycoprotein of the human immunodeficiency virus binds to the immunoQibulin-like domain of CD4," Nature, 334(6178): 159-162. cited by applicant

Laub et al. (2000) "A multiple transgenic mouse model with a partially humanized activation pathway for helper T cell responses," J. Immunol. Methods, 246(1-2):37-50. cited by applicant

Lauzurica P. and Krangel M.S. (1994) "Enhancer-dependent and -independent Steps in the Rearrangement of a Human T Cell Receptor Delta Transgene," J. Exp. Med., 179:43-55. cited by applicant

Lavial et al., (2008) "Molecular control of pluripotency and germ line competency in chicken embryonic stem cells," Cell Research, 18:s106 (1 page). cited by applicant

Lavial and Pain (2010) "Chicken embryonic stem cells as a non-mammalian embryonic stem cell model," Develop. Growth Differ., 52:101-114. cited by applicant

Law et al. (1994) "Human CD4 Restores Normal T Cell Development and Function in Mice Deficient in Murine CD4," J. Exp. Med., 179(4): 1233-1242. cited by applicant

Leahy (1995) "A structural view of CD4 and CD8," FASEB J., 9:17-25. cited by applicant

Leduc et al. (2000) "T Cell Development in TCR β Enhancer-Deleted Mice: Implications for $\alpha\beta$ T Cell Lineage Commitment and Differentiation," J. Immunol., 165:1364-1373. cited by applicant

Lee et al. (1982) "Sequence of an HLA-DR alpha-chain cDNA clone and intron-exon organization of the corresponding gene," Nature, 299:750-752. cited by applicant

Li et al. (2009) "Mamu-A*01/Kb transgenic and MHC Class I knockout mice as a tool for HIV vaccine development," Virology, 387:16-28. cited by applicant

Li et al. (2010) "Transgenic mice with a diverse human T Cell antigen receptor repertoire," Nature Med., 16(9):1029-1034, doi:10.1038/nm.2197. cited by applicant

Li et al. (2013) "Generation of transgenic mice with megabase-sized human yeast artificial chromosomes by yeast spheroplast-embryonic stem cell fusion," Nat. Protoc., 8(8):1567-1582, doi:10.1038/nprot.2013.093. cited by applicant

Li et al. (2016) "The Implication and Significance of Beta 2 Microglobulin: A Conservative Multifunctional Regulator," Chinese Medical Journal, 129(4):448-455. cited by applicant

Lie and Petropoulos (1998) "Advances in quantitative PCR technology: 5' nuclease assays," Curr. Opin. Biotech., 9:43-48. cited by applicant

Linder C. (2006) "Genetic Variables That Influence Phenotype," ILAR Journal, 47(2):132-140. cited by applicant

Linnenbach et al. (1980) "DNA-transformed murine teratocarcinoma cells: regulation of expression of simian virus 40 tumor antigen in stem versus differentiated cells," PNAS, 77(8):4875-4879. cited by applicant

Littman (1987) "The Structure of the CD4 and CD8 Genes," *Annual Review of Immunology*, 5:561-584. cited by applicant

Lizee et al. (2003) "Control of dendritic cell cross-presentation by the major histocompatibility complex class I cytoplasmic domain," *Nature Immunol.*, 4:1065-1073. cited by applicant

Lynch et al. (2009) "Novel MHC Class I Structures on Exosomes," *J. Immunol.*, 183:1884-1891. cited by applicant

Maddon et al. (1987) "Structure and expression of the human and mouse T4 genes," *PNAS*, 84(24):9155-9159. cited by applicant

Madsen et al. (1999) "A humanized model for multiple sclerosis using HLA-DR2 and a human T-cell receptor," *Nature Genet.*, 23:343-347. cited by applicant

Magliocca et al. (2006) "Undifferentiated Murine Embryonic Stem Cells Cannot Induce Portal Tolerance but May Possess Immune Privilege Secondary to Reduced Major Histocompatibility Complex Antigen Expression," *Stem Cells and Development*, 15(5):707-717. cited by applicant

Maksimenko et al. (2013) "Use of Transgenic Animals in Biotechnology: Prospects and Problems," *ACTNA Naturae*, 5(1):33-46. cited by applicant

Manz et al. (2009) "Renaissance for mouse models of human hematopoiesis and immunobiology," *Nature Immunology*, 10(10):1039-1042. cited by applicant

Marsh et al. (2010) "Nomenclature for factors of the HLA system," 2010, *Tissue Antigens*, 75:291-455. cited by applicant

Marten Hofker et al. (2003) "Transgenic mouse methods and protocols," *Methods in Molecular Biology*, 209:51-58. cited by applicant

Mendez et al. (1997) "Functional transplant of megabase human immunoglobulin loci recapitulates human antibody response in mice," *Nature Genet.*, 15(2):146-156. cited by applicant

Miao (2012) "Recent Advances and Applications of Transgenic Animal Technology," *Polymerase Chain Reaction*, Dr. Patricia Hernandez-Rodriguez (Ed.), ISBN: 978-953-51-0612-8, InTech, pp. 255-282, Available from: <http://www.intechopen.com/books/polymerase-chain-reaction/recent-advances-and-applications-of-transgenic-animal-technology>. cited by applicant

Moir et al. (1996) "Postbinding events mediated by human immunodeficiency virus type 1 are sensitive to modifications in the D4-transmembrane linker region of CD4," *J. Virology*, 70(11):8019-8028. cited by applicant

Moldovan et al. (2002) "CD4 Dimers Constitute the Functional Component Required for T Cell Activation," *J. Immunol.*, 169:6261-6268. cited by applicant

Mombaerts P et al. (1991) "Creation of a large genomic deletion at the T-cell antigen receptor β -subunit locus in mouse embryonic stem cells by gene targeting," *PNAS*, 88:3084-3087. cited by applicant

Mombaerts et al. (1992) "Mutations in T-cell antigen receptor genes α and β block thymocyte development at different stages," *Nature*, 360:225-231. cited by applicant

Mombaerts et al. (1993) "Spontaneous Development of Inflammatory Bowel Disease in T Cell Receptor Mutant Mice," *Cell*, 75:275-282. cited by applicant

Munoz et al. (2008) "Technical note; Conventional pluripotency markers are unspecific for bovine embryonic-derived cell-lines," *Theriogenology*, 69:1159-1164. cited by applicant

Murphy et al. (2008) "Janeway's Immunobiology" 7th ed., Garland Science, pp. 125-13B and 196-213. cited by applicant

Murphy and Silha (2000) "Unexpected and unexplained phenotypes in transgenic models," *Growth Hormone & IGF Research*, 10:233-235. cited by applicant

Musser G. (Encyclopedia Britannica, May 31, 2016) Rodent, Mammal, <http://www.britannica.com/animal/rodent>. cited by applicant

Nakayama et al. (1992) "Recent Duplication of the Two Human CD8 β -chain genes," *J. Immunol.*, 148:1919-1927. cited by applicant

Newberg et al. "Importance of MHC class I α 2 and α 3 domains in the recognition of self

and non-self MHC molecules,” *Journal of Immunology*, 1996, 156:2473-2480. cited by applicant

Nickerson et al. (1990) “Expression of HLA-B27 in Transgenic Mice Is Dependent on the Mouse H-20 Genes,” *J. Exp. Med.*, 172:1255-1261. cited by applicant

Noordzij et al. (2000) “N-terminal truncated human RAG1 proteins can direct T-cell receptor but not immunoglobulin gene rearrangements,” *Blood*, 96(1):203-209. cited by applicant

Norment et al. (1988) “A second subunit of CD8 is expressed in human T cells,” *EMBO J.*, 7(11):3433-3439. cited by applicant

Norment et al. (1989) “Alternatively Spliced mRNA Encodes a Secreted Form of Human CD8 α , Characterization of the Human CD8 α gene,” *J. Immunol.*, 142:3312-3319. cited by applicant

Nowak et al. (1992) “The optimal No. of major histocompatibility complex molecules in an individual,” *Immunology*, 89:10896-10899. cited by applicant

Ostrand-Rosenberg et al. (1991) “Abrogation of Tumorigenicity by MHC Class II Antigen Expression Requires the Cytoplasmic Domain of the Class II Molecule,” *J. Immunol.*, 147:2419-2422. cited by applicant

Pajot et al. (2004) “A mouse model of human adaptive immune functions: HLA-A2.1/HLA-DR1-transgenic H-2 class 1-/class 11-knockout mice,” *Eur. J. Immunol.*, 34:3060-3069. cited by applicant

Paris and Stout (2010) “Embryonic Stem Cells in Domestic Animals; Equine embryos and embryonic stem cells: Defining reliable markers of pluripotency,” *Theriogenology*, 74:516-524. cited by applicant

Parnes, et al. (1982) “Structure of Wild-Type and Mutant Mouse β .SUB.2.-Microglobulin Genes,” *Cell*, 29:661-669. cited by applicant

Pasare et al. (2001) “T cells in mice expressing a transgenic human TCR β chain get positively selected but cannot be activated in the periphery by signaling through TCR,” *Intl. Immunol.*, 13(1):53-62. cited by applicant

Pascolo, et al. (1997) “HLA-A2.1-restricted Education and Cytolytic Activity of CD8+ T Lymphocytes from β 2 Microglobulin (β 2m) HLA-A2.1 Monochain Transgenic H-2Db β 2m Double Knockout Mice,” *J. Exp. Med.*, 185:2043-2051. cited by applicant

Pascolo, et al. (2005) “HLA class I transgenic mice: development, utilisation and improvement,” *Expert Opinion Biol. Ther.*, 5(7):919-938. cited by applicant

Perarnau et al. (1988) “Human B2-microglobulin specifically enhances cell-surface expression of HLA class I molecules in transfected murine cells,” *J. Immunol.*, 141:1383-1389. cited by applicant

Petitte et al. (2004) “Avian pluripotent stem cells,” *Mechanisms of Development*, 121:1159-1168. cited by applicant

Pettersen et al. (1998) “The TCR-Binding Region of the HLA Class I α 2 Domain Signals Rapid Fast Independent Cell, Death: A Direct Pathway for T Cell-Mediated Killing of Target cells,” *J. Immunol.*, 160:4343-4352. cited by applicant

Pittet et al. (2003) “Alpha3 Domain Mutants of Peptide/MHC Class I Multimers Allow the Selective Isolation of High Avidity Tumor-Reactive CD8 T cells,” *J. Immunol.*, 171 :1844-1849. cited by applicant

Potter et al. (1989) “Substitution at residue 227 of H-2 class I molecules abrogates recognition by CD8-dependent, but not CD8-independent, cytotoxic T lymphocytes,” *Nature*, 337:73-75. cited by applicant

Poueymirou et al. (2007) “F0 generation mice fully derived from gene-targeted embryonic stem cells allowing immediate phenotypic analyses,” *Nature Biotech.*, 25:91-99. cited by applicant

Quinn et al. (1997) “Virus-Specific, CD8+ Major Histocompatibility Complex Class 1-Restricted Cytotoxic T Lymphocytes in Lymphocytic Choriomeningitis Virus-Infected B2-Microglobulin-Deficient Mice,” *J. Virol.*, 71 :8392-8396. cited by applicant

Rack et al. (1997) “A Chromosome 14q11/TCR α / δ Specific Yeast Artificial Chromosome Improves the Detection Rate and Characterization of Chromosome Abnormalities in T-Lymphoproliferative

Disorders,” *Blood*, 90(3):1233-1240. cited by applicant

Raffegerst et al. (2009) “Diverse Hematological Malignancies Including Hodgkin-Like Lymphomas Develop in Chimeric MHC Class II Transgenic Mice,” *PLoS ONE*, 4:e8539, 12 pages. cited by applicant

Reipert et al. (2009) “Opportunities and limitations of mouse models humanized for HLA class II antigens,” *Thrombosis and Haemostasis*, 7(Suppl. 1):92-97. cited by applicant

Ren et al. (2006) “Construction of bioactive chimeric MHC class I tetramer by expression and purification of human-murine chimeric MHC heavy chain and β 2M as a fusion protein in *Escherichia coli*,” *Protein Expression and Purification*, 50:171-78. cited by applicant

Rock et al. (2016) “Present Yourself! By MHC Class I and MHC Class II Molecules,” *Trends Immunol.*, 37(11):724-737. cited by applicant

Rodriguez-Cruz et al. (2011) “Natural Splice Variant of MHC Class I Cytoplasmic Tail Enhances Dendritic Cell-Induced CD8+ T-Cell, Responses and Boosts Anti-Tumor Immunity,” *PLoS ONE*, 6:e22939, 10 pages. cited by applicant

Rohrlich et al. (2003) “HLA-B*0702 transgenic, H-2K.SUP.b.D.SUP.b .doubleknockout mice: phenotypical and functional characterization in response to influenza virus,” *International Immunology*, 15(6):765-772. cited by applicant

Rosano et al. (2005) “The three-dimensional structure of B2 microglobulin: Results from X-ray crystallography,” *Biochim. Biophys. Acta*, 1753:85-91. cited by applicant

Rothe et al. (1993) Functional expression of a human TCR β gene in transgenic mice, *Intl. Immunol.*, 5(1):11-17. cited by applicant

Rowen et al., (1996) “The Complete 685-Kilobase DNA Sequence of the Human β T Cell Receptor Locus,” *Science*, 272:1755-1762. cited by applicant

Rubio et al. (2004) “Cross-linking of MHC class I molecules on human NK cells inhibits NK cell function, segregates MHC I from the NK cell synapse, and induces intracellular phosphotyrosines,” *J. Leukoc. Biol.*, 76:116-124. cited by applicant

Salter et al. (1989) “Polymorphism in the alpha3 domain of HLA-A molecules affects binding to CD8,” *Nature*, 338:345-347. cited by applicant

Salter et al. (1990) “A binding site for the T-cell co-receptor CD8 on the alpha 3 domain of HLA-A2,” *Nature*, 345:41-46. cited by applicant

Samberg et al. (1989) “The α 3 domain of major histocompatibility complex class I molecules plays a critical role in cytotoxic T lymphocyte stimulation,” *Eur. J. Immunol.*, 19(12):2349-2354. cited by applicant

Sanders et al. (1991) “Mutations in CD8 that Affect Interactions with HLA Class I and Monoclonal Anti-CD8 Antibodies,” *J. Exp. Med.*, 174:371-379. cited by applicant

Santagata et al. (2000) “The genetic and biochemical basis of Omenn syndrome,” *Immunological Reviews*, 178:64-74. cited by applicant

Scardino et al. (2003) “In vivo study of the GC90/IRIV vaccine for immune response and autoimmunity into a novel humanised transgenic mouse,” *British Journal of Cancer*, 89:199-205. cited by applicant

Scheer et al. (2013) “Generation and utility of genetically humanized mouse models.” *Drug Discovery Today*, 18(23/24):1200-1211. cited by applicant

Schwarz et al. (1996) “RAG mutations in human B cell negative SCID,” *Science*, 274(5284):97-99. cited by applicant

Sebzda et al. (1999) “Selection of the T Cell Repertoire,” *Annu. Rev. Immunol.*, 17:829-874. cited by applicant

Shankar Pradhan and Majumdar, (2016) “An Efficient Method for Generation of Transgenic Rats Avoiding Embryo Manipulation,” *Molecular Therapy—Nucleic Acids*, 5, e293:1-11 doi:10.1038/mtna.2016.9. cited by applicant

Shankarkumar (2004) “The Human Leukocyte Antigen (HLA) System,” *Int. J. Hum. Genet.*, 4:91-

103. cited by applicant

Sherman et al. (1992) "Selecting T Cell Receptors with High Affinity for Self-MHC by Decreasing the Contribution of CD8," *Science*, 258:815-818. cited by applicant

Shields et al. (1998) "Functional comparison of bovine, murine, and human β .SUB.2.-microglobulin: interactions with murine MHC I molecules," *Molecular Immunology*, 35:919-928. cited by applicant

Shin et al. (2006) "Surface expression of MHC class II in dendritic cells is controlled by regulated ubiquitination," *Nature*, 444:115-118. cited by applicant

Shinkai et al. (1992) "RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement," *Cell* 68(5):855-67. cited by applicant

Shinohara et al. (2007) "Active integration: new strategies for transgenesis," *Transgenic Res.*, 16:333-339. cited by applicant

Shiroishi et al. (2003) "Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G," *PNAS*, 100:8856-8861. cited by applicant

Shultz et al. (2007) "Humanized mice in translational biomedical research," *Nature Rev.*, 7:118-130. cited by applicant

Sigmund (2000) "Viewpoint: Are Studies in Genetically Altered Mice Out of Control?," *Arterioscler. Thromb. Vasc. Biol.*, p. 1425-1429. cited by applicant

Sims et al (2004) "Genetic Susceptibility to Ankylosing Spondylitis," *Cur. Mol. Med.*, 4(1):13-20. cited by applicant

Singer et al. (2008) "Lineage fate and intense debate: myths, models and mechanisms of CD4/CD8 lineage choice," *Nature Rev. Immunol.*, 8(10):788-801. cited by applicant

Sittig et al. (2016) "Genetic Background Limits Generalizability of Genotype-Phenotype Relationships," *Neuron*, 91(6):1253-1359. cited by applicant

Siu et al. (1994) "A transcriptional silencer controls the developmental expression of the CD4 gene," *EMBO J.*, 13(15):3570-3579. cited by applicant

Sleckman et al. (2000) "Mechanisms that direct ordered assembly of T cell receptor β locus V, D, and J gene segments," *PNAS*, 97(14):7975-7980. cited by applicant

Smiley et al. (1995) "Transgenic mice expressing MHC class II molecules with truncated A-beta cytoplasmic domains reveal signaling-independent defects in antigen presentation," *Internal. Immunol.*, 7:665-677. cited by applicant

Smiley et al. (1996) "Truncation of the class II beta-chain cytoplasmic domain influences the level of class II/invariant chain-derived peptide complexes," *PNAS*, 93:241-244. cited by applicant

Street et al. (2002) "Limitations of HLA-transgenic mice in presentation of HLA-restricted cytotoxic T-cell epitopes from endogenously processed human papillomavirus type 16 E7 protein," *Immunology*, 106:526-536. cited by applicant

Sundelin et al. (1988) "The Complete Amnio Acid Sequence of Rat β 2-Microglobulin," *Scand. J. Immunol.*, 27:195-199. cited by applicant

Takaki et al. (2006) "HLA-A*0201-Restricted T cells from Humanized NOD Mice Recognize Autoantigens of Potential Clinical Relevance to Type I Diabetes," *J. Immunol.*, 176:3257-3265. cited by applicant

Tanabe et al. (1989) "Analysis of xenoantigenicity of HLA Class I molecules by a complete series of human-mouse hybrid genes," *Transplantation*, 48:1, 135-140. cited by applicant

Taneja and David (1998) "HLA Transgenic Mice as Humanized Mouse Models of Disease and Immunity," *J. Clin. Invest.*, 101:921-926. cited by applicant

Taneja and David (2009) "Spontaneous autoimmune myocarditis and cardiomyopathy in HLA-DQ8.NODAb0 transgenic mice," *Journal of Autoimmunity* 33:260-269. cited by applicant

Taylor et al. (1993) "Human immunoglobulin transgenes undergo rearrangement, somatic mutation and class switching in mice that lack endogenous IgM," *Int. Immunol.*, 6(4):579-591. cited by

applicant

Theobald et al. (1995) "Targeting p53 as a general tumor antigen," PNAS, 92:11993-11997. cited by applicant

Tishon et al. (2000) "Transgenic Mice Expressing Human HLA and CD8 Molecules Generate HLA-Restricted Measles Virus Cytotoxic T Lymphocytes of the Same Specificity as Humans with Natural Measles Infection," Virology, 275:286-293. cited by applicant

Tong et al. (2010) "Production of p53 gene knockout rats by homologous recombination in embryonic stem cells," Nature Letters, 467:211-215. cited by applicant

Toyonaga et al. (1985) "Organization and sequences of the diversity, joining, and constant region genes of the human T-cell receptor beta chain," PNAS. 82(24):8624-8628. cited by applicant

Tran et al. (2006) "Additional human β 2-microglobulin curbs HLA-B27 misfolding and promotes arthritis and spondylitis without colitis in male HLA-B27-transgenic rats," Arthritis & Rheumatism, 54:1317-1327. cited by applicant

Ureta-Vidal et al. (1999) "Phenotypical and Functional Characterization of the CD8 + T Cell Repertoire of HLA-A2.1 Transgenic, H-2Kb^oDb^o Double Knockout Mice," J. Immunol., 163:2555-2560. cited by applicant

Valenzuela et al. (2003) "High-throughput engineering of the mouse genome coupled with high-resolution expression analysis," Nature Biotech., 21:652-659. cited by applicant

Vignali et al. (1992) "Species-specific Binding of CD4 to the Beta2 Domain of Major Histocompatibility Complex Class II Molecules," J. Exp. Med., 175:925-932. cited by applicant

Vignali et al. (1996) "The Two Membrane Proximal Domains of CD4 Interact with the T Cell Receptor," J. Exp. Med., 183:2097-2107. cited by applicant

Villa et al. (1998) "Partial V(D)J recombination activity leads to Omenn syndrome," Cell 93(5): 855-896. cited by applicant

Villa et al. (1999) "Omenn syndrome: a disorder of Rag1 and Rag2 genes," J. Clin. Immunol., 19(2):87-97. cited by applicant

Viney et al. (1992) "Generation of Monoclonal Antibodies Against a Human T Cell Receptor β chain Expressed in Transgenic Mice," Hybridoma, 11(6):701-714. cited by applicant

Vitiello et al. (1991) "Analysis of the HLA-restricted Influenza-specific Cytotoxic T Lymphocyte Response in 6 Transgenic Mice Carrying a Chimeric Human-Mouse Class I Major Histocompatibility Complex," J. Exp. Med., 173:1007-1015. cited by applicant

Vollmer et al. (2000) "Antigen contacts by Ni-reactive TCR: typical $\alpha\beta$ chain cooperation versus a chain-dominated specificity," IntL. Immunol., 12(12):1723-1731. cited by applicant

Vugmeyster et al. (1998) "Major histocompatibility complex (MHC) class I KbDb $-/-$ deficient mice possess functional CD8 + T cells and natural killer cells," PNAS, 95:12492-12497. cited by applicant

Wagner et al. (1994) "Antibodies generated from human immunoglobulin miniloci in transgenic mice," Nucleic Acids Res., 22(8):1389-1393. cited by applicant

Wagner et al. (1994) "Ligation of MHC Class I and Class II Molecules Can Lead to Heterologous Desensitization of Signal Transduction Pathways That Regulate Homotypic Adhesion in Human Lymphocytes," J. Immunol., 152:5275-5287. cited by applicant

Wagner et al. (1994) "The diversity of antigen-specific monoclonal antibodies from transgenic mice bearing human immunoglobulin gene miniloci," Eur. J. Immunol., 24(11):2672-2681. cited by applicant

Wang and Reinherz (2001) "Structural basis of T cell recognition of peptides bound to MHC molecules," Mol. Immunol., 38:1039-1049. cited by applicant

Wei et al. (1996) "Repertoire and Organization of Human T-Cell Receptor α Region Variable Genes," Short Communication, Genomics, 38:442-445. cited by applicant

Wen et al. (1998) "Induction of Insulitis by Glutamic Acid Decarboxylase Peptide-specific and HLA-DQ8-restricted CD4+ T Cells from Human DQ Transgenic Mice," J. Clin. Invest.,

102(5):947-957. cited by applicant

Willcox et al. (2003) "Crystal structure of HLA-A2 bound to LIR-1, a host and viral major histocompatibility complex receptor," *Nature Immunol.*, 4:913-919. cited by applicant

Willinger et al. (2011) "Improving human hemato-lymphoid-system mice by cytokine knock-in gene replacement," *Trands in Immunology*, 32(7):321-327. cited by applicant

Wong and Wen (2004) "What can the HLA transgenic mouse tell us about autoimmune diabetes?," *Diabetologia*, 47:1476-1487. cited by applicant

Woodle et al. (1997) "Anti-Human Class I MHC Antibodies Induce Apoptosis by a Pathway That Is Distinct from the Fas Antigen-Meidaled Pathway," *J. Immunol.*, 158:2156-2164. cited by applicant

Woods et al. (1994) "Human Major Histocompatibility Complex Class 11-Restricted T Cell, Responses in Transgenic Mice," *J. Exp. Med.*, 180:173-181. cited by applicant

Wooldridge et al. (2010) "MHC Class I Molecules with Superenhanced CD8 Binding Properties Bypass the Requirement for Cognate TCR Recognition and Nonspecifically Activate CTLs," *J. Immunol.*, 184:3357-3366. cited by applicant

Wu et al. (1997) "Dimeric association and segmental variability in the structure of human CD4," *Nature*, 387:527. cited by applicant

Yamamoto et al. (1994) "Functional Interaction between Human Histocompatibility Leukocyte Antigen (HLA) Class 27 II and Mouse CD4 Molecule in Antigen Recognition by T cells in HLA-DR and DQ Transgenic Mice," *J. Exp. Med.*, 180:165-171. cited by applicant

Yantha et al., (2010) "Unexpected Acceleration of Type 1 Diabetes by Transgenic Expression of B7-H1 in NOD Mouse Peri-Islet Glia," *Diabetes*, 59:2588-2596. cited by applicant

Yoshikai et al. (1985) "Organization and sequences of the variable, joining and constant region genes of the human T-cell receptor alpha-chain," *Nature*, 316(6031):837-840. cited by applicant

Zamoyska (1998) "CD4 and CD8: modulators of T cell receptor recognition of antigen and of immune responses?," *Curr. Opin. Immunol.*, 10:82-87. cited by applicant

Zhou et al., (2009) "Developing tTA transgenic rats for inducible and reversible gene expression," *International Journal of Biological Sciences*, 5:171-181. cited by applicant

Zhu et al. (2019) "Humanising the mouse genome piece by piece," *Nature Communications*, 10(1845):1-13. cited by applicant

Zijlstra et al. (1990) "B2-Microglobulin deficient mice lack CD4-8+ cytolytic T Cells," *Nature*, 344:742-746. cited by applicant

Zumla et al. (1992) "Co-expression of human T cell receptor chains with mouse CD3 on the cell surface of a mouse T cell hybridoma," *J. Immunol. Methods.*, 149(1):69-76. cited by applicant

Zumla et al. (1992) "Use of a murine T-cell hybridoma expressing human T-cell receptor alpha-and betagene products as a tool for the production of human T-cell receptor-specific monoclonal antibodies," *Human Immunol.*, 35(3):141-148. cited by applicant

International Search Report and Written Opinion for PCT/US2014/023076, mailed Jul. 18, 2014. cited by applicant

International Search Report and Written Opinion for PCT/US2012/062042, mailed Feb. 18, 2013. cited by applicant

International Search Report and Written Opinion for PCT/US2012/062029, mailed Feb. 28, 2013. cited by applicant

International Search Report and Written Opinion for PCT/US2014/023068, mailed Jul. 24, 2014. cited by applicant

International Search Report and Written Opinion for PCT/US2014/017395, mailed Jun. 2, 2014. cited by applicant

International Search Report and Written Opinion for PCT/US2014/017387, mailed Jun. 2, 2014. cited by applicant

International Search Report and Written Opinion for PCT/US2016/026260, mailed Aug. 11, 2016.

cited by applicant

Extended European Search Report with respect to Europe Application No. 17184955.7, mailed Nov. 24, 2017. cited by applicant

English Translation of Office Action with Respect to Japan Application No. 2014-539027 Mailed Sep. 6, 2016. cited by applicant

Statement of Relatedness under MPEP 2001.06 with Respect to U.S. Application Serial No. to be assigned, dated Dec. 14, 2021. cited by applicant

Landsverk, Ole., "Invariant chain and internalization of mature MHC class II from the cell surface," Department of Molecular Biosciences, University of Oslo, Masters Thesis; Published: Aug. 2005 (Year 2005) (59 pages). cited by applicant

Simonsen et al. "Polarized Transport of MHC Class II Molecules in Madin-Darby Canine Kidney Cells is Directed by Leucine-Based Signal In the Cytoplasmic Tail of the β chain," J. Immunol., 1999, 163(5):2540-2546. cited by applicant

Sønderstrup et al. (1999) HLA class II transgenic mice: models of the human CD4+ T-cell immune response, Immunological Reviews, 172(1):335-343. cited by applicant

Touloukian et al. "Identification of a MHC Class II-Restricted Human gp100 Epitope Using DR4-IE Transgenic Mice," Author Manuscript, J. Immunol, 2000, 164(7):3535-3542. cited by applicant

Statement of Relatedness under MPEP 2001.06 with Respect to U.S. Appl. No. 17/550,945 dated Jul. 18, 2024. cited by applicant

Dechiara et al. "VelociMouse: Fully ES Cell-Derived FO-Generation Mice Obtained from the Injection of ES Cells into Eight-Cell-Stage Embryos," Chapter 16, Gene Knockout Protocols, Second Edition, Methods in Molecular Biology, Humana Press, 530:311-324, (2009), 36 pages. cited by applicant

Mangalam et al. "HLA Class II Transgenic Mice Mimic Human Inflammatory Diseases," Advances in Immunology, 97:65-147, (2008) Chapter 2, 83 pages. cited by applicant

Parham, Peter "The Immune System" Third Edition, Garland Science, Taylor & Francis Group, London and New York, 2009, 24 pages. cited by applicant

Reveille and Bruce "MHC Class II and Non-MHC Genes in the Pathogenesis of Systemic Lupus Erythematosus," Chapter 4, Systemic Lupus Erythematosus, 109-150 (2004), 43 pages. cited by applicant

Wielders et al. "Generation of Double-Knockout Embryonic Stem Cells," Chapter 11, Gene Knockout Protocols, Second Edition, Methods in Molecular Biology, Humana Press, 530:205-218, (2009), 36 pages. cited by applicant

Cohen, et al. (2006) "Enhanced Antitumor Activity of Murine-Human Hybrid T-Cell Receptor (TCR) in Human Lymphocytes is Associated with Improved Pairing and TCR/CD3 Stability," Cancer Res., 66:8878-8886. cited by applicant

Kawamura et al. (2008) "Different Development of Myelin Basic Protein Agonist-and-Antagonist-Specific Human TCR Transgenic T Cells in the Thymus and Periphery," J. Immunol., 181(8):5462-5472. cited by applicant

Massimo et al. (2008) "Primer: Immunity and Autoimmunity," Diabetes, 57(11):2872-2882. cited by applicant

Williams and Barclay et al. (1988) "The Immunoglobulin Superfamily-Domains for Cell Surface Recognition," Ann. Rev. Immunol., 6:381-405. cited by applicant

Shiue et al. "A Second Chain of Human CD8 is Expressed on Peripheral Blood Lymphocytes," J. Exp. Med., Dec. 1988, 168:1993-2005. cited by applicant

Statement of Relatedness under MPEP 2001.06 with Respect to U.S. Appl. No. 17/550,945 dated Jan. 22, 2025. cited by applicant

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS (1) This application is a continuation of U.S. application Ser. No. 16/184,473, filed Nov. 8, 2018, now U.S. Pat. No. 11,224,208, issued on Jan. 18, 2022, which is a continuation of U.S. application Ser. No. 14/185,316, filed Feb. 20, 2014, now U.S. Pat. No. 10,154,658, issued Dec. 18, 2018, which claims benefit under 35 U.S.C. § 119 (e) of U.S. provisional patent application Ser. No. 61/767,811, filed Feb. 22, 2013, each of which application is hereby incorporated by reference in its entirety.

SEQUENCE LISTING

(1) The present specification makes reference to a sequence listing submitted in electronic Form as an ascii .txt file named "1700US03_ST25" on Mar. 17, 2022. The .txt file was generated on Mar. 17, 2022, and is 8 kb in size. The same is incorporated herein by reference in its entirety.

FIELD OF THE INVENTION

(2) Present invention relates to a genetically modified non-human animal, e.g., a rodent (e.g., a mouse or a rat), that expresses a human or humanized Major Histocompatibility Complex (MHC) class I and a human or humanized MHC class II molecules. The invention also relates to a genetically modified non-human animal, e.g., a mouse or a rat, that expresses a human or humanized MHC I protein (e.g., MHC I α chain) and a human or humanized MHC II protein (e.g., MHC II α and MHC II β chains), and further expresses a human or humanized β 2 microglobulin; as well as embryos, tissues, and cells expressing the same. The invention further provides methods for making a genetically modified non-human animal that expresses both human or humanized MHC class I and class II proteins, and/or β 2 microglobulin. Also provided are methods for identifying and evaluating peptides in the context of a humanized cellular immune system in vitro or in a genetically modified non-human animal, and methods of modifying an MHC locus of a non-human animal, e.g., a mouse or a rat, to express a human or humanized MHC I and a human or humanized MHC II proteins.

BACKGROUND OF THE INVENTION

(3) In the adaptive immune response, foreign antigens are recognized by receptor molecules on B lymphocytes (e.g., immunoglobulins) and T lymphocytes (e.g., T cell receptor or TCR). These foreign antigens are presented on the surface of cells as peptide fragments by specialized proteins, generically referred to as major histocompatibility complex (MHC) molecules. MHC molecules are encoded by multiple loci that are found as a linked cluster of genes that spans about 4 Mb. In mice, the MHC genes are found on chromosome 17, and for historical reasons are referred to as the histocompatibility 2 (H-2) genes. In humans, the genes are found on chromosome 6 and are called human leukocyte antigen (HLA) genes. The loci in mice and humans are polygenic; they include three highly polymorphic classes of MHC genes (class I, II and III) that exhibit similar organization in human and murine genomes (see FIG. 2 and FIG. 3, respectively).

(4) MHC loci exhibit the highest polymorphism in the genome; some genes are represented by >300 alleles (e.g., human HLA-DR β and human HLA-B). All class I and II MHC genes can present peptide fragments, but each gene expresses a protein with different binding characteristics, reflecting polymorphisms and allelic variants. Any given individual has a unique range of peptide fragments that can be presented on the cell surface to B and T cells in the course of an immune response.

(5) Both humans and mice have class I MHC genes (see FIG. 2 and FIG. 3). In humans, the

classical class I genes are termed HLA-A, HLA-B and HLA-C, whereas in mice they are H-2K, H-2D and H-2L. Class I molecules consist of two chains: a polymorphic α -chain (sometimes referred to as heavy chain) and a smaller chain called β 2-microglobulin (also known as light chain), which is generally not polymorphic (FIG. 1, left). These two chains form a non-covalent heterodimer on the cell surface. The α -chain contains three domains (α 1, α 2 and α 3). Exon 1 of the α -chain gene encodes the leader sequence, exons 2 and 3 encode the α 1 and α 2 domains, exon 4 encodes the α 3 domain, exon 5 encodes the transmembrane domain, and exons 6 and 7 encode the cytoplasmic tail. The α -chain forms a peptide-binding cleft involving the α 1 and α 2 domains (which resemble Ig-like domains) followed by the α 3 domain, which is similar to β 2-microglobulin.

(6) β 2 microglobulin is a non-glycosylated 12 kDa protein; one of its functions is to stabilize the MHC class I α -chain. Unlike the α -chain, the β 2 microglobulin does not span the membrane. The human β 2 microglobulin locus is on chromosome 15, while the mouse locus is on chromosome 2. The β 2 microglobulin gene consists of 4 exons and 3 introns. Circulating forms of β 2 microglobulin are present in serum, urine, and other body fluids; non-covalently MHC I-associated β 2 microglobulin can be exchanged with circulating β 2 microglobulin under physiological conditions.

(7) Class I MHC molecules are expressed on all nucleated cells, including tumor cells. They are expressed specifically on T and B lymphocytes, macrophages, dendritic cells and neutrophils, among other cells, and function to display peptide fragments (typically 8-10 amino acids in length) on the surface to CD8⁺ cytotoxic T lymphocytes (CTLs). CTLs are specialized to kill any cell that bears an MHC I-bound peptide recognized by its own membrane-bound TCR. When a cell displays peptides derived from cellular proteins not normally present (e.g., of viral, tumor, or other non-self origin), such peptides are recognized by CTLs, which become activated and kill the cell displaying the peptide.

(8) Both humans and mice also have class II MHC genes (see FIGS. 2 and 3). In humans, the classical MHC II genes are termed HLA-DP, HLA-DQ, and HLA-DR, whereas in mice they are H-2A and H-2E (often abbreviated as I-A and I-E, respectively). Additional proteins encoded by genes in the MHC II locus, HLA-DM and HLA-DO in humans, and H-2M and H-2O in mice, are not found on the cell surface, but reside in the endocytic compartment and ensure proper loading of MHC II molecules with peptides. Class II molecules consist of two polypeptide chains: α chain and β chain. The extracellular portion of the α chain contains two extracellular domains, α 1 and α 2; and the extracellular portion of the β chain also contains two extracellular domains, β 1 and β 2 (see FIG. 1, right). The α and the β chains are non-covalently associated with each other.

(9) MHC class II molecules are expressed on antigen-presenting cells (APCs), e.g., B cells, macrophages, dendritic cells, endothelial cells during a course of inflammation, etc. MHC II molecules expressed on the surface of APCs typically present antigens generated in intracellular vesicles to CD4⁺ T cells. In order to participate in CD4⁺ T cell engagement, the MHC class II complex with the antigen of interest must be sufficiently stable to survive long enough to engage a CD4⁺ T cell. When a CD4⁺ T helper cell is engaged by a foreign peptide/MHC II complex on the surface of APC, the T cell is activated to release cytokines that assist in immune response to the invader.

(10) Not all antigens will provoke T cell activation due to tolerance mechanisms. However, in some diseases (e.g., cancer, autoimmune diseases) peptides derived from self-proteins become the target of the cellular component of the immune system, which results in destruction of cells presenting such peptides. There has been significant advancement in recognizing antigens that are clinically significant (e.g., antigens associated with various types of cancer). However, in order to improve identification and selection of peptides that will provoke a suitable response in a human T cell, in particular for peptides of clinically significant antigens, there remains a need for in vivo and in vitro systems that mimic aspects of human immune system. Thus, there is a need for biological systems (e.g., genetically modified non-human animals and cells) that can display components of a

human immune system.

SUMMARY OF THE INVENTION

(11) A biological system for generating or identifying peptides that associate with human MHC class I proteins and chimeras thereof and bind CD8⁺ T cells, as well as peptides that associate with human MHC class II proteins and chimeras thereof and bind to CD4⁺ T cells, is provided. Non-human animals comprising non-human cells that express humanized molecules that function in the cellular immune response are provided. Humanized rodent loci that encode humanized MHC I and MHC II proteins are also provided. Humanized rodent cells that express humanized MHC molecules are also provided. In vivo and in vitro systems are provided that comprise humanized rodent cells, wherein the rodent cells express one or more humanized immune system molecules.

(12) In various embodiments, provided herein is a non-human animal comprising at an endogenous MHC locus a first nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein a human portion of the chimeric MHC I polypeptide comprises an extracellular domain of a human MHC I polypeptide; a second nucleotide sequence encoding a chimeric human/non-human MHC II α polypeptide, wherein a human portion of the chimeric human/non-human MHC II α polypeptide comprises an extracellular domain of a human MHC II α polypeptide; and a third nucleotide sequence encoding a chimeric human/non-human MHC II β polypeptide, wherein a human portion of the chimeric human/non-human MHC II β polypeptide comprises an extracellular domain of a human MHC II β polypeptide, wherein the non-human animal expresses functional chimeric human/non-human MHC I and MHC II proteins from its endogenous non-human MHC locus. In one embodiment, the animal does not express functional endogenous MHC I, II α , and/or II β polypeptides from the endogenous non-human MHC locus.

(13) In one aspect, the first nucleotide sequence is located at the endogenous non-human MHC I locus, the second nucleotide sequence is located at the endogenous non-human MHC II α locus, and the third nucleotide sequence is located at the endogenous non-human MHC II β locus. In one aspect, the first, second and/or third nucleotide sequence(s) are operably linked to endogenous non-human regulatory elements. In one aspect, the first nucleotide sequence is operably linked to endogenous non-human MHC I promoter and regulatory elements, the second nucleotide sequence is operably linked to endogenous non-human MHC II α promoter and regulatory elements, and the third nucleotide sequence is operably linked to endogenous non-human MHC II β promoter and regulatory elements.

(14) In one embodiment, the human portion of a chimeric MHC I polypeptide comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of the human MHC I polypeptide. In one aspect, a non-human portion of the chimeric MHC I polypeptide comprises transmembrane and cytoplasmic domains of an endogenous non-human MHC I polypeptide. The human MHC I polypeptide may be selected from the group consisting of HLA-A, HLA-B, and HLA-C. In one embodiment, the human MHC I polypeptide is HLA-A2. In another aspect, the human MHC I polypeptide is HLA-A3, HLA-B7, HLA-B27, HLA-Cw6, or any other MHC I molecule expressed in a human population. In an additional embodiment, a non-human animal of the invention further comprises at its endogenous non-human $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide, wherein the animal expresses the human or humanized $\beta 2$ microglobulin polypeptide.

(15) In one embodiment, the human MHC II α extracellular domain comprises human MHC II $\alpha 1$ and $\alpha 2$ domains. In another embodiment, the human MHC II β extracellular domain comprises human MHC II $\beta 1$ and $\beta 2$ domains. In one aspect, the non-human portion of a chimeric human/non-human MHC II α polypeptide comprises transmembrane and cytoplasmic domains of an endogenous non-human MHC II α polypeptide. In one aspect, the non-human portion of a chimeric human/non-human MHC II β polypeptide comprises transmembrane and cytoplasmic domains of an endogenous non-human MHC II β polypeptide. In one embodiment, the human portions of a chimeric human/mouse MHC II α and β polypeptides are derived from a human HLA

class II protein selected from the group consisting of HLA-DR, HLA-DQ, and HLA-DP. In one specific embodiment, the human portions of chimeric human/non-human MHC II α and β polypeptides are derived from a human HLA-DR4 protein. Alternatively, the human portions of chimeric human/non-human MHC II α and β polypeptides may be derived from human MHC II protein selected from HLA-DR2, HLA-DQ2.5, HLA-DQ8, or any other MHC II molecule expressed in a human population.

(16) In some aspects, a provided animal comprises two copies of the MHC locus containing the first, the second, and the third nucleotide sequences, while in other aspects, a provided animal comprises one copy of the MHC locus containing the first, the second, and the third nucleotide sequences. Thus, the animal may be homozygous or heterozygous for the MHC locus containing nucleotide sequences encoding chimeric human/non-human MHC I, MHC II α , and MHC II β polypeptides. In some embodiments of the invention, the genetically modified MHC locus, comprising nucleotide sequences encoding chimeric human/non-human MHC I, MHC II α , and MHC II β polypeptides described herein, is in the germline of the non-human animal.

(17) Also provided herein is an MHC locus comprising a first nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein a human portion of the chimeric MHC I polypeptide comprises an extracellular domain of a human MHC I polypeptide; a second nucleotide sequence encoding a chimeric human/non-human MHC II α polypeptide, wherein a human portion of the chimeric human/non-human MHC II α polypeptide comprises an extracellular domain of a human MHC II α polypeptide; and a third nucleotide sequence encoding a chimeric human/non-human MHC II β polypeptide, wherein a human portion of the chimeric human/non-human MHC II β polypeptide comprises an extracellular domain of a human MHC II β polypeptide. In some aspects, non-human portions of the chimeric MHC I, II α , and II β polypeptides comprise transmembrane and cytoplasmic domains of non-human MHC I, II α , and II β , respectively.

(18) In one embodiment, the genetically engineered non-human animal is a rodent. In one embodiment, the rodent is a rat or a mouse. In one embodiment, the rodent is a mouse. Thus, in one aspect, the first nucleotide sequence encodes a chimeric human/mouse MHC I polypeptide, and the mouse portion of the chimeric MHC I polypeptide is derived from H-2K, H-2D, or H-2L. In one specific embodiment, the mouse portion of the chimeric MHC I polypeptide is derived from H-2K. In one aspect, the second nucleotide sequence encodes a chimeric human/mouse MHC II α polypeptide, the third nucleotide sequence encodes a chimeric human/mouse MHC II β polypeptide, and the mouse portions of the chimeric MHC II α and β polypeptides are derived from H-2E or H-2A. In a specific embodiment, the mouse portions of the chimeric MHC II polypeptides are derived from H-2E.

(19) Thus, also provided herein is a genetically engineered mouse comprising at an endogenous MHC locus a first nucleotide sequence encoding a chimeric human/mouse MHC I polypeptide, wherein the human portion of the chimeric MHC I polypeptide comprises an extracellular domain of a human MHC I polypeptide; a second nucleotide sequence encoding a chimeric human/mouse MHC II α polypeptide, wherein the human portion of the chimeric human/non-human MHC II α polypeptide comprises an extracellular domain of a human MHC II α polypeptide; and a third nucleotide sequence encoding a chimeric human/mouse MHC II β polypeptide, wherein the human portion of the chimeric human/non-human MHC II β polypeptide comprises an extracellular domain of a human MHC II β polypeptide; wherein the mouse expresses functional chimeric human/mouse MHC I and MHC II proteins from its endogenous mouse MHC locus. In one specific embodiment, the first nucleotide sequence encodes a chimeric HLA-A2/H-2K polypeptide, the second nucleotide sequence encodes an α chain of a chimeric HLA-DR4/H-2E polypeptide, and the third nucleotide sequence encodes a β chain of a chimeric HLA-DR4/H-2E polypeptide, and the mouse expresses functional HLA-A2/H-2K and HLA-DR4/H-2E proteins. In an additional embodiment, the mouse further comprises at an endogenous β 2 microglobulin locus a nucleotide

sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide. In one embodiment, the mouse does not express functional endogenous MHC polypeptides from its endogenous MHC locus.

(20) Also provided herein are methods for generating a genetically modified non-human animal (e.g., rodent, e.g., mouse or rat) described herein. Thus, in one aspect, the invention provides a method of generating a genetically modified non-human animal comprising replacing at an endogenous non-human MHC II locus a nucleotide sequence encoding a non-human MHC II complex with a nucleotide sequence encoding a chimeric human/non-human MHC II complex to generate a first non-human animal; and replacing at an endogenous non-human MHC I locus a nucleotide sequence encoding a non-human MHC I polypeptide with a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide to generate a second non-human animal. In one aspect, the steps of replacing nucleotide sequences comprise homologous recombination in non-human ES cells, and the second non-human animal is generated by homologous recombination in ES cells bearing nucleotide sequences encoding chimeric human/non-human MHC II complex. The chimeric MHC II complex comprises chimeric human/non-human MHC II α and β polypeptides.

(21) In an alternative embodiment, the invention provides a method of generating a genetically modified non-human animal comprising replacing at an endogenous non-human MHC I locus a nucleotide sequence encoding a non-human MHC I polypeptide with a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide to generate a first non-human animal; and replacing at an endogenous non-human MHC II locus a nucleotide sequence encoding a non-human MHC II complex with a nucleotide sequence encoding a chimeric human/non-human MHC II complex to generate a second non-human animal. In one aspect, the steps of replacing nucleotide sequences comprise homologous recombination in non-human ES cells, and the second non-human animal is generated by homologous recombination in ES cells bearing a nucleotide sequence encoding chimeric human/non-human MHC I polypeptide.

(22) Also provided herein are cells, e.g., isolated antigen-presenting cells, derived from the non-human animals (e.g., rodents, e.g., mice or rats) described herein. Tissues and embryos derived from the non-human animals described herein are also provided.

(23) Any of the embodiments and aspects described herein can be used in conjunction with one another, unless otherwise indicated or apparent from the context. Other embodiments will become apparent to those skilled in the art from a review of the ensuing detailed description. The following detailed description includes exemplary representations of various embodiments of the invention, which are not restrictive of the invention as claimed. The accompanying figures constitute a part of this specification and, together with the description, serve only to illustrate embodiments and not to limit the invention.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 is a schematic drawing of the MHC I (left panel) and MHC II (right panel) class molecules expressed on the surface of a cell. The gray circles represent peptides bound in the peptide-binding clefts.

(2) FIG. 2 is a schematic representation (not to scale) of the relative genomic structure of the human HLA, showing class I, II and III genes.

(3) FIG. 3 is a schematic representation (not to scale) of the relative genomic structure of the mouse MHC, showing class I, II and III genes.

(4) FIG. 4 depicts the strategy for generating a humanized MHC locus comprising humanized MHC I and MHC II genes. In the particular embodiment depicted, the MHC locus of the generated

mouse comprises human HLA-A2 and HLA-DR4 sequences (H2-K.sup.+/1666 MHC-II.sup.+/1681). Large Targeting Vectors introduced into ES cells at each stage of humanization are depicted to the right of the arrows.

(5) FIG. 5 (A-D) is a schematic illustration (not to scale) of the strategy for generating a targeting vector comprising humanized I-E β and I-E α (i.e., H-2E β /HLA-DR β 1*04 and H-2E α /HLA-DR α *01 chimera, respectively). In FIG. 5C, the final humanized MHC II sequence from FIG. 5B is ligated between PI-SceI and I-CeuI restriction sites of the final construct from FIG. 5A, to generate a construct comprising humanized MHC II and exon 1 of I-E α from BALB/c. Pg=pseudogene; BHR=bacterial homologous recombination; CM=chloramphenicol; spec=spectinomycin; hyg=hygromycin; neo=neomycin; EP=electroporation. Triangles represent exons, filled triangles represent mouse exons from C57BL/6 mouse (with the exception of hashed triangles, which represent exon 1 of I-E α from BALB/c mouse) and open triangles represent human exons.

(6) FIG. 6 shows a schematic illustration, not to scale, of MHC class II I-E and I-A genes, showing knockout of the mouse locus using a hygromycin cassette, followed by introduction of a vector comprising a humanized I-E β and I-E α (i.e., H-2E β /HLA-DR β 1*04 and H-2E α /HLA-DR α *01 chimera, respectively). Open triangles represent human exons; filled triangles represent mouse exons. Probes used for genotyping are encircled.

(7) FIG. 7 shows a schematic illustration, not to scale, of Cre-mediated removal of the neomycin cassette of FIG. 6. Open triangles represent human exons; filled triangles represent mouse exons. Top two strands represent MHC II loci in humanized MHC II heterozygous mouse harboring a neomycin selection cassette, and bottom two strands represent MHC II loci in humanized MHC II heterozygous mouse with neomycin cassette removed.

(8) FIG. 8 is a schematic diagram (not to scale) of the targeting strategy used for making a chimeric H-2K locus that expresses an extracellular region of a human HLA-A2 protein. Mouse sequences are represented in black and human sequences are represented in white. L=leader, UTR=untranslated region, TM=transmembrane domain, CYT=cytoplasmic domain, HYG=hygromycin.

(9) FIG. 9 is a dot plot of in vivo expression of HLA-A2 and HLA-DR4 in heterozygous mice harboring chimeric HLA-A2/H-2K and HLA-DR4/H-2E loci. Steady state HLA-DR4 expression was low but present; this low expression was expected and the expression gets upregulated upon activation.

DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

Definitions

(10) The present invention provides genetically modified non-human animals (e.g., mice, rats, rabbits, etc.) that express both human or humanized MHC I and MHC II proteins; embryos, cells, and tissues comprising the same; methods of making the same; as well as methods of using the same. Unless defined otherwise, all terms and phrases used herein include the meanings that the terms and phrases have attained in the art, unless the contrary is clearly indicated or clearly apparent from the context in which the term or phrase is used.

(11) The term “conservative,” when used to describe a conservative amino acid substitution, includes substitution of an amino acid residue by another amino acid residue having a side chain R group with similar chemical properties (e.g., charge or hydrophobicity). Conservative amino acid substitutions may be achieved by modifying a nucleotide sequence so as to introduce a nucleotide change that will encode the conservative substitution. In general, a conservative amino acid substitution will not substantially change the functional properties of interest of a protein, for example, the ability of MHC I or MHC II to present a peptide of interest. Examples of groups of amino acids that have side chains with similar chemical properties include aliphatic side chains such as glycine, alanine, valine, leucine, and isoleucine; aliphatic-hydroxyl side chains such as serine and threonine; amide-containing side chains such as asparagine and glutamine; aromatic side chains such as phenylalanine, tyrosine, and tryptophan; basic side chains such as lysine, arginine,

and histidine; acidic side chains such as aspartic acid and glutamic acid; and, sulfur-containing side chains such as cysteine and methionine. Conservative amino acids substitution groups include, for example, valine/leucine/isoleucine, phenylalanine/tyrosine, lysine/arginine, alanine/valine, glutamate/aspartate, and asparagine/glutamine. In some embodiments, a conservative amino acid substitution can be a substitution of any native residue in a protein with alanine, as used in, for example, alanine scanning mutagenesis. In some embodiments, a conservative substitution is made that has a positive value in the PAM250 log-likelihood matrix disclosed in Gonnet et al. ((1992) Exhaustive Matching of the Entire Protein Sequence Database, Science 256:1443-45), hereby incorporated by reference. In some embodiments, the substitution is a moderately conservative substitution wherein the substitution has a nonnegative value in the PAM250 log-likelihood matrix. (12) Thus, also encompassed by the invention is a genetically modified non-human animal whose genome comprises a nucleotide sequence encoding a human or humanized MHC I and II polypeptides, wherein MHC I or MHC II the polypeptide comprises conservative amino acid substitutions in the amino acid sequence described herein.

(13) One skilled in the art would understand that in addition to the nucleic acid residues encoding a human or humanized MHC I or MHC II polypeptide described herein, due to the degeneracy of the genetic code, other nucleic acids may encode the polypeptide of the invention. Therefore, in addition to a genetically modified non-human animal that comprises in its genome a nucleotide sequence encoding MHC I and MHC II polypeptides with conservative amino acid substitutions, a non-human animal whose genome comprises a nucleotide sequence that differs from that described herein due to the degeneracy of the genetic code is also provided.

(14) The term “identity” when used in connection with sequence includes identity as determined by a number of different algorithms known in the art that can be used to measure nucleotide and/or amino acid sequence identity. In some embodiments described herein, identities are determined using a ClustalW v. 1.83 (slow) alignment employing an open gap penalty of 10.0, an extend gap penalty of 0.1, and using a Gonnet similarity matrix (MacVector™ 10.0.2, MacVector Inc., 2008). The length of the sequences compared with respect to identity of sequences will depend upon the particular sequences. In various embodiments, identity is determined by comparing the sequence of a mature protein from its N-terminal to its C-terminal. In various embodiments when comparing a chimeric human/non-human sequence to a human sequence, the human portion of the chimeric human/non-human sequence (but not the non-human portion) is used in making a comparison for the purpose of ascertaining a level of identity between a human sequence and a human portion of a chimeric human/non-human sequence (e.g., comparing a human ectodomain of a chimeric human/mouse protein to a human ectodomain of a human protein).

(15) The terms “homology” or “homologous” in reference to sequences, e.g., nucleotide or amino acid sequences, means two sequences which, upon optimal alignment and comparison, are identical in at least about 75% of nucleotides or amino acids, e.g., at least about 80% of nucleotides or amino acids, e.g., at least about 90-95% nucleotides or amino acids, e.g., greater than 97% nucleotides or amino acids. One skilled in the art would understand that, for optimal gene targeting, the targeting construct should contain arms homologous to endogenous DNA sequences (i.e., “homology arms”); thus, homologous recombination can occur between the targeting construct and the targeted endogenous sequence.

(16) The term “operably linked” refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. As such, a nucleic acid sequence encoding a protein may be operably linked to regulatory sequences (e.g., promoter, enhancer, silencer sequence, etc.) so as to retain proper transcriptional regulation. In addition, various portions of the chimeric or humanized protein of the invention may be operably linked to retain proper folding, processing, targeting, expression, and other functional properties of the protein in the cell. Unless stated otherwise, various domains of the chimeric or humanized protein of the invention are operably linked to each other.

(17) The term “MHC I complex” or the like, as used herein, includes the complex between the MHC I α chain polypeptide and the β 2-microglobulin polypeptide. The term “MHC I polypeptide” or the like, as used herein, includes the MHC I α chain polypeptide alone. The terms “MHC II complex,” “MHC II protein,” or the like, as used herein, include the complex between an MHC II α polypeptide and an MHC II β polypeptide. The term “MHC II α polypeptide” or “MHC II β polypeptide” (or the like), as used herein, includes the MHC II α polypeptide alone or MHC II β polypeptide alone, respectively. Similarly, the terms “HLA-DR4 complex”, “HLA-DR4 protein,” “H-2E complex,” “H-2E” protein,” or the like, refer to complex between α and β polypeptides. Typically, the terms “human MHC” and “HLA” are used interchangeably.

(18) The term “replacement” in reference to gene replacement refers to placing exogenous genetic material at an endogenous genetic locus, thereby replacing all or a portion of the endogenous gene with an orthologous or homologous nucleic acid sequence. As demonstrated in the Examples below, nucleic acid sequence of endogenous MHC locus was replaced by a nucleotide sequence comprising sequences encoding a portion of human MHC I polypeptide, specifically, encoding the extracellular portion of the MHC I polypeptide; as well as portions of human MHC II α and β polypeptides, specifically, encoding the extracellular portions of the MHC II α and β polypeptides.

(19) “Functional” as used herein, e.g., in reference to a functional polypeptide, refers to a polypeptide that retains at least one biological activity normally associated with the native protein. For example, in some embodiments of the invention, a replacement at an endogenous locus (e.g., replacement at an endogenous non-human MHC locus) results in a locus that fails to express a functional endogenous polypeptide, e.g., MHC I or MHC II polypeptide. Likewise, the term “functional” as used herein in reference to functional extracellular domain of a protein, refers to an extracellular domain that retains its functionality, e.g., in the case of MHC I or MHC II, ability to bind an antigen, ability to bind a T cell co-receptor, etc. In some embodiments of the invention, a replacement at the endogenous MHC locus results in a locus that fails to express an extracellular domain (e.g., a functional extracellular domain) of an endogenous MHC while expressing an extracellular domain (e.g., a functional extracellular domain) of a human MHC.

(20) Genetically Modified MHC Animals

(21) In various embodiments, the invention generally provides genetically modified non-human animals that comprise in their genome a nucleotide sequence encoding a human or humanized MHC I and MHC II polypeptides; thus, the animals express a human or humanized MHC I and MHC II polypeptides.

(22) MHC genes are categorized into three classes: class I, class II, and class III, all of which are encoded either on human chromosome 6 or mouse chromosome 17. A schematic of the relative organization of the human and mouse MHC classes is presented in FIGS. 2 and 3, respectively. The MHC genes are among the most polymorphic genes of the mouse and human genomes. MHC polymorphisms are presumed to be important in providing evolutionary advantage; changes in sequence can result in differences in peptide binding that allow for better presentation of pathogens to cytotoxic T cells.

(23) MHC class I protein comprises an extracellular domain (which comprises three domains: α .sub.1, α .sub.2, and α .sub.3), a transmembrane domain, and a cytoplasmic tail. The α .sub.1 and α .sub.2 domains form the peptide-binding cleft, while the α .sub.3 interacts with β 2-microglobulin.

(24) In addition to its interaction with β 2-microglobulin, the α .sub.3 domain interacts with the TCR co-receptor CD8, facilitating antigen-specific activation. Although binding of MHC class I to CD8 is about 100-fold weaker than binding of TCR to MHC class I, CD8 binding enhances the affinity of TCR binding. Wooldridge et al. (2010) MHC Class I Molecules with Superenhanced CD8 Binding Properties Bypass the Requirement for Cognate TCR Recognition and Nonspecifically Activate CTLs, *J. Immunol.* 184:3357-3366. Interestingly, increasing MHC class I binding to CD8 abrogated antigen specificity in CTL activation. Id.

(25) CD8 binding to MHC class I molecules is species-specific; the mouse homolog of CD8, Lyt-2,

was shown to bind H-2 D.sup.d molecules at the 13 domain, but it did not bind HLA-A molecules. Connolly et al. (1988) The Lyt-2 Molecule Recognizes Residues in the Class I 3 Domain in Allogeneic Cytotoxic T Cell Responses, *J. Exp. Med.* 168:325-341. Differential binding was presumably due to CDR-like determinants (CDR1- and CDR2-like) on CD8 that was not conserved between humans and mice. Sanders et al. (1991) Mutations in CD8 that Affect Interactions with HLA Class I and Monoclonal Anti-CD8 Antibodies, *J. Exp. Med.* 174:371-379; Vitiello et al. (1991) Analysis of the HLA-restricted Influenza-specific Cytotoxic T Lymphocyte Response in Transgenic Mice Carrying a Chimeric Human-Mouse Class I Major Histocompatibility Complex, *J. Exp. Med.* 173:1007-1015; and, Gao et al. (1997) Crystal structure of the complex between human CD8aa and HLA-A2, *Nature* 387:630-634. It has been reported that CD8 binds HLA-A2 in a conserved region of the 13 domain (at position 223-229). A single substitution (V245A) in HLA-A reduced binding of CD8 to HLA-A, with a concomitant large reduction in T cell-mediated lysis. Salter et al. (1989), Polymorphism in the $\alpha 3$ domain of HLA-A molecules affects binding to CD8, *Nature* 338:345-348. In general, polymorphism in the $\alpha 3$ domain of HLA-A molecules also affected binding to CD8. Id. In mice, amino acid substitution at residue 227 in H-2D.sup.d affected the binding of mouse Lyt-2 to H-2D.sup.d, and cells transfected with a mutant H-2 D.sup.d were not lysed by CD8+ T cells. Potter et al. (1989) Substitution at residue 227 of H-2 class I molecules abrogates recognition by CD8-dependent, but not CD8-independent, cytotoxic T lymphocytes, *Nature* 337:73-75.

(26) Therefore, due to species specificity of interaction between the MHC class I $\alpha 3$ domain and CD8, an MHC I complex comprising a replacement of an H-2K $\alpha 3$ domain with a human HLA-A2 $\alpha 3$ domain was nonfunctional in a mouse (i.e., in vivo) in the absence of a human CD8. In animals transgenic for HLA-A2, substitution of human $\alpha 3$ domain for the mouse $\alpha 3$ domain resulted in restoration of T cell response. Irwin et al. (1989) Species-restricted interactions between CD8 and the $\alpha 3$ domain of class I influence the magnitude of the xenogeneic response, *J. Exp. Med.* 170:1091-1101; Vitiello et al. (1991), *supra*.

(27) The transmembrane domain and cytoplasmic tail of mouse MHC class I proteins also have important functions. One function of MHC I transmembrane domain is to facilitate modulation by HLA-A2 of homotypic cell adhesion (to enhance or inhibit adhesion), presumably as the result of cross-linking (or ligation) of surface MHC molecules. Wagner et al. (1994) Ligation of MHC Class I and Class II Molecules Can Lead to Heterologous Desensitization of Signal Transduction Pathways That Regulate Homotypic Adhesion in Human Lymphocytes, *J. Immunol.* 152:5275-5287. Cell adhesion can be affected by mAbs that bind at diverse epitopes of the HLA-A2 molecule, suggesting that there are multiple sites on HLA-A2 implicated in modulating homotypic cell adhesion; depending on the epitope bound, the affect can be to enhance or to inhibit HLA-A2-dependent adhesion. Id.

(28) The cytoplasmic tail, encoded by exons 6 and 7 of the MHC I gene, is reportedly necessary for proper expression on the cell surface and for LIR1-mediated inhibition of NK cell cytotoxicity. Gruda et al. (2007) Intracellular Cysteine Residues in the Tail of MHC Class I Proteins Are Crucial for Extracellular Recognition by Leukocyte Ig-Like Receptor 1, *J. Immunol.* 179:3655-3661. A cytoplasmic tail is required for multimerization of at least some MHC I molecules through formation of disulfide bonds on its cysteine residues, and thus may play a role in clustering and in recognition by NK cells. Lynch et al. (2009) Novel MHC Class I Structures on Exosomes, *J. Immunol.* 183:1884-1891.

(29) The cytoplasmic domain of HLA-A2 contains a constitutively phosphorylated serine residue and a phosphorylatable tyrosine, although—in Jurkat cells—mutant HLA-A2 molecules lacking a cytoplasmic domain appear normal with respect to expression, cytoskeletal association, aggregation, and endocytic internalization. Gur et al. (1997) Structural Analysis of Class I MHC Molecules: The Cytoplasmic Domain Is Not Required for Cytoskeletal Association, Aggregation, and Internalization, *Mol. Immunol.* 34(2):125-132. Truncated HLA-A2 molecules lacking the

cytoplasmic domain are apparently normally expressed and associate with $\beta 2$ microglobulin. Id. (30) However, several studies have demonstrated that the cytoplasmic tail is critical in intracellular trafficking, dendritic cell (DC)-mediated antigen presentation, and CTL priming. A tyrosine residue encoded by exon 6 was shown to be required for MHC I trafficking through endosomal compartments, presentation of exogenous antigens, and CTL priming; while deletion of exon 7 caused enhancement of anti-viral CTL responses. Lizée et al. (2003) Control of Dendritic Cross-Presentation by the Major Histocompatibility Complex Class I Cytoplasmic Domain, *Nature Immunol.* 4:1065-73; Basha et al. (2008) MHC Class I Endosomal and Lysosomal Trafficking Coincides with Exogenous Antigen Loading in Dendritic Cells, *PLoS ONE* 3: e3247; and Rodriguez-Cruz et al. (2011) Natural Splice Variant of MHC Class I Cytoplasmic Tail Enhances Dendritic Cell-Induced CD8⁺ T-Cell Responses and Boosts Anti-Tumor Immunity, *PLoS ONE* 6:e22939.

(31) MHC class II complex comprises two non-covalently associated domains: an α chain and a β chain, also referred herein as an α polypeptide and a β polypeptide (FIG. 1, right). The protein spans the plasma membrane; thus it contains an extracellular domain, a transmembrane domain, and a cytoplasmic domain. The extracellular portion of the α chain includes $\alpha 1$ and $\alpha 2$ domains, and the extracellular portion of the β chain includes $\beta 1$ and $\beta 2$ domains. The $\alpha 1$ and $\beta 1$ domains form a peptide-binding cleft on the cell surface. Due to the three-dimensional conformation of the peptide-binding cleft of the MHC II complex, there is theoretically no upper limit on the length of the bound antigen, but typically peptides presented by MHC II are between 13 and 17 amino acids in length.

(32) In addition to its interaction with the antigenic peptides, the peptide-binding cleft of the MHC II molecule interacts with invariant chain (Ii) during the processes of MHC II complex formation and peptide acquisition. The α/β MHC II dimers assemble in the endoplasmic reticulum and associate with Ii chain, which is responsible for control of peptide binding and targeting of the MHC II into endocytic pathway. In the endosome, Ii undergoes proteolysis, and a small fragment of Ii, Class II-associated invariant chain peptide (CLIP), remains at the peptide-binding cleft. In the endosome, under control of HLA-DM (in humans), CLIP is exchanged for antigenic peptides.

(33) MHC II interacts with T cell co-receptor CD4 at the hydrophobic crevice at the junction between $\alpha 2$ and $\beta 2$ domains. Wang and Reinherz (2002) Structural Basis of T Cell Recognition of Peptides Bound to MHC Molecules, *Molecular Immunology*, 38:1039-49. When CD4 and T cell receptor bind the same MHC II molecule complexed with a peptide, the sensitivity of a T cell to antigen is increased, and it requires 100-fold less antigen for activation. See, Janeway's Immunobiology, 7th Ed., Murphy et al. eds., Garland Science, 2008, incorporated herein by reference.

(34) Numerous functions have been proposed for transmembrane and cytoplasmic domains of MHC II. In the case of cytoplasmic domain, it has been shown to be important for intracellular signaling, trafficking to the plasma membrane, and ultimately, antigen presentation. For example, it was shown that T cell hybridomas respond poorly to antigen-presenting cells (APCs) transfected with MHC II β chains truncated at the cytoplasmic domain, and induction of B cell differentiation is hampered. See, e.g., Smiley et al. (1996) Truncation of the class II β -chain cytoplasmic domain influences the level of class II/invariant chain-derived peptide complexes, *Proc. Natl. Acad. Sci. USA*, 93:241-44. Truncation of Class II molecules seems to impair cAMP production. It has been postulated that deletion of the cytoplasmic tail of MHC II affects intracellular trafficking, thus preventing the complex from coming across relevant antigens in the endocytic pathway. Smiley et al. (supra) demonstrated that truncation of class II molecules at the cytoplasmic domain reduces the number of CLIP/class II complexes, postulating that this affects the ability of CLIP to effectively regulate antigen presentation.

(35) It has been hypothesized that, since MHC II clustering is important for T cell receptor (TCR) triggering, if MHC II molecules truncated at the cytoplasmic domain were prevented from binding

cytoskeleton and thus aggregating, antigen presentation to T cells would be affected. Ostrand-Rosenberg et al. (1991) Abrogation of Tumorigenicity by MHC Class II Antigen Expression Requires the Cytoplasmic Domain of the Class II Molecule, *J. Immunol.* 147:2419-22. In fact, it was recently shown that HLA-DR truncated at the cytoplasmic domain failed to associate with the cytoskeleton following oligomerization. El Fakhy et al. (2004) Delineation of the HLA-DR Region and the Residues Involved in the Association with the Cytoskeleton, *J. Biol. Chem.* 279:18472-80. Importantly, actin cytoskeleton is a site of localized signal transduction activity, which can effect antigen presentation. In addition to association with cytoskeleton, recent studies have also shown that up to 20% of all HLA-DR molecules constitutively reside in the lipid rafts of APCs, which are microdomains rich in cholesterol and glycosphingolipids, and that such localization is important for antigen presentation, immune synapse formation, and MHC II-mediated signaling. See, e.g., Dolan et al. (2004) Invariant Chain and the MHC II Cytoplasmic Domains Regulate Localization of MHC Class II Molecules to Lipid Rafts in Tumor Cell-Based Vaccines, *J. Immunol.* 172:907-14. Dolan et al. suggested that truncation of cytoplasmic domain of MHC II reduces constitutive localization of MHC II to lipid rafts.

(36) In addition, the cytoplasmic domain of MHC II, in particular the β chain, contains a leucine residue that is subject to ubiquitination by ubiquitin ligase, membrane-associated RING-CH I (MARCH I), which controls endocytic trafficking, internalization, and degradation of MHC II; and it has been shown that MARCH-mediated ubiquitination ceases upon dendritic cell maturation resulting in increased levels of MHC II at the plasma membrane. Shin et al. (2006) Surface expression of MHC class II in dendritic cells is controlled by regulated ubiquitination, *Nature* 444:115-18; De Gassart et al. (2008) MHC class II stabilization at the surface of human dendritic cells is the result of maturation-dependent MARCH I down-regulation, *Proc. Natl. Acad. Sci. USA* 105:3491-96.

(37) Transmembrane domains of α and β chains of MHC II interact with each other and this interaction is important for proper assembly of class II MHC complex. Cosson and Bonifacio (1992) Role of Transmembrane Domain Interactions in the Assembly of Class II MHC Molecules, *Nature* 258:659-62. In fact, MHC II molecules in which the transmembrane domains of the α and β chains were replaced by the α chain of IL-2 receptor were retained in the ER and were barely detectable at the cell surface. Id. Through mutagenesis studies, conserved Gly residues at the α and β transmembrane domains were found to be responsible for MHC II assembly at the cell surface. Id. Thus, both transmembrane and cytoplasmic domains are crucial for the proper function of the MHC II complex.

(38) In various embodiments, provided herein is a genetically modified non-human animal, e.g., rodent (e.g., mouse or rat) comprising in its genome a nucleotide sequence encoding a human or humanized MHC I polypeptide and a nucleotide sequence encoding human or humanized MHC II protein. The MHC I nucleotide sequence may encode an MHC I polypeptide that is partially human and partially non-human, e.g., chimeric human/non-human MHC I polypeptide, and the MHC II nucleotide sequence may encode an MHC II protein that is partially human and partially non-human, e.g., chimeric human/non-human MHC II protein (e.g., comprising chimeric human/non-human MHC II α and β polypeptides). In some aspects, the animal does not express endogenous MHC I and II polypeptides, e.g., functional endogenous MHC I and II polypeptides.

(39) A genetically modified non-human animal comprising in its genome, e.g., at the endogenous locus, a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide is disclosed in U.S. patent application Ser. Nos. 13/661,159 and 13/793,812, which applications are incorporated herein by reference in their entireties. A genetically modified non-human animal comprising in its genome, e.g., at the endogenous locus, a nucleotide sequence encoding humanized, e.g., chimeric human/non-human MHC II polypeptides is disclosed in U.S. patent application Ser. Nos. 13/661,116 and 13/793,935, which applications are incorporated herein by reference in their entireties. Provided herein is a genetically modified non-human animal that

comprises at its endogenous MHC locus a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide and chimeric human/non-human MHC II polypeptides. It would be difficult to generate such animal by simple breeding of humanized MHC I and MHC II animals due to the close linkage of MHC I and MHC II genes on mouse chromosome 17 and human chromosome 6. Therefore, the present application also provides a novel method for generating genetically modified non-human animals comprising sequences encoding human or humanized MHC I and MHC II polypeptides, e.g., animals in which sequences encoding endogenous MHC I and II polypeptides are replaced by those encoding chimeric human/non-human MHC I and II polypeptides.

(40) Thus, in various embodiments provided herein is a genetically modified non-human animal comprising in its genome, e.g., at endogenous MHC locus, a first nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide, wherein a human portion of the chimeric MHC I polypeptide comprises an extracellular domain of a human MHC I polypeptide; a second nucleotide sequence encoding a chimeric human/non-human MHC II α polypeptide, wherein a human portion of the chimeric MHC II α polypeptide comprises an extracellular domain of a human MHC II α polypeptide; and a third nucleotide sequence encoding a chimeric human/non-human MHC II β polypeptide, wherein a human portion of the chimeric MHC II β polypeptide comprises an extracellular domain of a human MHC II β polypeptide; wherein the non-human animal expresses functional chimeric human/non-human MHC I and MHC II proteins from its endogenous non-human MHC locus. In one embodiment, the first, second, and/or third nucleotide sequences are located at the endogenous non-human MHC locus. In one embodiment, wherein the non-human animal is a mouse, the first, second, and/or third nucleotide sequences are located at the endogenous mouse MHC locus on mouse chromosome 17. In one embodiment, the first nucleotide sequence is located at the endogenous non-human MHC I locus. In one embodiment, the second nucleotide sequence is located at the endogenous non-human MHC II α locus. In one embodiment, the third nucleotide sequence is located at the endogenous non-human MHC II β locus.

(41) In one embodiment, the non-human animal only expresses the chimeric human/non-human MHC I, MHC II α and/or MHC II β polypeptides and does not express endogenous non-human MHC polypeptides (e.g., functional endogenous MHC I, II α and/or II β polypeptides) from the endogenous non-human MHC locus. In one embodiment, the animal described herein expresses a functional chimeric MHC I and a functional chimeric MHC II on the surface of its cells, e.g., antigen presenting cells, etc.

(42) In one embodiment, the chimeric human/non-human MHC I polypeptide comprises in its human portion a peptide binding domain of a human MHC I polypeptide. In one aspect, the human portion of the chimeric polypeptide comprises an extracellular domain of a human MHC I. In this embodiment, the human portion of the chimeric polypeptide comprises an extracellular domain of an α chain of a human MHC I. In one embodiment, the human portion of the chimeric polypeptide comprises $\alpha 1$ and $\alpha 2$ domains of a human MHC I. In another embodiment, the human portion of the chimeric polypeptide comprises $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains of a human MHC I.

(43) In one aspect, a human portion of the chimeric MHC II α polypeptide and/or a human portion of the chimeric MHC II β polypeptide comprises a peptide-binding domain of a human MHC II α polypeptide and/or human MHC II β polypeptide, respectively. In one aspect, a human portion of the chimeric MHC II α and/or β polypeptide comprises an extracellular domain of a human MHC II α and/or β polypeptide, respectively. In one embodiment, a human portion of the chimeric MHC II α polypeptide comprises $\alpha 1$ domain of a human MHC II α polypeptide; in another embodiment, a human portion of the chimeric MHC II α polypeptide comprises $\alpha 1$ and $\alpha 2$ domains of a human MHC II α polypeptide. In an additional embodiment, a human portion of the chimeric MHC II β polypeptide comprises $\beta 1$ domain of a human MHC II β polypeptide; in another embodiment, a human portion of the chimeric MHC II β polypeptide comprises $\beta 1$ and $\beta 2$ domains of a human MHC II β polypeptide.

(44) The human or humanized MHC I polypeptide may be derived from a functional human HLA molecule encoded by any of HLA-A, HLA-B, HLA-C, HLA-E, HLA-F, or HLA-G loci. The human or humanized MHC II polypeptide may be derived from a functional human HLA molecule encoded by any of HLA-DP, -DQ, and -DR loci. A list of commonly used HLA antigens and alleles is described in Shankarkumar et al. ((2004) The Human Leukocyte Antigen (HLA) System, *Int. J. Hum. Genet.* 4(2):91-103), incorporated herein by reference. Shankarkumar et al. also present a brief explanation of HLA nomenclature used in the art. Additional information regarding HLA nomenclature and various HLA alleles can be found in Holdsworth et al. (2009) The HLA dictionary 2008: a summary of HLA-A, -B, -C, -DRB1/3/4/5, and DQB1 alleles and their association with serologically defined HLA-A, —B, —C, -DR, and -DQ antigens, *Tissue Antigens* 73:95-170, and a recent update by Marsh et al. (2010) Nomenclature for factors of the HLA system, 2010, *Tissue Antigens* 75:291-455, both incorporated herein by reference. Thus, the human or humanized MHC I and/or II polypeptides may be derived from any functional human HLA molecules described therein.

(45) Of particular interest are human HLA molecules, specific polymorphic HLA alleles, known to be associated with a number of human diseases, e.g., human autoimmune diseases. In fact, specific polymorphisms in HLA loci have been identified that correlate with development of rheumatoid arthritis, type I diabetes, Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis, Graves' disease, systemic lupus erythematosus, celiac disease, Crohn's disease, ulcerative colitis, and other autoimmune disorders. See, e.g., Wong and Wen (2004) What can the HLA transgenic mouse tell us about autoimmune diabetes?, *Diabetologia* 47:1476-87; Taneja and David (1998) HLA Transgenic Mice as Humanized Mouse Models of Disease and Immunity, *J. Clin. Invest.* 101:921-26; Bakker et al. (2006), A high-resolution HLA and SNP haplotype map for disease association studies in the extended human MHC, *Nature Genetics* 38:1166-72 and Supplementary Information; and International MHC and Autoimmunity Genetics Network (2009) Mapping of multiple susceptibility variants within the MHC region for 7 immune-mediated diseases, *Proc. Natl. Acad. Sci. USA* 106:18680-85. Thus, the human or humanized MHC I and/or II polypeptides may be derived from a human HLA molecule known to be associated with a particular disease, e.g., autoimmune disease.

(46) In one specific aspect, the human or humanized MHC I polypeptide is derived from human HLA-A. In a specific embodiment, the HLA-A polypeptide is an HLA-A2 polypeptide (e.g., and HLA-A2.1 polypeptide). In one embodiment, the HLA-A polypeptide is a polypeptide encoded by an HLA-A*0201 allele, e.g., HLA-A*02:01:01:01 allele. The HLA-A*0201 allele is commonly used amongst the North American population. Although the present Examples describe this particular HLA sequence, any suitable HLA-A sequence is encompassed herein, e.g., polymorphic variants of HLA-A2 exhibited in human population, sequences with one or more conservative or non-conservative amino acid modifications, nucleic acid sequences differing from the sequence described herein due to the degeneracy of genetic code, etc.

(47) In another specific aspect, the human portion of the chimeric MHC I polypeptide is derived from human MHC I selected from HLA-B and HLA-C. In one aspect, it is derived from HLA-B, e.g., HLA-B27. In another aspect, it is derived from HLA-A3, -B7, -Cw6, etc.

(48) In one specific aspect, the human portions of the humanized MHC II α and β polypeptides described herein are derived from human HLA-DR, e.g., HLA-DR4. Typically, HLA-DR α chains are monomorphic, e.g., the α chain of HLA-DR complex is encoded by HLA-DRA gene (e.g., HLA-DR α *01 gene). On the other hand, the HLA-DR β chain is polymorphic. Thus, HLA-DR4 comprises an α chain encoded by HLA-DRA gene and a β chain encoded by HLA-DRB1 gene (e.g., HLA-DR β 1*04 gene). As described herein below, HLA-DR4 is known to be associated with incidence of a number of autoimmune diseases, e.g., rheumatoid arthritis, type I diabetes, multiple sclerosis, etc. In one embodiment of the invention, the HLA-DRA allele is HLA-DR α *01 allele, e.g., HLA-DR α *01:01:01:01. In another embodiment, the HLA-DRB allele is HLA-DR β 1*04, e.g.,

HLA-DRβ1*04:01:01. Although the present Examples describe these particular HLA sequences; any suitable HLA-DR sequences are encompassed herein, e.g., polymorphic variants exhibited in human population, sequences with one or more conservative or non-conservative amino acid modifications, nucleic acid sequences differing from the sequences described herein due to the degeneracy of genetic code, etc.

(49) The human portions of the chimeric MHC II α and/or β polypeptide may be encoded by nucleotide sequences of HLA alleles known to be associated with common human diseases. Such HLA alleles include, but are not limited to, HLA-DRB1*0401, -DRB1*0301, -DQA1*0501, -DQB1*0201, -DRB1*1501, -DRB1*1502, -DQB1*0602, -DQA1*0102, -DQA1*0201, -DQB1*0202, -DQA1*0501, and combinations thereof. For a summary of HLA allele/disease associations, see Bakker et al. (2006), *supra*, incorporated herein by reference.

(50) In one aspect, the non-human portion of a chimeric human/non-human MHC I, MHC II α and/or MHC II β polypeptide(s) comprises transmembrane and/or cytoplasmic domains of an endogenous non-human (e.g., rodent, e.g., mouse, rat, etc.) MHC I, MHC II α and/or MHC II β polypeptide(s), respectively. Thus, the non-human portion of the chimeric human/non-human MHC I polypeptide may comprise transmembrane and/or cytoplasmic domains of an endogenous non-human MHC I polypeptide. The non-human portion of a chimeric MHC II α polypeptide may comprise transmembrane and/or cytoplasmic domains of an endogenous non-human MHC II α polypeptide. The non-human portion of a chimeric human/non-human MHC II β polypeptide may comprise transmembrane and/or cytoplasmic domains of an endogenous non-human MHC II β polypeptide. In one aspect, the non-human animal is mouse, and a non-human portion of the chimeric MHC I polypeptide is derived from a mouse H-2K protein. In one aspect, the animal is a mouse, and non-human portions of the chimeric MHC II α and β polypeptides are derived from a mouse H-2E protein. Thus, a non-human portion of the chimeric MHC I polypeptide may comprise transmembrane and cytoplasmic domains derived from a mouse H-2K, and non-human portions of the chimeric MHC II α and β polypeptides may comprise transmembrane and cytoplasmic domains derived from a mouse H-2E protein. Although specific H-2K and H-2E sequences are contemplated in the Examples, any suitable sequences, e.g., polymorphic variants, conservative/non-conservative amino acid substitutions, etc., are encompassed herein.

(51) A chimeric human/non-human polypeptide may be such that it comprises a human or a non-human leader (signal) sequence. In one embodiment, the chimeric MHC I polypeptide comprises a non-human leader sequence of an endogenous MHC I polypeptide. In one embodiment, the chimeric MHC II α polypeptide comprises a non-human leader sequence of an endogenous MHC II α polypeptide. In one embodiment, the chimeric MHC II β polypeptide comprises a non-human leader sequence of an endogenous MHC II β polypeptide. In an alternative embodiment, the chimeric MHC I, MHC II α and/or MHC II β polypeptide(s) comprises a non-human leader sequence of MHC I, MHC II α and/or MHC II β polypeptide(s), respectively, from another non-human animal, e.g., another rodent or another mouse strain. Thus, the nucleotide sequence encoding the chimeric MHC I, MHC II α and/or MHC II β polypeptide may be operably linked to a nucleotide sequence encoding a non-human MHC I, MHC II α and/or MHC II β leader sequence, respectively. In yet another embodiment, the chimeric MHC I, MHC II α and/or MHC II β polypeptide(s) comprises a human leader sequence of human MHC I, human MHC II α and/or human MHC II β polypeptide, respectively (e.g., a leader sequence of human HLA-A2, human HLA-DRA and/or human HLA-DRβ1*04, respectively).

(52) A chimeric human/non-human MHC I, MHC II α and/or MHC II β polypeptide may comprise in its human portion a complete or substantially complete extracellular domain of a human MHC I, human MHC II α and/or human MHC II β polypeptide, respectively. Thus, a human portion may comprise at least 80%, preferably at least 85%, more preferably at least 90%, e.g., 95% or more of the amino acids encoding an extracellular domain of a human MHC I, human MHC II α and/or human MHC II β polypeptide (e.g., human HLA-A2, human HLA-DRA and/or human HLA-

DRβ1*04). In one example, substantially complete extracellular domain of the human MHC I, human MHC II α and/or human MHC II β polypeptide lacks a human leader sequence. In another example, the chimeric human/non-human MHC I, chimeric human/non-human MHC II α and/or the chimeric human/non-human MHC II β polypeptide comprises a human leader sequence. (53) Moreover, the chimeric MHC I, MHC II α and/or MHC II β polypeptide may be operably linked to (e.g., be expressed under the regulatory control of) endogenous non-human promoter and regulatory elements, e.g., mouse MHC I, MHC II α and/or MHC II β regulatory elements, respectively. Such arrangement will facilitate proper expression of the chimeric MHC I and/or MHC II polypeptides in the non-human animal, e.g., during immune response in the non-human animal.

(54) The genetically modified non-human animal may be selected from a group consisting of a mouse, rat, rabbit, pig, bovine (e.g., cow, bull, buffalo), deer, sheep, goat, chicken, cat, dog, ferret, primate (e.g., marmoset, rhesus monkey). For the non-human animals where suitable genetically modifiable ES cells are not readily available, other methods are employed to make a non-human animal comprising the genetic modification. Such methods include, e.g., modifying a non-ES cell genome (e.g., a fibroblast or an induced pluripotent cell) and employing nuclear transfer to transfer the modified genome to a suitable cell, e.g., an oocyte, and gestating the modified cell (e.g., the modified oocyte) in a non-human animal under suitable conditions to form an embryo.

(55) In one aspect, the non-human animal is a mammal. In one aspect, the non-human animal is a small mammal, e.g., of the superfamily Dipodoidea or Muroidea. In one embodiment, the genetically modified animal is a rodent. In one embodiment, the rodent is selected from a mouse, a rat, and a hamster. In one embodiment, the rodent is selected from the superfamily Muroidea. In one embodiment, the genetically modified animal is from a family selected from Calomyscidae (e.g., mouse-like hamsters), Cricetidae (e.g., hamster, New World rats and mice, voles), Muridae (true mice and rats, gerbils, spiny mice, crested rats), Nesomyidae (climbing mice, rock mice, with-tailed rats, *Malagasy* rats and mice), Platacanthomyidae (e.g., spiny dormice), and Spalacidae (e.g., mole rates, bamboo rats, and zokors). In a specific embodiment, the genetically modified rodent is selected from a true mouse or rat (family Muridae), a gerbil, a spiny mouse, and a crested rat. In one embodiment, the genetically modified mouse is from a member of the family Muridae. In one embodiment, the animal is a rodent. In a specific embodiment, the rodent is selected from a mouse and a rat. In one embodiment, the non-human animal is a mouse.

(56) In a specific embodiment, the non-human animal is a rodent that is a mouse of a C57BL strain selected from C57BL/A, C57BL/An, C57BL/GrFa, C57BL/KaLwN, C57BL/6, C57BL/6J, C57BL/6ByJ, C57BL/6NJ, C57BL/10, C57BL/10ScSn, C57BL/10Cr, and C57BL/Ola. In another embodiment, the mouse is a 129 strain selected from the group consisting of a strain that is 129P1, 129P2, 129P3, 129X1, 129S1 (e.g., 129S1/SV, 129S1/SvIm), 129S2, 129S4, 129S5, 129S9/SvEvH, 129S6 (129/SvEvTac), 129S7, 129S8, 129T1, 129T2 (see, e.g., Festing et al. (1999) Revised nomenclature for strain 129 mice, Mammalian Genome 10:836, see also, Auerbach et al (2000) Establishment and Chimera Analysis of 129/SvEv- and C57BL/6-Derived Mouse Embryonic Stem Cell Lines). In a specific embodiment, the genetically modified mouse is a mix of an aforementioned 129 strain and an aforementioned C57BL/6 strain. In another specific embodiment, the mouse is a mix of aforementioned 129 strains, or a mix of aforementioned BL/6 strains. In a specific embodiment, the 129 strain of the mix is a 129S6 (129/SvEvTac) strain. In another embodiment, the mouse is a BALB strain, e.g., BALB/c strain. In yet another embodiment, the mouse is a mix of a BALB strain and another aforementioned strain.

(57) In one embodiment, the non-human animal is a rat. In one embodiment, the rat is selected from a Wistar rat, an LEA strain, a Sprague Dawley strain, a Fischer strain, F344, F6, and Dark Agouti. In one embodiment, the rat strain is a mix of two or more strains selected from the group consisting of Wistar, LEA, Sprague Dawley, Fischer, F344, F6, and Dark Agouti.

(58) Thus, in one embodiment, the invention relates to a genetically modified mouse that comprises

in its genome a first nucleotide sequence encoding a chimeric human/mouse MHC I, a second nucleotide sequence encoding a chimeric human/mouse MHC II α , and a third nucleotide sequence encoding a chimeric human/mouse MHC II β polypeptides. A human portion of the chimeric MHC I, MHC II α , and MHC II β may comprise an extracellular domain of a human MHC I, MHC II α , and MHC II β , respectively. In one embodiment, the mouse expresses functional chimeric human/mouse MHC I, MHC II α , and MHC II β polypeptides from its endogenous mouse MHC locus. In one embodiment, the mouse does not express functional mouse MHC polypeptides, e.g., functional mouse MHC I, MHC II α , and MHC II β polypeptides, from its endogenous mouse MHC locus.

(59) In one embodiment, a human portion of the chimeric human/mouse MHC I polypeptide comprises a peptide binding domain or an extracellular domain of a human MHC I (e.g., human HLA-A, e.g., human HLA-A2, e.g., human HLA-A2.1). In some embodiments, the mouse does not express a peptide binding or an extracellular domain of an endogenous mouse MHC I polypeptide from its endogenous mouse MHC I locus. The peptide binding domain of the human MHC I may comprise $\alpha 1$ and $\alpha 2$ domains. Alternatively, the peptide binding domain of the human MHC I may comprise $\alpha 1$, $\alpha 2$, and $\alpha 3$ domains. In one aspect, the extracellular domain of the human MHC I comprises an extracellular domain of a human MHC I α chain. In one embodiment, the endogenous mouse MHC I locus is an H-2K (e.g., H-2Kb) locus, and the mouse portion of the chimeric MHC I polypeptide comprises transmembrane and cytoplasmic domains of a mouse H-2K (e.g., H-2Kb) polypeptide. Thus, in one embodiment, the mouse of the invention comprises at its endogenous mouse MHC I locus a nucleotide sequence encoding a chimeric human/mouse MHC I, wherein a human portion of the chimeric polypeptide comprises an extracellular domain of a human HLA-A2 (e.g., HLA-A2.1) polypeptide and a mouse portion comprises transmembrane and cytoplasmic domains of a mouse H-2K (e.g., H-2Kb) polypeptide, and a mouse expresses a chimeric human/mouse HLA-A2/H-2K protein. In other embodiment, the mouse portion of the chimeric MHC I polypeptide may be derived from other mouse MHC I, e.g., H-2D, H-2L, etc.; and the human portion of the chimeric MHC I polypeptide may be derived from other human MHC I, e.g., HLA-B, HLA-C, etc. In one aspect, the mouse does not express a functional endogenous H-2K polypeptide from its endogenous mouse H-2K locus.

(60) In one embodiment, a human portion of the chimeric human/mouse MHC II α polypeptide comprises a human MHC II α peptide binding or extracellular domain and a human portion of the chimeric human/mouse MHC II β polypeptide comprises a human MHC II β peptide binding or extracellular domain. In some embodiments, the mouse does not express a peptide binding or an extracellular domain of endogenous mouse α and/or β polypeptide from an endogenous mouse locus (e.g., H-2A and/or H-2E locus). In some embodiments, the mouse comprises a genome that lacks a gene that encodes a functional MHC class II molecule comprising an H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, H-2E α , and a combination thereof. The peptide-binding domain of the human MHC II α polypeptide may comprise $\alpha 1$ domain and the peptide-binding domain of the human MHC II β polypeptide may comprise a $\beta 1$ domain; thus, the peptide-binding domain of the chimeric MHC II complex may comprise human $\alpha 1$ and $\beta 1$ domains. The extracellular domain of the human MHC II α polypeptide may comprise $\alpha 1$ and $\alpha 2$ domains and the extracellular domain of the human MHC II β polypeptide may comprise $\beta 1$ and $\beta 2$ domains; thus, the extracellular domain of the chimeric MHC II complex may comprise human $\alpha 1$, $\alpha 2$, $\beta 1$ and $\beta 2$ domains. In one embodiment, the mouse portion of the chimeric MHC II complex comprises transmembrane and cytosolic domains of mouse MHC II, e.g. mouse H-2E (e.g., transmembrane and cytosolic domains of mouse H-2E α and β chains). Thus, in one embodiment, the mouse of the invention comprises at its endogenous mouse MHC II locus a nucleotide sequence encoding a chimeric human/mouse MHC II α , wherein a human portion of the chimeric MHC II α polypeptide comprises an extracellular domain derived from an α chain of a human MHC II (e.g., α chain of HLA-DR4) and a mouse portion comprises transmembrane and cytoplasmic domains derived from an α chain of a mouse MHC II (e.g., H-2E);

and a mouse comprises at its endogenous mouse MHC II locus a nucleotide sequence encoding a chimeric human/mouse MHC II β , wherein a human portion of the chimeric MHC II β polypeptide comprises an extracellular domain derived from a β chain of a human MHC II (e.g., β chain of HLA-DR4) and a mouse portion comprises transmembrane and cytoplasmic domains derived from a β chain of a mouse MHC II (e.g., H-2E); wherein the mouse expresses a chimeric human/mouse HLA-DR4/H-2E protein. In other embodiment, the mouse portion of the chimeric MHC II protein may be derived from other mouse MHC II, e.g., H-2A, etc.; and the human portion of the chimeric MHC II protein may be derived from other human MHC II, e.g., HLA-DQ, etc. In one aspect, the mouse does not express functional endogenous H-2A and H-2E polypeptides from their endogenous mouse loci (e.g., the mouse does not express H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea polypeptides).

(61) In a further embodiment, a non-human animal of the invention, e.g., a rodent, e.g., a mouse, comprises at an endogenous β 2 microglobulin locus a nucleotide sequence encoding a human or humanized β 2 microglobulin. β 2 microglobulin or the light chain of the MHC class I complex (also abbreviated " β 2M") is a small (12 kDa) non-glycosylated protein, that functions primarily to stabilize the MHC I α chain. Generation of human or humanized microglobulin animals is described in detail in U.S. patent application Ser. No. 13/661,159, and is incorporated herein by reference. A mouse comprising a humanized MHC locus as described in the present disclosure, and a human or humanized β 2 microglobulin locus as described in U.S. patent application Ser. No. 13/661,159, may be generated by any methods known in the art, e.g., breeding.

(62) Various other embodiments of a genetically modified non-human animal, e.g. rodent, e.g., rat or mouse, would be evident to one skilled in the art from the present disclosure and from the disclosure of U.S. patent application Ser. Nos. 13/661,159 and 13/661,116, incorporated herein by reference.

(63) In various aspects of the invention, the sequence(s) encoding a chimeric human/non-human MHC I and MHC II polypeptides are located at an endogenous non-human MHC locus (e.g., mouse H-2K and/or H-2E locus). In one embodiment, this results in a replacement of an endogenous MHC gene(s) or a portion thereof with a nucleotide sequence(s) encoding a human or humanized MHC I polypeptides. Since the nucleotide sequences encoding MHC I, MHC II α and MHC II β polypeptides are located in proximity to one another on the chromosome, in order to achieve the greatest success in humanization of both MHC I and MHC II in one animal, the MHC I and MHC II loci should be targeted sequentially. Thus, also provided herein are methods of generating a genetically modified non-human animal comprising nucleotide sequences encoding chimeric human/non-human MHC I. MHC II α and MHC II β polypeptides as described herein.

(64) In some embodiments, the method utilizes a targeting construct made using VELOCIGENE® technology, introducing the construct into ES cells, and introducing targeted ES cell clones into a mouse embryo using VELOCIMOUSE® technology, as described in the Examples.

(65) The nucleotide constructs used for generating non-human animals described herein are also provided. In one aspect, the nucleotide construct comprises: 5' and 3' non-human homology arms, a human DNA fragment comprising human MHC gene sequences (e.g., human HLA-A2 or human HLA-DR4 gene sequences), and a selection cassette flanked by recombination sites. In one embodiment, the human DNA fragment is a genomic fragment that comprises both introns and exons of a human MHC gene (e.g., human HLA-A2 or HLA-DR4 gene). In one embodiment, the non-human homology arms are homologous to a non-human MHC locus (e.g., MHC I or MHC II locus). Specific constructs are described in the Examples below (e.g., FIG. 6 for MHC II construct, MAID 1680; FIG. 8 for MHC I construct, MAID 1665), as well as in U.S. application Ser. Nos. 13/661,159 and 13/661,116, incorporated herein by reference.

(66) A selection cassette is a nucleotide sequence inserted into a targeting construct to facilitate selection of cells (e.g., ES cells) that have integrated the construct of interest. A number of suitable selection cassettes are known in the art. Commonly, a selection cassette enables positive selection

in the presence of a particular antibiotic (e.g., Neo, Hyg, Pur, CM, Spec, etc.). In addition, a selection cassette may be flanked by recombination sites, which allow deletion of the selection cassette upon treatment with recombinase enzymes. Commonly used recombination sites are loxP and Frt, recognized by Cre and Flp enzymes, respectively, but others are known in the art. In one embodiment, the selection cassette is located at the 5' end of the human DNA fragment. In another embodiment, the selection cassette is located at the 3' end of the human DNA fragment. In another embodiment, the selection cassette is located within the human DNA fragment. In another embodiment, the selection cassette is located within an intron of the human DNA fragment.

(67) In one embodiment, the 5' and 3' non-human homology arms comprise genomic sequence at 5' and 3' locations, respectively, of an endogenous non-human (e.g., murine) MHC class I or class II gene locus (e.g., 5' of the first leader sequence and 3' of the 03 exon of the mouse MHC I gene, or upstream of mouse H-2Ab1 gene and downstream of mouse H-2Ea gene). In one embodiment, the endogenous MHC class I locus is selected from mouse H-2K, H-2D and H-2L. In a specific embodiment, the endogenous MHC class I locus is mouse H-2K. In one embodiment, the endogenous MHC II locus is selected from mouse H-2E and H-2A. In one embodiment, the engineered MHC II construct allows replacement of both mouse H-2E and H-2A genes.

(68) Thus, in one embodiment, provided herein is a method of generating a genetically engineered non-human animal (e.g., rodent, e.g., rat or mouse) capable of expressing humanized MHC I and II proteins comprising replacing at an endogenous non-human MHC II locus a nucleotide sequence encoding a non-human MHC II complex with a nucleotide sequence encoding a chimeric human/non-human MHC II complex to generate a first non-human animal; and replacing at an endogenous non-human MHC I locus a nucleotide sequence encoding a non-human MHC I polypeptide with a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide to generate a second non-human animal. In one embodiment, the steps of replacing nucleotide sequences comprise homologous recombination in ES cells. In one embodiment, the second non-human animal is generated by homologous recombination in ES cells bearing nucleotide sequences encoding chimeric human/non-human MHC II complex. Alternatively, also provided herein is a method of generating a genetically engineered non-human animal (e.g., rodent, e.g., rat or mouse) capable of expressing humanized MHC I and II proteins comprising replacing at an endogenous non-human MHC I locus a nucleotide sequence encoding a non-human MHC I polypeptide with a nucleotide sequence encoding a chimeric human/non-human MHC I polypeptide to generate a first non-human animal; and replacing at an endogenous non-human MHC II locus a nucleotide sequence encoding a non-human MHC II complex with a nucleotide sequence encoding a chimeric human/non-human MHC II complex to generate a second non-human animal. In such embodiment, the second non-human animal is generated by homologous recombination in ES cells bearing a nucleotide sequence encoding chimeric human/non-human MHC I polypeptide.

(69) Upon completion of gene targeting, ES cells or genetically modified non-human animals are screened to confirm successful incorporation of exogenous nucleotide sequence of interest or expression of exogenous polypeptide. Numerous techniques are known to those skilled in the art, and include (but are not limited to) Southern blotting, long PCR, quantitative PCT (e.g., real-time PCR using TAQMAN®), fluorescence in situ hybridization, Northern blotting, flow cytometry, Western analysis, immunocytochemistry, immunohistochemistry, etc. In one example, non-human animals (e.g., mice) bearing the genetic modification of interest can be identified by screening for loss of mouse allele and/or gain of human allele using a modification of allele assay described in Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis, *Nature Biotech.* 21(6):652-659. Other assays that identify a specific nucleotide or amino acid sequence in the genetically modified animals are known to those skilled in the art.

(70) In one aspect, a cell that expresses a chimeric human/non-human MHC I and MHC II proteins

(e.g., HLA-A2/H-2K and HLA-DR4/H-2E proteins) is provided. In one embodiment, the cell comprises an expression vector comprising a chimeric MHC class I sequence and chimeric MHC class II sequence as described herein. In one embodiment, the cell is selected from CHO, COS, 293, HeLa, and a retinal cell expressing a viral nucleic acid sequence (e.g., a PERC.6™ cell). (71) A chimeric MHC II complex comprising an extracellular domain of HLA-DR4 described herein may be detected by anti-HLA-DR antibodies. Thus, a cell displaying chimeric human/non-human MHC II polypeptide may be detected and/or selected using anti-HLA-DR antibody. The chimeric MHC I complex comprising an extracellular domain of HLA-A2 described herein may be detected using anti-HLA-A, e.g., anti-HLA-A2 antibodies. Thus, a cell displaying a chimeric human/non-human MHC I polypeptide may be detected and/or selected using anti-HLA-A antibody. Antibodies that recognize other HLA alleles are commercially available or can be generated, and may be used for detection/selection.

(72) Although the Examples that follow describe a genetically engineered animal whose genome comprises a replacement of a nucleotide sequence encoding mouse H-2K, and H-2A and H-2E proteins with a nucleotide sequence encoding a chimeric human/mouse HLA-A2/H-2K and HLA-DR4/H-2E protein, respectively, one skilled in the art would understand that a similar strategy may be used to introduce chimeras comprising other human MHC I and II genes (other HLA-A, HLA-B, and HLA-C; and other HLA-DR, HLA-DP and HLA-DQ genes). Such animals comprising multiple chimeric human/non-human (e.g., human/rodent, e.g., human/mouse) MHC I and MHC II genes at endogenous MHC loci are also provided.

(73) In various embodiments, the genetically modified non-human animals described herein make cells, e.g., APCs, with human or humanized MHC I and II on the cell surface and, as a result, present peptides as epitopes for T cells in a human-like manner, because substantially all of the components of the complex are human or humanized. The genetically modified non-human animals of the invention can be used to study the function of a human immune system in the humanized animal; for identification of antigens and antigen epitopes that elicit immune response (e.g., T cell epitopes, e.g., unique human cancer epitopes), e.g., for use in vaccine development; for evaluation of vaccine candidates and other vaccine strategies; for studying human autoimmunity; for studying human infectious diseases; and otherwise for devising better therapeutic strategies based on human MHC expression.

EXAMPLES

(74) The invention will be further illustrated by the following nonlimiting examples. These Examples are set forth to aid in the understanding of the invention but are not intended to, and should not be construed to, limit its scope in any way. The Examples do not include detailed descriptions of conventional methods that would be well known to those of ordinary skill in the art (molecular cloning techniques, etc.). Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is indicated in Celsius, and pressure is at or near atmospheric.

Example 1: Engineering a Chimeric Human/Mouse MHC II Locus and Generation of Chimeric MHC II Mice

Example 1.1: Deletion of the Endogenous MHC Class II H-2A and H-2E Loci

(75) The targeting vector for introducing a deletion of the endogenous MHC class II H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea genes was made using VELOCIGENE® genetic engineering technology (see, e.g., U.S. Pat. No. 6,586,251 and Valenzuela et al., supra). Bacterial Artificial Chromosome (BAC) RP23-458i22 (Invitrogen) DNA was modified to delete the endogenous MHC class II genes H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea.

(76) Briefly, upstream and downstream homology arms were derived by PCR of mouse BAC DNA from locations 5' of the H-2Ab1 gene and 3' of the H-2Ea gene, respectively. As depicted in FIG. 5, these homology arms were used to make a cassette that deleted ~79 kb of RP23-458i22 comprising genes H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea of the MHC class II locus by bacterial

homologous recombination (BHR). This region was replaced with a hygromycin cassette flanked by lox66 and lox71 sites. The final targeting vector from 5' to 3' included a 34 kb homology arm comprising mouse genomic sequence 5' to the H-2Ab1 gene of the endogenous MHC class II locus, a 5' lox66 site, a hygromycin cassette, a 3' lox71 site and a 63 kb homology arm comprising mouse genomic sequence 3' to the H-2Ea gene of the endogenous MHC class II locus (MAID 5111, see FIG. 6).

(77) The BAC DNA targeting vector (described above) was used to electroporate mouse ES cells to create modified ES cells comprising a deletion of the endogenous MHC class II locus. Positive ES cells containing a deleted endogenous MHC class II locus were identified by the quantitative PCR assay using TAQMAN™ probes (Lie and Petropoulos (1998) Curr. Opin. Biotechnology 9:43-48). The upstream region of the deleted locus was confirmed by PCR using primers 5111U F (CAGAACGCCAGGCTGTAAC; SEQ ID NO:1) and 5111U R (GGAGAGCAGGGTCAGTCAAC; SEQ ID NO:2) and probe 5111U P (CACCGCCACTCACAGCTCCTTACA; SEQ ID NO:3), whereas the downstream region of the deleted locus was confirmed using primers 5111D F (GTGGGCACCATCTTCATCATTC; SEQ ID NO:4) and 5111D R (CTTCCTTTCCAGGGTGTGACTC; SEQ ID NO:5) and probe 5111D P (AGGCCTGCGATCAGGTGGCACCT; SEQ ID NO:6). The presence of the hygromycin cassette from the targeting vector was confirmed using primers HYGF (TGCGGCCGATCTTAGCC; SEQ ID NO:7) and HYGR (TTGACCGATTCTTGCGG; SEQ ID NO:8) and probe HYGP (ACGAGCGGGTTTCGGCCCATTC; SEQ ID NO:9). The nucleotide sequence across the upstream deletion point (SEQ ID NO:10) included the following, which indicates endogenous mouse sequence upstream of the deletion point (contained within the parentheses below) linked contiguously to cassette sequence present at the deletion point: (TTTGTAACA AAGTCTACCC AGAGACAGAT GACAGACTTC AGCTCCAATG CTGATTGGTT CCTCACTTGG GACCAACCCT) CTCGAGTACC GTTCGTATAA TGTATGCTAT ACGAAGTTAT ATGCATCCGG GTAGGGGAGG. The nucleotide sequence across the downstream deletion point (SEQ ID NO:11) included the following, which indicates cassette sequence contiguous with endogenous mouse sequence downstream of the deletion point (contained within the parentheses below): CCTCGACCTG CAGCCCTAGG ATAACCTCGT ATAATGTATG CTATACGAAC GGTAGAGCTC (CACAGGCATT TGGGTGGGCA GGGATGGACG GTGACTGGGA CAATCGGGAT GGAAGAGCAT AGAATGGGAG TTAGGGAAGA). Positive ES cell clones were then used to implant female mice using the VELOCIMOUSE® method (described below) to generate a litter of pups containing a deletion of the endogenous MHC class II locus.

(78) Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, e.g., U.S. Pat. No. 7,294,754 and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses, Nature Biotech. 25(1):91-99). Mice bearing a deletion of H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea genes in the endogenous MHC class II locus were identified by genotyping using a modification of allele assay (Valenzuela et al., supra) that detected the presence of the hygromycin cassette and confirmed the absence of endogenous MHC class II sequences.

(79) Mice bearing a deletion of H-2Ab1, H-2Aa, H-2Eb1, H-2Eb2, and H-2Ea genes in the endogenous MHC class II locus can be bred to a Cre deleter mouse strain (see, e.g., International Patent Application Publication No. WO 2009/114400) in order to remove any floxed hygromycin cassette introduced by the targeting vector that is not removed, e.g., at the ES cell stage or in the embryo. Optionally, the hygromycin cassette is retained in the mice.

Example 1.2: Generation of Large Targeting Vector (LTVEC) Comprising Humanized H-2Eb1 and H-2Ea Genes

(80) A targeting vector to introduce humanized MHC II sequences was designed as depicted in FIG. 5. Using VELOCIGENE® genetic engineering technology, Bacterial Artificial Chromosome

(BAC) RP23-458i22 DNA was modified in various steps to: (1) create a vector comprising a functional I-E α exon 1 from BALB/c H-2Ea gene (FIG. 5A); (2) create a vector comprising replacement of exons 2 and 3 of mouse I-E β gene with those of human DR β 1*04 and replacement of exons 2 and 3 of mouse I-E α with those of human DR α 1*01 (FIG. 5B); (3) create a vector carrying exons 2 and 3 of human DR β 1*04 amongst remaining mouse I-E β exons, and exons 2 and 3 of human DR α 1*01 amongst remaining mouse I-E α exons including a functional I-E α exon 1 from BALB/c mouse (step (1) (FIG. 5C); and (4) remove a cryptic splice site in the vector generated in (3) (FIG. 5D).

(81) Specifically, because in the C57Bl/6 mice, the I-E α gene is a pseudogene due to the presence of a non-functional exon 1, first, a vector comprising a functional I-E α exon 1 from BALB/c H-2Ea gene was created (FIG. 5A). RP23-458i22 BAC was modified by bacterial homologous recombination (1.BHR) to replace chloramphenicol resistance gene with that of spectinomycin. The resultant vector was further modified by BHR to replace the entire I-A and I-E coding region with a neomycin cassette flanked by recombination sites (2.BHR). Another round of BHR (3. BHR) with the construct comprising an exon encoding BALB/c I-E α leader (exon 1) and chloramphenicol gene flanked by PI-SceI and I-CeuI restriction sites resulted in a vector comprising a functional BALB/c H-2Ea exon 1.

(82) Independently, in order to generate a vector comprising replacement of exons 2 and 3 of mouse I-E β gene with those of human DR β 1*04 and replacement of exons 2 and 3 of mouse I-E α with those of human DR α 1*01, RP23-458i22 BAC was modified via several homologous recombination steps, 4. BHR—8. BHR (FIG. 5B). The resultant nucleic acid sequence was flanked by PI-SceI/I-CeuI restriction sites to allow ligation into the construct carrying BALB/c I-E α exon 1, mentioned above (FIG. 5C).

(83) The sequence of the final construct depicted in FIG. 5C contained a cryptic splice site at the 3' end of the BALB/c intron. Several BHR steps (11. BHR—12. BHR) followed by a deletion step were performed to obtain the final targeting vector (MAID 1680) that was used to electroporate into ES cells (FIG. 5D).

(84) In detail, the final targeting vector (MAID 1680), from 5' to 3', was comprised of a 5' mouse homology arm consisting of ~26 kb of mouse genomic sequence ending just upstream of the H-2Ab1 gene of the endogenous MHC class II locus; an ~59 kb insert containing the humanized MHC II β chain gene (humanized H-2Eb1 gene) and humanized MHC II α chain gene (humanized H-2Ea gene) and a floxed neomycin cassette; and a 3' mouse homology arm consisting of ~57 kb of mouse genomic sequence beginning just downstream of the H-2Ea gene of the endogenous MHC class II locus. The nucleotide sequence across the junction between the 5' arm and the insert (SEQ ID NO:12) included the following: (TGCTGATTGG TTCCTCACTT GGGACCAACC C) TAAGCTTTA TCTATGTTCGG GTGCGGAGAA AGAGGTAATG AAATGGCACA AGGAGATCAC ACACCCAAAC CAAACTCGCC, where the italicized sequence is a unique PI-SceI site, and mouse genomic sequence in the 5' homology arm is in parentheses. The nucleotide sequence across the junction between the insert and the 3' arm (SEQ ID NO:13) included the following: CACATCAGTG AGGCTAGAAT AAATTAAAAT CGCTAATATG AAAATGGGG (ATTTGTACCT CTGAGTGTGA AGGCTGGGAA GACTGCTTTC AAGGGAC), where the mouse genomic sequence in the 3' homology arm is in parentheses.

(85) Within the ~59 kb insert, the H-2Eb1 gene was modified as follows: a 5136 bp region of H-2Eb1, including the last 153 bp of intron1, exon 2, intron 2, exon 3, and the first 122 bp of intron 3, was replaced with the 3111 bp homologous region of human HLA-DRB1*04, including the last 148 bp of intron 1, exon 2, intron 2, exon 3, and the first 132 bp of intron 3. At the junction between the human and mouse sequences of intron 3, a cassette consisting of a 5' lox2372 site, UbC promoter, neomycin resistance gene, and a 3' lox2372 site, was inserted. The resulting gene encoded a chimeric HLA-DRB1*04/H-2Eb1 protein comprised of the mouse H-2Eb1 leader, the human β 1 and β 2 domains from DRB1*04, and the mouse transmembrane domain and cytoplasmic

tail. The nucleotide sequence across the mouse/human junction in intron 1 (SEQ ID NO:14) included the following: (TCCATCACTT CACTGGGTAG CACAGCTGTA ACTGTCCAGC CTG) GGTACCGAGC TCGGATCCAC TAGTAACGGC CGCCAGTGTG CTGGAATTC GCCCTTGATC GAGCTCCCTG GGCTGCAGGT GGTGGGCGTT GCGGGTGGGG CCGGTAA, where the italicized sequence is a multiple cloning site introduced during the cloning steps, and the mouse intron 1 sequences are in parentheses. The nucleotide sequence across the junction between the human intron 3 and neomycin cassette (SEQ ID NO:15) included the following: (ATCTCCATCA GAAGGGCACC GGT) ATA ACTT CGTATAAGGT ATCCTATACG AAGTTATATG CATGGCCTCC GCGCCGGGTT, where the 5' lox2372 site is italicized, and human intron 3 sequence is in parentheses. The nucleotide sequence across the junction between the neomycin cassette and mouse intron 3 (SEQ ID NO:16) included the following: ATA ACTTCGT ATAAGGTATC CTATACGAAG TTATCTCGAG (TGGCTTACAG GTAGGTGCGT GAAGCTTCTA CAAGCACAGT TGCCCCCTGG), where the 3' lox2372 site is italicized, and the mouse intron 3 sequence is in parentheses.

(86) Also within the ~59 kb insert, the H-2Ea gene was modified as follows: a 1185 bp region of H-2Ea, including the last 101 bp of intron1, exon 2, intron 2, exon 3, and the first 66 bp of intron 3, was replaced with the 1189 bp homologous region of human HLA-DRA1*01, including the last 104 bp of intron 1, exon 2, intron 2, exon 3, and the first 66 bp of intron 3. As described above, because exon 1 of the C57BL/6 allele of H-2Ea contains a deletion which renders the gene nonfunctional, H-2Ea exon 1 and the remainder of intron 1 were replaced with the equivalent 2616 bp region from the BALB/c allele of H-2Ea, which is functional. The resulting gene encoded a chimeric H-2Ea/HLA-DRA1*01 protein comprised of the mouse H-2Ea leader from BALB/c, the human $\alpha 1$ and $\alpha 2$ domains from DRA1*01, and the mouse transmembrane domain and cytoplasmic tail. The nucleotide sequence across the mouse/human junction in intron 1 (SEQ ID NO:17) included the following: (CTGTTTCTTC CCTAACTCCC ATTCTATGCT CTTCATCCC GA) CCGCGGCCCA ATCTCTCTCC ACTACTTCCT GCCTACATGT ATGTAGGT, where the italicized sequence is a restriction enzyme site introduced during the cloning steps, and the BALB/c intron 1 sequences are in parentheses. The nucleotide sequence across the human/mouse junction in intron 3 (SEQ ID NO:18) included the following: CAAGGTTTCC TCCTATGATG CTTGTGTGAA ACTCGGGGCC GGCC (AGCATTTAAC AGTACAGGGA TGGGAGCACA GCTCAC), where the italicized sequence is a restriction enzyme site introduced during the cloning steps, and the mouse intron 3 sequences are in parentheses. The nucleotide sequence across the C57BL/6-BALB/c junction 5' of exon 1 (SEQ ID NO:19) included the following: (GAAAGCAGTC TTCCCAGCCT TCACACTCAG AGGTACAAAT) CCCCATTTTC ATATTAGCGA 1TTTAATTTA TTCTAGCCTC, where the C57BL/6-specific sequences are in parentheses. The nucleotide sequence across the BALB/c-C57BL/6 junction 3' of exon 1 (SEQ ID NO:20) included the following: TCTTCCCTAA CTCCCATTCT ATGCTCTTCC ATCCCGA CCG CGG (CCCAATC TCTCTCCACT ACTTCTGCC TACATGTATG), where SacII restriction site is italicized, and C57BL/6 sequences are in parenthesis.

Example 1.3: Generation of Humanized MHC II Mice

(87) Simplified diagrams of the strategy for generating humanized MHC II mice using the vector of Example 1.2 are presented in FIG. 6.

(88) Specifically, MAID1680 BAC DNA (described above) was used to electroporate MAID5111 ES cells to create modified ES cells comprising a replacement of the endogenous mouse I-A and I-E loci with a genomic fragment comprising a chimeric human DR4/mouse I-E locus. Positive ES cells containing deleted endogenous I-A and I-E loci replaced by a genomic fragment comprising a chimeric human DR4/mouse I-E locus were identified by a quantitative PCR assay using TAQMANTM probes (Lie and Petropoulos, supra). The insertion of the human DR α sequences was confirmed by PCR using primers hDRA1F (CTGGCGGCTTGAAGAATTTGG; SEQ ID NO:21), hDRA1R (CATGATTTCCAGGTTGGCTTTGTC; SEQ ID NO:22), and probe hDRA1P

(CGATTGCCAGCTTTGAGGCTCAAGG; SEQ ID NO:23). The insertion of the human DR β sequences was confirmed by PCR using primers hDRB1F (AGGCTTGGGTGCTCCACTTG; SEQ ID NO:24), hDRB1R (GACCCTGGTGATGCTGGAAAC; SEQ ID NO:25), and probe hDRB1P (CAGGTGTAAACCTCTCCACTCCGAGGA; SEQ ID NO:26). The loss of the hygromycin cassette from the targeting vector was confirmed with primers HYGF (TGCGGCCGATCTTAGCC; SEQ ID NO:7) and HYGR (TTGACCGATTCCTTGCGG; SEQ ID NO:8) and probe HYGP (ACGAGCGGGTTCGGCCCATTC; SEQ ID NO:9).

(89) Positive ES cell clones were then used to implant female mice using the VELOCIMOUSE® method (supra) to generate a litter of pups containing a replacement of the endogenous I-A and I-E loci with a chimeric human DR4/mouse I-E locus. Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method. Mice bearing a chimeric human DR4/mouse I-E locus were identified by genotyping using a modification of allele assay (Valenzuela et al., supra) that detected the presence of a chimeric human DR4/mouse I-E locus.

(90) Mice bearing a chimeric human DR4/mouse I-E locus can be bred to a Cre deleter mouse strain (see, e.g., International Patent Application Publication No. WO 2009/114400) in order to remove any floxed neomycin cassette introduced by the targeting vector that is not removed, e.g., at the ES cell stage or in the embryo (See FIG. 7).

Example 2: Engineering a Chimeric Human/Mouse MHC I Locus and Generation of Chimeric MHC I Mice

(91) The mouse H-2K gene was humanized in a single step by construction of a unique targeting vector from human and mouse bacterial artificial chromosome (BAC) DNA using VELOCIGENE® technology (see, e.g., U.S. Pat. No. 6,586,251 and Valenzuela et al. (2003) High-throughput engineering of the mouse genome coupled with high-resolution expression analysis. Nat. Biotech. 21(6): 652-659). DNA from mouse BAC clone RP23-173k21 (Invitrogen) was modified by homologous recombination to replace the genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the mouse H-2K gene with human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ subunits of the human HLA-A gene (FIG. 8).

(92) Briefly, the genomic sequence encoding the mouse the $\alpha 1$, $\alpha 2$ and $\alpha 3$ subunits of the H-2K gene is replaced with the human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the human HLA-A*0201 gene in a single targeting event using a targeting vector comprising a hygromycin cassette flanked by loxP sites with a 5' mouse homology arm containing sequence 5' of the mouse H-2K locus including the 5' untranslated region (UTR; 5' homology arm is set forth in SEQ ID NO:27) and a 3' mouse homology arm containing genomic sequence 3' of the mouse H-2K $\alpha 3$ coding sequence (3' homology arm is set forth in SEQ ID NO:28).

(93) The final construct for targeting the endogenous H-2K gene locus from 5' to 3' included (1) a 5' homology arm containing ~200 bp of mouse genomic sequence 5' of the endogenous H-2K gene including the 5'UTR, (2) ~1339 bp of human genomic sequence including the HLA-A*0201 leader sequence, the HLA-A*0201 leader/ $\alpha 1$ intron, the HLA-A*0201 $\alpha 1$ exon, the HLA-A*0201 $\alpha 1$ - $\alpha 2$ intron, the HLA-A*0201 $\alpha 2$ exon, ~316 bp of the 5' end of the $\alpha 2$ - $\alpha 3$ intron, (3) a 5' loxP site, (4) a hygromycin cassette, (5) a 3' loxP site, (6) ~580 bp of human genomic sequence including ~304 bp of the 3' end of the $\alpha 2$ - $\alpha 3$ intron, the HLA-A*0201 $\alpha 3$ exon, and (7) a 3' homology arm containing ~200 bp of mouse genomic sequence including the intron between the mouse H-2K $\alpha 3$ and transmembrane coding sequences (see FIG. 8 for schematic representation of the H-2K targeting vector). The sequence of 149 nucleotides at the junction of the mouse/human sequences at the 5' of the targeting vector is set forth in SEQ ID NO:29, and the sequence of 159 nucleotides at the junction of the human/mouse sequences at the 3' of the targeting vector is set forth in SEQ ID NO:30. Homologous recombination with this targeting vector created a modified mouse H-2K locus containing human genomic DNA encoding the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the HLA-A*0201 gene operably linked to the endogenous mouse H-2K transmembrane and cytoplasmic domain

coding sequences which, upon translation, leads to the formation of a chimeric human/mouse MHC class I protein.

(94) The targeted BAC DNA was used to electroporate mouse F1H4 ES cells to create modified ES cells for generating mice that express a chimeric MHC class I protein on the surface of nucleated cells (e.g., T and B lymphocytes, macrophages, neutrophils). ES cells containing an insertion of human HLA sequences were identified by a quantitative TAQMAN™ assay. Specific primer sets and probes were designed for detecting insertion of human HLA sequences and associated selection cassettes (gain of allele, GOA) and loss of endogenous mouse sequences (loss of allele, LOA). Table 2 identifies the names and locations detected for each of the probes used in the quantitative PCR assays.

(95) TABLE-US-00001

TABLE	2	Probes	Used	For	Genotyping	Region	Detected	by
SEQ	Probe	Assay	Probe	Sequence	ID	NO	HYG	GOA
								Hygromycin
								cassette
ACGAGCGGGTTCGGCCCATTC	9	1665H1	GOA	Human	HLA-A2	α2-α3		
AGTCCTTCAGCCTCCACTCAGGTCAGG	31	intron	1665H2	GOA	Human	HLA-A2	α2	
exon TACCACCAGTACGCCTACGACGGCA	32	5112H2	GOA	Human	HLA-A2	α2-α3		
ATCCTGTACCAGAGAGTG	33	intron						

(96) The selection cassette may be removed by methods known by the skilled artisan. For example, ES cells bearing the chimeric human/mouse MHC class I locus may be transfected with a construct that expresses Cre in order to remove the “floxed” hygromycin cassette introduced by the insertion of the targeting construct containing human HLA-A*0201 gene sequences (See FIG. 8). The hygromycin cassette may optionally be removed by breeding to mice that express Cre recombinase. Optionally, the hygromycin cassette is retained in the mice.

(97) Targeted ES cells described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, e.g., U.S. Pat. No. 7,294,754 and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses Nature Biotech. 25(1):91-99).

VELOCIMICE® (F0 mice fully derived from the donor ES cell) independently bearing a chimeric MHC class I gene were identified by genotyping using a modification of allele assay (Valenzuela et al., supra) that detects the presence of the unique human HLA-A*0201 gene sequences.

Example 3: Generation and Characterization of Mice Comprising Chimeric MHC I and MHC II Genes

Example 3.1: Generation of Mice Comprising Chimeric MHC I and II Genes

(98) The strategy for generation of mice comprising chimeric MHC I and MHC II genes is depicted in FIG. 4. Specifically, MAID1665 BAC DNA (HLA-A2/H-2K BAC described above in Example 2) was used to electroporate MAID1680 ES cells (ES cells bearing a humanized MHC II gene described in Example 1) to create modified ES cells comprising chimeric human/mouse MHC I and MHC II genes. Positive ES cells containing both chimeric MHC I and II genes were identified by a quantitative PCR assay using TAQMAN™ probes (Lie and Petropoulos, supra) using primers and probes described in Examples 1 and 2 above.

(99) The HYG and NEO selection cassettes may be removed by methods known by the skilled artisan. For example, ES cells bearing the chimeric human/mouse MHC class I and II loci may be transfected with a construct that expresses Cre in order to remove the “floxed” hygromycin and neomycin cassettes introduced by the insertion of the targeting constructs (see FIGS. 4, 7, and 8). The selection cassettes may optionally be removed by breeding to mice that express Cre recombinase. Optionally, the selection cassette is retained in the mice.

(100) Targeted ES cells comprising both humanized MHC I and II described above were used as donor ES cells and introduced into an 8-cell stage mouse embryo by the VELOCIMOUSE® method (see, e.g., U.S. Pat. No. 7,294,754 and Poueymirou et al. (2007) F0 generation mice that are essentially fully derived from the donor gene-targeted ES cells allowing immediate phenotypic analyses Nature Biotech. 25(1):91-99). VELOCIMICE® (F0 mice fully derived from the donor ES

cell) bearing a chimeric MHC class I and class II genes were identified by genotyping using a modification of allele assay (Valenzuela et al., supra).

Example 3.2: Characterization of Mice Comprising Chimeric MHC I and II Genes

(101) Splens from WT or double heterozygous humanized HLA-A2/HLA-DR4 mice (“1666HET/1681 HET” or “H-2K.sup.+ /1666 MHC-II.sup.+ /1681”) were perfused with Collagenase D (Roche Bioscience) and erythrocytes were lysed with ACK lysis buffer. Cell surface expression of human HLA-A2 and HLA-DR4 was analyzed by FACS using fluorochrome-conjugated anti-CD3 (17A2), anti-CD19 (1D3), anti-HLA-A2 (BB7.2) and anti-HLA-DR (L243). Flow cytometry was performed using BD-Fortessa. Expression of both human HLA-A2 and HLA-DR4 were clearly detectable on the surface of CD19⁺ B cells (FIG. 9).

EQUIVALENTS

(102) Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

(103) Entire contents of all non-patent documents, patent applications and patents cited throughout this application are incorporated by reference herein in their entirety.

Claims

1. A genetically modified mouse genome comprising: (i) at an endogenous MHC I locus, a first nucleotide sequence comprising at least a part of a human MHC I gene that encodes $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of a classical human MHC I polypeptide and at least a part of a mouse MHC I gene that encodes transmembrane and cytoplasmic domains of a classical mouse MHC I polypeptide, such that the first nucleotide sequence encodes a chimeric human/mouse MHC I polypeptide that comprises the $\alpha 1$, $\alpha 2$ and $\alpha 3$ domains of the classical human MHC I polypeptide operably linked to the transmembrane and cytoplasmic domains of the classical mouse MHC I polypeptide, (ii) at an endogenous MHC II α locus, a second nucleotide sequence comprising at least a part of a human MHC II α gene that encodes an $\alpha 2$ domain of a classical human MHC II α polypeptide and at least a part of a mouse MHC II α gene that encodes transmembrane and cytoplasmic domains of a classical mouse MHC II α polypeptide, such that the second nucleotide sequence encodes a chimeric human/mouse MHC II α polypeptide that comprises the $\alpha 2$ domain of the classical human MHC II α polypeptide operably linked to the transmembrane and cytoplasmic domains of the classical mouse MHC II α polypeptide, and (iii) at an endogenous MHC II β locus, a third nucleotide sequence comprising at least a part of a human MHC II β gene that encodes a $\beta 2$ domain of a classical human MHC II β polypeptide and at least a part of a mouse MHC II β gene that encodes transmembrane and cytoplasmic domains of a classical mouse MHC II β polypeptide, such that the third nucleotide sequence encodes a chimeric human/mouse MHC II β polypeptide that comprises the $\beta 2$ domain of the classical human MHC II β polypeptide operably linked to the transmembrane and cytoplasmic domains of the classical mouse MHC II β polypeptide.
2. The mouse genome of claim 1, wherein the first, second and/or third nucleotide sequence(s) are operably linked to endogenous mouse regulatory elements.
3. The mouse genome of claim 1, wherein the classical human MHC I polypeptide is HLA-A or HLA-B.
4. The mouse genome of claim 1, further comprising at an endogenous mouse $\beta 2$ microglobulin locus a nucleotide sequence encoding a human or humanized $\beta 2$ microglobulin polypeptide.
5. The mouse genome of claim 1, wherein the first nucleotide sequence is operably linked to endogenous mouse MHC I promoter and regulatory elements at the endogenous MHC I locus, the second nucleotide sequence is operably linked to endogenous mouse MHC II α promoter and regulatory elements at the endogenous MHC II α locus, and the third nucleotide sequence is operably linked to endogenous mouse MHC II β promoter and regulatory elements at the

endogenous MHC II β locus.

6. The mouse genome of claim 1, wherein the classical human MHC II α and β polypeptides respectively are human HLA-DR α and β polypeptides or human HLA-DQ α and β polypeptides.
 7. The mouse genome of claim 1, wherein the classical mouse MHC I polypeptide is H-2K or H-2D.
 8. The mouse genome of claim 1, wherein the classical mouse MHC II α and β polypeptides respectively are H-2A α and β polypeptides.
 9. The mouse genome of claim 1, wherein the classical mouse MHC II α and β polypeptides respectively are H-2E α and β polypeptides.
 10. The mouse genome of claim 1, wherein the first nucleotide sequence encodes a chimeric HLA-A/H-2K polypeptide, the second nucleotide sequence encodes an α chain of a chimeric HLA-DR/H-2E polypeptide, and the third nucleotide sequence encodes a β chain of a chimeric HLA-DR/H-2E polypeptide.
 11. The mouse genome of claim 10, wherein the mouse genome further comprises at an endogenous B2 microglobulin locus a nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide.
 12. The mouse genome of claim 10, wherein the chimeric human/mouse MHC II α polypeptide further comprises an α 1 domain of the classical human MHC II α polypeptide and wherein the chimeric human/mouse MHC II β polypeptide further comprises a β 1 domain of the classical human MHC II β polypeptide.
 13. A mouse cell comprising the mouse genome of claim 12, wherein the mouse cell is an embryonic stem cell and comprises, at an endogenous mouse β 2 microglobulin locus, a nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide.
 14. A mouse cell comprising the mouse genome of claim 12, wherein the mouse cell is an antigen presenting cell (APC), and wherein the APC expresses on its surface the chimeric human/mouse MHC I polypeptide and a chimeric human/mouse MHC II complex comprising the chimeric human/mouse MHC II α and β polypeptides from the endogenous MHC loci.
 15. The mouse cell of claim 14, further comprising at an endogenous mouse β 2 microglobulin locus a nucleotide sequence encoding a human or humanized β 2 microglobulin polypeptide, wherein the APC expresses on its surface the chimeric human or humanized β 2 microglobulin polypeptide.
-