



US012416026B2

(12) **United States Patent**
Graycar et al.

(10) **Patent No.: US 12,416,026 B2**

(45) **Date of Patent: Sep. 16, 2025**

(54) **EXPRESSION OF BETA-GLUCOSIDASE IN YEAST FOR IMPROVED ETHANOL PRODUCTION**

WO 2015042064 A1 3/2015
WO 2015148272 A1 10/2015
WO 2016069541 A1 5/2016
WO 2018089333 A1 5/2018
WO 2018111792 A1 6/2018

(71) Applicant: **DANISCO US INC.**, Palo Alto, CA (US)

OTHER PUBLICATIONS

(72) Inventors: **Thomas P. Graycar**, Pacifica, CA (US); **Kerry Hollands**, Newark, DE (US)

Chica et al. Curr Opin Biotechnol. Aug. 2005;16(4):378-84. (Year: 2005).*

Singh et al. Curr Protein Pept Sci. 2017, 18, 1-11 (Year: 2017).*

Kizer et al. Appl Environ Microbiol. May 2008;74(10):3229-41. (Year: 2008).*

Prather et al. Curr Opin Biotechnol. Oct. 2008;19(5):468-74. (Year: 2008).*

(73) Assignee: **DANISCO US INC.**, Palo Alto, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 598 days.

Accession E3QPM6. Jan. 11, 2011 (Year: 2011).*

Wei et al., "Simultaneous Utilization of Cellobiose, Xylose, and Acetic Acid from Lignocellulosic Biomass for Biofuel Production by an Engineered Yeast Platform", ACS Synthetic Biology, vol. 4, Issue 6, Jan. 14, 2015.

Altschul et al., "Basic Local Alignment Search Tool", J. Mol. Biol., 1990, vol. 215, pp. 403-410.

Altschul et al., "Local Alignment Statistics", Methods in Enzymology, 1996, vol. 266, pp. 460-480.

Devereux et al., "A comprehensive set of sequence analysis programs for the VAX", Nucleic Acids Research, 1984, vol. 12, No. 1, pp. 387-395.

Duskova et al., "Two glycerol uptake systems contribute to the high osmotolerance of *Zygosaccharomyces rouxii*", Molecular Microbiology, Aug. 2015, vol. 97, Issue 3, pp. 541-559.

Eauclaire et al., "Combinatorial metabolic pathway assembly in the yeast genome with RNA-guided Cas9", J. Ind. Microbiol. Biotechnol., 2016, vol. 43, pp. 1001-1015.

Feng et al., "Progressive Sequence Alignment as a Prerequisite to Correct Phylogenetic Trees", J. Mol. Evol., 1987, vol. 25, pp. 351-360.

Ferreira et al., "A Member of the Sugar Transporter Family, Stt1p Is the Glycerol/H⁺ Symporter in *Saccharomyces cerevisiae*", Molecular Biology of the Cell, Apr. 2005, vol. 16, pp. 2068-2076.

Gao et al., "Fast hemicellulose quantification via a simple one-step acid hydrolysis", Biotechnology and Bioengineering, Jun. 2014, vol. 111, Issue 6, pp. 1088-1096.

Gombert et al., "Improving conversion yield of fermentable sugars into fuel ethanol in 1st generation yeast-based production processes", Current Opinion in Biotechnology, 2015, vol. 33, pp. 81-86.

Henikoff et al., "Amino acid substitution matrices from protein blocks", Proc. Natl. Acad. Sci. USA, Nov. 1992, vol. 89, pp. 10915-10919.

(21) Appl. No.: **17/638,250**

(22) PCT Filed: **Aug. 20, 2020**

(86) PCT No.: **PCT/US2020/047150**

§ 371 (c)(1),

(2) Date: **Feb. 25, 2022**

(87) PCT Pub. No.: **WO2021/041141**

PCT Pub. Date: **Mar. 4, 2021**

(65) **Prior Publication Data**

US 2022/0290189 A1 Sep. 15, 2022

Related U.S. Application Data

(60) Provisional application No. 62/893,374, filed on Aug. 29, 2019.

(51) **Int. Cl.**

C12P 7/06 (2006.01)

C12N 9/38 (2006.01)

C12P 19/14 (2006.01)

(52) **U.S. Cl.**

CPC **C12P 7/06** (2013.01); **C12N 9/2471** (2013.01); **C12P 19/14** (2013.01); **C12Y 302/01023** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

8,795,998 B2 8/2014 Pronk et al.
8,956,851 B2 2/2015 Argyros et al.
9,175,270 B2 11/2015 Nevoigt et al.
9,181,566 B2 11/2015 Dauner et al.
2010/0304438 A1 12/2010 Scott et al.

FOREIGN PATENT DOCUMENTS

WO 2008155665 A2 12/2008
WO 2012125951 A1 9/2012
WO 2013090053 A1 6/2013
WO 2015023989 A1 2/2015

(Continued)

Primary Examiner — Christian L Fronda

(57)

ABSTRACT

Described are compositions and methods relating to the expression of β -glucosidase in yeast for use in industrial ethanol production. The yeast demonstrates increased ethanol production and reduced production of unwanted side-products, including acetate. Such yeast is particularly useful for large-scale ethanol production from starch substrates.

12 Claims, No Drawings

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Higgins et al., "Fast and sensitive multiple sequence alignments on a microcomputer", *Cabios Communications*, 1989, vol. 5, No. 2, pp. 151-153.

Karlin et al., "Applications and statistics for multiple high-scoring segments in molecular sequences", *Proc. Natl. Acad. Sci. USA*, Jun. 1993, vol. 90, pp. 5873-5877.

Needleman et al., "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins", *J. Mol. Biol.*, 1970, vol. 48, pp. 443-453.

Pearson et al., "Improved tools for biological sequence comparison", *Proc. Natl. Acad. Sci. USA*, Apr. 1988, vol. 35, pp. 2444-2448.

Petersen et al., "SignalP 4.0: discriminating signal peptides from transmembrane regions", *Nature Methods*, 2011, vol. 8, pp. 785-786.

Sluiter et al., "Determination of Structural Carbohydrates and Lignin in Biomass: Laboratory Analytical Procedure", Technical

Report NREL/TP-510-42618, National Renewable Energy Laboratory, Golden, CO, Apr. 2008, 18 pages.

Smith et al., "Comparison of Biosequences", *Advances in Applied Mathematics*, 1981, vol. 2, pp. 482-489.

Thompson et al., "Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice", *Nucleic Acids Research*, 1994, vol. 22, No. 22, pp. 4673-4680.

Wei et al., "Simultaneous Utilization of Cellobiose, Xylose, and Acetic Acid from Lignocellulosic Biomass for Biofuel Production by an Engineered Yeast Platform", *ACS Synthetic Biology*, Jun. 19, 2015, vol. 4, No. 6, pp. 707-713.

Zhang et al., "Engineering of the glycerol decomposition pathway and cofactor regulation in an industrial yeast Improves ethanol production", *J. Ind. Microbiol. Biotechnol.*, 2013, vol. 40, pp. 1153-1160.

International Search Report from PCT Application No. PCT/US2020/047150 dated Oct. 29, 2020, 5 pages.

* cited by examiner

1

EXPRESSION OF BETA-GLUCOSIDASE IN YEAST FOR IMPROVED ETHANOL PRODUCTION

TECHNICAL FIELD

The present compositions and methods relate to the expression of β -glucosidase in yeast for use in industrial ethanol production. The yeast demonstrates increased ethanol production and reduced production of unwanted side-products, including acetate. Such yeast is particularly useful for large-scale ethanol production from starch substrates.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

The official copy of the sequence listing is submitted electronically via EFS-Web as an ASCII formatted sequence listing with a file named "20190829_NB41307USPSP_SeqLst.txt created on Aug. 29, 2019 and having a size of 62 of kbs kilobytes and is filed concurrently with the specification. The sequence listing contained in this ASCII formatted document is part of the specification and is herein incorporated by reference in its entirety.

BACKGROUND

First-generation yeast-based ethanol production converts sugars into fuel ethanol. The annual fuel ethanol production by yeast is about 90 billion liters worldwide (Gombert, A. K. and van Maris, A. J. (2015) *Curr. Opin. Biotechnol.* 33:81-86). It is estimated that about 70% of the cost of ethanol production is the feedstock. Since the production volume is so large, even small yield improvements have massive economic impact across the industry.

The Renewable Fuel Standard (RFS) is a federal program that requires a minimum volume of renewable fuels to be blended into transportation fuel sold in the United States. The RFS originated with the Energy Policy Act of 2005 and was extended and expanded in the Energy Independence and Security Act of 2007. In 2010 the Environmental Protection Agency (EPA) established a process for companies to petition for new fuel pathways to qualify for the (RFS) program. A fuel pathway is a specific combination of (1) a feedstock, (2) a production process and (3) a fuel type, wherein each combination of three components represents a separate fuel pathway. Qualifying fuel pathways are assigned one or more D-codes corresponding to the type of Renewable Identification Number (RIN) they are eligible to generate. Conventional renewable fuel (e.g., from corn) is D6, advanced biofuel is D5, biodiesel is D4 and cellulosic biofuel is D3 or D7. Cellulosic Biofuel (D-Codes 3 and 7) must be produced from cellulose, hemicellulose or lignin.

RINs are tradable regulatory credits that represent a quantity of qualifying renewable fuel. RINs are assigned after a producer reports the production of a gallon of fuel to the EPA. Blenders demonstrate compliance with the RFS by turning RINs over to the EPA once the gallon of fuel is blended into transportation fuel. Because the RFS requires increasing amounts of advanced biofuels (including cellulosic biofuels) as time progresses, RINs have different values depending on the fuel pathway from which they are generated. For example, a D3 RIN is currently worth more than a D6 RIN.

The current National Renewable Energy Laboratory (NREL) laboratory analytical procedure (LAP) for determi-

2

nation of structural carbohydrates and lignin in biomass is described by Sluiter, A. et al. ((2008) NREL Laboratory Analytical Procedure NREL/TP-510-42618. Golden, CO: National Renewable Energy Laboratory). The method is based on two-step acid hydrolysis, in which biomass is first hydrolyzed using 72 wt % sulfuric acid at 30° C. for 1 h, followed by dilution to 4 wt % sulfuric acid for further hydrolysis at 121° C. for 1 h under autoclave conditions. A faster, single-step method has been described in which biomass is hydrolyzed in 4 wt % sulfuric acid for further hydrolysis at 121° C. for 1 h under autoclave conditions (Gao, X. et al. (2014) *Biotechnology and Bioengineering* 111:1088-96). In both methods, the products of hydrolysis are analyzed by HPLC. A limitation of these methods is that they cannot distinguish glucose derived from starch as opposed to glucose derived from cellulose and, therefore cannot be used to determine the fraction of glucose derived from cellulose in a mixed starch-cellulose feedstock. The EPA established corn kernel fiber as a qualified crop residue on Jul. 18, 2014.

As corn ethanol producers attempt to utilize corn fiber, as well as corn starch, to produce ethanol, there is a financial incentive to characterize as much ethanol as possible as D3 biofuel. However, the EPA requires accuracy in accounting and producers that have non-accurately characterized their biofuel can be subject to penalties. Accordingly, the need exists for an accurate method for determining the source of ethanol when mixed feedstocks of starch and cellulosic components are used to produce biofuels.

SUMMARY

The present compositions and methods relate to modified yeast that expresses β -glucosidase. Aspects and embodiments of the compositions and methods are described in the following, independently-numbered, paragraphs.

1. In one aspect, modified yeast cells derived from parental yeast cells are provided, the modified cells comprising a genetic alteration that causes the modified cells to produce an increased amount of β -glucosidase polypeptides compared to the parental cells, wherein the modified cells produce during fermentation more ethanol and/or less acetate compared to the amount of ethanol and acetate produced by otherwise identical parental yeast cells.

2. In some embodiments of the modified cells of paragraph 1, the genetic alteration comprises the introduction into the parental cells of a nucleic acid capable of directing the expression of a β -glucosidase polypeptide to a level above that of the parental cell grown under equivalent conditions.

3. In some embodiments of the modified cells of paragraph 1, the genetic alteration comprises the introduction of an expression cassette for expressing a β -glucosidase polypeptide.

4. In some embodiments of the modified cells of any of paragraphs 1-3, the cells further comprise an exogenous gene encoding a carbohydrate processing enzyme.

5. In some embodiments, the modified cells of any of paragraphs 1-4 further comprise a PKL pathway.

6. In some embodiments, the modified cells of any of paragraphs 1-5 further comprise an alteration in the glycerol pathway and/or the acetyl-CoA pathway.

7. In some embodiments, the modified cells of any of paragraphs 1-6 further comprise an alternative pathway for making ethanol.

8. In some embodiments of the modified cells of any of paragraphs 1-7, the modified cells further make reduced amounts of DP2 and/or DP3 compared to otherwise identical parental cells.

9. In some embodiments of the modified cells of any of paragraphs 1-8, the β -glucosidase polypeptides are derived from *Glomerella graminicola*.

10. In some embodiments of the modified cells of any of paragraphs 1-9, the β -glucosidase polypeptides have: (a) the amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 2, (b) an amino acid sequence having at least 80% amino acid sequence identity to SEQ ID NO: 1 or SEQ ID NO: 2, (c) the amino acid sequence of a mature polypeptide encoded by any of SEQ ID NOs: 11-16, (d) the amino acid sequence of a mature polypeptide encoded by a nucleic acid having at least 80% nucleic acid identity to any of SEQ ID NOs: 11-16, or (e) the amino acid sequence of a mature polypeptide encoded by a nucleic acid that hybridizes, under stringent condition, to any of SEQ ID NOs: 11-16, or the complement, thereof.

11. In some embodiments of the modified cells of any of paragraphs 1-10, the cells are of a *Saccharomyces* spp.

12. In another aspect, a method for increasing the production of alcohol and/or decreasing the production of acetate from yeast cells grown on a carbohydrate substrate is provided, comprising: introducing into parental yeast cells a genetic alteration that increases the production of β -glucosidase polypeptides compared to the amount produced in the parental cells.

13. In some embodiments of the method of paragraph 12, the cells having the introduced genetic alteration are the modified cells are the cells of any of paragraphs 1-11.

14. In some embodiments of the method of paragraph 12 or 13, the increased production of alcohol is at least 0.5%, at least 1.0%, at least 2.0% or at least 3.0%.

15. In some embodiments of the method of any of paragraphs 12-14, the decreased production of acetate is at least 1.0%, at least 2.0%, at least 4.0% or at least 6.0%.

16. In some embodiments of the method of any of paragraphs 12-15, the cells having the introduced genetic alteration comprise an exogenous PKL pathway.

These and other aspects and embodiments of present modified cells and methods will be apparent from the description, including any accompanying Drawings/Figures.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO: 1 represents the predicted mature amino acid sequence of ABG54.

SEQ ID NO: 2 represents the predicted mature amino acid sequence of FAB, which is the same as SEQ ID NO: 135 of WO2012125951 (Kaper et al.).

SEQ ID NO: 3 represents the predicted mature amino acid sequence of Mg3A.

SEQ ID NO: 4 represents the predicted mature amino acid sequence of TrBGL1.

SEQ ID NO: 5 represents the predicted native signal sequence for ABG54.

SEQ ID NO: 6 represents the predicted native signal sequence for FAB.

SEQ ID NO: 7 represents the predicted native signal sequence for Mg3A.

SEQ ID NO: 8 represents the predicted native signal sequence for TrBGL1.

SEQ ID NO: 9 represents the predicted MFalpha signal sequence.

SEQ ID NO: 10 represents the predicted SUC2 signal sequence.

SEQ ID NO: 11 represents the nucleotide sequence encoding native-ABG54 in pYKH1127, GKH-0464, G3020, G3014, GKH-0737 and GKH-0732.

SEQ ID NO: 12 represents the nucleotide sequence encoding MFalpha-ABG54 in pYKH1139 and GKH-0459.

SEQ ID NO: 13 represents the nucleotide sequence encoding SUC2-ABG54 in GKH-0484.

SEQ ID NO: 14 represents the nucleotide sequence encoding native-FAB in pYKH1135 and GKH-0455.

SEQ ID NO: 15 represents the nucleotide sequence encoding MFalpha-FAB in pYKH1095 and GKH-0450.

SEQ ID NO: 16 represents the nucleotide sequence encoding SUC2-FAB in GKH-0466.

SEQ ID NO: 17 represents the nucleotide sequence encoding native-Mg3A in pYKH1097.

SEQ ID NO: 18 represents the nucleotide sequence encoding MFalpha-Mg3A in pYKH1096.

SEQ ID NO: 19 represents the nucleotide sequence encoding native-TrBGL1 in pYKH1099.

SEQ ID NO: 20 represents the nucleotide sequence encoding MFalpha-TrBGL1 in pYKH1098.

DETAILED DESCRIPTION

I. Definitions

Prior to describing the present yeast and methods in detail, the following terms are defined for clarity. Terms not defined should be accorded their ordinary meanings as used in the relevant art.

As used herein, the term "alcohol" refers to an organic compound in which a hydroxyl functional group (—OH) is bound to a saturated carbon atom.

As used herein, the terms "yeast cells," "yeast strains," or simply "yeast" refer to organisms from the phyla Ascomycota and Basidiomycota. Exemplary yeast is budding yeast from the order Saccharomycetales. Particular examples of yeast are *Saccharomyces* spp., including but not limited to *S. cerevisiae*. Yeast include organisms used for the production of fuel alcohol as well as organisms used for the production of potable alcohol, including specialty and proprietary yeast strains used to make distinctive-tasting beers, wines, and other fermented beverages.

As used herein, the phrase "engineered yeast cells," "variant yeast cells," "modified yeast cells," or similar phrases, refer to yeast that include genetic modifications and characteristics described herein. Variant/modified yeast do not include naturally occurring yeast.

As used herein, the terms "polypeptide" and "protein" (and their respective plural forms) are used interchangeably to refer to polymers of any length comprising amino acid residues linked by peptide bonds. The conventional one-letter or three-letter codes for amino acid residues are used herein and all sequence are presented from an N-terminal to C-terminal direction. The polymer can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art.

5

As used herein, functionally and/or structurally similar proteins are considered to be “related proteins,” or “homologs.” Such proteins can be derived from organisms of different genera and/or species, or different classes of organisms (e.g., bacteria and fungi), or artificially designed. Related proteins also encompass homologs determined by primary sequence analysis, determined by secondary or tertiary structure analysis, or determined by immunological cross-reactivity, or determined by their functions.

As used herein, the term “homologous protein” refers to a protein that has similar activity and/or structure to a reference protein. It is not intended that homologs necessarily be evolutionarily related. Thus, it is intended that the term encompass the same, similar, or corresponding enzyme (s) (i.e., in terms of structure and function) obtained from different organisms. In some embodiments, it is desirable to identify a homolog that has a quaternary, tertiary and/or primary structure similar to the reference protein. In some embodiments, homologous proteins induce similar immunological response(s) as a reference protein. In some embodiments, homologous proteins are engineered to produce enzymes with desired activity(ies).

The degree of homology between sequences can be determined using any suitable method known in the art (see, e.g., Smith and Waterman (1981) *Adv. Appl. Math.* 2:482; Needleman and Wunsch (1970) *J. Mol. Biol.*, 48:443; Pearson and Lipman (1988) *Proc. Natl. Acad. Sci. USA* 85:2444; programs such as GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package (Genetics Computer Group, Madison, WI); and Devereux et al. (1984) *Nucleic Acids Res.* 12:387-95).

For example, PILEUP is a useful program to determine sequence homology levels. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pair-wise alignments. It can also plot a tree showing the clustering relationships used to create the alignment. PILEUP uses a simplification of the progressive alignment method of Feng and Doolittle, (Feng and Doolittle (1987) *J. Mol. Evol.* 35:351-60). The method is similar to that described by Higgins and Sharp ((1989) *CABIOS* 5:151-53). Useful PILEUP parameters including a default gap weight of 3.00, a default gap length weight of 0.10, and weighted end gaps. Another example of a useful algorithm is the BLAST algorithm, described by Altschul et al. ((1990) *J. Mol. Biol.* 215:403-10) and Karlin et al. ((1993) *Proc. Natl. Acad. Sci. USA* 90:5873-87). One particularly useful BLAST program is the WU-BLAST-2 program (see, e.g., Altschul et al. (1996) *Meth. Enzymol.* 266:460-80). Parameters “W,” “T,” and “X” determine the sensitivity and speed of the alignment. The BLAST program uses as defaults a word-length (W) of 11, the BLOSUM62 scoring matrix (see, e.g., Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M⁵, N⁻⁴, and a comparison of both strands.

As used herein, the phrases “substantially similar” and “substantially identical,” in the context of at least two nucleic acids or polypeptides, typically means that a polynucleotide or polypeptide comprises a sequence that has at least about 70% identity, at least about 75% identity, at least about 80% identity, at least about 85% identity, at least about 90% identity, at least about 91% identity, at least about 92% identity, at least about 93% identity, at least about 94% identity, at least about 95% identity, at least about 96% identity, at least about 97% identity, at least about 98% identity, or even at least about 99% identity, or more, compared to the reference (i.e., wild-type) sequence. Percent sequence identity is calculated using CLUSTAL W algo-

6

rithm with default parameters. See Thompson et al. (1994) *Nucleic Acids Res.* 22:4673-4680. Default parameters for the CLUSTAL W algorithm are:

Gap opening penalty:	10.0
Gap extension penalty:	0.05
Protein weight matrix:	BLOSUM series
DNA weight matrix:	IUB
Delay divergent sequences %:	40
Gap separation distance:	8
DNA transitions weight:	0.50
List hydrophilic residues:	GPSNDQEKR
Use negative matrix:	OFF
Toggle Residue specific penalties:	ON
Toggle hydrophilic penalties:	ON
Toggle end gap separation penalty	OFF

Another indication that two polypeptides are substantially identical is that the first polypeptide is immunologically cross-reactive with the second polypeptide. Typically, polypeptides that differ by conservative amino acid substitutions are immunologically cross-reactive. Thus, a polypeptide is substantially identical to a second polypeptide, for example, where the two peptides differ only by a conservative substitution. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions (e.g., within a range of medium to high stringency).

As used herein, the term “hybridization” refers to the process by which one strand of nucleic acid forms a duplex with, i.e., base pairs with, a complementary strand, as occurs during blot hybridization techniques and PCR techniques. Stringent hybridization conditions are exemplified by hybridization under the following conditions: 65° C. and 0.1×SSC (where 1×SSC=0.15 M NaCl, 0.015 M Na₃ citrate, pH 7.0). Hybridized, duplex nucleic acids are characterized by a melting temperature (T_m), where one-half of the hybridized nucleic acids are unpaired with the complementary strand. Mismatched nucleotides within the duplex lower the T_m. A nucleic acid encoding a variant α-amylase may have a T_m reduced by 1° C.-3° C. or more compared to a duplex formed between the nucleotide of SEQ ID NO: 2 and its identical complement.

As used herein, the term “gene” is synonymous with the term “allele” in referring to a nucleic acid that encodes and directs the expression of a protein or RNA. Vegetative forms of filamentous fungi are generally haploid, therefore a single copy of a specified gene (i.e., a single allele) is sufficient to confer a specified phenotype. The term “allele” is generally preferred when an organism contains more than one similar genes, in which case each different similar gene is referred to as a distinct “allele.”

As used herein, the term “expressing a polypeptide” and similar terms refers to the cellular process of producing a polypeptide using the translation machinery (e.g., ribosomes) of the cell.

As used herein, an “expression cassette” refers to a DNA fragment that includes a promoter, and amino acid coding region and a terminator (i.e., promoter::amino acid coding region::terminator) and other nucleic acid sequence needed to allow the encoded polypeptide to be produced in a cell. Expression cassettes can be exogenous (i.e., introduced into a cell) or endogenous (i.e., extant in a cell).

As used herein, the terms “wild-type” and “native” are used interchangeably and refer to genes, proteins or strains found in nature, or that are not intentionally modified for the advantage of the presently described yeast.

As used herein, the term “protein of interest” refers to a polypeptide that is desired to be expressed in modified yeast. Such a protein can be an enzyme, a substrate-binding protein, a surface-active protein, a structural protein, a selectable marker, a signal transducer, a receptor, a transporter, a transcription factor, a translation factor, a co-factor, or the like, and can be expressed. The protein of interest is encoded by an endogenous gene or a heterologous gene (i.e., gene of interest) relative to the parental strain. The protein of interest can be expressed intracellularly or as a secreted protein.

As used herein, the terms “genetic manipulation” and “genetic alteration” are used interchangeably and refer to the alteration/change of a nucleic acid sequence. The alteration can include but is not limited to a substitution, deletion, insertion or chemical modification of at least one nucleic acid in the nucleic acid sequence.

As used herein, a “functional polypeptide/protein” is a protein that possesses an activity, such as an enzymatic activity, a binding activity, a surface-active property, a signal transducer, a receptor, a transporter, a transcription factor, a translation factor, a co-factor, or the like, and which has not been mutagenized, truncated, or otherwise modified to abolish or reduce that activity. Functional polypeptides can be thermostable or thermolabile, as specified.

As used herein, “a functional gene” is a gene capable of being used by cellular components to produce an active gene product, typically a protein. Functional genes are the antithesis of disrupted genes, which are modified such that they cannot be used by cellular components to produce an active gene product, or have a reduced ability to be used by cellular components to produce an active gene product.

As used herein, yeast cells have been “modified to prevent the production of a specified protein” if they have been genetically or chemically altered to prevent the production of a functional protein/polypeptide that exhibits an activity characteristic of the wild-type protein. Such modifications include, but are not limited to, deletion or disruption of the gene encoding the protein (as described, herein), modification of the gene such that the encoded polypeptide lacks the aforementioned activity, modification of the gene to affect post-translational processing or stability, and combinations, thereof.

As used herein, “aerobic fermentation” refers to growth in the presence of oxygen.

As used herein, “anaerobic fermentation” refers to growth in the absence of oxygen.

As used herein, the expression “end of fermentation” refers to the stage of fermentation when the economic advantage of continuing fermentation to produce a small amount of additional alcohol is exceeded by the cost of continuing fermentation in terms of fixed and variable costs. In a more general sense, “end of fermentation” refers to the point where a fermentation will no longer produce a significant amount of additional alcohol, i.e., no more than about 1% additional alcohol, or no more substrate left for further alcohol production.

As used herein, the phrase “degree of polymerization” (DP) refers to the number of anhydroglucopyranose units in a given saccharide. An examples of DP1 is the monosaccharides glucose. Examples of DP2 are the disaccharides maltose and isomaltose.

As used herein, the expression “carbon flux” refers to the rate of turnover of carbon molecules through a metabolic pathway. Carbon flux is regulated by enzymes involved in metabolic pathways, such as the pathway for glucose metabolism and the pathway for maltose metabolism.

As used herein, the singular articles “a,” “an” and “the” encompass the plural referents unless the context clearly dictates otherwise. All references cited herein are hereby incorporated by reference in their entirety. The following abbreviations/acronyms have the following meanings unless otherwise specified:

° C. degrees Centigrade

AA α -amylase

AADH acetaldehyde dehydrogenases

ADH alcohol dehydrogenase

bp base pairs

DNA deoxyribonucleic acid

ds or DS dry solids

DP degree of polymerization

EC enzyme commission

EtOH ethanol

g or gm gram

g/L grams per liter

GA glucoamylase

HPLC high performance liquid chromatography

hr or h hour

M molar

mg milligram

min minute

mL or ml milliliter

mM millimolar

N normal

n/a not applicable

n/d no data

nm nanometer

PCR polymerase chain reaction

PKL phosphoketolase

ppm parts per million

PTA phosphotransacetylase

RPM revolutions per minute

Δ relating to a deletion

μ g microgram

μ L and μ l microliter

μ M micromolar

II. Modified Yeast Cells Expressing β -Glucosidase

Described are modified yeast cells that express β -glucosidase, and methods of use, thereof. The yeast produces increased amounts of ethanol, decreased amounts of acetate, and offer additional advantages compared to conventional yeast. The additional ethanol results from the utilization of non-starch-derived β -D-linked glucan substrates, for example, cellobiose and other cellooligosaccharide substrates, and may qualify for D3 RIN credits. Expression of β -glucosidase in yeast was inspired by the observation that contacting stillage with a cocktail of eight glycosyl hydrolase enzymes and enriched for β -glucosidase, liberated more glucose compared to cocktails enriched for a different glycosyl hydrolase.

Yeast can be selective in their ability to express exogenous proteins, and expression of β -glucosidase is no exception. Expression of β -glucosidase in yeast was not straightforward and suitable yeast were not readily obtained. It is also surprising that the expression of β -glucosidase by an ethanologen, so far downstream of initial starch liquefaction, would lead to the observed benefits in ethanol and acetate production. Expression of β -glucosidase was also found to benefit ethanol and acetate production in combination with an exogenous PKL pathway and in combination with glucoamylase expression.

In some embodiments, the increase in ethanol production by the modified cells is an increase of at least 0.5%, at least 0.7%, at least 0.9%, at least 1.2%, at least 1.5%, at least 2.0%, at least 3.0%, or more, compared to the amount of ethanol produced by parental cells grown under the same conditions.

In some embodiments, the decrease in acetate production by the modified cells is an decrease of at least 0.5%, at least 1.0%, at least 2.0%, at least 3.0%, at least 4.0%, at least 5.0%, at least 6.0%, or more, compared to the amount of acetate produced by parental cells grown under the same conditions.

Preferably, increased β -glucosidase expression is achieved by genetic manipulation using sequence-specific molecular biology techniques, as opposed to chemical mutagenesis, which is generally not targeted to specific nucleic acid sequences. However, chemical mutagenesis is not excluded as a method for making modified yeast cells.

In some embodiments, the present compositions and methods involve introducing into yeast cells a nucleic acid capable of directing the expression, or over-expression, of a β -glucosidase polypeptide. Particular methods include but are not limited to (i) introducing an exogenous expression cassette for producing the polypeptide into a host cell, optionally in addition to an endogenous expression cassette, (ii) substituting an exogenous expression cassette with an endogenous cassette that allows the production of an increased amount of the polypeptide, (iii) modifying the promoter of an endogenous expression cassette to increase expression, (iv) increase copy number of the same or different cassettes for over-expression of β -glucosidase, and/or (v) modifying any aspect of the host cell to increase the half-life of the polypeptide in the host cell.

In some embodiments, the parental cell that is modified already includes a gene of interest, such as a gene encoding a selectable marker, carbohydrate-processing enzyme, or other polypeptide. In some embodiments, a gene of interest is subsequently introduced into the modified cells.

In some embodiments, the parental cell that is modified already includes an engineered pathway of interest, such as a PKL pathway to increase ethanol production, or any other pathway to increase alcohol production.

In some embodiments of the present compositions and methods, the amino acid sequence of the β -glucosidase polypeptide that is expressed in modified yeast cells has at least about 80%, at least about 85%, at least about 87%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or even at least about 99% identity, to SEQ ID NO: 1 or SEQ ID NO: 2.

In some embodiments of the present compositions and methods, β -glucosidase polypeptide has an the amino acid sequence of a mature polypeptide encoded by a nucleic acid having at least about 80%, at least about 85%, at least about 87%, at least about 90%, at least about 91%, at least about 92%, at least about 93%, at least about 94%, at least about 95%, at least about 96%, at least about 97%, at least about 98%, or even at least about 99% identity nucleic acid identity to any of SEQ ID NOs: 11-16, or the amino acid sequence of a mature polypeptide encoded by a nucleic acid that hybridizes, under stringent condition, to any of SEQ ID NOs: 11-16, or the complement, thereof.

III. Modified Yeast Cells Having Increased β -Glucosidase Expression in Combination with Genes of an Exogenous PKL Pathway

Increased expression of β -glucosidase can be combined with expression of genes in the PKL pathway to further

increase ethanol production. Engineered yeast cells having a heterologous PKL pathway have been previously described in WO2015148272 (Miasnikov et al.). These cells express heterologous phosphoketolase (PKL), phosphotransacetylase (PTA) and acetylating acetyl dehydrogenase (AADH), optionally with other enzymes, to channel carbon flux away from the glycerol pathway and toward the synthesis of acetyl-CoA, which is then converted to ethanol. Such modified cells are capable of increased ethanol production in a fermentation process when compared to otherwise-identical parent yeast cells.

IV. Combination of Increased β -Glucosidase Production with Other Mutations that Affect Alcohol Production

In some embodiments, in addition to expressing increased amounts of β -glucosidase polypeptides, optionally in combination with introducing an exogenous PKL pathway, the present modified yeast cells include additional beneficial modifications.

The modified cells may further include mutations that result in attenuation of the native glycerol biosynthesis pathway and/or reuse glycerol pathway, which are known to increase alcohol production. Methods for attenuation of the glycerol biosynthesis pathway in yeast are known and include reduction or elimination of endogenous NAD-dependent glycerol 3-phosphate dehydrogenase (GPD) or glycerol phosphate phosphatase activity (GPP), for example by disruption of one or more of the genes GPD1, GPD2, GPP1 and/or GPP2. See, e.g., U.S. Pat. No. 9,175,270 (Elke et al.), U.S. Pat. No. 8,795,998 (Pronk et al.) and U.S. Pat. No. 8,956,851 (Argyros et al.). Methods to enhance the reuse glycerol pathway by over expression of glycerol dehydrogenase (GCY1) and dihydroxyacetone kinase (DAK1) to convert glycerol to dihydroxyacetone phosphate (Zhang et al. (2013) *J. Ind. Microbiol. Biotechnol.* 40:1153-60).

The modified yeast may further feature increased acetyl-CoA synthase (also referred to acetyl-CoA ligase) activity (EC 6.2.1.1) to scavenge (i.e., capture) acetate produced by chemical or enzymatic hydrolysis of acetyl-phosphate (or present in the culture medium of the yeast for any other reason) and converts it to Ac-CoA. This partially reduces the undesirable effect of acetate on the growth of yeast cells and may further contribute to an improvement in alcohol yield. Increasing acetyl-CoA synthase activity may be accomplished by introducing a heterologous acetyl-CoA synthase gene into cells, increasing the expression of an endogenous acetyl-CoA synthase gene and the like.

In some embodiments the modified cells may further include a heterologous gene encoding a protein with NAD⁺-dependent acetylating acetaldehyde dehydrogenase activity and/or a heterologous gene encoding a pyruvate-formate lyase. The introduction of such genes in combination with attenuation of the glycerol pathway is described. e.g., in U.S. Pat. No. 8,795,998 (Pronk et al.). In some embodiments of the present compositions and methods the yeast expressly lacks a heterologous gene(s) encoding an acetylating acetaldehyde dehydrogenase, a pyruvate-formate lyase or both.

In some embodiments, the present modified yeast cells may further over-express a sugar transporter-like (STL) polypeptide to increase the uptake of glycerol (see. e.g., Ferreira et al. (2005) *Mol. Biol. Cell.* 16:2068-76; Dušková et al. (2015) *Mol. Microbiol.* 97:541-59 and WO 2015023989 A1) to increase ethanol production and reduce acetate.

11

In some embodiments, the present modified yeast cells further include a butanol biosynthetic pathway. In some embodiments, the butanol biosynthetic pathway is an isobutanol biosynthetic pathway. In some embodiments, the isobutanol biosynthetic pathway comprises a polynucleotide encoding a polypeptide that catalyzes a substrate to product conversion selected from the group consisting of: (a) pyruvate to acetolactate; (b) acetolactate to 2,3-dihydroxyisovalerate; (c) 2,3-dihydroxyisovalerate to 2-ketoisovalerate; (d) 2-ketoisovalerate to isobutyraldehyde; and (e) isobutyraldehyde to isobutanol. In some embodiments, the isobutanol biosynthetic pathway comprises polynucleotides encoding polypeptides having acetolactate synthase, keto acid reductoisomerase, dihydroxy acid dehydratase, ketoisovalerate decarboxylase, and alcohol dehydrogenase activity.

In some embodiments, the modified yeast cells comprising a butanol biosynthetic pathway further comprise a modification in a polynucleotide encoding a polypeptide having pyruvate decarboxylase activity. In some embodiments, the yeast cells comprise a deletion, mutation, and/or substitution in an endogenous polynucleotide encoding a polypeptide having pyruvate decarboxylase activity. In some embodiments, the polypeptide having pyruvate decarboxylase activity is selected from the group consisting of: PDC1, PDC5, PDC6, and combinations thereof. In some embodiments, the yeast cells further comprise a deletion, mutation, and/or substitution in one or more endogenous polynucleotides encoding FRA2, ALD6, ADH1, GPD2, BDH1, DLS1, DPB3, CPR1, MAL23C, MNN4, PAB1, TMN2, HAC1, PTC1, PTC2, OSM1, GIS1, CRZ1, HUG1, GDS1, CYB2P, SFC1, MVB12, LDB10, C5SD, GIC1, GIC2 and/or YMR226C.

V. Combination of Increased Expression β-Glucosidase with Other Beneficial Mutations

In some embodiments, in addition to increased expression of β-glucosidase polypeptides, optionally in combination with other genetic modifications that benefit alcohol production and/or acetate reduction, the present modified yeast cells further include any number of additional genes of interest encoding proteins of interest. Additional genes of interest may be introduced before, during, or after genetic manipulations that result in the increased production of β-glucosidase polypeptides. Proteins of interest, include selectable markers, carbohydrate-processing enzymes, and other commercially-relevant polypeptides, including but not limited to an enzyme selected from the group consisting of a dehydrogenase, a transketolase, a phosphoketolase, a transaldolase, an epimerase, a phytase, a xylanase, a β-glucanase, a phosphatase, a protease, an α-amylase, a β-amylase, a glucoamylase, a pullulanase, an isoamylase, a cellulase, a trehalase, a lipase, a pectinase, a polyesterase, a cutinase, an oxidase, a transferase, a reductase, a hemicellulase, a mannanase, an esterase, an isomerase, a pectinase, a lactase, a peroxidase and a laccase. Proteins of interest may be secreted, glycosylated, and otherwise-modified.

VI. Use of the Modified Yeast for Increased Alcohol Production

The present compositions and methods include methods for increasing alcohol production and/or reducing glycerol production, in fermentation reactions. Such methods are not limited to a particular fermentation process. The present engineered yeast is expected to be a “drop-in” replacement

12

for convention yeast in any alcohol fermentation facility. While primarily intended for fuel alcohol production, the present yeast can also be used for the production of potable alcohol, including wine and beer.

VII. Yeast Cells Suitable for Modification

Yeasts are unicellular eukaryotic microorganisms classified as members of the fungus kingdom and include organisms from the phyla Ascomycota and Basidiomycota. Yeast that can be used for alcohol production include, but are not limited to, *Saccharomyces* spp., including *S. cerevisiae*, as well as *Kluyveromyces*, *Lachancea* and *Schizosaccharomyces* spp. Numerous yeast strains are commercially available, many of which have been selected or genetically engineered for desired characteristics, such as high alcohol production, rapid growth rate, and the like. Some yeasts have been genetically engineered to produce heterologous enzymes, such as glucoamylase or α-amylase.

VII. Substrates and Products

Alcohol production from a number of carbohydrate substrates, including but not limited to corn starch, sugar cane, cassava, and molasses, is well known, as are innumerable variations and improvements to enzymatic and chemical conditions and mechanical processes. The present compositions and methods are believed to be fully compatible with such substrates and conditions.

Alcohol fermentation products include organic compound having a hydroxyl functional group (—OH) is bound to a carbon atom. Exemplary alcohols include but are not limited to methanol, ethanol, n-propanol, isopropanol, n-butanol, isobutanol, n-pentanol, 2-pentanol, isopentanol, and higher alcohols. The most commonly made fuel alcohols are ethanol, and butanol.

These and other aspects and embodiments of the present yeast strains and methods will be apparent to the skilled person in view of the present description. The following examples are intended to further illustrate, but not limit, the compositions and methods.

EXAMPLES

Example 1: Addition of β-Glucosidases to Stillage

Fermentation broth from submerged fermentation cultures expressing eight different glycosyl hydrolase enzymes, including ABG54 β-glucosidase, was prepared for analysis by filtration to remove cell mass using 96-well Millipore filter plates. Filtration was followed by exchange of each enzyme from culture broth into 20 mM sodium acetate buffer, pH 5.0, 0.005% TWEEN®-80, using 96-well ZEBATM Spin desalting plates (ThermoFisher, Cat. No.: 89807).

An Agilent HPLC 1290 INFINITY™ system was used to quantify the protein obtained from each culture using a Waters ACQUITY UPLC® C4BEH 300 column (1.7 μm, 1×50 mm). A six-minute program was used starting with an initial gradient from 5% to 33% acetonitrile (Sigma-Aldrich) in 0.5 minutes, followed by a gradient from 33% to 48% acetonitrile in 4.5 minutes, and then a step gradient to 90% acetonitrile. A protein standard curve based on purified *Trichoderma reesei* Bgl1 (referred to as TrBGL1, below) was used to quantify the enzyme sample preparations. Prior to running enzyme samples on the HPLC system all the proteins were first deglycosylated using endoglycosidase-H.

13

Whole stillage sampled from a grain ethanol production plant was centrifuged and the resulting supernatant passed through a 0.2 μ m membrane filter. A 20% volume of 100 mM sodium acetate buffer, pH 5.0, with 0.005% Tween 80 was added to the whole stillage substrate for enzyme activity assays. Glycosyl hydrolase enzymes were added to the filtered whole stillage supernatant to identify enzymes capable of releasing glucose from the soluble recalcitrant oligosaccharides.

Glycosyl hydrolase reactions with the soluble recalcitrant oligosaccharide substrate were performed in 96-well microtiter plates containing 150 μ l of the buffered whole stillage supernatant, a 3 μ g protein blend of all eight glycosyl hydrolases, and 6 μ g of each individual glycosyl hydrolase combined with the protein blend. Reaction plates were sealed, and incubated at 32° C. for 18 hours with 250 RPM mixing. Immediately following the incubation, the plates were unsealed and 150 μ l of 0.1 N sulfuric acid was mixed into the well contents to quench the hydrolysis reactions. The amount of glucose product generated by the enzymatic hydrolysis of recalcitrant oligosaccharides was measured using an ABTS assay for glucose.

An ABTS stock solution containing 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulfonic acid) di-ammonium salt (Sigma Aldrich) at 2.88 mg/ml, horseradish peroxidase (Sigma Aldrich) at 0.11 U/ml and glucose oxidase (OXYGO™ HP 5000L, DuPont) at 1.05 U/ml was prepared in 50 mM sodium phosphate buffer, pH 7. 95 μ l of the ABTS stock solution was transferred into the wells of a 96-well microtiter plate, 5 μ l of the quenched reaction plates was pipetted into the assay plates containing the ABTS solution. Assay plates were loaded into a microtiter plate reader set for a 3-min kinetic measurement at an absorbance setting of 405 m with a 9-sec read interval and a 60-second lag. A 5-sec shaking step was used prior to the kinetic measurement for mixing and to eliminate any air bubbles in the assay plate wells. The amount of glucose produced for each sample reaction was calculated using a standard curve generated with glucose standards and the same ABTS assay. The results are shown in Table 1. The enzyme composition enriched for Enzyme H (the β -glucosidase ABG54) produced the most glucose from a stillage substrate.

TABLE 1

Production of glucose from recalcitrant soluble oligosaccharides in whole stillage	
Enzyme	Glucose produced (mM)
Protein blend (PB)	7.99
PB + Enzyme A	8.37
PB + Enzyme B	8.16
PB + Enzyme C	8.17
PB + Enzyme D	8.15
PB + Enzyme E	8.22
PB + Enzyme F	8.08
PB + Enzyme G	8.10
PB + Enzyme H	9.65

Example 2: Materials and Methods for Examples 2-8

Liquefact Preparation

Liquefact (Ground corn slurry) was prepared by adding 600 ppm of urea, 0.124 SAPU/g ds FERMGENT™ 2.5x(acid fungal protease), 0.33 GAU/g ds variant *Trichoderma* glucoamylase (TrGA) and 1.46 SSCU/g ds *Aspergillus* α -amy-

14

lase (AkAA), adjusted to a pH of 4.8. For evaluation of strains expressing glucoamylase, the dose of glucoamylase was reduced to 0.1 GAU/g ds. In certain experiments, 5 g/L cellobiose was also added to the liquefact.

Serum Vial Assays

2 mL of YPD in 24-well plates were inoculated with yeast cells and the cells allowed to grow overnight, 5 ml prepared liquefact was aliquoted to serum vials (Chemglass, Catalog No.: CG-4904-01) and yeast was added to each vial for a final OD of about 0.2-0.4. The lids of the vials were screw on and punctured with a needle (BD, Catalog No.: 305111) for ventilation (to release CO₂), then incubated at 32° C. with shaking (200 RPM) for 55 hours.

Shake Flask Assays

100 μ l of concentrated yeast overnight culture was added to each of a number of shake flasks filled with 50 g prepared liquefact for a final OD of 0.3. The flasks were incubated at 32° C. with shaking (200 RPM) for 55 hours.

HPLC Analysis

Samples from serum vial and shake flask experiments were filtered through 0.2 μ m PTFE filters and the filtrates analyzed for acetate, ethanol, glycerol, glucose, DP2, DP3 and DP4+ content by HPLC (Waters e2695 series) using Bio-Rad Aminex HPX-87H columns at 65° C., with an isocratic flow rate of 0.6 ml/min in 0.01 N H₂SO₄ eluent. A 2.5 μ l sample injection volume was used. Calibration standards were used for quantification of acetate, ethanol, glycerol and glucose. Values are expressed in g/L.

Growth of Strains for β -Glucosidase Activity Determination

Yeast strains were inoculated into 2 ml synthetic complete medium lacking uracil (SC-ura) or YPD medium in 24-well plates and the cells allowed to grow overnight. Cultures were filtered through a 0.2 μ m filter plate (PALL AcroPrep Advance, GHP membrane) and the resulting filtrates assayed for β -glucosidase activity.

β -Glucosidase Activity Assay

A 4-nitrophenyl- β -D-glucopyranoside (pNPG) substrate solution was prepared by dissolving 0.006 g pNPG in 20 ml 0.05 M sodium acetate buffer (0.05 M sodium acetate, 0.1% (v/v) Polyethylene glycerol 8000, pH 4.8). 100 μ l pNPG substrate solution was mixed with 20 μ l filtered culture supernatant or β -glucosidase standard and incubated for 45 min at 30°C. The reaction was stopped by adding 50 μ l 1 M sodium carbonate and incubating for 1 min at room temperature, then absorbance at 405 m was measured. β -glucosidase activity in culture supernatants was calculated by comparison to a standard curve generated using serially diluted β -glucosidase standard.

Signal Sequence Prediction

The determination of signal sequence cleavage sites was predicted using SignalP version 4.1 (Petersen, T. N. et al. (2011) Nature Methods, 8:785-86) with default parameters. It will be understood that actual signal sequence cleavage sites, and therefore the N-termini of mature secreted polypeptides, may differ by a few amino acid residues or may vary within a population of secreted polypeptides.

Example 3: Screening for Expression of β -Glucosidases in *S. cerevisiae*

Genes coding for four different β -glucosidases (SEQ ID NOs: 1-4), codon optimized for *Saccharomyces cerevisiae*, were cloned between the SpeI and NotI restriction sites in pJT257 (described in U.S. Pat. No. 9,181,566). In the resulting plasmids, expression of the β -glucosidase is under control of the FBA1 promoter and FBA1 terminator from *S. cerevisiae* (see, e.g., WO2018/111792). A second set of

15

plasmids was constructed containing genes coding for the same four β -glucosidases but with the native signal sequence (SEQ ID NOs: 5-8) removed and replaced with the signal sequence from the *S. cerevisiae* mating factor α (MFalpha; SEQ ID NO: 9). Plasmids were designated as show in Table 2. Note that FAB is hybrid enzyme derived from portions of the β -glucosidases from *Fusarium vesticilloles*, *Rasamsonia emersonii* and *Hypocrea (Trichoderma) jecorina*, which is described in WO2012125951 (Kaper et al.) as SEQ ID NO: 135.

TABLE 2

β -glucosidase expression plasmids					
Plasmid name	β -glucosidase	SEQ ID NO	Signal sequence	SEQ ID NO	Source organism
pYKH1127	ABG54	1	native	5	<i>Gomerella graninicola</i>
pYKH1139			MFalpha	9	
pYKH1135	FAB	2	native	6	<i>Fusarium vesticilloles</i> /
pYKH1095			MFalpha	9	<i>Rasamsonia emersonii</i> /
					<i>Hypocrea jecorina</i>
pYKH1097	Mg3A	3	native	7	<i>Magnaporthe grisea</i>
pYKH1096			MFalpha	9	
pYKH1099	TrBGL1	4	native	8	<i>Hypocrea jecorina</i>
pYKH1098			MFalpha	9	

The plasmids listed in Table 2 were transformed into strain FG-ura3 and transformants were selected by growth on synthetic complete medium lacking uracil (SC-ura). FG-ura3 is a derivative of FERMAX™ Gold (hereafter abbreviated, "FG") in which the ura3 gene has been deleted. The construction of FG-ara3 is described in WO201811792A1. FG-ura3 was also transformed with the control plasmid pPOL00040 that lacks a β -glucosidase coding sequence. Transformants were grown overnight in SC-ura liquid medium, and β -glucosidase activity in the culture medium was measured as described in Example 2. The results from two such experiments are shown in Table 3.

TABLE 3

β -glucosidase activity measured in FG-ura3 transformed with β -glucosidase expression plasmids				
Plasmid	Enzyme	Signal sequence	A ₄₀₅ (Exp. 1)	A ₄₀₅ (Exp. 2)
pYKH1127	ABG54	native	0.107 $\pm$ 0.018	0.174 $\pm$ 0.023
pYKH1139		MFalpha	n/d	0.244 $\pm$ 0.055
pYKH1135	FAB	native	0.086 $\pm$ 0.004	0.156 $\pm$ 0.009
pYKH1095		MFalpha	0.152 $\pm$ 0.014	0.308 $\pm$ 0.057
pYKH1097	Mg3A	native	0.005 $\pm$ 0.001	n/d
pYKH1096		MFalpha	0.004 $\pm$ 0.001	n/d
pYKH1099	TrBGL1	native	0.004 $\pm$ 0.001	n/d
pYKH1098		MFalpha	0.004 $\pm$ 0.000	n/d
pPOL00040	none	n/a	0.003 $\pm$ 0.001	0.003 $\pm$ 0.001

A₄₀₅ was higher for strains containing plasmids encoding ABG54 or FAB than for the control strain transformed with pPOL00040. This demonstrates that strains encoding ABG54 or FAB express β -glucosidase activity. β -glucosidase activity was detected for strains expressing ABG54 or FAB with either their native signal sequences or the MFalpha signal sequence. In contrast, strains containing plasmids encoding Mg3A or TrBGL1 did not appear to produce β -glucosidase activity above that measured for the negative control strain transformed with pPOL00040.

16

Example 4: Strains with Integrated Expression Cassettes for ABG54 or FAB

Expression cassettes coding for ABG54 with its native signal sequence, ABG54 with the MFalpha signal sequence, FAB with its native signal sequence (SEQ ID NO:6) or FAB with the MFalpha signal sequence (SEQ ID NO:9) were amplified by PCR from plasmids pYKH1127, pYKH1139, pYKH1135 or pYKH1095, respectively, using primers that incorporate regions of homology to a target site at the PAM1

locus in *S. cerevisiae*. Each of the amplified DNA fragments was used as a donor DNA for CRISPR-mediated integration at the PAM1 locus in FG. Integration of the β -glucosidase expression cassettes was confirmed by colony PCR, and the resulting strains were designated as shown in Table 4.

Strains were also constructed in which the native signal sequence of ABG54 or FAB was replaced with the signal sequence from the *S. cerevisiae* invertase gene, SUC2 (SEQ ID NO: 10). These strains were constructed by transforming FG with two overlapping DNA fragments—one containing the *S. cerevisiae* FBA1 promoter joined to the SUC2 signal sequence, and the other containing the ABG54 or FAB coding sequence (without the native signal sequence) joined to the FBA1 terminator. These fragments were assembled and integrated at the PAM1 locus using the procedure described in EauClaire et. al. (2016) *J. Ind. Microbiol. Biotechnol.* 43:1001-15. Correct assembly and integration of the β -glucosidase expression cassettes was confirmed by colony PCR. The resulting strains were designated as shown in Table 4.

Four clones of each strain, together with the parental strain FG, were grown overnight in YPD liquid medium, and β -glucosidase activity in the resulting culture medium was measured as described in Example 2. The results are shown in Table 4.

TABLE 4

β -glucosidase activity secreted by strains encoding ABG54 or FAB, versus FG.			
Strain	β -glucosidase expressed	Signal sequence	A ₄₀₅
GKH-0464	ABG54	native	0.970 $\pm$ 0.031
GKH-0459		MFalpha	1.240 $\pm$ 0.107
GKH-0484		SUC2	0.922 $\pm$ 0.016
GKH-0455	FAB	native	0.248 $\pm$ 0.011
GKH-0450		MFalpha	0.666 $\pm$ 0.039
GKH-0466		SUC2	0.170 $\pm$ 0.053
FG	n/a	n/a	0.088 $\pm$ 0.012

A₄₀₅ was higher for the new strains with integrated β -glucosidase expression cassettes than for the parental FG strain. This indicates that strains GKH-0464, GKH-0459, GKH-0484, GKH-0455, GKH-0450 and GKH-0466 all express β -glucosidase activity.

Example 5. Ethanol Production by *S. cerevisiae* Strains Expressing β -Glucosidase in Vial Assays

Four clones of each strain listed in Table 4 were screened for ethanol production, relative to the parent strain FG. Strains were grown in corn liquefact in serum vials, and ethanol production was analyzed after 55 h fermentation as described in Example 2. The results are shown in Table 5.

TABLE 5

Performance of β -glucosidase expressing yeast strains versus FG in vial assays				
Strain	β -glucosidase	Signal sequence	Ethanol (g/L)	Ethanol increase vs FG (%)
GKH-0464	ABG54	native	142.41 $\pm$ 0.31	1.5
GKH-0459		MFalpha	141.35 $\pm$ 0.59	0.7
GKH-0484		SUC2	142.02 $\pm$ 0.92	1.2
GKH-0455	FAB	native	141.99 $\pm$ 0.12	1.2
GKH-0450		MFalpha	138.94 $\pm$ 0.41	-1.0
GKH-0466		SUC2	142.31 $\pm$ 0.33	1.4
FG	n/a	n/a	140.37 $\pm$ 0.40	n/a

The (3-glucosidase expressing strains GKH-0464, GKH-0459, GKH-0484, GKH-0455 and GKH-0466 resulted in up to 1.5% increased ethanol production compared to the parental FG strain.

Example 6: Ethanol Production by *S. cerevisiae* Strains Expressing β -Glucosidase in Shake Flask Assays

To further examine the benefits of β -glucosidase expression, the performance of strains GKH-0464 and GKH-0484 were analyzed more precisely using a shake flask assay. Strains GKH-0464, GKH-0484 or the parental strain FG were grown for 55 h in corn liquefact, and their fermentation products analyzed as described in Example 2. In a second set of flasks, approx. 5 g/L cellobiose was added prior to inoculation with the yeast. The results are shown in Table 6.

TABLE 6

Performance of β -glucosidase expressing yeast strains versus FG in shake flask assays with or without cellobiose								
Strain	ABG54 exp.	Cellobiose added	Ethanol (g/L)	Glycerol (g/L)	Acetate (g/L)	Glucose (g/L)	DP2 (g/L)	DP3 (g/L)
GKH-0464	Yes	No	146.16	16.01	0.88	2.12	4.15	1.25
GKH-0484	Yes	No	143.87	16.36	0.88	4.87	4.12	1.40
FG	No	No	142.02	16.25	0.92	5.73	4.24	1.58
GKH-0464	Yes	Yes	147.00	16.34	0.90	2.86	4.31	1.37
GKH-0484	Yes	Yes	143.36	16.39	0.92	7.58	4.64	1.50
FG	No	Yes	142.25	16.19	0.96	6.90	10.16	1.77

The β -glucosidase-expressing strains GKH-0464 and GKH-0484 resulted in up to 3.3% increased ethanol production compared to the parental FG strain. GKH-0464 and GKH-0484 also resulted in up to 6.3% decreased acetate, as well as decreased DP2 and decreased DP3 compared to the parental strain. For fermentations with FG, addition of approximately 5 g/L cellobiose to the liquefact prior to inoculation with the yeast resulted in a corresponding increase in DP2 at the end of fermentation. This indicates that the added cellobiose was not consumed during fermentation with FG. In contrast, for GKH-0464 and GKH-0484, the DP2 concentration at the end of fermentation was similar for liquefact with or without added cellobiose. This indicates that cellobiose can be consumed during fermentation with GKH-0464 and GKH-0484.

Example 7: Strains Expressing β -Glucosidase with the PKL Pathway and Glucoamylase

An ABG54 coding sequence, identical to that in plasmid pYKH1127, flanked by the *S. cerevisiae* FBA1 promoter and GPD1 terminator, was integrated at the JIP5 locus in *S. cerevisiae*. The amplified DNA fragment was used as a donor DNA for CRISPR-mediated integration at the JIP5 locus in two parental strains: (i) FG-PKL and (ii) FG-PKL-GA. FG-PKL is an engineered FG yeast having a heterologous phosphoketolase (PKL) pathway involving the expression of phosphoketolase (PKL), phosphotransacetylase (PTA) and acetylating acetyl dehydrogenase (AADH), as described in WO2015148272. FG-PKL-GA is the FG-PKL strain further engineered to express a variant of *Trichoderma* glucoamylase.

Integration of the ABG54 expression cassette into these strains was confirmed by colony PCR. The resulting strains, along with the parental strains lacking the ABG54 expression cassette, were grown in corn liquefact in serum vials, and their fermentation products analyzed as described in Example 2. The results are shown in Table 7.

TABLE 7

Performance of β -glucosidase expressing strains G3020 and G3014 versus parental strains in vial assays							
Strains	Features	Ethanol (g/L)	Glycerol (g/L)	Acetate (g/L)	Glucose (g/L)	DP2 (g/L)	DP3 (g/L)
FG-PKL	PKL pathway	145.27	12.84	1.77	2.17	4.12	1.38
G3020	PKL pathway + ABG54	145.87	12.61	1.69	1.48	3.18	0.84
FG-PKL-GA	PKL pathway + GA	146.73	11.64	1.64	1.49	4.27	1.47
G3014	PKL pathway + GA + ABG54	147.40	11.75	1.43	1.46	3.61	0.99

The β -glucosidase-expressing strains G3020 and G3014 resulted in slightly increased ethanol production (<0.5%) compared to their corresponding parental strains. However, G3020 and G3014 resulted in up to almost 13% decreased acetate, as well as decreased DP2 and decreased DP3 compared to their respective parental strains.

Example 8: Hybrid Yeast Strains Expressing β -Glucosidase

The ABG54 expression cassette from plasmid pYKH1127 was amplified by PCR using primers that incorporate regions of homology to a target site at the JEN1 locus in *S. cerevisiae*. The amplified DNA fragment was used as a donor DNA for CRISPR-mediated integration at the JEN1 locus in two parental strains: (i) DGY1- Δ and (ii) DGY1- Δ -GA. DGY1- Δ is a hybrid yeast strain produced by mating two commercially-available parental yeast strains and further modified by deletion of the YJL065c gene encoding Dls1 (see, e.g., WO2018089333). DGY1- Δ -GA is the DGY1- Δ strain further modified to express a glucoamylase.

Integration of the ABG54 expression cassette into these strains was confirmed by colony PCR. The resulting strains, along with the parental strains lacking the ABG54 expression cassette, were grown in corn liquefact in shake flasks, and their fermentation products analyzed as described in Example 2. The results are shown in Table 8.

TABLE 8

Performance of β -glucosidase-expressing hybrid strains versus parental strains							
Strain	Features	Ethanol (g/L)	Glycerol (g/L)	Acetate (g/L)	Glucose (g/L)	DP2 (g/L)	DP3 (g/L)
DGY1- Δ	Hybrid	143.64	13.41	1.62	0.23	2.89	1.25
GKH-0737	Hybrid + ABG54	144.72	13.47	1.58	0.33	2.72	0.45
DGY1- Δ -GA	Hybrid + GA	144.17	12.69	1.36	0.34	3.15	1.24
GKH-0732	Hybrid + GA + ABG54	145.46	12.68	1.26	0.18	4.02	0.80

The β -glucosidase-expressing strains GKH-0737 and GKH-0732 resulted in up to about 0.9% increased ethanol production compared to their corresponding parental strains and up to over 7% decreased acetate, as well as decreased DP3, compared to their respective parental strains.

Example 9: Comparison of ABG54 and FAB with Other β -Glucosidases In Vitro

Following the observation that ABG54 and FAB were the best expressed β -glucosidases in yeast (Example 3), an experiment similar to that described in Example 1 was

performed to compare the activity of ABG54 and FAB to other β -glucosidases in the aforementioned whole stillage assay.

As shown in Example 1, ABG54 was the best performing glycosyl hydrolase for releasing glucose from soluble recalcitrant oligosaccharides in whole stillage, leading to the impetus to express β -glucosidases in yeast. As shown in Example 3, ABG54 and FAB were shown to be the best expressing β -glucosidases for expression in yeast.

In an essentially a fill-circle experiment summarized by the results shown in Table 9, ABG54 and FAB are demonstrated to be better performing β -glucosidases than both Mg3A and TrBgl1 (the two other β -glucosidases tested in yeast) and superior to a blend of other GH-15 and GH-3 molecules. Notably and not surprisingly, all the enzymes selected and tested, herein, were considered best-in-class, otherwise they would not have been tested.

Regardless, the measurable superior performance of ABG54 and FAB compared to Mg3A and TrBgl1 (in particular) in no way explains the remarkable difference in expression in yeast. The differences in expression of the β -glucosidases are ordinals of magnitude greater than the difference in performance of the enzymes in vitro. These data confirm the tenet that expression of exogenous enzymes in yeast is often problematic, and that ABG54 and FAB are especially well expressed, and are additionally superior within their class. These results are shown in Table 9.

TABLE 9

Comparison of in vitro performance of β -glucosidases expressed in yeast	
Enzyme combination	Glucose produced (mM)
PB + ABG54	5.99
PB + FAB	5.61
PB + Mg3A	4.92
PB + TrBgl1	5.49
GH15 + GH3 blend	3.61

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 20

<210> SEQ ID NO 1

<211> LENGTH: 860

<212> TYPE: PRT

<213> ORGANISM: *Gomerella graminicola*

<400> SEQUENCE: 1

```

Ala Val Thr Thr Glu Arg Gln Leu His Lys Arg Asp Leu Ala Tyr Ser
1          5          10          15
Pro Pro Val Tyr Pro Ser Pro Trp Met Asp Pro Asn Ala Asp Gly Trp
20        25        30
Thr Asp Ala Tyr Ala Lys Ala Lys Asp Phe Val Ser Gln Leu Thr Leu
35        40        45
Leu Glu Lys Val Asn Leu Thr Thr Gly Val Gly Trp Gln Gly Asp Leu
50        55        60
Cys Val Gly Asn Val Gly Ser Val Pro Arg Leu Gly Leu Arg Gly Leu
65        70        75        80
Cys Met Gln Asp Gly Pro Val Gly Ile Arg Phe Ser Asp Tyr Asn Ser
85        90        95
Val Phe Pro Ser Gly Gln Thr Ala Ala Ala Thr Trp Asp Arg Glu Leu
100       105       110
Ile Tyr Arg Arg Ala Glu Ala Ile Gly Phe Glu His Arg Ala Lys Gly
115       120       125
Val Asp Val Val Leu Ala Pro Val Ala Gly Pro Ile Gly Arg Ala Pro
130       135       140
Ala Gly Gly Arg Asn Trp Glu Gly Phe Ser Ser Asp Pro Tyr Leu Thr
145       150       155       160
Gly Val Ala Met Ala Glu Ser Val Lys Gly Ile Gln Gln His Ala Ile
165       170       175
Ala Cys Ala Lys His Phe Ile Gly Asn Glu Gln Glu His Phe Arg Gln
180       185       190
Ala Pro Glu Ala Ile Gly Tyr Asn Tyr Thr Ile Asp Glu Ser Ile Ser
195       200       205
Ser Asn Ile Asp Asp Lys Thr Leu His Glu Leu Tyr Leu Trp Pro Phe
210       215       220
Gln Asp Ala Val Ala Ala Gly Val Gly Ser Phe Met Cys Ser Tyr Asn
225       230       235       240
Gln Val Asn Asn Ser Tyr Gly Cys Gln Asn Ser Lys Leu Met Asn Gly
245       250       255
Ile Leu Lys Asp Glu Leu Gly Phe Gln Gly Phe Ile Met Ser Asp Trp
260       265       270
Ala Ala Gln His Ala Gly Val Ala Thr Ala Val Ala Gly Leu Asp Met
275       280       285
Ala Met Pro Gly Asp Thr Ala Phe Asn Ser Gly Met Thr Phe Trp Gly
290       295       300
Thr Asn Leu Thr Val Ala Val Leu Asn Gly Thr Leu Pro Glu Tyr Arg
305       310       315       320
Leu Asp Asp Met Ala Met Arg Ile Met Ala Ala Phe Phe Lys Val Gly
325       330       335
Phe Glu Leu Asn Glu Val Pro Glu Ile Asn Phe Ser Ser Trp Thr Thr
340       345       350
Asp Thr Val Gly Pro Leu Gln Tyr Tyr Ala Lys Glu Asn Val Gln Val
355       360       365

```

Ile	Asn	Gln	His	Val	Asp	Val	Arg	Arg	Gly	Gln	Glu	His	Gly	Lys	Leu
370 375 380															
Ile	Arg	Glu	Ile	Ala	Ala	Lys	Ala	Thr	Val	Leu	Leu	Lys	Asn	Glu	Gly
385 390 395 400															
Ala	Leu	Pro	Leu	Lys	Lys	Pro	Lys	Phe	Leu	Ala	Val	Ile	Gly	Glu	Asp
405 410 415															
Ala	Gly	Pro	Asn	Leu	Ser	Gly	Pro	Asn	Gly	Cys	Ser	Asp	His	Gly	Cys
420 425 430 435															
Asn	Glu	Gly	Thr	Leu	Gly	Ala	Gly	Trp	Gly	Ser	Gly	Thr	Ser	Asn	Tyr
440 445															
Pro	Tyr	Leu	Ile	Thr	Pro	Asp	Gln	Ala	Leu	Gln	Ala	Arg	Ala	Val	Ala
450 455 460 465															
Glu	Gly	Ser	Arg	Tyr	Glu	Ser	Ile	Leu	Ser	Asn	Tyr	Asp	Phe	Ala	Ala
470 475 480															
Thr	Thr	Ala	Leu	Val	Thr	Gln	Pro	Asp	Ala	Thr	Ala	Ile	Val	Phe	Val
485 490 495															
Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	Asp	Val	Gly	Gly	Asn	Glu	Gly
500 505 510															
Asp	Arg	Gln	Asn	Leu	Thr	Leu	Trp	Asn	Gly	Gly	Asp	Glu	Leu	Val	Lys
515 520 525															
Asn	Val	Ala	Ala	Gly	Asn	Asn	Asn	Thr	Ile	Val	Val	Ile	His	Ser	Val
530 535 540															
Gly	Pro	Val	Leu	Leu	Ala	Asp	Met	Lys	Asn	Asn	Pro	Asn	Ile	Thr	Ala
545 550 555															
Ile	Val	Trp	Ala	Gly	Leu	Pro	Gly	Gln	Glu	Ser	Gly	Asn	Ser	Ile	Thr
560 565 570 575															
Asp	Val	Leu	Tyr	Gly	Asp	Val	Asn	Pro	Gly	Gly	Lys	Ser	Pro	Phe	Thr
580 585 590															
Trp	Gly	Pro	Thr	Arg	Glu	Ser	Tyr	Gly	Thr	Asp	Val	Leu	Tyr	Glu	Pro
595 600 605															
Asn	Asn	Gly	Glu	Gly	Ala	Pro	Gln	Asp	Asp	Phe	Ser	Glu	Gly	Val	Phe
610 615 620															
Ile	Asp	Tyr	Arg	Tyr	Phe	Asp	Arg	Ala	Thr	Ser	Gly	Ser	Asn	Glu	Thr
625 630 635															
Ser	Thr	Gly	Ala	Ala	Pro	Val	Tyr	Pro	Phe	Gly	Phe	Gly	Leu	Ser	Tyr
640 645 650 655															
Thr	Thr	Phe	Glu	Tyr	Ser	Asn	Leu	Val	Val	Thr	Pro	Lys	Glu	Ala	Gly
660 665 670															
Glu	Tyr	Thr	Pro	Thr	Thr	Gly	Val	Thr	Glu	Lys	Ala	Pro	Thr	Phe	Gly
675 680 685															
Asn	Tyr	Ser	Thr	Asp	Pro	Ala	Ala	Tyr	Val	Phe	Pro	Ser	Gly	Glu	Phe
690 695 700															
Arg	Tyr	Ile	Tyr	Asn	Phe	Ile	Tyr	Pro	Tyr	Leu	Asn	Thr	Thr	Asp	Ile
705 710 715 720															
Ser	Lys	Ser	Ala	Asn	Asp	Pro	Ala	Tyr	Gly	Gln	Thr	Ala	Asp	Glu	Phe
725 730 735															
Leu	Pro	Pro	Lys	Ala	Leu	Glu	Ser	Gly	Pro	Gln	Pro	Lys	His	Pro	Ala
740 745 750															
Ser	Gly	Ala	Pro	Gly	Gly	Asn	Pro	Gln	Leu	Trp	Asp	Val	Leu	Tyr	Thr
755 760 765															
Val	Thr	Ala	Thr	Ile	Thr	Asn	Lys	Gly	Asp	Val	Ala	Gly	Asp	Glu	Val
770 775 780															
Ala	Gln	Leu	Tyr	Val	Ser	Leu	Gly	Gly	Pro	Asn	Asp	Pro	Val	Lys	Val

-continued

785	790	795	800
Leu Arg Gly Phe Asp Arg Ile Gly Ile Ala Pro Gly Glu Ser Ala Thr			
	805	810	815
Phe Thr Ala Asp Ile Thr Arg Arg Asp Leu Ser Asn Trp Asp Thr Val			
	820	825	830
Ser Gln Asn Trp Val Ile Ser Lys Tyr Pro Lys Lys Val Trp Val Gly			
	835	840	845
Gly Ser Ser Arg Glu Leu Pro Leu Ser Ala Ser Leu			
	850	855	860

<210> SEQ ID NO 2

<211> LENGTH: 867

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: hybrid from *Fusarium vesticilloies* / *Rasamsonia emersonii* / *Hypocrea jecorina*

<400> SEQUENCE: 2

Ala Val Ala Leu Ala Ser Ala Val Pro Asp Thr Leu Ala Gly Val Lys			
1	5	10	15
Lys Ala Asp Ala Gln Lys Val Val Thr Arg Asp Thr Leu Ala Tyr Ser			
	20	25	30
Pro Pro His Tyr Pro Ser Pro Trp Met Asp Pro Asn Ala Val Gly Trp			
	35	40	45
Glu Glu Ala Tyr Ala Lys Ala Lys Ser Phe Val Ser Gln Leu Thr Leu			
	50	55	60
Met Glu Lys Val Asn Leu Thr Thr Gly Val Gly Trp Gln Gly Glu Arg			
	65	70	75
Cys Val Gly Asn Val Gly Ser Ile Pro Arg Leu Gly Met Arg Gly Leu			
	85	90	95
Cys Leu Gln Asp Gly Pro Leu Gly Ile Arg Leu Ser Asp Tyr Asn Ser			
	100	105	110
Ala Phe Pro Ala Gly Thr Thr Ala Gly Ala Ser Trp Ser Lys Ser Leu			
	115	120	125
Trp Tyr Glu Arg Gly Leu Leu Met Gly Thr Glu Phe Lys Glu Lys Gly			
	130	135	140
Ile Asp Ile Ala Leu Gly Pro Ala Thr Gly Pro Leu Gly Arg Thr Ala			
	145	150	155
Ala Gly Gly Arg Asn Trp Glu Gly Phe Thr Val Asp Pro Tyr Met Ala			
	165	170	175
Gly His Ala Met Ala Glu Ala Val Lys Gly Ile Gln Asp Ala Gly Val			
	180	185	190
Ile Ala Cys Ala Lys His Tyr Ile Ala Asn Glu Gln Glu His Phe Arg			
	195	200	205
Gln Ser Gly Glu Val Gln Ser Arg Lys Tyr Asn Ile Ser Glu Ser Leu			
	210	215	220
Ser Ser Asn Leu Asp Asp Lys Thr Met His Glu Leu Tyr Ala Trp Pro			
	225	230	235
Phe Ala Asp Ala Val Arg Ala Gly Val Gly Ser Val Met Cys Ser Tyr			
	245	250	255
Asn Gln Ile Asn Asn Ser Tyr Gly Cys Gln Asn Ser Lys Leu Leu Asn			
	260	265	270
Gly Ile Leu Lys Asp Glu Met Gly Phe Gln Gly Phe Val Met Ser Asp			
	275	280	285

Trp 290	Ala	Gln	His	Thr	Gly 295	Ala	Ala	Ser	Ala	Val 300	Ala	Gly	Leu	Asp
Met 305	Ser	Met	Pro	Gly	Asp 310	Thr	Ala	Phe	Asp	Ser 315	Gly	Tyr	Ser	Phe 320
Gly	Gly	Asn	Leu	Thr 325	Leu	Ala	Val	Ile	Asn 330	Gly	Thr	Val	Pro	Ala 335
Arg	Val	Asp	Asp	Met 340	Ala	Leu	Arg	Ile 345	Met	Ser	Ala	Phe	Phe 350	Lys 355
Gly	Lys	Thr 355	Ile	Glu	Asp	Leu	Pro 360	Asp	Ile	Asn	Phe	Ser 365	Ser	Trp 370
Arg	Asp 370	Thr	Phe	Gly	Phe 375	Val	His	Thr	Phe	Ala	Gln 380	Glu	Asn	Arg 385
Gln 385	Val	Asn	Phe	Gly 390	Val	Asn	Val	Gln	His 395	Asp	His	Lys	Ser	Ile 400
Arg	Glu	Ala	Ala	Ala 405	Lys	Gly	Ser	Val	Val 410	Leu	Lys	Asn	Thr	Gly 415
Leu	Pro	Leu	Lys 420	Asn	Pro	Lys	Phe	Leu	Ala 425	Val	Ile	Gly	Glu	Asp 430
Gly	Pro	Asn 435	Pro	Ala	Gly	Pro	Asn 440	Gly	Cys	Gly	Asp	Arg 445	Gly	Cys 450
Asn 450	Gly	Thr	Leu	Ala	Met 455	Ala	Trp	Gly	Ser	Gly	Thr 460	Ser	Gln	Phe 465
Tyr 465	Leu	Ile	Thr	Pro 470	Asp	Gln	Gly	Leu	Ser 475	Asn	Arg	Ala	Thr	Gln 480
Gly	Thr	Arg	Tyr	Glu 485	Ser	Ile	Leu	Thr	Asn 490	Asn	Glu	Trp	Ala	Ser 495
Gln	Ala	Leu	Val 500	Ser	Gln	Pro	Asn	Val 505	Thr	Ala	Ile	Val	Phe 510	Ala 515
Ala	Asp	Ser 515	Gly	Glu	Gly	Tyr	Ile 520	Glu	Val	Asp	Gly	Asn 525	Phe	Gly 530
Arg	Lys 530	Asn	Leu	Thr	Leu 535	Trp	Gln	Gln	Gly	Asp 540	Glu	Leu	Ile	Lys 545
Val 545	Ser	Ser	Ile	Cys 550	Pro	Asn	Thr	Ile	Val 555	Val	Leu	His	Thr	Val 560
Pro	Val	Leu	Leu	Ala 565	Asp	Tyr	Glu	Lys	Asn 570	Pro	Asn	Ile	Thr	Ala 575
Val	Trp	Ala	Gly 580	Leu	Pro	Gly	Gln	Glu 585	Ser	Gly	Asn	Ala	Ile 590	Ala 595
Leu	Leu	Tyr 595	Gly	Lys	Val	Ser	Pro 600	Gly	Arg	Ser	Pro	Phe 605	Thr	Trp 610
Arg	Thr 610	Arg	Glu	Ser	Tyr 615	Gly	Thr	Glu	Val	Leu	Tyr 620	Glu	Ala	Asn 625
Gly 625	Arg	Gly	Ala	Pro 630	Gln	Asp	Asp	Phe	Ser 635	Glu	Gly	Val	Phe	Ile 640
Tyr	Arg	His	Phe 645	Asp	Lys	Tyr	Asn	Ile	Thr 650	Pro	Ile	Tyr	Glu	Phe 655
His	Gly	Leu	Ser 660	Trp	Ser	Thr	Phe	Lys 665	Phe	Ser	Asn	Leu	His 670	Ile 675
Lys	Asn	Asn	Val 675	Gly	Pro	Met	Ser 680	Pro	Pro	Asn	Gly	Lys 685	Thr	Ile 690
Ala	Pro	Ser	Leu 690	Gly	Asn	Phe 695	Ser	Lys	Asn	Leu	Lys 700	Asp	Tyr	Gly 705
Pro	Lys	Asn	Val	Arg	Arg	Ile	Lys	Glu	Phe	Ile	Tyr	Pro	Tyr	Leu 710

-continued

705	710	715	720
Thr Thr Thr Ser Gly Lys Glu Ala Ser Gly Asp Ala His Tyr Gly Gln			
	725	730	735
Thr Ala Lys Glu Phe Leu Pro Ala Gly Ala Leu Asp Gly Ser Pro Gln			
	740	745	750
Pro Arg Ser Ala Ala Ser Gly Glu Pro Gly Gly Asn Arg Gln Leu Tyr			
	755	760	765
Asp Ile Leu Tyr Thr Val Thr Ala Thr Ile Thr Asn Thr Gly Ser Val			
	770	775	780
Met Asp Asp Ala Val Pro Gln Leu Tyr Leu Ser His Gly Gly Pro Asn			
	785	790	800
Glu Pro Pro Lys Val Leu Arg Gly Phe Asp Arg Ile Glu Arg Ile Ala			
	805	810	815
Pro Gly Gln Ser Val Thr Phe Lys Ala Asp Leu Thr Arg Arg Asp Leu			
	820	825	830
Ser Asn Trp Asp Thr Lys Lys Gln Gln Trp Val Ile Thr Asp Tyr Pro			
	835	840	845
Lys Thr Val Tyr Val Gly Ser Ser Ser Arg Asp Leu Pro Leu Ser Ala			
	850	855	860
Arg Leu Pro			
865			
<210> SEQ ID NO 3			
<211> LENGTH: 855			
<212> TYPE: PRT			
<213> ORGANISM: Magnaporthe grisea			
<400> SEQUENCE: 3			
His Pro Gly Asp Tyr Ser Lys Leu Glu Arg Arg Ala Val Ala Thr Ser			
1	5	10	15
Glu Pro His Tyr Pro Gln Pro Trp Met Asn Pro Asp Ala Asp Gly Trp			
	20	25	30
Gln Glu Ala Tyr Val Lys Ala Lys Asp Phe Val Ser Gln Met Thr Leu			
	35	40	45
Leu Glu Lys Val Asn Leu Thr Thr Gly Val Gly Trp Ala Ser Asp Leu			
	50	55	60
Cys Val Gly Asn Val Gly Ala Val Pro Arg Leu Gly Leu Arg Ser Leu			
	65	70	75
Cys Leu Gln Asp Ser Pro Thr Gly Val Arg Phe Ala Asp Trp Val Ser			
	85	90	95
Val Phe Pro Ala Gly Ile Thr Thr Gly Ala Thr Phe Asp Lys Gly Leu			
	100	105	110
Met Tyr Arg Arg Gly Gln Ala Met Gly Gln Glu Ala Lys Asp Lys Gly			
	115	120	125
Ile Asn Val Leu Leu Gly Pro Val Ala Gly Gly Leu Gly Arg Val Ala			
	130	135	140
Ala Gly Gly Arg Ala Trp Glu Ser Phe Gly Ala Asp Pro Val Leu Thr			
	145	150	155
Gly Tyr Gly Met Ile Glu Thr Ile Lys Gly Ile Gln Asp Thr Gly Val			
	165	170	175
Ile Ala Thr Ala Lys His Phe Ile Gly Asn Glu Gln Glu His Phe Arg			
	180	185	190
Gln Val Gly Glu Glu Arg Gly Arg Gly Val Asn Ile Ser Glu Ser Leu			
	195	200	205

Ser 210	Ser	Asn	Ile	Asp	Asp	Lys 215	Thr	Met	His	Glu	Leu 220	Tyr	Leu	Trp	Pro
Phe 225	Ala	Asp	Ala	Val	Arg 230	Ala	Gly	Val	Gly	Ser 235	Val	Met	Cys	Ser	Tyr 240
Thr	Gln	Val	Asn	Asn 245	Ser	Tyr	Gly	Cys	Gln 250	Asn	Ser	Lys	Leu	Leu 255	Asn
Gly	Leu	Leu	Lys 260	Asp	Glu	Leu	Gly	Phe	Gln 265	Gly	Phe	Val	Met	Ser	Asp
Trp	Gln	Ala	Gln	His	Thr	Gly	Ala 280	Ala	Ser	Ala	Ala 285	Ala	Gly	Leu	Asp
Met	Ser 290	Met	Pro	Gly	Asp	Thr 295	Glu	Phe	Asn	Thr	Gly 300	Leu	Ser	Phe	Trp
Gly 305	Ala	Asn	Leu	Thr	Leu 310	Ala	Val	Val	Asn	Gly 315	Thr	Val	Ala	Glu	Trp 320
Arg	Ile	Asp	Asp	Met 325	Ala	Met	Arg	Ile	Met 330	Ala	Ala	Phe	Phe	Lys 335	Val
Gly	Asn	Thr	Leu 340	Asp	Gln	Pro	Glu	Ile	Asn 345	Phe	Ser	Ser	Trp	Thr	Lys
Asp	Thr	Phe 355	Gly	Pro	Leu	His	Ser 360	Ser	Ser	Gly	Asn 365	Arg	Ile	Gln	Gln
Ile 370	Asn	Gln	His	Val	Asp	Val 375	Arg	Arg	Asp	His	Gly 380	Asn	Leu	Ile	Arg
Glu 385	Val	Ala	Ala	Lys 390	Gly	Thr	Val	Leu	Leu	Lys 395	Asn	Thr	Asn	Asn	Ala 400
Leu	Pro	Leu	Asn	Lys 405	Pro	Lys	Phe	Leu	Ala 410	Val	Ile	Gly	Asp	Asp	Ala 415
Gly	Ser	Asn	Pro 420	Arg	Gly	Pro	Asn	Gly 425	Cys	Pro	Asp	Arg	Gly 430	Cys	Leu
Leu	Gly 435	Thr	Leu	Gly	Met	Ala	Trp 440	Gly	Ser	Gly	Thr 445	Ala	Asp	Phe	Pro
Tyr 450	Leu	Ile	Thr	Pro	Asp	Ala 455	Ala	Leu	Gln	Ala	Gln 460	Ala	Ile	Glu	Asp
Gly 465	Thr	Arg	Tyr	Glu	Ser 470	Ile	Leu	Ser	Asn	Tyr 475	Ala	Thr	Ala	Gln	Thr 480
Gln	Ala	Leu	Val	Ser 485	Gln	Thr	Tyr	Ala	Thr 490	Ala	Ile	Val	Phe	Val 495	Ala
Ala	Ser	Ser	Gly 500	Glu	Gly	Tyr	Ile	Asp 505	Phe	Asp	Gly	Asn	Lys 510	Gly	Asp
Arg	Asn 515	Asn	Leu	Thr	Leu	Trp	Tyr 520	Asp	Gly	Asp	Ser	Leu 525	Val	Lys	Asn
Val 530	Ser	Ser	Val	Cys	Asn	Asn 535	Thr	Ile	Val	Val	Ile 540	His	Ser	Thr	Gly
Pro 545	Thr	Ile	Leu	Thr	Glu 550	Trp	Tyr	Asp	Asn	Pro 555	Asn	Val	Thr	Ala	Ile 560
Val	Trp	Ala	Gly	Val 565	Pro	Gly	Gln	Glu	Ser 570	Gly	Arg	Ala	Ile	Thr	Asp 575
Val	Leu	Tyr	Gly 580	Arg	Val	Asn	Pro	Ala 585	Gly	Arg	Ser	Pro	Phe	Thr	Trp
Gly	Lys 595	Thr	Arg	Glu	Ser	Tyr	Gly 600	Thr	Asp	Val	Met 605	Tyr	Lys	Pro	Asn
Asn 610	Gly	Asn	Glu	Ala	Pro	Gln 615	Gln	Asp	Tyr	Thr	Glu 620	Gly	Val	Phe	Ile
Asp	Tyr	Arg	His	Phe	Asp	Gln	Gln	Lys	Asp	Glu	Pro	Val	Tyr	Glu	Phe

-continued

625	630	635	640
Gly His Gly Leu Ser Tyr Thr Thr Phe Glu Tyr Ser Asn Ile Arg Val			
	645	650	655
Asp Lys Ala Pro Ala Ser Glu Tyr Lys Pro Thr Thr Gly Gln Thr Ile			
	660	665	670
Pro Ala Pro Val Phe Gly Ala Asn Val Ser Lys Asp Leu Ser Gln Tyr			
	675	680	685
Thr Phe Pro Ser Asp Glu Phe Pro His Ile Tyr Leu Phe Ile Tyr Pro			
	690	695	700
Tyr Leu Asn Thr Ser Ser Ser Gly Glu Glu Ala Ser Arg Asp Pro Lys			
	705	710	715
Tyr Gly Gly Thr Ala Glu Glu Phe Leu Pro Pro Lys Ala Leu Asp Gly			
	725	730	735
Ser Pro Gln Pro Leu Pro Arg Ala Ser Gly Lys Asn Ser Pro Gly Gly			
	740	745	750
Asn Arg Gln Leu Tyr Asp Thr Leu Tyr Thr Val Thr Ala Thr Ile Thr			
	755	760	765
Asn Thr Gly Lys Leu Val Gly Glu Glu Val Pro Gln Leu Tyr Val Ser			
	770	775	780
His Gly Gly Pro Glu Asp Pro Pro Val Val Leu Arg Gly Phe Glu Arg			
	785	790	795
Ile Arg Leu Asp Pro Gly Gln Ser Ala Thr Phe Lys Val Asp Leu Thr			
	805	810	815
Arg Arg Asp Val Ser Asn Trp Asp Val Lys Val Gln Asp Trp Val Ile			
	820	825	830
Ser Glu His Pro Lys Lys Val Phe Val Gly Ser Ser Ser Arg Lys Leu			
	835	840	845
His Leu Ser Ala Asp Leu Asn			
	850	855	

<210> SEQ ID NO 4

<211> LENGTH: 521

<212> TYPE: PRT

<213> ORGANISM: Hypocrea jecorina

<400> SEQUENCE: 4

Asp Ser His Ser Thr Ser Gly Ala Ser Ala Glu Ala Val Val Pro Pro			
1	5	10	15
Ala Gly Thr Pro Trp Gly Thr Ala Tyr Asp Lys Ala Lys Ala Ala Leu			
	20	25	30
Ala Lys Leu Asn Leu Gln Asp Lys Val Gly Ile Val Ser Gly Val Gly			
	35	40	45
Trp Asn Gly Gly Pro Cys Val Gly Asn Thr Ser Pro Ala Ser Lys Ile			
	50	55	60
Ser Tyr Pro Ser Leu Cys Leu Gln Asp Gly Pro Leu Gly Val Arg Tyr			
	65	70	75
Ser Thr Gly Ser Thr Ala Phe Thr Pro Gly Val Gln Ala Ala Ser Thr			
	85	90	95
Trp Asp Val Asn Leu Ile Arg Glu Arg Gly Gln Phe Ile Gly Glu Glu			
	100	105	110
Val Lys Ala Ser Gly Ile His Val Ile Leu Gly Pro Val Ala Gly Pro			
	115	120	125
Leu Gly Lys Thr Pro Gln Gly Gly Arg Asn Trp Glu Gly Phe Gly Val			
	130	135	140

-continued

```

Asp Pro Tyr Leu Thr Gly Ile Ala Met Gly Gln Thr Ile Asn Gly Ile
145                      150                      155                      160

Gln Ser Val Gly Val Gln Ala Thr Ala Lys His Tyr Ile Leu Asn Glu
                      165                      170                      175

Gln Glu Leu Asn Arg Glu Thr Ile Ser Ser Asn Pro Asp Asp Arg Thr
                      180                      185                      190

Leu His Glu Leu Tyr Thr Trp Pro Phe Ala Asp Ala Val Gln Ala Asn
195                      200                      205

Val Ala Ser Val Met Cys Ser Tyr Asn Lys Val Asn Thr Thr Trp Ala
210                      215                      220

Cys Glu Asp Gln Tyr Thr Leu Gln Thr Val Leu Lys Asp Gln Leu Gly
225                      230                      235                      240

Phe Pro Gly Tyr Val Met Thr Asp Trp Asn Ala Gln His Thr Thr Val
245                      250                      255

Gln Ser Ala Asn Ser Gly Leu Asp Met Ser Met Pro Gly Thr Asp Phe
260                      265                      270

Asn Gly Asn Asn Arg Leu Trp Gly Pro Ala Leu Thr Asn Ala Val Asn
275                      280                      285

Ser Asn Gln Val Pro Thr Ser Arg Val Asp Asp Met Val Thr Arg Ile
290                      295                      300

Leu Ala Ala Trp Tyr Leu Thr Gly Gln Asp Gln Ala Gly Tyr Pro Ser
305                      310                      315                      320

Phe Asn Ile Ser Arg Asn Val Gln Gly Asn His Lys Thr Asn Val Arg
325                      330                      335

Ala Ile Ala Arg Asp Gly Ile Val Leu Leu Lys Asn Asp Ala Asn Ile
340                      345                      350

Leu Pro Leu Lys Lys Pro Ala Ser Ile Ala Val Val Gly Ser Ala Ala
355                      360                      365

Ile Ile Gly Asn His Ala Arg Asn Ser Pro Ser Cys Asn Asp Lys Gly
370                      375                      380

Cys Asp Asp Gly Ala Leu Gly Met Gly Trp Gly Ser Gly Ala Val Asn
385                      390                      395                      400

Tyr Pro Tyr Phe Val Ala Pro Tyr Asp Ala Ile Asn Thr Arg Ala Ser
405                      410                      415

Ser Gln Gly Thr Gln Val Thr Leu Ser Asn Thr Asp Asn Thr Ser Ser
420                      425                      430

Gly Ala Ser Ala Ala Arg Gly Lys Asp Val Ala Ile Val Phe Ile Thr
435                      440                      445

Ala Asp Ser Gly Glu Gly Tyr Ile Thr Val Glu Gly Asn Ala Gly Asp
450                      455                      460

Arg Asn Asn Leu Asp Pro Trp His Asn Gly Asn Ala Leu Val Gln Ala
465                      470                      475                      480

Val Ala Gly Ala Asn Ser Asn Val Ile Val Val Val His Ser Val Gly
485                      490                      495

Ala Ile Ile Leu Glu Gln Ile Leu Ala Leu Pro Gln Val Lys Ala Val
500                      505                      510

Val Trp Ala Gly Leu Pro Ser Gln Glu
515                      520

```

<210> SEQ ID NO 5

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Gomerella graminicola

<400> SEQUENCE: 5

-continued

Met Arg Ser Gln Thr Leu Ala Val Ala Leu Leu Ala Ala Ala Asp Gln
1 5 10 15

Val Ala Ala

<210> SEQ ID NO 6
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: *Fusarium vesticilloides*

<400> SEQUENCE: 6

Met Lys Leu Asn Trp Val Ala Ala Ala Leu Ser Ile Gly Ala Ala Gly
1 5 10 15

Thr Asp Ser

<210> SEQ ID NO 7
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: *Magnaporthe grisea*

<400> SEQUENCE: 7

Met Arg Phe Ser Gly Ile Val Ala Thr Leu Val Ala Gly Ala Gly Val
1 5 10 15

Ser Ala

<210> SEQ ID NO 8
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: *Hypocrea jecorina*

<400> SEQUENCE: 8

Met Arg Tyr Arg Thr Ala Ala Ala Leu Ala Leu Ala Thr Gly Pro Phe
1 5 10 15

Ala Arg Ala

<210> SEQ ID NO 9
<211> LENGTH: 85
<212> TYPE: PRT
<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 9

Met Arg Phe Pro Ser Ile Phe Thr Ala Val Leu Phe Ala Ala Ser Ser
1 5 10 15

Ala Leu Ala Ala Pro Val Asn Thr Thr Thr Glu Asp Glu Thr Ala Gln
20 25 30

Ile Pro Ala Glu Ala Val Ile Gly Tyr Leu Asp Leu Glu Gly Asp Phe
35 40 45

Asp Val Ala Val Leu Pro Phe Ser Asn Ser Thr Asn Asn Gly Leu Leu
50 55 60

Phe Ile Asn Thr Thr Ile Ala Ser Ile Ala Ala Lys Glu Glu Gly Val
65 70 75 80

Ser Leu Asp Lys Arg
85

<210> SEQ ID NO 10
<211> LENGTH: 19
<212> TYPE: PRT
<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 10

-continued

Met Leu Leu Gln Ala Phe Leu Phe Leu Leu Ala Gly Phe Ala Ala Lys
1 5 10 15

Ile Ser Ala

<210> SEQ ID NO 11
<211> LENGTH: 2640
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

<400> SEQUENCE: 11

atgagatctc agacattagc ggtagcacta ttggcggcag cggaccaggt tgcagcagca	60
gtaacaactg agagacaact acataagaga gacttggcctt attctccacc tgtttacccc	120
tctccgtgga tggaccctaa tgctgacggt tggacggatg cttatgctaa agctaaggat	180
ttcgtaaacc aactgacgct gttagagaag gtaaacctaa cgactggcgt cggatggcag	240
ggtgatctat gtgtagggaa cgtggggagc gtgccaaggc tggggctaag aggtctgtgc	300
atgcaagatg gtcccgtagg aatacgttcc agcgactaca actctgtctt tcttcaggt	360
caaacggcag cggccacatg ggacagagag cttatttacc gtagggcgga agcaatcgg	420
tttgaacaca gggcgaaagg agtagacgtc gtcctagccc cggtcgcagg gcccataggt	480
agagccccag cagggggcag aaactgggaa ggattcagtt cagaccgta cctgacggga	540
gtggcgatgg ccgaatctgt aaaggggaatt caacaacacg ccatcgccctg tgccaaacat	600
tttataggaa acgaacagga gcatTTtagg caagcgcccg agggcaatagg gtacaattac	660
actatagacg aaagcatcag tagcaacatc gacgataaaa cactacatga attgtactta	720
tggccattcc aggaacgagc ggctgcagga gtgggctctt tcatgtgcag ctataaccaa	780
gtgaacaact catacggttg tcaaaatagc aaacttatga atggcatcct gaaggacgag	840
ttgggcttcc aagggttctat tatgtctgac tgggcagcac aacatgcagg agtcgctacc	900
gcggtagccg gattagatat ggctatgcct ggggatacag cttttaattc aggcgatgacg	960
ttctggggta ccaatctaac agtagccgtc ttaaatggaa ctctgcccga atatagactg	1020
gatgatatgg ccatgagaat tatggcggct tttttcaaa gtcgggtttga gctgaatgag	1080
gtgccggaaa taaatTTtag ctcatggacg actgacaccg ttggccccct acaatactat	1140
gcaaaagaga acgttcaggt tattaatcag cacgttgatg tcagaagagg ccaagagcac	1200
ggaaagtTga ttagagaaat agcggcgaaa gccactgtct tgttgaagaa tgagggagcc	1260
ttgccgtTga agaagccgaa gtttttgga gtgattggcg aggaacccgg tcccaacttg	1320
tctggtccta acgggtgcag tgatcatgga tgtaacgagg gtaccctagg cgaggatgg	1380
ggatcaggga cctcaaaacta tccctatcta atcaccctg atcaagcgct acaagcgct	1440
gctgtagcgg agggTtctag atacgaatct attctttcaa attacgactt tgcggccaca	1500
acagctcttg tgaccagcc tgacgccacc gctattgtct ttgttaacgc agattcaggt	1560
gaaggttata ttgatgtggg aggtaacgaa ggcgacaggc agaacctgac gctgtggaac	1620
ggcggcgatg agttggTgaa aaacgttgcc gcaggtaata acaataccat agtagtaatc	1680
cattcagtag gtccggtcct tctggcggat atgaaaaata accctaatat caccgctatt	1740
gtatgggctg gtttgccctg acaggagtct ggaaattcaa ttactgatgt tctgtacggg	1800
gatgtcaacc ccggagggaa atccccattc acgtggggac ctactcgtga atcctatggc	1860
acggacgtcc tgtacgagcc aaacaacggt gaaggggctc ctcaagatga ctttagcgaa	1920

-continued

```

ggagtgttca tgcattatcg ttactttgat cgtgccacgt ctgggtccaa tgaacgtcc 1980
acgggtgcgg cgccagttta cccatttggg ttgggcttt ttatactac gtttgagtac 2040
tctaactctg tggttacgcc gaaggaagcg ggtgaataca caccaactac tgggtgtgact 2100
gagaaggcgc ctacgttttg gaactacagc accgaccccg ctgcatacgt ctttccctca 2160
ggtgaattca ggtacattta taatttcatt tatccgtacc taaatactac agatatatct 2220
aagtctgcga atgacctgcg gtatgggcag acggccgatg agttctacc acctaaggcg 2280
cttgagagcg ggccgcagcc aaaacatcct gctagcggtg cccctggggg taatccacaa 2340
ttgtgggacg ttttgatata cgttacgccg acgattacga ataaaggaga tgttgccggc 2400
gatgaggttg ccaattgta tgtgtctctg ggaggtccaa atgacccgtg taagggtgta 2460
agaggttttg acaggatttg gatagctcca ggggaaagtg ccaacttttac ggctgacatt 2520
acgcgtcgtg acttgagcaa ctgggacacc gtctcacaaa attgggtgat tagtaaatat 2580
ccaaagaagg tttgggtcgg aggatcatca cgtgagttgc cgtatccgc aagcctatga 2640

```

```

<210> SEQ ID NO 12
<211> LENGTH: 2838
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

```

```

<400> SEQUENCE: 12

```

```

atgagattcc catcatcttt tactgcgcgc ttgttcgcgc ctagtccgcg tttggtgct 60
ccagttaaca ctactactga agacgaaact gcccaaatcc cagccgaagc cgtcattggt 120
tacttggtat tagaagggtg tttcgatggt gctgttttgc cattctccaa ctccaccaac 180
aacggtttgc tatttatcaa caccactatc gtttccatcg ctgctaagga ggaagggtgc 240
tcttttgata agagagctgt taccactgaa agacaattac acaagagaga cttagcttac 300
tctctccag tttaccatc tccatggatg gacccaaaac cgcacggttg gactgacgct 360
tacgctaagg ctaaggactt cgtctctcaa ttgactttgt tagaaaagg taaactgacc 420
actggtgtcg gttggcaagg tgacttatgt gtccgtaacg tgggttctgt cccacgtttg 480
ggtctaagag gtttgtgtat gcaagacggt ccagttggtg ttagattctc tgactacaac 540
tcagtttttc catctggtca aaccgctgcc gccacttggg acagagaatt gatctacaga 600
agagctgaag ctattggttt tgaacacaga gctaagggtg tcgatgtcgt tttggctcca 660
gtcgtggtgc caattggcag agctcctgcc ggtggttaga actgggaagg tttctctagc 720
gaccatact tgactggtgt tgccatggcc gaatctgtca agggatttca acaacacgct 780
atcgttgtgt ctaagcactt cataggtaac gaacaagaac acttcagaca agctccagaa 840
gctatcggtt acaattacac catcgatgaa tctatttcct ccaacatcga cgataagacc 900
ttgcatgaat tgtacttgtg gccctttccaa gatgctgttg ctgctggtgt cggtctcttc 960
atgtgttctt acaatcaagt caacaactct tacggttgtc aaaactctaa gttgatgaac 1020
ggtattttga aagacgaatt aggtttccaa ggtttcatca tgtctgattg ggctgctcaa 1080
cacgctgggt ttgccaccgc cgttgcgtgt ttggacatgg ctatgccagg tgacaccgct 1140
tttaactctg gcctgacett ctgggggtact aatttgactg tcgctgttct aaacggtacc 1200
ttgccagaat accgttttga cgacatggct atgagaatca tggtgctttt cttcaagggt 1260
ggtttcgaac taaacgaagt tccagaaatt aacttttctt cttggactac tgacaccgct 1320
ggaccattgc aatactacgc taaggaaaac gttcaagtca ttaaccaaca cgttgatggt 1380

```

-continued

```

agaagaggtc aagagcacgg taagttgata agagaaattg ctgctaaagc caccgtattg 1440
ctaaaaaacg aaggtgcttt gcccttgaag aagccaaagt tcttggtgtg catcggtgaa 1500
gatgccggtc caaacttgtc tgggtccaaac ggttgctctg atcacggttg taacgaaggt 1560
accttggtg ctggttggtg ttctggtacc tccaactacc cctacctaat caccacagat 1620
caagctttgc aagctagagc tgcctgtgaa ggttccagat acgaatctat cttgtctaata 1680
tacgacttcg ctgccaccac cgttttggtc acccaaccag acgccacagc tatcgtcttc 1740
gtcaacgctg actcgggtga aggttacatc gacgttggtg gtaacgaagg tgacagacaa 1800
aacttgactt tgtggaacgg tggtgacgag ttggtcaaaa acgtcgtgctc tggaataaac 1860
aacaccatcg ttgtatttca ctctgtaggt ccagtcttgt tagccgatat gaagaacaac 1920
ccaaacatca ccgtatcgt ttgggtcgtt ttaccaggtc aagaaagtg taactctatc 1980
actgacgtct tgaacgggtg cgtcaacccc ggtggttaagt caccattcac ctgggggtccc 2040
actagagaat cctacggtac tgacgtcttg tacgaacct acaacgggtg aggtgcccc 2100
caagacgact tctccgaagg tgttttcac gactacagat acttcgatag ggctacttct 2160
ggttctaata aaacctccac tgggtccgcc ccagtttacc cattcggttt cggtttgtct 2220
tacaccacct tcgaatactc caacttggtc gttaccccaa aggaggccgg tgaatacact 2280
ccaactactg gtgtcactga aaaggccctt acctcgggtg actactctac cgaccagct 2340
gcttacgtct tcccatccgg tgaattcaga tacatttaca acttcactta cccatactta 2400
aacaccactg acatctccaa gtcgccaac gaccagcct acggtcaaac tgccgatgaa 2460
ttcttaccac caaaggcctt ggaaagtgtt ccacaaccaa aacatccagc ctctggtgct 2520
ccagtggtga atccacaatt gtgggacgtt ttgtacactg tcaactgtac catcacaat 2580
aagggtgacg tcgctggcga cgaagttgct caattatacg tgtctctagg tggtcctaac 2640
gatcctgtca aggttctacg tggtttcgat cgtattggtg tcgtcctgg tgaatctgct 2700
actttcaccg ctgacattac cagaagagat ttgtctaact gggacaccgt tagtcaaac 2760
tgggtcatct ctaagtaccc aaagaagggt tgggttggtg gttctagcag agaattgcct 2820
ttatctgctt ccttgtaa 2838

```

```

<210> SEQ ID NO 13
<211> LENGTH: 2640
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

```

```

<400> SEQUENCE: 13

```

```

atgtactcc aagcattcct tttctgtta gcaggatttg ctgcaaaaat ctctgtgca 60
gtaacaactg agagacaact acataagaga gacttggtt attctccacc tgtttacccc 120
tctccgtgga tggaccctaa tgctgacggt tggacggatg cttatgctaa agctaaggat 180
ttcgtaaacc aactgacgct gttagagaag gtaaacctaa cgactggcgt cggtggcag 240
ggtgatctat gtgtagggaa cgtggggagc gtgccaaggc tggggctaag aggtctgtgc 300
atgcaagatg gtcccgtagg aatacgttc agcgactaca actctgtctt tccttcaggt 360
caaacggcag cggccacatg ggacagagag cttatttacc gtggggcgga agcaatcgg 420
tttgaacaca gggcgaaagg agtagacgtc gtccatagccc cggtcgcagg gccataggt 480
agagccccag cagggggcag aaactgggaa ggattcagtt cagaccgta cctgacggga 540

```

-continued

gtggcgatgg ccgaatctgt aaaggggaatt caacaacacg ccategcctg tgccaaacat	600
tttataggaa acgaacagga gcatttttagg caagcgcccc aggcaatagg gtacaattac	660
actatagacg aaagcatcag tagcaacatc gacgataaaa cactacatga attgtactta	720
tggccattcc aggacgcagt ggctgcagga gtgggctctt tcatgtgcag ctataaccaa	780
gtgaacaact catacggttg tcaaaatagc aaacttatga atggcatcct gaaggacgag	840
ttgggcttcc aaggtttcat tatgtctgac tgggcagcac aacatgcagg agtcgctacc	900
gcggtagccg gattagatat ggctatgcct ggggatacag cttttaattc aggcgatgacg	960
ttctggggta ccaatctaac agtagccgtc ttaaatggaa ctctgccga atatagactg	1020
gatgatatgg ccatgagaat tatggcggtc tttttcaaag tcgggtttga gctgaatgag	1080
gtgccggaat taaattttag ctcatggacg actgacaccg ttggccccct acaatactat	1140
gcaaagaga acgttcagggt tattaatcag cacgttgatg tcagaagagg ccaagagcac	1200
ggaaagttag ttatagaaat agcggcgaaa gccactgtct tgttgaagaa tgagggagcc	1260
ttgccgttga agaagccgaa gtttttgga gtgattggcg aggacgccgg tcccaacttg	1320
tctggtccta acgggtgcag tgatcatgga tgtaacgagg gtaccctagg cgaggatgg	1380
ggatcaggga cctcaaaacta tccctatcta atcaccctg atcaagcgct acaagcgct	1440
gctgtagcgg agggttctag atacgaatct attctttcaa attacgactt tgcggccaca	1500
acagctcttg tgaccagcc tgacgccacc gctattgtct ttgttaacgc agattcagggt	1560
gaagggatata ttgatgtggg aggtaacgaa ggcgacagcg agaacctgac gctgtggaac	1620
ggcgcgcatg agttggtaga aaacgttgcc gcaggaata acaataccat agtagtaac	1680
cattcagtag gtccggctct tctggcggat atgaaaaata accctaatat caccgctatt	1740
gtatgggctg gtttgcttgg acaggagtct ggaaattcaa ttaactgatgt tctgtacggg	1800
gatgtcaacc ccggaggga atccccattc acgtggggac ctactcgtga atcctatggc	1860
acggacgtcc tgtacgagcc aaacaacggt gaaggggctc ctcaagatga ctttagcgaa	1920
ggagtgttca tcgattatcg ttactttgat cgtgccacgt ctgggtccaa tgaaacgtcc	1980
acgggtgcgg cgccagttta cccatttggg tttgggcttt cttatactac gtttgagtac	2040
tctaactctg tcgttacgcc gaaggaagcg ggtgaataca caccaactac tgggtgtgact	2100
gagaaggcgc ctacgttttg gaactacagc accgaccccg ctgcatacgt ctttccctca	2160
ggtgaattca ggtacattta taatttcatt tatccgtacc taaatactac agatatatct	2220
aagtctgcga atgacctgc gtatgggcag acggccgatg agttcctacc acctaaggcg	2280
cttgagagcg ggccgcagcc aaaacatcct gctagcggty cccctggggg taatccacaa	2340
ttgtgggacg ttttgtatag cgttaccgcc acgattacga ataaaggaga tgttgccggc	2400
gatgaggttg cccaattgta tgtgtctctg ggaggtccaa atgaccccg taagggtgcta	2460
agaggttttg acaggatttg gatagctcca ggggaaagtg ccacttttac ggctgacatt	2520
acgcgtcgtg acttgagcaa ctgggacacc gtctcacaaa attgggtgat tagtaaatat	2580
ccaaagaagg tttgggtcgg aggatcatca cgtgagttgc cgctatccgc aagcctatga	2640

<210> SEQ ID NO 14

<211> LENGTH: 2661

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic DNA

<400> SEQUENCE: 14

-continued

atgaagctaa actgggtcgc tgcggctcta tctatcggag ctgcaggaac ggatagtga	60
gtcgtctcttg cctccgctgt accagacact cttgctggag taaagaaggc tgacgccccaa	120
aaggtgggta ctagagatac actggcctac tccccaccac actatccctc tccgtggatg	180
gacccaaatg ctgtgggggtg ggaggaagcg tatgcgaagg caaagagttt tgtatcccaa	240
cttaccttaa tggaaaaagt gaatttaact acaggagtcg gttggcaagg agaaagatgt	300
gttggaatg tgggcagtat acccagacta ggcatgaggg gactatgtct tcaggacgga	360
ccgtgggaa ttaggtctgc cgactacaat tcagcgtttc cagctggtag taccgtggg	420
gcgtcatgga gcaaaagtct gtggtacgaa cgtgggctat tgatgggtac tgaatttaag	480
gaaaaggga tcgatatcgc attaggtcct gcgactggac cgttaggcag gaccgccgcc	540
ggtggaagaa attgggaggg atttacggtt gatccgtata tggccggta tgcaatggcg	600
gaagccgtga aaggcataca agacgccggc gtcatagcct gcgccaagca ctacatagcc	660
aatgaacaag aacattttcg tcaatctgga gaagtccaaa gcagaaagta caacatcagc	720
gaatccctat catccaactt agacgacaag actatgcacg agttatatgc ttggccgttc	780
gccgatgccg tcagagctgg agtaggttct gtcattgtct cctataacca gattaacaac	840
agttatggat gtcaaaactc caaactactg aatgggatct taaaagatga gatgggtttt	900
caaggctttg taatgtctga ctgggcccgc cagcacactg gtgcggccag tgcagttgca	960
ggactagata tgagtatgcc aggcgataca gcgtttgatt ctggctacag tttctggggt	1020
ggtaatctaa ctttggcagt gatcaatggg acagtaccgg cctggagggt agatgatatg	1080
gcgttgagaa ttatgtccgc gttcttcaag gtgggtaaaa ctatagaaga cctgcctgat	1140
ataaatTTTT catcttggaac aagggatacg tttggattcg tacatacgtt tgcacaagag	1200
aaccgtgaac aagtgaattt tgggggtcaac gtacagcacg accacaagag ccacattaga	1260
gaagccgcgg ccaagggtac agtgggtgta aaaaacacgg ggtcccttcc acttaaaaac	1320
ccaaagtttt tagcagtaat aggggaagac gcgggaccaa atcctgctgg tcccaacggc	1380
tcgggagatc gtgggtgtga taatggaacg cttgctatgg catggggatc cggcacaagc	1440
cagttccctt atcttatcac cctgatcaa ggattaagca accgtgccac acaagatggt	1500
accaggtatg aaagcatctt gaccaacaat gagtgggcaa gcgtacaagc gctggtttct	1560
caaccaacg tgacagcaat tgtcttcgca aacgctgatt ctggagaagg ttatattgag	1620
gtagatggaa atttcgggga tagaaaaaac ctgacctgtg ggcagcaggg cgacgagctt	1680
ataaagaacg tatcttccat atgtccta atcgatagtag tcctacatac agtaggtccc	1740
gtcttattag cggactatga gaagaatccc aacatcacag ccatagtttg ggcggttta	1800
ccagggcagg agagtggtaa cgcgatcgcc gatttgctat atggaaaagt ctctccgggc	1860
aggagcccat tcacctgggg aagaacgagg gaatcctatg gcacagaggt actatatgaa	1920
gccaataacg gcagaggcgc accacaagat gacttttagc aggggtctt catcgattat	1980
cgtcactttg ataaatacaa catcacgccc atatacgagt tcggtcattg tttatcctgg	2040
agcacattta agttctctaa cctgcacatt cagaagaata acgtcgggccc aatgtcacct	2100
ccaaatggta aaacgatagc tgcaccttcc ttaggaaatt tttccaagaa cctaaaggat	2160
tatggcttcc ccaaaaacgt aagaagaatt aaagagttaa tttacccta tctgaacact	2220
actacttccg gaaaggaagc ctcaggtgac gcacattacg ggcaaacgc caaggaattt	2280
ttgcggcgag gtgcgctaga tggttctccg cagcccggtt ccgcagctag cggcgagccc	2340

-continued

```

ggtggaaaca gacagctgta tgatatttta tataccgtaa ccgctactat cacaaatata 2400
ggttcagtta tggatgacgc cgttccccag ctatatctgt cccatggggg gccaaatgaa 2460
cctccgaaag tgctgagagg ttttgataga atagagagaa ttgcaccggg tcagagcgtt 2520
actttcaagg cagacttgac cagaagggat ttaagtaact gggatacaaa aaaacaacaa 2580
tgggtcatca ccgattaccg gaagactggt tacgttggt caagctccag agacttgccg 2640
ttaagcgcca gactgccgta g 2661

```

```

<210> SEQ ID NO 15
<211> LENGTH: 2859
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

```

```

<400> SEQUENCE: 15

```

```

atgagattcc catctatctt caccgcgctc ttgttcgctg cctcttctgc tttggtgcc 60
ccagttaaca ccactactga agatgaaacc gctcaaatcc cagctgaagc tgctattggt 120
tacttggtact tagaagggtg tttcgatggt gctgttctac cattctctaa ttctaccaac 180
aacggcttgt tggtcatcaa caccactatc gtttccatcg ctgccaaagg agaaggtgtc 240
tctttggata aaagagctgt cgctttggcc tctgcggtc cagacactct agccggtgtc 300
aagaaggctg acgccccaaa ggtcgtgact agagacactt tggcttacag cccaccacac 360
taccatctc cttggatgga cccaaacgct gtcggttggg aagaagccta cgtaaggcc 420
aagtccttcg tctctcaatt gaccttgatg gaaaagggtc atttaaccac cggtgttggt 480
tggcaagggt aaagatgtgt cggtaacggt ggatccatcc ctagattggg tatgctggt 540
ttgtgtttgc aagaagggtc attgggtatc agattgtctg actacaactc agctttccct 600
gctggtacta ccgcggtgtc ttcttggtct aagtcctgtt ggtacgaaag aggtttgttg 660
atgggtaccg aatttaagga aaagggtatt gatatcgctt tgggtcctgc cacaggtcct 720
ttgggtagaa ctgctgctgg tggtagaaac tgggaagggt tcaccgtcga cccatacatg 780
gctggccacg ctatggccga agccgttaag ggtatccaag acgctggtgt tatcgcttgt 840
gctaagcact acattgcaa cgaacaagaa cactttagac aatctggtga agtccaatcc 900
agaaagtaca acatttctga atccttgtcc tccaatttag acgataagac tatgcacgag 960
ctatacgctt ggcccttcgc tgacgctggt agagctggtg ttggttcctg catgtgttcc 1020
tacaacaaaa ttaacaactc ttacggttgt caaaactcta agttgttgaa cggtattttg 1080
aaagacgaaa tgggtttcca aggtctcgtg atgtctgact gggccgctca acacactggt 1140
gctgcttccg ctgttctggt tctagacatg tctatgccag gtgatactgc cttcgactct 1200
ggttactctt tctgggtggt taacctaaac ttagctgtca ttaacggtac cgttccggcc 1260
tggagagtct atgacatggc tttgagaatc atgtctgctt tcttcaaggt tggttaagact 1320
atcgaagact tgcagacat taacttctct tcctggactc gtgacacctt cggtttcgtt 1380
cacactttcg ccaagaaaa cagggaacaa gtcaacttcg gtgtaaatgt tcaacacgac 1440
cataagtccc acatcagaga agctgctgct aagggttccg tcgttttgaa gaacaccggt 1500
tccttgccat tgaagaaccc aaagtttttg gctgtaatcg gtgaggacgc tgggtccaaac 1560
ccagctggtc caaatggttg cggtgataga ggttgatgata acggtactct agctatggct 1620
tggggttccg gcacttctca attcccttac ctaattaccc cagatcaagg tttatctaac 1680
agagctactc aagacggtac tagatacgaa tctattttga ctaataacga gtgggcttct 1740

```

-continued

```

gtccaagccc tagtttccca accaaaagtc accgctatcg ttttcgctaa cgccgactct 1800
gggtgaaggtt acattgaagt tgacggcaac tttggtgaca gaaagaactt gactttgtgg 1860
caacaaggtg acgaactaat caagaacgtt tcttctatct gtccaaacac catcgctcgtt 1920
ttgcataccg ttggtccagt tctattggct gactacgaaa agaaccctaaa cattaccgct 1980
atcgtttggg ctggtctacc aggtcaagaa tctggtaacg ccacgcccga tttgctatac 2040
ggtaaggttt cccaggtcgt tctccattc acctggggta gaaccagaga atccacggtt 2100
accgaagttt tgtacgaagc caacaacggt agaggtgctc cgcaagatga cttcagttaa 2160
gggtgttttca tcgactacag acacttcgat aagtacaaca ttactcccat ctacgaattc 2220
ggtcacggtt tatcctggtc caccttcaag ttcagtaact tgcacatcca aaagaacaat 2280
gtcggtccta tgtctccacc taacggcaag accatcgctg ctccatcttt gggaaacttc 2340
tccaaaaact tgaaggacta cggtttccca aagaacgtcc gtcgtattaa ggaattcatt 2400
taccataact taaacaccac tacctccggt aaggaagcct ctggtgatgc tctacacgga 2460
caaatgctta aagaattctt gccagctggt gccttggtat gttctccaca accaagatct 2520
gccgcttctg gtgaaccagg tggtaacaga caattatacg atatcttgta caccgtcact 2580
gccaccatca ccaacactgg ttcgctcatg gatgatgctg tcccacaatt atactgttca 2640
cacggtggtc caaacgaacc accaaaggtc ttgagaggtt tcgacagaat cgaaagaatt 2700
gccccaggtc aatctgttac attcaaggct gacttgacta gaagagactt atctaactgg 2760
gacaccaaga agcaacaatg ggctcattacc gattacccaa aaacgctcta cgttggttct 2820
tcttccagag atttgccatt gtcagctaga ttgccataa 2859

```

```

<210> SEQ ID NO 16
<211> LENGTH: 2661
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

```

```

<400> SEQUENCE: 16

```

```

atgtactacc aagcattcct tttctgttta gcaggatttg ctgccccaaat ctctgctgct 60
gtcgcttttg cctctgccgt tccagacact ctagccggtg tcaagaaggc tgacgcccac 120
aaggctcgtg ctagagacac tttggcttac agcccaccac actaccctac tccttggtatg 180
gacccaaaag ctgtcggttg ggaagaagcc tacgctaagg ccaagtcctt cgtctctcaa 240
ttgaccttga tggaaaaggc caatttaacc accggtgttg gttggcaagg tgaaagatgt 300
gtcggtaacg ttggatccat ccttagattg ggtatgcgtg gtttgtgttt gcaagacggt 360
ccattgggta tcagattgtc tgactacaac tcagctttcc ctgctggtac taccgcccgt 420
gcttcttggt ctaagtcctt gtggtacgaa agaggtttgt tgatgggtac cgaatttaag 480
gaaaagggtt ttgatatcgc tttgggtcct gccacaggtc ctttgggtag aactgctgct 540
ggtggttaga actgggaagg ttccaccgtc gaccataca tggctggcca cgctatggcc 600
gaagccgtta aggttatcca agacgctggt gttatcgctt gtgctaagca ctacattgcc 660
aacgaacaag aacactttag acaatctggt gaagtccaat ccagaaagta caacatttct 720
gaatccttgt cctccaattt agacgataag actatgcacg agctatacgc ttggcccttc 780
gctgacgctg ttgagctggt tgttggttcc gtcattgtgt cctacaacca aattaacaac 840
tcttacggtt gtcaaaactc taagttgttg aacggtatct tgaaagacga aatgggtttc 900

```

-continued

caaggcttcg tgatgtctga ctgggccgct caacacactg gtgtgtcttc cgtgttgtgt	960
ggtctagaca tgtctatgcc aggtgatact gccttcgact ctggttactc tttctgggggt	1020
ggtaacctaa ccttagctgt cattaacggt accgttccgg cctggagagt cgatgacatg	1080
gctttgagaa tcatgtctgc tttcttcaag gttggtgaaga ctatcgaaga cttgccagac	1140
attaacttct cttcctggac tctgtacacc ttcggtttcg ttcacacttt cgcccaagaa	1200
aacagggaac aagtcaactt cgggtgtaaat gttcaacacg accataagtc ccacatcaga	1260
gaagctgctg ctaagggttc cgtcgttttg aagaacaccg gttccttgcc attgaagaac	1320
ccaaagtttt tggctgtaat cggtgaggac gctggtccaa acccagctgg tccaaatggt	1380
tgcggtgata gaggttgtga taacggtact ctatctatgg cttgggggttc cggcacttct	1440
caattccctt acctaattac ccagatcaa ggtttatcta acagagctac tcaagacggt	1500
actagatacg aatctatttt gactaataac gagtgggctt ctgtccaagc cctagtttcc	1560
caaccaaagc tcaccgctat cgttttcgct aacgccgact ctggtgaagg ttacattgaa	1620
gttgacggca actttggtga cagaaagaac ttgactttgt ggcaacaagg tgacgaacta	1680
atcaagaacg tttcttctat ctgtccaaac accatcgctg ttttgcatag cgttggtcca	1740
gttctattgg ctgactacga aaagaacca aacattaccg ctatcgtttg ggctggtcta	1800
ccaggtcaag aatctggtaa cgccatcgcc gatttgctat acggtgaagg ttcccaggt	1860
cgttctccat tcacctgggg tagaaccaga gaatcctacg gtaccgaagt tttgtacgaa	1920
gccacaacg gttagaggtgc tccgcaagat gacttcagtg aaggtgtttt catcgactac	1980
agacacttcg ataagtacaa cattactccc atctacgaat tcggtcacgg tttatcctgg	2040
tccaccttca agttcagtaa cttgcacatc caaaagaaca atgtcggtcc tatgtctcca	2100
cctaacggca agaccatcgc tgtcccatct ttgggaaaact tctccaaaaa cttgaaggac	2160
tacggtttcc caaagaacgt ccgtcgattt aaggaattca tttaccata cttaaacacc	2220
actacctccg gtaagggaag cctctggtgat gctcactacg gacaaactgc taaagaattc	2280
ttgccagctg gtgccttgga tggttctcca caaccaagat ctgccgcttc tgggtgaacca	2340
ggtggttaaca gacaattata cgatatcttg tacaccgtca ctgccaccat caccaacact	2400
ggttccgtca tggatgatgc tgtcccacaa ttatacttgt cacacggtgg tccaaacgaa	2460
ccaccaaagg tcttgagagg tttcgacaga atcgaaagaa ttgcccagg tcaatctgtt	2520
acattcaagg ctgacttgac tagaagagac ttatctaact gggacaccaa gaagcaacaa	2580
tgggtcatta ccgattaccc aaaaaccgtc tacgttggtt cttcttcag agatttgcca	2640
ttgtcagcta gattgccata a	2661

<210> SEQ ID NO 17
 <211> LENGTH: 2622
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic DNA

<400> SEQUENCE: 17

atgagattct ctggtatcgt cgctacccta gtcgtggtg ccggtgtctc tgetcaccia	60
ggtgattact ctaagttgga aagaagagct gtcgtacct ccgaaccaca ctaccacaaa	120
ccatggatga acccagatgc cgacggttg caagaagctt acgtcaagge taaggacttc	180
gtttctcaaa tgacctgtt ggaaaaggtc aacttaacta ctggtgtcgg ttgggcttcc	240
gatttggtg tcggtaacgt cgggtgctgt ccacgtttgg gtttgagatc cctgtgctta	300

-continued

caagactctc caactgggtg cagattcgct gactggggtt ccgtgttccc agctgggtatc	360
accaccgggtg ccaccttcga caagggtcta atgtaccgta gaggtcaagc tatgggtcaa	420
gaagctaagg acaagggtat aaacgttttg ttgggtccag tcgccgggtg tttgggcaga	480
gttgctgctg gtggtagagc ttgggaatcc ttcggtgctg acccagtctt aactggttac	540
ggtatgattg aaactattaa gggatttcaa gacactggcg ttatcgctac tgctaagcac	600
ttcatcggta acgaacaaga acacttcaga caagttgggtg aagaacgtgg tagagggtgc	660
aacatctctg aatccttgtc ctctaacatt gatgataaga ctatgcatga actatacttg	720
tggccattcg ccgacgctgt tagagccggg gtcggctctg ttatgtgttc ctacactcaa	780
gttaacaact cctacgggtg tcaaaactct aagttgttga acggtctatt gaaggatgaa	840
ttaggcttcc aaggtttcgt tatgtccgac tggcaagctc aacacacggg tgccgcctct	900
gctgctgctg gtttggacat gtctatgcct ggtgacactg aattcaacac cggctctatcc	960
ttctgggggtg ccaacttgac tttggctgtt gttaacggca ccgttgccga atggagaatc	1020
gatgacatgg ctatgagaat catggctgct ttcttcaaag tcggtaacac cctagaccaa	1080
ccagaaatca acttctcttc ctggaccaag gatactttcg gtccattgca ctcttctagt	1140
ggtaacagaa tccaacaat taaccaacac gttgacgtca gaagagatca cggtaaccta	1200
atcagagaag ttgctgctaa gggtaactgtc ttgttgaaga acaccaacaa cgctttgcc	1260
ttgaataagc caaagttctt ggctgtcatt ggtgatgatg ctggttctaa ccccagaggt	1320
cctaacgggt gtccagatag aggatgtttg ttgggtacct tgggtatggc ttggggttct	1380
ggcactgctg acttcccata cttaattacc ccagacgctg ccttgcaagc ccaagccatc	1440
gaagacggta cccgttacga atctatcttg tctaattacg ccaccgccc aactcaagct	1500
ttggtctctc aaacatacgc cactgctatt gttttcgtcg ctgcctcttc cggcgaaggt	1560
tacatcgatt ttgacggcaa caagggtgac agaaacaatt tgactttgtg gtacgacgg	1620
gactctttgg ttaagaacgt ttcttctgtc tgtaacaaca ccatcgttgt tatccactct	1680
actggtccta ccactctgac cgaatggtag gacaacccta atgtcacgcg tatcgtctgg	1740
gctggtgttc cagggtcaaga atctggtaga gccatcactg acgttttgta cggtagagtt	1800
aaaccagctg gtcgttctcc attcaccctg ggttaagacta gagaatccta cggtagctgat	1860
gttatgtaca agccaaaacaa cggtaacgaa gctccacaac aagactacac cgaagggtgt	1920
ttcattgact acagacactt tgaccaacaa aaggatgaac cagtctacga atttgggtcac	1980
ggtttgtcct acacaacctt cgaatacagt aacatcaggg ttgataaggc tccagcctct	2040
gaatacaaac caactactgg tcaaacctac ccagctcctg ttttcggtgc taacgtttct	2100
aaggatttga gtcaatacac ttcccatcc gacgaatttc cacacatcta cttgtttatc	2160
taccgctact tgaacacctc ctcttctggt gaagaagctt ccagagaccc aaagtaacgg	2220
ggtaccgctg aggaattctt gccacctaag gccttagacg gttccccaca accattgcc	2280
agagcttctg gtaagaactc tccagggtgt aacagacaat tgtaacgacac cttatacact	2340
gttactgcc	2400
ccattaccaa taccggttaag ttggttggtg aagaagttcc acaactatac	2400
gtttccacg gtggtccaga agaccaccca gttgtgttgc gtggtttcga aagaattaga	2460
ttagaccag gtcaatccgc tactttcaag gttgatttga ccagaagaga cgtctctaac	2520
tgggacgtca aggtccaaga ctgggtcctc tccgagcctc caaagaaggt cttcgttggt	2580
tcctcctcta gaaagttgca cttgtctgct gacttgaact aa	2622

-continued

```

<210> SEQ ID NO 18
<211> LENGTH: 2823
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

<400> SEQUENCE: 18

atgagattcc catctatttt caccgctgtc ttgttcgctg cctcttctgc cttggtgct 60
ccagtcaata ccaccactga agacgaaacc gctcaaattc cagccgaagc tgttattggt 120
tacttggtgact tggaagggtga cttcgatgtc gctgttttgc cattctccaa ctccaccaac 180
aacggcttgc tattcatcaa caccactatc gcttctattg ctgctaagga agaagggtgt 240
tccttggaaca agcgtcatcc tgggtgattac tccaagttgg aaagaagagc tgtcgctacc 300
tccgaacctc actaccccca accatggatg aaccagacg ctgacggttg gcaagaagct 360
tacgtgaagg ccaaggattt cgtctcccaa atgacctat tagaaaagg taaacttaact 420
accggtgttg gttgggcctc tgatctatgt gtcggtaacg tcggtgctgt tccaagatta 480
ggcttgagat ctttgtgttt acaagattct ccaaccggtg tccgtttcgc tgactgggtt 540
tctgtcttcc cagctggcat caccactggt gctaccttcg acaagggttt aatgtacaga 600
agaggccaag ctatgggtca agaagccaag gacaaaggta tcaacgtctt gttgggtcca 660
gtcgccggtg gtttgggtag agttgctgct ggtggtagag cctgggagtc tttcggtgcc 720
gatccagtct tgacagggtta cggtatgatt gaaacaatta agggtatcca agacactggt 780
gttatcgcta ccgctaaaca cttcattggt aacgaacaag aacactttag acaagttggt 840
gaagaaagag gtagagggtg taacatctcc gaatccttgt cctctaacat tgatgacaag 900
actatgcacg aattgtactt atggccattc gctgacgctg ttcgtgcggg tgcgggttcc 960
gttatgtgtt cctacaccca agttaacaac tcttacggtt gtcaaaactc taagttattg 1020
aacggtttat tgaaggacga attgggtttc caagggttcg ttatgtccga ctggcaagct 1080
caacacacgg gtgctgcttc tgtgcccggc ggtttggata tgtccatgcc aggtgacacc 1140
gaattcaaca ctggactatc tttctggggt gctaacttga ccttggtgtg cgtaacgggt 1200
actgtcgctg aatggagaat agacgacatg gccatgagaa tcatggctgc tttctttaag 1260
gtcggtaaca ctttggatca acccgaaatc aacttctctt cttggactaa ggacactttc 1320
ggtccattgc actcttcttc tggtaacaga atccaacaaa taaaccaaca cgtggacggt 1380
agaagagacc acggttaactt aatcagagaa gttgccgcca aaggcacagt ttgctaaag 1440
aacactaaca acgcttttgc attgaacaag ccaaagtttt tggctgtcat aggtgatgat 1500
gctggtagca acccacgtgg tccaaatggt tgcccagaca gaggttggtt attgggtacc 1560
ttgggtatgg cctggggttc tggtagcgct gacttcccat acttaatcac tccagacgcc 1620
gctttgcaag ctcaagccat cgaagatggt actaggtacg aatccatcct gtccaactac 1680
gctactgtc aaacccaagc tttggtttct caaacctacg ctaccgccat cgtcttcgtc 1740
gctgcttcta gtggtgaagg ttacatcgat ttcgacggtg acaagggtga tagaaataac 1800
ttaactttgt ggtagacgg tgactctttg gtcaagaatg tttcctcgt ttgtaacaac 1860
accatcgctg taattcactc taccggccct actattttga ccgagtggta cgacaaccca 1920
aacgtaacgg ctatgctctg ggcgggtgtt ccgggtcaag aatctggtag agctatcact 1980
gacgttctat acggtagagt taacctgct ggtagatctc cattcacttg gggtaaagacc 2040
agagaatctt acggtactga cgtcatgtac aagccaaaca acggtaacga agctccacaa 2100

```

-continued

```

caagactaca ctgaagggtgt cttcatcgac tacagacact tcgaccaaca aaaggatgaa 2160
cctgtttacg aattcgggtca cggtttgtct tacaccacct tcgaatactc caacattaga 2220
gttgacaagg cccagcctc tgaatacaag ccaactaccg gtcaaaccat cccagctcct 2280
gtttttgggt ccaacgtttc aaaggatttg agccaataca ctttcccatc tgatgaattc 2340
cctcacatct acttggttcat ttaccatac ctaaatactt cctcctccgg tgaagaagct 2400
tctagagatc caaagtacgg tggcacgct gaagaattcc taccaccaa ggctttggac 2460
ggttcaccac aaccattgcc aagagcctcc ggtaagaact ctcccggtgg caatcgtaa 2520
ttgtacgaca ccttgtagac tgtaactgct accatcacta aactggtaa gttggtcggc 2580
gaggaagtcc cacaattgta cgtctctcac ggtggccctg aagaccacc agtcgtcttg 2640
agaggtttcg aaagaattag actagatcca ggtcaatctg ccacottcaa ggtcgacct 2700
acccgtagag atgtttctaa ctgggacgtt aaggccaag actgggttat ctccgaacac 2760
ccaaagaaag tttttgttgg ttcttcttcc cgtaagttgc attgtctgc cgacctaaac 2820
taa 2823

```

```

<210> SEQ ID NO 19
<211> LENGTH: 1623
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic DNA

```

```

<400> SEQUENCE: 19

```

```

atgagataca gaactgcgc tgccttgcc ttggccacg gtccattgc tagagctgat 60
agccactcca cttccggtgc ttcagctgaa gctgtgtcc caccagctgg tactccatgg 120
ggtactgctt acgataaagc caaggctgct ttggccaagc taaacttgca agacaaggtc 180
ggtattgtct ccggtgttgg ttggaacggc ggtccatgtg ttggaaacac atcccagct 240
tccaagattt cttacccttc tctatgttg caagacggtc cattgggtgt cagatactct 300
actggttcca ctgctttcac cccaggtgtt caagccgct ctacttggga cgttaacttg 360
atcagagaaa gaggtcaatt tatcgagaa gaagtcaagg cttctggtat ccacgttatt 420
ttaggtccag ttgctggtcc attgggtaag acccctcaag gtgtagaaa ctgggaaggt 480
ttcggcgtcg acccatactt gaccggtatc gccatgggtc aaaccatcaa cggtattcaa 540
tccgttgggt ttcaagccac tgccaagcac tacatcttga atgaacaaga attgaaccgt 600
gaaactatct cctccaacc tgacgataga actttgcacg aattgtacac ctggccattc 660
gctgatgccg tccaagctaa cgtcgcttcc gtaatgtgta gttacaacaa ggtaaacacc 720
acttgggctt gcgaagatca atacacatta caaacgtct tgaaggacca attgggtttc 780
ccaggttacg tcatgactga ctggaacgcc caacacacca ccgttcaaag tgccaattct 840
ggtttggaca tgtctatgcc aggtaccgac ttcaacggt acaacagact atggggtcca 900
gctttgacta acgctgttaa ctctaacc aa gttccaacct ccagagtcca cgacatggtt 960
actagaatct tggcgcgttg gtacttgacc ggtcaagatc aagccggtta cccatcttct 1020
aacatttcta gaaacgtcca aggtaaccac aagactaacg ttagagccat cgctagagac 1080
ggtatcgtct tgttgaagaa cgacgctaac atcttaccat tgaaaaagcc agcttccatc 1140
gctgttgtcg gttctgtctg tatcattggt aatcacgcta gaaactcccc atcctgtaac 1200
gacaagggtt gtgacgatgg tgccttaggt atgggttggg gttccggtgc tgtcaattac 1260

```

-continued

ccatacttcg ttgtccata cgtgctatc aataccagag cttcctctca aggtacacaa	1320
gttaccttgt ctaacaccga taacacctct tccggtgctt ctgcgcgtcg tggtaaggat	1380
gtcgctattg tcttcattac tgcgactct ggtgaagggt acatcacgt cgaaggtaac	1440
gctggtgacc gtaacaactt ggacccttgg cacaatggta acgctttggt tcaagcgtt	1500
gctggtgcta actctaactg tatcggtgc gtccattctg tgggtgctat tatcttagaa	1560
caaatcttgg cctaccaca agttaaggct gtcgtctggg ctggtttgcc atcccaagaa	1620
taa	1623

<210> SEQ ID NO 20
 <211> LENGTH: 1821
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic DNA

<400> SEQUENCE: 20

atgagattcc catctatctt cactgccgtc ttgttcgctg ccagttccgc ttggtgtgcc	60
ccagttaaca ccactaccga agatgaaacc gctcaaatcc ctgctgaagc tgcacgtggt	120
tacttagact tggagggtga cttcgacgtg gccgtcttgc cttctcttaa ctetaccaac	180
aacggtttgc tattcattaa cactaccatt gcttctatcg ccgccaagga agaaggtgtt	240
tctttggaca agagagactc tcaactccacc tctggtgctt ccgccaagc tgtgtgtccc	300
ccagctggta ctccatgggg taccgcttac gacaaggcta aggtgcttt ggccaagttg	360
aaactgcaag acaagggttg tctcggttcc ggtgtcggtt ggaacggtgg tccatgtgtc	420
ggtaatacct caccagcttc taagatctcc tacccttcat tgtgtttgca agacggccct	480
ttgggtgtca gatactcaac tggctctacc gccctcactc caggtgttca agctgcttcc	540
acttgggatg ttaacttgat cagagagaga ggtcaattca tcggtgaaga agtcaaggct	600
tctggtatc atgtcatcct aggtccagtt gccggtccat tgggcaagac cccacaagg	660
ggtagaaact gggagggttt cgggtgtggc ccatacttga ctggtatcgc tatgggtcaa	720
accatcaacg gtatccaatc tgcggtgtt caagctaccg ctaagcacta cttttgaa	780
gaacaagaat taaacagaga aacaatttct tctaattccag acgacagaac cttacacgaa	840
ctatacacct ggccatttgc cgaagctgtg caagccaacg ttgcctctgt tatgtgtct	900
tacaacaagg ttaacaccac ttgggtctgt gaagaccaat acactttgca aactgtttta	960
aaagatcaat tgggtttccc aggttacctg atgactgatt ggaacgctca acacactacc	1020
gtccaatccg ctaactctgg ttggacatg tccatgccag gtactgattt caacggtaac	1080
aaaccgttgt ggggtccagc ttgactaac gctgtcaact ccaaccaagt cccaacctcc	1140
agagttgacg acatgggtac acgtatcttg gctgcttgg acctaactgg tcaagaccaa	1200
gctggttacc catctttcaa catcagtaga aacgttcaag gtaaccacaa gaccaacgtc	1260
agagctattg cttagagatg tctcgtcttg ttaagaacg atgctaatat cttgccattg	1320
aagaagcccc cttctattgc tgtgtgtggt tccgctgcc tcatgtgtaa ccacgctaga	1380
aactctccat cttgtaacga caagggttgt gatgacggtg ctttgggtat ggggtgggt	1440
agcgtgtcgt tcaactaccc atactttgtc gctccatacg acgtatcaa caccagagct	1500
tcttctcaag gtacccaagt tacctgttcc aacactgaca acacttcctc tggcgctct	1560
gccgctagag gtaaggacgt tgcctatcgt ttcacaccg ccgattcttg tgaaggttac	1620
attacgctcg aaggtaacgc cggatgata aacaacttgg acccatggca caacggtaac	1680

-continued

```
gccttggtcc aagctgttgc tggtgctaac tccaatgtca tcgttgctgt tcaactctgtc 1740
ggagctatta ttttgaaca aatcttagct ttgcctcaag ttaaagctgt tgtttgggct 1800
ggtttgccat cccaagaata a 1821
```

What is claimed is:

1. Modified yeast cells derived from parental yeast cells, the modified yeast cells comprising a genetic alteration that causes the modified yeast cells to produce an increased amount of β -glucosidase polypeptides compared to the parental yeast cells, wherein the modified yeast cells produce during fermentation more ethanol and/or less acetate compared to the amount of ethanol and acetate produced by otherwise identical parental yeast cells, wherein the genetic alteration comprises introduction of an expression cassette for expressing a β -glucosidase polypeptide, and wherein the β -glucosidase polypeptides have:

- a) an amino acid sequence having at least 80% amino acid sequence identity to SEQ ID NO: 1 or SEQ ID NO: 2;
- b) the amino acid sequence of a mature polypeptide encoded by a nucleic acid having at least 80% nucleic acid sequence identity to any of SEQ ID NOs: 11-16; or
- c) the amino acid sequence of a mature polypeptide encoded by a nucleic acid that hybridizes, under stringent hybridization conditions comprising 65° C. and 0.1×SSC, to any of SEQ ID NOs: 11-16, or the complement, thereof.

2. The modified yeast cells of claim 1, further comprising an exogenous gene encoding a carbohydrate processing enzyme, wherein the carbohydrate processing enzyme is a dehydrogenase, a transketolase, a phosphoketolase, a transaldolase, an epimerase, a phytase, a xylanase, a β -glucanase, a phosphatase, a protease, an α -amylase, a β -amylase, a glucoamylase, a pullulanase, an isoamylase, a cellulase, a trehalase, a lipase, a pectinase, a polyesterase, a cutinase, an oxidase, a transferase, a reductase, a hemicellulase, a mannanase, an esterase, an isomerase, a pectinases, a lactase, a peroxidase, or a laccase.

3. The modified yeast cells of claim 1, further comprising an exogenous PKL pathway.

10 4. The modified yeast cells of claim 1, further comprising a reduction or elimination of an endogenous NAD-dependent 3-phosphate dehydrogenase (GPD); a disruption of one or more of genes GPD1, GPD2, GPP1, and/or GPP2; overexpression of glycerol dehydrogenase (GCV1); and/or
15 overexpression of dihydroxyacetone kinase (DAK1).

5. The modified yeast cells of claim 1, wherein the modified yeast cells further make reduced amounts of DP2 and/or DP3 compared to otherwise identical parental yeast cells.

20 6. The modified yeast cells of claim 1, wherein the β -glucosidase polypeptides are derived from *Glomerella graminicola*.

7. The modified yeast cells of claim 1, wherein the cells are of a *Saccharomyces* spp.

25 8. A method for increasing the production of alcohol and/or decreasing the production of acetate from yeast cells grown on a carbohydrate substrate, comprising: introducing into parental yeast cells a genetic alteration that increases the production of β -glucosidase polypeptides compared to the
30 amount produced in the parental yeast cells, wherein the cells having the introduced genetic alteration are the modified yeast cells of claim 1.

9. The method of claim 8, wherein the increased production of alcohol is at least 0.5%, at least 1.0%, at least 2.0%
35 or at least 3.0%.

10. The method of claim 8, wherein the decreased production of acetate is at least 1.0%, at least 2.0%, at least 4.0% or at least 6.0%.

11. The method of claim 8, wherein the modified yeast cells further comprise an exogenous PKL pathway.

40 12. The modified yeast cells of claim 1, further comprising a heterologous acetyl-CoA synthase gene and/or increased expression of an endogenous acetyl-CoA synthase gene.

* * * * *