

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent
Kind Code
Date of Patent
Inventor(s)

12396461
B2
August 26, 2025
Abe; Madoka et al.

Functional peptides having antimicrobial activity against phytopathogenic microorganisms

Abstract

The disclosure describes antimicrobial peptides exhibiting activity against a wide variety of plant pathogens are described. Functional peptides consisting of the amino acid sequence: MKVMIVVKTKVKNKKVAKMMVK (SEQ ID NO: 1) or an amino acid sequence having an identity of 85% or more to the amino acid sequence of SEQ ID NO: 1, or consisting of a partial peptide thereof are described.

Inventors:	Abe; Madoka (Nagoya, JP), Kondo; Satoshi (Miyoshi, JP), Shimada; Takehiko (Nagoya, JP), Hanada; Kosuke (Fukuoka, JP), Mochida; Keiko (Nagoya, JP), Tsuge; Takashi (Kasugai, JP)
Applicant:	TOYOTA JIDOSHA KABUSHIKI KAISHA (Toyota, JP); KYUSHU INSTITUTE OF TECHNOLOGY (Kitakyushu, JP); NATIONAL UNIVERSITY CORPORATION TOKAI NATIONAL HIGHER EDUCATION AND RESEARCH SYSTEM (Nagoya, JP); CHUBU UNIVERSITY EDUCATIONAL FOUNDATION (Kasugai, JP)
Family ID:	1000008779562
Assignee:	TOYOTA JIDOSHA KABUSHIKI KAISHA (Toyota, JP); KYUSHU INSTITUTE OF TECHNOLOGY (Kitakyushu, JP); NATIONAL UNIVERSITY CORPORATION TOKAI NATIONAL HIGHER EDUCATION AND RESEARCH SYSTEM (Nagoya, JP); CHUBU UNIVERSITY EDUCATIONAL FOUNDATION (Kasugai, JP)
Appl. No.:	17/436853
Filed (or PCT Filed):	March 06, 2020
PCT No.:	PCT/JP2020/009812
PCT Pub. No.:	WO2020/184468

PCT Pub. Date: September 17, 2020

Prior Publication Data

Document Identifier	Publication Date
US 20220174961 A1	Jun. 09, 2022

Foreign Application Priority Data

JP	2019-042748	Mar. 08, 2019
----	-------------	---------------

Publication Classification

Int. Cl.: C07K14/00 (20060101); A01N63/50 (20200101); A01P1/00 (20060101); A61K38/00 (20060101); A61K38/08 (20190101); A61K38/10 (20060101); A61P1/00 (20060101); C07K7/00 (20060101); C07K7/06 (20060101); C07K7/08 (20060101); C07K14/415 (20060101)

U.S. Cl.:

CPC A01N63/50 (20200101); A01P1/00 (20210801); C07K14/415 (20130101);

Field of Classification Search

CPC: C07K (14/415); C07K (14/00); C07K (7/06); C07K (7/00); C07K (7/08); A61K (38/08); A61K (38/00); A61K (38/10); C12N (15/8282); A61P (1/00); A01H (6/825); A01N (63/50)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4940840	12/1989	Suslow et al.	N/A	N/A
7041808	12/2005	Tomie et al.	N/A	N/A
7465783	12/2007	Gallois et al.	N/A	N/A
7713531	12/2009	Takakura	N/A	N/A
8865967	12/2013	English et al.	N/A	N/A
2012/0054911	12/2011	English et al.	N/A	N/A
2015/0224182	12/2014	Hunt	435/325	A61K 35/17

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2004-329215	12/2003	JP	N/A
2006-512052	12/2005	JP	N/A
4257119	12/2008	JP	N/A
5988200	12/2015	JP	N/A
03/033532	12/2002	WO	N/A

OTHER PUBLICATIONS

Primary Examiner: Ha; Julie

Attorney, Agent or Firm: Sughrue Mion, PLLC

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS

(1) This application is a National Stage of International Application No. PCT/JP2020/009812, filed Mar. 6, 2020, claiming priority based on Japanese Patent Application No. 2019-042748, filed Mar. 8, 2019.

TECHNICAL FIELD

(2) The present invention relates to a functional peptide exhibiting antimicrobial activity against plant pathogens such as plant pathogenic filamentous fungi, and also to an agricultural chemical comprising the functional peptide, and a method for controlling plant pathogens using the functional peptide.

BACKGROUND ART

(3) A reduction in crop yield due to plant disease contributes to the food problem. In order to increase the crop yield, it is necessary to prevent plants from plant pathogens by using pesticides. In recent years, what is called, environmental conservation-type agriculture, in which the used amounts of agricultural chemicals or chemical fertilizers comprising chemical substances as main ingredients have been reduced, has been promoted. Thus, it has been desired to develop a substance suitable for such environmental conservation-type agriculture, which is used instead of chemical substances.

(4) Such a substance is, for example, a peptidic substance consisting of 100 or less amino acids. Among others, defensin has been known as a peptide having antimicrobial activity that is derived from plants. It has been reported that defensin is a basic protein containing a large amount of cysteine and is present in many plant species. However, defensin is problematic in that it needs to maintain its higher-order structure in order to express antimicrobial activity, and also in that, since it is a relatively long peptide consisting of 40 or more amino acids, it is difficult to synthesize defensin and thus is expensive. Non Patent Literature 1 discloses the results obtained by synthesizing a partial peptide of rice-derived defensin and then studying the antimicrobial activity thereof. However, the partial peptide of defensin disclosed in Non Patent Literature 1 has exhibited only effects on rice blast disease, damping-off disease, rice Bakanae disease.

(5) Moreover, Patent Literature 1 discloses that a defensin gene derived from cabbage or Japanese mustard spinach belonging to *Brassicaceae* is introduced into a plant, so that combined disease resistance can be imparted to the plant. Patent Literature 2 discloses isolation of a peptide having antimicrobial activity derived from *Arabidopsis*. Patent Literature 3 discloses a peptide having antimicrobial activity against *Magnaporthe oryzae*, which is extracted from *Pleurotus cornucopiae* var. *citrinopileatus* that is one type of edible mushroom. Patent Literature 4 discloses 10 to 16 partial peptides having antimicrobial activity, which have been designed from rice-derived defensin-like proteins having antimicrobial activity. Patent Literature 5 discloses isolation of a peptide having antimicrobial activity derived from *Arabidopsis*. Patent Literature 6 discloses an antimicrobial peptide consisting of 40 to 66 amino acids, which has been prepared from the body

fluid of rhinoceros beetle larva. Patent Literature 7 discloses a method of performing DNA shuffling on a defensin gene to obtain an antimicrobial peptide against plant pathogens.

CITATION LIST

Patent Literature

(6) PTL 1: JP 2004-329215 A PTL 2: JP 2006-512052 A PTL 3: Japanese Patent No. 4257119 PTL 4: Japanese Patent No. 5988200 PTL 5: Japanese Patent No. 4445855 PTL 6: WO 2003/033532 PTL 7: U.S. Pat. No. 8,865,967

Non Patent Literature

(7) NPL 1: J. Pestic. Sci. 42(4), 172-175, 2017

SUMMARY OF INVENTION

Technical Problem

(8) However, a short functional peptide that has antimicrobial activity against a wide variety of plant pathogens and is easily synthesized has not yet been known. Hence, it is an object of the present invention to consider the aforementioned circumstances and to provide a functional peptide having antimicrobial activity against a wide variety of plant pathogens, a nucleic acid encoding the functional peptide, an agricultural chemical comprising the functional peptide, and a method for controlling plant pathogens using the functional peptide.

Solution to Problem

(9) In order to achieve the aforementioned object, the present inventor has conducted intensive studies. As a result, the present inventor has identified a peptide having antimicrobial activity against a wide variety of plant pathogens, from among peptides that have been specifically expressed in the xylem sap of tomatoes inoculated with wilt disease pathogens, thereby completing the present invention.

(10) The present invention encompasses the following. (1) A functional peptide consisting of the amino acid sequence: MKVMIVVKTKVKNKKVAKMMVK (SEQ ID NO: 1) or an amino acid sequence having an identity of 85% or more with respect to the amino acid sequence, or consisting of 5 or more consecutive amino acids in the amino acid sequence as set forth in SEQ ID NO: 1. (2) The functional peptide according to the above (1), wherein the 5 or more consecutive amino acids are within a range of 1 to 15 amino acid residues counted from the N-terminus of the amino acid sequence as set forth in SEQ ID NO: 1. (3) The functional peptide according to the above (1), wherein it consists of one amino acid sequence selected from the group consisting of the following (a) to (g): (a) amino acid sequence: MKVMIVVK (SEQ ID NO: 2), (b) amino acid sequence: KVMIVVKT (SEQ ID NO: 3), (c) amino acid sequence: VMIVVKTK (SEQ ID NO: 4), (d) amino acid sequence: MKVMI (SEQ ID NO: 5), (e) amino acid sequence: KVMIV (SEQ ID NO: 6), (f) amino acid sequence: VMIVV (SEQ ID NO: 7), and (g) amino acid sequence: VAKMM (SEQ ID NO: 8). (4) A nucleic acid encoding the functional peptide according to any one of the above (1) to (3). (5) A recombinant vector having the nucleic acid according to the above (4). (6) A transformant having the recombinant vector according to the above (5). (7) The transformant according to the above (6), wherein it is a plant cell or a plant body transformed with the recombinant vector. (8) A composition exhibiting sterilizing action or antimicrobial action against plant pathogens, comprising the functional peptide according to any one of the above (1) to (3). (9) An agricultural chemical comprising the functional peptide according to any one of the above (1) to (3). (10) The agricultural chemical according to the above (9), which is for sterilizing or antimicrobial use against plant pathogens. (11) The agricultural chemical according to the above (10), wherein the plant pathogens are one or more plant pathogenic filamentous fungi selected from the group consisting of *Botrytis cinerea*, *Colletotrichum destructivum*, *Colletotrichum orbiculare*, *Fusarium oxysporum*, and *Pyricularia oryzae*. (12) The agricultural chemical according to the above (10), wherein the plant pathogen is *Xanthomonas oryzae* pv. *oryzae* or *Pseudomonas syringae* pv. *tomato*.

Advantageous Effects of Invention

(11) The functional peptide according to the present invention has excellent antimicrobial activity against a wide variety of plant pathogens, and also, can be easily synthesized. As such, the functional peptide according to the present invention can be used, for example, as a composition or an agricultural chemical having sterilizing action or antimicrobial action against plant pathogens.

(12) In addition, by using a nucleic acid encoding the functional peptide according to the present invention, the functional peptide can be expressed in a host. A plant cell or a plant body expresses the above-described functional peptide, so that resistance to a wide variety of plant pathogens can be improved.

Description

BRIEF DESCRIPTION OF DRAWINGS

- (1) FIG. 1 shows photographs showing the results obtained by evaluating the antimicrobial activity of TsORF6-1 against five types of plant pathogenic filamentous fungi.
- (2) FIG. 2 is a photograph showing the results obtained by inoculating TsORF6-1, together with *Botrytis cinerea*, into the leaves of tomatoes.
- (3) FIG. 3 is a photograph showing the results obtained by inoculating TsORF6-1, together with *Colletotrichum destructivum*, into the leaves of Chinese cabbages.

DESCRIPTION OF EMBODIMENTS

(4) <Functional Peptide>

(5) The functional peptide according to the present invention consists of the amino acid sequence as set forth in SEQ ID NO: 1, MKVMIVVKTKVKNKKVAKMMVK, or an amino acid sequence having an identity of 85% or more with respect to the aforementioned amino acid sequence. Herein, peptides, which are contained in xylem saps collected from tomatoes infected with wilt disease and xylem saps collected from non-infected tomatoes, have been comprehensively analyzed, and as a result, the amino acid sequence as set forth in SEQ ID NO: 1 has been identified as a peptide having antimicrobial activity against a wide variety of plant pathogens. That is to say, the functional peptide according to the present invention, at least, has a function such as antimicrobial activity against plant pathogens. Accordingly, the functional peptide according to the present invention can be used as an agricultural chemical or an agricultural chemical composition, based on its antimicrobial activity against plant pathogens.

(6) The amino acid sequence having an identity of 85% or more with respect to the amino acid sequence as set forth in SEQ ID NO: 1 specifically means an amino acid sequence comprising a deletion, substitution or addition of 1 to 3 amino acids with respect to the amino acid sequence as set forth in SEQ ID NO: 1. The functional peptide according to the present invention has an identity of preferably 90% or more, and more preferably 95% or more, to the amino acid sequence as set forth in SEQ ID NO: 1. It is highly likely that an amino acid sequence having this range of identity to the amino acid sequence as set forth in SEQ ID NO: 1 exhibits antimicrobial activity against plant pathogens, as with the peptide consisting of the amino acids shown in SEQ ID NO: 1.

(7) In particular, the amino acid sequence comprising a deletion, substitution or addition of 1 to 3 amino acids with respect to the amino acid sequence as set forth in SEQ ID NO: 1 is not particularly limited, but the position in which 1 to 3 amino acids are deleted, substituted or added is preferably a position close to the C-terminus. More specifically, the position in which 1 to 3 amino acids are deleted, substituted or added is a position in a range from 11 to 22, preferably from 14 to 22, and more preferably 16 to 22, counted from the N-terminus. The position in which 1 to 3 amino acids are deleted, substituted or added is set to be within this range, so that antimicrobial activity against plant pathogens can be maintained at high.

(8) It is to be noted that an amino acid sequence different from the amino acid sequence as set forth in SEQ ID NO: 1 can be produced by introducing a desired mutation into a nucleic acid encoding

the amino acid sequence as set forth in SEQ ID NO: 1, for example, using a mutation introduction kit utilizing site-specific mutagenesis (e.g., Mutant-K (manufactured by Takara Bio, Inc.) or Mutant-G (manufactured by Takara Bio, Inc.)), or using LA PCR in vitro Mutagenesis Series Kit of Takara Bio, Inc.

(9) By the way, the functional peptide according to the present invention is not limited to a functional peptide consisting of the amino acid sequence as set forth in SEQ ID NO: 1 (a length of 22 amino acid residues), but it may also consist of consecutive 5 or more amino acids from the amino acid sequence as set forth in SEQ ID NO: 1. Hereafter, the peptide consisting of consecutive 5 or more amino acids from the amino acid sequence as set forth in SEQ ID NO: 1 is also referred to as a partial peptide in some cases. An example of this partial peptide may be, but is not limited to, a peptide consisting of consecutive 5 to 15 amino acids, preferably consecutive 5 to 10 amino acids, more preferably consecutive 5 to 8 amino acids, and further preferably consecutive 8 amino acids, selected from the amino acid sequence as set forth in SEQ ID NO: 1.

(10) In particular, such a partial peptide is selected from a range from 1 to 15 amino acid residues, preferably a range from 1 to 13 amino acid residues, and more preferably a range from 1 to 10 amino acid residues, counted from the N-terminus of the amino acid sequence as set forth in SEQ ID NO: 1. When the partial peptide has an amino acid sequence in the aforementioned range, it can exhibit more excellent antimicrobial activity.

(11) More specifically, the partial peptide is preferably one selected from the group consisting of the following (a) to (g). The following partial peptides can exhibit particularly excellent antimicrobial activity against plant pathogens: (a) amino acid sequence: MKVMIVVK (SEQ ID NO: 2), (b) amino acid sequence: KVMIVVKT (SEQ ID NO: 3), (c) amino acid sequence: VMIVVKT (SEQ ID NO: 4), (d) amino acid sequence: MKVMI (SEQ ID NO: 5), (e) amino acid sequence: KVMIV (SEQ ID NO: 6), (f) amino acid sequence: VMIVV (SEQ ID NO: 7), and (g) amino acid sequence: VAKMM (SEQ ID NO: 8).

(12) The functional peptide according to the present invention (including the aforementioned partial peptides) can be easily produced according to conventionally known methods. Examples of the method of synthesizing a peptide may include a solid phase synthesis method and a liquid phase synthesis method. With regard to the solid phase synthesis method, a Fmoc method (fluorenylmethyloxycarbonyl method), a tBoc method (t-butyloxycarbonyl method), etc. is applied, so that a functional peptide having a desired amino acid sequence can be synthesized.

(13) Moreover, the functional peptide according to the present invention can also be produced by producing a nucleic acid encoding the amino acid sequence, and then applying a genetic engineering method using the produced nucleic acid. That is, for example, a nucleic acid encoding the aforementioned functional peptide is chemically synthesized, the nucleic acid is then incorporated into an expression vector, and the thus obtained expression vector is then introduced into suitable host cells, so as to obtain a transformant. This transformant is cultured and is then allowed to generate a functional peptide, and the generated functional peptide may be then recovered from the transformant and/or the medium.

(14) The expression vector is not particularly limited, as long as it has a nucleic acid encoding a functional peptide and can express the functional peptide in any given host cells. Examples of the expression vector may include a plasmid vector and a viral vector. Besides, the host cells are not particularly limited, and bacteria such as *Escherichia coli* or *Bacillus subtilis*, and fungi such as yeast or filamentous fungi can be utilized. Moreover, as such host cells, animal cells such as insect cells or mammalian cells may be used. For example, when *Escherichia coli* are used as host cells, an expression vector comprising, at least, a promoter region, a start codon, a nucleic acid encoding the functional peptide according to the present invention, a stop codon, a terminator region and a replication origin can be used.

(15) The aforementioned transformant is cultured, and thereafter, the functional peptide according to the present invention can be recovered from the transformant and can be then purified. The

method of isolating and purifying a functional peptide is not particularly limited. Examples of the method of isolating and purifying a functional peptide may include, but are not limited to: methods of utilizing solubility, such as salting-out and a solvent precipitation method; methods of utilizing a difference in molecular weights, such as dialysis, ultrafiltration, gel filtration, and sodium dodecyl sulfate-polyacrylamide gel electrophoresis; methods of utilizing electric charges, such as ion exchange chromatography and hydroxylapatite chromatography; methods of utilizing specific affinity, such as affinity chromatography; methods of utilizing a difference in hydrophobicity, such as reverse phase high performance liquid chromatography; and methods of utilizing a difference in isoelectric points, such as isoelectric point electrophoresis.

(16) Furthermore, the functional peptide according to the present invention can also be produced, for example, as a histidine-tagged or Flag-tagged functional peptide, by a transformant. In this case, by using a substance having affinity for such a tag, the functional peptide can be more easily isolated and purified.

(17) Further, the functional peptide according to the present invention can be synthesized by utilizing, what is called, a cell-free protein synthesis system. In such a cell-free protein synthesis system, an extract, for example, from *Escherichia coli*, rabbit reticulocytes, or wheat germs can be utilized.

(18) <Antimicrobial Activity of Functional Peptide>

(19) The functional peptide according to the present invention has antimicrobial activity against a wide variety of plant pathogens. Herein, the antimicrobial activity includes both the activity of preventing infection with plant pathogens, and sterilizing activity against plant pathogens. Examples of plant pathogens, against which the functional peptide according to the present invention exhibits antimicrobial activity, may include a wide range of plant pathogenic filamentous fungi. More specifically, examples of the plant pathogens may include plant pathogenic filamentous fungi belonging to the genus *Botrytis*, plant pathogenic filamentous fungi belonging to the genus *Colletotrichum*, plant pathogenic filamentous fungi belonging to the genus *Fusarium*, and plant pathogenic filamentous fungi belonging to the genus *Pyricularia*.

(20) Examples of the plant pathogenic filamentous fungi belonging to the genus *Botrytis* may include *Botrytis aclada*, *Botrytis byssoides*, *Botrytis cinerea*, *Botrytis convoluta*, *Botrytis diospyri*, *Botrytis elliptica*, *Botrytis fabae*, *Botrytis galanthina*, *Botrytis gladiolorum*, *Botrytis paeoniae*, *Botrytis polyblastis*, *Botrytis* sp., *Botrytis squamosa*, and *Botrytis tulipae*.

(21) Examples of the plant pathogenic filamentous fungi belonging to the genus *Colletotrichum* may include *Colletotrichum actinidiicola*, *Colletotrichum acutatum*, *Colletotrichum aenigma*, *Colletotrichum ampelopsidis*, *Colletotrichum belamcandium*, *Colletotrichum boninense sensu lato*, *Colletotrichum capsici*, *Colletotrichum carthami*, *Colletotrichum caudatum*, *Colletotrichum chrysanthemi*, *Colletotrichum circinans*, *Colletotrichum coccodes*, *Colletotrichum coffeanum*, *Colletotrichum corchori*, *Colletotrichum crassipes*, *Colletotrichum cypripedii*, *Colletotrichum daphnicola*, *Colletotrichum dematium*, *Colletotrichum destructivum*, *Colletotrichum durionis*, *Colletotrichum echinocloeae*, *Colletotrichum elasticae*, *Colletotrichum fatsiae*, *Colletotrichum fiorinae*, *Colletotrichum fuscum*, *Colletotrichum gloeosporioides*, *Colletotrichum godetiae*, *Colletotrichum graminicola*, *Colletotrichum hibisci*, *Colletotrichum higginsianum*, *Colletotrichum horii*, *Colletotrichum hydrangeae*, *Colletotrichum kahawae*, *Colletotrichum karstii*, *Colletotrichum koyasuensis*, *Colletotrichum lappae*, *Colletotrichum liliacearum*, *Colletotrichum lili*, *Colletotrichum lindemuthianum*, *Colletotrichum malvarum*, *Colletotrichum medicaginis-denticulatae*, *Colletotrichum metake*, *Colletotrichum moricola*, *Colletotrichum morinum*, *Colletotrichum musae*, *Colletotrichum nigrum*, *Colletotrichum nymphaeae*, *Colletotrichum orbiculare*, *Colletotrichum panacicola*, *Colletotrichum paniculatae*, *Colletotrichum pehkinense*, *Colletotrichum phaseolorum*, *Colletotrichum sansevieriae*, *Colletotrichum sasicola*, *Colletotrichum siamense*, *Colletotrichum sophorae-japonicae*, *Colletotrichum spaethianum*, *Colletotrichum spinaciae*, *Colletotrichum sublineolum*, *Colletotrichum tabacum*, *Colletotrichum*

theobromicola, *Colletotrichum trichellum*, *Colletotrichum trifolii*, *Colletotrichum tropicale*, *Colletotrichum truncatum*, *Colletotrichum villosum*, and *Colletotrichum yoshinoi*.

(22) Examples of the plant pathogenic filamentous fungi belonging to the genus *Fusarium* may include *Fusarium acuminatum*, *Fusarium ananatum*, *Fusarium anguioides*, *Fusarium arthrosporioides*, *Fusarium asiaticum*, *Fusarium avenaceum*, *Fusarium brasiliense*, *Fusarium commune*, *Fusarium conglutinans* var. *betae*, *Fusarium cuneirostrum*, *Fusarium decemcellulare*, *Fusarium dimerum* var. *dimerum*, *Fusarium foetens*, *Fusarium fujikuroi*, *Fusarium graminearum*, *Fusarium guttiforme*, *Fusarium lactis*, *Fusarium lagenariae*, *Fusarium lateritium*, *Fusarium merismoides*, *Fusarium oxysporum*, *Fusarium pallidoroseum*, *Fusarium pallidum*, *Fusarium phaseoli*, *Fusarium phyllophilum*, *Fusarium poae*, *Fusarium proliferatum*, *Fusarium redolens*, *Fusarium ricini*, *Fusarium roseum*, *Fusarium solani*, *Fusarium striatum*, *Fusarium subglutinans*, and *Fusarium verticillioides*.

(23) Examples of the plant pathogenic filamentous fungi belonging to the genus *Pyricularia* may include *Pyricularia grisea*, *Pyricularia higginsii*, *Pyricularia oryzae*, *Pyricularia panici*, and *Pyricularia zingiberis*.

(24) Other than these, examples of the plant pathogens may include plant pathogens belonging to the genus *Alternaria*, plant pathogens belonging to the genus *Cladosporium*, plant pathogens belonging to the genus *Claviceps*, plant pathogens belonging to the genus *Sclerotinia*, plant pathogens belonging to the genus *Septoria*, plant pathogens belonging to the genus *Pseudoperonospora*, and plant pathogens belonging to the genus *Puccinia*.

(25) In particular, examples of plant pathogenic filamentous fungi, against which the functional peptide according to the present invention exhibits high antimicrobial activity, may include *Botrytis cinerea*, *Colletotrichum destructivum*, *Colletotrichum orbiculare*, *Fusarium oxysporum*, and *Pyricularia oryzae* (that is a synonym for *Magnaporthe oryzae*).

(26) Moreover, the functional peptide according to the present invention has antimicrobial activity, not only against plant pathogenic filamentous fungi, but also against plant pathogenic bacteria such as *Xanthomonas oryzae* pv. *oryzae* and *Pseudomonas syringae* pv. *tomato*.

(27) Examples of the plant pathogenic bacteria belonging to the genus *Xanthomonas* may include *Xanthomonas campestris* pv. *malloti*, *Xanthomonas axonopodis* pv. *phaseoli*, *Xanthomonas alfalfae* pv. *alfalfae*, *Xanthomonas arboricola* pv. *pruni*, *Xanthomonas translucens* pv. *poae*, *Xanthomonas campestris* pv. *fici*, *Xanthomonas oryzae* pv. *oryzae*, *Xanthomonas oryzae* pv. *oryzicola*, *Xanthomonas axonopodis* pv. *phaseoli*, *Xanthomonas translucens* pv. *translucens*, *Xanthomonas arboricola*, *Xanthomonas campestris* pv. *raphani*, *Xanthomonas campestris* pv. *campestris*, *Xanthomonas cucurbitae*, *Xanthomonas citri* subsp. *citri*, *Xanthomonas hortorum* pv. *hederae*, *Xanthomonas axonopodis* pv. *manihotis*, *Xanthomonas arboricola* pv. *juglandis*, *Xanthomonas campestris* pv. *nigromaculans*, *Xanthomonas albilineans*, *Xanthomonas axonopodis* pv. *vasculorum*, *Xanthomonas campestris*, *Xanthomonas cucurbitae*, *Xanthomonas axonopodis* pv. *glycines*, *Xanthomonas campestris* pv. *cannabis*, *Xanthomonas translucens* pv. *phleipratensis*, *Xanthomonas theicola*, *Xanthomonas vesicatoria*, *Xanthomonas euvesicatoria*, *Xanthomonas hortorum* pv. *carotae*, *Xanthomonas axonopodis* pv. *allii*, *Xanthomonas arboricola* pv. *celebensis*, *Xanthomonas axonopodis* pv. *ricini*, *Xanthomonas hortorum* pv. *pelargonii*, *Xanthomonas translucens* pv. *cerealis*, *Xanthomonas campestris* pv. *mangiferaeindicae*, *Xanthomonas vasicola* pv. *holcicola*, *Xanthomonas axonopodis* pv. *lespedezae*, *Xanthomonas translucens* pv. *cerealis*, *Xanthomonas hortorum*, *Xanthomonas axonopodis* pv. *Vitians*, and *Xanthomonas citri* subsp. *Malvacearum*.

(28) As mentioned above, since the functional peptide according to the present invention has antimicrobial action against a wide variety of plant pathogens, it can control plant diseases caused by plant pathogens, against which the present functional peptide exhibits antimicrobial action. Specific examples of the plant diseases, which can be controlled by the functional peptide according to the present invention, may include rice diseases such as rice blast disease, sesame

blight disease, sheath blight disease, rice Bakanae disease, and bacterial leaf blight. In addition, other examples of the plant diseases, which can be controlled by the functional peptide according to the present invention, may include, are not particularly limited to: powdery mildew, *fusarium* blight, rust, *Microdochium nivale*, *Typhula incarnata*, smut, bare smut, stinking smut, eye spot, leaf blight, blight, *Pyrenophora tritici-repentis*, *Rhynchosporium secalis*, *Pyrenophora teres*, *Cochliobolus sativus*, *Pyrenophora graminea*, seedling damping-off disease caused by *Rhizoctonia*, zonate leaf spot, *Puccinia polysora*, gray leaf spot disease, *Cercospora kikuchii*, anthracnose, melanose, brown spot, target spot, *sclerotinia* rot, anthracnose, *Alternaria solani*, scarlet rot, powdery scab, and mildew.

(29) <Agricultural Chemical and Composition Each Comprising Functional Peptide>

(30) Since the functional peptide according to the present invention has antimicrobial action against a wide variety of plant pathogens, it can be used as an agricultural chemical or an agricultural chemical composition. Herein, the agricultural chemical composition means a composition that can be used upon the production of an agricultural chemical. For example, the functional peptide according to the present invention can be utilized as a germicide having germicidal action against plant pathogens, in particular, against plant pathogenic filamentous fungi. More specifically, a functional peptide compound may be directly used as an agricultural chemical, but in general, the functional peptide compound may be mixed with a suitable solid carrier, liquid carrier, surfactant or other formulation auxiliary agents, so that it can be used in any given dosage form such as an emulsion, an EW preparation, a liquid agent, a suspending agent, a water-dispersible powder, a water-dispersible granule, a powder agent, a DL powder agent, a microparticle agent, a microparticle agent F, a granule, a tablet, an oil agent, an aerosol, a flowable agent, a dry flowable agent, and a microcapsule.

(31) Examples of the solid carrier may include: animal and plant powders, such as starch, activated carbon, soy flour, wheat flour, wood flour, fish flour, and powdered milk; and inorganic matter powders, such as talc, kaolin, bentonite, calcium carbonate, zeolite, diatomaceous earth, white carbon, clay, alumina, ammonium sulfate, and urea.

(32) Examples of the liquid carrier may include: water; alcohols such as isopropyl alcohol and ethylene glycol; ketones such as cyclohexanone and methylethylketone; ethers such as dioxane and tetrahydrofuran; aliphatic hydrocarbons such as kerosene and light oil; aromatic hydrocarbons such as xylene, trimethylbenzene, tetramethylbenzene, methylnaphthalene, and solvent naphtha; halogenated hydrocarbons such as chlorobenzene; acid amides such as dimethylacetamide; esters such as glycerin esters of fatty acids; nitriles such as acetonitrile; and sulfur-containing compounds such as dimethyl sulfoxide.

(33) Examples of the surfactant may include alkylbenzene sulfonic acid metal salts, dinaphthyl methane disulfonic acid metal salts, alcohol sulfuric acid ester salts, alkylaryl sulfonate, lignin sulfonate, polyoxyethylene glycol ether, polyoxyethylene alkylaryl ether, and polyoxyethylene sorbitan monoalkylate.

(34) Examples of other auxiliary agents that can be used herein may include: fixing agents or thickeners, such as carboxymethyl cellulose, gum Arabic, sodium alginate, guar gum, gum tragacanth, and polyvinyl alcohol; defoaming agents, such as metal soap; physical property improving agents, such as fatty acid, alkyl phosphate, silicone, and paraffin; and coloring agents.

(35) In general, various types of germicidal preparations or diluted products thereof can be applied by commonly used application methods, such as spraying (e.g., spraying, misting, atomizing, dusting, granule application, submerged application, box application, etc.), soil application (e.g., mixing into soil, irrigation, etc.), surface application (e.g., application, dressing, coating, etc.), immersion, use of poison baits, and smoke application.

(36) Furthermore, needless to say, a pesticide comprising, as an active ingredient, the functional peptide according to the present invention is sufficiently effective, even when the present functional peptide is used alone. However, as necessary, the functional peptide can be mixed with,

or can be used in combination with other fertilizers, agricultural chemicals such as insecticides, acaricides or nematocides, other pesticides, antiviral agents, attractants, herbicides, and plant growth regulators, and the like.

(37) The thus configured agricultural chemical can be applied to plants that may develop plant diseases due to the aforementioned plant pathogens. Examples of the target plants may include, but are not particularly limited to: agricultural crops, such as corn, rice, wheat, barley, rye, oats, sorghum, cotton, soybeans, peanuts, soba, sugar beet, rape, sunflower, sugar cane, tobacco, solanaceous vegetables (eggplant, tomato, pimiento, pepper, potato, etc.), cucurbitaceous vegetables (cucumber, pumpkin, zucchini, watermelon, melon, squash, etc.) cruciferous vegetables (Japanese radish, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, mustard greens, broccoli, cauliflower, etc.), asteraceae vegetables (burdock, crown daisy, artichoke, lettuce, etc.), liliaceae vegetables (leek, onion, garlic, asparagus, etc.), apiaceous vegetables (carrot, parsley, celery, red wolf, etc.), chenopodiaceae vegetables (spinach, chard, etc.), mint vegetables (Japanese basil, mint, basil, etc.), strawberry, sweet potato, yam, and taro; flowers; foliage plants; turfs; fruit trees, such as pome fruits (apple, pear, Japanese pear, quince, marmelo, etc.), stone fruits (peach, plum, nectarine, Japanese plum, cherry, apricot, prune, etc.), citrus fruits (citrus unshiu, orange, lemon, lime, grapefruit, etc.), nuts (chestnuts, walnuts, hazelnuts, almonds, pistachios, cashew nuts, macadamia nuts, etc.), berry fruits (blueberry, cranberry, blackberry, raspberry, etc.), grapes, persimmon, olive, loquat, banana, coffee, date palm, coconuts, etc.; and trees other than fruit trees, such as tea, mulberry, *Jatropha* (*Jatropha curcas*), flower trees, roadside trees (ash, birch, dogwood, *eucalyptus*, ginkgo, lilac, maple, oak, poplar, *Cercis chinensis*, *Liquidambar formosana*, plane tree, Japanese zelkova, *Thuja standishii*, fir tree, *Tsuga sieboldii*, *Juniperus rigida*, pine tree, spruce tree, and yew tree).

(38) Among others, the aforementioned agricultural chemical comprising the present functional peptide as an active ingredient is preferably applied to cruciferous plants, cucurbitaceous plants, and gramineous plants.

(39) <Transformed Plant>

(40) As mentioned above, the functional peptide according to the present invention has antimicrobial action against a wide variety of plant pathogens. Thus, a transformed plant that expresses the present functional peptide at a high level is characterized in that it exhibits resistance to the aforementioned plant diseases. The transformed plant includes both a transformed plant cell prepared by expressly introducing a nucleic acid encoding the aforementioned functional peptide into a predetermined plant cell, and a transformed plant body prepared by regenerating the transformed plant cell in a plant body. The transformed plant according to the present invention expresses the aforementioned functional peptide, so that the resistance of the transformed plant to the aforementioned plant pathogens is improved.

(41) More specifically, examples of the transformed plant may include, but are not particularly limited to: agricultural crops, such as corn, rice, wheat, barley, rye, oats, sorghum, cotton, soybeans, peanuts, soba, sugar beet, rape, sunflower, sugar cane, tobacco, solanaceous vegetables (eggplant, tomato, pimiento, pepper, potato, etc.), cucurbitaceous vegetables (cucumber, pumpkin, zucchini, watermelon, melon, squash, etc.) cruciferous vegetables (Japanese radish, turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, mustard greens, broccoli, cauliflower, etc.), asteraceae vegetables (burdock, crown daisy, artichoke, lettuce, etc.), liliaceae vegetables (leek, onion, garlic, asparagus, etc.), apiaceous vegetables (carrot, parsley, celery, red wolf, etc.), chenopodiaceae vegetables (spinach, chard, etc.), mint vegetables (Japanese basil, mint, basil, etc.), strawberry, sweet potato, yam, and taro; flowers; foliage plants; turfs; fruit trees, such as pome fruits (apple, pear, Japanese pear, quince, marmelo, etc.), stone fruits (peach, plum, nectarine, Japanese plum, cherry, apricot, prune, etc.), citrus fruits (citrus unshiu, orange, lemon, lime, grapefruit, etc.), nuts (chestnuts, walnuts, hazelnuts, almonds, pistachios, cashew nuts, macadamia nuts, etc.), berry fruits (blueberry, cranberry, blackberry, raspberry, etc.), grapes, persimmon, olive,

loquat, banana, coffee, date palm, coconuts, etc.; and trees other than fruit trees, such as tea, mulberry, *Jatropha* (*Jatropha curcas*), flower trees, roadside trees (ash, birch, dogwood, *eucalyptus*, ginkgo, lilac, maple, oak, poplar, *Cercis chinensis*, *Liquidambar formosana*, plane tree, Japanese zelkova, *Thuja standishii*, fir tree, *Tsuga sieboldii*, *Juniperus rigida*, pine tree, spruce tree, and yew tree).

(42) It is to be noted that a method for producing the transformed plant according to the present invention is not particularly limited, and as a summary, a transformed plant expressing the aforementioned functional peptide can be produced according to a method comprising incorporating a nucleic acid encoding the aforementioned functional peptide into an expression vector, and then introducing the obtained expression vector into a plant.

(43) The expression vector is configured to comprise a promoter enabling expression in a plant and a nucleic acid encoding the aforementioned functional peptide. As vectors used to produce the present expression vector, various types of conventionally known vectors can be used. For example, a plasmid, a phage, a cosmid or the like can be used, and a suitable vector can be selected from these, as appropriate, depending on a plant cell into which the vector is to be introduced, or an introduction method. Specific examples of such a vector may include vectors such as pBR322, pBR325, pUC19, pUC119, pBluescript, pBluescript SK, and pBI. In particular, when the method of introducing a vector into a plant body is a method using *Agrobacterium*, a pBI binary vector is preferably used. Specific examples of the pBI binary vector may include pBIG, pBIN19, pBI101, and pBI121.

(44) The promoter is not particularly limited, as long as it is a promoter enabling expressing a nucleic acid encoding the above-described functional peptide in a plant body. A known promoter can be preferably used. In particular, as a promoter, it is preferable to use a constant expression promoter constantly expressing a gene located downstream thereof in a plant body. Examples of such a promoter may include a cauliflower mosaic virus 35S promoter (CaMV35S), various types of actin gene promoters, various types of ubiquitin gene promoters, a napaline synthase gene promoter, a tobacco PR1a gene promoters, a tomato ribulose 1,5-diphosphate carboxylase oxygenase small subunit gene promoter, and a napin gene promoter. Among these promoters, a cauliflower mosaic virus 35S promoter, an actin gene promoter, or a ubiquitin gene promoter can be more preferably used. Using each of the above described promoters, when the promoter is introduced into a plant cell, any given gene can be strongly expressed therein.

(45) Moreover, as such a promoter, a promoter having the function of site-specifically expressing any given gene in a plant can also be used. As such promoters, any conventionally known promoters can be used. Using such promoters, the above-described functional peptide can be site-specifically expressed.

(46) Furthermore, as a promoter, a gene promoter inducing the expression of the gene in response to disease can be used. Examples of such a promoter may include a plant defensin 1.2 (PDF1.2) gene promoter, a pathogenesis-related 1 (PR1) gene promoter, a PR2 gene promoter, and a PR5 gene promoter. Using such a promoter, the above described functional peptide can be expressed specifically to the time of pathogenic infection.

(47) It is to be noted that the expression vector may comprise other DNA segments, in addition to a promoter and a nucleic acid encoding the above-described functional peptide. Such other DNA segments are not particularly limited, but examples thereof may include a terminator, a selective marker, an enhancer, and a nucleotide sequence for enhancing translation efficiency. In addition, the above-described expression vector may further have a T-DNA region. The T-DNA region can enhance the efficiency of gene introduction, when the above-described recombinant expression vector is introduced into a plant body, particularly using *Agrobacterium*.

(48) The transcription terminator is not particularly limited, as long as it functions as a transcription termination site, and a known transcription terminator may be used. For example, specifically, a transcription termination region of a napaline synthase gene (Nos terminator), a transcription

termination region of cauliflower mosaic virus 35S (CaMV35S terminator), and the like can be preferably used. Among these, a Nos terminator can be more preferably used. By disposing a transcription terminator at a suitable position in the above-described expression vector, generation of a phenomenon, by which an unnecessary long transcriptional product is synthesized, can be prevented after introduction of the vector into a plant cell.

(49) As a selective marker for selecting a transformant, for example, a drug resistance gene can be used. Specific examples of such a drug resistance gene may include drug resistance genes that are resistant to hygromycin, bleomycin, kanamycin, gentamycin, chloramphenicol, etc. Thus, by selecting a plant body that grows in a medium containing the above-described antibiotics using such a selective marker, a transformed plant body can be easily selected.

(50) A method of constructing the expression vector is not particularly limited, either. The above-described promoter and the above-described nucleic acid encoding the functional peptide, and also, as necessary, the above-described other DNA segments may be introduced in a predetermined order into the appropriately selected vector used to produce an expression vector. For instance, a nucleic acid encoding the above described functional peptide may be connected with a promoter (as necessary, with a transcription terminator, etc.) to construct an expression cassette, and this expression cassette may be then introduced into a vector. In construction of such an expression cassette, for example, the cleavage sites of individual DNA segments are set to be protruding ends that are complementary to one another, and they are then reacted using a ligation enzyme, so that the order of the DNA segments can be determined. When the expression cassette comprises a terminator, the positional order may be a promoter, a nucleic acid encoding the above-described functional peptide, and a terminator from the upstream thereof. Moreover, the types of reagents used to construct the expression vector, namely, the types of restriction enzymes, ligation enzymes, etc. are not particularly limited, either, and commercially available products may be selected, as appropriate, and may be then used.

(51) The aforementioned expression vector is introduced into a target plant according to a common transformation method. The method of introducing an expression vector into a plant cell (transformation method) is not particularly limited, and a conventionally known suitable method that depends on a plant cell can be applied. Specifically, for example, a method using *Agrobacterium* or a method of directly introducing an expression vector into a plant cell can be used. Examples of the method of directly introducing an expression vector into a plant cell that can be used herein may include a microinjection method, an electroporation method, a polyethylene glycol method, a particle cancer method, a protoplast fusion method, and a calcium phosphate method.

(52) Moreover, if a method of directly introducing DNA into a plant cell is adopted, it is sufficient for the DNA to comprise transcription units necessary for the expression of a target gene, such as, for example, a promoter and a transcription terminator, and also, a nucleic acid encoding the above-described functional peptide. Thus, it is not essential for the DNA to function as a vector. Furthermore, it may be sufficient even if DNA comprises only a coding region of the above-described functional peptide, without having transcription units, if the DNA can be integrated into the transcription units of a host and can express the functional peptide as a target therein.

(53) Examples of the plant cell, into which the above-described expression vector or an expression cassette that does not comprise the expression vector but comprises a nucleic acid encoding the target functional peptide is to be introduced, may include the cells, callus, and suspension cultured cells of individual tissues in plant organs such as flowers, leaves, and roots. The expression vector that is used herein may be constructed, as appropriate, depending on the type of a plant body to be produced. Otherwise, a versatile expression vector may have previously been constructed, and it may be then introduced into a plant cell.

(54) Tumor tissues, shoots, hairy roots and the like obtained as a result of the transformation can be directly used in cell culture, tissue culture or organ culture. Alternatively, according to the

conventionally known plant tissue culture method, a suitable concentration of plant hormone (auxin, cytokinin, gibberellin, abscisic acid, ethylene, brassinolide, etc.) is administered to such tumor tissues and the like, so as to regenerate a plant body.

(55) As a regeneration method, a method comprising transferring callus-like transformed cells into a medium containing different type and different concentration of hormone, then culturing the cells therein to form an adventive embryo, and then obtaining a complete plant body, is adopted.

Examples of the medium used herein may include an LS medium and an MS medium.

(56) Moreover, the transformed plant according to the present invention also includes progeny plants obtained by introducing an expression vector comprising a nucleic acid encoding the aforementioned functional peptide into host cells to obtain transformed plant cells, then regenerating a transformed plant body from the transformed plant cells, then obtaining plant seeds from the obtained transformed plant body, and then obtaining the progeny plants from the plant seeds. In order to obtain plant seeds from a transformed plant body, for example, a transformed plant body is collected from a rooting medium and is then transplanted into a pot in which water-containing soil has been placed, and thereafter, the plant body is allowed to grow under a constant temperature and to form flowers, and finally seeds are formed. Otherwise, in order to produce a plant body from a seed, for example, when seeds formed in a transformed plant body have become mature, the seeds are isolated, are then seeded on soil containing water, and are then allowed to grow under a constant temperature and constant illumination, thereby producing plant bodies. The thus produced plant expresses the aforementioned functional peptide, and thus exhibits excellent resistance to the aforementioned plant pathogens.

EXAMPLES

(57) Hereinafter, the present invention will be described in more detail in the following examples. However, the following examples are not intended to limit the technical scope of the present invention.

(58) In the present example, functional peptides having antimicrobial activity against plant pathogenic filamentous fungi and partial peptides thereof were identified in the following order. 1. Collection of xylem saps of tomatoes infected/not-infected with wilt disease 2. Proteomic analysis 3. Identification of peptides in the xylem saps 4. Synthesis of functional peptides 5. Evaluation of antimicrobial activity against plant pathogenic filamentous fungi

(59) Hereafter, the above-described procedures 1. to 5. will be successively explained.

(60) <1. Collection of Xylem Saps of Tomatoes Infected/not Infected with Wilt Disease>

(61) A tomato variety Ponterosa was allowed to grow in soil (Kumiai Nippi Engei Baido). A tomato seedling with two first leaves developed was dugged up, and the roots thereof were then washed with tap water. The roots were immersed in a bud cell suspension of tomato wilt disease pathogens (*Fusarium oxysporum* CK3-1 strain) (1×10^6 cells/ml) for approximately 30 seconds, and were then transplanted in a vinyl pot containing Kumiai Nippi Engei Baido. Thereafter, the roots were allowed to grow in an environment control room (25° C., light: 14 hours, dark: 10 hours). As a control group, a non-inoculated seedling was allowed to grow in the same manner as described above. The inoculated seedling and the non-inoculated seedling were cut at a site immediately below the first leaf, 7, 9, 11, and 13 days after the transplantation. The pot was laid down, and the cross-section of the stem was placed in a 2.0-ml plastic tube. The stem was left at rest at room temperature for 12 hours, and a xylem sap leaked from each cross-section was recovered. The amount of the recovered liquid was measured, and thereafter, the liquid was then filtrated through a cellulose filter (Minisart RC15, Sartrium Stedim), and was then preserved at -80° C.

(62) The concentration of a protein in the xylem sap was measured using Bio-Rad DC Protein Assay Kit (Bio-Rad). A calibration curve was produced using bovine serum albumin (BSA).

(63) <2. Proteomic Analysis>

(64) In-Gel Trypsin Digestion of Xylem Sap Protein

(65) To a xylem sap sample (12 μ l, approx. 10 μ g) collected from each of the seedlings inoculated

with and not-inoculated with tomato wilt disease pathogens, 4 µl of 4×sample buffer (Wako Pure Chemical Industries, Ltd.) was added, and electrophoresis of approximately 2.5 cm was then performed according to SDS-PAGE. Thereafter, the gel was divided into 6 gel sections, and the sections were further minced to sections with 1 mm square. In-gel digestion was carried out with reference to the method of Rosefeld et al. (1992) (Rosenfeld, J., Capdevielle, J., Guillemot, J. C. and Ferrara, P. (1992). In-gel digestion of proteins for internal sequence analysis after one- or twodimensional gel electrophoresis. *Anal. Biochem.* 203: 173-179).

(66) First, the gel was washed with Milli-Q water (ultrapure water) and 50 mM ammonium carbonate/50% acetonitrile each once, and was then dried and solidified with acetonitrile. 50 mM tris(2-carboxyethyl)phosphine was added to the resulting gel, and the obtained mixture was then left at rest at 60° C. for 10 minutes. Thereafter, 100 mM 2-iodoacetamide was added to the resultant, and the thus obtained mixture was then left at rest at room temperature for 1 hour. Thereafter, the resulting gel was washed with Milli-Q water (ultrapure water) and 50 mM ammonium carbonate/50% acetonitrile each once, and was then dried and solidified with acetonitrile. After that, Lys-C/trypsin (0.01 mg/ml, manufactured by Promega) was added to the gel to swell it. To the swollen gel, 20 µl of 50 mM ammonium carbonate was added, and the obtained mixture was then reacted at 37° C. for 16 hours. Using 20% trifluoroacetic acid, the reaction solution was adjusted to pH=2, and was then filtrated through 0.1-µm Durapore Membrane PVDF (manufactured by Merck Millipore), followed by vacuum concentration, and the resultant was dissolved in 0.1% trifluoroacetic acid/2% acetonitrile.

(67) Nano LC-MS/MS Analysis

(68) For nano-LC-MS/MS analysis, a system formed by combining Q Exactive Hybrid Quadrupole-Orbitrap Mass Spectrometer (manufactured by Thermo Fisher Scientific) with Dionex U3000 gradient pump (manufactured by Thermo Fisher Scientific) was used. A trap column (L-column ODS, 300 µm I.D.×5 mm, particle diameter: 5 µm, Chemicals Evaluation and Research Institute) and an analytical column (NTCC-360, 100 µm I.D.×125 mm, particle diameter: 3 µm, Nikkyo Technos Co., Ltd.) were used. Also, a mobile phase A (0.5% acetic acid) and a mobile phase B (0.5% acetic acid-containing 80% acetonitrile) were used. The flow rate was set at 0.5 µl/min, and the gradient was carried out with an acetonitrile concentration from 5% B to 35% B for 100 minutes, from 35% B to 100% B for 1 minute, 100% B for 3 minutes, and finally 5% B for 10 minutes.

(69) <3. Identification of Peptide in Xylem Sap>

(70) Proteins detected by the nano LC-MS/MS analysis were analyzed using Proteome Discoverer 2.0.0.802 (manufactured by Thermo Fisher Scientific). As database, tomato sORF database (ITAG2.4_gene_models_sORF), tomato protein database (ITAG2.4_protein), tomato wilt disease pathogen protein database (*Fusarium_oxysporum_f._sp._lycopersici_4827_2_protein*) were used. As a result, the following peptides were discovered: 4 peptides (named as TsORF 6, 11, 19, and 29), the non-inoculated tomato-specific expression of which was confirmed; 4 peptides (named as TsORF 8, 15, 18, and 30), the inoculated tomato-specific expression of which was confirmed; and 6 peptides (named as TsORF 10, 16, 17, 21, 24, and 25), which were expressed in both non-inoculated and inoculated tomatoes.

(71) The above-described 14 types of peptides identified in the present example were not registered in the above-described tomato protein database (ITAG2.4_protein) and tomato wilt disease pathogen protein database (*Fusarium_oxysporum_f._sp._lycopersici_4827_2_protein*), but were registered only in the tomato sORF database (ITAG2.4_gene_models_sORF). This tomato sORF database store information regarding frames (sORFs: small open reading frames) encoding peptides consisting of 10 to 100 amino acid residues, which were found by applying the method disclosed in sORF finder: a program package to identify small open reading frames with high coding potential K Hanada, K Akiyama, T Sakurai, T Toyoda, K Shinozaki, *SH Shiu Bioinformatics* 26 (3), 399-400 to tomatoes and then discovering the peptides in an intergenic region that reportedly did not have a

common ORF.

(72) <4. Synthesis of Functional Peptide>

(73) In the present example, among the identified peptides, a peptide consisting of 55 amino acid residues, which was named as “TsORF6”

(MKVMIVVKTKVKKNKKVAKMMVKEDEKNKRVITMKEELMVVAEEEEAYRRV RGGMGW: SEQ ID NO: 9) was used as an analytical target. This peptide was divided into the following three regions, and these three regions were each chemically synthesized.

(74) TABLE-US-00001 TsORF6-1: (SEQ ID NO: 1) MKVMIVVKTKVKKNKKVAKMMVK
TsORF6-2: (SEQ ID NO: 10) KMMVKEDEKNKRVITMKEELMV TsORF6-3: (SEQ ID NO: 11) EELMVVAEEEEAYRRVRGGMGW

(75) Moreover, in the present example, based on the amino acid sequence of TsORF6-1, partial peptides each consisting of 8 amino acid residues (Table 1) and partial peptides each consisting of 5 amino acid residues (Table 2) were designed, and individual partial peptides were chemically synthesized.

(76) TABLE-US-00002 TABLE 1 Peptide name Amino acid sequence SEQ ID NO TsORF6-1-MKVMIVVK SEQ ID NO: 2 8aa-1 TsORF6-1- KVMIVVKT SEQ ID NO: 3 8aa-2 TsORF6-1-VMIVVKTK SEQ ID NO: 4 8aa-3 TsORF6-1- MIVVKTKV SEQ ID NO: 12 8aa-4 TsORF6-1-IVVKTKVK SEQ ID NO: 13 8aa-5 TsORF6-1- VVKTKVKN SEQ ID NO: 14 8aa-6 TsORF6-1-VKTKVKNK SEQ ID NO: 15 8aa-7 TsORF6-1- KTKVKNKK SEQ ID NO: 16 8aa-8 TsORF6-1-TKVKNKKV SEQ ID NO: 17 8aa-9 TsORF6-1- KVKNKKVA SEQ ID NO: 18 8aa-10 TsORF6-1-VKNKKVAK SEQ ID NO: 19 8aa-11 TsORF6-1- KNKKVAKM SEQ ID NO: 20 8aa-12 TsORF6-1- NKKVAKMM SEQ ID NO: 21 8aa-13 TsORF6-1- KKVAKMMV SEQ ID NO: 22 8aa-14 TsORF6-1- KVAKMMVK SEQ ID NO: 23 8aa-15

(77) TABLE-US-00003 TABLE 2 Peptide name Amino acid sequence SEQ ID NO TsORF6-1-5aa-1 MKVMI SEQ ID NO: 5 TsORF6-1-5aa-2 KVMIV SEQ ID NO: 6 TsORF6-1-5aa-3 VMIVV SEQ ID NO: 7 TsORF6-1-5aa-4 MIVVK SEQ ID NO: 24 TsORF6-1-5aa-5 IVVKT SEQ ID NO: 25 TsORF6-1-5aa-6 VVKTK SEQ ID NO: 26 TsORF6-1-5aa-7 VKTKV SEQ ID NO: 27 TsORF6-1-5aa-8 KTKVK SEQ ID NO: 28 TsORF6-1-5aa-9 TKVKN SEQ ID NO: 29 TsORF6-1-5aa-10 KVKNK SEQ ID NO: 30 TsORF6-1-5aa-11 VKNKK SEQ ID NO: 31 TsORF6-1-5aa-12 KNKKV SEQ ID NO: 32 TsORF6-1-5aa-13 NKKVA SEQ ID NO: 33 TsORF6-1-5aa-14 KKVAK SEQ ID NO: 34 TsORF6-1-5aa-15 KVAKM SEQ ID NO: 35 TsORF6-1-5aa-16 VAKMM SEQ ID NO: 8

<5. Evaluation of Antimicrobial Activity Against Plant Pathogenic Filamentous Fungi>

5-1. Plate Antimicrobial Test

(78) In the present example, antimicrobial activity against *Botrytis cinerea*, *Colletotrichum destructivum*, *Colletotrichum orbiculare*, *Fusarium oxysporum*, and *Pyricularia oryzae* was evaluated by observing germination of spores. In the present example, first, 100 µl of a spore suspension (approx. 10^{sup.4} cells) of each of these filamentous fungi was dispensed in a 96-well microplate.

(79) Subsequently, the peptides to be tested (TsORF6-1, TsORF6-2, TsORF6-3, and partial peptides of TsORF6-1 (Tables 1 and 2)) were each added into individual wells. It is to be noted that the concentration of the peptide was set at 1 µM, 10 µM or 100 µM. Thereafter, the peptides were cultured at 25° C. for 1 day or 2 days.

(80) After completion the culture, each well was observed under a microscope, and the antimicrobial activity of the tested peptides was evaluated. In the present example, antimicrobial activity was defined to be levels 0 to 3 in an ascending order. As a result of the microscopic observation, a case where hyphae extended to the entire well and no antimicrobial activity was found is defined as “level 0,” a case where hyphae extended to less than a half of the well and thus the peptide inhibited the growth of the hyphae is defined as “level 1,” a case where ungerminated spores were present and the peptide partially inhibited the germination is defined as “level 2,” and a case where all spores were ungerminated and the peptide completely inhibited the germination is

defined as “level 3.”

(81) As an example, the microscopic observation results of TsORF6-1 are shown in FIG. 1. Besides, the photographs shown in the upper case of FIG. 1 indicate non-treated controls. As shown in FIG. 1, when TsORF6-1 was used, all of *Botrytis cinerea*, *Colletotrichum destructivum*, *Colletotrichum orbiculare*, *Fusarium oxysporum* and *Pyricularia oryzae* were completely inhibited in terms of germination.

(82) 5-2. *Botrytis cinerea* Leaf Inoculation Test

(83) In the present example, a suspension comprising *Botrytis cinerea* and a peptide to be tested was dropped on a plant leaf, and lesions were then observed by visual inspection to evaluate the antimicrobial activity of the tested peptide.

(84) First, the pores of *Botrytis cinerea* were suspended in a potato dextrose broth (PDB) to result in a spore concentration of 5×10^5 spores/ml, so as to prepare a PDB suspension spore solution. Subsequently, a peptide to be tested (TsORF6-1) was added to the PDB suspension spore solution, so that the final concentration of the peptide to be tested became 0 μ M, 1 μ M, 10 μ M or 100 μ M.

(85) Thereafter, 10 μ l each of the PDB suspension spore solution containing the peptide to be tested was dropped on the front side and the back side of the leaf of tomato (variety name: “Momotaro Fight”). In this condition, incubation was carried out at 25° C. for 2 days. Thereafter, the front and back sides of the leaf were observed by visual inspection. A photograph of the front and back sides of the leaf is shown in FIG. 2. As shown in FIG. 2, even if the PDB suspension spore solution containing 100 μ l TsORF6 was dropped on the leaf, a lesion of *Botrytis cinerea* was not observed. From these results, it became clear that 100 μ l TsORF6 exhibited extremely excellent antimicrobial activity against *Botrytis cinerea*.

(86) 5-3. *Colletotrichum destructivum* Leaf Inoculation Test

(87) In the present example, a suspension comprising *Colletotrichum destructivum* and a peptide to be tested was dropped on a plant leaf, and lesions were then observed by visual inspection to evaluate the antimicrobial activity of the tested peptide.

(88) First, the pores of *Colletotrichum destructivum* were suspended in water to result in a spore concentration of 5×10^5 spores/ml, so as to prepare a water suspension of spores. Subsequently, a peptide to be tested (TsORF6-1) was added to the water suspension of spores, so that the final concentration of the peptide to be tested became 0 μ M, 1 μ M, 10 μ M or 100 μ M.

(89) Thereafter, 10 μ l each of the water suspension of spores containing the peptide to be tested was dropped on the front side and the back side of the leaf of Chinese cabbage (variety name: “Kigokoro 85”). In this condition, incubation was carried out at 25° C. for 4 days. Thereafter, the front and back sides of the leaf were observed by visual inspection. A photograph of the front and back sides of the leaf is shown in FIG. 3. As shown in FIG. 3, even if the water suspension of spores containing 10 μ l or 100 μ l TsORF6-1 was dropped on the leaf, a lesion of *Colletotrichum destructivum* was not observed. From these results, it became clear that 10 μ l or 100 μ l TsORF6-1 exhibited extremely excellent antimicrobial activity against *Colletotrichum destructivum*.

(90) 5-4. Antimicrobial Tests Against Other Plant Pathogens

(91) In the present example, antimicrobial activity against *Ralstonia solanacearum*, *Erwinia carotovora* subsp. *carotovora*, *Xanthomonas oryzae* pv. *oryzae*, and *Pseudomonas syringae* pv. *tomato* was evaluated. In the present example, first, these microorganisms were cultured, and 100 μ l of the culture solution (OD.sub.600=0.1) was dispensed in a 96-well microplate.

(92) Subsequently, TsORF6-1 was added to each well to result in a final concentration of 100 μ M. In addition, as a negative control, water was added instead of TsORF6-1. Thereafter, the obtained mixture was cultured at 28° C. a whole day and night, and the value of OD.sub.600 was then measured. Using the measured OD.sub.600 value, the proliferation percentage was calculated according to the following equation:

[Proliferation percentage (%)] = $(1 - \text{OD.sub.600}[\text{peptide added}]) / \text{OD.sub.600}[\text{water added}] \times 100$

5-5. Results

acid sequence: MKVMIVVK (SEQ ID NO: 2), (b) amino acid sequence: KVMIVVKT (SEQ ID NO: 3), (c) amino acid sequence: VMIVVKT (SEQ ID NO: 4), (d) amino acid sequence: MKVMI (SEQ ID NO: 5), (e) amino acid sequence: KVMIV (SEQ ID NO: 6), (f) amino acid sequence: VMIVV (SEQ ID NO: 7), and (g) amino acid sequence: VAKMM (SEQ ID NO: 8) have extremely excellent germination inhibitory activity against plant pathogenic filamentous fungi. (100) Besides, partial peptides, which do not exhibit the germination inhibitory effects on the aforementioned plant pathogenic filamentous fungi in Table 3, are likely to exhibit the germination inhibitory effects, if various conditions such as a concentration are studied.

Claims

1. A composition comprising a functional peptide consisting of 5 or more consecutive amino acids in the amino acid sequence: MKVMIVVKT VKNKKVAKMMVK (SEQ ID NO: 1), the composition exhibiting sterilizing action or antimicrobial action against plant pathogens.
 2. An agricultural chemical comprising a functional peptide consisting of 5 or more consecutive amino acids in the amino acid sequence: MKVMIVVKT VKNKKVAKMMVK (SEQ ID NO: 1), for sterilizing or antimicrobial use against plant pathogens.
 3. The agricultural chemical according to claim 2, wherein the plant pathogens are one or more plant pathogenic filamentous fungi selected from the group consisting of *Botrytis cinerea*, *Colletotrichum destructivum*, *Colletotrichum orbiculare*, *Fusarium oxysporum*, and *Pyricularia oryzae*.
 4. The agricultural chemical according to claim 2, wherein the plant pathogens are *Xanthomonas oryzae* pv. *oryzae* or *Pseudomonas syringae* pv. *tomato*.
-