



US 20250281623A1

(19) **United States**

(12) **Patent Application Publication**
Eastman et al.

(10) **Pub. No.: US 2025/0281623 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **HETEROBIFUNCTIONAL COMPOUNDS AND
THEIR USE IN TREATING DISEASE**

(71) Applicant: **Halda Therapeutics OpCo, Inc.**, New
Haven, CT (US)

(72) Inventors: **Kyle J. Eastman**, Killingworth, CT
(US); **Katherine J. Kayser-Bricker**,
Branford, CT (US); **Michael**
Barrington Martin, Manchester, CT
(US); **James John Mousseau**, Salem,
CT (US); **Matthew Alexander Perry**,
Uncasville, CT (US); **David E. Puleo**,
Fairfield, CT (US); **Nilesh K. Zaware**,
Hamden, CT (US); **Samuel W. Gerritz**,
Guilford, CT (US)

(21) Appl. No.: **19/073,111**

(22) Filed: **Mar. 7, 2025**

Related U.S. Application Data

(60) Provisional application No. 63/562,801, filed on Mar.
8, 2024, provisional application No. 63/761,647, filed
on Feb. 21, 2025.

Publication Classification

(51) **Int. Cl.**

A61K 47/55 (2017.01)

A61K 47/54 (2017.01)

A61P 35/00 (2006.01)

C07D 495/14 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 47/55** (2017.08); **A61K 47/545**
(2017.08); **A61P 35/00** (2018.01); **C07D**
495/14 (2013.01)

(57)

ABSTRACT

The invention provides heterobifunctional compounds,
pharmaceutical compositions, and their use in treating dis-
ease, such as cancer.

HETEROBIFUNCTIONAL COMPOUNDS AND THEIR USE IN TREATING DISEASE

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of and priority to U.S. Provisional Patent Application Ser. No. 63/761,647, filed Feb. 21, 2025, and U.S. Provisional Patent Application Ser. No. 63/562,801, filed Mar. 8, 2024; the contents of each of which are hereby incorporated by reference in their entirety.

FIELD OF THE INVENTION

[0002] The invention provides heterobifunctional compounds, pharmaceutical compositions, and their use in treating disease, such as cancer.

BACKGROUND

[0003] Cancer continues to be a significant health problem despite the substantial research efforts and scientific advances reported in the literature for treating this disease. Solid tumors, including prostate cancer, breast cancer, and lung cancer remain highly prevalent among the world population. The incidence of prostate cancer increases with age, and with increasing longevity of human subjects, there

continues to be a corresponding rise in the number of patients suffering from prostate cancer. Breast cancer is one of the most common cancers among women and is a leading cause of death for women between ages 50-55. Lung cancer is a leading cause of death among cancer patients, where over 85% of lung cancers are non-small cell lung cancer (NSCLC). Many lung cancers are attributed to tobacco smoking. Current treatment options for these cancers are not effective for all patients and/or can have substantial adverse side effects.

[0004] New therapies are needed to address this unmet need in cancer therapy. In particular, new therapies are needed that achieve an anti-cancer effect through a different mechanism than commonly available therapies. Exemplary mechanisms for common anti-cancer therapies include (a) alkylation of DNA, which limits ability of the cell to reproduce, (b) topoisomerase inhibition, in which the therapeutic agent inhibits the activity of one or more topoisomerases, thereby limiting separation of strands of DNA, and (c) mitotic inhibition, where the therapeutic agent reduces ability of the cell to divide. New therapies that achieve an anti-cancer effect through a different mechanism present an opportunity to treat cancers more effectively and/or to treat cancers that have become resistant to currently available medicines.

[0005] The present invention addresses the foregoing needs and provides other related advantages.


SUMMARY

[0006] The invention provides heterobifunctional compounds, pharmaceutical compositions, and their use in treating disease, such as cancer. In particular, one aspect of the invention provides a collection of heterobifunctional compounds, such as a compound represented by Formula I:



or a pharmaceutically acceptable salt thereof, where the variables are as defined in the detailed description. Further description of additional collections of heterobifunctional compounds are described in the detailed description. The compounds may be part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

[0007] Another aspect of the invention provides a collection of heterobifunctional compounds, such as a compound represented by Formula II:



or a pharmaceutically acceptable salt thereof, where the variables are as defined in the detailed description. Further description of additional collections of heterobifunctional compounds are described in the detailed description. The compounds may be part of a pharmaceutical composition comprising a pharmaceutically acceptable carrier.

[0008] Another aspect of the invention provides a method of treating cancer. The method comprises administering to a patient in need thereof a therapeutically effective amount of a compound described herein, such as a compound of Formula I or II, to treat the cancer.

[0009] Another aspect of the invention provides a method of causing death of a cancer cell. The method comprises contacting a cancer cell with an effective amount of a compound described herein, such as a compound of Formula I or II, to cause death of the cancer cell.

DETAILED DESCRIPTION

[0010] The invention provides heterobifunctional compounds, pharmaceutical compositions, and their use in treating disease, such as cancer. The practice of the present invention employs, unless otherwise indicated, conventional techniques of organic chemistry, pharmacology, molecular biology (including recombinant techniques), cell biology, biochemistry, and immunology. Such techniques are explained in the literature, such as in "Comprehensive

Organic Synthesis" (B. M. Trost & I. Fleming, eds., 1991-1992); "Handbook of experimental immunology" (D. M. Weir & C. C. Blackwell, eds.); "Current protocols in molecular biology" (F. M. Ausubel et al., eds., 1987, and periodic updates); and "Current protocols in immunology" (J. E. Coligan et al., eds., 1991), each of which is herein incorporated by reference in its entirety.

[0011] Various aspects of the invention are set forth below in sections; however, aspects of the invention described in one particular section are not to be limited to any particular section. Further, when a variable is not accompanied by a definition, the previous definition of the variable controls.


Definitions

[0012] Compounds of the present invention include those described generally herein, and are further illustrated by the classes, subclasses, and species disclosed herein. As used herein, the following definitions shall apply unless otherwise indicated. These definitions apply regardless of whether a term is used by itself or in combination with other terms, unless otherwise indicated. Hence, the definition of "alkyl" applies to "alkyl" as well as the "alkyl" portions of "—O-alkyl" etc. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 75th Ed. Additionally, general principles of organic chemistry are described in "Organic Chemistry", Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry", 5th Ed., Ed.: Smith, M. B. and March, J., John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.


[0013] The term "aliphatic" or "aliphatic group", as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated or that contains one or more units of unsaturation, or a monocyclic hydrocarbon or bicyclic hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic (also referred to herein as "cycloaliphatic"), that has a single point of attachment to the rest of the molecule. Unless otherwise specified, aliphatic groups contain 1-6 aliphatic carbon atoms. In some embodiments, aliphatic groups contain 1-5 aliphatic carbon atoms. In other embodiments, aliphatic groups contain 1-4 aliphatic carbon atoms. In still other embodiments, aliphatic groups contain 1-3 aliphatic carbon atoms, and in yet other embodiments, aliphatic groups contain 1-2 aliphatic carbon atoms. In some embodiments, "cycloaliphatic" refers to a monocyclic C₃-C₆ hydrocarbon that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, that has a single point of attachment to the rest of the molecule. Suitable aliphatic groups include, but are not limited to, linear or branched, substituted or unsubstituted alkyl, alkenyl, alkynyl groups and hybrids thereof such as (cycloalkyl)alkyl, (cycloalkenyl)alkyl or (cycloalkyl)alkenyl.


[0014] As used herein, the term "bicyclic ring" or "bicyclic ring system" refers to any bicyclic ring system, i.e., carbocyclic or heterocyclic, saturated or having one or more units of unsaturation, having one or more atoms in common between the two rings of the ring system. Thus, the term includes any permissible ring fusion, such as ortho-fused or spirocyclic. As used herein, the term "heterobicyclic" is a subset of "bicyclic" that requires that one or more heteroa-

toms are present in one or both rings of the bicycle. Such heteroatoms may be present at ring junctions and are optionally substituted, and may be selected from nitrogen (including N-oxides), oxygen, sulfur (including oxidized forms such as sulfones and sulfonates), phosphorus (including oxidized forms such as phosphates), boron, etc. In some embodiments, a bicyclic group has 7-12 ring members and 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. As used herein, the term "bridged bicyclic" refers to any bicyclic ring system, i.e., carbocyclic or heterocyclic, saturated or partially unsaturated, having at least one bridge. As defined by IUPAC, a "bridge" is an unbranched chain of atoms or an atom or a valence bond connecting two bridgeheads, where a "bridgehead" is any skeletal atom of the ring system which is bonded to three or more skeletal atoms (excluding hydrogen). In some embodiments, a bridged bicyclic group has 7-12 ring members and 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur. Such bridged bicyclic groups are well known in the art and include those groups set forth below where each group is attached to the rest of the molecule at any substitutable carbon or nitrogen atom. Unless otherwise specified, a bridged bicyclic group is optionally substituted with one or more substituents as set forth for aliphatic groups. Additionally or alternatively, any substitutable nitrogen of a bridged bicyclic group is optionally substituted. Exemplary bridged bicyclic rings include:



[0015] Exemplary bridged bicyclics include:





[0016] The term “lower alkyl” refers to a C_{1-4} straight or branched alkyl group. Exemplary lower alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, and tert-butyl.

[0017] The term “lower haloalkyl” refers to a C_{1-4} straight or branched alkyl group that is substituted with one or more halogen atoms.

[0018] The term “heteroatom” means one or more of oxygen, sulfur, nitrogen, phosphorus, or silicon (including, any oxidized form of nitrogen, sulfur, phosphorus, or silicon; the quaternized form of any basic nitrogen or; a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl) or NR^+ (as in N-substituted pyrrolidinyl)).

[0019] The term “unsaturated,” as used herein, means that a moiety has one or more units of unsaturation.

[0020] As used herein, the term “bivalent C_{1-8} (or C_{1-6}) saturated or unsaturated, straight or branched, hydrocarbon chain”, refers to bivalent alkylene, alkenylene, and alkyne chains that are straight or branched as defined herein.

[0021] The term “alkylene” refers to a bivalent alkyl group. An “alkylene chain” is a polymethylene group, i.e., $-(CH_2)_n-$, wherein n is a positive integer, preferably from 1 to 6, from 1 to 4, from 1 to 3, from 1 to 2, or from 2 to 3. A substituted alkylene chain is a polymethylene group in which one or more methylene hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0022] The term “ $-(C_0 \text{ alkylene})-$ ” refers to a bond. Accordingly, the term “ $-(C_{0-3} \text{ alkylene})-$ ” encompasses a bond (i.e., C_0) and a $-(C_{1-3} \text{ alkylene})-$ group.

[0023] The term “alkenylene” refers to a bivalent alkenyl group. A substituted alkenylene chain is a polymethylene group containing at least one double bond in which one or more hydrogen atoms are replaced with a substituent. Suitable substituents include those described below for a substituted aliphatic group.

[0024] The term “halogen” means F, Cl, Br, or I.

[0025] The term “aryl” used alone or as part of a larger moiety as in “aralkyl,” “aralkoxy,” or “aryloxyalkyl,” refers to monocyclic or bicyclic ring systems having a total of five to fourteen ring members, wherein at least one ring in the system is aromatic and wherein each ring in the system contains 3 to 7 ring members. The term “aryl” may be used interchangeably with the term “aryl ring.” In certain embodiments of the present invention, “aryl” refers to an aromatic ring system which includes, but not limited to, phenyl, biphenyl, naphthyl, anthracyl and the like, which may bear one or more substituents. Also included within the scope of the term “aryl,” as it is used herein, is a group in which an aromatic ring is fused to one or more non-aromatic rings,

such as indanyl, phthalimidyl, naphthimidyl, phenanthridinyl, or tetrahydronaphthyl, and the like. The term “haloaryl” refers to an aryl group that is substituted with at least one halogen. Exemplary haloaryl groups include chlorophenyl (e.g., 3-chlorophenyl, 4-chlorophenyl), fluorophenyl, and the like. The term “phenylene” refers to a bivalent phenyl group.

[0026] The terms “heteroaryl” and “heteroar-,” used alone or as part of a larger moiety, e.g., “heteroaralkyl,” or “heteroaralkoxy,” refer to groups having 5 to 10 ring atoms, preferably 5, 6, or 9 ring atoms; having 6, 10, or 14 π electrons shared in a cyclic array; and having, in addition to carbon atoms, from one to five heteroatoms. The term “heteroatom” refers to nitrogen, oxygen, or sulfur, and includes any oxidized form of nitrogen or sulfur, and any quaternized form of a basic nitrogen. Heteroaryl groups include, without limitation, thienyl, furanyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, oxadiazolyl, thiazolyl, isothiazolyl, thiadiazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, indoliziny, purinyl, naphthyridinyl, and pteridinyl. The terms “heteroaryl” and “heteroar-,” as used herein, also include groups in which a heteroaromatic ring is fused to one or more aryl, cycloaliphatic, or heterocyclic rings, where unless otherwise specified, the radical or point of attachment is on the heteroaromatic ring or on one of the rings to which the heteroaromatic ring is fused. Nonlimiting examples include indolyl, isoindolyl, benzothienyl, benzofuranyl, dibenzofuranyl, indazolyl, benzimidazolyl, benzthiazolyl, quinolyl, isoquinolyl, cinnolyl, phthalazinyl, quinazolinyl, quinoxalinyl, 4H-quinoliziny, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, tetrahydroquinolyl, and tetrahydroisoquinolyl. A heteroaryl group may be mono- or bicyclic. The term “heteroaryl” may be used interchangeably with the terms “heteroaryl ring,” “heteroaryl group,” or “heteroaromatic,” any of which terms include rings that are optionally substituted. The term “heteroaralkyl” refers to an alkyl group substituted by a heteroaryl, wherein the alkyl and heteroaryl portions independently are optionally substituted. The term “haloheteroaryl” refers to a heteroaryl group that is substituted with at least one halogen. Exemplary haloheteroaryl groups include chloropyridine, fluoropyridine, chloropyrazole, fluoropyrazole, and the like. The term “heteroarylene” refers to a bivalent heteroaryl group. Similarly, the terms “pyrazolylene,” “imidazolylene,” and “pyrrolylene,” respectively refer to bivalent pyrazolyl, imidazolyl, and pyrrolyl groups. Similarly, the terms “pyridinylene” and “pyrimidinylene,” respectively refer to bivalent pyridinyl and pyrimidinyl groups.

[0027] As used herein, the terms “heterocycle,” “heterocyclic,” “heterocyclic radical,” and “heterocyclic ring” are used interchangeably and refer to a stable 5- to 7-membered monocyclic or 7-10-membered bicyclic heterocyclic moiety that is either saturated or partially unsaturated, and having, in addition to carbon atoms, one or more, preferably one to four, heteroatoms, as defined above. When used in reference to a ring atom of a heterocycle, the term “nitrogen” includes a substituted nitrogen. As an example, in a saturated or partially unsaturated ring having 0-3 heteroatoms selected from oxygen, sulfur or nitrogen, the nitrogen may be N (as in 3,4-dihydro-2H-pyrrolyl), NH (as in pyrrolidinyl), or ^+NR (as in N-substituted pyrrolidinyl).

[0028] A heterocyclic ring can be attached to its pendant group at any heteroatom or carbon atom that results in a

stable structure and any of the ring atoms can be optionally substituted. Examples of such saturated or partially unsaturated heterocyclic radicals include, without limitation, tetrahydrofuran-yl, tetrahydrothiophen-yl pyrrolidin-yl, piperidin-yl, pyrrolin-yl, tetrahydroquinolin-yl, tetrahydroisoquinolin-yl, decahydroquinolin-yl, oxazolidin-yl, piperazin-yl, dioxan-yl, dioxolan-yl, diazepin-yl, oxazepin-yl, thiazepin-yl, morpholin-yl, 2-oxa-6-azaspiro[3.3]heptane, and quinuclidin-yl. The terms “heterocycle,” “heterocycl-yl,” “heterocycl-yl ring,” “heterocyclic group,” “heterocyclic moiety,” and “heterocyclic radical,” are used interchangeably herein, and also include groups in which a heterocycl-yl ring is fused to one or more aryl, heteroaryl, or cycloaliphatic rings, such as indolin-yl, 3H-indolin-yl, chroman-yl, phenanthridin-yl, or tetrahydroquinolin-yl. A heterocycl-yl group may be mono- or bicyclic. The term “heterocycl-ylalk-yl” refers to an alkyl group substituted by a heterocycl-yl, wherein the alkyl and heterocycl-yl portions independently are optionally substituted. The term “heterocycl-ylene” refers to a bivalent heterocycl-yl group.

[0029] As used herein, the term “heterocycloalk-yl” refers to a saturated heterocycl-yl. The term “heterocycloalk-yl” refers to a bivalent heterocycloalk-yl group.

[0030] As used herein, the term “partially unsaturated” refers to a ring moiety that includes at least one double or triple bond. The term “partially unsaturated” is intended to encompass rings having multiple sites of unsaturation, but is not intended to include aryl or heteroaryl moieties, as herein defined.

[0031] As described herein, compounds of the invention may contain “optionally substituted” moieties. In general, the term “substituted,” whether preceded by the term “optionally” or not, means that one or more hydrogens of the designated moiety are replaced with a suitable substituent. Unless otherwise indicated, an “optionally substituted” group may have a suitable substituent at each substitutable position of the group, and when more than one position in any given structure may be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at every position. Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable,” as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, in certain embodiments, their recovery, purification, and use for one or more of the purposes disclosed herein.

[0032] Each optional substituent on a substitutable carbon is a monovalent substituent independently selected from halogen; $-(CH_2)_{0-4}R^\circ$; $-(CH_2)_{0-4}OR^\circ$; $-O(CH_2)_{0-4}R^\circ$; $-O(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}CH(OR^\circ)_2$; $-(CH_2)_{0-4}SR^\circ$; $-(CH_2)_{0-4}Ph$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}Ph$ which may be substituted with R° ; $-CH=CHPh$, which may be substituted with R° ; $-(CH_2)_{0-4}O(CH_2)_{0-1}pyridyl$ which may be substituted with R° ; $-NO_2$; $-CN$; $-N_3$; $-(CH_2)_{0-4}N(R^\circ)_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)R^\circ$; $-N(R^\circ)C(S)R^\circ$; $-(CH_2)_{0-4}N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)C(S)NR^\circ_2$; $-(CH_2)_{0-4}N(R^\circ)C(O)OR^\circ$; $-N(R^\circ)N(R^\circ)C(O)R^\circ$; $-N(R^\circ)N(R^\circ)C(O)NR^\circ_2$; $-N(R^\circ)N(R^\circ)C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)R^\circ$; $-C(S)R^\circ$; $-(CH_2)_{0-4}C(O)OR^\circ$; $-(CH_2)_{0-4}C(O)SR^\circ$; $-(CH_2)_{0-4}C(O)OSiR^\circ_3$; $-(CH_2)_{0-4}OC(O)R^\circ$; $-OC(O)(CH_2)_{0-4}SR^\circ$; $-SC(S)SR^\circ$; $-(CH_2)_{0-4}SC(O)R^\circ$; $-(CH_2)_{0-4}C(O)NR^\circ_2$; $-C(S)NR^\circ_2$;

$-C(S)SR^\circ$; $-SC(S)SR^\circ$; $-(CH_2)_{0-4}OC(O)NR^\circ_2$; $-C(O)N(OR^\circ)R^\circ$; $-C(O)C(O)R^\circ$; $-C(O)CH_2C(O)R^\circ$; $-C(NOR^\circ)R^\circ$; $-(CH_2)_{0-4}SSR^\circ$; $-(CH_2)_{0-4}S(O)_2R^\circ$; $-(CH_2)_{0-4}S(O)_2OR^\circ$; $-(CH_2)_{0-4}OS(O)_2R^\circ$; $-S(O)_2NR^\circ_2$; $-S(O)(NR^\circ)R^\circ$; $-S(O)_2N=C(NR^\circ)_2$; $-(CH_2)_{0-4}S(O)R^\circ$; $-N(R^\circ)S(O)_2NR^\circ_2$; $-N(R^\circ)S(O)_2R^\circ$; $-N(OR^\circ)R^\circ$; $-C(NH)NR^\circ_2$; $-P(O)_2R^\circ$; $-P(O)R^\circ_2$; $-OP(O)R^\circ_2$; $-OP(O)(OR^\circ)_2$; SiR°_3 ; $-(C_{1-4}$ straight or branched alkylene) $O-N(R^\circ)_2$; or $-(C_{1-4}$ straight or branched alkylene) $C(O)O-N(R^\circ)_2$.

[0033] Each R° is independently hydrogen, C_{1-6} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, $-CH_2$ -(5-6 membered heteroaryl ring), or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, notwithstanding the definition above, two independent occurrences of R° , taken together with their intervening atom(s), form a 3-12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, which may be substituted by a divalent substituent on a saturated carbon atom of R° selected from $=O$ and $=S$; or each R° is optionally substituted with a monovalent substituent independently selected from halogen, $-(CH_2)_{0-2}R^*$, $-(haloR^*)$, $-(CH_2)_{0-2}OH$, $-(CH_2)_{0-2}OR^*$, $-(CH_2)_{0-2}CH(OR^*)_2$, $-O(haloR^*)$, $-CN$, $-N_3$, $-(CH_2)_{0-2}C(O)R^*$, $-(CH_2)_{0-2}C(O)OH$, $-(CH_2)_{0-2}C(O)OR^*$, $-(CH_2)_{0-2}SR^*$, $-(CH_2)_{0-2}SH$, $-(CH_2)_{0-2}NH_2$, $-(CH_2)_{0-2}NHR^*$, $-(CH_2)_{0-2}NR^*_2$, $-NO_2$, $-SiR^*_3$, $-OSiR^*_3$, $-C(O)SR^*$, $-(C_{1-4}$ straight or branched alkylene) $C(O)OR^*$, or $-SSR^*$.

[0034] Each R^* is independently selected from C_{1-4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein each R^* is unsubstituted or where preceded by halo is substituted only with one or more halogens; or wherein an optional substituent on a saturated carbon is a divalent substituent independently selected from $=O$, $=S$, $=NNR^*_2$, $=NNHC(O)R^*$, $=NNHC(O)OR^*$, $=NNHS(O)_2R^*$, $=NR^*_2$, $=NOR^*$, $-O(C(R^*_2))_{2-3}O-$, or $-S(C(R^*_2))_{2-3}S-$, or a divalent substituent bound to vicinal substitutable carbons of an “optionally substituted” group is $-O(CR^*_2)_{2-3}O-$, wherein each independent occurrence of R^* is selected from hydrogen, C_{1-6} aliphatic or an unsubstituted 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0035] When R^* is C_{1-6} aliphatic, R^* is optionally substituted with halogen, $-R^*$, $-(haloR^*)$, $-OH$, $-OR^*$, $-O(haloR^*)$, $-CN$, $-C(O)OH$, $-C(O)OR^*$, $-NH_2$, $-NHR^*$, $-NR^*_2$, or $-NO_2$, wherein each R^* is independently selected from C_{1-4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein each R^* is unsubstituted or where preceded by halo is substituted only with one or more halogens.

[0036] An optional substituent on a substitutable nitrogen is independently $-R^\dagger$, $-NR^\dagger_2$, $-C(O)R^\dagger$, $-C(O)OR^\dagger$, $-C(O)C(O)R^\dagger$, $-C(O)CH_2C(O)R^\dagger$, $-S(O)_2R^\dagger$, $-S(O)_2NR^\dagger_2$, $-C(S)NR^\dagger_2$, $-C(NH)NR^\dagger_2$, or $-N(R^\dagger)S(O)_2R^\dagger$; wherein each RT is independently hydrogen, C_{1-6} aliphatic, unsubstituted $-OPh$, or an unsubstituted 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4

heteroatoms independently selected from nitrogen, oxygen, or sulfur, or, two independent occurrences of $R^{\dagger}$, taken together with their intervening atom(s) form an unsubstituted 3-12-membered saturated, partially unsaturated, or aryl mono- or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein when $R^{\dagger}$ is C_{1-6} aliphatic, RT is optionally substituted with halogen, $-R^{\dagger}$, $-(haloR^{\dagger})$, $-OH$, $-OR^{\dagger}$, $-O(haloR^{\dagger})$, $-CN$, $-C(O)OH$, $-C(O)OR^{\dagger}$, $-NH_2$, $-NHR^{\dagger}$, $-NR^{\dagger}_2$, or $-NO_2$, wherein each $R^{\dagger}$ is independently selected from C_{1-4} aliphatic, $-CH_2Ph$, $-O(CH_2)_{0-1}Ph$, or a 5-6-membered saturated, partially unsaturated, or aryl ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur, and wherein each $R^{\dagger}$ is unsubstituted or where preceded by halo is substituted only with one or more halogens.

[0037] As used herein, the term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like, and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66, 1-19, incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like.

[0038] Further, acids which are generally considered suitable for the formation of pharmaceutically useful salts from basic pharmaceutical compounds are discussed, for example, by P. Stahl et al., Camille G. (eds.) *Handbook of Pharmaceutical Salts. Properties, Selection and Use*. (2002) Zurich: Wiley-VCH; S. Berge et al., *Journal of Pharmaceutical Sciences* (1977) 66(1) 1-19; P. Gould, *International J. of Pharmaceutics* (1986) 33 201-217; Anderson et al., *The Practice of Medicinal Chemistry* (1996), Academic Press, New York; and in *The Orange Book* (Food & Drug Administration, Washington, D.C. on their website). These disclosures are incorporated herein by reference.

[0039] Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4}alkyl)_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium,

and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, loweralkyl sulfonate and aryl sulfonate.

[0040] Unless otherwise stated, structures depicted herein are also meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, Z and E double bond isomers, and Z and E conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. In certain embodiments, the compound is in the form of a single stereoisomer. In certain embodiments, a stereogenic center in a single stereoisomer compound has the R configuration. In certain embodiments, a stereogenic center in a single stereoisomer compound has the S configuration. Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structures including the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools, as probes in biological assays, or as therapeutic agents in accordance with the present invention. In certain embodiments, the compound depicted by a chemical structure herein has the natural distribution of isotopes (i.e., it is not isotopically enriched).

[0041] Diastereomeric mixtures can be separated into their individual diastereomers on the basis of their physical chemical differences by methods known to those skilled in the art, such as, for example, by chromatography and/or fractional crystallization. Enantiomers can be separated by converting the enantiomeric mixture into a diastereomeric mixture by reaction with an appropriate optically active compound (e.g., chiral auxiliary such as a chiral alcohol or Mosher's acid chloride), separating the diastereomers and converting (e.g., hydrolyzing) the individual diastereomers to the corresponding pure enantiomers. Alternatively, a particular enantiomer of a compound of the present invention may be prepared by asymmetric synthesis. Still further, where the molecule contains a basic functional group (such as amino) or an acidic functional group (such as carboxylic acid) diastereomeric salts are formed with an appropriate optically active acid or base, followed by resolution of the diastereomers thus formed by fractional crystallization or chromatographic means known in the art, and subsequent recovery of the pure enantiomers.

[0042] Individual stereoisomers of the compounds of the invention may, for example, be substantially free of other isomers, or may be admixed, for example, as racemates or with all other, or other selected, stereoisomers. Chiral center(s) in a compound of the present invention can have the S or R configuration as defined by the IUPAC 1974 Recommendations. Further, to the extent a compound described herein may exist as an atropisomer (e.g., substituted biaryls), all forms of such atropisomer are considered part of this invention.

[0043] Chemical names, common names, and chemical structures may be used interchangeably to describe the same structure. If a chemical compound is referred to using both a chemical structure and a chemical name, and an ambiguity exists between the structure and the name, the structure predominates. It should also be noted that any carbon as well as heteroatom with unsatisfied valences in the text, schemes, examples and tables herein is assumed to have the sufficient number of hydrogen atom(s) to satisfy the valences.

[0044] The terms “a” and “an” as used herein mean “one or more” and include the plural unless the context is inappropriate.

[0045] The term “alkyl” refers to a saturated straight or branched hydrocarbon, such as a straight or branched group of 1-12, 1-10, or 1-6 carbon atoms, referred to herein as C₁-C₁₂ alkyl, C₁-C₁₀ alkyl, and C₁-C₆ alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, etc.

[0046] The term “cycloalkyl” refers to a monovalent saturated cyclic, bicyclic, or bridged cyclic (e.g., adamantyl) hydrocarbon group of 3-12, 3-8, 4-8, or 4-6 carbons, referred to herein, e.g., as “C₃-C₆ cycloalkyl,” derived from a cycloalkane. Exemplary cycloalkyl groups include cyclohexyl, cyclopentyl, cyclobutyl, and cyclopropyl. The term “cycloalkylene” refers to a bivalent cycloalkyl group.

[0047] The term “haloalkyl” refers to an alkyl group that is substituted with at least one halogen. Exemplary haloalkyl groups include —CH₂F, —CHF₂, —CF₃, —CH₂CF₃, —CF₂CF₃, and the like. The term “chloroalkyl” refers to an alkyl group that is substituted with at least one chloro. The term “bromoalkyl” refers to an alkyl group that is substituted with at least one bromo. The term “haloalkylene” refers to a bivalent haloalkyl group.

[0048] The term “hydroxyalkyl” refers to an alkyl group that is substituted with at least one hydroxyl. Exemplary hydroxyalkyl groups include —CH₂CH₂OH, —C(H)(OH)CH₃, —CH₂C(H)(OH)CH₂CH₂OH, and the like.

[0049] The term “heteroalkyl” refers to an alkyl group in which one or more carbon atoms has been replaced by a heteroatom (e.g., N, O, or S). Exemplary heteroalkyl groups include —OCH₃, —CH₂OCH₃, —CH₂CH₂N(CH₃)₂, and —CH₂CH₂OH. The heteroalkyl group may contain, for example, from 2-4, 2-6, or 2-8 atoms selected from the group consisting of carbon and a heteroatom (e.g., N, O, or S). The phrase 3-8 membered heteroalkyl refers to a heteroalkyl group having from 3 to 8 atoms selected from the group consisting of carbon and a heteroatom. The term “heteroalkylene” refers to a bivalent heteroalkyl group.


[0050] The terms “alkenyl” and “alkynyl” are art-recognized and refer to unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double or triple bond respectively. The term “haloalkenyl” refers to an alkenyl group that is substituted with at least one halogen. The term “fluoroalkenyl” refers to an alkenyl group that is substituted with at least one fluoro. The term “nitroalkenyl” refers to an alkenyl group that is substituted with at least one nitro.

[0051] The term “carbocyclylene” refers to a bivalent cycloaliphatic group.

[0052] The terms “alkoxyl” or “alkoxy” are art-recognized and refer to an alkyl group, as defined above, having an oxygen radical attached thereto. Representative alkoxyl groups include methoxy, ethoxy, propoxy, tert-butoxy and the like. The term “haloalkoxyl” refers to an alkoxyl group that is substituted with at least one halogen. Exemplary haloalkoxyl groups include —OCH₂F, —OCHF₂, —OCF₃, —OCH₂CF₃, —OCF₂CF₃, and the like.


[0053] The term “oxo” is art-recognized and refers to a “=O” substituent. For example, a cyclopentane substituted with an oxo group is cyclopentanone.

[0054] The term “amino” is art-recognized and refers to both unsubstituted and substituted amines, e.g., a moiety that may be represented by the general formulas:



wherein R⁵⁰, R⁵¹, R⁵² and R⁵³ each independently represent a hydrogen, an alkyl, an alkenyl, —(CH₂)_m—R⁶¹, or R⁵⁰ and R⁵¹, taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure; R⁶¹ represents an aryl, a 3-7 membered cycloalkyl, a 4-7 membered cycloalkenyl, 5-10 membered heteroaryl, or 3-10 membered heterocyclyl; and m is zero or an integer in the range of 1 to 8.

[0055] The term “amido” is art-recognized and refers to both unsubstituted and substituted amides, e.g., a moiety that may be represented by the general formulas:



wherein R⁵⁰ and R⁵¹ each independently represent a hydrogen, an alkyl, an alkenyl, —(CH₂)_m—R⁶¹, or R⁵⁰ and R⁵¹, taken together with the N atom to which they are attached complete a heterocycle having from 4 to 8 atoms in the ring structure; R⁶¹ represents an aryl, a 3-7 membered cycloalkyl, a 4-7 membered cycloalkenyl, 5-10 membered heteroaryl, or 3-10 membered heterocyclyl; and m is zero or an integer in the range of 1 to 8; and R⁵² is an alkyl, an alkenyl, or —(CH₂)_m—R⁶¹.

[0056] The symbol “” indicates a point of attachment.

[0057] When any substituent or variable occurs more than one time in any constituent or the compound of the invention, its definition on each occurrence is independent of its definition at every other occurrence, unless otherwise indicated.

[0058] One or more compounds of the invention may exist in unsolvated as well as solvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. “Solvate” means a physical association of a compound of this invention with one or more

solvent molecules. This physical association involves varying degrees of ionic and covalent bonding, including hydrogen bonding. In certain instances the solvate will be capable of isolation, for example when one or more solvent molecules are incorporated in the crystal lattice of the crystalline solid. "Solvate" encompasses both solution-phase and isolatable solvates. Non-limiting examples of suitable solvates include ethanولات, methanولات, and the like. "Hydrate" is a solvate wherein the solvent molecule is H₂O.

[0059] As used herein, the terms "subject" and "patient" are used interchangeably and refer to organisms to be treated by the methods of the present invention. Such organisms preferably include, but are not limited to, mammals (e.g., murines, simians, equines, bovines, porcines, canines, felines, and the like), and most preferably includes humans.

[0060] The term "IC₅₀" is art-recognized and refers to the concentration of a compound that is required to achieve 50% inhibition of the target.

[0061] As used herein, the term "effective amount" refers to the amount of a compound sufficient to effect beneficial or desired results (e.g., a therapeutic, ameliorative, inhibitory or preventative result). An effective amount can be administered in one or more administrations, applications or dosages and is not intended to be limited to a particular formulation or administration route. As used herein, the term "treating" includes any effect, e.g., lessening, reducing, modulating, ameliorating or eliminating, that results in the improvement of the condition, disease, disorder, and the like, or ameliorating a symptom thereof.

[0062] As used herein, the term "pharmaceutical composition" refers to the combination of an active agent with a carrier, inert or active, making the composition especially suitable for diagnostic or therapeutic use in vivo or ex vivo.

[0063] As used herein, the term "pharmaceutically acceptable carrier" refers to any of the standard pharmaceutical carriers, such as a phosphate buffered saline solution, water, emulsions (e.g., such as an oil/water or water/oil emulsions), and various types of wetting agents. The compositions also can include stabilizers and preservatives. For examples of carriers, stabilizers and adjuvants, see e.g., Martin, Remington's Pharmaceutical Sciences, 15th Ed., Mack Publ. Co., Easton, PA [1975].

[0064] For therapeutic use, salts of the compounds of the present invention are contemplated as being pharmaceutically acceptable. However, salts of acids and bases that are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound.

[0065] In addition, when a compound of the invention contains both a basic moiety (such as, but not limited to, a pyridine or imidazole) and an acidic moiety (such as, but not limited to, a carboxylic acid) zwitterions ("inner salts") may be formed. Such acidic and basic salts used within the scope of the invention are pharmaceutically acceptable (i.e., non-toxic, physiologically acceptable) salts. Such salts of the compounds of the invention may be formed, for example, by reacting a compound of the invention with an amount of acid or base, such as an equivalent amount, in a medium such as one in which the salt precipitates or in an aqueous medium followed by lyophilization.


[0066] Throughout the description, where compositions are described as having, including, or comprising specific components, or where processes and methods are described as having, including, or comprising specific steps, it is

contemplated that, additionally, there are compositions of the present invention that consist essentially of, or consist of, the recited components, and that there are processes and methods according to the present invention that consist essentially of, or consist of, the recited processing steps.

[0067] As a general matter, compositions specifying a percentage are by weight unless otherwise specified.

I. Heterobifunctional Compounds

[0068] The invention provides heterobifunctional compounds. The compounds are generally represented by the following formula:




or a pharmaceutically acceptable salt thereof, wherein EPL is a moiety that binds to an effector protein selected from BRD4, CDK1, CDK2, CDK9, or GSPT1; L is a linker; and TPL is a moiety that binds to estrogen receptor.

[0069] The compounds may be used in the pharmaceutical compositions and therapeutic methods described herein. Exemplary compounds are described in the following sections, along with exemplary procedures for making the compounds.

Part A: Compounds of Formula I

[0070] One aspect of the invention provides a compound represented by Formula I:



(I)

[0071] or a pharmaceutically acceptable salt thereof, wherein:

[0072] EPL is a moiety that binds to an effector protein selected from BRD4, CDK1, CDK2, or CDK9;

[0073] L is a linker; and

[0074] TPL is a moiety that binds to an estrogen receptor.

[0075] The definitions of variables in Formula I above encompass multiple chemical groups. The application contemplates embodiments where, for example, i) the definition of a variable is a single chemical group selected from those chemical groups set forth above, ii) the definition of a variable is a collection of two or more of the chemical groups selected from those set forth above, and iii) the compound is defined by a combination of variables in which the variables are defined by (i) or (ii).

[0076] In certain embodiments, the compound is a compound of Formula I.


[0077] The compound may be further characterized according to, for example, the identity of L and/or TPL. Exemplary further embodiments for L and TPL are provided in Parts C and D below.

[0078] As generally defined above, EPL is a moiety that binds to an effector protein selected from BRD4, CDK1, CDK2, and CDK9. In certain embodiments, the EPL is a moiety that binds to BRD4. In certain embodiments, the


EPL is a moiety that binds to CDK1. In certain embodiments, the EPL is a moiety that binds to CDK2. In certain embodiments, the EPL is a moiety that binds to CDK9.

A. Moiety for BRD4


[0079] In certain embodiments, the EPL is a moiety that binds to bromodomain-containing protein 4 (BRD4). Exemplary compounds that bind to BRD4 are reported in the literature, including:




as described by Bradner, J. E., et al. in WO 2011/143669;




as described by Fidanze, S. D., et al. in WO 2017/177955.




as described by Wang, S., et al. in WO 2016/138332;




as described by Chen, L., et al. in *ACS Med Chem Lett* 2015, vol. 6(7), page 764;




described by Norris, D. J., et al. in WO 2015/100282.




as described by Yang, S. M., et al. in *Bioorg Med Chem Lett* 2018, vol. 28(21), page 3483;




as described by Ouyang, L., et al. in *J Med Chem* 2017, vol. 60(24), page 9990;




as described by Millan, D. S., et al. in *ACS Med Chem Lett* 2017, vol. 8(8), page 847;




as described in Law, R. P., et al., et al. in *J Med Chem* 2018, vol. 61(10), page 4317;




as described by Aktoudianakis, E., et al. in WO 2014/182929;




as described by Hu, Y., et al., et al. in WO 2018/086605;




as described by Chekler, E. L. P., et al. in WO 2017/037567;




as described by Fish, P. V., et al. in WO 2013/027168;




as described by Demont, E. H., et al. in WO 2011/054848;




as described by Ozer, H. G., et al. in *Cancer Discov* 2018, vol. 8(4), page 458;




as described by Andrews, F. H., et al. in *Proc Natl Acad Sci USA (PNAS)* 2017, vol. 114(7), page E1072;




as described by Zhang, M., et al. in *J Med Chem* 2018, vol. 61(7), page 3037;




as described by Embe, et al. in *ACS Chem Biol* 2014, vol. 9(5), page 1160;



as described by Xiang, Q., et al. in *ACS Med Chem Lett* 2018, vol. 9(3), page 262;




as described by Marineau, J. J., et al. in WO 2015/013635;




as described by Kharenko, O. A., et al. in *J Med Chem* 2018, vol. 61(18), page 8202.

[0080] In certain embodiments, the EPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.


[0081] In certain embodiments, the EPL has the following formula:



as described by Huegle, M., et al. in *J Med Chem* 2016, vol. 59(4), page 1518;



as described by Xue, X. Q., et al. in *Eur J Med Chem* 2018, vol. 152, page 542;



[0082] wherein:

[0083] R^1 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0084] R^2 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0085] R^3 represents independently for each occurrence hydrogen, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, or C_{1-4} alkoxy;

[0086] R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), $-(C_{1-6}$ alkylene)- $C(O)N(R^5)(R^6)$, $-(C_{1-6}$ alkylene)- $N(R^5)C(O)R^7$, $-(C_{1-6}$ alkylene)- CO_2R^7 , $-(C_{1-6}$ alkylene)- $OC(O)R^7$, $-(C_{1-6}$ alkylene)-cyano, $-(C_{1-6}$ alkylene)- $O-(C_{1-6}$ alkyl), C_{1-6} alkyl, C_{3-6} cycloalkyl, or hydrogen; or R^4 and R^5 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring;

[0087] R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl; or R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;

[0088] R^7 is C_{1-6} alkyl, $-(C_{1-6}$ alkylene)-(C_{3-6} cycloalkyl), or C_{3-6} cycloalkyl;

[0089] R^8 is hydrogen or C_{1-4} alkyl; and

[0090] m and n are independently 0, 1, or 2.

[0091] In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl. In certain embodiments, R^3 represents independently for each occurrence hydrogen or halo. In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^3 represents independently for each occurrence hydrogen or fluoro.


[0092] In certain embodiments, R^3 is hydrogen. In certain embodiments, R^3 represents independently for each occurrence halo. In certain embodiments, R^3 represents independently for each occurrence fluoro or chloro. In certain embodiments, R^3 is fluoro. In certain embodiments, R^3 is chloro. In certain embodiments, R^3 represents independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^3 is methyl. In certain embodiments, R^3 represents independently for each occurrence C_{1-4} haloalkyl. In certain embodiments, R^3 represents independently for each occurrence C_{1-4} alkoxy. In certain embodiments, R^3 is methoxy.

[0093] In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2.

[0094] In certain embodiments, R^1 is C_{1-4} alkyl, R^2 represents independently for each occurrence C_{1-4} alkyl, and m is 2. In certain embodiments, R^1 and R^2 are methyl, and m is 2.

[0095] In certain embodiments, each of variables R^1 , R^2 , R^4 , R^5 , R^6 , R^7 , R^8 , and m is as defined in the embodiments below, both singly, and in combination.

[0096] In certain embodiments, the EPL has the following formula:



[0097] wherein:

[0098] R^1 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0099] R^2 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0100] R^3 is hydrogen, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, or C_{1-4} alkoxy;

[0101] R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), $-(C_{1-6}$ alkylene)- $C(O)N(R^5)(R^6)$, $-(C_{1-6}$ alkylene)- $N(R^5)C(O)R^7$, $-(C_{1-6}$ alkylene)- CO_2R^7 , $-(C_{1-6}$ alkylene)- $OC(O)R^7$, $-(C_{1-6}$ alkylene)-cyano, $-(C_{1-6}$ alkylene)- $O-(C_{1-6}$ alkyl), C_{1-6} alkyl, C_{3-6} cycloalkyl, or hydrogen; or R^4 and R^8 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring;

[0102] R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl; or R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;

[0103] R^7 is C_{1-6} alkyl, $-(C_{1-6}$ alkylene)-(C_{3-6} cycloalkyl), or C_{3-6} cycloalkyl;

[0104] R^8 is hydrogen or C_{1-4} alkyl; and

[0105] m is 0, 1, or 2.

[0106] In certain embodiments, R^1 is C_{1-4} alkyl. In certain embodiments, R^1 is methyl. In certain embodiments, R^1 is C_{3-4} cycloalkyl.

[0107] In certain embodiments, R^2 represents independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^2 is methyl. In certain embodiments, R^2 represents independently for each occurrence C_{3-4} cycloalkyl.

[0108] In certain embodiments, R^3 is hydrogen or halo. In certain embodiments, R^3 is hydrogen or fluoro. In certain embodiments, R^3 is halo. In certain embodiments, R^3 is fluoro. In certain embodiments, R^3 is C_{1-4} alkyl. In certain embodiments, R^3 is C_{1-4} haloalkyl. In certain embodiments, R^3 is C_{1-4} alkoxy.

[0109] In certain embodiments, R^4 is $-\text{CH}_2-(\text{oxazolyl})$, methyl, or $-\text{CH}_2-(\text{thiazolyl})$. In certain embodiments, R^4 is $-\text{CH}_2-(\text{oxazol-2-yl})$, methyl, or $-\text{CH}_2-(\text{thiazol-2-yl})$. In certain embodiments, R^4 is $-\text{CH}_2-(\text{oxazolyl})$ or methyl. In certain embodiments, R^4 is $-\text{CH}_2-(\text{oxazol-2-yl})$ or methyl.

[0110] In certain embodiments, R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur). In certain embodiments, R^4 is $-\text{CH}_2-(5-6$ membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur). In certain

embodiments, R^4 is $-\text{CH}_2$ -(5-membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur). In certain embodiments, R^4 is $-\text{CH}_2$ -(oxazolyl) or $-\text{CH}_2$ -(thiazolyl). In certain embodiments, R^4 is $-\text{CH}_2$ -(oxazol-2-yl) or $-\text{CH}_2$ -(thiazol-2-yl). In certain embodiments, R^4 is $-\text{CH}_2$ -(oxazolyl). In certain embodiments, R^4 is $-\text{CH}_2$ -(oxazol-2-yl). In certain embodiments, R^4 is $-\text{CH}_2$ -(thiazolyl). In certain embodiments, R^4 is $-\text{CH}_2$ -(thiazol-2-yl).

[0111] In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})-\text{C}(\text{O})\text{N}(\text{R}^5)(\text{R}^6)$. In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{R}^5)\text{C}(\text{O})\text{R}^7$. In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})-\text{CO}_2\text{R}^7$. In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})-\text{OC}(\text{O})\text{R}^7$. In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})$ -cyano. In certain embodiments, R^4 is $-(\text{C}_{1-6} \text{ alkylene})-\text{O}-(\text{C}_{1-6} \text{ alkyl})$. In certain embodiments, R^4 is C_{1-6} alkyl. In certain embodiments, R^4 is methyl. In certain embodiments, R^4 is C_{3-6} cycloalkyl. In certain embodiments, R^4 is hydrogen. In certain embodiments, R^4 and R^8 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring. In certain embodiments, R^4 is methyl.

[0112] In certain embodiments, R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl. In certain embodiments, R^5 is hydrogen. In certain embodiments, R^5 is C_{1-6} alkyl. In certain embodiments, R^5 is C_{3-6} cycloalkyl. In certain embodiments, R^6 is hydrogen. In certain embodiments, R^6 is C_{1-6} alkyl. In certain embodiments, R^6 is C_{3-6} cycloalkyl. In certain embodiments, R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom.


[0113] In certain embodiments, R^7 is C_{1-6} alkyl. In certain embodiments, R^7 is $-(\text{C}_{1-6} \text{ alkylene})-(\text{C}_{3-6} \text{ cycloalkyl})$. In certain embodiments, R^7 is C_{3-6} cycloalkyl.

[0114] In certain embodiments, R^8 is hydrogen. In certain embodiments, R^8 is C_{1-4} alkyl.


[0115] In certain embodiments, m is 0. In certain embodiments, m is 1. In certain embodiments, m is 2.

[0116] In certain embodiments, R^1 is C_{1-4} alkyl, R^1 and R^3 are independently C_1 - C_6 alkyl, and m is 2. In certain embodiments, R^1 is C_{1-4} alkyl, R^2 represents independently for each occurrence C_{1-4} alkyl, and m is 2. In certain embodiments, R^1 and R^2 are methyl, and m is 2.


[0117] In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is



In certain embodiments, the EPL is




or




wherein R^3 is independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl; and R^4 is methyl, $-\text{CH}_2-(\text{oxazolyl})$, or $-\text{CH}_2-(\text{thiazolyl})$.


[0118] In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is




[0119] In certain embodiments, the EPL is




In certain embodiments, the EPL is




In certain embodiments, the EPL is



In certain embodiments, the EPL is



[0120] In certain embodiments, the EPL has the following formula that is substituted by one occurrence of R^{I-1} :



[0121] wherein:

[0122] R^{I-1} is a bond to L;

[0123] R^1 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0124] R^2 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0125] R^3 represents independently for each occurrence hydrogen, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, or C_{1-4} alkoxy;

[0126] R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), $-(C_{1-6}$ alkylene)- $C(O)N(R^5)(R^6)$, $-(C_{1-6}$ alkylene)- $N(R^5)C(O)R^7$, $-(C_{1-6}$ alkylene)- CO_2R^7 , $-(C_{1-6}$ alkylene)- $OC(O)R^7$, $-(C_{1-6}$ alkylene)-cyano, $-(C_{1-6}$ alkylene)- $O-(C_{1-6}$ alkyl), C_{1-6} alkyl, C_{3-6} cycloalkyl, or hydrogen; or R^4 and R^8 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring;


[0127] R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl; or R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;

[0128] R^7 is C_{1-6} alkyl, $-(C_{1-6}$ alkylene)-(C_{3-6} cycloalkyl), or C_{3-6} cycloalkyl;

[0129] R^8 is hydrogen or C_{1-4} alkyl; and

[0130] m and n are independently 0, 1, or 2.

[0131] In certain embodiments, the EPL has the following formula that is substituted by one occurrence of R^{I-1} :




[0132] wherein:

[0133] R^{I-1} is a bond to L;

[0134] R^1 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

- [0135] R^2 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;
- [0136] R^3 is hydrogen, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, or C_{1-4} alkoxy;
- [0137] R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), $-(C_{1-6}$ alkylene)-C(O)N(R^5)(R^6), $-(C_{1-6}$ alkylene)-N(R^5)C(O) R^7 , $-(C_{1-6}$ alkylene)-CO $2R^7$, $-(C_{1-6}$ alkylene)-OC(O) R^7 , $-(C_{1-6}$ alkylene)-cyano, $-(C_{1-6}$ alkylene)-O $-(C_{1-6}$ alkyl), C_{1-6} alkyl, C_{3-6} cycloalkyl, or hydrogen; or R^4 and R^8 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring;
- [0138] R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl; or R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;
- [0139] R^7 is C_{1-6} alkyl, $-(C_{1-6}$ alkylene)-(C_{3-6} cycloalkyl), or C_{3-6} cycloalkyl;
- [0140] R^8 is hydrogen or C_{1-4} alkyl; and
- [0141] m is 0, 1, or 2.
- [0142] In certain embodiments, the EPL has the following formula:



[0143] wherein:

- [0144] R^1 is phenyl or 6-membered heteroaryl containing 1 or 2 heteroatoms independently selected from nitrogen, oxygen, and sulfur; wherein the phenyl and heteroaryl are optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, hydroxyl, or C_{1-6} alkoxy;
- [0145] R^2 and R^3 each represent independently for each occurrence halo, C_{1-6} alkyl, C_{1-6} haloalkyl, C_{3-6} cycloalkyl, hydroxyl, C_{1-6} alkoxy, or cyano;
- [0146] m is 0, 1, or 2; and
- [0147] n is 0, 1, 2, 3, or 4.
- [0148] In certain embodiments, R^1 is phenyl optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, hydroxyl, or C_{1-6} alkoxy. In certain embodiments, R^1 is 6-membered heteroaryl containing 1 or 2 heteroatoms independently selected from nitrogen, oxygen, and sulfur; wherein the heteroaryl is optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_{1-6} alkyl, C_{1-6} haloalkyl, hydroxyl, or C_{1-6} alkoxy. In certain embodiments, R^1 is phenyl. In certain embodiments, R^1 is 6-membered heteroaryl containing 1 or 2 heteroatoms independently selected from nitrogen, oxygen, and sulfur.
- [0149] In certain embodiments, R^2 is halo. In certain embodiments, R^2 is C_{1-6} alkyl. In certain embodiments, R^2


is C_{1-6} haloalkyl. In certain embodiments, R^2 is C_{3-6} cycloalkyl. In certain embodiments, R^2 is hydroxyl. In certain embodiments, R^2 is C_{1-6} alkoxy. In certain embodiments, R^2 is cyano.

[0150] In certain embodiments, R^3 is halo. In certain embodiments, R^3 is C_{1-6} alkyl. In certain embodiments, R^3 is C_{1-6} haloalkyl. In certain embodiments, R^3 is C_{3-6} cycloalkyl. In certain embodiments, R^3 is hydroxyl. In certain embodiments, R^3 is C_{1-6} alkoxy. In certain embodiments, R^3 is cyano.


[0151] In certain embodiments, m is 0. In certain embodiments, m is 1. In certain embodiments, m is 2. In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2. In certain embodiments, n is 3. In certain embodiments, n is 4.

[0152] In certain embodiments, R^2 and R^3 are independently C_{1-6} alkyl, and m is 2.


[0153] In certain embodiments, the EPL is



In certain embodiments, the EPL is



[0154] In certain embodiments, the EPL has the following formula that is substituted by one occurrence of R^{U-1} :



[0155] wherein:

[0156] R^{I-1} is a bond to L;


[0157] R^1 is phenyl or 6-membered heteroaryl containing 1 or 2 heteroatoms independently selected from nitrogen, oxygen, and sulfur; wherein the phenyl and heteroaryl are optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, hydroxyl, or C_1 - C_6 alkoxy;

[0158] R^2 and R^3 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0159] m is 0, 1, or 2; and

[0160] n is 0, 1, 2, 3, or 4.

[0161] In certain embodiments, the EPL has the following formula:



[0162] wherein:

[0163] R^1 represents independently for each occurrence halo, C_1 - C_4 alkyl, or C_1 - C_4 haloalkyl;

[0164] R^2 is hydrogen, halo, C_1 - C_4 alkyl, or C_1 - C_4 haloalkyl;

[0165] R^3 is $-N(R^8)SO_2R^9$, $-SO_2N(R^8)_2$, $-SO_2R^9$, $-(C_{1-6} \text{ alkylene})-SO_2R^9$, C_{1-6} hydroxyalkyl, or a 4-7 membered saturated carbocyclic ring in which one CH_2 is replaced with SO_2 ;

[0166] R^4 is hydrogen, halo, or C_1 - C_4 alkyl;

[0167] R^5 is C_1 - C_4 alkyl or C_{3-4} cycloalkyl;

[0168] R^6 is hydrogen, C_1 - C_4 alkyl, or C_{3-4} cycloalkyl;

[0169] R^7 represents independently for each occurrence C_1 - C_4 alkyl or C_{3-4} cycloalkyl;

[0170] R^8 represents independently for each occurrence hydrogen or C_1 - C_4 alkyl; or two occurrences of R^8 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered heterocyclic ring containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen; or R^8 and R^9 are taken together with their intervening atoms to form a 5-7 membered ring containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur;

[0171] R^9 is C_{1-6} alkyl, C_{1-6} haloalkyl, $-(C_{1-6} \text{ alkylene})-(C_{3-6} \text{ cycloalkyl})$, or C_{3-6} cycloalkyl;

[0172] A^3 is phenylene, a 3-10 membered saturated monocyclic, bicyclic or spirocyclic carbocyclylene, or C_{1-6} alkylene; and

[0173] p and t are independently 0, 1, or 2.

[0174] In certain embodiments, R^1 represents independently for each occurrence halo. In certain embodiments, R^1

represents independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^1 represents independently for each occurrence C_{1-4} haloalkyl.

[0175] In certain embodiments, R^2 is hydrogen. In certain embodiments, R^2 is halo. In certain embodiments, R^2 is C_{1-4} alkyl. In certain embodiments, R^2 is C_{1-4} haloalkyl.

[0176] In certain embodiments, R^3 is $-N(R^8)SO_2R^9$. In certain embodiments, R^3 is $-SO_2N(R^8)_2$. In certain embodiments, R^3 is $-SO_2R^9$. In certain embodiments, R^3 is $-(C_{1-6} \text{ alkylene})-SO_2R^9$. In certain embodiments, R^3 is C_{1-6} hydroxyalkyl. In certain embodiments, R^3 is a 4-7 membered saturated carbocyclic ring in which one CH_2 is replaced with SO_2 .

[0177] In certain embodiments, R^4 is hydrogen. In certain embodiments, R^4 is halo. In certain embodiments, R^4 is C_{1-4} alkyl. In certain embodiments, R^5 is C_{1-4} alkyl. In certain embodiments, R^5 is C_{3-4} cycloalkyl. In certain embodiments, R^6 is hydrogen. In certain embodiments, R^6 is C_{1-4} alkyl. In certain embodiments, R^6 is C_{3-4} cycloalkyl. In certain embodiments, R^7 represents independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^7 represents independently for each occurrence C_{3-4} cycloalkyl.


[0178] In certain embodiments, R^8 represents independently for each occurrence hydrogen. In certain embodiments, R^8 represents independently for each occurrence C_{1-4} alkyl. In certain embodiments, two occurrences of R^8 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered heterocyclic ring containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen; or R^8 and R^9 are taken together with their intervening atoms to form a 5-7 membered ring containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

[0179] In certain embodiments, R^9 is C_{1-6} alkyl. In certain embodiments, R^9 is C_{1-6} haloalkyl. In certain embodiments, R^9 is $-(C_{1-6} \text{ alkylene})-(C_{3-6} \text{ cycloalkyl})$. In certain embodiments, R^9 is C_{3-6} cycloalkyl.

[0180] In certain embodiments, A^3 is phenylene. In certain embodiments, A^3 is a 3-10 membered saturated monocyclic, bicyclic or spirocyclic carbocyclylene. In certain embodiments, A^3 is C_{1-6} alkylene.

[0181] In certain embodiments, p is 0. In certain embodiments, p is 1. In certain embodiments, p is 2. In certain embodiments, t is 0. In certain embodiments, t is 1. In certain embodiments, t is 2.

[0182] In certain embodiments, the EPL has the following formula:



[0183] wherein:

[0184] R^1 represents independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, or hydroxyl;

[0185] R^2 represents independently for each occurrence hydrogen, C_1 - C_6 alkyl, or C_3 - C_6 cycloalkyl;

[0186] R^3 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl; and


[0187] n represents independently for each occurrence 0, 1, 2, or 3.

[0188] In certain embodiments, R^1 represents independently for each occurrence halo. In certain embodiments, R^1 represents independently for each occurrence C_1 - C_6 alkyl. In certain embodiments, R^1 represents independently for each occurrence C_1 - C_6 haloalkyl. In certain embodiments, R^1 represents independently for each occurrence C_3 - C_6 cycloalkyl. In certain embodiments, R^1 represents independently for each occurrence hydroxyl.


[0189] In certain embodiments, R^2 represents independently for each occurrence hydrogen. In certain embodiments, R^2 represents independently for each occurrence C_1 - C_6 alkyl. In certain embodiments, R^2 represents independently for each occurrence C_3 - C_6 cycloalkyl.

[0190] In certain embodiments, R^3 is C_1 - C_6 alkyl. In certain embodiments, R^3 is C_3 - C_6 cycloalkyl. In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2. In certain embodiments, n is 3.


[0191] In certain embodiments, the EPL is



[0192] In certain embodiments, the EPL is



[0193] In certain embodiments, the EPL has the following formula that is substituted by one occurrence of R^{1z-1} :



[0194] wherein:

[0195] R^{1z-1} is a bond to L;

[0196] R^1 represents independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, or hydroxyl;


[0197] R^2 represents independently for each occurrence hydrogen, C_1 - C_6 alkyl, or C_3 - C_6 cycloalkyl;

[0198] R^3 is C_1 - C_6 alkyl or C_3 - C_6 cycloalkyl; and


[0199] n represents independently for each occurrence 0, 1, 2, or 3.

B. Moiety for CDK1


[0200] In certain embodiments, the EPL is a moiety that binds to Cyclin-Dependent Kinase 1 (CDK1). Exemplary compounds that bind to CDK1 are reported in the literature, including:




as described by Sivakumar, M., et al. in WO2007/148158;




as described by Lucking, U., et al. in WO2005/037800;




as described by D'Alessio, R., et al. in WO2004/104007;




as described by Guzi, T. J., et al. in WO2005/077954;




as described by Wyatt, P. G., et al. in WO2005/012256;



as described by Brumby, T., et al. in WO2002/096888;




as described by Dumont, J. A., et al. in WO2000/044362;




as described by Wang, S., et al. in WO2013/156780;

as described by Wang, Z., et al. in WO2004/092139;




as described by Wang, S., et al. in WO2009/118567;



[0201] as described by Caligiuri, M., et al. in Chem Biol (London) 2005, 12(10): 1103.

[0202] In certain embodiments, the EPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.

[0203] In certain embodiments, the EPL has the following formula:




[0204] wherein:

[0205] R^1 , R^2 and R^3 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;


[0206] m and p each represent independently 0, 1, or 2; and

[0207] n is 0, 1 or 2.

[0208] In certain embodiments the EPL is



[0209] In certain embodiments, the EPL has the following formula:



[0210] wherein:


[0211] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0212] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C_1 - C_6 alkoxy),

[0213] m is 0, 1, or 2; and

[0214] n and p each represent independently 0, 1, 2, or 3.

[0215] In certain embodiments, the EPL has the following formula:



[0216] wherein:


[0217] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0218] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C_1 - C_6 alkoxy),

[0219] m is 0, 1, or 2; and

[0220] n and p each represent independently 0, 1, 2, or 3.

[0221] In certain embodiments, the EPL has the following formula:



[0222] wherein:

[0223] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;


[0224] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C₁-C₆ alkoxy),

[0225] m is 0, 1, or 2; and


[0226] n and p each represent independently 0, 1, 2, or 3.

C. Moiety for CDK2


[0227] In certain embodiments, the EPL is a moiety that binds to Cyclin-Dependent Kinase 2 (CDK2). Exemplary compounds that bind to CDK2 are reported in the literature, including:




as described by Lucking, U., et al. in WO2005/037800;




as described by D'Alessio, R., et al. in WO2004/104007;




as described by Misra, R. N., et al. in WO2001/044242;




as described by Sheldrake, P. W., et al. in WO2008/122767;




as described by Guzi, T. J., et al. in WO2005/077954;




as described by Wyatt, P. G., et al. in WO2005/012256;



described by Brumby, T., et al. in WO2002/096888;




as described by Dumont, J. A., et al. in WO2000/044362;



as described by Hao, M., et al. in WO2017/044858.

[0228] In certain embodiments, the EPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.

[0229] In certain embodiments, the EPL has the following formula:



[0230] wherein:

[0231] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;


[0232] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloal-

kyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C_1 - C_6 alkoxy),

[0233] m is 0, 1, or 2; and

[0234] n and p each represent independently 0, 1, 2, or 3.

[0235] In certain embodiments, the EPL has the following formula:



[0236] wherein:

[0237] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;


[0238] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloal-

kyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C_1 - C_6 alkoxy),

[0239] m is 0, 1, or 2; and

[0240] n and p each represent independently 0, 1, 2, or 3.

[0241] In certain embodiments, the EPL has the following formula:





[0242] wherein:

[0243] R^1 , R^2 and R^3 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0244] m and p each represent independently 0, 1, or 2; and

[0245] n is 0, 1 or 2.



[0246] In certain embodiments the EPL is




D. Moiety for CDK9

[0247] In certain embodiments, the EPL is a moiety that binds to Cyclin-Dependent Kinase 9 (CDK9). Exemplary compounds that bind to CDK9 are reported in the literature, including:


as described by Pike, K. G., et al. in WO2017001354;




as described by D'Alessio, R., et al. in WO2004/104007;



as described by Sivakumar, M., et al. in WO2007/148158;




as described by Blanchard, S., et al. in WO2007/058628;




as described by Lucking, U., et al. in WO2005/037800;


as described by Misra, R. N., et al. in WO2001/044242;




as described by Sheldrake, P. W. et al. in WO2008/122767;




as described by Brumby, T., et al. in WO2002/096888;




as described by Guzi, T. J., et al. in WO2005/077954;




as described by Dumont, J. A., et al. in WO2000/044362;




as described by Hao, M., et al. in WO2017/044858;



as described by Wyatt, P. G., et al. in WO2005/012256;




as described by Smith, C. D., et al. in WO2018/089902;



as described by Gao, Q., et al. in CN105111191.

[0248] In certain embodiments, the EPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.

[0249] In certain embodiments, the EPL has the following formula:



[0250] wherein:


[0251] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0252] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C₁-C₆ alkoxy),

[0253] m is 0, 1, or 2; and

[0254] n and p each represent independently 0, 1, 2, or 3.

[0255] In certain embodiments, the EPL has the following formula:



[0256] wherein:


[0257] R^1 and R^2 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0258] R^3 represents independently for each occurrence C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, C_1 - C_6 hydroxyalkyl, or $-(C_1$ - C_6 alkylene)-(C₁-C₆ alkoxy),

[0259] m is 0, 1, or 2; and

[0260] n and p each represent independently 0, 1, 2, or 3.

[0261] In certain embodiments, the EPL has the following formula:




[0262] wherein:

[0263] R^1 , R^2 and R^3 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;

[0264] m and p each represent independently 0, 1, or 2; and

[0265] n is 0, 1 or 2.

[0266] In certain embodiments, the EPL is




[0267] In certain embodiments, the EPL is selected from those depicted in the compounds in Tables 1, 3, 4, and 5, below. In certain embodiments, the EPL is selected from those depicted in the compounds in Tables 3 and 4, below. In certain embodiments, the EPL is selected from those depicted in the compounds in Table 1, below. In certain embodiments, the EPL is selected from those depicted in the compounds in Table 3, below. In certain embodiments, the EPL is selected from those depicted in the compounds in Table 4, below. In certain embodiments, the EPL is selected from those depicted in the compounds in Table 5, below.

[0268] The description above regarding the identity of the EPL, and in Parts C and D below regarding the identity of the TPL and linker, describe multiple embodiments relating to compounds of Formula I. The patent application specifically contemplates all combinations of the embodiments.

Part B: Compounds of Formula II


[0269] Another aspect of the invention provides a compound represented by the formula:




or a pharmaceutically acceptable salt thereof, wherein EPL is a moiety that binds to an effector protein selected from

Eukaryotic Peptide Chain Release Factor GTP-Binding Subunit ERF3A (GSPT1); L is a linker; and TPL is a moiety that binds to estrogen receptor.


[0270] Exemplary moieties that bind GSPT1 are reported in the literature, including:




as described in Luo, Y. et al., in WO2021047627.




as described in Gray, N. et al., in WO2020006264.




as described in Chan, K. et al., in US2020369679.




as described in Chan, K. et al., in WO2019241271.




as described in Chan, K. et al., in WO2019241274.




as described in Chan, K. et al., in WO2019241274.



as described in Chan, K. et al., in US2018298027.




as described in Muller, G. et al., in US2009142297.



as described in Hansen, J. et al., in WO2016007848.

[0271] In certain embodiments, the EPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.

[0272] In certain embodiments, the EPL has the following formula:



[0273] wherein:



[0274] R^{1a} is hydrogen, halo, or C_{1-4} alkyl;

[0275] R^{2a} and R^{3a} are independently hydrogen or C_{1-4} alkyl;


[0276] R^{4a} represents independently for each occurrence halo, C_{1-4} alkyl, C_{1-4} haloalkyl, hydroxyl, or C_{1-4} alkoxy; and

[0277] n is 0, 1, 2, or 3.

[0278] In certain embodiments, the EPL is one of the following:




In certain embodiments, the EPL is



In certain embodiments, EPL is selected from those depicted in the compounds in Table 2, below.

[0279] Another aspect of the invention provides a compound represented by Formula II:



[0280] or a pharmaceutically acceptable salt thereof, wherein:

[0281] R^1 is hydrogen, halo, or C_{1-4} alkyl;


[0282] R^2 and R^3 are independently hydrogen or C_{1-4} alkyl;

[0283] R^4 represents independently for each occurrence halo, C_{1-4} alkyl, C_{1-4} haloalkyl, hydroxyl, or C_{1-4} alkoxy;

[0284] n is 0, 1, 2, or 3;

[0285] L is a linker; and

[0286] TPL is represented by:




[0287] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0288] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy.


[0289] As generally defined above for Formula II, R^1 is hydrogen, halo, or C_{1-4} alkyl. In certain embodiments, R^1 is hydrogen. In certain embodiments, R^1 is halo. In certain embodiments, R^1 is C_{1-4} alkyl. In certain embodiments, R^1 , R^2 , and R^3 are hydrogen.

[0290] As generally defined above for Formula II, R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy. In certain embodiments, R^{1B} is hydroxyl. In certain embodiments, R^{1B} is halo. In certain embodiments, R^{1B} is C_{1-4} alkyl. In certain embodiments, R^{1B} is C_{1-4} alkoxy. In certain embodiments, R^{1C} is hydroxyl. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1C} is C_{1-4} alkoxy.

[0291] In certain embodiments, the compound is represented by one of the following:



-continued




[0292] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1B} and R^{1C} are hydroxyl. In certain embodiments, R^{1B} is halo, and R^{1C} is hydroxyl.


Part C: Exemplary Further Description of the TPL Component of Compounds of Formula I and Formula II

[0293] Compounds of Formula I and II may be further characterized according to, for example, the identity of the TPL component. As generally described above, the TPL is a moiety that binds to an estrogen receptor. Exemplary moieties for the TPL component are described in more detail below.


[0294] In certain embodiments, the TPL is a moiety that is an activator, inhibitor, and/or bind to the estrogen receptor (ER). Compounds that activate, inhibit, and/or bind to the ER are reported in the literature, which include:




as described in Bock, M. et al., in US20160347717.




as described in Palkowitz, A., in U.S. Pat. No. 5,488,058.




as described in Cameron, K. et al., in WO1995010513.




as described in Duan, S. et al., in WO2020125640.




as described in Nanjyo, S. et al., in Bioorg Med Chem 2019, 27(10): 1952.




as described in Wang, G. et al., in WO2020055973.




as described in Yang, F. et al., in WO2019223715.




as described in Wang, G. et al., in WO2020055973.




as described in Yang, F. et al., in WO2019223715.




as described in Ruenitz, P., in 219th Am Chem Soc (ACS) Natl Meet 2000-03-26/2000-03-30 San Francisco, United States Abst MEDI 330.




as described in Watanabe, N. et al., in Bioorg Med Chem Lett 2003, 13(24): 4317.




as described in Scott, J. et al., in ACS Med Chem Lett 2016, 7(1): 94.




as described in Scott, J. et al., in ACS Med Chem Lett 2016, 7(1): 94.




as described in Bouaboula, M. et al., in US2020392081.




as described in Dalton, J. et al., in WO2008091555.




as described in Watanabe, N. et al., in J Med Chem 2003, 46(19): 3961.



as described in Nanjyo, S. et al., in *Bioorg Med Chem* 2019, 27(10): 1952.



as described in Miller, C. et al., in *J Med Chem* 2001, 44(11): 1654.




as described in Cameron, K. et al., in WO1995010513.

[0295] In certain embodiments, the TPL is a moiety that binds to the estrogen receptor (ER). Exemplary compounds that bind to ER are reported in the literature, such as raloxifene, H3B-6545, and AZD9496. A radical of such compounds reported in the literature that bind ER are amenable for use in the present invention.

[0296] In certain embodiments, the TPL is a radical of one of the above compounds, which is attached to L through a modifiable oxygen, nitrogen, or carbon atom.

[0297] In certain embodiments, the TPL is




[0298] wherein:

[0299] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0300] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy.

[0301] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is halo. In certain embodiments, R^{1A} is C_{1-4} alkyl.


[0302] In certain embodiments, the TPL is




[0303] wherein R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy.

[0304] In certain embodiments, R^{1B} is hydroxyl. In certain embodiments, R^{1B} is halo. In certain embodiments, R^{1B} is C_{1-4} alkyl. In certain embodiments, R^{1B} is C_{1-4} alkoxy. In certain embodiments, R^{1C} is hydroxyl. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1C} is C_{1-4} alkoxy. In certain embodiments, R^{1B} and R^{1C} are hydroxyl. In certain embodiments, R^{1B} is halo, and R^{1C} is hydroxyl. In certain embodiments, R^{1B} and R^{1C} are methoxy.

[0305] In certain embodiments, the TPL is



[0306] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{1A} :



[0307] wherein:



[0308] R^{1A} is a bond to L;

[0309] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0310] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy.

[0311] In certain embodiments, the TPL is

In certain embodiments, the TPL is



[0312] wherein:

[0313] A^1 is phenylene or 6-membered heteroarylene containing 1 or 2 heteroatoms that are nitrogen;



[0314] R^{1C} and R^{1E} are each independently for each occurrence hydrogen, halo, C_{1-4} alkoxy, C_{1-4} alkyl, or C_{1-4} haloalkyl;

[0315] R^{1D} is hydroxyl, $-CO_2H$, or $-B(OH)_2$; and

[0316] q and r are independently 0, 1, 2, or 3.

[0317] In certain embodiments, A^1 is phenylene. In certain embodiments, A^1 is a 6-membered heteroarylene containing 1 or 2 heteroatoms that are nitrogen. In certain embodiments, A^1 is pyridinylene, pyrimidinylene, or pyridazinylene. In certain embodiments, A^1 is

In certain embodiments, the TPL is




wherein the attachment point adjacent to a nitrogen atom of A^1 is connected to L.


[0318] In certain embodiments, A^1 is para-phenylene, and R^{1D} is hydroxyl.

[0319] In certain embodiments, R^{1D} is hydroxyl. In certain embodiments, R^{1D} is $-\text{CO}_2\text{H}$. In certain embodiments, R^{1D} is $-\text{B}(\text{OH})_2$.


[0320] In certain embodiments, the TPL is




In certain embodiments, the TPL is




or



In certain embodiments, the TPL is










In certain embodiments the TPL is





or

[0321] In certain embodiments, the TPL is one of the following:






[0322] In certain embodiments, the TPL is one of the following:



[0323] In certain embodiments, the TPL is one of the following:



[0324] In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen or halo. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, methyl, or trifluoromethyl. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl.


[0325] In certain embodiments, R^{1C} is independently for each occurrence halo, C_{1-4} alkoxy, or C_{1-4} alkyl. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or halo. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or C_{1-4} alkoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or C_{1-4} alkyl. In certain embodiments, R^{1C} is independently for each occurrence halo or C_{1-4} alkoxy. In certain embodiments, R^{1C} is independently for each occurrence halo or C_{1-4} alkyl.

[0326] In certain embodiments, R^{1C} is independently for each occurrence halo. In certain embodiments, R^{1C} is independently for each occurrence C_{1-4} alkoxy. In certain embodiments, R^{1C} is independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^{1C} is independently for each occurrence C_{1-4} haloalkyl. In certain embodiments, R^{1C} is $-\text{CF}_3$. In certain embodiments, R^{1C} is independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl.

[0327] In certain embodiments, R^{1E} is independently for each occurrence halo, C_{1-4} alkoxy, or C_{1-4} alkyl. In certain embodiments, R^{1E} is independently for each occurrence hydrogen or halo. In certain embodiments, R^{1E} is independently for each occurrence hydrogen or C_{1-4} alkyl. In certain embodiments, R^{1E} is independently for each occurrence halo or C_{1-4} alkyl.

[0328] In certain embodiments, R^{1E} is independently for each occurrence halo. In certain embodiments, R^{1E} is independently for each occurrence C_{1-4} alkoxy. In certain embodiments, R^{1E} is independently for each occurrence C_{1-4} alkyl. In certain embodiments, R^{1E} is independently for each occurrence C_{1-4} haloalkyl. In certain embodiments, R^{1E} is $-\text{CF}_3$. In certain embodiments, R^{1E} is independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl.



[0329] In certain embodiments, the TPL is one of the following:



[0330] wherein R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl; and q and r are independently 0, 1, or 2.

[0331] In certain embodiments, q and r are independently 0 or 1. In certain embodiments, q and r are independently 1 or 2. In certain embodiments, q and r are independently 0, 1, or 2. In certain embodiments, q is 0. In certain embodiments, q is 1. In certain embodiments, q is 2. In certain embodiments, q is 3. In certain embodiments, q is 0 or 1. In certain embodiments, q is 1 or 2. In certain embodiments, q is 0, 1, or 2. In certain embodiments, r is 0. In certain embodiments, r is 1. In certain embodiments, r is 2. In certain embodiments, r is 3. In certain embodiments, r is 0 or 1. In certain embodiments, r is 1 or 2. In certain embodiments, r is 0, 1, or 2.

[0332] In certain embodiments, the TPL is one of the following:



[0333] wherein R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl.


[0334] In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence fluoro, chloro, methoxy, or methyl. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1C} and R^{1E} are hydrogen. In certain embodiments, R^{1C} and R^{1E} are fluoro.

[0335] In certain embodiments, R^{1C} is independently for each occurrence fluoro, chloro, methoxy, or methyl. In certain embodiments, R^{1C} is independently for each occurrence

hydrogen, fluoro, chloro, or methoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or methoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or methyl. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methoxy. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methyl. In certain embodiments, R^{1C} is methoxy.

[0336] In certain embodiments, R^{1E} is independently for each occurrence fluoro, chloro, methoxy, or methyl. In certain embodiments, R^{1E} is independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1E} is independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or methyl. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methyl. In certain embodiments, R^{1C} is methoxy.

[0337] In certain embodiments, the TPL is



[0338] wherein:

[0339] R^{1C} and R^{1E} are independently hydrogen, halo, or C_{1-4} alkyl; and


[0340] R^{1D} is hydroxyl.

[0341] In certain embodiments, R^{1C} and R^{1E} are independently hydrogen or halo. In certain embodiments, R^{1C} and R^{1E} are independently hydrogen or fluoro. In certain embodiments, R^{1C} and R^{1E} are hydrogen. In certain embodiments, R^{1C} and R^{1E} are fluoro.


[0342] In certain embodiments, R^{1C} is hydrogen or halo. In certain embodiments, R^{1C} is hydrogen or fluoro. In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is fluoro. In certain embodiments, R^{1C} is chloro. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1C} is methyl.

[0343] In certain embodiments, R^{1E} is hydrogen or halo. In certain embodiments, R^{1E} is hydrogen or fluoro. In certain embodiments, R^{1E} is hydrogen. In certain embodiments, R^{1E} is halo. In certain embodiments, R^{1E} is fluoro. In certain embodiments, R^{1E} is chloro. In certain embodiments, R^{1E} is C_{1-4} alkyl. In certain embodiments, R^{1E} is methyl.


[0344] In certain embodiments, the TPL is




In certain embodiments, the TPL is




In certain embodiments, the TPL is



In certain embodiments, the TPL is



[0345] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0346] wherein:

[0347] R^{II-1A} is a bond to L;

[0348] A^1 is phenyl or 6-membered heteroaryl containing 1 or 2 heteroatoms that are nitrogen;


[0349] R^{1C} and R^{1E} are each independently for each occurrence hydrogen, halo, C_{1-4} alkoxy, C_{1-4} alkyl, or C_{1-4} haloalkyl;

[0350] R^{1D} is hydroxyl, $-CO_2H$, or $-B(OH)_2$; and

[0351] q and r are independently 0, 1, 2, or 3.

In certain embodiments, each of the variables is independently as described and defined in embodiments above.

[0352] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0353] wherein:


[0354] R^{II-1A} is a bond to L;

[0355] R^{1C} and R^{1E} are independently hydrogen, halo, or C_{1-4} alkyl; and

[0356] R^{1D} is hydroxyl.

[0357] In certain embodiments, each of the variables is independently as described and defined in embodiments above.

[0358] In certain embodiments, the TPL is



[0359] wherein:


[0360] R^{1C} and R^{1E} are independently hydrogen, halo, or C_{1-4} alkyl;

[0361] R^{1D} is hydroxyl; and


[0362] R^{1F} is hydrogen or C_{1-4} alkyl.

[0363] In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1E} is hydrogen. In certain embodiments, R^{1E} is halo. In certain embodiments, R^{1E} is C_{1-4} alkyl. In certain embodiments, R^{1F} is hydrogen. In certain embodiments, R^{1F} is C_{1-4} alkyl.

[0364] In certain embodiments, the TPL is



[0365] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{1A-1D} :



[0366] wherein:


[0367] R^{1A-1D} is a bond to L;

[0368] R^{1C} and R^{1E} are independently hydrogen, halo, or C_{1-4} alkyl;

[0369] R^{1D} is hydroxyl; and

[0370] R^{1F} is hydrogen or C_{1-4} alkyl.

[0371] In certain embodiments, the TPL is



[0372] wherein:

[0373] R^{1C} represents independently for each occurrence hydrogen, halo, hydroxyl, or C_{1-4} alkyl;

[0374] R^{1D} is hydroxyl;

[0375] R^{1E} represents independently for each occurrence halo, hydroxyl, or C_{1-4} alkyl;


[0376] R^{1F} is hydrogen or C_{1-4} alkyl;

[0377] y is 1 or 2; and


[0378] z is 1, 2, or 3.

[0379] In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is hydroxyl. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1E} is halo. In certain embodiments, R^{1E} is fluoro. In certain embodiments, R^{1E} is hydroxyl. In certain embodiments, R^{1E} is C_{1-4} alkyl. In certain embodiments, R^{1F} is hydrogen. In certain embodiments, R^{1F} is C_{1-4} alkyl. In certain embodiments, R^{1F} is $-CH_3$. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, z is 1. In certain embodiments, z is 2. In certain embodiments, z is 3.


[0380] In certain embodiments, the TPL is




In certain embodiments, the TPL is




In certain embodiments, the TPL is




In certain embodiments, the TPL is




In certain embodiments the TPL is



In certain embodiments, the TPL is



[0381] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of $R^{T-1,4}$:



[0382] wherein:

[0383] $R^{T-1,4}$ is a bond to L;

[0384] R^{1C} represents independently for each occurrence hydrogen, halo, hydroxyl, or C_{1-4} alkyl;

[0385] R^{1D} is hydroxyl;


[0386] R^{1E} represents independently for each occurrence halo, hydroxyl, or C_{1-4} alkyl;

[0387] R^{1F} is hydrogen or C_{1-4} alkyl;

[0388] y is 1 or 2; and

[0389] z is 1, 2, or 3.

[0390] In certain embodiments, the TPL is



[0391] wherein:

[0392] R^{1C} is hydrogen, hydroxyl, or C_{1-4} alkyl;

[0393] R^{1D} is hydroxyl;


[0394] R^{1E} represents independently for each occurrence halo, hydroxyl, or C_{1-4} alkyl;

[0395] R^{1F} is hydrogen or C_{1-4} alkyl; and

[0396] y is 1, 2, or 3.


[0397] In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is hydroxyl. In certain embodiments, R^{1C} is C_{1-4} alkyl. In certain embodiments, R^{1E} is halo. In certain embodiments, R^{1E} is fluoro. In certain embodiments, R^{1E} is hydroxyl. In certain embodiments, R^{1E} is C_{1-4} alkyl.

In certain embodiments, R^{1F} is hydrogen. In certain embodiments, R^{1F} is C_{1-4} alkyl. In certain embodiments, R^{1F} is $-CH_3$. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, y is 3.



[0398] In certain embodiments, the TPL is F

[0399] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0400] wherein:

[0401] R^{II-1A} is a bond to L;

[0402] R^{1C} is hydrogen, hydroxyl, or C_{1-4} alkyl;


[0403] R^{1D} is hydroxyl;

[0404] R^{1E} represents independently for each occurrence halo, hydroxyl, or C_{1-4} alkyl;

[0405] R^{1F} is hydrogen or C_{1-4} alkyl; and

[0406] y is 1, 2, or 3.

[0407] In certain embodiments, the TPL is



[0408] wherein:


[0409] R^{1A} is hydrogen or C_{1-4} alkyl; and

[0410] R^{1B} is $-C(O)CH_3$, hydrogen, hydroxyl, or C_{1-4} alkyl.


[0411] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is C_{1-4} alkyl. In certain embodiments, R^{1B} is $-C(O)CH_3$. In certain embodiments, R^{1B} is hydro-

gen. In certain embodiments, R^{1B} is hydroxyl. In certain embodiments, R^{1B} is C_{1-4} alkyl.

[0412] In certain embodiments, the TPL is



[0413] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :




[0414] wherein:

[0415] R^{II-1A} is a bond to L;

[0416] R^{1A} is hydrogen or C_{1-4} alkyl; and

[0417] R^{1B} is $-C(O)CH_3$, hydrogen, hydroxyl, or C_{1-4} alkyl.

[0418] In certain embodiments, the TPL is




[0419] wherein:

[0420] R^{1A} represents independently for each occurrence hydrogen, C_{1-4} haloalkyl, C_{1-4} alkyl, or halo;


[0421] R^{1B} is hydrogen, halo, or C_{1-4} alkyl; and

[0422] y is 1, 2, or 3.

[0423] In certain embodiments, the TPL is




In certain embodiments, the TPL is




[0424] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is C_{1-4} haloalkyl. In certain embodiments, R^{1A} is C_{1-4} alkyl. In certain embodiments, R^{1A} is halo. In certain embodiments, R^{1B} is hydrogen. In certain embodiments, R^{1B} is halo. In certain embodiments, R^{1B} is C_{1-4} alkyl. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, y is 3.


[0425] In certain embodiments, the TPL is



In certain embodiments, the TPL is



[0426] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0427] wherein:


[0428] R^{III-1A} is a bond to L;

[0429] R^{1A} represents independently for each occurrence hydrogen, C_{1-4} haloalkyl, C_{1-4} alkyl, or halo;

[0430] R^{1B} is hydrogen, halo, or C_{1-4} alkyl; and

[0431] y is 1, 2, or 3.

[0432] In certain embodiments, the TPL is



[0433] wherein:

[0434] R^{1A} represents independently for each occurrence hydrogen, C_{1-4} haloalkyl, C_{1-4} alkyl, or halo;


[0435] R^{1B} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkyl;

[0436] y is 1, 2, or 3; and


[0437] z is 1 or 2.

[0438] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is C_{1-4} haloalkyl. In certain embodiments, R^{1A} is C_{1-4} alkyl. In certain embodiments, R^{1A} is halo. In certain embodiments, R^{1B} is hydrogen. In certain embodiments, R^{1B} is halo. In certain embodiments, R^{1B} is C_{1-4} alkyl. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, y is 3. In certain embodiments, z is 1. In certain embodiments, z is 2.

[0439] In certain embodiments, the TPL is



[0440] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{1A} :



[0441] wherein:

[0442] R^{1A} is a bond to L;


[0443] R^{1A} represents independently for each occurrence hydrogen, C_{1-4} haloalkyl, C_{1-4} alkyl, or halo;

[0444] R^{1B} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkyl;

[0445] y is 1, 2, or 3; and

[0446] z is 1 or 2.

[0447] In certain embodiments, the TPL is



[0448] wherein:

[0449] R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, halo, C_{1-4} alkyl, or C_{1-4} alkoxy;

[0450] R^{1B} is $-C(O)OH$, hydrogen, halo, hydroxyl, $-B(OH)_2$, or C_{1-4} alkyl;


[0451] q is 1 or 2; and

[0452] y is 1, 2, or 3.

[0453] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, chloro, methyl, or methoxy. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkoxy. In certain embodiments, R^{1A} represents indepen-

dently for each occurrence hydrogen or halo. In certain embodiments, R^{1A} represents independently for each occurrence C_{1-4} alkoxy. In certain embodiments, R^{1B} is $-C(O)OH$, $-OH$, or $-B(OH)_2$. In certain embodiments, R^{1C} represents independently for each occurrence hydrogen, fluoro, chloro, methyl, or methoxy. In certain embodiments, R^{1C} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkoxy. In certain embodiments, R^{1C} represents independently for each occurrence hydrogen or halo.

[0454] In certain embodiments, the TPL is



[0455] wherein:

[0456] R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, fluoro, chloro, methyl, or methoxy;


[0457] R^{1B} is $-C(O)OH$, $-OH$, or $-B(OH)_2$;

[0458] q is 1 or 2; and

[0459] y is 1, 2, or 3.

[0460] In certain embodiments, q is 1. In certain embodiments, q is 2.

[0461] In certain embodiments, the TPL is



[0462] wherein:


[0463] R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, fluoro, chloro, methyl, or methoxy; and

[0464] R^{1B} is $-C(O)OH$, $-OH$, or $-B(OH)_2$.

[0465] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} is hydrogen or fluoro. In certain embodiments, R^{1A} is methoxy. In certain embodiments, R^{1B} is $-B(OH)_2$. In certain embodiments, R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In

certain embodiments, R^{1C} is hydrogen or fluoro. In certain embodiments, R^{1C} is fluoro or chloro. In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is fluoro. In certain embodiments, R^{1C} is chloro. In certain embodiments, R^{1C} is methoxy. In certain embodiments, R^{1C} is methyl.

[0466] In certain embodiments the TPL is



[0467] wherein:

[0468] R^{1A} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkyl;

[0469] R^{1B} is $-C(O)OH$, hydrogen, halo, hydroxyl, or C_{1-4} alkyl; and


[0470] y is 1, 2, or 3.

[0471] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, chloro, or methyl. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen or halo. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen or chloro.

[0472] In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is halo. In certain embodiments, R^{1A} is fluoro. In certain embodiments, R^{1A} is chloro. In certain embodiments, R^{1A} is C_{1-4} alkyl. In certain embodiments, R^{1A} is methyl.

[0473] In certain embodiments, R^{1B} is $-C(O)OH$ or $-OH$. In certain embodiments, R^{1B} is $-C(O)OH$. In certain embodiments, R^{1B} is hydrogen. In certain embodiments, R^{1B} is halo. In certain embodiments, R^{1B} is hydroxyl. In certain embodiments, R^{1B} is C_{1-4} alkyl. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, y is 3.

[0474] In certain embodiments, the TPL is




[0475] wherein R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, fluoro, chloro, methyl, or methoxy.

[0476] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} is hydrogen or fluoro.


[0477] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen or chloro. In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is fluoro. In certain embodiments, R^{1A} is chloro.

[0478] In certain embodiments, R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} is hydrogen or fluoro. In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is fluoro.


[0479] In certain embodiments, the TPL is



In certain embodiments, the TPL is



[0480] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0481] wherein:

[0482] R^{III-1A} is a bond to L;


[0483] R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, halo, C_{1-4} alkyl, or C_{1-4} alkoxy;

[0484] R^{1B} is $-C(O)OH$, hydrogen, halo, hydroxyl, $-B(OH)_2$, or C_{1-4} alkyl;

[0485] q is 1 or 2; and

[0486] y is 1, 2, or 3.

[0487] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0488] wherein:


[0489] R^{III-1A} is a bond to L;

[0490] R^{1A} represents independently for each occurrence hydrogen, halo, or C_{1-4} alkyl;

[0491] R^{1B} is $-C(O)OH$, hydrogen, halo, hydroxyl, or C_{1-4} alkyl; and

[0492] y is 1, 2, or 3.

[0493] In certain embodiments, the TPL is



[0494] wherein:

[0495] R^{1A} is C_{1-4} alkyl;


[0496] R^{1B} is C_{1-4} alkoxy;

[0497] R^{1C} is hydrogen, halo, or C_{1-4} alkyl; and


[0498] R^{1D} is hydroxyl.

[0499] In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is halo. In certain embodiments, R^{1C} is C_{1-4} alkyl.

[0500] In certain embodiments, the TPL is



[0501] In certain embodiments, the TPL has the following formula that is substituted by one occurrence of R^{II-1A} :



[0502] wherein:

[0503] R^{III-1A} is a bond to L;

[0504] R^{1A} is C_{1-4} alkyl;

[0505] R^{1B} is C_{1-4} alkoxy;

[0506] R^{1C} is hydrogen, halo, or C_{1-4} alkyl; and

[0507] R^{1D} is hydroxyl.

[0508] In certain embodiments, the TPL is selected from those depicted in the compounds in Tables 1, 2, 3, 4, and 5, below. In certain embodiments, the TPL is selected from those depicted in the compounds in Table 1, below. In certain embodiments, the TPL is selected from those depicted in the compounds in Table 2, below. In certain embodiments, the TPL is selected from those depicted in the compounds in Table 3, below. In certain embodiments, the TPL is selected from those depicted in the compounds in Table 4, below. In certain embodiments, the TPL is selected from those depicted in the compounds in Table 5, below.

Additional Features

[0509] Compounds of Formula I and/or II may be further characterized according to the molecular weight of the TPL. In certain embodiments, the TPL has a molecular weight of less than 1500 Da, 1200 Da, 1000 Da, 800 Da, 600 Da, 400 Da, 300 Da, 200 Da, 150 Da, or 100 Da. Compounds of Formula II may be further characterized according to the molecular weight of the EPL. In certain embodiments, the EPL has a molecular weight of less than 1500 Da, 1200 Da, 1000 Da, 800 Da, 600 Da, 400 Da, 300 Da, 200 Da, 150 Da, or 100 Da.

Part D: Exemplary Further Description of the Linker (L) Component of Compounds of Formula I and II

[0510] Compounds of Formula I and II may be further characterized according to, for example, the identity of the linker (L) component. A variety of linkers are known to one of skill in the art and may be used in the heterobifunctional compounds described herein. For example, in certain embodiments, L comprises one or more optionally substituted groups selected from amino acids, polyether chains, aliphatic groups, and any combinations thereof. In certain embodiments, L consists of one or more optionally substituted groups selected from amino acids, polyether chains, aliphatic groups, and any combinations thereof. In certain embodiments, L consists of one or more groups selected from amino acids, polyether chains, aliphatic groups, and any combinations thereof.

[0511] In some embodiments, L is symmetrical. In some embodiments, L is asymmetric. In certain embodiments, L is a bond.

[0512] In certain embodiments, L is a covalent bond or a bivalent C_{1-30} saturated or unsaturated, straight or branched, hydrocarbon chain, wherein 1-15 methylene units of L are optionally and independently replaced by cyclopropylene, $-N(H)-$, $-N(C_{1-4} \text{ alkyl})-$, $-N(C_{3-5} \text{ cycloalkyl})-$, $-O-$, $-C(O)-$, $-OC(O)-$, $-C(O)O-$, $-S-$, $-S(O)-$, $-S(O)_2-$, $-S(O)_2N(H)-$, $-S(O)_2N(C_{1-4} \text{ alkyl})-$, $-S(O)_2N(C_{3-5} \text{ cycloalkyl})-$, $-N(H)C(O)-$, $-N(C_{1-4} \text{ alkyl})C(O)-$, $-N(C_{3-5} \text{ cycloalkyl})C(O)-$, $-C(O)N(H)-$, $-C(O)N(C_{1-4} \text{ alkyl})-$, $-C(O)N(C_{3-5} \text{ cycloalkyl})-$, phenylene, an 8-10 membered bicyclic arylene, a 4-7 membered saturated or partially unsaturated carbocyclylene, an 8-10 membered bicyclic saturated or partially unsaturated carbocyclylene, a 3-7 membered saturated or partially unsaturated heterocyclylene having 1-2 heteroatoms independently selected from nitrogen, oxygen, and sulfur, an 8-10 membered bicyclic saturated or partially unsaturated heterocyclylene having 1-2 heteroatoms independently selected from nitrogen, oxygen, and sulfur, a 5-6 membered heteroarylene having 1-4 heteroatoms independently selected from nitrogen, oxygen, and sulfur, or an 8-10 membered bicyclic heteroarylene having 1-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur.

[0513] In certain embodiments, L is a bivalent, saturated or unsaturated, straight or branched C_{1-60} hydrocarbon chain, wherein 0-20 methylene units of the hydrocarbon are independently replaced with $-O-$, $-S-$, $-N(R^{**})-$, $-OC(O)-$, $-C(O)O-$, $-S(O)-$, $-S(O)_2-$, $-N(R^{**})S(O)_2-$, $-S(O)_2N(R^{**})-$, $-N(R^{**})C(O)-$, $-C(O)N(R^{**})-$, $-OC(O)N(R^{**})-$, $-N(R^{**})C(O)O-$, optionally substituted 3-10 membered carbocyclyl, or optionally substituted 3-10 membered heterocyclyl containing 1, 2, 3, or 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur, wherein R^{**} represents independently for each occurrence hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl.

[0514] In certain embodiments, L is a bivalent, saturated or unsaturated, straight or branched C_{1-60} hydrocarbon chain, wherein 0-20 methylene units of the hydrocarbon are independently replaced with $-O-$, $-S-$, $-N(H)-$, $-N(C_{1-6} \text{ alkyl})-$, $-OC(O)-$, $-C(O)O-$, $-S(O)-$, $-S(O)_2-$, $-N(H)S(O)_2-$, $-N(C_{1-6} \text{ alkyl})S(O)_2-$, $-S(O)_2N(H)-$, $-S(O)_2N(C_{1-6} \text{ alkyl})-$, $-N(H)C(O)-$, $-N(C_{1-6} \text{ alkyl})C(O)-$, $-C(O)N(H)-$, $-C(O)N(C_{1-6} \text{ alkyl})-$, $-OC(O)N(H)-$, $-OC(O)N(C_{1-6} \text{ alkyl})-$, $-N(H)C(O)O-$, $-N(C_{1-6} \text{ alkyl})C(O)O-$, optionally substituted

3-10 membered carbocyclyl, or optionally substituted 3-10 membered heterocyclyl containing 1, 2, 3, or 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

[0515] In certain embodiments, L is a bivalent, saturated, straight or branched C_{3-30} hydrocarbon chain, wherein 0-15 methylene units of the hydrocarbon are independently replaced with $-O-$, $-N(H)-$, $-N(C_{1-6} \text{ alkyl})-$, $-OC(O)-$, $-C(O)O-$, $-N(H)C(O)-$, $-N(C_{1-6} \text{ alkyl})C(O)-$, $-C(O)N(H)-$, $-C(O)N(C_{1-6} \text{ alkyl})-$, 3-10 membered carbocyclyl, or 3-10 membered heterocyclyl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

[0516] In certain embodiments, L is a bivalent, saturated, straight or branched C_{3-30} hydrocarbon chain, wherein 0-15 methylene units of the hydrocarbon are independently replaced with $-O-$, $-N(H)-$, $-N(C_{1-6} \text{ alkyl})-$, $-OC(O)-$, $-C(O)O-$, $-N(H)C(O)-$, $-N(C_{1-6} \text{ alkyl})C(O)-$, $-C(O)N(H)-$, or $-C(O)N(C_{1-6} \text{ alkyl})-$.

[0517] In certain embodiments, L is a bivalent, saturated or unsaturated, straight or branched C_{1-60} hydrocarbon chain, wherein 0-20 methylene units of the hydrocarbon are independently replaced with $-O-$, $-S-$, $-N(H)-$, $-N(C_{1-6} \text{ alkyl})-$, $-OC(O)-$, $-C(O)O-$, $-S(O)-$, $-S(O)_2-$, $-N(H)S(O)_2-$, $-N(C_{1-6} \text{ alkyl})S(O)_2-$, $-S(O)_2N(H)-$, $-S(O)_2N(C_{1-6} \text{ alkyl})-$, $-N(H)C(O)-$, $-N(C_{1-6} \text{ alkyl})C(O)-$, $-C(O)N(H)-$, $-C(O)N(C_{1-6} \text{ alkyl})-$, $-OC(O)N(H)-$, $-OC(O)N(C_{1-6} \text{ alkyl})-$, $-N(H)C(O)O-$, $-N(C_{1-6} \text{ alkyl})C(O)O-$, optionally substituted 3-10 membered carbocyclyl, optionally substituted 8-11 membered spirocyclic heterocyclyl containing 1, 2, 3, or 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur, or optionally substituted 3-10 membered heterocyclyl containing 1, 2, 3, or 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

[0518] In certain embodiments, L is a bivalent, saturated, straight or branched C_{3-30} hydrocarbon chain, wherein 0-15 methylene units of the hydrocarbon are independently replaced with $-O-$, $-N(H)-$, $-N(C_{1-6} \text{ alkyl})-$, $-OC(O)-$, $-C(O)O-$, $-N(H)C(O)-$, $-N(C_{1-6} \text{ alkyl})C(O)-$, $-C(O)N(H)-$, $-C(O)N(C_{1-6} \text{ alkyl})-$, 3-10 membered carbocyclyl, optionally substituted 8-11 membered spirocyclic heterocyclyl containing 1, 2, 3, or 4 heteroatoms independently selected from nitrogen, oxygen, and sulfur, or 3-10 membered heterocyclyl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur.

[0519] In yet other embodiments, L comprises a polyethylene glycol chain ranging in size from about 1 to about 12 ethylene glycol units, from about 1 to about 10 ethylene glycol units, from about 2 to about 6 ethylene glycol units, from about 2 to about 5 ethylene glycol units, or from about 2 to about 4 ethylene glycol units. In yet other embodiments, L is a diradical of a polyethylene glycol chain ranging in size from about 1 to about 12 ethylene glycol units, from about 1 to about 10 ethylene glycol units, from about 2 to about 6 ethylene glycol units, from about 2 to about 5 ethylene glycol units, or from about 2 to about 4 ethylene glycol units.

[0520] In certain embodiments, L is a heteroalkylene having from 4 to 30 atoms selected from carbon, oxygen, nitrogen, and sulfur. In certain embodiments, L is a heteroalkylene having from 4 to 20 atoms selected from carbon, oxygen, nitrogen, and sulfur. In certain embodiments, L is a heteroalkylene having from 4 to 10 atoms selected from carbon, oxygen, nitrogen, and sulfur. In certain embodi-

ments, L is a heteroalkylene having from 4 to 30 atoms selected from carbon, oxygen, and nitrogen. In certain embodiments, L is a heteroalkylene having from 4 to 20 atoms selected from carbon, oxygen, and nitrogen. In certain embodiments, L is a heteroalkylene having from 4 to 10 atoms selected from carbon, oxygen, and nitrogen. In certain embodiments, L is a heteroalkylene having from 4 to 30 atoms selected from carbon and oxygen. In certain embodiments, L is a heteroalkylene having from 4 to 20 atoms selected from carbon and oxygen. In certain embodiments, L is a heteroalkylene having from 4 to 10 atoms selected from carbon and oxygen.

[0521] In additional embodiments, the L is an optionally substituted (poly)ethyleneglycol having between 1 and about 100 ethylene glycol units, between about 1 and about 50 ethylene glycol units, between 1 and about 25 ethylene glycol units, between about 1 and about 10 ethylene glycol units, between 1 and about 8 ethylene glycol units, between 1 and about 6 ethylene glycol units, between 2 and about 4 ethylene glycol units, or optionally substituted alkyl groups interspersed with optionally substituted, O, N, S, P or Si atoms. In certain embodiments, L is substituted with an aryl, phenyl, benzyl, alkyl, alkylene, or heterocycle group.

[0522] In certain embodiments, L is a bivalent, saturated or unsaturated, straight or branched C_{1-45} hydrocarbon chain, wherein 0-10 methylene units of the hydrocarbon are independently replaced with $-O-$, $-S-$, $-N(R^{**})-$, $-OC(O)-$, $-C(O)O-$, $-S(O)-$, $-S(O)_2-$, $-N(R^{**})S(O)_2-$, $-S(O)_2N(R^{**})-$, $-N(R^{**})C(O)-$, $-C(O)N(R^{**})-$, $-OC(O)N(R^{**})-$, $-N(R^{**})C(O)O-$, optionally substituted carbocyclyl, or optionally substituted heterocyclyl, wherein R^{**} represents independently for each occurrence hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl.

[0523] In certain embodiments, L is a bivalent, saturated or unsaturated, straight or branched C_{1-45} hydrocarbon chain, wherein 0-10 methylene units of the hydrocarbon are independently replaced with $-O-$, $-S-$, $-N(R^{**})-$, $-OC(O)-$, $-C(O)O-$, $-S(O)-$, $-S(O)_2-$, $-N(R^{**})S(O)_2-$, $-S(O)_2N(R^{**})-$, $-N(R^{**})C(O)-$, $-C(O)N(R^{**})-$, $-OC(O)N(R^{**})-$, $-N(R^{**})C(O)O-$, optionally substituted 3-10 membered carbocyclyl, or optionally substituted 3-10 membered heterocyclyl containing 1, 2, 3, or 4 heteroatoms selected from nitrogen, oxygen, and sulfur, wherein R^{**} represents independently for each occurrence hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl.

[0524] In certain embodiments, L has the formula $-N(R)-(optionally substituted 3-20 membered heteroalkylene)p-CH_2-C(O)-$, wherein R is hydrogen or optionally substituted C_{1-6} alkyl, and p is 0 or 1.

[0525] In certain embodiments, L has the formula $-N(R)-(3-20 membered heteroalkylene)_p-CH_2-C(O)-$; wherein the 3-20 membered heteroalkylene is optionally substituted with 1, 2, 3, or 4 substituents independently selected from halogen, C_1-C_6 haloalkyl, C_3-C_6 cycloalkyl, hydroxyl, and cyano; R is hydrogen or optionally substituted C_{1-6} alkyl; and p is 0 or 1.

[0526] In certain embodiments, L has the formula $-N(R)-(3-20 membered heteroalkylene)_p-CH_2-C(O)-$; wherein the 3-20 membered heteroalkylene is optionally substituted with 1, 2, or 3 substituents independently selected from halogen and C_1-C_6 haloalkyl; R is hydrogen or C_{1-6} alkyl; and p is 0 or 1.

[0527] In certain embodiments, L is $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{10-20}$

$alkylene)-O-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, or $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, where *** is a point of attachment to TPL.

[0528] In certain embodiments, L is $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{10-20} alkylene)-O-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{2-6}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{7-15} alkylene)-C(O)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{2-6}-(C_{1-6} alkylene)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{2-9} alkylene)-O-(C_{1-6} alkylene)-C(O)N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{2-6}-(C_{1-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, or $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, where *** is a point of attachment to TPL.

[0529] In certain embodiments, L is $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{1-6} alkylene)-N(C_{1-6} alkyl)C(O)-(C_{1-6} alkylene)^{***}$, $-N(H)-(C_{1-6} alkylene)-N(H)C(O)-(C_{1-6} alkylene)^{***}$, $-N(H)-(C_{2-6} alkylene)-^{***}$, $-N(H)-(C_{7-15} alkylene)-^{***}$, $-N(C_{1-6} alkyl)-(C_{2-6} alkylene)-^{***}$, $-N(C_{1-6} alkyl)-(C_{7-15} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{2-6}-(C_{1-6} alkylene)-^{***}$, $-N(H)-[(C_{2-4} alkylene)-O-]_{7-15}-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{1-6} alkylene)-(3-6 membered heterocycloalkylene)-(C_{1-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{1-6} alkylene)-(3-6 membered heterocycloalkylene)-(C_{1-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{2-6} alkylene)-N(H)-(C_{1-6} alkylene)-^{***}$, or $-N(H)-(C_{2-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, where *** is a point of attachment to TPL.

[0530] In certain embodiments, L is $-N(H)-[CH_2CH_2-O-]_{2-6}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-C(O)-^{***}$, $-N(H)-(C_{1-6} alkylene)-N(C_{1-6} alkyl)C(O)-(C_{1-6} alkylene)^{***}$, $-N(H)-(C_{1-6} alkylene)-N(H)C(O)-(C_{1-6} alkylene)^{***}$, $-N(H)-(C_{2-6} alkylene)-^{***}$, $-N(H)-(C_{7-15} alkylene)-^{***}$, $-N(C_{1-6} alkyl)-(C_{2-6} alkylene)-^{***}$, $-N(C_{1-6} alkyl)-(C_{7-15} alkylene)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{2-6}-(C_{1-6} alkylene)-^{***}$, $-N(H)-[CH_2CH_2-O-]_{7-15}-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{1-6} alkylene)-(3-6 membered heterocycloalkylene)-(C_{1-6} alkylene)-N(C_{1-6} alkyl)-(C_{1-6} alkylene)-^{***}$, $-N(H)-(C_{1-6} alkylene)-(3-6 membered heterocycloalkylene)-(C_{1-6}$

alkylene)-N(H)-(C₁₋₆ alkylene)-***, —N(H)-(C₂₋₆ alkylene)-N(H)-(C₁₋₆ alkylene)-***, or —N(H)-(C₂₋₆ alkylene)-N(C₁₋₆ alkyl)-(C₁₋₆ alkylene)-***, where *** is a point of attachment to TPL.

[0531] In certain embodiments, L is $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(H)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(H)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[(C}_{2-4}\text{ alkylene)-O-]}_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, or $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[(C}_{2-4}\text{ alkylene)-O-]}_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, where *** is a point of attachment to TPL.


[0532] In certain embodiments, L is $-\text{[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(H)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(H)(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[CH}_2\text{CH}_2-\text{O}]_{2-6}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)-(C}_{1-6}\text{ alkylene)-}^{***}$, $-\text{(C}_{1-9}\text{ alkylene)-C(O)N(H)-[CH}_2\text{CH}_2-\text{O}]_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)-(C}_{1-6}\text{ alkylene)-}^{***}$, or $-\text{(C}_{1-9}\text{ alkylene)-N(H)C(O)-[(CH}_2\text{CH}_2-\text{O})_{7-15}-\text{(C}_{1-6}\text{ alkylene)-N(C}_{1-6}\text{ alkyl)-(C}_{1-6}\text{ alkylene)-}^{***}$, where *** is a point of attachment to TPL.



[0533] In certain embodiments, L is $-\text{N}(\text{H})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{2-6}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{H})-\text{***}$, $-\text{N}(\text{H})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{7-15}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{H})-\text{***}$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{2-6}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{H})-\text{***}$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{7-15}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{H})-\text{***}$, $-\text{N}(\text{C}_{1-6} \text{ alkyl})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{2-6}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{C}_{1-6} \text{ alkyl})-\text{***}$, or $-\text{N}(\text{C}_{1-6} \text{ alkyl})-[(\text{C}_{2-4} \text{ alkylene})-\text{O}]_{7-15}-(\text{C}_{1-6} \text{ alkylene})-\text{N}(\text{C}_{1-6} \text{ alkyl})-\text{***}$, where *** is a point of attachment to TPL.

[0534] In certain embodiments, L is —N(H)—[CH₂CH₂—O]₂₋₆—(C₁₋₆ alkylene)—N(H)—***, —N(H)—[CH₂CH₂—O]₇₋₁₅—(C₁₋₆ alkylene)—N(H)—***, —N(C₁₋₆ alkyl)—[CH₂CH₂—O]₂₋₆—(C₁₋₆ alkylene)—N(H)—***, —N(C₁₋₆ alkyl)—[CH₂CH₂—O]₇₋₁₅—(C₁₋₆ alkylene)—N


(H)-***, —N(C₁₋₆ alkyl)-[CH₂CH₂-O-]₂₋₆-(C₁₋₆ alkylene)-N(C₁₋₆ alkyl)-***, or —N(C₁₋₆ alkyl)-[CH₂CH₂-O-]₇₋₁₅-(C₁₋₆ alkylene)-N(C₁₋₆ alkyl)-***, where *** is a point of attachment to TPL.

[0535] In some embodiments, L is one of the following:







-continued




-continued




-continued




-continued






-continued





-continued












-continued




-continued




-continued



-continued





wherein a dashed bond indicates a point of attachment.

[0536] In certain embodiments, L has the formula $-(C_{0-12} \text{ alkylene})-(\text{optionally substituted 3-40 membered heteroalkylene})-(C_{0-12} \text{ alkylene})-$. In certain embodiments, L is C_{4-14} alkylene. In certain embodiments, L is $-(CH_2)_{6-10}-$.

[0537] In certain embodiments, L is $-CH_2CH_2(OCH_2CH_2)-^{***}$, $-CH_2CH_2(OCH_2CH_2)_2-^{***}$,

$-CH_2CH_2(OCH_2CH_2)_3-^{***}$, $-CH_2CH_2(OCH_2CH_2)_4-^{***}$, $-CH_2CH_2(OCH_2CH_2)_5-^{***}$, $-CH_2CH_2(OCH_2CH_2)_6-^{***}$, $-CH_2CH_2(OCH_2CH_2)_7-^{***}$, $-CH_2CH_2(OCH_2CH_2)_8-^{***}$, $-CH_2CH_2(OCH_2CH_2)_9-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{10}-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{11}-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{12}-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{13}-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{14}-^{***}$, $-CH_2CH_2(OCH_2CH_2)_{15}-^{***}$, or $-CH_2CH_2(OCH_2CH_2)_{16-20}-^{***}$, where *** is a point of attachment to TPL.

[0538] In certain embodiments, L is $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{2-4}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{5-7}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{8-10}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{11-13}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{14-16}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{2-20} \text{ alkylene})-(OCH_2CH_2)_{17-20}-(C_{0-4} \text{ alkylene})-^{***}$, $-(C_{1-20} \text{ alkylene})-(OCH_2CH_2)_{1-10}-(C_{0-4} \text{ alkylene})-C(O)-^{***}$, or $-(C_{1-20} \text{ alkylene})-(OCH_2CH_2)_{11-20}-(C_{0-4} \text{ alkylene})-C(O)-^{***}$, where *** is a point of attachment to TPL.

[0539] In certain embodiments, L is $-O(CH_2CH_2O)_{2-4}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{5-7}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{8-10}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{11-13}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{14-16}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{16-20}-(C_{0-4} \text{ alkylene})-^{***}$, $-O(CH_2CH_2O)_{2-10}-(C_{0-4} \text{ alkylene})C(O)-^{***}$, or $-O(CH_2CH_2O)_{11-20}-(C_{0-4} \text{ alkylene})C(O)-^{***}$, where *** is a point of attachment to TPL.

[0540] In certain embodiments, L is $-(C_{0-20} \text{ alkylene})-(OCH_2CH_2)_{1-10}-(N(C_{1-4} \text{ alkyl}))^{***}$, $-(C_{0-20} \text{ alkylene})-(OCH_2CH_2)_{11-20}-(N(C_{1-4} \text{ alkyl}))^{***}$, $-(C_{0-20} \text{ alkylene})-(CH_2CH_2O)_{1-10}-(C_{2-10} \text{ alkylene})-(N(C_{1-4} \text{ alkyl}))^{***}$, or $-(C_{0-20} \text{ alkylene})-(CH_2CH_2O)_{11-20}-(C_{2-10} \text{ alkylene})-(N(C_{1-4} \text{ alkyl}))^{***}$, where *** is a point of attachment to TPL.

[0541] In certain embodiments, L is $-(C_{2-10} \text{ alkylene})-(OCH_2CH_2)_{2-4}-O-(3-6 \text{ membered saturated heterocyclylene containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen})-(C_{1-5} \text{ alkylene})-^{***}$, $-(C_{2-10} \text{ alkylene})-(3-6 \text{ membered saturated heterocyclylene containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen})-(C_{1-5} \text{ alkylene})-^{***}$, $-(C_{2-10} \text{ alkylene})-N(H)-(C_{1-5} \text{ alkylene})-^{***}$, $-(C_{2-10} \text{ alkylene})-N(C_{1-6} \text{ alkyl})-(C_{1-5} \text{ alkylene})-^{***}$, $-N(H)-(C_{1-5} \text{ alkylene})-^{***}$, $-(CH_2CH_2O)_{1-4}-(C_{1-4} \text{ alkylene})-^{***}$, $-(CH_2CH_2O)_{1-4}-(C_{1-4} \text{ alkylene})-N(C_{1-6} \text{ alkyl})-(C_{1-6} \text{ alkylene})-^{***}$, $-N(H)-(C_{2-6} \text{ alkylene})-(3-6 \text{ membered saturated heterocyclylene containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen})-(C_{1-5} \text{ alkylene})-^{***}$, $-N(H)-(C_{2-6} \text{ alkylene})-(OCH_2CH_2)_{1-4}-O-(3-6 \text{ membered saturated heterocyclylene containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen})-(C_{1-5} \text{ alkylene})-^{***}$, $-N(H)-[CH_2CH_2O]_{2-6}-(C_{1-6} \text{ alkylene})-N(C_{1-6} \text{ alkyl})-(C_{1-6} \text{ alkylene})-^{***}$, $-N(H)-[CH_2CH_2O]_{2-10}-(C_{1-6} \text{ alkylene})-^{***}$,

$N(H)-***$, $-(C_{1-10} \text{ alkylene})-N(C_{1-4} \text{ alkyl})-(C_{1-6} \text{ alkylene})-O-***$, or $-(C_{1-15} \text{ alkylene})-[O-(C_{1-6} \text{ alkylene})]_{1-8}-N(C_{1-4} \text{ alkyl})-(C_{1-6} \text{ alkylene})-O-***$, where *** is a point of attachment to TPL. In certain embodiments, L is $-O-(C_{1-6} \text{ alkylene})-O-(C_{1-6} \text{ alkylene})-O-(C_{1-6} \text{ alkylene})-(4 \text{ membered saturated monocyclic heterocyclyl containing 1 nitrogen atom})-N(H)-***$, where *** is a point of attachment to TPL. In certain embodiments, L is $-(C_{1-10} \text{ alkylene})-N(C_{1-4} \text{ alkyl})-(C_{1-6} \text{ alkylene})-O-***$, where *** is a point of attachment to TPL. In certain embodiments, L is $-(C_{1-15} \text{ alkylene})-[O-(C_{1-6} \text{ alkylene})]_{1-8}-N(C_{1-4} \text{ alkyl})-(C_{1-6} \text{ alkylene})-O-***$, where *** is a point of attachment to TPL.

[0545] In certain embodiments, L is.

[0546] (i) $-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$

[0547] (ii) $-(7-11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$, or

[0548] (iii) $-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic ring containing 0, 1, or 2 nitrogen atoms, wherein the ring is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$; wherein *** is the point of attachment to TPL.

[0549] In certain embodiments, L is.

[0550] (i) $-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$, or


[0551] (ii) $-(7-11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$;

[0552] wherein *** is the point of attachment to TPL.


[0553] In certain embodiments, L is $-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$; wherein *** is the point of attachment to TPL. In certain embodiments, L is $-(7-11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$; wherein *** is the point of attachment to TPL. In certain embodiments, L is $-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro})-(C_{1-4} \text{ alkylene})-(4-6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$; wherein *** is the point of attachment to TPL.

atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic ring containing 0, 1, or 2 nitrogen atoms, wherein the ring is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***; wherein *** is the point of attachment to TPL. In certain embodiments, L is $-(\text{azetidinylen})-CH_2-(\text{piperazinylen})$ or $-(\text{piperidinylen})-CH_2-(\text{azetidinylen})$.


[0554] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:





-continued




-continued




[0555] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:



[0556] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:



[0557] In certain embodiments, L is one of the following:




[0558] wherein: Z^{L1} is C(OH), C(H), C(F), or N;

[0559] Z^{L2} is independently for each occurrence N or C(H); and

[0560] ** is the point of attachment to TPL.

[0561] In certain embodiments, L is one of the following:




[0562] wherein: Z^{L1} is C(OH), C(H), C(F), or N;


[0563] Z^{L2} is independently for each occurrence N or C(H); and

[0564] ** is the point of attachment to TPL.


[0565] In certain embodiments, L is




wherein *** is the point of attachment to TPL. In certain embodiments, L is




wherein *** is the point of attachment to TPL. In certain embodiments, L is




wherein *** is the point of attachment to TPL. In certain embodiments, L is



wherein *** is the point of attachment to TPL. In certain embodiments, L is



wherein *** is the point of attachment to TPL. In certain embodiments, L is









wherein *** is the point of attachment to TPL.

[0566] In certain embodiments, Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H). In certain embodiments, Z^{L1} is C(OH), and Z^{L2} is N. In certain embodiments, Z^{L1} is C(OH), C(H), or


C(F). In certain embodiments, Z^{L1} is C(OH) or C(H). In certain embodiments, Z^{L1} is C(OH). In certain embodiments, Z^{L1} is C(H). In certain embodiments, Z^{L1} is C(F). In certain embodiments, Z^{L1} is N. In certain embodiments, Z^{L2} is independently for each occurrence N or C(H). In certain embodiments, Z^{L2} is N or C(H). In certain embodiments, Z^{L2} is N. In certain embodiments, Z^{L2} is C(H).

[0567] In certain embodiments, L is one of the following, wherein *** is the point of attachment to TPL:







[0568] In certain embodiments, L is




wherein *** is the point of attachment to TPL. In certain embodiments, L is




wherein *** is the point of attachment to TPL. In certain embodiments, L is one of the following, wherein *** is the point of attachment to TPL:





In certain embodiments, L is



wherein *** is the point of attachment to TPL.

[0569] In certain embodiments, L is selected from those depicted in the compounds in Table 1, 3, 4, and 5, below. In certain embodiments, L is selected from those depicted in the compounds in Table 1, below. In certain embodiments, L is selected from those depicted in the compounds in Table 2, below. In certain embodiments, L is selected from those depicted in the compounds in Table 3, below. In certain embodiments, L is selected from those depicted in the compounds in Table 4, below. In certain embodiments, L is selected from those depicted in the compounds in Table 5, below.


Part E: Exemplary More Specific Embodiments

[0570] Embodiments described above may be combined to provide more specific embodiments defining compounds. All combinations and permutations are contemplated.

[0571] For example, in certain embodiments, the compound of Formula I is a compound of Formula I-A or I-B, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A or I-B. In certain embodiments, the compound of each of Formula I-A and I-B is as defined and described in the following embodiments.

[0572] In certain embodiments, the compound of Formula I is represented by Formula I-A:

(I-A)



[0573] or a pharmaceutically acceptable salt thereof, wherein:

[0574] R^3 , R^{1C} , and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl;

[0575] R^4 is methyl, $-\text{CH}_2\text{-(oxazolyl)}$, or $-\text{CH}_2\text{-(thiazolyl)}$;

[0576] L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :

[0577] (i) $-(4\text{-}6\text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4}\text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, or}$

[0578] (ii) $-(7\text{-}11\text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4}\text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, and}$

[0579] n, q, and r are independently 0, 1, or 2.

[0580] In certain embodiments, the compound is a compound of Formula I-A.


[0581] In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^3 represents independently for each occurrence hydrogen or fluoro. In certain embodiments, R^3 represents independently for each occurrence fluoro or chloro. In certain embodiments, R^3 is hydrogen. In certain embodiments, R^3 is fluoro. In certain embodiments, R^3 is chloro. In certain embodiments, R^3 is methyl. In certain embodiments, R^3 is methoxy.

[0582] In certain embodiments, R^4 is methyl. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazolyl)}$ or $-\text{CH}_2\text{-(thiazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazol-2-yl)}$ or $-\text{CH}_2\text{-(thiazol-2-yl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazol-2-yl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(thiazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(thiazol-2-yl)}$.

[0583] In certain embodiments, L is $-(4\text{-}6\text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4}\text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, wherein *** is the point of attachment to the phenyl ring bearing } R^{1C}$. In certain embodiments, L is $-(7\text{-}11\text{ membered saturated spirocyclic heterocyclyl contain-}$

ing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***; wherein *** is the point of attachment to the phenyl ring bearing R^{1C}

[0584] In certain embodiments, L is one of the following:




[0585] wherein: Z^{L1} is C(OH), C(H), C(F), or N;


[0586] Z^{L2} is independently for each occurrence N or C(H); and

[0587] ** is the point of attachment to the phenyl ring bearing R^{1C}.


[0588] In certain embodiments, L is




wherein *** is the point of attachment to the phenyl ring bearing R^{1C}. In certain embodiments, L is




wherein *** is the point of attachment to the phenyl ring bearing R^{1C}. In certain embodiments, L is




-continued



wherein *** is the point of attachment to the phenyl ring bearing R^{1C}. In certain embodiments, L is




wherein *** is the point of attachment to the phenyl ring bearing R^{1C}. In certain embodiments, L is





wherein *** is the point of attachment to the phenyl ring bearing R^{1C}.













[0589] In certain embodiments, Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H). In certain embodiments, Z^{L1} is C(OH), and Z^{L2} is N. In certain embodiments, Z^{L1} is C(OH), C(H), or C(F). In certain embodiments, Z^{L1} is C(OH) or C(H). In certain embodiments, Z^{L1} is C(OH). In certain embodiments, Z^{L1} is C(H). In certain embodiments, Z^{L1} is C(F). In certain embodiments, Z^{L1} is N. In certain embodiments, Z^{L2} is independently for each occurrence N or C(H). In certain embodiments, Z^{L2} is N or C(H). In certain embodiments, Z^{L2} is N. In certain embodiments, Z^{L2} is C(H).

[0590] In certain embodiments, L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C}:








[0591] In certain embodiments, L is



wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is



wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :



[0592] In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence fluoro, chloro, methoxy, or methyl. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} and R^{1E} are each independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1C} and R^{1E} are hydrogen. In certain embodiments, R^{1C} and R^{1E} are fluoro.

[0593] In certain embodiments, R^{1C} is independently for each occurrence fluoro, chloro, methoxy, or methyl. In

certain embodiments, R^{1C} is independently for each occurrence hydrogen, fluoro, chloro, or methoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or methoxy. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methoxy. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methyl. In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is fluoro. In certain embodiments, R^{1C} is chloro. In certain embodiments, R^{1C} is methyl. In certain embodiments, R^{1C} is methoxy.

[0594] In certain embodiments, R^{1E} is independently for each occurrence fluoro, chloro, methoxy, or methyl. In certain embodiments, R^{1E} is independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1E} is independently for each occurrence hydrogen or fluoro. In certain embodiments, R^{1E} is hydrogen or fluoro. In certain embodiments, R^{1C} is independently for each occurrence hydrogen or methyl. In certain embodiments, R^{1C} is independently for each occurrence fluoro or methyl. In certain embodiments, R^{1E} is hydrogen. In certain embodiments, R^{1E} is fluoro. In certain embodiments, R^{1E} is chloro. In certain embodiments, R^{1E} is methyl. In certain embodiments, R^{1C} is methoxy.

[0595] In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2.

[0596] In certain embodiments, q and r are 0. In certain embodiments, q and r are independently 0 or 1. In certain embodiments, q and r are independently 1 or 2. In certain embodiments, q and r are independently 0, 1, or 2. In certain embodiments, q is 0. In certain embodiments, q is 1. In certain embodiments, q is 2. In certain embodiments, q is 3. In certain embodiments, q is 0 or 1. In certain embodiments, q is 1 or 2. In certain embodiments, q is 0, 1, or 2. In certain embodiments, r is 0. In certain embodiments, r is 1. In certain embodiments, r is 2. In certain embodiments, r is 3. In certain embodiments, r is 0 or 1. In certain embodiments, r is 1 or 2. In certain embodiments, r is 0, 1, or 2.


[0597] In certain embodiments:


[0598] R^3 represents independently for each occurrence hydrogen, fluoro, or chloro;

[0599] R^4 is methyl or $-\text{CH}_2-(\text{oxazolyl})$;

[0600] R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy; and

[0601] L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} .





[0602] wherein: Z^{L1} is C(OH), C(H), C(F), or N;

[0603] Z^{L2} is independently for each occurrence N or C(H); and

[0604] $**$ is the point of attachment to the phenyl ring bearing R^{1C}


[0605] In certain embodiments:

[0606] R^3 represents independently for each occurrence hydrogen or fluoro;

[0607] R^4 is methyl;


[0608] R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy; and

[0609] L is




wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} , Z^{L1} is C(OH), C(H), C(F), or N, and Z^{L2} is independently for each occurrence N or C(H).

[0610] In certain embodiments, L is




wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} , Z^{L1} is C(OH), C(H), C(F), or N, and Z^{L2} is N or C(H). In certain embodiments, L is



wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} , Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

[0611] In certain embodiments, R^4 is methyl, and L is




[0612] In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^3 represents independently for each occurrence hydrogen and fluoro, and R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy.


[0613] The description above describes multiple embodiments relating to compounds of Formula I-A. The patent application specifically contemplates all combinations of the embodiments.

[0614] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-1 or I-A-2, or a pharmaceutically acceptable salt thereof.

(I-A-1)



(I-A-2)



[0615] In certain embodiments, the compound is a compound of Formula I-A-1, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A-2, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A-1 or I-A-2. In certain embodiments, the compound is a compound of Formula I-A-1. In certain embodiments, the compound is a compound of Formula I-A-2.


[0616] In certain embodiments, each of variables R^3 , R^4 , L , R^{1C} , R^{1E} , n , q , and r in Formula I-A-1 and I-A-2 is as defined and described in embodiments above for compounds of Formula I-A, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-A-1 and I-A-2. The patent application specifically contemplates all combinations of the embodiments.

[0617] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-3* or I-A-4*, or a pharmaceutically acceptable salt thereof:


embodiments, the compound is a compound of Formula I-A-3*. In certain embodiments, the compound is a compound of Formula I-A-4*.

[0619] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-3 or I-A-4, or a pharmaceutically acceptable salt thereof:


(I-A-3)




(I-A-3*)



(I-A-4)




(I-A-4*)





[0620] In certain embodiments, the compound is a compound of Formula I-A-3, or a pharmaceutically acceptable salt thereof:

(I-A-3)



[0618] In certain embodiments, the compound is a compound of Formula I-A-3*, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A-4*, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A-3* or I-A-4*. In certain




[0621] In certain embodiments, the compound is a compound of Formula I-A-4, or a pharmaceutically acceptable salt thereof:



[0622] In certain embodiments, the compound is a compound of Formula I-A-3 or I-A-4. In certain embodiments, the compound is a compound of Formula I-A-3. In certain embodiments, the compound is a compound of Formula I-A-4.

[0623] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-3** or I-A-4**, or a pharmaceutically acceptable salt thereof:

[0625] In certain embodiments, the compound is a compound of Formula I-A-4**, or a pharmaceutically acceptable salt thereof:




[0626] In certain embodiments, the compound is a compound of Formula I-A-3** or I-A-4**. In certain embodiments, the compound is a compound of Formula I-A-3**. In certain embodiments, the compound is a compound of Formula I-A-4**.

[0627] In certain embodiments, each of variables R^3 , R^4 , L , R^{1C} , and R^{1E} in Formula I-A-3*, I-A-4*, I-A-3, I-A-4, I-A-3** and I-A-4** is as defined and described in embodiments above for compounds of Formula I-A, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-A-3*, I-A-4*, I-A-3, I-A-4, I-A-3** and I-A-4**. The patent application specifically contemplates all combinations of the embodiments.

[0624] In certain embodiments, the compound is a compound of Formula I-A-3**, or a pharmaceutically acceptable salt thereof:

[0628] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-5, or a pharmaceutically acceptable salt thereof.





[0629] wherein Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

[0630] In certain embodiments, the compound is a compound of Formula I-A-5. In certain embodiments, Z^{L1} is C(OH). In certain embodiments, Z^{L1} is C(H). In certain embodiments, Z^{L2} is N. In certain embodiments, Z^{L2} is C(H). In certain embodiments, Z^{L1} is C(OH), and Z^{L2} is N.

[0631] In certain embodiments, each of variables R³, R^{1C}, and R^{1E} in Formula I-A-5 is as defined and described in


embodiments above for compounds of Formula I-A, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-A-5. The patent application specifically contemplates all combinations of the embodiments.

[0632] In certain embodiments, the compound of Formula I or I-A is represented by Formula I-A-6 or I-A-7, or a pharmaceutically acceptable salt thereof.




[0633] wherein Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

[0634] In certain embodiments, the compound is a compound of Formula I-A-6, or a pharmaceutically acceptable salt thereof:



[0635] wherein Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

[0636] In certain embodiments, the compound is a compound of Formula I-A-7, or a pharmaceutically acceptable salt thereof:



[0637] wherein Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

[0638] In certain embodiments, the compound is a compound of Formula I-A-6 or I-A-7. In certain embodiments, the compound is a compound of Formula I-A-6. In certain embodiments, the compound is a compound of Formula I-A-7.

[0639] In certain embodiments, Z^{L1} is C(OH). In certain embodiments, Z^{L1} is C(H). In certain embodiments, Z^{L2} is N. In certain embodiments, Z^{L2} is C(H). In certain embodiments, Z^{L1} is C(OH), and Z^{L2} is N.


[0640] In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^3 represents independently for each occurrence hydrogen and

fluoro, and R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy.

[0641] In certain embodiments, each of variables R^3 , R^{1C} , and R^{1E} in Formula I-A-6 and I-A-7 is as defined and described in embodiments above for compounds of Formula

I-A, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-A-6 and I-A-7. The patent application specifically contemplates all combinations of the embodiments.

[0642] In certain embodiments, the compound of Formula I is represented by Formula I-B:



- [0643] or a pharmaceutically acceptable salt thereof, wherein:
- [0644] R^3 represents independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl;
- [0645] R^4 is methyl, $-\text{CH}_2\text{-(oxazolyl)}$, or $-\text{CH}_2\text{-(thiazolyl)}$;
- [0646] n is 0, 1, or 2;
- [0647] L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :
- [0648] (i) $-(4\text{-}6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$,
- [0649] (ii) $-(7\text{-}11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$, or
- [0650] (iii) $-(4\text{-}6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic ring containing 0, 1, or 2 nitrogen atoms, wherein the ring is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$,
- [0651] R^{1A} and R^{1C} each represents independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl;
- [0652] R^{1B} is $-\text{C(O)OH}$, $-\text{OH}$, or $-\text{B(OH)}_2$;
- [0653] q is 1 or 2; and
- [0654] y is 1, 2, or 3.
- [0655] In certain embodiments, the compound is a compound of Formula I-B.
- [0656] In certain embodiments, R^3 represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^3 represents independently for each occurrence hydrogen or fluoro. In certain embodiments, R^3 represents independently for each occurrence fluoro or chloro. In certain embodiments, R^3 is hydrogen. In certain embodiments, R^3 is fluoro. In certain embodiments, R^3 is chloro. In certain embodiments, R^3 is methyl. In certain embodiments, R^3 is methoxy.
- [0657] In certain embodiments, R^4 is methyl. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazolyl)}$ or $-\text{CH}_2\text{-(thiazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazol-2-yl)}$ or $-\text{CH}_2\text{-(thiazol-2-yl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(oxazol-2-yl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(thiazolyl)}$. In certain embodiments, R^4 is $-\text{CH}_2\text{-(thiazol-2-yl)}$.
- [0658] In certain embodiments, n is 0. In certain embodiments, n is 1. In certain embodiments, n is 2.
- [0659] In certain embodiments, L is.
- [0660] (i) $-(4\text{-}6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of$


hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, or


- [0661] (ii) $-(7\text{-}11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$;

[0662] wherein *** is the point of attachment to the phenyl ring bearing R^{1C}

[0663] In certain embodiments, L is $-(4\text{-}6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$; wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is $-(7\text{-}11 \text{ membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$; wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is $-(4\text{-}6 \text{ membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic ring containing 0, 1, or 2 nitrogen atoms, wherein the ring is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C}_{1-4} \text{ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)}\text{-***}$; wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is $-(\text{azetidinylen})\text{-CH}_2\text{-(piperazinylen or piperidinylen)}\text{-CH}_2\text{-(azetidinylen)}$.

[0664] In certain embodiments, L is one of the following:






[0665] wherein: Z^{L1} is C(OH), C(H), C(F), or N;

[0666] Z^{L2} is independently for each occurrence N or C(H); and

[0667] $**$ is the point of attachment to the phenyl ring bearing R^{1C}

[0668] In certain embodiments, L is one of the following:




[0669] wherein: Z^{L1} is C(OH), C(H), C(F), or N;


[0670] Z^{L2} is independently for each occurrence N or C(H); and

[0671] $**$ is the point of attachment to the phenyl ring bearing R^{1C} .


[0672] In certain embodiments, L is




wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is




wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is




wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is



wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is



wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is











wherein $***$ is the point of attachment to the phenyl ring bearing R^{1C}






[0673] In certain embodiments, Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H). In certain embodiments, Z^{L1} is C(OH), and

Z^{L2} is N. In certain embodiments, Z^{L1} is C(OH), C(H), or C(F). In certain embodiments, Z^{L1} is C(OH) or C(H). In certain embodiments, Z^{L1} is C(OH). In certain embodiments, Z^{L1} is C(H). In certain embodiments, Z^{L1} is C(F). In certain embodiments, Z^{L1} is N. In certain embodiments, Z^{L2} is independently for each occurrence N or C(H). In certain embodiments, Z^{L2} is N or C(H). In certain embodiments, Z^{L2} is N. In certain embodiments, Z^{L2} is C(H).





[0674] In certain embodiments, L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :




-continued





[0675] In certain embodiments, L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :







[0676] In certain embodiments, L is




wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is




wherein *** is the point of attachment to the phenyl ring bearing R^{1C} . In certain embodiments, L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :





In certain embodiments, L is



wherein *** is the point of attachment to the phenyl ring bearing R^{1C} .

[0677] In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro, and R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1A} represents independently for each occurrence hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen, fluoro, or chloro. In certain embodiments, R^{1A} is hydrogen or chloro. In certain embodiments, R^{1A} is hydrogen. In certain embodiments, R^{1A} is fluoro. In certain embodiments, R^{1A} is chloro. In certain embodiments, R^{1A} is methyl. In certain embodiments, R^{1A} is methoxy.

[0678] In certain embodiments, R^{1B} is $-\text{C}(\text{O})\text{OH}$ or $-\text{OH}$. In certain embodiments, R^B is $-\text{C}(\text{O})\text{OH}$. In certain embodiments, R^{1B} is hydroxyl. In certain embodiments, R^{1B} is $-\text{B}(\text{OH})_2$. In certain embodiments, R^{1C} represents independently for each occurrence hydrogen, fluoro, or methoxy. In certain embodiments, R^{1C} is hydrogen or fluoro. In certain


embodiments, R^{1C} is fluoro or chloro. In certain embodiments, R^{1C} is hydrogen. In certain embodiments, R^{1C} is fluoro. In certain embodiments, R^{1C} is chloro. In certain embodiments, R^{1C} is methoxy. In certain embodiments, R^{1C} is methyl.

[0679] In certain embodiments, q is 1. In certain embodiments, q is 2. In certain embodiments, y is 1. In certain embodiments, y is 2. In certain embodiments, y is 3.

[0680] The description above describes multiple embodiments relating to compounds of Formula I-B. The patent application specifically contemplates all combinations of the embodiments.

[0681] In certain embodiments, the compound of Formula I or Formula I-B is represented by Formula I-B-1, or a pharmaceutically acceptable salt thereof.


(I-B-1)



[0682] In certain embodiments, the compound is a compound of Formula I-B-1. In certain embodiments, each of variables R^3 , R^4 , L, R^{1A} , R^{1B} , R^{1C} , n, q, and y in Formula I-B-1 is as defined and described in embodiments above for compounds of Formula I-B, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-B-1. The patent application specifically contemplates all combinations of the embodiments.


[0683] In certain embodiments, the compound of Formula I or Formula I-B is represented by Formula I-B-2, or a pharmaceutically acceptable salt thereof.

(I-B-2)




[0684] In certain embodiments, the compound of Formula I or Formula I-B is represented by Formula I-B-3, or a pharmaceutically acceptable salt thereof:

(I-B-3)



[0685] In certain embodiments, the compound of Formula I or Formula I-B is represented by Formula I-B-4, or a pharmaceutically acceptable salt thereof:

(I-B-4)




[0686] In certain embodiments, the compound is a compound of Formula I-B-2. In certain embodiments, the compound is a compound of Formula I-B-3. In certain embodiments, the compound is a compound of Formula I-B-4.

[0687] In certain embodiments, each of variables R^3 , R^4 , L , R^{1A} , and R^{1C} in Formula I-B-2, I-B-3, and I-B-4 is as defined and described in embodiments above for compounds of Formula I-B, both singly and in combination. The description above describes multiple embodiments relating to compounds of Formula I-B-2, I-B-3, and I-B-4. The patent application specifically contemplates all combinations of the embodiments.


[0688] Another aspect of the invention provides a compound of Formula I-1:

(I-1)



[0689] or a pharmaceutically acceptable salt thereof, wherein:

[0690] EPL has the following formula:



[0691] wherein:

[0692] R^1 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0693] R^2 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0694] R^3 is hydrogen, halo, C_{1-4} alkyl, C_{1-4} haloalkyl, or C_{1-4} alkoxy;


[0695] R^4 is $-(C_{0-6}$ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), $-(C_{1-6}$ alkylene)- $C(O)N(R^5)(R^6)$, $-(C_{1-6}$ alkylene)- $N(R^5)C(O)R^7$, $-(C_{1-6}$ alkylene)- CO_2R^7 , $-(C_{1-6}$ alkylene)- $OC(O)R^7$, $-(C_{1-6}$ alkylene)-cyano, $-(C_{1-6}$ alkylene)- $O-(C_{1-6}$ alkyl), C_{1-6} alkyl, C_{3-6} cycloalkyl, or hydrogen; or R^4 and R^8 taken together with the carbon atom to which they are attached form a C_{3-5} saturated carbocyclic ring;

[0696] R^5 and R^6 are independently hydrogen, C_{1-6} alkyl, or C_{3-6} cycloalkyl; or R^5 and R^6 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;

[0697] R^7 is C_{1-6} alkyl, $-(C_{1-6}$ alkylene)-(C_{3-6} cycloalkyl), or C_{3-6} cycloalkyl;

[0698] R^8 is hydrogen or C_{1-4} alkyl; and

[0699] m is 0, 1, or 2;



[0700] wherein:

[0701] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0702] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy; and

[0703] L is one of the following:



[0704] (i) $-(5-6$ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, $-(4-6$ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C_{0-6} alkylene)-($5-6$ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***,



-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of hydroxyl)-(C₀₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 heteroatoms selected from nitrogen)-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₁₋₆ alkylene)-O-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-O-***, where *** is a point of attachment to TPL;

[0705] (ii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -N(H)-[O-(C₁₋₆ alkylene)-O]₁₋₆-(C₁₋₆ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, -N(H)-(C₁₋₁₀ alkylene)-O-***, -N(H)-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of fluoro)-(C₀₋₆ alkylene)-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, where *** is a point of attachment to TPL; or




[0706] (iii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, -(C₁₋₁₀ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, or -(C₁₋₁₅ alkylene)-[O-(C₁₋₆ alkylene)-]₁₋₈-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, where *** is a point of attachment to TPL.

[0707] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:







[0708] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:



[0709] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:






[0710] Another aspect of the invention provides a compound of Formula I-2:

[0716] m is 0, 1, or 2; and


[0717] n is 0, 1, 2, 3, or 4;

[0718] TPL is



[0711] or a pharmaceutically acceptable salt thereof, wherein:


[0712] EPL has the following formula:



[0713] wherein:

[0714] R^1 is phenyl or 6-membered heteroaryl containing 1 or 2 heteroatoms independently selected from nitrogen, oxygen, and sulfur; wherein the phenyl and heteroaryl are optionally substituted with 1, 2, or 3 substituents independently selected from halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, hydroxyl, or C_1 - C_6 alkoxy;

[0715] R^2 and R^3 each represent independently for each occurrence halo, C_1 - C_6 alkyl, C_1 - C_6 haloalkyl, C_3 - C_6 cycloalkyl, hydroxyl, C_1 - C_6 alkoxy, or cyano;



[0719] wherein:

[0720] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0721] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy; and L is one of the following:



[0722] (i) -(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, -(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of hydroxyl)-(C₀₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 heteroatoms selected from nitrogen)-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-


(C₀₋₆ alkylene)-O-***, —O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, —(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, —(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, —O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, —O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₁₋₆ alkylene)-O-***, or —O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-O-***, where *** is a point of attachment to TPL;

[0723] (ii) —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)—(C₀₋₆ alkylene)-***, —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—***, —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)-***, —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—***, —N(H)—[(C₁₋₆ alkylene)-O]₁₋₆—(C₁₋₆ alkylene)-N(C₁₋₄ alkyl)—(C₁₋₆ alkylene)—O—***, —N(H)—(C₁₋₁₀ alkylene)—O—***, —N(H)—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—***, —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—(4 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of fluoro)—(C₀₋₆ alkylene)-***, or —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)—N(H)—***, where *** is a point of attachment to TPL; or



[0724] (iii) —O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—O—(C₁₋₆ alkylene)—(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)—N(H)—***, —(C₁₋₁₀ alkylene)—N(C₁₋₄ alkyl)—(C₁₋₆ alkylene)—O—***, or —(C₁₋₁₅ alkylene)—[—O—(C₁₋₆ alkylene)—]₁₋₈—N(C₁₋₄ alkyl)—(C₁₋₆ alkylene)—O—***, where *** is a point of attachment to TPL.

[0725] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:







[0726] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:




[0727] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:






[0728] Another aspect of the invention provides a compound of Formula I-3:



[0729] or a pharmaceutically acceptable salt thereof, wherein:

[0730] EPL has the following formula:



[0731] wherein:

[0732] R^1 represents independently for each occurrence halo, C_{1-4} alkyl, or C_{1-4} haloalkyl;

[0733] R^2 is hydrogen, halo, C_{1-4} alkyl, or C_{1-4} haloalkyl;

[0734] R^3 is $-N(R^8)SO_2R^9$, $-SO_2N(R^8)_2$, $-SO_2R^9$, $-(C_{1-6} \text{ alkylene})-SO_2R^9$, C_{1-6} hydroxyalkyl, or a 4-7 membered saturated carbocyclic ring in which one CH_2 is replaced with SO_2 ;

[0735] R^4 is hydrogen, halo, or C_{1-4} alkyl;

[0736] R^5 is C_{1-4} alkyl or C_{3-4} cycloalkyl;

[0737] R^6 is hydrogen, C_{1-4} alkyl, or C_{3-4} cycloalkyl;

[0738] R^7 represents independently for each occurrence C_{1-4} alkyl or C_{3-4} cycloalkyl;


[0739] R^8 represents independently for each occurrence hydrogen or C_{1-4} alkyl; or two occurrences of R^8 are taken together with the nitrogen atom to which they are attached to form a 3-7 membered heterocyclic ring containing 1 or 2 heteroatoms independently selected from nitrogen and oxygen; or R^8 and R^9 are taken together with their intervening atoms to form a 5-7 membered ring containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur;

[0740] R^9 is C_{1-6} alkyl, C_{1-6} haloalkyl, $-(C_{1-6} \text{ alkylene})-(C_{3-6} \text{ cycloalkyl})$, or C_{3-6} cycloalkyl;

[0741] A^3 is phenylene, a 3-10 membered saturated monocyclic, bicyclic or spirocyclic carbocyclene, or C_{1-6} alkylene; and

[0742] p and t are independently 0, 1, or 2;

[0743] TPL is



[0744] wherein:

[0745] R^{1A} is hydrogen, halo, or C_{1-4} alkyl; and

[0746] R^{1B} and R^{1C} are independently hydroxyl, halo, C_{1-4} alkyl, or C_{1-4} alkoxy; and L is one of the following:



[0747] (i) $-(5-6 \text{ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$, $-(4-6 \text{ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-(C_{0-6} \text{ alkylene})-(5-6 \text{ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$, $-(4-6 \text{ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$, or $-(4-6 \text{ membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms})-***$.



nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of hydroxyl)-(C₀₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 heteroatoms selected from nitrogen)-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₁₋₆ alkylene)-O-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-O-***, where *** is a point of attachment to TPL;

[0748] (ii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -N(H)-[-(C₁₋₆ alkylene)-O-]₁₋₆-(C₁₋₆ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, -N(H)-(C₁₋₁₀ alkylene)-O-***, -N(H)-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of fluoro)-(C₀₋₆ alkylene)-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, where *** is a point of attachment to TPL; or

[0749] (iii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, -(C₁₋₁₀ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, or -(C₁₋₁₅ alkylene)-[-O-(C₁₋₆ alkylene)-]₁₋₈-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, where *** is a point of attachment to TPL.




[0750] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:







[0751] In certain embodiments, L is one of the following, where *** is a point of attachment

to TPL:




[0752] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:






[0753] Another aspect of the invention provides a compound of Formula I-4:



[0754] or a pharmaceutically acceptable salt thereof, wherein:

[0755] EPL has the following formula:



[0756] wherein:

[0757] R¹ is C₁₋₄ alkyl or C₃₋₄ cycloalkyl;

[0758] R² represents independently for each occurrence C₁₋₄ alkyl or C₃₋₄ cycloalkyl;

[0759] R³ is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ haloalkyl, or C₁₋₄ alkoxy;

[0760] R⁴ is —(C₀₋₆ alkylene)-(5-6 membered heteroaryl containing 1, 2, or 3 heteroatoms independently selected from nitrogen, oxygen, and sulfur), —(C₁₋₆

alkylene)-C(O)N(R⁵)(R⁶) —(C₁₋₆ alkylene)-N(R⁵)C(O)R⁷, —(C₁₋₆ alkylene)-CO₂R⁷, —(C₁₋₆ alkylene)-OC(O)R⁷, —(C₁₋₆ alkylene)-cyano, —(C₁₋₆ alkylene)-O—(C₁₋₆ alkyl), C₁₋₆ alkyl, C₃₋₆ cycloalkyl, or hydrogen; or R⁴ and R⁸ taken together with the carbon atom to which they are attached form a C₃₋₅ saturated carbocyclic ring;


[0761] R⁵ and R⁶ are independently hydrogen, C₁₋₆ alkyl, or C₃₋₆ cycloalkyl; or R⁵ and R⁶ are taken together with the nitrogen atom to which they are attached to form a 3-7 membered ring containing 1 nitrogen atom;

[0762] R⁷ is C₁₋₆ alkyl, —(C₁₋₆ alkylene)-(C₃₋₆ cycloalkyl), or C₃₋₆ cycloalkyl;

[0763] R⁸ is hydrogen or C₁₋₄ alkyl; and

[0764] m is 0, 1, or 2;

[0765] TPL is



[0766] wherein:

[0767] R^{1C} represents independently for each occurrence hydrogen, halo, hydroxyl, or C₁₋₄ alkyl;

[0768] R^{1D} is hydroxyl;

[0769] R^{1E} represents independently for each occurrence halo, hydroxyl, or C₁₋₄ alkyl;

[0770] R^{1F} is hydrogen or C₁₋₄ alkyl;

[0771] y is 1 or 2; and

[0772] z is 1, 2, or 3; and


[0773] L is one of the following:

[0774] (i) -(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***, -(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of hydroxyl)-(C₀₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -(4-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₀₋₆ alkylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 heteroatoms selected from nitrogen)-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -O-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-(C₀₋₆ alkylene)-O-***, -(C₂₋₃ alkynylene)-(7-11 membered spirocyclic heterocyclyl containing 1-2 nitrogen atoms)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-***, -O-(C₀₋₆ alkylene)-(5-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₁₋₆ alkylene)-O-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-O-***, where *** is a point of attachment to TPL;


[0775] (ii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(4-6 membered monocyclic heterocyclyl containing 1 nitrogen atom)-(C₀₋₆ alkylene)-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -N(H)-[O-(C₁₋₆ alkylene)-O]₁₋₆-(C₁₋₆ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, -N(H)-(C₁₋₁₀ alkylene)-O-***, -N(H)-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-***, -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom, wherein the heterocyclyl is substituted with 0 or 1 occurrences of fluoro)-(C₀₋₆ alkylene)-***, or -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, where *** is a point of attachment to TPL; or

[0776] (iii) -O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-O-(C₁₋₆ alkylene)-(4 membered monocyclic heterocyclyl containing 1 nitrogen atom)-N(H)-***, -(C₁₋₁₀ alkylene)-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, or -(C₁₋₁₅ alkylene)-[O-(C₁₋₆ alkylene)-]₁₋₈-N(C₁₋₄ alkyl)-(C₁₋₆ alkylene)-O-***, where *** is a point of attachment to TPL.











[0777] In certain embodiments, L is one of the following, where *** is a point of attachment





-continued



-continued




[0778] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:



[0779] In certain embodiments, L is one of the following, where *** is a point of attachment to TPL:

6, or I-A-7, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table



Part F: Exemplary Specific Compounds

[0780] Another aspect of the invention provides specific compounds of Formula I or II. In certain embodiments, the compound is a compound described in one of the embodiments below, a stereoisomer thereof, or a pharmaceutically acceptable salt of either of the foregoing. In certain embodiments, the compound is a compound described in one of the embodiments below, or a stereoisomer thereof.

[0781] In certain embodiments, the compound is a compound in Table 1, 2, 3, 4, or 5, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1, 2, 3, 4, or 5. In certain embodiments, the compound is a compound in Table 1, 3, 4, or 5, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1, 3, 4, or 5. In certain embodiments, the compound is a compound in Table 3, 4, or 5, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 3, 4, or 5.

[0782] In certain embodiments, the compound is a compound in Table 1 or 3, or a pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula I-A, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1 or 3, wherein the compound is a compound of Formula I-A-1, I-A-2, I-A-3, I-A-4, I-A-3*, I-A-4*, I-A-3**, I-A-4**, I-A-5, I-A-6, or I-A-7. In certain embodiments, the compound is a compound in Table 1 or 4, or a pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula I-B or I-B-1, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1 or 4, wherein the compound is a compound of Formula I-B or I-B-1.

1 or 3, wherein the compound is a compound of Formula I-A-1, I-A-2, I-A-3, I-A-4, I-A-3*, I-A-4*, I-A-3**, I-A-4**, I-A-5, I-A-6, or I-A-7. In certain embodiments, the compound is a compound in Table 1 or 4, or a pharmaceutically acceptable salt thereof, wherein the compound is a compound of Formula I-B or I-B-1, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1 or 4, wherein the compound is a compound of Formula I-B or I-B-1.

[0783] In certain embodiments, the compound is one of compounds III-244, III-245, III-250, III-251, III-275, III-276, III-307, III-308, III-311, III-312, III-321, III-322, III-327, III-328, or III-333 through III-346, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is one of compounds III-244, III-245, III-250, III-251, III-275, III-276, III-307, III-308, III-311, III-312, III-321, III-322, III-327, III-328, or III-333 through III-346. In certain embodiments, the compound is one of compounds III-245, III-251, III-276, III-307, III-311, III-321, III-328, III-334, III-336, III-338, III-339, III-341, III-344, or III-346, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is one of compounds III-245, III-251, III-276, III-307, III-311, III-321, III-328, III-334, III-336, III-338, III-339, III-341, III-344, or III-346.


[0784] In certain embodiments, the compound is a compound in Table 1, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 1.

TABLE 1

Com-
pound
No.

Structure

I-1



I-2





TABLE 1-continued


Com-
pound
No.

Structure

I-3



I-4



I-5





TABLE 1-continued


Com-
pound
No.

Structure


I-6



I-7



I-8



I-9





TABLE 1-continued


Compound
No.

Structure


I-10



I-11



I-12



I-13





TABLE 1-continued


Com-
pound
No.

Structure


I-14



I-15



I-16



I-17





TABLE 1-continued


Com-
pound
No.

Structure


I-18



I-19



I-20



I-21





TABLE 1-continued


Compound
No.

Structure


I-22



I-23



I-24



I-25





TABLE 1-continued


Compound
No.

Structure


I-26



I-27



I-28



I-29





TABLE 1-continued


Com-
pound
No.

Structure


I-30



I-31



I-32



I-33




TABLE 1-continued





Compound No.	Structure
I-34a	
	Stereoisomer I
I-34b	
	Stereoisomer II
I-35	
I-36	

TABLE 1-continued

Compound No.	Structure
I-37a	
I-37b	
I-38	
I-39	


Mixture of Stereoisomer I and Stereoisomer II

TABLE 1-continued


Compound
No.

Structure


I-40



I-41



I-42



I-43





TABLE 1-continued


Com-
pound
No.

Structure


I-44



I-45



I-46



I-47





TABLE 1-continued


Compound
No.

Structure


I-48



I-49



I-50



I-51





TABLE 1-continued


Compound
No.

Structure

I-52



I-53



I-54





TABLE 1-continued


Compound
No.

Structure

I-55



I-56



I-57





TABLE 1-continued


Compound
No.

Structure


I-58



I-59



I-60



I-61





TABLE 1-continued


Compound
No.

Structure


I-62



I-63



I-64



I-65




TABLE 1-continued





Compound No.	Structure
I-66	
I-67	

[0785] In certain embodiments, the compound is a compound in Table 2, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 2.

TABLE 2

Compound No.	Structure
II-1	

TABLE 2-continued

Compound No.	Structure
II-2	
II-3	
II-4	
II-5	

[0786] In certain embodiments, the compound is a compound in Table 3, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 3.

TABLE 3

Compound No.	Compound Structure
III-1	
III-2	
III-3	
III-4	

TABLE 3-continued





Compound No.	Compound Structure
III-5	 Chemical structure of Compound III-5: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring, which is further linked to a 4-(4-phenylpiperidin-1-yl)phenyl group. The latter is connected to a 4-phenylpiperidine ring, which is finally linked to a 4-phenylpiperidine ring. The structure includes a hydroxyl group (OH) on the final phenyl ring.
III-6	 Chemical structure of Compound III-6: Similar to III-5, but the stereochemistry at the chiral centers is different. The hydroxyl group (OH) is on the final phenyl ring.
III-7	 Chemical structure of Compound III-7: Similar to III-5, but the stereochemistry at the chiral centers is different. The hydroxyl group (HO) is on the final phenyl ring.
III-8	 Chemical structure of Compound III-8: Similar to III-5, but the stereochemistry at the chiral centers is different. The hydroxyl group (HO) is on the final phenyl ring.

TABLE 3-continued





Compound No.	Compound Structure
III-9	 Chemical structure of Compound III-9: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a methylene bridge to a piperidine ring, which is further connected to a 4-membered nitrogen-containing ring. This ring is linked via an ether oxygen to a phenyl ring, which is in turn connected to a 1-phenyl-2-(4-hydroxyphenyl)hexahydroindole system. The stereochemistry at the chiral center is (S).
III-10	 Chemical structure of Compound III-10: Similar to III-9, but the stereochemistry at the chiral center is (R).
III-11	 Chemical structure of Compound III-11: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a methylene bridge to a piperidine ring, which is further connected to a 4-membered nitrogen-containing ring. This ring is linked via a methylene bridge to a piperidine ring, which is further connected to a phenyl ring. This phenyl ring is linked via a methylene bridge to a 1-phenyl-2-(4-hydroxyphenyl)hexahydroindole system. The stereochemistry at the chiral center is (S).
III-12	 Chemical structure of Compound III-12: Similar to III-11, but the stereochemistry at the chiral center is (R).

TABLE 3-continued





Compound No.	Compound Structure
III-13	 Chemical structure of Compound III-13: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a methylene bridge to a cyclobutane ring, which is further connected to a piperidine ring. The piperidine ring is substituted with a 4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalene group.
III-14	 Chemical structure of Compound III-14: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a methylene bridge to a cyclobutane ring, which is further connected to a piperidine ring. The piperidine ring is substituted with a 4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalene group.
III-15	 Chemical structure of Compound III-15: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a methylene bridge to a cyclobutane ring, which is further connected to a piperidine ring. The piperidine ring is substituted with a 4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalene group.
III-16	 Chemical structure of Compound III-16: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a methylene bridge to a piperazine ring. The piperazine ring is further connected to a 4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalene group.

TABLE 3-continued





Compound No.	Compound Structure
III-17	 Chemical structure of Compound III-17: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide derivative. The benzothiazine core is substituted with a methyl group at position 7 and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group at position 6. The benzothiazine ring is also substituted with a methyl group at position 4 and a 1,2,4-triazol-5-yl group at position 3. The triazole ring is substituted with a methyl group at position 5 and a 1,2,4-triazol-5-yl group at position 4.
III-18	 Chemical structure of Compound III-18: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide derivative. The benzothiazine core is substituted with a methyl group at position 7 and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group at position 6. The benzothiazine ring is also substituted with a methyl group at position 4 and a 1,2,4-triazol-5-yl group at position 3. The triazole ring is substituted with a methyl group at position 5 and a 1,2,4-triazol-5-yl group at position 4.
III-19	 Chemical structure of Compound III-19: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide derivative. The benzothiazine core is substituted with a methyl group at position 7 and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group at position 6. The benzothiazine ring is also substituted with a methyl group at position 4 and a 1,2,4-triazol-5-yl group at position 3. The triazole ring is substituted with a methyl group at position 5 and a 1,2,4-triazol-5-yl group at position 4.
III-20	 Chemical structure of Compound III-20: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide derivative. The benzothiazine core is substituted with a methyl group at position 7 and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group at position 6. The benzothiazine ring is also substituted with a methyl group at position 4 and a 1,2,4-triazol-5-yl group at position 3. The triazole ring is substituted with a methyl group at position 5 and a 1,2,4-triazol-5-yl group at position 4.

TABLE 3-continued




Compound No.	Compound Structure
III-21	 Chemical structure of Compound III-21: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-((4-((4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-2-yl)phenyl)piperidin-1-yl)but-1-yn-1-yl)phenyl)-6-methyl-1,2,4-thiazole-5-yl group. The triazole ring is also substituted with a 2-furylmethyl group.
III-22	 Chemical structure of Compound III-22: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-((4-((4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-2-yl)phenyl)piperidin-1-yl)but-1-yn-1-yl)phenyl)-6-methyl-1,2,4-thiazole-5-yl group. The triazole ring is also substituted with a 2-furylmethyl group.
III-23	 Chemical structure of Compound III-23: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-((4-((4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-2-yl)phenyl)piperidin-1-yl)but-1-yn-1-yl)phenyl)-6-methyl-1,2,4-thiazole-5-yl group. The triazole ring is also substituted with a 2-furylmethyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-24	 Chemical structure of Compound III-24: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This core is linked via a biphenyl system to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a chain leading to another piperidine ring, which is connected to a biphenyl system and a tetralin derivative with a hydroxyl group.
III-25	 Chemical structure of Compound III-25: Similar to III-24, but the stereochemistry at the chiral center is different, indicated by a wedge bond.
III-26	 Chemical structure of Compound III-26: Similar to III-24, but the stereochemistry at the chiral center is different, indicated by a wedge bond.
III-27	 Chemical structure of Compound III-27: Similar to III-24, but the stereochemistry at the chiral center is different, indicated by a wedge bond.

TABLE 3-continued





Compound No.	Compound Structure
III-28	 Chemical structure of Compound III-28: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)pyrrolidin-1-yl group. The triazole ring is also substituted with a 4-methyl-5-(4-methyl-2-thienyl)phenyl group.
III-29	 Chemical structure of Compound III-29: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl group. The triazole ring is also substituted with a 4-methyl-5-(4-methyl-2-thienyl)phenyl group.
III-30	 Chemical structure of Compound III-30: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl group. The triazole ring is also substituted with a 4-methyl-5-(4-methyl-2-thienyl)phenyl group.
III-31	 Chemical structure of Compound III-31: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl group. The triazole ring is also substituted with a 4-methyl-5-(4-methyl-2-thienyl)phenyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-32	 Chemical structure of Compound III-32: A complex molecule featuring a central fused ring system (a thiazine derivative) substituted with a furan ring and a pyrazole ring. This central core is linked via an amine group to a piperazine ring, which is further connected to a 4-ethoxyphenyl group. The ethoxy group is linked to a biphenyl system, which is substituted with a hydroxyl group.
III-33	 Chemical structure of Compound III-33: A complex molecule featuring a central fused ring system (a thiazine derivative) substituted with a furan ring and a pyrazole ring. This central core is linked via an amine group to a piperazine ring, which is further connected to a 4-ethoxyphenyl group. The ethoxy group is linked to a biphenyl system, which is substituted with a hydroxyl group.
III-34	 Chemical structure of Compound III-34: A complex molecule featuring a central fused ring system (a thiazine derivative) substituted with a furan ring and a pyrazole ring. This central core is linked via an amine group to a piperazine ring, which is further connected to a 4-ethoxyphenyl group. The ethoxy group is linked to a biphenyl system, which is substituted with a hydroxyl group.
III-35	 Chemical structure of Compound III-35: A complex molecule featuring a central fused ring system (a thiazine derivative) substituted with a furan ring and a pyrazole ring. This central core is linked via an amine group to a piperazine ring, which is further connected to a 4-ethoxyphenyl group. The ethoxy group is linked to a biphenyl system, which is substituted with a hydroxyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-36	 Chemical structure of Compound III-36: A complex molecule featuring a thienopyrimidine core substituted with a 4-isoxazol-5-ylmethyl group, a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group.
III-37	 Chemical structure of Compound III-37: A complex molecule featuring a thienopyrimidine core substituted with a 4-isoxazol-5-ylmethyl group, a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group.
III-38	 Chemical structure of Compound III-38: A complex molecule featuring a thienopyrimidine core substituted with a 4-isoxazol-5-ylmethyl group, a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group.
III-39	 Chemical structure of Compound III-39: A complex molecule featuring a thienopyrimidine core substituted with a 4-isoxazol-5-ylmethyl group, a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidine-1-yl group.

TABLE 3-continued





Compound No.	Compound Structure
III-40	 Chemical structure of Compound III-40: A thiazine derivative with a 4-isoxazol-5-ylmethyl group at position 2, a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group at position 4, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidin-1-yl group at position 6. The thiazine ring has methyl groups at positions 5 and 7.
III-41	 Chemical structure of Compound III-41: A thiazine derivative with a 4-isoxazol-5-ylmethyl group at position 2, a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group at position 4, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidin-1-yl group at position 6. The thiazine ring has methyl groups at positions 5 and 7.
III-42	 Chemical structure of Compound III-42: A thiazine derivative with a 4-isoxazol-5-ylmethyl group at position 2, a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group at position 4, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidin-1-yl group at position 6. The thiazine ring has methyl groups at positions 5 and 7.
III-43	 Chemical structure of Compound III-43: A thiazine derivative with a 4-isoxazol-5-ylmethyl group at position 2, a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group at position 4, and a 4-(4-(2-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl)piperidin-1-yl group at position 6. The thiazine ring has methyl groups at positions 5 and 7.

TABLE 3-continued




Compound No.	Compound Structure
III-44	 Chemical structure of Compound III-44: A complex molecule featuring a central benzothiazine core. The core is substituted with a 1,2,4-triazole ring, a thiazole ring, and a phenyl ring. The phenyl ring is further substituted with a piperidine ring, which is connected via a chain to a 4-phenoxyphenyl group. This group is linked to a biphenyl system, which includes a 4-hydroxyphenyl ring.
III-45	 Chemical structure of Compound III-45: A complex molecule featuring a central benzothiazine core. The core is substituted with a 1,2,4-triazole ring, a thiazole ring, and a phenyl ring. The phenyl ring is further substituted with a piperidine ring, which is connected via a chain to a 4-phenoxyphenyl group. This group is linked to a biphenyl system, which includes a 4-hydroxyphenyl ring.
III-46	 Chemical structure of Compound III-46: A complex molecule featuring a central benzothiazine core. The core is substituted with a 1,2,4-triazole ring, a thiazole ring, and a phenyl ring. The phenyl ring is further substituted with a piperidine ring, which is connected via a chain to a 4-phenoxyphenyl group. This group is linked to a biphenyl system, which includes a 4-hydroxyphenyl ring.

TABLE 3-continued





Compound No.	Compound Structure
III-47	 Chemical structure of Compound III-47: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-(2-methyl-5-methylthiophen-2-yl) group, and a 4-(2-methyl-5-methylthiophen-2-yl) group. The triazole ring is linked via a methylene group to a 4-(2-methyl-5-methylthiophen-2-yl) group. This thiophene ring is further substituted with a 4-(2-methyl-5-methylthiophen-2-yl) group. The molecule is connected via a 2-(2-methyl-5-methylthiophen-2-yl) group to a 4-(2-methyl-5-methylthiophen-2-yl) group, which is linked to a 4-(2-methyl-5-methylthiophen-2-yl) group. The structure includes a 4-(2-methyl-5-methylthiophen-2-yl) group and a 4-(2-methyl-5-methylthiophen-2-yl) group.
III-48	 Chemical structure of Compound III-48: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-(2-methyl-5-methylthiophen-2-yl) group, and a 4-(2-methyl-5-methylthiophen-2-yl) group. The triazole ring is linked via a methylene group to a 4-(2-methyl-5-methylthiophen-2-yl) group. This thiophene ring is further substituted with a 4-(2-methyl-5-methylthiophen-2-yl) group. The molecule is connected via a 2-(2-methyl-5-methylthiophen-2-yl) group to a 4-(2-methyl-5-methylthiophen-2-yl) group, which is linked to a 4-(2-methyl-5-methylthiophen-2-yl) group. The structure includes a 4-(2-methyl-5-methylthiophen-2-yl) group and a 4-(2-methyl-5-methylthiophen-2-yl) group.
III-49	 Chemical structure of Compound III-49: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-(2-methyl-5-methylthiophen-2-yl) group, and a 4-(2-methyl-5-methylthiophen-2-yl) group. The triazole ring is linked via a methylene group to a 4-(2-methyl-5-methylthiophen-2-yl) group. This thiophene ring is further substituted with a 4-(2-methyl-5-methylthiophen-2-yl) group. The molecule is connected via a 2-(2-methyl-5-methylthiophen-2-yl) group to a 4-(2-methyl-5-methylthiophen-2-yl) group, which is linked to a 4-(2-methyl-5-methylthiophen-2-yl) group. The structure includes a 4-(2-methyl-5-methylthiophen-2-yl) group and a 4-(2-methyl-5-methylthiophen-2-yl) group.
III-50	 Chemical structure of Compound III-50: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-(2-methyl-5-methylthiophen-2-yl) group, and a 4-(2-methyl-5-methylthiophen-2-yl) group. The triazole ring is linked via a methylene group to a 4-(2-methyl-5-methylthiophen-2-yl) group. This thiophene ring is further substituted with a 4-(2-methyl-5-methylthiophen-2-yl) group. The molecule is connected via a 2-(2-methyl-5-methylthiophen-2-yl) group to a 4-(2-methyl-5-methylthiophen-2-yl) group, which is linked to a 4-(2-methyl-5-methylthiophen-2-yl) group. The structure includes a 4-(2-methyl-5-methylthiophen-2-yl) group and a 4-(2-methyl-5-methylthiophen-2-yl) group.

TABLE 3-continued





Compound No.	Compound Structure
III-51	 Chemical structure of Compound III-51: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.
III-52	 Chemical structure of Compound III-52: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.
III-53	 Chemical structure of Compound III-53: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.
III-54	 Chemical structure of Compound III-54: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-55	 Chemical structure of Compound III-55: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-isoxazolylmethyl group and a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group. This triazole is linked via a methylene bridge to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring. The piperidine is further connected to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring via a 2-(4-phenoxy)ethyl chain.
III-56	 Chemical structure of Compound III-56: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-isoxazolylmethyl group and a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group. This triazole is linked via a methylene bridge to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring. The piperidine is further connected to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring via a 2-(4-phenoxy)ethyl chain.
III-57	 Chemical structure of Compound III-57: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-isoxazolylmethyl group and a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group. This triazole is linked via a methylene bridge to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring. The piperidine is further connected to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring via a 2-(4-phenoxy)ethyl chain.
III-58	 Chemical structure of Compound III-58: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-isoxazolylmethyl group and a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group. This triazole is linked via a methylene bridge to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring. The piperidine is further connected to a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidine ring via a 2-(4-phenoxy)ethyl chain.

TABLE 3-continued




Compound No.	Compound Structure
III-59	 Chemical structure of Compound III-59. It features a central 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group, and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)methyl group. The latter is connected via a 1-azabicyclo[2.2.1]heptane ring.
III-60	 Chemical structure of Compound III-60. It features a central 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group, and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)methyl group. The latter is connected via a 1-azabicyclo[2.2.1]heptane ring.
III-61	 Chemical structure of Compound III-61. It features a central 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group, a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)thiophen-2-yl group, and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)methyl group. The latter is connected via a 1-azabicyclo[2.2.1]heptane ring.

TABLE 3-continued




Compound No.	Compound Structure
III-62	 Chemical structure of Compound III-62: A complex molecule featuring a central fused ring system (a pyrazole-thienopyrimidine derivative) substituted with a furan-2-ylmethyl group and a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The butyl chain is connected via an amine linkage to a piperidine ring.
III-63	 Chemical structure of Compound III-63: A complex molecule featuring a central fused ring system (a pyrazole-thienopyrimidine derivative) substituted with a furan-2-ylmethyl group and a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The butyl chain is connected via an amine linkage to a piperazine ring, which is further substituted with a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.
III-64	 Chemical structure of Compound III-64: A complex molecule featuring a central fused ring system (a pyrazole-thienopyrimidine derivative) substituted with a furan-2-ylmethyl group and a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group. The butyl chain is connected via an amine linkage to a piperidine ring, which is further substituted with a 4-(4-(2-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)butyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-65	 Chemical structure of Compound III-65: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 2-furylmethyl group. This triazole is linked via a methylene bridge to a 2-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidin-1-yl)phenyl group. The piperidine ring is part of a bicyclic system fused to a cyclobutane ring, which is connected via an amine bridge to a 2-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)ethoxy group.
III-66	 Chemical structure of Compound III-66: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 2-furylmethyl group. This triazole is linked via a methylene bridge to a 2-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidin-1-yl)phenyl group. The piperidine ring is part of a bicyclic system fused to a cyclobutane ring, which is connected via an amine bridge to a 2-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)ethoxy group.
III-67	 Chemical structure of Compound III-67: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 2-furylmethyl group. This triazole is linked via a methylene bridge to a 2-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidin-1-yl)phenyl group. The piperidine ring is part of a bicyclic system fused to a cyclobutane ring, which is connected via an amine bridge to a 2-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)ethoxy group.
III-68	 Chemical structure of Compound III-68: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 2-furylmethyl group. This triazole is linked via a methylene bridge to a 2-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidin-1-yl)phenyl group. The piperidine ring is part of a bicyclic system fused to a cyclobutane ring, which is connected via an amine bridge to a 2-(4-(2-((1R,2R)-2-phenyl-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)ethoxy group.

TABLE 3-continued





Compound No.	Compound Structure
III-69	 Chemical structure of Compound III-69: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. The triazole ring is connected via a methylene group to a 4-phenoxyphenyl group. This phenyl group is linked through an ether bridge to a 4-(4-((4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidine-1-yl group. The piperidine ring is further connected to a 4-phenylpiperidine-1-yl group, which is finally linked to a 4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl group. The naphthalene system has a hydroxyl group at the 6-position.
III-70	 Chemical structure of Compound III-70: Similar to III-69, but the stereochemistry at the chiral centers is different. The 4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl group is shown with a different configuration at the 1-position, indicated by a wedge bond.
III-71	 Chemical structure of Compound III-71: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. The triazole ring is connected via a methylene group to a 4-(4-((4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)oxy)phenyl)piperidine-1-yl group. This piperidine ring is further connected to a 4-phenylpiperidine-1-yl group, which is finally linked to a 4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl group. The naphthalene system has a hydroxyl group at the 6-position.
III-72	 Chemical structure of Compound III-72: Similar to III-71, but the stereochemistry at the chiral centers is different. The 4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl group is shown with a different configuration at the 1-position, indicated by a wedge bond.

TABLE 3-continued




Compound No.	Compound Structure
III-73	 Chemical structure of Compound III-73: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiophen-2-yl group. This triazole is linked via a chiral center (dashed bond) to a 4-isoxazol-5-yl group. The triazole is also connected to a 4-phenyl-1,2,4-triazole ring, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The structure includes a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.
III-74	 Chemical structure of Compound III-74: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiophen-2-yl group. This triazole is linked via a chiral center (dashed bond) to a 4-isoxazol-5-yl group. The triazole is also connected to a 4-phenyl-1,2,4-triazole ring, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The structure includes a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.
III-75	 Chemical structure of Compound III-75: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiophen-2-yl group. This triazole is linked via a chiral center (dashed bond) to a 4-isoxazol-5-yl group. The triazole is also connected to a 4-phenyl-1,2,4-triazole ring, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The structure includes a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-76	 Chemical structure of Compound III-76. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group. The triazole ring is connected via a methylene bridge to a piperazine ring. The piperazine ring is further connected via a methylene bridge to a piperidine ring. The piperidine ring is connected via a methylene bridge to a phenyl ring, which is in turn connected via a methylene bridge to a biphenyl system. The biphenyl system consists of a phenyl ring and a 4-hydroxyphenyl ring, with a chiral center indicated by a wedge bond.
III-77a	 Chemical structure of Compound III-77a. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group. The triazole ring is connected via a methylene bridge to a piperazine ring. The piperazine ring is further connected via a methylene bridge to a pyrrolidine ring. The pyrrolidine ring is connected via a methylene bridge to a phenyl ring, which is in turn connected via a methylene bridge to a biphenyl system. The biphenyl system consists of a phenyl ring and a 4-hydroxyphenyl ring, with a chiral center indicated by a wedge bond.
III-77b	 Chemical structure of Compound III-77b. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group. The triazole ring is connected via a methylene bridge to a piperazine ring. The piperazine ring is further connected via a methylene bridge to a pyrrolidine ring. The pyrrolidine ring is connected via a methylene bridge to a phenyl ring, which is in turn connected via a methylene bridge to a biphenyl system. The biphenyl system consists of a phenyl ring and a 4-hydroxyphenyl ring, with a chiral center indicated by a wedge bond.

TABLE 3-continued





Compound No.	Compound Structure
III-78	 Chemical structure of Compound III-78: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-thienyl group. This triazole is linked via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further connected to a 4-phenylazetidine ring, which is linked via a 2-ethoxyethyl chain to a 4-phenoxyphenyl group. This phenyl group is attached to a 1,2,3,4-tetrahydronaphthalene system, which has a hydroxyl group at the 1-position and a phenyl group at the 2-position.
III-79	 Chemical structure of Compound III-79: Similar to III-78, but the stereochemistry at the 2-position of the 1,2,3,4-tetrahydronaphthalene system is inverted, with the phenyl group shown with a dashed bond and the hydroxyl group at the 1-position.
III-80	 Chemical structure of Compound III-80: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-thienyl group. This triazole is linked via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further connected to a 4-phenylazetidine ring, which is linked via a 2-ethoxyethyl chain to a 4-phenoxyphenyl group. This phenyl group is attached to a 1,2,3,4-tetrahydronaphthalene system, which has a hydroxyl group at the 1-position and a phenyl group at the 2-position.
III-81	 Chemical structure of Compound III-81: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-thienyl group. This triazole is linked via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further connected to a 4-phenylazetidine ring, which is linked via a 2-ethoxyethyl chain to a 4-phenoxyphenyl group. This phenyl group is attached to a 1,2,3,4-tetrahydronaphthalene system, which has a hydroxyl group at the 1-position and a phenyl group at the 2-position.

TABLE 3-continued





Compound No.	Compound Structure
III-82	 Chemical structure of Compound III-82: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a 4-phenylene group to a 1,3-dioxolane ring. The dioxolane is further connected to a 1,4-bis(p-phenylene)bis(piperazine) moiety. One of the piperazine rings is substituted with a 4-phenylene group, which is in turn linked to a 1,2,3,4-tetrahydronaphthalen-1-yl group. This tetrahydronaphthalene moiety is substituted with a phenyl group and a 4-hydroxyphenyl group.
III-83	 Chemical structure of Compound III-83: Similar to III-82, but the 1,2,3,4-tetrahydronaphthalene moiety is substituted with a phenyl group and a 4-hydroxyphenyl group, with different stereochemistry at the chiral centers.
III-84	 Chemical structure of Compound III-84: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a 4-phenylene group to a 1,3-dioxolane ring. The dioxolane is further connected to a 1,4-bis(p-phenylene)bis(piperazine) moiety. One of the piperazine rings is substituted with a 4-phenylene group, which is in turn linked to a 1,2,3,4-tetrahydronaphthalen-1-yl group. This tetrahydronaphthalene moiety is substituted with a phenyl group and a 4-hydroxyphenyl group.
III-85	 Chemical structure of Compound III-85: Similar to III-84, but the 1,2,3,4-tetrahydronaphthalene moiety is substituted with a phenyl group and a 4-hydroxyphenyl group, with different stereochemistry at the chiral centers.

TABLE 3-continued





Compound No.	Compound Structure
III-86	 Chemical structure of Compound III-86: A complex molecule featuring a 4-methyl-5-methylthio-1H-imidazo[1,2-a]pyridine core. The core is substituted with a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenoxy group. The stereochemistry is indicated with a dashed bond for the phenyl group on the tetrahydronaphthalene system.
III-87	 Chemical structure of Compound III-87: A complex molecule featuring a 4-methyl-5-methylthio-1H-imidazo[1,2-a]pyridine core. The core is substituted with a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenoxy group. The stereochemistry is indicated with a dashed bond for the phenyl group on the tetrahydronaphthalene system.
III-88	 Chemical structure of Compound III-88: A complex molecule featuring a 4-methyl-5-methylthio-1H-imidazo[1,2-a]pyridine core. The core is substituted with a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenoxy group. The stereochemistry is indicated with a dashed bond for the phenyl group on the tetrahydronaphthalene system.
III-89	 Chemical structure of Compound III-89: A complex molecule featuring a 4-methyl-5-methylthio-1H-imidazo[1,2-a]pyridine core. The core is substituted with a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenoxy group. The stereochemistry is indicated with a dashed bond for the phenyl group on the tetrahydronaphthalene system.

TABLE 3-continued





Compound No.	Compound Structure
III-90	
III-91	
III-92	
III-93a	

TABLE 3-continued




Compound No.	Compound Structure
III-93b	
III-94a	
III-94b	

TABLE 3-continued





Compound No.	Compound Structure
III-95	 Chemical structure of Compound III-95: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide, 1,1-dimethyl-, 1,1-dioxide derivative. The benzothiazine core is substituted at the 4-position with a 4-(4-(2-(4-hydroxy-3-phenyl-3,4,4a,8a-tetrahydronaphthalen-1-yl)phenoxy)azetidin-1-yl)phenyl group. The azetidine ring is connected via a 2-azetidinyl chain to the benzothiazine core.
III-96	 Chemical structure of Compound III-96: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide, 1,1-dimethyl-, 1,1-dioxide derivative. The benzothiazine core is substituted at the 4-position with a 4-(4-(2-(4-hydroxy-3-phenyl-3,4,4a,8a-tetrahydronaphthalen-1-yl)phenoxy)azetidin-1-yl)phenyl group. The azetidine ring is connected via a 2-azetidinyl chain to the benzothiazine core.
III-97	 Chemical structure of Compound III-97: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide, 1,1-dimethyl-, 1,1-dioxide derivative. The benzothiazine core is substituted at the 4-position with a 4-(4-(2-(4-hydroxy-3-phenyl-3,4,4a,8a-tetrahydronaphthalen-1-yl)phenoxy)azetidin-1-yl)phenyl group. The azetidine ring is connected via a 2-azetidinyl chain to the benzothiazine core.
III-98	 Chemical structure of Compound III-98: A 7-methyl-2,3,4,5-tetrahydro-1H-benzothiazine-6-carboxamide, 1,1-dimethyl-, 1,1-dioxide derivative. The benzothiazine core is substituted at the 4-position with a 4-(4-(2-(4-hydroxy-3-phenyl-3,4,4a,8a-tetrahydronaphthalen-1-yl)phenoxy)azetidin-1-yl)phenyl group. The azetidine ring is connected via a 2-azetidinyl chain to the benzothiazine core.

TABLE 3-continued

Compound No.	Compound Structure
III-99	
III-100	
III-101	
III-102	

TABLE 3-continued




Compound No.	Compound Structure
III-103	 Chemical structure of Compound III-103: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group.
III-104	 Chemical structure of Compound III-104: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group.
III-105	 Chemical structure of Compound III-105: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-106	 Chemical structure of Compound III-106: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group.
III-107	 Chemical structure of Compound III-107: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group.
III-108	 Chemical structure of Compound III-108: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenoxy)piperidin-1-yl)benzyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-109	 Chemical structure of Compound III-109: A complex molecule featuring a central fused ring system (a pyrazolo[1,5-a]pyrimidine derivative) substituted with a thiazole ring and a furan ring. This central core is linked via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (OH) and a phenyl ring.
III-110	 Chemical structure of Compound III-110: A complex molecule featuring a central fused ring system (a pyrazolo[1,5-a]pyrimidine derivative) substituted with a thiazole ring and a furan ring. This central core is linked via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (OH) and a phenyl ring.
III-111	 Chemical structure of Compound III-111: A complex molecule featuring a central fused ring system (a pyrazolo[1,5-a]pyrimidine derivative) substituted with a thiazole ring and a furan ring. This central core is linked via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (HO) and a phenyl ring.
III-112	 Chemical structure of Compound III-112: A complex molecule featuring a central fused ring system (a pyrazolo[1,5-a]pyrimidine derivative) substituted with a thiazole ring and a furan ring. This central core is linked via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (HO) and a phenyl ring.

TABLE 3-continued





Compound No.	Compound Structure
III-113	
III-114	
III-115	
III-116	

TABLE 3-continued





Compound No.	Compound Structure
III-117	 Chemical structure of Compound III-117: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group and a 4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl group.
III-118	 Chemical structure of Compound III-118: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group and a 4-(4-(4-(4-hydroxy-4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butyl)piperidin-1-yl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl)phenyl group.
III-119	 Chemical structure of Compound III-119: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butyl)piperidin-1-yl)phenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl)phenyl group.
III-120	 Chemical structure of Compound III-120: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group and a 4-(4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(4-((1R,2R)-2-phenyl-2,3,4,5-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-121	 Chemical structure of Compound III-121: A complex molecule featuring a 7-methyl-2,4,6-trimethyl-1,2,4-triazolo[5,4-b]thiophene core. The triazole ring is substituted with a methyl group at position 7 and methyl groups at positions 2, 4, and 6. The thiophene ring is substituted with a methyl group at position 5 and a 4-(4-methoxy-4,5,6,7-tetrahydropyrimidin-2-yl)phenyl group at position 3. The phenyl ring is further substituted with a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group.
III-122	 Chemical structure of Compound III-122: A complex molecule featuring a 7-methyl-2,4,6-trimethyl-1,2,4-triazolo[5,4-b]thiophene core. The triazole ring is substituted with a methyl group at position 7 and methyl groups at positions 2, 4, and 6. The thiophene ring is substituted with a methyl group at position 5 and a 4-(4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl group at position 3.
III-123	 Chemical structure of Compound III-123: A complex molecule featuring a 7-methyl-2,4,6-trimethyl-1,2,4-triazolo[5,4-b]thiophene core. The triazole ring is substituted with a methyl group at position 7 and methyl groups at positions 2, 4, and 6. The thiophene ring is substituted with a methyl group at position 5 and a 4-(4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)phenyl group at position 3.

TABLE 3-continued




Compound No.	Compound Structure
III-124	 Chemical structure of Compound III-124: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group (via a dashed bond). This triazole is linked to a 2-methyl-4-methylthiophene ring, which is further connected to a phenyl ring. This phenyl ring is linked via a pyrrolidine ring to a piperazine ring. The piperazine ring is connected to a benzyl group, which is further linked to a 1-phenyl-2-phenyl-3-(4-hydroxyphenyl)propane moiety.
III-125	 Chemical structure of Compound III-125: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group (via a dashed bond). This triazole is linked to a 2-methyl-4-methylthiophene ring, which is further connected to a phenyl ring. This phenyl ring is linked via a piperidine ring to a piperazine ring. The piperazine ring is connected to a pyridine ring, which is further linked to a 1-phenyl-2-phenyl-3-(4-hydroxyphenyl)propane moiety.
III-126	 Chemical structure of Compound III-126: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group (via a dashed bond). This triazole is linked to a 2-methyl-4-methylthiophene ring, which is further connected to a phenyl ring. This phenyl ring is linked via a piperidine ring to a piperazine ring. The piperazine ring is connected to a pyridine ring, which is further linked to a 1-phenyl-2-phenyl-3-(4-hydroxyphenyl)propane moiety.

TABLE 3-continued





Compound No.	Compound Structure
III-127	 Chemical structure of Compound III-127: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via a methylene bridge to a 4-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-phenylpiperazine ring. The piperazine ring is linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring. The piperazine ring is also linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring.
III-128	 Chemical structure of Compound III-128: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via a methylene bridge to a 4-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-phenylpiperazine ring. The piperazine ring is linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring. The piperazine ring is also linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring.
III-129	 Chemical structure of Compound III-129: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via a methylene bridge to a 4-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-phenylpiperazine ring. The piperazine ring is linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring. The piperazine ring is also linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring.
III-130	 Chemical structure of Compound III-130: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via a methylene bridge to a 4-methyl-5-methylthiophene ring. The thiophene ring is further connected to a 4-phenylpiperazine ring. The piperazine ring is linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring. The piperazine ring is also linked via a methylene bridge to a 4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)piperazine ring.

TABLE 3-continued




Compound No.	Compound Structure
III-131	 Chemical structure of Compound III-131. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. The triazole ring is connected via a dashed bond to a 4-((4-((4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)methyl)piperidin-1-yl)methyl)azetidin-1-yl)phenyl group.
III-132	 Chemical structure of Compound III-132. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. The triazole ring is connected via a dashed bond to a 4-((4-((4-((4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)methyl)piperidin-1-yl)methyl)piperidin-1-yl)methyl)azetidin-1-yl)phenyl group.
III-133	 Chemical structure of Compound III-133. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. The triazole ring is connected via a dashed bond to a 4-((4-((4-((4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)methyl)piperidin-1-yl)methyl)azetidin-1-yl)methyl)piperidin-1-yl)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-134	 Chemical structure of Compound III-134. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiazole-5-yl group. The thiazole ring is connected via its 5-position to a 4-(azetidin-1-yl)phenyl group. This phenyl group is linked through the nitrogen of the azetidine ring to a carbonyl group, which is further connected to a piperidine ring. The piperidine ring is linked via its nitrogen to another carbonyl group, which is connected to a second azetidine ring. This second azetidine ring is linked through its nitrogen to a 4-phenylphenyl group. The phenyl ring of this group is further substituted with a 1-phenyl-2-(4-hydroxyphenyl)ethyl group.
III-135	 Chemical structure of Compound III-135. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiazole-5-yl group. The thiazole ring is connected via its 5-position to a 4-(azetidin-1-yl)phenyl group. This phenyl group is linked through the nitrogen of the azetidine ring to a carbonyl group, which is further connected to a piperidine ring. The piperidine ring is linked via its nitrogen to another carbonyl group, which is connected to a second azetidine ring. This second azetidine ring is linked through its nitrogen to a 4-phenylphenyl group. The phenyl ring of this group is further substituted with a 1-phenyl-2-(4-hydroxyphenyl)ethyl group.
III-136	 Chemical structure of Compound III-136. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methylthiazole-5-yl group. The thiazole ring is connected via its 5-position to a 4-(azetidin-1-yl)phenyl group. This phenyl group is linked through the nitrogen of the azetidine ring to a carbonyl group, which is further connected to a piperidine ring. The piperidine ring is linked via its nitrogen to another carbonyl group, which is connected to a second azetidine ring. This second azetidine ring is linked through its nitrogen to a 4-phenylphenyl group. The phenyl ring of this group is further substituted with a 1-phenyl-2-(4-hydroxyphenyl)ethyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-137	 Chemical structure of Compound III-137: A complex molecule featuring a 4-methyl-5-methylthio-1,2,4-triazole ring system. The triazole is substituted with a 4-(4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butan-1-one group. The triazole ring also has a methyl group at the 3-position and a methyl group at the 5-position.
III-138	 Chemical structure of Compound III-138: A complex molecule featuring a 4-methyl-5-methylthio-1,2,4-triazole ring system. The triazole is substituted with a 4-(4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butan-1-one group. The triazole ring also has a methyl group at the 3-position and a methyl group at the 5-position.
III-139	 Chemical structure of Compound III-139: A complex molecule featuring a 4-methyl-5-methylthio-1,2,4-triazole ring system. The triazole is substituted with a 4-(4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butan-1-one group. The triazole ring also has a methyl group at the 3-position and a methyl group at the 5-position.
III-140	 Chemical structure of Compound III-140: A complex molecule featuring a 4-methyl-5-methylthio-1,2,4-triazole ring system. The triazole is substituted with a 4-(4-((4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)butan-1-one group. The triazole ring also has a methyl group at the 3-position and a methyl group at the 5-position.

TABLE 3-continued




Compound No.	Compound Structure
III-141	 Chemical structure of Compound III-141: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a phenyl group, which is connected via a methylene bridge to a biphenyl system. The biphenyl system includes a hydroxyl group (OH) and a phenyl ring.
III-142	 Chemical structure of Compound III-142: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a phenyl group, which is connected via a methylene bridge to a biphenyl system. The biphenyl system includes a hydroxyl group (OH) and a phenyl ring.
III-143	 Chemical structure of Compound III-143: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a phenyl group, which is connected via a methylene bridge to a biphenyl system. The biphenyl system includes a hydroxyl group (OH) and a phenyl ring.

TABLE 3-continued




Compound No.	Compound Structure
III-144	 Chemical structure of Compound III-144: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a fluorine atom and a (4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)methyl group.
III-145	 Chemical structure of Compound III-145: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a methyl group and a (4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)methyl group.
III-146	 Chemical structure of Compound III-146: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a pyrrolidine ring. The pyrrolidine ring is further substituted with a (4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)methyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-147	 Chemical structure of Compound III-147: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a chiral center (dashed bond). This system is connected via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (OH) and a phenyl ring.
III-148	 Chemical structure of Compound III-148: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a chiral center (dashed bond). This system is connected via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (OH) and a phenyl ring.
III-149	 Chemical structure of Compound III-149: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a chiral center (dashed bond). This system is connected via a piperazine ring to a biphenyl system, which is further substituted with a hydroxyl group (OH) and a phenyl ring.

TABLE 3-continued





Compound No.	Compound Structure
III-150	 Chemical structure of Compound III-150: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a central carbon atom bonded to a nitrile group and a piperazine ring. The piperazine ring is further substituted with a phenyl group, which is connected to a biphenyl system with a hydroxyl group.
III-151	 Chemical structure of Compound III-151: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a central carbon atom bonded to a fluorine atom and a piperazine ring. The piperazine ring is further substituted with a phenyl group, which is connected to a biphenyl system with a hydroxyl group.
III-152	 Chemical structure of Compound III-152: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a central carbon atom bonded to a hydroxyl group and a piperazine ring. The piperazine ring is further substituted with a phenyl group, which is connected to a biphenyl system with a hydroxyl group.
III-153	 Chemical structure of Compound III-153: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a central carbon atom bonded to a hydroxyl group and a piperazine ring. The piperazine ring is further substituted with a phenyl group, which is connected to a biphenyl system with a hydroxyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-154	 Chemical structure of Compound III-154: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The triazole ring is substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The thiophene ring is substituted with a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.
III-155	 Chemical structure of Compound III-155: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The triazole ring is substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The thiophene ring is substituted with a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.
III-156	 Chemical structure of Compound III-156: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The triazole ring is substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group. The thiophene ring is substituted with a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group and a 4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-157	 Chemical structure of Compound III-157: A complex molecule featuring a thiazine-thienopyrimidine core. The thiazine ring is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The thienopyrimidine core is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The molecule also contains a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group.
III-158	 Chemical structure of Compound III-158: A complex molecule featuring a thiazine-thienopyrimidine core. The thiazine ring is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The thienopyrimidine core is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The molecule also contains a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group.
III-159	 Chemical structure of Compound III-159: A complex molecule featuring a thiazine-thienopyrimidine core. The thiazine ring is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The thienopyrimidine core is substituted with a methyl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group. The molecule also contains a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group and a 4-(4-(2-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1-phenylethyl)-1-piperidin-1-yl)-1H-1,2,3,4-tetrahydroquinoline-2-yl group.

TABLE 3-continued




Compound No.	Compound Structure
III-160	 Chemical structure of Compound III-160. It features a 1,3,4,5-tetramethyl-1H-thiazolo[5,4-c]pyrimidin-2-yl group connected via a methylene bridge to a 4-(azetidin-1-yl)phenyl group. This is further linked through a piperazine ring to a carbonyl group, which is connected to another azetidine ring. This second azetidine is linked to a 4-phenylphenyl group, which is in turn connected to a 4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl group.
III-161	 Chemical structure of Compound III-161. It features a 1,3,4,5-tetramethyl-1H-thiazolo[5,4-c]pyrimidin-2-yl group connected via a methylene bridge to a 4-(azetidin-1-yl)phenyl group. This is further linked through a piperazine ring to a carbonyl group, which is connected to another azetidine ring. This second azetidine is linked to a 4-phenylphenyl group, which is in turn connected to a 4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl group.
III-162	 Chemical structure of Compound III-162. It features a 1,3,4,5-tetramethyl-1H-thiazolo[5,4-c]pyrimidin-2-yl group connected via a methylene bridge to a 4-(azetidin-1-yl)phenyl group. This is further linked through a piperazine ring to a carbonyl group, which is connected to another azetidine ring. This second azetidine is linked to a 4-phenylphenyl group, which is in turn connected to a 4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl group.

TABLE 3-continued




Compound No.	Compound Structure
III-163	 Chemical structure of Compound III-163: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-methyl-5-methylthiophen-2-yl group and a 2-(furan-2-ylmethyl) group. This triazole is linked via a methylene bridge to a p-phenylene ring, which is further connected to a piperidine ring. The piperidine ring is linked via a hydroxymethyl group to another piperidine ring, which is connected to a p-phenylene ring. This p-phenylene ring is linked to a 1,2,3,4-tetrahydronaphthalene system, which has a hydroxyl group at the 6-position and a phenyl group at the 1-position.
III-164	 Chemical structure of Compound III-164: Similar to Compound III-163, but the stereochemistry at the chiral center connecting the triazole system to the p-phenylene ring is different, indicated by a dashed bond.
III-165	 Chemical structure of Compound III-165: A complex molecule featuring a 1,2,4-triazole ring substituted with a 2-methyl-5-methylthiophen-2-yl group and a 2-(furan-2-ylmethyl) group. This triazole is linked via a methylene bridge to a p-phenylene ring, which is further connected to a piperidine ring. The piperidine ring is linked via a hydroxymethyl group to another piperidine ring, which is connected to a p-phenylene ring. This p-phenylene ring is linked to a 1,2,3,4-tetrahydronaphthalene system, which has a hydroxyl group at the 6-position and a phenyl group at the 1-position.

TABLE 3-continued





Compound No.	Compound Structure
III-166	
III-167	
III-168	
III-169	

TABLE 3-continued




Compound No.	Compound Structure
III-170	 Chemical structure of Compound III-170: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with methyl groups. This is connected via a methylene bridge to a p-phenylene ring, which is further linked to a piperidine ring. The piperidine ring is connected to a carbonyl group, which is linked to a pyrrolidine ring. The pyrrolidine ring is connected to a p-phenylene ring, which is further linked to a biphenyl system. The biphenyl system is substituted with a hydroxyl group (OH) and a phenyl ring.
III-171	 Chemical structure of Compound III-171: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with methyl groups. This is connected via a methylene bridge to a p-phenylene ring, which is further linked to a piperidine ring. The piperidine ring is connected to a carbonyl group, which is linked to a pyrrolidine ring. The pyrrolidine ring is connected to a p-phenylene ring, which is further linked to a biphenyl system. The biphenyl system is substituted with a hydroxyl group (OH) and a phenyl ring.
III-172	 Chemical structure of Compound III-172: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with methyl groups. This is connected via a methylene bridge to a p-phenylene ring, which is further linked to a piperidine ring. The piperidine ring is connected to a carbonyl group, which is linked to a pyrrolidine ring. The pyrrolidine ring is connected to a p-phenylene ring, which is further linked to a biphenyl system. The biphenyl system is substituted with a hydroxyl group (OH) and a phenyl ring.

TABLE 3-continued




Compound No.	Compound Structure
III-173	 Chemical structure of Compound III-173: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)phenyl)piperidin-1-yl group.
III-174	 Chemical structure of Compound III-174: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)phenyl)piperidin-1-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)phenyl)piperidin-1-yl group.
III-175	 Chemical structure of Compound III-175: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)phenyl)piperidin-1-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxy-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)azetidin-1-yl)phenyl)piperidin-1-yl group.

TABLE 3-continued

Compound No.	Compound Structure
III-176	
III-177	
III-178	

TABLE 3-continued




Compound No.	Compound Structure
III-179	 Chemical structure of Compound III-179. It features a central 1,2,4-triazole ring substituted with a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The triazole ring is also substituted with a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group.
III-180	 Chemical structure of Compound III-180. It features a central 1,2,4-triazole ring substituted with a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The triazole ring is also substituted with a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group.
III-181	 Chemical structure of Compound III-181. It features a central 1,2,4-triazole ring substituted with a 4-methyl-5-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The triazole ring is also substituted with a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group. The structure includes a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group and a 4-(4-methyl-1,3,4-thiadiazol-2-yl)phenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-182	 Chemical structure of Compound III-182: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. This triazole is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(4-(2-hydroxy-3-(4-fluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)piperidin-1-yl)methyl group.
III-183	 Chemical structure of Compound III-183: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. This triazole is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(4-(2-(4-fluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)piperidin-1-yl)methyl group.
III-184	 Chemical structure of Compound III-184: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazol-2-yl group. This triazole is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(4-(2-(4-fluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)piperidin-1-yl)methyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-185	 Chemical structure of Compound III-185: A complex molecule featuring a 1,2,4-triazole-thienothiopyran core. The triazole ring is substituted with a methyl group and a dashed bond. The thiopyran ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected to a piperidine ring, which is further connected to a hydroxyl group (HO). The piperidine ring is also connected to a piperazine ring, which is connected to a 2,4-difluorophenyl group. The 2,4-difluorophenyl group is connected to a biphenyl system, which is further connected to a hydroxyl group (OH).
III-186	 Chemical structure of Compound III-186: A complex molecule featuring a 1,2,4-triazole-thienothiopyran core. The triazole ring is substituted with a methyl group and a dashed bond. The thiopyran ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected to a piperidine ring, which is further connected to a hydroxyl group (OH). The piperidine ring is also connected to a piperazine ring, which is connected to a 2,4-difluorophenyl group. The 2,4-difluorophenyl group is connected to a biphenyl system, which is further connected to a hydroxyl group (OH).
III-187	 Chemical structure of Compound III-187: A complex molecule featuring a 1,2,4-triazole-thienothiopyran core. The triazole ring is substituted with a methyl group and a dashed bond. The thiopyran ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected to a piperidine ring, which is further connected to a hydroxyl group (HO). The piperidine ring is also connected to a piperazine ring, which is connected to a 2,4-difluorophenyl group. The 2,4-difluorophenyl group is connected to a biphenyl system, which is further connected to a hydroxyl group (OH).

TABLE 3-continued




Compound No.	Compound Structure
III-188	 Chemical structure of Compound III-188: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The molecule is shown with stereochemistry: a dashed bond to the triazole ring and a solid bond to the thiophene ring.
III-189	 Chemical structure of Compound III-189: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The molecule is shown with stereochemistry: a dashed bond to the triazole ring and a dashed bond to the thiophene ring.
III-190	 Chemical structure of Compound III-190: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-hydroxy-4-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The molecule is shown with stereochemistry: a dashed bond to the triazole ring and a dashed bond to the thiophene ring.

TABLE 3-continued





Compound No.	Compound Structure
III-191	
III-192	
III-193	
III-194	

TABLE 3-continued

Compound No.	Compound Structure
III-195	
III-196	
III-197	

TABLE 3-continued




Compound No.	Compound Structure
III-198	
III-199	
III-200	

TABLE 3-continued




Compound No.	Compound Structure
III-201	 Chemical structure of Compound III-201: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a pyrazole ring. This core is linked via a methylene group to a piperidine ring. The piperidine ring is connected to a piperazine ring, which is in turn linked to a biphenyl system. One of the phenyl rings of the biphenyl system is substituted with a carboxylic acid group (COOH).
III-202	 Chemical structure of Compound III-202: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a pyrazole ring. This core is linked via a methylene group to a piperidine ring. The piperidine ring is connected to a piperazine ring, which is in turn linked to a biphenyl system. One of the phenyl rings of the biphenyl system is substituted with a hydroxyl group (OH).
III-203	 Chemical structure of Compound III-203: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a pyrazole ring. This core is linked via a methylene group to a piperidine ring. The piperidine ring is connected to a piperazine ring, which is in turn linked to a biphenyl system. One of the phenyl rings of the biphenyl system is substituted with a hydroxyl group (OH).

TABLE 3-continued

Compound No.	Compound Structure
III-204	
III-205	
III-206	

TABLE 3-continued

Compound No.	Compound Structure
III-207	
III-208	
III-209	

TABLE 3-continued




Compound No.	Compound Structure
III-210	 Chemical structure of Compound III-210: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl group. The triazole ring is substituted with a 4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)piperidin-1-yl)phenyl group.
III-211	 Chemical structure of Compound III-211: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-(4-phenyl-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)piperidin-1-yl)phenyl)azetidin-1-yl group. The triazole ring is substituted with a 4-(4-(4-(4-phenyl-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)piperidin-1-yl)phenyl)azetidin-1-yl group.
III-212	 Chemical structure of Compound III-212: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-(4-(4-phenyl-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)azetidin-1-yl)piperidin-1-yl)phenyl)azetidin-1-yl group. The triazole ring is substituted with a 4-(4-(4-(4-(4-phenyl-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)azetidin-1-yl)piperidin-1-yl)phenyl)azetidin-1-yl group.

TABLE 3-continued




Compound No.	Compound Structure
III-213	 Chemical structure of Compound III-213: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. The thiophene ring is substituted with two methyl groups and a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group.
III-214	 Chemical structure of Compound III-214: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. The thiophene ring is substituted with two methyl groups and a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group.
III-215	 Chemical structure of Compound III-215: A complex molecule featuring a 1,2,4-triazole-thiophene core. The triazole ring is substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. The thiophene ring is substituted with two methyl groups and a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group. The thiophene ring is also substituted with a 4-(4-hydroxy-4-(4-(2,2,6,6-tetrafluorophenyl)-1,2,3,6-tetrahydropyridin-1-yl)phenyl)phenyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-216	
III-217	
III-218	
III-219	

TABLE 3-continued




Compound No.	Compound Structure
III-220	
III-221	
III-222	

TABLE 3-continued




Compound No.	Compound Structure
III-223	
III-224	
III-225	

TABLE 3-continued





Compound No.	Compound Structure
III-226	 Chemical structure of Compound III-226: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The molecule includes a piperidine ring with a hydroxyl group and a piperazine ring.
III-227	 Chemical structure of Compound III-227: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The molecule includes a piperidine ring with a fluorine atom and a piperazine ring.
III-228	 Chemical structure of Compound III-228: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The molecule includes a piperidine ring and a piperazine ring.
III-229	 Chemical structure of Compound III-229: A complex molecule featuring a 1,2,4-triazole ring substituted with a 4-isoxazol-5-ylmethyl group and a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The molecule includes a piperidine ring and a piperazine ring.

TABLE 3-continued




Compound No.	Compound Structure
III-230	 Chemical structure of Compound III-230: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. The triazole ring is connected via a methylene group to a 4-phenyl-1,3-dioxane ring. This is further connected via a methylene group to a 4-phenyl-1,3-dioxane ring, which is then connected via a methylene group to a 4-phenyl-1,3-dioxane ring. The final 4-phenyl-1,3-dioxane ring is substituted with a 1-phenyl-2-phenyl-3-hydroxy-1,2,3,4-tetrahydronaphthalene group.
III-231	 Chemical structure of Compound III-231: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. The triazole ring is connected via a methylene group to a 4-phenyl-1,3-dioxane ring. This is further connected via a methylene group to a 4-phenyl-1,3-dioxane ring, which is then connected via a methylene group to a 4-phenyl-1,3-dioxane ring. The final 4-phenyl-1,3-dioxane ring is substituted with a 1-phenyl-2-phenyl-3-hydroxy-1,2,3,4-tetrahydronaphthalene group.
III-232	 Chemical structure of Compound III-232: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. The triazole ring is connected via a methylene group to a 4-phenyl-1,3-dioxane ring. This is further connected via a methylene group to a 4-phenyl-1,3-dioxane ring, which is then connected via a methylene group to a 4-phenyl-1,3-dioxane ring. The final 4-phenyl-1,3-dioxane ring is substituted with a 1-phenyl-2-phenyl-3-hydroxy-1,2,3,4-tetrahydronaphthalene group.

TABLE 3-continued




Compound No.	Compound Structure
III-233	 Chemical structure of Compound III-233. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. This dioxane ring is further connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is in turn connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The final dioxane ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is finally connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The structure is symmetrical, with the central dioxane ring being the most complex, featuring a 4-phenyl-1,3-dioxane ring and a 4-phenyl-1,3-dioxane ring.
III-234	 Chemical structure of Compound III-234. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. This dioxane ring is further connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is in turn connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The final dioxane ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is finally connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The structure is symmetrical, with the central dioxane ring being the most complex, featuring a 4-phenyl-1,3-dioxane ring and a 4-phenyl-1,3-dioxane ring.
III-235	 Chemical structure of Compound III-235. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. This dioxane ring is further connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is in turn connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The final dioxane ring is connected via a dashed bond to a 4-phenyl-1,3-dioxane ring, which is finally connected via a dashed bond to a 4-phenyl-1,3-dioxane ring. The structure is symmetrical, with the central dioxane ring being the most complex, featuring a 4-phenyl-1,3-dioxane ring and a 4-phenyl-1,3-dioxane ring.

TABLE 3-continued




Compound No.	Compound Structure
III-236	 Chemical structure of Compound III-236. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a p-phenylene ring, which is further linked to a piperidine ring. This piperidine ring is connected to a 1,4-bis(piperidin-1-yl)butane chain, which is then linked to a 2,6-difluorophenyl ring. The 2,6-difluorophenyl ring is connected to a 4-fluorophenyl ring, which is finally linked to a 4-hydroxyphenyl ring.
III-237	 Chemical structure of Compound III-237. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a p-phenylene ring, which is further linked to a piperidine ring. This piperidine ring is connected to a 1,4-bis(piperidin-1-yl)butane chain, which is then linked to a 2,6-difluorophenyl ring. The 2,6-difluorophenyl ring is connected to a 4-fluorophenyl ring, which is finally linked to a 4-hydroxyphenyl ring.
III-238	 Chemical structure of Compound III-238. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-thiazole ring. The triazole ring is connected via a dashed bond to a p-phenylene ring, which is further linked to a piperidine ring. This piperidine ring is connected to a 1,4-bis(piperidin-1-yl)butane chain, which is then linked to a 2,6-difluorophenyl ring. The 2,6-difluorophenyl ring is connected to a 4-fluorophenyl ring, which is finally linked to a 4-hydroxyphenyl ring.

TABLE 3-continued




Compound No.	Compound Structure
III-239	 Chemical structure of Compound III-239: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The molecule is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group.
III-240	 Chemical structure of Compound III-240: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The molecule is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group.
III-241	 Chemical structure of Compound III-241: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The thiophene ring is also substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group. The molecule is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group, which is further linked to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-1H-1,2,4-triazol-5-yl group.

TABLE 3-continued

Compound No.	Compound Structure
III-242	
III-243	
III-244	
III-245	

TABLE 3-continued




Compound No.	Compound Structure
III-246	 Chemical structure of Compound III-246: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1-hydroxy-2-(4-(2,2-diphenyl-1-hydroxyethyl)-3,5-difluorophenyl)ethyl piperidine derivative. The structure includes stereochemical indicators (wedges and dashes) at the chiral centers.
III-247	 Chemical structure of Compound III-247: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1-hydroxy-2-(4-(2,2-diphenyl-1-hydroxyethyl)-3,5-difluorophenyl)ethyl piperidine derivative. The structure includes stereochemical indicators (wedges and dashes) at the chiral centers.
III-248	 Chemical structure of Compound III-248: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1-hydroxy-2-(4-(2,2-diphenyl-1-hydroxyethyl)-3,5-difluorophenyl)ethyl piperidine derivative. The structure includes stereochemical indicators (wedges and dashes) at the chiral centers.

TABLE 3-continued




Compound No.	Compound Structure
III-249	 Chemical structure of Compound III-249: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This system is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(2-(4-fluorophenyl)-4-hydroxyphenyl)piperidin-1-yl)methyl group.
III-250	 Chemical structure of Compound III-250: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This system is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(2-(4-fluorophenyl)-4-hydroxyphenyl)piperidin-1-yl)methyl group.
III-251	 Chemical structure of Compound III-251: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This system is linked via a methylene bridge to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(2-(4-fluorophenyl)-4-hydroxyphenyl)piperidin-1-yl)methyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-252	 Chemical structure of Compound III-252: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is further substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring.
III-253	 Chemical structure of Compound III-253: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is further substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring.
III-254	 Chemical structure of Compound III-254: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is further substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring.
III-255	 Chemical structure of Compound III-255: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is further substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring. The 1,3-dioxane ring is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group. The 4-hydroxyphenyl group is connected to a 4-methoxyphenyl group via a methylene bridge, which is in turn connected to a 4-phenyl-1,3-dioxane ring.

TABLE 3-continued




Compound No.	Compound Structure
III-256	 Chemical structure of Compound III-256: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1,4-bis(piperidin-1-yl)benzene moiety. The second piperidine ring is further substituted with a 2,6-difluorophenyl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group.
III-257	 Chemical structure of Compound III-257: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1,4-bis(piperidin-1-yl)benzene moiety. The second piperidine ring is further substituted with a 2,6-difluorophenyl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group.
III-258	 Chemical structure of Compound III-258: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a p-phenylene group. This p-phenylene group is connected via a piperidine ring to a 1,4-bis(piperidin-1-yl)benzene moiety. The second piperidine ring is further substituted with a 2,6-difluorophenyl group and a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is also substituted with a 4-fluorophenyl group and a 4-hydroxyphenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-259	 Chemical structure of Compound III-259. The molecule features a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via its 4-position to a 2-methyl-5-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl)phenyl group. The phenyl ring is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group through a piperazine ring system.
III-260	 Chemical structure of Compound III-260. The molecule features a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via its 4-position to a 2-methyl-5-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl)phenyl group. The phenyl ring is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group through a piperazine ring system.
III-261	 Chemical structure of Compound III-261. The molecule features a 1,2,4-triazole ring substituted with a methyl group and a 4-isoxazol-5-ylmethyl group. This triazole is linked via its 4-position to a 2-methyl-5-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl)phenyl group. The phenyl ring is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group through a piperazine ring system.

TABLE 3-continued





Compound No.	Compound Structure
III-262	
III-263	
III-264	
III-265	

TABLE 3-continued




Compound No.	Compound Structure
III-266	 Chemical structure of Compound III-266: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group. This core is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group, which is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group.
III-267	 Chemical structure of Compound III-267: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group. This core is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group, which is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group.
III-268	 Chemical structure of Compound III-268: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group. This core is linked via a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group, which is further connected to a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2,6-difluorophenyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-269	 Chemical structure of Compound III-269: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (2-(4-(2-(4-hydroxy-3-(2,4-difluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)phenyl)ethyl)piperidin-1-yl group.
III-270	 Chemical structure of Compound III-270: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (2-(4-(2-(4-(2,4-difluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)phenyl)ethyl)piperidin-1-yl group.
III-271	 Chemical structure of Compound III-271: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (2-(4-(2-(4-(2,4-difluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)phenyl)ethyl)piperidin-1-yl group.
III-272	 Chemical structure of Compound III-272: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (2-(4-(2-(4-(2,4-difluorophenyl)-3,4-dihydroquinolin-1-yl)phenyl)phenyl)ethyl)piperidin-1-yl group.

TABLE 3-continued





Compound No.	Compound Structure
III-273	
III-274	
III-275	
III-276	

TABLE 3-continued




Compound No.	Compound Structure
III-277	 Chemical structure of Compound III-277: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperazine ring to a biphenyl system. The biphenyl system includes a 4-hydroxyphenyl group and a 2,6-difluorophenyl group.
III-278	 Chemical structure of Compound III-278: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperazine ring to a biphenyl system. The biphenyl system includes a 4-hydroxyphenyl group and a 2,6-difluorophenyl group.
III-279	 Chemical structure of Compound III-279: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperazine ring to a biphenyl system. The biphenyl system includes a 4-hydroxyphenyl group and a 2,6-difluorophenyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-280	 Chemical structure of Compound III-280. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a 4-phenylpiperazine ring to a 4-hydroxycyclohexyl group. The cyclohexyl group is further substituted with a 2,6-difluorophenyl ring and a 4-fluorophenyl ring. The piperazine ring is also substituted with a 4-hydroxycyclohexyl group.
III-281	 Chemical structure of Compound III-281. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a 4-phenylpiperazine ring to a 4-hydroxycyclohexyl group. The cyclohexyl group is further substituted with a 2,6-difluorophenyl ring and a 4-fluorophenyl ring. The piperazine ring is also substituted with a 4-hydroxycyclohexyl group.
III-282	 Chemical structure of Compound III-282. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. This triazole is linked via a 4-phenylpiperazine ring to a 4-hydroxycyclohexyl group. The cyclohexyl group is further substituted with a 2,6-difluorophenyl ring and a 4-fluorophenyl ring. The piperazine ring is also substituted with a 4-hydroxycyclohexyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-283	 Chemical structure of Compound III-283: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-thienyl group. This triazole is linked via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further substituted with a hydroxyl group and a (4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)methyl group. The tetrahydronaphthalene moiety is also substituted with a methyl group.
III-284	 Chemical structure of Compound III-284: Similar to III-283, but the tetrahydronaphthalene moiety is substituted with a fluorine atom at the 1-position and a hydroxyl group at the 4-position. The stereochemistry at the 1-position is indicated by a wedge bond.
III-285	 Chemical structure of Compound III-285: Similar to III-283, but the tetrahydronaphthalene moiety is substituted with a fluorine atom at the 1-position and a hydroxyl group at the 4-position. The stereochemistry at the 1-position is indicated by a dashed bond.
III-286	 Chemical structure of Compound III-286: Similar to III-284, but the tetrahydronaphthalene moiety is substituted with a fluorine atom at the 1-position and a hydroxyl group at the 4-position. The stereochemistry at the 1-position is indicated by a wedge bond.

TABLE 3-continued





Compound No.	Compound Structure
III-287	 Chemical structure of Compound III-287: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The pyridine ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The two phenyl rings are connected via a 1,4-bis(piperidin-2-yl)benzene linker.
III-288	 Chemical structure of Compound III-288: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The pyridine ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The two phenyl rings are connected via a 1,4-bis(piperidin-2-yl)benzene linker.
III-289	 Chemical structure of Compound III-289: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The pyridine ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The two phenyl rings are connected via a 1,4-bis(piperidin-2-yl)benzene linker.
III-290	 Chemical structure of Compound III-290: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The pyridine ring is substituted with a methyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl group. The two phenyl rings are connected via a 1,4-bis(piperidin-2-yl)benzene linker.

TABLE 3-continued





Compound No.	Compound Structure
III-291	 Chemical structure of Compound III-291: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is linked to a 1-hydroxy-2-(4-(2-fluorophenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)ethyl group.
III-292	 Chemical structure of Compound III-292: Similar to III-291, but the stereochemistry at the chiral center is inverted, resulting in a (1S,2S)-1-hydroxy-2-(4-(2-fluorophenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)ethyl group.
III-293	 Chemical structure of Compound III-293: Similar to III-291, but the stereochemistry at the chiral center is inverted, resulting in a (1R,2R)-1-hydroxy-2-(4-(2-fluorophenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)ethyl group.
III-294	 Chemical structure of Compound III-294: Similar to III-292, but the stereochemistry at the chiral center is inverted, resulting in a (1R,2R)-1-hydroxy-2-(4-(2-fluorophenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)ethyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-295	
III-296	
III-297	
III-298	

TABLE 3-continued





Compound No.	Compound Structure
III-299	
III-300	
III-301	
III-302	

TABLE 3-continued




Compound No.	Compound Structure
III-303	
III-304	
III-305	

TABLE 3-continued





Compound No.	Compound Structure
III-306	 Chemical structure of Compound III-306: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with two methyl groups and a (4-(4-chloro-2-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group. The triazole ring is also substituted with a (4-(4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group.
III-307	 Chemical structure of Compound III-307: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with two methyl groups and a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group. The triazole ring is also substituted with a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group.
III-308	 Chemical structure of Compound III-308: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with two methyl groups and a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group. The triazole ring is also substituted with a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group.
III-309	 Chemical structure of Compound III-309: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with two methyl groups and a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group. The triazole ring is also substituted with a (4-(4-(4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl group.

TABLE 3-continued

Compound No.	Compound Structure
III-310	 <chem>Cc1nc2c(ncn2C)c3cc(ccc3N1Cc4ccc(cc4)N5CCc6cc(ccc6)N5Cc7cc(OC)c(cc7F)C8Cc9ccccc9CC8O)O</chem>
III-311	 <chem>Cc1nc2c(ncn2C)c3cc(ccc3N1Cc4ccc(cc4)N5CCc6cc(ccc6)N5Cc7cc(OC)c(cc7F)C8Cc9ccccc9CC8O)O</chem>
III-312	 <chem>Cc1nc2c(ncn2C)c3cc(ccc3N1Cc4ccc(cc4)N5CCc6cc(ccc6)N5Cc7cc(OC)c(cc7F)C8Cc9ccccc9CC8O)O</chem>
III-313	 <chem>Cc1nc2c(ncn2C)c3cc(ccc3N1Cc4ccc(cc4)N5CCc6cc(ccc6)N5Cc7cc(OC)c(cc7F)C8Cc9ccccc9CC8O)O</chem>

TABLE 3-continued

Compound No.	Compound Structure
III-314	
III-315	
III-316	
III-317	

TABLE 3-continued





Compound No.	Compound Structure
III-318	
III-319	
III-320	
III-321	

TABLE 3-continued





Compound No.	Compound Structure
III-322	
III-323	
III-324	
III-325	

TABLE 3-continued

Compound No.	Compound Structure
III-326	
III-327	
III-328	
III-329	

TABLE 3-continued




Compound No.	Compound Structure
III-330	 Chemical structure of Compound III-330: A complex molecule featuring a 1,2,4-triazole-thienyl core substituted with methyl groups and a chiral center. This core is linked via a p-phenylene ring to a piperidine ring, which is further connected to a 1,4-bis(methoxy)phenyl ring. This ring is then linked to a 4-methylphenyl ring, which is finally connected to a 4-hydroxy-1,2,3,4-tetrahydronaphthalene moiety.
III-331	 Chemical structure of Compound III-331: Similar to III-330, but the p-phenylene ring is substituted with a fluorine atom at the 3-position. The stereochemistry at the chiral center is inverted (dashed bond).
III-332	 Chemical structure of Compound III-332: Similar to III-330, but the p-phenylene ring is substituted with a fluorine atom at the 3-position. The stereochemistry at the chiral center is solid (wedged bond).

TABLE 3-continued





Compound No.	Compound Structure
III-333	
III-334	
III-335	
III-336	

TABLE 3-continued





Compound No.	Compound Structure
III-337	
III-338	
III-339	
III-340	

TABLE 3-continued

Compound No.	Compound Structure
III-341	
III-342	
III-343	
III-344	

TABLE 3-continued





Compound No.	Compound Structure
III-345	 Chemical structure of Compound III-345: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a methyl group and a 4-fluorophenyl group. This core is linked via a piperidine ring (with a hydroxyl group) to a second piperidine ring, which is further connected to a 4-methoxyphenyl group. The 4-methoxyphenyl group is also linked to a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is substituted with a 2-fluorophenyl group and a hydroxyl group.
III-346	 Chemical structure of Compound III-346: Similar to III-345, but the stereochemistry at the chiral centers is different. The 1,2,3,4-tetrahydronaphthalen-1-yl group is substituted with a 2-fluorophenyl group and a hydroxyl group, and the 4-methoxyphenyl group is linked to it.
III-347	 Chemical structure of Compound III-347: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a methyl group and a 4-fluorophenyl group. This core is linked via a piperidine ring (with a hydroxyl group) to a second piperidine ring, which is further connected to a 4-methoxyphenyl group. The 4-methoxyphenyl group is also linked to a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is substituted with a 2-fluorophenyl group and a hydroxyl group.
III-348	 Chemical structure of Compound III-348: A complex molecule featuring a 1,2,4-triazole-thiophene core substituted with a methyl group and a 4-fluorophenyl group. This core is linked via a piperidine ring (with a hydroxyl group) to a second piperidine ring, which is further connected to a 4-methoxyphenyl group. The 4-methoxyphenyl group is also linked to a 1,2,3,4-tetrahydronaphthalen-1-yl group, which is substituted with a 2-fluorophenyl group and a hydroxyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-349	 Chemical structure of Compound III-349: A complex molecule featuring a 1,2,4-triazole-thienopyrimidine core. The triazole ring is substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxyphenyl)-4-(trifluoromethyl)phenyl)piperidin-1-yl)piperidin-1-yl)methyl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-(4-(4-hydroxyphenyl)-4-(trifluoromethyl)phenyl)piperidin-1-yl)piperidin-1-yl)methyl group.
III-350	 Chemical structure of Compound III-350: A complex molecule featuring a 1,2,4-triazole-thienopyrimidine core. The triazole ring is substituted with a methyl group and a 4-(4-(4-(4-(4-(4-fluorophenyl)piperidin-1-yl)piperidin-1-yl)methyl)piperidin-1-yl)methyl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-(4-(4-(4-fluorophenyl)piperidin-1-yl)piperidin-1-yl)methyl)piperidin-1-yl)methyl group.
III-351	 Chemical structure of Compound III-351: A complex molecule featuring a 1,2,4-triazole-thienopyrimidine core. The triazole ring is substituted with a methyl group and a 4-(4-(4-(4-(4-(4-(4-fluorophenyl)piperidin-1-yl)piperidin-1-yl)methyl)piperidin-1-yl)methyl)piperidin-1-yl)methyl group. The thiophene ring is substituted with a methyl group and a 4-(4-(4-(4-(4-(4-(4-fluorophenyl)piperidin-1-yl)piperidin-1-yl)methyl)piperidin-1-yl)methyl)piperidin-1-yl)methyl group.

TABLE 3-continued




Compound No.	Compound Structure
III-352	 Chemical structure of compound III-352: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a methylene bridge to a p-methoxyphenyl group. The p-methoxyphenyl group is linked to a piperidine ring, which is further connected to a piperazine ring. The piperazine ring is substituted with a 4-fluorophenyl group and a 4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl group.
III-353	 Chemical structure of compound III-353: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a methylene bridge to a p-methoxyphenyl group. The p-methoxyphenyl group is linked to a piperidine ring, which is further connected to a piperazine ring. The piperazine ring is substituted with a 4-fluorophenyl group and a 4-(4-fluorophenyl)-4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl group.
III-354	 Chemical structure of compound III-354: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is substituted with a methyl group. This system is connected via a methylene bridge to a p-fluorophenyl group. The p-fluorophenyl group is linked to a piperidine ring, which is further connected to a piperazine ring. The piperazine ring is substituted with a 4-(4-(trifluoromethyl)phenoxy)phenyl group and a 4-(4-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl group.

TABLE 3-continued





Compound No.	Compound Structure
III-355	
III-356	
III-357	
III-358	

TABLE 3-continued

Compound No.	Compound Structure
III-359	

[0787] In certain embodiments, the compound is a compound in Table 4, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 4.

TABLE 4

Compound No.	Compound Structure
IV-1	
IV-2	

TABLE 4-continued





Compound No.	Compound Structure
IV-3	 Chemical structure of compound IV-3: A 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group is connected via a methylene bridge to a 4-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenyl)piperazine. The piperazine ring is further connected via a 2-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenoxy)ethyl chain to a 4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)phenyl group, which is substituted with a carboxylic acid group.
IV-4	 Chemical structure of compound IV-4: A 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group is connected via a methylene bridge to a 4-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenyl)piperazine. The piperazine ring is further connected via a 2-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenoxy)ethyl chain to a 4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)phenyl group, which is substituted with a carboxylic acid group.
IV-5	 Chemical structure of compound IV-5: A 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group is connected via a methylene bridge to a 4-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenyl)piperazine. The piperazine ring is further connected via a 2-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenoxy)ethyl chain to a 4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)phenyl group, which is substituted with a carboxylic acid group.
IV-6	 Chemical structure of compound IV-6: A 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole-2-yl group is connected via a methylene bridge to a 4-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenyl)piperazine. The piperazine ring is further connected via a 2-(4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)-2-chlorophenoxy)ethyl chain to a 4-(2-(3,4-dichlorophenyl)-7,8,9,10-tetrahydro-7H-benzo[7,9]furan-6-yl)phenyl group, which is substituted with a carboxylic acid group.

TABLE 4-continued





Compound No.	Compound Structure
IV-7	
IV-8	
IV-9	
IV-10	

TABLE 4-continued





Compound No.	Compound Structure
IV-11	 Chemical structure of compound IV-11: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring, which is further linked to a 1,4-bis(4-chlorophenyl)piperazine system. The piperazine is connected to a 4-phenylpiperidine ring, which is finally linked to a 4-chlorophenyl ring substituted with a carboxylic acid group.
IV-12	 Chemical structure of compound IV-12: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring, which is further linked to a 1,4-bis(4-chlorophenyl)piperazine system. The piperazine is connected to a 4-phenylpiperidine ring, which is finally linked to a 4-chlorophenyl ring substituted with a carboxylic acid group.
IV-13	 Chemical structure of compound IV-13: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring, which is further linked to a 1,4-bis(4-chlorophenyl)piperazine system. The piperazine is connected to a 4-phenylpiperidine ring, which is finally linked to a 4-chlorophenyl ring substituted with a carboxylic acid group.
IV-14	 Chemical structure of compound IV-14: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring, which is further linked to a 1,4-bis(4-chlorophenyl)piperazine system. The piperazine is connected to a 4-phenylpiperidine ring, which is finally linked to a 4-chlorophenyl ring substituted with a carboxylic acid group.

TABLE 4-continued





Compound No.	Compound Structure
IV-15	
IV-16	
IV-17	
IV-18	

TABLE 4-continued





Compound No.	Compound Structure
IV-19	
IV-20	
IV-21	
IV-22	

TABLE 4-continued

Compound No.	Compound Structure
IV-23	
IV-24	
IV-25	

TABLE 4-continued




Compound No.	Compound Structure
IV-26	 Chemical structure of compound IV-26: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is substituted with a hydroxyl group and a (2-(2,6-difluorophenyl)-7-oxo-7,8,9,10-tetrahydrobenzo[b]furan-5-yl)methyl group.
IV-27	 Chemical structure of compound IV-27: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is substituted with a fluorine atom and a (2-(2,6-dichlorophenyl)-7-oxo-7,8,9,10-tetrahydrobenzo[b]furan-5-yl)methyl group.
IV-28	 Chemical structure of compound IV-28: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring, which is further substituted with a methyl group. This system is connected via a p-phenylene ring to a piperidine ring. The piperidine ring is substituted with a hydroxyl group and a (2-(2-fluorophenyl)-7-phenyl-7,8,9,10-tetrahydrobenzo[b]furan-5-yl)methyl group.

TABLE 4-continued




Compound No.	Compound Structure
IV-29	
IV-30	
IV-31	

TABLE 4-continued




Compound No.	Compound Structure
IV-32	
IV-33	
IV-34	

TABLE 4-continued




Compound No.	Compound Structure
IV-35	 Chemical structure of compound IV-35: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group.
IV-36	 Chemical structure of compound IV-36: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group.
IV-37	 Chemical structure of compound IV-37: A complex molecule featuring a 1,2,4-triazole-thienopyridine core. The triazole ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group. The thiophene ring is substituted with a methyl group and a 4-(4-(2-(4-chloro-1-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-2-fluorophenyl)-1-hydroxy-4-phenylpiperidin-1-yl)methylpiperidin-1-yl group.

TABLE 4-continued





Compound No.	Compound Structure
IV-38	
IV-39	
IV-40	
IV-41	

TABLE 4-continued




Compound No.	Compound Structure
IV-42	 Chemical structure of compound IV-42. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-(methylthio)phenyl group. The triazole ring is connected via a methylene bridge to a 4-phenylpiperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-hydroxy-4-(methylthio)piperidine ring. This hydroxy-piperidine ring is connected via a methylene bridge to a 4-phenylpiperidine ring, which is in turn connected via a methylene bridge to a 4-phenyl-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid moiety.
IV-43	 Chemical structure of compound IV-43. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-(methylthio)phenyl group. The triazole ring is connected via a methylene bridge to a 4-phenylpiperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-hydroxy-4-(methylthio)piperidine ring. This hydroxy-piperidine ring is connected via a methylene bridge to a 4-phenylpiperidine ring, which is in turn connected via a methylene bridge to a 4-phenyl-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid moiety.
IV-44	 Chemical structure of compound IV-44. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-(methylthio)phenyl group. The triazole ring is connected via a methylene bridge to a 4-phenylpiperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-hydroxy-4-(methylthio)piperidine ring. This hydroxy-piperidine ring is connected via a methylene bridge to a 4-phenylpiperidine ring, which is in turn connected via a methylene bridge to a 4-phenyl-1,2,3,4-tetrahydronaphthalene-1-carboxylic acid moiety.

TABLE 4-continued




Compound No.	Compound Structure
IV-45	
IV-46	
IV-47	

TABLE 4-continued





Compound No.	Compound Structure
IV-48	
IV-49	
IV-50	
IV-51	

TABLE 4-continued




Compound No.	Compound Structure
IV-52	 Chemical structure of compound IV-52: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The molecule is linked via a 1,4-bis(piperidin-4-yl)butane chain to a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group, which is further substituted with a carboxylic acid group.
IV-53	 Chemical structure of compound IV-53: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The molecule is linked via a 1,4-bis(piperidin-4-yl)butane chain to a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group, which is further substituted with a carboxylic acid group.
IV-54	 Chemical structure of compound IV-54: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The triazole ring is also substituted with a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group. The molecule is linked via a 1,4-bis(piperidin-4-yl)butane chain to a 4-(4-(4-chlorophenyl)-1,2,3,4-tetrahydro-1H-benzo[7,8-b]pyrido[4,3-d]pyridin-5-yl)phenyl group, which is further substituted with a carboxylic acid group.

TABLE 4-continued

Compound No.	Compound Structure
IV-55	
IV-56	
IV-57	
IV-58	

TABLE 4-continued




Compound No.	Compound Structure
IV-59	 Chemical structure of compound IV-59: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a 4-phenyl-1,3-dioxane linker to a 4-(4-chlorophenyl)-1,3-dioxane linker, which is further connected to a 4-(4-chlorophenyl)-1,3-dioxane linker, and finally to a 4-(4-chlorophenyl)-1,3-dioxane linker with a carboxylic acid group.
IV-60	 Chemical structure of compound IV-60: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a 4-phenyl-1,3-dioxane linker to a 4-(4-chlorophenyl)-1,3-dioxane linker, which is further connected to a 4-(4-chlorophenyl)-1,3-dioxane linker, and finally to a 4-(4-chlorophenyl)-1,3-dioxane linker with a carboxylic acid group.
IV-61	 Chemical structure of compound IV-61: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a 4-phenyl-1,3-dioxane linker to a 4-(4-chlorophenyl)-1,3-dioxane linker, which is further connected to a 4-(4-chlorophenyl)-1,3-dioxane linker, and finally to a 4-(4-chlorophenyl)-1,3-dioxane linker with a carboxylic acid group.

TABLE 4-continued

[illegible]

TABLE 4-continued





Compound No.	Compound Structure
IV-65	
IV-66	
IV-67	
IV-68	

TABLE 4-continued

Compound No.	Compound Structure
IV-69	
IV-70	
IV-71	
IV-72	

[0788] In certain embodiments, the compound is a compound in Table 5, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound in Table 5.

TABLE 5




Compound No.	Compound Structure
V-1	
V-2	
V-3	

TABLE 5-continued




Compound No.	Compound Structure
V-4	 Chemical structure of compound V-4: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected via a piperidine ring to a morpholine ring, which is further connected via an ethyl chain to a 2,6-difluorophenyl ring. This 2,6-difluorophenyl ring is connected to a 4-hydroxyphenyl ring, which is part of a larger fused ring system including a benzofuran moiety.
V-5	 Chemical structure of compound V-5: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected via a piperidine ring to a morpholine ring, which is further connected via an ethyl chain to a 2,6-difluorophenyl ring. This 2,6-difluorophenyl ring is connected to a 4-hydroxyphenyl ring, which is part of a larger fused ring system including a benzofuran moiety.
V-6	 Chemical structure of compound V-6: A complex molecule featuring a 1,2,4-triazole ring system fused to a thiophene ring. The thiophene ring is substituted with a methyl group and a phenyl ring. The phenyl ring is connected via a piperidine ring to a morpholine ring, which is further connected via an ethyl chain to a 2,6-difluorophenyl ring. This 2,6-difluorophenyl ring is connected to a 4-hydroxyphenyl ring, which is part of a larger fused ring system including a benzofuran moiety.

TABLE 5-continued




Compound No.	Compound Structure
V-7	 Chemical structure of compound V-7: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This thiophene ring is connected via a double bond to a phenyl ring. The phenyl ring is further connected to a piperidine ring, which is linked to a 3,4-dihydroquinoline ring. The 3,4-dihydroquinoline ring is connected to a 2,3,5-trifluorophenyl ring, which is in turn connected to a 4-hydroxyphenyl ring.
V-8	 Chemical structure of compound V-8: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This thiophene ring is connected via a double bond to a phenyl ring. The phenyl ring is further connected to a piperidine ring, which is linked to a 3,4-dihydroquinoline ring. The 3,4-dihydroquinoline ring is connected to a 2,3,5-trifluorophenyl ring, which is in turn connected to a 4-hydroxyphenyl ring.
V-9	 Chemical structure of compound V-9: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is substituted with a methyl group. This thiophene ring is connected via a double bond to a phenyl ring. The phenyl ring is further connected to a piperidine ring, which is linked to a 3,4-dihydroquinoline ring. The 3,4-dihydroquinoline ring is connected to a 2,3,5-trifluorophenyl ring, which is in turn connected to a 4-hydroxyphenyl ring.

TABLE 5-continued




Compound No.	Compound Structure
V-10	 Chemical structure of compound V-10: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a cyclohexane ring, which is then linked to a 2,4,6-trifluorophenyl group. The structure also includes a 4-hydroxyphenyl group and a 2,4,6-trifluorophenyl group.
V-11	 Chemical structure of compound V-11: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a cyclohexane ring, which is then linked to a 2,4,6-trifluorophenyl group. The structure also includes a 4-hydroxyphenyl group and a 2,4,6-trifluorophenyl group.
V-12	 Chemical structure of compound V-12: A complex molecule featuring a 1,2,4-triazole ring fused to a thiophene ring, which is further substituted with a phenyl group. This phenyl group is connected via a piperidine ring to a cyclohexane ring, which is then linked to a 2,4,6-trifluorophenyl group. The structure also includes a 4-hydroxyphenyl group and a 2,4,6-trifluorophenyl group.

TABLE 5-continued





Compound No.	Compound Structure
V-13	 Chemical structure of compound V-13: A complex molecule featuring a thiazine-sulfone core on the left, connected via a biphenyl linker to a piperidine ring. This piperidine is further linked to another piperidine ring, which is connected to a biphenyl system. The second biphenyl system is substituted with a 2,4,6-trifluorophenyl group and a 2-hydroxy-3-methyl-4H-chromene group.
V-14	 Chemical structure of compound V-14: A complex molecule featuring a thiazine-sulfone core on the left, connected via a biphenyl linker to a piperidine ring. This piperidine is further linked to a carbonyl group, which is connected to a piperidine ring. The second piperidine ring is connected to a biphenyl system, which is substituted with a 2,4,6-trifluorophenyl group and a 2-hydroxy-3-methyl-4H-chromene group.
V-15	 Chemical structure of compound V-15: A complex molecule featuring a thiazine-sulfone core on the left, connected via a biphenyl linker to a piperidine ring. This piperidine is further linked to a carbonyl group, which is connected to a piperidine ring. The second piperidine ring is connected to a biphenyl system, which is substituted with a 2,4,6-trifluorophenyl group and a 2-hydroxy-3-methyl-4H-chromene group.
V-16	 Chemical structure of compound V-16: A complex molecule featuring a thiazine-sulfone core on the left, connected via a biphenyl linker to a piperidine ring. This piperidine is further linked to a piperidine ring, which is connected to a biphenyl system. The second biphenyl system is substituted with a 2,4,6-trifluorophenyl group and a 2-hydroxy-3-methyl-4H-chromene group.

TABLE 5-continued




Compound No.	Compound Structure
V-17	 Chemical structure of compound V-17. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group.
V-18	 Chemical structure of compound V-18. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group.
V-19	 Chemical structure of compound V-19. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group. The triazole ring is also substituted with a methyl group and a 4-(4-(4-(2,6-dimethyl-1,2,4-triazol-4-yl)phenoxy)-2,6-difluorophenyl)phenyl group.

TABLE 5-continued





Compound No.	Compound Structure
V-20	 Chemical structure of compound V-20. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. The triazole ring is connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring.
V-21	 Chemical structure of compound V-21. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. The triazole ring is connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring.
V-22	 Chemical structure of compound V-22. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. The triazole ring is connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring.
V-23	 Chemical structure of compound V-23. It features a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazo[5,1-b]thien-2-yl group. The triazole ring is connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring. This piperidine ring is further connected via a methylene bridge to a 4-((4-(2,6-difluoro-4-hydroxyphenyl)-2-methylchroman-7-yl)phenyl)piperidine ring.

TABLE 5-continued





Compound No.	Compound Structure
V-24	
V-25	
V-26	
V-27	

TABLE 5-continued





Compound No.	Compound Structure
V-28	
V-29	
V-30	
V-31	

TABLE 5-continued




Compound No.	Compound Structure
V-32	 Chemical structure of Compound V-32: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further substituted with a hydroxyl group and a 4-(4-hydroxy-7-methyl-2,3-difluoro-6,7-dihydro-2H-benzofuro[3,2-c]pyridin-5-yl)phenyl group.
V-33	 Chemical structure of Compound V-33: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further substituted with a hydroxyl group and a 4-(4-hydroxy-7-methyl-2,3-difluoro-6,7-dihydro-2H-benzofuro[3,2-c]pyridin-5-yl)phenyl group.
V-34	 Chemical structure of Compound V-34: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-methyl-5-methylthio-1H-imidazole ring. This is connected via a methylene bridge to a 4-phenylpiperidine ring. The piperidine ring is further substituted with a hydroxyl group and a 4-(4-hydroxy-7-methyl-2,3-difluoro-6,7-dihydro-2H-benzofuro[3,2-c]pyridin-5-yl)phenyl group.

TABLE 5-continued




Compound No.	Compound Structure
V-35	 Chemical structure of Compound V-35: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. This triazole is linked via a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group, which is further connected to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The structure also includes a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group.
V-36	 Chemical structure of Compound V-36: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. This triazole is linked via a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group, which is further connected to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The structure also includes a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group.
V-37	 Chemical structure of Compound V-37: A complex molecule featuring a 1,2,4-triazole ring substituted with a methyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. This triazole is linked via a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group, which is further connected to a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group. The structure also includes a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group and a 4-(4-methyl-5-(4-methyl-1,3,4-thiazol-2-yl)-1H-1,2,4-triazol-1-yl)phenyl group.

TABLE 5-continued

Compound No.	Compound Structure
V-38	
V-39	
V-40	

Synthetic Methods

[0789] Methods for preparing compounds described herein are illustrated in the following synthetic Schemes. The Schemes are given for the purpose of illustrating the invention, and are not intended to limit the scope or spirit of the invention. Starting materials shown in the Schemes can be obtained from commercial sources or can be prepared based on procedures described in the literature.


[0790] In the Schemes, it is understood by one skilled in the art of organic synthesis that the functionality present on various portions of the molecule should be compatible with the reagents and reactions proposed. Substituents not compatible with the reaction conditions will be apparent to one skilled in the art, and alternate methods are therefore indicated (for example, use of protecting groups or alternative reactions). Protecting group chemistry and strategy is well

known in the art, for example, as described in detail in "Protecting Groups in Organic Synthesis", T. W. Greene and P. G. M. Wuts, 3rd edition, John Wiley & Sons, 1999, the entire contents of which are hereby incorporated by reference.

[0791] The synthetic route illustrated in Scheme 1 is a general method for preparing heterobifunctional compounds D. Coupling compound A (a precursor of TPL, for example, a discrete compound that is a target protein ligand) with L' (a precursor to linker L, containing functionality for coupling to the precursors of both TPL and EPL) affords intermediate B (wherein L" is a precursor to linker L that contains functionality for coupling to the EPL precursor). Coupling intermediate B with compound C (a precursor of EPL) affords heterobifunctional compound D. Alternatively, the order of coupling compounds A and C to L' may be

reversed, such that L' is first coupled with compound C, before being coupled to compound A.

SCHEME 1.



[0792] The coupling of compound A with L', and the coupling of intermediate B with compound C, can be accomplished with a wide variety of strategies. For example, amide coupling conditions can be employed when compound A (or compound C) is to be attached at a modifiable nitrogen atom and L' (or L'') contains a carboxylic acid group, or vice versa (i.e. compound A contains a carboxylic acid group and L' contains a nucleophilic amine nitrogen atom). Alternatively, reductive amination conditions can be employed when compound A (or compound C) is to be attached at a modifiable nitrogen atom and L' (or L'') contains an aldehyde group, or vice versa. Alternatively, nucleophilic substitution conditions can be employed when compound A (or compound C) is to be attached at a modifiable carbon, oxygen, or nitrogen atom (where the carbon atom may be activated, for example, with a bromide or sulfonate) and L' (or L'') contains a suitable coupling partner (for example, an olefin for a Heck coupling, a trialkylstannane for a Stille coupling, or a boronic acid or boronate ester for a Suzuki coupling, Buchwald-Hartwig amination, or Chan-Lam coupling), or vice versa.

[0793] It is understood by one skilled in the art of organic synthesis that protecting group strategies may be employed as necessary, for example, if L' contains two of the same functional group that are to be selectively coupled to compound A and compound C. For example, L' may contain, for example, both an unprotected carboxylic acid for coupling to compound A, and a carboxylic acid group that is protected (for example, as a methyl or benzyl ester) during the coupling with compound A and subsequently deprotected (for example, via basic hydrolysis of a methyl ester or hydrogenolysis of a benzyl ester) prior to coupling with compound C.

II. Therapeutic Applications

[0794] The heterobifunctional compounds described herein, such as a compound of Formula I or II, or other compounds in Section I, provide therapeutic benefits to patients suffering from cancer. Accordingly, one aspect of the invention provides a method of treating cancer. The method comprises administering to a patient in need thereof

a therapeutically effective amount of a compound described herein, such as a compound of Formula I or II, or other compounds in Section I, to treat the cancer. In certain embodiments, the particular compound of Formula I or II is a compound defined by one of the embodiments described above. In certain embodiments, the compound is a compound of Formula I, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I. In certain embodiments, the compound is a compound of Formula I-A or I-B, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A or I-B. In certain embodiments, the particular compound of Formula I, I-A, or I-B is a compound defined by one of the embodiments described above (including, for example, a compound of Formula I-A-1, I-A-2, I-A-3, I-A-4, etc.).

Cancer

[0795] In certain embodiments, the cancer is ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland carcinoma, bile duct and gallbladder cancers, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia. In certain embodiments, the cancer is ovarian cancer. In certain embodiments, the cancer is endometrial cancer. In certain embodiments, the cancer is cervical cancer.

[0796] In certain embodiments, the cancer is breast cancer. In certain embodiments, the cancer is advanced breast cancer. In certain embodiments, the cancer is metastatic breast cancer. In certain embodiments, the cancer is ER+ breast cancer or ER+/HER2- breast cancer. In certain embodiments, the cancer is ER+ advanced breast cancer, ER+ metastatic breast cancer ER+/HER2- advanced breast cancer, or ER+/HER2- metastatic breast cancer. In certain embodiments, the cancer is ER+ breast cancer. In certain embodiments, the cancer is ER+ advanced breast cancer or ER+ metastatic breast cancer. In certain embodiments, the cancer is ER+/HER2- breast cancer. In certain embodiments, the cancer is ER+/HER2- advanced breast cancer or ER+/HER2- metastatic breast cancer. In certain embodiments, the cancer is sporadic breast cancer. In certain embodiments, the cancer is Cowden disease.

[0797] In certain embodiments, the cancer has a mutant estrogen receptor alpha protein. In certain embodiments, the cancer has an estrogen receptor alpha protein containing one or more of the following mutations: D538G, Y537S, and L536R.

[0798] In certain embodiments, the cancer is squamous cell cancer, lung cancer including small cell lung cancer, non-small cell lung cancer, vulval cancer, thyroid cancer, adenocarcinoma of the lung and squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastric or stomach cancer including gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney or renal cancer, prostate cancer, hepatic carcinoma, anal carcinoma, penile carcinoma, and head and neck cancer. In certain embodiments, the cancer is at least one selected from

the group consisting of ALL, T-lineage Acute lymphoblastic Leukemia (T-ALL), T-lineage lymphoblastic Lymphoma (T-LL), Peripheral T-cell lymphoma, Adult T-cell Leukemia, Pre-B ALL, Pre-B Lymphomas, Large B-cell Lymphoma, Burkitts Lymphoma, B-cell ALL, Philadelphia chromosome positive ALL, Philadelphia chromosome positive CML, lymphoma, leukemia, multiple myeloma myeloproliferative diseases, large B cell lymphoma, or B cell Lymphoma.

[0799] In certain embodiments, the cancer is a solid tumor or leukemia. In certain other embodiments, the cancer is colon cancer, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, lung cancer, leukemia, bladder cancer, stomach cancer, cervical cancer, testicular cancer, skin cancer, rectal cancer, thyroid cancer, kidney cancer, uterus cancer, esophagus cancer, liver cancer, an acoustic neuroma, oligodendroglioma, meningioma, neuroblastoma, or retinoblastoma. In certain other embodiments, the cancer is small cell lung cancer, non-small cell lung cancer, melanoma, cancer of the central nervous system tissue, brain cancer, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, or diffuse large B-Cell lymphoma. In certain other embodiments, the cancer is breast cancer, colon cancer, small-cell lung cancer, non-small cell lung cancer, prostate cancer, renal cancer, ovarian cancer, leukemia, melanoma, or cancer of the central nervous system tissue. In certain other embodiments, the cancer is colon cancer, small-cell lung cancer, non-small cell lung cancer, renal cancer, ovarian cancer, renal cancer, or melanoma.

[0800] In certain embodiments, the cancer is a fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, or hemangioblastoma.

[0801] In certain embodiments, the cancer is a neuroblastoma, meningioma, hemangiopericytoma, multiple brain metastase, glioblastoma multiforms, glioblastoma, brain stem glioma, poor prognosis malignant brain tumor, malignant glioma, anaplastic astrocytoma, anaplastic oligodendroglioma, neuroendocrine tumor, rectal adenocarcinoma, Dukes C & D colorectal cancer, unresectable colorectal carcinoma, metastatic hepatocellular carcinoma, Kaposi's sarcoma, karotype acute myeloblastic leukemia, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cutaneous T-Cell lymphoma, cutaneous B-Cell lymphoma, diffuse large B-Cell lymphoma, low grade follicular lymphoma, metastatic melanoma, localized melanoma, malignant mesothelioma, malignant pleural effusion mesothelioma syndrome, peritoneal carcinoma, papillary serous carcinoma, gynecologic sarcoma, soft tissue sarcoma, scleroderma, cutaneous vasculitis, Langerhans cell histiocytosis, leiomyosarcoma, fibrodysplasia ossificans progressive, hormone refractory prostate cancer, resected high-risk soft tissue sarcoma, unresectable hepatocellular carcinoma, Waldenstrom's

macroglobulinemia, smoldering myeloma, indolent myeloma, fallopian tube cancer, androgen independent prostate cancer, androgen dependent stage IV non-metastatic prostate cancer, hormone-insensitive prostate cancer, chemotherapy-insensitive prostate cancer, papillary thyroid carcinoma, follicular thyroid carcinoma, medullary thyroid carcinoma, or leiomyoma.

[0802] In certain embodiments, the cancer is bone cancer, pancreatic cancer, skin cancer, cancer of the head or neck, cutaneous or intraocular melanoma, ovarian cancer, colon cancer, rectal cancer, cancer of the anal region, stomach cancer, gastrointestinal (gastric, colorectal, and duodenal), uterine cancer, carcinoma of the fallopian tubes, carcinoma of the endometrium, carcinoma of the cervix, carcinoma of the vagina, carcinoma of the vulva, Hodgkin's Disease, cancer of the esophagus, cancer of the small intestine, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, cancer of the adrenal gland, sarcoma of soft tissue, cancer of the urethra, cancer of the penis, prostate cancer, testicular cancer, chronic or acute leukemia, chronic myeloid leukemia, lymphocytic lymphomas, cancer of the bladder, cancer of the kidney or ureter, renal cell carcinoma, carcinoma of the renal pelvis, non-Hodgkins's lymphoma, spinal axis tumors, brain stem glioma, pituitary adenoma, adrenocortical cancer, gall bladder cancer, multiple myeloma, cholangiocarcinoma, fibrosarcoma, neuroblastoma, retinoblastoma, or a combination of one or more of the foregoing cancers.

[0803] In certain embodiments, the cancer is hepatocellular carcinoma, ovarian cancer, ovarian epithelial cancer, or fallopian tube cancer; papillary serous cystadenocarcinoma or uterine papillary serous carcinoma (UPSC); prostate cancer; testicular cancer; gallbladder cancer; hepatocholangiocarcinoma; soft tissue and bone synovial sarcoma; rhabdomyosarcoma; osteosarcoma; chondrosarcoma; Ewing sarcoma; anaplastic thyroid cancer; adrenocortical adenoma; pancreatic cancer; pancreatic ductal carcinoma or pancreatic adenocarcinoma; gastrointestinal/stomach (GIST) cancer; lymphoma; squamous cell carcinoma of the head and neck (SCCHN); salivary gland cancer; glioma, or brain cancer; neurofibromatosis-1 associated malignant peripheral nerve sheath tumors (MPNST); Waldenstrom's macroglobulinemia; or medulloblastoma.

[0804] In certain embodiments, the cancer is hepatocellular carcinoma (HCC), hepatoblastoma, colon cancer, rectal cancer, ovarian cancer, ovarian epithelial cancer, fallopian tube cancer, papillary serous cystadenocarcinoma, uterine papillary serous carcinoma (UPSC), hepatocholangiocarcinoma, soft tissue and bone synovial sarcoma, rhabdomyosarcoma, osteosarcoma, anaplastic thyroid cancer, adrenocortical adenoma, pancreatic cancer, pancreatic ductal carcinoma, pancreatic adenocarcinoma, glioma, neurofibromatosis-1 associated malignant peripheral nerve sheath tumors (MPNST), Waldenstrom's macroglobulinemia, or medulloblastoma.

[0805] In certain embodiments, the cancer is a solid tumor, such as a sarcoma, carcinoma, or lymphoma. In certain embodiments, the cancer is kidney cancer; hepatocellular carcinoma (HCC) or hepatoblastoma, or liver cancer; melanoma; breast cancer; colorectal carcinoma, or colorectal cancer; colon cancer; rectal cancer; anal cancer; lung cancer, such as non-small cell lung cancer (NSCLC) or small cell lung cancer (SCLC); ovarian cancer, ovarian epithelial cancer, ovarian carcinoma, or fallopian tube cancer; papillary

serous cystadenocarcinoma or uterine papillary serous carcinoma (UPSC); prostate cancer; testicular cancer; gallbladder cancer; hepatocholangiocarcinoma; soft tissue and bone synovial sarcoma; rhabdomyosarcoma; osteosarcoma; chondrosarcoma; Ewing sarcoma; anaplastic thyroid cancer; adrenocortical carcinoma; pancreatic cancer; pancreatic ductal carcinoma or pancreatic adenocarcinoma; gastrointestinal/stomach (GIST) cancer; lymphoma; squamous cell carcinoma of the head and neck (SCCHN); salivary gland cancer; glioma, or brain cancer; neurofibromatosis-1 associated malignant peripheral nerve sheath tumors (MPNST); Waldenstrom's macroglobulinemia; or medulloblastoma.

[0806] In certain embodiments, the cancer is renal cell carcinoma, hepatocellular carcinoma (HCC), hepatoblastoma, colorectal carcinoma, colorectal cancer, colon cancer, rectal cancer, anal cancer, ovarian cancer, ovarian epithelial cancer, ovarian carcinoma, fallopian tube cancer, papillary serous cystadenocarcinoma, uterine papillary serous carcinoma (UPSC), hepatocholangiocarcinoma, soft tissue and bone synovial sarcoma, rhabdomyosarcoma, osteosarcoma, chondrosarcoma, anaplastic thyroid cancer, adrenocortical carcinoma, pancreatic cancer, pancreatic ductal carcinoma, pancreatic adenocarcinoma, glioma, brain cancer, neurofibromatosis-1 associated malignant peripheral nerve sheath tumors (MPNST), Waldenstrom's macroglobulinemia, or medulloblastoma.

[0807] In certain embodiments, the cancer is hepatocellular carcinoma (HCC), hepatoblastoma, colon cancer, rectal cancer, ovarian cancer, ovarian epithelial cancer, ovarian carcinoma, fallopian tube cancer, papillary serous cystadenocarcinoma, uterine papillary serous carcinoma (UPSC), hepatocholangiocarcinoma, soft tissue and bone synovial sarcoma, rhabdomyosarcoma, osteosarcoma, anaplastic thyroid cancer, adrenocortical carcinoma, pancreatic cancer, pancreatic ductal carcinoma, pancreatic adenocarcinoma, glioma, neurofibromatosis-1 associated malignant peripheral nerve sheath tumors (MPNST), Waldenstrom's macroglobulinemia, or medulloblastoma.

[0808] In certain embodiments, the cancer is hepatocellular carcinoma (HCC). In some embodiments, the cancer is hepatoblastoma. In some embodiments, the cancer is colon cancer. In some embodiments, the cancer is rectal cancer. In some embodiments, the cancer is ovarian cancer, or ovarian carcinoma. In some embodiments, the cancer is ovarian epithelial cancer. In some embodiments, the cancer is fallopian tube cancer. In some embodiments, the cancer is papillary serous cystadenocarcinoma. In some embodiments, the cancer is uterine papillary serous carcinoma (UPSC). In some embodiments, the cancer is hepatocholangiocarcinoma. In some embodiments, the cancer is soft tissue and bone synovial sarcoma. In some embodiments, the cancer is rhabdomyosarcoma. In some embodiments, the cancer is osteosarcoma. In some embodiments, the cancer is anaplastic thyroid cancer. In some embodiments, the cancer is adrenocortical carcinoma. In some embodiments, the cancer is pancreatic cancer, or pancreatic ductal carcinoma. In some embodiments, the cancer is pancreatic adenocarcinoma. In some embodiments, the cancer is glioma. In some embodiments, the cancer is malignant peripheral nerve sheath tumors (MPNST). In some embodiments, the cancer is neurofibromatosis-1 associated MPNST. In some embodiments, the cancer is Waldenstrom's macroglobulinemia. In some embodiments, the cancer is medulloblastoma.

Causing Death of Cancer Cell

[0809] Another aspect of the invention provides a method of causing death of a cancer cell. The method comprises contacting a cancer cell with an effective amount of a compound described herein, such as a compound of Formula I or II, or other compounds in Section I, to cause death of the cancer cell. In certain embodiments, the compound is a compound of Formula I, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I. In certain embodiments, the compound is a compound of Formula I-A or I-B, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A or I-B. In certain embodiments, the particular compound of Formula I, I-A, or I-B is a compound defined by one of the embodiments described above (including, for example, a compound of Formula I-A-1, I-A-2, I-A-3, I-A-4, etc.).

[0810] In certain embodiments, the cancer cell is selected from ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland carcinoma, bile duct and gallbladder cancers, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia cell. In certain embodiments, the cancer cell is a breast cancer cell. In certain embodiments, the cancer cell is a uterine cancer cell. In certain embodiments, the cancer cell is one or more of the cancers recited in the section above entitled "Cancer."

Diseases Mediated by the Estrogen Receptor

[0811] Another aspect of the invention provides a method of treating a disease or condition mediated by the estrogen receptor. The method comprises administering to a patient in need thereof a therapeutically effective amount of a compound described herein, such as a compound of Formula I or II, or other compounds in Section I, to treat the disease or condition. In certain embodiments, the particular compound of Formula I or II is a compound defined by one of the embodiments described above, including, for example, a compound of Formula I, I-A, I-A-1, I-A-2, I-A-3, I-A-4, etc.). In certain embodiments, the disease or condition is mediated by the estrogen receptor alpha protein. In certain embodiments, the patient has a mutant estrogen receptor alpha protein. In certain embodiments, the patient has an estrogen receptor alpha protein containing one or more of the following mutations: D538G, Y537S, and L536R. In certain embodiments, the disease or condition is endometriosis.

Combination Therapies

[0812] The compounds useful within the methods of the invention may be used in combination with one or more additional therapeutic agents useful for treating any disease contemplated herein. These additional therapeutic agents may comprise compounds that are commercially available or synthetically accessible to those skilled in the art. These additional therapeutic agents are known to treat, prevent, or reduce the symptoms, of a disease or disorder contemplated herein.

[0813] Accordingly, in certain embodiments, the method further comprises administering to the subject an additional therapeutic agent that treats the disease contemplated herein.

[0814] In certain embodiments, administering the compound of the invention to the subject allows for administering a lower dose of the additional therapeutic agent as compared to the dose of the additional therapeutic agent alone that is required to achieve similar results in treating the disease contemplated herein. For example, in certain embodiments, the compound of the invention enhances the therapeutic activity of the additional therapeutic compound, thereby allowing for a lower dose of the additional therapeutic compound to provide the same effect.

[0815] A synergistic effect may be calculated, for example, using suitable methods such as, for example, the Sigmoid- E_{max} equation (Holford & Scheiner, 1981, Clin. Pharmacokinet. 6:429-453), the equation of Loewe additivity (Loewe & Muischnek, 1926, Arch. Exp. Pathol Pharmacol. 114:313-326) and the median-effect equation (Chou & Talalay, 1984, Adv. Enzyme Regul. 22:27-55). Each equation referred to above may be applied to experimental data to generate a corresponding graph to aid in assessing the effects of the drug combination. The corresponding graphs associated with the equations referred to above are the concentration-effect curve, isobologram curve and combination index curve, respectively.

[0816] In certain embodiments, the compound of the invention and the therapeutic agent are co-administered to the subject. In other embodiments, the compound of the invention and the therapeutic agent are coformulated and co-administered to the subject.

[0817] In certain embodiments, the compound is administered in combination with a second therapeutic agent having activity against cancer.

[0818] Accordingly, one aspect of the invention provides a method for treating cancer. The method comprises administering to a patient in need thereof a therapeutically effective amount of a compound described herein, such as a compound of Formula I or II, or other compounds in Section I, in combination with a second therapeutic agent having activity against cancer, in order to treat the cancer. In certain embodiments, the compound is a compound of Formula I, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I. In certain embodiments, the compound is a compound of Formula I-A or I-B, or a pharmaceutically acceptable salt thereof. In certain embodiments, the compound is a compound of Formula I-A or I-B. In certain embodiments, the particular compound of Formula I, I-A, or I-B is a compound defined by one of the embodiments described above (including, for example, a compound of Formula I-A-1, I-A-2, I-A-3, I-A-4, etc.).

[0819] In certain embodiments, the method further comprises administering a third therapeutic agent having activity against cancer. In certain embodiments, the method further comprises administering a fourth therapeutic agent having activity against cancer.

[0820] In certain embodiments, the second therapeutic agent is mitomycin, tretinoin, ribomustin, gemcitabine, vincristine, etoposide, cladribine, mitobronitol, methotrexate, doxorubicin, carboquone, pentostatin, nitracrine, zinzotatin, cetorelix, letrozole, raltitrexed, daunorubicin, fadrozole, fotemustine, thymalfasin, sobuzoxane, nedaplatin, cytarabine, bicalutamide, vinorelbine, vesnarinone, amino-

glutethimide, amsacrine, proglumide, elliptinium acetate, ketanserin, doxifluridine, etretinate, isotretinoin, streptozocin, nimustine, vindesine, flutamide, drogenil, butocin, carmofur, razoxane, sizofilan, carboplatin, mitolactol, tegafur, ifosfamide, prednimustine, picibanil, levamisole, teniposide, improsulfan, enocitabine, lisuride, oxymetholone, tamoxifen, progesterone, mepitiostane, epitioctanol, formestane, interferon-alpha, interferon-2 alpha, interferon-beta, interferon-gamma, colony stimulating factor-1, colony stimulating factor-2, denileukin diftitox, interleukin-2, and leutinizing hormone releasing factor.

[0821] In certain embodiments, the second therapeutic agent is an mTOR inhibitor, which inhibits cell proliferation, angiogenesis and glucose uptake. Approved mTOR inhibitors useful in the present invention include everolimus (Afinitor®, Novartis); temsirolimus (Torisel®, Pfizer); and sirolimus (Rapamune®, Pfizer).

[0822] In certain embodiments, the second therapeutic agent is a Poly ADP ribose polymerase (PARP) inhibitor. Approved PARP inhibitors useful in the present invention include olaparib (Lynparza®, AstraZeneca); rucaparib (Rubraca®, Clovis Oncology); and niraparib (Zejula®, Tesaro). Other PARP inhibitors being studied which may be used in the present invention include talazoparib (MDV3800/BMN 673/LT00673, Medivation/Pfizer/Biomarin); veliparib (ABT-888, AbbVie); and BGB-290 (BeiGene, Inc.).

[0823] In certain embodiments, the second therapeutic agent is a phosphatidylinositol 3 kinase (PI3K) inhibitor. Approved PI3K inhibitors useful in the present invention include idelalisib (Zydelig®, Gilead). Other PI3K inhibitors being studied which may be used in the present invention include alpelisib (BYL719, Novartis); taselisib (GDC-0032, Genentech/Roche); pictilisib (GDC-0941, Genentech/Roche); copanlisib (BAY806946, Bayer); duvelisib (formerly IPI-145, Infinity Pharmaceuticals); PQR309 (Piquar Therapeutics, Switzerland); and TGR1202 (formerly RP5230, TG Therapeutics).

[0824] In certain embodiments, the second therapeutic agent is a proteasome inhibitor. Approved proteasome inhibitors useful in the present invention include bortezomib (Velcade®, Takeda); carfilzomib (Kyprolis®, Amgen); and ixazomib (Ninlaro®, Takeda).

[0825] In certain embodiments, the second therapeutic agent is a histone deacetylase (HDAC) inhibitor. Approved HDAC inhibitors useful in the present invention include vorinostat (Zolinza®, Merck); romidepsin (Istodax®, Celgene); panobinostat (Farydak®, Novartis); and belinostat (Beleodaq®, Spectrum Pharmaceuticals). Other HDAC inhibitors being studied which may be used in the present invention include entinostat (SNDX-275, Syndax Pharmaceuticals) (NCT00866333); and chidamide (Epidaza®, HBI-8000, Chipscreen Biosciences, China).

[0826] In certain embodiments, the second therapeutic agent is a CDK inhibitor, such as a CDK 4/6 inhibitor. Approved CDK 4/6 inhibitors useful in the present invention include palbociclib (Ibrance®, Pfizer); and ribociclib (Kisqali®, Novartis). Other CDK 4/6 inhibitors being studied which may be used in the present invention include abemaciclib (Ly2835219, Eli Lilly); and trilaciclib (G1T28, G1 Therapeutics).

[0827] In certain embodiments, the second therapeutic agent is an indoleamine (2,3)-dioxygenase (IDO) inhibitor. IDO inhibitors being studied which may be used in the present invention include epacadostat (INCB024360,

Incyte); indoximod (NLG-8189, NewLink Genetics Corporation); capmanitib (INC280, Novartis); GDC-0919 (Genentech/Roche); PF-06840003 (Pfizer); BMS:F001287 (Bristol-Myers Squibb); Phy906/KD108 (Phytoceutica); and an enzyme that breaks down kynurenine (Kynase, Kyn Therapeutics).

[0828] In certain embodiments, the second therapeutic agent is a growth factor antagonist, such as an antagonist of platelet-derived growth factor (PDGF), or epidermal growth factor (EGF) or its receptor (EGFR). Approved PDGF antagonists which may be used in the present invention include olaratumab (Lartruvo®; Eli Lilly). Approved EGFR antagonists which may be used in the present invention include cetuximab (Erbix®; Eli Lilly); necitumumab (Portrazza®, Eli Lilly), panitumumab (Vectibix®, Amgen); and osimertinib (targeting activated EGFR, Tagrisso®, Astra-Zeneca).

[0829] In certain embodiments, the second therapeutic agent is an aromatase inhibitor. Approved aromatase inhibitors which may be used in the present invention include exemestane (Aromasin®, Pfizer); anastrozole (Arimidex®, AstraZeneca) and letrozole (Femara®, Novartis).

[0830] In certain embodiments, the second therapeutic agent is an antagonist of the hedgehog pathway. Approved hedgehog pathway inhibitors which may be used in the present invention include sonidegib (Odomzo®, Sun Pharmaceuticals); and vismodegib (Erivedge®, Genentech), both for treatment of basal cell carcinoma.

[0831] In certain embodiments, the second therapeutic agent is a folic acid inhibitor. Approved folic acid inhibitors useful in the present invention include pemetrexed (Alimta®, Eli Lilly).

[0832] In certain embodiments, the second therapeutic agent is a CC chemokine receptor 4 (CCR4) inhibitor. CCR4 inhibitors being studied that may be useful in the present invention include mogamulizumab (Poteligeo®, Kyowa Hakko Kirin, Japan).

[0833] In certain embodiments, the second therapeutic agent is an isocitrate dehydrogenase (IDH) inhibitor. IDH inhibitors being studied which may be used in the present invention include AG120 (Celgene; NCT02677922); AG221 (Celgene, NCT02677922; NCT02577406); BAY1436032 (Bayer, NCT02746081); IDH305 (Novartis, NCT02987010).

[0834] In certain embodiments, the second therapeutic agent is an arginase inhibitor. Arginase inhibitors being studied which may be used in the present invention include AEB1102 (pegylated recombinant arginase, Aeglea Biotherapeutics), which is being studied in Phase 1 clinical trials for acute myeloid leukemia and myelodysplastic syndrome (NCT02732184) and solid tumors (NCT02561234); and CB-1158 (Calithera Biosciences).

[0835] In certain embodiments, the second therapeutic agent is a glutaminase inhibitor. Glutaminase inhibitors being studied which may be used in the present invention include CB-839 (Calithera Biosciences).

[0836] In certain embodiments, the second therapeutic agent is an antibody that binds to tumor antigens, that is, proteins expressed on the cell surface of tumor cells. Approved antibodies that bind to tumor antigens which may be used in the present invention include rituximab (Rituxan®, Genentech/BiogenIdec); ofatumumab (anti-CD20, Arzerra®, GlaxoSmithKline); obinutuzumab (anti-CD20, Gazyva®, Genentech); ibritumomab (anti-CD20 and

Yttrium-90, Zevalin®, Spectrum Pharmaceuticals); daratumumab (anti-CD38, Darzalex®, Janssen Biotech), dinutuximab (anti-glycolipid GD2, Unituxin®, United Therapeutics); trastuzumab (anti-HER2, Herceptin®, Genentech); ado-trastuzumab emtansine (anti-HER2, fused to emtansine, Kadcyla®, Genentech); and pertuzumab (anti-HER2, Perjeta®, Genentech); and brentuximab vedotin (anti-CD30-drug conjugate, Adcetris®, Seattle Genetics).

[0837] In certain embodiments, the second therapeutic agent is a topoisomerase inhibitor. Approved topoisomerase inhibitors useful in the present invention include irinotecan (Onivyde®, Merrimack Pharmaceuticals); topotecan (Hycamtin®, GlaxoSmithKline). Topoisomerase inhibitors being studied which may be used in the present invention include pixantrone (Pixuvri®, CTI Biopharma).

[0838] In certain embodiments, the second therapeutic agent is a nucleoside inhibitor, or other therapeutic that interfere with normal DNA synthesis, protein synthesis, cell replication, or will otherwise inhibit rapidly proliferating cells. Such nucleoside inhibitors or other therapeutics include trabectedin (guanidine alkylating agent, Yondelis®, Janssen Oncology), mechlorethamine (alkylating agent, Valchlor®, Aktelion Pharmaceuticals); vincristine (Oncovin®, Eli Lilly; Vincasar®, Teva Pharmaceuticals; Marqibo®, Talon Therapeutics); temozolomide (prodrug to alkylating agent 5-(3-methyltriazene-1-yl)-imidazole-4-carboxamide (MTIC) Temodar®, Merck); cytarabine injection (ara-C, antimetabolic cytidine analog, Pfizer); lomustine (alkylating agent, CeeNU®, Bristol-Myers Squibb; Gleostine®, Next-Source Biotechnology); azacitidine (pyrimidine nucleoside analog of cytidine, Vidaza®, Celgene); omacetaxine mepe-succinate (cephalotaxine ester) (protein synthesis inhibitor, Synribo®; Teva Pharmaceuticals); asparaginase *Erwinia chrysanthemi* (enzyme for depletion of asparagine, Elspar®, Lundbeck; Erwinaze®, EUSA Pharma); eribulin mesylate (microtubule inhibitor, tubulin-based antimetotic, Halaven®, Eisai); cabazitaxel (microtubule inhibitor, tubulin-based antimetotic, Jevtana®, Sanofi-Aventis); capacetrine (thymidylate synthase inhibitor, Xeloda®, Genentech); bendamustine (bifunctional mechlorethamine derivative, believed to form interstrand DNA cross-links, Treanda®, Cephalon/Teva); ixabepilone (semi-synthetic analog of epothilone B, microtubule inhibitor, tubulin-based antimetotic, Ixempra®, Bristol-Myers Squibb); nelarabine (prodrug of deoxy-guanosine analog, nucleoside metabolic inhibitor, Arranon®, Novartis); clorafabine (prodrug of ribonucleotide reductase inhibitor, competitive inhibitor of deoxycytidine, Clolar®, Sanofi-Aventis); and trifluridine and tipiracil (thymidine-based nucleoside analog and thymidine phosphorylase inhibitor, Lonsurf®, Taiho Oncology).

[0839] In certain embodiments, the second therapeutic agent is a platinum-based therapeutic, also referred to as platins. Platins cause cross-linking of DNA, such that they inhibit DNA repair and/or DNA synthesis, mostly in rapidly reproducing cells, such as cancer cells. Approved platinum-based therapeutics which may be used in the present invention include cisplatin (Platinol®, Bristol-Myers Squibb); carboplatin (Paraplatin®, Bristol-Myers Squibb; also, Teva; Pfizer); oxaliplatin (Eloxitin® Sanofi-Aventis); and nedaplatin (Aqupla®, Shionogi). Other platinum-based therapeutics which have undergone clinical testing and may be used in the present invention include picoplatin (Poniard Pharmaceuticals); and satraplatin (JM-216, Agennix).

[0840] In certain embodiments, the second therapeutic agent is a taxane compound, which causes disruption of microtubules, which are essential for cell division. Approved taxane compounds which may be used in the present invention include paclitaxel (Taxol®, Bristol-Myers Squibb), docetaxel (Taxotere®, Sanofi-Aventis; Docefrez®, Sun Pharmaceutical), albumin-bound paclitaxel (Abraxane®; Abraxis/Celgene), and cabazitaxel (Jevtana®, Sanofi-Aventis). Other taxane compounds which have undergone clinical testing and may be used in the present invention include SID530 (SK Chemicals, Co.) (NCT00931008).

[0841] In certain embodiments, the second therapeutic agent is an inhibitor of anti-apoptotic proteins, such as BCL-2. Approved anti-apoptotics which may be used in the present invention include venetoclax (Venclexta®, AbbVie/Genentech); and blinatumomab (Blincyto®, Amgen). Other therapeutic agents targeting apoptotic proteins which have undergone clinical testing and may be used in the present invention include navitoclax (ABT-263, Abbott), a BCL-2 inhibitor (NCT02079740).

[0842] In certain embodiments, the second therapeutic agent is a selective estrogen receptor modulator (SERM), which interferes with the synthesis or activity of estrogens. Approved SERMs useful in the present invention include raloxifene (Evista®, Eli Lilly).

[0843] In certain embodiments, the second therapeutic agent is an inhibitor of interaction between the two primary p53 suppressor proteins, MDMX and MDM2. Inhibitors of p53 suppression proteins being studied which may be used in the present invention include ALRN-6924 (Aileron), a stapled peptide that equipotently binds to and disrupts the interaction of MDMX and MDM2 with p53. ALRN-6924 is currently being evaluated in clinical trials for the treatment of AML, advanced myelodysplastic syndrome (MDS) and peripheral T-cell lymphoma (PTCL) (NCT02909972; NCT02264613).

[0844] In certain embodiments, the second therapeutic agent is an inhibitor of transforming growth factor-beta (TGF-beta or TGFβ). Inhibitors of TGF-beta proteins being studied which may be used in the present invention include NIS793 (Novartis), an anti-TGF-beta antibody being tested in the clinic for treatment of various cancers, including breast, lung, hepatocellular, colorectal, pancreatic, prostate and renal cancer (NCT 02947165). In some embodiments, the inhibitor of TGF-beta proteins is fresolimumab (GC1008; Sanofi-Genzyme), which is being studied for melanoma (NCT00923169); renal cell carcinoma (NCT00356460); and non-small cell lung cancer (NCT02581787). Additionally, in some embodiments, the additional therapeutic agent is a TGF-beta trap, such as described in Connolly et al. (2012) *Int'l J. Biological Sciences* 8:964-978. One therapeutic compound currently in clinical trials for treatment of solid tumors is M7824 (Merck KGaA—formerly MSB0011459X), which is a bispecific, anti-PD-L1/TGFβ trap compound (NCT02699515); and (NCT02517398). M7824 is comprised of a fully human IgG1 antibody against PD-L1 fused to the extracellular domain of human TGF-beta receptor II, which functions as a TGFβ “trap.”

[0845] In certain embodiments, the second therapeutic agent is a cancer vaccine. In some embodiments, the cancer vaccine is selected from sipuleucel-T (Provenge®, Dendreon/Valeant Pharmaceuticals), which has been approved for treatment of asymptomatic, or minimally symptomatic

metastatic castrate-resistant (hormone-refractory) prostate cancer; and talimogene laherparepvec (Imlygic®, BioVex/Amgen, previously known as T-VEC), a genetically modified oncolytic viral therapy approved for treatment of unresectable cutaneous, subcutaneous and nodal lesions in melanoma. In some embodiments, the additional therapeutic agent is selected from an oncolytic viral therapy such as pexastimogene devacirepvec (PexaVec/JX-594, SillaJen/formerly Jennerex Biotherapeutics), a thymidine kinase-(TK-) deficient vaccinia virus engineered to express GM-CSF, for hepatocellular carcinoma (NCT02562755) and melanoma (NCT00429312); pelareorep (Reolysin®, Oncolytics Biotech), a variant of respiratory enteric orphan virus (reovirus) which does not replicate in cells that are not RAS-activated, in numerous cancers, including colorectal cancer (NCT01622543); prostate cancer (NCT01619813); head and neck squamous cell cancer (NCT01166542); pancreatic adenocarcinoma (NCT00998322); and non-small cell lung cancer (NSCLC) (NCT 00861627); enadenotucirev (NG-348, PsiOxus, formerly known as ColoAd1), an adenovirus engineered to express a full length CD80 and an antibody fragment specific for the T-cell receptor CD3 protein, in ovarian cancer (NCT02028117); metastatic or advanced epithelial tumors such as in colorectal cancer, bladder cancer, head and neck squamous cell carcinoma and salivary gland cancer (NCT02636036); ONCOS-102 (Targovax/formerly Oncos), an adenovirus engineered to express GM-CSF, in melanoma (NCT03003676); and peritoneal disease, colorectal cancer or ovarian cancer (NCT02963831); GLV-ONC1 (GLV-1h68/GLV-1h153, Genelux GmbH), vaccinia viruses engineered to express beta-galactosidase (beta-gal)/beta-glucuronidase or beta-gal/human sodium iodide symporter (hNIS), respectively, were studied in peritoneal carcinomatosis (NCT01443260); fallopian tube cancer, ovarian cancer (NCT 02759588); or CG0070 (Cold Genesys), an adenovirus engineered to express GM-CSF, in bladder cancer (NCT02365818).

[0846] In certain embodiments, the second therapeutic agent is an immune checkpoint inhibitor selected from a PD-1 antagonist, a PD-L1 antagonist, or a CTLA-4 antagonist. In some embodiments, a compound disclosed herein or a pharmaceutically acceptable salt thereof is administered in combination with nivolumab (anti-PD-1 antibody, Opdivo®, Bristol-Myers Squibb); pembrolizumab (anti-PD-1 antibody, Keytruda®, Merck); ipilimumab (anti-CTLA-4 antibody, Yervoy®, Bristol-Myers Squibb); durvalumab (anti-PD-L1 antibody, Imfinzi®, AstraZeneca); or atezolizumab (anti-PD-L1 antibody, Tecentriq®, Genentech). Other immune checkpoint inhibitors suitable for use in the present invention include REGN2810 (Regeneron), an anti-PD-1 antibody tested in patients with basal cell carcinoma (NCT03132636); NSCLC (NCT03088540); cutaneous squamous cell carcinoma (NCT02760498); lymphoma (NCT02651662); and melanoma (NCT03002376); pidilizumab (CureTech), also known as CT-011, an antibody that binds to PD-1, in clinical trials for diffuse large B-cell lymphoma and multiple myeloma; avelumab (Bavencio®, Pfizer/Merck KGaA), also known as MSB0010718C), a fully human IgG1 anti-PD-L1 antibody, in clinical trials for non-small cell lung cancer, Merkel cell carcinoma, mesothelioma, solid tumors, renal cancer, ovarian cancer, bladder cancer, head and neck cancer, and gastric cancer; and PDR001 (Novartis), an inhibitory antibody that binds to PD-1, in clinical trials for non-small cell lung cancer,

melanoma, triple negative breast cancer and advanced or metastatic solid tumors. Tremelimumab (CP-675,206; AstraZeneca) is a fully human monoclonal antibody against CTLA-4 that has been studied in clinical trials for a number of indications, including: mesothelioma, colorectal cancer, kidney cancer, breast cancer, lung cancer and non-small cell lung cancer, pancreatic ductal adenocarcinoma, pancreatic cancer, germ cell cancer, squamous cell cancer of the head and neck, hepatocellular carcinoma, prostate cancer, endometrial cancer, metastatic cancer in the liver, liver cancer, large B-cell lymphoma, ovarian cancer, cervical cancer, metastatic anaplastic thyroid cancer, urothelial cancer, fallopian tube cancer, multiple myeloma, bladder cancer, soft tissue sarcoma, and melanoma. AGEN-1884 (Agenus) is an anti-CTLA4 antibody that is being studied in Phase 1 clinical trials for advanced solid tumors (NCT02694822).

[0847] In certain embodiments, the third therapeutic agent is one of the second therapeutic agents described above. In certain embodiments, the fourth therapeutic agent is one of the second therapeutic agents described above.

[0848] Another aspect of the invention provides a compound described herein (such as a compound of Formula I, Formula II, or other compounds in Section I) for use in treating a medical disease, such a disease described herein (e.g., cancer).

[0849] Another aspect of the invention provides for the use of a compound described herein (such as a compound of Formula I, Formula II, or other compounds in Section I) in the manufacture of a medicament. In certain embodiments, the medicament is for treating a disease described herein, such as cancer.

[0850] Another aspect of the invention provides for the use of a compound described herein (such as a compound of Formula I, Formula II, or other compounds in Section I) for treating a medical disease, such a disease described herein (e.g., cancer).

III. Pharmaceutical Compositions and Dosing Considerations

[0851] As indicated above, the invention provides pharmaceutical compositions, which comprise a therapeutically effective amount of one or more of the compounds described above, formulated together with one or more pharmaceutically acceptable carriers (additives) and/or diluents. The pharmaceutical compositions may be specially formulated for administration in solid or liquid form, including those adapted for the following: (1) oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; (2) parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; (3) topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin; (4) intravaginally or intrarectally, for example, as a pessary, cream or foam; (5) sublingually; (6) ocularly; (7) transdermally; or (8) nasally. In certain embodiments, the invention provides a pharmaceutical composition comprising a compound described herein (e.g., a compound of Formula I) and a pharmaceutically acceptable carrier.

[0852] The phrase “therapeutically effective amount” as used herein means that amount of a compound, material, or composition comprising a compound of the present invention which is effective for producing some desired therapeutic effect in at least a sub-population of cells in an animal at a reasonable benefit/risk ratio applicable to any medical treatment.

[0853] The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

[0854] Wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the compositions.

[0855] Examples of pharmaceutically acceptable antioxidants include: (1) water soluble antioxidants, such as ascorbic acid, cysteine hydrochloride, sodium bisulfate, sodium metabisulfite, sodium sulfite and the like; (2) oil-soluble antioxidants, such as ascorbyl palmitate, butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT), lecithin, propyl gallate, alpha-tocopherol, and the like; and (3) metal chelating agents, such as citric acid, ethylenediamine tetraacetic acid (EDTA), sorbitol, tartaric acid, phosphoric acid, and the like.

[0856] Formulations of the present invention include those suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will vary depending upon the host being treated, the particular mode of administration. The amount of active ingredient which can be combined with a carrier material to produce a single dosage form will generally be that amount of the compound which produces a therapeutic effect. Generally, out of one hundred percent, this amount will range from about 0.1 percent to about ninety-nine percent of active ingredient, preferably from about 5 percent to about 70 percent, most preferably from about 10 percent to about 30 percent.

[0857] In certain embodiments, a formulation of the present invention comprises an excipient selected from the group consisting of cyclodextrins, celluloses, liposomes, micelle forming agents, e.g., bile acids, and polymeric carriers, e.g., polyesters and polyanhydrides; and a compound of the present invention. In certain embodiments, an aforementioned formulation renders orally bioavailable a compound of the present invention.

[0858] Methods of preparing these formulations or compositions include the step of bringing into association a compound of the present invention with the carrier and, optionally, one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing into association a compound of the present invention with liquid carriers, or finely divided solid carriers, or both, and then, if necessary, shaping the product.

[0859] Formulations of the invention suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia) and/or as mouth washes and the like, each containing a predetermined amount of a compound of the present invention as an active ingredient. A compound of the present invention may also be administered as a bolus, electuary or paste.

[0860] In solid dosage forms of the invention for oral administration (capsules, tablets, pills, dragees, powders, granules, trouches and the like), the active ingredient is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds and surfactants, such as poloxamer and sodium lauryl sulfate; (7) wetting agents, such as, for example, cetyl alcohol, glycerol monostearate, and non-ionic surfactants; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, zinc stearate, sodium stearate, stearic acid, and mixtures thereof; (10) coloring agents; and (11) controlled release agents such as crospovidone or ethyl cellulose. In the case of capsules, tablets and pills, the pharmaceutical compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-shelled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like.

[0861] A tablet may be made by compression or molding, optionally with one or more accessory ingredients. Compressed tablets may be prepared using binder (for example, gelatin or hydroxypropylmethyl cellulose), lubricant, inert diluent, preservative, disintegrant (for example, sodium starch glycolate or cross-linked sodium carboxymethyl cellulose), surface-active or dispersing agent. Molded tablets may be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

[0862] The tablets, and other solid dosage forms of the pharmaceutical compositions of the present invention, such as dragees, capsules, pills and granules, may optionally be scored or prepared with coatings and shells, such as enteric coatings and other coatings well known in the pharmaceutical-formulating art. They may also be formulated so as to provide slow or controlled release of the active ingredient therein using, for example, hydroxypropylmethyl cellulose in varying proportions to provide the desired release profile, other polymer matrices, liposomes and/or microspheres. They may be formulated for rapid release, e.g., freeze-dried. They may be sterilized by, for example, filtration through a bacteria-retaining filter, or by incorporating sterilizing

agents in the form of sterile solid compositions which can be dissolved in sterile water, or some other sterile injectable medium immediately before use. These compositions may also optionally contain opacifying agents and may be of a composition that they release the active ingredient(s) only, or preferentially, in a certain portion of the gastrointestinal tract, optionally, in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes. The active ingredient can also be in micro-encapsulated form, if appropriate, with one or more of the above-described excipients.

[0863] Liquid dosage forms for oral administration of the compounds of the invention include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active ingredient, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

[0864] Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, coloring, perfuming and preservative agents.

[0865] Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[0866] Formulations of the pharmaceutical compositions of the invention for rectal or vaginal administration may be presented as a suppository, which may be prepared by mixing one or more compounds of the invention with one or more suitable nonirritating excipients or carriers comprising, for example, cocoa butter, polyethylene glycol, a suppository wax or a salicylate, and which is solid at room temperature, but liquid at body temperature and, therefore, will melt in the rectum or vaginal cavity and release the active compound.

[0867] Formulations of the present invention which are suitable for vaginal administration also include pessaries, tampons, creams, gels, pastes, foams or spray formulations containing such carriers as are known in the art to be appropriate. Dosage forms for the topical or transdermal administration of a compound of this invention include powders, sprays, ointments, pastes, creams, lotions, gels, solutions, patches and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier, and with any preservatives, buffers, or propellants which may be required.

[0868] The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients, such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

[0869] Powders and sprays can contain, in addition to a compound of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays

can additionally contain customary propellants, such as chlorofluorohydrocarbons and volatile unsubstituted hydrocarbons, such as butane and propane.

[0870] Transdermal patches have the added advantage of providing controlled delivery of a compound of the present invention to the body. Such dosage forms can be made by dissolving or dispersing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate of such flux can be controlled by either providing a rate controlling membrane or dispersing the compound in a polymer matrix or gel.

[0871] Ophthalmic formulations, eye ointments, powders, solutions and the like, are also contemplated as being within the scope of this invention.

[0872] Pharmaceutical compositions of this invention suitable for parenteral administration comprise one or more compounds of the invention in combination with one or more pharmaceutically acceptable sterile isotonic aqueous or nonaqueous solutions, dispersions, suspensions or emulsions, or sterile powders which may be reconstituted into sterile injectable solutions or dispersions just prior to use, which may contain sugars, alcohols, antioxidants, buffers, bacteriostats, solutes which render the formulation isotonic with the blood of the intended recipient or suspending or thickening agents.

[0873] Examples of suitable aqueous and nonaqueous carriers which may be employed in the pharmaceutical compositions of the invention include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol, and the like), and suitable mixtures thereof, vegetable oils, such as olive oil, and injectable organic esters, such as ethyl oleate. Proper fluidity can be maintained, for example, by the use of coating materials, such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

[0874] These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms upon the subject compounds may be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid, and the like. It may also be desirable to include isotonic agents, such as sugars, sodium chloride, and the like into the compositions. In addition, prolonged absorption of the injectable pharmaceutical form may be brought about by the inclusion of agents which delay absorption such as aluminum monostearate and gelatin.

[0875] In some cases, in order to prolong the effect of a drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material having poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

[0876] Injectable depot forms are made by forming microencapsule matrices of the subject compounds in biodegradable polymers such as polylactide-polyglycolide. Depending on the ratio of drug to polymer, and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers

include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissue.

[0877] When the compounds of the present invention are administered as pharmaceuticals, to humans and animals, they can be given per se or as a pharmaceutical composition containing, for example, 0.1 to 99% (more preferably, 10 to 30%) of active ingredient in combination with a pharmaceutically acceptable carrier.

[0878] The preparations of the present invention may be given orally, parenterally, topically, or rectally. They are of course given in forms suitable for each administration route. For example, they are administered in tablets or capsule form, by injection, inhalation, eye lotion, ointment, suppository, etc. administration by injection, infusion or inhalation; topical by lotion or ointment; and rectal by suppositories. Oral administrations are preferred.

[0879] The phrases "parenteral administration" and "administered parenterally" as used herein means modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous, subcuticular, intraarticular, subcapsular, subarachnoid, intraspinal and intrasternal injection and infusion.

[0880] The phrases "systemic administration," "administered systemically," "peripheral administration" and "administered peripherally" as used herein mean the administration of a compound, drug or other material other than directly into the central nervous system, such that it enters the patient's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous administration.

[0881] These compounds may be administered to humans and other animals for therapy by any suitable route of administration, including orally, nasally, as by, for example, a spray, rectally, intravaginally, parenterally, intracisternally and topically, as by powders, ointments or drops, including buccally and sublingually.

[0882] Regardless of the route of administration selected, the compounds of the present invention, which may be used in a suitable hydrated form, and/or the pharmaceutical compositions of the present invention, are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art.

[0883] Actual dosage levels of the active ingredients in the pharmaceutical compositions of this invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient.

[0884] The selected dosage level will depend upon a variety of factors including the activity of the particular compound of the present invention employed, or the ester, salt or amide thereof, the route of administration, the time of administration, the rate of excretion or metabolism of the particular compound being employed, the rate and extent of absorption, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

[0885] A physician or veterinarian having ordinary skill in the art can readily determine and prescribe the effective amount of the pharmaceutical composition required. For example, the physician or veterinarian could start doses of the compounds of the invention employed in the pharmaceutical composition at levels lower than that required in order to achieve the desired therapeutic effect and gradually increase the dosage until the desired effect is achieved.

[0886] In general, a suitable daily dose of a compound of the invention will be that amount of the compound which is the lowest dose effective to produce a therapeutic effect. Such an effective dose will generally depend upon the factors described above. Preferably, the compounds are administered at about 0.01 mg/kg to about 200 mg/kg, more preferably at about 0.1 mg/kg to about 100 mg/kg, even more preferably at about 0.5 mg/kg to about 50 mg/kg. When the compounds described herein are co-administered with another agent (e.g., as sensitizing agents), the effective amount may be less than when the agent is used alone.

[0887] If desired, the effective daily dose of the active compound may be administered as two, three, four, five, six or more sub-doses administered separately at appropriate intervals throughout the day, optionally, in unit dosage forms. Preferred dosing is one administration per day.

[0888] The invention further provides a unit dosage form (such as a tablet or capsule) comprising a heterobifunctional compound described herein in a therapeutically effective amount for the treatment of a medical disorder described herein.

IV. Medical Kits

[0889] Another aspect of this invention is a kit comprising (i) a compound described herein, such as a compound of Formula I, and (ii) instructions for use, such as treating cancer.

EXAMPLES

[0890] The invention now being generally described, will be more readily understood by reference to the following examples, which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and is not intended to limit the invention.

General Methods

[0891] All reactions were carried out under an atmosphere of dry nitrogen or argon. Glassware was oven-dried prior to use. Unless otherwise indicated, common reagents or materials were obtained from commercial sources and used without further purification. N,N-Diisopropylethylamine (DIPEA) was obtained anhydrous by distillation over potassium hydroxide. Tetrahydrofuran (THF), Dichloromethane (CH_2Cl_2), and dimethylformamide (DMF) was dried by a PureSolv™ solvent drying system. PTLC refers to preparatory thin layer chromatographic separation. Abbreviations: HFIP (hexafluoroisopropanol), HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid. Flash column chromatography was performed using silica gel 60 (230-400 mesh). Analytical thin layer chromatography (TLC) was carried out on Merck silica gel plates with QF-254 indicator and visualized by UV or KMnO_4 .

[0892] ^1H and ^{13}C NMR spectra were recorded on an Agilent DD₂ 500 (500 MHz ^1H ; 125 MHz ^{13}C) or Agilent DD₂ 600 (600 MHz ^1H ; 150 MHz ^{13}C) or Agilent DD₂ 400 (400 MHz ^1H ; 100 MHz ^{13}C) spectrometer at room temperature. Chemical shifts were reported in ppm relative to the residual CDCl_3 (δ 7.26 ppm ^1H ; δ 77.0 ppm ^{13}C), CD_3OD (δ 3.31 ppm ^1H ; δ 49.00 ppm ^{13}C), or d_6 -DMSO (δ 2.50 ppm ^1H ; δ 39.52 ppm ^{13}C). NMR chemical shifts were expressed in ppm relative to internal solvent peaks, and coupling constants were measured in Hz. (bs=broad signal). In most cases, only peaks of the major rotamer are reported.

[0893] Mass spectra were obtained using Agilent 1100 series LC/MSD spectrometers. Analytical HPLC analyses were carried out on 250×4.6 mm C-18 column using gradient conditions (10-100% B, flow rate=1.0 mL/min, 20 min), or as described in the LC-MS Method tables.

[0894] Unless indicated otherwise, preparative HPLC was carried out on 250×21.2 mm C-18 column using gradient conditions (10-100% B, flow rate=10.0 mL/min, 20 min). The eluents used were: solvent A (H_2O with 0.1% TFA) and solvent B (CH_3CN with 0.1% TFA). Final products were typically purified via reversed-phase HPLC, PTLC, or flash column chromatography. The abbreviation “TFA” refers to trifluoroacetic acid.

LC-MS Method 01			
HPLC	Instrument	Agilent 1100 LC & Agilent G1956A	
	Software	Agilent Chemstation Rev. B. 04.03[54]	
	Column	Agilent ZORBAX 5 μm SB-Aq, 2.1*50 mm	
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)	
Gradient		Time(min)	Flow(mL/min)
		0.00	0.8
		0.40	0.8
		3.40	0.8
		3.90	0.8
		3.91	0.8
		4.00	1.0
		4.50	1.0
MS	Post time(min)	0	
	Column Temp	50° C.	
	Detector	DAD	
	Ionization source	ESI	
	Drying Gas	N_2	
		Drying Gas Flow	10(L/min)

-continued

	Nebulizer Pressure	40(psi)
	Drying Gas Temperature	350° C.
	Capillary Voltage	2500(V)
	MS Polarity	Positive
	MS Mode	Scan
	Mass Range	100-1500
LC-MS Method 5-95		
HPLC	Instrument	SHIMADZU LCMS-2020
	Software	LabSolution Version 5.93
	Column	Kinetex EVO C18 2.1 × 30 mm, 5 μm
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)
	Gradient	Time(min) B(%) Flow(mL/min)
		0.0 5 1.5
		0.80 95 1.5
		1.20 95 1.5
		1.21 5 1.5
		1.55 5 1.5
MS	Column Temp	50° C.
	Detector	PDA (220 nm&254 nm)
	Ionization source	ESI
	Drying Gas Flow	15(L/min)
	DL Voltage	120(v)
	Qarray DC Voltage	20(V)
	MS Polarity	Positive
	MS Mode	Scan
	Mass range	100-1000
LC-MS Method 10		
HPLC	Instrument	Agilent 1100 LC & Agilent G1956A
	Software	Agilent Chemstation Rev. B. 04.03[54]
	Column	Agilent ZORBAX 5 μm SB-Aq, 2.1*50 mm
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)
	Gradient	Time(min) B(%) Flow(mL/min)
		0.00 10 0.8
		0.40 10 0.8
		3.40 100 0.8
		3.90 100 0.8
		3.91 10 0.8
		4.00 10 1.0
		4.50 10 1.0
MS	Post time(min)	0
	Column Temp	50° C.
	Detector	DAD
	Ionization source	ESI
	Drying Gas	N ₂
	Drying Gas Flow	10(L/min)
	Nebulizer Pressure	40(psi)
	Drying Gas Temperature	350° C.
	Capillary Voltage	2500(V)
	MS Polarity	Positive
	MS Mode	Scan
	Mass Range	100-1500

-continued

LC-MS Method 25				
HPLC	Instrument	Agilent 1100 LC & Agilent G1956A		
	Software	Agilent Chemstation Rev. B. 04.03[54]		
	Column	Agilent ZORBAX 5 μm SB-Aq, 2.1*50 mm		
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)		
Gradient		Time(min)	B(%)	Flow(mL/min)
		0.00	25	0.8
		0.40	25	0.8
		3.40	100	0.8
		3.90	100	0.8
		3.91	25	0.8
		4.00	25	1.0
		4.50	25	1.0
MS	Post time(min)	0		
	Column Temp	50° C.		
	Detector	DAD		
	Ionization source	ESI		
	Drying Gas	N ₂		
	Drying Gas Flow	10(L/min)		
	Nebulizer Pressure	40(psi)		
	Drying Gas Temperature	350° C.		
	Capillary Voltage	2500(V) Positive		
	MS Polarity	Positive		
	MS Mode	Scan		
	Mass Range	100-1500		
LC-MS METHOD 40				
HPLC	Instrument	Agilent 1100 LC & Agilent G1956A		
	Software	Agilent Chemstation Rev. B. 04.03[16]		
	Column	Agilent ZORBAX 5 μm SB-Aq, 2.1*50 mm		
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)		
Gradient		Time(min)	B(%)	Flow(mL/min)
		0.00	40	0.8
		0.40	40	0.8
		3.40	100	0.8
		3.90	100	0.8
		3.91	40	0.8
		4.00	40	1.0
		4.50	40	1.0
MS	Post time(min)	0		
	Column Temp	50° C.		
	Detector	DAD(Agilent 1100)/ELSD(Agilent 1260 Infinity)		
	Ionization source	ESI		
	Drying Gas	N ₂		
	Drying Gas Flow	10(L/min)		
	Nebulizer Pressure	2070(Torr)		
	Drying Gas Temperature	350° C.		
	Capillary Voltage	2500(V) Positive		
LC-MS METHOD 100				
HPLC	Instrument	Agilent 1100 LC & Agilent G1956A		
	Software	Agilent Chemstation Rev. B. 04.03[16]		
	Column	Agilent ZORBAX 5 μm SB-Aq, 2.1*50 mm		
	Mobile Phase	A: 0.0375% TFA in water (v/v) B: 0.01875% TFA in Acetonitrile (v/v)		
Gradient		Time(min)	B(%)	Flow(mL/min)
		0.00	0	0.6
		0.40	0	0.6
		3.40	80	0.6


-continued

		3.90	100	0.6
		3.91	0	0.6
		4.00	0	1.0
		4.50	0	1.0
MS	Post time(min)	1.20		
	Column Temp	50° C.		
	Detector	DAD		
	Ionization source	ESI		
	Drying Gas	N2		
	Drying Gas Flow	10(L/min)		
	Nebulizer Pressure	2070(Torr)		
	Drying Gas Temperature	350° C.		
	Capillary Voltage	2500(V) Positive		
	MS Polarity	Positive		
	MS Mode	Scan		
	Mass Range	50-1500		

[0895] The following abbreviations are used herein: ACN: acetonitrile; Bn: benzyl; Boc: tert-butoxycarbonyl; DCM: dichloromethane; DIEA: diisopropylethylamine; DMF: dimethylformamide; DMSO: dimethylsulfoxide; EtOH: ethanol; EA or EtOAc: ethyl acetate; equiv. or eq.: molar equivalents; FA: formic acid; h: hour or hours; HATU: 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxide hexafluorophosphate; HPLC: high-pressure liquid chromatography; LCMS or LC-MS: liquid chromatography-mass spectrometry; MeCN: acetonitrile; MeOH: methanol; MS: mass spectrometry; NMP: N-methylpyrrolidone; NMR: nuclear magnetic resonance; PE: petroleum ether; rt: room temperature; TEA: triethylamine; TFA: trifluoroacetic acid; THF: tetrahydrofuran; TLC: thin-layer chromatography; psi: pounds-per-square inch; and Tos or Ts: p-toluenesulfonyl.

Example 1—Synthesis of Exemplary Compounds

[0896] The compounds set forth in the table below were prepared, according to the following general approach, a related synthetic strategy, or based on procedures described herein below.



[0897] Coupling compound A (a precursor of TPL, for example, a discrete compound that is a target protein ligand) with L' (a precursor to linker L, containing functionality for coupling to the precursors of both TPL and EPL) affords intermediate B (wherein L'' is a precursor to linker L that contains functionality for coupling to the EPL precursor). The coupling of compound A with L' can be accomplished using, for example, amide coupling conditions. Coupling intermediate B with compound C (a precursor of EPL) affords heterobifunctional compound D. The coupling of intermediate B with compound C can be accomplished using, for example, amide coupling conditions.

Compound No.	Structure
I-40	


-continued

Compound No.	Structure
I-41	
I-42	


[0898] Physical characterization data for compounds I-40, I-41, and I-42 is provided below.

Compound No.	Observed Mass	Retention Time (min)	LCMS Method
I-40	515.7, 1030.3	220-1.970	25
I-41	559.7, 1118.3	220-2.013	25
I-42	603.8, 1206.4	220-1.064	40

Example 2—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)butyl)acetamide (I-1)



-continued



[0899] Step 1: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-hydroxybutyl)acetamide. To a solution of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]acetic acid (219 mg, 546 μ mol, 1.0 equiv), 4-aminobutan-1-ol (48.7 mg, 546 μ mol, 50.7 μ L, 1.0 equiv), EDCI (209 mg, 1.09 mmol, 2.0 equiv) and HOBt (148 mg, 1.09 mmol, 2.0 equiv) in DMF (3 mL) was added DIEA (282 mg, 2.19 mmol, 381 μ L, 4.0 equiv) and the mixture was stirred at 25° C. for 16 h. The reaction mixture was diluted with water (10 mL) and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Unisil 3-100 C18 Ultra 150 $\times$ 50 mm $\times$ 3 μ m; mobile phase: [water (0.225% FA)-ACN]; B %: 30%-60%, 10 min) to give (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-hydroxybutyl)acetamide (130 mg, 49% yield) as a white solid. LC-MS: MS (ES⁻): RT=0.683 min, m/z=472.0 [M+H⁺].

[0900] Step 2: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-oxobutyl)acetamide. To a solution of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-


9-yl]-N-(4-hydroxybutyl)acetamide (50.0 mg, 106 μ mol, 1.0 equiv) in DMF (1 mL) was added DMP (67.4 mg, 159 μ mol, 49.2 μ L, 1.5 equiv) and the mixture was stirred at 25° C. for 1 h. The mixture was filtered to give (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-oxobutyl)acetamide (50 mg) as a light yellow oil. LC-MS: MS (ES⁻): RT=0.621 min, m/z=470.1 [M+H⁺].

[0901] Step 3: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)butyl)acetamide. To a solution of [4-[2-(ethylamino)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (58.3 mg, 106 μ mol, 1.0 equiv, TFA salt) and NaBH(OAc)₃ (225 mg, 1.06 mmol, 10 equiv) in DCM (1 mL) was added TEA (53.8 mg, 532 μ mol, 74.0 μ L, 5.0 equiv), then to the mixture solution was added a solution of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-oxobutyl)acetamide (50.0 mg, 106 μ mol, 1.0 equiv) in DMF (1 mL) and the mixture solution was stirred at 25° C. for 16 h. The reaction mixture was diluted with water (30 mL) and extracted with DCM (50 mL $\times$ 3). The combined organic layers were dried over Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to give a residue. The

residue was purified by prep-HPLC (column: Waters Xbridge 150×25 mm×5 um; mobile phase: [water (10 mM NH₄HCO₃)-ACN]; B %: 36%-66%, 9 min) to give compound (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)butyl)acetamide (I-1, 18.5 mg, 20% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.326 min, m/z=887.3 [M+H⁺], m/z=443.3 [M+2H⁺]/2; LCMS method: 40. ¹H NMR (400 MHz, DMSO-d₆) δ 9.76 (s, 2H), 8.17 (s, 1H), 7.63-7.61 (m, 2H), 7.45-7.42 (m, 4H), 7.33 (s, 1H), 7.32 (s, 1H), 7.17-7.15 (m, 2H), 6.89-6.87 (m, 3H), 6.68-6.65 (m, 2H), 4.51-4.50 (m, 1H), 4.03-4.00 (m, 2H), 3.32-3.30 (m, 2H), 3.21-3.18 (m, 4H), 2.73-2.67 (m, 2H), 2.58 (s, 3H), 2.50-2.44 (m, 2H), 2.37 (s, 3H), 1.42-1.41 (m, 4H), 1.09 (t, J=6.8 Hz, 3H).

Example 3—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(8-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)octyl)acetamide (I-2)

then to the mixture solution was added a solution of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(8-oxooctyl)acetamide (50.0 mg, 95.0 μmol, 1.0 equiv) in DMF (1 mL), the mixture solution was stirred at 25° C. for 1 h. The reaction mixture was diluted with water (30 mL), extracted with DCM (50 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Unisil 3-100 C18 Ultra 150×50 mm×3 um; mobile phase: [water (0.225% FA)-ACN]; B %: 28%-58%, 10 min) to give a crude product, then the crude product was purified by prep-HPLC (column: Unisil 3-100 C18 Ultra 150×50 mm×3 um; mobile phase: [water (0.225% FA)-ACN]; B %: 25%-55%, 10 min) to give 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatri-cyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[8-[ethyl-[2-[4-[6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl]phenoxy]ethyl]amino]octyl]acetamide (I-2, 27.8 mg, 31% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.695 min, m/z=943.3 [M+H⁺], m/z=472.3 [M+2H⁺]/2; LCMS method: 40. ¹H NMR (400 MHz, DMSO-d₆) δ 8.29 (s, 1H), 8.16-8.15 (m, 1H), 7.66-7.64 (m, 2H), 7.45-7.42 (m, 4H), 7.34-7.33 (m, 1H), 7.23 (s, 1H),




I-2

[0902] To a solution of [4-[2-(ethylamino)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophen-3-yl] methanone (52.0 mg, 95.0 μmol, 1.0 equiv, TFA salt) and NaBH(OAc)₃ (201 mg, 950 μmol, 10 equiv) in DCM (1 mL) was added TEA (48.1 mg, 475 μmol, 66.1 μL, 5.0 equiv),

7.17-7.15 (m, 2H), 7.00-6.88 (m, 3H), 6.68-6.66 (m, 2H), 4.52-4.48 (m, 1H), 4.03-4.00 (m, 2H), 3.24-3.16 (m, 2H), 3.14-3.12 (m, 4H), 2.73-2.71 (m, 2H), 2.58 (s, 3H), 2.50-2.39 (m, 5H), 1.61 (s, 3H), 1.41-1.33 (m, 4H), 1.23-1.21 (m, 8H), 1.93-0.90 (t, J=6.8 Hz, 3H).

Example 4—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxy phenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)ethoxy)ethoxy)ethyl)acetamide (I-3)



I-3

[0903] Step 1: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-oxoethoxy)ethoxy)ethyl)acetamide. To a solution of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-oxoethoxy)ethoxy)ethyl)


acetamide (55.0 mg, 103 μ mol, 1.0 equiv) in DMF (1.5 mL) was added DMP (110 mg, 258 μ mol, 80.0 μ L, 2.5 equiv). The mixture was stirred at 25° C. for 0.5 h. The reaction mixture was concentrated to give (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-oxoethoxy)ethoxy)ethyl)acet-


amide (54 mg) as a yellow oil. LC-MS: MS (ES⁺): RT=0.851 min, m/z=530.3 [M+H⁺].

[0904] Step 2: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)ethoxy)ethoxy)ethyl) acetamide. To a solution of [4-[2-(ethylamino)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxy-phenyl)benzothiophen-3-yl]methanone (50.0 mg, 91.3 μ mol, 1.0 equiv, TFA salt) in DCM (3 mL) was added TEA (92.4 mg, 913 μ mol, 127 μ L, 10 equiv) and NaBH(OAc)₃ (194 mg, 913 μ mol, 10 equiv), then (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-oxoethoxy)ethoxy)ethyl)acetamide (48.4 mg, 91.3 μ mol, 1.0 equiv) was added and the mixture was stirred at 20° C. for 0.5 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was purified by prep-HPLC (column: Waters Xbridge 150 $\times$ 25 mm $\times$ 5 μ m; mobile phase: [water (10 mM NH₄HCO₃)-ACN]; B %: 44%-74%, 9 min))

to give compound (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(2-(2-(2-(ethyl(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)amino)ethoxy)ethoxy)ethyl) acetamide (I-3, 20 mg, 23% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=2.064 min, m/z=947.2 [M+H⁺]; m/z=474.3 [M+2H⁺]/2 LCMS Method: 25. ¹H NMR (400 MHz, CD₃OD) δ 7.64 (d, J=8.8 Hz, 2H), 7.33-7.43 (m, 5H), 7.27 (d, J=2.1 Hz, 1H), 7.18 (d, J=8.6 Hz, 2H), 6.87 (dd, J=8.8, 2.2 Hz, 1H), 6.76 (d, J=8.9 Hz, 2H), 6.63 (d, J=8.5 Hz, 2H), 4.62 (dd, J=9.1, 5.1 Hz, 1H), 4.07 (t, J=5.7 Hz, 2H), 3.58-3.64 (m, 7H), 3.38-3.49 (m, 5H), 2.94 (t, J=5.5 Hz, 2H), 2.79 (t, J=5.6 Hz, 2H), 2.64-2.73 (m, 5H), 2.44 (s, 3H), 1.68 (s, 3H), 1.06 (t, J=7.1 Hz, 3H).

Example 5—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)-6,9,12,15-tetraoxa-3-azaheptadecan-17-yl) acetamide (I-4)






[0905] Step 1: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(14-hydroxy-3,6,9,12-tetraoxa-tetradecyl)acetamide. To a solution of 2-[2-[2-[2-(2-aminoethoxy)ethoxy]ethoxy]ethoxy]ethanol (200 mg, 569 μmol, 1.0 equiv, TFA salt) and 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]acetic acid (293 mg, 569 μmol, 1.0 equiv, TFA salt) in DMF (2 mL) was added DIEA (368 mg, 2.85 mmol, 496 μL, 5.0 equiv) and HATU (238 mg, 626 μmol, 1.1 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150x25 mmx10 um; mobile phase: [water (0.225% FA)-ACN]; B %: 33%-63%, 10 min) to give compound (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(14-hydroxy-3,6,9,12-tetraoxatetradecyl)acetamide (47.0 mg, 13% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.857 min, m/z=620.2 [M+H⁺].


[0906] Step 2: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(14-oxo-3,6,9,12-tetraoxatetradecyl) acetamide. To a solution of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo [8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-(2-hydroxy-ethoxy)ethoxy]ethoxy]ethoxy]ethyl]acetamide (47.0 mg, 75.8 μmol, 1.0 equiv) in DMF (1.5 mL) was added DMP (80.4 mg, 189 μmol, 58.7 μL, 2.5 equiv). The mixture was stirred at 20° C. for 1 h. The reaction mixture was concentrated to give (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-

yl)-N-(14-oxo-3,6,9,12-tetraoxatetradecyl) acetamide (46.5 mg) as a yellow oil. LC-MS: MS (ES⁺): RT=0.824 min, m/z=620 [M+H⁺]

[0907] Step 3: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[h]thiophene-3-carbonyl)phenoxy)-6,9,12,15-tetraoxa-3-azahexadecan-17-yl)acetamide. To a solution of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-(2-oxoethoxy) ethoxy]ethoxy]ethoxy]ethyl)acetamide (46.0 mg, 74.4 μmol , 1.0 equiv) and [4-[2-(ethylamino) ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (40.8 mg, 74.4 μmol , 1.0 equiv, TFA salt) in DCM (2 mL) was added $\text{NaBH}(\text{OAc})_3$ (158 mg, 744 μmol , 10 equiv) and TEA (75.3 mg, 744 μmol , 104 μL , 10 equiv). The mixture was stirred at 20° C. for 12 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150x25 mmx10 μm ; mobile phase:[water(0.225% FA)-ACN];B %: 25%-55%,10 min)) to give (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[h]thiophene-3-carbonyl)phenoxy)-6,9,12,15-tetraoxa-3-azahexadecan-17-yl)acetamide (1-4, 15.4 mg, 20% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.077 min, m/z =1035.3 [M+H⁺]; m/z =518.3 [M+2H⁺]/2 LCMS Method: 40. ¹H NMR (400 MHz, CD₃OD) δ 7.69-7.41 (m, 2H), 7.29-7.35 (m, 5H), 7.26-7.25 (m, 1H), 7.16-6.82 (m, 3H), 6.62-6.61 (m, 2H), 4.86-4.60 (m, 1H), 4.16 (s, 2H), 3.64-3.58 (m, 2H), 3.58-3.55 (m, 14H), 3.43-3.32 (m, 3H), 3.31-3.30 (m, 1H), 3.15 (s, 2H), 2.65-2.42 (m, 4H), 2.65 (s, 3H), 2.42 (s, 3H), 1.67 (s, 3H), 1.15-1.12 (t, J=7 Hz, 3H).

Example 6—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)-6,9,12,15,18,21-hexaoxa-3-azatricosan-23-yl) acetamide (I-5)






I-5

[0908] Step 1: Preparation of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11, 12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethoxy]ethoxy]ethyl]acetamide. To a solution of 2-[2-[2-[2-[2-(2-aminoethoxy)ethoxy]ethoxy]ethoxy]ethoxy]ethanol (200 mg, 455 μ mol, 1.0 equiv, TFA salt) and 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]acetic acid (234 mg, 455 μ mol, 1.0 equiv, TFA salt) in DMF (4 mL) was added DIEA (176 mg, 1.37 mmol, 238 μ L, 3.0 equiv) and HATU (190 mg, 501 μ mol, 1.1 equiv). The mixture was stirred at 20° C. for 0.1 h. The reaction mixture was concentrated. It was purified by prep-HPLC (column: Phenomenex Gemini-NX C18 75*30 mm*3 μ m; mobile phase: [water(0.225% FA)-ACN]; B %: 20%-50%, 7 min) to give 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11, 12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethoxy]ethyl]acetamide (50.0 mg, 70.6 μ mol, 16% yield) as a yellow gum. LC-MS: MS (ES⁺): RT=0.625 min, m/z=708.1 [M+H⁺].


[0909] Step 2: Preparation of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11, 12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-oxoethoxy)ethoxy]ethoxy]ethoxy]ethoxy]ethyl]acetamide. To a solution of 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-hydroxyethoxy)ethoxy]ethoxy]ethoxy]ethyl]acetamide (50.0 mg, 70.6 μ mol, 1.0 equiv) in DMF (1 mL) was added DMP (74.9 mg, 176 μ mol, 54.6 μ L, 2.5 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was concentrated to give 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-oxoethoxy)ethoxy]ethoxy]ethoxy]ethyl]acetamide (49.0 mg, 69.4 μ mol, 98% yield) as a yellow oil.

[0910] Step 3: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)-6,9,12,15,18,21-hexaoxa-3-azatricosan-23-yl)acetamide. To a solution of [4-[2-(ethylamino)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (40.0 mg, 73.1 μ mol, 1.0 equiv, TFA salt) in DCM (3 mL) was added TEA (73.9 mg, 731 μ mol, 102 μ L, 10 equiv) and NaBH(OAc)₃ (155 mg, 731 μ mol, 10 equiv), then 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[2-[2-[2-[2-[2-(2-oxoethoxy)ethoxy]ethoxy]ethoxy]ethyl]acetamide (49.0 mg, 69.4 μ mol, 1.0 equiv) was added and the mixture was stirred at 20° C. for 1 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The crude product was purified by prep-HPLC (column: Phenomenex luna C18 150*25 mm*10 μ m; mobile phase: [water (0.225% FA)-ACN]; B %: 25%-55%, 10 min and column: Phenomenex Gemini-NX C18 75*30 mm*3 μ m; mobile phase: [water(10 mM NH₄HCO₃)-ACN]; B %: 35%-65%, 8 min and column: Phenomenex Gemini-NX C18 75*30 mm*3 μ m; mobile phase: [water(10 mM NH₄HCO₃)-ACN]; B %: 35%-65%, 8 min) to give compound (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(3-ethyl-1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)-6,9,12,15,18,21-hexaoxa-3-azatricosan-23-yl)acetamide (I-5, 15.3 mg, 18% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.327 min, m/z=1123.4 [M+H⁺]; m/z=562.3 [M+2H⁺]/2; LCMS Method: 40. ¹H NMR (400 MHz, CD₃OD) δ 7.71 (d, J=9.0 Hz, 2H), 7.37-7.49 (m, 5H), 7.27 (d, J=2.3 Hz, 1H), 7.20 (d, J=8.8 Hz, 2H), 6.80-6.91 (m, 3H), 6.64 (d, J=8.8 Hz, 2H), 4.65 (d, J=5.3 Hz, 1H), 4.13 (s, 2H), 3.43-3.65 (m, 28H), 3.00 (s, 2H), 2.69 (s, 6H), 2.44 (s, 3H), 1.70 (s, 3H), 1.09 (t, J=7.6 Hz, 3H).

Example 7—Synthesis of (6S)-2,3,6,9-tetramethyl-4-(4-(3-(4-(6-(6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (1-6)



-continued



I-6

[0911] Step 1: Preparation of tert-butyl 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)-3',6'-dihydro-[3,4'-bipyridine]-1'-(2'H)-carboxylate. To a solution of (6S,8R)-6-(5-bromopyridin-2-yl)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline (700 mg, 1.65 mmol, 1.00 equiv), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydropyridine-1(2H)-carboxylate (610 mg, 1.98 mmol, 1.20 equiv), and cesium carbonate (1.61 g, 4.94 mmol, 3.00 equiv) in dioxane (12 mL) and water (3 mL) was added 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride (240 mg, 0.32 mmol, 0.20 equiv) and the mixture was stirred at 90° C. for 2 h under a nitrogen atmosphere. The reaction mixture was quenched with water (60 mL), and then extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (80 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica gel, petroleum ether/ethyl acetate=1/0 to 1/1) to afford tert-butyl 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)-3',6'-dihydro-[3,4'-bipyridine]-1'-(2'H)-carboxylate (550 mg, 63% yield) as a yellow oil. LCMS: MS (ES⁺): RT=0.530 min, m/z=528.3 [M+H]⁺.

[0912] Step 2: Preparation of tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidine-1-carboxylate. To a solution of tert-butyl 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)-3',6'-dihydro-[3,4'-bipyridine]-1'-(2'H)-carboxylate (550 mg, 1.04 mmol, 1.00 equiv) in methanol (15 mL) was added palladium on activated carbon (100 mg, 10% purity) and then the mixture was degassed and purged with hydrogen three times, and then the mixture was stirred at 25° C. for 5 h under a hydrogen (15 psi) atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to afford tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidine-1-carboxylate (426 mg, 77% yield) as a yellow oil, which was used in the next step without further purification. LCMS: MS (ES⁺): RT=0.544 min, m/z=530.3 [M+H]⁺.

[0913] Step 3: Preparation of (6S,8R)-8-methyl-6-(5-(piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline. To a solution of tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidine-1-carboxylate (426 mg, 0.80 mmol, 1.00 equiv) in dichloromethane (4.5 mL) was added trifluoro-

acetic acid (2.30 g, 20.19 mmol, 25.10 equiv) dropwise and the mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to afford (6S,8R)-8-methyl-6-(5-(piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (345 mg, 99% yield) as a white solid, which was used in the next step without further purification. LCMS: MS (ES⁺): RT=0.432 min, m/z=430.3 [M+H]⁺.


[0914] Step 4: Preparation of tert-butyl 3-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidine-1-carboxylate. To a solution of (6S,8R)-8-methyl-6-(5-(piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (345 mg, 0.80 mmol, 1.00 equiv), tert-butyl 3-oxopyrrolidine-1-carboxylate (446 mg, 2.41 mmol, 3.00 equiv) in dichloromethane (8 mL) was added sodium triacetoxy borohydride (510 mg, 2.41 mmol, 3.00 equiv) and the mixture was stirred at 25° C. for 1 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane:methanol=10:1) to afford tert-butyl 3-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidine-1-carboxylate (450 mg, 93% yield) as a white solid. LCMS: MS (ES⁺): RT=0.440 min, m/z=599.3 [M+H]⁺.


[0915] Step 5: Preparation of (6S,8R)-8-methyl-6-(5-(1-(pyrrolidin-3-yl)piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline. To a solution of tert-butyl 3-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidine-1-carboxylate (450 mg, 0.75 mmol, 1.00 equiv) in dichloromethane (4.5 mL) was added trifluoroacetic acid (2.30 g, 20.19 mmol, 26.87 equiv) dropwise and then the mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Waters Xbridge Prep OBD C18 150*40 mm*10 um; mobile phase: [water(NH₄HCO₃)-ACN]; gradient: 20%-50% B over 20 min] to afford (6S,8R)-8-methyl-6-(5-(1-(pyrrolidin-3-yl)piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (230 mg, 61% yield) as a white solid. LCMS: MS (ES⁺): RT=0.381 min, m/z=499.4 [M+H]⁺.

[0916] Step 6: Preparation of (6S)-2,3,6,9-tetramethyl-4-(4-(3-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of (6S,8R)-8-methyl-6-(5-(1-(pyrrolidin-3-yl)piperidin-4-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (150 mg, 0.30 mmol, 1.00 equiv), (S)-4-(4-chlorophenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (161 mg, 0.45 mmol, 1.50 equiv), cesium carbonate (294 mg, 0.90 mmol, 3.00 equiv) in dioxane (3 mL) was added (2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl) [2-(2'-amino-1,1'-biphenyl)]palladium(II) methanesulfonate (46 mg, 0.060 mmol, 0.20 equiv) and the mixture was stirred at 90° C. for 4 h under nitrogen atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column:

Phenomenex luna C18 150*25 mm*10 um; mobile phase: [water(FA)-ACN]; gradient:8%-38% B over 1 min] to give the crude product. The product was purified by SFC [column: DAICEL CHIRALCEL OX (250 mm*30 mm, 10 um); mobile phase: [CO₂-ACN/MeOH (0.1% NH₃H₂O)];B %:70%, isocratic elution mode] to give the crude product (peak 1). The crude product was further purified by prep-HPLC [column: Phenomenex luna C18 150 mm*25 mm*10 um; mobile phase: [water(FA)-ACN]; gradient: 8%-38% B over 1 min] to afford (6S)-2,3,6,9-tetramethyl-4-(4-(3-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperidin-1-yl)pyrrolidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-6, 10.37 mg, 4.17% yield) as a yellow solid. LCMS: MS (ES⁺): RT=0.981 min, m/z=819.6 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ 8.38 (s, 1H), 8.06 (s, 1H), 7.50 (d, J=8.0 Hz, 1H), 7.42-7.35 (m, 3H), 7.23 (s, 1H), 6.98 (d, J=8.8 Hz, 1H), 6.47 (d, J=8.4 Hz, 2H), 5.13 (s, 1H), 4.16-4.08 (m, 1H), 3.63-3.54 (m, 2H), 3.52-3.45 (m, 1H), 3.40-3.25 (m, 4H), 3.23-3.15 (m, 1H), 3.12-2.95 (m, 3H), 2.93-2.86 (m, 1H), 2.67 (s, 3H), 2.61-2.53 (m, 1H), 2.41 (s, 3H), 2.34-2.14 (m, 4H), 2.07 (d, J=6.4 Hz, 3H), 1.92-1.85 (m, 4H), 1.78 (s, 3H), 1.17 (d, J=6.4 Hz, 3H).

Example 8—Synthesis of (S)-2,3,6,9-tetramethyl-4-(4-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl) piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-7)






[0917] Step 1: Preparation of tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazine-1-carboxylate. To a solution of (6S,8R)-6-(5-bromopyridin-2-yl)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline (1.00 g, 2.35 mmol, 1.00 equiv), tert-butyl piperazine-1-carboxylate (1.75 g, 9.41 mmol, 4.00 equiv), and potassium tert-butoxide (1.06 g, 9.41 mmol, 4.00 equiv) in dioxane (20 mL) was added methanesulfonato(2-dicyclohexylphosphino-2,6-di-1-propoxy-1,1-biphenyl)(2-amino-1,1-biphenyl-2-yl)palladium(ii) (393 mg, 0.47 mmol, 0.20 equiv) and the mixture was stirred at 100° C. for 10 h under a nitrogen atmosphere. The reaction mixture was quenched with water (100 mL), and then extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (100 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica gel, petroleum ether/ethyl acetate=1/0 to 1/1) to afford tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazine-1-carboxylate (400 mg, 32% yield) as a yellow solid. LCMS: MS (ES⁺): RT=0.493 min, m/z=531.4 [M+H]⁺.


[0918] Step 2: Preparation of (6S,8R)-8-methyl-6-(5-(piperazin-1-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline. To a solution of tert-butyl 4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyri-

din-3-yl)piperazine-1-carboxylate (400 mg, 0.75 mmol, 1.00 equiv) in dichloromethane (4.5 mL) was added dropwise trifluoroacetic acid (2.30 g, 20.19 mmol, 26.79 equiv) and the mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Waters Xbridge 150×25 mm×5 μm; mobile phase: [water (ammonia hydroxide v/v)-ACN]; gradient:20%-500% B over 10 min] to afford (6S,8R)-8-methyl-6-(5-(piperazin-1-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline (130 mg, 40% yield) as a white solid. LCMS: MS (ES⁺): RT=0.585 min, m/z=431.2 [M+H]⁺.

[0919] Step 3: Preparation of (S)-2,3,6,9-tetramethyl-4-(4-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of (6S,8R)-8-methyl-6-(5-(piperazin-1-yl)pyridin-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline (35 mg, 0.08 mmol, 1.00 equiv), (S)-4-(4-chlorophenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (43 mg, 0.12 mmol, 1.50 equiv), and cesium carbonate (79 mg, 0.24 mmol, 3.00 equiv) in dioxane (1 mL) was added dicyclohexyl[2-(2,4,6-triisopropylphenyl)phenyl]phosphane; methanesulfonate; [2-[2-(methylamino)phenyl]phenyl]palladium(1+) (13 mg, 0.01 mol, 0.20 equiv) and the mixture was stirred at 90° C. for 2 h under nitrogen atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Phenomenex luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient:15%-45% B over 1 min] to give the crude product. The crude product was purified by SFC [column: DAICEL CHIRALCEL OD (250 mm×30 mm, 10 μm); mobile phase: [CO₂-ACN/EtOH (0.1% NH₃H₂O)]; B %: 60%, isocratic elution mode] to give a product that was further purified by prep-HPLC [column: Phenomenex luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient: 15%-45% B over 1 min] to afford (S)-2,3,6,9-tetramethyl-4-(4-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-7, 8.61 mg, 13.53% yield) as a yellow solid. LCMS: MS (ES⁺): RT=1.483 min, m/z=751.6 [M+H]⁺; LCMS Method: 10. ¹H NMR (400 MHz, CDCl₃) δ 8.21 (d, J=2.8 Hz, 1H), 8.07 (s, 1H), 7.43 (d, J=8.4 Hz, 2H), 7.31 (d, J=9.2 Hz, 1H), 7.24-7.18 (m, 2H), 6.94 (d, J=8.8 Hz, 1H), 6.89 (d, J=8.8 Hz, 2H), 5.08 (s, 1H), 4.15 (q, J=6.8 Hz, 1H), 3.66-3.56 (m, 1H), 3.46-3.38 (m, 4H), 3.38-3.30 (m, 5H), 3.29-3.18 (m, 1H), 3.08-2.99 (m, 1H), 2.93-2.85 (m, 1H), 2.68 (s, 3H), 2.42 (s, 3H), 2.08 (d, J=6.8 Hz, 3H), 1.76 (s, 3H), 1.16 (d, J=6.4 Hz, 3H).

Example 9—Synthesis of 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (1-8)





[0920] Step 1: Preparation of tert-butyl 3-(((2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate. To a solution of tert-butyl 3-(4-formylphenyl)azetidine-1-carboxylate (900 mg, 2 mmol, 1 equiv) and 2-(6-((tert-butyldimethylsilyl)oxy)-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyaniline (603 mg, 2 mmol, 1 equiv) in DCM (5 mL) was added AcOH (201.65 mg, 3.36 mmol, 1.5 equiv). The mixture was stirred at 30° C. for 2 h, then NaBH₃CN (281.36 mg, 4.48 mmol, 2 equiv) was added, and the mixture was stirred at 30° C. for 1 h. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150*40 mm* 15 um; mobile phase: [water(FA)-ACN]; gradient:55%-85% B over 15 min) to afford tert-butyl 3-(((2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate (610 mg, 52% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.608 min, m/z=515.3 [M+H⁺].

[0921] Step 2: Preparation of tert-butyl 3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate. To a solution of tert-butyl 3-(((2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate (380 mg, 0.74 mmol, 1 equiv) and acetaldehyde (5 M, 2.22 mL, 15 equiv) in MeOH (3 mL) was added NaBH₃CN (92.80 mg, 1.48 mmol, 2 equiv). The mixture was stirred at 20° C. for 2 hr. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether/ethyl acetate=3/1) to afford tert-butyl 3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate (362 mg, 84% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.26-7.21 (m, 2H), 7.16 (d, J=8.5 Hz, 3H), 6.90 (d, J=8.0 Hz, 1H), 6.80 (d, J=2.4 Hz, 1H), 6.71 (dd, J=2.4, 8.8 Hz, 1H), 6.66-6.59 (m, 2H), 4.31 (t, J=8.8 Hz, 2H), 4.01 (s, 2H), 3.94 (dd, J=6.4, 8.0 Hz, 2H), 3.81 (s, 3H), 3.74-3.56 (m, 2H), 3.03-2.90 (m, 2H), 2.85-2.75 (m, 2H), 2.75-2.64 (m, 2H), 1.84-1.69 (m, 2H), 1.62 (d, J=6.8 Hz, 2H), 1.48 (s, 9H), 0.96 (t, J=7.2 Hz, 3H). LC-MS: MS (ES⁺): RT=0.530 min, m/z=543.4 [M+H⁺].

[0922] Step 3: Preparation of 6-(2-((4-(azetidin-3-yl)benzyl)(ethyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol. To a solution of tert-butyl 3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidine-1-carboxylate (380 mg, 0.7 mmol, 1 equiv) in DCM (3 mL) was added TFA (1.77 g, 15.48 mmol, 22 equiv). The mixture was stirred at 25° C. for 1 hr. The reaction mixture was concentrated under reduced pressure to remove solvent to afford 6-(2-((4-(azetidin-3-yl)benzyl)(ethyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (310 mg, crude) as a yellow solid. LC-MS: MS (ES⁺): RT=0.414 min, m/z=443.2 [M+H⁺].

[0923] Step 4: Preparation of tert-butyl 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-4-hy-

droxypiperidine-1-carboxylate. tert-Butyl 1-oxa-6-azaspiro [2.5]octane-6-carboxylate (223 mg, 1 mmol, 1.5 equiv), 6-(2-((4-(azetidin-3-yl)benzyl)(ethyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (309 mg, 0.7 mmol, 1 equiv) and TEA (211 mg, 2.09 mmol, 3 equiv) were taken up into a microwave tube in ethyl alcohol (5 mL) and water (1 mL). The sealed tube was heated at 100° C. for 3 h under microwave irradiation. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether/ethyl acetate=0/1) to afford tert-butyl 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-4-hydroxypiperidine-1-carboxylate (210 mg, 44% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.456 min, m/z=656.5 [M+H⁺].


[0924] Step 5: Preparation of 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)piperidin-4-ol. To a solution of tert-butyl 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-4-hydroxypiperidine-1-carboxylate (110 mg, 0.16 mmol, 1 equiv) in DCM (2 mL) was added TFA (57.37 mg, 0.5 mmol, 3 equiv). The mixture was stirred at 20° C. for 0.5 hr. The reaction mixture was quenched by the addition of a saturated solution of aqueous sodium bicarbonate (10 mL) and then diluted with water (10 mL) and extracted with EA (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)piperidin-4-ol (166 mg, crude) as a yellow solid. LC-MS: MS (ES⁺): RT=0.398 min, m/z=556.4 [M+H⁺].


[0925] Step 6: Preparation of 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol. A mixture of 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)piperidin-4-ol (160 mg, 0.28 mmol, 1 equiv), (S)-4-(4-chlorophenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (123.29 mg, 0.34 mmol, 1.2 equiv), Cs₂CO₃ (281.41 mg, 0.86 mmol, 3 equiv), and dicyclohexyl-[2-(2,6-dimethoxyphenyl)phenyl]phosphane; methanesulfonate;(2-phenylanilino)palladium(1+) (22.46 mg, 0.03 mmol, 0.1 equiv) in dioxane (3 mL) was degassed and purged with N₂ 3 times, and then the mixture was stirred at 90° C. for 3 hr under an N₂ atmosphere. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM/MeOH=10/1) to afford 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl)phenyl)azetidin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol as a white solid. The


solid was purified by SFC (column: DAICEL CHIRALCEL OX (250 mm×30 mm, 10 μm); mobile phase: [CO₂-ACN/EtOH (0.1% NH₃H₂O)]; B %: 70%, isocratic elution mode) to afford 4-((3-(4-((ethyl(2-(6-hydroxy-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)amino)methyl) phenyl)azetidin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl) piperidin-4-ol (I-8, 40 mg, 46% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 9.01 (s, 1H), 7.27 (d, J=8.4 Hz, 2H), 7.21-7.12 (m, 5H), 6.90 (d, J=9.0 Hz, 2H), 6.83-6.77 (m, 2H), 6.66 (dd, J=2.8, 8.8 Hz, 1H), 6.55-6.45 (m, 2H), 4.17-4.07 (m, 2H), 3.98 (d, J=2.8 Hz, 2H), 3.71 (s, 3H), 3.65 (t, J=7.2 Hz, 2H), 3.56-3.46 (m, 5H), 3.14-3.07

(m, 4H), 2.87 (d, J=7.2 Hz, 2H), 2.73 (d, J=3.6 Hz, 2H), 2.57 (s, 3H), 2.40 (s, 5H), 1.84 (d, J=6.4 Hz, 3H), 1.70 (s, 3H), 1.62-1.41 (m, 6H), 1.27-1.19 (m, 1H), 0.87 (t, J=7.2 Hz, 3H). LC-MS: MS (ES⁺): RT=1.513 min, m/z=876.5 [M+H⁺]. LCMS Method: 5-95.

Example 10—Synthesis of (S)-6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl(4-(2-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)ethoxy)phenyl methanone (I-9)







[0926] Step 1: Preparation of tert-butyl-4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-(4-hydroxybutyl)piperidine-1-carboxylate (1 g, 3.89 mmol, 1 eq) in DCM (10 mL) were added TosCl (1.1 g, 5.77 mmol, 1.48 eq) and TEA (1.16 g, 11.50 mmol, 1.60 mL, 2.96 eq), DMAP (48.00 mg, 392.90 mol, 1.01e-1 eq). The mixture was stirred at 20° C. for 2 hours. The mixture was concentrated in vacuum. The residue was purified by column chromatography (0–30% ethyl acetate/petroleum ether) to give tert-butyl-4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate (1.5 g, 94% yield) as a colorless oil.

[0927] Step 2: Preparation of tert-butyl (S)-4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl piperidine-1-carboxylate. To a solution of tert-butyl-4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate (500 mg, 1.21 mmol, 1.64 eq) in MeCN (4 mL) and DMF (4 mL) were added Cs₂CO₃ (500 mg, 1.53 mmol, 2.07 eq) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (300 mg, 739.88 mol, 1 eq). The mixture was stirred at 60° C. for 12 hours. The reaction mixture was poured into water (10 mL). The aqueous phase was extracted with EA (30 mL×2). The combined organic phase was washed with brine (20 mL×3), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by column chromatography (0–100% ethyl acetate/petroleum ether). tert-butyl-4-[4-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]butyl]piperidine-1-carboxylate (400 mg, 84% yield) was obtained as a yellow oil.

[0928] Step 3: Preparation of 2-[[[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl-4-[4-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]butyl]piperidine-1-carboxylate (400 mg, 620.32 mol, 1 eq) in DCM (15 mL) was added TFA (7.68 g, 67.31 mmol, 5 mL, 108.51 eq). The mixture was stirred at 20° C. for 12 hours. The mixture was concentrated under vacuum. To the mixture was added NaHCO₃ to adjust the pH to 8–10. The mixture was poured into water (20 mL). The aqueous phase was extracted with DCM (20 mL×2). The combined organic phase was washed with brine (20 mL×2), dried with anhydrous Na₂SO₄, filtered and concentrated under vacuum. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 um; mobile phase: water (FA)-ACN; B %: 16%–46%, 10 min; Flow rate: 25 ml/min). 2-[[[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (200 mg, 59% yield) was obtained as a yellow solid.

[0929] Step 4: Preparation of (4-fluorophenyl)-[6-methoxy-2-(4-methoxyphenyl)benzothiophen-3-yl]methanone. To a solution of 6-methoxy-2-(4-methoxyphenyl)benzothiophene (13 g, 48.09 mmol, 1 eq) and 4-fluorobenzoylchloride (10 g, 63.07 mmol, 7.56 mL, 1.31 eq) in DCM (150 mL) was added portion-wise AlCl₃ (7.80 g, 58.50 mmol, 3.20 mL, 1.22 eq) at 0° C. The mixture was stirred at 25° C. for 1 hour. The reaction mixture was quenched by the addition of H₂O (150.0 mL) at 0° C., and then extracted with DCM (50.0 mL×3). The combined

organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , PE/EtOAc=1/0 to 8/1). Compound (4-fluorophenyl)-[6-methoxy-2-(4-methoxyphenyl) benzo thiophen-3-yl]methanone (12 g, 64% yield) was obtained as a yellow oil.

[0930] Step 5: Preparation of (4-fluorophenyl)-[6-hydroxy-2-(4-hydroxyphenyl)benzo thiophen-3-yl]methanone. To a solution of (4-fluorophenyl)-[6-methoxy-2-(4-methoxy phenyl)benzothiophen-3-yl]methanone (6 g, 15.29 mmol, 1 eq) in DCM (50 mL) was added dropwise BBr_3 (15.60 g, 62.27 mmol, 6 mL, 4.07 eq) at 0°C . and the mixture was stirred at 20°C . for 0.5 hour. The reaction mixture was quenched by the addition of sat. aq. NaHCO_3 (60.0 mL) at 0°C ., and then diluted with H_2O (10.0 mL) and extracted with DCM (40.0 mL $\times$ 2). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (0–30% ethyl acetate/petroleum ether). (4-fluorophenyl)-[6-hydroxy-2-(4-hydroxyphenyl) benzo thiophen-3-yl]methanone (5 g, 90% yield) was obtained as a yellow solid.

[0931] Step 6: Preparation of [6-benzyloxy-2-(4-benzyloxyphenyl)benzothiophen-3-yl]-(4-fluorophenyl)methanone. To a solution of (4-fluorophenyl)-[6-hydroxy-2-(4-hydroxy phenyl)benzothiophen-3-yl]methanone (5 g, 13.72 mmol, 1 eq) in CH_3CN (10 mL), THF (50 mL) were added K_2CO_3 (6 g, 43.41 mmol, 3.16 eq) and BnBr (10.08 g, 58.94 mmol, 7 mL, 4.30 eq). The mixture was stirred at 60°C . for 12 hours. The mixture was filtered and the filtrate was concentrated. The crude product was triturated with EA and PE to afford [6-benzyl oxy-2-(4-benzyloxyphenyl)benzothiophen-3-yl]-(4-fluorophenyl)methanone (5.9 g, 80% yield) as a yellow solid.

[0932] Step 7: Preparation of [6-benzyloxy-2-(4-benzyloxyphenyl)benzothiophen-3-yl]-(4-(2-hydroxyethoxy)phenyl)methanone. To a solution of ethylene glycol (700 mg, 11.28 mmol, 0.6 mL, 2.05 eq) in DMF (30 mL) was added NaH (550 mg, 13.75 mmol, 60% purity, 2.5 eq) at 0°C . After the addition, the mixture was stirred at 20°C . for 0.5 hours. Then [6-benzyloxy-2-(4-benzyloxyphenyl)benzothiophen-3-yl]-(4-fluorophenyl)methanone (3 g, 5.51 mmol, 1 eq) was added to the mixture and the resulting mixture was stirred at 20°C . for 0.5 hour. The mixture was then stirred at 90°C . for 12 hours. The reaction mixture was quenched by the addition of sat. aq. NH_4Cl (60.0 mL), and then diluted with H_2O (20.0 mL) and extracted with EtOAc (50.0 mL $\times$ 2). The combined organic layers were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (0–80% ethyl acetate/petroleum ether). [6-benzyloxy-2-(4-benzyloxy phenyl)benzothiophen-3-yl]-(4-(2-hydroxyethoxy)phenyl)methanone (1.4 g, 43% yield) was obtained as a yellow solid.

[0933] Step 8: Preparation of 2-[4-[6-benzyloxy-2-(4-benzyloxyphenyl) benzothiophene-3-carbonyl]phenoxy]ethyl-4-methylbenzenesulfonate. To a solution of [6-benzyloxy-2-(4-benzyloxyphenyl)benzothiophen-3-yl]-(4-(2-hydroxyethoxy)phenyl)methanone (1.4 g, 2.39 mmol, 1 eq) in DCM (15 mL) was added tosylchloride (700.00 mg, 3.67 mmol, 1.54 eq) and TEA (1.45 g, 14.37 mmol, 2 mL, 6.02 eq), DMAP (70.00 mg, 572.98 mol, 0.24 eq). The mixture was stirred at 20°C . for 12 hours. The mixture was concentrated in vacuum. The residue was purified by flash silica gel chromatography (0–50% Ethyl acetate/Petroleum ether). The residue was purified by a second time by column chromatography (gradient of 80–100% water (1% formic acid)/ACN). 2-(4-(6-(benzyloxy)-2-(4-(benzyloxy)phenyl)


benzo[b]thiophene-3-carbonyl)phenoxy)ethyl 4-methylbenzenesulfonate (800 mg, 45% yield) was obtained as a yellow solid.


[0934] Step 9: Preparation of 2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl 4-methylbenzenesulfonate. To a solution of 2-(4-(6-(benzyloxy)-2-(4-(benzyloxy)phenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl 4-methylbenzene sulfonate (500 mg, 674.87 mol, 1 eq) in THF (5 mL) was added Pd/C (100 mg, 10% w/w) under an N_2 atmosphere. The suspension was degassed and purged with H_2 3 times. The mixture was stirred under H_2 (50 psi) at 50°C . for 12 hours. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The crude product was used into the next step without further purification. 2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy) ethyl 4-methylbenzenesulfonate (400 mg, crude) was obtained as a yellow oil.

[0935] Step 10: Preparation of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl] methanone. To a solution of 2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl 4-methylbenzenesulfonate (400 mg, 713.47 mol, 1 eq) in CH_3CN (3 mL), and DMF (2 mL) was added LiBr (188.00 mg, 2.16 mmol, 54.34 L, 3.03 eq). The mixture was stirred at 60°C . for 12 hours. The mixture was concentrated in vacuum. The residue was purified by column chromatography (0–30% ethyl acetate/petroleum ether). [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzo thiophen-3-yl]methanone (300 mg, 90% yield) was obtained as a yellow oil.

[0936] Step 11: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)ethoxy)phenyl) methanone. To a solution of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzothiophen-3-yl]methanone (40 mg, 85.22 mol, 1 eq) in DMF (1 mL) were added DIEA (40 mg, 309.49 mol, 53.91 L, 3.63 eq) and 2-[[[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}.trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (20 mg, 36.7 mol, 0.4 eq). The mixture was stirred at 80°C . for 12 hours. The mixture was concentrated under vacuum. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150 $\times$ 25 mm $\times$ 10 μm ; mobile phase: water (FA)-ACN; B %: 27%–57%, 10 min; Flow rate: 25 ml/min). (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)ethoxy)phenyl) methanone (1–9, 20.41 mg, 24% yield) was obtained as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ (ppm) 8.20 (s, 1H), 8.03 (s, 1H), 7.65 (d, J=8.8 Hz, 2H), 7.33 (d, J=2.4 Hz, 1H), 7.28 (d, J=8.8 Hz, 2H), 7.24 (d, J=8.8 Hz, 1H), 7.17 (d, J=8.8 Hz, 2H), 7.12 (s, 1H), 6.91 (dd, J=8.8, 2.4 Hz, 4H), 6.84 (dd, J=8.8, 2.4 Hz, 1H), 6.67 (d, J=8.4 Hz, 2H), 4.61 (t, J=7.2 Hz, 1H), 4.07 (br t, J=5.6 Hz, 2H) 3.97 (br t, J=6.4 Hz, 2H), 3.92–3.85 (m, 1H), 3.83–3.77 (m, 1H), 2.86 (br d, J=11.2 Hz, 2H), 2.67–2.62 (m, 1H), 2.59 (s, 3H), 2.45 (br d, J=2.0 Hz, 1H), 2.41 (s, 3H), 2.36–2.30 (m, 1H), 1.95 (br t, J=10.4 Hz, 2H), 1.68 (br d, J=7.2 Hz, 1H), 1.64 (s, 3H), 1.59 (br d, J=12.0 Hz, 2H), 1.38 (br d, J=6.0 Hz, 2H), 1.24–1.05 (m, 5H). LC-MS: MS (ES⁺): RT=1.798 min, m/z=933.2[M+H⁺]. LCMS Method: 25.

Example 11—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl) methanone (I-10)





[0937] Step 1: Preparation of tert-butyl 2-(p-tolylsulfonyloxy)-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-hydroxy-7-azaspiro[3.5]nonane-7-carboxylate (1 g, 4.14 mmol, 1 eq) in DCM (10 mL) were added TosCl (1.2 g, 6.29 mmol, 1.52 eq), TEA (1.31 g, 12.93 mmol, 1.80 mL, 3.12 eq), DMAP (52.00 mg, 425.64 μ mol, 0.1 eq). The mixture was stirred at 20° C. for 12 hours. The mixture was concentrated in vacuum. The residue was purified by column chromatography (0~25% ethyl acetate/petroleum ether). Compound tert-butyl 2-(p-tolylsulfonyloxy)-7-azaspiro[3.5]nonane-7-carboxylate (1.3 g, 79% yield) was obtained as a black solid.

[0938] Step 2: Preparation of tert-butyl 2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-(p-tolylsulfonyloxy)-7-azaspiro[3.5]nonane-7-carboxylate (600 mg, 1.52 mmol, 1.54 eq) in MeCN (2 mL), DMF (4 mL) were added Cs₂CO₃ (650 mg, 1.99 mmol, 2.02 eq) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (400 mg, 986.50 μ mol, 1 eq). The mixture was stirred at 80° C. for 12 hours. The residue was poured into water (10 mL). The aqueous phase was extracted with EA (30 mL $\times$ 2). The combined organic phase was washed with brine (20 mL $\times$ 3), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by column chromatography (0~20% DCM/MeOH). Compound tert-butyl 2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]-7-azaspiro[3.5]nonane-7-carboxylate (450 mg, 73% yield) was obtained as a yellow solid.

[0939] Step 3: Preparation of 2-[[[(9S)-7-[4-(7-azaspiro[3.5]nonan-2-yloxy)phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 2-[4-[(9S)-4,


5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]-7-azaspiro[3.5]nonane-7-carboxylate (450 mg, 715.67 μ mol, 1 eq) in DCM (5 mL) was added TFA (10.75 g, 94.24 mmol, 7 mL, 131.67 eq). The mixture was stirred at 20° C. for 2 hours. The mixture was concentrated in vacuum. The mixture was added NaHCO₃ adjust to pH 8-10. The residue was poured into water (20 mL). The aqueous phase was extracted with DCM (20 mL $\times$ 2). The combined organic phase was washed with brine (20 mL $\times$ 2), dried with anhydrous Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150 $\times$ 25 mm $\times$ 10 μ m; mobile phase: water (FA)-ACN; B %: 19%-35%, 8 min; Flow rate: 25 ml/min). Compound 2-[[[(9S)-7-[4-(7-azaspiro[3.5]nonan-2-yloxy)phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (200 mg, 49% yield) was obtained as a white solid.

[0940] Step 4: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone. To a solution of 2-[[[(9S)-7-[4-(7-azaspiro[3.5]nonan-2-yloxy)phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (70 mg, 132.41 μ mol, 1 eq), [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl]methanone (77.00 mg, 164.06 μ mol, 1.24 eq) in DMF (1 mL) was added DIEA (74.20 mg, 574.13 μ mol, 0.1 mL, 4.34 eq). The mixture was stirred at 80° C. for 12 hours. The mixture was concentrated in vacuum. The residue was purified by prep-HPLC (FA condition: Instrument: ACS-WH-GX-E; column: Phenomenex Luna C18 150 $\times$ 25 mm $\times$ 10 μ m; mobile phase: water (FA)-ACN; B %: 25%-55%, 10 min; Flow rate: 25 ml/min; Column temperature: R. T.; Wavelength: 220 nm). (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-

(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl) methanone (I-10, 35.68 mg, 29% yield) was obtained as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆): δ (ppm) 8.27 (s, 1H), 8.03 (s, 1H), 7.65 (d, J=8.8 Hz, 2H), 7.33 (d, J=2.0 Hz, 1H), 7.25 (dd, J=11.2, 8.8 Hz, 3H), 7.17 (d, J=8.8 Hz, 2H), 7.12 (s, 1H), 6.91 (d, J=8.8 Hz, 2H), 6.86-6.80 (m, 3H), 6.67 (d, J=8.8 Hz, 2H), 4.73 (br t, J=6.8 Hz, 1H), 4.61 (t, J=7.2 Hz, 1H), 4.07 (br t, J=5.6 Hz, 2H), 3.93-3.73 (m, 3H), 2.62-2.61 (m, 1H), 2.59 (s, 3H), 2.41 (s, 3H), 2.40-2.23 (m, 6H), 1.73 (br dd, J=12.4, 6.0 Hz, 2H), 1.64 (s, 3H), 1.59-1.51 (m, 4H). LC-MS: MS (ES⁺): RT=1.687 min, m/z=917.4 [M+H⁺]. LCMS Method: 25.

Example 12—Synthesis of 4-(2-(4-(3-((1-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)piperidin-4-yl)methyl)azetidin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-11)

droxyphenyl)benzothiophen-3-yl]methanone (55.00 mg, 117.18 mol, 1.23 eq) in DMF (1 mL) was added DIEA (74.20 mg, 574.13 mol, 0.1 mL, 6.05 eq). The mixture was stirred at 80° C. for 12 hours. The mixture was concentrated under vacuum. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: water (FA)-ACN; B %: 23%-53%, 10 min; Flow rate: 25 ml/min). 4-(2-(4-(3-((1-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)piperidin-4-yl)methyl)azetidin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-11, 17.07 mg, 19% yield) was obtained as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 11.97 (br s, 1H), 9.88-9.69 (m, 1H), 7.65 (d, J=8.8 Hz, 2H), 7.52 (d, J=2.4 Hz, 1H), 7.36-7.32 (m, 2H), 7.27-7.22 (m, 3H), 7.19-7.14 (m, 2H), 6.91 (d, J=9.2 Hz, 2H), 6.85 (dd, J=8.8, 2.4 Hz, 1H), 6.79 (d, J=8.8 Hz, 2H), 6.72-6.66 (m, 3H), 6.34 (d, J=8.8 Hz, 2H), 6.23 (t, J=2.4 Hz, 1H), 5.04-4.89 (m, 1H), 4.08 (br t, J=5.6 Hz, 2H), 3.88 (br t, J=7.2 Hz, 2H), 3.54 (s, 3H), 2.86 (br d, J=11.2 Hz, 2H),




I-11


[0941] To a solution of 4-[5-(1-hydroxy-1-methyl-ethyl)-2-[4-[3-(4-piperidylmethyl) azetidin-1-yl]phenoxy]phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (50 mg, 94.94 mol, 1 eq), [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hy-

2.77-2.67 (m, 2H), 2.62 (br d, J=6.0 Hz, 2H), 1.96 (br t, J=10.8 Hz, 2H), 1.58-1.49 (m, 4H), 1.44 (s, 6H), 1.27-0.98 (m, 4H). LC-MS: MS (ES⁺): RT=1.528 min, m/z=915.4 [M+H⁺]. LCMS Method: 25.


Example 13—Synthesis of (5R,6S)-6-phenyl-5-(4-(4-((4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl) piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (I-12), (5R,6S)-6-phenyl-5-(4-(4-((4-((R)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl) piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (III-1), (5S,6R)-6-phenyl-5-(4-(4-((4-((R)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl) piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (III-2), and (5S,6R)-6-phenyl-5-(4-(4-((4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl) piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (III-3)




-continued



-continued



III-2



III-3

[0942] Step 1: Preparation of (+)-cis-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]trifluoro-methanesulfonate. To a solution of (i)-cis-4-(6-methoxy-2-phenyl-tetralin-1-yl)phenol (500 mg, 1.51 mmol, 1 eq) in THE (20 mL) was added LiHMDS (1 M, 2.50 mL, 1.65 eq) and 1,1,1-trifluoro-N-phenyl-N-(trifluoromethylsulfonyl)methanesulfonamide (800.00 mg, 2.24 mmol, 1.48 eq) under N₂ at 0° C. The mixture was stirred at 20° C. for 1 hour. The reaction mixture was quenched by addition H₂O 5 mL at 0° C., and then extracted with EA 8 mL (4 mL×2). The combined organic layers were washed with brine 5 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (0-15% Ethyl acetate/Petroleum ether). Compound (±)-cis-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]trifluoromethanesulfonate (650 mg, 93% yield) was obtained as a white solid.

[0943] Step 2: Preparation of (i)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine. A mixture of (i)-cis-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]trifluoromethanesulfonate (630.00 mg, 1.36 mmol, 1 eq), 4-(dimethoxymethyl) piperidine (315.00 mg, 1.98 mmol, 1.45 eq), SPhos Pd G3 (100.00 mg, 128.16 mol, 9.41e-2 eq), Cs₂CO₃ (1.10 g, 3.38 mmol, 2.48 eq) in dioxane (10 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 100° C. for 12 hours under N₂ atmosphere. The reaction mixture was diluted with brine 5 mL and extracted with EA 15 mL (5 mL×3). The combined organic layers were washed with brine 10 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (5:1 Petroleum ether: Ethyl acetate). Compound (±)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine (420 mg, 65% yield) was obtained as a brown gum.

[0944] Step 3: Preparation of (±)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde. To a solution of (i)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine (400 mg, 848.12 mol, 1 eq) in DCM (4 mL) was added BBr₃ (2.60 g, 10.38 mmol, 1 mL, 12.24 eq) at 0° C. The mixture was stirred at 20° C. for 1 hours. And then H₂O (4 mL) was added dropwise at 0° C. The resulting mixture was stirred at 20° C. for 2 hours. The reaction mixture was adjusted to pH 7-8 with aqueous

Na₂CO₃ and extracted with EA 15 mL (5 mL×3). The combined organic layers were washed with brine 10 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, Petroleum ether: Ethyl acetate=1:1). Compound (±)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (300 mg, 86% yield) was obtained as a brown solid.

[0945] Step 4: Preparation of tert-butyl 4-[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazine-1-carboxylate. A mixture of 2-[[[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (200 mg, 471.79 mol, 1 eq), tert-butyl piperazine-1-carboxylate (150.00 mg, 805.37 mol, 1.71 eq), SPhos Pd G3 (70.00 mg, 89.71 mol, 0.19 eq), Cs₂CO₃ (461.15 mg, 1.42 mmol, 3 eq) in dioxane (0.5 mL) was degassed and purged with N₂ three times, and then the mixture was stirred at 80° C. for 12 hours under N₂ atmosphere. The reaction mixture was diluted with H₂O 5 mL and extracted with EA 15 mL (5 mL×3). The combined organic layers were washed with brine 10 mL (5 mL×2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (10:1 DCM:MeOH). Compound tert-butyl 4-[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazine-1-carboxylate (210 mg, 78% yield) was obtained as a yellow solid.

[0946] Step 5: Preparation of 2-[[[4,5,13-trimethyl-7-(4-piperazin-1-ylphenyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 4-[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazine-1-carboxylate (200 mg, 348.61 mol, 1 eq) in DCM (2 mL) was added TFA (1.23 g, 10.77 mmol, 800 L, 30.89 eq). The mixture was stirred at 20° C. for 2 hours. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (FA condition: column: Phenomenex luna C18 150*25 mm*10 um; mobile phase: [water(FA)-ACN]; gradient: 8%-38% B over 9 min). Compound 2-[[[(9S)-4,5,13-trimethyl-7-(4-piper-


azin-1-ylphenyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl)methyl]oxazole (200 mg, 98% yield, TFA) was obtained as a yellow solid.


[0947] Step 6: Preparation of Four cis-Tetraline Diastereomers of 6-phenyl-5-(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol. To a solution of 2-[[4,5,13-trimethyl-7-(4-piperazin-1-ylphenyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl)methyl]oxazole (190 mg, 323.34 mol, 1 eq, TFA) and (±)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (150 mg, 364.49 mol, 1.13 eq) in MeOH (2 mL) was added AcOH (1.94 mg, 32.33 mol, 1.85 μL, 0.1 eq). After addition, the mixture was stirred at 20° C. for 0.5 hour, and then NaBH(OAc)₃ (137.06 mg, 646.68 mol, 2 eq), 1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (150 mg, 364.49 mol, 1.13 eq) was added. The resulting mixture was stirred at 20° C. for 12 hours. The reaction mixture was diluted with H₂O 5 mL and extracted with EA 15 mL (5 mL×3). The combined organic layers were washed with brine 10 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient: 21%-51% B over 10 min). A mixture of the four cis-tetraline diastereomers of 2-phenyl-1-[4-[4-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazin-1-yl)methyl]-1-piperidyl]

phenyl]tetralin-6-ol (100 mg, 115.06 mol, 35.58% yield) was obtained as a yellow solid.

[0948] Step 7: Preparation of Compounds I-12, III-1, III-2, and III-3. The yellow solid from the preceding step was separated by SFC (condition: column: (s,s) WHELK-01 (250 mm×30 mm, 10 μm); mobile phase: [CO₂-ACN/MeOH (0.1% NH₃H₂O)]; B %: 60%, isocratic elution mode). The second fraction was further separated by SFC (condition: column: DAICEL CHIRALCEL OD (250 mm×30 mm, 10 μm); mobile phase: [CO₂-ACN/MeOH (0.1% NH₃H₂O)]; B %: 55%, isocratic elution mode). The diastereomer that was found to be the most potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below) was obtained as a yellow solid (13.58 mg, 16% yield). ¹H NMR (400 MHz, CDCl₃) δ (ppm)=7.62 (s, 1H), 7.31 (br d, J=8.4 Hz, 2H), 7.19-7.09 (m, 3H), 7.04 (s, 1H), 6.85-6.75 (m, 5H), 6.70 (s, 1H), 6.61-6.50 (m, 3H), 6.29 (br d, J=8.4 Hz, 2H), 4.70 (t, J=7.2 Hz, 1H), 4.19 (br d, J=4.8 Hz, 1H), 4.15-3.99 (m, 2H), 3.54 (br d, J=8.8 Hz, 2H), 3.38-3.23 (m, 5H), 3.08-2.95 (m, 2H), 2.68-2.54 (m, 9H), 2.41 (s, 3H), 2.31 (br d, J=6.4 Hz, 2H), 2.19-2.14 (m, 1H), 1.84 (br d, J=13.6 Hz, 3H), 1.73 (s, 3H), 1.66 (br d, J=4.2 Hz, 1H), 1.40-1.27 (m, 2H). LC-MS: MS (ES⁺): RT=1.438 min, m/z=869.4 [M+H]⁺; LCMS Method: 25.

Example 14—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)ethoxy)phenyl)methanone (I-13)





I-13

[0949] Step 1: Preparation of tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-(2-hydroxyethyl)piperidine-1-carboxylate (1 g, 4.36 mmol, 1 eq) in DCM (20 mL) were added DMAP (106.55 mg, 872.16 mol, 0.2 eq), TosCl (1.66 g, 8.72 mmol, 2 eq) and TEA (1.32 g, 13.08 mmol, 1.82 mL, 3 eq). The mixture was stirred at 25° C. for 12 hours. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (0~21% Ethyl acetate/Petroleum ether). Compound tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate (1.5 g, 90% yield) was obtained as yellow oil.

[0950] Step 2: Preparation of tert-butyl 4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate (605.32 mg, 1.58 mmol, 1.6 eq) in CH₃CN (4 mL) and DMF (4 mL) was added Cs₂CO₃ (964.27 mg, 2.96 mmol, 3 eq) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (400 mg, 986.50 mol, 1 eq). The mixture was stirred at 60° C. for 2 hours. The reaction mixture was concentrated under reduced pressure to

remove solvent. The residue was diluted with H₂O 15 mL and extracted with EA 45 mL (15 mL×3). The combined organic layers were washed with brine 20 mL (10 mL×2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (0~100% Ethyl acetate/Petroleum ether). tert-butyl 4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate (468 mg, 77% yield) was obtained as yellow oil.


[0951] Step 3: Preparation of 2-[[[4-[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate (440 mg, 713.39 mol, 1 eq) in DCM (5 mL) was added TFA (1.54 g, 13.46 mmol, 1.00 mL, 18.87 eq). The mixture was stirred at 25° C. for 2 hours. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was diluted with NaHCO₃ 15 mL and extracted with DCM 45 mL (15 mL×3). The combined organic layers were dried over

Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient: 12%-42% B over 10 min). Compound 2-[[[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl)methyl]oxazole (368 mg, 99% yield) was obtained as a yellow solid.


[0952] Step 4: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)ethoxy)phenyl)methanone. To a solution of 2-[[[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl)methyl]oxazole (75 mg, 145.16 μmol, 1 eq) in DMF (1 mL) was added DIEA (56.28 mg, 435.49 μmol, 75.85 μL, 3 eq) and [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)methanone (81.76 mg, 174.20 μmol, 1.2 eq). The mixture was stirred at 80° C. for 12 hours. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (FA condition; column: Phenomenex

Luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient: 25%-55% B over 10 min). (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)ethoxy)phenyl)methanone (1-13, 27.6 mg, 21% yield) was obtained as a yellow gum. ¹H NMR (400 MHz, DMSO-d₆) δ (ppm) 9.95-9.54 (m, 1H), 8.04 (s, 1H), 7.65 (d, J=8.8 Hz, 2H), 7.34 (d, J=2.0 Hz, 1H), 7.29 (d, J=8.4 Hz, 2H), 7.25 (d, J=8.8 Hz, 1H), 7.20-7.16 (m, 2H), 7.13 (s, 1H), 6.93 (dd, J=8.8, 5.6 Hz, 4H), 6.85 (dd, J=8.8, 2.2 Hz, 1H), 6.68 (d, J=8.8 Hz, 2H), 4.62 (t, J=7.2 Hz, 1H), 4.12-3.99 (m, 4H), 3.92-3.85 (m, 1H), 3.84-3.77 (m, 1H), 2.87 (br d, J=10.8 Hz, 2H), 2.63 (br t, J=5.8 Hz, 2H), 2.60 (s, 3H), 2.42 (s, 3H), 1.97 (br t, J=10.8 Hz, 2H), 1.68-1.59 (m, 7H), 1.46-1.37 (m, 1H), 1.24-1.13 (m, 2H). LC-MS: MS (ES⁺): RT=1.663 min, m/z=905.4 [M+H⁺]. LCMS Method: 25.

Example 15—Synthesis of (S)-(2-(4-bromophenyl)-6-hydroxybenzo[b]thiophen-3-yl)(4-(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (I-14)



-continued



I-14

[0953] Step 1: Preparation of 1-(4-bromophenyl)-2-(3-methoxyphenyl)sulfanyl-ethanone. To a solution of KOH (88.82 g, 1.58 mol, 10 equiv) in EtOH (300 mL) was added 3-methoxybenzenethiol (24.41 g, 174.14 mmol, 21.40 mL, 1.1 equiv) at 25° C. The mixture was stirred at 25° C. for 30 min. Then the mixture was cooled to 0° C. and 2-bromo-1-(4-bromophenyl)ethanone (44 g, 158.31 mmol, 1 equiv) in EtOAc (300 mL) were added subsequently at 0° C. The mixture was stirred at 0° C. for 30 min. The reaction mixture was warmed to 25° C. and stirred for 4 h. The solvent was removed under reduced pressure. The residue was extracted with ethyl acetate (500 mL×2), washed with brine (300 mL),

and dried over anhydrous Na₂SO₄. The solvent was removed under reduced pressure. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate, 1/0 to 10/1) to afford 1-(4-bromophenyl)-2-(3-methoxyphenyl)sulfanyl-ethanone (43 g, 80% yield) was obtained as a white solid.

[0954] Step 2: Preparation of 2-(4-bromophenyl)-6-methoxy-benzothiophene. To a stirring solution of PPA (30 mL) at 80° C. was added 1-(4-bromophenyl)-2-(3-methoxyphenyl)sulfanyl-ethanone (3 g, 8.90 mmol, 1 equiv) in portions within 30 min. Then the reaction was heated to 90°

C. and stirred for 1.5 h. The solution was stirred at 135° C., for another 10 h. The reaction mixture was poured into ice water. The crude product was collected by filtration and the solid was washed with water to afford 2-(4-bromophenyl)-6-methoxy-benzothiophene (2.3 g, 80% yield) as a white solid.

[0955] Step 3: Preparation of [2-(4-bromophenyl)-6-methoxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone. To a solution of 2-(4-bromophenyl)-6-methoxy-benzothiophene (2.2 g, 6.89 mmol, 1 equiv) in DCM (20 mL) was added AlCl_3 (1.38 g, 10.34 mmol, 564.94 L, 1.5 equiv) and 4-fluorobenzoyl chloride (1.64 g, 10.34 mmol, 1.24 mL, 1.5 equiv) at 0° C. The mixture was stirred at 0° C. for 2 h. To a solution of NaHCO_3 (aq) (100 mL) was added the reaction mixture at 25° C., and then diluted with water 100 mL and extracted with EtOAc (100 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 (250 mm $\times$ 70 mm, 10 μm); mobile phase: [water (FA)-ACN]; gradient 30%-95% B over 45 min) to give [2-(4-bromophenyl)-6-methoxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (140 mg, 4% yield) as a white solid.

[0956] Step 4: Preparation of [2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone. To a solution of [2-(4-bromophenyl)-6-methoxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (140 mg, 317.24 μL , 1 equiv) in DCM (6 mL) was added BBr_3 (1 M, 317.24 μL , 1 equiv) at -50° C. under N_2 . Then the mixture was stirred at 25° C. for 2 h. The reaction was poured into water (70 mL). The aqueous phase was extracted with ethyl acetate (15 mL $\times$ 3). The reaction mixture was quenched by dropwise addition of NaHCO_3 (aq) 5 mL to pH 5 at 25° C. under N_2 , and then diluted with H_2O 20 mL and extracted with DCM (20 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , PE:EtOAc=3:1) to afford [2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (120 mg, 88% yield) as a yellow solid.

[0957] Step 5: Preparation of tert-butyl 4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-(4-hydroxybutyl)piperidine-1-carboxylate (2.5 g, 9.71 mmol, 1 equiv) in pyridine (20 mL) was added TosCl (5.56 g, 29.14 mmol, 3.0 equiv) at 0° C. Then the mixture was stirred at 25° C. for 4 h. The reaction mixture was partitioned between H_2O (100 mL) and EtOAc (100 mL $\times$ 3). The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate, 1/0 to 10/1) to afford tert-butyl 4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate (1.4 g, 35% yield) as a white oil.


[0958] Step 6: Preparation of tert-butyl 4-[4-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]butyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-[4-(p-tolylsulfonyloxy)butyl]piperidine-1-carboxylate (203.00 mg, 493.25 mol, 2.0 equiv) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-


yl]phenol (100 mg, 246.63 mol, 1 equiv) in DMF (3 mL) was added K_2CO_3 (68.17 mg, 493.25 mol, 2.0 equiv). The mixture was stirred at 100° C. for 3 h. The reaction mixture was partitioned between H_2O 50 mL and EtOAc (50 mL $\times$ 3). The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , DCM: MeOH=10:1) to afford tert-butyl 4-[4-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]butyl]piperidine-1-carboxylate (150 mg, 94% yield) as a yellow oil.

[0959] Step 7: Preparation of 2-[[4-[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 4-[4-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]butyl]piperidine-1-carboxylate (130 mg, 201.61 mol, 1 equiv) in DCM (3 mL) was added TFA (1.54 g, 13.46 mmol, 1.00 mL, 66.78 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was quenched by addition NaHCO_3 (aq) 30 mL at 25° C., and then diluted with H_2O (50 mL) and extracted with DCM (50 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue to afford 2-[[4-[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (100 mg, 91% yield) as a yellow oil.

[0960] Step 8: Preparation of (S)-(2-(4-bromophenyl)-6-hydroxybenzo[b]thiophen-3-yl)(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone. To a solution of 2-[[4-[(9S)-4,5,13-trimethyl-7-[4-[4-(4-piperidyl)butoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (60 mg, 110.15 mol, 1 equiv) and [2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (51.77 mg, 121.17 mol, 1.1 equiv) in DMSO (2 mL) was added DIEA (42.71 mg, 330.45 mol, 57.56 L, 3.0 equiv). The mixture was stirred at 110° C. for 12 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: YMC-Actus Triart C18 150 mm $\times$ 30 mm $\times$ 7 μm ; mobile phase: [water (FA)-ACN]; gradient 80%-100% B over 10 min) to afford (S)-(2-(4-bromophenyl)-6-hydroxybenzo[b]thiophen-3-yl)(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (1-14, 24.97 mg, 23% yield) as a yellow solid. ^1H NMR (400 MHz, DMSO-d_6) δ 9.87 (s, 1H), 8.03 (s, 1H), 7.57-7.48 (m, 4H), 7.37 (d, $J=2.0$ Hz, 1H), 7.32 (d, $J=8.8$ Hz, 2H), 7.28 (d, $J=8.8$ Hz, 2H), 7.23 (d, $J=8.8$ Hz, 1H), 7.12 (s, 1H), 6.92 (d, $J=8.8$ Hz, 2H), 6.88-6.80 (m, 3H), 4.66-4.58 (m, 1H), 4.01-3.95 (m, 2H), 3.94-3.85 (m, 3H), 3.84-3.76 (m, 1H), 2.81 (t, $J=12.0$ Hz, 2H), 2.59 (s, 3H), 2.41 (s, 3H), 1.74-1.66 (m, 4H), 1.63 (s, 3H), 1.53-1.36 (m, 3H), 1.27-1.22 (m, 2H), 1.15-1.02 (m, 2H). LC-MS: MS (ES^+): RT=3.536 min, $m/z=953.3$ [$\text{M}+\text{H}^+$], LC-MS Method: 25.

Example 16—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)phenyl)methanone (I-15)





I-15

[0961] Step 1: Preparation of tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-(2-hydroxyethyl)piperidine-1-carboxylate (24 g, 104.66 mmol, 1 equiv) in DCM (240 mL) was added 4-methylbenzenesulfonyl chloride (29.93 g, 156.99 mmol, 1.5 equiv) and DMAP (12.79 g, 104.66 mmol, 1 equiv). The mixture was stirred at 25° C. for 12 h. The reaction mixture was partitioned between H₂O 300 mL and DCM 300 mL. The organic phase was separated, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=30/1 to 10/1) to afford tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate (17 g, 42% yield) as a colorless oil.

[0962] Step 2: Preparation of afford tert-butyl 4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate. To a solution of tert-butyl 4-[2-(p-tolylsulfonyloxy)ethyl]piperidine-1-carboxylate (141.87 mg, 369.94 mol, 1.5 equiv) in MeCN (10 mL) was added K₂CO₃ (68.17 mg, 493.25 mol, 2 equiv) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (100 mg, 246.63 mol, 1 equiv). The mixture was stirred at 80° C. for 2 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to afford tert-butyl 4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate (130 mg, 85% yield) as a yellow oil.

[0963] Step 3: Preparation of 2-[[[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 4-[2-[4-[(9S)-4,


5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]piperidine-1-carboxylate (170 mg, 275.63 mol, 1 equiv) in DCM (3 mL) was added TFA (1.54 g, 13.46 mmol, 1 mL, 49 equiv). The mixture was stirred at 25° C. for 2 h. The reaction mixture was quenched by addition NaHCO₃ (aq) 10 mL at 25° C., adjust PH to 8, and then diluted with H₂O 20 mL and extracted with EA 40 mL (20 mL×2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 2-[[[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (140 mg, 98% yield) as a white solid.

[0964] Step 4: Preparation of [6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophen-3-yl]-[4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]-1-piperidyl]phenyl] methanone. To a solution of 2-[[[(9S)-4,5,13-trimethyl-7-[4-[2-(4-piperidyl)ethoxy]phenyl]-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (60 mg, 116.13 mol, 1 equiv) in DMSO (2 mL) was added DIEA (45.03 mg, 348.39 mol, 60.68 μL, 3 equiv) and (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl] methanone (61.85 mg, 116.13 mol, 1 equiv). The mixture was stirred at 110° C. for 12 h. The reaction mixture was partitioned between EtOAc (20 mL) and H₂O (20 mL). The organic phase was separated, washed with NaCl 20 mL (10 mL×2), filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to afford [6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophen-3-yl]-[4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]-1-piperidyl]phenyl] methanone (40 mg, 33% yield) as a white solid.


[0965] Step 5: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)phenyl)methanone. To a solution of [6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxy-phenyl)benzothiophen-3-yl]-[4-[4-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]ethyl]-1-piperidyl]phenyl]methanone (38 mg, 36.92 mol, 1 equiv) in MeOH (3 mL) and H₂O (0.3 mL) was added HCl (306.00 mg, 8.39 mmol, 0.3 mL, 227 equiv). The mixture was stirred at 60° C. for 0.5h. The reaction mixture was added HCl (4M) and adjust PH to 7, then partitioned between EA (10 mL) and H₂O (10 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: C18 150x30 mm; mobile phase: [water (FA)-ACN]; gradient of 60%-90% over 7 min) to afford

(S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethyl)piperidin-1-yl)phenyl)methanone (I-15, 16 mg, 49% yield) as a yellow solid. H NMR (400 MHz, DMSO-d₆) δ 9.72 (br dd, J=6.4, 2.4 Hz, 2H) 8.03 (s, 1H) 7.48-7.54 (m, 2H) 7.26-7.32 (m, 3H) 7.20 (d, J=8.4 Hz, 2H) 7.11-7.18 (m, 2H) 6.90-6.96 (m, 2H) 6.82 (dd, J=8.8, 2.8 Hz, 3H) 6.69 (d, J=8.8 Hz, 2H) 4.58-4.65 (m, 1H) 4.01-4.06 (m, 2H) 3.78-3.93 (m, 4H) 2.78-2.86 (m, 2H) 2.59 (s, 3H) 2.41 (s, 3H) 1.72-1.77 (m, 2H) 1.64 (s, 3H) 1.23 (br s, 5H). LC-MS: MS (ES⁺): RT=2.979 min, m/z=861.4 [M+H⁺]. LCMS Method: 01.

Example 17—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(7-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)heptyl)acetamide (I-16)



-continued



I-16

[0966] Step 1: Preparation of tert-butyl N-(7-hydroxyheptyl)carbamate. To a solution of 7-aminoheptan-1-ol (4 g, 30.48 mmol, 1 equiv) in THF (40 mL) was added (Boc)₂O (7.98 g, 36.58 mmol, 8.40 mL, 1.2 equiv). The mixture was stirred at 20° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (Silica, Eluent of 0–30% Ethyl acetate/Petroleum ether) to afford tert-butyl N-(7-hydroxyheptyl)carbamate (6.3 g, 89% yield) as a colorless oil.

[0967] Step 2: Preparation of tert-butyl N-[7-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophene-3-carbonyl]phenoxy]heptyl]carbamate. To a solution of tert-butyl N-(7-hydroxyheptyl)carbamate (260.60 mg, 1.13 mmol, 2 equiv) in THF (4 mL) was added portion wise NaH (56.33 mg, 1.41 mmol, 60% purity, 2.5 equiv) at 0° C. under N₂, the mixture was stirred at 0° C. for 0.5 h under N₂, then (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-yl]methanone (300 mg, 563.25 μmol, 1 equiv) was added. The mixture was stirred at 25° C. for 1 h under N₂. The reaction mixture was quenched by addition NH₄Cl (10


mL) at 25° C., and then extracted with EA 27 mL (9 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica, Eluent of 0–30% Ethyl acetate/Petroleum ether) to afford tert-butyl N-[7-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophene-3-carbonyl]phenoxy]heptyl]carbamate (300 mg, 71% yield) as a yellow solid.

[0968] Step 3: Preparation of [4-(7-aminoheptoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzothiophen-3-yl] methanone. Tert-butyl N-[7-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenoxy]heptyl]carbamate (300 mg, 403.26 μmol, 1 equiv) was added H₂O (0.25 mL), MeOH (2.5 mL) and HCl (0.25 mL). The mixture was stirred at 50° C. for 2 h. The reaction mixture was quenched by addition NaHCO₃ (9 mL) at 25° C., and then extracted with EtOAc (12 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue to afford [4-(7-aminoheptoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzothiophen-3-yl]methanone (230 mg, crude) as a yellow solid.


[0969] Step 4: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(7-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)heptyl)acetamide. To a solution of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetic acid (110 mg, 274.40 mol, 1 equiv) in DCM (2 mL) was added HOBT (55.61 mg, 411 mol, 1.5 equiv) and EDCI (78.90 mg, 411 mol, 1.5 equiv). Then the mixture was added the solution of [4-(7-aminoheptoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (195.75 mg, 411 mol, 1.5 equiv) in DCM (1 mL) and DIEA (106.39 mg, 823.19 mol, 143 L, 3 equiv). The mixture was stirred at 25° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=9:1). Then the residue was purified by prep-HPLC (column: Phenomenex Luna C18 150 mm×25 mm×10 um; mobile phase: [water(FA)-ACN]; gradient: 55%-85% B over 10 min) to afford (S)-2-

(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(7-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)heptyl)acetamide (I-16, 81.64 mg, 34% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 9.75 (d, J=17.2 Hz, 2H), 8.15 (t, J=5.6 Hz, 1H), 7.64 (d, J=8.8 Hz, 2H), 7.49-7.44 (m, 2H), 7.43-7.38 (m, 2H), 7.33 (d, J=2.4 Hz, 1H), 7.24 (d, J=8.8 Hz, 1H), 7.20-7.12 (m, 2H), 6.92-6.82 (m, 3H), 6.71-6.64 (m, 2H), 4.49 (dd, J=6.0, 8.0 Hz, 1H), 3.96 (t, J=6.4 Hz, 2H), 3.27-3.02 (m, 4H), 2.58 (s, 3H), 2.39 (s, 3H), 1.73-1.62 (m, 2H), 1.60 (s, 3H), 1.48-1.39 (m, 2H), 1.31 (d, J=3.6 Hz, 6H). LC-MS: MS (ES⁺): RT=2.766 min, m/z=858.4 [M+H⁺], LCMS Method: 25.

Example 18—Synthesis of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)acetamide (I-17)



-continued



[0970] Step 1: Preparation of Ethyl 2-[4-(tert-butoxycarbonylamino)butoxy]acetate. To a solution of tert-butyl N-(4-hydroxybutyl) carbamate (8 g, 42.27 mmol, 1.0 equiv) in DCM (100 mL) was added diacetoxyrhodium (467.08 mg, 2.11 mmol, 0.05 equiv) and ethyl 2-diazoacetate (48.23 g, 422.72 mmol, 44.45 mL, 10.0 equiv) at 0° C. The mixture was stirred at 25° C. for 12 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was purified by column chromatography (silica, 0~50% Ethyl acetate/Petroleum ether) to afford ethyl 2-[4-(tert-butoxycarbonylamino)butoxy]acetate (12.3 g, crude) as a yellow oil.

[0971] Step 2: Preparation of Tert-butyl N-[4-(2-hydroxyethoxy)butyl]carbamate. To a solution of ethyl 2-[4-(tert-butoxycarbonylamino)butoxy]acetate (12.3 g, 44.67 mmol, 1 equiv) in THF (100 mL) was added portion-wise LiAlH₄ (2.5 M, 44.67 mL, 2.5 equiv) at 0° C. under N₂. The mixture was stirred at 0° C. for 2 hr under N₂. The reaction mixture was quenched with water (4.5 mL), 15% NaOH (4.5 mL) and water (13.5 mL) in turn at 0° C. under N₂. The reaction

mixture was dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica, 0-50% Ethyl acetate/Petroleum ether) to afford tert-butyl N-[4-(2-hydroxyethoxy) butyl]carbamate (2.9 g, 27% yield) as a light yellow oil.

[0972] Step 3: Preparation of tert-butyl N-[4-[2-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethoxy]butyl]carbamate. To a solution of tert-butyl N-[4-(2-hydroxyethoxy) butyl]carbamate (262.82 mg, 1.13 mmol, 2 equiv) in THE (5 mL) was added portion-wise NaH (67.58 mg, 1.69 mmol, 60% purity, 3 equiv) at 0° C. under N₂, the mixture was stirred at 0° C. for 0.5 h under N₂, then (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophen-3-yl]methanone (300 mg, 563.25 mol, 1 equiv) was added. The mixture was stirred at 25° C. for 2 h under N₂. The reaction mixture was quenched by addition saturated NH₄Cl (50 mL) under N₂ at 25° C. and extracted with EtOAc (30 mL×3). The combined organic


layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica, 0-50% Ethyl acetate/Petroleum ether) to afford tert-butyl N-[4-[2-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophene-3-carbonyl]phenoxy]ethoxy]butyl]carbamate (400 mg, 95% yield) as a yellow oil.

[0973] Step 4: Preparation of [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone. To a solution of tert-butyl N-[4-[2-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethoxy]butyl]carbamate (250 mg, 335.16 mol, 1 equiv) in MeOH (3 mL), H_2O (0.3 mL) and HCl (0.3 mL). The mixture was stirred at 60° C. for 0.5h. The reaction mixture was concentrated under reduced pressure to afford [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (200 mg, crude, HCl) as a yellow oil.


[0974] Step 5: Preparation of (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)acetamide. To a solution of [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzothiophen-3-yl]methanone (115.40 mg, 224 mol, 1.2 equiv, HCl) and 2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]acetic acid (75 mg, 187 mol, 1 equiv) in DCM (2 mL) was added HOBT (37.92 mg, 280 mol, 1.5 eq), EDCI (53.80

mg, 280 mol, 1.5 eq) and DIEA (120.90 mg, 935 mol, 162.94 μL , 5 eq). The mixture was stirred at 25° C. for 12 h. The reaction mixture was diluted with H_2O (100 mL) and extracted with DCM (100 mL). The combined organic layers were washed with brine (15 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150 $\times$ 25 mm $\times$ 5 μm ; mobile phase: [water(FA)-ACN]; gradient: 53%-73% over 10 min) to afford (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)-N-(4-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)acetamide (I-17, 36 mg, 22% yield) as a yellow solid. ^1H NMR (400 MHz, DMSO-d_6) δ 9.79 (br d, $J=16.0$ Hz, 2H), 8.18 (br t, $J=5.6$ Hz, 1H), 7.64 (d, $J=8.8$ Hz, 2H), 7.49-7.44 (m, 2H), 7.43-7.38 (m, 2H), 7.34 (d, $J=2.0$ Hz, 1H), 7.24 (d, $J=8.8$ Hz, 1H), 7.19-7.14 (m, 2H), 6.91 (d, $J=8.8$ Hz, 2H), 6.84 (dd, $J=2.4$, 8.8 Hz, 1H), 6.70-6.64 (m, 2H), 4.49 (dd, $J=6.0$, 8.0 Hz, 1H), 4.14-4.05 (m, 2H), 3.70-3.63 (m, 2H), 3.44 (t, $J=6.4$ Hz, 2H), 3.28-3.16 (m, 2H), 3.16-3.04 (m, 2H), 2.58 (s, 3H), 2.39 (s, 3H), 1.59 (s, 3H), 1.56-1.43 (m, 4H). LC-MS: MS (ES^+): $\text{RT}=2.471$ min, $m/z=860.3$ [$\text{M}+\text{H}^+$]; LC-MS Method: 25.

Example 19—Synthesis of 4-(2-(4-(4-((1-(4-((1R,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-18)



-continued



I-18

[0975] Step 1: Preparation of Benzyl 4-[4-[4-(1-hydroxy-1-methyl-ethyl)-2-(6-methyl-7-oxo-1H-pyrrolo[2,3-c]pyridin-4-yl)phenoxy]phenyl]piperazine-1-carboxylate. A mixture of 4-[2-(4-bromophenoxy)-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (400 mg, 882 mol, 1 equiv), benzyl piperazine-1-carboxylate (600 mg, 2.72 mmol, 525.39 μL , 3.09 equiv), K_3PO_4 (561.90 mg, 2.65 mmol, 3 equiv), CuI (168.05 mg, 882 mol, 1 equiv) and 2-(2,6-dimethylanilino)-2-oxo-acetic acid (170.47 mg, 882 mol, 1 equiv) in DMSO (5 mL) was degassed and purged with N_2 three times, and then the mixture was stirred at 100°C for 12 hours under a N_2 atmosphere. The reaction mixture was diluted with H_2O (5 mL) and extracted with EA (5 mL $\times$ 3). The combined organic layers were washed with brine (5 mL $\times$ 2), dried over Na_2SO_4 , filtered

and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (Waters Xbridge C18 150 $\times$ 50 mm $\times$ 10 μm ; mobile phase: [water (NH_4HCO_3)-ACN]; gradient: 35-65% over 10 min) to afford benzyl 4-[4-[4-(1-hydroxy-1-methyl-ethyl)-2-(6-methyl-7-oxo-1H-pyrrolo[2,3-c]pyridin-4-yl)phenoxy]phenyl]piperazine-1-carboxylate (240 mg, 46% yield) as a white solid.


[0976] Step 2: Preparation of 4-[5-(1-hydroxy-1-methyl-ethyl)-2-(4-piperazin-1-ylphenoxy)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one. To a solution of benzyl 4-[4-[4-(1-hydroxy-1-methyl-ethyl)-2-(6-methyl-7-oxo-1H-pyrrolo[2,3-c]pyridin-4-yl)phenoxy]phenyl]piperazine-1-carboxylate (240 mg, 404 mol, 1 equiv) in MeOH (5 mL) was added Pd/C (50 mg, 47 mol, 10% purity) under N_2


atmosphere. The suspension was degassed and purged with H₂ three times. The mixture was stirred under H₂ (50 Psi) at 50° C. for 12 hours. The reaction mixture was filtered and concentrated under reduced pressure to give 4-[5-(1-hydroxy-1-methyl-ethyl)-2-(4-piperazin-1-ylphenoxy)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (180 mg, 97% yield) as a white solid.

[0977] Step 3: Preparation of 4-(2-(4-(4-((1-(4-((1R,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one. To a solution of 4-[5-(1-hydroxy-1-methyl-ethyl)-2-(4-piperazin-1-ylphenoxy)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (180.00 mg, 392 mol, 1.24 equiv) and 1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (130 mg, 315 mol, 1 equiv) in MeOH (1 mL) was added AcOH (18.97 mg, 315 mol, 18.08 μ L, 1 equiv). After addition, the mixture was stirred at 20° C. for 0.5 hour, and then NaBH(OAc)₃ (66.95 mg, 315 mol, 1 equiv), 1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (130 mg, 315 mol, 1 equiv) was added. The resulting mixture was stirred at 20° C. for 1 hour. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150 $\times$ 25 mm $\times$ 10 μ m; mobile phase: [water (FA)-ACN]; gradient: 21%-51% over 10 min). 4-(2-(4-(4-((1-(4-((1R,2S)-6-hydroxy-2-phenyl-1,2,3,4-

tetrahydronaphthalen-1-yl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-18, 100 mg, 37% yield) was obtained as a white solid. The white solid was further separated by SFC (column: DAICEL CHIRALPAK AS (250 mm $\times$ 30 mm $\times$ 10 μ m); mobile phase: [CO₂-ACN/i-PrOH (0.1% NH₃)]); 55% B, isocratic). 4-(2-(4-(4-((1-(4-((1R,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-4-yl)methyl)piperazin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-18, 22.3 mg, 27% yield) was isolated as an off-white solid. ¹H NMR (400 MHz, CDCl₃) δ (ppm)=10.05 (br s, 1H), 7.62 (d, J 2.4 Hz, 1H), 7.43-7.38 (m, 1H), 7.23 (t, J 2.8 Hz, 1H), 7.19-7.13 (m, 3H), 7.09 (s, 1H), 6.92 (d, J 8.8 Hz, 1H), 6.87-6.79 (m, 7H), 6.72 (d, J 2.8 Hz, 1H), 6.63-6.54 (m, 3H), 6.40 (t, J 2.4 Hz, 1H), 6.29 (d, J 8.8 Hz, 2H), 4.20 (d, J 4.8 Hz, 1H), 3.65 (s, 3H), 3.60-3.48 (m, 2H), 3.39-3.29 (m, 1H), 3.16 (br s, 4H), 3.08-2.97 (m, 2H), 2.66 (br d, J=18.8 Hz, 4H), 2.60-2.53 (m, 2H), 2.36 (br d, J 6.4 Hz, 2H), 2.18 (br d, J 4.8 Hz, 1H), 1.90-1.75 (m, 4H), 1.64 (s, 6H), 1.40-1.29 (m, 2H). LC-MS: MS (ES⁺): RT=1.510 min, m/z=854.4 [M+H]; LCMS Method: 25.

Example 20—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-19)





I-19

[0978] Step 1: Preparation of (9S)-7-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenyl]-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene. To a solution of tert-butyl 2-[[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-azaspiro[3.5]nonane-7-carboxylate (200 mg, 357 μ mol, 1 equiv) in DCM (3 mL) was added TFA (3.07 g, 27 mmol, 2.00 mL, 75 equiv). The mixture was stirred at 25° C. for 0.5 h. The reaction mixture was adjusted to pH 8-9 by NaHCO₃ (aq) and then diluted with H₂O (20 mL) and extracted with DCM (20 mL $\times$ 2). The combined organic layers were washed with brine (20 mL $\times$ 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford (9S)-7-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenyl]-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (160 mg, 97% yield) as a yellow solid.

[0979] Step 2: Preparation of 2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethyl 4-methylbenzenesulfonate. The hydrogenation flask was filled with N₂, and H₂ was slowly added in the N₂ atmosphere, then 2-[4-[6-benzoyloxy-2-(4-benzoyloxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethyl 4-methylbenzenesulfonate (430 mg, 580 μ mol, 1.0 equiv) and Pd/C (200 mg, 187 μ mol, 10% purity, 0.32 equiv) in THE (5 mL) was added

under N₂ atmosphere. The suspension was degassed and purged with H₂ three times. The mixture was stirred under H₂ (50 Psi) at 50° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, PE:EA=1:1) to afford 2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethyl 4-methylbenzenesulfonate (300 mg, 92% yield) as a white solid.


[0980] Step 3: Preparation of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone. To a solution of 2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]phenoxy]ethyl 4-methylbenzenesulfonate (300 mg, 535 μ mol, 1 equiv) in MeCN (5 mL) was added LiBr (140.81 mg, 1.62 mmol, 3.03 equiv). The mixture was stirred at 60° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=10/1 to 1/1) to afford [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (120 mg, 47% yield) as a white solid.


[0981] Step 4: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)ethoxyphenyl]methanone. To a solution of [4-(2-bromoethoxy)

phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (40 mg, 85 mol, 1 equiv) in DMF (1 mL) was added DIEA (38.55 mg, 298 mol, 52 L, 3.5 equiv) and (9S)-7-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenyl]-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (58.76 mg, 127 mol, 1.5 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150x25 mmx10 um; mobile phase: [water (FA)-I]; gradient: 28-58% over 9 min) to afford (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-19, 29 mg, 38% yield) as a yellow solid. ¹H NMR (400

MHz, DMSO-d₆) δ 9.80 (s, 1H) 9.75 (s, 1H) 7.70 (d, J=8.0 Hz, 2H) 7.31-7.36 (m, 3H) 7.25 (d, J=8.8 Hz, 1H) 7.18 (dd, J=8.0, 5.2 Hz, 4H) 6.96 (d, J=8.8 Hz, 2H) 6.85 (dd, J=8.8, 2.4 Hz, 1H) 6.64-6.70 (m, 2H) 4.34 (d, J=5.2 Hz, 2H) 4.20 (q, J=6.4 Hz, 1H) 3.43-3.46 (m, 2H) 3.37 (s, 4H) 2.66-2.74 (m, 3H) 2.59 (s, 3H) 2.39 (s, 3H) 1.86 (d, J=6.4 Hz, 3H) 1.74-1.80 (m, 1H) 1.68 (d, J=8.0 Hz, 3H) 1.61 (s, 3H) 1.44-1.56 (m, 2H) 1.22-1.29 (m, 2H). LC-MS: MS (ES⁺): RT=1.868 min, m/z=848.5 [M+H]⁺, LCMS Method: 25.

Example 21—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-((4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-20)





I-20


[0982] Step 1: Preparation of tert-butyl 2-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of 2-[[[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (500 mg, 1.18 mmol, 1 equiv) and tert-butyl 2-ethynyl-7-azaspiro[3.5]nonane-7-carboxylate (735.25 mg, 2.95 mmol, 2.5 equiv) in MeCN (15 mL) was added Cs₂CO₃ (768.59 mg, 2.36 mmol, 2.0 equiv) and DavePhos Pd G₃ (90.03 mg, 117.95 mol, 0.1 equiv) under N₂. The mixture was stirred at 90° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate 1/0 to 1/1) to afford tert-butyl 2-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate (700 mg, 93% yield) as a white solid.

[0983] Step 2: Preparation of 2-[[[(9S)-7-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole. To a solution of tert-butyl 2-[2-[4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate (350 mg, 549 mol, 1 equiv) in DCM (3 mL) was added TFA (1.54 g, 13.46 mmol, 1 mL, 24 equiv). The mixture was stirred at 25° C. for 1h. The reaction mixture was quenched by addition NaHCO₃ 25 mL at 25° C., and then diluted with H₂O (20 mL) and extracted with EtOAc (20 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 2-[[[(9S)-7-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]


phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (200 mg, 67% yield) as a brown solid.

[0984] Step 3: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-((4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl) methanone. To a solution of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (40 mg, 85 mol, 1 equiv) and 2-[[[(9S)-7-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (45.74 mg, 85 mol, 1 equiv) in DMF (1 mL) was added DIEA (40.00 mg, 309 mol, 53.91 μL, 3.63 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water (FA)-ACN]; gradient: 30-60% over 9 min) to afford (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-((4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl) methanone (I-20, 10.41 mg, 12% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 9.80 (s, 1H), 9.76 (s, 1H), 8.04 (s, 1H), 7.70 (d, J=8.8 Hz, 2H), 7.42-7.37 (m, 2H), 7.36-7.34 (m, 2H), 7.32 (s, 1H), 7.25 (d, J=8.8 Hz, 1H), 7.17 (d, J=8.8 Hz, 2H), 7.13 (s, 1H), 6.98 (d, J=8.8 Hz, 2H), 6.86 (J=2.4, 8.8 Hz, 1H), 6.71-6.65 (m, 2H), 4.68 (t, J=7.2 Hz, 1H), 4.39 (s, 2H), 3.98-3.77 (m, 2H), 3.54-3.39 (m, 3H), 3.06-2.89 (m, 2H), 2.60 (s, 3H), 2.41 (s, 3H), 2.38-2.32 (m, 1H), 2.18 (t, J=9.2 Hz, 1H), 2.00-1.92 (m, 3H), 1.88-1.72 (m, 3H), 1.60 (s, 3H), 1.32-1.17 (m, 2H). LC-MS: MS (ES⁺): RT=1.260 min, m/z=925.3 [M+H⁺]. LCMS Method: 5-95.

Example 22—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (I-21)



-continued



I-21

[0985] Step 1: Preparation of [4-[4-(4-hydroxybutyl)-1-piperidyl]phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone. To a solution of (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl) benzothiophen-3-yl]methanone (300 mg, 563 μmol, 1 equiv) in DMF (3 mL) was added Cs₂CO₃ (550.56 mg, 1.69 mmol, 3 equiv) and 4-(4-piperidyl)butan-1-ol (177.15 mg, 1.13 mmol, 2 equiv). The mixture was stirred at 50° C. for 12 h. LCMS showed that the desired compound was detected. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM: MeOH=9:1) to afford [4-[4-(4-hydroxybutyl)-1-piperidyl]

phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone (220 mg, 58% yield) as a brown solid.

[0986] Step 2: Preparation of 4-[1-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothio-
phene-3-carbonyl]phenyl]-4-piperidyl]butyl 4-methylben-
zenesulfonate. To a solution of [4-[4-(4-hydroxybutyl)-1-
piperidyl]phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-
tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]
methanone (200 mg, 298 mol, 1 equiv) in DCM (2 mL) was
added DMAP (3.65 mg, 30 mol, 0.1 equiv) and TEA (75.53
mg, 746 mol, 103.89 μ L, 2.5 equiv). Then the mixture was


added TosCl (85.38 mg, 448 mol, 1.5 equiv). The mixture was stirred at 25° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (Silica, 0-50% Ethyl acetate/Petroleum ether) to afford 4-[1-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenyl]-4-piperidyl]butyl 4-methylbenzenesulfonate (230 mg, 93% yield) as a yellow solid.

[0987] Step 3: Preparation of (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone. To a solution of 4-[1-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenyl]-4-piperidyl]butyl 4-methylbenzenesulfonate (101.62 mg, 123 mol, 1 equiv) in DMF (1.5 mL) was added K₂CO₃ (34.09 mg, 246 mol, 2 equiv) and 4-[(9S)-4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (50 mg, 123 mol, 1 equiv). The mixture was stirred at 80° C. for 3 h. The reaction mixture was quenched by addition H₂O (3 mL) at 25° C., and then extracted with EA (5 mL×3). The combined organic layers were washed with aqueous NaCl 15 mL (5 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to afford (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (100 mg, 77% yield) as a brown solid.

[0988] Step 4: Preparation of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (I-21, 32.74 mg, 39% yield) as a yellow solid.

trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone. To a solution of (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (100 mg, 94 mol, 1 equiv) in MeOH (1 mL) was added HCl (0.1 mL) and H₂O (0.1 mL). The mixture was stirred at 50° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150×25 mm×5 um; mobile phase: [water(FA)-ACN]; gradient: 65%-85% B over 10 min) to afford (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(4-(4-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)butyl)piperidin-1-yl)phenyl)methanone (I-21, 32.74 mg, 39% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 9.86-9.56 (m, 2H), 8.04 (s, 1H), 7.52 (d, J=8.8 Hz, 2H), 7.33-7.26 (m, 3H), 7.22-7.16 (m, 3H), 7.13 (s, 1H), 6.93 (d, J=9.2 Hz, 2H), 6.82 (dd, J=2.0, 8.8 Hz, 3H), 6.72-6.66 (m, 2H), 4.63 (t, J=7.2 Hz, 1H), 3.98 (t, J=6.4 Hz, 2H), 3.93-3.84 (m, 3H), 3.84-3.77 (m, 1H), 2.79 (t, J=11.6 Hz, 2H), 2.60 (s, 3H), 2.41 (s, 3H), 1.70 (d, J=8.4 Hz, 4H), 1.64 (s, 3H), 1.49-1.37 (m, 3H), 1.28-1.22 (m, 2H), 1.15-1.03 (m, 2H). LC-MS: MS (ES⁺): RT=2.815 min, m/z=889.4 [M+H⁺]; LCMS Method: 25.

Example 23—Synthesis of 4-(2-(4-(3-((1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenyl)piperidin-4-yl)methyl)azetidin-1-yl)phenoxy)-5-(2-hydroxy-propan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-22)




DCM:MeOH=10:1) to afford 4-[5-(1-hydroxy-1-methyl-ethyl)-2-[4-[3-[[1-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenyl]-4-piperidyl]methyl]pyridine-1-yl]phenoxy]phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridine-7-one (50 mg, 58% yield) as a green solid.

[0991] Step 3: Preparation of 4-(2-(4-(3-((1-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenyl)piperidin-4-yl)methyl)azetidin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridine-7-one. To a solution of 4-[5-(1-hydroxy-1-methyl-ethyl)-2-[4-[3-[[1-[4-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophene-3-carbonyl]phenyl]-4-piperidyl]methyl]pyridine-1-yl]phenoxy]phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridine-7-one (35 mg, 34 mol, 1 equiv) in THE (0.5 mL) was added H₂O (0.1 mL) and PPTS (11.00 mg, 44 mol, 1.3 equiv). The mixture was stirred at 60° C. for 0.5 h. The reaction mixture was added H₂O (10 mL), pH was adjusted to 9-10 by adding saturated NaHCO₃ (aq) and extracted with DCM (10 mL×3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and

concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150×25 mm×5 um; mobile phase: [water(FA)-ACN]; gradient: 50%-70% B over 10 min) to afford 4-(2-(4-(3-((1-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenyl)piperidin-4-yl)methyl)azetidin-1-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridine-7-one (I-22, 15 mg, 50% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 11.98 (s, 1H), 9.73 (d, 2H, J=9.2 Hz), 7.52 (dd, 3H, J=2.8, 5.2H), 7.38-7.30 (m, 2H), 7.28-7.15 (m, 5H), 6.86-6.76 (m, 5H), 6.74-6.65 (m, 3H), 6.34 (d, 2H, J=9.2H), 6.23 (t, 1H, J=2.4 Hz), 4.99 (s, 1H), 3.89 (t, 4H, J=7.2 Hz), 3.54 (s, 3H), 3.30 (s, 2H), 2.85-2.70 (m, 3H), 1.66 (d, 2H, J=11.6 Hz), 1.57-1.47 (m, 3H), 1.44 (s, 6H), 1.19-1.03 (m, 2H). LC-MS: MS (ES⁺): RT=2.111 min, m/z=871.5 [M+H⁺], LCMS Method: 10.


Example 24—Synthesis of (S)-4-(4-(2-(2-(3,5-difluoro-4-((1R,3R)-2-(2-fluoro-2-methyl propyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenoxy)ethoxy) ethoxy) phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-23)



[0992] To a solution of 3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenol (70 mg, 0.18 mmol, 1 equiv) and (S)-4-(4-(2-(2-bromoethoxy)ethoxy) phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (105.84 mg, 0.21 mmol, 1.2 equiv) in ACN (5 mL) was added K_2CO_3 (74.72 mg, 0.54 mmol, 3 equiv). The mixture was stirred at 80° C. for 10 h. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (10 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: YMC-Actus Triart C18 150 $\times$ 30 mm $\times$ 7 μ m; mobile phase: [water (FA)-ACN]; gradient: 55%-85% B over 10 min) to afford (S)-4-(4-(2-(2-(3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-1,3,4,9-tetrahydro-1H-pyrido[3,4-





b]indol-1-yl)phenoxy) ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-23, 70.65 mg, 46% yield) as a yellow solid. 1H NMR (400 MHz, DMSO- d_6) δ 10.50 (s, 1H), 7.42-7.31 (m, 3H), 7.17 (d, $J=7.6$ Hz, 1H), 7.04-6.87 (m, 4H), 6.67 (d, $J=11.2$ Hz, 2H), 5.11 (s, 1H), 4.20-4.07 (m, 5H), 3.83-3.75 (m, 4H), 3.55-3.47 (m, 2H), 2.91-2.77 (m, 3H), 2.58 (s, 3H), 2.39 (s, 3H), 1.85 (d, $J=6.4$ Hz, 3H), 1.64 (s, 3H), 1.22-1.01 (m, 9H). LC-MS: MS (ES $^+$): RT=2.125 min, m/z =680.4 [M-116], m/z =796.6; LC-MS Method: 25.

Example 25—Synthesis of 4-(2-(4-((7-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)-7-azaspiro[3.5]nonan-2-yl) methyl) phenoxy) -5-(2-hydroxypropan-2-yl) phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-24)






307

-continued



-continued



I-24

[0993] Step 1: Preparation of [4-(2, 2-dimethoxyethoxy)phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone. To a solution of 2,2-dimethoxyethanol (398.48 mg, 3.76 mmol, 2 equiv) in THF (11 ml) was added portion wise NaH (187.75 mg, 4.69 mmol, 60% purity, 2.5 equiv) at 0° C. under N₂, the mixture was stirred at 0° C. for 0.5 h under N₂, then (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone (1 g, 1.88 mmol, 1 equiv) was added. The mixture was stirred at 25° C. for 1 h under N₂. The reaction mixture was quenched by addition saturated NH₄Cl (10 ml) under N₂ at 25° C., and then extracted with EA (12 ml×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by flash silica gel chromatography (0-30% Ethyl acetate/Petroleum ether) to afford [4-(2, 2-dimethoxyethoxy)phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone (1 g, 86% yield) as a yellow oil.

[0994] Step 2: Preparation of [4-(2, 2-dimethoxyethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone. To a solution of [4-(2,2-dimethoxyethoxy)phenyl]-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzothiophen-3-yl]methanone (980 mg, 1.58 mmol, 1 equiv) in MeOH (10 ml) was added HCl (1 ml) and H₂O (1 ml). The mixture was stirred at 50° C. for 1 h. The reaction mixture was concentrated under reduced pressure to afford [4-(2, 2-dimethoxyethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (710 mg, crude) as a yellow solid.

[0995] Step 3: Preparation of 2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-carbonyl]phenoxy]acetaldehyde. To a solution of [4-(2,2-dimethoxyethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (710 mg, 1.58 mmol, 1 equiv) in dioxane (7.5 ml) was added HCl (12 M, 2.5 ml, 19 equiv). The mixture was stirred at 25° C. for 1 hr. The residue was purified by flash silica gel chromatography (0-50% Ethyl acetate/Petro-

leum ether) to afford tert-butyl 2-[(4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]phenoxy]acetaldehyde (340 mg, 53% yield) as a yellow solid.

[0996] Step 4: Preparation of tert-butyl 2-[(4, 4, 5, 5-tetramethyl-1, 3, 2-dioxaborolan-2-yl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of 4,4,5,5-tetramethyl-2-[(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methyl]-1,3,2-dioxaborolane (13.44 g, 50.14 mmol, 1.2 equiv) in THE (180 ml) was added LDA (2 M, 31.34 ml, 1.5 equiv) at -75°C . under N_2 atmosphere and the reaction mixture was stirred for 2 h. A solution of tert-butyl 2-oxo-7-azaspiro[3.5]nonane-7-carboxylate (10 g, 41.79 mmol, 1 equiv) in THE (20 ml) was added below -70°C . After 1 h, the mixture was stirred under N_2 at 25°C . for 15 h. The reaction mixture was quenched by addition H_2O (80 ml), and then extracted with EA (100 ml $\times$ 3). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate 1/0 to 20/1) to afford tert-butyl 2-[(4, 4, 5, 5-tetramethyl-1, 3, 2-dioxaborolan-2-yl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate (7.9 g, 52% yield) as a colorless oil.

[0997] Step 5: Preparation of tert-butyl 2-[(4-benzyloxyphenyl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-[(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate (2 g, 5.51 mmol, 1 equiv) and 1-benzyloxy-4-bromo-benzene (2.17 g, 8.26 mmol, 1.5 equiv) in dioxane (40 ml) and H_2O (10 ml) was added K_2CO_3 (1.52 g, 11.01 mmol, 2 equiv) and $\text{Pd}(\text{dppf})\text{Cl}$ (402.82 mg, 550 μmol , 0.1 equiv) was degassed and purged with N_2 three times, and then the mixture was stirred at 90°C . for 16 h under N_2 atmosphere. The reaction mixture was quenched by addition H_2O (20 ml), and then extracted with EA (30 ml $\times$ 3). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate, 1/0 to 20/1) to afford tert-butyl 2-[(4-benzyloxyphenyl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate (1.51 g, 65% yield) as a yellow oil.

[0998] Step 6: Preparation of tert-butyl 2-[(4-hydroxyphenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-[(4-benzyloxyphenyl)methylene]-7-azaspiro[3.5]nonane-7-carboxylate (1.51 g, 3.60 mmol, 1 equiv) in MeOH (20 ml) was added Pd/C (10%, 0.5 g) under N_2 atmosphere. The suspension was degassed and purged with H_2 three times. The mixture was stirred under H_2 (50 Psi) at 25°C . for 16 hr. The reaction mixture was filtered and concentrated under reduced pressure to give a residue that was diluted with H_2O (30 ml) and extracted with EA (30 ml $\times$ 3). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate, 1/0 to 10/1) to afford tert-butyl 2-[(4-hydroxyphenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate (0.88 g, 73% yield) as a colorless oil.

[0999] Step 7: Preparation of tert-butyl 2-[(4-(2-bromo-4-methoxycarbonylphenoxy)phenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-[(4-hydroxyphenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate (0.88 g, 2.66 mmol, 1 equiv) in DMF (50 ml) was added

Cs_2CO_3 (2.60 g, 7.97 mmol, 3 equiv) and methyl 3-bromo-4-fluoro-benzoate (804.32 mg, 3.45 mmol, 1.3 equiv). The mixture was stirred at 80°C . for 16 hr. The reaction mixture was quenched by addition H_2O (50 ml), and then extracted with EA (50 ml $\times$ 3). The combined organic layers were washed with brine, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate, 1/0 to 10/1) to afford tert-butyl 2-[(4-(2-bromo-4-methoxycarbonylphenoxy)phenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate (0.97 g, 67% yield) as a colorless oil.

[1000] Step 8: Preparation of 2-[(4-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-3-bromo-phenyl]propan-2-ol. A mixture of tert-butyl 2-[(4-(2-bromo-4-methoxycarbonylphenoxy)phenyl)methyl]-7-azaspiro[3.5]nonane-7-carboxylate (0.97 g, 1.78 mmol, 1 equiv) in DCM (50 ml) was degassed and purged with N_2 three times, and then to the mixture was added drop-wise MeMgBr (3 M, 8.91 ml, 15 equiv) at -70°C . under an N_2 atmosphere. The resulting mixture was stirred at 40°C . for 16 hours. The reaction mixture was quenched by adding to NH_4Cl :aq (100 ml) at $0-5^{\circ}\text{C}$. under N_2 , and then mixture was extracted with DCM:MeOH (10:1, 150 ml $\times$ 3). The organic layer was dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford 2-[(4-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-3-bromo-phenyl]propan-2-ol (700 mg, 88% yield) as a yellow gum.

[1001] Step 9: Preparation of 4-[2-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-c]pyridin-7-one. A mixture of 2-[(4-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-3-bromo-phenyl]propan-2-ol (700 mg, 1.58 mmol, 1 equiv), 6-methyl-1-(p-tolylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[2,3-c]pyridin-7-one (674.65 mg, 1.58 mmol, 1 equiv), K_2CO_3 (478.93 mg, 3.47 mmol, 2.2 equiv), [2-(2-aminophenyl)phenyl]palladium(1+); bis(1-adamantyl)-butyl-phosphane:methanesulfonate (114.71 mg, 157.51 μmol , 0.1 equiv) in THF (20 ml) and H_2O (2 ml) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 70°C . for 16 hr under N_2 atmosphere. The mixture was concentrated to afford 4-[2-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-c]pyridin-7-one (900 mg, 1.35 mmol, 85% yield) as a yellow oil.


[1002] Step 10: Preparation of 4-[2-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one. To a solution of 4-[2-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-c]pyridin-7-one (900 mg, 1.35 mmol, 1 equiv) in MeOH (40 ml) was added KOH (1.67 g, 29.7 mmol, 22 equiv). The mixture was stirred at 30°C . for 1 hr. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 (250 $\times$ 70 mm $\times$ 10 μm); mobile phase: [water(FA)-ACN]; gradient: 15%-45% B over 20 min) to afford 4-[2-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (290 mg, 41% yield) as a white solid.


[1003] Step 11: Preparation of 4-[2-[4-[[7-[2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]

phenoxy]ethyl]-7-azaspiro[3.5]nonan-2-yl)methyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one. To a solution of 4-[2-[4-(7-azaspiro [3.5]nonan-2-ylmethyl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo [2,3-c]pyridin-7-one (40 mg, 78.2 μ mol, 1 equiv) in MeOH (1 ml) and DCM (1 ml) was added NaOAc (12.83 mg, 156 μ mol, 2 equiv). The mixture was stirred at 25° C. for 1h, then the mixture was added 2-[4-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophene-3-carbonyl]phenoxy]acetaldehyde (79.04 mg, 195 μ mol, 2.5 equiv), the mixture was stirred at 25° C. for 1 h, then the mixture was added NaBH(OAc)₃ (49.71 mg, 234 μ mol, 3equiv) and the mixture was stirred at 25° C. for 1 h, after the mixture was added NaBH₃CN (9.83 mg, 156 μ mol, 2 equiv) and HOAc (104.90 mg, 1.75 mmol, 0.1 ml, 22.34 equiv), then the mixture was stirred at 25° C. for 14 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150×25 mm×5 μ m; mobile phase: [water(FA)-ACN]; gradient: 30%-50% B over 10 min) to afford 4-(2-(4-((7-(2-(4-(6-hydroxy-2-(4-hydroxy-

phenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)-7-azaspiro[3.5]nonan-2-yl)methyl]phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-24, 14.6 mg, 19% yield) as a yellow solid. ¹H NMR (400 MHz, methanol-d₄) δ =7.73 (d, J=8.8 Hz, 2H), 7.65 (d, J=2.4 Hz, 1H), 7.48 (dd, J=2.4, 8.6 Hz, 1H), 7.43 (d, J=8.8 Hz, 1H), 7.31 (d, J=2.8 Hz, 1H), 7.26 (d, J=2.0 Hz, 1H), 7.20-7.15 (m, 2H), 7.14 (s, 1H), 7.01 (d, J=8.4 Hz, 1H), 6.94 (d, J=8.4 Hz, 2H), 6.90-6.85 (m, 3H), 6.67-6.59 (m, 4H), 6.35 (d, J=2.8 Hz, 1H), 4.63-4.52 (m, 1H), 4.33-4.21 (m, 2H), 3.61-3.56 (m, 3H), 3.28-3.25 (m, 1H), 3.16-2.79 (m, 4H), 2.62-2.54 (m, 2H), 2.49-2.33 (m, 1H), 1.96-1.77 (m, 4H), 1.68 (br t, J=5.2 Hz, 2H), 1.59 (s, 6H), 1.49 (br dd, J=8.8, 11.6 Hz, 2H) LC-MS: MS (ES⁺): RT=2.095 min, m/z=900.3 [M+H⁺], LC-MS Method: 25.

Example 26—Synthesis of (S)-4-(4-(2-(2-(2-(3,5-difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenoxy)ethoxy)ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-25)





I-25

[1004] Step 1: Preparation of 3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenol. To a solution of 2-fluoro-N-[(1R)-2-(1H-indol-3-yl)-1-methyl-ethyl]-2-methyl-propan-1-amine (113 mg, 0.45 mmol, 1 equiv) and 2,6-difluoro-4-hydroxy-benzaldehyde (71.94 mg, 0.45 mmol, 1 equiv) in toluene (2 mL) was added AcOH (54.65 mg, 0.90 mmol, 2 equiv). The mixture was stirred at 80° C. for 16 h. The reaction mixture was quenched by addition aqueous saturated sodium bicarbonate (10 mL) and then diluted with water (10 mL) and extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate, 3/1) to afford 3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenol (136 mg, 71% yield) as a yellow solid.

[1005] Step 2: Preparation of 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethanol. To a solution of 3,5-difluoro-4-[(1R,3R)-2-(2-

fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenol (136 mg, 0.35 mmol, 1 equiv) and 2-[2-(2-bromoethoxy)ethoxy]ethanol (82.06 mg, 0.38 mmol, 1.1 equiv) in ACN (3 mL) was added K₂CO₃ (48.39 mg, 0.35 mmol, 1 equiv). The mixture was stirred at 80° C. for 10 h. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, Petroleum ether/Ethyl acetate, 1/1) to afford 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethanol (135 mg, 60% yield) as a yellow solid.


[1006] Step 3: Preparation of 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethyl 4-methylbenzenesulfonate. To a solution of 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethanol (132 mg, 0.25 mmol, 1 equiv) in

CH_2Cl_2 (3 mL) was added TosCl (62.84 mg, 0.30 mmol, 1.3 equiv), TEA (51.32 mg, 0.50 mmol, 2 equiv) and DMAP (3.10 mg, 25.36 mol, 0.1 eq). The mixture was stirred at 25° C. for 2 h. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (10 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , dichloromethane/methanol, 3/1) to afford 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethyl 4-methylbenzenesulfonate (98 mg, 51% yield) as a yellow oil.


[1007] Step 4: Preparation of (S)-4-(4-(2-(2-(2-(3,5-difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenoxy)ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of 2-[2-[2-[3,5-difluoro-4-[(1R,3R)-2-(2-fluoro-2-methyl-propyl)-3-methyl-1,3,4,9-tetrahydropyrido[3,4-b]indol-1-yl]phenoxy]ethoxy]ethoxy]ethyl 4-methylbenzenesulfonate (95 mg, 0.14 mmol, 1 equiv) and (S)-4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenol (47.65 mg, 0.14 mmol, 1 equiv) in ACN (5 mL) was added K_2CO_3 (58.37 mg, 0.4 mmol, 3 equiv). The mixture was stirred at 80° C. for 10 h. The

reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL $\times$ 3). The combined organic layers were washed with brine (10 mL $\times$ 3), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: YMC-Actus Triart C18 150 $\times$ 30 mm $\times$ 7 μm ; mobile phase: [water (FA)-ACN]; gradient: 50%-80% B over 10 min) to afford (S)-4-(4-(2-(2-(2-(3,5-difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenoxy)ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (I-25, 13.52 mg, 11% yield) as a yellow solid. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) 10.51 (s, 1H), 7.41-7.33 (m, 3H), 7.16 (d, $J=7.6$ Hz, 1H), 7.01-6.88 (m, 4H), 6.66 (d, $J=10.8$ Hz, 2H), 5.11 (d, $J=0.8$ Hz, 1H), 4.10 (d, $J=3.2$ Hz, 4H), 3.76-3.70 (m, 4H), 3.59 (s, 4H), 2.89-2.82 (m, 3H), 2.68-2.66 (m, 3H), 2.34-2.31 (m, 3H), 1.85 (d, $J=6.4$ Hz, 3H), 1.64 (s, 3H), 1.25-1.00 (m, 12H). LC-MS: MS (ES^+): RT=1.821 min, m/z =841.7 [$\text{M}+\text{H}^+$], LC-MS Method: 25.

Example 27—Synthesis of (6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone (I-26)



-continued



[1008] Step 1: Preparation of tert-butyl 2-methylsulfonyloxy-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-hydroxy-7-azaspiro[3.5]nonane-7-carboxylate (2 g, 8.29 mmol, 1 equiv) in DCM (30 mL) was added TEA (2.10 g, 20.72 mmol, 2.88 mL, 2.5 equiv) and MsCl (1.34 g, 11.70 mmol, 905 μ L, 1.41 equiv) was added at 0° C. The mixture was stirred at 20° C. for 2 h. The reaction mixture was poured into ice water (50 mL) and then diluted with DCM (50 mL) and extracted with DCM (50 mL $\times$ 2). The combined organic layers were washed with citric acid (100 mL) and brine (30 mL $\times$ 2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford tert-butyl 2-methylsulfonyloxy-7-azaspiro[3.5]nonane-7-carboxylate (2.6 g, 98% yield) as a white solid.

[1009] Step 2: Preparation of tert-butyl 2-[4-(4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenoxy]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of 4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenol (200 mg, 493 mol, 1 equiv) in DMF (7 mL) was added K₂CO₃ (136.34 mg, 986 mol, 2 equiv) and tert-butyl 2-methylsulfonyloxy-7-azaspiro[3.5]nonane-7-carboxylate (787.76 mg, 2.47 mmol, 5 equiv). The mixture was stirred at 100° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH, 10:1) to afford tert-butyl 2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonane-7-carboxylate (240 mg, 77% yield) as a white solid.


[1010] Step 3: Preparation of 2-((4-(4-((7-azaspiro[3.5]nonan-2-yl)oxy)phenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)methyl)oxazole. To a solution of tert-butyl 2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonane-7-carboxylate (240 mg, 381 mol, 1 equiv) in DCM (3 mL) was added TFA (3.68 g, 32.31 mmol, 2.40 mL, 84.65 equiv). The mixture was stirred at 25° C. for 2 h. The reaction mixture was quenched by addition NaHCO₃ (aq) (10 mL) at 25° C., adjusted pH to 8, and then diluted with H₂O (20 mL) and extracted with EA (20 mL $\times$ 2). The combined organic fractions were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 2-((4-(4-((7-azaspiro[3.5]nonan-2-yl)oxy)phenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)methyl)oxazole (200 mg, 99% yield) as a white solid.

[1011] Step 4: Preparation of (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone. To a solution of 2-((4-(4-((7-azaspiro[3.5]nonan-2-yl)oxy)phenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)methyl)oxazole (200 mg, 378 mol, 1 equiv) and (4-fluorophenyl)-[6-tetrahydropyran-2-yloxy-2-(4-tetrahydropyran-2-yloxyphenyl)benzo[b]thiophen-3-yl]methanone (201.50 mg, 378 mol, 1 equiv) in DMSO (2 mL) was added Cs₂CO₃ (246.52 mg, 756 mol, 2 equiv). The mixture was stirred at 100° C. for 12 h. The reaction mixture was partitioned between EA (50 mL) and


H₂O (50 mL). The organic phase was separated, washed with brine (30 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH, 10:1) to afford (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone (200 mg, 50% yield) as a white solid.

[1012] Step 5: Preparation of (6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone (isomer 1 and isomer 2). To a solution of (6-((tetrahydro-2H-pyran-2-yl)oxy)-2-(4-((tetrahydro-2H-pyran-2-yl)oxy)phenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone (150 mg, 144 mol, 1 equiv) in MeOH (0.1 mL) was added HCl (102 mg, 2.80 mmol, 0.1 mL, 100% purity, 19.4 equiv) and H₂O (1 mL). The mixture was stirred at 60° C. for 0.5h. The reaction mixture was quenched by addition NaHCO₃ (aq) (10 mL) at 25° C., adjust pH to 8, and then diluted with H₂O (20 mL) and extracted with EA (20 mL $\times$ 2). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150 $\times$ 25 mm $\times$ 10 μ m; mobile phase: [water (FA)-ACN]; gradient: 58%-88% B over 9 min) to afford a white solid. The racemate was resolved by SFC ((s,s)-WHELK-O1 (250 mm $\times$ 30 mm $\times$ 10 μ m); mobile phase: [CO₂-ACN/EtOH(0.1% NH₃H₂O)]; 65% B, isocratic elution mode) to give isomer 1 as a white solid (peak 1, 18 mg, 44% yield) and isomer 2 of (6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)-7-azaspiro[3.5]nonan-7-yl)phenyl)methanone as a white solid (peak 2, 20 mg, 49% yield). ¹H NMR (400 MHz, DMSO-d₆) δ 9.73 (br d, J=9.2 Hz, 2H) 8.03 (s, 1H) 7.52 (d, J=8.8 Hz, 2H) 7.31 (d, J=2.0 Hz, 1H) 7.27 (br d, J=8.8 Hz, 2H) 7.15-7.22 (m, 3H) 7.12 (s, 1H) 6.79-6.87 (m, 5H) 6.65-6.71 (m, 2H) 4.73-4.82 (m, 1H) 4.61 (t, J=7.6 Hz, 1H) 3.76-3.90 (m, 2H) 3.25 (br d, J=2.8 Hz, 2H) 2.59 (s, 3H) 2.52-2.54 (m, 2H) 2.41 (s, 5H) 1.77-1.84 (m, 2H) 1.53-1.65 (m, 7H). ¹H NMR (400 MHz, DMSO-d₆) δ 9.68-9.79 (m, 2H) 8.03 (d, J=0.8 Hz, 1H) 7.52 (d, J=8.8 Hz, 2H) 7.31 (d, J=2.0 Hz, 1H) 7.27 (d, J=8.4 Hz, 2H) 7.15-7.22 (m, 3H) 7.12 (d, J=0.8 Hz, 1H) 6.79-6.87 (m, 5H) 6.68 (d, J=8.8 Hz, 2H) 4.78 (br t, J=6.8 Hz, 1H) 4.61 (t, J=7.6 Hz, 1H) 3.75-3.92 (m, 2H) 3.26 (dt, J=4.0, 2.2 Hz, 2H) 2.59 (s, 3H) 2.54 (s, 2H) 2.39-2.45 (m, 5H) 1.80 (br dd, J=12.8, 6.4 Hz, 2H) 1.64 (s, 7H).

Example 28—Synthesis of (S)-2,3,6,9-tetramethyl-4-(4-(4-((6-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (1-27)



-continued



I-27

[1013] Step 1: Preparation of tert-butyl 4-(6-((8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridine-3-yl)piperazine-1-carboxylate. To a solution of (8R)-6-(5-bromopyridin-2-yl)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinoline (1.00 g, 2.35 mmol, 1.00 equiv), tert-butyl piperazine-1-carboxylate (1.31 g, 7.05 mmol, 3.00 equiv), potassium tert-butoxide (1.32 g, 11.76 mmol, 5.00 equiv), [2-(2-aminophenyl)phenyl]-methylsulfonyloxy-palladium; dicyclohexyl-[2-(2,6-diisopropoxyphenyl)phenyl] phosphane (393 mg, 0.47 mmol, 0.20 equiv) in dioxane (20 mL) was stirred at 100°C for 10 h under nitrogen atmosphere. The mixture was filtered and the filtrate was concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography

(petroleum ether/ethyl acetate=6/1 to 1/1) to give tert-butyl 4-(6-((8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridine-3-yl)piperazine-1-carboxylate (450 mg, 36% yield) as a yellow solid.

[1014] Step 2: Preparation of (8R)-8-methyl-6-(5-(piperazin-1-yl)pyridine-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline. To a solution of tert-butyl 4-(6-((8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-3H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridine-3-yl)piperazine-1-carboxylate (150 mg, 0.28 mmol, 1.00 equiv) in dichloromethane (1 mL) was added trifluoroacetic acid (767 mg, 6.73 mmol, 23.81 equiv). The mixture was stirred at 25°C for 10 min. Ethyl acetate (2 mL) and saturated sodium bicarbonate solution (2 mL) were

added, then the mixture was extracted with ethyl acetate (2 mL×3). The organic layers was washed with brine (2 mL×3), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give (8R)-8-methyl-6-(5-(piperazin-1-yl)pyridine-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (120 mg, 98% yield) as a red oil, which was used into the next step without further purification.


[1015] Step 3: Preparation of (S)-4-(4-(4-(dimethoxymethyl)piperidin-1-yl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. A mixture of (9S)-7-(4-chlorophenyl)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (100 mg, 0.28 mmol, 1.00 equiv), 4-(dimethoxymethyl)piperidine (223 mg, 1.40 mmol, 5.00 equiv), dicyclohexyl-[2-(2,6-dimethoxyphenyl)phenyl]phosphane; methanesulfonate; (2-phenylanilino)palladium(1+), (22 mg, 0.03 mmol, 0.10 equiv) and cesium carbonate (183 mg, 0.56 mmol, 2.00 equiv) in dioxane (1 mL) was degassed and purged with nitrogen for 3 times, and then the mixture was stirred at 90° C. for 3 h under nitrogen atmosphere. Ethyl acetate (5 mL) and water (5 mL) were added before the mixture was extracted with ethyl acetate (5 mL×3). The combined organic layers were washed with brine (5 mL×3), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to get the residue. The residue was purified by prep-TLC (dichloromethane/methanol=10/1) to afford (S)-4-(4-(4-(dimethoxymethyl)piperidin-1-yl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (100 mg, 73% yield) as a yellow oil.


[1016] Step 4: Preparation of (S)-1-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidine-4-carbaldehyde. To a solution of (S)-4-(4-(4-(dimethoxymethyl)piperidin-1-yl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (100 mg, 0.21 mmol, 1.00 equiv) in dichloromethane (1 mL) was added trifluoroacetic acid (768 mg, 6.73 mmol, 32 equiv). The mixture was stirred at 25° C. for 10 min. The mixture was concentrated under reduced pressure to give (S)-1-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperi-

dine-4-carbaldehyde (110 mg, 96% yield, trifluoroacetate) as a yellow oil, which was used into the next step without further purification.

[1017] Step 5: Preparation of (S)-2,3,6,9-tetramethyl-4-(4-(4-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of (S)-1-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidine-4-carbaldehyde (110 mg, 0.20 mmol, 1.00 equiv, trifluoroacetate) in dichloromethane (2 mL) was added sodium triacetoxyborohydride (128 mg, 0.60 mmol, 3.00 equiv) and (8R)-8-methyl-6-(5-(piperazin-1-yl)pyridine-2-yl)-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinoline (86 mg, 0.20 mmol, 1.00 equiv). The mixture was stirred at 25° C. for 12 h. The mixture was concentrated under reduced pressure to get the residue. The residue was purified by prep-TLC (dichloromethane/methanol=10/1) to get the crude product. The crude product was purified by prep-SFC (column: DAICEL CHIRALCEL OD (250 mm×30 mm×10 μm); mobile phase: [CO₂—I/EtOH (0.1% NH₃H₂O)]; 40/60 isocratic elution mode) give a crude product, and then the crude product was further purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water(FIACN)]; gradient: 10%-40% B over 8 min) to afford (S)-2,3,6,9-tetramethyl-4-(4-(4-(4-(6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-yl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (1-27, 38.7 mg, 22% yield) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 12.98 (s, 1H), 8.14-8.02 (m, 2H), 7.33-7.19 (m, 4H), 7.10 (d, J=8.8 Hz, 1H), 6.89 (d, J=8.8 Hz, 2H), 6.82 (d, J=8.8 Hz, 1H), 5.00 (s, 1H), 4.12 (q, J=6.2 Hz, 1H), 3.82-3.76 (m, 2H), 3.59-3.45 (m, 3H), 3.30-3.20 (m, 3H), 3.15 (s, 4H), 3.07-2.95 (m, 2H), 2.89-2.82 (m, 1H), 2.77-2.69 (m, 2H), 2.57 (s, 3H), 2.47-2.46 (m, 1H), 2.40 (s, 3H), 2.18 (d, J=6.4 Hz, 2H), 1.84 (d, J=6.4 Hz, 3H), 1.81-1.75 (m, 2H), 1.70 (s, 3H), 1.25-1.13 (m, 2H), 1.09 (d, J=6.4 Hz, 3H). LC-MS: MS (ES⁺): RT=1.289 min, m/z=848.6 [M+H]⁺, LC-MS Method: 10.

Example 29—Synthesis of (S)-N-(4-(2-(4-(2-(4-bromophenyl)-6-hydroxybenzo[b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetamide (I-28)





I-28

[1018] Step 1: Preparation of [2-(4-bromophenyl)-6-[tert-butyl(dimethyl)silyl]oxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone. To a solution of [2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (350 mg, 819 mol, 1.0 equiv) in DCM (5 mL) was added TBSCl (185.19 mg, 1.23 mmol, 151 μL , 1.5 equiv) and IMIDAZOLE (66.92 mg, 983 mol, 1.2 equiv). The mixture was stirred at 25°C. for 4 h. The reaction mixture was partitioned between H_2O 50 mL and EtOAc (50 mL $\times$ 3). The

organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=1/0 to 5/1) to afford [2-(4-bromophenyl)-6-[tert-butyl(dimethyl)silyl]oxy-benzothiophen-3-yl]-(4-fluorophenyl)methanone (300 mg, 67% yield) as a yellow solid.

[1019] Step 2: Preparation of tert-butyl N-[4-[2-[4-[2-(4-bromophenyl)-6-hydroxy-benzothiophene-3-carbonyl]phe-


noxy]ethoxy]butyl]carbamate. To a solution of tert-butyl N-[4-(2-hydroxyethoxy)butyl]carbamate (193.86 mg, 831 mol, 1.5 equiv) in THF (8 mL) was added NaH (44.32 mg, 1.11 mmol, 60% purity, 2 equiv) at 0° C. under N₂. The mixture was stirred at 0° C. for 1 h. Then added [2-(4-bromophenyl)-6-[tert-butyl(dimethyl)silyl]oxy-benzo thiophene-3-yl]-(4-fluorophenyl)methanone (300 mg, 554 mol, 1 equiv) to the mixture at 0° C. under N₂. The mixture was stirred at 60° C. for 11 h. The reaction mixture was quenched by addition H₂O (2 mL) at 25° C., and then diluted with H₂O (20 mL) and extracted with EtOAc (20 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to afford tert-butyl N-[4-[2-[4-[2-(4-bromophenyl)-6-hydroxy-benzothiophene-3-carbonyl]phenoxy]ethoxy]butyl]carbamate (210 mg, 59% yield) as a white solid.


[1020] Step 3: Preparation of [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]methanone. To a solution of tert-butyl N-[4-[2-[4-[2-(4-bromophenyl)-6-hydroxy-benzothiophene-3-carbonyl]phenoxy]ethoxy]butyl]carbamate (190 mg, 297 mol, 1 equiv) in DCM (6 mL) was added HCl/dioxane (4 M, 2 mL, 27 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was quenched by addition of NaHCO₃ (aq) (15 mL) at 25° C., and then diluted with H₂O (50 mL) and extracted with EtOAc (50 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue to afford [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]methanone (150 mg, 93% yield) as a yellow oil.

[1021] Step 4: Preparation of (S)—N-(4-(2-(4-(2-(4-bromophenyl)-6-hydroxybenzo [b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-

6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetamide. To a solution of [4-[2-(4-aminobutoxy)ethoxy]phenyl]-[2-(4-bromophenyl)-6-hydroxy-benzothiophen-3-yl]methanone (150 mg, 278 mol, 1 equiv) and (S)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetic acid (111.26 mg, 278 mol, 1 equiv) in DCM (5 mL) was added DIEA (143.48 mg, 1.11 mmol, 193 μL, 4.0 equiv), EDCI (159.61 mg, 832 mol, 3.0 equiv) and HOBt (56.25 mg, 416 mol, 1.5 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was partitioned between H₂O 50 mL and EtOAc (50 mL×3). The organic phase was separated, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1). The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient: 62%-92% B over 9 min) to afford (S)—N-(4-(2-(4-(2-(4-bromophenyl)-6-hydroxybenzo[b]thiophene-3-carbonyl)phenoxy)ethoxy)butyl)-2-(4-(4-chlorophenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)acetamide (I-28, 66.3 mg, 25% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 10.13-9.69 (m, 1H), 8.17 (t, J=5.6 Hz, 1H), 7.66 (d, J=8.8 Hz, 2H), 7.54-7.49 (m, 2H), 7.47-7.43 (m, 2H), 7.42-7.37 (m, 3H), 7.32-7.26 (m, 3H), 6.93 (d, J=9.2 Hz, 2H), 6.88 (dd, J=2.4, 8.8 Hz, 1H), 4.49 (dd, J=6.0, 8.4 Hz, 1H), 4.16-4.05 (m, 2H), 3.71-3.62 (m, 2H), 3.44 (s, 2H), 3.27-3.17 (m, 2H), 3.15-3.04 (m, 2H), 2.58 (s, 3H), 2.38 (s, 3H), 1.59 (s, 3H), 1.55-1.43 (m, 4H). LC-MS: MS (ES⁺): RT=1.494 min, m/z=924.4 [M+H⁺]. LCMS Method: 50.

Example 30—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-29)






[1022] Step 1: Preparation of tert-butyl (S)-2-((4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of (S)-4-(4-bromophenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (500 mg, 1.25 mmol, 1 equiv) and tert-butyl 2-ethynyl-7-azaspiro[3.5]nonane-7-carboxylate (777 mg, 3.11 mmol, 2.5 equiv) in MeCN (15 mL) was added DavePhos Pd G₃ (95 mg, 125 mol, 0.1 equiv) and Cs₂CO₃ (812 mg, 2.49 mmol, 2.0 equiv) under N₂. The mixture was stirred at 90° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to afford tert-butyl (S)-2-((4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonane-7-carboxylate (600 mg, 84% yield) as a white solid.

[1023] Step 2: Preparation of (S)-4-(4-((7-azaspiro[3.5]nonan-2-yl)ethynyl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of tert-butyl (S)-2-((4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonane-7-carboxylate (300 mg, 526 mol, 1 equiv) in DCM (3 mL) was added TFA (1.54 g, 13.5 mmol, 1 mL, 26 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was quenched by addition NaHCO₃ (15 mL) at 25° C., and then diluted with H₂O (20 mL) and extracted with DCM (20 mL×3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford (S)-4-(4-((7-azaspiro[3.5]nonan-2-yl)ethynyl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (240 mg, 97% yield) as a yellow oil.

[1024] Step 3: Preparation of (S)-(6-hydroxy-2-(4-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-((4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone. To a solution of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)methanone (60 mg, 127.84 mol, 1 equiv) and (S)-4-(4-((7-azaspiro[3.5]nonan-2-yl)ethynyl)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (60.04 mg, 127 mol, 1 equiv) in DMF (1.5 mL) was added DIEA (49.6 mg, 383 mol, 66.80 μL, 3.0 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was diluted with H₂O (50 mL) and extracted with EtOAc (50 mL×3). The combined organic phases dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water (FA)-ACN]; gradient: 30%-60% B over 9 min) to afford (S)-(6-hydroxy-2-(4-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-((4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)ethynyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-29, 33.6 mg, 29% yield) as a yellow solid. H NMR (400 MHz, DMSO-d₆) δ 9.87-9.62 (m, 2H), 8.14 (s, 1H), 7.65 (d, J=8.8 Hz, 2H), 7.40 (s, 3H), 7.33 (d, J=2.0 Hz, 1H), 7.24 (d, J=8.8 Hz, 1H), 7.17 (d, J=8.4 Hz, 2H), 6.92 (d, J=8.8 Hz, 2H), 6.84 (J=2.0, 8.8 Hz, 1H), 6.67 (d, J=8.4 Hz, 2H), 4.22 (q, J=6.8 Hz, 1H), 4.09 (t, J=5.6 Hz, 2H), 2.59 (s, 3H), 2.54 (s, 4H), 2.39 (s, 7H), 2.17 (t, J=10.0 Hz, 2H), 1.93-1.79 (m, 5H), 1.67-1.51 (m, 7H). LC-MS: MS (ES⁺): RT=1.960 min, m/z=858.3 [M+H⁺]. LCMS Method: 25.


Example 31—Synthesis of (S)-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yl)(4-(2-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl)methanone (I-30)




[1025] To a solution of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophen-3-yl]methanone (40 mg, 85.22 μ mol, 1 equiv) in DMF (1 mL) was added (S)-2-((4-(7-azaspiro[3.5]nonan-2-yl)methyl)phenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-6-yl)methyl)oxazole (44.89 mg, 85 μ mol, 1 equiv), DIEA (27.54 mg, 213 μ mol, 37 μ L, 2.5 equiv). The mixture was stirred at 100° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150 $\times$ 25 mm $\times$ 5 μ m; mobile phase: [water(FA)-ACN]; gradient: 23%-53% B over 10 min) to afford (S)-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophen-3-yl)(4-(2-

(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)ethoxy)phenyl) methanone (I-30, 46.4 mg, 59% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-*d*₆) δ 9.75 (d, *J*=18.4 Hz, 2H), 8.14 (s, 1H), 8.04 (s, 1H), 7.65 (d, *J*=8.8 Hz, 2H), 7.34 (d, *J*=2.4 Hz, 1H), 7.27-7.22 (m, 3H), 7.19-7.14 (m, 4H), 7.13 (s, 1H), 6.91 (d, *J*=8.8 Hz, 2H), 6.85 (dd, *J*=2.0, 8.8 Hz, 1H), 6.67 (d, *J*=8.8 Hz, 2H), 4.65 (t, *J*=7.6 Hz, 1H), 4.09 (d, *J*=5.2 Hz, 2H), 3.95-3.75 (m, 2H), 2.76-2.64 (m, 4H), 2.60 (s, 3H), 2.41 (s, 6H), 2.36-2.30 (m, 2H), 1.77 (t, *J*=9.6 Hz, 2H), 1.60 (s, 3H), 1.56-1.35 (m, 6H). LC-MS: MS (ES⁺): RT=1.978 min, *m/z*=915.4 [M+H⁺], LC-MS Method: 25.

Example 32—Synthesis of 4-(2-(4-((7-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo [b]thiophene-3-carbonyl)phenoxy)ethyl)-7-azaspiro[3.5]nonan-2-yl)ethynyl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-31)



-continued



I-31

[1026] Step 1: Preparation of tert-butyl 2-(methoxymethylene)-7-azaspiro [3.5]nonane-7-carboxylate. A mixture of methoxymethyl(triphenyl)phosphonium chloride (35.81 g, 105 mmol, 2 equiv) in THF (400 ml) added tBuOK (9.38 g, 84 mmol, 1.6 equiv) in THF (70 ml) at 0° C. under N₂ atmosphere, and then the mixture was stirred at 20° C. for 3 h under N₂ atmosphere. Add tert-butyl 2-oxo-7-azaspiro [3.5]nonane-7-carboxylate (12.5 g, 52 mmol, 1 equiv) in THF (30 ml) to the reaction mixture, then the reaction was stirred at 70° C. for 3 hours. The reaction mixture was quenched by addition H₂O (100 mL), and then extracted with EA (150 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 10/1). Compound tert-butyl 2-(methoxymethylene)-7-azaspiro [3.5]nonane-7-carboxylate (9.18 g, 65% yield) was obtained as a colorless oil.

[1027] Step 2: Preparation of tert-butyl 2-formyl-7-azaspiro [3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-(methoxymethylene)-7-azaspiro [3.5]nonane-7-carboxylate (21.9 g, 81.9 mmol, 1 equiv) in ACN (520 ml) and H₂O (130 ml) was added TFA (7.47 g, 65.5 mmol, 4.87 ml, 0.8 equiv). The mixture was stirred at 25° C. for 1.5 h. TLC (PE:EA=3:1) indicated Reactant 1 was consumed completely and one new spot formed. The mixture was adjusted to pH 9-10 with aqueous sodium bicarbonate solution, and then diluted extracted with EA (200 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄,

filtered and concentrated under reduced pressure to give tert-butyl 2-formyl-7-azaspiro [3.5]nonane-7-carboxylate (20.4 g, 98% yield) as a yellow oil which was used in the next step without further purification.

[1028] Step 3: Preparation of tert-butyl 2-ethynyl-7-azaspiro [3.5]nonane-7-carboxylate. To a stirred mixture of tert-butyl 2-formyl-7-azaspiro[3.5]nonane-7-carboxylate (10 g, 39.5 mmol, 1 equiv) and K₂CO₃ (10.91 g, 78.9 mmol, 2 equiv) in MeOH (300 ml) was added 1-diazo-1-dimethoxyphosphoryl-propan-2-one (11.37 g, 59.2 mmol, 1.5 equiv) in portions at 0° C. The reaction mixture was stirred for 4 hours at 25° C. TLC (PE:EA=5:1) indicated Reactant 1 was consumed completely and one new spot formed. The reaction mixture was concentrated under reduced pressure remove MeOH, and then diluted extracted with H₂O (80 mL) and EA (80 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0, 20/1) to afford tert-butyl 2-ethynyl-7-azaspiro [3.5]nonane-7-carboxylate (6.5 g, 66% yield) was obtained as a yellow oil.

[1029] Step 4: Preparation of tert-butyl 2-[2-[4-[tert-butyl (dimethyl) silyl]oxyphenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate. To tert-butyl-(4-iodophenoxy)-dimethyl-silane (4.95 g, 14.81 mmol, 1 equiv) in DCM (60 ml) was added TEA (4.50 g, 44.42 mmol, 6.18 ml, 3 equiv) and Pd(PPh₃)₂Cl₂ (1.04 g, 1.48 mmol, 0.1 equiv) and CuI (423.02 mg, 2.22 mmol, 0.15 equiv), then, tert-butyl 2-ethy-

nyl-7-azaspiro [3.5]nonane-7-carboxylate (4.8 g, 19.25 mmol, 1.3 equiv) in DCM (20 ml) was dropwise. The mixture was stirred at 25° C. for 16 h. The reaction mixture was quenched by addition H₂O (30 mL), and then extracted with DCM (50 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 20/1) to afford tert-butyl 2-[2-[4-[tert-butyl (dimethyl) silyl]oxyphenyl]ethynyl]-7-azaspiro [3.5]nonane-7-carboxylate (6.27 g, 92% yield) as a yellow oil.

[1030] Step 5: Preparation of tert-butyl 2-[2-(4-hydroxyphenyl) ethynyl]-7-azaspiro [3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-[2-[4-[tert-butyl (dimethyl) silyl]oxyphenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate (6.27 g, 13.76 mmol, 1 equiv) in THF (60 ml) was added TBAF (1 M, 20.64 ml, 1.5 equiv). The mixture was stirred at 25° C. for 16 h. The reaction mixture was quenched by addition H₂O (20 mL) and extracted with EtOAc (30 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 5/1) to afford tert-butyl 2-[2-(4-hydroxyphenyl) ethynyl]-7-azaspiro [3.5]nonane-7-carboxylate (4.38 g, 93% yield) as a colorless oil.

[1031] Step 6: Preparation of Compound tert-butyl 2-[2-[4-(2-bromo-4-methoxy carbonyl-phenoxy) phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate. To a solution of tert-butyl 2-[2-(4-hydroxyphenyl) ethynyl]-7-azaspiro [3.5]nonane-7-carboxylate (4.38 g, 12.83 mmol, 1 equiv) in DMF (100 ml) was added Cs₂CO₃ (12.54 g, 38.48 mmol, 3 equiv) and methyl 3-bromo-4-fluoro-benzoate (3.89 g, 16.7 mmol, 1.3 equiv). The mixture was stirred at 80° C. for 16 h. The reaction mixture was quenched by addition H₂O (50 mL), and extracted with EA (100 mL×3). The combined organic layers were washed with brine, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 5/1). Compound tert-butyl 2-[2-[4-(2-bromo-4-methoxycarbonyl-phenoxy) phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate (6.52 g, 91% yield) was obtained as a white solid.

[1032] Step 7: Preparation of 2-[4-[4-[2-(7-azaspiro [3.5]nonan-2-yl)ethynyl]phenoxy]-3-bromo-phenyl]propan-2-ol. A mixture of tert-butyl 2-[2-[4-(2-bromo-4-methoxycarbonyl-phenoxy)phenyl]ethynyl]-7-azaspiro[3.5]nonane-7-carboxylate (3 g, 5.41 mmol, 1 equiv) in DCM (100 ml) was degassed and purged with N₂ three times, and then the mixture was added MeMgBr (3 M, 9.02 mL, 5 equiv) at -70° C. dropwise under N₂ atmosphere. The resulting mixture was stirred at 40° C. for 16 hours. The reaction mixture was quenched by adding to NH₄Cl (aq) (200 mL) at 0-5° C. under N₂, and then extracted with DCM/MeOH (10:1, 100 mL×3). The organic layer was dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 2-[4-[4-[2-(7-azaspiro [3.5]nonan-2-yl)ethynyl]phenoxy]-3-bromo-phenyl]propan-2-ol (2.1 g, 85% yield) as a yellow gum.


[1033] Step 8: Preparation of 4-[2-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-pyridinedin-7-one. A mixture of 2-[4-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenoxy]-3-bromo-phenyl]propan-2-ol

(2.1 g, 4.62 mmol, 1 equiv), 6-methyl-1-(p-tolylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[2,3-pyridinedin-7-one (1.98 g, 4.62 mmol, 1 equiv), K₂CO₃ (1.41 g, 10.17 mmol, 2.2 equiv), [2-(2-aminophenyl)phenyl]palladium[bis(1-adamantyl)-butyl-phosphane]methanesulfonate (336.57 mg, 462 μmol, 0.1 equiv) in THE (50 ml) and H₂O (5 ml) was degassed and purged with N₂ three times, and then the mixture was stirred at 70° C. for 16 hr under N₂ atmosphere. The mixture was concentrated to give crude 4-[2-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-pyridinedin-7-one (2.7 g, 4.00 mmol, 86% yield) as a yellow oil and was used without further purification.


[1034] Step 9: Preparation of 4-[2-[4-[2-(7-azaspiro [3.5]nonan-2-yl)ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl) phenyl]-6-methyl-1H-pyrrolo[2,3-pyridinedin-7-one. To a solution of 4-[2-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-pyridinedin-7-one (2.7 g, 4.00 mmol, 1 equiv) in MeOH (120 ml) was added KOH (4.93 g, 87.89 mmol, 22 equiv). The mixture was stirred at 30° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex Luna C18 (250×70 mm×10 μm); mobile phase: [water(FIACN)]; gradient: 20%-50% B over 20 min) to give 4-[2-[4-[2-(7-azaspiro [3.5]nonan-2-yl) ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl) phenyl]-6-methyl-1H-pyrrolo[pyridineyridin-7-one (1 g, 47% yield) as a white solid.

[1035] Step 10: Preparation of 4-(2-(4-((7-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl) phenoxy)ethyl)-7-azaspiro[3.5]nonan-2-yl)ethynyl) phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one. To a solution of [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxy-phenyl)benzothiophen-3-yl]methanone (40 mg, 85.22 μmol, 1 equiv) and 4-[2-[4-[2-(7-azaspiro[3.5]nonan-2-yl)ethynyl]phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (44.46 mg, 85.2 μmol, 1 equiv) in DMF (1 ml) was added DIEA (39.98 mg, 309 μmol, 54 L, 3.63 equiv). The mixture was stirred at 100° C. for 2 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Waters Xbridge 150×25 mm×5 μm; mobile phase: [water (NH₄HCO₃)-ACN]; gradient: 48-78% B over 9 min) to provide 4-(2-(4-((7-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)-7-azaspiro[3.5]nonan-2-yl)ethynyl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (1-31, 19 mg, 24% yield) as a brown gum. ¹H NMR (400 MHz, DMSO-d₆) δ=12.03-11.94 (m, 1H), 9.85-9.62 (m, 2H), 7.67-7.59 (m, 3H), 7.47 (dd, J=2.4, 8.5 Hz, 1H), 7.33 (d, J=2.4 Hz, 1H), 7.27-7.14 (m, 7H), 7.02 (d, J=8.4 Hz, 1H), 6.94-6.88 (m, 2H), 6.86-6.81 (m, 1H), 6.77-6.71 (m, 2H), 6.70-6.64 (m, 2H), 6.22-6.16 (m, 1H), 5.08 (s, 1H), 4.06 (br t, J=6.0 Hz, 2H), 3.51-3.46 (m, 3H), 3.29-3.27 (m, 3H), 3.19-3.13 (m, 1H), 2.60-2.58 (m, 1H), 2.31-2.23 (m, 2H), 2.17-2.08 (m, 2H), 1.83-1.74 (m, 2H), 1.57-1.50 (m, 4H), 1.49-1.44 (m, 6H). LC-MS: MS (ES⁺): RT=2.057 min, m/z=910.6 [M+H⁺], LC-MS Method: 25.

Example 33—Synthesis of 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy) propyl)pyrrolidin-3-yl)oxy) phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylic acid (1-32)



326



I-32

[1036] Step 1: Preparation of 2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethyl 4-methylbenzenesulfonate. To a solution of 2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethan-1-ol (2.70 g, 13.22 mmol, 1.00 equiv) in dichloromethane (30 mL) was added triethylamine (4.01 g, 39.65 mmol, 3.00 equiv). Then was added p-toluenesulfonyl chloride (2.52 g, 13.22 mmol, 1.00 equiv) at 0° C., and the mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to afford a residue. The residue was purified by column chromatography (Silica gel, Petroleum ether/Ethyl acetate=1/0 to 1/1) to afford

2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethyl 4-methylbenzenesulfonate (500 mg, 10% yield) as a yellow oil.

[1037] Step 2: Preparation of (6S)-2,3,6,9-tetramethyl-4-(4-(2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. A mixture of 2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethyl 4-methylbenzenesulfonate (444 mg, 1.24 mmol, 1.50 equiv), (S)-4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenol (280 mg, 0.83 mmol, 1.00 equiv), potassium carbonate (343 mg, 2.48 mmol, 3.00 equiv) in N,N-dimethylformamide (5 mL) was

stirred at 110° C. for 1 h. The reaction mixture was quenched by the addition of water (20 mL), and then extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane/methanol=10:1, R_f =0.45) to afford (6S)-2,3,6,9-tetramethyl-4-(4-(2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (210 mg, 48% yield) as a yellow solid.

[1038] Step 3: Preparation of (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propan-1-ol. To a solution of (6S)-2,3,6,9-tetramethyl-4-(4-(2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (210 mg, 0.40 mmol, 1.00 equiv) in methanol (2 mL) was added tosyl chloride (6 mg, 0.04 mmol, 0.10 equiv) and the mixture was stirred at 25° C. for 0.5 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane/methanol=10:1) to afford (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propan-1-ol (170 mg, 96% yield) as a yellow solid.


[1039] Step 4: Preparation of (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl 4-methylbenzene sulfonate. To a solution of (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propan-1-ol (160 mg, 0.36 mmol, 1.00 equiv), p-toluene-sulfonyl chloride (138 mg, 0.73 mmol, 2.00 equiv) in dichloromethane (2 mL) was added triethylamine (147 mg, 1.45 mmol, 4.00 equiv) and the reaction mixture was stirred at 25° C. for 1 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane/methanol=10:1) to afford (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl 4-methylbenzenesulfonate (145 mg, 67% yield) as a yellow oil.

[1040] Step 5: Preparation of methyl 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylate. A mixture of methyl (S)-8-(2,4-dichlorophenyl)-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylate (51 mg, 0.10 mmol, 1.20 equiv), (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl 4-methylbenzenesulfonate (50 mg, 0.08 mmol, 1.00 equiv),

potassium iodide (13 mg, 0.8 mmol, 1.00 equiv), cesium carbonate (82 mg, 0.25 mmol, 3.00 equiv) in N,N-dimethylformamide (1 mL) was stirred at 100° C. for 1 h. The reaction mixture was quenched by water (10 mL), and then extracted with ethyl acetate (5 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Phenomenex Luna C18 150×25 mm×10 μ m; mobile phase: [water(FA)-ACN]; gradient: 35%-65% B over 10 min] to afford methyl 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylate (50 mg, 63% yield) as a yellow oil.


[1041] Step 6: Preparation of 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylic acid. To a solution of methyl 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylate (50 mg, 0.05 mmol, 1.00 equiv) in tetrahydrofuran (0.2 mL), methanol (0.2 mL) and water (0.2 mL) was added lithium hydroxide (6 mg, 0.16 mmol, 3.00 equiv) and the mixture was stirred at 25° C. for 1 h. The reaction mixture was quenched by aqueous hydrochloric acid (1 M, 5 mL), and then filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Welch Xtimate C18 150×25 mm×5 μ m; mobile phase: [water(FA)-ACN]; gradient: 28%-58% B over 10 min] to afford 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulene-3-carboxylic acid (1-32, 16.82 mg, 32% yield) as a white solid. ^1H NMR (400 MHz, CDCl_3) δ 7.98 (s, 1H), 7.82 (d, J=8.4 Hz, 1H), 7.43 (d, J=8.4 Hz, 2H), 7.39-7.34 (m, 1H), 7.09-6.97 (m, 2H), 6.92 (d, J=8.4 Hz, 1H), 6.88 (d, J=8.8 Hz, 2H), 6.76 (d, J=8.8 Hz, 2H), 6.57-6.51 (m, 2H), 4.93-4.74 (m, 1H), 4.21-4.04 (m, 3H), 3.78 (t, J=4.4 Hz, 2H), 3.62 (t, J=5.6 Hz, 3H), 3.44-3.21 (m, 1H), 3.09-2.85 (m, 6H), 2.68 (s, 3H), 2.41 (s, 3H), 2.32-2.24 (m, 4H), 2.09 (s, 3H), 1.99-1.94 (m, 4H), 1.71 (s, 3H). LCMS: MS (ES $^+$): RT=2.014 min, m/z=916.6 [M+H] $^+$; LC-MS Method: 25.

Example 34—Synthesis of 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)-N-(1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl)azetidin-3-yl)pyridin-3-amine (1-33)



328

-continued



[1042] Step 1: Preparation of 2-[2-(3-tetrahydropyran-2-yloxypropoxy)ethoxy]ethyl 4-methylbenzenesulfonate. To a solution of 2-(2-(3-((tetrahydro-2H-pyran-2-yl)oxy)propoxy)ethoxy)ethan-1-ol (1.00 g, 4.03 mmol, 1.00 equiv) in dichloromethane (10 mL) was added triethylamine (1.22 g, 12.08 mmol, 3.00 equiv) and p-toluenesulfonyl chloride (1.54 g, 8.05 mmol, 2.00 equiv) at 0° C. The mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (petroleum ether/ethyl acetate=10:1 to 1:1) to afford 2-[2-(3-tetrahydropyran-2-yloxypropoxy)ethoxy]ethyl 4-methylbenzenesulfonate (1.50 g, 92% yield) as a yellow oil.

[1043] Step 2: Preparation of (6S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(2-((tetrahydro-2H-pyran-2-yl)oxy)ethoxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of 2-[2-(3-tetrahydropyran-2-yloxypropoxy)ethoxy]ethyl 4-methylbenzenesulfonate (309 mg, 0.76 mmol, 2.00 equiv) and (S)-4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenol (130 mg, 0.38 mmol, 1.00 equiv) in N,N-dimethylformamide (5 mL) was added potassium carbonate (159 mg, 1.15 mmol, 3.00 equiv). The mixture was stirred at 110° C. for 1 h. Ethyl acetate (20 mL) and water (30 mL) were added and the organic layer collected. The aqueous layer was extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (30 mL×3), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to get the residue. The residue was purified by silica gel column chromatography (DCM/methanol=100:1 to 10:1) to afford (6S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(2-((tetrahydro-2H-pyran-2-yl)oxy)ethoxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (216 mg, 98% yield) as a yellow oil.


[1044] Step 3: Preparation of (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propan-1-ol. To a solution of (6S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(2-((tetrahydro-2H-pyran-2-yl)oxy)ethoxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (216 mg, 0.37 mmol, 1.00 equiv) in methanol (2 mL) was added p-toluenesulfonic acid (6 mg, 0.03 mmol, 0.10 equiv) and the mixture was stirred at 25° C. for 0.5 h. The reaction mixture was concentrated under reduced pressure to give a residue. The crude product was purified by prep-TLC (DCM:methanol=10:1) to afford (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propan-1-ol (140 mg, 76% yield) as a yellow oil.

[1045] Step 4: Preparation of (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl 4-methyl benzenesulfonate. To a solution of (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]


diazepin-4-yl)phenoxy)ethoxy)ethoxy)propan-1-ol (140 mg, 0.28 mmol, 1.00 equiv) in dichloromethane (2 mL) was added triethylamine (87 mg, 0.86 mmol, 3.00 equiv). Then p-toluenesulfonyl chloride (82 mg, 0.43 mmol, 1.50 equiv) was added at 0° C. The mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to give a residue. The crude product was purified by prep-TLC (DCM/methanol=10:1) to afford (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl 4-methylbenzenesulfonate (130 mg, 70% yield) as a yellow oil.

[1046] Step 5: Preparation of 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo[4,3-f]isoquinolin-6-yl)-N-(1-(3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl)azetidin-3-yl)pyridin-3-amine. To a solution of (S)-3-(2-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl 4-methyl-benzenesulfonate (118 mg, 0.18 mmol, 1.40 equiv) and N-(azetidin-3-yl)-6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-3,6,8,9-tetrahydropyrazolo[4,3-f]isoquinolin-6-yl)pyridin-3-amine (70 mg, 0.13 mmol, 1.00 equiv, trifluoroacetate) in acetonitrile (2 mL) was added potassium carbonate (54 mg, 0.39 mmol, 3.00 equiv) and potassium iodide (43 mg, 0.26 mmol, 2.00 equiv). The mixture was stirred at 80° C. for 1 h. The mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane:methanol=10:1). The obtained residue was further purified by preparative SFC (Column: Chiralcel OD-3 50×4.6 mm×3 μm; Mobile phase: Phase A for CO₂, and Phase B for IPA/acetonitrile (0.05% DEA); Gradient elution: 40% IPA+acetonitrile (0.05% DEA) in CO₂). The obtained residue was further purified by prep-HPLC (column: Phenomenex Luna C18 150×25 mm×10 μm; mobile phase: [water (formic acid)-acetonitrile]; gradient: 10%-40% B over 10 min) to give 6-((6S,8R)-8-methyl-7-(2,2,2-trifluoroethyl)-6,7,8,9-tetrahydro-2H-pyrazolo [4,3-f]isoquinolin-6-yl)-N-(1-(3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)propyl)azetidin-3-yl)pyridin-3-amine (1-33, 30 mg, 25% yield) as a white solid. ¹H NMR (400 MHz, DMSO-d₆) δ 12.97 (s, 1H), 8.04 (s, 1H), 7.72 (d, J=2.4 Hz, 1H), 7.37 (d, J=8.8 Hz, 2H), 7.21 (d, J=8.8 Hz, 1H), 7.01-6.89 (m, 3H), 6.86-6.74 (m, 2H), 6.21 (d, J=6.8 Hz, 1H), 4.93 (s, 1H), 4.20-4.07 (m, 3H), 3.93-3.87 (m, 1H), 3.76-3.70 (m, 2H), 3.64-3.51 (m, 6H), 3.50-3.43 (m, 6H), 3.13-2.95 (m, 3H), 2.87-2.79 (m, 1H), 2.75-2.70 (m, 1H), 2.58 (s, 3H), 2.38 (s, 3H), 1.85 (d, J=6.8 Hz, 3H), 1.64 (s, 3H), 1.52-1.42 (m, 2H), 1.08 (d, J=6.8 Hz, 3H). LC-MS: MS (ES⁺): RT=1.215 min, m/z=883.7 [M+H⁺], LC-MS Method: 25.

Example 35—Synthesis of (R)-6-(2-(ethyl(4-(2-((R)-3-(2-(2-(4-(S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol and (S)-6-(2-(ethyl(4-(2-((R)-3-(2-(2-(4-(S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (I-34a and I-34b)



-continued



[1047] Step 1: Preparation of (S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(((R)-pyrrolidin-3-yl)oxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of tert-butyl (R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidine-1-carboxylate (110 mg, 0.18 mmol, 1 equiv.) in DCM (3 mL) was added TFA (5.07 g, 44 mmol, 3.30 mL, 240 equiv.). The mixture was stirred at 20° C. for 1 hr. The reaction mixture was concentrated under reduced pressure to remove solvent to afford (S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(((R)-pyrrolidin-3-yl)oxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (92 mg, crude) as a yellow oil. LC-MS: MS (ES⁺): RT=0.384 min, m/z=496.3 [M+H⁺].

[1048] Step 2: Preparation of 2-(6-((tert-butyl)dimethylsilyloxy)-1,2,3,4-tetrahydronaphthalen-2-yl)-N-ethyl-5-methoxy-N-(4-(2-((R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)aniline. To a solution of (S)-2,3,6,9-tetramethyl-4-(4-(2-(2-(((R)-pyrrolidin-3-yl)oxy)ethoxy)ethoxy)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (92 mg, 0.18 mmol, 1 equiv) and 4-((2-(6-((tert-butyl)dimethylsilyloxy)-1,2,3,4-tetrahydronaphthalen-2-yl)-5-methoxyphenyl)(ethyl)amino)methyl)phenethyl 4-methylbenzenesulfonate (129.94 mg, 0.18 mmol, 1 equiv) in ACN (5 mL) was added DIEA (72 mg, 0.5 mmol, 3 equiv). The mixture was stirred at 80° C. for 12 hr. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM/MeOH=10/1) to afford 2-(6-


((tert-butyl)dimethylsilyloxy)-1,2,3,4-tetrahydronaphthalen-2-yl)-N-ethyl-5-methoxy-N-(4-(2-((R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)aniline (103 mg, 49% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 7.44 (d, J=8.8 Hz, 2H), 7.19-7.12 (m, 3H), 7.07 (d, J=8.0 Hz, 2H), 6.92-6.85 (m, 3H), 6.78 (d, J=2.8 Hz, 1H), 6.69 (dd, J=2.8, 8.8 Hz, 1H), 6.64-6.58 (m, 2H), 4.19-4.08 (m, 4H), 3.98 (s, 2H), 3.86 (t, J=4.8 Hz, 2H), 3.79 (s, 3H), 3.74-3.66 (m, 3H), 3.60 (q, J=5.2 Hz, 2H), 2.94-2.87 (m, 3H), 2.86-2.76 (m, 5H), 2.76-2.69 (m, 4H), 2.67 (s, 3H), 2.41 (s, 3H), 2.17-2.06 (m, 4H), 1.96-1.85 (m, 1H), 1.82-1.70 (m, 6H), 1.26 (s, 1H), 1.00 (s, 9H), 0.93 (t, J=7.2 Hz, 3H), 0.24-0.19 (m, 6H). LC-MS: MS (ES⁺): RT=0.493 min, m/z=289.2 [M+H⁺].

[1049] Step 3: Preparation of (R)-6-(2-(ethyl(4-(2-((R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol and (S)-6-(2-(ethyl(4-(2-((R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)amino)-4-methoxyphenyl)-5,6,7,8-tetrahydronaphthalen-2-ol. To a solution of 2-(6-((tert-butyl)dimethylsilyloxy)-1,2,3,4-tetrahydronaphthalen-2-yl)-N-ethyl-5-methoxy-N-(4-(2-((R)-3-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)pyrrolidin-1-yl)ethyl)benzyl)aniline (80 mg, 0.07 mmol, 1 equiv) in DMF (3 mL) was added CsF (59.37 mg, 0.39 mmol, 5 equiv). The mixture was stirred at 80° C. for 0.5 hr. The reaction mixture was diluted with water (20 mL) and extracted with EA (20 mL×3). The combined organic layers


were washed with brine (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient:26%-56% B over min) to afford a mixture of I-34a and I-34b (50 mg, 66% yield) as a white solid. The mixture was separated by SFC (column: Phenomenex-Cellulose-2 (250 mm×30 mm, 0.1 μm); mobile phase: [CO₂-ACN/MeOH(0.1% NH₃H₂O)]; B %:70%, isocratic elution mode) to afford I-34a (16.14 mg, 31% yield) as a white solid and I-34b (19.8 mg, 38% yield) as a white solid. ¹H NMR (I-34a) (400 MHz, DMSO-d₆) δ 7.75-7.64 (m, 1H), 7.37 (d, J=8.4 Hz, 2H), 7.15-7.03 (m, 5H), 6.95 (d, J=8.8 Hz, 2H), 6.84-6.75 (m, 2H), 6.66 (dd, J=2.4, 8.8 Hz, 1H), 6.52-6.46 (m, 2H), 4.18-4.08 (m, 5H), 3.96 (d, J=2.4 Hz, 3H), 3.75-3.71 (m, 4H), 3.70 (s, 3H), 3.55 (s, 3H), 2.89-2.83 (m, 4H), 2.72 (d, J=3.6 Hz, 4H), 2.69-2.65 (m, 4H), 2.64 (s, 2H), 2.33 (dd, J=1.6, 3.2 Hz, 2H), 1.94 (dd, J=6.0, 13.2 Hz, 2H), 1.85 (d, J=6.8 Hz, 3H), 1.64 (s, 3H),

1.29-1.20 (m, 2H), 0.87 (t, J=7.2 Hz, 5H). LC-MS: MS (ES⁺): RT=1.395 min, m/z=909.7 [M+H⁺], 25. ¹H NMR (I-34b) (400 MHz, DMSO-d₆) δ 7.75-7.62 (m, 1H), 7.37 (d, J=8.8 Hz, 2H), 7.15-7.03 (m, 5H), 6.95 (d, J=8.8 Hz, 2H), 6.82-6.76 (m, 2H), 6.66 (dd, J=2.4, 8.4 Hz, 1H), 6.52-6.47 (m, 2H), 4.17-4.08 (m, 5H), 3.96 (d, J=2.4 Hz, 3H), 3.75-3.71 (m, 4H), 3.70 (s, 3H), 3.55 (s, 3H), 2.88-2.83 (m, 4H), 2.73 (d, J=5.2 Hz, 4H), 2.66 (d, J=3.6 Hz, 4H), 2.63 (s, 2H), 2.33 (d, J=1.6 Hz, 2H), 1.97-1.90 (m, 2H), 1.85 (d, J=6.4 Hz, 3H), 1.64 (s, 3H), 1.27-1.21 (m, 2H), 0.89-0.82 (m, 5H). LC-MS: MS (ES⁺): RT=1.395 min, m/z=909.7 [M+H⁺], Method 25.

Example 36—Synthesis of 4-(2-(4-(8-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)-2,8-diazaspiro[4.5]decan-2-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (I-35)



-continued



I-35

[1050] Step 1: Preparation of tert-butyl 2-[4-[tert-butyl (dimethyl)silyl]oxyphenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate. A mixture of tert-butyl 2,8-diazaspiro[4.5]decane-8-carboxylate (1 g, 4.16 mmol, 1 equiv), tert-butyl-(4-iodophenoxy)-dimethyl-silane (1.39 g, 4.16 mmol, 1 equiv), SPhos Pd G3 (227.25 mg, 291.25 μ mol, 0.07 equiv), Cs₂CO₃ (3.39 g, 10.40 mmol, 2.5 equiv) in dioxane (20 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 80° C. for 12 h under N₂ atmosphere. The mixture was filtered off, filter-cake was washed by EtOAc (20 mL *3), and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel chromatography (20 g SepaFlash® Silica Flash Column,

Eluent of 3-10% Ethyl acetate/Petroleum ether gradient 20 mL/min) to afford tert-butyl 2-[4-[tert-butyl(dimethyl)silyl]oxyphenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate (1.2 g, 64% yield) as a colorless oil. ^1H NMR (400 MHz, CDCl_3) δ 6.79-6.71 (m, 2H), 6.47-6.39 (m, 2H), 3.55-3.45 (m, 2H), 3.42-3.35 (m, 2H), 3.32 (t, 2H, $J=6.8$ Hz), 3.11 (s, 2H), 1.87 (t, 2H, $J=7.2$ Hz), 1.63 (s, 1H), 1.58 (s, 3H), 1.47 (s, 9H), 0.98 (s, 9H), 0.16 (s, 6H).

[1051] Step 2: Preparation of tert-butyl 2-(4-hydroxyphenyl)-2,8-diazaspiro [4.5]decane-8-carboxylate. To a solution of tert-butyl 2-[4-[tert-butyl(dimethyl)silyl]oxyphenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate (1.2 g, 2.69 mmol, 1

equiv) in THE (20 mL) was added TBAF (1 M, 4.03 mL, 1.5 equiv) at 0° C. The mixture was stirred at 25° C. for 1 h. The mixture was concentrated under reduced pressure to afford tert-butyl 2-(4-hydroxy phenyl)-2,8-diazaspiro[4.5]decane-8-carboxylate (820 mg, crude) as brown solid.

[1052] Step 3: Preparation of tert-butyl 2-[4-(2-bromo-4-methoxycarbonyl-phenoxy)phenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate. To a solution of tert-butyl 2-(4-hydroxyphenyl)-2,8-diazaspiro[4.5]decane-8-carboxylate (800 mg, 2.41 mmol, 1 equiv) in MeCN (3 mL) was added Cs₂CO₃ (1.96 g, 6.02 mmol, 2.5 equiv) and methyl 3-bromo-4-fluoro-benzoate (392.55 mg, 1.68 mmol, 0.7 equiv). The mixture was stirred at 50° C. for 1 h. The mixture was filtered, the filter-cake was washed by EtOAc (10 mL×3), and the filtrate was concentrated under reduced pressure. The residue was purified by flash silica gel chromatography (20 g SepaFlash Silica Flash Column, eluent: 15~30% ethyl acetate/petroleum ether gradient 30 mL/min) to afford tert-butyl 2-[4-(2-bromo-4-methoxycarbonyl-phenoxy) phenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate (410 mg, 29% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 8.29 (d, 1H, J=2.4Hz), 7.82 (dd, 1H, J=2.0, 8.4 Hz), 7.02-6.94 (m, 2H), 6.71 (d, 1H, J=8.8 Hz), 6.60-6.50 (m, 2H), 3.90 (s, 3H), 3.57-3.47 (m, 2H), 3.45-3.36 (m, 4H), 3.18 (s, 2H), 1.92 (t, 2H, J=6.8 Hz), 1.65-1.59 (m, 4H), 1.48 (s, 9H). LC-MS: MS (ES⁺): RT=0.730 min, m/z=545.1; 547.1 [M+H⁺].

[1053] Step 4: Preparation of 2-[3-bromo-4-[4-(2,8-diazaspiro[4.5]decan-2-yl) phenoxy]phenyl]propan-2-ol. To a solution of tert-butyl 2-[4-(2-bromo-4-methoxycarbonyl-phenoxy) phenyl]-2,8-diazaspiro[4.5]decane-8-carboxylate (460 mg, 843.32 mol, 1 equiv) in THF (3 mL) was added MeMgBr (3 M, 5.62 mL, 20 equiv) at 0° C. under an N₂ atmosphere. The mixture was stirred at 40° C. for 1 h. The reaction mixture was quenched by adding saturated NH₄Cl (10 mL), and extracted with DCM:MeOH (10:1, 20 mL ×3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford 2-[3-bromo-4-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]phenyl]propan-2-ol (220 mg, 54% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.730 min, m/z=445.1; 447.1 [M+H⁺].


[1054] Step 5: Preparation of 4-(2-(4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1-tosyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one. A mixture of 2-[3-bromo-4-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]phenyl]propan-2-ol (100 mg, 224.52 mol, 1 equiv), 6-methyl-1-(p-tolylsulfonyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)pyrrolo[2,3-c]pyridin-7-one (96.16 mg, 224.52 mol, 1 equiv), [2-(2-aminophenyl)phenyl]palladium(1+);bis(1-adamantyl)-butyl-phosphane; methanesulfonate (16.35 mg, 22.45 mol, 0.1 equiv) and K₂CO₃ (77.58 mg, 561.30 mol, 2.5 equiv) in dioxane (3 mL) and H₂O (0.3 mL) was degassed and purged with N₂ 3 times, and then the mixture was stirred at 90° C. for 12 h under an


N₂ atmosphere. To the reaction mixture was added H₂O (10 mL) and the reaction mixture was extracted with DCM: MeOH=10:1 (10 mL ×3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to afford the crude desired product (105 mg) as a yellow solid.

[1055] Step 6: Preparation of 4-[2-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phe-nyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one. To a solution of 4-[2-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1-(p-tolylsulfonyl)pyrrolo[2,3-c]pyridin-7-one (100 mg, 149.96 mol, 1 equiv) in MeOH (3 mL) was added KOH (168.28 mg, 3.00 mmol, 20 equiv). The mixture was stirred at 30° C. for 1 h. The mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (column: Welch Xtimate C18 150×25 mm×5 um; mobile phase: [water(FA)-ACN]; gradient: 10%-40% B over 10 min) to afford 4-[2-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (45 mg, 55% yield) as an off-white solid. LC-MS: MS (ES⁺): RT=0.890 min, m/z=513.3 [M+H⁺].

[1056] Step 7: Preparation of 4-(2-(4-(8-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl) benzo[b]thiophene-3-carbonyl)phe-noxy)ethyl)-2,8-diazaspiro[4.5]decan-2-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one. To a solution of 4-[2-[4-(2,8-diazaspiro[4.5]decan-2-yl)phenoxy]-5-(1-hydroxy-1-methyl-ethyl)phenyl]-6-methyl-1H-pyrrolo[2,3-c]pyridin-7-one (40 mg, 78.03 mol, 1 equiv) in DMSO (0.5 mL) was added DIEA (30.25 mg, 234.08 mol, 40.77 μL, 3 equiv) and [4-(2-bromoethoxy)phenyl]-[6-hydroxy-2-(4-hydroxyphenyl)benzothiophen-3-yl]methanone (36.62 mg, 78.03 mol, 1 equiv). The mixture was stirred at 100° C. for 1 h. To the reaction mixture was added H₂O (20 mL), and it was extracted with EtOAc (20 mL ×3). The combined organic layers were washed with brine (10 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1), and then purified by prep-HPLC (column: Welch Xtimate C18 150*25 mm*5 um; mobile phase: [water(FA)-ACN]; gradient:20%-500% B over 10 min) to afford 4-(2-(4-(8-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophene-3-carbonyl)phenoxy)ethyl)-2,8-diazaspiro[4.5]decan-2-yl)phenoxy)-5-(2-hydroxypropan-2-yl)phenyl)-6-methyl-1,6-dihydro-7H-pyrrolo[2,3-c]pyridin-7-one (1-35, 17 mg, 23% yield) as a yellow solid. ¹H NMR (400 MHz, DMSO-d₆) δ 11.98 (s, 1H), 10.03-9.51 (m, 2H), 8.20 (s, 1H), 7.65 (d, 2H, J=8.8 Hz), 7.51 (d, 1H, J=2.4 Hz), 7.35-7.30 (m, 2H), 7.28-7.22 (m, 3H), 7.19-7.14 (m, 2H), 6.92 (d, 2H, J=7.2 Hz), 6.88-6.80 (m, 3H), 6.71-6.64 (m, 3H), 6.47 (d, 2H, J=9.2 Hz), 6.24 (t, 1H, J=2.4 Hz), 4.98 (s, 1H), 4.10 (t, 2H, J=5.6 Hz), 3.55 (s, 3H), 3.23 (t, 4H, J=6.8 Hz), 3.03 (s, 2H), 2.67 (t, 2H, J=5.6 Hz), 2.44-2.38 (m, 2H), 1.79 (t, 2H, J=6.4 Hz), 1.58-1.48 (m, 4H), 1.44 (s, 6H). LC-MS: MS (ES⁺): RT=1.917 min, m/z=901.7 [M+H⁺], LCMS Method: 25.

Example 37—Synthesis of 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy) propyl) pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-ol (I-36)





I-36

[1057] Step 1: Preparation of tert-butyl (S)-3-(4-(3-acetoxy-6,7-dihydro-5H-benzo[7]annulen-9-yl)phenoxy)pyrrolidine-1-carboxylate. To a solution of 9-(((trifluoromethyl)sulfonyl)oxy)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate (3.00 g, 8.56 mmol, 1.00 equiv), tert-butyl (S)-3-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenoxy)pyrrolidine-1-carboxylate (3.33 g, 8.56 mmol, 1.00 equiv), and cesium carbonate (5.58 g, 17.13 mmol, 2.00 equiv) in dioxane (24 mL) and water (6 mL) was added 1,1'-bis(diphenyl phosphino) ferrocene-palladium(II)dichloride (626 mg, 0.86 mmol, 0.10 equiv) and the mixture was stirred at 25° C. for 8 h under a nitrogen atmosphere. The reaction mixture was quenched with water (50 mL), and then extracted with ethyl acetate (30 mL×3). The combined organic layers were washed with brine (50 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica gel, petroleum ether/ethyl acetate=1/0 to 3/1) to afford tert-butyl (S)-3-(4-(3-acetoxy-6,7-dihydro-5H-benzo[7]annulen-9-yl)phenoxy)pyrrolidine-1-carboxylate (3.50 g, 88% yield) as a white solid. LCMS: MS (ES⁺): RT=0.758 min, m/z=486.4 [M+Na]⁺.

[1058] Step 2: Preparation of (S)-8-bromo-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate. A mixture of monopyridinium tribromide (379 mg, 1.19 mmol, 1.10 equiv) and tert-butyl (S)-3-(4-(3-acetoxy-6,7-dihydro-5H-benzo[7]annulen-9-yl)phenoxy)pyrrolidine-1-carboxylate (500 mg, 1.08 mmol, 1.00 equiv) in dichloromethane (15 mL) was stirred at 25° C. for 8 h. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (silica gel, petroleum ether/ethyl acetate=1/0 to 10/1) to afford (S)-8-bromo-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate (330 mg, 69% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.21 (d, J=8.4 Hz, 2H), 6.99 (d, J=2.0 Hz, 1H), 6.94-6.78 (m,

4H), 5.09 (s, 1H), 3.70-3.53 (m, 4H), 2.77 (t, J=6.8 Hz, 2H), 2.64-2.57 (m, 2H), 2.44 (dd, J=6.0, 14.4 Hz, 1H), 2.38-2.22 (m, 7H).

[1059] Step 3: Preparation of (S)-8-(2,4-dichlorophenyl)-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate. To a solution of (S)-8-bromo-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate (330 mg, 0.74 mmol, 1.00 equiv), (2,4-dichlorophenyl)boronic acid (156 mg, 0.82 mmol, 1.10 equiv), and cesium carbonate (486 mg, 1.49 mmol, 2.00 equiv) in dioxane (4 mL) and water (1 mL) was added 1,1'-bis(diphenylphosphino)ferrocene-palladium(II)dichloride (54 mg, 0.07 mmol, 0.10 equiv) and the mixture was stirred at 90° C. for 3 h under a nitrogen atmosphere. The reaction mixture was quenched with water (20 mL), and then extracted with ethyl acetate (10 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (dichloromethane:methanol=10:1) to afford (S)-8-(2,4-dichlorophenyl)-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate (320 mg, 84% yield) as a yellow solid. LCMS: MS (ES⁺): RT=0.625 min, m/z=510.1 [M+H]⁺.


[1060] Step 4: Preparation of 8-(2,4-dichlorophenyl)-9-(((S)-1-(3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate. A mixture of (S)-8-(2,4-dichlorophenyl)-9-(4-(pyrrolidin-3-yloxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl acetate (51 mg, 0.10 mmol, 1.20 equiv), (S)-3-(2-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl 4-methyl benzenesulfonate (50 mg, 0.08 mmol, 1.00 equiv), potassium iodide (13 mg, 0.08 mmol, 1.00 equiv), and cesium carbonate (82 mg, 0.25 mmol, 3.00 equiv) in N,N-dimethylformamide (1 mL) was stirred at 100° C. for 1


h. The reaction mixture was quenched with water (10 mL), and then extracted with ethyl acetate (5 mL×3). The combined organic layers were washed with brine (30 mL), dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Phenomenex luna C18 150×25 mm×10 μm; mobile phase: [water(FA)-ACN]; gradient:80%-100% B over 10 min] to afford 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl) acetate (50 mg, 63% yield) as a white solid. LCMS: MS (ES⁺): RT=0.656 min, m/z=930.4 [M+H]⁺.

[1061] Step 5: Preparation of 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl) acetate (50 mg, 0.05 mmol, 1.00 equiv) in water (0.2 mL), tetrahydrofuran (0.2 mL) and methanol (0.2 mL) was added lithium hydroxide (6 mg, 0.16 mmol, 3.00 equiv) and the mixture was stirred at 25° C. for 108 h. The reaction mixture was quenched with an aqueous hydrochloric acid solution (1M, 5 mL), and then filtered and concen-

trated under reduced pressure to give a residue. The residue was purified by prep-HPLC [column: Welch Xtimate C18 150×25 mm×5 μm; mobile phase: [water(FA)-ACN]; gradient:30%-60% B over 10 min] to afford 8-(2,4-dichlorophenyl)-9-(4-(((S)-1-(3-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)propyl)pyrrolidin-3-yl)oxy)phenyl)-6,7-dihydro-5H-benzo[7]annulen-3-yl) (I-36, 22.78 mg, 42.09% yield) as a white solid. LCMS: MS (ES⁺): RT=2.604 min, m/z=888.4 [M+H]⁺; 5-95. ¹H NMR (400 MHz, DMSO-d₆) δ 8.38 (s, 2H), 7.57 (d, J=2.0 Hz, 1H), 7.36 (d, J=8.4 Hz, 2H), 7.27-7.22 (m, 1H), 7.15 (d, J=8.4 Hz, 1H), 6.94 (d, J=8.4 Hz, 2H), 6.89 (d, J=2.4 Hz, 1H), 6.75-6.69 (m, 3H), 6.67-6.62 (m, 3H), 4.75 (t, J=7.6 Hz, 1H), 4.17-4.10 (m, 3H), 4.04 (t, J=6.0 Hz, 2H), 3.75-3.72 (m, 2H), 3.62 (s, 2H), 3.43 (s, 1H), 3.07-3.03 (m, 1H), 2.85-2.76 (m, 4H), 2.58 (s, 3H), 2.38 (s, 3H), 2.19-2.12 (m, 2H), 2.10-2.03 (m, 2H), 2.00-1.93 (m, 3H), 1.87-1.82 (m, 3H), 1.76-1.69 (m, 1H), 1.63 (s, 3H).

Example 38—Synthesis of (5S,6R)-6-phenyl-5-(4-(4-((4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol and (5R,6S)-6-phenyl-5-(4-(4-((4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (I-37a and I-37b)





[1062] Step 1: Preparation of (+)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine. A mixture of (i)-cis-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]trifluoromethanesulfonate (200 mg, 0.43 mmol, 1 equiv), 4-(dimethoxymethyl)piperidine (103 mg, 0.64 mmol, 1.5 equiv), CPHOS PD G₃ (69 mg, 0.08 mmol, 0.2 equiv), Xantphos (75 mg, 0.12 mmol, 0.3 equiv) and Cs₂CO₃ (281 mg, 0.86 mol, 2 equiv) in dioxane (5 mL) was degassed and purged with N₂ 3 times, and then the mixture was stirred at 95° C. for 12 h under an N₂ atmosphere. The

reaction mixture was quenched by addition H₂O (50 mL), and then extracted with ethyl acetate (200 mL). The combined organic layers were dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether/ethyl acetate=5:1). (+)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine (150 mg, 0.31 mmol, 73% yield) was obtained as a yellow oil. LC-MS: MS (ES⁺): RT=0.537 min, m/z=472.3 [M+H⁺].

[1063] Step 2: Preparation of ($\pm$)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde. To a solution of (i)-cis-4-(dimethoxymethyl)-1-[4-(6-methoxy-2-phenyl-tetralin-1-yl)phenyl]piperidine (150.00 mg, 0.31 mmol, 1 equiv) in DCM (2 mL) was added BBr_3 (1.30 g, 5.19 mmol, 0.5 mL, 16.32 equiv) at 0° C. The mixture was stirred at 20° C. for 3 h. And then H_2O (2 mL) was added dropwise at 0° C. The resulting mixture was stirred at 20° C. for 2 hours. The reaction mixture was adjusted to pH=7-8 with aqueous NaHCO_3 and extracted with EA 15 mL (5 mL*3). The combined organic layers were washed with brine (10 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to afford ($\pm$)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (120 mg, 0.29 mmol, 91% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.471 min, m/z=412.2 [M+H⁺].


[1064] Step 3: Preparation of tert-butyl (S)-4-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazine-1-carboxylate. A mixture of (S)-4-(4-chlorophenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (100 mg, 0.28 mmol, 1 equiv), tert-butyl piperazine-1-carboxylate (78 mg, 0.42 mmol, 1.5 equiv), SPhos Pd G₃ (21 mg, 0.02 mmol, 0.1 equiv), and Cs_2CO_3 (182 mg, 0.56 mmol, 2 equiv) in dioxane (10 mL) was degassed and purged with N_2 3 times, and then the mixture was stirred at 100° C. for 12 h under an N_2 atmosphere. The reaction mixture was partitioned between EtOAc (20 mL) and H_2O (20 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , DCM:MeOH=10:1) to afford tert-butyl (S)-4-(4-(2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazine-1-carboxylate (140 mg, 98% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.478 min, m/z=507.3 [M+H⁺].


[1065] Step 4: Preparation of (S)-2,3,6,9-tetramethyl-4-(4-(piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of tert-butyl 4-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2⁶]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazine-1-carboxylate (140 mg, 0.27 mmol, 1 equiv) in DCM (2 mL) was added HCl/dioxane (2 M, 3.50 mL, 25.33 equiv). The mixture was stirred at 25° C. for 12 h. The reaction mixture was concentrated under reduced pressure to remove solvent to afford (S)-2,3,6,9-tetramethyl-4-(4-(piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (120 mg, 98% yield, HCl) as a yellow solid. LC-MS: MS (ES⁺): RT=0.352 min, m/z=407.2 [M+H⁺].


[1066] Step 5: Preparation of (5S,6R)-6-phenyl-5-(4-(4-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol and (5R,6S)-6-phenyl-5-(4-(4-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperazin-1-yl)methyl)piperidin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (I-37a and I-37b). To a solution of (S)-2,3,6,9-tetramethyl-4-(4-(piperazin-1-yl)phenyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine (80 mg, 0.18 mmol, 1 equiv, HCl) in DCM (3 mL) was added NaOAc (148 mg, 1.81 mmol, 10 equiv) at 25° C. The mixture was stirred at 25° C. for 0.5 h. Then (+)-cis-1-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)phenyl]piperidine-4-carbaldehyde (81 mg, 0.19 mmol, 1.1 equiv) and $\text{NaBH}(\text{OAc})_3$

(57 mg, 0.27 mmol, 1.5 equiv) were added. The mixture was stirred at 25° C. for 12 hr. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was purified by prep-TLC (SiO_2 , DCM:MeOH=10:1) to afford 2-phenyl-1-[4-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2⁶]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperazin-1-yl]methyl]-1-piperidyl]phenyl]tetralin-6-ol (40 mg, 0.04 mmol, 27% yield) as a white solid. The stereoisomeric mixture was purified by prep-HPLC (column: DAICEL CHIRALPAK AD (250 mm×30 mm, 10 μm); mobile phase: [CO_2 -ACN/i-PrOH (0.1% $\text{NH}_3\text{H}_2\text{O}$)]); B %: 70%, isocratic elution mode) to give I-37a (about 22% yield) as a white solid and I-37b (about 22% yield) as a white solid. ¹H NMR (peak 1): (400 MHz, CHLOROFORM-d) δ ppm 1.29-1.38 (m, 3H) 1.56-1.68 (m, 2H) 1.76 (s, 3H) 1.83 (br d, J=13.6 Hz, 3H) 2.05-2.09 (m, 3H) 2.22-2.28 (m, 2H) 2.41 (s, 3H) 2.52-2.61 (m, 6H) 2.66-2.68 (m, 3H) 2.95-3.07 (m, 2H) 3.25 (br d, J=2.8 Hz, 4H) 3.50-3.59 (m, 2H) 4.11-4.21 (m, 2H) 6.29 (d, J=8.8 Hz, 2H) 6.55-6.62 (m, 3H) 6.72 (d, J=2.4 Hz, 1H) 6.79-6.86 (m, 5H) 7.11-7.18 (m, 3H) 7.38-7.44 (m, 2H) LC-MS (peak 1): MS (ES⁺): RT=1.035 min, m/z=802.5 [M+H⁺], LCMS Method 10. ¹H NMR (peak 2): (400 MHz, CHLOROFORM-d) δ ppm 1.28-1.38 (m, 3H) 1.56-1.69 (m, 2H) 1.76 (s, 3H) 1.83 (br d, J=14.4 Hz, 3H) 2.08 (d, J=6.8 Hz, 3H) 2.22-2.29 (m, 2H) 2.41 (s, 3H) 2.52-2.62 (m, 6H) 2.67 (s, 3H) 2.93-3.06 (m, 2H) 3.25 (br d, J=2.0 Hz, 4H) 3.55 (br dd, J=8.8, 3.2 Hz, 2H) 4.11-4.21 (m, 2H) 6.29 (d, J=8.8 Hz, 2H) 6.54-6.60 (m, 3H) 6.72 (d, J=2.4 Hz, 1H) 6.78-6.87 (m, 5H) 7.12-7.18 (m, 3H) 7.39 (d, J=8.8 Hz, 2H). LC-MS (peak 2): MS (ES⁺): RT=1.035 min, m/z=802.5 [M+H⁺], LCMS Method 10.

Example 39—Synthesis of (5R,6S)-6-phenyl-5-(4-(2-(4-((S)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (I-62) and (5R,6S)-6-phenyl-5-(4-(2-(4-((R)-2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (III-4)







[1067] Step 1: Preparation of 6-(2-trimethylsilylethoxy) tetralin-1-one. To a solution of 6-hydroxytetralin-1-one (21 g, 129.48 mmol, 1 equiv), 2-trimethylsilylethanol (19.14 g, 161.85 mmol, 23.20 mL, 1.25 equiv) and PPh_3 (42.45 g, 161.85 mmol, 1.25 equiv) in DCM (210 mL) was added DIAD (32.73 g, 161.85 mmol, 31.38 mL, 1.25 equiv) at -78°C . The mixture was stirred at 40°C for 12 h. The reaction mixture was partitioned between DCM (100 mL) and H_2O (100 mL). The organic phase was separated, dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether/ethyl acetate=10/1 to 3/1). 6-(2-trimethylsilylethoxy) tetralin-1-one (8.4 g, 32.01 mmol, 24% yield) was obtained as a yellow oil. LC-MS: MS (ES^+): RT=0.460 min, m/z =265.2 [$\text{M}+\text{H}^+$].

[1068] Step 2: Preparation of [6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate. To a mixture of 6-(2-trimethylsilylethoxy)tetralin-1-one (10.6 g, 40.39 mmol, 1 equiv) in THF (100 mL) was added LiHMDS (1 M, 60.59 mL, 1.5 equiv) dropwise at -78°C and the mixture was stirred at -78°C for 1 h under an N_2 atmosphere. Then 1,1,1-trifluoro-N-phenyl-N-(trifluoromethylsulfonyl)methanesulfonamide (15.87 g, 44.43 mmol, 1.1 equiv) in THF (30 mL) was added dropwise to the mixture. The mixture was stirred at 25°C for 12 h under an N_2 atmosphere. The reaction mixture was quenched by the addition of H_2O (100 mL) at 0°C under an N_2 atmosphere, stirred at 0°C for 20 min and then stirred at 25°C for 20 min. The reaction mixture was then diluted with H_2O (500 mL) and extracted with EtOAc (500 mL $\times$ 2). The combined organic layers were washed with brine (300 mL $\times$ 2), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether/ethyl acetate=20/1 to 3/1) to afford [6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (10.5 g, 26.62 mmol, 65% yield) as a white solid. LC-MS: MS (ES^+): RT=0.718 min.

[1069] Step 3: Preparation of 4-[6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol. A mixture of [6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (6.5 g, 16.48 mmol, 1 equiv), (4-hydroxyphenyl)boronic acid (2.73 g, 19.77 mmol, 1.2 equiv), K_2CO_3 (4.55 g, 32.95 mmol, 2 equiv), and $\text{Pd}(\text{dppf})\text{Cl}_2$ (1.21 g, 1.65 mmol, 0.1 equiv) in dioxane (65 mL) and

H_2O (13 mL) was degassed and purged with N_2 3 times, and then the mixture was stirred at 100°C for 2 hr under an N_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 (250 $\times$ 70 mm, 10 μm); mobile phase: [water (FA)-ACN]; gradient: 66%-96% B over 21 min) to afford 4-[6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.1 g, 73% yield) as a white solid. LC-MS: MS (ES^+): RT=0.686 min, m/z =339.2 [$\text{M}+\text{H}^+$].

[1070] Step 4: Preparation of 4-[2-bromo-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol. To a solution of 4-[6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.1 g, 12.11 mmol, 1 equiv) in MeCN (40 mL) was added NBS (2.05 g, 11.51 mmol, 0.95 equiv). The mixture was stirred at 20°C for 2 h. The reaction mixture was partitioned between EA (50 mL) and H_2O (50 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether/ethyl acetate=10/1 to 3/1) to afford 4-[2-bromo-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.3 g, 85% yield) as a white solid. LC-MS: MS (ES^+): RT=0.720 min.

[1071] Step 5: Preparation of 4-[2-phenyl-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol. A mixture of 4-[2-bromo-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.3 g, 10.30 mmol, 1 equiv), phenylboronic acid (1.51 g, 12.36 mmol, 1.2 equiv), K_2CO_3 (2.85 g, 20.60 mmol, 2 equiv), and $\text{Pd}(\text{dppf})\text{Cl}_2$ (753.78 mg, 1.03 mmol, 0.1 equiv) in dioxane (50 mL) and H_2O (10 mL) was degassed and purged with N_2 3 times, and then the mixture was stirred at 90°C for 3 hr under an N_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether/ethyl acetate=30/1 to 3/1) to afford 4-[2-phenyl-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.2 g, 98% yield) as a white solid. LC-MS: MS (ES^+): RT=0.793 min, m/z =415.2 [$\text{M}+\text{H}^+$].

[1072] Step 6: Preparation of 4-[(1R,2S)-2-phenyl-6-(2-trimethylsilylethoxy)tetralin-1-yl]phenol. To a hydrogenation vessel filled with N_2 was added Pd/C (500 mg, 469.84 mol, 10% purity, 4.64e-2 equiv), followed by 4-[2-phenyl-6-(2-trimethylsilylethoxy)-3,4-dihydronaphthalen-1-yl]phenol (4.2 g, 10.13 mmol, 1 equiv) in MeOH (40 mL). The suspension was degassed and purged with H_2 3 times. The mixture was stirred under H_2 (45 psi) at 25°C for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150 $\times$ 40 mm $\times$ 15 μm ; mobile phase: [water (FA)-ACN]; gradient: 70%-100% B over 15 min) and SFC (column: DAICEL CHIRALPAKAD (250 mm $\times$ 30 mm $\times$ 10 μm); mobile phase: [CO_2 -i-PrOH (0.1% $\text{NH}_3\text{H}_2\text{O}$)]); B %: 40%, isocratic elution mode) to afford 4-[(1R,2S)-2-phenyl-6-(2-trimethylsilylethoxy)tetralin-1-yl]phenol (280 mg, 672.07 μmol , 6% yield, $[\alpha]_D^{25}$: -287.21) as a white solid and 4-[(1S,2R)-2-phenyl-6-(2-trimethylsilylethoxy)tetralin-1-yl]phenol (280 mg, 672.07 μmol , 6% yield, $[\alpha]_D^{25}$: +356.42) as a white solid. LC-MS: MS (ES^+): RT=0.729 min.

[1073] Step 7: Preparation of [4-[(1R,2S)-2-phenyl-6-(2-trimethylsilylethoxy)tetralin-1-yl]phenyl]trifluoromethanesulfonate. To a solution of 4-[(1R,2S)-2-phenyl-6-(2-trimethylsilylethoxy)tetralin-1-yl]phenol (250 mg, 600.06 mol,

1 equiv) in DCM (5 mL) was added Ti_2O (338.60 mg, 1.20 mmol, 198.01 L, 2 equiv) and TEA (182.16 mg, 1.80 mmol, 250.56 μL , 3 equiv). The mixture was stirred at 25° C. for 1h. The reaction mixture was partitioned between EtOAc (30 mL) and H_2O (30 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , PE:EA=5:1) to afford 4-[(1R,2S)-2-phenyl-6-(2-trimethylsilylethoxy) tetralin-1-yl]phenyl]trifluoromethanesulfonate (280 mg, 510.31 mol, 85% yield) as a white solid.


[1074] Step 8: Preparation of trimethyl-[2-[(1R,2S)-2-phenyl-1-[4-[2-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-azaspiro[3.5]nonan-7-yl]phenyl]tetralin-6-yl]oxyethyl]silane. To a mixture of 2-[[[(9S)-7-[4-(7-azaspiro[3.5]nonan-2-ylmethyl)phenyl]-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-9-yl]methyl]oxazole (100 mg, 189.86 mol, 1 equiv), 4-[(1R,2S)-2-phenyl-6-(2-trimethylsilyl ethoxy)tetralin-1-yl]phenyl]trifluoromethanesulfonate (125.01 mg, 227.84 mol, 1.2 equiv), Cs_2CO_3 (185.58 mg, 569.59 mol, 3 equiv), and CPHOS PD G₃ (15.31 mg, 18.99 mol, 0.1 equiv) in dioxane (5 mL) was degassed and purged with N_2 3 times, and then the mixture was stirred at 90° C. for 12 h under an N_2 atmosphere. The reaction mixture was partitioned between EA (20 mL) and H_2O (20 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , DCM:MeOH=10:1) to afford trimethyl-[2-[(1R,2S)-2-phenyl-1-[4-[2-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-azaspiro[3.5]nonan-7-yl]phenyl]tetralin-6-yl]oxyethyl]silane (80 mg, 45% yield) as a white solid. LC-MS: MS (ES^+): RT=0.702 min, m/z =925.6 [$\text{M}+\text{H}^+$].

[1075] Step 9: Preparation of (1R,2S)-2-phenyl-1-[4-[2-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-azaspiro[3.5]nonan-7-yl]phenyl]tetralin-6-ol. To a solution of trimethyl-[2-[(1R,2S)-2-phenyl-1-[4-[2-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-


azaspiro[3.5]nonan-7-yl]phenyl]tetralin-6-yl]oxyethyl]silane (30 mg, 32.42 mol, 1 equiv) in DCM (3 mL) was added HCl/dioxane (4 M, 3.60 mL, 444.15 equiv). The mixture was stirred at 25° C. for 3 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150×25 mm×10 μm ; mobile phase: [water (FA)-ACN]; gradient: 48%-78% B over 9 min) to afford (5R,6S)-6-phenyl-5-(4-(2-(4-(2,3,9-trimethyl-6-(oxazol-2-ylmethyl)-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)benzyl)-7-azaspiro[3.5]nonan-7-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (6.76 mg, 25% yield) as a white solid. LC-MS: MS (ES^+): RT=0.785 min, m/z =825.5 [$\text{M}+\text{H}^+$].

[1076] Step 10: Preparation of I-62 and III-4. The residue from the prior step was purified by SFC (column: DAICEL CHIRALPAK IC (250 mm×30 mm×10 μm); mobile phase: [CO_2 -ACN/EtOH (0.1% $\text{NH}_3\text{H}_2\text{O}$)]); B %: 75%, isocratic elution mode) to afford compounds I-62 and III-4, each of the two diastereomers of (1R,2S)-2-phenyl-1-[4-[2-[[4-[4,5,13-trimethyl-9-(oxazol-2-ylmethyl)-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]methyl]-7-azaspiro[3.5]nonan-7-yl]phenyl]tetralin-6-ol. The diastereomer that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below), was obtained as a white solid (6.76 mg, 8.19 mol, 25% yield, 100% purity). ^1H NMR (400 MHz, CDCl_3) δ 7.64 (s, 1H), 7.30 (d, J =8.0 Hz, 2H), 7.17-7.12 (m, 3H), 7.09 (d, J =8.4 Hz, 2H), 7.06 (br s, 1H), 6.84-6.77 (m, 3H), 6.70 (d, J =2.4 Hz, 1H), 6.58 (br dd, J =2.8, 8.1 Hz, 3H), 6.28 (br d, J =8.4 Hz, 2H), 4.75-4.70 (m, 1H), 4.20 (br d, J =4.4 Hz, 1H), 4.14-4.03 (m, 2H), 3.41-3.26 (m, 1H), 3.10-2.87 (m, 6H), 2.71 (br d, J =7.6 Hz, 2H), 2.68 (s, 3H), 2.50-2.44 (m, 1H), 2.41 (s, 3H), 2.26-2.08 (m, 1H), 1.92-1.86 (m, 2H), 1.83-1.78 (m, 1H), 1.67 (s, 3H), 1.50-1.39 (m, 4H), 1.27 (s, 2H). LCMS (ES^+): RT=2.165 min, m/z =825.3 [$\text{M}+\text{H}^+$]; LCMS Method: 10.

Example 40—Synthesis of 3-(2,6-difluoro-4-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)phenyl)-4-methyl-2-phenyl-2H-chromen-6-ol (I-39) As a Mixture of Stereoisomer I and II



-continued



I-39

[1077] Step 1: Preparation of 2-[2-[3,5-difluoro-4-(4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-2H-chromen-3-yl)phenoxy]ethoxy]ethanol. To a solution of 2-[2-[tert-butyl(dimethyl)silyl]oxyethoxy]ethanol (97.41 mg, 0.44 mmol, 2 equiv) in DMSO (3 mL) was added NaH (17.68 mg, 0.4 mmol, 60% purity, 2 equiv) at 20° C. and the reaction mixture was stirred for 0.5 h. Then 4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-3-(2,4,6-trifluorophenyl)-2H-chromene (100 mg, 0.2 mmol, 1 equiv) was added and the mixture was stirred at 60° C. for 2.5 h. The reaction mixture

was quenched by the addition of water (50 mL) and then the reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (20 mL×3), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , petroleum ether/ethyl acetate=1/1) to afford 2-[2-[3,5-difluoro-4-(4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-2H-chromen-3-yl)phenoxy]

ethoxy]ethanol (52 mg, 42% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.645 min, m/z=561.2 [M+Na⁺].


[1078] Step 2: Preparation of 2-(2-(3,5-difluoro-4-(4-methyl-2-phenyl-6-((tetrahydro-2H-pyran-2-yl)oxy)-2H-chromen-3-yl)phenoxy)ethoxy)ethyl 4-methylbenzenesulfonate. To a solution of 2-[2-[3,5-difluoro-4-(4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-2H-chromen-3-yl)phenoxy]ethoxy]ethanol (140 mg, 0.26 mmol, 1 equiv) in DCM (3 mL) was added TosCl (99.12 mg, 0.5 mmol, 2 equiv), DMAP (3.18 mg, 0.025 mmol, 0.1 equiv) and TEA (78.91 mg, 0.78 mmol, 3 equiv). The mixture was stirred at 20° C. for 12 hr. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, petroleum ether/ethyl acetate=1/1) to afford 2-(2-(3,5-difluoro-4-(4-methyl-2-phenyl-6-((tetrahydro-2H-pyran-2-yl)oxy)-2H-chromen-3-yl)phenoxy)ethoxy)ethyl 4-methylbenzenesulfonate (135 mg, 60% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.703 min, m/z=691.3 [M-H⁺].

[1079] Step 3: Preparation of (6S)-4-(4-(2-(2-(3,5-difluoro-4-(4-methyl-2-phenyl-6-((tetrahydro-2H-pyran-2-yl)oxy)-2H-chromen-3-yl)phenoxy)ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine. To a solution of 2-[2-[3,5-difluoro-4-(4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-2H-chromen-3-yl)phenoxy]ethoxy]ethyl 4-methylbenzenesulfonate (150 mg, 0.21 mmol, 1 equiv) and 4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenol (73.28 mg, 0.21 mmol, 1 equiv) in DMF (3 mL) was added K₂CO₃ (89.77 mg, 0.64 mmol, 3 equiv). The mixture was stirred at 120° C. for 12 hr. The reaction mixture was diluted with water (20 mL) and extracted with ethyl acetate (20 mL×3). The combined organic layers were washed with brine (10 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM/MeOH=10/1) to afford (6S)-4-(4-(2-(2-(3,5-difluoro-4-(4-methyl-2-phenyl-6-((tetrahydro-2H-pyran-2-yl)oxy)-2H-chromen-3-yl)phenoxy)ethoxy)ethoxy)phenyl)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]


triazolo[4,3-a][1,4]diazepine (110 mg, 57% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.643 min, m/z=859.3 [M+H⁺]

[1080] Step 4: Preparation of 3-(2,6-difluoro-4-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)phenyl)-4-methyl-2-phenyl-2H-chromen-6-ol. To a solution of (9S)-7-[4-[2-[2-[3,5-difluoro-4-(4-methyl-2-phenyl-6-tetrahydropyran-2-yloxy-2H-chromen-3-yl)phenoxy]ethoxy]ethoxy]phenyl]-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (100 mg, 0.11 mmol, 1 equiv) in dioxane (3 mL) was added HCl (1 M, 2 mL, 17.18 equiv). The mixture was stirred at 20° C. for 1 hr. The reaction mixture was concentrated under reduced pressure to remove solvent. The residue was purified by prep-HPLC (column: YMC-Actus Triart C18 150 mm×30 mm×7 μm; mobile phase: [water (FA)-ACN]; gradient: 53%-83% B over 10 min) to afford 3-(2,6-difluoro-4-(2-(2-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenoxy)ethoxy)ethoxy)phenyl)-4-methyl-2-phenyl-2H-chromen-6-ol (I-39, 42.73 mg, 44% yield) as a mixture of stereoisomers, which were a white solid. ¹H NMR (400 MHz, DMSO-d₆) 7.50-7.39 (m, 2H), 7.31 (t, J=6.0 Hz, 2H), 7.24-7.07 (m, 4H), 7.05-6.98 (m, 1H), 6.92-6.81 (m, 3H), 6.75-6.66 (m, 1H), 6.49-6.26 (m, 2H), 6.06-5.86 (m, 1H), 5.35-5.25 (m, 1H), 4.22-4.11 (m, 3H), 4.10-3.93 (m, 4H), 3.93-3.80 (m, 3H), 3.75-3.53 (m, 2H), 2.69 (s, 3H), 2.41 (s, 3H), 2.11 (d, J=6.4 Hz, 3H), 2.01-1.93 (m, 3H), 1.91-1.79 (m, 3H), 1.72-1.70 (m, 2H), 1.68 (d, J=7.6 Hz, 2H). LC-MS: MS (ES⁺): RT=2.121 min, m/z=775.5 [M+H⁺]; LCMS Method: 25.





Example 41—Synthesis of 4-((4-(3,5-difluoro-4-((1S,2S)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-245) and 4-((4-(3,5-difluoro-4-((1R,2R)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-244)




-continued




-continued




-continued



Mixture of Two cis Diastereomers



III-245



III-244

[1081] Step 1: A mixture of 1-bromo-3,5-difluorobenzene (9 g, 46.63 mmol, 5.37 mL, 1 equiv), tert-butyl 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-3,6-dihydro-2H-pyridine-1-carboxylate (17.30 g, 55.96 mmol, 1.2 equiv), Cs_2CO_3 (45.58 g, 139.90 mmol, 3 equiv), and $\text{Pd}(\text{dppf})\text{Cl}_2$ (3.41 g, 4.66 mmol, 0.1 equiv) in dioxane (200 mL) and H_2O (40 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 100° C. for 12 h under N_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO_2 , PE:EA=3:1) to afford tert-butyl 4-(3,5-difluorophenyl)-3,6-dihydro-2H-pyridine-

1-carboxylate (13 g, 94% yield) as a yellow oil. LC-MS: MS (ES): RT=0.903 min, m/z =240.4 [M-55]; LCMS method 5-95.

[1082] Step 2: A mixture of tert-butyl 4-(3,5-difluorophenyl)-3,6-dihydro-2H-pyridine-1-carboxylate (13 g, 44.4 mmol, 1 equiv) and Pd/C (4.68 g, 4.40 mmol, 10% purity, 0.1 equiv) in THF (130 mL) was degassed and purged with H_2 for 3 times, and then the mixture was stirred at 20° C. for 12 h under H_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to afford tert-butyl 4-(3,5-difluorophenyl)piperidine-1-carboxylate (12.7 g, 97% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=0.922 min, m/z =242.2 [M-55]; LCMS method 5-95.

[1083] Step 3: A mixture of tert-butyl 4-(3,5-difluorophenyl)piperidine-1-carboxylate (4 g, 13.45 mmol, 1 equiv), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (5.12 g, 20.18 mmol, 1.5 equiv), 3,4,7,8-tetramethyl-1,10-phenanthroline (317 mg, 1.35 mmol, 0.1 equiv), and (1Z,5Z)-cycloocta-1,5-diene;2,4-dimethyl-BLAHbicyclo[1.1.0]butane (445 mg, 672 mol, 0.05 equiv) in THF (45 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 60° C. for 12 h under N₂ atmosphere. The reaction mixture was filtered, and the filtrate was concentrated to give crude compound as a yellow solid. The crude product tert-butyl 4-[3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperidine-1-carboxylate (5.6 g, crude) was used in the next step without further purification. LC-MS: MS (ES⁺): RT=0.993 min, m/z=368.1 [M-55]; LCMS method 5-95.

[1084] Step 4: A mixture of (6-benzyloxy-3,4-dihydronaphthalen-1-yl) trifluoromethanesulfonate (4.24 g, 11.02 mmol, 1 equiv), tert-butyl 4-[3,5-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperidine-1-carboxylate (5.6 g, 13.23 mmol, 1.2 equiv), Pd(dppf)Cl₂ (806 mg, 1.10 mmol, 0.1 equiv), and K₂CO₃ (4.57 g, 33.07 mmol, 3 equiv) in dioxane (50 mL) and H₂O (12 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 100° C. for 3 h under N₂ atmosphere. The reaction mixture was filtered and concentrated in vacuum. The residue was purified by prep-TLC (SiO₂, PE:EA=3:1) to afford tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3,5-difluoro-phenyl]piperidine-1-carboxylate (2.3 g, 39% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.090 min, m/z=476.1 [M-55]; LCMS method 5-95.

[1085] Step 5: To a solution of tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3,5-difluoro-phenyl]piperidine-1-carboxylate (2.3 g, 4.33 mmol, 1 equiv) in MeCN (20 mL) was added NBS (770 mg, 4.33 mmol, 1 equiv). The mixture was stirred at 20° C. for 2 h. The reaction mixture was partitioned between ethyl acetate 100 mL and water 50 mL. The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3,5-difluoro-phenyl]piperidine-1-carboxylate (2.2 g, 83% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=1.223 min, m/z=554.2 [M-55]; LCMS method 5-95.

[1086] Step 6: A mixture of (4-fluorophenyl)boronic acid (504 mg, 3.60 mmol, 1.1 equiv), tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3,5-difluoro-phenyl]piperidine-1-carboxylate (2 g, 3.28 mmol, 1 equiv), Pd(dppf)Cl₂ (239 mg, 327 mol, 0.1 equiv), and K₂CO₃ (1.13 g, 8.19 mmol, 2.5 equiv) in dioxane (30 mL) and H₂O (3 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 90° C. for 2 h under N₂ atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford tert-butyl 4-[4-[6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3,5-difluoro-phenyl]piperidine-1-carboxylate (2 g, 97% yield) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ=7.47-7.34 (m, 5H), 7.10-7.01 (m, 2H), 6.94-6.79 (m, 3H), 6.79-6.55 (m, 4H), 5.08 (s, 2H), 4.38-4.13 (m, 2H), 3.03 (t, J=7.8 Hz, 2H), 2.92-2.70 (m, 4H), 2.69-2.53 (m, 1H),

1.90-1.75 (m, 2H), 1.68-1.58 (m, 2H), 1.50 (s, 9H). LC-MS: MS (ES⁺): RT=1.238 min, m/z=526.3 [M-100]; LCMS method 5-95.

[1087] Step 7: To a solution of tert-butyl 4-[4-[6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3,5-difluoro-phenyl]piperidine-1-carboxylate (2 g, 3.20 mmol, 1 equiv) in MeOH (30 mL) was added Pd/C (680 mg, 639 mol, 10% purity, 0.2 equiv) and Pd(OH)₂ (448 mg, 639 mol, 20% purity, 0.2 equiv) under H₂ atmosphere. The suspension was degassed and purged with H₂ for 3 times. The mixture was stirred under H₂ (50 Psi.) at 50° C. for 12 h. The reaction mixture was filtered, and the filtrate was concentrated. The residue was purified by silica gel column chromatography (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford (±)-cis-tert-butyl 4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]phenyl]piperidine-1-carboxylate (1.7 g, 98% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=1.098 min, m/z=482.3 [M-55]; LCMS method 5-95.

[1088] Step 8: To a solution of (i)-cis-tert-butyl 4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]phenyl]piperidine-1-carboxylate (1.7 g, 3.16 mmol, 1 equiv) in DCM (20 mL) was added TIPSCl (1.22 g, 6.32 mmol, 1.35 mL, 2 equiv) and imidazole (645.81 mg, 9.49 mmol, 3 equiv) at 0° C. The mixture was stirred at 20° C. for 12 h. The reaction mixture was partitioned between DCM 200 mL and water 100 mL. The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford (±)-cis-tert-butyl 4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperidine-1-carboxylate (1 g, 45% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=1.280 min, m/z=594.4 [M-100]; LCMS method 5-95.

[1089] Step 9: To a solution of (i)-cis-tert-butyl 4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperidine-1-carboxylate (1 g, 1.44 mmol, 1 equiv) in DCM (6 mL) was added TFA (2.30 g, 20.19 mmol, 1.50 mL, 14.01 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between DCM 100 mL and water 50 mL. The organic phase was separated, filtered and concentrated under reduced pressure to afford (±)-cis-[1-[2,6-difluoro-4-(4-piperidyl)phenyl]-2-(4-fluorophenyl)tetralin-6-yl]oxy-triisopropyl-silane (855 mg, 99% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=0.954 min, m/z=594.4 [M+H⁺]; LCMS method 5-95.

[1090] Step 10: To a solution of (i)-cis-[1-[2,6-difluoro-4-(4-piperidyl)phenyl]-2-(4-fluorophenyl)tetralin-6-yl]oxy-triisopropyl-silane (1.7 g, 2.86 mmol, 1 equiv) in EtOH (20 mL) was added DIEA (1.11 g, 8.59 mmol, 1.50 mL, 3 equiv) and tert-butyl 1-oxa-6-azaspiro[2.5]octane-6-carboxylate (915 mg, 4.29 mmol, 1.5 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford (±)-cis-tert-butyl 4-[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl-4-hydroxy-piperidine-1-carboxylate (1.7 g, 73% yield) as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ=7.10-7.01 (m, 1H), 6.93-6.72 (m, 5H), 6.69-6.65 (m, 1H), 6.63-6.57 (m, 1H), 6.51-6.41 (m, 1H), 4.03-3.60 (m, 4H), 3.50-3.40 (m, 2H), 3.23-3.13 (m, 2H), 3.01-2.91 (m, 3H), 2.62-2.22 (m, 6H), 1.85-1.78 (m, 2H), 1.51-1.46 (m, 18H),

1.15-1.09 (m, 18H). LC-MS: MS (ES⁺): RT=1.012 min, m/z=807.5 [M+H⁺]; LCMS method 5-95.

[1091] Step 11: To a solution of (i)-cis-tert-butyl 4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]-4-hydroxy-piperidine-1-carboxylate (1 g, 1.24 mmol, 1 equiv) in DCM (5 mL) was added TFA (3.84 g, 33.66 mmol, 2.5 mL, 27.16 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between DCM 100 mL and NaHCO₃ 50 mL. The organic phase was separated, filtered and concentrated under reduced pressure to afford (±)-cis-4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]piperidin-4-ol (875 mg, 99% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=0.876 min, m/z=707.2 [M+H⁺]; LCMS method: 5-95.


[1092] Step 12: A mixture of (9S)-7-(4-chlorophenyl)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (326 mg, 913.49 mol, 1 equiv), (±)-cis-4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]piperidin-4-ol (775 mg, 1.10 mmol, 1.2 equiv), Cs₂CO₃ (892 mg, 2.74 mmol, 3 equiv) and SPhos Pd G₃ (71 mg, 91 mol, 0.1 equiv) in dioxane (10 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 90° C. for 3 h under N₂ atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford a mixture of the two cis-tetraline diastereomers of 4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperidin-4-ol (850 mg, 90% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.927 min, m/z=1026.9 [M+H⁺]; LCMS method 5-95.

[1093] Step 13: To a solution of the two cis-tetraline diastereomers of 4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperidin-4-ol (400 mg, 389 mol, 1 equiv) in THF (5 mL) was added TBAF (1 M, 1.00 mL, 2.57 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between EA 100 mL and water 50 mL. The


organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to afford a mixture of the two cis-tetraline diastereomers of 4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperidin-4-ol (300 mg, 88% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.761 min, m/z=871.3 [M+H⁺]; LCMS method: 5-95.

[1094] Step 14: A mixture of the two cis-tetraline diastereomers of 4-[[4-[3,5-difluoro-4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]phenyl]-1-piperidyl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]piperidin-4-ol (300 mg, 344 mol) was purified by prep-SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μm); mobile phase: [CO₂-ACN/i-PrOH (0.1% NH₃H₂O)]; B %:65%, isocratic elution mode) and (column: DAICEL CHIRALPAK IG (250 mm*30 mm, 10 μm); mobile phase: [CO₂-ACN/EtOH(0.1% NH₃H₂O)]; B %:65%, isocratic elution mode) to afford compounds III-245 and III-244. The second-eluting diastereomer, which was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below), was obtained as a yellow solid (45 mg, 16% yield). ¹H NMR (400 MHz, CDCl₃) δ=7.30 (d, J=8.8 Hz, 2H), 6.83-6.75 (m, 4H), 6.74-6.69 (m, 3H), 6.62 (d, J=2.4 Hz, 1H), 6.55-6.47 (m, 1H), 6.45-6.27 (m, 2H), 4.57 (d, J=5.6 Hz, 1H), 4.07 (d, J=6.8 Hz, 1H), 3.57-3.45 (m, 2H), 3.37-3.27 (m, 1H), 3.26-3.14 (m, 2H), 2.99-2.91 (m, 3H), 2.59 (s, 3H), 2.47-2.37 (m, 2H), 2.34 (s, 3H), 2.29-2.20 (m, 2H), 2.00 (d, J=6.8 Hz, 3H), 1.84-1.76 (m, 2H), 1.69 (s, 3H), 1.65-1.45 (m, 10H). LC-MS: MS (ES⁺): RT=1.530 min, m/z=871.3 [M+H⁺]; LCMS method 25.




Example 42—Synthesis of 4-((4-(2-fluoro-4-((1R,2S)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperazin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-251) and 4-((4-(2-fluoro-4-((1S,2R)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperazin-1-yl)methyl)-1-(4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-250)




350



-continued





[1095] Step 1: To a solution of 6-benzyloxytetralin-1-one (10 g, 39.63 mmol, 1.0 equiv) in dioxane (200 mL) and H₂O (50 mL) was added 1-fluoro-4-iodo-benzene (10.56 g, 47.56 mmol, 5.48 mL, 1.2 equiv), tritert-butylphosphonium tetrafluoroborate (2.30 g, 7.93 mmol, 0.2 equiv), Pd₂(dba)₃ (3.63 g, 3.96 mmol, 0.1 equiv) and NaOH (3.17 g, 79.27 mmol, 2.0 equiv). The mixture was stirred at 100° C. for 12 h. The reaction mixture was diluted with H₂O 100 mL and extracted with EA 150 mL (50 mL*3). The combined organic layers were washed with saturated NaCl aqueous (20 mL*2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to give 6-benzyloxy-2-(4-fluorophenyl)tetralin-1-one (12 g, 87% yield) as a brown solid. ¹H NMR (400 MHz, CDCl₃) δ 8.12-8.05 (m, 1H), 7.49-7.34 (m, 5H), 7.21-7.12 (m, 2H), 7.08-6.99 (m, 2H), 6.97-6.92 (m, 1H), 6.85-6.80 (m, 1H), 5.15 (s, 2H), 3.88-3.65 (m, 1H), 3.20-2.91 (m, 2H), 2.49-2.30 (m, 2H). LC-MS: MS (ES⁺): RT=0.667 min, m/z=347.1 [M+H⁺]; LCMS Method: 5-95.

[1096] Step 2: To a mixture of 6-benzyloxy-2-(4-fluorophenyl)tetralin-1-one (4.6 g, 13.28 mmol, 1.0 equiv) in THE (50 mL) was added NaH (1.06 g, 26.56 mmol, 60% purity, 2.0 equiv) at 0° C. for 0.5 h under N₂ atmosphere. Then 1,1,1-trifluoro-N-phenyl-N-(trifluoromethylsulfonyl) methanesulfonamide (13.05 g, 36.52 mmol, 2.75 equiv) was added to the mixture. The mixture was stirred at 25° C. for 12.5 h under N₂ atmosphere. The reaction mixture was quenched by addition saturated NH₄Cl (20 mL) under N₂

atmosphere, and then extracted with EA (30 mL*3). The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, PE/EA=100/1 to 1/1) to give [6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (10 g, 98% yield, 62% purity) as a yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 7.64-7.51 (m, 3H), 7.44-7.41 (m, 5H), 7.15-7.08 (m, 2H), 6.95-6.86 (m, 2H), 5.11 (s, 2H), 3.09-2.93 (m, 2H), 2.82-2.74 (m, 2H).

[1097] Step 3: To a solution of [6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (8.70 g, 11.27 mmol, 1.0 equiv) in DMAC (225 mL) and H₂O (45 mL) was added tert-butyl N-[2-fluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]carbamate (3.8 g, 11.27 mmol, 1.0 equiv), [2-(2-aminophenyl)phenyl]-methylsulfonyloxypalladium; dichloromethane; tricyclohexylphosphane (828.41 mg, 1.13 mmol, 0.1 equiv) and K₃PO₄ (7.18 g, 33.81 mmol, 3.0 equiv). The mixture was stirred at 60° C. for 12 h. The reaction mixture was diluted with H₂O 150 mL and extracted with EA (50 mL*3). The combined organic layers were washed with saturated NaCl aqueous (50 mL*2), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to give tert-butyl N-[4-[6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-2-fluorophenyl]carbamate (5.33 g, 88% yield) as

a yellow solid. LC-MS: MS (ES⁺): RT=1.227 min, m/z=562.3 [M+Na⁺]; LCMS Method: 5-95.

[1098] Step 4: To a solution of tert-butyl N-[4-[6-benzoyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-2-fluorophenyl]carbamate (5.3 g, 9.82 mmol, 1.0 equiv) in MeOH (60 mL) was added Pd/C (500 mg, 470 mol, 10% purity, 0.1 equiv) and Pd(OH)₂ (500 mg, 712 mol, 20% purity, 0.1 equiv) under N₂ atmosphere. The suspension was degassed and purged with H₂ for 3 times. The mixture was stirred under H₂ (50 psi) at 25° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give (±)-cis-tert-butyl 3-[(4-methoxycarbonylcyclohexyl)methyl]azetidine-1-carboxylate (4 g, 90% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.670 min, m/z=474.2 [M+Na⁺]; LCMS Method: 5-95.

[1099] Step 5: To a solution of (i)-cis-tert-butyl N-[2-fluoro-4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]phenyl]carbamate (2 g, 4.43 mmol, 1.0 equiv) in DCM (20 mL) was added TFA (3.07 g, 26.92 mmol, 2 mL, 6.08 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was added NaHCO₃ to adjust pH >7. To the reaction mixture was added water (2 mL) and the mixture was extracted with DCM (30 mL). The combined organic phase was washed with brine (3 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated in vacuum give (i)-cis-1-(4-amino-3-fluoro-phenyl)-2-(4-fluorophenyl)tetralin-6-ol (1.5 g, 96% yield) as a white oil. LC-MS: MS (ES⁺): RT=0.813 min, m/z=352.3 [M+H⁺]; LCMS Method: 5-95.

[1100] Step 6: To a solution of (i)-cis-1-(4-amino-3-fluoro-phenyl)-2-(4-fluorophenyl)tetralin-6-ol (1.5 g, 4.27 mmol, 1.0 equiv) in DCM (15 mL) was added TIPSCl (2.47 g, 12.81 mmol, 2.74 mL, 3.0 equiv) and imidazole (1.45 g, 21.34 mmol, 5.0 equiv). The mixture was stirred at 25° C. for 12 h. The reaction mixture was quenched by addition Na₂SO₃ (5 mL) at 0° C., and then diluted with H₂O 10 mL and extracted with EA 30 mL (10 mL×3). The combined organic layers were washed with saturated NaCl aqueous 30 mL (10 mL×3), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to give (±)-cis-2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]aniline (1.2 g, 2.36 mmol, 55% yield) as a white oil. ¹H NMR (400 MHz, CDCl₃) δ 6.93-6.84 (m, 2H), 6.81-6.74 (m, 4H), 6.67-6.61 (m, 1H), 6.45-6.38 (m, 1H), 6.11-6.03 (m, 1H), 5.98-5.92 (m, 1H), 4.17-4.10 (m, 1H), 3.39-3.29 (m, 1H), 3.06-2.95 (m, 2H), 2.20-2.05 (m, 1H), 1.86-1.72 (m, 1H), 1.33-1.23 (m, 3H), 1.20-1.05 (m, 18H). LC-MS: MS (ES⁺): RT=0.916 min, m/z=508.3 [M+H⁺]; LCMS Method: 5-95.

[1101] Step 7: To a solution of (i)-cis-2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]aniline (1 g, 1.97 mmol, 1.0 equiv) in THE (20 mL) was added NaNO₂ (271.78 mg, 3.94 mmol, 2.0 equiv) in H₂O (1 mL). Then was added H₂SO₄ (2 M, 2.95 mL, 3.0 equiv) at 0-5° C., the mixture was stirred at this temperature for 0.5 h. Then NaI (1.48 g, 9.85 mmol, 5.0 equiv) in H₂O (2.5 mL) was added into the mixture. Then mixture was stirred at 0-5° C. for 0.5 hr. Then the reaction mixture was added NaHCO₃ to adjust pH >7. Then the reaction mixture was added water (2 mL) and was extracted with DCM (5 mL). The combined organic phase was washed with brine (3 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated in vacuum to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 1/1) to

give (±)-cis-[1-(3-fluoro-4-iodo-phenyl)-2-(4-fluorophenyl)tetralin-6-yl]oxy-triisopropyl-silane (820 mg, 67% yield) as a white solid. LC-MS: MS (ES⁺): RT=1.280 min, m/z=619.2 [M+H⁺]; LCMS Method: 5-95.

[1102] Step 8: To a solution of (i)-cis-[1-(3-fluoro-4-iodo-phenyl)-2-(4-fluorophenyl)tetralin-6-yl]oxy-triisopropyl-silane (700 mg, 1.13 mmol, 1.0 equiv) in dioxane (15 mL) was added tert-butyl 4-hydroxy-4-(piperazin-1-ylmethyl)piperidine-1-carboxylate (677.60 mg, 2.26 mmol, 2.0 equiv), CPhos Pd G3 (182.49 mg, 226 mol, 0.2 equiv) and Cs₂CO₃ (1.11 g, 3.39 mmol, 3.0 equiv). The mixture was stirred at 90° C. for 3 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=100/1 to 0/1) to give (±)-cis-tert-butyl 4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxytetralin-1-yl]phenyl]piperazin-1-yl]methyl]-4-hydroxy-piperidine-1-carboxylate (240 mg, 27% yield) as a white solid. LC-MS: MS (ES⁺): RT=1.073 min, m/z=790.6 [M+H⁺]; LCMS Method: 5-95.

[1103] Step 9: To a solution of (i)-cis-tert-butyl 4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperazin-1-yl]methyl]-4-hydroxy-piperidine-1-carboxylate (240 mg, 304 mol, 1.0 equiv) in DCM (2 mL) was added TFA (307.00 mg, 2.69 mmol, 0.2 mL, 8.86 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was added NaHCO₃ to adjust pH >7. To the reaction mixture was added water (2 mL) and the mixture was extracted with DCM (5 mL). The combined organic phase was washed with brine (3 mL×3), dried over anhydrous Na₂SO₄, filtered and concentrated in vacuum to give (±)-cis-4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperazin-1-yl]methyl]piperidin-4-ol (209 mg, 99% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.711 min, m/z=690.4 [M+H⁺]; LCMS Method: 5-95.


[1104] Step 10: To a solution of (i)-cis-4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperazin-1-yl]methyl]piperidin-4-ol (209 mg, 308 mol, 1.0 equiv) in dioxane (8 mL) was added (9S)-7-(4-chlorophenyl)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (110 mg, 308 mol, 1.0 equiv), SPhos Pd G3 (48 mg, 63 mol, 0.2 equiv) and Cs₂CO₃ (301 mg, 925 mol, 3.0 equiv). The mixture was stirred at 90° C. for 3 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) to give a mixture of the two cis-tetraline diastereomers of 4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxytetralin-1-yl]phenyl]piperazin-1-yl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene-7-yl]phenyl]piperidin-4-ol (210 mg, 71% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.703 min, m/z=1010.3 [M+H⁺]; LCMS Method: 5-95.

[1105] Step 11: To a solution of the two cis-tetraline diastereomers of 4-[[4-[2-fluoro-4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]phenyl]piperazin-1-yl]methyl]-1-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene-7-yl]phenyl]piperidin-4-ol (210 mg, 208 mol, 1.0 equiv) in THE (1 mL) was added TBAF (1 M, 208 L, 1.0 equiv). The mixture was stirred at 25° C. for 2 h. The reaction mixture was filtered and concentrated under reduced pressure to give


a residue. The residue was purified by prep-TLC (SiO₂, DCM:MeOH=10:1) and SFC (Column: Chiralcel OD-350×4.6 mm I.D., 3 μm Mobile phase: Phase A for CO₂, and Phase B for IPA+ACN(0.05% DEA); Gradient elution: 40% IPA+ACN (0.05% DEA) in CO₂ Flow rate: 3 mL/min; Detector: PDA; Column Temp: 35°C; Back Pressure: 100 Bar) to give III-251 and III-250. The second-eluting diastereomer, which was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below), was obtained as a white solid (65.43 mg, 36% yield, 98.86% purity). ¹H NMR (400 MHz, CDCl₃) δ 7.42-7.35 (m, 2H), 6.95-6.82 (m, 4H), 6.81-6.70 (m, 4H), 6.65-6.49 (m, 2H), 6.17-6.06 (m, 2H), 4.19-4.10 (m, 2H), 3.3.63-3.52 (m, 2H), 3.39-3.31 (m, 1H), 3.30-3.19

(m, 2H), 3.09-2.93 (m, 6H), 2.86-2.75 (m, 4H), 2.67 (s, 3H), 2.41 (s, 5H), 2.14-2.05 (m, 4H), 1.77 (s, 6H), 1.70-1.66 (m, 2H). LCMS: MS (ES⁺): RT=1.493 min, m/z=854.3 [M+H⁺]. LCMS Method: 25.


Example 43—Synthesis of 4-((4-(4-((1S,2S)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-(((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-276) and 4-((4-(4-((1R,2R)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-(((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-275)




355




-continued



(or cis tetraline diastomer)



III-276



III-275

[1106] Step 1: A mixture of tert-butyl piperazine-1-carboxylate (2.98 g, 15 mmol, 1 equiv), 1-bromo-4-iodo-2-methoxy-benzene (5 g, 15 mmol, 1 equiv), $\text{Pd}_2(\text{dba})_3$ (1 g, 1 mmol, 0.1 equiv), Xantphos (924 mg, 1 mmol, 0.1 equiv) and Cs_2CO_3 (15 g, 47 mmol, 3 equiv) in DMF (50 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 100° C. for 12 h under N_2 atmosphere. The reaction mixture was diluted with water (20 mL) and then extracted with ethyl acetate (120 mL). The combined organic layer was washed with brine (30 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatog-

raphy (SiO_2 , Petroleum ether: Ethyl acetate=10/1) to give compound tert-butyl 4-(4-bromo-3-methoxy-phenyl)piperazine-1-carboxylate (4 g, 10.8 mmol, 67% yield) as a red solid, and used for next step directly. LC-MS: MS (ES^+): $\text{RT}=0.906$ min, $m/z=371.0$ [$\text{M}-100$]; LCMS Method: 5-95. **[1107]** Step 2: A mixture of tert-butyl 4-(4-bromo-3-methoxy-phenyl)piperazine-1-carboxylate (4 g, 10 mmol, 1 equiv), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (8 g, 32 mmol, 3 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2$ (788 mg, 1 mmol, 0.1 equiv), and potassium acetate (3 g, 32 mmol, 3 equiv) in dioxane (80 mL) was degassed and purged with N_2 for 3 times, and then

the mixture was stirred at 60° C. for 12 h under N₂ atmosphere. The reaction mixture was diluted with water (20 mL) and then extracted with ethyl acetate (60 mL×3). The combined organic layer was washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, Petroleum ether: Ethyl acetate=3/1) to give compound tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (4 g, 9 mmol, 88% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.883 min, m/z=419.4 [M+1]; LCMS Method: 5-95.

[1108] Step 3: A mixture of tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (2 g, 6 mmol, 1 equiv), [6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl] trifluoromethanesulfonate (3 g, 6 mmol, 1 equiv), tricyclohexylphosphane Pd G3 (460 mg, 627 mol, 0.1 equiv), K₃PO₄ (3 g, 18 mmol, 3 equiv) in H₂O (15 mL) and DMAC (75 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 60° C. for 12 h under N₂ atmosphere. The reaction mixture was diluted with water (20 mL) and then extracted with ethyl acetate (60 mL×3). The combined organic layer was washed with brine (30 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography (SiO₂, Petroleum ether: Ethyl acetate=5/1) to give compound tert-butyl 4-[4-[6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.8 g, 6 mmol, 97% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=1.058 min, m/z=621.4 [M+1]; LCMS Method: 5-95.

[1109] Step 4: Palladium on carbon (1.5 g, 1 mmol, 10% purity, 0.2 equiv) and Pd(OH)₂ (1.5 g, 2 mmol, 20% purity, 0.4 equiv) were added into a 250 mL single-necked round bottom flask under N₂, and then MeOH (80 mL) was added at 25° C. under N₂. After addition, tert-butyl 4-[4-[6-benzyloxy-2-(4-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.8 g, 6 mmol, 1 equiv) was added under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (50 psi) at 50° C. for 48 h. The residue was filtered and concentrated under reduced pressure to give (±)-cis-tert-butyl 4-[4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.2 g, 98% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.948 min, m/z=533.3 [M+1]; LCMS Method: 5-95.

[1110] Step 5: Racemic cis-tert-butyl 4-[4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (500 mg, 0.938 mmol) was purified by prep-SFC (column: DAICEL CHIRALPAK AD (250 mm×30 mm, 10 μm); mobile phase: [CO₂-EtOH (0.1% NH₃H₂O)]; B %: 40%, isocratic elution mode) to afford (1S,2S)- and (1R,2R)-tert-butyl 4-[4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. The faster-eluting enantiomer (146 mg, 274.10 mol, 29.20% yield, 99% ee) was obtained as a yellow oil, and the slower-eluting enantiomer (188 mg, 37% yield, 99% ee) was obtained as a yellow solid. LC-MS: MS (ES⁺): RT=1.025 min, m/z=533.3 [M+H⁺]; LCMS Method: 5-95.

[1111] The following Steps 6 through 11 describe further synthetic procedures conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[4-[2-(4-fluorophenyl)-6-hydroxy-

tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the diastereomer of compounds III-276 and III-275 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of cis-tert-butyl 4-[4-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-276 and III-275.

[1112] Step 6: To a solution of tert-butyl 4-[4-(1S,2S)-[2-(4-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (188 mg, 352 mol, 1 equiv) in DCM (2 mL) was added TIPSCl (136 mg, 705 mol, 151 μL, 2 equiv) and imidazole (72 mg, 1.06 mmol, 3 equiv) at 0° C. The mixture was stirred at 20° C. for 12 h. The reaction mixture was partitioned between dichloromethane (100 mL) and water (50 mL). The organic phase was separated, filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, Petroleum ether: Ethyl acetate=3:1) to afford tert-butyl 4-[4-(1S,2S)-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (240 mg, 98% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=1.338 min, m/z=689.5 [M+1]; LCMS Method: 5-95.




[1113] Step 7: To a solution of tert-butyl 4-[4-(1S,2S)-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (240 mg, 348 mol, 1 equiv) in DCM (2 mL) was added TFA (1.54 g, 13.46 mmol, 1 mL, 38 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between dichloromethane (100 mL) and water (50 mL). The organic phase was separated, filtered and concentrated under reduced pressure to afford [2-(1S,2S)-[4-(4-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-yl]oxy-triisopropyl-silane or its enantiomer (205 mg, 99% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=1.046 min, m/z=589.5 [M+1]; LCMS Method: 5-95.


[1114] Step 8: To a solution of [2-(1S,2S)-[4-(4-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-yl]oxy-triisopropyl-silane or its enantiomer (180 mg, 305 mol, 1 equiv) in EtOH (3 mL) was added DIEA (118 mg, 917 mol, 159 μL, 3 equiv) and tert-butyl 1-oxa-6-azaspiro[2.5]octane-6-carboxylate (97 mg, 458 μmol, 1.5 equiv). The mixture was stirred at 80° C. for 2 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-TLC (SiO₂, Petroleum ether: Ethyl acetate=3:1) to afford tert-butyl 4-[4-[4-(1S,2S)-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]-3-methoxy-phenyl]piperazin-1-yl]methyl-4-hydroxy-piperidine-1-carboxylate or its enantiomer (186 mg, 75% yield) as a yellow oil. LC-MS: MS (ES⁺): RT=0.986 min, m/z=802.2 [M+1]; LCMS Method: 5-95.

[1115] Step 9: To a solution of tert-butyl 4-[4-(1S,2S)-[4-[2-(4-fluorophenyl)-6-triisopropylsilyloxy-tetralin-1-yl]-3-methoxy-phenyl]piperazin-1-yl]methyl-4-hydroxy-piperidine-1-carboxylate or its enantiomer (140 mg, 174 mol, 1 equiv) in DCM (1 mL) was added TFA (2.15 g, 18.85 mmol, 1.40 mL, 107 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between aqueous NaHCO₃ (30 mL) and dichloromethane (50 mL). The organic phase was separated, filtered and concentrated under

359

-continued





[1118] Step 1: To a solution of 6-benzyloxytetralin-1-one (5 g, 19 mmol, 1.0 equiv) and 1-fluoro-2-iodo-benzene (6.6 g, 29 mmol, 3.4 mL, 1.5 equiv) in dioxane (40.0 mL) and H₂O (10.0 mL) was added Pd₂(dba)₃ (1.8 g, 1 mmol, 0.1 equiv), NaOH (1.5 g, 39 mmol, 2.0 equiv) and tritert-butylphosphonium tetrafluoroborate (1.1 g, 3 mmol, 0.2 equiv) in one portion at 25° C. Then the resulting mixture was heated to 100° C. and stirred under N₂ atmosphere for 12 h. The residue was diluted with water (10.0 mL) and extracted with ethyl acetate (20.0 mL*3). The combined organic layers were washed with saturated aqueous NaCl solution (10.0 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=20/1 to 1/1) to give 6-benzyloxy-2-(2-fluorophenyl)tetralin-1-one (6.4 g, 93% yield) as yellow solid. LC-MS: MS (ES⁺): RT=0.653 min, m/z=347.1 [M+H⁺]. LCMS Method: 5-95.

[1119] Step 2: To a mixture of 6-benzyloxy-2-(2-fluorophenyl)tetralin-1-one (6 g, 17 mmol, 1.0 equiv) in THE (60.0 mL) was added NaH (1.3 g, 34 mmol, 60% purity, 2.0 equiv) at 0° C. for 0.5 h under N₂ atmosphere. Then 1,1,1-trifluoro-N-phenyl-N-(trifluoromethylsulfonyl) methanesulfonamide (18.5 g, 51 mmol, 3.0 equiv) was added to the mixture. The mixture was stirred at 25° C. for 12 h under N₂ atmosphere. The reaction mixture was quenched with saturated aqueous NH₄Cl (50.0 mL) solution at 0° C. dropwise. And the resulting mixture was extracted with ethyl acetate (30.0 mL*3). The combined organic layers

were dried over anhydrous Na₂SO₄, filtered and concentrated in vacuum. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=1/0 to 5/1) to give [6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (3.9 g, 47% yield) as white solid. ¹H NMR (400 MHz, DMSO-d₆): δ 7.52-7.45 (m, 4H), 7.43-7.34 (m, 3H), 7.33-7.24 (m, 3H), 7.09-6.98 (m, 2H), 5.16 (s, 2H), 2.96 (s, 2H), 2.76-2.66 (m, 2H).

[1120] Step 3: To a solution of [6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (2.9 g, 6 mmol, 1.0 equiv) and tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (2.5 g, 6 mmol, 1.0 equiv) in H₂O (8.0 mL) and toluene (32.0 mL) was added K₃PO₄ (3.8 g, 18 mmol, 3.0 equiv), SPhos Pd G₃ (472 mg, 606 mol, 0.1 equiv) in one portion at 25° C. Then the resulting mixture was heated to 60° C. and stirred under N₂ atmosphere for 12 h. The residue was diluted with water (30.0 mL) and extracted with ethyl acetate (30.0 mL*3). The combined organic layers were washed with saturated aqueous NaCl solution (30.0 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=5/1 to 3/1) to give tert-butyl 4-[4-[6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.5 g, 93% yield) as white solid. LC-MS: MS (ES⁺): RT=0.795 min, m/z=621.5 [M+H⁺]. LCMS Method: 5-95.

[1121] Step 4: To a solution of tert-butyl 4-[4-[6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.5 g, 5 mmol, 1.0 equiv) in THE (35.0 mL) was added Pd/C (6.0 g, 5 mmol, 10% purity, 1.0 equiv) and Pd(OH)₂ (3.9 g, 5 mmol, 20% purity, 1.0 equiv) under N₂ atmosphere. The suspension was degassed and purged with H₂ for 3 times. The mixture was stirred under H₂ (50 Psi) at 40° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The crude product was triturated with MeOH at 25° C. for 30 min to give (±)-cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (2.1 g, 69% yield) as white solid. LC-MS: MS (ES⁺): RT=3.038 min, m/z=533.4 [M+H⁺]. LCMS Method: 5-95.

[1122] Step 5: Racemic cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (700 mg, 1 mmol, 1.0 equiv) was purified by preparative SFC (column: DAICEL CHIRALPAK AD(250 mm*30 mm,10 um);mobile phase: [CO₂—IPA(0.1% NH₃H₂O)];B %:25%, isocratic elution mode) to give (1S,2S)- and (1R,2R)-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. The slower-eluting enantiomer (340 mg) was obtained as a white solid. LC-MS: MS (ES⁺): RT=0.655 min, m/z=533.2 [M+H⁺]; LCMS Method: 5-95.


[1123] The following Steps 6 and 7 describe further synthetic procedures conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the diastereomer of compounds III-307 and III-308 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-307 and III-308.

[1124] Step 6: To a solution of tert-butyl 4-[4-[(1S,2S)-2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phe-


nyl]piperazine-1-carboxylate or its enantiomer (150 mg, 281 mol, 1.0 equiv) in DCM (1.0 mL) was added TFA (307 mg, 2 mmol, 0.2 mL, 9.5 equiv). The mixture was stirred at 25° C. for 1 h. The mixture was concentrated to give a residue. The crude (1S,2S)-2-(2-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (121 mg, crude) as white solid was used into the next step without further purification. LC-MS: MS (ES⁺): RT=0.536 min, m/z=433.2[M+H⁺]. LCMS Method: 5-95.


[1125] Step 7: To a solution of (1S,2S)-2-(2-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (121 mg, 279 mol, 1.0 equiv) in EtOH (4.0 mL) was added DIEA (180 mg, 1 mmol, 243 L, 5.0 equiv) and 6-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (303 mg, 699 mol, 2.5 equiv) at 0° C. The mixture was stirred at 80° C. for 12 h. The mixture was concentrated to give a residue. The residue was purified by preparative HPLC (column: Waters Xbridge C18 150*50 mm*10 um;mobile phase: [H₂O(10 mM NH₄HCO₃)-ACN]; gradient: 58%-88% B over 11.0 min) to give compound III-307 or III-308 (118 mg, 47% yield, 97% purity) as white solid. ¹H NMR (400 MHz, DMSO-d₆): δ 9.29-8.86 (m, 1H), 7.26 (s, 2H), 7.16-7.04 (m, 2H), 6.89 (s, 2H), 6.80-6.73 (m, 1H), 6.59-6.51 (m, 2H), 6.46-6.40 (m, 1H), 6.36-6.21 (m, 3H), 6.06 (s, 1H), 4.72 (s, 1H), 4.23 (s, 1H), 4.10 (s, 1H), 3.54-3.42 (m, 6H), 3.30 (s, 3H), 3.14 (s, 2H), 3.00 (s, 3H), 2.95 (s, 5H), 2.60 (s, 3H), 2.57 (s, 2H), 2.47 (s, 1H), 2.38 (s, 3H), 2.28 (s, 1H), 1.83 (s, 3H), 1.64 (s, 3H), 1.52 (s, 3H). LC-MS: MS (ES⁺): RT=1.779 min, m/z=866.7[M+H⁺]. LCMS method: 25.

Example 45—Synthesis of 4-((4-(2-fluoro-4-((1S,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-5-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-311) and 4-((4-(2-fluoro-4-((1R,2R)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-5-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-312)



-continued





[1126] Step 1: A mixture of 1-bromo-4-chloro-2-fluoro-5-methoxy-benzene (5 g, 20.8 mmol, 1.0 equiv), tert-butyl piperazine-1-carboxylate (3.89 g, 20.9 mmol, 1.0 equiv), [1-(2-diphenylphosphanyl-1-naphthyl)-2-naphthyl]-diphenyl-phosphane (2.60 g, 4.18 mmol, 0.2 equiv), $\text{Pd}_2(\text{dba})_3$ (1.91 g, 2.09 mmol, 0.1 equiv) and t-BuONa (4.01 g, 41.8 mmol, 2.0 equiv) in toluene (50 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 60° C. for 12 h under N_2 atmosphere. The mixture was filtered and concentrated to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum ether: Ethyl acetate from 50/1 to 5/1) to give tert-butyl 4-(4-chloro-2-fluoro-5-methoxy-phenyl)piperazine-1-carboxylate (6.6 g, 92% yield) as a yellow solid. ^1H NMR (400 MHz, CDCl_3): δ 7.08 (d, $J=11.6$ Hz, 1H), 6.50 (d, $J=7.6$ Hz, 1H), 3.86 (s, 3H), 3.65-3.55 (m, 4H), 3.08-2.96 (m, 4H), 1.49 (s, 9H). LC-MS: MS (ES^+): RT=0.684 min, $m/z=345.1$ [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1127] Step 2: A mixture of tert-butyl 4-(4-chloro-2-fluoro-5-methoxy-phenyl)piperazine-1-carboxylate (8.5 g, 24.7 mmol, 1.0 equiv), BPD (12.5 g, 49.3 mmol, 2.0 equiv), XPhos (1.76 g, 3.70 mmol, 0.15 equiv), $\text{Pd}_2(\text{dba})_3$ (1.13 g, 1.23 mmol, 0.05 equiv) and KOAc (7.26 g, 74.0 mmol, 3.0 equiv) in dioxane (100 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 110° C. for 12 h under N_2 atmosphere. The mixture was filtered and concentrated to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum

ether: Ethyl acetate from 50/1 to 3/1) to give tert-butyl 4-[2-fluoro-5-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (9.4 g, 87% yield) as a yellow solid. ^1H NMR (400 MHz, CDCl_3): δ 7.34 (d, $J=13.2$ Hz, 1H), 6.39 (d, $J=6.8$ Hz, 1H), 3.80 (s, 3H), 3.66-3.54 (m, 4H), 3.14-3.02 (m, 4H), 1.48 (s, 9H), 1.33 (s, 12H). LC-MS: MS (ES^+): RT=0.713 min, $m/z=437.3$ [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1128] Step 3: To a solution of tert-butyl 4-[2-fluoro-5-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (3.41 g, 7.82 mmol, 1.2 equiv) in toluene (40 mL) and H_2O (10 mL) and was added K_3PO_4 (4.15 g, 19.6 mmol, 3.0 equiv), (6-benzyloxy-2-phenyl-3,4-dihydronaphthalen-1-yl) trifluoromethanesulfonate (3 g, 6.52 mmol, 1.0 equiv) and SPhos Pd G_3 (508 mg, 652 μmol , 0.1 equiv). The mixture was stirred at 60° C. for 12 h under N_2 atmosphere. The mixture was filtered and concentrated to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum ether: Ethyl acetate from 50/1 to 5/1) to give tert-butyl 4-[4-(6-benzyloxy-2-phenyl-3,4-dihydronaphthalen-1-yl)-2-fluoro-5-methoxy-phenyl]piperazine-1-carboxylate (2.4 g, 59% yield) as a yellow solid. ^1H NMR (400 MHz, CDCl_3): δ 7.47-7.42 (m, 2H), 7.39 (t, $J=7.6$ Hz, 2H), 7.36-7.30 (m, 1H), 7.15-7.08 (m, 2H), 7.08-7.01 (m, 3H), 6.85 (d, $J=1.6$ Hz, 1H), 6.72-6.59 (m, 3H), 6.37 (d, $J=7.2$ Hz, 1H), 5.07 (s, 2H), 3.60 (t, $J=4.8$ Hz, 4H), 3.48 (s, 3H), 3.11-2.94 (m, 6H), 2.79 (m, $J=3.6$, 7.6 Hz, 2H), 1.50 (s, 9H). LC-MS: MS (ES^+): RT=0.727 min, $m/z=621.3$ [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1129] Step 4: To a solution of tert-butyl 4-[4-(6-benzyloxy-2-phenyl-1,3,4-dihydronaphthalen-1-yl)-2-fluoro-5-methoxyphenyl]piperazine-1-carboxylate (2.4 g, 3.87 mmol, 1.0 equiv) in THE (24 mL) and MeOH (12 mL) was added Pd/C (1 g, 10% purity) and Pd(OH)₂ (1 g) under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (50 psi) at 40° C. for 16 h. The mixture was filtered and concentrated to give a residue. The crude product was triturated with PE:EA=5/1 (36 mL) at 25° C. for 1 h. The mixture was filtered to give (±)-cis-tert-butyl 4-[2-fluoro-4-(6-hydroxy-2-phenyl-tetralin-1-yl)-5-methoxy-phenyl]piperazine-1-carboxylate (1.7 g, 83% yield) as a white solid. ¹H NMR (400 MHz, CDCl₃): δ 7.15-7.02 (m, 3H), 6.70-6.82 (m, 3H), 6.69 (d, J=2.0 Hz, 1H), 6.55-6.63 (m, 1H), 6.37 (d, J=13.6 Hz, 1H), 6.06 (d, J=7.2 Hz, 1H), 5.34-5.14 (m, 1H), 4.81 (d, J=5.2 Hz, 1H), 3.57 (s, 4H), 3.31 (m, J=5.2, 11.6 Hz, 1H), 3.09-2.88 (m, 9H), 2.24 (m, J=6.4, 12.8 Hz, 1H), 1.80-1.63 (m, 2H), 1.49 (s, 9H). LC-MS: MS (ES⁺): RT=3.255 min, m/z=533.4 [M+H⁺]. LCMS Method: 5-95.


[1130] Step 5: Racemic cis-tert-butyl 4-[2-fluoro-4-(6-hydroxy-2-phenyl-tetralin-1-yl)-5-methoxy-phenyl]piperazine-1-carboxylate (700 mg, 1.31 mmol, 1.0 equiv) was purified by preparative SFC (column: DAICEL CHIRAL-CEL OD (250 mm*30 mm, 10 μm); mobile phase: [CO₂—IPA (0.1% NH₃H₂O)]; B %:55%, isocratic elution mode) to give (1S,2S)- and (1R,2R)-tert-butyl 4-[2-fluoro-4-[6-hydroxy-2-phenyl-tetralin-1-yl]-5-methoxy-phenyl]piperazine-1-carboxylate. The faster-eluting enantiomer was isolated (340 mg, 49% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.651 min, m/z=533.3 [M+H⁺]; LCMS Method: 5-95.


[1131] The following Steps 6 and 7 describe further synthetic procedures conducted on the faster-eluting enantiomer of cis-tert-butyl 4-[2-fluoro-4-(6-hydroxy-2-phenyl-tetralin-1-yl)-5-methoxy-phenyl]piperazine-1-carboxylate. This faster-eluting enantiomer produced the diastereomer of compounds III-311 and III-312 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The slower-eluting enantiomer of cis-tert-butyl 4-[2-fluoro-4-(6-hydroxy-2-phenyl-tetralin-1-yl)-5-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-311 and III-312.

[1132] Step 6: To a solution of tert-butyl 4-[2-fluoro-4-[(1S,2S)-6-hydroxy-2-phenyl-tetralin-1-yl]-5-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (150 mg, 281 mol, 1.0 equiv) in DCM (1.5 mL) was added TFA (461 mg, 4.04 mmol, 0.3 mL, 14.3 equiv). The mixture was stirred at 25° C. for 1 h. The mixture was concentrated to give (1S,2S)-1-(5-fluoro-2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol or its enantiomer (120 mg, 99% yield) as a colorless oil. LC-MS: MS (ES⁺): RT=0.489 min, m/z=433.2 [M+H⁺]. LCMS Method: 5-95.


[1133] Step 7: To a solution of 6-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (241 mg, 555 mol, 2.0 equiv) in EtOH (3 mL) was added DIEA (179 mg, 1.39 mmol, 0.2 mL, 5.0 equiv) and (1S,2S)-1-(5-fluoro-2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol or its enantiomer (120 mg, 277 mol, 1.0 equiv). The mixture was stirred at 80° C. for 12 h. The mixture was concentrated to give a residue. The residue was purified by preparative HPLC (column: Waters Xbridge C18 150*25 mm*5 μm; mobile phase: [H₂O (10 mM NH₄HCO₃)-ACN]; gradient: 64%-94% B over 9.0 min) to give compound III-311 or III-312 (74 mg, 30% yield, 98% purity) as an off-white solid. ¹H NMR (400 MHz, CDCl₃): δ 7.38 (d, J=8.8 Hz, 2H), 7.12-7.04 (m, 3H), 6.86 (d, J=9.2 Hz, 2H), 6.81-6.74 (m, 3H), 6.69 (d, J=2.4 Hz, 1H), 6.65-6.55 (m, 1H), 6.35 (d, J=13.6 Hz, 1H), 6.06 (d, J=7.6 Hz, 1H), 5.56-5.11 (m, 1H), 4.80 (d, J=5.2 Hz, 1H), 4.15 (q, J=6.8 Hz, 1H), 3.70-3.50 (m, 2H), 3.34-3.20 (m, 3H), 3.12-2.95 (m, 9H), 2.80 (d, J=4.0 Hz, 4H), 2.67 (s, 3H), 2.46-2.35 (m, 5H), 2.30-2.19 (m, 1H), 2.08 (d, J=6.8 Hz, 3H), 1.82-1.65 (m, 8H). LC-MS: MS (ES⁺): RT=1.796 min, m/z=866.7 [M+H⁺]. LCMS method: 25.

Example 46—Synthesis of 4-((4-((1S,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-321) and 4-((4-((1R,2R)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-322)





-continued



[1134] Step 1: A mixture of tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (4.80 g, 11.4 mmol, 1.0 equiv), (6-benzyloxy-3,4-dihydronaphthalen-1-yl) trifluoromethanesulfonate (5.29 g, 13.7 mmol, 1.2 equiv), K_2CO_3 (4.76 g, 34.4 mmol, 3.0 equiv) and $Pd(dppf)Cl_2$ (834 mg, 1.15 mmol, 0.1 equiv) in dioxane (80.0 mL) and H_2O (20.0 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at $100^\circ C$. for 3 h under N_2 atmosphere. The mixture was filtered, and the filtrate was concentrated under reduced pressure to remove solvent and give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=20/1 to 3/1) to obtain compound tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (4.88 g, 8.80 mmol, 77% yield) as a pale yellow solid. LC-MS: MS (ES^+): RT=0.763 min, m/z =527.3 [$M+H$] $^+$.

[1135] Step 2: To a solution of tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (4.88 g, 8.80 mmol, 1.0 equiv) in MeCN (50.0 mL) was added NBS (1.48 g, 8.33 mmol, 0.9 equiv) solution in MeCN (50.0 mL) dropwise. The mixture was stirred at $25^\circ C$. for 1 h. The reaction mixture was quenched by addition of saturated NH_4Cl (aq, 100 mL) at $0^\circ C$., and then extracted with EtOAc (100 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=10/1) to obtain compound tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (4.20 g, 6.94 mmol, 75% yield) as a colorless oil. LC-MS: MS (ES^+): RT=0.783 min, m/z =607.2 [$M+H$] $^+$.

len-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (4.20 g, 6.94 mmol, 75% yield) as a colorless oil. LC-MS: MS (ES^+): RT=0.783 min, m/z =607.2 [$M+H$] $^+$.

[1136] Step 3: A mixture of tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (1.40 g, 2.31 mmol, 1.0 equiv), 4,4,5,5-tetramethyl-2-phenyl-1,3,2-dioxaborolane (708 mg, 3.47 mmol, 1.5 equiv), K_2CO_3 (799 mg, 5.78 mmol, 2.5 equiv) and $Pd(dppf)Cl_2$ (169 mg, 231 μ mol, 0.1 equiv) in dioxane (10.0 mL) and H_2O (1.0 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at $90^\circ C$. for 2 h under N_2 atmosphere. The reaction mixture was treated under reduced pressure to remove solvent and give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=10/1) to obtain compound tert-butyl 4-[4-(6-benzyloxy-2-phenyl-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (1.41 g, crude) as a pale yellow solid. LC-MS: MS (ES^+): RT=0.800 min, m/z =603.4 [$M+H$] $^+$; LCMS Method: 5-95.

[1137] Step 4: A solution of tert-butyl 4-[4-(6-benzyloxy-2-phenyl-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (1.4 g) in MeOH (14.0 mL) and THE (14.0 mL) was prepared. To a fixed bed flow reactor was added 5% $Pd(OH)_2/Al_2O_3$. The H_2 back pressure regulator was adjusted to 1.0 MPa, and the flow rate of H_2 was 30 mL/min. Then the solution was pumped (0.3 mL/min) through fixed bed reactor (set at $50^\circ C$.). The flow process continued for 2 h. The resultant reaction solution was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex

luna C18 150×40 mm×15 μm; mobile phase: [H₂O (0.225% FA)-ACN]; gradient: 60%-90% B over 15.0 min) to obtain compound (±)-cis-tert-butyl 4-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (765 mg, 1.47 mmol, 55% yield) as a pale yellow solid. LC-MS: MS (ES⁺): RT=0.690 min, m/z=515.3 [M+H]⁺; LCMS Method: 5-95.

[1138] Step 5: The compound (±)-cis-tert-butyl 4-[4-(6-hydroxy-2-phenyl-tetralin-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (765 mg, 1.47 mmol, 1.0 equiv) was separated by Chiral SFC (column: DAICEL CHIRALCEL OD(250 mm×30 mm, 10 μm); mobile phase: [CO₂-IPA(0.1% NH₃H₂O)]; B %: 50%, isocratic elution mode; RT=1.785 and 1.994 min.) to obtain (1S,2S)- and (1R,2R)-tert-butyl 4-[4-[6-hydroxy-2-phenyl-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. The faster-eluting enantiomer was isolated (282 mg, 510 μmol) as a pale yellow solid; and the slower-eluting enantiomer was isolated (301 mg, 550 μmol) as a pale yellow solid. LC-MS: MS (ES⁺): RT=0.691 min, m/z=515.3 [M+H]⁺ (isomer-1); 0.690 min, m/z=515.3 [M+H]⁺ (isomer-2); LCMS Method: 5-95.


[1139] The following Steps 6 and 7 describe further synthetic procedures conducted on the faster-eluting enantiomer of cis-tert-butyl 4-[4-[6-hydroxy-2-phenyl-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This faster-eluting enantiomer produced the diastereomer of compounds III-321 and III-322 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The slower-eluting enantiomer of cis-tert-butyl 4-[4-[6-hydroxy-2-phenyl-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-321 and III-322.

[1140] Step 6: To a solution of tert-butyl 4-[4-[(1S,2S)-6-hydroxy-2-phenyl-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (100 mg, 194 μmol, 1.0 equiv) in DCM (1.0 mL) was added TFA (0.5 mL). The mixture was stirred at 25° C. for 0.5 h. The reaction mixture


was concentrated under reduced pressure to obtain compound (1S,2S)-1-(2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol or its enantiomer (110 mg, crude, TFA salt) as a green oil. LC-MS: MS (ES⁺): RT=0.563 min, m/z=415.2 [M+H]⁺; LCMS Method: 5-95.

[1141] Step 7: To a solution of (1S,2S)-1-(2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol or its enantiomer (110 mg, 208 μmol, 1.0 equiv, TFA salt) and 6-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (246 mg, 250 μmol, 1.2 equiv) in EtOH (5.0 mL) was added DIEA (80.7 mg, 624 μmol, 109 μL, 3.0 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was treated under reduced pressure to remove solvent and give a residue. The residue was purified by prep-HPLC (column: Waters Xbridge C18 150×25 mm×5 μm; mobile phase: [H₂O(10 mM NH₄HCO₃)-ACN]; gradient: 53%-83% B over 15.0 min) to obtain compound III-321 or III-322 (65.4 mg, 74.3 μmol, 36% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=2.592 min, m/z=848.8 [M+H]⁺; LCMS Method: 01. ¹H NMR (400 MHz, CD₃OD) δ=7.31 (br d, J=8.4 Hz, 2H), 7.05-7.00 (m, 3H), 6.94 (br d, J=8.9 Hz, 2H), 6.75-6.69 (m, 2H), 6.64-6.57 (m, 2H), 6.54-6.41 (m, 2H), 6.35 (dd, J=1.7 Hz, 1H), 6.15 (d, J=1.7 Hz, 1H), 4.75 (br d, J=5.1 Hz, 1H), 4.24-4.14 (m, 1H), 3.60 (br dd, J=4.1 Hz, 2H), 3.28-3.18 (m, 3H), 3.09-2.92 (m, 9H), 2.77-2.66 (m, 7H), 2.46-2.35 (m, 5H), 2.34-2.22 (m, 1H), 1.96 (d, J=6.7 Hz, 3H), 1.78 (dt, J=3.9 Hz, 2H).

Example 47—Synthesis of 1-(3-fluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-((1S,2S)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-328) and 1-(3-fluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-((1R,2R)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-327)



-continued



[1142] Step 1: A mixture of (9S)-7-(4-chloro-2-fluorophenyl)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (2.1 g, 5.6 mmol, 1.0 equiv), 4-(triisopropylsilyloxymethyl)piperidin-4-ol (1.9 g, 6.7 mmol, 1.2 equiv), SPhos Pd G3 (874 mg, 1.1 mmol, 0.2 equiv), Cs₂CO₃ (5.5 g, 16.8 mmol, 3.0 equiv) in dioxane (20.0 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 90° C. for 12 h under N₂ atmosphere. The reaction mixture was partitioned between Ethyl acetate 200 mL and H₂O 50 mL. The organic phase was separated, washed with NaCl 50 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The reaction mixture was filtered and concentrated under reduced pressure to give compound 1-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(triisopropylsilyloxymethyl)piperidin-4-ol (3.5 g, 5.6 mmol, 99% yield) as a brown solid. LC-MS: MS (ES⁺): RT=0.623 min, m/z=626.4 [M+1]⁺. LCMS Method: 5-95.

[1143] Step 2: To a solution of 1-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(triisopropylsilyloxymethyl)piperidin-4-ol (3.5 g, 5.6 mmol, 1.0 equiv) in THE (35 mL) was added TBAF (1.00 M, 5.6 mL, 1.0 equiv) at 0-25° C. The mixture was stirred at 0-25° C. for 12 h. The reaction mixture was partitioned between Ethyl acetate 200 mL and H₂O 50 mL. The organic phase was separated, washed with NaCl 50 mL, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=0/1 to Ethyl acetate:MeOH=10:1) to give compound 1-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)piperidin-4-ol (1.4 g, 3.0 mmol, 53% yield) as a yellow solid. LC-MS: MS (ES⁺): RT=0.478 min, m/z=470.2 [M+1]⁺. LCMS Method: 5-95.


[1144] Step 3: To a solution of 1-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)piperidin-4-ol (200 mg, 426 mol, 1.0 equiv) and 1,1,2,2,3,3,4,4,4-nonafluorobutane-1-sulfonyl fluoride (386 mg, 1.3 mmol, 225 μL, 3.0 equiv) in THE (3.0 mL) was added DBU (324 mg, 2.1 mmol, 321 L, 5.0 equiv). The mixture was stirred at 25° C. for 1 h. The mixture was quenched with H₂O (10 mL) and extracted with DCM 20 mL (10 mL×2). The combined organic layers were dried over anhydrous sodium sulfate, filtered and concentrated under reduced pressure to give a residue. The crude product was used into the next step without purification. Compound

6-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (200 mg, crude) was obtained as a brown oil. LC-MS: MS (ES⁺): RT=0.467 min, m/z=452.2 [M+1]⁺. LCMS Method: 5-95.


[1145] Step 4: To a solution of 6-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (80.0 mg, 177 mol, 1.0 equiv) and the slower-eluting enantiomer of (5S,6S)- or (5R,6R)-6-(4-fluorophenyl)-5-(2-methoxy-4-(piperazin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol (76.6 mg, 177 mol, 1.0 equiv; prepared as described in Example 43) in EtOH (2 mL) was added DIEA (22.9 mg, 177 mol, 30.8 μL, 1.0 equiv). The mixture was stirred at 80° C. for 12 h. It was filtered. The residue was purified by prep-HPLC (column: Waters xbridge 150*25 mm 10 μm; mobile phase: [H₂O(10 mM NH₄HCO₃)-ACN]; gradient: 68-88% B over 8.0 min) (column: Phenomenex Luna C18 150*25 mm*10 μm; mobile phase: [H₂O(0.225% FA)-ACN]; gradient: 3%-33% B over 10.0 min) to afford compound III-328 or III-327 (34.92 mg, 39.50 mol, 22% yield) as a yellow solid. ¹H NMR (400 MHz, CD₃OD) δ 7.33 (br t, J=8.9 Hz, 1H), 6.82-6.69 (m, 5H), 6.64-6.57 (m, 3H), 6.51-6.43 (m, 2H), 6.36 (dd, J=2.1, 8.5 Hz, 1H), 6.18 (d, J=2.0 Hz, 1H), 4.72 (br d, J=5.4 Hz, 1H), 4.22 (q, J=6.7 Hz, 1H), 3.67-3.57 (m, 2H), 3.25 (br d, J=11.8 Hz, 3H), 3.11 (s, 3H), 3.06 (br s, 4H), 3.00-2.93 (m, 2H), 2.73 (br d, J=4.5 Hz, 4H), 2.67 (s, 3H), 2.43-2.36 (m, 5H), 2.31-2.17 (m, 1H), 1.97 (d, J=6.9 Hz, 3H), 1.81-1.73 (m, 2H), 1.71-1.60 (m, 6H). LC-MS: MS (ES⁺): RT=2.674 min, m/z=884.7 [M+H]⁺; LCMS Method: 01. This diastereomer of compounds III-328 and III-327 was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below).

[1146] The faster-eluting enantiomer of cis-6-(4-fluorophenyl)-5-(2-methoxy-4-(piperazin-1-yl)phenyl)-5,6,7,8-tetrahydronaphthalen-2-ol was used in a similar reaction to afford the other one of compounds III-327 and III-328.

Example 48—Synthesis of 4-((4-((1S,2S)-2-(2,4-difluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-334) and 4-((4-((1R,2R)-2-(2,4-difluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)-1-(4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-333)






370



371

-continued



[1147] Step 1: A mixture of (6-benzyloxy-3,4-dihydronaphthalen-1-yl) trifluoromethanesulfonate (1.8 g, 4.68 mmol, 1.0 equiv), tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (2.55 g, 6 mmol, 1.3 equiv), K_2CO_3 (1.62 g, 12 mmol, 2.5 equiv), and $Pd(dppf)Cl_2$ (171 mg, 234 μ mol, 0.05 equiv) in dioxane (20 mL) and H_2O (4 mL) in one portion at 25° C. Then the resulting mixture was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 90° C. for 2 h under N_2 atmosphere. The residue was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=1/0 to 20/1) to give tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (2.0 g, 81% yield) as a white solid. 1H NMR (400 MHz, $CDCl_3$): δ 7.53-7.30 (m, 7H), 7.15-7.07 (m, 1H), 6.84 (s, 1H), 6.67 (s, 2H), 6.63-6.51 (m, 1H), 5.93-5.95 (m, 1H), 5.06 (s, 2H), 3.77-3.56 (m, 6H), 3.30-3.15 (m, 4H), 2.92-2.80 (m, 2H), 2.47-2.36 (m, 2H), 1.52 (s, 9H). LC-MS: MS (ES^+): RT=0.767 min, m/z =527.3 [$M+H^+$]. LCMS method: 5-95.

[1148] Step 2: To a solution of tert-butyl 4-[4-(6-benzyloxy-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (1.80 g, 3 mmol, 1.0 equiv) in ACN (20 mL) was added NBS (547 mg, 3 mmol, 0.9 equiv). The mixture was stirred at 25° C. for 1 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=1/0 to 20/1) to give tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (1.20 g, 58% yield) as a white solid. LC-MS: MS (ES^+): RT=0.777 min, m/z =605.2 [$M+H^+$]. LCMS method: 5-95.

[1149] Step 3: A mixture of tert-butyl 4-[4-(6-benzyloxy-2-bromo-3,4-dihydronaphthalen-1-yl)-3-methoxy-phenyl]piperazine-1-carboxylate (500 mg, 826 μ mol, 1.0 equiv), 2-(2,4-difluorophenyl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (208 mg, 867 μ mol, 1.05 equiv), K_2CO_3 (285 mg, 2 mmol, 2.5 equiv), and $Pd(dppf)Cl_2$ (30 mg, 41 μ mol, 0.05 equiv) in dioxane (5 mL) and H_2O (0.5 mL) in one portion at 25° C. Then the resulting mixture was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 90° C. for 2 h under N_2 atmosphere. The residue was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 150*40 mm* 15 μ m; mobile phase: [H_2O (0.225% FA)-ACN]; gradient:90%-100% B over 15.0 min) to give tert-butyl 4-[4-[6-benzyloxy-2-(2,4-difluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (420 mg, 80% yield) as a white solid. LC-MS: MS (ES^+): RT=0.797 min, m/z =639.3 [$M+H^+$]. LCMS method: 5-95.

[1150] Step 4: Pd/C (200 mg, 10% purity), $Pd(OH)_2$ (200 mg, 10% purity) was added into a flask under N_2 , tert-butyl 4-[4-[6-benzyloxy-2-(2,4-difluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (420 mg, 657 μ mol, 1.0 equiv) in THF (4 mL) and MeOH (2 mL) was added into the mixture. The suspension was degassed and purged with H_2 for 3 times. The mixture was stirred under H_2 (50 Psi) at 40° C. for 48 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The crude ($\pm$)-cis-tert-butyl 4-[4-[2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-

phenyl]piperazine-1-carboxylate (360 mg, crude) as a white solid was used into the next step without further purification. LC-MS: MS (ES^+): RT=0.682 min, m/z =551.4 [$M+H^+$]. LCMS method: 5-95.


[1151] Step 5: The ($\pm$)-cis-tert-butyl 4-[4-[2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (360 mg, crude) was purified by preparative SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μ m); mobile phase: [CO_2 -IPA(0.1% NH_3H_2O)]; B %:35%, isocratic elution mode) to give (1S,2S)- and (1R, 2R)-tert-butyl 4-[4-[2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. The slower-eluting enantiomer (150 mg, 42% yield) was obtained as a white solid. LC-MS: MS (ES^+): RT=0.689 min, m/z =551.2 [$M+H^+$]. LCMS method: 5-95.

[1152] The following Steps 6 and 7 describe further synthetic procedures conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the diastereomer of compounds III-333 and III-334 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-333 and III-334.




[1153] Step 6: To a solution of tert-butyl 4-[4-[(1S,2S)-2-(2,4-difluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (150 mg, 272 μ mol, 1.0 equiv) in DCM (1.5 mL) was added TFA (767 mg, 7 mmol, 0.5 mL, 25.0 equiv). The mixture was stirred at 25° C. for 2 h. The mixture was diluted with $NaHCO_3$ (10 mL), then extracted with DCM (10 mL *3). The combined organic layers were washed with H_2O (10 mL *2), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The crude (1S,2S)-2-(2,4-difluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (122 mg, crude) as a yellow solid was used into the next step without further purification. LC-MS: MS (ES^+): RT=0.548 min, m/z =451.1 [$M+H^+$]. LCMS method: 5-95.

[1154] Step 7: To a solution of (1S,2S)-2-(2,4-difluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (122 mg, 271 μ mol, 1.0 equiv) in EtOH (4 mL) was added DIEA (105 mg, 812 μ mol, 141 L, 3.0 equiv) and 6-[4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (141 mg, 325 μ mol, 1.2 equiv) at 0° C. The mixture was stirred at 80° C. for 12 h. The mixture was concentrated to give a residue. The residue was purified by preparative HPLC (column: Waters Xbridge C18 150*50 mm*10 μ m; mobile phase: [H_2O (10 mM NH_4HCO_3)-ACN]; gradient:60%-90% B over 11.0 min) to give compound III-334 or III-333 (114 mg, 48% yield) as a yellow solid. 1H NMR (400 MHz, $CDCl_3$): δ 7.43-7.35 (m, 2H), 6.93-6.84 (m, 2H), 6.81-6.72 (m, 2H), 6.70 (s, 1H), 6.63-6.57 (m, 1H), 6.53-6.43 (m, 2H), 6.34-6.24 (m, 2H), 6.07 (s, 1H), 5.35 (s, 1H), 4.88 (s, 1H), 4.24-4.07 (m, 1H), 3.73-3.51 (m, 3H), 3.33-3.23 (m, 2H), 3.20 (s, 3H), 3.17-3.06 (m, 4H), 3.05-2.98 (m, 2H), 2.78 (br s, 4H), 2.68 (s, 3H), 2.47-2.33 (m, 5H), 2.31-2.16 (m, 1H), 2.13-2.05 (m, 3H), 1.78 (s, 3H), 1.62 (s, 6H). LC-MS: MS (ES^+): RT=1.84 min, m/z =884.8 [$M+H^+$]. LCMS method: 25.

Example 49—Synthesis of 4-((4-(3-chloro-4-((1S, 2S)-6-hydroxy-2-(4-methoxyphenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperazin-1-yl)methyl)-1-((4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-336) and 4-((4-(3-chloro-4-((1R,2R)-6-hydroxy-2-(4-methoxyphenyl)-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperazin-1-yl)methyl)-1-((4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-335)




374



-continued

or



III-335

[1155] Step 1: A mixture of tert-butyl piperazine-1-carboxylate (10 g, 53 mmol, 1 equiv), 1-bromo-2-chloro-4-iodo-benzene (20.45 g, 64 mmol, 1.2 equiv), $\text{Pd}_2(\text{dba})_3$ (4.92 g, 5.37 mmol, 0.1 equiv), Xantphos (9.32 g, 16.11 mmol, 0.3 equiv) and NaOtBu (15.48 g, 161.07 mmol, 3 equiv) in Tol. (180 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 60° C. for 12 h under N_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum ether: Ethyl acetate from 1/0 to 3/1) to afford desired product tert-butyl 4-(4-bromo-3-chloro-phenyl)piperazine-1-carboxylate (19 g, 94% yield) as a yellow oil. LC-MS: MS (ES^+): RT=0.643 min, m/z =320.9 [$\text{M}-55$]. LCMS method: 5-95.

[1156] Step 2: A mixture of tert-butyl 4-(4-bromo-3-chloro-phenyl)piperazine-1-carboxylate (15 g, 39.93 mmol, 1 equiv), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (20.28 g, 79.85 mmol, 2 equiv), KOAc (7.84 g, 79.85 mmol, 2 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2$ (5.84 g, 7.99 mmol, 0.2 eq) and 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (20.28 g, 79.85 mmol, 2 equiv) in dioxane (200 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 90° C. for 3 h under N_2 atmosphere. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by silica column chromatography on silica gel (Petroleum ether: Ethyl acetate from 1/0 to 10/1) to afford tert-butyl 4-[3-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (16 g, 94% yield) as yellow oil. ^1H NMR (400 MHz, CDCl_3): δ =7.63 (d, J=8.4 Hz, 1H), 6.86 (d, J=2.4 Hz, 1H), 6.75 (dd, J=2.4, 8.4 Hz, 1H), 3.60-3.57 (m, 4H), 3.24-3.21 (m, 4H), 1.50 (s, 9H), 1.36 (s, 12H). LC-MS: MS (ES^+): RT=0.711 min, m/z =423.1 [$\text{M}+\text{H}^+$]. LCMS method: 5-95.

[1157] Step 3: A mixture of [6-benzyloxy-2-(4-methoxyphenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (6.54 g, 13.33 mmol, 1 equiv), tert-butyl 4-[3-chloro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (6.2 g, 14.67 mmol, 1.1 equiv), tricyclohexylphosphane Pd G3 (980 mg, 1.33 mmol, 0.1 equiv) and K_3PO_4 (8.49 g, 40.00 mmol, 3 equiv) in H_2O

(20 mL) and DMAC (100 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 60° C. for 12 h under N_2 atmosphere. The reaction mixture was filtered and concentrated in vacuum. The residue was purified by prep-TLC (SiO_2 , PE:EA=3:1) to afford tert-butyl 4-[4-[6-benzyloxy-2-(4-methoxyphenyl)-3,4-dihydronaphthalen-1-yl]-3-chloro-phenyl]piperazine-1-carboxylate (3.6 g, 42% yield) as a yellow solid. LC-MS: MS (ES^+): RT=0.806 min, m/z =637.1 [$\text{M}+\text{H}^+$]. LCMS method: 5-95.

[1158] Step 4: To a solution of tert-butyl 4-[4-[6-benzyloxy-2-(4-methoxyphenyl)-3,4-dihydronaphthalen-1-yl]-3-chloro-phenyl]piperazine-1-carboxylate (3.45 g, 5.41 mmol, 1 equiv) in MeOH (15 mL) and THE (15 mL) was added LiCl (229 mg, 5.41 mmol, 110 μL , 1 equiv) and Pd/C (1.15 g, 1.08 mmol, 10% purity, 0.2 equiv) under N_2 atmosphere. The suspension was degassed and purged with H_2 for 3 times. The mixture was stirred under H_2 (50 Psi) at 45° C. for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to afford ($\pm$)-cis-tert-butyl 4-[3-chloro-4-[6-hydroxy-2-(4-methoxyphenyl)tetralin-1-yl]phenyl]piperazine-1-carboxylate (600 mg, 20% yield) as a yellow oil. ^1H NMR (400 MHz, CDCl_3): δ =7.11 (s, 2H), 6.98-6.85 (m, 3H), 6.80-6.74 (m, 2H), 6.64-6.58 (m, 3H), 4.13-4.06 (m, 1H), 3.67 (s, 3H), 3.54-3.46 (m, 5H), 3.06-3.01 (m, 4H), 2.86-2.80 (m, 1H), 2.58-2.51 (m, 1H), 2.33-2.25 (m, 1H), 1.94-1.89 (m, 1H), 1.41 (s, 9H). LC-MS: MS (ES^+): RT=0.684 min, m/z =549.2 [$\text{M}+\text{H}^+$]. LCMS method: 5-95.

[1159] Step 5: Racemic cis-tert-butyl 4-[3-chloro-4-[6-hydroxy-2-(4-methoxyphenyl)tetralin-1-yl]phenyl]piperazine-1-carboxylate (600 mg, 1.09 mmol) was purified by prep-SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μm); mobile phase: [CO_2 -EtOH (0.1% $\text{NH}_3\text{H}_2\text{O}$)]); B %:45%, isocratic elution mode) to afford (1S,2S)- and (1R,2R)-tert-butyl 4-[3-chloro-4-[6-hydroxy-2-(4-methoxyphenyl)tetralin-1-yl]phenyl]piperazine-1-carboxylate. The slower-eluting enantiomer (238 mg, 39% yield) was obtained as a yellow solid. LC-MS: MS (ES^+): RT=0.682 min, m/z =549.3 [$\text{M}+\text{H}^+$]. LCMS method: 5-95.

[1160] The following Steps 6 and 7 describe further synthetic procedures conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[3-chloro-4-[6-hydroxy-2-(4-methoxyphenyl)tetralin-1-yl]phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the


diastereomer of compounds III-335 and III-336 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of *cis*-*tert*-butyl 4-[3-chloro-4-[6-hydroxy-2-(4-methoxyphenyl) tetralin-1-yl]phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-335 and III-336.

[1161] Step 6: To a solution of *tert*-butyl 4-[3-chloro-4-[(1*S*,2*S*)-6-hydroxy-2-(4-methoxyphenyl)tetralin-1-yl]phenyl]piperazine-1-carboxylate or its enantiomer (119 mg, 216 mol, 1 equiv) in DCM (1 mL) was added TFA (767 mg, 6.73 mmol, 0.5 mL, 31.06 equiv). The mixture was stirred at 20° C. for 0.5 h. The reaction mixture was partitioned between NaHCO₃ 30 mL and DCM 50 mL. The organic phase was separated, filtered and concentrated under reduced pressure to afford desired product (1*S*,2*S*)-1-(2-chloro-4-piperazin-1-yl-phenyl)-2-(4-methoxyphenyl)tetralin-6-ol or its enantiomer (97 mg, 99% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.480 min, *m/z*=449.1 [M+H⁺]. LCMS method: 5-95.


[1162] Step 7: To a solution of 6-[4-[(9*S*)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (281 mg, 648 mol, 3 equiv) in EtOH (3 mL) was added DIEA (83 mg, 648 mol, 112 μL, 3 equiv) and (1*S*,2*S*)-1-(2-chloro-4-piperazin-1-yl-phenyl)-2-(4-

methoxyphenyl)tetralin-6-ol or its enantiomer (97 mg, 216 mol, 1 equiv). The mixture was stirred at 80° C. for 2 h. The mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Waters xbridge 150*25 mm 10 μm; mobile phase: [H₂O (10 mM NH₄HCO₃)-ACN]; gradient: 55%-85% B over 15.0 min) to afford compound III-336 or III-335 (46 mg, 23% yield) as a yellow solid. ¹H NMR (400 MHz, MeOD): δ=7.32 (d, *J*=6.0 Hz, 2H), 6.96 (d, *J*=7.6 Hz, 2H), 6.71 (s, 2H), 6.67-6.57 (m, 7H), 6.51-6.41 (m, 1H), 4.80-4.75 (m, 1H), 4.29-4.15 (m, 1H), 3.72 (s, 3H), 3.65-3.55 (m, 2H), 3.29-3.21 (m, 3H), 3.10-3.05 (m, 4H), 3.04-2.94 (m, 2H), 2.76-2.66 (m, 7H), 2.45-2.36 (m, 5H), 2.00-1.974 (m, 3H), 1.87-1.74 (m, 3H), 1.70-1.5 (m, 6H). LC-MS: MS (ES⁺): RT=2.625 min, *m/z*=882.7 [M+H⁺]. LCMS Method: 01.

Example 50—Synthesis of 4-((4-(3,5-difluoro-4-((1*S*,2*S*)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl)-1-(3-fluoro-4-((6*S*)-2,3,6,9-tetramethyl-6H-thieno[3,2-*f*][1,2,4]triazolo[4,3-*a*][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (III-338) and 4-((4-(3,5-difluoro-4-((1*R*,2*R*)-2-(4-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)phenyl)piperidin-1-yl)methyl)-1-(3-fluoro-4-((6*S*)-2,3,6,9-tetramethyl-6H-thieno[3,2-*f*][1,2,4]triazolo[4,3-*a*][1,4]diazepin-4-yl)phenyl)piperidin-4-ol (II-337)



-continued




[1163] Step 1: To a mixture of (1S,2S)-1-[2,6-difluoro-4-(4-piperidyl)phenyl]-2-(4-fluorophenyl)tetralin-6-ol or its enantiomer (106 mg, 0.24 mmol, 1.0 equiv) in EtOH (5 mL) was added DIEA (156 mg, 1.21 mmol, 0.21 mL, 5.0 equiv) and 6-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (144 mg, 0.32 mmol, 1.3 equiv) under N₂. The mixture was stirred at 80° C. for 15 h under N₂. The reaction mixture was concentrated in vacuum to give a residue. The residue was purified by Prep-HPLC (column: Waters xbridge 150*25 mm 10 um; mobile phase: [H₂O (10 mM NH₄HCO₃)-ACN]; gradient: 65%-95% B over 9.0 min) to give compound III-338 or III-337 (114.21 mg, 127.81 mol, 53% yield, 99.5% purity) as a white solid. LCMS: MS(ES⁺): RT=2.701 min, m/z=889.7 [M+H⁺]. LCMS method: 01. ¹H NMR (400 MHz, DMSO-d₆): δ 9.11 (s, 1H), 7.28 (t, J=8.4 Hz, 1H), 6.98-6.86 (m, 4H), 6.77 (dd, J₁=1.6 Hz, J₂=8.8 Hz, 1H), 6.69-6.52 (m, 5H), 6.42 (dd, J=2.0, J₂=8.0 Hz, 1H), 4.54 (d, J=6.0 Hz, 1H), 4.19 (s, 1H), 4.16-4.09 (m, 1H), 3.57-3.39 (m, 3H), 3.14 (t, J=11.2 Hz, 2H), 2.95-2.93 (m, 4H), 2.57 (s,

3H), 2.39-2.29 (m, 4H), 2.25 (s, 2H), 2.22-2.11 (m, 3H), 1.84 (d, J=6.8 Hz, 3H), 1.76-1.73 (m, 1H), 1.64 (s, 3H), 1.62-1.58 (m, 4H), 1.53-1.41 (m, 4H). This diastereomer of compounds III-337 and III-338 was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below).


[1164] The other enantiomer of cis-1-[2,6-difluoro-4-(4-piperidyl)phenyl]-2-(4-fluorophenyl)tetralin-6-ol was used in a similar reaction to afford the other one of compounds III-337 and III-338.

Example 51—Synthesis of 1-(3-fluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-(4-((1S,2S)-2-(3-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-344) and 1-(3-fluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-(4-((1R,2R)-2-(3-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-343)




378

-continued



-continued



[1165] Step 1: A mixture of 6-benzyloxytetralin-1-one (8.0 g, 31.7 mmol, 1.0 equiv), 1-fluoro-3-iodo-benzene (8.5 g, 38.1 mmol, 4.5 mL, 1.2 equiv), NaOH (1.3 g, 31.7 mmol, 1.0 equiv), $\text{Pd}(\text{dba})_3$ (2.9 g, 3.2 mmol, 0.1 equiv) and tritert-butylphosphonium tetrafluoroborate (1.8 g, 6.3 mmol, 0.2 equiv) in dioxane (80.0 mL) and H_2O (20.0 mL) was degassed and purged with N_2 for 3 times, and then the mixture was stirred at 100°C . for 12 h under N_2 atmosphere. The reaction mixture was partitioned between Ethyl acetate (300 mL) and H_2O (300 mL $\times$ 2). The organic phase was separated, washed with brine (300 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Welch Ultimate XB— NH_2 250*50 mm*10 um;mobile phase: [Heptane-EtOH(0.1% $\text{NH}_3\cdot\text{H}_2\text{O}$)]gradient:1%-30% B over 15.0 min) to give the 6-benzyloxy-2-(3-fluorophenyl)tetralin-1-one (3.1 g, 9.0 mmol, 28% yield) as a yellow solid. LC-MS: MS (ES^+): RT=0.703 min, m/z =347.1 [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1166] Step 2: To a solution of 6-benzyloxy-2-(3-fluorophenyl)tetralin-1-one (3.1 g, 9.0 mmol, 1.0 equiv) in THF (31.0 mL) was added NaH (716 mg, 17.9 mmol, 60% purity, 2.0 equiv) at 0°C . under N_2 , the mixture was stirred at 25°C . for 30 min, then the 1,1,1-trifluoro-N-phenyl-N-(trifluoromethylsulfonyl)methanesulfonamide (6.4 g, 17.9 mmol, 2.0 equiv) was added to the mixture at 0°C . The mixture was stirred at 25°C . for 12 h under N_2 . The reaction mixture was quenched with saturated aqueous NH_4Cl (10 mL) solution at 0°C . dropwise. And the resulting mixture was extracted with Ethyl acetate (70 mL $\times$ 3). The organic phase was separated, washed with brine (100 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=1/0 to 20/1) to give the [6-benzyloxy-2-(3-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (3.3 g, 6.9 mmol, 77% yield) as a white solid. LC-MS: MS (ES^+): RT=0.736 min, m/z =501.2 [$\text{M}+\text{Na}^+$]. LCMS Method: 5-95.

[1167] Step 3: A mixture of [6-benzyloxy-2-(3-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (3.0 g, 6.3 mmol, 1.0 equiv), tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (3.2 g, 7.5 mmol, 1.2 equiv), $\text{Pd}(\text{dppf})\text{Cl}_2$ (459 mg, 627 μmol , 0.1 equiv), Cs_2CO_3 (4.1 g, 12.5 mmol, 2.0 equiv) in dioxane (30.0 mL) and H_2O (3.0 mL) was

degassed and purged with N_2 for 3 times, and then the mixture was stirred at 80°C . for 1 h under N_2 atmosphere. The reaction mixture was partitioned between Ethyl acetate (100 mL) and H_2O (100 mL $\times$ 2). The organic phase was separated, washed with brine (100 mL), dried over Na_2SO_4 , filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Phenomenex luna C18 250*50 mm*10 um;mobile phase: [H_2O (0.225% FA)-ACN];gradient:70%-100% B over 22.0 min) to give the tert-butyl 4-[4-[6-benzyloxy-2-(3-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (2.0 g, 3.2 mmol, 51% yield, 99% purity) as a yellow solid. LC-MS: MS (ES^+): RT=0.731 min, m/z =621.3 [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1168] Step 4: Pd/C (1.0 g, 940 μmol , 10% purity, 2.92e-1 equiv) and $\text{Pd}(\text{OH})_2$ (1.0 g, 1.4 mmol, 20% purity, 4.42e-1 equiv) was added into a 75 mL single-necked round bottom flask under N_2 , and then MeOH (10.0 mL) was added at 25°C . under N_2 . After addition, tert-butyl 4-[4-[6-benzyloxy-2-(3-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (2.0 g, 3.2 mmol, 1.0 equiv) in THF (10.0 mL) was added under N_2 . The suspension was degassed under vacuum and purged with H_2 several times. The mixture was stirred under H_2 (50 PSI) at 50°C . for 12 h. The reaction mixture was filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO_2 , Petroleum ether/Ethyl acetate=20/1 to 10/1) to give the ($\pm$)-cis-tert-butyl 4-[4-[2-(3-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (1.5 g, 2.8 mmol, 87% yield) as a yellow solid. LC-MS: MS (ES^+): RT=0.653 min, m/z =533.2 [$\text{M}+\text{H}^+$]. LCMS Method: 5-95.

[1169] Step 5: The residue was purified by SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μm); mobile phase: [CO_2 -EtOH(0.1% $\text{NH}_3\cdot\text{H}_2\text{O}$)]B %:45%, isocratic elution mode) to give the (1S,2S)- and (1R,2R)-tert-butyl 4-[4-[2-(3-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. The faster-eluting enantiomer (Rt=1.195 min) was isolated (730 mg, 1.4 mmol, 49% yield) as a yellow solid, and the slower-eluting enantiomer (Rt=1.437 min) was isolated (720 mg, 1.4 mmol, 48% yield) as a yellow solid.

[1170] The following Steps 6 and 7 describe further synthetic procedures conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[4-[2-(3-fluorophenyl)-6-hydroxy-


tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the diastereomer of compounds III-343 and III-344 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of *cis*-tert-butyl 4-[4-[2-(3-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same steps to afford the other one of compounds III-343 and III-344.

[1171] Step 6: To a solution of tert-butyl 4-[4-[(1*S*,2*S*)-2-(3-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (720 mg, 1.4 mmol, 1.0 equiv) in DCM (6.0 mL) was added TFA (2.0 mL). The mixture was stirred at 25° C. for 1 h. The reaction was added saturated aqueous NaHCO₃ solution to adjust the pH>7 and the reaction mixture was partitioned between DCM (80 mL) and H₂O (80 mL×2). The organic phase was separated, washed with brine (80 mL), dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue to give the (1*S*,2*S*)-2-(3-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (crude) as a white solid.

[1172] Step 7: A mixture of (1*S*,2*S*)-2-(3-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (109 mg, 252 μmol, 1.2 equiv), 6-[3-fluoro-4-[(9*S*)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (95 mg, 210 μmol, 1.0 equiv) and




DIEA (136 mg, 1.1 mmol, 183 μL, 5.0 equiv) in EtOH (2.0 mL) was degassed and purged with N₂ for 3 times, and then the mixture was stirred at 80° C. for 12 h under N₂ atmosphere. The reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Waters xbridge 150×25 mm 10 μm; mobile phase: [H₂O(10 mM NH₄HCO₃)-ACN]; gradient: 62%-82% B over 8.0 min) to give compound III-344 or III-343 (90 mg, 99 μmol, 47% yield, 97% purity) as a yellow solid. LC-MS: MS (ES⁺): RT=2.655 min, m/z=884.7 [M+H⁺]. LCMS Method: 01. ¹H NMR: (400 MHz, CD₃OD) δ=7.40-7.28 (m, 1H), 7.11-7.00 (m, 1H), 6.84-6.71 (m, 2H), 6.68-6.55 (m, 4H), 6.53-6.41 (m, 2H), 6.41-6.30 (m, 2H), 6.19 (d, J=2.0 Hz, 1H), 4.76 (d, J=5.2 Hz, 1H), 4.28-4.16 (m, 1H), 3.70-3.53 (m, 2H), 3.29-3.22 (m, 3H), 3.10 (s, 3H), 3.06 (s, 4H), 3.00-2.90 (m, 2H), 2.73 (d, J=4.4 Hz, 4H), 2.68 (s, 3H), 2.41-2.34 (m, 5H), 2.30-2.19 (m, 1H), 1.97 (d, J=6.8 Hz, 3H), 1.84-1.71 (m, 2H), 1.70-1.58 (m, 6H).

Example 52—Synthesis of 1-(2-fluoro-4-((6*S*)-2,3,6,9-tetramethyl-6H-thieno[3,2-*f*][1,2,4]triazolo[4,3-*a*][1,4]diazepin-4-yl)phenyl)-4-((4-((1*S*,2*S*)-2-(2-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-346) and 1-(2-fluoro-4-((6*S*)-2,3,6,9-tetramethyl-6H-thieno[3,2-*f*][1,2,4]triazolo[4,3-*a*][1,4]diazepin-4-yl)phenyl)-4-((4-((1*R*,2*R*)-2-(2-fluorophenyl)-6-hydroxy-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-345)




381

-continued



III-346

or



[1173] Step 1: To a solution of [6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]trifluoromethanesulfonate (3.0 g, 6.3 mmol, 1.0 equiv) and tert-butyl 4-[3-methoxy-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl]piperazine-1-carboxylate (3.7 g, 8.8 mmol, 1.4 equiv) in dioxane (30.0 mL) and H₂O (3.0 mL) was added Cs₂CO₃ (4.1 g, 12.5 mmol, 2.0 equiv) and Pd(dppf)Cl₂ (459 mg, 627 mol, 0.1 equiv). The mixture was stirred at 80° C. for 1 h under N₂. The resulting mixture was partitioned between EtOAc (50 mL) and H₂O (50 mL), the H₂O phase was separated, washed twice with EtOAc (50 mL), then the organic phase was separated, dried over Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=20/1 to 8/1). Compound tert-butyl 4-[4-[6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.7 g, 5.9 mmol, 94% yield, 99% purity) was obtained as a white solid. LC-MS: MS (ES⁺): RT=0.733 min, m/z=621.3 [M+H]⁺. LCMS method: 5-95.

[1174] Step 2: Pd/C (1.8 g, 1.7 mmol, 10% purity, 0.3 equiv) and Pd(OH)₂ (1.8 g, 2.6 mmol, 20% purity, 0.4 equiv) was added into a 250 mL single-necked round bottom flask under N₂, and then THE (30.0 mL) and MeOH (30.0 mL) was added at 25° C. under N₂. After addition, tert-butyl 4-[4-[6-benzyloxy-2-(2-fluorophenyl)-3,4-dihydronaphthalen-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (3.7 g, 6.0 mmol, 1.0 equiv) was added under N₂. The suspension was degassed under vacuum and purged with H₂ several times. The mixture was stirred under H₂ (50 psi) at 50° C. for 12 h. The reaction mixture was filtered and the filtrate was concentrated and then triturated with MeOH at 25° C. for 8 h. Compound (1S,2S)-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (2.4 g, 4.5 mmol, 76% yield, 99% purity) was obtained as a white solid. LC-MS: MS (ES⁺): RT=0.643 min, m/z=533.3 [M+H]⁺. LCMS Method: 5-95.

[1175] Step 3: Compound (±)-cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate (2.4 g, 4.5 mmol) was separated by SFC (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 um); mobile phase: [CO₂-EtOH(0.1% NH₃H₂O)]; B %:30%, isocratic elution mode) to afford (1S,2S)- and (1R,2R)-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate.

Each of the faster-eluting and slower-eluting enantiomers were isolated as white solids (710 mg, 1.3 mmol, 47% yield, 99% purity). LC-MS: MS (ES⁺): RT=0.641 min, m/z=533.3 [M+H]⁺; LCMS Method: 5-95.


[1176] The following Steps 4 and 5 were conducted on the slower-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate. This slower-eluting enantiomer produced the diastereomer of compounds III-345 and III-346 that was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The faster-eluting enantiomer of cis-tert-butyl 4-[4-[2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate was carried through the same step to afford the other one of compounds III-345 and III-346.


[1177] Step 4: To a solution of tert-butyl 4-[4-[(1S,2S)-2-(2-fluorophenyl)-6-hydroxy-tetralin-1-yl]-3-methoxy-phenyl]piperazine-1-carboxylate or its enantiomer (710 mg, 1.3 mmol, 1.0 equiv) in DCM (9.0 mL) was added TFA (3.0 mL). The mixture was stirred at 25° C. for 1 h. The reaction mixture was concentrated under reduced pressure to remove solvent. The resulting mixture was basified with saturated aqueous NaHCO₃ solution to adjust pH=8, and then diluted with DCM (30 mL) and extracted with DCM (30 mL×2). The combined organic layers were washed with DCM (20 mL), dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to give a residue. Compound (1S,2S)-2-(2-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (560 mg, 1.3 mmol, 97% yield) was obtained as a yellow solid and used in the next step without further purification. LC-MS: MS (ES⁺): RT=0.843 min, m/z=433.2 [M+H]⁺; LCMS method: 5-95.

[1178] Step 5: To a solution of (1S,2S)-2-(2-fluorophenyl)-1-(2-methoxy-4-piperazin-1-yl-phenyl)tetralin-6-ol or its enantiomer (120 mg, 277 mol, 1.3 equiv) and 6-[3-fluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2⁶]]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (95 mg, 210 mol, 1.0 equiv) in EtOH (3.0 mL) was added DIEA (82 mg, 631 mol, 110 μL, 3.0 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was concentrated under reduced pressure to remove solvent and purified by prep-HPLC (column: Waters xbridge 150*25 mm 10 um; mobile phase: [H₂O(10 mM

NH_4HCO_3)-ACN]; gradient: 63%-83% B over 8.0 min). Compound III-346 or III-345 (80 mg, 87 mol, 41% yield, 96% purity) was obtained as an off-white solid. LC-MS: MS (ES^+): $\text{RT}=2.626$ min, $m/z=884.6$ [$\text{M}+\text{H}$] $^+$; LCMS method: 01. ^1H NMR (400 MHz, CD_3OD) $\delta=7.37\text{--}7.29$ (m, 1H), 7.11-7.03 (m, 1H), 7.02-6.95 (m, 1H), 6.82-6.77 (m, 1H), 6.72-6.66 (m, 1H), 6.62-6.56 (m, 3H), 6.50-6.43 (m, 2H), 6.37-6.25 (m, 2H), 6.12 (d, $J=2.0$ Hz, 1H), 4.26-4.18 (m, 1H), 3.66-3.56 (m, 3H), 3.29-3.21 (m, 2H), 3.16-2.88 (m, 10H), 2.74-2.65 (m, 7H), 2.40-2.26 (m, 5H), 1.97 (d, $J=6.8$ Hz, 3H), 1.81-1.73 (m, 2H), 1.69-1.57 (m, 6H).

Example 53—Synthesis of 1-(2,5-difluoro-4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-((1S,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-339) and 1-(2,5-difluoro-4-((S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-((1R,2R)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-3-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-340)





[1179] Step 1: To a solution of 4-(triisopropylsilyloxymethyl)piperidin-4-ol (2 g, 10 mmol, 1 equiv), (9S)-7-(4-bromo-2,5-difluorophenyl)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaene (3 g, 6 mmol, 1 equiv) in dioxane (50 mL) was added Cs₂CO₃ (6 g, 20 mmol, 3 equiv) and Cphos Pd G₃ (553 mg, 686 mol, 0.1 equiv). The mixture was stirred at 90° C. for 3 h. The combined organic phase was filtered and concentrated under vacuum to give a residue. The residue was purified by column chromatography (SiO₂, Petroleum ether/Ethyl acetate=0/1) to give compound 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(triisopropylsilyloxymethyl)piperidin-4-ol (3 g, crude) as a white solid. LC-MS: MS (ES⁺): RT=0.733 min, m/z=644.4 [M-55]. LCMS Method: 5-95.

[1180] Step 2: To a solution of 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(triisopropylsilyloxymethyl)piperidin-4-ol (2 g, 4 mmol, 1 equiv) in THE (30 mL) was added TBAF (1 M, 6 mL, 1.5 equiv). The mixture was stirred at 25° C. for 0.5 h. The combined organic phase was filtered and concentrated under vacuum to give a residue. The residue was purified by column chromatography (SiO₂, Dichloromethane:Methanol=10/1) to give compound 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)pip-

eridin-4-ol (1 g, 2 mmol, 59% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.464 min, m/z=488.2 [M-100]. LCMS Method: 5-95.



[1181] Step 3: 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)piperidin-4-ol (1.3 g) was purified by prep-SFC to remove impurities, including any minor amounts of enantiomeric material (column: DAICEL CHIRALPAK AD (250 mm*30 mm, 10 μm); mobile phase: [CO₂-IPA:ACN=4:1 (0.1% NH₃H₂O)]; B %: 45%, isocratic elution mode) to give compound 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)piperidin-4-ol (950 mg, 1 mmol, 73% yield) as a white solid. LC-MS: MS (ES⁺): RT=0.481 min, m/z=488.1 [M+1]. LCMS Method: 5-95.

[1182] Step 4: To a solution of 1-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-4-(hydroxymethyl)piperidin-4-ol (400 mg, 820 μmol, 1 equiv), 1,1,2,2,3,3,4,4,4-nonafluorobutane-1-sulfonyl fluoride (371 mg, 1 mmol, 216 μL, 1 equiv) in THE (4 mL) was added DBU (249 mg, 1 mmol, 247 μL, 2 equiv). The mixture was stirred at 25° C. for 1 h. The combined organic layers were filtered and concentrated in vacuum and used directly. LC-MS: MS (ES⁺): RT=0.501 min, m/z=470.1 [M+1]. LCMS Method: 5-95.


[1183] Step 5: To a solution of 6-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (95 mg, 202 mol, 1 equiv) and the enantiomer of cis-1-(2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol prepared in Step 6 of Example 46 (83 mg, 202 mol, 1 equiv) in EtOH (2 mL) was added DIEA (78 mg, 606 mol, 105 μ L, 3 equiv). The mixture was stirred at 80° C. for 12 h. The combined organic layers were filtered and concentrated under reduced pressure to give a residue. The residue was purified by prep-HPLC (column: Waters xbridge 150*25 mm 10 μ m; mobile phase: [H₂O (10 mM NH₄HCO₃)-ACN]; gradient: 55%-85% B over 15.0 min) to give compound III-339 or III-340 (49 mg, 27% yield, 99% purity) as a white solid. ¹H NMR (400 MHz, MeOD) ¹H NMR (400 MHz, METHANOL-d₄) δ =7.30-7.16 (m, 1H), 7.12-7.00 (m, 3H), 6.82-6.70 (m, 3H), 6.68-6.60 (m, 2H), 6.58-6.44 (m, 2H), 6.42-6.32 (m, 1H), 6.18 (d, J=2.1 Hz, 1H), 4.76 (d, J=5.4 Hz, 1H), 4.32-4.20 (m, 1H), 3.42-3.34 (m, 2H), 3.28-3.12 (m, 3H), 3.11 (d, J=4.2 Hz, 4H), 3.02 (s, 5H), 2.82-2.72 (m, 4H), 2.68 (s, 3H), 2.48-2.36 (m, 5H), 2.36-2.24 (m, 1H), 2.00 (d, J=6.8 Hz, 3H), 1.92-1.80 (m,

2H), 1.78-1.64 (m, 6H). LC-MS: MS (ES⁺): RT=2.927 min, m/z=884.1 [M+1]. LCMS Method: 25. This diastereomer of compounds III-339 and III-340 was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The other diastereomer of compounds III-339 and III-340 was prepared from the other enantiomer of cis-1-(2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol.

Example 54—Synthesis of 1-(2,5-difluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-(2-fluoro-4-((1S,2S)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-5-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-341) and 1-(2,5-difluoro-4-((6S)-2,3,6,9-tetramethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepin-4-yl)phenyl)-4-((4-(2-fluoro-4-((1R,2R)-6-hydroxy-2-phenyl-1,2,3,4-tetrahydronaphthalen-1-yl)-5-methoxyphenyl)piperazin-1-yl)methyl)piperidin-4-ol (III-342)



-continued



III-342

[1184] Step 1: To a solution of the enantiomer of cis-1-(5-fluoro-2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol prepared in Step 6 of Example 45 (87 mg, 201 μ mol, 1 equiv) in EtOH (2 mL) was added DIEA (129 mg, 1.01 mmol, 175 μ L, 5 equiv) and 6-[2,5-difluoro-4-[(9S)-4,5,9,13-tetramethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0^{2,6}]trideca-2(6),4,7,10,12-pentaen-7-yl]phenyl]-1-oxa-6-azaspiro[2.5]octane (94 mg, 201 μ mol, 1 equiv). The mixture was stirred at 80° C. for 12 h. The reaction mixture was filtered. The residue was purified by prep-HPLC (neutral condition; column: Waters Xbridge C18 150*25 mm*5 μ m; mobile phase: [H₂O(10 mM NH₄HCO₃)-ACN]; gradient: 50%-80% B over 15.0 min) to give compound III-341 or III-342 (51 mg, 28% yield). ¹H NMR (400 MHz, CDCl₃): δ 7.34-7.27 (m, 1H), 7.16-7.03 (m, 3H), 6.85-6.73 (m, 3H), 6.70 (d, J=2.4 Hz, 1H), 6.62-6.48 (m, 2H), 6.36 (d, J=13.2 Hz, 1H), 6.05 (d, J=7.2 Hz, 1H), 5.70-5.18 (m, 1H), 4.80 (d, J=5.2 Hz, 1H), 4.17 (q, J=6.6 Hz, 1H), 3.41-3.23 (m, 4H), 3.22-3.05 (m, 5H), 3.04-2.99 (m, 2H), 2.97 (s, 3H), 2.89-2.75 (m, 2H), 2.68 (s, 3H), 2.54-2.41 (m, 1H), 2.39 (s, 3H), 2.30-2.20 (m, 1H), 2.09 (d, J=6.8 Hz, 3H), 1.77 (s, 3H), 1.75-1.56 (m, 8H). LC-MS: MS (ES⁺): RT=2.962 min, m/z=902.6 [M+H⁺]. LCMS method: 01. This diastereomer of compounds III-341 and III-342 was found to be the more potent inhibitor of cell growth of T-Rex 293 cells having increased expression of estrogen receptor alpha protein (as described in the Example 56, below). The other diastereomer of compounds III-339 and III-340 was prepared from the other enantiomer of cis-1-(5-fluoro-2-methoxy-4-piperazin-1-yl-phenyl)-2-phenyl-tetralin-6-ol.

Example 55—Synthesis of Additional Compounds in Tables 2, 3, 4, and 5

[1185] All compounds in Tables 2, 3, 4, and 5 that do not have a synthetic procedure described in the preceding examples were synthesized by analogy to the procedures described above. Observed mass data (m/z values) obtained from LC/MS of the compounds is provided in Table 6, below.

TABLE 6

Compound No.	Observed Mass (m/z)
I-34a	909.7
I-34b	909.7
I-37a or	802.5
I-37b	802.7
I-57	839.5

TABLE 6-continued

Compound No.	Observed Mass (m/z)
I-59	862.6
I-60	868.7
II-1	1234.6
II-2	1318.7
II-3	970.4
II-4	1230.7
II-5	1146.4
I-12,	869.7
III-1,	869.4
III-2, or	869.4
III-3	869.6
I-62 or	825.6
III-4	825.6
III-5 or	802.7
III-6	802.6
III-7 or	775.6
III-8	775.6
III-9 or	803.7
III-10	803.7
III-11 or	842.7
III-12	842.7
I-44 or	758.4
III-13	758.4
III-14 or	758.4
III-15	758.4
III-16 or	749.6
III-17	749.6
I-47 or	768.8
III-18	768.7
I-46,	835.6
III-19,	835.6
III-20, or	835.6
III-21	835.6
III-22 or	800.4
III-23	800.4
III-24 or	817.7
III-25	817.7
III-26 or	817.4
III-27	817.6
III-28	816.3
III-29	859.3
III-30	862.3
III-31	842.3
III-32	858.4
III-33	844.3
III-34	844.4
III-35	899.3
III-36	870.4
III-37	873.3
III-38	884.4
III-39	846.4

TABLE 6-continued

Compound No.	Observed Mass (m/z)
III-40	828.3
III-41	871.3
III-42	887.4
III-43	858.4
III-44	858.4
III-45	858.4
III-46	848.3
III-47	844.3
III-48	872.3
III-49	872.3
III-50	885.4
III-51	856.3
III-52	844.3
III-53	844.3
III-54	844.3
III-55	884.3
III-56	844.4
III-57	830.4
III-58	844.3
III-59	856.3
III-60	886.4
III-61	830.3
III-62	858.4
III-63	899.4
III-64	870.3
III-65	870.4
III-66	846.4
III-67 or	774.4
III-68	774.4
III-69 or	803.4
III-70	803.4
III-71,	869.4
III-72,	869.4
III-73, or	869.4
III-74	869.4
III-75 or	816.4
III-76	816.7
III-77a or b	843.4
III-78 or	789.7
III-79	789.7
III-80	788.7
III-81	846.7
III-82 or	801.7
III-83	801.7
III-84 or	789.6
III-85	789.6
III-86	817.8
III-87	842.6
III-88	842.7
III-89	774.6
III-90	760.5
III-91 or	870.7
III-92	870.7
III-93a or b	899.7
III-94a or b	803.5
III-95 or	789.7
III-96	789.7
III-97	408
III-98	773.5
III-99 or	761.5
III-100	381.3
III-101 or	775.6
III-102	775.6
III-103 or	859.7
III-104	859.6
III-105	899.7
III-106	803.7
III-107	759.7
III-108	407.9
III-109 or	860.6
III-110	860.7
III-111 or	845.7
III-112	845.7
III-113 or	871.7
III-114	871.6

TABLE 6-continued

Compound No.	Observed Mass (m/z)
III-115	787.7
III-116	773.6
III-117	759.5
III-118	818.6
III-119	816.5
III-120	759.5
III-121	832.7
III-122 or	804.5
III-123	804.5
III-124	841.6
III-125 or	870.8
III-126	870.7
III-127 or	871.6
III-128	871.6
III-129	787.6
III-130	831.6
III-131	842.8
III-132	801.7
III-133	843.7
III-134	857.6
III-135	857.7
III-136	844.8
III-137	801.6
III-138	774.6
III-139	788.6
III-140	885.6
III-141	774.3
III-142	788.6
III-143	886.6
III-144	819.6
III-145	816.6
III-146	844.6
III-147	844.7
III-148	773.4
III-149	884.6
III-150	827.6
III-151	860.6
III-152	817.6
III-153	817.6
III-154 or	884.6
III-155	884.6
III-156	855.6
III-157	861.4
III-158	790.4
III-159	790.3
III-160	858.6
III-161	909.6
III-162	910.6
III-163 or	885.4
III-164	885.4
III-165	858.4
III-166	859.4
III-167	857.4
III-168	925.7
III-169	925.6
III-170	788.4
III-171	858.6
III-172	857.4
III-173	857.4
III-174	855.4
III-175	924.5
III-176	924.5
III-177	859.6
III-178	857.4
III-179	925.5
III-180	926.5
III-181	927.7
III-182 or	854.6
III-183	854.6
III-184 or	837.6
III-185	836.6
III-186 or	854.6
III-187	854.6
III-188 or	846.6
III-189	846.6

TABLE 6-continued

Compound No.	Observed Mass (m/z)
III-190	857.7
III-191	856.6
III-192	921.3
III-193	921.6
III-194,	922.6
III-195,	922.7
III-196, or	922.7
III-197	922.6
III-198,	914.6
III-199,	914.7
III-200, or	914.6
III-201	914.6
III-202	924.8
III-203	861.7
III-204 or	883.6
III-205	883.6
III-206	858.7
III-207,	904.7
III-208,	904.6
III-209, or	904.7
III-210	904.7
III-211	925.7
III-212	861.7
III-213 or	950.6
III-214	949.7
III-215	869.6
III-216	869.6
III-217 or	821.6
III-218	821.6
III-219	815.5
III-220	815.6
III-221	815.7
III-222	859.4
III-223,	888.7
III-224,	888.7
III-225, or	888.7
III-226	888.7
III-227	927.7
III-228	829.7
III-229	842.6
III-230	843.8
III-231 or	886.6
III-232	886.6
III-233	842.7
III-234	841.6
III-235	859.6
III-236 or	873.6
III-237	873.6
III-238 or	877.5
III-239	877.4
III-240	831.6
III-241	831.6
III-242	814.4
III-243	814.4
III-244 or	871.4
III-245	871.5
III-246 or	855.7
III-247	854.3
III-248 or	854.3
III-249	855.3
III-250 or	855.3
III-251	854.5
III-252 or	861.1
III-253	861.1
III-254 or	867.5
III-255	867.2
III-256 or	873.3
III-257	873.3
III-258 or	922.4
III-259	922.3
III-260 or	814.5
III-261	813.6
III-262 or	833.6
III-263	833.6
III-264	847.6

TABLE 6-continued

Compound No.	Observed Mass (m/z)
III-265	878.4
III-266 or	911.7
III-267	911.6
III-268	831.5
III-269 or	873.6
III-270	873.6
III-271 or	873.6
III-272	873.6
III-273 or	873.5
III-274	873.6
III-275 or	866.4
III-276	866.4
III-277 or	851.5
III-278	851.6
III-279 or	885.6
III-280	885.6
III-281 or	851.5
III-282	851.6
III-283 or	851.6
III-284	851.6
III-285 or	851.6
III-286	851.7
III-287 or	851.7
III-288	851.6
III-289 or	855.7
III-290	855.6
III-291 or	855.8
III-292	855.6
III-293 or	872.7
III-294	873.5
III-295 or	869.6
III-296	869.5
III-297 or	851.6
III-298	851.5
III-299 or	922.6
III-300	922.6
III-301 or	871.6
III-302	872.4
III-303 or	873.5
III-304	873.4
III-305 or	870.8
III-306	870.8
III-307 or	867.7
III-308	867.7
III-309 or	867.9
III-310	867.7
III-311 or	867.8
III-312	867.8
III-313 or	868.7
III-314	869.7
III-315 or	851.8
III-316	851.7
III-317 or	839.7
III-318	839.7
III-319 or	885.6
III-320	885.7
III-321 or	848.5
III-322	849.5
III-323 or	839.7
III-324	839.8
III-325 or	839.7
III-326	839.7
III-327 or	885.8
III-328	884.7
III-329 or	846.9
III-330	846.8
III-331 or	864.5
III-332	865.8
III-333 or	885.7
III-334	885.8
III-335 or	882.8
III-336	883.7
III-337 or	890.7
III-338	890.7
III-339 or	885.6

TABLE 6-continued

Compound No.	Observed Mass (m/z)
III-340	885.7
III-341 or	903.7
III-342	903.7
III-343 or	885.7
III-344	885.7
III-345 or	885.7
III-346	885.7
III-347	828.3
III-348	917.7
III-349	917.8
III-350	919.8
III-351	919.7
III-352	885.6
III-353	885.6
III-354	935.7
III-355	935.7
III-356	885.8
III-357	885.7
III-358	902.7
III-359	902.7
IV-1	912.6
IV-2	928.6
IV-3	912.6
IV-4	977.6
IV-5	927.6
IV-6	911.6
IV-7	926.6
IV-8	951.6
IV-9	928.6
IV-10	884.6
IV-11	912.6
IV-12	911.5
IV-13	951.3
IV-14	979.4
IV-15	928.6
IV-16	859.6
IV-17	830.6
IV-18	893.6
IV-19	900.6
IV-20	865.6
IV-21	929.6
IV-22	855.6
IV-23	855.6
IV-24	882.6
IV-25	920.7
IV-26	894.5
IV-27	930.6
IV-28	877.6
IV-29	899.6
IV-30 or	995.4
IV-31	995.6
IV-32	882.6
IV-33	877.6
IV-34	912.5
IV-35	910.6
IV-36	855.6
IV-37	877.4
IV-38	902.4
IV-39	814.3
IV-40	850.2
IV-41	854.3
IV-42	870.3
IV-43	870.3
IV-44	882.3
IV-45	900.4
IV-46	899.2
IV-47	855.1
IV-48	896.3
IV-49	963.3
IV-50	826.6
IV-51	893.5
IV-52	928.5
IV-53	928.5
IV-54	917.6
IV-55	843.6

TABLE 6-continued

Compound No.	Observed Mass (m/z)
IV-56	916.5
IV-57	892.6
IV-58	906.6
IV-59	890.5
IV-60	978.5
IV-61	928.6
IV-62	853.6
IV-63	859.6
IV-64	932.6
IV-65	922.7
IV-66	888.4
IV-67	895.7
IV-68	925.7
IV-69	894.4
IV-70	920.6
IV-71	947.5
IV-72	909.5
V-1	961.4
V-2	945.4
V-3	933.4
V-4 or	885.6
V-5	885.6
V-6	831.6
V-7,	871.5
V-8,	871.5
V-9, or	871.6
V-10	871.6
V-11	871.4
V-12 or	869.4
V-13	869.4
V-14 or	885.6
V-15	885.6
V-16	872.6
V-17	872.6
V-18	870.6
V-19 or	911.6
V-20	911.7
V-21	843.6
V-22 or	870.4
V-23	870.4
V-24 or	871.4
V-25	871.4
V-26 or	886.4
V-27	886.4
V-28	887.4
V-29 or	886.4
V-30	886.3
V-31 or	886.5
V-32	886.6
V-33 or	953.4
V-34	953.4
V-35 or	911.6
V-36	911.6
V-37 or	921.7
V-38	921.7
V-39 or	921.8
V-40	921.7

Example 56—Cellular Growth Inhibition Assay Using T-Rex 293 Cells

[1186] Exemplary compounds were tested for the ability to inhibit the proliferation of the following types of cells: (i) a T-Rex 293 cell line having increased expression of estrogen receptor alpha protein from exposure to doxycycline and (ii) a T-Rex 293 cell line lacking increased expression of estrogen receptor alpha protein. Experimental procedures and results are provided below.

Part I—Experimental Procedure

[1187] The following types of cells were prepared for this experiment: (i) a T-Rex 293 cell line having increased

expression of estrogen receptor alpha protein from exposure to doxycycline and (ii) a T-Rex 293 cell line lacking increased expression of estrogen receptor alpha protein. The ability of the test compounds to inhibit proliferation of the foregoing cell types was evaluated according to the procedures set forth below.

[1188] The doxycycline-inducible estrogen receptor alpha protein expressing cell line was established using the following protocol: T-Rex 293 cells were purchased from Invitrogen (Cat #R71007) and transfected using Lipofectamine 2000 with the wild-type estrogen receptor alpha protein sequence cloned into the pcDNA4/TO vector. Transfected cells were selected using 400 µg/mL Zeocin (Invitrogen Cat #R25001). Following selection, single clones were raised and maintained in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% w/w Tetracycline-free fetal bovine serum (FBS) and 250 µg/mL Zeocin. Clones were analyzed for expression of estrogen receptor alpha protein in the presence and absence of 3 ng/mL doxycycline (Sigma Cat #D9891), and a single doxycycline-inducible clone (hereinafter "SC") was selected for use in downstream assays.

[1189] The SC cells were seeded on poly-D-lysine coated, black clear-bottom 384-well plates at 2500/well, in 25 µL Phenol Red Free Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% w/w charcoal-dextran treated fetal bovine serum (FBS) and 1% w/w pen-strep, with or without 3 ng/mL doxycycline. Pen-Strep is a commercially available mixture of penicillin G and streptomycin, which is used in mammalian cell culture media to prevent bacterial contamination. Phenol Red Free Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% w/w charcoal-dextran treated fetal bovine serum (FBS) and 1% w/w pen-strep, with or without 3 ng/mL doxycycline is herein referred to as Treatment Medium. Following seeding of cells in the plates, the plates were spun at 300×g for 30 seconds, then equilibrated to room temperature for 30 minutes, and then deposited in a humidified tissue culture incubator maintained at 37° C. with 5% CO₂.

[1190] At 24 hours after seeding of the cells, dilutions of test compound were prepared in DMSO and dissolved in Treatment Medium, to achieve a final DMSO concentration of 0.5% w/w, thereby providing the Test Compound Solution. A 25 µL aliquot of the Test Compound Solution was added to cells in the well plates. An equal volume of a solution containing DMSO and Treatment Medium was used as a negative control. Following treatment of cells with Test Compound Solution or said equal volume of a solution containing DMSO and Treatment Medium, the plates were spun at 300×g for 30 seconds, and then left in an incubator for 72 hours.

[1191] At the end of the treatment duration, cell viability was quantified with CellTiter-Glo 2.0 reagent (Promega Cat #G9243). For this purpose, plates were equilibrated to room temperature for 30 minutes, and then 25 µL of CellTiter-Glo 2.0 reagent was added to cells in the plate wells. Plates were then agitated on a shaker for two minutes at 500 rpm and subsequently incubated at room temperature for 10 minutes. Following incubation, the plates were spun at 3000×g for 30 seconds, then sealed with an optical adhesive cover, and luminescence readings were measured with an EnVision Plate Reader (Perkin Elmer).

[1192] Data was normalized using zero luminescence for baseline. A four-parameter non-linear regression curve fit

was applied to dose-response data in GraphPad Prism data analysis software to determine the half-maximal growth inhibitory concentration (GI₅₀) for each test compound.

Part II—Results

[1193] Results showing ability of exemplary compounds to inhibit the proliferation of T-Rex 293 cells having, or lacking, increased expression of estrogen receptor alpha protein are provided in Tables 7, 7-A, and 7-B, below, where columns labeled "(+) Dox TRex ERα" or "(+) Dox" refer to results in the experiment where compound was exposed to T-Rex 293 cells having increased expression of estrogen receptor alpha protein due to exposing the cells to doxycycline. Columns labeled "(−) Dox TRex ERα" or "(−) Dox" refer to results in the experiment where compound was exposed to T-Rex 293 cells lacking increased expression of estrogen receptor alpha protein. The symbol "++++" indicates a GI₅₀ less than 0.05 µM. The symbol "+++" indicates a GI₅₀ in the range of 0.05 µM to 0.5 µM. The symbol "++" indicates a GI₅₀ in the range of greater than 0.5 µM to 2.5 µM. The symbol "+" indicates a GI₅₀ greater than 2.5 µM. The symbol "N/A" indicates that no data was available.

[1194] Unless indicated otherwise, for compounds depicted in Tables 2-5 above having one or more stereogenic center(s) where stereochemistry is not defined using a dashed or wedged bond, the results reported in Tables 7, 7-A, and 7-B below are for a mixture of compounds that are diastereomers.

TABLE 7

Compound No.	(+) Dox TRex ERα	(−) Dox TRex ERα
I-1	+++	+++
I-2	+++	+++
I-3	+++	++
I-4	++++	++
I-5	++++	++
I-6	+++	+++
I-7	+++	+++
I-8	+++	+++
I-9	++++	+++
I-10	++++	++++
I-11	++++	++++
I-13	++++	+++
I-14	+++	+
I-15	++++	++++
I-16	+	+
I-17	++	++
I-18	++++	+++
I-19	++++	+++
I-20	++++	++++
I-21	++++	+++
I-22	++++	+++
I-23	++	++
I-24	+++	+++
I-25	+++	+++
I-26	++++	+++
I-27	++++	++
I-28	+++	++
I-29	++++	++++
I-30	++++	++++
I-31	++++	+++
I-32	++++	++
I-33	++++	+
I-35	++++	++++
I-36	++	++
I-39	+++	++

TABLE 7-A

Compound No.	(+) Dox	(-) Dox
I-34a	+++	++
I-34b	++++	++
I-37a or	+++	++
I-37b	++	++
I-57	++++	++
I-59	+++	++
I-60	+++	++
I-12,	++++	++
III-1,	+	+
III-2, or	+	+
III-3	+++	++
I-62 or	+++	++
III-4	++	+
III-5 or	++++	++
III-6	+++	++
III-7 or	+++	++
III-8	++	++
III-9 or	++++	+++
III-10	++++	+++
III-11 or	++++	+++
III-12	++++	++
I-44 or	+++	++
III-13	++	+
III-14 or	+++	++
III-15	++	++
III-16 or	+++	+++
III-17	++	++
I-47 or	+++	+
III-18	+	+
I-46,	++++	+
III-19,	+	+
III-20, or	+	+
III-21	++	+
III-22 or	++++	++
III-23	++++	+
III-24 or	++++	+++
III-25	++++	++
III-26 or	++++	+++
III-27	++++	+++
III-28	+++	+++
III-29	++++	+++
III-30	++++	+++
III-31	+	+++
III-32	++++	+++
III-33	++++	+++
III-34	++++	+++
III-35	++++	+++
III-36	++++	+++
III-37	++++	+++
III-38	++++	+++
III-39	++++	+++
III-40	+++	+++
III-41	++++	+++
III-42	++++	+++
III-43	++++	+++
III-44	+++	++
III-45	++++	+++
III-46	++++	++++
III-47	+++	+++
III-48	++++	+++
III-49	++++	+++
III-50	++++	+++
III-51	++++	+++
III-52	+++	+++
III-53	+++	+++
III-54	+++	+++
III-55	+++	++
III-56	+++	+++
III-57	+++	+++
III-58	++++	+++
III-59	++++	+++
III-60	+++	+++
III-61	++++	+++
III-62	++++	+++
III-63	++++	+++

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
III-64	+++	+++
III-65	++++	+++
III-66	+++	+++
III-67 or	++++	++++
III-68	+++	++
III-69 or	+++	+++
III-70	++++	+++
III-71,	+	+
III-72,	++	++
III-73, or	++++	++
III-74	++++	+++
III-75 or	+++	+++
III-76	+++	+++
III-77a or b	++++	+++
III-78 or	+++	+++
III-79	+++	+++
III-80	+++	++
III-81	+++	+
III-82 or	+++	++
III-83	+++	++
III-84 or	++++	++
III-85	+++	++
III-86	+++	++
III-87	++++	+++
III-88	++++	+++
III-89	++++	++
III-90	++	+
III-91 or	+++	++
III-92	+++	+++
III-93a or b	+++	+++
III-94a or b	++++	+++
III-95 or	++++	+++
III-96	+++	+++
III-97	++++	++
III-98	++++	++
III-99 or	+++	+++
III-100	+++	+++
III-101 or	++++	+++
III-102	+++	+++
III-103 or	++++	+++
III-104	++++	+++
III-105	++++	+++
III-106	++++	++
III-107	+++	++
III-108	++++	+++
III-109 or	++++	+++
III-110	++++	++++
III-111 or	++++	+++
III-112	+++	+++
III-113 or	++++	+++
III-114	++++	+++
III-115	+++	++
III-116	+	+
III-117	++	++
III-118	++++	++
III-119	+++	++
III-120	+	++
III-121	++	+
III-122 or	++	+
III-123	++++	++
III-124	+++	++
III-125 or	++++	+++
III-126	++++	+++
III-127 or	+++	++
III-128	++++	+++
III-129	+++	++
III-130	+++	++
III-131	++++	++
III-132	+++	++
III-133	++++	+++
III-134	++++	+++
III-135	++++	++
III-136	+	+
III-137	++	++
III-138	++	+

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
III-139	++	++
III-140	+++	++
III-141	+++	++
III-142	+++	+++
III-143	+	+
III-144	+++	+
III-145	++	++
III-146	++	++
III-147	+++	+++
III-148	++++	++
III-149	++++	+++
III-150	++++	++
III-151	++++	+
III-152	++++	++
III-153	++++	++
III-154 or	+	+
III-155	++++	+++
III-156	++++	++++
III-157	++++	++
III-158	++++	+++
III-159	++++	++++
III-160	++++	+++
III-161	++++	+++
III-162	++++	+++
III-163 or	+	+
III-164	++++	+++
III-165	++++	+++
III-166	++++	+++
III-167	++++	+++
III-168	++++	+++
III-169	++++	+++
III-170	++++	+++
III-171	++++	+++
III-172	++++	+++
III-173	++++	+++
III-174	++++	+++
III-175	++++	+++
III-176	++++	+++
III-177	++++	+++
III-178	++++	+++
III-179	++++	++++
III-180	++++	+++
III-181	++++	+++
III-182 or	++++	++
III-183	++++	++
III-184 or	++++	+++
III-185	++++	++
III-186 or	++++	++
III-187	++++	+++
III-188 or	++++	+++
III-189	+	+
III-190	++++	++
III-191	++++	+++
III-192	+	+
III-193	+++	++
III-194,	+	+
III-195,	+	+
III-196, or	++++	+++
III-197	++++	+++
III-198,	+	+
III-199,	+	+
III-200, or	++++	+++
III-201	+++	+++
III-202	++++	+++
III-203	++++	++
III-204 or	++++	++
III-205	+++	++
III-206	++++	++
III-207,	+	+
III-208,	+	+
III-209, or	++++	+++
III-210	++++	+++
III-211	++++	+++
III-212	++++	+++
III-213 or	+	+

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
III-214	++++	+++
III-215	++++	+
III-216	+	+
III-217 or	++++	+++
III-218	++++	+++
III-219	++++	+++
III-220	++++	+++
III-221	++++	+++
III-222	++++	+
III-223,	++	++
III-224,	+	+
III-225, or	++++	+++
III-226	++++	+++
III-227	++++	+
III-228	++++	++
III-229	++++	++
III-230	++++	+
III-231 or	+	+
III-232	++++	+
III-233	++++	++
III-234	++++	++
III-235	++++	++
III-236 or	++++	++
III-237	+++	++
III-238 or	+++	++
III-239	++++	+++
III-240	++++	+++
III-241	++++	+++
III-242	++++	++
III-243	++++	++
III-244 or	+	+
III-245	++++	++
III-246 or	++++	++
III-247	++++	+++
III-248 or	++++	++
III-249	+++	++
III-250 or	+++	+++
III-251	++++	++
III-252 or	+++	++
III-253	++++	++
III-254 or	++++	++
III-255	++++	++
III-256 or	++++	++
III-257	+++	++
III-258 or	+++	+++
III-259	++++	++
III-260 or	++++	++
III-261	++++	++
III-262 or	+++	++
III-263	++++	++
III-264	++++	++
III-265	+++	+
III-266 or	++++	++
III-267	+++	+
III-268	++++	++
III-269 or	+++	+++
III-270	++++	++
III-271 or	+++	+++
III-272	++++	++
III-273 or	+++	+++
III-274	++++	++
III-275 or	+++	+++
III-276	++++	++
III-277 or	++++	++
III-278	+++	++
III-279 or	++++	++
III-280	+++	++
III-281 or	+++	++
III-282	++++	+++
III-283 or	+++	++
III-284	+++	++
III-285 or	+++	+++
III-286	++++	++
III-287 or	++++	++
III-288	+++	++

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
III-289 or	++++	++
III-290	++++	+++
III-291 or	++++	++
III-292	+++	+++
III-293 or	++++	++
III-294	+++	+++
III-295 or	++	++
III-296	+++	++
III-297 or	+++	+++
III-298	+++	+++
III-299 or	++++	+++
III-300	++++	+++
III-301 or	++	++
III-302	+++	++
III-303 or	+++	+++
III-304	++++	++
III-305 or	++++	++
III-306	+++	++
III-307 or	++++	++
III-308	+++	+++
III-309 or	+++	+++
III-310	++++	++
III-311 or	++++	++
III-312	+++	+++
III-313 or	++++	++
III-314	+++	++
III-315 or	++	++
III-316	++++	++
III-317 or	++++	+++
III-318	+++	+++
III-319 or	++++	+++
III-320	++++	++
III-321 or	++++	++
III-322	+++	+++
III-323 or	++++	++
III-324	+++	+++
III-325 or	+++	++
III-326	++	++
III-327 or	+++	+++
III-328	++++	++
III-329 or	+++	+++
III-330	++++	+++
III-331 or	++	++
III-332	++++	+++
III-333 or	+++	++
III-334	++++	++
III-335 or	+++	++
III-336	++++	++
III-337 or	+++	++
III-338	++++	++
III-339 or	++++	+
III-340	+++	+++
III-341 or	++++	+
III-342	+++	+++
III-343 or	+++	+++
III-344	++++	++
III-345 or	++++	+++
III-346	++++	++
IV-1	++++	+
IV-2	+++	++
IV-3	++++	++
IV-4	+++	++
IV-5	++++	+
IV-6	+++	++
IV-7	+++	++
IV-8	++++	++
IV-9	++++	++
IV-10	+++	++
IV-11	++++	++
IV-12	+++	++
IV-13	++++	+++
IV-14	++++	++
IV-15	+	+
IV-16	++++	++
IV-17	++++	++

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
IV-18	++++	++
IV-19	++++	++
IV-20	++++	++
IV-21	++++	++
IV-22	++++	+++
IV-23	++++	+++
IV-24	++++	++
IV-25	++++	+++
IV-26	++++	++
IV-27	+++	++
IV-28	++++	++
IV-29	++++	+++
IV-30 or	++	++
IV-31	++++	+++
IV-32	++++	+++
IV-33	++++	++
IV-34	++++	+++
IV-35	++++	++
IV-36	++++	+++
IV-37	++++	++
IV-38	++++	++
IV-39	++++	+
IV-40	++++	+
IV-41	++++	+++
IV-42	++++	+++
IV-43	++++	+++
IV-44	++++	++
IV-45	++++	++++
IV-46	++++	++
IV-47	++++	++
IV-48	++++	++
IV-49	++++	++
IV-50	++++	++
IV-51	++++	++
IV-52	+++	++
IV-53	++++	++
IV-54	++++	+++
IV-55	++++	++
IV-56	++++	+++
IV-57	++++	++
IV-58	++++	++
IV-59	++++	++
IV-60	++++	+++
IV-61	++++	++
IV-62	++++	++
IV-63	++++	++
IV-64	+++	++
V-1	+++	+++
V-2	+++	++
V-3	+++	+++
V-4 or	++	++
V-5	+++	++
V-6	+++	++
V-7,	++++	+++
V-8,	++++	++
V-9, or	+	++
V-10	++	++
V-11	+++	++
V-12 or	++++	+++
V-13	+++	+++
V-14 or	+++	++
V-15	++++	+++
V-16	++	++
V-17	+++	+++
V-18	+++	+++
V-19 or	++++	++
V-20	++++	++
V-21	++++	+++
V-22 or	++++	++
V-23	++++	++
V-24 or	+	+
V-25	+	+
V-26 or	++++	+++
V-27	++++	++
V-28	++++	+++

TABLE 7-A-continued

Compound No.	(+) Dox	(-) Dox
V-29 or	+++	+++
V-30	+++	+++
V-31 or	++++	+++
V-32	++++	++
V-33 or	++++	+++
V-34	+++	+++
V-35 or	++++	+++
V-36	++++	++

TABLE 7-B

Compound No.	(+) Dox	(-) Dox
III-347	++++	++++
III-348	++	++
III-349	++++	++
III-350	+	+
III-351	++++	+
III-352	++++	++
III-353	+++	++
III-354	+++	++
III-355	++++	++
III-356	++++	++
III-357	+++	++
III-358	++++	++
III-359	+++	++
IV-65	++++	++
IV-66	++++	++
IV-67	++++	++
IV-68	++++	++
IV-69	++++	++
IV-70	++++	++
IV-71	++++	++
IV-72	++++	++
V-37 or	++++	++
V-38	+++	+++
V-39 or	+++	+++
V-40	++++	++

Example 57—Cellular Growth Inhibition Assay Using MCF7 Cells

[1195] Exemplary compounds were tested for the ability to inhibit the proliferation of MCF7 cells. Experimental procedures and results are provided below.

Part I—Experimental Procedure

[1196] MCF7 cells (ATCC #HTB-22 or ECACC #86012803) were cultured in RPMI 1640 medium supplemented with 10% heat-inactivated fetal bovine serum, 1% pen-strep, and 0.2 U/ml insulin. Cells were seeded in 25 μ L of RPMI 1640 medium supplemented with 10% heat-inactivated fetal bovine serum, 1% pen-strep, 0.2 U/ml insulin in poly-D-lysine treated, black clear-bottom 384-well plates at 225 cells/well for MCF7 cells purchased from ECACC or 1,500 cells/well for MCF7 cells purchased from ATCC. Following seeding, plates were spun at 300 $\times$ g for 30 seconds and cultured at 37° C. with 5% CO₂ in a humidified tissue culture incubator.

[1197] At 24 hours after seeding, compounds were titrated in 100% DMSO and diluted in RPMI 1640 medium supplemented with 10% heat-inactivate fetal bovine serum, 1% pen-strep, and 0.2 U/ml insulin. A 25 μ L aliquot of the compound/media mixture was added to cells, bringing the total volume in each well to 50 μ L. An equal volume of

DMSO without compound was used as a negative control. After treatment, plates were spun at 300 $\times$ g for 30 seconds, then cultured at 37° C. with 5% CO₂ for six days in a humidified tissue culture incubator.

[1198] On Day 6 of treatment, cell viability was quantified with CellTiter-Glo 2.0 reagent (Promega). Plates were equilibrated to room temperature for 30 minutes, then 25 μ L of CellTiter-Glo 2.0 reagent was added to cells, bringing the total volume in each well to 75 μ L. After reagent was added, plates were mixed on a shaker for two minutes at 500 rpm and then incubated at room temperature for 10 minutes. Following incubation, plates were spun at 3,000 $\times$ g for 30 seconds, sealed with an optical adhesive cover, and luminescence readings were measured with an EnVision Plate Reader.

[1199] Data was normalized to 0 luminescence for baseline. A four-parameter, non-linear regression curve fit was applied to dose-response data in GraphPad Prism data analysis software to determine the half maximal growth inhibitory concentration (GI₅₀) for each compound.

Part II—Results

[1200] Results showing ability of exemplary compounds to inhibit the proliferation of MCF7 cells purchased from ATCC are provided in Table 8-A, below. Results showing ability of exemplary compounds to inhibit the proliferation of MCF7 cells purchased from ECACC are provided in Table 8-B, below. The symbol “++++” indicates a GI₅₀ less than 0.05 μ M. The symbol “+++” indicates a GI₅₀ in the range of 0.05 μ M to 0.5 μ M. The symbol “++” indicates a GI₅₀ in the range of greater than 0.5 μ M to 2.5 μ M. The symbol “+” indicates a GI₅₀ greater than 2.5 M.

[1201] Unless indicated otherwise, for compounds depicted in Tables 2-5 above having one or more stereogenic center(s) where stereochemistry is not defined using a dashed or wedged bond, the results reported in Tables 8-A and 8-B below are for a mixture of compounds that are diastereomers.

TABLE 8-A

Compound No.	GI ₅₀
I-1	+++
I-2	+++
I-3	+++
I-4	+++
I-5	+++
I-6	+++
I-7	+++
I-8	++++
I-9	++++
I-10	++++
I-11	++++
I-13	++++
I-14	++
I-15	++++
I-16	+++
I-17	+++
I-19	++++
I-20	++++
I-21	++++
I-22	++++
I-23	+++
I-24	++++
I-25	++++
I-26	++++
I-27	+++
I-28	+++

TABLE 8-A-continued

Compound No.	GI ₅₀
I-29	++++
I-30	++++
I-31	++++
I-32	+++
I-33	++++
I-34a	+++
I-34b	+++
I-35	++++
I-36	++
I-37a or	+++
I-37b	++
I-39	+++
I-57	++++
I-59	+++
I-60	+++
II-1	++++
II-2	++++
II-3	++++
II-4	++++
II-5	++++
I-12,	++++
III-1,	++
III-2, or	++
III-3	+++
I-62 or	+
III-4	+
III-5 or	++++
III-6	+++
III-7 or	+++
III-8	+++
III-9 or	+++
III-10	+++
III-11 or	+++
III-12	+++
I-44 or	++
III-13	++
III-14 or	++
III-15	++
III-16 or	+++
III-17	++
I-47 or	++
III-18	++
I-46,	+++
III-19,	+
III-20, or	+
III-21	++
III-22 or	+++
III-23	+++
III-24 or	+++
III-25	++++
III-26 or	+++
III-27	+++
III-28	+++
III-29	+++
III-30	+++
III-31	+++
III-32	+++
III-33	+++
III-34	+++
III-35	++++
III-36	++++
III-37	++++
III-38	++++
III-39	++++
III-40	++++
III-42	+++
III-43	+++
III-44	+++
III-45	+++
III-46	++++
III-47	+++
III-48	++++
III-49	++++
III-50	++++
III-51	++++

TABLE 8-A-continued

Compound No.	GI ₅₀
III-52	+++
III-53	+++
III-54	+++
III-55	+++
III-56	+++
III-57	+++
III-58	+++
III-59	+++
III-60	+++
III-61	++++
III-62	+++
III-63	+++
III-64	+++
III-65	+++
III-66	+++
III-67 or	++++
III-68	+++
III-69 or	+++
III-70	++++
III-71,	+
III-72,	++
III-73, or	++++
III-74	+++
III-75 or	+++
III-76	+++
III-77a or b	++++
III-78 or	+++
III-79	+++
III-80	+
III-81	+++
III-82 or	+++
III-83	+++
III-84 or	+++
III-85	+++
III-86	+++
III-87	++++
III-88	++++
III-89	+++
III-90	+++
III-91 or	++++
III-92	+++
III-93a or b	++++
III-94a or b	+++
III-95 or	++++
III-96	+++
III-97	++++
III-98	+++
III-99 or	+++
III-100	+++
III-101 or	+++
III-102	+++
III-103 or	++++
III-104	++++
III-105	++++
III-106	+++
III-107	+++
III-108	++++
III-109 or	+++
III-110	++++
III-111 or	+++
III-112	+++
III-113 or	++++
III-114	+++
III-115	+++
III-116	++
III-117	+++
III-118	+++
III-119	+++
III-120	+++
III-121	++
III-122 or	++
III-123	+++
III-124	+++
III-125 or	+++
III-126	+++

TABLE 8-A-continued

Compound No.	GI ₅₀
III-127 or	+++
III-128	+++
III-129	+++
III-130	+++
III-131	+++
III-132	+++
III-133	+++
III-134	++++
III-135	++++
III-136	+
III-137	+++
III-138	++
III-139	+++
III-140	++++
III-141	+++
III-142	+++
III-143	+
III-144	++
III-145	+++
III-146	++++
III-147	+++
III-148	++++
III-149	++++
III-150	+++
III-151	++++
III-152	+++
III-153	+++
III-154 or	+++
III-155	++++
III-156	++++
III-157	+++
III-158	++++
III-159	++++
III-160	++++
III-161	++++
III-162	++++
III-163 or	+++
III-164	++++
III-165	++++
III-166	+++
III-167	++++
III-168	++++
III-169	+++
III-170	+++
III-171	+++
III-172	+++
III-173	++++
III-174	++++
III-175	++++
III-176	++++
III-177	+++
III-178	++++
III-179	++++
III-180	++++
III-181	++++
III-182 or	+++
III-183	+++
III-184 or	+++
III-185	++++
III-186 or	++++
III-187	+++
III-188 or	+++
III-189	+++
III-190	+++
III-191	+++
III-192	+
III-193	+++
III-194,	+
III-195,	++
III-196, or	++++
III-197	+++
III-198,	+
III-199,	+
III-200, or	++++
III-201	+++

TABLE 8-A-continued

Compound No.	GI ₅₀
III-202	++++
III-203	++++
III-204 or	+++
III-205	+++
III-206	++++
III-207,	++
III-208,	++
III-209, or	+++
III-210	++++
III-211	++++
III-212	++++
III-213 or	+
III-214	++++
III-215	++++
III-216	+++
III-217 or	+++
III-218	+++
III-219	+++
III-220	+++
III-221	++++
III-222	+++
III-223,	++
III-224,	+++
III-225, or	+++
III-226	++++
III-227	+++
III-228	+++
III-229	+++
III-230	+++
III-231 or	++
III-232	++++
III-233	+++
III-234	+++
III-235	++++
III-236 or	++++
III-237	+++
III-238 or	+++
III-239	++++
III-240	++++
III-241	++++
III-242	++++
III-243	++++
III-244 or	+++
III-245	++++
III-246 or	++++
III-247	+++
III-248 or	++++
III-249	+++
III-250 or	+++
III-251	++++
III-252 or	++
III-253	++++
III-254 or	+++
III-255	+++
III-256 or	++++
III-257	+++
III-258 or	+++
III-259	++++
III-260 or	+++
III-261	+++
III-262 or	+++
III-263	++++
III-264	++++
III-265	+++
III-266 or	++++
III-267	+++
III-268	++++
III-269 or	+++
III-270	+++
III-271 or	+++
III-272	++++
III-273 or	+++
III-274	++++
III-275 or	+++
III-276	++++

TABLE 8-A-continued

Compound No.	GI ₅₀
III-277 or	++++
III-278	+++
III-279 or	++++
III-280	+++
III-281 or	++
III-282	++++
III-283 or	+++
III-284	+++
III-285 or	+++
III-286	++++
III-287 or	+++
III-288	+++
III-289 or	++++
III-290	+++
III-291 or	++++
III-292	+++
III-293 or	++++
III-294	+++
III-295 or	+++
III-296	+++
III-297 or	+++
III-298	+++
III-299 or	++++
III-300	+++
III-301 or	+++
III-302	+++
III-303 or	+++
III-304	++++
III-305 or	++++
III-306	+++
III-307 or	++++
III-308	+++
III-309 or	+++
III-310	++++
III-311 or	++++
III-312	+++
III-317 or	+++
III-318	+++
III-339 or	++++
III-340	+++
III-341 or	++++
III-342	+++
III-343 or	++++
III-344	++++
III-345 or	++++
III-346	++++
III-347	++++
IV-1	+++
IV-2	+++
IV-3	+++
IV-4	++++
IV-5	+++
IV-6	+++
IV-7	+++
IV-8	++++
IV-9	++++
IV-10	+++
IV-11	+++
IV-12	+++
IV-13	++++
IV-14	++++
IV-15	++
IV-16	++++
IV-17	++++
IV-18	++++
IV-19	+++
IV-20	++++
IV-21	+++
IV-22	++++
IV-23	++++
IV-24	++++
IV-25	++++
IV-26	++++
IV-27	+++
IV-28	++++

TABLE 8-A-continued

Compound No.	GI ₅₀
IV-29	++++
IV-30 or	+++
IV-31	++++
IV-32	+++
IV-33	++++
IV-34	++++
IV-35	++++
IV-36	++++
IV-37	++++
IV-38	++++
IV-39	+++
IV-40	++++
IV-41	++++
IV-42	++++
IV-43	++++
IV-44	++++
IV-45	++++
IV-46	+++
IV-47	++++
IV-48	++++
IV-49	++++
IV-50	++++
IV-51	++++
IV-52	+++
IV-53	++++
IV-54	++++
IV-55	++++
IV-56	++++
IV-57	++++
IV-58	++++
IV-59	++++
IV-60	++++
IV-61	++++
IV-62	++++
IV-63	++++
IV-64	++++
IV-65	+++
V-1	+++
V-2	+++
V-3	+++
V-4 or	+++
V-5	+++
V-6	+++
V-7,	+++
V-8,	+++
V-9, or	+++
V-10	++
V-11	+++
V-12 or	+++
V-13	+++
V-14 or	+++
V-15	+++
V-16	+++
V-17	+++
V-18	+++
V-19 or	++++
V-20	+++
V-21	++++
V-22 or	+++
V-23	+++
V-24 or	+
V-25	+
V-26 or	+++
V-27	+++
V-28	++++
V-29 or	+++
V-30	+++
V-31 or	+++
V-32	+++
V-33 or	+++
V-34	+++
V-35 or	++++
V-36	+++
V-37 or	++++
V-38	+++

TABLE 8-A-continued

Compound No.	GI ₅₀
V-39 or	+++
V-40	++++

TABLE 8-B

Compound No.	GI ₅₀
III-244 or	+++
III-245	++++
III-250 or	+++
III-251	++++
III-275 or	++++
III-276	++++
III-307 or	++++
III-308	+++
III-311 or	++++
III-312	++++
III-313 or	+++
III-314	+++
III-315 or	+++
III-316	++++
III-317 or	++++
III-318	+++
III-319 or	+++
III-320	++++
III-321 or	++++
III-322	+++
III-323 or	++++
III-324	+++
III-325 or	+++
III-326	++
III-327 or	+++
III-328	++++
III-329 or	+++
III-330	++++
III-331 or	++
III-332	++++
III-333 or	+++
III-334	++++
III-335 or	+++
III-336	++++
III-337 or	+++
III-338	++++
III-339 or	++++
III-340	+++
III-341 or	++++
III-342	+++
III-343 or	+++
III-344	++++
III-345 or	+++
III-346	++++
III-348	++
III-349	++++
III-350	+++
III-351	++++
III-352	++++
III-353	+++
III-354	+++
III-355	++++
III-356	+++
III-357	+++
III-358	++++
III-359	+++
IV-66	++++
IV-67	++++
IV-68	++++
IV-69	++++
IV-70	++++
IV-71	++++
IV-72	++++

Example 58—Assay for Binding Affinity to BRD4-BD1

[1202] Exemplary compounds were tested for ability to bind to BRD4-BD 1. Experimental procedures and results are provided below.

Part I—Experimental Procedure

[1203] Compounds were tested using a bromoKdELECT assay. T7 phage strains displaying bromodomains were grown in parallel in 24-well blocks in an *E. coli* host derived from the BL21 strain. *E. coli* were grown to log-phase and infected with T7 phage from a frozen stock (multiplicity of infection=0.4) and incubated with shaking at 32° C. until lysis (90-150 minutes). The lysates were centrifuged (5,000×g) and filtered (0.2 m) to remove cell debris. Streptavidin-coated magnetic beads were treated with biotinylated small molecule or acetylated peptide ligands for 30 minutes at room temperature to generate affinity resins for bromodomain assays. The liganded beads were blocked with excess biotin and washed with blocking buffer (SeaBlock (Pierce), 1% BSA, 0.05% Tween 20, 1 mM DTT) to remove unbound ligand and to reduce nonspecific phage binding. Binding reactions were assembled by combining bromodomains, liganded affinity beads, and test compounds in 1× binding buffer (17% SeaBlock, 0.33× PBS, 0.04% Tween 20, 0.02% BSA, 0.004% Sodium azide, 7.4 mM DTT). Test compounds were prepared as 1000× stocks in 100% DMSO. Kds were determined using an 11-point 3-fold compound dilution series with one DMSO control point. All compounds for Kd measurements are distributed by acoustic transfer (non-contact dispensing) in 100% DMSO. The compounds were then diluted directly into the assays such that the final concentration of DMSO was 0.09%. All reactions performed in polypropylene 384-well plates. Each was a final volume of 0.02 ml. The assay plates were incubated at room temperature with shaking for 1 hour and the affinity beads were washed with wash buffer (1× PBS, 0.05% Tween 20). The beads were then resuspended in elution buffer (1× PBS, 0.05% Tween 20, 2 M non-biotinylated affinity ligand) and incubated at room temperature with shaking for 30 minutes. The bromodomain concentration in the eluates was measured by qPCR.

Part II—Results

[1204] Results showing ability of exemplary compounds to bind to BRD4-BD1 are provided in Table 9 below. The symbol “++++” indicates a K_d less than 0.05 μM. The symbol “+++” indicates a K_d in the range of 0.05 μM to 0.5 μM. The symbol “++” indicates a K_d in the range of greater than 0.5 μM to 2.5 μM. The symbol “+” indicates a K_d greater than 2.5 μM. The symbol “N/A” indicates that no data was available.

TABLE 9

Compound No.	K _d
I-2	+++
I-3	++++
I-5	++++

Example 59—Assay for Binding Affinity to BRD4-BD2

[1205] Exemplary compounds were tested for ability to bind to BRD4-BD2. Experimental procedures and results are provided below.

Part I—Experimental Procedure

[1206] Compounds were tested using a bromoKdELECT assay. T7 phage strains displaying bromodomains were grown in parallel in 24-well blocks in an *E. coli* host derived from the BL21 strain. *E. coli* were grown to log-phase and infected with T7 phage from a frozen stock (multiplicity of infection=0.4) and incubated with shaking at 32° C. until lysis (90-150 minutes). The lysates were centrifuged (5,000×g) and filtered (0.2 m) to remove cell debris. Streptavidin-coated magnetic beads were treated with biotinylated small molecule or acetylated peptide ligands for 30 minutes at room temperature to generate affinity resins for bromodomain assays. The liganded beads were blocked with excess biotin and washed with blocking buffer (SeaBlock (Pierce), 1% BSA, 0.05% Tween 20, 1 mM DTT) to remove unbound ligand and to reduce nonspecific phage binding. Binding reactions were assembled by combining bromodomains, liganded affinity beads, and test compounds in 1× binding buffer (17% SeaBlock, 0.33× PBS, 0.04% Tween 20, 0.02% BSA, 0.004% Sodium azide, 7.4 mM DTT). Test compounds were prepared as 1000× stocks in 100% DMSO. Kds were determined using an 11-point 3-fold compound dilution series with one DMSO control point. All compounds for Kd measurements are distributed by acoustic transfer (non-contact dispensing) in 100% DMSO. The compounds were then diluted directly into the assays such that the final concentration of DMSO was 0.09%. All reactions performed in polypropylene 384-well plates. Each was a final volume of 0.02 ml. The assay plates were incubated at room temperature with shaking for 1 hour and the affinity beads were washed with wash buffer (1× PBS, 0.05% Tween 20). The beads were then resuspended in elution buffer (1× PBS, 0.05% Tween 20, 2 M non-biotinylated affinity ligand) and incubated at room temperature with shaking for 30 minutes. The bromodomain concentration in the eluates was measured by qPCR.

Part II—Results

[1207] Results showing ability of exemplary compounds to bind to BRD4-BD2 are provided in Table 10 below. The symbol “++++” indicates a K_d less than 0.05 μ M. The symbol “+++” indicates a K_d in the range of 0.05 μ M to 0.5 μ M. The symbol “++” indicates a K_d in the range of greater than 0.5 μ M to 2.5 μ M. The symbol “+” indicates a K_d greater than 2.5 μ M. The symbol “N/A” indicates that no data was available.

TABLE 10

Compound No.	K_d
I-2	++++
I-3	++++

Example 60—Assay for Binding Affinity to CDK1

[1208] Exemplary compounds were tested for ability to bind to CDK1. Experimental procedures and results are provided below.

Part I—Experimental Procedure

[1209] Compounds were tested using a KdELECT assay. Kinase-tagged T7 phage strains were prepared in an *E. coli* host derived from the BL21 strain. *E. coli* were grown to log-phase and infected with T7 phage and incubated with shaking at 32° C. until lysis. The lysates were centrifuged and filtered to remove cell debris. The remaining kinases were produced in HEK-293 cells and subsequently tagged with DNA for qPCR detection. Streptavidin-coated magnetic beads were treated with biotinylated small molecule ligands for 30 minutes at room temperature to generate affinity resins for kinase assays. The liganded beads were blocked with excess biotin and washed with blocking buffer (SeaBlock (Pierce), 1% BSA, 0.05% Tween 20, 1 mM DTT) to remove unbound ligand and to reduce non-specific binding. Binding reactions were assembled by combining kinases, liganded affinity beads, and test compounds in 1× binding buffer (20% SeaBlock, 0.17× PBS, 0.05% Tween 20, 6 mM DTT). Test compounds were prepared as 11 IX stocks in 100% DMSO. Kd values were determined using an 11-point 3-fold compound dilution series with three DMSO control points. All compounds for Kd measurements are distributed by acoustic transfer (non-contact dispensing) in 100% DMSO. The compounds were then diluted directly into the assays such that the final concentration of DMSO was 0.9%. All reactions performed in polypropylene 384-well plate. Each was a final volume of 0.02 mL. The assay plates were incubated at room temperature with shaking for 1 hour and the affinity beads were washed with wash buffer (1× PBS, 0.05% Tween 20). The beads were then resuspended in elution buffer (1× PBS, 0.05% Tween 20, 0.5 μ M nonbiotinylated affinity ligand) and incubated at room temperature with shaking for 30 minutes. The kinase concentration in the eluates was measured by qPCR.

Part II—Results

[1210] Results showing ability of exemplary compounds to bind to CDK1 are provided in Table 11 below. The symbol “++++” indicates a K_d less than 0.05 μ M. The symbol “+++” indicates a K_d in the range of 0.05 μ M to 0.5 μ M. The symbol “++” indicates a K_d in the range of greater than 0.5 μ M to 2.5 μ M. The symbol “+” indicates a K_d greater than 2.5 μ M. The symbol “N/A” indicates that no data was available.

TABLE 11

Compound No.	K_d
I-40	+++
I-42	+++

INCORPORATION BY REFERENCE


[1211] The entire disclosure of each of the patent documents and scientific articles referred to herein is incorporated by reference for all purposes.

EQUIVALENTS

[1212] The invention may be embodied in other specific forms without departing from the spirit or essential characteristics thereof. The foregoing embodiments are therefore to be considered in all respects illustrative rather than limiting the invention described herein. Scope of the inven-

tion is thus indicated by the appended claims rather than by the foregoing description, and all changes that come within the meaning and range of equivalency of the claims are intended to be embraced therein.

1. A compound represented by Formula I-A:




or a pharmaceutically acceptable salt thereof, wherein: R^3 , R^{1C} , and R^{1E} are each independently for each occurrence hydrogen, fluoro, chloro, methoxy, or methyl; R^4 is methyl, $-\text{CH}_2\text{-(oxazolyl)}$, or $-\text{CH}_2\text{-(thiazolyl)}$; L is one of the following, wherein *** is the point of attachment to the phenyl ring bearing R^{1C} :

- (i) -(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-*** or
- (ii) -(7-11 membered saturated spirocyclic heterocyclyl containing 1 or 2 nitrogen atoms, wherein the heterocyclyl is substituted with 0 or 1 occurrence of hydroxyl or fluoro)-(C₁₋₄ alkylene)-(4-6 membered saturated monocyclic heterocyclyl containing 1 or 2 nitrogen atoms)-***; and

n , q , and r are independently 0, 1, or 2.

2. The compound of claim 1, wherein L is one of the following:




wherein: Z^{L1} is C(OH), C(H), C(F), or N;


Z^{L2} is independently for each occurrence N or C(H); and

*** is the point of attachment to the phenyl ring bearing R^{1C} .

3. The compound of claim 2, wherein R^4 is methyl, and L is



4. The compound of claim 1, wherein the compound is represented by Formula I-A-6, or a pharmaceutically acceptable salt thereof:






wherein Z^{L1} is C(OH) or C(H), and Z^{L2} is N or C(H).

5. The compound of claim 4, wherein the compound is a compound of Formula I-A-6.





6. The compound of claim 5, wherein R^{1C} and R^{1E} are each independently for each occurrence hydrogen, fluoro, or methoxy.





7. The compound of claim 6, wherein R^3 represents independently for each occurrence hydrogen or fluoro.


8. A compound selected from the following, or a pharmaceutically acceptable salt thereof:







-continued












-continued




-continued






9. The compound of claim 8, wherein the compound is




or a pharmaceutically acceptable salt thereof.

10. The compound of claim 8, wherein the compound is




11. The compound of claim 8, wherein the compound is




or a pharmaceutically acceptable salt thereof.

12. The compound of claim 8, wherein the compound is




or a pharmaceutically acceptable salt thereof.

13. A compound represented by



or a pharmaceutically acceptable salt thereof.

14. The compound of claim 13, wherein the compound is



15. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

16. A pharmaceutical composition comprising a compound of claim 9 and a pharmaceutically acceptable carrier.

17. A pharmaceutical composition comprising a compound of claim 10 and a pharmaceutically acceptable carrier.

18. A pharmaceutical composition comprising a compound of claim 13 and a pharmaceutically acceptable carrier.

19. A pharmaceutical composition comprising a compound of claim 14 and a pharmaceutically acceptable carrier.

20. A method of treating cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1 to treat the cancer, wherein the cancer is ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland carcinoma, bile duct cancer, gallbladder cancer, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia.

21. A method of treating cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 9 to treat the cancer, wherein the cancer is ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland carcinoma, bile duct cancer, gallbladder cancer, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia.

22. A method of treating cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 13 to treat the cancer, wherein the cancer is ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland

carcinoma, bile duct cancer, gallbladder cancer, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia.

23. A method of treating breast cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 1 to treat the breast cancer.

24. A method of treating breast cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 9 to treat the breast cancer.

25. A method of treating breast cancer, comprising administering to a patient in need thereof a therapeutically effective amount of a compound of claim 13 to treat the breast cancer.

26. The method of claim 24, wherein the patient is a human.

27. The method of claim 25, wherein the patient is a human.

28. A method of causing death of a cancer cell, comprising contacting a cancer cell with an effective amount of a compound of claim 1 to cause death of the cancer cell, wherein the cancer cell is selected from an ovarian cancer, uterine cancer, endometrial cancer, cervical cancer, prostate cancer, testicular cancer, breast cancer, brain cancer, lung cancer, oral cancer, esophageal cancer, head and neck cancer, stomach cancer, colon cancer, rectal cancer, skin cancer, sebaceous gland carcinoma, bile duct cancer, gallbladder cancer, liver cancer, pancreatic cancer, bladder cancer, urinary tract cancer, kidney cancer, eye cancer, thyroid cancer, lymphoma, or leukemia cell.

29. A method of causing death of a breast cancer cell, comprising contacting a breast cancer cell with an effective amount of a compound of claim 9 to cause death of the breast cancer cell.

30. A method of causing death of a breast cancer cell, comprising contacting a breast cancer cell with an effective amount of a compound of claim 13 to cause death of the breast cancer cell.

* * * * *