



US012398399B2

(12) **United States Patent**
De Both et al.

(10) **Patent No.: US 12,398,399 B2**

(45) **Date of Patent: Aug. 26, 2025**

(54) **SHOOT REGENERATION BY
OVEREXPRESSION OF CHK GENES**

(71) Applicant: **Keygene N.V.**, Wageningen (NL)

(72) Inventors: **Michiel Theodoor Jan De Both**,
Wageningen (NL); **Rik Hubertus
Martinus Op Den Camp**, Wageningen
(NL); **Bjorn Alexander Kloosterman**,
Wageningen (NL); **Bernarda Gerharda
Johanna Fierens-Onstenk**, Wageningen
(NL)

(73) Assignee: **Keygene N.V.**, Wageningen (NL)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 174 days.

(21) Appl. No.: **17/062,221**

(22) Filed: **Oct. 2, 2020**

(65) **Prior Publication Data**

US 2021/0079415 A1 Mar. 18, 2021

Related U.S. Application Data

(63) Continuation of application No.
PCT/EP2019/058614, filed on Apr. 5, 2019.

(30) **Foreign Application Priority Data**

Apr. 5, 2018 (EP) 18165895

(51) **Int. Cl.**
C12N 15/82 (2006.01)

(52) **U.S. Cl.**
CPC **C12N 15/8262** (2013.01); **C12N 15/8295**
(2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,026,530 B2 * 4/2006 Benfey C07K 14/415
536/23.6
9,127,073 B2 * 9/2015 Schmulling C07K 14/415
2006/0185030 A1 * 8/2006 Sheen C07K 14/415
800/278

FOREIGN PATENT DOCUMENTS

EP 2 272 862 A1 1/2011
WO WO-02/099079 A2 12/2002
WO WO-2006057832 A2 * 6/2006 C12N 15/8295

OTHER PUBLICATIONS

Kim et al 2006 (PNAS 103:3, p. 814-819) (Year: 2006).
Nishimura et al 2004 (The Plant Cell 16: p. 1365-77), "Histidine
Kinase Homologs That Act as Cytokinin Receptors Possess Over-

lapping Functions in the Regulation of Shoot and Root Growth in
Arabidopsis." (Year: 2004).*

Zuo et al 2002 (Current Opinions in Biotechnology 13: p. 173-80)
"Marker-free transformation: increasing transformation frequency
by the use of regeneration-promoting genes" (Year: 2002).*

Hwang et al 2001 (Nature 413: p. 383-389) (Year: 2001).*

Debnath 2005 (HortScience 40(1): p. 189-192) (Year: 2005).*

International Search Report issued in PCT/EP2019/058614 dated
May 24, 2019, 5 pages.

Riefler, Michael, et al., "Arabidopsis Cytokinin Receptor Mutants
Reveal Functions in Shoot Growth, Leaf Senescence, Seed Size,
Germination, Root Development, and Cytokinin Metabolism," The
Plant Cell, vol. 18(1), Jan. 2006, pp. 40-54, XP-002362525.

Banno et al., "Overexpression of *Arabidopsis* ESR1 Induces Initia-
tion of Shoot Regeneration", The Plant Cell, vol. 13, Dec. 2001, pp.
2609-2618 (10 pages).

Duclercq et al., "De novo shoot organogenesis: from art to science",
Trends in Plant Science, vol. 16, No. 11, Nov. 2011 (10 pages).

Mok et al., "Cytokinin Metabolism and Action", Annu. Rev. Plant.
Physiol. Plant Mol. Biol., vol. 52, 2001, pp. 89-118 (30 pages).

Quiroz-Figueroa et al., "Embryo production through somatic
embryogenesis can be used to study cell differentiation in plants",
Plant Cell Tiss Organ Cult, vol. 86, 2006, pp. 285-301 (17 pages).

Srinivasan et al., "Heterologous expression of the Baby Boom
AP2/ERF transcription factor enhances the regeneration capacity of
tobacco (*Nicotiana tabacum* L.)", Planta, vol. 225, 2007, pp.
341-351 (11 pages).

Hill et al., "Enhancing plant regeneration in tissue culture A
molecular approach through manipulation of cytokinin sensitivity",
Plant Signaling and Behavior, vol. 8(10), Oct. 2013, pp. 1-10.

* cited by examiner

Primary Examiner — Matthew R Keogh

(74) *Attorney, Agent, or Firm* — Sunit Talapatra; Foley &
Lardner LLP

(57) **ABSTRACT**

Genetic modification of plants is hampered by the limited
capacity of plant cells to regenerate. The current invention
solves this problem by introducing or increasing the expres-
sion of a histidine kinase in a plant cell. Preferred histidine
kinases are at least one of CHK2, CHK3 and CHK4. The
invention therefore concerns a method for improving a
cytokinin-induced regeneration capacity of a plant cell,
wherein the method comprises a step of increasing or
introducing the expression of a histidine kinase in the plant
cell. The invention further pertains to a method for regen-
erating a plant, wherein the method comprises a step of
introducing or increasing the expression of a histidine kinase
and to a plant obtainable from such method. Moreover, the
method concerns the use of at least one of CHK2, CHK3 and
CHK4 for improving a cytokinin-induced regeneration
capacity of a plant.

17 Claims, No Drawings

Specification includes a Sequence Listing.

1

SHOOT REGENERATION BY OVEREXPRESSION OF CHK GENES

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of International Application No. PCT/EP2019/058614 filed Apr. 5, 2019, which claims the benefit of and priority to European Application No. 18165895.6 filed Apr. 5, 2018, both of which are hereby incorporated by reference herein in their entireties.

SEQUENCE LISTING

The instant application contains a Sequence Listing which is being submitted in ASCII format via EFS-WEB and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Apr. 5, 2019, is named 085342-3800SequenceListing.txt and is 96 kb.

FIELD OF THE INVENTION

The present invention relates to the field of molecular plant biology, in particular to the field of plant regeneration. The invention concerns methods for improving the regeneration capacity of plant cells.

BACKGROUND

Plant breeding aims to improve the productivity and performance of crop plants through selection and recombination of useful and superior traits, and to improve such plant traits by genetic technologies. These technologies are often faced with a central technical challenge: how to regenerate from a single cell back to a fertile plant. This is true both for techniques aimed at making genetic changes (mutagenesis, genome editing, genetic transformation) and for those affecting the genome as a whole (DH production, polyploidisation, somatic hybridisations). This is because the genetic or genomic changes realised occur in only one or a small number of cells, and never in all cells of an organism at once in exactly the same manner. Therefore, an important limitation of these current techniques is the identification and singling out of the desired cells, and subsequently growing them back to complete and fertile plants.

Techniques of genome improvement are hampered by the potential of in vitro regeneration of modified single cells into whole fertile plants. Regeneration typically passes through a stage whereby a single cell first undergoes sustained cell division to form a multicellular structure or callus. Subsequently, under the influence of exogenously supplied plant growth regulators, cells in the multicellular mass form organized structures. Hence regeneration involves at least two sequential critical steps, i.e. a step of cell division followed by a step of differentiation to form organized structures. The first step one may involve callus formation. The second step must involve de novo meristem formation.

In vitro plant regeneration follows one of two alternative pathways, organogenesis and somatic embryogenesis, both of which rely on the induction by plant growth regulators (Duclercq et al., 2011, TIPS 16: 597). Monocotyledonous and dicotyledonous plants can use either both or only one of these two alternative pathways to regenerate.

A common pathway of plant regeneration in dicotyledonous plants is organogenesis, in which de novo apical meristems are formed from undifferentiated cells. These meristems usually grow out to form shoots, which are then

2

dissected from the underlying cell mass and induced to form roots. Organogenesis is typically induced in culture media containing cytokinins, or a mixture of auxins or cytokinins in which the cytokinins are often predominantly present. Cytokinins are a group of plant growth regulators or phytohormones, derivatives of adenine, and capable of promoting cell division (Mok and Mok, 2001, Annu. Rev. Plant Physiol. Plant Mol. Biol. 52: 89-118). The group encompasses naturally occurring cytokinins such as zeatin, and synthetic cytokinins such as kinetin and 6-benzylaminopurine (6-BAP).

Another regeneration pathway, often found in monocotyledonous and woody species, is through somatic embryogenesis, whereby undifferentiated cells in a callus, under proper conditions, generate embryogenic cells that ultimately form structures that resemble zygotic embryos, the so-called somatic embryos (Quiroz-Figueroa et al., 2006, PCTOC 86: 285-301). These somatic embryos are then capable of being converted into small plantlets, either spontaneous or under low concentrations of plant growth regulators. Embryogenic potential in callus or cell suspensions is typically induced by auxins, a group of naturally occurring or synthetic plant growth regulators.

A major bottleneck is the generally limited capacity of plant cells to regenerate into fertile and healthy plants. Regeneration potential is highly dependent on plant species, on variety and on tissue origin. Even with established protocols, the fraction of cells successfully regenerating to plants is usually quite low (Srinivasan et al., 2007, Planta 225: 341-351). Plant species or varieties in which the regeneration fails or the efficiency is poor are considered recalcitrant. Examples of crop species recalcitrant in organogenesis are pepper, soybean, cucumber and sugar beet.

In the past, recalcitrance has often been addressed by trial and error approaches through empirical variations in tissue culture conditions (media composition, light, temperature), and met with limited success. Protocols developed in this way are still dependent on genotype and laboratory conditions, and therefore to some extent unpredictable.

Regeneration through organogenesis or somatic embryogenesis can be enhanced by the ectopic expression of transgenes. An example of such approach is the overexpression of ESR1 (an AP2/EREBP transcription factor) in *Arabidopsis* (Banno et al., 2001, Plant Cell 13: 2609-2618), resulting in enhanced shoot regeneration from root explants. Another example is the overexpression of BBM, an AP2/ERF transcription factor, in tobacco (Srinivasan et al., 2007, Planta 225: 341-351), resulting in increased shoot formation and a higher competence for somatic embryogenesis. However, the constitutive expression of these transcription factors results in morphological and developmental defects, and these therefore require the controlled expression by exogenously supplied inducers (estradiol-inducible and dexamethasone-inducible system, respectively). The requirement of controlled expression thus currently limits the use of transgenes for regeneration.

Hence, there is still a need in the art for improving the generation capacity of a plant, especially for improving the generation capacity of a plant having a low or insufficient regeneration capacity. In particular, there is also a need in the art to improve the regeneration capacity of a plant, without requiring the controlled expression of transgenes.

SUMMARY OF THE INVENTION

In a first aspect, the invention pertains to a method for improving a cytokinin-induced regeneration capacity of a

3

plant cell, wherein the method comprises a step of increasing or introducing the expression of a histidine kinase in the plant cell and wherein the histidine kinase is at least one of CHK2, CHK3 and CHK4, wherein preferably the histidine kinase is at least one of CHK2 and CHK4, preferably wherein the histidine kinase is CHK4.

In one embodiment, the histidine kinase is encoded by a nucleotide sequence having at least 50% sequence identity with at least one of SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3, wherein preferably the nucleotide sequence has at least 50% sequence identity with SEQ ID NO: 3.

Preferably, the regeneration capacity of the plant cell is improved as compared to an identical plant cell not having an increased or introduced expression of the histidine kinase.

In an embodiment, the amino acid sequence of the histidine kinase has at least 50% sequence identity with at least one of SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6, wherein preferably the amino acid sequence of the histidine kinase has at least 50% sequence identity with SEQ ID NO: 6.

Preferably, the expression of the histidine kinase is transiently increased or introduced into the plant cell.

In one embodiment of the invention, the expression of the histidine kinase is continuously increased or introduced in the plant cell.

Preferably, the plant cell is obtainable from a plant selected from the group consisting of barley, cabbage, canola, cassava, cauliflower, chicory, cotton, cucumber, eggplant, grape, hot pepper, lettuce, maize, melon, oilseed rape, potato, pumpkin, rice, rye, *sorghum*, squash, sugar cane, sugar beet, sunflower, sweet pepper, tomato, water melon, wheat, zucchini, soybean, *chrysanthemum* and *Arabidopsis*.

In a further embodiment, the cytokinin is an adenine-type cytokinin, wherein preferably the adenine-type cytokinin is selected from the group consisting of kinetin, zeatin, trans-zeatin, cis-zeatin, dihydrozeatin, 6-benzylaminopurine and 2iP.

In a second aspect, the invention concerns a method for regenerating a plant comprising the steps of:

incubating the plant cell as defined herein in a medium comprising a cytokinin; and

allowing the plant cell to regenerate into a plant.

Preferably the medium comprises at least one further plant hormone, wherein preferably the one further plant hormone is an auxin.

In an embodiment, the plant cell is part of a multicellular tissue, preferably a callus tissue, a plant organ or an explant.

Preferably the explant is at least one of a hypocotyl explant, a stem explant, a cotyledon explant, a root explant, a leaf explant, a flower explant and a meristematic tissue.

In a further embodiment, the concentration cytokinin in the medium is about 100-3000 ng/ml, wherein preferably the concentration cytokinin is about 200-600 ng/ml.

In an embodiment of the invention, the improved cytokinin-induced regeneration capacity is selected from the group of an improved meristem formation, an improved adventitious shoot formation, an improved inflorescence formation, an improved somatic embryo formation, an improved root formation, an improved elongation of adventitious shoots and an improved regeneration of a complete plant.

In a third aspect, the invention relates to a plant or plant part obtainable by the method of the invention as defined herein, wherein preferably the plant part is a seed, a fruit or a non-propagating material.

4

In another aspect, the invention pertains to an expression construct comprising a first nucleotide sequence having at least 50% sequence identity with SEQ ID NO: 3 and a second nucleotide sequence having at least 50% sequence identity with SEQ ID NO: 1, wherein preferably at least one of the first and the second nucleotide sequence is operably linked to a regulatory element.

In a further aspect, the invention concerns the use of a CHK2, CHK3 and/or CHK4 histidine kinase or the expression construct as defined herein, for improving a cytokinin-induced regeneration capacity of a plant.

Definitions

Various terms relating to the methods, compositions, uses and other aspects of the present invention are used throughout the specification and claims. Such terms are to be given their ordinary meaning in the art to which the invention pertains, unless otherwise indicated. Other specifically defined terms are to be construed in a manner consistent with the definition provided herein.

It is clear for the skilled person that any methods and materials similar or equivalent to those described herein can be used for practising the present invention.

Methods of carrying out the conventional techniques used in methods of the invention will be evident to the skilled worker. The practice of conventional techniques in molecular biology, biochemistry, computational chemistry, cell culture, recombinant DNA, bioinformatics, genomics, sequencing and related fields are well-known to those of skill in the art and are discussed, for example, in the following literature references: Sambrook et al. *Molecular Cloning. A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N. Y., 1989; Ausubel et al. *Current Protocols in Molecular Biology*, John Wiley & Sons, New York, 1987 and periodic updates; and the series *Methods in Enzymology*, Academic Press, San Diego.

The singular terms "a," "an," and "the" include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to "a cell" includes a combination of two or more cells, and the like. The indefinite article "a" or "an" thus usually means "at least one".

The term "and/or" refers to a situation wherein one or more of the stated cases may occur, alone or in combination with at least one of the stated cases, up to with all of the stated cases.

As used herein, the term "about" is used to describe and account for small variations. For example, the term can refer to less than or equal to $\pm(+ \text{ or } -)$ 10%, such as less than or equal to $\pm 5\%$, less than or equal to $\pm 4\%$, less than or equal to $\pm 3\%$, less than or equal to $\pm 2\%$, less than or equal to $\pm 1\%$, less than or equal to $\pm 0.5\%$, less than or equal to $\pm 0.1\%$, or less than or equal to $\pm 0.05\%$. Additionally, amounts, ratios, and other numerical values are sometimes presented herein in a range format. It is to be understood that such range format is used for convenience and brevity and should be understood flexibly to include numerical values explicitly specified as limits of a range, but also to include all individual numerical values or sub-ranges encompassed within that range as if each numerical value and sub-range is explicitly specified. For example, a ratio in the range of about 1 to about 200 should be understood to include the explicitly recited limits of about 1 and about 200, but also to include individual ratios such as about 2, about 3, and about 4, and sub-ranges such as about 10 to about 50, about 20 to about 100, and so forth.

The term “comprising” is construed as being inclusive and open ended, and not exclusive. Specifically, the term and variations thereof mean the specified features, steps or components are included. These terms are not to be interpreted to exclude the presence of other features, steps or components.

The terms “protein” or “polypeptide” are used interchangeably and refer to molecules consisting of a chain of amino acids, without reference to a specific mode of action, size, 3 dimensional structure or origin. A “fragment” or “portion” of a protein may thus still be referred to as a “protein.” An “isolated protein” is used to refer to a protein which is no longer in its natural environment, for example in vitro or in a recombinant bacterial or plant host cell.

“Plant” refers to either the whole plant or to parts of a plant, such as cells, tissue or organs (e.g. pollen, seeds, gametes, roots, leaves, flowers, flower buds, anthers, fruit, etc.) obtainable from the plant, as well as derivatives of any of these and progeny derived from such a plant by selfing or crossing.

“Plant cell(s)” include protoplasts, gametes, suspension cultures, microspores, pollen grains, etc., either in isolation or within a tissue, organ or organism. The plant cell can e.g. be part of a multicellular structure, such as a callus, meristem, plant organ or an explant.

“Similar conditions” for culturing the plant/plant cells means among other things the use of a similar temperature, humidity, nutrition and light conditions, and similar irrigation and day/night rhythm.

The terms “homology”, “sequence identity” and the like are used interchangeably herein. Sequence identity is herein defined as a relationship between two or more amino acid (polypeptide or protein) sequences or two or more nucleotide (polynucleotide) sequences, as determined by comparing the sequences. In the art, “identity” also means the degree of sequence relatedness between amino acid or nucleic acid sequences, as the case may be, as determined by the match between strings of such sequences. “Similarity” between two amino acid sequences is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. “Identity” and “similarity” can be readily calculated by known methods. The percentage sequence identity/similarity can be determined over the full length of the sequence.

“Sequence identity” and “sequence similarity” can be determined by alignment of two peptide or two nucleotide sequences using global or local alignment algorithms, depending on the length of the two sequences. Sequences of similar lengths are preferably aligned using a global alignment algorithms (e.g. Needleman Wunsch) which aligns the sequences optimally over the entire length, while sequences of substantially different lengths are preferably aligned using a local alignment algorithm (e.g. Smith Waterman). Sequences may then be referred to as “substantially identical” or “essentially similar” when they (when optimally aligned by for example the programs GAP or BESTFIT using default parameters) share at least a certain minimal percentage of sequence identity (as defined below). GAP uses the Needleman and Wunsch global alignment algorithm to align two sequences over their entire length (full length), maximizing the number of matches and minimizing the number of gaps. A global alignment is suitably used to determine sequence identity when the two sequences have similar lengths. Generally, the GAP default parameters are used, with a gap creation penalty=50 (nucleotides)/8 (proteins) and gap extension penalty=3 (nucleotides)/2 (pro-

teins). For nucleotides the default scoring matrix used is nwsgapdna and for proteins the default scoring matrix is Blossum62 (Henikoff & Henikoff, 1992, PNAS 89, 915-919). Sequence alignments and scores for percentage sequence identity may be determined using computer programs, such as the GCG Wisconsin Package, Version 10.3, available from Accelrys Inc., 9685 Scranton Road, San Diego, CA 92121-3752 USA, or using open source software, such as the program “needle” (using the global Needleman Wunsch algorithm) or “water” (using the local Smith Waterman algorithm) in EmbossWIN version 2.10.0, using the same parameters as for GAP above, or using the default settings (both for ‘needle’ and for ‘water’ and both for protein and for DNA alignments, the default Gap opening penalty is 10.0 and the default gap extension penalty is 0.5; default scoring matrices are Blossum62 for proteins and DNATool for DNA). When sequences have a substantially different overall lengths, local alignments, such as those using the Smith Waterman algorithm, are preferred.

Alternatively percentage similarity or identity may be determined by searching against public databases, using algorithms such as FASTA, BLAST, etc. Thus, the nucleic acid and protein sequences of the present invention can further be used as a “query sequence” to perform a search against public databases to, for example, identify other family members or related sequences. Such searches can be performed using the BLASTn and BLASTx programs (version 2.0) of Altschul, et al. (1990) J. Mol. Biol. 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, wordlength=12 to obtain nucleotide sequences homologous to nucleic acid molecules of the invention. BLAST protein searches can be performed with the BLASTx program, score=50, wordlength=3 to obtain amino acid sequences homologous to protein molecules of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) Nucleic Acids Res. 25(17): 3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., BLASTx and BLASTn) can be used. See the homepage of the National Center for Biotechnology Information at <http://www.ncbi.nlm.nih.gov/>.

The terms “nucleic acid construct”, “nucleic acid vector”, “vector” and “expression construct” are used interchangeably herein and is herein defined as a man-made nucleic acid molecule resulting from the use of recombinant DNA technology. The terms “nucleic acid construct” and “nucleic acid vector” therefore does not include naturally occurring nucleic acid molecules although a nucleic acid construct may comprise (parts of) naturally occurring nucleic acid molecules.

The vector backbone may for example be a binary or superbinary vector (see e.g. U.S. Pat. No. 5,591,616, US 2002138879 and WO 95/06722), a co-integrate vector or a T-DNA vector, as known in the art and as described elsewhere herein, into which a chimeric gene is integrated or, if a suitable transcription regulatory sequence is already present, only a desired nucleic acid sequence (e.g. a coding sequence, an antisense or an inverted repeat sequence) is integrated downstream of the transcription regulatory sequence. Vectors can comprise further genetic elements to facilitate their use in molecular cloning, such as e.g. selectable markers, multiple cloning sites and the like.

The term “gene” means a DNA fragment comprising a region (transcribed region), which is transcribed into an RNA molecule (e.g. an mRNA) in a cell, operably linked to suitable regulatory regions (e.g. a promoter). A gene will

usually comprise several operably linked fragments, such as a promoter, a 5' leader sequence, a coding region and a 3' non-translated sequence (3' end) comprising a polyadenylation site.

"Expression of a gene" refers to the process wherein a DNA region which is operably linked to appropriate regulatory regions, particularly a promoter, is transcribed into an RNA, which is biologically active, i.e. which is capable of being translated into a biologically active protein or peptide.

The term "operably linked" refers to a linkage of polynucleotide elements in a functional relationship. A nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter, or rather a transcription regulatory sequence, is operably linked to a coding sequence if it affects the transcription of the coding sequence. Operably linked may mean that the DNA sequences being linked are contiguous.

"Promoter" refers to a nucleic acid fragment that functions to control the transcription of one or more nucleic acids. A promoter fragment is preferably located upstream (5') with respect to the direction of transcription of the transcription initiation site of the gene, and is structurally identified by the presence of a binding site for DNA-dependent RNA polymerase, transcription initiation site(s) and can further comprise any other DNA sequences, including, but not limited to transcription factor binding sites, repressor and activator protein binding sites, and any other sequences of nucleotides known to one of skill in the art to act directly or indirectly to regulate the amount of transcription from the promoter.

Optionally the term "promoter" may also include the 5' UTR region (5' Untranslated Region) (e.g. the promoter may herein include one or more parts upstream of the translation initiation codon of transcribed region, as this region may have a role in regulating transcription and/or translation). A "constitutive" promoter is a promoter that is active in most tissues under most physiological and developmental conditions. An "inducible" promoter is a promoter that is physiologically (e.g. by external application of certain compounds) or developmentally regulated. A "tissue specific" promoter is only active in specific types of tissues or cells.

The term "regeneration" is herein defined as the formation of a new tissue and/or a new organ from a single plant cell, a callus, an explant, a tissue or from an organ. A preferred regeneration pathway is organogenesis, i.e. the formation of new organs from (undifferentiated) cells. Preferably, the regeneration is at least one of ectopic apical meristem formation, shoot regeneration and root regeneration. The regeneration as defined herein can preferably concern at least de novo shoot formation. For example, regeneration can be the regeneration of a(n) (elongated) hypocotyl explant towards a(n) (inflorescence) shoot.

Regeneration may further include the formation of a new plant from a single plant cell or from e.g. a callus, an explant, a tissue or an organ. The regeneration process can occur directly from parental tissues or indirectly, e.g. via the formation of a callus.

The term "conditions that allow for regeneration" is herein understood as an environment wherein a plant cell or tissue can regenerate. Such conditions include at minimum a suitable temperature (i.e. between 0° C.-60° C.), nutrition, day/night rhythm, irrigation and the plant hormone cytokinin. These conditions can include one, two or three of the conditions specified in the examples. Furthermore, "optimal

conditions that allow for regeneration" are those environmental conditions that allow for a maximum regeneration of the plant cells.

The terms "introduced expression" and "de novo expression" are used interchangeably herein.

DETAILED DESCRIPTION OF THE INVENTION

The inventors discovered that introducing or enhancing the expression of a histidine kinase in a plant cell increases the regeneration capacity of this plant cell when grown in a medium comprising a cytokinin. The regeneration potential surprisingly increased in comparison to unmodified plant cells grown in the same cytokinin-comprising medium. Introducing a histidine kinase in a plant cell thus significantly augments the effect of the cytokinin. This enhanced effect of increasing the expression of a histidine kinase in combination with growing the cells on cytokinin-comprising medium surprisingly increased the maximum ability of a cell to regenerate. The inventors thus discovered that introducing a histidine kinase into a plant cell can transform a plant cell having a poor regeneration capacity into a plant cell that can regenerate, when grown in a cytokinin-comprising medium.

Without wishing to be bound to any theory, histidine kinase genes such as CHK genes encoding the cytokinin receptor molecules, seem to render plants more sensitive, and consequently more responsive to cytokinin, resulting in greatly improved shoot regeneration capacity. CHK genes previously have been implicated in the cytokinin signal transduction (Kakimoto, 1996, *Science* 274: 982-985), and enhanced expression of CHK genes have been associated with functions attributed to cytokinin response, but not regeneration capacity (Deng et al., 2010, *Plant Cell* 22: 1232-1248).

WO02/099079, EP2272862, and Riefler M et al, (*The Plant Cell*, Vol. 18 (2006): p 40-54) concern cytokinin response regulators and uses thereof. These documents are however silent on the combination of expressing a histidine kinase and maintaining the cells in cytokinin-comprising medium. In particular, the inventors now show that expressing a histidine kinase in cells in combination with maintaining the cells in cytokinin-comprising medium can push the cells above their natural maximum ability to regenerate.

In a first aspect, the invention therefore pertains to a method for improving a cytokinin-induced regeneration capacity of a plant cell, wherein the method comprises a step of increasing or introducing the expression of a histidine kinase in the plant cell.

The plant cell may be a protoplast. A cytokinin-induced regeneration is defined herein as the regeneration of a plant cell when exposed to at least a cytokinin, e.g. the regeneration response of a plant cell when exposed to a cytokinin under conditions that allow for the regeneration of the plant cell.

The cytokinin-induced regeneration capacity of a plant cell is herein understood as the maximum potential of a plant cell to regenerate when exposed to at least the plant hormone cytokinin under conditions, preferably optimal conditions, that allow for the regeneration of the plant cell. It is well-known in the art that plant cells have a certain inherent maximum potential to regenerate in a medium comprising a cytokinin. For example, the maximum regeneration potential of a plant cell obtained from e.g. a tobacco plant is usually higher than the maximum regeneration potential of a plant cell obtained from e.g. a cucumber plant. The regeneration capacity of plant cells may be assessed by

determining the number of de novo formed shoots on a multicellular tissue or tissues. As a non-limiting example upon induction of regeneration, tobacco multicellular tissues may provide for a higher number of de novo formed shoots per multicellular tissue as compared to the same number of cucumber multicellular tissues. A preferred multicellular tissue is a callus or explant.

In addition or alternatively, the maximum potential of a plant cell to regenerate can for example be measured by comparing the amount of starting material with the amount of material that is regenerated after exposure to cytokinin. As a non-limiting example, the number of multicellular tissues that form the starting material can be determined. Subsequently after exposure to at least a cytokinin under conditions, preferably optimal conditions that allow for regeneration, the number of multicellular tissues that are regenerated can be determined. The percentage of regenerated multicellular tissues is herein understood to be the cytokinin-induced regeneration capacity of the plant cell. As a non-limiting example, the starting material can be a hypocotyl explant, such as an elongated hypocotyl explant, and/or the regenerated tissue can be a shoot, such as an inflorescence shoot.

The cytokinin-induced regeneration may be at least one of meristem formation, adventitious shoot formation, inflorescence formation, somatic embryo formation, root formation, elongation of adventitious shoots and regeneration of a complete plant. As a non-limiting example, the number of de novo formed shoots per multicellular tissue may be determined.

The skilled person understands that there are other similar approaches known in the art to determine the cytokinin-induced regeneration capacity.

Preferably, the cytokinin-induced regeneration capacity of the plant cell is improved as compared to a control preferably as tested under the same, or substantially the same conditions, or optimal conditions. Preferably, said control is an identical plant cell or substantial identical plant cell that has the same genetic background as compared to the plant cell obtained or obtainable by the method of the invention, with the exception of the modified or newly introduced histidine kinase. In other words, preferably, the control is a plant cell that is genetically identical to the plant cell obtained by the method of the invention before the step of increasing or introducing the expression of a histidine kinase of the method of the invention. Thus, preferably the identical plant cell or substantially identical plant cell has the same genetic background as the plant cell having an improved regeneration capacity, but does not have an introduced or increased expression of the histidine kinase as defined herein.

The cytokinin-induced regeneration capacity of a plant cell is the maximum potential of the plant cell to regenerate under conditions, preferably optimal conditions, that allow for regeneration. The optimal conditions preferably includes at least an optimal concentration of a cytokinin.

The regeneration capacity can be expressed as the percentage of the starting material that regenerates. As a non-limiting example, if 7 of the 360 explants regenerate under conditions that allow for regeneration, the regeneration capacity is $(7/360 \times 100\%)$ 1.9%. Expression of a histidine kinase as defined herein allows for an improved regeneration capacity. As a non-limiting example, introduced or increased expression of a histidine kinase can increase the number of regenerated explants to e.g. 160 when grown under similar or substantially the same regeneration conditions. The regeneration capacity has increased to $(160/360 \times 100\%)$ to

44.4%. Hence, the regeneration capacity has improved $(44.4 - 1.9) / 42.5\%$ (see e.g. Table 2). As a further non-limiting example, the regeneration capacity of a plant cell can be expressed as the number of shoots formed per explant at a certain time point. For example 183 days after induced regeneration, 93 shoots can be harvested from 200 explants that have an introduced or increased expression of a histidine kinase (see also table 4), e.g. the regeneration capacity at day 183 is $(93/200 \times 100\%)$ 46.5%. At the same time point 64 shoots can be harvested from 200 control explants, e.g. the regeneration capacity at day 183 is $(64/200 \times 100\%)$ 32%. The regeneration capacity thus has improved $((46.5 - 32) / 32 \times 100\%)$ 45% (see e.g. Table 4).

As a non-limiting example, the improvement in cytokinin-induced regeneration capacity can be expressed as the difference in regeneration capacity between a plant cell having an increased or introduced expression of the histidine kinase as defined herein and an identical cell not having said increased or introduced expression. Preferably, the regeneration capacity of the plant cell is improved at least about 1%, 2%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or about 100% as compared to an identical plant cell not having an increased or introduced expression of the histidine kinase.

The skilled person understands that there are other suitable approaches in the art to determine an improvement in cytokinin-induced regeneration capacity, which can be equally used in the method of the invention.

A regenerated plant cell has preferably at least one of meristem formation or de novo meristem formation, shoot formation or adventitious shoot formation, inflorescence formation, somatic embryo formation, root formation, elongation of adventitious shoots and the regeneration of a complete plant.

The method can further comprise, for example, testing the plant cell for an improved regeneration capacity. The improved regeneration capacity is preferably improved compared to a control, wherein the control is preferably an identical plant cell not having an increased or introduced expression of a histidine kinase as defined herein.

As a non-limiting example, the testing step may be performed by determining the number of de novo formed shoots. Hence, the method of the invention may further comprise a step of determining, e.g. counting, the number of de novo formed shoots. The number of de novo formed shoots can be determined in relation to the number of multicellular tissues, preferably the number of explants, prior to shoot formation. Alternatively or in addition, the number of meristems may be determined or any tissues derived therefrom. It is however understood herein that the formation of callus or "green callus" is not indicative of the number of shoots that can be formed. The formation of callus can be an indicator of proliferation, but not an indicator of organisation/differentiation.

As a further non-limiting example, the testing step may be performed by determining the number of explants that regenerate. Hence, the method of the invention may further comprise a step of determining, e.g. counting, the number of explants that have regenerated. The number of regenerated explants can be determined in relation to the number of explants that formed the starting material.

A histidine kinase for use in the invention is a protein that is capable of transferring a phosphate group to a histidine residue on a specific substrate. Preferably, the histidine kinase for use in the invention is a transmembrane protein and can act as a cellular receptor for cytokinin. Upon

binding the cytokinin, the histidine kinase can initiate a signal transduction resulting in the activation of Type B ARRs (*Arabidopsis* response regulators) and subsequent cytokinin-regulated transcription.

The histidine kinase may be a CHASE-domain containing histidine kinase (CHK). These receptors preferably display a complex multidomain structure with a N-terminal part including preferably at least two hydrophobic membrane-spanning domains (TM) that border an extracytosolic sensing domain referred to as CHASE (Cyclase/Histidine kinase Associated Sensory Extracellular) as well as a cytoplasmic C-terminal part containing a catalytic histidine kinase (HK) domain and both receiver and pseudo-receiver domains (REC and REC-like, respectively). The HK domain is preferably composed of an HK dimerization and phosphoacceptor domain (HisKA) and an HK catalytic domain called the HK-like ATPase domain (HATPase) (Daudu et al, "CHASE-Containing Histidine Kinase Receptors in Apple Tree: From a Common Receptor Structure to Divergent Cytokinin Binding Properties and Specific Functions", Front Plant Sci. (2017); 8: 1614).

The histidine kinase for use in the invention can be a protein that is native to the plant cell, e.g. an endogenous histidine kinase protein that is expressed, or overexpressed, in the cell. Hence, the endogenous protein can be a protein that is encoded in the genome of a wild-type plant cell and its expression is introduced or enhanced. In addition or alternatively, additional copies of the endogenously encoded protein can be introduced into the plant cell.

In an embodiment, the histidine kinase for use in the invention is not native, i.e. is foreign, to the plant cell. Such exogenous protein may be a homologous protein. The protein can be derived from the same subgenus, genus, tribe, subfamily, family, order and/or clade. Alternatively, the protein can be derived from a different subgenus, genus, tribe, subfamily, family, order and/or clade. The histidine kinase protein can be an artificial protein, e.g. a protein that does not occur in nature but fulfils the same or similar function as a naturally-occurring histidine kinase.

In a preferred embodiment, the histidine kinase is at least one of CHK2, CHK3 and CHK4. Preferably the histidine kinase is at least one of CHK2 and CHK4. In an embodiment, the histidine kinase is CHK2. In another embodiment, the histidine kinase is CHK4. It is envisioned herein that the expression of a combination of histidine kinases is introduced or increased in the plant cell. For example, the expression of at least two different CHK2 proteins, at least two different CHK3 and/or at least two different CHK4 proteins is increased or introduced in the plant cell. Alternatively or in addition, the plant cell may have an increased or introduced expression of at least a CHK2 and CHK4 protein, at least a CHK2 and CHK3 protein, or at least a CHK3 and CHK4 protein. The plant cell can have an increased or introduced expression of at least a CHK2, CHK3 and a CHK4 protein as defined herein.

The histidine kinase of the invention may be encoded by a nucleotide sequence having at least 50% sequence identity with at least one of SEQ ID NO: 1, SEQ ID NO:2 and SEQ ID NO: 3. Preferably, the nucleotide sequence encoding the CHK2 can have at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 1. SEQ ID NO: 1 is the *Arabidopsis* CHK2 coding sequence, also annotated as AHK2.

Preferably, the nucleotide sequence encoding the CHK3 can have at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity

with SEQ ID NO: 2. SEQ ID NO: 2 is the *Arabidopsis* CHK3 coding sequence, also annotated as AHK3.

Preferably, the nucleotide sequence encoding the CHK4 preferably has at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 3. SEQ ID NO: 3 is the *Arabidopsis* CHK4 coding sequence, also annotated as AHK4, CRE1, WOL1, WOODEN LEG 1.

In one embodiment, the nucleotide sequence encoding the CHK2 protein is, or is derived from the gene At5g35750 (SEQ ID NO: 7), a homolog thereof or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with At5g35750 or its homolog. The percentage identity can be determined over the full length of the genomic sequence. Alternatively the percentage identity can be determined over the full length of the coding sequence of the gene.

Examples of homologs include *Glycine max* (Soybean) (GLYMA02G47611 or GLYMA14G01040), *Oryza sativa* (Rice) (0510G0362300, OSJNBA0058E19.1 or OSJNBA0073L01.1), *Populus trichocarpa* (Black Cottonwood) (POPTR_0014516260 G), *Solanum lycopersicum* (Tomato) (SOLYC07G047770.2) and *Vitis vinifera* (Grape) (VIT_1250057 G00690). A preferred homolog is *Solanum lycopersicum* SOLYC07G047770.2.

Preferably, the nucleotide sequence encoding the CHK2 protein is, or is derived from the gene SOLYC07G047770.2 (SEQ ID NO: 30), a homolog thereof and/or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SOLYC07G047770.2. The percentage identity can be determined over the full length of the genomic sequence. Alternatively the percentage identity can be determined over the full length of the coding sequence of the gene.

In one embodiment, the sequence encoding the CHK3 protein is, or is derived from the gene At1g27320 (SEQ ID NO: 8), a homolog thereof or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with At1g27320 or its homolog. The percentage identity can be determined over the full length of the genomic sequence. Alternatively the percentage identity can be determined over the full length of the coding sequence of the gene.

Examples of homologs include *Brachypodium distachyon* (Purple false brome) (BRAD12G59127), *Glycine max* (Soybean) (GLYMA05G28070 or GLYMA08G11060), *Oryza sativa* (Rice) (B1455F06.33), *Populus trichocarpa* (Black Cottonwood) (HK3A or POPTR_0003516950 G), *Solanum lycopersicum* (Tomato) (SOLYC05G015610.2) and *Vitis vinifera* (Grape) (VIT_0150010 G03780).

In one embodiment, the sequence encoding the CHK4 protein is, or is derived from, the gene At2g01830 (SEQ ID NO: 9), a homolog thereof, or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with At2g01830 or its homolog. The percentage identity can be determined over the full length of the genomic sequence. Alternatively the percentage identity can be determined over the full length of the coding sequence of the gene.

Examples of homologs include *Brachypodium distachyon* (Purple false brome) (BRAD11G10660), *Glycine max* (Soybean) (GLYMA02G09550, GLYMA05G34310, GLYMA07G19620, GLYMA07G27540 or GLYMA08G05370), *Physcomitrella patens* (Moss) (CKI3A, CKI3B, CKI3C, CRE1, CRE2, CRE3, PHY-PADRAFT_162473, PHYPADRAFT_169293, PHY-PADRAFT_172225, PHYPADRAFT_229095 or PHY-

13

PADRAFT_68053), *Populus trichocarpa* (Black Cottonwood) (CRE1B, POPTR_0008513720 G), *Solanum lycopersicum* (Tomato) (SOLYC04G008110.2), *Sorghum bicolor* (Sorghum) (SB01G010070) and *Vitis vinifera* (Grape) (VIT_01S0011G06190)

The sequence encoding the histidine kinase may be codon-optimized for expression in plant cells, preferably codon-optimized for expression in the plant cell of the method of the invention. As a non-limiting example, the expressed or de novo expressed histidine kinase can be an endogenous protein while the sequence encoding this endogenous protein is an exogenous, codon-optimized, sequence. Alternatively, the codon-optimized sequence can encode a histidine kinase that is exogenous for the plant cell.

In one embodiment, the amino acid sequence of the histidine kinase has at least 50% sequence identity with at least one of SEQ ID NO: 4, SEQ ID NO: 5 and SEQ ID NO: 6.

The amino acid sequence encoding the CHK2 can have at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 4. SEQ ID NO: 4 is the *Arabidopsis* CHK2 protein.

Alternatively, the nucleotide sequence encoding the CHK3 can have at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 5. SEQ ID NO: 5 is the *Arabidopsis* CHK3 protein.

Preferably, the nucleotide sequence encoding the CHK4 preferably has at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SEQ ID NO: 6. SEQ ID NO: 6 is the *Arabidopsis* CHK4 protein.

In one embodiment, the CHK2 amino acid sequence is or is derived from AT5G35750.1 (AHK2), a homolog thereof, or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with AHK2 or its homolog. A preferred homolog is *Solanum lycopersicum* CHK2 (SlyCHK2) having a sequence of SEQ ID NO: 31, and/or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with SlyCHK2.

Examples of homologs include *Glycine max* (Soybean) (K7KBR1 or K7M476), *Oryza sativa* (Rice) (Q0IY65, Q9AUQ0 or Q8S6P5), *Populus trichocarpa* (Black Cottonwood) (B9IAR0), *Solanum lycopersicum* (Tomato) (K4CEY3) and *Vitis vinifera* (Grape) (F6HHM7).

In one embodiment, the CHK3 amino acid sequence is or is derived from AT1G27320 (AHK3), a homolog thereof, or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with AHK3 or its homolog.

Examples of homologs include *Brachypodium distachyon* (Purple false brome) (I1HUP8), *Glycine max* (Soybean) (I1K3M7 or I1KS30), *Oryza sativa* (Rice) (Q5JJP1), *Populus trichocarpa* (Black Cottonwood) (B9GML7 or B9GZP2), *Solanum lycopersicum* (Tomato) (K4BYS7) and *Vitis vinifera* (Grape) (D7IAZ7).

In one embodiment, the CHK4 amino acid sequence is or is derived from AT2G01830 (AHK4), a homolog thereof, or a sequence having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98%, 99% or 100% sequence identity with AHK4 or its homolog.

Examples of homologs include *Brachypodium distachyon* (Purple false brome) (I1GNZ3), *Glycine max* (Soybean) (K7K767, K7KRH0, K7L210, K7L2C5 or I1KQE9), *Physcomitrella patens* (Moss) (A9RME1, A9SJM1, A9T3T9,

14

A9S5U9, A9TAF3, A9TKN3, A9S2L4, A9TCH3, A9TVM0, A9SEU1 or A9RME0), *Populus trichocarpa* (Black Cottonwood) (B9HVS3 or B9HJJ3), *Solanum lycopersicum* (Tomato) (K4BNW7), *Sorghum bicolor* (Sorghum) (C5WN04) and *Vitis vinifera* (Grape) (F6HFB2)

In an embodiment of the invention, the histidine kinase having increased or introduced expression is a functional histidine kinase. A functional histidine kinase is preferably fulfilling the same or similar function in a plant cell as the function of a histidine kinase having amino acid sequence SEQ ID NO: 4, SEQ ID NO: 5 or SEQ ID NO: 6 in *Arabidopsis thaliana*. A CHK2 as defined herein is preferably fulfilling the same or similar function in a plant cell as the function of a protein having amino acid sequence of SEQ ID NO: 4 in *Arabidopsis thaliana*. A CHK3 as defined herein is preferably fulfilling the same or similar function in a plant cell as the function of a protein having amino acid sequence of SEQ ID NO: 5 in *Arabidopsis thaliana*. A CHK4 as defined herein is preferably fulfilling the same or similar function in a plant cell as the function of a protein having amino acid sequence of SEQ ID NO: 6 in *Arabidopsis thaliana*. In context of the invention, a functional histidine kinase, or CHASE-domain containing histidine kinase, is preferably capable of binding cytokinin and initiates a signal transduction that results in cytokinin-regulated transcription. Alternatively or in addition, the functional histidine kinase can also be defined as a histidine kinase that is capable of improving the cytokinin-induced regeneration capacity of a plant cell upon increased or introduced expression.

The method can comprise, for example, a step of genetically engineering the plant, plant protoplast or plant cell to overexpress or express de novo a histidine kinase protein. In one embodiment, the expression of the histidine kinase is transiently increased or introduced into the plant cell. It is well-known in the art how to transiently increase or introduce the expression of a protein and the invention is not limited to any specific method.

As a non-limiting example, the method can comprise transforming a plant cell with a vector or expression construct comprising a recombinant nucleic acid encoding the histidine kinase as described herein, preferably as an expression construct of the ninth aspect as defined herein. Such vector or expression construct preferably does not integrate into the plant genome. As a result, the histidine kinase protein transcribed from the expression construct will be temporarily expressed. Introduction of a nucleic acid encoding the histidine kinase can be performed using any suitable means known in the art, for example such as described in WO2009/082190. In some embodiments, the method comprises contacting the plant protoplast or plant cell with an *Agrobacterium* strain comprising the vector to introduce the recombinant nucleic acid into the plant protoplast or plant cell.

Alternatively or in addition, the plant cells can be transformed with a histidine kinase protein as defined herein. The introduction of the histidine kinase protein can be performed by any suitable means known to the skilled person.

In one embodiment, the genome of the plant cell can be modified to transiently express or overexpress the histidine kinase protein as defined herein. As a non-limiting example, an expression cassette can be introduced into the genome of a plant cell, wherein the expression cassette at least comprises an inducible promoter and a sequence encoding the histidine kinase. The histidine kinase can for example be expressed upon the presence of an inducer. The inducer can bind to a transactivator, e.g. an introduced transactivator, which transactivator initiates or augments the expression of

the histidine kinase, for example by binding to the inducible promoter that is operably linked to the sequence encoding the histidine kinase.

In an embodiment, the expression of the histidine kinase is continuously increased or introduced in the plant cell.

Continuous overexpression or continuous de novo expression of the histidine kinase protein can be achieved by, for example, inserting at least one additional copy of an endogenous gene encoding the histidine kinase protein into the genome of a plant, plant protoplast or plant cell. Further ways are modulating promoter and/or further regulating sequences that are operably linked to an endogenous histidine kinase-encoding sequence, resulting in enhanced or introduced expression. These regulating sequences can include genomic as well as epigenomic regulators.

In an embodiment, the genome of the plant is modified to overexpress or de novo express a histidine kinase as defined herein by modifying a genomic promoter fragment having at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or 100% sequence identity with at least one of SEQ ID NO: 10, SEQ ID NO: 11 and SEQ ID NO: 12.

In one embodiment, the genome of the plant cell is modified to overexpress or de novo express a histidine kinase as defined herein. As a non-limiting example, regulatory sequences surrounding a gene encoding an endogenous histidine kinase as defined herein can be modified to increase or induce the expression of the histidine kinase. These regulatory sequences can be located less than about 10 kb, 8 kb, 5 kb, 3 kb, 2 kb, 1 kb, 800 bp, 600 bp, 400 bp, 200 bp, 100 bp or less than about 50 bp upstream from the start codon encoding the histidine kinase. Alternatively, these regulatory sequences can be located less than about 10 kb, 8 kb, 5 kb, 3 kb, 2 kb, 1 kb, 800 bp, 600 bp, 400 bp, 200 bp, 100 bp or less than about 50 bp downstream from the start codon encoding the histidine kinase.

In an embodiment, the regulatory sequences modified to increase or induce the expression of an endogenous histidine kinase as defined herein can be located less than about 10 kb, 8 kb, 5 kb, 3 kb, 2 kb, 1 kb, 800 bp, 600 bp, 400 bp, 200 bp, 100 bp or less than about 50 bp upstream of a sequence having at least about 50%, 60%, 70%, 80%, 90%, 95%, 98% or 100% sequence identity with SEQ ID NO: 7.

In an embodiment, the regulatory sequences modified to increase or induce the expression of an endogenous histidine kinase as defined herein can be located less than about 10 kb, 8 kb, 5 kb, 3 kb, 2 kb, 1 kb, 800 bp, 600 bp, 400 bp, 200 bp, 100 bp or less than about 50 bp upstream of a sequence having a sequence with at least about 50%, 60%, 70%, 80%, 90%, 95%, 98% or 100% sequence identity with SEQ ID NO: 8.

In an embodiment, the regulatory sequences modified to increase or induce the expression of an endogenous histidine kinase as defined herein can be located less than about 10 kb, 8 kb, 5 kb, 3 kb, 2 kb, 1 kb, 800 bp, 600 bp, 400 bp, 200 bp, 100 bp or less than about 50 bp upstream of a sequence having a sequence with at least about 50%, 60%, 70%, 80%, 90%, 95%, 98% or 100% sequence identity with SEQ ID NO: 9.

In an embodiment, the plant genome can be modified at specific position using for example KeyBase®, targeted nucleotide exchange (TNE), oligo-directed mutagenesis (ODM) or Oligonucleotide Directed Targeted Mutagenesis (ODTM) to increase or introduce the expression of a histidine kinase as defined herein.

In some embodiment, methods for editing the genome of the plant cell to introduce or increase the expression of an endogenous histidine kinase as defined herein includes, but

is not limited to, the use of specific nucleases such as the CRISPR system, ZFNs or TALENs. Non-limiting examples of specific nucleases of the CRISPR system include Cas9, Cpf1 and CasX.

Instead or in addition to introducing or increasing the expression of an endogenous histidine kinase, an exogenous histidine kinase protein can be stably expressed in a plant cell. For example, a nucleic acid encoding a histidine kinase, e.g. an exogenous histidine kinase, can be stably inserted into the genome of a plant cell. The expression of such histidine kinase can be regulated by sequences operably linked to the introduced sequence. These regulating sequences can be endogenous to the cell, or may be introduced e.g. together with the introduction of the nucleic acid encoding the histidine kinase. These regulating sequences can include genomic as well as epigenomic regulators.

The method of the invention may further comprise a step of testing overexpression or de novo expression of the histidine kinase of the method of the invention. In other words, the invention also provides for a method for improving a cytokinin-induced regeneration capacity of a plant cell comprising the steps of:

- a) increasing or introducing the expression of a histidine kinase as defined herein in the plant cell; and,
- b) detecting the expression level of said histidine kinase in the plant cell.

Methods for testing overexpression or de novo expression of the histidine kinase protein include, but are not limited to, PCR analysis, sequencing of genomic DNA, sequencing of mRNA transcript, analyzing mRNA transcript levels (Northern-blot analysis), analyzing copy number (Southern blot analysis), etc. Preferably, the method of the invention results in at least 2%, 5%, 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 100% increase of expression of the histidine kinase as compared to the control as defined herein.

In an embodiment, the plant cell is obtained or obtainable from a plant that is known to have an inefficient regeneration capacity. An insufficient regeneration capacity is herein understood as the regeneration capacity is less than about 25%, 20%, 15%, 10%, 8%, 5%, 4%, 3%, 2%, or 1%, preferably when the plant cell is kept under defined conditions, preferably optimal conditions, that allow for regeneration. Examples of plants having an inefficient regeneration capacity include, but are not limited to sweet pepper, cucumber and melon.

The plant cell for use in the invention may be obtained or obtainable from a plant that is incapable to regenerate under conditions that should allow for regeneration.

In one embodiment, the plant cell is obtained or obtainable from a plant selected from the group consisting of barley, cabbage, canola, cassava, cauliflower, chicory, cotton, cucumber, eggplant, grape, hot pepper, lettuce, maize, melon, oilseed rape, potato, pumpkin, rice, rye, *sorghum*, squash, sugar cane, sugar beet, sunflower, sweet pepper, tomato, water melon, wheat, zucchini, soybean, *chrysanthemum* and *Arabidopsis*.

In one embodiment, the plant cell is obtained or obtainable from a plant selected from the group consisting of sweet pepper, cucumber, melon, soybean, *chrysanthemum* and *Arabidopsis*.

In an embodiment, the plant cell is obtained or obtainable from a plant selected from the group consisting of sweet pepper, cucumber and melon.

In one embodiment, the plant cell is obtained or obtainable from tomato or pepper, preferably the plant cell is obtained or obtainable from *Solanum lycopersicum* or *Cap-sicum annuum*.

In a second aspect, the invention pertains to a method for regenerating a plant comprising the steps of i) incubating a plant cell obtained or obtainable by the method of the first aspect of the invention in a medium comprising a cytokinin as defined herein; and ii) allowing the plant cell to regenerate into a plant.

The invention therefore also pertains to a method for regenerating a plant comprising the steps of:

- a) increasing or introducing the expression of a histidine kinase as defined herein in the plant cell;
- b) optionally detecting the expression level of said histidine kinase in the plant cell and optionally selecting a plant cell having an increased or introduced expression of the histidine kinase;
- c) incubating the plant cell, optionally the plant cell selected in step b), in a medium comprising a cytokinin as defined herein; and
- d) allowing the plant cell to regenerate into a plant.

In one embodiment, the plant cell is a plant cell as specified in the first aspect of the invention. The plant cell can be obtained or obtainable from a plant selected from the group consisting of barley, cabbage, canola, cassava, cauliflower, chicory, cotton, cucumber, eggplant, grape, hot pepper, lettuce, maize, melon, oilseed rape, potato, pumpkin, rice, rye, *sorghum*, squash, sugar cane, sugar beet, sunflower, sweet pepper, tomato, water melon, wheat, zucchini, soybean, *chrysanthemum* and *Arabidopsis*.

The method of the invention relates to improving a cytokinin-induced regeneration capacity of a plant cell, wherein the method comprises a step of increasing or introducing the expression of a histidine kinase protein. Cytokinins (CK) are a class of plant growth substances (phytohormones) that promote cell division, or cytokinesis, in plant roots and shoots. Cytokinins can travel up the xylem and promote lateral growth. They are involved primarily in cell growth and differentiation, but can also affect apical dominance, axillary bud growth, and leaf senescence.

The cytokinin for use in the invention can be an adenine-type cytokinin or a phenylurea-type cytokinin. Similarly, the cytokinin can be a naturally produced phytohormone or can be a synthesized compound. The adenine-type cytokinin can be a phytohormone that is synthesized in at least one of roots, seeds and fruits. In addition, cambium and other actively dividing tissues can also synthesize cytokinins.

A non-limiting example of a naturally occurring adenine-type cytokinin is Zeatin as well as its metabolic precursor 2iP. Non-limiting examples of synthetic adenine-type cytokinins are kinetin and 6-benzylaminopurine (BAP). Substituted urea compounds, such as thidiazuron and CPPU do not occur in plants but can act as cytokinins in tissue culture.

The adenine-type cytokinin can be selected from the group consisting of kinetin, zeatin, trans-zeatin, cis-zeatin, dihydrozeatin, 6-benzylaminopurine and 2iP, and combinations thereof. The phenylurea-type cytokinin can be diphenylurea or thidiazuron.

In one embodiment, the plant cell as defined herein is incubated in a medium comprising a cytokinin as defined herein under conditions that allow for the regeneration of the plant cell. The plant cell can be incubated in the medium comprising a cytokinin for at least about 1, 2, 3, 4, 5, 6, 7, 8, 9 or about 10 days, under conditions that allow for regeneration. The plant cell can be incubated in the medium comprising a cytokinin for at least about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or about 12 weeks, under conditions that allow for regeneration.

In one embodiment, the plant cell as defined herein is incubated in a medium comprising a cytokinin for at least 5, 6, 7, 8, 9 weeks.

In one embodiment, the regenerated plant cell can be further developed into a cotyledonary embryo. The terms "cotyledonary embryo" and "cotyledonary stage embryo" can be used interchangeably herein. The cotyledonary embryo is preferably a cotyledonary stage somatic embryo. In an embodiment, all cells of the cotyledonary embryo express or overexpress the histidine kinase as defined herein. The mutagenized cotyledonary embryo produced by the method of the invention can be further developed into a plantlet. The plantlet can further be developed into a plant.

In an embodiment of the invention, the method therefore comprises a further step of regenerating a plantlet and/or a plant from the plant cell.

In one embodiment, the plant cell as defined herein is incubated in a medium comprising a cytokinin. The cytokinin in the medium can be only one type of cytokinin. Alternatively, the plant cell can also be incubated in a mixture of cytokinins. For example, the plant cell can be incubated in a medium comprising at least 1, 2, 3, 4 or 5 different types of cytokinins.

A mixture of at least two cytokinins is for example selected from the group consisting of kinetin and zeatin, kinetin and trans-zeatin, kinetin and cis-zeatin, kinetin and dihydrozeatin, kinetin and 6-benzylaminopurine, kinetin and 2iP, zeatin and trans-zeatin, zeatin and cis-zeatin, zeatin and dihydrozeatin, zeatin and 6-benzylaminopurine, zeatin and 2iP, trans-zeatin, and cis-zeatin, trans-zeatin and dihydrozeatin, trans-zeatin and 6-benzylaminopurine, trans-zeatin and 2iP, cis-zeatin and dihydrozeatin, cis-zeatin and 6-benzylaminopurine, cis-zeatin and 2iP, dihydrozeatin and 6-benzylaminopurine, dihydrozeatin and 2iP, and 6-benzylaminopurine and 2iP. The skilled person understands that (an) additional type(s) of cytokinin(s) can be added to the mixture of the at least two cytokinins. It is known in the art that the type of added cytokinin is dependent on the type of plant cell and the skilled person can straightforwardly select the suitable cytokinin(s).

In one embodiment, the medium can contain additional compounds that promote the regeneration of the plant cell. Such additional compounds can for example be one or more growth regulators, e.g. one or more plant hormones. Plant hormones (also known as phytohormones or 'plant growth substances') are chemicals that can regulate plant growth. A plant hormone can affect at least one of plant shape, seed growth, flowering time, the sex of flowers, senescence of leaves, and senescence of fruits. Alternatively or in addition, plant hormone can affect at least one of upward growth of tissues, downward growth of tissues, leaf formation, stem growth, fruit development, fruit ripening, plant longevity, and plant death.

The further plant hormone for use in the method of the invention can be at least one of an auxin, abscisic acid, ethylene and a gibberellin. Alternatively or in addition, the further plant hormone can be at least one of a brassinosteroid, salicylic acid, a jasmonate, a plant peptide hormone, a polyamine, nitric oxide, a strigolactones, a Karrikin and triacontanol.

The at least one further plant hormone can be an auxin. Auxins are a class of plant hormones that can have morphogen-like characteristics. The auxin can be an endogenously synthesized auxin. The endogenously synthesized auxin can be selected from the group consisting of indole-

3-acetic acid (IAA), 4-chloroindole-3-acetic acid, phenylacetic acid, indole-3-butyric acid and indole-3-propionic acid.

The auxin can be a synthetic auxin, e.g. an auxin analog. The synthetic auxin can be at least one of 1-naphthaleneacetic acid, 2,4-dichlorophenoxyacetic acid (2,4-D), α -Naphthalene acetic acid (α -NAA), 2-Methoxy-3,6-dichlorobenzoic acid (dicamba), 4-Amino-3,5,6-trichloropicolinic acid (tordon or picloram), 1-naphthaleneacetic acid (NAA), indole-3-butyric acid (IBA) and 2,4,5-trichlorophenoxyacetic acid (2,4,5-T). The auxin can be 1-naphthaleneacetic acid (NAA).

In addition or alternatively, the further plant hormone can be at least a gibberellin. The gibberellin can be a 19-carbon gibberellin or a 20-carbon gibberellin. The gibberellin can be a dihydroxylated gibberellin. The gibberellin can be at least one of GA1, GA3, GA4 and GA7.

The concentration of the at least one further plant hormone (e.g. auxin) in the medium can be the same or similar to the concentration cytokinin (e.g. a ratio of about 1:1). Alternatively, the concentration of the at least one further plant hormone can be lower than the concentration cytokinin in the medium. The ratio of the at least one further plant hormone to cytokinin can be about 0.9:1.0; 0.8:1.0; 0.7:1.0; 0.6:1.0; 0.5:1.0; 0.4:1.0; 0.3:1.0; 0.2:1.0; 0.1:1.0; 0.01:1.0 or even about 0.001:1.0.

Alternatively, the concentration of the at least one further plant hormone can be higher than the concentration cytokinin in the medium. The ratio of the at least one further plant hormone to cytokinin can be about 1:0.9; 1:0.8; 1:0.7; 1:0.6; 1:0.5; 1:0.4; 1:0.3; 1:0.2; 1:0.1; 1:0.01 or even about 1:0.001.

The concentration cytokinin in the medium can be at least about 50 ng/ml, 75 ng/ml, 100 ng/ml, 125 ng/ml, 150 ng/ml, 175 ng/ml, 200 ng/ml, 225 ng/ml, 250 ng/ml, 275 ng/ml, 300 ng/ml, 325 ng/ml, 350 ng/ml, 375 ng/ml, 400 ng/ml, 425 ng/ml, 450 ng/ml, 475 ng/ml, 500 ng/ml, 525 ng/ml, 550 ng/ml, 575 ng/ml, 600 ng/ml, 625 ng/ml, 650 ng/ml, 675 ng/ml, 700 ng/ml, 750 ng/ml, 800 ng/ml, 850 ng/ml, 900 ng/ml, 950 ng/ml, 1000 ng/ml, 1100 ng/ml, 1200 ng/ml, 1300 ng/ml, 1400 ng/ml, 1500 ng/ml, 1750 ng/ml, 2000 ng/ml, 2225 ng/ml, 2500 ng/ml, 2750 ng/ml, 3000 ng/ml, 3250 ng/ml, 3500 ng/ml, 3750 ng/ml, 4000 ng/ml, 4250 ng/ml, 4500 ng/ml, 4750 ng/ml or at least about 5000 ng/ml.

Alternatively or in addition, the concentration cytokinin in the medium can at most about 50 ng/ml, 75 ng/ml, 100 ng/ml, 125 ng/ml, 150 ng/ml, 175 ng/ml, 200 ng/ml, 225 ng/ml, 250 ng/ml, 275 ng/ml, 300 ng/ml, 325 ng/ml, 350 ng/ml, 375 ng/ml, 400 ng/ml, 425 ng/ml, 450 ng/ml, 475 ng/ml, 500 ng/ml, 525 ng/ml, 550 ng/ml, 575 ng/ml, 600 ng/ml, 625 ng/ml, 650 ng/ml, 675 ng/ml, 700 ng/ml, 750 ng/ml, 800 ng/ml, 850 ng/ml, 900 ng/ml, 950 ng/ml, 1000 ng/ml, 1100 ng/ml, 1200 ng/ml, 1300 ng/ml, 1400 ng/ml, 1500 ng/ml, 1750 ng/ml, 2000 ng/ml, 2225 ng/ml, 2500 ng/ml, 2750 ng/ml, 3000 ng/ml, 3250 ng/ml, 3500 ng/ml, 3750 ng/ml, 4000 ng/ml, 4250 ng/ml, 4500 ng/ml, 4750 ng/ml or about 5000 ng/ml.

In an embodiment, the concentration cytokinin in the medium is in the range of about 50-5000 ng/ml, 75-4000 ng/ml, 100-3000 ng/ml, 125-2000 ng/ml, 130-1500 ng/ml, 140-1250 ng/ml, 150-1000 ng/ml, 175-800 ng/ml, 200-600 ng/ml or about 250-500 ng/ml.

The concentration cytokinin in the medium is preferably a concentration that is optimal to allow regeneration of a plant cell. The skilled person knows how to establish such optimal concentrations, e.g. through routine experimental

tion or these concentrations have been described previously in the art. A preferred optimal concentration is about 1000 ng/ml.

The medium for incubating the plant cell as defined herein can be a liquid medium or a solid medium. The medium is preferably sterile.

In the method and use of the invention as defined herein in the different aspects, the plant cell can be part of a multicellular tissue. A plant multicellular tissue can comprise differentiated cells. Alternatively or in addition, a multicellular tissue can comprise undifferentiated cells. In an embodiment, all cells of the multicellular tissue have an increased or introduced expression of a histidine kinase as defined herein. Alternatively, about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or about 99% of the cells of the multicellular tissue have an increased or de novo expression of a histidine kinase as defined herein.

The multicellular tissue can be a callus tissue, a plant organ or an explant.

The plant cell having induced or introduced expression of a histidine kinase as defined herein can be part of a callus. A callus is a group of undifferentiated cells, preferably derived from adult cells. Callus cells can be capable of undergoing embryogenesis and formation of an entirely new plant. A plant callus is considered a growing mass of unorganized plant parenchyma cells. Callus can be produced from a single differentiated cell, and callus cells can be totipotent, being able to regenerate the whole plant body. The plant callus can be derived from a somatic tissue or tissues, e.g. a tissue that is available for explant culture. The cells that give rise to callus and somatic embryos preferably undergo rapid division and/or are partially undifferentiated such as meristematic tissue. The callus cell used in the method of the invention can be friable or compact. In addition or alternatively the callus cell can be rooty, shooty, or embryogenic callus (Ikeuchi M, Plant Cell. 2013 September; 25(9): 3159-3173).

In an embodiment, the plant cell having an increased or introduced expression of a histidine kinase as defined herein can be part of a plant organ. The plant organ can be a vegetative organ or a reproductive organ. A vegetative organ can be derived from the shoot system or root system. The organ can be at least one of roots, stems and leaves. A reproductive plant organ can be selected from the group consisting of flower, seed, fruit, cone, sori, strobili and gametophores.

In an embodiment, the plant cell having an increased or introduced expression of a histidine kinase as defined herein can be part of an explant. An explant can be defined herein as a sample obtained from a part of a plant. The plant sample can be placed on a solid culture medium or liquid medium. Explants can be taken from many different parts of a plant, including portions of shoots, leaves, stems, flowers, roots, single undifferentiated cells and from mature cells. The cells preferably contain living cytoplasm and nuclei and are able to de-differentiate and resume cell division. An explant can be, or can be obtainable or obtained from, a meristematic end of a plant, such as e.g. the stem tip, axillary bud tip or root tip. In one embodiment, the explant is selected from the group consisting of a hypocotyl explant, a stem explant, a cotyledon explant, a root explant, a leaf explant, a flower explant and a meristematic tissue. In one embodiment, the explant is a hypocotyl explant.

The plant regenerated by the method of the invention may subsequently be crossed to remove the increased or introduced expression of a histidine kinase as defined herein.

21

Hence the method may comprise a step of crossing the regenerated plant to remove the previously increased or introduced expression of a histidine kinase. The plant may be crossed with a plant of a different species or of the same species and progeny that no longer has an increased or introduced expression of a histidine kinase may be selected.

In a third aspect, the invention pertains to a plant or plant part obtainable or obtained by the method of the invention as defined herein. The plant or plant part can be obtainable or obtained by the process of organogenesis or somatic embryogenesis. Plant cells obtained from the plant or plant part have an increased or introduced expression of a histidine kinase as defined herein. The plant obtainable or obtained by the method of the invention is preferably selected from the group consisting of barley, cabbage, canola, cassava, cauliflower, chicory, cotton, cucumber, eggplant, grape, hot pepper, lettuce, maize, melon, oilseed rape, potato, pumpkin, rice, rye, *sorghum*, squash, sugar cane, sugar beet, sunflower, sweet pepper, tomato, water melon, wheat, zucchini, soybean, *chrysanthemum* and *Arabidopsis*. The plant, plant part or plant cell can have an increased or introduced expression of the histidine kinase as defined herein in all cells. Alternatively about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or about 99% of the plant cells have an increased or introduced expression of the histidine kinase as defined herein. In one embodiment, the plant part is a seed, a fruit or a non-propagating material.

In an embodiment, the plant or the plant cell of the invention comprises a genetic modification, wherein the modification results in an increased or introduced expression of a histidine kinase, as compared to a control plant. Preferred genetic modifications are indicated herein above. The plant or plant cell of the invention preferably comprises a nucleic acid of the eight aspect as defined herein and/or an expression construct of the ninth aspect as defined herein.

In a fourth aspect, the invention concerns a product derived from the plant or plant part obtainable or obtained by the method of the invention, e.g. fruits, leaves, plant organs, plant fats, plant oils, plant starch, and plant protein fractions, either crushed, milled or still intact, mixed with other materials, dried, frozen, and so on. These products may be non-propagating. Preferably, said plant product comprises a genetic modification, wherein the modification results in an increased or introduced expression of a histidine kinase, as compared to a control plant. Preferred genetic modifications are indicated herein above. Said plant product may comprise a nucleic acid of the eight aspect as defined herein and/or an expression construct of the ninth aspect as defined herein. Preferably, these products comprise at least fractions of said genetic modification, nucleic acid and/or construct, which allows to assess that the plant product is derived from a plant obtained by the method of the first and/or second aspect of the invention as defined herein.

In a fifth aspect, the invention concerns progeny of the plant cell, plantlet or plant obtainable or obtained by the method of the invention. Hence, the plant cells of the progeny have an increased or induced expression of a histidine kinase as defined herein and have an improved cytokinin-induced regeneration capacity. Said progeny may comprise the genetic modification, nucleic acid and/or an expression construct of the plant or plant part of the third aspect of the invention as defined herein.

In a sixth aspect, the invention pertains to the use of a CHK2, CHK3 and/or CHK4 histidine kinase, as defined herein for improving a cytokinin-induced regeneration capacity of a plant. In one embodiment, the invention

22

concerns the use of a CHK2 and/or CHK4 histidine kinase as defined herein for improving a cytokinin-induced regeneration capacity of a plant. In a further embodiment, the invention concerns the use of at least a CHK4 histidine kinase as defined herein for improving a cytokinin-induced regeneration capacity of a plant. The plant is preferably a plant as defined herein above.

In a seventh aspect, the invention pertains to the use of a CHK2, CHK3 and/or CHK4 histidine kinase as defined herein above in a method for improving a cytokinin-induced regeneration capacity of a plant. Preferably, the CHK2, CHK3 and/or CHK4 histidine kinase is used in a method as defined in the first and/or second aspect of the invention, for improving a cytokinin-induced regeneration capacity of a plant.

In an eighth aspect, the invention concerns a nucleic acid encoding a histidine kinase as defined herein. In a preferred embodiment, the nucleic acid encodes for at least one of CHK2, CHK3 and CHK4 as defined herein. Preferably the nucleic acid encodes for least one of CHK2 and CHK4 as defined herein. Preferably, the nucleic acid encodes for CHK4 as defined herein.

In one embodiment, the sequence of the nucleic acid can have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with at least one of SEQ ID NO: 1, SEQ ID NO: 2 and SEQ ID NO: 3.

Alternatively, the sequence of the nucleic acid can have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with at least one of SEQ ID NO: 7, SEQ ID NO: 8 and SEQ ID NO: 9.

The nucleic acid can be isolated from its natural environment. In addition or alternatively, at least 1, 2, 3 or 4 nucleotides can flank the nucleic acid as defined herein, wherein said nucleotides do not flank the nucleic acid in a natural environment, i.e. resulting in a non-naturally occurring nucleic acid.

In a ninth aspect, the invention pertains to an expression construct for the expression of a histidine kinase as defined herein. In an embodiment, the expression construct comprises the sequence of a nucleic acid as defined herein above in the eighth aspect. The expression construct can comprise the sequence of at least 1, 2, 3, 4 or 5 nucleic acids as defined herein above in the eighth aspect.

The expression construct can comprise the sequence of at least two or more copies of the same nucleic acid and/or can comprise the sequence of at least two different nucleic acids as defined herein.

As a non-limiting example, the expression construct can comprise at least the sequence of a nucleic acid having at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with SEQ ID NO: 1 as well as the sequence of a nucleic acid having at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with SEQ ID NO: 3,

As another non-limiting example, the expression construct can comprise at least the sequence of a nucleic acid having at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with SEQ ID NO: 7 as well as the sequence of a nucleic acid having at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with SEQ ID NO: 9,

The expression construct can further comprise one or more regulatory elements that are operably linked to the

nucleic acid sequence or nucleic acid sequences as defined herein. A preferred regulatory element is a promoter. The promoter for expression in a plant cells is herein understood as a promoter that is active in plants or plant cells, i.e. the promoter has the general capability to drive transcription within a plant or plant cell.

The promoter can be a constitutive promoter, an inducible promoter or a tissue specific promoter. It is understood herein that a “constitutive” promoter is a promoter that is active in most tissues under most physiological and developmental conditions. An “inducible” promoter is a promoter that is physiologically (e.g. by external application of certain compounds) or developmentally regulated. A “tissue specific” promoter is only active in specific types of tissues or cells.

The promoter can be a caulimovirus promoter, such as a cauliflower mosaic virus (CaMV) promoter, a nopaline synthase promoter or an octopine synthase promoter.

The promoter can have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with at least one of SEQ ID NO: 10, SEQ ID NO: 11, SEQ ID NO: 12 and SEQ ID NO: 13. Preferably, the promoter can have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 98% or about 100% sequence identity with SEQ ID NO: 12.

The present invention has been described above with reference to a number of exemplary embodiments. Modifications and alternative implementations of some parts or elements are possible, and are included in the scope of protection as defined in the appended claims.

EXAMPLES

Example 1: Cloning of the *Arabidopsis* CHK2 Gene

The coding sequence of the *Arabidopsis* CHK2 gene (AT5G35750) was amplified by PCR using CHK2 specific forward primer 14_04109 and reverse primer 14_04110 from *Arabidopsis thaliana* Col-0 genomic DNA as template. A second PCR amplification was performed with primers 14_04116 (forward) and 14_04117 (reverse) to incorporate Gateway cloning attB sites at the ends of the fragments. The amplification products were purified from gel and cloned into pDONR221 donor vector (Invitrogen™) via standard Gateway BP reaction cloning (thermofisher).

The 2.0 kb promoter sequence upstream of the AtCHK2 gene (AT5G35750) was synthesized by gene synthesis (ThermoFisher Scientific). The synthesized fragment was PCR-amplified using forward primer 15_00083 and reverse primer 15_00084, and cloned into entry vector pENTR™ 5'-TOPO using the pENTR™ 5'-TOPO® TA Cloning® Kit (Invitrogen™).

Example 2: Cloning of the *Arabidopsis* CHK4 Gene

The coding sequence of the *Arabidopsis* CHK4 gene (AT2G01830) was amplified by PCR using CHK4 specific forward primer 14_04106 and reverse primer 14_04107 from *Arabidopsis thaliana* Col-0 genomic DNA as template. A second amplification was performed with primers 14_04114 (forward) and 14_04115 (reverse) to incorporate Gateway cloning attB sites at the ends of the fragments. The amplification products were purified from gel and cloned

into pDONR221 donor vector (Invitrogen™) via standard Gateway BP reaction cloning (thermofisher).

The 2.0 kb promoter sequence upstream of the AtCHK4 gene (AT2G01830) was synthesized by gene synthesis (ThermoFisher Scientific). The synthesized fragment was PCR-amplified using forward primer 15_00087 and reverse primer 15_00088, and cloned into entry vector pENTR™ 5'-TOPO using the pENTR™ 5'-TOPO® TA Cloning® Kit (Invitrogen™).

Example 3: Construction of Expression Constructs pKG9785 and pKG9791

An expression construct for the *Arabidopsis* CHK2 gene under control of its native 2 kb promoter was obtained by combining the CHK2 promoter fragment and the CHK2 coding sequence fragment from the entry vectors described in Example 1 into multisite Gateway destination vector pK7m24gw,3 through a single step LR cloning reaction (ThermoFisher).

An expression construct for the *Arabidopsis* CHK4 gene under control of its native 2 kb promoter was obtained by combining the CHK4 promoter fragment and the CHK4 coding sequence fragment from the entry vectors described in Example 2 into multisite Gateway destination vector pK7m24gw,3 through a single step LR cloning reaction (ThermoFisher).

The resulting plasmid constructs containing either CHK coding sequence under control of its native promoter were checked using restriction enzyme digestion with the combination NcoI and SacI, or the combination XbaI and HindIII, prior to transformation to *Agrobacterium tumefaciens* strain GV3101 via electroporation. The final constructs were named pKG9785 for the CHK2 expression cassette and pKG9791 for the CHK4 expression cassette, respectively. Individual *Agrobacterium* colonies were selected and used in plant transformation to obtain transgenic lines.

Example 4: Transformation of *Arabidopsis*

Transgenic plants with stable integration of either CHK construct (pKG9785 or pKG9791) were obtained by transforming *Arabidopsis thaliana* Col-0 plants using the floral dip method according to the protocol of Clough and Bent (Plant J. 16 (6): 735-743, 1998). Positive transformants were obtained by sterilizing T1 progeny seed using vapor phase (Clough and Bent, 1998, Plant J. 16 (6): 735-743), germinating the seeds in vitro on 0.5MS10 culture medium (which is half strength MS medium according to Murashige and Skoog, 1962, Physiol. Plant. 15: 473-497 containing 100 g·L⁻¹ sucrose) supplemented with 50 mg·L⁻¹ kanamycin, and selecting kanamycin-resistant seedlings.

Multiple independent kanamycin-resistant seedlings for each construct were subsequently transferred to soil and selfed to obtain T2 seeds. Seeds were sterilized using vapor phase and sown on 0.5MS10 plates supplemented with 50 mg·L⁻¹ kanamycin. After two weeks, T3 lines of which the seedlings did not segregate for kanamycin resistance or sensitivity were considered homozygous and used in the regeneration assays. Expression levels of the transgenes in selected transgenic lines was checked by qRT-PCR using RNA extracted from whole in vitro seedlings and compared to the expression levels of the endogenous CHK genes in non-transformed Col-0 control plants. The primers used for qRT-PCR were 17_03894 (forward primer) and 17_03895 (reverse primer) for AtCHK4, and 17_03890 (forward) and

17_03891 (reverse) for AtCHK2. For each of constructs, homozygous plant lines were selected with a moderate to strong expression level to be tested for their regeneration capacity.

Example 5: Regeneration Assays of *Arabidopsis*
Lines Expressing CHK Genes

The regeneration assays used are based on the shoot initiation assays described by To et al. (Plant Cell 16: 658-671, 2004, which incorporated herein by reference). *Arabidopsis* seedlings of the tested lines were grown on 0.5MS10 culture medium in square plates placed vertically in the dark for 3-4 days and then in dim light for 3 days to produce elongated and firm hypocotyls. Hypocotyls of around 7 mm in length were excised from the seedlings. Hypocotyl explants were transferred to MS medium (Murashige and Skoog, 1962, Physiol. Plant. 15: 473-497) with 1% (w/v) sucrose and 0.4% (w/v) phytigel containing kinetin (300 ng·mL⁻¹) and NAA (100 ng·mL⁻¹) and kept for 7 weeks at 25° C. at 16 h light/8 h dark. For each transgenic line 10 individual explants were tested in single square plates that also included five non-transformed (negative) control explants of *Arabidopsis* Col-0 and five explants of the *Arabidopsis* arr3,4,5,6,8,9 hexuple mutant serving as a positive control for regeneration. This mutant is known to regenerate at high efficiencies under these conditions (To et al., 2004, Plant Cell 16: 658-671). In each experiment four independent plates were assayed and the experiment was conducted in duplicate. Shoot regeneration rates were determined based on the number of inflorescence shoot that were formed after 7 weeks of growth.

Regeneration experiments were monitored, imaged and scored over seven weeks. All transgenic lines displayed

enhanced regeneration in comparison to the control Col-0 line (Table 2). Regeneration efficiency was expressed as the percentage of explants displaying inflorescence shoot formation out of the total number of explants cultured. It is clear from Table 2 that the regeneration efficiency of plant material expressing CHK2 or CHK4 genes under control of their native promoters outperforms the regeneration efficiency of the positive control (hexuple arr mutant). Overall it is clear that enhanced expression of either CHK2 or CHK4 genes improves the regeneration capacity.

The percentage amino acid identity between CHK2, CHK3 and CHK4 is indicated in Table 3.

TABLE 1A

Sequences	
SEQ ID NO:	Sequence
1	Arabidopsis thaliana Nucleotide sequence CHK2
2	Arabidopsis thaliana Nucleotide sequence CHK3
3	Arabidopsis thaliana Nucleotide sequence CHK4
4	Arabidopsis thaliana Amino acid sequence CHK2
5	Arabidopsis thaliana Amino acid sequence CHK3
6	Arabidopsis thaliana Amino acid sequence CHK4
7	Arabidopsis thaliana Genomic sequence CHK2
8	Arabidopsis thaliana Genomic sequence CHK3
9	Arabidopsis thaliana Genomic sequence CHK4
10	Arabidopsis thaliana promoter sequence CHK2
11	Arabidopsis thaliana promoter sequence CHK3
12	Arabidopsis thaliana promoter sequence CHK4
13	Arabidopsis thaliana promoter sequence CHK2 with SNP

TABLE 1B

primer sequences		
SEQ ID NO:	Primer name	Sequence
14	pAtCHK2_F	TAATTTGAATATTTATATTCAATTTTCATACATAT
15	pAtCHK2_R	TTCGACTCCTAATCTCAGATTCA
16	pAtCHK4_F	CCAATTCACGTTAAATCTATCTCTTG
17	pAtCHK4_R	CACTTCAAATGTAGGTATTCCATTTT
18	AtCHK2_F_CDS	ATGTCCTATACTTGTGAGCTCTTGAA
19	AtCHK2_R_CDS	TTAACAAGGTTCAAGAATCTTGC
20	AtCHK2_F_CDS_attB1F	GGGGACAAGTTTGTACAAAAAGCAGGCTATGTCTATA ACTTGTGAGCTCTTGAA
21	AtCHK2_R_CDS_attB2R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTAAACAAG GTTCAAGAATCTTGC
22	AtCHK4_F_CDS	ATGAGAAGAGATTTTGTGTATAATAATAATGC
23	AtCHK4_R_CDS	TTACGACGAAGGTGAGATAGGA
24	AtCHK4_F_CDS_attB1F	GGGGACAAGTTTGTACAAAAAGCAGGCTATGAGAAG AGATTTTGTGTATAATAATAATGC
25	AtCHK4_R_CDS_attB2R	GGGGACCACTTTGTACAAGAAAGCTGGGTTTACGACG AAGGTGAGATAGGA
26	Q_CHK4_F	TAAGCCTGTTAGAGCTGCTGTG
27	Q_CHK4_R	TCTTTCAAACGCAGCAGCTG

TABLE 1B-continued

primer sequences		
SEQ ID NO: Primer name	Sequence	
28 Q_CHK2_F	GCTACAGAAGAACAGCGTGTG	
29 Q_CHK2_R	TGTTTGCTGGCAAGTTGGTG	

TABLE 2

Regeneration efficiencies of Arabidopsis hypocotyl explants expressing CHK alleles on medium with kinetin and NAA			
Type of CHK alleles	# explants	# regener.	% regener.
Effect of CHK alleles			
AtCHK2 (pKG9785)	79	40	50.6
AtCHK4 (pKG9791)	68	43	63.2
Controls			
Arabidopsis arr3,4,5,6,8,9 hexuple mutant	360	160	44.4
Col-0 negative control	360	7	1.9

TABLE 3

Percentage amino acid sequence identity between CHK2, CHK3 and CHK4			
	CHK2	CHK3	CHK4
CHK2	100%	58%	54%
CHK3		100%	53%
CHK4			100%

Example 6: Identifying the Tomato CHK2 Orthologue

An orthology search was carried out with a query of the AtCHK2 and AtCHK4 genes (AT5G35750 and AT2G0183) in 15 plant proteome databases using multiple protein sequence alignment software JackHMMER (<http://hmmer.org>, version 3.2.1; June 2018; Johnson et al., BMC Bioinformatics, 11: 431, 2010, doi: 10.1186/1471-2105-11-431). JackHMMER blast searches were supplemented with a MEME Suite motif-based sequence analysis (<http://meme-suite.org/index.html>; Meme Suite 4.12.0., June 2017; Bailey et al., Nucl. Acids Res. 37: W2302-W208, 2009, doi.org/10.1093/nar/gkp335) to identify and visualize known protein domains. The orthology search revealed single candidate genes in the tomato genome for both AtCHK2 and AtCHK4 genes as queries, with amino acid identities of >50%. These tomato genes are Solyc07g047770 (SlyCHK2) and Solyc07g008110 (SlyCHK4) as most likely homologs of AtCHK2 and AtCHK4, respectively. Solyc07g047770 (SlyCHK2) was selected for further work.

Example 7: Construction of a Tomato CHK Expression Cassette

A functional expression cassette of SlyCHK2 containing 2 kb of its native promoter sequence was designed in silico.

The nopaline synthase terminator and Gateway adapters were appended to the sequence to create a cloning fragment for direct cloning in a Gateway destination vector. The synthesis of the cassette was ordered from GeneART (www.thermofisher.com). The full sequence of the cassette is given in SEQ ID NO: 32. The fragment was introduced in Gateway binary destination vector pKm43GW (Karimi et al., Plant Physiol. 145: 1144-1154, 2007, doi.org/10.1104/pp.107.106989) through a single step LR cloning reaction (www.thermofisher.com). pKm43GW is a Gateway multi-site binary destination vector containing streptomycin and spectinomycin resistance markers for bacterial selection, and an nptII gene for selection of plant tissue on kanamycin. The resulting plasmid construct was named pKG10867 and was cloned in *E. coli* and checked by restriction enzyme digestion using the combination EcoRV and NheI. Miniprep plasmid DNA was electroporated to *Agrobacterium tumefaciens* strain GV3101.

Example 8: Tomato Transformation

The SlyCHK2 under its native promoter was introduced into tomato cultivar Moneyberg-Plus (TMV-resistant) by *Agrobacterium*-mediated transformation. Approximately 50 tomato seeds were sterilized and germinated on % MS10 medium for 11 days. Cotyledon explants from the seedlings were dissected and precultured for 24 h on 2N1B medium (=M520 medium containing 2 mg·l⁻¹ NAA and 1 mg·l⁻¹ BAP) supplemented with 40 µg·l⁻¹ acetosyringone. The explants were submerged in a suspension of *Agrobacterium tumefaciens* GV3101 carrying pKG10867 grown overnight in TY medium containing 20 mg·l⁻¹ streptomycin and 50 mg·l⁻¹ spectinomycin, and diluted to OD₆₀₀ 0.138. The explants were blotted dry and cocultivated for 2 days on 2N1B plates with 40 µg·l⁻¹ acetosyringone. Subsequently, the explants were transferred to selective medium MS20ZVCK consisting of MS20 medium with 1 mg·l⁻¹ zeatin, 200 mg·l⁻¹ vancomycin, 200 mg·l⁻¹ cefotaxim and 100 mg·l⁻¹ kanamycin and cultivated at 25° C. and 3000 lux (16/8 h photoperiod) in a growth chamber. The explants were subcultured every 3 weeks onto fresh medium. When callus had formed on the selective medium, the callus was subcultured and the original explants were discarded.

Example 9: Recording Transformation Efficiencies

Tomato callus subcultured every 3 weeks onto fresh medium MS20ZCVK started to produce shoot meristems and shoots from day 114 of the experiment. Shoots over 5 mm in length were taken off the calli, and transferred to

rooting medium consisting of MS20 without any additions. The accumulated number of shoots harvested in this way was recorded (Table 4) and compared to similarly cultivated tomato Moneyberg cotyledon explants that had not been contacted with *Agrobacterium tumefaciens* and were grown on medium without kanamycin. It is clear that shoots harvested from kanamycin-resistant callus after pKG10867-transformation appear in larger numbers and faster than shoots regenerating from control tomato explants. This effect is attributed to the ectopic expression of the SlyCHK2 gene.

TABLE 4

Shoot regeneration efficiencies of tomato Moneyberg cotyledon explants transformed with SlyCHK2 (pKG10867), recorded as the cumulative number of shoots harvested from transformed calli over time.

	#	day 114	day 136	day 156	day 183
Moneyberg pKG10867	200	1	26	47	93
Moneyberg control	200	1	7	29	64

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 32

<210> SEQ ID NO 1

<211> LENGTH: 3531

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 1

```

atgtctataa cttgtgagct cttgaatctt acttcaaaga aagctaagaa gtcgtcgagc   60
agtgacaaga aatggctaaa gaagcctctc ttcttcctga ttttgtgtgg ctctttggta   120
attgttttgg ttatgttctt acggttaggt agaagtcaga aggaggagac agattcttgt   180
aatggagaag agaaagtgtt gtatagacat caaatgtca caagaagtga gattcatgat   240
ttggtctctt tgttctctga ttcagatcag gtaacatcct ttgaatgtca taaggaatca   300
agccctggaa tgtggacaaa ctatggtatt acatgttccc tgagtgtgcg ttctgataaa   360
caagagacta gagggcttcc ctggaatctt ggcttaggac attctatctc atcaacatct   420
tgtatgtgtg gtaatcttga accgatttta cagcaacctg aaaaccttga ggaagaaaac   480
catgaagaag ggctggagca ggggtttgtca tcgtatttaa gaaatgcatg gtggtgtcta   540
atccttggtg tgttagtgtg ccataagatt tatgtatctc attctaaagc acgaggtgag   600
aggaaagaga agtacatctt gcaagaggct ttagctccaa agaagcagca acaacgtgct   660
cagacttctt ctagaggggc tggaagatgg aggaagaata tccttctcct tgggtatttta   720
ggaggagttt ccttctctgt ttggtggttt tgggacacta atgaggagat cataatgaaa   780
aggagggaga ctttgccaaa catgtgtgac gaacgagcac gtgttttaca agatcagttc   840
aatgttagct tgaaccatgt tcatgccttg tctattcttg tatctacatt tcatcatggt   900
aaaatcccat ctgccattga tcagagaaca tttgaagaat atactgagag aacaaacttt   960
gagaggccac ttactagtgg tgtagcgtat gctttgaaag tcccacactc agaaagagag  1020
aaatttgaaa aggagcatgg atgggcaata aagaaaatgg aaactgagga ccagacagtt  1080
gtacaagatt gtgttctctga aaactttgat ccgcaccca ttcaagacga atacgcgcca  1140
gttatatttg ctcaagaaac tgtttcccat attgtatcgg tcgacatgat gtctggagaa  1200
gaagaccgtg aaaacatctt acgggcaagg gcatcaggaa aaggagtgtt aacatctcca  1260
tttaagcttc ttaagtcaaa tcatcttggg gttgtgttga cctttgctgt ctatgacacg  1320
agcctaccgc ctgatgctac agaagaacag cgtgttgaag caactattgg gtaccttggg  1380
gcatcatatg atatgccate gctggtggag aaacttcttc accaacttgc cagcaaacag  1440
acaattgctg tggatgttta cgacacaact aacacttcag gtctaataaa aatgtatggc  1500
tcagaaattg gggatataag tgagcagcat ataagtagcc ttgatttttg tgatccatca  1560
aggaacctat agatgcattg caggtttaag cataaacttc ccattccctg gacagcgata  1620

```


-continued

acaccgtcga tcttagttct ggttattact tttcttgttg gttatatttt atatgaagcc	1680
atcaaccgaa ttgcgacagt tgaagaggat tgtcagaaga tgagggaact caaagctcgt	1740
gctgaggccg ctgacattgc aaagtcacag ttcctagcaa ctgtttctca tgagatacgg	1800
actccgatga atggagtttt aggaatgctg aaaatgctga tggacaccga tcttgatgcg	1860
aagcagatgg actatgcgca aactgctcat ggcagtgagg aggatcttac atcactaata	1920
aatgaggttc ttgatcaggc aaagattgaa tccggaaggc tcgagcttga aaatgtgcct	1980
tttgatatgc gttttattct tgataatgtt tcactctccc tctctggcaa ggcaaatgaa	2040
aaaggaattg agttggccgt ttatgtttct agtcaagttc ctgatgttgt agtcggtgat	2100
ccgagtcggt tccggcagat cattacaaac ctggttggaa actcaatcaa attcacacag	2160
gaaaggggac acatatattat ctcaagtgcac ctgacagatg aggtaaagga gcctcttact	2220
attgaagacg cagtgcataa acagcgacta gctttaggat gcagcgagtc cggtgagaca	2280
gttagcgggt ttcctgcggt aaatgcatgg ggaagctgga agaatttcaa gacatgttac	2340
agtactgaga gtcagaattc tgatcaaatc aaattgctag ttacagtgga ggacactgga	2400
gttggcatac ctgtggatgc acaaggccga atcttcacac cttttatgca agccgacagt	2460
tccacatcgc ggacttatgg tggaaactgc ataggtttga gtataagcaa acgtttggtt	2520
gaactcatgc aaggagagat ggggtttgtg agtgagcccg ggataggcag tactttttca	2580
tttactggag ttttcgggaa agcagaaaca aatacgtcga ttactaagct ggaacgattc	2640
gatctagcta ttcaggagtt tacaggattg agagcattag ttattgataa cagaaacatt	2700
agagcagagg tcaccaggta cgaacttcgg agactgggaa tatctgcaga cattgtttca	2760
agtctgagaa tggcatgcac ttgttgtatc agcaaattag aaaatttggc tatgattcta	2820
atagacaaag acgcctggaa caaggaagaa ttttcagtac ttgacgagtt gtttaccgca	2880
agcaaagtaa cctttacaag agtcccaaag atttttcttt tggcaacttc tgcaactctt	2940
actgagcgca gtgagatgaa gtctactggt ctcatcgatg aggtggtgat aaagcctctt	3000
cggatgagtg tcttaatatg ttgcttgcaa gaaacccttg tcaatggcaa gaagaggcaa	3060
ccgaacagac agcgaagaaa tcttgacac ttgctaagag aaaaacagat tctggttgtg	3120
gatgataatc ttgtgaacag acgagttgca gaaggtgcac ttaagaaata tggagctatt	3180
gttacatcgc ttgagagtgg caaagctgca ttggcaatgc ttaagccgcc tcataacttc	3240
gatgcttgct tcatggatct ccagatgcct gaaatggatg gatttgaagc gacaaggaga	3300
gtccgtgagc tggagaggga aatcaataag aaaatagctt ctggagaagt ttcagctgaa	3360
atgttctgta aatttagtag ttggcacgtc ccgatattag caatgacagc agatgttatt	3420
caggctactc atgaagaatg catgaaatgt ggaatggatg gttatgtatc aaaaccgttt	3480
gaagaggaag tgctctacac agcggtagca agattctttg aaccttgtaa a	3531

<210> SEQ ID NO 2

<211> LENGTH: 3111

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 2

atgagtctgt tccatgtgct aggggtttggt gtcaagattg ggcactctct ctggatgcta	60
tgctgctggt ttgtttcttg gttcgttgat aatgggatcg aggacaagtc tggctcttta	120
gttggctctg tcggtgatct tgagaagact aagatgacta cgttgaagaa gaagaacaag	180
atgtggttct ggaataagat ctctagcagc ggactcaaga tcccagttt ctcttatcag	240

-continued

tttcttggt	ctgttaaatt	caacaaggcg	tggtaggagga	agcttggtgt	ggtttgggtt	300
gtcttctggg	tcttggtctc	tatttggtacg	ttttggtact	ttagctcgca	agctatggag	360
aagagggaaa	agacgctagc	tagtatgtgt	gatgagagag	ctcgtatgct	gcaggatcag	420
ttcaacgtta	gcataaatca	tgttcaagcc	atgtctatct	tgatctcaac	cttccaccat	480
ggcaagattc	cttctgctat	cgatcagaga	acattctcag	agtaactga	tagaacttcc	540
tttgagaggc	ctcttactag	cggggtagct	tatgctatga	gggtgctcca	ttcagagagg	600
gaagagttcg	agaggcaaca	aggttggtact	attaggaaga	tgtattctct	tgaacaaaac	660
ccagttcaca	aggatgacta	tgacctggaa	gctttggaac	catccctgt	ccaagaagag	720
tacgtccag	tcctctttgc	tcaggacact	gtttctcacg	ttgtttctct	cgatatgctg	780
tctgggaaa	aagatcgtga	aaacgttttg	cgggccagga	gttcaggtaa	aggggttttg	840
acagctcctt	tccattgat	aaagacaaat	agacttgggg	tgatcctgac	atttgcagtg	900
tacaagagag	atctcccctc	caatgcaacg	ccaaaagaga	gaattgaggc	tactaacggg	960
tatctcgggg	gagtgtttga	cattgagtcc	ctggtagaaa	acttgcctca	acagctgggt	1020
agcaagcaaa	cgattcttgt	caatgtgtac	gatatcacca	atcactctca	accgattagc	1080
atgtatggta	caaagtgtgc	ggctgatggg	ttggaacgtg	ttagtccact	aatctttggc	1140
gatccattga	gaaagcatga	gatgcgttgc	agatttaagc	agaaaccacc	atggccagtg	1200
ctatcaatgg	tgacatcatt	cggatccctt	gtgattgctg	tacttgttgc	acataatac	1260
cacgcaaccg	ttagtcgaat	acacaaagtt	gaagaagatt	gtgataaaat	gaagcagctc	1320
aagaaaaagg	ctgaagcagc	agatgttgca	aagtcacagt	tccttgccac	tgtttcacat	1380
gaaatcagaa	ctccaatgaa	tgggtgttcta	ggaatgttgc	atatgcttat	ggacacagag	1440
ttagatgtta	cgcaacagga	ttagttagg	accgcacagg	caagtggaaa	agctttagtc	1500
tcgctaataa	atgaggtttt	ggaccaagca	aagattgaat	ctggaaagct	tgaacttgag	1560
gagggtcggt	ttgatattgag	aggaatatta	gatgatgtcc	tgctactctt	ctctagcaag	1620
tcccaacaaa	aggggggtgga	gttggcagta	tacatatctg	atcgtgttcc	agatatgtta	1680
attggtgatc	ctggggagggt	tcgacaaata	ctcacaaatc	ttatgggtta	ttccattaag	1740
ttcactgaga	aaggacacat	ctttgttaact	gttcatttgg	tggatgagct	atttgaatct	1800
atcgatggag	agacagcatc	atctccggaa	agtacactga	gtgggcttcc	agttgcagac	1860
cggcagagga	gctgggaaaa	ctttaaagct	ttcagctcca	acgggcatcg	gagctttgaa	1920
ccatctcccc	ctgatataaa	cctaactcgtc	tcagttgagg	atactggcgt	agggatccct	1980
gtagaagcgc	agtcctgtat	ttttacgcct	ttcatgcaag	tcggaccatc	catatccagg	2040
acgcattggag	gcacaggaat	tggacttagc	ataagcaaat	gtctagttag	actgatgaag	2100
ggagaaattg	gattctcgag	tactcccaag	gttgggtcca	cattcacatt	tactgctgta	2160
ttttccaatg	ggatgcaacc	agctgaaaga	aagaatgaca	acaaccagcc	catattctcg	2220
gaattccggg	gcataaaagc	tgtggttggt	gaccataggc	ctgcaagggc	aaaagtctcg	2280
tggtagcatt	ttcagcgtct	tgggaattcga	gtcgaagtag	ttccacgtgt	tgaacaggct	2340
ctacattatc	tgaagattgg	tactaccact	gtgaatatga	tactcataga	gcaagaaata	2400
tggaaatagg	aagcagatga	tttcattaaa	aagctacaga	aagaccctct	tttcctttct	2460
cctaagttga	ttttgttagc	aaactcagta	gaatcgtcaa	tatcagaggc	tttatgcacc	2520
ggtatagatc	ctccaatagt	gatagtgaag	ccattgaggg	cgagtatgct	agcagcaact	2580

-continued

ttgcagaggg gattgggtat tggaatcaga gaaccacctc aacacaaggg acctcctgct	2640
ttgattctca ggaatcttct ccttggtaga aaaattttaa tcgtggatga taacaacgta	2700
aacctcagag tggcagcggg agctctgaaa aagtaaggag ctgatgtggt ctgcgctgag	2760
agtgggataa aggcaatctc attgcttaag ccacctcacg agtttgatgc ttgcttcacg	2820
gacattcaga tgccagaaat ggatggattt gaagctacaa ggagaatacg agatatggaa	2880
gaggagatga acaagagaat aaagaatggg gaggccttga tagtagagaa cggttaacaaa	2940
acaagctggc atcttccggg attagcaatg acggcagatg tgatccaagc aacgcatgag	3000
gaatgtctga agtgtggaat ggatgggtat gtatcaaac catttgaagc agagcagctg	3060
tacaggggaag tttctcgctt tttcaattcg ccttcagata cagaatcata a	3111

<210> SEQ ID NO 3

<211> LENGTH: 3243

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 3

atgagaagag attttgtgta taataataat gcaatgttca atcctctcac aactcattac	60
agctcagata tgaactgggc actcaacaat catcaagaag aagaagaaga gccacgaaga	120
attgaaattt ctgattccga gtcactagaa aacttgaaaa gcagcgattt ttatcaactg	180
ggtggtggtg gtgctctgaa ttcgtcagaa aagccgagaa agatcgattt ttggcgctcg	240
gggttgatgg gttttgcgaa gatgcagcag cagcaacagc ttcagcattc agtggcggtg	300
aagatgaaca ataataataa taacgatcta atgggtaata aaaaagggtc aactttcata	360
caagaacatc gagcattgtt accaaaagct ttgattctgt ggatcatcat tgttgggttt	420
ataagcagtg ggatttatca gtggatggat gatgctaata agattagaag ggaagaggtt	480
ttggtcagca tgtgtgatca aagagctaga atgttcagag atcaatttag tgtagtggtt	540
aatcatgttc atgctttggc tattctcgtc tccacttttc attaccacaa gaacccttct	600
gcaattgatc aggagacatt tgcggagtac acggcaagaa cagcatttga gagaccgttg	660
ctaagtggag tggcttatgc tgaaaaagtt gtgaattttg agaggagat gtttgagcgg	720
cagcacaatt ggggtataaa gacaatggat agaggagagc cttcaccggt tagggatgag	780
tatgctcctg ttatattctc tcaagatagt gtctcttacc ttgagtcact cgatatgatg	840
tcaggcgagg aggatcgtga gaatattttg cgagctagag aaaccggaaa agctgtcttg	900
actagccctt ttaggttggtt ggaaactcac catctcgag ttgtgttgac attccctgtc	960
tacaagtctt ctcttctgta aaatccgact gtcgaagagc gtattgcagc cactgcaggg	1020
taccttggtg gtgcgtttga tgtggagtct ctagtogaga atttacttgg tcagcttget	1080
ggtaaccaag caatagtgtg gcattgtgat gatatcacca atgcatcaga tccacttgtc	1140
atgtatggta atcaagatga agaagccgac agatctctct ctcatgagag caagctcgat	1200
tttgagagacc ccttcaggaa acataagatg atatgcaggt accacaaaaa ggcaccaata	1260
ccgttgaatg tgctcacaac tgtgccattg ttctttgcga ttggtttctt ggtgggttat	1320
atactgtatg gtgcagctat gcacatagta aaagtcgaag atgatttcca tgaaatgcaa	1380
gagcttaaaag ttcgagcaga agctgtgat gtcgctaaat cgcagtttct tgctaccgtg	1440
tctcagaga tcaggacacc aatgaatggc attctcggaa tgccttctat gctcctagat	1500
acagaactaa gctcgacaca gagagattac gctcaaaccg ctcaagtatg tggtaaagct	1560
ttgattgcat tgataaatga ggttcttgat cgcgccaaaga ttgaagctgg aaagctggag	1620

-continued

ttggaatcag taccatttga tatccgttca atattggatg atgtccttcc tctattctct	1680
gaggagtcaa ggaacaaaag cattgagctc gcggttttcg ttccagacaa agtaccagag	1740
atagtcaaag gagattcagg gagatttaga cagataatca taaacctgtg tggaaattcg	1800
gttaaattca cagagaaagg acatatcttt gttaaagtcc atcttgcgga acaatcaaaa	1860
gatgaatctg aaccgaaaaa tgcattgaat ggtggagtgt ctgaagaaat gatcgttggt	1920
tccaaacagt caagttacaa cacattgagc gggtacgaag ctgctgatgg tcggaatagc	1980
tgggattcat tcaagcattt ggtctctgag gagcagtcac tatcggaggt tgatatttct	2040
agcaatgtta ggcttatggt ttcaatcgaa gacacgggta ttggaatccc ttagttgctg	2100
caaggccctg tgtttatgcc gtttatgcaa gcagatagct cgacttcaag aaactatgga	2160
ggtactggta ttggtttgag tataagcaag tgtcttggtg aacttatgctg tggtcagata	2220
aatttcataa gccggcctca tattggaagc acgttctggt tcacggctgt ttagagaaa	2280
tcgataaat gcagtgcat taacctatg aagaaacctc atgtggaaca cttgccttct	2340
acttttaag gaatgaaagc tatagttggt gatgctaagc ctgtagagc tgctgtgact	2400
agataccata tgaagagact cggaatcaat gttgatgtcg tgacaagtct caaaaccgct	2460
gttggtgcag ctgctgctgt tgaagaaac ggttctctcc tcccaacaaa accgcaactt	2520
gatatgatct tagtagagaa agattcatgg atttcaactg aagataatga cttagagatt	2580
cgttttattga attcaagaac caacggaaac gttcatcaca agtctccgaa actagctcta	2640
ttcgcaacaa acatcacaaa ttcggagttc gacagagcta aatccgcagg atttgcatat	2700
acggtaataa tgaaccggtt aagagcaagc atgattgggg cgtgtctgca acaagttctc	2760
gagctgagaa aaacaagaca acaacatcca gaaggatcat caccgcgaac tctcaagagc	2820
ttgcttacag ggaagaagat tcttggtggt gatgataata tagttaacag gagagtagct	2880
gcaggagctc tcaagaaatt tggagcagaa gtggtttgtg cagagagtgg tcaagttgct	2940
ttgggtttgc ttcagattcc acacacttcc gatgcttgct tcatggatat tcaaagcca	3000
cagatggagc gatttgaagc aactcgtcag ataagaatga tggagaagga aactaaagag	3060
aagacaaatc tcgaatggca tttaccgatt cttagcgatga ctgcggatgt gatacacgcg	3120
acctacgagg aatgtctgaa aagtgggatg gatggttacg tctccaaacc tttgaagaa	3180
gagaatctct ataaatccgt tgccaaatca ttcaaaccta atcctatctc accttcgtcg	3240
taa	3243

<210> SEQ ID NO 4

<211> LENGTH: 1176

<212> TYPE: PRT

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 4

Met	Ser	Ile	Thr	Cys	Glu	Leu	Leu	Asn	Leu	Thr	Ser	Lys	Lys	Ala	Lys
1				5					10					15	

Lys	Ser	Ser	Ser	Ser	Asp	Lys	Lys	Trp	Leu	Lys	Lys	Pro	Leu	Phe	Phe
				20				25					30		

Leu	Ile	Leu	Cys	Gly	Ser	Leu	Val	Ile	Val	Leu	Val	Met	Phe	Leu	Arg
		35					40					45			

Leu	Gly	Arg	Ser	Gln	Lys	Glu	Glu	Thr	Asp	Ser	Cys	Asn	Gly	Glu	Glu
	50					55					60				

Lys	Val	Leu	Tyr	Arg	His	Gln	Asn	Val	Thr	Arg	Ser	Glu	Ile	His	Asp
65					70					75					80

Leu	Val	Ser	Leu	Phe	Ser	Asp	Ser	Asp	Gln	Val	Thr	Ser	Phe	Glu	Cys
			85						90			95			
His	Lys	Glu	Ser	Ser	Pro	Gly	Met	Trp	Thr	Asn	Tyr	Gly	Ile	Thr	Cys
			100						105			110			
Ser	Leu	Ser	Val	Arg	Ser	Asp	Lys	Gln	Glu	Thr	Arg	Gly	Leu	Pro	Trp
			115						120			125			
Asn	Leu	Gly	Leu	Gly	His	Ser	Ile	Ser	Ser	Thr	Ser	Cys	Met	Cys	Gly
			130						135			140			
Asn	Leu	Glu	Pro	Ile	Leu	Gln	Gln	Pro	Glu	Asn	Leu	Glu	Glu	Glu	Asn
			145						150			155			
His	Glu	Glu	Gly	Leu	Glu	Gln	Gly	Leu	Ser	Ser	Tyr	Leu	Arg	Asn	Ala
			165						170			175			
Trp	Trp	Cys	Leu	Ile	Leu	Gly	Val	Leu	Val	Cys	His	Lys	Ile	Tyr	Val
			180						185			190			
Ser	His	Ser	Lys	Ala	Arg	Gly	Glu	Arg	Lys	Glu	Lys	Val	His	Leu	Gln
			195						200			205			
Glu	Ala	Leu	Ala	Pro	Lys	Lys	Gln	Gln	Gln	Arg	Ala	Gln	Thr	Ser	Ser
			210						215			220			
Arg	Gly	Ala	Gly	Arg	Trp	Arg	Lys	Asn	Ile	Leu	Leu	Leu	Gly	Ile	Leu
			225						230			235			
Gly	Gly	Val	Ser	Phe	Ser	Val	Trp	Trp	Phe	Trp	Asp	Thr	Asn	Glu	Glu
			245						250			255			
Ile	Ile	Met	Lys	Arg	Arg	Glu	Thr	Leu	Ala	Asn	Met	Cys	Asp	Glu	Arg
			260						265			270			
Ala	Arg	Val	Leu	Gln	Asp	Gln	Phe	Asn	Val	Ser	Leu	Asn	His	Val	His
			275						280			285			
Ala	Leu	Ser	Ile	Leu	Val	Ser	Thr	Phe	His	His	Gly	Lys	Ile	Pro	Ser
			290						295			300			
Ala	Ile	Asp	Gln	Arg	Thr	Phe	Glu	Glu	Tyr	Thr	Glu	Arg	Thr	Asn	Phe
			305						310			315			
Glu	Arg	Pro	Leu	Thr	Ser	Gly	Val	Ala	Tyr	Ala	Leu	Lys	Val	Pro	His
			325						330			335			
Ser	Glu	Arg	Glu	Lys	Phe	Glu	Lys	Glu	His	Gly	Trp	Ala	Ile	Lys	Lys
			340						345			350			
Met	Glu	Thr	Glu	Asp	Gln	Thr	Val	Val	Gln	Asp	Cys	Val	Pro	Glu	Asn
			355						360			365			
Phe	Asp	Pro	Ala	Pro	Ile	Gln	Asp	Glu	Tyr	Ala	Pro	Val	Ile	Phe	Ala
			370						375			380			
Gln	Glu	Thr	Val	Ser	His	Ile	Val	Ser	Val	Asp	Met	Met	Ser	Gly	Glu
			385						390			395			
Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Ala	Ser	Gly	Lys	Gly	Val
			405						410			415			
Leu	Thr	Ser	Pro	Phe	Lys	Leu	Leu	Lys	Ser	Asn	His	Leu	Gly	Val	Val
			420						425			430			
Leu	Thr	Phe	Ala	Val	Tyr	Asp	Thr	Ser	Leu	Pro	Pro	Asp	Ala	Thr	Glu
			435						440			445			
Glu	Gln	Arg	Val	Glu	Ala	Thr	Ile	Gly	Tyr	Leu	Gly	Ala	Ser	Tyr	Asp
			450						455			460			
Met	Pro	Ser	Leu	Val	Glu	Lys	Leu	Leu	His	Gln	Leu	Ala	Ser	Lys	Gln
			465						470			475			
Thr	Ile	Ala	Val	Asp	Val	Tyr	Asp	Thr	Thr	Asn	Thr	Ser	Gly	Leu	Ile
			485						490			495			

Lys	Met	Tyr	Gly 500	Ser	Glu	Ile	Gly	Asp 505	Ile	Ser	Glu	Gln	His 510	Ile	Ser
Ser	Leu	Asp 515	Phe	Gly	Asp	Pro	Ser 520	Arg	Asn	His	Glu	Met 525	His	Cys	Arg
Phe	Lys 530	His	Lys	Leu	Pro	Ile 535	Pro	Trp	Thr	Ala	Ile 540	Thr	Pro	Ser	Ile
Leu 545	Val	Leu	Val	Ile	Thr	Phe 550	Leu	Val	Gly	Tyr 555	Ile	Leu	Tyr	Glu	Ala 560
Ile	Asn	Arg	Ile	Ala 565	Thr	Val	Glu	Glu	Asp 570	Cys	Gln	Lys	Met	Arg 575	Glu
Leu	Lys	Ala	Arg 580	Ala	Glu	Ala	Ala	Asp 585	Ile	Ala	Lys	Ser 590	Gln	Phe	Leu
Ala	Thr 595	Val	Ser	His	Glu	Ile	Arg 600	Thr	Pro	Met	Asn	Gly 605	Val	Leu	Gly
Met 610	Leu	Lys	Met	Leu	Met	Asp 615	Thr	Asp	Leu	Asp 620	Ala	Lys	Gln	Met	Asp
Tyr 625	Ala	Gln	Thr	Ala	His 630	Gly	Ser	Gly	Lys	Asp 635	Leu	Thr	Ser	Leu	Ile 640
Asn	Glu	Val	Leu 645	Asp	Gln	Ala	Lys	Ile	Glu 650	Ser	Gly	Arg	Leu	Glu 655	Leu
Glu	Asn	Val 660	Pro	Phe	Asp	Met	Arg	Phe 665	Ile	Leu	Asp	Asn 670	Val	Ser	Ser
Leu 675	Leu	Ser	Gly	Lys	Ala	Asn 680	Glu	Lys	Gly	Ile	Glu	Leu 685	Ala	Val	Tyr
Val 690	Ser	Ser	Gln	Val	Pro	Asp 695	Val	Val	Val	Gly	Asp 700	Pro	Ser	Arg	Phe
Arg 705	Gln	Ile	Ile	Thr	Asn 710	Leu	Val	Gly	Asn 715	Ser	Ile	Lys	Phe	Thr	Gln 720
Glu	Arg	Gly	His 725	Ile	Phe	Ile	Ser	Val	His 730	Leu	Ala	Asp	Glu	Val 735	Lys
Glu	Pro	Leu 740	Thr	Ile	Glu	Asp	Ala	Val 745	Leu	Lys	Gln	Arg 750	Leu	Ala	Leu
Gly	Cys 755	Ser	Glu	Ser	Gly	Glu	Thr 760	Val	Ser	Gly	Phe 765	Pro	Ala	Val	Asn
Ala 770	Trp	Gly	Ser	Trp	Lys	Asn 775	Phe	Lys	Thr	Cys	Tyr 780	Ser	Thr	Glu	Ser
Gln 785	Asn	Ser	Asp	Gln	Ile 790	Lys	Leu	Leu	Val 795	Thr	Val	Glu	Asp	Thr	Gly 800
Val	Gly	Ile	Pro 805	Val	Asp	Ala	Gln	Gly	Arg 810	Ile	Phe	Thr	Pro	Phe 815	Met
Gln	Ala	Asp 820	Ser	Ser	Thr	Ser	Arg	Thr 825	Tyr	Gly	Gly	Thr 830	Gly	Ile	Gly
Leu	Ser 835	Ile	Ser	Lys	Arg	Leu	Val 840	Glu	Leu	Met	Gln	Gly 845	Glu	Met	Gly
Phe 850	Val	Ser	Glu	Pro	Gly	Ile 855	Gly	Ser	Thr	Phe	Ser 860	Phe	Thr	Gly	Val
Phe 865	Gly	Lys	Ala	Glu	Thr 870	Asn	Thr	Ser	Ile	Thr 875	Lys	Leu	Glu	Arg	Phe 880
Asp	Leu	Ala	Ile 885	Gln	Glu	Phe	Thr	Gly	Leu 890	Arg	Ala	Leu	Val	Ile 895	Asp
Asn	Arg	Asn 900	Ile	Arg	Ala	Glu	Val	Thr 905	Arg	Tyr	Glu	Leu 910	Arg	Arg	Leu
Gly	Ile	Ser	Ala	Asp	Ile	Val	Ser	Ser	Leu	Arg	Met	Ala	Cys	Thr	Cys

-continued

915	920	925
Cys Ile Ser Lys Leu Glu Asn Leu Ala Met Ile Leu Ile Asp Lys Asp 930 935 940		
Ala Trp Asn Lys Glu Glu Phe Ser Val Leu Asp Glu Leu Phe Thr Arg 945 950 955 960		
Ser Lys Val Thr Phe Thr Arg Val Pro Lys Ile Phe Leu Leu Ala Thr 965 970 975		
Ser Ala Thr Leu Thr Glu Arg Ser Glu Met Lys Ser Thr Gly Leu Ile 980 985 990		
Asp Glu Val Val Ile Lys Pro Leu Arg Met Ser Val Leu Ile Cys Cys 995 1000 1005		
Leu Gln Glu Thr Leu Val Asn Gly Lys Lys Arg Gln Pro Asn Arg 1010 1015 1020		
Gln Arg Arg Asn Leu Gly His Leu Leu Arg Glu Lys Gln Ile Leu 1025 1030 1035		
Val Val Asp Asp Asn Leu Val Asn Arg Arg Val Ala Glu Gly Ala 1040 1045 1050		
Leu Lys Lys Tyr Gly Ala Ile Val Thr Cys Val Glu Ser Gly Lys 1055 1060 1065		
Ala Ala Leu Ala Met Leu Lys Pro Pro His Asn Phe Asp Ala Cys 1070 1075 1080		
Phe Met Asp Leu Gln Met Pro Glu Met Asp Gly Phe Glu Ala Thr 1085 1090 1095		
Arg Arg Val Arg Glu Leu Glu Arg Glu Ile Asn Lys Lys Ile Ala 1100 1105 1110		
Ser Gly Glu Val Ser Ala Glu Met Phe Cys Lys Phe Ser Ser Trp 1115 1120 1125		
His Val Pro Ile Leu Ala Met Thr Ala Asp Val Ile Gln Ala Thr 1130 1135 1140		
His Glu Glu Cys Met Lys Cys Gly Met Asp Gly Tyr Val Ser Lys 1145 1150 1155		
Pro Phe Glu Glu Glu Val Leu Tyr Thr Ala Val Ala Arg Phe Phe 1160 1165 1170		
Glu Pro Cys 1175		

<210> SEQ ID NO 5

<211> LENGTH: 1036

<212> TYPE: PRT

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 5

Met Ser Leu Phe His Val Leu Gly Phe Gly Val Lys Ile Gly His Leu 1 5 10 15
Phe Trp Met Leu Cys Cys Trp Phe Val Ser Trp Phe Val Asp Asn Gly 20 25 30
Ile Glu Asp Lys Ser Gly Leu Leu Val Gly Ser Val Gly Asp Leu Glu 35 40 45
Lys Thr Lys Met Thr Thr Leu Lys Lys Lys Asn Lys Met Trp Phe Trp 50 55 60
Asn Lys Ile Ser Ser Ser Gly Leu Lys Ile Pro Ser Phe Ser Tyr Gln 65 70 75 80
Phe Leu Gly Ser Val Lys Phe Asn Lys Ala Trp Trp Arg Lys Leu Val 85 90 95

-continued

Val	Val	Trp	Val	Val	Phe	Trp	Val	Leu	Val	Ser	Ile	Trp	Thr	Phe	Trp
			100					105					110		
Tyr	Phe	Ser	Ser	Gln	Ala	Met	Glu	Lys	Arg	Lys	Glu	Thr	Leu	Ala	Ser
		115					120					125			
Met	Cys	Asp	Glu	Arg	Ala	Arg	Met	Leu	Gln	Asp	Gln	Phe	Asn	Val	Ser
	130					135					140				
Met	Asn	His	Val	Gln	Ala	Met	Ser	Ile	Leu	Ile	Ser	Thr	Phe	His	His
145					150					155					160
Gly	Lys	Ile	Pro	Ser	Ala	Ile	Asp	Gln	Arg	Thr	Phe	Ser	Glu	Tyr	Thr
			165					170						175	
Asp	Arg	Thr	Ser	Phe	Glu	Arg	Pro	Leu	Thr	Ser	Gly	Val	Ala	Tyr	Ala
			180					185					190		
Met	Arg	Val	Leu	His	Ser	Glu	Arg	Glu	Glu	Phe	Glu	Arg	Gln	Gln	Gly
		195					200					205			
Trp	Thr	Ile	Arg	Lys	Met	Tyr	Ser	Leu	Glu	Gln	Asn	Pro	Val	His	Lys
	210					215					220				
Asp	Asp	Tyr	Asp	Leu	Glu	Ala	Leu	Glu	Pro	Ser	Pro	Val	Gln	Glu	Glu
225				230					235						240
Tyr	Ala	Pro	Val	Ile	Phe	Ala	Gln	Asp	Thr	Val	Ser	His	Val	Val	Ser
			245						250					255	
Leu	Asp	Met	Leu	Ser	Gly	Lys	Glu	Asp	Arg	Glu	Asn	Val	Leu	Arg	Ala
		260						265					270		
Arg	Ser	Ser	Gly	Lys	Gly	Val	Leu	Thr	Ala	Pro	Phe	Pro	Leu	Ile	Lys
		275					280					285			
Thr	Asn	Arg	Leu	Gly	Val	Ile	Leu	Thr	Phe	Ala	Val	Tyr	Lys	Arg	Asp
	290					295					300				
Leu	Pro	Ser	Asn	Ala	Thr	Pro	Lys	Glu	Arg	Ile	Glu	Ala	Thr	Asn	Gly
305				310						315					320
Tyr	Leu	Gly	Gly	Val	Phe	Asp	Ile	Glu	Ser	Leu	Val	Glu	Asn	Leu	Leu
			325						330					335	
Gln	Gln	Leu	Ala	Ser	Lys	Gln	Thr	Ile	Leu	Val	Asn	Val	Tyr	Asp	Ile
			340					345					350		
Thr	Asn	His	Ser	Gln	Pro	Ile	Ser	Met	Tyr	Gly	Thr	Asn	Val	Ser	Ala
		355					360					365			
Asp	Gly	Leu	Glu	Arg	Val	Ser	Pro	Leu	Ile	Phe	Gly	Asp	Pro	Leu	Arg
	370					375					380				
Lys	His	Glu	Met	Arg	Cys	Arg	Phe	Lys	Gln	Lys	Pro	Pro	Trp	Pro	Val
385					390					395					400
Leu	Ser	Met	Val	Thr	Ser	Phe	Gly	Ile	Leu	Val	Ile	Ala	Leu	Leu	Val
			405						410					415	
Ala	His	Ile	Ile	His	Ala	Thr	Val	Ser	Arg	Ile	His	Lys	Val	Glu	Glu
			420					425					430		
Asp	Cys	Asp	Lys	Met	Lys	Gln	Leu	Lys	Lys	Lys	Ala	Glu	Ala	Ala	Asp
		435					440					445			
Val	Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr
			450				455				460				
Pro	Met	Asn	Gly	Val	Leu	Gly	Met	Leu	His	Met	Leu	Met	Asp	Thr	Glu
465					470					475					480
Leu	Asp	Val	Thr	Gln	Gln	Asp	Tyr	Val	Arg	Thr	Ala	Gln	Ala	Ser	Gly
			485					490						495	
Lys	Ala	Leu	Val	Ser	Leu	Ile	Asn	Glu	Val	Leu	Asp	Gln	Ala	Lys	Ile
			500					505					510		
Glu	Ser	Gly	Lys	Leu	Glu	Leu	Glu	Glu	Val	Arg	Phe	Asp	Leu	Arg	Gly

-continued

515					520					525					
Ile	Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser	Ser	Lys	Ser	Gln	Gln	Lys
530						535					540				
Gly	Val	Glu	Leu	Ala	Val	Tyr	Ile	Ser	Asp	Arg	Val	Pro	Asp	Met	Leu
545					550					555					560
Ile	Gly	Asp	Pro	Gly	Arg	Phe	Arg	Gln	Ile	Leu	Thr	Asn	Leu	Met	Gly
				565					570					575	
Asn	Ser	Ile	Lys	Phe	Thr	Glu	Lys	Gly	His	Ile	Phe	Val	Thr	Val	His
			580					585					590		
Leu	Val	Asp	Glu	Leu	Phe	Glu	Ser	Ile	Asp	Gly	Glu	Thr	Ala	Ser	Ser
		595					600					605			
Pro	Glu	Ser	Thr	Leu	Ser	Gly	Leu	Pro	Val	Ala	Asp	Arg	Gln	Arg	Ser
	610					615					620				
Trp	Glu	Asn	Phe	Lys	Ala	Phe	Ser	Ser	Asn	Gly	His	Arg	Ser	Phe	Glu
625					630					635					640
Pro	Ser	Pro	Pro	Asp	Ile	Asn	Leu	Ile	Val	Ser	Val	Glu	Asp	Thr	Gly
				645					650					655	
Val	Gly	Ile	Pro	Val	Glu	Ala	Gln	Ser	Arg	Ile	Phe	Thr	Pro	Phe	Met
			660					665					670		
Gln	Val	Gly	Pro	Ser	Ile	Ser	Arg	Thr	His	Gly	Gly	Thr	Gly	Ile	Gly
		675					680						685		
Leu	Ser	Ile	Ser	Lys	Cys	Leu	Val	Gly	Leu	Met	Lys	Gly	Glu	Ile	Gly
	690					695					700				
Phe	Ser	Ser	Thr	Pro	Lys	Val	Gly	Ser	Thr	Phe	Thr	Phe	Thr	Ala	Val
705					710					715					720
Phe	Ser	Asn	Gly	Met	Gln	Pro	Ala	Glu	Arg	Lys	Asn	Asp	Asn	Asn	Gln
				725					730					735	
Pro	Ile	Phe	Ser	Glu	Phe	Arg	Gly	Met	Lys	Ala	Val	Val	Val	Asp	His
			740					745					750		
Arg	Pro	Ala	Arg	Ala	Lys	Val	Ser	Trp	Tyr	His	Phe	Gln	Arg	Leu	Gly
	755						760					765			
Ile	Arg	Val	Glu	Val	Val	Pro	Arg	Val	Glu	Gln	Ala	Leu	His	Tyr	Leu
	770					775					780				
Lys	Ile	Gly	Thr	Thr	Thr	Val	Asn	Met	Ile	Leu	Ile	Glu	Gln	Glu	Ile
785					790					795					800
Trp	Asn	Arg	Glu	Ala	Asp	Asp	Phe	Ile	Lys	Lys	Leu	Gln	Lys	Asp	Pro
				805					810					815	
Leu	Phe	Leu	Ser	Pro	Lys	Leu	Ile	Leu	Leu	Ala	Asn	Ser	Val	Glu	Ser
			820					825					830		
Ser	Ile	Ser	Glu	Ala	Leu	Cys	Thr	Gly	Ile	Asp	Pro	Pro	Ile	Val	Ile
		835					840					845			
Val	Lys	Pro	Leu	Arg	Ala	Ser	Met	Leu	Ala	Ala	Thr	Leu	Gln	Arg	Gly
	850					855					860				
Leu	Gly	Ile	Gly	Ile	Arg	Glu	Pro	Pro	Gln	His	Lys	Gly	Pro	Pro	Ala
865					870					875					880
Leu	Ile	Leu	Arg	Asn	Leu	Leu	Leu	Gly	Arg	Lys	Ile	Leu	Ile	Val	Asp
				885					890					895	
Asp	Asn	Asn	Val	Asn	Leu	Arg	Val	Ala	Ala	Gly	Ala	Leu	Lys	Lys	Tyr
			900					905					910		
Gly	Ala	Asp	Val	Val	Cys	Ala	Glu	Ser	Gly	Ile	Lys	Ala	Ile	Ser	Leu
		915					920					925			
Leu	Lys	Pro	Pro	His	Glu	Phe	Asp	Ala	Cys	Phe	Met	Asp	Ile	Gln	Met
	930					935					940				

-continued

Pro Glu Met Asp Gly Phe Glu Ala Thr Arg Arg Ile Arg Asp Met Glu
 945 950 955 960

Glu Glu Met Asn Lys Arg Ile Lys Asn Gly Glu Ala Leu Ile Val Glu
 965 970 975

Asn Gly Asn Lys Thr Ser Trp His Leu Pro Val Leu Ala Met Thr Ala
 980 985 990

Asp Val Ile Gln Ala Thr His Glu Glu Cys Leu Lys Cys Gly Met Asp
 995 1000 1005

Gly Tyr Val Ser Lys Pro Phe Glu Ala Glu Gln Leu Tyr Arg Glu
 1010 1015 1020

Val Ser Arg Phe Phe Asn Ser Pro Ser Asp Thr Glu Ser
 1025 1030 1035

<210> SEQ ID NO 6
 <211> LENGTH: 1080
 <212> TYPE: PRT
 <213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 6

Met Arg Arg Asp Phe Val Tyr Asn Asn Asn Ala Met Phe Asn Pro Leu
 1 5 10 15

Thr Thr His Tyr Ser Ser Asp Met Asn Trp Ala Leu Asn Asn His Gln
 20 25 30

Glu Glu Glu Glu Glu Pro Arg Arg Ile Glu Ile Ser Asp Ser Glu Ser
 35 40 45

Leu Glu Asn Leu Lys Ser Ser Asp Phe Tyr Gln Leu Gly Gly Gly Gly
 50 55 60

Ala Leu Asn Ser Ser Glu Lys Pro Arg Lys Ile Asp Phe Trp Arg Ser
 65 70 75 80

Gly Leu Met Gly Phe Ala Lys Met Gln Gln Gln Gln Gln Leu Gln His
 85 90 95

Ser Val Ala Val Lys Met Asn Asn Asn Asn Asn Asn Asp Leu Met Gly
 100 105 110

Asn Lys Lys Gly Ser Thr Phe Ile Gln Glu His Arg Ala Leu Leu Pro
 115 120 125

Lys Ala Leu Ile Leu Trp Ile Ile Ile Val Gly Phe Ile Ser Ser Gly
 130 135 140

Ile Tyr Gln Trp Met Asp Asp Ala Asn Lys Ile Arg Arg Glu Glu Val
 145 150 155 160

Leu Val Ser Met Cys Asp Gln Arg Ala Arg Met Leu Gln Asp Gln Phe
 165 170 175

Ser Val Ser Val Asn His Val His Ala Leu Ala Ile Leu Val Ser Thr
 180 185 190

Phe His Tyr His Lys Asn Pro Ser Ala Ile Asp Gln Glu Thr Phe Ala
 195 200 205

Glu Tyr Thr Ala Arg Thr Ala Phe Glu Arg Pro Leu Leu Ser Gly Val
 210 215 220

Ala Tyr Ala Glu Lys Val Val Asn Phe Glu Arg Glu Met Phe Glu Arg
 225 230 235 240

Gln His Asn Trp Val Ile Lys Thr Met Asp Arg Gly Glu Pro Ser Pro
 245 250 255

Val Arg Asp Glu Tyr Ala Pro Val Ile Phe Ser Gln Asp Ser Val Ser
 260 265 270

Tyr Leu Glu Ser Leu Asp Met Met Ser Gly Glu Glu Asp Arg Glu Asn

-continued

275					280					285					
Ile	Leu	Arg	Ala	Arg	Glu	Thr	Gly	Lys	Ala	Val	Leu	Thr	Ser	Pro	Phe
290						295					300				
Arg	Leu	Leu	Glu	Thr	His	His	Leu	Gly	Val	Val	Leu	Thr	Phe	Pro	Val
305					310					315					320
Tyr	Lys	Ser	Ser	Leu	Pro	Glu	Asn	Pro	Thr	Val	Glu	Glu	Arg	Ile	Ala
				325					330					335	
Ala	Thr	Ala	Gly	Tyr	Leu	Gly	Gly	Ala	Phe	Asp	Val	Glu	Ser	Leu	Val
			340					345					350		
Glu	Asn	Leu	Leu	Gly	Gln	Leu	Ala	Gly	Asn	Gln	Ala	Ile	Val	Val	His
		355					360					365			
Val	Tyr	Asp	Ile	Thr	Asn	Ala	Ser	Asp	Pro	Leu	Val	Met	Tyr	Gly	Asn
	370					375					380				
Gln	Asp	Glu	Glu	Ala	Asp	Arg	Ser	Leu	Ser	His	Glu	Ser	Lys	Leu	Asp
385					390					395					400
Phe	Gly	Asp	Pro	Phe	Arg	Lys	His	Lys	Met	Ile	Cys	Arg	Tyr	His	Gln
				405					410					415	
Lys	Ala	Pro	Ile	Pro	Leu	Asn	Val	Leu	Thr	Thr	Val	Pro	Leu	Phe	Phe
			420					425					430		
Ala	Ile	Gly	Phe	Leu	Val	Gly	Tyr	Ile	Leu	Tyr	Gly	Ala	Ala	Met	His
		435					440					445			
Ile	Val	Lys	Val	Glu	Asp	Asp	Phe	His	Glu	Met	Gln	Glu	Leu	Lys	Val
	450					455					460				
Arg	Ala	Glu	Ala	Ala	Asp	Val	Ala	Lys	Ser	Gln	Phe	Leu	Ala	Thr	Val
465					470					475					480
Ser	His	Glu	Ile	Arg	Thr	Pro	Met	Asn	Gly	Ile	Leu	Gly	Met	Leu	Ala
				485					490					495	
Met	Leu	Leu	Asp	Thr	Glu	Leu	Ser	Ser	Thr	Gln	Arg	Asp	Tyr	Ala	Gln
			500						505				510		
Thr	Ala	Gln	Val	Cys	Gly	Lys	Ala	Leu	Ile	Ala	Leu	Ile	Asn	Glu	Val
		515					520						525		
Leu	Asp	Arg	Ala	Lys	Ile	Glu	Ala	Gly	Lys	Leu	Glu	Leu	Glu	Ser	Val
	530					535					540				
Pro	Phe	Asp	Ile	Arg	Ser	Ile	Leu	Asp	Asp	Val	Leu	Ser	Leu	Phe	Ser
545					550					555					560
Glu	Glu	Ser	Arg	Asn	Lys	Ser	Ile	Glu	Leu	Ala	Val	Phe	Val	Ser	Asp
				565					570					575	
Lys	Val	Pro	Glu	Ile	Val	Lys	Gly	Asp	Ser	Gly	Arg	Phe	Arg	Gln	Ile
			580					585					590		
Ile	Ile	Asn	Leu	Val	Gly	Asn	Ser	Val	Lys	Phe	Thr	Glu	Lys	Gly	His
		595					600						605		
Ile	Phe	Val	Lys	Val	His	Leu	Ala	Glu	Gln	Ser	Lys	Asp	Glu	Ser	Glu
	610					615					620				
Pro	Lys	Asn	Ala	Leu	Asn	Gly	Gly	Val	Ser	Glu	Glu	Met	Ile	Val	Val
625					630					635					640
Ser	Lys	Gln	Ser	Ser	Tyr	Asn	Thr	Leu	Ser	Gly	Tyr	Glu	Ala	Ala	Asp
				645					650					655	
Gly	Arg	Asn	Ser	Trp	Asp	Ser	Phe	Lys	His	Leu	Val	Ser	Glu	Glu	Gln
				660				665					670		
Ser	Leu	Ser	Glu	Phe	Asp	Ile	Ser	Ser	Asn	Val	Arg	Leu	Met	Val	Ser
		675					680					685			
Ile	Glu	Asp	Thr	Gly	Ile	Gly	Ile	Pro	Leu	Val	Ala	Gln	Gly	Arg	Val
	690					695					700				

Phe	Met	Pro	Phe	Met	Gln	Ala	Asp	Ser	Ser	Ser	Thr	Ser	Arg	Asn	Tyr	Gly
705					710							715				720
Gly	Thr	Gly	Ile	Gly	Leu	Ser	Ile	Ser	Lys	Cys	Leu	Val	Glu	Leu	Met	
			725						730					735		
Arg	Gly	Gln	Ile	Asn	Phe	Ile	Ser	Arg	Pro	His	Ile	Gly	Ser	Thr	Phe	
			740					745					750			
Trp	Phe	Thr	Ala	Val	Leu	Glu	Lys	Cys	Asp	Lys	Cys	Ser	Ala	Ile	Asn	
		755					760					765				
His	Met	Lys	Lys	Pro	Asn	Val	Glu	His	Leu	Pro	Ser	Thr	Phe	Lys	Gly	
	770					775					780					
Met	Lys	Ala	Ile	Val	Val	Asp	Ala	Lys	Pro	Val	Arg	Ala	Ala	Val	Thr	
					790					795					800	
Arg	Tyr	His	Met	Lys	Arg	Leu	Gly	Ile	Asn	Val	Asp	Val	Val	Thr	Ser	
			805						810					815		
Leu	Lys	Thr	Ala	Val	Val	Ala	Ala	Ala	Ala	Phe	Glu	Arg	Asn	Gly	Ser	
			820					825					830			
Pro	Leu	Pro	Thr	Lys	Pro	Gln	Leu	Asp	Met	Ile	Leu	Val	Glu	Lys	Asp	
		835					840					845				
Ser	Trp	Ile	Ser	Thr	Glu	Asp	Asn	Asp	Ser	Glu	Ile	Arg	Leu	Leu	Asn	
	850					855					860					
Ser	Arg	Thr	Asn	Gly	Asn	Val	His	His	Lys	Ser	Pro	Lys	Leu	Ala	Leu	
				865		870				875					880	
Phe	Ala	Thr	Asn	Ile	Thr	Asn	Ser	Glu	Phe	Asp	Arg	Ala	Lys	Ser	Ala	
			885						890					895		
Gly	Phe	Ala	Asp	Thr	Val	Ile	Met	Lys	Pro	Leu	Arg	Ala	Ser	Met	Ile	
			900					905					910			
Gly	Ala	Cys	Leu	Gln	Gln	Val	Leu	Glu	Leu	Arg	Lys	Thr	Arg	Gln	Gln	
		915					920					925				
His	Pro	Glu	Gly	Ser	Ser	Pro	Ala	Thr	Leu	Lys	Ser	Leu	Leu	Thr	Gly	
	930					935					940					
Lys	Lys	Ile	Leu	Val	Val	Asp	Asp	Asn	Ile	Val	Asn	Arg	Arg	Val	Ala	
	945				950					955					960	
Ala	Gly	Ala	Leu	Lys	Lys	Phe	Gly	Ala	Glu	Val	Val	Cys	Ala	Glu	Ser	
			965					970						975		
Gly	Gln	Val	Ala	Leu	Gly	Leu	Leu	Gln	Ile	Pro	His	Thr	Phe	Asp	Ala	
			980					985					990			
Cys	Phe	Met	Asp	Ile	Gln	Met	Pro	Gln	Met	Asp	Gly	Phe	Glu	Ala	Thr	
		995					1000					1005				
Arg	Gln	Ile	Arg	Met	Met	Glu	Lys	Glu	Thr	Lys	Glu	Lys	Thr	Asn		
	1010					1015					1020					
Leu	Glu	Trp	His	Leu	Pro	Ile	Leu	Ala	Met	Thr	Ala	Asp	Val	Ile		
	1025					1030					1035					
His	Ala	Thr	Tyr	Glu	G											

```
<210> SEQ ID NO 7
<211> LENGTH: 4595
<212> TYPE: DNA
<213> ORGANISM: Arabidopsis thaliana
```

-continued

<400> SEQUENCE: 7

atgtctataa cttgtgagct cttgaatcctt acttcaaaga aagctaagaa gtcgtcgagc	60
agtgacaaga aatggctaaa gaagcctctc ttcttctga ttttgtgtgg ctctttggta	120
attgttttgg ttatgttctt acggttaggt agaagtcaga aggaggagac agattcttgt	180
aatggagaag agaaagtgtt gtatagacat caaaatgtca caagaagtga gattcatgat	240
ttggtctctt tgttctctga ttcagatcag gtaattgcat gaattgactt gtatttgttg	300
aaattgagct tttgaatacg caccagattt gccttaaagt gtaacagttt ctctgtattt	360
gttgttaggt acatcctttg aatgtcataa ggaatcaagc cctggaatgt ggacaaacta	420
tggtattaca tgttccctga gtgtgcgttc tgataaaca gagactagag ggcttccctg	480
gaatcttggc ttaggacatt ctatctcatc aacatcttgt atgtgtggta atcttgaacc	540
ggtaagataa tattcgattc gacaacatgt gaaggaaatg ctttaagatt tgggtgtttag	600
ctagtcttca caaaagttta ttgtgaatgt gtttggttat gagtagattt tacagcaacc	660
tgaaaacctt gaggaagaaa accatgaaga agggctggag cagggtttgt catcgtattt	720
aagaaatgca tgggtggtgc taatccttgg tgtgttagtg tgccataaga tttatgtatc	780
tcattctaaa gcacgagggt agaggaaaga gaaagtacat ctgcaagagg ctttagctcc	840
aaagaagcag caacaacgtg ctgagacttc ttctagaggg gctggaagat ggaggaagaa	900
tatccttctc cttggtattt taggaggagt ttccttctct gtttggtggt tttgggacac	960
taatgaggag atcataatga aaaggaggga gactttggca aacatgtgtg acgaacgagc	1020
acgtgtttta caagatcagt tcaatgtag cttgaaccat gttcatgcct tgtctattct	1080
tgtatctaca tttcatcatg gtaaaatccc atctgccatt gatcagggtga tgttttttct	1140
ttactgctaa atacattttg tgtctcaagt ttatgtttaa atcatcaact tctgttacat	1200
ttacagagaa catttgaaga atatactgag agaacaaact ttgagaggcc acttactagt	1260
ggtgtagcgt atgctttgaa agtccacac tcagaaagag agaaatttga aaaggagcat	1320
ggatgggcaa taaagaaaat ggaaactgag gaccagacag ttgtacaaga ttgtgttctt	1380
gaaaactttg atccgcaccc gattcaagac gaatacgcgc cagttatatt tgctcaagaa	1440
actgtttccc atattgtatc ggtcgacatg atgtctggag aagtcagtaa cgtctaaaag	1500
tttcttgaac tattttgcca aaccaatgtc cttaaaagag aattcaaaag tctaactatt	1560
ttgcaggaag accgtgaaaa catcttacgg gcaagggcat caggaaaagg agtgtaaca	1620
tctccattta agcttcttaa gtcaaatcat cttggtgttg tgttgacctt tgctgtctat	1680
gacacgagcc tacgcctga tgtacagaa gaacagcgtg ttgaagcaac tattgggtac	1740
tacttctact taaaatgatt ctaggactga agaaattgaa cctatgtaac aaagaatgat	1800
ctctggacca gaaaatatta ataagataca cttaaaaaa ggtaccttgg tgcatcatat	1860
gatatgccat cgctggtgga gaaacttctt caccaacttg ccagcaaaac gacaattgct	1920
gtggatgttt acgacacaac taacacttca ggtctaataa aaatgtatgg ctcaaaaatt	1980
gggatataa gtgagcagca tataagtagc cttgattttg gtgatccatc aaggaacct	2040
gagatgcatt gcaggtagt tagctctaac tgttatggta catttttata agatatgttt	2100
cacttctctg ttttctgaag tatgaaagtg gcatttttca tttacagggt taagcataaa	2160
cttcccttc cctggacagc gataacaccg tegatcttag ttctggttat tacttttctt	2220
gttggttata ttttatatga agccatcaac cgaattgcga cagttgaaga ggattgtcag	2280
aagatgaggg aactcaaagc tcgtgctgag gccgctgaca ttgcaaagtc acagggtgatc	2340

-continued

tttgtgaatc atatatgtctc aaaagcttta cttgtttttt cactgaaatg cttcttattt	2400
tgcagttcct agcaactggt tctcatgaga tacggactcc gatgaatgga gttttaggta	2460
ctttttctac ttatccttgg tcaatattgc atgttctctt aaaatcagct gaaacgttta	2520
agcattttgt tacaggaatg ctgaaaatgc tgatggacac cgatcttgat gcgaagcaga	2580
tggactatgc gcaaaactgct catggcagtg ggaaggatct tacatcacta ataaatgagg	2640
ttcttgatca ggcaaagatt gaatccggaa ggctcgagct tgaaaatgtg ccttttgata	2700
tgcgttttat tcttgataat gtttcacttc tctctcttgg caaggcaaat gaaaaaggaa	2760
ttgaggata attataaact gcatgacctt ctactttctt aatgttttca tatggcaaac	2820
aaatccata tgtaatgaaa tgttgacttc ttggtacagt tggccgttta tgtttctagt	2880
caagtctctg atgttgtagt cggtgatccg agtcggttcc ggcagatcat tacaaacctg	2940
gttggaact caatcaaagt aatttactcc ttacttctta cagaacaaca ggcttcacga	3000
atctctacta ttacagtact catttggtat ttacttaata acagttcaca caggaaaggg	3060
gacacatatt tatctcagtg caccttgtag atgaggtaaa ggagcctctt actattgaag	3120
acgcagtgct aaaacagcga ctagcttttag gatgcagcga gtccggtgag acagttagcg	3180
ggtttctctg ggtaaatgca tggggaagct ggaagaattt caagacatgt tacagtactg	3240
agagtcagaa ttctgatcaa atcaaattgc tagttacagt ggaggacact ggagtggca	3300
tacctgtgga tgcacaaggc cgaatcttca caccttttat gcaagccgac agttccacat	3360
cgcggactta tgggtggaact ggcataaggt tgagtataag caaacgtttg gttgaactca	3420
tgcaaggaga gatggggttt gtgagtgagc ccgggatagg cagtactttt tcatttactg	3480
gagttttcgg gaaagcagaa acaaatacgt cgattactaa gctggaacga ttcgatctag	3540
ctattcagga gtttacagga ttgagagcat tagttattga taacagaaac attagagcag	3600
aggtcaccag gtacgaactt cgggagactg gaatatctgc agacattgtt tcaagtctga	3660
gaatggcatg cacttggtgt atcaggtact ttacgtacat tagtgtctgt ctgtctttag	3720
agattatagt gagttcacta aaagcgtttt attgttctgg aatctttgca gcaaattaga	3780
aaatttggct atgattctaa tagacaaaga cgcctggaac aaggaagaat tttcagtact	3840
tgacgagttg tttacccgaa gcaaagtaac ctttacaaga gtcccaaaga tttttctttt	3900
ggcaacttct gcaactctta ctgagcgag tgagatgaag tctactggtc tcatcgatga	3960
ggtggtgata aagcctcttc ggatgagtggt cttaatatgt tgcttgcaag aaacccttgt	4020
caatggcaag aagaggcaac cgaacagaca gcgaagaaat cttggacact tgctaagaga	4080
aaaacagatt ctggttggtg atgataatct tgtgaacaga cgagttgcag aagggtgact	4140
taagaaatat ggagctattg ttacatgcgt tgagagtggc aaagctgcat tggcaatgct	4200
taagccgct cataacttcg atgcttgctt catggatctc cagatgcctg aaatggatgg	4260
gtagataact atttttcaat cttatctctt cagcttggtc atttcttga tgcgtcctt	4320
ggtaacttta taattttttg cgcagatttg aagcgacaag gagagtccgt gagctggaga	4380
gggaaatcaa taagaaaata gcttctggag aagtttcagc tgaaatgttc tgtaaattta	4440
gtagttggca cgtcccgata tttagcaatga cagcagatgt tattcaggct actcatgaag	4500
aatgcatgaa atgtggaatg gatggttatg tatcaaaacc gtttgaagag gaagtgcctt	4560
acacagcggg agcaagattc tttgaacctt gttaa	4595

-continued

<211> LENGTH: 4248

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 8

```

atgagtctgt tccatgtgct aggggttgggt gtcaagattg ggcattctctt ctggatgcta      60
tgctgctggt ttgtttcttg gtctgtgat  aatgggacgc aggacaagtc tggctcttcta      120
gttgctctcg tcggtgatct tgagaagact aagatgacta cgttgaagaa gaagaacaag      180
atgtggttct ggaataagat ctctagcagc ggactcaaga tcccagagttt ctcttatcag      240
tttcttggct ctgttaaatt caacaaggcg tggtaggagga agcttgtggt ggtttgggtt      300
gtctctggtg tcttgggtct tatttggacg ttttggtagt ttagctcgca agctatggag      360
aagaggaaag agacgctagc tagtatgtgt gatgagagag ctctgatgct gcaggatcag      420
ttcaacgcta gcataaatca tgttcaagcc atgtctatct tgatctcaac ctccaccat      480
ggcaagattc cttctgctat cgatcaggta ttctcgattc cgcattctct tgtctcttaa      540
tctgtttgga ttgatgcctg agatctagtc ttgctttagt tacataagtt gttctctcgg      600
tgtagtatat gaacggctct tggatgatgg tagctttttt cattcaactt gctctcgaat      660
ttcagttagg atttacaatt ttgctgattg ttgcttacag tctttgtatt tagctaacac      720
ttggtagctt gattcctatc ttcttgagta ttagtgaaca gtaactaaca ttaagaagct      780
ttgtgtagag aacatttctc gagtacactg atagaacttc ctttgagagg cctcttacta      840
gctgggttagc ttatgctatg aggggtgctc attcagagag ggaagagttc gagaggcaac      900
aagggttgac tattaggaag atgtattctc ttgaacaaaa cccagttcac aaggatgact      960
atgacctgga agcttttgaa ccataccctg tccaagaaga gtacgctcca gtcattcttg     1020
ctcaggacac tgtttctcac gttgtttctc tcgatatgct gtctgggaaa gtaagcttct     1080
ttactgggtc taattttact gtttttatgt aatttactat catggtctac actcactacc     1140
gggcgaatcc tgcgacatg gctagtccat atttcaatct cggcttaccg attaagttgt     1200
tttgatcatt ttattttaaag ttttagagca tgttttaact gatacttatt ctcaactaat     1260
tcctcttgat aggaagatcg tgaaaacgtt ttgcgggcca ggagttcagg taaaggggtt     1320
ttgacagctc ctttccatt gataaagaca aatagacttg gggtagacct gacatttgca     1380
gtgtacaaga gagatctccc ctccaatgca acgcaaaaag agagaattga ggtactaac     1440
gggtgtgtag cataggcgct taaataaata ataacgcata agttttacaa cattattttt     1500
gctcattcat ctcttttgat gccctgaag gtatctcggg ggagtgtttg acattgagtc     1560
cctggtagaa aacttgcttc aacagctggc tagcaagcaa acgattcttg tcaatgtgta     1620
cgatataacc aatcactctc aaccgattag catgtatggt acaaatgtgt cggtgatgg     1680
gttgaacgt  gttagtccac taatctttgg cgatccattg agaaagcatg agatgcgttg     1740
caggtagctg cagttggcac atacatatgt ctgtaatttc ttcttgtttg caagaatcca     1800
ggtgctaaca ttttgtgtg agcttcttcc tctttgtaga tttaagcaga aaccaccatg     1860
gccagtgcta tcaatggtga catcattcgg tatccttgtg attgcgttac ttgttgcaac     1920
tataatccac gcaaccgtta gtcgaataca caaagttgaa gaagattgtg ataaatgaa     1980
gcagctcaag aaaaaggctg aagcagcaga tgttgcaaaag tcacaggtaa atacatcatc     2040
ctcagttgat aaatttccac agcatattaa actttctatg acagagagaa gagttgtaat     2100
ctaacttttt tttatgcagt tccttgccac tgtttcacat gaaatcagaa ctccaatgaa     2160
tgggtgttcta ggtgagtatc gaaatggcca ctatgttgcc tccttctctc acctatgcc     2220

```

-continued

atgaatat	tctgaagaac	aaacttaacc	aaatattctc	atgttgactt	tgatctgggg	2280
tgtaggaatg	ttgcatatgc	ttatggacac	agagttagat	gttacgcaac	aggattatgt	2340
taggaccgca	caggcaagtg	gaaaagcttt	agtctcgcta	ataaatgagg	ttttggacca	2400
agcaaagatt	gaatctggaa	agcttgaact	tgaggagggtg	cggtttgatt	tgagaggaa	2460
attagatgat	gtcctgtcac	tcttctctag	caagtcccaa	caaaaggggg	tgagagtaac	2520
ttactatatg	atctgcaaag	caagggttgt	aactgatagc	aagtctttct	tactgatttt	2580
ttgagtgttg	ctgatgaacg	cagttggcag	tatacatatc	tgatcgtgtt	ccagatatgt	2640
taattgggtga	tcctggggagg	tttcgacaaa	tactcacaaa	tcttatgggt	aattccatta	2700
aggtaaaact	ttttattatg	ttttttctg	accattcctt	gcatccgagt	tgatcaacgg	2760
agctgatcat	ttatatatat	tctggcagtt	cactgagaaa	ggacacatct	ttgtaactgt	2820
tcatttgggtg	gatgagctat	ttgaatctat	cgatggagag	acagcatcat	ctccgaaaag	2880
tacactgagt	gggcttcag	ttgcagaccg	gcagaggagc	tgggaaaact	ttaaagcttt	2940
cagctccaac	gggcacgga	gctttgaacc	atctccccct	gatataaacc	taatcgtctc	3000
agttgaggat	actggcgtag	ggatccctgt	agaagcgag	tcccgatttt	ttacgccttt	3060
catgcaagtc	ggaccatcca	tatccaggac	gcattggaggc	acaggaattg	gacttagcat	3120
aagcaaatgt	ctagttggac	tgatgaaggg	agaaattgga	ttctcgagta	ctcccagggt	3180
tgggtccaca	ttcacattta	ctgctgtatt	ttccaatggg	atgcaaccag	ctgaaagaaa	3240
gaatgacaac	aaccagccca	tattctcgga	attccggggc	atgaaagctg	tggttgtgga	3300
ccataggcct	gcaagggcaa	aagtctcgtg	gtaccatttt	cagcgtcttg	gaattcgagt	3360
cgaagtagtt	ccacgtgttg	aacaggtctt	acattatctg	aagattggta	ctaccactgt	3420
gaatatgata	ctcatagagc	aagaaatatg	gaatagggaa	gcagatgatt	tcattaaaaa	3480
gctacagaaa	gacctctttt	tcctttctcc	taagttgatt	ttgttagcaa	actcagtaga	3540
atcgtcaata	tcagaggctt	tatgcaccgg	tatagatcct	ccaatagtga	tagtgaaacc	3600
attgagggcg	agtatgctag	cagcaacttt	gcagagggga	ttgggtattg	gaatcagaga	3660
accacctcaa	cacaagggac	ctcctgcttt	gattctcagg	aatcttctcc	ttggtagaaa	3720
aattttaatc	gtggatgata	acaacgtaaa	cctcagagtg	gcagcgggag	ctctgaaaaa	3780
gtacggagct	gatgtgtgtc	gcgctgagag	tgggataaag	gcaatctcat	tgcttaagcc	3840
acctcacgag	tttgatgctt	gcttcatgga	cattcagatg	ccagaaatgg	atgggtatgc	3900
ctgattggta	tactagtttt	tttgaaaagt	tcgaaatatg	taataagaaa	attgaaatgt	3960
ttttctgccc	tgtctttctg	cagatttgaa	gctacaagga	gaatacagag	tatggaagag	4020
gagatgaaca	agagaataaa	gaatggggag	gctttgatag	tagagaacgg	taacaaaaca	4080
agctggcatc	ttccggtatt	agcaatgacg	gcagatgtga	tccaagcaac	gcagagggaa	4140
tgtctgaagt	gtggaatgga	tgggtatgta	tcaaaacat	ttgaagcaga	gcagctgtac	4200
aggggaagttt	ctcgcttttt	caattcgcct	tcagatacag	aatcataa		4248

<210> SEQ ID NO 9

<211> LENGTH: 4685

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 9

atgagaagag	atgttgtgta	taataataat	gcaatgttca	atcctctcac	aactcattac	60
------------	------------	------------	------------	------------	------------	----

-continued

aggttaactaa aataatttct ccatgtgctt gcttattagt cgttcttctt aatgttatgt	120
ttctctctgt gttctttctt tcttttggtca aagctttaat ttttttctta ttgttggtt	180
tgagacagtg aacatagcta tgttcttggt ccaataataa acaatcacgc ctgtaagag	240
cttatgattg attagtgtgt ttttttagtat taattaattt ctctgacaat aattacttag	300
tttttaattc ttctctgtaa gaaacctttg gaaactgagc aaagttgctt cttttgagaa	360
ccatgcggtt ctttctctct tttgttcttg aattcgcaaa aacatgtctt tttctgtcta	420
caggtttcta gggtttggtt ctgtactata aactatgttt atggtaacat tcttaatcat	480
aactacacta ccaatgcttt tatgttatat gtatgcaaaa aaggctctaa cttttgtttt	540
ctttcactat tgtttcttct tttgttctct attgtttag ctcagatatg aactgggcac	600
tcaacaatca tcaagaagaa gaagaagagc cacgaagaat tgaaatttct gattccgagt	660
cactagaaaa cttgaaaagc agcgattttt atcaactggg tgggtggtgt gctctgaatt	720
cgtcagaaaa gccgagaaag atcgattttt ggcgttcggg gttgatgggt ttgccaaga	780
tgcagcagca gcaacagctt cagcattcag tggcggtgaa gatgaacaat aataataata	840
acgatctaata gggtaataaa aaagggtcaa ctttcataca agaacatcga gcattgttac	900
caaaagcttt gattctgttg atcatcattg ttgggtttat aagcagtggt atttatcagt	960
ggatggatga tgctaataag attagaaggg aagaggtttt ggtcagcatg tgtgatcaaa	1020
gagctagaat gttgcaggat caatttagtg ttagtgtaa tcatgttcat gctttggcta	1080
ttctcgtctc cacttttcat taccacaaga acccttctgc aattgatcag gtcagtgtgt	1140
gctttattaa aagctcaaag ctttggtgaa atttttgatt cataaatgca aagttgactt	1200
tttttggaac cagaagtttt tgaattttac tcttttatgt ttggttattg agtaaatgaa	1260
gggaaattta ggcctttgct ttactcattg gtttggtggt aattttgcag gagacatttg	1320
cggagtacac ggcaagaaca gcatttgaga gaccgttgct aagtgagtg gcttatgctg	1380
aaaaagttgt gaattttgag agggagatgt ttgagcggca gcacaattgg gttataaaga	1440
caatggatag aggagagcct tcaccgggta gggatgagta tgctctggtt atattctctc	1500
aagatagtgt ctcttacctt gagtcactcg atatgatgtc aggcgaggtat atataaacat	1560
tatcagacac ttatggtttt gtttttctcg ttaagtatcc gacctgcacc tcttctgaat	1620
tttggtctatc ttttctgcag gaggatcgtg agaatatatt gcgagctaga gaaacggaa	1680
aagctgtctt gactagccct tttaggttgt tggaaactca ccatctcgga gttgtgttga	1740
cattccctgt ctacaagtct tctcttctcg aaaatccgac tgtcgaagag cgtattgcag	1800
ccactgcagg gtaaaagtcc catctgatta ctggagcaaa cttttgcttg tgatcatgta	1860
aaaagagatt ttcgagattt catgatttgt ttatgttgt gatcatgtaa aggaaagagg	1920
ttttttgaga tttcaagatt ggtttgtgtg taaagtaaag attgatgctt gaatttgaaa	1980
tgcagggtacc ttggtgtgtc gtttgatgtg gagtctctag tcgagaattt acttggtcag	2040
cttgctggta accaagcaat agttgtgcat gtgtatgata tcaccaatgc atcagatcca	2100
cttgtcatgt atggaatca agatgaagaa gccgacagat ctctctctca tgagagcaag	2160
ctcgattttg gagaccctt caggaaacat aagatgatat gcaggtagca cacagaaaat	2220
cttaaagcta aaatactctc tgatatctaa aaactgatat cttgtttgaa caatttcgag	2280
tgcagggtacc accaaaaggc accaataccg ttgaatgtgc tcacaactgt gccattgttc	2340
tttgcgattg gtttcttggt gggttatata ctgtatggtg cagctatgca catagtaaaa	2400
gtcgaagatg atttccatga aatgcaagag cttaaagttc gacgagaagc tgctgatgtc	2460

-continued

gctaaatcgc agtttcttgc taccgtgtct caccgagatca ggacaccaat gaatggcatt	2520
ctcgggtacta catcttcaaa caaatccag tactgttgcc ttccatcttt ggtttctcaa	2580
actgcctcaa agattctgtt ttaaaaattt tctgcaggaa tgcttgctat gctcctagat	2640
acagaactaa gctcgacaca gagagattac gctcaaaccg ctcaagtatg tggtaaagct	2700
ttgattgcat tgataaatga ggttcttgat cgcgccaaga ttgaagctgg aaagctggag	2760
ttggaatcag taccatttga tatccgttca atattggatg atgtccttcc tctattctct	2820
gaggagtcaa ggaacaaaag cattgaggta agttttcaat ccaaggaata aagatacatc	2880
tatttgagat tgtggctgct aatttcttga ttccaaattt ggcagctcgc ggttttcgtt	2940
tcagacaaag taccagagat agtcaaagga gattcaggga gatttagaca gataatcata	3000
aaccttgttg gaaattcggg taaagttagt ccttatctca atctctgctt ttgttggcag	3060
cttgtgtata tgatttttgt ttgatcaaga tgttgtcttt ttgacagtcc acagagaaa	3120
gacatatctt tgttaaagtc catcttgccg aacaatcaaa agatgaatct gaaccgaaaa	3180
atgcattgaa tgggtgagtg tctgaagaaa tgatcgttgt ttccaaacag tcaagttaca	3240
acacattgag cggttacgaa gctgctgatg gtcggaatag ctgggattca ttcaagcatt	3300
tggctcttga ggagcagtc tcatcggagt ttgatatttc tagcaatgtt aggcttatgg	3360
tttcaatcga agacacgggt attggaatcc ctttagttgc gcaaggccgt gtgtttatgc	3420
cgtttatgca agcagatagc tcgacttcaa gaaactatgg aggtactggg attggtttga	3480
gtataagcaa gtgtcttgtt gaacttatgc gtggtcagat aaatttcata agccggcctc	3540
atattggaag cacgttcttg ttacaggctg ttttagagaa atgcgataaa tgcagtgcga	3600
ttaaccatat gaagaaacct aatgtggaac acttgcttcc tactttttaa ggaatgaaag	3660
ctatagttgt tgatgctaag cctgtagtag ctgctgtgac tagataccat atgaaaagac	3720
tcggaatcaa tgttgatgtc gtgacaagtc tcaaaaccgc tgttggttga gctgctgcgt	3780
ttgaaagaaa cggttctcct ctcccaacaa aaccgcaact tgatatgac tttagtagaga	3840
aagattcatg gatttcaact gaagataatg actcagagat tcgtttattg aattcaagaa	3900
ccaacggaaa cgttcacac aagtctccga aactagctct attcgcaaca aacatcacia	3960
attcggagtt cgacagagct aaatccgcag gatttgcaga tacggtaata atgaaaccgt	4020
taagagcaag catgattggg gcgtgtctgc aacaagttct cgagctgaga aaaacaagac	4080
aacaacatcc agaaggatca tcaccgcgaa ctctcaagag cttgcttaca gggaagaaga	4140
ttcttggtgt tgatgataat atagttaaca ggagagtagc tgcaggagct ctcaagaaat	4200
ttggagcaga agtggtttgt gcagagagtg gtcaagttgc tttgggtttg cttcagattc	4260
cacacacttt cgatgcttgc ttcatggata ttcaaatgcc acagatggac gggtaactaa	4320
atcacaacta atactatttt tggattattc ggtttttgaa ttgttcggta tgaaaaattt	4380
cccgccaaat tataactgac ttggtttggg tcggattcgg ttaaaatctt ttataaactt	4440
cattgattct cagatttgaa gcaactcgtc agataagaat gatggagaag gaaactaaag	4500
agaagacaaa tctcgaatgg catttaccga ttctagcgat gactgcggat gtgatacacg	4560
cgacctacga ggaatgtctg aaaagtggga tggatgggta cgtctccaaa ccttttgaag	4620
aagagaatct ctataaatcc gttgccaaat cattcaaacc taatcctatc tcaccttcgt	4680
cgtaa	4685

-continued

<211> LENGTH: 2000

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 10

```

taatttgaat atttatattc aatttcatac atattatattt aaaatttata attttatata      60
ctaaaataat taataacata ttagatTTTT taatgtgtat gatgtgattc aatttttaat      120
acaagttgga aaatcctaga attgacgggt tttaatcaat attaatacaa gatggtatag      180
tacaggtgaa actctaaaat taacaatcaa aatacaagta tataatttta aacattaaac      240
aatataaata atatcaacaa aacatataca actagcagtt tactatacat ttaaaatcaa      300
ataaaataat atcatatatt tgtaatcaac tttaacaataa aagtttttag tttaactaaa      360
atataagcct acctgctgta taatgcaagt taaaattata aataagcaga ttaccatcac      420
gatggatgta ccttctcaca tacacaccta tcaaaaactt aaaacctttc aaaacctcct      480
ggatatattc aattaggatt cttcattcat agaatcatta ttccaagggt tttgggatgt      540
cttatgaaat gattttggca gaagaggatg taaatttttt tgaataaaaag gtattcaaca      600
aaacatatgt tttattggga ttttttagat aattaaatca gagaataggg ctgggcatat      660
gagtaaccca ttgaggttcg ggtagaaccc gttcgggttt gggattaatg ggtatatgat      720
ttactaccta atagaataat tctaaatatt cggttcgggt cgggtccagt tgggtttggg      780
tcagtttgga taaaagttt caggccctaa aaatactcaa aaaaatagaa acgagtattt      840
tgtatgtaga tagtgcgtct ctaaacaaaa aaaactttta aatttattaa gttttaataa      900
atttaattat atttgaatat attttactaa attttgacaa atataacaaa taattttgat      960
aaacataaca aacataacaa agtatacaat gtaattttata caaaatatta atgcccggtgc    1020
tatacagcac gggttatgat ctagtccctaa tatattttca aagtcaaata caattttaaa    1080
cagaacttct tacataaaaa taatcagaat cataaatgta actgggagac aaagatacat    1140
aaactgataa aaagaaataa gatgtacata tagaatcaaa tcaataacca cttctcgtaa    1200
aaccactca tcgttatatt ttatacaaga taagataaga cttaagagta gatgttaaac    1260
tatgctaattg aagcaattta aaaacaccaa aaagacccaa caaaaaacaa ataacaatga    1320
gtaggtttta ggagaagcat gtattgattc ttggtatagt tgagtttatt tatattttca    1380
ttagccaatc actttttcta atttaactaa aaatgttata ataccttccc cacctagatg    1440
atgagataat catggaccca aaaacctcag tttttaaaaa tgtttattca ccaaagaca    1500
aaaacaataa taataatcaa aagtagagta aaagaagaga gaagtgacat tctttttcct    1560
tagctcataa tttctctaac aaatgcttct tttatccaaa ctctgatcat tgagtccagt    1620
tgggtaatga gagacaccag ctgggtttct tttgtttaag ctcaaaaaca tctaaaacag    1680
agaaagcttt gtcatttttc tttaccctga aaccatgaaa gaagataaag ctttgcctct    1740
tttctctttt cgtttctttt actctcaaca atgtcttctt tctcccattg caacaaccac    1800
tgggttttgt tcatcttgct ctcttgctgc ttaagtaaga aaaaagaaaa aagctttgctg    1860
tttgggacta tgtcacagct tctgttctt cttgtggatg gtttcaagaa ctcaactatg    1920
agattgttgt tgatagagtt ttagtttttg gtttgaagag atctgaattt cttcagctga    1980
atctgagatt aggagtcgaa                                2000

```

<210> SEQ ID NO 11

<211> LENGTH: 2000

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

-continued

<400> SEQUENCE: 11

```

tctggatcca tctctttctt cgtcttcctt tttgatcctt agctcttgcg tegtgtttt    60
tcaacaggcg agagaggatc gagagggtta gataatttac tgaaaaatta tacgaatgga    120
ccaagactag agatactgaa atgagtgttt gacgatactt taaatgggccc ttatatTTTT    180
caagttgggc ccgtaatat atgggacaat caacagtata gctcaattgg taaattgact    240
aacttcattt gattcctaca ccatttaaag atttttaato aaattctgat ataactaaga    300
taatatttta aagcatttaa ctaacatgga aaattatgaa atgaaatatt aattttggat    360
agaaaattta aaacatttat ctttgtgaaa taaaaagaaa gtctcttata atataacttg    420
taaagtaato ctttaaatc taaagatct aaaacaaaat gagttaaatc ctatgtctat    480
gtgttgaatc tggaagatcg aggtggaaga aaaaggatca tgattcatgg gtgagagaac    540
aaccgttgat tgggtgtgtt caactttcaa gcaatcccat tttaaaaaa taaatttatg    600
taagcaattg caaacaaat gaaatgatta tgctttaaga tatcgacacg tgtaaacacg    660
atgacgtacc aagaaaataa atgattctgt attttgtttt agccatttgt tttttcttta    720
aaggaaagtg tatgttgaaa ttgaaaaaac aaaccgagtt atatatatag agttatttga    780
ctaaagaagg aagctttggt taaactaact tgtataatc ttttgtgtag tttagttttg    840
ataaaatggc caaacaattt tggtcacttt accaactact gcaaaaaaac cattttcata    900
ataataaatt gtaacacgcc ttttattttt attattggag ggtgagaact gagaccatat    960
atgcctttgt tccctctctg tgggtccctt tagccttttt tttctctttg cctgcgcctg    1020
ttagtgtttt ttcttttata tataaataga ggtaagttat tttaccaatc atttggtgtt    1080
attaatagta gtattaatga ggttctaaac ttgtttataa aaatcagatc ttactaaaca    1140
aactcaaaga atctcaacta tgaacatttt tcttctagaa tcgagtttta acatcatcat    1200
cagcaggata taattttttg tgtttgaatt ttatctgtat tttttttagg agtacataag    1260
cagcttaaga ttaaaatata tattaatgga agtatgaatg aaaaaagaaa agaaagataa    1320
tcagataata gagagagaga gagagagaag gaagcagaca tttgtcttcc accatattct    1380
tctcatcttt tccgatgata gattcttctt cttcttcttc ttctactact acttctgtct    1440
ctctttctta agtttctcca ttgataatt cactcaagtc tccagctcct cttcttcttc    1500
tcgttgcttt atcgtctctc ttctactcat catcatcaca ctcaaagatt tctgttgat    1560
tggagcaagc tctagcgaat ttctgtactc ccttgtaga attttagttt cctgttttag    1620
atttcgagga atctcgtctt ggaaacttta agctcctgtt caaagtttct acctttgttt    1680
attttgtttt tgttttcttc tctcttatct ttcgtcatct ttgattaagt aaccactttg    1740
atctctcaat ttcagccttt gggttgtcta atctgacacg ttctctgttc ttagctttcc    1800
ttttctcttc tcgtttgttg gaagcctaca aggattaatt attaaatacc caaccccaaa    1860
aaattcttcc tttttgatc tctgattcgt ccttttttgt acatgggttc tgtttggttg    1920
ttagagcttc ttgctgagtt gaattgagct ggttgcaatc actctgttcc ttgggggttg    1980
atcgtgtatt caagtgggtg                                     2000

```

<210> SEQ ID NO 12

<211> LENGTH: 2000

<212> TYPE: DNA

<213> ORGANISM: Arabidopsis thaliana

<400> SEQUENCE: 12

-continued

```

ccaattcacg ttaaatctat ctcttgctct tgettcctcc aaaaaaaaaa aaaaatcatt    60
cccagatcca tcgatatgaa attgtataga aaaaatggta ttcgatccaa gtttattgtc    120
ttctatTTTT cttagggttaa ttccacttta ttccagattc attgtttgtt ttttctttct    180
cggaagagca caatgtgagt ttcactggcc tctgttataa acatatatag aaatctgtaa    240
caaaaatcat tactaaaatt ctgtgacatg tgcagcgatc aaagaatcaa tagcggaaaa    300
agaaactaca ctgcattcat ctatgactga aagcttctga tcaagccatg aaattaaggt    360
atcccaaaca cgtatcttct ctatgtttat caatcttgct ttaagttcta attctgcata    420
tttcaaagga accatacaag tgttcctaaa atccatttga atattcaaaa acttctctca    480
aatatcatgt agttatagaa gctactgtct ctaagcgcac gagagaaagc tacacaaccc    540
acgtcagttt ccatctacac atataaggta ataataatat tttcatgtat ctttaataat    600
agctctatgt ttttttctgt atttttcatt ataaaactca taactatggt atcatttaat    660
atggtactaa tttaatggga ttgatttact attgcctcaa acatgtaata atttaatgat    720
tttttgTTTT taacgtTTTT agaaattcat gagcatttta aatttggttg taggtcataa    780
caatttgcta ttacaaaaaa aagaaacact ctaaaataa taaaaaatag tttaccgtat    840
aatactagta gtaataaat aatttgattg ttattcataa attttgaatt ctaaaatctc    900
ctgaatcaac tcatgcaatt gtcttaagaa ttacacgtgg ataaatcatg ggcttatgag    960
tcaggcccat ttaaccgggg tattttcgta gttaagagac tagaatgggt ggtatttcag   1020
gtaaaaggtc tatggggcca gatctgcgct ttgtcgcgat gtcattatcg ccaaagatat   1080
gcgatagcga ctctcgtaca aagtctctca ctcacctata ttttttgTTT tcttatattt   1140
caacaaaaaa acgTTTTatt ttcttttggt tgtaagtaaa aaacaaaaac aaacgTTTT   1200
atttctaaag ttcagaaaaa ttattttatac caaggaaaaa atagataata aattttgaga   1260
agttggtgac tatatattac ttcacttatt caagaaattt aaacatggta aatgttactt   1320
taaatgttaa atgatgtata agaaatgtaa tgaaatgtaa taaatgtagt tttaaagatg   1380
ttttaattag taagacaaac ctagttagtg tcacaataat tatatttttt tttttgtcat   1440
ccaaaattat taaagctcaa gtaaaccaat cctgagggat attatttaca aatgtgatat   1500
gatgcggttc ggtgcggatc ttccgcgcca aattatacgc ttttatatta gcattataaa   1560
aaattataga taaagagaag tttgtgaatt cttcattgtc gctttgcaat ttctctaaat   1620
acacagtaaa taccgacaat tcggttagag aaaatatatc tatttcgtat aataatgtta   1680
actttgagga gattttggtt aaaataataa cttttgttgg atggatcata tcatgagcca   1740
ttaagaaaaa gtccaaaact tttcttcttc aaagttggac tcaagttaga aaaagaaaaa   1800
agagctagag agatataaaa atgaaaagaa agttcatggc aaaaaactga tatagacaga   1860
gacacagaga gagagaaacg tatctgaaga aaatctaaaa aattcgattc aatttttttc   1920
ttacttttaa aagcaaaaaa tctcactaaa acaaaagaag aagaaagaag aaagaaaatg   1980
gaatacctac atttgaagtg                                     2000

```

```

<210> SEQ ID NO 13
<211> LENGTH: 2000
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Modified AtCHK2 promoter with SNP

<400> SEQUENCE: 13

```

```

taatttgaat atttatattc aatttcatac atattatttt aaaatttata attttatata    60

```

-continued

ctaaaaataat taataacata ttagatTTTT taatgtgtat gatgtgattc aatttttaat	120
acaagttgga aaatcctaga attgacgggt ttaaatcaat attaatacaa gatggtatag	180
tacaggtgaa actctaaaat taacaatcaa aatacaagta tataatttta aacattaaac	240
aatataaata atatcaacaa aacatatata actagcagtt tactatacat ttaaaatcaa	300
ataaaataat atcatatatt tgtaatcaac tttaacaataa aagtttttag tttaactaaa	360
atataagcct acctgctgta taatgcaagt taaaattata aataagcaga ttaccatcac	420
gatggatgta ccttctcaca tacacaccta tcaaaaactt aaaacctttc aaaacctctt	480
ggtatattca aattaggatt cttcattcat agaatcatta ttccaagggt ttgggatgt	540
cttatgaaat gattttggca gaagaggatg taaatttttt tgaataaaag gtattcaaca	600
aaacatatgt ttatttggga ttttttagat aattaaatca gagaataggg ctgggcataat	660
gagtaacca ttcgagttcg ggtagaaccg gttcgggttt gggattaatg ggtatatgat	720
ttactaccta atagaataat tctaaatatt cgggtcgggt cgggtccagt tgggtttggg	780
tcagtttgga taaaaagttt caggccctaa aaatactcaa aaaaatagaa acgagtattt	840
tgtagttaga tagtgcgtct ctaaacaaaa aaaactttta aatttattaa gttttaataa	900
atttaattat atttgaatat attttactaa attttgacaa atataacaaa taattttgat	960
aaacataaca aacataacaa agtatataat gtaatttata caaaatatta atgccgtgc	1020
tatacagcac gggttatgat ctagtcctaa tatattttca aagtcaaata caattttaaa	1080
cagaacttct tacataaaaa taatcagaat cataaatgta actgggagac aaagatacat	1140
aaactgataa aaagaaataa gatgtacata tagaatcaaa tcaataccca cttctcgtaa	1200
aaaccactca tcgttatatt ttatacaaga taagataaga cttaagagta gatgttaaac	1260
tatgctaata aagcaattta aaaacaccaa aaaggccaaa caaaaaacaa ataacaatga	1320
gtaggtttta ggagaagcat gtattgattc ttggtatagt tgagtttatt tatattttca	1380
ttagccaatc actttttcta atttaactaa aaatgttata atacctccc cacctagatg	1440
atgagataat catggaccca aaaacctcag tttttaaaaa tgttttattca ccaaaagaca	1500
aaaacaataa taataatcaa aagtagagta aaagaagaga gaagtgcacat tctttttctt	1560
tagctcataa tttctctaac aaatgcttct ttatccaaa ctctgatcat tgagtccagt	1620
tgggtaatga gagacaccag ctgggtttct tttgtttaag ctcaaaaaca tctaaaacag	1680
agaaagcttt gtcatttttc ttaccctga aaccatgaaa gaagataaag ctttgcctt	1740
tttctctttt cgtttctttt actctcaaca atgtcttctt tctccattg caacaaccac	1800
tgggtttggt tcactctgtc ctcttgctgc ttaagtaaga aaaaagaaaa aagctttgct	1860
tttgggacta tgtcacagct tcttgttctt cttgtggatg gtttcaagaa ctcaactatg	1920
agattgttgt tgatagagtt ttagtttttg gtttgaagag atctgaattt cttcagctga	1980
atctgagatt aggagtcgaa	2000

<210> SEQ ID NO 14

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: pAtCHK2_F

<400> SEQUENCE: 14

taatttgaat atttatattc aatttcatac atat

34

-continued

```

<210> SEQ ID NO 15
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: pAtCHK2_R

<400> SEQUENCE: 15

ttcgactcct aatctcagat tca                                     23

<210> SEQ ID NO 16
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: pAtCHK4_F

<400> SEQUENCE: 16

ccaattcacg ttaaattctat ctcttg                                 26

<210> SEQ ID NO 17
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: pAtCHK4_R

<400> SEQUENCE: 17

cacttcaaat gtaggtattc catttt                                   26

<210> SEQ ID NO 18
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: AtCHK2_F_CDS

<400> SEQUENCE: 18

atgtctataa cttgtgagct cttgaa                                  26

<210> SEQ ID NO 19
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: AtCHK2_R_CDS

<400> SEQUENCE: 19

ttaacaaggt tcaaagaatc ttgc                                    24

<210> SEQ ID NO 20
<211> LENGTH: 55
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: AtCHK2_F_CDS_attB1F

<400> SEQUENCE: 20

ggggacaagt ttgtacaaaa aagcaggcta tgtctataac ttgtgagctc ttgaa 55

<210> SEQ ID NO 21
<211> LENGTH: 53
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: AtCHK2_R_CDS_attB2R

```

-continued

<400> SEQUENCE: 21

ggggaccact ttgtacaaga aagctggggt taacaagggt caaagaatct tgc 53

<210> SEQ ID NO 22

<211> LENGTH: 32

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: AtCHK4_F_CDS

<400> SEQUENCE: 22

atgagaagag attttgtgta taataataat gc 32

<210> SEQ ID NO 23

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: AtCHK4_R_CDS

<400> SEQUENCE: 23

ttacgacgaa ggtgagatag ga 22

<210> SEQ ID NO 24

<211> LENGTH: 61

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: AtCHK4_F_CDS_attB1F

<400> SEQUENCE: 24

ggggacaagt ttgtacaaaa aagcaggcta tgagaagaga ttttgtgtat aataataatg 60

c 61

<210> SEQ ID NO 25

<211> LENGTH: 51

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: AtCHK4_R_CDS_attB2R

<400> SEQUENCE: 25

ggggaccact ttgtacaaga aagctggggt tacgacgaag gtgagatagg a 51

<210> SEQ ID NO 26

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Q_CHK4_F

<400> SEQUENCE: 26

taagcctgtt agagctgctg tg 22

<210> SEQ ID NO 27

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Q_CHK4_R

<400> SEQUENCE: 27

tctttcaaac gcagcagctg 20

-continued

<210> SEQ ID NO 28
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Q_CHK2_F

<400> SEQUENCE: 28

gctacagaag aacagcgtgt tg 22

<210> SEQ ID NO 29
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Q_CHK2_R

<400> SEQUENCE: 29

tgtttgctgg caagttgggtg 20

<210> SEQ ID NO 30
<211> LENGTH: 3708
<212> TYPE: DNA
<213> ORGANISM: Solanum lycopersicum

<400> SEQUENCE: 30

atgtccttaa attgcaaatt gttgggcatg aatgggagct tctcttcaag ttttaggttg 60
aagaaggcca gagagtcttt gccaggacca agttgtgggt ggaaatggca aaggaagctt 120
ttgtttttat ggctcatttt ctttgaatt ggattttttt ggttggtcat tagcttaaatt 180
ggtgatgttt atagttggaa gaaggaggcg tcagaactaa atgaagataa atcttatttt 240
cttcttgaac gtttcaatgt cagcaaggaa caaattcaag atttagctac cttattcttt 300
gaaaaagatc agatttcac cttggaatgc tcaaaagttc ataagcacga aatgcctatg 360
agtactacca ttacttgct tctcaagggtg ctaggttcgg agagtctgga atatgaacag 420
catgagatgg tttagacaa tattgaagca gaaggccagt gccagttcc agatgaggag 480
acactgaaaa atagtgcac ttacttagat gaaaagtcgt taccatttgt ttgcatcgt 540
ttatcttctc taatctcaac agatcctaag ttctttgaaa agaaagcatc acaataaga 600
gaagttggga attttaacc agagcattgt gatagtattg ccttctgttt cacaaaaacta 660
tgctggtggg tccttcttgg gattgtgatc agttggaaga tacttctgtt atgtgcaaaa 720
ggtggggaac atcaacagaa cggattcatt cagcagcaac cactctctca acaacttcat 780
ccactacagc agttgcagca gcagcaagtt cagatttcct gtcgaactgg tggaaagtgg 840
aggaaaaagg ctcttggtat ctttgtcttt ggtgggggtga ctctagccat ctggttgat 900
ttgtacctga gtgcagacat tgcattgagg aggaaagaaa cactgacaag catgtgtgac 960
gaacgagcgc gaatgttgca ggaccagttc aacgtaagca tgaaccatgt tcatgcattg 1020
gctattcttg tttccacatt tcaccatgga aaacaacctt ctgcaataga ccagaaaact 1080
tttgaagaat atactgagag aacagcttcc gaaaggccac ttacaagtgg tgttgccat 1140
gctttaagag ttcgtcactc agagagagaa gagtttgaga agctgcatgg gtggactata 1200
aagaaaaatgg aagcagagga ccaaaactta ggacatgatt atatccctgc aaacttggat 1260
cctgctcctg atcaagatga atatgcgcct gtcatatattt cacaacaaac agtttcccat 1320
atcgtctcta ttgatatgat gtctggaaag gaagatcgtg agaacatttt gcgagcgagg 1380
gcttctggca aggggggttt gacgtcaccg tttaagctat tgaaatccaa tcacctgggt 1440

-continued

gttggtccta cattedcggt ctataatact catctccttc cttatgctac cccggtggac	1500
cgcacaaatg ctactgttgg gtacattggt gcttcttacg atgttccatc attagttgaa	1560
aagcttctgc agcagcttgc gagcaaacaa actattgtgg taaatgttta tgatacaact	1620
aacaagtttg ctccaattaa aatgtatggc atggatgaga acgacacagg attacttcac	1680
gttagcaacc ttgatttttg ggatcctgca aggagccacg agatgcattg caggttcaag	1740
cagaaacctt ctccaccttg gactgcaata actctatctg tcggagtcct tgttatcact	1800
ctgcttattg gtcatactct ccatgctgcc atcaacagaa ttgctgaagt tgagggtcag	1860
tatcaggaaa tgatggagct caaacatcgt gctgaggctg cagatatagc aaaatcacag	1920
tttcttgcaa cggtttctca tgaatcagg actccaatga atggtgtttt aggcatgctt	1980
cagatgctca tggacacgaa ccttgacctc acgcaactgg attatgcaca aactgctctt	2040
agtagtggga atgaactgat atcattgatt aatgaggtgc tggatcaggc taagattgaa	2100
tcaggaaggc ttgagctgga ggtgttctct tttgatctcc gtgctgaact tgataatgtt	2160
tcacgctctc tctcgggaaa atctcataaa aaagggttg agttggctgt ttatgtttct	2220
gacctagtcc cagaagttgt tatcggagat tcaggacggt tcaagcagat aattacgaat	2280
ctagtggaa actcagttaa gttcacaaat gacaaaggc acatatttgt cacagtacat	2340
ttagctgatg aagtgaggaa cctcatgat gtgacggatg aagtcttgaa acaaagctta	2400
acctttgttc aagaaaggtc aaatgcctct tggaaacct tcagtgggtt tctgttagtt	2460
gacagatggc aaagtgtgca aaagtttgat aggtctgagc gcacagagga agaagtggga	2520
aagatcaagt tgtagtgac catagaagac actggtgtgg gaattcctct tgaagcacag	2580
gcccgcattt tcacaccttt tatgcaggct gacagttcaa catctcgaa atatggcgga	2640
acagggatag gattgagcat tagcaaacgt ttggttagacc tcatgggggg agaaataggc	2700
tttttcagtg aacctggcag agggagtaca ttttcattca ctgcagcctt cacaagagga	2760
gaagaaggtt ctttagagcg taaatggaaa cagtatgac cagcttttcc agaatttcgt	2820
gggttaagag cattggtggt tgatgataaa agcattaggg cagtggtcac aagataccat	2880
ttgcagagac tgggtatatg tgtcaacata acttcacaa tgcgttcagc atgctcatat	2940
ctctctaatt actccaatac tagttcgttg gaacatttag ccgtgggttt tgttgaccaa	3000
gatagttggg ataagaagc tctcttgca ctaagtaata tgttgaaaga ggtcagacca	3060
aatggctcaa ctactacttt ggggaagcct ccaaaaatct gtctcttatg tatgaacttt	3120
atggagagag ctgagcttaa aatagctgga attgtggatc acgtgttaac aaagcctgtg	3180
cggttgagtg cggttgataac atgtgttcaa gaagccattg gctgccaaa taagaagcaa	3240
gtgactcaac ctacaactct tggagtttg ctgacaggaa agcatatatt ggtggtagat	3300
gacaataatg taaacaggag agtggcagaa ggtgctctaa agaagtatgg ggcgatagtg	3360
acctgtgtag acagtggaaa ggtgctttg acacatctta atccaccaca caagtttgat	3420
gcttggttta tggacctcca aatgcctgaa atggatgggt ttgaagcgac acgacaaata	3480
cgtaatctag aaaacaaata taatgaaaag gtcaattctg gtgagttatt ttccggcatg	3540
tctgccagag tggctcattg gcacacacca atattagcaa tgacagcaga tgtaattcaa	3600
gcaacaaatg aagagtgcgt gaagtgcggt atggatgatt atgtatcaaa accgtttgaa	3660
aaagggcagc tttattcaac agtggcacgc ttctttgggt caggttga	3708

-continued

```

<211> LENGTH: 1235
<212> TYPE: PRT
<213> ORGANISM: Solanum lycopersicum

<400> SEQUENCE: 31

Met Ser Leu Asn Cys Lys Leu Leu Gly Met Asn Gly Ser Phe Ser Ser
 1          5          10          15

Ser Phe Arg Leu Lys Lys Ala Arg Glu Ser Leu Pro Gly Pro Ser Cys
 20          25          30

Gly Trp Lys Trp Gln Arg Lys Leu Leu Phe Leu Trp Leu Ile Phe Phe
 35          40          45

Gly Ile Gly Phe Phe Trp Leu Val Ile Ser Leu Asn Gly Asp Val Tyr
 50          55          60

Ser Trp Lys Lys Glu Ala Ser Glu Leu Asn Glu Asp Lys Ser Tyr Phe
 65          70          75          80

Leu Leu Glu Arg Phe Asn Val Ser Lys Glu Gln Ile Gln Asp Leu Ala
 85          90          95

Thr Leu Phe Phe Glu Lys Asp Gln Ile Ser Ser Leu Glu Cys Ser Lys
100          105          110

Val His Lys His Glu Met Pro Met Ser Thr Thr Ile Thr Cys Leu Leu
115          120          125

Lys Val Leu Gly Ser Glu Ser Leu Glu Tyr Glu Gln His Glu Met Val
130          135          140

Val Asp Asn Ile Glu Ala Glu Gly Gln Cys Pro Val Pro Asp Glu Glu
145          150          155          160

Thr Leu Lys Asn Ser Asp Ile Ser Leu Asp Glu Lys Ser Leu Pro Phe
165          170          175

Val Leu His Arg Leu Ser Ser Leu Ile Ser Thr Asp Pro Lys Phe Phe
180          185          190

Glu Lys Lys Ala Ser Gln Ile Arg Glu Val Gly Asn Phe Asn Pro Glu
195          200          205

His Cys Asp Ser Ile Ala Phe Cys Phe Thr Lys Leu Cys Trp Trp Val
210          215          220

Leu Leu Gly Ile Val Ile Ser Trp Lys Ile Leu Leu Cys Ala Lys
225          230          235          240

Gly Gly Glu His Gln Gln Asn Gly Phe Ile Gln Gln Gln Pro Leu Ser
245          250          255

Gln Gln Leu His Pro Leu Gln Gln Leu Gln Gln Gln Val Gln Ile
260          265          270

Ser Cys Arg Thr Gly Gly Lys Trp Arg Lys Lys Ala Leu Val Ile Phe
275          280          285

Val Phe Gly Gly Val Thr Leu Ala Ile Trp Leu Tyr Leu Tyr Leu Ser
290          295          300

Ala Asp Ile Ala Leu Arg Arg Lys Glu Thr Leu Thr Ser Met Cys Asp
305          310          315          320

Glu Arg Ala Arg Met Leu Gln Asp Gln Phe Asn Val Ser Met Asn His
325          330          335

Val His Ala Leu Ala Ile Leu Val Ser Thr Phe His His Gly Lys Gln
340          345          350

Pro Ser Ala Ile Asp Gln Lys Thr Phe Glu Glu Tyr Thr Glu Arg Thr
355          360          365

Ala Phe Glu Arg Pro Leu Thr Ser Gly Val Ala Tyr Ala Leu Arg Val
370          375          380

Arg His Ser Glu Arg Glu Glu Phe Glu Lys Leu His Gly Trp Thr Ile

```

-continued

385					390						395					400
Lys	Lys	Met	Glu	Ala	Glu	Asp	Gln	Thr	Leu	Gly	His	Asp	Tyr	Ile	Pro	
				405					410					415		
Ala	Asn	Leu	Asp	Pro	Ala	Pro	Asp	Gln	Asp	Glu	Tyr	Ala	Pro	Val	Ile	
			420					425					430			
Phe	Ser	Gln	Gln	Thr	Val	Ser	His	Ile	Val	Ser	Ile	Asp	Met	Met	Ser	
		435					440					445				
Gly	Lys	Glu	Asp	Arg	Glu	Asn	Ile	Leu	Arg	Ala	Arg	Ala	Ser	Gly	Lys	
	450					455					460					
Gly	Val	Leu	Thr	Ser	Pro	Phe	Lys	Leu	Leu	Lys	Ser	Asn	His	Leu	Gly	
465					470					475				480		
Val	Val	Leu	Thr	Phe	Ala	Val	Tyr	Asn	Thr	His	Leu	Leu	Pro	Tyr	Ala	
			485						490					495		
Thr	Pro	Val	Asp	Arg	Ile	Asn	Ala	Thr	Val	Gly	Tyr	Ile	Gly	Ala	Ser	
			500					505					510			
Tyr	Asp	Val	Pro	Ser	Leu	Val	Glu	Lys	Leu	Leu	Gln	Gln	Leu	Ala	Ser	
	515						520					525				
Lys	Gln	Thr	Ile	Val	Val	Asn	Val	Tyr	Asp	Thr	Thr	Asn	Lys	Phe	Ala	
	530					535					540					
Pro	Ile	Lys	Met	Tyr	Gly	Met	Asp	Glu	Asn	Asp	Thr	Gly	Leu	Leu	His	
545					550					555					560	
Val	Ser	Asn	Leu	Asp	Phe	Gly	Asp	Pro	Ala	Arg	Ser	His	Glu	Met	His	
			565						570					575		
Cys	Arg	Phe	Lys	Gln	Lys	Pro	Ser	Pro	Pro	Trp	Thr	Ala	Ile	Thr	Leu	
		580						585					590			
Ser	Val	Gly	Val	Leu	Val	Ile	Thr	Leu	Leu	Ile	Gly	His	Ile	Phe	His	
	595						600					605				
Ala	Ala	Ile	Asn	Arg	Ile	Ala	Glu	Val	Glu	Gly	Gln	Tyr	Gln	Glu	Met	
	610					615					620					
Met	Glu	Leu	Lys	His	Arg	Ala	Glu	Ala	Ala	Asp	Ile	Ala	Lys	Ser	Gln	
625					630					635					640	
Phe	Leu	Ala	Thr	Val	Ser	His	Glu	Ile	Arg	Thr	Pro	Met	Asn	Gly	Val	
			645						650					655		
Leu	Gly	Met	Leu	Gln	Met	Leu	Met	Asp	Thr	Asn	Leu	Asp	Leu	Thr	Gln	
		660						665					670			
Leu	Asp	Tyr	Ala	Gln	Thr	Ala	Leu	Ser	Ser	Gly	Asn	Glu	Leu	Ile	Ser	
	675						680					685				
Leu	Ile	Asn	Glu	Val	Leu	Asp	Gln	Ala	Lys	Ile	Glu	Ser	Gly	Arg	Leu	
	690					695					700					
Glu	Leu	Glu	Ala	Val	Pro	Phe	Asp	Leu	Arg	Ala	Glu	Leu	Asp	Asn	Val	
705				710						715				720		
Ser	Ser	Leu	Phe	Ser	Gly	Lys	Ser	His	Lys	Lys	Gly	Ile	Glu	Leu	Ala	
			725						730					735		
Val	Tyr	Val	Ser	Asp	Leu	Val	Pro	Glu	Val	Val	Ile	Gly	Asp	Ser	Gly	
			740					745					750			
Arg	Phe	Lys	Gln	Ile	Ile	Thr	Asn	Leu	Val	Gly	Asn	Ser	Val	Lys	Phe	
	755						760					765				
Thr	Asn	Asp	Lys	Gly	His	Ile	Phe	Val	Thr	Val	His	Leu	Ala	Asp	Glu	
	770					775					780					
Val	Arg	Asn	Pro	His	Asp	Val	Thr	Asp	Glu	Val	Leu	Lys	Gln	Ser	Leu	
785					790					795					800	
Thr	Phe	Val	Gln	Glu	Arg	Ser	Asn	Ala	Ser	Trp	Asn	Thr	Phe	Ser	Gly	
			805						810						815	

-continued

Phe Pro Val Val Asp Arg Trp Gln Ser Trp Gln Lys Phe Asp Arg Leu
 820 825 830

Ser Ser Thr Glu Glu Glu Val Gly Lys Ile Lys Leu Leu Val Thr Ile
 835 840 845

Glu Asp Thr Gly Val Gly Ile Pro Leu Glu Ala Gln Ala Arg Ile Phe
 850 855 860

Thr Pro Phe Met Gln Ala Asp Ser Ser Thr Ser Arg Thr Tyr Gly Gly
 865 870 875 880

Thr Gly Ile Gly Leu Ser Ile Ser Lys Arg Leu Val Asp Leu Met Gly
 885 890 895

Gly Glu Ile Gly Phe Phe Ser Glu Pro Gly Arg Gly Ser Thr Phe Ser
 900 905 910

Phe Thr Ala Ala Phe Thr Arg Gly Glu Glu Gly Ser Leu Glu Arg Lys
 915 920 925

Trp Lys Gln Tyr Asp Pro Ala Phe Pro Glu Phe Arg Gly Leu Arg Ala
 930 935 940

Leu Val Val Asp Asp Lys Ser Ile Arg Ala Val Val Thr Arg Tyr His
 945 950 955 960

Leu Gln Arg Leu Gly Ile Cys Val Asn Ile Thr Ser Thr Met Arg Ser
 965 970 975

Ala Cys Ser Tyr Leu Ser Asn Tyr Ser Asn Thr Ser Ser Leu Glu His
 980 985 990

Leu Ala Val Val Phe Val Asp Gln Asp Ser Trp Asp Lys Glu Thr Ser
 995 1000 1005

Leu Ala Leu Ser Asn Met Leu Lys Glu Val Arg Pro Asn Gly Ser
 1010 1015 1020

Thr Thr Thr Leu Gly Lys Pro Pro Lys Ile Cys Leu Leu Cys Met
 1025 1030 1035

Asn Phe Met Glu Arg Ala Glu Leu Lys Ile Ala Gly Ile Val Asp
 1040 1045 1050

His Val Leu Thr Lys Pro Val Arg Leu Ser Ala Leu Ile Thr Cys
 1055 1060 1065

Val Gln Glu Ala Ile Gly Cys Gln Asn Lys Lys Gln Val Thr Gln
 1070 1075 1080

Pro Thr Thr Leu Gly Ser Leu Leu Thr Gly Lys His Ile Leu Val
 1085 1090 1095

Val Asp Asp Asn Asn Val Asn Arg Arg Val Ala Glu Gly Ala Leu
 1100 1105 1110

Lys Lys Tyr Gly Ala Ile Val Thr Cys Val Asp Ser Gly Lys Ala
 1115 1120 1125

Ala Leu Thr His Leu Asn Pro Pro His Lys Phe Asp Ala Cys Phe
 1130 1135 1140

Met Asp Leu Gln Met Pro Glu Met Asp Gly Phe Glu Ala Thr Arg
 1145 1150 1155

Gln Ile Arg Asn Leu Glu Asn Lys Tyr Asn Glu Lys Val Asn Ser
 1160 1165 1170

Gly Glu Leu Phe Ser Gly Met Ser Ala Arg Val Ala His Trp His
 1175 1180 1185

Thr Pro Ile Leu Ala Met Thr Ala Asp Val Ile Gln Ala Thr Asn
 1190 1195 1200

Glu Glu Cys Val Lys Cys Gly Met Asp Asp Tyr Val Ser Lys Pro
 1205 1210 1215

-continued

Phe Glu Lys Gly Gln Leu Tyr Ser Thr Val Ala Arg Phe Phe Gly
1220 1225 1230

Ser Gly
1235

<210> SEQ ID NO 32
<211> LENGTH: 6097
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: expression cassette SlyCHK2

<400> SEQUENCE: 32

```

ggggacaact ttgtatagaa aagttgagtt tatcaatatt ttgagatttt tatatatatta      60
cattagtaat tttcttaata ataactcca gaatttataa ttattttatt aattactcta      120
ccatcgaata ttttttattg gttgttctaa ttcccttgtc actttcaact tgtaattgac      180
caaaagaaat attttagctt agataagttt cttacatcat atttcccttc tttttttata      240
tgctttattt gtataggagt ttattatttt ttatatagtg tttggagctt atattaaatt      300
tggatcgcac actgtaaaga ttcatataaa gtgatgcttc gaacaagttt ttctctatat      360
ctaagctcga aattattttt ctccatacac gaaactcaaa tccgaaatat ttaattaaag      420
tgaatcagtt tcaccccaat accataacca taaatcatgt tgattagttg ctatcaactt      480
ctttcttcat gttcaaatgt tttttttaat gaaaaggat taataattca catgcattca      540
tcaactagct taatttaatt ttattttgat aaataattca ataacaattt ctactacaat      600
tgactcattc aaggatttta catataatgc aaactcacct atatgtattt attaagcaca      660
caatagtgtt ttagaatcac aagaaatcat ataatgatgg tcaaactaat catgtctttt      720
taatattaaa agtgattagt tttatctaac actattaaac ataccaattc atttattatt      780
atattaaata atattattcc ccttattttt ttttttagta atttaaaaca cattaggggt      840
attattaggg agaagtgggg ttggagagat gaggtgcatt agccagccaa ctgggatgcc      900
cacatcatca attatcttcc aagaataaaag tgtaggataa gcatctcaag tagccggcgc      960
atttttagtac ctagcccttg ctttattatt attttttcac gccgattaat cgcgatctt     1020
ttgtagtctg tacgccgtat tatcaaaagc atagatgctt ttttttttta aaaataaatt     1080
tattttttaa ttttaattta tttttgttct tttctccatg tgaaatgaac aaatacaact     1140
ggctagctct tcccacaacc atcccttttc ttttttttcc tttcttttcc tcctctccca     1200
ttctttatat ttattatgcc tagccatata tttatatttt gttcgtttga tttatgtgat     1260
gtttattgat tagatgttaa aattttaaaa tataattata tttataatat tttaaattat     1320
tcttaaaaaa ataaatttta aatatttttt tatcaaataa gtgagagcaa acatgcgata     1380
gcgttattaa attctgataa gtaagaaaaa atatgaattt aaaaataaac atgtaatata     1440
aattagaata aaaaagtatt attttttcta atcgaaggca atatttcgag gggaaaaaac     1500
ctatatagac ctatgtgaga tgattttgaa ttagtaagaa tccaatgtgc ccgacatcga     1560
atgttttttt tttaaaaata tttacagtta cattatattg aaactgaaaa gggacagtga     1620
aggaatggag ctatcttaaa aagcaaagaa aactatcttt ttattatttt tactatatag     1680
acaagaaaag aaatatggta acaaaaattt gcatattgtg gaagtgtaaa catttgagaa     1740
atgcataatt ttcggggaaa gatgatgac gaaatttttg agatatatct taattaaatt     1800
aaggacatat tattttaaat atatattttt ttgtaatttt gtgcattctt ttgatgtacg     1860
tgacatttaa atattttcca catgcctcaa ttgcgtgaaa ttatggagtg tgtcacgtaa     1920

```

-continued

gacaaaaaat atacaaaata ataataaaat aatttaagaa taataaaaacc tcaatataat	1980
taatgtgtat atatgaaatt ttaatcataa tttaagaggt atttgtatga gcttttctgc	2040
tctgagtggg tttgctctta agttttcaag gcattttttg aagatatgta ggtggattct	2100
attaagatg tccttaaatt gcaaatgtt gggcatgaat gggagcttct cttcaagttt	2160
taggttgaag aaggccagag agtcctttgcc aggaccaagt tgtgggtgga aatggcaaag	2220
gaagcttttg tttttatggc tcatttttctt tggaattgga tttttttggg tggtcattag	2280
cttaaatggt gatgtttata gttggaagaa ggaggcgtca gaactaatg aagataaatc	2340
ttattttctt cttgaacgtt tcaatgtcag caaggaacaa attcaagatt tagctacctt	2400
attctttgaa aaagatcaga tttcatcctt ggaatgtca aaagttcata agcacgaaat	2460
gcctatgagt actaccatta cttgccttct caaggtgcta gggtcgaga gtctggaata	2520
tgaacagcat gagatggttg tagacaatat tgaagcagaa ggcagtgcc cagttccaga	2580
tgaggagaca ctgaaaaata gtgacatttc actagatgaa aagtcgttac catttgtttt	2640
gcatcgttta tcttctctaa tctcaacaga tcttaagtcc tttgaaaaga aagcatcaca	2700
aataagagaa gttgggaatt ttaaccaga gcattgtgat agtattgcct tctgtttcac	2760
aaaactatgc tgggtgggtcc ttcttgggat tgtgatcagt tggaagatac ttctgttatg	2820
tgcaaaaggt ggggaacatc aacagaacgg attcattcag cagcaaccac tctctcaaca	2880
acttcacca ctacagcagt tgcagcagca gcaagttcag atttctgtc gaactgggtg	2940
aaagtggagg aaaaaggctc ttgttatctt tgtctttggg ggggtgactc tagccatctg	3000
gttgtatttg tacctgagtg cagacattgc attgaggagg aaagaaacac tgacaagcat	3060
gtgtgacgaa cgagcgcgaa tgttgcagga ccagttcaac gtaagcatga accatgttca	3120
tgcataggct attcttgttt ccacatttca ccatggaaaa caacctctcg caatagacca	3180
gaaaactttt gaagaatata ctgagagAAC agctttcgaa agggcaactta caagtgggtg	3240
tgctatgct ttaagagttc gtcactcaga gagagaagag ttgagaagc tgcatgggtg	3300
gactataaag aaaatggaag cagaggacca aacttttaga catgattata tccctgcaaa	3360
cttggatcct gctcctgac aagatgaata tgcgcctgtc atattttcac aacaaacagt	3420
ttcccatatc gtctctattg atatgatgtc tggaaaggaa gatcgtgaga acattttgcg	3480
agcgagggtc tctggcaagg gggttttgac gtcaccgttt aagctattga aatccaatca	3540
cctgggtgtt gttcttacat ttgcggtcta taatactcat ctcttctctt atgctacccc	3600
ggtggaccgc atcaatgcta ctgttgggta cattgggtgct tcttacgatg ttccatcatt	3660
agttgaaaag cttctgcagc agcttgcgag caaacaaact attgtggtaa atgtttatga	3720
tacaactaac aagtttgcct caattaaaat gtatggcatg gatgagaacg acacaggatt	3780
acttcacgtt agcaaccttg attttgggga tcttgcaagg agccacgaga tgcattgcag	3840
gttcaagcag aaacctctc caccctggac tgcaataact ctatctgtcg gagtccttgt	3900
tatcactctg cttattggtc atatcttcca tgctgccatc aacagaattg ctgaagttga	3960
gggtcagtat caggaaatga tggagctcaa acatcgtgct gaggctgcag atatagcaaa	4020
atcacagttt cttgcaacgg tttctcatga aatcaggact ccaatgaatg gtgttttagg	4080
catgcttcag atgctcatgg acacgaacct tgacctaacg caactggatt atgcacaaac	4140
tgctcttagt agtgggaatg aactgatatc attgattaat gaggtgctgg atcaggctaa	4200
gattgaatca ggaaggcttg agctggaggc tgttctcttt gatctccgtg ctgaacttga	4260

-continued

taatgtttca tgcgtcttct cgggaaaatc tcataaaaaa gggattgagc agttggctgt	4320
ttatgtttct gacctagtc cagaagttgt tatcgagat tcaggacggt tcaagcagat	4380
aattacgaat ctagttagaa actcagttaa gttcacaaat gacaaagggc acatatttgt	4440
cacagtacat ttagctgatg aagttaggaa ccctcatgat gtgacggatg aagtcttgaa	4500
acaaagctta acctttgttc aagaaaggtc aaatgcatct tggaatacct tcagtgggtt	4560
tcctgtagtt gacagatggc aaagttggca aaagtttgat aggctgagca gcacagagga	4620
agaagtggga aagatcaagt tgttagtgac catagaagac actggtgtgg gaattcctct	4680
tgaagcacag gcccgcatct tcacaccttt tatgcaggct gacagttcaa catctcgaa	4740
atatggcgga acagggatag gattgagcat tagcaaacgt ttggtagacc tcatgggggg	4800
agaaataggc tttttcagtg aacctggcag agggagtaca ttttcattca ctgcagcctt	4860
cacaagagga gaagaagggt ctttagagcg taaatggaaa cagtatgatc cagcttttcc	4920
agaatttcgt gggtaagag cattggtggt tgatgataaa agcattaggg cagtggtcac	4980
aagataccat ttgcagagac tgggtatatg tgtcaacata acttcacaa tgcgttcagc	5040
atgctcatat ctctctaatt actccaatac tagttcgttg gaacatttag ccgtggtttt	5100
tgttgaccaa gatagtggg ataagaaac ttctcttgca ctaagtaata tgttgaaaga	5160
ggtcagacca aatggctcaa ctactacttt ggggaagcct ccaaaaatct gtctcttatg	5220
tatgaacttt atggagagag ctgagcttaa aatagctgga attgtggatc acgtgttaac	5280
aaagcctgtg cggttgagtg cggttgataac atgtgttcaa gaagccattg gctgccaaaa	5340
taagaagcaa gtgactcaac ctacaactct tggagtttg ctgacaggaa agcatatatt	5400
ggtggtgatg gacaataatg taaacaggag agtggcagaa ggtgctctaa agaagtatgg	5460
ggcgatagtg acctgtgtag acagtggaaa ggctgctttg acacatctta atccaccaca	5520
caagtttgat gcttgtttta tggacctcca aatgcctgaa atggatgggt ttgaagcgac	5580
acgacaaata cgtaacttag aaaacaaata taatgaaaag gtcaattctg gtgagttatt	5640
ttccggcatg tctgccagag tggctcattg gcacacacca atattagcaa tgacagcaga	5700
tgtaattcaa gcaacaaatg aagagtgcgt gaagtgcggt atggatgatt atgtatcaaa	5760
accgtttgaa aaagggcagc tttattcaac agtggcacgc ttctttgggt cagggtgaga	5820
tcgttcaaac atttggcaat aaagtttctt aagattgaat cctgttgccg gtcttgcgat	5880
gattatcata taatttctgt tgaattacgt taagcatgta ataattaaca tgtaatgcat	5940
gacgttattt atgagatggg tttttatgat tagagtccg caattatata tttataacgc	6000
gatagaaaac aaaatatagc gcgcaaaata ggataaatta tcgcgcgcgg tgtcatctat	6060
gttactagat cgggggacaac tttgtataat aaagttg	6097

The invention claimed is:

1. A method for improving cytokinin-induced de novo shoot formation of a plant cell, comprising:

- (a) increasing or introducing expression of at least one a histidine kinase in the plant cell;
- (b) inducing de novo formation of a number of shoots and elongation of the number of shoots by exposing the plant cell of (a) to a cytokinin-comprising medium;
- (c) dissecting a de novo formed shoot; and
- (d) inducing root formation of the dissected shoot, wherein the number of de novo formed shoots is increased as compared to an identical cell not having an increased or introduced expression of the histidine

kinase, and wherein the histidine kinase has at least 80% sequence identity with SEQ ID NO: 31.

2. The method according to claim 1, wherein the amino acid sequence of the histidine kinase has at least 90% sequence identity with SEQ ID NO: 31.

3. The method according to claim 1, wherein the amino acid sequence of the histidine kinase has at least 95% sequence identity with SEQ ID NO: 31.

4. The method according to claim 1, wherein the histidine kinase is encoded by a nucleotide sequence having at least 50% sequence identity with SEQ ID NO: 30.

5. The method according to claim 1, wherein the expression of the histidine kinase is transiently increased or introduced into the plant cell.

95

6. The method according to claim 1, wherein the expression of the histidine kinase is continuously increased or introduced in the plant cell.

7. The method according to claim 1, wherein the plant cell is from a plant selected from the group consisting of barley, 5
cabbage, canola, cassava, cauliflower, chicory, cotton, cucumber, eggplant, grape, hot pepper, lettuce, maize, melon, oilseed rape, potato, pumpkin, rice, rye, *sorghum*, squash, sugar cane, sugar beet, sunflower, sweet pepper, tomato, water melon, wheat, zucchini, soybean, *chrysanthemum* and *Arabidopsis*. 10

8. The method according to claim 1, wherein the cytokinin is an adenine-type cytokinin.

9. The method according to claim 8, wherein the adenine-type cytokinin is selected from the group consisting of 15
kinetin, zeatin, trans-zeatin, cis-zeatin, dihydrozeatin, 6-benzylaminopurine and 2iP.

10. The method according to claim 1, further comprising:
(a) incubating the plant cell having increased or introduced expression of the histidine kinase in the medium 20
comprising a cytokinin plant hormone; and
(b) allowing the plant cell to regenerate into a plant.

96

11. The method according to claim 5, wherein the medium comprises at least one further plant hormone.

12. The method according to claim 11, wherein the one further plant hormone is an auxin.

13. The method according to claim 10, wherein the plant cell is part of at least one of a multicellular tissue, a callus tissue, a plant organ, an explant, a hypocotyl explant, a stem explant, a cotyledon explant, a root explant, a leaf explant, a flower explant and a meristematic tissue.

14. A method according to claim 10, wherein the concentration of cytokinin in the medium is 100-3000 ng/ml.

15. The method according to claim 14, wherein the concentration of cytokinin in the medium is 200-600 ng/ml.

16. The method according to claim 1, wherein the plant cell is incubated in a medium comprising a cytokinin for at least 5 weeks prior to dissecting a de novo formed shoot.

17. The method according to claim 1, wherein the plant cell is a plant cell of a recalcitrant plant.

* * * * *