



US012397053B2

(12) **United States Patent
Coffin**

(10) **Patent No.: US 12,397,053 B2**

(45) **Date of Patent: *Aug. 26, 2025**

(54) **ENGINEERED VIRUS**

(71) Applicant: **Replimune Limited**, Oxfordshire (GB)

(72) Inventor: **Robert Stuart Coffin**, London (GB)

(73) Assignee: **Replimune Limited**, Oxfordshire (GB)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **18/163,837**

(22) Filed: **Feb. 2, 2023**

(65) **Prior Publication Data**

US 2023/0212531 A1 Jul. 6, 2023

Related U.S. Application Data

(63) Continuation of application No. 17/164,635, filed on Feb. 1, 2021, now Pat. No. 12,049,647, which is a continuation of application No. 16/068,830, filed as application No. PCT/GB2017/050038 on Jan. 9, 2017, now Pat. No. 10,947,513.

(30) **Foreign Application Priority Data**

Jan. 8, 2016	(GB)	1600380
Jan. 8, 2016	(GB)	1600381
Jan. 8, 2016	(GB)	1600382

(51) **Int. Cl.**

A61K 35/763	(2015.01)
A61K 39/00	(2006.01)
A61K 39/395	(2006.01)
A61K 45/06	(2006.01)
A61P 35/00	(2006.01)
C07K 14/005	(2006.01)
C07K 14/535	(2006.01)
C07K 16/28	(2006.01)
C12N 7/00	(2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/39558** (2013.01); **A61K 35/763** (2013.01); **A61K 39/3955** (2013.01); **A61K 45/06** (2013.01); **A61P 35/00** (2018.01); **C07K 14/005** (2013.01); **C07K 14/535** (2013.01); **C07K 16/2818** (2013.01); **C12N 7/00** (2013.01); **A61K 2039/505** (2013.01); **C12N 2710/16621** (2013.01); **C12N 2710/16622** (2013.01); **C12N 2710/16632** (2013.01); **C12N 2710/16633** (2013.01); **C12N 2710/16643** (2013.01); **C12N 2740/13022** (2013.01)

(58) **Field of Classification Search**

CPC G01N 2333/165; G01N 2470/00; G01N 2470/04; G01N 33/543; G01N 2474/00; A61K 35/763; A61K 39/39558; A61K 48/00; A61K 38/193; A61K 2039/5254; A61K 2039/55522; A61K 2039/5256; A61K 2039/505; A61K 39/3955; A61K

45/06; A61P 35/00; A61P 37/02; A61P 37/04; A61P 9/10; A61P 31/20; C07K 14/005; C07K 14/535; C12N 2710/16632; C12N 2710/16643; C12N 15/869; C12N 15/86; C12N 2710/10032; C12N 15/8695; C12N 2710/16633

See application file for complete search history.

(56)

References Cited

U.S. PATENT DOCUMENTS

5,122,458 A	6/1992	Post et al.
5,168,062 A	12/1992	Stinski
5,288,641 A	2/1994	Roizman
5,328,688 A	7/1994	Roizman
5,385,839 A	1/1995	Stinski
5,599,691 A	2/1997	Roizman
5,602,007 A	2/1997	Dunn et al.
5,698,531 A	12/1997	Nabel et al.
5,824,318 A	10/1998	Mohr et al.
5,846,707 A	12/1998	Roizman
6,040,169 A	3/2000	Brown et al.
6,071,692 A	6/2000	Roizman
6,120,773 A	9/2000	Roizman
6,172,047 B1	1/2001	Roizman et al.
6,297,219 B1	10/2001	Nabel et al.
6,340,673 B1	1/2002	Roizman et al.
6,423,528 B1	7/2002	Brown et al.
6,428,968 B1	8/2002	Molnar-Kimber et al.
6,649,157 B2	11/2003	Coffey et al.
6,770,274 B1	8/2004	Martuza et al.
7,063,835 B2	6/2006	Coffin

(Continued)

FOREIGN PATENT DOCUMENTS

EP	1235853 B1	7/2009
JP	2013511549 A	4/2013

(Continued)

OTHER PUBLICATIONS

Jean-François Fonteneau et al. Oncoimmunology. 2016; 5(1): e1066961, published online Aug. 12, 2015. doi: 10.1080/2162402X.2015.1066961.*

Hu et al. Cancer Therapy: Clinical, published on Nov. 22, 2006, pp. 6737-6747.*

NIH, National Cancer Institute (NCI)'s Dictionary of Cancer terms on line searched by on Sep. 25, 2024. p. 1-1.*

Fielding et al. "A hyperfusogenic gibbon apeleukemia envelope glycoprotein: targeting of a cytotoxic gene by ligand display", Hum Gene Ther. Apr. 10, 2000;11(6):817-26.

Majid et al. Recombinant Vesicular Stomatitis Virus (VSV) and Other Strategies in HCV Vaccine Designs and Immunotherapy. Tan SL, (Ed.) Hepatitis C Viruses: Genomes and Molecular Biology, Ch. 15. Norfolk (UK): Horizon Bioscience (2006).

(Continued)

Primary Examiner — Bao Q Li

(74) Attorney, Agent, or Firm — Norton Rose Fulbright US LLP

(57)

ABSTRACT

The present invention relates to oncolytic virus comprising: (i) a GM-CSF-encoding gene; and (ii) an immune co-stimulatory pathway activating molecule or an immune co-stimulatory pathway activating molecule-encoding gene.

30 Claims, 33 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited**U.S. PATENT DOCUMENTS**

7,223,593	B2	5/2007	Coffin	
7,537,924	B2	5/2009	Coffin	
7,749,745	B2	7/2010	Johnson et al.	
7,981,669	B2	7/2011	Coffin et al.	
8,273,568	B2	9/2012	Martuza et al.	
8,277,818	B2	10/2012	Coffin	
8,361,978	B2	1/2013	Rabkin et al.	
8,470,577	B2	6/2013	Johnson et al.	
8,679,830	B2	3/2014	Coffin et al.	
8,680,068	B2	3/2014	Coffin	
8,703,120	B2	4/2014	Martuza et al.	
8,871,193	B2	10/2014	Johnson et al.	
8,986,672	B2	3/2015	Zhang et al.	
9,487,581	B2	11/2016	Abate et al.	
9,492,482	B2	11/2016	Beech et al.	
9,789,182	B2	10/2017	Graziano et al.	
9,827,307	B2	11/2017	Rabkin et al.	
9,868,961	B2	1/2018	Allison et al.	
10,039,796	B2	8/2018	Zhang et al.	
10,287,252	B2	5/2019	Cowley et al.	
10,301,600	B2	5/2019	Coffin	
10,555,981	B2	2/2020	Silvestre et al.	
10,570,377	B2	2/2020	Coffin	
10,612,005	B2	4/2020	Coffin	
10,626,377	B2	4/2020	Coffin	
10,765,710	B2	9/2020	Zitvogel et al.	
10,947,513	B2	3/2021	Coffin	
11,427,810	B2 *	8/2022	Coffin	C07K 16/2818
11,473,063	B2 *	10/2022	Coffin	C07K 14/005
2003/0091537	A1	5/2003	Coffin	
2008/0014175	A1	1/2008	Hallahan et al.	
2010/0297072	A1	11/2010	DePinho	
2011/0044953	A1	2/2011	Allison et al.	
2013/0202639	A1	8/2013	Kousoulas et al.	
2014/0154216	A1	6/2014	Coffin	
2014/0271677	A1	9/2014	Palese et al.	
2015/0232812	A1	8/2015	Coffin	
2015/0283234	A1	10/2015	Graziano et al.	
2016/0040186	A1	2/2016	Liu	
2021/0252135	A1	8/2021	Coffin	
2021/0254019	A1	8/2021	Coffin	
2022/0056480	A1	2/2022	Bell et al.	

FOREIGN PATENT DOCUMENTS

JP	2015/508156	A	3/2015	
JP	2016509045	A	3/2016	
WO	97/12623	A1	4/1997	
WO	9830707	A2	7/1998	
WO	2001/53505	A2	7/2001	
WO	2001/53506	A2	7/2001	
WO	2005/011715	A1	2/2005	
WO	2006/002394	A2	1/2006	
WO	2006/048749	A1	5/2006	
WO	2007/052029	A1	5/2007	
WO	2007/123737	A2	11/2007	
WO	2010042189	A2	4/2010	
WO	2011063309	A1	5/2011	
WO	2011/118866	A1	9/2011	
WO	2012/038606	A1	3/2012	
WO	2013/038066	A1	3/2013	
WO	2013112942	A1	8/2013	
WO	WO2014022138	A2 *	2/2014	A61K 39/395
WO	WO2014036412	A2 *	3/2014	A61K 39/00
WO	WO2014036412	A3 *	3/2014	C07K 16/28
WO	2014/066532	A1	5/2014	
WO	2014128235	A1	8/2014	
WO	2015032755	A1	3/2015	
WO	2015/059303	A1	4/2015	
WO	2015/077624	A1	5/2015	
WO	2015066042	A1	5/2015	
WO	2015/128313	A1	9/2015	
WO	2015/153417	A1	10/2015	
WO	2016/008976	A1	1/2016	
WO	2016/118865	A1	7/2016	

WO	2017/118864	A1	7/2017
WO	2017/118866	A1	7/2017
WO	2017/118867	A1	7/2017
WO	2017/181420	A1	10/2017
WO	2018127713	A1	7/2018

OTHER PUBLICATIONS

Malhotra et al. Use of an Oncolytic Virus Secreting GM-CSF as Combined Oncolytic and Immunotherapy for Treatment of Colorectal and Hepatic Adenocarcinomas, 141(4) Surgery 520-529 (Apr. 2007).

Marabelle et al., "Intratumoral Anti-CTLA-4 Therapy: Enhancing Efficacy While Avoiding Toxicity", Clin Cancer Res. 2013, 19(19):5261-3.

McDonald et al. A measles virus vaccine strain derivative as a novel oncolytic agent against breast cancer, 99 Breast Cancer Research and Treatment 177-184 (2006).

Msaouel et al. Attenuated oncolytic Measles Virus strains as cancer therapeutics, 13(9) Curr. Pharm. Biotechnol. 1732-41 (Jul. 1, 2012).

Murata et al: "X40 costimulation synergizes with GM-CSF whole-cell vaccination to overcome established CD8+ T cell tolerance to an endogenous tumor antigen", J Immunol. Jan. 15, 2006;176(2):974-83.

Nakamori et al. Potent Antitumor Activity After Systemic Delivery of a Doubly Fusogenic Oncolytic Herpes Simplex Virus Against Metastatic Prostate Cancer, 60 The Prostate 53-60 (2004).

Nakano et al., Journal of Japan Surgical Society, 2001,102, Extra Issue, p. 82, No. SF4e-4.

Office Action issued in European Patent Application No. 1770385, dated May 21, 2019.

Oliveira et al. Poxvirus Host Range Genes and Virus-Host Spectrum: A Critical Review, 9(11) Viruses 2017 331 (Nov. 7, 2017).

Output from antibodies-online.com search for CTLA-4 Antibodies (performed Nov. 24, 2021), available at <https://www.antibodies-online.com/search.php#5qk9>.

Output from Antibodypedia search for CTLA-4 Antibodies (performed Nov. 24, 2021), available at <https://www.antibodypedia.com/gene/19961/CTLA4>.

Output from Biocompare search for CTLA-4 Antibodies (performed Nov. 24, 2021), available at <https://www.biocompare.com/Search-Antibodies/?search=CTLA-4&said=0>.

Output from the National Institutes of Health (NIH) National Center for Biotechnology Information (NCBI) Taxonomy Browser searches for "herpesviridae", "poxviridae", "adenoviridae", "retroviridae", "rhabdoviridae", "paramyxoviridae", and "reoviridae" (performed Nov. 3, 2021), available at: <https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Root>.

Patentee's response to EPO communication dtd Sep. 25, 2009, EP No. 17701910.6.

Pentcheva-Hoang et al. B7-1 and B7-2 Selectively Recruit CTLA-4 and CD28 to the Immunological Synapse, 21 Immunity 401-413 (Sep. 2004).

Petition for Post-Grant Review of U.S. Pat. No. 10,947,513, filed Dec. 15, 2021 with the TTAB, Petitioner—Transgene and Bioinvent International AB.

Piasecki et al., "Talilmogene laherparepvec increases the anti-tumor efficacy of the anti-PD-1 Abstract, Apr. 19, 2015 Immune checkpoint blockade," AACR Annual Meeting Presentation.

Reese, "Abstract IA24: New frontiers in oncolytic virus therapy," Cancer Immunology Research, 2016, 4(11):1A24-1A24.

Reoviridae Information from Virus Pathogen Resource (ViPR) retrieved on Nov. 4, 2021, available at <https://www.viprbrc.org/brc/aboutPathogen.spg?decorator=reo>.

Ribas, Clinical Development of the Anti-CTLA-4 Antibody Tremelimumab, 37(5) Seminars in Oncology 450-454 (Oct. 2010).

Riedel et al. Components and Architecture of the Rhabdovirus Ribonucleoprotein Complex, 12(9) Viruses 2020 959 (Aug. 2020).

Robbins et al; "Viral Vectors for Gene Therapy"; Pharmacol, Ther.; vol. 80, No. 1; pp. 35-47; 1998.

Robinson et al., "Novel Immunocompetent Murine Tumor Model for Evaluation of Conditionally Replication-Competent (Oncolytic) Murine Adenoviral Vectors," Journal of Virology, 2009, 83(8):3450-3462.

(56)

References Cited**OTHER PUBLICATIONS**

- Rojas et al. Defining Effective Combinations of Immune Checkpoint Blockade and Oncolytic Virotherapy, 21(24) Clin. Cancer Res. 5543-51 (Dec. 2015).
- Saha et al. The Adenovirus Genome Contributes to the Structural Stability of the Virion, 6(9) Viruses 2014 3563-3583 (Sep. 24, 2014).
- Salzberg, Open questions: How many genes do we have? 16 BMC Biology 94 (Aug. 20, 2018).
- Schirrmann et al., "Transient Production of scFv-Fc Fusion Proteins in Mammalian Cells", Antibody Engineering, 2010, vol. 2; Chapter 30, p. 387-398, © Springer-Verlag Berlin Heidelberg.
- Senzer et al., "Phase II Clinical Trial of a Granulocyte-Macrophage Colony-Stimulating Factor-Encoding, Second-Generation Oncolytic herpesvims in Patients with Unresectable Metastatic Melanoma" Journal of Clinical Oncology, 2009, 27(34):5763-5771.
- Shan et al., "Characterization of scFv-Ig Constructs Generated from the Anti-CD20 mAb 1F5 Using Linker Peptides of Varying Lengths", Journal of Immunology, 1999, 162:6589-6595.
- Sharp and Li, The codon adaptation index—a measure of directional synonymous codon usage bias, and its potential applications, 15(3) Nucleic Acids Research 1281-95 (1987).
- Simpson et al., "Combination of a Fusogenic Glycoprotein, Prodrug activation, and Oncolytic Herpes Simplex Virus for Enhanced Local Tumor Control," Cancer Research, 2006, 66(9):4835-4842.
- Singh et al. Oncolytic viruses & their specific targeting to tumour cells, 136 Indian J. Med. Res. 571-584 (Oct. 2012).
- Sinkovics and Horvath, Natural and genetically engineered viral agents for oncolysis and gene therapy of human cancers, 56 Arch. Immunol. Ther. Exp. 3-59 (2008).
- Smith et al. Studies on the Use of Viruses in the Treatment of Carcinoma of the Cervix, 9(6) Cancer 1211-18 (Nov.-Dec. 1956).
- Sokolowski et al., "Oncolytic virotherapy using herpes simplex vims: how far have we come?" Oncolytic Virotherapy, 2015, 4:207-219.
- Species list extracted from International Committee on Taxonomy of Viruses (ICTV) Master Species List (Jul. 20, 2021), available at: <https://talk.ictvonline.org/taxonomy/vmr/>.
- Statement of Grounds of Opposition from the Opponent, Margaret Dixon Limited, dated Jun. 7, 2021, EP3400293 (EP Appl. No. 17701910.6).
- Study Details for Clinical Trial NCT02272855 "A Study of Combination Treatment With HF10 and Ipilimumab in Patients With Unresectable or Metastatic Melanoma", last updated Sep. 26, 2018, available at: <https://clinicaltrials.gov/ct2/show/NCT02272855>.
- Study Details for Clinical Trial NCT02620423 "Study of Pembrolizumab with REOLYSIN® and Chemotherapy in Patients With Advanced Pancreatic Adenocarcinoma", last updated Sep. 13, 2018, available at: <https://clinicaltrials.gov/ct2/show/NCT02620423>.
- Sumimoto et al. "GM-CSF and B7-1 (CD80) co-stimulatory signals co-operate in the induction of effective anti-tumor Immunity in syngeneic mice", Int J Cancer. Nov. 14, 1997;73(4):556-61.
- Summary of Characteristics of Commercial Viral Vectors from ThermoFisher Scientific, retrieved Nov. 4, 2021, available at <https://www.thermofisher.com/us/en/home/references/gibco-cell-culture-basics/transfection-basics/gene-delivery-technologies/viral-delivery/viral-vectors.html>.
- Tan et al. Combination therapy of oncolytic herpes simplex virus HF10 and bevacizumab against experimental model of human breast carcinoma xenograft, 136 Int. J. Cancer 1718-30 (2015).
- Terada K. et al., "Development of a rapid method to generate multiple oncolytic HSV vectors Gene Therapy, vol. 13, No. 8, (Apr. 1, 2006), pp. 705-714 and their in vivo evaluation using syngeneic mouse tumor models".
- Tesfay et al. PEGylation of Vesicular Stomatitis Virus Extends Virus Persistence in Blood Circulation of Passively Immunized Mice, 87(7) Journal of Virology 3752-59 (Apr. 2013).
- Third Party Submission submitted in Related U.S. Appl. No. 16/068,816, dated Jul. 16, 2019.
- Third Party Submission submitted in Related U.S. Appl. No. 16/068,823, dated Jul. 18, 2019.
- Third Party Submission submitted in Related U.S. Appl. No. 16/068,826, dated Aug. 7, 2019.
- Third Party Submission submitted in Related U.S. Appl. No. 16/068,830, dated Jul. 18, 2019.
- Todo, Tomoki, Armed oncolytic herpes simplex viruses for brain tumor therapy, 208-213, Cell Adhesion* Migration 2:3, Jul./Aug./Sep. 2008.
- Van den Wollenberg et al. Replicating reoviruses with a transgene replacing the codons for the head domain of the viral spike, 22 Gene Therapy 267-279 (2015).
- Wennier et al. Bugs and Drugs: Oncolytic Virotherapy in Combination with Chemotherapy, 13(9) Curr. Pharm. Biotechnol. 1817-33 (Jul. 2012).
- Wertz et al. Adding genes to the RNA genome of vesicular stomatitis virus: positional effects on stability of expression, 76(15) J. Virol. 7642-50 (Aug. 2002).
- Willemsen and Zwart, On the stability of sequences inserted into viral genomes, 5(2) Virus Evolution vez045 (Jul. 2019).
- Yan et al., "Developing Novel Oncolytic Adenoviruses through Bioselection," Journal of Virology, 2003, 77(4):2640-2650.
- Yang et al. Cascade regulation of vaccinia virus gene expression is modulated by multistage promoters, 447(1-2) Virology 213-220 (Dec. 2013).
- Yen et al. Vaccinia virus infection & temporal analysis of virus gene expression: Part 2, 2009(26) J. Vis. Exp. 1169 (Apr. 2009).
- Yi et al Cancer Res 2007, 67 20 10027-10037.
- Yo, Y-T et al: "Coexpression of Flt3 ligand and GM-CSF genes modulates immune responses induced by HER2/neu DNA vaccine", Cancer Gene Ther. Nov. 2007;14(11):904-17.
- Ahlers et al: "A push-pull approach to maximize vaccine efficacy: abrogating suppression with an IL-13 inhibitor while augmenting help with granulocyte/macrophage colony-stimulating factor and CD40L", Proc Natl Acad Sci USA, Oct. 1, 2002;99(20):13020-5.
- Ahmed et al, Intratumoral expression of a fusogenic membrane glycoprotein enhances the efficacy of replicating adenovirus therapy, 10 Gene Therapy 1663-71 (2003).
- Allison et al., "For Their Discovery of Cancer Therapy by Inhibition of Negative Immune in Physiology of Medicine Regulation"; The Nobel Assembly at Karolinska Institutet; 2018 Nobel Prize.
- Altschul, S F et al (1990) J Mol Biol 215:403-10.
- Altschul, S.F. (1993) J Mol Evol 36:290-300.
- Annex A—WO 2017/118864—Figures 3 and 4 published Jul. 13, 2017.
- Asada, Treatment of Human Cancer with Mumps Virus, 34(6) Cancer 1907-28 (Dec. 1974).
- Assal et al: "Emerging targets in cancer immunotherapy: beyond CTLA-4 and PD-1", Immunotherapy. 2015;7(11):1169-86.
- Balvay et al. Translational control of retroviruses, 5 Nature Reviews Microbiology 128-140 (Feb. 2007).
- Bateman et al. Cancer Res. Mar. 1, 2000;60(6):1492-7.
- Bateman et al. Cancer Res. Nov. 15, 2002;62(22):6566-78.
- Bauzon and Hermiston, 2014. Front. Immunol., 5(74): 1-10.
- Belsham and Sonenberg, RNA-protein interactions in regulation of picomavirus RNA translation, 60(3) Microbiological Reviews 499-511 (Sep. 1996).
- Bett et al. Packaging capacity and stability of human adenovirus type 5 vectors, 67(10) J. Virol. 5911-21 (Oct. 1993).
- BLAST analysis (publicly available through the National Centre for Biotechnology Information—<http://www.ncbi.nlm.nih.gov/>).
- Blechacz et al. Engineered Measles Virus as a Novel Oncolytic Viral Therapy System for Hepatocellular Carcinoma, 44(6) Hepatology 1465-77 (Dec. 2006).
- Brochu-Lafontaine and Lemay, Addition of exogenous polypeptides on the mammalian reovirus outer capsid using reverse genetics, 179 J. Virol. Methods 342-350 (2012).
- Capece et al: "Targeting costimulatory molecules to improve anti-tumor immunity", J Biomed Biotechnol, 2012; 2012:926321.
- Carson et al., "Oncolytic Herpes Simplex Virus 1 (HSV-1) Vectors: Increasing Treatment Efficacy and Range Through Strategic Virus Design", Drugs Future. 2010;35(3): 183-195.

(56)

References Cited**OTHER PUBLICATIONS**

- Carter et al. Identification of an overprinting gene in Merkel cell polyomavirus provides evolutionary insight into the birth of viral genes, 110(31) *Proceedings of the National Academy of Sciences* 12744-49 (Jul. 2013).
- Cell Signaling Technology; Immune Checkpoint Signaling in the Tumor Microenvironment1; Mar. 2018.
- Chen et al., "Dual silencing of Bcl-2 and Survivin by HSV-1 vector shows better antitumor efficacy in higher PKR phosphorylation tumor cells in vitro and in vivo", *Cancer Gene Ther* 22, 380-386; 2015.
- Choi et al. Polymeric oncolytic adenovirus for cancer gene therapy, 219 *Journal of Controlled Release* 181-191 (2015).
- Choi et al., "Concurrent delivery of GM-CSF and B7-1 using an oncolytic adenovirus elicits potent antitumor effect", *Gene Therapy* (2006) 13, 1010-1020 & 2006 Nature Publishing Group.
- Choi et al., "Strengthening of antitumor immune memory and prevention of thymic atrophy mediated by adenovirus expressing IL-12 and GM-CSF", *Gene Therapy* (2012) 19, 711-723 & 2012 Macmillan Publishers.
- Chou et al., "Mapping of Herpes Simplex Virus-1 Neurovirulence to ?134.5, a Gene Nonessential for Growth in Culture," *Science*, 1990, 250(4985):1262-1266.
- Compilation of Virus Information from Swiss Institute of Bioinformatics retrieved on Nov. 3, 2021, available at <https://viralzone.expasy.org/>.
- Croyle et al. PEGylation of a Vesicular Stomatitis Virus G Pseudotyped Lentivirus Vector Prevents Inactivation in Serum, 78(2) *Journal of Virology* 912-921 (Jan. 2004).
- Danthinne and Imperiale, Production of first generation adenovirus vectors: a review, 7 *Gene Therapy* 1707-14 (2000).
- Declaration of Dr. Sylvia D. Hall-Ellis dated Nov. 29, 2021 and Curriculum vitae.
- Declaration of John C. Bell, Ph.D. dated Dec. 14, 2021 and Curriculum vitae.
- Deguchi et al. Combination of the Tumor Angiogenesis Inhibitor Bevacizumab and Intratumoral Oncolytic Herpes Virus Injections as a Treatment Strategy for Human Gastric Cancers, 59(118) *Hepatogastroenterology* 1844-50 (Sep. 2012).
- Devereux et al (1984) *Nucleic Acids Research* 12, p. 387-395.
- Dias et al., 2012. *Gene Ther.*, 19: 988-998.
- Diefenbach et al., "Oncolytic virotherapy using herpes simplex virus: how far have we come?" *Oncolytic Virotherapy*, Nov. 1, 2015 (Nov. 1, 2015), p. 207.
- Dikstein, The unexpected traits associated with core promoter elements, 2(5) *Transcription* 201-206 (Sep. 2011).
- Documents filed on Jul. 9, 2018 in U.S. Appl. No. 16/068,830, including original application, preliminary amendment, application data sheet, search report, and transmittal form.
- Donovan-Banfield et al. Deep splicing plasticity of the human adenovirus type 5 transcriptome drives virus evolution, 3 *Communications Biology* (2020) 124.
- Du et al. "Tumor-specific oncolytic adenoviruses expressing granulocyte macrophage colony-stimulating factor or anti-CTLA4 antibody for the treatment of cancers", *Cancer Gene Therapy*, vol. 21, No. 8, Jul. 18, 2014 (Jul. 18, 2014), pp. 340-348.
- Ebert et al. Syncytia Induction Enhances the Oncolytic Potential of Vesicular Stomatitis Virus in Virotherapy for Cancer, 64 *Cancer Research* 3265-3270 (May 2004).
- Engeland et al. CTLA-4 and PD-L1 Checkpoint Blockade Enhances Oncolytic Measles Virus Therapy, 22(11) *Molecular Therapy* 1949-59 (Nov. 2014).
- Excerpts from S. Baron (Ed.), *Medical Microbiology*, 4th. Ed. University of Texas Medical Branch at Galveston (1996).
- Fransen et al., "Controlled Local Delivery of CTLA-4 Blocking Antibody Induces CD8+ T-Cell-Dependent Tumor Eradication and Decreases Risk of Toxic Side Effects" *Clin Cancer Res.* 2013, 19(19):5381-9.
- Fu et al. Expression of a Fusogenic Membrane Glycoprotein by an Oncolytic Herpes Simplex Virus Potentiates the Viral Antitumor Effect, 7(6) *Molecular Therapy* 748-754 (Jun. 2003).
- Fukuhara et al. Triple Gene-Deleted Oncolytic Herpes Simplex Virus Vector Double-Armed with Interleukin 18 and Soluble B7-1 Constructed by Bacterial Artificial Chromosome-Mediated System, 65(23) *Cancer Res.* 10663-68 (Dec. 2005).
- Gangi et al., "The safety of talimogene laherparepvec for the treatment of advanced melanoma", *Expert Opinion on Drug Safety*, Dec. 28, 2016 (Dec. 28, 2016), pp. 1-5.
- Gao et al: "Recombinant vesicular stomatitis virus targeted to Her2/neu combined with anti-CTLA4 antibody eliminates implanted mammary tumors", *Cancer Gene Ther.* Jan. 2009;16(1):44-52.
- Gibney et al., "Preliminary results from a phase A study of INCB024360 combined with ipilimumab (ipi) in patients (pts) with melanoma." 2014 ASCO Annual Meeting, No. 3010.
- Grandi, et al., *Cancer Gene Therapy* (2010) 17, 655-663 (Year: 2010).
- Gri et al: "X40 ligand-transduced tumor cell vaccine synergizes with GM-CSF and requires CD40-Apc signaling to boost the host T cell antitumor response", *J Immunol.* Jan. 1, 2003;170(1):99-106.
- Guedan et al. GALVexpression enhances the therapeutic efficacy of an oncolytic adenovirus by inducing cell fusion and enhancing virus distribution, 19 *Gene Therapy* 1048-57 (2012).
- Gómez-Treviño et al. Effects of adenovirus-mediated SV5 fusogenic glycoprotein expression on tumor cells, 5 *J. Gene Med.* (2003) 483-492.
- Haswell et al *Eur J Immunol* 2001 31 3094-3100.
- Heinkoff and Heinkoff (1992) *Proc. Natl. Acad. Sci. USA* 89:10915-10919.
- Herpesviridae Information from Virus Pathogen Resource (ViPR) retrieved on Nov. 4, 2021, available at: <https://www.viprbrc.org/brc/aboutPathogen.spg?decorato=Hierpes>.
- Heterologous Expression. In Binder, Hirokawa and Windorst (eds.)—*Encyclopedia of Neuroscience*. (2009) Springer, Berlin, Heidelberg https://doi.org/10.1007/978-3-540-29678-2_2190.
- Hillier et al. Genomics in *C. elegans*: so many genes, such a little worm, 15 *Genome Research* 1651-60 (2005).
- Ho et al. Unconventional viral gene expression mechanisms as therapeutic targets, 593 *Nature* 362-371 (May 2021).
- Hoffmann et al. *World J Gastroenterol.* Jun. 14, 2007;13(22):3063-70.
- Hoffmann et al. *World J Gastroenterol.* Mar. 28, 2008 14(12):1842-1850.
- Hoggmann et al. W.J. G 2007, Jun. 14, 2013 (22), pp. 3063-30700.
- Hooren et al., "Abstract B103: Intravesical administration of CTLA-4 blocking monoclonal antibodies as a means to optimize bladder cancer therapy", *Cancer Immunol Res.* 2016,4 (11_Supplement): B103.
- Hooren et al., "Local checkpoint inhibition of CTLA-4 as a monotherapy or in combination with anti-PD1 prevents the growth of murine bladder cancer" *Eur J Immunol.* 2017,47(2):385-393.
- Hu et al. "A simplified system for generating oncolytic adenovirus vector carrying one or two transgenes", *Cancer Gene Therapy* (2008) 15, 173-182 & 2008 Nature Publishing Group.
- Huang et al., *Mol Ther.* Feb. 2010, vol. 18, No. 2, pp. 264-274.
- Hurwitz et al: "CTLA-4 blockade synergizes with tumor-derived granulocyte-macrophage colony-stimulating factor for treatment of an experimental mammary carcinoma", *Proc Natl Acad Sci USA*, Aug. 18, 1998;95(17):10067-71.
- IGI Global "What is Heterologous Expression" retrieved from <https://www.igiglobal.com/dictionary/heterologousexpression/49470>.
- Inouye et al., "Codon optimization of genes for efficient protein expression in mammalian cells by selection of only preferred human codons", *Protein Expression and Purification*, 2015, 109:47-54.
- International Search Report and Written Opinion issued in International Patent Application No. PCT/GB2017/050036, dated Apr. 26, 2017.
- International Search Report and Written Opinion issued in International Patent Application No. PCT/GB2017/050037, dated Apr. 25, 2017.

(56)

References Cited**OTHER PUBLICATIONS**

- International Search Report and Written Opinion issued in International Patent Application No. PCT/GB2017/050038, dated Apr. 24, 2017.
- International Search Report for International Patent Application No. PCT/EP2015/066263, mailed from European Patent Office Oct. 7, 2015.
- International Search Report for International Patent Application No. PCT/FI2009/051025, mailed from European Patent Office Mar. 24, 2010.
- Ishihara et al. Systemic CD8+ T Cell-Mediated Tumoricidal Effects by Intratumoral Treatment of Oncolytic Herpes Simplex Virus with the Agonistic Monoclonal Antibody for Murine Glucocorticoid-Induced Tumor Necrosis Factor Receptor, 9(8) PLoS One e104669 (Aug. 2014).
- Ishikawa et al. STING regulates intracellular DNA-mediated, type I interferon-dependent innate immunity, 461 Nature 788-792 (Oct. 8, 2009).
- Jacobs et al. HSV-1 based vectors for gene therapy of neurological diseases and brain tumors Part II Vector Systems and Applications, 1(5) Neoplasia 402-416 (Nov. 1999).
- Jacobs et al. Vaccinia Virus Vaccines: Past, Present and Future, 84(1) Antiviral Res. 1-13 (Oct. 2009).
- John et al. Oncolytic Virus and Anti-4-IBB Combination Therapy Elicits Strong Antitumor Immunity against Established Cancer, 72(7) Cancer Research 1651-60 (Apr. 2012).
- Kanagavelu et al PlosOne 2014, 9, 2, e90100.
- Kanagavelu et al Vaccine 2012 30 691-701.
- Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5787.
- Kasuya et al., Journal of Japan Surgical Society, 2006, 107, Extra Issue (2), p. 369, No. PS-005-8.
- Kaufman et al: "Oncolytic viruses: a new class of immunotherapy drugs", Nat Rev Drug Discov, vol. 14, 642-662 (Sep. 2015).
- Kaufmann et al. Chemovirotherapy of Malignant Melanoma with a Targeted and Armed Oncolytic Measles Virus, 133 Journal of Investigative Dermatology 1034-42 (2013).
- Kelly and Russell, History of Oncolytic Viruses: Genesis to Genetic Engineering, 15(4) Molecular Therapy 651-659 (Apr. 2007).
- Kim et al Cancer Res 2009, 69, 21, 8516-8525.
- Kleinpeter et al. Vectorization in an oncolytic vaccinia virus of an antibody, a Fab and a scFv against programmed cell death-1 (PD-1) allows their intratumoral delivery and an improved tumor-growth inhibition, 5(10) Oncoimmunology e1220467 (2016).
- Le Boeuf et al. Synergistic Interaction Between Oncolytic Viruses Augments Tumor Killing, 18(5) Molecular Therapy 888-895 (May 2010).
- Lee et al. Enhanced Antitumor Effect of Oncolytic Adenovirus Expressing Interleukin-12 and B7-1 in an Immunocompetent Murine Model, 12(19) Clin. Cancer Res. 5859-68 (Oct. 2006).
- Lee et al: "Oncolytic potential of E1B 55 kDa-deleted YKL-1 recombinant adenovirus: correlation with p53 functional status" Int J Cancer (2000) 88: 454-463.
- Li et al. Int. J. Cancer 2008, 123: 493-499.
- Li, B et al: "Established B16 tumors are rejected following treatment with GM-CSF-secreting tumor cell immunotherapy in combination with anti-4-1 BB mAb", Clin Immunol. Oct. 2007;125(1):76-87.
- Li, B et al: "Anti-programmed death-1 synergizes with granulocyte macrophage colony-stimulating factor-secreting tumor cell immunotherapy providing therapeutic benefit to mice with established tumors", Clin Cancer Res. Mar. 1, 2009;15(5):1623-34.
- Lipson and Drake, Ipilimumab: An Anti-CTLA-4 Antibody for Metastatic Melanoma, 17(22) Clin. Cancer Res. 6958-62 (Nov. 2011).
- List of known isolates within each virus family extracted from NCBI Taxonomy Browser Output of Ex. 1023, dtd Nov. 3, 2021.
- Liu et al., "ICP34.5 deleted herpes simplex virus with enhanced oncolytic, immune stimulating, and anti-tumour properties," Gene Therapy, 2003, 10(4):292-303.
- Loskog, Angelica, "Immunostimulatory Gene Therapy Using Oncolytic Viruses as Vehicles," Viruses, 2015, 7:5780-5791.
- Lundstrom, New frontiers in oncolytic viruses: optimizing and selecting for virus strains with improved efficacy, 12 Biologics: Targets and Therapy 43-60 (2018).
- Ma et al. Oncolytic herpes simplex virus and immunotherapy, 19 BMC Immunology 40 (2018).
- Maclean et al., "Herpes simplex virus type 1 deletion variants 1714 and 1716 pinpoint neurovirulence-related sequences in Glasgow strain 17 + between immediate early gene 1 and the 'a' sequence," Journal of General Virology, 1991, 72:631-639.
- EPO Opposition "Opponent's Response in opposition proceedings against Replimune's European Patent EP 3400291", provided by the European Patent Office on May 4, 2023.
- Fonteneau et al., "Oncolytic immunotherapy: The new clinical outbreak", Oncoimmunology, 2016, 5:1:e1066961.
- Japanese Notice of Rejection mailed Feb. 28, 2023 during examination of related JP Patent Appl. No. 2019-537074.
- Marcos et al., "Mapping of the RNA promoter of Newcastle disease virus", Virology, vol. 331, Issue 2, 2005, pp. 396-406.
- Noton and Fearn, "Initiation and regulation of paramyxovirus transcription and replication", Virology, 2015, 479-480, 545- 554.
- Alekseenko et al: "Therapeutic properties of a vector carrying the HSV thymidine kinase and GM-CSF genes and delivered as a complex with a cationic copolymer", Journal of Translational Medicine (2015) 13:78.
- Amendment and Reply to Accompany Request for Continued Examination dated Sep. 26, 2018, U.S. Appl. No. 15/325,576, filed Jan. 11, 2017, 65 pages.
- Amendment and Reply to Pursuant to 37 CFR §1.112 dated Aug. 26, 2019, U.S. Appl. No. 15/325,576, filed Jan. 11, 2017, 34 pages.
- Applicant-Initiated Interview Summary dated Jan. 29, 2021 for U.S. Appl. No. 16/068,830, filed Jul. 9, 2018, 2 pages.
- Bell, John C., Transcript of John C. Bell, Sep. 23, 2022, 87 pages.
- Chiocca, E.A., Curriculum Vitae of E. Antonio Chiocca, M.D., Ph.D., Oct. 8, 2019, 92 pages.
- Chiocca, E.A., Declaration of E. Antonio Chiocca, M.D., Ph.D., FAANS, Sep. 28, 2022, 35 pages.
- Chiocca, E.A., Transcript of E. Antonio Chiocca, M.D., Ph.D., Nov. 30, 2022, 237 pages.
- Correction of Notice of Allowability dated Feb. 1, 2021 for U.S. Appl. No. 16/068,830, filed Jul. 9, 2018, 2 pages.
- Decision Granting Institution of Past-Grant Review dated Jun. 16, 2022, Paper 16, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 27 pages.
- Demonstrative Exhibits of Petitioners, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, Exhibit 1107, for Oral Argument Date Mar. 17, 2023, 56 pages.
- Disclaimer in Patent Under 37 CFR 1.321(a) filed Mar. 15, 2022 for U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 6 pages.
- Final Written Decision dated May 25, 2023, Paper 38, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 61 pages.
- Guo, Z.S. et al., "Rapid Generation of Multiple Loci-Engineered Marker-free Poxvirus and Characterization of a Clinical-Grade Oncolytic Vaccinia Virus", Molecular Therapy: Methods & Clinical Development, vol. 7, Dec. 2017, Pittsburgh, PA, USA, pp. 112-122.
- Guse, K. et al., "Antiangiogenic Arming of an Oncolytic Vaccinia Virus Enhances Antitumor Efficacy in Renal Cell Cancer Models", Journal of Virology, 84(2), Jan. 2010, pp. 856-866.
- Lun, X.Q. et al., "Efficacy of Systemically Administered Oncolytic Vaccinia Virotherapy for Malignant Gliomas Is Enhanced by Combination Therapy with Rapamycin or Cyclophosphamide", Clinical Cancer Research 15(8), 2009, pp. 2777-2788.
- Patent Owner Response dated Sep. 28, 2022, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 58 pages.
- Patent Owner Sur Reply dated Feb. 1, 2023, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 36 pages.
- Patent Owner's Demonstrative Exhibits, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, Exhibit 2024, Replimune Limited, 75 pages.

(56)

References Cited**OTHER PUBLICATIONS**

Patent Owner's Objections to Petitioners Evidence dated Jul. 1, 2022, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 33 pages.

Patent Owner's Preliminary Response dated Mar. 22, 2022, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 41 pages.

Patent Owner's Supplemental Brief Regarding Xerox and Intel dated Mar. 31, 2023, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 8 pages.

Petitioners Additional Briefing Regarding Xerox and Intel dated Mar. 31, 2023, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 8 pages.

Petitioners' Reply to Owner's Response dated Dec. 20, 2022, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 42 pages.

Petitioners' Reply to Patent Owner's Preliminary Response dated Apr. 14, 2022, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 8 pages.

Record of Oral Hearing Held Mar. 17, 2023 dated May 22, 2023, Paper 37, AIA Review No. PGR2022-00014, U.S. Pat. No. 10,947,513 issued Mar. 16, 2021, 95 pages.

Reply under 37 CFR 1.111 dated Apr. 30, 2020, U.S. Appl. No. 16/068,830, filed Jan. 9, 2017, 11 pages.

Response to Rule 312 Communication and Applicant-Initiated Interview Summary dated Jan. 13, 2021 for U.S. Appl. No. 16/068,830, filed Jul. 9, 2018, 4 pages.

Rintoul, J.L., et al. "A Selectable and Excisable Marker System for the Rapid Creation of Recombinant Poxviruses", *PloS one*, 6(9), 2011, e24643, pp. 1-12.

Semmrch, M. et al., "Vectorized Treg-depleting α CTLA-4 elicits antigen cross- presentation and CD8+ T cell immunity to reject 'cold' tumors", *Journal for ImmunoTherapy of Cancer*, 10(1), 2022, 36 pages.

Semmrch, M. et al., "Vectorized Treg-depleting α CTLA-4 elicits antigen cross-presentation and CD8+ T cell immunity to reject "cold" tumors", *BioInvent International AB, Lund, Sweden, Transgene S.A., Illkirch-Graffenstaden, France, Abstract #746*, 1 page.

Third Party Submissions Under 37 CFR §1.290 dated Jul. 30, 2019 for U.S. Appl. No. 16/068,830, filed Jul. 9, 2018, 2 pages.

Thomas, S. et al., "Development of a new fusion-enhanced oncolytic immunotherapy platform based on herpes simplex virus type 1", *Journal for Immuno Therapy of Cancer*, 7:214, 2019, 17 pages.

Yamamoto, S., et al., "Imaging immediate-early and strict-late promoter activity during oncolytic herpes simplex virus type 1 infection and replication in tumors", *Gene Therapy*, 13, 2006, pp. 1731-1736.

Yang, S. et al., "Development of optimal bicistronic lentiviral vectors facilitates high-level TCR gene expression and robust tumor cell recognition", *Gene Therapy*, 15(21), Nov. 2008, 20 pages.

Thorne, "Next-Generation Oncolytic Vaccinia Vectors," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 14, pp. 205-215, 2012.

Dhar et al., "Syrian Hamster Tumor Model to Study Oncolytic ADS-Based Vectors," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 4, pp. 53-63, 2012.

Doronin et al., "Construction of Targeted and Armed Oncolytic Adenoviruses," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 3, pp. 35-52, 2012.

Fournier et al., "Analysis of Three Properties of Newcastle Disease Virus for Fighting Cancer: Tumor-Selective Replication, Antitumor Cytotoxicity, and Immunostimulation," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 13, pp. 177-204, 2012.

Gimenez-Alejandre et al., "Construction of Capsid-Modified Adenoviruses by Recombination in Yeast and Purification of Iodixanol-Gradient," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 2, pp. 21-34, 2012.

Goldufsky et al., "Oncolytic virus therapy for cancer," *Oncolytic Virotherapy*, 2:31-46, 2013.

Jeon et al. *Journal of Virological Methods*, 2022, vol. 299, pp. 1-7.

Rajiani et al. *Molecular Therapy*, published on May 2015, vol. 23, Supplement 1, S30.

Takara Bio, 2000 URL: <https://www.takarabio.com/documents/Nector%20Documents/PT3155-5.pdf>; Accessed Apr. 20, 2022 (Year: 2000).

Shmulevitz et al., "Exploring Host Factors that Impact Reovirus Replication, Dissemination, and Reovirus-Induced Cell Death in Cancer Versus Normal Cells in Culture," In: *Oncolytic Viruses: Methods and Protocols*, edited by David H. Kirm et al., Humana Press, vol. 797, Chapter 12, pp. 163-176, 2012.

Stedman's Medical dictionary, 2000, lines 1-3 (Year: 2000).

Woller et al. "Viral infection of tumors overcomes resistance to PD-1-immunotherapy by broadening neoantigenome-directed T-cell responses", *Molecular Therapy*, vol. 23, No. 10, 1630-1640, 2015.

* cited by examiner

Figure 1




Figure 2




Figure 3




Figure 4A




Figure 4B




Figure 4C





Figure 4D






Figure 4F




Figure 5B





Figure 5C







Figure 5E






Figure 6




Figure 7A




Figure 7B




Figure 8A




Figure 8B




Figure 9




Figure 10






Figure 11A




Virus: 5 injections of 5×10^6 pfu into the right flank tumor only
Anti-PD1: 10mg/kg i.p. Q3Dx9 (clone RMP1-14; BioXCell)

Figure 11B




Virus: 5 injections of 5×10^6 pfu into the right flank tumor only
 Anti-CTLA-4: 3mg/kg i.p. Q3Dx9 (clone 9D9; BioXCell)

Figure 11C




Virus: 5 injections of 5×10^6 pfu into the right flank tumor only
Anti-PD1: 10mg/kg i.p. Q3Dx9 (clone RMP1-14; BioXCell)
1-MT: 5mg/ml in drinking water (1-MT alone has no effect)

Figure 12A




Virus: 3 injections of 5×10^6 pfu into the right flank tumor only
Anti-PD1: 10mg/kg i.p. Q3Dx9 (clone RMP1-14; BioXCell)

Figure 12B



Virus: 3 injections of 5×10^5 pfu into the right flank tumor only
Anti-PD1: 10mg/kg i.p. Q3Dx9 (clone RMP1-14; BioXCell)

Figure 12C



Virus: 3 injections of 5×10^4 pfu into the right flank tumor only
 Anti-PD1: 10mg/kg i.p. Q3Dx9 (clone RMP1-14; BioXCell)

Figure 12D




Figure 13






Figure 14




Human A549 lung cancer tumors in nude mice (no immune effect)
3 injections of Virus 16 or Virus 19 over 1wk of vehicle or the indicated dose of virus (N=10/group)

Figure 15



Rat 9L glioma tumors in immune competent rats
Virus: 5×10^5 pfu injected 3x/wk into the right tumor only for 3 wks

Figure 16



Mouse A20 tumors in immune competent mice (dual flank tumors)
Viruses: 5×10^4 pfu of the indicated virus injected 3x into the right tumor only

ENGINEERED VIRUS**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation of U.S. Non-Provisional patent application Ser. No. 17/164,635 filed Feb. 1, 2021, which is a continuation of U.S. Non-Provisional patent application Ser. No. 16/068,830 filed Jul. 9, 2018, which is a national phase application under 35 U.S.C. § 371 to International Application No. PCT/GB2017/050038 filed Jan. 9, 2017, which claims the benefit of priority to Great Britain Patent Application Serial Nos. 1600380.8, 1600381.6 and 160382.4 all filed Jan. 8, 2016, all of which are incorporated herein by reference in their entirety.

INCORPORATION OF SEQUENCE LISTING

The instant application contains a Sequence Listing, named "SL_KEMPP0086USC2.XML" (78,944 bytes; created Jan. 31, 2023) which has been submitted electronically in ST.26 format and is hereby incorporated by reference in its entirety.

FIELD OF THE INVENTION

The invention relates to an oncolytic immunotherapeutic agent and to the use of the oncolytic immunotherapeutic agent in treating cancer.

BACKGROUND TO THE INVENTION

Viruses have a unique ability to enter cells at high efficiency. After entry into cells, viral genes are expressed and the virus replicates. This usually results in the death of the infected cell and the release of the antigenic components of the cell as the cell ruptures as it dies. As a result, virus mediated cell death tends to result in an immune response to these cellular components, including both those derived from the host cell and those encoded by or incorporated into the virus itself and enhanced due to the recognition by the host of so called damage associated molecular patterns (DAMPs) which aid in the activation of the immune response.

Viruses also engage with various mediators of the innate immune response as part of the host response to the recognition of a viral infection through e.g. toll-like receptors and cGAS/STING signalling and the recognition of pathogen associated molecular patterns (PAMPs) resulting in the activation of interferon responses and inflammation which are also immunogenic signals to the host. These immune responses may result in the immunogenic benefit to cancer patients such that immune responses to tumor antigens provide a systemic overall benefit resulting in the treatment of tumors which have not been infected with the virus, including micro-metastatic disease, and providing vaccination against relapse.

The combined direct ('oncolytic') effects of the virus, and immune responses against tumor antigens (including non-self 'neo-antigens', i.e. derived from the particular mutated genes in individual tumors) is termed 'oncolytic immunotherapy'.

Viruses may also be used as delivery vehicles ('vectors') to express heterologous genes inserted into the viral genome in infected cells. These properties make viruses useful for a variety of biotechnology and medical applications. For example, viruses expressing heterologous therapeutic genes

may be used for gene therapy. In the context of oncolytic immunotherapy, delivered genes may include those encoding specific tumor antigens, genes intended to induce immune responses or increase the immunogenicity of antigens released following virus replication and cell death, genes intended to shape the immune response which is generated, genes to increase the general immune activation status of the tumor, or genes to increase the direct oncolytic properties (i.e. cytotoxic effects) of the virus. Importantly, viruses have the ability to deliver encoded molecules which are intended to help to initiate, enhance or shape the systemic anti-tumor immune response directly and selectively to tumors, which may have benefits of e.g. reduced toxicity or of focusing beneficial effects on tumors (including those not infected by the virus) rather than off-target effects on normal (i.e. non-cancerous) tissues as compared to the systemic administration of these same molecules or systemic administration of other molecules targeting the same pathways.

It has been demonstrated that a number of viruses including, for example, herpes simplex virus (HSV) have utility in the oncolytic treatment of cancer. HSV for use in the oncolytic treatment of cancer must be disabled such that it is no longer pathogenic, but can still enter into and kill tumor cells. A number of disabling mutations to HSV, including disruption of the genes encoding ICP34.5, ICP6, and/or thymidine kinase, have been identified which do not prevent the virus from replicating in culture or in tumor tissue in vivo, but which prevent significant replication in normal tissue. HSVs in which only the ICP34.5 genes have been disrupted replicate in many tumor cell types in vitro, and replicate selectively in tumor tissue, but not in surrounding tissue, in mouse tumor models. Clinical trials of ICP34.5 deleted, or ICP34.5 and ICP6 deleted, HSV have also shown safety and selective replication in tumor tissue in humans.

As discussed above, an oncolytic virus, including HSV, may also be used to deliver a therapeutic gene in the treatment of cancer. An ICP34.5 deleted virus of this type additionally deleted for ICP47 and encoding a heterologous gene for GM-CSF has also been tested in clinical trials, including a phase 3 trial in melanoma in which safety and efficacy in man was shown. GM-CSF is a pro-inflammatory cytokine which has multiple functions including the stimulation of monocytes to exit the circulation and migrate into tissue where they proliferate and mature into macrophages and dendritic cells. GM-CSF is important for the proliferation and maturation of antigen presenting cells, the activity of which is needed for the activation of an anti-tumor immune response. The trial data demonstrated that tumor responses could be seen in injected tumors, and to a lesser extent in uninjected tumors. Responses tended to be highly durable (months-years), and a survival benefit appeared to be achieved in responding patients. Each of these indicated engagement of the immune system in the treatment of cancer in addition to the direct oncolytic effect. However, this and other data with oncolytic viruses generally showed that not all tumors respond to treatment and not all patients achieve a survival advantage. Thus, improvements to the art of oncolytic therapy are clearly needed.

Recently it has been shown that oncolytic immunotherapy can result in additive or synergistic therapeutic effects in conjunction with immune checkpoint blockade (i.e. inhibition or 'antagonism' of immune checkpoint pathways, also termed immune co-inhibitory pathways). Checkpoint (immune inhibitory pathway) blockade is intended to block host immune inhibitory mechanisms which usually serve to prevent the occurrence of auto-immunity. However, in cancer

patients these mechanisms can also serve to inhibit the induction of or block the potentially beneficial effects of any immune responses induced to tumors.

Systemic blockade of these pathways by agents targeting CTLA-4, PD-1 or PD-L1 have shown efficacy in a number of tumor types, including melanoma and lung cancer. However, unsurprisingly, based on the mechanism of action, off target toxicity can occur due to the induction of auto-immunity. Even so, these agents are sufficiently tolerable to provide considerable clinical utility. Other immune co-inhibitory pathway and related targets for which agents (mainly antibodies) are in development include LAG-3, TIM-3, VISTA, CSF1R, IDO, CEACAM1, CD47. Optimal clinical activity of these agents, for example PD1, PDL1, LAG-3, TIM-3, VISTA, CSF1R, IDO, CD47, CEACAM1, may require systemic administration or presence in all tumors due to the mechanism of action, i.e. including targeting of the interface of immune effector cells with tumors or other immune inhibitory mechanisms in/of tumors. In some cases, more localised presence in e.g. just some tumors or in some lymph nodes may also be optimally effective, for example agents targeting CTLA-4.

An alternative approach to increasing the anti-tumor immune response in cancer patients is to target (activate) immune co-stimulatory pathways, i.e. in contrast to inhibiting immune co-inhibitory pathways. These pathways send activating signals into T cells and other immune cells, usually resulting from the interaction of the relevant ligands on antigen presenting cells (APCs) and the relevant receptors on the surface of T cells and other immune cells. These signals, depending on the ligand/receptor, can result in the increased activation of T cells and/or APCs and/or NK cells and/or B cells, including particular subtypes, increased differentiation and proliferation of T cells and/or APCs and/or NK cells and/or B cells, including particular subtypes, or suppression of the activity of immune inhibitory T cells such as regulatory T cells. Activation of these pathways would therefore be expected to result in enhanced anti-tumor immune responses, but it might also be expected that systemic activation of these pathways, i.e. activation of immune responses generally rather than anti-tumor immune responses specifically or selectively, would result in considerable off target toxicity in non-tumor tissue, the degree of such off target toxicity depending on the particular immune co-stimulatory pathway being targeted. Nevertheless agents (mainly agonistic antibodies, or less frequently the soluble ligand to the receptor in question) targeting immune co-stimulatory pathways, including agents targeting GITR, 4-1-BB, OX40, CD40 or ICOS, and intended for systemic use (i.e. intravenous delivery) are in or have been proposed for clinical development.

For many of these approaches targeting immune co-inhibitory or co-inhibitory pathways to be successful, pre-existing immune responses to tumors are needed, i.e. so that a pre-existing immune response can be potentiated or a block to an anti-tumor immune response can be relieved. The presence of an inflamed tumor micro-environment, which is indicative of such an ongoing response, is also needed. Pre-existing immune responses to tumor neo-antigens appear to be particularly important for the activity of immune co-inhibitory pathway blockade and related drugs. Only some patients may have an ongoing immune response to tumor antigens including neoantigens and/or an inflamed tumor microenvironment, both of which are required for the optimal activity of these drugs. Therefore, oncolytic agents which can induce immune responses to tumor antigens, including neoantigens, and/or which can induce an inflamed

tumor microenvironment are attractive for use in combination with immune co-inhibitory pathway blockade and immune potentiating drugs. This likely explains the promising combined anti-tumor effects of oncolytic agents and immune co-inhibitory pathway blockade in mice and humans that have so far been observed.

The above discussion demonstrates that there is still much scope for improving oncolytic agents and cancer therapies utilising oncolytic agents, anti-tumor immune responses and drugs which target immune co-inhibitory or co-stimulatory pathways.

SUMMARY OF THE INVENTION

The invention provides oncolytic viruses expressing GM-CSF and at least one molecule targeting an immune co-stimulatory pathway. GM-CSF aids in the induction of an inflammatory tumor micro-environment and stimulates the proliferation and maturation of antigen presenting cells, including dendritic cells, aiding the induction of an anti-tumor immune responses. These immune responses are amplified through activation of an immune co-stimulatory pathway or pathways using an immune co-stimulatory pathway activating molecule or molecules also delivered by the oncolytic virus.

The use of an oncolytic virus to deliver molecules targeting immune co-stimulatory pathways to tumors focuses the amplification of immune effects on anti-tumor immune responses, and reduces the amplification of immune responses to non-tumor antigens. Thus, immune cells in tumors and tumor draining lymph nodes are selectively engaged by the molecules activating immune co-stimulatory pathways rather than immune cells in general. This results in enhanced efficacy of immune co-stimulatory pathway activation and anti-tumor immune response amplification, and can also result in reduced off target toxicity. It is also important for focusing the effects of combined systemic immune co-inhibitory pathway blockade and immune co-stimulatory pathway activation on tumors, i.e. such that the amplified immune responses from which co-inhibitory blocks are released are antitumor immune responses rather than responses to non-tumor antigens.

The invention utilizes the fact that, when delivered by an oncolytic virus, the site of action of co-stimulatory pathway activation and of GM-CSF expression is in the tumor and/or tumor draining lymph node, but the results of such activation (an amplified systemic anti-tumor-immune response) are systemic. This targets tumors generally, and not only tumors to which the oncolytic virus has delivered the molecule or molecules targeting an immune co-stimulatory pathway or pathways and GM-CSF. Oncolytic viruses of the invention therefore provide improved treatment of cancer through the generation of improved tumor focused immune responses. The oncolytic virus of the invention also offers improved anti-tumor immune stimulating effects such that the immune-mediated effects on tumors which are not destroyed by oncolysis, including micro-metastatic disease, are enhanced, resulting in more effective destruction of these tumors, and more effective long term anti-tumor vaccination to prevent future relapse and improve overall survival.

Anti-tumor efficacy is improved when an oncolytic virus of the invention is used as a single agent and also when the virus is used in combination with other anti-cancer modalities, including chemotherapy, treatment with targeted agents, radiation and, in preferred embodiments, immune

checkpoint blockade drugs (i.e. antagonists of an immune co-inhibitory pathway) and/or agonists of an immune co-stimulatory pathway.

Accordingly, the present invention provides an oncolytic virus comprising: (i) a GM-CSF-encoding gene; and (ii) an immune co-stimulatory pathway activating molecule or immune co-stimulatory pathway activating molecule-encoding gene. The virus may encode more than one immune co-stimulatory pathway activating molecule/gene.

The immune co-stimulatory pathway activating molecule is preferably GITRL, 4-1-BBL, OX40L, ICOSL or CD40L or a modified version of any thereof or a protein capable of blocking signaling through CTLA-4, for example an antibody which binds CTLA-4. Examples of modified versions include agonists of a co-stimulatory pathway that are secreted rather than being membrane bound, and/or agonists modified such that multimers of the protein are formed.

The virus may be a modified clinical isolate, such as a modified clinical isolate of a virus, wherein the clinical isolate kills two or more tumor cell lines more rapidly and/or at a lower dose in vivo than one or more reference clinical isolates of the same species of virus.

The virus is preferably a herpes simplex virus (HSV), such as HSV1. The HSV typically does not express functional ICP34.5 and/or functional ICP47 and/or expresses the US11 gene as an immediate early gene.

The invention also provides:

a pharmaceutical composition comprising a virus of the invention and a pharmaceutically acceptable carrier or diluent;

the virus of the invention for use in a method of treating the human or animal body by therapy;

the virus of the invention for use in a method of treating cancer, wherein the method optionally comprises administering a further anti-cancer agent;

a product of manufacture comprising a virus of the invention in a sterile vial, ampoule or syringe;

a method of treating cancer, which comprises administering a therapeutically effective amount of a virus or a pharmaceutical composition of the invention to a patient in need thereof, wherein the method optionally comprises administering a further anti-cancer agent which is optionally an antagonist of an immune co-inhibitory pathway, or an agonist of an immune co-stimulatory pathway;

use of a virus of the invention in the manufacture of a medicament for use in a method of treating cancer, wherein the method optionally comprises administering a further anti-cancer agent which is optionally an antagonist of an immune co-inhibitory pathway, or an agonist of an immune co-stimulatory pathway;

a method of treating cancer, which comprises administering a therapeutically effective amount of an oncolytic virus, an inhibitor of the indoleamine 2,3-dioxygenase (IDO) pathway and a further antagonist of an immune co-inhibitory pathway, or an agonist of an immune co-stimulatory pathway to a patient in need thereof.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 depicts the structure of an exemplary virus of the invention that comprises a gene encoding GM-CSF and a gene encoding CD40L.

FIG. 2 shows the differential abilities of the eight top ranking HSV1 clinical isolate strains as assessed by crystal violet staining 24 hours or 48 hours after infection with a MG of 0.1, 0.01 or 0.001 as indicated in the Figure to kill

Fadu, SK-mel-28, A549, HT1080, MIA-PA-CA-2, HT29 and MDA-MB-231 human tumor cell lines. The virus strains ranked first and second on each cell line are indicated. The virus RH018A was ranked first on each of the Fadu, HT1080, MIA-PA-CA-2 and HT29 cell lines and second on each of the SK-mel-28, A549 and MDA-MB-231 cell lines. RH004A was ranked joint first with RH018A and RH015A on the HT29 cell line, first on the SK-mel-28 and A549 cell lines and second on the Fadu cell line. RH023A was ranked first on the MDA-MB-231 cell line and second on the HT1080 cell line. RH031A was ranked second on each of the MIA-PA-CA-2 and HT29 cell lines. RH040A was ranked joint second on the HT29 cell line.

FIG. 3 shows a comparison between strain RH018A, the strain ranked first of all the strains tested, with an 'average' strain from the screen (i.e. strain RH065A). Approximately 10 fold less of strain RH018A was needed to kill an equal proportion of cells than was needed of strain RH065A as shown by crystal violet staining 24 or 48 hours post infection with MOIs of 0.1, 0.01 and 0.001 in SK-mel-28, HT1080, MDA-MB-231, Fadu, MIA-PA-CA-2 and A549 cell lines.

FIGS. 4A-4F and 5A-5F depict structures of HSV1 viruses modified by the deletion of ICP34.5 and ICP47 such that the US11 gene is under control of the ICP457 immediate early promoter and containing heterologous genes in the ICP34.5 locus. The viruses were constructed using the RH018A strain unless otherwise stated in the Figure.

FIG. 6 shows the results of an ELISA to detect expression of human or mouse GM-CSF in supernatants from BHK cells infected with virus 16 (mGM-CSF and GALVR-), virus 17 (hGM-CSF and GALVR-) and virus 19 (mGM-CSF).

FIGS. 7A-7B is a comparison between the cell-killing abilities of strain RH018A in which ICP34.5 is deleted and which expresses GALVR- and GFP (virus 10) with a virus that expresses only GFP (virus 12) as determined by crystal violet staining in three cell lines at low magnification.

FIGS. 8A-8B is a comparison between the cell-killing abilities of strain RH018A in which ICP34.5 and ICP47 are deleted and which expresses GALVR- and GM-CSF (virus 17) with a prior art strain with the same modifications as determined by crystal violet staining in four cell lines.

FIG. 9 shows the effectiveness of Virus 16 (ICP34.5 and ICP47 deleted expressing GALVR- and mGM-CSF) in treating mice harbouring A20 lymphoma tumors in both flanks. Tumors on the right flanks were injected with the virus or vehicle and the effects on tumor size was observed for 30 days. The virus was effective against both injected tumors and non-injected tumors.

FIG. 10 demonstrates the effects of Virus 15 (ICP34.5 and ICP47 deleted expressing GALVR- and GFP) and Virus 24 (ICP34.5 and ICP47 deleted expressing GFP) on rat 9L cells in vitro as assessed by crystal violet staining. The virus expressing GALV (Virus 15) showed enhanced killing of rat 9L cells in vitro as compared to a virus which does not express GALV (Virus 24).

FIGS. 11A-11C shows the antitumor effects of Virus 16 in Balb/c mice harboring mouse CT26 tumors in the left and right flanks. Groups of 10 mice were then treated with: Vehicle (3 injections into right flank tumors every other day); 5x10exp6 pfu of Virus 16 (mRP1) injected in the right flank tumor every other day; anti-mouse PD1 alone (10 mg/kg i.p. every three days, BioXCell clone RMP1-14); anti-mouse CTLA-4 (3 mg/kg i.p. every three days, BioXCell clone 9D9); anti-mouse PD1 together with Virus 16; anti-mouse CTLA4 together with Virus 16; 1-methyl tryptophan (1-MT; IDO inhibitor (5 mg/ml in drinking water)); anti-mouse PD1 together with 1-methyl tryptophan; or

anti-mouse PD1 together with 1-methyl tryptophan and Virus 16. Effects on tumor size were observed for a further 30 days. Greater tumor reduction was seen in animals treated with combinations of virus and checkpoint blockade than with the single treatment groups. FIG. 11A shows that using Virus 16 and anti-PD1 in combination has a better anti-tumor effect than using either anti-PD1 or the virus alone. FIG. 11B shows that the anti-tumor effect of Virus 16 in combination with anti-CTLA-4 was better than the anti-tumor effect of either Virus 16 or anti-CTLA-4 alone. FIG. 11C shows that enhanced tumor reduction was observed using Virus 16 together with both anti-PD1 and IDO inhibition as compared to anti-PD1 and 1-MT inhibition in the absence of the virus.

FIGS. 12A-12D shows the enhanced anti-tumor activity of Virus 16 in combination with immune checkpoint blockade in mouse A20 tumors in both flanks of Balb/c mice as compared to either virus alone or checkpoint blockade alone (anti-PD1).

FIG. 13 shows the structure of ICP34.5 and ICP47 deleted viruses expressing GALVR-, GM-CSF and codon optimized anti-mouse or anti-human CTLA-4 antibody constructs (secreted scFv molecules linked to human or mouse IgG1 Fc regions). The scFvs contain the linked ($[G_4S]_3$) light and heavy variable chains from antibody 9D9 (US2011044953; mouse version) and from ipilimumab (US20150283234; human version). The resulting structure of the CTLA-4 inhibitor is also shown.

FIG. 14 shows anti-tumor effects of Virus 16 and Virus 19 in a human xenograft model (A549). There were three injections of Virus 16, Virus 19 or of vehicle over one week at three different dose levels (N=10/group). The doses of the viruses used is indicated. The anti-tumor effects of Virus 16 which expresses GALV were better than those of Virus 19 which does not express GALV.

FIG. 15 demonstrates the effects of viruses of the invention expressing GALVR- on 9L cells in the flanks of Fischer 344 rats. The following treatments were administered to groups of rats (ten per group), into one flank of each rat only three times per week for three weeks: 50 μ l of vehicle; 50 μ l of 10^7 pfu/ml of Virus 19 (expresses mGM-CSF but not GALV R-); or 50 μ l of 10^7 pfu/ml of Virus 16 (expresses both mouse GM-CSF and GALV-R-). Effects on tumor growth were then observed for a further 30 days. Superior tumor control and shrinkage was observed with the virus expressing GM-CSF and GALV-R- as compared to the virus expressing GM-CSF alone.

FIG. 16 shows the anti-tumor effects of viruses expressing anti-mCTLA-4 (virus 27), mCD40L (virus 32), mOX40L (virus 35), m4-2BBL (virus 33), each also with mGM-CSF and GALV-R- compared to virus 16 (expresses GALV and mGM-CSF).

BRIEF DESCRIPTION OF THE SEQUENCE LISTING

SEQ ID NO: 1 is the nucleotide sequence of mouse GM-CSF.

SEQ ID NO: 2 is the nucleotide sequence of a codon optimized version of mouse GM-CSF.

SEQ ID NO: 3 is the nucleotide sequence of human GM-CSF.

SEQ ID NO: 4 is the nucleotide sequence of a codon optimized version of human GM-CSF.

SEQ ID NO: 5 is the amino acid sequence of mouse GM-CSF.

SEQ ID NO: 6 is the amino acid sequence of human GM-CSF.

SEQ ID NO: 7 is the nucleotide sequence of GALV-R-.

SEQ ID NO: 8 is the nucleotide sequence of a codon optimized version of GALV-R- (the first three nucleotides are optional)

SEQ ID NO: 9 is the amino acid sequence of GALV-R-.

SEQ ID NO: 10 is the nucleotide sequence of a codon optimized version of a human membrane bound version of CD40L.

SEQ ID NO: 11 is the amino acid sequence of a human membrane bound version of CD40L.

SEQ ID NO: 12 is the nucleotide sequence of a codon optimized version of a multimeric secreted version of human CD40L.

SEQ ID NO: 13 is the amino acid sequence of a multimeric secreted version of human CD40L.

SEQ ID NO: 14 is the nucleotide sequence of a codon optimized version of a multimeric secreted version of mouse CD40L.

SEQ ID NO: 15 is the amino acid sequence of a multimeric secreted version of mouse CD40L.

SEQ ID NO: 16 is a codon optimized version of the nucleotide sequence of wild-type human CD40L.

SEQ ID NO: 17 is the amino acid sequence of wild-type human CD40L.

SEQ ID NO: 18 is a codon optimized version of the nucleotide sequence of wild-type mouse CD40L.

SEQ ID NO: 19 is the amino acid sequence of wild-type mouse CD40L.

SEQ ID NO: 20 is the nucleotide sequence of a codon optimized version of murine 4-1BBL.

SEQ ID NO: 21 is the nucleotide sequence of a codon optimized version of human 4-1BBL.

SEQ ID NO: 22 is the nucleotide sequence of a codon optimized version of secreted mouse 4-1BBL.

SEQ ID NO: 23 is the nucleotide sequence of a codon optimized version of human secreted 4-1BBL.

SEQ ID NO: 24 is the nucleotide sequence of a codon optimized version of murine GITRL.

SEQ ID NO: 25 is the nucleotide sequence of a codon optimized version of human GITRL.

SEQ ID NO: 26 is the nucleotide sequence of a codon optimized version of secreted murine GITRL.

SEQ ID NO: 27 is the nucleotide sequence of a codon optimized version of secreted human GITRL.

SEQ ID NO: 28 is the nucleotide sequence of a codon optimized version of murine OX40L.

SEQ ID NO: 29 is the nucleotide sequence of a codon optimized version of human OX40L.

SEQ ID NO: 30 is the nucleotide sequence of a codon optimized version of secreted murine OX40L.

SEQ ID NO: 31 is the nucleotide sequence of a codon optimized version of secreted human OX40L.

SEQ ID NO: 32 is the nucleotide sequence of a codon optimized version of murine ICOSL.

SEQ ID NO: 33 is the nucleotide sequence of a codon optimized version of human ICOSL.

SEQ ID NO: 34 is the nucleotide sequence of a murine scFv CTLA-4 antibody. The first six and last eight nucleotides are restriction sites added for cloning purposes.

SEQ ID NO: 35 is the nucleotide sequence of a murine scFv CTLA-4 antibody. The first six and last eight nucleotides are restriction sites added for cloning purposes.

SEQ ID NO: 36 is the nucleotide sequence of the CMV promoter.

SEQ ID NO: 37 is the nucleotide sequence of the RSV promoter.

SEQ ID NO: 38 is the nucleotide sequence of BGH polyA.

SEQ ID NO: 39 is the nucleotide sequence of SV40 late polyA.

SEQ ID NO: 40 is the nucleotide sequence of the SV40 enhancer promoter.

SEQ ID NO: 41 is the nucleotide sequence of rabbit beta-globulin (RBG) polyA.

SEQ ID NO: 42 is the nucleotide sequence of GFP.

SEQ ID NO: 43 is the nucleotide sequence of the MoMuLV LTR promoter.

SEQ ID NO: 44 is the nucleotide sequence of the EF1a promoter.

SEQ ID NO: 45 is the nucleotide sequence of HGH polyA.

DETAILED DESCRIPTION OF THE INVENTION

Oncolytic Virus

The virus of the invention is oncolytic. An oncolytic virus is a virus that infects and replicates in tumor cells, such that the tumor cells are killed. Therefore, the virus of the invention is replication competent. Preferably, the virus is selectively replication competent in tumor tissue. A virus is selectively replication competent in tumor tissue if it replicates more effectively in tumor tissue than in non-tumor tissue. The ability of a virus to replicate in different tissue types can be determined using standard techniques in the art.

The virus of the invention may be any virus which has these properties, including a herpes virus, pox virus, adenovirus, retrovirus, rhabdovirus, paramyxovirus or reovirus, or any species or strain within these larger groups. Viruses of the invention may be wild type (i.e. unaltered from the parental virus species), or with gene disruptions or gene additions. Which of these is the case will depend on the virus species to be used. Preferably the virus is a species of herpes virus, more preferably a strain of HSV, including strains of HSV1 and HSV2, and is most preferably a strain of HSV1. In particularly preferred embodiments the virus of the invention is based on a clinical isolate of the virus species to be used. The clinical isolate may have been selected on the basis of it having particular advantageous properties for the treatment of cancer.

The clinical isolate may have surprisingly good anti-tumor effects compared to other strains of the same virus isolated from other patients, wherein a patient is an individual harbouring the virus species to be tested. The virus strains used for comparison to identify viruses of the invention may be isolated from a patient or an otherwise healthy (i.e. other than harboring the virus species to be tested) volunteer, preferably an otherwise healthy volunteer. HSV1 strains used to identify a virus of the invention are typically isolated from cold sores of individuals harboring HSV1, typically by taking a swab using e.g. Virocult (Sigma) brand swab/container containing transport media followed by transport to the facility to be used for further testing.

After isolation of viruses to be compared from individuals, stocks of the viruses are typically prepared, for example by growing the isolated viruses on BHK or vero cells. Preferably, this is done following no more than 3 cycles of freeze thaw between taking the sample and it being grown on, for example, BHK or vero cells to prepare the virus stock for further use. More preferably the virus sample has undergone 2 or less than 2 cycles of freeze thaw prior to

preparation of the stock for further use, more preferably one cycle of freeze thaw, most preferably no cycles of freeze thaw. Lysates from the cell lines infected with the viruses prepared in this way after isolation are compared, typically by testing for the ability of the virus to kill tumor cell lines in vitro. Alternatively, the viral stocks may be stored under suitable conditions, for example by freezing, prior to testing. Viruses of the invention have surprisingly good anti-tumor effects compared to other strains of the same virus isolated from other individuals, preferably when compared to those isolated from >5 individuals, more preferably >10 other individuals, most preferably >20 other individuals.

The stocks of the clinical isolates identified for modification to produce viruses of the invention (i.e. having surprisingly good properties for the killing of tumor cells as compared to other viral strains to which they were compared) may be stored under suitable conditions, before or after modification, and used to generate further stocks as appropriate.

A clinical isolate is a strain of a virus species which has been isolated from its natural host. The clinical isolate has preferably been isolated for the purposes of testing and comparing the clinical isolate with other clinical isolates of that virus species for a desired property, in the case of viruses of the invention that being the ability to kill human tumor cells. Clinical isolates which may be used for comparison also include those from clinical samples present in clinical repositories, i.e. previously collected for clinical diagnostic or other purposes. In either case the clinical isolates used for comparison and identification of viruses of the invention will preferably have undergone minimal culture in vitro prior to being tested for the desired property, preferably having only undergone sufficient culture to enable generation of sufficient stocks for comparative testing purposes. As such, the viruses used for comparison to identify viruses of the invention may also include deposited strains, wherein the deposited strain has been isolated from a patient, preferably an HSV1 strain isolated from the cold sore of a patient.

The virus may be a modified clinical isolate, wherein the clinical isolate kills two or more tumor cell lines more rapidly and/or at a lower dose in vitro than one or more reference clinical isolate of the same species of virus. Typically, the clinical isolate will kill two or more tumor cell lines within 72 hours, preferably within 48 hours, more preferably within 24 hours, of infection at multiplicities of infection (MOI) of less than or equal to 0.1, preferably less than or equal to an MOI of 0.01, more preferably less than or equal to an MOI of 0.001. Preferably the clinical isolate will kill a broad range of tumor cell lines, such as 2, 3, 4, 5, 6, 7, 8, 9, 10 or, for example, all of the following human tumor cell lines: U87MG (glioma), HT29 (colorectal), LNCaP (prostate), MDA-MB-231 (breast), SK-MEL-28 (melanoma), Fadu (squamous cell carcinoma), MCF7 (breast), A549 (lung), MIAPACA-2 (pancreas), CAPAN-1 (pancreas), HT1080 (fibrosarcoma).

Thus, the virus of the invention may be capable of killing cells from two or more, such as 3, 4, 5, 6, 7 or more, different types of tumor such as two or more, such as 3, 4, 5, 6, 7 or more, solid tumors, including but not limited to colorectal tumor cells, prostate tumor cells, breast tumor cells, ovarian tumor cells, melanoma cells, squamous cell carcinoma cells, lung tumor cells, pancreatic tumor cells, sarcoma cells and/or fibrosarcoma cells.

Tumor cell line killing can be determined by any suitable method. Typically, a sample is first isolated from a patient, preferably, in the case of HSV1, from a cold sore, is used to

11

infect BHK cells, or another suitable cell line such as vero cells. Positive samples are typically identified by the presence of a cytopathic effect (CPE) 24-72 hours post infection, such as 48 hours post infection, and confirmed to be the target viral species by, for example, immunohistochemistry or PCR. Viral stocks are then generated from the positive samples. A sample from the viral stock is typically tested and compared to other samples generated in the same way using swabs from different patients. Testing may be carried out by determining the level of CPE achieved at a range of multiplicity of infection (MOI) and at various times post infection.

For example, cell lines at 80% confluency may be infected with the viral sample at MOI of 1, 0.1, 0.01 and 0.001 and duplicate plates incubated for 24 and 48 hours at 37° C., 5% CO₂ prior to determination of the extent of viral cell killing. This may be determined by, for example, fixing the cells with glutaraldehyde and staining with crystal violet using standard methods. The level of cell lysis may then be assessed by standard methods such as gross observation, microscopy (cell counts) and photography. The method may be repeated with the cells being incubated for shorter time periods, such as 8, 12 or 16 hours, or longer time periods, such as 72 hours, before cell killing is determined, or at additional MOIs such as 0.0001 or less.

Growth curve experiments may also be conducted to assess the abilities of different clinical isolates to replicate in tumor cell lines in vitro. For example, cell lines at 80% confluency may be infected with the viral sample at MOI of 1, 0.1, 0.01 and 0.001 are incubated at 37° C., 5% CO₂ and the cells lysed, typically by freeze/thawing, at 0, 8, 16, 24 and 48 hours post infection prior to determination of the extent of viral cell killing. This may be determined by, for example, assessing viral titres by a standard plaque assay.

A clinical isolate of the invention can kill infected tumor cell lines more rapidly and/or at a lower MOI than the other clinical isolates to which it is compared, preferably 2, 3, 4, 5 or 10 or more, other clinical isolates of the same virus species. The clinical isolate of the invention typically kills a 10%, 25% or 50% greater proportion of the tumor cells present at a particular MOI and time point than at least one, preferably 2, 3, 4, 5 or 10 or more, other clinical isolates of the same virus type at the same MOI and time point to which it was compared. The clinical isolate of the invention typically kills the same or a greater proportion of tumor cells at a MOI that is half or less than half that of the MOI at which one or more, preferably 2, 3, 4, 5, 10 or 15 or more, other clinical isolates of the same virus species used for the comparison at the same time point, typically at 12, 24 and/or 48 hours, kills the same proportion of tumor cells. Preferably, a clinical isolate of the invention typically kills the same or a greater proportion of tumor cells at a MOI that is 5 or 10 times lower than the MOI at which one or more, preferably 2, 3, 4, 5, 10 or 15 or more, other clinical isolates of the same virus used for the comparison at the same time point, typically at 12, 24 and/or 48 hours kills the same proportion of tumor cells. The improved tumor cell killing abilities of a virus of the invention are typically achieved compared to at least 50%, 75% or 90% of the other clinical isolates of the same viral species used for the comparison. The virus is preferably compared to at least 4 other virus strains, such as, for example, 7, 9, 19, 39 or 49 other virus strains of the same species.

The isolated strains may be tested in batches, for example of 4-8 viral strains at a time, on, for example, 4-8 of the tumor cell lines at a time. For each batch of experiments, the degree of killing achieved is ranked on each cell line for the

12

best (i.e. least surviving cells at each time point/MOI) to the worst (i.e. most surviving cells for each time point/MOI) for the viruses being compared in that experiment. The virus strains from each experiment which perform the best across the range of tumor cell line tested (i.e. that consistently ranked as one of the best at killing the cell lines) may then be compared head to head in further experiments using other clinical isolates and/or other tumor cell lines to identify the best virus strains in the total of, for example, >20 virus strains sampled. Those ranked as the best overall are the viruses of the invention.

In a preferred embodiment, the virus of the invention is a strain selected from:

strain RH018A having the provisional accession number ECCAC 16121904;
strain RH004A having the provisional accession number ECCAC 16121902;
strain RH031A having the provisional accession number ECCAC 16121907;
strain RH040B having the provisional accession number ECCAC 16121908;
strain RH015A having the provisional accession number ECCAC 16121903;
strain RH021A having the provisional accession number ECCAC 16121905;
strain RH023A having the provisional accession number ECCAC 16121906; and
strain RH047A having the provisional accession number ECCAC 16121909.

More preferably, the virus of the invention is a strain selected from:

strain RH018A having the provisional accession number ECCAC 16121904;
strain RH004A having the provisional accession number ECCAC 16121902;
strain RH031A having the provisional accession number ECCAC 16121907;
strain RH040B having the provisional accession number ECCAC 16121908; and
strain RH015A having the provisional accession number ECCAC 16121903;

Most preferably, the virus of the invention is strain RH018A having the accession number EACC 16121904.

An HSV of the invention is capable of replicating selectively in tumors, such as human tumors. Typically, the HSV replicates efficiently in target tumors but does not replicate efficiently in non-tumor tissue. This HSV may comprise one or more mutations in one or more viral genes that inhibit replication in normal tissue but still allow replication in tumors. The mutation may, for example, be a mutation that prevents the expression of functional ICP34.5, ICP6 and/or thymidine kinase by the HSV.

In one preferred embodiment, the ICP34.5-encoding genes are mutated to confer selective oncolytic activity on the HSV. Mutations of the ICP34.5-encoding genes that prevent the expression of functional ICP34.5 are described in Chou et al. (1990) Science 250:1262-1266, Maclean et al. (1991) J. Gen. Virol. 72:631-639 and Liu et al. (2003) Gene Therapy 10:292-303, which are incorporated herein by reference. The ICP6-encoding gene and/or thymidine kinase-encoding gene may also be inactivated, as may other genes provided that such inactivation does not prevent the virus infecting or replicating in tumors.

The HSV may contain a further mutation or mutations which enhance replication of the HSV in tumors. The resulting enhancement of viral replication in tumors not only results in improved direct 'oncolytic' tumor cell killing by

the virus, but also enhances the level of heterologous (i.e. a gene inserted into the virus, in the case of viruses of the invention genes encoding GM-CSF and an immune co-stimulatory pathway activating molecule(s)) gene expression and increases the amount of tumor antigen released as tumor cells die, both of which may also improve the immunogenic properties of the therapy for the treatment of cancer. For example, in a preferred embodiment of the invention, deletion of the ICP47-encoding gene in a manner that places the US11 gene under the control of the immediate early promoter that normally controls expression of the ICP47 encoding gene leads to enhanced replication in tumors (see Liu et al., 2003, which is incorporated herein by reference).

Other mutations that place the US11 coding sequence, which is an HSV late gene, under the control of a promoter that is not dependent on viral replication may also be introduced into a virus of the invention. Such mutations allow expression of US11 before HSV replication occurs and enhance viral replication in tumors. In particular, such mutations enhance replication of an HSV lacking functional ICP34.5-encoding genes.

Accordingly, in one embodiment the HSV of the invention comprises a US11 gene operably linked to a promoter, wherein the activity of the promoter is not dependent on viral replication. The promoter may be an immediate early (IE) promoter or a non-HSV promoter which is active in mammalian, preferably human, tumor cells. The promoter may, for example, be a eukaryotic promoter, such as a promoter derived from the genome of a mammal, preferably a human. The promoter may be a ubiquitous promoter (such as a promoter of β -actin or tubulin) or a cell-specific promoter, such as tumor-specific promoter. The promoter may be a viral promoter, such as the Moloney murine leukaemia virus long terminal repeat (MMLV LTR) promoter or the human or mouse cytomegalovirus (CMV) IE promoter. HSV immediate early (IE) promoters are well known in the art. The HSV IE promoter may be the promoter driving expression of ICP0, ICP4, ICP22, ICP27 or ICP47.

The genes referred to above the functional inactivation of which provides the property of tumor selectivity to the virus may be rendered functionally inactive by any suitable method, for example by deletion or substitution of all or part of the gene and/or control sequence of the gene or by insertion of one or more nucleic acids into or in place of the gene and/or the control sequence of the gene. For example, homologous recombination methods, which are standard in the art, may be used to generate the virus of the invention. Alternatively bacterial artificial chromosome (BAC)-based approaches may be used.

As used herein, the term "gene" is intended to mean the nucleotide sequence encoding a protein, i.e. the coding sequence of the gene. The various genes referred to above may be rendered non-functional by mutating the gene itself or the control sequences flanking the gene, for example the promoter sequence. Deletions may remove one or more portions of the gene, the entire gene or the entire gene and all or some of the control sequences. For example, deletion of only one nucleotide within the gene may be made, resulting in a frame shift. However, a larger deletion may be made, for example at least about 25%, more preferably at least about 50% of the total coding and/or non-coding sequence. In one preferred embodiment, the gene being rendered functionally inactive is deleted. For example, the entire gene and optionally some of the flanking sequences may be removed from the virus. Where two or more copies

of the gene are present in the viral genome both copies of the gene are rendered functionally inactive.

A gene may be inactivated by substituting other sequences, for example by substituting all or part of the endogenous gene with a heterologous gene and optionally a promoter sequence. Where no promoter sequence is substituted, the heterologous gene may be inserted such that it is controlled by the promoter of the gene being rendered non-functional. In an HSV of the invention it is preferred that the ICP34.5 encoding-genes are rendered non-functional by the insertion of a heterologous gene or genes and a promoter sequence or sequences operably linked thereto, and optionally other regulatory elements such as polyadenylation sequences, into each the ICP34.5-encoding gene loci.

A virus of the invention is used to express GM-CSF and an immune co-stimulatory pathway activating molecule in tumors. This is typically achieved by inserting a heterologous gene encoding GM-CSF and a heterologous gene encoding the immune co-stimulatory pathway activating molecule in the genome of a selectively replication competent virus wherein each gene is under the control of a promoter sequence. As replication of such a virus will occur selectively in tumor tissue, expression of the GM-CSF and the immune co-stimulatory activating protein by the virus is also enhanced in tumor tissue as compared to non-tumor tissue in the body. Enhanced expression occurs where expression is greater in tumors as compared to other tissues of the body. Proteins expressed by the oncolytic virus would also be expected to be present in oncolytic virus-infected tumor draining lymph nodes, including due to trafficking of expressed protein and of virus in and on antigen presenting cells from the tumor. Accordingly, the invention provides benefits of expression of both GM-CSF and an immune co-stimulatory pathway activating molecule selectively in tumors and tumor draining lymph nodes combined with the anti-tumor effect provided by oncolytic virus replication.

The virus of the invention comprises GM-CSF. The sequence of the gene encoding GM-CSF may be codon optimized so as to increase expression levels of the respective proteins in target cells as compared to if the unaltered sequence is used.

The virus of the invention comprises one or more immune co-stimulatory pathway activating molecules and/or one or more genes encoding an immune co-stimulatory pathway activating molecule. Immune co-stimulatory pathway activating molecules include proteins and nucleic acid molecules (e.g. aptamer sequences). Examples of immune co-stimulatory pathway activating molecules include CD40 ligand, GITR ligand, 4-1-BB ligand, OX40 ligand, ICOS ligand, flt3 ligand, TL1A, CD30 ligand, CD70 and single chain antibodies targeting the respective receptors for these molecules (CD40, GITR, 4-1-BB, OX40, ICOS, flt3, DR3, CD30, CD27). The CD40L, GITRL, 4-1-BBL, OX40L, ICOSL, flt3L, TL1A, CD30L or CD70L may be a modified version of any thereof, such as a soluble version.

Activators of immune co-stimulatory pathways include mutant or wild type, soluble, secreted and/or membrane bound ligands, and agonistic antibodies including single chain antibodies. Viruses of the invention preferably encode one or more of CD40L, ICOSL, 4-1-BBL, GITRL or OX40L.

The inhibitor of a co-inhibitory pathway may be a CTLA-4 inhibitor. The CTLA-4 inhibitor is typically a molecule such as a peptide or protein that binds to CTLA-4 and reduces or blocks signaling through CTLA-4, such as by reducing activation by B7. By reducing CTLA-4 signalling,

the inhibitor reduces or removes the block of immune stimulatory pathways by CTLA-4.

The CTLA-4 inhibitor is preferably an antibody or an antigen binding fragment thereof. The term "antibody" as referred to herein includes whole antibodies and any antigen binding fragment (i.e., "antigen-binding portion") or single chains thereof. An antibody refers to a glycoprotein comprising at least two heavy (H) chains and two light (kappa) (L) chains inter-connected by disulfide bonds, or an antigen binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR). The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e.g., effector cells) and the first component (C1q) of the classical complement system.

The antibody is typically a monoclonal antibody. The antibody may be a chimeric antibody. The antibody is preferably a humanised antibody and is more preferably a human antibody.

The term "antigen-binding fragment" of an antibody refers to one or more fragments of an antibody that retain the ability to specifically bind to CTLA-4. The antigen-binding fragment also retains the ability to inhibit CTLA-4 and hence to reduce or remove the CTLA-4 blockade of a stimulatory immune response. Examples of suitable fragments include a Fab fragment, a F(ab')₂ fragment, a Fab' fragment, a Fd fragment, a Fv fragment, a dAb fragment and an isolated complementarity determining region (CDR). Single chain antibodies such as scFv and heavy chain antibodies such as VHH and camel antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. In a preferred embodiment, the antibody is an scFv. Examples of suitable scFv molecules are disclosed in, for example, WO2007/123737 and WO2014/066532, which are incorporated herein by reference. The scFv may be encoded by the nucleotide sequence shown in SEQ ID NO: 34 the nucleotide sequence shown in SEQ ID NO: 35.

Viruses of the invention may encode one or more immune co-stimulatory pathway activating molecules, preferably 1, 2, 3 or 4 immune co-stimulatory pathway activating molecules, more preferably 1 or 2 immune co-stimulatory pathway activating molecules.

For example, the virus may comprise genes encoding:

CD40L and one or more of ICOSL, 4-1-BBL, GITRL, OX40L and a CTLA-4 inhibitor;
 ICOSL and one or more of CD40L, 4-1-BBL, GITRL, OX40L and a CTLA-4 inhibitor;
 4-1-BBL and one or more of CD40L, ICOSL, GITRL, OX40L and a CTLA-4 inhibitor;
 GITRL and one or more of CD40L, ICOSL, 4-1-BBL, OX40L and a CTLA-4 inhibitor;
 OX40L and one or more of CD40L, ICOSL, 4-1-BBL, GITRL and a CTLA-4 inhibitor;
 a CTLA-4 inhibitor and one or more of CD40L, ICOSL, 4-1-BBL, GITRL and OX40L.

The sequence of the gene encoding the immune co-stimulatory activating molecule may be codon optimized so

as to increase expression levels of the respective protein(s) in target cells as compared to if the unaltered sequence is used.

The virus of the invention may comprise one or more further heterologous genes in addition to GM-CSF and an immune co-stimulatory pathway activating molecule, including, in a preferred embodiment, a fusogenic protein such as GALVR-.

The fusogenic protein may be any heterologous protein capable of promoting fusion of a cell infected with the virus of the invention to another cell. A fusogenic protein, preferably a wild type or modified viral glycoprotein (i.e. modified to increase its fusogenic properties), is a protein which is capable in inducing the cell to cell fusion (syncytia formation) of cells in which it is expressed. Examples of fusogenic glycoproteins include VSV-G, syncitin-1 (from human endogenous retrovirus-W (HERV-W)) or syncitin-2 (from HERVFRDE1), paramyxovirus SV5-F, measles virus-H, measles virus-F, RSV-F, the glycoprotein from a retrovirus or lentivirus, such as gibbon ape leukemia virus (GALV), murine leukemia virus (MLV), Mason-Pfizer monkey virus (MPMV) and equine infectious anemia virus (EIAV) with the R transmembrane peptide removed (R-versions). In a preferred embodiment the fusogenic protein is from GALV and has the R- peptide removed (GALV-R-).

The virus of the invention may optionally comprise multiple copies of the fusogenic protein-encoding gene, preferably 1 or 2 copies. The virus may comprise two or more different fusogenic proteins, including any of the fusogenic proteins listed above.

The fusogenic protein or proteins optionally expressed by a virus of the invention may be identical to a naturally occurring protein, or may be a modified protein.

The fusogenic protein-encoding gene (fusogenic gene) may have a naturally occurring nucleic acid sequence or a modified sequence. The sequence of the fusogenic gene may, for example, be modified to increase the fusogenic properties of the encoded protein, or to provide codon optimisation and therefore increase the efficiency of expression of the encoded protein.

The invention also provides a virus, such as a pox virus or a HSV, preferably HSV1, which expresses at least three heterologous genes, wherein each of the three heterologous genes is driven by a different promoter selected from the CMV promoter, the RSV promoter, the EF1a promoter, the SV40 promoter and a retroviral LTR promoter. The virus may, for example, express four heterologous genes, wherein each of the four heterologous genes is driven by a different promoter selected from the CMV promoter, the RSV promoter, the EF1a promoter, the SV40 promoter and a retroviral LTR promoter. The retroviral LTR is preferably from MMLV (SEQ ID NO:43), also known as MoMuLV. The heterologous genes may be terminated by poly adenylation sequences. The poly adenylation sequences may be the same or different. Preferably each heterologous gene is terminated by a different poly adenylation sequence, which is preferably selected from the BGH, SV40, HGH and RBG poly adenylation sequences.

The invention also provides a virus, such as a pox virus or a HSV, preferably HSV1, which expresses at least three heterologous genes, wherein each of the three heterologous genes is terminated by a different poly adenylation sequence selected from the BGH, SV40, HGH and RBG poly adenylation sequences. The virus may, for example, express four heterologous genes terminated by each of the BGH, SV40, HGH and RBG poly adenylation sequences, respectively.

Production of Virus

Viruses of the invention are constructed using methods well known in the art. For example plasmids (for smaller viruses and single and multiple genome component RNA viruses) or BACs (for larger DNA viruses including herpes viruses) encoding the viral genome to be packaged, including the genes encoding the fusogenic and immune stimulating molecules under appropriate regulatory control, can be constructed by standard molecular biology techniques and transfected into permissive cells from which recombinant viruses can be recovered.

Alternatively, in a preferred embodiment plasmids containing DNA regions flanking the intended site of insertion can be constructed, and then co-transfected into permissive cells with viral genomic DNA such that homologous recombination between the target insertion site flanking regions in the plasmid and the same regions in the parental virus occur. Recombinant viruses can then be selected and purified through the loss or addition of a function inserted or deleted by the plasmid used for modification, e.g. insertion or deletion of a marker gene such as GFP or lacZ from the parental virus at the intended insertion site. In a most preferred embodiment the insertion site is the ICP34.5 locus of HSV, and therefore the plasmid used for manipulation contains HSV sequences flanking this insertion site, between which are an expression cassette encoding GM-CSF and the immune co-stimulatory pathway activating molecule. In this case, the parental virus may contain a cassette encoding GFP in place of ICP34.5 and recombinant virus plaques are selected through the loss of expression of GFP. In a most preferred embodiment the US11 gene of HSV is also expressed as an IE gene. This may be accomplished through deletion of the ICP47-encoding region, or by other means.

The GM-CSF encoding sequences and immune co-stimulatory pathway activating molecule encoding sequences are inserted into the viral genome under appropriate regulatory control. This may be under the regulatory control of natural promoters of the virus species of the invention used, depending on the species and insertion site, or preferably under the control of heterologous promoters. Suitable heterologous promoters include mammalian promoters, such as the IEF2a promoter or the actin promoter. More preferred are strong viral promoters such as the CMV IE promoter, the RSV LTR, the MMLV LTR, other retroviral LTR promoters, or promoters derived from SV40. Preferably each exogenous gene (e.g. encoding the GM-CSF and immune co-stimulatory pathway activating molecule) will be under separate promoter control, but may also be expressed from a single RNA transcript, for example through insertion of an internal ribosome entry sites (IRES) between protein coding sequences. RNA derived from each promoter is typically terminated using a polyadenylation sequence (e.g. mammalian sequences such as the bovine growth hormone (BGH) poly A sequence, synthetic polyadenylation sequences, the rabbit betaglobin polyadenylation sequence, or viral sequences such as the SV40 early or late polyadenylation sequence).

Pharmaceutical Compositions

The invention provides a pharmaceutical composition comprising the virus and a pharmaceutically acceptable carrier or diluent. Suitable carriers and diluents include isotonic saline solutions, for example phosphate-buffered saline. The composition may further comprise other constituents such as sugars or proteins to improve properties such as stability of the product. Alternatively a lyophilized formulation may be used, which is reconstituted in a pharmaceutically acceptable carrier or diluent before use.

The choice of carrier, if required, is frequently a function of the route of delivery of the composition. Within this invention, compositions may be formulated for any suitable route and means of administration. Pharmaceutically acceptable carriers or diluents are those used in compositions suitable for intra-tumoral administration, intravenous/intraarterial administration, administration into the brain or administration into a body cavity (e.g. bladder, pleural cavity or by intraperitoneal administration). The composition may be administered in any suitable form, preferably as a liquid.

The present invention also provides a product of manufacture comprising a virus of the invention in a sterile vial, ampoule or syringe.

Medical Uses/Methods of Treatment

The invention provides the virus of the invention for use in the treatment of the human or animal body by therapy, particularly for use in a method of treating cancer. The cancer is typically in a mammal, preferably in a human. The virus kills infected tumour cells by lysis and by causing infected tumor cells to fuse with one another. The virus of the invention also elicits a systemic anti-tumor immune response, augmented through the expression of GM-CSF and the immune co-stimulatory pathway activating molecule, which also kills cancer cells.

The invention also provides a method of treating cancer, the method comprising administering a therapeutically effective amount of the virus of the invention to an individual in need thereof.

The invention additionally provides the use of the virus of the invention in the manufacture of a medicament for treating cancer.

The virus of the invention is particularly useful in treating any solid tumor including any adenocarcinoma, carcinoma, melanoma or sarcoma. For example, the virus of the invention is useful in treating head and neck, prostate, breast, ovarian, lung, liver, endometrial, bladder, gall bladder, pancreas, colon, kidney, stomach/gastric, esophageal, or cervical cancers, mesothelioma, melanoma or other skin cancer, lymphoma, glioma or other cancer of the nervous system, or sarcomas such as soft tissue sarcoma.

The virus of the invention may be used to treat malignant tumors, including tumors that have metastasised from the site of the original tumor. In this embodiment, the virus may be administered to the primary tumor or to one or more secondary tumors.

The virus of the invention may be administered in combination with other therapeutic agents, including chemotherapy, targeted therapy, immunotherapy (including immune checkpoint blockade, i.e. administration of one or more antagonist of an immune co-inhibitory pathway, and/or one or more agonist of an immune co-stimulatory pathway) and/or in combination with radiotherapy and/or in combination with any combination of these. The therapeutic agent is preferably an anti-cancer agent.

The virus of the invention may be administered in combination with a second virus, such as a second oncolytic virus.

For example, the therapeutic agent may comprise an immunogen (including a recombinant or naturally occurring antigen, including such an antigen or combination of antigens delivered as DNA or RNA in which it/they are encoded), to further stimulate an immune response, such as a cellular or humoral immune response, to tumor cells, particularly tumor neoantigens. The therapeutic agent may be an agent intended to increase or potentiate an immune response, such as a cytokine, an agent intended to inhibit an

immune checkpoint pathway or stimulate an immune potentiating pathway or an agent which inhibits the activity of regulatory T cells (Tregs) or myeloid derived suppressor cells (MDSCs).

The therapeutic agent may be an agent known for use in an existing cancer therapeutic treatment. The therapeutic agent may be radiotherapy or a chemotherapeutic agent. The therapeutic agent may be selected from cyclophosphamide, alkylating-like agents such as cisplatin or melphalan, plant alkaloids and terpenoids such as vincristine or paclitaxel (Taxol), antimetabolites such as 5-fluorouracil, topoisomerase inhibitors type I or II such as camptothecin or doxorubicin, cytotoxic antibiotics such as actinomycin, anthracyclines such as epirubicin, glucocorticoids such as triamcinolone, inhibitors of protein, DNA and/or RNA synthesis such as methotrexate and dacarbazine, histone deacetylase (HDAC) inhibitors, or any other chemotherapy agent.

The therapeutic agent may be one, or a combination of: immunotherapeutics or immunomodulators, such as TLR agonists; agents that down-regulate T-regulatory cells such as cyclophosphamide; or agents designed to block immune checkpoints or stimulate immune potentiating pathways, including but not limited to monoclonal antibodies, such as a CTLA-4 inhibitor, a PD-1 inhibitor, a PD-L1 inhibitor, a LAG-3 inhibitor, a TIM-3 inhibitor, a VISTA inhibitor, a CSF1R inhibitor, an IDO inhibitor, a CEACAM1 inhibitor, a GITR agonist, a 4-1-BB agonist, a KIR inhibitor, a SLAMF7 inhibitor, an OX40 agonist, a CD40 agonist, an ICOS agonist or a CD47 inhibitor. In a preferred embodiment, the therapeutic agent is a CTLA-4 inhibitor such as an anti-CTLA-4 antibody, a PD1 inhibitor, such as an anti-PD-1 antibody or a PD-L1 inhibitor such as an anti-PD-L1 antibody. Such inhibitors, agonists and antibodies can be generated and tested by standard methods known in the art.

Immunotherapeutic agents may also include bi-specific antibodies, cell based-therapies based on dendritic cells, NK cells or engineered T cells such CAR-T cells or T cells expressing engineered T cell receptors. Immunotherapeutic agents also include agents that target a specific genetic mutation which occurs in tumors, agents intended to induce immune responses to specific tumor antigens or combinations of tumor antigens, including neoantigens and/or agents intended to activate the STING/cGAS pathway, TLR or other innate immune response and/or inflammatory pathway, including intra-tumoral agents.

For example, a virus of the invention may be used: in combination with dacarbazine, a BRAF inhibitor and or CTLA-4, PD1 or PD-L1 blockade to treat melanoma; in combination with taxol, doxorubicin, vinorelbine, cyclophosphamide and/or gemcitabine to treat breast cancer; in combination with 5-fluorouracil and optionally leucovorin, irinotecan and/or oxaliplatin to treat colorectal cancer; in combination with taxol, carboplatin, vinorelbine and/or gemcitabine, PD-1 or PD-L1 blockade to treat lung cancer; in combination with cisplatin and/or radiotherapy to treat head and neck cancer.

The therapeutic agent may be an inhibitor of the indoleamine 2,3-dioxygenase (IDO) pathway. Examples of IDO inhibitors include epacadostat (INCB024360), 1-methyl-tryptophan, Indoximod (1-methyl-D-tryptophan), GDC-0919 or F001287.

The mechanism of action of IDO in suppressing anti-tumor immune responses may also suppress immune responses generated following oncolytic virus therapy. IDO expression is induced by toll like receptor (TLR) activation and interferon- γ both of which may result from oncolytic

virus infection. One embodiment of the use of oncolytic virus therapy for cancer treatment includes combination of an oncolytic virus, including a virus expressing GM-CSF and an immune co-stimulatory pathway activating molecule or molecules with an inhibitor of the IDO pathway and optionally one or more antagonist of an immune co-inhibitory pathway and/or one or more agonist of an immune co-stimulatory pathway, including those targeting CTLA-4, PD-1 and/or PD-L1.

The invention also provides a method of treating cancer, which comprises administering a therapeutically effective amount of an oncolytic virus, an inhibitor of the indoleamine 2,3-dioxygenase (IDO) pathway and a further antagonist of an immune co-inhibitory pathway, and/or an agonist of an immune co-stimulatory pathway to a patient in need thereof.

The oncolytic virus is preferably a modified clinical isolate. The oncolytic virus is preferably a pox virus, more preferably a HSV, such as a HSV1 and/or a HSV rendered functionally inactive for ICP34.5 and/or ICP47.

The oncolytic virus may express an immune stimulating molecule, such as GM-CSF and/or a co-stimulatory pathway encoding molecule such as CD40L, GITRL, OX40L, 4-1-BBL or ICO5L, and/or an inhibitor of CTLA-4, and/or a fusogenic protein, such as the GALV fusogenic glycoprotein with the R sequence mutated or deleted.

The further antagonist of an immune co-inhibitory pathway is preferably an antagonist of CTLA-4, an antagonist of PD1 or an antagonist of PD-L1. For example, the further antagonist of an immune co-inhibitory pathway may be an inhibitor of the interaction between PD1 and PD-L1.

Where a therapeutic agent and/or radiotherapy is used in conjunction with a virus of the invention, administration of the virus and the therapeutic agent and/or radiotherapy may be contemporaneous or separated by time. The composition of the invention may be administered before, together with or after the therapeutic agent or radiotherapy. The method of treating cancer may comprise multiple administrations of the virus of the invention and/or of the therapeutic agent and/or radiotherapy. In preferred embodiments, in the case of combination with immune checkpoint blockade or other immune potentiating agents, the virus of the invention is administered once or multiple times prior to the concurrent administration of the immune checkpoint blockade or other immune potentiating agent or agents thereafter, or concurrent with the administration of the immune checkpoint blockade or other immune potentiating agent or agents without prior administration of the virus of the invention.

The virus of the invention may be administered to a subject by any suitable route. Typically, a virus of the invention is administered by direct intra-tumoral injection. Intra-tumoral injection includes direct injection into superficial skin, subcutaneous or nodal tumors, and imaging guided (including CT, MRI or ultrasound) injection into deeper or harder to localize deposits including in visceral organs and elsewhere. The virus may be administered into a body cavity, for example into the pleural cavity, bladder or by intra-peritoneal administration. The virus may be injected into a blood vessel, preferably a blood vessel supplying a tumor.

Therapeutic agents which may be combined with a virus of the invention can be administered to a human or animal subject in vivo using a variety of known routes and techniques. For example, the composition may be provided as an injectable solution, suspension or emulsion and administered via parenteral, subcutaneous, oral, epidermal, intradermal, intramuscular, interarterial, intraperitoneal, intravenous injection using a conventional needle and syringe, or

using a liquid jet injection system. The composition may be administered topically to skin or mucosal tissue, such as nasally, intratracheally, intestinally, sublingually, rectally or vaginally, or provided as a finely divided spray suitable for respiratory or pulmonary administration. In preferred

embodiments, the compositions are administered by intravenous infusion, orally, or directly into a tumor. The virus and/or therapeutic agent may be administered to a subject in an amount that is compatible with the dosage composition that will be therapeutically effective. The administration of the virus of the invention is for a "therapeutic" purpose. As used herein, the term "therapeutic" or "treatment" includes any one or more of the following as its objective: the prevention of any metastasis or further metastasis occurring; the reduction or elimination of symptoms; the reduction or complete elimination of a tumor or cancer; an increase in the time to progression of the patient's cancer; an increase in time to relapse following treatment; or an increase in survival time.

Therapeutic treatment may be given to Stage I, II, III, or IV cancers, preferably Stage II, III or IV, more preferably Stage III or IV, pre- or post-surgical intervention (i.e. following recurrence or incomplete removal of tumors following surgery), preferably before any surgical intervention (either for resection of primary or recurrent/metastatic disease), or following recurrence following surgery or following incomplete surgical removal of disease, i.e. while residual tumor remains.

Therapeutic treatment may be carried out following direct injection of the virus composition into target tissue which may be the tumor, into a body cavity, or a blood vessel. As a guide, the amount of virus administered is in the case of HSV in the range of from 10^4 to 10^{10} pfu, preferably from 10^5 to 10^9 pfu. In the case of HSV, an initial lower dose (e.g. 10^4 to 10^7 pfu) may be given to patients to seroconvert patients who are seronegative for HSV and boost immunity in those who are seropositive, followed by a higher dose then being given thereafter (e.g. 10^6 to 10^9 pfu). Typically up to 20 ml of a pharmaceutical composition consisting essentially of the virus and a pharmaceutically acceptable suitable carrier or diluent may be used for direct injection into tumors, or up to 50 ml for administration into a body cavity (which may be subject to further dilution into an appropriate diluent before administration) or into the bloodstream. However for some oncolytic therapy applications larger or smaller volumes may also be used, depending on the tumor and the administration route and site.

The routes of administration and dosages described are intended only as a guide since a skilled practitioner will be able to determine readily the optimum route of administration and dosage. The dosage may be determined according to various parameters, especially according to the location of the tumor, the size of the tumor, the age, weight and condition of the patient to be treated and the route of administration. Preferably the virus is administered by direct injection into the tumor or into a body cavity. The virus may also be administered by injection into a blood vessel. The optimum route of administration will depend on the location and size of the tumor. Multiple doses may be required to achieve an immunological or clinical effect, which, if required, will be typically administered between 2 days to 12 weeks apart, preferably 3-days to 3 weeks apart. Repeat doses up to 5 years or more may be given, preferably for up to one month to two years dependent on the speed of response of the tumor type being treated and the response of a particular patient, and any combination therapy which may also be being given.

The following Examples illustrate the invention.

Example 1. Construction of a Virus of the Invention

The virus species used to exemplify the invention is HSV, specifically HSV1. The strain of HSV1 used for exemplification is identified through the comparison of more than 20 primary clinical isolates of HSV1 for their ability to kill a panel of human tumor-derived cell lines and choosing the virus strain with the greatest ability to kill a broad range of these rapidly, and at low dose. Tumor cell lines used for this comparison include U87MG (glioma), HT29 (colorectal), LNCaP (prostate), MDA-MB-231 (breast), SK-MEL-28 (melanoma), Fadu (squamous cell carcinoma), MCF7 (breast), A549 (lung), MIAPACA-2 (pancreas), CAPAN-1 (pancreas), and/or HT1080 (fibrosarcoma).

Specifically, each primary clinical isolate of HSV is titrated onto each of the cell lines used for screening at MOIs such as 1, 0.1, 0.01 and 0.001 and assessed for the extent of cell death at time points such as 24 and 48 hrs at each dose. The extent of cell killing may be assessed by e.g. microscopic assessment of the proportion of surviving cells at each time point, or e.g. a metabolic assay such as an MTT assay. The exemplary virus of the invention is then constructed by deletion of ICP47 from the viral genome using homologous recombination with a plasmid containing regions flanking HSV1 nucleotides 145300 to 145582 (HSV1 nucleotides 145300 to 145582 being the sequences to be deleted; HSV1 strain 17 sequence Genbank file NC 001806.2) between which are encoded GFP. GFP expressing virus plaques are selected, and GFP then removed by homologous recombination with the empty flanking regions and plaques which do not express GFP are selected. This results in an ICP47 deleted virus in which US11 is expressed as an IE protein as it is now under the control of the ICP47 promoter. ICP34.5 is then deleted using homologous recombination with a plasmid containing regions flanking HSV1 nucleotides 124953 to 125727 (HSV1 nucleotides 124953 to 125727 being the sequences to be deleted; HSV1 strain 17 sequence Genbank file NC 001806.2) between which GFP is encoded. GFP expressing virus plaques are again selected, and GFP then removed by homologous recombination with the same flanking regions but between which are now an expression cassette comprising a codon optimized version of the mouse GM-CSF sequence, a codon optimized version of the GALV R- sequence and codon optimized version of mouse soluble multimeric CD40L driven by a CMV, an RSV and an SV40 promoter. Non-GFP expressing plaques are selected.

The structure of the resulting virus is shown in FIG. 1. The mGM-CSF, CD40L and GALV-R- sequences are shown in SEQ ID NOs 2, 14 and 8 respectively. The structure of the resulting virus is confirmed by restriction digestion and Southern blot, GM-CSF and CD40L expression is confirmed by ELISA, and GALV-R- expression is confirmed by infection of human HT1080 tumor cells and the observation of syncytial plaques.

Viruses are also constructed using similar procedures which only have inserted the gene for GALV-R- or mouse GM-CSF and GALV-R-, but without CD40L. The structures of these viruses are also shown in FIG. 1.

For human use, hGM-CSF and hCD40L are used, the sequence for codon optimised versions of which are shown in SEQ ID NO 4 and 13.

23

Example 2. The Effect of the Combined Expression
of GM-CSF and an Immune Co-Stimulatory
Pathway Activating Molecule from an Oncolytic
Virus in Mouse Tumor Models

The GALV R- protein causes cell to cell fusion in human cells but not in mouse cells because the PiT-1 receptor required for cell fusion to occur has a sequence in mice which does not allow cell fusion to occur. As a result mouse tumor cells expressing human PiT-1 are first prepared using methods standard in the art. Human PiT-1 is cloned into a lentiviral vector also comprising a selectable marker gene. The vector is transfected into target CT26 mouse colorectal cancer tumor cells and clones resistant to the selectable marker are selected to generate CT26/PiT-1 cells. PiT-1 expression is confirmed by western blotting in untransfected cells and in cells transfected with the PiT-1 expressing lentivirus and by transfection of a plasmid expressing GALV-R- and confirmation that cell fusion occurs.

The utility of the invention is demonstrated by administering CT26/PiT-1 cells into both flanks of Balb/c mice and allowing the CT26/PiT-1 tumors to grow to approximately 0.5 cm in diameter.

The following treatments are then administered to groups of mice (five per group), into one flank of each mouse only 3 times per week for two weeks:

- 50 μ l of saline (1 group);
- 50 μ l of 10^5 pfu/ml, 10^6 pfu/ml, or 10^7 pfu/ml of the ISV with only GALVR- inserted (3 groups);
- 50 μ l of 10^5 pfu/ml, 10^6 pfu/ml, or 10^7 pfu/ml of the HSV with only GALVR- and mouse GM-CSF inserted (3 groups);
- 50 μ l of 10^5 pfu/ml, 10^6 pfu/ml, or 10^7 pfu/ml of the virus with GALVR- and both mouse GM-CSF and CD40L inserted (3 groups).

Effects on tumor growth are then observed for up to one month. Superior tumor control and shrinkage in both injected and uninjected tumors with the virus expressing GM-CSF and CD40L as compared to the other groups is observed, including through an improved dose response curve.

Example 3. The Effect of Combined Expression of
GM-CSF and an Immune Co-Stimulatory Pathway
Activating Molecule from an Oncolytic Virus on
the Therapeutic Effect of Immune Checkpoint
Blockade in Mouse Tumor Models

The experiment in Example 2 above is repeated but mice are additionally dosed bi-weekly by the intra-peritoneal route with an antibody targeting mouse PD-1 (10 mg/kg; Bioxcell RMP-1-14 on the same days as virus dosing) or an antibody targeting mouse CTLA-4 (10 mg/kg; Bioxcell 9H10 on the same days as virus dosing). An additional group of mice is added which receive no antibody treatment. More specifically, groups of mice receive (1) saline, (2) HSV with GALVR- inserted as in Example 2, (3) HSV with GM-CSF and GALV-R- inserted as in Example 2, (4) HSV with GM-CSF, CD40L and GALV-R- inserted as in Example 2, (5) PD-1 antibody, (6) CTLA-4 antibody, (7) HSV with GALV-R- inserted plus PD-1 antibody, (8) HSV with GALV-R- inserted gene plus CTLA-4 antibody, (9) HSV with GM-CSF and GALV-R- and PD-1 antibody or (10) HSV with GM-CSF and GALV-R- and CTLA-4 antibody (11) HSV with GM-CSF, CD40L and GALV-R- and PD-1 antibody or (12) HSV with GM-CSF, CD40L and GALV-R- and CTLA-4 antibody. Superior tumor control and shrinkage in

24

both injected and uninjected tumors with the virus expressing GM-CSF and CD40L together with the anti-PD-1 antibody or the anti-CTLA-4 antibody as compared to the other groups is observed, including through an improved dose response curve.

Example 4. Collection of Clinical Isolates

The virus species used to exemplify the invention is HSV, specifically HSV1. To exemplify the invention, 181 volunteers were recruited who suffered from recurrent cold sores. These volunteers were given sample collection kits (including Sigma Virovult collection tubes), and used these to swab cold sores when they appeared following which these samples were shipped to Replimune, Oxford UK. From June 2015-February 2016, swabs were received from 72 volunteers. A sample of each swab was used to infect BHK cells. Of these 36 live virus samples were recovered following plating out and growth on BHK cells. These samples are detailed in Table 1.

TABLE 1

Details of Tested Swab Samples & Result	
Sample Number	Virus retrieved
RH001A	No
RH001B	
RH002A	Yes
RH003A	No
RH004A	Yes
RH004B	
RH005A	No
RH005B	
RH006A	No
RH006B	
RH007A	Yes
RH007B	
RH007C	
RH008A	No
RH008B	
RH008C	
RH009A	No
RH009B	
RH010A	No
RH011A	No
RH011B	
RH011C	
RH012A	No
RH013A	No
RH014A	Yes
RH014B	
RH015A	Yes
RH016A	No
RH016B	
RH017A	Yes
RH018A	Yes
RH018B	
RH018C	
RH019A	No
RH019B	
RH019C	
RH020A	Yes-RH020A only
RH020B	
RH020C	
RH021A	Yes
RH021B	
RH022A	Yes
RH022B	
RH023A	Yes
RH024A	No
RH025A	Yes-RH025B only
RH025B	
RH026A	Yes
RH027A	No
RH027B	

25

TABLE 1-continued

Details of Tested Swab Samples & Result	
Sample Number	Virus retrieved
RH027C	
RH028A	No
RH028B	
RH028C	
RH029A	No
RH030A	No
RH031A	Yes-RH031A to
RH031B	RH031D
RH031C	
RH031D	
RH031E	
RH031F	
RH032A	No
RH033A	No
RH033B	
RH033C	
RH034A	No
RH034B	
RH034C	
RH035A	No
RH036A	Yes
RH037A	Yes
RH038A	Yes
RH039A	No
RH039B	
RH039C	
RH040A	Yes
RH040B	
RH040C	
RH041A	Yes
RH042A	Yes
RH043A	No
RH043B	
RH043C	
RH044A	No
RH045A	No
RH046A	Yes
RH047A	Yes-RH047A and
RH047B	RH047C
RH047C	
RH048A	No
RH049A	No
RH049B	
RH049C	
RH050A	No
RH051A	Yes
RH051B	
RH052A	Yes-RH052A only
RH052B	
RH053A	No
RH054A	No
RH055A	No
RH055B	
RH056A	Yes
RH057A	No
RH058A	Yes
RH058B	
RH059A	No
RH060A	No
RH061A	Yes
RH062A	No
RH063A	No
RH064A	Yes
RH065A	Yes
RH065B	
RH066A	No
RH067A	No
RH067B	
RH068A	No-contaminated
RH069A	No
RH069A	
RH070A	Yes
RH071A	Yes
RH072A	No
RH073A	Yes
RH073B	

26

TABLE 1-continued

Details of Tested Swab Samples & Result	
Sample Number	Virus retrieved
RH074A	No
RH074B	
RH075A	No
RH076A	No
RH078A	No
RH078B	
RH079B	Yes
RH079B	
RH080A	No
RH081A	Yes
RH082A	No
RH082B	
RH083A	Yes
RH083B	
RH084A	Yes
RH084B	
RH084C	
RH085A	No
RH086A	No
RH087A	Yes-RH078B only
RH087B	

Designations A, B, C etc. indicate multiple swabs from the same volunteer.

Example 5. Identification of Clinical Isolates with Improved Anti-Tumor Effects

The abilities of the primary clinical isolates of HSV1 to kill a panel of human tumor-derived cell lines was tested. The tumor cell lines used for this comparison were HT29 (colorectal), MDA-MB-231 (breast), SK-MEL-28 (melanoma), Fadu (squamous cell carcinoma), MCF7 (breast), A549 (lung), MIAPACA-2 (pancreas) and HT1080 (fibrosarcoma). The cell lines were used to test for the level of CPE achieved at a range of MOI and times post infection for each of the primary clinical isolates.

Experiments were conducted in parallel using 5 to 8 of the new viruses strains at the same time. The virus strains were plated out in duplicate at a range of MOIs (0.001-1), and the extent of CPE following crystal violet staining was assessed at 24 and 48 hours following infection. The viral strains which were most effective at killing the tumor cell lines were scored, and the most effective two or three strains from each screen of 5-8 strains were identified and compared in parallel in a further experiment to identify the top strains for further development.

The initial screens demonstrated substantial variability in the ability of the different strains to kill the different tumor cell lines. Of an initial 29 strains tested, 8 strains of interest were identified in the initial screens for further comparison. These were strains RH004A, RH015A, RH018A, RH021A, RH1023A, RH31A, RH040A, and RH047A.

The 8 strains for further comparison were tested in parallel on the panel of tumor cell lines, and their relative ability to kill these tumor cell lines was assessed following crystal violet staining and observation for CPE. FIG. 2 shows a representative dose point and MOI for these viruses on each of the viruses on each of the cell lines demonstrating the differential ability of the viruses to kill the target tumor cell lines observed.

There was substantial variation amongst the strains, and it was found that while a particular strain may be particularly effective at killing one cell line, it is not necessarily particularly effective at killing other cell lines too, further demonstrating the degree of variability in the ability of clinical strains of HSV to kill tumor cells of different types.

FIG. 3 also indicates which of the virus strains was both best and second best at killing each of the cell lines, enabling the virus strains to be rank ordered as to their overall relative ability to kill the panel of cell lines as a whole. This analysis demonstrated that strains RH004A, RH015A, RH018A, RH031A and RH040A were relatively more effective than the other strains, and these five strains were chosen for potential further development as oncolytic agents. Of these top five strains, the relative rank order based on their abilities to kill across the panel of cell lines was RH1018A>RH004A>RH031A>RH040A>RH015A.

More specifically, in these experiments, the tumor cell lines were used to seed multi-well tissue culture plates so that they were about 80% confluent on the day of infection. Representative wells from each tumor cell line were trypsinised and the number of cells in the well determined. These cell counts are used to determine the volume of each clinical isolate required to give an MOI of 1, 0.1, 0.01 and 0.001. Separate wells of a tumor cell line were infected with the clinical isolate at these MOI. All infections are carried out in quadruplicate. Duplicate wells were incubated for 24 hours and duplicate wells were incubated for 48 hours, both at 37° C., 5% CO₂, prior to fixation of the cells with glutaraldehyde and staining with crystal violet. The level of cell lysis was then assessed by gross observation, microscopy (cell counts) and photography.

Strain RH018A, the strain ranked first of all the strains tested was compared to an 'average' strain from the screen (i.e. a strain which was not in the top 8, but was also not in the group of strains which were least effective and killing the panel of tumor cell lines). This comparison showed that Strain RH018A was approximately 10 fold more effective than this average strain (Strain RH065A) at killing the tumor cell lines (i.e. approximately 10 fold less of Strain RH018A was needed to kill an equal proportion of cells than was needed of Strain RH065A). This is shown in FIG. 3.

Example 6. Modification of Clinical Isolates

In this Example the clinical isolates selected in Example 5 were modified by deletion of ICP34.5 from the viral genome using homologous recombination with a plasmid containing regions flanking the ICP34.5 encoding gene (nucleotides 143680-145300 and 145,582-147,083; HSV1 strain 17 sequence Genbank file NC 001806.2) between which are encoded GFP and the GALV-R-fusogenic glycoprotein. The structure of this virus, (Virus 10) is shown in FIGS. 4A-4F.

Additional viruses based on Strain RH018A were also constructed in which both ICP34.5 and ICP47 (using flanking regions containing nucleotides 123464-124953 and 125727-126781; HSV1 strain 17 sequence Genbank file NC 001806.2) were deleted (resulting in placement of US11 under the control of the ICP47 promoter). To construct these viruses, GFP expressing virus plaques, with GFP expressed in place of ICP47 were first selected. GFP was then removed by homologous recombination with the empty flanking regions, and plaques not expressing GFP were selected. This resulted in an ICP47 deleted virus in which US11 is expressed as an IE protein as it is now under the control of the ICP47 promoter. ICP34.5 was then deleted using homologous recombination with a plasmid containing regions flanking HSV1 nucleotides 143680-145300 and 145,582-147,083; HSV1 strain 17 sequence Genbank file NC 001806.2) between which GFP is encoded. GFP expressing virus plaques were again selected, and GFP then removed by homologous recombination with the same

flanking regions but between which are now an expression cassette comprising the genes to be inserted. The viruses that were constructed are shown in FIGS. 1, 4 and 5. These included a codon optimized version of the mouse GM-CSF sequence and a codon optimized version of the GALV R-sequence driven by the CMV IE promoter and RSV promoter respectively, in a back to back orientation and again selecting virus plaques which do not express GFP. This virus construction was performed using methods which are standard in the art.

The mGM-CSF and GALV-R- sequences are shown in SEQ ID NOs 2 and 8 respectively. The structure of the resulting virus was confirmed by PCR, GM-CSF expression was confirmed by ELISA, and GALV-R- expression was confirmed by infection of human HT1080 tumor cells and the observation of syncytial plaques.

For human use, hGM-CSF is used, the sequence for a codon optimised version of which is shown in SEQ ID NO 4. The structure of this virus is shown in FIGS. 4A-4F. Expression of mouse or human GM-CSF from viruses 16, 17 and 19 is shown in FIG. 6.

Example 7. A Virus of the Invention Modified for Oncolytic Use and Expressing a Fusogenic Glycoprotein Shows Enhanced Tumor Cell Killing In Vitro as Compared to a Virus which does not Express a Fusogenic Glycoprotein

Virus 10 (see FIGS. 4A-4F), based on clinical Strain RH018A in which ICP34.5 is deleted and which expresses GALV-R- and GFP, was compared in vitro to a virus which expresses only GFP Virus 12). Virus 10 showed enhanced killing on a panel of human tumor cell lines as compared to Virus 12, as shown in FIGS. 7A-7B.

Example 8. A Virus of the Invention Modified for Oncolytic Use Shows Enhanced Tumor Cell Killing as Compared to a Similarly Modified Virus which is not of the Invention

Virus 17 (see FIGS. 4A-4F), based on clinical Strain RH018A in which ICP34.5 and ICP47 are deleted and which expresses GALV-R- and GM-CSF, was compared in vitro to a known virus which was also deleted for ICP34.5 and ICP47 but which was not derived from a strain of the invention and which expresses only GM-CSF. Virus 17 showed enhanced killing on a panel of human tumor cell lines as compared to the previous virus, as shown in FIGS. 8A-8B.

Example 9. A Virus of the Invention Modified for Oncolytic Use Effectively Treats Mouse Tumors In Vivo

Virus 16 was tested in mice harboring A20 lymphoma tumors in the left and right flanks. One million tumor cells were first implanted in both flanks of Balb/c mice and tumors allowed to grow to 0.5-0.7 cm in diameter. Tumors on the right flank were then injected 3 times (every other day) with either vehicle (10 mice) or 5x10exp6 pfu of Virus 16 (10 mice), and effects on tumor size observed for a further 30 days. This demonstrated that both injected and uninjected tumors were effectively treated with Virus 16 (see FIG. 9).

Example 10. The Effect of the Combined Expression of a Fusogenic Protein and an Immune Stimulatory Molecule from an Oncolytic Virus of the Invention in a Rat Tumor Model

The GALV R- protein causes cell to cell fusion in human cells but not in mouse cells. However, GALV R- does cause fusion in rat cells.

The utility of the invention was further demonstrated by administering 9L cells into the flanks of Fischer 344 rats and allowing the 9L tumors to grow to approximately 0.5 cm in diameter.

The following treatments were then administered to groups of rats (ten per group), into one flank only of each rat three times per week for three weeks:

- 50 μ l of vehicle;
- 50 μ l of 10^7 pfu/ml of Virus 19 (expresses mGM-CSF but not GALV R-);
- 50 μ l of 10^7 pfu/ml of Virus 16 (expresses both mouse GM-CSF and GALV-R-).

Effects on tumor growth were then observed for a further ≈ 30 days. This demonstrated superior tumor control and shrinkage with the virus expressing GALV-R- in both injected and uninjected tumors, demonstrating improved systemic effects. This is shown in FIG. 15. FIG. 10 shows that a virus expressing GALV (Virus 15) also shows enhanced killing of rat 91 cells in vitro as compared to a virus which does not express GALV (Virus 24).

Example 11. A Virus of the Invention Modified for Oncolytic Use is Synergistic with Immune Checkpoint Blockade in Mouse Tumor Models

Virus 16 was tested in mice harboring CT26 tumors in the left and right flanks. One million tumor cells were first implanted in both flanks of Balb/c mice and tumors allowed to grow to 0.5-0.6 cm in diameter.

Groups of 10 mice were then treated with:

- Vehicle (3 injections into right flank tumors every other day);
- 5×10^6 pfu of Virus 16 injected in the right flank tumor every other day;
- anti-mouse PD1 alone (10 mg/kg i.p. every three days, BioXCell clone RMP1-14);
- anti-mouse CTLA-4 (3 mg/Kg i.p every three days, BioXCell clone 9D9);
- anti-mouse PD1 together with Virus 16;
- anti-mouse CTLA4 together with Virus 16;
- 1-methyl tryptophan (IDO inhibitor (5 mg/ml in drinking water));
- anti-mouse PD1 together with 1-methyl tryptophan;
- anti-mouse PD1 together with 1-methyl tryptophan and Virus 16;

Effects on tumor size were observed for a further 30 days. A greater tumor reduction in animals treated with combinations of virus and checkpoint blockade was demonstrated than in animals treated with the single treatment groups (see FIGS. 11A-11C). Enhanced tumor reduction with Virus 16 together with both anti-PD1 and IDO inhibition was also demonstrated as compared to Virus 16 together with only anti-PD1 (see FIGS. 11A-11C).

Enhanced activity of Virus 16 in combination with immune checkpoint blockade was also seen in A20 tumors (FIGS. 12A-12D).

Example 12. The Effect of the Expression of a Fusogenic Protein Front an Oncolytic Virus of the Invention in Human Xenograft Models in Immune Deficient Mice

The GALV R- protein causes cell to cell fusion in human cells but not in mouse cells. However, human xenograft tumors grown in immune deficient mice can be used to assess the effects of GALV expression on anti-tumor efficacy.

The utility of the invention was therefore further demonstrated by administering A549 human lung cancer cells into the flanks of nude mice and allowing the tumors to grow to approximately 0.5 cm in diameter.

The following treatments were then administered to groups of mice (ten per group), into tumor containing flank of each mouse three times over one week:

- 50 μ l of vehicle;
- 50 μ l of 10^7 pfu/ml of Virus 16 (expresses both mouse GM-CSF and GALV-R-);
- 50 μ l of 10^6 pfu/ml of Virus 16;
- 50 μ l of 10^5 pfu/ml of Virus 16;
- 50 μ l of 10^7 pfu/ml of Virus 19 (expresses only mouse GM-CSF);
- 50 μ l of 10^6 pfu/ml of Virus 19;
- 50 μ l of 10^5 pfu/ml of Virus 19.

Effects on tumor growth were then observed for a further ≈ 30 days. This experiment demonstrated superior tumor control and shrinkage with the virus expressing GALV-R- in both tumor models (see FIG. 14).

Example 13. Expression of Two Immune Stimulatory Molecules from a Virus Expressing a Fusogenic Protein

Viruses similar to the GALV-R- and mGM-CSF expressing virus described above (Virus 16) were constructed, but additionally expressing mouse versions of CD40L (virus 32), ICOSL (virus 36), OX40L (virus 35), 4-1BBL (virus 33) and GITRL (virus 34). Here, instead of using a plasmid containing ICP34.5 flanking regions and an expression cassette comprising GM-CSF and GALV-R- driven by a CMV and an RSV promoter, a plasmid containing ICP34.5 flanking regions and an expression cassette comprising GM-CSF, GALV and the additional proteins driven by a CMV, an RSV and an MMLV promoter respectively were used for recombination with a virus containing GM-CSF, GALV and GFP inserted into ICP34.5. Non-GFP expressing plaques were again selected. Correct insertion was confirmed by PCR, and expression by western blotting and/or ELISA for the additional inserted gene. These viruses are shown in FIGS. 5A-5F. Similarly, viruses expressing anti-mouse and anti-human CTLA-4 in addition to GALV and mGM-CSF were also constructed (Viruses 27 and 31 in FIGS. 5A-5F and see also FIG. 13). Effects of viruses expressing anti-mouse CTLA-4 (virus 27), mCD40L (virus 32), m4-1BBL (virus 33) or mOX40L (virus 35) in addition to mGM-CSF and GALV-R- in vivo is shown in FIG. 16 which showed enhanced activity in A20 tumors as compared to virus 16 (expresses mGM-CSF and GALV-R-). In these experiments tumors were induced in both flanks of mice, and virus or vehicle injected only into the right flank tumor. The dose of virus used was 5×10^4 pfu (50 μ l of 1×10^6 pfu/ml in each case), given three times over one week. This dose level of virus is subtherapeutic for uninjected tumors for virus 16, which allows the benefits of the delivery of the additional molecules encoded by viruses 27, 32, 33 and 35 to clearly be seen.

Deposit Information

The following HSV1 strains were deposited at the ECACC, Culture Collections, Public Health England, Porton Down, Salisbury, SP4 0JG, United Kingdom on 19 Dec. 2016 by Replimune Limited and were allocated the indicated provisional accession numbers:

- RH004A—Provisional Accession Number 16121902
- RH015A—Provisional Accession Number 16121903
- RH018A—Provisional Accession Number 16121904
- RH021A—Provisional Accession Number 16121905
- RH023A—Provisional Accession Number 16121906
- RH031A—Provisional Accession Number 16121907
- RH040B—Provisional Accession Number 16121908
- RH047A—Provisional Accession Number 16121909.

SEQUENCE LISTING

Sequence total quantity: 45

SEQ ID NO: 1 moltype = DNA length = 426
 FEATURE Location/Qualifiers
 source 1..426
 mol_type = other DNA
 organism = Mus musculus

SEQUENCE: 1
 atgtggctgc agaatttact ttctctgggc attgtggtct acagcctctc agcaccacc 60
 cgctcaccca tcaactgtcac ccggccttgg aagcatgtag aggccatcaa agaagccctg 120
 aacctcctgg atgacatgcc tgtcacattg aatgaagagg tagaagtcgt ctctaacgag 180
 ttctccttca agaagctaac atgtgtgcag acccgctga agatattcga gcagggtcta 240
 cggggcaatt tcaccaaact caagggcgcc ttgaacatga cagccagcta ctaccagaca 300
 tactgcccc caactccgga aacggactgt gaaacacaag ttaccaccta tgcggatttc 360
 atagacagcc ttaaaacctt tctgactgat atcccccttg aatgcaaaaa accagtccaa 420
 aatga 426

SEQ ID NO: 2 moltype = DNA length = 426
 FEATURE Location/Qualifiers
 source 1..426
 mol_type = other DNA
 organism = Mus musculus

SEQUENCE: 2
 atgtggctcc agaacctcct ctctctcggt atcgctggtt attcactctc cgcacctact 60
 cgctcaccta tcaactgtcac cagaccttgg aagcacgtgg aggccatcaa ggaggctctg 120
 aacctgctgg acgatatgcc agtgaccttg aacgaggagg tggagggtgt gagcaacgag 180
 ttctccttta agaagctgac ctgctgacag acaaggctga agatcttcga gcagggctg 240
 agaggaaact ttaccaagct gaagggcgcc ctgaacatga ccgcttctta ctaccagaca 300
 tactgcccc ctaccccgga gacagactgt gagacacagg tgaccacata cgcgacttc 360
 attgatagcc tgaaaacatt cctgaccgac attccatttg agtgaagaa gcccgtcag 420
 aagtaa 426

SEQ ID NO: 3 moltype = DNA length = 435
 FEATURE Location/Qualifiers
 source 1..435
 mol_type = other DNA
 organism = Homo sapiens

SEQUENCE: 3
 atgtggctgc agagcctgct gctcttgggc actgtggcct gcagcatctc tgcaccgcgc 60
 cgctcgccca gcccagcac gcagccctgg gagcatgtga atgccaatca ggaggcccg 120
 cgtctcctga acctgagtag agacactgct gctgagatga atgaaacagt agaagtcac 180
 tcagaaatgt ttgacctcca ggagccgacc tgcctacaga cccgcctgga gctgtacaag 240
 cagggcctgc ggggcagcct caccaagctc aagggcccct tgaccatgat ggccagccac 300
 tacaagcagc actgcccctc aaccccgga acttctctgt caaccacgat tatcaccttt 360
 gaaagtttca aagagaacct gaaggacttt ctgcttgta tccccttga ctgctgggag 420
 ccagtcagg agtga 435

SEQ ID NO: 4 moltype = DNA length = 435
 FEATURE Location/Qualifiers
 source 1..435
 mol_type = other DNA
 organism = Homo sapiens

SEQUENCE: 4
 atgtggctgc agtccctgct gctgctgggc accgtcgctt gttctatttc cgcaccgcga 60
 aggtcaccaa gtccatctac tcagccttgg gagcacgtga acgcaatca ggaggcacgg 120
 cggctgctga acctgagcgg ggacaccgcc gccgagatga acgagacagt ggaagtgtac 180
 agcgagatgt tcgacttgca ggagccaccc tgcctgcaga caaggctgga gctgtacaag 240
 cagggcctgc gccgctctct gaccaagctg aagggccac tgacaatgat ggccagccac 300
 tataagcagc actgcccccc taccctcgag acaagctgtg ccaccacgat catcacattc 360
 gagtccttta aggagaacct gaaggatttt ctgctggta ttccatttga ttgttgggag 420
 cccgtccagg agtaa 435

SEQ ID NO: 5 moltype = AA length = 141
 FEATURE Location/Qualifiers
 source 1..141
 mol_type = protein
 organism = Mus musculus

SEQUENCE: 5
 MWLQNLFLPLG IVVYSLSAPT RSPITVTRPW KHVEAIKEAL NLLDDMPVTL NEEVEVVSNE 60
 FSFKKLTCVQ TRLKIFEQGL RGNFTKLKGA LNMTASYVQT YCPPTPETDC ETQVTYADF 120
 IDSLKTFLTD IPFECKKPVQ K 141

SEQ ID NO: 6 moltype = AA length = 144
 FEATURE Location/Qualifiers
 source 1..144
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 6
 MWLQSLLLLG TVACISAPA RSPSPSTQPW EHVNAIQEAR RLLNLSRDTA AEMNETVEVI 60

-continued

```

SEMFDLQEPT CLQTRLELYK QGLRGSLSLTKL KGPLTMMASH YKQHCPTPE TSCATQIITF 120
ESFKENLKDF LLVIPDFCWE PVQE 144

SEQ ID NO: 7          moltype = DNA length = 2010
FEATURE              Location/Qualifiers
source                1..2010
                     mol_type = other DNA
                     organism = Gibbon leukemia virus

SEQUENCE: 7
atggtattgc tgcctgggtc catgcttctc acctcaaac tgcaccacct tcggcaccag 60
atgagtcctg ggagctggaa aagactgata atcctcttaa gctgcgtatt cggcggcggc 120
gggacgagtc tgcaaaataa gaacccccac cagcccatga cctcacttg gcaggtaactg 180
tcccaactgt gagagcttgt ctgggataca aaggcagtc agcccccttg gacttggttg 240
ccacactta aacctgatgt atgtgccttg gcggctagtc ttgagtcctg ggatatcccg 300
ggaaaccgatg tctcgtcctc taaacgagtc agacctccgg actcagacta tactgcccgt 360
tataagcaaa tcacctgggg agccataggg tgcagctacc ctccgggctag gactagaatg 420
gcaagctctc cctctacgt atgtcccccg gatggccgga ccttttcaga agctagaagg 480
tgccggggggc tagaatccct atactgtaaa gaatgggatt gtgagaccac ggggaccggt 540
tattggctat ctaaatccctc aaaaagacctc ataactgtaa aatgggacca aaatagcgaa 600
tggactcaaa aattttcaaca gtgtcaccag accggctggg gtaacccctc taaaatagat 660
ttcacagaca aaggaaaatt atccaaggac tggataacgg gaaaaacctg gggattaaga 720
ttctatgtgt ctggacatcc agcgctacag ttcaccattc gcttaaaaat caccaacatg 780
ccagctgttg cagtaggttc tgacctcgtc cttgtggaac aaggacctcc tagaacgttc 840
ctcgtctctc cacctcctct tcccccaagg gaagcgccac cgccatctct ccccgactct 900
aaactccacag cctcggcgac tagtgcaaca actcccacgg tgagaaaaac aattgttacc 960
ctaaacactc cgctcccccac cacaggcgac agactttttg atcttgtgca gggggccttc 1020
ctaaccttaa atgtaccaca cccagggggc actgagctct gctggctttg tttggccatg 1080
ggccccccctt attatgaagc aatagcctca tcaggagagg tcgcctactc caccgacctt 1140
gaccgggtgc cttgggggac ccaaggaaaag ctcacctcca ctgaggtctc agggacacgg 1200
ttgtgcatag gaaaggtgcc ctttaccatc cagcatctct gcaatcagac cctatccatc 1260
aattcctcgg gagaccatca gtactgtctc cctccaacc atagctgggt ggcttgacg 1320
actggcctca ccccttgctc ctccacctca gtttttaac agactagaga tttctgtatc 1380
caggtccagc tgattcctcg catctattac tatcctgaag aagttttgtt acaggcctat 1440
gacaattctc accccaggac taaaagagag gctgtctcac ttaccctagc tgtttactg 1500
gggttgggaa tcacggcggg aataggtact ggttcaactg ccttaattaa aggacctata 1560
gacctccagc aaggcctgac aagcctccag atcgccatag atgtgacct cggggccctc 1620
caagactcag tcagcaagtt agaggactca ctgacttccc tgcagaggt agtgctccaa 1680
aataggagag gccttgactt gctgtttcta aaagaagggt gcctctgtgc gggcctaaag 1740
gaagagtgtc gtttttacat agaccactca ggtgcagtag gggactccat gaaaaaactc 1800
aaagaaaaaa tggataaaaag acagttagag cgccagaaaa gccaaaactg gtaggaagga 1860
tgggtcaata actcccttg gtccactacc ctgctatcaa ccatcgctgg gcccctatta 1920
ctcctccttc tgttgctcat cctcgggcca tgcacatcaa ataagttagt tcaatccatc 1980
aatgatagga taagtgcagt taaaatttaa 2010

SEQ ID NO: 8          moltype = DNA length = 2013
FEATURE              Location/Qualifiers
source                1..2013
                     mol_type = other DNA
                     organism = Gibbon leukemia virus

SEQUENCE: 8
accatggctc tgctgcctgg gtctatgtct ctgacttcta acctgcacca cctgcgacac 60
cagatgtctc ccgggtcatg gaacgggctg atcatcctgc tgagctgcgt gttcggagga 120
ggaggcacct cctgcagaaa caagaatcct caccagccaa tgacctgac atggcagggt 180
ctgtcccaga caggcgacgt ggtgtgggat accaaggcag tgcagccacc ttggacatgg 240
tggcccaacc tgaagcctga cgtgtgcgcc ctggccgcct cctggagtc ttgggacatc 300
ccgggcacag acgtgagcag cgtcaagagg gtgagaccac ccgactctga ttatacagcc 360
gcctacaagc agatcacctg gggcgccatc ggctgtagct atcctcgggc ccgcacaagg 420
atggccagct ccacctttta cgtgtgccca cgcaacggaa ggacctgtc tgaggcaagg 480
agatgtggcg gcctggagag cctgtattgc aaggagtggg attgtgagac cacaggcaca 540
ggctactggc tgtctaagtc tagcaaggac ctgatcaccg tgaagtggga tcagaacagc 600
gagtggaacac agaagttcca cagatgccac cagaccggct ggtgtaatcc cctgaagatc 660
gactttacag ataagggcaa gctgtccaag gactggatca ccggcaagac atggggcctg 720
agattctacg tgtctggcca cctggcgtg cagtttataa tccgggtgaa gatcaccaac 780
atgccagcag tggcagtggg accagacctg gtgctgggtg agcagggacc tccacgcacc 840
tccctggccc tgcctccctc actgccccct agggaggccc caccctctag cctgcccgat 900
tctaacagca cagccctggc cactccgcc cagaccctca cagtgcgcaa gaccatcgtg 960
acactgaata cccaccccc taccacaggc gacaggctgt tcatctggt gcaggggccc 1020
ttctcgacac tgaaagccac caatcctggc gcaaccgaga gctgctggct gtgcctggct 1080
atgggcccac cctactatga ggcaatcgcc tctctggag aggtggcata ttccacagac 1140
ctggatagat gcagatgggg caccagggc aagctgacct tgacagaggt gtctggccac 1200
ggcctgtgca tcggcaaggt gccattcaca caccagcacc tgtgcaacca gacctgagc 1260
atcaatagct ccggcgacca ccagtacctg ctgccaagca accactcctg gtgggcacgc 1320
tccacaggac tgaccccatg tctgagcacc agcgtgttca accagaccag agacttttgt 1380
atccagggtg agctgtacct tattcccag aggaggtgct gctgcaggcc 1440
tatgataatt cccacccaag aacaaaggag gaggccgtgt ccttgacctt ggcctgtgct 1500
ctgggactgg gaatcacagc aggaatcggc acaggcagca ccgccctgat caagggacca 1560
atcgacctgc agcagggact gacctccctg cagatcgcca tcgacgccga tctgagagcc 1620
ctgcaggaca cgtgtgcca gctggaggat tctctgacct cctgagcgca ggtgggtgct 1680
cagaacagga ggggcctgga cctgctgttc ctgaaggagg gaggactgtg cgccgcctg 1740

```


-continued

```

aaggaggagt gctgttttta tategaccac tctggcgccg tgcgggtag catgaagaag 1800
ctgaaggaga agctggataa gcgccagctg gagaggcaga agagccagaa ttggtacgag 1860
ggctggttca acaattcccc ctggtttacc acaactgctg ctaccatcgc aggacctctg 1920
ttattactgc tgctgctgct gatccctggg ccatgtatca tcaacaagct ggtgcagttt 1980
atcaacgacc gaatctccgc agtgaaaatc taa 2013

```

```

SEQ ID NO: 9      moltype = AA length = 669
FEATURE          Location/Qualifiers
source           1..669
                  mol_type = protein
                  organism = Gibbon leukemia virus

```

```

SEQUENCE: 9
MVLPGSMLL  TSNLHHLRHQ MSPGSWKRLI  ILLSCVFGGG  GTSLQKNKPH  QPMTLTWQVL  60
SQTGDVVWDT KAVQPPPTWW  PTLKPDVCL  AASLESWDIP  GTDVSSSKRV  RPPDSDYTAA  120
YKQITWGAIG CSYPRARTRM ASSTFYVCPR DGRTLSEARR  CGGLESLYCK  EWCETTTGTG  180
YWLKSSKSDL ITVKWDQNSE  WTQKFQQCHQ  TGWCNPLKID  FTDKGKLSKD  WITGKTWGLR  240
FYVSGHPGVQ FTIRLKITNM  PAVAVGPDV  LVEQGPPRTS  LALPPPLPPR  EAPPPSLPDS  300
NSTALATSAQ TPTVRKTIVT LNTPPPPTGD  RLFDLVQGAF  LTLNATNPGA  TESCWLCLAM  360
GPPYVEAIAS SGEVAYSTD  DRCRWGTQ GK  LTLTEVSGHG  LCIGKVPFTH  QHLCNQTL SI  420
NSSGDHQYLL PSNHSWWACS  TGLTPCLSTS  VFNQTRDFCI  QVQLIPRIYY  YPEEVLLQAY  480
DNSHPRTKRE AVSLTLAVLL  GLGITAGIGT  GSTALIKGPI  DLQQGLTSLQ  IAIDADLRAL  540
QDSVSKLEDS LTVSLSEVVL  NRRGLDLLFL  KEGGLCAALK  EECFYIDHS  GAVRDSMKKL  600
KEKLDKRQLE RQKSNQWYEG  WFNNSPWFTT  LLSTIAGPLL  LLLLLLILGP  CIINKLVQFI  660
NDRISAVKI  669

```

```

SEQ ID NO: 10     moltype = DNA length = 759
FEATURE          Location/Qualifiers
misc_feature      1..759
                  note = Homo sapiens
source           1..759
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 10
atgatcgaga cctacaatca gacaagccca cggtcgcgcg caaccggact gcctatcagc 60
atgaagatct tcatgtacct gctgaccgtg tttctgatca cacagatgat cggctccgcc 120
ctgttcgcgcg tgtatctgca caggagactg gacaagatcg aggatgagcg caatctgcac 180
gaggacttcg tgtttatgaa gaccatccag cggtgcaaca caggcgagag gagcctgtct 240
ctgctgaatt gtgaggagat caagtcccag ttcgagggct ttgtgaagga tatcatgctg 300
aacaaggagg agacaaagaa ggacgaggat ccacagatcg cagcacacgt ggtgtccgag 360
gcaaaactcta atgcgcgcag cgtgctgcag tgggccaaga agggctacta taccatgaag 420
tctaactctg tgacactgga gaatggcaag cagctgaccg tgaagaggca gggcctgtac 480
tatatctatg cccagggtgac attctgctct aacagagagg caagctccca ggcacccttc 540
atcgtgggac tgtggctgaa gccctctagc ggcagcgaga ggatcctgct gaaggccgcc 600
aatacccact cctctagcca gctgtgcgag cagcagtgca tccacctggg aggcgtgttc 660
gagctgcagc ctggagccag cgtgttcgtg aacgtgacag acccatctca ggtgagccac 720
ggcaccggct tcacaagctt tggcctgctg aagctgtga 759

```

```

SEQ ID NO: 11     moltype = AA length = 252
FEATURE          Location/Qualifiers
REGION          1..252
                  note = Homo sapiens
source           1..252
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 11
MIETYNQTSR RSAATGLPIS MKIFMYLLTV FLITQMIGSA LFAVYLHRLR DKIEDERNLH 60
EDFVFMKTIQ RCNTGERSLS LLNCEEIKSQ FEGFVKDIDL NKEETKKDED PQIAAHVVSE 120
ANSNAASVLQ WAKKGYITMK SNLVTLENGK QLTVKRQGLY YIIAQVTFCS NREASSQAPF 180
IVGLWLKPSG GSERILLKAA NTHSSSQLCE QQSIIHLGGVF ELQPGASVVF NVTDPQSQVSH 240
GTGFTSFGLL KL 252

```

```

SEQ ID NO: 12     moltype = DNA length = 1416
FEATURE          Location/Qualifiers
source           1..1416
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 12
atgtgcctt  ttctgagcat  gctgggtgctg  ctgtgagcag  ctctgggaaa  cctgggagcc  60
gagatgaaga  gcctgtccca  gagatctgtg  cctaacacct  gcacactggt  catgtgcagc  120
cccaccgaga  atggactgcc  tggaaaggac  ggaagggatg  gaaggaggag  ccctcggggc  180
gagaaggcgc  acccaggact  gcctggacca  atgggactga  gcggactgca  gggaccaaca  240
ggacctgtgg  gaccaaagg  agagaacgga  tccgccggag  agccaggccc  taagggcgag  300
aggggcctgt  ctggccccc  tggcctgcca  ggcacccag  gcccgcgcg  caaggagggc  360
ccatccggca  agcagggcaa  tatcgcccc  cagggcaagc  ctggcccaaa  gggcgaggca  420
ggaccaaagg  gagaagtggg  agcacctggc  atgcagggat  ccaccggagc  aaagggatct  480
acaggaccaa  agggcgagcg  cggcgcccc  ggcgtgcagg  gcgcccccg  caatgcagga  540
gcagcaggac  cagcaggacc  tgcaggccca  cagggcgcgc  ctggctctag  gggcccaccc  600
ggcctgaagg  gcgacagggg  agtgcctggc  gataggggca  tcaagggaga  gagcggactg  660
ccagattccg  ccgcctgag  gcagcagatg  gaggccctga  agggcaagct  gcagaggctg  720

```

-continued

```

gaggtggcct tctccacta ccagaaggcc gccctgttcc cagacggcca caggagactg 780
gacaagatcg aggtgagcg caacctgcac gaggatctcg tgtttatgaa gacctaccag 840
agatgcaaca caggcgagcg gtctctgagc ctgctgaatt gtgaggagat caagtctcag 900
ttcgagggct ttgtgaagga catcatgctg aacaaggagg agaccaagaa ggagaatagc 960
ttcgagatgc agaagggcga tcagaatccc cagatcgagc cacacgtgat cagcgaggca 1020
agctccaaga ccacatccgt gctgcagtgg gccgagaagg gctactatac catgtccaac 1080
aatctggtga cactggagaa cggcaagcag ctgaccgtga agagacaggg cctgtactat 1140
atctatgccc aggtgacatt ctgctctaatt cgggaggcct ctagccaggg cctttttatc 1200
gcctctctgt gcctgaagag cccaggcaga ttcgagcgga tccctgtcag gcccgccaac 1260
acccactcct ctgccaagcc atgcggagag cagagcatcc acctgggagg cgtgttcgag 1320
ctgcagccag gagctccgt gtttgtgaat gtgacagacc catcccaggt gtctcacgga 1380
accggttca catcctttgg cctgctgaag ctgtga 1416

```

```

SEQ ID NO: 13      moltype = AA  length = 471
FEATURE           Location/Qualifiers
source            1..471
                  mol_type = protein
                  organism = Homo sapiens

```

```

SEQUENCE: 13
MLPFLSMLVL LVQPLGNLGA EMKSLSQRSV PNTCTLVMCS PTENGLPGRD GRDGREGPRG 60
EKGDPLPLPG MGLSGLQGPT GPVGPKEGNG SAGEPGPKGE RGLSGPPGLP GIPGPAGKEG 120
PSGKQGNIGP QGKPGPKGEA GPKGEVGPAG MQGSTGAKGS TGPKGERGAP GVQGAPGNAG 180
AAGPAGPAGP QGAPGSRGPP GLKGDRGVPG DRGIKGESGL PDSAALRQOM EALKGKLQRL 240
EVAFSHYQKA ALFPDGHRRLL DKIEDERNLH EDFVFMKTIQ RCNTERGSLN LLNCEEIKSQ 300
FEGFVKDIDL NKEETKKENS FEMQKGDQNP QIAAHVISEA SSKTSTVLQW AEKGYTMSN 360
NLVTLENGKQ LTVKRQGLYY IYQVTFCSN REASSQAPFI ASLCLKSPGR FERILLRAAN 420
THSSAKPCQQ QSIHLGGVFE LQPGASVFFN VTDPSQVSHG TGFTSFGLLK L 471

```

```

SEQ ID NO: 14      moltype = DNA  length = 1412
FEATURE           Location/Qualifiers
source            1..1412
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 14
atgtgcctct tctgagcat gctggtgctg ctggtgcagc ctctgggcaa tctggggcgc 60
gagatgaagt cctctgtctc gagggagcgt ccaaacaccc gcacactggt catgtgctct 120
ccaaaccgaga atggactgcc aggaagggac ggaagagatg gaagggaggg accaagggga 180
gagaagggcg acctggact gcctggacca atgggactgt ccggactgca gggaccaaca 240
ggcctctgtg gaccaaaggg agagaatgga agcgcgggag agccaggacc taagggagag 300
aggggcctgt ccggccccc cggcctgccc ggcaccccag gcccgcggcg caaggagggc 360
cctctggcca agcagggcaa catcggacca caggggcaagc ctggacaaaa gggagaggca 420
ggaccaaagg gagaagtggg agcaccggcg atgcagggca gcaccggagc aaagggatcc 480
accggcccta agggagagag agggagaccc ggagtgacag gcgccccagg caatgcagga 540
gcagcaggac cagcaggacc tgcaggccca cagggcgccc caggcagccg gggccccacc 600
ggcctgaagg gcgacagggg agtgccaggc gataggggca tcaagggaga gtccggactg 660
ccagactctg ccgcctctgag gcagcagatg gaggccctga agggcaagct gcagaggctg 720
gaggtggcct tctccacta ccagaaggcc gccctgttcc cagacggaca caggagactg 780
gataaggttg agggaggagt gaacctgcac gaggatctcg tgttcatcaa gaagtgaag 840
agggtgcaaca agggcgaggg cagcctgtcc ctgctgaatt gtgaggagat gcggcgccag 900
ttcgaggacc tgggaaagga tatcacctcg aacaaggagg agaagaagga gaattctttt 960
gagatgcaga ggggcgacga ggatcctcag atcgagcac acgtggtgtc cgaggcaaac 1020
tctaagtcgg ccagcgtgct gcaagtgggc aagaagggct actataccat gaagtctaac 1080
ctggctcatg tggagaatgg caagcagctg acagtgaaga gagaggccct gtactacgtg 1140
tacacccagg tgacattctc cagcaacaga gagcccagct cccagcgccc ttttatcgtg 1200
ggcctgtggc tgaagccctc tatcggaagc gagaggatcc tgcgaaaggc agccaatacc 1260
cactctagct cccagctgtg cgagcagcag tccgtgcacc tgggaggcgt gttcgagctg 1320
caggcaggag caagcgtgtt cgtgaacgga cagaggccag ccagggtcatc cacagagtgg 1380
gcttctctag ctttggcctg ctgaagctgt ga 1412

```

```

SEQ ID NO: 15      moltype = AA  length = 470
FEATURE           Location/Qualifiers
source            1..470
                  mol_type = protein
                  organism = Mus musculus

```

```

SEQUENCE: 15
MLPFLSMLVL LVQPLGNLGA EMKSLSQRSV PNTCTLVMCS PTENGLPGRD GRDGREGPRG 60
EKGDPLPLPG MGLSGLQGPT GPVGPKEGNG SAGEPGPKGE RGLSGPPGLP GIPGPAGKEG 120
PSGKQGNIGP QGKPGPKGEA GPKGEVGPAG MQGSTGAKGS TGPKGERGAP GVQGAPGNAG 180
AAGPAGPAGP QGAPGSRGPP GLKGDRGVPG DRGIKGESGL PDSAALRQOM EALKGKLQRL 240
EVAFSHYQKA ALFPDGHRRLL DKVEEVNLH EDFVFIKKL RCNKGESLS LLNCEEMRRQ 300
FEDLVKDIIL NKEEKKENSF EMQRGDEDPQ IAAHVSEAN SNAASVLQWA KKGYYTMSN 360
LVMLENGKQL TVKREGLYYV YTVTFCSNR EPSSQRPFIV GLWLKPSIGS ERILLKAANT 420
HSSSQLCEQQ SVHLGGVFEL QAGASVFFNV TEASQVIHRV GFSSFGLLKL 470

```

```

SEQ ID NO: 16      moltype = DNA  length = 786
FEATURE           Location/Qualifiers
source            1..786
                  mol_type = other DNA
                  organism = Homo sapiens

```

-continued

SEQUENCE: 16

atgatcgaaa	catacaacca	aacttctccc	cgatctgcgg	ccactggact	gcccacagc	60
atgaaaattt	ttaatgtatt	acttactgtt	tttcttatca	cccagatgat	tgggtcagca	120
ctttttgctg	tgtatcttca	tagaagggtg	gacaagatag	aagatgaaag	gaatcttcat	180
gaagattttg	tattcoatga	aacgatacac	agatgcaaca	caggagaaag	atccttatcc	240
ttactgaact	gtgaggagat	taaaagccag	tttgaaggct	ttgtgaagga	tataatgtta	300
aacaaagagg	agacgaagaa	agaaaacagc	tttgaatgac	aaaaagggtg	tcagaatcct	360
caaattgcgg	cacatgtcat	aagtggagcc	agcagtaaaa	caacatctgt	gttacagtgg	420
gctgaaaaag	gatactacac	catgagcaac	aacttggtaa	ccctggaaaa	tgggaaacag	480
ctgaccgtta	aaagacaagg	actctattat	atctatgccc	aagtacacct	ctgttccaat	540
cgggaagctt	cgagtcaagc	tccatttata	gccagcctct	gcctaaagtc	ccccggtaga	600
ttcgagagaa	tcttactcag	agctgcaaat	accacagtt	cgcacaaacc	ttgcgggcaa	660
caatccattc	acttgggagg	agtatttgaa	ttgcaaccag	gtgcttcggt	gtttgtcaat	720
gtgactgatc	caagccaagt	gagccatggc	actggcttca	cgtcctttgg	cttactcaaa	780
ctctga						786

SEQ ID NO: 17 moltype = AA length = 261
 FEATURE Location/Qualifiers
 source 1..261
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 17

MIETYNQTS	RSAATGLPIS	MKIFMYLLTV	FLITQMIGSA	LFAVYLHRR	DKIEDERNLH	60
EDFVPMKTI	RCNTGERSLS	LLNCEEIKSQ	FEFVKDIDL	NKEETKKENS	FEMQKGDQNP	120
QIAAHVISEA	SSKTTSVLQW	AEKGYTMSN	NLVTLENGKQ	LTVKRQGLYY	IYAQVTFCSN	180
REASSQAPFI	ASLCLKSPGR	FERILLRAAN	THSSAKPCGQ	QSIHLGGVFE	LQPGASVFN	240
VTDPQVSHG	TGFTSFGLLK	L				261

SEQ ID NO: 18 moltype = DNA length = 783
 FEATURE Location/Qualifiers
 source 1..783
 mol_type = other DNA
 organism = Mus musculus

SEQUENCE: 18

atgatagaaa	catacagcca	accttcccc	agatccgtgg	caactggact	tccagcgagc	60
atgaagattt	ttaatgtatt	acttactgtt	ttccttatca	cccaaatgat	tggatctgtg	120
ctttttgctg	tgtatcttca	tagaagattg	gataaggctg	aagaggaaag	aaaccttcat	180
gaagattttg	tattcataaa	aaagctaaag	agatgcaaca	aaggagaagg	atctttatcc	240
ttgctgaact	gtgaggagat	gagaaggcaa	tttgaagacc	ttgtcaagga	tataacgtta	300
aacaaagaag	agaaaaaaga	aaacagcttt	gaaatgcaaa	gaggtgatga	ggatcctcaa	360
attgcagcac	acgttgtaag	cgaagccaac	agtaatgcag	catccgttct	acagtgggcc	420
aagaaaggat	attataccat	gaaaagcaac	ttggtaaatg	ttgaaaatgg	gaaacagctg	480
acgggttaaa	gagaaggact	ctattatgtc	tacactcaag	tcaccttctg	ctctaactcg	540
gagccttcga	gtcaacgccc	attcatcgtc	ggcctctggc	tgaagcccag	cagtggatct	600
gagagaatct	tactcaaggc	ggcaaatacc	cacagttcct	cccagctttg	cgagcagcag	660
tctgttcact	tggcgaggag	gtttgaatta	caagctgggtg	cttctgtgtt	tgtaaacgtg	720
actgaagcaa	tccaagtgat	ccacagagtt	ggcctctcat	cttttggtct	actcaaaactc	780
tga						783

SEQ ID NO: 19 moltype = AA length = 260
 FEATURE Location/Qualifiers
 source 1..260
 mol_type = protein
 organism = Mus musculus

SEQUENCE: 19

MIETYSQSP	RSVATGLPAS	MKIFMYLLTV	FLITQMIGSV	LFAVYLHRR	DKVEEVNLH	60
EDFVFIKKL	RCNKGESLS	LLNCEEMRRQ	FEDLVKDITL	NKEEKKENS	EMQRGDEDPQ	120
IAAHVVSEAN	SNAASVLQWA	KKGYTMSN	LVMLENGKQL	TVKREGLYYV	YTQVTFCSNR	180
EPSSQRPFIV	GLWLKPSSGS	ERILLKAANT	HSSSQLCEQQ	SVHLGGVFEL	QAGASVFN	240
TEASQVIHRV	GFSSFGLLKL					260

SEQ ID NO: 20 moltype = DNA length = 930
 FEATURE Location/Qualifiers
 source 1..930
 mol_type = other DNA
 organism = Mus musculus

SEQUENCE: 20

atggatcagc	acacactgga	cgtggaggat	accgctgacg	ctaggcaccc	agctggcacc	60
tctgcccctt	ctgatgccgc	tctgctgcgc	gacacaggac	tgctggccga	tgccgctctg	120
ctgtctgaca	cagtgcggcc	aaccaacgcc	gctctgccaa	ccgatgtctg	ttacctgtct	180
gtgaacgtga	gggtacagaga	ggctgcttgg	ccacctgccc	tgaacttctg	cagccgccac	240
cctaagctgt	acggcctggt	ggccttggtg	ctgctgtctg	tgategctgc	ttgcgtgcca	300
atctttacc	ggacagagcc	acgccccgct	ctgacaatca	ccacatcccc	caacctgggc	360
accagggaga	acaacgccga	tcaggtgaca	ccagtgtctc	acatcggtgc	ccccaacacc	420
acacagcagg	gaagcccagt	gttcgccaa	ctgctggcta	agaaccaggc	cagcctgtgc	480
aacaccacac	tgaactggca	cagccaggac	ggagctggaa	gctcctacct	gtcccagggc	540
ctgagatacg	aggaggataa	gaaggagctg	gtggtggact	ccctggact	gtactacgtg	600
ttcctggagc	tgaagctgtc	tccaaccttt	acaaacaccg	gccacaaggt	gcagggatgg	660
gtgtctctgg	tgctgcaggc	taagccccag	gtggacgatt	tcgataacct	ggcctgacc	720

-continued

```

gtggagctgt ttcctttag catggagaa aagctggtg acaggtcttg gagccagctg 780
ctgctgtctga aggttggcca caggctgtcc gtgggactga gagcctacct gcacggcgcc 840
caggatgctt acagagactg ggagctgagc taccctaaca ccacatcctt cggactgttt 900
ctggtgaagc ctgacaaccc atgggagtga 930

```

```

SEQ ID NO: 21      moltype = DNA length = 765
FEATURE           Location/Qualifiers
source            1..765
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 21
atggagtacg cctctgacgc cagcctggat ccagaggccc cttggccacc tgcaccaagg 60
gcccgcgcct gccgcgtgct gccctgggcc ctggtggccc gccctgtatt actgtgtctg 120
ctggcgcgccg cctgcgcgtt gttcctggca tgtccttggg ccgtgagcgg agccagagcc 180
tccccaggct ctgcgcgcag cctcggctg agagagggac cagagctgtc cccagacgat 240
ccagcaggcc tgctggacct gaggcaggga atgtttgccc agctggtggc ccagaacgtg 300
ctgctgatat acggccccct gtccctgtac tctgacctg gccctggcgg cgtgtctctg 360
accggcgccc tgagctataa ggaggataca aaggagctgg tgggtggcaa ggccggcgtg 420
tactacgtgt tcttccagct ggagctgagg agagtgtgtg caggagaggg ctctggaagc 480
gtgtcccttg gcagccccct gcagccccg caggagccgc cgcctgggcc 540
ctgacccttg acctgccacc agccagctcc gaggcaagga attccgcctt cggctttcag 600
ggcagactgc tgcactgtc tgcggacag aggtgtggg tgcacctgca caccgaggcc 660
agggccccgc acgcattgca gctgacccag ggagcaacag tgcgtggcct gttccgcgtg 720
acacctgaga tcccagcagg cctgcctagc ccacggtccg agtga 765

```

```

SEQ ID NO: 22      moltype = DNA length = 1389
FEATURE           Location/Qualifiers
source            1..1389
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 22
atgtgtcctt tctgttccat gctggtgctg ctggtgcagc cactgggcaa cctgggagcc 60
gagatgaagt ctctgagcca gcgcagcgtg cctaaccact gcacactggt catgtgtctc 120
cctacagaga acggcctgcc aggaagggac ggaagagatg gaaggagggg accaagggga 180
gagaaggcgc acccggact ccttggaaca atgggactga gcggcctgca gggaccaacc 240
ggccccgtgg gacctaaggg agagaacgga tccgctggag agccaggacc taagggagag 300
agaggactgt ctggaccacc tggactgcca ggaatcccag gaccagctgg caaggaggga 360
ccatccggca agcagggaaa catcggacca cagggaaaagc ctggaccaaa gggagaggct 420
ggacctaaag gagaagtggg cgcgccagga atgcagggct ctacaggagc taagggcagc 480
accggaccaa agggagagag gggagccccg ggaagtgcagg gagccccctg caacgctgga 540
gccgtggccc cagcggagac cgttggccct cagggagccc ccggtcttag gggaccacca 600
ggcctgaagg gagacagagg cgtgccccga gatcggggca tcaagggaga gagcggcctg 660
cctgaactcg ccgtcttgag acagcagatg gaggtctcta agggcaagct gcagcggctg 720
gaggtggcct tctcccacta ccagaaggcc gctctgttct ctgacggaa gacagagccc 780
aggctctgtc tgacctcac cacatctcca aacctgggca caagagagaa caacgccgat 840
caggtgaccc ccgtgtctca catcgatgc cctaaccacca cacagcaggg cagccccctg 900
tttgccaagc tctgggctaa gaaccaggcc agcctgtgca acaccacact gaactggcac 960
tcccaggatg gccgcggaag ctccctacct tctcagggcc tgcggtagca ggaggacaag 1020
aaggagctgg tgggtgatag ccaggcctg tactacgtgt tccaggagct gaagctgtcc 1080
ccacacctta caaacaccgg acacaagggt cagggatggg tgagcctggt gctgcaggct 1140
aagccccagg tggacgattt cgacaacctg gccctgaccg tggagctgtt tcttctgtct 1200
atggagaaca atgtgttgga tagatcctgg agccagctgc tgcgtctgaa ggctggacac 1260
cgctgagcgc tgggcttgag ggcttacctg cacggagctc agggacgtta cagggatagg 1320
gagctgtcct accctaacac cacatcttct ggctgttctc tgggtgaagcc agacaacccc 1380
tgggagtga 1389

```

```

SEQ ID NO: 23      moltype = DNA length = 1389
FEATURE           Location/Qualifiers
source            1..1389
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 23
atgtgtctgt tccctgtgtc cgccttggtg ctgctgaccc agcctctggg ctacctggag 60
gccagatga agactatctc tcaccggaca atgccaaagc cctgcacact ggtcatgtgc 120
agcagcgtgg agtctggcct gccagggaag gacggaaggg atggaaggga gggacctaga 180
ggcagagaag gcgaccttgg cctgccagga cgacaggac aggcaggaaat gcccgcccag 240
gccggccccg tgggacctaa gggcgacaa cgaagcgtgg gagagccagg accaaagggc 300
gataccggcc cttcgggacc acctggacca ccaggcgtgc ctggcccagc cggcagggag 360
ggcctctctg gcaagcaggg caatatcgcc ccacagggca agcccggccc taagggcgag 420
gccggcccca agggcggaag gggcgccccct ggcctgcagg gaagcgccgg agcccgcggc 480
ctggccggac ctaagggcga gagaggcgtg cctggagaga ggggcgtgcc aggaaacaca 540
ggcgcagcag gatctgccgg agcaatggga ccccagggca gccctggcgc caggggccct 600
ccaggcctga agggcgacaa gggcatccca ggcgataagg gagcaaaagg agagagcggc 660
ctgccagatg tggcctcctc gcgccagcag gtggaggccc tgcagggcca ggtgcagcac 720
ctgcaggccc gttctctca gtacaagaag gtggagctgt ttcaaaacgg cgcctgcccc 780
tgggcccgtg gccggagccc gccctcccca ggctctgccc ccagccctag gctgcgcgag 840
ggaccagagc tgagcccaga cgatccagca ggctctgctg acctgagaca gggaatgttc 900
gccagctggg tggcccagaa tgtgtgtgtg atcgacggcc cactgtctct gtactctgat 960
ccaggcctgg ccggcgtgtc cctgaccggc ggctgtctct ataaggagga tacaaggag 1020

```

-continued

```

ctggtggtgg ccaaggccgg cgtgtactac gtgttcttcc agctggagct gaggagagtg 1080
gtggcaggag agggatccgg atctgtgagc ctggccctgc acctgcagcc cctgcggtcc 1140
gcccagggag ccgcccctct ggccctgacc gtggacctgc cacctgcctc tagcgaggca 1200
cgcaattccg ccttcggctt tcaggggccg ctgctgcacc tgtctgccgg acagagactg 1260
ggagtgcacc tgcacaccga gggccggggc agacacgcct ggcagctgac ccaggggagca 1320
acagtgtctg gcctgttttag ggtgacacct gagatcccag ccggcctgcc aagccccccg 1380
tccgagtga                                     1389

```

```

SEQ ID NO: 24      moltype = DNA length = 522
FEATURE           Location/Qualifiers
source            1..522
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 24
atggaggaga tgccctctgag ggagagctcc ccacagaggg ccgagagatg caagaagagc 60
tggtgtgtgt gcatcgtggc tctgtgtgtg atgtgtgtgt gctctctggg caccctgac 120
tacacaagcc tgaagccaac cgcctcgag tctgttatgg tgaagttcga gctgtctagc 180
tccaagtggc acatgacatc ccccaagcct cactgcgtga acaccacatc tgacggaaag 240
ctgaagatcc tgcagagcgg cactacctg atctacggac aggtcatccc cgtggacaag 300
aagtacatca aggataacgc ccctttcgtg gtgcagatct acaagaagaa cgacgtgtctg 360
cagacactga tgaacgattt tcagatcctg cccatcgccg gagtgtacga gctgcacgct 420
ggcgacaaca tctacctgaa gttcaactcc aaggatcaca tccagaagac caacacatac 480
tggggaatca tctctgatgc agatctgccc tttatctctt ga 522

```

```

SEQ ID NO: 25      moltype = DNA length = 600
FEATURE           Location/Qualifiers
source            1..600
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 25
atgaccctgc acccaagccc catcacatgc gagttcctgt tttctaccgc cctgatcagc 60
ccaaagatgt gccctgagcca cctggagaat atgcccctgt cccactctcg gacacaggga 120
ccccagagaa gctcctggaa gctgtggctg tctgtctcta tcgtgatgct gctgttctcg 180
tgcagctttt cctggctgat ctctcatctt ctgcagctgg agacagccaa ggagccttgc 240
atggccaagt ttggccctct gccatccaag tggcagatgg cctctagcga gcccccttgc 300
gtgaacaagg tgagcgactg gaagctggag atcctgcaga acggcctgta cctgatctat 360
ggccaggttg cccccaacgc caattacaac gacgtggccc ctttcgaggt gcggtgtgat 420
aagaacaagg atatgatcca gaccctgaca aataagtcta agatccagaa cgtgggcccgc 480
acatacagagc tgcacgtggg cgacaccatc gacctgatct tcaacagcga gcaccagggtg 540
ctgaagaaca atacatatgt gggcatcatc ctgctggcca acccccagtt tatctcctga 600

```

```

SEQ ID NO: 26      moltype = DNA length = 1164
FEATURE           Location/Qualifiers
source            1..1164
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 26
atgtgccttt tctctgtctat gctggtgctg ctgtgacgac cactgggcaa cctggggagcc 60
gagatgaaga gctgtgtccc gagatccgtg cccaacacct gcacactggt catgtgtctc 120
cctaccgaga acggcctgac aggaagggag ggaagagatg gaagggaggg acctcgggga 180
gagaagggcg acccaggact gcttgaccca atgggactga gcggcctgca gggaccaaca 240
ggccccgttg gacctaaagg agagaacgga agcgccggag agccaggacc taaggggagag 300
aggggactgt ccgggaccacc tggactgcct ggaatcccag gaccagctgg caaggaggga 360
ccatccggca agcagggaaa ctacggacca cagggaaaag ctggaccaaa gggagaggct 420
ggaccaaaag gagaagtggg cgctcctgga atgcagggct ccaccggagc caagggtctc 480
acaggaccaa aaggagagag gggagctccc ggagtgcagg gagcccctgg caacgctgga 540
gccgctggcc cagccggacc cgctggccct cagggagccc caggcagcag gggaccaccc 600
ggcctgaagg gcgacagggg cgtgccagga gataggggca tcaagggaga gtctggcctg 660
ccagacagcg ccgctctgag acagcagatg gagggccctga agggcaagct gcagcggctg 720
gaggtggctt tctccacta ccagaaggcc gctctgtttc cagatggcag cctgaagccc 780
accgccatcg agtccctgat ggtgaagttt gagctgagct cctctaagtg gcacatgaca 840
tctcccaagc ctcactgcgt gaacaccaca tctgacggca agctgaagat cctgcagagc 900
ggcactacc tgatctacgg ccaggtcatc cccgtggaca agaagtacat caaggataac 960
gcccctttcg tggcgccgtg ctacaagaag aacgacgtgc tgcagacact gatgaacgat 1020
tttcagatcc tgccaatcgg cggagtgtac gagctgcacg ctggcgacaa catctacctg 1080
aagttcaact ctaaggatca catccagaag accaacacat actggggcat catcctgatg 1140
ccagatctgc cctttatcag ctga 1164

```

```

SEQ ID NO: 27      moltype = DNA length = 1152
FEATURE           Location/Qualifiers
source            1..1152
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 27
atgtgtgtgt tctgtgtgtc tgccttggtg ctgtgacccc agccactggg ctacctggag 60
gccagatgta agacctattc ccaccgcaca atgccttctg cctgcacact ggtcatgtgc 120
agcagcgttg agagcgccct gccaggaagg gacggaagag atggaaggga gggaccacga 180
ggcgagaagg gcgacctgtg cctgccagga gcagcaggac aggcaggaaat gccaggccag 240
gccggccccc tgggccctaa gggcgacaat ggtaccgtgg gagagccagg accaaagggc 300

```

-continued

```

gataccggccc cttctggacc acctggacca ccaggcgtgc ctggaccagc aggaagagag 360
ggacctctgg gcaagcaggg aaacatcgga ccacagggca agccaggccc taaggcgag 420
gcccggcccca agggcgcaagt gggcgccccc ggcatgcagg gatccgcccg agccaggggc 480
ctggccgggac ctaaggggcga gcgcggcgctg cctggagaga ggggcggtgc aggaataca 540
ggcgagcgag gatctgcccg agcaatggga ccacagggca gcccggcgcc cagaggccct 600
ccaggcctga agggcgacaa ggggaatccct ggcgataagg gagcaagggg agagagcggc 660
ctgccagacg tggcctccct gaggcagcag gtggaggccc tgcaggggaca ggtgcagcac 720
ctgcaggccg ccttcagcca gtacaagaag gtggagctgt ttccaaatgg cgagacagcc 780
aaggagccct gcatggccaa gttcggccca ctgccagca agtggcagat ggcccttagc 840
gagccccctt gcgtgaacaa ggtgagcgat tggaaagtgg agatcctgca gaacggcctg 900
tacctgatct atggccaggt gggccccaac gccaaattaca acgacgtggc cccttttgag 960
gtgcggctgt ataagaacaa ggatatgatc cagaccctga caaataagtc taagatccag 1020
aacgtgggag gcacctacga gctgcacgtg ggcgacacaa tcgacctgat cttcaacagc 1080
gagcaccagg tgctgaagaa caatacatat tggggcatca tctgtctggc caacccccag 1140
tttatctcct ga                                     1152

```

```

SEQ ID NO: 28      moltype = DNA length = 597
FEATURE           Location/Qualifiers
source            1..597
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 28
atggaggggc agggagtgc gcccctggat gagaacctgg agaacggctc cgggcctcgc 60
ttcaagtggg agaagaccct gcggtctgtg gtgtctggaa tcaagggcgc cggaatgctg 120
ctgtgcttta tctacgtgtg cctgcagctg agctcctctc ccgccaagga tccccctatc 180
cagaggctga gaggagctgt gaccagggtg gaggacggac agctgttcac cagctcctac 240
aagaacagat accagacaat ggagggtgcag aacaacagcg tggtcaccaa gtgtgatggc 300
ctgtacatca tctacctgaa gggatccttc ttccaggagg tgaagatcga cctgcacttt 360
cgggaggatc acaaccaaat ctctatcccc atgctgaacg acggcaggag aatcgtgttc 420
acagtgtgtg ccagcctggc ttttaaggac aaggtgtacc tgaccgtgaa cggcccagat 480
acactgtgcg agcactgca gatcaacgac ggagagctga tctgtgtgca gctgaccctc 540
ggctactgtg ctccagaggg atcttaccac agcacagtga accaggtgac cctgtga 597

```

```

SEQ ID NO: 29      moltype = DNA length = 552
FEATURE           Location/Qualifiers
source            1..552
                  mol_type = other DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 29
atggagaggg tgcagccctt ggaggagaaac gtgggaaatg ccgcccggcc tagattcgag 60
aggaacaagc tgcgtctggt ggcctctgtg atccagggcc tgggcctgct gctgtgcttc 120
acctacatct gctcgcactt ttctgcccct caggtgagcc acagataccc ccgcattccag 180
agcatcaagg tgcagttcac caggtataag aaggagaagg gctttatcct gacatccag 240
aaggaggacg agatcatgaa ggtgcagaac aattctgtga tcatcaactg cgatggcttc 300
tacctgatct cctcgaaggg ctatttttct caggaaagta atatcagcct gcaactatcag 360
aaggacgagg agcactgttt tcagctgaag aaggtgcgga gcgtgaattc cctgatgggtg 420
gccagcctga cctacaagga caaggtgtat ctgaacgtga ccacagataa tacatccctg 480
gacgatttcc acgtgaacgg cggcgagctg atcctgatcc accagaatcc cggcgagttt 540
tgctgtctgt ga                                     552

```

```

SEQ ID NO: 30      moltype = DNA length = 1215
FEATURE           Location/Qualifiers
source            1..1215
                  mol_type = other DNA
                  organism = Mus musculus

```

```

SEQUENCE: 30
atgtgcctct tctgtccat gctggtgctg ctgtgacgac ctctgggcaa cctgggagcc 60
gagatgaagt ctctgagcca gagatccgtg ccaaacacct gcacactggt catgtgctct 120
cccaccgaga acggcctgct tggaaaggac ggaagagatg gaaggagggg accccgggga 180
gagaaggggc atcctggact gccaggacct atgggactga gcggcctgca gggaccaaca 240
ggccccgtgg gacctaaggg agagaacgga agcgccggag agccaggacc aaagggagag 300
aggggactgt ccggccacc tggactgcct ggaatccctg gaccagctgg caaggaggga 360
ccttcgggca agcagggaaa catcggacca cagggaaggc caggacctaa gggagaggct 420
ggaccaaaag gagaagtggg cgctcccgga atgcagggct ctaccggagc caaggcgacg 480
acaggacctt agggagagag gggagctcca ggagtgcagg gagcccccgg caacgctgga 540
gctgctggac cagctggacc agctggccct caggagagcc caggctctag gggaccacca 600
ggcctgaagg gcgacagggg cgtgccagga gataggggca tcaagggaga gagcgccctg 660
ccagattccg ccgctctgag acagcagatg gagggcctga agggcaagct gcagcggtg 720
gaggtggcct tcagccacta ccagaaggcc gctctgttct ctgacggcag ctctctcca 780
gccaaaggat cctcaatcca gcggctgctg ggagctgtga ccaggtgcga ggatggccag 840
ctgttcatac gctcctacaa gaacgagtac cagacaatgg aggtgcagaa caactctgtg 900
gtcatcaagt gtgacggcct gtacatcatc tacctgaagg gcagcttctt tcaggagggtg 960
aagatcgacc tgcactttag agaggatcac aacccaatct ccattcccat gctgaacgac 1020
ggcaggagaa tctgtttcac cctgtgtgct tctctggcct ttaaggacaa ggtgtacctg 1080
accgtgaacg ccccgatcac actgtgcgag cacctgcaga tcaacgacgg cgagctgatc 1140
gtggtgcagc tgacccctgg atactgtgct ccagagggct cctaccactc tacagtgaac 1200
caggtgcctc tgtga                                     1215

```

```

SEQ ID NO: 31      moltype = DNA length = 1170

```

-continued

FEATURE
source

Location/Qualifiers
1..1170
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 31

```

atgctgctgt tccctgtgag cgccttgggt ctgctgaccc agccactggg ctacctggag 60
gccgagatga agacctattc ccacagaaca atgccttctg cctgcacact ggctcatgtg 120
agcagcggtg agtcogggct gccaggaagg gacggcagag atggcaggga gggccccagg 180
ggcgagaagg gcgaccccggt cctgcttggg gcagcaggcc agggccggcat gccaggccag 240
gccggcccgag tggggcccaa gggcgacaa ggcagcgtgg gcgagcccg gctaaggggc 300
gataccggcc cctcogggcc ccttgccca cccggcgtgc caggaccagc aggaaggggag 360
ggaccactgg gcaagcaggg caatatcgga cctcagggca agcctggacc aaaggggagag 420
gcaggaccaa agggagaagt gggcgccctt ggcatgcagg gatctgccc agcccggggc 480
ctggccggcc ccaagggcga gagaggcgtg cccggcgaga gggcggtgcc tggcaacaca 540
ggcgccggcc gctcggcggt cgcctgggga cctcagggct ctccaggagc caggaggccct 600
ccaggcctga agggcgacaa ggggaatccct ggcgataagg gagcaagggg agagagcggc 660
ctggcagacg tggcctccct gggcgagcag gtggaggccc tgcaggggca ggtgcagcac 720
ctgcaggccg ccttcagcca gtacaagaag gtggagctgt ttccaatagg cgtgtctcac 780
cgtaaccacg ggaaccagag catcaagggt cagttcaccg agtataagaa ggagaagggc 840
tctatcctga catctcgaaa ggaggacgag atcatgaagg tgcagaacaa tagcgtgatc 900
atcaactcgg atggtctcta cctgatcagc ctgaagggtt attttccca ggaagtgaat 960
atctctctgc actatcagaa ggaatgaggag cctctgtttc agctgaagaa ggtgagatct 1020
gtgaacagcc tgatgggtgc ctcctgacc tacaaggaca aggtgtatct gaacctgacc 1080
acagataata catctctgga cgatttcac gtgaacggcg gcgagctgat cctgatccac 1140
cagaatcccg gcgagttttg cgtgctgtga 1170

```

SEQ ID NO: 32

FEATURE
source

moltype = DNA length = 969
Location/Qualifiers
1..969
mol_type = other DNA
organism = Mus musculus

SEQUENCE: 32

```

atgcagctga agtgctccatg cttcgtgtcc ctgggaacaa gacagccgt ctggaagaaa 60
ctgcacgtga gctcgggctt ctttagcggc ctggggctgt ttctgctgct gctgtctagt 120
ctgtgcggcg gctcggcaga gactgaagtc ggagccatgg tgggcagtaa cgtggtcctg 180
tcatgcatcg accacacccg acggcatttc aacctgtctg gcctgtacgt gtattggcag 240
attgagaatc ccgaagtgtc agtcacctac tatctgcctt acaagagccc agggatcaac 300
gtggactcaa gctataaaaa tagggggcac ctgtccctgg attctatgaa gcagggaaac 360
ttcagcctgt acctgaaaaa tgtgacccct caggacacac aggaagttcac ttgtcgcgtc 420
tttatgaaca ctgcaaccga actggtgaag attctggagg aagtgggtcc gctgagagtc 480
gcagccaact ttagcactcc tgtgatctct accagtgaat cctctaatcc agggccaggag 540
cggacatata cttgcatgtc taagaacgga tccccgaac ctaatctgta ttggatcaac 600
accacagaca atagtctgat tgataccgct ctgcagaaca atacagtcta cctgaacaag 660
ctggggctgt atgacgtgat ctctactctg cggctgccat ggaccagtag aggagatgtg 720
ctgtgtctcg tggagaacgt ggcctgcac cagaatatca cctcaattag ccaggctgag 780
tcctttaccg gcaacaatcc caggagacac ataacaatga actgaaagtg 840
ctggtgccag tgggtggcgt cctgggtgca gcagctttcg tgtcttttat catctacaga 900
aggacccgcc ctcaccgctc atacactgga cctaagaccg tgcagctgga actgacagac 960
catgtctga

```

SEQ ID NO: 33

FEATURE
source

moltype = DNA length = 909
Location/Qualifiers
1..909
mol_type = other DNA
organism = Homo sapiens

SEQUENCE: 33

```

atgcgtctgg gttcacctgg tctgctgttt ctgctgtttt caagtctgcg tgctgatact 60
caggagaagg aagtcggggt tatggtcgga agtgacgtgg agctgtcatg cgcttgtccc 120
gaagggtccc ggttcgacct gaacgatgtc tacgtgtatt ggcagacctc tgagagtaag 180
accgtggtca cataccacat cctcagaaac tccagcctgg aaaatgtgga ttcaagggtat 240
cggaaacagag cctgcatgtc cctgctggc atgctgcggg gagacttctc tctgagactg 300
tttaatgtga caccacagga tgagcagaaa ttccattgcc tggctcctgtc acagtcctctg 360
ggatttcagg aggtgctgag tgtcgaagtg actctgcacg tcgcccgtaa tttctccgtg 420
cctgtgtgta gccgaccaca tagccctctc caggacgagc tgacctttac atgtacttcc 480
atcaacgggt ccccccgccc taacgtgtac tggattaaca agactgacaa tagcctgctg 540
gatcaggcac tgcagaacga caccgtgttt ctgaatatgc gaggactgta cgatgtggtc 600
agcgtcctgc gtattgccag gaccccatct gtgaacatcg ggtgctgtat tgagaacgtc 660
ctgctgcagc agaactctgac agtggggagc cagactggta atgacatcg cgagagggat 720
aagattaccg aaaaccccggt gagtacaggc gagaagaacg cagccacatg gtcaatcctg 780
gctgtgctgt gcctgctggg ggtcgtggct gtcgcaattg gctgggtgtg ccgcatcggt 840
tgtctgcagc actcttatgc cggctgcttg gcagtgaagt cagagactga actgacggcg 900
catgtctaa

```

SEQ ID NO: 34

FEATURE
source

moltype = DNA length = 1574
Location/Qualifiers
1..1574
mol_type = other DNA
organism = Mus musculus

SEQUENCE: 34

-continued

```

cttaagatgg aaactgatac tctgtctgtc tgggtgtctc tctctgtgggt gctgtgttca 60
actgggggaca ttctgacgggc tgacattgtg atgacccaga ccacactgag cctgcccgtg 120
tccctgggagc accaggccag catctcctgc cggagctccc agtctatcgt gcacagcaac 180
ggaaacacat acctggagtg gtatctgcag aagcctggcc agtcccaaa gctgtgtatc 240
tacaagggtt ccaacagggt cagcggcggt cctgaccgct tttctggaag cggctccgga 300
acagatttca cctgaagat cagcagggtg gagcgtgagg acctgggctg gtactactgc 360
ttccagggat cccacgtgcc ttacacctt ggccgaggca caaagctgga gatcaagaga 420
gcccagtgctg ctccaaccgt gctctggaagc ggaggcgggg gttctggagg cgggtgggagc 480
gggtggcggag ggtctgaggc taagctgcag gagagcggcc ccgtgctggt gaagcctgga 540
gccagcgtga agatgtcctg taaggcttct ggatacacct tcacagacta ctacatgaac 600
tgggtgaagc agagccacgg caagtccctg gagtggatcg gactgatcaa ccttacaac 660
ggcgacacct cttacaacca gaagttaaag ggcaaggcca cctgacagct ggataagtct 720
agctccaccg cttacatgga gctgaacagc ctgacatccg aggtattctgc cgtgtactac 780
tgtgtcaggt actaagaaag ctgggtcgcc agggaaacat gatcacccgtg 840
tccacagcca agaccacacc ccttagcgtg taccctctgg ctcttaggtc tagcagaggc 900
tgcaagccat gcatctgtac cgtgcccagc gtgagcagcg tgttcatctt tccaccaag 960
cccaaggagc tctgacacct cacactgacc cctaagggtg catgctgggt ggtggatatt 1020
agcaaggagc atccagaggt gcagttctcc tgggttctgg acgattgtga ggtgcacacc 1080
gcccagacac agccaaggga ggagcagttc aactccacct ttagatccgt gctgagctg 1140
cccatcctgc accaggactg gctgaacgga aaggagttca agtcccgggt gaactccgcc 1200
gcttttctctg ctccaatcga gaagaccatc tctaagacaa agggccgcc aaaggctcca 1260
caggtgtaca ccatccctcc acccaaggag cagatggcta aggataaggt gagcctgacc 1320
tgtatgatca cagacttctt tcccagggat atcacagtgg agtggcagtg gaacggacag 1380
ctgcccagga actacaagaa caccagcca atcatggaca cagatggctc ttactctgtg 1440
tacagcaagc tgaacgtgca gaagtctaac tgggaggtcg gcaacacctt cacctgcagc 1500
gtgctgcagc aaggtctcca taatcaccac accgaaaaga gcctcagtc cagccctggg 1560
aatgaggcg cgcc 1574

```

```

SEQ ID NO: 35      moltype = DNA length = 1484
FEATURE            Location/Qualifiers
source             1..1484
                   mol_type = other DNA
                   organism = Homo sapiens

```

```

SEQUENCE: 35
cttaagatgg aaactgacac cctgtctgtc tgggtcctgc tctgtgtgggt gctgtgatcc 60
accggcgata tctgtctgtc ccagctctct ggccacactga gtctgtcacc aggggagcga 120
gcaacactgt cttgtagagc cagccagctc gtgggaagct cctacctggc ttggtatcag 180
cagaagccag gccaggcacc caggtctgtc atctacggag ccttcagccg ggccactggc 240
attccagaca ggttctcttg aagtggctca gggaccgact tcacctgac catcagccga 300
ctggagcccg aagactctgc cgtgtactat tgccagcagt acggtcttag tcttggact 360
tttgacaggc gcaccaaagt ggagatcaag cgcggcgggg gaggtctctg gggaggcggg 420
agtggaggcg ggggatcaca ggtccagctg gtggaagcgg cggggggagt ggtccagcca 480
ggccggagcc tgcggctgag cgtcgccgct tcaggattca cattttcaag ctataccatg 540
cactgggtcc ggcaggcacc aggggaaggga ctggagtggg tgaccttcat cagctatgac 600
ggcaacaaca agtattacgc tgattccgtg aaaggagggt ttaccattag ccgcgacaac 660
tccaaaaata cactgtacct gcagatgaac agcctgcggg ccgaggatac tgctatctac 720
tattgcgcaa gaacccgggt gctgggaccc ttgactattt ggggcccagg gactctggtc 780
accgtgtcct ctgataagac acacacatgc cctccctgtc ctgcaccaga gctgtggggc 840
gggcatcccg tgttctctgt tccaccaag cctaagacaa cctgatgat cagccggaca 900
cctgaagtca tttgcgtggt cgtggacgtg agtcacgagg atccagaagt caagttaac 960
tggtagctgg atggcgtcga ggtgcataat gccaaagcca aacctcgcca ggaacagtac 1020
aatagcacat atcgagtcgt gatccctctg actgtgtctg atcaggattg gctgaacggc 1080
aaagagtata agtgcaaaagt gagcaataag gcactgcctg cccaatcga gaaaacaatt 1140
tccaaggcta aaggccagcc cagggaaacct caggtgtaca ctctgcctcc aagtgcgcag 1200
gaaatgacca agaaccaggt gaccttgacc tgtctggtga aagggttcta tccatcagac 1260
attgagtgag agtgggaaag caatggacag cccgaaaaaa attacaagac cacacccct 1320
gtgctggaca gcgatggctc cttcttctgt tattctaagc tgactgtgga taaaagtgcg 1380
tggcagcagg ggaacgtctt tagctgttcc gtgatgcag aggtctgca caatcattac 1440
acacagaagt cctctgagttc gtcacccggc aaatgaggcg cgcc 1484

```

```

SEQ ID NO: 36      moltype = DNA length = 632
FEATURE            Location/Qualifiers
misc_feature        1..632
note = CMV promoter
source              1..632
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 36
gttgacattg attattgact agttattaat agtaatcaat tacgggggtca ttagttcata 60
gcccataatg ggagttccgc gttacataac ttacggtaaa tggcccgcct ggctgaccgc 120
ccaacgacct ccgcccattg acgtcaataa tgacgtatgt tccatagta acgccaatag 180
ggactttcca ttgacgtcaa tgggtggagt atttacggta aactgcccc ttggcagtac 240
atcaagtgtg tcatatgcca agtacgcccc ctattgacgt caatgacggt aaatggcccg 300
cctggcatta ttgccagtac atgaccttat gggacttttc tacttgccag tacatctacg 360
tattagtcat cgctattacc atgggtgatg ggttttggca gtacatcaat gggcgtggat 420
agcggtttga ctacggggga ttccaagtc tccaccccat tgacgtcaat gggagtttgt 480
tttggcacca aaatcaacgg gactttccaa aatgtcgtaa caactccgcc ccattgacgc 540
aaatggcgcg taggcgtgta cgggtggagg tctatataag cagagctctc tggttaacta 600
gagaaccac tgcttactgg cttatcgaaa tt 632

```


-continued

```

SEQ ID NO: 37          moltype = DNA length = 394
FEATURE               Location/Qualifiers
misc_feature          1..394
                      note = RSV promoter
source                1..394
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 37
tgtacgggcc agatatacgc gtatctgagg ggactagggt gtgtttaggc gaaaagcggg 60
gcttcgggtg tacgcggtta ggagtcacct caggatatag tagtttcgct ttgcatagg 120
gagggggaaa tgtagtctta tgcaatacac ttgtagtctt gcaacatggt aacgatgagt 180
tagcaacatg cttacaagg agagaaaaag caccgtgcat gccgattggt ggaagtaagg 240
tggtaacgat gtgcottatt aggaaggcaa cagacaggtc tgacatggat tggacgaacc 300
actgaattcc gcattgcaga gataattgta tttaagtgcc tagctcgata caataaacgc 360
catttgacca ttcaccacat tgggtgacac ctcc 394

SEQ ID NO: 38          moltype = DNA length = 188
FEATURE               Location/Qualifiers
misc_feature          1..188
                      note = BGH polyA
source                1..188
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 38
ctgtgccttc tagttgccag ccattctgtt ttggccctc ccccgctgct tcttgaccc 60
tggaaagggtg cactcccact gtcccttctc aataaaatga ggaaattgca tcgcattgtc 120
tgagtaggtg tcattctatt ctggggggtg ggggtgggca ggacagcaag ggggaggatt 180
gggaagac 188

SEQ ID NO: 39          moltype = DNA length = 249
FEATURE               Location/Qualifiers
misc_feature          1..249
                      note = SV40 late polyA
source                1..249
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 39
gacatgataa gatacattga tgagtttga caaaccacaa ctagaatgca gtgaaaaaaa 60
tgctttattt gtgaaatttg tgatgctatt gctttatttg tgaatttgt gatgctattg 120
ctttatttgt aaccattata agctgcaata aacaagttaa caacaacaat tgcattcatt 180
ttatgtttca ggttcagggtg gaggtgtggg aggtttttta aagcaagtaa aacctctaca 240
aatgtggtg 249

SEQ ID NO: 40          moltype = DNA length = 345
FEATURE               Location/Qualifiers
misc_feature          1..345
                      note = SV40 enhancer promoter
source                1..345
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 40
gctgtggaat gtgtgtcagt tagggtgtgg aaagtcccca ggctccccag caggcagaag 60
tatgcaaagc atgcatctca attagtcagc aaccagggtg ggaaagtccc caggctcccc 120
agcaggcaga agtatgcaaa gcatgcatct caattagtca gcaaccatag tcccgccctc 180
aactccgccc atcccgcccc taactccgcc cagttccgcc cattctccgc cccatggctg 240
actaattttt tttatttatg cagaggccga ggccgcctcg gcctctgagc tattccagaa 300
gtagtgagga ggcttttttg gaggcctagg cttttgcaaa aagct 345

SEQ ID NO: 41          moltype = DNA length = 99
FEATURE               Location/Qualifiers
misc_feature          1..99
                      note = Rabbit beta-globin polyA
source                1..99
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 41
gacctctggc taataaagga aatttatatt cattgcaata gtgtgttgga attttttgtg 60
tctctcactc ggaaggacat atgggagggc aaatcattt 99

SEQ ID NO: 42          moltype = DNA length = 723
FEATURE               Location/Qualifiers
misc_feature          1..723
                      note = GFP
source                1..723
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 42
accatggtga gcaagggcga ggagctgttc accgggggtg tgcccatcct ggctcgagctg 60

```

-continued

```

gacggcgacg taaacggcca caagttcagc gtgtccggcg agggcgaggg cgatgccacc 120
tacggcaagc tgacctgaa gttcatctgc accacggcca agctgccgt gccctggccc 180
accctcgtga ccacctgac ctacggcgtg cagtgtctca gccctaccc cgaccacatg 240
aagcagcagc acttcttcaa gtccgcatg cccgaaggct acgtccagga gcgcaccatc 300
ttcttcaagg acgaocggcaa ctacaagacc cgcgcgaggg tgaagttcga gggcgacacc 360
ctgggtgaacc gcctcgagct gaagggcctc gacttcaagg aggaocggcaa catcctgggg 420
cacaagctgg agtacaacta caacagccac aacgtctata tcatggccga caagcagaag 480
aacggcatca aggtgaactt caagatccgc cacaacatcg aggaocggcag cgtgcagctc 540
gccgaccact accagcagaa caccctcctc ggcgacggcc cctgtctgct gcccgacaac 600
cactacctga gcacccagtc gcgcttgagc aaagacccca acgagaagcg cgatcacatg 660
gtcctgctgg agttcgtgac cgcgcgaggg atcactctcg gcattggacga gctgtacaag 720
taa 723

```

```

SEQ ID NO: 43      moltype = DNA length = 454
FEATURE            Location/Qualifiers
misc_feature        1..454
                    note = MoMuLV LTR
source              1..454
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 43
ttaattaagt aacgccattt tgcaaggcat ggaaaaatac ataactgaga atagagaagt 60
tcagatcaag gtcaggaaca gatggaacag ctgaatatgg gccaaacagg atatctgttg 120
taagcagttc ctgccccggc tcagggccaa gaacagatgg aacagctgaa tatgggccaa 180
acaggatata tgtggtaagc agttcctgcc cgggctcagg gccagaaca gatgggtccc 240
agatcgggtc cagccctcag cagtttctag agaaccatca gatgtttcca gggtgcccca 300
aggacctgaa atgacctgtg gccttatttg aactaaccaa tcagttcgtc tctcgtttct 360
gttcgcgcgc ttctgtctcc cgagctcaat aaaagagccc acaaccctc actcggggcg 420
ccagtcctcc gattgactga gtcgccgct taag 454

```

```

SEQ ID NO: 44      moltype = DNA length = 1349
FEATURE            Location/Qualifiers
misc_feature        1..1349
                    note = EF1alpha promoter
source              1..1349
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 44
ttaattaaga gtaattcata caaaaggact cgccctcgcc ttggggaatc ccagggaccg 60
tcgttaaact cccactaacg tagaaccacg agatcgctgc gttccccccc cctcaccgcg 120
ccgctctcgt catcactgag gtggagaaga gcatgcgtga ggctccgggtg cccgtcagtg 180
ggcagagcgc acatcgccca cagtccccga gaagtggggg ggaggggtcg gcaattgaac 240
cggtgcttag agaagggtgc ggggggtaaa ctgggaaagt gatgtcgtgt actggctccg 300
cctttttccc gaggggtggg gagaaccgta tataagtgcg tagtcgccg tgaacgttct 360
ttttcgcaac gggtttgcg ccagaacaca ggtaagtgcg gtgtgtggtt cccgcggggc 420
tggcctcttt acgggttatg gcccttgctg gcctgaatt acttccacgc ccttggtctg 480
agtacgtgat tcttgatccc gagcttcggg ttggaagtgg gtggagagat tcgaggcctt 540
gcggttaagg agcccttcg cctcgtgctt gagttgaggg ctggcttggg cgctggggcc 600
gccgcgtgcg aatctggtgg caccttcgcg cctgtctcgc tgccttcgat aagtctctag 660
ccatttaaaa tttttgatga cctgtcgcga cgcttttttt ctggcaagat agtcttgtaa 720
atgcggggca agatctgcac actcgtattt cggtttttgg ggcgcggggc ggcgacgggg 780
cccgctgcgc ccagcgcaca tgttcggcga ggccggggcct gcgagcgcgg ccaccgagaa 840
tcggacgggg gtagtctcaa gctggccggc ctgctctggt gcctggcctc gcgccccgct 900
gtatcgcccc gccctggggc gcaaggtcgg cccggtcggc accagttcgg tgagcggaaa 960
gatggccgct tcccggccct gctgcaggga gctcaaaatg gaggacgcgg cgctcggggag 1020
agcggggcgg tgagtcaacc acacaaagga aaagggcctt tccgtctca gccgtcgctt 1080
catgtgactc cacggagtac cgggcgccgt ccaggcacct cgattagttc tcgagctttt 1140
ggagtacgct gtcttttagt tggggggagg ggttttatgc gatggagttt ccccaactg 1200
agtgggtgga gactgaagtt aggcacgctt ggcacttgat gtaattctcc ttggaatttg 1260
ccctttttga gtttggatct tggttcattc tcaagcctca gacagtggtt caaagttttt 1320
ttcttcattc tcaggtgtcg tgacttaag 1349

```

```

SEQ ID NO: 45      moltype = DNA length = 481
FEATURE            Location/Qualifiers
misc_feature        1..481
                    note = HGH polyA
source              1..481
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 45
gacgggtggc atccctgtga cccctcccca gtgcctctcc tggccctgga agttgccact 60
ccagtgccta ccagccttgt cctaataaaa ttaagttgca tcattttgtc tgactaggtg 120
tcctctata atattatggg gtggaggggg gtggtatgga gcaaggggca agttgggaag 180
acaacctgta gggcctgcgg ggtctatttg gaaccaagct ggagtgcagt ggcacaatct 240
tggctcactg caatctccgc ctctggggtt caagcgatcc tctgacctca gctcccgcg 300
ttgttgggat tccaggcatg catgaccagg ctcagctaat ttttgttttt ttggtagaga 360

```

-continued

cggggtttca ccatattggc caggettggtc tccaactcct aatctcaggt gatctaccca	420
ccttgccctc ccaaattgct gggattacag gegtgaacca ctgtccctt cctgtcctt	480
t	481

The invention claimed is:

1. A method for treating injected and non-injected solid tumors in a subject in need thereof, the method comprising:

administering to the subject an effective amount of an oncolytic herpes virus comprising inserted in its genome (i) a heterologous gene encoding granulocyte-macrophage colony-stimulating factor (GM-CSF) and (ii) a heterologous gene encoding an anti-cytotoxic T-lymphocyte-associated antigen 4 (CTLA-4) antibody,

wherein the oncolytic virus is administered by intra-tumoral injection, wherein the administration induces the regression of injected and non-injected solid tumors, and wherein injected solid tumors are solid tumors into which the oncolytic virus is injected, and non-injected solid tumors are solid tumors which are not injected with the oncolytic virus.

2. The method of claim 1, wherein the solid tumor is an adenocarcinoma, carcinoma, melanoma, or sarcoma.

3. The method of claim 1, comprising administering multiple doses of 10^4 to 10^{10} pfu of the oncolytic virus to the subject.

4. The method of claim 3, wherein the multiple doses of the oncolytic virus are administered to the subject between 3 days to 3 weeks apart.

5. The method of claim 1, wherein the method further comprises administering an anti-programmed cell death ligand 1 (PD1) antibody to the subject.

6. The method of claim 5, comprising administering the oncolytic virus to the subject prior to administering the anti-PD1 antibody to the subject.

7. The method of claim 6, comprising administering the oncolytic virus to the subject once prior to administering the anti-PD1 antibody to the subject.

8. The method of claim 6, comprising administering the oncolytic virus multiple times to the subject prior to administering the anti-PD1 antibody to the subject.

9. The method of claim 5, comprising administering the oncolytic virus and anti-PD1 antibody to the subject concurrently.

10. The method of claim 1, wherein the herpes virus is HSV-1.

11. A method for inducing a systemic anti-tumor response to regress solid tumors in a subject in need thereof, the method comprising:

administering to the subject an effective amount of an oncolytic herpes virus comprising inserted in its genome (i) a heterologous gene encoding GM-CSF and (ii) a heterologous gene encoding an anti-CTLA-4 antibody wherein the oncolytic virus is administered by intra-tumoral injection, wherein the administration induces the systemic anti-tumor response in the subject such that injected and non-injected solid tumors are regressed, and wherein injected solid tumors are solid tumors into which the oncolytic virus is injected, and non-injected solid tumors are solid tumors which are not injected with the oncolytic virus.

12. The method of claim 11, wherein the solid tumor is an adenocarcinoma, carcinoma, melanoma, or sarcoma.

13. The method of claim 11, comprising administering multiple doses of 10^4 to 10^{10} pfu of the oncolytic virus to the subject.

14. The method of claim 13, wherein the multiple doses of the oncolytic virus are administered to the subject between 3 days to 3 weeks apart.

15. The method of claim 11, wherein the method further comprises administering an anti-PD1 antibody to the subject.

16. The method of claim 15, comprising administering the oncolytic virus to the subject prior to administering the anti-PD1 antibody to the subject.

17. The method of claim 16, comprising administering the oncolytic virus to the subject once prior to administering the anti-PD1 antibody to the subject.

18. The method of claim 16, comprising administering the oncolytic virus multiple times to the subject prior to administering the anti-PD1 antibody to the subject.

19. The method of claim 15, comprising administering the oncolytic virus and anti-PD1 antibody to the subject concurrently.

20. The method of claim 11, wherein the herpes virus is HSV-1.

21. A method for treating one or more solid tumors in a subject in need thereof, the method comprising:

administering to the subject an effective amount of an oncolytic herpes virus comprising inserted in its genome (i) a heterologous gene encoding GM-CSF and (ii) a heterologous gene encoding an anti-CTLA-4 antibody, wherein the oncolytic virus is administered by injection into a solid tumor, and wherein the administration induces regression of the solid tumors in the subject.

22. The method of claim 21, wherein the solid tumor is an adenocarcinoma, carcinoma, melanoma, or sarcoma.

23. The method of claim 21, comprising administering multiple doses of 10^4 to 10^{10} pfu of the oncolytic virus to the subject.

24. The method of claim 23, wherein the multiple doses of the oncolytic virus are administered to the subject between 3 days to 3 weeks apart.

25. The method of claim 21, wherein the method further comprises administering an anti-PD1 antibody to the subject.

26. The method of claim 25, comprising administering the oncolytic virus to the subject prior to administering the anti-PD1 antibody to the subject.

27. The method of claim 26, comprising administering the oncolytic virus to the subject once prior to administering the anti-PD1 antibody to the subject.

28. The method of claim 26, comprising administering the oncolytic virus multiple times to the subject prior to administering the anti-PD1 antibody to the subject.

29. The method of claim 25, comprising administering the oncolytic virus and anti-PD1 antibody to the subject concurrently.

30. The method of claim 21, wherein the herpes virus is HSV-1.

* * * * *