



US 20250277226A1

(19) **United States**(12) **Patent Application Publication**
Jankowitsch et al.(10) **Pub. No.: US 2025/0277226 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **IMPROVED BIOTECHNOLOGICAL
METHOD FOR PRODUCING GUANIDINO
ACETIC ACID (GAA) BY INACTIVATION OF
AN AMINO ACID EXPORTER**(71) Applicant: **Evonik Operations GmbH**, Essen
(DE)(72) Inventors: **Frank Jankowitsch**, Sassenberg (DE);
Kay Marin, Borgholzhausen (DE);
Frank Schneider, Halle (DE); **Brigitte
Bathe**, Salzkotten (DE)(73) Assignee: **Evonik Operations GmbH**, Essen
(DE)(21) Appl. No.: **18/562,157**(22) PCT Filed: **May 10, 2022**(86) PCT No.: **PCT/EP2022/062663**

§ 371 (c)(1),

(2) Date: **Nov. 17, 2023**(30) **Foreign Application Priority Data**

May 21, 2021 (EP) 21175138.3

Nov. 16, 2021 (EP) 21208485.9

Publication Classification(51) **Int. Cl.****C12N 15/77** (2006.01)**C07K 14/245** (2006.01)**C07K 14/34** (2006.01)**C12N 9/00** (2006.01)**C12N 9/06** (2006.01)**C12N 9/10** (2006.01)**C12N 9/88** (2006.01)**C12N 15/70** (2006.01)**C12P 13/04** (2006.01)**C12P 13/10** (2006.01)**C12R 1/15** (2006.01)**C12R 1/19** (2006.01)(52) **U.S. Cl.**CPC **C12N 15/77** (2013.01); **C07K 14/245**(2013.01); **C07K 14/34** (2013.01); **C12N****9/0016** (2013.01); **C12N 9/1007** (2013.01);**C12N 9/1018** (2013.01); **C12N 9/1051**(2013.01); **C12N 9/88** (2013.01); **C12N 9/93**(2013.01); **C12N 15/70** (2013.01); **C12P 13/04**(2013.01); **C12P 13/10** (2013.01); **C12R****2001/15** (2021.05); **C12R 2001/19** (2021.05);**C12Y 104/01002** (2013.01); **C12Y 201/01002**(2013.01); **C12Y 201/03003** (2013.01); **C12Y****201/04001** (2013.01); **C12Y 403/02001**(2013.01); **C12Y 603/04005** (2013.01); **C12Y****603/04016** (2013.01)

(57)

ABSTRACT

A microorganism is transformed to be capable of producing guanidinoacetic acid (GAA) having an inactivated amino acid exporter. The microorganism is used in a method for the fermentative production of GAA. Moreover, creatine is produced by a method of fermentative production.

Specification includes a Sequence Listing.

IMPROVED BIOTECHNOLOGICAL METHOD FOR PRODUCING GUANIDINO ACETIC ACID (GAA) BY INACTIVATION OF AN AMINO ACID EXPORTER

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a National Stage entry under § 371 of International Application No. PCT/EP20221062663, filed on May 10, 2022, and which claims the benefit of priority to European Patent Application No. 21175138.3, filed on May 21, 2021, and European Patent Application No. 21208485.9, filed on Nov. 16, 2021. The content of each of these applications is hereby incorporated by reference in its entirety.

REFERENCE TO A SEQUENCE LISTING

[0002] The present invention is accompanied by an ASCII text file as a computer readable form containing the sequence listing entitled, “2024-09-30-005779USPCT-Sequence-List-as-filed.txt”, created on Sep. 30, 2024, with a file size of 55,912 bytes, the content of which is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

Field of the Invention

[0003] The present invention relates to a microorganism transformed to be capable of producing guanidinoacetic acid (GAA) and to a method for the fermentative production of GAA using such microorganism. The present invention also relates to a method for the fermentative production of creatine.

Description of Related Art

[0004] GAA is an organic compound used as animal feed additive (WO2005120246 A1/US2011257075 A1). GAA is a natural precursor of creatine. Therefore, the supplementation of GAA allows for an optimal supply of creatine in the organism.

[0005] The present invention pertains to a method to produce GAA by a fermentative process using industrial feed stocks (e.g. ammonia, ammonium salts and glucose or sugar containing substrates) as starting material. In biological systems GAA and ornithine are formed from arginine and glycine as starting materials by the catalytic action of an L-arginine:glycine-amidino transferase (AGAT; EC 2.1.4.1), which is the first step in creatine biosynthesis:



[0006] Guthmiller et al. (J Biol Chem. 1994 Jul. 1; 269 (26):17556-60) have characterized a rat kidney AGAT by cloning and heterologously expressing the enzyme in *E. coli*. Muenchhoff et al. (FEBS Journal 277 (2010) 3844-3860) report the first characterization of an AGAT from a prokaryote also by cloning and heterologously expressing the enzyme in *E. coli*. For the production of GAA from L-arginine and glycine by a whole cell catalyst Zhang et al. have designed a reconstituted ornithine cycle in *Escherichia coli* by introducing a heterogenous AGAT from different species

(e.g. *Homo sapiens*, *Cylindrospermopsis raciborskii*, *Moorea producens*) and by introducing a citrulline synthesis module (e.g. overexpression of carAB, argF and argI) and an arginine synthesis module (e.g. overexpression of argG, argH; introduction of aspA) into *Escherichia coli* (Yiwen Zhang, Hang Zhou, Yong Tao, and Baixue Lin, ACS Synth. Biol. 2020, 9, 2066-2075).

[0007] Several approaches for increasing the production of one of the starting materials in GAA synthesis, i.e. L-arginine, in microorganisms, particularly bacteria, are also known from literature. An overview for the metabolic engineering of *Corynebacterium glutamicum* (*C. glutamicum*) for L-arginine production is provided by Park et al. (NATURE COMMUNICATIONS/DOI: 10.1038/ncomms5618) They propose random mutagenesis and screening for L-arginine producers of already L-arginine producing *C. glutamicum* strains, e.g. of ATCC 21831 (Nakayama and Yoshida 1974, U.S. Pat. No. 3,849,250 A) and stepwise rational metabolic engineering based on system-wide analysis of metabolism results in a gradual increase in L-arginine production throughout the strain engineering steps. Yim et al. (J Ind Microbiol Biotechnol (2011) 38.1911-1920) could show that inactivation of the argR, gene coding for the central repressor protein ArgR controlling the L-arginine biosynthetic pathway, by disrupting the chromosomal argR gene in *C. glutamicum* leads to an improved arginine-producing strain. Ginesy et al. (Microbial Cell Factories (2015) 14.29) report the successful engineering of *E. coli* for enhanced arginine production Among other, they proposed the deletion of the argR repressor gene.

[0008] Kurahashi et al. (EP1057893 A1) report on methods for increasing the L-arginine producing ability of a microorganism by enhancing L-arginine biosynthesis enzymes utilizing recombinant DNA techniques, e.g. by utilizing a microorganism belonging to the genus *Corynebacterium* or *Brevibacterium* which is made to harbor a recombinant DNA comprising a vector DNA and a DNA fragment containing genes for acetylornithine deacetylase, N-acetylglutamic acid-γ-semialdehyde dehydrogenase, N-acetyl glutamokinase and argininosuccinase derived from a microorganism belonging to the genus *Escherichia*. For an improved L-arginine production the authors further propose a microorganism which is enhanced in an activity of intracellular glutamate dehydrogenase (GDH) and which has an L-arginine producing ability.

[0009] A method of using a genetic recombinant strain, wherein a gene which inhibits the expression of arginine-biosynthesizing operon argR was inactivated has been reported by Suga et al. (U.S. Pat. No. 7,160,705 B2). In particular, the deletion in argR, which controls the arginine operon, has been considered as an important factor in arginine production. In a microorganism of *Corynebacterium*, the argCJBDFR gene, which is involved in arginine biosynthesis, is constituted in the form of an operon and is subjected to feedback-inhibition by intracellular arginine (Sakanyan et al., Microbiology, 142:9-108, 1996), thus imposing a limitation on its high yield L-arginine production. The arginine operon is an operon consisting of genes encoding enzymes involved in the mechanism of L-arginine biosynthesis, and in particular, arginine operon consists of genes encoding enzymes constituting the cyclic steps of L-arginine biosynthesis. Specifically, the arginine operon consists of N-acetylglutamyl phosphate reductase (ArgC), glutamate N-acetyltransferase (ArgJ), N-acetylglutamate

kinase (ArgB), acetylomithine aminotransferase (ArgD), ornithine carbamoyltransferase (ArgF), and the arginine repressor (ArgR). These enzymes are involved in the continuous enzyme reactions of L-arginine biosynthesis.

[0010] According to the literature, the amino acid exporter LysE catalyzes the cellular export of L-lysine, but also of L-arginine and L-citrulline. LysG activates the transcription of the gene *lysE*, which encodes LysE the amino acid exporter. LysG requires a co-inducer such as L-lysine, L-arginine, L-citrulline or L-histidine (Lubitz et al. (2016). "Roles of export genes *cmgA* and *lysE* for the production of L-arginine and L-citrulline by *Corynebacterium glutamicum*." Appl Microbiol Biotechnol 100(19): 8465-8474). Lubitz et al. use the *C. glutamicum* strain ARG2 (Peters-Wendisch et al (2014) Engineering biotin prototrophic *Corynebacterium glutamicum* strains for amino acid, diamine and carotenoid production. J Biotechnol. doi:10.1016/j.jbiotec.2014.01.023) which is characterized by the inactivation of the *argR* gene by deletion in combination with the plasmid-based expression of a feedback-resistant allele of ArgB (*ArgB^{br}*). For this strain carrying both modifications the authors describe the accumulation of L-arginine in the supernatant of the cultures. Furthermore, Lubitz et al describe that in such a strain the inactivation of the genes *lysE* and *cmg*, encoding for membrane proteins, results in a reduction of the arginine formation.

[0011] Ginesy et al. (M. Ginesy et al., Microbiol Cell Factories (2015) 14:29, DOI 10.1186/s12934-015-0211-y) could show that arginine production with an *E. coli* producing strain having among other a deleted *argR* repressor gene can be increased by overexpression of the arginine exporter system.

[0012] However, in order to obtain relatively high L-arginine concentrations in the cell it is necessary to prevent L-arginine exportation. The amino acid exporter LysE counteracts the intracellular arginine concentration and reduces the substrate availability by efficiently transporting the substrate arginine from the cell. In addition, the citrulline from arginine biosynthesis is also secreted into the medium by an active LysE exporter. LysE is regulated by the transcriptional activator LysG (Bellmann, A., et al. (2001). "Expression control and specificity of the basic amino acid exporter LysE of *Corynebacterium glutamicum*." Microbiology (Reading) 147 (Pt 7): 1765-1774).

[0013] Fan Wenchao discloses a method for the production of creatine by fermentation of non-pathogenic microorganisms, such as *Corynebacterium glutamicum* (CN106065411 A). The microorganism has the following biotransformation functions: glucose conversion to L-glutamic acid; conversion of L-glutamic acid to N-acetyl-L-glutamic acid; conversion of N-acetyl-L-glutamic acid to N-acetyl-L-glutamic acid semialdehyde; conversion of N-acetyl-L-glutamic acid semialdehyde to N-acetyl-L-ornithine; conversion of N-acetyl-L-ornithine to L-ornithine; conversion of L-ornithine to L-citrulline; conversion of L-citrulline to arginino-succinic acid; conversion of arginino-succinic acid to L-arginine; conversion of L-arginine to guanidinoacetic acid; and, finally, conversion of guanidinoacetic acid to creatine Fan Wenchao proposes, that the microorganism overexpresses one or more enzymes selected from the group consisting of N-acetylglutamate synthase, N-acetylomithine- δ -aminotransferase, N-acetylomithinase, omithine-carbamoyl transferase, argininosuccinate synthetase, glycine amidino-transferase (EC: 2.1. 4.1),

and guanidinoacetate N-methyltransferase (EC: 2.1.1.2). The microorganism overexpresses preferably glycine aminotransferase (L-arginine:glycine amidinotransferase) and guanidinoacetate N-methyltransferase.

SUMMARY OF THE INVENTION

[0014] The problem underlying the present invention is to provide an improved microorganism transformed to be capable of producing guanidinoacetic acid (GAA) and to a method for the fermentative production of GAA using such microorganism.

DETAILED DESCRIPTION OF THE INVENTION

[0015] The problem is solved by a microorganism having an increased ability to provide L-arginine compared with the ability of the wildtype microorganism and comprising at least one gene coding for a protein having the function of an L-arginine:glycine amidinotransferase and having a decreased activity of a protein having the function of an arginine exporter compared with the activity of the respective protein in the wildtype microorganism at the same status of the cell cycle.

[0016] This means that the protein having the function of an arginine exporter in the microorganism according to the present invention shows at any time and circumstances during the cell cycle an artificially designed decreased activity compared with the activity of the respective protein in the wildtype microorganism at the same time and circumstances during the cell cycle.

[0017] The microorganism according to the present invention is preferably a genetically modified organism (GMO) that does not naturally occur. In a GMO the genetic material has been altered using genetic engineering techniques. Preferably, at least one gene coding for a protein having the function of an L-arginine:glycine amidinotransferase has been introduced using genetic engineering techniques. Preferably, also the decreased activity of the protein having the function of an arginine exporter compared with the activity of the respective protein in the wildtype microorganism has been achieved using genetic engineering techniques.

[0018] Proteins having the function of an L-arginine:glycine amidinotransferase (AGAT) belong to the amidinotransferase family. The amidinotransferase family comprises glycine (EC:2.1.4.1) and inosamine (EC.2.1.4.2) amidinotransferases, enzymes involved in creatine and streptomycin biosynthesis respectively. This family also includes arginine deiminases, EC:3.5.3.6. These enzymes catalyse the reaction: $\text{arginine} + \text{H}_2\text{O} \rightleftharpoons \text{citrulline} + \text{NH}_3$. Also found in this family is the *Streptococcus* anti tumour glycoprotein. Enzymes or proteins with an L-arginine:glycine-amidinotransferase (AGAT) activity are also described to possess a conserved domain that belongs to the PFAM Family: Amidinotransf (PF02274) (Marchler-Bauer A et al. (2017), "CDD/SPARCLE: functional classification of proteins via subfamily domain architectures.", Nucleic Acids Res. 45 (D1):D200-D203.) as described also in the following publications: Pissowotzki K et al., Mol Gen Genet 1991; 231: 113-123 (PUBMED:1661369 EPMC:1661369); D'Hooghe I et al., J Bacteriol 1997; 179:7403-7409 (PUBMED: 9393705 EPMC:9393705); Kanaoka M et al., Jpn J Cancer Res 1987; 78:1409-1414 (PUBMED.3123442 EPMC. 3123442). Particular examples of AGATs are those of

Moorea producens, *Homo sapiens*, *Rattus norvegicus*, *Galeopterus variegatus*, and of *Cylindrospermopsis raciborskii*.

[0019] In the context of the present invention, a microorganism having an improved ability to provide L-arginine means a microorganism producing or recycling L-arginine in excess of its own need. This property may be achieved by selection of microorganisms that are natural L-arginine producers or may have acquired the ability to produce L-arginine by mutation. Examples for such L-arginine producing microorganisms are e.g. *C. glutamicum* ATCC 21831 or those disclosed by Park et al. (NATURE COMMUNICATIONS DOI: 10.1038/ncomms5618) or by Ginesy et al. (Microbial Cell Factories (2015) 14:29).

[0020] In one embodiment of the present invention, the argR gene coding for the arginine responsive repressor protein ArgR in the microorganism according to the present invention is attenuated or deleted.

[0021] The activity of an enzyme having the function of a carbamoyl phosphate synthase (EC 6.3.4.16, e.g. CarAB) in the microorganism according to the present invention may be increased compared to the respective enzymic activity in the wildtype microorganism. This may be achieved by a mutation and/or overexpression of a gene coding for the enzyme having the function of a carbamoyl phosphate synthase.

[0022] Furthermore, in the microorganism according to the present invention at least one or more of the genes coding for an enzyme of the biosynthetic pathway of L-ornithine and L-arginine, comprising argF/argF2 coding for an ornithine carbamoyl transferase, argG coding for an argininosuccinate synthetase and argH coding for an argininosuccinate lyase may be overexpressed.

[0023] Additionally or alternatively, at least one or more of the genes coding for an enzyme of the biosynthetic pathway of L-ornithine and L-arginine, comprising gdh coding for a glutamate dehydrogenase, argJ coding for an ornithine acetyltransferase, argB coding for an acetyl glutamate kinase, argC coding for an acetylglutamylphosphate reductase and argD coding for an acetylornithine aminotransferase, may be overexpressed in the microorganism according to the present invention.

[0024] Overexpression of a gene is generally achieved by increasing the copy number of the gene and/or by functionally linking the gene with a strong promoter and/or by enhancing the ribosomal binding site and/or by codon usage optimization of the start codon or of the whole gene or a combination comprising a selection of all methods mentioned above.

[0025] In the microorganism according to the present invention the gene coding for the protein having the function of an L-arginine:glycine amidinotransferase may be heterologous.

[0026] The microorganism according to the present invention is preferably recombinant and the gene coding for the protein having the function of an L-arginine:glycine amidinotransferase (AGAT) is preferably heterologous.

[0027] A heterologous gene means that the gene has been inserted into a host organism which does not naturally have this gene. Insertion of the heterologous gene in the host is performed by recombinant DNA technology. Microorganisms that have undergone recombinant DNA technology are called transgenic, genetically modified or recombinant.

[0028] In the microorganism of the present invention the gene coding for a protein having the function of an L-argi-

nine:glycine amidinotransferase may further be overexpressed. Overexpression of a gene is generally achieved by increasing the copy number of the gene and/or by functionally linking the gene with a strong promoter and/or by enhancing the ribosomal binding site and/or by codon usage optimization of the start codon or of the whole gene or a combination comprising a selection or all methods mentioned above.

[0029] The protein having the function of an L-arginine:glycine amidinotransferase (AGAT) encoded by at least one respective gene in the microorganism of the present invention may e.g. comprise an amino acid sequence which is at least 70% identical, preferably 80% or at least 90% identical to the amino acid sequence according to SEQ ID NO. 13, i.e. the AGAT of *Moorea producens* 30 ("AGAT_Mpr"). In a further embodiment of the present invention the amino acid sequence of the L-arginine:glycine amidinotransferase is identical to amino acid sequence according to SEQ ID NO. 13 (cf. Database UniProt, 15 Feb. 2017, "Glycine amidinotransferase", XP055706853, EBI accession no. UNIPROT: A0A1D8TKD3). The sequence of wildtype DNA coding for the *Moorea producens* AGAT is SEQ ID NO: 12, the corresponding DNA sequence that has been codon optimized for *C. glutamicum* is SEQ ID NO: 14.

[0030] The protein having the function of an L-arginine:glycine amidinotransferase encoded by at least one respective gene in the microorganism of the present invention may e.g. also comprise an amino acid sequence which is at least 70% homologous, preferably at least 80% or at least 90% identical to the amino acid sequence of the AGAT of *Cylindrospermopsis raciborskii* ATW205 (J Muenchhoff et al., FEBS Journal 277 (2010) 3844-3860).

[0031] The protein having the function of an L-arginine:glycine amidinotransferase encoded by at least one respective gene in the microorganism of the present invention may comprise an amino acid sequence which is at least 70% homologous, preferably at least 80% or at least 90% identical to the amino acid sequence of the AGAT of *Galeopterus variegatus*.

[0032] The protein having the function of an L-arginine:glycine amidinotransferase in the microorganism of the present invention may comprise an amino acid sequence which is at least 70% homologous, preferably at least 80% or at least 90% homologous to the amino acid sequence of the AGAT of *Homo sapiens*, e.g. the AGAT of *Homo sapiens* itself (A. Humm, Biochem. J. (1997) 322, 771-776) or the AGAT of *Rattus norvegicus*.

[0033] Generally, the overexpression of a gene, according to the present invention, is achieved by increasing the copy number of the gene and/or by an enhancement of regulatory factors, e.g. by functionally linking the gene with a strong promoter and/or by enhancing the ribosomal binding site and/or by codon usage optimization of the start codon or of the whole gene. The enhancement of such regulatory factors which positively influence gene expression can, for example, be achieved by modifying the promoter sequence upstream of the structural gene in order to increase the effectiveness of the promoter or by completely replacing said promoter with a more effective or a so-called strong promoter. Promoters are located upstream of the gene. A promoter is a DNA sequence consisting of about 40 to 50 base pairs and which constitutes the binding site for an RNA polymerase holoenzyme and the transcriptional start point, whereby the strength of expression of the controlled poly-

nucleotide or gene can be influenced. Generally, it is possible to achieve an overexpression or an increase in the expression of genes in bacteria by selecting strong promoters, for example by replacing the original promoter with strong, native (originally assigned to other genes) promoters or by modifying certain regions of a given, native promoter (for example its so-called -10 and -35 regions) towards a consensus sequence, e.g. as taught by M. Patek et al. (Microbial Biotechnology 6 (2013), 103-117) for *C. glutamicum*. An example for a “strong” promoter is the superoxide dismutase (sod) promoter (“Psod”; Z. Wang et al., Eng. Life Sci. 2015, 15, 73-82). A “functional linkage” is

frequency. Instead, an organism will typically show certain preferences for specific codons meaning that these codons are found more frequently in the coding sequence of transcribed genes of an organism. If a certain gene foreign to its future host, i.e. from a different species, should be expressed in the future host organism the coding sequence of said gene should then be adjusted to the codon usage of said future host organism (i.e. codon usage optimization).

[0035] Table 1 shows the different names of enzymes involved in or contributing to arginine biosynthesis in different species, i.e. *E. coli*, *C. glutamicum* and *Pseudomonas putida* (*P. putida*)

TABLE 1

Names of Enzymes in different species					
Enzyme name	Alias	EC Number	<i>E. coli</i>	<i>C. glutamicum</i>	<i>P. putida</i>
glutamate	NADP-SPECIFIC GLUTAMATE	1.4.1.4	gdhA	gdh	gdhA/ gdhB
dehydrogenase	DEHYDROGENASE	(1.4.1.2 in <i>P. putida</i>)			
N-acetyl glutamokinase	ACETYLGLUTAMATE KINASE	2.7.2.8	argB	argB	argB
N-acetylglutamyl phosphate reductase	N-acetyl-gamma- glutamylphosphate reductase	1.2.1.38	argC	argC	argC
N-acetylglutamic acid- γ-semialdehyde dehydrogenase	ACETYLORNITHINE AMINOTRANSFERASE (in e.c. bifunctional acetylornithine aminotransferase/ succinyldiaminopimelate aminotransferase)	2.6.1.11/ 2.6.1.17	argD	argD	aruC
acetylornithine deacetylase	bifunctional acetylornithine deacetylase/GLUTAMATE N-ACETYLTRANSFERASE	2.3.1.35/ 2.3.1.1		argJ	argJ
acetylornithine deacetylase		3.5.1.16	argE		argE
	GLUTAMATE N- ACETYLTRANSFERASE	2.3.1.1	argA		argA
ornithine carbamoyltransferase	ornithine carbamoyltransferase 1	2.1.3.3	argI	argF/ argF2	arcB/ argF
Argininosuccinate synthetase	ARGININOSUCCINATE SYNTHASE	6.3.4.5	argG	argG	argG
Argininosuccinate lyase	ARGININOSUCCINATE LYASE	4.3.2.1	argH	argH	argH
carbamoyl-phosphate synthase	carbamoyl-phosphate synthase	6.3.5.5	carAB	carAB	carAB
carbamate kinase	carbamate kinase	2.7.2.2	ybcF/ yqeA		arcC

understood to mean the sequential arrangement of a promoter with a gene, which leads to a transcription of the gene. **[0034]** The genetic code is degenerated which means that a certain amino acid may be encoded by a number of different triplets. The term codon usage refers to the observation that a certain organism will typically not use every possible codon for a certain amino acid with the same

[0036] In the microorganism of the present invention the gene coding for the protein having the function of an arginine exporter may be inactivated or deleted. Furthermore, in the microorganism of the present invention, a gene coding for a transcriptional activator of the gene coding for the protein having the function of an arginine exporter may be deleted.

[0037] The microorganism of the present invention may belong to the genus *Corynebacterium*, preferably *Corynebacterium glutamicum* (*C. glutamicum*), or to the genus *Enterobacteriaceae*, preferably *Escherichia coli* (*E. coli*), or to the genus *Pseudomonas*, preferably *Pseudomonas putida* (*P. putida*).

[0038] In *Corynebacterium glutamicum* the gene coding for the protein having the function of an arginine exporter is *lysE* and the gene coding for the transcriptional activator is *lysG*. In *Escherichia coli* the gene coding for the protein having the function of an arginine exporter is *argO* (*ybjE*). In *Pseudomonas putida* the protein having the function of an arginine exporter is *lysE*.

[0039] The above-mentioned problem is further solved by a method for the fermentative production of guanidino acetic acid (GAA), comprising the steps of a) cultivating the microorganism according to the present invention as defined above in a suitable medium under suitable conditions, and b) accumulating GAA in the medium to form an GAA containing fermentation broth.

[0040] The method according to the present invention may further comprise adding glycine and/or adding L-arginine and/or adding L-ornithine to the medium. Preferably, the medium is supplemented with glycine in a concentration ranging from 0.1 to 300 g glycine/l medium, preferably 0.82 g glycine/l medium, and/or with L-arginine to obtain a concentration ranging from 0.1 to 200 g L-arginine/l medium, preferably 1.9 g L-arginine/l medium.

[0041] The method of the present invention may further comprise the step of isolating GAA from the fermentation broth.

[0042] The method according to the present invention may further comprise the step of drying and/or granulating the GAA containing fermentation broth.

[0043] The present invention further concerns a microorganism as defined above, further comprising a gene coding for an enzyme having the activity of a guanidinoacetate N-methyltransferase (EC: 2.1.1.2). Preferably, the gene coding for an enzyme having the activity of a guanidinoacetate N-methyltransferase is overexpressed.

[0044] The present invention also concerns a method for the fermentative production of creatine, comprising the steps of a) cultivating the microorganism according to the present invention comprising a gene coding for an enzyme having the activity of a guanidinoacetate N-methyltransferase in a suitable medium under suitable conditions, and b) accumulating creatine in the medium to form a creatine containing fermentation broth.

[0045] Preferably, the method further comprises isolating creatine from the creatine containing fermentation broth, creatine may be extracted from fermentation broth by isoelectric point method and/or ion exchange method. Alternatively, creatine can be further purified by a method of recrystallization in water.

EXPERIMENTAL SECTION

a) MATERIALS and METHODS

Chemicals

[0046] Kanamycin solution from *Streptomyces kanamyceticus* was purchased from Sigma Aldrich (St. Louis, USA, Cat. no. K0254). If not stated otherwise, all other chemicals

were purchased analytically pure from Merck (Darmstadt, Germany), Sigma Aldrich (St. Louis, USA) or Carl-Roth (Karlsruhe, Germany).

Cultivation for Cell Proliferation

[0047] If not stated otherwise, cultivation/incubation procedures were performed as follows herewith:

[0048] a. LB broth (MILLER) from Merck (Darmstadt, Germany; Cat. no. 110285) was used to cultivate *E. coli* strains in liquid medium. The liquid cultures (10 ml liquid medium per 100 ml Erlenmeyer flask with 3 baffles) were incubated in the Infors HT Multitron standard incubator shaker from Infors GmbH (Bottmingen, Switzerland) at 30° C. and 200 rpm.

[0049] b. LB agar (MILLER) from Merck (Darmstadt, Germany, Cat. no. 110283) was used for cultivation of *E. coli* strains on agar plates. The agar plates were incubated at 30° C. in an INCU-Line® mini incubator from VWR (Radnor, USA).

[0050] c. Brain heart infusion broth (BHI) from Merck (Darmstadt, Germany, Cat. no. 110493) was used to cultivate *C. glutamicum* strains in liquid medium. The liquid cultures (10 ml liquid medium per 100 ml Erdenmeyer flask with 3 baffles) were incubated in the Infors HT Multitron standard incubator shaker from Infors GmbH (Bottmingen, Switzerland) at 30° C. and 200 rpm.

[0051] d. Brain heart agar (BHI-agar) from Merck (Darmstadt, Germany, Cat. no. 113825) was used for cultivation of *C. glutamicum* strains on agar plates. The agar plates were incubated at 30° C. in an incubator from Heraeus Instruments with Kelvitron® temperature controller (Hanau, Germany).

[0052] e. For cultivating *C. glutamicum* after electroporation, BHI-agar (Merck, Darmstadt, Germany, Cat. no. 113825) was supplemented with 134 g/l sorbitol (Carl Roth GmbH+Co. KG, Karlsruhe, Germany), 2.5 g/l yeast extract (Oxoid/ThermoFisher Scientific, Waltham, USA, Cat. no. LP0021) and 25 mg/l kanamycin. The agar plates were incubated at 30° C. in an incubator from Heraeus Instruments with Kelvitron® temperature controller (Hanau, Germany).

Determining Optical Density of Bacterial Suspensions

[0053] a. The optical density of bacterial suspensions in shake flask cultures was determined at 600 nm (OD600) using the Bio-Photometer from Eppendorf AG (Hamburg, Germany).

[0054] b. The optical density of bacterial suspensions produced in the Wouter Duetz (WDS) micro fermentation system (24-Well Plates) was determined at 660 nm (OD660) with the GENios™ plate reader from Tecan Group AG (Männedorf, Switzerland).

Centrifugation

[0055] a. Bacterial suspensions with a maximum volume of 2 ml were centrifuged in 1.5 ml or 2 ml reaction tubes (e.g. Eppendorf Tubes® 3810X) using an Eppendorf 5417 R benchtop centrifuge (5 min. at 13.000 rpm).

[0056] b. Bacterial suspensions with a maximum volume of 50 ml were centrifuged in 15 ml or 50 ml centrifuge tubes

(e.g. Falcon™ 50 ml Conical Centrifuge Tubes) using an Eppendorf 5810 R benchtop centrifuge for 10 min. at 4.000 rpm.

DNA Isolation

[0057] Plasmid DNA was isolated from *E. coli* cells using the QIAprep Spin Miniprep Kit from Qiagen (Hilden, Germany, Cat. No. 27106) according to the instructions of the manufacturer.

Polymerase Chain Reaction (PCR)

[0058] PCR with a proof reading (high fidelity) polymerase was used to amplify a desired segment of DNA for Sanger sequencing or DNA assembly. Non-proof-reading polymerase Kits were used for determining the presence or absence of a desired DNA fragment directly from *E. coli* or *C. glutamicum* colonies. a. The Phusion® High-Fidelity DNA Polymerase Kit (Phusion Kit) from New England BioLabs Inc. (Ipswich, USA, Cat. No. M0530) was used for template-correct amplification of selected DNA regions according to the instructions of the manufacturer (see Table 2).

TABLE 2

Thermocycling conditions for PCR with Phusion® High-Fidelity DNA Polymerase Kit from New England BioLabs Inc. PCR Program			
Step	Time [min.:sec.]	T [° C.]	Description
1	00:30	98	Initial denaturation step
2	00:05	98	Denaturation step
3	00:30	60	Annealing step
4	00:xx	72	Elongation step 1 min. per kb DNA Repeat step 2 to 4: 35 x
5	05:00	72	Final elongation step
6	Hold	4	Cooling step

[0059] b. Taq PCR Core Kit (Taq Kit) from Qiagen (Hilden, Germany, Cat. No. 201203) was used to amplify a desired segment of DNA in order to confirm its presence. The kit was used according to the instructions of the manufacturer (see Table 3).

TABLE 3

Thermocycling conditions for PCR with Taq PCR Core Kit (Taq Kit) from Qiagen. PCR Program			
Step	Time [min.:sec.]	T [° C.]	Description
1	05:00	94	Initial denaturation step
2	00:30	94	Denaturation step
3	00:30	52	Annealing step
4	01:20	72	Elongation step 1 min. per kb DNA Repeat step 2 to 4: 35 x
5	04:00	72	Final elongation step
6	Hold	4	Cooling step

[0060] c. SapphireAmp® Fast PCR Master Mix (Sapphire Mix) from Takara Bio Inc (Takara Bio Europe S.A.S., Saint-Germain-en-Laye, France, Cat. No. RR350A/B) was used as an alternative to confirm the presence of a desired

segment of DNA in cells taken from *E. coli* or *C. glutamicum* colonies according to the instructions of the manufacturer (see Table 4).

TABLE 4

Thermocycling conditions for PCR with SapphireAmp® Fast PCR Master Mix (Sapphire Mix) from Takara Bio Inc. PCR Program			
Step	Time [min.:sec.]	T [° C.]	Description
1	01:00	94	Initial denaturation step
2	00:05	98	Denaturation step
3	00:05	55	Annealing step
4	00:05	72	Elongation step Repeat step 2 to 4: 30 x
5	04:00	72	Final elongation step
6	Hold	4	Cooling step

[0061] d. All oligonucleotide primers were synthesized by Eurofins Genomics GmbH (Ebersberg, Germany) using the phosphoramidite method described by McBride and Caruthers (1983).

[0062] e. As PCR template either a suitably diluted solution of isolated plasmid DNA or of total DNA isolated from a liquid culture or the total DNA contained in a bacterial colony (colony PCR) was used. For said colony PCR the template was prepared by taking cell material with a toothpick from a colony on an agar plate and placing the cell material directly into the PCR reaction tube. The cell material was heated for 10 sec. with 800 W in a microwave oven type Mikrowave & Grill from SEVERIN Elektrogeräte GmbH (Sundem, Germany) and then the PCR reagents were added to the template in the PCR reaction tube.

[0063] f. All PCR reactions were carried out in PCR cyclers type Mastercycler or Mastercycler nexus gradient from Eppendorf AG (Hamburg, Germany).

Restriction Enzyme Digestion of DNA

[0064] For restriction enzyme digestions either “FastDigest restriction endonucleases (FD)” (ThermoFisher Scientific, Waltham, USA) or restriction endonucleases from New England BioLabs Inc. (Ipswich, USA) were used. The reactions were carried out according to the instructions of the manufacturer’s manual.

Determining the Sizes of DNA Fragments

[0065] a. The sizes of small DNA fragments (<1000 bps) were usually determined by automatic capillary electrophoresis using the QIAxcel from Qiagen (Hilden, Germany).

[0066] b. If DNA fragments needed to be isolated or if the DNA fragments were >1000 bps DNA was separated by TAE agarose gel electrophoresis and stained with GelRed® Nucleic Acid Gel Stain (Biotium, Inc., Fremont, Canada). Stained DNA was visualized at 302 nm.

Purification of PCR Amplificates and Restriction Fragments

[0067] PCR amplificates and restriction fragments were cleaned up using the QIAquick PCR Purification Kit from Qiagen (Hilden, Germany; Cat. No. 28106), according to the manufacturer’s instructions. DNA was eluted with 30 µl 10 mM Tris*HCl (pH 8.5).

Determining DNA Concentration

[0068] DNA concentration was measured using the Nano-Drop Spectrophotometer ND-1000 from PEQLAB BioTechnology GmbH, since 2015 VWR brand (Erlangen, Germany).

Assembly Cloning

[0069] Plasmid vectors were assembled using the “NEBuilder HiFi DNA Assembly Cloning Kit” purchased from New England BioLabs Inc. (Ipswich, USA, Cat. No. E5520). The reaction mix, containing the linear vector and at least one DNA insert, was incubated at 50° C. for 60 min. 0.5 µl of Assembly mixture was used for each transformation experiment.

Chemical Transformation of *E. coli*

[0070] For plasmid cloning, chemically competent “NEB® Stable Competent *E. coli* (High Efficiency)” (New England BioLabs Inc., Ipswich, USA, Cat. No. C3040) were transformed according to the manufacturers protocol. Successfully transformed cells were selected on LB agar supplemented with 25 mg/l kanamycin.

Transformation of *C. glutamicum*

[0071] Transformation of *C. glutamicum* with plasmid-DNA was conducted via electroporation using a “Gene Pulser Xcell” (Bio-Rad Laboratories GmbH, Feldkirchen, Germany) as described by Ruan et al. (2015). Electroporation was performed in 1 mm electroporation cuvettes (Bio-Rad Laboratories GmbH, Feldkirchen, Germany) at 1.8 kV and a fixed time constant set to 5 ms. Transformed cells were selected on BHI-agar containing 134 g/l sorbitol, 2.5 g/l Yeast Extract and 25 mg/l kanamycin.

C. glutamicum Strains

[0072] *Corynebacterium glutamicum* ATCC13032 (DSM 20300, Kinoshita S, Uda S, Shimono M., J. Gen. Appl. Microbiol. 1957; 3(3): 193-205), the *C. glutamicum* wild type strain, is commercially available at the American Type Culture Collection (ATCC) or at the DSMZ-German Collection of Microorganisms and Cell Cultures GmbH.

[0073] *Corynebacterium glutamicum* ATCC21831, an L-arginine producing *C. glutamicum* strain (U.S. Pat. No. 3,849,250 A), is commercially available at the American Type Culture Collection (ATCC).

Determining Nucleotide Sequences

[0074] Nucleotide sequences of DNA molecules were determined by Eurofins Genomics GmbH (Ebersberg, Germany) by cycle sequencing, using the dideoxy chain termination method of Sanger et al. (Proceedings of the National Academy of Sciences USA 74, 5463-5467, 1977). Clone-manager Professional 9 software from Scientific & Educational Software (Denver, USA) was used to visualise and evaluate the sequences.

Glycerol Stocks of *E. coli* and *C. glutamicum* Strains

[0075] For long time storage of *E. coli*- and *C. glutamicum* strains glycerol stocks were prepared. Selected *E. coli* clones were cultivated in 10 ml LB medium supplemented with 2 g/l glucose. Selected *C. glutamicum* clones were cultivated in 10 ml twofold concentrated BHI medium supplemented with 2 g/l glucose. Media for growing plasmid containing *E. coli*- and *C. glutamicum* strains were supplemented with 25 mg/l kanamycin. The medium was contained in 100 ml Erlenmeyer flasks with 3 baffles. It was inoculated with a loop of cells taken from a colony. The culture was then

incubated for 18 h at 30° C. and 200 rpm. After said incubation period 1.2 ml 85% (v/v) sterile glycerol were added to the culture. The obtained glycerol containing cell suspension was then aliquoted in 2 ml portions and stored at –80° C.

Gaa Production in Shake Flask Cultivations

[0076] Shake flask cultivation in 250 mL baffled Erlenmeyer flasks was used to assess the GAA-production of the strains.

[0077] Precultures of the strains were done in 10 ml seed medium (SM). The medium was contained in a 100 ml Erlenmeyer flask. It was inoculated with 100 µl of a glycerol stock culture and the culture was incubated for 24 h at 30° C. and 200 rpm. The composition of the seed medium (SM) is shown in Table 5. Kanamycin was added to cultures were needed to retain plasmids.

TABLE 5

Seed medium (SM)	
Components	Concentration (g/l)
(NH ₄) ₂ SO ₄	20
Yeast extract FM902 (Angel Yeast Co., LTD, Hubei, P.R.China)	20
KH ₂ PO ₄	1
Mg-Sulfat * 7 H ₂ O	0.5
Glucose	20
Kanamycin	0.025
pH = 7.0	

[0078] After said incubation period the optical densities OD600 of the precultures were determined. The volume, needed to inoculate 2.5 ml of production medium (PM) to an OD600 of 0.5, was sampled from the preculture, centrifuged (1 min at 8000 g) and the supernatant was discarded. Cells were then resuspended in 200 µl of production medium.

[0079] The main cultures were started by inoculating the 2.4 ml production medium (PM) containing wells of the 24 Well WDS-Plate with each 100 µl of the resuspended cells from the precultures. The composition of the production medium (PM) is shown in Table 6.

TABLE 6

Production medium (PM)	
Components	Concentration (g/l)
Yeast extract FM902 (Angel Yeast Co., LTD, Hubei, P.R.China)	8
(NH ₄) ₂ SO ₄	40
KH ₂ PO ₄	1.5
MgSO ₄ * 7 H ₂ O	0.5
FeSO ₄ * 7 H ₂ O	0.02
MnSO ₄ * H ₂ O	0.02
Calcium carbonate	20
Glucose	100
Kanamycin	0.025
L-arginine	5 mM
Glycine	20
pH = 7.2	

[0080] The main cultures were incubated for 48 h at 30° C. and 200 rpm in an Infors HT Multitron standard incubator shaker from Infors GmbH (Bottmingen, Switzerland) until complete consumption of glucose. The glucose concentra-

tion in the suspension was analyzed with the blood glucose-meter OneTouch Vita® from LifeScan (Johnson & Johnson Medical GmbH, Neuss, Germany).

[0081] After cultivation the culture suspensions were transferred to 50 ml centrifuge tubes (e.g. Falcon™ 50 ml Conical Centrifuge Tubes). A part of the culture suspension was suitably diluted to measure the OD₆₆₀. Another part of the culture was centrifuged and the concentration of GAA in the supernatant was analyzed as described below.

Quantification of GAA

[0082] Samples were analyzed with an analyzing system from Agilent, consisting of a HPLC “Infinity 1260” coupled with a mass analyzer “Triple Quad 6420” (Agilent Technologies Inc., Santa Clara, USA). Chromatographic separation was done on the Atlantis HILIC Silica column, 4,6λ250 mm, 5 μm (Waters Corporation, Milford, USA) at 35° C. Mobile phase A was water with 10 mM ammonium formate and 0.2% formic acid. Mobile phase B was a mixture of 90% acetonitrile and 10% water, 10 mM ammonium formate were added to the mixture. The HPLC system was started with 100% B, followed by a linear gradient for 22 min and a constant flow rate of 0.6 mL/min to 66% B. The mass analyzer was operated in the ESI positive ionization mode. For detection of GAA the m/z values were monitored by using an MRM fragmentation [M+H]⁺ 118–76. The limit of quantification (LOQ) for GAA was fixed to 7 ppm.

B) Experimental Results

Example 1: Cloning of the Plasmid pK19mobsacB-ΔlysEG for the Chromosomal Deletion of the Genes lysE and lysG in *C.* *Glutamicum* Based Strains

[0083] The gene lysE codes for an exporter protein that catalyzes the efflux of L-Lysine, L-arginine and L-citrulline in *Corynebacterium glutamicum*. The expression of lysE is positively regulated by the gene product of lysG. Both genes are located next to each other but are transcribed divergently.

[0084] In order to inactivate the transporter protein LysE and the positive regulator protein LysG, the plasmid pK19mobsacB-ΔlysEG (SEQ ID NO: 1) was constructed as described for pK18mobsacB-ΔlysEG in Vrljic et. al 1996 (Vrljic, M., et al. (1996). “A new type of transporter with a new type of cellular function: L-lysine export from *Corynebacterium glutamicum*.” Mol Microbiol 22(5): 815-826; <https://doi.org/10.1046/j.1365-2958.1996.01527.x>).

Example 2: Chromosomal Deletion of the Gene argR in ATCC13032

[0085] To improve intracellular L-Arginine formation and L-Arginine recycling from L-Ornithine, the gene argR coding for the central repressor protein ArgR controlling the L-arginine biosynthetic pathway was inactivated.

[0086] Therefore, the plasmid pK18mobsacB_DargR was constructed as follows. Plasmid pK18mobsacB (Schäfer, 1994) was cut using XbaI and the linearized vector DNA (5721 bps) was purified using the “QIAquick Gel Extraction Kit” (Qiagen GmbH, Hilden, Germany).

[0087] For constructing the insert, two DNA fragments were created by PCR with the following pairs of primers (genomic DNA of ATCC13032 as template):

[0088] DargR_If (SEQ ID NO: 2), +DargR_Ir (SEQ ID NO: 3)=left homology arm (983 bps)

[0089] DargR_rf (SEQ ID NO: 4), +DargR_jr (SEQ ID NO: 5)=left homology arm (984 bps)

[0090] The product DNAs were purified using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0091] The linearized plasmid and the PCR products were then assembled using the “NEBuilder HiFi DNA Assembly Cloning Kit” (New England BioLabs Inc., Ipswich, USA, Cat. No. E5520). The resulting deletion vector was named pK18mobsacB_DargR. It was verified by restriction enzyme digestion and DNA sequencing.

[0092] For deleting the argR gene, pK18mobsacB_DargR was transformed into ATCC13032 by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0093] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 μl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for 48 h at 33° C. Colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR and DNA sequencing. The resulting strain was named ATCC13032_DargR.

Example 3: Chromosomal Insertion of the Sod Promoter Upstream of the carAB Operon in ATCC13032_DargR

[0094] To improve the production of L-arginine, the strong sod-promoter was inserted into the genome of ATCC13032_DargR upstream of the carAB operon. Therefore, the plasmid pK18mobsacB_Psod-carAB was constructed as follows. pK18mobsacB was cut using EcoRI+HindIII and the linearized vector DNA (5670 bps) was cut out of an agarose gel. The DNA was extracted using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0095] For constructing the insert, three DNA fragments were created by PCR with the following pairs of primers (genomic DNA of ATCC13032 as template):

[0096] PsodcarAB-LA-F (SEQ ID NO: 6)+PsodcarAB-LA-R (SEQ ID NO: 7)=left homology arm (1025 bps)

[0097] PsodcarAB-F (SEQ ID NO: 8)+PsodcarAB-R (SEQ ID NO: 9)=sod-promoter (250 bps)

[0098] PsodcarAB-RA-F (SEQ ID NO: 10)+PsodcarAB-RA-R (SEQ ID NO: 11)=right homology arm (944 bps)

[0099] The product DNAs were purified using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany). The linearized plasmid and the PCR products were

then assembled using the “NEBuilder HiFi DNA Assembly Cloning Kit” (New England BioLabs Inc., Ipswich, USA, Cat. No. E5520). Proper plasmid clones were identified by restriction digestion and DNA sequencing.

[0100] The resulting plasmid pK18mobsacB_Psod-carAB was then transformed into ATCC13032_DargR by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0101] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 µl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for 48 h at 33° C. The colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR and DNA sequencing for the appropriate integration of the sod promoter. The resulting strain was named ATCC13032_DargR_Psod-carAB.

Example 4: Chromosomal Deletion of the Genes lysE and lysG in ATCC13032_DargR_Psod-carAB

[0102] The gene lysE codes for an exporter protein that catalyzes the efflux of L-Lysine, L-arginine and L-citrulline in *Corynebacterium glutamicum*. The expression of lysE is positively regulated by the gene product of lysG. Both genes are located next to each other but are transcribed divergently. For deleting the genes lysE and lysG gene, pK19mobsacB_DlysEG (see example 1) was transformed into ATCC13032_DargR_Psod-carAB by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0103] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 µl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for 48 h at 33° C. Colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR and DNA

sequencing. The resulting strain was named ATCC13032_DargR_Psod-carAB_DlysEG.

Example 5: Cloning of the Gene AGAT-Mp Coding for an L-Arginine:Glycine Amidinotransferase (AGAT, EC 2.1.4.1) from *Moorea producens*

[0104] *Moorea producens* is a filamentous cyanobacterium. The genome of the *Moorea producens* strain PAL-8-15-08-1 was published by Leao et al. (Leao T, Castelgo G, Korobeynikov A, Monroe E A, Podell S, Glukhov E, Allen E E, Gerwick W H, Gerwick L. Proc Natl Acad Sci USA. 2017 Mar. 21; 114(12):3198-3203. doi: 10.1073/pnas.1618556114; Genbank accession Number CP017599.1). It contains an open reading frame coding for a L-arginine:glycine amidinotransferase (AGAT, EC 2.1.4.1; locus_tag BJP34_00300 shown in SEQ ID NO: 12). SEQ ID NO: 13 shows the derived amino acid sequence (Genbank accession Number WP_070390602).

[0105] Using the software tool “Optimizer” (<http://genomes.urv.es/OPTIMIZER/>) this amino acid sequence was translated back into a DNA sequence optimized for the codon usage of *C. glutamicum* (SEQ ID NO: 14).

[0106] A segment of the optimized gene, consisting of base pairs 13-1142, was expanded with a BsmBI restriction site at its 5'-end. At the 3'-end a second stop-codon, the lysS-terminator from *C. glutamicum* and a BsmBI restriction site were added. The resulting DNA sequence (SEQ ID NO: 15) was ordered for gene synthesis from Invitrogen/Genent (Thermo Fisher Scientific, Waltham, USA) and it was delivered as part of a cloning plasmid with an ampicillin resistance gene (designated as pMA-RQ_AGAT_Mp_opt). A second DNA segment was designed, consisting of a sequence for assembly cloning, a promoter sequence, a ribosomal binding site and the first 81 nucleotides of the optimized AGAT-Mp gene. The sequence was ordered as a linear DNA string for gene synthesis from Invitrogen/Genent (Thermo Fisher Scientific, Waltham, USA; SEQ ID NO: 16).

Example 6: Cloning of the Expression Plasmid pLIB_pBL1_AGAT-Mp

[0107] The *E. coli*-*C. glutamicum* shuttle plasmid pLIB_pBL1 has the replication origin from pBL1, the pSC101 replication origin, a kanamycin resistance gene and the BioBricks Terminator BBa_B1006 downstream of a NotI restriction site (SEQ ID NO: 17). It was digested using the restriction endonuclease NotI and the DNA was purified with the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0108] Using the primers AGAT_f (SEQ ID NO: 18) and AGAT_r (SEQ ID NO: 19) a DNA fragment was amplified by PCR with pMA-RQ_AGAT_Mp_opt as a template. The PCR-product was purified using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0109] The linearized plasmid, the promoter containing DNA-string and the PCR product were assembled using the “NEBuilder HiFi DNA Assembly Cloning Kit” (New England BioLabs Inc., Ipswich, USA, Cat. No. E5520). The assembly product was transformed into “NEB Stable Competent *E. coli* (High Efficiency)” (New England Biolabs, Ipswich, USA) and cells were grown on LB agar containing 25 mg/l kanamycin. Proper plasmid clones were identified

by restriction digestion and DNA sequencing. The resulting plasmid was named pLIB_pBL1_AGAT-Mp.

Example 7: Chromosomal Expression of the Arginine Biosynthesis Genes *argF*, *argG*, *argH* Under The Control of the Strong Constitutive Promoter Pg3

[0110] For enhancing the activities of *ArgF*, *ArgG* and *ArgH*, additional copies of the corresponding genes were inserted into the genome. A synthetic operon was designed, consisting of Pg3, *argF*, *argG*, *argH* and flanking regions for genomic integration (SEQ ID NO: 20).

[0111] The DNA sequence was ordered for gene synthesis from Invitrogen/Genecart (Thermo Fisher Scientific, Waltham, USA) and it was delivered as part of a cloning plasmid with an ampicillin resistance gene (designated as pMA-RQ_argFGH).

[0112] Using the primers *argFGH_f* (SEQ ID NO: 21) and *argFGH_r* (SEQ ID NO: 22) a DNA fragment was amplified by PCR with pMA-RQ_argFGH as a template. The PCR-product was purified using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0113] The plasmid pK18mobsacB (Schäfer, A. et al., Gene. 1994 Jul. 22; 145(1):69-73. doi: 10.1016/0378-1119(94)90324-7) was cut using EcoRI+HindIII and the linearized vector DNA (5870 bps) was cut out of an agarose gel. The DNA was extracted using the “QIAquick PCR Purification Kit” (Qiagen GmbH, Hilden, Germany).

[0114] The linearized plasmid and the PCR-product were assembled using the “NEBuilder HiFi DNA Assembly Cloning Kit” (New England BioLabs Inc., Ipswich, USA, Cat. No. E5520). The assembly product was transformed into “NEB Stable Competent *E. coli* (High Efficiency)” (New England Biolabs, Ipswich, USA) and cells were grown on LB agar containing 25 mg/l kanamycin. Proper plasmid clones were identified by restriction digestion and DNA sequencing. The resulting plasmid was named pK18_IBcg0054::Pg3-argFGH (SEQ ID NO: 23).

[0115] For inserting the synthetic operon, pK18_IBcg0054::Pg3-argFGH was transformed into ATCC13032_DargR by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0116] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 µl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for 48 h at 33° C. Colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR and DNA sequencing. The resulting strain was named ATCC13032_DargR_IBcg0054::Pg3-argFGH.

Example 8: Chromosomal Deletion of the Genes *lysE* and *lysG* in ATCC13032_DargR_IBcg0054::Pg3-argFGH

[0117] The gene *lysE* codes for an exporter protein that catalyzes the efflux of L-Lysine, L-arginine and L-citrulline in *Corynebacterium glutamicum*. The expression of *lysE* is positively regulated by the gene product of *lysG*. Both genes are located next to each other but are transcribed divergently.

[0118] For deleting the genes *lysE* and *lysG* gene, pK19mobsacB_DlysEG (see example 1) was transformed into ATCC13032_DargR_IBcg0054::Pg3-argFGH by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0119] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 µl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for 48 h at 33° C. Colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR and DNA sequencing. The resulting strain was named ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG.

Example 9: Chromosomal Deletion of the Genes *lysE* and *lysG* in ATCC21831_DlysEG

[0120] The gene *lysE* codes for an exporter protein that catalyzes the efflux of L-Lysine, L-arginine and L-citrulline in *Corynebacterium glutamicum*. The expression of *lysE* is positively regulated by the gene product of *lysG*. Both genes are located next to each other but are transcribed divergently.

[0121] For deleting the genes *lysE* and *lysG* gene, pK19mobsacB_DlysEG (see example 1) was transformed into ATCC21831_DlysEG by electroporation. Chromosomal integration (resulting from a first recombination event) was selected by plating on BHI agar supplemented with 134 g/l sorbitol, 2.5 g/l yeast extract and 25 mg/l kanamycin. The agar plates were incubated for 48 h at 33° C.

[0122] Individual colonies were transferred onto fresh agar plates (with 25 mg/l kanamycin) and incubated for 24 h at 33° C. Liquid cultures of these clones were cultivated for 24 h at 33° C. in 10 ml BHI medium contained in 100 ml Erlenmeyer flasks with 3 baffles. To isolate clones that have encountered a second recombination event, an aliquot was taken from each liquid culture, suitably diluted and plated (typically 100 to 200 µl) on BHI agar supplemented with 10% saccharose. These agar plates were incubated for

48 h at 33° C. Colonies growing on the saccharose containing agar plates were then examined for kanamycin sensitivity. To do so a toothpick was used to remove cell material from the colony and to transfer it onto BHI agar containing 25 mg/l kanamycin and onto BHI agar containing 10% saccharose. The agar plates were incubated for 60 h at 33° C. Clones that proved to be sensitive to kanamycin and resistant to saccharose were examined by PCR. The resulting strain was named ATCC2183_DlysEG.

Example 10: Transformation of *C. glutamicum*
Strains with pLIB_pBL1_AGAT-Mp

[0123] The following strains of *C. glutamicum* were transformed with pLIB_pBL1_AGAT-Mp by electroporation and plasmid containing cells were selected with 25 mg/l kanamycin. The resulting plasmid containing strains are shown in Table 7.

TABLE 7

List of plasmid-containing <i>C. glutamicum</i> strains		
Plasmid	Recipient strain	Resulting strain
pLIB_pBL1	ATCC13032	ATCC13032/pLIB_pBL1
pLIB_pBL1_AGAT-Mp	ATCC13032	ATCC13032/pLIB_pBL1_AGAT-Mp
pLIB_pBL1_AGAT-Mp	ATCC13032_DargR_Psod-carAB	ATCC13032_DargR_Psod-carAB/pLIB_pBL1_AGAT-Mp
pLIB_pBL1_AGAT-Mp	ATCC13032_DargR_Psod-carAB_DlysEG	ATCC13032_DargR_Psod-carAB_DlysEG/pLIB_pBL1_AGAT-Mp
pLIB_pBL1_AGAT-Mp	ATCC13032_DargR_IBcg0054::Pg3-argFGH	ATCC13032_DargR_IBcg0054::Pg3-argFGH/pLIB_pBL1_AGAT-Mp
pLIB_pBL1_AGAT-Mp	ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG	ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG/pLIB_pBL1_AGAT-Mp
pLIB_pBL1	ATCC21831	ATCC21831/pLIB_pBL1
pLIB_pBL1_AGAT-Mp	ATCC21831	ATCC21831/pLIB_pBL1_AGAT-Mp
pLIB_pBL1_AGAT-Mp	ATCC21831_DlysEG	ATCC21831_DlysEG/pLIB_pBL1_AGAT-Mp

C. glutamicum ATCC13032: commonly used wild type strain (Kinoshita et al., J. Gen. Appl. Microbiol. 1957; 3(3): 193-205)
 ATCC13032_DargR_Psod-carAB: Increased ability to produce L-arginine due to the reduced activity of the ArgR regulator protein and integration of the strong sod promoter upstream of carAB in ATCC13032.
 ATCC13032_DargR_Psod-carAB_DlysEG: Increased ability to produce L-arginine due to the reduced activity of the ArgR regulator protein and integration of the strong sod promoter upstream of carAB in ATCC13032. Reduced export activity of L-arginine.
 ATCC13032_DargR_IBcg0054::Pg3-argFGH: Increased ability to produce L-arginine due to the reduced activity of the ArgR regulator protein and genomic integration of an argFGH expression cassette under the control of the strong promoter Pg3 in ATCC13032.
 ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG: Increased ability to produce L-arginine due to the reduced activity of the ArgR regulator protein and genomic integration of an argFGH expression cassette under the control of the strong promoter Pg3 in ATCC13032. Reduced export activity of L-arginine.
C. glutamicum strain ATCC21831 (Park et al., Nat Commun. 2014 Aug. 5; 5: 4618) synthesizes L-arginine from primary substrates like ammonia and glucose.
C. glutamicum strain ATCC21831_DlysEG (Park et al., Nat Commun. 2014 Aug. 5; 5: 4618) synthesizes L-arginine from primary substrates like ammonia and glucose. Reduced export activity of L-arginine

Example 11: Impact of Increased Ability to
Produce L-Arginine and Reduced Arginine Export
on GAA Production

[0124] To assess the combined impact of the increased ability to produce L-arginine and reduced L-arginine export, on GAA production, strains ATCC13032, ATCC13032/pLIB_pBL1, ATCC13032/pLIB_pBL1_AGAT-Mp, ATCC13032_DargR_Psod-carAB/pLIB_pBL1_AGAT-Mp, ATCC13032_DargR_Psod-carAB_DlysEG/pLIB_pBL1_AGAT-Mp, ATCC13032_DargR_IBcg0054::Pg3-argFGH/pLIB_pBL1_AGAT-Mp and ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG/pLIB_pBL1_AGAT-Mp were cultivated in the Wouter Duetz system in production medium, and the resulting GAA titers were determined.

TABLE 8

Combined impact of increased ability to produce L-arginine and reduced Arginine export on GAA production.	
Strain	GAA (g/L)
ATCC13032	0
ATCC13032/pLIB_pBL1	0
ATCC13032/pLIB_pBL1_AGAT-Mp	1.1
ATCC13032_DargR_Psod-carAB/pLIB_pBL1_AGAT-Mp	2.2
ATCC13032_DargR_Psod-carAB_DlysEG/pLIB_pBL1_AGAT-Mp	2.3
ATCC13032_DargR_IBcg0054::Pg3-argFGH/pLIB_pBL1_AGAT-Mp	1.7
ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG/pLIB_pBL1_AGAT-Mp	1.9

[0125] As shown in Table 8, ATCC13032_DargR_Psod-carAB/pLIB_pBL1_AGAT-Mp and ATCC13032_DargR_IBcg0054::Pg3-argFGH/pLIB_pBL1_AGAT-Mp having a polynucleotide coding for the AGAT from *Moorea producens*, a deleted argR gene and increased expression of carAB or argFGH genes produced 2.2 g/l or 1.7 g/l of GAA respectively.

[0126] Strains ATCC13032_DargR_Psod-carAB_DlysEG/pLIB_pBL1_AGAT-Mp and ATCC13032_DargR_IBcg0054::Pg3-argFGH_DlysEG/pLIB_pBL1_AGAT-Mp also have a polynucleotide coding for the AGAT from *Moorea producens*, a deleted argR gene and increased expression of carAB or argFGH genes. In addition, the genes lysEG are inactivated and they have reduced L-arginine export activity. These strains produced 2.3 g/l and 1.9 g/l GAA respectively, which is improved as compared to strains without reduced L-arginine export.

[0127] We conclude that the combination of enzymatic AGAT activity, increased ability to provide L-arginine and reducing L-arginine export improves GAA production.

Example 12: Impact of Increased Ability to Produce L-Arginine and Reduced Arginine Export on GAA Production Using the L-Arginine Producer Strain ATCC21831

[0128] To assess the combined impact of the increased ability to produce L-arginine and reduced L-arginine export, on GAA production, strains ATCC21831, ATCC21831/pLIB_pBL1 and ATCC21831/pLIB_pBL1_AGAT-Mp were cultivated in the Wouter Duetz system in production medium, and the resulting GAA titers were determined.

TABLE 9

Combined impact of increased ability to produce L-arginine and reduced Arginine export on GAA production using the L-Arginine producer strain ATCC21831.	
Strain	g/L
ATCC21831	0.0
ATCC21831/pLIB_pBL1	0.0
ATCC21831_DlysEG/pLIB_pBL1	0.0
ATCC21831/pLIB_pBL1_AGAT-Mp	5.0
ATCC21831_DlysEG/pLIB_pBL1_AGAT-Mp	5.6

[0129] As shown in Table 9, ATCC21831, ATCC21831/pLIB_pBL1 and ATCC21831_DlysEG/pLIB_pBL1 are not able to produce GAA.

[0130] Strain ATCC21831/pLIB_pBL1_AGAT-Mp also carries a polynucleotide coding for the AGAT from *Moorea producens* and produces 5.0 g/l GAA. Furthermore, as shown in Table 8, the strain ATCC13032/pLIB_pBL1_AGAT-Mp that carries a polynucleotide coding for the AGAT from *Moorea producens*, but that is not capable to provide additional L-arginine, only produces 1.1 g/L of GAA.

[0131] Strain ATCC21831_DlysEG/pLIB_pBL1_AGAT-Mp also carries a polynucleotide coding for the AGAT from *Moorea producens*, but additionally the genes lysEG are inactivated in this strain and they have reduced L-arginine export activity. This strain produces 5.6 g/l of GAA, which is improved as compared to the strain without reduced L-arginine export.

[0132] We conclude that the combination of enzymatic AGAT activity, increased ability to provide L-arginine and reducing L-arginine export improves GAA production.

Overview Sequences

- [0133] SEQ ID NO: 1: Shows the DNA sequence coding for the plasmide pK19mobsacB-DlysEG which was constructed as described in Vrljic et. al 1996 (Vrljic, M., et al. (1996). "A new type of transporter with a new type of cellular function: L-lysine export from *Corynebacterium glutamicum*." Mol Microbiol 22(5): 815-826)
- [0134] SEQ ID NO: 2: Shows the DNA sequence coding for the primer DargR_If
- [0135] SEQ ID NO: 3: Shows the DNA sequence coding for the primer DargR_Ir
- [0136] SEQ ID NO: 4: Shows the DNA sequence coding for the plasmid DargR_rf

[0137] SEQ ID NO: 5: Shows the DNA sequence coding for the primer DargR_rr

[0138] SEQ ID NO: 6: Shows the DNA sequence coding for the primer PsodcarAB-LA-F

[0139] SEQ ID NO: 7: Shows the DNA sequence coding for the primer PsodcarAB-LA-R

[0140] SEQ ID NO: 8: Shows the DNA sequence coding for the primer PsodcarAB-F

[0141] SEQ ID NO: 9: Shows the DNA sequence coding for the primer PsodcarAB-R

[0142] SEQ ID NO: 10: Shows the DNA sequence coding for the primer PsodcarAB-RA-F

[0143] SEQ ID NO: 11: Shows the DNA sequence coding for the primer PsodcarAB-RA-R

[0144] SEQ ID NO: 12: Shows the open reading frame coding for a L-arginine:glycine amidinotransferase (AGAT, EC 2.1.4.1; locus_tag BJP34_00300)

[0145] SEQ ID NO: 13: Shows the amino acid sequence derived from SEQ ID No: 12 (Genbank accession Number WP_070390602)

[0146] SEQ ID NO: 14: Shows the DNA sequence coding for a L-arginine:glycine amidinotransferase from *Moorea producens* (AGAT, EC 2.1.4.1) optimized for the codon usage of *C. glutamicum*.

[0147] SEQ ID NO: 15: Shows the DNA sequence coding for a L-arginine:glycine amidinotransferase from *Moorea producens* (AGAT, EC 2.1.4.1) optimized for the codon usage of *C. glutamicum* derived from SEQ ID No: 14 with a segment, consisting of base pairs 13-1142, expanded with a BsmBI restriction site at its 5'-end. Furthermore, at the 3'-end a second stop-codon, the lysS-terminator from *C. glutamicum* and a BsmBI restriction site were added.

[0148] SEQ ID NO: 16: DNA segment consisting of a sequence for assembly cloning, a promoter sequence, a ribosomal binding site and the first 81 nucleotides of the optimized AGAT-Mp gene.

[0149] SEQ ID NO: 17: Shows the DNA sequence of the shuttle plasmid pLIB_pBL1 has the replication origin from pBL1, the pSC101 replication origin, a kanamycin resistance gene and the BioBricks Terminator BBa_B1006 downstream of a NotI restriction site

[0150] SEQ ID NO: 18: Shows the DNA sequence of the primer AGAT_f

[0151] SEQ ID NO: 19: Shows the DNA sequence of the primer AGAT_r

[0152] SEQ ID NO: 20 Shows the DNA sequence of promotor Pg3-argFGH

[0153] SEQ ID NO: 21 Shows the DNA sequence of the primer argFGH_f

[0154] SEQ ID NO: 22 Shows the DNA sequence of the primer argFGH_r

[0155] SEQ ID NO: 23 Shows the DNA sequence of the plasmid pK18-IBcg0054::Pg3-argFGH

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1

<211> LENGTH: 6544

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(6544)

<223> OTHER INFORMATION: pK19mobsacB-DlysEG

<220> FEATURE:

<223> OTHER INFORMATION: plasmid constructed as described for
pK18mobsacB-lysEG in Vrljic et. al 1996

<400> SEQUENCE: 1

```
cgataagcta gcttcacgct gccgcaagca ctcaggcgcg aagggtgct aaaggaagcg      60
gaacacgtag aaagccagtc cgcagaaacg gtgctgaccc cggatgaatg tcagctactg      120
ggctatctgg acaagggaaa acgcaagcgc aaagagaaag caggtagctt gcagtgggct      180
tacatggcga tagctagact gggcgggttt atggacagca agcgaaccgg aattgccagc      240
tggggcgccc tctggtaagg ttgggaagcc ctgcaaagta aactggatgg ctttcttgcc      300
gccaaaggatc tgatggcgca ggggatcaag atctgatcaa gagacaggat gaggatcggt      360
tcgcatgatt gaacaagatg gattgcacgc aggttctccg gccgcttggg tggagaggct      420
attcggctat gactgggcac aacagacaat cggtgctctt gatgccgccg tgttcgggct      480
gtcagcgtag gggcgcccg tttcttttgt caagaccgac ctgtccggtg ccctgaatga      540
actcaagac gaggcagcgc ggctatctg gctggccacg acgggcttcc cttgcgcagc      600
tgtgctcgac gttgtcactg aagcgggaag ggactggctg ctattgggag aagtgcgggg      660
gcaggatctc ctgtcatctc acctgtctcc tgccgagaaa gtatccatca tggctgatgc      720
aatgcggcgg ctgcatacgc ttgatccggc tacctgccca ttcgaccacc aagcgaaaca      780
tcgcatcgag cgagcacgta ctcgatgga agccggctct gtcgatcagg atgatctgga      840
cgaagagcat caggggctcg cgccagccga actgttcgcc aggtcaagg cgcggatgcc      900
cgacggcgag gatctcgtag tgaccatgg cgatgcctgc ttgccgaata tcatggtgga      960
aaatggccgc ttttctggat tcactgactg tggcgggctg ggtgtggcgg accgctatca     1020
ggacatagcg ttggtatccc gtgatattgc tgaagagctt ggcggcgaat gggctgaccg     1080
cttctcgtg ctttacggta tcgccgctcc cgattcgtag cgcacgcct tctatcgctt     1140
tcttgacgag ttcttctgag cgggactctg ggggttcgta gaggatcgat cttttttaac     1200
ccatcacata tacctgccgt tcaactattat ttagtgaaat gagatattat gatattttct     1260
gaattgtgat taaaaaggca actttatgcc catgcaacag aaactataaa aaatacagag     1320
aatgaaaaga aacagataga ttttttagtt ctttaggccc gtagtctgca aatcctttta     1380
tgattttcta tcaaacaaaa gaggaataa gaccagttag aatccaaacg agagtctaata     1440
agaatgaggt cgaagagtaa atcgcgccgg tttgttactg ataaagcagg caagacctaa     1500
aatgtgtaaa gggcaaatgt tatacttttg cgtcaccctt tacatatttt aggtcttttt     1560
ttattgtgag taactaactt gccatcttca aacaggaggg ctggaagaag cagaccgcta     1620
acacagtaca taaaaaagga gacatgaacg atgaacatca aaaagtttgc aaaacaagca     1680
acagtattaa ctttacttac cgcactgctg gcaggaggcg caactcaagc gtttgcgaaa     1740
gaaacgaacc aaaagccata taaggaaaca tacggcattt cccatattac acgcatgat     1800
```

-continued

atgctgcaaa	tccctgaaca	gcaaaaaaat	gaaaaaatatc	aagtttctga	atttgattcg	1860
tccacaatta	aaaatatctc	ttctgcaaaa	ggcctggacg	tttgggacag	ctggccatta	1920
caaaacgctg	acggcactgt	cgcaaaactat	cacggctacc	acatcgctctt	tgcattagcc	1980
ggagatccta	aaaatgcgga	tgacacatcg	atttacatgt	tctatcaaaa	agtcggcgaa	2040
acttctattg	acagctggaa	aaacgctggc	cgctcttcta	aagacagcga	caaattcgat	2100
gcaaatgatt	ctatcctaaa	agaccaaaaca	caagaatggg	caggttcagc	cacatttaca	2160
tctgacggaa	aaatcgtttt	attctacact	gatttctccg	gtaaacatta	cggcaaaaca	2220
acactgacaa	ctgcacaagt	taacgtatca	gcacagaca	gctctttgaa	catcaacggg	2280
gtagaggatt	ataaatcaat	ctttgacggg	gacggaaaaa	cgtatcaaaa	tgtacagcag	2340
ttcatcgatg	aaggcaacta	cagctcaggc	gacaaccata	cgctgagaga	tctctactac	2400
gtagaagata	aaggccacaa	atacttagta	tttgaagcaa	acactggaac	tgaagatggc	2460
taccaaggcg	aagaatcttt	atttaacaaa	gcatactatg	gcaaaagcac	atcattcttc	2520
cgtaagaaa	gtcaaaaact	tctgcaaaagc	gataaaaaac	gcacggctga	gtagcaaac	2580
ggcgctctcg	gtatgattga	gctaaacgat	gattacacac	tgaaaaaagt	gatgaaaccg	2640
ctgattgcat	ctaacacagt	aacagatgaa	attgaacgcg	cgaacgtctt	taaaatgaac	2700
ggcaaatggg	acctgttcac	tgactcccg	ggatcaaaaa	tgacgattga	cggcattacg	2760
tctaacgata	tttacatgct	tggttatgtt	tctaattctt	taactggccc	atacaagccg	2820
ctgaacaaaa	ctggccttgt	gttaaaaaatg	gatcttgatc	ctaacgatgt	aacctttact	2880
tactcacact	tcgctgtacc	tcaagcgaaa	ggaaacaatg	tcgtgattac	aagctatatg	2940
acaaacagag	gattctacgc	agacaaacaa	tcaacgtttg	cgccgagctt	cctgctgaac	3000
atcaaaaggca	agaaaacatc	tgttgtcaaa	gacagcatcc	tgaacaagg	acaattaaca	3060
gttaacaaat	aaaaacgcaa	aagaaaatgc	cgatgggtac	cgagcgaaat	gaccgaccaa	3120
gcgacgcccc	acctgccatc	acgagatttc	gattccaccg	ccgccttcta	tgaaagggtg	3180
ggcttcggaa	tcgttttccg	ggacgccc	gcggacgtgc	tcatagtcca	cgacgcccgt	3240
gattttgtag	ccctggccga	cggccagcag	gtaggcccag	aggctcatgc	cggccgcccgc	3300
cgctttttcc	tcaatcgctc	ttcgttcgtc	tggaaaggcag	tacaccttga	taggtgggct	3360
gcccctctcg	gttggttggt	tttcatcagc	catccgcttg	ccctcatctg	ttacgccggc	3420
ggtagccggc	cagcctcgca	gagcaggatt	cccgttgagc	accgccagggt	gcgaataagg	3480
gacagtgaag	aaggaaacac	cgctcgccgg	tgggcctact	tcacctatcc	tgccccgctg	3540
acgccgttgg	atacaccaag	gaaagtctac	acgaaccctt	tggcaaaatc	ctgtatatcg	3600
tcgcaaaaaag	gatggatata	ccgaaaaaat	cgctataatg	accccgaagc	agggttatgc	3660
agcggaaaaag	cgctgcttcc	ctgctgtttt	gtggaatata	taccgactgg	aaacaggcaa	3720
atgcaggaaa	ttactgaact	gaggggacag	gcgagagacg	atgccaaaga	gctcctgaaa	3780
atctcgataa	ctcaaaaaat	acgcccggta	gtgatcttat	ttcattatgg	tgaaagtggg	3840
aacctcttac	gtgccgatca	acgtctcatt	ttcgccaaaa	gttggcccag	ggcttcccgg	3900
tatcaacagg	gacaccagga	tttatttatt	ctgcgaagtg	atcttccgtc	acaggatttt	3960
attcggcgca	aagtgcgtcg	ggtgatgctg	ccaacttact	gatttagtgt	atgatggtgt	4020
ttttgaggtg	ctccagtggc	ttctgtttct	atcagctcct	gaaaatctcg	ataactcaaa	4080

-continued

aaatacgcgc	ggtagtgtac	ttatttcatt	atggtgaaag	ttggaacctc	ttacgtgccg	4140
atcaacgtct	cattttcgcc	aaaagttggc	ccagggttc	ccggtatcaa	cagggaaccc	4200
aggatttatt	tattctgcga	agtgtacttc	cgtcacaggt	atttattcgg	cgcaaagtgc	4260
gtcgggtgat	gctgccaact	tactgattta	gtgtatgatg	gtgtttttga	ggtgctccag	4320
tggtttctgt	ttctatcagg	gctggatgat	cctccagcgc	ggggatctca	tgctggagtt	4380
cttcgcccac	cccaaaagga	tctagggtgaa	gatccttttt	gataatctca	tgacccaaat	4440
cccttaacgt	gagttttcgt	tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	4500
ttcttgagat	cctttttttc	tgcgcgtaat	ctgctgcttg	caaacaaaaa	aaccacgcgt	4560
accagcgggtg	gtttgtttgc	cggatcaaga	gctaccaact	ctttttccga	aggttaactgg	4620
cttcagcaga	gcgcagatac	caaatactgt	ccttctagt	tagccgtagt	taggccacca	4680
cttcaagaac	tctgtagcac	cgctacata	cctcgctctg	ctaactctgt	taccagtggc	4740
tgctgccagt	ggcgataagt	cgtgtcttac	cgggttgga	tcaagacgat	agttaccgga	4800
taaggcgag	cggtcgggct	gaacgggggg	ttcgtgcaca	cagcccagct	tgagcggaac	4860
gaacctaccc	gaactgagat	acctacagcg	tgagcattga	gaaagcgcca	cgcttccga	4920
aggagaaaa	gcgcagaggt	atccggtgaa	cggcagggtc	ggaacaggag	agcgacagag	4980
ggagcttcca	gggggaaacg	cctggtatct	ttatagtcct	gtcgggtttc	gccacctctg	5040
acttgagcgt	cgatttttgt	gatgctcgtc	aggggggcgg	agcctatgga	aaaacgccag	5100
caacgcggcc	tttttacggt	tcttgccctt	ttgctggcct	tttgcacaca	tgttctttcc	5160
tgcgttatcc	cctgattctg	tggataaccg	tattaccgcc	tttgagttag	ctgataaccg	5220
tcgcccagc	cgaacgaccg	agcgacagca	gtcagttagc	gaggaagcgg	aagagcgccc	5280
aatacgcaaa	ccgcctctcc	ccgcgcgttg	gccgattcat	taatgcagct	ggcacgacag	5340
gtttcccgac	tggaaagcgg	gcagttagcg	caacgcaatt	aatgtgagtt	agctcactca	5400
ttaggcacc	caggctttac	actttatgct	tccggtcgt	atgttggtg	gaattgtgag	5460
cggataacaa	tttcacacag	gaaacagcta	tgacctgat	tacgccaaag	ttgcatgcct	5520
gcaggtgcac	tctagaggat	ccccatttgc	tgaagggtgt	actctgcctg	gccccattcc	5580
tgccggcgaa	gaagtgaaaa	accctgaacc	ttttcagaag	taactaaggc	cgcaatccct	5640
cgattgtctg	atcaacgacg	gcgtctgtga	gtctagctag	agatctagat	tccaggcgcc	5700
atcgttgcca	atacatcggt	gtgtcaatgg	gtatctcacc	gaggaggatc	acttctctctg	5760
cttttagcat	gggagcagct	tgggtttcgg	gaagaagtcc	ccaaccaagg	cgcgtttgtg	5820
tttatcgcg	gcgtcgcg	gcaatacggc	gacaccggac	ggtggatttt	cgccgctggc	5880
gcgttcgcgg	caagcctgat	ctgggtcccc	ctgggtgggt	tcggcgagc	agcattgtca	5940
cgcccgctgt	ccagccccaa	ggtgtggcgc	tggatcaacg	tcgtcgtggc	agttgtgatg	6000
accgcattgg	ccatcaaaact	gatgttgatg	ggttagtttt	cgcggttttt	ggaatcggtg	6060
gccttcgccc	aaatgttgat	gccggcgctg	tgggaaatct	catcgatcgc	ctccaaactcg	6120
gcgtcagaaa	actccaagtt	gttgagtga	tcaaggctgt	tgtccagctg	ctcaactgac	6180
gaagcaccaa	tcaatgcact	ggtcacggta	tccgcgcctg	actctccttg	ctcgcgagc	6240
acccatgcaa	gcgccatctg	cgcaagtga	tgcccggtt	cctggcgcat	gtcattgagc	6300
ttgcggacca	tatcaatatt	gttcacgttc	aacatgcctt	cagacaggga	cttaccctgg	6360

-continued

```

ctggcgcggg aacctctctgg aattcactgg ccgtcgtttt acaacgtcgt gactgggaaa 6420
acctggcggt tacccaactt aatcgccctg cagcacatcc ccccttcgcc agctggcgta 6480
atagcgaaga ggcccgccacc gatcgccctt cccaacagtt gcgcagcctg aatggcgaat 6540
ggcg 6544

```

```

<210> SEQ ID NO 2
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(44)
<223> OTHER INFORMATION: DargR_lf
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer DargR_lf
      (genomic DNA of ATCC13032 as template)

```

```

<400> SEQUENCE: 2

```

```

tcgagctcgg taccggggga tcctctggct gatgatgatc tcac 44

```

```

<210> SEQ ID NO 3
<211> LENGTH: 45
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(45)
<223> OTHER INFORMATION: DargR_lr
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer DargR_lr
      (genomic DNA of ATCC13032 as template)

```

```

<400> SEQUENCE: 3

```

```

ctagggatag tagacacttc acaagtgtct tacctcggtt gggttg 45

```

```

<210> SEQ ID NO 4
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(47)
<223> OTHER INFORMATION: DargR_rf
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the plasmid DargR_rf
      (genomic DNA of ATCC13032 as template)

```

```

<400> SEQUENCE: 4

```

```

cttgtgaagt gtctactatc cctagaatta ctcagcgggc gcaccac 47

```

```

<210> SEQ ID NO 5
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(46)
<223> OTHER INFORMATION: DargR_rr
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer DargR_rr
      (genomic DNA of ATCC13032 as template)

```

```

<400> SEQUENCE: 5

```

```

agcttgcatg cctgcaggtc gactctgcac gacggttcag ctcttc 46

```

-continued

<210> SEQ ID NO 6
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(46)
<223> OTHER INFORMATION: PsodcarAB-LA-F
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer
PsodcarAB-LA-F (genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 6

ggaaacagct atgacatgat tacgcgggta tcgcggaatc cgtatg 46

<210> SEQ ID NO 7
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(25)
<223> OTHER INFORMATION: PsodcarAB-LA-R
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer
PsodcarAB-LA-R (genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 7

ttaagcgttt tgtgcaactc cgtct 25

<210> SEQ ID NO 8
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(50)
<223> OTHER INFORMATION: PsodcarAB-F
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer PsodcarAB-F
(genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 8

agacggagtt gcacaaaacg cttaaacctt acttagctgc caattattcc 50

<210> SEQ ID NO 9
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(50)
<223> OTHER INFORMATION: PsodcarAB-R
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer PsodcarAB-R
(genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 9

ggtaggtggt ggtgtcttta ctcatgggta aaaaatcctt tcgtaggttt 50

<210> SEQ ID NO 10
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature

-continued

<222> LOCATION: (1) .. (25)
<223> OTHER INFORMATION: PsodcarAB-RA-F
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer
PsodcarAB-RA-F (genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 10

atgagtaaag acaccaccac ctacc 25

<210> SEQ ID NO 11
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1) .. (46)
<223> OTHER INFORMATION: PsodcarAB-RA-R
<220> FEATURE:
<223> OTHER INFORMATION: DNA sequence coding for the primer
PsodcarAB-RA-R (genomic DNA of ATCC13032 as template)

<400> SEQUENCE: 11

gttgtaaaac gacggccagt gccaccggtg atgtggttct tcaactg 46

<210> SEQ ID NO 12
<211> LENGTH: 1146
<212> TYPE: DNA
<213> ORGANISM: Moorea producens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1) .. (1146)
<223> OTHER INFORMATION: L-arginine:glycine amidinotransferase

<400> SEQUENCE: 12

atgtcggaaa aaattgttaa ttcctggaat gaatgggatg aattggaaga aatggtggta 60
ggaattgcag actatgctag ctttgaacca aaagaaccag ggaatcatcc gaaattaaga 120
aatcaaaatt tagcggaaat cattccttc cccagtggac ctaaagacc taaagtcctt 180
gaaaaagcta atgaagaatt aaatggactg gcttatttat taaaagacca cgatgtgata 240
gtaagaagac ccgaaaaaat tgattttact aaatctctaa aaacacctta ctttgaagtt 300
gcaaatacat actgtggagt ctgtcctcgg gatgtcatga ttacctttgg gaatgaaatc 360
atggaagcga ctatgtcgaa gagagctaga ttttttgaat acttacctta ccggaaattg 420
gtctatgaat attggaataa agacgagcat atgatttgga atgtgctgcc taaaccgact 480
atgcaggata gtatgtatct agagaatttc tgggagctgt ctttagaaga acgattttaag 540
cgtatgcatg attttgaatt ttgtattaca caagatgaag taatttttga tgcggctgac 600
tgtagcagat taggaaaagga tatattagtt caggaatcga tgacaacaaa tagaacagga 660
attcgggtggt taaaaaagca cctagaacca agaggatttc gggttcacc tgttcatttt 720
ccccttgatt tttcccctc acacattgac tgtacgtttg ttcctttgcg accaggtcct 780
attttgacaa accctgaaag acctatacgg gaagaggagg agaagatttt taaagagaat 840
ggctgggagt tgatcacagt tcctcaaccg acttgctcga atgatgaaat gccaatgttt 900
tgccagtcca gtaagtgggt gtcaatgaat gttctgagta tatcacgac aaagggttatc 960
tgtgaggaaa gagaaaaacc tctccaagaa ttgttgagata agcatggatt tgaggttttt 1020
cctttacct ttagacatgt ctttgaattt ggggggtctt ttcattgtgc aacttgggat 1080
attcgccgaa aaggtgagtg tgaagattat ttaccaaatt taaactatca accgatttgt 1140

-continued

gggtaa

1146

<210> SEQ ID NO 13
 <211> LENGTH: 381
 <212> TYPE: PRT
 <213> ORGANISM: Moorea producens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(381)
 <223> OTHER INFORMATION: L-arginine:glycine amidinotransferase

<400> SEQUENCE: 13

Met Ser Glu Lys Ile Val Asn Ser Trp Asn Glu Trp Asp Glu Leu Glu
 1 5 10 15
 Glu Met Val Val Gly Ile Ala Asp Tyr Ala Ser Phe Glu Pro Lys Glu
 20 25 30
 Pro Gly Asn His Pro Lys Leu Arg Asn Gln Asn Leu Ala Glu Ile Ile
 35 40 45
 Pro Phe Pro Ser Gly Pro Lys Asp Pro Lys Val Leu Glu Lys Ala Asn
 50 55 60
 Glu Glu Leu Asn Gly Leu Ala Tyr Leu Leu Lys Asp His Asp Val Ile
 65 70 75 80
 Val Arg Arg Pro Glu Lys Ile Asp Phe Thr Lys Ser Leu Lys Thr Pro
 85 90 95
 Tyr Phe Glu Val Ala Asn Gln Tyr Cys Gly Val Cys Pro Arg Asp Val
 100 105 110
 Met Ile Thr Phe Gly Asn Glu Ile Met Glu Ala Thr Met Ser Lys Arg
 115 120 125
 Ala Arg Phe Phe Glu Tyr Leu Pro Tyr Arg Lys Leu Val Tyr Glu Tyr
 130 135 140
 Trp Asn Lys Asp Glu His Met Ile Trp Asn Ala Ala Pro Lys Pro Thr
 145 150 155 160
 Met Gln Asp Ser Met Tyr Leu Glu Asn Phe Trp Glu Leu Ser Leu Glu
 165 170 175
 Glu Arg Phe Lys Arg Met His Asp Phe Glu Phe Cys Ile Thr Gln Asp
 180 185 190
 Glu Val Ile Phe Asp Ala Ala Asp Cys Ser Arg Leu Gly Lys Asp Ile
 195 200 205
 Leu Val Gln Glu Ser Met Thr Thr Asn Arg Thr Gly Ile Arg Trp Leu
 210 215 220
 Lys Lys His Leu Glu Pro Arg Gly Phe Arg Val His Pro Val His Phe
 225 230 235 240
 Pro Leu Asp Phe Phe Pro Ser His Ile Asp Cys Thr Phe Val Pro Leu
 245 250 255
 Arg Pro Gly Leu Ile Leu Thr Asn Pro Glu Arg Pro Ile Arg Glu Glu
 260 265 270
 Glu Glu Lys Ile Phe Lys Glu Asn Gly Trp Glu Leu Ile Thr Val Pro
 275 280 285
 Gln Pro Thr Cys Ser Asn Asp Glu Met Pro Met Phe Cys Gln Ser Ser
 290 295 300
 Lys Trp Leu Ser Met Asn Val Leu Ser Ile Ser Pro Thr Lys Val Ile
 305 310 315 320
 Cys Glu Glu Arg Glu Lys Pro Leu Gln Glu Leu Leu Asp Lys His Gly

-continued

	325		330		335	
Phe	Glu	Val	Phe	Pro	Leu	Pro
	340					
			Phe	Arg	His	Val
			345			
					Phe	Glu
					350	
					Phe	Gly
						Gly
Ser	Phe	His	Cys	Ala	Thr	Trp
	355					
			Asp	Ile	Arg	Arg
			360			
					Lys	Gly
					365	
					Glu	Cys
						Glu
Asp	Tyr	Leu	Pro	Asn	Leu	Asn
	370					
					Tyr	Gln
					375	
					Pro	Ile
					Cys	Gly
					380	

<210> SEQ ID NO 14
 <211> LENGTH: 1146
 <212> TYPE: DNA
 <213> ORGANISM: Moorea producens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1) .. (1146)
 <223> OTHER INFORMATION: L-arginine:glycine amidinotransferase codon
 optimized for C. glutamicum

<400> SEQUENCE: 14

atgtcagaaa agattgttaa ctcttggaaac gaatgggacg aacttgaaga aatggtggtt	60
ggcatcgcgg actacgcttc cttcgagcca aaagaaccag gtaaccaccc taagctacgc	120
aaaccagaacc tggccgagat catcccatc ccatccggcc caaaggaccc aaaggttctc	180
gaaaaagcaa acgaagagct gaacggcctc gcatacctcc taaaggacca cgatgttatc	240
gtccgcccgc cagaaaaaat tgatttcaact aagtcacctt aaaccccata cttcgaggtc	300
gctaaccagt actgcccgtt ttgccacgc gacgttatga ttaccttcgg taatgagatt	360
atggaggcta ccatgagcaa gcgtgcacgc ttcttcgagt acttgccata ccgtaagctc	420
gtctacgagt actggaacaa ggatgagcat atgatctgga acgctgctcc aaagcctacg	480
atgcaggact ccatgtactt ggagaacttc tgggagctgt ccctcgagga gcgcttcaag	540
cgcattgcacg acttcgaatt ttgcattacc caggacgagg ttattcttca cgcagctgac	600
tgctcccgcc tgggcaagga catcctggtc caggagtcca tgaccaccaa ccgcaccggc	660
attcgttggt tgaagaagca cctcgaaact cgcggtttcc gcgttcaccc agtcacctc	720
cctctcgact tcttcccttc tcacattgac tgcactttcg tccctctccg tcccgccctc	780
atccttacca acccagagcg cccaatccgc gaagaagaag aaaagatctt caaggagaac	840
ggctgggagc ttatcacctg ccacaaacct acctgctcca acgacgaaat gcctatgttc	900
tgccagtcct cgaagtggct cagcatgaac gtcccttcta tctccccaac aaaggttatt	960
tgccaagagc gtgaaaaagc actacaggaa ctctgggaca agcacggctt cgaagtcttc	1020
ccactcccat tccgccacgt ttctgaattc ggccggtcct ttcactgcgc caattgggac	1080
atccgccgca agggcgcaatg cgaagactac ctcccaaacc tgaactacca gcctatctgc	1140
ggctaa	1146

<210> SEQ ID NO 15
 <211> LENGTH: 1264
 <212> TYPE: DNA
 <213> ORGANISM: Moorea producens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1) .. (1264)
 <223> OTHER INFORMATION: L-arginine:glycine amidinotransferase segment

<400> SEQUENCE: 15

-continued

```

agcgctccgtc tectctcgga acgaatggga cgaacttgaa gaaatggtgg ttggcatcgc      60
ggactacgct tccttcgagc caaaagaacc aggtaaccac cctaagctac gcaaccagaa      120
ctgggccgag atcatcccat tccatcccg cccaaaggac ccaaaggttc tcgaaaaagc      180
aaacgaagag ctgaacggcc tcgcatacct cctaaaggac cagcatgtta tcgtccgccg      240
cccagaaaaa attgatttca ctaagtcct taaaaccca tacttcgagg tcgctaacca      300
gtactgcggc gtttgcacac gcgacgttat gattaccttc ggtaatgaga ttatggaggc      360
taccatgagc aagcgtgcac gcttcttcga gtacttgcca taccgtaagc tcgtctacga      420
gtactggaac aaggatgagc atatgatctg gaacgctgct ccaaagccta cgatgcagga      480
ctccatgtac ttggagaact tctgggagct gtccctcgag gagcgcttca agcgcatgca      540
cgacttcgaa ttttgatta cccaggacga ggttatcttc gacgcagctg actgctcccg      600
cctgggcaag gacatctcgg tccaggagtc catgaccacc aaccgcaccg gcattcgttg      660
gttgaagaag cacctegaac ctgcgggttt ccgcgttcac ccagtcactc tccctctcga      720
cttcttcctt tctcacattg actgcacttt cgtccctctc cgtcccgcc tcactcttac      780
caaccagag cgcccaatcc gcgaagaaga agaaaagatc ttcaaggaga acggctggga      840
gcttatcacc gtccacaac ctacctgctc caacgacgaa atgcctatgt tctgccagtc      900
ctcgaagtgg ctcagcatga acgtcctttc tatctcccca acaaaggtta ttgcgaaga      960
gcgtgaaaag ccactacagg aactcctgga caagcacggc ttcgaagtct tccactccc     1020
attccgccac gttttcgaat tcggcggtc ctttactgc gccacttggg acatccgccg     1080
caagggcgaa tgcaagact acctccaaa cctgaactac cagcctatct gcggctaata     1140
aggcgcgctt tttgaatcct ccaaacctgt ggatagaaca atctatccac aggttttcag     1200
cctgcaggca tgcaagcttg gctgttttg cggtagagag aagattttca gcgagacgat     1260
catg                                             1264

```

```

<210> SEQ ID NO 16
<211> LENGTH: 351
<212> TYPE: DNA
<213> ORGANISM: Moorea producens
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1) .. (351)
<223> OTHER INFORMATION: L-arginine:glycine amidinotransferase promoter
        containing segment

```

```

<400> SEQUENCE: 16
ccgagtccaa tgtgtccgtg ttcacctcag cggccgcacc ggtcggggta ccttgacaaa      60
tactttccaa tgggggtacaa tggcgagtg ccatccactc cgagggtgat accagcctcc     120
tcggtggaat gatcatccgc gtcggagacg aagtaattga cggcagcacc tcgggcaaac     180
tcgagcgtct gcgggcaagc ttcgcataaa gacacgacga attagacaac attagtaatg     240
ctggaagaaa caaccgagag caggaagaac atgtcagaaa agattgttaa ctctggaac     300
gaatgggacg aactgaaga aatggtggtt ggcacgcgg actacgcttc c                 351

```

```

<210> SEQ ID NO 17
<211> LENGTH: 6903
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature

```

-continued

<222> LOCATION: (1) .. (6903)
<223> OTHER INFORMATION: pLIB_pBL1
<220> FEATURE:
<223> OTHER INFORMATION: E. coli-C. glutamicum shuttle plasmid pLIB_pBL
1 has the replication origin from pBL 1, the pSC101 replication
origin, a kanamycin resistance gene and the BioBricks Terminator
BBa_B1006 downstream of a *NatI* restriction site

<400> SEQUENCE: 17

gacagtaaga cgggtaagcc tgttgatgat accgctgcct tactgggtgc attagccagt 60
ctgaatgacc tgtcacggga taatccgaag tggtcagact ggaaaatcag agggcaggaa 120
ctgcagaaca gcaaaaagtc agatagcacc acatagcaga cccgccataa aacgcctga 180
gaagcccgctg acgggctttt cttgtattat gggtagtttc cttgcatgaa tccataaaag 240
gcgcctgtag tgccatttac cccattcac tgccagagcc gtgagcgag cgaactgaat 300
gtcacgaaaa agacagcgac tcagggtgcct gatggtcgga gacaaaagga atattcagcg 360
at ttgcccga gcttgcgagg gtgctactta agcctttagg gttttaaggt ctgttttgta 420
gaggagcaaa cagcgtttgc gacatccttt tgtaatactg cggaactgac taaagtagtg 480
agttatacac agggctggga tctattcttt ttattctttt ttattctttc ttattcttat 540
aaattataac cacttgaata taaacaaaaa aaacacacaa aggtctagcg gaatttacag 600
agggcttagc agaatttaca agttttccag caaaggctta gcagaattta cagataccca 660
caactcaaag gaaaaggact agtaattatc attgactagc ccatctcaat tggatatagt 720
attaaaaatc cctagaccaa ttgagatgta tgtctgaatt agttgttttc aaagcaatg 780
aactagcgat tagtcgctat gacttaacgg agcatgaaac caagctaatt ttatgctgtg 840
tggcactact caaccccacg attgaaaacc ctacaaggaa agaacggacg gtatcgttca 900
cttataacca atacgctcag atgatgaaca tcagtaggga aaatgcttat ggtgtattag 960
ctaaagcaac cagagagctg atgacgagaa ctgtggaaat caggaatcct ttggttaaag 1020
gctttgagat tttccagtg acaaaactatg ccaagttctc aagcgaaaaa ttagaattag 1080
tttttagtga agagatatgt ccttatcttt tccagttaaa aaaattcata aaatataatc 1140
tggaacatgt taagtctttt gaaaacaaat actctatgag gatttatgag tggttattaa 1200
aagaactaac acaaaagaaa actcacaagg caaatataga gattagcctt gatgaattta 1260
agttcatgtt aatgcttgaa aataactacc atgagtttaa aaggcttaac caatggggtt 1320
tgaaaccaat aagtaaagat ttaaacactt acagcaatat gaaattgggtg gttgataagc 1380
gaggccgccc gactgatacg ttgattttcc aagttgaact agatagacaa atggatctcg 1440
taaccgaact tgagaacaac cagataaaaa tgaatggtga caaaatacca acaaccatta 1500
catcagatc ctacctacat aacggactaa gaaaaaact acacgatgct ttaactgcaa 1560
aaattcagct caccagtttt gagggcaaat ttttgagtga catgcaaagt aagcatgatc 1620
tcaatgggtc gttctcatgg ctacgcaaaa aacaacgaac cacactagag aacatactgg 1680
ctaaatacgg aaggatctga ggttcttatg gctcttgat ctatcagtga agcatcaaga 1740
ctaacaacaa aaagtagaac aactgttcac cggtacatat caaagggaaa actgtccata 1800
tgccacagatg aaaaagggtg aaaaaagata gatacatcag agcttttacg agtttttggt 1860
gcattcaaag ctgttcacca tgaacagatc gacaatgtaa cagatgaaca gcatgtaaca 1920
cctaatagaa cagggtgaaac cagtaaaaca aagcaactag aacatgaaat tgaacacctg 1980

-continued

agacaacttg ttacagctca acagtcacac atagacagcc tgaaacaggc gatgctgctt	2040
atcgaatcaa agctgccgac aacacgggag ccagtgcgc ctcccgtagg gaaaaaatca	2100
tggcaattct ggaagaaata gcgccattcg ccattcaggc tgectgcagg gaaagccacg	2160
ttgtgtctca aaatctctga tgttacattg cacaagataa aaatatatca tcatgaacaa	2220
taaaactgtc tgcttacata aacagtaata caaggggtgt tatgagccat attcaacggg	2280
aaacgtcttg ctcgaggcgc cgattaaatt ccaacatgga tgctgattta tatgggtata	2340
aatgggctcg cgataatgtc gggcaatcag gtgcgacaat ctatcgattg tatgggaagc	2400
ccgatgcgcc agagtgtgtt ctgaaacatg gcaaaggtag cgttgccaat gatgttacag	2460
atgagatggt cagactaaac tggtgcagcg aatttatgcc tcttcgcacc atcaagcatt	2520
ttatccgtac tctgatgat gcattggttac tcaccactgc gatccccggg aaaacagcat	2580
tccaggtatt agaagaatat cctgattcag gtgaaaatat tggtgatgcg ctggcagtgt	2640
tctgcgcgcg gttgcattcg attcctgttt gtaattgtcc ttttaacage gatcgcgat	2700
ttcgtctcgc tcaggcgcaa tcacgaatga ataacgggtt gggtgatgcg agtgattttg	2760
atgacgagcg taatggctgg cctgttgaac aagtctgga agaaatgcat aagcttttgc	2820
cattctcacc ggattcagtc gtcactcatg gtgatttctc acttgataac cttatttttg	2880
acgaggggaa attaataggt tgtattgatg ttggacgagt cggaatcgca gaccgatacc	2940
aggatcttgc catcctatgg aactgcctcg gtgagtttct tccttcatta cagaaacggc	3000
tttttcaaaa atatggtatt gataatcctg atatgaataa attgcagttt catttgatgc	3060
tcgatgagtt tttctaata gaattgggta attggttgta aactggcag agcattacgc	3120
tgacttgacg ggacgggggc tttgttgaat aaatcgaact tttgctgagt tgaaggatca	3180
gatcacgcat cttcccgaca acgcagaccg ttccgtggca aagcaaaagt tcaaatcac	3240
caactggtcc acctacaaca aagctctcat caaccgtggc tcctcactt tctggctgga	3300
tgatgggctc gcaggcctca gcgcgattca ggcctgggat gagtcagcaa caccctcttc	3360
acgagggcaga tctcagcgcc cccccccccc tagcttgtct acgtctgatg ctttgaatcg	3420
gacggacttg ccgatcttgt atcggtgat ttttcctcg tttgccact ttttaatggt	3480
ggccgggggtg agagctacgc gggcggcgac ctgctgcgct gtgatccaat attcggggtc	3540
gttcaactggt tcccccttct gatttctggc atagaagaac ccccgtaga tgtgtgggtc	3600
cgggggttgc tgatttttgc gagacttctc gcgcaattcc cttagctagg tgaaaacacc	3660
atgaaacact agggaaaacac ccataaaca ccattaggg cagtagggcg gcttcttctg	3720
ctagggcttg catttgggcg gtgatctggt ctttagcgtg tgaaagtgtg tcgtagggtg	3780
cgtgctcaat gcaactgaac gtcacgtcat ttaccgggtc acggtgggca aagagaacta	3840
gtgggttaga cattgttttc ctgctgtgct gtggtggtga gcttttctag ccgctcggt	3900
aacgcggcga tcataaactc ttggagggtt tcaccgttct gcatgcctgc gcgcttcag	3960
tctcacgta gtgcaaaagg aacgcgtgcg gtgaccacga cgggcttagc ctttgccctgc	4020
gcttctagtg cttcgatggt ggcttgtgcc tgcgcttgct gcgcctgtag tgcctgttga	4080
gcttcttgta gttgctgttc tagctgtgcc ttggttgcca tgctttaaga ctctagtagc	4140
tttctgcga tatgtcatgc gcatgcgtag caaacattgt cctgcaactc attcattatg	4200
tgcaagtctc ctgttactag tcgtacatac tcataattac ctagtctgca tgcagtgcat	4260

-continued

gcacatgcag	tcatgtcgtg	ctaattgtga	aaacatgtac	atgcagattg	ctgggggtgc	4320
agggggcgga	gccacctgt	ccatgcgggg	tgtggggtt	gccccgccg	tacagacagt	4380
gagcaccggg	gcacctagtc	gcggatcccc	cccctaggt	tcggacacgt	aacctccca	4440
tgtcgatgca	aatctttaac	attgagtacg	ggttaagctg	cacgcatagc	caagctaggc	4500
ggccacaaaa	caccactaaa	aattaatagt	tcttagacaa	gacaaacccc	cgtgcgagct	4560
accaactcat	atgcacgggg	gccacataac	ccgaagggtt	ttcaattgac	aaccatagca	4620
ctagctaaga	caacgggcac	aacacccgca	caaactcgca	ctgcgcaacc	ccgcacaaca	4680
tcgggtctag	gtaacactga	aatagaagt	aacacctcta	aggaaccgca	ggtcaatgag	4740
ggttctaagg	tactcgcgc	tagggcggtg	cgtaggcaaa	acgtcatgta	caagatcacc	4800
aatagtaagg	ctctggcggg	gtgccatagg	tggcgacggg	acgaagctgt	tgcggtgtcc	4860
tggctgtcta	acggtgtctc	gcagtttgag	ggtctgcaaa	actctcactc	tcgctggggg	4920
tcacctctgg	ctgaattgga	agtcacgggc	gaacgccgca	ttgagctggc	tattgctact	4980
aagaatcact	tggcgggcg	tggcgcgctc	atgatgtttg	tgggcaactgt	tcgacacaac	5040
cgtctacagt	catttgcgca	ggttgaagcg	ggtattaaga	ctgcgtactc	ttcgatgggtg	5100
aaaacatctc	agtggaagaa	agaacgtgca	cggtagcggg	tggagcacac	ctatagtga	5160
tatgaggtca	cagactcttg	ggcgaaacgt	tggcaactgc	accgcaacat	gctgtgttcc	5220
ttggatcgct	cactgtctga	cgatgaactc	aaggcggttg	aggattccat	gttttcccgc	5280
tggctcgtcg	gtgtggttaa	ggcgggtatg	gacgcgccac	tgcgtgagca	cgggggtcaaa	5340
cttgatcagg	tgtctacctg	gggtggagac	gctgcgaaaa	tggcaacct	cctcgctaag	5400
ggcatgtctc	aggaactgac	tggctccgct	actaaaaccc	cgtctaaggg	gtcgtacacg	5460
cgtttcaga	tgttggatat	gttggccgat	caaagcgcg	ccggcgagga	tatggacgct	5520
gttttggtgg	ctcgggtggc	tgagtatgag	gttggttcta	aaaacctgcg	ttcgtcctgg	5580
tcacgtgggg	ctaagcgtgc	tttgggcatt	gattacatag	acgctgatgt	acgtcgtgaa	5640
atggaagaag	aactgtacaa	gctcgccggt	ctggaagcac	cggaaacgggt	cgaatcaacc	5700
cgcgttgctg	ttgctttggt	gaagcccgat	gattggaaac	tgattcagtc	tgatttcgcg	5760
gttaggcagt	acgttctaga	ttgcgtggat	aaggctaagg	acgtggccgc	tcgcgaacgt	5820
gtcgctaagt	agggtctggc	aagtctgggt	gtggattcca	ccccgtgcac	gatcgttatg	5880
gatgatgtgg	acttggacgc	ggtctgcct	actcatgggg	acgctactaa	gcgtgatctg	5940
aatgcggcgg	tgttcgcggg	taatgagcag	actattcttc	gcaccacta	aaagcggcat	6000
aaaccccggt	cgatattttg	tgcgatgaat	ttatgggtcaa	tgtcgcgggg	gcaaactatg	6060
atgggtcttg	ttgttgacaa	tggtgatatt	catcaggaat	ggaactgtca	tgctgttatg	6120
tgcttggtgc	ctaatacaag	ctggggacaa	tgggttgccc	cgttgatctg	atctagtctg	6180
gattggcggg	gcttcactgt	atctgggggt	ggcatcgtga	atagattgca	caccgtagt	6240
ggcagtgtgc	acaccatagt	ggccatgagc	accaccaccc	ccagggaacg	cgaaggcgcg	6300
aagctctcgc	cctggtgcgg	ctcggagatc	aagcaatccg	gcgtcgcccg	gagccgggac	6360
tactgccgcc	gctcctgccg	ccagcggggc	tacgagggcc	ggcgccagcg	cgaggcgatc	6420
gtgtccgcgc	tggcgctggc	agtcgctcgc	cgagatacgt	cacgtgacga	aatgcagcag	6480
ccttccattc	cgtcacgtga	cgaactcgg	gccgcaggtc	agagcacggg	tcgcgccgct	6540

-continued

ccggccctgc cggacccccg gcatcccgca agaggcccg cagtaccggc ataaccaagc	6600
ctatgcctac agcatccagg gtgacgggtgc cgaggatgac gatgagcgca ttgttagatt	6660
tcatacacgg tgccctgactg cgtttagcaat ttaactgtga taaactaccg cattaaagct	6720
tatccgagtc caatgtgtcc gtgttcacct cagcggccgc gttaacttaa ggtaccgaat	6780
tctaagcttc acaaaaaaaaa accccgcccc tgacaggcgcg gggttttttt tactcagcat	6840
gtggatggcc agtggtcgga ccgagcgctt taattaatta gcatggtgac acaagcacag	6900
tat	6903

<210> SEQ ID NO 18
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(20)
 <223> OTHER INFORMATION: AGAT_f
 <220> FEATURE:
 <223> OTHER INFORMATION: primer AGAT_f

<400> SEQUENCE: 18

tggttgcat cgcgactac	20
---------------------	----

<210> SEQ ID NO 19
 <211> LENGTH: 57
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(57)
 <223> OTHER INFORMATION: AGAT_r
 <220> FEATURE:
 <223> OTHER INFORMATION: primer AGAT_r

<400> SEQUENCE: 19

gcttagaatt cggtacctta agttaacgcy gccgcttatt agccgcagat aggctgg	57
--	----

<210> SEQ ID NO 20
 <211> LENGTH: 6026
 <212> TYPE: DNA
 <213> ORGANISM: Corynebacterium glutamicum
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(6026)
 <223> OTHER INFORMATION: Pg3-argFGH

<400> SEQUENCE: 20

ttccggctcg tatgttgtgt ggaattgtga gcggataaca atttcacaca ggaacacgct	60
atgaccatga ttacgaattg cgtggcaatc aggctttacc tcagcgaaga attcgacgac	120
gccacccgcy tctacaccgt ccgatccttt gatgccgcaa ctgaaagcat cgtggtggat	180
gtggttcaac accaccacga aagccccatg atgcgctggt cagacaccgt aaaaaatcaac	240
gacaccctcg tgctcacagg accccgcccc cactttgtca tccccgaagg cgaacaagca	300
gcactcttcc ttgatgacac cgccatcccc gctctcgccg ctattttggga tcaatggcca	360
acagatcttc gtggcaaagg atgggtgtgc actgacgac cgcagcctt cgatgaacta	420
cccagcatcg acggactgga actgaacctg ctgcgcgcy gatcagatcc aactgttcag	480
ccacttgccc aacaggcata tgacctggaa aaccagaaa cttacgtggt gtgggcagcc	540

-continued

ggcgagcgag	atgaaataaa	atccatccgc	aggcacttcc	gcaagcaggt	gggattggaa	600
aaagatgcag	tggccgtggt	tgggtactgg	aaatacaaca	ccaccaacac	tcagatcgat	660
gcagtccgca	aagaaaaacta	catgaagatg	ctctctgaag	ggctacagct	ggaaaaacttc	720
gacgacctct	cattggagat	ttaaggggtc	gagtttttag	aatgggtggt	tgcattgatt	780
ggcgtctgat	gagctcttag	aagcgattct	gggagggcat	gatttttggg	attcggagct	840
gaatgtggac	tgattttggg	ggaaccagac	tggcgtgccc	aaatgtcgtg	tcttcagtcc	900
taggtcaata	ttacaggcca	taaaactaatg	cttatcgacg	tacccccgca	cgtgcccact	960
ccgcgtgagc	ggtagcttgcg	gcacgcacac	gtgcgcgttg	ctccgcaacc	cgcacgcccg	1020
cggttccacc	gcgcgttgca	cgggaagcca	ctgcaccatc	aatggtgagc	acttcccgtta	1080
cctctggggg	cagacgtgca	tcaacgccac	tgagttcttc	atcagtgaga	tcgataaggt	1140
ccactcccct	ggactccgcg	atccgcacgc	aagcgccgga	tgcttcgtgt	gcctcacgga	1200
atggaacgcc	ctggcgcacc	atccactcag	ccaagtcggt	ggcaaggggtg	aaacctgctg	1260
gtgcaagttc	acgcatgcgc	tcggtgttga	aggtaaggt	ggaaaccaa	ccagtcattg	1320
cagggagcag	caggttgagc	tgccgcacgc	aatctacgat	tggttcctta	tcttcctgca	1380
ggtcgcggtt	gtacgctaaa	ggctgtgcct	tcagggtagc	cagcagaccg	gtgaggttac	1440
cgatcaagcg	accagacttg	ccacgggtca	gctctgccac	gtcagggttc	ttcttctgcg	1500
gcatgattga	gctgcctgtg	gaccaggaat	cagacaaggt	gatgtaacca	aattctgggg	1560
tgcccatgc	gatgatttct	tcagccaagc	gggacatata	cactgcaagc	tgccgcagca	1620
cgaaggcggg	ttcagatgcg	aaatcgccgg	agctgggtggc	atcaatggag	ttatctgctg	1680
cgggaatcaaa	gccgagttct	tcagcgattg	cttcagggtt	gagcttcaaa	gaggaaaccag	1740
caagtgcgcc	ggaaccgtaa	ggagacaccg	caagacgctt	gtccaggtca	cggatacgat	1800
caatatcgcg	cagcaaaggc	tgtgcgtgtg	ccagcagctg	gtgtgccaga	aggaccggct	1860
gagctgcctg	gaagtgggtc	ttgcctggca	tgatcgccgc	tgcatgtgcc	ttagcttggg	1920
cgtcaggggc	gtcgacaagc	tcgggtgttc	ccagcgcgat	gtcgcgcact	gcgtcgcgga	1980
cccacatgcg	gaacagggtt	gccacctggt	cgttcgggga	acgaccagcg	cgcagacggc	2040
cggccaccto	aggaccaacg	cgttcaatca	gaccgcgttc	catcgccgcc	tgacatacct	2100
catcagaagg	cagcggaccg	aagggtccgt	cggcgacatc	cttgcccagc	tgatcaagcc	2160
cagccagcat	ggtggctaga	tcttcatcag	aaagtagatc	tgcttggtgc	aaaaccttgg	2220
cgtgtgcctt	ggaggccaac	acatcataag	gggccaaaac	ccagtcgaaa	tgagtggaga	2280
cactcaaggc	gaacatggcc	tcggaggggtc	caccggagaa	gcggccgccc	cacagcgcac	2340
cttcattggt	tccgtgctgt	tccatgtggt	gttcttctac	tttctaata	cttgaagttc	2400
tagtctggat	gcttgaagg	gtggcttagt	tggtgccagc	ttcgcgatcg	cgcttgtag	2460
cgatcttgga	ggacagaccg	tgacgtgga	caaagccctt	agccagggtc	tggtcgaagg	2520
tgctgcgggt	gtcgtaggtg	gccaggttga	agtcgtacag	ggagtggctg	gaacgacgac	2580
cattgatggt	gatggaacct	gcgtgcagaa	ccatgcggat	atcgccgggtg	acgtgctcct	2640
gggtggaatc	aatgaacgcg	tccagggagc	gcttcagagg	tccgaaccac	aggccgtcgt	2700
atacttccto	agcccaacgt	gcgtcaacgc	cgcgcttgta	gcgagccagt	tcgcgctcga	2760
tggtgacatc	ttccaaagcc	tcgtgagcct	taatcagtcg	gattgcgcct	gggtgcttcgt	2820

-continued

agatttcgcg	ggacttgatg	cccacgagac	ggctctcaac	catgtcaagg	cggccaacgc	2880
cctgtgcgc	tgcacgacgg	ttcagctctt	caatagcctg	cagtacggag	actggacggc	2940
catcgatgga	gactggcttg	ccaccctcga	aggagatgat	gaectcatct	ggagcggttac	3000
ccagagctgg	atcctcggtg	tatgcgtaga	tgtccttggg	tggagcattc	cacagatctt	3060
ccaggtaacc	ggctctcaata	gcgcggcccc	agacgttctg	gtcgatggag	aatggggact	3120
tcacggactg	ctcgattgga	acgttggtct	cctcggcgaa	ggcgatagcc	ttgtcgcggg	3180
tccatgcgaa	gtcacgagca	ggcgcaatga	tctccaggtt	tggatcggtg	tccatgaagc	3240
cgacctcgaa	acgaacctgg	tcgttgccct	taccagtgc	gccgtgtgca	acgtgggtac	3300
cgttgaactg	cttgccagcc	tcaacgaggt	gcttgacgat	cagtggcgcg	gagattgcag	3360
aaaccagtg	gtactgcttc	atgtacatgc	cgtttgcctt	gatggttggc	aggcagtact	3420
cctcagcgaa	ctcactcttt	gcatacaaca	cgatggactc	agctgcaccc	gcaccaatg	3480
cacgctggcg	aacgttgctc	atgttctctc	cacctggccc	caggctgagg	gaaactgcga	3540
tgacttcacc	atcaatcatc	ttcttcaggt	atggaattgc	cacagtgggt	tccagaccgc	3600
cggagtatgc	aagaacgatg	cggttagtca	tgaggtgtgc	tccttcattt	taagtctctt	3660
gccagccgtg	ttattcttac	ctcggtcgtg	tggccagcag	ccacaccagc	agtgtttctt	3720
gagcgtggag	gcggttttct	gcttcacga	aaactttgga	cgctgggtcca	tcaatcacgg	3780
aggctgccac	ttctttgcca	cggtaggcag	gaaggcagtg	caggaagatg	gcgccgtcgt	3840
tagctttcgc	catgacctca	tcgttgacct	ggtaaggaac	gaaagggtgtg	gtgcgatcga	3900
tgccgtcgtt	ttccataccc	atggataccc	aggatcgggt	gatgacaaca	tcggcgccgg	3960
caacctcgct	gaggctgtcg	gtgacaacaa	ccttcgcgcc	ggtttctctg	ccacgctttt	4020
ccgcgcgcct	cacgaattcc	gcacgaggct	ggaaccttc	aggagcgatg	atggaaatat	4080
ccatgccccg	ggtggcaaa	ccaatcatgt	aggagtgggc	catgttggtg	tcgccatcgc	4140
ccaggtagac	agccttctta	cccttaaggc	ctgctggggc	ttcttcaggg	ctgaggtttt	4200
ccacgatagt	ctgcagatca	gccagaatct	ggcatgggtg	cagatcatcg	gacaaggagt	4260
tcaccagcgg	cacagtggac	gtctccgcca	tggcgtggaa	attgctgtgt	gcgtaggtgc	4320
gccacacaat	tgtctccacg	tagcgggaca	atacagctgc	gggtgcctgc	agggactcgc	4380
ccttaccat	ctgtgagcta	ccggaatcca	cgacgatggc	gtgtccaccc	aaatgagcga	4440
tgcccgctgc	gaaggagaag	cgagtacgag	ttgaagtctt	atcaaaaaga	actgcaacgg	4500
actttggctc	ctcgagtgga	cgtccgaaa	acggcgctgc	cttgagcttt	gcggctaggg	4560
tcaaaacctc	tgctgtctct	gcaggggtga	gatcatcatc	agccagaaaa	tggcgaaacct	4620
gtggttgatg	agtcataagt	ttgagtcctt	tatgcgattg	tctcggcagt	tcgattgtag	4680
gaaatgcaat	gtgtcaagca	gttaggtcaa	tcagagcatc	ggctactcaa	gaattcttga	4740
gtgacagaaa	cccaaggggc	agtatggcga	attttttcaa	ctcacgcacc	aaacctactc	4800
acattggata	attcgtcgag	tcaactccgt	gtcaacctgt	cctcaacttt	agcggggaaa	4860
caaagatttc	gcagcagatg	caaaatcata	cagacacatt	caaaaatgga	cattgacccc	4920
aaacgggggc	atttccaaca	cttgcgctaa	atagtttttg	cagggaaacga	acctgaaatc	4980
gttcgcacta	cgatctgtct	ccccgaatgg	ggtaatttat	gaactacatt	acacccccgt	5040
gaatccacaa	attaaagtct	cagatttgaa	gccaatgtga	tgcaactcac	actggcttca	5100

-continued

```

atggtaatcc caataggctc ggtcgttttt aggcttcaat gcatgcctgt ctggtttctc 5160
ggctctcaaa ccgaacagga gtgcccacgg ctggcaaaaa gttcagttct tgggcacgct 5220
agtctggttt tacctttttg agtgccccga actgccaaac gagagtgcc atgaggggct 5280
aaatctggat ccgcccgcac aaaacaactg ccgcagcaca acgaatcaac gctgtgctgc 5340
ggcagtcact tttttgaagg attaacgcca acccatggtc atgagtaggc cgatgatcat 5400
cagaccgaag ccgatgccat agttccatgc accaagatca gccatgaatg ggatctgtgg 5460
gcccacgagg tagttaatga tcaaccaggc taggcccagc atcatgaagg caaacatgat 5520
gacctgttac cacattgggg ttccggcgga attgatctta accgggggtc ggtttgcgct 5580
tgggttgctt gaaaccggtg cggctctggt tttagttact cttgcctttg gcattgttca 5640
caccttcata atttattgtc ttgaaaaaac cccagcaact ggtacttattc ttgtttgagc 5700
ttaacgtatc agttgccggg ccagctacct tgacgggttc cttgttggtt attgcacgag 5760
tgcagcgaga tcgaattcga agagtgcgac ttgcacttgg gcgtctttgc ggaagagggt 5820
tgcagggggt ggggattgga atccgatttt gttttgatcc acgagggctg cgggtgtggat 5880
ggggtcgccc acgatcaggg attgatctgg ggcggtccag ccagcagcgc ggagggcact 5940
gatggcctgt tcggtgaagc ttggcactgg ccgtcgtttt acaacgtcgt gactgggaaa 6000
accctggcgt tacccaactt aatcgc 6026

```

```

<210> SEQ ID NO 21
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(18)
<223> OTHER INFORMATION: argFGH_f
<220> FEATURE:
<223> OTHER INFORMATION: primer argFGH_f

```

```

<400> SEQUENCE: 21

```

```

tccggctcgt atgttgtg 18

```

```

<210> SEQ ID NO 22
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: argFGH_r
<220> FEATURE:
<223> OTHER INFORMATION: primer argFGH_r

```

```

<400> SEQUENCE: 22

```

```

tggcactggc cgtcgtttta c 21

```

```

<210> SEQ ID NO 23
<211> LENGTH: 11554
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(11554)
<223> OTHER INFORMATION: pK18_IBcg0054::Pg3-argFGH
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (2321)..(2321)

```

-continued

<223> OTHER INFORMATION: n is a, c, g, or t

<220> FEATURE:

<223> OTHER INFORMATION: plasmid pK18_IBcg0054::Pg3-argFGH

<400> SEQUENCE: 23

```

cgataagcta gcttcacgct gccgcaagca ctcagggcgc aagggtgct aaaggaagcg      60
gaacacgtag aaagccagtc cgcagaaacg gtgctgaccc cggatgaatg tcagctactg      120
ggctatctgg acaagggaaa acgcaagcgc aaagagaaag caggtagctt gcagtgggct      180
tacatggcga tagctagact gggcggtttt atggacagca agcgaaccgg aattgccagc      240
tggggcgccc tctggttaag ttgggaagcc ctgcaaagta aactggatgg ctttcttgcc      300
gccaaaggatc tgatggcgca ggggatcaag atctgatcaa gagacaggat gaggatcggt      360
tcgcatgatt gaacaagatg gattgcacgc aggttctccg gccgcttggg tggagaggct      420
attcggctat gactgggcac aacagacaat cggctgctct gatgccgcg tgttccggct      480
gtcagcgcgag gggcgcccg ttctttttgt caagaccgac ctgtccgggt ccttgaatga      540
actccaagac gaggcagcgc ggctatctg gctggccacg acgggcggtt cttgcgcagc      600
tgtgtctgac gttgtcactg aagcgggaag ggactggctg ctattgggcg aagtgccggg      660
gcaggatctc ctgtcatctc accttgctcc tgccgagaaa gtatccatca tggctgatgc      720
aatgcggcgg ctgcatacgc ttgatccggc tacctgccc ttcgaccacc aagcgaaca      780
tcgcatcgag cgagcacgta ctcggatgga agccgggtct gtcatcagg atgatctgga      840
cgaagagcat caggggctcg cgccagccga actgttcgcc aggtcaagg cgcggatgcc      900
cgacggcgag gatctcgctg tgaccatgg cgatgcctgc ttgccgaata tcatggtgga      960
aaatggccgc tttcttgat tcactgactg tggccggctg ggtgtggcgg accgctatca    1020
ggacatagcg ttgggtaccc gtgatattgc tgaagagctt ggcggcgaat gggctgaccg    1080
cttctctgtg ctttacggta tcgccgctcc cgattcgcag cgcctgcct tctatcgct    1140
tcttgacgag ttcttctgag cgggactctg gggttcgcta gaggatcgat cctttttaac    1200
ccatcacata tacctgccgt tcactattat ttagtgaaat gagatattat gatattttct    1260
gaattgtgat taaaaaggca actttatgcc catgcaacag aaactataaa aaatacagag    1320
aatgaaaaga aacagataga ttttttagtt ctttaggccc gtagtctgca aatcctttta    1380
tgattttcta tcaacaaaaa gagggaaaata gaccagttgc aatccaaacg agagtctaata    1440
agaatgaggt cgaagagtaa atcgccgggg ttgttactg ataaagcagg caagacctaa    1500
aatgtgtaaa gggcaaagtg tatactttgg cgtcaccctt tacatatctt aggtcttttt    1560
ttattgtgag taactaactt gccatcttca aacaggaggg ctggaagaag cagaccgcta    1620
acacagtaca taaaaaagga gacatgaacg atgaacatca aaaagtgtgc aaacaagca    1680
acagtattaa cctttactac cgcactgctg gcaggaggcg caactcaagc gtttgcgaaa    1740
gaaacgaacc aaaagccata taaggaaaca tacggcattt cccatattac acgccatgat    1800
atgctgcaaa tcctgaaca gcaaaaaaat gaaaaatgc aagtttctga atttgattcg    1860
tccacaatta aaaatatctc ttctgcaaaa ggcttgacg tttgggacag ctggccatta    1920
caaacgctg acggcactgt cgcaactat cacggctacc acatcgtctt tgcattagcc    1980
ggagatccta aaaatgcgga tgacacatcg atttacatg tctatcaaaa agtcggcgaa    2040
acttctattg acagctggaa aaacgctggc cgcgtcttta aagacagcga caaattcgat    2100

```

-continued

gcaaatgatt	ctatcctaaa	agaccaaaca	caagaatggt	caggttcagc	cacatttaca	2160
tctgacggaa	aatccggtt	attctacact	gatttctccg	gtaaacatta	cggcaaaaca	2220
acactgacaa	ctgcacaagt	taacgtatca	gcacagaca	gctctttgaa	catcaacggt	2280
gtagaggatt	ataaatcaat	ctttgacggt	gacggaaaaa	ncgtatcaaa	atgtacagca	2340
gttcacgat	gaaggcaact	acagctcagg	cgacaacat	acgctgagag	atcctcacta	2400
cgtagaagat	aaaggccaca	aatacttagt	atttgaagca	aacactggaa	ctgaagatgg	2460
ctaccaaggc	gaagaatctt	tatttaacaa	agcatactat	ggcaaaagca	catcattctt	2520
cgtcaagaa	agtcaaaaac	tcttgcaaag	cgataaaaaa	cgcacggctg	agttagcaaa	2580
cggcgctctc	ggtatgattg	agctaaacga	tgattacaca	ctgaaaaaag	tgatgaaacc	2640
gctgattgca	tctaacacag	taacagatga	aattgaacgc	gcgaacgtct	ttaaaatgaa	2700
cggcaaatgg	tacctgttca	ctgactcccg	cggatcaaaa	atgacgattg	acggcattac	2760
gtctaacgat	atttcatgc	ttgggttatgt	ttctaattct	ttaactggcc	catacaagcc	2820
gctgaacaaa	actggccttg	tgtaaaaaat	ggatcttgat	cctaacgatg	taacctttac	2880
ttactcacac	ttcgctgtac	ctcaagcgaa	aggaaacaat	gtcgtgatta	caagctatat	2940
gacaaacaga	ggattctacg	cagacaaaca	atcaacgttt	gcgccgagct	tctgctgaa	3000
catcaaaggc	aagaaaacat	ctgttgtaa	agacagcatc	cttgaacaag	gacaattaac	3060
agttaacaaa	taaaaacgca	aaagaaaatg	cggatgggta	ccgagcgaaa	tgaccgacca	3120
agcgacgccc	aacctgccat	cacgagattt	cgattccacc	gccgccttct	atgaaagggt	3180
gggcttcgga	atcgttttcc	gggacgccct	cgcggacgtg	ctcatagtcc	acgacgcccg	3240
tgattttgta	gccctggcgg	acggccagca	ggtaggccga	caggctcatg	ccggccgccc	3300
ccgccttttc	ctcaatcgct	cttcgttcgt	ctggaaggca	gtacaccttg	atagggtggc	3360
tgcccttcct	ggttggttg	gtttcatcag	ccatccgctt	gccctcatct	gttacgccgg	3420
cggtagcccg	ccagcctcgc	agagcaggat	tcccgttgag	caccgccagg	tgcaataaag	3480
ggacagtga	gaaggaaac	ccgctcgcgg	gtgggcctac	ttcacctatc	ctgccccgct	3540
gacgccgttg	gatacaccaa	ggaaagtcta	cacgaaccct	ttggcaaaat	cctgtatatc	3600
gtgcgaaaaa	ggatggatat	accgaaaaaa	tcgctataat	gacccgaaag	cagggttatg	3660
cagcggaaaa	gcgctgcttc	cctgctgttt	tgtggaatat	ctaccgactg	gaaacaggca	3720
aatgcaggaa	attactgaac	tgaggggaca	ggcgagagac	gatgccaaag	agctcctgaa	3780
aatctcgata	actcaaaaaa	tacgcccggt	agtgatctta	tttcattatg	gtgaaagttg	3840
gaacctctta	cgtgccgac	aacgtctcat	tttcgccaaa	agttggccca	gggcttcccg	3900
gtatcaacag	ggacaccagg	atttatattat	tctgcgaagt	gatcttccgt	cacaggtatt	3960
tattcggcgc	aaagtgcgtc	gggtgatgct	gccaaacttac	tgatttagtg	tatgatgggtg	4020
tttttgaggt	gctccagtgg	cttctgtttc	tatcagctcc	tgaaaatctc	gataactcaa	4080
aaaatacgcc	cggtagtgat	cttatctcat	tatggtgaaa	gttggaacct	cttacgtgcc	4140
gatcaacgct	tcatttttcg	caaaagtgg	cccagggtct	cccggatatca	acagggacac	4200
caggatttat	ttattctgcg	aagtgatctt	ccgtcacagg	tatttattcg	gcgcaaagtg	4260
cgtcgggtga	tgctgccaac	ttactgattt	agtgatgat	ggtgtttttg	aggtgctcca	4320
gtggcttctg	ttctatcag	ggctggatga	tcctccagcg	cggggatctc	atgctggagt	4380

-continued

tcttcgccca	ccccaaaagg	atctaggtga	agatcctttt	tgataatctc	atgacaaaaa	4440
tcccttaacg	tgagttttcg	ttccactgag	cgtcagaccc	cgtagaaaag	atcaaaggat	4500
cttcttgaga	tccttttttt	ctgcgcgtaa	tctgctgctt	gcaaacaaaa	aaaccaccgc	4560
taccagcggg	ggtttggttg	cggatcaag	agctaccaac	tctttttccg	aaggtaactg	4620
gcttcagcag	agcgcagata	ccaaatactg	tccttctagt	gtagccgtag	ttaggccacc	4680
acttcaagaa	ctctgtagca	cgcctacat	acctcgctct	gctaatactg	ttaccagtgg	4740
ctgctgccag	tggcgataag	tcgtgtctta	ccgggttgga	ctcaagacga	tagttaccgg	4800
ataaggcgca	gcggtcgggc	tgaacggggg	gttcgtgcac	acagcccagc	ttggagcgaa	4860
cgaactacac	cgaactgaga	tacctacagc	gtgagcattg	agaaagcgcc	acgcttcccg	4920
aaggagagaaa	ggcggacagg	tatccggtaa	gcggcagggt	cggaacagga	gagcgcacga	4980
gggagcttcc	agggggaaac	gcctggtatc	tttatagtc	tgctgggttt	cgccacctct	5040
gacttgagcg	tcgatttttg	tgatgctcgt	cagggggggc	gagcctatgg	aaaaacgcca	5100
gcaacgcggc	ctttttacgg	ttcctggcct	tttgctggcc	ttttgctcac	atgttctttc	5160
ctgcgttata	ccctgattct	gtggataaac	gtattaccgc	ctttgagtga	gctgataacc	5220
ctcgcgcgag	ccgaacgacc	gagcgcagcg	agtcagtggg	cgagggaagc	gaagagcgcc	5280
caatacgcga	accgcctctc	cccgcgcgtt	ggccgattca	ttaatgcagc	tggcagcaca	5340
ggtttcccca	ctggaagcgc	ggcagtgagc	gcaacgcaat	taatgtgagt	tagctcactc	5400
attaggcacc	ccaggcttta	cactttatgc	ttccggctcg	tatgttggtg	ggaattgtga	5460
gcggataaca	atttcacaca	ggaacagct	atgaccatga	ttacgaattg	cgtggcaatc	5520
aggctttacc	tcagcgaaga	attcgacgac	gccaccgcgc	tctacaccgt	ccgatccttt	5580
gatgccgcaa	ctgaagcat	cgtggtggat	gtggttcaac	accaccacga	aagccccatg	5640
atgcgctggt	cagacaccgt	aaaaatcaac	gacaccctcg	tgctcaccgg	accccgccca	5700
cactttgtca	tccccgaagg	cgaacaagca	gcaactcttc	ttgatgacac	cgccatcccc	5760
gctctgcgcg	ctatttttga	tcaatggcca	acagatcttc	gtggcaaaag	atgggttgtc	5820
actgacgata	cgcagcctt	cgatgaacta	cccagcatcg	acggactgga	actgaacctg	5880
ctcgcgccgg	gatcagatcc	aactgttcag	ccacttgccc	aacaggcata	tgacctggaa	5940
aaccagagaa	cttacgtggt	gtgggcagcc	ggcgcagcag	atgaaataaa	atccatccgc	6000
aggcaacttc	gcaagcagg	gggattggaa	aaagatgcag	tggccgtggt	tgggtactgg	6060
aaatacaaca	ccaccaacac	tcagatcgat	gcagtccgca	aagaaaacta	catgaagatg	6120
ctctctgaag	ggctacagct	ggaaaacttc	gacgacctct	cattggagat	ttaaggggtc	6180
gagtttttag	aatgggtggt	tgcattgatt	ggcgtctgat	gagctcttag	aagcgattct	6240
gggagggcat	gatttttggg	attcggagct	gaatgtggac	tgattttggg	ggaaccagac	6300
tggcgtgccc	aaatgtcgtg	tcttcagtcc	taggtcaata	ttacaggcca	taaactaatg	6360
cttatcgacg	tacccccgca	cgtgcccaact	cgcgctgagc	ggtaacttgc	gcatcgacac	6420
gtgcgcgttg	ctccgcaacc	cgcacgcccc	cggttccacc	gcgcgttgca	cggaagacca	6480
ctgcaccatc	aatggtgagc	acttcccgtg	cctctggggg	cagacgtgca	tcaacgccac	6540
tgagttcttc	atcagtgaga	tcgataaggt	ccactcccct	ggactccgcg	atccgcacgc	6600
aagcgccgga	tgttctggtg	gcctcacgga	atggaacgcc	ctggcgcacc	atccactcag	6660

-continued

ccaagtcggt	ggcaaggggtg	aaacctgctg	gtgcaagttc	acgcatgcgc	tcggtgttga	6720
aggtcaaggt	ggaacacaaa	ccagtcattg	cagggagcag	caggttgagc	tgcgccacgg	6780
aatctacgat	tggttcctta	tcttctgca	ggtcgcggtt	gtacgctaaa	ggctgtgcct	6840
tcagggtagc	cagcagaccg	gtgaggttac	cgatcaagcg	accagacttg	ccacgggtca	6900
gctctgccac	gtcaggggtc	ttcttctgcg	gcatgattga	gctgcctgtg	gaccaggaat	6960
cagacaaggt	gatgtaacca	aattctgggg	tgccaccatgc	gatgatttct	tcagccaagc	7020
gggacataac	cactgcaagc	tgccgcagca	cgaaggcggg	ttcagatgcg	aaatcgcggg	7080
agctgggtgg	atcaatggag	ttatctgctg	cggaaacaaa	gccgagttct	tcagcgattg	7140
cttcagggtt	gagcttcaaa	gaggaaccag	caagtgcgcc	ggaaccgtaa	ggagacaccg	7200
caagacgctt	gtccagggtc	cggatacgat	caatatcgcg	cagcaaaggc	tgtgcgtgtg	7260
ccagcagctg	gtgtgccaga	aggaccggct	gagctgcctg	gaagtgggtc	ttgcctggca	7320
tgatcgcgcc	tgcatgtgcc	ttagcttggg	cgctgagggc	gtcgacaagc	tcggtgtgtc	7380
ccagcgcgat	gtcgcgcact	gcgtgcggga	cccacatgcg	gaacagggtt	gccacctggt	7440
cgttgcggga	acgaccagcg	cgcagacggc	cgccaccctc	aggaccaacg	cggatcaatca	7500
gaccgcgttc	catcgcgccg	tgccatccct	catcagaagg	cagcggaccg	aagggtccgt	7560
cggcgacatc	cttgcccagc	tgatcaagcc	cagccagcat	ggtggctaga	tcttcacag	7620
aaagtagatc	tgcttggtgc	aaaaccttgg	cggtgcctt	ggaggccaac	acatcataag	7680
ggggcaaaa	ccagtcgaaa	tgagtggaga	cactcaaggc	gaacatggcc	tcggagggtc	7740
caccggagaa	gcggccgccc	cacagcgcac	cttcattggt	tccgtgctgt	tccatgtggt	7800
gttctcttac	tttctaaata	cttgaagttc	tagtctggat	gcttgaaaag	gtggcttagt	7860
tgttgccagc	ttcgcgatcg	cgcttggttag	cgatcttgga	ggacagaccg	tgccagctgga	7920
caaagccctt	agccagggtc	tggtcgaaag	gtcgccggg	gtcgtaggta	gccaggttga	7980
agtcgtacag	ggagtggctg	gaacgacgac	cattgatggt	gatggaacct	gcgtgcagaa	8040
ccatgcggat	atcgccgggtg	acgtgctcct	gggtggaatc	aatgaacgcg	tcaggggagc	8100
gcttcagagg	tccgaaccac	aggccgtcgt	atacttcctc	agcccaacgt	gcgtcaacgc	8160
cgcgcttgta	gcgagccagt	tcgcgctcga	tggtgacatc	ttccaaagcc	tcgtgagcct	8220
taatcagtgc	gattgcgcct	ggtgcttctg	agatttcgcg	ggacttgatg	cccacgagac	8280
ggtcctcaac	catgtcaagg	cggccaacgc	cctgtgcgcc	tgccagacgg	ttcagctctt	8340
caatagcctg	cagtacggag	actggacggc	catcgatgga	gactggcttg	ccaccctcga	8400
aggagatgat	gacctcatct	ggagcggttac	ccagagctgg	atcctcggtg	tatgcgtaga	8460
tgctcttggt	tggagcattc	cacagatctt	ccaggttaacc	ggtctcaata	gcgcggcccc	8520
agacgttctg	gtcgatggag	aatggggact	tcacggactg	ctcgattgga	acgttgttct	8580
cctcggcgaa	ggcgatagcc	ttgtcgcggg	tccatgcgaa	gtccagagca	ggtgcaatga	8640
tctccagggt	tggatcggtg	tccatgaagc	cgacctcgaa	acgaacctgg	tcgttgccct	8700
taccagtgc	gccgtgtgca	acgtgggtac	cggtgaactg	cttgccagcc	tcaacgaggt	8760
gcttgacgat	cagtggggcg	gagattgcag	aaaccagtg	gtactgcttc	atgtacatgc	8820
cgtttgcctt	gatggttggc	aggcagtact	cctcagcgaa	ctcatccttt	gcatcaacaa	8880
cgatggactc	agctgcaccg	gcatccaatg	cacgctggcg	aacgttgttc	atgttctctc	8940

-continued

cacctggcc	caggtcgagg	gaaactgcga	tgacttcacc	atcaatcacc	ttcttcaggt	9000
atggaattgc	cacagtgggtg	tccagaccgc	cggagtatgc	aagaacgatg	cggttagtca	9060
tgaggtgtgc	tccttcattt	taagtctctt	gccagccgtg	ttattcttac	ctcggtggt	9120
tggccagcag	ccacaccagc	agtgttttct	gagcgtggag	gcggttttct	gcttcacga	9180
aaactttgga	cgttggtcca	tcaatcacgg	aggctgccac	ttctttgcca	cggtaggcag	9240
gaaggcagtg	caggaagatg	gcgccgtcgt	tagctttcgc	catgacctca	tcgttgacct	9300
ggtaaggaac	gaaaggtgtg	gtgcgatcga	tgcctcgtt	ttccataccc	atggataccc	9360
aggatcgggt	gatgacaaca	tcggcgccgg	caacctcgtc	gaggctgtcg	gtgacaacaa	9420
ccttcgcgcc	ggtttctctg	ccacgctttt	cgcgcgcgtc	cacgaattcc	gcacgaggct	9480
ggaaccttc	aggagcgatg	atggaaatat	ccatgccgcg	ggtggcaaag	ccaatcatgt	9540
aggagtgtgc	catgttgttg	tcgccatcgc	ccaggtaaac	agccttctta	cccttaaggc	9600
ctgctggggc	ttcttcaggg	ctgaggtttt	ccacgatagt	ctgcagatca	gccagaatct	9660
ggcatgggtg	cagatcatcg	gacaaggagt	tcaccagcgg	cacagtggac	gtctccgcca	9720
tggcgtgga	attgctgtgt	gcgtaggtgc	gccacacaat	tgcttccacg	tagcgggaca	9780
atacagctgc	ggtgtctctc	agggactcgc	ccttaccat	ctgtgagcta	ccggaatcca	9840
cgacgatggc	gtgtccaccc	aaatgagcga	tgcccgcgtc	gaaggagaag	cgagtacgag	9900
ttgaagtctt	atcaaaaaga	actgcaacgg	actttgggtc	ctcgagtgga	cgctccgaaa	9960
acggcgcgtc	cttgagcttt	gcggctaggg	tcaaaacctc	tgctgctct	gcaggggtga	10020
gatcatcacc	agccagaaaa	tggcgaaact	gtggttgtga	agtcataagt	ttgagtcctt	10080
tatgcgattg	tctcggcagt	tcgattgtag	gaaatgcaat	gtgtcaagca	gttaggtcaa	10140
tcagagcacc	ggtcactcaa	gaattcttga	gctgacagaa	cccaagggcg	agtatggcga	10200
attttttcaa	ctcagccacc	aaacctactc	acattggata	attcgtcgag	tcaactccgt	10260
gtcaacctgt	cctcaacttt	agcggggaaa	caaagatttc	gcagcagatg	caaaatcata	10320
cagacacatt	caaaaatgga	cattgacccc	aaacgggggc	atttccaaca	cttgcgctaa	10380
atagtttttg	cagggaaacga	acctgaaatc	gttcgcacta	cgatctgctc	cccgaatgg	10440
ggtaattat	gaactacatt	acacccccgt	gaatccacaa	attaaagtct	cagatttgaa	10500
gccaatgtga	tgcaactcac	actggttcca	atggtaatcc	caataggctc	ggctggtttt	10560
aggcttcaat	gcctgcctgt	ctggtttctc	ggctctcaaa	ccgaacagga	gtgcccacgg	10620
ctggcaaaaa	gttcagttct	tgggcacgct	agtctgggtt	tacctttttg	agtgcgccga	10680
actgccaaac	gagagtgcgc	atgaggggct	aaatctggat	ccgcccgcac	aaaacaactg	10740
ccgcagcaca	acgaatcaac	gctgtgctgc	ggcagtcact	tttttgaagg	attaacgcca	10800
acccatggtc	atgagtaggc	cgatgatcat	cagaccgaag	ccgatgccat	agttccatgc	10860
accaagatca	gccatgaatg	ggatctgtgg	gcccacgagg	tagttaatga	tcaaccaggc	10920
taggcccagc	atcatgaagg	caaacatgat	gacctgttac	cacattgggg	ttccggcgga	10980
attgatctta	accgggggtgc	ggtttgcgct	tgggttgctt	gaaaccgggtg	cggctctcgtt	11040
tttagttact	cttgcccttg	gcattgttca	caccttcata	atttattgtc	ttggaaaaac	11100
cccagcaact	ggtacttacc	ttgtttgagc	ttaacgtatc	agttgccggg	ccagctacct	11160
tgacgggttc	cttggtggct	attgcacgag	tgacgcgaga	tcgaattcga	agagtcgcac	11220

-continued

ttgcacttgg gcgtctttgc ggaagagggt tgcaggggtt ggggattgga atccgatttt	11280
gttttgatcc acgagggctg cggtgtggat ggggtcgccg acgatcaggg attgatctgg	11340
ggcgggtccag ccagcagcgc ggagggcact gatggcctgt tcggtgaage ttggcactgg	11400
ccgtcgtttt acaacgtcgt gactgggaaa accctggcgt tacccaactt aatcgccctg	11460
cagcacatcc ccccttcgcc agctggcgta atagcgaaga ggcccgacc gatcgccctt	11520
cccaacagtt gcgcagcctg aatggcgaat ggcg	11554

1. A microorganism having an increased ability to provide L-arginine compared with the ability of the wildtype microorganism, comprising:

at least one gene coding for a protein having the function of an L-arginine:glycine amidinotransferase and having a decreased activity of a protein having the function of an arginine exporter compared with the activity of the respective protein in the wildtype microorganism at the same status of the cell cycle.

2. The microorganism of claim 1, wherein the activity of the arginine responsive repressor protein ArgR is attenuated or deleted.

3. The microorganism of claim 1 having increased activities of an enzyme having the function of a carbamoyl phosphate synthase compared to the respective enzymic activity in the wildtype microorganism.

4. The microorganism of claim 3, wherein the increased activity of the enzyme having the function of a carbamoyl phosphate synthase is achieved by a mutation and/or overexpression of a gene coding for the enzyme having the function of a carbamoyl phosphate synthase.

5. The microorganism of claim 1, wherein at least one or more of the genes coding for an enzyme of the biosynthetic pathway of L-ornithine and L-arginine, comprising argF/argF2 coding for an ornithine carbamoyl transferase, argG coding for an argininosuccinate synthetase and argH coding for an argininosuccinate lyase is overexpressed.

6. The microorganism of claim 1, wherein a gene gdh coding for a glutamate dehydrogenase, is overexpressed.

7. The microorganism of claim 1, wherein the gene coding for the protein having the function of an L-arginine:glycine amidinotransferase is heterologous.

8. The microorganism of claim 1, wherein the protein having the function of an L-arginine:glycine amidinotransferase comprises an amino acid sequence which is at least 70% identical to the amino acid sequence according to SEQ ID NO: 13.

9. The microorganism of claim 1, wherein the gene coding for the protein having the function of an arginine exporter is inactivated or deleted.

10. The microorganism of claim 1, wherein a gene coding for a transcriptional activator of the gene coding for the protein having the function of an arginine exporter is deleted.

11. The microorganism of claim 1, wherein the microorganism belongs to the genus *Corynebacterium*, to the genus *Enterobacteriaceae* or to the genus *Pseudomonas*.

12. The microorganism of claim 11, wherein the microorganism is *Corynebacterium glutamicum* and the gene coding for the protein having the function of an arginine exporter is lysE and the gene coding for the transcriptional activator is lysG.

13. The microorganism of claim 11 wherein the microorganism is *Escherichia coli* and the gene coding for the protein having the function of an arginine exporter is argO (ybjE).

14. The microorganism of claim 11 wherein the microorganism is *Pseudomonas putida* and the protein having the function of an arginine exporter is lysE.

15. The method for the fermentative production of guanidino acetic acid (GAA), comprising:

- a) cultivating the microorganism as defined in claim 1 in a suitable medium under suitable conditions, and
- b) accumulating GAA in the medium to form a GAA containing fermentation broth.

16. The method of claim 15, further comprising:

isolating GAA from the GAA containing fermentation broth.

17. The method of claim 15, further comprising:

drying and/or granulating the GAA containing fermentation broth.

18. The microorganism as claimed in claim 1, further comprising:

a gene coding for an enzyme having the activity of a guanidinoacetate N-methyltransferase.

19. The microorganism of claim 18, wherein the gene coding for an enzyme having the activity of a guanidinoacetate N-methyltransferase is overexpressed.

20. The method for the fermentative production of creatine, comprising:

- a) cultivating the microorganism as defined in claim 18 in a suitable medium under suitable conditions, and
- b) accumulating creatine in the medium to form a creatine containing fermentation broth.

21. The method of claim 20, further comprising isolating creatine from the creatine containing fermentation broth.

* * * * *