



US 20250276998A1

(19) **United States**(12) **Patent Application Publication**
Seyfert et al.(10) **Pub. No.: US 2025/0276998 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **NOVEL DAROBACTIN DERIVATIVES***A61P 31/04* (2006.01)*C12N 15/70* (2006.01)(71) Applicant: **Helmholtz-Zentrum Fur
Infektionsforschung GmbH,**
Braunschweig (DE)(52) **U.S. Cl.**CPC *C07K 7/06* (2013.01); *A61K 38/08*
(2013.01); *A61P 31/04* (2018.01); *C12N 15/70*
(2013.01)(72) Inventors: **Carsten E. Seyfert**, Braunschweig
(DE); **Fabian T. Panter**, Braunschweig
(DE); **Rolf Müller**, Braunschweig (DE);
Selina Deckarm, Braunschweig (DE);
Sebastian Gross, Braunschweig (DE);
Domen Scherzer, Braunschweig (DE);
Chantal Bader, Braunschweig (DE);
Jennifer Herrmann, Braunschweig
(DE)

(57)

ABSTRACTThe present invention relates to darobactin derivatives of
formula (I), methods of their use, and a method of produc-
tion.(21) Appl. No.: **18/277,870**(22) PCT Filed: **Feb. 18, 2022**(86) PCT No.: **PCT/EP2022/054063**

§ 371 (c)(1),

(2) Date: **Aug. 18, 2023**(30) **Foreign Application Priority Data**

Feb. 19, 2021 (EP) 21158189.7

Jul. 22, 2021 (EP) 21187173.6




Publication Classification(51) **Int. Cl.***C07K 7/06* (2006.01)*A61K 38/08* (2019.01)**Specification includes a Sequence Listing.**

Fig. 2



NOVEL DAROBACTIN DERIVATIVES

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a U.S. national phase application, pursuant to 35 U.S.C. § 371, of PCT Application No. PCT/EP2022/054063, filed Feb. 18, 2022, which claims priority from European Patent Application No. 21158189.7, filed on Feb. 19, 2021 and European Application No. 21187173.6, filed on Jul. 22, 2021. The entire contents of these applications are incorporated herein by reference in their entirety.

INCORPORATION BY REFERENCE OF SEQUENCE LISTING


[0002] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 19, 2024, is named 348710_01501_SequenceListing.txt and is 58,400 bytes in size.

[0003] The present invention relates to novel derivatives of darobactin, methods of their use, and a method of production.

[0004] The emerging number of antibiotic-resistant bacterial pathogens leads to increasing mortality rates in humans. Many deaths are caused by the especially troublesome bacteria belonging to the ESKAPE panel (vancomycin-resistant *Enterococci*, *Staphylococcus aureus*, *Klebsiella pneumoniae*, *Acinetobacter baumannii*, *Pseudomonas aeruginosa* and *Enterobacter* species). The World Health Organization recommended focusing on the discovery and development of new antibiotics with anti-Gram-negative activity.


[0005] Darobactin A is a ribosomally-produced peptide antibiotic that selectively kills Gram-negative pathogens, including bacteria from the ESKAPE panel (Imai, Y., Meyer, K. J., Iinishi, A. et al. A new antibiotic selectively kills Gram-negative pathogens. *Nature* 576, 459-464 (2019); WO 2020/018173). However, the production yield of Darobactin A in hitherto known producer strains is poor, which limits the possibilities to perform structure engineering or isolate sufficient amounts of compound needed for potential semi-synthesis approaches.

[0006] In one aspect, the present invention provides compounds of formula (I) or a salt thereof:



wherein


[0007] R^1 , R^2 , R^4 and R^5 are independently selected from H, CH_3 , CH_2OH , one of the following groups:



[0008] $-CH_2-SR^{14}$, wherein R^{14} is independently selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted;

[0009] or $-CH_2-Ind$, wherein Ind is an optionally substituted indole group;

[0010] R^3 is selected from H, OH, SH, COOH, CONH₂, one of the following groups:



[0011] —SR^{3,4}, wherein R^{3,4} is selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted;

[0012] or -Ind, wherein Ind is an optionally substituted indole group;

[0013] R⁶ is a hydrogen atom or a methyl group (especially a hydrogen atom);

[0014] R⁷ is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom);

[0015] R⁸ is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group; and

[0016] m is an integer of from 0 to 3 (especially 0 or 1; preferably 0);

[0017] R^{6d} is a hydrogen atom or a methyl group (especially a hydrogen atom);


[0018] R^{7,4} is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom);

[0019] $R^{8,d}$ is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group; and


[0020] p is an integer of from 0 to 3 (especially 0 or 1; preferably 0);

[0021] with the proviso, that R⁵ is not a group of formula —CH₂—phenyl if R⁴ is a group of formula —CH₂—OH or —CH₂—CH₂—CH₂—NH—C(=NH)—NH₂.

[0022] Preferably, R⁵ is selected from H, CH₃, CH₂OH or one of the following groups:




-continued




[0023] Further preferably, R⁵ is a group of formula —CH₂-Ind, wherein Ind is an optionally substituted indole group.

[0024] Moreover preferably, R⁵ is a group of formula —CH₃.

[0025] Further preferably, R^5 is a group of the following formula:




[0026] Moreover preferably, the group of formula $-\text{CH}_2-$ Ind is a group of the following formula:




wherein R⁹ is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom); R¹⁰ is a hydrogen atom or a methyl group (especially a hydrogen atom); R¹¹ is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).

[0027] Especially preferably, R^5 is a group of the following formula:




wherein R⁹ is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom); R¹⁰ is a hydrogen atom or a methyl group (especially a hydrogen atom); R¹¹ is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).

[0028] In some preferred embodiments, R^5 is a group of the following formula:




[0029] wherein R^9 is a hydrogen or halogen atom (especially a hydrogen atom); R^{10} is a hydrogen atom or a methyl group (especially a hydrogen atom); R^{11} is, at each occasion, independently selected from a halogen atom (especially fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).

[0030] Further especially preferably, R^5 is a group of the following formula:





[0031] In some embodiments, the group of formula $-\text{CH}_2-\text{SR}^{1A}$ can preferably be:



[0032] Further preferably, R^1 is selected from the following groups: $-\text{CH}_2-\text{CONH}_2$, $-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}_2-\text{CONH}_2$ and $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$; and more preferably R^1 is a group of $-\text{CH}_2-\text{CONH}_2$.


[0033] Preferably, R^2 is selected from CH_3 , CH_2OH , one of the following groups:



[0034] $-\text{CH}_2-\text{SR}^{1A}$, wherein R^{1A} is independently selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted.


[0035] Moreover preferably, R^2 is selected from the following groups: $-\text{CH}_3$, $-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{SH}$, $-\text{CH}_2-\text{CONH}_2$, $-\text{CH}_2-\text{CH}_2-\text{CONH}_2$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$, and $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$, such as $-\text{CH}_3$, $-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{SH}$, and $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$, or such as $-\text{CH}_2-\text{OH}$ and $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$.


[0036] Moreover preferably, R^2 is selected from the following groups: $-\text{CH}_2-\text{OH}$, $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$ and $-\text{CH}_3$, or represents a group:




[0037] Further preferably, R^3 is a hydrogen atom or selected from the following groups: $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2$ and $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$. In some preferred embodiments, R^3 is a group: $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2$ or $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$, preferably the group: $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2$.

[0038] R^4 can preferably represent a hydrogen atom, CH_3 , CH_2OH , one of the following groups:







[0039] $-\text{CH}_2-\text{SR}^{1A}$, wherein R^{1A} is independently selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted (e.g. a group:



[0040] Further preferably, R^4 is CH_3 , CH_2OH , or selected from one of the following groups:




such as CH_3 , CH_2OH , $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}_2$, $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$, $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$ or



or such as CH_3 , CH_2OH , $-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$, or $-\text{CH}_2-\text{CH}(\text{CH}_3)-\text{OH}$.

[0041] Moreover preferably, R^4 is selected from the following groups: $-\text{CH}_2-\text{OH}$, $-\text{CH}_3$ and $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$.

[0042] Further preferably, R^4 is a group of formula $-\text{CH}_2-\text{SR}^{4A}$, wherein R^{4A} is selected from a hydrogen atom, an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted. A preferred example of a group of formula $-\text{CH}_2-\text{SR}^{4A}$ is:



[0043] In some embodiments, R^6 is a methyl group or a hydrogen atom, preferably a hydrogen atom. In some embodiments, R^{6A} is a methyl group or a hydrogen atom, preferably a hydrogen atom. In some embodiments, each of R^6 and R^{6A} is a hydrogen atom.

[0044] In some embodiments, R^7 is a hydrogen or halogen atom, preferably a hydrogen atom.

[0045] In some embodiments, R^{7A} is a hydrogen or halogen atom, preferably a hydrogen atom.

[0046] In some embodiments, each of R^7 and R^{7A} is a hydrogen atom.

[0047] In some embodiments, each of R^6 , R^{6A} , R^7 , and R^{7A} is a hydrogen atom.

[0048] In some embodiments, R^8 is a halogen atom, preferably a fluor or chlor atom, and more preferably a fluor atom. In some embodiments, R^{8A} is a halogen atom, preferably a fluor or chlor atom, and more preferably a fluor atom. In some embodiments, one of R^8 and R^{8A} is present.


[0049] In some embodiments, m is 0, 1, or 2, preferably 0 or 1, more preferably 0.

[0050] In some embodiments, p is 0, 1, or 2, preferably 0 or 1, more preferably 0.

[0051] In some embodiments, n is 0, 1, 2 or 3, preferably 0, 1 or 2, more preferably 0 or 1, especially 0.


[0052] Moreover preferred are compounds of formula (Ia) or a salt thereof:

(Ia)




wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^{6A} , R^{7A} , R^{8A} , m and p are as defined above.

[0053] Further preferred are compounds of formula (Ia') or a salt thereof:




wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 , R^7 , R^8 , R^{6A} , R^{7A} , R^{8A} , m and p are as defined above.

[0054] Moreover preferred are compounds of formula (Ib) or a salt thereof:




wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined above.

[0055] Further preferred are compounds of formula (Ib') or a salt thereof:




wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined above.

[0056] Moreover preferred are compounds of formula (Ib'') or a salt thereof:




wherein R^1 , R^2 , R^3 , R^4 and R^5 are as defined above.

[0057] In some embodiments, the compound of formula (I) is a compound of formula (1):



or a salt (especially a pharmaceutically acceptable salt) thereof, wherein R^2 and R^4 are as defined above.

[0058] In some embodiments, the compound of formula (1) is a compound of formula (1a) or formula (1a'):



or a salt (especially a pharmaceutically acceptable salt) thereof, wherein R^2 and R^4 are as defined above.

[0059] Compounds including suitable combinations of preferred embodiments of the compounds according to the invention, or a salt thereof, are particularly preferred; e.g. a compound or salt thereof including a combination of preferred embodiments of residues R^1 , R^2 , R^3 , R^4 and R^5 as disclosed herein. In other words, the present invention specifically encompasses all possible combinations of residues as indicated above, which result in a stable compound.

[0060] The most preferred compounds of the present invention are the compounds disclosed in the examples, or a salt thereof.

[0061] The expression alkyl refers to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 20 carbon atoms, preferably from 1 to 10 carbon atoms, especially from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms, for example a methyl (Me, CH_3), ethyl (Et), n-propyl (nPr), iso-propyl (iPr), n-butyl (nBu), iso-butyl (iBu), sec-butyl (sBu), tert-butyl (tBu), n-pentyl, iso-pentyl, n-hexyl, 2,2-dimethylbutyl or n-octyl group.

[0062] Especially preferred alkyl groups are C_{1-6} alkyl groups; moreover preferred alkyl groups are C_{1-4} alkyl groups.

[0063] The expression C_{1-6} alkyl refers to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 6 carbon atoms. The expression C_{1-4} alkyl refers

to a saturated, straight-chain or branched hydrocarbon group that contains from 1 to 4 carbon atoms. Examples are the methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl or the tert-butyl group.

[0064] The expressions alkenyl and alkynyl refer to at least partially unsaturated, straight-chain or branched hydrocarbon groups that contain from 2 to 20 carbon atoms, preferably from 2 to 10 carbon atoms, especially from 2 to 6 (e.g. 2, 3 or 4) carbon atoms, for example an ethenyl (vinyl), propenyl (allyl), iso-propenyl, butenyl, ethynyl (acetylenyl), propynyl (e.g. propargyl), butynyl, isoprenyl or hex-2-enyl group. Preferably, alkenyl groups have one or two (especially preferably one) double bond(s), and alkynyl groups have one or two (especially preferably one) triple bond(s).

[0065] Furthermore, the terms alkyl, alkenyl and alkynyl refer to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl) such as, for example, a 2,2,2-trichloroethyl or a trifluoromethyl group.

[0066] The expression heteroalkyl refers to an alkyl, alkenyl or alkynyl group in which one or more (preferably 1 to 8; especially preferably 1, 2, 3 or 4) carbon atoms have been replaced by an oxygen, nitrogen, phosphorus, boron, selenium, silicon or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom) or by a SO or a SO_2 group. The expression heteroalkyl furthermore refers to a carboxylic acid or to a group derived from a carboxylic acid, such as, for example, acyl, acylalkyl, alkoxycarbonyl, acyloxy, acyloxyalkyl, carboxyalkylamide or alkoxycarbonyloxy. Furthermore, the term heteroalkyl refers to groups in which one or more hydrogen atoms have been replaced by a halogen atom (preferably F or Cl).

[0067] Preferably, a heteroalkyl group contains from 1 to 12 carbon atoms and from 1 to 8 heteroatoms selected from oxygen, nitrogen and sulfur (especially oxygen and nitrogen). Especially preferably, a heteroalkyl group contains from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms and 1, 2, 3 or 4 (especially 1, 2 or 3) heteroatoms selected from oxygen, nitrogen and sulfur (especially oxygen and nitrogen). The term C₁-C₁₀ heteroalkyl refers to a heteroalkyl group containing from 1 to 10 carbon atoms and 1, 2, 3, 4, 5 or 6 heteroatoms selected from O, S and/or N (especially O and/or N). The term C₁-C₆ heteroalkyl refers to a heteroalkyl group containing from 1 to 6 carbon atoms and 1, 2, 3 or 4 heteroatoms selected from O, S and/or N (especially O and/or N). The term C₁-C₄ heteroalkyl refers to a heteroalkyl group containing from 1 to 4 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and/or N (especially O and/or N).

[0068] Further preferably, the expression heteroalkyl refers to an alkyl group as defined above (straight-chain or branched) in which one or more (preferably 1 to 6; especially preferably 1, 2, 3 or 4) carbon atoms have been replaced by an oxygen, sulfur or nitrogen atom or a CO group or a SO group or a SO₂ group; this group preferably contains from 1 to 6 (e.g. 1, 2, 3 or 4) carbon atoms and 1, 2, 3 or 4 (especially 1, 2 or 3) heteroatoms selected from oxygen, nitrogen and sulfur (especially oxygen and nitrogen); this group may preferably be substituted by one or more (preferably 1 to 6; especially preferably 1, 2, 3 or 4) fluorine, chlorine, bromine or iodine atoms or OH, =O, SH, =S, NH₂, =NH, N₃, CN or NO₂ groups.

[0069] Examples of heteroalkyl groups are groups of formulae: R^a-O-Y^a-, R^a-S-Y^a-, R^a-SO-Y^a-, R^a-SO₂-Y^a-, R^a-N(R^b)-SO₂-Y^a-, R^a-SO₂-N(R^b)-Y^a-, R^a-N(R^b)-Y^a-, R^a-CO-Y^a-, R^a-O-CO-Y^a-, R^a-CO-O-Y^a-, R^a-CO-N(R^b)-Y^a-, R^a-N(R^b)-CO-Y^a-, R^a-O-CO-N(R^b)-Y^a-, R^a-N(R^b)-CO-O-Y^a-, R^a-N(R^b)-CO-N(R^c)-Y^a-, R^a-O-CO-O-Y^a-, R^a-N(R^b)-C(=NR^d)-N(R^c)-Y^a-, R^a-CS-Y^a-, R^a-O-CS-Y^a-, R^a-CS-O-Y^a-, R^a-CS-N(R^b)-Y^a-, R^a-N(R^b)-CS-Y^a-, R^a-O-CS-N(R^b)-Y^a-, R^a-N(R^b)-CS-O-Y^a-, R^a-N(R^b)-CS-N(R^c)-Y^a-, R^a-O-CS-O-Y^a-, R^a-S-CO-Y^a-, R^a-CO-S-Y^a-, R^a-S-CO-N(R^b)-Y^a-, R^a-N(R^b)-CO-S-Y^a-, R^a-S-CO-O-Y^a-, R^a-O-CO-S-Y^a-, R^a-S-CO-S-Y^a-, R^a-S-CS-Y^a-, R^a-CS-S-Y^a-, R^a-S-CS-N(R^b)-Y^a-, R^a-N(R^b)-CS-S-Y^a-, R^a-S-CS-O-Y^a-, R^a-O-CS-S-Y^a-, wherein R^a being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^b being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^c being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group; R^d being a hydrogen atom, a C₁-C₆ alkyl, a C₂-C₆ alkenyl or a C₂-C₆ alkynyl group and Y^a being a bond, a C₁-C₆ alkylene, a C₂-C₆ alkenylene or a C₂-C₆ alkynylene group, wherein each heteroalkyl group contains at least one carbon atom and one or more hydrogen atoms may be replaced by fluorine or chlorine atoms.

[0070] Specific examples of heteroalkyl groups are methoxy, trifluoromethoxy, ethoxy, n-propyloxy, iso-propyloxy, n-butoxy, tert-butyloxy, methoxymethyl, ethoxymethyl, -CH₂CH₂OH, -CH₂OH, -SO₂Me, -NHAc, methoxyethyl, 1-methoxyethyl, 1-ethoxyethyl, 2-methoxy-

ethyl or 2-ethoxyethyl, methylamino, ethylamino, propylamino, isopropylamino, dimethylamino, diethylamino, isopropylethylamino, methylamino methyl, ethylamino methyl, diisopropylamino ethyl, methylthio, ethylthio, isopropylthio, enol ether, dimethylamino methyl, dimethylamino ethyl, acetyl, propionyl, butyryloxy, acetyloxy, methoxycarbonyl, ethoxycarbonyl, propionyloxy, acetylamino or propionylamino, carboxymethyl, carboxyethyl or carboxypropyl, N-ethyl-N-methylcarbamoyl or N-methyl-carbamoyl. Further examples of heteroalkyl groups are nitrile (-CN), isonitrile, cyanate, thiocyanate, isocyanate, isothiocyanate and alkylnitrile groups.

[0071] The expression cycloalkyl refers to a saturated or partially unsaturated (for example, a cycloalkenyl group) cyclic group that contains one or more rings (preferably 1 or 2), and contains from 3 to 14 ring carbon atoms, preferably from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms. The expression cycloalkyl refers furthermore to groups in which one or more hydrogen atoms have been replaced by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups, thus, for example, cyclic ketones such as, for example, cyclohexanone, 2-cyclohexenone or cyclopentanone. Further specific examples of cycloalkyl groups are a cyclopropyl, cyclobutyl, cyclopentyl, spiro[4,5]decanyl, norbornyl, cyclohexyl, cyclopentenyl, cyclohexadienyl, decalyl, bicyclo[4.3.0]nonyl, tetraline, cyclopentylcyclohexyl, fluorocyclohexyl or cyclohex-2-enyl group. Preferably, the expression cycloalkyl refers to a saturated cyclic group that contains one or more rings (preferably 1 or 2), and contains from 3 to 14 ring carbon atoms, preferably from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms.

[0072] The expression heterocycloalkyl refers to a cycloalkyl group as defined above in which one or more (preferably 1, 2 or 3) ring carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom) or a SO group or a SO₂ group. A heterocycloalkyl group has preferably 1 or 2 ring(s) and 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms (preferably selected from C, O, N and S). The expression heterocycloalkyl refers furthermore to groups that are substituted by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups. Examples are a piperidyl, prolinyl, imidazolidinyl, piperazinyl, morpholinyl (e.g. -N(CH₂CH₂)₂O), urotropinyl, pyrrolidinyl, tetrahydrothiophenyl, tetrahydropyranyl, tetrahydrofuryl or 2-pyrazolinyl group and also lactams, lactones, cyclic imides and cyclic anhydrides.

[0073] The expression alkylcycloalkyl refers to groups that contain both cycloalkyl and alkyl, alkenyl or alkynyl groups in accordance with the above definitions, for example alkylcycloalkyl, cycloalkylalkyl, alkylcycloalkenyl, alkenylcycloalkyl and alkynylcycloalkyl groups. An alkylcycloalkyl group preferably contains a cycloalkyl group that contains one or two rings and from 3 to 10 (especially 3, 4, 5, 6 or 7) ring carbon atoms, and one or two alkyl, alkenyl or alkynyl groups (especially alkyl groups) having 1 or 2 to 6 carbon atoms.

[0074] The expression heteroalkylcycloalkyl refers to alkylcycloalkyl groups as defined above in which one or more (preferably 1, 2 or 3) carbon atoms have been replaced by an oxygen, nitrogen, silicon, selenium, phosphorus or sulfur atom (preferably by an oxygen, sulfur or nitrogen atom) or a SO group or a SO₂ group. A heteroalkylcycloalkyl

group preferably contains 1 or 2 rings having from 3 to 10 (especially 3, 4, 5, 6 or 7) ring atoms, and one or two alkyl, alkenyl, alkynyl or heteroalkyl groups (especially alkyl or heteroalkyl groups) having from 1 or 2 to 6 carbon atoms. Examples of such groups are alkylheterocycloalkyl, alkylheterocycloalkenyl, alkenylheterocycloalkyl, alkynylheterocycloalkyl, heteroalkylcycloalkyl, heteroalkylheterocycloalkyl and heteroalkylheterocycloalkenyl, the cyclic groups being saturated or mono-, di- or tri-unsaturated.

[0075] The expression aryl refers to an aromatic group that contains one or more rings and from 6 to 14 ring carbon atoms, preferably from 6 to 10 (especially 6) ring carbon atoms. The expression aryl refers furthermore to groups that are substituted by fluorine, chlorine, bromine or iodine atoms or by OH, SH, NH₂, N₃ or NO₂ groups. Examples are the phenyl (Ph), naphthyl, biphenyl, 2-fluorophenyl, anil-nyl, 3-nitrophenyl or 4-hydroxyphenyl group.

[0076] The expression heteroaryl refers to an aromatic group that contains one or more rings and from 5 to 14 ring atoms, preferably from 5 to 10 (especially 5 or 6 or 9 or 10) ring atoms, comprising one or more (preferably 1, 2, 3 or 4) oxygen, nitrogen, phosphorus or sulfur ring atoms (preferably O, S or N). The expression heteroaryl refers furthermore to groups that are substituted by fluorine, chlorine, bromine or iodine atoms or by OH, SH, N₃, NH₂ or NO₂ groups. Examples are pyridyl (e.g. 4-pyridyl), imidazolyl (e.g. 2-imidazolyl), phenylpyrrolyl (e.g. 3-phenylpyrrolyl), thiazolyl, isothiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, oxadiazolyl, thiadiazolyl, indolyl, indazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, 4-hydroxypyridyl (4-pyridonyl), 3,4-hydroxypyridyl (3,4-pyridonyl), oxazolyl, isoxazolyl, triazolyl, tetrazolyl, isoxazolyl, indazolyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzthiazolyl, pyridazinyl, quinolinyl, isoquinolinyl, pyrrolyl, purinyl, carbazolyl, acridinyl, pyrimidyl, 2,3'-bifuryl, pyrazolyl (e.g. 3-pyrazolyl) and isoquinolinyl groups.

[0077] The expression aralkyl refers to groups containing both aryl and also alkyl, alkenyl, alkynyl and/or cycloalkyl groups in accordance with the above definitions, such as, for example, arylalkyl, arylalkenyl, arylalkynyl, arylcycloalkyl, arylcycloalkenyl, alkylaryl cycloalkyl and alkylaryl cycloalkenyl groups. Specific examples of aralkyls are phenylcyclopentyl, cyclohexylphenyl as well as groups derived from toluene, xylene, mesitylene, styrene, benzyl chloride, o-fluorotoluene, 1H-indene, tetraline, dihydronaphthalene, indanone, cumene, fluorene and indane. An aralkyl group preferably contains one or two aromatic ring systems (especially 1 or 2 rings), each containing from 6 to 10 carbon atoms and one or two alkyl, alkenyl and/or alkynyl groups containing from 1 or 2 to 6 carbon atoms and/or a cycloalkyl group containing 3, 4, 5, 6 or 7 ring carbon atoms.

[0078] The expression heteroaralkyl refers to groups containing both aryl and/or heteroaryl groups and also alkyl, alkenyl, alkynyl and/or heteroalkyl and/or cycloalkyl and/or heterocycloalkyl groups in accordance with the above definitions. A heteroaralkyl group preferably contains one or two aromatic ring systems (especially 1 or 2 rings), each containing from 5 or 6 to 9 or 10 ring atoms (preferably selected from C, N, O and S) and one or two alkyl, alkenyl and/or alkynyl groups containing 1 or 2 to 6 carbon atoms and/or one or two heteroalkyl groups containing 1 to 6 carbon atoms and 1, 2 or 3 heteroatoms selected from O, S and N and/or one or two cycloalkyl groups each containing 3, 4, 5, 6 or 7 ring carbon atoms and/or one or two

heterocycloalkyl groups, each containing 3, 4, 5, 6 or 7 ring atoms comprising 1, 2, 3 or 4 oxygen, sulfur or nitrogen atoms.

[0079] Examples are arylheteroalkyl, arylheterocycloalkyl, arylheterocycloalkenyl, arylalkylheterocycloalkyl, arylalkenylheterocycloalkyl, arylalkynylheterocycloalkyl, arylalkylheterocycloalkenyl, heteroarylalkyl, heteroarylalkenyl, heteroarylalkynyl, heteroarylheteroalkyl, heteroarylcycloalkyl, heteroarylcycloalkenyl, heteroarylheterocycloalkyl, heteroarylheterocycloalkenyl, heteroarylalkylcycloalkyl, heteroaryl-alkylheterocycloalkenyl, heteroarylheteroalkylcycloalkyl, heteroarylheteroalkylcycloalkenyl and heteroarylheteroalkylheterocycloalkyl groups, the cyclic groups being saturated or mono-, di- or tri-unsaturated. Specific examples are a tetrahydroisoquinolinyl, benzoyl, phthalidyl, 2- or 3-ethylindolyl, 4-methylpyridino, 2-, 3- or 4-methoxyphenyl, 4-ethoxyphenyl, 2-, 3- or 4-carboxyphenylalkyl group.

[0080] As already stated above, the expressions cycloalkyl, heterocycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, aryl, heteroaryl, aralkyl and heteroaralkyl also refer to groups that are substituted by fluorine, chlorine, bromine or iodine atoms or by OH, =O, SH, =S, NH₂, =NH, N₃ or NO₂ groups.

[0081] The term halogen refers to F, Cl, Br or I. Preferred halogens are F, Cl and Br. Especially preferred halogens are F and Cl.

[0082] The term “optionally substituted” refers to a group, which is unsubstituted or substituted by one or more (especially by one, two or three; preferably by one or two) substituents.

[0083] If a group comprises more than one substituent, these substituents are independently selected, i.e., they may be the same or different.

[0084] Examples for substituents are fluorine, chlorine, bromine and iodine and OH, SH, NH₂, —SO₃H, —SO₂NH₂, —COOH, —COOMe, —COMe (Ac), —NHSO₂Me, —SO₂NMe₂, —CH₂NH₂, —NHAc, —SO₂Me, —CONH₂, —CN, —NHCONH₂, —NHC(NH)NH₂, —NOHCH₃, —N₃ and —NO₂ groups. Further examples of substituents are C₁-C₁₀ alkyl, C₂-C₁a alkenyl, C₂-C₁a alkynyl, C₁-C₁₀ heteroalkyl, C₃-C₁₈ cycloalkyl, C₁-C₁₇ heterocycloalkyl, C₄-C₂₀ alkylcycloalkyl, C₁-C₁₉ heteroalkylcycloalkyl, C₆-C₁₈ aryl, C₁-C₁₇ heteroaryl, C₇-C₂₀ aralkyl and C₁-C₁₉ heteroaralkyl groups; especially C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₁-C₆ heteroalkyl, C₃-C₁₀ cycloalkyl, C₁-C₉ heterocycloalkyl, C₄-C₁₂ alkylcycloalkyl, C₁-C₁₁ heteroalkylcycloalkyl, C₆-C₁₀ aryl, C₁-C₉ heteroaryl, C₇-C₁₂ aralkyl and C₁-C₁₁ heteroaralkyl groups, further preferably C₁-C₆ alkyl and C₁-C₆ heteroalkyl groups.

[0085] When an aryl, heteroaryl, cycloalkyl, alkylcycloalkyl, heteroalkylcycloalkyl, heterocycloalkyl, aralkyl or heteroaralkyl group contains more than one ring, these rings may be bonded to each other via a single or double bond or these rings may be annulated or fused or bridged.

[0086] Owing to their substitution, the compounds of the present invention may contain one or more centers of chirality. The present invention therefore includes both all pure enantiomers and all pure diastereomers and also mixtures thereof in any mixing ratio. The present invention moreover also includes all cis/trans-isomers of the compounds of the present invention and also mixtures thereof.

The present invention moreover includes all tautomeric forms of the compounds of the present invention.

[0087] The present invention further provides pharmaceutical compositions comprising a compound or a salt thereof according to the present invention and optionally one or more carrier substances and/or one or more adjuvants and/or one or more further active pharmaceutical ingredient(s).

[0088] The present invention furthermore provides compounds (or a salt thereof) or pharmaceutical compositions as described herein for use as a medicament, e.g. for use in the prophylaxis or treatment of a bacterial infection; especially for use in the prophylaxis or treatment of a bacterial infection caused by Gram-negative bacteria.

[0089] The compounds and/or compositions described herein can also be useful in the prophylaxis treatment of infections by bacteria, which are susceptible or multidrug resistant, polymyxin resistant mutant, carbapenam-resistant bacteria, methicillin-resistant *Staphylococcus aureus*, vancomycin-resistant *Enterococci* or multi-drug resistant *Neisseria gonorrhoeae*.

[0090] Examples of Gram-negative bacteria include, but are not limited to, *Escherichia coli*, *Pseudomonas aeruginosa*, *Candidatus liberibacter*, *Agrobacterium tumefaciens*, *Branhamella catarrhalis*, *Citrobacter di versus*, *Enterobacter aerogenes*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Salmonella typhimurium*, *Neisseria meningitidis*, *Serratia marcescens*, *Shigella sonnei*, *Shigella boydii*, *Neisseria gonorrhoeae*, *Acinetobacter baumannii*, *Salmonella enteritidis*, *Fusobacterium nucleatum*, *Veillonella parvula*, *Actinobacillus actinomycetemcomitans*, *Aggregatibacter actinomycetemcomitans*, *Porphyromonas gingivalis*, *Helicobacter pylori*, *Francisella tularensis*, *Yersinia pestis*, *Vibrio cholera*, *Morganella morganii*, *Ectwardsiella tarda*, *Campylobacter jejuni*, or *Haemophilus influenza*, *Enterobacter cloacae* and numerous others. Other notable groups of Gram-negative bacteria include the cyanobacteria, spirochaetes, green sulfur and green non-sulfur bacteria.

[0091] The present invention further provides a compound (or a salt thereof) as described herein or a pharmaceutical composition as defined herein for the preparation of a medicament for use in the prophylaxis or treatment of a bacterial infection; especially for use in the prophylaxis or treatment of a bacterial infection caused by Gram-negative bacteria.

[0092] Examples of salts of sufficiently basic compounds are salts of physiologically acceptable mineral acids like hydrochloric, hydrobromic, sulfuric and phosphoric acid; or salts of organic acids like methanesulfonic, p-toluenesulfonic, lactic, acetic, trifluoroacetic, citric, succinic, fumaric, maleic and salicylic acid. Further, a sufficiently acidic compound may form alkali or earth alkali metal salts, for example sodium, potassium, lithium, calcium or magnesium salts; ammonium salts; or organic base salts, for example methylamine, dimethylamine, trimethylamine, triethylamine, ethylenediamine, ethanolamine, choline hydroxide, meglumin, piperidine, morpholine, tris-(2-hydroxyethyl) amine, lysine or arginine salts; all of which are also further examples of salts of the compounds described herein.

[0093] The compounds described herein may be solvated, especially hydrated. The solvation/hydration may occur during the process of production or as a consequence of the hygroscopic nature of the initially water-free compounds. The solvates and/or hydrates may e.g. be present in solid or liquid form.

[0094] The therapeutic use of the compounds described herein, their salts, as well as formulations and pharmaceutical compositions also lie within the scope of the present invention.

[0095] In general, the compounds and pharmaceutical compositions described herein will be administered by using the established and acceptable modes known in the art.

[0096] For oral administration, such therapeutically useful agents can be administered by one of the following routes: oral, e.g. as tablets, dragees, coated tablets, pills, semisolids, soft or hard capsules, for example soft and hard gelatine capsules, aqueous or oily solutions, emulsions, suspensions or syrups, parenteral including intravenous, intramuscular and subcutaneous injection, e.g. as an injectable solution or suspension, rectal as suppositories, by inhalation or insufflation, e.g. as a powder formulation, as microcrystals or as a spray (e.g. liquid aerosol), transdermal, for example via an transdermal drug delivery system (TDDS) such as a plaster containing the active ingredient or intranasal. For the production of such tablets, pills, semisolids, coated tablets, dragees and hard, e.g. gelatine, capsules the therapeutically useful product may be mixed with pharmaceutically inert, inorganic or organic excipients as are e.g. lactose, sucrose, glucose, gelatine, malt, silica gel, starch or derivatives thereof, talc, stearic acid or their salts, dried skim milk, and the like. For the production of soft capsules, one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat, and polyols. For the production of liquid solutions, emulsions or suspensions or syrups one may use as excipients e.g. water, alcohols, aqueous saline, aqueous dextrose, polyols, glycerin, lipids, phospholipids, cyclodextrins, vegetable, petroleum, animal or synthetic oils. Especially preferred are lipids and more preferred are phospholipids (preferred of natural origin, especially preferred with a particle size between 300 to 350 nm) preferred in phosphate buffered saline (pH=7 to 8, preferred 7.4). For suppositories, one may use excipients as are e.g. vegetable, petroleum, animal or synthetic oils, wax, fat and polyols. For aerosol formulations, one may use compressed gases suitable for this purpose, e.g. oxygen, nitrogen and carbon dioxide. The pharmaceutically useful agents may also contain additives for conservation, stabilization, e.g. UV stabilizers, emulsifiers, sweetener, aromatizers, salts to change the osmotic pressure, buffers, coating additives and antioxidants.

[0097] In general, in the case of oral or parenteral administration to adult humans weighing approximately 80 kg, a daily dosage of about 1 mg to about 10,000 mg, preferably from about 5 mg to about 1,000 mg, should be appropriate, although the upper limit may be exceeded when indicated. The daily dosage can be administered as a single dose or in divided doses, or for parenteral administration, it may be given as continuous infusion or subcutaneous injection.

[0098] According to a moreover preferred embodiment, the present invention provides a method for treating a bacterial infection, which comprises administering to a subject in need of such treatment a therapeutically effective amount of a compound as described herein, or a salt thereof.

[0099] According to a further preferred embodiment, the present invention provides a method of treating, ameliorating, or preventing a bacterial infection in a subject in need thereof, comprising administering to the subject a therapeutically effective amount of at least one compound described herein, a salt thereof, or a pharmaceutical composition

comprising at least one of the (specific) compounds, or a salt thereof, described herein. Administration of the compound may be topical, such as subcutaneous, transdermal, rectal, intravaginal, intranasal, intrabronchial, intraocular, or intraural. Alternatively, administration may be systemic, such as oral administration. In still other alternatives, administration may be parenteral, intravenous, intramuscular, or intraperitoneal.

[0100] As used herein, the term “administration” can also include administering a combination of compounds. Thus, administration may be in the form of dosing an organism with a compound or combination of compounds, such that the organism’s circulatory system will deliver a compound or combination of compounds to the target area, including but not limited to a cell or cells, synaptic junctions and circulation. Administration may also mean that a compound or combination of compounds is placed in direct contact with an organ, tissue, area, region, cell or group of cells, such as but not limited to direct injection of the combination of compounds.

[0101] In select embodiments, a combination of compounds can be administered, and thus the individual compounds can also be said to be co-administered with one another. As used herein, “co-administer” indicates that each of at least two compounds is administered during a time frame wherein the respective periods of biological activity or effects overlap. Thus, the term co-administer includes sequential as well as coextensive administration of the individual compounds, at least one of which is a compound of the present invention. Accordingly, “administering” a combination of compounds according to some of the methods of the present invention includes sequential as well as coextensive administration of the individual compounds of the present invention. Likewise, the phrase “combination of compounds” indicates that the individual compounds are co-administered, and the phrase “combination of compounds” does not mean that the compounds must necessarily be administered contemporaneously or coextensively. In addition, the routes of administration of the individual compounds need not be the same.

[0102] The invention also relates to a combination preparation or pharmaceutical composition containing at least one compound according to the invention, or a salt thereof, and at least one further (different) active pharmaceutical ingredient. The combination preparation of the invention can be used as a medicament, in particular in the treatment or prophylaxis of a bacterial infection with Gram-negative bacteria or Gram-negative and -positive bacteria.

[0103] Preferably, in the combination preparation or pharmaceutical composition of the invention the further active pharmaceutical ingredient is another antibiotic. The other antibiotic can be selected from the group consisting of β -lactam antibiotics, including penams, carbapenams, oxapenams, penems, carbapenems, monobactams, cepheams, carbacephems, oxacephems, and monobactams; aminoglycoside antibiotics, including amikacin, arbekacin, astromycin, bekanamycin, dibekacin, framycetin, gentamicin, hygromycin B, isepamicin, kanamycin, neomycin, netilmicin, paromomycin, paromomycin sulfate, ribostamycin, sisomicin, spectinomycin, streptomycin, tobramycin, and verdamycin; quinolone antibiotics, including ciprofloxacin, enoxacin, gatifloxacin, grepafloxacin, levofloxacin, lomefloxacin, moxifloxacin, nalidixic acid, norfloxacin, ofloxacin, sparfloxacin, temafloxacin, and trovafloxacin; or glycopeptide antibiotics, e.g. vancomycin, telavancin, bleomycin, ramoplanin, and decaplanin; linezolid; or daptomycin.

[0104] As used herein, the terms “treat” and “treatment” refer to a slowing of or a reversal of the progress of the disease or infection. Treating a disease includes treating a symptom and/or reducing the symptoms of the disease or infection. The terms “prophylaxis” and “preventing” refer to a slowing of the disease or of the onset of the disease, infection or the symptoms thereof. Prophylaxis of, or preventing, a disease or infection can include stopping the onset of the disease, infection or symptom thereof.


[0105] As used herein, the term “subject” may be an animal, vertebrate animal, mammal, rodent (e.g., a guinea pig, a hamster, a rat, a mouse), a murine (e.g., a mouse), a canine (e.g., a dog), a feline (e.g., a cat), an equine (e.g., a horse), a primate, a simian (e.g., a monkey or ape), a monkey (e.g., marmoset, a baboon), an ape (e.g., gorilla, chimpanzee, orangutan, gibbon), or a human.

[0106] As used herein, the term “dosage unit” refers to a physically discrete unit, such as a capsule or tablet suitable as a unitary dosage for a subject. Each unit contains a predetermined quantity of a compound of the invention, or a salt thereof, which was discovered or believed to produce the desired pharmacokinetic profile, which yields the desired therapeutic effect. The dosage unit is composed of a compound of the present invention, or a salt thereof, in association with at least one pharmaceutically acceptable carrier, salt, excipient or a combination thereof. The term “dose” or “dosage” refers to the amount of active ingredient that an individual takes or is administered at one time.

[0107] The term “therapeutically effective amount” refers to the amount sufficient to produce a desired biological effect in a subject. Accordingly, a therapeutically effective amount of a compound may be an amount which is sufficient to treat or prevent a disease or infection, and/or delay the onset or progression of a disease or infection, and/or alleviate one or more symptoms of the disease or infection, when administered to a subject suffered from or susceptible to that disease or infection. A “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” herein refers to a non-API (where API refers to Active Pharmaceutical Ingredient) substances such as disintegrators, binders, fillers, and lubricants used in formulating pharmaceutical products. They are generally safe for administering to humans.

[0108] The term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term “vehicle” refers to a diluent, adjuvant, excipient, or carrier with which a compound of the invention is administered. Such pharmaceutical vehicles can be liquids, such as water and oils, and the like.

[0109] In a further aspect, the present invention provides a method for the production of a compound of formula (Ic), or a compound of formula (Ic') or (Ic''):



(Ic)

[0117] The culturing step can, for example, be performed in liquid culture, by growing the respective recombinant host (e.g., microorganism, e.g. bacterial host cell) in media containing one or several different carbon sources, and one or different nitrogen sources. Also salts are essential for growth and production. Suitable carbon sources are different mono-, di-, and polysaccharides like maltose, glucose or carbon from amino acids like peptones. Nitrogen sources are ammonium, nitrate, urea, chitin or nitrogen from amino acids. The following inorganic ions support the growth or are essential in synthetic media: Mg-ions, Ca-ions, Fe-ions, Mn-ions, Zn-ions, K-ions, sulfate-ions, Cl-ions, phosphate-ions. Preferred compositions of nutrient media used in the production method of the invention are described in more detail in the examples. In an embodiment, the recombinant host is a microorganism, e.g., a bacterium such as a bacterium of the species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*), or a yeast such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*. In one embodiment, the recombinant host can be a bacterial host cell, e.g. an *Escherichia coli* or a *Lactobacillus* cell, preferably an *E. coli* cell, and more preferably an *E. coli* BL21(DE3) cell. Preferred embodiments include *E. coli* BL21 (DE3) pNOSodarA-E-D9 (deposited at DSMZ on 29 Jan. 2021; DSM 33798) and *E. coli* BL21 (DE3) pNOSodarA-E-D17 (deposited at DSMZ on 29 Jan. 2021; DSM 33799). Temperatures for growth and production are between 15° C. to 40° C., preferred temperatures are between 25° C. and 35° C., especially at 30° C. The pH of the culture solution is from 5 to 8, preferably a pH of 6.9 to 7.2, more preferable a pH of about 7.1.

[0118] The compound may be recovered from the fermentation broth by resin absorption and eluted from the resin by washing with solvents of various polarities. Purification may be furthered by chromatographic separation such as high performance liquid chromatography (HPLC) or reverse-phase high performance liquid chromatography (RP-HPLC).

[0119] Compounds of formula (Ic), (Ic') or (Ic''), i.e. darobactin A and darobactin derivatives, are ribosomally synthesized and post-translationally modified peptide compounds (RiPPs). The artificial synthetic or recombinant biosynthetic gene cluster (BGC) encoding said peptide compounds comprises (consists of) genes with the following predicted functions: darA or modified variant thereof: darobactin propeptide or propetide of darobactin derivative; darBCD: ABC-type trans envelope exporter; and darE: radical S-adenosyl methionine (SAM) methyltransferase (proposed to catalyze the cyclization reactions to connect W₃-K₅ and W₁-W₃). The bacterial host cell comprises at least one synthetic or recombinant nucleic acid sequence, which encodes such a synthetic or recombinant BGC of a compound of formula (Ic), (Ic') or (Ic''), and thus being capable of synthesizing a compound according to formula (Ic), (Ic') or (Ic''). The synthetic or recombinant nucleic acid sequence utilized in the production method of the present

invention, may, in addition to the nucleic acid sequence encoding the BGC, also include regulatory sequences, such as promoter and translation initiation and termination sequences, and can further include sequences that facilitate stable maintenance in a host cell, i.e., sequences that provide the function of an origin of replication or facilitate integration into host cell chromosomal or other DNA by homologous recombination. The synthetic or recombinant nucleic acid sequence utilized in the production method of the present invention is a synthetic or recombinant nucleic acid sequence, which encodes a BGC of a compound of formula (Ic) (or of formula (Ic'), or of formula (Ic'')), and has:




[0120] (i) a sequence identity to the full-length sequence of SEQ ID NO. 1 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%, or

[0121] (ii) a sequence completely complementary to any nucleic acid sequence of (i).



[0122] The compounds of the invention described herein, i.e. darobactin derivatives, are ribosomally synthesized and post-translationally modified peptide compounds (RiPPs).

[0123] These peptide compounds can be produced using any method. For example, the compounds can be produced by chemical synthesis. Alternatively, peptide compounds described herein can be produced by standard recombinant technology using heterologous expression vectors encoding polypeptides. Expression vectors can be introduced into host cells, e.g., by transformation or transfection, for expression of the encoded polypeptide, which then can be purified. Expression systems that can be used for small or large scale production of polypeptides include, without limitation, microorganisms such as bacteria (e.g., species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*) transformed with recombinant bacteriophage DNA, plasmid DNA, or cosmid DNA expression vectors containing the nucleic acid molecules described herein, and yeast (e.g., *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*) transformed with recombinant yeast expression vectors containing the nucleic acid molecules described herein. Useful expression systems also include insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus) containing the nucleic acid molecules described herein, and plant cell systems infected with recombinant virus expression vectors (e.g., tobacco mosaic virus) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing the nucleic acid molecules described herein. Peptide compounds of this invention also can be produced using mammalian expression system harboring recombinant expression constructs containing promoters derived from the genome of mammalian cells (e.g., the metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter and the cytomegalovirus promoter), along with the nucleic acids described herein. Peptide compounds of this invention can have an N-terminal or C-terminal tag.


[0124] In one embodiment, described herein is a method for the production of a compound of formula (I'), (Ib) or (Ib'):



wherein each of R^{22} and R^{44} independently of one another, represents H, CH_3 , CH_2OH or a group:



-continued



wherein the method comprises the steps of:

[0125] (a) providing a recombinant host capable of producing said compound of formula (I'), (1b) or (1b'), wherein said recombinant host harbors at least one

synthetic or recombinant nucleic acid encoding a bio-synthetic gene cluster (BGC) of the compound, which BGC has:




[0126] (i) a sequence identity to the full-length sequence of SEQ ID NO. 44 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or


[0127] (ii) a sequence completely complementary to any nucleic acid sequence of (i);

[0128] (b) cultivating said recombinant host for a time sufficient for said recombinant host to produce the compound of formula (I'), (1b) or (1b');

[0129] (c) isolating the compound from said recombinant host or from cultivation supernatant, thereby producing the compound of formula (I'), (1b) or (1b').


[0130] In one embodiment, described herein is a method for the production of a compound of formula (II), (IIa) or (IIa'):






wherein


[0131] R^2 and R^4 each independently represents H, CH_3 , CH_2OH or a group:




[0132] R^3 is a group



[0133] R^5 is a group of the following formula:



[0134] wherein R^9 is a hydrogen or halogen atom; R^{10} is a hydrogen atom or a methyl group; R^{11} is, at each occasion, independently selected from a halogen atom (especially fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0); or a group of formula $-CH_3$; or a group of the following formula:



and

[0135] wherein the method comprises the steps of:

[0136] (a) providing a recombinant host capable of producing said compound of formula (II) (or (IIa), (IIa') or (IIa'')), wherein said recombinant host harbors at least one synthetic or recombinant nucleic acid encoding a biosynthetic gene cluster (BGC) of the compound, which BGC has:


[0137] (i) a sequence identity to the full-length sequence of SEQ ID NO. 86 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or

[0138] (ii) a sequence completely complementary to any nucleic acid sequence of (i);

[0139] (b) cultivating said recombinant host for a time sufficient for said recombinant host to produce the compound of formula (II) (or (IIa), (IIa') or (IIa''));


[0140] (c) isolating the compound from said recombinant host or from cultivation supernatant, thereby producing the compound of formula (II) (or (IIa), (IIa') or (IIa'')).

[0141] In some embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^5 is a group of the following formula:




[0142] wherein R^9 is a hydrogen atom; R^{10} is a hydrogen atom; R^{11} is, at each occasion, independently selected from a halogen atom (especially fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).


[0143] In some embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^2 is a group of the following formula:




[0144] In some embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^4 is a group of the following formula:




[0145] In some embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^3 is a group:



[0146] In some embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^3 is a group:



[0147] In some preferred embodiments of the compound of formula (II), (IIa), (IIa') or (IIa''), R^3 is a group:



[0148] In some embodiments, the production method of the compound of formula (II), (IIa), (IIa') or (IIa'') optionally comprises feeding of one or more halogenated tryptophan(s), preferably halogenated L-tryptophan(s), including, for example, commercially available 5-chloro-L-tryptophan, 7-chloro-L-tryptophan, 6-fluoro-L-tryptophan or 7-fluor-L-tryptophan, to the fermentation broth during cultivation. The one or more halogenated tryptophan(s) can be fed periodically in different concentrations known to the person skilled in the art, and total end concentration of the halogenated tryptophan(s) can, for example, be between 0.5 mM and 10 mM.

[0149] In some embodiments, the production method of the compound of formula (II), (IIa), (IIa') or (IIa'') optionally comprises feeding of one or more halogenated tryptophan(s), preferably halogenated L-tryptophan(s), including, for example, commercially available 5-chloro-L-tryptophan, 7-chloro-L-tryptophan, 6-fluoro-L-tryptophan or 7-fluor-L-tryptophan, to the fermentation broth during cultivation. The one or more halogenated tryptophan(s) can be fed periodically in different concentrations known to the person skilled in the art, and total end concentration of the halogenated tryptophan(s) can, for example, be between 0.5 mM and 10 mM.

[0150] In some embodiments, the production method of the compound of formula (II), (IIa), (IIa') or (IIa'') optionally comprises the cultivation of a host cell which contains vector pUC18-zeo-mx8-corP-trpAB described herein, and the combined feeding of serine (preferably L-serine) and one or more halogenated indol(s) to the fermentation broth during cultivation. In some embodiment, the production method of the compound of formula (II), (IIa), (IIa') or (IIa'') optionally comprises the cultivation of a recombinant host which contains at least one synthetic or recombinant nucleic

acid encoding a biosynthetic gene cluster (BGC) of the compound as described herein, wherein said BGC further includes the *trpAB* gene of pSTB7 37845 upstream of the *darE* gene in the BGC, and the feeding of serine (preferably L-serine) and one or more halogenated indol(s) to the fermentation broth during cultivation of the recombinant host. The serine and the one or more halogenated indol(s) can be fed periodically in different concentrations known to the person skilled in the art. For example, serine may be fed in a concentration of 0.5 mM and 15 mM (preferably 8-12 mM, especially 10 mM) and the halogenated indol may be fed in a concentration of 5-10 mM.

[0151] As used herein, the term “recombinant host” is intended to refer to a host, the genome of which has been augmented by at least one incorporated DNA sequence. Such DNA sequences include but are not limited to genes that are not naturally present, DNA sequences that are not normally transcribed into RNA or translated into a protein (“expressed”), and other genes or DNA sequences which one desires to introduce into the non-recombinant host. It will be appreciated that typically the genome of a recombinant host described herein is augmented through the stable introduction of one or more recombinant genes. However, autonomous or replicative plasmids or vectors can also be used within the scope of this invention. Moreover, the present invention can be practiced using a low copy number, e.g., a single copy, or high copy number (as exemplified herein) plasmid or vector. “Introduced,” or “augmented” as used herein, are known in the art to mean introduced or augmented by the hand of man.

[0152] Generally, the introduced DNA is not originally resident in the host that is the recipient of the DNA, but it is within the scope of the invention to isolate a DNA segment from a given host, and to subsequently introduce one or more additional copies of that DNA into the same host, e.g., to enhance production of the product of a gene or alter the expression pattern of a gene. In some instances, the introduced DNA will modify or even replace an endogenous gene or DNA sequence by, e.g., homologous recombination or site-directed mutagenesis. Suitable recombinant hosts include microorganisms such as bacteria (e.g., species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*), or a yeast such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*), as well as mammalian, insect or plant cell expression systems.

[0153] A synthetic or recombinant nucleic acid encoding a biosynthetic gene cluster (BGC) of the peptide compound described herein includes the coding sequence for that polypeptide, operably linked, in sense orientation, to one or more regulatory regions suitable for expressing the polypeptide. Because many microorganisms are capable of expressing multiple gene products from a polycistronic mRNA, multiple polypeptides can be expressed under the control of a single regulatory region for those microorganisms, if desired. A coding sequence and a regulatory region are considered to be operably linked when the regulatory region and coding sequence are positioned so that the regulatory region is effective for regulating transcription or translation of the sequence. Typically, the translation initiation site of the translational reading frame of the coding

sequence is positioned between one and about fifty nucleotides downstream of the regulatory region for a monocistronic gene.

[0154] “Regulatory region” refers to a nucleic acid having nucleotide sequences that influence transcription or translation initiation and rate, and stability and/or mobility of a transcription or translation product. Regulatory regions include, without limitation, promoter sequences, enhancer sequences, response elements, protein recognition sites, inducible elements, protein binding sequences, 5' and 3' untranslated regions (UTRs), transcriptional start sites, termination sequences, polyadenylation sequences, introns, and combinations thereof. A regulatory region typically includes at least a core (basal) promoter. A regulatory region also may include at least one control element, such as an enhancer sequence, an upstream element or an upstream activation region (UAR). The choice of regulatory regions to be included depends upon several factors, including, but not limited to, efficiency, selectability, inducibility, desired expression level, and preferential expression during certain culture stages. It is a routine matter for one of skill in the art to modulate the expression of a coding sequence by appropriately selecting and positioning regulatory regions relative to the coding sequence. It will be understood that more than one regulatory region may be present, e.g., introns, enhancers, upstream activation regions, transcription terminators, and inducible elements.

[0155] In many cases, the coding sequence for a polypeptide described herein is identified in a species other than the recombinant host, i.e., is a heterologous nucleic acid. The term “heterologous nucleic acid” as used herein, refers to a nucleic acid introduced into a recombinant host, wherein said nucleic acid is not naturally present in said host. Thus, if the recombinant host is a microorganism, the coding sequence can be from other prokaryotic or eukaryotic microorganisms, from plants or from animals. In some case, however, the heterologous nucleic acid may contain a sequence (or parts thereof) that is native to the host and is being reintroduced into that organism. A native sequence can often be distinguished from the naturally occurring sequence by the presence of non-natural sequences linked to the exogenous nucleic acid, e.g., non-native regulatory sequences flanking a native sequence in a recombinant nucleic acid construct. In addition, stably transformed exogenous nucleic acids typically are integrated at positions other than the position where the native sequence is found.

[0156] It will be appreciated that because of the degeneracy of the genetic code, a number of nucleic acids can encode a particular peptide compound; i.e., for many amino acids, there is more than one nucleotide triplet that serves as the codon for the amino acid. Thus, codons in the coding sequence for a given peptide compound can be modified such that optimal expression in a particular host is obtained, using appropriate codon bias tables for that host (e.g., microorganism). As isolated nucleic acids, these modified sequences can exist as purified molecules and can be incorporated into a vector or a virus for use in constructing modules for recombinant nucleic acid constructs.

[0157] Recombinant hosts described herein express darobactin derivatives, e.g. peptide compounds of formula (I), (Ia) or (Ia') as described above. In one aspect, the present invention relates to a recombinant host harboring a heterologous nucleic acid encoding a darobactin derivative described herein. In particular, the recombinant host

described herein harbors a heterologous nucleic acid encoding a peptide compound of formula (I), (Ia) or (Ia'), wherein said heterologous nucleic acid is operably linked to a regulatory region allowing expression in said recombinant host. In an embodiment, the heterologous nucleic acid is an artificial synthetic or recombinant BGC encoding said peptide compounds. Such artificial synthetic or recombinant BGC comprises darA (modified) and darE, which genes have the following predicted functions: modified darA variant (i.e. propeptide of darobactin derivative described herein) and darE: radical S-adenosyl methionine (SAM) methyltransferase (proposed to catalyze the cyclization reactions to connect W₃-K₅ and W₁-W₃). In one embodiment, the artificial synthetic or recombinant BGC further comprises darBCD: having the (predicted) function of an ABC-type trans envelope exporter.

[0158] The recombinant host, e.g. microorganism such as bacterial host cell or yeast cell, comprises at least one synthetic or recombinant nucleic acid sequence, which encodes such a synthetic or recombinant BGC of a compound of formula (I), (Ia) or (Ia'), and thus being capable of synthesizing a compound according to formula (I), (Ia) or (Ia'). The synthetic or recombinant nucleic acid sequence utilized in the production method of the present invention, may, in addition to the nucleic acid sequence encoding the BGC, also include regulatory sequences, such as promoter and translation initiation and termination sequences, and can further include sequences that facilitate stable maintenance in a host cell, i.e., sequences that provide the function of an origin of replication or facilitate integration into host cell chromosomal or other DNA by homologous recombination. The synthetic or recombinant nucleic acid sequence utilized in the production method of the present invention is a synthetic or recombinant nucleic acid sequence, which encodes a BGC of a compound of formula (I) (or of formula (Ia), or of formula (Ia')), and has:

[0159] (i) a sequence identity to the full-length sequence of SEQ ID NO. 44 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%, or

[0160] (ii) a sequence completely complementary to any nucleic acid sequence of (i).

[0161] The term "identity" as used herein refers to a property of sequences that measures their similarity or relationship. Identity is measured by dividing the number of identical residues by the total number of residues and multiplying the product by 100.

[0162] The phrases "nucleic acid" or "nucleic acid sequence" as used herein refer to an oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA of genomic or synthetic origin, which may be single-stranded or double-stranded and may represent a sense or antisense strand, natural or synthetic in origin. "Oligonucleotide" includes either a single stranded polydeoxynucleotide or two complementary polydeoxynucleotide strands that may be chemically synthesized. Such synthetic oligonucleotides have no 5' phosphate and thus will not ligate to another oligonucleotide without adding a phosphate with an ATP in the presence of a kinase. A synthetic oligonucleotide can ligate to a fragment that has not been dephosphorylated. A "coding sequence" of or a "nucleotide sequence encoding" a particular polypeptide or protein, is a nucleic acid sequence which is transcribed and translated into a polypeptide or protein when placed under the control of appropriate regulatory sequences. The nucleic acids used to prac-

tice this invention may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed/generated recombinantly. Techniques for the manipulation of nucleic acids, such as, e.g., subcloning, labeling probes (e.g., random-primer labeling using Klenow polymerase, nick translation, amplification), sequencing, hybridization and the like are well described in the scientific and patent literature, see, e.g., Sambrook, et al, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, N. Y., Vols. 1-3, (1989); *Current Protocols in Molecular Biology*, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); *Laboratory Techniques In Biochemistry And Molecular Biology: Hybridization With Nucleic Acid Probes*, Part I. Theory and Nucleic Acid Preparation, Tijssen, ed. Elsevier, N.Y. (1993). A nucleic acid encoding a polypeptide of the invention is assembled in appropriate phase with a leader sequence capable of directing secretion of the translated polypeptide or fragment thereof.

[0163] The term "isolated" as used herein means that the material, e.g., a nucleic acid, a polypeptide, a vector, a cell, is removed from its original environment, e.g., the natural environment if it is naturally occurring. For example, a naturally-occurring polynucleotide or polypeptide present in a living animal is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides can be part of a vector and/or such polynucleotides or peptides can be part of a composition and still be isolated in that such vector or composition is not part of its natural environment.

[0164] The term "synthetic" as used herein means that the material, e.g., a nucleic acid, has been synthesized in vitro by well-known chemical synthesis techniques, as described in, e.g., Adams (1983) *J. Am. Chem. Soc.* 105:661; Belousov (1997) *Nucleic Acids Res.* 25:3440-3444; Frenkel (1995) *Free Radic. Biol. Med.* 19:373-380; Blommers (1994) *Biochemistry* 33:7886-7896; Narang (1979) *Meth. Enzymol.* 68:90; Brown (1979) *Meth. Enzymol.* 68:109; Beaucage (1981) *Tetra. Lett.* 22: 1859.

[0165] The term "recombinant" means that the nucleic acid is adjacent to a "backbone" nucleic acid to which it is not adjacent in its natural environment. Backbone molecules according to the invention include nucleic acids such as cloning and expression vectors, e.g., plasmids, used to maintain or manipulate a nucleic acid insert of interest. Recombinant peptide compounds of the invention, generated from these nucleic acids can be individually isolated or cloned and tested for a desired activity. Any suitable recombinant expression system can be used, including bacterial expression systems.

[0166] Also provided is a vector comprising at least one heterologous nucleic acid, e.g., artificial synthetic or recombinant BGC encoding a peptide compound described herein. The vector may be a cloning vector, an expression vector or an artificial chromosome.

[0167] As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. Vectors, including cloning and expression vectors, comprise a heterologous nucleic acid described herein or a functional equivalent thereof. Heterologous nucleic acids described herein can be incorporated into a recombinant replicable vector, for example a cloning or expression vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus, the invention

also provides a method of making polynucleotides of the invention by introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells are generally known to the person skilled in the art, and examples of suitable host cells are described herein. The vector into which the expression cassette or heterologous nucleic acid described herein is inserted may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of the vector will often depend on the host cell into which it is to be introduced. A variety of cloning and expression vectors for use with prokaryotic and eukaryotic hosts are described by Sambrook, et al, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, N. Y., (1989).

[0168] A vector according to the invention may be an autonomously replicating vector, i.e. a vector which exists as an extra-chromosomal entity, the replication of which is independent of chromosomal replication, e. g. a plasmid. Also provided herein is an isolated and purified plasmid comprising at least one synthetic or recombinant nucleic acid sequence, which encodes an artificial biosynthetic gene cluster (BGC) of a compound of formula (Ic) (or of formula (Ic'), or of formula (Ic'')), and has:

[0169] (i) a sequence identity to the full-length sequence of SEQ ID NO. 1 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or

[0170] (ii) a sequence completely complementary to any nucleic acid sequence of (i).

[0171] In one embodiment, provided herein is an isolated and purified plasmid comprising at least one synthetic or recombinant nucleic acid sequence, which encodes an artificial BGC encoding a peptide compound of formula (II) (or (IIa), (IIa') or (IIa'')) described herein above, and has:

[0172] (i) a sequence identity to the full-length sequence of SEQ ID NO. 86 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or

[0173] (ii) a sequence completely complementary to any nucleic acid sequence of (i).

[0174] Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

[0175] In a preferred embodiment, provided herein is an isolated and purified plasmid comprising at least one synthetic or recombinant nucleic acid sequence, which encodes an artificial BGC encoding a peptide compound of formula (1') (or (1b), or (1b')) described herein above, and has:

[0176] (i) a sequence identity to the full-length sequence of SEQ ID NO. 44 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or

[0177] (ii) a sequence completely complementary to any nucleic acid sequence of (i).

[0178] Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

[0179] The term "plasmid" refers to a circular double stranded DNA loop into which additional DNA segments can be ligated. A plasmid may have a bacterial origin of replication, or may contain an autonomously replicating sequence (ARS) derived from a yeast chromosome, and thus

be capable of autonomous replication in a bacterial, or yeast, host cell into which it is introduced. Alternatively, the plasmid may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the genomic DNA into which it has been integrated. The plasmid can be used to replicate the synthetic or recombinant nucleic acid utilized in the invention in a compatible host cell. The plasmid may be recovered from the host cell, and suitable host cells are described below. Moreover, the plasmid may be capable of directing the expression of genes to which it is operatively linked. The plasmid into which the synthetic or recombinant nucleic acid sequence utilized in the invention (expression cassette) is inserted may be any plasmid which may conveniently be subjected to recombinant DNA procedures, and the choice of the plasmid will often depend on the host cell into which it is to be introduced. A variety of cloning and expression plasmids for use with bacterial and yeast hosts are described by Sambrook, et al, *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, N. Y., (1989). A plasmid according to the invention may be used in vitro, for example, to transfect or transform a host cell. The plasmid of the invention may comprise one, two or more, for example three, four or five, of the synthetic or recombinant nucleic acid sequences utilized in the production method of the present invention, for example for overexpression. The isolated and purified plasmid of the invention comprises a synthetic or recombinant nucleic acid sequence utilized in the production method of the present invention in a form suitable for expression of the nucleic acid in a bacterial host cell (e.g., bacterium of a species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*), or a yeast such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*), which means that the plasmid may include one or more regulatory sequences, selected on the basis of the host cells to be used for expression, which is operationally linked to the nucleic acid sequence to be expressed. Particularly useful examples of host cells include bacterial cells of the species *E. coli*, *Bacillus*, *Corynebacteria*, *Lactobacillus* or *Streptomyces*, such as *E. coli* BL21(DE3); and yeast cells of the species *S. cerevisiae*, *Pichia pastoris* or *Yarrowia lipolytica*, such as *Pichia pastoris* GS115, *Pichia pastoris* X-33, *Pichia pastoris* KM71 and *Yarrowia lipolytica* Po1h (CLIB 882). The plasmids of the invention described herein can be designed for expression of the BGC encoding the peptide compound or the peptide compound of interest in a host cell. Suitable host cells are discussed further in Goeddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Representative examples of appropriate hosts are generally known in the art, and examples thereof are described herein. Appropriate culture media and conditions for host cells are also known in the art.

[0180] Within a plasmid, such as the isolated and purified plasmid of the invention, "operationally linked" is intended to mean that the nucleic acid sequence of interest (i.e. the BGC) is linked to the regulatory sequence(s) in a manner which allows for expression of the nucleic acid sequence (e.g., in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell), i.e. the term "operationally linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A

regulatory sequence such as a promoter, enhancer or other expression regulation signal “operationally linked” to a coding sequence is positioned in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences or the sequences are arranged so that they function in concert for their intended purpose, for example transcription initiates at a promoter and proceeds through the DNA sequence encoding the polypeptide.

[0181] The term “regulatory sequence” or “control sequence” is intended to include promoters, operators, enhancers, attenuators and other expression control elements (e.g., polyadenylation signal). Such regulatory sequences are described, for example, in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). The term regulatory or control sequence includes those sequences, which direct constitutive expression of a nucleotide sequence in many types of host cells and those, which direct expression of the nucleotide sequence only in a certain host cell (e.g. genus-specific regulatory sequences).

[0182] A plasmid or expression construct for a given host cell may thus comprise the following elements operationally linked to each other in a consecutive order from the 5'-end to 3'-end relative to the coding strand of the sequence encoding a polypeptide for production by the method of the invention: (i) a promoter sequence capable of directing transcription of the nucleotide sequence encoding the polypeptide in the given host cell; (ii) optionally, a signal sequence capable of directing secretion of the polypeptide from the given host cell into a culture medium; (iii) optionally, a sequence encoding for a C-terminal, N-terminal or internal epitope tag sequence or a combination of the aforementioned allowing purification, detection or labeling of the polypeptide; (iv) a nucleic acid sequence of the invention encoding a polypeptide of the invention; and preferably also (v) a transcription termination region (terminator) capable of terminating transcription downstream of the nucleotide sequence encoding the polypeptide.

[0183] Particular named bacterial promoters include *lad*, *ntpII*, *lacZ*, *pTuf*, *Eftu*, *T3*, *T7*, *SP6*, *K1F*, *tac*, *tet*, *gpt*, *p_{BAD}*, *lambda P_R*, *P_L* and *trp*. Examples of *Saccharomyces cerevisiae* promoters include *P_{ADH1}*, *P_{GAL1}*, *P_{CYC1}*, *P_{GP}* (also called *P_{TDH3}*), *P_{ZEV}*, *P_{ACT1}*, *P_{HXK1}*, *P_{YGR243}*, *P_{HXT4}*, *P_{HXT7}*, *P_{TEF1}*, *P_{TPI1}*, *P_{PGK1}*, *P_{TDH3}* and *P_{PYK1}*. In a preferred embodiment, the *S. cerevisiae* promoter is *P_{ADH1}*. Examples of *Pichia pastoris* promoters include *P_{AOX1}*, *P_{GAP}*, *P_{ADH3}*, *P_{PGK1}*, *P_{DAS}*, *P_{FLD1}*, and *P_{PEX8}*; and preferred embodiments include *P_{AOX1}*, *P_{GAP}*, *P_{ADH3}*, and *P_{PGK1}*. Examples of *Yarrowia lipolytica* promoters include *PmLEU2*, *pPOX2*, *pEYK1*, *pLIP2*, *pXPR2* and *pLEU2*; and preferred embodiments include *PmLEU2*. Examples of *Streptomyces albus* and *Streptomyces lividans* promoters include *KasOp*, *psf*, *ermEp*, *SF14p*, *PtpA*, *PkasO^{*}_{R15}*, and *Ptac*. In a preferred embodiment the *S. albus* and *S. lividans* promoters are *KasOp*, *psf* and *ermEp*.

[0184] Selection of the appropriate plasmid and promoter is well within the level of ordinary skill in the art. Downstream of the synthetic or recombinant nucleic acid sequence utilized in the production method of the present invention there may be a 3' untranslated region containing one or more transcription termination sites (e. g. a terminator). The origin of the terminator is less critical. The terminator can, for example, be native to the DNA sequence encoding the

peptide compound. Preferably, the terminator is endogenous to the host cell (in which the nucleotide sequence encoding the polypeptide is to be expressed). In the transcribed region, a ribosome binding site for translation may be present. The coding portion of the mature transcripts expressed by the constructs will include a translation initiating AUG (or TUG or GUG in prokaryotes) at the beginning and a termination codon appropriately positioned at the end of the peptide compound to be translated or processed.

[0185] Enhanced expression of a polynucleotide of the invention or a heterologous nucleic acid described herein may also be achieved by the selection of heterologous regulatory regions, e.g. promoter, secretion leader and/or terminator regions, which may serve to increase expression and, if desired, secretion levels of the protein of interest from the expression host and/or to provide for the inducible control of the expression of the peptide compound. It will be appreciated by those skilled in the art that the design of the plasmid can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The plasmids of the invention can be introduced into the recombinant host, e.g., microorganism (e.g., bacterial host cells) to thereby produce peptides, encoded by nucleic acids as described herein. Preferably, plasmids of the invention are capable of autonomous replication in a microorganism, such as a bacterium (e.g., bacterium of the species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*), or a yeast such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*). Preferably, plasmids of the invention are capable of autonomous replication in bacteria of the species *Lactobacillus* or *E. coli*, preferably *E. coli*.

[0186] The plasmids of the invention can be designed for expression of the BGC or the peptide compound of interest in bacterial cells such as *E. coli* or *Bacillus* strains, e.g., *Lactobacillus*. Suitable host cells are discussed further in Goeddel, Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, CA (1990). Representative examples of appropriate hosts are described hereafter. Appropriate culture media and conditions for the above-described host cells are known in the art.

[0187] As set out above, the term “control sequences” or “regulatory sequences” is defined herein to include at least any component, which may be necessary and/or advantageous for the expression of a peptide compound described herein. Any control sequence may be native or foreign to the nucleic acid sequences encoding a BGC for production of a peptide compound described herein. Such control sequences may include, but are not limited to, a promoter, a leader, optimal translation initiation sequences (as described in Kozak, 1991, J. Biol. Chem. 266:19867-19870), a secretion signal sequence, a pro-peptide sequence, a polyadenylation sequence, a transcription terminator. At a minimum, the control sequences typically include a promoter, and transcriptional and translational stop signals. A stably transformed microorganism is one that has had one or more DNA fragments introduced such that the introduced molecules are maintained, replicated and segregated in a growing culture. Stable transformation may be due to multiple or single chromosomal integration(s) or by (an) extrachromosomal element(s) such as (a) plasmid vector(s). A plasmid vector is capable of directing the expression of peptide compounds encoded by particular DNA fragments. Expression may be

constitutive or regulated by inducible (or repressible) promoters that enable high levels of transcription of functionally associated DNA fragments encoding specific peptide compounds.

[0188] Plasmids of the invention may also include a selectable marker gene to allow for the selection of bacterial strains that have been transformed, e.g., genes, which render the bacteria resistant to drugs such as chloramphenicol, erythromycin, kanamycin, neomycin, apramycin, tetracycline, zeocin, hygromycin, as well as ampicillin and other penicillin derivatives like carbenicillin. Selectable markers can also include biosynthetic genes, such as those in the histidine, tryptophan and leucine biosynthetic pathways.

[0189] The appropriate polynucleotide sequence may be inserted into the plasmid by a variety of procedures. In general, the polynucleotide sequence is ligated to the desired position in the plasmid following digestion of the insert and the vector with appropriate restriction endonucleases. Alternatively, blunt ends in both the insert and the vector may be ligated. A variety of cloning techniques are disclosed in Ausubel et al. Current Protocols in Molecular Biology, John Wiley 503 Sons, Inc. 1997 and Sambrook et al. Molecular Cloning: A Laboratory Manual 2nd Ed., Cold Spring Harbor Laboratory Press (1989). The polynucleotide sequence may also be cloned using homologous recombination techniques including in vitro as well as in vivo recombination. Such procedures and others are deemed to be within the scope of those skilled in the art.

[0190] Further provided is a recombinant host comprising a heterologous nucleic acid encoding a darobactin derivative of the invention, i.e. a compound of formula (I), (Ia) or (Ia') as described above. In an embodiment, the recombinant host is a microorganism, e.g., a bacterium such as a bacterium of the species of *E. coli*, *Bacillus*, *Corynebacteria* (e.g. *C. glutamicum*), *Lactobacillus* (e.g. *Lactococcus lactis*) or *Streptomyces* (e.g. *S. albus*, *S. lividans*), or a yeast such as *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*. The heterologous nucleic acid may be part of a vector, such as a plasmid, contained in the microorganism. The vector, e.g. an expression vector, such as a plasmid, can be introduced into the microorganism, e.g. a bacterial host cell or yeast host cell of a species described herein, by any of a variety of techniques known to those skilled in the art, including transformation or transfection. The invention also provides bacterial host cells, i.e. transformed cells comprising at least one synthetic or recombinant nucleic acid sequence, which encodes a biosynthetic gene cluster for production of a compound of formula (Ic) (or formula (Id) or (Ie)), or a plasmid of the invention. The host cell may be any bacterial host cell familiar to those skilled in the art, e.g. an *Escherichia coli* or *Lactobacillus* cell, preferably an *E. coli* cell, and more preferably an *E. coli* BL21(DE3) cell. The plasmid can be introduced into the bacterial host cell using any of a variety of techniques, including transformation or transfection. Particular methods include calcium phosphate transfection, DEAE-Dextran mediated transfection, lipofection, or electroporation (Davis, L., Dibner, M., Battey, L., Basic Methods in Molecular Biology, (1986)).

[0191] Where appropriate, the engineered host cells can be cultured in conventional nutrient media modified as appropriate for activating promoters, selecting transformants or amplifying the synthetic or recombinant nucleic acid utilized in the production method of the invention. Following transformation of a suitable host strain and growth of the host

strain to an appropriate cell density, the selected promoter may be induced by appropriate means (e.g., temperature shift or chemical induction) and the cells may be cultured for an additional period to allow them to produce the desired peptide compound or derivatives thereof. The culture conditions such as temperature, pH and the like, are apparent to the ordinarily skilled artisan, and examples are described herein. The host cell, when cultured under suitable conditions, is capable of producing darobactin A or darobactin derivatives, i.e. peptide compounds according to formula (Ic) (or (Ic'), or (Ic'')) or any of the other compounds described herein (especially darobactin derivatives of the present invention), that it otherwise does not produce, or produces at a significantly lower level, in the absence of a modification according to the invention. Cells can be harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract is retained for further purification. Microbial cells employed for expression of proteins can be disrupted by any convenient method, including freeze-thaw cycling, sonication, mechanical disruption, or use of cell lysing agents. Such methods are well known to those skilled in the art. The expressed peptide compound or derivatives thereof can be recovered and purified from recombinant cell cultures by methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite chromatography and lectin chromatography. If desired, high performance liquid chromatography (HPLC) can be employed for final purification steps.

[0192] The terms "protein", "polypeptide", "peptide" as used herein define an organic compound made of two or more amino acid residues arranged in a linear chain or cyclic structure, wherein individual amino acids in the organic compound are linked by peptide bonds, i.e. an amide bond formed between adjacent amino acid residues. By convention, the primary structure of a protein is reported starting from the amino-terminal (N) end to the carboxyl-terminal (C) end.

BRIEF DESCRIPTION OF THE FIGURES

[0193] FIG. 1 graphically depicts the general structure of a darobactin expression vector, i.e. plasmid, of the invention. Table 1 below summarizes the restriction sites (R-sites) that are present in the expression vector for modification by cloning techniques.

TABLE 1

Restriction sites (R-sites) with appropriate enzyme and function		
R-site	Enzyme	Function
R ₂	MauBI	pNOSO cloning
R ₃	AgeI	Exchange/introduction of element(s)
R ₄	MreI	T7 promoter exchange
R ₅	XmaII	T7 promoter exchange
R ₆	NdeI	T7 promoter & darA exchange
R ₇	PstI	darA exchange
R ₈	BamHI	darA exchange
R ₉	NotI	ntpII promoter exchange
R ₁₀	XhoI	ntpII promoter exchange
R ₁₁	SpeI	introduction of alternative and/or additional genes
R ₁₂	PacI	introduction of alternative and/or additional genes
R ₁₃	SgrDI	pNOSO cloning

[0194] FIG. 2 graphically depicts a biosynthetic gene cluster (BGC) responsible for the production of darobactin A, and further shows the DNA sequence encoding the darobactin A propeptide core sequence, its corresponding linear propeptide amino acid sequence and the applied numbering of the amino acids.

EXAMPLES

Generation of Expression Vector pNOSOdarA-E-DA

[0195] Plasmid expression vector pNOSOdarA-E-DA (deposited at DSMZ on 10 Feb. 2021; DSM 33801) was generated from pNOSO vector and an artificial synthetic sequence comprising a BGC for the production of darobactin A. Both, pNOSO and the synthetic sequence comprising the BGC, were designed in silico, then chemically synthesized and assembled using restriction/ligation-based cloning techniques. The pNOSO vector harbours p15A ori, kanamycin resistance cassette (KanR), CEN6/ARS4 (yeast ori) and URA3 (yeast counter selection marker). The artificial synthetic sequence comprises lactose-dependent repressor gene (*lacI*), T7 promoter (P_{T7}), constitutive promoter (P_{nptII}), termination cassettes (t_{D1}) and a BGC of darobactin A. Additionally, point mutations were introduced into pNOSO and the synthetic BGC to generate unique R-sites for cloning purposes. After chemical synthesis of the vector pNOSO and the artificial synthetic sequence comprising the darobactin BGC, the pNOSOdarA-E-DA expression construct was generated by restriction/ligation-based cloning techniques. The CEN6/ARS4 sequence, the URA3 sequence and the *relE* gene were removed during generation of plasmid expression vector pNOSOdarA-E-DA. The detailed vector organization is graphically depicted in FIG. 1; for details of the R-sites see Table 1 above.

Generation of Artificial darA Gene Fragments and Plasmid Expression Vectors

[0196] For the generation of new artificial darobactin derivatives with altered amino acid sequences compared to darobactin A, the propeptide sequences in the darA gene were modified in silico by exchange of nucleotides and/or codons. Artificial darA gene fragments containing the modified propeptide sequences were chemically synthesized and cloned into pNOSOdarA-E-DA by replacing darA with a modified darA gene fragment using restriction/ligation-based cloning techniques. Using this strategy, plasmid expression vectors with altered propeptide sequences in the darA gene were generated (e.g. pNOSOdarA-E-DB to pNOSOdarA-E-DE and pNOSOdarA-E-D1 to pNOSOdarA-E-D17). Table 2 shows modified propeptide DNA sequences of artificial darA gene fragments and their corresponding propeptide amino acid sequence, which can be used for the production of darobactin derivatives.

TABLE 2

Propeptide DNA and protein sequences of darobactin derivatives.		
Darobactin derivative	DNA Propeptide core sequence	Propeptide amino acid sequence
B	TGGAAGTGGACCAACGATTC	WNWTKRF
C	TGGTCATGGTCAAGAAGCTTC	WSWSRSF

TABLE 2-continued

Propeptide DNA and protein sequences of darobactin derivatives.		
Darobactin derivative	DNA Propeptide core sequence	Propeptide amino acid sequence
D	TGGAAGTGGTCAAGAAGCTTC	WNWSRSF
E	TGGTCATGGTCAAAAAGCTTC	WSWSKSF
D1	TGGTCATGGACCAACGATTC	WSWTKRF
D2	TGGTCATGGACCAAGAAGCTTC	WSWTKRF
D4	TGGAAGTGGACCAAAAAGCTTC	WNWTKSF
D5	TGGTGCTGGTCAAAAAGCTTC	WCWSKSF
D6	TGGAAGTGGTGCAAAAAGCTTC	WNWCKSF
D7	TGGTGCTGGTGCAAAAAGCTTC	WCWCKSF
D8	TGGAAGTGGTCAAAATGCTTC	WNWSKCF
D9	TGGAAGTGGTCAAAAAGCTGG	WNWSKSW
D10	TGGAAGTGGTCAAAAAGCTAC	WNWSKSY
D11	TGGCAGTGGTCAAAAAGCTTC	WQWSKSF
D12	TGGACCTGGTCAAAAAGCTTC	WTWSKSF
D13	TGGGCATGGTCAAAAAGCTTC	WAWSKSF
D14	TGGAAGTGGGCAAAAAGCTTC	WNWAKSF
D15	TGGAAGTGGTCAGCAAGCTTC	WNWSASF
D16	TGGAAGTGGTCAAAAAGCTTTC	WNWSKAF
D17	TGGAAGTGGTCAAAAAGCGCA	WNWSKSA

[0197] The correctness of the generated plasmids was verified by restriction hydrolysis followed by agarose gel electrophoresis and DNA sequencing. Designation of the plasmid expression vectors corresponds to nomenclature of the darobactin derivatives, e.g., pNOSOdarA-E-D9 harbours the propeptide sequence of darobactin D9 (i.e. WNWSKSW) and pNOSOdarA-E-D17 harbours the propeptide sequence of darobactin derivative 17 (i.e. WNWSKSA); both of which have been deposited at DSMZ on 10 Feb. 2021; DSM 33802 (pNOSOdarA-E-D9) and DSM 33803 (pNOSOdarA-E-D17)).

Heterologous Expression and Production

[0198] For heterologous production of darobactin A (DA) and darobactin derivatives, a plasmid expression vector as described above was transformed into *E. coli* BL21 (DE3) cells by electroporation, and the respective transformed host cells were grown under controlled conditions as, for example, described below.

Fermentation/Cultivation of Host Cells

Day 1

[0199] Inoculation of a single clone of *E. coli* BL21 (DE3) cells harbouring plasmid pNOSodarA-E-DA or pNOSodarA-E with modified darA gene in 10 ml LB medium including 30 µg/ml Kanamycin (Kan30)

Day 2

[0200] Inoculation of 50 ml FM medium (1.254% K₂HPO₄, 0.231% KH₂PO₄, 0.4% D(+)-glucose, 0.1% NH₄Cl, 1.2% yeast extract, 0.5% NaCl and 0.0492% MgSO₄(*7H₂O); pH 7.1) including Kan30 and 1 mg/l VitaminB12 with 1% (v/v) of preculture

[0201] Incubation for 3 days at 30° C., 180 rpm (Infors HT)

Separation and Retaining of Compounds

Day 5

[0202] Harvest supernatant by centrifugation at 8,000 g for 10 min at 4° C.

[0203] Incubation of supernatant with 0.05% XAD16N resin (Sigma Aldrich) (w/v) under agitation for 2 h

[0204] Harvest of XAD16N resin and elution of bound compounds from XAD16N resin with 10 ml 80% Methanol under shaking for 1.5 h

[0205] Drying of eluate in a rotary evaporator

[0206] Solubilization of evaporated extract in 0.5 ml 50/50 methanol/water mixture

[0207] HPLC-MS analysis to verify the production of peptide compound of interest, e.g., darobactin A or darobactin derivative; characteristic ions for darobactins in tandem MS mode are m/z 160.075 and a prominent [M-H₂O+2H]²⁺ ion.

[0208] In case of production of compounds for antimicrobial testing with the minimum inhibitory concentration assays described below, no antibiotic was added to the expression cultures as a selection marker.

[0209] HPLC purification was carried out on a Waters Autopurifier (Eschborn, Germany) high pressure gradient system, equipped with 2545 binary gradient module, SFO system fluidics organizer, 2767 sample manager and a 2998 photodiode array detector coupled to a 3100 single quadrupole mass spectrometer operated in positive ion mode. Source and voltage settings for the MS were as follows: mass range, m/z 300-1000; scan duration, 1 s; points per Dalton, 4; capillary voltage, 3.5 kV; cone voltage, 30 V; extractor voltage 3 V; RF lens, 0.1 V; source temperature 120° C.; desolvation temperature, 250° C.; desolvation gas flow, 400 L/h; cone gas flow, 50 L/h; ion counting threshold, 30. Separation was carried out on a Waters X-Bridge prep C-18 5 µm ODB, 150×19 mm column using ACN+0.1% FA as B and H₂O+0.1% FA as A and a flow rate of 25 ml/min. Compound detection was carried out by detection of the [M+2H]²⁺. Second purification step was carried out on the same instrument on an Phenomenex Kinetex 5u Biphenyl 100A 250×21.2 mm using ACN+0.1% FA as B and H₂O+0.1% FA as A and a flow rate of 25 ml/min. Where necessary, a Dionex Ultimate 3000 SDLC low pressure gradient system on a Waters CSH Phenyl-Hexyl 250 mm×10 mm 5 µm dp column with the eluents H₂O+0.1% FA as A and ACN+0.1% FA as B, a flow rate of 5 ml/min and a column thermostated at 30° C. was used for further purification of peptide compounds of interest. Peptide compounds of interest, e.g. darobactin A and darobactin derivatives, can be detected by UV absorption at 254 nm.

[0210] Following the above protocol, darobactin A was, for example, heterologously produced by expression of a modified artificial plasmid expression vector comprising propeptide sequence A (see, e.g., FIG. 2) in *E. coli* BL21 (DE3) cells, and obtained as a slightly yellow solid.

Example Compounds

[0211] Exemplified example compounds obtained by heterologous expression of the modified BGCs according to the present invention are shown in Table 3 below.

TABLE 3


Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²
A		483.70

TABLE 3-continued




Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²
B		525.25
B*		525.25
D		497.71

TABLE 3-continued




Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²
E		470.20
D1		511.75
D1*		511.75

TABLE 3-continued





Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²
D2		491.21
D2*		491.21
D4		490.71
D4*		490.71


TABLE 3-continued

Produced example compounds	
Darobactin/ Darobactin derivative	Mass (m/z)* [M + 2H] ²
D9	503.21
D10	491.70
D11	490.71
D12	477.21

TABLE 3-continued

Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²
D12*	The chemical structure of D12* is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, and a 2-amino-3-hydroxybutyryl group. The structure is highly symmetrical and contains multiple amide and hydroxyl groups.	477.21
D14	The chemical structure of D14 is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, and a 2-amino-3-hydroxybutyryl group. The structure is highly symmetrical and contains multiple amide and hydroxyl groups.	475.71
D15	The chemical structure of D15 is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, and a 2-amino-3-hydroxybutyryl group. The structure is highly symmetrical and contains multiple amide and hydroxyl groups.	455.17
D16	The chemical structure of D16 is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, a 2-amino-3-hydroxybutyryl group, and a 2-amino-3-hydroxybutyryl group. The structure is highly symmetrical and contains multiple amide and hydroxyl groups.	475.71

TABLE 3-continued

Produced example compounds		
Darobactin/ Darobactin derivative	Chemical structure based on exact mass and MS2 fragmentation pattern	Mass (m/z)* [M + 2H] ²⁺
D17		445.69

*All observed [M + 2H]²⁺ masses correlate with the expected [M + 2H]²⁺ masses.

Assessment of Antimicrobial Activity of Darobactin A and Darobactin Derivative 9

[0212] All microorganisms were handled according to standard procedures as recommended by the depositor. Strains were obtained from the German Collection of Cell Cultures and Microorganisms (*Klebsiella pneumoniae* DSM-30104, *Acinetobacter baumannii* DSM-30008), the American Type Culture Collection (*Staphylococcus aureus* ATCC-29213, *Escherichia coli* ATCC-25922, *Enterococcus*

growth was observed. An extract of sterile fermentation medium (FM medium) and *E. coli* BL21 (DE3) cells harbouring plasmid expression vector pNOSO without BGC were used as negative controls. Results of darobactin A and darobactin derivative 9 produced in accordance with the invention compared to data published in Imai et al. (loc. cit.) are summarized in Table 4 below. Negative controls showed no inhibitory effect on the tested pathogen strains (data not shown) thereby confirming that antimicrobial activity is attributable to the tested compounds.

TABLE 4

Antimicrobial activity of darobactin A and darobactin derivative 9 produced in accordance with the method of the invention compared to darobactin A activity data published in Imai et al.				
	<i>K. pneumoniae</i> DSM-30104	<i>P. aeruginosa</i> PA01	<i>E. coli</i> ATCC25922	<i>A. baumannii</i> DSM-30008
MIC [μg/ml] of darobactin A (produced by method of invention)	4	4	2	4
MIC [μg/ml] of darobactin D9 (produced by method of invention)	8	0.5	2	0.5
MIC [μg/ml] of darobactin A (Imai et al. 2019)	4	2	2	4

faecalis ATCC-29212), or were part of our internal strain collection (*Pseudomonas aeruginosa* PA01).

[0213] All samples were tested according to standard procedures to establish minimum inhibitory concentration (MIC) by the microbroth dilution method. Bacteria were grown on solid media (CASO agar) and single colonies were collected with sterile cotton swabs and resuspended in saline solution (0.9% NaCl) to obtain McFarland 0.5. The bacterial suspension was diluted 1:100 in cation-adjusted Mueller-Hinton broth to obtain a final inoculum of approximately 10⁶ CFU/mL. Serial dilutions of the test samples were prepared in cation-adjusted Mueller-Hinton broth in sterile 96-well microtiter plates in a total volume of 75 μL. The prepared cell suspension was added (75 μL) and the plates were incubated at either 30° C. or 37° C. for at least 24 h under static conditions. Plates were then checked visually and MIC was determined as lowest concentration where no visible

[0214] As demonstrated above, darobactin A produced by the method of the invention has at least the same antimicrobial activity against Gram-negative bacteria as reported by Imai et al. Consequently, the production method of the invention provides a viable and efficient way to produce active compounds of interest selectively in high yields. Moreover, the novel darobactin D9 has comparable or even higher activity against the tested Gram-negative bacteria than Darobactin A.

Assessment of Antimicrobial Activity of Darobactin Derivatives

[0215] Crude extracts of darobactin derivatives obtained by the heterologous expression and separation protocol described above were tested for their activity against exemplified Gram-negative bacteria with a minimum inhibitory concentration (MIC) assay. The presence of darobactin derivatives in the crude extracts was verified by checking the expected masses via LC-MS.

[0216] For the MIC assay, 10 μ l of crude extract of each test sample, diluted in 50% methanol/H₂O, was added each into a well of the first row of a 96 well plate. After evaporation of the methanol/H₂O solvent, until only about 4 μ l solvent was remaining in each well of the first row, 71 μ l cation-adjusted Mueller-Hinton broth medium was added to each of said wells, so that each well in the first row of the 96 well plate had a volume of 75 μ l. Then, 75 μ l of cation-adjusted Mueller-Hinton broth medium was added to each well and the samples contained in the first row were mixed with the 75 μ l medium/extract mixture. Starting from the 150 μ l mixtures in the first row, serial dilutions of the respective mixtures were prepared in corresponding subjacent rows of the 96 well plate. Bacterial suspensions of each strain to be tested were diluted 1:100 in cation-adjusted Mueller-Hinton broth to obtain a final inoculum of approximately 10⁶ CFU/mL. 75 μ l of each bacterial suspension to be tested was added to the respective crude extract samples in the wells of the 96 well plate and the plates were incubated at either 30° C. or 37° C. for at least 24 h under static conditions. Plates were then checked visually and MIC was determined as lowest concentration where no visible growth was observed. In order to quantify the activity of the obtained darobactin derivatives, their relative activity in relation to the activity of darobactin A was determined. The amount of darobactin derivative present in the extract test sample was calculated based on the area under the curve (AUC) of the mass peak as automatically calculated by software DataAnalysis 4.2 and extracted from the LC-MS measurements of each of the crude extract samples. Using the AUC of the mass peak of darobactin A (DA) as calibration standard, i.e. ratio between AUC of darobactin derivative and darobactin A was correlated to amount of compound present in sample. The results of activities of tested darobactin derivatives against exemplified Gram-negative bacteria as compared to the activity of darobactin A are summarized in Table 5 below. A value of '+++' indicates that the derivative has the same antimicrobial activity as darobactin A, a value of 0 (no activity) to '++' indicates a lower activity than darobactin A, and a value of '++++' to '+++++' indicates a higher activity than darobactin A.

[0217] As demonstrated above, darobactin derivatives have an excellent antimicrobial activity against Gram-negative bacteria, and some show selectivity against certain pathogens.

Artificial Darobactin 9 (D9) Derivatives

[0218] For the generation of new artificial darobactin 9 (D9) derivatives with altered amino acid sequences, the propeptide sequences in the darA gene were modified in silico by exchange of certain nucleotides and/or codons. Artificial darA gene fragments containing the modified propeptide sequences were prepared by overlap extension polymerase chain reaction (OE-PCR) using plasmid expression vector pNOSOdarA-E-D9 (DSM 33802), which harbours the propeptide sequence of D9 (i.e. WNWSKSW), as template. To insert the required mutations into the propeptide sequence, specific reverse primers (denoted as pr2darXX_rv with XX indicating the number of the designated darobactin derivative; e.g., pr2dar22_rv includes substitutions (mismatches compared to template sequence) required for the DNA propeptide core sequence of darobactin derivative 22), which included the substitutions and a complementary sequence with primer pr3dartrp_fw at their 5' end, were designed. A first DNA fragment was generated using primer pr1dar_fw and one of pr2darXX_rv. Primer pr1dar_fw binds to a region located in the T7 lac promoter of the plasmid expression vector pNOSOdarA-E-D9 (DSM 33802), and primer pr2darXX_rv binds to the core region of darA, contains the required substitutions encoding the desired darobactin derivative and carries a region of 13 base pairs that overlaps with primer pr3dartrp_fw. A second DNA fragment was generated using primers pr3dartrp_fw and pr4dar_rv. Primer pr4dar_rv binds in front of the termination region at the end of the intergenic region between darA and darB of the plasmid expression vector pNOSOdarA-E-D9, and primer pr3dartrp_fw binds to the core region of darA. The respective first DNA fragment (containing the muta-

TABLE 5

Relative activity of tested compounds against exemplified Gram-negative bacteria				
Darobactin / Derivative	<i>Klebsiella</i> <i>pneumoniae</i> DSM-30104	<i>Pseudomonas</i> <i>aeruginosa</i> PA01	<i>E. coli</i> ATCC25922	<i>Acinetobacter</i> <i>baumannii</i> DSM-30008
A	+++	+++	+++	+++
B	+++++	+++++	+++++	+++++
B*	+++++	+++++	+++++	+++++
D	0	++++	++++	++++
2	0	++++	++++	++++
2*	0	++++	++++	++++
4	++	+++	+++	+++
4*	++	+++	+++	+++
6	++++	+++++	+++++	+++++
8	0	+++++	+++++	+++++
9	+++	+++	++++	++++
10	0	+	0	0
11	++++	+++++	+++++	+++++
12	0	+++++	+++++	+++++
12*	0	+++++	+++++	+++++
14	+++	++	+++	+++
16	+	+	+	+

tions) was then combined with the second DNA fragment, and a third PCR reaction with primers pr1dar_fw and pr4dar_rv was performed to thereby generate a fusion fragment, due to PCR based overlap extension, harbouring the modified darA propeptide core sequence and unique restriction sites on the 3' and 5' end, respectively. Primer sequences are listed in Table 6; substitutions are shown in bold.

TABLE 6	
Oligonucleotide primers used for OE-PCR	
Primer	Sequence
pr1dar_fw	GGTGATGTCGCGCATATAGG
pr3dartrp_fw	TGGCAGGAAATTTAAAGCTTATCCCAT
pr4dar_rv	TGGGGATCCTCAGGACTGCAG
pr2dar22_rv	AAATTTCTGCCACCGTTTAGTCCAGTTCC
pr2dar23_rv	AAATTTCTGCCAATGTTTAGTCCAGTTCC
pr2dar24_rv	AAATTTCTGCCATGATTTAGTCCAGTTCC
pr2dar25_rv	AAATTTCTGCCACGCTTTAGTCCAGTTCC
pr2dar30_rv	AAATTTCTGCCATGATTTGCCCCAGTTCC
pr2dar31_rv	AAATTTCTGCCACCGTTTCGCCAGTTCC
pr2dar32_rv	AAATTTCTGCCACCGTTTACACCAGTTCC
pr2dar33_rv	AAATTTCTGCCATGATTTACACCAGTTCC
pr2dar34_rv	AAATTTCTGCCACGCTTTACACCAGTTCC
pr2dar36_rv	AAATTTCTGCCATTTTTAGTCCAGTTCC
pr2dar37_rv	AAATTTCTGCCAAGTTTGTACAGTTCC
pr2dar38_rv	AAATTTCTGCCAAGTTTTAGTCCAGTTCC
pr2dar39_rv	AAATTTCTGCCACCGCGAGTCCAGTTCCA

[0219] The product of the ‘overlap extension’ PCR reaction was gel-purified and the obtained DNA fragment was cloned into plasmid pNOSodarA-E-D9 by replacing the sequence encoding darobactin derivative D9 with the modified darA gene fragment using restriction/ligation-based cloning techniques. Using this strategy, plasmid expression vectors with altered propeptide sequences in the darA gene were generated (e.g. pNOSodarA-E-D22 to pNOSodarA-E-D25, pNOSodarA-E-D30 to pNOSodarA-E-D34 and pNOSodarA-E-D36 to pNOSodarA-E-D39). Table 7 shows modified propeptide DNA sequences of modified darA gene fragments and their corresponding propeptide amino acid core sequence. The nucleotide substitutions compared to D9-encoding darA core sequence are shown in bold and the designation of the plasmid expression vectors corresponds to the nomenclature of the darobactin derivatives, e.g., pNOSodarA-E-D22 harbours the propeptide sequence of darobactin 22 (D22). Correctness of the plasmid sequences was verified by restriction hydrolysis followed by agarose gel electrophoresis and sanger sequencing at LGS genomics.

TABLE 7		
Propeptide DNA and protein sequences of darobactin derivatives.		
Darobactin derivative	DNA propeptide core sequence	Propeptide amino acid sequence
22	TGG AAC TGG ACT AAA CGG TGG	WNWTKRW
23	TGG AAC TGG ACT AAA CAT TGG	WNWTKHW
24	TGG AAC TGG ACT AAA TCA TGG	WNWTKSW
25	TGG AAC TGG ACT AAA GCG TGG	WNWTKAW
30	TGG AAC TGG GCG AAA TCA TGG	WNWAKSW
31	TGG AAC TGG GCG AAA CGG TGG	WNWAKRW
32	TGG AAC TGG TGT AAA CGG TGG	WNWCKRW
33	TGG AAC TGG TGT AAA TCA TGG	WNWCKSW
34	TGG AAC TGG TGT AAA GCG TGG	WNWCKAW
36	TGG AAC TGG ACT AAA AAA TGG	WNWTKKW
37	TGG AAC TGG TCA AAA ACT TGG	WNWSKRW
38	TGG AAC TGG ACT AAA ACT TGG	WNWTKRW
39	TGG AAC TGG ACT CGG CGG TGG	WNWTRRW

Heterologous Expression and Production

[0220] For heterologous production of the darobactin derivatives, one of the plasmid expression vectors described above, i.e. pNOSodarA-E-D22 to pNOSodarA-E-D25, pNOSodarA-E-D30 to pNOSodarA-E-D34 and pNOSodarA-E-D36 to pNOSodarA-E-D39, were separately transformed into *E. coli* BL21 (DE3) cells by electroporation, and the respective recombinant strains were cultivated to investigate their darobactin production profile as described below.

Fermentation/Cultivation of Host Cells

Day 1

[0221] Inoculation of a single clone of *E. coli* BL21 (DE3) cells harbouring plasmid pNOSodarA-E with modified darA gene in 10 ml LB medium including 30 µg/ml Kanamycin (Kan30) and incubation overnight at 30° C., 180 rpm (Infors HT)

Day 2

[0222] Inoculation of 50 ml FM medium (1.254% K₂HPO₄, 0.231% KH₂PO₄, 0.4% D(+)-glucose, 0.1% NH₄Cl, 1.2% yeast extract, 0.5% NaCl and 0.0492% MgSO₄(*7H₂O), pH 7.1) including Kan30 and 1 mg/l VitaminB12 with 1% (v/v) of preculture

[0223] Incubation for 3 days at 30° C., 180 rpm (Infors HT)

Separation and Retaining of Compounds

Day 5

- [0224] Harvest supernatant by centrifugation at 8,000 g for 10 min at 4° C.
- [0225] Incubation of supernatant with 0.05% XAD16N resin (Sigma Aldrich) (w/v) under agitation for 2 h
- [0226] Harvest of XAD16N resin and elution of bound compounds from XAD16N resin with 10 ml 80% Methanol under shaking for 1.5 h
- [0227] Drying of eluate in a rotary evaporator
- [0228] Solubilization of evaporated extract in 0.5 ml 50/50 methanol/water mixture; or in 0.5 ml 100% water
- [0229] HPLC-MS analysis to verify the production of peptide compound of interest, e.g., darobactin A or darobactin derivative; characteristic ions for darobactins in tandem MS mode are m/z 160.075 and a prominent $[M-H_2O+2H]^{2+}$ ion.
- [0230] In case of production of compounds for antimicrobial testing with the minimum inhibitory concentration assays described below, no antibiotic was added to the expression cultures as a selection marker.
- [0231] HPLC purification was carried out on a Waters Autopurifier (Eschborn, Germany) high pressure gradient system, equipped with 2545 binary gradient module, SFO system fluidics organizer, 2767 sample manager and a 2998 photodiode array detector coupled to a 3100 single quadrupole mass spectrometer operated in positive ion mode. Source and voltage settings for the MS were as follows: mass range, m/z 300-1000; scan duration, 1 s; points per Dalton, 4; capillary voltage, 3.5 kV; cone voltage, 30 V; extractor voltage 3 V; RF lens, 0.1 V; source temperature

120° C., desolvation temperature, 250° C.; desolvation gas flow, 400 L/h; cone gas flow, 50 L/h; ion counting threshold, 30. Separation was carried out on a Waters X-Bridge prep C-18 5 μ m ODB, 150×19 mm column using ACN+0.1% FA as B and H₂O+0.1% FA as A and a flow rate of 25 ml/min. Compound detection was carried out by detection of the $[M+2H]^{2+}$. Second purification step was carried out on the same instrument on an Phenomenex Kinetex 5u Biphenyl 100A 250×21.2 mm using ACN+0.1% FA as B and H₂O+0.1% FA as A and a flow rate of 25 ml/min. Where necessary, a Dionex Ultimate 3000 SDLC low pressure gradient system on a Waters CSH Phenyl-Hexyl 250 mm×10 mm 5 μ m dp column with the eluents H₂O+0.1% FA as A and ACN+0.1% FA as B, a flow rate of 5 ml/min and a column thermostated at 30° C. was used for further purification of peptide compounds of interest. Peptide compounds of interest, e.g. darobactin A and darobactin derivatives, can be detected by UV absorption at 254 nm.

[0232] Following the above protocol, darobactin 22 was, for example, heterologously produced by expression of a modified artificial plasmid expression vector pNOSodarA-E-D22 comprising propeptide sequence D22 (see, Table 2 above) in *E. coli* BL21(DE3) cells.

Example Compounds

[0233] Exemplified compounds obtained by heterologous expression of the modified BGCs according to the present invention are shown in Tables 8 and 9 below. It was found that darobactin derivatives with sulfur-containing amino acids carry a protection group (proved via Nuclear Magnetic Resonance (NMR) spectroscopy).

TABLE 8


Produced example compounds D22-D25*, D30, D31, and D36-D39		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) $[M + 2H]^{+2}$
D22		544.76

TABLE 8-continued

Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (<i>m/z</i>) [M + 2H] ⁺
D22*		544.76
D23		535.23
D23*		535.23

TABLE 8-continued








Produced example compounds D22-D25*, D30, D31, and D36-D39		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D24		510.22
D24*		510.22
D25		502.22
D25*		502.22

TABLE 8-continued

Produced example compounds D22-D25*, D30, D31, and D36-D39		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ⁺ 2
D36*	The chemical structure of D36* is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 4-aminobutyl group, a 2-hydroxypropyl group, a 2-aminopropyl group, and a 2-aminobutyl group. The molecule also contains a 2-aminobutyl group and a 2-aminopropyl group. The structure is highly branched and contains multiple amide and ester linkages.	530.75
D37	The chemical structure of D37 is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 4-aminobutyl group, a 2-hydroxypropyl group, a 2-aminopropyl group, and a 2-aminobutyl group. The molecule also contains a 2-aminobutyl group and a 2-aminopropyl group. The structure is highly branched and contains multiple amide and ester linkages.	510.22
D37*	The chemical structure of D37* is a complex polycyclic molecule. It features a central benzimidazole core. Attached to this core are several side chains: a 4-aminobutyl group, a 2-hydroxypropyl group, a 2-aminopropyl group, and a 2-aminobutyl group. The molecule also contains a 2-aminobutyl group and a 2-aminopropyl group. The structure is highly branched and contains multiple amide and ester linkages.	510.22

TABLE 8-continued

Produced example compounds D22-D25*, D30, D31, and D36-D39		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D38		517.22
D38*		517.22
D39		558.75

*All observed [M + 2H]²⁺ masses correlate with the expected [M + 2H]²⁺ masses.

TABLE 9A



Produced example compounds PD32, PD33 and PD34 and PD32*, PD33* and PD34* with sulfur-containing amino acid		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass, MS2 fragmentation pattern and NMR	Observed mass* (m/z) [M + 2H] ²⁺
PD32		641.25
PD32*		641.25

TABLE 9A-continued





Produced example compounds PD32, PD33 and PD34 and PD32*, PD33* and PD34* with sulfur-containing amino acid		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass, MS2 fragmentation pattern and NMR	Observed mass* (m/z) [M + 2H] ²⁺
PD33		606.72
PD33*		606.72

TABLE 9A-continued

Produced example compounds PD32, PD33 and PD34 and PD32*, PD33* and PD34* with sulfur-containing amino acid		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass, MS2 fragmentation pattern and NMR	Observed mass* (m/z) [M + 2H] ²⁺
PD34		598.72
PD34*		598.72

[0234] Darobactins D32 to D34 and D32* to D34* shown in Table 9B can be prepared from the protected compounds PD32 to PD34* by following standard protocols for the cleavage of disulfide bonds using dithiobutylamine (DTBA), tris (2-carboxyethyl) phosphine hydrochloride (TCEP-HCl) or mercaptoethanol or dithiothreitol (DTT). Such standard procedures are generally known in the art, and have, for example, been described by Kirley, Terence L. et al. (2016), Selective disulfide reduction for labeling and enhancement of Fab antibody fragments, *Biochemical and Biophysical*

Research Communications 480 (4), pp. 752-757 (DOI: 10.1016/j.bbrc.2016.10.128)-Lukesh, John C. et al. (2012), A Potent, Versatile Disulfide-Reducing Agent from Aspartic Acid, *Journal of the American Chemical Society* 134 (9), pp. 4057-4059 (DOI: 10.1021/ja211931f); and Yu, Shukun et al. (2020), Comparative study of protease hydrolysis reaction demonstrating Normalized Peptide Bond Cleavage Frequency and Protease Substrate Broadness Index, *PLoS ONE* 15 (9), e0239080 (DOI: 10.1371/journal.pone.0239080)

TABLE 9B




Compounds D32-D34 and D32*-D34*		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D32		545.73
D32*		545.73
D33		511.20

TABLE 9B-continued

Compounds D32-D34 and D32*-D34*		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D33*	The chemical structure of D33* is a complex polypeptide derivative. It features a central benzimidazole core. On the left, there is a tryptophan residue with a 5-amino group. The main chain includes several amide bonds, a thiol group (-SH), and a hydroxyl group (-OH). On the right, there is a tryptophan residue with a 7-amino group. The structure is highly branched and contains multiple functional groups.	511.20
D34	The chemical structure of D34 is similar to D33* but with a different side chain on the right. It features a central benzimidazole core. On the left, there is a tryptophan residue with a 5-amino group. The main chain includes several amide bonds, a thiol group (-SH), and a hydroxyl group (-OH). On the right, there is a tryptophan residue with a 7-amino group. The structure is highly branched and contains multiple functional groups.	503.20
D34*	The chemical structure of D34* is similar to D34 but with a different side chain on the right. It features a central benzimidazole core. On the left, there is a tryptophan residue with a 5-amino group. The main chain includes several amide bonds, a thiol group (-SH), and a hydroxyl group (-OH). On the right, there is a tryptophan residue with a 7-amino group. The structure is highly branched and contains multiple functional groups.	503.20

Generation of Halogenated Darobactin Derivatives by Feeding Halogenated Tryptophan During Heterologous Expression and Production

[0235] For heterologous production of halogenated darobactin derivatives, for example commercially available 5-chloro-L-tryptophan, 7-chloro-L-tryptophan, 6-fluoro-L-tryptophan or 7-fluor-L-tryptophan can be fed periodically

in different concentrations (total end concentration can be for example between 0.5 mM and 10 mM) to the fermentation broth during cultivation of *E. coli* BL21-Gold (DE3) (Agilent Technologies). *E. coli* was previously transformed with pNOSodarA-E-DX, where X stands for any generated expression plasmid. The cultivation and feeding periods were performed as described below:

Fermentation/Cultivation of Host Cells

Day 1

[0236] Inoculation of a single clone of *E. coli* BL21-Gold (DE3) (Agilent Technologies) cells harbouring plasmid pNOSodarA-E-DX with modified darA gene, in 10 ml LB medium including 30 µg/ml Kanamycin (Kan30) and incubation overnight at 30° C., 180 rpm (Infors HT)

Day 2

[0237] Inoculation of 20 ml FM medium (1.254% K₂HPO₄, 0.231% KH₂PO₄, 0.4% D(+)-glucose, 0.1% NH₄Cl, 1.2% yeast extract, 0.5% NaCl and 0.0492% MgSO₄(*7H₂O), pH 7.1) including Kan30 and 1 mg/l VitaminB12 with 1% (v/v) of preculture

[0238] Preparation of halogenated tryptophan stocks by dissolving appropriate mass in dH₂O. For example: Dissolving of 10 mg 6-fluoro-L-tryptophan in 1.2 ml dH₂O to reach an end concentration of approximately 2.1 mM in 20 mL of production culture and storage of the solution at 4° C.

[0239] Addition of 0.4 µM IPTG (end concentration) to induce T7/ac promoter at OD₆₀₀ of 0.8 to 1.2 and first feeding of 1/6 (v/v) of prepared halogenated tryptophan stock

[0240] 2 h after first feeding, second feeding with 1/6 (v/v) of prepared halogenated tryptophan stock

Day 3

[0241] 18 h after last feeding, 1/6 (v/v) of prepared halogenated tryptophan stock was fed

[0242] 6 h after last feeding, 1/6 (v/v) of prepared halogenated tryptophan stock was fed

Day 4

[0243] 18 h after last feeding, 1/6 (v/v) of prepared halogenated tryptophan stock was fed

[0244] 6 h after last feeding, 1/6 (v/v) of prepared halogenated tryptophan stock was fed to reach total concentration of for example 0.5 mM to 10 mM, depending of chosen stock concentration.

[0245] Complete cultivation performed at 30° C. and 180 rpm (Infors HT)

[0246] Alternative for use of commercially available halogenated tryptophan for feeding to generate halogenated darobactin is the use of *E. coli* pUC18-zeo-mx8-corP-trpAB to produce halogenated tryptophan out of serine and halogenated indole. Vector pUC18-zeo-mx8-corP-trpAB was generated out of pUC18-zeo-mx8-corP (Pogorevc, D. et al. Production optimization and biosynthesis revision of coralpyronin A, a potent anti-filarial antibiotic. *Metab. Eng.* 55, 201-211; 10.1016/j.ymben.2019.07.010 (2019)) and trpAB gene of pSTB7 37845 (ATCC) after restriction digestion with NsiI and NdeI and followed ligation according to standard molecular biology techniques. Co-expression of pNOSodarA-E-DX and pUC18-zeo-mx8-corP-trpAB in *E. coli* BL21-Gold (DE3) (Agilent Technologies) lead to production of halogenated tryptophan through feeding of for example 10 mM serine and 5 to 10 mM desired indole. This system is flexible and the trpAB gene can also be cloned upstream darE into the BGC of Darobactin via unique restriction sites SpeI and PacI (R₁₁ and R₁₂, shown in Table

1). The trpAB gene, encodes tryptophan synthase enzymes TrpA and TrpB, catalyzing generation of tryptophan out of serine and indole. This allows modifications of darobactin with all commercially available or synthetically generated indoles that can be incorporated into darobactin during translation.

Separation and Retaining of Compounds

Day 5

[0247] In theory, after feeding with halogenated tryptophan, the appropriate halogenated tryptophan could be incorporated at each position where a tryptophan is assumed according to genetic code. For example, derivatives of Darobactin 9 with a 6-fluoro-L-tryptophan can carry the 6-fluoro-L-tryptophan at position 1, 3 or 7, at position 1 and 3, 1 and 7, 3 and 7, or at position 1, 3 and 7 of the respective core peptide amino acid (see FIG. 2). The naming of these halogenated compounds follows the pattern “darobactin derivative” (e.g. darobactin 9)-“position of fluorine on tryptophan according to IUPAC” (e.g. 6F for 6-fluoro-L-tryptophan), followed by the position of the halogenated tryptophan in the amino acid of the core peptide (e.g. 1, 3 or 7). As an example, darobactin 9-6F1 denotes Darobactin 9 with a 6-fluoro-L-tryptophan at position 1. In case of multiple halogenated tryptophans, positions are separated by hyphenation, e.g. darobactin 9-6F1-7 denoting Darobactin 9 with a 6-fluoro-L-tryptophan at positions 1 and 7. To check which new derivatives were produced, compounds were extracted and analyzed with HPLC-MS and HPLC-MS-MS detailed described below:

[0248] Harvest supernatant by centrifugation at 8,000 g for 10 min at 4° C.

[0249] Incubation of supernatant with 2 ml XAD16N resin (Sigma Aldrich) (w/v) under agitation for 2 h

[0250] Harvest of XAD16N resin and elution of bound compounds from XAD16N resin with 10 ml 80% Methanol under shaking for 1.5 h

[0251] Drying of eluate in a rotary evaporator

[0252] Solubilization of evaporated extract in 0.2 ml 50/50 methanol/water mixture; or in 0.2 ml 100% water

[0253] HPLC-MS analysis to verify the production of peptide compound of interest, e.g., darobactin 9-F1 or darobactin derivative; characteristic ions for darobactins in tandem MS mode are m/z 160.075 and a prominent [M-H₂O+2H]²⁺ ion.

[0254] As an example, production of darobactin 9-6F derivatives was performed according to described protocol by feeding 6-fluoro-L-tryptophan during heterologous expression and production. According to HPLC-MS and MS-MS analysis, darobactin 9-6F1, darobactin 9-6F7 and darobactin 9-6F1-7 were produced in high amounts. In some embodiments, darobactin 9-6F1 and 9-6F7 were preferably produced, especially darobactin 9-6F1. Further examples for the production of halogenated darobactin derivatives described herein are darobactin 9-7F1, darobactin 9-7F7, darobactin 9-5CI1 and 9-5CI7. Confirmation of production was achieved by mass, isotope pattern and MS-MS fragmentation pattern, showing incorporation of halogenated tryptophans, after or before purification according to described procedure.

Purification of Halogenated Darobactams

[0255] HPLC purification was carried out on a Waters Autopurifier (Eschborn, Germany) high pressure gradient system, equipped with 2545 binary gradient module, SFO system fluidics organizer, 2767 sample manager and a 2998 photodiode array detector coupled to a 3100 single quadrupole mass spectrometer operated in positive ion mode. Source and voltage settings for the MS were as follows: mass range, m/z 300-1000; scan duration, 1 s; points per Dalton, 4; capillary voltage, 3.5 kV; cone voltage, 30 V; extractor voltage 3 V; RF lens, 0.1 V; source temperature 120° C., desolvation temperature, 250° C.; desolvation gas flow, 400 L/h; cone gas flow, 50 L/h; ion counting threshold, 30. Separation was carried out on a Waters XBridge® Prep C-18 5 µm ODB™, 150×19 mm column using ACN+0.1% FA as B and H₂O+0.1% FA as A and a flow rate of 25 ml/min. Compound detection was carried out by detection of the [M+2H]²⁺.

[0256] Second purification step was carried out on a Thermo Scientific™ Dionex™ UltiMate™ 3000 SDLC low pressure gradient system on a Waters XSelect® Peptide CSH C18 OBD™ Prep 130 Å, 5 µm, 10 mm×250 mm column. Used eluents were H₂O+0.1% FA as A and ACN+0.1% FA as B, a flow rate of 6 ml/min and a column thermostated at 45° C. was used for further purification of peptide compounds of interest. Peptide compounds of interest, e.g. darobactin 9-6F1 can be detected by UV absorption at 254 nm or 280 nm. The two isomers darobactin 9-6F1 and darobactin 9-6F7 elute very close together, the first eluting isomer is the 6F7 variant, which is produced in less amounts, followed by darobactin 9-6F1, which is the main isomer. Exemplified compounds obtained by heterologous expression and production of the modified BGCs according to the present invention are shown in Table 10 below.

TABLE 10



Produced halogenated example compounds D9-6F1, D9-6F7, D9-6F1-7, D9-7F7, D9-5C11 and D9-5C17		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D9-6F1		512.21
D9-6F7		512.21

TABLE 10-continued





Produced halogenated example compounds D9-6F1, D9-6F7, D9-6F1-7, D9-7F7, D9-5Cl1 and D9-5Cl7		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D9-6F1-7		521.21
D9-7F7		512.21
D9-5Cl1		520.19

TABLE 10-continued

Produced halogenated example compounds D9-6F1, D9-6F7, D9-6F1-7, D9-7F7, D9-5Cl1 and D9-5Cl7		
Darobactin derivative	Chemical structure based on amino acid sequence of core peptide, mass and MS2 fragmentation pattern	Observed mass* (m/z) [M + 2H] ²⁺
D9-5Cl7		520.19

*All observed [M + 2H]²⁺ masses correlate with the expected [M + 2H]²⁺ masses.

Assessment of Antimicrobial Activity of Darobactin Derivatives

[0257] Crude extracts containing the respective darobactin derivatives obtained by the heterologous expression and separation protocol described above were tested for their activity against exemplified Gram-negative bacteria in standard microbroth dilution assays. The presence of darobactin derivatives in the crude extracts was verified by checking the expected masses via LC-HRMS.

[0258] All microorganisms were handled according to standard procedures as recommended by the depositor. Strains were obtained from the German Collection of Cell Cultures and Microorganisms (*Klebsiella pneumoniae* DSM-30104, *Acinetobacter baumannii* DSM-30007 and *A. baumannii* DSM-30008), the American Type Culture Collection (*Escherichia coli* ATCC-25922), or were part of our internal strain collection (*Pseudomonas aeruginosa* PAO1, PA14 and PA14ΔmexAB).

[0259] For the MIC assay, 10 μl of crude extract of each test sample, diluted in 50% methanol/H₂O, was added each into a well of the first row of a 96 well plate. After evaporation of the methanol/H₂O solvent, until only about 4 μl solvent was remaining in each well of the first row, 71 μl cation-adjusted Mueller-Hinton broth medium was added to each of said wells, so that each well in the first row of the 96 well plate had a volume of 75 μl. Then, 75 μl of cation-adjusted Mueller-Hinton broth medium was added to each well and the samples contained in the first row were mixed with the 75 μl medium/extract mixture. Starting from the 150 μl mixtures in the first row, serial dilutions of the respective mixtures were prepared in corresponding subsequent rows of the 96 well plate. Bacterial suspensions of each strain to be tested were diluted 1:100 in cation-adjusted

Mueller-Hinton broth to obtain a final inoculum of approximately 10⁶ CFU/mL. 75 μl of each bacterial suspension to be tested was added to the respective crude extract samples in the wells of the 96 well plate and the plates were incubated at either 30° C. or 37° C. for at least 24 h under static conditions. Plates were then checked visually and MIC was determined as lowest concentration where no visible growth was observed. In order to quantify the activity of the obtained darobactin derivatives, their relative activity in relation to the activity of darobactin A was determined. The amount of darobactin derivative present in the extract test sample was calculated based on the area under the curve (AUC) of the mass peak as automatically calculated by software Data Analysis 4.2 and extracted from the LC-MS measurements of each of the crude extract samples. Concentration factor of the tested extract was calculated by division of the concentration (100× caused by the extraction process from 50 ml culture to 0.5 ml extract) by the dilution factor in which visible growth of the respective strain was no longer observed. For example, if the darobactin derivative-containing crude extract in column 1 (of a standard 96 well plate) resulted in growth inhibition of the respective strain from wells A1 to E1, but not anymore in F1, the concentration factor score 0.42× was assigned to the darobactin derivative. If the darobactin derivative-containing crude extract in column 2 resulted in growth inhibition only from wells A2 to B2, but not anymore in C2, we assigned the concentration factor score 3.34×, meaning that the crude extract in column 1 showed stronger antibacterial activity (assuming that the compound concentration is similar). The results of activities of tested darobactin derivatives against exemplified Gram-negative bacteria as compared to the activity of darobactin A and D9 are summarized in Table 11 below.

TABLE 11

Activity of crude extracts of tested compounds against exemplified Gram negative bacteria					
Darobactin	Concentration factor of crude extract at which full growth inhibition is observed				AUC ratio
in crude extract	<i>K. pneumoniae</i> DSM-30104	<i>P. aeruginosa</i> PAO1	<i>E. coli</i> ATCC25922	<i>A. baumannii</i> DMS-30008	(derivative/ D9)
9	0.42x	0.11x	0.42x	0.21x	1
22	0.83x	0.05x	0.21x	0.05x	0.33
22*	0.83x	0.05x	0.21x	0.05x	0.33
23	0.83x	0.11x	0.21x	0.21x	0.38
23*	0.83x	0.11x	0.21x	0.21x	0.38
24	0.83x	0.05x	0.21x	0.21x	1.44
24*	0.83x	0.05x	0.21x	0.21x	1.44
25	3.34x	0.05x	3.34x	0.42x	1.89
25*	3.34x	0.05x	3.34x	0.42x	1.89
30	1.67x	0.11x	0.83x	0.83x	0.29
31	0.83x	0.05x	0.21x	0.05x	0.41
32	—	1.67x	6.67x	6.67x	0.02
32*	—	1.67x	6.67x	6.67x	0.02
33	6.67x	0.83x	1.67x	1.67x	0.14
33*	6.67x	0.83x	1.67x	1.67x	0.14
34	—	1.67x	3.34x	6.67x	0.13
34*	—	1.67x	3.34x	6.67x	0.13
P32	—	1.67x	6.67x	6.67x	0.02
P32*	—	1.67x	6.67x	6.67x	0.02
P33	6.67x	0.83x	1.67x	1.67x	0.14
P33*	6.67x	0.83x	1.67x	1.67x	0.14
P34	—	1.67x	3.34x	6.67x	0.13
P34*	—	1.67x	3.34x	6.67x	0.13
36	3.34x	0.05	0.83x	0.11x	0.31
36*	3.34x	0.05	0.83x	0.11x	0.31
37	1.67x	0.11x	0.42x	0.42x	1.43
37*	1.67x	0.11x	0.42x	0.42x	1.43
38	1.67x	0.11x	0.42x	0.42x	1.98
38*	1.67x	0.11x	0.42x	0.42x	1.98
39	0.42x	0.11x	0.42x	0.05x	0.52x

[0260] Selected examples (darobactins 9, 22, 23, 22*, 23*) were tested for their antibacterial activity in microbroth dilution assays as described above. Compounds were tested in serial dilution (0.03-64 µg/mL) and determined MICs are given in Table 12, which demonstrate superior activity of the new derivatives against *Acinetobacter baumannii* and *Pseudomonas aeruginosa*.

TABLE 12

Antibacterial activity (MIC) of pure darobactin derivatives 9, 22, 23, 22* and 23* in comparison to darobactin A						
Bacterium	Darobactin derivatives; MIC [µg/mL]					
	A	9	22	22*	23	23*
<i>A. baumannii</i> DSM-30008	4	1-2	0.25	0.25	1-2	1-2
<i>A. baumannii</i> DSM-30007	16	2	2	2	4	4
<i>E. coli</i> ATCC 25922	2	1-2	1-2	1-2	8	8
<i>K. pneumoniae</i> DSM-30104	2-4	1-2	2-4	2-4	4-8	4-8

TABLE 12-continued

Antibacterial activity (MIC) of pure darobactin derivatives 9, 22, 23, 22* and 23* in comparison to darobactin A						
Bacterium	Darobactin derivatives; MIC [µg/mL]					
	A	9	22	22*	23	23*
<i>P. aeruginosa</i> PAO1	0.5-1	0.125	0.5	0.5	1	1
<i>P. aeruginosa</i> PA14	16	2	2	2	4	4
<i>P. aeruginosa</i> PA14ΔmexAB	4	1	1-2	1-2	2-4	2-4

[0261] Activity data of exemplified darobactin derivatives 31, 32, 36 to 38, 32*, 36*-38*, 9-6F1, and 9-6F7 are summarized in Table 13 below. The compounds were tested for their antibacterial activity in microbroth dilution assays as described above. In particular, compounds were tested in serial dilution (0.03-64 µg/mL) and the determined MICs demonstrate comparable or superior microbial activity of the new derivatives.

TABLE 13

Antibacterial activity (MIC) of pure darobactin derivatives 31, 32, 36 to 38, 32*, 36*-38*, 9-6F1, and 9-6F7 in comparison to darobactin A												
Bacterium	Darobactin derivatives; MIC [μ g/mL]											
	A	31	32	36	37	38	32*	36*	37*	38*	9-6F1	9-6F7
<i>A. baumannii</i> DSM-30008	4	0.5-1	1	0.5	4-8	4	1	0.5	4-8	4	1	1
<i>A. baumannii</i> DSM-30007	16	2	1-2	2-4	16	8	1-2	2-4	16	8	2-4	2-4
<i>E. coli</i> ATCC 25922	2	1	2	2-4	4-8	8	2	2-4	4-8	8	1	0.5
<i>K. pneumoniae</i> DSM-30104	2-4	2-4	8	16	16-32	16-32	8	16	16-32	16-32	1	0.5-1
<i>P. aeruginosa</i> PAO1	0.5-1	0.25	2	0.25-0.5	2	1	2	0.25-0.5	2	1	1	0.5
<i>P. aeruginosa</i> PA14 DSM-19882	16	2-4	8	2-4	8-16	8-16	8	2-4	8-16	8-16	8	8
<i>P. aeruginosa</i> PA14AmexAB	4	1-2	4	2-4	8-16	8	4	2-4	8-16	8	4	4-8

[0262] As demonstrated above, darobactin derivatives described herein have an excellent antimicrobial activity against Gram-negative bacteria, and some show selectivity

against certain pathogens. Antibacterial activity of certain darobactin derivatives is superior against clinical isolates of *A. baumannii*, shown in Table 14 below.

TABLE 14

Antibacterial activity (MIC) of pure darobactin derivatives 9 and 22* in comparison to darobactin A and globomycin					
Bacterium	Darobactin derivatives; MIC [μ g/mL]				
	DA	D9	D22	D22*	Globomycin
<i>A. baumannii</i> 038 (OXA-23)	16	4	0.5	0.5	32
<i>A. baumannii</i> 045 (OXA-58)	16-32	8	0.5	0.5	64
<i>A. baumannii</i> 046 (OXA-40)	8-16	4	0.5	0.5	32
<i>A. baumannii</i> 047 (OXA-235)	16-32	4-8	0.25	0.25	64
<i>A. baumannii</i> 070 (NDM-1)	8	2	0.0625	0.0625	16
<i>A. baumannii</i> 054 (OXA-51-ISAba1)	16-32	8	0.125	0.125	>64

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 89

<210> SEQ ID NO 1

<211> LENGTH: 8131

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Biosynthetic Gene Cluster (BGC)

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1767)..(1769)

<223> OTHER INFORMATION: n is a, g, c or t
each nucleotide is a nucleotide which is part of a base triplet
coding for an amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1773)..(1784)

<223> OTHER INFORMATION: n is a, g, c or t
each nucleotide is a nucleotide which is part of a base triplet
coding for an amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (2073)..(2075)

<223> OTHER INFORMATION: n is a, g, c or t
each nucleotide is a nucleotide which is part of a base triplet

-continued

```

      coding for an amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (2079)..(2090)
<223> OTHER INFORMATION: n is a, g, c or t
      each nucleotide is a nucleotide which is part of a base triplet
      coding for an amino acid

<400> SEQUENCE: 1

cgcgcgcgat tctaaccggt ataagaaata aagtgagcg cgcgcgatta tccctaggtc      60
actgcccgtt ttccagtcgg gaaacctgtc gtgccagctg cattaatgaa tcggccaacg    120
cgcggggaga ggcggtttgc gtattgggcg ccaggggtgt tttcttttc accagtgaga    180
cgggcaacag ctgattgccc ttcaccgcct ggccctgaga gagttgcagc aagcgggtcca    240
cgctggtttg ccccgagcagg cgaaaatcct gtttgatggt ggtaacggc gggatataac    300
atgagctgtc ttccgtatcg tcgtatccca ctaccgagat gtccgcacca acgcgcagcc    360
cggaactcgt aatggcgcgc attgcgccc gcgccatctg atcgttggca accagcatcg    420
cagtgggaac gatgccctca ttcagcattt gcatggtttg ttgaaaaccg gacatggcac    480
tccagtcgcc ttcccggtcc gctatcggct gaatttgatt gcgagtgaga tatztatgcc    540
agccagccag acgcagacgc gccgagacag aacttaatgg gcccgctaac agcgcgattt    600
gctggtgacc caatgcgacc agatgtcca cgccagtcg cgtaccgtct tcatgggaga    660
aaataatact gttgatgggt gtcgtgtcag agacatcaag aaataacgcc ggaacattag    720
tgcaggcgag ttccacagca atggcaccct ggtcatccag cggatagtta atgatcagcc    780
cactgacgcg ttgcgcgaga agattgtgca ccgccgcttt acaggcttcg acgccgcttc    840
gttctaccat cgacaccacc acgctggcac ccagttgac ggcgcgagat ttaatcgccg    900
cgacaatttg cgacggcgcg tgcaggggca gactggaggt ggcaacgcca atcagcaacg    960
actgtttgcc cgccagttgt tgtgccacgc ggttgggaat gtaattcagc tccgccatcg   1020
ccgcttcacc tttttccgcg gttttcgag aaacgtggct ggctggttc accacgcggg   1080
aaacgggtctg ataagagaca ccggcatact ctgcgacatc gtataacggt actggtttca   1140
cattcaccac cctgaattga ctctcttccg ggcgctatca tgccataacc cgaaaggttt   1200
tgcgccatcc gatggtgtcc gggatctcga cgctctccct tatgcgactc ctgcattagc   1260
aagcagccca gtagtagggt gaggccgttg agcaccgccg ccgcaaggaa tggatgatgc   1320
aaggagatgg cgcccaacag tcccccgccc acggggcctg ccaccatacc cagccgaaa   1380
caagcgctca tgagcccgaa gtggcgagcc cgatcttccc catcggtgat gtcggcgata   1440
taggcgcgag caaccgcacc tgtggcgccg gtgatgccg ccacgatgcg tccggcgtag   1500
aggatcgaga tcgatctcga tcccgcgaaa ttaatacgac tcatatagg ggaattgtga   1560
gcggataaca attcccctct agaaaataat ttgtttaact ttaagaagga gatatacata   1620
tgcataatac cttaaatgaa accggtaaaa ctcaagaagc actcaattct cttgctgcat   1680
cattcaaaga gactgaactc tcaattactg ataaagcact aaacgaatta agcaataaac   1740
ctaagatccc tgagatcacg gctcggnnt ggnnnnnnnn nnnncaggaa atttaaagct   1800
tatcccatca gggtatttta tttctctgaa aaaacaatgt ccattgggat aaacaatcaa   1860
gttggttcagt taaaataaat aattttttga accttaatca tagggtaatc ctatgtatta   1920
tccgggatta ccctaaatac ttctataaag taaatataaa aatttgaata atacagataa   1980

```

-continued

gtttttttaa cctgaatgga aggtttgtat tttatcatta aacacaacca ccttttacca	2040
ctaataattat tggatttttt catttcaaga tgaaaatata atagatttca gtaactaaaa	2100
agggtaatag cctgtcccag caggtgttac tgatgcaggc aaataaaata gccctaatag	2160
gatgggcttt aagtaaggaa cggtagccat aactgcagtc ctgaggatcc ccaccctgca	2220
agggcccga accgcaaggt tccgggccct tcgcctttta tcagggttatt tgcggccgct	2280
ggacagcaag cgaaccgga ttgccagctg gggcgccctc tggttaagggt ggggaagccct	2340
gcaaagtaaa ctggatggct ttcttgccgc caaggatctg atggcgagg ggatcaagat	2400
ctgatcaaga gacaggataa ggaggtacag attatgaatg ctattaacat ctcagaatcc	2460
atgaaacgtt tcaatatatt ggtatcgata ctctcagtat caatagcaat attagcaatt	2520
tatttacttt ctactgcctt ctattcagat gctcgagttg ataactatta tcaagataat	2580
ggcaacattt accgggtgga aaccaccttt aatttgccca atggagaaaa agtgagatcg	2640
gcaaaatcgc cattgccgct catagacgag ctagaaaaag ataaaaggat aaatcgtgca	2700
gactactttt tcaaaactaa cacgacgatt gagttacaag gaaaaagat caccaaagtt	2760
cccggtgttg cagtcagtc tcaattcctg aataccctgt ctccctttta gaaaatcgat	2820
aatatcctgg gagcaaatga gatttacatc acaaaagagt tcaataatcg ctatctcggc	2880
ctggaaaatc ccaaaggaaa atctattgtt ctggataatg aaacctatat catcaaggat	2940
atcgtagaaa aacgtcatga ttcgagtcct aatatgaatg ctattatttc atttaagcca	3000
acacttatca aaaactacaa taaagaaata ttcaattggg atgatacaca tgcttatatt	3060
tttataaaac tctctgatga aaaatcaata tccaatcgcg atgcgcttct taatcatatt	3120
attgcaacca aagcccccaa tcttcaggc gctccattta ctgccagtga atttattcac	3180
ctatcattaa aaaaatcaaa agatattcat taccaagatg gattgcctga tgaaatttct	3240
attggtttat caaaggagat aatttatagc tcatatatcg cgttgcatt tatcttattt	3300
tccacattaa ccaactatta taatttaact tctgctgatt cagcagaaaa aagggatatt	3360
tatcagttaa agaaagcgt aggcgcttca actttcaata ttatttgcca ttactgcgg	3420
gccatgttga tcaaaataat cctttccacc cttatctttt tggtacttat cataactata	3480
tttaatgtct ctctctatat aaataatata ttaccctat tagaagcaga cagtttaata	3540
tatctaggga tattttcagt tgttattgcc tttatcttta tttcatcaat acatttttca	3600
ttctcattac gtttttatat ttatcaaaga aacagaaaaa tggatatgag atatgaatca	3660
acccttgctt cttgggttag gaaatcatca ttggctattc aattacttgt ttccggaata	3720
tctatttatt tagtcaccgg attaaccaat caatatattc acctgttgga ttacatact	3780
ccttatgata atgccaagac aatatcaata gcaattaata acgataataa taaaaaatca	3840
actattgatg aattaagcta tgagtttatt agaaaaata atacttcaa aataaccctc	3900
agtaattggc gaccatttga tatgtcaaga gaaaatatca ctattaatta ttccggccaa	3960
caaacacagga ataactatat ctcttcaa atgttattactg cggatgaaaa tttactgat	4020
gtgtggagaa tgaagatatt agccgggtgt gataaacgag tatatcagag tgataatgct	4080
gatgtagtcc atgcattagt gacaaaagaa tttttaaaac agaattggaat tctcaaatat	4140
gatgatattt ttaacaacta ctattgttac acaatggagg ataaaaagat ccaaatcaga	4200
tttattcaga ttatagatga tttcaattta ggtgctgtag atgaaccatt caggccaata	4260

-continued

gttgttttca tcaaaaaaga tcatggaaaa tatgcttcgc tgaattttaa caatatgaaa	4320
aacctttctc cggtaacttaa cgaactatca aaacagggtt ttgataataa tgaaattgat	4380
ttaaccagtc acttatttta ctcttattat gataattatt taaaagtaat aaaattatca	4440
tctgccgtcg ctattttctc ggttctatta ttagtcacgc cctgctttac taccagcatg	4500
actgacttta acgcgatgaa aaggagggtta agtataatgg aatctattgg tggttctatt	4560
tacaccaatc tcattcactt tttatcgaaa agtgcaccc cgataatcgc atcaattttc	4620
gtggcatata tatcaggaag aactatttta gtttatctgc ttgatgaata tagtgtgatt	4680
tacaccgatt cttttgtata cagtgcctac gccctatctg ctacaatgat attcaccgtt	4740
attataatga tgactaccta tttggtgaat tatagaaaat tatccgccac taaatttttg	4800
tgagattaat tatggatata gaaatcagaa aaaaaggcat acataacaga gagaaattac	4860
tcattctcat tttctttatc gcttttattt catttggtgt atttattatc tattatgtcg	4920
tttcggctaa agacgctttc gtgaatgaaa atgaagttac ccagtttcag gtcagcagca	4980
aagtagatag tgatattctg aaaaccaggg cagttgtggt tcctgataaa agtgtctcta	5040
tctcaaccga gataggggga atcggtaccg atataatgaa aaaaccatca caagatgttt	5100
taaagaatga agaaatcgtg aaattatcta attttaattt cacattgaat aattcttcga	5160
tgttagcgga cgtgacagat aaactcaata atttaataaa cataagaata aacttgcaat	5220
ctgactatcg ggatatcaat aaccgattct tagaagccgc caaagaacta aaagaagtag	5280
aagataaatt aaagcgatat cagcatcttg tacataaaaa ttatatttct aaagaaacta	5340
tgcccgattt aaggattaaa agagattatt ggcacgatgt ttatctatc tacaaaaaat	5400
taaaaaatga taaagatcgt gatatatcaa aacagttaaa ggaaatcgac gagtttgcg	5460
aaaaacagag aaaactctct gacatcatcg aaaatgggtt cgagcagctt tcaataaaat	5520
ccctatcgc cggaatatc agttcattag atttaatttt aggacaacgg ttaaacctg	5580
gcgataaaat agctattggt gatgatttat caaactttta tttgaaatca gaaatcaacg	5640
aatactattt aaacaaaata acataccatt catctgccag ttaaatctat aacaatggaa	5700
aaataccttt actggtgaaa ttaatttcat ctgaagtga taatggaaca ttcaaagtca	5760
gatttgaatt ggaagataaa aaaacaatta acttcaaacg ggggcaatcc gttgatgtca	5820
tgattaaact tgatgaagat cggaaaaatat tctccgtacc ttcactatg gtgttttcta	5880
ttgacaataa aaactatgtg tttgtttacc atccagataa aaaaattgcc cgacgtattc	5940
aagtttccac cggccaagat aatggttctg atattgagat caaaagtggt gttactgaag	6000
gacaaacgct cgttagcttt ggtaagaata aattagttaa taatgatata gtaaggattg	6060
aataaaaatga ttagcatgat gaatatctgt aaatcatata aaacgaaatt cattcaaaca	6120
aatgttttga atgacataaa tctcaatata gataaaggcg aatttatttc tataatggga	6180
gcgtcagggc cagggaaatc aacattactg aatgtcattg gcatgtttga aacaatcgat	6240
agcgggtcaat taacactgaa caatagcaat attttaacaa tgaaatatc tgaaaaata	6300
tcattcagac gagaatttat tgggttatatt tttcaatcat tcaatctgtt acccaatttg	6360

-continued

```

acggtatttg aaaacataga gttaccgta aaatacagag gtttctctaa aaaacagcgg 6420
aaaagcaaag tatttgatgc aataaatagt ttgggattag aaaatcggga aaatcataaa 6480
ccgatacaat tatctggtgg acaacagcaa cgtgttgcta tcgccagagc aatgatagca 6540
aaccaccacca ttttattagc tgatgaacca acggggaacc tggatagcgt aaatggtcag 6600
aatattttat cttcacttaa ggagctgaat gaaaacgcta cgacgattgt tatggtaacc 6660
cattccgtag aagcggcagc gttttctgac agaattataa cgatgaggga tggccacttg 6720
cttggttgag cataaataac ataataaagg agttctttat ggacacaata atccccataa 6780
aatattttaga ttcagacgaa tcacgattc ttaagaaatc atctaaaatt aactacaggc 6840
aattagcttg cagaattatc ggtgaaatct ccgccgaaaa aatattagat gatgatgaac 6900
tggttttata taataaagaa atcagtatac atttcagccc tgaaattatt aatgctaata 6960
aattagttgt ggtgtgtaaa gccaccaggc ttgcaattt aagatgcact tattgtcact 7020
cctgggcaga aggaaaagga aataccttaa cattcttcaa tttaatgcgt tccattcacc 7080
gtttcttctc cctaccgaat atcaagcgat ttgaattcgt ctggcatggc ggccaagtaa 7140
cgttgttgaa tgtaattac ttaagaaac tcactctggt acaggaacaa tttaaaaaac 7200
cggatcaagt tatcaccaat tcggtacaga caaatgccgt caatattcct gaagattggt 7260
tagtgttctc caaagggtatt ggaatggggg taggaataag cgttgatggt attccggaaa 7320
tacacgatag caggagatta gattacagag gaaggccaac atccccataa gtcgcggcaa 7380
gtatgaaaaa gttaagaagt tatggcatac cttacggtgc gcttatcgtc gtcgaccgcg 7440
atgtttatga atcaaatata gaaaaaatgc tctcttattt ttacgaaatc ggtttaacgg 7500
atattgaatt tctgaatatt gtcccagata accgatgcca gccgggtgat gatcctggag 7560
gaagttatat aacttaccat aactatatta atttccttct taagggtttc cgtgtctggt 7620
ggaatgggta tcaaggcaaa atcaatattc gcttggttga cggatttatt gacagtatca 7680
aatcgtecca aaagaaaatg tcagattggt attgggcggg taactgttct caggaaataa 7740
tcacattaga acctaatggt acggtatcag catgtgataa atatgttggt gctgaaggga 7800
ataattatgg ttcgattatt gataatgatc ttgggaattt actatctaaa tcaaatacaa 7860
ataaggatca tcttaagag gaaatggaat cttatgaaaa aatgcataa tgtaaatggt 7920
ttcatttggt taatggtgga tgcccacacg atcgagtgc caacaggaag cacaatccaa 7980
attatgatgg ttcattgtgt ggaaccggcg gtttgttgga gacaataaaa caaaccatcg 8040
cggcgtaaat tctaactagt tctgattaa ttaaccaccc tgcaaggggc cggaaccgca 8100
aggttcggg cccttcgct tttcgtcgac g 8131

```

```

<210> SEQ ID NO 2
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Photorhabdus kharii
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin A

<400> SEQUENCE: 2

```

```

tgg aac tgg tca aaa agc ttc
Trp Asn Trp Ser Lys Ser Phe
1 5

```

21

-continued

<210> SEQ ID NO 3
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Photorhabdus khanii

<400> SEQUENCE: 3

Trp Asn Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin B
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin B

<400> SEQUENCE: 4

tgg aac tgg acc aaa cga ttc 21
Trp Asn Trp Thr Lys Arg Phe
1 5

<210> SEQ ID NO 5
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 5

Trp Asn Trp Thr Lys Arg Phe
1 5

<210> SEQ ID NO 6
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin C
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin C

<400> SEQUENCE: 6

tgg tca tgg tca aga agc ttc 21
Trp Ser Trp Ser Arg Ser Phe
1 5

<210> SEQ ID NO 7
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 7

Trp Ser Trp Ser Arg Ser Phe
1 5

<210> SEQ ID NO 8
<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin D
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin D

<400> SEQUENCE: 8

tgg aac tgg tca aga agc ttc 21
Trp Asn Trp Ser Arg Ser Phe
1 5

<210> SEQ ID NO 9
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 9

Trp Asn Trp Ser Arg Ser Phe
1 5

<210> SEQ ID NO 10
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin E
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin E

<400> SEQUENCE: 10

tgg tca tgg tca aaa agc ttc 21
Trp Ser Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 11
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 11

Trp Ser Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 12
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 1
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 1

<400> SEQUENCE: 12

tgg tca tgg acc aaa cga ttc 21
Trp Ser Trp Thr Lys Arg Phe
1 5

-continued

<210> SEQ ID NO 13
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 13

Trp Ser Trp Thr Lys Arg Phe
1 5

<210> SEQ ID NO 14
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 2
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 2

<400> SEQUENCE: 14

tgg tca tgg acc aga agc ttc 21
Trp Ser Trp Thr Arg Ser Phe
1 5

<210> SEQ ID NO 15
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 15

Trp Ser Trp Thr Arg Ser Phe
1 5

<210> SEQ ID NO 16
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 4
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 4

<400> SEQUENCE: 16

tgg aac tgg acc aaa agc ttc 21
Trp Asn Trp Thr Lys Ser Phe
1 5

<210> SEQ ID NO 17
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 17

Trp Asn Trp Thr Lys Ser Phe
1 5

-continued

<210> SEQ ID NO 18
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 5
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 5

<400> SEQUENCE: 18

tgg tgc tgg tca aaa agc ttc 21
Trp Cys Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 19
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 19

Trp Cys Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 20
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 6
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 6

<400> SEQUENCE: 20

tgg aac tgg tgc aaa agc ttc 21
Trp Asn Trp Cys Lys Ser Phe
1 5

<210> SEQ ID NO 21
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 21

Trp Asn Trp Cys Lys Ser Phe
1 5

<210> SEQ ID NO 22
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 7
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 7

<400> SEQUENCE: 22

-continued

tgg tgc tgg tgc aaa agc ttc	21
Trp Cys Trp Cys Lys Ser Phe	
1 5	

<210> SEQ ID NO 23
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 23

Trp Cys Trp Cys Lys Ser Phe
1 5

<210> SEQ ID NO 24
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 8
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 8

<400> SEQUENCE: 24

tgg aac tgg tca aaa tgc ttc	21
Trp Asn Trp Ser Lys Cys Phe	
1 5	

<210> SEQ ID NO 25
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 25

Trp Asn Trp Ser Lys Cys Phe
1 5

<210> SEQ ID NO 26
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 9
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 9

<400> SEQUENCE: 26

tgg aac tgg tca aaa agc tgg	21
Trp Asn Trp Ser Lys Ser Trp	
1 5	

<210> SEQ ID NO 27
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 27

-continued

Trp Asn Trp Ser Lys Ser Trp
1 5

<210> SEQ ID NO 28
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 10
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 10

<400> SEQUENCE: 28

tgg aac tgg tca aaa agc tac 21
Trp Asn Trp Ser Lys Ser Tyr
1 5

<210> SEQ ID NO 29
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 29

Trp Asn Trp Ser Lys Ser Tyr
1 5

<210> SEQ ID NO 30
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 11
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 11

<400> SEQUENCE: 30

tgg cag tgg tca aaa agc ttc 21
Trp Gln Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 31
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 31

Trp Gln Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 32
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 12
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 12

-continued

<400> SEQUENCE: 32

tgg acc tgg tca aaa agc ttc 21
Trp Thr Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 33

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 33

Trp Thr Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 34

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 13

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (21)

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 13

<400> SEQUENCE: 34

tgg gca tgg tca aaa agc ttc 21
Trp Ala Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 35

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 35

Trp Ala Trp Ser Lys Ser Phe
1 5

<210> SEQ ID NO 36

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 14

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (21)

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 14

<400> SEQUENCE: 36

tgg aac tgg gca aaa agc ttc 21
Trp Asn Trp Ala Lys Ser Phe
1 5

<210> SEQ ID NO 37

<211> LENGTH: 7

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 37

Trp Asn Trp Ala Lys Ser Phe
1 5<210> SEQ ID NO 38
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 15
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 15

<400> SEQUENCE: 38

tgg aac tgg tca gca agc ttc 21
Trp Asn Trp Ser Ala Ser Phe
1 5<210> SEQ ID NO 39
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 39

Trp Asn Trp Ser Ala Ser Phe
1 5<210> SEQ ID NO 40
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 16
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 16

<400> SEQUENCE: 40

tgg aac tgg tca aaa gct ttc 21
Trp Asn Trp Ser Lys Ala Phe
1 5<210> SEQ ID NO 41
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 41

Trp Asn Trp Ser Lys Ala Phe
1 5<210> SEQ ID NO 42
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 17
<220> FEATURE:

-continued

```

<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 17

<400> SEQUENCE: 42

tgg aac tgg tca aaa agc gca                21
Trp Asn Trp Ser Lys Ser Ala
1                      5

<210> SEQ ID NO 43
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 43

Trp Asn Trp Ser Lys Ser Ala
1                      5

<210> SEQ ID NO 44
<211> LENGTH: 8131
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Biosynthetic Gene Cluster (BGC)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1773)..(1775)
<223> OTHER INFORMATION: n is a, g, c or t
      each nucleotide is a nucleotide which is part of a base triplet
      coding for an amino acid
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1779)..(1781)
<223> OTHER INFORMATION: n is a, g, c or t
      each nucleotide is a nucleotide which is part of a base triplet
      coding for an amino acid

<400> SEQUENCE: 44

cgcgcgcgat tctaaccggt ataagaaata aagtggagcg cggcgatta tccctaggtc      60
actgccccgt ttccagtcgg gaaacctgtc gtgccagctg cattaatgaa tcggccaacg      120
cgcggggaga ggcgggttgc gtattgggcg ccagggtggt tttcttttc accagtgaga      180
cgggcaacag ctgattgccc ttcaccgcct ggccctgaga gagttgcagc aagcgggtcca      240
cgctgggttg ccccgagcagg cgaaaatcct gtttgatggt ggtaaacggc gggatataac      300
atgagctgtc ttcggtatcg tcgtatccca ctaccgagat gtccgcacca acgcgcagcc      360
cggactcggg aatggcgcg c attgcgcca gcgccatctg atcgttggca accagcatcg      420
cagtggaac gatgccctca ttcagcattt gcatgggttg ttgaaaaccg gacatggcac      480
tccagtcgcc ttcccggttc gctatcggct gaatttgatt gcgagtgaga tatttatgcc      540
agccagccag acgcagacgc gccgagacag aacttaatgg gcccgctaac agcgcgattt      600
gtgtgtgacc caatgcgacc agatgtcca cgcccagtcg cgtaccgtct tcatgggaga      660
aaataatact gttgatgggt gtctggtcag agacatcaag aaataacgcc ggaacattag      720
tgcaggcagc ttccacagca atggcatcct ggtcatccag cggatagtta atgatcagcc      780
cactgacgcg ttgcgcgaga agattgtgca ccgcgcttt acaggcttcg acgccgcttc      840
gttctacat cgacaccacc acgctggcac ccagttgata ggcgcgagat ttaatcgccg      900
cgacaatttg cgacggcgcg tgcagggcc a gactggaggt ggcaacgcc atcagcaacg      960

```


-continued

actgtttgcc	cgccagttgt	tgtgccacgc	ggttggaat	gtaattcagc	tccgccatcg	1020
ccgcttccac	tttttccgcg	gttttcgcag	aaacgtggct	ggcctggttc	accacgcggg	1080
aaacggtctg	ataagagaca	ccggcatact	ctgcgacatc	gtataacggt	actggtttca	1140
cattcaccac	cctgaattga	ctctcttccg	ggcgctatca	tgcataccg	cgaaagggtt	1200
tgcgccattc	gatgggtgcc	gggatctcga	cgtctccct	tatgcgactc	ctgcattagg	1260
aagcagccca	gtagtaggtt	gaggccgttg	agcaccgcg	ccgcaaggaa	tggtgcatgc	1320
aaggagatgg	cgcccaacag	tcccccgcc	acggggcctg	ccaccatacc	cacgcgaaa	1380
caagcgctca	tgagcccgaa	gtggcgagcc	cgatcttccc	catcggtgat	gtcggcgata	1440
taggcgccag	caaccgcacc	tgtggcgccg	gtgatgcgg	ccacgatgcg	tccggcgtag	1500
aggatcgaga	tcgatctcga	tccgcgaaa	ttaatacgac	tactatagg	ggaattgtga	1560
gcggataaca	attccctct	agaaataatt	ttgtttaact	ttaagaagga	gatatacata	1620
tgcataatac	cttaaatgaa	accgttaaaa	ctcaagaagc	actcaattct	cttgctgcat	1680
cattcaaaga	gactgaactc	tcaattactg	ataaagcact	aaacgaatta	agcaataaac	1740
ctaagatccc	tgagatcacg	gcctggaact	ggnnnaaann	ntggcaggaa	atttaaagct	1800
tatcccatca	ggttatttta	tttctcgaa	aaaacaatgt	ccattgggat	aaacaatcaa	1860
gttgttcagt	taaaaataat	aattttttga	accttaatca	tagggtaatc	ctatgtatta	1920
tcggggatta	ccctaaatac	tttcataaag	taaatataaa	aatttgaata	atacagataa	1980
gtttttttaa	cctgaatgga	aggtttgat	tttatcatta	aacacaacca	cctttttacca	2040
ctaataattat	tggatttttt	catttcaaga	tgaaaatata	atagatttca	gtaactaaaa	2100
agggtaatag	cctgtcccg	cagggtgtac	tgatgcaggc	aaataaaata	gccctaatag	2160
gatgggcttt	aagtaaggaa	cggtagccat	aactgcagtc	ctgaggatcc	ccaccctgca	2220
agggcccgga	accgcaaggt	tccgggcct	tcgcctttta	tcagggttatt	tgcggccgct	2280
ggacagcaag	cgaaccggaa	tgccagctg	gggcgcctc	tggttaaggtt	gggaagccct	2340
gcaaagtaaa	ctggatggct	ttcttgccgc	caaggatctg	atggcgcagg	ggatcaagat	2400
ctgatcaaga	gacaggataa	ggaggtagag	attatgaatg	ctattaacat	ctcagaatcc	2460
atgaaacgtt	tcaatatatt	ggtatcgata	ctctcagtat	caatagcaat	attagcaatt	2520
tatttacttt	ctactgcctt	ctattcagat	gctcgagttg	ataactatta	tcaagataat	2580
ggcaacattt	accgggtgga	aaccaccttt	aatttgccca	atggagaaaa	agtgagatcg	2640
gcaaaatcgc	cattgcccgt	catagacgag	ctagaaaaag	ataaaaggat	aaatcgtgca	2700
gactactttt	tcaaaactta	cacgacgatt	gagttacaag	gaaaaaagat	caccaaaagt	2760
cccggtttg	cagtcagtca	tcaattcctg	aataccctgt	ctccctttaa	gaaaatcgat	2820
aatatcctgg	gagcaaatga	gatttacatc	acaaaagagt	tcaataatcg	ctatctcggc	2880
ctggaaaatc	ccaaaggaaa	atctattggt	ctggataatg	aaacctatat	catcaaggat	2940
atcgtagaaa	aacgtcatga	ttcgagtcct	aatatgaatg	ctattatttc	atttaagcca	3000
acacttatca	aaaactacaa	taaagaaata	ttcaattggt	atgatacaca	tgcttatatt	3060
tttataaaac	tctctgatga	aaaatcaata	tccaatcgcg	atgcgcttct	taatcatatt	3120
attgcaacca	aagcccccaa	tcttcaggc	gtccatttta	ctgccagtga	atttattcac	3180
ctatcattaa	aaaaatcaaa	agatattcat	taccaagatg	gattgcctga	tgaaatttct	3240

-continued

attggtttat	caaaggagat	aatttatagc	tcatatatcg	cggtcgcat	tatcttattt	3300
tccacattaa	ccaactatta	taatttaact	tctgctgatt	cagcagaaaa	aagggatatt	3360
tatcagttaa	agaagcgct	aggcgcttca	actttcaata	ttatttgcca	ttcactgccg	3420
gccatgttga	tcaaaataat	cctttccacc	cttatctttt	tggtacttat	cataactata	3480
tttaatgtct	ctctctatat	aaataatata	tttaccctat	tagaagcaga	cagtttaata	3540
tatctaggga	tattttcagt	tggtattgcc	tttatcttta	tttcatcaat	acatttttca	3600
ttctcattac	gtttttatat	ttatcaaaga	aacagaaaaa	tggtatgag	atatgaatca	3660
acccttgctt	cttggtttag	gaaatcatca	ttggctattc	aattacttgt	ttccggaata	3720
tctattttatt	tagtcaccgg	attaaccaat	caatatattc	accttggtga	tttacatact	3780
ccttatgata	atgccaaagc	aatatcaata	gcaattaata	acgataataa	taaaaaatca	3840
actattgatg	aattaagcta	tgagtttata	agaaaaatac	atacttcaaa	aataaccctc	3900
agtaattggc	gaccatttga	tatgtcaaga	gaaaatatca	ctattaatta	ttccggccaa	3960
caaaacagga	ataactatat	ctcttcaaat	gttattactg	cggtatgaaa	tttactgat	4020
gtgtggagaa	tgaagatatt	agccggtggt	gataaacgag	tatatcagag	tgataatgct	4080
gatgtagtcc	atgcattagt	gacaaaagaa	tttttaaaac	agaatggaat	tctcaaatat	4140
gatgatattt	ttaacaacta	ctattggtac	acaatggagg	ataaaaagat	ccaaatcaga	4200
tttattcaga	ttatagatga	tttcaattta	ggtgctgtag	atgaaccatt	caggccaata	4260
gtgtgtttca	tcaaaaaaga	tcatggaaaa	tatgcttcgc	tgaatttaaa	caatatgaaa	4320
aaacctttctc	cggtacttaa	cgaactatca	aaacagggtt	ttgataataa	tgaaattgat	4380
ttaaccagtc	acttatttta	ctcttattat	gataattatt	taaaagtaat	aaaattatca	4440
tctgcgctcg	ctattttctc	gggtctatta	ttagtcacgc	cctgctttac	taccagcatg	4500
actgacttta	acgcgatgaa	aaggagggtta	agtataatgg	aatctattgg	tggttctatt	4560
tacaccaatc	tcattcactt	tttatcgaaa	agtgtcatcc	cgataatcgc	atcaattttc	4620
gtggcatata	tatcaggaag	aactatttta	gtttatctgc	ttgatgaata	tagtgtgatt	4680
tacaccgatt	cttttgtata	cagtgtctatc	gccctatctg	ctacaatgat	attcaccggt	4740
attataatga	tgactaccta	tttggtgaat	tatagaaaat	tatccgccac	taaatttttg	4800
tgagattaat	tatggatata	gaaatcagaa	aaaaaggcat	acataacaga	gagaaattac	4860
tcattctcat	tttctttatc	gcttttattt	catttggtgt	atttattatc	tattatgtcg	4920
tttcggctaa	agacgctttc	gtgaatgaaa	atgaagttac	ccagtttcag	gtcagcagca	4980
aagtagatag	tgatattctg	aaaaccaggg	cagttgtggt	tcctgataaa	agtgtctcta	5040
tctcaaccga	gataggggga	atcggtaccg	atataatgaa	aaaaccatca	caagatgttt	5100
taaagaatga	agaaatcgtg	aaattatcta	attttaattt	cacattgaat	aattcttcga	5160
tgtagcgga	cgtgacagat	aaactcaata	atttaataaa	cataagaata	aacttgcaat	5220
ctgactatcg	ggatatcaat	aaccgattct	tagaagccgc	caaagaacta	aaagaagtag	5280
aagataaatt	aaagcgatat	cacgatcttg	tacataaaaa	ttatatttct	aaagaaacta	5340
tgtagcggtt	aaggattaaa	agagattatt	ggcagcatgt	ttatctattc	tacaaaaaat	5400
taaaaaatga	taaagatcgt	gatatatcaa	aacagttaaa	ggaaatcgac	gagtttgtcg	5460
aaaaacagag	aaaactctct	gacatcatcg	aaaatgggtt	cgagcagctt	tcaataaaat	5520

-continued

ccctatcgc	cgtaatatc	agttcattag	athtaatttt	aggacaacgg	ttaaaacctg	5580
gcgataaaat	agctattgtt	gatgatttat	caaactttta	ttttgaatca	gaaatcaacg	5640
aatactattt	aaacaaaata	acataccatt	catctgccag	tttaactctat	aacaatggaa	5700
aaataccttt	actggtgaaa	ttaatttcat	ctgaagtga	taatggaaca	ttcaaagtca	5760
gatttgaatt	ggaagataaa	aaaacaatta	acttcaaacg	ggggcaatcc	gttgatgtca	5820
tgattaaact	tgatgaagat	cggaaaatat	tctccgtacc	ttcatctatg	gtgttttcta	5880
ttgacaataa	aaactatgtg	ttgttttacc	atccagataa	aaaaattgcc	cgacgtattc	5940
aagtttccac	cggccaagat	aatggttctg	atattgagat	caaaagtggg	gttactgaag	6000
gacaaacgct	cgtagctttt	ggtaagaata	aattagttaa	taatgataca	gtaaggattg	6060
aataaaatga	ttagcatgat	gaatatctgt	aatcatata	aaacgaaatt	cattcaaaca	6120
aatgttttga	atgacataaa	tctcaatata	gataaaggcg	aatttatttc	tataatggga	6180
gcgtcagggt	cagggaatc	aacattactg	aatgtcattg	gcatgtttga	aacaatcgat	6240
agcggtaaat	taacactgaa	caatagcaat	atthtaacaa	tgaatatctc	tgaaaaata	6300
tcattcagac	gagaatttat	tggttatatt	tttcaatcat	tcaatctgtt	acccaatttg	6360
acggtatttg	aaaacataga	gttaccgtta	aaatacagag	gttttcctaa	aaaacagcgg	6420
aaaagcaaag	tatttgatgc	aataaatagt	tttggttag	aaaatcgga	aaatcataaa	6480
ccgatacaat	tatctggtgg	acaacagcaa	cgtgttgcta	tcgccagagc	aatgatagca	6540
aaacccacca	ttttattagc	tgatgaacca	acggggaacc	tgatagcgt	aatggtcag	6600
aatattttat	cttcacttaa	ggagctgaat	gaaaacggta	cgacgattgt	tatggtaacc	6660
cattccgtag	aagcggcagc	gttttctgac	agaatattaa	cgatgagga	tgccacttg	6720
cttggttgag	cataaataac	ataataaagg	agttctttat	ggacacaata	atccccataa	6780
aatattttaga	ttcagacgaa	tcatcgattc	ttaagaaatc	atctaaaatt	aactacaggc	6840
aattagcttg	cagaattatc	ggtgaaatct	ccgccgaaaa	aatattagat	gatgatgaac	6900
tggttttata	taataaagaa	atcagtatac	atthcagccc	tgaattatt	aatgctaata	6960
aattagtgtg	ggtgttgaaa	gccaccaggc	tttgcaattt	aagatgcact	tattgtcact	7020
cctgggcaga	aggaaaagga	aataccttaa	cattcttcaa	tttaatgcgt	tccattcacc	7080
gtttcttatac	cctaccgaat	atcaagcgat	ttgaattcgt	ctggcatggc	ggcgaagtaa	7140
cgttggtgaa	tgtaatttac	tttaagaaac	tcattctggt	acaggaacaa	tttaaaaaac	7200
cggatcaagt	tatcaccaat	tcggtacaga	caaatgccgt	caatattcct	gaagattggt	7260
tagtggtcct	caaaggtatt	ggaatggggg	taggaataag	cgttgatggt	attccggaaa	7320
tacacgtag	caggagatta	gattacagag	gaaggccaac	atccccataa	gtcgcggcaa	7380
gtatgaaaaa	gttaagaagt	tatggcatat	cttacgggtc	gcttatcgtc	gtcgaccgcg	7440
atgtttatga	atcaaatata	gaaaaaatgc	tctcttattt	ttacgaaatc	ggtttaacgg	7500
atattgaatt	tctgaatatt	gtcccagata	accgatgcca	gccgggtgat	gatcctggag	7560
gaagtatat	aacttaccat	aactatatta	atthcctttc	taagggtttc	cgtgtctggt	7620
ggaatgggta	tcaaggcaaa	atcaatattc	gcttggttga	cggatttatt	gacagtatca	7680
aatcgtccca	aaagaaaatg	tcagattgtt	attggggcgg	taactgttct	caggaaataa	7740
tcacattaga	acctaagtgt	acggtatcag	catgtgataa	atatgttggt	gctgaaggga	7800

-continued

ataattatgg ttcgattatt gataatgac ttgggaattt actatctaaa tcaaatacaa 7860
ataaggatca tcttaaagag gaaatggaat cttatgaaaa aatgcacaa tgtaaatggt 7920
ttcatttgtg taatgggtga tgcccacacg atcgagtgcac caacaggaag cacaatccaa 7980
attatgatgg ttcattgtgt ggaaccggcg gtttggtgga gacaataaaa caaacatcg 8040
cggcgtaaat tctaactagt tcctgattaa ttaaccaccc tgcaagggcc cggaaccgca 8100
agggtccggg cccttcgect ttctgctgac g 8131

<210> SEQ ID NO 45
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward Primer - pridar_fw for overlap
extension PCR

<400> SEQUENCE: 45

ggtgatgtcg gcgatatagg 20

<210> SEQ ID NO 46
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Forward Primer - pr3dartrp_fw for overlap
extension PCR

<400> SEQUENCE: 46

tggcaggaaa tttaaagctt atcccat 27

<210> SEQ ID NO 47
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr4dar_rv for overlap
extension PCR

<400> SEQUENCE: 47

tggggatcct caggactgca g 21

<210> SEQ ID NO 48
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar22_rv containing
mutations of D22

<400> SEQUENCE: 48

aaatttcctg ccaccgttta gtccagttcc 30

<210> SEQ ID NO 49
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar23_rv containing
mutations of D23

<400> SEQUENCE: 49

-continued

aaatttcctg ccaatgttta gtccagttcc 30

<210> SEQ ID NO 50
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar24_rv containing
mutations of D24

<400> SEQUENCE: 50

aaatttcctg ccatgattta gtccagttcc 30

<210> SEQ ID NO 51
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar25_rv containing
mutations of D25

<400> SEQUENCE: 51

aaatttcctg ccacgcttta gtccagttcc 30

<210> SEQ ID NO 52
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar30_rv containing
mutations of D30

<400> SEQUENCE: 52

aaatttcctg ccatgatttc gccagttcc 30

<210> SEQ ID NO 53
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar31_rv containing
mutations of D31

<400> SEQUENCE: 53

aaatttcctg ccacggtttc gccagttcc 30

<210> SEQ ID NO 54
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar32_rv containing
mutations of D32

<400> SEQUENCE: 54

aaatttcctg ccacggttta caccagttcc 30

<210> SEQ ID NO 55
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar33_rv containing
mutations of D33

-continued

<400> SEQUENCE: 55

aaatttcctg ccatgattta caccagttcc

30

<210> SEQ ID NO 56

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Reverse Primer - pr2dar34_rv containing mutations of D34

<400> SEQUENCE: 56

aaatttcctg ccacgcttta caccagttcc

30

<210> SEQ ID NO 57

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Reverse Primer - pr2dar36_rv containing mutations of D36

<400> SEQUENCE: 57

aaatttcctg ccatttttta gtccagttcc

30

<210> SEQ ID NO 58

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Reverse Primer - pr2dar37_rv containing mutations of D37

<400> SEQUENCE: 58

aaatttcctg ccaagttttt gaccagttcc

30

<210> SEQ ID NO 59

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Reverse Primer - pr2dar38_rv containing mutations of D38

<400> SEQUENCE: 59

aaatttcctg ccaagtttta gtccagttcc

30

<210> SEQ ID NO 60

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 9

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1) .. (21)

<223> OTHER INFORMATION: Propetide core sequence - Darobactin 9

<400> SEQUENCE: 60

tgg aac tgg tca aaa agc tgg

21

Trp Asn Trp Ser Lys Ser Trp

1

5

<210> SEQ ID NO 61

-continued

<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 61

Trp Asn Trp Ser Lys Ser Trp
1 5

<210> SEQ ID NO 62
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 22

<400> SEQUENCE: 62

tgg aac tgg act aaa cgg tgg 21
Trp Asn Trp Thr Lys Arg Trp
1 5

<210> SEQ ID NO 63
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 63

Trp Asn Trp Thr Lys Arg Trp
1 5

<210> SEQ ID NO 64
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 23

<400> SEQUENCE: 64

tgg aac tgg act aaa cat tgg 21
Trp Asn Trp Thr Lys His Trp
1 5

<210> SEQ ID NO 65
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 65

Trp Asn Trp Thr Lys His Trp
1 5

<210> SEQ ID NO 66
<211> LENGTH: 21

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 24

<400> SEQUENCE: 66

tgg aac tgg act aaa tca tgg 21
Trp Asn Trp Thr Lys Ser Trp
1 5

<210> SEQ ID NO 67
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 67

Trp Asn Trp Thr Lys Ser Trp
1 5

<210> SEQ ID NO 68
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 25

<400> SEQUENCE: 68

tgg aac tgg act aaa gcg tgg 21
Trp Asn Trp Thr Lys Ala Trp
1 5

<210> SEQ ID NO 69
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 69

Trp Asn Trp Thr Lys Ala Trp
1 5

<210> SEQ ID NO 70
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 30

<400> SEQUENCE: 70

tgg aac tgg gcg aaa tca tgg 21
Trp Asn Trp Ala Lys Ser Trp
1 5

-continued

<210> SEQ ID NO 71
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 71

Trp Asn Trp Ala Lys Ser Trp
1 5

<210> SEQ ID NO 72
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 31

<400> SEQUENCE: 72

tgg aac tgg gcg aaa cgg tgg 21
Trp Asn Trp Ala Lys Arg Trp
1 5

<210> SEQ ID NO 73
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 73

Trp Asn Trp Ala Lys Arg Trp
1 5

<210> SEQ ID NO 74
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 32

<400> SEQUENCE: 74

tgg aac tgg tgt aaa cgg tgg 21
Trp Asn Trp Cys Lys Arg Trp
1 5

<210> SEQ ID NO 75
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 75

Trp Asn Trp Cys Lys Arg Trp
1 5

-continued

<210> SEQ ID NO 76
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 33

<400> SEQUENCE: 76

tgg aac tgg tgt aaa tca tgg 21
Trp Asn Trp Cys Lys Ser Trp
1 5

<210> SEQ ID NO 77
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 77

Trp Asn Trp Cys Lys Ser Trp
1 5

<210> SEQ ID NO 78
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 34

<400> SEQUENCE: 78

tgg aac tgg tgt aaa gcg tgg 21
Trp Asn Trp Cys Lys Ala Trp
1 5

<210> SEQ ID NO 79
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 79

Trp Asn Trp Cys Lys Ala Trp
1 5

<210> SEQ ID NO 80
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 36

<400> SEQUENCE: 80

-continued

tgg aac tgg act aaa aaa tgg 21
Trp Asn Trp Thr Lys Lys Trp
1 5

<210> SEQ ID NO 81
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 81

Trp Asn Trp Thr Lys Lys Trp
1 5

<210> SEQ ID NO 82
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 37

<400> SEQUENCE: 82

tgg aac tgg tca aaa act tgg 21
Trp Asn Trp Ser Lys Thr Trp
1 5

<210> SEQ ID NO 83
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 83

Trp Asn Trp Ser Lys Thr Trp
1 5

<210> SEQ ID NO 84
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Propetide core sequence
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 38

<400> SEQUENCE: 84

tgg aac tgg act aaa act tgg 21
Trp Asn Trp Thr Lys Thr Trp
1 5

<210> SEQ ID NO 85
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 85

-continued

Trp Asn Trp Thr Lys Thr Trp
1 5

<210> SEQ ID NO 86
 <211> LENGTH: 8131
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Biosynthetic Gene Cluster (BGC)
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1773)..(1775)
 <223> OTHER INFORMATION: n is a, g, c or t
 each nucleotide is a nucleotide which is part of a base
 triplet coding for an amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1776)..(1778)
 <223> OTHER INFORMATION: each nucleotide is a nucleotide which is part
 of a base triplet coding for arginine or lysine, including agg,
 aga, aag, aaa, cga, cgc, cgg and cgt
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1779)..(1781)
 <223> OTHER INFORMATION: n is a, g, c or t
 each nucleotide is a nucleotide which is part of a base triplet
 coding for an amino acid
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1782)..(1784)
 <223> OTHER INFORMATION: each nucleotide is a nucleotide which is part
 of a base triplet coding for tryptophan, alanine or histidine,
 including tgg, gca, gcc, gcg, gct, cac and cat
 <400> SEQUENCE: 86
 cgcgcgcgat tctaaccggt ataagaaata aagtggagcg cggcgcgatta tccctaggtc 60
 actgcccgcgt ttccagtcgg gaaacctgtc gtgccagctg cattaatgaa tcggccaacg 120
 cgcggggaga ggcggtttgc gtattgggcg ccagggtggt tttcttttc accagtgaga 180
 cgggcaacag ctgattgcc ttcaccgcct ggccctgaga gagttgcagc aagcgggtcca 240
 cgctgggttg ccccgagcagg cgaaaatcct gtttgatggt ggtaaacggc gggatataac 300
 atgagctgtc ttccgtatcg tcgtatccca ctaccgagat gtccgcacca acgcgcagcc 360
 cggactcggg aatggcgcg attgcgcca gcgccatctg atcgttggca accagcatcg 420
 cagtgggaac gatgccctca ttcagcattt gcatgggttg ttgaaaaccg gacatggcac 480
 tccagtcgcc ttcccggtcc gctatcggct gaatttgatt gcgagtgaga tatttatgcc 540
 agccagccag acgcagacgc gccgagacag aacttaatgg gcccgctaac agcgcgattt 600
 gctggtgacc caatgcgacc agatgctcca cgcccagtcg cgtaccgtct tcatgggaga 660
 aaataatact gttgatgggt gtcgtgtcag agacatcaag aaataacgcc ggaacattag 720
 tgcaggcagc ttccacagca atggcaccct ggtcatccag cggatagtta atgatcagcc 780
 cactgacgcy ttgcgcgaga agattgtgca ccgccgcttt acaggcttcg acgccgcttc 840
 gttctaccat cgacaccacc acgctggcac ccagttgatc ggcgcgagat ttaatcgccg 900
 cgacaatttg cgacggcgcy tgcagggccca gactggaggt ggcaacgcca atcagcaacg 960
 actgtttgcc cgccagtgtg tgtgccacgc ggttgggaat gtaattcagc tccgccatcg 1020
 ccgcttccac tttttccgc gttttcgag aaacgtggct ggctgggttc accacgcggy 1080
 aaacggtctg ataagagaca ccggcatact ctgcgacatc gtataacggt actggtttca 1140
 cattcaccac cctgaattga ctctcttcg ggcgctatca tgccataccg cgaaagggtt 1200

-continued

tgcgccattc gatggtgtcc gggatctcga cgctctccct tatgcgactc ctgcattagg	1260
aagcagccca gtagtaggtt gaggccgttg agcaccgccg ccgcaaggaa tggatgcattgc	1320
aaggagatgg cgcccaacag tcccccgccc acggggcctg ccaccatacc caccgcgaaa	1380
caagcgctca tgagcccgaa gtggcgagcc cgatcttccc catcggtgat gtcggcgata	1440
taggcgccag caaccgcacc tgtggcgccg gtgatgccgg ccacgatgcg tccggcgtag	1500
aggatcgaga tcgatctcga tcccgcgaaa ttaatacgac tcaactatagg ggaattgtga	1560
gcggataaca attccctctc agaaataatt ttgtttaact ttaagaagga gatatacata	1620
tgcataatac cttaaatgaa accgttaaaa ctcaagaagc actcaattct cttgctgcat	1680
cattcaaaga gactgaactc tcaattactg ataaagcact aaacgaatta agcaataaac	1740
ctaagatccc tgagatcacg gcttggaact gnnnnnnnnn nnnncaggaa atttaaagct	1800
tatcccatca gggtatttta tttctctgaa aaaacaatgt ccattgggat aaacaatcaa	1860
gttgttcagt taaaaataat aattttttga accttaatac tagggtaatc ctatgtatta	1920
tccgggatta ccctaaatac ttccataaag taaatataaa aatttgaata atacagataa	1980
gtttttttta cctgaatgga aggtttgtat tttatcatta aacacaacca ccttttacca	2040
ctaataattat tggatttttt catttcaaga tgaaaaata atagatttca gtaactaaaa	2100
agggtaatat cctgtcccag caggtgttac tgatgcagcg aaataaaata gccctaatag	2160
gatgggcttt aagtaaggaa cggtagccat aactgcagtc ctgaggatcc ccaccctgca	2220
agggcccgga accgcaaggt tccgggccct tcgcctttta tcagggttatt tgcggccgct	2280
ggacagcaag cgaaccgga ttgccagctg gggcgccctc tggttaaggtt gggaagccct	2340
gcaaagtaaa ctggatggct ttcttgccgc caaggatctg atggcgcagg ggatcaagat	2400
ctgatcaaga gacaggataa ggaggtacag attatgaatg ctattaacat ctcaaatcc	2460
atgaaacggt tcaatatatt ggtatcgata ctctcagtat caatagcaat attagcaatt	2520
tatttacttt ctactgcctt ctattcagat gctcgagttg ataactatta tcaagataat	2580
ggcaacattt accgggtgga aaccaccttt aatttgccca atggagaaaa agtgagatcg	2640
gcaaaatcgc cattgccgct catagacgag ctagaaaaag ataaaaggat aaatcgtgca	2700
gactactttt tcaaaactaa cagcagcatt gagttacaag gaaaaaagat caccaaagtt	2760
cccgtgtttg cagtcagtc tcaattcctg aataccctgt ctccctttta gaaaatcgat	2820
aatatcctgg gagcaaatga gatttacatc acaaaagagt tcaataatcg ctatctcggc	2880
ctggaaaatc ccaaaggaaa atctattgtt ctggataatg aaacctatat catcaaggat	2940
atcgtagaaa aacgtcatga ttcgagcttt aatatgaatg ctattatttc atttaagcca	3000
acacttatca aaaactacaa taaagaaata ttcaattggg atgatacaca tgcttatatt	3060
tttataaaac tctctgatga aaaatcaata tccaatcgcg atgcgcttct taatcatatt	3120
attgcaacca aagcccccga tcttcaggc gctccattta ctgccagtga atttattcac	3180
ctatcattaa aaaaatcaaa agatattcat taccaagatg gattgcctga tgaaatttct	3240
attggtttat caaaggagat aatttatagc tcatatatcg cgttcgcatt tatcttattt	3300
tccacattaa ccaactatta taatttaact tctgctgatt cagcagaaaa aagggatatt	3360
tatcagttaa agaaagcgt aggcgcttca actttcaata ttatttgcca ttcactgccg	3420
gccatgttga tcaaaataat ctttccacc ctatctttt tgttacttat cataactata	3480

-continued

tttaatgtct	ctctctatat	aaataatata	tttacctat	tagaagcaga	cagtttaata	3540
tatctagga	tattttcagt	tggtattgcc	tttatcttta	tttcatcaat	acatttttca	3600
ttctcattac	gttttttatat	ttatcaaaaga	aacagaaaaa	tggtatagag	atatgaatca	3660
accttgctct	cttggttttag	gaaatcatca	ttggctattc	aattacttgt	ttccggaata	3720
tctatttatt	tagtcaccgg	attaaccaat	caatatattc	accttggtga	ttacatact	3780
ccttatgata	atgccaagac	aatatcaata	gcaattaata	acgataataa	taaaaaatca	3840
actattgatg	aattaagcta	tgagtttata	agaaaaatac	atacttcaaa	aataaccctc	3900
agtaattggc	gaccatttga	tatgtcaaga	gaaaatatca	ctattaatta	ttccggccaa	3960
caaaacagga	ataactatat	ctcttcaaat	gttattactg	cggatgaaaa	tttactgat	4020
gtgtggagaa	tgaagatatt	agccggtgtg	gataaacgag	tatatcagag	tgataatgct	4080
gatgtagtcc	atgcattagt	gacaaaagaa	tttttaaaac	agaatggaat	tctcaaatat	4140
gatgatattt	ttaacaacta	ctattggtac	acaatggagg	ataaaaagat	ccaaatcaga	4200
tttattcaga	ttatagatga	tttcaattta	ggtgctgtag	atgaaccatt	caggccaata	4260
gttgttttca	tcaaaaaaga	tcatggaaaa	tatgcttcgc	tgaatttaaa	caatatgaaa	4320
aacctttctc	cggactttaa	cgaactatca	aaacagggtt	tgataataa	tgaaattgat	4380
ttaaccagtc	acttatttta	ctcttattat	gataattatt	taaaagtaat	aaaattatca	4440
tctgccgtcg	ctattttctc	ggttctatta	ttagtcatcg	cctgctttac	taccagcatg	4500
actgacttta	acgcgatgaa	aaggagggtta	agtataatgg	aatctattgg	tggttctatt	4560
tacaccaatc	tcattcactt	tttatcgaaa	agtgtcatcc	cgataatcgc	atcaattttc	4620
gtggcatata	tatcaggaag	aactatttta	gtttatctgc	ttgatgaata	tagtgtgatt	4680
tacaccgatt	cttttgtata	cagtgtctatc	gccctatctg	ctacaatgat	attcaccggt	4740
attataatga	tgactacctt	tttgggtgaat	tatagaaaat	tatccgccac	taaatttttg	4800
tgagattaat	tatggatata	gaaatcagaa	aaaaaggcat	acataacaga	gagaaattac	4860
tcattctcat	tttctttatc	gcttttattt	catttggtgt	atttattatc	tattatgtcg	4920
tttcggctaa	agacgctttc	gtgaatgaaa	atgaagttac	ccagtttcag	gtcagcagca	4980
aagtagatag	tgatattctg	aaaaccaggg	cagttgtggt	tcctgataaa	agtgctctta	5040
tctcaaccga	gataggggga	atcggtaccg	atataatgaa	aaaaccatca	caagatgttt	5100
taaagaatga	agaaatcgtg	aaattatcta	attttaattt	cacattgaat	aattcttcga	5160
tgtagcgga	cgtgacagat	aaactcaata	atttaataaa	cataagaata	aacttgcaat	5220
ctgactatcg	ggatatacat	aaccgattct	tagaagccgc	caaagaacta	aaagaagtag	5280
aagataaatt	aaagcgatat	cacgatcttg	tacataaaaa	ttatatctct	aaagaaacta	5340
tgtagcgatt	aaggattaaa	agagattatt	ggcacgatgt	ttatctattc	tacaaaaaat	5400
taaaaaatga	taaagatcgt	gatatatcaa	aacagttaaa	ggaaatcgac	gagtttgtcg	5460
aaaaacagag	aaaactctct	gacatcatcg	aaaatgggtt	cgagcagctt	tcaataaaat	5520
cccttatcgc	cggtaatatc	agttcattag	atttaatttt	aggacaacgg	ttaaaacctg	5580
gcgataaaat	agctattggt	gatgatttat	caaactttta	ttttgaatca	gaaatcaacg	5640
aatactattt	aaacaaaata	acataccatt	catctgccag	tttaactctat	aacaatggaa	5700
aaataccttt	actggtgaaa	ttaatttcat	ctgaagtga	taatggaaca	ttcaagtca	5760

-continued

gatttgaatt ggaagataaa aaaacaatta acttcaaacg ggggcaatcc gttgatgtca	5820
tgattaaact tgatgaagat cggaaaaatat tctccgtacc ttcactatg gtgttttcta	5880
ttgacaataa aaactatgtg tttgtttacc atccagataa aaaaattgcc cgacgtattc	5940
aagtttccac cggccaagat aatggttctg atattgagat caaaagtggt gttactgaag	6000
gacaaacgct cgtagctttt ggtaagaata aattagttaa taatgatata gtaaggattg	6060
aataaaatga ttagcatgat gaatatctgt aaatcatata aaacgaaatt cattcaaaca	6120
aatgttttga atgacataaa tctcaatata gataaaggcg aatttatttc tataatggga	6180
gcgtcagggg cagggaatc aacattactg aatgtcattg gcattgttga aacaatcgat	6240
agcggtaaat taacactgaa caatagcaat attttaacaa tgaaatattc tgaaaaata	6300
tcattcagac gagaatttat tggttatatt tttcaatcat tcaatctgtt acccaatttg	6360
acggtatttg aaaacataga gttaccgtta aaatacagag gttttcctaa aaaacagcgg	6420
aaaagcaaaag tatttgatgc aataaatagt tttggattag aaaatcggga aaatcataaa	6480
ccgatacaat tatctggtgg acaacagcaa cgtgttgcta tcgccagagc aatgatagca	6540
aacccccacca ttttattagc tgatgaacca acggggaacc tggatagcgt aaatggtcag	6600
aatattttat cttcacttaa ggagctgaat gaaaacggta cgacgattgt tatggtaacc	6660
cattccgtag aagcggcagc gttttctgac agaattataa cgatgaggga tggccacttg	6720
cttgtttgag cataaataac ataataaagg agttctttat ggacacaata atccccataa	6780
aatattttaga ttcagacgaa tcatcgattc ttaagaaatc atctaaaatt aactacaggc	6840
aattagcttg cagaattatc ggtgaaatct ccgccgaaaa aatattagat gatgatgaac	6900
tggttttata taataaagaa atcagtatac atttcagccc tgaaattatt aatgctaata	6960
aattagtgtg ggtgtgtaaa gccaccaggc tttgcaattt aagatgcact tattgtcact	7020
cctgggcaga aggaaaagga aataccttaa cattcttcaa tttaatgcgt tccattcacc	7080
gtttcttato cctaccgaat atcaagcgat ttgaattcgt ctggcatggc ggcgaaagtaa	7140
cgttgttgaa tgtaattac tttaaagaaac tcatctggtt acaggaacaa tttaaaaaac	7200
cggatcaagt tatcaccaat tcggtacaga caaatgccgt caatattcct gaagattggt	7260
tagtgttcct caaaggtatt ggaatggggg taggaataag cgttgatggt attccggaaa	7320
tacacgatag caggagatta gattacagag gaaggccaac atccccataa gtcgcggcaa	7380
gtatgaaaaa gttaagaagt tatggcatac cttacggtgc gcttatcgtc gtcgaccgcg	7440
atgtttatga atcaaatata gaaaaaatgc tctcttattt ttacgaaatc ggtttaacgg	7500
atattgaatt tctgaatatt gtcccagata accgatgcca gccgggtgat gatcctggag	7560
gaagttatat aacttaccat aactatatta atttcctttc taagggtttc cgtgtctggt	7620
ggaatgggta tcaaggcaaa atcaatattc gcttgtttga cggatttatt gacagtatca	7680
aatcgtccca aaagaaaaatg tcagattggt attggggcgg taactgttct caggaaataa	7740
tcacattaga acctaattggt acggtatcag catgtgataa atatgttggt gctgaaggga	7800
ataattatgg ttcgattatt gataatgatc ttgggaattt actatctaaa tcaaatacaa	7860
ataaggatca tottaaagag gaaatggaat cttatgaaaa aatgcacaa tgtaaatggt	7920

-continued

```

ttcatttggtg taatggtgga tgccacacg atcgagtgac caacaggaag cacaatccaa 7980
attatgatgg ttcattgtgt ggaaccggcg gtttggtgga gacaataaaa caaaccatcg 8040
cgcgtaaat tctaactagt tctgattaa ttaaccaccc tgcaagggcc cggaaccgca 8100
aggttcggg cccttcgcct ttcgtcgac g 8131

```

```

<210> SEQ ID NO 87
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Peptide Derivative - Darobactin 39
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(21)
<223> OTHER INFORMATION: Propetide core sequence - Darobactin 39

```

```

<400> SEQUENCE: 87

```

```

tgg aac tgg act cgg cgg tgg 21
Trp Asn Trp Thr Arg Arg Trp
1 5

```

```

<210> SEQ ID NO 88
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

```

```

<400> SEQUENCE: 88

```

```

Trp Asn Trp Thr Arg Arg Trp
1 5

```

```

<210> SEQ ID NO 89
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Reverse Primer - pr2dar39_rv containing
mutations of D39

```

```

<400> SEQUENCE: 89


```

```

aaatttctcg ccaccgccga gtccagttcc a 31


```

1. Compound of formula (I) or a salt thereof:




wherein

R^1 , R^2 , and R^4 are independently selected from H, CH_3 , CH_2OH , one of the following groups:




$-CH_2-SR^{1,4}$, wherein $R^{1,4}$ is independently selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted; or $-CH_2-Ind$, wherein Ind is an optionally substituted indole group;

R^3 is selected from H, OH, SH, COOH, $CONH_2$, one of the following groups:



-continued



$-SR^{3,4}$, wherein $R^{3,4}$ is selected from an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted;

or -Ind, wherein Ind is an optionally substituted indole group;

R^5 is a group of formula $-CH_2-Ind$, wherein Ind is an optionally substituted indole group

R^6 is a hydrogen atom or a methyl group (especially a hydrogen atom);

R^7 is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom);

R^8 is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group; and

m is an integer of from 0 to 3 (especially 0 or 1; preferably 0);


$R^{6,4}$ is a hydrogen atom or a methyl group (especially a hydrogen atom);

$R^{7,4}$ is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom);

$R^{8,4}$ is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group; and


p is an integer of from 0 to 3 (especially 0 or 1; preferably 0).

2. The compound or a salt thereof according to claim 1, wherein the group of formula $-\text{CH}_2\text{-Ind}$ is a group of the following formula:




wherein R^9 is a hydrogen atom, a halogen atom, a hydroxy group or a methoxy group (especially a hydrogen atom); R^{10} is a hydrogen atom or a methyl group (especially a hydrogen atom); R^{11} is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group (especially a halogen atom, preferably fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).

3. The compound or a salt thereof according to claim 1, wherein the group of formula $-\text{CH}_2\text{-SR}^{1,4}$ is:




4. The compound or a salt thereof according to claim 1, wherein R^1 is selected from the following groups: $-\text{CH}_2\text{-CONH}_2$, $-\text{CH}_2\text{-OH}$, $-\text{CH}_2\text{-CH}_2\text{-CONH}_2$ and $-\text{CH}_2\text{-CH}(\text{CH}_3)\text{-OH}$ (especially wherein R^1 is a group of $-\text{CH}_2\text{-CONH}_2$).

5. The compound or a salt thereof according to claim 1, wherein R^2 is selected from the following groups: $-\text{CH}_2\text{-OH}$, $-\text{CH}_2\text{-CH}(\text{CH}_3)\text{-OH}$ and $-\text{CH}_3$, or one of the following groups:







-continued



(especially wherein R^2 is selected from the following groups: $-\text{CH}_3$, $-\text{CH}_2\text{-OH}$, $-\text{CH}_2\text{-SH}$, $-\text{CH}_2\text{-CONH}_2$, $-\text{CH}_2\text{-CH}_2\text{-CONH}_2$, $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-NH-C(=NH)-NH}_2$, and $-\text{CH}_2\text{-CH}(\text{CH}_3)\text{-OH}$).







6. The compound or a salt thereof according to claim 1, wherein R^3 is a hydrogen atom or selected from the following groups: $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-NH}_2$ and $-\text{CH}_2\text{-CH}_2\text{-NH-C(=NH)-NH}_2$ (especially wherein R^3 is the group: $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-NH}_2$).

7. The compound or a salt thereof according to claim 1, wherein R^4 is selected from the following groups: $-\text{CH}_2\text{-OH}$, $-\text{CH}_3$ and $-\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-NH-C(=NH)-NH}_2$; one of the following groups:





or

wherein R^4 is a group of formula $-\text{CH}_2-\text{SR}^{4A}$, wherein R^{4A} is selected from a hydrogen atom, an alkyl, an alkenyl, an alkynyl, a heteroalkyl, a cycloalkyl, a heterocycloalkyl, an alkylcycloalkyl, a heteroalkylcycloalkyl, an aryl, a heteroaryl, an aralkyl or a heteroaralkyl group, all of which groups may optionally be substituted (especially wherein R^4 is selected from the following groups: CH_3 , CH_2OH ,




-continued



preferably R^4 is selected from $-\text{CH}_2-\text{OH}$, $-\text{CH}_3$ and $-\text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}(=\text{NH})-\text{NH}_2$.

8. The compound or a salt thereof according to claim 1, wherein R^5 is a group of the following formula:




wherein R^9 is a hydrogen atom; R^{10} is a hydrogen atom; R^{11} is, at each occasion, independently selected from a halogen atom, a hydroxy group or a methoxy group (especially a halogen atom, preferably fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0).

9. The compound or a salt thereof according to claim 1, wherein each of R^6 , R^{6A} , R^7 , and R^{7A} is a hydrogen atom.

10. The compound or a salt thereof according to claim 1, wherein R^8 and/or R^{8A} is a halogen atom (especially a fluor or chlor atom, preferably a fluor atom).

11. The compound or a salt thereof according to claim 1, wherein the compound has structural formula (Ia):




wherein $R^1, R^2, R^3, R^4, R^5, R^6, R^7, R^8, R^{6A}, R^{7A}, R^{8A}, m$ and p are as defined in any one of the preceding claims.

12. Pharmaceutical composition comprising a compound or a salt thereof according to claim 1 and optionally i) one or more carrier substances and/or ii) one or more adjuvants and/or iii) one or more further active pharmaceutical ingredient(s).


13. (canceled)

14. A method of producing a compound according to claim 1, wherein the compound is a compound of formula (II):




wherein


R^2 and R^4 each independently represents H, CH_3 , CH_2OH or a group:




-continued



R^3 is a group:



R⁵ is a group of the following formula:



wherein R⁹ is a hydrogen or halogen atom (especially a hydrogen atom); R¹⁰ is a hydrogen atom or a methyl group (especially a hydrogen atom); R¹¹ is, at each occasion, independently selected from a halogen atom (especially fluor or chlor) and n is an integer of from 0 to 4 (especially 0 or 1; preferably 0)

and

wherein the method comprises the steps of:

- (a) providing a recombinant host capable of producing said compound of formula (II) (or (IIa), (IIa') or (IIa'')), wherein said recombinant host harbors at least one synthetic or recombinant nucleic acid encoding a bio-synthetic gene cluster (BGC) of the compound, which BGC has:
 - (i) a sequence identity to the full-length sequence of SEQ ID NO. 86 of at least 95%, 96%, 97%, 98%, 98.5%, 99%, or 99.5% to 100%; or
 - (ii) a sequence completely complementary to any nucleic acid sequence of (i);
- (b) cultivating said recombinant host for a time sufficient for said recombinant host to produce the compound of formula (II) (or (IIa), (IIa') or (IIa'')); and
- (c) isolating the compound from said recombinant host or from cultivation supernatant, thereby producing the compound of formula (II) (or (IIa), (IIa') or (IIa'')).

15. A recombinant host comprising a heterologous nucleic acid encoding a compound of formula (II) as defined in claim 14.

16. The recombinant host of claim 15, wherein the recombinant host is a microorganism, preferably a bacterium or yeast, such as a bacterial cell of the species *E. coli*, such as *E. coli* BL21(DE3) (especially DSM 33798 and/or DSM 33799); *Corynebacteria*, such as *C. glutamicum*; *Bacillus*; *Lactobacillus*, such as *Lactococcus lactis*; or *Streptomyces*, such as *S. albus* and *S. lividans*; or a yeast cell of the species *Saccharomyces cerevisiae*, *Saccharomyces pombe*, *Pichia pastoris* or *Yarrowia lipolytica*.

17. A vector comprising at least one nucleic acid as defined in claim 14, such as a vector capable of autonomous replication in bacteria or yeast, preferably an isolated and purified plasmid capable of autonomous replication in bacteria; preferably capable of autonomous replication in bacteria of the species *Lactobacillus*, such as *Lactococcus actis*, or *E. coli*, more preferably capable of autonomous replication in bacteria of the species *E. coli*, especially DSM 33802 and/or DSM 33803.

18.-19. (canceled)

20. A method of treating a subject suffering from or susceptible to a bacterial infection comprising: administering to the subject an effective amount of compound of claim 1.

21. The method of claim 20 wherein the bacterial infection is caused by Gram-negative bacteria.

22. The method of claim 20 wherein the subject is identified as suffering from or susceptible to a bacterial infection, and the compound is administered to the identified subject.

23. The method of claim 21 wherein the subject is identified as suffering from or susceptible to a bacterial infection, and the compound is administered to the identified subject

* * * * *