

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12409132
Kind Code	B2
Date of Patent	September 09, 2025
Inventor(s)	Plakogiannis; Fotios M. et al.

Transdermal delivery of dronabinol

Abstract

Provided is a transdermal drug delivery system comprising dronabinol. The dronabinol transdermal delivery system provides a drug plasma concentration at predetermined rate for a predetermined period of time, offering a simplified therapeutic regimen by decreasing dosing frequency for the treatment and/or prevention of nausea and/or vomiting associated with, for example, chemotherapy.

Inventors: Plakogiannis; Fotios M. (Whitestone, NY), Lather; Tamanna (Jersey City, NJ), Modi; Nisarg (Jersey City, NJ), Borovinskaya; Marina (East Brunswick, NJ)

Applicant: Pike Therapeutics USA, Inc. (Wilmington, DE)

Family ID: 72964852

Assignee: Pike Therapeutics USA, Inc. (Wilmington, DE)

Appl. No.: 17/221667

Filed: April 02, 2021

Prior Publication Data

Document Identifier	Publication Date
US 20210220325 A1	Jul. 22, 2021

Publication Classification

Int. Cl.: A61K31/352 (20060101); A61K9/00 (20060101); A61K9/70 (20060101); A61P1/08 (20060101); A61P3/04 (20060101)

U.S. Cl.:

CPC A61K9/0021 (20130101); A61K9/0014 (20130101); A61K9/703 (20130101); A61K9/7084 (20130101); A61K31/352 (20130101); A61P1/08 (20180101); A61P3/04

Field of Classification Search

CPC: A61K (9/0021); A61K (9/0014); A61K (9/703); A61K (9/7084); A61K (31/352); A61P (3/04); A61P (1/08)

USPC: 514/455

References Cited**U.S. PATENT DOCUMENTS**

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5804592	12/1997	Volicer	N/A	N/A
6113940	12/1999	Brooke et al.	N/A	N/A
6245347	12/2000	Zhang et al.	N/A	N/A
6328992	12/2000	Brooke et al.	N/A	N/A
6689379	12/2003	Bracht	N/A	N/A
6946150	12/2004	Whittle	N/A	N/A
7025992	12/2005	Whittle et al.	N/A	N/A
7544676	12/2008	Dolle et al.	N/A	N/A
7592328	12/2008	Jarho et al.	N/A	N/A
7622140	12/2008	Whittle et al.	N/A	N/A
7671052	12/2009	Dolle et al.	N/A	N/A
7709536	12/2009	Whittle	N/A	N/A
7807711	12/2009	Korthout et al.	N/A	N/A
7968594	12/2010	Guy et al.	N/A	N/A
8071641	12/2010	Weiss et al.	N/A	N/A
8211946	12/2011	Whittle	N/A	N/A
8246981	12/2011	Patel et al.	N/A	N/A
8293786	12/2011	Stinchcomb et al.	N/A	N/A
8435556	12/2012	Stinchcomb et al.	N/A	N/A
8449908	12/2012	Stinchcomb et al.	N/A	N/A
8481085	12/2012	Musty et al.	N/A	N/A
8603515	12/2012	Whittle	N/A	N/A
8642645	12/2013	Kelly	N/A	N/A
8771760	12/2013	Guy et al.	N/A	N/A
8992908	12/2014	Smith et al.	N/A	N/A
9029423	12/2014	Whittle	N/A	N/A
9034395	12/2014	Whittle et al.	N/A	N/A
9205063	12/2014	Guy et al.	N/A	N/A
9304134	12/2015	Smith	N/A	N/A
9375417	12/2015	Smith et al.	N/A	N/A
9533942	12/2016	Stinchcomb et al.	N/A	N/A
9603887	12/2016	Kelly	N/A	N/A
9763912	12/2016	Chen et al.	N/A	N/A
9833433	12/2016	Chen et al.	N/A	N/A
9918961	12/2017	Hearn et al.	N/A	N/A
9949937	12/2017	Guy et al.	N/A	N/A

9956183	12/2017	Guy et al.	N/A	N/A
9956184	12/2017	Guy et al.	N/A	N/A
9956185	12/2017	Guy et al.	N/A	N/A
9956186	12/2017	Guy et al.	N/A	N/A
9957321	12/2017	Smith et al.	N/A	N/A
9962340	12/2017	Weimann	N/A	N/A
9962341	12/2017	Stott et al.	N/A	N/A
10004684	12/2017	Whittle et al.	N/A	N/A
10028904	12/2017	Smith et al.	N/A	N/A
10092525	12/2017	Guy et al.	N/A	N/A
10098867	12/2017	Javid et al.	N/A	N/A
10098895	12/2017	Chang et al.	N/A	N/A
10111840	12/2017	Guy et al.	N/A	N/A
10118006	12/2017	Davidson et al.	N/A	N/A
10137095	12/2017	Guy et al.	N/A	N/A
10155018	12/2017	Jenn	N/A	N/A
10172809	12/2018	Aung-Din	N/A	N/A
10195159	12/2018	Whittle et al.	N/A	N/A
10213390	12/2018	Bonn-Miller et al.	N/A	N/A
10272125	12/2018	Weimann	N/A	N/A
10278996	12/2018	Avidov et al.	N/A	N/A
10307392	12/2018	Kariman	N/A	N/A
10314792	12/2018	Sebree et al.	N/A	N/A
10383816	12/2018	Aung-Din	N/A	N/A
10413521	12/2018	Hearn et al.	N/A	N/A
10420809	12/2018	Crowley	N/A	N/A
10471022	12/2018	Bonn-Miller et al.	N/A	N/A
10538373	12/2019	Whittle	N/A	N/A
10555927	12/2019	Jenn	N/A	N/A
10568848	12/2019	Sebree et al.	N/A	N/A
RE47885	12/2019	Strinchcomb et al.	N/A	N/A
10588869	12/2019	Weimann	N/A	N/A
10588871	12/2019	Fracassi et al.	N/A	N/A
10588974	12/2019	Leone-Bay et al.	N/A	N/A
10603288	12/2019	Guy et al.	N/A	N/A
10617733	12/2019	Kelly	N/A	N/A
10632064	12/2019	Aung-Din	N/A	N/A
10660872	12/2019	Sarne	N/A	N/A
10675240	12/2019	Smith et al.	N/A	N/A
10675264	12/2019	Green et al.	N/A	N/A
10695287	12/2019	Robbins et al.	N/A	N/A
10709671	12/2019	Guy et al.	N/A	N/A
10709673	12/2019	Guy	N/A	N/A
10709674	12/2019	Guy et al.	N/A	N/A
10709748	12/2019	Witowski et al.	N/A	N/A
10716766	12/2019	Aung-Din	N/A	N/A
10751299	12/2019	Ghalili	N/A	N/A
10758497	12/2019	Bonn-Miller et al.	N/A	N/A
10758514	12/2019	Liu et al.	N/A	N/A
10799545	12/2019	Weimann	N/A	N/A

10807777	12/2019	Whittle	N/A	N/A
10828266	12/2019	Aung-Din	N/A	N/A
10849860	12/2019	Guy et al.	N/A	N/A
10864189	12/2019	Borok	N/A	N/A
10869842	12/2019	Summers	N/A	N/A
10881606	12/2020	Schmitz et al.	N/A	N/A
10918608	12/2020	Guy et al.	N/A	N/A
10918686	12/2020	Siurkus	N/A	N/A
10945967	12/2020	Song	N/A	N/A
10966939	12/2020	Guy et al.	N/A	N/A
11026896	12/2020	Fitzsimmons et al.	N/A	N/A
11052055	12/2020	Kochinke	N/A	N/A
11065209	12/2020	Guy et al.	N/A	N/A
11065227	12/2020	Stott et al.	N/A	N/A
11096905	12/2020	Guy et al.	N/A	N/A
11116730	12/2020	Fracassi et al.	N/A	N/A
11147799	12/2020	Kopsky et al.	N/A	N/A
11154516	12/2020	Guy et al.	N/A	N/A
11154517	12/2020	Guy et al.	N/A	N/A
11160795	12/2020	Guy et al.	N/A	N/A
11207292	12/2020	Guy et al.	N/A	N/A
2002/0019421	12/2001	Biberman	N/A	N/A
2002/0111377	12/2001	Stinchcomb	N/A	N/A
2004/0033254	12/2003	Song et al.	N/A	N/A
2004/0138293	12/2003	Werner et al.	N/A	N/A
2005/0042271	12/2004	Xiong et al.	N/A	N/A
2005/0070596	12/2004	Baker et al.	N/A	N/A
2006/0039959	12/2005	Wessling	N/A	N/A
2006/0135599	12/2005	Symonds et al.	N/A	N/A
2007/0060638	12/2006	Olmstead et al.	N/A	N/A
2007/0072939	12/2006	Kupper	N/A	N/A
2007/0083186	12/2006	Carter et al.	N/A	N/A
2008/0112895	12/2007	Kottayil et al.	N/A	N/A
2009/0197941	12/2008	Guy et al.	N/A	N/A
2009/0298929	12/2008	Jarho et al.	N/A	N/A
2010/0035978	12/2009	Guy et al.	N/A	N/A
2010/0168108	12/2009	Dolle et al.	N/A	N/A
2010/0184848	12/2009	Wine et al.	N/A	N/A
2010/0273895	12/2009	Stinchcomb et al.	N/A	N/A
2010/0286098	12/2009	Robson et al.	N/A	N/A
2010/0317729	12/2009	Guy et al.	N/A	N/A
2011/0021617	12/2010	Korthout et al.	N/A	N/A
2011/0052694	12/2010	Stinchcomb et al.	N/A	N/A
2011/0097393	12/2010	Al-Ghananeem	N/A	N/A
2012/0034293	12/2011	Stinchcomb et al.	N/A	N/A
2013/0022687	12/2012	Fitzgerald, Jr. et al.	N/A	N/A
2013/0122077	12/2012	Al-Ghananeem	N/A	N/A
2013/0210786	12/2012	Howson	N/A	N/A
2013/0245110	12/2012	Guy et al.	N/A	N/A
2013/0253449	12/2012	Yoshitake et al.	N/A	N/A

2014/0039043	12/2013	Musty et al.	N/A	N/A
2014/0314757	12/2013	Sanchez et al.	N/A	N/A
2015/0265720	12/2014	Levine et al.	N/A	N/A
2015/0297556	12/2014	Smith	N/A	N/A
2015/0342902	12/2014	Vangara et al.	N/A	N/A
2015/0343071	12/2014	Vangara et al.	N/A	N/A
2016/0000843	12/2015	Lowe et al.	N/A	N/A
2016/0022627	12/2015	Smith	N/A	N/A
2016/0039591	12/2015	Kinzer	N/A	N/A
2016/0220593	12/2015	Anastassov et al.	N/A	N/A
2016/0228385	12/2015	Sievers et al.	N/A	N/A
2016/0256411	12/2015	Aung-Din	N/A	N/A
2016/0271252	12/2015	Vangara et al.	N/A	N/A
2016/0338974	12/2015	Aung-Din	N/A	N/A
2016/0361271	12/2015	Weimann	N/A	N/A
2017/0042791	12/2016	Ghalili et al.	N/A	N/A
2017/0071870	12/2016	Weimann	N/A	N/A
2017/0202895	12/2016	Hugh	N/A	N/A
2017/0273914	12/2016	Knudsen	N/A	N/A
2017/0306013	12/2016	Clark et al.	N/A	N/A
2018/0004994	12/2017	Bottazzi et al.	N/A	N/A
2018/0021247	12/2017	Ghalili et al.	N/A	N/A
2018/0042842	12/2017	Whittle et al.	N/A	N/A
2018/0042845	12/2017	Sinai et al.	N/A	N/A
2018/0071210	12/2017	Wilkhu et al.	N/A	N/A
2018/0078512	12/2017	Weimann	N/A	N/A
2018/0169035	12/2017	Eyal	N/A	N/A
2018/0284402	12/2017	Hoag	N/A	N/A
2018/0289665	12/2017	Turner et al.	N/A	N/A
2018/0311180	12/2017	Kochinke	N/A	N/A
2018/0311181	12/2017	Kochinke	N/A	N/A
2018/0311184	12/2017	Hoag	N/A	N/A
2018/0318529	12/2017	Davidson et al.	N/A	N/A
2018/0353463	12/2017	Winnicki	N/A	N/A
2018/0360757	12/2017	Doroudian et al.	N/A	N/A
2018/0369191	12/2017	Muscarella	N/A	N/A
2019/0023780	12/2018	Smith et al.	N/A	N/A
2019/0083388	12/2018	Gutterman et al.	N/A	N/A
2019/0105298	12/2018	Eyal	N/A	N/A
2019/0110981	12/2018	Weimann	N/A	N/A
2019/0117617	12/2018	Kariman	N/A	N/A
2019/0125779	12/2018	Ziburkus et al.	N/A	N/A
2019/0133994	12/2018	Smith et al.	N/A	N/A
2019/0134121	12/2018	Bermudez et al.	N/A	N/A
2019/0167583	12/2018	Shah	N/A	N/A
2019/0201372	12/2018	McKay	N/A	N/A
2019/0224118	12/2018	Navon et al.	N/A	N/A
2019/0224140	12/2018	Guy et al.	N/A	N/A
2019/0231826	12/2018	Avidov et al.	N/A	N/A
2019/0255014	12/2018	Gardner	N/A	N/A

2019/0298683	12/2018	Friedman	N/A	N/A
2019/0314297	12/2018	Gallily	N/A	N/A
2019/0321355	12/2018	Anavi-Goffer	N/A	N/A
2019/0321426	12/2018	Gallily	N/A	N/A
2019/0328884	12/2018	Jones, Jr. et al.	N/A	N/A
2019/0365667	12/2018	Wright et al.	N/A	N/A
2020/0038421	12/2019	Anastassov et al.	N/A	N/A
2020/0054887	12/2019	Levin	N/A	N/A
2020/0078332	12/2019	Leone-Bay et al.	N/A	N/A
2020/0085816	12/2019	Raz	N/A	N/A
2020/0093755	12/2019	Biro et al.	N/A	N/A
2020/0108027	12/2019	Whalley et al.	N/A	N/A
2020/0129741	12/2019	Kawwas et al.	N/A	N/A
2020/0138771	12/2019	Diez et al.	N/A	N/A
2020/0138773	12/2019	Jenn	N/A	N/A
2020/0163980	12/2019	Dellinger	N/A	N/A
2020/0170963	12/2019	Tich et al.	N/A	N/A
2020/0188324	12/2019	Sebree et al.	N/A	N/A
2020/0188348	12/2019	Sinai et al.	N/A	N/A
2020/0206184	12/2019	Robson et al.	N/A	N/A
2020/0214995	12/2019	Sebree et al.	N/A	N/A
2020/0215136	12/2019	Naheed	N/A	N/A
2020/0237683	12/2019	Whalley et al.	N/A	N/A
2020/0253919	12/2019	Raz et al.	N/A	N/A
2020/0261376	12/2019	Yu et al.	N/A	N/A
2020/0261404	12/2019	Raz et al.	N/A	N/A
2020/0276132	12/2019	Weimann	N/A	N/A
2020/0297656	12/2019	Guy et al.	N/A	N/A
2020/0330379	12/2019	Singh et al.	N/A	N/A
2020/0338041	12/2019	Smith et al.	N/A	N/A
2020/0338151	12/2019	Witowski et al.	N/A	N/A
2020/0345653	12/2019	Hansen et al.	N/A	N/A
2020/0345655	12/2019	Heinzerling et al.	N/A	N/A
2020/0345657	12/2019	Lurya et al.	N/A	N/A
2020/0345685	12/2019	Otiko	N/A	N/A
2020/0352849	12/2019	Rotunda	N/A	N/A
2020/0352901	12/2019	Raber et al.	N/A	N/A
2020/0360299	12/2019	Bonn-Miller et al.	N/A	N/A
2020/0384048	12/2019	Kariman	N/A	N/A
2020/0384049	12/2019	Kariman	N/A	N/A
2021/0015740	12/2020	Greenspan et al.	N/A	N/A
2021/0023044	12/2020	Spiros	N/A	N/A
2021/0030777	12/2020	Maida	N/A	N/A
2021/0052512	12/2020	Guy et al.	N/A	N/A
2021/0052545	12/2020	Jones, Jr. et al.	N/A	N/A
2021/0059976	12/2020	Wilkhu et al.	N/A	N/A
2021/0069333	12/2020	Diez et al.	N/A	N/A
2021/0077421	12/2020	Sebree et al.	N/A	N/A
2021/0100737	12/2020	Plakogiannis	N/A	A61P 1/08
2021/0106540	12/2020	Plakogiannis et al.	N/A	N/A

2021/0137833	12/2020	Wang et al.	N/A	N/A
2021/0145764	12/2020	Lephart	N/A	N/A
2021/0177937	12/2020	Triebel et al.	N/A	N/A
2021/0186860	12/2020	Weimann	N/A	N/A
2021/0196669	12/2020	Schleider et al.	N/A	N/A
2021/0220325	12/2020	Plakogiannis et al.	N/A	N/A
2021/0236417	12/2020	Plakogiannis	N/A	A61K 9/7084
2021/0244680	12/2020	Kassab et al.	N/A	N/A
2021/0244683	12/2020	Chaiyasate	N/A	N/A
2021/0244684	12/2020	Ghalili et al.	N/A	N/A
2021/0290565	12/2020	Guy et al.	N/A	N/A
2021/0308070	12/2020	Summers	N/A	N/A
2021/0308072	12/2020	Wright et al.	N/A	N/A
2021/0379011	12/2020	Guynn	N/A	N/A
2021/0386684	12/2020	Weimann	N/A	N/A
2021/0386685	12/2020	Weimann	N/A	N/A
2021/0401766	12/2020	Rhodes et al.	N/A	N/A
2021/0401770	12/2020	Fracassi et al.	N/A	N/A
2021/0401771	12/2020	Guy et al.	N/A	N/A
2022/0062211	12/2021	Stott et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2859930	12/2015	CA	N/A
2978605	12/2015	CA	N/A
2609912	12/2012	EP	N/A
2011505382	12/2010	JP	N/A
2001003668	12/2000	WO	N/A
2001064149	12/2000	WO	N/A
2002064109	12/2001	WO	N/A
2002069993	12/2001	WO	N/A
2002089945	12/2001	WO	N/A
2003105800	12/2002	WO	N/A
2004016246	12/2003	WO	N/A
2006041841	12/2005	WO	N/A
2006044645	12/2005	WO	N/A
2006124698	12/2005	WO	N/A
2006044645	12/2005	WO	N/A
2007001891	12/2006	WO	N/A
2008024408	12/2007	WO	N/A
2008024490	12/2007	WO	N/A
2008039179	12/2007	WO	N/A
2008063625	12/2007	WO	N/A
2008129258	12/2007	WO	N/A
2009020666	12/2008	WO	N/A
2009073633	12/2008	WO	N/A
2010126501	12/2009	WO	N/A
2011026144	12/2010	WO	N/A
2013108254	12/2012	WO	N/A
2015025312	12/2014	WO	N/A

2015161165	12/2014	WO	N/A
2016090287	12/2015	WO	N/A
2016141056	12/2015	WO	N/A
2018048789	12/2017	WO	N/A
2018071452	12/2017	WO	N/A
2018071581	12/2017	WO	N/A
2018135943	12/2017	WO	N/A
2019058261	12/2018	WO	N/A
2019130215	12/2018	WO	N/A
2019210287	12/2018	WO	N/A
2020016581	12/2019	WO	N/A
2020016582	12/2019	WO	N/A
2020123625	12/2019	WO	N/A
2020128767	12/2019	WO	N/A
2020136593	12/2019	WO	N/A
2020142692	12/2019	WO	N/A
2020152438	12/2019	WO	N/A
2020157569	12/2019	WO	N/A
2020183350	12/2019	WO	N/A
2020198252	12/2019	WO	N/A
2020198883	12/2019	WO	N/A
2020257538	12/2019	WO	N/A
2021003467	12/2020	WO	N/A
2021023351	12/2020	WO	N/A
2021055499	12/2020	WO	N/A
2021070120	12/2020	WO	N/A
2021177940	12/2020	WO	N/A
2021236782	12/2020	WO	N/A
2022118290	12/2021	WO	N/A

OTHER PUBLICATIONS

Alper, Kenneth et al., “The ibogaine medical subculture”, Journal Of Ethnopharmacology, vol. 115(1), p. 9-24, 2008. cited by applicant

ASI., “Adhesives Transdermal Delivery Systems”, 2005, retrieved from AdhesivesMag.com, Nov. 1, 2005, retrieved from Web Archive Oct. 5, 2016, <https://web.archive.org/web/20161005045648/https://www.adhesivemag.com/articles/86012-adhesives-in-transdermal-drug-delivery-systems>. cited by applicant

Beug, Michael et al., “Psilocybin and Psilocin Levels in Twenty Species From Seven Genera of Wild Mushrooms in the Pacific Northwest, U.S.A.”, Journal of Ethnopharmacology vol. 5(3), p. 271-285, 1982. cited by applicant

Cameron, Lindsay et al., “Psychedelic Microdosing: Prevalence and Subjective Effects”, Journal of Psychoactive Drugs vol. 52(2), p. 113-122, 2020. cited by applicant

Center for Drug Evaluation and Research, Application No. 210365Orig1s000; Non Clinical Reviews; Submission date: Jun. 23, 2017. cited by applicant

Glick, Stanley et al., “18-Methoxycoronaridine (18-MC) and ibogaine: Comparison of Antiaddictive Efficacy Toxicity, and Mechanisms of Action”, Annals of the New York Academy of Sciences, vol. 914(1) p. 369-386, 2006. cited by applicant

International Preliminary Report on Patentability for PCT/IB2021/057474 dated Mar. 2, 2023. cited by applicant

International Preliminary Report on Patentability for PCT/IB2021/057483 dated Mar. 2, 2023. cited

by applicant

International Search Report and Written Opinion for International Application No. PCT/US2022/053271 mailed Jul. 12, 2022. cited by applicant

International Search Report for PCT/IB2022/053271 mailed Jul. 12, 2022. cited by applicant

International Search Report mailed on Dec. 9, 2021 for PCT/IB2021/057474. cited by applicant

Jensen, Bjorn et al., “Medical Marijuana and Chronic Pain: a Review of Basic Science and Clinical Evidence”, Curr Pain Headache Rep. 2015. cited by applicant

Leehy, M. et al., “Safety and Tolerability of Cannabidiol in Parkinson's Disease: An Open Label, Dose-Escalation Study, Cannabis and Cannabinoid Research”, 2020, vol. 5, No. 4. cited by applicant

McPartland, J. et al., “Care and Feeding of the Endocannabinoid System: A Systematic Review of Potential Clinical Interventions that Upregulate the Endocannabinoid System”. PLOS One, Mar. 12, 2014, vol. 9, Issue 3. cited by applicant

Ohlsson, Per-Ingvar, “Lacetoperoxidase, a dithionite ion dismutase” Eur. J. Biochem, 1984, vol. 142, pp. 233-238. cited by applicant

Polito, Vince, “A systematic study of microdosing psychedelics”, PLOS, vol. 14(2), p. 1-26, 2019. cited by applicant

Shapiro, L., “Children, but Linked to Side Effects Analysis: Clinicians should consider adverse effects before treatment”, Dravet Syndrome News, Aug. 30, 2022. cited by applicant

Sinha V.R. et al., “Permeation Enhancers for Transdermal Drug Delivery”, Drug Development and Industrial Pharmacy, vol. 26, p. 1131-1140, 2000. cited by applicant

Taylor, L. “A Phase I, Randomized, Double-Blind, Placebo-Controlled, Single Ascending Dose, Multiple Dose and Food Effect Trial of the Safety, Tolerability and Pharmacokinetics of Highly Purified Cannabidiol in Healthy Subjects”, CNS Drugs, 2018, vol. 32, pp. 1053-1067. cited by applicant

Valenta, Claudia et al., “The use of polymers for dermal and transdermal delivery” European Journal of Pharmaceutics and Biopharmaceutics, pp. 279-289, vol. 58 (2), 2004. cited by applicant

Touitou et al., Transdermal delivery of tetrahydrocannabinol. Int J Pharmaceutics. Apr. 1988;43(1-2):9-15.4. cited by applicant

Communication pursuant to Article 94(3) EPC for European Application No. 20793917.4 dated Jun. 25, 2024. cited by applicant

English Translation of Japanese Patent Application 2011505382 filed Dec. 1, 2008. cited by applicant

International Preliminary Report on Patentability for International Application No. PCT/US2020/054070 dated Apr. 5, 2022. cited by applicant

International Search Report and Written Opinion for International Application No. PCT/US2020/054070 dated Jan. 1, 2021. cited by applicant

Office Action for Japanese Patent Application No. 2022-520408 issued on Sep. 24, 2024. cited by applicant

Second Office Action for Chinese Patent Application No. 20280077942.3 issued on Sep. 19, 2024. cited by applicant

Beal, J. E. et al., Dronabinol as a Treatment for Anorexia Associated with Weight Loss in Patients with AIDS' Journal of Pain and Symptom Management, 1995, vol. 10, pp. 89-97. cited by applicant

Office Action for Australian Patent Application No. 2020358869 dated Jun. 19, 2025. cited by applicant

Office Action for Japanese Patent Application No. 2022-520408 dated May 16, 2025 (with English translation). cited by applicant

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS

(1) This application is a continuation-in-part of, and claims the benefit of and priority to, International Application No. PCT/US2020/054070, filed on Oct. 2, 2020, which claims the benefit of U.S. Provisional Application No. 62/910,255, filed Oct. 3, 2019, both of which are incorporated by reference herein in their entirety for all purposes.

TECHNICAL FIELD

(2) The present disclosure relates to treatment of nausea and vomiting, such as nausea and vomiting commonly experienced by patients undergoing chemotherapy.

BACKGROUND

(3) Nausea and vomiting are the common side effects of cancer chemotherapy. Drugs such as dronabinol and 5-HT₃ receptor antagonists are recommended for the treatment of nausea and vomiting associated with cancer chemotherapy. At times, to prevent nausea and vomiting one or more drugs can be used (See Herrsted J., et al., 2016 Updated MASCC/ESMO Consensus Recommendations: Prevention of Nausea and Vomiting Following High Emetic Risk Chemotherapy. *Support Care Cancer*. 2017 January; 25(1): 277-288. Epub 2016 Jul. 22).

(4) Dronabinol is a synthetic form of delta-9-tetrahydrocannabinol (delta-9-THC) in sesame oil. Dronabinol is approved for use in treating emesis, and is available as an oral capsule (MARINOL) and as an oral solution (SYNDROS). As stated in the package insert or drug label for the oral capsule of dronabinol the drug "is indicated for the treatment of nausea and vomiting associated with cancer chemotherapy in patients who have failed to respond adequately to conventional antiemetic treatments. It is also indicated for the treatment of anorexia associated with weight loss in patients with AIDS" (See Label for NDA Marinol, Labeling, Label action date Jun. 21, 2006 accessed on Jul. 13, 2017, hereafter "Label").

(5) Many shortcomings are associated with the dronabinol oral capsule. The first challenge is the dosage regimen. For an antiemetic effect the following dosage regimen is recommended: 1 to 3 hours before the chemotherapy a first dose of 5 mg/m² is administered orally. After chemotherapy drug is administered every 2 to 4 hours in total 4 to 6 doses per day. Depending on the clinical response, if the dose is not sufficiently effective then the dose is increased by increments of 2.5 mg/m² up to about 15 mg/m². This dosage regimen is inconvenient for patients who are already experiencing nausea and vomiting. A second challenge is that the pharmacologic response is dose related and there is interpatient variability. A third challenge is that at maximum dose the likelihood of disturbing psychiatric symptoms increases. A fourth challenge is that after oral administration dronabinol undergoes first pass hepatic metabolism and has high lipid solubility; therefore, of the administered dose 10%-20% reaches systemic circulation. Another challenge is that at room temperature the active ingredient dronabinol is unstable in the capsules, and so the capsules are packaged in a closed container and recommended storage is in refrigerator or between 8° C.-15° C. (See FDA Label

https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/018651s029lbl.pdf). The dronabinol capsule is stable for only 3 months at room temperature. This is due to synthetic delta-9-THC exhibiting rapid oxidation and acid and basic degradation. Moreover, synthetic delta-9-THC also degrades more rapidly in light and at higher temperatures the degradation rate increases. Accordingly, if dronabinol capsules are not stored refrigerated and in the original container, the concentration of delta-9-THC in the capsules will decrease, potentially below the therapeutically required concentration.

(6) Dronabinol is a synthetic form of delta-9-tetrahydrocannabinol (delta-9-THC) in sesame oil. Previous studies have been performed with naturally derived delta-9-THC from marijuana plant *Cannabis sativa* L. However, it is difficult to extract pure delta-9-THC from a plant source, such as *sativa* L. This is caused by an adulteration of the extract by small amounts of another active cannabidiols (such as cannabitol, delta-8-THC, cannabidiol and cannabichromene) present in final product. Accordingly, the amount of THC in final extract is dependent on the extraction process which is important due to the psychoactive properties of THC, among other considerations. Alternatively, the synthetic form of delta-9-THC is developed under a more controlled procedure as opposed to the extraction process for its naturally derived counterpart. For example, synthetically produced delta-9-THC consists essentially of uncontaminated delta-9-THC that is not adulterated by the presence of other active cannabidiols. Therefore, the synthetic version of delta-9-THC is able to provide improved transdermal permeability as compared to previously published work with naturally derived delta-9-THC which demonstrates a maximum flux of 0.01 $\mu\text{g}/\text{cm}^2/\text{hr}$ through rat skin for naturally extracted delta-9-THC (U.S. Pat. No. 6,503,532).

(7) U.S. Pat. No. 6,328,992 discloses the preparation of transdermal delivery of cannabinoids. However, the disclosure describes different cannabinoids in combination and does not describe the use of pure synthetic delta-9-THC alone. Furthermore, the '992 patent inventors describe the use of permeability enhancers which are not pharmaceutical grade and not included in U.S. Food and Drug Administration Inactive Ingredient Listing (FDA IIG). Therefore the disclosure of the '992 does not provide a composition with any pharmaceutical utility. Furthermore, the '992 patent describes the use of rat skin with 50% ethanol in receiving media for the transdermal permeability testing. However, ethanol destroys the dermis skin structure and can increase the in-vitro flux value thus providing results that are not germane to transdermal patches for pharmaceutical applications. For example, ethanol treated rat skin is almost ten time more permeable than human cadaver skin.

(8) U.S. Pat. No. 8,449,908 provides delivery of the cumulative amount of 10000 ng THC in 96 hrs through the human cadaver skin. This amount represents the flux of 60 $\text{ng}/\text{cm}^2/\text{hr}$. The patch area can be calculated using following equation:

$$\begin{aligned}\text{In-Vitro Flux}(\text{ng}/\text{cm}^2/\text{hr}) &= (\text{Css}(\text{ng}/\text{mL}) * \text{CL}(\text{L}/\text{kg} * \text{hr}) * \text{BW}(\text{Kg})) / \text{Patch Area}(\text{cm}^2) \\ \text{Patch Area}(\text{cm}^2) &= (\text{Css}(\text{ng}/\text{mL}) * \text{CL}(\text{L}/\text{kg} * \text{hr}) * \text{BW}(\text{kg})) / \text{In-Vitro Flux}(\text{ng}/\text{cm}^2/\text{hr}) = \\ &= (1.38 * 0.2 * 70 * 1000) / 60 = 322 \text{ cm}^2\end{aligned}$$

(9) In order to deliver a therapeutic dosage of 5 mg/day of delta-9-THC, a transdermal formulation would need to cover at least 325 cm^2 surface area of the patient's skin. This is an impractical patch size for any transdermal drug delivery system (TDDS).

(10) Therefore, there is a need for an improved drug delivery system of dronabinol which can overcome above stated drawbacks associated with oral administration and naturally derived delta-9-THC. As provided herein, transdermal delivery of dronabinol comprising synthetic delta-9-THC can address the challenges associated with oral drug delivery.

BRIEF SUMMARY

(11) The following aspects and embodiments thereof described and illustrated below are meant to be exemplary and illustrative, not limiting in scope.

(12) In one aspect, a pharmaceutical composition comprising dronabinol in a dosage form for transdermal delivery is provided.

(13) In an embodiment, the pharmaceutical composition comprises no additional antiemetic drug.

(14) In an embodiment, dronabinol is in a form selected from the group consisting of co-crystals, amorphous, coated, crystalline, a salt, an isomer, a solid solution, a prodrug, an analog, a derivative, a metabolite, a solution, synthetic, an ethanol solution, and a naturally derived delta-9-tetrahydrocannabinol.

(15) In an embodiment, dronabinol is in the composition at between about 0.01%-95% w/w or between about 0.01%-95% w/v.

(16) In an embodiment, dronabinol is selected from a group consisting of amorphous dronabinol,

crystalline dronabinol, co-crystals of dronabinol, coated dronabinol, and ethanolic solution of dronabinol in the range of 0.01%-95% w/w or w/v.

(17) In an embodiment, dronabinol is in a salt form.

(18) In an embodiment, the composition is formulated as transdermal liquid formulation, transdermal semisolid formulation and/or transdermal polymer matrix formulation.

(19) In an embodiment, a carrier or an ingredient in effective amount either alone or in combinations thereof is included in the composition. In an embodiment, the carrier or ingredient is selected from the group consisting of solvents, gelling agents, polymers, penetration enhancers, emollients, skin irritation reducing agents, buffering agents, pH stabilizers, solubilizers, suspending agents, dispersing agents, stabilizers, plasticizers, surfactants, antioxidants, and oxidants.

(20) In an embodiment, the carrier or ingredient is in the composition in a range of between about 0.01%-95% w/w or w/v.

(21) In an embodiment, the pharmaceutical composition is formulated as a transdermal patch.

(22) In an embodiment, the transdermal patch is selected from the group consisting of a reservoir patch, a micro-reservoir patch, a matrix patch, a pressure sensitive adhesive patch, and an extended release transdermal film.

(23) In an embodiment, the pharmaceutical composition is formulated as microneedles.

(24) In an embodiment, the microneedles are formulated as a transdermal patch.

(25) In an embodiment, a method for the treatment and/or prevention and/or control of nausea and/or vomiting associated with cancer chemotherapy comprises selecting a patient in need of treatment and/or prevention and/or control of nausea and/or vomiting associated with cancer chemotherapy; and topically applying or instructing to topically apply the pharmaceutical composition as described herein, wherein the topically applying is performed at least once in a day.

(26) In an embodiment, the composition is a liquid formulation and/or a semisolid formulation, wherein the topically applying is done two to six times in a day, once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week.

(27) In an embodiment, the composition is topically applied once in two days, once in three days, once in four days, once in five days, once in six days, once in a week, or once in ten days.

(28) In an embodiment, topically applying provides a constant rate of delivery of the active components of the transdermal patch over a time period.

(29) In an embodiment, topically applying provides a steady absorption rate of dronabinol over a time period.

(30) In an embodiment, topically applying achieves a constant blood serum level of dronabinol over a time period.

(31) In an embodiment, topically applying achieves a reduced variability in blood serum level of dronabinol over a time period relative to oral administration of a dronabinol over the time period.

(32) In an embodiment, topically applying achieves a plasma concentration of dronabinol in a therapeutic range over a period of time.

(33) In an embodiment, as method for the treatment and/or prevention and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS comprises selecting a patient in need of the treatment and/or prevention and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS; topically applying or instructing to topically apply the pharmaceutical composition described herein; wherein applying achieves topical delivery of dronabinol for the treatment and/or prevention and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS.

(34) In an embodiment, the pharmaceutical composition is topically applied once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week, or once in ten days.

(35) In an embodiment, the pharmaceutical composition is topically applied two to six times in a day, once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week.

(36) In an embodiment, the pharmaceutical composition is a liquid formulation or a semisolid formulation.

(37) In an embodiment, the pharmaceutical composition is a transdermal delivery system.

(38) In addition to the exemplary aspects and embodiments described above, further aspects and embodiments will become apparent by reference to the drawings and by study of the following descriptions.

(39) Additional embodiments of the present methods and compositions, and the like, will be apparent from the following description, drawings, examples, and claims. As can be appreciated from the foregoing and following description, each and every feature described herein, and each and every combination of two or more of such features, is included within the scope of the present disclosure provided that the features included in such a combination are not mutually inconsistent. In addition, any feature or combination of features may be specifically excluded from any embodiment of the present disclosure. Additional aspects and advantages of the present disclosure are set forth in the following description and claims, particularly when considered in conjunction with the accompanying examples and drawings.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 is a graph showing the plasma concentration of a drug as a function of time, in hours, when delivered via the oral route (open circles) or the transdermal route (closed circles).

DETAILED DESCRIPTION

(2) In transdermal drug delivery, a transdermal patch or transdermal composition is applied topically to the skin surface. Throughout the duration of topical application of a transdermal patch or transdermal composition drug is continuously released and delivered through the intact skin (via transcellular, intercellular and trans-appendageal routes) to achieve systemic effect. Therefore, once applied the transdermal composition or transdermal patch can deliver drug into systemic circulation throughout the day or even for more than one day depending on the duration of its application, which can be up to a week or longer.

(3) Transdermal delivery can reduce the dosing frequency of dronabinol which is currently administered orally 4-6 times a day. Through transdermal delivery, transdermal compositions or transdermal formulations or transdermal patch of dronabinol can be applied topically to skin thereby delivering the drug throughout the duration of topical application. Depending on the requirement, duration of topical application and can be once in day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week. Therefore, transdermal delivery can overcome the multiple dose regimen of oral delivery by reducing the dosing frequency.

(4) Moreover, in transdermal drug delivery drug is delivered slowly and continuously throughout the duration of topical application hence there are no peaks and troughs in drug plasma concentration which are associated with multiple dose administration in a day. Therefore, by transdermal delivery of dronabinol patients can have the therapeutic effect of the drug for extended period of time without drastic changes in drug plasma concentration.

(5) In transdermal delivery drug is delivered into systemic circulation through the skin, it escapes the first pass hepatic metabolism therefore to achieve the desired therapeutic activity less drug is required, resulting into less adverse effects or side effects. Dronabinol has high lipid solubility and after oral administration undergoes hepatic first pass metabolism therefore of the administered dose

10%-20% reaches systemic circulation hence, as compared to oral dose with transdermal delivery small dose of dronabinol can give the desired therapeutic effects and can also overcome the disturbing psychiatric symptoms associated with high dose.

(6) Also, the transdermal delta-9-THC delivery described herein provides almost 10 times more in-vitro permeability through human cadaver skin using FDA IIG database for approved drug products listed ingredients, as compared to previous systems.

(7) For example, according to FDA labels for Marinol (dronabinol capsule) were given to healthy volunteers (n=34; 20-25 years) twice a day under fasted condition, the following pharmacokinetic (PK) parameters were found (see Label).

(8) TABLE-US-00001 TABLE 1 Summary of Marinol® PK parameters BID Mean (S.D.) PK Parameters Value Dose C.sub.max Median T.sub.max AUC (0-12) (mg) ng/ml (range) hr ng * hr/mL 2.5 1.32 (0.62) 1.00 (0.50-4.00) 2.88 (1.57) 5 2.96 (1.81) 2.50 (0.50-4.00) 6.16 (1.85) 10 7.88 (4.54) 1.50 (0.50-3.50) 15.2 (5.52)

(9) Additionally, according to the PK parameters, the maximum plasma concentration for oral delivery of 5 mg/day by dronabinol capsule is 1.32 ng/ml and for oral delivery of 20 mg/day the maximum plasma concentration is 7.88 ng/mL. The oral delivery PK values demonstrate a peak and valley in plasma concentration over time. In contrast to oral delivery, transdermal drug delivery delivers the drug molecule at predetermined rate and maintains constant average plasma concentration over time (FIG. 1).

(10) The average plasma concentration is calculated in the following manner:

Average Plasma Concentration for 5 mg/day= $AUC_{sub.(0-t)}/t=2.88/12=0.24$ ng/ml
20 mg/day= $15.2/12=1.27$ ng/ml

(11) Therefore, transdermal drug delivery systems (TDDS) are calculated to deliver delta-9-THC, at average plasma concentrations of 0.24-1.27 ng/mL.

(12) For the prevention of nausea and vomiting, the oral route is often not the most convenient. Patients are already experiencing nausea and vomiting and if they vomit shortly after the drug administration uncertainty remains whether the dose was absorbed or is vomited. On the other hand, transdermally dronabinol will be delivered through the skin which completely eliminates these kinds of uncertainties.

(13) Moreover, there are various additional side effects, such as Central Nervous System (CNS) side effects, related to oral dronabinol capsule administration. The side effects are dose related and therefore patients receiving the oral dosage require close monitoring so that the dosage can be adjusted and reduced as needed.

(14) Alternatively, the TDDS systems provided herein provide constant drug delivery at a predetermined and defined input rate. Therefore, the currently provided TDDS systems provide a constant average plasma concentration due to constant input rates and do not exhibit PK peaks and valleys in plasma concentration associated with orally delivered delta-9-THC. Accordingly, the currently provided TDDS systems are also not as likely to cause CNS side effects as compared to the orally delivered counterparts. For example, adverse effects of delta-9-THC are not induced with a dosage of 7 mg/m.sup.2 of delta-9-THC as shown in the following examples. Dosages greater than 7 mg/m.sup.2 of delta-9-THC had no CINV effect and had more side effects.

(15) Furthermore, oral dosages must be divided by being provided 4 to 6 times per day instead of taking once a day in order to avoid adverse effects. Additionally, oral dosage undergoes first pass metabolism and only 10-20% of the oral dose becomes available in plasma, resulting an inefficient delivery method and a large loss of the active pharmaceutical. Moreover, in order to achieve therapeutic effect from any drug molecule, the drug should have continuous delivery to achieve steady state plasma concentration or a plasma concentration in therapeutic window (between minimum and maximum therapeutic effective concentration). Therefore, the TDDS systems for delta-9-THC provided herein address long felt and unmet needs in the field of cannabidiol based pharmaceuticals.

(16) Also, transdermal delivery is easy, noninvasive and convenient. Administration of transdermal patch or transdermal composition does not require medical supervision as patients can topically apply the transdermal patch or transdermal composition themselves.

(17) With respect to dronabinol it is expected that interpatient variability in pharmacologic response will be less with the transdermal delivery as drug plasma concentration can be controlled by controlling the rate of drug delivery from transdermal composition or transdermal patch. Alpha half-life of dronabinol is about 4 hours, by oral administration it gets rapidly metabolized in the body. In contrast, with transdermal delivery small amount of dronabinol can be delivered for longer duration than oral administration. Transdermal formulations of dronabinol also provide more abuse deterrence than immediate release dosage forms.

(18) Moreover, in case of any adverse effect, side effect or emergency transdermal delivery gives the liberty to terminate the therapy anytime by taking off the transdermal patch or transdermal composition from skin.

(19) As per above stated reasons for the prevention of nausea and vomiting associated with cancer chemotherapy transdermal delivery can provide patient friendly, simplified and convenient therapeutic regimen over traditional delivery systems. Transdermal delivery can reduce the dosing frequency of dronabinol. Depending on the necessity, dosing frequency can be once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week.

(20) For the prevention of nausea and vomiting associated with cancer chemotherapy patients are prescribed with a variety of different drugs, some of which are administered multiple times a day. By formulating the antiemetic drug(s) in a transdermal patch or transdermal composition provides a much-simplified dosage regimen to patients who are already weak and tired due to chemotherapy. Depending on the necessity, dosing frequency of the transdermal patch or transdermal composition containing the drug can be once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week. It would be a great addition to the patient compliance.

(21) Dronabinol is an unstable drug. Stability of dronabinol can be improved by preparation of dronabinol's co crystals, coating the dronabinol, dronabinol's crystals, amorphous form, and/or inert substances, and the like.

(22) Transdermal drug delivery composition, that in one embodiment, are in the form of a transdermal patch or delivery system comprising dronabinol are contemplated . . . Transdermal delivery can provide drug plasma concentration at predetermined rate for a predetermined period of time with a simplified therapeutic regimen by decreasing dosing frequency. Preferably, without any limitation dronabinol is selected from group such as dronabinol's co crystals, dronabinol's amorphous form, coated dronabinol, dronabinol's crystalline form.

(23) As used herein, "dronabinol" refers to all forms of dronabinol either alone or in combinations thereof, for example, in following forms but not limited to such as free base or salts or isomers or amorphous or crystalline or co-crystals or solid solution or prodrugs or analogs or derivatives or metabolites or coated form or natural extract of delta-9-tetrahydrocannabinol or synthetic delta-9-tetrahydrocannabinol, solution of dronabinol. For example, dronabinol's free form or its salts or its isomers or its amorphous form or its crystalline form or its co crystals or its solid solution or its prodrugs or its analogs or its derivatives or its metabolites or its coated form or natural extract of delta-9-tetrahydrocannabinol or synthetic delta-9-tetrahydrocannabinol, ethanolic solution of dronabinol. Dronabinol may be in the form of a pharmaceutically acceptable salt, such as an acid addition salt or a base salt, or a solvate thereof, including a hydrate thereof. Suitable acid addition salts are formed from acids which form non-toxic salts. Chemical Name: (6aR-trans)-6a,7,8,10a-tetrahydro-6,6,9-trimethyl-3-pentyl-6H-dibenzo[b,d]pyran-1-ol Empirical Formula:

C.sub.21H.sub.30O.sub.2 Molecular weight: 314.47 Structure: Formula 1

(24) ##STR00001##

(25) Dronabinol is synthetic delta-9-tetrahydrocannabinol. Delta-9-tetrahydrocannabinol is a natural component, present in *Cannabis sativa* L

(26) In embodiments of the disclosure, without any limitation, a preferable form of dronabinol is selected from group such as dronabinol co-crystals, dronabinol amorphous form, coated dronabinol, dronabinol crystalline form. Dronabinol's co-crystals, coated dronabinol, dronabinol's crystalline form may be prepared.

(27) Amorphous forms of the drug do not have a definite structure. An amorphous form of the drug has higher solubility as compared to a crystalline form. Different techniques and methods are used to make amorphous form of drugs.

(28) As stated in regulatory classification of pharmaceutical co-crystals guidance for industry "Co-crystals are crystalline materials composed of two or more different molecules, typically drug and co-crystal formers ("coformers"), in the same crystal lattice" (See U.S. Department of Health and Human Services Food and Drug Administration Center for Drug Evaluation and Research (CDER), Regulatory Classification of Pharmaceutical Co-Crystals Guidance for Industry Draft Guidance, Pharmaceutical Quality/CMC Revision 1, August 2016

<https://www.fda.gov/downloads/Drugs/Guidances/UCM516813.pdf>, accessed on Jul. 12, 2017).

Different methods are available for the preparation of co-crystals. Each drug has distinct chemical structure and physicochemical properties therefore, it is difficult to predict the success rate of co-crystallization reaction. Studies under various experimental conditions are done to determine an approach to form co-crystals of drug (See Nate Schultheiss, Ann Newman. Pharmaceutical Cocystals and their Physicochemical Properties. Cryst Growth Des. 2009 Jun. 3; 9 (6): 2950-2967).

(29) Coating of the drug can be done with polymer or other excipients. Different techniques are used for the coating of drug. Stability of the drug can also be increased by encapsulation.

(30) As used herein, the term "pharmaceutically acceptable salts" includes acid addition salts or addition salts of free bases. The term "pharmaceutically acceptable salts" of the dronabinol within its scope all the possible isomers and their mixtures, and any pharmaceutically acceptable metabolite, bio-precursor and/or pro-drug, such as, for example, a compound which has a structural formula different from the one of the compounds of the disclosure, and yet is directly or indirectly converted in vivo into a compound of the disclosure, upon administration to a subject, such as a mammal, particularly a human being.

(31) In one embodiment, dronabinol is incorporated into the transdermal system in the form of a pharmaceutically acceptable salt form, either as a single salt, as combinations of salts, or as a combination of the base form and one or more salt forms. Examples of the various forms of dronabinol include but not limited to such as free base, salts, racemic form, isomers, amorphous, crystalline, co-crystals, solid solution, prodrugs, analogs, derivatives, metabolites, solutions, hydrates. Therapeutic agents may be in the form of a pharmaceutically acceptable salt, such as an acid addition salt or a base salt, or a solvate thereof, including a hydrate thereof. Suitable acid addition salts are formed from acids which form non-toxic salts and without any limitation examples are acetate, hydrochloride, hydrobromide, hydroiodide, sulphate, bisulphate, nitrate, phosphate, hydrogen phosphate, sodium phosphate, maleate, fumarate, lactate, tartrate, citrate, gluconate, succinate, saccharate, benzoate, methanesulphonate, ethanesulphonate, benzenesulphonate, p-toluenesulphonate and pamoate salts. Suitable base salts are formed from bases which form non-toxic salts and without any limitation examples are the sodium, potassium, aluminum, calcium, magnesium, zinc and diethanolamine salts.

(32) As used herein, the terms "subject" and "patient" are used interchangeably. As used herein, the term "patient" refers to an animal, preferably a mammal such as a non-primate (e.g., cows, pigs, horses, cats, dogs, rats etc.) and a primate (e.g., monkey and human), and most preferably a human. In some embodiments, the subject is a non-human animal such as a farm animal (e.g., a horse, pig, or cow) or a pet (e.g., a dog or cat). In a specific embodiment, the subject is a human.

(33) As used herein, the term “agent” refers to any molecule, compound, methodology and/or substance for use in the prevention, treatment, management and/or diagnosis of a disease or condition.

(34) As used herein, the term “effective amount” refers to the amount of a therapy that is sufficient to result in the prevention of the development, recurrence, or onset of a disease or condition, and one or more symptoms thereof, to enhance or improve the prophylactic effect(s) of another therapy, reduce the severity, the duration of a disease or condition, ameliorate one or more symptoms of a disease or condition, prevent the advancement of a disease or condition, cause regression of a disease or condition, and/or enhance or improve the therapeutic effect(s) of another therapy.

(35) As used herein, the phrase “pharmaceutically acceptable” means approved by a regulatory agency of the federal or a state government, or listed in the U.S. Pharmacopeia, European Pharmacopeia, or other generally recognized pharmacopeia for use in animals, and more particularly, in humans.

(36) As used herein, the term “therapeutic agent” refers to any molecule, compound, and/or substance that is used for treating and/or managing a disease or disorder.

(37) As used herein, the terms “therapies” and “therapy” can refer to any method(s), composition(s), and/or agent(s) that can be used in the prevention, treatment and/or management of a disease or condition, or one or more symptoms thereof. In certain embodiments, the terms “therapy” and “therapies” refer to small molecule therapy.

(38) The term “derivative” or “derivatized” as used herein includes chemical modification of a compound of the disclosure, or pharmaceutically acceptable salts thereof or mixtures thereof. That is, a “derivative” may be a functional equivalent of a compound of the disclosure, which is capable of inducing the improved pharmacological functional activity in a given subject.

(39) As used herein, the terms “composition” and “formulation” are used interchangeably.

(40) As used herein, the term “topical delivery” means delivery of drug into systemic circulation through the skin.

(41) Transdermal Compositions

(42) According to certain embodiments, transdermal compositions described herein are for the prevention and/or treatment of nausea and/or vomiting associated with cancer chemotherapy.

(43) According to certain embodiments described herein, the pharmaceutical composition or transdermal formulation contains dronabinol in a form selected from a group such as co-crystals, amorphous form, crystalline form, its coated form, its solution, and its salts which can be anhydrous and/or hydrous alone or in combinations thereof. More preferably the transdermal formulation may include dronabinol selected from following forms: amorphous or co-crystals or crystalline or coated or its ethanolic solution either alone or in combination of thereof.

(44) One embodiment of the present disclosure can be a transdermal drug delivery system which may include without any limitation a transdermal formulation, a transdermal patch, a topical formulation, microneedles, iontophoresis, or a metered dose transdermal spray.

(45) Transdermal formulation which includes liquids for example without any limitation like solutions, suspensions, dispersions, emulsion. Transdermal formulation includes semisolids for example without any limitations like gels, ointments, emulsions, creams, suspension, paste, lotion, balm. Liquid formulation and/or gel formulation incorporated in transdermal patch is preferred. Transdermal formulations which include a polymer matrix can be, without any limitation, an adhesive matrix or a non-adhesive matrix.

(46) Without any limitation, transdermal patch may include all transdermal drug delivery systems stated in art preferably but not limited to reservoir patch, matrix patch, bilayer matrix patch, multilayer matrix patch, microreservoir patch, adhesive systems, transdermally applicable tape and other.

(47) In certain embodiments of the present disclosure, a transdermal patch comprises dronabinol contained in a reservoir or a matrix, and an adhesive which allows the transdermal patch to adhere

to the skin, allowing the passage of the dronabinol from the transdermal patch through the skin of the patient. The transdermal delivery system can be occlusive, semi-occlusive or non-occlusive, and can be adhesive or non-adhesive.

(48) In some embodiments, the transdermal patches provide for a constant rate of delivery of the active components of the transdermal patch over a predetermined time period. In some embodiments, the predetermined time period is about 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 7 days, 8 to 13 days, two weeks, or 15 days.

(49) In yet further embodiments, the transdermal patches described herein provide a steady absorption rate of the active components of the transdermal patches by the patient over a predetermined time. In some embodiments, the predetermined time period is 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 7 days, 8 to 13 days, two weeks, or 15 days.

(50) In yet further embodiments, the transdermal patches described herein provide a constant blood serum level of the active components of the transdermal patches in a patient over a predetermined time. In some embodiments, the predetermined time period is about 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 7 days, 8 to 13 days, two weeks, or 15 days.

(51) In yet further embodiments, the transdermal patches described herein provide a plasma concentration of the active components of the transdermal patches in a therapeutic range in a patient over a predetermined time. In some embodiments, the predetermined time period is about 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 7 days, 8 to 13 days, two weeks, or 15 days.

(52) In yet further embodiments, the transdermal patches described herein allow for reduced variability in dosage of active components in a patient over a predetermined time. In some embodiments, the predetermined time period is about 24 hours, 48 hours, 72 hours, 96 hours, 120 hours, 144 hours, 7 days, 8 to 13 days, two weeks, or 15 days.

(53) The topical formulation stated in the art which include, for example without any limitation, semisolids such as ointment, cream, emulsion, micro emulsion, nano emulsion, paste, balms, gels, lotions, mousses. Liquids such as solutions, suspensions, micro suspension, nano suspension, dispersions, nano dispersion etc. Sprays, aerosols, magma, etc. The topical formulation comprising dronabinol can be topically applied to the skin surface for transdermal delivery of dronabinol.

(54) The transdermal formulation and/or topical formulation of some embodiments of the present disclosure may include carriers or ingredients in effective amount either alone or in combinations thereof without any limitation to the following carriers or ingredients such as solvents, gelling agents, polymers, biodegradable polymers, penetration enhancers, emollients, skin irritation reducing agents, buffering agents, pH stabilizers, solubilizers, suspending agents, dispersing agents, stabilizers, plasticizers, tackifiers, surfactants, volatile chemicals, antioxidants, oxidants, chelating agents, complexing agents, diluents, excipients, material to prepare patch, material to prepare a matrix patch or a reservoir patch.

(55) Dronabinol may be dissolved, suspended, dispersed or uniformly mixed in the above stated single carrier, mixture of carriers and combinations of carrier.

(56) The desired optimum transdermal and/or topical formulation of dronabinol may comprise without any limitation to following carriers as stated from Example 1 to Example 11 either alone or in combinations thereof.

EXAMPLES

Example 1

(57) A transdermal formulation and/or topical formulation comprises solvents, alone or in combinations thereof, including but not limited to alcohol C1-C20 such as but not limited to (methanol, ethanol, isopropyl alcohol, butanol, propanol etc.), polyhydric alcohols, glycols such as but not limited to (propylene glycol, polyethylene glycol, dipropylene glycol, hexylene glycol, butylene glycol, glycerine etc.), derivative of glycols, pyrrolidone such as but not limited to (N methyl 2-pyrrolidone, 2-pyrrolidone etc.), sulfoxides such as but not limited to (dimethyl sulfoxide,

decymethylsulfoxide etc), dimethylisobutylidene, mineral oils, vegetable oils, water, polar solvents, semi polar solvents, non-polar solvents, volatile chemicals which can be used to make matrix patch such as but not limited to (ethanol, propanol, ethyl acetate, acetone, methanol, dichloromethane, chloroform, toluene, IPA), acids such as but not limited to acetic acid, lactic acid, levulinic acid, fatty acids, bases and others. The mentioned solvents are, in an embodiment, in the range of 0.01%-95% w/w or w/v.

Example 2

(58) A transdermal formulation and/or topical formulation comprises gelling agents and/or thickening and/or suspending agents, alone or in combinations, including but not limited to natural polymers, polysaccharides and its derivatives such as but not limited to (agar, alginic acid and derivatives, cassia tora, collagen, gelatin, gellum gum, guar gum, pectin, potassium, or sodium carageenan, tragacanth, xanthan, gum copal, chitosan, resin etc.), semisynthetic polymers and its derivatives such as without any limitation to cellulose and its derivatives (methylcellulose, ethyl cellulose, carboxymethyl cellulose, hydroxylpropyl cellulose, hydroxylpropylmethyl cellulose etc.), synthetic polymers and its derivatives such as without any limitation to carboxyvinyl polymers or carbomers (carbopol 940, carbopol 934, carbopol 971p NF), polyethylene, and its copolymers etc, clays such as but not limited to (silicates, bentonite), silicon dioxide, polyvinyl alcohol, acrylic polymers (eudragit), acrylic acid esters, polyacrylate copolymers, polyacrylamide, polyvinyl pyrrolidone homopolymer and polyvinyl pyrrolidone copolymers such as but not limited to (PVP, Kollidon 30, Poloxamer), isobutylene, ethyl vinyl acetate copolymers, natural rubber, synthetic rubber, pressure sensitive adhesives such as silicone polymers such as but not limited to (bio-psa 4302, bio-psa 4202 etc.), acrylic pressure sensitive adhesives such as but not limited to (DURO-TAK 87-2156, DURO-TAK 387-2287, etc.), polyisobutylene such as but not limited to (polyisobutylene low molecular weight, polyisobutylene medium molecular weight, polyisobutylene 35000 mw, etc), acrylic copolymers, rubber based adhesives, hot melt adhesives, styrene-butadiene copolymers, bentonite, all water and/or organic solvent swellable polymers, etc. The mentioned agents are, in an embodiment, in the range of 0.1% 70% w/w or w/v.

Example 3

(59) The transdermal formulation and/or topical formulation of the disclosure may comprise permeation enhancers known to those skilled in the art either alone or in combination thereof without any limitation to the following, such as sulfoxides, and similar chemicals such as but not limited to (dimethylsulfoxide, dimethylacetamide, dimethylformamide, decymethylsulfoxide, dimethylisobutylidene etc), azone, pyrrolidones such as but not limited to (N-methyl-2-pyrrolidone, 2-pyrrolidone etc.), esters, fatty acid esters such as but not limited to (propylene glycol monolaurate, butyl ethanoate, ethyl ethanoate, isopropyl myristate, isopropyl palmitate, methyl ethanoate, decyl oleate, glycerol monooleate, glycerol monolaurate, lauryl laurate etc.), fatty acids such as but not limited to (capric acid, caprylic acid, lauric acid, oleic acid, myristic acid, linoleic acid, stearic acid, palmitic acid etc.), alcohols, fatty alcohols and glycols such as but not limited to (oleyl alcohol, nathanol, dodecanol, propylene glycol, glycerol etc.), ethers alcohol such as but not limited to (diethylene glycol monoethyl ether), urea, triglycerides such as but not limited to triacetin, polyoxyethylene fatty alcohol ethers, polyoxyethylene fatty acid esters, esters of fatty alcohols, essential oils, surfactant type enhancers such as but not limited to (brij, sodium lauryl sulfate, tween, polysorbate), terpene, terpenoids and all penetration or permeation enhancers referred in the book "Percutaneous Penetration Enhancers" (Eric W. Smith, Howard I. Maibach, 2005. Nov, CRC press). The mentioned agents are, in an embodiment, in the range of 0.01%-95% w/w or w/v.

Example 4

(60) A transdermal formulation and/or topical formulation comprises plasticizers alone or in combination thereof, including without any limitation glycerol and its esters, phosphate esters, glycol derivatives, sugar alcohols, sebacic acid esters, citric acid esters, tartaric acid esters, adipate, phthalic acid esters, triacetin, oleic acid esters and all the plasticizers which can be used in

transdermal drug delivery system referred in the book "Handbook of Plasticizers" (George Wypych, 2004, Chem Tec Publishing). The mentioned agents are, in an embodiment, in the range of 0.01%-95% w/w or w/v.

Example 5

(61) A transdermal formulation and/or topical formulation comprises emollients, humectants, and/or skin irritation reducing agents either alone or in combinations thereof without any limitation to following like petrolatum, lanolin, mineral oil, dimethicone, zinc oxide, glycerin, propylene glycol and others. The mentioned agents are, in an embodiment, in the range of 0.01%-95% w/w or w/v.

Example 6

(62) A transdermal formulation and/or topical formulation comprises solubilizers, surfactants, emulsifying agents, dispersing agents alone or in combination thereof without any limitation to following: surfactants, including anionic, cationic, nonionic and amphoteric surfactants, such as the sorbitan oleates under the trade name SPAN such as but not limited to SPAN 80, SPAN 20 etc. and the polysorbates such as but not limited to Polysorbate 20, Polysorbate 40, Polysorbate 60, Polysorbate 80 etc., propylene glycol monocaprylate type I, propylene glycol monocaprylate type II, propylene glycol dicaprylate, medium chain triglycerides, propylene glycol monolaurate type II, linoleoyl polyoxyl-6 glycerides, oleoyl-polyoxyl-6-glycerides, lauroyl polyoxyl-6-glycerides, polyglyceryl-3-dioleate, diethylene glycol monoethyl ether, propylene glycol monolaurate type I, polyglyceryl-3-dioleate, caprylocaproyl polyoxyl-8 glycerides etc, cyclodextrins and others. The mentioned agents are, in an embodiment, in the range of 0.01% 95% w/w or w/v.

Example 7

(63) Different techniques and ingredients can be used to increase the stability and/or solubility of dronabinol in the formulation such as without any limitation coating, encapsulation, microencapsulation, nanoencapsulation, lyophilization, chelating agents, complexing agents, etc.

Example 8

(64) A transdermal formulation and/or topical formulation comprises auxiliary pH buffering agents and pH stabilizers and similar compounds known to those skilled in the art to maintain the appropriate pH of formulation preferably in the range of 4.0-8.0 either alone or in combination thereof without any limitation to following such as phosphate buffer, acetate buffer, citrate buffer, etc., acids such as but not limited to (carboxylic acids, inorganic acids, sulfonic acids, vinylogous carboxylic acids, fatty acids, and others), base such as but not limited to (sodium hydroxide, potassium hydroxide, ammonium hydroxide, triethylamine, sodium carbonate, sodium bicarbonate) etc. The mentioned agents are, in an embodiment, in the range of 0.01%-30% w/w or w/v.

Example 9

(65) A transdermal formulation and/or topical formulation comprises antioxidants, such as but not limited to, sodium metabisulfite, citric acid, ascorbic acid, BHA, and BHT, oxidizing agents, stabilizers, discoloring agents, preservatives and similar compounds or chemicals to provide a stable formulation. The mentioned agents are, in an embodiment, in the range of 0.01%-50% w/w or w/v.

Example 10

(66) A transdermal formulation and/or topical formulation with dronabinol formulated in ointment and/or cream base, gels, lotions, and other topical formulations.

Example 11

(67) Materials to make the transdermal delivery system of the disclosure in patch form known to those skilled in the art, for example, such as but not limited to reservoir patch, matrix patch, drug in adhesives, transdermal films and may include, such as but are not limited to polymers, copolymers, derivatives, backing film, release membranes, release liners, etc. either alone or in combinations thereof. Pressure sensitive adhesives (such as but not limited to silicone polymers, rubber based

through the skin and into receptor compartment. It is important to confirm that the receptor fluid is always in contact with the skin. The receptor compartment was emptied at 24 hr intervals for assay of delta-9-THC and replaced with fresh receptor solution. In order to maintain the sink condition in receptor compartment, it is important keep the delta-9-THC concentration in receptor compartment less than 10% of its solubility. The experimental conditions are provided in Table 3:

(75) TABLE-US-00003 TABLE 3 Experimental Condition for In-vitro Permeability testing De-ionized water + 0.5% Brij-O(20) + Receiving Media 0.01% Sodium Azide Receiving Media Volume (mL) 13 Sample Volume (mL) 13 Sampling Interval (hr) 24 Franz-cell diffusion area (sqcm) 1.76 Membrane Type Human Cadaver Skin

(76) Flux of THC through the human cadaver skin was measured for a minimum period of 96 Hrs (4 days) and results of the flux measurement are provided in Table 4.

(77) TABLE-US-00004 TABLE 4 THC Flux Results 001 002 006 007 008 009 Total Amount of THC 98700 154400 64200 122700 90900 102800 Permeated at 96 hrs (ng/cm.sup.2) Flux (ng/cm.sup.2/hr) 1000 1600 700 1300 950 1070 Patch Area (cm.sup.2) (5 20 12 28 15 21 18 mg/day)

Example 13

(78) Additional synthetic delta-9-THC formulations for transdermal delivery (Formulation Nos. 010 through 018) were prepared by mixing ingredients as shown in Table 5:

(79) TABLE-US-00005 TABLE 5 Transdermal Synthetic delta-9-THC formulation nos. 010 to 018
010 011 012 013 014 015 016 017 018 Ingredients (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) THC 1.3 2.0 2.8 1.3 2.0 2.8 2.0 2.8 2.0 Ethanol 27.3 42.4 58.7 27.3 42.4 58.7 42.4 PG 64.3 50 34.6 71.4 55.6 38.5 — — — Water 7.1 5.6 3.9 — — — — — Dimethyl — — — — — 55.6 38.5 55.6 Sulfoxide (DMSO) Abbreviations: THC = tetrahydrocannabinol; PG = propylene glycol.

(80) Synthetic delta-9-THC formulations for transdermal delivery (010-018) were prepared by the same procedure described in Example 12. Flux measurement was also performed as described in Example 12. The experimental conditions are the same as provided in Table 3 of Example 12.

(81) Flux of THC through the human cadaver skin was measured for a minimum period of 96 Hrs (4 days) and results of the flux measurement experiments are provided in Table 6.

(82) TABLE-US-00006 TABLE 6 THC Flux Results Formulation No. 010 011 012 013 014 015 016 017 018 Total 75600 94900 102800 54000 145500 204600 35200 35800 53400 Amount of (31%) (20%) (52%) (33%) (63%) (43%) (22%) (1%) (6%) THC Permeated at 96 hrs (ng/cm.sup.2) Flux 800 1000 1070 600 1500 2100 400 400 600 (ng/cm.sup.2/hr) Patch Area 24 19 18 32 13 9 48 48 32 (cm.sup.2) (5 mg/day)

Example 14

(83) Additional synthetic delta-9-THC formulations for transdermal delivery (Formulation Nos. 019 to 027) were prepared by mixing ingredients as shown in Table 7.

(84) TABLE-US-00007 TABLE 7 Transdermal Synthetic delta-9-THC formulation nos. 019-027
019 020 021 022 023 024 025 026 027 Ingredients (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) (% W/V) THC 2.0 2.0 2.0 2.0 2.0 2.0 2.0 2.0 2.0 Ethanol 42.4 42.4 42.4 42.4 42.4 42.4 42.4 PG 44.4 44.4 44.4 44.4 44.4 44.4 44.4 44.4 NMP 11.1 Lauric acid 11.1 Oleic Acid 11.1 Lauryl 11.1 Lactate Oleyl 11.1 Alcohol GMO 11.1 Lactic Acid 11.1 HPC 11.1 (Klucel ®) Abbreviations: THC = tetrahydrocannabinol; PG = propylene glycol; NMP = n-methyl pyrrolidone; GMO = glycerol monooleate; HPC = hydroxypropyl cellulose.

(85) Synthetic delta-9-THC formulations for transdermal delivery (019-027) were prepared by the same procedure described in Example 12. Flux measurement was also performed as described in Example 12. The experimental conditions are the same as provided in Table 3 of Example 12.

(86) Flux of THC through the human cadaver skin was measured for a minimum period of 72 hours (3 days) and results of the flux measurement experiments are provided in Table 8.

(87) TABLE-US-00008 TABLE 8 THC Flux Results Formulation No. 019 020 021 022 023 024

025 026 027 Total 56300 29600 20500 5100 5100 26100 36400 31300 Amount of (10%) (82%) (14%) (29%) (22%) (34%) (92%) (19%) (16%) THC Permeated at 72 hrs (ng/cm.sup.2) Flux 800 400 300 70 70 400 500 400 400 (ng/cm.sup.2/hr) Patch Area 24 48 64 276 276 48 38 48 48 (cm.sup.2) (5 mg/day)

Example 15

(88) Additional synthetic delta-9-THC formulations for transdermal delivery (Formulation Nos. 028 to 034) were prepared by mixing ingredients as shown in Table 9:

(89) TABLE-US-00009 TABLE 9 Transdermal Synthetic delta-9-THC formulation nos. 028 to 034

	032	034	028	029	030	031	(% 033	(% Ingredients	(% W/V)	(% W/V)	(% W/V)	(% W/V)	W/V)	(% W/V)	W/V)	THC	2.0	2.0	2.0	2.0	2.0	2.0	2.0	Ethanol	42.4	42.4	42.4	42.4	42.4	42.4	42.4	PG	27.8
27.8	27.8	27.8	27.8	27.8	27.8	27.8	1,3	Butanediol	27.8	Dipropylene	Glycol	27.8	Hexylene	Glycol	27.8	PEG-400	27.8	Tween-20	27.8	Tween-80	27.8	Span-20	27.8										

(90) Synthetic delta-9-THC formulations for transdermal delivery (formulation nos. 028 to 034) were prepared by the same procedure described in Example 12. Flux measurement was also performed as described in Example 12. The experimental conditions are the same as provided in Table 3 of Example 12.

(91) Flux of THC through the human cadaver skin was measured for a minimum period of 120 hours (5 days) and results of the flux measurement experiments are provided in Table 10.

(92) TABLE-US-00010 TABLE 10 Transdermal Synthetic delta-9-THC formulations Formulation Nos. 028 029 030 031 032 033 034 Total Amount of 103400 86400 115300 61400 7400 30100 43800 THC Permeated at (9%) (17%) (53%) (43%) (80%) (137%) (47%) 120 Hrs (ng/cm.sup.2) Flux (ng/cm.sup.2/hr) 900 700 1000 500 60 300 400 Patch Area (cm.sup.2) 22 28 19 38 322 64 48 (5 mg/day)

Example 16

(93) Additional synthetic delta-9-THC formulations for transdermal delivery patches (Formulation Nos. 035 to 038) were prepared by mixing ingredients as shown in Table 11:

(94) TABLE-US-00011 TABLE 11 Transdermal Synthetic delta-9-THC formulation nos. 035 to 038

	035	036	037	038	Ingredients	(% W/W)	(% W/W)	(% W/W)	(% W/W)	THC	2.0	2.0	2.0	2.0	PG	27.8	27.8	27.8	27.8	Hexylene	Glycol	27.8	27.8	27.8	27.8	Durotak	9301	42.4	Durotak	2516	42.4	Durotak	2207	42.4	Silicone	Adhesive	42.4

(95) To prepare a transdermal patch containing synthetic delta-9-THC, all of the components from Table 11, with the exception of the THC, were mixed together with stirring for 18 hours. Next, the THC was added 30 minutes before spreading the formulation. The formulation was spread using a commercial benchtop spreader. Specifically, the formulation matrix is evenly spread onto an 8×14 inch sheet of release liner (such as 3M 9744) to a thickness of 0.5 mm. The sheet is then place in an oven at 100° F. for one hour to evaporate off the ethyl acetate and ethanol adhesive solvent. An opaque backing membrane (such as 3M 9730 NR film) with low permeability to oxygen, for inhibition of photo and oxidative degradation, is then carefully applied to the sheet by hand to avoid formation of bubbles and voids. A circular die (1.5 inches diameter) was used to cut patches (7 cm.sup.2) for subsequent studies.

(96) The general procedure for flux measurements of transdermal formulations in the examples above was as follows. The human cadaver skin, stored at -80° C., was thawed at room temperature in PBS, and visually inspected for defects before use. Transdermal flux was measured using standard Franz diffusion cells composed of a cylindrical donor compartment and a separate water jacketed cylindrical receptor compartment with the volume of 13 mL. The human cadaver skin was clamped between the two compartments with the dermis side facing toward the receptor compartment. The general procedure for flux measurement of the transdermal adhesive patch is as follows. The release liner is peeled off the patch and the adhesive surface is applied to a piece of human cadaver skin (Example 16, Table 11 only). The transdermal patch was adhered to the skin with the patch on the side of the skin in contact with the donor compartment. The receptor

compartment was filled with receptor medium, held at constant temperature, and constantly stirred to collect the THC as it diffuses from the adhered patch, through the skin and into receptor compartment. It was confirmed that the receptor fluid was always in contact with the skin. The receptor compartment was emptied at 24 hour intervals for assay of delta-9-THC and replaced with fresh receptor solution. In order to maintain the sink condition in receptor compartment, the delta-9-THC concentration in the receptor compartment was maintained at less than 10% of its solubility. The experimental conditions are the same as provided in Table 3 of Example 12.

Example 17

(97) Synthetic delta-9-THC (THC) formulations for transdermal delivery (Formulation Nos. 040 through 042) were prepared by mixing ingredients as shown in Table 12:

(98) TABLE-US-00012 TABLE 12 Transdermal Synthetic delta-9-THC formulation no. 040 to 042

THC—040	THC—041	THC—042	Excipients (Solution)	(Solution)	(Gel)	THC in ethanol (20% w/w)
13.1%	12.8%	13.6%	Ethanol	43.2%	38.6%	41.9%
			Propylene Glycol	43.7%	38.6%	43.4%
			Hydramol	10.0%	—	—
			Klucel HF	—	—	1.0%

(99) The following examples used composition THC_040 as an example for preparing a transdermal patch. The above excipients except THC were mixed together by stirring for 18 hours and then the THC in ethanol was added into the above excipient mixture.

(100) The general procedure for flux measurement of transdermal formulation was conducted as follows. The human cadaver skin, stored at -80°C , was thawed at room temperature in PBS, and visually inspected for defects before using in the study. Transdermal flux was measured using standard Franz diffusion cells composed of a cylindrical donor compartment and a separate water jacketed cylindrical receptor compartment with the volume of 13 ml. The human cadaver skin was clamped between the two compartments with the dermis side facing toward the receptor compartment. The receptor compartment was filled with receptor medium, held at constant temperature, and constantly stirred to collect the THC as it diffuses through the skin and into receptor compartment, making sure that that the receptor fluid was always in contact with the skin. The receptor compartment was emptied at 24 hr intervals for assay of delta-9-THC and replaced with fresh receptor solution. In order to maintain the sink condition in receptor compartment, the delta-9-THC concentration in receptor compartment was kept at less than 10% of its solubility. The experimental conditions were as follow:

(101) TABLE-US-00013 Experimental Condition for In-vitro Permeability testing Receiving Media De-ionize water + 0.5% Brij-O(20) + 0.01% Sodium Azide Receiving Media Volume (ml) 13 Sample Volume (ml) 13 Sampling Interval (hr) 24, 48, 72, 96, 120 Franz-cell diffusion area (sqcm) 1.76 Membrane Type Human Cadaver Skin

(102) TABLE-US-00014 TABLE 13 Flux is measure for a minimum period of 120 Hrs (5 days)

THC—040	THC—041	THC—042	Avg Flux 0-24 hr, $\mu\text{g/sqcm/hr}$	0.11	0.00	0.00	(% RSD)	Avg Flux 24-48 hr, $\mu\text{g/sqcm/hr}$	0.72	0.59	2.16	(% RSD)	(173.21)	(62.53)	(62.29)	Avg Flux 48-72 hr, $\mu\text{g/sqcm/hr}$	1.55	0.83	2.93	(% RSD)	(50.46)	(12.42)	(33.55)	Avg Flux 72-96 hr, $\mu\text{g/sqcm/hr}$	2.15	0.66	3.03	(% RSD)	(49.90)	(39.91)	(10.27)	Avg Flux 96-120 hr, $\mu\text{g/sqcm/hr}$	1.66	0.22	2.03	(% RSD)	(39.69)	(87.14)	(10.04)	Avg Flux 24-120 hr, $\mu\text{g/sqcm/hr}$	1.24	0.46	2.03	(% RSD)	(97.00)	(40.40)	(29.04)
---------	---------	---------	---	------	------	------	---------	--	------	------	------	---------	----------	---------	---------	--	------	------	------	---------	---------	---------	---------	--	------	------	------	---------	---------	---------	---------	---	------	------	------	---------	---------	---------	---------	---	------	------	------	---------	---------	---------	---------

Example 18

(103) Synthetic delta-9-THC (THC) formulations for transdermal delivery (Formulation Nos. 043 through 0048) were prepared by mixing ingredients as shown in Table 14:

(104) TABLE-US-00015 TABLE 14 THC Matrix System

THC	THC	THC	THC	THC	THC	THC
Excipients 043-R1	044-R1	045-R1	046	047	048	THC
4.05%	4.05%	4.05%	7.81%	7.86%	7.79%	Oleic acid
—	4.45%	—	—	—	—	Propylene
17.00%	12.55%	12.16%	—	14.94%	15.19%	Glycol Isopropyl
—	4.46%	—	—	—	—	Palmitate
—	—	—	15.17%	—	—	Lactic Acid
—	—	14.85%	15.92%	14.86%	GMO	—
—	—	—	4.84%	4.87%	4.95%	Hydramol
—	—	—	—	—	—	—
5.49%	5.20%	5.28%	BIO-	78.95%	78.95%	79.32%
—	—	—	—	—	—	PSA 4501 DURO-
—	—	—	—	—	—	—

(107) TABLE-US-00016 TABLE 15 Experimental Condition for In-vitro Permeability testing De-ionized water + 0.5% Brij-O(20) + Receiving Media 0.01% Sodium Azide Receiving Media Volume (mL) 13 Sample Volume (mL) 13 Sampling Interval (hr) 24, 48, 72, 96, 120, 144 Franz-cell diffusion area (sqcm) 1.76 Membrane Type Human Cadaver Skin

(109) TABLE-US-00017

	TABLE 16	THC Flux Results	THC	THC	THC	THC	THC	THC	043-R1
	044-R1	045-R1	046	047	048	Avg Flux 0-	0.00	0.28	0.00
						0.00	0.00	0.00	N/A
						24 hr, µg/sqcm/	(173.21)	hr	
(% RSD)						Avg Flux 24-	0.00	1.01	0.00
						0.00	0.00	0.00	N/A
						48 hr, µg/sqcm/	(16.39)	hr	(% RSD)
						Avg Flux 48-	0.00	0.96	0.52
						0.00	0.00	N/A	72 hr, µg/sqcm/
						(12.86)	(82.26)	hr	(% RSD)
						Avg Flux 72-	0.00	1.07	0.84
						0.00	0.00	N/A	96 hr, µg/sqcm/
						(13.81)	(2.82)	hr	(% RSD)
						Avg Flux 96-	0.00	1.22	0.94
						0.00	0.00	N/A	120 hr, µg/sqcm/
						(15.51)	(2.75)	hr	(% RSD)
						Avg Flux 120-	0.00	1.03	0.56
						0.00	0.00	0.00	N/A
						144 hr, µg/sqcm/	(12.46)	(86.67)	hr
						(% RSD)	Avg Flux 24-	0.00	0.93
							0.43	0.00	0.00
						N/A	144		
						hr, µg/sqcm/	(21.54)	(33.55)	hr
						(% RSD)			

(110) The effect of gelling agents and their concentration on the permeation of THC through human cadaver skin. THC gel formulation can be gelled by gelling agents including but not limited to, natural polymers such as natural polymers, polysaccharides and its derivatives such as but not limited to (agar, alginic acid and derivatives, cassia tora, collagen, gelatin, gellum gum, guar gum, pectin, potassium or sodium carrageenan, tragacanth, xanthum gum, copal, starch, chitosan, resin etc.), synthetic polymers and its derivatives such as without any limitation to carboxyvinyl polymers or carbomers (carbopol 940, carbopol 934, carbopol 971), polyethylene and its copolymers etc. clays such as silicate etc. polyvinyl alcohol, polyacrylamide, polyvinyl pyrrolidone homopolymer and polyvinyl pyrrolidone copolymers (PVP, Poloxamer), acrylic acid its ester, polyacrylate copolymers, isobutylene, ethylene vinyl acetate copolymers, natural rubbers, synthetic rubbers such as styrene-diene copolymers, styrene-butadiene block copolymers, isoprene block copolymers, acrylonitrile butadiene rubber, butyl rubber or neoprene rubber, as well as pressure

sensitive adhesive based on silicone, or “hot-melt adhesive”. In addition, other than human cadaver skin, clobazam can be evaluated with other artificial membranes including but not limited to cellulose membrane, silicone membranes (polydimethylsiloxane), liposome coated membranes, solid-supported liquid membranes, lecithin organogel membrane and other. Besides the gel formulation of clobazam, other dosage forms including but not limited to ointment, creams, emulsion, liposomes, etc. may be used.

Example 20

(111) The effect of enhancers or solubilizers on the flux of clobazam through human cadaver skin was evaluated. The desire optimum composition of clobazam gel formulation contained dimethylsulfoxide (DMSO), dimethylisorbide (DMI), Lactic acid, Tween-20, highly purified diethylene glycol monoethyl ether (Transcutol P), dipropylene glycol, polyethylene glycol-400, propylene glycol (PG), Hexylene Glycol (HG), Lauroglycol-90. Apart from above mentioned enhancers and/or solubilizers, the clobazam transdermal delivery can be influenced by enhancers and/or solubilizers including but not limited water, sulfoxides, and similar as but not limited to (dimethylsulfoxide, dimethylacetamide, chemicals such dimethylformamide, decylmethylsulfoxide, dimethylisorbide etc), azone, pyrrolidones such as but not limited to (N-methyl-2-pyrrolidone, 2-pyrrolidone etc), esters such as but not limited to (Propylene glycol monolaurate, butyl ethanoate, ethyl ethanoate, isopropyl myristate, isopropyl palmitate, methyl ethanoate, decyl oleate, glycerol monooleate, glycerol monolaurate, lauryl laurate etc), fatty acids such as but not limited to (capric acid, caprylic acid, lauric acid, oleic acid, myristic acid, linoleic acid, stearic acid, palmitic acid etc), alcohols, fatty alcohols and glycols such as but not limited to (oleyl alcohol, nathanol, dodecanol, propylene glycol, glycerol etc), ethers such as but not limited to (diethylene glycol monoethyl ether), urea, polyoxyethylene fatty alcohol ethers, polyoxyethylene fatty acid esters, esters of fatty alcohols, esters of long chain fatty acids with methyl, ethyl or isopropyl alcohol, esters of fatty alcohols with acetic acid, lactic acid, as well as oleic acid diethanolamine, essential oils, terpene and terpenoids such as but not limited to (terpineol, limonene, thymol, cineole etc), surfactant type enhancers (polysorbate 80, polysorbate 20 etc.), liposomes, niosomes, transferomes, ethanosomes, polysorbate such as but not limited to (polysorbate 20, polysorbate 40, polysorbate 60, polysorbate 80 etc), span such as but not limited to (span 80, span 20 etc), surfactants such as (anionic, cationic, nonionic and amphoteric), propylene glycol monocaprylate type I, propylene glycol monocaprylate type II, propylene glycol dicaprylate, medium chain triglycerides, propylene glycol monolaurate type II, linoleoyl polyoxyl-6 glycerides, Caprylic glyceride, oleoyl-polyoxyl-6-glycerides, lauroyl polyoxyl-6-glycerides, polyglyceryl-3-dioleate, diethylene glycol monoethyl ether, propylene glycol monolaurate type I etc, cyclodextrins, polyhydric alcohol, especially 1,2-propanediol, butanediol, glycerine, polyethylene glycol (m.w. 100 and higher), Dimethyl Sulfoxide, Dimethyl Isosorbide, tetrahydrofurfuryl alcohol, diethyl toluamide, monoisopropylidene glycerine and others Solubilizers, surfactants, emulsifying agents, dispersing agents and similar compounds or chemicals known to those skilled in the art can be used either alone or in combination thereof

Example 21

(112) Oral bioavailability of THC is only 10-20%. For our calculation purpose, we took an average bioavailability of 15% 17. Accordingly, the actual dose delivering to patient upon oral delivery is described in Table 17:

(113) TABLE-US-00018 TABLE 17 Theoretical dose required from Transdermal Dosage form.

Transdermal Dose range	Oral Dose (mg/day)	5 mg/day	0.75	10 mg/day	1.5	20 mg/day	3
Flux Required=Dose/Surface area=3.0 mg/day/surface area=3000 ug/24 hr/50 sqcm=2.5 ug/sqcm/hr							

Accordingly, 50 sqcm patch with 2.5 µg/sqcm/hr flux will deliver 3 mg of drug in one day through transdermal route which is equivalent to 20 mg/day oral dose. According to table, The maximum dose of approved dronabinol is 20 mg/day which is feasible to deliver through this formulation.

Claims

1. A method for the treatment and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS, comprising: selecting a patient in need of the treatment and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS; topically applying or instructing to topically apply a pharmaceutical composition in the form of a transdermal matrix patch comprising: dronabinol, wherein the concentration of the dronabinol is from about 1% to about 15% w/w; wherein the pharmaceutical composition further comprises: about 10% to about 17% w/w of a solvent comprising propylene glycol; about 4% to about 10% w/w of a penetration enhancer comprising oleic acid; about 50% to about 80% w/w of a silicone pressure sensitive adhesive; at least one suspending agent comprising silicon dioxide; an antioxidant comprising BHT, wherein the transdermal matrix patch provides an average flux of the dronabinol of about 0.43 to about 0.93 $\mu\text{g}/\text{cm}^2/\text{hr}$ over at least 6 days, wherein said applying achieves topical delivery of dronabinol for the treatment and/or control of nausea and/or vomiting associated with cancer chemotherapy and/or anorexia associated with weight loss in patients with AIDS.
 2. The method of claim 1, wherein the pharmaceutical composition is topically applied once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week, or once in ten days.
 3. The method of claim 1, wherein the pharmaceutical composition is topically applied two to six times in a day, once in a day, once in two days, once in three days, once in four days, once in five days, once in six days, once in a week.
 4. The method of claim 1, wherein the pharmaceutical composition is a transdermal delivery system.
-