

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12385021
Kind Code	B2
Date of Patent	August 12, 2025
Inventor(s)	Jiang; Xiwen et al.

Heat-resistant DNA polymerase mutant with high amplification activity

Abstract

The present invention provides a heat-resistant DNA polymerase mutant with high amplification activity. Particularly, the present invention uses protein directed evolution technology to construct a random mutation library for the polymerase active domain of Taq enzyme, and gradually adds screening pressure, so that unsuitable mutations will be eliminated naturally, and mutations with dominant traits will gradually accumulate. Finally, a series of amino acid sites and their mutations that are critical to Taq enzyme amplification and polymerization performance will be selected, and a Taq enzyme mutant with high amplification activity will be obtained.

Inventors:	Jiang; Xiwen (Guangdong, CN), Liu; Aishan (Guangdong, CN)
Applicant:	Daan Gene Co., Ltd. (Guangdong, CN)
Family ID:	78373147
Assignee:	DAAN GENE CO., LTD. (Guangdong, CN)
Appl. No.:	16/957276
Filed (or PCT Filed):	April 30, 2020
PCT No.:	PCT/CN2020/088341
PCT Pub. No.:	WO2021/217597
PCT Pub. Date:	November 04, 2021

Prior Publication Data

Document Identifier	Publication Date
US 20220325259 A1	Oct. 13, 2022

Publication Classification

Int. Cl.: C12N9/12 (20060101)

U.S. Cl.:

CPC C12N9/1252 (20130101); C12Y207/07007 (20130101);

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
9523085	12/2015	Hogrefe	N/A	C12N 9/1252
2005/0112637	12/2004	Chatterjee	435/6.12	C12N 9/1252
2019/0225951	12/2018	Meng et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
101528919	12/2008	CN	N/A
107475216	12/2016	CN	N/A
109486788	12/2018	CN	N/A
109957557	12/2018	CN	N/A
111032863	12/2019	CN	N/A
2009010251	12/2008	WO	N/A
2011014885	12/2010	WO	N/A
WO-2012097318	12/2011	WO	C12N 9/1252
2016183294	12/2015	WO	N/A

OTHER PUBLICATIONS

Suzuki et al., Proc. Natl. Acad. Sci. USA, vol. 93, pp. 9670-9675, Sep. 1996. cited by examiner
Ngo et al. in The Protein Folding Problem and Tertiary Structure Prediction, 1994, Merz et al. (ed.), Birkhauser, Boston, MA, pp. 433 and 492-495. cited by examiner

Wang et al., "A novel strategy to engineer DNA polymerases for enhanced processivity and improved performance in vitro", Nucleic Acids Research, vol. 32, No. 3, Published online Feb. 18, 2004, pp. 1197-1207. cited by applicant

Suzuki et al., "Thermus Aquaticus DNA Polymerase | Mutants With Altered Fidelity. Interacting Mutations in the O-helix", The Journal of Biological Chemistry, vol. 275, No. 42, Oct. 20, 2000, pp. 32728-32735. cited by applicant

Yamagami et al., "Mutant Taq DNA polymerases with improved elongation ability as a useful reagent for genetic engineering", Frontiers in Microbiology, Sep. 3, 2014, vol. 5, Article 461, 11 pages provided. cited by applicant

Zhang et al., "Direct DNA Amplification From Crude Clinical Samples Using a PCR Enhancer Cocktail and Novel Mutants of Taq", Journal of Molecular Diagnostics, vol. 12, No. 2, Mar. 2010, 10 pages. cited by applicant

Vainshtein et al., "Peptide rescue of an N-terminal truncation of the Stoffel fragment of taq DNA polymerase", Protein Science, accepted Jun. 25, 1996, pp. 1785-1792. cited by applicant
International Search Report (with English translation) and Written Opinion issued in PCT/CN2020/088341, dated Jan. 27, 2021, 13 pages provided. cited by applicant
The extended European search report issued in European Application No. 20761492.6, dated Feb. 17, 2022, 9 pages provided. cited by applicant
RecName: Full=DNA polymerase I, thermostable ; AltName: Full=Taq polymerase 1, GenBank, created Feb. 1, 1991, 6 pages, cited in ISR. cited by applicant
Yamagami et al., "Mutant Taq DNA polymerases with improved elongation ability as a useful reagent for genetic engineering", Frontiers in Microbiology, vol. 5 Issue No. SEP, published 2014, 10 pages, cited in EESR. cited by applicant
Arezi et al., "Compartmentalized self-replication under fast PCR cycling conditions yields Taq DNA polymerase mutants with increased DNA-binding affinity and blood resistance", Frontiers in Microbiology, Published Aug. 14, 2014, 10 pages, cited in EESR. cited by applicant

Primary Examiner: Hutson; Richard G

Attorney, Agent or Firm: HSML P.C.

Background/Summary

TECHNICAL FIELD

(1) The present invention belongs to the field of biotechnology. Specifically, the present invention relates to a heat-resistant DNA polymerase mutant with high amplification activity.

BACKGROUND

(2) Taq enzyme is a heat-resistant DNA polymerase derived from a heat-resistant bacterium, *Thermus aquaticus*, with a molecular weight of 94 KDa, which has an optimal reaction temperature range of 75–80° C., an active half-life at 95° C. of 40 minutes and 5'-3'exonuclease activity in the presence of magnesium ions. Because of its resistance to high temperature, it is widely used in polymerase chain reaction (PCR) and is the enzyme of first choice for nucleic acid amplification and detection and other reactions. The commercial Taq enzyme is cloned and expressed using an *E. coli* prokaryotic expression system. Increasing sensitivity, accuracy, and durability of PCR reactions are required in modern molecular biological detection technology, and wild-type Taq enzyme cannot fulfill the requirement of practical applications. In order to make it more suitable for certain technologies, many attempts have been made in mutation of the Taq enzyme sequence, such as adding a DNA binding domain to obtain a stronger extension activity (Wang Y (2004). A novel strategy to engineer DNA polymerases for enhanced processivity and improved performance in vitro. Nucleic Acids Res 32, 1197-1207); site-directed mutagenesis and deletion of domains is performed to obtain a higher fidelity (Suzuki M, Yoshida S, Adman E T, Blank A, Loeb L A (2000) *Thermus Aquaticus* DNA polymerase I mutants with altered fidelity. Interacting mutations in the 0-Helix. J Biol Chem 275:32728-32735), a higher DNA polymerization activity (Mutant Taq DNA polymerases with improved elongation ability as a useful reagent for genetic engineering. Front Microbiol 5:461. doi: 10.3389/fmicb.2014.00461), tolerance to a high concentration of inhibitors (Zhang Z, Kermekchiev M B, Barnes W M (2010) Direct DNA amplification from crude clinical samples using a PCR enhancer cocktail and novel mutants of Taq. J Mol Diagn 12:152-161), a reduced 5'-3'exonuclease activity (Vainshtein I, Atrazhev A, Eom S H, Elliott J F, Wishart D S, Malcolm B A (1996) Peptide rescue of an N-Terminal truncation of the Stoffel fragment of Taq DNA polymerase. Protein Sci 5:51785-51792).

(3) There are several ways to transform Taq enzyme. 1: Adding a domain to obtain a new property. For example, adding a single-stranded binding domain (SSB) or DNA binding protein Sso7 to enhance the binding of Taq enzyme to primers and template DNA, so that it has stronger extension ability and continuous synthesis ability, and is suitable for amplification reaction of long DNA fragments. However, adding a domain will directly increase the molecular weight of Taq enzyme, which may reduce the solubility and stability of Taq enzyme. Then the yield of prokaryotic expression production is reduced. 2: Removing non-essential domains on Taq enzyme. For example, the 5'-3'exonuclease domain (first 280 amino acids of N-terminus of Taq enzyme) is deleted, Taq enzyme only retains the active region of the nucleic acid polymerase, which reduces the possibility of degradation of primers and template DNA by a high concentration of Taq enzyme in order to achieve the purpose of improving the polymerization activity of Taq enzyme. However, the Taq enzyme mutant obtained from this method does not have 5'-3'exonuclease activity, so it is not suitable for quantitative PCR reaction based on the Taq man probe method, and the scope of application is limited. 3: Site-directed mutation method. Site-directed mutation is performed on the amino acids in the active site, magnesium ion binding site, and DNA binding site to increase the affinity of each site for substrates, templates, and primers, thereby improving tolerance to various inhibitors. Due to the complexity of protein structure, some amino acids far from the active site may also affect the overall structure of the enzyme. Therefore, it is difficult to modify the enzyme as a whole with mutation of only amino acids in certain active sites. Moreover, it is difficult to predict the effect of mutations at various sites on the overall structure with existing computer simulation technology. The workload of preparing mutants by site-directed mutation and screening mutants is very large, its efficiency is low, and some sites that significantly affect the activity cannot be identified.

SUMMARY OF INVENTION

(4) The object of the present invention is to provide a heat-resistant DNA polymerase mutant with high amplification activity.

(5) In the first aspect of the present invention, there is provided a mutated DNA polymerase that is mutated at one or more sites selected from the group consisting of: V453, F495, E507, K508, T509, A518, S624, Y672, E734, R737, F749, T757, L764, H785, wherein the amino acid residues are numbered based on SEQ ID NO. 2.

(6) In another preferred example, the activity of the mutated DNA polymerase is at least 1.5 times, preferably at least 2 times; more preferably at least 3 times larger than that of wild-type DNA polymerase (SEQ ID NO: 2).

(7) In another preferred example, the amino acid sequence of wild-type DNA polymerase is set forth in SEQ ID NO: 2.

(8) In another preferred example, the amino acid sequence of the mutated DNA polymerase has at least 80% homology, more preferably at least 90% homology, most preferably at least 95% homology, such as at least 96%, 97%, 98%, 99% homology to SEQ ID NO: 2.

(9) In another preferred example, the mutated DNA polymerase is selected from the group consisting of mutants 1-20:

(10) TABLE-US-00001 Mutant No. Mutated amino acid 1 E507A, K508L, E734E, F749K 2 K508L, V453A, R737K 3 E734G 4 F749G, K508L, L764K 5 E507Q, T757S 6 H785G 7 S624T, F749V 8 E734F, F749V 9 K508L, R737W, Y672R 10 E507H, H785L 11 A518Q, E734M 12 F495R, F749T 13 K508L, F749T, E734F 14 R737P, S624K 15 T757W, V453G, E507M 16 F749E, H785G, F495G 17 E734F, Y672P 18 T509L, H785K 19 E734G, T757S, L764Q 20 K508L, V453A, A518Q

(11) In another preferred example, the number of mutation sites in the mutated DNA polymerase is 1-4, preferably 2 or 3.

(12) In another preferred example, the mutated DNA polymerase is selected from each specific mutant enzyme listed in Table 2.

- (13) In another preferred example, the mutated DNA polymerase includes the mutation sites of each specific mutant enzyme listed in Table 2.
- (14) In another preferred example, the mutated DNA polymerase is mutated based on the wild-type DNA polymerase as set forth in SEQ ID NO: 2, and the mutated DNA polymerase includes a mutation site selected from the group consisting of: (1) E507A, K508L, E734E, F749K; (2) K508L, V453A, R737K (3) E734G (4) F749G, K508L, L764K (5) E507Q, T757S (6) H785G (7) S624T, F749V (8) E734F, F749V (9) K508L, R737W, Y672R (10) E507H, H785L (11) A518Q, E734M (12) F495R, F749T (13) K508L, F749T, E734F (14) R737P, S624K (15) T757W, V453G, E507M (16) F749E, H785G, F495G (17) E734F, Y672P (18) T509L, H785K (19) E734G, T757S, L764Q; and (20) K508L, V453A, A518Q.
- (15) In the second aspect of the present invention, there is provided a polynucleotide molecule encoding the mutated DNA polymerase according to the first aspect of the present invention.
- (16) In the third aspect of the present invention, there is provided a vector containing the nucleic acid molecule according to the second aspect of the present invention.
- (17) In the fourth aspect of the present invention, there is provided a host cell containing the vector according to the first aspect of the present invention or a chromosome integrated with the nucleic acid molecule according to the second aspect of the present invention.
- (18) In another preferred example, the host cell is a prokaryotic cell or an eukaryotic cell.
- (19) In another preferred example, the prokaryotic cell is *E. coli*.
- (20) In another preferred example, the eukaryotic cell is a yeast cell.
- (21) In the fifth aspect of the present invention, there is provided a method for preparing the mutated DNA polymerase according to the first aspect of the present invention, comprising the steps of: (i) culturing the host cell according to the fourth aspect of the present invention under suitable conditions to express the mutated DNA polymerase; and (ii) isolating the mutated DNA polymerase.
- (22) In another preferred example, the temperature for culturing the host cell in step (i) is 20° C.-40° C., preferably 25° C.-37° C., such as 35° C.
- (23) In the sixth aspect of the present invention, there is provided a kit comprising the mutated DNA polymerase according to the first aspect of the present invention.
- (24) It should be understood that, within the scope of the present invention, the above technical features of the present invention and the technical features specifically described below (e.g., in embodiments) can be combined with each other, thereby forming a new or preferred technical solution. As space is limited, not every technical solution will be illustrated herein.
-

Description

DETAILED DESCRIPTION

- (1) After extensive and intensive research, the inventor has screened out a series of amino acid sites and mutations that play a key role in Taq enzyme amplification performance and polymerization performance, and has obtained Taq enzyme mutants with high amplification performance using protein directed evolution technology to construct a random mutation library for the polymerase active domain of Taq enzyme, and gradually applying screening pressure to eliminate unsuitable mutations naturally and to accumulate mutations with dominant traits. On this basis, the present invention has been completed.
- (2) Before describing the present invention, it should be understood that the present invention is not limited to the specific methods and experimental conditions, as such methods and conditions may vary. It should also be understood that the terminology as used herein is for the purpose of describing specific embodiments and is not intended to be limiting, and the scope of the present invention will be limited only by the appended claims.

(3) Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by those skill in the art to which this invention belongs. As used herein, when used in reference to a recited value, the term “about” means that the value can vary from the recited value by no more than 1%. For example, as used herein, the expression “about 100” includes 99 and 101 and all values therebetween (e.g., 99.1, 99.2, 99.3, 99.4, etc.).

(4) Although any methods and materials similar or equivalent to those described in the present invention can be used in the practice or testing of the present invention, the preferred methods and materials are exemplified herein.

(5) Taq Enzyme

(6) Taq enzyme is widely used in polymerase chain reaction (PCR) and is the enzyme of first choice for reactions such as nucleic acid amplification and detection. The commercial Taq enzyme is cloned and expressed using an *E. coli* prokaryotic expression system.

(7) The DNA sequence of the wild-type Taq enzyme is as follows:

(8) TABLE-US-00002 (SEQ ID NO: 1)

ATGCGTGGCATGCTGCCGCTTTTCGAGCCTAAGGGACGCGTTCTTCT
TGTGGATGGACATCATCTGGCGTACCGTACCTTTCATGCCCTGAAGGGCC
TGACCACTTCGCGTGGGGAACCCGTGCAAGCAGTTTATGGATTGCGCCAAA
TCGTTACTTAAGGCTCTGAAGGAGGATGGTGATGCGGTCATTGTTGTGTT
CGACGCAAAAGCTCCCTCGTTCCGTCACGAGGCCTACGGCGGCTATAAAG
CTGGGCGTGCACCCACACCTGAGGATTTTCCCCGGCAACTTGCTTTGATA
AAGGAATTAGTAGACCTGTTAGGCCTGGCGCGGTTAGAAGTGCCGGGGTTA
CGAAGCAGATGACGTCTTGGCTAGTTTAGCGAAAAAGGCTGAAAAAGAG
GGATATGAAGTGCGGATCCTGACCGCGGATAAAGATCTGTATCAACTGTT
GTCCGACCGTATTCACGTGCTTCATCCGGAGGGCTACTTGATAACCCCGG
CTTGGCTGTGGGAGAAATATGGGCTGCGTCCAGATCAGTGGGCTGATTAT
CGTGCACTTACAGGCGATGAATCTGATAATCTTCCCGGCGTCAAGGGGAT
TGGTGAGAAAACCGCCCGTAAACTTTTGGAGGAGTGGGGCAGCTTGGAG
GCGCTGTTGAAGAATCTGGATCGTTTGAAACCCGCTATACGGGAAAAAAT
CTTGGCGCACATGGACGACTTAAACTGTCTTGGGACCTGGCGAAAGTTC
GTACTGATTTGCCGCTGGAGGTGCGACTTTGCGAAGCGTCGCGAGCCCGAT
CGTGAACGTCTTCGCGCATTTCTGGAGCGTTTAGAATTTGGCTCCCTGTTG
CATGAGTTTGGTTTGCTTGAAAGCCCGAAGGCACTTGAGGAAGCTCCTTG
GCCTCCGCCTGAGGGCGCTTTTGTGCGATTGTCTTGAGCCGTAAAGAAC
CGATGTGGGCGGACTTACTGGCCCTTGCTGCTGCTCGTGGGGGTCGCGTG
CATCGCGCACCGGAGCCATACAAAGCACTTCGTGACCTTAAAGAAGCCCG
TGGCTTGTTGGCAAAAGATTTAAGTGTCCTGGCTTTACGCGAGGGGCTTGG
GCTTACCACCGGGAGATGATCCGATGCTTTTGGCCTATCTGCTGGACCCG
AGCAACACGACTCCAGAGGGCGTTGCCCCGTGTTATGGCGGAGAATGGA
CGGAGGAGGCGGGAGAGCGCGCAGCGTTAAGCGAGCGTCTGTTTGCTAA
TCTGTGGGGACGCTTAGAGGGAGAGGAGCGCCTGTTGTGGTTGTACCGTG
AAGTGGAACGGCCGCTGAGTGCAAGTGTTAGCTCACATGGAAGCAACCGG
GGTGCGGCTGGACGTTGCGTATTTGCGTGCGCTGTCGTTAGAGGTGCGGG
AGGAAATAGCCCGTCTGGAGGCCGAAGTATTCGTTTGGCTGGCCATCCT
TTCAACCTGAACAGTCGGGATCAGCTGGAACGTGTACTTTTTGATGAACT
GGGGCTGCCCCGCCATCGGCAAAACCGAAAAAACCGGCAAACGTAGCACC
TCTGCGGCAGTGCTGGAAGCGTTACGTGAAGCTCATCCGATTGTGGAGAA
AATTCTGCAATATCGCGAATTGACGAAACTGAAGAGCACCTATATTGATC
CGCTGCCAGACTTAATTCACCCCCGTACCGGACGGTTGCATACCCGCTTC
AACCAGACCGCGACGGCGACAGGGCGGCTGAGTAGCAGCGATCCGAACC
TGCAAAACATTCCCGTGCGTACCCCGCTGGGTCAGCGTATTCGCCGTGCT

TTCAATTGCGTGTGTTGGCCACCTGTCTGGGCGACGAAAACCTTAATACGCG
TGTTTCAAGAAGGTCGTGACATACATACTGAAACCGCGTCCTGGATGTTT
GGAGTCCCACGGGAGGCTGTTCGATCCTCTTATGCGTCGTGCCGCCAAAAC
AATTAACCTTCGGAGTTCTGTACGGCATGTCTGGCACATCGTTTATCACAGG
AACTGGCGATTCCGTATGAAGAAGCGCAGGCCTTCATAGAACGTTATTTTC
CAATCATTCCCCAAGGTGCGGGCCTGGATTGAGAAGACCCTGGAAGAGG
GCCGTCTGTCGTGGCTATGTAGAGACTCTGTTCTGGACGTCTGGCGGTATGTA
CCCGATCTTGAGGCCCGTGTGAAGTCCGTTCTGTGAGGCAGCAGAACGTAT
GGCGTTTAACATGCCAGTCCAGGGCACAGCGGCGGACCTGATGAAATTA
GCTATGGTTAAGCTGTTTCCGCGTTTGGGAAGAAATGGGCGCTCGTATGCT
GTTACAGGTTTCATGACGAGTTAGTATTAGAAGCACCGAAGGAGCGTGCCG
AAGCCGTGGCCCGGTTAGCCAAAGAGGTAATGGAAGGCGTCTACCCCCTT
GCAGTCCCGCTTGAAGTCGAAGTTGGCATAGGGGAAGACTGGTTATCTGC GAAGGAA

(9) The amino acid sequence of the wild-type Taq enzyme is as follows:

(10) TABLE-US-00003 (SEQ ID NO: 2)

MRGMLPLFEPKGRVLLVDGHHLAYRTFHALKGLTTSRGEVPQAVYGFAK
SLLKALKEDGDAVIVVFDKAPSRHEAYGGYKAGRPTPEDFPRQLAL
IKELVDLLGLARLEVPGYEADDVLASLAKKAEKEGYEVRILTADKDLYQ
LLSDRIHVLHPEGYLITPAWLWEKYGLRPDQWADYRALTGDESDNLPV
KGIGECTARKLLEEWGSLEALLKNLDRLKPAIREKILAHMDDLKLSWDL
AKVRTDLPLEVDFAKRREPDRERLRAFLERLEFGSLLHEFGLLESPKAL
EEAPWPPPEGAFFVGFVLSRKEPMWADLLALAAARGGRVHRAPEPYKALR
DLKEARGLLAKDLSVLALREGLGLPPGDDPMLLAYLLDPSNTTPEGVAR
RYGGEWTEEAGERAAALSERLFANLWGRLEGEERLLWLYREVERPLSAVL
AHMEATGVRLDVAYLRALSLEVAEEIARLEAEVFRLAGHPFNLSRDQL
ERVLFDELGLPAIGKTEKTGKRSTSAVLEALREAHPIVEKILQYRELT
KLKSTYIDPLPDLIHPRTGRLHTRFNQTATATGRLSSSDPNLQNIPIVRT
PLGQRIRRAFIAEEGWLLVALDYSQIELRVLAHLSGDENLIRVFQEGRD
IHTETASWMFGVPREAVDPLMRRAAKTINFGVLYGMSAHRLSQELAIPY
EEAQAFIERYFQSFPKVRAWIEKTLEEGRRRGYVETLFGRRRYVPDLEA
RVKSVREAERMAFNMPVQGTAAADLMKLMVKLFPRLEEMGARMMLLQVH
DELVLEAPKERAEAVARLAKEVMIEGVYPLAVPLEVEVGIGEDWLSAKE

(11) The amino acid positions and mutation modes that are highly related to Taq enzyme amplification activity are identified through directed evolution in the present invention. Related mutated amino acid sites include: V453, F495, E507, K508, T509, A518, S624, Y672, E734, R737, F749, T757, L764, and H785, and the amino acid residues are numbered based on SEQ ID NO: 2. Mutation of the above amino acid position to any other amino acid may produce a Taq enzyme mutant with higher activity. Preferred mutant form includes: E507A/Q/H/M, K508L, E734G/F/M, F749K/G/V/T/E, L764K/Q, V453A/G, R737K/W/P, T757S/W, H785G/L/K, S624T/K, Y672R/P, A518Q, F495G/R, T509L.

(12) The amino acid sites and mutation modes thereof highly related to Taq enzyme activity are screened from a random mutation library with directed evolution technology in the present invention. The number of mutants is 10⁵ times more than that of the site-directed mutation, which is more conducive to identify the mutation sites with synergistic effects that cannot be predicted by existing computer simulation techniques. Furthermore, based on the principle of directed evolution, the accumulated dominant traits are those most suitable for the added screening conditions, so the mutants obtained are certainly the optimal individuals among all mutants.

(13) In a preferred embodiment of the present invention, the activity of the mutant DNA polymerase provided by the present invention is at least 1.2 times, preferably at least 1.3 times,

more preferably at least 1.5 times, such as more than 2 times higher than that of the wild-type DNA polymerase (SEQ ID NO: 2).

(14) In a preferred embodiment of the present invention, the method of assaying the activity of the mutated DNA polymerase and wild-type DNA polymerase (SEQ ID NO: 2) is as follows:

(15) TABLE-US-00004 PCR reaction solution: pET28a vector 100 pg Taq enzyme mutant or wild-type Taq enzyme 10 ng 10XTaq enzyme reaction solution (100 mM 2 ul Tris, 500 mM KCl, 100 mM (NH₄)₂SO₄, pH 8.0) pET28_F primer 4 pmol pET28_R primer 4 pmol dNTPs (2.5 mM) 2 uL ddH₂O added to 20 ul PCR program: 95° C. for 5 minutes, 30 cycles (95° C. for 15 seconds, 55° C. for 15 seconds, 72° C. for 10 seconds), 4° C. ∞

(16) The PCR product was purified by ethanol precipitation, the light absorption of the product was measured at 260 nm, and the total amount of PCR product (ng) corresponding to each cycle was calculated.

(17) Wherein, the following pair of primers was used in the PCR reaction:

(18) TABLE-US-00005 pET28_F primer: (SEQ ID NO: 9)

TACGGTTAACCTTTGAATCA pET28_R primer: (SEQ ID NO: 10)

GTTACCTGGTTAACTGTACT.

(19) Dividing the total amount of PCR products obtained using Taq enzyme mutants by the total amount of PCR products obtained using wild-type Taq enzymes is the activity multiple of Taq enzyme mutants compared to wild-type Taq enzymes.

(20) Those skilled in the art can use conventional methods to obtain the sequence of the Taq enzyme gene of the present invention, for example, by complete artificial synthesis or PCR synthesis. A preferred method of synthesis is asymmetric PCR. The method of asymmetric PCR uses a pair of primers of unequal amounts to produce a large amount of single-stranded DNA (ssDNA) after PCR amplification. This pair of primers are called unrestricted primer and restricted primer, respectively, and their ratio is generally 50-100:1. In the first 10-15 cycles of the PCR reaction, the amplification product is mainly double-stranded DNA, but when the restricted primer (low concentration primer) is depleted, the PCR guided by the unrestricted primer (high concentration primer) will produce a large amount of single-stranded DNA. The primers used in PCR can be appropriately selected according to the sequence information of the present invention disclosed herein, and can be synthesized by conventional methods. The amplified DNA/RNA fragments can be separated and purified by conventional methods such as gel electrophoresis.

(21) The Taq enzyme of the present invention can be expressed or produced by conventional recombinant DNA technology, comprising the steps of: (1) Transforming or transducing a suitable host cell with a polynucleotide encoding the protein of the present invention, or with a recombinant expression vector containing the polynucleotide; (2) Culturing the host cell in a suitable culture medium; (3) Isolating and purifying the protein of interest from the culture medium or cell to obtain the Taq enzyme.

(22) Methods well known by those skilled in the art can be used to construct an expression vector containing the DNA sequence encoding the Taq enzyme of the present invention and suitable transcription/translation control signals, preferably a commercially available vector, pET28. These methods include in vitro recombinant DNA technology, DNA synthesis technology, in vivo recombinant technology, etc. The DNA sequence can be effectively linked to an appropriate promoter in an expression vector to guide mRNA synthesis. The expression vector also includes a ribosome binding site for translation initiation and a transcription terminator. In addition, the expression vector preferably contains one or more selectable marker genes to provide a phenotypic trait for selection of transformed host cells.

(23) The recombinant vector includes in the 5' to 3' direction: a promoter, a gene of interest and a terminator. If desired, the recombinant vector may also include the following elements: a protein purification tag, 3' polynucleotide signal, an untranslated nucleic acid sequence, a transporting and targeting nucleic acid sequence, a selection marker (antibiotic resistance gene, fluorescent protein,

etc.), an enhancer, or an operator.

(24) Methods for preparing recombinant vectors are well known to those skill in the art. The expression vector may be a bacterial plasmid, bacteriophage, yeast plasmid, plant cell virus, mammalian cell virus, or other vector. To sum up, any plasmid and vector can be used as long as it is able to replicate and stabilize in a host.

(25) Those skilled in the art can construct a vector containing the promoter of the present invention and/or the gene sequence of interest using well-known methods. These methods comprise in vitro recombinant DNA technology, DNA synthesis technology, in vivo recombinant technology etc.

(26) The expression vector of the present invention can be used to transform an appropriate host cell so that the host transcribes the target RNA or expresses the target protein. The host cell may be a prokaryotic cell, such as *E. coli*, *Corynebacterium glutamicum*, *Brevibacterium flavum*, *Streptomyces*, *Agrobacterium*; or a lower eukaryotic cell, such as a yeast cell; or a higher eukaryotic cell, such as a plant cell. Those skilled in the art know how to select appropriate vectors and host cells. Transformation of host cells with recombinant DNA can be performed using conventional techniques well known to those skilled in the art. When the host is a prokaryote (such as *E. coli*), it can be treated with CaCl₂ method or electroporation. When the host is a eukaryote, the following DNA transfection methods can be used: calcium phosphate co-precipitation method, conventional mechanical methods (such as microinjection, electroporation, liposome packaging, etc.). The plant can also be transformed using methods such as *Agrobacterium* transformation or gene gun transformation, such as leaf disc method, immature embryo transformation method, flower bud soaking method, etc. The transformed plant cells, tissues or organs can be regenerated into plants using conventional methods to obtain transgenic plants.

(27) Term “operably linked” means that the target gene to be transcribed for expression is linked to its control sequence in a manner conventional in the art for expression.

(28) Culturing of Engineering Bacteria and Fermentation Production of Target Protein

(29) After obtaining the engineered cells, the engineered cells can be cultured under suitable conditions to express the protein encoded by the gene sequence of the present invention.

Depending on the host cell, the culture medium used in the culture can be selected from a variety of conventional media and cultured under conditions suitable for the growth of the host cell. When the host cell grows to an appropriate cell density, the selected promoter is induced by an appropriate method (such as temperature conversion or chemical induction), and the cell is cultured for another period of time.

(30) In the present invention, conventional fermentation conditions can be used. Representative conditions comprise (but are not limited to): (a) Regarding temperature, the fermentation and induction temperature of Taq enzyme is maintained at 25-37° C.; (b) Regarding pH during the induction period, the pH during the induction period is controlled at 3-9; (c) Regarding dissolved oxygen (DO), DO is controlled at 10-90%, and the maintenance of dissolved oxygen can be achieved by the input of oxygen/air mixed gas; (d) Regarding the supplements, the type of the supplements should preferably include carbon sources such as glycerin, methanol, and glucose, which can be supplemented alone or mixed; (e) Regarding IPTG concentration during the induction period, conventional induction concentrations can be used in the present invention, usually the IPTG concentration is controlled at 0.1-1.5 mM; (f) Regarding the induction time, without any particular limitation, it is usually 2-20 hours, preferably 5-15 hours.

(31) The target protein Taq enzyme of the present invention is present within *E. coli* cells, the host cells are collected by a centrifuge, and then the host cells are crushed by high-pressure, mechanical force, enzymatic hydrolysis or other cell disruption methods, preferably the high-pressure method to release recombinant proteins. The host cell lysate can be preliminarily purified by methods such as flocculation, salting out, and ultrafiltration, and then purified by chromatography, ultrafiltration, etc., or can be directly purified by chromatography.

(32) Chromatography techniques include cation exchange chromatography, anion exchange

chromatography, gel filtration chromatography, hydrophobic chromatography, affinity chromatography, etc. Common chromatography methods comprise:

(33) 1. Anion Exchange Chromatography:

(34) Anion exchange chromatography media include (but are not limited to): Q-Sepharose®, DEAE-Sepharose®. If the salt concentration of the fermentation sample is high, which affects the binding to the ion exchange medium, the salt concentration needs to be reduced before ion exchange chromatography. The sample can be exchanged with an equilibration buffer by dilution, ultrafiltration, dialysis, gel filtration chromatography, etc. until it is similar to the corresponding ion exchange column equilibration solution system, and then the sample is loaded for gradient elution of salt concentration or pH.

(35) 2. Hydrophobic Chromatography:

(36) Hydrophobic chromatography media include (but are not limited to): Phenyl-Sepharose™, Butyl-Sepharose™, Octyle-Sepharose™. The salt concentration of the sample is increased by adding NaCl, (NH₄)₂SO₄, etc., then the sample is loaded, and eluted by reducing the salt concentration. heteroproteins with large differences in hydrophobicity is removed by hydrophobic chromatography.

(37) 3. Gel Filtration Chromatography

(38) Hydrophobic chromatography media include (but are not limited to): Sephacryl®, Superdex™, Sephadex®. The buffer system is exchanged by gel filtration chromatography, or it is further purified.

(39) 4. Affinity Chromatography

(40) Affinity chromatography media include (but are not limited to): HiTrap™ HeparinHPColumns.

(41) 5. Membrane Filtration

(42) Ultrafiltration media include: organic membranes such as polysulfone membranes, inorganic membranes such as ceramic membranes, and metal membranes. The purpose of purification and concentration may be achieved with membrane filtration.

The Advantages of the Present Invention are

(43) (1) The heat-resistant DNA polymerase mutant with high amplification activity of the present invention has a significant increase in the amount of products from amplification at the same number of PCR cycles as the wild-type Taq enzyme. (2) The heat-resistant DNA polymerase mutant with high amplification activity of the present invention can produce the same amount of product from amplification in a shorter time under the same conditions as the wild-type Taq enzyme, thereby increasing the efficiency of detection.

(44) The present invention will be further described in detail below in conjunction with specific examples. It should be understood that these examples are only used to illustrate the present invention and not to limit the scope of the present invention. The experimental methods without detailed conditions in the following examples are generally in accordance with conventional conditions such as those described in Sambrook. J et al. "Guide to Molecular Cloning Laboratory" (translated by Huang Peitang et al., Beijing: Science Press, 2002), or in accordance with the conditions recommended by the manufacturer. Unless otherwise illustrated, percentages and parts are calculated by weight. The experimental materials and reagents used in the following examples can be obtained commercially unless otherwise illustrated.

Example 1: Construction of Random Mutant Plasmid of Taq Enzyme

(45) The DNA sequence of polymerase active domain of Taq enzyme (coding sequence of amino acids 423-831) was amplified by low-fidelity PCR (Error-PCR) with a mutation rate of 0.3%, and then the remaining coding sequence (sequence of amino acids 1-423) was ligated and cloned into pET28a prokaryotic expression vector to obtain a random mutant plasmid of Taq enzyme. Specific steps were as follows: 1) Using Taq-pET28a plasmid as a template, and designing primer T (1-423) to amplify the Taq (1-423) fragment.

(46) TABLE-US-00006 Taq (1-423) DNA Seq (SEQ ID NO: 3)
ATGCGTGGCATGCTGCCGCTTTTCGAGCCTAAGGGACGCGTTCTTCT
TGTGGATGGACATCATCTGGCGTACCGTACCTTTCATGCCCTGAAGGGCC
TGACCACTTCGCGTGGGGAACCCGTGCAAGCAGTTTATGGATTCGCCAAA
TCGTTACTTAAGGCTCTGAAGGAGGATGGTGATGCGGTCATTGTTGTGTT
CGACGCAAAAGCTCCCTCGTTCCGTCACGAGGCCTACGGCGGCTATAAAG
CTGGGCGTGACCCACACCTGAGGATTTTCCCCGGCAACTTGCTTTGATA
AAGGAATTAGTAGACCTGTTAGGCCTGGCGCGGTTAGAAGTGCCGGGTTA
CGAAGCAGATGACGTCTTGGCTAGTTTAGCGAAAAAAGGCTGAAAAAGAG
GGATATGAAGTGCGGATCCTGACCGCGGATAAAGATCTGTATCAACTGTT
GTCCGACCGTATTACAGTGCTTCATCCGGAGGGCTACTTGATAACCCCGG
CTTGGCTGTGGGAGAAATATGGGCTGCGTCCAGATCAGTGGGCTGATTAT
CGTGCACTTACAGGCGATGAATCTGATAATCTTCCCCGGCGTCAAGGGGAT
TGGTGAGAAAACCGCCCGTAAACTTTTGGAGGAGTGGGGCAGCTTGGAGG
CGCTGTTGAAGAATCTGGATCGTTTGAAACCCGCTATACGGGAAAAAATC
TTGGCGCACATGGACGACTTAAAACTGTCTTGGGACCTGGCGAAAGTTCG
TACTGATTTGCCGCTGGAGGTCGACTTTGCGAAGCGTCGCGAGCCCGATC
GTGAACGTCTTCGCGCATTCTGGAGCGTTTAGAATTTGGCTCCCTGTTG
CATGAGTTTGGTTTGCTTGAAAGCCCGAAGGCACTTGAGGAAGCTCCTTG
GCCTCCGCCTGAGGGCGCTTTTGTGCGATTGTCTTGAGCCGTAAAGAAC
CGATGTGGGCGGACTTACTGGCCCTTGCTGCTGCTCGTGGGGGTCGCGTG
CATCGCGCACCGGAGCCATACAAAGCACTTCGTGACCTTAAAGAAGCCCG
TGGCTTGTTGGCAAAGATTTAAGTGTCTTGGCTTTACGCGAGGGGCTTGG
GCTTACCACCGGGAGATGATCCGATGCTTTTGGCCTATCTGCTGGACCCG
AGCAACACGACTCCAGAGGGCGTTGCCCGTCGTTATGGCGGAGAATGGA
CGGAGGAGGCGGGAGAGCGCGCAGCGTTAAGCGAGCGTCTGTTTGCTAA
TCTGTGGGGACGCTTAGAGGGAGAG T1-423_PF: (SEQ ID NO.: 4) 5'
ATATCATATGCGTGGCATGCTGCCGCTTTT 3' T1-423_PR: (SEQ ID NO.: 5) 5'
GCATGAATTCCGTCTCCTCTCCCTCTAAGC 3'

PCR Reaction System and Program:

(47) TABLE-US-00007 plasmid Taq-pET28a 10 ng T1-423_PF primer 4 pmol T1-423_PR primer 4 pmol 2.5 mM dNTP 2 ul 10X reaction buffer 2 ul KAPA HiFi DNA Polymerase 5 U ddH2O added to total volume of 20 ul PCR program: 95° C. for 3 minutes, 25 cycles (95° C. for 30 seconds, 60° C. for 30 seconds, 72° C. for 1 minute), 72° C. for 3 minutes, storage at 4° C.

(48) The PCR product was purified and recovered with DNA gel recovery kit, digested with NdeI and XhoI, and ligated into pET28a vector. The sequence was confirmed by sequencing to be correct, and the resulting plasmid was named as Taq (1-423)-pET28. 2) Using Taq-pET28a plasmid as a template, using Clontech Diversify® PCR Random Mutagenesis Kit (Takara Bio, Dalian, PT3393-2), and designing primers (TMu_F/R) to amplify Taq (423-822) fragments.

(49) TABLE-US-00008 Taq (423-832) DNA Seq (SEQ ID NO.: 6)
GGAGAGGAGCGCCTGTTGTGGTTGTACCGTGAAGTGGAACGGCCGC
TGAGTGCAAGTGTAGCTCACATGGAAGCAACCGGGGTGCGGCTGGACGTT
GCGTATTTGCGTGCGCTGTCGTTAGAGGTCGCGGAGGAAATAGCCCGTCT
GGAGGCCGAAGTATTCCGTTTGGCTGGCCATCCTTTCAACCTGAACAGTC
GGGATCAGCTGGAACGTGTACTTTTTGATGAACTGGGGCTGCCCCGCCATC
GGCAAAACCGAAAAAACCGGCAAACGTAGCACCTCTGCGGCAGTGCTGG
AAGCGTTACGTGAAGCTCATCCGATTGTGGAGAAAATTCTGCAATATCGC
GAATTGACGAAACTGAAGAGCACCTATATTGATCCGCTGCCAGACTTAAT
TCACCCCCGTACCGGACGGTTGCATACCCGCTTCAACCAGACCGCGACGG
CGACAGGGCGGCTGAGTAGCAGCGATCCGAACCTGCAAAACATTCCCGT

GCGTACCCTGGGTACGCGTTCGCGCTGGCTTTTCATTGCCGAGGAAG
 GCTGGCTGCTGGTCGCGCTGGACTACTCGCAAATCGAATTGCGTGTGTTG
 GCCCACCTGTCGGGCGACGAAAACCTTAATACGCGTGTTCAGAAGGTCG
 TGACATACATACTGAAACCGCGTCCTGGATGTTTGGAGTCCCACGGGAGG
 CTGTCGATCCTCTTATGCGTCGTGCCGCCAAAACAATTAACCTTCGGAGTT
 CTGTACGGCATGTTCGGCACATCGTTTATCACAGGAAGTGGCGATTCCGTA
 TGAAGAAGCGCAGGCCTTCATAGAACGTTATTTCCAATCATTCCCCAAGG
 TGCGGGCCTGGATTGAGAAGACCCTGGAAGAGGGCCGTCGTCGTGGCTAT
 GTAGAGACTCTGTTTCGGACGTCGGCGGTATGTACCCGATCTTGAGGCCCG
 TGTGAAGTCCGTTTCGTGAGGCAGCAGAACGTATGGCGTTTAACATGCCAG
 TCCAGGGCACAGCGGCGGACCTGATGAAATTAGCTATGGTTAAGCTGTTT
 CCGCGTTTGGAAAGAAATGGGCGCTCGTATGCTGTTACAGGTTTCATGACGA
 GTTAGTATTAGAAGCACCGAAGGAGCGTGCCGAAGCCGTGGCCCGGTTA
 GCCAAAGAGGTAATGGAAGGCGTCTACCCCTTGCAGTCCCGCTTGAAGT
 CGAAGTTGGCATAGGGGAAGACTGGTTATCTGCGAAGGAATAA TMu_F: (SEQ ID
 NO.: 7) 5' GGAGAGGAGCGCCTGTTGTGGTTGT 3' TMu_R: (SEQ ID NO.: 8) 5'
 TTATTCCTTCGCAGATAACCAGTCT 3'

(50) PCR reaction system and program:

(51) TABLE-US-00009 10* Titanium Taq buffer 5 ul dGTP (2 mM) 1 ul 50* Diversify dNTP Mix 1 ul TMu_F primer 10 pmol TMu_R primer 10 pmol Titanium Taq 1 ul ddH.sub.2O added to 50 ul 95° C. for 3 minutes, 25 cycles (95° C. for 30 seconds, 60° C. for 30 seconds, 68° C. for 2 minutes), 68° C. for 5 minutes, storage for 4° C.

(52) The PCR product was digested with BsmBI and XhoI, then ligated with Taq (1-423)-pET28 plasmid digested with BsmBI and XhoI, BL21 (DE3) expression host bacteria was transformed with the ligated product, and the number of transformants was counted.

Example 2: Expression and Directed Evolution Screening of Taq Enzyme Mutants

(53) BL21 (DE3) expression strain was transformed with the Taq enzyme mutant plasmid and induced to express Taq enzyme mutation library. The BL21 (DE3) induced expression bacteria containing Taq enzyme mutant library were dispersed and packaged by emulsion PCR system, and subjected to PCR reaction to amplify and obtain DNA containing Taq enzyme mutant fragments. Then, the DNA fragments produced from amplification by emulsion PCR were subjected to high-fidelity PCR secondary amplification using Taq enzyme-specific primers, and the amplified DNA products were re-cloned into the pET28a expression vector to complete a screening process. Then, the screening process of emulsion PCR-secondary high-fidelity PCR-cloning into pET28a expression vector was repeated, and the extension time of emulsion PCR in each screening was gradually shortened to accumulate mutant populations with high extension activity and high amplification activity. The specific steps were as follows: 1) The transformant obtained in Example 1 was taken and inoculated into LB medium, incubated at 37° C. for 6 hours with shaking, isopropylthiogalactoside (IPTG) was added to a final concentration of 0.1 mM, and incubated at 37° C. for 3 hours. The cells were collected by centrifugation, washed twice with ddH.sub.2O, and finally resuspended with ddH.sub.2O. The light absorption (OD600 value) of the cell solution was determined at 600 nm, and it was diluted to the final concentration of OD600=1.0 with ddH.sub.2O. 2) Preparation of oil phase solution

(54) TABLE-US-00010 Tween-80 200 ul Triton X-100 25 ul Mineral oil 10 ml

(55) The above three reagents were combined and mixed evenly. 3) Preparation of aqueous phase reaction solution

(56) The cell resuspension solution prepared in step 1) with OD600=1.0 was diluted 100 times with ddH.sub.2O to prepare the following reaction solution

(57) TABLE-US-00011 10XTaq enzyme reaction solution (100 mM 13 ul Tris, 500 mM KCl, 100 mM (NH.sub.4).sub.2SO.sub.4, pH 8.0) BSA (100 mg/ml) 26 uL pET28_F primer 10 pmol

pET28_R primer 10 pmol dNTPs (2.5 mM) 26 uL Diluted cell resuspension solution 26 ul dd H.sub.2O added to 260 ul

(58) TABLE-US-00012 pET28_F primer: (SEQ ID NO.: 9)

TACGGTTAACCCTTTGAATCA pET28_R primer: (SEQ ID NO.: 10)

GTTACCTGGTTAAACTGTACT 4) Preparation of emulsion system

(59) 200 ul of aqueous phase+400 ul of oil phase were taken and mixed in a 2 ml tube, shaken at high speed for 10 minutes on a vortex shaker. 5 of PCR tubes were taken, each was filled with 100 ul of mixed solution. PCR program: 95° C. for 5 minutes, 25 cycles (95° C. for 30 seconds, 55° C. for 30 seconds, 72° C. for 2 minutes), 72° C. for 5 minutes, 4° C. ∞ 5) The emulsion PCR product was transferred to a 1.5 ml tube, with 166 ul of water-saturated ether added, vortexed for 30 seconds, centrifuged at 12000 rpm for 10 minutes, and the lower liquid phase was removed, allowed to stand at room temperature for 10 minutes until the ether volatilized. The liquid product was extracted with phenol-chloroform method, and then recovered by ethanol precipitation overnight. 6) High-fidelity PCR secondary amplification products

(60) The product of step 4) was used as a template for secondary PCR amplification

(61) TABLE-US-00013 The product of step 4) 2 ul Taq_F primer 4 pmol Taq_R primer 4 pmol dd H.sub.2O 6 ul KAPA HiFi mix 10 ul PCR program: 95° C. for 5 minutes, 20 cycles (95° C. for 30 seconds, 62° C. for 30 seconds, 72° C. for 2 minutes), 72° C. for 5 minutes, 4° C. ∞

(62) TABLE-US-00014 Taq_F primer: (SEQ ID NO.: 11)

ATGCGTGGCATGCTGCCGCTTTTCGAGCCTAAGGGACG Taq_R primer: (SEQ ID NO.: 12) TTCCTTCGCAGATAACCAGTCTTCCCCTATGCCAACTTCGAC 7) The PCR

product was purified with DNA product purification recovery kit, and then relinked to pET28a expression vector. So far, one round of screening was completed. 8) Steps (1)-(6) were repeated for the transformants relinked to pET28a vector, the conditions of emulsion PCR was changed according to the program in the table below, and selection pressure was gradually added to the mutation library.

(63) The second round of screening: 95° C. for 5 minutes, 25 cycles (95° C. for 30 seconds, 55° C. for 30 seconds, 72° C. for 1.5 minutes), 72° C. for 5 minutes, 4° C. ∞

(64) The third round of screening: 95° C. for 5 minutes, 20 cycles (95° C. for 30 seconds, 55° C. for 30 seconds, 72° C. for 1.5 minutes), 72° C. for 5 minutes, 4° C. ∞

(65) The fourth round of screening: 95° C. for 5 minutes, 15 cycles (95° C. for 30 seconds, 55° C. for 30 seconds, 72° C. for 30 seconds), 72° C. for 5 minutes, 4° C. ∞

(66) After 4 rounds of screening, the resulting Taq enzyme mutant transformants were used in the high-throughput screening of Example 3.

Example 3: High-Throughput Screening of Taq Enzyme Mutants

(67) 384 clones were randomly picked out from the mutation library obtained in Example 2, after culturing and inducing for expression, their amplification activity was tested by high-throughput PCR reaction, and 20 mutants with high amplification activity were selected among them. The detailed steps were as follows: 1) 384 single clones were selected, inoculated into LB medium, incubated at 37° C. for 6 hours, with isopropylthiogalactoside (IPTG) added at a final concentration of 0.1 mM, and incubated at 37° C. for 3 hours. 2) The cells after induction culture were collected by centrifugation, with lysate containing 0.1 mg/ml lysozyme (50 Mm Tris, 50 Mm NaCl, 5% glycerol pH8.5) added. The cells were resuspended, incubated at 37° C. for 10 minutes, and heated at 75° C. for 30 minutes. Then they were centrifuged at 12000 rpm for 10 minutes, and the supernatant was taken. 3) A 96-well PCR plate was taken with the following reaction components added to each well:

(68) TABLE-US-00015 10XTaq enzyme reaction solution (100 mM 2 ul Tris, 500 mM KCl, 100 mM (NH.sub.4).sub.2SO.sub.4, pH 8.0) pET28_F primer 4 pmol pET28_R primer 4 pmol dNTPs (2.5 mM) 2 uL Treated supernatant 1 ul ddH2O added to 20 ul PCR program: 95° C. for 5 minutes, 20 cycles (95° C. for 30 seconds, 62° C. for 30 seconds, 72° C. for 60 seconds), 4° C. ∞

(69) 5 ul of the PCR product was subjected to agarose gel electrophoresis, the 20 clones with the highest yield were selected by comparing the yield of the PCR product in the supernatant prepared by each clone. The amplification yield of each mutant was 1.2 times to 2 times more than that of the wild-type.

Example 4: Confirmation of the Mutation Sites in the Dominant Taq Enzyme Mutant

(70) The DNA sequence of the Taq enzyme mutant selected in Example 3 was sequenced to determine the mutation in its amino acid sequence, and the high-frequency mutation sites and their mutation forms were counted.

(71) TABLE-US-00016 TABLE 1 Mutant No. Mutated amino acid 1 E507A, K508L, E734E, F749K 2 K508L, V453A, R737K 3 E734G 4 F749G, K508L, L764K 5 E507Q, T757S 6 H785G 7 S624T, F749V 8 E734F, F749V 9 K508L, R737W, Y672R 10 E507H, H785L 11 A518Q, E734M 12 F495R, F749T 13 K508L, F749T, E734F 14 R737P, S624K 15 T757W, V453G, E507M 16 F749E, H785G, F495G 17 E734F, Y672P 18 T509L, H785K 19 E734G, T757S, L764Q 20 K508L, V453A, A518Q

(72) The 20 mutants with better amplification activity were sequenced, and their amino acid mutations were counted as shown in the above table. V453, F495, E507, K508, T509, A518, S624, Y672, E734, R737, F749, T757, L764, and H785 were repeated at high frequency in 20 mutants, proving that these mutations had a significant effect on the amplification activity of Taq enzyme.

Example 5: Comparison of Mutated Taq Enzyme and Wild-Type Taq Enzyme

(73) The Taq enzyme mutants were taken, and after expression and purification, the following amplification test was performed with wild-type Taq enzyme:

(74) TABLE-US-00017 pET28a vector 100 pg Taq enzyme mutant/Wild-type Taq enzyme 10 ng 10XTaq enzyme reaction solution (100 mM 2 ul Tris, 500 mM KCl, 100 mM

(NH.sub.4).sub.2SO.sub.4, pH 8.0) pET28_F primer 4 pmol pET28_R primer 4 pmol dNTPs (2.5 mM) 2 uL ddH.sub.2O added to 20 ul PCR program: 95° C. for 5 minutes, n cycles (95° C. for 15 seconds, 55° C. for 15 seconds, 72° C. for 10 seconds), 4° C. ∞

(75) The above reaction solutions were prepared, PCR amplification was performed for 15, 20, 25, 30 cycles, precipitated with ethanol to purify the PCR products. The light absorption of the products at 260 nm was measured, and the total amount of PCR products (ng) corresponding to each cycle number was thereby calculated. The results were as follows:

(76) TABLE-US-00018 TABLE 2 Number of PCR amplification cycles Taq enzyme 15 20 25 30 Taq enzyme 104.35 604.25 1124.31 1643.24 mutant 1 Taq enzyme 61.89 412.58 701.39 981.95 mutant 2 Taq enzyme 74.07 493.80 839.46 1175.25 mutant 3 Taq enzyme 80.41 536.04 911.26 1275.76 mutant 4 Taq enzyme 65.30 435.33 740.05 1036.08 mutant 5 Taq enzyme 83.82 558.78 949.92 1329.89 mutant 6 Taq enzyme 59.45 396.34 673.78 943.29 mutant 7 Taq enzyme 55.55 370.35 629.60 881.44 mutant 8 Taq enzyme 66.27 441.82 751.10 1051.54 mutant 9 Taq enzyme 76.51 510.05 867.08 1213.91 mutant 10 Taq enzyme 72.61 484.06 822.90 1152.05 mutant 11 Taq enzyme 84.79 565.27 960.97 1345.35 mutant 12 Taq enzyme 78.94 526.29 894.69 1252.57 mutant 13 Taq enzyme 71.63 477.56 811.85 1136.59 mutant 14 Taq enzyme 62.86 419.08 712.44 997.42 mutant 15 Taq enzyme 63.35 422.33 717.96 1005.15 mutant 16 Taq enzyme 86.74 578.27 983.06 1376.28 mutant 17 Taq enzyme 53.60 357.36 607.51 850.51 mutant 18 Taq enzyme 68.22 454.82 773.19 1082.47 mutant 19 Taq enzyme 85.28 568.52 966.49 1353.08 mutant 20 Wild-type Taq 52.43 324.87 527.64 649.21 enzyme

(77) From the above results, it can be seen that the amount of products amplified by Taq enzyme mutants 1 to 20 at the same number of PCR cycles is significantly higher than that of wild-type Taq enzyme. Wherein, the amount of product obtained when mutant 1 was amplified for 20 cycles was already equivalent to that obtained by wild-type Taq enzyme for 30 cycles of amplification; under the same 30 amplification cycles, the amount of product obtained by mutant 1 is 2.5 times more than that of wild-type Taq enzyme.

Example 6: Use of Mutated Taq Enzyme in a Novel Coronavirus SARS-CoV-2 Fluorescent

Quantitative PCR Detection Kit

(78) The Taq enzyme mutants 1 #, 6 #, and 17 # were selected, and the reaction system was prepared according to the table below

(79) TABLE-US-00019 NC (ORF1ab/N) PCR reaction solution A 17 ul Taq enzyme mutant/wild-type Taq enzyme 10 ng MMLV reverse transcriptase 200 U RNase Inhibitor 20 U NC(ORF1ab/N) positive quality control nucleic acid extract 5 ul

(80) Wherein NC (ORF1ab/N) PCR reaction solution A, nucleic acid extract of NC (ORF 1 ab/N) positive quality control were both provided from 2019 Novel Coronavirus (2019-nCoV) ORFlab N nucleic acid detection kit (PCR-fluorescent probe method) (Sun Yat-sen University, Daan Gene Co., Ltd.). MMLV reverse transcriptase and RNase Inhibitor are both prepared by Sun Yat-sen University, Daan Gene Co., Ltd. PCR program was set as follows:

(81) TABLE-US-00020 Number Temperature Running Fluorescent data Step of cycles (° C.) time acquisition 1 1 50 00:15:00 2 1 95 00:15:00 3 45 94 00:00:15 55 00:00:45 ✓

(82) The ct values of NC (ORFlab/N) positive quality control with different concentration gradients amplified by each Taq enzyme mutant and wild-type Taq enzyme were as follows:

(83) TABLE-US-00021 Concentration of NC(ORF1ab/N) positive quality control (copies/ml) 10.sup.6 10.sup.5 10.sup.4 10.sup.3 Taq enzyme 22.5 25.16 28.24 31.54 mutant 1 Taq enzyme 23.14 26.41 29.55 33.01 mutant 6 Taq enzyme 24.09 27.65 30.97 34.52 mutant 17 Wild-type Taq 25.61 28.97 31.26 34.97 enzyme

(84) From the above results, it can be seen that the performance of Taq enzyme mutants on SARS-CoV-2 fluorescent quantitative PCR detection kit is significantly improved compared to the wild-type Taq enzyme.

(85) All documents mentioned in the present invention are cited as references in this application, just as each document is individually cited as a reference. In addition, it should be understood that, after reading the above content of the present invention, those skilled in the art can make various changes or modifications to the present invention, and these equivalent forms also fall within the scope defined by the appended claims of the present application.

Claims

1. A mutated DNA polymerase, wherein the mutated DNA polymerase is mutated based on the wild-type DNA polymerase as set forth in SEQ ID NO: 2, wherein the amino acid sequence of the mutated DNA polymerase has at least 98% homology to SEQ ID NO: 2, and the mutated DNA polymerase comprises the mutations: F495R and F749T (Mutant No. 12).
 2. A method for preparing the mutated DNA polymerase of claim 1, wherein the method comprises the steps of: (i) culturing a host cell under suitable conditions to express the mutant DNA polymerase; and (ii) isolating the mutated DNA polymerase, wherein the host cell contains a chromosome integrated with the nucleic acid molecule encoding the mutated DNA polymerase of claim 1.
 3. A kit, wherein the kit comprises the mutated DNA polymerase of claim 1.
-