



US012383555B2

(12) **United States Patent**
Centore et al.

(10) **Patent No.:** **US 12,383,555 B2**

(45) **Date of Patent:** **Aug. 12, 2025**

(54) **METHODS OF TREATING CANCERS**

(71) Applicant: **Foghorn Therapeutics Inc.**,
Cambridge, MA (US)

(72) Inventors: **Richard C. Centore**, Wakefield, MA
(US); **Lan Xu**, Wellesley, MA (US);
David L. Lahr, Watertown, MA (US);
Ammar Adam, Cambridge, MA (US)

(73) Assignee: **FOGHORN THERAPEUTICS INC.**,
Cambridge, MA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 496 days.

(21) Appl. No.: **17/325,937**

(22) Filed: **May 20, 2021**

(65) **Prior Publication Data**

US 2022/0079940 A1 Mar. 17, 2022

Related U.S. Application Data

(60) Provisional application No. 63/027,759, filed on May
20, 2020.

(51) **Int. Cl.**

A61K 31/501 (2006.01)

A61K 35/02 (2015.01)

A61K 35/04 (2006.01)

A61K 39/00 (2006.01)

A61K 45/06 (2006.01)

A61P 35/02 (2006.01)

A61P 35/04 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/501** (2013.01); **A61K 39/00**
(2013.01); **A61K 45/06** (2013.01); **A61P 35/02**
(2018.01); **A61P 35/04** (2018.01)

(58) **Field of Classification Search**

CPC **A61P 35/02**; **A61P 35/04**
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,788,341 A	4/1957	Schwyzner et al.
3,717,642 A	2/1973	Von Strandtmann
4,109,496 A	8/1978	Allemann et al.
4,650,796 A	3/1987	George et al.
4,868,103 A	9/1989	Stavrianopoulos et al.
5,223,409 A	6/1993	Ladner et al.
5,225,422 A	7/1993	Nagata et al.
5,283,317 A	2/1994	Saifer et al.
5,631,169 A	5/1997	Lakowicz et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN	103038231 A	4/2013
CN	105473141 A	4/2016

(Continued)

OTHER PUBLICATIONS

Advani, A Phase 1 study of imatinib mesylate in combination with
cytarabine and daunorubicin for c-kit positive relapsed acute myeloid
leukemia, *Leukemia Research*, 2010, 34, pp. 1622-1626 (Year:
2010).*

(Continued)

Primary Examiner — Renee Claytor

Assistant Examiner — Andrew P Lee


(74) *Attorney, Agent, or Firm* — Clark & Elbing LLP

(57) **ABSTRACT**

The present invention relates to methods and compositions
for the treatment of BAF-related disorders such as acute
myeloid leukemia.

13 Claims, 6 Drawing Sheets

Specification includes a Sequence Listing.



(56)

References Cited**U.S. PATENT DOCUMENTS**

5,677,158 A 10/1997 Zhou et al.
 5,801,030 A 9/1998 McVey et al.
 5,858,358 A 1/1999 June et al.
 5,883,223 A 3/1999 Gray
 6,180,612 B1 1/2001 Hockensmith et al.
 6,309,634 B1 10/2001 Bankiewicz et al.
 6,352,694 B1 3/2002 June et al.
 6,551,786 B2 4/2003 Manfredi
 6,683,058 B1 1/2004 Tuszynski
 6,692,964 B1 2/2004 June et al.
 6,716,662 B2 4/2004 Akai
 6,797,514 B2 9/2004 Berenson et al.
 6,867,041 B2 3/2005 Berenson et al.
 6,887,466 B2 5/2005 June et al.
 6,905,680 B2 6/2005 June et al.
 6,905,681 B1 6/2005 June et al.
 6,905,874 B2 6/2005 Berenson et al.
 6,995,011 B2 2/2006 Itoh et al.
 7,067,318 B2 6/2006 June et al.
 7,144,575 B2 12/2006 June et al.
 7,172,869 B2 2/2007 June et al.
 7,175,843 B2 2/2007 June et al.
 7,205,103 B2 4/2007 Emerson
 7,232,566 B2 6/2007 June et al.
 7,572,631 B2 8/2009 Berenson et al.
 8,324,367 B2 12/2012 Kaemmerer et al.
 8,642,660 B2 2/2014 Goldfarb
 8,697,359 B1 4/2014 Zhang
 8,703,761 B2 4/2014 Forster et al.
 8,771,945 B1 7/2014 Zhang
 8,795,965 B2 8/2014 Zhang
 8,865,406 B2 10/2014 Zhang et al.
 8,946,268 B2 2/2015 Lau et al.
 9,072,052 B2 6/2015 Griffin et al.
 9,126,985 B2 9/2015 Kley et al.
 9,353,051 B2 5/2016 Byrd et al.
 9,410,943 B2 8/2016 Kadoch et al.
 9,546,206 B2 1/2017 Ring et al.
 9,546,296 B2 1/2017 Wang et al.
 9,636,323 B2 5/2017 Lin et al.
 9,656,959 B2 5/2017 Ni et al.
 9,694,084 B2 7/2017 Bradner et al.
 9,708,338 B2 7/2017 Yukimasa et al.
 9,708,348 B2 7/2017 Castro et al.
 9,932,340 B2 4/2018 Dai et al.
 10,105,420 B2 10/2018 Kadoch et al.
 10,131,637 B2 11/2018 Abdel-Meguid et al.
 10,207,998 B2 2/2019 Derbyshire et al.
 10,239,888 B2 3/2019 Bradner et al.
 10,266,850 B2 4/2019 Doudna et al.
 10,464,925 B2 11/2019 Bradner et al.
 10,472,376 B2 11/2019 Yamamoto et al.
 10,646,575 B2 5/2020 Phillips et al.
 10,660,968 B2 5/2020 Phillips et al.
 10,669,253 B2 6/2020 Bradner et al.
 10,849,982 B2 12/2020 Phillips et al.
 10,905,768 B1 2/2021 Phillips et al.
 10,976,320 B2 4/2021 Dykhuizen et al.
 11,419,859 B2 8/2022 Agresta
 11,485,732 B2 11/2022 Vaswani et al.
 11,497,752 B2 11/2022 Anthony et al.
 11,639,345 B2 5/2023 Bloch et al.
 11,773,085 B2 10/2023 Zhou et al.
 11,793,802 B2 10/2023 Bearss et al.
 11,851,445 B2 12/2023 Ruppel et al.
 11,865,114 B2 1/2024 Ramachandra et al.
 2002/0022018 A1 2/2002 Curiel et al.
 2002/0037281 A1 3/2002 Davidson et al.
 2002/0106632 A1 8/2002 Manfredi
 2003/0022375 A1 1/2003 Itoh et al.
 2003/0027335 A1 2/2003 Ruley et al.
 2004/0216178 A1 10/2004 Jones et al.
 2005/0079512 A1 4/2005 Emerson et al.
 2005/0130919 A1 6/2005 Xu et al.
 2006/0058255 A1 3/2006 Chen et al.

2006/0121005 A1 6/2006 Berenson et al.
 2007/0105181 A1 5/2007 Pope et al.
 2008/0221157 A1 9/2008 Chakravarty et al.
 2009/0163545 A1 6/2009 Goldfarb
 2010/0048565 A1 2/2010 Frenkel et al.
 2010/0197621 A1 8/2010 Henry et al.
 2010/0284990 A1 11/2010 Kaemmerer et al.
 2011/0230486 A1 9/2011 Lau et al.
 2012/0034867 A1 2/2012 Griffin et al.
 2012/0035244 A1 2/2012 Chinnaiyan et al.
 2012/0308484 A1 12/2012 Szalay et al.
 2013/0034867 A1 2/2013 Bomgarden et al.
 2014/0068797 A1 3/2014 Doudna et al.
 2014/0287931 A1 9/2014 Bernards et al.
 2015/0018315 A1 1/2015 Kley et al.
 2015/0057169 A1 2/2015 Siu et al.
 2015/0376139 A1 12/2015 Abdel-Meguid et al.
 2016/0032402 A1 2/2016 Jagani et al.
 2016/0130663 A1 5/2016 Kohno et al.
 2016/0176916 A1 6/2016 Bradner et al.
 2016/0200721 A1 7/2016 Yukimasa et al.
 2016/0347708 A1 12/2016 Ebright et al.
 2017/0174688 A1 6/2017 Dai et al.
 2018/0086720 A1 3/2018 Albrecht et al.
 2018/0105500 A1 4/2018 Derbyshire et al.
 2018/0140722 A1 5/2018 Willis et al.
 2018/0258491 A1 9/2018 Jagani et al.
 2018/0328913 A1 11/2018 Kadoch et al.
 2020/0069669 A1 3/2020 Grassian et al.
 2020/0206344 A1 7/2020 Kadoch et al.
 2020/0261434 A1 8/2020 Choe et al.
 2021/0009568 A1 1/2021 Zhou et al.
 2021/0038611 A1 2/2021 Anthony et al.
 2021/0171958 A1 6/2021 Chan et al.
 2021/0230154 A1 7/2021 Vaswani et al.
 2021/0230190 A1 7/2021 Ruppel et al.
 2021/0251988 A1 8/2021 Zhou et al.
 2021/0260171 A1 8/2021 Zhou et al.
 2021/0388040 A1 12/2021 Kadoch et al.
 2022/0016083 A1 1/2022 Centore et al.
 2022/0079940 A1 3/2022 Centore et al.
 2022/0098190 A1 3/2022 Ruppel et al.
 2022/0119378 A1 4/2022 Anthony et al.
 2022/0396604 A1 12/2022 Kadoch et al.
 2023/0035235 A1 2/2023 Kadoch et al.
 2023/0079819 A1 3/2023 Vaswani et al.
 2023/0121497 A1 4/2023 Vaswani et al.
 2023/0129003 A1 4/2023 Vaswani et al.
 2023/0138480 A1 5/2023 Anthony et al.
 2023/0145003 A1 5/2023 Wilson et al.
 2023/0149414 A1 5/2023 Anthony et al.
 2024/0101550 A1 3/2024 Vaswani et al.
 2024/0158387 A1 5/2024 Vaswani et al.
 2024/0189318 A1 6/2024 Huang

FOREIGN PATENT DOCUMENTS

CN 104530013 B 6/2016
 CN 107531668 A 1/2018
 EA 202192101 A1 12/2021
 JP 2008-505963 A 2/2008
 JP 2011-507910 A 3/2011
 JP 2011-528016 A 11/2011
 JP 2016-520515 A 7/2016
 WO WO-94/10300 A1 5/1994
 WO WO-95/30761 A2 11/1995
 WO WO-2000/024392 A1 5/2000
 WO WO-00/59888 A1 10/2000
 WO WO-00/59905 A1 10/2000
 WO WO-2005/039643 A2 5/2005
 WO WO-2005/112620 A2 12/2005
 WO WO-2006/005941 A1 1/2006
 WO WO-2006/051063 A1 5/2006
 WO WO-2008/022396 A1 2/2008
 WO WO-2009/086303 A2 7/2009
 WO WO-2010/007046 A2 1/2010
 WO WO-2011/115998 A2 9/2011
 WO WO-2012/085650 A1 6/2012
 WO WO-2013/116663 A1 8/2013

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO-2013/116682	A1	8/2013
WO	WO-2014/150395	A1	9/2014
WO	WO-2015/005473	A1	1/2015
WO	WO-2015/002230	A1	1/2015
WO	WO-2015/103317	A1	7/2015
WO	WO-2015/120320	A1	8/2015
WO	WO-2015/121688	A1	8/2015
WO	WO-2016/054491	A1	4/2016
WO	WO-2016/138114	A1	9/2016
WO	WO-2016/160718	A1	10/2016
WO	WO-2016/207212	A1	12/2016
WO	WO-2017/024318	A1	2/2017
WO	WO-2017/060470	A1	4/2017
WO	WO-2017/087885	A1	5/2017
WO	WO-2017/118734	A1	7/2017
WO	WO-2017/158381	A1	9/2017
WO	WO-2018/148443	A1	8/2018
WO	WO-2018/160636	A1	9/2018
WO	WO-2018/175324	A1	9/2018
WO	WO-2019/040098	A1	2/2019
WO	WO-2019/138017	A1	7/2019
WO	WO-2019/142192	A1	7/2019
WO	WO-2019/152437	A1	8/2019
WO	WO-2019/152440	A1	8/2019
WO	WO-2019/226915	A1	11/2019
WO	WO-2020/035779	A1	2/2020
WO	WO-2020/081556	A2	4/2020
WO	WO-2020/081588	A1	4/2020
WO	WO-2020/106915	A1	5/2020
WO	WO-2020/127685	A1	6/2020
WO	WO-2020/160100	A1	8/2020
WO	WO-2020/160180	A1	8/2020
WO	WO-2021/081032	A1	4/2021
WO	WO-2021/155262	A1	8/2021
WO	WO-2021/155264	A1	8/2021
WO	WO-2021/155316	A1	8/2021
WO	WO-2021/155320	A1	8/2021
WO	WO-2021/155321	A2	8/2021
WO	WO-2021/183218	A1	9/2021
WO	WO-2021/236080	A1	11/2021
WO	WO-2022/192621	A1	9/2022
WO	WO-2022/198043	A1	9/2022
WO	WO-2023/009834	A2	2/2023
WO	WO-2023/196560	A1	10/2023
WO	WO-2023/196565	A1	10/2023
WO	WO-2023/196567	A2	10/2023
WO	WO-2024/024428	A1	2/2024
WO	WO-2024/031875	A1	2/2024
WO	WO-2024/086577	A1	4/2024

OTHER PUBLICATIONS

Godwin, Gemtuzumab ozogamicin in acute myeloid leukemia, *Leukemia*, 2017, 31, pp. 1855-1868 (Year: 2017).*

Danziger, Automated site-directed drug design: a general algorithm for knowledge acquisition about hydrogen-bonding regions at protein surfaces, 1989, *Proc. R. Soc. Lond.*, 236, pp. 101-113 (Year: 1989).*

Simone, *Oncology: Introduction*, Textbook of Medicine, 1997, 20(1), pp. 1004-1010 (Year: 1997).*

Adamo et al., "The oncogene ERG: a key factor in prostate cancer," *Oncogene* 35(4):403-14 (Jan. 28, 2016).

Anders et al., "HTSeq—a Python framework to work with high-throughput sequencing data," *Bioinformatics* 31(2):166-9 (Jan. 15, 2015).

Asangani et al., "Therapeutic targeting of BET bromodomain proteins in castration-resistant prostate cancer," available in PMC Dec. 12, 2014. Published in final edited form as: *Nature*. 510(7504):278-82 (2014) (44 pages).

Attard et al., "Duplication of the fusion of TMPRSS2 to ERG sequences identifies fatal human prostate cancer," available in PMC Feb. 24, 2009. Published in final edited form as: *Oncogene*. 27(3):253-63 (2008) (19 pages).

Basuyaux et al., "The Ets transcription factors interact with each other and with the c-Fos/c-Jun complex via distinct protein domains in a DNA-dependent and-independent manner," *J Biol Chem*. 272(42):26188-95 (1997).

Bendall et al., "Prevention of amino acid conversion in SILAC experiments with embryonic stem cells," *Mol Cell Proteomics*. 7(9):1587-97 (2008).

Berger et al., "Androgen-induced differentiation and tumorigenicity of human prostate epithelial cells," *Cancer Res*. 64(24):8867-75 (2004).

Börnø et al., "Genome-wide DNA methylation events in TMPRSS2-ERG fusion-negative prostate cancers implicate an EZH2-dependent mechanism with miR-26a hypermethylation," *Cancer Discov*. 2(11):1024-35 (2012).

Camuzeaux et al., "Imaging Erg and Jun transcription factor interaction in living cells using fluorescence resonance energy transfer analyses," *Biochem Biophys Res Commun*. 332(4):1107-14 (2005).
Cancer Genome Atlas Research Network, "The Molecular Taxonomy of Primary Prostate Cancer," *Cell*. 163(4):1011-25 (2015) (16 pages).

Carell et al., "A novel procedure for the synthesis of libraries containing small organic molecules," *Angew Chem Int Ed Engl*. 33(20):2059-2061 (1994).

Carell et al., "A Solution-Phase Screening Procedure for the Isolation of Active Compounds from a Library of Molecules," *Angewandte Chemie International Edition in English*. 33(20): 2061-2064 (1994).

Chen et al., "ETS factors reprogram the androgen receptor cistrome and prime prostate tumorigenesis in response to PTEN loss," available in PMC Feb. 1, 2014. Published in final edited form as: *Nat Med*. 19(8):1023-9 (2013) (21 pages).

Chng et al., "A transcriptional repressor co-regulatory network governing androgen response in prostate cancers," *EMBO J*. 31(12):2810-23 (2012).

Cho et al., "An unnatural biopolymer," *Science*. 261(5126):1303-1305 (1993).

Cull et al., "Screening for receptor ligands using large libraries of peptides linked to the C terminus of the lac repressor," *Proc Natl Acad Sci*. 89(5):1865-1869. (1992).

Cwirla et al., "Peptides on phage: A vast library of peptides for identifying ligands," *Proc Natl Acad Sci*. 87:6378-6382 (1990).

Delattre et al., "Gene fusion with an ETS DNA-binding domain caused by chromosome translocation in human tumours," *Nature*. 359(6391):162-5 (1992).

Demicheli et al., "TMPRSS2:ERG gene fusion associated with lethal prostate cancer in a watchful waiting cohort," *Oncogene* 26:4596-4599 (2007).

Devlin et al., "Random Peptide Libraries: a Source of Specific Protein Binding Molecules," *Science*. 249(4967):404-406 (1990).

DeWitt et al., "Diversomers": an approach to nonpeptide, nonoligomeric chemical diversity," *Proc Natl Acad Sci*. 90(15):6909-6913 (1993).

Dobin et al., "STAR: ultrafast universal RNA-seq aligner," *Bioinformatics*. 29(1):15-21 (2013).

Donaldson et al., "Solution structure of the ETS domain from murine Ets-1: a winged helix-turn-helix DNA binding motif," *EMBO J*. 15(1):125-34 (1996).

Erb et al., "Recursive deconvolution of combinatorial chemical libraries," *Proc Natl Acad Sci*. 91(24):11422-11426 (1994).

Extended European Search Report for European Application No. 19748410.8, dated Sep. 24, 2021 (10 pages).

Extended European Search Report for European Application No. 19887386.1, dated Dec. 5, 2022 (18 pages).

Extended European Search Report for European Application No. 20749261.2, dated Oct. 18, 2022 (8 pages).

Felici et al., "Selection of antibody ligands from a large library of oligopeptides expressed on a multivalent exposition vector," *J Mol Biol*. 222(2):301-10 (1991).

Feng et al., "GFOLD: a generalized fold change for ranking differentially expressed genes from RNA-seq data," *Bioinformatics*. 28(21):2782-8 (2012).

Fodor et al., "Multiplexed biochemical assays with biological chips," *Nature*. 364:555-556 (1993).

(56)

References Cited**OTHER PUBLICATIONS**

Gallop et al., "Applications of Combinatorial Technologies to Drug Discovery. 1. Background and Peptide Combinatorial Libraries," *J Med Chem* 37(9):1233-51 (1994).

Gene Ontology Consortium, "Gene Ontology Consortium: going forward," *Nucleic Acids Res.* 43(Database issue): D1049-56 (2015).

Helgeson et al., "Characterization of TMPRSS2:ETV5 and SLCA4A3:ETV5 gene fusions in prostate cancer," *Cancer Res.* 68(1):73-80 (2008).

Ho et al., "An embryonic stem cell chromatin remodeling complex, esBAF, is essential for embryonic stem cell self-renewal and pluripotency," *Proc Natl Acad Sci U S A.* 106(13):5181-6 (2009).

Houghten et al., "The use of synthetic peptide combinatorial libraries for the identification of bioactive peptides," *Biotechniques.* 13(3):412-21 (1992).

Ichikawa et al., "An RNA-binding protein gene, TLS/FUS, is fused to ERG in human myeloid leukemia with t(16;21) chromosomal translocation," *Cancer Res.* 54(11):2865-8 (1994).

International Preliminary Report on Patentability for International Application No. PCT/US2016/062911 issued May 22, 2018 (13 pages).

International Preliminary Report on Patentability for International Application No. PCT/US2019/015722, issued Aug. 4, 2020 (6 pages).

International Preliminary Report on Patentability for International Application No. PCT/US2020/033829, issued Nov. 17, 2022 (6 pages).

International Search Report and Written Opinion for International Application No. PCT/US19/15722, mailed May 31, 2019 (12 pages).

International Search Report and Written Opinion for International Application No. PCT/US2016/062911 dated Mar. 3, 2017 (16 pages).

International Search Report and Written Opinion for International Application No. PCT/US2020/015605, mailed Jun. 16, 2020 (15 pages).

International Search Report and Written Opinion for International Application No. PCT/US2020/015723, mailed Jul. 2, 2020 (15 pages).

International Search Report and Written Opinion for International Application No. PCT/US2020/033829, mailed Aug. 17, 2020 (14 pages).

International Search Report and Written Opinion for International Application No. PCT/US2022/019506, dated Jun. 7, 2022 (9 pages).

International Search Report and Written Opinion for International Application No. PCT/US21/15876, mailed on Apr. 7, 2021 (23 pages).

International Search Report and Written Opinion for International Application No. PCT/US21/15878, dated Jun. 4, 2021 (10 pages).

International Search Report and Written Opinion for International Patent Application No. PCT/US2019/062525, mailed Feb. 18, 2020 (14 pages).

Kadoch et al., "Mammalian SWI/SNF chromatin remodeling complexes and cancer: Mechanistic insights gained from human genomics," *Sci Adv.* 1(5):e1500447 (2015) (17 pages).

Kadoch et al., "Proteomic and bioinformatic analysis of mammalian SWI/SNF complexes identifies extensive roles in human malignancy," *Nat Genet.* 45(6):592-601 (2013) (11 pages).

Kadoch et al., "Reversible Disruption of mSWI/SNF (BAF) Complexes by the SS18-SSX Oncogenic Fusion in Synovial Sarcoma," available in PMC May 16, 2013, published in final edited form as: *Cell.* 153(1):71-85 (2013) (26 pages).

Karim et al., "The ETS-domain: a new DNA-binding motif that recognizes a purine-rich core DNA sequence," *Genes Dev.* 4(9):1451-3 (1990).

Klezovitch et al., "A causal role for ERG in neoplastic transformation of prostate epithelium," *Proc Natl Acad Sci U S A.* 105(6):2105-10 (2008).

Kumar-Sinha et al., "Recurrent gene fusions in prostate cancer," available in PMC Jul. 16, 2009. Published in final edited form as: *Nat Rev Cancer.* 8(7):497-511 (2008) (29 pages).

Kunderfranco et al., "ETS transcription factors control transcription of EZH2 and epigenetic silencing of the tumor suppressor gene Nkx3.1 in prostate cancer," *PLoS One.* 5(5):e10547 (2010) (17 pages).

Lam et al., "A new type of synthetic peptide library for identifying ligand-binding activity," *Nature.* 354(6348):82-84 (1991).

Lam, "Application of combinatorial library methods in cancer research and drug discovery," *Anticancer Drug Des.* 12(3):145-67 (1997).

Langmead et al., "Fast gapped-read alignment with Bowtie 2," available in PMC Apr. 1, 2013. Published in final edited form as: *Nat Methods.* 9(4):357-9 (2012) (8 pages).

Link et al., "Targeting the BAF57 SWI/SNF subunit in prostate cancer: a novel platform to control androgen receptor activity," *Cancer Res.* 68(12):4551-8 (2008).

Love et al., "Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2," *Genome Biol.* 15(12):550 (2014) (21 pages).

Lupien et al., "FoxA1 translates epigenetic signatures into enhancer-driven lineage-specific transcription," *Cell.* 132(6):958-70 (2008).

Machanic et al., "MEME-ChIP: motif analysis of large DNA datasets," *Bioinformatics.* 27(12):1696-7 (2011).

Mackereth et al., "Diversity in structure and function of the Ets family PNT domains," *J Mol Biol.* 342(4):1249-64 (2004).

Madura et al., "N-recognition/Ubc2 interactions in the N-end rule pathway," *J Biol Chem.* 268(16):12046-54 (1993).

McBride et al., "Disruption of mammalian SWI/SNF and polycomb complexes in human sarcomas: mechanisms and therapeutic opportunities," *J Pathol.* 244(5): 638-649 (Apr. 2018).

Mel  t et al., "The human transcriptome across tissues and individuals," available in PMC Aug. 24, 2015. Published in final edited form as: *Science.* 348(6235):660-5 (2015) (12 pages).

Mounir et al., "ERG signaling in prostate cancer is driven through PRMT5-dependent methylation of the Androgen Receptor," *Elife.* 5:e13964 (2016) (19 pages).

Nagaich et al., "Rapid periodic binding and displacement of the glucocorticoid receptor during chromatin remodeling," *Mol Cell.* 14(2):163-74 (2004).

Nam et al., "Expression of the TMPRSS2:ERG fusion gene predicts cancer recurrence after surgery for localised prostate cancer," *Br J Cancer.* 97(12):1690-5 (2007).

Office Action for Chinese Patent Application No. 201980023925.9, dated Apr. 20, 2022 (13 pages).

Ong et al., "A practical recipe for stable isotope labeling by amino acids in cell culture (SILAC)," *Nat Protoc.* 1(6):2650-60 (2006).

Papillon et al., "Discovery of Orally Active Inhibitors of Brahma Homolog (BRM)/SMARCA2 ATPase Activity for the Treatment of Brahma Related Gene 1 (BRG1)/SMARCA4-Mutant Cancers," *J Med Chem.* 61(22):10155-72 (Nov. 2018).

Partial Supplementary European Search Report for European Patent Application No. 19887386.1, dated Jul. 20, 2022 (23 pages).

Paulo et al., "FLI1 is a novel ETS transcription factor involved in gene fusions in prostate cancer," *Genes Chromosomes Cancer.* 51(3):240-9 (2012).

Petrovics et al., "Frequent overexpression of ETS-related gene-1 (ERG1) in prostate cancer transcriptome," *Oncogene.* 24(23):3847-52 (2005).

Pomerantz et al., "The androgen receptor cistrome is extensively reprogrammed in human prostate tumorigenesis," available in PMC May 1, 2016. Published in final edited form as: *Nat Genet.* 47(11):1346-51 (2015) (17 pages).

Prensner et al., "The long noncoding RNA SChLAP1 promotes aggressive prostate cancer and antagonizes the SWI/SNF complex," available in PMC May 1, 2014. Published in final edited form as: *Nat Genet.* 45(11):1392-8 (2013) (26 pages).

PubChem CID 117640569, "N-[2-[[4-(4-Methoxyphenyl)-1,3-thiazol-2-yl]amino]-2-oxoethyl]-1,3-thiazole-5-carboxamide," <https://pubchem.ncbi.nlm.nih.gov/compound/117640569>, created Feb. 23, 2016 (9 pages).

PubChem CID 56442706, "1-(4-Methoxyphenyl)-N-[2-oxo-2-[4-(1,2,4-triazol-1-yl) anilino]ethyl]pyrazole-3-carboxamide," <https://pubchem.ncbi.nlm.nih.gov/compound/56442706>, created Jan. 25, 2012 (8 pages).

(56)

References Cited

OTHER PUBLICATIONS

- PubChem CID 91946137, "N-[2-[(1-Ethylpyrazol-3-yl)amino]-2-oxoethyl]-1-methylpyrazole-3-carboxamide," <https://pubchem.ncbi.nlm.nih.gov/compound/91946137>, created Oct. 22, 2015 (8 pages).
- PubChem Compound Summary for CID 155037309, dated Dec. 19, 2020 (9 pages).
- PubChem Compound Summary for CID No. 136572628, "4-Chloro-N-[2-(cyclopentylamino)-2-oxoethyl]-5-nitro-1H-pyrazole-3-carboxamide," created Jan. 24, 2019, <<https://pubchem.ncbi.nlm.nih.gov/compound/136572628>>, (7 pages).
- PubChem Compound Summary for CID No. 49726797, "N-Methyl-N-(2-oxo-2-[(4-pyridin-2-yl)thiazol-2-yl]aminoethyl)-1H-indole-3-carboxamide," created Nov. 27, 2010, <<https://pubchem.ncbi.nlm.nih.gov/compound/49726797>>, (8 pages).
- PubChem Compound Summary for CID No. 91945707, "N-[2-[(4,5-Dimethyl-1,3-thiazol-2-yl)amino]-2-oxoethyl]-1-methylpyrazole-3-carboxamide," created Oct. 22, 2015 <<https://pubchem.ncbi.nlm.nih.gov/compound/91945707>>, (8 pages).
- PubChem Compound Summary for PubChem CID 49726798, "N-(2-[(4-Furan-2-yl)thiazol-2-yl]amino)-2-oxoethyl-N-methyl-1H-indole-3-carboxamide," created Nov. 27, 2010 <<https://pubchem.ncbi.nlm.nih.gov/compound/49726798>> (8 pages).
- PubChem Compound Summary for SID 172131678, dated Dec. 9, 2014 (8 pages).
- Pubchem, "Compound Summary for CID 108452511," <<https://pubchem.ncbi.nlm.nih.gov/compound/108452511>>, created Jan. 15, 2016, retrieved Jan. 4, 2021 (7 pages).
- Pubchem, "Compound Summary for CID 2955118," <<https://pubchem.ncbi.nlm.nih.gov/compound/2955118>>, created Jul. 29, 2005, retrieved Mar. 22, 2017 (13 pages).
- Pubchem, "Compound Summary for CID 7325930," <<https://pubchem.ncbi.nlm.nih.gov/compound/7325930>>, created Jul. 29, 2006, retrieved Mar. 22, 2017 (11 pages).
- Pubchem, "Compound Summary for CID 970466," <<https://pubchem.ncbi.nlm.nih.gov/compound/970466>>, created Jul. 9, 2005, retrieved Mar. 22, 2017 (11 pages).
- Quinlan et al., "BEDTools: a flexible suite of utilities for comparing genomic features," *Bioinformatics*. 26(6):841-2 (2010).
- Rajput et al., "Frequency of the TMPRSS2:ERG gene fusion is increased in moderate to poorly differentiated prostate cancers," *J Clin Pathol*. 60(11):1238-43 (2007).
- Rappsilber et al., "Protocol for micro-purification, enrichment, pre-fractionation and storage of peptides for proteomics using StageTips," *Nat Protoc*. 2(8):1896-906 (2007).
- Scott et al., "Searching for peptide ligands with an epitope library," *Science*. 249(4967):386-390 (1990).
- Shen et al., "The SWI/SNF ATPase Brm is a gatekeeper of proliferative control in prostate cancer," *Cancer Res*. 68(24):10154-62 (Dec. 2008).
- Shi et al., "Role of SWI/SNF in acute leukemia maintenance and enhancer-mediated Myc regulation," *Genes Dev*. 27(24):2648-62 (Dec. 2013).
- Siegel et al., "Cancer statistics, 2015," *CA Cancer J Clin*. 65(1):5-29 (2015).
- Subramanian et al., "Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles," *Proc Natl Acad Sci USA*. 102(43):15545-50 (2005).
- Sun et al., "TMPRSS2-ERG fusion, a common genomic alteration in prostate cancer activates C-MYC and abrogates prostate epithelial differentiation," available in PMC Oct. 4, 2020. Published in final edited form as: *Oncogene*. 27(40):5348-53 (2008) (12 pages).
- Tikdari et al., "Reaction of 2-Aminothiazoles with 5-Oxazolones," *ChemInform*. 18(47):Abstract 199 (1987) (1 page).
- Tikdari et al., "Reaction of 2-Aminothiazoles with 5-Oxazolones," *Indian Journal of Chemistry* 26B:478-9 (May 1987).
- Tomlins et al., "Recurrent fusion of TMPRSS2 and ETS transcription factor genes in prostate cancer," *Science*. 310(5748):644-8 (2005).
- Tomlins et al., "Role of the TMPRSS2-ERG gene fusion in prostate cancer," *Neoplasia*. 10(2):177-88 (2008) (21 pages).
- Tomlins et al., "TMPRSS2:ETV4 gene fusions define a third molecular subtype of prostate cancer," *Cancer Res*. 66(7):3396-400 (2006).
- Tuoc et al., "Chromatin regulation by BAF170 controls cerebral cortical size and thickness," *Dev Cell*. 25(3):256-69 (May 2013).
- Vachtenheim et al., "SWI/SNF chromatin remodeling complex is critical for the expression of microphthalmia-associated transcription factor in melanoma cells," *Biochemical and Biophysical Research Communications*. 392(3):454-459 (2010).
- Varambally et al., "The polycomb group protein EZH2 is involved in progression of prostate cancer," *Nature*. 419(6907):624-9 (2002).
- Vela et al., "Discovery of Enhancers of the Secretion of Leukemia Inhibitory Factor for the Treatment of Multiple Sclerosis," *J Biomol Screen*. 21(5):437-45 (Jun. 2016).
- Verger et al., "Identification of amino acid residues in the ETS transcription factor Erg that mediate Erg-Jun/Fos-DNA ternary complex formation," *J Biol Chem*. 276(20):17181-9 (2001).
- Wollenick et al., "Synthetic transactivation screening reveals ETV4 as broad coactivator of hypoxia-inducible factor signaling," *Nucleic Acids Res*. 40(5):1928-43 (2012).
- Yang et al., "EZH2, an epigenetic driver of prostate cancer," *Protein Cell*. 4(5):331-41 (2013).
- Yildirim et al., "Mbd3/NURD complex regulates expression of 5-hydroxymethylcytosine marked genes in embryonic stem cells," *Cell*. 147(7):1498-510 (2011).
- Yu et al., "An integrated network of androgen receptor, polycomb, and TMPRSS2-ERG gene fusions in prostate cancer progression," *Cancer Cell*. 17(5):443-54 (2010).
- Yu et al., "Direct recruitment of polycomb repressive complex 1 to chromatin by core binding transcription factors," *Mol Cell*. 45(3):330-43 (2012).
- Zervos et al., "Mxi1, a protein that specifically interacts with Max to bind Myc-Max recognition sites," *Cell*. 72(2):223-32 (1993).
- Zhang et al., "Model-based analysis of ChIP-Seq (MACS)," *Genome Biol*. 9(9):R137 (2008) (9 pages).
- Zong et al., "ETS family transcription factors collaborate with alternative signaling pathways to induce carcinoma from adult murine prostate cells," *Proc Natl Acad Sci U S A*. 106(30):12465-70 (Jul. 2009).
- Zuckermann et al., "Discovery of Nanomolar Ligands for 7-Transmembrane G-Protein-Coupled Receptors from a Diverse N-(Substituted)glycine Peptoid Library," *J Med Chem*. 37(17):2678-2685 (1994).
- Zvarec et al., "5-Benzylidenetherhodanine and 5-benzylidene-2,4-thiazolidinedione based antibacterials," *Bioorg Med Chem Lett*. 22(8):2720-2 (2012).
- U.S. Appl. No. 18/281,022, Vaswani et al.
- U.S. Appl. No. 18/282,279, Huang, Liye.
- U.S. Appl. No. 18/373,518, Huang et al.
- Alazawi, "Foghorn Therapeutics," Blackseed Bio, last updated Mar. 4, 2022, retrieved Jul. 24, 2023, from <<https://blackseedbio.com/reports/fhtx#pipeline>> (26 pages).
- Dominguez et al. "Beyond editing: Repurposing CRISPR-Cas9 for Precision Genome Regulation and Interrogation," available in PMC Jun. 27, 2016, published in final edited form as: *Nat Rev Mol Cell Biol*. 17(1):5-15 (Jan. 2016) (24 pages).
- Fathi et al., "Differentiation syndrome with lower-intensity treatments for acute myeloid leukemia," *Am J Hematol*. 96(6):735-46 (Jun. 1, 2021) (13 Pages).
- Gaj et al. "ZFN, Talen and CRISPR/Cas-based Methods for Genome Engineering," available in PMC Jul. 1, 2014, published in final edited form as: *Trends Biotechnol*. 31(7):397-405 (Jul. 2013) (20 pages).
- International Search Report and Written Opinion for International Application No. PCT/US2023/017839, mailed Sep. 6, 2023 (11 pages).
- International Search Report and Written Opinion for International Application No. PCT/US23/17821, mailed Jun. 30, 2023 (9 pages).
- International Search Report and Written Opinion for International Application No. PCT/US23/17829, mailed Aug. 23, 2023 (15 pages).
- Kadoch et al. "Reversible Disruption of mSWI/SNF (BAF) Complexes by the SS18-SSX Oncogenic Fusion in Synovial Sarcoma,"

(56)

References Cited**OTHER PUBLICATIONS**

available in PMC May 16, 2013, published in final edited form as: Cell. 153(1):71-85 (Mar. 2013) (26 pages).

Tsai et al. "Dimeric CRISPR RNA-guided FokI Nucleases for Highly Specific Genome Editing," available in PMC Dec. 1, 2014, published in final edited form as: Nat Biotechnol. 32(6):569-576 (Jun. 2014) (22 pages).

Database Registry, RN 878254-76-3, entered Mar. 28, 2006 (1 page).

Database Registry, RN 1004932-80-2, entered Feb. 21, 2008 (1 page).

Database Registry, RN 1175782-23-6, entered Aug. 26, 2009 (1 page).

Database Registry, RN 1315743-98-6, entered Aug. 11, 2011 (1 page).

Centore et al., "Abstract 1224: Discovery of novel BAF inhibitors for the treatment of transcription factor-driven cancers," Poster Presentations—Proffered abstracts, Cancer Research 81(13_Supplement):1224 (Jul. 1, 2021) (2 pages).

Extended European Search Report for European Application No. 21748261.1, dated Jan. 29, 2024 (17 pages).

International Preliminary Report on Patentability for International Application No. PCT/US2022/019506, issued Sep. 12, 2023 (6 pages).

International Preliminary Report on Patentability for International Application No. PCT/US2022/020943, issued Sep. 12, 2023 (5 pages).

International Search Report and Written Opinion for International Application No. PCT/US2023/077088, mailed Mar. 4, 2024 (17 pages).

International Search Report and Written Opinion for PCT/US2022/020943, dated Jun. 14, 2022 (17 pages).

Mill et al., "RUNX1-targeted therapy for AML expressing somatic or germline mutation in RUNX1," Blood 134(1):59-73 (Jul. 4, 2019).

Partial Supplementary European Search Report for European Patent Application No. 20936213.6, dated Feb. 8, 2024 (20 pages).

Ramos et al., "Current Approaches in the Treatment of Relapsed and Refractory Acute Myeloid Leukemia," J Clin Med. 4(4):665-95 (Apr. 2015).

STN Registry Database, RN 1010893-05-6, entered Mar. 30, 2008 (2 pages).

STN Registry Database, RN 1049271-26-2, entered Sep. 14, 2008 (2 pages).

STN Registry Database, RN 1081662-32-9, entered Dec. 8, 2008 (2 pages).

STN Registry Database, RN 1209112-42-4, entered Mar. 12, 2010 (2 pages).

STN Registry Database, RN 1246047-75-5, entered Oct. 12, 2010 (2 pages).

STN Registry Database, RN 1308280-67-2, entered Jun. 9, 2011 (2 pages).

STN Registry Database, RN 1351682-19-3, entered Dec. 22, 2011 (2 pages).

STN Registry Database, RN 1401558-47-1, entered Oct. 22, 2012 (2 pages).

STN Registry Database, RN 1455783-72-8, entered Oct. 6, 2013 (2 pages).

STN Registry Database, RN 1576383-94-2, entered Mar. 31, 2014 (2 pages).

STN Registry Database, RN 1586193-45-4, entered Apr. 17, 2014 (2 pages).

STN Registry Database, RN 1827759-12-5, entered Dec. 13, 2015 (2 pages).

STN Registry Database, RN 1831899-24-1, entered Dec. 17, 2015 (2 pages).

STN Registry Database, RN 1839545-15-1, entered Dec. 31, 2015 (2 pages).

STN Registry Database, RN 923768-18-7, entered Feb. 28, 2007 (2 pages).

STN Registry Database, RN 923809-79-4, entered Feb. 28, 2007 (2 pages).

STN Registry Database, RN 931893-54-8, entered Apr. 23, 2007 (2 pages).

STN Registry Database, RN 932130-00-2, entered Apr. 24, 2007 (2 pages).

STN Registry Database, RN 938283-11-5, entered Jun. 22, 2007 (2 pages).

"FLI1 gene," MedlinePlus, published May 1, 2012, <<https://medlineplus.gov/genetics/>> (3 pages).

Adam et al., International Application No. PCT/US2024/024407, filed Apr. 12, 2024 by applicant Foghorn Therapeutics Inc. (42 pages).

Adam et al., International Application No. PCT/US2024/024428, filed Apr. 12, 2024 by applicant Foghorn Therapeutics Inc. (40 pages).

Boulay et al., "Cancer-Specific Retargeting of BAF Complexes by a Prion-like Domain," Cell 171(1):163-78 (Sep. 21, 2017) (36 pages).

CAS RN: 1223164-86-0; STN entry date: May 14, 2010; N-[2-[[4-(3-Fluoro-4-methoxyphenyl)-2-thiazolyl]amino]-2-oxoethyl]-2-methyl-3-furancarboxamide (1 page).

CAS RN: 1300403-14-8; STN entry date: May 25, 2011; 5-Methyl-N-[2-oxo-2-[(5-phenyl-2-pyridinyl)amino]ethyl]-2-thiophenecarboxamide (1 page).

CAS RN: 924410-17-3; STN entry date: Mar. 2, 2007; 5-Methyl-N-[2-oxo-2-[(4-phenyl-2-thiazolyl)amino]ethyl]-2-thiophenecarboxamide (1 page).

CAS RN: 924420-04-2; STN entry date: Mar. 2, 2007; 5-Methyl-N-[2-oxo-2-[[4-(4-pyridinyl)-2-thiazolyl]amino]ethyl]-2-thiophenecarboxamide (1 page).

Chandler et al., "ARID1a-DNA interactions are required for promoter occupancy by SWI/SNF," Mol Cell Biol. 33(2):265-80 (Jan. 2013).

Chattopadhyay et al., "Uveal melanoma: From diagnosis to treatment and the science in between," Cancer. 122(15):2299-2312 (26 pages) (Aug. 2016).

Database Registry, RN 1323331-37-8, entered Aug. 25, 2011 (1 page).

Database Registry, RN 1323542-96-6, entered Aug. 26, 2011 (1 page).

Database Registry, RN 1324163-01-0, entered Aug. 28, 2011 (1 page).

Database Registry, RN 1327304-26-6, entered Sep. 2, 2011 (1 page).

Fadul et al., "EWS/FLI utilizes NKX2-2 to repress mesenchymal features of Ewing sarcoma," Genes Cancer 6(3-4):129-43 (Mar. 2015).

Grohar et al., "Ecteinascidin 743 interferes with the activity of EWS-FLI1 in Ewing sarcoma cells," Neoplasia 13(2):145-53 (Feb. 2011).

Herrero-Martín et al., "Stable interference of EWS-FLI1 in an Ewing sarcoma cell line impairs IGF-1/IGF-1R signalling and reveals TOPK as a new target," Br J Cancer 101(1):80-90 (Jul. 7, 2009).

Hohmann et al., "Sensitivity and engineered resistance of myeloid leukemia cells to BRD9 inhibition," Nat Chem Biol. 12(9): 672-679 (Sep. 2016) (12 pages).

International Preliminary Report on Patentability for International Application No. PCT/US2019/056312, mailed Apr. 14, 2021 (10 pages).

International Search Report and Written Opinion for International Application No. PCT/US18/000339 dated Jan. 28, 2019 (13 pages).

International Search Report and Written Opinion for International Application No. PCT/US19/56312, dated Jan. 14, 2020 (14 pages).

International Search Report and Written Opinion for International Application No. PCT/US19/56365 dated Jan. 30, 2020 (8 pages).

International Search Report and Written Opinion for International Application No. PCT/US2024/024407, mailed Jun. 24, 2024 (15 pages).

International Search Report and Written Opinion for International Application No. PCT/US2024/024428, mailed Jul. 16, 2024 (18 pages).

(56)

References Cited**OTHER PUBLICATIONS**

Kedage et al., "An Interaction with Ewing's Sarcoma Breakpoint Protein EWS Defines a Specific Oncogenic Mechanism of ETS Factors Rearranged in Prostate Cancer," *Cell Rep.* 17(5):1289-301 (Oct. 25, 2016) (14 pages).

Michel et al., "Abstract PR15: BRD9 defines a novel mammalian SWI/SNF (BAF) complex configuration which supports proliferation in AML," *Clin Cancer Res.* 23(24_Suppl) Abstract PR15 (2017) (4 pages).

Piel et al., International Application No. PCT/US2024/031875, filed May 31, 2024 by applicant Foghorn Therapeutics Inc. (47 pages).

Rago et al., "Exquisite Sensitivity to Dual BRG1/BRM ATPase Inhibitors Reveals Broad SWI/SNF Dependencies in Acute Myeloid Leukemia," *Mol Cancer Res.* 20(3):361-72 (Mar. 1, 2022).

Riggi et al., "EWS-FLI1 utilizes divergent chromatin remodeling mechanisms to directly activate or repress enhancer elements in Ewing sarcoma," *Cancer Cell* 26(5):668-81 (Nov. 10, 2014).

Sankar et al., "Promiscuous partnerships in Ewing's sarcoma," *Cancer Genet.* 204(7):351-65 (Jul. 2011).

STN Registry Database, CAS RN 858073-83-3, Albany Molecular Research, Inc., entered Aug. 3, 2005 (1 page).

Takigami et al., "Synthetic siRNA targeting the breakpoint of EWS/Flt-1 inhibits growth of Ewing sarcoma xenografts in a mouse model," *Int J Cancer* 128(1):216-26 (Jan. 1, 2011).

Wu et al., "Targeting the chromatin remodeling enzyme BRG1 increases the efficacy of chemotherapy drugs in breast cancer cells," *Oncotarget* 7(19):27158-75 (May 10, 2016).

Collins et al., "Abstract 2122: The dual BRM/BRG1 (SMARCA2/4) inhibitor FHD-286 induces differentiation in preclinical models of AML," *Cancer Res.* 83(7_Supplement) (Apr. 2023) (5 pages).

Fiskus et al., "Pre-Clinical Efficacy of Targeting Baf Complexes through Inhibition of the Dual ATPases BRG1 and BRM By FHD-286 in Cellular Models of AML of Diverse Genetic Background," *Blood.* 140(Supplement 1):8819-20 (Nov. 2022) (15 pages).

Hentemann, "Abstract ND14: Pharmacological profile and anti-tumor properties of FHD-286: A novel BAF inhibitor for the treatment of transcription factor-driven cancers," *Cancer Res.* 82(12_Supplement): ND14 (Jun. 2022) (4 pages).

International Search Report and Written Opinion for International Patent Application No. PCT/US2024/031875, mailed Oct. 18, 2024 (15 pages).

International Search Report and Written Opinion for PCT/US2024/050660, mailed Nov. 26, 2024 (12 pages).

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6

METHODS OF TREATING CANCERS

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Nov. 5, 2021 is named 51121-049002_Sequence_Listing_9_13_21_ST25 and is 43,669 bytes in size.

BACKGROUND

The invention relates to methods for modulating BRG1- or BRM-associated factors (BAF) complexes for use in the treatment of acute myeloid leukemia (AML). In particular, the invention relates to methods for treatment of disorders associated with BAF complex function.

Chromatin regulation is essential for gene expression, and ATP-dependent chromatin remodeling is a mechanism by which such gene expression occurs. The human Switch/Sucrose Non-Fermentable (SWI/SNF) chromatin remodeling complex, also known as BAF complex, has two SWI2-like ATPases known as BRG1 (Brahma-related gene-1) and BRM (Brahma). The transcription activator BRG1, also known as ATP-dependent chromatin remodeler SMARCA4, is encoded by the SMARCA4 gene on chromosome 19. BRG1 is overexpressed in some cancer tumors and is needed for cancer cell proliferation. BRM, also known as probable global transcription activator SNF2L2 and/or ATP-dependent chromatin remodeler SMARCA2, is encoded by the SMARCA2 gene on chromosome 9 and has been shown to be essential for tumor cell growth in cells characterized by loss of BRG1 function mutations. Deactivation of BRG and/or BRM results in downstream effects in cells, including cell cycle arrest and tumor suppression.

AML is a cancer of the myeloid line of blood cells. AML is characterized by the rapid growth of abnormal cells that build up in the bone marrow and blood and interfere with normal blood cells. AML is generally considered incurable in about 65% of subjects under 60 years old and about 90% of subjects over 60 years old. Typical survival of older subjects with health too poor for intensive chemotherapy is 5- to 10-months. The five-year survival rate for AML is about 25% overall.

SUMMARY OF THE INVENTION

The present invention features methods to treat AML, e.g., in a subject in need thereof.

In one aspect, the invention features a method of treating AML in a subject in need thereof, the method including administering to the subject an effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM.

In another aspect, the invention features a method of reducing proliferation of AML in a subject in need thereof, the method including administering (e.g., oral administration) to the subject an effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM in the tumor.

In another aspect, the invention features a method of suppressing metastatic progression of AML in a subject, the method including administering (e.g., oral administration) an effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM.

In another aspect, the invention features a method of suppressing metastatic colonization of AML in a subject, the

method including administering (e.g., oral administration) an effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM.

In another aspect, the invention features a method of reducing the level and/or activity of BRG1 and/or BRM in an AML cell, the method including contacting the cell with an effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM in the cell.

In some embodiments of any of the above aspects, the AML cell is in a subject.

In some embodiments of any of the above aspects, the effective amount of the agent reduces the level and/or activity of BRG1 by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference. In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRG1 by at least 50% (e.g., 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference. In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRG1 by at least 90% (e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%).

In some embodiments, the effective amount of the agent reduces the level and/or activity of BRG1 by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference for at least 12 hours (e.g., 14 hours, 16 hours, 18 hours, 20 hours, 22 hours, 24 hours, 30 hours, 36 hours, 48 hours, 72 hours, or more).

In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRG1 by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference for at least 4 days (e.g., 5 days, 6 days, 7 days, 14 days, 28 days, or more).

In some embodiments of any of the above aspects, the effective amount of the agent reduces the level and/or activity of BRM by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference. In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRM by at least 50% (e.g., 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference. In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRM by at least 90% (e.g., 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%).

In some embodiments, the effective amount of the agent reduces the level and/or activity of BRM by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference for at least 12 hours (e.g., 14 hours, 16 hours, 18 hours, 20 hours, 22 hours, 24 hours, 30 hours, 36 hours, 48 hours, 72 hours, or more). In some embodiments, the effective amount of the agent that reduces the level and/or activity of BRM by at least 5% (e.g., 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, or 95%) as compared to a reference for at least 4 days (e.g., 5 days, 6 days, 7 days, 14 days, 28 days, or more).

In some embodiments, the AML expresses BRG1 and/or BRM protein and/or the cell or subject has been identified as expressing BRG1 and/or BRM. In some embodiments, the AML has an elevated expression of BRG1 and/or BRM. In some embodiments, the AML expresses BRG1 protein and/or the cell or subject has been identified as expressing BRG1. In some embodiments, the AML has an elevated

expression of BRG1. In some embodiments, the AML expresses BRM protein and/or the cell or subject has been identified as expressing BRM. In some embodiments, the AML has an elevated expression of BRM.

In some embodiments, the AML is acute promyelocytic leukemia (APL). In some embodiments, the AML arises from a pre-existing myelodysplastic syndrome or myeloproliferative disease. In some embodiments, the AML is treatment-related AML, e.g., AML arising after chemotherapy for another previous malignancy. In some embodiments, the AML is AML with recurrent genetic abnormalities. In some embodiments, the AML is AML with myelodysplasia-related changes. In some embodiments, the AML is therapy-related myeloid neoplasms. In some embodiments, the AML is myeloid sarcoma. In some embodiments, the AML is myeloid proliferations related to Down syndrome. In some embodiments, the AML is blastic plasmacytoid dendritic cell neoplasm. In some embodiments, the AML is AML minimally differentiated. In some embodiments, the AML is AML without maturation. In some embodiments, the AML is AML with granulocytic maturation. In some embodiments, the AML is myelomonocytic together with bone marrow eosinophilia. In some embodiments, the AML is acute monoblastic leukemia. In some embodiments, the AML is acute erythroid leukemia, e.g., erythroleukemia or pure erythroid leukemia. In some embodiments, the AML is acute megakaryoblastic leukemia. In some embodiments, the AML is acute basophilic leukemia.

In some embodiments, the cytogenetics of the AML are t(8;21), t(15;17), or inv(16). In some embodiments the cytogenetics of the AML are normal, +8, +21, +22, del(7q), del(9q), or abnormal 11q23. In some embodiments, the cytogenetics of the AML are -5, -7, del(5q), abnormal 3q, or complex cytogenetics.

In some embodiments, the cancer is metastatic (e.g., the cancer has spread to the liver). The metastatic cancer can include cells exhibiting migration and/or invasion of migrating cells and/or include cells exhibiting endothelial recruitment and/or angiogenesis. In some embodiments, the effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM is an amount effective to inhibit metastatic colonization of the AML to the liver.

In some embodiments, the AML harbors a DNMT3A mutation. In some embodiments, the AML harbors an FLT3-ITD mutation. In some embodiments, the AML harbors a NPM1 mutation. In some embodiments, the AML harbors a CEBPA mutation (e.g., a biallelic CEBPA mutation). In some embodiments, the AML harbors a c-KIT mutation. In some embodiments, the AML harbors a RUNX1 mutation. In some embodiments, the AML harbors an ASXL1 mutation. In some embodiments, the AML harbors a TP53 mutation. In some embodiments, the AML harbors a DNMT3A mutation. In some embodiments, the AML harbors an IDH1 mutation. In some embodiments, the AML harbors an IDH2 mutation.

In some embodiments, the subject is over 60 years old. In some embodiments, the subject has elevated levels of lactate dehydrogenase.

In some embodiments, the anticancer therapy and the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell are administered within 28 days of each other and each in an amount that together are effective to treat the subject.

In some embodiments, the subject or cancer has and/or has been identified as having a BRG1 loss of function mutation. In some embodiments, the subject or cancer has

and/or has been identified as having a BRM loss of function mutation. In some embodiments, the cancer harbors a BRG1 T910M mutation.

In some embodiments, the method further includes administering induction chemotherapy (e.g., cytarabine, an anthracycline such as daunorubicin, arsenic trioxide, all-trans-retinoic acid, or combinations thereof). In some embodiments, the method includes administering a 7-day infusion of cytarabine and a 3-day intravenous push of an anthracycline such as daunorubicin. In some embodiments, the method further includes administering consolidation therapy (e.g., an allogenic stem cell transplantation, immunotherapy such as histamine dihydrochloride and interleukin 2, or combinations thereof). In some embodiments, the method further includes administering a hemopoietic stem cell transplant, gemtuzumab ozogamicin, or combinations thereof. In some embodiments, the method further includes administering venetoclax, gilteritinib, or combinations thereof.

In some embodiments, the cancer is resistant to one or more chemotherapeutic or cytotoxic agents (e.g., the cancer has been determined to be resistant to chemotherapeutic or cytotoxic agents such as by genetic markers, or is likely to be resistant, to chemotherapeutic or cytotoxic agents such as a cancer that has failed to respond to a chemotherapeutic or cytotoxic agent). In some embodiments, the cancer has failed to respond to one or more chemotherapeutic or cytotoxic agents. In some embodiments, the cancer is resistant or has failed to respond to cytarabine, an anthracycline such as daunorubicin, arsenic trioxide, all-trans-retinoic acid, histamine dihydrochloride, interleukin 2, gemtuzumab ozogamicin, dacarbazine, temozolomide, cisplatin, treosulfan, fotemustine, IMCgp100, a CTLA-4 inhibitor (e.g., ipilimumab), a PD-1 inhibitor (e.g., Nivolumab or pembrolizumab), a PD-L1 inhibitor (e.g., atezolizumab, avelumab, or durvalumab), a mitogen-activated protein kinase (MEK) inhibitor (e.g., selumetinib, binimetinib, or tametinib), and/or a protein kinase C (PKC) inhibitor (e.g., sotrastaurin or LXS196, also known as IDE196). In some embodiments, the cancer is resistant or has failed to respond to cytarabine, an anthracycline such as daunorubicin, arsenic trioxide, all-trans-retinoic acid, histamine dihydrochloride, interleukin 2, and/or gemtuzumab ozogamicin. In some embodiments, the cancer is resistant or has failed to respond to venetoclax, gilteritinib, or combinations thereof.

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a small molecule compound, an antibody, an enzyme, and/or a polynucleotide.

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is an enzyme, e.g., a clustered regularly interspaced short palindromic repeats (CRISPR)-associated protein such as CRISPR-associated protein 9 (Cas9), CRISPR-associated protein 12a (Cas12a), a zinc finger nuclease (ZFN), a transcription activator-like effector nuclease (TALEN), or a meganuclease.


In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a polynucleotide, e.g., an antisense nucleic acid, a short interfering RNA (siRNA), a short hairpin RNA (shRNA), a microRNA (miRNA), a CRISPR/Cas 9 nucleotide, or a ribozyme.

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a small molecule compound, e.g., a small molecule BRG1 and/or BRM inhibitor. In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a

5

small molecule compound, e.g., a small molecule BRG1 inhibitor. In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a small molecule compound, e.g., a small molecule BRM inhibitor or a degrader.


In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula I:



Formula I

wherein m is 0, 1, 2, 3, or 4; X¹ is N or CH; and each R¹ is, independently, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino.

In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula II:



Formula II


wherein R² is phenyl that is substituted with hydroxy and that is optionally substituted with one or more groups independently selected from the group consisting of halo, cyano, trifluoromethyl, trifluoromethoxy, C₁-₃ alkyl, and C₁-₃ alkoxy; R³ is selected from the group consisting of —Rᵃ, —O—Rᵃ, —N(Rᵃ)₂, —S(O)₂Rᵃ, and —C(O)—N(Rᵃ)₂; each Rᵃ is, independently, selected from the group consisting of hydrogen, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein each C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from the group consisting of Rᵇ, oxo, halo, —NO₂, —N(Rᵇ)₂, —CN, —C(O)—N(Rᵇ)₂, —S(O)—N(Rᵇ)₂, —S(O)₂—N(Rᵇ)₂, —O—Rᵇ, —S—Rᵇ, —O—C(O)—Rᵇ, —C(O)—Rᵇ, —C(O)—ORᵇ, —S(O)—Rᵇ, —S(O)₂—Rᵇ, —N(Rᵇ)—C(O)—Rᵇ, —N(Rᵇ)—S(O)—Rᵇ, —N(Rᵇ)—C(O)—N(Rᵇ)₂, and —N(Rᵇ)—S(O)₂—Rᵇ; each Rᵇ is, independently, selected from the group consisting of hydrogen, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, C₁-₆ alkoxy, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein each C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, C₁-₆ alkoxy, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from RC; or two Rᵇ are

6

taken together with the nitrogen to which they are attached to form a heterocyclyl that is optionally substituted with one or more groups independently selected from the group consisting of oxo, halo and C₁-₃ alkyl that is optionally substituted with one or more groups independently selected from the group consisting of oxo and halo; each RC is, independently, selected from the group consisting of oxo, halo, —NO₂, —N(Rᵈ)₂, —CN, —C(O)—N(Rᵈ)₂, —S(O)—N(Rᵈ)₂, —S(O)₂—N(Rᵈ)₂, —S—Rᵈ, —O—C(O)—Rᵈ, —C(O)—Rᵈ, —C(O)—ORᵈ, —S(O)—Rᵈ, —S(O)₂—Rᵈ, —N(Rᵈ)—C(O)—Rᵈ, —N(Rᵈ)—S(O)—Rᵈ, —N(Rᵈ)—C(O)—N(Rᵈ)₂, —N(Rᵈ)—S(O)—Rᵈ, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein any C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from the group consisting of Rᵈ, oxo, halo, —NO₂, —N(Rᵈ)₂, —CN, —C(O)—N(Rᵈ)₂, —S(O)—N(Rᵈ)₂, —S(O)₂—N(Rᵈ)₂, —O—Rᵈ, —S—Rᵈ, —O—C(O)—Rᵈ, —C(O)—Rᵈ, —C(O)—ORᵈ, —S(O)—Rᵈ, —S(O)₂—Rᵈ, —N(Rᵈ)—C(O)—Rᵈ, —N(Rᵈ)—S(O)—Rᵈ, —N(Rᵈ)—C(O)—N(Rᵈ)₂, and —N(Rᵈ)—S(O)₂—Rᵈ; each Rᵈ is, independently, selected from the group consisting of hydrogen, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, carbocyclyl, and carbocyclyl(C₁-₃ alkyl); R⁴ is H, C₁-₆ alkyl, or —C(=O)—C₁-₆ alkyl; and R⁵ is H or C₁-₆ alkyl.

Compounds of Formula II may be synthesized by methods known in the art, e.g., those described in U.S. Patent Publication No. 2018/0086720, the synthetic methods of which are incorporated by reference.


In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula III:



Formula III

wherein R⁶ is halo, e.g., fluoro or chloro; R⁷ is hydrogen, optionally substituted amino, or optionally substituted C₁-₆ alkyl; and R⁸ is optionally substituted C₆-₁₀ aryl or optionally substituted C₂-₉ heteroaryl.










In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of any one of compounds 1-16:



1

7

-continued




8

-continued

2

11

5




3

10

12

15

4




20

13

5

25




6

30

14

35




15

7

40

8

45




16

50

9

55




10

60

65 In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula IV:

9


Formula IV



wherein R¹ is absent, H, optionally substituted C₁-C₆ acyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶;



is 5- or 6-membered heteroarylene; each of R² and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl; R³ is H or optionally substituted C₁-C₆ alkyl; and R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; or R³ and R⁴, together with the carbon atom to which each is attached, form an optionally substituted C₃-C₆ cycloalkyl; R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸; R⁷ and R⁸ are, independently, optionally substituted C₁-C₆ alkyl; Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or



A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene; L is absent, —O—, optionally substituted C₁-C₆ alkylene, optionally substituted C₁-C₆ heteroalkylene, optionally substituted C₂-C₆ alkenylene, optionally substituted C₂-C₆ heteroalkenylene, optionally substituted C₂-C₆ alkynylene, optionally substituted C₂-C₆ heteroalkynylene, optionally substituted C₂-C₉ heterocyclylene, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkylene, optionally substituted C₂-C₉ heteroarylene, or optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkylene; and B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl, or a pharmaceutically acceptable salt thereof.

In some embodiments,




is 6-membered heteroarylene. In some embodiments,



is 5-membered heteroarylene.


10

In some embodiments,




where each of X, Y, and Z is, independently, N or CH.

In some embodiments, the compound of Formula IV has the structure of Formula IVa:



wherein each of X, Y, and Z is, independently, N or CH; R¹ is H, optionally substituted C₁-C₆ acyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶; each of R², R³, and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl; R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸; each of R⁷ and R⁸ is, independently, optionally substituted C₁-C₆ alkyl; Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or




A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene; L is absent, —O—, optionally substituted C₁-C₆ alkylene, optionally substituted C₁-C₆ heteroalkylene, optionally substituted C₂-C₆ alkenylene, optionally substituted C₂-C₆ heteroalkenylene, optionally substituted C₂-C₆ alkynylene, optionally substituted C₂-C₆ heteroalkynylene, optionally substituted C₂-C₉ heterocyclylene, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkylene, optionally substituted C₂-C₉ heteroarylene, or optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkylene; and B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl, or a pharmaceutically acceptable salt thereof.

In some embodiments, X, Y, and Z are CH; X is N and Y and Z are CH; Z is N and X and Y are CH; Y is N and X and Z are CH; X is CH and Y and Z are N; Z is CH and X and Y are N; Y is CH and X and Z are N; or X, Y, and Z are N.

In some embodiments, the compound of Formula IV has the structure of Formula IVb:

11


Formula IVb



or a pharmaceutically acceptable salt thereof.

In some embodiments, the compound of Formula IV has the structure of Formula IVc:


Formula IVc



or a pharmaceutically acceptable salt thereof.

In some embodiments, the compound of Formula IV has the structure of Formula IVd:


Formula IVd



or a pharmaceutically acceptable salt thereof.


In some embodiments, the compound of Formula IV has the structure of Formula IVe:

Formula IVe



or a pharmaceutically acceptable salt thereof.

In some embodiments,




where X' is O or S; Y' is N or CH; and Z' is N or CH.


In some embodiments, the compound of Formula IVa has the structure of Formula Va:

12

Formula V



wherein W is C or N; X' is O, S, or N—CH₃; Y' is N or CH; Z' is N or CH; R¹ is absent, H, optionally substituted C₁-C₆ acyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶; each of R², R³, and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl; R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸; each of R⁷ and R⁸ is, independently, optionally substituted C₁-C₆ alkyl; Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or




A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene; L is absent, —O—, optionally substituted C₁-C₆ alkylene, optionally substituted C₁-C₆ heteroalkylene, optionally substituted C₁-C₆ alkenylene, optionally substituted C₂-C₆ heteroalkenylene, optionally substituted C₂-C₆ alkynylene, optionally substituted C₂-C₆ heteroalkynylene, optionally substituted C₂-C₉ heterocyclylene, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkylene, optionally substituted C₂-C₉ heteroarylene, or optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkylene; and B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl, or a pharmaceutically acceptable salt thereof.

In some embodiments, X' is O, Y' is CH, and Z' is N; X' is S, Y' is CH, and Z' is N; X' is O, Y' is N, and Z' is CH; X' is S, Y' is N, and Z' is CH; X' is O, Y' is N, and Z' is N; or X' is S, Y' is N, and Z' is N.

In some embodiments, the compound of Formula V has the structure of Formula Va:

Formula Va




or a pharmaceutically acceptable salt thereof.

In some embodiments, the compound of Formula II has the structure of Formula Vb:





13

Formula Vb



or a pharmaceutically acceptable salt thereof.

In some embodiments, the small molecule compound, or pharmaceutically acceptable salt thereof is any one of compounds 17-20 having the structure:



In some embodiments, the small molecule compound, or a pharmaceutically acceptable salt thereof is a degrader. In some embodiments, the degrader has the structure of Formula VI:

C-L-D

Formula VI



wherein C is a BRG1 and/or BRM binding moiety; L is a linker; and D is a degradation moiety, or a pharmaceutically acceptable salt thereof. In some embodiments, the degradation moiety is a ubiquitin ligase moiety. In some embodiments, the ubiquitin ligase binding moiety includes Cereblon ligands, IAP (Inhibitors of Apoptosis) ligands, mouse

14

double minute 2 homolog (MDM2), hydrophobic tag, or von Hippel-Lindau ligands, or derivatives or analogs thereof.


In some embodiments, A includes the structure of any one of Formula I-V, or any one of compounds 1-20.

In some embodiments, the hydrophobic tag includes a diphenylmethane, adamantane, or tri-Boc arginine, i.e., the hydrophobic tag includes the structure:




In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula A:

Formula A



wherein X^1 is CH_2 , O, S, or NR^1 , wherein R^1 is H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; X^2 is $C=O$, CH_2 , or




R^3 and R^4 are, independently, H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; m is 0, 1, 2, 3, or 4; and each R^2 is, independently, halogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocycl, optionally substituted C_2 - C_9 heterocycl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_6 alkenyl, optionally

15

substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino, or a pharmaceutically acceptable salt thereof.


In some embodiments, the ubiquitin ligase binding moiety includes the structure:



or is a derivative or an analog thereof, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula B:


Formula B



wherein each R⁴, R^{4'}, and R⁷ is, independently, H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; R⁵ is optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₁-C₆ alkyl C₃-C₁₀ carbocyclyl, or optionally substituted C₁-C₆ alkyl C₆-C₁₀ aryl; R⁶ is H, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₁-C₆ alkyl C₃-C₁₀ carbocyclyl, or optionally substituted C₁-C₆ alkyl C₆-C₁₀ aryl; n is 0, 1, 2, 3, or 4; each R⁸ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino; and each R⁹ and R¹⁰ is, independently, H, halogen, optionally substituted C₁-C₆ alkyl, or optionally substituted C₆-C₁₀ aryl, wherein R^{4'} or R⁵ includes a bond to the linker, or a pharmaceutically acceptable salt thereof.

16


In some embodiments, the ubiquitin ligase binding moiety includes the structure:



or is a derivative or analog thereof, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula C:

Formula C




wherein each R¹¹, R¹³, and R¹⁵ is, independently, H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; R¹² is optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₁-C₆ alkyl C₃-C₁₀ carbocyclyl, or optionally substituted C₁-C₆ alkyl C₆-C₁₀ aryl; R¹⁴ is optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₁-C₆ alkyl C₃-C₁₀ carbocyclyl, or optionally substituted C₁-C₆ alkyl C₆-C₁₀ aryl; p is 0, 1, 2, 3, or 4; each R¹⁶ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally

17


substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino; q is 0, 1, 2, 3, or 4; and each R¹⁷ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure:



or is a derivative or an analog thereof, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula D:




wherein each R¹⁸ and R¹⁹ is, independently, H, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₁-C₆ alkyl C₃-C₁₀ carbocyclyl, or optionally substituted C₁-C₆ alkyl C₆-C₁₀ aryl; r₁ is 0, 1, 2, 3, or 4; each R²⁰ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino; r₂ is 0, 1, 2,

18


3, or 4; and each R²¹ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure:



or is a derivative or an analog thereof, or a pharmaceutically acceptable salt thereof.

In some embodiments, the linker has the structure of Formula V:




wherein A¹ is a bond between the linker and A; A² is a bond between B and the linker; B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl, optionally substituted C₁-C₃ heteroalkyl, O, S, S(O)₂, and NR^N; R^N is hydrogen, optionally substituted C₁₋₄ alkyl, optionally substituted C₂₋₄ alkenyl, optionally substituted C₂₋₄ alkynyl, optionally substituted C₂₋₆ heterocyclyl, optionally substituted C₆₋₁₂ aryl, or optionally substituted C₁₋₇ heteroalkyl; C¹ and C² are each, independently, selected from carbonyl, thiocarbonyl, sulfonyl, or phosphoryl; f, g, h, i, j, and k are each, independently, 0 or 1; and D is optionally substituted C₁₋₁₀ alkyl, optionally substituted C₂₋₁₀ alkenyl, optionally substituted C₂₋₁₀ alkynyl, optionally substituted C₂₋₆ heterocyclyl, optionally substituted C₆₋₁₂ aryl, optionally substituted C₂-C₁₀ polyethylene glycol, or optionally substituted C₁₋₁₀ heteroalkyl, or a chemical bond linking A¹-(B¹)_f-(C¹)_g-(B²)_h to -(B³)_i-(C²)_j-(B⁴)_k-A².

In some embodiments, D is optionally substituted C₂-C₁₀ polyethylene glycol. In some embodiments, C¹ and C² are each, independently, a carbonyl or sulfonyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl, optionally substituted C₁-C₃ heteroalkyl, O, S, S(O)₂, and NR^N; R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl or optionally substituted C₁-C₃ heteroalkyl. In some embodiments, j is 0. In some embodiments, k is 0. In some embodiments, j and k are each, independently, 0. In some embodiments, f, g, h, and i are each, independently, 1.

19


In some embodiments, the linker of Formula VII has the structure of Formula VIIa:



wherein A¹ is a bond between the linker and A, and A² is a bond between B and the linker.

In some embodiments, D is optionally substituted C₁₋₁₀ alkyl. In some embodiments, C¹ and C² are each, independently, a carbonyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl, optionally substituted C₁-C₃ heteroalkyl, O, S, S(O)₂, and NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁-C₂ alkyl, O, S, S(O)₂, and NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B¹ and B⁴ each, independently, is optionally substituted C₁-C₂ alkyl. In some embodiments, B¹ and B⁴ each, independently, is C₁ alkyl. In some embodiments, B² and B⁴ each, independently, is NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B² and B⁴ each, independently, is NH. In some embodiments, f, g, h, i, j, and k are each, independently, 1.

In some embodiments, the linker of Formula VII has the structure of Formula VIIb:



wherein A¹ is a bond between the linker and A, and A² is a bond between B and the linker.

Chemical Terms

For any of the following chemical definitions, a number following an atomic symbol indicates that total number of atoms of that element that are present in a particular chemical moiety. As will be understood, other atoms, such as hydrogen atoms, or substituent groups, as described herein, may be present, as necessary, to satisfy the valences of the atoms. For example, an unsubstituted C₂ alkyl group has the formula —CH₂CH₃. When used with the groups defined herein, a reference to the number of carbon atoms includes the divalent carbon in acetal and ketal groups but does not include the carbonyl carbon in acyl, ester, carbonate, or carbamate groups. A reference to the number of oxygen, nitrogen, or sulfur atoms in a heteroaryl group only includes those atoms that form a part of a heterocyclic ring.

The term “acyl,” as used herein, represents a hydrogen or an alkyl group that is attached to a parent molecular group through a carbonyl group, as defined herein, and is exemplified by formyl (i.e., a carboxyaldehyde group), acetyl, trifluoroacetyl, propionyl, and butanoyl. Exemplary unsubstituted acyl groups include from 1 to 6, from 1 to 11, or from 1 to 21 carbons.

20

The term “alkyl,” as used herein, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of 1 to 20 carbon atoms (e.g., 1 to 16 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms).

An alkylene is a divalent alkyl group. The term “alkenyl,” as used herein, alone or in combination with other groups, refers to a straight chain or branched hydrocarbon residue having a carbon-carbon double bond and having 2 to 20 carbon atoms (e.g., 2 to 16 carbon atoms, 2 to 10 carbon atoms, 2 to 6, or 2 carbon atoms).

The term “alkynyl,” as used herein, alone or in combination with other groups, refers to a straight chain or branched hydrocarbon residue having a carbon-carbon triple bond and having 2 to 20 carbon atoms (e.g., 2 to 16 carbon atoms, 2 to 10 carbon atoms, 2 to 6, or 2 carbon atoms).

The term “amino,” as used herein, represents —N(R^{N1})₂, wherein each R^{N1} is, independently, H, OH, NO₂, N(R^{N2})₂, SO₂OR^{N2}, SO₂R^{N2}, SOR^{N2}, an N-protecting group, alkyl, alkoxy, aryl, arylalkyl, cycloalkyl, acyl (e.g., acetyl, trifluoroacetyl, or others described herein), wherein each of these recited R^{N1} groups can be optionally substituted; or two R^{N1} combine to form an alkylene or heteroalkylene, and wherein each R^{N2} is, independently, H, alkyl, or aryl. The amino groups of the compounds described herein can be an unsubstituted amino (i.e., —NH₂) or a substituted amino (i.e., —N(R^{N1})₂).

The term “aryl,” as used herein, refers to an aromatic mono- or polycarbocyclic radical of 6 to 12 carbon atoms having at least one aromatic ring. Examples of such groups include, but are not limited to, phenyl, naphthyl, 1,2,3,4-tetrahydronaphthyl, 1,2-dihydronaphthyl, indanyl, and 1H-indenyl.

The term “arylalkyl,” as used herein, represents an alkyl group substituted with an aryl group. Exemplary unsubstituted arylalkyl groups are from 7 to 30 carbons (e.g., from 7 to 16 or from 7 to 20 carbons, such as C₁-C₆ alkyl C₆-C₁₀ aryl, C₁-C₁₀ alkyl C₆-C₁₀ aryl, or C₁-C₂₀ alkyl C₆-C₁₀ aryl), such as, benzyl and phenethyl. In some embodiments, the alkyl and the aryl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective groups.

The term “azido,” as used herein, represents a —N₃ group.

The term “bridged polycycloalkyl,” as used herein, refers to a bridged polycyclic group of 5 to 20 carbons, containing from 1 to 3 bridges.

The term “cyano,” as used herein, represents a —CN group.

The term “carbocyclyl,” as used herein, refers to a non-aromatic C₃-C₁₂ monocyclic, bicyclic, or tricyclic structure in which the rings are formed by carbon atoms. Carbocyclyl structures include cycloalkyl groups and unsaturated carbocyclyl radicals.

The term “cycloalkyl,” as used herein, refers to a saturated, non-aromatic, monovalent mono- or polycarbocyclic radical of 3 to 10, preferably 3 to 6 carbon atoms. This term is further exemplified by radicals such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, norbornyl, and adamantyl.

The term “halogen,” as used herein, means a fluorine (fluoro), chlorine (chloro), bromine (bromo), or iodine (iodo) radical.

The term “heteroalkyl,” as used herein, refers to an alkyl group, as defined herein, in which one or more of the constituent carbon atoms have been replaced by nitrogen, oxygen, or sulfur. In some embodiments, the heteroalkyl group can be further substituted with 1, 2, 3, or 4 substituent

21

groups as described herein for alkyl groups. An example of a heteroalkyl group is an "alkoxy", which, as used herein, refers alkyl-O— (e.g., methoxy and ethoxy). A heteroalkylene is a divalent heteroalkyl group. The term "heteroalkenyl," as used herein, refers to an alkenyl group, as defined herein, in which one or more of the constituent carbon atoms have been replaced by nitrogen, oxygen, or sulfur. In some embodiments, the heteroalkenyl group can be further substituted with 1, 2, 3, or 4 substituent groups as described herein for alkenyl groups. Examples of heteroalkenyl groups are an "alkenoxy" which, as used herein, refers alkenyl-O—. A heteroalkenylene is a divalent heteroalkenyl group. The term "heteroalkynyl," as used herein, refers to an alkynyl group, as defined herein, in which one or more of the constituent carbon atoms have been replaced by nitrogen, oxygen, or sulfur. In some embodiments, the heteroalkynyl group can be further substituted with 1, 2, 3, or 4 substituent groups as described herein for alkynyl groups. Examples of heteroalkynyl groups are an "alkynoxy" which, as used herein, refers alkynyl-O—. A heteroalkynylene is a divalent heteroalkynyl group.

The term "heteroaryl," as used herein, refers to an aromatic mono- or polycyclic radical of 5 to 12 atoms having at least one aromatic ring containing 1, 2, or 3 ring atoms selected from nitrogen, oxygen, and sulfur, with the remaining ring atoms being carbon. One or two ring carbon atoms of the heteroaryl group may be replaced with a carbonyl group. Examples of heteroaryl groups are pyridyl, pyrazolyl, benzoxazolyl, benzoimidazolyl, benzothiazolyl, imidazolyl, oxazolyl, and thiazolyl.

The term "heteroarylalkyl," as used herein, represents an alkyl group substituted with a heteroaryl group. Exemplary unsubstituted heteroarylalkyl groups are from 7 to 30 carbons (e.g., from 7 to 16 or from 7 to 20 carbons, such as C₁-C₆ alkyl C₂-C₉ heteroaryl, C₁-C₁₀ alkyl C₂-C₉ heteroaryl, or C₁-C₂₀ alkyl C₂-C₉ heteroaryl). In some embodiments, the alkyl and the heteroaryl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective groups.

The term "heterocyclyl," as used herein, refers a mono- or polycyclic radical having 3 to 12 atoms having at least one ring containing 1, 2, 3, or 4 ring atoms selected from N, O or S, wherein no ring is aromatic. Examples of heterocyclyl groups include, but are not limited to, morpholinyl, thiomorpholinyl, furyl, piperazinyl, piperidinyl, pyranyl, pyrrolidinyl, tetrahydropyranyl, tetrahydrofuranlyl, and 1,3-dioxan-yl.

The term "heterocyclylalkyl," as used herein, represents an alkyl group substituted with a heterocyclyl group. Exemplary unsubstituted heterocyclylalkyl groups are from 7 to 30 carbons (e.g., from 7 to 16 or from 7 to 20 carbons, such as C₁-C₆ alkyl C₂-C₉ heterocyclyl, C₁-C₁₀ alkyl C₂-C₉ heterocyclyl, or C₁-C₂₀ alkyl C₂-C₉ heterocyclyl). In some embodiments, the alkyl and the heterocyclyl each can be further substituted with 1, 2, 3, or 4 substituent groups as defined herein for the respective groups.

The term "hydroxyalkyl," as used herein, represents alkyl group substituted with an —OH group.

The term "hydroxyl," as used herein, represents an —OH group.

The term "N-protecting group," as used herein, represents those groups intended to protect an amino group against undesirable reactions during synthetic procedures. Commonly used N-protecting groups are disclosed in Greene, "Protective Groups in Organic Synthesis," 3rd Edition (John Wiley & Sons, New York, 1999). N-protecting groups include, but are not limited to, acyl, aryloyl, or carbamyl

22

groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxycarbonyl, α-chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl, and chiral auxiliaries such as protected or unprotected D, L, or D, L-amino acids such as alanine, leucine, and phenylalanine; sulfonyl-containing groups such as benzenesulfonyl, and p-toluenesulfonyl; carbamate forming groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α-dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxy carbonyl, t-butyloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, and phenylthiocarbonyl, arylalkyl groups such as benzyl, triphenylmethyl, and benzyloxymethyl, and silyl groups, such as trimethylsilyl. Preferred N-protecting groups are alloc, formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, alanil, phenylsulfonyl, benzyl, t-butyloxycarbonyl (Boc), and benzyloxycarbonyl (Cbz).

The term "nitro," as used herein, represents an —NO₂ group.

The term "thiol," as used herein, represents an —SH group.

The alkyl, alkenyl, alkynyl, heteroalkyl, heteroalkenyl, heteroalkynyl, carbocyclyl (e.g., cycloalkyl), aryl, heteroaryl, and heterocyclyl groups may be substituted or unsubstituted. When substituted, there will generally be 1 to 4 substituents present, unless otherwise specified. Substituents include, for example: alkyl (e.g., unsubstituted and substituted, where the substituents include any group described herein, e.g., aryl, halo, hydroxy), aryl (e.g., substituted and unsubstituted phenyl), carbocyclyl (e.g., substituted and unsubstituted cycloalkyl), halogen (e.g., fluoro), hydroxyl, heteroalkyl (e.g., substituted and unsubstituted methoxy, ethoxy, or thioalkoxy), heteroaryl, heterocyclyl, amino (e.g., NH₂ or mono- or dialkyl amino), azido, cyano, nitro, or thiol. Aryl, carbocyclyl (e.g., cycloalkyl), heteroaryl, and heterocyclyl groups may also be substituted with alkyl (unsubstituted and substituted such as arylalkyl (e.g., substituted and unsubstituted benzyl)).

Compounds described herein can have one or more asymmetric carbon atoms and can exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates, or mixtures of diastereoisomeric racemates. The optically active forms can be obtained for example by resolution of the racemates, by asymmetric synthesis or asymmetric chromatography (chromatography with a chiral adsorbent or eluant). That is, certain of the disclosed compounds may exist in various stereoisomeric forms. Stereoisomers are compounds that differ only in their spatial arrangement. Enantiomers are pairs of stereoisomers whose mirror images are not superimposable, most commonly because they contain an asymmetrically substituted carbon atom that acts as a chiral center. "Enantiomer" means one of a pair of molecules that are mirror images of each other and are not

superimposable. Diastereomers are stereoisomers that are not related as mirror images, most commonly because they contain two or more asymmetrically substituted carbon atoms and represent the configuration of substituents around one or more chiral carbon atoms. Enantiomers of a compound can be prepared, for example, by separating an enantiomer from a racemate using one or more well-known techniques and methods, such as, for example, chiral chromatography and separation methods based thereon. The appropriate technique and/or method for separating an enantiomer of a compound described herein from a racemic mixture can be readily determined by those of skill in the art. "Racemate" or "racemic mixture" means a compound containing two enantiomers, wherein such mixtures exhibit no optical activity; i.e., they do not rotate the plane of polarized light. "Geometric isomer" means isomers that differ in the orientation of substituent atoms in relationship to a carbon-carbon double bond, to a cycloalkyl ring, or to a bridged bicyclic system. Atoms (other than H) on each side of a carbon-carbon double bond may be in an E (substituents are on 25 opposite sides of the carbon-carbon double bond) or Z (substituents are oriented on the same side) configuration. "R," "S," "S*," "R*," "E," "Z," "cis," and "trans," indicate configurations relative to the core molecule. Certain of the disclosed compounds may exist in atropisomeric forms. Atropisomers are stereoisomers resulting from hindered rotation about single bonds where the steric strain barrier to rotation is high enough to allow for the isolation of the conformers. The compounds described herein may be prepared as individual isomers by either isomer-specific synthesis or resolved from an isomeric mixture. Conventional resolution techniques include forming the salt of a free base of each isomer of an isomeric pair using an optically active acid (followed by fractional crystallization and regeneration of the free base), forming the salt of the acid form of each isomer of an isomeric pair using an optically active amine (followed by fractional crystallization and regeneration of the free acid), forming an ester or amide 35 of each of the isomers of an isomeric pair using an optically pure acid, amine or alcohol (followed by chromatographic separation and removal of the chiral auxiliary), or resolving an isomeric mixture of either a starting material or a final product using various well known chromatographic methods. When the stereochemistry of a disclosed compound is named or depicted by structure, the named or depicted stereoisomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by weight relative to the other stereoisomers. When a single enantiomer is named or depicted by structure, the depicted or named enantiomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by weight optically pure. When a single diastereomer is named or depicted by structure, the depicted or named diastereomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by weight pure. Percent optical purity is the ratio of the weight of the enantiomer or over the weight of the enantiomer plus the weight of its optical isomer. Diastereomeric purity by weight is the ratio of the weight of one diastereomer or over the weight of all the diastereomers. When the stereochemistry of a disclosed compound is named or depicted by structure, the named or depicted stereoisomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by mole fraction pure relative to the other stereoisomers. When a single enantiomer is named or depicted by structure, the depicted or named enantiomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by mole fraction pure. When a single diastereomer is named or depicted by structure, the depicted or named diastereomer is at least 60%, 70%, 80%, 90%, 99%, or 99.9% by mole fraction pure.

Percent purity by mole fraction is the ratio of the moles of the enantiomer or over the moles of the enantiomer plus the moles of its optical isomer. Similarly, percent purity by moles fraction is the ratio of the moles of the diastereomer or over the moles of the diastereomer plus the moles of its isomer. When a disclosed compound is named or depicted by structure without indicating the stereochemistry, and the compound has at least one chiral center, it is to be understood that the name or structure encompasses either enantiomer of the compound free from the corresponding optical isomer, a racemic mixture of the compound, or mixtures enriched in one enantiomer relative to its corresponding optical isomer. When a disclosed compound is named or depicted by structure without indicating the stereochemistry and has two or more chiral centers, it is to be understood that the name or structure encompasses a diastereomer free of other diastereomers, a number of diastereomers free from other diastereomeric pairs, mixtures of diastereomers, mixtures of diastereomeric pairs, mixtures of diastereomers in which one diastereomer is enriched relative to the other diastereomer(s), or mixtures of diastereomers in which one or more diastereomer is enriched relative to the other diastereomers. The invention embraces all of these forms.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Methods and materials are described herein for use in the present disclosure; other, suitable methods and materials known in the art can also be used. The materials, methods, and examples are illustrative only and not intended to be limiting. All publications, patent applications, patents, sequences, database entries, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control.

Definitions

In this application, unless otherwise clear from context, (i) the term "a" may be understood to mean "at least one"; (ii) the term "or" may be understood to mean "and/or"; and (iii) the terms "including" and "including" may be understood to encompass itemized components or steps whether presented by themselves or together with one or more additional components or steps.

As used herein, the terms "about" and "approximately" refer to a value that is within 10% above or below the value being described. For example, the term "about 5 nM" indicates a range of from 4.5 to 5.5 nM.

As used herein, the term "administration" refers to the administration of a composition (e.g., a compound or a preparation that includes a compound as described herein) to a subject or system. Administration to an animal subject (e.g., to a human) may be by any appropriate route. For example, in some embodiments, administration may be bronchial (including by bronchial instillation), buccal, enteral, interdermal, intra-arterial, intradermal, intragastric, intramedullary, intramuscular, intranasal, intraperitoneal, intrathecal, intratumoral, intravenous, intraventricular, mucosal, nasal, oral, rectal, subcutaneous, sublingual, topical, tracheal (including by intratracheal instillation), transdermal, vaginal, and vitreal.

As used herein, the term "BAF complex" refers to the BRG1-associated or HBRM-associated factors complex in a human cell.

As used herein, the term "BRG1 loss of function mutation" refers to a mutation in BRG1 that leads to the protein

having diminished activity (e.g., at least 1% reduction in BRG1 activity, for example 2%, 5%, 10%, 25%, 50%, or 100% reduction in BRG1 activity). Exemplary BRG1 loss of function mutations include, but are not limited to, a homozygous BRG1 mutation and a deletion at the C-terminus of BRG1.

As used herein, the term “BRG1 loss of function disorder” refers to a disorder (e.g., cancer) that exhibits a reduction in BRG1 activity (e.g., at least 1% reduction in BRG1 activity, for example 2%, 5%, 10%, 25%, 50%, or 100% reduction in BRG1 activity).

As used herein, the term “BRM loss of function mutation” refers to a mutation in BRM that leads to the protein having diminished activity (e.g., at least 1% reduction in BRM activity, for example 2%, 5%, 10%, 25%, 50%, or 100% reduction in BRM activity). Exemplary BRM loss of function mutations include, but are not limited to, a homozygous BRM mutation and a deletion at the C-terminus of BRM.

As used herein, the term “BRM loss of function disorder” refers to a disorder (e.g., cancer) that exhibits a reduction in BRM activity (e.g., at least 1% reduction in BRM activity, for example 2%, 5%, 10%, 25%, 50%, or 100% reduction in BRG1 activity).

As used herein, the terms “GBAF complex” and “GBAF” refer to a SWI/SNF ATPase chromatin remodeling complex in a human cell. GBAF complex subunits may include, but are not limited to, ACTB, ACTL6A, ACTL6B, BICRA, BICRAL, BRD9, SMARCA2, SMARCA4, SMARCC1, SMARCD1, SMARCD2, SMARCD3, and SS18.

The term “cancer” refers to a condition caused by the proliferation of malignant neoplastic cells, such as tumors, neoplasms, carcinomas, sarcomas, leukemias, and lymphomas.

As used herein, a “combination therapy” or “administered in combination” means that two (or more) different agents or

treatments are administered to a subject as part of a defined treatment regimen for a particular disease or condition. The treatment regimen defines the doses and periodicity of administration of each agent such that the effects of the separate agents on the subject overlap. In some embodiments, the delivery of the two or more agents is simultaneous or concurrent and the agents may be co-formulated. In some embodiments, the two or more agents are not co-formulated and are administered in a sequential manner as part of a prescribed regimen. In some embodiments, administration of two or more agents or treatments in combination is such that the reduction in a symptom, or other parameter related to the disorder is greater than what would be observed with one agent or treatment delivered alone or in the absence of the other. The effect of the two treatments can be partially additive, wholly additive, or greater than additive (e.g., synergistic). Sequential or substantially simultaneous administration of each therapeutic agent can be effected by any appropriate route including, but not limited to, oral routes, intravenous routes, intramuscular routes, and direct absorption through mucous membrane tissues. The therapeutic agents can be administered by the same route or by different routes. For example, a first therapeutic agent of the combination may be administered by intravenous injection while a second therapeutic agent of the combination may be administered orally.

As used herein, the term “BRG1” refers to ATP-dependent chromatin remodeler SMARCA4. BRG1 is a component of the BAF complex, a SWI/SNF ATPase chromatin remodeling complex. Human BRG1 is encoded by the SMARCA4 gene on chromosome 19, a nucleic acid sequence of which is set forth in SEQ ID NO: 1 (GenBank Accession No.: NM_001128849.1 (mRNA); www.ncbi.nlm.nih.gov/nuccore/NM_001128849.1?report=fasta).

```

GGCGGGGAGGCGCGGGAAGTCGACGGCGCGCGGCTCCTGCAGGAGGCCACTGTCTGCAGTCCCGT
GAAGATGTCCACTCCAGACCCACCCCTGGGCGGAACCTCTCGGCCAGGTCTTCCCCGGGCCCTGGCCCT
TCCCTTGGAGCATGCTGGGCCCTAGCCCGGGTCCCTCGCCGGGCTCCGCCACAGCATGATGGGGCCCA
GCCCAGGGCCCGCCTCAGCAGGACACCCCATCCCCACCCAGGGGCTGGAGGGTACCCTCAGGACAACAT
GCACCAGATGCACAAGCCCATGGAGTCCATGCATGAGAAGGGCATGTCCGACGACCCGCGCTACAACCAG
ATGAAAGGAATGGGGATGCGGTGAGGGGCGCATGCTGGGATGGGGCCCCGCCAGCCCCATGGACCAGC
ACTCCCAAGTTATCCCTCGCCCTGGGTGGCTCTGAGCATGCCTCTAGTCCAGTTCCAGCCAGTGGCCC
GTCCTCGGGGCCCCAGATGTCTTCCGGGCCAGGAGTGCCCGCTGGATGGTGTGACCCCCAGGCCTTG
GGGAGCAGAAACCGGGGCCAACCCATTAAACAGAACAGCTGCACCAGCTCAGAGCTCAGATCATGG
CCTACAAGATGCTGGCCAGGGGGCAGCCCTCCCCGACCACCTGCAGATGGCGGTGAGGGCAAGCGGCC
GATGCCCGGGATGCAGCAGCAGATGCCAACGCTACCTCCACCCTCGGTGTCCGCAACAGGACCCGCCCT
GGCCCTGGCCCTGGCCCCGGCCCGGGTCCCGGCCCGGCACCTCCAAATTACAGCAGGCCTCATGGTATGG
GAGGGCCCAACATGCCTCCCCAGGACCTCGGGCGTGCCCCCGGGATGCCAGGCCAGCCTCTGGAGG
GCCTCCAAGCCCTGGCCTGAAGGACCCATGGCGAATGCTGCTGCCCCACGAGCACCCCTCAGAAGCTG
ATTCCCCCGCAGCCAACGGGCGGCCCTTCCCCCGCGCCCTGCGTCCACCCCGCGCTCGCCCGTGA
TGCCACCGCAGACCCAGTCCCCCGGGCAGCCGGCCAGCCCGCGCCCATGGTGCCACTGCACCAGAAGCA
GAGCCGCATCACCCCATCCAGAAGCCGCGGGGCTCGACCTGTGGAGATCTGCAGGAGCGCGAGTAC
AGGCTGCAGGCTCGCATCGCACACCGAATTCAGGAACCTTGAAAACCTTCCCGGGTCCCTGGCCGGGATT

```

-continued

TGCGAACCAAGCGACCATTGAGCTCAAGGCCCTCAGGCTGCTGAACCTCCAGAGGCAGCTGCGCCAGGA
GGTGGTGGTGTGCTATGCGGAGGGACACAGCGCTGGAGACAGCCCTCAATGCTAAGGCCTACAAGCGCAGC
AAGCGCCAGTCCCTGCGGAGGCCCGCATCACTGAGAAGCTGGAGAAGCAGCAGAAGATCGAGCAGGAGC
GCAAGCGCCGGCAGAAGCACCAGGAATACCTCAATAGCATTCTCCAGCATGCCAAGGATTTCAAGGAATA
TCACAGATCCGTACAGGCAAAATCCAGAAGCTGACCAAGGCAGTGGCCACGTACCATGCCAACACGGAG
CGGGAGCAGAAGAAGAGAACGAGCGGATCGAGAAGGAGCGCATGCGGAGGCTCATGGCTGAAGATGAGG
AGGGGTACCGCAAGCTCATCGACCAGAAGAAGGACAAGCGCCTGGCCTACCTCTTGACGACAGACAGCA
GTACGTGGCTAACCTACGGAGCTGGTGCAGCAGCACAAGGCTGCCAGGTGCCAAGGAGAAAAAGAAG
AAAAAGAAAAAGAAGGAGCAGAAAATGCAGAAGGACAGACGCCCTGCCATTGGGCCGGATGGCGAGCCTC
TGGACGAGACCAGCCAGATGAGCGACCTCCCGGTGAAGGTGATCCACGTGGAGAGTGGGAAGATCCTCAC
AGGCACAGATGCCCCAAAGCGGGCAGCTGGAGGCCCTGGCTCGAGATGAACCCGGGGTATGAAGTAGCT
CCGAGGTCTGATAGTGAAGAAAGTGGCTCAGAAGAAGAGGAGGAGGAGGAGGAGAGCAGCCGAGG
CAGCACAGCCTCCACCTGCCCGTGGAGGAGAAGAAGAAGATTCCAGATCCAGACAGCGATGACGTCTC
TGAGGTGAGCGCGGCACATCATTGAGAATGCCAAGCAAGATGTCGATGATGAATATGGCGTGTCCAG
GCCCTTGACGTGGCCTGCAGTCTACTATGCCGTGGCCCATGCTGTCACTGAGAGAGTGGACAAGCAGT
CAGCGCTTATGGTCAATGGTGTCTCAAACAGTACCAGATCAAAGGTTTGGAGTGGCTGGTGTCCCTGTA
CAACAACAACCTGAACGCGCATCCTGGCCGACGAGATGGGCCTGGGGAAGACCATCCAGACCATCGCGCTC
ATCAGTACCTCATGGAGCACAAACGCATCAATGGGCCCTTCTCATCATCGTGCCTCTCTCAACGTGT
CCAACCTGGCGTACGAGTTTGACAAGTGGGCCCCCTCCGTGGTGAAGGTGTCTTACAAGGGATCCCCAGC
AGCAAGAGCGGCTTTGTCCCCAGCTCCGGAGTGGGAAGTTCAACGTCTTGCTGACGACGTACGAGTAC
ATCATCAAAGACAAGCACATCTCGCCAAGATCCGTGGAAGTACATGATTGTGGACGAAGGTACCCGCA
TGAAGAACCACCACTGCAAGCTGACGCAGGTGCTCAACACGCACTATGTGGACCCCGCCGCTGCTGCT
GACGGGCACACCGCTGCAGAACAGCTTCCCGAGCTCTGGGCGCTGCTCAACTTCTGCTGCCACCATC
TTCAAGAGCTGCAGCACCTTCGAGCAGTGGTTTAACGCACCCCTTGCCATGACCGGGGAAAAGGTGGACC
TGAATGAGGAGGAAACCATTCTCATCATCCGGCGTCTCCACAAAGTGTGCGGCCCTTCTTGCTCCGACG
ACTCAAGAAGGAAGTGCAGGCCAGTTGCCCGAAAAGGTGGAGTACGTCAATGAGTGCACATGCTCTGCG
CTGCAGCGAGTGTCTTACCGCCACATGCAGGCCAAGGGCGTGTGCTGACTGATGGCTCCGAGAAGGACA
AGAAGGGCAAAGGCGGCACCAAGACCTGATGAACACCATCATGCAGCTGCGGAAGATCTGCAACCACCC
CTACATGTTCCAGCACATCGAGGAGTCTTTTCCGAGCACTTGGGGTTCACTGGCGGCATGTTCCAAAGG
CTGGACCTGTACCGAGCCTCGGGTAAATTGAGCTTCTTGATAGAATTCTTCCAAACTCCGAGCAACCA
ACCACAAAGTGTGCTGTTCTGCCAAATGACCTCCCTCATGACCATCATGGAAGATTACTTTGCGTATCG
CGCCTTTAAATACCTCAGGCTTGATGGAACACGAAGGCGGAGGACCGGGCATGCTGCTGAAAACCTTC
AACGAGCCCGGCTCTGAGTACTTCTCTTCTGCTCAGCACCGGGCTGGGGGGCTCGGCCTGAACCTCC
AGTCGGCAGACACTGTGATCATTTTGTACAGCGACTGGAATCCTCACAGGACCTGCAAGCGCAGGACCG
AGCCCAACGCATCGGCAGCAGAACGAGGTGCGTGTGCTCCGCCTCTGCACCGTCAACAGCGTGGAGGAG
AAGATCCTAGCTGCAGCCAAGTACAAGCTCAACGTGGACCAGAAGGTGATCCAGGCCGCATGTTTCGACC
AGAAGTCTCCAGCCATGAGCGGCGCGCCTTCTGTCAGGCCATCCTGGAGCACGAGGAGCAGGATGAGAG
CAGACACTGCAGCACGGGCGAGCGGAGTGCCAGCTTCGCCCACACTGCCCCCTCCGCCAGCGGGCGTCAAC
CCCGACTTGGAGGAGCCACCTCTAAAGGAGGAAGACGAGGTGCCGACGACGAGACCGTCAACCAGATGA
TCGCCCCGCACGAGGAGGAGTTTGATCTGTTTCATGCGCATGGACCTGGACCGCAGGCGGAGGAGGCCG
CAACCCCAAGCGGAAGCGCGCCTCATGGAGGAGGACGAGCTCCCTCGTGGATCATCAAGGACGACGCG

- continued

GAGGTGGAGCGGCTGACCTGTGAGGAGGAGGAGAGAAGATGTTGGCCGTGGCTCCCGCCACCGCAAGG
 AGGTGGACTACAGCGACTCACTGACGGAGAAGCAGTGGCTCAAGAAAATTACAGGAAAAGATATCCATGA
 CACAGCCAGCAGTGTGGCAGCTGGGCTACAATTCCAGCGTGGCTTCAGTTCTGCACACGTGCGTCAAAG
 GCCATCGAGGAGGGCAGCTGGAGGAGATCGAAGAGGAGGTCCGGCAGAAGAAATCATCACGGAAGCGCA
 AGCGAGACAGCGACGCCGGCTCCTCCACCCGACCACCAGCACCCGAGCCGCGACAAGGACGACGAGAG
 CAAGAAGCAGAAGAAGCGCGGGCGGCCCTGCCGAGAACTCTCCCTAACCCACCCAACCTCACCAG
 AAGATGAAGAAGATTGTGGATCGCGTGATCAAGTACAAGGACAGCAGCAGTGGACGTGAGCTCAGCGAGG
 TCTTCATCCAGCTGCCCTCGCGAAAGGAGCTGCCCGAGTACTACGAGCTCATCCGCAAGCCCGTGGACTT
 CAAGAAGATAAAGGAGCGCATTTCGAACCAAGTACCGCAGCTCAACGACCTAGAGAAGGACGTGATG
 CTCTGTGCCAGAAGCAGCAGACCTTCAACCTGGAGGGCTCCCTGATCTATGAAGACTCCATCGTCTTGC
 AGTCGGTCTTACCAGCGTGGCGCAGAAAATCGAGAAGGAGGATGACAGTGAAGCGAGGAGAGTGAGGA
 GGAGGAAGAGGGCGAGGAGGAAGGCTCCGAATCCGAATCTCGTCCGTCAAAGTGAAGATCAAGCTTGGC
 CGGAAGGAGAAGGCACAGGACCGGTGAAGGGCGGCGCGCGCGGAGCCGAGGGTCCCGAGCCAAGC
 CGGTCTGTGAGTGACGATGACAGTGAGGAGGAACAAGAGGAGGACCGCTCAGGAAGTGGCAGCGAAGAAGA
 CTGAGCCCCGACATTCAGTCTCGACCCCGAGCCCTCGTTCAGAGCTGAGATGGCATAGGCCTTAGCA
 GTAACGGGTAGCAGCAGATGTAGTTTCAGACTTGGAGTAAACTGTATAAACAAGAATCTTCCATATT
 TATACAGCAGAGAAGCTGTAGACTGTTTGTGACTGGCCCTGTCTGGCATCAGTAGCATCTGTAACAGC
 ATTAAGTGTCTTAAGAGAGAGAGAGAGAATTCGAATGGGGAACACACGATACCTGTTTTCTTTTCC
 GTTGTGCGCAGTACTGTGCGCCGAGTTTGGAGTCACTGTAGTTAAGTGTGATGCATGTGCGTACCG
 TCCACTCTCTACTGTATTTATTGGACAGGTCAGACTCGCCGGGGGCCGCGAGGGTATGTCAGTGT
 CACTGGATGTCAAACAGTAATAAATAAACCAACAACAAACGACAGCCAAAAA

The term “BRG1” also refers to natural variants of the wild-type human BRG1 protein, such as proteins having at least 85% identity (e.g., 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.9% identity, or more) to an amino acid sequence of wild-type BRG1, which is set forth in SEQ ID NO: 2 (UniProt Accession No.: P51532; www.uniprot.org/uniprot/P51532.fasta).

SEQ ID NO: 2
 MSTPDPPLGGTTPRGPSPGPGSPGAMLGPSPGSPGSAHSMGSPGPG
 PSAGHIP IPTQGGPGYPQDNMHQMHKPMESMHEKMSDDPRYNQMGKMG
 RSGGHAGMGPPSPMDQHSQGYPSPLGGSEHASSVPASGSPSGPQMSS
 GPGGAPLDGADPQALGQQNRGPTPFNQNLHQLRAQIMAYKMLARGQPL
 PDHLQMAVQGKRPMPGMQQMPTLPPSVSATGPGPGPGPGPGGPA
 PPNYSRPHMGMPGNMPPGPGSPVPPGMPGQPPGPPKPWPEGPMANAAA
 PTSTPQKLIPPQPTGRPSAPPAPVPPAASPVMPPTQSPGQPAQPAPMV
 PLHQKQSRITPIQKPRGLDPVEILQEREYRLQARIAHRIQELNLPGL
 AGDLRTKATIELKALRLNLFQRQLRQEVVCMRRDTALETALNAKAYKR
 SKRQSLREARITEKLEKQKI EQERKRRQKHQEYLNLSILQHAKEKEYH
 RSVTGKIQLTKAVATYHANTEREQKENERIEKERMRLMADEEGYR
 KLIDQKDKRLAYLLQQTDEYVANLTENVRHKAQVAKEKKKKKKKK

- continued

AENAEGQTPAIGPDGEPLDETSQMSDLPVKVIHVESGKILGTDPKAG
 40 QLEAWLEMPNGYEVAPRSDSEESGSEEEEEEEQQAAPPTLPVEE
 KKKIPDPDSDDVSEVDARHIIENAKQDVDDDEYGVSQLARGLQSYAYA
 HAVTERVDKQSALMVNGLKQYQIKGLEWLVSLYNNNLNGILADEMGLG
 45 KTIQTIALITYLMEHKRINGPFLIIVPLSTLSNWAYEFDKWPVSVKVS
 YKGS PAARRAFVPQLRSGKENVLLTTYEYIIKDKHILAKIRWKYMIVDE
 GHRMKNHHCKLTQVLNTHYVAPRRLLTGTPLQNKLPWALLNELLPT
 50 IFKSCSTFEQWENAPFAMTGEKVDLNEETILIIIRLHKVLRPFLRLRL
 KKEVEAQLPEKVEYVIKCDMSALQVRLYRHMQAKGVLLTDGSEKDKKGK
 GGTKILMNTIMQLRKICNHPYMFQHI EESFSEHLGFTGGIVQGLDLYRA
 55 SGKFELLDRI LPKLRATNHKVLLFCQMTSLMTIMEDYFAYRGEKYLRLD
 GITKAEDRGMLLKT FNPGSEYFIFLLSTRAGGLGLNLQSADTVIIFDS
 DWNPHQDLQAQDRAHRIQQNEVRVRLRCTVNSVEEKILAAKYKLNVD
 QKVIQAGMFDQKSSSHERRAFLQAILEHEEQDES RHCTSGSGSASFAHT
 60 APPPAGVNPDL EEPPLKEEDEV PDET VNMQIARHEEFDLFMRMDLDR
 RREEARNPKRKPRLMEEDLP SWIIKDDAEVERLICEEEEEKMFGRGSR
 HRKEVDYSDSLTEKQWLKAI EEGTLEEEI EEEVRQKKSSRKRKRSDAGS
 65 STPTTSTRSRDKDDESKKQKKRGRPPAEKLSNPNNLT KMKKIVDAVI

31

-continued

KYKDSSSGRQLSEVFIQLPSRKELPEYYELIRKPVDFKKIKERIRNHKY
 RSLNDLEKDVMLLCQNAQTFNLEGLIYEDSIVLQSVFTSVRQKIEKED
 DSEGESEEEEEEGESESRSVKVKKILGRKEKAQDRLKGRRRPS
 RGSRAKPVVDDDDSEEEQEEDRSGSGSEED .

32

As used herein, the term “BRM” refers to probable global transcription activator SNF2L2. BRM is a component of the BAF complex, a SWI/SNF ATPase chromatin remodeling complex. Human BRM is encoded by the SMARCA2 gene on chromosome 9, a nucleic acid sequence of which is set forth in SEQ ID NO: 3 (GenBank Accession No.: NM_003070.4, www.ncbi.nlm.nih.gov/nuccore/NM_003070.4?report=fasta).

SEQ ID NO: 3

GCGTCTTCCGCGCCCCGCGAGGAGGCGAGGGTGGGACGCTGGGCGGAGCCCGAGTTTAGGAAGAGGAGG
 GGACGGCTGTCATCAATGAAGTCATATTCATAATCTAGTCCTCTCTCCCTCTGTTTCTGTACTCTGGGTG
 ACTCAGAGAGGGAAGAGATTAGCCAGCACACTCCTCGCGAGCAAGCATTACTCTACTGACTGGCAGAGA
 CAGGAGAGGTAGATGTCCACGCCACAGACCCCTGGTGCGATGCCCCACCCAGGGCCTTCGCGGGGCGCTG
 GGCTTCCCCTGGGCCAATTCTTGGGCCTAGTCCAGGACCAGGACCATCCCCAGGTTCCGTCCACAGCAT
 GATGGGGCCAAGTCTTGACCTCCAAGTGTCTCCCATCTATGCCGACGATGGGGTCCACAGACTTCCCA
 CAGGAAGGCATGCATCAATGCATAAGCCCATCGATGGTATACATGACAAGGGGATTGTAGAAGACATCC
 ATTGTGGATCCATGAAGGGCACTGGTATGCGACCACCTCACCAGGCATGGGCGCTCCCGAGAGTCCAAT
 GGATCAACACAGCCAAGTTATATGTACCACACCCATCTCCATTAGGAGCCCCAGAGCACGTCTCCAGC
 CCTATGTCTGGAGGAGGCCAACTCCACCTCAGATGCCACCAAGCCAGCCGGGGGCGCTCATCCAGGTG
 ATCCGCAGGCCATGAGCCAGCCCAACAGAGGTCCCTCACCTTTCAGTCTGTCCAGCTGCATCAGCTTCG
 AGCTCAGATTTTAGCTTATAAATGCTGGCCGAGGCCAGCCCCCTCCCGAAACGCTGCAGCTTGAGTC
 CAGGGGAAAAGGACGTTGCCTGGCTTGAGCAACAACAGCAGCAGCAACAGCAGCAGCAGCAGCAGCAGC
 AGCAGCAGCAGCAGCAGCAACAGCAGCCGAGCAGCAGCCGCGCAACACAGCAGCAGCAACAACAGCA
 GCCGGCCCTTGTTAACTACAACAGACCATCTGGCCCGGGCCGAGCTGAGCGGCCCGAGCACCCGCGAG
 AAGCTGCCGTGCCCGCGCCCGCGCGGCCCTCGCCCGCGCCCCCGCAGCCGCGCAGCCGCGCGCGG
 CCGCAGTGCCCGGCCCTCAGTGCCGCGAGCCGCCCCGGGGCAGCCCTCGCCGCTCTCCAGCTGCAGCA
 GAAGCAGAGCCGCATCAGCCCCATCCAGAAACCGCAAGGCCTGGACCCCGTGAAATTCTGCAAGAGCGG
 GAATACAGACTTCAGGCCCGCATAGCTCATAGGATACAAGAACTGGAAAATCTGCCTGGCTCTTTGCCAC
 CAGATTTAAGAACCAAGCAACCGTGGAATAAAAGCACTTCGGTTACTCAATTTCCAGCGTCAGCTGAG
 ACAGGAGGTGGTGGCTGCATGCGCAGGGACACGACCTGGAGACGGCTCTCAACTCCAAAGCATACAAA
 CGGAGCAAGCGCCAGACTCTGAGAGAAGCTCGCATGACCGAGAAGCTGGAGAAGCAGCAGAAGATTGAGC
 AGGAGAGGAAACGCCGTGAGAAACACAGGAATACCTGAACAGTATTTTGCAACATGCAAAAGATTTTAA
 GGAATATCATCGGTCTGTGCGCGAAAGATCCAGAAGCTCTCAAAGCAGTGGAACCTTGGCATGCCAAC
 ACTGAAAGAGAGCAGAAGAAGGAGACAGAGCGGATTGAAAAGGAGAGAATGCGGCAGCTGATGGCTGAAG
 ATGAGGAGGGTTATAGAAAACCTGATTGATCAAAAGAAAGACAGGCGTTTAGCTTACCTTTTGAGCAGAC
 CGATGAGTATGTAGCCAATCTGACCAATCTGGTTTGGGAGCACAAGCAAGCCAGGCAGCCAAAGAGAAG
 AAGAAGAGGAGGAGGAGGAAGAAGAAGGCTGAGGAGAATGCAGAGGTGGGGAGTCTGCCCTGGGACCGG
 ATGAGAGAGCCATAGATGAGAGCAGCCAGATGAGTGACCTCCCTGTCAAAGTGACTCACACAGAAACCGG
 CAAGGTTCTGTTCCGACCAGAAGCACCCAAAGCAAGTCAGCTGGACGCTGGCTGGAAATGAATCTGGT
 TATGAAGTTGCCCTAGATCTGACAGTGAAGAGAGTGATTCTGATTATGAGGAAGAGGATGAGGAAGAAG
 AGTCCAGTAGGCAGGAACCGAAGAGAAAATACTCTGGATCCAAATAGCGAAGAAGTTTCTGAGAAGGA
 TGCTAAGCAGATCATTTAGACAGCTAAGCAAGACGTGGATGATGAATACAGCATGCAGTACAGTGCCAGG
 GGCTCCAGTCTTACTACACCGTGGCTCATGCCATCTCGGAGAGGGTGGAGAAACAGTCTGCCCTCCTAA

- continued

TTAATGGGACCCTAAAGCATTACCAGCTCCAGGGCTTGAATGGATGGTTTCCCTGTATAATAACAACTT
GAACGGAATCTTAGCCGATGAAATGGGGCTTGGAAGACCATAACAGACCATTGCACTCATCACTTATCTG
ATGGAGCACAAAAGACTCAATGGCCCTATCTCATCATTTGCCCTTCGACTCTATCTAACTGGACAT
ATGAATTTGACAAATGGGCTCCTTCTGTGGTGAAGATTTCTTACAAGGGTACTCCTGCCATGCGTCGCTC
CCTTGTCCTCCAGCTACGGAGTGGCAAATTCATGTCTCTTGACTACTTATGAGTATATTATAAAAGAC
AAGCACATCTTTGCAAAGATTCGGTGGAAATACATGATAGTGGACGAAGGCCACCGAATGAAGAATCACC
ACTGCAAGCTGACTCAGGTCTTGAACACTCACTATGTGGCCCCAGAAAGGATCCTCTTGACTGGGACCCC
GCTGCAGAATAAGCTCCCTGAACCTTGGGCCCTCCTCAACTTCCTCTCCCAACAATTTTAAAGAGCTGC
AGCACATTTGAACAATGGTTCAATGCTCCATTTGCCATGACTGGTGAAAGGGTGGACTTAAATGAAGAAG
AAACTATATTGATCATCAGGCGTCTACATAAGGTGTTAAGACCATTTTACTAAGGAGACTGAAGAAAGA
AGTTGAATCCAGCTTCCCGAAAAAGTGAATATGTGATCAAGTGTGACATGTGAGCTCTGCAGAAGATT
CTGTATCGCCATATGCAAGCCAAAGGGATCCTTCTCACAGATGGTTCTGAGAAAGATAAGAAGGGGAAAG
GAGGTGCTAAGACACTTATGAACACTATTATGCAGTTGAGAAAAATCTGCAACCACCCATATATGTTTCA
GCACATTGAGGAATCCTTTGCTGAACACCTAGGCTATTCAAATGGGGTCATCAATGGGGCTGAACGTAT
CGGGCTCAGGGAAGTTTGAGCTGCTTGATCGTATTCTGCCAAAATTGAGAGCGACTAATCACCAGGTGC
TGCTTTTCTGCCAGATGACATCTCTCATGACCATCATGGAGGATTATTTTGCTTTTCGGAACCTCCTTTA
CCTACGCCTTGATGCGACCCACCAAGTCTGAAGATCGTGCTGCTTTGCTGAAGAAATTCATGAACCTGGA
TCCCAGTATTTTCAATTTCTTGCTGAGCACAAAGAGCTGGTGGCCTGGGCTTAAATCTTCAGGCAGCTGATA
CAGTGGTCATCTTTGACAGCGACTGGAATCCTCATCAGGATCTGCAGGCCAAGACCGAGCTCACCGCAT
CGGCAGCAGAAACGAGGTCCGGTACTGAGGCTCTGTACCGTGAACAGCGTGGAGGAAAAGATCCTCGCG
GCCGCAAATACAAGCTGAACGTGGATCAGAAAGTGATCCAGGCGGGCATGTTTGACCAAAGTCTTCAA
GCCACGAGCGGAGGGCATTCTGCAGGCCATCTTGGAGCATGAGGAGGAAAATGAGGAAGAAGATGAAGT
ACCGGACGATGAGACTCTGAACCAAATGATTGCTCGACGAGAAGAAGAAATTTGACCTTTTATGCGGATG
GACATGGACCGCGGAGGGAAGATGCCCGGAACCGAAACGGAAGCCCCGTTTAAATGGAGGAGGATGAGC
TGCCCTCCTGGATCATTAAAGGATGACGCTGAAGTAGAAAGGCTCACCTGTGAAGAAGAGGAGGAGAAAAT
ATTTGGGAGGGGTCCCGCCAGCGCGTGACGTGGACTACAGTGACGCCCTCACGGAGAAGCAGTGGCTA
AGGGCCATCGAAGACGGCAATTTGAGGAAATGGAAGAGGAAGTACGGCTTAAAGAGCGAAAAAGACGAA
GAAATGTGGATAAAGATCCTGCAAAAGAAAGATGTGAAAAAGCTAAGAAGAGAAGAGCCGCCCTCCCGC
TGAGAAACTGTACCAAATCCCCCAAACCTGACAAAGCAGATGAACGCTATCATCGATACTGTGATAAAC
TACAAAGATAGGTGTAACGTGGAGAAGGTGCCAGTAATTCTCAGTTGGAATAGAAGGAAACAGTTTCAG
GGCGACAGCTCAGTGAAGTCTTCATTCAGTTACCTTCAAGGAAAGAATTACCAGAATACTATGAATTAAT
TAGGAAGCCAGTGGATTTCAAAAAATAAAGGAAAGGATTCGTAATCATAAGTACCGGAGCCTAGGCGAC
CTGGAGAAGGATGTATGCTTCTCTGTCAACGCTCAGACGTTCAACCTGGAGGGATCCCAGATCTATG
AAGACTCCATCGTCTTACAGTCAGTGTTTAAGAGTGCCCGGCAGAAAATTGCCAAGAGGAAGAGAGTGA
GGATGAAAGCAATGAAGAGGAGGAAGGAAGATGAAGAAGAGTCAGAGTCCGAGGCAAATCAGTCAAG
GTGAAAAATTAAGCTCAATAAAAAAGATGACAAAGGCCGGGACAAAGGAAAGGCAAGAAAAGGCCAAATC
GAGGAAAAGCCAAACCTGTAGTGAGCGATTTTGACAGCGATGAGGAGCAGGATGAACGTGAACAGTCAGA
AGGAAGTGGGACGGATGATGAGTGATCAGTATGGACCTTTTCTCTGGTAGAACTGAATTCCTTCTCTCC
CTGTCTCATTTCTACCCAGTGAGTTTCAATTTGTATATAGGCACTGGGTTGTTTCTATATCATCATCGTCT
ATAAACTAGCTTTAGGATAGTGCCAGACAAACATATGATATCATGGTGTA AAAACACACATACACAA
ATATTTGTAACATATTGTGACCAAATGGGCCTCAAAGATTGAGATTGAAACAAACAAAAGCTTTTGATG

- continued

GAAAATATGTGGGTGGATAGTATATTTCTATGGGTGGGTCTAATTTGGTAACGGTTTGATTGTGCCTGGT
 TTTATCACCTGTTTCTAGATGAGAAGATTTTTGTCTTTGTAGCACTGATAACCAGGAGAAGCCATTAAAG
 CCACTGGTTATTTTATTTTTCATCAGGCAATTTTCGAGGTTTTTATTGTTTCGGTATTGTTTTTTTACAC
 TGTGTACATATAAGCACTTTAATAGGTGATAAATGTACAGTAGTTAGATTTCACCTGCATATACATT
 TTCCATTTTATGCTCTATGATCTGAACAAAAGCTTTTGAATTGTATAAGATTATGTCTACTGTAAACA
 TTGCTTAATTTTTTTGCTCTTGATTAAAAAAGTTTTGTTGAAAGCGCTATTGAATATTGCAATCTAT
 ATAGTGATTGGATGGCTCTTTTGTCAACCCTGATCTCCTATGTTACCAATGTGTATCGTCTCCTTCTCC
 CTAAAGTGACTTAATCTTTGCTTTCTTTGCACAATGTCTTTGGTTGCAAGTCATAAGCCTGAGGCAAAT
 AAAATCCAGTAATTTGAAGAATGTGGTGTGGTGTCTTCTTAATAAGAAATAATTAGCTTGACAAA.

The term “BRM” also refers to natural variants of the wild-type human BRM protein, such as proteins having at least 85% identity (e.g., 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.9% identity, or more) to an amino acid sequence of wild-type BRM, which is set forth in SEQ ID NO: 4 (Uniprot Accession No.: P51531; www.uniprot.org/uniprot/P51531.fasta).

SEQ ID NO: 4

MSTPTDPGAMPHPGSPGPGSPGPGPILGSPGPGSPGSPVHSMGPGSPGP
 PSVSHPMPTMGSTDFPQEGMHQMHKPIDIGHDKGIVEDIHCGSMKGTGMR
 PPHPGMGPQSPMDQHSQGYMSPHPSPLGAPEHVSSPMSGGPTTPQMP
 SPQALIPGDPQAMSPNRPSPFSPVQLHLRAQILAYKMLARGQPLPE
 TLQLAVQKRTLPGLQQQQQQQQQQQQQQQQQQQQPQQPPQPQTQQ
 QQQPALVYNRPSPGPELSPSTPQKLPVPAPGGRPSAPPAAQPPAA
 AVPGSPVPQAPGQSPVLQLQQKQSRISPIQKPQGLDPVEILQEREYRL
 QARIAHRIQELNLPGLPDLRTKATVELKALRLNFRQRLRQEVVACM
 RRDTTLETALNSKAYKRSKRQTLREARMTEKLEKQKIEQERKRQKHQ
 YLNSILQHAQDFKEYHRSVAGKIQKLSKAVATWHANTEREQKKETERIEK
 ERMRLMAEDEEGYRLIDQKDRRLAYLLQQTDEYVANLTNLVWEHKQA
 QAAKEKKKRRRRKKAEENAEAGESALPGDGEPIDESSQMSDLPVKVTH
 ETGKVLFGPEAPKASQLDAWLEMNPGYEVAPRSDSESDSDYEEDEEEEE
 SSRQETEEKILLDPNSEEVSEKDAKQIIETAKQDVDEYSMQYSARGSQS
 YVTVAHAISERVEKQSALLINGTLKHYQLQGLEWMVSLYNNNLNGILADE
 MGLGKTIQTIATITYLMEHKRLNGPYLIIVPLSTLSNWTYEFKWNAPSVV
 KISYKGTAMRRSLVPQLRSGKENVLLTTYEYIIKDKHILAKIRWKYMI
 DEGRMKNHCKLIQVLNTHYVAPRRILLIGTLPQNKLPWALLNFFLLP
 TIFKSCSTFEQWFNAPPAMTGERVDLNEETILIIIRLHKVLRPFLRLRL
 KKEVESQLPEKVEYVIKCDMSALQKILYRHMQAKGILLTDGSEKDKKGG
 GAKTLMNTIMQLRKICNHYPMFQHIIESFAEHLGYSNGVINGAELYRAG
 KFELLDRIPLKLRATNHRVLLFCQMTSLMTIMEDYFAFRNPLYLRDGLT
 KSEDRAALLKKENEPGSQYIFLLSTRAGGLNLQAADTWIFSDWNPH
 QDLQAQDRAHRIGQQNEVRVLRLLCTVNSVEEKILAAKYKLNVDQKVIQA

- continued

GMFDQKSSSHERRAFLQAILLEHEEENEEDEVPPDETNLQMIARREEEFD
 20 LFMRMMDRRREDARNPKRKPRLMEDELPSWIIKDDAEVERLTCEEEEE
 KIFGRGSRQRDQVSDALTEKQWLRAIEDGNLEEMEEVRLKKRKRNRN
 VDKDPKEDVEKAKRRRPPAEKLSNPPKLTQKMNAIIDTVINYKDR
 25 NVEKVPNSQLEIEGNSSGRQLSEVFIQLPSRKELPEYYELIRKPVDFKK
 IKERIRNHKYSRLGDLEKDVMLLCHNAQTFNLEGSQIYEDSIVLQSVFKS
 ARQKIAKEEESDESNEEEEEDEEESSEAKSVKVKIKLNKDDKGRDK
 30 GKGGKRPNGKAKPVVSDFDSDDEEQDEREQSEGSGTDDE.

As used herein, the term “degrader” refers to a small molecule compound including a degradation moiety, wherein the compound interacts with a protein (e.g., BRG1 and/or BRM) in a way which results in degradation of the protein, e.g., binding of the compound results in at least 5% reduction of the level of the protein, e.g., in a cell or subject.

As used herein, the term “degradation moiety” refers to a moiety whose binding results in degradation of a protein, e.g., BRG1 and/or BRM. In one example, the moiety binds to a protease or a ubiquitin ligase that metabolizes the protein, e.g., BRG1 and/or BRM.

By “determining the level of a protein” is meant the detection of a protein, or an mRNA encoding the protein, by methods known in the art either directly or indirectly. “Directly determining” means performing a process (e.g., performing an assay or test on a sample or “analyzing a sample” as that term is defined herein) to obtain the physical entity or value. “Indirectly determining” refers to receiving the physical entity or value from another party or source (e.g., a third-party laboratory that directly acquired the physical entity or value). Methods to measure protein level generally include, but are not limited to, western blotting, immunoblotting, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), immunoprecipitation, immunofluorescence, surface plasmon resonance, chemiluminescence, fluorescent polarization, phosphorescence, immunohistochemical analysis, matrix-assisted laser desorption/ionization time-of-flight (MALDI-TOF) mass spectrometry, liquid chromatography (LC)-mass spectrometry, microcytometry, microscopy, fluorescence activated cell sorting (FACS), and flow cytometry, as well as assays based on a property of a protein including, but not limited to, enzymatic activity or interaction with other protein partners. Methods to measure mRNA levels are known in the art.

By “modulating the activity of a BAF complex” is meant altering the level of an activity related to a BAF complex

37

(e.g., GBAF), or a related downstream effect. The activity level of a BAF complex may be measured using any method known in the art, e.g., the methods described in Kadoch et al, Cell 153:71-85 (2013), the methods of which are herein incorporated by reference.

By “reducing the activity of BRG1 and/or BRM” is meant decreasing the level of an activity related to a BRG1 and/or BRM, or a related downstream effect. A non-limiting example of inhibition of an activity of BRG1 and/or BRM is decreasing the level of a BAF complex (e.g., GBAF) in a cell. The activity level of BRG1 and/or BRM may be measured using any method known in the art. In some embodiments, an agent which reduces the activity of BRG1 and/or BRM is a small molecule BRG1 and/or BRM inhibitor. By “reducing the level of BRG1 and/or BRM” is meant decreasing the level of BRG1 and/or BRM in a cell or subject. The level of BRG1 and/or BRM may be measured using any method known in the art.


As used herein, the term “inhibiting BRG and/or BRM” refers to blocking or reducing the level or activity of the ATPase catalytic binding domain or the bromodomain of the protein. BRG1 and/or BRM inhibition may be determined using methods known in the art, e.g., a BRG and/or BRM ATPase assay, a Nano DSF assay, or a BRG1 and/or BRM Luciferase cell assay.

By “level” is meant a level of a protein, or mRNA encoding the protein, as compared to a reference. The reference can be any useful reference, as defined herein. By a “decreased level” or an “increased level” of a protein is meant a decrease or increase in protein level, as compared to a reference (e.g., a decrease or an increase by about 5%, about 10%, about 15%, about 20%, about 25%, about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, about 60%, about 65%, about 70%, about 75%, about 80%, about 85%, about 90%, about 95%, about 100%, about 150%, about 200%, about 300%, about 400%, about 500%, or more; a decrease or an increase of more than about 10%, about 15%, about 20%, about 50%, about 75%, about 100%, or about 200%, as compared to a reference; a decrease or an increase by less than about 0.01-fold, about 0.02-fold, about 0.1-fold, about 0.3-fold, about 0.5-fold, about 0.8-fold, or less; or an increase by more than about 1.2-fold, about 1.4-fold, about 1.5-fold, about 1.8-fold, about 2.0-fold, about 3.0-fold, about 3.5-fold, about 4.5-fold, about 5.0-fold, about 10-fold, about 15-fold, about 20-fold, about 30-fold, about 40-fold, about 50-fold, about 100-fold, about 1000-fold, or more). A level of a protein may be expressed in mass/vol (e.g., g/dL, mg/mL, µg/mL, ng/mL) or percentage relative to total protein or mRNA in a sample.

As used herein, the term “inhibitor” refers to any agent which reduces the level and/or activity of a protein (e.g., BRG1 and/or BRM). Non-limiting examples of inhibitors include small molecule inhibitors, degraders, antibodies, enzymes, or polynucleotides (e.g., siRNA).

As used herein, the term “LXS196,” refers to the PKC inhibitor having the structure:

38



or a pharmaceutically acceptable salt thereof.

As used herein, the terms “effective amount,” “therapeutically effective amount,” and “a sufficient amount” of an agent that reduces the level and/or activity of BRG1 and/or BRM (e.g., in a cell or a subject) described herein refer to a quantity sufficient to, when administered to the subject, including a human, effect beneficial or desired results, including clinical results, and, as such, an “effective amount” or synonym thereto depends on the context in which it is being applied. For example, in the context of treating cancer, it is an amount of the agent that reduces the level and/or activity of BRG1 and/or BRM sufficient to achieve a treatment response as compared to the response obtained without administration of the agent that reduces the level and/or activity of BRG1 and/or BRM. The amount of a given agent that reduces the level and/or activity of BRG1 and/or BRM described herein that will correspond to such an amount will vary depending upon various factors, such as the given agent, the pharmaceutical formulation, the route of administration, the type of disease or disorder, the identity of the subject (e.g., age, sex, and/or weight) or host being treated, and the like, but can nevertheless be routinely determined by one of skill in the art. Also, as used herein, a “therapeutically effective amount” of an agent that reduces the level and/or activity of BRG1 and/or BRM of the present disclosure is an amount which results in a beneficial or desired result in a subject as compared to a control. As defined herein, a therapeutically effective amount of an agent that reduces the level and/or activity of BRG1 and/or BRM of the present disclosure may be readily determined by one of ordinary skill by routine methods known in the art. Dosage regimen may be adjusted to provide the optimum therapeutic response.

The term “inhibitory RNA agent” refers to an RNA, or analog thereof, having sufficient sequence complementarity to a target RNA to direct RNA interference. Examples also include a DNA that can be used to make the RNA. RNA interference (RNAi) refers to a sequence-specific or selective process by which a target molecule (e.g., a target gene, protein, or RNA) is down-regulated. Generally, an interfering RNA (“iRNA”) is a double-stranded short-interfering RNA (siRNA), short hairpin RNA (shRNA), or single-stranded micro-RNA (miRNA) that results in catalytic degradation of specific mRNAs, and also can be used to lower or inhibit gene expression.

The terms “short interfering RNA” and “siRNA” (also known as “small interfering RNAs”) refer to an RNA agent, preferably a double-stranded agent, of about 10-50 nucleotides in length, the strands optionally having overhanging ends comprising, for example 1, 2 or 3 overhanging nucleotides (or nucleotide analogs), which is capable of directing or mediating RNA interference. Naturally-occurring siR-

NAs are generated from longer dsRNA molecules (e.g., >25 nucleotides in length) by a cell's RNAi machinery (e.g., Dicer or a homolog thereof).

The term "shRNA", as used herein, refers to an RNA agent having a stem-loop structure, comprising a first and second region of complementary sequence, the degree of complementarity and orientation of the regions being sufficient such that base pairing occurs between the regions, the first and second regions being joined by a loop region, the loop resulting from a lack of base pairing between nucleotides (or nucleotide analogs) within the loop region.

The terms "miRNA" and "microRNA" refer to an RNA agent, preferably a single-stranded agent, of about 10-50 nucleotides in length, preferably between about 15-25 nucleotides in length, which is capable of directing or mediating RNA interference. Naturally-occurring miRNAs are generated from stem-loop precursor RNAs (i.e., pre-miRNAs) by Dicer. The term "Dicer" as used herein, includes Dicer as well as any Dicer ortholog or homolog capable of processing dsRNA structures into siRNAs, miRNAs, siRNA-like or miRNA-like molecules. The term microRNA ("miRNA") is used interchangeably with the term "small temporal RNA" ("stRNA") based on the fact that naturally-occurring miRNAs have been found to be expressed in a temporal fashion (e.g., during development).

The term "antisense," as used herein, refers to a nucleic acid comprising a polynucleotide that is sufficiently complementary to all or a portion of a gene, primary transcript, or processed mRNA, so as to interfere with expression of the endogenous gene (e.g., BRG1 and/or BRM). "Complementary" polynucleotides are those that are capable of base pairing according to the standard Watson-Crick complementarity rules. Specifically, purines will base pair with pyrimidines to form a combination of guanine paired with cytosine (G:C) and adenine paired with either thymine (A:T) in the case of DNA, or adenine paired with uracil (A:U) in the case of RNA. It is understood that two polynucleotides may hybridize to each other even if they are not completely complementary to each other, provided that each has at least one region that is substantially complementary to the other.

The term "antisense nucleic acid" includes single-stranded RNA as well as double-stranded DNA expression cassettes that can be transcribed to produce an antisense RNA. "Active" antisense nucleic acids are antisense RNA molecules that are capable of selectively hybridizing with a primary transcript or mRNA encoding a polypeptide having at least 80% sequence identity (e.g., 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.9% identity, or more) with the targeted polypeptide sequence (e.g., a BRG1 and/or BRM polypeptide sequence). The antisense nucleic acid can be complementary to an entire coding strand, or to only a portion thereof. In some embodiments, an antisense nucleic acid molecule is antisense to a "coding region" of the coding strand of a nucleotide sequence. The term "coding region" refers to the region of the nucleotide sequence comprising codons that are translated into amino acid residues. In some embodiments, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence. The term "noncoding region" refers to 5' and 3' sequences that flank the coding region that are not translated into amino acids (i.e., also referred to as 5' and 3' untranslated regions). The antisense nucleic acid molecule can be complementary to the entire coding region of mRNA, or can be antisense to only a portion of the coding or noncoding region of an mRNA. For example, the antisense oligonucleotide can be complementary to the region sur-

rounding the translation start site. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides in length.

"Percent (%) sequence identity" with respect to a reference polynucleotide or polypeptide sequence is defined as the percentage of nucleic acids or amino acids in a candidate sequence that are identical to the nucleic acids or amino acids in the reference polynucleotide or polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid or amino acid sequence identity can be achieved in various ways that are within the capabilities of one of skill in the art, for example, using publicly available computer software such as BLAST, BLAST-2, or Megalign software. Those skilled in the art can determine appropriate parameters for aligning sequences, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For example, percent sequence identity values may be generated using the sequence comparison computer program BLAST. As an illustration, the percent sequence identity of a given nucleic acid or amino acid sequence, A, to, with, or against a given nucleic acid or amino acid sequence, B, (which can alternatively be phrased as a given nucleic acid or amino acid sequence, A that has a certain percent sequence identity to, with, or against a given nucleic acid or amino acid sequence, B) is calculated as follows:

$$100 \text{ multiplied by (the fraction } X/Y)$$

where X is the number of nucleotides or amino acids scored as identical matches by a sequence alignment program (e.g., BLAST) in that program's alignment of A and B, and where Y is the total number of nucleic acids in B. It will be appreciated that where the length of nucleic acid or amino acid sequence A is not equal to the length of nucleic acid or amino acid sequence B, the percent sequence identity of A to B will not equal the percent sequence identity of B to A.

The term "pharmaceutical composition," as used herein, represents a composition containing a compound described herein formulated with a pharmaceutically acceptable excipient, and manufactured or sold with the approval of a governmental regulatory agency as part of a therapeutic regimen for the treatment of disease in a mammal. Pharmaceutical compositions can be formulated, for example, for oral administration in unit dosage form (e.g., a tablet, capsule, caplet, gelcap, or syrup); for topical administration (e.g., as a cream, gel, lotion, or ointment); for intravenous administration (e.g., as a sterile solution free of particulate emboli and in a solvent system suitable for intravenous use); or in any other pharmaceutically acceptable formulation.

A "pharmaceutically acceptable excipient," as used herein, refers any ingredient other than the compounds described herein (for example, a vehicle capable of suspending or dissolving the active compound) and having the properties of being substantially nontoxic and non-inflammatory in a patient. Excipients may include, for example: antiadherents, antioxidants, binders, coatings, compression aids, disintegrants, dyes (colors), emollients, emulsifiers, fillers (diluents), film formers or coatings, flavors, fragrances, glidants (flow enhancers), lubricants, preservatives, printing inks, sorbents, suspending or dispersing agents, sweeteners, and waters of hydration. Exemplary excipients include, but are not limited to: butylated hydroxytoluene (BHT), calcium carbonate, calcium phosphate (dibasic), calcium stearate, croscarmellose, crosslinked polyvinyl pyrrolidone, citric acid, crospovidone, cysteine, ethylcellulose,

gelatin, hydroxypropyl cellulose, hydroxypropyl methylcellulose, lactose, magnesium stearate, maltitol, mannitol, methionine, methylcellulose, methyl paraben, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, povidone, pregelatinized starch, propyl paraben, retinyl palmitate, shellac, silicon dioxide, sodium carboxymethyl cellulose, sodium citrate, sodium starch glycolate, sorbitol, starch (corn), stearic acid, sucrose, talc, titanium dioxide, vitamin A, vitamin E, vitamin C, and xylitol.

As used herein, the term “pharmaceutically acceptable salt” means any pharmaceutically acceptable salt of the compound of any of the compounds described herein. For example, pharmaceutically acceptable salts of any of the compounds described herein include those that are within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and animals without undue toxicity, irritation, allergic response and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well known in the art. For example, pharmaceutically acceptable salts are described in: Berge et al., *J. Pharmaceutical Sciences* 66:1-19, 1977 and in *Pharmaceutical Salts: Properties, Selection, and Use*, (Eds. P. H. Stahl and C. G. Wermuth), Wiley-VCH, 2008. The salts can be prepared in situ during the final isolation and purification of the compounds described herein or separately by reacting a free base group with a suitable organic acid.

The compounds described herein may have ionizable groups so as to be capable of preparation as pharmaceutically acceptable salts. These salts may be acid addition salts involving inorganic or organic acids or the salts may, in the case of acidic forms of the compounds described herein, be prepared from inorganic or organic bases. Frequently, the compounds are prepared or used as pharmaceutically acceptable salts prepared as addition products of pharmaceutically acceptable acids or bases. Suitable pharmaceutically acceptable acids and bases and methods for preparation of the appropriate salts are well-known in the art. Salts may be prepared from pharmaceutically acceptable non-toxic acids and bases including inorganic and organic acids and bases. Representative acid addition salts include acetate, adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptonate, glycerophosphate, hemisulfate, heptonate, hexanoate, hydrobromide, hydrochloride, hydroiodide, 2-hydroxyethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, toluenesulfonate, undecanoate, and valerate salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, and magnesium, as well as nontoxic ammonium, quaternary ammonium, and amine cations, including, but not limited to ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, and ethylamine.

By a “reference” is meant any useful reference used to compare protein or mRNA levels. The reference can be any sample, standard, standard curve, or level that is used for comparison purposes. The reference can be a normal reference sample or a reference standard or level. A “reference sample” can be, for example, a control, e.g., a predetermined negative control value such as a “normal control” or a prior sample taken from the same subject; a sample from a normal

healthy subject, such as a normal cell or normal tissue; a sample (e.g., a cell or tissue) from a subject not having a disease; a sample from a subject that is diagnosed with a disease, but not yet treated with a compound described herein; a sample from a subject that has been treated by a compound described herein; or a sample of a purified protein (e.g., any described herein) at a known normal concentration. By “reference standard or level” is meant a value or number derived from a reference sample. A “normal control value” is a pre-determined value indicative of non-disease state, e.g., a value expected in a healthy control subject. Typically, a normal control value is expressed as a range (“between X and Y”), a high threshold (“no higher than X”), or a low threshold (“no lower than X”). A subject having a measured value within the normal control value for a particular biomarker is typically referred to as “within normal limits” for that biomarker. A normal reference standard or level can be a value or number derived from a normal subject not having a disease or disorder (e.g., cancer); a subject that has been treated with a compound described herein. In preferred embodiments, the reference sample, standard, or level is matched to the sample subject sample by at least one of the following criteria: age, weight, sex, disease stage, and overall health. A standard curve of levels of a purified protein, e.g., any described herein, within the normal reference range can also be used as a reference.

As used herein, the term “subject” refers to any organism to which a composition in accordance with the invention may be administered, e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include any animal (e.g., mammals such as mice, rats, rabbits, non-human primates, and humans). A subject may seek or be in need of treatment, require treatment, be receiving treatment, be receiving treatment in the future, or be a human or animal who is under care by a trained professional for a particular disease or condition.

As used herein, the terms “treat,” “treated,” or “treating” mean both therapeutic treatment and prophylactic or preventative measures wherein the object is to prevent or slow down (lessen) an undesired physiological condition, disorder, or disease, or obtain beneficial or desired clinical results. Beneficial or desired clinical results include, but are not limited to, alleviation of symptoms; diminishment of the extent of a condition, disorder, or disease; stabilized (i.e., not worsening) state of condition, disorder, or disease; delay in onset or slowing of condition, disorder, or disease progression; amelioration of the condition, disorder, or disease state or remission (whether partial or total), whether detectable or undetectable; an amelioration of at least one measurable physical parameter, not necessarily discernible by the patient; or enhancement or improvement of condition, disorder, or disease. Treatment includes eliciting a clinically significant response without excessive levels of side effects. Treatment also includes prolonging survival as compared to expected survival if not receiving treatment.

As used herein, the terms “variant” and “derivative” are used interchangeably and refer to naturally-occurring, synthetic, and semi-synthetic analogues of a compound, peptide, protein, or other substance described herein. A variant or derivative of a compound, peptide, protein, or other substance described herein may retain or improve upon the biological activity of the original material.

The details of one or more embodiments of the invention are set forth in the description below. Other features, objects,

and advantages of the invention will be apparent from the description and from the claims.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph illustrating inhibition of cell proliferation of several cancer cell lines by a BRG1/BRM inhibitor (compound 17).

FIG. 2 is a graph illustrating inhibition of cell proliferation of several cancer cell lines by a BRG1/BRM inhibitor, compound 18.

FIG. 3 is a graph illustrating the area under the curves (AUCs) calculated from dose-response curves for cancer cell lines treated with a BRG1/BRM inhibitor.

FIG. 4 is a graph illustrating in vivo inhibition of AML proliferation by a BRG1/BRM inhibitor.

FIG. 5 is a graph illustrating in vivo inhibition of AML proliferation by a BRG1/BRM inhibitor.

FIG. 6 is a graph illustrating in vivo inhibition of AML proliferation by a BRG1/BRM inhibitor.

DETAILED DESCRIPTION

The present inventors have found that depletion of BRG1 and/or BRM in AML results in decreased proliferation of the cancer cells.

Accordingly, the invention features methods and compositions useful for the inhibition of the activity of the BRG1 and/or BRM, e.g., for the treatment of AML. The invention further features methods and compositions useful for inhibition of the activity of the BRG1 and/or BRM protein, e.g., for the treatment of AML, e.g., in a subject in need thereof. Exemplary methods are described herein.

BRG1 and/or BRM-Reducing Agents

Agents described herein that reduce the level and/or activity of BRG1 and/or BRM in a cell may be an antibody, a protein (such as an enzyme), a polynucleotide, or a small molecule compound. The agents reduce the level of an activity related to BRG1 and/or BRM, or a related downstream effect, or reduce the level of BRG1 and/or BRM in a cell or subject.

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is an enzyme, a polynucleotide, or a small molecule compound such as a small molecule BRG1 and/or BRM inhibitor.

Antibodies

The agent that reduces the level and/or activity of BRG1 and/or BRM can be an antibody or antigen binding fragment thereof. For example, an agent that reduces the level and/or activity of BRG1 and/or BRM described herein is an antibody that reduces or blocks the activity and/or function of BRG1 and/or BRM through binding to BRG1 and/or BRM.

The making and use of therapeutic antibodies against a target antigen (e.g., BRG1 and/or BRM) is known in the art. See, for example, the references cited herein above, as well as Zhiqiang An (Editor), *Therapeutic Monoclonal Antibodies: From Bench to Clinic*. 1st Edition. Wiley 2009, and also Greenfield (Ed.), *Antibodies: A Laboratory Manual*. (Second edition) Cold Spring Harbor Laboratory Press 2013, for methods of making recombinant antibodies, including antibody engineering, use of degenerate oligonucleotides, 5'-RACE, phage display, and mutagenesis; antibody testing and characterization; antibody pharmacokinetics and pharmacodynamics; antibody purification and storage; and screening and labeling techniques.

Polynucleotides

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM is a polynucleotide. In some embodiments, the polynucleotide is an inhibitory RNA molecule, e.g., that acts by way of the RNA interference (RNAi) pathway. An inhibitory RNA molecule can decrease the expression level (e.g., protein level or mRNA level) of BRG1 and/or BRM. For example, an inhibitory RNA molecule includes a short interfering RNA (siRNA), short hairpin RNA (shRNA), and/or a microRNA (miRNA) that targets full-length BRG1 and/or BRM. A siRNA is a double-stranded RNA molecule that typically has a length of about 19-25 base pairs. A shRNA is a RNA molecule including a hairpin turn that decreases expression of target genes via RNAi. A microRNA is a non-coding RNA molecule that typically has a length of about 22 nucleotides. miRNAs bind to target sites on mRNA molecules and silence the mRNA, e.g., by causing cleavage of the mRNA, destabilization of the mRNA, or inhibition of translation of the mRNA. Degradation is caused by an enzymatic, RNA-induced silencing complex (RISC).

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM is an antisense nucleic acid. Antisense nucleic acids include antisense RNA (asRNA) and antisense DNA (asDNA) molecules, typically about 10 to 30 nucleotides in length, which recognize polynucleotide target sequences or sequence portions through hydrogen bonding interactions with the nucleotide bases of the target sequence (e.g., BRG1 and/or BRM). The target sequences may be single- or double-stranded RNA, or single- or double-stranded DNA.

In some embodiments, the polynucleotide decreases the level and/or activity of a negative regulator of function or a positive regulator of function. In other embodiments, the polynucleotide decreases the level and/or activity of an inhibitor of a positive regulator of function.

A polynucleotide of the invention can be modified, e.g., to contain modified nucleotides, e.g., 2'-fluoro, 2'-o-methyl, 2'-deoxy, unlocked nucleic acid, 2'-hydroxy, phosphorothioate, 2'-thiouridine, 4'-thiouridine, 2'-deoxyuridine. Without being bound by theory, it is believed that certain modification can increase nuclease resistance and/or serum stability, or decrease immunogenicity. The polynucleotides mentioned above, may also be provided in a specialized form such as liposomes, microspheres, or may be applied to gene therapy, or may be provided in combination with attached moieties. Such attached moieties include polycations such as polylysine that act as charge neutralizers of the phosphate backbone, or hydrophobic moieties such as lipids (e.g., phospholipids, cholesterol, etc.) that enhance the interaction with cell membranes or increase uptake of the nucleic acid. These moieties may be attached to the nucleic acid at the 3' or 5' ends and may also be attached through a base, sugar, or intramolecular nucleoside linkage. Other moieties may be capping groups specifically placed at the 3' or 5' ends of the nucleic acid to prevent degradation by nucleases such as exonuclease, RNase, etc. Such capping groups include hydroxyl protecting groups known in the art, including glycols such as polyethylene glycol and tetraethylene glycol. The inhibitory action of the polynucleotide can be examined using a cell-line or animal based gene expression system of the present invention in vivo and in vitro. In some embodiments, the polynucleotide decreases the level and/or activity or function of BRG1 and/or BRM. In other embodiments, the polynucleotide inhibits expression of BRG1 and/or BRM. In other embodiments, the polynucleotide increases degradation of BRG1 and/or BRM and/or

decreases the stability (i.e., half-life) of BRG1 and/or BRM. The polynucleotide can be chemically synthesized or transcribed in vitro.

Inhibitory polynucleotides can be designed by methods well known in the art. siRNA, miRNA, shRNA, and asRNA molecules with homology sufficient to provide sequence specificity required to uniquely degrade any RNA can be designed using programs known in the art, including, but not limited to, those maintained on websites for Thermo Fisher Scientific, the German Cancer Research Center, and The Ohio State University Wexner Medical Center. Systematic testing of several designed species for optimization of the inhibitory polynucleotide sequence can be routinely performed by those skilled in the art. Considerations when designing interfering polynucleotides include, but are not limited to, biophysical, thermodynamic, and structural considerations, base preferences at specific positions in the sense strand, and homology. The making and use of inhibitory therapeutic agents based on non-coding RNA such as ribozymes, RNase P, siRNAs, and miRNAs are also known in the art, for example, as described in Sioud, RNA Therapeutics: Function, Design, and Delivery (Methods in Molecular Biology). Humana Press 2010. Exemplary inhibitory polynucleotides, for use in the methods of the invention, are provided in Table 1, below. In some embodiments, the inhibitory polynucleotides have a nucleic acid sequence with at least 50% (e.g., at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%) sequence identity to the nucleic acid sequence of an inhibitory polynucleotide in Table 1. In some embodiments, the inhibitory polynucleotides have a nucleic acid sequence with at least 70% sequence identity (e.g., 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.9% identity, or more) to the nucleic acid sequence of an inhibitory polynucleotide in Table 1.

Construction of vectors for expression of polynucleotides for use in the invention may be accomplished using conventional techniques which do not require detailed explanation to one of ordinary skill in the art. For generation of efficient expression vectors, it is necessary to have regulatory sequences that control the expression of the polynucleotide. These regulatory sequences include promoter and enhancer sequences and are influenced by specific cellular factors that interact with these sequences, and are well known in the art.

Gene Editing

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM is a component of a gene editing system. For example, the agent that reduces the level and/or activity of BRG1 and/or BRM introduces an alteration (e.g., insertion, deletion (e.g., knockout), translocation, inversion, single point mutation, or other mutation) in BRG1 and/or BRM. In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM is a nuclease. Exemplary gene editing systems include the zinc finger nucleases (ZFNs), Transcription Activator-Like Effector-based Nucleases (TALENs), and the clustered regulatory interspaced short palindromic repeat (CRISPR) system. ZFNs, TALENs, and CRISPR-based methods are described, e.g., in Gaj et al., *Trends Biotechnol.* 31(7):397-405 (2013).

CRISPR refers to a set of (or system including a set of) clustered regularly interspaced short palindromic repeats. A CRISPR system refers to a system derived from CRISPR and Cas (a CRISPR-associated protein) or other nuclease

that can be used to silence or mutate a gene described herein. The CRISPR system is a naturally occurring system found in bacterial and archeal genomes. The CRISPR locus is made up of alternating repeat and spacer sequences. In naturally-occurring CRISPR systems, the spacers are typically sequences that are foreign to the bacterium (e.g., plasmid or phage sequences). The CRISPR system has been modified for use in gene editing (e.g., changing, silencing, and/or enhancing certain genes) in eukaryotes. See, e.g., Wiedenheft et al., *Nature* 482(7385):331-338 (2012). For example, such modification of the system includes introducing into a eukaryotic cell a plasmid containing a specifically-designed CRISPR and one or more appropriate Cas proteins. The CRISPR locus is transcribed into RNA and processed by Cas proteins into small RNAs that include a repeat sequence flanked by a spacer. The RNAs serve as guides to direct Cas proteins to silence specific DNA/RNA sequences, depending on the spacer sequence. See, e.g., Horvath et al., *Science* 327(5962):167-170 (2010); Makarova et al., *Biology Direct* 1:7 (2006); Pennisi, *Science* 341(6148):833-836 (2013). In some examples, the CRISPR system includes the Cas9 protein, a nuclease that cuts on both strands of the DNA. See, e.g., Id.

In some embodiments, in a CRISPR system for use described herein, e.g., in accordance with one or more methods described herein, the spacers of the CRISPR are derived from a target gene sequence, e.g., from a BRG1 and/or BRM sequence. In some embodiments, in a CRISPR system for use described herein, e.g., in accordance with one or more methods described herein, the spacers of the CRISPR are derived from a target gene sequence, e.g., from a BRG1 sequence. In some embodiments, in a CRISPR system for use described herein, e.g., in accordance with one or more methods described herein, the spacers of the CRISPR are derived from a target gene sequence, e.g., from a BRM sequence.

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM includes a guide RNA (gRNA) for use in a CRISPR system for gene editing. Exemplary gRNAs, for use in the methods of the invention, are provided in Table 1, below. In embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM includes a ZFN, or an mRNA encoding a ZFN, that targets (e.g., cleaves) a nucleic acid sequence (e.g., DNA sequence) of BRG1 and/or BRM. In embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM includes a TALEN, or an mRNA encoding a TALEN, that targets (e.g., cleaves) a nucleic acid sequence (e.g., DNA sequence) of BRG1 and/or BRM. In embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM includes a TALEN, or an mRNA encoding a TALEN, that targets (e.g., cleaves) a nucleic acid sequence (e.g., DNA sequence) of BRG1. In embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM includes a TALEN, or an mRNA encoding a TALEN, that targets (e.g., cleaves) a nucleic acid sequence (e.g., DNA sequence) of BRM.

For example, the gRNA can be used in a CRISPR system to engineer an alteration in a gene (e.g., BRG1 and/or BRM). In other examples, the ZFN and/or TALEN can be used to engineer an alteration in a gene (e.g., BRG1 and/or BRM). Exemplary alterations include insertions, deletions (e.g., knockouts), translocations, inversions, single point mutations, or other mutations. The alteration can be introduced in the gene in a cell, e.g., in vitro, ex vivo, or in vivo. In some embodiments, the alteration decreases the level and/or activity of (e.g., knocks down or knocks out) BRG1 and/or BRM,

47

e.g., the alteration is a negative regulator of function. In yet another example, the alteration corrects a defect (e.g., a mutation causing a defect), in BRG1 and/or BRM. In yet another example, the alteration corrects a defect (e.g., a mutation causing a defect), in BRG1. In yet another example, the alteration corrects a defect (e.g., a mutation causing a defect), in BRM.

In certain embodiments, the CRISPR system is used to edit (e.g., to add or delete a base pair) a target gene, e.g., BRG1 and/or BRM. In other embodiments, the CRISPR system is used to introduce a premature stop codon, e.g., thereby decreasing the expression of a target gene. In yet other embodiments, the CRISPR system is used to turn off a target gene in a reversible manner, e.g., similarly to RNA interference. In embodiments, the CRISPR system is used to direct Cas to a promoter of a target gene, e.g., BRG1 and/or BRM, thereby blocking an RNA polymerase sterically. In embodiments, the CRISPR system is used to direct Cas to a promoter of a target gene, e.g., BRG1, thereby blocking an RNA polymerase sterically. In embodiments, the CRISPR system is used to direct Cas to a promoter of a target gene, e.g., BRM, thereby blocking an RNA polymerase sterically.

In some embodiments, a CRISPR system can be generated to edit BRG1 and/or BRM using technology described in, e.g., U.S. Publication No. 20140068797; Cong et al., *Science* 339(6121):819-823 (2013); Tsai, *Nature Biotechnol.*, 32(6):569-576 (2014); and U.S. Pat. Nos. 8,871,445; 8,865,406; 8,795,965; 8,771,945; and 8,697,359.

In some embodiments, the CRISPR interference (CRISPRi) technique can be used for transcriptional repression of specific genes, e.g., the gene encoding BRG1 and/or BRM. In CRISPRi, an engineered Cas9 protein (e.g., nuclease-null dCas9, or dCas9 fusion protein, e.g., dCas9-KRAB or dCas9-SID4x fusion) can pair with a sequence specific guide RNA (sgRNA). The Cas9-gRNA complex can block RNA polymerase, thereby interfering with transcription elongation. The complex can also block transcription initiation by interfering with transcription factor binding. The CRISPRi method is specific with minimal off-target effects and is multiplexable, e.g., can simultaneously repress more than one gene (e.g., using multiple gRNAs). Also, the CRISPRi method permits reversible gene repression. In some embodiments, CRISPR-mediated gene activation (CRISPRa) can be used for transcriptional activation, e.g., of one or more genes described herein, e.g., a gene that inhibits BRG1 and/or BRM. In the CRISPRa technique, dCas9 fusion proteins recruit transcriptional activators. For example, dCas9 can be used to recruit polypeptides (e.g., activation domains) such as VP64 or the p65 activation domain (p65D) and used with sgRNA (e.g., a single sgRNA or multiple sgRNAs), to activate a gene or genes, e.g., endogenous gene(s). Multiple activators can be recruited by using multiple sgRNAs—this can increase activation efficiency. A variety of activation domains and single or multiple activation domains can be used. In addition to engineering dCas9 to recruit activators, sgRNAs can also be engineered to recruit activators. For example, RNA aptamers can be incorporated into a sgRNA to recruit proteins (e.g., activation domains) such as VP64. In some examples, the synergistic activation mediator (SAM) system can be used for transcriptional activation. In SAM, MS2 aptamers are added to the sgRNA. MS2 recruits the MS2 coat protein (MCP) fused to p65AD and heat shock factor 1 (HSF1). The CRISPRi and CRISPRa techniques are described in greater detail, e.g., in Dominguez et al., *Nat. Rev. Mol. Cell Biol.* 17(1):5-15 (2016), incorporated herein by reference.


48

Small Molecule Compounds

In some embodiments of the invention, the agent that reduces the level and/or activity of BRG1 and/or BRM in a cell is a small molecule compound. In some embodiments, the small molecule compound is a structure of Formula I-III.

In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula I:

Formula I




wherein m is 0, 1, 2, 3, or 4;

X¹ is N or CH; and

each R¹ is, independently, halogen, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₁₀ carbocyclyl, optionally substituted C₂-C₉ heterocyclyl, optionally substituted C₆-C₁₀ aryl, optionally substituted C₂-C₉ heteroaryl, optionally substituted C₂-C₆ alkenyl, optionally substituted C₂-C₆ heteroalkenyl, hydroxy, thiol, or optionally substituted amino.

In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula II:

Formula II



wherein R² is phenyl that is substituted with hydroxy and that is optionally substituted with one or more groups independently selected from the group consisting of halo, cyano, trifluoromethyl, trifluoromethoxy, C₁-₃ alkyl, and C₁-₃ alkoxy;

R³ is selected from the group consisting of —Rᵃ, —O—Rᵃ, —N(Rᵃ)₂, —S(O)₂Rᵃ, and —C(O)—N(Rᵃ)₂;

each Rᵃ is, independently, selected from the group consisting of hydrogen, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein each C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from the group consisting of Rᵇ, oxo, halo, —NO₂, —N(Rᵇ)₂, —CN, —C(O)—N(Rᵇ)₂, —S(O)—N(Rᵇ)₂, —S(O)₂—N(Rᵇ)₂, —O—Rᵇ, —S—Rᵇ, —O—C(O)—Rᵇ, —C(O)—Rᵇ, —C(O)—ORᵇ, —S(O)—Rᵇ, —S(O)₂—Rᵇ, —N(Rᵇ)—C(O)—Rᵇ, —N(Rᵇ)—S(O)—Rᵇ, —N(Rᵇ)—C(O)—N(Rᵇ)₂, and —N(Rᵇ)—S(O)₂—Rᵇ;

each Rᵇ is independently selected from the group consisting of hydrogen, C₁-₆ alkyl, C₂-₆ alkenyl, C₂-₆ alkynyl, C₁-₆ alkoxy, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein each C₁-₆ alkyl, C₂-₆

49

alkenyl, C₂₋₆ alkynyl, C₁₋₆ alkoxy, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from RC; or two R^b are taken together with the nitrogen to which they are attached to form a heterocyclyl that is optionally substituted with one or more groups independently selected from the group consisting of oxo, halo and C₁₋₃ alkyl that is optionally substituted with one or more groups independently selected from the group consisting of oxo and halo;

each RC is independently selected from the group consisting of oxo, halo, —NO₂, —N(R^d)₂, —CN, —C(O)—N(R^d)₂, —S(O)—N(R^d)₂, —S(O)₂—N(R^d)₂, —S—R^d, —O—C(O)—R^d, —C(O)—R^d, —C(O)—OR^d, —S(O)—R^d, —S(O)₂—R^d, —N(R^d)—C(O)—R^d, —N(R^d)—S(O)—R^d, —N(R^d)—C(O)—N(R^d)₂, —N(R^d)—S(O)₂—R^d, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl, wherein any C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, 3-15 membered carbocyclyl, and 3-15 membered heterocyclyl is optionally substituted with one or more groups independently selected from the group consisting of R^d, oxo, halo, —NO₂, —N(R^d)₂, —CN, —C(O)—N(R^d)₂, —S(O)—N(R^d)₂, —S(O)₂—N(R^d)₂, —O—R^d, —S—R^d, —O—C(O)—R^d, —C(O)—R^d, —C(O)—R^d, —S(O)—R^d, —S(O)₂—R^d, —N(R^d)—C(O)—R^d, —N(R^d)—S(O)—R^d, —N(R^d)—C(O)—N(R^d)₂, and —N(R^d)—S(O)₂—R^d;


each R^d is independently selected from the group consisting of hydrogen, C₁₋₆ alkyl, C₂₋₆ alkenyl, C₂₋₆ alkynyl, carbocyclyl, and carbocyclyl(C₁₋₃ alkyl)-;

R^d is H, C₁₋₆ alkyl, or —C(=O)—C₁₋₆ alkyl; and

R^d is H or C₁₋₆ alkyl.

Compounds of Formula II may be synthesized by methods known in the art, e.g., those described in U.S. Patent Publication No. 2018/0086720, the synthetic methods of which are incorporated by reference.

In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula III:












wherein R⁶ is halo, e.g., fluoro or chloro;

R⁷ is hydrogen, optionally substituted amino, or optionally substituted C₁₋₃ alkyl; and

R⁸ is optionally substituted C₆₋₁₀ aryl or optionally substituted C₂₋₉ heteroaryl.


In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of any one of compounds 1-16:

50



51


-continued



52

In some embodiments, the small molecule BRG1 and/or BRM inhibitor is a compound, or pharmaceutically acceptable salt thereof, having the structure of Formula IV:

Formula IV



where R¹ is absent, H, optionally substituted C₁-C₆ acyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶;



is 5- or 6-membered heteroarylene;


each of R² and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl;

R³ is H or optionally substituted C₁-C₆ alkyl; and R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl; or R³ and R⁴, together with the carbon atom to which each is attached, form an optionally substituted C₃-C₆ cycloalkyl;

R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸;

R⁷ and R⁸ are, independently, optionally substituted C₁-C₆ alkyl;

Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or



A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene;

L is absent, —O—, optionally substituted C₁–C₆ alkylene, optionally substituted C₁–C₆ heteroalkylene, optionally substituted C₂–C₆ alkenylene, optionally substituted C₂–C₆ heteroalkenylene, optionally substituted C₂–C₆ alkynylene, optionally substituted C₂–C₆ heteroalkynylene, optionally substituted C₂–C₉ heterocyclylene, optionally substituted C₂–C₉ heterocyclyl C₁–C₆ alkylene, optionally substituted C₂–C₉ heteroarylene, or optionally substituted C₂–C₉ heteroaryl C₁–C₆ alkylene; and

B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl, or a pharmaceutically acceptable salt thereof.

53

In some embodiments,




is 6-membered heteroarylene. In some embodiments,



is 5-membered heteroarylene.


In some embodiments,



where each of X, Y, and Z is, independently, N or CH.

In some embodiments, the compound of Formula IV has the structure of Formula IVa:

Formula IVa



where each of X, Y, and Z is, independently, N or CH;

R¹ is H, optionally substituted C₁-C₆ acyl, optionally


substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶;

each of R², R³, and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl;

R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl;

R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸; each of R⁷ and R⁸ is, independently, optionally substituted C₁-C₆ alkyl;

Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or



A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene;

L is absent, —O—, optionally substituted C₁-C₆ alkylene, optionally substituted C₁-C₆ heteroalkylene, optionally substituted C₂-C₆ alkenylene, optionally substituted

54


C₂-C₆ heteroalkenylene, optionally substituted C₂-C₆ alkynylene, optionally substituted C₂-C₆ heteroalkynylene, optionally substituted C₂-C₉ heterocyclylene, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkylene, optionally substituted C₂-C₉ heteroarylene, or optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkylene; and

B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl, or a pharmaceutically acceptable salt thereof.

In some embodiments, X, Y, and Z are CH; X is N and Y and Z are CH; Z is N and X and Y are CH; Y is N and X and Z are CH; X is CH and Y and Z are N; Z is CH and X and Y are N; Y is CH and X and Z are N; or X, Y, and Z are N.

In some embodiments, the compound of Formula IV has the structure of Formula IVb:


Formula IVb



or a pharmaceutically acceptable salt thereof.


In some embodiments, the compound of Formula IV has the structure of Formula IVc:

Formula IVc



or a pharmaceutically acceptable salt thereof. In some embodiments, the compound of Formula IV has the structure of Formula Ic:


Formula IVd



or a pharmaceutically acceptable salt thereof.

In some embodiments, the compound of Formula IV has the structure of Formula IVe:


Formula IVe



or a pharmaceutically acceptable salt thereof.

55


In some embodiments,



where X' is O or S; Y' is N or CH; and Z' is N or CH.

In some embodiments, the compound of Formula IVa has the structure of Formula V:

Formula V



where

W is C or N;

X' is O, S, or N—CH₃;

Y' is N or CH;

Z' is N or CH;


R¹ is absent, H, optionally substituted C₁-C₆ acyl, optionally substituted C₁-C₆ alkyl, optionally substituted C₁-C₆ heteroalkyl, optionally substituted C₃-C₈ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or —SO₂R⁶;

each of R², R³, and R⁵ is, independently, H or optionally substituted C₁-C₆ alkyl;

R⁴ is H, optionally substituted C₁-C₆ alkyl, or optionally substituted C₁-C₆ heteroalkyl;

R⁶ is optionally substituted C₁-C₆ alkyl or —NR⁷R⁸; each of R⁷ and R⁸ is, independently, optionally substituted C₁-C₆ alkyl;

Het is optionally substituted 5-membered heteroarylene, optionally substituted 6-membered heteroarylene, or



A is optionally substituted C₆-C₁₀ arylene, optionally substituted C₂-C₉ heterocyclylene, or optionally substituted C₂-C₉ heteroarylene;

L is absent, —O—, optionally substituted C₁-C₆ alkylene, optionally substituted C₁-C₆ heteroalkylene, optionally substituted C₁-C₆ alkenylene, optionally substituted C₂-C₆ heteroalkenylene, optionally substituted C₂-C₆ alkynylene, optionally substituted C₂-C₆ heteroalkynylene, optionally substituted C₂-C₉ heterocyclylene, optionally substituted C₂-C₉ heterocyclyl C₁-C₆ alkylene, optionally substituted C₂-C₉ heteroarylene, or optionally substituted C₂-C₉ heteroaryl C₁-C₆ alkylene; and

B is H, halogen, cyano, optionally substituted C₆-C₁₀ aryl, optionally substituted C₃-C₁₀ cycloalkyl, optionally substituted C₂-C₉ heterocyclyl, or optionally substituted C₂-C₉ heteroaryl,


or a pharmaceutically acceptable salt thereof.

56

In some embodiments, X' is O, Y' is CH, and Z' is N; X' is S, Y' is CH, and Z' is N; X' is O, Y' is N, and Z' is CH; X' is S, Y' is N, and Z' is CH; X' is O, Y' is N, and Z' is N; or X' is S, Y' is N, and Z' is N.

In some embodiments, the compound of Formula V has the structure of Formula Va:


Formula Va



or a pharmaceutically acceptable salt thereof.




In some embodiments, the compound of Formula II has the structure of Formula Vb:

Formula Vb




or a pharmaceutically acceptable salt thereof.

In some embodiments, the small molecule compound, or pharmaceutically acceptable salt thereof is any one of compounds 17-20 having the structure:



57

-continued



20

5

10

In some embodiments, the small molecule compound, or a pharmaceutically acceptable salt thereof is a degrader. In some embodiments, the degrader has the structure of Formula VI:



C-L-D

Formula VI

wherein C is a BRG1 and/or BRM binding moiety; L is a linker; and D is a degradation moiety, or a pharmaceutically acceptable salt thereof. In some embodiments, the degradation moiety is a ubiquitin ligase moiety. In some embodiments, the ubiquitin ligase binding moiety includes Cereblon ligands, IAP (Inhibitors of Apoptosis) ligands, mouse double minute 2 homolog (MDM2), hydrophobic tag, or von Hippel-Lindau ligands, or derivatives or analogs thereof.

In some embodiments, A includes the structure of any one of Formula I-V, or any one of compounds 1-20.


In some embodiments, the hydrophobic tag includes a diphenylmethane, adamantane, or tri-Boc arginine, i.e., the hydrophobic tag includes the structure:




In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula A:

58

Formula A





wherein X^1 is CH_2 , O, S, or NR^1 , wherein R^1 is H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; X^2 is $C=O$, CH_2 , or



R^3 and R^4 are, independently, H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; m is 0, 1, 2, 3, or 4; and each R^2 is, independently, halogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_2 - C_9 heterocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 heteroalkenyl, hydroxy, thiol, or optionally substituted amino,

or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure:




or is a derivative or an analog thereof, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula B:


59

Formula B




wherein each R^4 , R^4 , and R^7 is, independently, H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; R^5 is optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_1 - C_6 alkyl C_3 - C_{10} carbocyclyl, or optionally substituted C_1 - C_6 alkyl C_6 - C_{10} aryl; R^6 is H, optionally substituted C_1 - C_6 alkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_1 - C_6 alkyl C_3 - C_{10} carbocyclyl, or optionally substituted C_1 - C_6 alkyl C_6 - C_{10} aryl; n is 0, 1, 2, 3, or 4; each R^8 is, independently, halogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_2 - C_9 heterocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 heteroalkenyl, hydroxy, thiol, or optionally substituted amino; and each R^9 and R^{10} is, independently, H, halogen, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_6 - C_{10} aryl, wherein R^4 or R^5 includes a bond to the linker, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure:



60


-continued



or is a derivative or analog thereof, or a pharmaceutically acceptable salt thereof.


In some embodiments, the ubiquitin ligase binding moiety includes the structure of Formula C:

Formula C



wherein each R^{11} , R^{13} , and R^{15} is, independently, H, optionally substituted C_1 - C_6 alkyl, or optionally substituted C_1 - C_6 heteroalkyl; R^{12} is optionally substituted C_1 - C_6 alkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_1 - C_6 alkyl C_3 - C_{10} carbocyclyl, or optionally substituted C_1 - C_6 alkyl C_6 - C_{10} aryl; R^{14} is optionally substituted C_1 - C_6 alkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_1 - C_6 alkyl C_3 - C_{10} carbocyclyl, or optionally substituted C_1 - C_6 alkyl C_6 - C_{10} aryl; p is 0, 1, 2, 3, or 4; each R^{16} is, independently, halogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_2 - C_9 heterocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_6 alkenyl, optionally substituted C_2 - C_6 heteroalkenyl, hydroxy, thiol, or optionally substituted amino; q is 0, 1, 2, 3, or 4; and each R^{17} is, independently, halogen, optionally substituted C_1 - C_6 alkyl, optionally substituted C_1 - C_6 heteroalkyl, optionally substituted C_3 - C_{10} carbocyclyl, optionally substituted C_2 - C_9 heterocyclyl, optionally substituted C_6 - C_{10} aryl, optionally substituted C_2 - C_9 heteroaryl, optionally substituted C_2 - C_6 alkenyl, hydroxy, thiol, or optionally substituted amino, or a pharmaceutically acceptable salt thereof.

In some embodiments, the ubiquitin ligase binding moiety includes the structure:




63

wherein A¹ is a bond between the linker and A, and A² is a bond between B and the linker.

In some embodiments, D is optionally substituted C₁₋₁₀ alkyl. In some embodiments, C¹ and C² are each, independently, a carbonyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁₋₂ alkyl, optionally substituted C₁₋₃ heteroalkyl, O, S, S(O)₂, and NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B¹, B², B³, and B⁴ each, independently, is selected from optionally substituted C₁₋₂ alkyl, O, S, S(O)₂, and NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B¹ and B⁴ each, independently, is C₁ alkyl. In some embodiments, B² and B⁴ each, independently, is NR^N, wherein R^N is hydrogen or optionally substituted C₁₋₄ alkyl. In some embodiments, B² and B⁴ each, independently, is NH. In some embodiments, f, g, h, i, j, and k are each, independently, 1.

In some embodiments, the linker of Formula V has the structure of Formula VIIb:

Formula VIIb



wherein A¹ is a bond between the linker and A, and A² is a bond between B and the linker.

Pharmaceutical Uses

The compounds described herein are useful in the methods of the invention and, while not bound by theory, are believed to exert their desirable effects through their ability to modulate the level, status, and/or activity of a BAF complex, e.g., by inhibiting the activity or level of the BRG1 and/or BRM proteins in a cell within the BAF complex in a mammal.

An aspect of the present invention relates to methods of treating disorders related to BRG1 and/or BRM proteins such as AML in a subject in need thereof. In some embodiments, the compound is administered in an amount and for a time effective to result in one of (or more, e.g., two or more, three or more, four or more of): (a) reduced tumor size, (b) reduced rate of tumor growth, (c) increased tumor cell death (d) reduced tumor progression, (e) reduced number of metastases, (f) reduced rate of metastasis, (g) decreased tumor recurrence (h) increased survival of subject, and (i) increased progression free survival of a subject.

Treating cancer can result in a reduction in size or volume of a tumor. For example, after treatment, tumor size is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or greater) relative to its size prior to treatment. Size of a tumor may be measured by any reproducible means of measurement. For example, the size of a tumor may be measured as a diameter of the tumor.

Treating cancer may further result in a decrease in number of tumors. For example, after treatment, tumor number is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or greater) relative to number prior to treatment. Number of tumors may be measured by any reproducible means of measurement, e.g., the number of

64

tumors may be measured by counting tumors visible to the naked eye or at a specified magnification (e.g., 2×, 3×, 4×, 5×, 10×, or 50×).

Treating cancer can result in a decrease in number of metastatic nodules in other tissues or organs distant from the primary tumor site (e.g., in the liver). For example, after treatment, the number of metastatic nodules is reduced by 5% or greater (e.g., 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or greater) relative to number prior to treatment. The number of metastatic nodules may be measured by any reproducible means of measurement. For example, the number of metastatic nodules may be measured by counting metastatic nodules visible to the naked eye or at a specified magnification (e.g., 2×, 10×, or 50×).

Treating cancer can result in inhibition or slowing of the metastatic progression of the cancer. For example, a patient may be administered an amount of an agent that reduces the activity or level of the BRG1 and/or BRM that is effective to inhibit metastasis of the cancer to other parts of the body (e.g., a patient having uveal melanoma that has metastasized (e.g., to the liver)). An agent may be administered in an adjuvant or neo-adjuvant setting, such as prior to or subsequent to surgical resection of the cancer, and result in a decrease incidence of metastasis of the cancer.

Treating cancer can result in an increase in average survival time of a population of subjects treated according to the present invention in comparison to a population of untreated subjects. For example, the average survival time is increased by more than 30 days (more than 60 days, 90 days, or 120 days). An increase in average survival time of a population may be measured by any reproducible means. An increase in average survival time of a population may be measured, for example, by calculating for a population the average length of survival following initiation of treatment with the compound described herein. An increase in average survival time of a population may also be measured, for example, by calculating for a population the average length of survival following completion of a first round of treatment with a pharmaceutically acceptable salt of a compound described herein.

Treating cancer can also result in a decrease in the mortality rate of a population of treated subjects in comparison to an untreated population. For example, the mortality rate is decreased by more than 2% (e.g., more than 5%, 10%, or 25%). A decrease in the mortality rate of a population of treated subjects may be measured by any reproducible means, for example, by calculating for a population the average number of disease-related deaths per unit time following initiation of treatment with a pharmaceutically acceptable salt of a compound described herein. A decrease in the mortality rate of a population may also be measured, for example, by calculating for a population the average number of disease-related deaths per unit time following completion of a first round of treatment with a pharmaceutically acceptable salt of a compound described herein.

Combination Therapies

A method of the invention can be used alone or in combination with an additional therapeutic agent, e.g., other agents that treat cancer or symptoms associated therewith, or in combination with other types of therapies to treat cancer. In combination treatments, the dosages of one or more of the therapeutic compounds may be reduced from standard dosages when administered alone. For example, doses may be determined empirically from drug combinations and permutations or may be deduced by isobolographic analysis (e.g.,

Black et al., *Neurology* 65:S3-S6 (2005)). In this case, dosages of the compounds when combined should provide a therapeutic effect.

In some embodiments, the second therapeutic agent is a chemotherapeutic agent (e.g., a cytotoxic agent or other chemical compound useful in the treatment of cancer). These include alkylating agents, antimetabolites, folic acid analogs, pyrimidine analogs, purine analogs and related inhibitors, vinca alkaloids, epipodophyllotoxins, antibiotics, L-Asparaginase, topoisomerase inhibitors, interferons, platinum coordination complexes, anthracenedione substituted urea, methyl hydrazine derivatives, adrenocortical suppressant, adrenocorticosteroides, progestins, estrogens, antiestrogen, androgens, antiandrogen, and gonadotropin-releasing hormone analog. Also included is 5-fluorouracil (5-FU), leucovorin (LV), irinotecan, oxaliplatin, capecitabine, paclitaxel, and doxorubicin. Non-limiting examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; calyculin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammall and calicheamicin omegall (see, e.g., *Agnew, Chem. Intl. Ed Engl.* 33:183-186 (1994)); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, anthramycin, azaserine, bleomycin, cactinomycin, carabacin, caminomycin, carzinophilin, chromomycinis, dactinomycin, daunorubicin, detorubicin, 6-di-azo-5-oxo-L-norleucine, ADRIAMYCIN® (doxorubicin, including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofof, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epiandrostanol, mepitiostane, testosterone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestabucil; bisantrene; edatraxate; defofamine; demecol-

cine; diaziquone; elfomithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, OR); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL® (paclitaxel; Bristol-Myers Squibb Oncology, Princeton, NJ), ABRAXANE®, cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, IL), and TAXOTERE® doxorubicin (Rhône-Poulenc Rorer, Antony, France); chlorambucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum coordination complexes such as cisplatin, oxaliplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (e.g., CPT-11); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Two or more chemotherapeutic agents can be used in a cocktail to be administered in combination with the first therapeutic agent described herein. Suitable dosing regimens of combination chemotherapies are known in the art and described in, for example, Saltz et al., *Proc. Am. Soc. Clin. Oncol.* 18:233a (1999), and Douillard et al., *Lancet* 355(9209):1041-1047 (2000).

In some embodiments, the second therapeutic agent is a therapeutic agent which is a biologic such as a cytokine (e.g., interferon or an interleukin (e.g., IL-2)) used in cancer treatment. In some embodiments the biologic is an anti-angiogenic agent, such as an anti-VEGF agent, e.g., bevacizumab (AVASTIN®). In some embodiments the biologic is an immunoglobulin-based biologic, e.g., a monoclonal antibody (e.g., a humanized antibody, a fully human antibody, an Fc fusion protein or a functional fragment thereof) that agonizes a target to stimulate an anti-cancer response, or antagonizes an antigen important for cancer. Such agents include RITUXAN® (rituximab); ZENAPAX® (daclizumab); SIMULECT® (basiliximab); SYNAGIS® (palivizumab); REMICADE® (infliximab); HERCEPTIN® (trastuzumab); MYLOTARG® (gemtuzumab ozogamicin); CAMPATH® (alemtuzumab); ZEVALIN® (ibritumomab tiuxetan); HUMIRA® (adalimumab); XOLAIR® (omalizumab); BEXXAR® (tositumomab-1-131); RAPTIVA® (efalizumab); ERBITUX® (cetuximab); AVASTIN® (bevacizumab); TYSABRI® (natalizumab); ACTEMRA® (tocilizumab); VECTIBIX® (panitumumab); LUCENTIS® (ranibizumab); SOLIRIS® (eculizumab); CIMZIA® (certolizumab pegol); SIMPONI® (golimumab); ILARIS® (canakinumab); STELARA® (ustekinumab); ARZERRA® (ofatumumab); PROLIA® (denosumab); NUMAX® (motavizumab); ABTHRAX® (raxibacumab); BENLYSTA® (belimumab); YERVOY® (ipilimumab); ADCETRIS® (brentuximab vedotin); PERJETA® (pertuzumab); KADCYLA® (ado-trastuzumab emtansine); and GAZYVA® (obinituzumab). Also included are antibody-drug conjugates.

In some embodiments, the second agent is dacarbazine, temozolomide, cisplatin, treosulfan, fotemustine, IMCgp100, a CTLA-4 inhibitor (e.g., ipilimumab), a PD-1 inhibitor (e.g., Nivolumab or pembrolizumab), a PD-L1 inhibitor (e.g., atezolizumab, avelumab, or durvalumab), a

mitogen-activated protein kinase (MEK) inhibitor (e.g., selumetinib, binimetinib, or tametinib), and/or a protein kinase C (PKC) inhibitor (e.g., sotrastaurin or LXS196).

In some embodiments, the second agent is a mitogen-activated protein kinase (MEK) inhibitor (e.g., selumetinib, binimetinib, or tametinib) and/or a protein kinase C (PKC) inhibitor (e.g., sotrastaurin or LXS196).

In some embodiments, the second agent is cytarabine, an anthracycline such as daunorubicin, arsenic trioxide, all-trans-retinoic acid, or a combination thereof. In some

embodiments, the second agent is an immunotherapy such as histamine dihydrochloride and interleukin 2. In some embodiments, the second agent is gemtuzumab ozogamicin.

The second agent may be a therapeutic agent which is a non-drug treatment. For example, the second therapeutic

agent is radiation therapy, thermotherapy, photocoagulation, cryotherapy, hyperthermia, surgical excision of tumor, and/or a stem cell transplant (e.g., an allogenic stem cell transplant or hemapoeitic stem cell transplant).

The second agent may be a checkpoint inhibitor. In one embodiment, the inhibitor of checkpoint is an inhibitory antibody (e.g., a monospecific antibody such as a monoclonal antibody). The antibody may be, e.g., humanized or fully human. In some embodiments, the inhibitor of checkpoint is a fusion protein, e.g., an Fc-receptor fusion protein. In some

embodiments, the inhibitor of checkpoint is an agent, such as an antibody, that interacts with a checkpoint protein. In some embodiments, the inhibitor of checkpoint is an agent, such as an antibody, that interacts with the ligand of a checkpoint protein. In some embodiments, the inhibitor of

checkpoint is an inhibitor (e.g., an inhibitory antibody or small molecule inhibitor) of CTLA-4 (e.g., an anti-CTLA4 antibody or fusion a protein such as ipilimumab/YERVOY® or tremelimumab). In some embodiments, the inhibitor of

checkpoint is an inhibitor (e.g., an inhibitory antibody or small molecule inhibitor) of PD-1 (e.g., nivolumab/OP-DIVO®; pembrolizumab/KEYTRUDA®; pidilizumab/CT-011). In some embodiments, the inhibitor of checkpoint is an

inhibitor (e.g., an inhibitory antibody or small molecule inhibitor) of PDL1 (e.g., atezolizumab, avelumab, durvalumab, MPDL3280A/RG7446; MEDI4736; MSB0010718C; BMS 936559). In some embodiments, the

inhibitor of checkpoint is an inhibitor (e.g., an inhibitory antibody or Fc fusion or small molecule inhibitor) of PDL2 (e.g., a PDL2/Ig fusion protein such as AMP 224). In some

embodiments, the inhibitor of checkpoint is an inhibitor (e.g., an inhibitory antibody or small molecule inhibitor) of B7-H3 (e.g., MGA271), B7-H4, BTLA, HVEM, TIM3, GAL9, LAG3, VISTA, KIR, 2B4, CD160, CGEN-15049, CHK 1, CHK2, A2aR, B-7 family ligands, or a combination

thereof.

In some embodiments, the anti-cancer therapy is a T cell adoptive transfer (ACT) therapy. In some embodiments, the T cell is an activated T cell. The T cell may be modified to express a chimeric antigen receptor (CAR). CAR modified T (CAR-T) cells can be generated by any method known in the art. For example, the CAR-T cells can be generated by introducing a suitable expression vector encoding the CAR to a T cell. Prior to expansion and genetic modification of the T cells, a source of T cells is obtained from a subject. T cells

tissue, cord blood, thymus tissue, tissue from a site of infection, ascites, pleural effusion, spleen tissue, and tumors. In certain embodiments of the present invention, any number of T cell lines available in the art, may be used. In some

embodiments, the T cell is an autologous T cell. Whether prior to or after genetic modification of the T cells to express a desirable protein (e.g., a CAR), the T cells can be activated and expanded generally using methods as described, for

example, in U.S. Pat. Nos. 6,352,694; 6,534,055; 6,905,680; 6,692,964; 5,858,358; 6,887,466; 6,905,681; 7,144,575; 7,067,318; 7,172,869; 7,232,566; 7,175,843; 5,883,223; 6,905,874; 6,797,514; 6,867,041; and U.S. Patent Application Publication No. 20060121005.

In any of the combination embodiments described herein, the first and second therapeutic agents are administered simultaneously or sequentially, in either order. The first therapeutic agent may be administered immediately, up to 1 hour, up to 2 hours, up to 3 hours, up to 4 hours, up to 5 hours, up to 6 hours, up to 7 hours, up to, 8 hours, up to 9

hours, up to 10 hours, up to 11 hours, up to 12 hours, up to 13 hours, 14 hours, up to hours 16, up to 17 hours, up 18 hours, up to 19 hours up to 20 hours, up to 21 hours, up to 22 hours, up to 23 hours up to 24 hours or up to 1-7, 1-14, 1-21 or 1-30 days before or after the second therapeutic

agent.

Delivery of Anti-BRG1 and/or BRM Agents

A variety of methods are available for the delivery of anti-BRG1 and/or BRM agents to a subject including viral and non-viral methods.

Viral Delivery Methods

In some embodiments, the agent that reduces the level and/or activity of BRG1 and/or BRM is delivered by a viral vector (e.g., a viral vector expressing an anti-BRG1 and/or BRM agent). Viral genomes provide a rich source of vectors that can be used for the efficient delivery of exogenous genes into a mammalian cell. Viral genomes are particularly useful vectors for gene delivery because the polynucleotides contained within such genomes are typically incorporated into the nuclear genome of a mammalian cell by generalized or specialized transduction. These processes occur as part of the natural viral replication cycle, and do not require added proteins or reagents in order to induce gene integration. Examples of viral vectors include a retrovirus (e.g., Retroviridae family viral vector), adenovirus (e.g., Ad5, Ad26, Ad34, Ad35, and Ad48), parvovirus (e.g., adeno-associated viruses), coronavirus, negative strand RNA viruses such as orthomyxovirus (e.g., influenza virus), rhabdovirus (e.g., rabies and vesicular stomatitis virus), paramyxovirus (e.g., measles and Sendai), positive strand RNA viruses, such as picornavirus and alphavirus, and double-stranded DNA viruses including adenovirus, herpesvirus (e.g., Herpes Simplex virus types 1 and 2, Epstein-Barr virus, cytomegalovirus, replication deficient herpes virus), and poxvirus (e.g., vaccinia, modified vaccinia Ankara (MVA), fowlpox and canarypox). Other viruses include Norwalk virus, togavirus, flavivirus, reoviruses, papovavirus, hepadnavirus, human papilloma virus, human foamy virus, and hepatitis virus, for example. Examples of retroviruses include: avian leukosis-sarcoma, avian C-type viruses, mammalian C-type, B-type viruses, D-type viruses, oncoretroviruses, HTLV-BLV group, lentivirus, alpharetrovirus, gammaretrovirus, spumavirus (Coffin, J. M., *Retroviridae: The viruses and their replication*, Virology (Third Edition) Lippincott-Raven, Philadelphia, 1996). Other examples include murine leukemia viruses, murine sarcoma viruses, mouse mammary tumor virus, bovine leukemia virus, feline leukemia virus, feline sarcoma virus, avian leukemia virus, human T cell

leukemia virus, baboon endogenous virus, Gibbon ape leukemia virus, Mason Pfizer monkey virus, simian immunodeficiency virus, simian sarcoma virus, Rous sarcoma virus and lentiviruses. Other examples of vectors are described, for example, in U.S. Pat. No. 5,801,030, the teachings of which are incorporated herein by reference.

Exemplary viral vectors include lentiviral vectors, AAVs, and retroviral vectors. Lentiviral vectors and AAVs can integrate into the genome without cell divisions, and both types have been tested in pre-clinical animal studies. Methods for preparation of AAVs are described in the art e.g., in U.S. Pat. Nos. 5,677,158, 6,309,634, and 6,683,058, each of which is incorporated herein by reference. Methods for preparation and in vivo administration of lentiviruses are described in US 20020037281 (incorporated herein by reference). Preferably, a lentiviral vector is a replication-defective lentivirus particle. Such a lentivirus particle can be produced from a lentiviral vector comprising a 5' lentiviral LTR, a tRNA binding site, a packaging signal, a promoter operably linked to a polynucleotide signal encoding the fusion protein, an origin of second strand DNA synthesis and a 3' lentiviral LTR.

Retroviruses are most commonly used in human clinical trials, as they carry 7-8 kb, and have the ability to infect cells and have their genetic material stably integrated into the host cell with high efficiency (see, e.g., WO 95/30761; WO 95/24929, each of which is incorporated herein by reference). Preferably, a retroviral vector is replication defective. This prevents further generation of infectious retroviral particles in the target tissue. Thus, the replication defective virus becomes a "captive" transgene stably incorporated into the target cell genome. This is typically accomplished by deleting the gag, env, and pol genes (along with most of the rest of the viral genome). Heterologous nucleic acids are inserted in place of the deleted viral genes. The heterologous genes may be under the control of the endogenous heterologous promoter, another heterologous promoter active in the target cell, or the retroviral 5' LTR (the viral LTR is active in diverse tissues).

These delivery vectors described herein can be made target-specific by attaching, for example, a sugar, a glycolipid, or a protein (e.g., an antibody to a target cell receptor).

Reversible delivery expression systems may also be used. The Cre-loxP or FLP/FRT system and other similar systems can be used for reversible delivery-expression of one or more of the above-described nucleic acids. See WO2005/112620, WO2005/039643, US20050130919, US20030022375, US20020022018, US20030027335, and US20040216178. In particular, the reversible delivery-expression system described in US20100284990 can be used to provide a selective or emergency shut-off.

Non-Viral Delivery Methods

Several non-viral methods exist for delivery of anti-BRG1 and/or BRM agents including polymeric, biodegradable microparticle, or microcapsule delivery devices known in the art. For example, a colloidal dispersion system may be used for targeted delivery an anti-BRG1 and/or BRM agent described herein. Colloidal dispersion systems include macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. Liposomes are artificial membrane vesicles that are useful as delivery vehicles in vitro and in vivo. It has been shown that large unilamellar vesicles (LUV), which range in size from 0.2-4.0 μm can encapsulate a substantial percentage of an aqueous buffer containing large macromolecules.

The composition of the liposome is usually a combination of phospholipids, usually in combination with steroids, especially cholesterol. Other phospholipids or other lipids may also be used. The physical characteristics of liposomes depend on pH, ionic strength, and the presence of divalent cations.

Lipids useful in liposome production include phosphatidyl compounds, such as phosphatidylglycerol, phosphatidylcholine, phosphatidylserine, phosphatidyl-ethanolamine, sphingolipids, cerebrosides, and gangliosides. Exemplary phospholipids include egg phosphatidylcholine, dipalmitoylphosphatidylcholine, and distearoyl-phosphatidylcholine. The targeting of liposomes is also possible based on, for example, organ-specificity, cell-specificity, and organelle-specificity and is known in the art. In the case of a liposomal targeted delivery system, lipid groups can be incorporated into the lipid bilayer of the liposome in order to maintain the targeting ligand in stable association with the liposomal bilayer. Various linking groups can be used for joining the lipid chains to the targeting ligand. Additional methods are known in the art and are described, for example in U.S. Patent Application Publication No. 20060058255.

Pharmaceutical Compositions

The pharmaceutical compositions described herein are preferably formulated into pharmaceutical compositions for administration to human subjects in a biologically compatible form suitable for administration in vivo.

The compounds described herein may be used in the form of the free base, in the form of salts, solvates, and as prodrugs. All forms are within the methods described herein. In accordance with the methods of the invention, the described compounds or salts, solvates, or prodrugs thereof may be administered to a patient in a variety of forms depending on the selected route of administration, as will be understood by those skilled in the art. The compounds described herein may be administered, for example, by oral, parenteral, buccal, sublingual, nasal, rectal, patch, pump, intratumoral, or transdermal administration and the pharmaceutical compositions formulated accordingly. Parenteral administration includes intravenous, intraperitoneal, subcutaneous, intramuscular, transepithelial, nasal, intrapulmonary, intrathecal, rectal, and topical modes of administration. Parenteral administration may be by continuous infusion over a selected period of time.

A compound described herein may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or it may be enclosed in hard or soft shell gelatin capsules, or it may be compressed into tablets, or it may be incorporated directly with the food of the diet. For oral therapeutic administration, a compound described herein may be incorporated with an excipient and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, and wafers. A compound described herein may also be administered parenterally. Solutions of a compound described herein can be prepared in water suitably mixed with a surfactant, such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, DMSO, and mixtures thereof with or without alcohol, and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms. Conventional procedures and ingredients for the selection and preparation of suitable formulations are described, for example, in Remington's Pharmaceutical Sciences (2012, 22nd ed.) and in The United States Pharmacopeia: The National Formulary (USP 41 NF36), published in 2018. The pharmaceutical forms suitable for injectable use include

71

sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases the form must be sterile and must be fluid to the extent that may be easily administered via syringe. Compositions for nasal administration may conveniently be formulated as aerosols, drops, gels, and powders. Aerosol formulations typically include a solution or fine suspension of the active substance in a physiologically acceptable aqueous or non-aqueous solvent and are usually presented in single or multidose quantities in sterile form in a sealed container, which can take the form of a cartridge or refill for use with an atomizing device. Alternatively, the sealed container may be a unitary dispensing device, such as a single dose nasal inhaler or an aerosol dispenser fitted with a metering valve which is intended for disposal after use. Where the dosage form includes an aerosol dispenser, it will contain a propellant, which can be a compressed gas, such as compressed air or an organic propellant, such as fluorochlorohydrocarbon. The aerosol dosage forms can also take the form of a pump-atomizer. Compositions suitable for buccal or sublingual administration include tablets, lozenges, and pastilles, where the active ingredient is formulated with a carrier, such as sugar, acacia, tragacanth, gelatin, and glycerine. Compositions for rectal administration are conveniently in the form of suppositories containing a conventional suppository base, such as cocoa butter. A compound described herein may be administered intratumorally, for example, as an intratumoral injection. Intratumoral injection is injection directly into the tumor vasculature and is specifically contemplated for discrete, solid, accessible tumors. Local, regional, or systemic administration also may be appropriate. A compound described herein may advantageously be contacted by administering an injection or multiple injections to the tumor, spaced for example, at approximately, 1 cm intervals. In the case of surgical intervention, the present invention may be used preoperatively, such as to render an inoperable tumor subject to resection. Continuous administration also may be applied where appropriate, for example, by implanting a catheter into a tumor or into tumor vasculature.

The compounds described herein may be administered to an animal, e.g., a human, alone or in combination with pharmaceutically acceptable carriers, as noted herein, the proportion of which is determined by the solubility and chemical nature of the compound, chosen route of administration, and standard pharmaceutical practice.

Dosages

The dosage of the compounds described herein, and/or compositions including a compound described herein, can vary depending on many factors, such as the pharmacodynamic properties of the compound; the mode of administration; the age, health, and weight of the recipient; the nature and extent of the symptoms; the frequency of the treatment, and the type of concurrent treatment, if any; and the clearance rate of the compound in the animal to be treated. One of skill in the art can determine the appropriate dosage based on the above factors. The compounds described herein may be administered initially in a suitable dosage that may be adjusted as required, depending on the clinical response. In general, satisfactory results may be obtained when the compounds described herein are administered to a human at a daily dosage of, for example, between 0.05 mg and 3000 mg (measured as the solid form).

Alternatively, the dosage amount can be calculated using the body weight of the patient. For example, the dose of a compound, or pharmaceutical composition thereof, administered to a patient may range from 0.1-50 mg/kg.

72


Kits

The invention also features kits including (a) a pharmaceutical composition including an agent that reduces the level and/or activity of BRG1 and/or BRM in a cell or subject described herein, and (b) a package insert with instructions to perform any of the methods described herein. In some embodiments, the kit includes (a) a pharmaceutical composition including an agent that reduces the level and/or activity of BRG1 and/or BRM in a cell or subject described herein, (b) an additional therapeutic agent (e.g., an anti-cancer agent), and (c) a package insert with instructions to perform any of the methods described herein.

EXAMPLES


Example 1. Compound 17

Synthesis of Compound 17: BRG1/BRM Inhibitor Compound 17 has the Structure







Compound 17 was synthesized as shown in Scheme 1 below.

Scheme 1. Synthesis of Compound 17




73

-continued




Step 1: Preparation of
2-bromo-1-(3-bromophenyl)ethanone (Intermediate
B)



To a solution of 1-(3-bromophenyl)ethanone (132.45 mL, 1.00 mol) in CHCl_3 (250 mL) was added Br_2 (77.70 mL, 1.51 mol) in a dropwise manner at 20°C . under N_2 (g). The reaction mixture was subsequently stirred at 80°C . After 1 h, the mixture was cooled to room temperature and concentrated to give intermediate B (279.27 g) as yellow oil, which was used for next step directly.


74

Step 2: Preparation of
4-(3-bromophenyl)thiazol-2-amine (Intermediate C)



To a solution of thiourea (229.23 g, 3.01 mol) in EtOH (1.5 L) was added intermediate B (279 g, 1.00 mol). The reaction mixture was stirred at 85°C . After 2 h, the mixture was cooled to room temperature and concentrated in vacuum to give an oil. The oil was carefully poured into saturated aqueous NaHCO_3 (2 L). The resulting basic (pH ~ 8) solution was extracted with ethyl acetate (3 $\times$ 2 L). The combined organic layers were washed with brine (4 L), dried over Na_2SO_4 , filtered, and concentrated to give an oil. The oil was purified by column chromatography (SiO_2 , PE:EA=5:1) to give intermediate C (130.00 g, 494.40 mmol, 49.25% yield) as a yellow solid. LCMS (ESI) m/z : $[\text{M}+\text{H}]^+=257.0$. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.98 (m, 1H), 7.79 (d, $J=8.0$ Hz, 1H), 7.44-7.42 (m, 1H), 7.33-7.29 (m, 1H), 7.14 (s, 1H), 7.09 (s, 2H).


Step 3: Preparation of 4-[3-(4-pyridyl)phenyl]thiazol-2-amine (Intermediate E)



Intermediate C (20.00 g, 78.40 mmol), 4-pyridylboronic acid (28.9 g, 239.18 mmol), dichloro[1,1'-bis(di-*t*-butylphosphino)ferrocene]palladium(II) (2.56 g, 3.92 mmol) and K_3PO_4 (66.56 g, 313.56 mmol) were diluted in 1,4-dioxane (240 mL) and water (24 mL). The mixture was purged with N_2 (g) three times and then stirred at 80°C . After 7 h, the reaction mixture was cooled to room temperature and water (800 mL) was added. The mixture was extracted with EtOAc (3 $\times$ 800 mL). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated in vacuo. The resulting oil was stirred over a mixture of dichloromethane (30 mL) and MTBE (100 mL). After stirring for 5 min, the precipitate was filtered and washed with MTBE (10 mL) to give intermediate E (16.20 g, 61.17 mmol, 78.03% yield) as a yellow solid. LCMS (ESI) m/z : $[\text{M}+\text{H}]^+=254.0$.


75

Step 4: Preparation (S)-4-(methylthio)-1-oxo-1-((4-(3-(pyridin-4-yl)phenyl)thiazol-2-yl)amino)butan-2-aminium chloride (Intermediate G)



To a mixture of intermediate E (12.60 g, 49.74 mmol) and (2S)-2-(tert-butoxycarbonylamino)-4-methylsulfanylbutanoic acid (18.60 g, 74.61 mmol) in dichloromethane (900 mL) was added EEDQ (24.60 g, 99.48 mmol). After stirring for 2 h at room temperature, the reaction mixture was concentrated in vacuo. The residue was triturated with dichloromethane (100 mL) followed by MeOH (200 mL) to give the intermediate G (11.70 g, 23.73 mmol, 47.71% yield, ee % = 99.44%) as white solids. LCMS (ESI) m/z: [M+H]⁺ = 485.1. ¹H NMR (400 MHz, DMSO) δ 12.39 (s, 1H), 8.68-8.66 (m, 2H), 8.30 (s, 1H), 8.02-7.99 (m, 1H), 7.83 (s, 1H), 7.76-7.74 (m, 3H), 7.61-7.57 (m, 1H), 7.28 (d, J=7.6 Hz, 1H), 4.31-4.30 (m, 1H), 2.65-2.44 (m, 2H), 2.06 (s, 3H), 2.01-1.85 (m, 2H), 1.38 (s, 9H).

Step 5: Preparation of (S)-4-(methylthio)-1-oxo-1-((4-(3-(pyridin-4-yl)phenyl)thiazol-2-yl)amino)butan-2-aminium chloride (Intermediate H)



A mixture of intermediate G (11.50 g, 23.73 mmol) in MeOH (50 mL) was added a solution of 4 M HCl in 1,4-dioxane (100 mL). After stirring for 1 h at room temperature, the mixture was poured into MTBE (1000 mL). The resulting precipitates were filtered to give the intermediate H (9.99 g, 23.73 mmol, 100.00% yield, HCl salt) as a yellow solid. LCMS (ESI) m/z: [M+H]⁺ = 385.0

76


Step 6: Preparation of 4-amino-N-[(1S)-3-methylsulfanyl-1-[[4-[3-(4-pyridyl)phenyl]thiazol-2-yl]carbamoyl]propyl]benzamide (compound 17)

5

G

10

15



17

To a mixture of intermediate H (4.00 g, 9.50 mmol) and 4-aminobenzoic acid (1.30 g, 9.50 mmol) in DMF (40 mL) was sequentially added N,N-diisopropylethylamine (6.62 mL, 38.01 mmol), EDCI (2.73 g, 14.25 mmol) and HOBt (1.93 g, 14.25 mmol). The solution was stirred at 25° C. for 14 h and subsequently poured into water (200 mL). The resulting precipitates were collected by filtration. The solids were triturated in MeOH (200 mL) and filtered. The solids were further purified by column chromatography (SiO₂, DCM:MeOH=80:1-20:1) to give compound 17 (2.13 g, 4.19 mmol, 44.11% yield, ee % = 99.28%) as white solids. LCMS (ESI) m/z: [M+H]⁺ = 504.0. ¹H NMR (400 MHz, DMSO) δ 12.40 (s, 1H), 8.68-8.66 (m, 2H), 8.31-8.30 (m, 1H), 8.22 (d, J=7.2 Hz, 1H), 8.02-7.99 (m, 1H), 7.82 (s, 1H), 7.76-7.74 (m, 3H), 7.67-7.63 (m, 2H), 7.61-7.57 (m, 1H), 6.58-6.54 (m, 2H), 5.67 (s, 2H), 4.72-4.67 (m, 1H), 2.65-2.54 (m, 2H), 2.12-2.06 (m, 5H).

ATPase Activity of Compound 17

The ATPase catalytic activity of BRM or BRG-1 in the presence of compound 17 was measured by the in vitro biochemical assay using ADP-Glo™ (Promega, V9102). The ADP-Glo™ kinase assay is performed in two steps once the reaction is complete. The first step is to deplete any unconsumed ATP in the reaction. The second step is to convert the reaction product ADP to ATP, which will be utilized by the luciferase to generate luminescence and be detected by a luminescence reader, such as Envision.

The assay reaction mixture (10 μL) contains 30 nM of BRM or BRG1, 20 nM salmon sperm DNA (from Invitrogen, UltraPure™ Salmon Sperm DNA Solution, cat #15632011), and 400 μM of ATP in the ATPase assay buffer, which comprises of 20 mM Tris, pH 8, 20 mM MgCl₂, 50 mM NaCl, 0.1% Tween-20, and 1 mM fresh DTT (Pierce™ DTT (Dithiothreitol), cat #20290). The reaction is initiated by the addition of the 2.5 μL ATPase solution to 2.5 μL ATP/DNA solution on low volume white Proxiplate-384 plus plate (PerkinElmer, cat #6008280) and incubates at room temperature for 1 hour. Then following addition of 5 μL of ADP-Glo™ Reagent provided in the kit, the reaction incubates at room temperature for 40 minutes. Then 10 μL of Kinase Detection Reagent provided in the kit is added to convert ADP to ATP, and the reaction incubates at room temperature for 60 minutes. Finally, luminescence measurement is collected with a plate-reading luminometer, such as Envision.

BRM and BRG1 were synthesized from high five insect cell lines with a purity of greater than 90%. Compound 17 was found to have an IP₅₀ of 10.4 nM against BRM and 19.3 nM against BRG1 in the assay.

Example 2. Effects of BRG1/BRM ATPase
Inhibition on the Growth of Cancer Cell Lines

Procedure: Uveal melanoma cell lines (92-1, MP41, MP38, MP46), prostate cancer cell lines (LNCAP), lung cancer cell lines (NCIH1299), and immortalized embryonic kidney lines (HEK293T) were plated into 96 well plates with growth media (see Table 1). BRG1/BRM ATPase inhibitor, compound 17, was dissolved in DMSO and added to the cells in a concentration gradient from 0 to 10 micromolar at the time of plating. Cells were incubated at 37° C. for 3 days. After 3 days of treatment, the media was removed from the cells, and 30 microliters of TrypLE (Gibco) was added to cells for 10 minutes. Cells were detached from the plates and resuspended with the addition of 170 microliters of growth media. Cells from two DMSO-treated control wells were counted, and the initial number of cells plated at the start of the experiment, were re-plated into fresh-compound containing plates for an additional four days at 37° C. At day 7, cells were harvested as described above.

On day 3 and day 7, relative cell growth was measured by the addition of Cell-titer glo (Promega), and luminescence was measured on an Envision plate reader (Perkin Elmer). The concentration of compound 17 at which each cell line's growth was inhibited by 50% (GI₅₀) was calculated using Graphpad Prism and is plotted in FIG. 1.

For multiple myeloma cell lines (OPM2, MM1S, LP1), ALL cell lines (TALL1, JURKAT, RS411), DLBCL cell lines (SUDHL6, SUDHL4, DB, WSUDLCL2, PFEIFFER), AML cell lines (OCIAML5), MDS cell lines (SKM1), ovarian cancer cell lines (OV7, TYKNU), esophageal cancer cell lines (KYSE150), rhabdoid tumor lines (RD, G402, G401, HS729, A204), liver cancer cell lines (HLF, HLE, PLCRPF5), and lung cancer cell lines (SW1573, NCIH2444), the above methods were performed with the following modifications: Cells were plated in 96 well plates, and the next day, BRG1/BRM ATPase inhibitor, compound 17, was dissolved in DMSO and added to the cells in a concentration gradient from 0 to 10 micromolar. At the time of cell splitting on days 3 and 7, cells were split into new 96 well plates, and fresh compound was added four hours after re-plating.

Table 1 lists the tested cell lines and growth media used.

TABLE 1

Cell Lines and Growth Media		
Cell Line	Source	Growth Media
92-1	SIGMA	RPMI1640 + 20% FBS
A204	ATCC	McCoy's 5A + 10% FBS
DB	ATCC	RPMI1640 + 10% FBS
G401	ATCC	McCoy's 5A + 10% FBS
G402	ATCC	McCoy's 5A + 10% FBS
HEK293T	ATCC	DMEM + 10% FBS
HLE	JCRB	DMEM + 10% FBS
HLF	JCRB	DMEM + 10% FBS
HS729	ATCC	DMEM + 10% FBS
JURKAT	ATCC	RPMI1640 + 10% FBS
KYSE150	DSMZ	RPMI1640/Ham's F12 + 10% FBS
LNCAP	ATCC	RPMI1640 + 10% FBS
LP1	DSMZ	IMDM + 20% FBS
MM1S	ATCC	RPMI1640 + 10% FBS
MP38	ATCC	RPMI1640 + 20% FBS
MP41	ATCC	RPMI1640 + 20% FBS
MP46	ATCC	RPMI1640 + 20% FBS
NCIH1299	ATCC	RPMI1640 + 10% FBS
NCIH2444	ATCC	RPMI1640 + 20% FBS
OCIAML5	DSMZ	alpha-MEM + 20% FBS + 10 ng/ml GM-CSF
OPM2	DSMZ	RPMI1640 + 10% FBS


TABLE 1-continued

Cell Lines and Growth Media		
Cell Line	Source	Growth Media
OV7	ECACC	DMEM/Ham's F12 (1:1) + 2 mM Glutamine + 10% FBS + 0.5 ug/ml hydrocortisone + 10 ug/ml insulin
PFEIFFER	ATCC	RPMI1640 + 10% FBS
PLCRPF5	ATCC	EMEM + 10% FBS
RD	ATCC	DMEM + 10% FBS
RS411	ATCC	RPMI1640 + 10% FBS
SKM1	JCRB	RPMI1640 + 10% FBS
SUDHL4	DSMZ	RPMI1640 + 10% FBS
SUDHL6	ATCC	RPMI1640 + 20% FBS
SW1573	ATCC	DMEM + 10% FBS
TALL1	JCRB	RPMI1640 + 10% FBS
TYKNU	JCRB	EMEM + 20% FBS
WSUDLCL2	DSMZ	RPMI1640 + 10% FBS

Results: As shown in FIG. 1, the AML cell line was more sensitive to BRG1/BRM inhibition than the other tested cell lines. Inhibition of the AML cell lines was maintained through day 7.


Example 3. Synthesis of Compound 18

BRG1/BRM Inhibitor compound 18 has the structure:




Compound 18 was synthesized as shown in Scheme 2 below.

Scheme 2. Synthesis of Compound 18



79

Step 1: Preparation of (S)-1-(methylsulfonyl)-N-(4-(methylthio)-1-oxo-1-((4-(3-(pyridin-4-yl)phenyl)thiazol-2-yl)amino)butan-2-yl)-1H-pyrrole-3-carboxamide (compound 18)



To a mixture of (S)-4-(methylthio)-1-oxo-1-((4-(3-(pyridin-4-yl)phenyl)thiazol-2-yl)amino)butan-2-aminium chloride (2.00 g, 4.75 mmol) and 1-methylsulfonylpyrrole-3-carboxylic acid (0.899 g, 4.75 mmol) in DMF (20 mL) was added EDCI (1.37 g, 7.13 mmol), HOBt (0.963 g, 7.13 mmol), and N,N-diisopropylethylamine (3.31 mL, 19.00 mmol). After stirring for 3 h, the mixture was poured into water (100 mL) and the resulting precipitates were filtered. The solids were triturated in MeOH (20 mL) and the precipitate was collected by filtration. The solids were re-dissolved in DMSO (10 mL) and poured into MeOH (50 mL). The precipitates were filtered and lyophilized to give Compound 18 (2.05 g, 3.66 mmol, 77.01% yield) as white solids. LCMS (ESI) m/z $[M+H]^+=555.9$. 1H NMR (400 MHz, DMSO) δ 12.49 (s, 1H), 8.68-8.66 (m, 2H), 8.46 (d, $J=7.2$ Hz, 1H), 8.31-8.30 (m, 1H), 8.02-8.00 (m, 1H), 7.94-7.96 (m, 1H), 7.83 (s, 1H), 7.73-7.74 (m, 3H), 7.61-7.57 (m, 1H), 7.31-7.29 (m, 1H), 6.79-6.77 (m, 1H), 4.74-4.69 (m, 1H), 3.57 (s, 3H), 2.67-2.53 (m, 2H), 2.13-2.01 (m, 5H). SFC: AS-3-MeOH (DEA)-40-3 mL-35T-1cm, $t=0.932$ min, ee %=100%.

Example 4. Effects of BRG1/BRM ATPase Inhibition on the Growth of Cancer Cell Lines

Procedure: All cell lines described above in Example 2 were also tested as described above with compound 18. In addition, the following cell lines were also tested as follows. Briefly, for Ewing's sarcoma cell lines (CADOES1, RDES, SKES1), retinoblastoma cell lines (WERIRB1), ALL cell lines (REH), AML cell lines (KASUMI1), prostate cancer cell lines (PC3, DU145, 22RV1), melanoma cell lines (SH4, SKMEL28, WM115, COLO829, SKMEL3, A375), breast cancer cell lines (MDAMB415, CAMA1, MCF7, BT474, HCC1419, DU4475, BT549), B-ALL cell lines (SUPB15), CML cell lines (K562, MEG01), Burkitt's lymphoma cell lines (RAMOS2G64C10, DAUDI), mantle cell lymphoma cell lines (JEK01, REC1), bladder cancer cell lines (HT1197), and lung cancer cell lines (SBC5), the above methods were performed with the following modifications: Cells were plated in 96 well plates, and the next day, BRG1/BRM ATPase inhibitor, compound 18, was dissolved in DMSO and added to the cells in a concentration gradient from 0 to 10 micromolar. At the time of cell splitting on days

80

3 and 7, cells were split into new 96 well plates, and fresh compound was added four hours after re-plating.

Table 2 lists the tested cell lines and growth media used.

TABLE 2

Cell Lines and Growth Media			
Cell Line	Source	Growth Media	
22RV1	ATCC	RPMI1640 + 10% FBS	
A375	ATCC	DMEM + 10% FBS	
BT474	ATCC	HybriCare medium + 1.5 g/L sodium bicarbonate + 10% FBS	
BT549	ATCC	RPMI1640 + 0.023 IU/ml insulin + 10% FBS	
CADOES1	DSMZ	RPMI1640 + 10% FBS	
CAMA1	ATCC	EMEM + 10% FBS	
COLO829	ATCC	RPMI1640 + 10% FBS	
DAUDI	ATCC	RPMI1640 + 10% FBS	
DU145	ATCC	EMEM + 10% FBS	
DU4475	ATCC	RPMI1640 + 10% FBS	
HCC1419	ATCC	RPMI1640 + 10% FBS	
HT1197	ATCC	EMEM + 10% FBS	
JEK01	ATCC	RPMI1640 + 20% FBS	
K562	ATCC	IMDM + 10% FBS	
KASUMI1	ATCC	RPMI1640 + 10% FBS	
MCF7	ATCC	EMEM + 0.01 mg/ml bovine insulin + 10% FBS	
MDAMB415	ATCC	Leibovitz's L-15 + 2 mM L-glutamine + 10 mcg/ml insulin + 10 mcg/ml glutathione + 15% FBS	
MEG01	ATCC	RPMI1640 + 10% FBS	
PC3	ATCC	F-12K + 10% FBS	
RAMOS2G64C10	ATCC	RPMI1640 + 10% FBS	
RDES	ATCC	RPMI1640 + 15% FBS	
REC1	ATCC	RPMI1640 + 10% FBS	
REH	ATCC	RPMI1640 + 10% FBS	
SBC5	JCRB	EMEM + 10% FBS	
SH4	ATCC	DMEM + 10% FBS	
SKES1	ATCC	McCoy's 5A + 15% FBS	
SKMEL28	ATCC	EMEM + 10% FBS	
SKMEL3	ATCC	McCoy's 5A + 15% FBS	
SUPB15	ATCC	IMDM + 4 mM L-glutamine + 1.5 g/L sodium bicarbonate + 0.05 mM 2-mercaptoethanol + 20% FBS	
WERIRB1	ATCC	RPMI1640 + 10% FBS	
WM115	ATCC	EMEM + 10% FBS	

Results: As shown in FIG. 2, the AML cell lines were more sensitive to BRG1/BRM inhibition than the other tested cell lines. Inhibition of the AML cell lines was maintained through day 7.

Example 5. Effects of BRG1/BRM ATPase Inhibition on the Growth of Cancer Cell Lines

Procedure: A pooled cell viability assay was performed using PRISM (Profiling Relative Inhibition Simultaneously in Mixtures) as previously described ("High-throughput identification of genotype-specific cancer vulnerabilities in mixtures of barcoded tumor cell lines", Yu et al, Nature Biotechnology 34, 419-423, 2016), with the following modifications. Cell lines were obtained from the Cancer Cell Line Encyclopedia (CCLE) collection and adapted to RPMI-1640 medium without phenol red, supplemented with 10% heat-inactivated fetal bovine serum (FBS), in order to apply a unique infection and pooling protocol to such a big compendium of cell lines. A lentiviral spin-infection protocol was executed to introduce a 24 nucleotide-barcode in each cell line, with an estimated multiplicity of infection (MOI) of 1 for all cell lines, using blasticidin as selection marker. Over 750 PRISM cancer cell lines stably barcoded were then


81

pooled together according to doubling time in pools of 25. For the screen execution, instead of plating a pool of 25 cell lines in each well as previously described (Yu et al.), all the adherent or all the suspension cell line pools were plated together using T25 flasks (100,000 cells/flask) or 6-well plates (50,000 cells/well), respectively. Cells were treated with either DMSO or compound in a 8-point 3-fold dose response in triplicate, starting from a top concentration of 10 μ M. As control for assay robustness, cells were treated in parallel with two previously validated compounds, the pan-Raf inhibitor AZ-628, and the proteasome inhibitor bortezomib, using a top concentration of 2.5 μ M and 0.039 μ M, respectively.

Following 3 days of treatment with compounds, cells were lysed, genomic DNA was extracted, barcodes were amplified by PCR and detected with Next-Generation Sequencing. Cell viability was determined by comparing the counts of cell-line specific barcodes in treated samples to those in the DMSO-control and Day 0 control. Dose-response curves were fit for each cell line and corresponding area under the curves (AUCs) were calculated and compared to the median AUC of all cell lines (FIG. 3). Cell lines with AUCs less than the median were considered most sensitive.


Example 6. Synthesis of Compound 19

BRG1/BRM inhibitor compound 19 has the structure:




Compound 19 was synthesized as shown in Scheme 3 below.

Scheme 3. Synthesis of Compound 19




82

-continued



Step 1: Preparation of 6-fluoropyridine-2-carbonyl chloride (Intermediate L)




To a cooled (0° C.) solution of 6-fluoropyridine-2-carboxylic acid (50.00 g, 354.36 mmol) in dichloromethane (500 mL) and N,N-dimethylformamide (0.26 mL, 3.54 mmol) was added oxalyl chloride (155.10 mL, 1.77 mol). After complete addition of oxalyl chloride, the reaction mixture was warmed to room temperature and stirred for an additional 0.5 h. The mixture was subsequently concentrated

83


in vacuo to give intermediate L (56.50 g) as white solids, which were used to next step without further purification.

Step 2: Preparation of
2-chloro-1-(6-fluoro-2-pyridyl)ethanone
(Intermediate M)



To a cooled (0° C.) mixture of intermediate L (56.00 g, 351.00 mmol) in 1,4-dioxane (800 mL) was added in a dropwise manner a solution of 2 M trimethylsilyl diazomethane in hexanes (351 mL). The resulting reaction mixture was stirred at 25° C. for 10 h. The reaction mixture was subsequently quenched with a solution of 4 M HCl in 1,4-dioxane (500 mL). After stirring for 2 h, the reaction solution was concentrated in vacuo to give an oil. The residue was diluted with saturated aqueous NaHCO₃ (500 mL) and extracted with EtOAc (3×200 mL). The combined organic layers were washed with brine (2×300 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give intermediate M (35.50 g) as white solids, which was used to next step directly. LCMS (ESI) m/z: [M+H]⁺=173.8.


Step 3: Preparation of
4-(6-fluoro-2-pyridyl)thiazol-2-amine (Intermediate O)



To a solution of intermediate M (35.50 g, 204.53 mmol) and thiourea (14.01 g, 184.07 mmol) in a mixture of MeOH (250 mL) and water (250 mL) at room temperature was added NaF (3.56 g, 84.82 mmol). After stirring for 30 min, the reaction mixture was partially concentrated in vacuo to remove MeOH. The resulting solution was acidified to pH ~3 with 2 M aqueous HCl and extracted with EtOAc (3×200 mL). The combined organic layers were discarded, and the aqueous phase was basified with saturated aqueous NaHCO₃ (500 mL). After stirring for 30 min, the aqueous phase was extracted with EtOAc (3×325 mL). The combined organic layers were washed with brine (3×225 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The solids were triturated with petroleum ether (300 mL), stirred at 25° C. for 10 min, and filtered. The solids were dried under vacuum to give intermediate O (28.00 g, 143.43 mmol, 70.13% yield, 100% purity) as white solids. LCMS (ESI) m/z: [M+H]⁺=195.8; ¹H NMR (400 MHz, DMSO-d₆) δ 8.00-7.96 (m, 1H), 7.72 (d, J=7.2 Hz, 1H), 7.24 (s, 1H), 7.16 (s, 2H), 7.02 (d, J=8.0 Hz, 1H).


84

Step 4: Preparation of tert-butyl N-[2-[[4-(6-fluoro-2-pyridyl)thiazol-2-yl]amino]-2-oxo-ethyl]carbamate
(Intermediate P)




To a solution of N-Boc-glycine (5.92 g, 33.81 mmol), HATU (12.86 g, 33.81 mmol), and N,N-diisopropylethylamine (21.41 mL, 122.94 mmol) in dichloromethane (100 mL) was added intermediate O (6.00 g, 30.74 mmol). After stirring for 2 h, the reaction mixture was concentrated. The resulting oil was diluted with water (100 mL) and subsequently extracted with EtOAc (4×60 mL). The combined organic layers were washed with brine (2×100 mL), dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give solids. The solids were triturated with a 1:1 mixture of petroleum ether and MeOH (40 mL). After stirring at 25° C. for 20 minutes, the suspension was filtered, and the filter cake was washed with MTBE (20 mL). The solids were dried in vacuo to give intermediate P (7.70 g, 21.63 mmol, 70.4% yield, 99.0% purity) as white solids. LCMS (ESI) m/z: [M+H]⁺=353.1.


Step 5: Preparation of 2-((4-(6-fluoropyridin-2-yl)thiazol-2-yl)amino)-2-oxoethan-1-aminium chloride
(Intermediate Q)



A solution of intermediate P (5.40 g, 15.32 mmol) in 4 M HCl in 1,4-dioxane (35 mL) was stirred at 25° C. for 1.5 h. The reaction mixture was subsequently concentrated under vacuum to give intermediate Q (4.42 g) as white solids, which were used to next step directly without further purification. LCMS (ESI) m/z: [M+H]⁺=252.9.


Step 6: Preparation of 1-tert-butyl-N-[2-[[4-(6-fluoro-2-pyridyl)thiazol-2-yl]amino]-2-oxo-ethyl]pyrrole-3-carboxamide (Intermediate S)





87

88




Step 1: Preparation of 6-fluoropyridine-2-carboxyl chloride (Intermediate B)

40


Step 2: Preparation of 2-chloro-1-(6-fluoro-2-pyridyl)ethanone (Intermediate C)

B 45

C



50



To a cooled (0° C.) solution of 6-fluoropyridine-2-carboxylic acid (50.0 g, 354 mmol) in dichloromethane (500 mL) and N,N-dimethylformamide (0.26 mL, 3.54 mmol) was added oxalyl chloride (155 mL, 1.77 mol). After complete addition of oxalyl chloride, the reaction mixture was warmed to room temperature. After 0.5 hours, the mixture was concentrated under vacuum to give Intermediate B (56.50 g) as a white solid, which was used in the next step without further purification.

55

60


65

To a cooled (0° C.) mixture of Intermediate B (56.0 g, 351 mmol) in 1,4-dioxane (800 mL) was added in a dropwise manner a solution of 2M trimethylsilyl diazomethane in hexanes (351 mL, 702 mmol). The resulting reaction mixture was stirred at 25° C. for 10 h. The reaction mixture was subsequently quenched with a solution of 4M HCl in 1,4-dioxane (500 mL, 2.0 mol). After stirring for 2 h, the reaction solution was concentrated under vacuum to give an oil. The residue was diluted with saturated aqueous NaHCO₃ and extracted three times with ethyl acetate. The combined organic layers were washed twice with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to give Intermediate C (35.5 g) as a white solid, which was used to next step directly.

LCMS (ESI) m/z: [M+H]⁺=173.8.

89

Step 3: Preparation of
4-(6-fluoro-2-pyridyl)thiazol-2-amine (Intermediate
E)




To a solution of Intermediate C (35.5 g, 205 mmol) and thiourea (14.0 g, 184 mmol) in a mixture of methanol (250 mL) and water (250 mL) at room temperature was added NaF (3.56 g, 84.8 mmol). After stirring for 0.5 h, the reaction mixture was partially concentrated under vacuum to remove MeOH, and the resulting solution was acidified to pH ~3 with aqueous 2M HCl. After 15 minutes, the solution was extracted three times with ethyl acetate. The organic layers were discarded and the aqueous phase was alkalized with saturated aqueous NaHCO₃ and stirred for 30 minutes, and extracted three times with ethyl acetate. The combined organic layers were washed three times with brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure. The residue was triturated with petroleum ether and stirred at 25° C. for 10 minutes and filtered. The resultant solids were dried under vacuum to give Intermediate E (28.0 g, 143 mmol, 70.1% yield, 100% purity) as a white solid.

LCMS (ESI) m/z: [M+H]⁺=195.8.

¹H NMR (400 MHz, DMSO-d₆) δ 8.00-7.96 (m, 1H), 7.72 (d, J=7.2 Hz, 1H), 7.24 (s, 1H), 7.16 (s, 2H), 7.02 (d, J=8.0 Hz, 1H).

Step 4: Preparation of 4-[6-(cis-2,6-dimethylmorpholin-4-yl)-2-pyridyl]thiazol-2-amine (Intermediate G)



Ten separate mixtures of Intermediate E (2.00 g, 10.3 mmol), cis-2,6-dimethylmorpholine (3.54 g, 30.7 mmol), and DIPEA (5.35 mL, 30.7 mmol) in dimethyl sulfoxide (10 mL) were stirred in parallel at 120° C. under N₂ atmosphere. After 36 h, the reaction mixtures were combined and added dropwise to water. The resulting suspension was filtered and the filter cake was washed three times with water and once with petroleum ether, then dried over under reduced pressure to give Intermediate G (25.5 g, 87.8 mmol, 95.2% yield) as a yellow solid.


LCMS (ESI) m/z: [M+H]⁺=291.2.

¹H NMR (400 MHz, DMSO-d₆) δ 7.56-7.54 (m, 1H), 7.17 (s, 1H), 7.13 (d, J=7.6 Hz, 1H), 7.01 (s, 2H), 6.72 (d,

90

J=8.8 Hz, 1H), 4.26-4.15 (m, 2H), 3.67-3.55 (m, 2H), 2.38-2.34 (m, 2H), 1.17 (d, J=6.4 Hz, 6H).

Step 5: Preparation of tert-butyl N-[(1S)-2-[[4-[6-(cis-2,6-dimethylmorpholin-4-yl)-2-pyridyl]thiazol-2-yl]amino]-1-(methoxymethyl)-2-oxo-ethyl]carbamate (Intermediate I)




To a solution of Intermediate G (12.0 g, 41.3 mmol) and (2S)-2-(tert-butoxycarbonylamino)-3-methoxy-propanoic acid (10.9 g, 49.6 mmol) in dichloromethane (60 mL) was added EEDQ (12.3 g, 49.6 mmol). After stirring at room temperature for 16 h, the reaction mixture was concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate=2:1 to 3:2) to give Intermediate I (20.0 g, 40.7 mmol, 98.5% yield) as a yellow gum.

LCMS (ESI) m/z: [M+H]⁺=492.2.

¹H NMR (400 MHz, DMSO-d₆) δ 12.37 (s, 1H), 7.78 (s, 1H), 7.64-7.60 (m, 1H), 7.25 (d, J=7.2 Hz, 1H), 7.16 (d, J=7.2 Hz, 1H), 6.79 (d, J=8.4 Hz, 1H), 4.50-4.48 (m, 1H), 4.25 (d, J=11.6 Hz, 2H), 3.70-3.51 (m, 4H), 3.26 (s, 3H), 2.44-2.40 (m, 2H), 1.39 (s, 9H), 1.18 (d, J=6.4 Hz, 6H).

Step 6: Preparation of (S)-4-(4-(6-(cis-2,6-dimethylmorpholino)pyridin-2-yl)thiazol-2-yl)-1-methoxy-3-oxobutan-2-aminium chloride (Intermediate J)



To a solution of 4M HCl in 1,4-dioxane (200 mL, 800 mmol) was added a solution of Intermediate I (20.0 g, 40.7 mmol) in dichloromethane (50 mL). After stirring at room temperature for 2 h, the mixture was diluted with methyl tert-butyl ether resulting in a suspension. The solid was collected by filtration, washed twice with methyl tert-butyl ether, and dried in vacuo to give Intermediate J (19.0 g) as a yellow solid, which was used in the next step without further purification.

LCMS (ESI) m/z: [M+H]⁺=392.3.


¹H NMR (400 MHz, DMSO-d₆) δ 13.44-12.30 (m, 1H), 8.65 (d, J=4.4 Hz, 3H), 7.87 (s, 1H), 7.66-7.64 (m, 1H), 7.25

91


(d, J=7.2 Hz, 1H), 6.83 (d, J=8.8 Hz, 1H), 4.39-4.30 (m, 1H), 4.25 (d, J=11.6 Hz, 2H), 3.94-3.86 (m, 1H), 3.85-3.77 (m, 1H), 3.69-3.57 (m, 2H), 3.31 (s, 3H), 2.43 (m, 2H), 1.18 (d, J=6.4 Hz, 6H).

Preparation of
1-(methylsulfonyl)-1H-pyrrole-3-carboxylic acid
(Intermediate K)

1-(methylsulfonyl)-1H-pyrrole-3-carboxylic acid was synthesized as shown in Scheme 5 below.



Step A: Preparation of tert-butyl
1H-pyrrole-3-carboxylate (Intermediate N)



To a mixture of tert-butyl-prop-2-enoate (78.6 mL, 542 mmol) and 1-(isocyanomethylsulfonyl)-4-methylbenzene (106 g, 542 mmol) in THE (1300 mL) was added 60% NaH in mineral oil (25.97 g, 649 mmol) slowly at 30° C. over 1 hour and then heated to 70° C. After 2 h, the reaction mixture was poured into saturated aqueous NH₄Cl solution and extracted three times with ethyl acetate. The combined organic phase was washed twice with brine, dried over


92

anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to afford a residue. The residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate=20:1 to 3:1) to afford Intermediate N (41.5 g, 236 mmol, 43% yield) as a yellow solid.

LCMS (ESI) m/z [M+Na]⁺=180.4.

¹H NMR (400 MHz, CDCl₃) δ 8.36 (br s, 1H), 7.35-7.25 (m, 1H), 6.71-6.62 (m, 1H), 6.59-6.49 (m, 1H), 1.48 (s, 9H).


Step B: Preparation of tert-butyl
1-methylsulfonylpyrrole-3-carboxylate (Intermediate O)



To a cooled solution (0° C.) of Intermediate N (40.5 g, 242 mmol) in THE (1500 mL) was added a 1 M solution of NaHMDS (484 mL, 484 mmol). After stirring at 0° C. for 30 min, methanesulfonyl chloride (28.1 mL, 363 mmol) was slowly added and the mixture was warmed to 30° C. After 16 h, the reaction mixture was slowly poured into saturated aqueous NH₄Cl solution and extracted three times with ethyl acetate. The combined organic layers were washed twice with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to afford a residue. The residue was purified by silica gel chromatography (petroleum ether:ethyl acetate=10:1) to afford a yellow solid. The yellow solid was triturated with methyl tert-butyl ether at room temperature, stirred for 20 minutes, filtered, and dried in vacuum to afford Intermediate O (25.7 g, 105 mmol, 43% yield) as a white solid.

¹H NMR (400 MHz, CDCl₃) δ 7.66-7.64 (m, 1H), 7.10-7.08 (m, 1H), 6.73-6.71 (m, 1H), 3.21 (s, 3H), 1.56 (s, 9H).

Step C: Preparation of
1-methylsulfonylpyrrole-3-carboxylic acid
(Intermediate K)




To a mixture of Intermediate O (25.7 g, 105 mmol) in 1,4-dioxane (100 mL) was added a 4M solution of HCl in 1,4-dioxane (400 mL, 1.6 mol) at 15° C. After stirring at 15° C. for 14 h, the reaction mixture was concentrated under reduced pressure to afford a residue. The residue was triturated with methyl tert-butyl ether at 15° C. for 16 h. The mixture was filtered and dried in vacuum to afford Intermediate K (18.7 g, 98.8 mmol, 94% yield) as a white solid.

LCMS (ESI) m/z [M+H]⁺=189.8.

¹H NMR (400 MHz, methanol-d₄) δ 7.78-7.77 (m, 1H), 7.25-7.23 (m, 1H), 6.72-6.70 (m, 1H), 3.37 (s, 3H).

93

Step 7: Preparation of N—((S)-1-((4-(6-(cis-2,6-dimethylmorpholino)pyridin-2-yl)thiazol-2-yl)amino)-3-methoxy-1-oxopropan-2-yl)-1-(methylsulfonyl)-1H-pyrrole-3-carboxamide



To a solution of 1-methylsulfonylpyrrole-3-carboxylic acid (Intermediate K) (2.43 g, 12.9 mmol), EDCI (2.69 g, 14.0 mmol), HOBt (1.89 g, 14.0 mmol), and DIPEA (10.2 mL, 58.4 mmol) in dichloromethane (50 mL) was added Intermediate J (5.00 g, 11.7 mmol). After stirring at room temperature for 4 h, the reaction mixture was concentrated under reduced pressure. The residue was diluted with water and extracted three times with ethyl acetate. The combined organic layers were washed three times with saturated aqueous NH_4Cl , once with brine, dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to give a residue. The residue was purified by silica gel column chromatography (petroleum ether:ethyl acetate=1:1 to 1:2). The residue was triturated with methyl tert-butyl ether. After 0.5 h, the suspension was filtered, the filter cake was washed with methyl tert-butyl ether, and dried in vacuo. The solid was dissolved in dimethyl sulfoxide (12 mL) and added dropwise to water (800 mL). The suspension was filtered to give wet filter cake. The filter cake was suspended in water and stirred at room temperature. After 1 hour, the solid was collected by filtration, washed three times with water and dried in vacuo to give N—((S)-1-((4-(6-(cis-2,6-dimethylmorpholino)pyridin-2-yl)thiazol-2-yl)amino)-3-methoxy-1-oxopropan-2-yl)-1-(methylsulfonyl)-1H-pyrrole-3-carboxamide (3.9 g, 6.93 mmol, 59.3% yield) as a white solid.

LCMS (ESI) m/z : $[\text{M}+\text{H}]^+=563.1$.

^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 12.49 (br s, 1H), 8.51 (d, $J=7.2$ Hz, 1H), 7.98-7.97 (m, 1H), 7.78 (s, 1H), 7.67-7.57

94

(m, 1H), 7.29-7.27 (m, 1H), 7.26 (d, $J=7.2$ Hz, 1H), 6.88-6.74 (m, 2H), 4.94-4.91 (m, 1H), 4.25 (d, $J=11.6$ Hz, 2H), 3.77-3.67 (m, 2H), 3.63-3.62 (m, 2H), 3.57 (s, 3H), 3.31 (s, 3H), 2.44-2.38 (m, 2H), 1.18 (d, $J=6.0$ Hz, 6H).

5

Example 9. BRM/BRG1 Inhibition Causes Tumor Growth Delay in an OCI-AML2 Tumor that Carries a DNMT3A Mutation

Method: SCID mice (Charles River, Wilmington) were inoculated subcutaneously on the right flank with the single cell suspension of OCI-AML2 human AML tumor cells (1×10^7) in 100 μL in RPMI-1640 media. When tumor size reach $\sim 100 \text{ mm}^3$, the mice were randomized into either Vehicle group (20% HP- β -CD) or Treatment group (Compound 20 at 1.5 mg/kg daily for 21 days per oral route). All dose volumes were adjusted by body weights in terms of mg/kg.

Results: As shown in FIG. 6, treatment with Compound 20 at 1.5 mg/kg inhibited the growth of OCI-AML2 tumor. All treatments were well tolerated based on % body weight change observed.

OTHER EMBODIMENTS

All publications, patents, and patent applications mentioned in this specification are incorporated herein by reference in their entirety to the same extent as if each individual publication, patent, or patent application was specifically and individually indicated to be incorporated by reference in its entirety. Where a term in the present application is found to be defined differently in a document incorporated herein by reference, the definition provided herein is to serve as the definition for the term.

While the invention has been described in connection with specific embodiments thereof, it will be understood that invention is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure that come within known or customary practice within the art to which the invention pertains and may be applied to the essential features hereinbefore set forth, and follows in the scope of the claims.

Other embodiments are in the claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1

<211> LENGTH: 5589

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

ggcgggggag ggcgggggaa gtcgacggcg cggcgggctc ctgcaggagg ccactgtctg      60
cagctcccggt gaagatgtcc actccagacc caccctggg cggaactcct cggccaggtc      120
cttccccggg ccctggccct tccctggag ccatgctggg ccctagcccg ggtccctcgc      180
cgggctccgc ccacagcatg atggggccca gccagggcc gccctcagca ggacacccca      240
tccccacca ggggctgga ggggtaccctc aggacaacat gcaccagatg cacaagccca      300
tggagtcctat gcatgagaag ggcattgctgg acgacccgcg ctacaaccag atgaaaggaa      360
tggggatgctg gtcagggggc catgctggga tggggccccc gccagcccc atggaccagg      420

```

-continued

actcccaagg ttacccctcg cccctgggtg gctctgagca tgectctagt ccagttccag	480
ccagtggccc gtcttcgggg ccccagatgt cttccgggcc aggaggtgcc ccgctggatg	540
gtgctgaccc ccaggccttg gggcagcaga accggggccc aacccattt aaccagaacc	600
agctgcacca gctcagagct cagatcatgg cctacaagat gctggccagg gggcagcccc	660
tccccgacca cctgcagatg gcggtgcagg gcaagcggcc gatgccggg atgcagcagc	720
agatgccaac gctacctcca cctcgggtgt ccgcaacagg acccggccct ggccctggcc	780
ctggccccgg cccgggtccc gggccggcac ctccaaatta cagcaggcct catggtatgg	840
gagggcccaa catgctccc ccaggaccct cgggcgtgcc ccccgggatg ccaggccagc	900
ctcctggagg gcctcccaag ccttggcctg aaggacccat gggaatgct gctgccccca	960
cgagcacccc tcagaagctg attccccgc agccaacggg ccgcccctcc cccgcgcccc	1020
ctgcctgcc acccgccgcc tcgcccgtga tgccaccgca gaccagtc ccggggcagc	1080
cggcccgacc cgcccccatt gtgccactgc accagaagca gagccgcatc acccccatcc	1140
agaagccgcg gggcctcgac cctgtggaga tcttgcagga gcgcgagtac aggttgcagg	1200
ctcgcatcgc acaccgaatt caggaacttg aaaaccttc cgggtccctg gccggggatt	1260
tgcaaccaa agcgaccatt gagctcaagg cctcaggct gctgaactc cagaggcagc	1320
tgcccgagga ggtggtggtg tgcatcgga gggacacagc gctggagaca gccctcaatg	1380
ctaaggccta caagcgcagc aagcgccagt ccttgcgcga gggccgcatc actgagaagc	1440
tggagaagca gcagaagatc gagcaggagc gcaagcgccg gcagaagcac caggaatacc	1500
tcaatagcat tctccagcat gccaggatt tcaaggaata tcacagatcc gtcacaggca	1560
aaatccagaa gctgaccaag gcagtggcca cgtaccatgc caacacggag cgggagcaga	1620
agaaagagaa cgagcggatc gagaaggagc gcatcgagg gctcatggct gaagatgagg	1680
aggggtaccg caagctcatc gaccagaaga aggacaagcg cctggcctac ctcttgcagc	1740
agacagacga gtacgtggct aacctcacgg agctggtgcg gcagcacaag gctgcccagg	1800
tcgccaagga gaaaaagaag aaaaagaaaa agaagaaggc agaaaatgca gaaggacaga	1860
cgctgccat tgggcccgat ggcgagcctc tggacgagac cagccagatg agcgacctcc	1920
cggatgaagt gatccacgtg gagagtggga agatcctcac aggcacagat gcccacaaag	1980
ccgggcagct ggaggcctgg ctcgagatga acccggggta tgaagtagct ccgaggtctg	2040
atagtgaaga aagtggctca gaagaaggg aagaggagga ggagggaagag cagccgcagg	2100
cagcacagcc tcccacctg cccgtggagg agaagaaga gattccagat ccagacagcg	2160
atgacgtctc tgagggtggc gcgcggcaca tcattgagaa tgccaagcaa gatgtcgatg	2220
atgaatatgg cgtgtcccag gcccttgac gtggcctgca gtctactat gccgtggccc	2280
atgctgtcac tgagagagtg gacaagcagt cagcgttat ggtcaatggt gtccctcaaac	2340
agtaccagat caaaggtttg gagtggtggt gtgcctgta caacaacaac ctgaacggca	2400
tcttggccga cgagatgggc ctggggaaga ccatccagac catcgcgctc atcacgtacc	2460
tcatggagca caaacgcac aatgggcctt tctcatcat cgtgcctctc tcaacgctgt	2520
ccaactgggc gtacgagttt gacaagtggg cccctccgt ggtgaagggtg tcttacaagg	2580
gatccccagc agcaagacgg gcctttgtcc cccagctccg gagtgggaag tcaacgtct	2640
tgctgacgac gtacgagtac atcatcaaag acaagcacat cctcgccaag atccgttgga	2700
agtacatgat tgtggacgaa ggtcacgca tgaagaacca cactgcaag ctgacgcagg	2760

-continued

tgctcaacac gcaactatgtg gcaccccgcc gcctgctgct gacgggcaca ccgctgcaga	2820
acaagcttcc cgagctctgg gcgctgctca acttcctgct gccaccatc ttcaagagct	2880
gcagcacctt cgagcagtggt tttaacgcac cctttgccat gaccggggaa aaggtggacc	2940
tgaatgagga ggaaaccatt ctcatcatcc ggcgctctcca caaagtgtg cggcccttct	3000
tgctccgacg actcaagaag gaagtcgagg ccagttgcc cgaaaagggt gagtacgtca	3060
tcaagtgcga catgtctgcg ctgcagcgag tgctctaccg ccacatgcag gccaggggcg	3120
tgctgctgac tgatggctcc gagaaggaca agaagggcaa aggcggcacc aagaccctga	3180
tgaacaccat catgcagctg cggaagatct gcaaccaccc ctacatgttc cagcacatcg	3240
aggagtcctt ttccgagcac ttgggggttca ctggcgccat tgtccaagggt ctggacctgt	3300
accgagcctc gggtaaattt gagcttcttg atagaattct tcccaaactc cgagcaacca	3360
accacaaaagt gctgctgttc tgccaaatga cctccctcat gaccatcatg gaagattact	3420
ttgcgtatcg cggttttaaa tacctcaggc ttgatggaac cacgaaggcg gaggaccggg	3480
gcatgctgct gaaaaccttc aacgagcccg gctctgagta cttcatcttc ctgctcagca	3540
cccggtctgg ggggctcggc ctgaacctcc agtcggcaga cactgtgatc atttttgaca	3600
gcgactggaa tcctcaccag gacctgcaag cgcaggaccg agccaccgc atcgggcagc	3660
agaacgaggt gcgtgtgctc cgcctctgca ccgtcaacag cgtggaggag aagatcctag	3720
ctgcagccaa gtacaagctc aacgtggacc agaaggatgat ccaggccggc atgttcgacc	3780
agaagtcttc cagccatgag cggcgcgctt tcttgcaggc catcctggag cagcaggagc	3840
aggatgagag cagacactgc agcacgggca gcggcagtg cagcttcgcc cacttgccc	3900
ctccgccagc gggcgtaaac ccgacttggt aggagccacc tctaaaggag gaagacgagg	3960
tgcccgacga cgagaccgtc aaccagatga tcgcccgga cgaggaggag ttgatctgt	4020
tcatgcgcat ggacctggac cgcaggcgcg aggaggcccg caacccaag cggaagccgc	4080
gcctcatgga ggaggacgag ctccctcgt ggatcatcaa ggacgacgag gaggtggagc	4140
ggctgacctg tgaggaggag gaggagaaga tgttcggccg tggtcccg caccgcaagg	4200
agggtgacta cagcgactca ctgacggaga agcagtggct caagaaaatt acaggaaaag	4260
atatccatga cacagccagc agtgtggcac gtgggttaca attccagcgt ggcttcagt	4320
tctgcacacg tgcgtcaaag gccatcgagg agggcacgct ggaggagatc gaagaggagg	4380
tcgggcagaa gaaatcatca cggaagcgca agcgagacag cgacgccggc tcctccacc	4440
cgaccaccag caccgcgagc cgcgacaagg acgacgagag caagaagcag aagaagcgcg	4500
ggcgcccgcc tgccgagaaa ctctcccta accacccaa cctcaccaag aagatgaaga	4560
agattgtgga tgccgtgatc aagtacaagg acagcagcag tggacgtcag ctacgcgagg	4620
tcttcatcca gctgccctcg cgaaaggagc tgcccgagta ctacgagctc atccgcaagc	4680
ccgtggactt caagaagata aaggagcgca ttcgcaacca caagtaccgc agcctcaacg	4740
acctagagaa ggacgtcatg ctctgtgccc agaacgcaca gacctcaac ctggagggt	4800
ccctgatcta tgaagactcc atcgctctgc agtcggtctt caccagcgtg cggcagaaaa	4860
tcgagaagga ggatgacagt gaaggcgagg agagtgagga ggagggaagag ggcgaggagg	4920
aaggctccga atccgaatct cgttcgtca aagtgaagat caagcttggc cggaaggaga	4980
aggcacagga ccggtgaag ggcggccggc ggcggccgag ccgagggtcc cgagccaagc	5040
cggtcgtgag tgacgatgac agtgaggagg aacaagagga ggaccgctca ggaagtggca	5100
cggaagaaga ctgagcccg acattccagt ctcgaccccg agccctcgt tccagagctg	5160

-continued

```

agatggcata ggccttagca gtaacgggta gcagcagatg tagtttcaga cttggagtaa 5220
aactgtataa acaaaagaat ctcccatatt tatacagcag agaagctgta ggactgtttg 5280
tgactggccc tgtcctggca tcagtagcat ctgtaacagc attaactgtc ttaaagagag 5340
agagagagaa ttccgaattg gggaacacac gatacctgtt tttcttttcc gttgtgggca 5400
gtactgttgc gccgcagttt ggagtcactg tagttaagtg tggatgcatg tgcgtcaccg 5460
tccactccto ctactgtatt ttattggaca ggtcagactc gccggggggc cggcgagggg 5520
atgtcagtgt cactggatgt caaacagtaa taaattaac caacaacaaa acgcacagcc 5580
aaaaaaaaa 5589

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1647
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Met Ser Thr Pro Asp Pro Pro Leu Gly Gly Thr Pro Arg Pro Gly Pro
1      5      10     15
Ser Pro Gly Pro Gly Pro Ser Pro Gly Ala Met Leu Gly Pro Ser Pro
20     25     30
Gly Pro Ser Pro Gly Ser Ala His Ser Met Met Gly Pro Ser Pro Gly
35     40     45
Pro Pro Ser Ala Gly His Pro Ile Pro Thr Gln Gly Pro Gly Gly Tyr
50     55     60
Pro Gln Asp Asn Met His Gln Met His Lys Pro Met Glu Ser Met His
65     70     75     80
Glu Lys Gly Met Ser Asp Asp Pro Arg Tyr Asn Gln Met Lys Gly Met
85     90     95
Gly Met Arg Ser Gly Gly His Ala Gly Met Gly Pro Pro Pro Ser Pro
100    105    110
Met Asp Gln His Ser Gln Gly Tyr Pro Ser Pro Leu Gly Gly Ser Glu
115    120    125
His Ala Ser Ser Pro Val Pro Ala Ser Gly Pro Ser Ser Gly Pro Gln
130    135    140
Met Ser Ser Gly Pro Gly Gly Ala Pro Leu Asp Gly Ala Asp Pro Gln
145    150    155    160
Ala Leu Gly Gln Gln Asn Arg Gly Pro Thr Pro Phe Asn Gln Asn Gln
165    170    175
Leu His Gln Leu Arg Ala Gln Ile Met Ala Tyr Lys Met Leu Ala Arg
180    185    190
Gly Gln Pro Leu Pro Asp His Leu Gln Met Ala Val Gln Gly Lys Arg
195    200    205
Pro Met Pro Gly Met Gln Gln Gln Met Pro Thr Leu Pro Pro Pro Ser
210    215    220
Val Ser Ala Thr Gly Pro Gly Pro Gly Pro Gly Pro Gly Pro Gly Pro
225    230    235    240
Gly Pro Gly Pro Ala Pro Pro Asn Tyr Ser Arg Pro His Gly Met Gly
245    250    255
Gly Pro Asn Met Pro Pro Pro Gly Pro Ser Gly Val Pro Pro Gly Met
260    265    270
Pro Gly Gln Pro Pro Gly Gly Pro Pro Lys Pro Trp Pro Glu Gly Pro
275    280    285

```

Met 290	Asn	Ala	Ala	Ala	Ala	Pro 295	Thr	Ser	Thr	Pro	Gln 300	Lys	Leu	Ile	Pro
Pro 305	Gln	Pro	Thr	Gly	Arg 310	Pro	Ser	Pro	Ala	Pro 315	Pro	Ala	Val	Pro	Pro 320
Ala	Ala	Ser	Pro	Val 325	Met	Pro	Pro	Gln	Thr 330	Gln	Ser	Pro	Gly	Gln 335	Pro
Ala	Gln	Pro	Ala	Pro	Met	Val	Pro	Leu	His 345	Gln	Lys	Gln	Ser	Arg	Ile
Thr	Pro	Ile	Gln	Lys	Pro	Arg	Gly 360	Leu	Asp	Pro	Val	Glu 365	Ile	Leu	Gln
Glu 370	Arg	Glu	Tyr	Arg	Leu	Gln 375	Ala	Arg	Ile	Ala	His 380	Arg	Ile	Gln	Glu
Leu 385	Glu	Asn	Leu	Pro	Gly 390	Ser	Leu	Ala	Gly	Asp 395	Leu	Arg	Thr	Lys	Ala 400
Thr	Ile	Glu	Leu	Lys 405	Ala	Leu	Arg	Leu	Leu 410	Asn	Phe	Gln	Arg	Gln 415	Leu
Arg	Gln	Glu	Val	Val	Val	Cys	Met	Arg	Arg	Asp	Thr	Ala	Leu	Glu	Thr
Ala	Leu	Asn	Ala	Lys	Ala	Tyr	Lys 440	Arg	Ser	Lys	Arg	Gln 445	Ser	Leu	Arg
Glu 450	Ala	Arg	Ile	Thr	Glu	Lys 455	Leu	Glu	Lys	Gln	Gln	Lys	Ile	Glu	Gln
Glu 465	Arg	Lys	Arg	Arg	Gln 470	Lys	His	Gln	Glu	Tyr 475	Leu	Asn	Ser	Ile	Leu 480
Gln	His	Ala	Lys	Asp 485	Phe	Lys	Glu	Tyr	His 490	Arg	Ser	Val	Thr	Gly 495	Lys
Ile	Gln	Lys	Leu	Thr	Lys	Ala	Val	Ala 505	Thr	Tyr	His	Ala	Asn	Thr	Glu
Arg	Glu	Gln	Lys	Lys	Glu	Asn 520	Glu	Arg	Ile	Glu	Lys	Glu 525	Arg	Met	Arg
Arg	Leu	Met	Ala	Glu	Asp 535	Glu	Glu	Gly	Tyr	Arg	Lys 540	Leu	Ile	Asp	Gln
Lys 545	Lys	Asp	Lys	Arg	Leu 550	Ala	Tyr	Leu	Leu	Gln 555	Gln	Thr	Asp	Glu	Tyr 560
Val	Ala	Asn	Leu	Thr 565	Glu	Leu	Val	Arg	Gln 570	His	Lys	Ala	Ala	Gln 575	Val
Ala	Lys	Glu	Lys 580	Lys	Lys	Lys	Lys	Lys 585	Lys	Lys	Lys	Ala	Glu 590	Asn	Ala
Glu	Gly	Gln	Thr	Pro	Ala	Ile	Gly 600	Pro	Asp	Gly	Glu	Pro 605	Leu	Asp	Glu
Thr	Ser	Gln	Met	Ser	Asp 615	Leu	Pro	Val	Lys	Val	Ile 620	His	Val	Glu	Ser
Gly 625	Lys	Ile	Leu	Thr	Gly 630	Thr	Asp	Ala	Pro	Lys 635	Ala	Gly	Gln	Leu	Glu 640
Ala	Trp	Leu	Glu	Met 645	Asn	Pro	Gly	Tyr	Glu 650	Val	Ala	Pro	Arg	Ser	Asp 655
Ser	Glu	Glu	Ser	Gly 660	Ser	Glu	Glu	Glu	Glu 665	Glu	Glu	Glu	Glu	Glu 670	Glu
Gln	Pro	Gln	Ala	Ala	Gln	Pro	Pro	Thr	Leu 680	Pro	Val	Glu 685	Glu	Lys	Lys
Lys 690	Ile	Pro	Asp	Pro	Asp	Ser 695	Asp	Asp	Val	Ser	Glu 700	Val	Asp	Ala	Arg
His	Ile	Ile	Glu	Asn	Ala	Lys	Gln	Asp	Val	Asp	Asp	Glu	Tyr	Gly	Val

-continued

705	710	715	720
Ser Gln Ala Leu Ala Arg Gly Leu Gln Ser Tyr Tyr Ala Val Ala His			
	725	730	735
Ala Val Thr Glu Arg Val Asp Lys Gln Ser Ala Leu Met Val Asn Gly			
	740	745	750
Val Leu Lys Gln Tyr Gln Ile Lys Gly Leu Glu Trp Leu Val Ser Leu			
	755	760	765
Tyr Asn Asn Asn Leu Asn Gly Ile Leu Ala Asp Glu Met Gly Leu Gly			
	770	775	780
Lys Thr Ile Gln Thr Ile Ala Leu Ile Thr Tyr Leu Met Glu His Lys			
	785	790	795
Arg Ile Asn Gly Pro Phe Leu Ile Ile Val Pro Leu Ser Thr Leu Ser			
	805	810	815
Asn Trp Ala Tyr Glu Phe Asp Lys Trp Ala Pro Ser Val Val Lys Val			
	820	825	830
Ser Tyr Lys Gly Ser Pro Ala Ala Arg Arg Ala Phe Val Pro Gln Leu			
	835	840	845
Arg Ser Gly Lys Phe Asn Val Leu Leu Thr Thr Tyr Glu Tyr Ile Ile			
	850	855	860
Lys Asp Lys His Ile Leu Ala Lys Ile Arg Trp Lys Tyr Met Ile Val			
	865	870	875
Asp Glu Gly His Arg Met Lys Asn His His Cys Lys Leu Thr Gln Val			
	885	890	895
Leu Asn Thr His Tyr Val Ala Pro Arg Arg Leu Leu Leu Thr Gly Thr			
	900	905	910
Pro Leu Gln Asn Lys Leu Pro Glu Leu Trp Ala Leu Leu Asn Phe Leu			
	915	920	925
Leu Pro Thr Ile Phe Lys Ser Cys Ser Thr Phe Glu Gln Trp Phe Asn			
	930	935	940
Ala Pro Phe Ala Met Thr Gly Glu Lys Val Asp Leu Asn Glu Glu Glu			
	945	950	955
Thr Ile Leu Ile Ile Arg Arg Leu His Lys Val Leu Arg Pro Phe Leu			
	965	970	975
Leu Arg Arg Leu Lys Lys Glu Val Glu Ala Gln Leu Pro Glu Lys Val			
	980	985	990
Glu Tyr Val Ile Lys Cys Asp Met Ser Ala Leu Gln Arg Val Leu Tyr			
	995	1000	1005
Arg His Met Gln Ala Lys Gly Val Leu Leu Thr Asp Gly Ser Glu			
	1010	1015	1020
Lys Asp Lys Lys Gly Lys Gly Gly Thr Lys Thr Leu Met Asn Thr			
	1025	1030	1035
Ile Met Gln Leu Arg Lys Ile Cys Asn His Pro Tyr Met Phe Gln			
	1040	1045	1050
His Ile Glu Glu Ser Phe Ser Glu His Leu Gly Phe Thr Gly Gly			
	1055	1060	1065
Ile Val Gln Gly Leu Asp Leu Tyr Arg Ala Ser Gly Lys Phe Glu			
	1070	1075	1080
Leu Leu Asp Arg Ile Leu Pro Lys Leu Arg Ala Thr Asn His Lys			
	1085	1090	1095
Val Leu Leu Phe Cys Gln Met Thr Ser Leu Met Thr Ile Met Glu			
	1100	1105	1110
Asp Tyr Phe Ala Tyr Arg Gly Phe Lys Tyr Leu Arg Leu Asp Gly			
	1115	1120	1125

-continued

Thr	Thr	Lys	Ala	Glu	Asp	Arg	Gly	Met	Leu	Leu	Lys	Thr	Phe	Asn
1130						1135					1140			
Glu	Pro	Gly	Ser	Glu	Tyr	Phe	Ile	Phe	Leu	Leu	Ser	Thr	Arg	Ala
1145						1150					1155			
Gly	Gly	Leu	Gly	Leu	Asn	Leu	Gln	Ser	Ala	Asp	Thr	Val	Ile	Ile
1160						1165					1170			
Phe	Asp	Ser	Asp	Trp	Asn	Pro	His	Gln	Asp	Leu	Gln	Ala	Gln	Asp
1175						1180					1185			
Arg	Ala	His	Arg	Ile	Gly	Gln	Gln	Asn	Glu	Val	Arg	Val	Leu	Arg
1190						1195					1200			
Leu	Cys	Thr	Val	Asn	Ser	Val	Glu	Glu	Lys	Ile	Leu	Ala	Ala	Ala
1205						1210					1215			
Lys	Tyr	Lys	Leu	Asn	Val	Asp	Gln	Lys	Val	Ile	Gln	Ala	Gly	Met
1220						1225					1230			
Phe	Asp	Gln	Lys	Ser	Ser	Ser	His	Glu	Arg	Arg	Ala	Phe	Leu	Gln
1235						1240					1245			
Ala	Ile	Leu	Glu	His	Glu	Glu	Gln	Asp	Glu	Ser	Arg	His	Cys	Ser
1250						1255					1260			
Thr	Gly	Ser	Gly	Ser	Ala	Ser	Phe	Ala	His	Thr	Ala	Pro	Pro	Pro
1265						1270					1275			
Ala	Gly	Val	Asn	Pro	Asp	Leu	Glu	Glu	Pro	Pro	Leu	Lys	Glu	Glu
1280						1285					1290			
Asp	Glu	Val	Pro	Asp	Asp	Glu	Thr	Val	Asn	Gln	Met	Ile	Ala	Arg
1295						1300					1305			
His	Glu	Glu	Glu	Phe	Asp	Leu	Phe	Met	Arg	Met	Asp	Leu	Asp	Arg
1310						1315					1320			
Arg	Arg	Glu	Glu	Ala	Arg	Asn	Pro	Lys	Arg	Lys	Pro	Arg	Leu	Met
1325						1330					1335			
Glu	Glu	Asp	Glu	Leu	Pro	Ser	Trp	Ile	Ile	Lys	Asp	Asp	Ala	Glu
1340						1345					1350			
Val	Glu	Arg	Leu	Thr	Cys	Glu	Glu	Glu	Glu	Glu	Lys	Met	Phe	Gly
1355						1360					1365			
Arg	Gly	Ser	Arg	His	Arg	Lys	Glu	Val	Asp	Tyr	Ser	Asp	Ser	Leu
1370						1375					1380			
Thr	Glu	Lys	Gln	Trp	Leu	Lys	Ala	Ile	Glu	Glu	Gly	Thr	Leu	Glu
1385						1390					1395			
Glu	Ile	Glu	Glu	Glu	Val	Arg	Gln	Lys	Lys	Ser	Ser	Arg	Lys	Arg
1400						1405					1410			
Lys	Arg	Asp	Ser	Asp	Ala	Gly	Ser	Ser	Thr	Pro	Thr	Thr	Ser	Thr
1415						1420					1425			
Arg	Ser	Arg	Asp	Lys	Asp	Asp	Glu	Ser	Lys	Lys	Gln	Lys	Lys	Arg
1430						1435					1440			
Gly	Arg	Pro	Pro	Ala	Glu	Lys	Leu	Ser	Pro	Asn	Pro	Pro	Asn	Leu
1445						1450					1455			
Thr	Lys	Lys	Met	Lys	Lys	Ile	Val	Asp	Ala	Val	Ile	Lys	Tyr	Lys
1460						1465					1470			
Asp	Ser	Ser	Ser	Gly	Arg	Gln	Leu	Ser	Glu	Val	Phe	Ile	Gln	Leu
1475						1480					1485			
Pro	Ser	Arg	Lys	Glu	Leu	Pro	Glu	Tyr	Tyr	Glu	Leu	Ile	Arg	Lys
1490						1495					1500			
Pro	Val	Asp	Phe	Lys	Lys	Ile	Lys	Glu	Arg	Ile	Arg	Asn	His	Lys
1505						1510					1515			

-continued

Tyr Arg Ser Leu Asn Asp Leu Glu Lys Asp Val Met Leu Leu Cys
 1520 1525 1530
 Gln Asn Ala Gln Thr Phe Asn Leu Glu Gly Ser Leu Ile Tyr Glu
 1535 1540 1545
 Asp Ser Ile Val Leu Gln Ser Val Phe Thr Ser Val Arg Gln Lys
 1550 1555 1560
 Ile Glu Lys Glu Asp Asp Ser Glu Gly Glu Glu Ser Glu Glu Glu
 1565 1570 1575
 Glu Glu Gly Glu Glu Glu Gly Ser Glu Ser Glu Ser Arg Ser Val
 1580 1585 1590
 Lys Val Lys Ile Lys Leu Gly Arg Lys Glu Lys Ala Gln Asp Arg
 1595 1600 1605
 Leu Lys Gly Gly Arg Arg Arg Pro Ser Arg Gly Ser Arg Ala Lys
 1610 1615 1620
 Pro Val Val Ser Asp Asp Asp Ser Glu Glu Glu Gln Glu Glu Asp
 1625 1630 1635
 Arg Ser Gly Ser Gly Ser Glu Glu Asp
 1640 1645

<210> SEQ ID NO 3

<211> LENGTH: 5892

<212> TYPE: RNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

```

gcgtcttccg cgcgcccgagg aggaggcgag ggtgggacgc tgggaggagc ccgagtttag      60
gaagaggagg ggaaggctgt catcaatgaa gtcatatcca taatctagtc ctctctccct      120
ctgtttctgt actctgggtg actcagagag ggaagagatt cagccagcac actcctcgcg      180
agcaagcatt actctactga ctggcagaga caggagaggt agatgtccac gcccacagac      240
cctggtgcga tgccccaccc agggccttcg ccggggcctg ggccttcccc tgggccaatt      300
cttgggccta gtccaggacc aggaccatcc ccagggttccg tccacagcat gatggggcca      360
agtctggac ctccaagtgt ctcccatcct atgccgacga tgggggtccac agacttccca      420
caggaaggca tgcatacaat gcataagccc atcgatggta tacatgacaa ggggattgta      480
gaagacatcc attgtggatc catgaagggc actggtatgc gaccacctca ccagagcatg      540
ggccctcccc agagtccaat ggatcaacac agccaaggtt atatgtcacc acaccatct      600
ccattaggag cccagagaca cgtctccagc cctatgtctg gaggaggccc aactccacct      660
cagatgccac caagccagcc gggggccctc atcccaggtg atccgagggc catgagccag      720
cccaacagag gtccctcacc ttccagtcct gtccagctgc atcagcttcg agtcagatt      780
ttagcttata aaatgctggc ccgaggccag cccctccccg aaacgctgca gcttgagtc      840
caggggaaaa ggacgttgcc tggttgtag caacaacagc agcagcaaca gcagcagcag      900
cagcagcagc agcagcagca gcagcagcaa cagcagccgc agcagcagcc gccgcaacca      960
cagacgcagc aacaacagca gccggccctt gttaactaca acagaccatc tggcccgggg      1020
ccggagctga gcggcccag cccccgcag aagctgccgg tgcccgcgcc cggcgcccg      1080
ccctcgcccg cgccccccgc agccgcgcag ccgcccgcgg ccgcagtgcc cgggcccctca      1140
gtgccgcagc cggccccggg gcagccctcg cccgtcctcc agctgcagca gaagcagagc      1200
cgcatcagcc ccatacagaa accgcaaggc ctggaccccc tggaaattct gcaagagcgg      1260
gaatacagac ttcaggcccc catagctcat aggatacaag aactggaaaa tctgcctggc      1320

```

-continued

tctttgccac	cagatttaag	aaccaaagca	accgtggaac	taaaagcact	tcggttactc	1380
aatttccagc	gtcagctgag	acaggaggtg	gtggcctgca	tgcgcaggga	cacgaccctg	1440
gagacggctc	tcaactccaa	agcatacaaa	cggagcaagc	gccagactct	gagagaagct	1500
cgcctgaccg	agaagctgga	gaagcagcag	aagattgagc	aggagaggaa	acgccgtcag	1560
aaacaccagg	aatacctgaa	cagtattttg	caacatgcaa	aagattttaa	ggaatatcat	1620
cggctctgtg	cgggaaagat	ccagaagctc	tccaaagcag	tggcaacttg	gcatgccaac	1680
actgaaagag	agcagaagaa	ggagacagag	cggattgaaa	aggagagaat	gcggcgactg	1740
atggctgaag	atgaggaggg	ttatagaaaa	ctgattgatc	aaaagaaaga	caggcgttta	1800
gcttaccttt	tgcagcagac	cgatgagtat	gtagccaatc	tgaccaatct	ggtttgggag	1860
cacaagcaag	cccaggcagc	caaagagaag	aagaagagga	ggaggaggaa	gaagaaggct	1920
gaggagaatg	cagagggtgg	ggagtctgcc	ctgggaccgg	atggagagcc	catagatgag	1980
agcagccaga	tgagtgcact	cctgtcaaaa	gtgactcaca	cagaaacogg	caaggttctg	2040
ttcggaccag	aagcacccaa	agcaagtcag	ctggacgcct	ggctggaaat	gaatcctggt	2100
tatgaagtgt	cccctagatc	tgacagttaa	gagagtgttt	ctgattatga	ggaagaggat	2160
gaggaagaag	agtcacgtag	gcaggaaacc	gaagagaaaa	tactcctgga	tccaaatagc	2220
gaagaagttt	ctgagaagga	tgctaagcag	atcattgaga	cagctaagca	agacgtggat	2280
gatgaataca	gcctgcagta	cagtgccagg	ggctcccagt	cctactacac	cgtggctcat	2340
gccatctcgg	agagggtgga	gaaacagtct	gccctcctaa	ttaatgggac	cctaaagcat	2400
taccagctcc	agggcctgga	atggatgggt	tccctgtata	ataacaactt	gaacggaatc	2460
ttagccgatg	aaatggggct	tggaaagacc	atacagacca	ttgcactcat	catttatctg	2520
atggagcaca	aaagactcaa	tggcccttat	ctcatcattg	ttcccctttc	gactctatct	2580
aactggacat	atgaatttga	caaatgggct	ccttctgtgg	tgaagatttc	ttacaagggt	2640
actcctgcca	tgcgtcgtct	cctgttcccc	cagctacgga	gtggcaaatt	caatgtcctc	2700
ttgactactt	atgagtatat	tataaaagac	aagcacattc	ttgcaaagat	tcggtggaaa	2760
tacatgatag	tggacgaagg	ccaccgaatg	aagaatcacc	actgcaagct	gactcaggtc	2820
ttgaacactc	actatgtggc	cccagaagg	atcctcttga	ctgggacccc	gctgcagaat	2880
aagctccctg	aactctgggc	cctcctcaac	ttcctcctcc	caacaatttt	taagagctgc	2940
agcactttg	aacaatgggt	caatgctcca	tttgccatga	ctggtgaaag	ggtggactta	3000
aatgaagaag	aaactatatt	gatcatcagg	cgtctacata	aggtgttaag	accattttta	3060
ctaaggagac	tgaagaaaga	agttgaatcc	cagcttcccg	aaaaagtgga	atatgtgatc	3120
aagtgtgaca	tgctcagctc	gcagaagatt	ctgtatcgcc	atatgcaagc	caaggggatc	3180
cttctcacag	atggtttctg	gaaagataag	aaggggaaag	gaggtgctaa	gacacttatg	3240
aacactatta	tgcagttgag	aaaaatctgc	aaccacccat	atatgtttca	gcacattgag	3300
gaatcctttg	ctgaacacct	aggetattca	aatggggcca	tcaatggggc	tgaactgtat	3360
cgggcctcag	ggaagtttga	gctgcttgat	cgtattctgc	caaaattgag	agcgactaat	3420
caccgagtgc	tgtttttctg	ccagatgaca	tctctcatga	ccatcatgga	ggattatttt	3480
gcttttcgga	acttccctta	cctacgcctt	gatggcacca	ccaagtctga	agatcgtgct	3540
gctttgtctg	agaaattcaa	tgaacctgga	tcccagtatt	tcattttctt	gctgagcaca	3600
agagctgggt	gcctgggctt	aaatcttcag	gcagctgata	cagtggctcat	ctttgacagc	3660
gactggaatc	ctcatcagga	tctgcaggcc	caagaccgag	ctcaccgcac	cgggcagcag	3720

-continued

```

aacgaggtcc gggacttgag gctctgtacc gtgaacagcg tggaggaaaa gatcctcgcg 3780
gccgcaaaat acaagctgaa cgtggatcag aaagtgatcc aggcgggcat gtttgaccaa 3840
aagtcttcaa gccacgagcg gagggcattc ctgcaggcca tcttgagca tgaggaggaa 3900
aatgaggaag aagatgaagt accggacgat gagactctga accaaatgat tgctcgacga 3960
gaagaagaat ttgacctttt tatgcggtat gacatggacc ggcggaggga agatgcccg 4020
aacccgaaac ggaagccccg tttaatggag gaggatgagc tgccctcctg gatcattaag 4080
gatgacgctg aagtagaaag gctcacctgt gaagaagagg aggagaaaa atttgggagg 4140
gggtcccgcc agcgccgtga cgtggactac agtgacgccc tcacggagaa gcagtggcta 4200
agggccatcg aagacggcaa tttggaggaa atggaagagg aagtacggct taagaagcga 4260
aaaagacgaa gaaatgtgga taaagatcct gcaaaagaag atgtgaaaa agctaagaag 4320
agaagaggcc gccctcccg tgagaaactg tcaccaaac ccccaaaact gacaaagcag 4380
atgaacgcta tcacgatac tgtgataaac tacaagata ggtgtaacgt ggagaagggtg 4440
cccgtaatt ctgagttgga aatagaagga aacagttcag ggcgacagct cagtgaagtc 4500
ttcattcagt taccttcaag gaaagaatta ccagaatact atgaattaat taggaagcca 4560
gtggatttca aaaaaataaa ggaaggatt cgtaatcata agtaccggag cctaggcgac 4620
ctggagaagg atgtcatgct tctctgtcac aacgctcaga cgttcaacct ggagggatcc 4680
cagatctatg aagactccat cgtcttacag tcagtgttta agagtgcgc gcagaaaatt 4740
gccaaagagg aagagagtga ggtatgaagc aatgaagagg aggaagagga agatgaagaa 4800
gagtcagagt ccgaggcaaa atcagtcagg gtgaaaatta agctcaataa aaaagatgac 4860
aaaggccggg acaaagggaagg aggcagaaa aggcacaaac gaggaaaagc caaacctgta 4920
gtgagcgatt ttgacagcga tgaggagcag gatgaacgtg aacagtcaga aggaagtggg 4980
acggatgatg agtgatcagt atggacctt ttccttggtg gaactgaatt ccttctctcc 5040
ctgtctcatt tctaccagtg gagttcattt gtcataatagg cactgggttg tttctatatc 5100
atcatcgctc ataaactagc tttaggatag tgccagacaa acatagtata tcatggtgta 5160
aaaaacacac acatacacaa atatttgtaa catatttgta ccaaatgggc ctcaaagatt 5220
cagattgaaa caaacaaaaa gcttttgatg gaaaatatgt ggggtgatag tatatttcta 5280
tgggtgggtc taatttggtg acggtttgat tgtgcctggt tttatcacct gttcagatga 5340
gaagattttt gtcttttgta gcaactgataa ccaggagaag ccattaaaag ccactgggta 5400
ttttattttt catcaggcaa ttttcgaggt ttttatgtgt tcggtattgt tttttacac 5460
tgtgtacatc ataagcaact ttaatagggt ataaatgtac agtagttaga tttcacctgc 5520
atatacattt ttccatttta tgctctatga tctgaacaaa agctttttga attgtataag 5580
atztatgtct actgtaaaca ttgcttaatt tttttgctct tgatttaaaa aaaagttttg 5640
ttgaaagcgc tattgaatat tgcaatctat atagtgtatt ggatggcttc ttttgcacc 5700
ctgatctcct atgttaccaa tgtgtatcgt ctcttctctc ctaaagtga cttaatcttt 5760
gctttctttg cacaatgtct ttggttgcaa gtcataagcc tgaggcaaat aaaattccag 5820
taatttcgaa gaatgtggtg ttggtgcttt cctaataaag aaataattta gcttgacaaa 5880
aaaaaaaaaa aa 5892

```

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

```

Met Ser Thr Pro Thr Asp Pro Gly Ala Met Pro His Pro Gly Pro Ser
1      5      10      15

Pro Gly Pro Gly Pro Ser Pro Gly Pro Ile Leu Gly Pro Ser Pro Gly
20      25      30

Pro Gly Pro Ser Pro Gly Ser Val His Ser Met Met Gly Pro Ser Pro
35      40      45

Gly Pro Pro Ser Val Ser His Pro Met Pro Thr Met Gly Ser Thr Asp
50      55      60

Phe Pro Gln Glu Gly Met His Gln Met His Lys Pro Ile Asp Gly Ile
65      70      75      80

His Asp Lys Gly Ile Val Glu Asp Ile His Cys Gly Ser Met Lys Gly
85      90      95

Thr Gly Met Arg Pro Pro His Pro Gly Met Gly Pro Pro Gln Ser Pro
100     105     110

Met Asp Gln His Ser Gln Gly Tyr Met Ser Pro His Pro Ser Pro Leu
115     120     125

Gly Ala Pro Glu His Val Ser Ser Pro Met Ser Gly Gly Gly Pro Thr
130     135     140

Pro Pro Gln Met Pro Pro Ser Gln Pro Gly Ala Leu Ile Pro Gly Asp
145     150     155     160

Pro Gln Ala Met Ser Gln Pro Asn Arg Gly Pro Ser Pro Phe Ser Pro
165     170     175

Val Gln Leu His Gln Leu Arg Ala Gln Ile Leu Ala Tyr Lys Met Leu
180     185     190

Ala Arg Gly Gln Pro Leu Pro Glu Thr Leu Gln Leu Ala Val Gln Gly
195     200     205

Lys Arg Thr Leu Pro Gly Leu Gln Gln Gln Gln Gln Gln Gln Gln
210     215     220

Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Gln Pro Gln
225     230     235     240

Gln Gln Pro Pro Gln Pro Gln Thr Gln Gln Gln Gln Gln Pro Ala Leu
245     250     255

Val Asn Tyr Asn Arg Pro Ser Gly Pro Gly Pro Glu Leu Ser Gly Pro
260     265     270

Ser Thr Pro Gln Lys Leu Pro Val Pro Ala Pro Gly Gly Arg Pro Ser
275     280     285

Pro Ala Pro Pro Ala Ala Ala Gln Pro Pro Ala Ala Ala Val Pro Gly
290     295     300

Pro Ser Val Pro Gln Pro Ala Pro Gly Gln Pro Ser Pro Val Leu Gln
305     310     315     320

Leu Gln Gln Lys Gln Ser Arg Ile Ser Pro Ile Gln Lys Pro Gln Gly
325     330     335

Leu Asp Pro Val Glu Ile Leu Gln Glu Arg Glu Tyr Arg Leu Gln Ala
340     345     350

Arg Ile Ala His Arg Ile Gln Glu Leu Glu Asn Leu Pro Gly Ser Leu
355     360     365

Pro Pro Asp Leu Arg Thr Lys Ala Thr Val Glu Leu Lys Ala Leu Arg
370     375     380

Leu Leu Asn Phe Gln Arg Gln Leu Arg Gln Glu Val Val Ala Cys Met
385     390     395     400

```

-continued

Arg	Arg	Asp	Thr	Thr	Leu	Glu	Thr	Ala	Leu	Asn	Ser	Lys	Ala	Tyr	Lys	
				405					410					415		
Arg	Ser	Lys	Arg	Gln	Thr	Leu	Arg	Glu	Ala	Arg	Met	Thr	Glu	Lys	Leu	
			420					425					430			
Glu	Lys	Gln	Gln	Lys	Ile	Glu	Gln	Glu	Arg	Lys	Arg	Arg	Gln	Lys	His	
		435					440					445				
Gln	Glu	Tyr	Leu	Asn	Ser	Ile	Leu	Gln	His	Ala	Lys	Asp	Phe	Lys	Glu	
	450					455					460					
Tyr	His	Arg	Ser	Val	Ala	Gly	Lys	Ile	Gln	Lys	Leu	Ser	Lys	Ala	Val	
465					470					475					480	
Ala	Thr	Trp	His	Ala	Asn	Thr	Glu	Arg	Glu	Gln	Lys	Lys	Glu	Thr	Glu	
			485						490					495		
Arg	Ile	Glu	Lys	Glu	Arg	Met	Arg	Arg	Leu	Met	Ala	Glu	Asp	Glu	Glu	
			500					505					510			
Gly	Tyr	Arg	Lys	Leu	Ile	Asp	Gln	Lys	Lys	Asp	Arg	Arg	Leu	Ala	Tyr	
		515					520					525				
Leu	Leu	Gln	Gln	Thr	Asp	Glu	Tyr	Val	Ala	Asn	Leu	Thr	Asn	Leu	Val	
	530					535					540					
Trp	Glu	His	Lys	Gln	Ala	Gln	Ala	Ala	Lys	Glu	Lys	Lys	Lys	Arg	Arg	
545					550					555					560	
Arg	Arg	Lys	Lys	Lys	Ala	Glu	Glu	Asn	Ala	Glu	Gly	Gly	Glu	Ser	Ala	
				565					570					575		
Leu	Gly	Pro	Asp	Gly	Glu	Pro	Ile	Asp	Glu	Ser	Ser	Gln	Met	Ser	Asp	
			580					585					590			
Leu	Pro	Val	Lys	Val	Thr	His	Thr	Glu	Thr	Gly	Lys	Val	Leu	Phe	Gly	
		595					600					605				
Pro	Glu	Ala	Pro	Lys	Ala	Ser	Gln	Leu	Asp	Ala	Trp	Leu	Glu	Met	Asn	
	610					615					620					
Pro	Gly	Tyr	Glu	Val	Ala	Pro	Arg	Ser	Asp	Ser	Glu	Glu	Ser	Asp	Ser	
625					630					635					640	
Asp	Tyr	Glu	Glu	Glu	Asp	Glu	Glu	Glu	Glu	Ser	Ser	Arg	Gln	Glu	Thr	
			645						650					655		
Glu	Glu	Lys	Ile	Leu	Leu	Asp	Pro	Asn	Ser	Glu	Glu	Val	Ser	Glu	Lys	
			660					665					670			
Asp	Ala	Lys	Gln	Ile	Ile	Glu	Thr	Ala	Lys	Gln	Asp	Val	Asp	Asp	Glu	
		675					680					685				
Tyr	Ser	Met	Gln	Tyr	Ser	Ala	Arg	Gly	Ser	Gln	Ser	Tyr	Tyr	Thr	Val	
	690					695					700					
Ala	His	Ala	Ile	Ser	Glu	Arg	Val	Glu	Lys	Gln	Ser	Ala	Leu	Leu	Ile	
705					710					715					720	
Asn	Gly	Thr	Leu	Lys	His	Tyr	Gln	Leu	Gln	Gly	Leu	Glu	Trp	Met	Val	
			725						730					735		
Ser	Leu	Tyr	Asn	Asn	Asn	Leu	Asn	Gly	Ile	Leu	Ala	Asp	Glu	Met	Gly	
			740					745					750			
Leu	Gly	Lys	Thr	Ile	Gln	Thr	Ile	Ala	Leu	Ile	Thr	Tyr	Leu	Met	Glu	
		755					760					765				
His	Lys	Arg	Leu	Asn	Gly	Pro	Tyr	Leu	Ile	Ile	Val	Pro	Leu	Ser	Thr	
		770				775					780					
Leu	Ser	Asn	Trp	Thr	Tyr	Glu	Phe	Asp	Lys	Trp	Ala	Pro	Ser	Val	Val	
785					790					795					800	
Lys	Ile	Ser	Tyr	Lys	Gly	Thr	Pro	Ala	Met	Arg	Arg	Ser	Leu	Val	Pro	
			805						810					815		
Gln	Leu	Arg	Ser	Gly	Lys	Phe	Asn	Val	Leu	Leu	Thr	Thr	Tyr	Glu	Tyr	

820										825					830				
Ile	Ile	Lys	Asp	Lys	His	Ile	Leu	Ala	Lys	Ile	Arg	Trp	Lys	Tyr	Met				
835						840					845								
Ile	Val	Asp	Glu	Gly	His	Arg	Met	Lys	Asn	His	His	Cys	Lys	Leu	Thr				
850						855					860								
Gln	Val	Leu	Asn	Thr	His	Tyr	Val	Ala	Pro	Arg	Arg	Ile	Leu	Leu	Thr				
865						870					875								
Gly	Thr	Pro	Leu	Gln	Asn	Lys	Leu	Pro	Glu	Leu	Trp	Ala	Leu	Leu	Asn				
885						890					895								
Phe	Leu	Leu	Pro	Thr	Ile	Phe	Lys	Ser	Cys	Ser	Thr	Phe	Glu	Gln	Trp				
900						905					910								
Phe	Asn	Ala	Pro	Phe	Ala	Met	Thr	Gly	Glu	Arg	Val	Asp	Leu	Asn	Glu				
915						920					925								
Glu	Glu	Thr	Ile	Leu	Ile	Ile	Arg	Arg	Leu	His	Lys	Val	Leu	Arg	Pro				
930						935					940								
Phe	Leu	Leu	Arg	Arg	Leu	Lys	Lys	Glu	Val	Glu	Ser	Gln	Leu	Pro	Glu				
945						950					955								
Lys	Val	Glu	Tyr	Val	Ile	Lys	Cys	Asp	Met	Ser	Ala	Leu	Gln	Lys	Ile				
965						970					975								
Leu	Tyr	Arg	His	Met	Gln	Ala	Lys	Gly	Ile	Leu	Leu	Thr	Asp	Gly	Ser				
980						985					990								
Glu	Lys	Asp	Lys	Lys	Gly	Lys	Gly	Gly	Ala	Lys	Thr	Leu	Met	Asn	Thr				
995						1000					1005								
Ile	Met	Gln	Leu	Arg	Lys	Ile	Cys	Asn	His	Pro	Tyr	Met	Phe	Gln					
1010						1015					1020								
His	Ile	Glu	Glu	Ser	Phe	Ala	Glu	His	Leu	Gly	Tyr	Ser	Asn	Gly					
1025						1030					1035								
Val	Ile	Asn	Gly	Ala	Glu	Leu	Tyr	Arg	Ala	Ser	Gly	Lys	Phe	Glu					
1040						1045					1050								
Leu	Leu	Asp	Arg	Ile	Leu	Pro	Lys	Leu	Arg	Ala	Thr	Asn	His	Arg					
1055						1060					1065								
Val	Leu	Leu	Phe	Cys	Gln	Met	Thr	Ser	Leu	Met	Thr	Ile	Met	Glu					
1070						1075					1080								
Asp	Tyr	Phe	Ala	Phe	Arg	Asn	Phe	Leu	Tyr	Leu	Arg	Leu	Asp	Gly					
1085						1090					1095								
Thr	Thr	Lys	Ser	Glu	Asp	Arg	Ala	Ala	Leu	Leu	Lys	Lys	Phe	Asn					
1100						1105					1110								
Glu	Pro	Gly	Ser	Gln	Tyr	Phe	Ile	Phe	Leu	Leu	Ser	Thr	Arg	Ala					
1115						1120					1125								
Gly	Gly	Leu	Gly	Leu	Asn	Leu	Gln	Ala	Ala	Asp	Thr	Val	Val	Ile					
1130						1135					1140								
Phe	Asp	Ser	Asp	Trp	Asn	Pro	His	Gln	Asp	Leu	Gln	Ala	Gln	Asp					
1145						1150					1155								
Arg	Ala	His	Arg	Ile	Gly	Gln	Gln	Asn	Glu	Val	Arg	Val	Leu	Arg					
1160						1165					1170								
Leu	Cys	Thr	Val	Asn	Ser	Val	Glu	Glu	Lys	Ile	Leu	Ala	Ala	Ala					
1175						1180					1185								
Lys	Tyr	Lys	Leu	Asn	Val	Asp	Gln	Lys	Val	Ile	Gln	Ala	Gly	Met					
1190						1195					1200								
Phe	Asp	Gln	Lys	Ser	Ser	Ser	His	Glu	Arg	Arg	Ala	Phe	Leu	Gln					
1205						1210					1215								
Ala	Ile	Leu	Glu	His	Glu	Glu	Glu	Asn	Glu	Glu	Glu	Asp	Glu	Val					
1220						1225					1230								

-continued


Pro Asp	Asp Glu Thr	Leu Asn	Gln Met	Ile Ala	Arg	Arg Glu Glu
1235		1240			1245	
Glu Phe	Asp Leu Phe	Met Arg	Met Asp	Met Asp	Arg	Arg Arg Glu
1250		1255			1260	
Asp Ala	Arg Asn Pro	Lys Arg	Lys Pro	Arg Leu	Met	Glu Glu Asp
1265		1270			1275	
Glu Leu	Pro Ser Trp	Ile Ile	Lys Asp	Asp Ala	Glu	Val Glu Arg
1280		1285			1290	
Leu Thr	Cys Glu Glu	Glu Glu	Glu Lys	Ile Phe	Gly	Arg Gly Ser
1295		1300			1305	
Arg Gln	Arg Arg Asp	Val Asp	Tyr Ser	Asp Ala	Leu	Thr Glu Lys
1310		1315			1320	
Gln Trp	Leu Arg Ala	Ile Glu	Asp Gly	Asn Leu	Glu	Glu Met Glu
1325		1330			1335	
Glu Glu	Val Arg Leu	Lys Lys	Arg Lys	Arg Arg	Arg	Asn Val Asp
1340		1345			1350	
Lys Asp	Pro Ala Lys	Glu Asp	Val Glu	Lys Ala	Lys	Lys Arg Arg
1355		1360			1365	
Gly Arg	Pro Pro Ala	Glu Lys	Leu Ser	Pro Asn	Pro	Pro Lys Leu
1370		1375			1380	
Thr Lys	Gln Met Asn	Ala Ile	Ile Asp	Thr Val	Ile	Asn Tyr Lys
1385		1390			1395	
Asp Arg	Cys Asn Val	Glu Lys	Val Pro	Ser Asn	Ser	Gln Leu Glu
1400		1405			1410	
Ile Glu	Gly Asn Ser	Ser Gly	Arg Gln	Leu Ser	Glu	Val Phe Ile
1415		1420			1425	
Gln Leu	Pro Ser Arg	Lys Glu	Leu Pro	Glu Tyr	Tyr	Glu Leu Ile
1430		1435			1440	
Arg Lys	Pro Val Asp	Phe Lys	Lys Ile	Lys Glu	Arg	Ile Arg Asn
1445		1450			1455	
His Lys	Tyr Arg Ser	Leu Gly	Asp Leu	Glu Lys	Asp	Val Met Leu
1460		1465			1470	
Leu Cys	His Asn Ala	Gln Thr	Phe Asn	Leu Glu	Gly	Ser Gln Ile
1475		1480			1485	
Tyr Glu	Asp Ser Ile	Val Leu	Gln Ser	Val Phe	Lys	Ser Ala Arg
1490		1495			1500	
Gln Lys	Ile Ala Lys	Glu Glu	Glu Ser	Glu Asp	Glu	Ser Asn Glu
1505		1510			1515	
Glu Glu	Glu Glu Glu	Asp Glu	Glu Glu	Ser Glu	Ser	Glu Ala Lys
1520		1525			1530	
Ser Val	Lys Val Lys	Ile Lys	Leu Asn	Lys Lys	Asp	Asp Lys Gly
1535		1540			1545	
Arg Asp	Lys Gly Lys	Gly Lys	Lys Arg	Pro Asn	Arg	Gly Lys Ala
1550		1555			1560	
Lys Pro	Val Val Ser	Asp Phe	Asp Ser	Asp Glu	Glu	Gln Asp Glu
1565		1570			1575	
Arg Glu	Gln Ser Glu	Gly Ser	Gly Thr	Asp Asp	Glu	
1580		1585			1590	

121


The invention claimed is:

1. A method of treating acute myeloid leukemia that harbors a DNMT3A, RUNX1, or ASXL1 mutation in a subject in need thereof, the method comprising administering to the subject an effective amount of an agent, wherein the agent is a small molecule compound having the structure:

Formula IVa



wherein each of X, Y, and Z is, independently, N or CH; wherein R1 is absent, H, C₁-C₆ acyl, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₁-C₆ heteroalkyl, C₂-C₉ heterocyclyl, or —SO₂R⁶; each of R² and R⁵ is, independently, H or C₁-C₆ alkyl; R³ is H or C₁-C₆ alkyl; and R⁴ is H, C₁-C₆ alkyl, or C₁-C₆ heteroalkyl; or R³ and R⁴, together with the carbon atom to which each is attached, form a C₃-C₆ cycloalkyl; R⁶ is C₁-C₆ alkyl or —NR⁷R⁸; R⁷ and R⁸ are, independently, C₁-C₆; Het is 5-membered heteroarylene, 6-membered heteroarylene, or



A is C₆-C₁₀ arylene, C₂-C₉ heterocyclylene, or C₂-C₉ heteroarylene;

L is absent, —O—, C₁-C₆ alkylene, C₁-C₆ heteroalkylene, C₂-C₆ alkenylene, C₂-C₆ heteroalkenylene, C₂-C₆ alkynylene, C₂-C₆ heteroalkynylene, C₂-C₉ heterocyclylene, C₂-C₉ heterocyclyl C₁-C₆ alkylene, C₂-C₉ heteroarylene, or C₂-C₉ heteroaryl C₁-C₆ alkylene, wherein the C₂-C₉ heterocyclylene is optionally substituted with C₁-C₆ alkyl;

and B is H, halogen, cyano, C₆-C₁₀ aryl, C₃-C₁₀ cycloalkyl, C₂-C₉ heterocyclyl, or C₂-C₉ heteroaryl, wherein the C₂-C₉ heterocyclyl is optionally substituted with C₁-C₆ alkyl;

or a pharmaceutically acceptable salt thereof.

2. The method of claim 1, wherein the acute myeloid leukemia is metastatic.

3. The method of claim 1, wherein the method further comprises administering induction chemotherapy.

4. The method of claim 3, wherein the induction chemotherapy comprises cytarabine, an anthracycline, or combinations thereof.

122

5. The method of claim 1, wherein the method further comprises administering consolidation therapy.

6. The method of claim 5, wherein the consolidation therapy comprises an allogenic stem cell transplantation or immunotherapy.

7. The method of claim 1, wherein the method further comprises administering a hemopoietic stem cell transplant, gemtuzumab ozogamicin, or a combination thereof.

8. The method of claim 1, wherein the subject or cancer has or has been identified as having a BRG1 or BRM loss of function mutation.

9. The method of claim 1, wherein the acute myeloid leukemia has failed to respond to or progressed after administration of one or more chemotherapeutic or cytotoxic agents.

10. The method of claim 1, wherein the acute myeloid leukemia is resistant to, or predicted to be resistant to one or more chemotherapeutic agents.

11. The method of claim 9, wherein the one or more chemotherapeutic or cytotoxic agents is cytarabine, an anthracycline or gemtuzumab ozogamicin.

12. The method of claim 1, wherein the effective amount of the agent reduces the level or activity of BRG1 or BRM by at least 5% as compared to a reference.

13. The method of claim 1, wherein the small molecule compound has the structure:

