

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12398143
Kind Code	B2
Date of Patent	August 26, 2025
Inventor(s)	Giovannini; Riccardo et al.

Imidazopyrazine derivatives and the use thereof as medicament

Abstract

The present invention relates to novel imidazopyrazines of general formula A, processes for their preparation, pharmaceutical compositions containing them and their use in therapy, particularly in the treatment or prevention of conditions having an association with NR2B negative allosteric modulating properties. ##STR00001##

Inventors:	Giovannini; Riccardo (Biberach an der Riss, DE), Wiedenmayer; Dieter (Biberach an der Riss, DE)
Applicant:	Boehringer Ingelheim International GmbH (Ingelheim am Rhein, DE)
Family ID:	1000008780563
Assignee:	Boehringer Ingelheim International GmbH (Ingelheim am Rhein, DE)
Appl. No.:	17/616205
Filed (or PCT Filed):	June 03, 2020
PCT No.:	PCT/EP2020/065253
PCT Pub. No.:	WO2020/245137
PCT Pub. Date:	December 10, 2020

Prior Publication Data

Document Identifier	Publication Date
US 20220306637 A1	Sep. 29, 2022

Foreign Application Priority Data

EP	19178045	Jun. 04, 2019
----	----------	---------------

Publication Classification

Int. Cl.: C07D487/04 (20060101); A61K47/00 (20060101); A61P43/00 (20060101); C07D241/38 (20060101); C07D519/00 (20060101)

U.S. Cl.:

CPC C07D487/04 (20130101); A61K47/00 (20130101); A61P43/00 (20180101); C07D241/38 (20130101); C07D519/00 (20130101);

Field of Classification Search

CPC: A61K (47/00); C07D (241/38); C07D (487/04); C07D (519/00); A61P (43/00)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
10030026	12/2017	Shapiro	N/A	N/A
10849907	12/2019	Giovannini et al.	N/A	N/A
2002/0055519	12/2001	Thompson et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2002080928	12/2001	WO	N/A
2003010159	12/2002	WO	N/A
2004048364	12/2003	WO	N/A
200610969	12/2005	WO	N/A
2006113471	12/2005	WO	N/A
2007067511	12/2006	WO	N/A
2010088408	12/2009	WO	N/A
2012128582	12/2011	WO	N/A
2014060398	12/2013	WO	N/A
2015080904	12/2014	WO	N/A
2015130905	12/2014	WO	N/A
2016006593	12/2015	WO	N/A
2016029146	12/2015	WO	N/A
2016029146	12/2015	WO	N/A
2016049165	12/2015	WO	N/A
2016100349	12/2015	WO	N/A
2017066368	12/2016	WO	N/A
2019110703	12/2018	WO	N/A

OTHER PUBLICATIONS

Wermuth, Molecular variations based in isosteric replacements, The Practice of Medicinal Chemistry, 1996, 203-237. (Year: 1996). cited by examiner

International Search Report and Written Opinion for PCT 2018/083728 mailed Feb. 4, 2019. cited by applicant

Murrough, "Antidepressant Efficacy of Ketamine in Treatment-Resistant Major Depression: A Two

Site Randomized Controlled Trial”, Am. J. Psychiatry, 2013, vol. 170, p. 1134-1142. cited by applicant

Singh, “Intravenous Eskatamine in Adult Treatment-Resistant Depression: A double-Blind, Double-Randomization, Placebo Controlled Study”, Society of Biological Psychiatry, vol. 80, 2016, p. 424-431. cited by applicant

Berman, “Antidepressant effects of Ketamine in depressed patients”, Biological Psychiatry, vol. 47, 2000, p. 351-354. cited by applicant

Krystal, “Subanesthetic Effects of the Noncompetitive NMDA Antagonist, Ketamine in Humans”, Arch. Gen. Psychiatry, 1994, vol. 51, p. 199-214. cited by applicant

Paoletti, NMDA receptor subunit diversity: impact on receptor properties, synaptic plasticity and disease, Nature Reviews, vol. 14, 2013. cited by applicant

Miller, “GluN2B-containing NMDA receptors regulate depression-like behavior and are critical for the rapid antidepressant actions of ketamine”, eLife3e03581, 2014. cited by applicant

Kiselycznyk, “NMDA receptor subunits and associated signaling molecules mediating anti-depressant related effects of NMDA-GluN2B antagonism”, Bhav. Nrain Res. 2015, p. 89-95. cited by applicant

Jimenez-Sanchez, “The Role of GluN2A and GluN2B Subunits on the effects of NMDA receptor Antagonists in modeling Schizophrenia and treating Refractory Depression”, Neuropsychopharmacology, 2014. cited by applicant

Taylor, “Absolute Oral Bioavailability of Traxoprodil in Cytochrome P450 2D6 Extensive and Poor Metabolisers”, Clin. Pharmacokinet, 2006, vol. 45, p. 989-1001. cited by applicant

Addy, “Single dose Administration of MK-0657, an NR2B-Selective NMDA Antagonist”, J. of Clinical Pharmacology, 2009, p. 856-864. cited by applicant

Layton, “Discovery of 3-Substituted Aminocyclopentanes as Potent and Orally Bioavailable NR2B Subtype-Selective NMDA Antagonists”, ACS Chem. Neuroscience, 2011. cited by applicant

Traynelis, Glutamate Receptor Ion Channels: Structure, Regulation and Function, Pharmacology reviews, 2010, vol. 62. cited by applicant

Chaffey, NMDA receptor subtypes, Current Anesthesia and Critical Care, 2008, vol. 19, p. 183-201. cited by applicant

Mony, Allosteric modulators of NR-2B-containing NMDA receptors, British Journal of Pharmacology, vol. 157, 2009. cited by applicant

Preskorn, An Innovative Design to Establish Proof of Concept of the Antidepressant effects of the NR2B Subunit Selective N-Methyl D-Aspartate Antagonist CP-101, 606, Journal of Clinical Pharmacology, vol. 28, 2008. cited by applicant

Beinat, Insights into Structure related activity relationships, Current Medicinal Chem, 2010. vol. 17, p. 4166-4190. cited by applicant

Serafini, The Role of Ketamine in Treatment resistant Depression, Current Neuropsychopharmacology, 2014, vol. 10, p. 444-461. cited by applicant

Sanchez, The role of GluN2A and GluN2B Subunits on the Effects of NMDA Receptor Antagonists in Modeling Schizophrenia and Treating Refractory Depression, Neuropsychopharmacology, vol. 30, 2014. cited by applicant

Beinat, Insights into Structure-Activity Relationships, Current Medicinal Chem., vol. 17, 2010. cited by applicant

Niu et al., “Synthesis of C8-alkyl-substituted purine analogues by direct alkylation of 8-H purines with tetrahydrofuran catalyzed by CoCl₂ 6H₂O”, Chinese chem. letters, 2017, 28, pp. 105-108. cited by applicant

Weed et al., “Negative Allosteric Modulators Selective for the NR2B Subtype of the NMDA Receptor Impair Cognition in Multiple Domains”, Neuropsychopharmacology, 2016, 41, pp. 568-577. cited by applicant

Abstract in English for WO2016006593, dated Jan. 14, 2016. cited by applicant

Background/Summary

(1) The present invention relates to novel imidazopyrazines of general formula A

(2) ##STR00002##

processes for their preparation, pharmaceutical compositions containing them and their use in therapy, particularly in the treatment or prevention of conditions having an association with NR2B negative allosteric modulating properties.

(3) The compounds of the invention according to general formula A show NR2B negative allosteric modulating properties.

(4) Extensive studies over the past twenty years have indicated that N-methyl-D-aspartate receptors (NMDA) play a relevant role in Alzheimer's disease, Parkinson's disease, dyskinesia, stroke, motor neuron disease, psychosis, epilepsy, anxiety, schizophrenia and pain.

(5) The non-selective NMDA receptor antagonist ketamine, (racemic as well as the S enantiomer), a medication mainly used for starting and maintaining anaesthesia, has demonstrated over the last years clinical efficacy in treating major depressive disorder (MDD) at subanaesthetic doses (Murrough et al. 2013, Am J Psychiatry. 170: 1134; Singh et al. 2016, Biol Psychiatry. 80: 424). More precisely, ketamine elicits a rapid onset of efficacy which lasts several days in MDD patients insufficiently responding to standard drug therapy (Berman et al. 2000. Biol Psychiatry 47:351, Serafini et al. 2014. Curr. Neuropharmacol.12:444). However, non-selective NMDA receptor antagonists have a range of undesirable effects which limit their application. In particular dissociative and psychogenic side effects are prominent for the non-selective NMDA receptor antagonists such as ketamine (Krystal et al. 1994. Arch. Gen. Psychiatry 51:199). In the early 1990s, it was found that multiple NMDA receptor subtypes exist, which contain different NR2(A-D) subunits (Paoletti et al., 2013 Nat Rev. Neurosci 14:383). More recently, NR2B subtype selective NMDA receptor negative allosteric modulators (NR2B NAM) have raised interest and have shown potential in a wide range of clinical indications, such as attention, emotion, mood, and pain, as well as being involved in a number of different human disorders (Mony et. al. 2009. Br. J. Pharmacol. 157:1301; Chaffey et al., Current Anaesthesia & Critical Care 19, 183). In particular, NR2B NAM have also demonstrated antidepressant efficacy in the early stage of clinical trials (Preskorn et al. 2008. J Clin Psychopharmacol 70:58). Preclinical studies using NR2B NAM as well as applying various transgenic mice strains have shown that NR2B containing NMDA-receptors are mediating the positive effect of ketamine in e.g. the Forced Swim Test (Miller et al. 2014 eLife 3:e03581; Kiselycznyk et al. 2015, Behav Brain Res, 287:89). Furthermore, selective NR2B NAM have advantages over unselective NMDA receptor antagonists, such as ketamine, due to greatly diminished dissociative and psychotomimetic side effects (Jimenez-Sanchez et al. 2014. Neuropsychopharmacology 39:2673). NR2B NAM described to date have exhibited drawbacks with regard to their receptor pharmacology and/or to other drug properties which have limited potential use in human drug therapy (Taylor, et al., 2006, Clin Pharmacokinet. 45: 989; Addy et al. 2009 J of Clinical Pharmacology 49:856)).

(6) WO2016/29146 discloses compounds of formula (I)

(7) ##STR00003##

that are inhibitors of methionyl-tRNA synthetase (MetRS) being useful as antibiotics.

(8) Formula (I) in WO2016/29146 encompasses the specific examples 1734, 1744, 1745, 1757

1758, 1785 and 1790 which exhibit a benzimidazole or imidazopyridine substructure.

(9) The compounds of the present invention have surprisingly been found to be potent NR2B negative allosteric modulators (see table 1), whereas the specific examples 1734, 1744, 1745, 1757, 1758, 1785 and 1790 of WO2016/29146 show rather poor negative allosteric modulation of the NR2B ion channel or no activity at all (see table 2).

(10) Further, the compounds of the present invention show good membrane permeability and low to moderate in vitro efflux (see table 3 for MDCK assay MDR1 (p-GP)). Therefore, compounds of the present invention are expected to show a favorable brain penetration which is required for efficacious CNS medicaments.

(11) The MDCK assays provide information on the potential of a compound to pass the blood brain barrier. Permeability measurements across polarized, confluent MDCK-MDR1 cell monolayers grown on permeable filter supports are used as an in vitro absorption model: apparent permeability coefficients (PE) of the compounds across the MDCK-MDR1 cell monolayers are measured (pH 7.4, 37° C.) in apical-to-basal (AB) and basal-to-apical (BA) transport direction. The AB permeability (PEAB) represents drug absorption from the blood into the brain and the BA permeability (PEBA) drug efflux from the brain back into the blood via both, passive permeability as well as active transport mechanisms mediated by efflux and uptake transporters that are expressed on the MDCK-MDR1 cells, predominantly by the overexpressed human MDR1. Identical or similar permeabilities in both transport directions indicate passive permeation, vectorial permeability points to additional active transport mechanisms. Higher PEBA than PEAB (PEBA/PEAB>5) indicates the involvement of active efflux mediated by MDR1, which might compromise the goal to achieve sufficient brain exposure. Therefore, this assay provides valuable support for selection of compounds applicable for further in vivo testing. High permeability not limited by efflux at the blood brain barrier is a favourable characteristic for compounds that are to be used for drugs acting primarily in the CNS. Consequently, to ensure high permeability at the blood brain barrier, it is highly preferred to minimize the efflux (efflux<5) at MDR1 transporter.

(12) Further, the compounds of the present invention are metabolically stable in human liver microsomes (see table 4, metabolic stability). Therefore, compounds of the present invention are expected to have a favorable in vivo clearance and thus the desired duration of action in humans.

(13) Stability in human liver microsomes refers to the susceptibility of compounds to biotransformation in the context of selecting and/or designing drugs with favorable pharmacokinetic properties. The primary site of metabolism for many drugs is the liver. Human liver microsomes contain the cytochrome P450s (CYPs), and thus represent a model system for studying drug metabolization in vitro. Enhanced stability in human liver microsomes is associated with several advantages, including increased bioavailability and adequate half-life, which can enable lower and less frequent dosing of patients. Thus, enhanced stability in human liver microsomes is a favorable characteristic for compounds that are to be used for drugs.

(14) Consequently, compounds of the present invention must be more viable for human use.

(15) The objective technical problem is thus to provide potent NR2B negative allosteric modulators.

(16) The present invention provides novel imidazopyrazines of formula A

(17) ##STR00004##

in which R^{sup.1} represents phenyl which is optionally substituted with 1 to 3 substituents selected from the group consisting of fluoro, chloro, methyl, ethyl, cyclopropyl, F_{sub.2}H_{sub.2}C—, F_{sub.2}H_{sub.2}C—, F_{sub.3}C—; or a salt thereof, particularly a pharmaceutically acceptable salt thereof.

(18) In another embodiment, in the general formula A R^{sup.1} represents phenyl which is optionally substituted with 1 or 2 substituents selected from the group consisting of fluoro, methyl.

(19) In another embodiment, in the general formula A R^{sup.1} represents

(20) ##STR00005##

(21) The present invention provides novel imidazopyrazines of general formula A that

unexpectedly are potent NR2B negative allosteric modulators.

(22) Another aspect of the invention refers to compounds according to formula A as NR2B negative allosteric modulators having appropriate membrane permeability and low to moderate in vitro efflux.

(23) Another aspect of the invention refers to compounds according to formula A as NR2B negative allosteric modulators having high metabolic stability in human liver microsomes.

(24) Another aspect of the invention refers to compounds according to formula A as NR2B negative allosteric modulators having appropriate membrane permeability, low to moderate in vitro efflux and high metabolic stability in human liver microsomes.

(25) Another aspect of the invention refers to pharmaceutical compositions, containing at least one compound according to formula A optionally together with one or more inert carriers and/or diluents.

(26) A further aspect of the present invention refers to compounds according to formula A, for the use in the prevention and/or treatment of disorders associated with NR2B negative allosteric modulators.

(27) Another aspect of the invention refers to processes of manufacture of the compounds of the present invention.

(28) Preparation

(29) The following schemes shall illustrate generally how to manufacture the compounds according to general formula A and the corresponding intermediate compounds by way of example. The abbreviated substituents may be as defined above if not defined otherwise within the context of the schemes.

(30) ##STR00006##

(31) ##STR00007##

(32) Alternatively, the synthesis can be performed using racemic morpholine-2,4-dicarboxylic acid 4-tert-butyl ester as starting material.

(33) Scheme 1 and 2 can be successfully used for gram scale synthesis of the final compounds starting from 40 mmoles of the desired substituted morpholine (racemic or the S enantiomer), using an excess of the desired substituted benzyl alcohol, DIPEA (3 equivalents), the needed coupling agent such as CDI and DMF as solvent.

(34) An alternative gram scale synthesis can be performed using the corresponding morpholine (racemic or the S enantiomer; 40 mmol), TEA (2.5 equivalents), a slight excess of the required imidoilylcarbonate and a 1/1 (volume/volume) mixture CH₂Cl₂/THF as solvent. The final compounds can be obtained with good to high enantiomeric excess; alternatively, chiral separation can be applied to obtain the pure enantiomers.

(35) In schemes 1 and 2, the substituent R^{sup.1} has the meaning as defined for general formula A, all embodiments of the invention that directly refer thereto and specifically the meaning as defined in the claims.

GENERAL DEFINITIONS

(36) Terms not specifically defined herein should be given the meanings that would be given to them by one skilled in the art in light of the disclosure and the context.

(37) In case a compound of the present invention is depicted in form of a chemical name as well as a formula, the formula shall prevail in case of any discrepancy.

(38) An asterisk may be used in sub-formulas to indicate the bond which is connected to the core molecule or to the substituent to which it is bound as defined.

(39) The term “substituted” as used herein means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's viable valence number is not exceeded, and that the substitution results in a stable compound.

(40) Stereochemistry:

(41) Unless specifically indicated, throughout the specification and the appended claims, a given

chemical formula or name shall encompass rotamers, tautomers and all stereo, optical and geometrical isomers (e.g. enantiomers, diastereoisomers, E/Z isomers etc.) and racemates thereof, as well as mixtures in different proportions of the separate enantiomers, mixtures of diastereoisomers, or mixtures of any of the foregoing forms where such isomers and enantiomers exist, as well as salts, including pharmaceutically acceptable salts thereof.

(42) Salts:

(43) The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use without excessive toxicity, irritation, allergic response, or other problem or complication, and commensurate with a reasonable benefit/risk ratio.

(44) As used herein, “pharmaceutically acceptable salts” refer to derivatives of the disclosed compounds wherein the parent compound forms a salt or a complex with an acid or a base.

(45) Examples for acids forming a pharmaceutically acceptable salt with a parent compound containing a basic moiety include mineral or organic acids such as benzenesulfonic acid, benzoic acid, citric acid, ethanesulfonic acid, fumaric acid, gentisic acid, hydro-bromic acid, hydrochloric acid, maleic acid, malic acid, malonic acid, mandelic acid, methanesulfonic acid, 4-methyl-benzenesulfonic acid, phosphoric acid, salicylic acid, succinic acid, sulfuric acid or tartaric acid.

(46) Examples for cations and bases forming a pharmaceutically acceptable salt with a parent compound containing an acidic moiety include Na.sup.+, K.sup.+, Ca.sup.2+, Mg.sup.2+, NH₄.sup.+, L-arginine, 2,2'-iminobisethanol, L-lysine, N-methyl-D-glucamine or tris(hydroxymethyl)-amino-methane.

(47) The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a sufficient amount of the appropriate base or acid in water or in an organic diluent like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile, or a mixture thereof. Salts of other acids than those mentioned above which for example are useful for purifying or isolating the compounds of the present invention (e.g. trifluoro-acetate salts) also comprise a part of the invention.

BIOLOGICAL ASSAYS AND DATA

List of Abbreviations

(48) DMEM Dulbecco's Modified Eagle's Medium FBS fetal Bovine Serum FLIPR fluorometric imaging plate reader HEK293 cell line derived from human embryonic kidney cells HEPES hydroxyethyl-piperazineethane-sulfonic acid buffer MDCK Madin-Darby canine kidney MDR1 Multi drug resistance protein 1 p-GP p-Glycoprotein

In-Vitro Effect:

Determination of In Vitro Pharmacological Activity

(49) The activity of the compounds of the invention may be demonstrated using the following in vitro NMDA NR1/NR2b cell assays:

(50) Method:

(51) A human HEK293 cell line with tetracyclin-inducible expression of NMDA NR1/NR2B receptor was used as a test system for compound efficacy and potency. The cell line was purchased from ChanTest, Catalog #CT6121. Compound activity was determined by measuring the effect of compounds on intracellular calcium concentration induced by glycine/glutamate agonism in a FLIPRtetra system (Molecular Devices).

(52) Cell Culture:

(53) The cells were obtained as frozen cells in cryo-vials and stored until use at -150° C. Cells were grown in culture medium (DMEM/F12, 10% FBS, 5 µg/mL Blasticidin, 150 µg/mL Zeozin, 500 µg/mL Geneticin). It is important that density does not exceed 80% confluence. For sub-culturing the cells were detached from flasks by Versene. For the assay, cells were detached, washed twice with induction medium (DMEM/F.sub.12 without glutamine, 10% FBS, 2 µg/mL

Tetracycline, 2 mM MEmamine) and seeded to 384 well pure coat amine plates (BD 359324, 50000 cells per well in 50 μ l) 48 h prior to assay in induction medium.

(54) Compound Preparation

(55) The test compounds were dissolved in 100% DMSO at a concentration of 10 mM and in a first step diluted in DMSO to a concentration of 5 mM, followed by serial dilution steps in 100% DMSO. Dilution factor and number of dilution steps may vary according to needs. Typically 8 different concentrations by 1:5 dilutions were prepared in duplicate, further intermediate dilutions (1:37.5) of the substances were carried out with aqueous assay buffer (137 mM NaCl, 4 mM KCl, 1.8 mM CaCl, 10 mM HEPES, 10 mM Glucose, pH 7.4) resulting in a compound concentration 3 times above the final test concentration and DMSO at 2.7% resulting in 0.9% final DMSO concentration in the assay.

(56) FLIPR Assay:

(57) At the assay day cells were washed 3 $\times$ with assay buffer, 10 μ L buffer remained in the wells after washing. 10 μ L Ca kit loading buffer (AAT Bioquest) was added to the cells and the plates were incubated with lid for 60 minutes at r.t. 20 μ l assay buffer containing 60 μ M glycine (20 μ M final) and 3 μ M glutamate (1 μ M final) was added to column 1-23. Fluorescence (indicating the calcium influx as a result of the NR1/NR2B ion channel activation) was read on the FLIPRtetra device for 60 seconds to monitor the glutamate induced effects. After 2 minutes 20 μ L of compound or controls (row 1-22) in assay buffer were carefully added to the wells. Fluorescence was read on the FLIPR tetra device for additional 6 minutes to monitor the compound induced effects after activation by agonists. The average of 2 measurements at 5 minutes and 5 min 10 seconds after compound addition is calculated and further used for IC₅₀ calculations. Each assay microtiter plate contained wells (in column 23 or 24) with DMSO controls instead of compound as controls for glycine/glutamate induced fluorescence (high controls) and wells with 1 μ M of a reference NR2b NAM as low controls (Compound 22; reference: Layton, Mark E et al, ACS Chemical Neuroscience 2011, 2(7), 352-362).

(58) Data Evaluation and Calculation:

(59) The output file of the reader contains the well number and measured average fluorescence units. For data evaluation and calculation, the measurement of the low control was set as 0% control and the measurement of the high control was set as 100% control. The IC₅₀ values were calculated using the standard 4 parameter logistic regression formula. Calculation: $[y=(a-d)/(1+(x/c)^b)+d]$, a=low value, d=high value; x=conc M; c=IC₅₀ M; b=slope.

(60) NR2B negative allosteric modulators covered by general structure A and exhibiting a low IC_{sub}.50 value are preferred.

(61) TABLE-US-00001 TABLE 1 In vitro affinity of the compounds of the present invention as obtained in the FLIPR assay Example number IC₅₀ [nM] 1 90 3 725 4 382 5 477 10 735 12 210

(62) TABLE-US-00002 TABLE 2 In vitro affinity of the closest prior art compounds (examples 1734, 1744, 1745, 1757, 1758, 1785 and 1790 in WO2016/29146) as obtained in the same FLIPR assay as compounds in table 1 Example number in WO2016/29146 IC₅₀ [nM] 1734 >8885 1744 >8889 1745 >8898 1757 >8900 1758 >8884 1785 6200 1790 >8887

MDCK Assay MDR-1 (p-GP)

(63) Apparent permeability coefficients (Papp) of the compounds across the MDCK-MDR1 monolayers (MDCKII cells transfected with human MDR1 cDNA expression plasmid) are measured in apical-to-basal (AB) and basal-to-apical (BA) direction.

(64) MDCK-MDR1 cells (6 $\times$ 10^{sup.5} cells/cm^{sup.2}) are seeded on filter inserts (Corning, Transwell, polycarbonate, 0.4 μ m pore size) and cultured for 9 to 10 days. Compounds dissolved in DMSO stock solution (1-20 mM) are diluted with HTP-4 aqueous buffer (128.13 mM NaCl, 5.36 mM KCl, 1 mM MgSO₄, 1.8 mM CaCl_{sub}.2), 4.17 mM NaHCO_{sub}.3, 1.19 mM Na₂HPO₄, 0.41 mM NaH₂PO₄, 15 mM HEPES, 20 mM glucose, pH 7.4) supplemented with 0.25% BSA to prepare the transport solutions (final concentration: 1 or 10 μ M, final DMSO $\leq$ 0.5%). The

transport solution is applied to the apical or basolateral donor side for measuring A-B or B-A permeability, respectively. The receiver side contains HTP-4 buffer supplemented with 0.25% BSA. Samples are collected at the start and end of experiment from the donor and at various time intervals for up to 2 hours also from the receiver side for concentration measurement by HPLC-MS/MS.

(65) Sampled receiver volumes are replaced with fresh receiver solution. Efflux ratio is calculated dividing the Papp (b-a) values by the Papp (a-b) values. Results are shown in table 3.

(66) TABLE-US-00003 TABLE 3 Papp (a-b) mean Ex. [10–6 cm/s] efflux ratio 1 33 1.6 3 61 0.7 4 57 0.8 5 40 1.4 10 23 2.9 12 31 1.8

The experimental results above show that compounds of the present invention are potent NR2B NAMs having good membrane permeability and low to moderate in vitro efflux.

Metabolic Stability

(67) The metabolic degradation of the test compound was assayed at 37° C. with pooled human liver microsomes. The final incubation volume of 60 l per time point contains TRIS buffer pH 7.6 at room temperature (0.1 M), magnesium chloride (5 mM aqueous solution), microsomal protein (1 mg/mL for human) and the test compound at a final concentration of 1 µM. Following a short preincubation period at 37° C., the reactions were initiated by addition of betanicotinamide adenine dinucleotide phosphate, reduced form (NADPH, 1 mM), and terminated by transferring an aliquot into solvent after different time points. After centrifugation (10000 g, 5 min), an aliquot of the supernatant was assayed by LC-MS/MS for the amount of parent compound. The half-life was determined by the slope of the semi-logarithmic plot of the concentration-time profile. Results are shown in Table 4.

(68) TABLE-US-00004 TABLE 4 Half-life - t½ [min] Ex. human liver microsomes 1 31.5 3 8 4 >130 5 >130 10 >130 12 36

The present invention provides compounds according to formula A that unexpectedly result in a favorable combination of the following key parameters: 1) NR2B negative allosteric modulation, 2) favorable stability in human liver microsomes, and 3) moderate to low in vitro efflux at MDR1 transporter.

Pharmaceutical Composition

(69) Suitable preparations for administering the compounds of the present invention will be apparent to those with ordinary skill in the art and include for example tablets, pills, capsules, suppositories, lozenges, troches, solutions, syrups, elixirs, sachets, injectables, inhalatives, powders, etc. The content of the pharmaceutically active compound(s) may vary in the range from 0.1 to 95 wt.-%, preferably 5.0 to 90 wt.-% of the composition as a whole.

(70) Suitable tablets may be obtained, for example, by mixing a compound of the present invention with known excipients, for example inert diluents, carriers, disintegrants, adjuvants, surfactants, binders and/or lubricants and pressing the resulting mixture to form tablets.

(71) Use in Treatment/Method of Use

(72) Human therapeutic applications of NR2B NAM have been summarized in reviews by Traynelis et al. (Traynelis et al., Pharmacology Reviews, 2010, 62:405), Beinat et al. (Beinat et al., Current Medicinal Chemistry, 2010, 17:4166) and Mony et al. (Mony et al., British J. Pharmacology, 2009, 157:1301).

(73) The present invention relates to compounds which are useful in the treatment of psychiatric disorders, diseases and conditions wherein negative allosteric modulation of NR2B is of therapeutic benefit, including: (1) mood disorders and mood affective disorders; (2) schizophrenia spectrum disorders; (3) neurotic, stress-related and somatoform disorders including anxiety disorders; (4) disorders of psychological development; (5) behavioral syndromes associated with physiological disturbances and physical factors; (6) substance-related and addictive disorders; (7) disease associated with symptoms of negative and positive valence.

(74) In view of their pharmacological effect, compounds of the present invention are suitable for

use in the treatment of a disorder, disease or condition selected from the list consisting of (1) treatment of mood disorders and mood affective disorders including bipolar disorder I depressed, hypomanic, manic and mixed form; bipolar disorder II; depressive disorders, such as single depressive episode or recurrent major depressive disorder, minor depressive disorder, depressive disorder with postpartum onset, depressive disorders with psychotic symptoms; major depressive disorder with or without concomitant anxious distress, mixed features, melancholic features, atypical features, mood-congruent psychotic features, mood-incongruent psychotic features, catatonia. (2) treatment of mood disorders belonging to the schizophrenia spectrum and other psychotic disorders including schizophrenia and schizoaffective disorder with associated negative and cognitive symptoms. (3) treatment of disorders belonging to the neurotic, stress-related and somatoform disorders including anxiety disorders, general anxiety disorder, panic disorder with or without agoraphobia, specific phobia, social phobia, chronic anxiety disorders; obsessive compulsive disorder; reaction to severe stress and adjustment disorders, such as post-traumatic stress disorder; other neurotic disorders such as depersonalisation-derealisation syndrome. (4) treatment of disorders of psychological development including pervasive developmental disorders, including Asperger's syndrome and Rett's syndrome, autistic disorders, childhood autism and overactive disorder associated with mental retardation and stereotyped movements, specific developmental disorder of motor function, specific developmental disorders of scholastic skills, attention deficit/hyperactivity disorder. (5) treatment of behavioral syndromes associated with physiological disturbances and physical factors including mental and behavioural disorders associated with the puerperium, including postnatal and postpartum depression; eating disorders, including anorexia nervosa and bulimia nervosa and other impulse control disorders. (6) treatment of disorders of substance-related and addictive disorders, which are substance use disorders induced by alcohol, *cannabis*, hallucinogen, stimulant, hypnotic, tobacco. (7) treatment of disease associated with symptoms of negative and positive valence including anhedonia, sustained threat and loss, suicidal ideation.

(75) As used herein, unless otherwise noted, the terms “treating”, “treatment” shall include the management and care of a human subject or human patient for the purpose of combating a disease, condition, or disorder and includes the administration of a compound of the present invention to prevent the onset of the symptoms or complications, alleviate the symptoms or complications, or eliminate the disease, condition, or disorder.

(76) As used herein, unless otherwise noted, the term “prevention” shall include (a) reduction in the frequency of one or more symptoms; (b) reduction in the severity of one or more symptoms; (c) the delay or avoidance of the development of additional symptoms; and/or (d) delay or avoidance of the development of the disorder or condition.

(77) According to another aspect, the present invention provides a compound of formula A or a pharmaceutically acceptable salt thereof for use in the treatment and/or prevention of the above mentioned conditions.

(78) According to another aspect, the present invention provides a compound of formula A according to any one of the preceding aspects characterized in that the compound of formula A is used in addition to behavioural therapy, TMS (transcranial magnetic stimulation), ECT (electroconvulsive therapy) and other therapies.

(79) Combination Therapy

(80) Compounds according to the present invention can be combined with other treatment options known to be used in the art in connection with a treatment of any of the indications the treatment of which is in the focus of the present invention.

(81) According to another aspect, the present invention provides a compound of formula A according to any one of the preceding aspects characterized in that the compound of formula A is administered in addition to treatment with one or more antidepressant selected from the list consisting of duloxetine, escitalopram, bupropion, venlafaxine, desvenlafaxine, sertraline,

paroxetine, fluoxetine, vortioxetine, mirtazapine, citalopram, vilazodone, trazodone, amitriptyline, clomipramine, agomelatine, levomilnacipran, lithium, doxepin, nortriptyline. The term “antidepressant” shall mean any pharmaceutical agent or drug which can be used to treat depression or diseases associated with depressive symptoms.

(82) According to another aspect, the present invention provides a compound of formula A according to any one of the preceding aspects characterized in that the compound of formula A is administered in addition to treatment with one or more antipsychotic selected from the list consisting of aripiprazole, paliperidone palmitate, lurasidone, quetiapine, risperidone, olanzapine, paliperidone, brexpiprazole, clozapine, asenapine, chlorpromazine, haloperidol, cariprazine, ziprasidone, amisulpride, iloperidone, fluphenazine, blonanserin, aripiprazole lauroxil. The term “antipsychotic” shall mean any pharmaceutical agent or drug which can be used to treat diseases associated with psychotic or depressive symptoms.

(83) According to another aspect, the present invention provides a compound of formula A according to any one of the preceding aspects characterized in that the compound of formula A is administered in addition to treatment with one or more psychostimulant selected from the list consisting of lisdexamfetamine, methylphenidate, amphetamine, dexamfetamine, dexmethylphenidate, armodafinil, modafinil. The term “psychostimulant” shall mean any pharmaceutical agent or drug which can be used to treat diseases like mood disorders, or impulse control disorders.

(84) According to another aspect, the present invention provides a compound of formula A according to any one of the preceding aspects characterized in that the compound of formula A is administered in addition to treatment with nootropics selected from the list consisting of oxiracetam, piracetam, or the natural product St John's-wort.

(85) According to another aspect, the present invention provides a compound of formula A which is administered in addition to treatment with one or more antidepressant, antipsychotic, psychostimulant, nootropics or natural product according to any one of the preceding aspects characterized in that the combination of compound of formula A and one or more antidepressant, antipsychotic, psychostimulant, nootropics or natural product is used in addition to behavioural therapy, TMS (transcranial magnetic stimulation), ECT (electroconvulsive therapy) and other therapies.

Description

EXPERIMENTAL SECTION

Abbreviations

(1) ACN acetonitrile APCI Atmospheric pressure chemical ionization Boc tert-butyloxycarbonyl CDI 1,1'-carbonyldiimidazole CO₂ Carbon Dioxide d day DA Diode Array DAD Diode Array Detector DCM dichloromethane DIPE diisopropylether DIPEA diisopropylethylamine DMF dimethylformamide ee, e.e. enantiomeric excess ELSD Evaporative Light Scattering Detector ESI electrospray ionization (in MS) EtOAc ethylacetate EtOH ethanol Exp. example h hour(s) HATU O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium-hexafluorophosphate HPLC high performance liquid chromatography HPLC-MS coupled high performance liquid chromatography-mass spectrometry IPA Isopropanol M molar (mol/L) MeOH methanol min minute(s) MS mass spectrometry MW molecular weight NH₃ ammonia PSI Pound per square inch rt room temperature R_t retention time scCO₂ supercritical CO₂ Sol Solvent solv solvent TBTU O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate TEA triethylamine TFA trifluoroacetic acid THE tetrahydrofuran TLC thin-layer chromatography SFC Supercritical fluid chromatography

General Analytics

(2) All reactions were carried out using commercial grade reagents and solvents. NMR spectra were recorded on a Bruker AVANCE IIIHD 400 MHz instrument using Top-Spin 3.2 p16 software. Chemical shifts are given in parts per million (ppm) downfield from internal reference trimethylsilane in 6 units. Selected data are reported in the following manner: chemical shift, multiplicity, coupling constants (J), integration. Analytical thin-layer chromatography (TLC) was carried out using Merck silica gel 60 F254 plates. All compounds were visualized as single spots using short wave UV light. Low resolution mass spectra were obtained using a liquid chromatography mass spectrometer (LCMS) that consisted of an Agilent 1100 series LC coupled to a Agilent 6130 quadru-pole mass spectrometer (electrospray positive ionization).

(3) Methods:

(4) For solvent mixtures used for HPLC-MS methods and Chiral SFC analytical methods, % solvent are given as volume percent of the corresponding solvent.

(5) HPLC-MS methods:

(6) TABLE-US-00005 Method 1 Method Name: Z011_S03 Device description: Agilent 1200 with DA- and MS-Detector Column: XBridge C18_3.0 × 30 mm_2.5 μm Column producer: Waters Description:

(7) TABLE-US-00006 Method Name: Z011_S03 Gradient/ % Sol Back Solvent [Water % Sol Flow Temp pressure Time [min] 0.1% NH.sub.3] [Acetonitrile] [ml/min] [° C.] [PSI] 0.0 97.0 3.0 2.2 60.0 0.2 97.0 3.0 2.2 60.0 1.2 0.0 100.0 2.2 60.0 1.25 0.0 100.0 3.0 60.0 1.4 0.0 100.0 3.0 60.0

Chiral SFC analytical methods:

(8) TABLE-US-00007 Method 2: I_C2_20_MeOH_NH.sub.3 —001 Method Name: I_C2_20_MEOH_NH.sub.3 —001 Device description: Agilent 1260 SFC with DAD and ELSD Column: Lux ® Cellulose-2_4.6 × 250 mm_5 μm Column producer: Phenomenex

(9) TABLE-US-00008 Gradient/ % Sol Back Solvent % Sol [MeOH Flow Temp pressure Time [min] [scCO₂] 20 mM NH.sub.3] [ml/min] [° C.] [PSI] 0.0 80.0 20.0 4.0 40.0 2175.0 10.0 80.0 20.0 4.0 40.0 2175.0

(10) TABLE-US-00009 Method 3: I_C2_40_IPA_NH.sub.3 —001 Method Name: I_C2_40_IPA_NH.sub.3 —001 Device description: Agilent 1260 SFC with DAD and ELSD Column: Lux ® Cellulose-2_4.6 × 250 mm_5 μm Column producer: Phenomenex

(11) TABLE-US-00010 Gradient/ % Sol Back Solvent % Sol [IPA Flow Temp pressure Time [min] [scCO₂] 20 mM NH.sub.3] [ml/min] [° C.] [PSI] 0.0 60.0 40.0 4.0 40.0 2175.0 10.0 60.0 40.0 4.0 40.0 2175.0

(12) TABLE-US-00011 Method 4: I_SA_20_MEOH_NH.sub.3 —001 Method Name: I_SA_20_MEOH_NH.sub.3 —001 Device description: Agilent 1260 SFC with DAD and MS Column: CHIRAL ART ® Amylose SA_4.6 × 250 mm_5 μm Column producer: YMC

(13) TABLE-US-00012 Gradient/ % Sol Back Solvent % Sol [MeOH Flow Temp pressure Time [min] [scCO₂] 20 mM NH.sub.3] [ml/min] [° C.] [PSI] 0.0 80.0 20.0 4.0 40.0 2175.0 10.0 80.0 20.0 4.0 40.0 2175.0

Preparation of Intermediates:

Example 1a

(14) TABLE-US-00013  S-Morpholin-2,4-dicarboxylic acid 4-tertbutylester (1.26 g, 5.45 mmol) was dissolved in DMF (20 ml). Then HATU (2.28 g, 5.99 mmol) was added, followed by Pyrazine-2,3-diamine (600 mg, 5.45 mmol, CAS No. 13134-31-1) and TEA (1.52 ml, 10.9 mmol). The reaction mixture was stirred 20 h at room temperature before the work-up: The reaction mixture was filtered with a glass filter and the filtrate was concentrated in vacuo. The residue was treated with 40 mL H.sub.2O and stirred for 2 h. The obtained precipitate was filtered off and dried in the air. Obtained 1.60 g of the desired product. HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.76 MS: 324 (M + H).sup.+

Example 2a—Method A (According to Scheme 1)

(15) TABLE-US-00014  A mixture of example 1a (1.60 g, 4.95 mmol),

potassium carbonate (2.74 g, 19.8 mmol) and isopropanol (50 ml) was stirred for 6 h at 80° C. The precipitate was filtered off, washed with isopropanol and the filtrate was concentrated in vacuo. The residue was dried overnight. Obtained 1.50 g of the desired product. HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.58 MS: 306 (M + H).sup.+

Example 2a—Method B (Direct Cyclisation, According to Scheme 2)

(16) TABLE-US-00015  S-Morpholine-2,4-dicarboxylic acid 4-tertbutylester (3.15 g, 13.62 mmol) was dissolved in DMF (30 ml). Then HATU (5.7 g, 15 mmol) was added, followed by Pyrazine-2,3-diamine (1.5 g, 13.62 mmol, CAS No. 13134-31-1) and TEA (3.8 ml, 27.2 mmol). The reaction mixture was stirred 7 h at room temperature before the work-up: The reaction mixture was filtered with a glass filter and the filtrate was concentrated in vacuo. The residue was treated with 100 mL H.sub.2O and stirred for 12 h. The obtained precipitate was filtered off and dried in the air. Obtained 2.0 g of the desired cyclised product. HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.76 MS: 306 (M + H).sup.+ Chiral SFC; Method: I_C2_20_MEOH_NH3_001; ee: 100% R.sub.t [min]: 3.36 min

Example 3a

(17) TABLE-US-00016  Example 2a (2 g, 6.55 mmol) was dissolved in DCM (120 ml), HCl in dioxane (4 M, 9.83 ml, 39.3 mmol) was added and the reaction mixture was stirred 3.5 h at rt. The solvents were evaporated in vacuo and the obtained residue was used as such in the next step. Obtained 1.8 g of the desired product as a salt. Chiral SFC; Method: I_C2_20_MEOH_NH3_001; MS: 206 (M + H).sup.+ R.sub.t [min]: 6.48 min ee: 100%

EXEMPLARY EMBODIMENTS

Example 1

(18) (2-Fluoro-4-methylphenyl)methanol (151 mg; 1.08 mmol, CAS No. 252004-38-9) and CDI (175 mg; 1.08 mmol) were mixed together in DMF (3 ml); the reaction mixture was heated at 50° C. during 30 minutes. Example 3a (150 mg; 0.54 mmol) and DIPEA (0.28 ml; 1.62 mmol) were then added in sequence and the reaction mixture was stirred 2 hours at rt. The reaction mixture was diluted with 3 ml of a mixture MeOH/Water (1/1; volume/volume) before being filtered and separated via semipreparative HPLC.

(19) The residue was triturated with 3 mL DIPE, filtered and dried in the air. Obtained 157 mg of the desired compound.

(20) TABLE-US-00017 Example 1  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.68 MS: 372 (M + H).sup.+ Chiral SFC; Method: I_SA_20_MEOH_NH.sub.3_001 Rt [min]: 4.13 ee: 92.4%

Example 3

(21) (2-Fluoro-3-methylphenyl)methanol (151 mg; 1.08 mmol, CAS No. 307975-03-7) and CDI (175 mg; 1.08 mmol) were mixed together in DMF (3 ml); the reaction mixture was heated at 50° C. during 30 minutes. Example 3a (150 mg; 0.54 mmol) and DIPEA (0.28 ml; 1.62 mmol) were then added in sequence and the reaction mixture was stirred 16 hours at rt. The reaction mixture was diluted with 3 ml of a mixture MeOH/Water (1/1; volume/volume) before being filtered and separated via semipreparative HPLC. Obtained 80 mg of the desired compound.

(22) TABLE-US-00018 Example 3  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.69 MS: 372 (M + H).sup.+ Chiral SFC; Method: I_SA_20_MEOH_NH.sub.3_001 Rt [min]: 3.3 ee: 94.4%

Example 4

(23) (2-Fluorophenyl)methanol (116 µl; 1.08 mmol, CAS No. 446-51-5) and CDI (175 mg; 1.08 mmol) were mixed together in DMF (3 ml); the reaction mixture was heated at 50° C. during 30 minutes. Example 3a (150 mg; 0.54 mmol) and DIPEA (0.28 ml; 1.62 mmol) were then added in sequence and the reaction mixture was stirred 3 hours at rt.

(24) The reaction mixture was diluted with 3 ml of a mixture MeOH/Water (1/1; vol-ume/volume) before being filtered and separated via semipreparative HPLC. Obtained 98 mg of the desired

compound.

(25) TABLE-US-00019 Example 4  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.63 MS: 358 (M + H).sup.+ Chiral SFC; Method: I_SA_20_MEOH_NH.sub.3_001 Rt [min]: 3.37 ee: 100%

Example 5

(26) Example 5 was synthesised in analogy to Example 4. Starting materials: Example 3a (150 mg, 0.54 mmol) and (2,4-Difluorophenyl)methanol (121 μ l, 1.08 mmol, CAS No. 56456-47-4). The crude was purified via semipreparative HPLC. Obtained 112 mg of the desired compound.

(27) TABLE-US-00020 Example 5  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.64 MS: 376 (M + H).sup.+ Chiral SFC; Method: I_SA_20_MEOH_NH.sub.3_001 Rt [min]: 3.18 ee: 100%

Example 10

(28) (4-Fluorophenyl)methanol (157 μ l, 1.44 mmol, CAS No. 459-56-3) was dissolved in DMF (6 ml); TEA (0.3 ml; 2.16 mmol) was added followed by Bis-[1,2,4]triazol-1-yl-methanone (236 mg, 1.44 mmol) and Example 3a (200 mg, 0.72 mmol). The reaction mixture was stirred 2 h at 50° C. before being diluted with H.sub.2O/MeOH (4 ml; 1/1; volume/volume mixture), filtered and purified with semipreparative HPLC.

(29) Obtained 150 mg of the desired compound.

(30) TABLE-US-00021 Example 10  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.61 MS: 358 (M + H).sup.+ Chiral SFC; Method : I_SA_20_MEOH_NH.sub.3_001 Rt [min]; 4.25 ee: 100%

Example 12

(31) Example 12 was synthesised in analogy to Example 10. Starting materials: Example 3a (200 mg, 0.72 mmol) and (4-Methylphenyl)methanol (176 mg, 1.44 mmol, CAS No. 589-18-4). The reaction mixture was purified via semipreparative HPLC. Obtained 117 mg of the desired compound.

(32) TABLE-US-00022 Example 12  HPLC-MS; Method: Z011_S03; R.sub.t [min]: 0.67 MS: 354 (M + H).sup.+ Chiral SFC; Method: I_C2_40_IPA_NH.sub.3_001 Rt [min]; 3.65 ee: 66.7%

Claims

1. A compound of formula A ##STR00018## in which R.sub.1 represents phenyl which is optionally substituted with 1 to 3 substituents selected from the group consisting of fluoro, chloro, methyl, ethyl, cyclopropyl, F.sub.2HC—, FH.sub.2C—, and F.sub.3C—; or a pharmaceutically acceptable salt thereof.
2. The compound according to claim 1, or pharmaceutically acceptable salt thereof, wherein R.sub.1 represents phenyl which is optionally substituted with 1 or 2 substituents selected from the group consisting of fluoro and methyl.
3. The compound according to claim 1, or pharmaceutically acceptable salt thereof, wherein R.sub.1 represents ##STR00019##
4. An (S)-enantiomer of a compound according to claim 1, selected from the group consisting of: TABLE-US-00023 Exp 1      
5. A pharmaceutically acceptable salt of a compound according to claim 1.
6. A pharmaceutical composition comprising the compound according to claim 1 in admixture with a pharmaceutically acceptable adjuvant, diluent and/or carrier.
7. A method for treating and/or preventing a psychiatric disease and/or disorder, comprising administering to a patient in need thereof the compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein the disease and/or disorder is: bipolar disorder I

depressed, hypomanic, manic or mixed form; bipolar disorder II; single depressive episode, recurrent major depressive disorder, minor depressive disorder, depressive disorder with postpartum onset, depressive disorders with psychotic symptoms; or major depressive disorder with or without concomitant anxious distress, mixed features, melancholic features, atypical features, mood-congruent psychotic features, mood-incongruent psychotic features, or catatonia.

8. The method according to claim 7, characterized in that the compound is administered in combination with another antidepressant drug.

9. The method according to claim 7, wherein the patient is further undergoing behavioural therapy.
