



US012398374B2

(12) **United States Patent**
Fan et al.

(10) **Patent No.: US 12,398,374 B2**

(45) **Date of Patent: Aug. 26, 2025**

(54) **METHOD FOR PREPARING INDUCED PLURIPOTENT STEM CELLS BY REPROGRAMMING SOMATIC CELLS**

(71) Applicant: **ZHEJIANG HUODE BIOENGINEERING COMPANY LIMITED**, Hangzhou (CN)

(72) Inventors: **Jing Fan**, Hangzhou (CN); **Fang Ren**, Hangzhou (CN); **Anxin Wang**, Hangzhou (CN)

(73) Assignee: **ZHEJIANG HUODE BIOENGINEERING COMPANY LIMITED**, Zhejiang (CN)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 811 days.

(21) Appl. No.: **17/597,970**

(22) PCT Filed: **Jul. 31, 2020**

(86) PCT No.: **PCT/CN2020/106327**

§ 371 (c)(1),

(2) Date: **Jan. 31, 2022**

(87) PCT Pub. No.: **WO2021/018296**

PCT Pub. Date: **Feb. 4, 2021**

(65) **Prior Publication Data**

US 2022/0325248 A1 Oct. 13, 2022

(30) **Foreign Application Priority Data**

Aug. 1, 2019 (CN) 201910707125.0

(51) **Int. Cl.**

C12N 5/074 (2010.01)

C07K 14/47 (2006.01)

C07K 14/71 (2006.01)

C12N 5/10 (2006.01)

C12N 9/12 (2006.01)

C12N 15/85 (2006.01)

C12N 15/86 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 5/0696** (2013.01); **C07K 14/4705** (2013.01); **C12N 5/10** (2013.01); **C12N 15/85** (2013.01); **C12N 15/86** (2013.01); **C07K 14/71** (2013.01); **C12N 9/1205** (2013.01); **C12N 2501/01** (2013.01); **C12N 2501/15** (2013.01); **C12N 2501/603** (2013.01); **C12N 2501/605** (2013.01); **C12N 2501/727** (2013.01); **C12N 2506/11** (2013.01); **C12N 2506/1307** (2013.01); **C12N 2510/00** (2013.01); **C12N 2760/18843** (2013.01); **C12N 2820/60** (2013.01)

(58) **Field of Classification Search**

CPC **C12N 5/0696**; **C12N 5/10**; **C12N 15/85**; **C12N 2501/01**; **C12N 2501/603**; **C12N 2501/605**; **C12N 2510/00**; **C12N**

2760/18843; **C12N 2820/60**; **C12N 9/1205**; **C12N 2501/15**; **C12N 2501/415**; **C12N 2501/727**; **C12N 2506/11**; **C12N 2506/1307**; **C12N 2533/52**; **C12N 2533/90**; **C12N 15/86**; **C07K 14/4705**; **C07K 14/71**
USPC 435/456, 461, 325, 357, 366, 372, 377; 530/358; 536/23.1
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

10,231,999 B2 3/2019 Lee et al.
2019/0024047 A1 1/2019 Matsumoto et al.

FOREIGN PATENT DOCUMENTS

CN 104673741 A * 6/2015
CN 109679921 A 4/2019
EP 3272858 A1 1/2018
WO WO 2009/117439 A2 9/2009
WO WO 2012/087965 A2 6/2012
WO WO 2015/003643 A1 1/2015

OTHER PUBLICATIONS

Li et al. 2017, Cell Stem Cell, vol. 21, 264-273. (Year: 2017).
Chichagova et al. 2015, Method in Molecular Biology, vol. 1353, 285-307. (Year: 2015).
Storm et al. 2007, Journal of Biological Chemistry, vol. 282(9), 6265-6273. (Year: 2007).
Zhao et al. 2018, Journal of Experimental Zoology Part B: Molecular and Developmental Evolution, vol. 330(8), 406-416. (Year: 2018).
Zhou & Sun 2019, Current Molecular Medicine, vol. 19, 77-90. (Year: 2019).
Chinese Search Report for Chinese Patent Application No. 2020800064401, dated Aug. 18, 2022, 2 pages.
Extended European Search Report for Application No. 20847071.6, dated Sep. 15, 2022, 10 pages.

(Continued)

Primary Examiner — Maria G Leavitt

Assistant Examiner — Katie L Pennington

(74) Attorney, Agent, or Firm — Sughrue Mion, PLLC

(57) **ABSTRACT**

The present invention provides a method for preparing induced pluripotent stem cells through somatic cell reprogramming and induced pluripotent stem cells obtained therefrom. The present method comprises introducing the factors Oct4 and Nanog as reprogramming-inducing factors into somatic cells to perform reprogramming; followed by culturing the partially or fully reprogrammed somatic cells in a medium comprising specific chemical inducing agents to obtain induced pluripotent stem cells. In the present invention, the combination of different forms of reprogramming-inducing factors and three small-molecule compounds as chemical inducing agents can significantly improve the reprogramming efficiency of human somatic cells and reduce the tumorigenicity of the obtained induced pluripotent stem cells.

15 Claims, 6 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Ichida et al., "A-Small-Molecule Inhibitor of Tgf-beta Signaling Replaces Sox2 in Reprogramming by Inducing Nanog," *Cell Stem Cell*, 2009, 5:491-503.

Lewandowski et al., "Techniques of Human Embryonic Stem Cell and Induced Pluripotent Stem Cell Derivation," *Arch Immunol Ther Exp*, 2016, 64:349-370.

Li et al., "Generation of Human-Induced Pluripotent Stem Cells in the Absence of Exogenous Sox2," *Stem Cells* 2009, 27:2992-3000.

Li et al., "Generation of iPSCs from mouse fibroblasts with a single gene, Oct4, and small molecules," *Cell Res*, 2011, 21:196-204.

Xie et al., "Pharmacological Reprogramming of Somatic Cells for Regenerative Medicine," *Acc Chem Res*, 2017, 50:1202-1211.

Zhou et al., "Research Progress on Somatic Cell Reprogramming Mechanism," *Chin J Cell Biol*, 2019, 41(5): 805-821.

Extended European Search Report dated Sep. 30, 2024 from the European Patent Office in Application No. 21896169.6.


Adelya A Galiakberova et al., "Neural Stem Cells and Methods for Their Generation From Induced Pluripotent Stem Cells in vitro", *Frontiers in Cell and Developmental Biology*, Oct. 2020, vol. 8, Article 815, pp. 1-20 (20 pages total).

Jeong Su Byun et al., "Rapid differentiation of astrocytes from human embryonic stem cells", *Neuroscience Letters*, 2020, vol. 716, No. 134681, pp. 1-7 (7 pages total).

Jin-Chong Xu et al., "Cultured networks of excitatory projection neurons and inhibitory interneurons for studying human cortical neurotoxicity", *Neurotoxicity*, 2016, vol. 8, Issue 333, pp. 1-14 (14 pages total).

Su-Chun Zhang et al., "In vitro differentiation of transplantable neural precursors from human embryonic stems cells", *nature biotechnology*, 2001, vol. 19, pp. 1129-1133 (5 pages total).

* cited by examiner






Figure 2




Figure 3




Figure 4




Figure 5




Figure 6

1

METHOD FOR PREPARING INDUCED PLURIPOTENT STEM CELLS BY REPROGRAMMING SOMATIC CELLS

This application is a national stage filing under 35 U.S.C. § 371 of International Application No. PCT/CN2020/106327, filed on Jul. 31, 2020, which claims priority of Chinese Application No. 201910707125.0, filed on Aug. 1, 2019. The contents of each of these applications are incorporated herein by reference.

TECHNICAL FIELD

The present disclosure belongs to the technical field of stem cells, and particularly relates to a method of preparing induced pluripotent stem cells through somatic cell reprogramming, as well as the induced pluripotent stem cells obtained therefrom.

BACKGROUND

In 2006, Yamanaka's team developed a cocktail consisting of four genes, namely Oct4, Sox2, Klf4 and c-Myc, which was used to successfully reprogram terminally differentiated skin fibroblasts into induced pluripotent stem cells (iPSCs) by viral infection. One year later, James Thomson successfully reprogrammed human fibroblasts into iPSCs by episomal plasmid transfection using a different combination of four factors (Oct4, Sox2, Nanog and Lin28). The above methods, which are relatively simple and stable in technology, break through the ethical limitations of using human embryonic stem cells in medicine, capable of solving the problem of immune rejection in cell transplantation therapy, and greatly expand the application potential of stem cell technology in clinical medicine. In addition, iPSCs technology and functional cells obtained by subsequent differentiation have great potential value in areas such as cell replacement therapy, pathogenesis research, and new drug screening.

However, with the development of iPSCs reprogramming technology, the drawbacks and problems of iPSCs reprogramming by exogenous introduction of transcription factors have gradually become prominent, such as incomplete reprogramming of some cells, low reprogramming efficiency, and oncogenicity of the introduced proto-oncogenes. Therefore, researchers are trying to find alternatives to the four factors and to optimize the reprogramming system, such as replacing oncogenic transcription factors, adding transcription factors or compounds that enhance reprogramming efficiency, or even completely giving up transcription factors and switching to small molecule compounds to induce somatic cell reprogramming, so as to avoid the potential defects to a certain extent.

At present, a variety of small molecule compounds have been found to induce the generation of pluripotent stem cells. Some of these small molecules directly improve the reprogramming efficiency of recipient cells by inhibiting genomic methylation; some of them affect specific signaling pathways, rendering the intermediate transitional cells and partially reprogrammed cells generated during the reprogramming process to be transformed into stably and completely induced pluripotent stem cells. For example, TGF β can induce the expression of EMT-related gene SNAIL in the early stage of reprogramming, thereby inducing EMT and hindering the reprogramming process; on the contrary, TGF β receptor inhibitors can improve the reprogramming efficiency and can be used to replace Sox2 and c-Myc in

2

Yamanaka four factors; cyclic AMP (cAMP) agonists can increase Oct4 expression; glycogen synthase kinase (GSK) inhibitors can increase Nanog expression, thereby improving the reprogramming efficiency of specific cells. Some methods even use a cocktail of purely chemical small molecules without involving any exogenously introduced transcription factors. However, such methods have relatively low reprogramming efficiency and are time-consuming, which cannot meet clinical needs. Some of the methods have not been used for the reprogramming of human cells.

To date, researchers have tried reprogramming techniques on many different types of cells and have achieved success. iPSCs can be obtained by reprogramming using genes, RNAs or proteins of the four factors.

Collectively the characteristics of the above methods, it is important and meaningful to provide a preparation method of adding a combination of reprogramming-inducing factors in different types of vectors and small molecule compounds, so as to improve the reprogramming efficiency, and to obtain iPSCs by safely and effectively reprogramming human somatic cells.

SUMMARY OF THE INVENTION

One purpose of the present invention is to provide a method of preparing induced pluripotent stem cells (iPSCs) by reprogramming of somatic cells.

Another purpose of the present invention is to provide induced pluripotent stem cells (iPSCs) prepared by the method as above.

The present inventors found that by introducing only two transcription factors as reprogramming-inducing factors, accompanied with three small molecule compounds as chemical inducing agents, a simpler and more efficient reprogramming method with broader application potentials as compared to the prior art can be achieved on somatic cells from different origins of the human body, and iPSCs were successfully obtained therefrom, thus completing the present invention.

Accordingly, in a first aspect, the present invention provides a method for preparing induced pluripotent stem cells through somatic cell reprogramming, comprising the following steps: (1) introducing Oct4 and Nanog as reprogramming-inducing factors into somatic cells to perform reprogramming; (2) culturing partially or completely reprogrammed somatic cells obtained in step (1) in the presence of chemical inducing agents to obtain induced pluripotent stem cells (iPSCs), wherein the chemical inducing agents comprise a TGF β receptor inhibitor, a cyclic AMP (cAMP) agonist and a glycogen synthase kinase (GSK) inhibitor.

Preferably, the TGF β receptor inhibitor is 616452, and/or the cyclic AMP (cAMP) agonist is Forskolin, and/or the glycogen synthase kinase (GSK) inhibitor is TD114-2.

In the method of the present invention, preferably, Oct4 and Nanog as the reprogramming-inducing factors can be introduced into somatic cells in the form of their DNAs, in the form of their RNAs, or in the form of their protein products. In the method of the present invention, preferably, the somatic cells can be any cells known in the art, for example, skin-derived cells, blood-derived cells, and can also include urine-derived cells, liver cells, epithelial cells, gastric cells, keratinocytes and the like. In a more preferred embodiment, the skin-derived cells may be skin fibroblasts. In another more preferred embodiment, the blood-derived cells may be erythroid progenitor cells.

In the method of the present invention, preferably, the somatic cells are derived from human, that is, human somatic cells.

In the method of the present invention, preferably, the working concentration of the TGF β receptor inhibitor such as 616452 is 0.1-20 μ M, more preferably 5-10 μ M, and still more preferably 5 μ M.

In the method of the present invention, preferably, the working concentration of the cyclic AMP (cAMP) agonist such as Forskolin is 0.1-50 μ M, more preferably 2-20 μ M, and still more preferably 10 μ M.

In the method of the present invention, preferably, the working concentration of the glycogen synthase kinase (GSK) inhibitor such as TD114-2 is 0.1-20 μ M, more preferably 2-10 μ M, still more preferably 5 μ M.

In a specific embodiment, in step (1), reprogramming is conducted by transfecting the vector(s) comprising the coding sequences of the reprogramming-inducing factors Oct4 and Nanog, respectively, into somatic cells by electroporation or chemical transfection. For example, the transfection can be performed by electroporation. Specifically, the electroporation method may include adding reprogramming vector(s) comprising the inducing factors Oct4 and Nanog to cell suspension, transferring into electroporation cuvette after mixing, placing the electroporation cuvette in the electroporation apparatus, and performing electroporation.

In another specific embodiment, in step (1), reprogramming is conducted by infecting the somatic cells with a virus comprising Oct4 and Nanog reprogramming-inducing factors. Preferably, the virus is Sendai Virus.

In the method of the present invention, preferably, the aforesaid specific chemical inducing agents are added to the culture medium from Day 2 after the introduction of the reprogramming-inducing factors.

In a second aspect, the present invention provides an induced pluripotent stem cell which is obtained by the method of the first aspect.

In a third aspect, the present invention relates to a combination of exogenously introduced two inducing factors and three chemical inducing agents, for use in reprogramming somatic cells to prepare iPSCs, wherein the exogenously introduced two inducing factors are Oct4 and Nanog, and the three chemical inducing agents consist of a TGF β receptor inhibitor such as 616452, a cyclic AMP (cAMP) agonist such as Forskolin, and a glycogen synthase kinase (GSK) inhibitor such as TD114-2. Specifically, the Oct4 and Nanog can be introduced into somatic cells in the form of their DNAs, RNAs or proteins.

In the present invention, different forms of reprogramming-inducing factors in combination with the three small molecule compounds as chemical inducing agents can significantly improve the reprogramming efficiency of human somatic cells and reduce the tumorigenicity of the obtained iPSCs.

DESCRIPTION OF DRAWINGS

FIG. 1A shows two brightfield images of iPSCs obtained by the methods as described in Example 1, in which the upper panel of FIG. 1A shows the iPSC colonies after reprogramming for about 12 days, and the lower panel of FIG. 1A shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture. FIG. 1B shows two brightfield images of iPSCs obtained by the methods as described in Example 2, in which the upper panel of FIG. 1B shows the iPSC colonies after reprogramming for about 12 days, and the lower panel

of FIG. 1B shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture. FIG. 1C shows two brightfield images of iPSCs obtained by the methods as described in Example 3, in which the upper panel of FIG. 1C shows the iPSC colonies after reprogramming for about 12 days, and the lower panel of FIG. 1C shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture.

FIGS. 2A-E are the detection results of flow cytometry, showing the expression of pluripotent cell markers by iPSCs obtained by reprogramming of skin fibroblasts. (A) SSEA4; (B) Tra-1-81; (C) Tra-1-60; (D) Oct4; (E) Nanog.

FIGS. 3A-B show the results of (A) immunofluorescence staining and (B) qPCR identification of differentiation of three germ layers from iPSCs obtained by reprogramming skin fibroblasts.

FIG. 4A shows two brightfield images of iPSCs obtained by the methods as described in Example 5, in which the upper panel of FIG. 4A shows the iPSC colonies after reprogramming for about 12 days, and the lower panel of FIG. 4A shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture. FIG. 4B shows two brightfield images of iPSCs obtained by the methods as described in Example 6, in which the upper panel of FIG. 4B shows the iPSC colonies after reprogramming for about 12 days, and the lower panel of FIG. 4B shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture. FIG. 4C shows two brightfield images of iPSCs obtained by the methods as described in Example 7, in which the upper panel of FIG. 4C shows the iPSC colonies after reprogramming for about 12 days, and the lower panel of FIG. 4C shows the iPSC colonies after picking single colonies and inoculating into a feeder-free system for culture.

FIGS. 5A-E are the detection results of flow cytometry, showing the expression of pluripotent cell markers by iPSCs obtained by reprogramming of erythroid progenitor cells. (A) SSEA4; (B) Tra-1-81; (C) Tra-1-60; (D) Oct4; (E) Nanog.

FIGS. 6A-B show the results of (A) immunofluorescence staining and (B) qPCR identification of differentiation of three germ layers from iPSCs obtained by reprogramming erythroid progenitor cells.

DETAILED DESCRIPTION

In the method of the present invention, only Oct4 and Nanog are used as reprogramming-inducing factors, and reprogramming is performed by introducing them into somatic cells. Both Oct4 and Nanog are transcription factors that play important roles in maintaining pluripotency. A variety of transcription factors have been identified in the prior art that can be used to induce the reprogramming of somatic cells into induced pluripotent stem cells, such as Oct4, c-Myc, Sox2, and Klf4 consisting the Yamanaka four-factor combination as described above, and Oct4, Sox2, Nanog and Lin28 consisting the James Thomson four-factor combination. But the method of the present invention uses only Oct4 in the Yamanaka four factors and Nanog in the James Thomson four factors. In other words, the method of the present invention does not use any transcription factors other than Oct4 and Nanog as reprogramming-inducing factors.

Oct4 and Nanog can be introduced by a method known in the art for introducing transcription factors. Such methods

include, but are not limited to, infecting somatic cells by introducing virus comprising recombinant DNA vector(s), mRNAs or RNAs of the nucleotides encoding Oct4 and Nanog so as to allow the expression of Oct4 and Nanog inducing factors, or by directly introducing Oct4 and Nanog in the form of proteins into somatic cells.

In one embodiment of the method of the present invention, the two reprogramming-inducing factors are introduced into somatic cells in the form of DNA. Specifically, a nucleotide sequence encoding Oct4 and a nucleotide sequence encoding Nanog can be introduced into somatic cells. In a specific embodiment, the nucleotide sequence encoding Oct4 comprises the nucleotide sequence as shown in SEQ ID NO: 6 or is consisted of the nucleotide sequence as shown in SEQ ID NO: 6; or comprises a nucleotide sequence that has at least 90% homology to the nucleotide sequence of SEQ ID NO: 6 and encodes Oct4; or is consisted of a nucleotide sequence that has at least 90% homology to the nucleotide sequence of SEQ ID NO: 6 and encodes Oct4. In a specific embodiment, the nucleotide sequence encoding Nanog comprises the nucleotide sequence as shown in SEQ ID NO: 15 or is consisted of the nucleotide sequence as shown in SEQ ID NO: 15; or comprises a nucleotide sequence that has at least 90% homology to the nucleotide sequence of SEQ ID NO: 15 and encodes Nanog; or is consisted of a nucleotide sequence that has at least 90% homology to the nucleotide sequence of SEQ ID NO: 15 and encodes Nanog.

The nucleotide sequence encoding Oct4 and the nucleotide sequence encoding Nanog can be placed in the same vector or in different vectors. When placed in the same vector, the nucleotide sequence encoding Oct4 and the nucleotide sequence encoding Nanog may be under the control of the same or different regulatory sequences. Regulatory sequences can be selected according to the type of target cell. In a specific embodiment, the nucleotide sequence encoding Oct4 and the nucleotide sequence encoding Nanog are placed in the same vector, such as pcDNA3.1. Additional elements, such as the coding sequence of EBNA1 and OriP sequence, may be comprised in the recombinant vector to increase the efficiency of plasmid replication in cells. In a specific embodiment, the sequence of the recombinant vector used in the present invention is shown in SEQ ID NO: 13 or SEQ ID NO: 24.

Methods for delivering vector(s) comprising nucleotides of interest into somatic cells are known in the art and include, but are not limited to, electroporation, gene gun, lipofection, calcium-mediated transfection. In a specific embodiment, electroporation is used.

In the method of the present invention, the above-mentioned inducing factors Oct4 and Nanog can be expressed in vitro to obtain corresponding proteins, which are then introduced into differentiated cells, thereby achieving the object of the present invention. Techniques for introducing proteins into cells are well known in the art and include, but are not limited to, Tat-delivery and related techniques, electroporation (nucleofection), protein and cellular ligand binding.

Those of ordinary skill in the art can also understand that the DNA sequences of the above-mentioned Oct4 and Nanog inducing factors can be transcribed in vitro, and the obtained mRNAs are directly introduced into differentiated cells to express the corresponding proteins in the cells, thus achieving the purpose of the present invention. Reprogramming can also be performed by infecting somatic cells with Sendai Virus comprising RNAs of the two inducing factors.

Sendai Virus is a non-integrating virus that will not integrate into the genome of infected cells and thus has relatively high safety.

After introduction of the reprogramming-inducing factors Oct4 and Nanog into somatic cells, cells were cultured in the presence of one or more of a TGF β receptor inhibitor, a cyclic AMP (cAMP) agonist, and a glycogen synthase kinase (GSK) inhibitor as chemical inducing agent(s) to generate iPSCs. In preferred embodiments, a TGF β receptor inhibitor, a cyclic AMP (cAMP) agonist, and a glycogen synthase kinase (GSK) inhibitor are used.

In a preferred embodiment, the TGF β receptor inhibitor is 616452, whose chemical name is 2-[3-(6-methyl-2-pyridinyl)-1H-pyrazol-4-yl]-1,5-naphthyridine (CAS No: 446859-33-2). Preferably, the TGF β receptor inhibitor is used at a working concentration of 0.1-20 μ M, more preferably 5-10 μ M, still more preferably 5 μ M. For example, the TGF β receptor inhibitor is added to the medium at such concentration.

In a preferred embodiment, the cyclic AMP (cAMP) agonist is Forskolin, whose chemical name is [(3R,4aR,5S,6S,6aS,10S,10aR,10bS)-3-ethenyl-6,10,10b-trihydroxy-3,4a,7,7,10a-pentamethyl-1-oxo-5,6,6a,8,9,10-hexahydro-2H-benzo[f]chromen-5-yl]acetate (CAS No: 66575-29-9). Preferably, the cAMP agonist is used at a working concentration of 0.1-50 μ M, more preferably 2-20 μ M, still more preferably 10 μ M. For example, the cAMP agonist is added to the medium at such concentration.

In a preferred embodiment, the glycogen synthase kinase (GSK) inhibitor is TD114-2, whose chemical name is 6,7,9,10,12,13,15,16,18,19-Decahydro-5,29:20,25-dimetheno-26H-dibenzo [n,t]pyrrolo[3,4-q][1,4,7,10,13,22] tetraoxa diazacyclotetracosine-26,28(27H)-dione (CAS No: 436866-52-3). Preferably, the GSK inhibitor is used at a working concentration of 0.1-20 μ M, more preferably 2-10 μ M, still more preferably 5 μ M. For example, the GSK inhibitor is added to the medium at such concentration.

The medium used in the method of the present invention can be selected by a person skilled in the art based on existing knowledge depending on the type of cells to be cultured.

EXAMPLES

In order to facilitate understanding of the present invention, the technical solutions of the present invention will be further exemplified below through specific examples. Those of ordinary skill in the art can understand that the present invention is not limited to the described examples, and those of ordinary skill in the art can make modifications to the examples based on the teachings of the disclosure. Such modifications are also included within the scope of the present invention as defined by the appended claims.

The experimental methods in the following examples are conventional methods unless otherwise specified.

Example 1. Inducing the Reprogramming of Skin Cells by Using a Vector Encoding Reprogramming Factors

1.1 Human-derived skin tissue was placed into a petri dish, and was quickly and repeatedly rinsed for 4 times with phosphate buffered saline (PBS) pre-cooled at 4° C.

1.2 The skin tissue was processed with sterilized ophthalmic scissors and a scalpel to remove the subcutaneous white fat, remove the epidermis and subcutaneous tissue, and leave the dermis layer to obtain pretreated skin tissues. The

pretreated skin tissue was transferred in the medium, and cut into small pieces to obtain skin tissue pieces with neat edges.

1.3 3 mL of fetal bovine serum (FBS) was evenly added to each well of a 6-well cell culture plate. The skin tissue pieces were added and incubated in the incubator for 0.5-1 h to make the tissues adhere to the bottom of the culture plate.

1.4 1 mL of DMEM (Gibco) medium comprising 20% (v/v) FBS was added to each well of the 6-well cell culture plate, and the plate was put back into the incubator for culture.

1.5 The medium was exchanged when fibroblasts migrated out of the skin tissue pieces as observed under the microscope. The amount of DMEM medium was increased to 3 mL, and the medium was exchanged every 1-3 days.

1.6 When the fibroblasts converged to the edge of each well, the spent medium was discarded. After washing twice with PBS, 1 mL of 0.25% trypsin-EDTA (GIBCO) was added, and the plate was placed in a cell culture incubator for about 4-6 min.

When the cells became round, detached and floated, DMEM medium comprising 20% (v/v) FBS was immediately added to terminate the digestion. The cells were pipetted several times gently and transferred into a 15 mL centrifuge tube, and centrifuged at 200 g for 3 min. The supernatant was aspirated and discarded. The cell culture medium was added to the tube and mixed thoroughly. The cells were transferred to a culture flask, which was placed in a cell incubator to continue culturing.

1.7 When the second-generation skin fibroblasts reached 80%-90% confluence, the old medium was discarded. The cells were washed twice with PBS, followed by digestion via adding 3 mL of 0.25% trypsin-EDTA (Gibco), until the cells were dispersed. DMEM medium comprising 20% (v/v) FBS was added to stop digestion. The cells were pipetted several times gently and transferred into a 15 mL centrifuge tube, and centrifuged at 200 g for 3 min. The supernatant was discarded. A suitable amount of PBS was added to resuspend the cells, which were counted by using a hemocytometer.

1.8 About 8×10^5 skin fibroblasts were taken for centrifuge at 200 g for 3 min. The cells were washed twice with PBS and once with OPTI-MEM (Gibco). The supernatant was discarded. Corresponding electroporation reagents were added according to the instructions of the Celtrix Kit. Cells were resuspended. The reprogramming vector comprising the inducing factors Oct4 and Nanog were added to the cell suspension, mixed well and transferred to an electroporation cuvette. The electroporation cuvette was placed in an electroporation apparatus, and electroporation was conducted at 430 V, 30 ms.

Specifically, the reprogramming vector was constructed as follows: ① Using the plasmid (Addgen #20922) as the template, KOD-Plus-Neo (TOYOBO #KOD-401) high-fidelity enzyme and primers F1/R1, F2/R2, F3/R3 were used to perform the amplification, resulting in the fragment of EF-1 α promoter, Oct4 coding sequence and Nanog coding sequence, respectively (see Table 1 below for the primer sequences and the sequences obtained by the amplification). ② Using the pcDNA3.1(-) plasmid as template, amplification was performed by using KOD-Plus-Neo (TOYOBO #KOD-401) high-fidelity enzyme and primers F4/R4 to obtain the pcDNA3.1 fragment (see Table 1 below for the primer sequences and the sequence obtained by the amplification). ③ Using NEBuilder HiFi DNA Assembly Bundle for Large Fragments (NEB #E2623), the pcDNA3.1 was ligated with the EF-1 α promoter, the Oct4 coding sequence and the Nanog coding sequence by homologous recombination,

so that the coding sequences of Oct4 and Nanog were under the control of the EF-1 α promoter, resulting in the target vector, namely pcDNA3.1-EF-Oct4-Nanog (SEQ ID NO: 13).

1.9 After the electroporation, the electroporation cuvette was taken out. The cell suspension was quickly aspirated and added to Matrigel (Corning)-coated cell culture plate comprising 2 mL of DMEM medium comprising 20% (v/v) FBS added in advance. On Day 2, the medium was changed to 2-3 mL of TeSR-E7 medium (Stem Cell Technologies). Static culture was conducted at 37° C. in cell incubator. The medium was completely exchanged every day since then. Starting from Day 2, medium being used was supplemented with a combination of small molecule compounds, in which the working concentrations of the compounds were as follows. The working concentration of TGF β receptor inhibitor 616452 was 5 μ M. The working concentration of cyclic AMP (cAMP) agonist Forskolin was 10 μ M. The working concentration of glycogen synthase kinase (GSK) inhibitor TD114-2 was 5 μ M.

After reprogramming for about 12 days, iPSC colonies with cell morphology distinct from those of skin fibroblasts were observed (see the upper panel of FIG. 1A). Single colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 1A).

Example 2. Inducing the Reprogramming of Skin Cells by Using a Vector Encoding Reprogramming Factors

The same protocol was used as described in steps 1.1-1.9 of Example 1. Fibroblasts derived from human skin tissue were used for conducting the method of the present invention. As compared to Example 1, the construction of the reprogramming vector in step 1.8 is different. On the basis of the construct of Example 1, EBNA1 and OriP elements were added, so that the replication efficiency of the plasmid could be enhanced in somatic cells based on the EBNA1/OriP system. The construction method is described in detail as follows.

① Using plasmid (Addgen #20922) as the template, KOD-Plus-Neo (TOYOBO #KOD-401) high-fidelity enzyme and primers F5/R5, F6/R6, F7/R7, F8/R8, F9/R9 were used to perform the amplification, resulting in the fragment of EF-1 α promoter, Oct4 coding sequence, Nanog coding sequence, EBNA1 coding sequence and OriP sequence, respectively (see Table 1 below for the primer sequences and the sequences obtained by the amplification). ② Using the pcDNA3.1(-) plasmid as the template, amplification was performed by using KOD-Plus-Neo (TOYOBO #KOD-401) high-fidelity enzyme and primers F10/R10 to obtain the pcDNA3.1 fragment (see Table 2 below for the primer sequences and the sequence obtained by the amplification). ③ Using NEBuilder HiFi DNA Assembly Bundle for Large Fragments (NEB #E2623), the pcDNA3.1 was ligated with the EF-1 α promoter, the Oct4 coding sequence, the Nanog coding sequence, the EBNA1 coding sequence and the OriP sequence by homologous recombination, resulting in the target vector, namely pcDNA3.1-EF-Oct4-Nanog-EBNA1-OriP (SEQ ID NO: 24).

After reprogramming for about 12 days, iPSC colonies with cell morphology distinct from those of skin fibroblasts were observed (see the upper panel of FIG. 1B). Single

colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 1B).

Example 3. Inducing the Reprogramming of Skin Cells by Using a Virus Comprising RNAs of Reprogramming-Inducing Factors

The same protocol was used as described in steps 1.1-1.6 of Example 1. Fibroblasts derived from human skin tissue were used for conducting the method of the present invention. Different from steps 1.7-1.9 of Example 1, reprogramming of iPSCs was performed by using a virus comprising RNAs of reprogramming-inducing factors as described in step 3.7 below.

3.7 When the second-generation skin fibroblasts reached 80%-90% confluence, Sendai Virus comprising Oct4 and Nanog reprogramming-inducing factors was mixed with the cells and cultured for 2 days. The medium was discarded. 2 mL of DMEM medium with 20% (v/v) FBS was added to culture, and the medium was changed every other day. After culturing for 5 days, 1 mL of 0.25% trypsin-EDTA (GIBCO) was added, followed by culture in a cell incubator for about 4-6 min. When the cells became round, detached and floated, they were added to Vitronectin (Gibco)-coated cell culture plates for culture. Starting from Day 2 of the culture, medium being used was supplemented with a combination of small molecule compounds, in which the working concentrations of the compounds were as follows. The working concentration of TGF β receptor inhibitor 616452 was 5 μ M. The working concentration of cyclic AMP (cAMP) agonist Forskolin was 10 μ M. The working concentration of glycogen synthase kinase (GSK) inhibitor TD114-2 was 5 μ M.

After reprogramming for about 12 days, iPSC colonies with cell morphology distinct from those of skin fibroblasts were observed (see the upper panel of FIG. 1C). Single colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 1C).

Example 4. Identification of iPSCs Obtained by Reprogramming Skin Fibroblasts

The iPSCs obtained by reprogramming skin fibroblasts were characterized and identified using different methods, including flow cytometry, immunofluorescence staining and quantitative PCR (qPCR).

The obtained iPSCs were analyzed by flow cytometry for the following molecular markers: SSEA4, Tra-1-81, Tra-1-60, Oct4 and Nanog. As shown in FIG. 2, the iPSCs obtained by the method of the present invention expressed the markers of human pluripotent stem cells including SSEA4, Tra-1-81, Tra-1-60, Oct4 and Nanog, which proved that the obtained cells possessed the characteristics of pluripotent stem cells.

To verify the totipotency, the obtained iPSCs were induced to differentiate into three germ layers in vitro. The expression of molecular markers specific for the three germ layers was detected by immunofluorescence staining and qPCR. As shown in FIG. 3A, results of the immunofluorescence experiments using three markers (endoderm: SOX17;

mesoderm: CDX2; ectoderm: PAX6) showed that the obtained iPSCs could differentiate into cells of all three germ layers. FIG. 3B shows the expression of molecular markers specific for three germ layers (endoderm: SOX17, mesoderm: MIXL1, and ectoderm: PAX6) detected by qPCR, in which the iPSCs expressed the molecular markers of pluripotent stem cell, OCT4 and TRA-1-81, at high levels (consistent with the aforementioned results of flow cytometry assay), while cells differentiated into three germ layers expressed the molecular markers specific for each germ layer (****p<0.0001, Student's t-test). These results indicated that the obtained iPSCs were totipotent.

Example 5. Inducing the Reprogramming of Peripheral Blood Cells by Using a Vector Encoding Reprogramming Factors

5.1 10 mL of human peripheral blood was collected, from which the erythroid progenitor cells were enriched and expanded by using RosetteSep™ kit and SepMate™ kit (Stem Cell Technologies). Specifically, 10 mL of blood was transferred from the blood collection tube to a regular centrifuge tube, to which 50 μ L of RosetteSep™ Human Progenitor Cell Basic Pre-Enrichment Cocktail was added and mixed thoroughly. Then the tubes were placed at room temperature for 10 min.

5.2 3.5 mL of Lymphoprep™ was added to a SepMate centrifuge tube along the central hole. After incubating at room temperature for 10 minutes, 10 mL of PBS+2% FBS was added and mixed thoroughly. 5 mL of blood was added along the wall of the SepMate centrifuge tube. Immediately after centrifuge at 1200 g for 10 min, the yellow supernatant was poured carefully into a new regular centrifuge tube (be careful not to pour in the unwanted cells at bottom), and centrifuged at 300 g for 8 min.

5.3 After centrifugation, the supernatant was discarded. The cells were resuspended with 0.5 mL of StemSpan™ SFEM II medium and counted for the total number. 2 mL of cells at a density of 5×10^6 /mL was inoculated into each well of a 6-well plate, and cultured at 37° C., 5% CO₂ in cell incubator.

5.4 Day 1: the cell suspension was transferred to a new 6-well plate to remove adherent unwanted cells. Each well was supplemented with 0.5 mL StemSpan™ SFEM II medium, and incubated at 37° C., 5% CO₂ in cell incubator.

5.5 Day 2, Day 4, Day 6 and Day 8: Cell suspension from each well was collected into a centrifuge tube. Centrifuge was conducted at 400 g for 5 min, and the supernatant was discarded. StemSpan™ SFEM II medium was added to resuspend the cells. After pipetting for 3-4 times, 2 mL was inoculated into each well of a new 6-well plate, shaken thoroughly and cultured at 37° C., 5% CO₂ in cell incubator.

5.6 Day 9: The number of erythroid progenitor cells significantly increased. Further, the erythroid progenitor cells rapidly enriched on Day 10. When the number of cells reached 2×10^6 , electroporation could be performed. The cell suspension was collected and centrifuged at 400 g for 5 min. The supernatant was discarded. According to the instructions of the Cellectra Kit, the corresponding electroporation reagents were added to resuspend the cells. Into the cell suspension, the reprogramming vector comprising inducing factors Oct4 and Nanog as constructed in Example 1, namely pcDNA3.1-EF-Oct4-Nanog (SEQ ID NO: 13), was added. After mixing thoroughly, the mixture was added to a electroporation cuvette, which was placed in electroporation apparatus to perform the electroporation at 820 V, 20 ms.

11

5.7 After the electroporation, the electroporation cuvette was taken out. The cell suspension was quickly aspirated and added to Matrigel (Corning)-coated cell culture plate comprising StemSpan™ SFEM II added in advance. On Day 2, 1 mL StemSpan™ SFEM II medium was supplemented to each well. Static culture was conducted at 37° C. in cell incubator. On Day 3 and Day 5, 1 mL ReproTeSR medium (Stem Cell Technologies) was supplemented to each well. On Day 7, it could be observed that cells began to grow adherent to the wall. The spent medium was discarded. Afterwards, the medium was replaced with 2 mL of fresh ReproTeSR medium for each well every day. Starting from Day 2, medium being used was supplemented with a combination of small molecule compounds, in which the working concentrations of the compounds were as follows. The working concentration of TGFβ receptor inhibitor 616452 was 5 μM. The working concentration of cyclic AMP (cAMP) agonist Forskolin was 10 μM. The working concentration of glycogen synthase kinase (GSK) inhibitor TD114-2 was 5 μM. After reprogramming for about 12 days, iPSC colonies derived from adherent growth of the cell mass in suspension culture could be observed (see the upper panel of FIG. 4A). Single colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 4A).

Example 6. Inducing the Reprogramming of
Peripheral Blood Cells by Using a Vector Encoding
Reprogramming Factors

The same protocol was used as described in steps 5.1-5.7 of Example 5. Erythroid progenitor cells derived from human peripheral blood were used for conducting the method of the present invention. Different from Example 5, the reprogramming vector used in step 5.6 was replaced by the reprogramming vector comprising inducing factors Oct4 and Nanog as constructed in Example 2, namely pcDNA3.1-EF-Oct4-Nanog-EBNA1-OriP.

After reprogramming for about 12 days, iPSC colonies derived from adherent growth of the cell mass in suspension culture could be observed (see the upper panel of FIG. 4B). Single colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 4B).

Example 7. Inducing the Reprogramming of
Peripheral Blood Cells by Using a Virus
Comprising RNAs of Reprogramming-Inducing
Factors

The same protocol was used as described in steps 5.1-5.7 of Example 5. Erythroid progenitor cells derived from human peripheral blood were used for conducting the method of the present invention. Different from steps 5.6-5.7 of Example 5, reprogramming of iPSCs was performed by using a virus comprising RNAs of reprogramming-inducing factors as described in step 7.6 below.

7.6 On Day 9: The number of erythroid progenitor cells significantly increased. The cell suspension was collected and centrifuged at 400 g for 5 min. The supernatant was

12

discarded. Sendai Virus comprising Oct4 and Nanog reprogramming-inducing factors in the form of RNAs was added to the cells and mixed. The cells were cultured for 2 days. The medium was discarded. 2 mL of ReproTeSR medium was added. Afterwards, the medium was replaced with 2 mL of fresh ReproTeSR medium for each well every day. Starting from Day 2, medium was supplemented with a combination of small molecule compounds, in which the working concentrations of the compounds were as follows. The working concentration of TGFβ receptor inhibitor 616452 was 5 μM. The working concentration of cyclic AMP (cAMP) agonist Forskolin was 10 μM. The working concentration of glycogen synthase kinase (GSK) inhibitor TD114-2 was 5 μM. After reprogramming for about 12 days, iPSC colonies derived from adherent growth of the cell mass in suspension culture could be observed (see the upper panel of FIG. 4C). Single colonies were picked and inoculated into a feeder-free system for culture. These iPSC colonies formed by cells with relatively small size were compact and well-defined iPSC colonies, in which the cells had a relatively large nuclei and high nucleocytoplasmic ratio, showing a typical iPSC morphology (see the lower panel of FIG. 4C).

Example 8. Identification of iPSCs Obtained by
Reprogramming Erythroid Progenitor Cells

The iPSCs obtained by reprogramming erythroid progenitor cells were characterized and identified using different methods, including flow cytometry, immunofluorescence staining and quantitative PCR (qPCR).

The obtained iPSCs were analyzed by flow cytometry for the following molecular markers: SSEA4, Tra-1-81, Tra-1-60, Oct4 and Nanog. As shown in FIG. 5, the iPSCs obtained by the method of the present invention expressed the markers of human pluripotent stem cells including SSEA4, Tra-1-81, Tra-1-60, Oct4 and Nanog, which proved that the obtained cells possessed the characteristics of pluripotent stem cells.

To verify the totipotency, the obtained iPSCs were induced to differentiate into three germ layers in vitro. The expression of molecular markers specific for the three germ layers was detected by immunofluorescence staining and qPCR. As shown in FIG. 6A, results of the immunofluorescence experiments using three markers (endoderm: SOX17; mesoderm: CDX2; ectoderm: PAX6) showed that the obtained iPSCs could differentiate into cells of all three germ layers. FIG. 6B shows the expression of molecular markers specific for three germ layers (endoderm: SOX17, mesoderm: MIXL1, and ectoderm: PAX6) detected by qPCR, in which the iPSCs expressed the molecular markers of pluripotent stem cell, OCT4 and TRA-1-81, at high levels (consistent with the aforementioned results of flow cytometry assay), while cells differentiated into three germ layers expressed the molecular markers specific for each germ layer (****p<0.0001, Student's t-test). These results indicated that the obtained iPSCs were totipotent.

TABLE 1

Products of PCR amplification				
No.	Gene name	Primer name	5'-3'	Sequence after bp amplification
1	Ef-1α	F1	SEQ ID NO: 1	35 SEQ ID NO: 3
		R1	SEQ ID NO: 2	35

13

TABLE 1-continued

Products of PCR amplification				
No.	Gene name	Primer name	5'-3'	Sequence after bp amplification
2	Oct4	F2	SEQ ID NO: 4	20 SEQ ID NO: 6
		R2	SEQ ID NO: 5	20
3	Nanog	F3	SEQ ID NO: 7	92 SEQ ID NO: 9
		R3	SEQ ID NO: 8	35
4	pcDNA3.1	F4	SEQ ID NO: 10	20 SEQ ID NO: 12
		R4	SEQ ID NO: 11	20

TABLE 2

Products of PCR amplification				
No.	Gene name	Primer name	5'-3'	Sequence after bp amplification
1	EF-1 α	F5, identical to F1	SEQ ID NO: 1	35 SEQ ID NO: 3
		R5, identical to R1	SEQ ID NO: 2	35
2	Oct4	F6, identical to F2	SEQ ID NO: 4	20 SEQ ID NO: 6
		R6, identical to R2	SEQ ID NO: 5	20
3	Nanog	F7, identical to F3	SEQ ID NO: 7	92 SEQ ID NO: 15
		R7	SEQ ID NO: 14	92

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 24

<210> SEQ ID NO 1
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ef-1 α F1 primer

<400> SEQUENCE: 1

atacgactca ctataggctc cgggtgccgt cagtg 35

<210> SEQ ID NO 2
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ef-1 α R1 primer

<400> SEQUENCE: 2

caggtgtccc gccatcacga cacctgaaat ggaag 35

<210> SEQ ID NO 3
 <211> LENGTH: 1192
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Ef-1 α amplicon

<400> SEQUENCE: 3

atacgactca ctataggctc cgggtgccgt cagtgggcag agcgcacatc gccacagtc 60
 cccgagaagt tggggggagg ggtcggaat tgaaccgggt cctagagaag gtggcgcggg 120
 gtaaaactggg aaagtgatgt cgtgtactgg ctccgccttt ttcccagggt tgggggagaa 180

14

TABLE 2-continued

Products of PCR amplification				
No.	Gene name	Primer name	5'-3'	Sequence after bp amplification
4	EBNA1	F8	SEQ ID NO: 16	38 SEQ ID NO: 18
		R8	SEQ ID NO: 17	85
5	OriP	F9	SEQ ID NO: 19	101 SEQ ID NO: 21
		R9	SEQ ID NO: 20	34
6	pcDNA3.1	F10	SEQ ID NO: 22	20 SEQ ID NO: 23
		R10, identical to R4	SEQ ID NO: 11	20

The method for preparing iPSCs of the present invention reduces the number of transcription factors required for reprogramming to two, reduces the tumorigenicity, and improves the transformation efficiency through the combination of three small molecule compounds. In general, the method of the present invention is simple, efficient and easy to operate. The iPSCs prepared by this method would be more suitable for clinical translations and applications.

The above are only the preferred embodiments of the present invention. It should be pointed out that for those skilled in the art, without departing from the principles of the present invention, several improvements and modifications can be made. These improvements and modifications should also be regarded as within the protection scope of the present invention.

-continued

ccgtatatataa gtgcagtagt cgcctggaac gttctttttc gcaacgggtt tgccgccaga	240
acacaggtaa gtgccgtgtg tggttcccg cggcctggcc tctttacggg ttatggccct	300
tgcggtgcctt gaattacttc cacctggctc cagtaagtga ttcttgatcc cgagctggag	360
ccagggggcg gccttgccgt ttaggagccc ctccgctcg tgcttgagtt gaggcctggc	420
ctgggcgctg gggccgccc gtgcgaatct ggtggcacct tcgcgcctgt ctgctgctt	480
tcgataagtc tctagccatt taaaattttt gatgaacctg tgcgacgctt tttttctggc	540
aagatagtct tgtaaattcg gcccaggatc tgcacactgg tatttcggtt tttgggccc	600
cggccggcga cggggcccgt gcgtccacgc gcacatgttc ggcgaggcgg ggccctgcgag	660
cgcggccacc gagaatcgga cgggggtagt ctcaagctgg cggcctgct ctggtgctg	720
gcctcgcgcc gccgtgtatc gcccgcctt gggcggcaag gctggcccgg tcggcaccag	780
ttgcgtgagc ggaagatgg ccgcttccc gcccctgctc agggggctca aaatggagga	840
cgcggcgctc gggagagcgg gcgggtgagt caccacaca aaggaaaagg gcctttccgt	900
cctcagccgt cgcttcatgt gactccacgg agtacgggc gccgtccagg cactcgtt	960
agttctggag cttttggagt acgtcgtctt taggttggg ggaggggtt tatgcatgg	1020
agtttcccca cactgagtgg gtggagactg aagttaggcc agcttggaac ttgatgta	1080
tctcgttga atttgccctt ttgagtttg gatcttggtt cattctcaag cctcagacag	1140
tggttcaaa tttttttctt ccatttcagg tgctgtgatg gcgggacacc tg	1192

<210> SEQ ID NO 4
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: OCT4 F2 primer

<400> SEQUENCE: 4

atggcgggac acctggcttc	20
-----------------------	----

<210> SEQ ID NO 5
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: OCT4 R2 primer

<400> SEQUENCE: 5

tcagtttgaa tgcattgggag	20
------------------------	----

<210> SEQ ID NO 6
 <211> LENGTH: 1083
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: OCT4 amplicon

<400> SEQUENCE: 6

atggcgggac acctggcttc ggatttcgcc ttctcgcccc ctccaggtgg tggaggtgat	60
gggccagggg ggccggagcc gggctgggtt gatcctcgga cctggctaag cttccaaggc	120
cctcctggag ggccaggaat cgggcggggg gttggggcag gctctgaggt gtgggggatt	180
cccccatgcc ccccgccgta tgagttctgt ggggggatgg cgtactgtgg gcccaggtt	240
ggagtggggc tagtgcccca aggcggcttg gagacctctc agcctgaggg cgaagcagga	300
gtcggggtgg agagcaactc cgatggggcc tccccggagc cctgcacctg caccctggt	360

-continued

```

gccgtgaagc tggagaagga gaagctggag caaaacccgg aggagtccca ggacatcaaa 420
gctctgcaga aagaactcga gcaatttgcc aagctcctga agcagaagag gatcaccctg 480
ggatatacac aggcgatgt ggggctcacc ctgggggttc tatttgggaa ggtattcagc 540
caaacgacca tctgccgtt tgaggctctg cagcttagct tcaagaacat gtgtaagctg 600
cgcccttgc tgcagaagt ggtggaggaa gctgacaaca atgaaaatct tcaggagata 660
tgcaaagcag aaacctcgt gcaggcccg aagagaaagc gaaccagtat cgagaaccga 720
gtgagaggca acctggagaa tttgttctg cagtgccga aaccacact gcagcagatc 780
agccacatcg ccagcagct tgggctcag aaggatgtg tccgagtgtg gttctgtaac 840
cggcgccaga agggcaagcg atcaagcagc gactatgcac aacgagagga ttttgaggct 900
gctgggtctc ctttctcagg gggaccagtg tcctttctc tggcccagg gcccatctt 960
ggtacccag gctatgggag ccctcacttc actgcactgt actcctcgt ccctttcct 1020
gagggggaag cctttccccc tgtctccgtc accactctgg gctctccat gcattcaaac 1080
tga 1083

```

```

<210> SEQ ID NO 7
<211> LENGTH: 92
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NANOG F3 primer

```

```

<400> SEQUENCE: 7

```

```

atgcattcaa actgagctac taacttcagc ctgctgaagc aggctggaga cgtggaggag 60
aaccctggac ctatgagtgt ggatccagct tg 92

```

```

<210> SEQ ID NO 8
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NANOG R3 primer

```

```

<400> SEQUENCE: 8

```

```

tgcaataaac aagtttcaca cgtcttcagg ttgca 35

```

```

<210> SEQ ID NO 9
<211> LENGTH: 1005
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NANOG amplicon

```

```

<400> SEQUENCE: 9

```

```

atgcattcaa actgagctac taacttcagc ctgctgaagc aggctggaga cgtggaggag 60
aaccctggac ctatgagtgt ggatccagct tgtcccaaa gcttgcttg ctttgaagca 120
tccgactgta aagaatcttc acctatgcct gtgatttgtg ggctgaaga aaactatcca 180
tccttgcaaa tgtcttctgc tgagatgcct cacacggaga ctgtctctcc tcttcttcc 240
tccatggatc tgcttattca ggacagccct gattcttcca ccagtccca aggcaaacaa 300
cccacttctg cagagaagag tgtcgcaaaa aaggaagaca aggtcccgt caagaaacag 360
aagaccagaa ctgtgttctc ttccaccag ctgtgtgtac tcaatgatag atttcagaga 420
cagaaatacc tcagcctcca gcagatgcaa gaactctcca acatcctgaa cctcagctac 480

```

-continued

aaacaggtga agacctggtt ccagaaccag agaatgaaat ctaagaggtg gcagaaaaac	540
aactggccga agaatagcaa tgggtgtgacg cagaaggcct cagcacctac ctaccccagc	600
ctttactctt cctaccacca gggatgacct gtgaaccgga ctgggaacct tccaatgtgg	660
agcaaccaga cctggaacaa ttcaacctgg agcaaccaga cccagaacat ccagtcctgg	720
agcaaccact cctggaacac tcagacctgg tgcacccaat cctggaacaa tcaggcctgg	780
aaacagtcctt totataactg tggagaggaa tctctgcagt cctgcatgca gttccagcca	840
aattctcctg ccagtgactt ggaggtgcc ttggaagctg ctggggaagg ccttaatgta	900
atacagcaga ccactaggtta ttttagtact ccacaaacca tggatttatt cctaaactac	960
tccatgaaca tgcaacctga agacgtgtga aacttggtta ttgca	1005

<210> SEQ ID NO 10
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1 F4 primer

<400> SEQUENCE: 10

aacttggtta ttgcagctta	20
-----------------------	----

<210> SEQ ID NO 11
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1 R4 primer

<400> SEQUENCE: 11

tatagtgagt cgtattaatt	20
-----------------------	----

<210> SEQ ID NO 12
 <211> LENGTH: 3204
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1 amplicon

<400> SEQUENCE: 12

aacttggtta ttgcagctta taatgggtac aaataaagca atagcatcac aaatttcaca	60
aataaagcat ttttttcact gcattctagt tgtggtttgt ccaaactcat caatgtatct	120
tatcatgtct gtataccgtc gacctctagc tagagcttgg cgtaatcatg gtcatactgt	180
tttctgtgt gaaattgtta tccgtcaca attccacaca acatacagag cggaagcata	240
aagtgtaaag cctgggggtgc ctaatgagtg agctaaactca cattaattgc gttgcgtcga	300
ctgcccgtt tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc	360
gcggggagag gcggttttgc tattggggcg tcttcgctt cctcgtcac tgactcgtg	420
cgctcggtcg ttcgggtgcg gcgagcggtc tcagctcact caaaggcggg aatacgggta	480
tccacagaat caggggataa cgcaggaaag aacatgtgag caaaggcca gcaaaggcc	540
aggaaccgta aaaaggccgc gttgctggcg tttttccata ggctccgccc ccctgacgag	600
catcacaaaa atcgacgtc aagtcagagg tggcgaaacc cgacaggact ataaagatac	660
caggcggttc cccctggaag ctcctcgtg cgctctcctg ttcgacct gcegttacc	720
ggatacctgt ccgcctttct cccttcggga agcgtggcg tttctcatag ctcacgtgt	780
aggtatctca gttcgggtga ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc	840

-continued

gttcagcccg accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccggttaaga	900
cacgacttat cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta	960
ggcggtgcta cagagtctct gaagtggtgg cctaactacg gctacactag aagaacagta	1020
tttggatatc gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga	1080
tcgggcaaac aaaccaccgc tggtagcggg ttttttgttt gcaagcagca gattacgcgc	1140
agaaaaaag gatctcaaga agatcctttg atcttttcta cggggtctga cgctcagtgg	1200
aacgaaaact cacgttaagg gattttggtc atgagattat caaaaaggat cttcacctag	1260
atccttttaa attaaaaatg aagttttaaa tcaatctaaa gtatatatga gtaaaacttg	1320
tctgacagtt accaatgctt aatcagttag gcacctatct cagcgatctg tctatttcgt	1380
tcatccatag ttgcctgact ccccgctgtg tagataacta cgatacggga gggcttacca	1440
tctggcccca gtgctgcaat gataccgga gacccacgct caccggctcc agatttatca	1500
gcaataaacc agccagccgg aagggccgag cgcagaagtg gtcttgcaac tttatccgcc	1560
tccatccagt ctattaattg ttgccgggaa gctagagtaa gtagttcgcc agttaatagt	1620
ttgcgcaacg ttgttgccat tgctacaggc atcgtggtgt cacgctcgtc gtttggtatg	1680
gcttcattca gctcgggttc ccaacgatca aggcgagtta catgatcccc catgttggtc	1740
aaaaaagcgg ttagctcctt cggctcctcg atcgttgta gaagtaagtt ggccgcagtg	1800
ttatcactca tgggtatggc agcactgcat aattctctta ctgtcatgcc atccgtaaga	1860
tgcttttctg tgactgttga gtactcaacc aagtcattct gagaatagtg tatgcggcga	1920
cagagttgct cttgccggc gtcaatacgg gataatacgg cgccacatag cagaacttta	1980
aaagtgtca tcattggaaa acgttctctg gggcgaaaac tctcaaggat cttaccgctg	2040
ttgagatcca gttcgatgta acccactcgt gcacccaact gatcttcagc atcttttact	2100
ttcaccagcg tttctgggtg agcaaaaaca ggaaggcaaa atgccgcaaa aaaggaata	2160
agggcgacac ggaaatgttg aatactcata ctcttcttt ttcaatatta ttgaagcatt	2220
tatcagggtt attgtctcat gagcggatac atatttgaat gtatttagaa aaataaaca	2280
ataggggttc gcgcacatt tccccgaaaa gtgccacctg acgtcgacgg atcgggagat	2340
ctcccgatcc cctatggtgc actctcagta caatctgctc tgatgccgca tagttaagcc	2400
agtatctgct cctgctgtgt gtgttgaggg tcgctgagta gtgcgcgagc aaaaattaa	2460
ctacaacaag gcaaggcttg accgacaatt gcatgaagaa tctgcttagg gttaggcgtt	2520
ttgcgctgct tcgcgatgta cgggacgat atacgcgttg acattgatta ttgactagtt	2580
attaatagta atcaattacg gggtcattag ttcataagccc atatatggag ttccgcgtta	2640
cataacttac ggtaaatggc ccgcctggct gaccgcccaa cgacccccgc ccattgacgt	2700
caataatgac gtatgttccc atagtaacgc caatagggac tttccattga cgtcaatggg	2760
tggagtattt acggtaaaact gccacttg cagtacatca agtgtatcat atgccagta	2820
cgccccctat tgacgtcaat gacggtaaat ggccgcctg gcattatgcc cagtacatga	2880
ccttatggga ctttctact tggcagtaca tctacgtatt agtcacgct attaccatgg	2940
tgatgcggtt ttggcagtac atcaatgggc gtggatagcg gtttgactca cggggatttc	3000
caagtctcca cccattgac gtcaatggga gtttgtttg gcacaaaaat caacgggact	3060
ttccaaaatg tcgtaacaac tccgccccat tgacgcaaat gggcggtagg cgtgtacggt	3120
gggaggtcta tataagcaga gctctctggc taactagaga acccactgct tactggctta	3180

-continued

 tcgaaattaa tacgactcac tata 3204

<210> SEQ ID NO 13
 <211> LENGTH: 6424
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1-EF-OCT4-NANOG

<400> SEQUENCE: 13

gacggatcgg gagatctccc gatcccctat ggtgcactct cagtacaatc tgctctgatg	60
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggctcgt gagtagtgcg	120
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc	180
ttagggttag gcgttttgcg ctgcttcgcg atgtacgggc cagatatacg cgttgacatt	240
gattattgac tagttattaa tagtaataca ttacggggtc attagttcat agcccatata	300
tggagtccg cgttacataa cttacggtaa atggcccgc tggtgacccg cccaacgacc	360
cccgccatt gacgtcaata atgacgtatg tcccatagt aacgccaata gggactttcc	420
attgacgtca atgggtggag tatttacggg aaactgccca cttggcagta catcaagtgt	480
atcatatgcc aagtagcccc cctattgacg tcaatgacgg taaatggccc gcttggcatt	540
atgccagta catgacctta tgggactttc ctacttgcca gtacatctac gtattagtca	600
tcgctattac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggtttg	660
actcacgggg atttccaaagt ctccacccca ttgacgtcaa tgggagtttg ttttggcacc	720
aaaatcaacg ggactttcca aaatgtcgta acaactccgc cccattgacg caaatgggcg	780
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaacca	840
ctgcttactg gcttatcgaa attaatacga ctcaactatg gctccggtgc ccgtcagtgg	900
gcagagcgca catcgccac agtccccgag aagttggggg gaggggtcgg caattgaacc	960
ggtgcctaga gaaggtggcg cggggtaaac tgggaaagtg atgtcgtgta ctggctccgc	1020
ctttttcccg aggggtgggg agaaccgtat ataagtgcag tagtcgcgt gaacgttctt	1080
tttcgcaacg ggtttgccgc cagaacacag gtaagtgcg tgtgtggttc ccgcgggcct	1140
ggcctcttta cgggttatgg ccttgcgtg ccttgaatta cttccacctg gctccagtac	1200
gtgattcttg atccccgact ggagccaggg gcgggccttg cgctttagga gccccttcgc	1260
ctcgtgcttg agttgaggcc tggcctgggc gctggggccg ccgcgtgcga atctggtggc	1320
accttcgcgc ctgtctcgtg gctttcgata agtctctagc catttaaaat ttttgatgac	1380
ctgctgcgac gctttttttc tggcaagata gtcttgtaaa tgcggggccag gatctgcaca	1440
ctggtatttc ggtttttggg ccgcggccg gcgacggggc ccgtgcgtcc cagcgcacat	1500
gttcggcgag gcggggcctg cgagcgcggc caccgagaat cggacggggg tagtctcaag	1560
ctggccgggc tgctctggtg cctggcctcg cgcgcctgtg tatcgcctcc cctggggcgg	1620
caaggctggc ccggtcggca ccagttgctg gagcggaaaag atggccgctt cccggccctg	1680
ctccaggggg ctcaaaatgg aggacgcggc gctcgggaga gcgggcgggt gagtcaacca	1740
cacaaaggaa aagggccttt ccgtcctcag ccgtcgcttc atgtgactcc acggagtacc	1800
ggggcccgct caggcacctc gatttagttct ggagcttttg gagtacgtcg tctttagggt	1860
ggggggaggg gttttatcg atggagtctt cccacactga gtgggtggag actgaagtta	1920
ggccagcttg gcacttgatg taattctcgt tggaaattgc cctttttgag tttggatctt	1980
ggttcattct caagcctcag acagtgggtc aaagtttttt tcttcatttt caggtgtcgt	2040

-continued

gatggcggga cacctggett cggatttcgc ettcctgccc cctccagggtg gtggagggtga	2100
tgggccaggg gggccggagc cgggctgggt tgatcctcgg acctggctaa gcttccaagg	2160
ccctcctgga gggccaggaa tcgggcccgg gggtgggcca ggctctgagg tgtgggggat	2220
tcccccatgc ccccgccgt atgagttctg tggggggatg gcgtactgtg ggccccagg	2280
tggagtggg ctagtgcctc aaggcggctt ggagacctct cagcctgagg gcgaagcagg	2340
agtcggggtg gagagcaact ccgatggggc ctcgccggag cctgcaccg tcaccctcg	2400
tgccgtgaag ctggagaagg agaagctgga gcaaaaccg gaggagtccc aggacatcaa	2460
agctctgcag aaagaactcg agcaatttgc caagctcctg aagcagaaga ggatcacct	2520
gggatataca caggccgatg tggggctcac cctgggggtt ctatttggga aggtattcag	2580
ccaaacgacc atctgcgct ttgaggctct gcagcttagc ttcaagaaca tgtgtaagct	2640
gcggcccttg ctgcagaagt ggggtggagga agctgacaac aatgaaaatc ttcaggagat	2700
atgcaaagca gaaacctcg tgcaggcccg aaagagaaag cgaaccagta tcgagaaccg	2760
agtgagaggc aacctggaga attgttctt gcagtgcctg aaaccacac tgcagcagat	2820
cagccacatc gccagcagc ttgggctcga gaaggatgtg gtccgagtgt ggttctgtaa	2880
ccggcgccag aagggaagc gatcaagcag cgactatgca caacgagagg attttgaggc	2940
tgctgggtct cctttctcag ggggaccagt gtcctttctt ctggcccag ggccccattt	3000
tggtacccca ggctatggga gccctcactt cactgcactg tactcctcgg tccctttccc	3060
tgagggggaa gcctttcccc ctgtctccgt caccactctg ggctctccca tgcattcaaa	3120
ctgagctact aacttcagcc tgtgtaagca ggctggagac gtggaggaga acctggacc	3180
tatgagtgtg gatccagctt gtccccaag cttgccttgc tttgaagcat ccgactgtaa	3240
agaatcttca cctatgcctg tgatttgtgg gcctgaagaa aactatccat ccttgcaaat	3300
gtcttctgct gagatgcctc acacggagac tgtctctctt ctctctctt ccatggatct	3360
gcttattcag gacagccctg attcttcac cagtcccaaa ggcaaacac ccacttctgc	3420
agagaagagt gtcgcaaaaa aggaagacaa ggtcccgtc aagaacaga agaccagaac	3480
tgtgttctct tccaccagc tgtgtgtact caatgataga tttcagagac agaaatacct	3540
cagcctccag cagatgcaag aactctccaa catcctgaac ctgagctaca aacaggtgaa	3600
gacctgggtc cagaaccaga gaatgaaatc taagagggtg cagaaaaaca actggccgaa	3660
gaatagcaat ggtgtgacgc agaaggctc agcacctacc taccocagcc tttactctt	3720
ctaccaccag ggatgcctgg tgaaccgac tgggaacctt ccaatgtgga gcaaccagac	3780
ctggaacaat tcaacctgga gcaaccagac ccagaacatc cagtccctgga gcaaccactc	3840
ctggaacact cagacctggt gcaccaatc ctggaacaat caggcctgga acagtccctt	3900
ctataactgt ggagaggaat ctctgcagtc ctgcatgcag ttcagccaa attctcctgc	3960
cagtgacttg gaggtgcct tggaagctgc tggggaaggc cttaatgtaa tacagcagac	4020
cactagggtat tttagtactc cacaacccat ggatttattc ctaaactact ccatgaacat	4080
gcaacctgaa gacgtgtgaa acttgtttat tgcagcttat aatgggttaca aataaagcaa	4140
tagcatcaca aatttcacaa ataaagcatt tttttcactg cattctagtt gtggtttgc	4200
caaatcatc aatgtatctt atcatgtctg tataccgtcg acctctagct agagcttggc	4260
gtaatcatgg tcatagctgt ttctgtgtg aaattgttat ccgctcaca ttccacacaa	4320
catacgagcc ggaagcataa agtgtaaagc ctgggggtgc taatgagtga gctaactcac	4380

-continued

attaattgcg ttgcgctcac tgcccgcctt ccagtcggga aacctgtcgt gccagctgca	4440
ttaatgaato ggccaacgcg cggggagagg cggtttgctg attgggcgct cttccgcttc	4500
ctcgctcact gactcgctgc gctcggtcgt tcggctgcgg cgagcggat cagctcactc	4560
aaaggcggta atacggttat ccacagaatc aggggataac gcaggaaaga acatgtgagc	4620
aaaaggccag caaaaggcca ggaaccgtaa aaaggccgcg ttgctggcgt tttccatag	4680
gctccgcccc cctgacgagc atcacaaaaa tcgacgctca agtcagaggt ggcgaaaccc	4740
gacaggacta taaagatacc aggcgtttcc ccctggaagc tccctcgtgc gctctcctgt	4800
tccgaccctg ccgcttaccg gatacctgtc cgcctttctc ccttcgggaa gcgtggcgct	4860
ttctcatagc tcacgctgta ggtatctcag ttcggtgtag gtcgttcgct ccaagctggg	4920
ctgtgtgcac gaaccccccg ttcagcccgga ccgctgcgcc ttatccggta actatcgtct	4980
tgagtccaac ccggtaaagc acgacttacc gccactggca gcagccactg gtaacaggat	5040
tagcagagcg aggtatgtag gcgggtgtac agagttcttg aagtgggtggc ctaactacgg	5100
ctacactaga agaacagtat ttggtatctg cgtctcgtcg aagccagtta ccttcggaaa	5160
aagagttggt agctcttgat ccggcacaaca aaccaccgct ggtagcgggt tttttgtttg	5220
caagcagcag attacgcgca gaaaaaaagg atctcaagaa gatcctttga tctttctac	5280
ggggtctgac gctcagtgga acgaaaaactc acgttaaggg attttggtca tgagattatc	5340
aaaaaggatc ttcacctaga tctttttaa ttaaaaatga agttttaa atcaatctaaag	5400
tatatatgag taaacttggt ctgacagtta ccaatgctta atcagtgagg cacctatctc	5460
agcgatctgt ctatttcgtt catccatagt tgctgactc cccgtcgtgt agataactac	5520
gatacgggag ggcttaccat ctggccccag tgctgcaatg ataccgcgag acccacgctc	5580
accggctcca gatattatcag caataaacca gccagccgga agggccgagc gcagaagtgg	5640
tcctgcaact ttatccgct ccatccagtc tattaattgt tgccgggaag ctagagtaag	5700
tagttcgcca gttaatagtt tgcgcaacgt tgttgccatt gctacaggca tcgtggtgtc	5760
acgctcgtcg tttggtatgg cttcattcag ctccgggtcc caacgatcaa ggcgagttac	5820
atgatcccc atgttgtgca aaaaagcggg tagctccttc ggtcctccga tcgttgtcag	5880
aagtaagttg gccgcagtg taccactcat ggttatggca gcactgcata attctcttac	5940
tgatcatgcca tccgtaagat gcttttctgt gactgggtgag tactcaacca agtcattctg	6000
agaatagtgt atgcggcgac cgagttgctc ttgcccgcg tcaatacggg ataataccgc	6060
gccacatagc agaactttaa aagtgtcat cattggaaaa cgttcttcgg ggcgaaaact	6120
ctcaaggatc ttaccgctgt tgagatccag ttcgatgtaa cccactcgtg cacccaactg	6180
atcttcagca tcttttactt tcaccagcgt ttctgggtga gcaaaaacag gaaggcaaaa	6240
tgccgcaaaa aagggaataa gggcgacacg gaaatgtga atactcatac tcttcctttt	6300
tcaatattat tgaagcattt atcagggtta ttgtctcatg agcggataca tatttgaatg	6360
tatttagaaa aataaaca aa taggggttcc gcgcacattt ccccgaaaag tgccacctga	6420
cgtc	6424

<210> SEQ ID NO 14

<211> LENGTH: 92

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: NANOG R7 primer

<400> SEQUENCE: 14

-continued

```

ccctctgtca gacataggtc cagggttctc ctccacgtct ccagcctgct tcagcaggct    60
gaagttagta gctcacacgt cttcaggttg ca                                  92

```

```

<210> SEQ ID NO 15
<211> LENGTH: 1062
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: NANOG amplicon

```

```

<400> SEQUENCE: 15

```

```

atgcattcaa actgagctac taacttcagc ctgctgaagc aggctggaga cgtggaggag    60
aaccctggac ctatgagtgt ggatccagct tgtcccaaaa gcttgccctg ctttgaagca    120
tccgactgta aagaatcttc acctatgcct gtgattttgtg ggcctgaaga aaactatcca    180
tccttgcaaa tgtcttctgc tgagatgcct cacacggaga ctgtctctcc tcttccttcc    240
tccatggatc tgcttattca ggacagccct gattcttcca ccagtcctca aggcaaacaa    300
cccactttctg cagagaagag tgtcgcaaaa aaggaagaca aggtcccgtt caagaaacag    360
aagaccagaa ctgtgtttctc ttccaccag ctgtgtgtac tcaatgatag atttcagaga    420
cagaaatacc tcagcctcca gcagatgcaa gaactctcca acatcctgaa cctcagctac    480
aaacaggtga agacctggtt ccagaaccag agaatgaaat ctaagagggt gcagaaaaac    540
aactggccga agaatagcaa tgggtgtgacg cagaaggcct cagcacctac ctacccacg    600
ctttactctt cctaccacca gggatgcctg gtgaaccga ctgggaacct tccaatgtgg    660
agcaaccaga cctggaacaa ttcaacctgg agcaaccaga ccagaaacat ccagtcctgg    720
agcaaccact cctggaacac tcagacctgg tgcacccaat cctggaacaa tcaggcctgg    780
aacagtcctt tctataactg tggagaggaa tctctgcagt cctgcatgca gttccagcca    840
aattctcctg ccagtgactt ggaggctgcc ttggaagctg ctggggaagg ccttaatgta    900
atacagcaga ccactaggta ttttagtact ccacaaacca tggatttatt cctaaactac    960
tccatgaaca tgcaacctga agacgtgtga gctactaact tcagcctgct gaagcaggct   1020
ggagacgtgg aggagaaccc tggacctatg tctgacgagg gg                               1062

```

```

<210> SEQ ID NO 16
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EBNA1 F8 primer

```

```

<400> SEQUENCE: 16

```

```

atgtctgacg aggggccagg tacaggacct ggaaatgg                                38

```

```

<210> SEQ ID NO 17
<211> LENGTH: 85
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EBNA1 R8 primer

```

```

<400> SEQUENCE: 17

```

```

ttatttgtga aatttgtgat gctattgctt tatttgtaac cattataagc tgcaataaac    60
aagtttcact cctgcccttc ctcac                                           85

```

```

<210> SEQ ID NO 18

```

-continued

```

<211> LENGTH: 1991
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: EBNA1 amplicon

<400> SEQUENCE: 18
atgtctgacg aggggccagg tacaggacct ggaaatggcc taggagagaa gggagacaca      60
tctggaccag aaggctccgg cggcagtgga cctcaaagaa gagggggtga taaccatgga      120
cgaggacggg gaagaggacg aggacgagga ggcggaagac caggagcccc gggcggctca      180
ggatcagggc caagacatag agatggtgtc cggagacccc aaaaacgtcc aagttgcatt      240
ggctgcaaag ggaccocagg tggaacagga gcaggagcag gacgggaggg ggcaggagca      300
ggaggggcag gagcaggagg aggggacagga gcaggaggag gggcaggagg ggcaggaggg      360
gcaggagggg caggagcagg aggaggggca ggagcaggag gaggggcagg aggggcagga      420
ggggcaggag caggaggagg ggcaggagca ggaggagggg caggaggggc aggagcagga      480
ggaggggcag gaggggacag aggggacagga gcaggaggag gggcaggagc aggagggagg      540
gcaggagggg caggagcagg aggaggggca ggaggggcag gaggggcagg agcaggaggga      600
ggggcaggag caggaggggc aggaggggca ggaggggcag gagcaggagg ggcaggagca      660
ggagggaggg caggaggggc aggaggggca ggagcaggag gggcaggagc aggaggggca      720
ggagcaggag gggcaggagc aggaggggca ggaggggcag gagcaggagg ggcaggaggg      780
gcaggagcag gaggggcagg aggggacagga gcaggaggag gggcaggagg ggcaggagca      840
ggagggaggg caggaggggc aggagcagga ggggcaggag gggcaggagc aggaggggca      900
ggaggggcag gagcaggagg ggcaggaggg gcaggagcag gaggaggggc aggagcagga      960
ggggcaggag caggagggtg aggccggggt cgaggaggca gtggaggccg gggtcgagga      1020
ggtagtggag gccggggtcg aggaggtagt ggaggccgcc ggggtagagg acgtgaaaga      1080
gccagggggg gaagtcgtga aagagccagg gggagaggtc gtggacgtgg agaaaagagg      1140
cccaggagtc ccagtagtca gtcacatcat tccgggtctc caccgcgcag gccccctcca      1200
ggtagaaggc catttttcca ccctgtaggg gaagccgatt attttgaata ccaccaagaa      1260
ggtggcccag atggtgagcc tgacgtgccc cggggagcga tagagcaggg ccccgcatat      1320
gacccaggag aaggcccaag cactggaccc cggggtcagg gtgatggagg caggcgcaaa      1380
aaaggagggt ggtttgaaa gcatcgtggt caaggagggt ccaaccgaa atttgagaac      1440
attgcagaag gtttaagagc tctcctggct aggagtcacg tagaaaggac taccgacgaa      1500
ggaacttggg tcgccggtgt gttcgtatat ggaggtagta agacctccct ttacaacctt      1560
aggcgaggaa ctgcccttgc tattccacaa tgtcgtctta caccattgag tcgtctcccc      1620
tttggaatgg cccctggacc cggcccacaa cctggccgcg taagggagtc cattgtctgt      1680
tatttcattg tctttttaca aactcatata tttgctgagg ttttgaagga tgcgattaag      1740
gaccttggtt tgacaaagcc cgctcctacc tgcaatatca gggtgactgt gtgcagcttt      1800
gacgatggag tagatttgcc tcctcggttt ccacctatgg tggaaggggc tgcgcgggag      1860
ggtgatgacg gagatgacgg agatgaagga ggtgatggag atgaggggtga ggaagggcag      1920
gagtgaaact tgtttattgc agcttataat ggttacaaat aaagcaatag catcacaaat      1980
ttcacaaata a                                     1991

```

```

<210> SEQ ID NO 19
<211> LENGTH: 101

```


-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OriP F9 primer

<400> SEQUENCE: 19

aaatttcaca aataaagcat ttttttcact gcattctagt tgtggtttgt ccaaactcat      60
caatgtatct tatcatgtct gaacgggtag catatgcttc c                               101

<210> SEQ ID NO 20
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OriP R9 primer

<400> SEQUENCE: 20

gaggtcgacg gtataaggaa aaggacaagc agcg                                     34

<210> SEQ ID NO 21
<211> LENGTH: 1885
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: OriP amplicon

<400> SEQUENCE: 21

aaatttcaca aataaagcat ttttttcact gcattctagt tgtggtttgt ccaaactcat      60
caatgtatct tatcatgtct gaacgggtag catatgcttc ccgggtagta gtatatacta      120
tccagactaa ccctaattca atagcatatg ttacccaacg ggaagcatat gctatogaat      180
tagggtagtg aaaaggggtc taaggaaacg cgatatctcc caccatga gctgtcacgg      240
ttttatttac atggggtcag gattccacga gggtagtgaa ccattttagt cacaagggca      300
gtggctgaag atcaaggagc gggcagtgaa ctctcctgaa tcttcgctg cttcttcatt      360
ctccttcgtt tagctaatag aataactgct gagttgtgaa cagtaagggt tatgtgaggt      420
gctcgaaaac aaggtttcag gtgacgcccc cagaataaaa tttggacggg ggggttcagt      480
gtggcattgt gctatgacac caatataacc ctcacaaaacc ccttgggcaa taaatactag      540
tgtaggaatg aaacattctg aatatcttta acaatagaaa tccatggggg ggggacaagc      600
cgtaaagact ggatgtccat ctcacacgaa tttatggcta tgggcaacac ataatacctag      660
tgcaatatga tactgggggtt attaagatgt gtcccaggca gggaccaaga cagggtgaacc      720
atgttggttac actctatttg taacaagggg aaagagagtg gacgccgaca gcagcggact      780
ccactgggtg tctctaacac ccccgaaaat taaacggggc tccacgcaa tggggcccat      840
aaacaaagac aagtgggcac tctttttttt gaaattgtgg agtgggggca cgcgtcagcc      900
cccacacgcc gccctgcggt tttggactgt aaaataagggt tgtaataact tggtgattg      960
taaccccgct aaccactgcg gtcaaaccac ttgccacaaa aaccactaat ggcaccccg      1020
ggaatacctg cataagtagg tgggcggggc aagatagggg cgcgattgct gcgatctgga      1080
ggacaaatta cacacacttg cgctgagcg ccaagcacag ggttggtggg cctcatattc      1140
acgaggtcgc tgagagcacg gtgggctaata gttgccatgg gtagcatata ctacccaaat      1200
atctggatag catatgctat cctaacttat atctgggtag cataggctat cctaacttat      1260
atctgggtag catatgctat cctaacttat atctgggtag tatatgctat cctaatttat      1320
atctgggtag cataggctat cctaacttat atctgggtag catatgctat cctaacttat      1380

```

-continued

atctgggtag tatatgctat cctaactctgt atccgggtag catatgctat cctaataagag	1440
attagggttag tatatgctat cctaatttat atctgggtag catatactac ccaaatatct	1500
ggatagcata tgctatccta atctatatct gggtagcata tgctatccta atctatatct	1560
gggtagcata ggctatccta atctatatct gggtagcata tgctatccta atctatatct	1620
gggtagtata tgctatccta atttatatct gggtagcata ggctatccta atctatatct	1680
gggtagcata tgctatccta atctatatct gggtagtata tgctatccta atctgtatcc	1740
gggtagcata tgctatccta atctatatct agtcagcata tgatacccag tagtagagtg	1800
ggagtgcctat cctttgcata tgccgccacc tcccaagggg gcgtgaattt tcgctgcttg	1860
tccttttctt tataaccgtcg acctc	1885

<210> SEQ ID NO 22
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1 F10 primer

<400> SEQUENCE: 22

tataaccgtcg acctctagct	20
------------------------	----

<210> SEQ ID NO 23
 <211> LENGTH: 3073
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pcDNA3.1 amplicon

<400> SEQUENCE: 23

tataaccgtcg acctctagct agagcttggc gtaatcatgg tcatagctgt ttctgtgtg	60
aaattgttat ccgctcacia ttccacacia catacgagcc ggaagcataa agtgtaaagc	120
ctgggggtgc taatgagtga gctaactcac attaatgctg ttgcgctcac tgcccgttt	180
ccagtcggga aacctgtcgt gccagctgca ttaatgaatc ggccaacgcg cggggagagg	240
cggtttgcgt attgggcgct ctccgcttc ctgcgctact gactcgctgc gctcggtcgt	300
tcggctgcgg cgagcggtat cagctcactc aaaggcggtg atacggttat ccacagaatc	360
aggggataac gcaggaaaga acatgtgagc aaaaggccag caaaaggcca ggaaccgtaa	420
aaaggccgcg ttgctggcgt ttttccatag gctccgcccc cctgacgagc atcacaaaaa	480
tcgacgctca agtcagaggt ggcgaaaccc gacaggacta taaagatacc aggcgtttcc	540
ccctggaagc tcctcgctgc gctctcctgt tccgacctg ccgcttaccg gatacctgtc	600
cgcctttctc ccttcgggaa gcgtggcgct ttctcatagc tcacgctgta ggtatctcag	660
ttcggtgtag gtcgttcgct ccaagctggg ctgtgtgcac gaaccccccg ttcagcccga	720
ccgctgcgcc ttatccggta actatcgtct tgagtccaac ccggtgaagc acgacttacc	780
gccactggca gcagccactg gtaacaggat tagcagagcg aggtatgtag gcggtgctac	840
agagttcttg aagtgggtgc ctaactacgg ctacactaga agaacagtat ttggtatctg	900
cgtctgctg aagccagtta ccttcggaaa aagagttggt agctcttgat ccggcaaaca	960
aaccaccgct ggtagcgggt tttttgttg caagcagcag attacgcgca gaaaaaagg	1020
atctcaagaa gatcctttga tctttctac ggggtctgac gctcagtga acgaaaactc	1080
acgttaaggg attttgtgca tgagattatc aaaaaggatc ttcacctaga tcctttttaa	1140
ttaaaaatga agtttttaaat caatctaaag tatatatgag taaacttggt ctgacagtta	1200

-continued

```

ccaatgctta atcagtgagg cacctatctc agcgatctgt ctatttcggt catccatagt 1260
tgctgactc cccgtcgtgt agataactac gatacgggag ggcttaccat ctggccccag 1320
tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag caataaacca 1380
gccagccgga agggccgagc gcagaagtgg tcttgcaact ttatccgect ccaccagtc 1440
tattaattgt tgccgggaag ctagagtaag tagttcgcca gttaatagtt tgcgcaacgt 1500
tgttgccatt gctacaggca tcgtggtgtc acgctcgtcg tttggtagtg cttcattcag 1560
ctccgggttc caacgatcaa ggcgagttac atgatcccc atgttggtga aaaaagcgg 1620
tagctccttc ggtcctccga tcgttgtagc aagtaagttg gccgcagtgt taccactcat 1680
ggttatggca gcactgcata attctcttac tgtcatgcca tccgtaagat gcttttctgt 1740
gactggtagg tactcaacca agtcattctg agaatagtg atgcggcgac cgagttgctc 1800
ttgcccggcg tcaatacggg ataataccgc gccacatagc agaactttaa aagtgtcat 1860
cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc ttaccgctgt tgagatccag 1920
ttcgatgtaa cccactcgtg caccacaactg atcttcagca tcttttactt tcaccagcgt 1980
ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa aagggaataa gggcgacacg 2040
gaaatgttga atactcctac tcttcctttt tcaatattat tgaagcattt atcaggggta 2100
ttgtctcatg agcggatata tatttgaatg tatttagaaa aataacaaa taggggttcc 2160
gcgcacattt ccccgaaaag tgccacctga cgtcgacgga tcgggagatc tcccgatccc 2220
ctatggtagc ctctcagtag aatctgctct gatgcgcgat agttaagcca gtatctgctc 2280
cctgcttggt tggtggaggt cgctgagtag tgccgagca aaatttaagc tacaacaagg 2340
caagcettga ccgacaattg catgaagaat ctgcttaggg ttaggcgttt tgcgctgctt 2400
cgcatgtac gggccagata tacgcgttga cattgattat tgactagtta ttaatagtaa 2460
tcaattacgg ggtcattagt tcatagccca tatatggagt tccgcgttac ataacttacg 2520
gtaaatggcc cgcttggtg accgccaac gacccccgcc cattgacgct aataatgacg 2580
tatgttccca tagtaacgcc aatagggact ttccattgac gtcaatgggt ggagtattta 2640
cggtaaaactg cccacttggc agtacatcaa gtgtatcata tgccaagtac gccccctatt 2700
gacgtcaatg acggtaaatg gccgcctgg cattatgccc agtacatgac cttatgggac 2760
tttctactt ggcagtacat ctacgtatta gtcacgcta ttaccatggt gatgcgggtt 2820
tggcagtaca tcaatgggag tggatagcgg tttgactcac ggggatttcc aagtctccac 2880
cccattgacg tcaatgggag tttgttttg caccaaaac aacgggactt tccaaaatgt 2940
cgtaacaact ccgccccatt gacgcaaatg ggcggtaggc gtgtacgggt ggaggtctat 3000
ataagcagag ctctctggct aactagagaa cccactgctt actggcttat cgaaattaat 3060
acgactcact ata 3073

```

<210> SEQ ID NO 24

<211> LENGTH: 10196

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pcDNA3.1-EF-OCT4-NANOG-EBNA1-OriP

<400> SEQUENCE: 24

```

gacggatcgg gagatctccc gatcccctat ggtgcactct cagtacaatc tgctctgatg 60
ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggctcgt gagtagtgcg 120

```

-continued

cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc	180
ttagggtttag gcgttttgcg ctgcttcgcg atgtacgggc cagatatagc cggtgacatt	240
gattattgac tagttattaa tagtaataa ttacggggtc attagttcat agcccatata	300
tggagtccg cggtacataa cttacggtaa atggcccgcc tggctgaccg cccaacgacc	360
ccgcccatt gacgtcaata atgacgtatg ttcccatagt aacgccaata gggactttcc	420
attgacgtca atgggtggag tatttacggt aaactgccca cttggcagta catcaagtgt	480
atcatatgcc aagtacgccc cctattgacg tcaatgacgg taaatggccc gcctggcatt	540
atgccagta catgacctta tgggactttc ctacttgga gtacatctac gtattagtca	600
tcgtattac catggtgatg cggttttggc agtacatcaa tgggcgtgga tagcggttg	660
actcacggg atttccaagt ctccaccca ttgacgtcaa tgggagtttg ttttggcacc	720
aaaatcaacg ggactttcca aaatgtcgta acaactccg cccattgacg caaatgggcg	780
gtaggcgtgt acggtgggag gtctatataa gcagagctct ctggctaact agagaacca	840
ctgcttactg gcttatcgaa attaatacga ctactatag gctccggtgc ccgtcagtgg	900
gcagagcgca catcgccac agtcccag agttggggg gaggggtcgg caattgaacc	960
ggtgcctaga gaagggtggc cggggtaaac tgggaaagt atgtcgtgta ctggctccgc	1020
ctttttccg aggggtgggg agaaccgat ataagtgcag tagtcgccgt gaacgttctt	1080
tttcgcaacg ggtttgccgc cagaacacag gtaagtgcg tgtgtggttc ccgcgggcct	1140
ggcctcttta cgggttatgg ccttgcgtg cettgaatta cttccacctg gctccagtac	1200
gtgattcttg atcccgagct ggagccagg gcgggccttg cgcttagga gcccttcgc	1260
ctcgtgcttg agttgagcc tggcctgggc gctggggccg ccgcgtgcga atctggtggc	1320
accttcgcgc ctgtctcgt gctttcgata agtctctagc catttaaaat ttttgatgac	1380
ctgctgcgac gctttttttc tggcaagata gtcttgtaaa tgcgggccag gatctgcaca	1440
ctggtatttc ggtttttggg ccgcggccg gcgacggggc ccgtgcgtcc cagcgacat	1500
gttcggcgag gcggggcctg cgagcgcggc caccgagaat cggacggggg tagtctcaag	1560
ctggccggcc tgccttggtg cctggcctcg ccgcgcctg tatcgcccc ccctgggcgg	1620
caagctggc ccggtcgca ccagtgcgt gagcggaag atggccgctt ccgggccctg	1680
ctccaggggg ctcaaatgg aggacgcgc gcctcggaaga gcgggcgggt gattcaccca	1740
cacaaaggaa aagggccttt ccgtcctcag ccgtcgcttc atgtgactcc acggagtacc	1800
gggcgcctc caggcacctc gatttagttc ggagcttttg gactacgtcg tctttaggtt	1860
ggggggaggg gttttatgag atggagtctt ccacactga gtgggtggag actgaagtta	1920
ggccagcttg gcacttgatg taattctcgt tggaaattgc ctttttgag tttgatctt	1980
ggttcattct caagcctcag acagtgggtt aaagtttttt tcttcattt cagggtgcgt	2040
gatggcggga cacctggctt cggatttcgc cttctcgccc cctccagggt gtggagggtga	2100
tgggccaggg gggccggagc cgggctgggt tgatcctcgg acctggctaa gcttccaagg	2160
ccctcctgga gggccaggaa tcgggcccgg ggttgggcca ggctctgagg tgtgggggat	2220
tcccccatgc ccccgccgt atgagttctg tgggggatg gcgtactgtg ggccccaggt	2280
tggagtggg ctagtcccc aaggcggctt ggagacctct cagcctgagg gcgaagcagg	2340
agtcggggtg gagagcaact ccgatgggc ctcgccggag ccctgcaccg tcaccttg	2400
tgccgtgaag ctggagaagg agaagctgga gcaaaacccg gaggagctcc aggacatcaa	2460
agctctgcag aaagaactcg agcaatttgc caagctcctg aagcagaaga ggatcacct	2520

-continued

gggatataca caggccgatg tggggctcac cctgggggtt ctatttggga aggtattcag	2580
ccaaacgacc atctgccgct ttgaggtctt gcagcttagc ttcaagaaca tgtgtaagct	2640
gcggcccttg ctgcagaagt ggggtggagga agctgacaac aatgaaaac ttcaggagat	2700
atgcaaagca gaaacctcg tgcaggcccg aaagagaaag cgaaccagta tcgagaaccg	2760
agtgagaggc aacctggaga atttgttctt gcagtgcccg aaaccacac tgcagcagat	2820
cagccacatc gccacgagc ttgggctcga gaaggatgtg gtccgagtgt ggttctgtaa	2880
ccggcgccag aagggcaagc gatcaagcag cgactatgca caacgagagg attttgaggc	2940
tgttgggtct cctttctcag ggggaccagt gtctttctt ctggccccag ggccccattt	3000
tggtagccca ggctatggga gcctcactt cactgcactg tactcctcgg tccctttccc	3060
tgagggggaa gcctttcccc ctgtctccgt caccactctg ggctctccca tgcattcaaa	3120
ctgagctact aacttcagcc tgtgaagca ggctggagac gtggaggaga accctggacc	3180
tatgagtgtg gatccagctt gtcccaaaag cttgccttgc tttgaagcat ccgactgtaa	3240
agaatcttca cctatgcctg tgatttgtgg gcctgaagaa aactatccat ccttgcaaat	3300
gtcttctgct gagatgcctc acacggagac tgtctctct cttccttctt ccatggatct	3360
gcttattcag gacagccctg attcttccac cagtcccaaa ggcaacaac ccacttctgc	3420
agagaagagt gtcgcaaaaa aggaagacaa ggtcccggtc aagaaacaga agaccagaac	3480
tgtgttctct tccaccagc tgtgtgtact caatgataga ttccagagac agaaatacct	3540
cagcctccag cagatgcaag aactctccaa catcctgaac ctcagctaca aacaggtgaa	3600
gacctggttc cagaaccaga gaatgaaac taagaggtgg cagaaaaaca actggccgaa	3660
gaatagcaat ggtgtgacgc agaaggctc agcacctacc taccacagc tttactctt	3720
ctaccaccag ggatgcctgg tgaaccgac tgggaacctt ccaatgtgga gcaaccagac	3780
ctggaacaat tcaacctgga gcaaccagac ccagaacatc cagtctgga gcaaccactc	3840
ctggaacact cagacctggt gcaccaatc ctggaacaat caggcctgga acagtccctt	3900
ctataactgt ggagaggaat cctgcagtc ctgcatgcag ttccagccaa attctcctgc	3960
cagtgacttg gaggtgcct tggaaactgc tggggaaggc cttaatgtaa tacagcagac	4020
cactaggtat tttagtact cacaaccat ggatttatt ctaaactact ccatgaacat	4080
gcaacctgaa gacgtgtgag ctactaactt cagcctgctg aagcaggctg gagacgtgga	4140
ggagaacctt ggacctatgt ctgacgagg gccaggtaga ggacctggaa atggcctagg	4200
agagaaggga gacacatctg gaccagaagg ctccggcggc agtggacctc aaagaagagg	4260
gggtgataac catggacgag gacggggaag aggacgagga cgaggaggcg gaagaccagg	4320
agccccgggc ggctcaggat caggggccaag acatagagat ggtgtccgga gacccccaaa	4380
acgtccaagt tgcattggct gcaaaggac ccacggtgga acaggagcag gagcaggagc	4440
gggaggggca ggagcaggag gggcaggagc aggaggaggg gcaggagcag gaggaggggc	4500
aggaggggca ggaggggag gaggggcagg agcaggagga ggggcaggag caggagggag	4560
ggcaggaggg gcaggaggag gggcaggagc aggaggggca ggagcaggag gaggggcagg	4620
aggggcagga gcaggaggag gggcaggagc ggcaggaggg gcaggagcag gaggaggggc	4680
aggagcagga ggaggggcag gaggggcagg agcaggagga ggggcaggag gggcaggagc	4740
ggcaggagca ggaggaggag caggagcagg aggggcagga ggggcaggag gggcaggagc	4800
aggaggggca ggagcaggag gaggggcagg aggggcagga ggggcaggag caggaggggc	4860

-continued

aggagcagga ggggcaggag caggaggggc aggagcagga ggggcaggag gggcaggagc	4920
aggaggggca ggaggggag gagcaggag ggagggagg gcaggagcag gaggaggggc	4980
aggaggggca ggagcaggag gaggggcagg aggggcagga gcaggagggg caggaggggc	5040
aggagcagga ggggcaggag gggcaggagc aggaggggca ggaggggag gagcaggag	5100
aggggcagga gcaggagggg caggagcagg aggtggaggc cggggtcgag gaggcagtgg	5160
agggcggggg cgaggaggta gtggaggccg gggtcgagga ggtagtggag gccgcgggg	5220
tagaggacgt gaaagagcca gggggggaag tcgtgaaaga gccaggggga gaggtcgtgg	5280
acgtggagaa aagaggccca ggagtcagg tagtcagta tcatcatccg ggtctccacc	5340
gcgcaggccc cctccaggta gaaggccatt tttccaccct gtaggggaag ccgattattt	5400
tgaataccac caagaagggt gccagatgg tgagcctgac gtgccccgg gagcgataga	5460
gcaggggccc gcagatgacc caggagaagg cccaagcact ggaccccggt gtcagggtga	5520
tggaggcagg cgcaaaaaag gaggggtggt tggaaagcat cgtgggtcaag gaggttccaa	5580
cccgaattt gagaacattg cagaagggtt aagagctctc ctggctagga gtcacgtaga	5640
aaggactacc gacgaaggaa cttgggtcgc cgggtgtgtc gtatatggag gtagtaagac	5700
ctccctttac aacctaggc gaggaactgc ccttgctatt ccacaatgc gtcttacacc	5760
attgagtcgt ctccctttg gaatggccc tggacccggc ccacaacctg gcccgctaag	5820
ggagtcatt gtctgttatt tcattgtctt ttacaaaact catatatgtg ctgaggtttt	5880
gaaggatgag attaaggacc ttgttatgac aaagcccgt cctacctgca atatcagggt	5940
gactgtgtgc agctttgagc atggagtaga ttgcctccc tggtttccac ctatggtgga	6000
aggggctgcc gcggagggtg atgacggaga tgacggagat gaaggagggt atggagatga	6060
gggtgaggaa gggcaggagt gaaactgtt tattgcagct tataatggtt acaataaag	6120
caatagcatc acaaatttca caaataaagc atttttttca ctgcattcta gttgtggttt	6180
gtccaaaact atcaatgtat cttatcatgt ctgaacgggt agcatatgct tcccggtag	6240
tagtatatac tatccagact aaccttaatt caatagcata tggtacccaa cgggaagcat	6300
atgctatcga attagggtta gtaaaagggt cctaaggaa agcgatatct cccaccccat	6360
gagctgtcac ggttttattt acatgggtc aggattccac gagggtagtg aaccatttta	6420
gtcacaaggg cagtggctga agatcaagga gcgggcagtg aactctcctg aatcttcgcc	6480
tgcttcttca ttctccttcg tttagctaag agaataactg ctgagttgtg aacagtaagg	6540
tgtatgtgag gtgctcgaaa acaaggttc aggtgacgcc ccagaataa aatttgagc	6600
gggggttcag tgggtgcatt gtgctatgac accaatataa ccttcacaaa cccctgggc	6660
aataaatact agtgtaggaa tgaaacattc tgaatatctt taacaataga aatccatggg	6720
gtggggacaa gccgtaaaga ctggatgtcc atctcacacg aatttatggc tatgggcaac	6780
acataatcct agtgcaatat gatactgggg ttattaagat gtgtcccagg cagggaacca	6840
gacaggtgaa ccatgttgtt aactctatt tgtaacaagg ggaaagagag tggacgccga	6900
cagcagcgga ctccactggt tgtctctaac acccccgaat attaaacggg gctccacgcc	6960
aatggggccc ataaacaaag acaagtggc actctttttt ttgaaattgt ggagtggggg	7020
cacgcgtcag ccccccacag ccgcccctgc gttttggact gtaaaataag ggtgtaataa	7080
cttggtgat tgtaaccccg ctaaccactg cggtaaac accctgcccac aaaaccacta	7140
atggcaccac ggggaatacc tgcataagta ggtggggggg ccaagatagg ggcgcgattg	7200
ctgcgatctg gaggacaaat tacacacact tgcgcctgag cgccaagcac aggggtgttg	7260

-continued

gtctcatat tcacgaggtc gctgagagca cggtaggcta atgttgccat gggtagcata	7320
tactacccaa atatctggat agcatatgct atcctaactc atatctgggt agcataggct	7380
atcctaactc atatctgggt agcatatgct atcctaactc atatctgggt agtatatgct	7440
atcctaattt atatctgggt agcataggct atcctaactc atatctgggt agcatatgct	7500
atcctaactc atatctgggt agtatatgct atcctaactc gtatccgggt agcatatgct	7560
atcctaatag agattagggt agtatatgct atcctaattt atatctgggt agcatatact	7620
acccaaatat ctggatagca tatgctatcc taatctatat ctgggtagca tatgctatcc	7680
taatctatat ctgggtagca taggctatcc taatctatat ctgggtagca tatgctatcc	7740
taatctatat ctgggtagta tatgctatcc taatttatat ctgggtagca taggctatcc	7800
taatctatat ctgggtagca tatgctatcc taatctatat ctgggtagta tatgctatcc	7860
taatctgtat ccgggtagca tatgctatcc tcatgcatat acagtcagca tatgataccc	7920
agtagtagag tgggagtgct atcctttgca tatgccgcca cctcccaagg gggcgtgaat	7980
tttcgtgct tgtccttttc cttataccgt cgacctctag ctagagcttg gcgtaatcat	8040
ggctcatagct gtttctgtg tgaaattgtt atccgctcac aattccacac aacatacgag	8100
ccggaagcat aaagtgtaaa gccctgggtg cctaatagag gagctaactc acattaattg	8160
cgttgcgctc actgccgct tccagtcgg gaaacctgtc gtgccagctg cattaatgaa	8220
tcggccaacg cgcggggaga ggcgggttgc gtattggcg ctcttcgct tcctcgtca	8280
ctgactcgct gcgctcggtc gttcggctgc ggcgagcgg atcagctcac tcaaaggcgg	8340
taatacgggt atccacagaa tcaggggata acgcaggaaa gaacatgtga gcaaaaggcc	8400
agcaaaaggc caggaaccgt aaaaaggcgg cgttgctggc gtttttccat aggctccgcc	8460
ccccgacga gcatcacaaa aatcgacgct caagtcagag gtggcgaaac ccgacaggac	8520
tataaagata ccaggcggtt cccctggaa gctccctcgt gcgctctcct gttccgaccc	8580
tgccgcttac cggataacctg tccgccttc tcccttcggg aagcgtggcg ctttctcata	8640
gctcacgctg taggtatctc agttcgggtg aggtcggtcg ctccaagctg ggctgtgtgc	8700
acgaaccccc cgttcagccc gaccgctgcg ccttatccgg taactatcgt cttgagtcca	8760
acccggtaag acacgactta tcgccactgg cagcagccac tggtaacagg attagcagag	8820
cgaggatatg aggcgggtct acagagttct tgaagtgggt gcctaactac ggctacacta	8880
gaagaacagt atttggatc tgcgctctgc tgaagccagt taccttcgga aaaagagttg	8940
gtagctcttg atccggcaaa caaaccaccg ctggtagcgg tttttttgtt tgcaagcagc	9000
agattacgcg cagaaaaaaa ggatctcaag aagatccttt gatcttttct acggggtctg	9060
acgctcagtg gaacgaaaac tcacgttaag ggattttggt catgagatta tcaaaaagga	9120
tcttcaccta gatcctttta aattaaat gaagttttaa atcaatctaa agtatatatg	9180
agtaaacttg gtctgacagt taccaatgct taatcagtg ggcacctatc tcagcgatct	9240
gtctatcttg ttcattcata gttgcctgac tccccgtcgt gtagataact acgatacggg	9300
agggettacc atctggcccc agtgcgcaa tgataccgag agacccacgc tcaccggctc	9360
cagatttata agcaataaac cagccagccg gaagggccga gcgcagaagt ggtcctgcaa	9420
ctttatccgc ctccatccag tctattaatt gttgcggga agctagagta agtagttcgc	9480
cagttaatag tttgcgcaac gttgttgcca ttgtacagg catcggtggg tcacgctcgt	9540
cgtttggat ggcttcattc agctccggtt cccaacgata aaggcgagtt acatgatccc	9600

-continued

```

ccatgttggtg caaaaaagcg gttagctcct tcggtcctcc gatcgttgtc agaagtaagt 9660
tggccgcagt gttatcactc atgggttatgg cagcactgca taattctctt actgtcatgc 9720
catccgtaag atgcttttct gtgactgggtg agtactcaac caagtcattc tgagaatagt 9780
gtatgcggcg accgagttgc tcttgcccgg cgtcaatacg ggataatacc gcgccacata 9840
gcagaacttt aaaagtgtc atcattggaa aacgtttctc ggggcgaaaa ctctcaagga 9900
tcttaccgct gttgagatcc agttcgatgt aaccactcg tgcaccaac tgatcttcag 9960
catcttttac tttcaccagc gtttctgggt gagcaaaaac aggaaggcaa aatgccgcaa 10020
aaaagggaaat aagggcgaca cggaaatgtt gaatactcat actcttcctt tttcaatatt 10080
attgaagcat ttatcagggt tattgtctca tgagcggata catatttgaa tgtatttaga 10140
aaaataaaca aataggggtt ccgcgcacat ttccccgaaa agtgccacct gacgtc 10196

```

The invention claimed is:

1. A method of generating mammalian induced pluripotent stem cells (iPSCs), the method comprising:

(1) introducing one or more nucleic acids encoding the reprogramming-inducing factors Oct4 and Nanog into mammalian fibroblasts or mammalian erythroid progenitor cells to obtain reprogrammed fibroblasts or erythroid progenitor cells;

(2) culturing the reprogrammed fibroblasts or erythroid progenitor cells obtained in step (1) in the presence of chemical inducing agents to obtain induced pluripotent stem cells (iPSCs), wherein the chemical inducing agents are consisting of a TGFB receptor inhibitor, a cyclic AMP (cAMP) agonist, and a glycogen synthase kinase (GSK) inhibitor,

wherein only Oct4 and Nanog are used as reprogramming-inducing factors.

2. The method according to claim 1, wherein the reprogramming-inducing factors Oct4 and Nanog are introduced in the mammalian fibroblasts or mammalian erythroid progenitor cells in the form of DNA thereof or in the form of mRNA thereof.

3. The method according to claim 1, wherein the TGFB receptor inhibitor is 616452, and/or the cyclic AMP (cAMP) agonist is Forskolin, and/or the glycogen synthase kinase (GSK) inhibitor is TD114-2.

4. The method according to claim 3, wherein the working concentration of the TGFB receptor inhibitor 616452 is 0.1-20 μ M; and/or the working concentration of the cyclic AMP (cAMP) agonist Forskolin is 0.1-50 μ M; and/or the working concentration of the glycogen synthase kinase (GSK) inhibitor TD114-2 is 0.1-20 μ M.

5. The method according to claim 1, wherein the mammalian fibroblast or mammalian erythroid progenitor cell is isolated from skin cells, blood cells, urine cells, liver cells, epithelial cells, or gastric cells.

6. The method according to claim 1, wherein the mammalian fibroblasts or mammalian erythroid progenitor cells are isolated from human.

7. The method according to claim 1, wherein in step (1), the introducing one or more nucleic acids encoding the reprogramming-inducing factors Oct4 and Nanog into the mammalian fibroblasts or mammalian erythroid progenitor cells comprises introducing:

a) a single recombinant vector comprising both nucleotide sequences encoding the reprogramming-inducing factors Oct4 and Nanog, or

b) two recombinant vectors, wherein one recombinant vector comprises a nucleic acid sequence encoding Oct4 and the other recombinant vector comprises a nucleic acid sequence encoding Nanog.

8. The method according to claim 1, wherein in step (1), the introducing one or more nucleic acids encoding the reprogramming-inducing factors Oct4 and Nanog is performed by infecting the mammalian fibroblasts or mammalian erythroid progenitor cells with:

a) a single recombinant Sendai virus comprising a nucleotide sequence encoding Oct4 and a nucleotide sequence encoding Nanog, or

b) two recombinant Sendai viruses, wherein one virus comprises the nucleotide sequence encoding Oct4 and the other virus comprises the nucleotide sequence encoding Nanog.

9. The method according to claim 8, wherein at least one of the nucleotide sequence encoding Oct4 and the nucleotide sequence encoding Nanog is RNA.

10. The method according to claim 4, wherein the working concentration of 616452 is 5-10 μ M.

11. The method according to claim 4, wherein the working concentration of 616452 is 5 μ M.

12. The method according to claim 4, wherein the working concentration of Forskolin is 2-20 μ M.

13. The method according to claim 4, wherein the working concentration of Forskolin is 10 μ M.

14. The method according to claim 4, wherein the working concentration of TD114-2 is 2-10 μ M.

15. The method according to claim 4, wherein the working concentration of TD114-2 is 5 μ M.

* * * * *