



US012398191B2

(12) **United States Patent**
Veatch et al.

(10) **Patent No.: US 12,398,191 B2**

(45) **Date of Patent: Aug. 26, 2025**

(54) **BRAF-SPECIFIC TCRS AND USES THEREOF**

(56) **References Cited**

(71) Applicants: **FRED HUTCHINSON CANCER CENTER**, Seattle, WA (US);
UNIVERSITY OF WASHINGTON,
Seattle, WA (US)

U.S. PATENT DOCUMENTS

(72) Inventors: **Joshua Veatch**, Seattle, WA (US);
Stanley R. Riddell, Sammamish, WA
(US); **Sylvia Lee**, Seattle, WA (US)

5,283,173	A	2/1994	Fields et al.
5,420,032	A	5/1995	Marshall et al.
5,468,614	A	11/1995	Fields et al.
6,410,319	B1	6/2002	Raubitschek et al.
6,833,252	B1	12/2004	Dujon et al.
7,446,191	B2	11/2008	Jensen
7,514,537	B2	4/2009	Jensen
8,697,359	B1	4/2014	Zhang
8,822,647	B2	9/2014	Jensen
9,574,000	B2	2/2017	Langermann et al.
2004/0002092	A1	1/2004	Arnould et al.
2004/0087025	A1	5/2004	June et al.
2006/0078552	A1	4/2006	Arnould et al.
2006/0153826	A1	7/2006	Arnould et al.
2006/0206949	A1	9/2006	Arnould et al.
2007/0117128	A1	5/2007	Smith et al.
2010/0065818	A1	3/2010	Kim et al.
2011/0189141	A1	8/2011	Kieback et al.
2011/0243972	A1	10/2011	Jaffee
2011/0301073	A1	12/2011	Gregory et al.
2014/0068797	A1	3/2014	Doudna et al.
2014/0186843	A1	7/2014	Zhang et al.
2015/0337369	A1	11/2015	Davis et al.

(73) Assignees: **FRED HUTCHINSON CANCER CENTER**, Seattle, WA (US);
UNIVERSITY OF WASHINGTON,
Seattle, WA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 1344 days.

(21) Appl. No.: **16/638,339**

(22) PCT Filed: **Aug. 10, 2018**

(86) PCT No.: **PCT/US2018/046350**

§ 371 (c)(1),

(2) Date: **Feb. 11, 2020**

(87) PCT Pub. No.: **WO2019/033057**

PCT Pub. Date: **Feb. 14, 2019**

FOREIGN PATENT DOCUMENTS

WO	WO 9709433	A1	3/1997
WO	2007/002811	A2	1/2007
WO	WO 2013025779	A1	2/2013
WO	WO 2014031687	A1	2/2014
WO	WO 2015071474	A2	5/2015

(Continued)

(65) **Prior Publication Data**

US 2020/0223899 A1 Jul. 16, 2020

US 2021/0238250 A9 Aug. 5, 2021

Related U.S. Application Data

(60) Provisional application No. 62/544,695, filed on Aug. 11, 2017.

(51) **Int. Cl.**

A61K 39/395 (2006.01)

A61K 35/17 (2015.01)

A61K 40/11 (2025.01)

A61K 40/32 (2025.01)

A61K 40/42 (2025.01)

A61P 35/00 (2006.01)

C07K 14/725 (2006.01)

C12N 5/0783 (2010.01)

C12N 7/00 (2006.01)

C12N 15/86 (2006.01)

A61K 38/00 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 14/7051** (2013.01); **A61K 35/17** (2013.01); **A61K 40/11** (2025.01); **A61K 40/32** (2025.01); **A61K 40/4251** (2025.01); **A61P 35/00** (2018.01); **C12N 5/0636** (2013.01); **C12N 7/00** (2013.01); **C12N 15/86** (2013.01); **A61K 38/00** (2013.01); **A61K 2239/57** (2023.05); **C12N 2740/10043** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

OTHER PUBLICATIONS

Dunn et al., "Directed evolution of human T cell receptor CDR2 residues by phage display dramatically enhances affinity for cognate peptide-MHC without increasing apparent cross-reactivity," *Protein Science* 15:710-721, Jan. 10, 2006.

Lee et al., "Current management and novel agents for malignant melanoma," *Journal of Hematology & Oncology* 5(3):1-7, 2012.

Legut et al., "CRISPR-mediated TCR replacement generates superior anticancer transgenic T cells," *blood* 131(3):311-322, Jan. 18, 2018.

Sharkey et al., "CD4+ T-Cell Recognition of Mutated B-RAF in Melanoma Patients Harboring the V599E Mutation," *Cancer Research* 64(5):1595-1599, Mar. 2004.

(Continued)

Primary Examiner — Yunsoo Kim

(74) *Attorney, Agent, or Firm* — Seed IP Law Group LLP

(57) **ABSTRACT**

The present disclosure provides binding proteins, such as TCRs, that specifically bind various tumor associated antigens (including human BRAFV600E epitope), cells expressing such antigen specific binding proteins, nucleic acids encoding the same, and compositions for use in treating diseases or disorders in which cells express BRAFV600E, such as in cancer.

16 Claims, 20 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	2015/095895	A1	6/2015
WO	WO 2016040724	A1	3/2016
WO	WO 2016054638	A1	4/2016
WO	2016/069647	A1	5/2016
WO	WO 2016134333	A1	8/2016
WO	2016/161273	A1	10/2016
WO	WO 2017021526	A1	2/2017
WO	2017/096331	A1	6/2017

OTHER PUBLICATIONS

Veatch et al., "Tumor-infiltrating BRAF^{V600E}-specific CD4⁺ T cells correlated with complete clinical response in melanoma," *The Journal of Clinic Investigation* 128(4):1563-1568, Apr. 2018.

Walseng et al., "A TCR-based Chimeric Antigen Receptor," *Scientific Reports* 7:1-10, Sep. 2017.

GenBank, "T-cell receptor alpha chain, partial [*Homo sapiens*]," Accession No. BAN84546.1, Nov. 21, 2013. (3 pages).

Kobayashi et al., "A new cloning and expression system yields and validates TCRs from blood lymphocytes of patients with cancer within 10 days," *Nature Medicine* 19(11), Nov. 2013. (30 pages).

Abbas et al., "Cellular and Molecular Immunology," Third Edition, pp. 149, 155, 250, 266, 413, 1997. (7 pages).

Altschul et al., "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs," *Nucleic Acids Research* 25(17):3389-3402, Jul. 16, 1997.

Argast et al., "I-Ppol and I-Crel Homing Site Sequence Degeneracy Determined by Random Mutagenesis and Sequential in Vitro Enrichment," *J. Mol. Biol.* 280:345-353, 1998.

Ashworth et al., "Computational redesign of endonuclease DNA binding and cleavage specificity," *Nature* 441(7093):656-659, Jun. 1, 2006 (NIH Public Access Author Manuscript, available in PMC Dec. 9, 2010) (11 pages).

Belfort et al., "Homing endonucleases: keeping the house in order," *Nucleic Acids Research* 25(17):3379-3388, 1997.

Brentjens et al., "Genetically Targeted T Cells Eradicates Systemic Acute Lymphoblastic Leukemia Xenografts," *Clin Cancer Res* 13(18):5426-5435, Sep. 15, 2007.

Cavallo et al., "2011: the immune hallmarks of cancer," *Cancer Immunol Immunother* 60:319-326, Jan. 26, 2011.

Chapuis et al., "Abstract LB-136: IL-21-derived melanoma-reactive CTL combined with anti-CTLA4 persist, acquire central memory characteristics, and mediate tumor regression in patients with metastatic melanoma," *Cancer Research* 72(8 Suppl):2012. (4 pages).

Chapuis et al., "Transferred melanoma-specific CD8⁺ T cells persist, mediate tumor regression, and acquire central memory phenotype," *PNAS* 109(12):4592-4597, Mar. 20, 2012.

Chapuis et al., "Transferred WT1-reactive CD8⁺ T cells can mediate antileukemic activity and persist in post-transplant patients," *Sci Transl Med* 5(174):174ra27, Feb. 27, 2013 (NIH Public Access Author manuscript, available in PMC Jun. 11, 2013) (25 pages).

Chevalier et al., "Design, Activity, and Structure of a Highly Specific Artificial Endonuclease," *Molecular Cell* 10:895-905, Oct. 2002.

Chothia, et al., "The outline structure of the T-cell ab receptor," *The EMBO Journal* 7(12):3745-3755, 1988.

Cole et al., "CD8: Adhesion Molecule, Co-Receptor and Immunomodulator," *Cellular & Molecular Immunology* 1(2):81-88, Apr. 2004.

Dangaj et al., "Novel Recombinant Human B7-H4 Antibodies Overcome Tumoral Immune Escape to Potentiate T-Cell Antitumor Responses," *Cancer Res* 73(15):4820-4829, Aug. 1, 2013.

Desjarlais et al., "Use of a zinc-finger consensus sequence framework and specificity rules to design specific DNA binding proteins," *Proc. Natl. Acad. Sci. USA* 90:2256-2260, Mar. 1993.

Dossett et al., "Adoptive Immunotherapy of Disseminated Leukemia With TCR-transduced, CD8⁺ T Cells Expressing a Known Endogenous TCR," *Molecular Therapy* 17(4):742-749, 2009.

Dudley et al., "Adoptive Transfer of Cloned Melanoma-Reactive T Lymphocytes for the Treatment of Patients with Metastatic Melanoma," *Journal of Immunotherapy* 24(4):363-373, 2001.

Dujon et al., "Mobile introns: definition of terms and recommended nomenclature," *Gene* 82:115-118, 1989.

Dunbar et al., "ANARCI: antigen receptor numbering and receptor classification," *Bioinformatics* 32(2):298-300, 2016.

Engels et al., "Retroviral Vectors for High-Level Transgene Expression in T Lymphocytes," *Human Gene Therapy* 14:1155-1168, Aug. 10, 2003.

Epinat et al., "A novel engineered meganuclease induces homologous recombination in yeast and mammalian cells," *Nucleic Acids Research* 31(11):2952-2962, 2003.

Floros et al., "Anticancer Cytokines: Biology and Clinical Effects of IFN- α 2, IL-2, IL-15, IL-21, and IL-12," *Semin Oncol.* 42(4):539-548, Aug. 2015 (HHS Public Access Author manuscript, available in PMC Aug. 1, 2016) (17 pages).

Frasca et al., "BRAF(V600E) mutation and the biology of papillary thyroid cancer," *Endocrine-Related Cancer* 15:191-205, 2008.

Gao et al., "Molecular interactions of coreceptor CD8 and MHC class I: the molecular basis for functional coordination with the T-cell receptor," *Immunology Today* 21(12):630-636, Dec. 2000.

GenBank, "*Homo sapiens* B-Raf proto-oncogene, serine/threonine kinase (BRAF), mRNA," Accession No. NM_004333.4, Jul. 17, 2017. (6 pages).

Gimble et al., "Substrate Recognition and Induced DNA Distortion by the PI-SceI Endonuclease, an Enzyme Generated by Protein Splicing," *J. Mol. Biol.* 263:163-180, 1996.

Green et al., "Mitochondria and Apoptosis," *Science* 281: 1309-1312, Aug. 28, 1998.

Gros et al., "Prospective identification of neoantigen-specific lymphocytes in the peripheral blood of melanoma patients," *Nat Med.* 22(4):433-438, Apr. 2016 (HHS Public Access Author manuscript, available in PMC Aug. 25, 2020) (19 pages).

Hanahan et al., "Hallmarks of Cancer: The Next Generation," *Cell* 144:646-674, Mar. 4, 2011.

Hanahan et al., "The Hallmarks of Cancer," *Cell* 100:57-70, Jan. 7, 2000.

Harris et al., "Adoptive T Cell Therapies: A Comparison of T Cell Receptors and Chimeric Antigen Receptors," *Trends in Pharmacological Sciences* 37(3):220-230, Mar. 2016.

Ho et al., "In vitro methods for generating CD8⁺ T-cell clones for immunotherapy from the naive repertoire," *Journal of Immunological Methods* 310:40-52, 2006.

Janeway et al., "Immunobiology: The Immune System in Health and Disease," 3rd Ed., Current Biology Publications, p. 4:33, 1997. (14 pages).

Jasin, "Genetic manipulation of genomes with rare-cutting endonucleases," *TIG* 12(6):224-228, Jun. 1996.

Jinek et al., "A Programmable Dual-RNA-Guided DNA Endonuclease in Adaptive Bacterial Immunity," *Science* 337:816-821, Aug. 17, 2012.

Johnson et al., "Gene therapy with human and mouse T-cell receptors mediates cancer regression and targets normal tissues expressing cognate antigen," *Blood* 114(3):535-546, Jul. 16, 2009. (13 pages).

Jones et al., "Lentiviral Vector Design for Optimal T Cell Receptor Gene Expression in the Transduction of Peripheral Blood Lymphocytes and Tumor-Infiltrating Lymphocytes," *Human Gene Therapy* 20:630-640, Jun. 2009.

Jores et al., "Resolution of hypervariable regions in T-cell receptor β chains by a modified Wu-Kabat index of amino acid diversity," *Proc. Natl. Acad. Sci. USA* 87:9138-9142, Dec. 1990.

Kabat et al., "Sequences of Proteins of Immunological Interest," Fifth Edition, U.S. Department of Health and Human Services, 1991. (84 pages).

Kim et al., "Analysis of the Paired TCR α - and β -chains of Single Human T Cells," *PLoS ONE* 7(5):e37338, May 2012. (12 pages).

Koboldt et al., "VarScan 2: Somatic mutation and copy No. alteration discovery in cancer by exome sequencing," *Genome Research* 22:568-576, 2012.

(56)

References Cited

OTHER PUBLICATIONS

- Kreiter et al., "Increased Antigen Presentation Efficiency by Coupling Antigens to MHC Class I Trafficking Signals," *The Journal of Immunology* 180:309-318, 2008 (12 pages).
- Kuball et al., "Facilitating matched pairing and expression of TCR chains introduced into human T cells," *Blood* 109(6):2331-2338, Mar. 15, 2007.
- Kyrgidis et al., "Melanoma: Stem cells, sun exposure and hallmarks for carcinogenesis, molecular concepts and future clinical implications," *Journal of Carcinogenesis* 9:3, Apr. 1, 2010. (29 pages).
- Larsson et al., "Local co-administration of gene-silencing RNA and drugs in cancer therapy: State-of-the art and therapeutic potential," *Cancer Treatment Reviews* 55:128-135, 2017.
- Leen et al., "Improving T Cell Therapy for Cancer," *Annu. Rev. Immunol.* 25:243-265, 2007.
- Lefranc et al., "IMGT unique numbering for immunoglobulin and T cell receptor variable domains and Ig superfamily V-like domains," *Developmental and Comparative Immunology* 27:55-77, 2003.
- Lehninger, Lehninger (eds.), *Biochemistry, The Molecular Basis of Cell Structure and Function*, Worth Publishers, Inc., New York, N.Y., 1975, pp. 45-52. (12 pages).
- Li et al., "Fast and accurate short read alignment with Burrows-Wheeler transform," *Bioinformatics* 25(14):1754-1760, 2009.
- Li et al., "RSEM: accurate transcript quantification from RNA-Seq data with or without a reference genome," *BMC Bioinformatics* 12:323, 2011. (16 pages).
- Li, "Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM," 2013. (3 pages).
- Lim et al., "Hepatitis B virus nuclear export elements: RNA stem-loop α and β , key parts of the HBV post-transcriptional regulatory element," *RNA Biology* 13(9):743-747, 2016. (6 pages).
- Liu et al., "Selective inhibition of IDO1 effectively regulates mediators of antitumor immunity," *Blood* 115(17):3520-3530, Apr. 29, 2010.
- Manici et al., "Melanoma Cells Present a MAGE-3 Epitope to CD4+ Cytotoxic T Cells in Association with Histocompatibility Leukocyte Antigen DR11," *J. Exp. Med.* 189(5):871-876, Mar. 1, 1999.
- Mautino et al., "NLG919, a novel indoleamine-2,3-dioxygenase (IDO)-pathway inhibitor drug candidate for cancer therapy," American Association for Cancer Research 104th Annual Meeting, Apr. 6-10, 2013 (Abstract only). (1 page).
- Munson et al., "Identification of shared TCR sequences from T cells in human breast cancer using emulsion RT-PCR," *PNAS* 113(29):8272-8277, Jul. 19, 2016.
- Penix et al., "Two Essential Regulatory Elements in the Human Interferon γ Promoter Confer Activation Specific Expression in T Cells," *The Journal of Experimental Medicine* 178:1483-1496, Nov. 1993.
- Perler et al., "Protein splicing elements: inteins and exteins—a definition of terms and recommended nomenclature," *Nucleic Acids Research* 22(7): 1125-1127, 1994.
- Petersdorf et al., "The International Histocompatibility Working Group in Hematopoietic Cell Transplantation," *Int J Immunogenet* 40(1):Feb. 2013 (NIH Public Access Author Manuscript, available in PMC Feb. 1, 2014) (15 pages).
- Porteus et al., "Gene targeting using zinc finger nucleases," *Nature Biotechnology* 23(8):967-973, Aug. 2005.
- Pâques et al., "Meganucleases and DNA Double-Strand Break-Induced Recombination: Perspectives for Gene Therapy," *Current Gene Therapy* 7:49-66, 2007.
- Quezada et al., "Tumor-reactive CD4+ T cells develop cytotoxic activity and eradicate large established melanoma after transfer into lymphopenic hosts," *J. Exp. Med.* 207(3):637-650, Mar. 15, 2010.
- Ramos et al., "Oncotator: cancer variant annotation tool," *Hum Mutat.* 36(4):E2423-E2429, Apr. 2015 (HHS Public Access Author manuscript, available in PMC Jul. 10, 2020) (9 pages).
- Redmond et al., "Single-cell TCRseq: paired recovery of entire T-cell alpha and beta chain transcripts in T-cell receptors from single-cell RNAseq," *Genome Medicine* 8:80, 2016 (13 pages).
- Ren et al., "Multiplex Genome Editing to Generate Universal CAR T Cells Resistant to PD1 Inhibition," *Clin Cancer Res* 23(9):2255-2266, May 1, 2017. (13 pages).
- Riddell et al., "The use of anti-CD3 and anti-CD28 monoclonal antibodies to clone and expand human antigen-specific T cells," *Journal of Immunological Methods* 128:189-201, 1990.
- Robins et al., "Comprehensive assessment of T-cell receptor B-chain diversity in $\alpha\beta$ T cells," *Blood* 114(19):4099-4107, Nov. 5, 2009.
- Robins et al., "Overlap and Effective Size of the Human CD8+ T Cell Receptor Repertoire," *Science Translational Medicine* 2(47):1-9, Sep. 1, 2010.
- Robins et al., "Ultra-sensitive detection of rare T cell clones," *Journal of Immunological Methods* 375:14-19 (2012).
- Sadelain et al., *The Basic Principles of Chimeric Antigen Receptor Design*, *Cancer Discovery* 3(4):388-398, 2013.
- Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, 1989. (3 pages).
- Scatchard et al., "The Attractions of Proteins for Small Molecules And Ions," *Annals New York Academy of Sciences* 1:660-672, 1949.
- Schmitt et al., "T Cell Receptor Gene Therapy for Cancer," *Human Gene Therapy* 20:1240-1248, 2009.
- Scholten et al., "Codon modification of T cell receptors allows enhanced functional expression in transgenic human T cells," *Clinical Immunology* 119:135-145, 2006.
- Shi et al., "Acquired Resistance and Clonal Evolution in Melanoma during BRAF Inhibitor Therapy," *Cancer Discov* 4(1):80-93, 2013.
- Sommermeier et al., "Chimeric antigen receptor-modified T cells derived from defined CD8+ and CD4+ subsets confer superior antitumor reactivity in vivo," *Leukemia* 30(2):492-500, Feb. 2016 (HHS Public Access Author manuscript, available in PMC May 18, 2016) (20 pages).
- Stone et al., "A novel T cell receptor single-chain signaling complex mediates antigen-specific T cell activity and tumor control," *Cancer Immunol Immunother* 63:1163-1176, Aug. 1, 2014.
- Stone et al., "Role of T cell receptor affinity in the efficacy and specificity of adoptive T cell therapies," *Frontiers in Immunology* 4(244):1-16, Aug. 2013.
- Stromnes et al., "Re-adapting T cells for cancer therapy: from mouse models to clinical trials," *Immunological Reviews* 257:145-164, 2014.
- Sun et al., "Defective CD8 T Cell Memory Following Acute Infection Without CD4 T Cell Help," *Science* 300(5617):339-342, Apr. 11, 2003 (NIH Public Access Author Manuscript, available in PMC Nov. 17, 2009) (7 pages).
- Sussman et al., "Isolation and Characterization of New Homing Endonuclease Specificities at Individual Target Site Positions," *J. Mol. Biol.* 342:31-41, 2004.
- Terentis et al., "The Selenazal Drug Ebselen Potently Inhibits Indoleamine 2,3-Dioxygenase by Targeting Enzyme Cysteine Residues," *Biochemistry* 49(3):591-600, 2010.
- Thompson et al., "cis-Acting Sequences Required for Inducible Interleukin-2 Enhancer Function Bind a Novel Ets-Related Protein, Elf-1," *Molecular and Cellular Biology* 12(3):1043-1053, Mar. 1992.
- Till et al., "Adoptive immunotherapy for indolent non-Hodgkin lymphoma and mantle cell lymphoma using genetically modified autologous CD20-specific T cells," *Blood* 112(6):2261-2271, Sep. 15, 2008.
- Todd et al., "Transcription of the Interleukin 4 Gene Is Regulated by Multiple Promoter Elements," *J. Exp. Med.* 177:1663-1674, Jun. 1993.
- Torikai et al., "A foundation for universal T-cell based immunotherapy: T cells engineered to express a CD19-specific chimeric-antigen-receptor and eliminate expression of endogenous TCR," *Blood* 119(24):5697-5705, Jun. 14, 2012.
- Torikai et al., "Genetic editing of HLA expression in hematopoietic stem cells to broaden their human application," *Scientific Reports* 6:21757, Feb. 23, 2016.
- Torikai et al., "Toward eliminating HLA class I expression to generate universal cells from allogeneic donors," *Blood* 122(8):1341-1349, Aug. 22, 2013.

(56)

References Cited



OTHER PUBLICATIONS



Tran et al., "Cancer Immunotherapy Based on Mutation-Specific CD4+ T Cells in a Patient with Epithelial Cancer," *Science* 344(6184):641-645, May 9, 2014. (6 pages).
 UniProt, "BRAF_Human," sequence ID No. P15056, downloaded Nov. 2, 2022. (24 pages).
 UniProt, "CD4_Human," sequence ID No. P01730, downloaded Nov. 2, 2022. (16 pages).
 UniProt, "CD8A_Human," sequence ID No. P01732, downloaded Nov. 2, 2022. (9 pages).
 UniProt, "CD8B_Human," sequence ID No. P10966, downloaded Nov. 2, 2022. (7 pages).
 Van der Auwera et al., "From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices pipeline," *Curr Protoc Bioinformatics* 11(1110):11.10.1-11.10.33 (NIH Public Access Author Manuscript, available in PMC Nov. 25, 2014) (43 pages).
 Wang et al., "Optimizing Adoptive Polyclonal T Cell Immunotherapy of Lymphomas, Using a Chimeric T Cell Receptor Possessing CD28 and CD137 Costimulatory Domains," *Human Gene Therapy* 18:712-725, Aug. 2007.
 Warren et al., "Exhaustive T-cell repertoire sequencing of human peripheral blood samples reveals signatures of antigen selection and a directly measured repertoire size of at least 1 million clonotypes," *Genome Research* 21:790-797 2011.



Williams et al., "Interleukin-2 signals during priming are required for secondary expansion of CD8+ memory T cells," *Nature* 441(7095):890-893, Jun. 15, 2006 (NIH Public Access Author Manuscript, available in PMC Nov. 11, 2009) (12 pages).
 Wilson, "Analyzing Biomolecular Interactions," *Science* 295:2103, 2105, Mar. 15, 2002. (3 pages).
 Wolfe et al., "Analysis of Zinc Fingers Optimized via Phage Display: Evaluating the Utility of a Recognition Code," *J. Mol. Biol.* 285:1917-1934 (1999).
 Wolff et al., "Monoclonal Antibody Homodimers: Enhanced Antitumor Activity in Nude Mice," *Cancer Research* 53:2560-2565, Jun. 1, 1993.
 Xie et al., "sgRNAcas9: A Software Package for Designing CRISPR sgRNA and Evaluating Potential Off-Target Cleavage Sites," *PLoS ONE* 9(6):e100448, Jun. 23, 2014. (9 pages).
 Yee et al., "Adoptive T cell therapy using antigen-specific CD8+ T cell clones for the treatment of patients with metastatic melanoma: In vivo persistence, migration, and antitumor effect of transferred T cells," *PNAS* 99(25):16168-16173, Dec. 10, 2002.
 Henkart et al., "Chapter 36: Cytotoxic T-Lymphocytes," *Fundamental Immunology*, Lippincott Williams & Wilkins, Philadelphia, PA, 2003, pp. 1127-1150.
 Coffin, "Chapter 58: Retroviridae: The viruses and their replication," *Fundamental Virology*, Third Edition, edited by Fields et al., Lippincott-Raven Publishers, Philadelphia, 1996, 1767-1847.







FIG. 1A

*FIG. 1B**FIG. 1C*

*FIG. 1D**FIG. 1E*

*FIG. 1F**FIG. 1G*

*FIG. 1H**FIG. 1I*

*FIG. 2A**FIG. 2B*












FIG. 2C



*FIG. 2F**FIG. 2G*

*FIG. 2H**FIG. 2I*

*FIG. 2J**FIG. 2K*




FIG. 2L




FIG. 2M




FIG. 2N




FIG. 20




FIG. 3A




FIG. 3B




FIG. 3B (Continued)







FIG. 3C

*FIG. 4A**FIG. 4B*

*FIG. 5*




FIG. 6B




FIG. 6A

1

BRAF-SPECIFIC TCRS AND USES THEREOF**STATEMENT OF GOVERNMENT INTEREST**

This invention was made with government support under CA015704 awarded by the National Institutes of Health. The government has certain rights in the invention.

STATEMENT REGARDING SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is 360056_454USPC_SEQUENCE_LISTING.txt. The text file is 126 KB, was created on Aug. 9, 2020, and is being submitted electronically via EFS-Web.

BACKGROUND

Adoptive transfer of tumor-specific T cells is an appealing strategy to eliminate existing tumors and requires the establishment of a robust population of antigen-specific T cells in vivo to eliminate existing tumor and prevent recurrences (Stromnes et al., *Immunol. Rev.* 257:145, 2014). In recent years, there is increasing evidence that immune responses to antigens created by mutations in cancer can be recognized by T cells and that these T cells can mediate clinical responses to treatment with adoptive cell therapy and immune checkpoint inhibitors. Antigens that arise from such mutations are particularly appealing targets for immunotherapies due to being completely specific for the cancer relative to normal tissue, and also because they lack central tolerance mechanisms that could limit T cell function against other antigen types.

However, although administration of autologous or engineered allogeneic tumor-specific CD8⁺ cytotoxic T lymphocytes (CTLs) can mediate direct anti-tumor activity in select patients (Chapuis et al., *Cancer Res.* 72:LB-136, 2012; Chapuis et al., *Sci. Transl. Med.* 5:174ra127, 2013; Chapuis et al., *Proc. Nat'l. Acad. Sci. U.S.A.* 109:4592, 2012)²⁻⁴, identifying and isolating tumor-reactive T cells with desired characteristics is a laborious and complex endeavor (see Stone and Kranz, *Frontiers Immunol.* 4:244, 2013; Chapuis et al., 2013; Schmitt et al., *Hum. Gene Ther.* 20:1240, 2009; Ho et al., *J. Immunol. Methods* 310:40, 2006). Further, the variability in the avidity of the CTLs isolated from each patient or donor limits the anti-tumor efficacy in clinical trials (Chapuis et al., 2013). Moreover, most antigen-specific mutations that lead to immune responses are found only in the cancer of one individual, and not in multiple patients.

There is a clear need for alternative antigen-specific TCR immunotherapies directed against various cancers, such as hairy cell leukemia, malignant melanoma, thyroid, lung, and colon cancers. In particular, TCR immunotherapies targeting antigens that are both cancer-specific and widely prevalent in cancers are needed. Presently disclosed embodiments address these needs and provide other related advantages.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1I show the identification and characterization of patient-derived CD4⁺ T cells specific for BRAF^{V600E}. (1A) Positron emission tomography showing recurrent tumor in left iliac region (left) and left thigh (right). (1B-1D) Specificity and HLA restriction of BRAF^{V600E}-specific T

2

cells: (1B) IFN- γ production by the patient-derived T cell line incubated with autologous B cells pulsed with wildtype and mutant BRAF peptide; (1C) recognition of autologous B cells pulsed with mutant BRAF peptide or transfected with mRNA encoding mutant or wildtype BRAF sequences; (1D) recognition of autologous B cells pulsed with mutant BRAF peptide in the presence or absence of HLA blocking antibodies. (1E) Recognition by BRAF^{V600E}-specific CD4⁺ T cells of the B-LCL line 1331, which is matched at HLA-DQ with the patient, and the HLA-mismatched B-LCL line VAVY, prior to and after transduction with HLA-DRB1*0404 (DR4) or HLA-DQB1*0302/DQA1*03 (DQ3). (1F) IFN- γ release by patient-derived BRAF^{V600E}-specific T cells incubated with allogeneic B-LCL cell lines expressing HLA DQB1*03 alleles and pulsed with the indicated amount of 21-mer BRAF^{V600E} peptide. Three technical replicates were performed. (1G) IFN- γ release by patient-derived BRAF^{V600E}-specific T cells incubated with autologous B cells pulsed with BRAF^{V600E} peptide or the indicated tumor cell lines with and without pretreatment with human IFN- γ 500 U/ml for 3 days. (1H, 1I) Expression (mean fluorescence intensity) of HLA-DQ (1H) and HLA-DR (1I) on tumor cell lines with and without IFN- γ pretreatment, quantitated by flow cytometry relative to the isotype control. Experiments were performed in technical duplicate or triplicate as indicated, and are representative of two independent experiments.

FIGS. 2A-2O show the specificity of CD8⁺ T cells in TIL (tumor infiltrating lymphocytes) and TCR sequencing of T cell clonotypes in blood after adoptive transfer. (2A) IFN- γ production by TIL incubated with autologous CD40L-activated B cells pulsed with 13 pools of peptides encompassing 40 20-mer peptides containing the 20 nonsynonymous mutations present in the patient's melanoma by elispot assay. The final concentration of each peptide in the assay was 10 μ g/ml. Three technical replicates were performed. (2B-2F) IFN- γ production by TILs incubated with B cells transduced with tandem minigenes encompassing 29 non-synonymous mutations or the coding sequences from self-antigens Tyrosinase, Mage A3, Mart1, SSX2, and GP100 in the presence of brefeldin A (2B-2E) or pulsed with tumor-associated self-antigens (2F). The final concentration of each peptide in the assay was 10 μ g/ml. Three technical replicates were performed. (2G) Frequency of TCR V β sequences in peripheral blood mononuclear cells after mock stimulation, BRAF^{V600E} peptide stimulation, or after sorting IFN- γ secreting cells following BRAF^{V600E} peptide re-stimulation. (2H) TCR V β clonotypes of BRAF-specific T cells quantitated by TCR β sequencing of pre-treatment blood, tumor single cell suspension, and the TIL product infused into the patient. (2I) TCRV β sequences in TIL product ranked by prevalence (right-hand portion of graph). (2J) Frequency of the top 34 TIL TCR V β clonotypes from (2I) in pre-treatment blood and post-treatment blood obtained at 10 and 24 months. (2K) Frequency of TCR V β clonotypes of CD4⁺ BRAF^{V600E} and CD8⁺ T cells specific for the specified antigens in pre-treatment and post-treatment blood. (2L-2O) TCR β sequencing on the TIL and T cells from TIL incubated with autologous B cells and tiled peptides. Tiled peptides spanned (2L) Tyrosinase, (2M) Mart1, (2N) Mage A3, and (2O) TRP2 and sorted by IFN- γ capture. Antigen-specific TCR β sequences enriched in the sorted cells are marked with a box.

FIGS. 3A-3C show phenotypic analysis for BRAF-specific T cells following treatment with tumor-infiltrating lymphocytes (TILs). (3A) Dump channel gating scheme for excluding monocytes (CD14⁺), B cells (CD19⁺), and dead

3

cells (ViaProbe) from TIL. Viable CD4⁺ T cells were plotted against tetramer and CD45RA. (3B) CD45RA⁺ memory cells (88.1% of CD4⁺) that were tetramer-positive and -negative were plotted against the indicated cell surface markers. Numbers indicate the percentage of cells in the gated regions or the percentage of tetramer-positive cells for each marker. (3C) Intracellular cytokine staining of activated (CD154⁺) BRAF-specific T cells.

FIGS. 4A-4B show that a synthetic TCR derived from the dominant V α and V β sequences of melanoma-responsive patient TILs recognizes cells expressing BRAF^{V600E}. (4A) Frequency of TCRBV α sequences in peripheral blood mononuclear cells after mock stimulation, BRAF^{V600E} stimulation, or after sorting IFN- γ secretion cells following BRAF^{V600E} peptide re-stimulation. (4B) IFN- γ production by CD4⁺ T cells from two normal donors transduced with a synthetic TCR construct and incubated with an HLA-DQB1*0302 B cell line 1331 pulsed with BRAF^{V600E} peptide or transfected with mRNA encoding mutant or wildtype BRAF sequences. N=2 or 3 technical replicates as indicated.

FIG. 5 shows IFN- γ production by CD4⁺ T cells transduced with one of four TCRs (pJV88-pJV91) made using TCR β and TCR α genes identified in the patient TILs. Left, no antigen peptide. Right, antigen peptide.

FIGS. 6A and 6B show the effects of CRISPR-mediated knockout of endogenous TCR sequences on expression of a heterologous BRAF-specific TCR in primary human CD4⁺ T cells. (6A) Stimulated T cells were transfected with Cas9-RNPs targeting TCRA and TCRB and transduced with DNA encoding BRAF^{V600E}-specific TCR. BRAF-specific TCR expression was measured by Vbeta3.1 and TCR expression was measured using anti-CD3. Top panels: unmodified cells without (left) or with (right) CRISPR-mediated TCR knockout. Bottom panels: transduced cells without (left) or with (right) CRISPR-mediated TCR knockout. (6B) Tetramer staining: T cells modified with BRAF^{V600E}-specific TCR with or without deletion of endogenous TCR were compared to untransduced cells (left-most peaks) and a patient-derived antigen-specific T cell clone (right-most peaks) for binding to tetramer.

DETAILED DESCRIPTION

In certain aspects, the present disclosure provides binding proteins, such as T cell receptors (TCRs), that are capable of specifically binding to a BRAF^{V600E} peptide antigen, such as a BRAF^{V600E} peptide antigen associated with a major histocompatibility complex (MHC) (e.g., human leukocyte antigen, HLA). Binding proteins of this disclosure are useful in, for example, therapies to treat hyperproliferative diseases, such as cancer, characterized by BRAF^{V600E} expression.

By way of background, antigens created by cancer-associated mutations are appealing targets for therapeutic intervention, but are generally unique to an individual patient. Thus, antigens caused by essential “driver” mutations of cancer are of interest since they are both specific to cancer cells and occur at high frequencies in patient populations. BRAF protein is involved in cell growth signaling, while mutant BRAF is implicated in a number of cancers (see, e.g., Frasca et al., *Endocrine-Related Cancer* 15:191(2008)). In particular, the substitution mutation V600E (BRAF^{V600E}) arising in exon 15 of the BRAF gene, activates BRAF to drive a growth signaling pathway that is an early event in carcinogenesis. This mutation is found in all instances of hairy cell leukemia, about half of malignant melanoma

4

cases, and significant numbers of patients with advanced thyroid, lung and colon cancer.

The compositions and methods described herein will in certain embodiments have therapeutic utility for the treatment of diseases and conditions associated with BRAF^{V600E} expression. Such diseases include various forms of hyperproliferative disorders, such as hairy cell leukemia, melanoma, thyroid cancers including poorly differentiated thyroid cancer, non-small cell lung cancer, colorectal cancer, papillary cancer, non-Hodgkin lymphoma, glioblastoma, and pilocytic astrocytoma, breast cancer, ovarian cancer, Langerhans cell histiocytosis, and sarcomas (e.g., fibrosarcoma (fibroblastic sarcoma), Dermatofibrosarcoma protuberans (DFSP), osteosarcoma, rhabdomyosarcoma, Ewing’s sarcoma, a gastrointestinal stromal tumor, Leiomyosarcoma; angiosarcoma (vascular sarcoma), Kaposi’s sarcoma, liposarcoma, pleomorphic sarcoma, and synovial sarcoma). Non-limiting examples of these and related uses are described herein and include in vitro, ex vivo and in vivo stimulation of BRAF^{V600E} antigen-specific T cell responses, such as by the use of recombinant T cells expressing TCR specific for a BRAF^{V600E} peptide.

Prior to setting forth this disclosure in more detail, it may be helpful to an understanding thereof to provide definitions of certain terms to be used herein. Additional definitions are set forth throughout this disclosure.

In the present description, any concentration range, percentage range, ratio range, or integer range is to be understood to include the value of any integer within the recited range and, when appropriate, fractions thereof (such as one tenth and one hundredth of an integer), unless otherwise indicated. Also, any number range recited herein relating to any physical feature, such as polymer subunits, size or thickness, are to be understood to include any integer within the recited range, unless otherwise indicated. As used herein, the term “about” means $\pm 20\%$ of the indicated range, value, or structure, unless otherwise indicated. It should be understood that the terms “a” and “an” as used herein refer to “one or more” of the enumerated components. The use of the alternative (e.g., “or”) should be understood to mean either one, both, or any combination thereof of the alternatives. As used herein, the terms “include,” “have” and “comprise” are used synonymously, which terms and variants thereof are intended to be construed as non-limiting.

In addition, it should be understood that the individual compounds, or groups of compounds, derived from the various combinations of the structures and substituents described herein, are disclosed by the present application to the same extent as if each compound or group of compounds was set forth individually. Thus, selection of particular structures or particular substituents is within the scope of the present disclosure.

The term “consisting essentially of” is not equivalent to “comprising,” and refers to the specified materials or steps of a claimed invention, or to those that do not materially affect the basic characteristics of a claimed invention. For example, a protein domain, region, or module (e.g., a binding domain, hinge region, linker module) or a protein (which may have one or more domains, regions, or modules) “consists essentially of” a particular amino acid sequence when the amino acid sequence of a domain, region, module, or protein includes extensions, deletions, mutations, or a combination thereof (e.g., amino acids at the amino- or carboxy-terminus or between domains) that, in combination, contribute to at most 20% (e.g., at most 15%, 10%, 8%, 6%, 5%, 4%, 3%, 2% or 1%) of the length of a domain, region, module, or protein and do not substantially affect (i.e., do not

reduce the activity by more than 50%, such as no more than 40%, 30%, 25%, 20%, 15%, 10%, 5%, or 1%) the activity of the domain(s), region(s), module(s), or protein (e.g., the target binding affinity of a binding protein).

As used herein, an “immune system cell” means any cell of the immune system that originates from a hematopoietic stem cell in the bone marrow, which gives rise to two major lineages, a myeloid progenitor cell (which give rise to myeloid cells such as monocytes, macrophages, dendritic cells, megakaryocytes and granulocytes) and a lymphoid progenitor cell (which give rise to lymphoid cells such as T cells, B cells and natural killer (NK) cells, including Natural Killer T (NK-T) cells). Exemplary immune system cells include a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁺ CD8⁺ double negative T cell, a $\gamma\delta$ T cell, a regulatory T cell, a natural killer cell, a natural killer T cell, and a dendritic cell. Macrophages and dendritic cells may be referred to as “antigen presenting cells” or “APCs,” which are specialized cells that can activate T cells when a major histocompatibility complex (MHC) receptor on the surface of the APC complexed with a peptide interacts with a TCR on the surface of a T cell.

“Major histocompatibility complex” (MHC) refers to glycoproteins that deliver peptide antigens to a cell surface. MHC class I molecules are heterodimers having a membrane spanning α chain (with three domains) and a non-covalently associated β 2 microglobulin. MHC class II molecules are composed of two transmembrane glycoproteins, α and β , both of which span the membrane. Each chain has two domains. MHC class I molecules deliver peptides originating in the cytosol to the cell surface, where a peptide:MHC complex is recognized by CD8⁺ T cells. MHC class II molecules deliver peptides originating in the vesicular system to the cell surface, where a peptide:MHC complex is recognized by CD4⁺ T cells. Human MHC is referred to as human leukocyte antigen (HLA). HLA-II types include DP, DM, DOA, DOB, DQ, and DR. Numerous alleles encoding the subunits of the various HLA types are known, including, for example, HLA-DQA1*03, HLA-DQB1*0301, HLA-DQB1*0302, HLA-DQB1*0303. In certain embodiments, a binding protein according to the present disclosure is capable of recognizing a BRAF^{V600E} peptide complexed with HLA-DQ. In certain embodiments, the HLA complex comprises HLA-DQB1*0301, *0302, or *0303. In certain embodiments, the HLA complex comprises HLA-DQB1*0302. In further embodiments, the HLA complex comprises HLA-DQA1*03.

A “T cell” or “T lymphocyte” is an immune system cell that matures in the thymus and produces T cell receptors (TCRs). T cells can be naïve (not exposed to antigen; increased expression of CD62L, CCR7, CD28, CD3, CD127, and CD45RA, and decreased expression of CD45RO as compared to T_{CM}), memory T cells (T_M) (antigen-experienced and long-lived), and effector cells (antigen-experienced, cytotoxic). T_M can be further divided into subsets of: central memory T cells (T_{CM}; increased expression of CD62L, CCR7, CD28, CD127, CD45RO, and CD95, and decreased expression of CD54RA as compared to naïve T cells); and effector memory T cells (T_{EM}; decreased expression of CD62L, CCR7, CD28, CD45RA, and increased expression of CD127 as compared to naïve T cells or T_{CM}).

Effector T cells (T_E) refers to antigen-experienced CD8⁺ cytotoxic T lymphocytes that have decreased expression of CD62L, CCR7, CD28, and are positive for granzyme and perforin as compared to T_{CM}. Helper T cells (T_H) are CD4⁺ cells that influence the activity of other immune cells by

releasing cytokines. CD4⁺ T cells can activate and suppress an adaptive immune response, and which of those two functions is induced will depend on presence of other cells and signals. T cells can be collected using known techniques, and the various subpopulations or combinations thereof can be enriched or depleted by known techniques, such as by affinity binding to antibodies, flow cytometry, or immunomagnetic selection. Other exemplary T cells include regulatory T cells, such as CD4⁺ CD25⁺ (Foxp3⁺) regulatory T cells and Treg17 cells, as well as Tr1, Th3, CD8⁺ CD28⁺, and Qa-1 restricted T cells.

“T cell receptor” (TCR) refers to an immunoglobulin superfamily member (having a variable binding domain, a constant domain, a transmembrane region, and a short cytoplasmic tail; see, e.g., Janeway et al., *Immunobiology: The Immune System in Health and Disease*, 3rd Ed., Current Biology Publications, p. 4:33, 1997) capable of specifically binding to an antigen peptide bound to a MHC receptor. A TCR can be found on the surface of a cell or in soluble form and generally is comprised of a heterodimer having α and β chains (also known as TCR α and TCR β , respectively), or γ and δ chains (also known as TCR γ and TCR δ , respectively). Like immunoglobulins, the extracellular portion of TCR chains (e.g., α -chain, β -chain) contain two immunoglobulin domains: a variable domain (e.g., α -chain variable domain or V α , β -chain variable domain or V β ; typically amino acids 1 to 116 based on Kabat numbering (Kabat et al., “Sequences of Proteins of Immunological Interest, US Dept. Health and Human Services, Public Health Service National Institutes of Health, 1991, 5th ed.)) at the N-terminus; and one constant domain (e.g., α -chain constant domain or C α , typically amino acids 117 to 259 based on Kabat, β -chain constant domain or C β , typically amino acids 117 to 295 based on Kabat) adjacent to the cell membrane. Also, like immunoglobulins, the variable domains contain complementary determining regions (CDRs) separated by framework regions (FRs) (see, e.g., Jores et al., *Proc. Nat'l Acad. Sci. U.S.A.* 87:9138, 1990; Chothia et al., *EMBO J.* 7:3745, 1988; see also Lefranc et al., *Dev. Comp. Immunol.* 27:55, 2003). TCR variable domain sequences can be aligned to a numbering scheme (e.g., Kabat, EU, International Immunogenetics Information System (IMGT) and Aho), which can allow equivalent residue positions to be annotated and for different molecules to be compared using Antigen receptor Numbering And Receptor Classification (ANARCI) software tool (2016, Bioinformatics 15:298-300). A numbering scheme provides a standardized delineation of framework regions and CDRs in the TCR variable domains.

In certain embodiments, a TCR is found on the surface of T cells (or T lymphocytes) and associates with the CD3 complex. The source of a TCR as used in the present disclosure may be from various animal species, such as a human, mouse, rat, rabbit or other mammal.

“CD3” is known in the art as a multi-protein complex of six chains (see, Abbas and Lichtman, 2003; Janeway et al., p. 172 and 178, 1999). In mammals, the complex comprises a CD3 γ chain, a CD3 δ chain, two CD3 ϵ chains, and a homodimer of CD3 ζ chains. The CD3 γ , CD3 δ , and CD3 ϵ chains are highly related cell surface proteins of the immunoglobulin superfamily containing a single immunoglobulin domain. The transmembrane regions of the CD3 γ , CD3 δ , and CD3 ϵ chains are negatively charged, which is a characteristic that allows these chains to associate with the positively charged T cell receptor chains. The intracellular tails of the CD3 γ , CD3 δ , and CD3 ϵ chains each contain a single conserved motif known as an immunoreceptor tyrosine-based activation motif or ITAM, whereas each CD3 ζ

chain has three ITAMs. Without wishing to be bound by theory, it is believed that the ITAMs are important for the signaling capacity of a TCR complex. CD3 as used in the present disclosure may be from various animal species, including human, mouse, rat, or other mammals.

As used herein, "TCR complex" refers to a complex formed by the association of CD3 with TCR. For example, a TCR complex can be composed of a CD3 γ chain, a CD3 δ chain, two CD3 ϵ chains, a homodimer of CD3 ζ chains, a TCR α chain, and a TCR β chain. Alternatively, a TCR complex can be composed of a CD3 γ chain, a CD3 δ chain, two CD3 ϵ chains, a homodimer of CD3 ζ chains, a TCR γ chain, and a TCR δ chain.

A "component of a TCR complex," as used herein, refers to a TCR chain (i.e., TCR α , TCR β , TCR γ or TCR δ), a CD3 chain (i.e., CD3 γ , CD3 δ , CD3 ϵ or CD3 ζ), or a complex formed by two or more TCR chains or CD3 chains (e.g., a complex of TCR α and TCR β , a complex of TCR γ and TCR δ , a complex of CD3 ϵ and CD3 δ , a complex of CD3 γ and CD3 ϵ , or a sub-TCR complex of TCR α , TCR β , CD3 γ , CD3 δ , and two CD3 ϵ chains).

"CD4" refers to an immunoglobulin co-receptor glycoprotein that assists the TCR in communicating with antigen-presenting cells (see, Campbell & Reece, *Biology* 909 (Benjamin Cummings, Sixth Ed., 2002); UniProtKB P01730). CD4 is found on the surface of immune cells such as T helper cells, monocytes, macrophages, and dendritic cells, and includes four immunoglobulin domains (D1 to D4) that are expressed at the cell surface. During antigen presentation, CD4 is recruited, along with the TCR complex, to bind to different regions of the MHCII molecule (CD4 binds MHCII β 2, while the TCR complex binds MHCII α 1/ β 1). Without wishing to be bound by theory, it is believed that close proximity to the TCR complex allows CD4-associated kinase molecules to phosphorylate the immunoreceptor tyrosine activation motifs (ITAMs) present on the cytoplasmic domains of CD3. This activity is thought to amplify the signal generated by the activated TCR in order to produce various types of T helper cells.

As used herein, the term "CD8 co-receptor" or "CD8" means the cell surface glycoprotein CD8, either as an alpha-alpha homodimer or an alpha-beta heterodimer. The CD8 co-receptor assists in the function of cytotoxic T cells (CD8 $^{+}$) and functions through signaling via its cytoplasmic tyrosine phosphorylation pathway (Gao and Jakobsen, *Immunol. Today* 21:630-636, 2000; Cole and Gao, *Cell. Mol. Immunol.* 1:81-88, 2004). In humans, there are five (5) different CD8 beta chains (see UniProtKB identifier P10966) and a single CD8 alpha chain (see UniProtKB identifier P01732).

The term "variable region" or "variable domain" refers to the domain of a TCR α -chain or β -chain (or γ -chain and δ -chain for $\gamma\delta$ TCRs) that is involved in binding of the TCR to antigen. The variable domains of the α -chain and β -chain (V α and V β , respectively) of a native TCR generally have similar structures, with each domain comprising four generally conserved framework regions (FRs) and three CDRs. The V α domain is encoded by two separate DNA segments, the variable gene segment and the joining gene segment (V-J); the V β domain is encoded by three separate DNA segments, the variable gene segment, the diversity gene segment, and the joining gene segment (V-D-J). A single V α or V β domain may be sufficient to confer antigen-binding specificity. Furthermore, TCRs that bind a particular antigen may be isolated using a V α or V β domain from a TCR that binds the antigen to screen a library of complementary V α or V β domains, respectively.

The terms "complementarity determining region," and "CDR," are synonymous with "hypervariable region" or "HVR," and are known in the art to refer to non-contiguous sequences of amino acids within TCR variable regions, which confer antigen specificity and/or binding affinity. In general, there are three CDRs in each α -chain variable region (α CDR1, α CDR2, α CDR3) and three CDRs in each β -chain variable region (β CDR1, β CDR2, β CDR3). CDR3 is thought to be the main CDR responsible for recognizing processed antigen. CDR1 and CDR2 mainly interact with the MHC. In certain embodiments, a binding protein of the present disclosure comprises an α CDR1, an α CDR2, and/or an α CDR3 amino acid sequence of a V α domain as set forth in any one of SEQ ID NOs:1-4. In certain embodiments, a binding protein of the present disclosure comprises a β CDR1, a β CDR2, and/or a β CDR3 amino acid sequence of a V β domain as set forth in any one of SEQ ID NOs:5-7.

"Antigen" or "Ag" as used herein refers to an immunogenic molecule that provokes an immune response. This immune response may involve antibody production, activation of specific immunologically-competent cells (e.g., T cells), or both. An antigen (immunogenic molecule) may be, for example, a peptide, glycopeptide, polypeptide, glycopolypeptide, polynucleotide, polysaccharide, lipid or the like. It is readily apparent that an antigen can be synthesized, produced recombinantly, or derived from a biological sample. Exemplary biological samples that can contain one or more antigens include tissue samples, tumor samples, cells, biological fluids, or combinations thereof. Antigens can be produced by cells that have been modified or genetically engineered to express an antigen. Exemplary antigens include BRAF^{V600E}.

The term "epitope" or "antigenic epitope" includes any molecule, structure, amino acid sequence or protein determinant that is recognized and specifically bound by a cognate binding molecule, such as an immunoglobulin, T cell receptor (TCR), chimeric antigen receptor, or other binding molecule, domain or protein. Epitopic determinants generally contain chemically active surface groupings of molecules, such as amino acids or sugar side chains, and can have specific three dimensional structural characteristics, as well as specific charge characteristics.

A "binding domain" (also referred to as a "binding region" or "binding moiety"), as used herein, refers to a molecule or portion thereof (e.g., peptide, oligopeptide, polypeptide, protein) that possesses the ability to specifically and non-covalently associate, unite, or combine with a target (e.g., BRAF^{V600E}). A binding domain includes any naturally occurring, synthetic, semi-synthetic, or recombinantly produced binding partner for a biological molecule, a molecular complex (i.e., complex comprising two or more biological molecules), or other target of interest. Exemplary binding domains include single chain immunoglobulin variable regions (e.g., scTCR, scFv), receptor ectodomains, ligands (e.g., cytokines, chemokines), or synthetic polypeptides selected for their specific ability to bind to a biological molecule, a molecular complex or other target of interest.

As used herein, "specifically binds" or "specific for" refers to an association or union of a binding protein (e.g., TCR receptor) or a binding domain (or fusion protein thereof) to a target molecule with an affinity or K_a (i.e., an equilibrium association constant of a particular binding interaction with units of 1/M) equal to or greater than 10^5 M $^{-1}$ (which equals the ratio of the on-rate [k_{on}] to the off-rate [k_{off}] for this association reaction), while not significantly associating or uniting with any other molecules or components in a sample. Binding proteins or binding domains (or

fusion proteins thereof) may be classified as “high-affinity” binding proteins or binding domains (or fusion proteins thereof) or as “low-affinity” binding proteins or binding domains (or fusion proteins thereof). “High-affinity” binding proteins or binding domains refer to those binding proteins or binding domains having a K_d of at least 10^7 M⁻¹, at least 10^8 M⁻¹, at least 10^9 M⁻¹, at least 10^{10} M⁻¹, at least 10^{11} M⁻¹, at least 10^{12} M⁻¹, or at least 10^{13} M⁻¹. “Low-affinity” binding proteins or binding domains refer to those binding proteins or binding domains having a K_d of up to 10^7 M⁻¹, up to 10^6 M⁻¹, up to 10^5 M⁻¹. Alternatively, affinity may be defined as an equilibrium dissociation constant (K_d) of a particular binding interaction with units of M (e.g., 10^{-5} M to 10^{-13} M).

In certain embodiments, a receptor or binding domain may have “enhanced affinity,” which refers to selected or engineered receptors or binding domains with stronger binding to a target antigen than a wild type (or parent) binding domain. For example, enhanced affinity may be due to a K_a (equilibrium association constant) for the target antigen that is higher than the wild type binding domain, due to a K_d (dissociation constant) for the target antigen that is less than that of the wild type binding domain, due to an off-rate (k_{off}) for the target antigen that is less than that of the wild type binding domain, or a combination thereof. In certain embodiments, enhanced affinity TCRs may be codon optimized to enhance expression in a particular host cell, such as T cells (Scholten et al., *Clin. Immunol.* 119:135, 2006).

A variety of assays are known for identifying binding domains of the present disclosure that specifically bind a particular target, as well as determining binding domain or fusion protein affinities, such as Western blot, ELISA, analytical ultracentrifugation, spectroscopy and surface plasmon resonance (Biacore®) analysis (see, e.g., Scatchard et al., *Ann. N.Y. Acad. Sci.* 51:660, 1949; Wilson, *Science* 295:2103, 2002; Wolff et al., *Cancer Res.* 53:2560, 1993; and U.S. Pat. Nos. 5,283,173, 5,468,614, or the equivalent). Assays for assessing affinity or apparent affinity or relative affinity are also known. In certain examples, apparent affinity for a TCR is measured by assessing binding to various concentrations of tetramers, for example, by flow cytometry using labeled tetramers. In some examples, apparent K_D of a TCR is measured using 2-fold dilutions of labeled tetramers at a range of concentrations, followed by determination of binding curves by non-linear regression, apparent K_D being determined as the concentration of ligand that yielded half-maximal binding.

The term “BRAF^{V600E}-specific binding protein” refers to a protein or polypeptide that specifically binds to a BRAF^{V600E} peptide antigen or a BRAF^{V600E} peptide antigen:HLA complex, e.g., on a cell surface, and does not bind a HLA complex on a cell surface comprising a BRAF peptide not containing the BRAF^{V600E} mutation.

In certain embodiments, a BRAF^{V600E}-specific binding protein binds a BRAF^{V600E}-containing peptide:HLA complex (or BRAF^{V600E}-containing peptide:MHC complex) with a K_d of less than about 10^{-8} M, less than about 10^{-9} M, less than about 10^{-10} M, less than about 10^{-11} M, less than about 10^{-12} M, or less than about 10^{-13} M, or with an affinity that is about the same as, at least about the same as, or is greater than at or about the affinity exhibited by an exemplary BRAF^{V600E}-specific binding protein provided herein, such as any of the BRAF^{V600E}-specific TCRs provided herein, for example, as measured by the same assay. In certain embodiments, a BRAF^{V600E}-specific binding protein comprises a BRAF^{V600E}-specific immunoglobulin superfamily binding protein or binding portion thereof.

The term “BRAF^{V600E} binding domain” or “BRAF^{V600E} binding fragment” refers to a domain or portion of a BRAF^{V600E}-specific binding protein responsible for the specific BRAF^{V600E} binding. A BRAF^{V600E}-specific binding domain alone (i.e., without any other portion of a BRAF^{V600E}-specific binding protein) can be soluble and can bind to BRAF^{V600E} (e.g., in complex with an MHC receptor molecule or functional fragment thereof) with a K_d of less than about 10^{-8} M, less than about 10^{-9} M, less than about 10^{-10} M, less than about 10^{-11} M, less than about 10^{-12} M, or less than about 10^{-13} M. Exemplary BRAF^{V600E}-specific binding domains include BRAF^{V600E}-specific scTCR (e.g., single chain $\alpha\beta$ TCR proteins such as V α -L-V β , V β -L-V α , V α -C α -L-V α , or V α -L-V β -C β , wherein V α and V β are TCR α and β variable domains respectively, C α and C β are TCR α and β constant domains, respectively, and L is a linker) and scFv fragments as described herein, which can be derived from an anti-BRAF^{V600E} TCR or antibody.

Principles of antigen processing by antigen presenting cells (APC) (such as dendritic cells, macrophages, lymphocytes or other cell types), and of antigen presentation by APC to T cells, including major histocompatibility complex (MHC)-restricted presentation between immunocompatible (e.g., sharing at least one allelic form of an MHC gene that is relevant for antigen presentation) APC and T cells, are well established (see, e.g., Murphy, Janeway’s Immunobiology (8th Ed.) 2011 Garland Science, NY; chapters 6, 9 and 16). For example, processed antigen peptides originating in the cytosol (e.g., tumor antigen, intracellular pathogen) are generally from about 7 amino acids to about 11 amino acids in length and will associate with class I MHC molecules, whereas peptides processed in the vesicular system (e.g., bacterial, viral) will generally vary in length from about 10 amino acids to about 25 amino acids and associate with class II MHC molecules.

“BRAF^{V600E} antigen” or “BRAF^{V600E} peptide antigen” or “BRAF^{V600E}-containing peptide antigen” refers to a naturally or synthetically produced portion of a BRAF protein ranging in length from about 7 amino acids to about 20 amino acids and comprising the V600E substitution mutation (e.g., a peptide from BRAF⁵⁹⁷⁻⁶⁰³, or BRAF⁵⁹⁰⁻⁶¹⁰, that includes a glutamic acid substituted for a valine at the residue corresponding to position 600 of the full-length wild-type BRAF; see, e.g., Uniprot entry no. P15056 and NCBI Reference identifier NP_004324.2), which can form a complex with a MHC (e.g., HLA) molecule and such a complex can bind with a binding protein specific for a BRAF^{V600E} peptide:MHC (e.g., HLA) complex. Exemplary BRAF^{V600E} peptide antigens include those having the amino acid sequence set forth in SEQ ID NO.: 38 or 39.

A “linker” refers to an amino acid sequence that connects two proteins, polypeptides, peptides, domains, regions, or motifs and may provide a spacer function compatible with interaction of the two sub-binding domains so that the resulting polypeptide retains a specific binding affinity (e.g., scTCR) to a target molecule or retains signaling activity (e.g., TCR complex). In certain embodiments, a linker is comprised of about two to about 35 amino acids, for instance, or about four to about 20 amino acids or about eight to about 15 amino acids or about 15 to about 25 amino acids.

“Junction amino acids” or “junction amino acid residues” refer to one or more (e.g., about 2-10) amino acid residues between two adjacent motifs, regions or domains of a polypeptide, such as between a binding domain and an adjacent constant domain or between a TCR chain and an adjacent self-cleaving peptide. Junction amino acids may

result from the construct design of a fusion protein (e.g., amino acid residues resulting from the use of a restriction enzyme site during the construction of a nucleic acid molecule encoding a fusion protein).

An “altered domain” or “altered protein” refers to a motif, region, domain, peptide, polypeptide, or protein with a non-identical sequence identity to a wild type motif, region, domain, peptide, polypeptide, or protein (e.g., a wild type TCR α chain, TCR β chain, TCR α constant domain, TCR β constant domain) of at least 85% (e.g., 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, 99.9%).

As used herein, “nucleic acid” or “nucleic acid molecule” refers to any of deoxyribonucleic acid (DNA), ribonucleic acid (RNA), oligonucleotides, fragments generated, for example, by the polymerase chain reaction (PCR) or by in vitro translation, and fragments generated by any of ligation, scission, endonuclease action, or exonuclease action. In certain embodiments, the nucleic acids of the present disclosure are produced by PCR. Nucleic acids may be composed of monomers that are naturally occurring nucleotides (such as deoxyribonucleotides and ribonucleotides), analogs of naturally occurring nucleotides (e.g., α -enantiomeric forms of naturally-occurring nucleotides), or a combination of both. Modified nucleotides can have modifications in or replacement of sugar moieties, or pyrimidine or purine base moieties. Nucleic acid monomers can be linked by phosphodiester bonds or analogs of such linkages. Analogs of phosphodiester linkages include phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoranilidate, phosphoramidate, and the like. Nucleic acid molecules can be either single stranded or double stranded.

The term “isolated” means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally occurring nucleic acid or polypeptide present in a living animal is not isolated, but the same nucleic acid or polypeptide, separated from some or all of the co-existing materials in the natural system, is isolated. Such nucleic acid could be part of a vector and/or such nucleic acid or polypeptide could be part of a composition (e.g., a cell lysate), and still be isolated in that such vector or composition is not part of the natural environment for the nucleic acid or polypeptide. The term “gene” means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (“leader and trailer”) as well as intervening sequences (introns) between individual coding segments (exons).

As used herein, the term “recombinant” refers to a cell, microorganism, nucleic acid molecule, or vector that has been genetically engineered by human intervention—that is, modified by introduction of a heterologous nucleic acid molecule, or refers to a cell or microorganism that has been altered such that expression of an endogenous nucleic acid molecule or gene is controlled, deregulated, deleted, attenuated, or constitutive. Human generated genetic alterations may include, for example, modifications that introduce nucleic acid molecules (which may include an expression control element, such as a promoter) that encode one or more proteins or enzymes, or other nucleic acid molecule additions, deletions, substitutions, or other functional disruption of or addition to a cell’s genetic material. Exemplary modifications include those in coding regions or functional fragments thereof of heterologous or homologous polypeptides from a reference or parent molecule.

As used herein, “mutation” refers to a change in the sequence of a nucleic acid molecule or polypeptide molecule as compared to a reference or wild-type nucleic acid molecule or polypeptide molecule, respectively. A mutation can result in several different types of change in sequence, including substitution, insertion or deletion of nucleotide(s) or amino acid(s). In certain embodiments, a mutation is a substitution of one or three codons or amino acids, a deletion of one to about 5 codons or amino acids, or a combination thereof.

A “conservative substitution” is recognized in the art as a substitution of one amino acid for another amino acid that has similar properties. Exemplary conservative substitutions are well known in the art (see, e.g., WO 97/09433 at page 10; Lehninger, *Biochemistry*, 2nd Edition; Worth Publishers, Inc. NY, NY, pp. 71-77, 1975; Lewin, *Genes IV*, Oxford University Press, NY and Cell Press, Cambridge, MA, p. 8, 1990).

The term “construct” refers to any polynucleotide that contains a recombinant nucleic acid molecule. A construct may be present in a vector (e.g., a bacterial vector, a viral vector) or may be integrated into a genome. A “vector” is a nucleic acid molecule that is capable of transporting another nucleic acid molecule. Vectors may be, for example, plasmids, cosmids, viruses, a RNA vector or a linear or circular DNA or RNA molecule that may include chromosomal, non-chromosomal, semi-synthetic or synthetic nucleic acid molecules. Exemplary vectors are those capable of autonomous replication (episomal vector) or expression of nucleic acid molecules to which they are linked (expression vectors).

Viral vectors include retrovirus, adenovirus, parvovirus (e.g., adeno-associated viruses), coronavirus, negative strand RNA viruses such as orthomyxovirus (e.g., influenza virus), rhabdovirus (e.g., rabies and vesicular stomatitis virus), paramyxovirus (e.g., measles and Sendai), positive strand RNA viruses such as picornavirus and alphavirus, and double-stranded DNA viruses including adenovirus, herpesvirus (e.g., Herpes Simplex virus types 1 and 2, Epstein-Barr virus, cytomegalovirus), and poxvirus (e.g., vaccinia, fowlpox and canarypox). Other viruses include Norwalk virus, togavirus, flavivirus, reoviruses, papovavirus, hepadnavirus, and hepatitis virus, for example. Examples of retroviruses include avian leukosis-sarcoma, mammalian C-type, B-type viruses, D type viruses, HTLV-BLV group, lentivirus, spumavirus (Coffin, J. M., *Retroviridae: The viruses and their replication*, In *Fundamental Virology*, Third Edition, B. N. Fields et al., Eds., Lippincott-Raven Publishers, Philadelphia, 1996).

“Lentiviral vector,” as used herein, means HIV-based lentiviral vectors for gene delivery, which can be integrative or non-integrative, have relatively large packaging capacity, and can transduce a range of different cell types. Lentiviral vectors are usually generated following transient transfection of three (packaging, envelope and transfer) or more plasmids into producer cells. Like HIV, lentiviral vectors enter the target cell through the interaction of viral surface glycoproteins with receptors on the cell surface. On entry, the viral RNA undergoes reverse transcription, which is mediated by the viral reverse transcriptase complex. The product of reverse transcription is a double-stranded linear viral DNA, which is the substrate for viral integration into the DNA of infected cells.

The term “operably linked” refers to the association of two or more nucleic acid molecules on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding

sequence when it is capable of affecting the expression of that coding sequence (i.e., the coding sequence is under the transcriptional control of the promoter). "Unlinked" means that the associated genetic elements are not closely associated with one another and the function of one does not affect the other.

As used herein, "expression vector" refers to a DNA construct containing a nucleic acid molecule that is operably-linked to a suitable control sequence capable of effecting the expression of the nucleic acid molecule in a suitable host. Such control sequences include a promoter to effect transcription, an optional operator sequence to control such transcription, a sequence encoding suitable mRNA ribosome binding sites, and sequences which control termination of transcription and translation. The vector may be a plasmid, a phage particle, a virus, or simply a potential genomic insert. Once transformed into a suitable host, the vector may replicate and function independently of the host genome, or may, in some instances, integrate into the genome itself. In the present specification, "plasmid," "expression plasmid," "virus" and "vector" are often used interchangeably.

The term "expression", as used herein, refers to the process by which a polypeptide is produced based on the encoding sequence of a nucleic acid molecule, such as a gene. The process may include transcription, post-transcriptional control, post-transcriptional modification, translation, post-translational control, post-translational modification, or any combination thereof.

The term "introduced" in the context of inserting a nucleic acid molecule into a cell, means "transfection", or "transformation" or "transduction" and includes reference to the incorporation of a nucleic acid molecule into a eukaryotic or prokaryotic cell wherein the nucleic acid molecule may be incorporated into the genome of a cell (e.g., chromosome, plasmid, plastid, or mitochondrial DNA), converted into an autonomous replicon, or transiently expressed (e.g., transfected mRNA).

As used herein, "heterologous" nucleic acid molecule, construct or sequence refers to a nucleic acid molecule or portion of a nucleic acid molecule that is not native to a host cell, but may be homologous to a nucleic acid molecule or portion of a nucleic acid molecule from the host cell. The source of the heterologous nucleic acid molecule, construct or sequence may be from a different genus or species. In certain embodiments, a heterologous nucleic acid molecule is added (i.e., is not endogenous or native) to a host cell or host genome by, for example, conjugation, transformation, transfection, electroporation, or the like, wherein the added molecule may integrate into the host genome or exist as extra-chromosomal genetic material (e.g., as a plasmid or other form of self-replicating vector), and may be present in multiple copies. In addition, "heterologous" refers to a non-native enzyme, protein or other activity encoded by a heterologous polynucleotide introduced into the host cell, even if the host cell encodes a homologous protein or activity.

As described herein, more than one heterologous nucleic acid molecule can be introduced into a host cell as separate nucleic acid molecules, as a plurality of individually controlled genes, as a polycistronic nucleic acid molecule, as a single nucleic acid molecule encoding a fusion protein, or any combination thereof. For example, as disclosed herein, a host cell can be modified to express two or more heterologous nucleic acid molecules encoding desired binding proteins specific for a BRAF^{V600E} antigen peptide (e.g., TCR α and TCR β). When two or more heterologous nucleic acid molecules are introduced into a host cell, it is under-

stood that the two or more heterologous nucleic acid molecules can be introduced as a single nucleic acid molecule (e.g., on a single vector), on separate vectors, integrated into the host chromosome at a single site or multiple sites, or any combination thereof. The number of referenced heterologous nucleic acid molecules or protein activities refers to the number of encoding nucleic acid molecules or the number of protein activities, not the number of separate nucleic acid molecules introduced into a host cell.

As used herein, the term "endogenous" or "native" refers to a gene, protein, or activity that is normally present in a host cell. Moreover, a gene, protein or activity that is mutated, overexpressed, shuffled, duplicated or otherwise altered as compared to a parent gene, protein or activity is still considered to be endogenous or native to that particular host cell. For example, an endogenous control sequence from a first gene (e.g., promoter, translational attenuation sequences) may be used to alter or regulate expression of a second native gene or nucleic acid molecule, wherein the expression or regulation of the second native gene or nucleic acid molecule differs from normal expression or regulation in a parent cell.

The term "homologous" or "homolog" refers to a molecule or activity found in or derived from a host cell, species or strain. For example, a heterologous polynucleotide may be homologous to a native host cell gene, and may optionally have an altered expression level, a different sequence, an altered activity, or any combination thereof.

"Sequence identity," as used herein, refers to the percentage of amino acid residues in one sequence that are identical with the amino acid residues in another reference polypeptide sequence after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. The percentage sequence identity values can be generated using the NCBI BLAST2.0 software as defined by Altschul et al. (1997) "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs", Nucleic Acids Res. 25:3389-3402, with the parameters set to default values.

As used herein, a "hematopoietic progenitor cell" is a cell that can be derived from hematopoietic stem cells or fetal tissue and is capable of further differentiation into mature cells types (e.g., immune system cells). Exemplary hematopoietic progenitor cells include those with a CD24^{Lo} Lin⁻ CD117⁺ phenotype or those found in the thymus (referred to as progenitor thymocytes).

As used herein, the term "host" refers to a cell (e.g., T cell) or microorganism targeted for genetic modification with a heterologous nucleic acid molecule to produce a polypeptide of interest (e.g., an anti-BRAF^{V600E} TCR). In certain embodiments, a host cell may optionally already possess or be modified to include other genetic modifications that confer desired properties related or unrelated to, e.g., biosynthesis of the heterologous protein (e.g., inclusion of a detectable marker; deleted, altered or truncated endogenous TCR; or increased co-stimulatory factor expression). In certain embodiments, a host cell is a human hematopoietic progenitor cell transduced with a heterologous nucleic acid molecule encoding a TCR α chain specific for a BRAF^{V600E} antigen peptide.

As used herein, "hyperproliferative disorder" refers to excessive growth or proliferation as compared to a normal or undiseased cell. Exemplary hyperproliferative disorders include tumors, cancers, neoplastic tissue, carcinoma, sarcoma, malignant cells, pre-malignant cells, as well as non-neoplastic or non-malignant hyperproliferative disorders

15

(e.g., adenoma, fibroma, lipoma, leiomyoma, hemangioma, fibrosis, restenosis, as well as autoimmune diseases such as rheumatoid arthritis, osteoarthritis, psoriasis, inflammatory bowel disease, or the like).

Binding Proteins Specific for BRAF^{V600E} Peptide:HLA Complexes

BRAF (also known as B-RAF1, BRAF1, NS7, RAFB1, B-Raf, B-Raf proto-oncogene, and serine/threonine kinase) refers to a 766-amino acid protein encoded by the BRAF gene. The transcript sequence for human wild-type BRAF is set forth in NCBI Reference identifier NM_004333.4 (SEQ ID NO:78), and the protein sequence is set forth in NCBI Reference identifier NP_004324.2 (SEQ ID NO:36). BRAF is a member of the Raf kinase family of growth signal transduction protein kinases, and plays a role in regulating the MAP kinase/ERKs signaling pathway, which affects cell division, differentiation, and secretion. In terms of structure, BRAF is composed of three conserved domains characteristic of Raf kinases: a Ras-GTP-binding self-regulatory domain; a serine-rich hinge region; and a catalytic kinase domain that phosphorylates a consensus sequence on protein substrates (CR1, CR2, and CR3, respectively). Active B-Raf forms dimers.

A mutant form of BRAF kinase comprising a V600E mutation (BRAF^{V600E}) is an oncogenic driver present in numerous neoplastic conditions, including 40% of melanoma cases, 10% of colorectal cancer cases, and 1% of non-small cell lung cancer cases, and confers constitutive signaling that promotes tumor cell growth and survival. Small molecule BRAF inhibitors have some efficacy in melanoma, but resistance evolves by recruitment of alternative signaling pathways without loss of expression of BRAF^{V600E} protein. See Shi et al., *Cancer Disc.* 4(1):80 (2014).

In certain aspects, the present disclosure provides a binding protein comprising: (a) a T cell receptor (TCR) α chain variable (V α) domain having a CDR3 amino acid sequence set forth in any one of SEQ ID NOS:29-32, or a CDR3 amino acid sequence set forth in any one of SEQ ID NOS:29-32 with up to five amino acid substitutions, insertions, and/or deletions, and a TCR β chain variable (V β) domain; (b) a V α domain, and a V β domain having a CDR3 amino acid sequence as set forth in any one of SEQ ID NOS:33-35, or a CDR3 amino acid sequence set forth in any one of SEQ ID NOS:33-35 with up to five amino acid substitutions, insertions, and/or deletions; or (c) a V α domain of (a) and a V β domain of (b), wherein the binding protein is capable of specifically binding to a HLA complex on a cell surface comprising a BRAF peptide containing a BRAF^{V600E} mutation, and does not bind a HLA complex on a cell surface comprising a BRAF peptide not containing the BRAF^{V600E} mutation. In certain embodiments, the HLA complex comprises HLA-DQ.

In certain embodiments, (a) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:29 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:33, (b) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:30 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:34, (c) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:31 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:34, (d) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:32 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:35, (e) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:29 and the V β domain comprises the CDR3 amino acid

16

sequence of SEQ ID NO:34, (f) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:29 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:35, (g) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:30 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:33, (h) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:30 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:35, (i) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:31 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:33, (j) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:31 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:35, (k) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:32 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:33, or (l) the V α domain comprises the CDR3 amino acid sequence of SEQ ID NO:32 and the V β domain comprises the CDR3 amino acid sequence of SEQ ID NO:34.

Peptide-MHC complexes, such as BRAF^{V600E} peptide:HLA complexes, are recognized by and bound by a TCR through the V α and V β domains. During lymphocyte development, V α exons are assembled from different variable and joining gene segments (V-J), and V β exons are assembled from different variable, diversity, and joining gene segments (V-D-J). The TCR α chromosomal locus has 70-80 variable gene segments and 61 joining gene segments. The TCR β chromosomal locus has 52 variable gene segments, and two separate clusters of each containing a single diversity gene segment, together with six or seven joining gene segments. Functional V α and V β gene exons are generated by the recombination of a variable gene segment with a joining gene segment for V α , and a variable gene segment with a diversity gene segment and a joining gene segment for V β .

The V α and V β domains each comprise three hypervariable loops, also referred to as complementary determining regions (CDRs) that contact the peptide-MHC complex. CDR1 and CDR2 are encoded within the variable gene segment, whereas CDR3 is encoded by the region spanning the variable and joining segments for V α , or the region spanning variable, diversity, and joining segments for V β . Compared with CDR1 and CDR2, CDR3 is significantly more diverse because of the addition and loss of nucleotides during the recombination process.

TCR variable domain sequences can be aligned to a numbering scheme (Kabat, Chothia, Enhanced Chothia, and Aho), allowing equivalent residue positions to be annotated and for different molecules to be compared using ANARCI software tool (2016, Bioinformatics 15:298-300). A numbering scheme provides a standardized delineation of framework regions and CDRs in the TCR variable domains.

Accordingly, CDR1 and CDR2 sequences may be deduced from the corresponding variable gene segments (e.g., TCRBV28-01, TCRAV21-01, TCRAV26-01, TCRAV12-02 alleles). In certain embodiments, (a) the V α CDR3 amino acid sequence comprises the amino acid sequence set forth in SEQ ID NO:29 and the V α domain further comprises a CDR1 amino acid sequence and CDR2 amino acid sequence encoded by the polynucleotide sequence set forth in SEQ ID NO:11, (b) the V α CDR3 amino acid sequence comprises the amino acid sequence set forth in SEQ ID NO:30 and the V α domain further comprises a CDR1 amino acid sequence and CDR2 amino acid sequence encoded by the polynucleotide sequence set forth in SEQ ID NO:12, (c) the V α CDR3 amino acid sequence comprises the amino acid sequence set forth in SEQ ID

NO:31 and the V α domain further comprises a CDR1 amino acid sequence and CDR2 amino acid sequence encoded by the polynucleotide sequence set forth in SEQ ID NO:13, or (d) the V α CDR3 amino acid sequence comprises the amino acid sequence set forth in SEQ ID NO:32 and the V α domain further comprises a CDR1 amino acid sequence and CDR2 amino acid sequence encoded by the polynucleotide sequence set forth in SEQ ID NO:13. In certain embodiments, the V β CDR3 amino acid sequence comprises the amino acid sequence set forth in any one of SEQ ID NOS:33-35 and the V β domain further comprises a CDR1 amino acid sequence and CDR2 amino acid sequence encoded by the polynucleotide sequence set forth in SEQ ID NO:8.

Methods of identifying binding pairs of TCR V α and V β domains include, for example, those described in PCT Patent Publication No. WO 2016/161273; Redmond et al., 2016, *Genome Med.* 8: 80; Munson et al., 2016, *Proc. Natl. Acad. Sci.* 113:8272-7; Kim et al., 2012, *PLoS ONE* 7:e37338 (each of the methods from which are incorporated by reference in its entirety). Accordingly, a V α domain for the BRAF^{V600E}-specific V β domains described herein (e.g., a V β domain comprising CDR3 as set forth in any one of SEQ ID NOS:33-35), or vice versa, may be identified.

A BRAF^{V600E}-specific binding protein described herein may possess one or more amino acid substitutions, deletions, or insertions relative to a naturally occurring binding protein (e.g., TCR). Conservative substitutions of amino acids are known and may occur naturally or may be introduced when the binding protein or TCR is recombinantly produced. Amino acid substitutions, deletions, and insertions may be introduced into a protein using mutagenesis methods (see, e.g., Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3d ed., Cold Spring Harbor Laboratory Press, NY, 2001). Oligonucleotide-directed site-specific (or segment specific) mutagenesis procedures may be employed to provide an altered polynucleotide that has particular codons altered according to the substitution, deletion, or insertion desired. Alternatively, random or saturation mutagenesis techniques, such as alanine scanning mutagenesis, error prone polymerase chain reaction mutagenesis, and oligonucleotide-directed mutagenesis may be used to prepare immunogen polypeptide variants (see, e.g., Sambrook et al., *supra*).

A variety of criteria known in the art indicate whether an amino acid that is substituted at a particular position in a peptide or polypeptide is conservative (or similar). For example, a similar amino acid or a conservative amino acid substitution is one in which an amino acid residue is replaced with an amino acid residue having a similar side chain. Similar amino acids may be included in the following categories: amino acids with basic side chains (e.g., lysine, arginine, histidine); amino acids with acidic side chains (e.g., aspartic acid, glutamic acid); amino acids with uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine, histidine); amino acids with nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan); amino acids with beta-branched side chains (e.g., threonine, valine, isoleucine), and amino acids with aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan). Proline, which is considered more difficult to classify, shares properties with amino acids that have aliphatic side chains (e.g., leucine, valine, isoleucine, and alanine). In certain circumstances, substitution of glutamine for glutamic acid or asparagine for aspartic acid may be considered a similar substitution in that glutamine and asparagine are

amide derivatives of glutamic acid and aspartic acid, respectively. As understood in the art "similarity" between two polypeptides is determined by comparing the amino acid sequence and conserved amino acid substitutes thereto of the polypeptide to the sequence of a second polypeptide (e.g., using GENEWORKS, Align, the BLAST algorithm, or other algorithms described herein and practiced in the art).

Accordingly, in certain embodiments, the binding protein of the instant disclosure comprises a V α domain that is at least about 90% (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or at least about 99.9%) identical to the amino acid sequence set forth in any one of SEQ ID NOS:1-4, and comprises a V β domain that is at least about 90% (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or at least about 99.9%) identical to the amino acid sequence set forth in any one of SEQ ID NOS:5-7, provided that (a) at least three or four of the CDRs have no mutations and (b) the CDRs that do have mutations have only up to three amino acid substitutions, insertions, deletions or combinations thereof. In certain embodiments, the V α domain comprises or consists of the amino acid sequence set forth in any one of SEQ ID NOS:1-4. In certain embodiments, the V β domain comprises or consists of the amino acid sequence set forth in any one of SEQ ID NOS:5-7.

In particular embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:1 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:5.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:1 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:6.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:1 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:7.

In particular embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:2 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:5.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:2 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:6.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:2 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:7.

In particular embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:3 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:5.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:3 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:6.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:3 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:7.

In particular embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:4 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:5.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:4 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:6.

In other embodiments, the V α domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:4 and the V β domain comprises or consists of the amino acid sequence set forth in SEQ ID NO:7.

In further embodiments, a BRAF^{V600E}-specific binding protein is a TCR, an antigen-binding fragment of a TCR, or a chimeric antigen receptor. A "chimeric antigen receptor" (also called a CAR) is a fusion protein comprising an antigen binding domain (e.g., obtained or derived from an immunoglobulin or immunoglobulin-like molecule, such as an scFv derived from an antibody or TCR specific for a cancer antigen, or an antigen-binding domain obtained or derived from a killer immunoreceptor from an NK cell) linked to a transmembrane domain and one or more intracellular signaling domains (optionally containing co-stimulatory domain(s)) (see, e.g., Sadelain et al., *Cancer Discov.*, 3(4): 388-398, 2013; see also Harris and Kranz, *Trends Pharmacol. Sci.*, 37(3): 220-230, 2016; Stone et al., *Cancer Immunol. Immunother.*, 63(11):1163-1176, 2014). In certain embodiments, a binding protein comprises a CAR comprising a BRAF^{V600E}-specific TCR binding domain (see, e.g., Walseng et al., *Scientific Reports* 7:10713, 2017; the TCR CAR constructs and methods of which are hereby incorporated by reference in their entirety). Methods of making CARs are described, for example, in U.S. Pat. Nos. 6,410, 319; 7,446,191; U.S. Patent Publication No. 2010/065818; U.S. Pat. No. 8,822,647; PCT Publication No. WO 2014/031687; U.S. Pat. No. 7,514,537; and Brentjens et al., *Clin. Cancer Res.* 13:5426, 2007.

In certain embodiments, the antigen-binding fragment of the TCR comprises a single chain TCR (scTCR), which comprises both the TCR V α and V β domains TCR, but only a single TCR constant domain (C α or C β). In certain embodiments, the antigen-binding fragment of the TCR, or chimeric antigen receptor is chimeric (e.g., comprises amino acid residues or motifs from more than one donor or species), humanized (e.g., comprises residues from a non-human organism that are altered or substituted so as to reduce the risk of immunogenicity in a human), or human. Binding proteins according to the present disclosure, e.g., TCRs, may further comprise a TCR constant domain, e.g., joined to the C-terminus of a V α domain, a V β domain, or both. A TCR β -chain constant domain may be encoded by a TRBC1 gene or TRBC2 gene, and a TCR α -chain may be encoded by a TRAC gene. In certain embodiments, the TCR comprises an α chain constant (C α) domain having at least 90% (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or at least about 99.9%) sequence identity to the amino acid sequence set forth in SEQ ID NO:25. In a particular embodiment, the C α domain comprises the amino acid sequence set forth in SEQ ID NO:25. In certain embodiments, the TCR comprises a β chain (C β) constant domain having at least 90% (e.g., at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or at least about 99.9%) sequence identity to the amino acid sequence set forth in SEQ ID NO:26. In a particular embodiment, C β domain comprises the amino acid sequence set forth in SEQ ID NO:26.

In certain embodiments, a binding protein comprises a α -chain (comprised of a V α domain and a C α domain) comprising the amino acid sequence set forth in any one of

SEQ ID NOS:55-58. In certain embodiments, a binding protein comprises a β -chain (comprised of a V β domain and a C β domain) comprising the amino acid sequence set forth in any one of SEQ ID NOS:59-61.

Methods useful for isolating and purifying recombinantly produced soluble binding proteins (e.g., TCRs), by way of example, may include obtaining supernatants from suitable host cell/vector systems that secrete the recombinant soluble TCR into culture media and then concentrating the media using a commercially available filter. Following concentration, the concentrate may be applied to a single suitable purification matrix or to a series of suitable matrices, such as an affinity matrix or an ion exchange resin. One or more reverse phase HPLC steps may be employed to further purify a recombinant polypeptide. These purification methods may also be employed when isolating an immunogen from its natural environment. Methods for large scale production of one or more of the isolated/recombinant soluble TCR described herein include batch cell culture, which is monitored and controlled to maintain appropriate culture conditions. Purification of the soluble TCR may be performed according to methods described herein and known in the art and that comport with laws and guidelines of domestic and foreign regulatory agencies.

In certain embodiments, nucleic acid molecules encoding a binding protein (e.g., a TCR) specific for a BRAF^{V600E} peptide:HLA complex) are used to transfect/transduce a host cell (e.g., a T cell) for use in adoptive transfer therapy. Advances in TCR sequencing have been described (e.g., Robins et al., *Blood* 114:4099, 2009; Robins et al., *Sci. Translat. Med.* 2:47ra64, 2010; Robins et al., (September 10) *J. Imm. Meth.* Epub ahead of print, 2011; Warren et al., *Genome Res.* 21:790, 2011) and may be employed in the course of practicing embodiments according to the present disclosure. Similarly, methods for transfecting/transducing T cells with desired nucleic acids have been described (e.g., U.S. Patent Application Pub. No. US 2004/0087025) as have adoptive transfer procedures using T cells of desired antigen-specificity (e.g., Schmitt et al., *Hum. Gen.* 20:1240, 2009; Dossett et al., *Mol. Ther.* 17:742, 2009; Till et al., *Blood* 112:2261, 2008; Wang et al., *Hum. Gene Ther.* 18:712, 2007; Kuball et al., *Blood* 109:2331, 2007; US 2011/0243972; US 2011/0189141; Leen et al., *Ann. Rev. Immunol.* 25:243, 2007), such that adaptation of these methodologies to the presently disclosed embodiments is contemplated, based on the teachings herein, including those directed to TCRs specific for BRAF^{V600E} peptide antigens complexed with an HLA receptor.

The BRAF^{V600E}-specific binding proteins or domains as described herein may be functionally characterized according to any of a large number of art-accepted methodologies for assaying T cell activity, including determination of T cell binding, activation or induction and also including determination of T cell responses that are antigen-specific. Examples include determination of T cell proliferation, T cell cytokine release, antigen-specific T cell stimulation, MHC restricted T cell stimulation, CTL activity (e.g., by detecting ⁵¹Cr release from pre-loaded target cells), changes in T cell phenotypic marker expression, and other measures of T-cell functions. Procedures for performing these and similar assays are may be found, for example, in Lefkovits (*Immunology Methods Manual: The Comprehensive Sourcebook of Techniques*, 1998). See, also, *Current Protocols in Immunology*; Weir, *Handbook of Experimental Immunology*, Blackwell Scientific, Boston, MA (1986); Mishell and Shigii (eds.) *Selected Methods in Cellular Immunology*, Freeman

Publishing, San Francisco, CA (1979); Green and Reed, *Science* 281:1309 (1998) and references cited therein.

"MHC-peptide tetramer staining" refers to an assay used to detect antigen-specific T cells, which features a tetramer of MHC molecules, each comprising an identical peptide having an amino acid sequence that is cognate (e.g., identical or related to) at least one antigen (e.g., BRAF^{V600E}), wherein the complex is capable of binding T cell receptors specific for the cognate antigen. Each of the MHC molecules may be tagged with a biotin molecule. Biotinylated MHC/peptides are tetramerized by the addition of streptavidin, which can be fluorescently labeled. The tetramer may be detected by flow cytometry via the fluorescent label. In certain embodiments, an MHC-peptide tetramer assay is used to detect or select enhanced affinity TCRs of the instant disclosure.

Levels of cytokines may be determined according to methods described herein and practiced in the art, including for example, ELISA, ELISPOT, intracellular cytokine staining, and flow cytometry and combinations thereof (e.g., intracellular cytokine staining and flow cytometry). Immune cell proliferation and clonal expansion resulting from an antigen-specific elicitation or stimulation of an immune response may be determined by isolating lymphocytes, such as circulating lymphocytes in samples of peripheral blood cells or cells from lymph nodes, stimulating the cells with antigen, and measuring cytokine production, cell proliferation and/or cell viability, such as by incorporation of tritiated thymidine or non-radioactive assays, such as MTT assays and the like. The effect of an immunogen described herein on the balance between a Th1 immune response and a Th2 immune response may be examined, for example, by determining levels of Th1 cytokines, such as IFN- γ , IL-12, IL-2, and TNF- β , and Type 2 cytokines, such as IL-4, IL-5, IL-9, IL-10, and IL-13.

In further aspects, the present disclosure provides compositions comprising a binding protein according to the present disclosure and a pharmaceutically acceptable carrier, diluent, or excipient. Pharmaceutically acceptable excipients are biologically compatible vehicles, e.g., physiological saline, which are described in greater detail herein, that are suitable for administration to a human or other non-human mammalian subject.

Antigen presentation by immune cells (e.g., dendritic cells, phagocytes, and B cells) is determined in part by the HLA complexes present on the cells. Without wishing to be bound by theory, it is believed that HLA proteins encoded by different HLA alleles can vary in their ability to present particular antigen peptides and interact with immune cell proteins (e.g., TCRs). For example, a given antigen peptide may be presented by HLA-DQ complexes, but not HLA-DR complexes, or vice versa. In certain embodiments, a binding protein according to the present disclosure is capable of recognizing a BRAF^{V600E} peptide complexed with HLA-DQ. In certain embodiments, the HLA complex comprises HLA-DQB1*0301, *0302, or *0303. In certain embodiments, the HLA complex comprises HLA-DQB1*0302. In further embodiments, the HLA complex comprises HLA-DQA1*03.

Peptide antigens targeted by binding proteins according to the present disclosure can also vary on size depending on, for example, the type of HLA molecule presenting the antigen. Generally, HLA Class I complexes present peptides that are about 8-10 amino acids length, while HLA Class II complexes present peptides that are about 15-24 amino acids in length, though the peptides may be shorter or longer than these general lengths. Accordingly, in certain embodiments,

a BRAF^{V600E} peptide specifically bound by a binding protein of the present disclosure comprises from about 7 to about 27 amino acids, from about 10 to about 25 amino acids, or from about 12 to about 20 amino acids, or from about 15 to about 19 amino acids. In particular embodiments, the BRAF^{V600E} peptide comprises the amino acid sequence set forth in SEQ ID NO:38 or 39.

Polynucleotides Encoding BRAF^{V600E}-Specific Binding Proteins and Related Vectors

In yet further aspects, isolated polynucleotides and expression vectors that encode binding proteins according to the present disclosure are provided. Construction of an expression vector that is used for genetically engineering and producing a binding protein or TCR specific for a BRAF^{V600E} peptide of interest can be accomplished by using any suitable molecular biology engineering techniques known in the art. To obtain efficient transcription and translation, a polynucleotide in each recombinant expression construct includes at least one appropriate expression control sequence (also called a regulatory sequence), such as a leader sequence and particularly a promoter operably (i.e., operatively) linked to the nucleotide sequence encoding the immunogen.

Certain embodiments relate to polynucleotides that encode the binding proteins provided herein, such as binding proteins (e.g., TCRs or CARs) specific for a BRAF^{V600E} peptide:HLA complex. A nucleic acid may be a single- or a double-stranded DNA, cDNA or RNA in any form, and may include a positive and a negative strand of the nucleic acid which complement each other, including anti-sense DNA, cDNA and RNA. Also included are siRNA, microRNA, RNA-DNA hybrids, ribozymes, and other various naturally occurring or synthetic forms of DNA or RNA. It will be appreciated that a polynucleotide of the present disclosure can vary (i.e., comprise a different nucleotide sequence) as compared to a reference polynucleotide sequence disclosed herein and still encode a same amino acid or polypeptide, due to, for example, the degeneracy of the genetic code. In certain embodiments, a polynucleotide encoding, for example, a binding protein or a portion thereof, a self-cleaving peptide, a linker peptide; or a binding protein-encoding construct, may have at least about 80% (e.g., at least about 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, or at least about 99.9%) identity to a polynucleotide according to any one of SEQ ID NOs:8-24; 27; 28; 44-48; 62-68; and 73-77.

In any of the aforementioned embodiments, a polynucleotide encoding a binding protein of the present disclosure is codon optimized for efficient expression in a target host cell.

Certain embodiments include polynucleotides of this disclosure contained in a vector. An exemplary vector may comprise a polynucleotide capable of transporting another polynucleotide to which it has been linked, or which is capable of replication in a host organism. Some examples of vectors include plasmids, viral vectors, cosmids, and others. Some vectors may be capable of autonomous replication in a host cell into which they are introduced (e.g. bacterial vectors having a bacterial origin of replication and episomal mammalian vectors), whereas other vectors may be integrated into the genome of a host cell or promote integration of the polynucleotide insert upon introduction into the host cell and thereby replicate along with the host genome (e.g., lentiviral vector, retroviral vector). Additionally, some vectors are capable of directing the expression of genes to which they are operatively linked (these vectors may be referred to

as "expression vectors"). According to related embodiments, it is further understood that, if one or more agents (e.g., polynucleotides encoding binding proteins or recombinant TCRs specific for BRAF^{V600E}, or variants thereof, as described herein) are co-administered to a subject, that each agent may reside in separate or the same vectors, and multiple vectors (each containing a different agent or the same agent) may be introduced to a cell or cell population or administered to a subject.

In certain embodiments, polynucleotides encoding binding proteins specific for a BRAF^{V600E} peptide:HLA complex may be operatively linked to certain elements of a vector. For example, polynucleotide sequences that are needed to effect the expression and processing of coding sequences to which they are ligated may be operatively linked. Expression control sequences may include appropriate transcription initiation, termination, promoter and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequences); sequences that enhance protein stability; and possibly sequences that enhance protein secretion. Expression control sequences may be operatively linked if they are contiguous with the gene of interest and expression control sequences that act in trans or at a distance to control the gene of interest. In certain embodiments, polynucleotides encoding binding proteins of the instant disclosure are contained in an expression vector that is a viral vector, such as a lentiviral vector or a γ -retroviral vector.

In certain embodiments, expression vectors are provided comprising a polynucleotide encoding a binding protein of the present disclosure, wherein the polynucleotide is operably linked to an expression control sequence (e.g., a promoter). In certain embodiments, the vector is capable of delivering the polynucleotide to a host cell. In certain embodiments, the host cell is a hematopoietic progenitor cell or a human immune system cell. In further embodiments, the immune system cell is a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁺ CD8⁺ double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a dendritic cell, or any combination thereof. In certain embodiments, the immune system cell is a CD4⁺ T cell. In certain embodiments, the T cell is a naïve T cell, a central memory T cell, an effector memory T cell, or any combination thereof.

In any of the embodiments herein, the vector is a viral vector. In certain embodiments, the viral vector is a lentiviral vector or a γ -retroviral vector.

Host Cells

In still further aspects, host cells are provided that comprise a heterologous polynucleotide according to the present disclosure, wherein the host cell expresses on its cell surface a binding protein encoded by the heterologous polynucleotide (i.e., expresses a binding protein according to the present disclosure). In particular embodiments, an expression vector is delivered to an appropriate cell, for example, a T cell or an antigen-presenting cell, i.e., a cell that displays a peptide/MHC complex on its cell surface (e.g., a dendritic cell). In certain embodiments, a host cell (e.g., a T cell, NK cell, or NK-T cell) lacks a CD8 co-receptor or a CD4 co-receptor and the encoded binding protein is capable of binding a BRAF^{V600E} antigen:HLA complex in the absence of a CD4 or CD8 co-receptor. In certain embodiments, the host cell is a hematopoietic progenitor cell or a human immune system cell. For example, the immune system cell can be a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁺ CD8⁺ double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a natural

killer T cell, a dendritic cell, or any combination thereof. In some embodiments, the encoded binding protein comprises a MHCII-restricted TCR binding domain and the host cell (e.g., a CD8⁺ T cell) comprises a polynucleotide encoding a heterologous CD4⁺ co-receptor.

In certain embodiments, wherein a T cell is the host, the T cell can be naïve, a central memory T cell, an effector memory T cell, a stem cell memory T cell, or any combination thereof. In certain embodiments, the T cell is a CD4⁺ T cell, a CD8⁺ T cell, or both. The expression vectors introduced into the host cells may also include, for example, lymphoid tissue-specific transcriptional regulatory elements (TREs), such as a B lymphocyte, T lymphocyte, or dendritic cell specific TREs. Lymphoid tissue specific TREs are known in the art (see, e.g., Thompson et al., *Mol. Cell. Biol.* 12:1043, 1992; Todd et al., *J. Exp. Med.* 177:1663, 1993; Penix et al., *J. Exp. Med.* 178:1483, 1993).

A host cell may include any individual cell or cell culture which may receive a vector or the incorporation of nucleic acids or express proteins. The term also encompasses progeny of the host cell, whether genetically or phenotypically the same or different. Suitable host cells may depend on the vector and may include mammalian cells, animal cells, human cells, simian cells, insect cells, yeast cells, and bacterial cells. These cells may be induced to incorporate the vector or other material by use of a viral vector, transformation via calcium phosphate precipitation, DEAE-dextran, electroporation, microinjection, or other methods. See, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual* 2d ed. (Cold Spring Harbor Laboratory, 1989).

Accordingly, in one aspect, a host cell is provided that comprises a heterologous polynucleotide or an expression vector according to the present disclosure, wherein the host cell expresses on its cell surface a binding protein encoded by the heterologous polynucleotide. In certain embodiments, the portion of the heterologous polynucleotide that encodes the V α domain is at least about 80% identical to the polynucleotide sequence set forth in any one of SEQ ID NOS:18-21. In certain embodiments, the portion of the heterologous polynucleotide that encodes the V β domain is at least about 80% identical to the polynucleotide sequence set forth in any one of SEQ ID NOS:22-24.

In certain embodiments, the portion of the heterologous polynucleotide that encodes the V α domain comprises or consists of the polynucleotide sequence set forth in any one of SEQ ID NOS:18-21. In certain embodiments, the portion of the heterologous polynucleotide that encodes the V β domain comprises or consists of the polynucleotide sequence set forth in any one of SEQ ID NOS:22-24.

In certain embodiments, the portion of the heterologous polynucleotide that encodes the V α domain comprises or consists of the polynucleotide sequence set forth in any one of SEQ ID NOS:18-21, and the portion of the heterologous polynucleotide that encodes the V β domain comprises or consists of the polynucleotide sequence set forth in any one of SEQ ID NOS:22-24.

In certain embodiments, the portion of the heterologous polynucleotide that encodes the V α domain is linked to a portion that encodes a TCR α -chain constant domain, wherein the portion that encodes the α -chain constant domain comprises or consists of a sequence that is at least about 80% identical to the polynucleotide sequence set forth in SEQ ID NO:27.

In certain embodiments, the portion of the heterologous polynucleotide that encodes the V β domain is linked to a portion that encodes a TCR β -chain constant domain, wherein the portion that encodes the β -chain constant

domain comprises or consists of a sequence that is at least about 80% identical to the polynucleotide sequence set forth in SEQ ID NO:28.

In particular embodiments, the portion of the polynucleotide that encodes the V_{α} domain comprises or consists of SEQ ID NO:18 and the portion that encodes the V_{β} domain comprises or consists of SEQ ID NO:22.

In other embodiments, the portion of the polynucleotide that encodes the V_{α} domain comprises or consists of SEQ ID NO:19 and the portion that encodes the V_{β} domain comprises or consists of SEQ ID NO:23.

In other embodiments, the portion of the polynucleotide that encodes the V_{α} domain comprises or consists of SEQ ID NO:20 and the portion that encodes the V_{β} domain comprises or consists of SEQ ID NO:23.

In still other embodiments, the portion of the polynucleotide that encodes the V_{α} domain comprises or consists of SEQ ID NO:21 and the portion that encodes the V_{β} domain comprises or consists of SEQ ID NO:24.

In any of the herein described embodiments, a host cell (e.g., a T cell) expressing a BRAF^{V600E}-specific binding protein of the present disclosure is capable of producing an interferon when co-cultured with an antigen-presenting cell presenting or expressing a BRAF^{V600E}-containing antigen. In certain embodiments, the produced interferon comprises interferon-gamma (IFN- γ). In some embodiments, the target cell has been pulsed with a peptide or polypeptide comprising or consisting of the BRAF^{V600E}-containing antigen. In some embodiments, the target cell has been transfected with a polynucleotide (e.g., DNA, cDNA, or mRNA) encoding a polypeptide or peptide comprising or consisting of the BRAF^{V600E}-containing antigen. In certain embodiments, a host cell of the present disclosure produces IFN- γ when cultured with an antigen-presenting cell that has been pulsed with a BRAF^{V600E}-containing antigen at a concentration of about 0.005 μ g/mL to about 10 μ g/mL antigen. In further embodiments, the host cell produces at least about 1,000 pg/mL IFN- γ when cultured with an antigen-presenting cell that has been pulsed with a BRAF^{V600E}-containing antigen at a concentration of at least about 0.1 μ g/mL antigen. In some embodiments, the host cell produces at least about 1,000 pg/mL IFN- γ when cultured with an antigen-presenting cell pulsed with a BRAF^{V600E}-containing antigen at a concentration of at least about 0.1, 0.2, 0.3, 0.4, or 0.5 μ g/mL antigen. In some embodiments, the host cell produces from about 1,000 pg/mL to about 10,000 pg/mL IFN- γ when cultured with an antigen-presenting cell pulsed with a BRAF^{V600E}-containing antigen at a concentration of about 0.5 μ g/mL antigen to about 10 μ g/mL antigen. In some embodiments, a target cell comprises a B cell. In further embodiments, the B cell expresses an HLA-DQ allele. In still further embodiments, the B cell is of B-LCL line 1331.

In certain embodiments, a portion of the polynucleotide encodes a self-cleaving peptide and is disposed between a TCR α -chain-encoding portion and a TCR β -chain-encoding portion. Self-cleaving peptides useful for expression of separable polypeptides by a single vector are known in the art and include, for example, Porcine teschovirus-1 2A (P2A) peptide, Thoseaasigna virus 2A (T2A) peptide, Equine rhinitis A virus (ERAV) 2A (E2A) peptide, and Foot-and-mouth disease virus 2A (F2A) peptide. Accordingly, in certain embodiments, the portion of the heterologous polynucleotide that encodes the self-cleaving peptide comprises or consists of the polynucleotide sequence set forth in any one of SEQ ID NOS:44-48. In further embodi-

ments, the encoded self-cleaving peptide comprises or consists of the amino acid sequence set forth in any one of SEQ ID NOS:49-52.

In certain embodiments, the host cell is a hematopoietic progenitor cell or a human immune system cell. In certain embodiments, the immune system cell is a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁻ CD8⁻ double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a dendritic cell, or any combination thereof. In certain embodiments, the immune system cell is a T cell. In particular embodiments, T cell is a naïve T cell, a central memory T cell, an effector memory T cell, or any combination thereof. In further embodiments, the T cell is a CD4⁺ T cell.

In certain embodiments, the binding protein or TCR expressed by the T cell is capable of more efficiently associating with a CD3 protein, a CD4 protein, or both, as compared to endogenous TCR. In certain embodiments, the binding protein or TCR higher surface expression on a T cell as compared to endogenous TCR.

In any of the foregoing embodiments, a host cell that comprises a heterologous polynucleotide encoding a BRAF^{V600E}-specific binding protein is an immune cell which is modified to reduce or eliminate expression of one or more endogenous genes that encode a polypeptide product selected from PD-1, LAG-3, CTLA4, TIM3, TIGIT, an HLA molecule, a TCR molecule, or any component or combination thereof. Without wishing to be bound by theory, certain endogenously expressed immune cell proteins may downregulate the immune activity of a modified immune host cell (e.g., PD-1, LAG-3, CTLA4, TIGIT), or may compete with a heterologous binding protein of the present disclosure for expression by the host cell, association with TCR complex components (e.g., CD3 proteins), or may interfere with the binding activity of a heterologously expressed binding protein of the present disclosure (e.g., an endogenous TCR that binds to a non-BRAF^{V600E} antigen or a non-BRAF^{V600E} antigen:HLA complex and interferes with binding of a presently disclosed binding protein to a BRAF^{V600E} antigen) and interferes with the immune host cell binding a target cell that expresses BRAF^{V600E} antigen), or any combination thereof. Further, endogenous proteins (e.g., immune host cell proteins, such as an HLA) expressed on a donor immune cell to be used in a cell transfer therapy may be recognized as foreign by an allogeneic recipient, which may result in elimination or suppression of the donor immune cell by the allogeneic recipient.

Accordingly, decreasing or eliminating expression or activity of such endogenous genes or proteins can improve the activity, tolerance, and persistence of the host cells in an autologous or allogeneic host setting, and allows universal administration of the cells (e.g., to any recipient regardless of HLA type). In certain embodiments, a modified host immune cell is a donor cell (e.g., allogeneic) or an autologous cell. In certain embodiments, a modified immune host cell of this disclosure comprises a chromosomal gene knockout of one or more of a gene that encodes PD-1, LAG-3, CTLA4, TIM3, TIGIT, an HLA component (e.g., a gene that encodes an α 1 macroglobulin, an α 2 macroglobulin, an α 3 macroglobulin, a β 1 macroglobulin, or a β 2 macroglobulin), or a TCR component (e.g., a gene that encodes a TCR variable region or a TCR constant region) (see, e.g., Torikai et al., *Nature Sci. Rep.* 6:21757 (2016); Torikai et al., *Blood* 119(24):5697 (2012); and Torikai et al., *Blood* 122(8):1341 (2013) the gene editing techniques, compositions, and adoptive cell therapies of which are herein incorporated by reference in their entirety; e.g., SEQ ID NOS:142-149). As used herein, the term "chromosomal gene knockout" refers

to a genetic alteration in a host cell that prevents production, by the host cell, of a functionally active endogenous polypeptide product. Alterations resulting in a chromosomal gene knockout can include, for example, introduced non-sense mutations (including the formation of premature stop codons), missense mutations, gene deletion, and strand breaks, as well as the heterologous expression of inhibitory nucleic acid molecules that inhibit endogenous gene expression in the host cell.

In certain embodiments, a chromosomal gene knock-out or gene knock-in is made by chromosomal editing of a host cell. Chromosomal editing can be performed using, for example, endonucleases. As used herein “endonuclease” refers to an enzyme capable of catalyzing cleavage of a phosphodiester bond within a polynucleotide chain. In certain embodiments, an endonuclease is capable of cleaving a targeted gene thereby inactivating or “knocking out” the targeted gene. An endonuclease may be a naturally occurring, recombinant, genetically modified, or fusion endonuclease. The nucleic acid strand breaks caused by the endonuclease are commonly repaired through the distinct mechanisms of homologous recombination or non-homologous end joining (NHEJ). During homologous recombination, a donor nucleic acid molecule may be used for a donor gene “knock-in”, for target gene “knock-out”, and optionally to inactivate a target gene through a donor gene knock in or target gene knock out event. NHEJ is an error-prone repair process that often results in changes to the DNA sequence at the site of the cleavage, e.g., a substitution, deletion, or addition of at least one nucleotide. NHEJ may be used to “knock-out” a target gene. Examples of endonucleases include zinc finger nucleases, TALE-nucleases, CRISPR-Cas nucleases, meganucleases, and megaTALs.

As used herein, a “zinc finger nuclease” (ZFN) refers to a fusion protein comprising a zinc finger DNA-binding domain fused to a non-specific DNA cleavage domain, such as a FokI endonuclease. Each zinc finger motif of about 30 amino acids binds to about 3 base pairs of DNA, and amino acids at certain residues can be changed to alter triplet sequence specificity (see, e.g., Desjarlais et al., *Proc. Natl. Acad. Sci.* 90:2256-2260, 1993; Wolfe et al., *J. Mol. Biol.* 285:1917-1934, 1999). Multiple zinc finger motifs can be linked in tandem to create binding specificity to desired DNA sequences, such as regions having a length ranging from about 9 to about 18 base pairs. By way of background, ZFNs mediate genome editing by catalyzing the formation of a site-specific DNA double strand break (DSB) in the genome, and targeted integration of a transgene comprising flanking sequences homologous to the genome at the site of DSB is facilitated by homology directed repair. Alternatively, a DSB generated by a ZFN can result in knock out of target gene via repair by non-homologous end joining (NHEJ), which is an error-prone cellular repair pathway that results in the insertion or deletion of nucleotides at the cleavage site. In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, made using a ZFN molecule.

As used herein, a “transcription activator-like effector nuclease” (TALEN) refers to a fusion protein comprising a TALE DNA-binding domain and a DNA cleavage domain, such as a FokI endonuclease. A “TALE DNA binding domain” or “TALE” is composed of one or more TALE repeat domains/units, each generally having a highly conserved 33-35 amino acid sequence with divergent 12th and 13th amino acids. The TALE repeat domains are involved in binding of the TALE to a target DNA sequence. The divergent amino acid residues, referred to as the Repeat Variable

Diresidue (RVD), correlate with specific nucleotide recognition. The natural (canonical) code for DNA recognition of these TALEs has been determined such that an HD (histidine-aspartic acid) sequence at positions 12 and 13 of the TALE leads to the TALE binding to cytosine (C), NG (asparagine-glycine) binds to a T nucleotide, NI (asparagine-isoleucine) to A, NN (asparagine-asparagine) binds to a G or A nucleotide, and NG (asparagine-glycine) binds to a T nucleotide. Non-canonical (atypical) RVDs are also known (see, e.g., U.S. Patent Publication No. US 2011/0301073, which atypical RVDs are incorporated by reference herein in their entirety). TALENs can be used to direct site-specific double-strand breaks (DSB) in the genome of T cells. Non-homologous end joining (NHEJ) ligates DNA from both sides of a double-strand break in which there is little or no sequence overlap for annealing, thereby introducing errors that knock out gene expression. Alternatively, homology directed repair can introduce a transgene at the site of DSB providing homologous flanking sequences are present in the transgene. In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, and made using a TALEN molecule.

As used herein, a “clustered regularly interspaced short palindromic repeats/Cas” (CRISPR/Cas) nuclease system refers to a system that employs a CRISPR RNA (crRNA)-guided Cas nuclease to recognize target sites within a genome (known as protospacers) via base-pairing complementarity and then to cleave the DNA if a short, conserved protospacer associated motif (PAM) immediately follows 3' of the complementary target sequence. CRISPR/Cas systems are classified into three types (i.e., type I, type II, and type III) based on the sequence and structure of the Cas nucleases. The crRNA-guided surveillance complexes in types I and III need multiple Cas subunits. Type II system, the most studied, comprises at least three components: an RNA-guided Cas9 nuclease, a crRNA, and a trans-acting crRNA (tracrRNA). The tracrRNA comprises a duplex forming region. A crRNA and a tracrRNA form a duplex that is capable of interacting with a Cas9 nuclease and guiding the Cas9/crRNA:tracrRNA complex to a specific site on the target DNA via Watson-Crick base-pairing between the spacer on the crRNA and the protospacer on the target DNA upstream from a PAM. Cas9 nuclease cleaves a double-stranded break within a region defined by the crRNA spacer. Repair by NHEJ results in insertions and/or deletions which disrupt expression of the targeted locus. Alternatively, a transgene with homologous flanking sequences can be introduced at the site of DSB via homology directed repair. The crRNA and tracrRNA can be engineered into a single guide RNA (sgRNA or gRNA) (see, e.g., Jinek et al., *Science* 337:816-21, 2012). Further, the region of the guide RNA complementary to the target site can be altered or programmed to target a desired sequence (Xie et al., *PLOS One* 9:e100448, 2014; U.S. Pat. Appl. Pub. No. US 2014/0068797, U.S. Pat. Appl. Pub. No. US 2014/0186843; U.S. Pat. No. 8,697,359, and PCT Publication No. WO 2015/071474; each of which is incorporated by reference). In certain embodiments, a gene knockout comprises an insertion, a deletion, a mutation or a combination thereof, and made using a CRISPR/Cas nuclease system.

Exemplary gRNA sequences and methods of using the same to knock out endogenous genes that encode immune cell proteins include those described in Ren et al., *Clin. Cancer Res.* 23(9):2255-2266 (2017), the gRNAs, CAS9 DNAs, vectors, and gene knockout techniques of which are hereby incorporated by reference in their entirety.

In some embodiments, a gene knockout comprises a CRISPR-mediated gene knockout of a TCR α -chain constant region locus (C α), a TCR β -chain constant region locus (C β), or both. In certain embodiments, a gRNA sequence targeting a TCR C α locus comprises the nucleotide sequence AGAGTCTCTCAGCTGGTACA (SEQ ID NO:136). In certain embodiments, a gRNA sequence targeting a TCR C α locus comprises the nucleotide sequence TGTGCTAGACATGAGGTCTA (SEQ ID NO:137). In certain embodiments, a gRNA sequence targeting a TCR C β locus comprises the nucleotide sequence GCAGTATCTG-GAGTCATTGA (SEQ ID NO:138). In certain embodiments, a gRNA sequence targeting a TCR C β locus comprises the nucleotide sequence GGAGAATGACGAGTGGACCC (SEQ ID NO:139). In some embodiments, a gene knockout comprises a CRISPR-mediated gene knockout of a human β 2M locus. In certain embodiments, a gRNA sequence targeting a human β 2M locus comprises the nucleotide sequence CGCGAGCACAGCTAAGGCCA (SEQ ID NO:140). In some embodiments, a gene knockout comprises a CRISPR-mediated gene knockout of a PD-1 locus. In certain embodiments, a gRNA sequence targeting a PD-1 comprises the nucleotide sequence GGCCAGGATGGTCTTAGGT (SEQ ID NO:141).

As used herein, a "meganuclease," also referred to as a "homing endonuclease," refers to an endodeoxyribonuclease characterized by a large recognition site (double stranded DNA sequences of about 12 to about 40 base pairs). Meganucleases can be divided into five families based on sequence and structure motifs: LAGLIDADG, GIY-YIG, HNH, His-Cys box and PD-(D/E)XK. Exemplary meganucleases include I-SceI, I-CeuI, PI-PspI, PI-Sce, I-SceIV, I-CsmI, I-PanI, I-SceII, I-PpoI, I-SceIII, I-CreI, I-TevI, I-TevII and I-TevIII, whose recognition sequences are known (see, e.g., U.S. Pat. Nos. 5,420,032 and 6,833,252; Belfort et al., *Nucleic Acids Res.* 25:3379-3388, 1997; Dujon et al., *Gene* 82:115-118, 1989; Perler et al., *Nucleic Acids Res.* 22:1125-1127, 1994; Jasin, *Trends Genet.* 12:224-228, 1996; Gimble et al., *J. Mol. Biol.* 263:163-180, 1996; Argast et al., *J. Mol. Biol.* 280:345-353, 1998).

In certain embodiments, naturally-occurring meganucleases may be used to promote site-specific genome modification of a target selected from PD-1, LAG3, TIM3, CTLA4, TIGIT, an HLA-encoding gene, or a TCR component-encoding gene. In other embodiments, an engineered meganuclease having a novel binding specificity for a target gene is used for site-specific genome modification (see, e.g., Porteus et al., *Nat. Biotechnol.* 23:967-73, 2005; Sussman et al., *J. Mol. Biol.* 342:31-41, 2004; Epinat et al., *Nucleic Acids Res.* 31:2952-62, 2003; Chevalier et al., *Molec. Cell* 10:895-905, 2002; Ashworth et al., *Nature* 441:656-659, 2006; Paques et al., *Curr. Gene Ther.* 7:49-66, 2007; U.S. Patent Publication Nos. US 2007/0117128; US 2006/0206949; US 2006/0153826; US 2006/0078552; and US 2004/0002092). In further embodiments, a chromosomal gene knockout is generated using a homing endonuclease that has been modified with modular DNA binding domains of TALENs to make a fusion protein known as a megaTAL. MegaTALs can be utilized to not only knock-out one or more target genes, but to also introduce (knock in) heterologous or exogenous polynucleotides when used in combination with an exogenous donor template encoding a polypeptide of interest, such as a TCR α chain, TCR β chain or both, wherein the knocked-in TCR produced by the cell is specific for a BRAF^{V600E} antigen or peptide.

In certain embodiments, a chromosomal gene knockout comprises an inhibitory nucleic acid molecule that is introduced into a host cell (e.g., an immune cell) comprising a heterologous polynucleotide encoding an antigen-specific receptor that specifically binds to a tumor associated antigen, wherein the inhibitory nucleic acid molecule encodes a target-specific inhibitor and wherein the encoded target-specific inhibitor inhibits endogenous gene expression (i.e., of PD-1, TIM3, LAG3, CTLA4, TIGIT, an HLA component, or a TCR component, or any combination thereof) in the host immune cell.

A chromosomal gene knockout can be confirmed directly by DNA sequencing of the host immune cell following use of the knockout procedure or agent. Chromosomal gene knockouts can also be inferred from the absence of gene expression (e.g., the absence of an mRNA or polypeptide product encoded by the gene) following the knockout.

Methods of Treatment

In some aspects, methods of the instant disclosure are for treating a hyperproliferative disorder, wherein the methods comprise administering to human subject in need thereof a composition comprising a binding protein specific for human BRAF^{V600E} or a host cell according to the present disclosure.

In certain aspects, the instant disclosure is directed to methods for treating a hyperproliferative disorder or a condition characterized by BRAF^{V600E} expression by administering to human subject in need thereof a composition comprising a binding protein or a host cell expressing a binding protein specific for a BRAF^{V600E} peptide:HLA complex according to any the aforementioned binding proteins.

The presence of a hyperproliferative disorder or malignant condition in a subject refers to the presence of dysplastic, cancerous and/or transformed cells in the subject, including, for example neoplastic, tumor, non-contact inhibited or oncogenically transformed cells, or the like (e.g., solid cancers; hematologic cancers including lymphomas and leukemias, such as acute myeloid leukemia, chronic myeloid leukemia, etc. such as renal, gastric, ovarian, and colorectal cancers), which are known in the art and for which criteria for diagnosis and classification are established (e.g., Hanahan and Weinberg, *Cell* 144:646, 2011; Hanahan and Weinberg, *Cell* 100:57, 2000; Cavallo et al., *Canc. Immunol. Immunother.* 60:319, 2011; Kyrigideis et al., *J. Carcinog.* 9:3, 2010). In particular, for example, hairy cell leukemia, melanoma, non-small cell lung cancer, colorectal cancer, papillary cancer, and thyroid cancer, such as poorly differentiated thyroid cancer. Accordingly, in further embodiments, there are provided methods for treating a hyperproliferative disorder or other condition associated with BRAF^{V600E} expression, including hairy cell leukemia, melanoma, thyroid cancer such as poorly differentiated thyroid cancer, non-small cell lung cancer, colorectal cancer, papillary cancer, non-Hodgkin lymphoma, adenocarcinoma of the lung, and brain tumors including glioblastoma and pilocytic astrocytomas.

As used herein, the terms, "treat" and "treatment," refer to medical management of a disease, disorder, or condition of a subject (i.e., patient, host, who may be a human or non-human animal) (see, e.g., Stedman's Medical Dictionary). In general, an appropriate dose and treatment regimen provide one or more of a binding protein or a BRAF^{V600E} peptide:HLA complex or a host cell expressing the same, and optionally an adjunctive therapy (e.g., a cytokine such as IL-2, IL-15, IL-21 or any combination thereof), in an amount sufficient to provide therapeutic or prophylactic

benefit. Therapeutic or prophylactic benefit resulting from therapeutic treatment or prophylactic or preventative methods include, for example an improved clinical outcome, wherein the object is to prevent or retard or otherwise reduce (e.g., decrease in a statistically significant manner relative to an untreated control) an undesired physiological change or disorder, or to prevent, retard or otherwise reduce the expansion or severity of such a disease or disorder. Beneficial or desired clinical results from treating a subject include abatement, lessening, or alleviation of symptoms that result from or are associated the disease or disorder to be treated; decreased occurrence of symptoms; improved quality of life; longer disease-free status (i.e., decreasing the likelihood or the propensity that a subject will present symptoms on the basis of which a diagnosis of a disease is made); diminishment of extent of disease; stabilized (i.e., not worsening) state of disease; delay or slowing of disease progression; amelioration or palliation of the disease state; and remission (whether partial or total), whether detectable or undetectable; or overall survival.

"Treatment" can also mean prolonging survival when compared to expected survival if a subject were not receiving treatment. Subjects in need of the methods and compositions described herein include those who already have the disease or disorder, as well as subjects prone to have or at risk of developing the disease or disorder. Subjects in need of prophylactic treatment include subjects in whom the disease, condition, or disorder is to be prevented (i.e., decreasing the likelihood of occurrence or recurrence of the disease or disorder). The clinical benefit provided by the compositions (and preparations comprising the compositions) and methods described herein can be evaluated by design and execution of in vitro assays, preclinical studies, and clinical studies in subjects to whom administration of the compositions is intended to benefit, as described in the examples.

Cells expressing the binding protein as described herein may be administered to a subject in a pharmaceutically or physiologically acceptable or suitable excipient or carrier. Pharmaceutically acceptable excipients are biologically compatible vehicles, e.g., physiological saline, which are described in greater detail herein, that are suitable for administration to a human or other non-human mammalian subject.

A therapeutically effective dose, in the context of adoptive cell therapy, is an amount of host cells (expressing a binding protein according to the present disclosure) used in adoptive transfer that is capable of producing a clinically desirable result (i.e., a sufficient amount to induce or enhance a specific T cell immune response against cells expressing BRAF^{V600E} (e.g., a cytotoxic T cell response) in a statistically significant manner) in a treated human or non-human mammal. As is well known in the medical arts, the dosage for any one patient depends upon many factors, including the patient's size, weight, body surface area, age, the particular therapy to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Doses will vary, but a preferred dose for administration of a host cell comprising a recombinant expression vector as described herein is about 10⁷ cells/m², about 5×10⁷ cells/m², about 10⁸ cells/m², about 5×10⁸ cells/m², about 10⁹ cells/m², about 5×10⁹ cells/m², about 10¹⁰ cells/m², about 5×10¹⁰ cells/m², or about 10¹¹ cell s/m².

In any of the presently disclosed embodiments, a unit dose can comprise T cells, e.g., CD4⁺ T cells, CD8⁺ T cells, or both, wherein the T cells can comprise bulk T cells, naïve T cells, stem cell memory T cells, central memory T cells, or

effector memory T cells. In certain embodiments, a unit dose comprises BRAF^{V600E}-specific CD4⁺ T cells and does not comprise CD8⁺ T cells. In other embodiments, a unit dose comprises BRAF^{V600E}-specific CD8⁺ T cells, which may be engineered to express a heterologous CD4⁺ co-receptor, and optionally does not comprise CD4⁺ T cells. In certain embodiments, a unit dose comprises BRAF^{V600E}-specific CD4⁺ T host cells of the present disclosure and further comprises CD4⁺ T cells or CD8⁺ T cells (e.g., allogeneic or autologous, modified (e.g., to express a heterologous protein such as a TCR, a CAR, a CD4 co-receptor, a CD8 co-receptor, or any combination thereof) or unmodified) that have binding specificity for one or more other antigens or antigen-HLA complexes, such as, for example: a BRAF^{V600E} antigen, a BRAF antigen that does not comprise a V600E mutation; a BRAF^{V600E}-containing antigen that associates with a different HLA than does the binding protein of a presently disclosed host cell in the unit dose; or a different antigen that is associated with hyperproliferative disease or disorder; e.g., NY-ESO-1, SSX-2, Tyrosinase, TMG1-4, GP100, MAGE-A3, MART1, ROR1, EGFR, EGFRvIII, EGP-2, EGP-40, GD2, GD3, HPV E6, HPV E7, Her2, L1-CAM, Lewis A, Lewis Y, MUC1, MUC16, PSCA, PSMA, CD19, CD20, CD22, CD56, CD23, CD24, CD30, CD33, CD37, CD44v7/8, CD38, CD56, CD123, CA125, c-MET, FcRH5, WT-1, folate receptor α , VEGF- α , VEGFR1, VEGFR2, IL-13R α 2, IL-11R α , MAGE-A1, PSA, ephrin A2, ephrin B2, an NKGD2D, NY-ESO-1, TAG-72, mesothelin, NY-ESO, 5T4, BCMA, FAP, Carbonic anhydrase 9, ERBB2, or CEA, or any combination thereof).

In certain embodiments, a unit dose comprises (i) a composition comprising at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 85%, at least about 90%, or at least about 95% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells (i.e., has less than about 50%, less than about 40%, less than about 30%, less than about 20%, less than about 10%, less than about 5%, or less than about 1% the population of naïve T cells present in a unit dose as compared to a patient sample having a comparable number of PBMCs).

In some embodiments, a unit dose comprises (i) a composition comprising at least about 50% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 50% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In further embodiments, a unit dose comprises (i) a composition comprising at least about 60% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 60% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In still further embodiments, a unit dose comprises (i) a composition comprising at least about 70% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 70% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In some embodiments, a unit dose comprises (i) a composition comprising at least about 80% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 80% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or

substantially no naïve T cells. In some embodiments, a unit dose comprises (i) a composition comprising at least about 85% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 85% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells. In some embodiments, a unit dose comprises (i) a composition comprising at least about 90% engineered CD4⁺ T cells, combined with (ii) a composition comprising at least about 90% engineered CD8⁺ T cells, in about a 1:1 ratio, wherein the unit dose contains a reduced amount or substantially no naïve T cells.

In any of the embodiments described herein, a unit dose comprises equal, or approximately equal numbers of engineered CD45RA⁻ CD3⁺ CD8⁺ and engineered CD45RA⁻ CD3⁺ CD4⁺ T_M cells.

Pharmaceutical compositions may be administered in a manner appropriate to the disease or condition to be treated (or prevented) as determined by persons skilled in the medical art. An appropriate dose and a suitable duration and frequency of administration of the compositions will be determined by such factors as the health condition of the patient, size of the patient (i.e., weight, mass, or body area), the type and severity of the patient's disease, the particular form of the active ingredient, and the method of administration. In general, an appropriate dose and treatment regimen provide the composition(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit (such as described herein, including an improved clinical outcome, such as more frequent complete or partial remissions, or longer disease-free and/or overall survival, or a lessening of symptom severity). For prophylactic use, a dose should be sufficient to prevent, delay the onset of, or diminish the severity of a disease associated with disease or disorder. Prophylactic benefit of the immunogenic compositions administered according to the methods described herein can be determined by performing pre-clinical (including in vitro and in vivo animal studies) and clinical studies and analyzing data obtained therefrom by appropriate statistical, biological, and clinical methods and techniques, all of which can readily be practiced by a person skilled in the art.

A condition associated with BRAF^{V600E} expression includes any disorder or condition in which a BRAF^{V600E}-driven cellular or molecular event is present, and typically manifests in overgrowth of diseased cells relative to normal cells. Some conditions associated with BRAF^{V600E} expression may include acute as well as chronic disorders and diseases, such as those pathological conditions that predispose the subject to a particular disorder.

Some examples of conditions associated with BRAF^{V600E} expression include hyperproliferative disorders, which refer to states of activated and/or proliferating cells (which may also be transcriptionally overactive) in a subject including tumors, neoplasms, cancer, malignancy, etc. In addition to activated or proliferating cells, the hyperproliferative disorder may also include an aberration or dysregulation of cell death processes, whether by necrosis or apoptosis. Such aberration of cell death processes may be associated with a variety of conditions, including cancer (including primary, secondary malignancies as well as metastasis), or other conditions.

According to certain embodiments, a cancer that is characterized by BRAF^{V600E} expression may be treated through the use of compositions and methods disclosed herein. Furthermore, "cancer" may refer to any accelerated proliferation of cells, including solid tumors, ascites tumors, blood or lymph or other malignancies; connective tissue

malignancies; metastatic disease; minimal residual disease following transplantation of organs or stem cells; multi-drug resistant cancers, primary or secondary malignancies, angiogenesis related to malignancy, or other forms of cancer. Also contemplated within the presently disclosed embodiments are specific embodiments wherein only one of the above types of disease is included, or where specific conditions may be excluded regardless of whether or not they are characterized by BRAF^{V600E} expression.

Certain methods of treatment or prevention contemplated herein include administering a host cell (which may be autologous, allogeneic or syngeneic) comprising a desired polynucleotide as described herein that is stably integrated into the chromosome of the cell. For example, such a cellular composition may be generated ex vivo using autologous, allogeneic or syngeneic immune system cells (e.g., T cells, antigen-presenting cells, natural killer cells) in order to administer a desired, BRAF^{V600E}-targeted T-cell composition to a subject as an adoptive immunotherapy. In certain embodiments, the host cell is a hematopoietic progenitor cell or a human immune cell. In certain embodiments, the immune system cell is a CD4⁺ T cell, a CD8⁺ T cell, a CD4⁺ CD8⁻ double-negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a natural killer T cell, a dendritic cell, or any combination thereof. In certain embodiments, the immune system cell is a naïve T cell, a central memory T cell, a stem cell memory T cell, an effector memory T cell, or any combination thereof. In particular embodiments, the cell is a CD4⁺ T cell.

As used herein, administration of a composition or therapy refers to delivering the same to a subject, regardless of the route or mode of delivery. Administration may be effected continuously or intermittently, and parenterally. Administration may be for treating a subject already confirmed as having a recognized condition, disease or disease state, or for treating a subject susceptible to or at risk of developing such a condition, disease or disease state. Co-administration with an adjunctive therapy may include simultaneous and/or sequential delivery of multiple agents in any order and on any dosing schedule (e.g., BRAF^{V600E}-specific recombinant (i.e., engineered) host cells with one or more cytokines; immunosuppressive therapy such as calcineurin inhibitors, corticosteroids, microtubule inhibitors, low dose of a mycophenolic acid prodrug, or any combination thereof).

In certain embodiments, a plurality of doses of a recombinant host cell as described herein is administered to the subject, which may be administered at intervals between administrations of about two to about four weeks. In further embodiments, a cytokine (e.g., IL-2, IL-15, IL-21) is administered sequentially, provided that the subject was administered the recombinant host cell at least three or four times before cytokine administration. In certain embodiments, the cytokine is administered concurrently with the host cell. In certain embodiments, the cytokine is administered subcutaneously.

In still further embodiments, the subject being treated is further receiving immunosuppressive therapy, such as calcineurin inhibitors, corticosteroids, microtubule inhibitors, low dose of a mycophenolic acid prodrug, or any combination thereof. In yet further embodiments, the subject being treated has received a non-myeloablative or a myeloablative hematopoietic cell transplant, wherein the treatment may be administered at least two to at least three months after the non-myeloablative hematopoietic cell transplant.

An effective amount of a therapeutic or pharmaceutical composition refers to an amount sufficient, at dosages and for periods of time needed, to achieve the desired clinical

results or beneficial treatment, as described herein. An effective amount may be delivered in one or more administrations. If the administration is to a subject already known or confirmed to have a disease or disease-state, the term “therapeutic amount” may be used in reference to treatment, whereas “prophylactically effective amount” may be used to describe administering an effective amount to a subject that is susceptible or at risk of developing a disease or disease-state (e.g., recurrence) as a preventative course.

The level of a CTL immune response may be determined by any one of numerous immunological methods described herein and routinely practiced in the art. The level of a CTL immune response may be determined prior to and following administration of any one of the herein described BRAF^{V600E}-specific binding proteins expressed by, for example, a T cell. Cytotoxicity assays for determining CTL activity may be performed using any one of several techniques and methods routinely practiced in the art (see, e.g., Henkart et al., “Cytotoxic T-Lymphocytes” in *Fundamental Immunology*, Paul (ed.) (2003 Lippincott Williams & Wilkins, Philadelphia, PA), pages 1127-50, and references cited therein).

Antigen-specific T cell responses are typically determined by comparisons of observed T cell responses according to any of the herein described T cell functional parameters (e.g., proliferation, cytokine release, CTL activity, altered cell surface marker phenotype, etc.) that may be made between T cells that are exposed to a cognate antigen in an appropriate context (e.g., the antigen used to prime or activate the T cells, when presented by immunocompatible antigen-presenting cells) and T cells from the same source population that are exposed instead to a structurally distinct or irrelevant control antigen. A response to the cognate antigen that is greater, with statistical significance, than the response to the control antigen signifies antigen-specificity.

A biological sample may be obtained from a subject for determining the presence and level of an immune response to a BRAF^{V600E}-containing antigen peptide as described herein. A “biological sample” as used herein may be a blood sample (from which serum or plasma may be prepared), biopsy specimen, body fluids (e.g., lung lavage, ascites, mucosal washings, synovial fluid), bone marrow, lymph nodes, tissue explant, organ culture, or any other tissue or cell preparation from the subject or a biological source. Biological samples may also be obtained from the subject prior to receiving any immunogenic composition, which biological sample is useful as a control for establishing baseline (i.e., pre-immunization) data.

The pharmaceutical compositions described herein may be presented in unit-dose or multi-dose containers, such as sealed ampoules or vials. Such containers may be frozen to preserve the stability of the formulation until. In certain embodiments, a unit dose comprises a recombinant host cell as described herein at a dose of about 10⁷ cells/m² to about 10¹¹ cells/m². The development of suitable dosing and treatment regimens for using the particular compositions described herein in a variety of treatment regimens, including e.g., parenteral or intravenous administration or formulation.

If the subject composition is administered parenterally, the composition may also include sterile aqueous or oleaginous solution or suspension. Suitable non-toxic parenterally acceptable diluents or solvents include water, Ringer's solution, isotonic salt solution, 1,3-butanediol, ethanol, propylene glycol or polyethylene glycols in mixtures with water. Aqueous solutions or suspensions may further comprise one or more buffering agents, such as sodium acetate, sodium

citrate, sodium borate or sodium tartrate. Of course, any material used in preparing any dosage unit formulation should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compounds may be incorporated into sustained-release preparation and formulations. Dosage unit form, as used herein, refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit may contain a predetermined quantity of recombinant cells or active compound calculated to produce the desired therapeutic effect in association with an appropriate pharmaceutical carrier.

In general, an appropriate dosage and treatment regimen provides the active molecules or cells in an amount sufficient to provide therapeutic or prophylactic benefit. Such a response can be monitored by establishing an improved clinical outcome (e.g., more frequent remissions, complete or partial, or longer disease-free survival) in treated subjects as compared to non-treated subjects. Increases in preexisting immune responses to a tumor protein generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or cytokine assays, which are routine in the art and may be performed using samples obtained from a subject before and after treatment.

In still further aspects, unit dose forms comprising host cells according to the present disclosure are provided.

Methods according to this disclosure may further include administering one or more additional agents to treat the disease or disorder in a combination therapy. For example, in certain embodiments, a combination therapy comprises administering a BRAF^{V600E}-specific binding protein (or an engineered host cell expressing the same) with (concurrently, simultaneously, or sequentially) an immune checkpoint inhibitor. In some embodiments, a combination therapy comprises administering a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) with an agonist of a stimulatory immune checkpoint agent. In further embodiments, a combination therapy comprises administering a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) with a secondary therapy, such as chemotherapeutic agent, a radiation therapy, a surgery, an antibody, or any combination thereof.

As used herein, the term “immune suppression agent” or “immunosuppression agent” refers to one or more cells, proteins, molecules, compounds or complexes providing inhibitory signals to assist in controlling or suppressing an immune response. For example, immune suppression agents include those molecules that partially or totally block immune stimulation; decrease, prevent or delay immune activation; or increase, activate, or up regulate immune suppression. Exemplary immunosuppression agents to target (e.g., with an immune checkpoint inhibitor) include PD-1, PD-L1, PD-L2, LAG3, CTLA4, B7-H3, B7-H4, CD244/2B4, HVEM, BTLA, CD160, TIM3, GAL9, KIR, PVR1G (CD112R), PVRL2, adenosine, A2aR, immunosuppressive cytokines (e.g., IL-10, IL-4, IL-1RA, IL-35), IDO, arginase, VISTA, TIGIT, LAIR1, CEACAM-1, CEACAM-3, CEACAM-5, Treg cells, or any combination thereof.

An immune suppression agent inhibitor (also referred to as an immune checkpoint inhibitor) may be a compound, an antibody, an antibody fragment or fusion polypeptide (e.g., Fc fusion, such as CTLA4-Fc or LAG3-Fc), an antisense molecule, a ribozyme or RNAi molecule, or a low molecular weight organic molecule. In any of the embodiments disclosed herein, a method may comprise administering a

BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) with one or more inhibitor of any one of the following immune suppression components, singly or in any combination.

In certain embodiments, a BRAF^{V600E}-specific binding protein is used in combination with a PD-1 inhibitor, for example a PD-1-specific antibody or binding fragment thereof, such as pidilizumab, nivolumab (Keytruda, formerly MDX-1106), pembrolizumab (Opdivo, formerly MK-3475), MEDI0680 (formerly AMP-514), AMP-224, BMS-936558 or any combination thereof. In further embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with a PD-L1 specific antibody or binding fragment thereof, such as BMS-936559, durvalumab (MEDI4736), atezolizumab (RG7446), avelumab (MSB0010718C), MPDL3280A, or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with a LAG3 inhibitor, such as LAG525, IMP321, IMP701, 9H12, BMS-986016, or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein is used in combination with an inhibitor of CTLA4. In particular embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with a CTLA4 specific antibody or binding fragment thereof, such as ipilimumab, tremelimumab, CTLA4-Ig fusion proteins (e.g., abatacept, belatacept), or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with a B7-H3 specific antibody or binding fragment thereof, such as enoblituzumab (MGA271), 376.96, or both. A B7-H4 antibody binding fragment may be a scFv or fusion protein thereof, as described in, for example, Dangaj et al., *Cancer Res.* 73:4820, 2013, as well as those described in U.S. Pat. No. 9,574,000 and PCT Patent Publication Nos. WO/201640724A1 and WO 2013/025779A1.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of CD244.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of BLTA, HVEM, CD160, or any combination thereof. Anti CD-160 antibodies are described in, for example, PCT Publication No. WO 2010/084158.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of TIM3.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of Gal9.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of adenosine signaling, such as a decoy adenosine receptor.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of A2aR.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of KIR, such as lirilumab (BMS-986015).

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of an inhibitory cytokine (typically, a cytokine other than TGFβ) or Treg development or activity.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an IDO inhibitor, such as levo-1-methyl tryptophan, epacadostat (INCB024360; Liu et al., *Blood* 115:3520-30, 2010), ebiselen (Terentis et al., *Biochem.* 49:591-600, 2010), indoximod, NLG919 (Mautino et al., American Association for Cancer Research 104th Annual Meeting 2013; Apr. 6-10, 2013), 1-methyl-tryptophan (1-MT)-tira-pazamine, or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an arginase inhibitor, such as N(omega)-Nitro-L-arginine methyl ester (L-NAME), N-omega-hydroxy-nor-L-arginine (nor-NOHA), L-NOHA, 2(S)-amino-6-boronoheptanoic acid (ABH), S-(2-boronoethyl)-L-cysteine (BEC), or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of VISTA, such as CA-170 (Curis, Lexington, Mass.).

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of TIGIT such as, for example, COM902 (Compugen, Toronto, Ontario Canada), an inhibitor of CD155, such as, for example, COM701 (Compugen), or both.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of PVRIG, PVRL2, or both. Anti-PVRIG antibodies are described in, for example, PCT Publication No. WO 2016/134333. Anti-PVRL2 antibodies are described in, for example, PCT Publication No. WO 2017/021526.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with a LAIR1 inhibitor.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an inhibitor of CEACAM-1, CEACAM-3, CEACAM-5, or any combination thereof.

In certain embodiments, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) is used in combination with an agent that increases the activity (i.e., is an agonist) of a stimulatory immune checkpoint molecule. For example, a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) can be used in combination with a CD137 (4-1BB) agonist (such as, for example, urelumab), a CD134 (OX-40) agonist (such as, for example, MEDI6469, MEDI6383, or MEDI0562), lenalidomide, pomalidomide, a CD27 agonist (such as, for example,

CDX-1127), a CD28 agonist (such as, for example, TGN1412, CD80, or CD86), a CD40 agonist (such as, for example, CP-870,893, rhuCD40L, or SGN-40), a CD122 agonist (such as, for example, IL-2) an agonist of GITR (such as, for example, humanized monoclonal antibodies described in PCT Patent Publication No. WO 2016/054638), an agonist of ICOS (CD278) (such as, for example, GSK3359609, mAb 88.2, JTX-2011, Icos 145-1, Icos 314-8, or any combination thereof). In any of the embodiments disclosed herein, a method may comprise administering a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) with one or more agonist of a stimulatory immune checkpoint molecule, including any of the foregoing, singly or in any combination.

In certain embodiments, a combination therapy comprises a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) and a secondary therapy comprising one or more of: an antibody or antigen binding-fragment thereof that is specific for a cancer antigen expressed by the non-inflamed solid tumor, a radiation treatment, a surgery, a chemotherapeutic agent, a cytokine, RNAi, or any combination thereof.

In certain embodiments, a combination therapy method comprises administering a BRAF^{V600E}-specific binding protein and further administering a radiation treatment or a surgery. Radiation therapy is well-known in the art and includes X-ray therapies, such as gamma-irradiation, and radiopharmaceutical therapies. Surgeries and surgical techniques appropriate to treating a given cancer or non-inflamed solid tumor in a subject are well-known to those of ordinary skill in the art.

In certain embodiments, a combination therapy method comprises administering a BRAF^{V600E}-specific binding protein of the present disclosure (or an engineered host cell expressing the same) and further administering a chemotherapeutic agent. A chemotherapeutic agent includes, but is not limited to, an inhibitor of chromatin function, a topoisomerase inhibitor, a microtubule inhibiting drug, a DNA damaging agent, an antimetabolite (such as folate antagonists, pyrimidine analogs, purine analogs, and sugar-modified analogs), a DNA synthesis inhibitor, a DNA interactive agent (such as an intercalating agent), and a DNA repair inhibitor. Illustrative chemotherapeutic agents include, without limitation, the following groups: anti-metabolites/anticancer agents, such as pyrimidine analogs (5-fluorouracil, fluoruridine, capecitabine, gemcitabine and cytarabine) and purine analogs, folate antagonists and related inhibitors (mercaptopurine, thioguanine, pentostatin and 2-chlorodeoxyadenosine (cladribine)); antiproliferative/antimitotic agents including natural products such as vinca alkaloids (vinblastine, vincristine, and vinorelbine), microtubule disruptors such as taxane (paclitaxel, docetaxel), vincristin, vinblastin, nocodazole, epothilones and navelbine, epididophyllotoxins (etoposide, teniposide), DNA damaging agents (actinomycin, amsacrine, anthracyclines, bleomycin, busulfan, camptothecin, carboplatin, chlorambucil, cisplatin, cyclophosphamide, Cytosan, dactinomycin, daunorubicin, doxorubicin, epirubicin, hexamethylmelamineoxalipatin, iphosphamide, melphalan, mechlorethamine, mitomycin, mitoxantrone, nitrosourea, plicamycin, procarbazine, taxol, taxotere, temozolamide, teniposide, triethylmethiophosphoramide and etoposide (VP 16)); antibiotics such as dactinomycin (actinomycin D), daunorubicin, doxorubicin (adriamycin), idarubicin, anthracyclines, mitoxantrone, bleomycins, plicamycin (mithramycin) and mitomycin; enzymes (L-asparaginase which systemically

metabolizes L-asparagine and deprives cells which do not have the capacity to synthesize their own asparagine); antiplatelet agents; antiproliferative/antimitotic alkylating agents such as nitrogen mustards (mechlorethamine, cyclophosphamide and analogs, melphalan, chlorambucil), ethylenimines and methylmelamines (hexamethylmelamine and thiotepea), alkyl sulfonates-busulfan, nitrosoureas (carmustine (BCNU) and analogs, streptozocin), trazenes-dacarbazine (DTIC); antiproliferative/antimitotic antimetabolites such as folic acid analogs (methotrexate); platinum coordination complexes (cisplatin, carboplatin), procarbazine, hydroxyurea, mitotane, aminoglutethimide; hormones, hormone analogs (estrogen, tamoxifen, goserelin, bicalutamide, nilutamide) and aromatase inhibitors (letrozole, anastrozole); anticoagulants (heparin, synthetic heparin salts and other inhibitors of thrombin); fibrinolytic agents (such as tissue plasminogen activator, streptokinase and urokinase), aspirin, dipyridamole, ticlopidine, clopidogrel, abciximab; antimigratory agents; antisecretory agents (breveldin); immunosuppressives (cyclosporine, tacrolimus (FK-506), sirolimus (rapamycin), azathioprine, mycophenolate mofetil); anti-angiogenic compounds (TNP470, genistein) and growth factor inhibitors (vascular endothelial growth factor (VEGF) inhibitors, fibroblast growth factor (FGF) inhibitors); angiotensin receptor blocker; nitric oxide donors; anti-sense oligonucleotides; antibodies (trastuzumab, rituximab); chimeric antigen receptors; cell cycle inhibitors and differentiation inducers (tretinoin); mTOR inhibitors, topoisomerase inhibitors (doxorubicin (adriamycin), amsacrine, camptothecin, daunorubicin, dactinomycin, eniposide, epirubicin, etoposide, idarubicin, irinotecan (CPT-11) and mitoxantrone, topotecan, irinotecan), corticosteroids (cortisone, dexamethasone, hydrocortisone, methylprednisolone, prednisone, and prednisolone); growth factor signal transduction kinase inhibitors; mitochondrial dysfunction inducers, toxins such as Cholera toxin, ricin, Pseudomonas exotoxin, Bordetella pertussis adenylate cyclase toxin, or diphtheria toxin, and caspase activators; and chromatin disruptors.

Cytokines are increasingly used to manipulate host immune response towards anticancer activity. See, e.g., Floros & Tarhini, *Semin. Oncol.* 42(4):539-548, 2015. Cytokines useful for promoting immune anticancer or antitumor response include, for example, IFN- α , IL-2, IL-3, IL-4, IL-10, IL-12, IL-13, IL-15, IL-16, IL-17, IL-18, IL-21, IL-24, and GM-CSF, singly or in any combination with the binding proteins or cells expressing the same of this disclosure.

Another cancer therapy approach involves reducing expression of oncogenes and other genes needed for growth, maintenance, proliferation, and immune evasion by cancer cells. RNA interference, and in particular the use of microRNAs (miRNAs) and small inhibitory RNAs (siRNAs) provides an approach for knocking down expression of cancer genes (see, e.g., Larsson et al., *Cancer Treat. Rev.* 16(55): 128-135, 2017), which can be used in combination with the binding proteins or cells expressing the same of this disclosure.

In any of the embodiments disclosed herein, any of the therapeutic agents (e.g. a BRAF^{V600E}-specific binding protein or engineered host cell, an inhibitor of an immune suppression component, an agonist of a stimulatory immune checkpoint molecule, an antitumor lymphocyte, a chemotherapeutic agent, a radiation therapy, a surgery, a cytokine, or an inhibitory RNA) may be administered once or more than once to the subject over the course of a treatment, and, in combinations, may be administered to the subject in any

41

order or any combination. An appropriate dose, suitable duration, and frequency of administration of a therapeutic agent will be determined by such factors as a condition of the patient; size, type, spread, growth, and severity of the tumor or cancer; particular form of the active ingredient; and the method of administration.

EXAMPLES

Example 1

Identification of CD4⁺ BRAF^{V600E}-Specific T Cells in a Melanoma Patient

A 52-year-old man presented with stage IIIC, BRAF^{V600E}-mutated melanoma originating on the left foot and was treated with wide excision, lymph node dissection and adjuvant ipilimumab. Shortly before completing one year of ipilimumab, he required resection of three in-transit

42

metastases in his left leg, and an additional in-transit metastasis 3 months later. He subsequently progressed with a 3 cm left iliac nodal metastasis and soft tissue nodular FDG avid lesion in the left thigh (FIG. 1A). The iliac node was resected for whole exome sequencing and expansion of TIL, and the patient subsequently received TIL infusion following lymphodepleting chemotherapy. The left thigh lesion resolved and the patient remains free of disease 27 months after therapy.

Whole exome and RNA sequencing of purified tumor cells and normal tissue identified only 20 nonsynonymous missense mutations, shown in Table 1: (columns 2 and 3 from left); columns 4-6 from left: 27-mer peptides encompassing the encoded nonsynonymous mutations and the presence of the mutation in DNA or RNA-Seq (columns 4-6 from left); (columns 7 and 8 from left) engineered 20-mer peptides comprising the mutation; and RNA-seq expression in units of transcripts per million (TPM).

TABLE 1

Nonsynonymous Mutations in Purified Tumor Cells and Normal Tissue from a Melanoma Patient								
symbol	sub.nt	sub.aa	27-mer aa sequence	Mut'n in DNA	Mut'n in RNA	20-mer peptide 1	20-mer peptide 2	TPM in RNA
AP1M1	AP1M1.A>C	p.I295L	VIEKHSRSRI EYMLKAKSQ FKRRSTAN (SEQ ID NO: 79)	yes	yes	VIEKHSRS RIEYMLK AKSQF (SEQ ID NO: 80)	SRIEYML KAKSQFK RRSTAN (SEQ ID NO: 81)	61.33
BRAF	BRAF.A>T	p.V600E	DLTVKIGDF GLATEKSRW SGSHQFEQL (SEQ ID NO: 37)	yes	yes	DLTVKIG DFGLATE KSRWSG (SEQ ID NO: 38)	DFGLA KSRWSGE HQFEQL (SEQ ID NO: 39)	10.83
DCAF6	DCAF6.G>A	p.A419T	EQFLQPSTSS TMSTQAHST SSPTESPH (SEQ ID NO: 82)	yes	yes	EQFLQPST SSTMTQ AHSTS (SEQ ID NO: 83)	TSSTMT QAHSTSS PTESPH (SEQ ID NO: 84)	41.11
GTF2H4	GTF2H4.C>T	p.T319M	FIVVETNYRL YAYMESELQ IALIALFS (SEQ ID NO: 85)	yes	yes	FIVVETN YRLYAY MESELQI (SEQ ID NO: 86)	YRLYAY MESELQI ALIALFS (SEQ ID NO: 87)	49.95
NBPF12	NBPF12.A>Gp	E2471G	DSCQPYRSSF YALGEKHVG FSLDVGEI (SEQ ID NO: 88)	yes	yes	DSCQPYR SSFYALG EKHVG (SEQ ID NO: 89)	SSFYALG EKHVGFS LDVGEI (SEQ ID NO: 90)	12.5
ORC3	ORC3.A>C	p.I236L	ESFATKVLQ DFIILSSQHL HEFPLILI (SEQ ID NO: 91)	yes	yes	ESFATKV LQDFIILS SQHLH (SEQ ID NO: 92)	LQDFIILS SQHLHEF PLILI (SEQ ID NO: 93)	19.92
ROR1	ROR1.A>G	p.N53S	LVPTSSWNIS SELSKDSYLT LDEPMNN (SEQ ID NO: 94)	yes	yes	LVPTSSW NISSELSK DSYLT (SEQ ID NO: 95)	NISSELSK DSYLTLD EPMNN (SEQ ID NO: 96)	4.69
SF3B1	ZNF700.T>G	p.T358A	QMGGSTPVL TPGKAPIGTP AMNMATPT (SEQ ID NO: 97)	yes	yes	QMGGSTP VLTPGKA PIGTPA (SEQ ID NO: 98)	VLTPGKA PIGTPAM NMATPT (SEQ ID NO: 99)	105.2

TABLE 1-continued

Nonsynonymous Mutations in Purified Tumor Cells and Normal Tissue from a Melanoma Patient								
symbol	sub.nt	sub.aa	27-mer aa sequence	Mut'n in DNA	Mut'n in RNA	20-mer peptide 1	20-mer peptide 2	TPM in RNA
UNKL	UNKL.C>T	p.V154I	AHGPLDLRP PVCDIRELQ AQEALQNGQ (SEQ ID NO: 100)	yes	yes	AHGPLDL RPPVCDIR ELQQAQ (SEQ ID NO: 101)	RPPVCDIR ELQQAQEA LQNGQ (SEQ ID NO: 102)	6.72
ZNF700	ZNF700.T>G	p.F287L	GEKPYECSK CDKALHSSS SYHRHERSH (SEQ ID NO: 103)	yes	yes	GEKPYEC SKCDKAL HSSSSY (SEQ ID NO: 104)	SKCDKAL HSSSSYH RHERSH (SEQ ID NO: 105)	8.69
NVL	NVL.T>G	p.T370P	APCIIFIDEID AIPPKREVAS KDMERR (SEQ ID NO: 106)	yes	yes	APCIIFIDE IDAIPPKR EVA (SEQ ID NO: 107)	DEIDAIPP KREVASK (SEQ ID NO: 108)	18.42
MATN1	MATN1.T>G	p.T153P	SRSPDISKVV IVVPDGRPQ DSVQDVSA (SEQ ID NO: 109)	yes	yes	SRSPDISK VVIVVPD GRPQD (SEQ ID NO: 110)	KVVIVVP DGRPQDS VQDVSA (SEQ ID NO: 111)	0.14
CTNNA2	CTNNA2.A>C	p.N351H	VRQALQDL SEYMHNTGR KEKGDPLNI (SEQ ID NO: 112)	yes	no	VRQALQD LLSEYMH NTGRKE (SEQ ID NO: 113)	LLSEYMH NTGRKEK GDPLNI (SEQ ID NO: 114)	2.36
GET4	GET4.T>G	p.L65R	RYMSQS KHT EARERMYSG ALLFFSHGQ (SEQ ID NO: 115)	yes	no	RYMSQSK HTEARER MYSGAL (SEQ ID NO: 116)	HTEARER MYSGALL FFSHGQ (SEQ ID NO: 117)	26.97
NTNG1	NTNG1.G>A	p.V271I	TVTDLRIRLL RPAIGEIVFD ELHLARY (SEQ ID NO: 118)	yes	yes	TVTDLRIR LLRPAIGE IFVD (SEQ ID NO: 119)	RLLRPAIG EIFVDELH LARY (SEQ ID NO: 120)	0.73
SPTBN5	SPTBN5.G>A	p.T3127I	TLLLDALWT TKAAIAESQ DYGQDLEGV (SEQ ID NO: 121)	yes	yes	TLLLDALW LTTKAAI AESQDY (SEQ ID NO: 122)	LTTKAAI AESQDYG QDLEGV (SEQ ID NO: 123)	1.11
DPP6	DPP6.C>T	p.5113L	LLVILVICSLI VTLVILLTPA EDNSLS (SEQ ID NO: 124)	yes	no	LLVILVIC SLIVTLVI LLTP (SEQ ID NO: 125)	CSLIVTLV ILLTPAED NSLS (SEQ ID NO: 126)	0.03
HIAT1	HIAT1.G>T	p.G93C	VKGLLSFLS APLICALSDV WGRKSFLL (SEQ ID NO: 127)	yes	no	VKGLLSF LSAPLICA LSDVW (SEQ ID NO: 128)	LSAPLICA LSDVWGR KSFLL (SEQ ID NO: 129)	46.71
ITGA4	ITGA4.G>T	p.V359F	GSGAVMNA METNLFGSD KYAARFGES I (SEQ ID NO: 130)	yes	no	GSGAVM NAMETNL FGSDKYA (SEQ ID NO: 131)	AMETNLF GSDKYAA RFGESI (SEQ ID NO: 132)	4.59
MYO1A	MYO1A.C>T	p.V1017I	SVRFKENS AVKVIQGPA GGDNSKLRY (SEQ ID NO: 133)	yes	no	SVRFKEN SVAVKVI QGPAGG (SEQ ID NO: 134)	SVAVKVI QGPAGGD NSKLRY (SEQ ID NO: 135)	0.05

45

Also determined, but not shown in Table 1, were chromosomal positions (using GRCh37/hg19 reference assembly) and the variant allele frequency (VAF) for each mutation. The VAF for the BRAF^{V600E} mutation was 35.4%.

T cell responses to these potential neoantigens were evaluated by stimulating peripheral blood mononuclear cells (PBMC) obtained from the patient after TIL therapy with a pool of peptides flanking each of the 20 mutations. No CD8⁺ T cell responses to the candidate neoantigens were detected; however, a CD4⁺ T cell response specific for 20-mer peptides encompassing BRAF^{V600E} was identified. The BRAF^{V600E}-reactive cells were purified by IFN- γ capture and shown to recognize autologous B cells pulsed with mutant but not wildtype BRAF peptide, confirming specificity for the mutant peptide (FIG. 1B). To determine whether BRAF^{V600E} is processed and presented by class II MHC⁺ APC, autologous B cells were transfected with mRNA encoding wildtype or mutant BRAF sequences targeted to the endosome. The T cells recognized B cells expressing mutant, but not wildtype BRAF (FIG. 1C).

Recognition was blocked by anti-HLA-DQ but not anti-class I or anti HLA-DR antibodies, identifying HLA-DQ as the likely restricting allele (FIG. 1D). Analysis of multiple B cell lines of known genotype suggested restriction by HLA-DQA1*03 paired with HLA-DQB1*03, with weak recognition of DQB1*0301 and stronger recognition of DQB1*0302 and DQB1*0303 (FIG. 1F and Table 2).

TABLE 2

IFN- γ production by BRAF ^{V600E} -specific CD4 ⁺ T cells following incubation with allogeneic B-LCL cell lines expressing different class II alleles and in the presence or absence of antigen.				
Cell line	BRAF V600E Peptide	Mean IFN-G pg/ml	HLA DRB1	HLA DQB1
1331	-	10	404	302
	+	41791		
CFS	-	10	0401, 0101	0301, 0501
	+	1547		
DEM	-	23	0401, 1602	0302, 0502
	+	29873		
DEU	-	6	401	301
	+	10359		
FAL	-	9	0403, 0801	03BG, 0402
	+	31620		
BM14	-	536	401	302
	+	42832		
DMB	-	38	0101, 1501	0501, 0602
	+	7		
DLM	-	26	0403, 0801	03BG, 0402
	+	39388		

46

TABLE 2-continued

IFN- γ production by BRAF ^{V600E} -specific CD4 ⁺ T cells following incubation with allogeneic B-LCL cell lines expressing different class II alleles and in the presence or absence of antigen.				
Cell line	BRAF V600E Peptide	Mean IFN-G pg/ml	HLA DRB1	HLA DQB1
AMM	-	6	0802, 1501	0402, 06WG
	+	9		
CLC	-	9	0301, 1104	02AB, 0301
	+	6		
BP	-	35	1601, 1101	0502, 0301
	+	36		
JWP	-	12	0701, 0701	02AB, 0303
	+	359		
DAH2	-	17	09, 1501	0303, 06W6
	+	49784		
VRM	-	6	0701, 10	0303, 0501
	+	188		

The complete patient HLA typing was as follows: A*11:01:01/A*24:02:01; B*15:01:01/B*40:01:02; C*03:03:01/C*03:04:01; DPA1*01:03:01/DPA1*01:03:01; DPB1*04:01:01/DPB1*04:01:01; DQA1*03:01:01/DQA1*03:02; DQB1*03:02:01/DQB1*03:03:02; DRB1*04:03:01/DRB1*09:01:02; DRB4*01:03:01/DRB4*01:03:02.

B-LCL transfected with HLA-DQA1*03 or DQB1*0302, but not the closely linked HLA-DRB1*04, were recognized by BRAF^{V600E}-specific CD4⁺ T cells when pulsed with the mutant peptide, confirming the HLA restriction (FIG. 1E).

Recognition of three melanoma cell lines with an HLA-DQB1*0302 and BRAF^{V600E} genotype was tested. One tumor cell line that expressed the greatest amount of HLA-DQ was recognized by the BRAF^{V600E}-specific CD4⁺ T cells demonstrating that the epitope can be presented directly by tumor cells (FIGS. 1G-II). Tumor-specific CD4⁺ T cells can have anti-tumor activity through direct cell killing and cytokine release (see, e.g., Quezada et al., *J. Exp. Med.* 207(3):637-650 (2010); Manici et al., *J. Exp. Med.* 189(5):871-876 (1999)), but a major role is to support the development and function of CD8⁺ T cells by licensing APC and producing cytokines (see, e.g., Sun and Bevan, *Science* 300(5617):339-342 (2003); Williams et al., *Nature* 441(7095):890-893 (2006)). Although adoptively transferred TIL contained BRAF^{V600E}-specific CD4⁺ T cells, CD8⁺ T cells were the prevalent population in TIL (Table 3). Briefly, the final TIL product infused into the patient was analyzed by flow cytometry for phenotype and, following stimulation with PMA/Ionomycin, was stained intracellularly for cytokines. Percentages shown in Table 3 are percentages of CD45⁺ cells (top) or CD4 or CD8 cells (bottom).

TABLE 3

Phenotype of Final TIL product						
% of live	% of CD45					
	CD45	CD3 T	T δ T	NKT	CD8T	CD4T Treg
	99.9	99.7	0.04	0.014	93.9	3.4 0.52
% of CD45						
	CD8P1	CD8 TIM3	CD8 TM	CD8 naïve	CD8 EMRA	CD8EM
	51.6	93.4	0.002	0.003	36.5	58.6

TABLE 3-continued

Phenotype of Final TIL product			
% of CD8 or CD4			
CD8 IFN- γ +	CD4 IL17+	CD4 IL22	CD4 IFN- γ +
99.3	3.6	0.42	98.8

Moreover, a majority of IFN- γ produced by stimulation of multiple independent TIL cultures with autologous tumor was blocked by a HLA class I blocking antibody (Table 4).

TABLE 4

Tumor specificity and class I blocking of initial TIL cultures												
	Pool T	Pool R	Pool B	Pool A	Pool S	Pool 1	Pool 2	Pool 3	Pool 4	Pool 5	Pool 6	Frag. 12
Tumor	2179.50	752.12	2993.64	4427.63	2313.73	3409.99	7213.00	3722.50	3866.23	2914.58	3916.23	2.44
Tumor + Class I Block	94.32	11.95	67.49	38.78	67.88	75.74	394.68	258.88	86.28	369.32	513.78	2.44
Media only	2.44	52.71	67.56	2.44	2.44	40.87	16.52	131.40	2.44	2.44	21.22	2.44
PMA/Iono	2010.14	360.33	2563.22	2929.14	1203.62	2038.37	1565.83	1454.78	2757.56	2012.09	2155.09	2.44
% Blocking	96%	98%	98%	99%	97%	98%	95%	93%	98%	87%	87%	0%

No IFN- γ production was observed when TIL were cultured with autologous B cells pulsed with pools of peptides that included all of the 20 non-synonymous mutations identified by tumor exome sequencing (FIG. 2A), but IFN- γ was produced after co-culture with B cells pulsed with peptides from lineage-restricted self-antigens (tyrosinase, Mart-1, TRP2) and a cancer testes antigen (Mage A3), which are known targets of T cells in melanoma (see, e.g., Gros et al. *Nat. Med.* 2016) (FIG. 2F), or transfected with tandem minigenes encompassing 29 non-synonymous mutations or the coding sequences of Tyrosinase, Mage-A3, Mart1, SSX2, and GP100 (FIGS. 2B-2E). Thus the patient TIL contained BRAF^{V600E}-specific CD4⁺ T cells and a diverse CD8⁺ T cell response to self-antigens.

Clinical Protocol

The patient was enrolled for TIL generation under an FDA-approved IND and a clinical protocol approved by the Institutional Review Board of Fred Hutchinson Cancer Research Center (FHCRC 2643; NCT01807182). Patients with stage IV melanoma, or stage III unlikely to be cured by surgery, >18 years of age, with an ECOG $\leq$ 1, with a site of metastatic disease that could be safely resected or biopsied, were eligible. TIL were expanded from tumor fragments in 6,000 IU/ml recombinant IL-2 (Proleukin; Novartis), using methodologies developed at the Surgery Branch of the National Cancer Institute (e.g., Dudley et al., *J. Immunother.* 24(4):363-373 (2002)). TIL cultures were selected based on cell growth and autologous tumor reactivity as determined by IFN- γ secretion following co-culture with autologous tumor cells. The TIL were cryopreserved until needed for use, then thawed and further expanded using a rapid expansion protocol, as previously-described (Riddell and Greenberg, *J. Immunological Methods* 128(2):189-201 (1990)). The expanded TIL were administered to the patient following a lymphodepleting chemotherapy regimen of cyclophosphamide 60 mg/kg/day $\times$ 2 days, then fludarabine

25 mg/m²/day $\times$ 5 days. Within 24 hours of the TIL infusion, the patient received high-dose IL-2 at 600,000 IU/kg IV every 8 hours, for a total of 9 doses. Tumor responses were

assessed using RECIST version 1.1 with CT and MRI at weeks 6, 12, and 24, then every 3-6 months, at the discretion of the primary provider.

Nucleic Acid Preparation for Exome Capture and RNA sequencing

Post-treatment blood was used to isolate non-tumor DNA. A single-cell suspension derived from the iliac nodal tumor recurrence was flow sorted (propidium iodide negative and CD45 negative) to deplete abundant infiltrating lymphocytes and enrich for neoplastic cells. Normal tissue and sorted tumor cells were processed with the Qiagen DNA/RNA AllPrep Micro kit to isolate DNA for exome capture, with RNA reserved for subsequent RNA-seq profiling. Genomic DNA concentration was quantified on an Invitrogen Qubit® 2.0 Fluorometer (Life Technologies-Invitrogen, Carlsbad, CA, USA) and Trinean DropSense96 spectrophotometer (Caliper Life Sciences, Hopkinton, MA).

Whole Exome Sequencing

Exome sequencing libraries were prepared using the Agilent SureSelectXT Reagent Kit and exon targets isolated using the Agilent All Human Exon v6 (Agilent Technologies, Santa Clara, CA, USA). 200 ng of genomic DNA was fragmented using a Covaris LE220 focused-ultrasonicator (Covaris, Inc., Woburn, MA, USA) and libraries prepared and captured on a Sciclone NGSx Workstation (PerkinElmer, Waltham, MA, USA). Library size distributions were validated using an Agilent 2200 TapeStation. Additional library QC, blending of pooled indexed libraries, and cluster optimization was performed using Life Technologies' Invitrogen Qubit® 2.0 Fluorometer.

The resulting libraries were sequenced on an Illumina HiSeq 2500 using a paired-end 100 bp (PE100) strategy. Image analysis and base calling was performed using Illumina's Real Time Analysis v1.18 software, followed by "demultiplexing" of indexed reads and generation of FASTQ files using Illumina's bcl2fastq Conversion Software v1.8.4 (<http://support.illumina.com/downloads/>

bcl2fastq_conversion_software_184.html). Read pairs passing standard Illumina quality filters were retained for further analysis, yielding 77M read pairs for the tumor and 89M read pairs for the normal. Paired reads were aligned to the human genome reference (GRCh37/hg19) with the BWA-MEM short-read aligner (see, e.g., Li, H., *arXiv preprint arXiv:1303.3997* (2013); Li and Ruddin, *Bioinformatics* 25(14):1754-1760). The resulting alignment files, in standard BAM format, were processed by Picard 2.0.1 and GATK 3.5[37] for quality score recalibration, indel realignment, and duplicate removal according to recommended best practices (see, Auwera et al., *Current Protocols in Bioinformatics* pp. 11.10.1-11.10.33 (2013)).

Three independent software packages were used to call somatic mutations from the analysis-ready tumor and normal BAM files: MuTect 1.1.7[39], Strelka 1.0.14[40], and VarScan.v2.4.1 (Koboldt et al., *Genome Res.* 22(3):568-576 (2012)). Variant calls from all tools, in VCF format, were annotated with Oncotator (Ramos et al., *Human Mutation* 36(4):E2423-E2429 (2015)). Missense somatic variants were combined and annotated further, including wild-type and variant peptide sequences, to form an integrated summary from which candidate peptides were chosen for synthesis.

RNA-Seq Data Processing

To rank candidate peptides by observed expression level, RNA-seq was performed on flow-sorted tumor cells from the same single cell suspension. RNA-seq libraries were prepared from total RNA using the TruSeq RNA Sample Prep v2 Kit (Illumina, Inc., San Diego, CA, USA) and a Sciclone NGSx Workstation (PerkinElmer, Waltham, MA, USA). Library size distributions were validated using an Agilent 2200 TapeStation (Agilent Technologies, Santa Clara, CA, USA). Additional library QC, blending of pooled indexed libraries, and cluster optimization was performed using Life Technologies' Invitrogen Qubit® 2.0 Fluorometer (Life Technologies-Invitrogen, Carlsbad, CA, USA). The library was sequenced on an Illumina HiSeq 2500 to generate 133M 50 nt paired reads (PESO). Reads were aligned to a RefSeq derived reference transcriptome with RSEM 1.2.19 (see Li and Dewey, *BMC Bioinformatics* 12(1):323 (2011)). Gene-level expression values from RSEM, in TPM units, were added to the summary of missense somatic variants.

T Cell Culture

Initial stimulations were performed with overlapping 20-mer crude peptides obtained from Elim Biopharma, with 2 peptides spanning each mutation with the mutated residue at position +7 or +13 of the 20 amino acid sequence. Subsequent experiments were performed with >80% purity 21 mer peptides with V600 (wildtype) or E600 (mutant) at position +11. Cryopreserved PBMC were thawed and rested overnight in CTL (RPMI media with L-glutamine and HEPES (Gibco) supplemented with 10% human serum (produced in house), 50 μ M beta-mercaptoethanol, penicillin and streptomycin, 4 mM L-glutamine and 2 ng/ml recombinant human IL-7 (Peprotech). The following morning, PBMC were washed and stimulated at 10^6 cells in 5 ml CTL per well of a 6 well plate with a pool of 1 μ g/ml of each peptide without cytokines. Recombinant IL-2 (Peprotech) was added to a final concentration of 10 U/ml on day +3, and half media changes with supplemental IL-2 were performed on days +3, +6, and +9. On day +13, cells were used in an ELISA and cytokine staining assays.

Antigen-specific T cell enrichment was carried out by staining live cells for secreted IFN- γ using the IFN- γ secretion assay APC (Miltenyi) following the manufacturer's

instructions, and using autologous B cells as antigen presenting cells pulsed with 10 μ g/ml 21-mer BRAF mutant peptide. CD4⁺ IFN- γ secreting cells were sorted on a FACS Aria2. Sorted cells were rested in CTL supplemented with 10 ng/ml human IL-15 for 5 days, then expanded using a rapid expansion protocol described previously (Riddell and Greenberg, *supra*). Antigen-specific T cells were further enriched by sorting for V β 3.1 positive, CD4⁺ cells by staining with anti-V β 3.1 (Thermo Scientific, cat. no. TCR2740), expanded, and cryopreserved at day 13 or 14 after expansion. Cryopreserved cells were thawed and rested overnight in CTL supplemented with 10 U/ml IL-2 prior to assays.

Antigen-Presenting Cells

Autologous B cells were isolated from fresh or thawed PBMC using magnetic beads coated with antibodies recognizing CD19 (Miltenyi, cat. no. 130-050-301) and magnetic positive selection according to the manufacturer's instructions (Miltenyi, cat. no. 130-042-401). Primary B cells were incubated in a 1:1 ratio with NIH 3T3 cells expressing humanCD40L for 7 days in B cell medium supplemented with 200 U/ml human IL-4 (Peprotech) as described (see Tran et al., *Science* 344(3184):641-645 (2014)). B cells were subsequently harvested and restimulated with 3T3 CD40L and fresh medium every 3 days. B cells were used in assays at day +3 of stimulation 2 or 3.

Cytokine Release Assays

In ELISA assays, 50,000 effector T cells were incubated in 96 well round bottom plates with 100,000 B cells or B-LCL lines and 10 μ g/ml or specific concentrations of peptides in RPMI (Gibco) supplemented with 5% heat inactivated fetal bovine serum. IFN- γ in supernatants was quantitated using the ready set go human IFN- γ ELISA kit (ebiosciences) in technical triplicate. HLA blocking experiments were carried out with 20 μ g/ml antibody anti class I (Biolegend, cat. no. 311411) anti-HLA DR (clone L243, cat. no. 307611) and HLA-DQ (Abcam, clone spv-13, cat. no. ab23632) added 1 hour prior to adding peptide. For elispot assays, 50,000 tumor infiltrating lymphocytes were incubated with 200,000 autologous B cells pulsed with peptide pools at a final concentration of 10 μ g/ml of each peptide in CTL medium using the human IFN- γ ELISpot-Pro kit (Mabtech) and developed using the manufacturer's instructions.

HLA Identification

LCL cell lines 1331, DUCAF, VAVY, BM14, DEM and DEU were utilized. For co-culture assays, LCL cell lines were pulsed with 10 μ g/ml of BRAF mutant peptide or DMSO control for 4 hours and then washed 3 times with PBS prior to ELISA assay. For identification of specific class II alleles, codon optimized linear DNA fragments encoding HLA-DRB1 0404 protein or the HLA-DQB1 0302 protein linked by a T2A skip sequence to HLA-DQA1 0301 protein were synthesized using Genestings™ (Life Sciences) and cloned into the vector MP71 (Engels et al., *Hum. Gene Ther.* 14(12):1155-1168 (2003)) linearized with NotI and EcoRI (Thermo Fisher) using the NEBuilder cloning kit (New England Biolabs) and sequence verified. Retroviral transduction was performed as described by Sommermeyer et al., (*Leukemia*, 2015)) into the VAVY cell line homozygous for HLA DRB1 0301, DQA1 0501, and DQB1 0201 (Research cell bank). Cells positive for DRB1 0404 were sorted on a FACS Aria2 sorter using the antibody DRB1-PE (Biolegend, cat. no. 362303) and cells positive for DQB1 03 DQA1 03 were sorted using the anti DQ antibody clone HLADQ1-FITC (Biolegend, cat. no. 318104). Assays were performed with and without pulsing with BRAF^{V600E} peptide. ELISA

experiments were performed in technical duplicate or triplicate as indicated and are representative of two independent experiments.

Example 2

Identification of TCR Gene Usage by TIL

Deep sequencing was performed to identify TCR gene usage in BRAF^{V600E}-specific T cells and other T cells in the TIL. Three TCR clonotypes showed marked expansion after stimulation of post-treatment PBMC with BRAF^{V600E} peptide, and these sequences were further enriched after IFN- γ capture, indicating their specificity for the mutant peptide (FIG. 2G). TCR V β sequencing of tumor, TIL, and PBMC obtained prior to TIL infusion identified all 3 TCR V β clones in the tumor, and 2 of 3 in TIL. All 3 TCR V β sequences were below the level of detection in pre-treatment PBMC indicating enrichment at the tumor site (FIG. 2H). A total of 34 common V β sequences collectively made up >50% of the TIL product (FIG. 2I). Only 5 of these 34 clones were detected in the pretreatment blood, with 4 at very low frequency (FIG. 2J). To assess TCR gene usage of CD8⁺ T cells recognizing each of the 4 lineage-specific or C/T antigens, IFN- γ capture was used to sort these cells from TIL and assess TCR V β usage. Seven different V β sequences in the sorted cells were identified and represented 4.7% of the T cells in the TIL product (FIGS. 2L-2O). These 7 clonotypes and one of the BRAF-specific clones were detected in blood obtained 10 and 24 months post-treatment (FIGS. 2J and 2K). RNA expression targeted to the endosome was carried out using the method described by Kreiter et al. (*J. Immunol.* 180(1):309-318 (2008)) where antigens are targeted to the endosome by fusion of the antigen to class I MHC sorting signals. The mRNA expression construct pJV57 was constructed by gene synthesis (Geneart, Life Sciences), which contained a T7 promoter fused to the N terminal 25 amino acids of the human HLA-B gene, followed by a BamHI restriction site, the coding sequence of enhanced GFP, an AgeI restriction site, the C terminal 55 amino acids of the human HLA-B gene, followed by the human beta globin untranslated region followed by a 30 nucleotide poly A tail followed by a SapI restriction site directing cleavage in the poly A tail. Construct pJV84 was cloned by ligating the following into AgeI/BamHI digested pJV57: annealed oligonucleotides (Ultramers, Integrated DNA Technologies) encoding BRAF amino acids 575-624 flanked by a 5' AgeI and 3' BamHI site containing the E600 substitution. Construct pJV85 was made by ligating annealed oligonucleotides (Ultramers, Integrated DNA Technologies) encoding BRAF amino acids 575-624 flanked by a 5' AgeI and 3' BamHI site containing the wildtype V600 amino acid. pJV84 and pJV85 were then linearized with SapI (Thermo Fisher) and mRNA was in vitro transcribed using the Highscribe T7 ARCA mRNA kit (New England Biolabs) and purified by lithium precipitation according to the manufacturer's instructions. mRNA was electroporated into CD40L stimulated B cells 16 hours prior to co-culture experiments as described by Tran et al. (supra).

Example 3

Phenotypic Characterization of TIL Product

Phenotypic analysis of BRAF-specific clones was performed following TIL treatment (FIGS. 3A-3C). BRAF^{V600E}-specific CD4⁺ T cells showed an effector

memory phenotype (CD45RA⁺CCR7⁺CD27⁺KLRG1⁺) and expressed low levels of PD-1 (FIGS. 3A, 3B). The majority of BRAF^{V600E}-specific cells expressed CXCR3 and CCR4. A fraction of the cells also expressed the skin-homing marker cutaneous lymphocyte-associated antigen (CLA). BRAF^{V600E} peptide-activated cells produced IFN- γ , TNF- α , IL-4, and IL-21 (FIG. 3C), sometimes in combination (data not shown). Taken together, these data suggest that circulating BRAF-specific CD4⁺ T cells after TIL infusion have a mixed Th1/Th2 phenotype, consistent with an established memory cellular immune response to mutated BRAF in melanoma.

Example 4

Construction and Testing of BRAF^{V600E} TCRs

Durable remissions in melanoma after adoptive transfer of self-antigen reactive CD8⁺ T cells alone are exceedingly rare (see, e.g., Johnson et al., *Blood* 114(3):535-546 (2009); Yee et al., *PNAS* 99(25):16168-16173 (2002); Dudley et al., *J. Immunother.* 24(4):363-373 (2001)). Without wishing to be bound by theory, it is believed that the BRAF^{V600E}-specific CD4⁺ T cells provide direct antitumor effects and aid the persistence and function of self-antigen reactive CD8⁺ T cells against a tumor that contained few neoantigens. The HLA-DQA1*03/DQB1*03 restricting allele for the BRAF^{V600E}-specific CD4⁺ T cells is present in 29% of individuals in the International Histocompatibility Workgroup database (Petersdorf, E., *personal communication, International Histocompatibility Working Group in Hematopoietic Cell Transplantation*. 2017), and isolation of the BRAF^{V600E}-specific TCR genes from this patient can facilitate adoptive therapy for patients with BRAF mutant tumors with TCR engineered T cells. TCR V α sequencing on samples with varying levels of BRAF-reactive clones identified four (4) TCR V α sequences that correlated in frequency with the three (3) TCR V β sequences (FIG. 2G).

Four (4) TCRs from the three (3) TCRB alleles and four (4) TRCA alleles identified were constructed and tested for antigen-specific response (FIGS. 4A, 4B, and 5). The dominant TCR from the patient (pJV88) was expressed in CD4⁺ T cells from two healthy donors and conferred specificity to cells expressing BRAF^{V600E} but not wildtype BRAF sequences (FIG. 2H). The TCR pJV90, which was made from the second most dominant V β clone from the patient and one of the possible alpha chains, showed some activity but also possibly nonspecific baseline activation.

TCR V β and V α Sequencing

DNA from clinical samples was isolated using the Qiagen DNeasy or Qiaamp micro DNA kits according to the manufacturer's instructions. TCRB sequencing was carried out using the human TCRB sequencing kit (Adaptive Biotechnology) following the manufacturer's instructions and sequenced using a MiSeq (Fred Hutchinson Cancer Research Center Genomics core) with data analysis carried out by Adaptive biotechnology software. TCRA sequencing was carried out using the human TCRA sequencing service (Adaptive Biotechnology).

T Cell Receptor Construction

TCR construction was in the vector PRRL (Jones et al., *Hum. Gene Ther.* 20(6):630-640 (2009)), which was further modified by introducing six point mutations into the start codon and putative promoter region of the woodchuck hepatitis virus X protein as in Lim and Brown, *RNA Biology* 13(9):743-747 (2016)) with the beta chain followed by a P2A translational skip sequence followed by the alpha chain

with cysteines introduced to facilitate pairing (see Kuball et al., *Blood* 109(6):2331-2338 (2007)). A codon-optimized DNA fragment containing the TRBV28 and CDR3 and TRBJ1-3 sequences followed by TCRB1 sequence with a cysteine substituted at residue 57 followed by a P2A skip sequence and the TRAV21 and CDR3 sequences followed by TRAJ43 and TRAC sequences was synthesized as a genestring (Life Sciences) and cloned using the NEBuilder cloning kit (New England Biolabs) into the vector PRRL-SIN linearized with PstI and AscI (Thermo Fisher) and sequence verified. One week after transduction, cells were sorted based on Vbeta3.1 expression using antibody clone 8F10 (Thermo Scientific, cat. no. TCR2740) and expanded via rapid expansion as described above. T cells were used in assays or cryopreserved on day 14 of the rapid expansion.

Example 5

CRISPR-Mediated Deletion of Endogenous TCR Leads to Increased Expression of Transgenic BRAF-Specific TCR

Following gene transfer of synthetic T cell receptor sequences into T cells, the transferred TCR alpha and beta chains may compete with endogenous TCR subunits for expression and signaling machinery. To investigate whether deletion of endogenous TCR increased expression of the transferred TCR, stimulated T cells were first transfected with CRISPR-Cas9 ribonucleoproteins with guide RNA sequences directing cleavage of the endogenous TCR alpha (guide rna AGAGTCTCTCAGCTGGTACA; SEQ ID NO:136) and TCR beta (guide rna: GGAGAATGACGAGTGGACCC; SEQ ID NO:139) constant region genes (from Ren et al., *Clin. Cancer Res.*

23(9):2255-2266 (2017)). 12 uM Cas9 protein (IDT) 20 uM guide RNA (IDT) with electroporation enhancer (IDT) was assembled and 3 ul was added to 2e6 primary human T cells 2 days following stimulation with antiCD3/antiCD28 Dynabeads (Thermo-Fisher) in nucleofection buffer P3 (Lonza), and nucleofected using an AMAXA 4D nucleofection electroporator (Lonza) using program EH-115. This combination led to >99% reduction in cells with TCR expression, as measured by staining of the CD3 component of the TCR receptor complex (FIG. 6A). When gene deletion of the endogenous TCR was combined with gene transfer of the BRAF^{V600E}-specific TCR increased expression of the transferred T cell receptor on the cell surface was observed, as measured by tetramer staining (FIG. 6B).

The various embodiments described above can be combined to provide further embodiments. All of the U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, including U.S. Provisional Patent Application No. 62/544,695, filed Aug. 11, 2017, are incorporated herein by reference, in their entirety. Aspects of the embodiments can be modified, if necessary to employ concepts of the various patents, applications and publications to provide yet further embodiments.

These and other changes can be made to the embodiments in light of the above-detailed description. In general, in the following claims, the terms used should not be construed to limit the claims to the specific embodiments disclosed in the specification and the claims, but should be construed to include all possible embodiments along with the full scope of equivalents to which such claims are entitled. Accordingly, the claims are not limited by the disclosure.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 149

<210> SEQ ID NO 1

<211> LENGTH: 137

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR1 alpha chain variable domain

<400> SEQUENCE: 1

Met Glu Thr Leu Leu Gly Leu Leu Ile Leu Trp Leu Gln Leu Gln Trp
1 5 10 15

Val Ser Ser Lys Gln Glu Val Thr Gln Ile Pro Ala Ala Leu Ser Val
20 25 30

Pro Glu Gly Glu Asn Leu Val Leu Asn Cys Ser Phe Thr Asp Ser Ala
35 40 45

Ile Tyr Asn Leu Gln Trp Phe Arg Gln Asp Pro Gly Lys Gly Leu Thr
50 55 60

Ser Leu Leu Leu Ile Gln Ser Ser Gln Arg Glu Gln Thr Ser Gly Arg
65 70 75 80

Leu Asn Ala Ser Leu Asp Lys Ser Ser Gly Arg Ser Thr Leu Tyr Ile
85 90 95

Ala Ala Ser Gln Pro Gly Asp Ser Ala Thr Tyr Leu Cys Ala Val Arg
100 105 110

Arg Gly Asn Asn Asp Met Arg Phe Gly Ala Gly Thr Arg Leu Thr Val
115 120 125

-continued

Lys Pro Asn Ile Gln Asn Pro Asp Pro
130 135

<210> SEQ ID NO 2
<211> LENGTH: 135
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-1 alpha chain variable domain

<400> SEQUENCE: 2

Met Arg Leu Val Ala Arg Val Thr Val Phe Leu Thr Phe Gly Thr Ile
1 5 10 15

Ile Asp Ala Lys Thr Thr Gln Pro Thr Ser Met Asp Cys Ala Glu Gly
20 25 30

Arg Ala Ala Asn Leu Pro Cys Asn His Ser Thr Ile Ser Gly Asn Glu
35 40 45

Tyr Val Tyr Trp Tyr Arg Gln Ile His Ser Gln Gly Pro Gln Tyr Ile
50 55 60

Ile His Gly Leu Lys Asn Asn Glu Thr Asn Glu Met Ala Ser Leu Ile
65 70 75 80

Ile Thr Glu Asp Arg Lys Ser Ser Thr Leu Ile Leu Pro His Ala Thr
85 90 95

Leu Arg Asp Thr Ala Val Tyr Tyr Cys Ile Val Arg Ala Tyr Ser Gly
100 105 110

Tyr Ser Thr Leu Thr Phe Gly Lys Gly Thr Met Leu Leu Val Ser Pro
115 120 125

Asp Ile Gln Asn Pro Asp Pro
130 135

<210> SEQ ID NO 3
<211> LENGTH: 142
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-2 alpha chain variable domain

<400> SEQUENCE: 3

Met Lys Ser Leu Arg Val Leu Leu Val Ile Leu Trp Leu Gln Leu Ser
1 5 10 15

Trp Val Trp Ser Gln Gln Lys Glu Val Glu Gln Asn Ser Gly Pro Leu
20 25 30

Ser Val Pro Glu Gly Ala Ile Ala Ser Leu Asn Cys Thr Tyr Ser Asp
35 40 45

Arg Gly Ser Gln Ser Phe Phe Trp Tyr Arg Gln Tyr Ser Gly Lys Ser
50 55 60

Pro Glu Leu Ile Met Phe Ile Tyr Ser Asn Gly Asp Lys Glu Asp Gly
65 70 75 80

Arg Phe Thr Ala Gln Leu Asn Lys Ala Ser Gln Tyr Val Ser Leu Leu
85 90 95

Ile Arg Asp Ser Gln Pro Ser Asp Ser Ala Thr Tyr Leu Cys Ala Val
100 105 110

Ile Thr Leu Asn Asn Asn Ala Gly Asn Met Leu Thr Phe Gly Gly Gly
115 120 125

Thr Arg Leu Met Val Lys Pro His Ile Gln Asn Pro Asp Pro
130 135 140

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 138
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 alpha chain variable domain

<400> SEQUENCE: 4

```
Met Lys Ser Leu Arg Val Leu Leu Val Ile Leu Trp Leu Gln Leu Ser
 1             5             10             15

Trp Val Trp Ser Gln Gln Lys Glu Val Glu Gln Asn Ser Gly Pro Leu
      20             25             30

Ser Val Pro Glu Gly Ala Ile Ala Ser Leu Asn Cys Thr Tyr Ser Asp
      35             40             45

Arg Gly Ser Gln Ser Phe Phe Trp Tyr Arg Gln Tyr Ser Gly Lys Ser
 50             55             60

Pro Glu Leu Ile Met Phe Ile Tyr Ser Asn Gly Asp Lys Glu Asp Gly
 65             70             75             80

Arg Phe Thr Ala Gln Leu Asn Lys Ala Ser Gln Tyr Val Ser Leu Leu
      85             90             95

Ile Arg Asp Ser Gln Pro Ser Asp Ser Ala Thr Tyr Leu Cys Ala Val
      100            105            110

Thr Ser Asn Ala Gly Lys Ser Thr Phe Gly Asp Gly Thr Thr Leu Thr
      115            120            125

Val Lys Pro Asn Ile Gln Asn Pro Asp Pro
      130            135
```

<210> SEQ ID NO 5
 <211> LENGTH: 132
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 beta chain variable domain

<400> SEQUENCE: 5

```
Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1             5             10             15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
      20             25             30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
      35             40             45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50             55             60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65             70             75             80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
      85             90             95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
      100            105            110

Asn Glu Gly Asn Ser Gly Asn Thr Ile Tyr Phe Gly Glu Gly Ser Trp
      115            120            125

Leu Thr Val Val
      130
```

<210> SEQ ID NO 6

-continued

```

<211> LENGTH: 130
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2 beta chain variable
domain

```

```

<400> SEQUENCE: 6

```

```

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1             5             10             15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20             25             30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35             40             45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50             55             60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65             70             75             80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85             90             95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
 100            105            110

Gly Ala Arg Gln Ile Pro Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr
 115            120            125

Val Val
 130

```

```

<210> SEQ ID NO 7
<211> LENGTH: 140
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR3 beta chain variable
domain

```

```

<400> SEQUENCE: 7

```

```

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1             5             10             15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20             25             30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35             40             45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50             55             60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65             70             75             80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85             90             95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
 100            105            110

Ser Leu Ser Ala Ala Gly Gly Gly Tyr Gly Tyr Thr Phe Asn Tyr Gly
 115            120            125

Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr Val Val
 130            135            140

```

```

<210> SEQ ID NO 8
<211> LENGTH: 287
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRBV28-01

<400> SEQUENCE: 8

gatgtgaaag taaccagag ctcgagatat ctagtcaaaa ggacgggaga gaaagttttt    60
ctggaatgtg tccaggatat ggaccatgaa aatatgttct ggtatcgaca agaccaggt    120
ctggggctac ggctgatcta tttctcatat gatgttaaaa tgaaagaaaa aggagatatt    180
cctgaggggt acagtgtctc tagagagaag aaggagcgct tctccctgat tctggagtcc    240
gccagcacca accagacatc tatgtacctc tgtgccagca gtttatg                287

<210> SEQ ID NO 9
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRBJ01-03

<400> SEQUENCE: 9

ctctggaaac accatatatt ttggagaggg aagttggctc actgtttag                50

<210> SEQ ID NO 10
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRBJ01-02

<400> SEQUENCE: 10

ctaactatgg ctacaccttc ggttcgggga ccaggttaac cgtttag                48

<210> SEQ ID NO 11
<211> LENGTH: 279
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAV21-01

<400> SEQUENCE: 11

aaacaggagg tgacgcagat tctgcagct ctgagtgtcc cagaaggaga aaacttggtt    60
ctcaactgca gtttactga tagcgctatt tacaacctcc agtggtttag gcaggacctt    120
gggaaaggtc tcacatctct gttgcttatt cagtcaagtc agagagagca aacaagtgga    180
agacttaatg cctcgctgga taaatcatca ggacgtagta ctttatacat tgcagcttct    240
cagcctggtg actcagccac ctacctctgt gctgtgagg                279

<210> SEQ ID NO 12
<211> LENGTH: 277
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAV26-01

<400> SEQUENCE: 12

gatgctaaga ccaccagcc cccctccatg gattgcgctg aaggaagagc tgcaaacctg    60
ccttgtaatc actctacat cagtggaaat gagtatgtgt attggtatcg acagattcac    120
tcccaggggc cacagtatat cattcatggt ctaaaaaaca atgaaaccaa tgaaatggcc    180
tctctgatca tcacagaaga cagaaagtcc agcaccttga tctgccccca cgctacgctg    240
agagacactg ctgtgtacta ttgcatcgtc agagtcg                277

```

-continued

```

<210> SEQ ID NO 13
<211> LENGTH: 277
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAV12-02

<400> SEQUENCE: 13

cagaaggagg tggagcagaa ttctggaccc ctcagtgttc cagagggagc cattgcctct      60
ctcaactgca cttacagtga ccgagggttc cagtccttct tctggtacag acaatattct      120
gggaaaagcc ctgagttgat aatgttcata tactccaatg gtgacaaaga agatggaagg      180
tttacagcac agctcaataa agccagccag tatgtttctc tgctcatcag agactcccag      240
cccagtgatt cagccaccta cctctgtgcc gtgaaca                                277

<210> SEQ ID NO 14
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAJ43-01

<400> SEQUENCE: 14

acaataacaa tgacatgcgc ttggagcag ggaccagact gacagtaaaa ccaa              54

<210> SEQ ID NO 15
<211> LENGTH: 60
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAJ11-01

<400> SEQUENCE: 15

tgaattcagg atacagcacc ctcaccttg ggaaggggac tatgcttcta gtctctccag      60

<210> SEQ ID NO 16
<211> LENGTH: 63
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAJ39-01

<400> SEQUENCE: 16

tgaataataa tgcaggcaac atgctcacct ttggaggggg aacaaggtta atggtcaaac      60
ccc                                                63

<210> SEQ ID NO 17
<211> LENGTH: 59
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCRAJ27-01

<400> SEQUENCE: 17

taacaccaat gcaggcaaat caacctttgg ggatgggact acgctcactg tgaagccaa      59

<210> SEQ ID NO 18
<211> LENGTH: 409
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR1 alpha chain variable
domain

```


-continued

<400> SEQUENCE: 18

atggaaacac tgctgggect gctgatectg ttgctgcaac tgcaatgggt gtccagcaag	60
caagaagtga ctcagatccc tgccgctctg tctgtgcctg agggcgaaaa cctggctctg	120
aactgcagct tcaccgacag cgccatctac aacctgcagt ggttcaggca ggatccaggc	180
aagggcctga catccctgct gctgattcag agcagccaga gagagcagac cagcggcaga	240
ctgaatgcc gcttgataa gtctccggc agaagcacc tgtatatgc cgtttctcag	300
ccaggcgata ggcacacata tctgtgtgcc gttcggagag gcaacaacga catgagattc	360
ggagccggca ccagactgac cgtgaagccc aacattcaga accccgatc	409

<210> SEQ ID NO 19

<211> LENGTH: 403

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR2-1 alpha chain variable domain

<400> SEQUENCE: 19

atgagactgg tggccagagt gacagtgttc ctgaccttcg gcaccatcat cgacgccaag	60
accacacagc ccaccagcat ggattgtgcc gagggcagag ctgccaacct gccttgtaat	120
cacagcacca tcagcgga caggtacgtg tactggtaca ggcagatcca ctctcagggc	180
cctcagtaca tcattccagg cctgaagaac aacgagacaa acgagatggc ctctctgac	240
atcaccgagg accgcaagag cagcaccctg attctgcctc acgccacact gagagacacc	300
gccgtgtact actgtatcgt gcgggcctac agcggctaca gcacactgac atttggaag	360
ggcaccatgc tgctcgtgtc ccagacatt cagaaccccg atc	403

<210> SEQ ID NO 20

<211> LENGTH: 424

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR2-2 alpha chain variable domain

<400> SEQUENCE: 20

atgaagtccc tgagagtgtc gctgggtcct ctgtggctgc agctgtcttg ggtctgggtcc	60
cagcagaaag aggtggaaca gaacagcggc cctctgtctg ttctgaagg cgctatcgcc	120
tctctgaatt gcacctacag cgacagaggc agccagagct tctctggta tcggcagtac	180
agcggcaaga gcccagagct gatcatgttc atttacagca acggcgacaa agaggacggc	240
cggtttacag ccagctgaa caaggccagc cagtacgtgt cactgctgat cagagacagc	300
cagcctagcg acagcgccac ctatctgtgt gccgtgatca ccctgaacaa caacgccggc	360
aacatgctga cctttggcgg cggaacacgg ctgatgggta agccccacat tcagaacccc	420
gac	424

<210> SEQ ID NO 21

<211> LENGTH: 412

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR3 alpha chain variable domain

<400> SEQUENCE: 21

-continued

atgaagtccc tgagagtgtc gctgggtcacc ctgtgggtgc agctgtcttg ggtctgggtcc	60
cagcagaaaag aggtggaaca gaacagcggc cctctgtccg ttctgaagg cgccattgct	120
agcctgaatt gcacctacag cgaccggggc agccagagct tctctggta taggcagtac	180
agcggcaaga gccccgagct gatcatgttc atctacagca acggcgacaa agaggacggc	240
cggtttacag cccagctgaa caaggccagc cagtacgtgt cactgctgat cagagacagc	300
cagcctagcg acagcgccac ctatctgtgt gccgtgacaa gcaatgccg caagtcacaca	360
tttggcgacg gcacaaccct gaccgtgaag cccaacattc agaaccgccg tc	412

<210> SEQ ID NO 22
 <211> LENGTH: 396
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 beta chain variable domain

<400> SEQUENCE: 22

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttg cctgggtggac	60
gtgaaagtga cccagagcag cagatacttg gtcaagagaa ccggcgagaa ggtgttcttg	120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg	180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaagg cgacatcccc	240
gagggctaca gcgtgtccag agagaagaaa gagcggttca gcctgacct ggaaagcgcc	300
agcaccaacc agaccagcat gtacctgtgc gccagcaacg agggcaatag cggaacacc	360
atctacttcg gcgaaggcag ctggctgacc gtggtg	396

<210> SEQ ID NO 23
 <211> LENGTH: 390
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2 beta chain variable domain

<400> SEQUENCE: 23

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttg cctgggtggac	60
gtgaaagtga cccagagcag cagatacttg gtcaagagaa ccggcgagaa ggtgttcttg	120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg	180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaagg cgacatcccc	240
gagggctaca gcgtgtccag agagaagaaa gagcggttca gcctgacct ggaaagcgcc	300
agcaccaacc agaccagcat gtacctgtgt gccagcggcg ccagacagat cccttacaca	360
tttggctccg gcaccagact gaccgtggtg	390

<210> SEQ ID NO 24
 <211> LENGTH: 420
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 beta chain variable domain

<400> SEQUENCE: 24

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttg cctgggtggac	60
gtgaaagtga cccagagcag cagatacttg gtcaagagaa ccggcgagaa ggtgttcttg	120

-continued

```

gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg 180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc 240
gagggctaca gcgtgtccag agagaagaaa gagcgggtca gcctgatactt ggaaagcgcc 300
agcaccaacc agaccagcat gtacctgtgt gccagctctc tgtctgctgc cggcggaggg 360
tatggctaca ccttcaatta cggctacaca ttcggcagcg gcaccagact gacctgggtg 420

```

```

<210> SEQ ID NO 25
<211> LENGTH: 134
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR alpha chain constant
domain

```

```

<400> SEQUENCE: 25

```

```

Ala Val Tyr Gln Leu Arg Asp Ser Lys Ser Ser Asp Lys Ser Val Cys
1          5          10          15

```

```

Leu Phe Thr Asp Phe Asp Ser Gln Thr Asn Val Ser Gln Ser Lys Asp
20          25          30

```

```

Ser Asp Val Tyr Ile Thr Asp Lys Cys Val Leu Asp Met Arg Ser Met
35          40          45

```

```

Asp Phe Lys Ser Asn Ser Ala Val Ala Trp Ser Asn Lys Ser Asp Phe
50          55          60

```

```

Ala Cys Ala Asn Ala Phe Asn Asn Ser Ile Ile Pro Glu Asp Thr Phe
65          70          75          80

```

```

Phe Pro Ser Pro Glu Ser Ser Cys Asp Val Lys Leu Val Glu Lys Ser
85          90          95

```

```

Phe Glu Thr Asp Thr Asn Leu Asn Phe Gln Asn Leu Ser Val Ile Gly
100         105         110

```

```

Phe Arg Ile Leu Leu Leu Lys Val Ala Gly Phe Asn Leu Leu Met Thr
115        120        125

```

```

Leu Arg Leu Trp Ser Ser
130

```

```

<210> SEQ ID NO 26
<211> LENGTH: 177
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR beta chain constant
domain

```

```

<400> SEQUENCE: 26

```

```

Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro
1          5          10          15

```

```

Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu
20          25          30

```

```

Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn
35          40          45

```

```

Gly Lys Glu Val His Ser Gly Val Cys Thr Asp Pro Gln Pro Leu Lys
50          55          60

```

```

Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser Arg Leu
65          70          75          80

```

```

Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe Arg Cys
85          90          95

```

```

Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr Gln Asp
100         105         110

```

-continued

Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp Gly Arg
 115 120 125

Ala Asp Cys Gly Phe Thr Ser Val Ser Tyr Gln Gln Gly Val Leu Ser
 130 135 140

Ala Thr Ile Leu Tyr Glu Ile Leu Leu Gly Lys Ala Thr Leu Tyr Ala
 145 150 155 160

Val Leu Val Ser Ala Leu Val Leu Met Ala Met Val Lys Arg Lys Asp
 165 170 175

Phe

<210> SEQ ID NO 27
 <211> LENGTH: 407
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR alpha chain constant domain

<400> SEQUENCE: 27

ctgcagtgtgta ccagctgcgg gacagcaaga gcagcgacaa gacgtgtgc ctgttcaccg	60
acttcgacag ccagaccaac gtgtcccaga gcaaggacag cgacgtgtac atcaccgata	120
agtgcgtgct ggacatgcgg agcatggact tcaagagcaa cagcgccgtg gcctggtcca	180
acaagagcga cttgcctgc gccaacgcct tcaacaacag cattatcccc gaggacacat	240
tcttcccaag ccccgagagc agctgcgacg tgaagctggt ggaaaagagc ttcgagacag	300
acaccaacct gaacttcag aacctcagcg tgatcggctt ccggtatcctg ctgctgaagg	360
tgccgggctt caacctgctg atgacctgc ggctgtggtc cagctga	407

<210> SEQ ID NO 28
 <211> LENGTH: 454
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR beta chain constant domain

<400> SEQUENCE: 28

gaagatctga acaaagtgtt ccctccagag gtggccgtgt tcgagccttc tgaggccgag	60
atcagccaca cacagaaagc cacactcgtg tgccctggcca ccggcttttt tcccgatcac	120
gtggaactgt cttggtgggt caacggcaaa gaggtgcaca gcggcgtctg taccgatcct	180
cagcctctga aagagcagcc cgctctgaac gactccagat actgcctgag cagcaggctg	240
agagtgtccg ccacctctctg gcagaacccc agaaaccact tcagatgccca ggtgcagttc	300
tacggcctga gcgagaacga tgagtggacc caggatagag ccaagcctgt gacacagatc	360
gtgtctgccg aagcctgggg cagagccgat tgtggttta cctccgtgtc ctatcagcag	420
ggcgtgctgt ctgccaccat cctgtatgag atcc	454

<210> SEQ ID NO 29
 <211> LENGTH: 12
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 alpha chain CDR3

<400> SEQUENCE: 29

Cys Ala Val Arg Arg Gly Asn Asn Asp Met Arg Phe
1 5 10

-continued

<210> SEQ ID NO 30
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-1 alpha chain CDR3

<400> SEQUENCE: 30

Cys Ile Val Arg Ala Tyr Ser Gly Tyr Ser Thr Leu Thr Phe
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-2 alpha chain CDR3

<400> SEQUENCE: 31

Cys Ala Val Ile Thr Leu Asn Asn Asn Ala Gly Asn Met Leu Thr Phe
1 5 10 15

<210> SEQ ID NO 32
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR3 alpha chain CDR3

<400> SEQUENCE: 32

Cys Ala Val Thr Ser Asn Ala Gly Lys Ser Thr Phe
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR1 beta chain CDR3

<400> SEQUENCE: 33

Cys Ala Ser Asn Glu Gly Asn Ser Gly Asn Thr Ile Tyr Phe
1 5 10

<210> SEQ ID NO 34
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-1 beta chain CDR3

<400> SEQUENCE: 34

Cys Ala Ser Gly Ala Arg Gln Ile Pro Tyr Thr Phe
1 5 10

<210> SEQ ID NO 35
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR3 beta chain CDR3

<400> SEQUENCE: 35

Cys Ala Ser Ser Leu Ser Ala Ala Gly Gly Gly Tyr Gly Tyr Thr Phe
1 5 10 15

-continued

```

<210> SEQ ID NO 36
<211> LENGTH: 766
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence Human BRAF

<400> SEQUENCE: 36

Met Ala Ala Leu Ser Gly Gly Gly Gly Gly Ala Glu Pro Gly Gln
 1             5             10             15

Ala Leu Phe Asn Gly Asp Met Glu Pro Glu Ala Gly Ala Gly
 20             25             30

Ala Ala Ala Ser Ser Ala Ala Asp Pro Ala Ile Pro Glu Glu Val Trp
 35             40             45

Asn Ile Lys Gln Met Ile Lys Leu Thr Gln Glu His Ile Glu Ala Leu
 50             55             60

Leu Asp Lys Phe Gly Gly Glu His Asn Pro Pro Ser Ile Tyr Leu Glu
 65             70             75             80

Ala Tyr Glu Glu Tyr Thr Ser Lys Leu Asp Ala Leu Gln Gln Arg Glu
 85             90             95

Gln Gln Leu Leu Glu Ser Leu Gly Asn Gly Thr Asp Phe Ser Val Ser
 100            105            110

Ser Ser Ala Ser Met Asp Thr Val Thr Ser Ser Ser Ser Ser Leu
 115            120            125

Ser Val Leu Pro Ser Ser Leu Ser Val Phe Gln Asn Pro Thr Asp Val
 130            135            140

Ala Arg Ser Asn Pro Lys Ser Pro Gln Lys Pro Ile Val Arg Val Phe
 145            150            155            160

Leu Pro Asn Lys Gln Arg Thr Val Val Pro Ala Arg Cys Gly Val Thr
 165            170            175

Val Arg Asp Ser Leu Lys Lys Ala Leu Met Met Arg Gly Leu Ile Pro
 180            185            190

Glu Cys Cys Ala Val Tyr Arg Ile Gln Asp Gly Glu Lys Lys Pro Ile
 195            200            205

Gly Trp Asp Thr Asp Ile Ser Trp Leu Thr Gly Glu Glu Leu His Val
 210            215            220

Glu Val Leu Glu Asn Val Pro Leu Thr Thr His Asn Phe Val Arg Lys
 225            230            235            240

Thr Phe Phe Thr Leu Ala Phe Cys Asp Phe Cys Arg Lys Leu Leu Phe
 245            250            255

Gln Gly Phe Arg Cys Gln Thr Cys Gly Tyr Lys Phe His Gln Arg Cys
 260            265            270

Ser Thr Glu Val Pro Leu Met Cys Val Asn Tyr Asp Gln Leu Asp Leu
 275            280            285

Leu Phe Val Ser Lys Phe Phe Glu His His Pro Ile Pro Gln Glu Glu
 290            295            300

Ala Ser Leu Ala Glu Thr Ala Leu Thr Ser Gly Ser Ser Pro Ser Ala
 305            310            315            320

Pro Ala Ser Asp Ser Ile Gly Pro Gln Ile Leu Thr Ser Pro Ser Pro
 325            330            335

Ser Lys Ser Ile Pro Ile Pro Gln Pro Phe Arg Pro Ala Asp Glu Asp
 340            345            350

His Arg Asn Gln Phe Gly Gln Arg Asp Arg Ser Ser Ser Ala Pro Asn
 355            360            365

```

Val	His	Ile	Asn	Thr	Ile	Glu	Pro	Val	Asn	Ile	Asp	Asp	Leu	Ile	Arg
370						375				380					
Asp	Gln	Gly	Phe	Arg	Gly	Asp	Gly	Gly	Ser	Thr	Thr	Gly	Leu	Ser	Ala
385				390				395						400	
Thr	Pro	Pro	Ala	Ser	Leu	Pro	Gly	Ser	Leu	Thr	Asn	Val	Lys	Ala	Leu
				405				410						415	
Gln	Lys	Ser	Pro	Gly	Pro	Gln	Arg	Glu	Arg	Lys	Ser	Ser	Ser	Ser	Ser
		420						425				430			
Glu	Asp	Arg	Asn	Arg	Met	Lys	Thr	Leu	Gly	Arg	Arg	Asp	Ser	Ser	Asp
		435				440						445			
Asp	Trp	Glu	Ile	Pro	Asp	Gly	Gln	Ile	Thr	Val	Gly	Gln	Arg	Ile	Gly
450						455				460					
Ser	Gly	Ser	Phe	Gly	Thr	Val	Tyr	Lys	Gly	Lys	Trp	His	Gly	Asp	Val
465				470						475				480	
Ala	Val	Lys	Met	Leu	Asn	Val	Thr	Ala	Pro	Thr	Pro	Gln	Gln	Leu	Gln
				485				490						495	
Ala	Phe	Lys	Asn	Glu	Val	Gly	Val	Leu	Arg	Lys	Thr	Arg	His	Val	Asn
		500						505				510			
Ile	Leu	Leu	Phe	Met	Gly	Tyr	Ser	Thr	Lys	Pro	Gln	Leu	Ala	Ile	Val
515						520						525			
Thr	Gln	Trp	Cys	Glu	Gly	Ser	Ser	Leu	Tyr	His	His	Leu	His	Ile	Ile
530						535				540					
Glu	Thr	Lys	Phe	Glu	Met	Ile	Lys	Leu	Ile	Asp	Ile	Ala	Arg	Gln	Thr
545				550						555				560	
Ala	Gln	Gly	Met	Asp	Tyr	Leu	His	Ala	Lys	Ser	Ile	Ile	His	Arg	Asp
				565				570						575	
Leu	Lys	Ser	Asn	Asn	Ile	Phe	Leu	His	Glu	Asp	Leu	Thr	Val	Lys	Ile
		580						585				590			
Gly	Asp	Phe	Gly	Leu	Ala	Thr	Val	Lys	Ser	Arg	Trp	Ser	Gly	Ser	His
		595				600						605			
Gln	Phe	Glu	Gln	Leu	Ser	Gly	Ser	Ile	Leu	Trp	Met	Ala	Pro	Glu	Val
610						615				620					
Ile	Arg	Met	Gln	Asp	Lys	Asn	Pro	Tyr	Ser	Phe	Gln	Ser	Asp	Val	Tyr
625				630						635				640	
Ala	Phe	Gly	Ile	Val	Leu	Tyr	Glu	Leu	Met	Thr	Gly	Gln	Leu	Pro	Tyr
				645				650						655	
Ser	Asn	Ile	Asn	Asn	Arg	Asp	Gln	Ile	Ile	Phe	Met	Val	Gly	Arg	Gly
		660						665				670			
Tyr	Leu	Ser	Pro	Asp	Leu	Ser	Lys	Val	Arg	Ser	Asn	Cys	Pro	Lys	Ala
675						680						685			
Met	Lys	Arg	Leu	Met	Ala	Glu	Cys	Leu	Lys	Lys	Lys	Arg	Asp	Glu	Arg
690						695				700					
Pro	Leu	Phe	Pro	Gln	Ile	Leu	Ala	Ser	Ile	Glu	Leu	Leu	Ala	Arg	Ser
705				710						715				720	
Leu	Pro	Lys	Ile	His	Arg	Ser	Ala	Ser	Glu	Pro	Ser	Leu	Asn	Arg	Ala
		725						730				735			
Gly	Phe	Gln	Thr</												

```
<210> SEQ ID NO 37
<211> LENGTH: 27
<212> TYPE: PRT
```

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence 27-mer BRAFV600E peptide

<400> SEQUENCE: 37

Asp Leu Thr Val Lys Ile Gly Asp Phe Gly Leu Ala Thr Glu Lys Ser
 1 5 10 15

Arg Trp Ser Gly Ser His Gln Phe Glu Gln Leu
 20 25

<210> SEQ ID NO 38
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence 20-mer BRAFV600E peptide

<400> SEQUENCE: 38

Asp Leu Thr Val Lys Ile Gly Asp Phe Gly Leu Ala Thr Glu Lys Ser
 1 5 10 15

Arg Trp Ser Gly
 20

<210> SEQ ID NO 39
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence 20-mer BRAFV600E peptide

<400> SEQUENCE: 39

Asp Phe Gly Leu Ala Thr Glu Lys Ser Arg Trp Ser Gly Ser His Gln
 1 5 10 15

Phe Glu Gln Leu
 20

<210> SEQ ID NO 40
 <211> LENGTH: 266
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence HLADRB1

<400> SEQUENCE: 40

Met Val Cys Leu Lys Leu Pro Gly Gly Ser Cys Met Thr Ala Leu Thr
 1 5 10 15

Val Thr Leu Met Val Leu Ser Ser Pro Leu Ala Leu Ala Gly Asp Thr
 20 25 30

Arg Pro Arg Phe Leu Trp Gln Leu Lys Phe Glu Cys His Phe Phe Asn
 35 40 45

Gly Thr Glu Arg Val Arg Leu Leu Glu Arg Cys Ile Tyr Asn Gln Glu
 50 55 60

Glu Ser Val Arg Phe Asp Ser Asp Val Gly Glu Tyr Arg Ala Val Thr
 65 70 75 80

Glu Leu Gly Arg Pro Asp Ala Glu Tyr Trp Asn Ser Gln Lys Asp Leu
 85 90 95

Leu Glu Gln Arg Arg Ala Ala Val Asp Thr Tyr Cys Arg His Asn Tyr
 100 105 110

Gly Val Gly Glu Ser Phe Thr Val Gln Arg Arg Val Glu Pro Lys Val
 115 120 125

Thr Val Tyr Pro Ser Lys Thr Gln Pro Leu Gln His His Asn Leu Leu

-continued

130	135	140			
Val Cys Ser Val Ser Gly Phe Tyr Pro Gly Ser Ile Glu Val Arg Trp					
145	150	155	160		
Phe Arg Asn Gly Gln Glu Glu Lys Ala Gly Val Val Ser Thr Gly Leu					
165	170	175			
Ile Gln Asn Gly Asp Trp Thr Phe Gln Thr Leu Val Met Leu Glu Thr					
180	185	190			
Val Pro Arg Ser Gly Glu Val Tyr Thr Cys Gln Val Glu His Pro Ser					
195	200	205			
Val Thr Ser Pro Leu Thr Val Glu Trp Arg Ala Arg Ser Glu Ser Ala					
210	215	220			
Gln Ser Lys Met Leu Ser Gly Val Gly Gly Phe Val Leu Gly Leu Leu					
225	230	235	240		
Phe Leu Gly Ala Gly Leu Phe Ile Tyr Phe Arg Asn Gln Lys Gly His					
245	250	255			
Ser Gly Leu Gln Pro Thr Gly Phe Leu Ser					
260	265				

<210> SEQ ID NO 41
 <211> LENGTH: 261
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence HLADQB10302

<400> SEQUENCE: 41

Met Ser Trp Lys Lys Ala Leu Arg Ile Pro Gly Gly Leu Arg Val Ala					
1	5	10	15		
Thr Val Thr Leu Met Leu Ala Met Leu Ser Thr Pro Val Ala Glu Gly					
20	25	30			
Arg Asp Ser Pro Glu Asp Phe Val Tyr Gln Phe Lys Gly Met Cys Tyr					
35	40	45			
Phe Thr Asn Gly Thr Glu Arg Val Arg Leu Val Thr Arg Tyr Ile Tyr					
50	55	60			
Asn Arg Glu Glu Tyr Ala Arg Phe Asp Ser Asp Val Gly Val Tyr Arg					
65	70	75	80		
Ala Val Thr Pro Leu Gly Pro Pro Ala Ala Glu Tyr Trp Asn Ser Gln					
85	90	95			
Lys Glu Val Leu Glu Arg Thr Arg Ala Glu Leu Asp Thr Val Cys Arg					
100	105	110			
His Asn Tyr Gln Leu Glu Leu Arg Thr Thr Leu Gln Arg Arg Val Glu					
115	120	125			
Pro Thr Val Thr Ile Ser Pro Ser Arg Thr Glu Ala Leu Asn His His					
130	135	140			
Asn Leu Leu Val Cys Ser Val Thr Asp Phe Tyr Pro Ala Gln Ile Lys					
145	150	155	160		
Val Arg Trp Phe Arg Asn Asp Gln Glu Glu Thr Thr Gly Val Val Ser					
165	170	175			
Thr Pro Leu Ile Arg Asn Gly Asp Trp Thr Phe Gln Ile Leu Val Met					
180	185	190			
Leu Glu Met Thr Pro Gln Arg Gly Asp Val Tyr Thr Cys His Val Glu					
195	200	205			
His Pro Ser Leu Gln Asn Pro Ile Ile Val Glu Trp Arg Ala Gln Ser					
210	215	220			
Glu Ser Ala Gln Ser Lys Met Leu Ser Gly Ile Gly Gly Phe Val Leu					

-continued

225	230	235	240
Gly Leu Ile Phe Leu Gly Leu Gly Leu Ile Ile His His Arg Ser Gln			
	245	250	255
Lys Gly Leu Leu His			
	260		

<210> SEQ ID NO 42
 <211> LENGTH: 255
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence HLADQA103

<400> SEQUENCE: 42

Met Ile Leu Asn Lys Ala Leu Met Leu Gly Ala Leu Ala Leu Thr Thr			
1	5	10	15
Val Met Ser Pro Cys Gly Gly Glu Asp Ile Val Ala Asp His Val Ala			
	20	25	30
Ser Tyr Gly Val Asn Leu Tyr Gln Ser Tyr Gly Pro Ser Gly Gln Tyr			
	35	40	45
Ser His Glu Phe Asp Gly Asp Glu Glu Phe Tyr Val Asp Leu Glu Arg			
	50	55	60
Lys Glu Thr Val Trp Gln Leu Pro Leu Phe Arg Arg Phe Arg Arg Phe			
	65	70	75
Asp Pro Gln Phe Ala Leu Thr Asn Ile Ala Val Leu Lys His Asn Leu			
	85	90	95
Asn Ile Val Ile Lys Arg Ser Asn Ser Thr Ala Ala Thr Asn Glu Val			
	100	105	110
Pro Glu Val Thr Val Phe Ser Lys Ser Pro Val Thr Leu Gly Gln Pro			
	115	120	125
Asn Thr Leu Ile Cys Leu Val Asp Asn Ile Phe Pro Pro Val Val Asn			
	130	135	140
Ile Thr Trp Leu Ser Asn Gly His Ser Val Thr Glu Gly Val Ser Glu			
	145	150	155
Thr Ser Phe Leu Ser Lys Ser Asp His Ser Phe Phe Lys Ile Ser Tyr			
	165	170	175
Leu Thr Phe Leu Pro Ser Ala Asp Glu Ile Tyr Asp Cys Lys Val Glu			
	180	185	190
His Trp Gly Leu Asp Glu Pro Leu Leu Lys His Trp Glu Pro Glu Ile			
	195	200	205
Pro Thr Pro Met Ser Glu Leu Thr Glu Thr Val Val Cys Ala Leu Gly			
	210	215	220
Leu Ser Val Gly Leu Val Gly Ile Val Val Gly Thr Val Leu Ile Ile			
	225	230	235
Arg Gly Leu Arg Ser Val Gly Ala Ser Arg His Gln Gly Pro Leu			
	245	250	255

<210> SEQ ID NO 43
 <211> LENGTH: 538
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence HLADQB10302-P2A-HLADQBA103

<400> SEQUENCE: 43

Met Ser Trp Lys Lys Ala Leu Arg Ile Pro Gly Gly Leu Arg Val Ala			
1	5	10	15

Thr	Val	Thr	Leu	Met	Leu	Ala	Met	Leu	Ser	Thr	Pro	Val	Ala	Glu	Gly
			20					25					30		
Arg	Asp	Ser	Pro	Glu	Asp	Phe	Val	Tyr	Gln	Phe	Lys	Gly	Met	Cys	Tyr
		35					40					45			
Phe	Thr	Asn	Gly	Thr	Glu	Arg	Val	Arg	Leu	Val	Thr	Arg	Tyr	Ile	Tyr
	50					55					60				
Asn	Arg	Glu	Glu	Tyr	Ala	Arg	Phe	Asp	Ser	Asp	Val	Gly	Val	Tyr	Arg
65					70					75					80
Ala	Val	Thr	Pro	Leu	Gly	Pro	Pro	Ala	Ala	Glu	Tyr	Trp	Asn	Ser	Gln
				85					90					95	
Lys	Glu	Val	Leu	Glu	Arg	Thr	Arg	Ala	Glu	Leu	Asp	Thr	Val	Cys	Arg
			100					105					110		
His	Asn	Tyr	Gln	Leu	Glu	Leu	Arg	Thr	Thr	Leu	Gln	Arg	Arg	Val	Glu
	115						120					125			
Pro	Thr	Val	Thr	Ile	Ser	Pro	Ser	Arg	Thr	Glu	Ala	Leu	Asn	His	His
	130					135					140				
Asn	Leu	Leu	Val	Cys	Ser	Val	Thr	Asp	Phe	Tyr	Pro	Ala	Gln	Ile	Lys
145				150						155					160
Val	Arg	Trp	Phe	Arg	Asn	Asp	Gln	Glu	Glu	Thr	Thr	Gly	Val	Val	Ser
				165					170					175	
Thr	Pro	Leu	Ile	Arg	Asn	Gly	Asp	Trp	Thr	Phe	Gln	Ile	Leu	Val	Met
		180						185					190		
Leu	Glu	Met	Thr	Pro	Gln	Arg	Gly	Asp	Val	Tyr	Thr	Cys	His	Val	Glu
	195						200					205			
His	Pro	Ser	Leu	Gln	Asn	Pro	Ile	Ile	Val	Glu	Trp	Arg	Ala	Gln	Ser
	210					215					220				
Glu	Ser	Ala	Gln	Ser	Lys	Met	Leu	Ser	Gly	Ile	Gly	Gly	Phe	Val	Leu
225					230					235					240
Gly	Leu	Ile	Phe	Leu	Gly	Leu	Gly	Leu	Ile	Ile	His	His	Arg	Ser	Gln
				245					250					255	
Lys	Gly	Leu	Leu	His	Gly	Ser	Gly	Ala	Thr	Asn	Phe	Ser	Leu	Leu	Lys
		260						265					270		
Gln	Ala	Gly	Asp	Val	Glu	Glu	Asn	Pro	Gly	Pro	Met	Ile	Leu	Asn	Lys
	275						280					285			
Ala	Leu	Met	Leu	Gly	Ala	Leu	Ala	Leu	Thr	Thr	Val	Met	Ser	Pro	Cys
	290					295					300				
Gly	Gly	Glu	Asp	Ile	Val	Ala	Asp	His	Val	Ala	Ser	Tyr	Gly	Val	Asn
305				310						315					320
Leu	Tyr	Gln	Ser	Tyr	Gly	Pro	Ser	Gly	Gln	Tyr	Ser	His	Glu	Phe	Asp
				325					330					335	
Gly	Asp	Glu	Glu	Phe	Tyr	Val	Asp	Leu	Glu	Arg	Lys	Glu	Thr	Val	Trp
		340						345					350		
Gln	Leu	Pro	Leu	Phe	Arg	Arg	Phe	Arg	Arg	Phe	Asp	Pro	Gln	Phe	Ala
	355						360					365			
Leu	Thr	Asn	Ile	Ala	Val	Leu									

-continued

435	440	445	
Lys Ser Asp His Ser Phe Phe Lys Ile Ser Tyr Leu Thr Phe Leu Pro			
450	455	460	
Ser Ala Asp Glu Ile Tyr Asp Cys Lys Val Glu His Trp Gly Leu Asp			
465	470	475	480
Glu Pro Leu Leu Lys His Trp Glu Pro Glu Ile Pro Thr Pro Met Ser			
	485	490	495
Glu Leu Thr Glu Thr Val Val Cys Ala Leu Gly Leu Ser Val Gly Leu			
	500	505	510
Val Gly Ile Val Val Gly Thr Val Leu Ile Ile Arg Gly Leu Arg Ser			
	515	520	525
Val Gly Ala Ser Arg His Gln Gly Pro Leu			
530	535		
 <210> SEQ ID NO 44			
<211> LENGTH: 66			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic sequence Porcine teschovirus-1 2A			
(P2A) peptide			
 <400> SEQUENCE: 44			
ggcagcggcg ccaccaactt tagcctgctg aaacaggctg gcgacgtgga agagaacccc			60
ggacct			66
 <210> SEQ ID NO 45			
<211> LENGTH: 66			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic sequence Porcine teschovirus-1 2A			
(P2A) peptide			
 <400> SEQUENCE: 45			
ggctctggcg ccaccaactt tagcctgctg aaacaggctg gcgacgtgga agagaacccc			60
ggacct			66
 <210> SEQ ID NO 46			
<211> LENGTH: 63			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic sequence Thoseaasigna virus 2A (T2A)			
peptide			
 <400> SEQUENCE: 46			
ggaagcggag agggcagagg aagtctgcta acatgcggtg acgtcgagga gaatcctgga			60
cct			63
 <210> SEQ ID NO 47			
<211> LENGTH: 69			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic sequence Equine rhinitis A virus			
(ERAV) 2A (E2A) peptide			
 <400> SEQUENCE: 47			
ggaagcggac agtgtactaa ttatgctctc ttgaaattgg ctggagatgt tgagagcaac			60
cctggacct			69

-continued

<210> SEQ ID NO 48
 <211> LENGTH: 75
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence Foot-and-Mouth disease virus
 2A (F2A) peptide

<400> SEQUENCE: 48

ggaagcggag tgaacacagac ttggaatttt gaccttctca agttggcggg agacgtggag 60
 tccaaccctg gacct 75

<210> SEQ ID NO 49
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence Porcine teschovirus-1 2A
 (P2A)

<400> SEQUENCE: 49

Gly Ser Gly Ala Thr Asn Phe Ser Leu Leu Lys Gln Ala Gly Asp Val
 1 5 10 15
 Glu Glu Asn Pro Gly Pro
 20

<210> SEQ ID NO 50
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence Thoseaasigna virus 2A (T2A)
 peptide

<400> SEQUENCE: 50

Leu Glu Gly Gly Gly Glu Gly Arg Gly Ser Leu Leu Thr Cys Gly Asp
 1 5 10 15
 Val Glu Glu Asn Pro Gly Pro Arg
 20

<210> SEQ ID NO 51
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence Equine rhinitis A virus
 (ERAV) 2A (E2A) peptide

<400> SEQUENCE: 51

Gln Cys Thr Asn Tyr Ala Leu Leu Lys Leu Ala Gly Asp Val Glu Ser
 1 5 10 15
 Asn Pro Gly Pro
 20

<210> SEQ ID NO 52
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence Foot-and-Mouth disease virus
 2A (F2A) peptide

<400> SEQUENCE: 52

Gly Ser Gly Val Lys Gln Thr Leu Asn Phe Asp Leu Leu Lys Leu Ala

-continued

```

1           5           10           15
Gly Asp Val Glu Ser Asn Pro Gly Pro
      20             25

<210> SEQ ID NO 53
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence Glycine-Serine linker

<400> SEQUENCE: 53

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1           5           10           15

<210> SEQ ID NO 54
<211> LENGTH: 18
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence Glycine-Serine linker

<400> SEQUENCE: 54

Gly Ser Thr Ser Gly Gly Gly Ser Gly Gly Gly Ser Gly Gly Gly Gly
1           5           10           15

Ser Ser

<210> SEQ ID NO 55
<211> LENGTH: 271
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR1 alpha chain (full-
length)

<400> SEQUENCE: 55

Met Glu Thr Leu Leu Gly Leu Leu Ile Leu Trp Leu Gln Leu Gln Trp
1           5           10           15

Val Ser Ser Lys Gln Glu Val Thr Gln Ile Pro Ala Ala Leu Ser Val
      20             25             30

Pro Glu Gly Glu Asn Leu Val Leu Asn Cys Ser Phe Thr Asp Ser Ala
      35             40             45

Ile Tyr Asn Leu Gln Trp Phe Arg Gln Asp Pro Gly Lys Gly Leu Thr
      50             55             60

Ser Leu Leu Leu Ile Gln Ser Ser Gln Arg Glu Gln Thr Ser Gly Arg
      65             70             75             80

Leu Asn Ala Ser Leu Asp Lys Ser Ser Gly Arg Ser Thr Leu Tyr Ile
      85             90             95

Ala Ala Ser Gln Pro Gly Asp Ser Ala Thr Tyr Leu Cys Ala Val Arg
      100            105            110

Arg Gly Asn Asn Asp Met Arg Phe Gly Ala Gly Thr Arg Leu Thr Val
      115            120            125

Lys Pro Asn Ile Gln Asn Pro Asp Pro Ala Val Tyr Gln Leu Arg Asp
      130            135            140

Ser Lys Ser Ser Asp Lys Ser Val Cys Leu Phe Thr Asp Phe Asp Ser
      145            150            155            160

Gln Thr Asn Val Ser Gln Ser Lys Asp Ser Asp Val Tyr Ile Thr Asp
      165            170            175

Lys Cys Val Leu Asp Met Arg Ser Met Asp Phe Lys Ser Asn Ser Ala
      180            185            190

```

-continued

Val Ala Trp Ser Asn Lys Ser Asp Phe Ala Cys Ala Asn Ala Phe Asn
 195 200 205
 Asn Ser Ile Ile Pro Glu Asp Thr Phe Phe Pro Ser Pro Glu Ser Ser
 210 215 220
 Cys Asp Val Lys Leu Val Glu Lys Ser Phe Glu Thr Asp Thr Asn Leu
 225 230 235 240
 Asn Phe Gln Asn Leu Ser Val Ile Gly Phe Arg Ile Leu Leu Leu Lys
 245 250 255
 Val Ala Gly Phe Asn Leu Leu Met Thr Leu Arg Leu Trp Ser Ser
 260 265 270

<210> SEQ ID NO 56
 <211> LENGTH: 269
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2-1 alpha chain
 (full-length)
 <400> SEQUENCE: 56

Met Arg Leu Val Ala Arg Val Thr Val Phe Leu Thr Phe Gly Thr Ile
 1 5 10 15
 Ile Asp Ala Lys Thr Thr Gln Pro Thr Ser Met Asp Cys Ala Glu Gly
 20 25 30
 Arg Ala Ala Asn Leu Pro Cys Asn His Ser Thr Ile Ser Gly Asn Glu
 35 40 45
 Tyr Val Tyr Trp Tyr Arg Gln Ile His Ser Gln Gly Pro Gln Tyr Ile
 50 55 60
 Ile His Gly Leu Lys Asn Asn Glu Thr Asn Glu Met Ala Ser Leu Ile
 65 70 75 80
 Ile Thr Glu Asp Arg Lys Ser Ser Thr Leu Ile Leu Pro His Ala Thr
 85 90 95
 Leu Arg Asp Thr Ala Val Tyr Tyr Cys Ile Val Arg Ala Tyr Ser Gly
 100 105 110
 Tyr Ser Thr Leu Thr Phe Gly Lys Gly Thr Met Leu Leu Val Ser Pro
 115 120 125
 Asp Ile Gln Asn Pro Asp Pro Ala Val Tyr Gln Leu Arg Asp Ser Lys
 130 135 140
 Ser Ser Asp Lys Ser Val Cys Leu Phe Thr Asp Phe Asp Ser Gln Thr
 145 150 155 160
 Asn Val Ser Gln Ser Lys Asp Ser Asp Val Tyr Ile Thr Asp Lys Cys
 165 170 175
 Val Leu Asp Met Arg Ser Met Asp Phe Lys Ser Asn Ser Ala Val Ala
 180 185 190
 Trp Ser Asn Lys Ser Asp Phe Ala Cys Ala Asn Ala Phe Asn Asn Ser
 195 200 205
 Ile Ile Pro Glu Asp Thr Phe Phe Pro Ser Pro Glu Ser Ser Cys Asp
 210 215 220
 Val Lys Leu Val Glu Lys Ser Phe Glu Thr Asp Thr Asn Leu Asn Phe
 225 230 235 240
 Gln Asn Leu Ser Val Ile Gly Phe Arg Ile Leu Leu Leu Lys Val Ala
 245 250 255
 Gly Phe Asn Leu Leu Met Thr Leu Arg Leu Trp Ser Ser
 260 265

-continued

```

<210> SEQ ID NO 57
<211> LENGTH: 276
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-2 alpha chain
      (full-length)

<400> SEQUENCE: 57
Met Lys Ser Leu Arg Val Leu Leu Val Ile Leu Trp Leu Gln Leu Ser
 1             5             10             15
Trp Val Trp Ser Gln Gln Lys Glu Val Glu Gln Asn Ser Gly Pro Leu
      20             25             30
Ser Val Pro Glu Gly Ala Ile Ala Ser Leu Asn Cys Thr Tyr Ser Asp
      35             40             45
Arg Gly Ser Gln Ser Phe Phe Trp Tyr Arg Gln Tyr Ser Gly Lys Ser
 50             55             60
Pro Glu Leu Ile Met Phe Ile Tyr Ser Asn Gly Asp Lys Glu Asp Gly
 65             70             75             80
Arg Phe Thr Ala Gln Leu Asn Lys Ala Ser Gln Tyr Val Ser Leu Leu
      85             90             95
Ile Arg Asp Ser Gln Pro Ser Asp Ser Ala Thr Tyr Leu Cys Ala Val
      100            105            110
Ile Thr Leu Asn Asn Asn Ala Gly Asn Met Leu Thr Phe Gly Gly Gly
      115            120            125
Thr Arg Leu Met Val Lys Pro His Ile Gln Asn Pro Asp Pro Ala Val
      130            135            140
Tyr Gln Leu Arg Asp Ser Lys Ser Ser Asp Lys Ser Val Cys Leu Phe
      145            150            155            160
Thr Asp Phe Asp Ser Gln Thr Asn Val Ser Gln Ser Lys Asp Ser Asp
      165            170            175
Val Tyr Ile Thr Asp Lys Cys Val Leu Asp Met Arg Ser Met Asp Phe
      180            185            190
Lys Ser Asn Ser Ala Val Ala Trp Ser Asn Lys Ser Asp Phe Ala Cys
      195            200            205
Ala Asn Ala Phe Asn Asn Ser Ile Ile Pro Glu Asp Thr Phe Phe Pro
      210            215            220
Ser Pro Glu Ser Ser Cys Asp Val Lys Leu Val Glu Lys Ser Phe Glu
      225            230            235            240
Thr Asp Thr Asn Leu Asn Phe Gln Asn Leu Ser Val Ile Gly Phe Arg
      245            250            255
Ile Leu Leu Leu Lys Val Ala Gly Phe Asn Leu Leu Met Thr Leu Arg
      260            265            270
Leu Trp Ser Ser
      275

<210> SEQ ID NO 58
<211> LENGTH: 272
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR3 alpha chain (full-
      length)

<400> SEQUENCE: 58
Met Lys Ser Leu Arg Val Leu Leu Val Ile Leu Trp Leu Gln Leu Ser
 1             5             10             15
Trp Val Trp Ser Gln Gln Lys Glu Val Glu Gln Asn Ser Gly Pro Leu

```


-continued

20						25						30					
Ser	Val	Pro	Glu	Gly	Ala	Ile	Ala	Ser	Leu	Asn	Cys	Thr	Tyr	Ser	Asp		
35						40						45					
Arg	Gly	Ser	Gln	Ser	Phe	Phe	Trp	Tyr	Arg	Gln	Tyr	Ser	Gly	Lys	Ser		
50						55						60					
Pro	Glu	Leu	Ile	Met	Phe	Ile	Tyr	Ser	Asn	Gly	Asp	Lys	Glu	Asp	Gly		
65						70						75					
Arg	Phe	Thr	Ala	Gln	Leu	Asn	Lys	Ala	Ser	Gln	Tyr	Val	Ser	Leu	Leu		
85						90						95					
Ile	Arg	Asp	Ser	Gln	Pro	Ser	Asp	Ser	Ala	Thr	Tyr	Leu	Cys	Ala	Val		
100						105						110					
Thr	Ser	Asn	Ala	Gly	Lys	Ser	Thr	Phe	Gly	Asp	Gly	Thr	Thr	Leu	Thr		
115						120						125					
Val	Lys	Pro	Asn	Ile	Gln	Asn	Pro	Asp	Pro	Ala	Val	Tyr	Gln	Leu	Arg		
130						135						140					
Asp	Ser	Lys	Ser	Ser	Asp	Lys	Ser	Val	Cys	Leu	Phe	Thr	Asp	Phe	Asp		
145						150						155					
Ser	Gln	Thr	Asn	Val	Ser	Gln	Ser	Lys	Asp	Ser	Asp	Val	Tyr	Ile	Thr		
165						170						175					
Asp	Lys	Cys	Val	Leu	Asp	Met	Arg	Ser	Met	Asp	Phe	Lys	Ser	Asn	Ser		
180						185						190					
Ala	Val	Ala	Trp	Ser	Asn	Lys	Ser	Asp	Phe	Ala	Cys	Ala	Asn	Ala	Phe		
195						200						205					
Asn	Asn	Ser	Ile	Ile	Pro	Glu	Asp	Thr	Phe	Phe	Pro	Ser	Pro	Glu	Ser		
210						215						220					
Ser	Cys	Asp	Val	Lys	Leu	Val	Glu	Lys	Ser	Phe	Glu	Thr	Asp	Thr	Asn		
225						230						235					
Leu	Asn	Phe	Gln	Asn	Leu	Ser	Val	Ile	Gly	Phe	Arg	Ile	Leu	Leu	Leu		
245						250						255					
Lys	Val	Ala	Gly	Phe	Asn	Leu	Leu	Met	Thr	Leu	Arg	Leu	Trp	Ser	Ser		
260						265						270					
<210> SEQ ID NO 59																	
<211> LENGTH: 309																	
<212> TYPE: PRT																	
<213> ORGANISM: Artificial Sequence																	
<220> FEATURE:																	
<223> OTHER INFORMATION: Synthetic sequence TCR1 beta chain (full-length)																	
<400> SEQUENCE: 59																	
Met	Gly	Ile	Arg	Leu	Leu	Cys	Arg	Val	Ala	Phe	Cys	Phe	Leu	Ala	Val		
1				5				10				15					
Gly	Leu	Val	Asp	Val	Lys	Val	Thr	Gln	Ser	Ser	Arg	Tyr	Leu	Val	Lys		
20				25				30									
Arg	Thr	Gly	Glu	Lys	Val	Phe	Leu	Glu	Cys	Val	Gln	Asp	Met	Asp	His		
35				40				45									
Glu	Asn	Met	Phe	Trp	Tyr	Arg	Gln	Asp	Pro	Gly	Leu	Gly	Leu	Arg	Leu		
50				55				60									
Ile	Tyr	Phe	Ser	Tyr	Asp	Val	Lys	Met	Lys	Glu	Lys	Gly	Asp	Ile	Pro		
65				70				75				80					
Glu	Gly	Tyr	Ser	Val	Ser	Arg	Glu	Lys	Lys	Glu	Arg	Phe	Ser	Leu	Ile		
85				90				95									
Leu	Glu	Ser	Ala	Ser	Thr	Asn	Gln	Thr	Ser	Met	Tyr	Leu	Cys	Ala	Ser		
100				105				110									

-continued

```

Asn Glu Gly Asn Ser Gly Asn Thr Ile Tyr Phe Gly Glu Gly Ser Trp
   115                               120               125

Leu Thr Val Val Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala
   130                               135               140

Val Phe Glu Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr
   145                               150               155               160

Leu Val Cys Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser
   165                               170               175

Trp Trp Val Asn Gly Lys Glu Val His Ser Gly Val Cys Thr Asp Pro
   180                               185               190

Gln Pro Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu
   195                               200               205

Ser Ser Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn
   210                               215               220

His Phe Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu
   225                               230               235               240

Trp Thr Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu
   245                               250               255

Ala Trp Gly Arg Ala Asp Cys Gly Phe Thr Ser Val Ser Tyr Gln Gln
   260                               265               270

Gly Val Leu Ser Ala Thr Ile Leu Tyr Glu Ile Leu Leu Gly Lys Ala
   275                               280               285

Thr Leu Tyr Ala Val Leu Val Ser Ala Leu Val Leu Met Ala Met Val
   290                               295               300

Lys Arg Lys Asp Phe
305

```

```

<210> SEQ ID NO 60
<211> LENGTH: 307
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-1 and 2-2 beta chain
(full-length)

```

```

<400> SEQUENCE: 60

```

```

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1           5           10           15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20           25           30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35           40           45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50           55           60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65           70           75           80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85           90           95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
100           105           110

Gly Ala Arg Gln Ile Pro Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr
115           120           125

Val Val Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe
130           135           140

Glu Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val
145           150           155           160

```

-continued

Cys Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp
 165 170 175
 Val Asn Gly Lys Glu Val His Ser Gly Val Cys Thr Asp Pro Gln Pro
 180 185 190
 Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser
 195 200 205
 Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe
 210 215 220
 Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr
 225 230 235 240
 Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp
 245 250 255
 Gly Arg Ala Asp Cys Gly Phe Thr Ser Val Ser Tyr Gln Gln Gly Val
 260 265 270
 Leu Ser Ala Thr Ile Leu Tyr Glu Ile Leu Leu Gly Lys Ala Thr Leu
 275 280 285
 Tyr Ala Val Leu Val Ser Ala Leu Val Leu Met Ala Met Val Lys Arg
 290 295 300
 Lys Asp Phe
 305

<210> SEQ ID NO 61
 <211> LENGTH: 317
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 beta chain (full-length)
 <400> SEQUENCE: 61

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1 5 10 15
 Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20 25 30
 Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35 40 45
 Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50 55 60
 Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65 70 75 80
 Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85 90 95
 Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
 100 105 110
 Ser Leu Ser Ala Ala Gly Gly Gly Tyr Gly Tyr Thr Phe Asn Tyr Gly
 115 120 125
 Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr Val Val Glu Asp Leu Asn
 130 135 140
 Lys Val Phe Pro Pro Glu Val Ala Val Phe Glu Pro Ser Glu Ala Glu
 145 150 155 160
 Ile Ser His Thr Gln Lys Ala Thr Leu Val Cys Leu Ala Thr Gly Phe
 165 170 175
 Phe Pro Asp His Val Glu Leu Ser Trp Trp Val Asn Gly Lys Glu Val
 180 185 190
 His Ser Gly Val Cys Thr Asp Pro Gln Pro Leu Lys Glu Gln Pro Ala

-continued

195	200	205
Leu Asn Asp Ser Arg Tyr Cys	Leu Ser Ser Arg	Leu Arg Val Ser Ala
210	215	220
Thr Phe Trp Gln Asn Pro Arg	Asn His Phe Arg	Cys Gln Val Gln Phe
225	230	235 240
Tyr Gly Leu Ser Glu Asn Asp	Glu Trp Thr Gln Asp	Arg Ala Lys Pro
	245	250 255
Val Thr Gln Ile Val Ser Ala	Glu Ala Trp Gly Arg	Ala Asp Cys Gly
	260	265 270
Phe Thr Ser Val Ser Tyr Gln	Gln Gly Val Leu Ser	Ala Thr Ile Leu
	275	280 285
Tyr Glu Ile Leu Leu Gly Lys	Ala Thr Leu Tyr	Ala Val Leu Val Ser
	290	295 300
Ala Leu Val Leu Met Ala Met	Val Lys Arg Lys	Asp Phe
305	310	315

<210> SEQ ID NO 62
 <211> LENGTH: 816
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 alpha chain (full-length)

<400> SEQUENCE: 62

```

atggaaacac tgtgggacct gctgacctg ttggtgcaac tgcaatgggt gtccagcaag    60
caagaagtga ctcagatccc tgccgctctg tctgtgacct agggcgaaaa cctggctctg    120
aactgcagct tcaccgcagc cgccatctac aacctgcagt gggtcaggca ggatccaggc    180
aaggggctga catcctgctt gctgattcag agcagccaga gagagcagac cagcggcaga    240
ctgaatgcca gcctggataa gtcctccggc agaagcacc tgtatatcgc cgcttctcag    300
ccaggcgata gcgccacata tctgtgtgcc gtccggagag gcaacaacga catgagattc    360
ggagccggga ccagactgac cgtgaagccc aacattcaga accccgatcc tgcagtgtac    420
cagctgctgg acagcaagag cagcgacaag agcgtgtgcc tgttcaccga ctctgacagc    480
cagaccaacg tgtccagag caaggacagc gacgtgtaca tcaccgataa gtgcgtgctg    540
gacatgcgga gcattggactt caagagcaac agcggcgtgg cctgggtccaa caagagcgac    600
ttgcctgcgc ccaacgcctt caacaacagc attatcccg aggacacatt ctccaagc    660
cccagagaca gctgcgacgt gaagctgggt gaaaagagct tcgagacaga caccaacctg    720
aacttcaga acctcagcgt gatcggttc cggatcctgc tgctgaaggt ggccgggttc    780
aacctgctga tgacctgcg gctgtggtcc agctga                                816

```

<210> SEQ ID NO 63
 <211> LENGTH: 810
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2-1 alpha chain (full-length)

<400> SEQUENCE: 63

```

atgagactgg tggccagagt gacagtgttc ctgaccttcg gcaccatcat cgacgccaag    60
accacacagc ccaccagcat ggattgtgcc gagggcagag ctgccaaact gccttgtaat    120
cacagcacca tcagcgga caagtagctg tactgggtaca ggagatcca ctctcagggc    180

```

-continued

cctcagtaca tcatccacgg cctgaagaac aacgagacaa acgagatggc ctctctgac	240
atcaccgagg accgcaagag cagcaccctg attctgcctc acgccacact gagagacacc	300
gccgtgtact actgtatcgt gccggcctac agcggctaca gcacactgac atttggaag	360
ggcaccatgc tgctcgtgtc cccagacatt cagaaccccg atcctgcagt gtaccagctg	420
cgggacagca agagcagcga caagagcgtg tgcctgttca ccgacttcga cagccagacc	480
aacgtgtccc agagcaagga cagcgacgtg tacatcacgg ataagtgcgt gctggacatg	540
cggagcatgg acttcaagag caacagcgcc gtggcctggt ccaacaagag cgacttcgcc	600
tgcgccaacg ccttcaacaa cagcattatc cccgaggaca cattcttccc aagccccgag	660
agcagctgcg acgtgaagct ggtggaaaag agcttcgaga cagacaccaa cctgaacttc	720
cagaacctca gcgtgatcgg cttccggatc ctgctgctga aggtggccgg cttcaacctg	780
ctgatgaccc tgcggctgtg gtcacgtga	810

<210> SEQ ID NO 64
 <211> LENGTH: 831
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2-2 alpha chain
 (full-length)

<400> SEQUENCE: 64

atgaagtccc tgagagtgtc gctggtcac ctgtggctgc agctgtcttg ggtctggtcc	60
cagcagaaaag aggtggaaca gaacagcggc cctctgtctg ttctgaagg cgctatcgcc	120
tctctgaatt gcacctacag cgacagaggc agccagagct tcttctggta tcggcagtac	180
agcggcaaga gccccagct gatcatgttc atctacagca acggcgacaa agaggacggc	240
cgggtttacag cccagctgaa caaggccagc cagtacgtgt cactgctgat cagagacagc	300
cagcctagcg acagcgccac ctatctgtgt gccgtgatca ccctgaacaa caacgccggc	360
aacatgctga cctttggcgg cggaacacgg ctgatggta agccccacat tcagaacccc	420
gatectgcag tgtaccagct gccggacagc aagagcagcg acaagagcgt gtgcctgttc	480
accgacttcg acagccagac caacgtgtcc cagagcaagg acagcgacgt gtacatcac	540
gataagtgcg tgctggacat gcggagcatg gacttcaaga gcaacagcgc cgtggcctgg	600
tccaacaaga gcgacttcgc ctgcgccaac gccttcaaca acagcattat ccccaggac	660
acattcttcc caagccccga gacgagctgc gacgtgaagc tgggtgaaaa gagcttcgag	720
acagacacca acctgaactt ccagaacctc agcgtgatcg gcttccggat cctgctgctg	780
aaggtggccg gcttcaacct gctgatgacc ctgcggctgt ggtccagctg a	831

<210> SEQ ID NO 65
 <211> LENGTH: 819
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 alpha chain (full-length)

<400> SEQUENCE: 65

atgaagtccc tgagagtgtc gctggtcac ctgtggctgc agctgtcttg ggtctggtcc	60
cagcagaaaag aggtggaaca gaacagcggc cctctgtccg ttctgaagg cgccattgct	120
agcctgaatt gcacctacag cgaccggggc agccagagct tcttctggta taggcagtac	180
agcggcaaga gccccagct gatcatgttc atctacagca acggcgacaa agaggacggc	240

-continued

cggtttacag cccagctgaa caaggccagc cagtacgtgt cactgctgat cagagacagc	300
cagcctagcg acagcgccac ctatctgtgt gccgtgacaa gcaatgccgg caagtccaca	360
tttggcgacg gcacaacct gaccgtgaag cccaacattc agaaccgccga tcctgcagtg	420
taccagctgc gggacagcaa gagcagcgac aagagcgtgt gcctgttcac cgacttcgac	480
agccagacca acgtgtccca gagcaaggac agcgacgtgt acatcaccga taagtgcgtg	540
ctggacatgc ggagcatgga cttcaagagc aacagcgccg tggcctggtc caacaagagc	600
gacttcgcct gcgccaacgc cttcaacaac agcattatcc ccgaggacac attcttccca	660
agccccgaga gcagctgcga cgtgaagctg gtggaaaaga gcttcgagac agacaccaac	720
ctgaacttcc agaacctcag cgtgatcggc ttccggatcc tgctgctgaa ggtggccggc	780
ttcaacctgc tgatgacct gcggctgtgg tccagctga	819

<210> SEQ ID NO 66
 <211> LENGTH: 927
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 beta chain (full-length)

<400> SEQUENCE: 66

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttg cctggtggac	60
gtgaaagtga cccagagcag cagatacttg gtcaagagaa ccggcgagaa ggtgttcttg	120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg	180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaagg cgacatcccc	240
gagggttaca gcgtgtccag agagaagaaa gagcgggttca gcctgatect ggaaagcgcc	300
agcaccaacc agaccagcat gtacctgtgc gccagcaacg agggcaatag cggaacacc	360
atctacttcc gcgaaggcag ctggctgacc gtggtggaag atctgaacaa agtgttccct	420
ccagaggtgg ccgtgttcga gccttctgag gccagatca gccacacaca gaaagccaca	480
ctcgtgtgcc tggccaccgg cttttttccc gatcacgtgg aactgtcttg gtgggtcaac	540
ggcaaagagg tgcacagcgg cgtctgtacc gatecctcag ctctgaaaga gcagcccgct	600
ctgaacgact ccagatactg cctgagcagc aggtctgagag tgtccgccac cttctggcag	660
aaaccacagaa accacttcag atgccagggt cagttctacg gcctgagcga gaacgatgag	720
tggagccagg atagagccaa gcctgtgaca cagatcgtgt ctgccgaagc ctggggcaga	780
gccgatttgt gctttacctc cgtgtcctat cagcagggcg tgctgtctgc caccatctg	840
tatgagatcc tgctgggcaa agccactctg tacgccgtgc tgggtgtctgc cctggtgctg	900
atggccatgg tcaagcggaa ggatttt	927

<210> SEQ ID NO 67
 <211> LENGTH: 921
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2-1 and 2-2 beta chain (full-length)

<400> SEQUENCE: 67

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttg cctggtggac	60
gtgaaagtga cccagagcag cagatacttg gtcaagagaa ccggcgagaa ggtgttcttg	120

-continued

gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg	180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc	240
gagggctaca gcgtgtccag agagaagaaa gagcgggtca gcctgacccg gaaaagcgcc	300
agcaccaacc agaccagcat gtacctgtgt gccagcggcg ccagacagat cccttacaca	360
tttggtccg gcaccagact gaccgtgggtg gaagatctga acaaagtgtt ccctccagag	420
gtggccgtgt tcgagccttc tgaggccgag atcagccaca cacagaaagc cacactcgtg	480
tgcttgcca ccggtttttt tcccgatcac gtggaactgt cttggtgggt caacggcaaa	540
gaggtgcaca gcgcgctctg taccgatcct cagcctctga aagagcagcc cgctctgaac	600
gactccagat actgctgag cagcaggctg agagtgtccg ccacctctg gcagaacccc	660
agaaaccact tcagatgcca ggtgcagttc tacggcctga gcgagaacga tgagtggacc	720
caggatagag ccaagcctgt gacacagatc gtgtctgccg aagcctgggg cagagccgat	780
tgtggcttta cctcgtgtc ctatcagcag ggcgtgctgt ctgccaccat cctgtatgag	840
atcctgctgg gcaaagccac tctgtacgcc gtgctggtgt ctgccctggt gctgatggcc	900
atggtcaagc ggaaggattt t	921

<210> SEQ ID NO 68
 <211> LENGTH: 951
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 beta chain (full-length)
 <400> SEQUENCE: 68

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttgg cctggtggac	60
gtgaaagtga cccagagcag cagatacctg gtcaagagaa ccggcgagaa ggtgttccctg	120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg	180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc	240
gagggctaca gcgtgtccag agagaagaaa gagcgggtca gcctgacccg gaaaagcgcc	300
agcaccaacc agaccagcat gtacctgtgt gccagctctc tgtctgctgc cggcggaggc	360
tatggctaca ccttcaatta cggtacaca ttcggcagcg gcaccagact gaccgtggtg	420
gaagatctga acaaagtgtt ccctccagag gtggccgtgt tcgagccttc tgaggccgag	480
atcagccaca cacagaaagc cacactcgtg tgcttgcca ccggtttttt tcccgatcac	540
gtggaactgt cttggtgggt caacggcaaa gaggtgcaca gcgcgctctg taccgatcct	600
cagcctctga aagagcagcc cgctctgaac gactccagat actgcctgag cagcaggctg	660
agagtgtccg ccacctctg gcagaacccc agaaaccact tcagatgcca ggtgcagttc	720
tacggcctga gcgagaacga tgagtggacc caggatagag ccaagcctgt gacacagatc	780
gtgtctgccg aagcctgggg cagagccgat tgtggcttta cctcgtgtc ctatcagcag	840
ggcgtgctgt ctgccaccat cctgtatgag atcctgctgg gcaaagccac tctgtacgcc	900
gtgctggtgt ctgccctggt gctgatggcc atggtcaagc ggaaggattt t	951

<210> SEQ ID NO 69
 <211> LENGTH: 602
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR1 (beta-P2A-alpha)

-continued

<400> SEQUENCE: 69

```

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
1           5           10           15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
20           25           30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
35           40           45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
50           55           60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
65           70           75           80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
85           90           95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
100          105          110

Asn Glu Gly Asn Ser Gly Asn Thr Ile Tyr Phe Gly Glu Gly Ser Trp
115          120          125

Leu Thr Val Val Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala
130          135          140

Val Phe Glu Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr
145          150          155          160

Leu Val Cys Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser
165          170          175

Trp Trp Val Asn Gly Lys Glu Val His Ser Gly Val Cys Thr Asp Pro
180          185          190

Gln Pro Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu
195          200          205

Ser Ser Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn
210          215          220

His Phe Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu
225          230          235          240

Trp Thr Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu
245          250          255

Ala Trp Gly Arg Ala Asp Cys Gly Phe Thr Ser Val Ser Tyr Gln Gln
260          265          270

Gly Val Leu Ser Ala Thr Ile Leu Tyr Glu Ile Leu Leu Gly Lys Ala
275          280          285

Thr Leu Tyr Ala Val Leu Val Ser Ala Leu Val Leu Met Ala Met Val
290          295          300

Lys Arg Lys Asp Phe Gly Ser Gly Ala Thr Asn Phe Ser Leu Leu Lys
305          310          315          320

Gln Ala Gly Asp Val Glu Glu Asn Pro Gly Pro Met Glu Thr Leu Leu
325          330          335

Gly Leu Leu Ile Leu Trp Leu Gln Leu Gln Trp Val Ser Ser Lys Gln
340          345          350

Glu Val Thr Gln Ile Pro Ala Ala Leu Ser Val Pro Glu Gly Glu Asn
355          360          365

Leu Val Leu Asn Cys Ser Phe Thr Asp Ser Ala Ile Tyr Asn Leu Gln
370          375          380

Trp Phe Arg Gln Asp Pro Gly Lys Gly Leu Thr Ser Leu Leu Leu Ile
385          390          395          400

Gln Ser Ser Gln Arg Glu Gln Thr Ser Gly Arg Leu Asn Ala Ser Leu
405          410          415

```


-continued

Asp Lys Ser Ser Gly Arg Ser Thr Leu Tyr Ile Ala Ala Ser Gln Pro
 420 425 430

Gly Asp Ser Ala Thr Tyr Leu Cys Ala Val Arg Arg Gly Asn Asn Asp
 435 440 445

Met Arg Phe Gly Ala Gly Thr Arg Leu Thr Val Lys Pro Asn Ile Gln
 450 455 460

Asn Pro Asp Pro Ala Val Tyr Gln Leu Arg Asp Ser Lys Ser Ser Asp
 465 470 475 480

Lys Ser Val Cys Leu Phe Thr Asp Phe Asp Ser Gln Thr Asn Val Ser
 485 490 495

Gln Ser Lys Asp Ser Asp Val Tyr Ile Thr Asp Lys Cys Val Leu Asp
 500 505 510

Met Arg Ser Met Asp Phe Lys Ser Asn Ser Ala Val Ala Trp Ser Asn
 515 520 525

Lys Ser Asp Phe Ala Cys Ala Asn Ala Phe Asn Asn Ser Ile Ile Pro
 530 535 540

Glu Asp Thr Phe Phe Pro Ser Pro Glu Ser Ser Cys Asp Val Lys Leu
 545 550 555 560

Val Glu Lys Ser Phe Glu Thr Asp Thr Asn Leu Asn Phe Gln Asn Leu
 565 570 575

Ser Val Ile Gly Phe Arg Ile Leu Leu Leu Lys Val Ala Gly Phe Asn
 580 585 590

Leu Leu Met Thr Leu Arg Leu Trp Ser Ser
 595 600

<210> SEQ ID NO 70
 <211> LENGTH: 598
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR2-1 (beta-P2A-alpha)

<400> SEQUENCE: 70

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1 5 10 15

Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20 25 30

Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35 40 45

Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50 55 60

Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65 70 75 80

Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85 90 95

Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
 100 105 110

Gly Ala Arg Gln Ile Pro Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr
 115 120 125

Val Val Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe
 130 135 140

Glu Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val
 145 150 155 160

Cys Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp
 165 170 175

Val	Asn	Gly	Lys	Glu	Val	His	Ser	Gly	Val	Cys	Thr	Asp	Pro	Gln	Pro
			180				185						190		
Leu	Lys	Glu	Gln	Pro	Ala	Leu	Asn	Asp	Ser	Arg	Tyr	Cys	Leu	Ser	Ser
			195				200						205		
Arg	Leu	Arg	Val	Ser	Ala	Thr	Phe	Trp	Gln	Asn	Pro	Arg	Asn	His	Phe
			210				215						220		
Arg	Cys	Gln	Val	Gln	Phe	Tyr	Gly	Leu	Ser	Glu	Asn	Asp	Glu	Trp	Thr
			225				230						235		
Gln	Asp	Arg	Ala	Lys	Pro	Val	Thr	Gln	Ile	Val	Ser	Ala	Glu	Ala	Trp
			245							250			255		
Gly	Arg	Ala	Asp	Cys	Gly	Phe	Thr	Ser	Val	Ser	Tyr	Gln	Gln	Gly	Val
			260				265						270		
Leu	Ser	Ala	Thr	Ile	Leu	Tyr	Glu	Ile	Leu	Leu	Gly	Lys	Ala	Thr	Leu
			275				280						285		
Tyr	Ala	Val	Leu	Val	Ser	Ala	Leu	Val	Leu	Met	Ala	Met	Val	Lys	Arg
			290				295						300		
Lys	Asp	Phe	Gly	Ser	Gly	Ala	Thr	Asn	Phe	Ser	Leu	Leu	Lys	Gln	Ala
			305				310						315		
Gly	Asp	Val	Glu	Glu	Asn	Pro	Gly	Pro	Met	Arg	Leu	Val	Ala	Arg	Val
			325							330			335		
Thr	Val	Phe	Leu	Thr	Phe	Gly	Thr	Ile	Ile	Asp	Ala	Lys	Thr	Thr	Gln
			340				345						350		
Pro	Thr	Ser	Met	Asp	Cys	Ala	Glu	Gly	Arg	Ala	Ala	Asn	Leu	Pro	Cys
			355				360						365		
Asn	His	Ser	Thr	Ile	Ser	Gly	Asn	Glu	Tyr	Val	Tyr	Trp	Tyr	Arg	Gln
			370				375						380		
Ile	His	Ser	Gln	Gly	Pro	Gln	Tyr	Ile	Ile	His	Gly	Leu	Lys	Asn	Asn
			385				390						395		
Glu	Thr	Asn	Glu	Met	Ala	Ser	Leu	Ile	Ile	Thr	Glu	Asp	Arg	Lys	Ser
			405							410			415		
Ser	Thr	Leu	Ile	Leu	Pro	His	Ala	Thr	Leu	Arg	Asp	Thr	Ala	Val	Tyr
			420				425						430		
Tyr	Cys	Ile	Val	Arg	Ala	Tyr	Ser	Gly	Tyr	Ser	Thr	Leu	Thr	Phe	Gly
			435				440						445		
Lys	Gly	Thr	Met	Leu	Leu	Val	Ser	Pro	Asp	Ile	Gln	Asn	Pro	Asp	Pro
			450				455						460		
Ala	Val	Tyr	Gln	Leu	Arg	Asp	Ser	Lys	Ser	Ser	Asp	Lys	Ser	Val	Cys
			465				470						475		
Leu	Phe	Thr	Asp	Phe	Asp	Ser	Gln	Thr	Asn	Val	Ser	Gln	Ser	Lys	Asp
			485							490			495		
Ser	Asp	Val	Tyr	Ile	Thr	Asp	Lys	Cys	Val	Leu	Asp	Met	Arg	Ser	Met
			500				505						510		
Asp	Phe	Lys	Ser	Asn	Ser	Ala	Val	Ala	Trp	Ser	Asn	Lys	Ser	Asp	Phe
			515				520						525		
Ala	Cys	Ala	Asn	Ala	Phe	Asn	Asn	Ser	Ile	Ile	Pro	Glu	Asp	Thr	Phe
			530				535						540		
Phe	Pro	Ser	Pro	Glu	Ser	Ser	Cys	Asp	Val	Lys	Leu	Val	Glu	Lys	Ser
			545				550						555		
Phe	Glu	Thr	Asp	Thr	Asn	Leu	Asn	Phe	Gln	Asn	Leu	Ser	Val	Ile	Gly
			565				570						575		
Phe	Arg	Ile	Leu	Leu	Lys	Val	Ala	Gly	Phe	Asn	Leu	Leu	Met	Thr	

-continued

Leu Arg Leu Trp Ser Ser
595

<210> SEQ ID NO 71
<211> LENGTH: 605
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TCR2-2 amino acid
(beta-P2A-alpha)

<400> SEQUENCE: 71

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
1 5 10 15
Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
20 25 30
Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
35 40 45
Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
50 55 60
Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
65 70 75 80
Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
85 90 95
Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser
100 105 110
Gly Ala Arg Gln Ile Pro Tyr Thr Phe Gly Ser Gly Thr Arg Leu Thr
115 120 125
Val Val Glu Asp Leu Asn Lys Val Phe Pro Pro Glu Val Ala Val Phe
130 135 140
Glu Pro Ser Glu Ala Glu Ile Ser His Thr Gln Lys Ala Thr Leu Val
145 150 155 160
Cys Leu Ala Thr Gly Phe Phe Pro Asp His Val Glu Leu Ser Trp Trp
165 170 175
Val Asn Gly Lys Glu Val His Ser Gly Val Cys Thr Asp Pro Gln Pro
180 185 190
Leu Lys Glu Gln Pro Ala Leu Asn Asp Ser Arg Tyr Cys Leu Ser Ser
195 200 205
Arg Leu Arg Val Ser Ala Thr Phe Trp Gln Asn Pro Arg Asn His Phe
210 215 220
Arg Cys Gln Val Gln Phe Tyr Gly Leu Ser Glu Asn Asp Glu Trp Thr
225 230 235 240
Gln Asp Arg Ala Lys Pro Val Thr Gln Ile Val Ser Ala Glu Ala Trp
245 250 255
Gly Arg Ala Asp Cys Gly Phe Thr Ser Val Ser Tyr Gln Gln Gly Val
260 265 270
Leu Ser Ala Thr Ile Leu Tyr Glu Ile Leu Leu Gly Lys Ala Thr Leu
275 280 285
Tyr Ala Val Leu Val Ser Ala Leu Val Leu Met Ala Met Val Lys Arg
290 295 300
Lys Asp Phe Gly Ser Gly Ala Thr Asn Phe Ser Leu Leu Lys Gln Ala
305 310 315 320
Gly Asp Val Glu Glu Asn Pro Gly Pro Met Lys Ser Leu Arg Val Leu
325 330 335
Leu Val Ile Leu Trp Leu Gln Leu Ser Trp Val Trp Ser Gln Gln Lys
340 345 350

-continued

Glu Val Glu Gln Asn Ser Gly Pro Leu Ser Val Pro Glu Gly Ala Ile
 355 360 365
 Ala Ser Leu Asn Cys Thr Tyr Ser Asp Arg Gly Ser Gln Ser Phe Phe
 370 375 380
 Trp Tyr Arg Gln Tyr Ser Gly Lys Ser Pro Glu Leu Ile Met Phe Ile
 385 390 395 400
 Tyr Ser Asn Gly Asp Lys Glu Asp Gly Arg Phe Thr Ala Gln Leu Asn
 405 410 415
 Lys Ala Ser Gln Tyr Val Ser Leu Leu Ile Arg Asp Ser Gln Pro Ser
 420 425 430
 Asp Ser Ala Thr Tyr Leu Cys Ala Val Ile Thr Leu Asn Asn Asn Ala
 435 440 445
 Gly Asn Met Leu Thr Phe Gly Gly Gly Thr Arg Leu Met Val Lys Pro
 450 455 460
 His Ile Gln Asn Pro Asp Pro Ala Val Tyr Gln Leu Arg Asp Ser Lys
 465 470 475 480
 Ser Ser Asp Lys Ser Val Cys Leu Phe Thr Asp Phe Asp Ser Gln Thr
 485 490 495
 Asn Val Ser Gln Ser Lys Asp Ser Asp Val Tyr Ile Thr Asp Lys Cys
 500 505 510
 Val Leu Asp Met Arg Ser Met Asp Phe Lys Ser Asn Ser Ala Val Ala
 515 520 525
 Trp Ser Asn Lys Ser Asp Phe Ala Cys Ala Asn Ala Phe Asn Asn Ser
 530 535 540
 Ile Ile Pro Glu Asp Thr Phe Phe Pro Ser Pro Glu Ser Ser Cys Asp
 545 550 555 560
 Val Lys Leu Val Glu Lys Ser Phe Glu Thr Asp Thr Asn Leu Asn Phe
 565 570 575
 Gln Asn Leu Ser Val Ile Gly Phe Arg Ile Leu Leu Leu Lys Val Ala
 580 585 590
 Gly Phe Asn Leu Leu Met Thr Leu Arg Leu Trp Ser Ser
 595 600 605

<210> SEQ ID NO 72
 <211> LENGTH: 476
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence TCR3 amino acid
 (beta-P2A-alpha)

<400> SEQUENCE: 72

Met Gly Ile Arg Leu Leu Cys Arg Val Ala Phe Cys Phe Leu Ala Val
 1 5 10 15
 Gly Leu Val Asp Val Lys Val Thr Gln Ser Ser Arg Tyr Leu Val Lys
 20 25 30
 Arg Thr Gly Glu Lys Val Phe Leu Glu Cys Val Gln Asp Met Asp His
 35 40 45
 Glu Asn Met Phe Trp Tyr Arg Gln Asp Pro Gly Leu Gly Leu Arg Leu
 50 55 60
 Ile Tyr Phe Ser Tyr Asp Val Lys Met Lys Glu Lys Gly Asp Ile Pro
 65 70 75 80
 Glu Gly Tyr Ser Val Ser Arg Glu Lys Lys Glu Arg Phe Ser Leu Ile
 85 90 95
 Leu Glu Ser Ala Ser Thr Asn Gln Thr Ser Met Tyr Leu Cys Ala Ser

-continued

100					105					110					
Ser	Leu	Ser	Ala	Ala	Gly	Gly	Gly	Tyr	Gly	Tyr	Thr	Phe	Asn	Tyr	Gly
	115						120						125		
Tyr	Thr	Phe	Gly	Ser	Gly	Thr	Arg	Leu	Thr	Val	Val	Glu	Asp	Leu	Asn
	130						135					140			
Lys	Val	Phe	Pro	Pro	Glu	Val	Ala	Val	Phe	Glu	Pro	Ser	Glu	Ala	Glu
	145					150				155					160
Ile	Ser	His	Thr	Gln	Lys	Ala	Thr	Leu	Val	Cys	Leu	Ala	Thr	Gly	Phe
				165					170						175
Phe	Pro	Asp	His	Val	Glu	Leu	Ser	Trp	Trp	Val	Asn	Gly	Lys	Glu	Val
			180					185					190		
His	Ser	Gly	Val	Cys	Thr	Asp	Pro	Gln	Pro	Leu	Lys	Glu	Gln	Pro	Ala
		195					200					205			
Leu	Asn	Asp	Ser	Arg	Tyr	Cys	Leu	Ser	Ser	Arg	Leu	Arg	Val	Ser	Ala
	210					215					220				
Thr	Phe	Trp	Gln	Asn	Pro	Arg	Asn	His	Phe	Arg	Cys	Gln	Val	Gln	Phe
	225				230					235					240
Tyr	Gly	Leu	Ser	Glu	Asn	Asp	Glu	Trp	Thr	Gln	Asp	Arg	Ala	Lys	Pro
				245					250					255	
Val	Thr	Gln	Ile	Val	Ser	Ala	Glu	Ala	Trp	Gly	Arg	Ala	Asp	Cys	Gly
			260						265				270		
Phe	Thr	Ser	Val	Ser	Tyr	Gln	Gln	Gly	Val	Leu	Ser	Ala	Thr	Ile	Leu
		275					280					285			
Tyr	Glu	Ile	Leu	Leu	Gly	Lys	Ala	Thr	Leu	Tyr	Ala	Val	Leu	Val	Ser
	290					295					300				
Ala	Leu	Val	Leu	Met	Ala	Met	Val	Lys	Arg	Lys	Asp	Phe	Gly	Ser	Gly
	305				310					315					320
Ala	Thr	Asn	Phe	Ser	Leu	Leu	Lys	Gln	Ala	Gly	Asp	Val	Glu	Glu	Asn
				325					330					335	
Pro	Gly	Pro	Met	Lys	Ser	Leu	Arg	Val	Leu	Leu	Val	Ile	Leu	Trp	Leu
			340					345					350		
Gln	Leu	Ser	Trp	Val	Trp	Ser	Gln	Gln	Lys	Glu	Val	Glu	Gln	Asn	Ser
		355					360					365			
Gly	Pro	Leu	Ser	Val	Pro	Glu	Gly	Ala	Ile	Ala	Ser	Leu	Asn	Cys	Thr
	370					375					380				
Tyr	Ser	Asp	Arg	Gly	Ser	Gln	Ser	Phe	Phe	Trp	Tyr	Arg	Gln	Tyr	Ser
	385				390					395					400
Gly	Lys	Ser	Pro	Glu	Leu	Ile	Met	Phe	Ile	Tyr	Ser	Asn	Gly	Asp	Lys
				405					410					415	
Glu	Asp	Gly	Arg	Phe	Thr	Ala	Gln	Leu	Asn	Lys	Ala	Ser	Gln	Tyr	Val
			420					425					430		
Ser	Leu	Leu	Ile	Arg	Asp	Ser	Gln	Pro	Ser	Asp	Ser	Ala	Thr	Tyr	Leu
		435					440					445			
Cys	Ala	Val	Thr	Ser	Asn	Ala	Gly	Lys	Ser	Thr	Phe	Gly	Asp	Gly	Thr
	450					455					460				
Thr	Leu	Thr	Val	Lys	Pro	Asn	Ile	Gln	Asn	Pro	Asp				
	465				470					475					

<210> SEQ ID NO 73

<211> LENGTH: 1809

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR1 (beta-P2A-alpha)

-continued

<400> SEQUENCE: 73

```

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttgg cctggtggac    60
gtgaaagtga cccagagcag cagataacctg gtcaagagaa ccggcgagaa ggtgttccctg    120
gaatgcgtgc aggacatgga ccacgagaat atgttcttgt acagacagga ccccggcctg    180
ggcctgagac tgatctactt cagctacgac gtgaagtga aggaaaaggg cgacatcccc    240
gagggctaca gcgtgtccag agagaagaaa gagcgggttca gcctgatact ggaaagcgcc    300
agcaccaacc agaccagcat gtacctgtgc gccagcaacg agggcaatag cggaacacc    360
atctacttgc gcgaaggcag ctggctgacc gtggtggaag atctgaacaa agtgttcct    420
ccagaggtgg ccgtgttcga gcttctgag gccgagatca gccacacaca gaaagccaca    480
ctcgtgtgcc tggccaccgg cttttttccc gatcacgtgg aactgtcttg gtgggtcaac    540
ggcaaagagg tgcacagcgg cgtctgtacc gatcctcagc ctctgaaaga gcagcccgct    600
ctgaacgact ccagataactg cctgagcagc aggtgagag tgctccgccac cttctggcag    660
aaccaccagaa accacttcag atgccagggt cagttctacg gcctgagcga gaacgatgag    720
tggacccagg atagagccaa gcctgtgaca cagatcgtgt ctgccgaagc ctggggcaga    780
gccgatttgt gctttacctc cgtgtcctat cagcaggcgg tgctgtctgc caccatcctg    840
tatgagatcc tgctgggcaa agccactctg tacgcctgct tgggtgtctgc cctggtgctg    900
atggccatgg tcaagcggaa ggattttggc agcggcgcca ccaactttag cctgtgaaa    960
caggctggcg acgtggaaga gaaccccgga cctatggaaa cactgtgggg cctgtgatc   1020
ctgtggctgc aactgcaatg ggtgtccagc aagcaagaag tgactcagat ccctgccgct   1080
ctgtctgtgc ctgagggcga aaacctggtc ctgaactgca gcttcaccga cagcgccatc   1140
tacaacctgc agtgggttcag gcaggatcca ggcaagggcc tgacatccct gctgtgatt   1200
cagagcagcc agagagagca gaccagcggc agactgaatg ccagcctgga taagtcctcc   1260
ggcagaagca ccctgtatat cgcgccttct cagccaggcg atagcgccac atatctgtgt   1320
gccgttcgga gaggaacaa cgacatgaga ttccggagccg gcaccagact gaccgtgaag   1380
cccaacattc agaaccccgga tcctgcagtg taccagctgc gggacagcaa gagcagcgac   1440
aagagcgtgt gcctgttcac cgacttcgac agccagacca acgtgtccca gagcaaggac   1500
agcgacgtgt acatcaccga taagtgcgtg ctggacatgc ggagcatgga cttcaagagc   1560
aacagcgccg tggcctggtc caacaagagc gacttcgcct gcgccaacgc cttcaacaac   1620
agcattatcc ccgaggacac attcttccca agccccgaga gcagctgcga cgtgaagctg   1680
gtggaaaaga gcttcgagac agacaccaac ctgaacttcc agaacctcag cgtgatcggc   1740
ttccggatcc tgctgtgtaa ggtggccggc ttcaacctgc tgatgacctc gcggctgtgg   1800
tccagctga                                     1809

```

<210> SEQ ID NO 74

<211> LENGTH: 1797

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR2-1 (beta-P2A-alpha)

<400> SEQUENCE: 74

```

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggctgttgg cctggtggac    60
gtgaaagtga cccagagcag cagataacctg gtcaagagaa ccggcgagaa ggtgttccctg    120
gaatgcgtgc aggacatgga ccacgagaat atgttcttgt acagacagga ccccggcctg    180

```

-continued

```

ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc 240
gagggctaca gcgtgtccag agagaagaaa gagcgggttca gcctgatact ggaaagcgcc 300
agcaccaacc agaccagcat gtacctgtgt gccagcggcg ccagacagat cccttacaca 360
tttggtccg gcaccagact gaccgtggtg gaagatctga acaaagtgtt ccctccagag 420
gtggccgtgt tcgagccttc tgaggccgag atcagccaca cacagaaagc cactctgtg 480
tgctggcca ccggtttttt tccgatcac gtggaactgt cttggtgggt caacggcaaa 540
gaggtgcaca gcggcgtctg taccgatcct cagcctctga aagagcagcc cgctctgaac 600
gactccagat actgctgag cagcaggctg agagtgtccg ccacctctg gcagaacccc 660
agaaaccact tcagatgcc ggtgcagttc tacggcctga gcgagaacga tgagtggacc 720
caggatagag ccaagcctgt gacacagatc gtgtctgccg aagcctgggg cagagccgat 780
tgtgcttta cctcgtgtc ctatcagcag ggcgtgtgt ctgccaccat cctgtatgag 840
atcctgctgg gcaaaaccac tctgtacgcc gtgctgggt ctgccctggt gctgatggcc 900
atggtcaagc ggaaggattt tggctctggc gccaccaact ttagcctgct gaaacaggct 960
ggcgacgtgg aagagaaccc cggacctatg agactggtgg ccagagtgc agtgttctg 1020
acctcggca ccatcatcga cgccaagacc acacagccca ccagcatgga ttgtgccgag 1080
ggcagagctg ccaacctgcc ttgtaatcac agcaccatca gcggcaacga gtacgtgtac 1140
tggtacaggc agatccactc tcagggccct cagtacatca tccacggcct gaagaacaa 1200
gagacaaacg agatggctc tctgatcacc accgaggacc gcaagagcag caccctgatt 1260
ctgctcacg ccacactgag agacaccgcc gtgtactact gtatcgtgcg ggctacagc 1320
ggctacagca cactgacatt tggcaagggc accatgtgc tcgtgtccc agacattcag 1380
aaccctgatc ctgcagtga ccagctgcgg gacagcaaga gcagcgcaa gagcgtgtgc 1440
ctgttcacg acttcgacag ccagaccaac gtgtccaga gcaaggacag cgacgtgtac 1500
atcaccgata agtgctgtgt ggacatgcgg agcatggact tcaagagcaa cagcgccgtg 1560
gcctggtcca acaagagcga ctcgcctgc gccaacgcct tcaacaacag cattatcccc 1620
gaggacacat tcttccaag ccccgagagc agctgcgacg tgaagctggt ggaaaagagc 1680
ttcgagacag acaccaacct gaacttcag aacctcagcg tgatcgctt ccgatactg 1740
ctgctgaagg tggcggctt caacctgctg atgacctgc ggctgtggtc cagctga 1797

```

<210> SEQ ID NO 75

<211> LENGTH: 1818

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR2-2 amino acid
(beta-P2A-alpha)

<400> SEQUENCE: 75

```

atgggaatta gactgtgtg cagagtggc ttctgttcc tggctgttg cctggtggac 60
gtgaaagtga ccagagcag cagatactg gtcaagagaa ccgcgagaa ggtgttctg 120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccgccctg 180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc 240
gagggctaca gcgtgtccag agagaagaaa gagcgggttca gcctgatact ggaaagcgcc 300
agcaccaacc agaccagcat gtacctgtgt gccagcggcg ccagacagat cccttacaca 360
tttggtccg gcaccagact gaccgtggtg gaagatctga acaaagtgtt ccctccagag 420

```

-continued

```

gtggccgtgt tcgagccttc tgaggccgag atcagccaca cacagaaagc cacactcgtg 480
tgccctggcca ccggtctttt tcccgatcac gtggaactgt cttggtgggt caacggcaaa 540
gaggtgcaca gcgcgctctg taccgatcct cagcctctga aagagcagcc cgctctgaac 600
gactccagat actgctgag cagcaggctg agagtgtccg ccacctctg gcagaacccc 660
agaaaccact tcagatgccca ggtgcagttc tacggcctga gcgagaacga tgagtggacc 720
caggatagag ccaagcctgt gacacagatc gtgtctgccg aagcctgggg cagagccgat 780
tgtggcttta cctcgtgtc ctatcagcag ggcgtgctgt ctgccaccat cctgtatgag 840
atcctgctgg gcaaagccac tctgtacgcc gtgctgggtg ctgccctggg gctgatggcc 900
atggtcaagc ggaaggattt tggtctgtgc gccaccaact ttagcctgct gaaacaggct 960
ggcgacgtgg aagagaaccc cggacctatg aagtcctga gagtgtgct ggtcctctg 1020
tggtgcagc tgtcttgggt ctggtcccag cagaaagagg tggaacagaa cagcggcct 1080
ctgtctgttc ctgaaggcgc tatcgctct ctgaattgca cctacagcga cagaggcagc 1140
cagagcttct tctggtatcg gcagtacagc ggcaagagcc ccgagctgat catgttcac 1200
tacagcaacg gcgacaaaga ggacggccgg ttacagccc agctgaacaa ggccagccag 1260
tacgtgtcac tgctgatcag agacagccag cctagcgaca gcgccaccta tctgtgtgcc 1320
gtgatcacc tcgaacaaca cgcggcaac atgtgacct ttggcggcgg aacacggctg 1380
atggttaagc cccacattca gaacccgat cctgcagtgt accagctgcg ggacagcaag 1440
agcagcgaca agagcgtgtg cctgttcacc gacttcgaca gccagaccaa cgtgtcccag 1500
agcaaggaca gcgacgtgta catcacgat aagtgcgtgc tggacatgcg gagcatggac 1560
ttcaagagca acagcgccgt ggctgtgtcc aacaagagcg acttcgcctg cgccaacgcc 1620
ttcaacaaca gcattatccc cgaggacaca ttcttcccaa gcccgcgag cagctgcgac 1680
gtgaagctgg tggaaaagag cttcgagaca gacaccaacc tgaacttcca gaacctcagc 1740
gtgatcggct tccggtacct gctgtgaag gtggccggct tcaacctgct gatgacctg 1800
cggctgtggt ccagctga 1818

```

<210> SEQ ID NO 76

<211> LENGTH: 1429

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence TCR3 amino acid
(beta-P2A-alpha)

<400> SEQUENCE: 76

```

atgggaatta gactgctgtg cagagtggcc ttctgcttcc tggtgttgg cctggtggac 60
gtgaaagtga cccagagcag cagatacctg gtcaagagaa ccgcgagaa ggtgttctg 120
gaatgcgtgc aggacatgga ccacgagaat atgttctggt acagacagga ccccggcctg 180
ggcctgagac tgatctactt cagctacgac gtgaagatga aggaaaaggg cgacatcccc 240
gagggctaca gcgtgtccag agagaagaaa gagcggttca gcctgatcct ggaaagcgcc 300
agcaccaacc agaccagcat gtacctgtgt gccagctctc tgtctgctgc cggcggaggc 360
tatggctaca ccttcaatta cggctacaca ttccgcagcg gcaccagact gacctgggtg 420
gaagatctga acaaagtgtt cctccagag gtggcctgt tcgagccttc tgaggccgag 480
atcagccaca cacagaaagc cacactcgtg tgccctggcca ccggtctttt tcccgatcac 540
gtggaactgt cttggtgggt caacggcaaa gaggtgcaca gcgcgctctg taccgatcct 600

```


-continued

cagcctctga aagagcagcc cgctctgaac gactccagat actgcctgag cagcaggctg	660
agagtgtccg ccaccttctg gcagaacccc agaaaccact tcagatgcca ggtgcagttc	720
tacggcctga gcgagaacga tgagtggacc caggatagag ccaagcctgt gacacagatc	780
gtgtctgccg aagcctgggg cagagccgat tgtggcttta cctccgtgtc ctatcagcag	840
ggcgtgctgt ctgccaccat cctgtatgag atcctgctgg gcaaagccac tctgtacgcc	900
gtgtcgtgtg ctgccctggt gctgatggcc atggccaagc ggaaggattt tggcagcggc	960
gccaccaact ttgacctgct gaaacaggct ggcgacgtgg aagagaaccc cggacctatg	1020
aagtccctga gagtgtgct ggtcctctg ttgctgcagc tgtcttgggt ctggtcccag	1080
cagaagagg tggaacagaa cagcggccct ctgtccgttc ctgaaggcgc cattgctagc	1140
ctgaattgca cctacagcga ccggggcagc cagagcttct tctggtatag gcagtacagc	1200
ggcaagagcc ccgagctgat catgttcac tacagcaacg gcgacaaaga ggacggcccg	1260
tttacagccc agctgaacaa ggccagccag tacgtgtcac tgctgatcag agacagccag	1320
cctagcgaca gcgccacct tctgtgtgcc gtgacaagca atgccggcaa gtccacattt	1380
ggcgacggca caacctgac cgtgaagccc aacattcaga accccgac	1429

<210> SEQ ID NO 77

<211> LENGTH: 8298

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence pRRLSIN-P2A-aWPRE-PST1

<400> SEQUENCE: 77

tcgaggggaat gaaagacccc acctgtaggt ttggcaagct agcttaagta acgccatttt	60
gcaaggcatg gaaaatacat aactgagaat agagaagttc agatcaagggt taggaacaga	120
gagacagcag aatatgggcc aaacaggata tctgtggtaa gcagttcctg ccccggtca	180
gggccaagaa cagatggtcc ccagatgcgg tcccgcctc agcagtttct agagaacct	240
cagatgttcc cagggtgccc caaggacctg aaaatgaccc tgtgccttat ttgaactaac	300
caatcagttc gcttctcgct tctgttcgcy cgcttctgct ccccgagctc aataaaagag	360
cccacaaccc ctcaactcggc gcgccggcca ccatgagcaa ccaggtgctg tgctgcgtgg	420
tgctgtgttt cctggggccc aacaccgtgg acggcggcat caccagagc cccaagtacc	480
tgttccggaa agagggccag aacgtcacc tgagctgcga gcagaacctg aaccacgacg	540
ccatgtactg gtacagacag gaccccgac agggcctgcg gctgatctac tacagccaga	600
tcgtgaacga cttccagaag ggagatatcg ccgagggcta cagcgtgtcc agagagaaga	660
aagagtcctt ccactgacc gtgaccagcg ccagaagaa cccacccgc ttctacctgt	720
gcgccagctc tcctggcgcc ctgtacgagc agtacttcgg ccctggcacc cggtgacag	780
tgaccgagga cctgaagaac gtgttcccc cagaggtggc cgtgttcgag cctagcgagg	840
ccgagatcag ccacaccag aaagccaccc tcgtgtgcct ggccaccggc ttttaccgcg	900
accacgtgga actgtcttg tggttcaacg gcaaagaggt gcacagcggc gtctgcaccg	960
acccccagcc cctgaaagag cagcccggcc tgaacgacag ccggtactgt ctgagcagca	1020
gactgagagt gtccgccacc ttctggcaga acccccgaa ccaactcaga tgccaggtgc	1080
agttctacgg cctgagcgag aacgacgagt ggacccagga ccgggccaag cccgtgaccc	1140
agatcgtgtc tgctgaggcc tggggcagag ccgattgcgg cttcaccagc gagagctacc	1200

-continued

agcagggcgt gctgagcgcc accatcctgt acgagatcct gctgggcaag gccaccctgt	1260
acgccgtgct ggtgtccgcc ctggtgctga tggccatggt caagcggaag gacagccggg	1320
gcggttccgg agccacgaac ttctctctgt taaagcaagc aggagacgtg gaagaaaacc	1380
ccggtcccat gaccagcatc cgggcccgtgt tcctcttctt gtggtgcag ctggacctcg	1440
tcaacggcga gaacgtggaa cagcacccca gcaccctgag cgtgcaggaa ggcgacagcg	1500
ccgtcatcaa gtgcacctac agcgactccg ccagcaacta cttccccctg tacaagcagg	1560
aactgggcaa gcgccccag ctgatcatcg acatccggtc caacgtgggc gagaagaagg	1620
accagcggat cgccgtgacc ctgaacaaga ccgccaagca cttcagcctg cacatcaccc	1680
agacacagcc cgaggactcc gccgtgtact tctgtgccgc caccgaggac taccagctga	1740
tctggggagc cggcaccaag ctgatcatta agcccgacat ccagaacccc gaccctgcag	1800
tgtaccagct cggggacagc aagagcagcg acaagagcgt gtgcctgttc accgacttcg	1860
acagccagac caacgtgtcc cagagcaagg acagcgacgt gtacatcacc gataagtgcg	1920
tgctggacat cgggagcatg gacttcaaga gcaacagcgc cgtggcctgg tccaacaaga	1980
gcgacttcgc ctgcgccaac gccttcaaca acagcattat ccccgaggac acattcttcc	2040
caagccccga gagcagctgc gacgtgaagc tgggtgaaaa gagcttcgag acagacacca	2100
acctgaactt ccagaacctc agcgtgatcg gcttccggat cctgctgctg aaggtggccg	2160
gcttcaacct gctgatgacc ctgcggctgt ggtccagctg agtcgacaat caacctctgg	2220
attacaaaat ttgtgaaaga ttgactggta ttcttaacta tgttgcctct ttacgctat	2280
gtggatacgc tgccttaatg cctttgtatc atgctattgc ttcccgatg gctttcattt	2340
tctctcctt gtataaatcc tgggtgctgt ctctttatga ggagtgttg cccgttgtca	2400
ggcaacgtgg cgtggtgtgc actgtgttg ctgacgcaac ccccaactgg tggggcattg	2460
ccaccacctg tcagctcctt tccgggactt tcgctttccc cctccctatt gccacggcgg	2520
aactcatcgc cgcctgcctt gcccgctgct ggacaggggc tcggctgttg ggcactgaca	2580
attccgtggg gttgtcgggg aaatcatcgt cctttccttg gctgctcgcc tgtgttgcca	2640
cctggattct gcgcgggacg tccttctgct acgtcccttc ggccctcaat ccagcggacc	2700
ttccttcccg cggcctgctg ccggctctgc ggccctcttc gcgtcttcgc cttcgccctc	2760
agacgagtcg gatctccctt tgggcgcct ccccgccctg aattcgagct cggtaacctt	2820
aagaccaatg acttacaagg cagctgtaga tcttagccac tttttaaag aaaagggggg	2880
actggaaggg ctaattcact cccaacgaag acaagatctg ctttttgctt gtactgggtc	2940
tctctgggta gaccagatct gagcctggga gctctctggc taactaggga acccaactgct	3000
taagcctcaa taaagcttgc cttgagtgct tcaagtagtg tgtgcccgtc tgttgtgtga	3060
ctctggtaac tagagatccc tcagacccct ttagtcagtg tggaaaatct ctgacgtag	3120
tagttcatgt catcttatta ttcagtattt ataacttgca aagaaatgaa tatcagagag	3180
tgagaggaa cttgtttattg cagcttataa tgggtacaaa taaagcaata gcatcacaaa	3240
tttcacaaat aaagcatttt ttcaactgca ttctagtgtg ggtttgtcca aactcatcaa	3300
tgtatcttat catgtctggc tctagctatc ccgcccctaa ctccgccag ttccgcccat	3360
tctccgcccc atggctgact aatttttttt atttatgcag aggcggaggg cgcctcggcc	3420
tctgagctat tccagaagta gtgaggaggc ttttttgag gcctaggctt ttgcgtcgag	3480
acgtacccaa ttccgcttat agtgagtcgt attacggcgc ctcactggcc gtcgttttac	3540
aacgtcgtga ctgggaaaac cctggcggtta cccaacttaa tcgccttgca gcacatcccc	3600

-continued

ctttcgccag	ctggcgtaat	agcgaagagg	ccgcacccga	tcgcccctcc	caacagttgc	3660
gcagcctgaa	tggcgaatgg	cgcgacgcgc	cctgtagcgg	cgcattaagc	gcggcgggtg	3720
tggtgggttac	gcgcagcgtg	accgctacac	ttgccagcgc	cctagcgcgc	gctcccttcg	3780
ctttcttccc	ttcctttctc	gccacgttcg	ecggctttcc	ccgtcaagct	ctaaatcggg	3840
ggctcccttt	agggttccga	tttagtgctt	tacggcacct	cgaccccaaa	aaacttgatt	3900
agggtgatgg	ttcacgtagt	gggccatcgc	cctgatagac	ggtttttcgc	cctttgacgt	3960
tggagtccac	gttctttaat	agtggactct	tggtccaaac	tggaacaaca	ctcaacccta	4020
tctcgggtcta	ttcttttgat	ttataaggga	ttttgccgat	ttcggcctat	tggttaaaaa	4080
atgagctgat	ttaacaaaaa	tttaacgcga	attttaacaa	aatattaacg	tttacaattt	4140
cccagggtggc	acttttcggg	gaaatgtgcg	cggaaacct	atttgtttat	ttttctaaat	4200
acattcaaat	atgtatccgc	tcatgagaca	ataacctga	taaatgcttc	aataatattg	4260
aaaaaggaag	agtatgagta	ttcaacattt	ccgtgtcgc	cttattccct	tttttcgggc	4320
attttgccct	cctgtttttg	ctcaccacga	aacgctgggtg	aaagtaaaag	atgctgaaga	4380
tcagttgggt	gcacgagtgg	gttacatcga	actggatctc	aacagcggta	agatccttga	4440
gagttttcgc	cccgaagaac	gttttccaat	gatgagcact	tttaaagtcc	tgctatgtgg	4500
cgcggtatta	tcccgatttg	acgccgggca	agagcaactc	ggtcgcgcga	tacactattc	4560
tcagaatgac	ttggttgagt	actcaccagt	cacagaaaag	catcttacgg	atggcatgac	4620
agtaagagaa	ttatgcagtg	ctgccataac	catgagtgat	aacactgcgg	ccaacttact	4680
tctgacaaag	atcggaggac	cgaaggagct	aaccgctttt	ttgcacaaca	tggggggatca	4740
tgtaactcgc	cttgatcggt	gggaaccgga	gctgaatgaa	gccataccaa	acgacgagcg	4800
tgacaccacg	atgcctgtag	caatggcaac	aacgttcgc	aaactattaa	ctggcgaaact	4860
acttactcta	gcttccggcg	aacaattaat	agactggatg	gaggcggata	aagttgcagg	4920
accacttctg	cgtcgggcgc	ttcgggtcgg	ctggtttatt	gctgataaat	ctggagccgg	4980
tgagcgtggg	tctcggcgta	tcattgcagc	actggggcca	gatggtaagc	cctcccgat	5040
cgtagttatc	tacacgacgg	ggagtcaggc	aactatggat	gaacgaaata	gacagatcgc	5100
tgagataggt	gcctcactga	ttaagcattg	gtaactgtca	gaccaagttt	actcatatat	5160
acttttagatt	gatttaaaac	ttcattttta	atttaaaagg	atctaggtga	agatcccttt	5220
tgataatctc	atgacaaaaa	tcctttaacg	tgagttttcg	ttccactgag	cgtcagaccc	5280
cgtagaaaag	atcaaaggat	cttcttgaga	tccttttttt	ctgcgcgtaa	tctgctgctt	5340
gcaaacaaaa	aaaccacgcg	taccagcggg	ggtttgtttg	ccgatcaag	agctaccaac	5400
tcttttttcg	aaggtaactg	gcttcagcag	agcgcagata	ccaaatactg	tccttctagt	5460
gtagccgtag	ttaggccacc	acttcaagaa	ctctgtagca	ccgcctacat	acctcgetct	5520
gctaatacctg	ttaccagtgg	ctgctgccag	tggcgataag	tcgtgtctta	ccgggttgga	5580
ctcaagacga	tagttaccgg	ataaggcgca	gcggtcgggc	tgaacggggg	gttcgtgcac	5640
acagcccagc	ttggagcgaa	cgaacctacac	cgaactgaga	tacctacagc	gtgagctatg	5700
agaaagcgcc	acgcttcccg	aaggggagaaa	ggcggacagg	tatccggtaa	gcggcagggg	5760
cggaacagga	gagcgcacga	gggagcttcc	agggggaaac	gcctgggtatc	tttatagtcc	5820
tgtcgggttt	cgccacctct	gacttgagcg	tcgatttttg	tgatgctcgt	caggggggcg	5880
gagcctatgg	aaaaacgcca	gcaacgcggc	ctttttacgg	ttcctggcct	tttgcgggcc	5940

-continued

ttttgctcac atgtttcttc ctgcgttate ccctgattct gtggataacc gtattaccgc	6000
ctttgagtga gctgataccg ctgcgcgcag ccgaacgacc gagcgcagcg agtcagtgag	6060
cgaggaaagc gaagagcgcc caatacgcaa accgcctctc cccgcgcgtt ggccgattca	6120
ttaatgcagc tggcacgaca ggtttcccgga ctggaaagcg ggcagtgagc gcaacgcaat	6180
taatgtgagt tagctcactc attaggcacc ccaggcttta cactttatgc ttccggctcg	6240
tatgttgtgt ggaattgtga gcggataaca atttcacaca ggaaacagct atgacctga	6300
ttacgccaaag cgcgcaatta accctcacta aagggaacaa aagctggagc tgcaagctta	6360
atgtagtctt atgcaatact ctgtagtct tgcaacatgg taacgatgag ttagcaacat	6420
gccttacaag gagagaaaaa gcaccgtgca tgccgattgg tggaagtaag gtggtacgat	6480
cgtgccttat taggaaggca acagacgggt ctgacatgga ttggacgaac cactgaattg	6540
ccgcattgca gagatatgtt atttaagtgc ctagctcgat acaataaacg ggtctctctg	6600
gttagaccag atctgagcct gggagctctc tggctaacta gggaacccac tgcttaagcc	6660
tcaataaagc ttgccttgag tgcttcaagt agtgtgtgcc cgtctgttgt gtgactctgg	6720
taactagaga tccctcagac ccttttagtc agtgtggaaa atctctagca gtggcgcccc	6780
aaacgggacc tgaagcgcaa agggaaacca gagctctctc gacgcaggac tcggcttgct	6840
gaagcgcgca cggcaagagg cgaggggagg cgactggtga gtacgcaaa aattttgact	6900
agcggaggct agaaggagag agatgggtgc gagagcgta gtattaaagc ggggagaatt	6960
agatcgcgat gggaaaaaat tcgggttaag ccagggggaa agaaaaata taaattaaaa	7020
catatagtat gggcaagcag ggagctagaa cgattcgag ttaatcctgg cctgttagaa	7080
acatcagaag gctgtagaca aatactggga cagctacaac catcccttca gacaggatca	7140
gaagaactta gatcattata taatacagta gcaacctct atttgtgtga tcaaaggata	7200
gagataaaag acaccaagga agcttttagac aagatagagg aagagcaaaa caaaagtaag	7260
accaccgcac agcaagcggc cgctgatctt cagacctgga ggaggagata tgagggacaa	7320
ttggagaagt gaattatata aatataaagt agtaaaaatt gaaccattag gagtagcacc	7380
caccaaggca aagagaagag tggtcgagag agaaaaaga gcagtgggaa taggagcttt	7440
gttccttggg ttcttgggag cagcaggaag cactatgggc gcagcctcaa tgacgctgac	7500
ggtacaggcc agacaattat tgtctggtat agtgcagcag cagaacaatt tgcaggggc	7560
tattgagcgc caacagcatc tgttgcaact cacagtctgg ggcacaaagc agctccaggc	7620
aagaatcctg gctgtggaaa gatacctaaa ggatcaacag ctctgggga tttggggttg	7680
ctctggaaaa ctcatthtga ccactgctgt gccttggaat gctagttgga gtaataaatc	7740
tctggaacag atttggaatc acacgacctg gatggagtgg gacagagaaa ttaacaatta	7800
cacaagctta atacactcct taattgaaga atcgcaaac cagcaagaaa agaataaaca	7860
agaattattg gaattagata aatgggcaag tttgtggaat tggtttaaca taacaaattg	7920
gctgtggtat ataaaattat tcataatgat agtaggaggc ttggtagggt taagaatagt	7980
ttttgctgta ctttctatag tgaatagagt taggcaggga tattcccat tatcgtttca	8040
gaccacacct ccaacccga ggggacccga caggcccgaa ggaatagaag aagaaggtag	8100
agagagagac agagacagat ccattcgatt agtgaacgga tctcgacggt atcggttaac	8160
ttttaaaaga aaagggggga ttggggggta cagtgcaggg gaaagaatag tagacataat	8220
agcaacagac atacaaacta aagaattaca aaaacaaatt acaaaaattc aaaattttat	8280
cgatcacgag actagccg	8298

-continued

```

<210> SEQ ID NO 78
<211> LENGTH: 2949
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence Human BRAF

<400> SEQUENCE: 78
cgctccctt cccctcccc gcccgacagc ggccgctcgg gcccggctc tcggttataa    60
gatggcggcg ctgagcgggtg gcggtgggtg gcgcgaggag cggggccagg ctctgttcaa    120
cggggacatg gagcccgagg ccggcgccgg cgccggcgcc cgggcctctt cggtgcgga    180
ccctgccatt ccggaggagg tgtggaatat caaacaaatg attaagtga cacaggaaca    240
tatagaggcc ctattggaca aatttggtgg ggagcataat ccaccatcaa tatatctgga    300
ggcctatgaa gaatacacca gcaagctaga tgcactcaa caaagagaac aacagttatt    360
ggaatctctg gggaacggaa ctgatttttc tgtttctagc tctgcatcaa tggataccgt    420
tacatcttct tcctcttcta gcttttcagt gctaccttca tctctttcag tttttcaaaa    480
tcccacagat gtggcacgga gcaaccccaa gtcaccacaa aaacctatcg ttagagtctt    540
cctgccaac aaacagagga cagtgtgacc tgcaagggtg ggagttacag tccgagacag    600
tctaagaaa gcaactgatg tgagagggtc aatcccagag tgctgtgctg tttacagaat    660
tcaggatgga gagaagaac caattggttg ggacactgat atttctggc ttactggaga    720
agaattgcat gtggaagtgt tggagaatgt tccacttaca acacacaact ttgtacgaaa    780
aacgttttct acctagcat tttgtgactt ttgtcgaaag ctgcttttcc aggggttccg    840
ctgtcaaaaca tgtggttata aatttcacca gcgttgtagt acagaagttc cactgatgtg    900
tgttaattat gaccaacttg atttgctgtt tgtctccaag ttctttgaac accaccaat    960
accacaggaa gaggggtcct tagcagagac tgccctaaca tctggatcat ccccttcgc    1020
acccgcctcg gactotattg ggcccaaat tctcaccagt ccgtctcctt caaatccat    1080
tccaattcca cagcccttcc gaccagcaga tgaagatcat cgaaatcaat ttgggcaacg    1140
agaccgatcc tcacagctc ccaatgtgca tataaacaca atagaacctg tcaatattga    1200
tgacttgatt agagaccaag gatttcgtgg tgatggagga tcaaccacag gtttgtctgc    1260
tccccccct gcctcattac ctgggtcact aactaacgtg aaagccttac agaaatctcc    1320
aggacctcag cgagaaaagg agtcatcttc atcctcagaa gacaggaatc gaatgaaaac    1380
acttggtaga cgggactcga gtgatgattg ggagattcct gatgggcaga ttacagtggg    1440
acaaagaatt ggatctggat catttggaac agtctacaag ggaaagtggc atggtgatgt    1500
ggcagtgaat atgttgatg tgacagcacc tacacctcag cagttacaag ccttcaaaaa    1560
tgaagtagga gtactcagga aaacacgaca tgtgaatct ctactcttca tgggctattc    1620
cacaagcca caactggcta ttgttacca gtggtgtgag ggctccagct tgtatcacca    1680
tctccatctc attgagacca aatttgagat gatcaaatct atagatattg cagcagagac    1740
tgcacagggc atggattact tacacgcaa gtcaatctc cacagagacc tcaagagtaa    1800
taatataatt cttcatgaag acctcacagt aaaaataggt gattttggtc tagctacagt    1860
gaaatctcga tggagtgggt cccatcagtt tgaacagttg tctggatcca ttttgtggat    1920
ggcaccagaa gtcacagaa tgcaagataa aaatccatag agctttcagt cagatgtata    1980
tgcatttgga attgttctgt atgaattgat gactggacag ttaccttatt caaacatcaa    2040

```

-continued

```

caacagggac cagataatTT ttatggtggg acgaggatac ctgtctccag atctcagtaa 2100
ggtacggagt aactgtccaa aagccatgaa gagattaatg gcagagtgcc tcaaaaagaa 2160
aagagatgag agaccactct tccccaaat tctcgctctc attgagctgc tggcccgctc 2220
attgccaaaa attcaccgca gtgcatacaga accctccttg aatcgggctg gtttccaaac 2280
agaggatTTT agtctatatg cttgtgcttc tccaaaaaca cccatccagg cagggggata 2340
tggtgcgTTT cctgtccact gaaacaaatg agtgagagag ttcaggagag tagcaacaaa 2400
agggaaaata atgaacatat gtttgcttat atgttaaatt gaataaaata ctctctTTTT 2460
ttttaaggTg aaccaaagaa cacttgTgtg gttaaagact agatataatt tttcccaaaa 2520
ctaaaattta tacttaacat tggattTTta acatccaagg gttaaaatac atagacattg 2580
ctaaaattg gcagagcctc ttctagaggc ttactttctc gttccgggtt tgtatcattc 2640
acttggttat tttagtagt aaacttcagt ttctcatgca acttttgTtg ccagctatca 2700
catgtccact agggactcca gaagaagacc ctacctatgc ctgtgTttgc aggtgagaag 2760
ttggcagtcg gttagcctgg gttagataag gcaaactgaa cagatctaatt ttaggaagtc 2820
agtagaattt aataattcta ttattattct taataatttt tctataacta tttctTTtta 2880
taacaatttg gaaaatgtgg atgtctTTta tttccttgaa gcaataaaact aagtttcttt 2940
ttataaaaaa

```

```

<210> SEQ ID NO 79
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence AP1M1

```

```

<400> SEQUENCE: 79

```

```

Val Ile Glu Lys His Ser His Ser Arg Ile Glu Tyr Met Leu Lys Ala
1             5             10            15
Lys Ser Gln Phe Lys Arg Arg Ser Thr Ala Asn
                20            25

```

```

<210> SEQ ID NO 80
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence AP1M1

```

```

<400> SEQUENCE: 80

```

```

Val Ile Glu Lys His Ser His Ser Arg Ile Glu Tyr Met Leu Lys Ala
1             5             10            15
Lys Ser Gln Phe
                20

```

```

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence AP1M1

```

```

<400> SEQUENCE: 81

```

```

Ser Arg Ile Glu Tyr Met Leu Lys Ala Lys Ser Gln Phe Lys Arg Arg
1             5             10            15
Ser Thr Ala Asn
                20

```

-continued

<210> SEQ ID NO 82
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence DCAF6

<400> SEQUENCE: 82

Glu Gln Phe Leu Gln Pro Ser Thr Ser Ser Thr Met Ser Thr Gln Ala
1 5 10 15

His Ser Thr Ser Ser Pro Thr Glu Ser Pro His
20 25

<210> SEQ ID NO 83
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence DCAF6

<400> SEQUENCE: 83

Glu Gln Phe Leu Gln Pro Ser Thr Ser Ser Thr Met Ser Thr Gln Ala
1 5 10 15

His Ser Thr Ser
20

<210> SEQ ID NO 84
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence DCAF6

<400> SEQUENCE: 84

Thr Ser Ser Thr Met Ser Thr Gln Ala His Ser Thr Ser Ser Pro Thr
1 5 10 15

Glu Ser Pro His
20

<210> SEQ ID NO 85
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GTF2H4

<400> SEQUENCE: 85

Phe Ile Val Val Glu Thr Asn Tyr Arg Leu Tyr Ala Tyr Met Glu Ser
1 5 10 15

Glu Leu Gln Ile Ala Leu Ile Ala Leu Phe Ser
20 25

<210> SEQ ID NO 86
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GTF2H4

<400> SEQUENCE: 86

Phe Ile Val Val Glu Thr Asn Tyr Arg Leu Tyr Ala Tyr Met Glu Ser
1 5 10 15

Glu Leu Gln Ile

-continued

20

<210> SEQ ID NO 87
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GTF2H4

<400> SEQUENCE: 87

Tyr Arg Leu Tyr Ala Tyr Met Glu Ser Glu Leu Gln Ile Ala Leu Ile
1 5 10 15

Ala Leu Phe Ser
20

<210> SEQ ID NO 88
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NBPF12

<400> SEQUENCE: 88

Asp Ser Cys Gln Pro Tyr Arg Ser Ser Phe Tyr Ala Leu Gly Glu Lys
1 5 10 15

His Val Gly Phe Ser Leu Asp Val Gly Glu Ile
20 25

<210> SEQ ID NO 89
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NBPF12

<400> SEQUENCE: 89

Asp Ser Cys Gln Pro Tyr Arg Ser Ser Phe Tyr Ala Leu Gly Glu Lys
1 5 10 15

His Val Gly Phe
20

<210> SEQ ID NO 90
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NBPF12

<400> SEQUENCE: 90

Ser Ser Phe Tyr Ala Leu Gly Glu Lys His Val Gly Phe Ser Leu Asp
1 5 10 15

Val Gly Glu Ile
20

<210> SEQ ID NO 91
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ORC3

<400> SEQUENCE: 91

Glu Ser Phe Ala Thr Lys Val Leu Gln Asp Phe Ile Ile Leu Ser Ser
1 5 10 15

-continued

Gln His Leu His Glu Phe Pro Leu Ile Leu Ile
20 25

<210> SEQ ID NO 92
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ORC3

<400> SEQUENCE: 92

Glu Ser Phe Ala Thr Lys Val Leu Gln Asp Phe Ile Ile Leu Ser Ser
1 5 10 15

Gln His Leu His
20

<210> SEQ ID NO 93
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ORC3

<400> SEQUENCE: 93

Leu Gln Asp Phe Ile Ile Leu Ser Ser Gln His Leu His Glu Phe Pro
1 5 10 15

Leu Ile Leu Ile
20

<210> SEQ ID NO 94
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ROR1

<400> SEQUENCE: 94

Leu Val Pro Thr Ser Ser Trp Asn Ile Ser Ser Glu Leu Ser Lys Asp
1 5 10 15

Ser Tyr Leu Thr Leu Asp Glu Pro Met Asn Asn
20 25

<210> SEQ ID NO 95
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ROR1

<400> SEQUENCE: 95

Leu Val Pro Thr Ser Ser Trp Asn Ile Ser Ser Glu Leu Ser Lys Asp
1 5 10 15

Ser Tyr Leu Thr
20

<210> SEQ ID NO 96
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ROR1

<400> SEQUENCE: 96

Asn Ile Ser Ser Glu Leu Ser Lys Asp Ser Tyr Leu Thr Leu Asp Glu
1 5 10 15

-continued

Pro Met Asn Asn
20

<210> SEQ ID NO 97
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence SF3B1

<400> SEQUENCE: 97

Gln Met Gly Gly Ser Thr Pro Val Leu Thr Pro Gly Lys Ala Pro Ile
1 5 10 15
Gly Thr Pro Ala Met Asn Met Ala Thr Pro Thr
20 25

<210> SEQ ID NO 98
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence SF3B1

<400> SEQUENCE: 98

Gln Met Gly Gly Ser Thr Pro Val Leu Thr Pro Gly Lys Ala Pro Ile
1 5 10 15
Gly Thr Pro Ala
20

<210> SEQ ID NO 99
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence SF3B1

<400> SEQUENCE: 99

Val Leu Thr Pro Gly Lys Ala Pro Ile Gly Thr Pro Ala Met Asn Met
1 5 10 15
Ala Thr Pro Thr
20

<210> SEQ ID NO 100
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence UNKL

<400> SEQUENCE: 100

Ala His Gly Pro Leu Asp Leu Arg Pro Pro Val Cys Asp Ile Arg Glu
1 5 10 15
Leu Gln Ala Gln Glu Ala Leu Gln Asn Gly Gln
20 25

<210> SEQ ID NO 101
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence UNKL

<400> SEQUENCE: 101

Ala His Gly Pro Leu Asp Leu Arg Pro Pro Val Cys Asp Ile Arg Glu

-continued

1	5	10	15
---	---	----	----

Leu Gln Ala Gln
20

<210> SEQ ID NO 102
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence UNKL

<400> SEQUENCE: 102

Arg Pro Pro Val Cys Asp Ile Arg Glu Leu Gln Ala Gln Glu Ala Leu
1 5 10 15

Gln Asn Gly Gln
20

<210> SEQ ID NO 103
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence ZNF700

<400> SEQUENCE: 103

Gly Glu Lys Pro Tyr Glu Cys Ser Lys Cys Asp Lys Ala Leu His Ser
1 5 10 15

Ser Ser Ser Tyr His Arg His Glu Arg Ser His
20 25

<210> SEQ ID NO 104
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence ZNF700

<400> SEQUENCE: 104

Gly Glu Lys Pro Tyr Glu Cys Ser Lys Cys Asp Lys Ala Leu His Ser
1 5 10 15

Ser Ser Ser Tyr
20

<210> SEQ ID NO 105
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence ZNF700

<400> SEQUENCE: 105

Ser Lys Cys Asp Lys Ala Leu His Ser Ser Ser Ser Tyr His Arg His
1 5 10 15

Glu Arg Ser His
20

<210> SEQ ID NO 106
 <211> LENGTH: 27
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence NVL

<400> SEQUENCE: 106

-continued

Ala Pro Cys Ile Ile Phe Ile Asp Glu Ile Asp Ala Ile Pro Pro Lys
1 5 10 15

Arg Glu Val Ala Ser Lys Asp Met Glu Arg Arg
20 25

<210> SEQ ID NO 107
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NVL

<400> SEQUENCE: 107

Ala Pro Cys Ile Ile Phe Ile Asp Glu Ile Asp Ala Ile Pro Pro Lys
1 5 10 15

Arg Glu Val Ala
20

<210> SEQ ID NO 108
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NVL

<400> SEQUENCE: 108

Asp Glu Ile Asp Ala Ile Pro Pro Lys Arg Glu Val Ala Ser Lys Asp
1 5 10 15

Met Glu Arg Arg
20

<210> SEQ ID NO 109
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MATN1

<400> SEQUENCE: 109

Ser Arg Ser Pro Asp Ile Ser Lys Val Val Ile Val Val Pro Asp Gly
1 5 10 15

Arg Pro Gln Asp Ser Val Gln Asp Val Ser Ala
20 25

<210> SEQ ID NO 110
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MATN1

<400> SEQUENCE: 110

Ser Arg Ser Pro Asp Ile Ser Lys Val Val Ile Val Val Pro Asp Gly
1 5 10 15

Arg Pro Gln Asp
20

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MATN1

<400> SEQUENCE: 111

-continued

Lys Val Val Ile Val Val Pro Asp Gly Arg Pro Gln Asp Ser Val Gln
1 5 10 15

Asp Val Ser Ala
20

<210> SEQ ID NO 112
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence CTNNA2

<400> SEQUENCE: 112

Val Arg Gln Ala Leu Gln Asp Leu Leu Ser Glu Tyr Met His Asn Thr
1 5 10 15

Gly Arg Lys Glu Lys Gly Asp Pro Leu Asn Ile
20 25

<210> SEQ ID NO 113
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence CTNNA2

<400> SEQUENCE: 113

Val Arg Gln Ala Leu Gln Asp Leu Leu Ser Glu Tyr Met His Asn Thr
1 5 10 15

Gly Arg Lys Glu
20

<210> SEQ ID NO 114
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence CTNNA2

<400> SEQUENCE: 114

Leu Leu Ser Glu Tyr Met His Asn Thr Gly Arg Lys Glu Lys Gly Asp
1 5 10 15

Pro Leu Asn Ile
20

<210> SEQ ID NO 115
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GET4

<400> SEQUENCE: 115

Arg Tyr Met Ser Gln Ser Lys His Thr Glu Ala Arg Glu Arg Met Tyr
1 5 10 15

Ser Gly Ala Leu Leu Phe Phe Ser His Gly Gln
20 25

<210> SEQ ID NO 116
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GET4

-continued

<400> SEQUENCE: 116

Arg Tyr Met Ser Gln Ser Lys His Thr Glu Ala Arg Glu Arg Met Tyr
1 5 10 15

Ser Gly Ala Leu
 20

<210> SEQ ID NO 117
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence GET4

<400> SEQUENCE: 117

His Thr Glu Ala Arg Glu Arg Met Tyr Ser Gly Ala Leu Leu Phe Phe
1 5 10 15

Ser His Gly Gln
 20

<210> SEQ ID NO 118
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NTNG1

<400> SEQUENCE: 118

Thr Val Thr Asp Leu Arg Ile Arg Leu Leu Arg Pro Ala Ile Gly Glu
1 5 10 15

Ile Phe Val Asp Glu Leu His Leu Ala Arg Tyr
 20 25

<210> SEQ ID NO 119
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NTNG1

<400> SEQUENCE: 119

Thr Val Thr Asp Leu Arg Ile Arg Leu Leu Arg Pro Ala Ile Gly Glu
1 5 10 15

Ile Phe Val Asp
 20

<210> SEQ ID NO 120
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence NTNG1

<400> SEQUENCE: 120

Arg Leu Leu Arg Pro Ala Ile Gly Glu Ile Phe Val Asp Glu Leu His
1 5 10 15

Leu Ala Arg Tyr
 20

<210> SEQ ID NO 121
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence SPTBN5

-continued

<400> SEQUENCE: 121

Thr Leu Leu Leu Asp Ala Trp Leu Thr Thr Lys Ala Ala Ile Ala Glu
1 5 10 15

Ser Gln Asp Tyr Gly Gln Asp Leu Glu Gly Val
20 25

<210> SEQ ID NO 122

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence SPTBN5

<400> SEQUENCE: 122

Thr Leu Leu Leu Asp Ala Trp Leu Thr Thr Lys Ala Ala Ile Ala Glu
1 5 10 15

Ser Gln Asp Tyr
20

<210> SEQ ID NO 123

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence SPTBN5

<400> SEQUENCE: 123

Leu Thr Thr Lys Ala Ala Ile Ala Glu Ser Gln Asp Tyr Gly Gln Asp
1 5 10 15

Leu Glu Gly Val
20

<210> SEQ ID NO 124

<211> LENGTH: 27

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence DPP6

<400> SEQUENCE: 124

Leu Leu Val Ile Leu Val Ile Cys Ser Leu Ile Val Thr Leu Val Ile
1 5 10 15

Leu Leu Thr Pro Ala Glu Asp Asn Ser Leu Ser
20 25

<210> SEQ ID NO 125

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence DPP6

<400> SEQUENCE: 125

Leu Leu Val Ile Leu Val Ile Cys Ser Leu Ile Val Thr Leu Val Ile
1 5 10 15

Leu Leu Thr Pro
20

<210> SEQ ID NO 126

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic sequence DPP6

<400> SEQUENCE: 126

Cys Ser Leu Ile Val Thr Leu Val Ile Leu Leu Thr Pro Ala Glu Asp
1 5 10 15

Asn Ser Leu Ser
20

<210> SEQ ID NO 127

<211> LENGTH: 27

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence HIAT1

<400> SEQUENCE: 127

Val Lys Gly Leu Leu Ser Phe Leu Ser Ala Pro Leu Ile Cys Ala Leu
1 5 10 15

Ser Asp Val Trp Gly Arg Lys Ser Phe Leu Leu
20 25

<210> SEQ ID NO 128

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence HIAT1

<400> SEQUENCE: 128

Val Lys Gly Leu Leu Ser Phe Leu Ser Ala Pro Leu Ile Cys Ala Leu
1 5 10 15

Ser Asp Val Trp
20

<210> SEQ ID NO 129

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence HIAT1

<400> SEQUENCE: 129

Leu Ser Ala Pro Leu Ile Cys Ala Leu Ser Asp Val Trp Gly Arg Lys
1 5 10 15

Ser Phe Leu Leu
20

<210> SEQ ID NO 130

<211> LENGTH: 27

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic sequence ITGA4

<400> SEQUENCE: 130

Gly Ser Gly Ala Val Met Asn Ala Met Glu Thr Asn Leu Phe Gly Ser
1 5 10 15

Asp Lys Tyr Ala Ala Arg Phe Gly Glu Ser Ile
20 25

<210> SEQ ID NO 131

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ITGA4

<400> SEQUENCE: 131

Gly Ser Gly Ala Val Met Asn Ala Met Glu Thr Asn Leu Phe Gly Ser
1 5 10 15
Asp Lys Tyr Ala
20

<210> SEQ ID NO 132
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence ITGA4

<400> SEQUENCE: 132

Ala Met Glu Thr Asn Leu Phe Gly Ser Asp Lys Tyr Ala Ala Arg Phe
1 5 10 15
Gly Glu Ser Ile
20

<210> SEQ ID NO 133
<211> LENGTH: 27
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MYO1A

<400> SEQUENCE: 133

Ser Val Arg Phe Lys Glu Asn Ser Val Ala Val Lys Val Ile Gln Gly
1 5 10 15
Pro Ala Gly Gly Asp Asn Ser Lys Leu Arg Tyr
20 25

<210> SEQ ID NO 134
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MYO1A

<400> SEQUENCE: 134

Ser Val Arg Phe Lys Glu Asn Ser Val Ala Val Lys Val Ile Gln Gly
1 5 10 15
Pro Ala Gly Gly
20

<210> SEQ ID NO 135
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence MYO1A

<400> SEQUENCE: 135

Ser Val Ala Val Lys Val Ile Gln Gly Pro Ala Gly Gly Asp Asn Ser
1 5 10 15
Lys Leu Arg Tyr
20

<210> SEQ ID NO 136
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TRAC-gRNA-1

<400> SEQUENCE: 136

agagtctctc agctggtaca 20

<210> SEQ ID NO 137
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TRAC-gRNA-2

<400> SEQUENCE: 137

tgtgctagac atgaggtcta 20

<210> SEQ ID NO 138
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TRBC-gRNA-1

<400> SEQUENCE: 138

gcagtatctg gagtcattga 20

<210> SEQ ID NO 139
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence TRBC-gRNA-2

<400> SEQUENCE: 139

ggagaatgac gagtggaccc 20

<210> SEQ ID NO 140
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence B2M-gRNA

<400> SEQUENCE: 140

cgcgagcaca gctaaggcca 20

<210> SEQ ID NO 141
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence PD-1-gRNA

<400> SEQUENCE: 141

ggccaggatg gttcttaggt 20

<210> SEQ ID NO 142
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic sequence sgRNA Forward Oligo:
TRAC_sgRNA_pLenti_F1

<400> SEQUENCE: 142

-continued

caccggagaa tcaaaatcgg tgaat	25
<210> SEQ ID NO 143 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence sgRNA Reverse Oligo: TRAC_sgRNA_pLenti_R1	
<400> SEQUENCE: 143	
aaacattcac cgattttgat tctcc	25
<210> SEQ ID NO 144 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence PD1_sgRNA_F1	
<400> SEQUENCE: 144	
caccgcagtt gtgtgacacg gaag	24
<210> SEQ ID NO 145 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence PD1_sgRNA_R1	
<400> SEQUENCE: 145	
aaaccttcg tgtcacacaa ctgc	24
<210> SEQ ID NO 146 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence CTLA4_sgRNA_F1	
<400> SEQUENCE: 146	
caccggcaaa ggtgagtgag acttt	25
<210> SEQ ID NO 147 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence CTLA4_sgRNA_R1	
<400> SEQUENCE: 147	
aaacaaagtc tcaactcacct ttgcc	25
<210> SEQ ID NO 148 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic sequence LAG3_sgRNA_F1	
<400> SEQUENCE: 148	
caccggtttc tgcagccgct ttggg	25
<210> SEQ ID NO 149 <211> LENGTH: 25 <212> TYPE: DNA	

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic sequence LAG3_sgRNA_R1

<400> SEQUENCE: 149

aaaccccaaa gcggtgcag aaacc

25

What is claimed is:

1. An isolated polynucleotide encoding a binding protein, wherein the binding protein comprises:

- (a) a T cell receptor (TCR) α chain variable ($V\alpha$) domain comprising the $V\alpha$ domain amino acid sequence set forth in SEQ ID NO.:1; and
- (b) a TCR β chain variable ($V\beta$) domain comprising the $V\beta$ domain amino acid sequence set forth in SEQ ID NO.:5,

wherein the binding protein is capable of specifically binding to a HLA complex on a cell surface comprising a BRAF peptide containing a BRAF^{V600E} mutation and does not bind a HLA complex on a cell surface comprising a BRAF peptide not containing the BRAF^{V600E} mutation,

wherein the BRAF^{V600E} peptide consists of the amino acid sequence set forth in SEQ ID NO: 38 or 39,

wherein the HLA complex comprises (1) HLA-DQA1*03 and (2) HLA-DQB1*0301, HLA-DQB1*0302, or HLA-DQB1*0303, wherein:

- (i) the binding protein comprises a TCR, wherein the $V\alpha$ domain is comprised in a TCR α chain that further comprises a constant ($C\alpha$) domain comprising an introduced cysteine mutation, and the $V\beta$ domain is comprised in a TCR β chain that further comprises a constant ($C\beta$) domain comprising an introduced cysteine mutation, wherein the introduced cysteine mutations promote pairing of the TCR α chain with the TCR β chain when the binding protein is expressed in a T cell;
- (ii) the polynucleotide comprises a sequence encoding a self-cleaving peptide disposed between a sequence encoding the $V\beta$ domain and a sequence encoding the $V\alpha$ domain;
- (iii) the polynucleotide is codon optimized for expression in a human T cell; or
- (iv) the binding protein comprises a single chain TCR (scTCR) or a chimeric antigen receptor (CAR).

2. The polynucleotide according to claim 1, wherein the $V\alpha$ domain is comprised in a TCR α chain that further comprises a constant ($C\alpha$) domain comprising an introduced cysteine mutation, and the $V\beta$ domain is comprised in a TCR β chain that further comprises a constant ($C\beta$) domain comprising an introduced cysteine mutation, wherein the introduced cysteine mutations promote pairing of the TCR α chain with the TCR β chain when the binding protein is expressed in a T cell.

3. The polynucleotide according to claim 1, wherein the polynucleotide is codon optimized for expression in a human T cell.

4. The polynucleotide according to claim 1, wherein the binding protein is a TCR, an antigen-binding fragment of a TCR, or a chimeric antigen receptor.

5. The polynucleotide according to claim 4, wherein the binding protein is a TCR and comprises (i) an α chain constant ($C\alpha$) domain comprising the amino acid sequence

of SEQ ID NO:25 and (ii) a β chain ($C\beta$) constant domain comprising the amino acid sequence of SEQ ID NO: 26.

6. An isolated polynucleotide encoding a TCR, wherein the TCR comprises (i) a TCR α chain that comprises the amino acid sequence set forth in SEQ ID NO: 55 and (ii) a TCR β chain that comprises the amino acid sequence set forth in SEQ ID NO: 59.

7. An expression vector, comprising a polynucleotide encoding a binding protein, wherein the binding protein comprises:

- (a) a T cell receptor (TCR) α chain variable ($V\alpha$) domain comprising the $V\alpha$ domain amino acid sequence set forth in SEQ ID NO.:1; and
- (b) a TCR β chain variable ($V\beta$) domain comprising the $V\beta$ domain amino acid sequence set forth in SEQ ID NO.:5,

wherein the binding protein is capable of specifically binding to a HLA complex on a cell surface comprising a BRAF peptide containing a BRAF^{V600E} mutation and does not bind a HLA complex on a cell surface comprising a BRAF peptide not containing the BRAF^{V600E} mutation,

wherein the BRAF^{V600E} peptide consists of the amino acid sequence set forth in SEQ ID NO: 38 or 39,

wherein the HLA complex comprises (1) HLA-DQA1*03 and (2) HLA-DQB1*0301, HLA-DQB1*0302, or HLA-DQB1*0303,

and wherein:

- (i) the expression vector (1) comprises an expression control element that is operably coupled to the polynucleotide and is heterologous to the polynucleotide, and/or (2) comprises a viral vector, a bacterial vector, a plasmid, or a cosmid;
- (ii) the binding protein comprises a TCR, wherein the $V\alpha$ domain is comprised in a TCR α chain that further comprises a constant ($C\alpha$) domain comprising an introduced cysteine mutation, and the $V\beta$ domain is comprised in a TCR β chain that further comprises a constant ($C\beta$) domain comprising an introduced cysteine mutation, wherein the introduced cysteine mutations promote pairing of the TCR α chain with the TCR β chain when the binding protein is expressed in a T cell;
- iii) the polynucleotide comprises a sequence encoding a self-cleaving peptide disposed between a sequence encoding the $V\beta$ domain and a sequence encoding the $V\alpha$ domain;
- (iv) the polynucleotide is codon optimized for expression in a human T cell; and/or
- (v) the binding protein comprises a single chain TCR (scTCR) or a chimeric antigen receptor (CAR).

8. A host cell, comprising a polynucleotide according to claim 1, wherein the host cell expresses on its cell surface the encoded binding protein.

9. A host cell comprising a heterologous polynucleotide encoding a binding protein, wherein the encoded binding protein comprises:

169

(a) a T cell receptor (TCR) α chain variable ($V\alpha$) domain comprising the $V\alpha$ domain amino acid sequence set forth in SEQ ID NO:1, and

(b) a TCR β chain variable ($V\beta$) domain comprising the $V\beta$ domain amino acid sequence set forth in SEQ ID NO:5,

wherein the binding protein is capable of specifically binding to a HLA complex on a cell surface comprising a BRAF peptide containing a BRAF^{V600E} mutation and does not bind a HLA complex on a cell surface comprising a BRAF peptide not containing the BRAF^{V600E} mutation,

wherein the BRAF^{V600E} peptide consists of the amino acid sequence set forth in SEQ ID NO: 38 or 39,

wherein the HLA complex comprises (1) HLA-DQA1*03 and (2) HLA-DQB1*0301, HLA-DQB1*0302, or HLA-DQB1*0303.

10. The host cell according to claim 9, wherein the host cell comprises a CD4+ T cell, a CD8+ T cell, a CD4- CD8- double negative T cell, a $\gamma\delta$ T cell, a natural killer cell, a dendritic cell, or any combination thereof.

170

11. The host cell according to claim 9, wherein the host cell is a hematopoietic progenitor cell or an immune system cell.

12. The host cell according to claim 9, wherein the encoded binding protein comprises: a TCR α chain that comprises or consists of the amino acid sequence set forth in SEQ ID NO.:55; and a TCR β chain that comprises or consists of the amino acid sequence set forth in SEQ ID NO.:59.

13. A CD4+ or CD8+ T cell encoding the amino acid sequence of SEQ ID NO.:69.

14. The host cell according to claim 8, wherein the host cell comprises a T cell.

15. A vector comprising the polynucleotide of claim 1, wherein the vector comprises a lentiviral vector or a retroviral vector.

16. The host cell according to claim 9, wherein the binding protein is a TCR and the host cell comprises a CD4+ T cell.

* * * * *