



US 20250276070A1

(19) **United States**
(12) **Patent Application Publication** (10) **Pub. No.: US 2025/0276070 A1**
Julien et al. (43) **Pub. Date: Sep. 4, 2025**

(54) **OPTIMIZED MULTABODY CONSTRUCTS, COMPOSITIONS, AND METHODS**

Publication Classification

(71) Applicants: **The Hospital for Sick Children,**
Toronto (CA); **THE GOVERNING COUNCIL OF THE UNIVERSITY OF TORONTO,** Toronto (CA)
(72) Inventors: **Jean-Philippe Julien,** Toronto (CA);
Eduarne Rujas Diez, Vitoria (ES);
Bebhinn Treanor, Toronto (CA); **Tian Tian Zhao,** Toronto (CA)

(51) **Int. Cl.**
A61K 47/64 (2017.01)
A61K 47/68 (2017.01)
C07K 16/46 (2006.01)
(52) **U.S. Cl.**
CPC **A61K 47/64** (2017.08); **A61K 47/68**
(2017.08); **C07K 16/46** (2013.01); **C07K**
2317/71 (2013.01); **C07K 2319/00** (2013.01)

(21) Appl. No.: **18/578,239**
(22) PCT Filed: **Jul. 12, 2022**
(86) PCT No.: **PCT/CA2022/051084**
§ 371 (c)(1),
(2) Date: **Jan. 10, 2024**

Related U.S. Application Data

(60) Provisional application No. 63/220,920, filed on Jul. 12, 2021, provisional application No. 63/289,016, filed on Dec. 13, 2021.

(57) **ABSTRACT**

In aspects, a self-assembled polypeptide complex comprises (a) a plurality of first fusion polypeptides, each first fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises a Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, and (b) a plurality of second fusion polypeptides, each second fusion polypeptide comprising (1) an antigen-binding antibody fragment linked to (2) a nanocage monomer or subunit thereof.

Specification includes a Sequence Listing.

Human ferritin light chain (hFTL)

N-term

C-term

N-hFTL (residues 1-90)	C-hFTL (residues 91-175)
------------------------	--------------------------




FIG. 1A




FIG. 1B




FIG. 1C




FIG. 1D




FIG. 2A




FIG. 2B





FIG. 3A

**FIG. 3B**




FIG. 3C




FIG. 3D





FIG. 3E

**FIG. 3F**





FIG. 4A

**FIG. 4B**





FIG. 4C

**FIG. 4D**




FIG. 4E




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 7




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 8D




FIG. 9A




FIG. 9B




FIG. 10A




FIG. 10B




FIG. 11




FIG. 12




FIG. 13




FIG. 14




FIG. 15




FIG. 16




FIG. 17




FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22

OPTIMIZED MULTABODY CONSTRUCTS, COMPOSITIONS, AND METHODS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application is a U.S. National Stage Application of International Patent Application No. PCT/CA2022/051084, filed Jul. 12, 2022, which claims the benefit of and priority to U.S. Provisional Patent Application Nos. 63/220,920, filed Jul. 12, 2021, and 63/289,016, filed Dec. 13, 2021, the entire contents of each of which are hereby incorporated by reference in their entirety for all purposes.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in XML format and is hereby incorporated by reference in its entirety. Said XML copy, created on Jul. 8, 2024, is named "RBT-006WOUS_SL.xml" and is 64,353 bytes in size.

BACKGROUND

[0003] Therapeutics based on antibodies or antibody fragments are being developed for a variety of uses, e.g., for treating various diseases or conditions.

[0004] However, antibody-based therapeutics may need to be tuned so that they have desirable characteristics (for example, desirable biodistribution, half-lives, etc.) after administration to a subject. Some antibody-based therapeutics have a format that differs from that of a native immunoglobulin molecule. For example, in some therapeutics, an antibody or antibody fragment is fused to another polypeptide; in some, the antibody or antibody fragment(s) is in a configuration or has a valence not found in nature. These antibody-based therapeutics may also need to be tuned.

[0005] Thus, there remains a need for optimized antibody-based therapeutics.

SUMMARY

[0006] The present invention addresses this need with the provision of a suite of optimized self-assembled polypeptide complexes comprising antibody fragments. Depending on the desired characteristics (e.g., pharmacokinetic characteristics), a particular self-assembled polypeptide complex or set of complexes can be chosen for use. For example, in certain embodiments, provided are self-assembled polypeptide complexes that have half-lives or other pharmacokinetic characteristics similar to that of IgG molecules.

[0007] In accordance with an aspect, there is provided a self-assembled polypeptide complex comprising

[0008] (a) a plurality of first fusion polypeptides, each first fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises a Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, and

[0009] (b) a plurality of second fusion polypeptides, each second fusion polypeptide comprising (1) an antigen-binding antibody fragment linked to (2) a nanocage monomer or subunit thereof.

[0010] In an aspect, (1) if the Fc polypeptide is an IgG1 Fc polypeptide, the antigen-binding fragment is not a Fab fragment that binds to SARS-CoV-2; and/or (2) if the

nanocage monomer is a mouse ferritin monomer and the Fc polypeptide is a mouse IgG2a Fc polypeptide, the antigen-binding antibody fragment is not a Fab fragment that binds to CD19.

[0011] In an aspect, the nanocage monomer is a ferritin monomer.

[0012] In an aspect, the ferritin monomer is a ferritin light chain.

[0013] In an aspect, the self-assembled polypeptide complex does not comprise any ferritin heavy chains or subunits of ferritin heavy chains.

[0014] In an aspect, the ferritin monomer is a human ferritin.

[0015] In an aspect, the Fc polypeptide is an IgG1 Fc polypeptide.

[0016] In an aspect, the Fc polypeptide is an IgG2 Fc polypeptide.

[0017] In an aspect, the Fc polypeptide is a single-chain Fc (scFc).

[0018] In an aspect, the Fc polypeptide is an Fc monomer.

[0019] In an aspect, the antigen-binding antibody fragment comprises a light chain variable domain and a heavy chain variable domain.

[0020] In an aspect, the antigen-binding antibody fragment is a Fab fragment.

[0021] In an aspect, each second fusion polypeptide does not comprise any CH2 or CH3 domains.

[0022] In an aspect, the one or more mutations comprise a mutation or set of mutations associated with altered binding to FcRn.

[0023] In an aspect, the mutation or set of mutations comprises a mutation at one or more of the following residues: M252, I253, S254, T256, K288, M428, and N434, or combinations thereof, wherein numbering is according to the EU index.

[0024] In an aspect, the mutation or set of mutations comprises mutations at M428 and N434, wherein numbering is according to the EU index.

[0025] In an aspect, the mutation or set of mutations comprises M428L and N434S mutations, wherein numbering is according to the EU index.

[0026] In an aspect, the altered binding to FcRn is decreased binding to FcRn.

[0027] In an aspect, the mutation or set of mutations associated with decreased binding to FcRn is selected from the group consisting of I253A, I253V, and K288A, and combinations thereof, wherein numbering is according to the EU index.

[0028] In an aspect, the one or more mutations comprise a mutation or set of mutations associated with altered effector function.

[0029] In an aspect, the Fc polypeptide is an IgG1 Fc polypeptide, and wherein the mutation or set of mutations comprises a mutation at one or more the following residues: L234, L235, G236, G237, P329, and A330, or combinations thereof, wherein numbering is according to the EU index.

[0030] In an aspect, the altered effector function is decreased effector function.

[0031] In an aspect, the mutation or set of mutations associated with decreased effector function is selected from the group consisting of LALA (L234A/L235A), LALAP (L234A/L235A/P329G), G236R, G237A, and A330L, wherein numbering is according to the EU index.

[0032] In an aspect, the nanocage monomer or subunit thereof is a ferritin monomer subunit, and

[0033] a. each first fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is N-half-ferritin; or

[0034] b. each first fusion polypeptide comprises a ferritin monomer subunit which is N-half ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin.

[0035] In an aspect, the self-assembled polypeptide complex is characterized by a 1:1 ratio of first fusion polypeptides to second fusion polypeptides.

[0036] In an aspect, within each first fusion polypeptide, the Fc polypeptide is linked to the nanocage monomer or subunit thereof via an amino acid linker.

[0037] In an aspect, within each first fusion polypeptide, the Fc polypeptide is linked to the nanocage monomer or subunit thereof at the N-terminus of the nanocage monomer or subunit thereof.

[0038] In an aspect, within each second fusion polypeptide, the antigen-binding antibody fragment is linked to the nanocage monomer or subunit thereof via an amino acid linker.

[0039] In an aspect, within each second fusion polypeptide, the antigen-binding antibody fragment is linked to the nanocage monomer or subunit thereof at the N-terminus of the nanocage monomer or subunit thereof.

[0040] In an aspect, the self-assembled polypeptide complex further comprises a plurality of third fusion polypeptides, each third fusion polypeptide comprising (1) an antigen-binding antibody fragment linked to (2) a nanocage monomer or a subunit thereof, wherein the third fusion polypeptide is different than the second fusion polypeptide.

[0041] In an aspect, the antigen-binding antibody fragment within the third fusion polypeptide comprises a light chain variable domain and a heavy chain variable domain.

[0042] In an aspect, the antigen-binding antibody fragment within the third fusion polypeptide is a Fab fragment.

[0043] In an aspect, each third fusion polypeptide does not comprise any CH2 or CH3 domains.

[0044] In an aspect, the antigen-binding antibody fragment of the second fusion polypeptide is capable of binding a first epitope, the antigen-binding fragment of the third fusion polypeptide is capable of binding a second epitope, and the first epitopes and second epitopes are distinct and non-overlapping.

[0045] In an aspect, first epitopes and second epitopes are from the same protein.

[0046] In an aspect, the self-assembled polypeptide complex comprises a total of 24 to 48 fusion polypeptides.

[0047] In an aspect, the self-assembled polypeptide complex comprises a total of at least 24 fusion polypeptides.

[0048] In an aspect, the self-assembled polypeptide complex comprises a total of at least 32 fusion polypeptides.

[0049] In an aspect, the self-assembled polypeptide complex has a total of about 32 fusion polypeptides.

[0050] In an aspect, the half-life of the self-assembled polypeptide complex is at least 3 days, at least 5 days, at least 7 days, at least 10 days, at least 14 days, at least 21 days, or at least 28 days when administered to a subject in need thereof.

[0051] In an aspect, the self-assembled polypeptide complex is characterized in that, after administration of a com-

position comprising the self-assembled polypeptide complex, the self-assembled polypeptide complex has a half-life substantially similar to that of a reference IgG molecule administered by the same route of administration and in a similar composition.

[0052] In an aspect, the reference IgG molecule is an antibody from which the antigen-binding antibody fragment within the second fusion polypeptide is derived or is an antibody from which the antigen-binding antibody fragment within the third fusion polypeptide is derived.

[0053] In an aspect, the half-life of the self-assembled polypeptide complex is from about 3 to about 35 days when administered to a subject in need thereof.

[0054] In an aspect, the self-assembled polypeptide complex is detectable in serum after at least 3 days, at least 5 days, at least 7 days, at least 10 days, at least 14 days, at least 21 days, or at least 28 days after administration to a subject in need thereof.

[0055] In an aspect, the area-under-the-curve (AUC) of the self-assembled polypeptide complex is at least 10 day·μg/mL, at least 25 day·μg/mL, at least 50 day·μg/mL, at least 100 day·μg/mL, at least 200 day·μg/mL, at least 300 day·μg/mL, at least 400 day·μg/mL, at least 500 day·μg/mL, at least 750 day·μg/mL, at least 1000 day·μg/mL, at least 1500 day·μg/mL, at least 2000 day·μg/mL, at least 2500 day·μg/mL, at least 3000 day·μg/mL, at least 4000 day·μg/mL, at least 5000 day·μg/mL, at least 6000 day·μg/mL, at least 7000 day·μg/mL, or at least 8000 day·μg/mL when administered to a subject in need thereof.

[0056] In an aspect, the area-under-the-curve (AUC) of the self-assembled polypeptide complex is from about 10 to about 8000 day·μg/mL when administered to a subject in need thereof.

[0057] In an aspect, the maximum concentration (C_{max}) of the self-assembled polypeptide complex is at least 10 μg/mL, at least 25 μg/mL, at least 50 μg/mL, at least 100 μg/mL, at least 250 μg/mL, at least 500 μg/mL, at least 750 μg/mL, at least 1 mg/mL, at least 10 mg/mL, at least 25 mg/mL, at least 50 mg/mL, at least 75 mg/mL, at least 100 mg/mL, at least 250 mg/mL, at least 500 mg/mL, or at least 750 mg/mL when administered to a subject in need thereof.

[0058] In an aspect, the maximum concentration (C_{max}) of the self-assembled polypeptide complex is from about 10 μg/mL to about 750 mg/mL when administered to a subject in need thereof.

[0059] In an aspect, the subject is human.

[0060] In an aspect, administration to the subject is by parenteral administration.

[0061] In an aspect, administration to the subject is by subcutaneous administration, intravenous administration, intramuscular administration, intranasal administration, or by inhalation.

[0062] In an aspect, the self-assembled polypeptide complex is characterized in that the self-assembled polypeptide complex induces antibody-dependent cellular phagocytosis (ADCP) in an in vitro model of ADCP.

[0063] In an aspect, the ADCP is induced at a level of at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, or at least 60% internalization of target.

[0064] In accordance with an aspect, there is provided a method comprising administering a composition comprising the self-assembled polypeptide complex described herein to a mammalian subject.

[0065] In an aspect, the subject is human.

[0066] In an aspect, the method comprises administration by a systemic route.

[0067] In an aspect, the systemic route comprises subcutaneous, intravenous, or intramuscular injection, inhalation, or intranasal administration.

[0068] In an aspect, after administration, the half-life of the self-assembled polypeptide complex in the mammalian subject is at least 3 days, at least 5 days, at least 7 days, at least 10 days, at least 14 days, at least 21 days, or at least 28 days.

[0069] In an aspect, after administration, the half-life of the self-assembled polypeptide complex in the mammalian subject is from 3 to 35 days.

[0070] In an aspect, after administration, the area-under-the-curve (AUC) of the self-assembled polypeptide complex in the mammalian subject is at least 10 day· $\mu\text{g/mL}$, at least 25 day· $\mu\text{g/mL}$, at least 50 day· $\mu\text{g/mL}$, at least 100 day· $\mu\text{g/mL}$, at least 200 day· $\mu\text{g/mL}$, at least 300 day· $\mu\text{g/mL}$, at least 400 day· $\mu\text{g/mL}$, at least 500 day· $\mu\text{g/mL}$, at least 750 day· $\mu\text{g/mL}$, at least 1000 day· $\mu\text{g/mL}$, at least 1500 day· $\mu\text{g/mL}$, at least 2000 day· $\mu\text{g/mL}$, at least 2500 day· $\mu\text{g/mL}$, at least 3000 day· $\mu\text{g/mL}$, at least 4000 day· $\mu\text{g/mL}$, at least 5000 day· $\mu\text{g/mL}$, at least 6000 day· $\mu\text{g/mL}$, at least 7000 day· $\mu\text{g/mL}$, or at least 8000 day· $\mu\text{g/mL}$.

[0071] In an aspect, after administration, the area-under-the-curve (AUC) of the self-assembled polypeptide complex in the mammalian subject is from about 10 to about 8000 day· $\mu\text{g/mL}$.

[0072] In an aspect, after administration, the maximum concentration (C_{max}) of the self-assembled polypeptide complex in the mammalian subject is at least 10 $\mu\text{g/mL}$, at least 25 $\mu\text{g/mL}$, at least 50 $\mu\text{g/mL}$, at least 100 $\mu\text{g/mL}$, at least 250 $\mu\text{g/mL}$, at least 500 $\mu\text{g/mL}$, at least 750 $\mu\text{g/mL}$, at least 1 mg/mL, at least 10 mg/mL, at least 25 mg/mL, at least 50 mg/mL, at least 75 mg/mL, at least 100 mg/mL, at least 250 mg/mL, at least 500 mg/mL, or at least 750 mg/mL.

[0073] In an aspect, after administration, the maximum concentration (C_{max}) of the self-assembled polypeptide complex in the mammalian subject is from about 10 $\mu\text{g/mL}$ to about 750 mg/mL.

[0074] In accordance with an aspect, there is provided a fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises a Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, wherein the one or more mutations comprise a mutation or set of mutations associated with altered binding to FcRn and/or with altered effector function.

[0075] In an aspect, the nanocage monomer is a ferritin monomer.

[0076] In an aspect, the ferritin monomer is a ferritin light chain.

[0077] In an aspect, the fusion polypeptide does not comprise any ferritin heavy chains or subunits of ferritin heavy chains.

[0078] In an aspect, the ferritin monomer is a human ferritin.

[0079] In an aspect, the Fc polypeptide is an IgG1 Fc polypeptide.

[0080] In an aspect, the Fc polypeptide is an IgG2 Fc polypeptide.

[0081] In an aspect, the Fc polypeptide is a single-chain Fc (scFc).

[0082] In an aspect, the mutation or set of mutations comprises a mutation at one or more of the following residues: M252, I253, S254, T256, K288, M428, and N434, or combinations thereof, wherein numbering is according to the EU index.

[0083] In an aspect, the altered binding to FcRn is decreased binding to FcRn.

[0084] In an aspect, the mutation or set of mutations associated with decreased binding to FcRn is selected from the group consisting of I253A, I253V, and K288A, and combinations thereof, wherein numbering is according to the EU index.

[0085] In an aspect, the Fc polypeptide is an IgG1 Fc polypeptide, and wherein the mutation or set of mutations comprises a mutation at one or more of the following residues: L234, L235, G236, G237, P329, and A330, or combinations thereof, wherein numbering is according to the EU index.

[0086] In an aspect, the altered effector function is decreased effector function.

[0087] In an aspect, the mutation or set of mutations associated with decreased effector function is selected from the group consisting of LALA (L234A/L235A), LALAP (L234A/L235A/P329G), G236R, G237A, and A330L, wherein numbering is according to the EU index.

[0088] In an aspect, the nanocage monomer or subunit thereof is a ferritin monomer subunit, and

[0089] a. each first fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is N-half-ferritin; or

[0090] b. each first fusion polypeptide comprises a ferritin monomer subunit which is N-half ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin.

[0091] In an aspect, within each first fusion polypeptide, the Fc polypeptide is linked to the nanocage monomer or subunit thereof via an amino acid linker.

[0092] In an aspect, within each first fusion polypeptide, the Fc polypeptide is linked to the nanocage monomer or subunit thereof at the N-terminus of the nanocage monomer or subunit thereof.

BRIEF DESCRIPTION OF THE DRAWINGS

[0093] FIG. 1A is a diagrammatic representation of human ferritin light chain (hFTL) and exemplary N-half ferritin and C-half ferritin molecules.

[0094] FIGS. 1B, 1C, and 1D are diagrammatic representations of exemplary Multabodies formation of the disclosure.

[0095] FIG. 2A are negative stain electron micrographs of T-01 MB, T-02 MB, and T-01 MB.v2, each containing wild-type IgG1 Fc.

[0096] FIG. 2B are negative stain electron micrographs of T-01 MB containing various Fc.

[0097] FIGS. 3A and 3B provide biolayer interferometry (BLI) concentration-response curves for the binding of T-01 MB containing various Fc to human FcRn at pH 5.6 and pH 7.4, respectively.

[0098] FIGS. 3C, 3D, and 3E provide BLI concentration-response curves for the binding of T-01 MB containing various Fc to human Fc γ RIIa, human Fc γ RIIb, and Fc γ RI, respectively.

[0099] FIG. 3F provides BLI concentration-response curves for the binding of T-01 MB.v2 containing wild-type IgG1 Fc or Fc IgG1 with M428L/N434S (LS) mutations to human Fc receptors.

[0100] FIG. 4A provides BLI concentration-response curves for the binding of T-01 MB, T-02 MB, and T-01 MB.v2 to target/epitope of PGDM1400, N49P7, 10E8v4, or iMab, as included as Fab in the Multabodies.

[0101] FIGS. 4B, 4C, and 4D provide BLI concentration-response curves for the binding of T-01 MB containing various Fc to the target/epitope of PGDM1400 (BG5050 SOSIP.664 D368R), N49P7 (93TH057), and 10E8v4 (gp41 MPER), respectively.

[0102] FIG. 4E provides BLI concentration-response curves for the binding of PGDM1400, N49P7, 10E8v4, or iMab IgG to BG5050 SOSIP.664 D368R, 93TH057, gp41 MPER, and CD4.

[0103] FIG. 5A illustrates an experiment described in Example 4 and conducted using CB17/Icr-Prkdc^{scid}/IcrIco-Crl immunodeficient (SCID) mice. FIGS. 5B, 5C, 5D, and 5E illustrates serum levels of testing Multabodies or IgG1 control following subcutaneous administration in SCID mice.

[0104] FIG. 6A illustrates an experiment described in Example 4 and conducted using NOD/Shi-scid/IL-2R^γ null immunodeficient (NCG) mice. FIG. 6B illustrates serum levels of testing Multabodies or IgG1 control following subcutaneous administration in NCG mice.

[0105] FIG. 6C provides body weights of NCG mice upon administration of testing Multabodies or IgG1 control. Mean values $\pm$ SD for n=3 mice are displayed.

[0106] FIG. 7 illustrates the dose-dependent phagocytosis induced by testing Multabodies or controls, quantified and represented as the percentage increase in the internalization of 93TH057-coated microspheres compared to non-coated microspheres. *P<0.05, ***P<0.001 and P ****<0.0001. n=four biologically-independent samples.

[0107] FIGS. 8A, 8B, and 8C illustrate the breadth and median IC₅₀ values (μ g/mL) of testing Multabodies (T-01 MB, T-01 MB.v2, and T-02 MB., respectively) (diamond), parental antibodies PGDM1400, N49P7, and 10E8v4 (circles), IgG1 control (triangle), and N6/PGDM1400x10E8v4 trisppecific antibody (inverted triangle), determined with a TZM-bl assay described in Example 6.

[0108] FIG. 8D illustrates the dose-dependent neutralization of HIV-1 by T-01 MB containing various Fc, determined with a TZM-bl assay described in Example 6.

[0109] FIG. 9A illustrates the inhibition of CXCR4-tropic HIV-1 isolate IIIB infection of PBMCs by T-01 MB, T-01 MB.v2, or IgG1 control. The mean values $\pm$ SD for three technical replicates are shown. FIG. 9B shows the percent of viable cells following T-01 MB, T-01 MB.v2, or IgG1 control relative to untreated control cells.

[0110] FIGS. 10A and 10B show 4-week stability under temperature stress conditions (40° C.) of T-01 MB and T-01 MB.v2. See Example 8.

[0111] FIG. 11. HIV-1 bNAb multimerization increases neutralization potency. Schematic of the self-assembly of (A) apoferritin (24 subunits) and (B) single-chain Fab-apoferritin fusions. Fab light chain (LC) and heavy chain (HC) are shown in light and dark pink, respectively, and are connected to the N terminus of the light chain of human apoferritin (grey) through a GGS-like flexible linker (dark).

(C) Schematic representation of different Fab densities displayed on human apoferritin. Co-transfection of scFab-human apoferritin-encoding plasmids together with unconjugated apoferritin at ratios of 1:4 (dark yellow), 1:1 (black), 4:1 (blue) and 1:0 (red) resulted in molecules with different scFab valency, as confirmed by elution volumes in size exclusion chromatography and less unconjugated apoferritin in SDS-PAGE. Negative stain electron micrographs of the samples with the lowest and highest scFab valency are shown (scale bar 50 nm). (D) Avidity effect on neutralization of five bNAbs against a five-PsV panel (PVO.04, JRCSEF, BG505 T332N, THRO4156.18 and t278-50). Fold potency increase was calculated as the parental IgG IC₅₀ (μ g/mL) divided by the Fab-apoferritin fusion IC₅₀ (μ g/mL). Fold potency increase analyses were omitted in the following cases: N49P7-t278-50, VRC01-T278-50 and 10-1074-THRO4156.18 due to neutralization resistance. Bars ($\pm$ SD) represent the mean value from n=3 biologically independent samples.

[0112] FIG. 12. Characterization of scFab-apoferritin fusions. (A) SDS-PAGE bands corresponding to scFab-apoferritin and unconjugated apoferritin were quantified by densitometry using the ImageJ software (rsb.info.nih.gov/ij/). Intensity plots of the bands in each lane (yellow box) are shown (B). (C) The approximate number of scFabs displayed on the particles was calculated as follows: intensity of scFab band/total intensity, and compared with the theoretical numbers inferred from DNA ratios used for co-transfection.

[0113] FIG. 13. Design, assembly and neutralization profile of HIV Multabodies against a 14-PsV panel. (A) Schematic of the human apoferritin split design that drives hetero-dimerization of scFab-human apoferritin subunits. (B) Size exclusion chromatography in-line with multi-angle light scattering of 24-mer PGDM1400 scFab-apoferritin particles (black) and T-01 MB (dark red). The molar mass of each elution peak (lines under UV absorbance) is shown in MDa. (C) Negative stain electron micrographs of T-01 MB (scale bar 50 nm). (D) Concentration-response curves for binding of T-01 MB to multiple epitopes. PGDM1400, N49P7 and 10E8 binding sites are colored in red, blue and pink, respectively on the surface representation of the HIV-1 Env trimer (grey). Red lines represent raw data; black lines represent global fits. (E) Breadth (cutoff IC₅₀ set at 10 μ g/mL) and median IC₅₀ values (μ g/mL) of T-01 MB (red diamond), parental bNAbs (white circles), IgG combination (gray triangle) and the N6/PGDM1400x10E8v4 tri-specific antibody (black triangle). The 14-PsV panel was selected based on susceptibility and resistance to the parental IgGs. (F) Individual IC₅₀ values (μ g/mL) to each PsV variant. The solid line denotes the median neutralization IC₅₀ of all 14 viral strains. Those pseudoviruses that show the highest neutralization resistance are highlighted by a red box. IC₅₀ values in (D) and (E) were calculated from three biological replicates.

[0114] FIG. 14. Multabody affinity-purification scheme. Protein A and Protein L sequential affinity purification. Binding to Protein A enriches for Multabodies with Fc (green), while Protein L enriches for Multabodies with the kappa chain Fab PGDM1400 (blue). Complementation of the two halves of the apoferritin split design ensures the presence of N49P7/iMab (orange) and 10E8v4 scFabs (pink) (fused to C-ferritin) during the protein A purification step. An alanine-to-proline point mutation at position 12 of the

kappa chain of iMab was introduced to disrupt binding to Protein L. Gel filtration is performed to separate any aggregated material or non-assembled component.

[0115] FIG. 15. Generation of a Multabody that cross-targets the HIV-1 Env and the CD4 receptor. (A) Schematic representation of MB components. (B) Size exclusion chromatography in-line with multi-angle light scattering of 24-mer PGDM1400 scFab-apoferritin particles (black) and T-02 MB (blue). The molar mass of each sample is shown in MDa. (C) Negative stain electron micrographs (scale bar 50 nm) and (D) binding profile of T-02 MB. (E) Breadth (cutoff IC_{50} set at 10 $\mu\text{g/mL}$) and median IC_{50} values ($\mu\text{g/mL}$) of T-02 MB (red diamond) in comparison to its parental bNAbs (white circles) and IgG combination (gray triangle). The mean IC_{50} values ($\mu\text{g/mL}$) derived from three biological replicates for those pseudoviruses that show the highest neutralization resistance to T-02 MB are shown.

[0116] FIG. 16. Biophysical characterization of HIV-1 Multabodies. Comparison of the T_m and T_{agg} temperatures of T-01/T-02 MB, 12-mer ferritin fusions, parental IgGs and the N6/PGDM1400x10 E8v4 tri-specific antibody.

[0117] FIG. 17. Binding characteristics of IgGs binding to four different antigens. BLI response curves of IgG binding to 93TH057 gp120, BG505 SOSIP.664_D368R, MPER peptide and CD4 immobilized onto Ni-NTA biosensors. The BG505 SOSIP.664_D368R trimer and 93TH057 gp120 monomer were selected as epitope-specific ligands for PGDM1400 and N49P7, respectively. Red lines represent raw data; black lines represent global fits.

[0118] FIG. 18. Engineering and biophysical characterization of Multabody v2. (A) The second-generation Multabody design displays two distinct features in comparison to the original Multabody design: 1) the Fc (green) is fused to the C terminus of the second half of apoferritin in the split ferritin design; and 2) the single-chain Fc domain (green), fused to the C terminus of an apoferritin half protomer is reverted to a monomeric Fc chain. Dimerization of each Fc in MB.v2 drives assembly of four Fabs (two Fab2 and two Fab3—bottom row) while only one Fab is assembled per Fc into the previous MB version (top row). (B) Negative stain electron micrographs of T-01 MB.v2 (scale bar 50 nm). (C) Concentration-response curves for binding of T-01 MB.v2 to multiple epitopes. Red lines represent raw data; black lines represent global fits. (D) Comparison of T_{agg} , and (E) long-term stability under temperature stress conditions (10 mg/ml; 40° C.) of the two different Multabody versions. PsV neutralization (mean values $\pm$ SD for two technical replicates) comparison at week 0 vs week 4 is shown.

[0119] FIG. 19. Multabody v2 features. (A) Modification of the fusion of the Fc (green) from the N terminus of N-ferritin (top row) to the C terminus of C-ferritin (bottom row) inverts the orientation of the Fc in the Multabody. (B) Fc dimerization of two Fc chains fused to the C terminus of two independent ferritin subunits at the 4-fold symmetry axes of the apoferritin nanocage acts as an additional driver of Multabody v2 assembly.

[0120] FIG. 20. Fine-tuning of Fc on Multabody for IgG-like characteristics. (A) Concentration-response curves for pH dependent binding to human FcRn by T-01 MB and T-01 MB.v2. (B) Comparison of the FcRn apparent binding affinities (K_D) at acidic pH between MBs and an IgG1. $n=3$ biologically independent samples are shown. Apparent K_D lower than 10^{-12} M (dash black line) is beyond the instrument detection limit. (C) Concentration-response curves to

the high-affinity human Fc γ RI (top), and low-affinity human Fc γ RIIa (bottom). (D) Dose-dependent phagocytosis determined as a percentage increase in internalization of 93TH057-coated fluorescent microspheres compared to a no-antibody control. Anti-human FcR binding inhibitor antibody was added to block Fc-mediated internalization (dark red). Data were analyzed by two-way ANOVA with Tukey's multiple comparisons test. Each group was compared to the IgG negative control. * $P<0.05$, *** $P<0.001$ and P **** <0.0001 . IgG and MB samples with no affinity for the antigen coated beads were added as control samples. $n=4$ biologically-independent samples. (E) Serum levels after subcutaneous administration of 5 mg/kg of Multabodies or parental IgG mixture in female NOD/Shi-scid/IL-2R γ null (NCG) immunodeficient mice. (F) Body weight upon administration of 5 mg/kg of molecules in NCG mice. Mean values $\pm$ SD for $n=3$ mice are shown in (E) and (F).

[0121] FIG. 21. Broad and potent neutralization by Multabody v2 against extended HIV-1 PsV panels. (A) Breadth and median IC_{50} values ($\mu\text{g/mL}$) of T-01 Multabody versions (different shades of red diamond), individual IgGs (black circles) and IgG mixture (blue triangle) against a 25-PsV panel with 56% of PsV variants resistant to PGDM1400 neutralization. (B) Individual IC_{50} values ($\mu\text{g/mL}$) for each PsV variant. IC_{50} values in (A) and (B) were calculated from three biological replicates. (C) Potency (IC_{50})-breadth (left) and potency (IC_{80})-breadth (right) curve comparison of T-01 Multabody versions, as well as parental IgGs and an IgG mixture against an extended multiclade panel of 118 HIV-1 PsV variants. (D) Individual IC_{50} (left) and IC_{80} (right) values for each PsV variant in (C). Yellow dots correspond to IC_{50} values from PsVs that are highly resistant to PGDM1400 neutralization. The solid line in (B) and (D) denotes the median IC_{50} neutralization titer of all viral strains in each panel.

[0122] FIG. 22. PsV neutralization and inhibition of primary PBMC infection by Multabodies. (A) Molar potency (IC_{50})-breadth (left) and molar potency (IC_{80})-breadth (right) graphs comparing T-01 Multabody versions, as well as parental IgGs and an IgG mixture against a panel of 118 HIV-1 PsV variants. (B) Measurement of HIV-1 replication via p24 detection in PBMC culture supernatant derived from three different blood donors. Data shown represent HIV-1 replication levels at 7 days post-infection with the CXCR4-tropic HIV-1 isolate IIIB. Mean values $\pm$ SD for three technical replicates are shown. (C) Impact of IgG mix and Multabody treatments on cell viability, showing the percent of viable cells relative to untreated control cells under the same experimental conditions as in (B).

DETAILED DESCRIPTION

[0123] The inventors have previously described self-assembled polypeptide complexes comprising fusion polypeptides that comprise antibody fragments. These self-assembled polypeptide complexes can be designed and adapted for a variety of therapeutic purposes. For example, as described herein, a self-assembled polypeptide complex comprising both Fab- and Fc-containing fusion polypeptides may be used to target cells expressing an antigen to which the Fab can bind, and the Fc portion may mediate interactions with other molecules in the body.

[0124] In many situations, it may be desirable to optimize the behavior of these self-assembled polypeptide complexes after administration, e.g., by modulating characteristics such

as half-life and/or ability to mediate antibody-mediated effects. Techniques for optimizing IgG molecules are known in the art. For example, both half-life and the certain antibody-mediated effects can be modulated using modifications to the Fc region of an IgG molecule.

[0125] However, the inventors have surprisingly discovered that, in some situations, the techniques for optimizing IgG molecules, when applied to the inventors' self-assembled polypeptide complexes, may have unexpected effects. As one example, the inventors have discovered that an Fc modification typically associated with reduced FcRn binding (and reduced half-life) in the context of an IgG1 molecule actually conferred more desirable bioavailability characteristics in the context of the inventors' self-assembled polypeptide complexes.

[0126] Utilizing these and other insights, the inventors have developed a suite of a self-assembled polypeptide complexes, each of which is optimized for a particular desired outcome, and related methods.

[0127] For example, in certain embodiments, after administration to a subject, a provided self-assembled polypeptide complex has one or more pharmacokinetic features similar to that of a reference IgG molecule (e.g., an IgG molecule whose class matches the class of an Fc chain within an Fc polypeptide within the self-assembled polypeptide complex). For example, in some embodiments, a self-assembled polypeptide complex as disclosed herein has a similar bioavailability to that of reference IgG molecule. In some embodiments, a self-assembled polypeptide complex as disclosed herein has a similar half-life to that of reference IgG molecule.

[0128] In certain embodiments, after administration to a subject, a provided self-assembled polypeptide complex induces antibody-dependent cellular phagocytosis (ADCP).

Definitions

[0129] The terms “about” and “approximately,” when used herein in reference to a value, are used interchangeably and refer to a value that is similar to the referenced value. In general, those skilled in the art, familiar with the context, will appreciate the relevant degree of variation encompassed by “about” or “approximately” in that context. For example, in some embodiments, the terms “about” and “approximately” may encompass a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less of the referred value.

[0130] As used herein, the terms “alter,” “altered,” “decrease,” “decreased,” “increase,” “increased,” or “reduction,” “reduced,” (e.g., in reference to certain outcomes or effects) have meanings relative to a reference level. In some embodiments, in the context of discussing mutations in an Fc chain or Fc polypeptide, the reference level is a level known or as determined with an IgG that does not contain the referenced mutation(s) in the Fc region.

[0131] The terms “ferritin” and “apoferritin” are used interchangeably herein and generally refer to a polypeptide (e.g., a ferritin chain) that is capable of assembling into a ferritin complex which typically comprises 24 protein subunits. In some embodiments, the ferritin is a human ferritin, e.g., a human ferritin light chain, e.g., a human ferritin light chain having at least 85% sequence identity to SEQ ID NO:1

or UniProt P02792. In some embodiments, the ferritin is a wild-type ferritin. For example, the ferritin may be a wild-type human ferritin.

[0132] The term “ferritin monomer,” is used herein to refer to a single chain of a ferritin that, in the presence of other ferritin chains, is capable of self-assembling into a polypeptide complex comprising a plurality of ferritin chains, e.g., 24 or more ferritin chains.

[0133] As used herein, the term “linker” is used to refer to an entity that connects two or more elements to form a multi-element agent. For example, those of ordinary skill in the art appreciate that a polypeptide (e.g., fusion polypeptide) whose structure includes two or more functional or organizational domains often includes a stretch of amino acids between such domains that links them to one another. In some embodiments, a polypeptide comprising a linker element has an overall structure of the general form S1-L-S2, wherein S1 and S2 may be the same or different and represent two domains associated with one another by the linker (L). In some embodiments, the linker is an “amino acid linker,” that is, it comprises amino acid residues, e.g., an amino acid linker may comprise at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more amino acid residues. In some embodiments, a linker is characterized in that it tends not to adopt a rigid three-dimensional structure, but rather provides flexibility to the polypeptide.

[0134] The term “multispecific,” as used herein, refers to the characteristic of having at least two binding sites at which at least two different binding partners, e.g., an antigen or receptor (e.g., Fc receptor), can bind. For example, a polypeptide complex that comprises at least two Fab fragments, wherein each of the two Fab fragments is capable of binding to a different antigen, is “multispecific.” As an additional example, a polypeptide complex that comprises an Fc fragment (which is capable of binding to an Fc receptor) and a Fab fragment (which is capable of binding to an antigen) is “multispecific.”

[0135] The term “multivalent,” as used herein, refers to the characteristic of having at least two binding sites at which a binding partner, e.g., an antigen or receptor (e.g., Fc receptor), can bind. The binding partners that can bind to the at least two binding sites may be the same or different.

[0136] The term “nanocage monomer,” as used herein, refers to a single chain of a polypeptide that is capable of self-assembling with other nanocage monomers to form a self-assembled polypeptide complex comprising a plurality of nanocage monomers. In some embodiments, the nanocage monomer is selected from monomers of ferritin, apoferritin, encapsulin, sulfur oxygenase reductase (SOR), lumazine synthase, pyruvate dehydrogenase, carboxysome, vault proteins, GroEL, heat shock protein, E2P coat protein, MS2 coat protein, fragments thereof, and variants thereof.

[0137] The term “polypeptide,” as used herein, generally has its art-recognized meaning of a polymer of at least three amino acids, e.g., linked to each other by peptide bonds. Those of ordinary skill in the art will appreciate that the term “polypeptide” is intended to be sufficiently general as to encompass not only polypeptides having a complete sequence recited herein, but also to encompass polypeptides that represent functional fragments (i.e., fragments retaining at least one activity) of such complete polypeptides. Moreover, those of ordinary skill in the art understand that protein

sequences generally tolerate some substitution without destroying activity. Thus, any polypeptide that retains activity and shares at least about 30-40% overall sequence identity, often greater than about 50%, 60%, 70%, or 80%, and further usually including at least one region of much higher identity, often greater than 90% or even 95%, 96%, 97%, 98%, or 99% in one or more highly conserved regions, usually encompassing at least 3-4 and often up to 20 or more amino acids, with another polypeptide of the same class, is encompassed within the relevant term “polypeptide” as used herein. Polypeptides may contain L-amino acids, D-amino acids, or both and may contain any of a variety of amino acid modifications or analogs known in the art. Useful modifications include, e.g., terminal acetylation, amidation, methylation, etc. In some embodiments, proteins may comprise natural amino acids, non-natural amino acids, synthetic amino acids, and combinations thereof.

[0138] The term “self-assembled,” when used in reference to a macromolecular complex (e.g., a polypeptide complex), refers to the spontaneous formation of that complex when sufficient constituents of the complex (e.g., fusion polypeptides) to be formed are present. In some embodiments, complexes self-assemble in physiological conditions, or in a buffer (e.g., a solution) that corresponds to physiological conditions.

[0139] As used herein, the term “subject” to an organism, typically a mammal (e.g., a human). In some embodiments, a subject is suffering from or susceptible to a relevant disease, disorder or condition. In some embodiments, a subject displays one or more symptoms or characteristics of a disease, disorder or condition. In some embodiments, a subject is someone with one or more features characteristic of susceptibility to or risk of a disease, disorder, or condition. In some embodiments, a subject is a patient. In some embodiments, a subject is a subject to whom diagnosis and/or therapy is and/or has been administered.

[0140] As used herein, the term “treatment” (also “treat” or “treating”) refers to any administration of a therapy that partially or completely alleviates, ameliorates, relieves, inhibits, delays onset of, reduces severity of, and/or reduces incidence of one or more symptoms, features, and/or causes of a particular disease, disorder, and/or condition. In some embodiments, such treatment may be of a subject who does not exhibit signs of the relevant disease, disorder and/or condition and/or of a subject who exhibits only early signs of the disease, disorder, and/or condition. Alternatively, or additionally, such treatment may be of a subject who exhibits one or more established signs of the relevant disease, disorder and/or condition. In some embodiments, treatment may be of a subject who has been diagnosed as suffering from the relevant disease, disorder, and/or condition. In some embodiments, treatment may be of a subject known to have one or more susceptibility factors that are statistically correlated with increased risk of development of the relevant disease, disorder, and/or condition.

A. Fusion Polypeptides

[0141] In many embodiments, fusion polypeptides compatible with compositions and methods disclosed herein generally comprise a nanocage monomer or subunit thereof linked to either an Fc polypeptide or to an antigen-binding antibody fragment. Within the fusion polypeptide, the Fc polypeptide or the antigen-binding antibody fragment may be linked to the nanocage monomer or subunit thereof at a

particular terminus of the nanocage monomer or subunit thereof, e.g., the N-terminus or the C-terminus. In some embodiments, the Fc polypeptide or antigen-binding antibody fragment is linked via an amino acid linker, such as a linker as described herein.

[0142] In some embodiments, (1) if the Fc polypeptide is an IgG1 Fc polypeptide, the antigen-binding fragment is not an Fab fragment that binds to SARS-CoV-2; and/or (2) if the nanocage monomer is a mouse ferritin monomer and the Fc polypeptide is a mouse IgG2a Fc polypeptide, the antigen-binding antibody fragment is not an Fab fragment that binds to CD19.

1. Nanocage Monomers and Subunits Thereof

[0143] In some embodiments, the nanocage monomer is a ferritin monomer.

[0144] The term “ferritin monomer,” is used herein to refer to a single chain of a ferritin that, in the presence of other ferritin chains, is capable of self-assembling into a polypeptide complex comprising a plurality of ferritin chains, e.g., 24 or more ferritin chains. In some embodiments, the ferritin monomer is a ferritin light chain. In some embodiments, the ferritin monomer does not include a ferritin heavy chain or other ferritin components capable of binding to iron.

[0145] In some embodiments, each fusion polypeptide within the self-assembled polypeptide complex comprises a ferritin light chain or a subunit of a ferritin light chain. In these embodiments, the self-assembled polypeptide complex does not comprise any ferritin heavy chains or subunits of ferritin heavy chains.

[0146] In some embodiments, the ferritin monomer is a human ferritin chain, e.g., a human ferritin light chain, e.g., a human ferritin light chain having the sequence of at least residues 2-175 of SEQ ID NO: 1. In some embodiments, the ferritin monomer is a mouse ferritin chain.

[0147] A “subunit” of a ferritin monomer refers to a portion of a ferritin monomer that is capable of spontaneously associating with another, distinct subunit of a ferritin monomer, so that the subunits together form a ferritin monomer, which ferritin monomer, in turn, is capable of self-assembling with other ferritin monomers to form a polypeptide complex.

[0148] In some embodiments, the ferritin monomer subunit comprises approximately half of a ferritin monomer. As used herein, the term “N-half ferritin” refers to approximately half of a ferritin chain, which half comprises the N-terminus of the ferritin chain. As used herein, the term “C-half ferritin” refers to approximately half a ferritin chain, which half comprises the C-terminus of the ferritin chain. The exact point at which a ferritin chain may be divided to form the N-half ferritin and the C-half ferritin may vary depending on the embodiment. In the context of ferritin monomer subunits based on human ferritin light chain, for example, the halves may be divided at a point that corresponds to a position from about position 75 to about position 100 of SEQ ID NO:1 (or a substantial portion thereof). For example, in some embodiments, an N-half ferritin based on a human ferritin light chain has an amino acid sequence corresponding to residues 1-95 of SEQ ID NO: 1 (or a substantial portion thereof), and a C-half ferritin based on a human ferritin light chain has an amino acid sequence corresponding to residues 96-175 of SEQ ID NO: 1 (or a substantial portion thereof).

[0149] In some embodiments, the halves are divided at a point that corresponds to a position from about position 85 to about position 92 of SEQ ID NO:1. For example, in some embodiments, an N-half ferritin based on a human ferritin light chain has an amino acid sequence corresponding to residues 1-90 of SEQ ID NO:1, and a C-half ferritin based on a human ferritin light chain has an amino acid sequence corresponding to residues 91-175 of SEQ ID NO:1.

2. Fc Polypeptides

[0150] In certain embodiments, fragment crystallizable (Fc) polypeptides comprise Fc chains that each have one or more mutations relative to a reference Fc chain of the same Ig class. As explained further herein below, the reference Fc chain may be of, e.g., the IgG1 or IgG2 class.

[0151] Unless otherwise noted, numbering of mutations within an antibody fragment, e.g., an Fc polypeptide, throughout this disclosure is according to the EU index.

[0152] In some embodiments, the Fc polypeptide is a human IgG Fc polypeptide, that is, except for mutations noted herein, the Fc polypeptide comprises an Fc chain that is substantially similar to that of an Fc chain within a wild type human IgG.

[0153] In some embodiments, the Fc polypeptide is an IgG1 Fc polypeptide (e.g., a human IgG1 Fc polypeptide), that is, except for mutations noted herein, the Fc polypeptide comprises an Fc chain that has an amino acid sequence that is substantially similar to that of the chains within a wild type IgG1 Fc. In some embodiments, the wild type IgG1 Fc is a human IgG1 Fc, in which each Fc chain has an amino acid sequence of SEQ ID NO:5.

[0154] For example, an IgG1 Fc polypeptide may comprise an Fc chain with an amino acid sequence that is at least 85%, at least 87.5%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to that of an Fc chain within a wild-type IgG1 Fc. In some embodiments, an IgG1 Fc polypeptide comprises an Fc chain that comprises the Fc mutations specifically described for that IgG1 Fc polypeptide, but has an amino acid sequence that is otherwise 100% identical to an Fc chain within a wild type IgG1 Fc. In some embodiments, the Fc polypeptide comprises an Fc chain that has an amino acid sequence that differs by at least one, at least two, at least three, or at least four amino acid residues from the sequence of SEQ ID NO:5. In some embodiments, the Fc polypeptide comprises an Fc chain that has an amino acid sequence that differs by no more than ten, no more than nine, no more than eight, no more than seven, no more than six, no more than five, or no more than four amino acid residues from the sequence of SEQ ID NO:5.

[0155] In some embodiments, the Fc polypeptide is an IgG2 Fc polypeptide, (e.g., a human IgG2 Fc polypeptide), that is, except for mutations noted herein, the Fc polypeptide comprises an Fc chain that has an amino acid sequence that is substantially similar to that of the chains within a wild type IgG2 Fc. In some embodiments, the wild type IgG2 Fc is a human IgG2 Fc, in which each Fc chain has an amino acid sequence of SEQ ID NO:46.

[0156] For example, an IgG2 Fc polypeptide may comprise an Fc chain with an amino acid sequence that is at least 85%, at least 87.5%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% identical to that of an Fc chain within a wild-type IgG2a Fc. In some embodiments,

an IgG2 Fc polypeptide comprises an Fc chain that comprises the Fc mutations specifically described for that IgG2 Fc polypeptide, but has an amino acid sequence that is otherwise 100% identical to an Fc chain within a wild type IgG2 Fc. In some embodiments, the Fc polypeptide comprises an Fc chain that has an amino acid sequence that differs by at least one, at least two, at least three, or at least four amino acid residues from the sequence of SEQ ID NO:46. In some embodiments, the Fc polypeptide comprises an Fc chain that has an amino acid sequences that differs by no more than ten, no more than nine, no more than eight, no more than seven, no more than six, no more than five, or no more than four amino acid residues from the sequence of SEQ ID NO:46.

[0157] In some embodiments, the Fc polypeptide is a single chain Fc (scFc), which comprises two Fc chains linked together by a covalent linker, e.g., via an amino acid linker.

[0158] In some embodiments, the Fc polypeptide is an Fc monomer, e.g., a single Fc chain having only one CH2 domain (second constant Ig domain of the heavy chain) and only one CH3 domain (third constant Ig domain of the heavy chain), which single Fc chain is typically capable of dimerizing with another single Fc chain.

[0159] In some embodiments, the one or more mutations comprises a mutation or set of mutations associated with an altered characteristic as further described herein. By "associated with," it is meant that the mutation or set of mutations has been previously characterized, in the context of antibodies such as IgG antibodies, as conferring the altered characteristic (e.g., altered binding to FcRn, altered effector function, etc.) By "altered" it is meant that the characteristic (e.g., binding to an Fc receptor (e.g., FcRn)), is different than that observed without the mutation or set of mutations.

[0160] For example, in some embodiments, the altered characteristic comprises altered binding to an Fc receptor.

[0161] In some embodiments, the altered characteristic comprising altered binding to FcRn.

[0162] For example, the mutation or set of mutations associated with altered binding to FcRn may comprise a mutation at one or more residues selected from: M252, I253, S254, T256, K288, M428, N434, or combinations thereof.

[0163] In some embodiments, the altered binding to an Fc receptor comprises decreased binding to FcRn (e.g., decreased binding relative to a reference level corresponding to that level observed without the one or more mutations). For example, in some embodiments, the one or more mutations comprise a mutation or set of mutations associated with decreased binding to FcRn, e.g., I253A, I253V, K288A, or combinations thereof.

[0164] In some embodiments, the one or more mutations comprises a mutation or set of mutations associated with altered effector function, e.g., altered binding to an Fc receptor associated with effector function (e.g., Fcγ receptors such as FcγRI, FcγRII, or FcγRIIIb).

[0165] For example, the one or more mutations may comprise a mutation or set of mutations at one or more residues selected from: L234, L235, G236, G237, P329, A330, and combinations thereof.

[0166] In some embodiments, the altered binding to an Fc receptor comprises decreased effector function, e.g., LALA (L234A/L235A), LALAP (L234A/L235A/P329G), G236R, G237A, A330L, or combinations thereof.

3. Antigen-Binding Antibody Fragments

[0167] In certain embodiments, the antigen-binding antibody fragment comprises a heavy chain variable region (e.g., a V_H). In certain embodiments, the antigen-binding antibody fragment comprises a heavy chain variable domain (e.g., V_H) and a light chain variable domain (e.g., a V_L or V_K). In certain embodiments, the antigen-binding antibody fragment comprises a Fab which comprises a heavy chain variable domain (e.g., V_H) and a light chain variable domain (e.g., a V_L or V_K).

[0168] In certain embodiments, the antigen-binding antibody fragment does not comprise any domains from the Fc region, e.g., does not comprise any CH2 or CH3 domains.

[0169] In some embodiments, the antigen-binding fragment binds to an antigen on an infectious disease agent, e.g., a virus.

[0170] In some embodiments, the antigen-binding antibody fragment binds to an antigen on a target cell, e.g., a cancer cell or an immune cell.

[0171] In embodiments where multiple types of fusion polypeptides having antigen-binding antibody fragments are used, the antigen-binding antibody fragments in the various types of fusion polypeptides may be capable of binding to the same epitope, or they may be capable of binding to epitopes that are distinct and non-overlapping. In some embodiments where the epitopes are distinct and non-overlapping, the epitopes are from the same protein.

4. Linkers

[0172] In certain embodiments, linkers are used within fusion polypeptides and/or within single-chain molecules such as scFcs. In some embodiments, the linker is an amino acid linker. For example, a linker as employed herein may comprise from about 1 to about 100 amino acid residues, e.g., about 1 to about 70, about 2 to about 70, about 1 to about 30, or about 2 to about 30 amino acid residues. In some embodiments, the linker comprises at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, or at least 10 amino acid residues.

[0173] In certain embodiments, the linker comprises a glycine-serine sequence, e.g., a $(G_nS)_m$ sequence (e.g., GGS, GGGG (SEQ ID NO:48), or GGGGS (SEQ ID NO:49) sequence) that is present in at least 1, at least 2, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, or at least 14 copies within the linker.

B. Self-Assembled Polypeptide Complexes

[0174] In one aspect, provided are self-assembled polypeptide complexes comprising a plurality of fusion polypeptides as disclosed herein. Generally, provided self-assembled polypeptide complexes comprise (a) a plurality of first fusion polypeptides, each first fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises an Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, and (b) a plurality of second fusion polypeptides, each second fusion polypeptide comprising (1) an antigen-binding antibody fragment linked to (2) a nanocage monomer or subunit thereof.

[0175] In some embodiments, the nanocage monomer is a ferritin monomer, and each fusion polypeptide within the

self-assembled polypeptide complex comprises a ferritin light chain or a subunit of a ferritin light chain. In these embodiments, the self-assembled polypeptide complex does not comprise any ferritin heavy chains, subunits of ferritin heavy chains, or other ferritin components capable of binding to iron.

[0176] In some embodiments, the nanocage monomer or subunit thereof is a ferritin monomer subunit, and (a) each first fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is N-half-ferritin; or (b) each first fusion polypeptide comprises a ferritin monomer subunit which is N-half ferritin and each second fusion polypeptide comprises a ferritin monomer subunit which is C-half-ferritin.

[0177] In some embodiments, the self-assembled polypeptide complex comprises from 24 to 48 fusion polypeptides in total. In some embodiments, the self-assembled polypeptide complex comprises 24 fusion polypeptides in total. In some embodiments, the self-assembled polypeptide complex comprises more than 24 fusion polypeptides, e.g., at least 26, at least 28, at least 30, at least 32 fusion polypeptides, at least 34 fusion polypeptides, at least 36 fusion polypeptides, at least 38 fusion polypeptides, at least 40 fusion polypeptides, at least 42 fusion polypeptides, at least 44 fusion polypeptides, at least 46 fusion polypeptides, or at least 48 fusion polypeptides in total. In some embodiments, the self-assembled polypeptide complex comprises about 32 fusion polypeptides.

[0178] In some embodiments, the self-assembled polypeptide complex comprises at least 4, at least 5, least 6, at least 7, or at least 8 first fusion polypeptides.

[0179] In some embodiments, the self-assembled polypeptide complex comprises at least 4, at least 5, least 6, at least 7, or at least 8 second fusion polypeptides.

[0180] In some embodiments, the self-assembled polypeptide complex further comprises at least 4, at least 5, least 6, at least 7, at least 8, at least 9, at least 10, least 11, at least 12, at least 13, at least 14, at least 15, or at least 16 third fusion polypeptides.

[0181] In some embodiments, the self-assembled polypeptide complex comprises a ratio of approximately 1:1, 1:2, 1:3, or 1:4 of first fusion polypeptides to all other fusion polypeptides.

Pharmacokinetic Characteristics

[0182] In certain embodiments, when administered to a subject in need thereof, a provided self-assembled polypeptide complex, has one or more pharmacokinetic features similar to that of a reference IgG molecule (e.g., an IgG molecule whose class matches the class of an Fc chain within an Fc polypeptide of a first fusion polypeptide within the self-assembled polypeptide complex). In some embodiments, the ranges for the pharmacokinetic characteristics discussed herein (e.g., half-life, AUC, and/or C_{max}) are obtained when the self-assembled polypeptide complex is administered to a human subject. In some embodiments, the ranges for the pharmacokinetic characteristics discussed herein are obtained when the self-assembled polypeptide complex is administered via a systemic route, e.g., via intravenous or subcutaneous administration.

[0183] In some embodiments, a self-assembled polypeptide complex as disclosed herein has a similar half-life to that of reference IgG molecule. The reference IgG molecule

[0186] In some embodiments, a self-assembled polypeptide complex as disclosed herein has a similar bioavailability to that of reference IgG molecule. For example, in some embodiments, after administration to a subject in need thereof, the self-assembled polypeptide complex has a maximum concentration (C_{max}) of from about 10 µg/mL to about 750 mg/mL, about 25 µg/mL to about 750 mg/mL, about 50 µg/mL to about 750 mg/mL, about 75 µg/mL to about 750 mg/mL, about 100 µg/mL to about 750 mg/mL, about 250 µg/mL to about 750 mg/mL, about 500 µg/mL to about 750 mg/mL, about 750 µg/mL to about 750 mg/mL, about 1 mg/mL to about 750 mg/mL, about 10 mg/mL to about 750 mg/mL, about 25 mg/mL to about 750 mg/mL, about 50 mg/mL to about 750 mg/mL, about 75 mg/mL to about 750 mg/mL, about 100 mg/mL to about 750 mg/mL, about 250 mg/mL to about 750 mg/mL, about 500 mg/mL to about 750 mg/mL, about 10 µg/mL to about 500 mg/mL, about 25 µg/mL to about 500 mg/mL, about 50 µg/mL to about 500 mg/mL, about 75 µg/mL to about 500 mg/mL, about 100 µg/mL to about 500 mg/mL, about 250 µg/mL to about 500 mg/mL, about 500 µg/mL to about 500 mg/mL, about 750 µg/mL to about 500 mg/mL, about 1 mg/mL to about 500 mg/mL, about 10 mg/mL to about 500 mg/mL, about 25 mg/mL to about 500 mg/mL, about 50 mg/mL to about 500 mg/mL, about 75 mg/mL to about 500 mg/mL, about 100 mg/mL to about 500 mg/mL, about 250 mg/mL to about 500 mg/mL, about 10 µg/mL to about 250 mg/mL, about 25 µg/mL to about 250 mg/mL, about 50 µg/mL to about 250 mg/mL, about 75 µg/mL to about 250 mg/mL, about 100 µg/mL to about 250 mg/mL, about 250 µg/mL to about 250 mg/mL, about 500 µg/mL to about 250 mg/mL, about 750 µg/mL to about 250 mg/mL, about 1 mg/mL to about 250 mg/mL, about 10 mg/mL to about 250 mg/mL, about 25 mg/mL to about 250 mg/mL, about 50 mg/mL to about 250 mg/mL, about 75 mg/mL to about 250 mg/mL, about 100 mg/mL to about 250 mg/mL, about 10 µg/mL to about 100

μg/mL, about 25 μg/mL to about 100 mg/mL, about 50
 mg/mL to about 100 mg/mL, about 75 μg/mL to about 100
 mg/mL, about 100 μg/mL to about 100 mg/mL, about 250
 μg/mL to about 100 mg/mL, about 500 μg/mL to about 100
 mg/mL, about 750 μg/mL to about 100 mg/mL, about 1
 mg/mL to about 100 mg/mL, about 10 mg/mL to about 100
 mg/mL, about 25 mg/mL to about 100 mg/mL, about 50
 mg/mL to about 100 mg/mL, about 75 mg/mL to about 100
 mg/mL, about 10 μg/mL to about 75 mg/mL, about 25
 μg/mL to about 75 mg/mL, about 50 μg/mL to about 75
 mg/mL, about 75 μg/mL to about 75 mg/mL, about 100
 μg/mL to about 75 mg/mL, about 250 μg/mL to about 75
 mg/mL, about 500 μg/mL to about 75 mg/mL, about 750
 μg/mL to about 75 mg/mL, about 1 mg/mL to about 75
 mg/mL, about 10 mg/mL to about 75 mg/mL, about 25
 mg/mL to about 75 mg/mL, about 50 mg/mL to about 75
 mg/mL, about 10 μg/mL to about 50 mg/mL, about 25
 μg/mL to about 50 mg/mL, about 50 μg/mL to about 50
 mg/mL, about 75 μg/mL to about 50 mg/mL, about 100
 μg/mL to about 50 mg/mL, about 250 μg/mL to about 50
 mg/mL, about 500 μg/mL to about 50 mg/mL, about 750
 μg/mL to about 50 mg/mL, about 1 mg/mL to about 50
 mg/mL, about 10 mg/mL to about 50 mg/mL, about 25
 mg/mL to about 50 mg/mL, about 10 μg/mL to about 25
 mg/mL, about 25 μg/mL to about 25 mg/mL, about 50
 μg/mL to about 25 mg/mL, about 75 μg/mL to about 25
 mg/mL, about 100 μg/mL to about 25 mg/mL, about 250
 μg/mL to about 25 mg/mL, about 500 μg/mL to about 25
 mg/mL, about 750 μg/mL to about 25 mg/mL, about 1
 mg/mL to about 25 mg/mL, about 10 mg/mL to about 25
 mg/mL, about 10 μg/mL to about 10 mg/mL, about 25
 μg/mL to about 10 mg/mL, about 50 μg/mL to about 10
 mg/mL, about 75 μg/mL to about 10 mg/mL, about 100
 μg/mL to about 10 mg/mL, about 250 μg/mL to about 10
 mg/mL, about 500 μg/mL to about 10 mg/mL, about 750
 μg/mL to about 10 mg/mL, about 1 mg/mL to about 10
 mg/mL, about 10 μg/mL to about 1 mg/mL, about 25 μg/mL
 to about 1 mg/mL, about 50 μg/mL to about 1 mg/mL, about
 75 μg/mL to about 1 mg/mL, about 100 μg/mL to about 1
 mg/mL, about 250 μg/mL to about 1 mg/mL, about 500
 μg/mL to about 1 mg/mL, about 750 μg/mL to about 1
 mg/mL, about 10 μg/mL to about 750 μg/mL, about 25
 μg/mL to about 750 μg/mL, about 50 μg/mL to about 750
 μg/mL, about 75 μg/mL to about 750 μg/mL, about 100
 μg/mL to about 750 μg/mL, about 250 μg/mL to about 750
 μg/mL, about 500 μg/mL to about 750 μg/mL, about 10
 μg/mL to about 500 μg/mL, about 25 μg/mL to about 500
 μg/mL, about 50 μg/mL to about 500 μg/mL, about 75
 μg/mL to about 500 μg/mL, about 100 μg/mL to about 500
 μg/mL, about 250 μg/mL to about 500 μg/mL, about 10
 μg/mL to about 250 μg/mL, about 25 μg/mL to about 250
 μg/mL, about 50 μg/mL to about 250 μg/mL, about 75
 μg/mL to about 250 μg/mL, about 100 μg/mL to about 250
 μg/mL, about 10 μg/mL to about 100 μg/mL, about 25
 μg/mL to about 100 μg/mL, about 50 μg/mL to about 100
 μg/mL, about 75 μg/mL to about 100 μg/mL, about 10
 μg/mL to about 75 μg/mL, about 25 μg/mL to about 75
 μg/mL, about 50 μg/mL to about 75 μg/mL, about 10 μg/mL
 to about 50 μg/mL, about 25 μg/mL to about 50 μg/mL, or
 about 10 μg/mL to about 25 μg/mL. In some embodiments,
 after administration to a subject in need thereof, the self-
 assembled polypeptide complex has a maximum concentra-
 tion (C_{max}) of at least 10 μg/mL, at least 25 μg/mL, at least
 50 μg/mL, at least 100 μg/mL, at least 250 μg/mL, at least

500 µg/mL, at least 750 µg/mL, at least 1 mg/mL, at least 10 mg/mL, at least 25 mg/mL, at least 50 mg/mL, at least 75 mg/mL, at least 100 mg/mL, at least 250 mg/mL, at least 500 mg/mL, or at least 750 mg/mL when administered to a subject in need thereof.

Functional Effects

[0187] In certain embodiments, a provided self-assembled polypeptide complex is capable of antibody-dependent cellular phagocytosis (ADCP). In some embodiments, the ADCP is induced at a level of at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, or at least 60% internalization of target. Methods of measuring ADCP are known in the art and include, for example, in vitro assays that utilize a macrophage cell line and a target.

C. Methods of Treatment

[0188] In one aspect, provided are methods that may be useful for treating, ameliorating, or preventing a disease or a condition (e.g., an infectious disease, cancer, or an autoimmune disease), generally comprising a step of administering a composition comprising a self-assembled polypeptide complex of the present disclosure to a subject.

[0189] In some embodiments, the subject is a mammal, e.g., a human.

[0190] Compositions for administration to subjects generally comprise a self-assembled polypeptide complex as disclosed herein. In some embodiments, such compositions further comprise a pharmaceutically acceptable excipient.

[0191] Compositions may be formulated for administration for any of a variety of routes of administration, including systemic routes (e.g., oral, intravenous, intraperitoneal, subcutaneous, or intramuscular administration).

EXAMPLES

Example 1. Expression and Analysis of Representative Multabodies

[0192] In this Example, Multabodies (MBs) were formed by fusion proteins generated by fusing together a human ferritin light chain (hFTL, SEQ ID NO:1), N-terminal fragment (residues 1-90) of hFTL (N_hFTL, SEQ ID NO:2), or C-terminal fragment (residues 91-175) of hFTL (C_hFTL, SEQ ID NO:3) with a single-chain Fab (scFab) and/or a fragment crystallizable region (Fc) or single chain Fc-dimer (scFc), via a linker, such as a (Gly-Ser)_m peptide linker as described herein. The different formats that were generated are illustrated in FIG. 1.

[0193] T-01 MB: genes encoding fusion proteins (1) scFab of HIV-neutralizing antibody PGDM1400 fused to the N-terminus of hFTL (PGDM1400-hFTL, SEQ ID NO:27), (2) scFc fused to the N-terminus of N_hFTL (scFc-N_hFTL, SEQ ID NO:34), (3) scFab of HIV-neutralizing antibody N49P7 fused to the N-terminus of C_hFTL (N49P7-C_hFTL, SEQ ID NO:28), and (4) scFab of HIV-neutralizing antibody 10E8v4 fused to the N-terminus of C_hFTL (10E8v4-C_hFTL, SEQ ID NO:30) were prepared, mixed at a molar ratio of 4:2:1:1 and transiently transfected into HEK 293F cells for the production and formation of T-01 MB. See, FIG. 1B.

[0194] T-02 MB: genes encoding fusion proteins (1) PGDM1400-hFTL (SEQ ID NO:27), (2) scFc-N_hFTL

(SEQ ID NO:34), (3) scFab of anti-CD4 antibody ibalizumab (iMab) fused to the N-terminus of C_hFTL (iMab-C_hFTL, SEQ ID NO:31), and (4) 10E8v4-C_hFTL (SEQ ID NO:30) were prepared, mixed at a molar ratio of 4:2:1:1, and transiently transfected into HEK 293F cells for the production and formation of T-02 MB. See, FIG. 1B.

[0195] T-01 MB.v2: genes encoding fusion proteins (1) PGDM1400-hFTL (SEQ ID NO:27), (2) scFab of N49P7 fused to the N-terminus of N_hFTL (N49P7-N_hFTL, SEQ ID NO:29), and (3) an Fc monomer fused to the C-terminus of 10E8-C_hFTL (10E8v4-C_hFTL-Fc, SEQ ID NO:32) were prepared, mixed at a molar ratio of 3:1:1, and transiently transfected into HEK 293F cells for the production and formation of T-01 MB.v2. See, FIG. 1D. (As described for “T-01 MB” above, 10E8-C_hFTL contains the scFab of HIV-neutralizing antibody 10E8v4 fused to the N-terminus of C_hFTL. Thus, the construct 10E8v4-C_hFTL-Fc (SEQ ID NO:32) contains a C-half ferritin with the Fab3 at the N-terminus of the C-half ferritin and an Fc monomer at the C-terminus of the C-half ferritin.)

[0196] The Multabodies described in this Example had wild-type (WT) Fc or engineered IgG1 Fc. Such an engineered IgG1 Fc contained any one or more mutations of L234A, L235A, K288A, I253V, I253A, P329G, M428L, N434S, or any combinations thereof (according to the EU numbering scheme). For example, T-01 MB IgG1 K288A had engineered IgG1 Fc with the K288A mutation; T-01 MB IgG1 µLALAP had engineered IgG1 Fc with mutations of L234A, L235A, and P329G; T-01 MB.v2 IgG1 µLS had engineered IgG1 Fc with mutations of M428L and N434S.

[0197] The Multabodies were purified using Protein A affinity chromatography, optionally followed by Protein L affinity chromatography. Fractions containing the Multabodies were concentrated and further purified by size-exclusion chromatography (SEC) in sodium phosphate buffer. After SEC purification, negative-stain electron microscopy (EM) and/or SEC with inline multi-angle light scattering (SEC-MALS) was used to evaluate the size of the formed Multabodies.

Example 2. Binding of Multabodies Towards Fc Receptors Determined by Biolayer Interferometry

[0198] The binding kinetics and affinity to Fc receptors—human Fcγ receptor I (hFcγRI), hFcγRIIa, and hFcγRIIb—of Multabodies comprising different Fc mutations were determined by biolayer interferometry (BLI) using an Octet RED96 BLI system (Pall ForteBio).

[0199] Briefly, His-tagged Fc receptors were loaded onto Ni-NTA biosensors to reach a signal response of 0.8 nm. The association rates were measured by transferring the loaded biosensors to wells containing serial dilutions of the test Multabodies (20-10-5-2.5-1.25-0.65 nM) or IgG1 control (250-125-62.5-31.2-15.6-7.8 nM), with a contact time of 180 s. The IgG1 control is a cocktail of PGDM1400, N49P7, and 10E8v4 antibodies, all with wild-type IgG1 backbone. In order to assess the potential of the Multabodies to undergo endosomal recycling, their bindings to the hFcRn/β2-microglobulin complex were measured at both physiological pH (7.4) and acidic pH (5.6).

[0200] Fc mutations of IgG1 backbone evaluated in Multabodies include: K288A, I253V, and I253A that decrease antibody binding to FcRn; P329G, LALA (L234A, L235A), and LALAP (L234A, L235A and P329G) that decrease

antibody binding to FcγRs; and combinations thereof (Numberings are according to the EU numbering scheme.)

[0201] Representative examples for the relevant segment of the resulting sensorgrams are provided in FIGS. 3A, 3B, 3C, 3D, 3E, and 3F. The values determined for k_{on} , k_{off} , and the resulting equilibrium dissociation constant (K_D) for the Multabodies are summarized in Tables 1 and 2.

[0202] At acidic pH (5.6), T-01 MB with wild-type IgG1 Fc binds to FcRn with over 1000-fold higher affinity compared to the IgG1 control; the same is also observed for T-01 MB with K288A, I253V, P329G, LALA, LALAP, K288A+P329G, or K288A+LALAP IgG1 Fc mutations. T-01 MB with I253A or I253A+LALAP IgG1 Fc mutations have a similar binding ability to FcRn at pH 5.6 compared to the IgG1 control. At physiological pH (7.4), T-01 MB with wild-IgG1 Fc, P329G IgG1 Fc mutation, LALA IgG1 Fc mutations, or LALAP IgG1 Fc mutations show measurable binding to FcRn.

[0203] T-01 MB with wild-type IgG1 Fc binds to hFcγRI, hFcγRIIa, and hFcγRIIb with over 1000-fold higher affinity compared to the IgG1 control. Multabodies with P329G, LALA, or LALAP IgG1 Fc mutation(s) show reduced binding or non-binding to the tested Fcγ receptors.

[0204] T-01 MB.v2 with wild-type IgG1 Fc or with LS Fc mutations has a FcRn binding profile more similar to IgG1, with comparable binding ability to human FcRn at acidic pH and no binding at physiological pH. Further, T-01 MB.v2 with type IgG1 Fc or LS mutations shows reduced binding to the high affinity FcγRI and no binding to the low affinity Fcγ receptors tested, similar to T-01 MB containing the LALAP mutations.

[0205] Among the tested Fc mutations and mutation combinations, the I253A+LALAP IgG1 Fc mutation combination adjusts the Fc receptor binding profile of the Multabody (in T-01 MB format) to IgG1-like.

TABLE 1

Kinetic constants and affinities to FcRn of Multabodies determined by BLI			
Multabody	FcRn, pH 5.6		
	k_{on} [$M^{-1} \times s^{-1}$]	k_{off} [s^{-1}]	K_D [M]
IgG1 control	5.80E+05	1.07E-03	1.85E-09
T-01 MB IgG1 WT*	8.29E+05	<1.0E-07	<1.0E-12
T-01 MB IgG2 WT	9.19E+05	<1.0E-07	<1.0E-12
T-01 MB IgG4 WT	2.93E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 I253A	8.22E+05	2.01E-03	2.44E-09
T-01 MB IgG1 I253V	1.76E+06	<1.0E-07	<1.0E-12
T-01 MB IgG1 K288A	8.49E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 P329G*	9.08E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 L234A L235A (LALA)*	8.09E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALA P329G (LALAP)*	8.38E+05	<1.0E-07	<1.0E-12
T-01 MB IgG2 K288A P329G	8.71E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALAP K288A	8.42E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALAP I253A	7.39E+05	1.29E-03	1.75E-09
T-01 MB.v2 IgG1 WT	7.39E+05	1.79E-03	2.42E-09
T-01 MB.v2 IgG1 M428L N434S (LS)	2.94E+05	1.25E-03	4.23E-09

*Multabody shows residual binding to FcRn at pH 7.4

TABLE 2

Kinetic constants and affinities to Fcγ receptors of Multabodies determined by BLI									
Multabody	FcγRI			FcγRIIa			FcγRIIb		
	kon [$M^{-1} \times s^{-1}$]	koff [s^{-1}]	KD [M]	kon [$M^{-1} \times s^{-1}$]	koff [s^{-1}]	KD [M]	kon [$M^{-1} \times s^{-1}$]	koff [s^{-1}]	KD [M]
IgG1 control	4.28E+04	1.28E-04	2.98E-09	2.39E+05	3.52E-03	1.48E-08	7.21E+04	2.54E-03	3.53E-08
T-01 MB IgG1 WT	6.72E+05	<1.0E-07	<1.0E-12	9.63E+05	<1.0E-07	<1.0E-12	1.19E+06	<1.0E-07	<1.0E-12
T-01 MB IgG2 WT	7.58E+05	1.54E-03	2.03E-09	4.33E+05	2.63E-04	6.06E-10	6.37E+05	1.03E-03	1.61E-09
T-01 MB IgG4 WT	8.35E+05	<1.0E-07	<1.0E-12	1.14E+06	4.45E-04	3.91E-10	1.37E+06	3.05E-04	2.23E-10
T-01 MB IgG1 I253A	6.05E+05	<1.0E-07	<1.0E-12	7.63E+05	<1.0E-07	<1.0E-12	1.39E+06	<1.0E-07	<1.0E-12
T-01 MB IgG1 I253V	7.56E+05	<1.0E-07	<1.0E-12	1.09E+06	<1.0E-07	<1.0E-12	1.39E+06	<1.0E-07	<1.0E-12
T-01 MB IgG1 K288A	5.78E+05	<1.0E-07	<1.0E-12	1.07E+06	<1.0E-07	<1.0E-12	1.25E+06	<1.0E-07	<1.0E-12
T-01 MB IgG1 P329G	6.65E+05	2.49E-05	3.75E-11	2.22E+05	5.53E-05	2.49E-10	—	—	No binding
T-01 MB IgG1 LALA	5.58E+05	1.66E-05	2.97E-11	5.81E+05	1.21E-04	2.08E-10	6.49E+05	3.94E-04	6.07E-10

TABLE 2-continued

Kinetic constants and affinities to Fcγ receptors of Multabodies determined by BLI									
Multabody	FcγRI			FcγRIIa			FcγRIIb		
	kon [M ⁻¹ × s ⁻¹]	koff [s ⁻¹]	KD [M]	kon [M ⁻¹ × s ⁻¹]	koff [s ⁻¹]	KD [M]	kon [M ⁻¹ × s ⁻¹]	koff [s ⁻¹]	KD [M]
T-01 MB IgG1 LALAP	6.39E+05	4.37E-04	6.84E-10	—	—	No binding	—	—	No binding
T-01 MB IgG2 K288A P329G	6.94E+05	9.13E-05	1.32E-10	—	—	No binding	—	—	No binding
T-01 MB IgG1 LALAP K288A	6.01E+05	1.61E-04	2.69E-10	—	—	No binding	—	—	No binding
T-01 MB IgG1 LALAP I253A	8.53E+05	6.56E-04	7.68E-10	—	—	No binding	—	—	No binding
T-01 MB.v2 IgG1 WT	3.12E+05	<1.0E-07	<1.0E-12	—	—	No binding	—	—	No binding
T-01 MB.v2 IgG1 LS	4.31E+05	<1.0E-07	<1.0E-12	—	—	No binding	—	—	No binding

Example 3. Target Binding of Multabodies Determined by Biolayer Interferometry

[0206] The binding kinetics and affinity to the respective target of PGDM1400, N49P7, 10E8v4, or iMab, as included as Fab in the Multabodies, were determined by BLI using an Octet RED96 BLI system (Pall ForteBio).

[0207] The experiments were performed similarly as described in Example 2, except that Hi-tagged targets—BG5050 SOSIP.664 D368R, gp120 subunit 93TH057, gp41 membrane-proximal external region (MPER), and soluble CD4 for PGDM1400, N49P7, 10E8v4, and iMab, respec-

tively—were loaded onto Ni-NTA biosensors to reach a signal response of 0.8 nm. The loaded biosensors were then titrated with test Multabodies, PGDM1400, N49P7, 10E8v4, or iMab antibody (with a wild-type IgG1 backbone), or a cocktail (IgG1 control) of PGDM1400, N49P7, and 10E8v4 antibodies (all with a wild-type IgG1 backbone) at various concentrations.

[0208] Representative examples for the relevant segment of the resulting sensorgrams are provided in FIGS. 4A, 4B, 4C, 4D), and 4E. The values determined for k_{on} , k_{off} , and the resulting equilibrium dissociation constant (K_D) for the Multabodies are summarized in Tables 3 and 4.

TABLE 3

Kinetic constants and affinities to target binding of Multabodies determined by BLI									
Multabody	BG5050 SOSIP.664 D368R			93TH057			gp41 MPER		
	k_{on} [M ⁻¹ × s ⁻¹]	k_{off} [s ⁻¹]	K_D [M]	k_{on} [M ⁻¹ × s ⁻¹]	k_{off} [s ⁻¹]	K_D [M]	k_{on} [M ⁻¹ × s ⁻¹]	k_{off} [s ⁻¹]	K_D [M]
IgG1 control	1.00E+05	7.21E-04	7.20E-09	8.02E+04	1.08E-04	1.34E-09	4.16E+04	1.15E-04	2.77E-09
T-01 MB IgG1 WT	8.03E+05	<1.0E-07	<1.0E-12	3.33E+05	<1.0E-07	<1.0E-12	3.87E+05	<1.0E-07	<1.0E-12
T-01 MB IgG2 WT	9.79E+05	<1.0E-07	<1.0E-12	4.65E+05	<1.0E-07	<1.0E-12	4.85E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 I253A	7.36E+05	<1.0E-07	<1.0E-12	4.25E+05	<1.0E-07	<1.0E-12	4.21E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 K288A	1.04E+06	<1.0E-07	<1.0E-12	4.58E+05	<1.0E-07	<1.0E-12	5.00E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 P329G	9.78E+05	<1.0E-07	<1.0E-12	5.44E+05	<1.0E-07	<1.0E-12	2.42E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALA	1.18E+06	<1.0E-07	<1.0E-12	5.83E+05	<1.0E-07	<1.0E-12	1.35E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALAP	9.39E+05	<1.0E-07	<1.0E-12	4.56E+05	<1.0E-07	<1.0E-12	5.25E+05	<1.0E-07	<1.0E-12
T-01 MB IgG2 K288A P329G	1.08E+06	<1.0E-07	<1.0E-12	6.51E+05	<1.0E-07	<1.0E-12	4.36E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALAP K288A	1.25E+06	<1.0E-07	<1.0E-12	6.21E+05	<1.0E-07	<1.0E-12	3.36E+05	<1.0E-07	<1.0E-12
T-01 MB IgG1 LALAP I253A	1.11E+06	<1.0E-07	<1.0E-12	6.68E+05	<1.0E-07	<1.0E-12	2.25E+05	<1.0E-07	<1.0E-12
T-01 MB.v2 IgG1 WT	1.08E+06	<1.0E-07	<1.0E-12	6.13E+05	<1.0E-07	<1.0E-12	1.83E+05	<1.0E-07	<1.0E-12
T-01 MB.v2 IgG1 LS	9.15E+05	<1.0E-07	<1.0E-12	4.98E+05	<1.0E-07	<1.0E-12	1.80E+05	<1.0E-07	<1.0E-12

TABLE 4

Kinetic constants and affinities to target binding of Multabodies determined by BLI					
		BG5050 SOSIP.664			
		D368R	93TH057	gp41 MPER	CD4
T-01 MB	k_{on} [$M^{-1} \times s^{-1}$]	8.12E+05	3.67E+05	3.54E+05	—
	k_{off} [s^{-1}]	<1.0E-07	<1.0E-07	<1.0E-07	—
	K_D [M]	<1.0E-12	<1.0E-12	<1.0E-12	No binding
T-02 MB	k_{on} [$M^{-1} \times s^{-1}$]	8.93E+05	—	4.16E+05	5.08E+05
	k_{off} [s^{-1}]	<1.0E-07	—	<1.0E-07	<1.0E-07
	K_D [M]	<1.0E-12	No binding	<1.0E-12	<1.0E-12
PGDM1400	k_{on} [$M^{-1} \times s^{-1}$]	6.90E+03	—	—	—
	k_{off} [s^{-1}]	2.62E-04	—	—	—
	K_D [M]	3.80E-08	No binding	No binding	No binding
N49P7	k_{on} [$M^{-1} \times s^{-1}$]	—	1.53E+05	—	—
	k_{off} [s^{-1}]	—	1.11E-03	—	—
	K_D [M]	No binding	1.39E-10	No binding	No binding
10E8v4	k_{on} [$M^{-1} \times s^{-1}$]	—	—	6.51E+04	—
	k_{off} [s^{-1}]	—	—	1.90E-04	—
	K_D [M]	No binding	No binding	2.92E-09	No binding
iMab	k_{on} [$M^{-1} \times s^{-1}$]	—	—	—	8.54E+04
	k_{off} [s^{-1}]	—	—	—	<1.0E-07
	K_D [M]	No binding	No binding	No binding	<1.0E-12

Example 4. Pharmacokinetics of Multabodies in Mice

[0209] Analyses of the pharmacokinetics of Multabodies with wild-type Fc or engineered IgG1 Fc were performed in mice.

[0210] Test Multabodies or IgG1 control were injected into five CB17/Icr-Prkdc^{scid}/IcrIcoCrI immunodeficient (SCID) mice/group on day 0 at a dose of 5 mg/kg. The IgG1 control is a cocktail of PGDM1400, N49P7, and 10E8v4 antibodies, all with a wild-type IgG1 backbone. Serum samples were taken from day 1 and every two days for 9 days. On day 10, an additional dose of 5 mg/kg was administered and serum samples collected on day 11 and day 15.

[0211] In addition, Multabodies containing wild-type Fc, LALAP+I253A IgG1 Fc mutation combination, or M428L+N434S (LS) Fc mutation combination were subcutaneously injected in NOD/Shi-scid/IL-2R γ null immunodeficient (NCG) mice (3 mice/group) at a single dose of 5 mg/kg. Blood samples were collected at multiple time points following the injection. IgG1 control was tested in parallel.

[0212] Levels of circulating Multabodies was assessed by ELISA. Briefly, 96-well plates were coated with 50 μ L of His-tagged antigens recognized by the Fabs in the Multabodies at 0.5 μ g/mL. Serum/blood samples were diluted and added to the wells. Bound agents were detected using HRP-Protein A as a secondary molecule. The chemiluminescence signal was quantified using a Microplate Reader. A calibration curve with standard protein dilutions was prepared.

[0213] FIGS. 5B-5E and FIG. 6B show plots of the plasma concentration over time for tested Multabodies. LALAP+I235A, LALAP+K288A, and K288A+P329G mutation combinations were able to restore the serum level of T-01 MB to similar to the IgG1 control; LS mutation combination was able to restore the serum level of T-01 MB.v2 to similar to the IgG1 control. Accordingly, T-01 MB IgG1 μ LALAP I235A, T-01 MB IgG1 μ LALAP K288A, T-01 MB IgG1 K288A P329G, and T-01 MB.v2 IgG1 μ LS display antibody-like pharmacokinetics or favorable pharmacokinetic profiles.

Example 5. Assessment of Multabody-Induced Antibody-Dependent Cellular Phagocytosis

[0214] The potential of selected Multabodies to induce antibody-dependent cellular phagocytosis (ADCP) was assessed using THP-1 cell line.

[0215] Red fluorescent FluoSpheres NeutrAvidin microspheres were coated with biotinylated 93TH057 gp120 (antigen of N49P7) and incubated with the T-01 MB IgG1 LALAP I253A or T-01 MB.v2 IgG1 μ LS or an IgG1 cocktail of PGDM1400, N49P7, and 10E8v4 IgG1 antibodies at various concentrations for 2 h at 37° C., followed by the addition of 200 μ L THP-1 cells at 5×10^4 cells/well. After 16 h, cells were pelleted and washed with PBS. The viability of cells was determined using Live/Dead Fixable Violet Stain. Cells washed with PBS were fixed with 2% paraformaldehyde for 20 min at room temperature, pelleted, and washed once with FACS buffer (PBS+10% FBS, 0.5 mM EDTA). Cells were subsequently analyzed using a BD LSR II Flow Cytometer, and data analyzed using FlowJo. IgG1 and Multabody controls with no affinity for 93TH057 gp120 were tested in parallel; FcR binding inhibitor antibody (Invitrogen, 14-9161-73) was used to block Fc-mediated internalization.

[0216] The results are depicted in FIG. 7. The phagocytosis was quantified and represented as the percentage increase in the internalization of 93TH057-coated microspheres compared to non-coated microspheres. T-01 MB.v2 IgG1 μ LALAP I253A induces a dose-dependent ADCP, comparable to that of the IgG1 control, despite of the low binding to Fc γ RI (see Example 2).

Example 6. Assessment of Multabody-Mediated Neutralization of HIV-1

[0217] The ability of selected Multabodies to neutralize HIV-1 was assessed using the TZM-bl assay, which measures HIV-1 neutralization as a function of reductions in HIV-1 Tat-regulated firefly luciferase (Luc) reporter gene expression after a single round of infection with Env-pseudotyped viruses.

[0218] Briefly, HIV-1 pseudotyped viruses were generated by co-transfection of 293T cells with the HIV-1 subtype B

backbone NL4-3.Luc.R-E plasmid, the plasmid encoding the full-length Env clone. Test Multabodies, antibodies of the Fabs included in Multabodies, IgG1 control-1 (a cocktail of PGDM1400, N49P7, and 10E8v4 IgG1 antibodies), IgG1 control-2 (a cocktail of PGDM1400, iMab, and 10E8v4 IgG1 antibodies), or N6/PGDM1400x10 E8v4 trispecific antibody (directed to the CD4bs, V1V2 apex, and MPER binding sites) were incubated with a 10-15% tissue culture infectious dose of pseudovirus for 1 h at 37° C. prior to a 44-72 h incubation with the cells transfected with the

the IC₈₀ of T-01 MB.v2 confirmed its superior neutralization propensity over both the individual IgGs and the IgG cocktail, neutralizing 96% of all viral strains tested with a median IC₈₀ value of 0.005 µg/mL (2.2 pM) (FIG. 21C-D, FIG. 22A and Table 9). Importantly, Multabodies also blocked infection of primary peripheral blood mononuclear cells (PBMCs) with the replication-competent CXCR4-tropic HIV-1 IIIB strain (FIG. 22B), showing enhanced potency over the matched IgG mix, and without any impact on cell viability (FIG. 22C).

TABLE 5

Neutralization of HIV by Multabodies				
Multabody	14-PsV Panel		25-PsV Panel	
	Median IC ₅₀ [µg/mL]	% Virus Neutralized	Median IC ₅₀ [µg/mL]	% Virus Neutralized
PGDM1400 IgG1	0.096	71	10	44
10E8.v4 IgG1	1.8	86	1.72	88
N49P7 IgG1	0.48	79	1.24	80
iMab IgG1	0.12	93	—	—
IgG1 Control-1	0.086	100	0.56	88
IgG1 Control-2	0.10	100	—	—
N6/PGDM1400x10E8v4	0.13	100	—	—
T-01 MB (IgG1)	0.009	93	1.9	84
T-01 MB.v2 (IgG1)	0.0049	100	0.07	100
T-02 MB (IgG1)	0.008	100	—	—

pseudotyped viruses. Virus neutralization was monitored by adding Britelite plus reagent (PerkinElmer) to the cells and measuring luminescence in relative light units (RLUs) using a Synergy Neo2 Multi-Mode Assay Microplate Reader. Test articles were assayed against a single pseudovirus or a panel of 14 or 25 pseudoviruses (14- or 25-PsV panel). The 25-PsV panel included the strains in the 14-PsV panel, with the addition of 11 HIV-1 strains highly resistant to PDGM1400 in the 14-PsV panel. Thus, the 25-PsV panel contains 56% of PsV variants resistant (cutoff IC₅₀ set at 10 µg/mL) to PDGM1400 IgG neutralization.

[0219] Exemplary results are shown in FIGS. 8A, 8B, 8C, and 8D, and the values determined for IC₅₀ and breadth of neutralization are summarized in Tables 5 and 6. The Multabodies display a decrease of approximately one and two orders of magnitude in the median IC₅₀ values compared to the IC₅₀ values of the IgG cocktails and the tri-specific antibody, respectively. T-01 MB.v2, in which the neutralization profile of antibodies N49P7 and 10E8v4 were more dominant than in other multabodies, achieved 100% neutralization in the 25-PsV panel.

[0220] When tested against an extended multiclade panel of 118 PsV, T-01 MB.v2 matched the pan-neutralization breadth of the corresponding IgG cocktail (100% virus coverage, cutoff IC₅₀ set at 10 µg/mL), yet displayed a remarkable neutralization potency (FIG. 21C-D, FIG. 22A and Table 9). Specifically, the IgG cocktail and T-01 MB were only able to neutralize 9% and 8% of the PsV with an IC₅₀ value of 0.001 µg/mL, respectively, while in the case of T-01 MB.v2, 50% of the PsVs were still neutralized with an IC₅₀ value of only 0.001 µg/mL (FIG. 21C). Remarkably, Multabodies achieved a median IC₅₀ value of only 0.0009 µg/mL (0.4 pM) and hence achieved pan-neutralization 32- and 490-fold more potently in mass and molarity, respectively, compared to the IgG cocktail (FIG. 21D). In addition,

TABLE 6

Neutralization of HIV by Multabodies	
Multabody	IC ₅₀
T-01 MB IgG1	0.014 nM
T-01 MB IgG2	0.018 nM
T-01 MB IgG1 P329G	0.0094 nM
T-01 MB IgG1 LALA	0.0041 nM
T-01 MB IgG1 LALAP	0.012 nM
T-01 MB IgG1 K288A	0.014 nM
T-01 MB IgG1 I253A	0.015 nM
T-01 MB IgG1 K288A P329G	0.0042 nM
T-01 MB IgG1 LALAP K288A	0.0060 nM
T-01 MB IgG1 LALAP I253A	0.0074 nM

Example 7. Assessment of Inhibition of HIV-1 Infection by Multabody

[0221] The ability of selected Multabodies to inhibit HIV-1 infection was assessed using human peripheral blood mononuclear cells (PBMCs).

[0222] Briefly, PBMCs were obtained from three healthy blood donors and activated with phytohemagglutinin (PHA) in the presence of recombinant human IL-2 in complete RPMI medium supplemented with 10% fetal bovine serum (FBS) for 72 h prior to HIV-1 infection. The laboratory CXCR4-tropic HIV-1 isolate IIIB was incubated with testing Multabodies or an IgG1 control for 1 h at room temperature and then added to the activated PBMCs in triplicates. The IgG1 control is a cocktail of PGDM1400, N49P7, and 10E8v4 antibodies, all with an wild-type IgG1 backbone. Infected cells were cultured in the presence or absence of the testing Multabodies or antibody controls at doses ranging from 0.01-10 ug/mL. The levels of HIV-1 replication were assessed at day 7 post-infection by measuring the extracellular release of p24 Gag protein in cell-free culture super-

natants using a high-sensitivity AlphaLISA p24 detection kit on a BioTEK Synergy Plate Reader, according to the manufacturer's protocols. Cell viability was also assessed on day 7 of infection by fixing the cells in 2% PFA, and the absolute number of cells was counted by flow cytometry using a BD LSRFortessa (Becton Dickinson).

[0223] Exemplary results are shown in FIGS. 9A and 9B. The Multabodies (T-01 MB and T-01 MB.v2) were able to inhibit the infection of primary PBMCs by the replication-competent CXCR4-tropic HIV-1 isolate IIIB, with enhanced potency compared to the IgG1 control and without any impact on cell viability.

Example 8. Characterization of Thermostability of Multabody

[0224] The melting temperature (T_m) and aggregation temperature (T_{agg}) of the Multabodies and reference molecules (parental antibodies PGDM1400, N49P7, 10E8v4, and iMab; PGDM1400x10 E8v4 trispecific antibody) were determined using a UNit system. Samples were concentrated to 1.0 mg/mL and subjected to a thermal ramp from 25 to 95° C. with 1° C. increments. T_m was obtained by measuring the barycentric mean fluorescence; T_{agg} was determined as the temperature at which 50% increase in the static light scattering at a 266 nm wavelength relative to baseline was observed. The average and the standard error of 3 independent measurements were calculated using the UNit analysis software. Table 7 summarizes the T_m and T_{agg} .

[0225] Stability of Multabodies was further analyzed under accelerated conditions. Samples were concentrated to 10 mg/mL and incubated at 40° C. for four weeks. Each week, the percentage of properly folded protein was calculated based on the soluble content from SEC. Multabodies were highly stable under these conditions, with over 70% of the sample remaining soluble for 30 days. (See FIG. 10A).

[0226] Additionally, a sample before (week 0) and after (week 4) the incubation period was assessed in a PsV neutralization assay to compare the biological function of the molecules. Stability was further confirmed by only a modest loss in neutralization potency at week 4 in comparison to their potency at week 0. (See FIG. 10B).

[0227] Tested Multabodies have similar thermostability compared to the reference molecules and are stable for at least four weeks when stored at 40° C. with minimal loss in neutralization potency.

TABLE 7

Melting temperature (T_m) and aggregation temperature (T_{agg}) of Multabodies		
Multabody	T_m [° C.]	T_{agg} [° C.]
PGDM1400 IgG1	53	67
10E8.v4 IgG1	69	72
N49P7 IgG1	65	81
iMab IgG1	68	76
T-01 MB IgG1 WT	63	68
T-02 MB IgG1 WT	66	69
T-01 MB.v2 IgG1 WT	65	68

Example 9. Engineering Pan-HIV-1 Neutralization Potency Through Multi-Specific Antibody Avidity

Abstract

[0228] Deep-mining of B-cell repertoires of HIV-1 infected individuals has resulted in the isolation of dozens of HIV-1 broadly neutralizing antibodies (bNAbs). Yet, it remains uncertain whether any such bNAbs alone are sufficiently broad and potent to deploy therapeutically. Here, we engineered HIV-1 bNAbs for their combination on a single multi-specific and avid molecule via direct genetic fusion of their Fab fragments to the human apoferritin light chain. The resulting molecule demonstrated a remarkable median IC_{50} value of 0.0009 μ g/mL and 100% neutralization coverage of a broad HIV-1 pseudovirus panel (118-isolates) at a 4 μ g/mL cut-off—a 32-fold enhancement in viral neutralization potency compared to a cocktail of the corresponding HIV-1 bNAbs. Importantly, Fc incorporation on the molecule and engineering to modulate Fc receptor binding resulted in IgG-like bioavailability in vivo. This robust plug-and-play antibody design is relevant against indications where multi-specificity and avidity are leveraged simultaneously to mediate optimal biological activity.

[0229] The high genetic diversity of HIV-1 continues to be a major barrier to the development of therapeutics for prevention and treatment. Here, we describe the design of an antibody platform that allows assembly of a highly avid, multi-specific molecule that targets simultaneously the most conserved epitopes on the HIV-1 Envelope glycoprotein. The combined multi-valency and multi-specificity translates into extraordinary neutralization potency and pan-neutralization of HIV-1 strains, surpassing that of the most potent anti-HIV broadly neutralizing antibody cocktails.

INTRODUCTION

[0230] Despite decades of research, no effective vaccine or cure exists for the human immunodeficiency virus type I (HIV-1). However, the fact that a small proportion of HIV-1 infected individuals develop antibodies with exceptional neutralization potency across circulating HIV-1 isolates highlights the potential for antibody-mediated control of HIV-1. Since the first-generation of broadly neutralizing antibodies (bNAbs) 2F5 (1), 4E10 (2, 3), 2G12 (4) and b12 (5, 6) were isolated, the number of bNAbs has dramatically increased due to implementation of new technologies such as Env-specific single B cell sorting (7-9), antibody cloning and high-throughput neutralization assays (10-13), and, more recently, proteomic deconvolution (14). Several HIV-1 bNAbs have now been described that primarily target six conserved sites on the trimeric HIV Envelope glycoprotein (Env), including the V1/V2 loops at the trimer apex, V3 loop glycans, the CD4 binding site (CD4bs), the gp120-gp41 interface, the Env silent face and the membrane-proximal external region (MPER) (7, 9, 11-20).

[0231] Interest in bNAbs as therapeutic agents in the fight against HIV-1 arise from the potent antiviral activity observed in challenge studies in macaques (21-25) and humanized mice (26-29), and from the reduced viremia achieved in infected humans following infusion of bNAbs (30-34). In addition, antibodies possess key advantages in comparison to oral antiretroviral therapy (ART): they have longer circulating half-lives, and can form immune complexes that enhance host immunity to the virus. These

observations have led to the clinical evaluation of antibody-based therapies to confer protection against HIV-1 acquisition through passive administration of bNAbs (35), and efforts to control and/or clear HIV-1 in infected individuals (31-33).

[0232] The recent Antibody Mediated Prevention (AMP) trials explored the ability of bNAb VRC01 to confer passive immunity against HIV-1. In these studies, antibody breadth and potency inferred from TZM-bl neutralization assays were proposed as effective predictors of antibody efficacy in humans. Specifically, an IC_{80} value below 1 $\mu\text{g}/\text{mL}$ was established as the potency threshold that a biotherapeutic needs to achieve in order to confer protection against a specific HIV-1 strain (35). VRC01 only met that threshold against 30% of HIV-1 strains in the trials and hence, failed to confer broad protection, highlighting a critical need for more potent and broadly acting molecules. While such breadth of coverage could be achieved by administration of multiple bNAbs, despite recent IgG engineering efforts (36-40), potency may still limit the therapeutic efficacy of antibody cocktails.

[0233] Here, we overcome the immense sequence diversity of HIV-1 with extraordinary neutralization potency by engineering the human apoferritin subunit to drive multimerization of three different HIV-1 bNAbs on a single molecule. The resulting MULTi-specific, multi-Affinity anti-BODY (Multabody) was able to achieve pan-neutralization (100% virus coverage) with a median IC_{50} value of 0.0009 $\mu\text{g}/\text{mL}$ (0.4 pM). The Multabody design described herein represents a robust and powerful plug-and-play platform to multimerize antibodies in order to enhance their neutralization of HIV-1 across the broadest range of isolates.

Materials and Methods

[0234] Expression and purification of Fab-only apoferritin multimers. Genes encoding the light chain of human apoferritin and the scFab-human apoferritin fusions were synthesized and cloned by GeneArt (Life Technologies) into the pHLsec expression vector. 200 mL of HEK 293F cells (Thermo Fisher Scientific) were seeded at a density of 0.8×10^6 cells/mL in Freestyle expression media and incubated with 125 rpm oscillation at 37° C., 8% CO_2 , and 70% humidity in a Multitron Pro shaker (Infors HT). Within 24 h of seeding, cells were transiently transfected using 50 μg of filtered DNA preincubated for 10 min at room temperature (RT) with the transfection reagent FectoPRO (Polyplus Transfections) at a 1:1 ratio. Plasmids encoding scFab-human apoferritin and human apoferritin were mixed at a ratio of 1:4, 1:1, 4:1 and 1:0. After 6-7 days, cell suspensions were harvested by centrifugation at $5000 \times g$ for 15 min and the supernatants filtered through a 0.22 μm Steritop filter (EMD Millipore). The particles were purified by affinity chromatography to the Fab and eluted after a wash. Fractions containing protein were pooled, concentrated and loaded onto a Superose 6 10/300 GL size exclusion column (GE Healthcare) in 20 mM sodium phosphate, pH 8.0, 150 mM NaCl.

[0235] Design, expression and purification of Multabodies. Genes encoding scFab and scFc fragments linked to half ferritin were generated by deletion of residues 1 to 90 (C-Ferritin) and 91 to 175 (N-Ferritin) of the light chain of human apoferritin. Furthermore, protein L binding specificity for iMab-C-Ferritin was disrupted by site-directed mutagenesis of alanine 12 of the antibody light chain to a proline

residue (69). Transient transfection of T-01 MB in HEK 293F cells was obtained by mixing 66 μg of plasmids PGDM1400 scFab-human apoferritin: scFc-N-Ferritin: N49P7 scFab-C-Ferritin: 10E8v4 scFab-C-Ferritin in a 4:2:1:1 ratio. In the case of the T-02 MB, plasmid N49P7 scFab-C-Ferritin was substituted by iMab scFab-C-Ferritin. In the case of the T-01 MB.v2, 63 μg of plasmids PGDM1400 scFab-human apoferritin: N49P7 scFab-N-Ferritin: 10E8v4 scFab-C-Ferritin-Fc in a 3:1:1 ratio were used. The DNA mixture was filtered and incubated at RT with 60 μL of FectoPRO before adding to the cell culture. Based on the necessary hetero-oligomerization to drive self-assembly, purification of Multabodies with the four components was achieved by a two-step affinity purification: protein A HP column (GE Healthcare) with 20 mM Tris pH 8.0, 3 M MgCl_2 and 10% glycerol elution buffer (Fc binding) and protein L (GE Healthcare) (PGDM1400 binding, as 10E8 and N49P7 do not bind to Protein L, and iMab-protein L binding was disrupted by the A12P mutation). A buffer exchange step was performed between both affinity chromatography steps using a PD-10 desalting column (GE Healthcare). Fractions containing the protein were concentrated and further purified by gel filtration on a Superose 6 10/300 GL column (GE Healthcare) in 20 mM sodium phosphate pH 8.0, 150 mM NaCl.

[0236] Negative-stain electron microscopy. 3 μL of Multabody at a concentration of approximately 0.02 mg/mL was added to a carbon-coated copper grid for 30 s and stained with 3 μL of 2% uranyl formate. Staining excess was immediately removed from the grid using Whatman No. 1 filter paper and an additional 3 μL of 2% uranyl formate was added for 20 s. Grids were imaged using a field-emission FEI Tecnai F20 electron microscope operating at 200 kV and equipped with an Orius charge-coupled device (CCD) camera (Gatan Inc).

[0237] Biolayer interferometry. Binding kinetics measurements were conducted using an Octet RED96 BLI system (Pall ForteBio) in PBS pH 7.4, 0.01% BSA and 0.002% Tween. A unique His-tagged ligand for each of the Multabody components and Fc receptors was selected and loaded onto Ni-NTA biosensors to reach a signal response of 0.8 nm. Association rates were measured by transferring the loaded biosensors to wells containing serial dilutions of the Multabodies (10-5-2.5-1.25-0.65-0.32 nM) or IgGs (500-250-125-62.5-31.2-15.6 nM). Dissociation rates were measured by dipping the biosensors into buffer-containing wells. The duration of each of these two steps was 180 s. To achieve selective binding to PGDM1400, a D368R mutation in the CD4bs of the BG5050 SOSIP.664 trimer was introduced and, consequently, binding of N49P7 to this antigen was disrupted. Similarly, the gp120 subunit 93TH057, soluble CD4 and hFcRn in complex with P2-microglobulin were produced as the ligands for N49P7, iMab and Fc, respectively. Binding to 10E8 was tested using a His-tagged MPER peptide

(SEQ ID NO: 47)

(HHHHHHNEQELLELDKWASLWNWFNTNWLWYIKKKK,

purchased from GenScript). Recombinantly expressed hFcγRI and hFcγRIIa were used to measure binding affinities of the IgGs and Multabodies with effector function silencing mutations. Ni-NTA purification followed by size exclusion chromatography in 20 mM phosphate, pH 8.0, 150

mM NaCl buffer was used for purification of BG5050 SOSIP.664 D368R, CD4, 93TH057, hFcRn, hFcγRI and hFcγRIIa.

[0238] Size-exclusion chromatography in-line with multi-angle light scattering (SEC-MALS). A MiniDAWN TREOS and an Optilab T-rEX refractometer (Wyatt) were used in-line with an Agilent Technologies 1260 infinity II HPLC. 50 μg of 24-mer PGDM1400 scFab-ferritin fusion, T-01 MB and T-02 MB were loaded onto a Superose 6 10/300 (GE Healthcare) column in 20 mM sodium phosphate, pH 8.0, 150 mM NaCl. Data collection and analysis were performed using the ASTRA software (Wyatt).

[0239] Stability measurements. The melting temperature (T_m) and aggregation temperature (T_{agg}) of Multabodies, parental IgGs, the 12-mer homo-oligomeric Fabs and Fc and the N6/PGDM1400x10 E8v4 tri-specific antibody were determined using a UNit system (Unchained Labs). T_m was obtained by measuring the barycentric mean (BCM) fluorescence, while T_{agg} was determined as the temperature at which 50% increase in the static light scattering at a 266 nm wavelength relative to baseline was observed. Samples were concentrated to 1.0 mg/mL and subjected to a thermal ramp from 25 to 95° C. with 1° C. increments. The average and the standard error of three independent measurements were calculated using the UNit analysis software.

[0240] Stability was further analyzed under accelerated stress conditions. Multabodies diluted in 20 mM sodium phosphate pH 8.0, 150 mM NaCl, were concentrated to 10 mg/mL and incubated at 40° C. for four weeks. Each week, the percentage of properly folded protein was calculated based on the soluble content from SEC. A sample before (week 0) and after (week 4) the incubation period was assessed in a PsV neutralization assay to compare the functional activity of the molecules.

[0241] Virus production and TZM-bl neutralization assays. A panel of 14 HIV-1 pseudotyped viruses was generated by co-transfection of 293T cells with the HIV-1 subtype B backbone NL4-3.Luc.R⁺E plasmid (AIDS Research and Reference Reagent Program (ARRRP)) and the plasmid encoding the full-length Env clone, as previously described (45). HIV isolates X2088.c09, ZM106.9 and 3817.v2.c59 were kindly provided by the Collaboration for AIDS Vaccine Discovery (CAVD), and pCNE8, 1632_S2_B10, THR04156.18, 278-50, ZM197M.PB7, SF162, t257-31, Du422.1 and BG505 from NIH ARRRP. Mutation T332N in the BG505 Env expression vector was introduced by site-directed mutagenesis using the KOD-Plus mutagenesis kit (Toyobo, Osaka, Japan). The extended 25 HIV-1 pseudotyped panel was generated by adding HIV isolates p1054.TC4.1499, 6535, ZM214M.PL15, AC10.29, p16845, P6244_13.B5.4576, pM246F_C1G, TRJ04551, QH0692 and pCAAN5342 obtained from NIH ARRRP. Neutralization was determined in a single-cycle neutralization assay using the standard TZM-bl neutralization assay (45). Briefly, IgGs and Multabodies were incubated with a 10-15% tissue culture infectious dose of pseudovirus for 1 h at 37° C. prior to a 44-72 h incubation with TZM-bl cells. Virus neutralization was monitored by adding Britelite plus reagent (PerkinElmer) to the cells and measuring luminescence in relative light units (RLUs) using a Synergy Neo2 Multi-Mode Assay Microplate Reader (Biotek Instruments). HIV-1 Env pseudoviruses in the extended multiclade panel of 118 PsVs were generated by transfection in 293T cells of Env expression plasmids with full-length, Env-defective HIV

genome SG3dEnv. HIV-1 pseudoviruses were incubated with Multabodies (primary concentration of 10 μg/mL and titrated 6-fold seven times) for 1 h at 37° C. before TZM-bl cells were added. Luciferase expression was quantified 48 h after infection upon cell lysis and the addition of luciferin substrate (Promega). For the neutralization assays done with parental IgGs, historical data at the Center for Virology and Vaccine Research, Harvard Medical School was used (primary concentration of 50 μg/mL and titrated 5-fold seven times). A cutoff limit of 10 g/mL was used to determine antibody breadth.

[0242] Antibody-dependent phagocytosis. 5 μL of red fluorescent Neutravidin microspheres (Invitrogen, F8775) were washed twice with PBS+0.1% BSA and incubated with 10 μg of biotinylated 93TH057 antigen. Biotinylation was performed using the EZ-link Sulfo-NHS biotinylation kit (Thermo Scientific, 2143) following the manufacturer instructions. The final volume was brought to 200 μL with PBS/0.1% BSA and incubated overnight with rotation at 4° C. Beads were washed twice before use to remove unbound protein, and resuspended in 200 μL per 5 μL of unlabeled bead volume.

[0243] Immune complexes were formed by incubating 93TH057-coated fluorescent beads (10 μL per sample) with 10 μL of 1, 5 and 10 μg of Multabody or antibody preparations for 2 h at 37° C. THP-1 cells (ATCC TIB-202) were maintained at fewer than 5×10⁵ cells/mL in RPMI+10% FBS (Wisent); and added to immune complexes at a concentration of 5×10⁴ cells/well (in 200 μL) before incubation for 16 h at 37° C., 5% CO₂. After incubation, cells were pelleted and washed with PBS before staining with Live Dead Fixable Violet stain (Invitrogen, L34995) according to the manufacturer's protocol. Cells were washed with PBS and fixed with 2% paraformaldehyde for 20 min at RT. Fixed cells were pelleted and washed once with FACS buffer (PBS+10% FBS, 0.5 mM EDTA) and analyzed on a LSRII Flow Cytometer (BD Biosciences). Data was analyzed in FlowJo (BD Biosciences, Ashland, OR), and phagocytosis was quantified as a percentage increase in phagocytosis compared to 93TH057-coated beads in the absence of antibody. Anti-human FcR binding inhibitor antibody (Invitrogen, 14-9161-73) was added to the indicated samples at the recommended concentration as an additional control.

[0244] PBMC infection. Peripheral blood mononuclear cells (PBMCs) were obtained from three healthy blood donors, with all donors providing written informed consent. The study was approved by the University of Toronto's Research Ethics Board (protocol #00037384). Blood was collected in heparinized vacutainers (BD Biosciences) and PBMCs were subsequently isolated using density centrifugation with Lymphoprep (StemCell Technologies, Cat #07861). PBMCs were activated with phytohemagglutinin (PHA; Gibco) in the presence of recombinant human IL-2 (50 U/mL) in complete RPMI medium (Wisent), containing 10% fetal bovine serum (FBS, Wisent), streptomycin at 100 μg/mL and penicillin at 100 U/mL for 72 h prior to HIV-1 infection. After three days of activation, HIV-1 infection of cells was performed by addition of the CXCR4-tropic laboratory isolate IIIB (150 pg of p24 Gag antigen per well) to triplicate cultures of activated PBMCs in round-bottom 96-well plates seeded with 2×10⁵ cells per well in RPMI+10% FBS+25 U/mL of IL-2. Before overlaying the cells with virus, Multabodies (T-01 MB and T-01 MB.v2) or IgG cocktail were pre-incubated with virus for 1 h at RT. Infected

cells were cultured in the presence/absence of Multabody or antibody controls at doses ranging from 0.01-10 $\mu\text{g/mL}$, as indicated. The levels of HIV-1 replication were assessed by measuring the extracellular release of p24 Gag protein in cell-free culture supernatants tested at day 7 post-infection using a high-sensitivity AlphaLISA p24 detection kit (PerkinElmer, Waltham, MA) on a BioTEK Synergy Plate Reader, according to the manufacturer's protocols.

[0245] Cell viability and flow cytometry. On day 7 of infection, cells were fixed in 2% PFA and harvested for viability testing via absolute counting by flow cytometry performed using a BD LSRFortessa (Becton Dickinson). Cell viability was determined by comparison of the total live-gated cell counts in Multabody- or antibody-treated wells to the number of cells recovered from untreated control wells. Cell viability data was analyzed using FACS-Diva.

[0246] Pharmacokinetic studies. In vivo studies were performed using three 6-week-old female NOD/Shi-scid/IL-2R γ null (NCG strain code 572, Charles River Laboratories) immunodeficient mice per group. Mice were hosted by groups of 4/6 individuals. Each mouse was uniquely identified. Animals were housed in a ventilated cage (type II (16 $\times$ 19 $\times$ 35 cm, floor area=500 cm²)) under the following controlled conditions: 22 $^{\circ}$ C., 55% humidity and 12:12-hour light dark cycle 7 am: 7 pm. This study was reviewed and approved by the local ethic committee (CELEAG). T-01 MB composed of the scFab of antibodies PGDM1400, N49P7 and 10E8v4 and scFc fragments of IgG1 Fc containing i) no mutations and ii) the effector function silencing mutations L234A, L235A and P329G (LALAP) and the I253A mutation were used in the study. In addition, T-01 MB.v2 composed of the same antibody specificities with i) no Fc mutations and ii) with the half-life extension mutations (M428L/N434S) in the IgG1 Fc was included. Mice received a single subcutaneous injection of 5 mg/kg of Multabodies or the control samples (an IgG mixture matching the Fab specificity of the Multabody) in 200 μL of PBS (pH 7.5). Blood samples were collected at multiple time points and serum samples were assessed for levels of circulating antibodies by ELISA. Briefly, 96-well Pierce Nickel Coated Plates (Thermo Fisher) were coated with 50 μL at 0.5 $\mu\text{g/mL}$ of each of the His₆-tagged antigens recognized by the MB: BG5050 D368R SOSIP.664 trimer, gp120 subunit 93TH057 and MPER peptide, to determine circulating sample concentrations using reagent-specific standard curves for IgGs and Multabodies. HRP-Protein A (Invitrogen) was used as a secondary molecule and the chemiluminescence signal was quantified using the Epoch 2 Microplate spectrophotometer with the software Biotek Gen5 3.03.

Results

Potency of HIV-1 bNAbs can be Enhanced with Avidity

[0247] Apoferritin is a spherical nanocage of approximately 6 nm hydrodynamic radius formed by the self-oligomerization of 24 identical subunits (FIG. 11A). To investigate the impact of multi-valency on neutralization potency, we used the self-assembly properties of the light chain of human apoferritin to multimerize fragments of antigen binding (Fabs) derived from the most potent and broad HIV-1 bNAbs, which target different HIV-1 Env epitopes. Apoferritin subunits were genetically fused to single-chain Fabs (scFabs). scFabs were generated using flexible linkers between the light and heavy chains to ensure

correct Fab heterodimerization. Apoferritin self-assembly drove multimerization of the scFab and displayed the antibody fragments at the nanocage periphery (FIG. 11B). Different densities of multimerized Fabs were achieved by co-transfection of scFab-human apoferritin-encoding plasmids together with different ratios of non-genetically modified human apoferritin (FIG. 11C, FIG. 12). The ability of the scFab-apoferritin fusions to block HIV-1 infection were compared to the corresponding IgGs using a small HIV-1 pseudovirus (PsV) panel (FIG. 11D). Strikingly, PGDM1400, one of the most potent anti-HIV bNAb described to date, showed 10- to 40-fold higher neutralization potency when multimerized via the light chain of apoferritin compared to its conventional IgG format. bNAb 10-1074 also showed a considerable improvement in neutralization potency (4- to 40-fold), whereas bNAbs 10E8, N49P7, and VRC01 showed no effect or more modest enhancements.

Multabodies Potently and Broadly Neutralize HIV-1

[0248] In view of these results, we sought to increase the coverage of PGDM1400 using our previously described Multabody platform based on an apoferritin split design (41). The strategy consists on the separation of the four-helix apoferritin subunit into two halves (N-ferritin and C-ferritin) and their N-terminal fusion to scFabs of different specificities (FIG. 13A). This approach allows inclusion of a higher number of Fabs on the surface of the nanocage resulting in a final molecule with higher avidity. In addition, the design allows the efficient combination of three different antibody specificities as well as a fragment crystallizable (Fc) to endow the molecule with IgG-like properties, such as ease of purification leveraging Protein A affinity (FIG. 14). Specifically, we combined scFab PGDM1400 with scFabs of the near-pan neutralizing antibodies 10E8v4 (a modified 10E8 with improved solubility (42)) and N49P7, and the single-chain construct of the Fc (scFc) of human IgG1 isotype (FIG. 13A). To explore whether a Multabody could also be designed that cross-targets the HIV-1 Env and its primary receptor, CD4, we replaced N49P7 with Ibalizumab (iMab), a CD4-directed post-attachment inhibitor that has been shown to effectively inhibit HIV-1 entry (43, 44) (FIG. 15A). The resulting tri-specific Multabodies, termed T-01 MB and T-02 MB, respectively, formed highly-decorated and homogeneous particles of around 2.4 MDa (FIG. 13B-C, FIG. 15B-C) with similar thermostability as the corresponding IgGs (FIG. 16). Epitope engagement by the tri-specific Multabodies was assessed in binding kinetics experiments using epitope-specific molecules: BG505 SOSIP D368R (PGDM1400), 93TH057 gp120/CD4 (N49P7/iMab), and a MPER peptide (10E8v4) (FIG. 17). Binding to the three epitope-specific antigens with high apparent binding affinities and no detectable dissociation confirms the presence of the three antibody specificities in the Multabodies (FIG. 13d, FIG. 15d).

[0249] Neutralization potency and breadth of the Multabodies was first assessed against a panel of 14 PsVs in a standardized in vitro TZM-bl neutralization assay (45). The 14-PsV panel was designed to include low-sensitivity PsVs with at least one PsV resistant to each bNAb being evaluated (cutoff IC₅₀ set at 10 $\mu\text{g/mL}$). The IC₅₀ value and breadth of the Multabodies were compared to (i) each individual IgG, (ii) an IgG cocktail that contains the same relative amount of each IgG present in the Multabody and (iii) the

N6/PGDM1400×10E8v4 tri-specific antibody (46). T-01 MB and T-02 MB displayed 93% and 100% breadth (cutoff IC₅₀ set at 10 µg/mL) against this panel with a median IC₅₀ value of 0.009 µg/mL (3.9 pM) and 0.008 µg/mL (3.5 pM), respectively (FIG. 13E, FIG. 15E and Table 8). As such, there was a decrease of approximately one and two orders of magnitude in the median IC₅₀ values when calculated in g/mL and nM for the Multabodies compared to the IC₅₀ values of the IgG cocktails and the tri-specific antibody, respectively. Inspection of individual IC₅₀ values revealed that PsVs that are resistant to PGDM1400 IgG neutralization were also less sensitive to the Multabodies (FIG. 13F, FIG. 15E and Table 8). These data suggested that the neutralization property of the Multabodies is heavily dependent on one out of the three antibody specificities within the particle, in this case PGDM1400.

MB.v2 contains a higher number of Fabs per Fc homodimer. To attain this, a monomeric Fc fragment (i.e. one Fc chain) and a scFab are positioned at the C terminus and the N terminus of the C-ferritin half, respectively (FIG. 18A bottom, FIG. 19A). As a result, dimerization of a functional Fc homodimer drives assembly of the MB.v2 particle together with split ferritin complementation and ferritin subunit oligomerization (FIG. 18A, FIG. 19B). Importantly, homodimerization to form one functional Fc ensures assembly of four Fabs different from PGDM1400 (i.e. two Fab2 and two Fab3), thus favoring a more balanced avidity for each of the three Fabs in the fully-assembled MB.v2.

[0251] The optimized Multobody design was tested in the T-01 background (PGDM1400, N49P7, 10E8v4) that targets three epitopes on HIV-1 Env. The resulting Multobody (T-01 MB.v2) assembled into well-formed spherical particles with

TABLE 8

IC ₅₀ of individual and IgG mixtures, PGDM1400/N6x10E8v4 tri-specific and HIV-1 Multabodies against a 14-PsV panel and 25-PsV panel (additional 11 HIV-1 strains highly resistant to PGDM1400).										
Virus ID	Neutralization IC ₅₀ µg/mL						Neutralization IC ₅₀ µg/mL			
	PGDM1400 IgG	10E8v4 IgG	N49P7 IgG	IgG mix (T-01)	T-01 MB	T-01 MB.v2	Tri-specific	iMab IgG	IgG mix (T-02)	T-02 MB
CRF02_AG (Clone 257)	0.012	1.5	1.2	0.021	0.0037	0.0016	0.028	0.095	0.0078	0.0015
CRF02_AG (Clone 278)	0.16	1.8	>10	0.16	0.0090	0.0016	1.2	0.67	0.14	0.0081
X1632	0.018	2.0	>10	0.042	0.0045	0.0029	0.030	>10	0.036	0.0080
ZM106.9	0.013	>10	0.18	0.016	0.0089	0.0011	0.017	0.20	0.014	0.0031
JRC5F	0.0074	1.8	0.13	0.008	0.0054	0.0057	0.026	0.064	0.014	0.0040
pTHRO4156	0.092	0.37	7.5	0.13	0.058	0.011	0.31	0.14	0.17	0.033
3817.V2	>10	3.10	3.3	7.2	>10	0.070	1.2	0.18	0.31	1.8
PVO, clone 4	1.26	6.30	0.28	0.85	0.33	0.022	1.3	0.11	0.40	0.12
X2088	>10	>10	0.38	1.5	5.2	0.11	0.56	0.13	0.66	3.1
Du422	>10	0.55	>10	0.25	2.5	0.29	0.19	0.079	0.31	0.70
ZM197M.PB7	0.10	0.19	0.58	0.031	0.0096	0.0049	0.073	0.042	0.062	0.0044
pCNE8	0.0027	0.96	0.28	0.012	0.0032	0.0011	0.024	0.062	0.015	0.0013
BG505 T332N	0.0032	1.36	0.21	0.0088	0.0045	0.0016	0.059	0.12	0.013	0.0026
SF162	>10	3.03	0.15	0.56	1.3	0.39	0.33	7.5	1.4	2.5
Median (14-PsV panel)	0.096	1.8	0.48	0.086	0.009	0.0049	0.13	0.12	0.10	0.008
% Virus Neutralized	71	86	79	100	93	100	100	93	100	100
p1054.TC4.1499	>10	0.48	2.9	5.7	6.4	0.391	µg/mL			
6535, clone 3	>10	0.25	0.85	5.4	1.9	0.06				
ZM214M.PL15	>10	3.26	3.7	>10	3.3	0.552				
AC10.29	3.05	0.8	>10	0.15	0.0069	0.0032				
p16845	>10	0.29	6.65	0.32	3.5	0.15				
p6244_13.B5.4576	>10	0.28	0.76	0.79	5.7	0.55				
pZM246F_C1G	>10	1.72	0.3	0.56	8.7	0.39				
TRJO4551	>10	1.77	0.36	1.2	7.7	0.32				
QH0692	>10	1.25	5.6	6	>10	1.68				
NL4.3	>10	>10	>10	>10	>10	1.8				
pCAAN5342.A2	>10	3.31	8.5	>10	>10	0.74				
Median (25-PsV panel)	10	1.72	1.24	0.56	1.9	0.07				
% Virus Neutralized	44	88	80	88	84	100				

Engineering the Apoferritin Scaffold

[0250] To further improve the neutralization properties of the Multobody, we introduced some modifications to its design and made a second-generation version (MB.v2). In the original MB, the scFc is located at the N terminus of the N-ferritin half, and only one Fab, either Fab2 or Fab3, is incorporated in the Multobody per each functional Fc homodimer (FIG. 18A, top). In comparison, the optimized

no significant differences in morphology compared to the previously characterized T-01 MB (FIG. 18B). Antigen binding to BG505 SOSIP D368R, 93TH057 gp120, and MPER peptide confirmed correct folding of the three Fab specificities in T-01 MB.v2 (FIG. 18C). In addition, the new Multobody version preserves the same high thermal stability reported for T-01 MB, with a T_{agg} value of 67° C. (FIG. 18D). Multabodies were concentrated to 10 mg/mL and subjected to an accelerated stability test by incubating them

at 40° C. for four weeks. Assessment of the amount of soluble protein over time revealed that the Multabodies were highly stable under these conditions, with over 70% of the sample remaining soluble for 30 days. Stability was further confirmed by only a modest loss in neutralization potency observed for the Multabodies at week 4 in comparison to their potency at week 0 (FIG. 18E).

Pharmacokinetics of Multabodies is Similar to Corresponding IgGs

[0252] The antibody Fc domain has the capacity to interact with a variety of receptors, including Fc gamma receptors (FcγR's) and the neonatal Fc receptor (FcRn) conferring effector functions and in vivo half-life, respectively. However, Fc avidity can negatively impact the circulation time of molecules with multiple Fc fragments (41, 47). Indeed, T-01 MB showed strong binding to Fc receptors including to human FcRn at physiological pH (FIG. 20A-B), and high and low affinity FcγR's (FIG. 20C). Hence, we introduced the unique combination of LALAP (L234A, L235A and P329G) and I253A mutations in the Fc of T-01 MB to decrease binding to FcγR and FcRn, respectively, and achieve comparable binding observed for an IgG1 molecule (FIG. 20A-C). T-01 MB.v2 showed a more similar binding profile to IgG1, with comparable binding to human FcRn at acidic pH and no binding at physiological pH, even in the case of the half-life extension mutations LS (M428L/N434S) (FIG. 20A-B). Binding of T-01 MB.v2 to FcγR's yielded a low binding profile similar to that obtained with the LALAP FcR-silencing mutations in T-01 MB (FIG. 20C). The different binding patterns observed for the two MB versions are likely due to the different arrangement of the Fc fragments within the molecules (FIG. 18A, FIG. 19A). Despite low FcγRI binding, phagocytosis experiments using antigen-coated beads showed that both Multabody formats induced Fc-dependent internalization in THP-1 cells at levels similar to those achieved with the corresponding IgG mixture (FIG. 20D).

[0253] Next, we examined the in vivo bioavailability of both Multabody formats with and without engineered-Fc. A single dose of 5 mg/kg was administered subcutaneously in NOD/Shi-scid/IL-2R^γ null (NCG) immunodeficient mice and the amount of each molecule in the sera was measured every two days for 15 consecutive days. As expected from the in vitro characterization, only Fc-engineered Multabod-

ies that had an IgG-like binding profile showed days of in vivo exposure with a similar rate of decay as the parental IgG cocktail (FIG. 20E). Multabody administration was well tolerated with no decrease in body weight (FIG. 20F) or visible adverse effects.

Extraordinary Potency and Pan-Neutralization Breadth Achieved by MB.v2

[0254] We assessed the neutralization profile of T-01 MB.v2 against a PsV panel generated through addition of 11 HIV-1 strains highly resistant to PGDM1400 to our previous panel. The resulting 25-PsV panel contains 56% of PsV variants resistant (cutoff IC₅₀ set at 10 μg/mL) to PGDM1400 IgG neutralization (FIG. 21A-B). As expected, breadth and potency of the T-01 MB was greatly affected in the presence of PGDM1400 resistant PsVs (FIG. 21A-B, Table 8). However, as engineered, the neutralization profile of antibodies N49P7 and 10E8v4 were more dominant in T-01 MB.v2 allowing this optimized Multabody to achieve pan-neutralization while preserving the enhanced neutralization potency previously observed for this type of molecule (FIG. 21A-B, Table 8). When tested against an extended multiclade panel of 118 PsV, T-01 MB.v2 matched the pan-neutralization breadth of the corresponding IgG cocktail (100% virus coverage, cutoff IC₅₀ set at 10 μg/mL), yet displayed a remarkable neutralization potency (FIG. 21C-D, FIG. 22A and Table 9). Specifically, the IgG cocktail and T-01 MB were only able to neutralize 9% and 8% of the PsV with an IC₅₀ value of 0.001 μg/mL, respectively, while in the case of T-01 MB.v2, 50% of the PsVs were still neutralized with an IC₅₀ value of only 0.001 μg/mL (FIG. 21C). Remarkably, Multabodies achieved a median IC₅₀ value of only 0.0009 μg/mL (0.4 pM) and hence achieved pan-neutralization 32- and 490-fold more potently in mass and molarity, respectively, compared to the IgG cocktail (FIG. 21D). In addition, the IC₈₀ of T-01 MB.v2 confirmed its superior neutralization propensity over both the individual IgGs and the IgG cocktail, neutralizing 96% of all viral strains tested with a median IC₈₀ value of 0.005 μg/mL (2.2 pM) (FIG. 21C-D, FIG. 22A and Table 9). Importantly, Multabodies also blocked infection of primary peripheral blood mononuclear cells (PBMCs) with the replication-competent CXCR4-tropic HIV-1 IIIB strain (FIG. 22B), showing enhanced potency over the matched IgG mix, and without any impact on cell viability (FIG. 22C)

TABLE 9

Potency of parental and IgG mixtures and T-01 Multabody versions against a 118-PsV panel.

Virus ID	Clade	PGDM1400 IgG		10E8 IgG		N49P7 IgG		IgG mix
		IC ₅₀	IC ₈₀	IC ₅₀	IC ₈₀	IC ₅₀	IC ₈₀	
6535.3	B	>10	>10	0.081	1.32	0.186	0.598	0.526
QH0692.42	B	>10	>10	0.442	4.046	0.663	2.411	1.296
SC422661.8	B	0.436	6.530	0.445	2.715	0.114	0.320	0.160
PVO.4	B	0.484	6.487	0.865	7.197	0.107	0.365	0.653
TRO.11	B	0.283	1.376	0.034	0.250	0.044	0.160	0.092
AC10.0.29	B	0.105	0.869	0.132	1.534	5.034	>10	0.054
RHPA4259.7	B	0.447	2.194	0.728	5.275	0.083	0.246	0.156
THRO4156.18	B	0.072	0.348	0.156	0.884	2.341	>10	0.032
REJO4541.67	B	0.028	0.992	0.203	1.686	0.015	0.049	0.013
TRJO4551.58	B	>10	>10	0.475	4.783	0.104	0.351	0.768
WITO4160.33	B	<0.001	0.008	0.094	1.226	0.084	0.248	0.002

TABLE 9-continued

Potency of parental and IgG mixtures and T-01 Multibody versions against a 118-PsV panel.								
CAAN5342.A2	B	>10	>10	1.284	9.281	0.409	1.211	2.861
WEAU_d15_410_787	B (T/F)	0.103	1.366	6.039	>10	0.045	0.117	0.050
1006_11_C3_1601	B (T/F)	4.318	>10	0.172	2.999	0.044	0.158	0.311
1054_07_TC4_1499	B (T/F)	>10	>10	0.046	0.305	0.494	2.308	0.767
1056_10_TA11_1826	B (T/F)	3.258	>10	0.134	1.028	0.141	0.468	0.191
1012_11_TC21_3257	B (T/F)	0.049	0.230	0.244	4.211	0.046	0.125	0.045
6240_08_TA5_4622	B (T/F)	>10	>10	0.756	5.591	0.457	1.728	1.673
6244_13_B5_4576	B (T/F)	>10	>10	0.086	0.809	0.111	0.386	0.522
62357_14_D3_4589	B (T/F)	>10	>10	0.119	1.060	0.025	0.102	0.603
SC05_8C11_2344	B (T/F)	1.264	3.587	0.665	2.736	0.164	0.403	0.423
Du156.12	C	0.002	0.008	0.034	0.144	0.032	0.086	<0.0006
Du172.17	C	5.179	>10	0.109	0.837	0.102	0.668	0.110
Du422.1	C	0.694	>10	0.351	1.883	4.496	>10	0.375
ZM197M.PB7	C	0.063	0.285	0.148	0.802	0.109	0.347	0.016
ZM214M.PL15	C	>10	>10	0.896	5.021	0.213	0.953	0.941
ZM233M.PB6	C	<0.001	0.001	0.442	2.393	0.226	0.774	<0.0006
ZM249M.PL1	C	0.016	0.701	1.511	6.294	0.039	0.134	0.005
ZM53M.PB12	C	0.002	0.011	3.248	>10	0.251	0.800	0.001
ZM109F.PB4	C	0.057	0.382	0.295	1.888	0.034	0.153	0.013
ZM135M.PL	C	>10	>10	0.171	1.186	0.200	0.960	0.227
10a								
CAP45.2.00.G3	C	<0.001	0.002	0.120	1.927	0.060	0.290	0.002
CAP210.2.00.E8	C	0.034	0.514	0.798	3.892	1.817	6.485	0.057
HIV-001428-2.42	C	<0.001	0.027	0.809	7.298	0.006	0.020	0.006
HIV-0013095-2.11	C	0.003	0.012	0.031	0.213	0.073	0.253	0.001
HIV-16055-2.3	C	<0.001	0.003	0.722	6.495	0.057	0.280	0.001
HIV-16845-2.22	C	>10	>10	0.068	0.495	0.950	3.935	0.049
Ce1086_B2	C (T/F)	>10	>10	1.162	6.934	0.089	0.568	0.442
Ce0393_C3	C (T/F)	0.015	0.044	1.179	7.423	0.195	0.802	0.096
Ce1176_A3	C (T/F)	0.587	>10	0.867	3.923	0.204	0.589	0.102
Ce2010_F5	C (T/F)	>10	>10	1.844	8.612	0.130	0.547	1.397
Ce0682_E4	C (T/F)	0.010	0.043	1.215	7.709	0.041	0.131	0.012
Ce1172_H1	C (T/F)	0.021	0.060	0.722	4.818	8.499	>10	0.005
Ce2060_G9	C (T/F)	0.001	0.007	1.019	5.030	0.661	2.547	0.002
Ce703010054_2A2	C (T/F)	0.015	0.260	2.299	8.579	0.132	0.466	0.009
BF1266.431a	C (T/F)	0.003	0.011	1.070	8.418	0.057	0.199	0.005
246F C1G	C (T/F)	>10	>10	2.217	9.937	0.094	0.255	0.815
249M B10	C (T/F)	0.039	0.786	1.271	9.050	0.081	0.377	0.008
ZM247v1(Rev-)	C (T/F)	0.019	0.051	0.506	6.745	0.062	0.214	0.015
7030102001E5(Rev-)	C (T/F)	>10	>10	1.635	>10	0.184	0.832	2.318
1394C9G1(Rev-)	C(T/F)	<0.001	0.004	0.963	5.327	3.841	>10	0.001
Ce704809221_1B3	C (T/F)	0.280	1.099	0.058	0.319	0.141	0.932	0.071
CNE19	BC	0.001	0.007	0.666	6.485	0.009	0.041	0.001
CNE20	BC	0.004	0.014	1.246	6.577	0.028	0.100	0.005
CNE21	BC	<0.001	0.002	0.410	3.181	0.756	6.020	0.001
CNE17	BC	0.007	0.021	1.114	5.255	0.126	0.460	0.007
CNE30	BC	>10	>10	1.560	9.842	0.180	0.609	1.135
CNE52	BC	1.147	9.970	0.304	3.056	0.008	0.023	0.132
CNE53	BC	0.077	0.246	0.326	1.916	0.035	0.124	0.027
CNE58	BC	0.004	0.014	1.289	8.268	0.029	0.067	<0.0006
MS208.A1	A	<0.001	0.139	0.609	3.717	0.120	0.479	0.004
Q23.17	A	<0.001	0.003	1.637	7.040	0.052	0.210	<0.0006
Q461.e2	A	0.089	0.572	0.786	3.638	0.261	0.922	0.041
Q769.d22	A	0.002	0.018	1.667	10.000	0.025	0.111	0.001
Q259.d2.17	A	0.166	>10	3.956	>10	0.332	2.474	0.890
Q842.d12	A	<0.001	0.002	1.824	7.270	0.021	0.060	<0.0006
0260.v5.c36	A	0.033	0.175	6.452	>10	0.354	0.884	0.011
3415.v1.cl	A	0.036	0.327	2.481	12.306	0.160	0.550	0.068
3365.v2.e20	A	0.004	0.031	1.496	4.796	0.070	0.186	0.026
191955_A11	A (T/F)	0.001	0.021	0.680	3.532	1.811	8.870	0.001
191084 B7-19	A (T/F)	<0.001	0.007	1.653	8.138	0.047	0.142	<0.0006
9004SS_A3_4	A (T/F)	<0.001	0.002	1.160	6.761	0.125	0.390	<0.0006
T257-31	CRF02_AG	0.002	0.031	0.880	5.779	0.158	0.619	<0.0006
928-28	CRF02_AG	0.220	1.246	0.384	1.962	0.167	0.579	0.017
263-8	CRF02_AG	0.020	0.103	0.196	1.728	0.056	0.201	0.002
T250-4	CRF02_AG	<0.001	<0.001	1.346	8.130	2.070	>10	<0.0006

TABLE 9-continued

Potency of parental and IgG mixtures and T-01 Multabody versions against a 118-PsV panel.								
T251-18	CRF02_AG	8.615	>10	0.310	1.993	0.178	0.620	0.376
T278-50	CRF02_AG	0.574	4.942	0.949	7.460	>10	>10	0.049
T255-34	CRF02_AG	0.654	>10	0.778	4.207	0.053	0.262	0.078
211-9	CRF02_AG	0.088	0.393	1.091	6.508	0.220	0.767	0.024
235-47	CRF02_AG	0.005	0.032	0.384	2.072	0.045	0.300	<0.0006
620345.c01	CRF01_AE	0.009	>10	0.455	6.002	>10	>10	0.594
CNE8	CRF01_AE	<0.001	0.010	0.100	0.927	0.037	0.125	0.001
C1080.c03	CRF01_AE	<0.001	<0.001	0.061	0.731	0.114	0.765	<0.0006
R2184.c04	CRF01_AE	0.003	0.012	0.272	2.455	0.011	0.031	0.007
R1166.c01	CRF01_AE	1.431	6.921	0.274	2.614	0.092	0.313	0.144
R3265.c06	CRF01_AE	0.013	0.041	2.534	>10	0.048	0.282	0.025
C2101.c01	CRF01_AE	0.006	0.021	1.921	8.260	0.060	0.340	0.011
C3347.c11	CRF01_AE	0.011	0.051	0.016	0.167	0.024	0.067	0.012
C4118.c09	CRF01_AE	0.002	0.005	1.622	9.064	0.043	0.147	0.003
CNE5	CRF01_AE	<0.001	<0.001	1.408	9.637	0.157	0.513	0.001
BJOX009000.02.4	CRF01_AE	0.443	1.287	1.570	9.954	0.113	0.506	0.108
BJOX015000.11.5	CRF01_AE (T/F)	0.520	>10	0.051	1.909	0.036	0.157	0.345
BJOX010000.06.2	CRF01_AE (T/F)	0.605	5.870	0.341	5.293	0.085	0.592	0.074
BJOX025000.01.1	CRF01_AE (T/F)	0.087	0.573	0.107	6.325	0.038	0.236	0.053
BJOX028000.10.3	CRF01_AE (T/F)	0.072	0.963	0.665	7.361	0.020	0.123	0.021
X1193_c1	G	0.219	0.781	0.497	4.195	0.016	0.060	0.113
P0402_c2_11	G	0.010	0.033	0.172	3.288	0.014	0.061	0.013
X1254_c3	G	>10	>10	2.338	8.836	2.969	>10	6.824
X2088_c9	G	>10	>10	>10	>10	0.116	0.338	2.221
X2131_C1_B5	G	0.128	0.568	0.189	0.974	0.081	0.273	0.045
P1981_C5_3	G	0.104	4.601	0.057	0.285	0.113	0.274	0.030
X1632_S2_B10	G	0.055	1.445	1.976	9.684	>10	>10	0.028
3016.v5.c45	D	>10	>10	0.476	4.393	0.221	1.037	0.581
A07412M1.vrc12	D	0.032	0.127	0.576	6.054	0.108	0.645	0.012
231965.c01	D	0.025	1.280	5.648	>10	0.054	0.193	0.036
231966.c02	D	0.001	0.004	0.191	3.130	0.102	0.430	0.002
3817.v2.c59	CD	>10	>10	1.181	5.977	0.716	3.726	2.878
6480.v4.c25	CD	>10	>10	1.572	>10	0.031	0.113	0.151
6952.v1.c20	CD	0.501	>10	0.343	1.907	>10	>10	0.075
6811.v7.c18	CD	>10	>10	2.539	>10	0.229	0.992	2.862
89-F1_2_25	CD	0.001	0.005	1.112	10.000	>10	>10	0.003
3301.v1.c24	AC	0.048	0.245	1.341	7.095	0.025	0.069	0.023
6041.v3.c23	AC	0.013	0.289	2.333	9.766	0.047	0.222	0.022
6540.v4.c1	AC	0.004	0.028	1.899	10.000	>10	>10	0.008
6545.v4.c1	AC	0.003	0.039	1.778	8.229	2.001	>10	0.011
0815.v3.c3	ACD	>10	>10	0.332	2.868	0.016	0.053	0.116
3103.v3.c10	ACD	0.671	1.998	>10	>10	0.158	0.509	0.220
MuL V	Neg.	>10	>10	>10	>10	>10	>10	>10
Control	Control							
Median		0.056	0.570	0.725	5.265	0.110	0.396	0.029
% Virus Neutralized		81	71	98	91	95	88	100

Virus ID	Clade	IgG mix		T-01 MB		T-01 MB.v2	
		IC ₈₀	IC ₅₀	IC ₈₀	IC ₅₀	IC ₈₀	IC ₅₀
6535.3	B	3.614	0.715	>10	0.009	0.434	
QH0692.42	B	6.343	>10	>10	1.520	9.309	
SC422661.8	B	0.604	0.176	0.889	0.017	0.081	
PVO.4	B	2.329	0.157	1.151	0.017	0.112	
TRO.11	B	0.341	0.111	0.826	0.006	0.067	
AC10.0.29	B	0.273	0.019	0.108	0.001	0.010	
RHPA4259.7	B	0.731	0.053	0.372	0.010	0.067	
THRO4156.18	B	0.121	0.023	0.104	0.003	0.016	
REJO4541.67	B	0.130	0.001	0.006	<0.00004	0.0005	
TRJO4551.58	B	3.517	8.675	>10	0.298	2.489	
WITO4160.33	B	0.014	0.002	0.013	0.0001	0.001	
CAAN5342.A2	B	>10	>10	>10	1.254	7.486	
WEAU_d15_410_787	B (T/F)	0.811	0.001	0.007	0.0003	0.002	
1006_11_C3_1601	B (T/F)	1.498	0.013	0.059	0.001	0.006	
1054_07_TC4_1499	B (T/F)	3.645	>10	>10	1.843	>10	
1056_10_TA11_1826	B (T/F)	1.065	0.478	1.917	0.022	0.187	
1012_11_TC21_3257	B (T/F)	0.184	0.023	0.156	0.001	0.015	
6240_08_TA5_4622	B (T/F)	8.310	0.704	3.199	0.035	0.261	

TABLE 9-continued

Potency of parental and IgG mixtures and T-01 Multibody versions against a 118-PsV panel.						
6244__13__ B5_4576	B (T/F)	2.654	>10	>10	0.605	4.641
62357__14__ D3_4589	B (T/F)	2.220	5.734	>10	0.005	0.032
SC05__8C11__ 2344	B (T/F)	1.544	2.278	>10	0.011	0.054
Du156.12	C	<0.0006	0.051	0.215	0.0004	0.001
Du172.17	C	0.752	1.298	8.551	0.005	0.022
Du422.1	C	3.455	6.972	>10	0.021	1.216
ZM197M.PB7	C	0.065	0.183	0.895	0.001	0.003
ZM214M.PL15	C	>10	>10	>10	2.352	>10
ZM233M.PB6	C	0.001	0.019	0.081	0.0005	0.001
ZM249M.PL1	C	0.056	0.122	0.824	0.0006	0.002
ZM53M.PB12	C	0.007	0.0004	0.002	<0.00004	0.0003
ZM109F.PB4	C	0.084	0.006	0.022	0.001	0.005
ZM135M.PL 10a	C	1.080	2.467	>10	0.091	1.290
CAP45.2.00.G3	C	0.005	0.0004	0.002	0.0001	0.0007
CAP210.2.00.E8	C	0.555	0.410	6.302	0.0008	0.005
HIV-001428-2.42	C	0.050	0.001	0.003	0.0003	0.001
HIV-0013095-2.11	C	0.005	0.001	0.004	0.0002	0.0008
HIV-16055-2.3	C	0.004	0.0005	0.004	0.00004	0.001
HIV-16845-2.22	C	0.272	9.848	>10	0.385	3.286
Ce1086__B2	C (T/F)	2.899	3.797	>10	0.014	0.328
Ce0393__C3	C (T/F)	0.614	0.118	2.008	0.0006	0.003
Ce1176__A3	C (T/F)	0.574	0.065	0.442	0.0008	0.003
Ce2010__F5	C (T/F)	5.243	>10	>10	0.957	9.262
Ce0682__E4	C (T/F)	0.053	0.256	1.583	0.0008	0.004
Ce1172__H1	C (T/F)	0.016	0.095	0.445	0.0003	0.002
Ce2060__G9	C (T/F)	0.005	0.001	0.004	<0.00004	0.0002
Ce703010054__2A2	C (T/F)	0.060	0.003	0.011	0.0006	0.002
BF1266.431a	C (T/F)	0.013	0.0009	0.003	0.00005	0.0005
246F C1G	C (T/F)	2.791	8.146	>10	0.207	1.437
249M B10	C (T/F)	0.026	0.311	1.096	0.0002	0.0020
ZM247v1 (Rev-)	C (T/F)	0.043	0.004	0.017	0.0006	0.004
7030102001E5 (Rev-)	C (T/F)	>10	>10	>10	0.871	>10
1394C9G1 (Rev-)	C (T/F)	0.008	0.042	0.448	0.0004	0.002
Ce704809221__1B3	C (T/F)	0.242	0.014	0.104	0.002	0.010
CNE19	BC	0.005	0.001	0.004	0.0002	0.001
CNE20	BC	0.014	0.072	0.321	0.0004	0.002
CNE21	BC	0.004	0.0004	0.003	0.0001	0.0007
CNE17	BC	0.021	0.206	1.029	0.001	0.007
CNE30	BC	4.984	>10	>10	1.315	6.682
CNE52	BC	0.431	0.022	0.103	0.004	0.019
CNE53	BC	0.097	0.008	0.028	0.001	0.005
CNE58	BC	0.001	0.025	0.111	0.0002	0.001
MS208.A1	A	0.116	0.507	3.255	0.001	0.007
Q23.17	A	0.137	0.0003	0.002	0.0001	0.0005
Q461.e2	A	0.217	0.011	0.052	0.0003	0.004
Q769.d22	A	0.018	0.002	0.008	0.0001	0.001
Q259.d2.17	A	>10	0.006	0.310	0.00004	0.018
Q842.d12	A	0.002	0.0006	0.002	<0.00004	0.0002
0260.v5.c36	A	0.045	0.006	0.022	0.0006	0.003
3415.v1.cl	A	0.352	0.006	0.028	0.0002	0.002
3365.v2.c20	A	0.131	0.004	0.021	0.0003	0.003
191955__A11	A (T/F)	0.006	0.151	1.213	0.0004	0.002
191084 B7-19	A (T/F)	0.004	0.0006	0.003	<0.00004	0.0001
9004SS__A3__4	A (T/F)	<0.0006	0.052	0.219	0.0001	0.0007
T257-31	CRF02_AG	0.016	0.002	0.007	0.0001	0.0008
928-28	CRF02_AG	0.120	0.027	0.092	0.002	0.010
263-8	CRF02_AG	0.020	0.011	0.036	0.001	0.005
T250-4	CRF02_AG	<0.0006	0.0004	0.002	0.0001	0.0003
T251-18	CRF02_AG	1.859	0.716	4.276	0.049	0.316
T278-50	CRF02_AG	0.282	0.012	0.055	0.001	0.008
T255-34	CRF02_AG	0.433	0.003	0.036	0.0003	0.004
211-9	CRF02_AG	0.097	0.016	0.074	0.002	0.006
235-47	CRF02_AG	0.005	0.004	0.014	0.0009	0.002
620345.c01	CRF01_AE	>10	0.039	>10	0.003	3.278
CNE8	CRF01_AE	0.006	0.001	0.005	0.0002	0.001
C1080.c03	CRF01_AE	<0.0006	0.0007	0.002	0.0001	0.0003
R2184.c04	CRF01_AE	0.020	0.004	0.012	0.0003	0.002
R1166.c01	CRF01_AE	0.555	0.139	0.465	0.012	0.057
R3265.c06	CRF01_AE	0.113	0.003	0.018	0.0008	0.004
C2101.c01	CRF01_AE	0.036	0.005	0.031	0.0009	0.004
C3347.c11	CRF01_AE	0.039	0.007	0.029	0.0006	0.003

TABLE 9-continued

Potency of parental and IgG mixtures and T-01 Multabody versions against a 118-PsV panel.						
C4118.c09	CRF01__AE	0.007	0.002	0.006	0.0002	0.0007
CNE5	CRF01__AE	0.003	0.001	0.004	0.0002	0.0007
BJOX009000.02.4	CRF01__AE	0.432	0.039	0.236	0.005	0.023
BJOX015000.11.5	CRF01__AE (T/F)	1.649	0.013	0.088	0.001	0.007
BJOX010000.06.2	CRF01__AE (T/F)	1.262	0.022	0.153	0.001	0.007
BJOX025000.01.1	CRF01__AE (T/F)	0.365	0.036	0.281	0.003	0.018
BJOX028000.10.3	CRF01__AE (T/F)	0.151	0.014	0.174	0.0007	0.008
X1193__cl	G	0.490	0.045	0.218	0.002	0.015
P0402__c2__11	G	0.035	0.006	0.025	0.0008	0.005
X1254__c3	G	>10	1.688	>10	0.141	7.410
X2088__c9	G	8.326	3.978	>10	0.184	1.379
X2131__C1__B5	G	0.164	0.021	0.098	0.002	0.014
P1981__C5__3	G	0.137	0.010	0.044	0.001	0.004
X1632__S2__B10	G	0.275	0.002	0.009	<0.00004	0.0003
3016.v5.c45	D	5.079	4.892	>10	0.099	2.159
A07412M1.vrc12	D	0.040	0.003	0.017	0.0003	0.002
231965.c01	D	0.160	0.005	0.022	0.0009	0.006
231966.c02	D	0.005	0.003	0.009	0.0004	0.002
3817.v2.c59	CD	>10	3.879	>10	0.388	>10
6480.v4.c25	CD	0.694	0.787	4.400	0.072	0.470
6952.v1.c20	CD	0.358	0.020	0.365	0.001	0.019
6811.v7.c18	CD	>10	>10	>10	3.030	>10
89-F1__2__25	CD	0.008	0.001	0.004	0.0001	0.0005
3301.v1.c24	AC	0.080	0.007	0.035	0.0008	0.008
6041.v3.c23	AC	0.148	0.004	0.014	0.0007	0.003
6540.v4.cl	AC	0.081	0.004	0.014	0.0004	0.003
6545.v4.c1	AC	0.236	0.003	0.009	0.0002	0.0009
0815.v3.c3	ACD	0.520	5.031	>10	0.311	2.426
3103.v3.c10	ACD	1.013	0.030	0.080	0.001	0.008
MuL V	Neg.	>10	>10	>10	>10	>10
	Control					
Median		0.155	0.022	0.106	0.0009	0.005
% Virus Neutralized		93	92	80	100	96

DISCUSSION

[0255] The recent AMP clinical trials have highlighted the anticipated importance of both potency and breadth of bNAbs to be viable therapeutics capable of protecting against HIV-1 infection. Leveraging the principle of antibody avidity used in our previously-described Multabody technology to improve the potency of antibodies against SARS-CoV-2 (41), here we have engineered a second-generation Multabody platform to deliver exceptional neutralization breadth and potency against the vast sequence diversity of HIV-1.

[0256] The most distinctive feature of the optimized Multabody design in comparison to the first generation Multabody format (41) is the relative number of Fabs that self-associate per functional Fc domain. In contrast to the 1:1 Fc: 10E8v4/N49P7 ratio imposed by the design of the previously described Multabody platform, in the design of T-01 MB.v2, two N49P7 Fabs and two 10E8v4 Fabs are incorporated into the MB per dimeric Fc. The higher number of these two Fabs in the optimized Multabody favors their avidity and, consequently, their greater contribution to the neutralization signature of the particle. This is in contrast to the T-01 MB, which primarily relies on the neutralization properties of PGDM1400. The more balanced contribution of each of the antibodies is reflected in the better functional properties of T-01 MB.v2, which displayed cross-clade neutralization coverage of 100% at a median IC_{50} value of 0.0009 μ g/mL. In addition, viral infection by 83% of the 118-pseudoviruses tested was blocked by T-01 MB.v2 with an IC_{80} value below 1 μ g/ml, which has been recently proposed as the potency threshold required to confer in vivo

protection in humans (35). However, it remains unclear whether that predictor of protection should be considered in mass or molarity. Indeed, despite a similar hydrodynamic radius and geometrical size for the Multabody compared to an IgM (41), the Multabody is approximately 10 $\times$ heavier in molar mass compared to an IgG. Therefore, if molarity is the relevant in vivo measure associated with protection, then T-01 MB.v2 exhibits an extraordinarily low median IC_{50} value of 0.4 pM. Correspondingly, a potency of IC_{80} below 6.7 nM (1 μ g/ml molar equivalent for an IgG) was achieved in 96% of the 118-HIV-1 PsV strains. These remarkable neutralization properties surpass those obtained with previously described bispecific and tri-specific antibodies (46, 48-50). In these antibody formats, the limited avidity precludes combination of both high avidity and multi-specificity and, consequently, potency and breadth are restricted to that of the parental mAbs.

[0257] There is a growing trend in the field of biotherapeutics toward the development of molecules with high valency. Strategies range from the generation of dodecavalency IgM-like molecules (51, 52) upon addition of the mu-tailpiece of IgM to the constant region of IgG, to the design of alternative antibody formats. Among them are fusions of Fabs in a linear head-to-tail manner (53), appended IgGs (54-56), or diabody combination in tandem (Tamdabs) (57) or fused to the CH3 of an IgG (di-diabody) (58). In addition, the use of multimerization scaffolds such as p53 (59), leucine zipper helixes (60), streptavidin (61), barnase-barstar modules (62), viral-like nanoparticles (63) and, more recently, de novo antibody cage-forming proteins (64), have been employed to overcome the limitation of IgG bivalency and improve the bioactivity of antibodies.

Although attractive, these approaches face different challenges for their successful development as therapeutic agents. Multimeric antibody formats that rely on variable fragments (Fv) of antibodies are often associated with low stability and, consequently, a high propensity to aggregate (65). Furthermore, dissociation of non-covalent fusions dictated by the affinity constant of the complex can limit the in vivo long-term stability of the molecule. In sharp contrast, Multabodies build on full IgG components (Fab and Fc) that are fused to the thermostable, functionally-silent human apoferritin light chain scaffold, and thus are highly stable IgG-like molecules even under thermal stress. A mouse surrogate Multabody previously administered subcutaneously in immuno-competent C57BL/6 mice showed undetectable levels of anti-drug antibodies similarly to its parent IgG, providing proof-of-principle for the potentially low intrinsic immunogenicity of the Multabody platform (41). Future studies in higher organisms will help determine the immunogenicity of Multabodies encoded by human-derived sequences, which we propose might be dictated predominantly by the properties of the underlying antibody sequences.

[0258] Bioavailability of large biologics is an additional challenge associated with engineered approaches to increase avidity (63). Multabodies have been engineered to include Fc domains and hence enable FcRn-mediated recycling of the molecule. As a consequence of Fc avidity, binding of T-01 MB to FcRn at pH 6.0 was improved by orders of magnitude, however, the simultaneous affinity improvement at pH 7.4 limits the application of this strategy towards enhancing half-life, and several mutations were needed to decrease Fc binding to FcRn and FcγR in order to not surpass the binding affinity observed for IgGs. Such enhanced Fc receptor avidity was not observed in the case of T-01 MB.v2, where fusion of the Fc chains at the C-terminus of apoferritin leads to the formation of particles with inverted and more distantly located Fc domains and, consequently, reduced Fc avidity. Similar to a previous study with a mouse surrogate Multabody (41), Fc avidity modulation strategies successfully resulted in Multabody molecules with remarkably similar rate of decay over time as the parental IgG cocktail. In addition to the favorable pharmacokinetic profile, Multabodies in both formats that possess residual binding to FcγR induced Fc-mediated phagocytosis in vitro to levels similar to the parental IgG mixture, at least in a THP-1 system that co-expresses both FcγRI and FcγRIIa (66). Future experiments will be needed to fully characterize the capacity of the Multabodies to trigger immune effector functions and their in vivo implications.

[0259] From the limited number of antibody specificities we characterized in this study, we observed that antibodies targeting epitopes located at the apex of the HIV-1 Env trimer such as PGDM1400 seem to experience the greatest benefit to neutralization potency when formulated as Multabodies. This increase in potency was less apparent as the epitope was located closer to the viral membrane, as in the case of 10E8. The dependence on epitope location for potency enhancement could be further impacted by the low surface spike density (67), the arrangement of those sparse Env trimers on the HIV-1 surface (68) or by the accessibility of certain epitopes that may be more or less sterically occluded. In light of this, it will be interesting to explore how the potency of antibodies against viruses with higher surface densities and closely spaced spikes can be enhanced

by the Multabody platform, and further determine the impact of epitope location on potency enhancement mediated by avidity.

[0260] Replacement of N49P7 with iMab—a CD4-directed post-attachment inhibitor—resulted in a functional Multabody with potent neutralization activity, demonstrating that cross-targeting of viral epitopes and cellular receptors can be attained with this type of particle. This data raises the interesting possibility that the Multabody technology could also be implemented in other fields that promote the binding of receptors across separate entities, such as cell-cell interactions in immunotherapy. Overall, our protein engineering study demonstrates the versatility of human apoferritin as a modular nanocage to build antibody avidity and multi-specificity jointly towards enhanced functionality.

REFERENCES

- [0261]** 1. A. J. Conley, et al., Neutralization of divergent human immunodeficiency virus type 1 variants and primary isolates by IAM-41-2F5, an anti-gp41 human monoclonal antibody. *Proc. Natl. Acad. Sci. U.S.A* 91, 3348-3352 (1994).
- [0262]** 2. G. Stiegler, et al., A Potent Cross-Clade Neutralizing Human Monoclonal Antibody against a Novel Epitope on gp41 of Human Immunodeficiency Virus Type 1. *AIDS Res. Hum. Retroviruses* 17, 1757-65 (2001).
- [0263]** 3. M. B. Zwick, et al., Broadly Neutralizing Antibodies Targeted to the Membrane-Proximal External Region of Human Immunodeficiency Virus Type 1 Glycoprotein gp41. *J Virol.* 75, 10892-905 (2001).
- [0264]** 4. A. Buchacher, et al., Generation of Human Monoclonal Antibodies against HIV-1 Proteins; Electrofusion and Epstein-Barr Virus Transformation for Peripheral Blood Lymphocyte Immobilization. *AIDS Res. Hum. Retroviruses* 10, 359-69 (1994).
- [0265]** 5. C. F. Barbas, et al., Recombinant human Fab fragments neutralize human type 1 immunodeficiency virus in vitro. *Proc. Natl. Acad. Sci. U.S.A* 89, 9339-9343 (1992).
- [0266]** 6. D. R. Burton, et al., Efficient neutralization of primary isolates of HIV-1 by a recombinant human monoclonal antibody. *Science* 266, 1024-1027 (1994).
- [0267]** 7. X. Wu, et al., Rational design of envelope identifies broadly neutralizing human monoclonal antibodies to HIV-1. *Science* 329, 856-861 (2010).
- [0268]** 8. J. F. Scheid, et al., Broad diversity of neutralizing antibodies isolated from memory B cells in HIV-infected individuals. *Nature* 458, 636-640 (2009).
- [0269]** 9. D. Sok, et al., Recombinant HIV envelope trimer selects for quaternary-dependent antibodies targeting the trimer apex. *Proc. Natl. Acad. Sci. U.S.A* 111, 17624-17629 (2014).
- [0270]** 10. L. M. Walker, et al., Broad neutralization coverage of HIV by multiple highly potent antibodies. *Nature* 477, 466-470 (2011).
- [0271]** 11. N. A. Doria-Rose, et al., Developmental pathway for potent V1V2-directed HIV-neutralizing antibodies. *Nature* 509, 55-62 (2014).
- [0272]** 12. J. Huang, et al., Broad and potent neutralization of HIV-1 by a gp41-specific human antibody. *Nature* 491, 406-412 (2012).
- [0273]** 13. L. M. Walker, et al., Broad and potent neutralizing antibodies from an African donor reveal a new HIV-1 vaccine target. *Science* 326, 285-289 (2009).

- [0274] 14. M. M. Sajadi, et al., Identification of Near-Pan-neutralizing Antibodies against HIV-1 by Deconvolution of Plasma Humoral Responses. *Cell* 173, 1783-1795.e14 (2018).
- [0275] 15. J. F. Scheid, et al., Sequence and Structural Convergence of Broad and Potent HIV Antibodies That Mimic CD4 Binding. *Science* 333, 1633-1637 (2011).
- [0276] 16. T. Schoofs, et al., Broad and Potent Neutralizing Antibodies Recognize the Silent Face of the HIV Envelope. *Immunity* 50, 1513-1529.e9 (2019).
- [0277] 17. J. Huang, et al., Identification of a CD4-Binding-Site Antibody to HIV that Evolved Near-Pan Neutralization Breadth. *Immunity* 45, 1108-1121 (2016).
- [0278] 18. R. Pejchal, et al., A potent and broad neutralizing antibody recognizes and penetrates the HIV glycan shield. *Science* 334, 1097-1103 (2011).
- [0279] 19. C. Blattner, et al., Structural delineation of a quaternary, cleavage-dependent epitope at the gp41-gp120 interface on intact HIV-1 env trimers. *Immunity* 40, 669-680 (2014).
- [0280] 20. H. Mouquet, et al., Complex-type N-glycan recognition by potent broadly neutralizing HIV antibodies. *Proc. Natl. Acad. Sci. U.S.A* 109, E3268-77 (2012).
- [0281] 21. T. W. Baba, et al., Human neutralizing monoclonal antibodies of the IgG1 subtype protect against mucosal simian-human immunodeficiency virus infection. *Nat. Med.* 6, 200-206 (2000).
- [0282] 22. A. J. Hessel, et al., Effective, low-titer antibody protection against low-dose repeated mucosal SHIV challenge in macaques. *Nat. Med.* 15, 951-954 (2009).
- [0283] 23. A. J. Hessel, et al., Broadly neutralizing human anti-HIV antibody 2G12 is effective in protection against mucosal SHIV challenge even at low serum neutralizing titers. *PLoS Pathog.* 5, e1000433 (2009).
- [0284] 24. R. Hofmann-Lehmann, et al., Postnatal pre- and postexposure passive immunization strategies: Protection of neonatal macaques against oral simian-human immunodeficiency virus challenge. *J. Med. Primatol.* 31, 109-119 (2002).
- [0285] 25. R. Hofmann-Lehmann, et al., Postnatal passive immunization of neonatal macaques with a triple combination of human monoclonal antibodies against oral simian-human immunodeficiency virus challenge. *J. Virol.* 75, 7470-7480 (2001).
- [0286] 26. Y. U. Van Der Velden, et al., Short Communication: Protective Efficacy of Broadly Neutralizing Antibody PGDM1400 Against HIV-1 Challenge in Humanized Mice. *AIDS Res. Hum. Retroviruses* 34, 790-793 (2018).
- [0287] 27. F. Klein, et al., HIV therapy by a combination of broadly neutralizing antibodies in humanized mice. *Nature* 492, 118-122 (2012).
- [0288] 28. M. Deruaz, et al., Protection of humanized mice from repeated intravaginal HIV challenge by passive immunization: A model for studying the efficacy of neutralizing antibodies in vivo. *J. Infect. Dis.* 214, 612-616 (2016).
- [0289] 29. J. A. Horwitz, et al., HIV-1 suppression and durable control by combining single broadly neutralizing antibodies and antiretroviral drugs in humanized mice. *Proc. Natl. Acad. Sci. U.S.A* 110, 16538-16543 (2013).
- [0290] 30. S. Mehndru, et al., Adjunctive Passive Immunotherapy in Human Immunodeficiency Virus Type 1-Infected Individuals Treated with Antiviral Therapy during Acute and Early Infection. *J. Virol.* 81, 11016-11031 (2007).
- [0291] 31. M. Caskey, et al., Viraemia suppressed in HIV-1-infected humans by broadly neutralizing antibody 3BNC117. *Nature* 522, 487-491 (2015).
- [0292] 32. M. Caskey, et al., Antibody 10-1074 suppresses viremia in HIV-1-infected individuals. *Nat. Med.* 23, 185-191 (2017).
- [0293] 33. R. M. Lynch, et al., Virologic effects of broadly neutralizing antibody VRC01 administration during chronic HIV-1 infection. *Sci. Transl. Med.* 7, 319ra206 (2015).
- [0294] 34. J. E. Ledgerwood, et al., Safety, pharmacokinetics and neutralization of the broadly neutralizing HIV-1 human monoclonal antibody VRC01 in healthy adults. *Clin. Exp. Immunol.* 182, 289-301 (2015).
- [0295] 35. L. Corey, et al., Two Randomized Trials of Neutralizing Antibodies to Prevent HIV-1 Acquisition. *N. Engl. J. Med.* 384, 1003-1014 (2021).
- [0296] 36. R. S. Rudicell, et al., Enhanced Potency of a Broadly Neutralizing HIV-1 Antibody In Vitro Improves Protection against Lentiviral Infection In Vivo. *J. Virol.* 88, 12669-12682 (2014).
- [0297] 37. Y. D. Kwon, et al., Surface-Matrix Screening Identifies Semi-specific Interactions that Improve Potency of a Near Pan-reactive HIV-1-Neutralizing Antibody. *Cell Rep.* 22, 1798-1809 (2018).
- [0298] 38. E. Rujas, et al., Functional Optimization of Broadly Neutralizing HIV-1 Antibody 10E8 by Promotion of Membrane Interactions. *J. Virol.* 92, e02249-17 (2018).
- [0299] 39. E. Rujas, et al., Affinity for the Interface Underpins Potency of Antibodies Operating In Membrane Environments. *Cell Rep.* 32, 108037 (2020).
- [0300] 40. R. Diskin, et al., Increasing the potency and breadth of an HIV antibody by using structure-based rational design. *Science* 334, 1289-1293 (2011).
- [0301] 41. E. Rujas, et al., Multivalency transforms SARS-CoV-2 antibodies into broad and ultrapotent neutralizers. *Nat Commun* 12, 3661 (2021).
- [0302] 42. Y. D. Kwon, et al., Optimization of the Solubility of HIV-1-Neutralizing Antibody 10E8 through Somatic Variation and Structure-Based Design. *J. Virol.* 90, 5899-5914 (2016).
- [0303] 43. J. M. Jacobson, et al., Safety, pharmacokinetics, and antiretroviral activity of multiple doses of ibalizumab (formerly TNX-355), an anti-CD4 monoclonal antibody, in human immunodeficiency virus type 1-infected adults. *Antimicrob. Agents Chemother.* 53, 450-457 (2009).
- [0304] 44. D. R. Kuritzkes, et al., Antiretroviral Activity of the Anti-CD4 Monoclonal Antibody TNX-355 in Patients Infected with HIV Type 1. *J. Infect. Dis.* 189, 286-291 (2004).
- [0305] 45. D. C. Montefiori, Measuring HIV neutralization in a luciferase reporter gene assay. *Methods Mol. Biol.* 485, 395-405 (2009).
- [0306] 46. L. Xu, et al., Trispecific broadly neutralizing HIV antibodies mediate potent SHIV protection in macaques. *Science* 358, 85-90 (2017).
- [0307] 47. O. S. Qureshi, et al., Multivalent Fc γ -receptor engagement by a hexameric Fc-fusion protein triggers Fc γ -receptor internalisation and modulation of Fc γ -receptor functions. *Sci. Rep.* 7 (2017).

- [0308] 48. M. Asokan, et al., Bispecific Antibodies Targeting Different Epitopes on the HIV-1 Envelope Exhibit Broad and Potent Neutralization. *J Virol.* 89, 12501-12512 (2015).
- [0309] 49. S. N. Khan, et al., Targeting the HIV-1 Spike and Coreceptor with Bi- and Trispecific Antibodies for Single-Component Broad Inhibition of Entry. *J Virol.* 92, e00384-18 (2018).
- [0310] 50. J. J. Steinhardt, et al., Rational design of a trispecific antibody targeting the HIV-1 Env with elevated anti-viral activity. *Nat. Commun.* 9 (2018).
- [0311] 51. R. I. Smith, M. J. Coloma, S. L. Morrison, Addition of a mu-tailpiece to IgG results in polymeric antibodies with enhanced effector functions including complement-mediated cytolysis by IgG4. *J. Immunol.* 154, 2226-36 (1995).
- [0312] 52. T. Olafsen, B. I. Rasmussen, L. Norderhaug, Ø. S. Bruland, I. Sandlie, IgM secretory tailpiece drives multimerisation of bivalent scFv fragments in eukaryotic cells. *Immunotechnology* 4, 141-153 (1998).
- [0313] 53. K. Miller, et al., Design, Construction, and In Vitro Analyses of Multivalent Antibodies. *J. Immunol.* 170, 4854-4861 (2003).
- [0314] 54. C. Wu, et al., Simultaneous targeting of multiple disease mediators by a dual-variable-domain immunoglobulin. *Nat. Biotechnol.* 25, 1290-1297 (2007).
- [0315] 55. C. Klein, W. Schaefer, J. T. Regula, The use of CrossMab technology for the generation of bi- and multispecific antibodies. *MAbs* 8, 1010-1020 (2016).
- [0316] 56. A. Steinmetz, et al., CODV-Ig, a universal bispecific tetravalent and multifunctional immunoglobulin format for medical applications. *MAbs* 8, 867-878 (2016).
- [0317] 57. S. M. Kipriyanov, et al., Bispecific tandem diabody for tumor therapy with improved antigen binding and pharmacokinetics. *J. Mol. Biol.* 293, 41-56 (1999).
- [0318] 58. D. Lu, et al., Di-diabody: A novel tetravalent bispecific antibody molecule by design. *J. Immunol. Methods* 279, 219-232 (2003).
- [0319] 59. S. Kubetzko, E. Balic, R. Waibel, U. Zangemeister-Wittke, A. Pluckthun, PEGylation and multimerization of the anti-p185HER-2 single chain Fv fragment 4D5: Effects on tumor targeting. *J. Biol. Chem.* 281, 35186-35201 (2006).
- [0320] 60. P. Pack, K. Müller, R. Zahn, A. Pluckthun, Tetravalent miniantibodies with high avidity assembling in *Escherichia coli*. *J. Mol. Biol.* 246, 28-34 (1995).
- [0321] 61. S. M. Kipriyanov, et al., Affinity enhancement of a recombinant antibody: Formation of complexes with multiple valency by a single-chain Fv fragment-core streptavidin fusion. *Protein Eng.* 9, 203-211 (1996).
- [0322] 62. S. M. Deyev, R. Waibel, E. N. Lebedenko, A. P. Schubiger, A. Pluckthun, Design of multivalent complexes using the barnase barstar module. *Nat. Biotechnol.* 21, 1486-1492 (2003).
- [0323] 63. M. A. G. Hoffmann, et al., Nanoparticles presenting clusters of CD4 expose a universal vulnerability of HIV-1 by mimicking target cells. *Proc. Natl. Acad. Sci. U.S.A* 117, 18719-18728 (2020).
- [0324] 64. R. Divine, et al., Designed proteins assemble antibodies into modular nanocages. *Science* 372, eabd99947 (2021).
- [0325] 65. A. Wörn, A. Pluckthun, Different equilibrium stability behavior of scFv fragments: Identification, classification, and improvement by protein engineering. *Biochemistry* 38, 8739-8750 (1999).
- [0326] 66. H. B. Fleit, C. D. Kobasiuk, The Human Cell Line THP-1 Expresses Fc-gamma-RI and Fc-gamma-RII. *J Leukoc Biol* 49, 556-565 (1991).
- [0327] 67. P. Zhu, et al., Electron tomography analysis of envelope glycoprotein trimers on HIV and simian immunodeficiency virus virions. *Proc. Natl. Acad. Sci. U.S.A* 100, 15812-15817 (2003).
- [0328] 68. J. Chojnacki, et al., Maturation-dependent HIV-1 surface protein redistribution revealed by fluorescence nanoscopy. *Science* 338, 524-528 (2012).
- [0329] 69. M. Graille, et al., Complex between *Peptostreptococcus magnus* protein L and a human antibody reveals structural convergence in the interaction modes of Fab binding proteins. *Structure* 9, 679-687 (2001).
- [0330] 70. T. Zhou, et al., Structural basis for broad and potent neutralization of HIV-1 by antibody VRC01. *Science* 329, 811-817 (2010).
- [0331] 71. E. O. Freed, D. J. Myers, R. Risser, Mutational analysis of the cleavage sequence of the human immunodeficiency virus type 1 envelope glycoprotein precursor gp160. *J. Virol.* 63, 4670-4675 (1989).

SEQUENCE LISTING

[0332] Underlining within sequences indicate linker sequences; bolding within sequences indicate ferritin or ferritin subunit sequences; boxed and bolded residues indicate residues that are mutated relative to a reference molecule, e.g. relative to an IgG1 Fc.

hFTL

SEQ ID NO: 1

MSSQIRQNYSTDVEAAVNSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEKREG

YERLLKMQNQRGGRALFQDIKKPAEDEWGKTPDAMKAAMALEKKLNQALLDLHALGSARTDP

HLCDFLETHFLDEEVKLIKMGDHLTNLHRLGGPEAGLGEYLFERLTLRHD

N hFTL

SEQ ID NO: 2

MSSQIRQNYSTDVEAAVNSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEKREG

YERLLKMQNQRGGRALFQDIKKPAEDW

SEO ID NO:3

HRLGGPEAGLGEYLFERLTRHD

SEO ID NO:4

HRLGGPEAGLGEYLFERLTLLKHD

SEO ID NO:5

YSKLTVDKSRWOOGNVFSCSVMHEALHNHYTOKSLSLSPGK

SEQ ID NO: 6

GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGDKTHTCPPCPAPEL

NVFSCSVMHEALHNHYTOKSLSLSPGK

SEQ ID NO: 7

SKLRVEKKNWVERNSYSCSVHEGLHNHHTTKSF SRTPGK

SEO ID NO: 8

GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGVECPPCPAPPVAGPSVFL

HEALHNHYTOKSLSLSPGK

SEO ID NO: 9

IgG4 scFc

SEO ID NO:10

PGDM1400 IgG1 HC

SEO ID NO:11

PGDM1400 LC

SEQ ID NO:12

PGDM1400 HC (Fab)

SEO ID NO:13

QAQLVQSGPEVRKPGTSVKVSCKAPGNTLTKTYDLHWVRSPVGQGLWGMWISHEGDKKVVIVE
RFKAKVTIDWDRSTNTAYLQLSGLTSGDTAVYYCAKSKHRLRDYALYDDDGALNWAVDVDY
LSNLEFWGQGTAVTVSSASTKGPSVFPLAPSSKSTSGGTAALGLCKYDYFPEPVTVSWNSGA
LTSGVHTFPFVAVLOSSGLYSLSSVVTVPSSSLGTOTYICNVNHKPSNTKVDKKVEPKSC

PGDM1400 scFab

SEO ID NO:14

DFVLTSQSPHSLSVTPGESASISCKSSHSLIHGDRNNYLAWYVOKPGRSPOLLITLASSRASG
VPDRFSGSGSKDKFTLKISRVEDVGTYYCQMGRSPWTFGQGTVDIKRTVAAPSVFIFPP
PSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDSSTYLSSTLT

SEO ID NO:15

SEO ID NO:16

SEO ID NO:17

SEQ ID NO:18

SEO ID NO:19

EVRLVESGGGLVKPGGSLRLSCSASGFDNAWMTWRQPPGKGLEWVGRITGPGEQWSVDY
AESVKGRFTISRDNKNTLYLEMMNVRTEDTGYYFCARTGKYIDFWSGYPPEGEYFQDWGQG

GVPDRFSGSGSGTDEFTLTISSSVOAEDVAVYYCOOYYSYRTFEGGGTKLEIKRTVAAPSVEFIEP

-continued

PSDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT
SKADYEKHKVYACEVTHQGLSSPVTKEENRGEC

iMab HC (Fab)

SEQ ID NO:25

QVQLQQSGPEVVKPGASVKMSCKASGYTFTSYVIHWVRQKPGQGLDWIGYINPYNDGTDYDE
KFKGKATLTSDTSTAYMELSSLRSEDTAVYYCAREKDNATGAWFAYWGQGLTVTVSSAS
TKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYS
LSSVVTVPSSSLGTQTYTCNVDPKPSNTKVDKRVESKYC

iMab scFab

SEQ ID NO:26

DIVMTQSPDLSAVSLGERVTMNCSSQSLLYSTNQKNYLAWYQQKPGOSPKLLIYWASTRES
GVPDRFSGSGSGTDFTLTITSSVQAEDVAVYYCQYYSYRTFGGGTKEIKRTVAAPSVFIFP
PSDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT
SKADYEKHKVYACEVTHQGLSSPVTKEENRGECGGGGSGGGSGGGSGGGSGGGSGGGG
SGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGVQLQQSGPEVVKPGASVKMS
CKASGYTFTSYVIHWVRQKPGQGLDWIGYINPYNDGTDYDEKFKGKATLTSDTSTAYMEL
SSLRSEDTAVYYCAREKDNATGAWFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA
ALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSC

PGDM1400-hFTL

SEQ ID NO:27

DFVLTQSPHSLVTPGESASISCKSSSHSLIHGDRNNYLAWYVQKPGRSPOLLIYLASSRASG
VPDRFSGSGSDKDFTLKISRVEDTVGTYYCMQGRESPTWTFGQGTKVDIKRTVAAPSVFIFP
PSDEQLKSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDSKDYSLSTLT
SKADYEKHKVYACEVTHQGLSSPVTKEENRGECGGGGSGGGSGGGSGGGSGGGSGGGG
SGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGAQLVQSGPEVRKPGTSVKVS
CKAPGNTLTKTYDLHWVRVSPGQGLWMGWI SHEGDKKIVVERFKAKVTIDWDRSTNTAYLQL
SGLTSGDTAVYYCAKGSKHLRDYALYDDDGALNWAVDVYLSNLEFWGQGTAVTVSSASTK
GPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSL
SVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDGGGGSGGGSGGGSGGGSGGGG
SGGSSQIRQNYSTDVEAVNSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEEKR
EGYERLLKMQRGGRALFQDIKKPAEDEVGKTPDAMKAAMALEKKLNQALLDLHALGSART
DPHLCDFLETHFLDEEVKLIKMGDHLTNLHRLGGPEAGLGEYLFERLTLRHD

N49P7-C hFTL

SEQ ID NO:28

QSALTQPRSVSASPGQSVTISCTGTHNLVSWCQHOPGRAPKLLIYDENKRPSGVPDRESGSG
SGGTASLTITGLQDDDAEYFCWAYEAFGGGKLTVLGQPKAAPSVLTLEPPSSEELQANKAT
LVCLISDFYPGAVTVAWKADSSPVKAGVETTPSKQSNKYAASSYLSLTPEQWKSHRSYSC
QVTHEGSTVEKTVAPTECGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGG
SGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGADLVQSGAVVKKPGDSVRI SCAQGYRFPDIYI IHWI
RRAPGQGPEWMGMNPMGGQVNI PWKFGQGRVSMTRDTSIETAFDLRLGLKSDDTAVYYCVRD
RSNGSGKRFESSNWFLDLWGRGTAVTIQSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF

PEPVTVSWNSGALTSVHTFPFVAVLQSSGLYLSVVTPVSSSLGTQTYICNVNHPKPSNTKVD
KKVEPKSCGGGGSGGGSGGGSGGGSGGGSGGGSGGGKTPDAMKAAMALEKKLNQALLDLHAL
GSARTDPHLCDFLETHFLDEEVKLIKMGDHLTNLHRLGGPEAGLGEYLFERLTRHD

N49P7-N hFTL

SEO ID NO:29

QSALTQPRSVSASPGQSVTISCTGTHNLVSWCQHQPGRAPKLLIYDENKRPSGVPDRESGSG
 SGGTASLTITGLQDDDDAEYFCWAYEAPGGGTKLTVLGQPKAAPSVTLEPPSSSEELQANKAT
 LVCLISDFYPGAVTVAWKADSPVKAGVETTTPSKOSNNKYAASSYLSLTPEQWKSHRYSYC
 QVTHEGSTVEKTVAPTECGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGG
SGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGG
 SGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGG
 RRAPQGQPEWMGWMNPMGGQVNI PWKFQGRVSMTRDTSIETAFLDLRGLKSDDTAVYYCVRD
 RNSGSGKRFESSNWFLDLWGRGTAVTIQSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYF
 PEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVTPSSSLGTQTYICNVNHKPSNTKVD
 KKVEPKSCGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGG
 YTYLSLGFYFDRDDVALEGVSHFFRELAEEKREGYERLLKMONORGGGRALFODIKKPAEDEW

10E8v4-C hFTL

SEQ ID NO:30

S E L T Q D P A V S V A L K Q I V T I T C R G D S L R S H Y A S W Y Q K K P G Q A P V L L F Y G K N N R P S G I P D R E S G
S A S G N R A S L T I T G A Q A E D E A D Y Y C S S R D K S G S R L S V F G G G T K L T V L S Q P K A A P S V T L F P P S S
E E L Q A N K A T L V C L I S D F Y P G A V T V A W K A D S S P V K A G V E T T P S K O S N N K Y A A S S Y L S L T P E Q
W K S H R S Y S C Q V T H E G S T V E K T V A P T E C G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S
G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S E V R L V E S G G L V K P G G S L R L S C S A S G F
D F D N A W M T W R Q P P K G L E W V G R I T G P G E G S V D Y A E S V K G R F T I S R D N T K N T L Y L E M N N V R
T E D T G Y Y F C A R T G K Y Y D F W S G Y P P G E E Y F Q D W G Q T L V I V S S A S T K G P S V F P L A P S S K S T S G
G T A A L G C L V K D Y F P E P V T V S W N S G A L T S G V H T F P A V L Q S S G L Y S L S S V V T V P S S S L G T Q T Y I
C N V N H K P S N T K V D K K V E P K S C G G G S G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S G G G G S
E K K L N Q A L L D L H A L G S A R T D P H L C D F L E T H F L D E E V K L I K M G D H L T N L H R L G G P E A G L G E Y
L F E R L T L R H D

iMab-C hFTL

SEO ID NO:31

DIVMTQSPDLSLAVSLGERVTMNCSSQSLLYSTNOKNYLAWYQQKPGGSKLLIYWASTRES
GVDPDRSGSGSGTDFTLTISSVQAEDVAVYYCQQYYSYRTFGGGTKLEIKRTVAAPSVEIFP
PSDEQLKSGTASVVCLLNFFYPREAKVQWKVDNALQSGNSQESVTEQDSKSDSTYLSLSTLT
SKADYEKHKVYACEVTHQGLSPVTKSENREGCGGGGSGGGGSGGGGSGGGGSGGGGSGGGG
SGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSGGGGSQVLQQSGPEVVKPGASVKMS
CKASGYFTTSYVIHWVRQKPGQLDWIGYINPYNDGDYDEKFKGKATLSTDTSTAYMEL
SSLRSEDTA VYYCAREKDNYATGAWFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTA
ALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSVVTVTPSSSLGTQTYICNV
NHKPSNTKVDKKVEPKSCGGGGSGGGGSGGGGSGGGGSGGGGSGGGGGKTPDAMKAMALEKK
LNQALLDLHALGSARTDPHLCDFLETHFLDEEVKLIIKKMGDHLTNLHRLGGPEAGLGLEYLFE
RLTLEHD

[illegible][illegible]

IgG1_scFc-N_hFTL

SEQ ID NO: 34

DKTHTCPPCPAPELLGGPSVFLFPPPKDTLMISRTPEVTCVVDVSHEDPEVKENWYVDGV

EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQPR

EPQVYITLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLVSDGSFFL

YSKLTVDKSRWQQGNVSCFVSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSG

-continued

GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGGDKTHTCPPCPAPEL
 LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY
 NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
 TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFFLYSKLTVDKSRWQQG
 NVFSCSVMEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGGGSSQIRQNY
STDVEAAVNSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEEKREGYERLLKMQN
QRGGRALFQDIKKPAEDEW

IgG2_scFc-N_hFTL

SEQ ID NO:35

VECPPCPAPPVAGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHN
 AKTKPREEQFNSTFRVSVLTVVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQV
 YTLPPSREEMTKNQVSLTCLVKGFYPSDISVEWESNGQPENNYKTTPMLDSGSFFLYSKL
 TVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGGGG
 GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGVECPPCPAPPVAGPSVEL
 FPPKPKDTLMISRTPEVTCVVVDVSHEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTFRVVS
 VLTVTVHQDWLNGKEYKCKVSNKGLPAPIEKTISKTKGQPREPQVYTLPPSREEMTKNQVSLT
 CLVKGFYPSDISVEWESNGQPENNYKTTPMLDSGSFFLYSKLTVDKSRWQQGNVFCSCVM
 HEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGGGSGGGSSQIRQNYSTDVEAAV
NLSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEEKREGYERLLKMQNORGGRALF
QDIKKPAEDEW

IgG4_scFc-N_hFTL

SEQ ID NO:36

KYGPPCPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSEQEDPEVQFNWYVDGV
 EVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPR
 EPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFEL
 YSRLTVDKSRWQEGNVFCSCVMHEALHNHYTQKSLSLGLKGGGSGGGSGGGSGGGSGGGSGG
 GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGKYGPPCPCPAPEF
 LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSEQEDPEVQFNWYVDGVEVHNAKTKPREEQF
 NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEM
 TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFFLYSRLTVDKSRWQEG
 NVFSCSVMEALHNHYTQKSLSLGLKGGGSGGGSGGGSGGGSGGGSGGGSSQIRQNY
STDVEAAVNSLVNLYLQASYTYLSLGFYFDRDDVALEGVSHFFRELAEEKREGYERLLKMON
QRGGRALFQDIKKPAEDEW

IgG1_P329G_scFc-N_hFTL

SEQ ID NO:37

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGV
 EVHNAKTKPREEQNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR
 EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGSFFL
 YSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGG
 GGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGDKTHTCPPCPAPEL
 LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY

NSTYRVVSVLTVLHODWLNQKEYKCKVSNKALGAPIEKTIISKAKGPREPOVYTLPPSREEM

IqGI LALA scFc-N hFTL

SEO ID NO:38

DKTHTCPPCPAP**E**AAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV

[illegible]

AGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEOY

NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM
 TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSGGSFFLYSKLTVDKSRWQQG
 NVFSCSVMHIALHNYHTQKSLSLSPGKGGGSGGGSGGGSGGGSGGGSGGGSSQIRQNY
 STDVEAAVNSLVNLYLQASYYTSLSLGFYFDRDDVALEGVSHFFRELAEEKREGYERLLKMON
 ORGGRALFODIKKPAEDEW

IqG1 LALAP scFc-N hFTL

SEO ID NO:39

DKTHTCPPCPAPEAAGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGV

[illegible]

AGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY

NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKAL**G**APIEKTISKAKGQPREPQVYTLPPSREEM
 TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQG
 NVFSCSVMHAEALHNYHTQKSLSLSPGKGGGGGGGGGGGGGGGGGGGGGGGGSS**Q**IRONY
 STDVEAAVNSLVNLYLQASYYTSLSLGFFYFDRDDVALEGVSHFFRELAEKREGYERLLKMON
 QRGGRALFQDIKKPAEDEW

IgG1 K288A scFc-N hFTL

SEO ID NO:40

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKENWYVDGV

EVHNA**D**TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYCKVSNKALPAPIEKTI SKAKGQPR
EPQVYITLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFL
YSKLTVDKSRWQQGNVSCSVMHEALHNHYTQKSLSLSPGKGGGGSGGGGSGGGGSGGGGSG

-continued

GGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGDKTHTCPPCPAPEL

LGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA**AT**KPREEQY

NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM

TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFFLYSKLTVDKSRWQQG

NVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGSS**QIRQNY**

STDVEAAVNSLVNLYLQASYYTSLGLGYFDRDDVALEGVSHFFRELAEEKREGYERLLKMOM

QRGGRALFQDIKKPAEDEV

IgG1_1253A_scFc-N_hFTL

SEQ ID NO: 41

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLM**AS**RTPEVTCVVVDVSHEDPECKFNWYVDGV

EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR

EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFFL

YSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGG

GGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGDKTHTCPPCPAPEL

LGGPSVFLFPPKPKDTLM**AS**RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY

NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM

TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFFLYSKLTVDKSRWQQG

NVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGSS**QIRQNY**

STDVEAAVNSLVNLYLQASYYTSLGLGYFDRDDVALEGVSHFFRELAEEKREGYERLLKMOM

QRGGRALFQDIKKPAEDEV

IgG1_1253V_scFc-N_hFTL

SEQ ID NO: 42

DKTHTCPPCPAPELLGGPSVFLFPPKPKDTLM**AS**RTPEVTCVVVDVSHEDPEVKFNWYVDGV

EVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPR

EPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFFL

YSKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGG

GGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGSGGGDKTHTCPPCPAPEL

LGGPSVFLFPPKPKDTLM**AS**RTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQY

NSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSREEM

TKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFFLYSKLTVDKSRWQQG

NVFCSCVMHEALHNHYTQKSLSLSPGKGGGGSGGGSGGGSGGGSGGGSGGGSS**QIRONY**

STDVEAAVNSLVNLYLQASYYTSLGLGYFDRDDVALEGVSHFFRELAEEKREGYERLLKMOM

QRGGRALFQDIKKPAEDEV

IgG1_LALAP_1253A_scFc-N_hFTL

SEQ ID NO: 43

DKTHTCPPCPAPE**AA**GGPSVFLFPPKPKDTLM**AS**RTPEVTCVVVDVSHEDPEVKFNWYVDG

VEVHNAKTKPREEQYNSTYRVVSVLTVLHQDQLNGKEYKCKVSNKAL**GA**PIEKTISKAKGQPR

REPQVYTLPPSREEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSGSEFF

AKTKPREEOFNSTFRVSVLTVVHODWLNQKEYKCKVSNKGLPAPIEKTISKTKGPREPOV

-continued

YTLPPSREEMTKNQVSLTCLVKGFYPSDISVEWESNGQPENNYKTPPMLDSDGSFFLYSKL

TVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK

His tagged MPER peptide

SEQ ID NO:47

HHHHHHNEQEELLELDKWASLWNWENITNWLWYIKKKK

Example GS repeat

SEQ ID NO:48

GGGS

Example GS repeat

SEQ ID NO:49

GGGGS

EQUIVALENTS/OTHER EMBODIMENTS

[0333] While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of

the invention following, in general, the principles of the invention and including such departures from the present disclosure that come within known or customary practice within the art to which the invention pertains and may be applied to the essential features herein before set forth.

SEQUENCE LISTING

Sequence total quantity: 49

SEQ ID NO: 1 moltype = AA length = 175
FEATURE Location/Qualifiers
source 1..175
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 1
MSSQIRQNYS TDVEAAVNSL VNYLYQASYT YLSLGFYFDR DDVALEGVSH FFRELAEEKR 60
EGYERLLKMQ NQRGGRALFQ DIKKPAEDEV GKTPDAMKAA MALEKKLNQA LLDLHALGSA 120
RTDPLHCDFL ETHFLDEEVK LIKKMGDHLT NLHRLGGPEA GLGEYLFERL TLRHD 175

SEQ ID NO: 2 moltype = AA length = 90
FEATURE Location/Qualifiers
source 1..90
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 2
MSSQIRQNYS TDVEAAVNSL VNYLYQASYT YLSLGFYFDR DDVALEGVSH FFRELAEEKR 60
EGYERLLKMQ NQRGGRALFQ DIKKPAEDEV 90

SEQ ID NO: 3 moltype = AA length = 85
FEATURE Location/Qualifiers
source 1..85
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 3
GKTPDAMKAA MALEKKLNQA LLDLHALGSA RTDPLHCDFL ETHFLDEEVK LIKKMGDHLT 60
NLHRLGGPEA GLGEYLFERL TLRHD 85

SEQ ID NO: 4 moltype = AA length = 85
FEATURE Location/Qualifiers
source 1..85
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 4
GKTPDAMKAA MALEKKLNQA LLDLHALGSA RTDPLHCDFL ETHFLDEEVK LIKKMGDHLT 60
NLHRLGGPEA GLGEYLFERL TLKHD 85

SEQ ID NO: 5 moltype = AA length = 227
FEATURE Location/Qualifiers
source 1..227
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 5
DKTHTCPPCP APELLGGPSV FLFPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAKTK PREEQNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120

-continued

GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTPPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGK		227

SEQ ID NO: 6	moltype = AA	length = 523
FEATURE	Location/Qualifiers	
source	1..523	
	mol_type = protein	
	organism = synthetic	construct

SEQUENCE: 6						
DKTHTCPPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTPPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGKGGG	GSGGGGSGGG	240
GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGDKTH	300
TCPPCPAPPEL	LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	360
HNAKTKPREE	QYNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKALPAPIEK	TISKAKQQR	420
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	480
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGK		523

SEQ ID NO: 7	moltype = AA	length = 226
FEATURE	Location/Qualifiers	
source	1..226	
	mol_type = protein	
	organism = synthetic	construct

SEQUENCE: 7						
KPCPPCKCPA	PNLLGGPSVF	IFPPKIKDVL	MISLSPIVTC	VVVDVSEDDP	DVQISWFVNN	60
VEVHTAQTQT	HREDYNSTLR	VVSALPIQHQ	DWMSGKEFKC	KVNNKDLFAP	IERTISKPKG	120
SVRAPQVYVL	PPPEEEMTKK	QVTLTCMVT	FMPEDIYVEW	TNNGKTELNY	KNTEPVLDS	180
GSYFMYSKLR	VEKKNWVERN	SYSCSVVHEG	LHNHHTTKSF	SRTPGK		226

SEQ ID NO: 8	moltype = AA	length = 515
FEATURE	Location/Qualifiers	
source	1..515	
	mol_type = protein	
	organism = synthetic	construct

SEQUENCE: 8						
VECPPCPAPP	VAGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVQ	FNWYVDGVEV	60
HNAKTKPREE	QFNSTFRVVS	VLTVVHQDWL	NGKEYKCKVS	NKGLPAPIEK	TISKTKGQPR	120
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDISVEWESN	GQPENNYKTT	PPMLDSGGSF	180
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGKGGGSGGG	GSGGGGSGGG	240
GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GGVECPPCPA	300
PPVAGPSVFL	PPPCKPKDTLM	ISRTPEVTCV	VVDVSHEDPE	VQFNWYVDGV	EVHNAKTKPR	360
EEQFNSTFRV	VSVLTVVHQD	WLNKEYKCK	VSNKGLPAPI	EKTISKTKGQ	PREPQVYTL	420
PSREEMTKNQ	VSLTCLVKGF	YPSDISVEWE	SNGQPENNYK	TTPMLDSGD	SFFLYSKLTV	480
DKSRWQQGNV	FSCSVMEAL	HNHYTQKSL	LSPGK			515

SEQ ID NO: 9	moltype = AA	length = 227
FEATURE	Location/Qualifiers	
source	1..227	
	mol_type = protein	
	organism = synthetic	construct

SEQUENCE: 9						
KYGPPCPSCP	APEFLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSQED	PEVQFNWYVD	60
GVEVHNAKTK	PREEQFNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKGLPS	SIEKTISKAK	120
GQPREPQVYT	LPPSQEEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTPPVLDS	180
DGSFFLYSRL	TVDKSRWQEG	NVFSCSVME	ALHNHYTQKS	LSLSLGK		227

SEQ ID NO: 10	moltype = AA	length = 523
FEATURE	Location/Qualifiers	
source	1..523	
	mol_type = protein	
	organism = synthetic	construct

SEQUENCE: 10						
KYGPPCPSCP	APEFLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSQED	PEVQFNWYVD	60
GVEVHNAKTK	PREEQFNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKGLPS	SIEKTISKAK	120
GQPREPQVYT	LPPSQEEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTPPVLDS	180
DGSFFLYSRL	TVDKSRWQEG	NVFSCSVME	ALHNHYTQKS	LSLSLGKGGG	GSGGGGSGGG	240
GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGSGGG	GSGGGGKYGP	300
PCPSCPAPEF	LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSQEDPEVQ	FNWYVDGVEV	360
HNAKTKPREE	QFNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKGLPSSIEK	TISKAKQQR	420
EPQVYTLPPS	QEEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	480
FLYSRLTVDK	SRWQEGNVFS	CSVMHEALHN	HYTQKSLSL	LGK		523

SEQ ID NO: 11	moltype = AA	length = 471
FEATURE	Location/Qualifiers	
source	1..471	

-continued

```

mol_type = protein
organism = synthetic construct

SEQUENCE: 11
QAQLVQSGPE VRKPGTSVKV SCKAPGNTLK TYDLHWVRSV PGQGLQWMGW ISHEGDKKVI 60
VERFKAKVTI DWDRSTNTAY LQLSGLTSGD TAVYYCAKGS KHLRLDYALY DDDGALNWAV 120
DVDYLSNLEF WGQGTAVTVS SASTKGPSVF PLAPSSKSTS GGTAALGCLV KDYPPEPVTV 180
SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ TYICNVNHKP SNTKVDKKVE 240
PKSCDKTHTC PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN 300
WYVDGVEVHN AKTKPREEQY NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI 360
SKAKGQPREP QVYTLPPSRE EMTKNQVSLT CLVKGFPYSD IAVEWESNGQ PENNYKTTTP 420
VLDSGGSFPL YSKLTVDKSR WQQGNVFCSS VMHEALHNNH TQKSLSLSPG K 471

SEQ ID NO: 12      moltype = AA length = 219
FEATURE           Location/Qualifiers
source            1..219
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 12
DFVLTQSPHS LSVTPGESAS ISCKSSHSLI HGDRNNYLAW YVQKPGRSPQ LLIYLASSRA 60
SGVPDRFSGS GSDKDFTLKI SRVETEDVGT YYCMQGRES P WTFGQGTQVD IKRTVAAPSV 120
FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL 180
SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC 219

SEQ ID NO: 13      moltype = AA length = 244
FEATURE           Location/Qualifiers
source            1..244
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 13
QAQLVQSGPE VRKPGTSVKV SCKAPGNTLK TYDLHWVRSV PGQGLQWMGW ISHEGDKKVI 60
VERFKAKVTI DWDRSTNTAY LQLSGLTSGD TAVYYCAKGS KHLRLDYALY DDDGALNWAV 120
DVDYLSNLEF WGQGTAVTVS SASTKGPSVF PLAPSSKSTS GGTAALGCLV KDYPPEPVTV 180
SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ TYICNVNHKP SNTKVDKKVE 240
PKSC 244

SEQ ID NO: 14      moltype = AA length = 533
FEATURE           Location/Qualifiers
source            1..533
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 14
DFVLTQSPHS LSVTPGESAS ISCKSSHSLI HGDRNNYLAW YVQKPGRSPQ LLIYLASSRA 60
SGVPDRFSGS GSDKDFTLKI SRVETEDVGT YYCMQGRES P WTFGQGTQVD IKRTVAAPSV 120
FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL 180
SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGECG GGGSGGGGSG GGGSGGGGSG 240
GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG AQLVQSGPEV 300
RKPPTSVMKVS CKAPGNTLKT YDLHWVRSVP GQGLQWMGWI SHEGDKKVIV ERFKAKVTID 360
WDRSTNTAYL QLSGLTSGDT AVYYCAKGS KHLRLDYALY DDDGALNWAV DVDYLSNLEF 420
GQGTAVTVSS ASTKGPSVFP LAPSSKSTSG GTAAALGCLVK DYFPEPVTVS WNSGALTSGV 480
HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT TYICNVNHKPS NTKVDKKVEP KSC 533

SEQ ID NO: 15      moltype = AA length = 457
FEATURE           Location/Qualifiers
source            1..457
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 15
ADLVQSGAVV KKPQDSVRIS CEAQGYRFPD YIIHWIRRAP GQGPWMGWM NPMGGQVNIP 60
WKFQGRVSM T RDTSIETAF L DLRGLKSDDT AVYYCVRDRS NGSGKRFESS NWFLDLWGRG 120
TAVTIQSAST KGPSVFPLAP SSKSTSGGTA ALGCLVKDYF PEPVTVSWNS GALTSGVHTF 180
PAVLQSSGLY SLSSVTVPS SSLGTQTYIC NVNHKPSNTK VDKKVEPKSC DKTHTCPPCP 240
APELLGGPSV FLPPPKPKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD GVEVHNATK 300
PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK GQPREPQVYT 360
LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSKL 420
TVDKSRWQQG NVFSCSVME ALHNHYTQKS LSLSPGK 457

SEQ ID NO: 16      moltype = AA length = 204
FEATURE           Location/Qualifiers
source            1..204
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 16
QSALTQPRSV SASPGQSVTI SCTGTHNLVS WCQHQPGRAP KLLIYDFNKR PSGVPDRFSG 60
SGSGGTASLT ITGLQDDDDA EYFCWAYEAF GGGTKLTVLG QPKAAPSVTL FPPSSEELQA 120
NKATLVCLIS DFYPGAVTV WKAADSPVKA GVETTPSKQ SNNKYAASSY LSLTPQWKS 180
HRSYSCQVTH EGSTVEKTV PTEC 204

```

-continued

SEQ ID NO: 17 moltype = AA length = 230
 FEATURE Location/Qualifiers
 source 1..230
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 17
 ADLVQSGAVV KKPGDSVRIS CEAQGYRFPD YIIHWIRRAP GQGPEWMGWM NPMGGQVNIP 60
 WKFGQGRVSM RDTSIETAFD DLRGLKSDDT AVYYCVRDRS NGSGRKFESS NWFLDLWGRG 120
 TAVTIQSAST KGPSVFPLAP SSKSTSGGTA ALGCLVKDYF PEPVTVSWNS GALTSGVHTF 180
 PAVLQSSGLY SLSSVVTVPS SSLGTQTYIC NVNHKPSNTK VDKKVEPKSC 230

SEQ ID NO: 18 moltype = AA length = 504
 FEATURE Location/Qualifiers
 source 1..504
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 18
 QSALTQPRSV SASPGQSVTI SCTGTHNLVS WCQHQPGRAP KLLIYDFNKR PSGVPDRFSG 60
 SGSGGTASLT ITGLQDDDDA EYFCWAYEAF GGGTKLTVLG QPKAAPSVTL FPPSSEELQA 120
 NKATLVCLIS DFYPGAVTVA WKADSSPVKA GVETTPSKQ SNNKYAASSY LSLTPEQWKS 180
 HRSYSCQVTH EGSTVEKTVA PTECGGGSG GGGSGGGSG GGGSGGGSG GGGSGGGSG 240
 GGGSGGGSG GGGSGGGSG GGGSGGGSG GGGSGGGSG GGGSGGGSG GGGSGGGSG 300
 RFPDYIIHWI RRAPQGQPEW MGWMNPMGGQ VNI PWKFQGR VSMTRDTSIE TAFDLRLGLK 360
 SDDTAVYYCV RDRSNGSGKR FESSNWFLDL WGRGTAVTIQ SASTKGPSVF PLAPSSKSTS 420
 GGTAALGCLV KDYPPEPVTV SWNSGALTSV VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ 480
 TYICNVNHKP SNTKVDKKVE PKSC 504

SEQ ID NO: 19 moltype = AA length = 461
 FEATURE Location/Qualifiers
 source 1..461
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 19
 EVRLVESGGG LVKPGGSLRL SCSASGFDFD NAWMTWVRQP PGKGLEWVGR ITGPGEWGSV 60
 DYAESVKGRF TISRDNKTNT LYLEMNNVRT EDTGYFFCAR TGKYYDFWSG YPPGEEYFQD 120
 WQGGTLVIVS SASTKGPSVF PLAPSSKSTS GGTAALGCLV KDYPPEPVTV SWNSGALTSV 180
 VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ TYICNVNHKP SNTKVDKKVE PKSCDKHTC 240
 PPCPAPELLG GPSVFLFPPK PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN 300
 AKTKPREEQY NSTYRVSVL TVLHQDWLNG KEYCKVSNK ALPAPIEKTI SKAKGQPREP 360
 QVYTLPPSRD ELTKNQVSLT CLVKGFYPSD IAVEWESNGQ PENNYKTPP VLDSDGSFFL 420
 YSKLTVDKSR WQQGNVPSCS VMHEALHNNH YQKSLSLSPG K 461

SEQ ID NO: 20 moltype = AA length = 214
 FEATURE Location/Qualifiers
 source 1..214
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 20
 SELTQDPAVS VALKQTVTIT CRGDSLRSHY ASWYQKKPGQ APVLLFYGKN NRPSGIPDRF 60
 SGSASGNRAS LTITGAQAE EADYYCSSRD KSGRSLSVFG GGTKLTVLSQ PKAAPSVTLF 120
 PPSSEELQAN KATLVCLISD FYPGAVTVAW KADSSPVKAG VETTPSKQS NNNKYAASSYL 180
 SLTPEQWKSH RSYSCQVTHE GSTVEKTVA TECS 214

SEQ ID NO: 21 moltype = AA length = 234
 FEATURE Location/Qualifiers
 source 1..234
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 21
 EVRLVESGGG LVKPGGSLRL SCSASGFDFD NAWMTWVRQP PGKGLEWVGR ITGPGEWGSV 60
 DYAESVKGRF TISRDNKTNT LYLEMNNVRT EDTGYFFCAR TGKYYDFWSG YPPGEEYFQD 120
 WQGGTLVIVS SASTKGPSVF PLAPSSKSTS GGTAALGCLV KDYPPEPVTV SWNSGALTSV 180
 VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ TYICNVNHKP SNTKVDKKVE PKSC 234

SEQ ID NO: 22 moltype = AA length = 517
 FEATURE Location/Qualifiers
 source 1..517
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 22
 SELTQDPAVS VALKQTVTIT CRGDSLRSHY ASWYQKKPGQ APVLLFYGKN NRPSGIPDRF 60
 SGSASGNRAS LTITGAQAE EADYYCSSRD KSGRSLSVFG GGTKLTVLSQ PKAAPSVTLF 120
 PPSSEELQAN KATLVCLISD FYPGAVTVAW KADSSPVKAG VETTPSKQS NNNKYAASSYL 180
 SLTPEQWKSH RSYSCQVTHE GSTVEKTVA TECEGGGGSG GSGGGGGSG GSGGGGGSG 240
 GSGGGGGSG GSGGGGGSG GSGGGGGSG GSGGGGGSG GSGGGGGSG GSGGGGGSG 300

-continued

```

LRLSCSASGF DFDNAWMTWV RQPPGKGLEW VGRITGPGEG WSVDYAESVK GRFTISRDN 360
KNTLYLEMNN VRTEDTGYF CARTGKYDF WSGYPPGEEY FQDWGQGLTV IVSSASTKGP 420
SVFPLAPSSK STSGGTAALG CLVKDYFPEP VTVSWNSGAL TSGVHTFPFV LQSSGLYSLS 480
SVVTVPSSSL GTQYICNVN HKPSNTKVDK KVEPKSC 517

```

```

SEQ ID NO: 23      moltype = AA length = 452
FEATURE           Location/Qualifiers
source            1..452
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 23
QVQLQSGPE VVKPGASVKM SCKASGYTFT SYVIHWVRQK PGQGLDWIGY INPYNDGTDY 60
DEKFKGKATL TSDTSTSTAY MELSSLRSED TAVYYCAREK DNYATGAWFA YWGQGLTVTV 120
SSASTKGPSV FPLAPCSRST SESTAALGCL VKDYFPEPVT VSWNSGALTS GVHTFPAVLQ 180
SSGLYSLSLV VTPSSSLGT KTYTCNVNDHK PSNTKVDKRV ESKYCDKTHT CPPCPAPELL 240
GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPREEQ 300
YNSTYRVVSV LTVLHQDWLNL GKEYKCKVSN KALPAPIEKT ISKAKGQPRE PQVYTLPPSR 360
EEMTKNQVSL TCVKGFPYPS DIAVEWESNG QPENNYKTP PVLDSDGSFF LYSKLTVDKS 420
RWQQGNVFCF SVMHEALHNNH YTKQSLSLSP GK 452

```

```

SEQ ID NO: 24      moltype = AA length = 219
FEATURE           Location/Qualifiers
source            1..219
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 24
DIVMTQSPDS LAVSLGERVT MNCKSSQSLL YSTNQKNYLA WYQQKPGQSP KLLIYWASTR 60
ESGVPDRFSG SGSGTDFTLT ISSVQAEDVA VYYCQYYYSY RTFGGGTKLE IKRTVAAPSV 120
FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL 180
SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGEC 219

```

```

SEQ ID NO: 25      moltype = AA length = 225
FEATURE           Location/Qualifiers
source            1..225
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 25
QVQLQSGPE VVKPGASVKM SCKASGYTFT SYVIHWVRQK PGQGLDWIGY INPYNDGTDY 60
DEKFKGKATL TSDTSTSTAY MELSSLRSED TAVYYCAREK DNYATGAWFA YWGQGLTVTV 120
SSASTKGPSV FPLAPCSRST SESTAALGCL VKDYFPEPVT VSWNSGALTS GVHTFPAVLQ 180
SSGLYSLSLV VTPSSSLGT KTYTCNVNDHK PSNTKVDKRV ESKYC 225

```

```

SEQ ID NO: 26      moltype = AA length = 514
FEATURE           Location/Qualifiers
source            1..514
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 26
DIVMTQSPDS LAVSLGERVT MNCKSSQSLL YSTNQKNYLA WYQQKPGQSP KLLIYWASTR 60
ESGVPDRFSG SGSGTDFTLT ISSVQAEDVA VYYCQYYYSY RTFGGGTKLE IKRTVAAPSV 120
FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL 180
SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGECG GGGSGGGGSG GGGSGGGGSG 240
GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG VQLQSGPEV 300
VVKPGASVKM SCKASGYTFT SYVIHWVRQK PGQGLDWIGY INPYNDGTDY DEKFKGKATLT 360
SDTSTSTAYM ELSSLRSED AVYYCAREK DNYATGAWFA WQGLTVTVS SASTKGPSVF 420
PLAPSSKSTS GGTAAALGCLV KDYFPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSLVV 480
TVPSSSLGTQ TYICNVNHPK SNTKVDKKVE PKSC 514

```

```

SEQ ID NO: 27      moltype = AA length = 735
FEATURE           Location/Qualifiers
source            1..735
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 27
DFVLTQSPHS LSVTPGESAS ISCKSSSLI HGDRNNYLAW YVQKPGRSPQ LLIYLASSRA 60
SGVPDRFSGS GSDKDFTLTKI SRVETEDVGT YYCMQGRESF WTFGQGTQVD IKRTVAAPSV 120
FIFPPSDEQL KSGTASVVCL LNNFYPREAK VQWKVDNALQ SGNSQESVTE QDSKDYSTYSL 180
SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGECG GGGSGGGGSG GGGSGGGGSG 240
GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG AQLVQSGPEV 300
RKPGETSVKVS CKAPGNTLKT YDLHWVRVSP GQGLQWMGIW SHEGDKKIV ERFKAKVTID 360
WDRSTNTAYL QLSGLTSGDT AVYYCAKGSK HRLRDYALYD DDGALNWAVD VDYLSNLEFV 420
GQGTAVTVSS ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV 480
HTFPAVLQSS GLYSLSSVVT VPSSSLGTQT YICNVNHPKS NTKVDKKVEP KSCDGGGGSG 540
GGGSGGGGSG GGGSGGGGSG GSSQIRQNYT TDVEAAVNSL NLYLQASYT YLSLGFYFDR 600
DDVALEGVSH FPRELAEEKR EGYERLLKMQ NQRGGRALFQ DIKPAEDEV GKTDPAMKAA 660
MALEKKLNQA LLDLHALGSA RTDPHLCDFL ETHFLDEEVK LIKKMGDHLT NLHRLGGPEA 720

```


-continued

GLGEYLFERL TLRHD 735

SEQ ID NO: 28 moltype = AA length = 616
 FEATURE Location/Qualifiers
 source 1..616
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 28
 QSALTQPRSV SASPGQSVTI SCTGTHNLVS WCQHQPGRAP KLLIYDFNKR PSGVPDRFSG 60
 SGSGGTASLT ITGLQDDDDA EYFCWAYEAF GGGTKLTVLG QPKAAPSVTL FPPSSEELQA 120
 NKATLVCLIS DFYPGAIVTA WKADSSPVKA GVETTTPSKQ SNNKYAASSY LSLTPEQWKS 180
 HRSYSCQVTH EGSTVEKTVA PTECGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG 240
 GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGADLVQS GAVVKKPGDS VRISCEAQGY 300
 RFPDYIIHWI RRAPQGQPEW MGWMNPMGGQ VNIPWKFQGR VSMTRDTSIE TAFLDLRGLK 360
 SDDTAVYYCV RDRSNGSGKR FESSNWFLDL WGRGTAVTIQ SASTKGPSVF PLAPSSKSTS 420
 GGTAALGCLV KDYPPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ 480
 TYICNVNHPK SNTKVDKKVE PKSCGGGGSG GGGSGGGGSG GGGSGGGGSG GGGTPDAMKA 540
 AMALEKKLNQ ALLDLHALGS ARTDPLHCLD LETHFLDEEV KLIKMGDHL TNLHRLGGPE 600
 AGLGEYLFER LTLRHD 616

SEQ ID NO: 29 moltype = AA length = 620
 FEATURE Location/Qualifiers
 source 1..620
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 29
 QSALTQPRSV SASPGQSVTI SCTGTHNLVS WCQHQPGRAP KLLIYDFNKR PSGVPDRFSG 60
 SGSGGTASLT ITGLQDDDDA EYFCWAYEAF GGGTKLTVLG QPKAAPSVTL FPPSSEELQA 120
 NKATLVCLIS DFYPGAIVTA WKADSSPVKA GVETTTPSKQ SNNKYAASSY LSLTPEQWKS 180
 HRSYSCQVTH EGSTVEKTVA PTECGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG 240
 GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGADLVQS GAVVKKPGDS VRISCEAQGY 300
 RFPDYIIHWI RRAPQGQPEW MGWMNPMGGQ VNIPWKFQGR VSMTRDTSIE TAFLDLRGLK 360
 SDDTAVYYCV RDRSNGSGKR FESSNWFLDL WGRGTAVTIQ SASTKGPSVF PLAPSSKSTS 420
 GGTAALGCLV KDYPPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSSVV TVPSSSLGTQ 480
 TYICNVNHPK SNTKVDKKVE PKSCGGGGSG GGGSGGGGSG GGGSGGGGSG GSSQIRQNY 540
 TDVEAAVNSL VNYLQASYT YLSLGFYFDR DDVALEGVSH FFRELAEEKR EGYERLLKMK 600
 NQRGGRALFQ DIKPAEDEW 620

SEQ ID NO: 30 moltype = AA length = 630
 FEATURE Location/Qualifiers
 source 1..630
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 30
 SELTQDPAVS VALKQTVTIT CRGDSLRSHY ASWYQKKPGQ APVLLFYGKN NRPSGIPDRF 60
 SGASAGNRAS LTITGAQAEAD EADYYCSSRD KSGSRLSVFG GGTCLTVLSQ PKAAPSVTLF 120
 PPSSEELQAN KATLVCLISD FYPGAVTVAW KADSSPVKAG VETTTPSKQS NNKYAASSYL 180
 SLTPEQWKSH RSYSCQVTHE GSTVEKTVA TEGCGGGGSG GGGSGGGGSG GGGSGGGGSG 240
 GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSEVRLVES GGLVVKPGGS 300
 LRLSCSASGF DFDNAWMTWV RQPPGKLEW VGRITGPGEW WSDVYAESVK GRFTISRDN 360
 KNTLYLEMMN VRTEDTGYFF CARTGKYDF WSGYPPGEEY FQDWGGQTLV IVSSASTKGP 420
 SVFPLAPSSK STSGGTAALG CLVKDYFPEP VTVSWNSGAL TSGVHTFPAV LQSSGLYSL 480
 SVVTVPSSSL GTQTYICNVN HKPSNTKVDK KVEPKSCGGG GSGGGGSGGG GSGGGGSGGG 540
 GSGGGGKTPD AMKAAMALEK KLNQALLDLH ALGSARTDPH LCDPLETHFL DEEVKLIKMK 600
 GDHLTNLHRL GGPEAGLGEY LFERLTLRHD 630

SEQ ID NO: 31 moltype = AA length = 627
 FEATURE Location/Qualifiers
 source 1..627
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 31
 DIVMTQSPDS LAVSLGERVT MNCKSSQSLL YSTNQKNYLA WYQKPGQSP KLLIYWASTR 60
 ESGVPDRFSG SSGTDFTLT ISSVQAEDVA VYCCQYYSY RTFGGGTKLE IKRTVAAPSV 120
 FIFPPSDEQL KSGTASVCL LNNFYPREAK VQWKVDNALQ SGNQSQESVTE QDSKDYSL 180
 SSTLTLSKAD YEKHKVYACE VTHQGLSSPV TKSFNRGECG GGGSGGGGSG GGGSGGGGSG 240
 GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG GGGSGGGGSG VQLQSGPEV 300
 VKPGASVKMS CKASGYFTS YVIHWVRQKP GQGLDWIGYI NPYNDGTDYD EKFKGKATLT 360
 SDTSTSTAYM ELSSLRSEDV AVYYCAREKD NYATGAWFAY WQGGTLVTVS SASTKGPSVF 420
 PLAPSSKSTS GGTAALGCLV KDYPPEPVTV SWNSGALTSG VHTFPAVLQS SGLYSLSSVV 480
 TVPSSSLGTQ TYICNVNHPK SNTKVDKKVE PKSCGGGGSG GGGSGGGGSG GGGSGGGGSG 540
 GGGKTPDAMK AAMALEKKLN QALLDLHALG SARTDPLHCLD FLETHFLDEE VKLIKMGDHL 600
 LTNLHRLGGP EAGLGEYLFE RLTLRHD 627

SEQ ID NO: 32 moltype = AA length = 883
 FEATURE Location/Qualifiers

-continued

source 1..883
mol_type = protein
organism = synthetic construct

SEQUENCE: 32

SELTQDPAVS	VALKQTVTIT	CRGDSLRSY	ASWYQKKPGQ	APVLLFYGKN	NRPSGIPDRF	60
SGSASGNRAS	LTITGAQAE	EADYYCSSRD	KSGSRLSVFG	GGTKLTVLSQ	PKAAPSVTLF	120
PPSSEELQAN	KATLVCLISD	FYPGAVTVAW	KADSSPVKAG	VETTTPSKQS	NNKYAASSYL	180
SLTPEQWKSH	RSYSCQVTHE	GSTVEKTVAP	TECGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	240
GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSEVRLVES	GGGLVKPGGS	300
LRLSCSASGF	DFDNAWMTWV	RQPPGKGLEW	VGRITGPGE	WSVDYAESVK	GRFTISRDN	360
KNTLYLEMN	VRTEDTGYFF	CARTGKYDF	WSGYPPGEEY	FQDWGQGTLV	IVSSASTKGP	420
SVFPLAPSSK	STSGGTAALG	CLVKDYFPEP	VTVSWNSGAL	TSGVHTFP	LQSSGLYSL	480
SVVTVPSSSL	GTQTYICNVN	HKPSNTKV	KVEPKSCGGG	GGSGGGGSGG	GGSGGGGSGG	540
GGSGGGKTPD	AMKAAMALEK	KLNLQALLDLH	ALGSARTDPH	LCDFLETHFL	DEEVKLIK	600
GDHLTNLHRL	GGPEAGLGEY	LFERLTLRHD	GGSGGGGSGG	GGSGGGGSGG	GGSGGDKTH	660
TCPPCPAPEL	LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	720
HNAKTKPRE	QYNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKALPAPIEK	TISKAKGQPR	780
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	840
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGK		883

SEQ ID NO: 33 moltype = AA length = 883
FEATURE Location/Qualifiers
source 1..883
mol_type = protein
organism = synthetic construct

SEQUENCE: 33

SELTQDPAVS	VALKQTVTIT	CRGDSLRSY	ASWYQKKPGQ	APVLLFYGKN	NRPSGIPDRF	60
SGSASGNRAS	LTITGAQAE	EADYYCSSRD	KSGSRLSVFG	GGTKLTVLSQ	PKAAPSVTLF	120
PPSSEELQAN	KATLVCLISD	FYPGAVTVAW	KADSSPVKAG	VETTTPSKQS	NNKYAASSYL	180
SLTPEQWKSH	RSYSCQVTHE	GSTVEKTVAP	TECGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	240
GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSEVRLVES	GGGLVKPGGS	300
LRLSCSASGF	DFDNAWMTWV	RQPPGKGLEW	VGRITGPGE	WSVDYAESVK	GRFTISRDN	360
KNTLYLEMN	VRTEDTGYFF	CARTGKYDF	WSGYPPGEEY	FQDWGQGTLV	IVSSASTKGP	420
SVFPLAPSSK	STSGGTAALG	CLVKDYFPEP	VTVSWNSGAL	TSGVHTFP	LQSSGLYSL	480
SVVTVPSSSL	GTQTYICNVN	HKPSNTKV	KVEPKSCGGG	GGSGGGGSGG	GGSGGGGSGG	540
GGSGGGKTPD	AMKAAMALEK	KLNLQALLDLH	ALGSARTDPH	LCDFLETHFL	DEEVKLIK	600
GDHLTNLHRL	GGPEAGLGEY	LFERLTLRHD	GGSGGGGSGG	GGSGGGGSGG	GGSGGDKTH	660
TCPPCPAPEL	LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	720
HNAKTKPRE	QYNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKALPAPIEK	TISKAKGQPR	780
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	840
FLYSKLTVDK	SRWQQGNVFS	CSVLHEALHS	HYTQKSLSL	PGK		883

SEQ ID NO: 34 moltype = AA length = 639
FEATURE Location/Qualifiers
source 1..639
mol_type = protein
organism = synthetic construct

SEQUENCE: 34

DKHTCPCPCP	APELLGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAKTK	PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALPA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTPPVLD	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCVMHE	ALHNHYTQKS	LSLSPGKGGG	GGSGGGGSGG	240
GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGDKTH	300
TCPPCPAPEL	LGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	360
HNAKTKPRE	QYNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKALPAPIEK	TISKAKGQPR	420
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	480
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGKGGGGSGG	GGSGGGGSGG	540
GGSGGGGSGG	SSQIRQNYST	DVEAAVNSLV	NLYLQASYTY	LSLGFYFDRD	DVALEGVSHF	600
FRELAEEKRE	GYERLLKMQN	QRGGRALFQD	IKKPAEDEV			639

SEQ ID NO: 35 moltype = AA length = 631
FEATURE Location/Qualifiers
source 1..631
mol_type = protein
organism = synthetic construct

SEQUENCE: 35

VECPPCPAPP	VAGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVQ	FNWYVDGVEV	60
HNAKTKPRE	QFNSTFRVVS	VLTVVHQDWL	NGKEYKCKVS	NKGLPAPIEK	TISKTKGQPR	120
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDISVEWESN	GQPENNYKTT	PPMLDSGGSF	180
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGKGGGGSGG	GGSGGGGSGG	240
GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSGGGGSGG	GGSECPPCPA		300
PPVAGPSVFL	PPPCKPKDTLM	ISRTPEVTCV	VVDVSHEDPE	VQFNWYVDGV	EVHNAKTKPR	360
EEQFNSTFRV	VSVLTVVHQD	WLNKEYKCK	VSNKGLPAPI	EKTISKTKGQ	PREPQVYTL	420
PSREEMTKNQ	VSLTCLVKGF	YPSDISVEWE	SNGQPENNYK	TTPPMLDSG	SFFLYSKLTV	480
DKSRWQQGNV	FSCSVMEAL	HNHYTQKSL	LSPGKGGGGS	GGGGGGGGGS	GGGGGGGGGS	540
GGSSQIRQNY	STDVEAAVNS	LVNLYLQASY	TYLSLGFYFD	RDDVALEGV	HFRELAEEK	600

-continued

 REGYERLLKM QNQRGGRALF QDIKKPAEDE W 631

SEQ ID NO: 36 moltype = AA length = 639
 FEATURE Location/Qualifiers
 source 1..639
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 36
 KYGPPCPSPC APEFLGGPSV FLFPKPKKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD 60
 GVEVHNATK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK 120
 GQPREPQVYT LPPSQEEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPPVLDS 180
 DGSFFLYSRL TVDKSRWQEG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
 GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGKYP 300
 PCPSCPAPAF LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSQEDPEVQ FNWYVDGVEV 360
 HNAKTKPREE QFNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKGLPSSIEK TISKAKGQPR 420
 EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
 FLYSRLTVDK SRWQEGNVFS CSVMHEALHN HYTKQSLSL LSGGGGSGGG GSGGGGSGGG 540
 GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
 FRELAEEKRE GYERLLKMQN QRGRALFQD IKKPAEDEW 639

SEQ ID NO: 37 moltype = AA length = 639
 FEATURE Location/Qualifiers
 source 1..639
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 37
 DKHTCPCPPC APELLGGPSV FLFPKPKKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNATK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALGA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPPVLDS 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
 GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
 TCPPCPAPEL LGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
 HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALGAPIEK TISKAKGQPR 420
 EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
 FLYSKLTVDK SRWQGNVFS CSVMHEALHN HYTKQSLSL PGKGGGSGGG GSGGGGSGGG 540
 GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
 FRELAEEKRE GYERLLKMQN QRGRALFQD IKKPAEDEW 639

SEQ ID NO: 38 moltype = AA length = 639
 FEATURE Location/Qualifiers
 source 1..639
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 38
 DKHTCPCPPC APEAAGGPSV FLFPKPKKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNATK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPPVLDS 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
 GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
 TCPPCPAPEA AGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
 HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALGAPIEK TISKAKGQPR 420
 EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
 FLYSKLTVDK SRWQGNVFS CSVMHEALHN HYTKQSLSL PGKGGGSGGG GSGGGGSGGG 540
 GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
 FRELAEEKRE GYERLLKMQN QRGRALFQD IKKPAEDEW 639

SEQ ID NO: 39 moltype = AA length = 639
 FEATURE Location/Qualifiers
 source 1..639
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 39
 DKHTCPCPPC APEAAGGPSV FLFPKPKKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
 GVEVHNATK PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALGA PIEKTISKAK 120
 GQPREPQVYT LPPSREEMTK NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPPVLDS 180
 DGSFFLYSKL TVDKSRWQGG NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
 GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
 TCPPCPAPEA AGGPSVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
 HNAKTKPREE QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALGAPIEK TISKAKGQPR 420
 EPQVYTLPPS REEMTKNQVS LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
 FLYSKLTVDK SRWQGNVFS CSVMHEALHN HYTKQSLSL PGKGGGSGGG GSGGGGSGGG 540
 GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
 FRELAEEKRE GYERLLKMQN QRGRALFQD IKKPAEDEW 639

SEQ ID NO: 40 moltype = AA length = 639
 FEATURE Location/Qualifiers

-continued

```

source                1..639
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 40
DKTHTCPCPCP APELLGGPSV FLFPKPKKDT LMISRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAATK  PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT  LPPSREEMTK  NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL  TVDKSRWQQG  NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
TCPPCPAPEL  LGGPSVFLFP  PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
HNAATKPREE  QYNSTYRVVS  VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR 420
EPQVYTLPPS  REEMTKNQVS  LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
FLYSKLTVDK  SRWQQGNVFS  CSVMHEALHN HTQKSLSLs PGKGGGSGGG GSGGGGSGGG 540
GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
FRELAEKRE  GYERLLKMQN  QRGGRALFQD IKKPAEDEV 639

SEQ ID NO: 41          moltype = AA length = 639
FEATURE               Location/Qualifiers
source                1..639
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 41
DKTHTCPCPCP APELLGGPSV FLFPKPKKDT LMASRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAATK  PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT  LPPSREEMTK  NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL  TVDKSRWQQG  NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
TCPPCPAPEL  LGGPSVFLFP  PKPKDTLMAS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
HNAATKPREE  QYNSTYRVVS  VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR 420
EPQVYTLPPS  REEMTKNQVS  LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
FLYSKLTVDK  SRWQQGNVFS  CSVMHEALHN HTQKSLSLs PGKGGGSGGG GSGGGGSGGG 540
GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
FRELAEKRE  GYERLLKMQN  QRGGRALFQD IKKPAEDEV 639

SEQ ID NO: 42          moltype = AA length = 639
FEATURE               Location/Qualifiers
source                1..639
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 42
DKTHTCPCPCP APELLGGPSV FLFPKPKKDT LMVSRTPPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAATK  PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALPA PIEKTISKAK 120
GQPREPQVYT  LPPSREEMTK  NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL  TVDKSRWQQG  NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
TCPPCPAPEL  LGGPSVFLFP  PKPKDTLMVS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
HNAATKPREE  QYNSTYRVVS  VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR 420
EPQVYTLPPS  REEMTKNQVS  LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
FLYSKLTVDK  SRWQQGNVFS  CSVMHEALHN HTQKSLSLs PGKGGGSGGG GSGGGGSGGG 540
GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
FRELAEKRE  GYERLLKMQN  QRGGRALFQD IKKPAEDEV 639

SEQ ID NO: 43          moltype = AA length = 639
FEATURE               Location/Qualifiers
source                1..639
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 43
DKTHTCPCPCP APEAAGGPSV FLFPKPKKDT LMASRTPEVT CVVVDVSHED PEVKFNWYVD 60
GVEVHNAATK  PREEQYNSTY RVVSVLTVLH QDWLNGKEYK CKVSNKALGA PIEKTISKAK 120
GQPREPQVYT  LPPSREEMTK  NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTPPVLD 180
DGSFFLYSKL  TVDKSRWQQG  NVFSCSVMHE ALHNHYTQKS LSLSPGKGGG GSGGGGSGGG 240
GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGSGGG GSGGGGDKTH 300
TCPPCPAPEA  AGGPSVFLFP  PKPKDTLMAS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV 360
HNAATKPREE  QYNSTYRVVS  VLTVLHQDWL NGKEYKCKVS NKALGAPIEK TISKAKGQPR 420
EPQVYTLPPS  REEMTKNQVS  LTCLVKGFYP SDIAVEWESN GQPENNYKTT PPVLDSDGSF 480
FLYSKLTVDK  SRWQQGNVFS  CSVMHEALHN HTQKSLSLs PGKGGGSGGG GSGGGGSGGG 540
GSGGGGSGGG SSQIRQNYST DVEAAVNSLV NLYLQASYTY LSLGFYFDRD DVALEGVSHF 600
FRELAEKRE  GYERLLKMQN  QRGGRALFQD IKKPAEDEV 639

SEQ ID NO: 44          moltype = AA length = 639
FEATURE               Location/Qualifiers
source                1..639
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 44

```

-continued

DKTHTCPPCP	APEAAGGPSV	FLFPPKPKDT	LMISRTPEVT	CVVVDVSHED	PEVKFNWYVD	60
GVEVHNAATK	PREEQYNSTY	RVVSVLTVLH	QDWLNGKEYK	CKVSNKALGA	PIEKTISKAK	120
GQPREPQVYT	LPPSREEMTK	NQVSLTCLVK	GFYPSDIAVE	WESNGQPENN	YKTTTPPVLDS	180
DGSFFLYSKL	TVDKSRWQQG	NVFSCSVME	ALHNHYTQKS	LSLSPGKGGG	GGGGGGGGGG	240
GGGGGGGGGG	GGGGGGGGGG	GGGGGGGGGG	GGGGGGGGGG	GGGGGGGGGG	GGGGGGDKTH	300
TCPPCPAPEA	AGGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVK	FNWYVDGVEV	360
HNAATKPREE	QYNSTYRVVS	VLTVLHQDWL	NGKEYKCKVS	NKALGAPIEK	TISKAKGQPR	420
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDIAVEWESN	GQPENNYKTT	PPVLDSGGSF	480
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGKGGGGSGG	GGSGGGGGSG	540
GGSGGGGGSG	SSQIRQNYST	DVEAAVNSLV	NLYLQASYTY	LSLGFYFDRD	DVALEGVSHF	600
FRELAEEKRE	GYERLLKMQN	QRGGRALFQD	IKKPAEDEW			639

SEQ ID NO: 45 moltype = AA length = 631
 FEATURE Location/Qualifiers
 source 1..631
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 45

VECPPCPAPP	VAGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVQ	FNWYVDGVEV	60
HNAATKPREE	QFNSTFRVVS	VLTVVHQDWL	NGKEYKCKVS	NKGLGAPIEK	TISKTKGQPR	120
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDISVEWESN	GQPENNYKTT	PPMLDSGGSF	180
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGKGGGGSGG	GGSGGGGGSG	240
GGSGGGGGSG	GGSGGGGGSG	GGSGGGGGSG	GGSGGGGGSG	GGSGGGGGSG	GGVECPPCPA	300
PPVAGPSVFL	PPPCKPKDTL	ISRTPEVTCV	VVDVSHEDPE	VQFNWYVDGV	EVHNAATKPR	360
EEQFNSTFRV	VSVLTVVHQD	WLNKKEYKCK	VSNKGLGAPI	EKTISKTKGQ	PREPQVYTL	420
PSREEMTKNQ	VSLTCLVKGF	YPSDISVEWE	SNGQPENNYK	TTPPMLDSDG	SFFLYSKLTV	480
DKSRWQQGNV	PSCSVMEAL	HNHYTQKSL	LSPGKGGGGG	GGGGSGGGGG	GGGGSGGGGG	540
GGSSQIRQNY	STDVEAAVNS	LVNLYLQASY	TYLSLGFYFD	RDDVALEGVS	HFFRELAEEK	600
REGYERLLKM	QNQRGGRALF	QDIKKPAEDE	W			631

SEQ ID NO: 46 moltype = AA length = 223
 FEATURE Location/Qualifiers
 source 1..223
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 46

VECPPCPAPP	VAGPSVFLFP	PKPKDTLMIS	RTPEVTCVVV	DVSHEDPEVQ	FNWYVDGVEV	60
HNAATKPREE	QFNSTFRVVS	VLTVVHQDWL	NGKEYKCKVS	NKGLGAPIEK	TISKTKGQPR	120
EPQVYTLPPS	REEMTKNQVS	LTCLVKGFYP	SDISVEWESN	GQPENNYKTT	PPMLDSGGSF	180
FLYSKLTVDK	SRWQQGNVFS	CSVMHEALHN	HYTQKSLSL	PGK		223

SEQ ID NO: 47 moltype = AA length = 37
 FEATURE Location/Qualifiers
 source 1..37
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 47

HHHHHHNEQE	LLELDKWASL	WNWFNITNWL	WYIKKKK			37
------------	------------	------------	---------	--	--	----

SEQ ID NO: 48 moltype = AA length = 4
 FEATURE Location/Qualifiers
 source 1..4
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 48

GGGS						4
------	--	--	--	--	--	---

SEQ ID NO: 49 moltype = AA length = 5
 FEATURE Location/Qualifiers
 source 1..5
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 49

GGGS						5
------	--	--	--	--	--	---

1. A self-assembled polypeptide complex comprising
 - (a) a plurality of first fusion polypeptides, each first fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises a Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, and
 - (b) a plurality of second fusion polypeptides, each second fusion polypeptide comprising (1) an antigen-binding antibody fragment linked to (2) a nanocage monomer or subunit thereof.
2. The self-assembled polypeptide complex of claim 1, wherein (1) if the Fc polypeptide is an IgG1 Fc polypeptide, the antigen-binding fragment is not a Fab fragment that binds to SARS-CoV-2; and/or (2) if the nanocage monomer is a mouse ferritin monomer and the Fc polypeptide is a mouse IgG2a Fc polypeptide, the antigen-binding antibody fragment is not a Fab fragment that binds to CD19.
3. The self-assembled polypeptide complex of claim 1, wherein the nanocage monomer is a ferritin monomer.
4. The self-assembled polypeptide complex of claim 3, wherein the ferritin monomer is a ferritin light chain.
5. The self-assembled polypeptide complex of claim 4, which does not comprise any ferritin heavy chains or subunits of ferritin heavy chains.
6. (canceled)
7. The self-assembled polypeptide complex of claim 1, wherein the Fc polypeptide is an IgG1 Fc polypeptide.
8. (canceled)
9. The self-assembled polypeptide complex of claim 1, wherein the Fc polypeptide is a single-chain Fc (scFc).
10. (canceled)
14. The self-assembled polypeptide complex of claim 1, wherein the one or more mutations comprise a mutation or set of mutations associated with altered binding to FcRn.
15. The self-assembled polypeptide complex of claim 14, wherein the mutation or set of mutations comprises a mutation at one or more of the following residues: M252, I253, S254, T256, K288, M428, and N434, or combinations thereof, wherein numbering is according to the EU index.
16. The self-assembled polypeptide complex of claim 15, wherein the mutation or set of mutations comprises mutations at M428 and N434, wherein numbering is according to the EU index.
17. The self-assembled polypeptide complex of claim 16, wherein the mutation or set of mutations comprises M428L and N434S mutations, wherein numbering is according to the EU index.
18. The self-assembled polypeptide complex of claim 14, wherein the altered binding to FcRn is decreased binding to FcRn.
19. The self-assembled polypeptide complex of claim 18, wherein the mutation or set of mutations associated with decreased binding to FcRn is selected from the group consisting of I253A, I253V, and K288A, and combinations thereof, wherein numbering is according to the EU index.
20. The self-assembled polypeptide complex of claim 1, wherein the one or more mutations comprise a mutation or set of mutations associated with altered effector function.
21. The self-assembled polypeptide complex of claim 20, wherein the Fc polypeptide is an IgG1 Fc polypeptide, and wherein the mutation or set of mutations comprises a mutation at one or more of the following residues: L234, L235,

G236, G237, P329, and A330, or combinations thereof, wherein numbering is according to the EU index.

22. The self-assembled polypeptide complex of claim 20, wherein the altered effector function is decreased effector function.

23. The self-assembled polypeptide complex of claim 22, wherein the mutation or set of mutations associated with decreased effector function is selected from the group consisting of LALA (L234A/L235A), LALAP (L234A/L235A/P329G), G236R, G237A, and A330L, wherein numbering is according to the EU index.

24.-53. (canceled)

54. A method comprising administering a composition comprising the self-assembled polypeptide complex of claim 1 to a mammalian subject.

55.-63. (canceled)

64. A fusion polypeptide comprising (1) an Fc polypeptide linked to (2) a nanocage monomer or subunit thereof, wherein the Fc polypeptide comprises a Fc chain having one or more mutations relative to a reference Fc chain of the same Ig class, wherein the one or more mutations comprise a mutation or set of mutations associated with altered binding to FcRn and/or with altered effector function.

65. The fusion polypeptide of claim 64, wherein the nanocage monomer is a ferritin monomer.

66. The fusion polypeptide of claim 65, wherein the ferritin monomer is a ferritin light chain.

67. The fusion polypeptide of claim 66, which does not comprise any ferritin heavy chains or subunits of ferritin heavy chains.

68. The fusion polypeptide of claim 65, wherein the ferritin monomer is a human ferritin.

69. The fusion polypeptide of claim 64, wherein the Fc polypeptide is an IgG1 Fc polypeptide.

70. (canceled)

71. The fusion polypeptide of claim 64, wherein the Fc polypeptide is a scFc.

72. The fusion polypeptide of claim 6, wherein the mutation or set of mutations comprises a mutation at one or more of the following residues: M252, I253, S254, T256, K288, M428, and N434, or combinations thereof, wherein numbering is according to the EU index.

73. The fusion polypeptide of claim 64, wherein the altered binding to FcRn is decreased binding to FcRn.

74. The fusion polypeptide of claim 73, wherein the mutation or set of mutations associated with decreased binding to FcRn is selected from the group consisting of I253A, I253V, and K288A, and combinations thereof, wherein numbering is according to the EU index.

75. The fusion polypeptide of claim 64, wherein the Fc polypeptide is an IgG1 Fc polypeptide, and wherein the mutation or set of mutations comprises a mutation at one or more of the following residues: L234, L235, G236, G237, P329, and A330, or combinations thereof, wherein numbering is according to the EU index.

76. The fusion polypeptide of claim 64, wherein the altered effector function is decreased effector function.

77. The fusion polypeptide of claim 76, wherein the mutation or set of mutations associated with decreased

effector function is selected from the group consisting of LALA (L234A/L235A), LALAP (L234A/L235A/P329G), G236R, G237A, and A330L, wherein numbering is according to the EU index.

78.-80. (canceled)

* * * * *