



US 20250277036A1

(19) **United States**

(12) **Patent Application Publication**
MONSEUR et al.

(10) **Pub. No.: US 2025/0277036 A1**

(43) **Pub. Date: Sep. 4, 2025**

(54) **ANTI-CD8 ALPHA DEPLETING ANTIBODIES**

A61P 9/10 (2006.01)

(71) Applicant: **POLYGON THERAPEUTICS,**
Levallois-perret (FR)

G01N 33/68 (2006.01)

(52) **U.S. CL.**

(72) Inventors: **Tatiana MONSEUR,** Levallois-perret
(FR); **Mohamed ABOU ALL,**
Levallois-perret (FR)

CPC *C07K 16/2815* (2013.01); *A61P 9/10*
(2018.01); *G01N 33/6854* (2013.01); *A61K*
2039/505 (2013.01); *C07K 2317/33* (2013.01);
C07K 2317/73 (2013.01); *C07K 2317/92*
(2013.01); *G01N 2333/70517* (2013.01)

(73) Assignee: **POLYGON THERAPEUTICS,**
Levallois-perret (FR)

(21) Appl. No.: **19/209,867**

(57)

ABSTRACT

(22) Filed: **May 16, 2025**

Related U.S. Application Data

(63) Continuation of application No. PCT/EP2023/
082168, filed on Nov. 17, 2023.

Novel anti-CD8 α (cluster of differentiation 8 alpha) anti-
bodies and antigen-binding fragments thereof, a fusion
protein including the anti-CD8 α antibodies and antigen-
binding fragments thereof, and a nucleic acid molecule
encoding the anti-CD8 α antibodies and antigen-binding
fragments thereof. Also, pharmaceutical compositions
including the anti-CD8 α antibodies and antigen-binding
fragments thereof. Further, methods using the anti-CD8 α
antibodies and antigen-binding fragments thereof for treat-
ing CD8-related diseases and in vitro methods in which
CD8 α -expressing cells are depleted in a sample or in which
CD8 α is detected or quantified in a sample.

(30) Foreign Application Priority Data

Nov. 17, 2022 (EP) 22306689.5


Specification includes a Sequence Listing.

Publication Classification

(51) **Int. Cl.**
C07K 16/28 (2006.01)
A61K 39/00 (2006.01)

Human IgG1 antibody clones

Human CD8 α






FIG. 1A




FIG. 1B




FIG. 1C




FIG. 2A




FIG. 2B







FIG. 2C


HPB-ALL cells




Irrelevant IgG1

Commercial anti-CD8 α
antibody



Primary CD8⁺ T cells

Irrelevant IgG1

Commercial anti-CD8 α
antibody

Control WT mice


Transgenic mice
expressing hCD8 α transgene

FIG. 6A




FIG. 6B

FIG. 7A




FIG. 7B




FIG. 7C




FIG. 8A




FIG. 8B







FIG. 8C








FIG. 10A




FIG. 10B




FIG. 10C




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 12C




FIG. 13A




FIG. 13B




FIG. 13C




FIG. 14

ANTI-CD8 ALPHA DEPLETING ANTIBODIES**REFERENCE TO RELATED APPLICATIONS**

[0001] This application is a continuation of International Patent Application No. PCT/EP2023/082168, filed on Nov. 17, 2023, which claims the benefit of EP 22306689.5, filed Nov. 22, 2022, each of which is incorporated by reference herein in its entirety.

REFERENCE TO A SEQUENCE LISTING

[0002] In accordance with 37 CFR § 1.831, the present specification makes reference to a Sequence Listing submitted electronically as a .xml file named “2490 PCT—Sequence listing.xml”. The .xml file was generated on Oct. 4, 2023 and is 45,256 bytes in size. The entire contents of the Sequence Listing are hereby incorporated by reference.

FIELD

[0003] The present invention relates to anti-CD8a (cluster of differentiation 8 alpha) antibodies and antigen-binding fragments thereof.

BACKGROUND

[0004] CD8α (cluster of differentiation 8 alpha) is a glycoprotein, member of the immunoglobulin superfamily. CD8α either homodimerizes with another CD8α protein, or heterodimerizes with a CD8β (cluster of differentiation 8 beta) protein to form the dimeric CD8 molecule expressed at the surface of CD8+ cytotoxic T cells. On CD8+ T cells, the dimeric CD8 molecule serves as a co-receptor for the T-cell receptor (TCR) to recognize antigens displayed by an antigen presenting cell in the context of class I MHC molecules. Therefore, the dimeric CD8 molecule plays a role of interaction reinforcement between the T cell and the antigen-presenting cell.

[0005] Both CD8α and CD8β proteins are constituted of an immunoglobulin-like extracellular domain, and of a long glycosylated peptide chain which serves both as transmembrane and intracellular domains. Thus, the dimeric CD8 molecule is also important for transmitting signals to the T cell through its intracellular domain which recruits the tyrosine kinase p56^{lck}, important for the phosphorylation cascade taking place after antigen recognition.

[0006] CD8+ cytotoxic T cells have been shown to mediate the clearance of cancer cells, cells infected by intracellular pathogens (such as viruses or bacteria), as well as damaged cells. When exposed to infected or dysfunctional cells, CD8+ cytotoxic T cells express death-inducing ligands, and they also release cytotoxins such as perforin, granzymes, and granulysin, which will induce the apoptosis or lysis of the target cells. Therefore, CD8+ cytotoxic T cells play an important role in the immune response and the fight against intracellular pathogens and tumor cells.

[0007] However, CD8+ cytotoxic T cells have also been involved in the development of various diseases and disorders, and it may thus be advantageous in certain circumstances to be able to deplete CD8+ cytotoxic T cells in a controlled manner. Therefore, there is a need for a new therapeutic tool specifically targeting CD8+ cytotoxic T cells that would allow to transiently deplete CD8+ cytotoxic T cells in a controlled manner.

[0008] Consequently, the Inventors herein aimed at developing novel antibodies that specifically bind CD8α and

allow the depletion of CD8α-expressing cells, mainly CD8+ cytotoxic T cells. Advantageously, the anti-CD8α antibodies as described herein are capable of inducing a fast and almost complete depletion of CD8α-expressing cells, followed by a controlled repopulation of CD8α-expressing cells. Thus, the anti-CD8α antibodies as described herein would make a valuable therapeutic tool for transiently depleting CD8α-expressing cells.

SUMMARY

[0009] The present invention relates to an isolated anti-CD8α antibody, or an antigen-binding fragment thereof, wherein:

[0010] a) the variable region of the heavy chain (VH) of said isolated anti-CD8α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

[0011] V_H-CDR1: NX₁X₂MN, wherein

[0012] X₁ is N or Y,

[0013] X₂ is D or A or Y,

[0014] V_H-CDR2: X₃ISGSSX₄YIX₅YADFVKG (SEQ ID NO: 1), wherein

[0015] X₃ is D or S,

[0016] X₄ is S or R,

[0017] X₅ is D or G or Y,

[0018] V_H-CDR3:

SSX₆X₇X₈X₉YX₁₀X₁₁X₁₂X₁₃MDV (SEQ ID NO: 2), wherein

[0019] X₆ is N or no amino acid,

[0020] X₇ is Y or no amino acid,

[0021] X₈ is Y or G or no amino acid,

[0022] X₉ is D or S or no amino acid,

[0023] X₁₀ is S or N or F,

[0024] X₁₁ is A or G,

[0025] X₁₂ is S or D or N,

[0026] X₁₃ is A or G; and

[0027] b) the variable region of the light chain (VL) of said isolated anti-CD8α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

[0028] VL-CDR1: AGTSSDVGGX₁₄X₁₅X₁₆VS (SEQ ID NO: 3), wherein

[0029] X₁₄ is G or N or Y,

[0030] X₁₅ is S or Y,

[0031] X₁₆ is S or Y,

[0032] V_L-CDR2: X₁₇DSX₁₈RPS (SEQ ID NO: 4), wherein

[0033] X₁₇ is Q or S or Y,

[0034] X₁₈ is Y or S,

[0035] VL-CDR3: SSX₁₉TX₂₀YSTRV (SEQ ID NO: 5), wherein

[0036] X₁₉ is Y or D,

[0037] X₂₀ is Y or Q or S.

[0038] In some embodiments, the isolated anti-CD8α antibody, or antigen-binding fragment thereof, comprises:

a) a VH comprising the three following CDRs:

-V_H-CDR1:

(SEQ ID NO: 6)

NNAMN,

(SEQ ID NO: 12)

NYDMN,

or -continued

NYVMN; (SEQ ID NO: 18)

-V_H-CDR2: (SEQ ID NO: 7)

DISGSSRYIGYADFVKG, (SEQ ID NO: 13)

DISGSSSYIDYADFVKG, or (SEQ ID NO: 19)

SISGSSRYIYYADFVKG; (SEQ ID NO: 8)

-V_H-CDR3: (SEQ ID NO: 14)

SSNYDYDYNADAMDV, (SEQ ID NO: 20)

SSYSGSGMDV, or (SEQ ID NO: 9)

SSGSYFGNAMDV; and (SEQ ID NO: 15)

b) a VL comprising the three following CDRs: (SEQ ID NO: 21)

-V_L-CDR1: (SEQ ID NO: 10)

AGTSSDVGGNSYVS, (SEQ ID NO: 16)

AGTSSDVGGSSVS, or (SEQ ID NO: 22)

AGTSSDVGGYYSVS; (SEQ ID NO: 11)

-V_L-CDR2: (SEQ ID NO: 17)

SDSSRPS, (SEQ ID NO: 23)

QDSYRPS, or (SEQ ID NO: 6)

YDSSRPS (SEQ ID NO: 7)

-V_L-CDR3: (SEQ ID NO: 11)

SSYTQYSTRV, (SEQ ID NO: 17)

SSYTYSTRV, or (SEQ ID NO: 23)

SSDTSYSTRV.

[0039] In some embodiments, the isolated anti-CD8α antibody, or antigen-binding fragment thereof, comprises the six following CDRs:

-V_H-CDR1: (SEQ ID NO: 6)

NNAMN,

V_H-CDR2: (SEQ ID NO: 7)

DISGSSRYIGYADFVKG,

-continued

V_H-CDR3: (SEQ ID NO: 8)

SSNYDYDYNADAMDV,

V_L-CDR1: (SEQ ID NO: 9)

AGTSSDVGGNSYVS,

VL-CDR2: (SEQ ID NO: 10)

SDSSRPS, and

V_L-CDR3: (SEQ ID NO: 11)

SSYTQYSTRV; or

-V_H-CDR1: (SEQ ID NO: 12)

NYDMN,

V_H-CDR2: (SEQ ID NO: 13)

DISGSSSYIDYADFVKG,

V_H-CDR3: (SEQ ID NO: 14)

SSYSGSGMDV,

V_L-CDR1: (SEQ ID NO: 15)

AGTSSDVGGSSVS,

V_L-CDR2: (SEQ ID NO: 16)

QDSYRPS, and

V_L-CDR3: (SEQ ID NO: 17)

SSYTYSTRV; or

-V_H-CDR1: (SEQ ID NO: 18)

NYVMN,

V_H-CDR2: (SEQ ID NO: 19)

SISGSSRYIYYADFVKG,

V_H-CDR3: (SEQ ID NO: 20)

SSGSYFGNAMDV,

V_L-CDR1: (SEQ ID NO: 21)

AGTSSDVGGYYSVS,

V_L-CDR2: (SEQ ID NO: 22)

YDSSRPS, and

V_L-CDR3: (SEQ ID NO: 23)

SSDTSYSTRV.

[0040] In some embodiments, the isolated anti-CD8α antibody, or antigen-binding fragment thereof, comprises a variable region of the heavy chain (V_H) comprising a sequence as set forth in SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28, or a sequence having at least 80% identity with SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28. In some embodiments, the isolated anti-CD8α

antibody, or antigen-binding fragment thereof, comprises a variable region of the light chain (VL) comprising a sequence as set forth in SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29, or a sequence having at least 80% identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0041] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises:

a V_H comprising a sequence as set forth in SEQ ID NO: 24, or a sequence having at least 80% identity with SEQ ID NO: 24, and a VL comprising a sequence as set forth in SEQ ID NO: 25, or a sequence having at least 80% identity with SEQ ID NO: 25; or

[0042] a V_H comprising a sequence as set forth in SEQ ID NO: 26, or a sequence having at least 80% identity with SEQ ID NO: 26, and a VL comprising a sequence as set forth in SEQ ID NO: 27, or a sequence having at least 80% identity with SEQ ID NO: 27; or

[0043] a V_H comprising a sequence as set forth in SEQ ID NO: 28, or a sequence having at least 80% identity with SEQ ID NO: 28, and a VL comprising a sequence as set forth in SEQ ID NO: 29, or a sequence having at least 80% identity with SEQ ID NO: 29.

[0044] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is a CD8 α -expressing cell depleting antibody.

[0045] The present invention also relates to a fusion protein comprising said anti-CD8 α antibody, or antigen-binding fragment thereof; and to a nucleic acid molecule encoding said anti-CD8 α antibody, or antigen-binding fragment thereof, or said fusion protein.

[0046] The present invention also relates to a pharmaceutical composition comprising said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, said fusion protein, and/or said nucleic acid molecule, and at least one pharmaceutically acceptable excipient.

[0047] The present invention also relates to said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, said fusion protein, said nucleic acid molecule, or said pharmaceutical composition, for use as a medicament.

[0048] The present invention also relates to said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, said fusion protein, said nucleic acid molecule, or said pharmaceutical composition, for use in the treatment of a CD8 $^-$ related disease. In some embodiments, the CD8 $^-$ related disease is a cardiovascular disease.

[0049] The present invention also relates an in vitro method for depleting CD8 α -expressing cells in a sample, comprising a step of contacting said sample with the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein, or with the fusion protein as described herein. The present invention also relates to an in vitro method for detecting and/or quantifying CD8 α in a sample, cell, tissue or organ, comprising contacting said sample, cell, tissue or organ with the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein, or with the fusion protein as described herein.

BRIEF DESCRIPTION OF THE DRAWINGS

[0050] FIG. 1A-C is a combination of graphs showing the dose-response ELISA of the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) in the human IgG1 format on immobilized recombinant human CD8 α , mouse CD8 α and human CD4. 12 different concentrations

were tested (75, 25, 8.3, 2.8, 0.93, 0.31, 0.103, 0.034, 0.011, 0.0038, 0.0013 and 0.00042 μ g/mL). All antibodies had a mean molecular mass of about 150 kDa. An irrelevant human IgG1 antibody named 13R4 was included as negative control. 1A: human CD8 α . 1B: mouse CD8 α , 1C: human CD4.

[0051] FIG. 2A-C is a combination of graphs showing the dose-response ELISA of the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) in the rat IgG2b format on immobilized recombinant human CD8 α , mouse CD8 α and human CD4. 12 different concentrations were tested (75, 25, 8.3, 2.8, 0.93, 0.31, 0.103, 0.034, 0.011, 0.0038, 0.0013 and 0.00042 μ g/mL). All antibodies had a mean molecular mass of about 150 kDa. 2A: human CD8 α . 2B: mouse CD8 α , 2C: human CD4.

[0052] FIG. 3A-E is a combination of flow cytometry plots showing the binding of the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) in the human IgG1 format onto CD8 $^+$ cells (HPB-ALL cells). Irrelevant human IgG1 13R4 antibody was used as a negative control and a commercial mouse anti-human CD8 α antibody was used as a positive control. Antibodies were used at a concentration of 3.3 μ g/mL. Light grey represents cells alone (back ground signal), darker grey represents cells with one of the antibodies as indicated (binding signal).

[0053] 3A represents MP08-R3-F08 human IgG1 clone.

[0054] 3B represents MP08-R3-C11 human IgG1 clone.

[0055] 3C represents MP09-R3-D03 human IgG1 clone.

[0056] 3D represents irrelevant human IgG1 13R4 antibody control.

[0057] 3E represents a commercial mouse anti-human CD8 α antibody.

[0058] FIG. 4A-E is a combination of flow cytometry plots showing the binding of the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) in the human IgG1 format onto CD8 $^-$ cells (H1299 cells). Irrelevant human IgG1 13R4 antibody was used as a negative control and a commercial mouse anti-human CD8 α antibody was used as a positive control. Antibodies were used at a concentration of 3.3 μ g/mL. Light grey represents cells alone (background signal), darker grey represents cells with one of the antibodies as indicated (binding signal).

[0059] 4A represents MP08-R3-F08 human IgG1 clone.

[0060] 4B represents MP08-R3-C11 human IgG1 clone.

[0061] 4C represents MP09-R3-D03 human IgG1 clone.

[0062] 4D represents irrelevant human IgG1 13R4 antibody control.

[0063] 4E represents a commercial mouse anti-human CD8 α antibody.

[0064] FIG. 5A-E is a combination of flow cytometry plots showing the binding of the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) in the human IgG1 format onto human primary CD8 $^+$ T cells. Irrelevant human IgG1 13R4 antibody was used as a negative control and a commercial mouse anti-human CD8 α antibody was used as a positive control. Antibodies were used at a concentration of 1.1 μ g/mL. Light grey represents cells alone (background signal), darker grey represents cells with one of the antibodies as indicated (binding signal).

[0065] 5A represents MP08-R3-F08 human IgG1 clone.

[0066] 5B represents MP08-R3-C11 human IgG1 clone.

[0067] 5C represents MP09-R3-D03 human IgG1 clone.

[0068] 5D represents irrelevant human IgG1 13R4 antibody control.

[0069] 5E represents a commercial mouse anti-human CD8 α antibody.

[0070] FIG. 6A-B is a combination of flow cytometry plots showing the percentage of cells expressing the human CD8 α transgene from CD45 $^{+}$ cells in control WT mice (6A) and transgenic mice expressing the human CD8 α transgene (6B).

[0071] FIG. 7A-C is a combination of graphs showing the depletion and subsequent repopulation of human CD8 α -expressing cells in mice after injection with antibody MP08-R3-F08 (7A), MP08-R3-C11 (7B) or MP09-R3-D03 (7C) at a dose of 50 μ g, 100 μ g or 200 μ g as indicated.

[0072] FIG. 8A-C is a combination of histograms showing the depletion and subsequent repopulation of human CD8 α -expressing cells in mice after injection with a dose of 200 μ g (8A), 100 μ g (8B) or 50 μ g (8C) of antibody MP08-R3-F08, MP08-R3-C11 or MP09-R3-D03 as indicated.

[0073] FIG. 9A-E is a combination of flow cytometry plots showing the percentage of mouse CD8 α^{+} cells in control C56B16 mice after injection with a control IgG1 isotype antibody (9A), MP08-R3-C11 (9B), MP08-R3-F08 (9C) or MP09-R3-D03 (9D) and of a histogram showing the quantification 1 hour and 7 days after injection (9E).

[0074] FIG. 10A-C is a combination of fluorescent images showing the confirmation screen on fixed cells (10A-B) and on non-fixed cells (10C) for anti-CD8 α antibody clone MP08-R3-F08 (10A & C) and control antibody Rituximab (10B & C). Each spot corresponds to cells transfected with an expression vector expressing proteins as indicated.

[0075] FIG. 11A-B is a combination of histograms showing the depletion of CD8 α -expressing cells in Cynomolgus monkeys after injection with anti-CD8 α antibody. 11A represents the percentage of CD8 $^{+}$ cells within CD45 $^{+}$ cells in Cynomolgus monkeys which received 20 mg/kg body-weight of anti-CD8 α antibody clone MP08-R3-F08, MP08-R3-C11 or MP09-R3-D03 as indicated. 11B represents the percentage of CD8 $^{+}$ cells within CD45 $^{+}$ cells in Cynomolgus monkeys which received 7.5 mg/kg or 20 mg/kg body-weight of anti-CD8 α antibody clone MP08-R3-F08, as indicated.

[0076] FIG. 12A-C is a combination of flow cytometry plots showing the percentage of CD45 $^{+}$ hCD8 α^{+} cells in the blood of control transgenic mice expressing the human CD8 α transgene which received saline solution (12A) and transgenic mice expressing the human CD8 α transgene which received anti-CD8 α antibody clone MP08-R3-F08 (12B), 3 days after injection, and of a histogram showing the quantification (12C). Males are represented by black circles and females by inverted black triangle. Control n=9 vs MP08-R3-F08 n=7. (**p<0.001).

[0077] FIG. 13A-C is a combination of flow cytometry plots showing the percentage of CD45 $^{+}$ hCD8 α^{+} cells in the spleen of control transgenic mice expressing the human CD8 α transgene which received saline solution (13A) and transgenic mice expressing the human CD8 α transgene which received anti-CD8 α antibody clone MP08-R3-F08 (13B), 3 days after injection, and of a histogram showing the quantification (13C). Males are represented by black circles and females by inverted black triangle. Control n=9 vs MP08-R3-F08 n=7. (**p<0.001).

[0078] FIG. 14 is a histogram showing the percentage of necrosis measured by staining with 2,3,5-triphenyltetrazolium chloride (TTC) on heart sections from control transgenic mice expressing the human CD8 α transgene which

received saline solution or transgenic mice expressing the human CD8 α transgene which received anti-CD8 α antibody clone MP08-R3-F08, 3 days after induction of myocardial infarction. Males are represented by black circles and females by inverted black triangle. Control n=9 vs MP08-R3-F08 n=7. (**p<0.01).

DETAILED DESCRIPTION

[0079] In the present invention, the following terms have the following meanings:

[0080] “About” preceding a figure encompasses plus or minus 10%, or less, of the value of said figure. It is to be understood that the value to which the term “about” refers is itself also specifically, and preferably, disclosed.

[0081] “Affinity” is used to define the strength of an antibody-antigen complex. Affinity measures the strength of interaction between an antigen and an antibody, or an antigen-binding fragment thereof. It may be expressed by an affinity constant K_A or by a dissociation constant K_D .

[0082] “Anti-CD8 α ” (also sometimes referred to as “anti-CD8 α^{+} ”), when qualifying an antibody, or an antigen-binding fragment thereof, means that said antibody, or antigen-binding fragment thereof, specifically binds to or specifically recognizes CD8 α (cluster of differentiation 8 alpha).

[0083] “Antibody (Ab)” and “immunoglobulin (Ig)” may be used interchangeably and refer to a protein having a combination of two heavy chains (H chains) and two light chains (L chains). In particular, the term “antibodies” refers to such assemblies which have significant known specific immunoreactive activity to an antigen of interest (e.g., CD8 α , in particular human CD8 α). As explained elsewhere herein, “specificity” for human CD8 α , does not exclude cross-reactivity with orthologs of human CD8 α , such as, for example, with simian or murine CD8 α . As mentioned above, antibodies and immunoglobulins comprise light and heavy chains, with or without an interchain covalent linkage between them. Basic immunoglobulin structures in vertebrate systems are relatively well-understood. The generic term “immunoglobulin” comprises five distinct classes of immunoglobulins that can be distinguished biochemically: IgG, IgM, IgA, IgD, and IgE. IgG immunoglobulins comprise two identical light chains with a molecular weight of about 23 kDa, and two identical heavy chains with a molecular weight of about 53-70 kDa. The four chains are joined by disulfide bonds in a “Y” configuration wherein the light chains bracket the heavy chains starting at the mouth of the “Y” and continuing through the variable region. The light chains of an immunoglobulin are classified as either kappa (κ) or lambda (λ). Each heavy chain class may be bonded with either a κ or λ light chain. In general, the light and heavy chains are covalently bonded to each other, and the “tail” regions of the two heavy chains are bonded to each other by covalent disulfide linkages or non-covalent linkages when the immunoglobulins are generated either by hybridomas, B cells or genetically engineered host cells. In the heavy chains, the amino acid sequence runs from the N-terminus at the forked ends of the Y configuration to the C-terminus at the bottom of each chain. Those skilled in the art will appreciate that heavy chains are classified as gamma (γ), mu (μ), alpha (α), delta (δ), or epsilon (ϵ) with some subclasses among them (e.g., 71-74). It is the nature of the heavy chain that determines the “class” of the antibody as IgG, IgM, IgA, IgD, or IgE, respectively. The immunoglobulin subclasses or “isotypes” (e.g., IgG1, IgG2, IgG3,

IgG4, IgA1, etc.) are well-characterized and are known to confer functional specialization. The variable region of an antibody allows the antibody to selectively recognize and specifically bind to an epitope on an antigen. That is, the light chain variable region (VL) and heavy chain variable region (VH) of an antibody combine to form the variable region that defines a three-dimensional antigen-binding site. This quaternary antibody structure thus forms the antigen-binding site present at the end of each arm of the “Y”. More specifically, the antigen-binding site is defined by three complementarity determining regions (CDRs) on each of the VH and VL.

[0084] “Antigen-binding fragment” of an antibody, which is interchangeable with the term “antigen-binding domain” of an antibody, refers to a part or region of an antibody, or immunoglobulin, which comprises fewer amino acid residues than a whole antibody, or immunoglobulin, and which is capable of binding to an antigen and/or of competing with a whole antibody for antigen binding (e.g., for specific binding to CD8 α). Examples of antigen-binding antibody fragments include, but are not limited to, Fab, Fab', F(ab')₂, Fv fragments, scFv fragments, disulfide-linked Fvs (sdFv), a Fd fragment consisting of the VH and CH1 domains, linear antibodies, single domain antibodies such as sdAb (either VL or VH), camelid VHH domains, multi-specific antibodies formed from antibody fragments such as a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region, and an isolated CDR of an antibody.

[0085] Papain digestion of antibodies produces two identical antigen-binding fragments, called “Fab” fragments, and a residual “Fc” fragment, a designation reflecting the ability to crystallize readily. A Fab fragment consists of one entire L chain (variable region of the L chain (VL) and constant domain of the L chain (CL)), along with part of one H chain consisting of the variable region of the H chain (V_H) and the first constant domain of the heavy chain (CH1). Each Fab fragment is monovalent with respect to antigen binding, i.e., it has a single antigen-binding site.

[0086] Pepsin treatment of an antibody yields a single large F(ab')₂ fragment that roughly corresponds to two disulfide-linked Fab fragments having divalent antigen-binding activity and is still capable of crosslinking antigen.

[0087] Fab' fragments differ from Fab fragments by having additional few amino acid residues at the carboxy terminus of the CH1 domain including one or more cysteines from the antibody hinge region.

[0088] Fab'-SH is the designation for Fab' in which the cysteine residue(s) of the constant domains bear(s) a free thiol group.

[0089] F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments that have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

[0090] “Antigen” or “Ag” as used herein refers to a protein (e.g., CD8 α), which is specifically recognized by an antibody or antibody-binding fragment thereof.

[0091] “CD8”, or “cluster of differentiation 8” refer to a transmembrane glycoprotein notably expressed on cytotoxic T lymphocytes. The CD8 molecule serves as co-receptor for the T-cell receptor (TCR) on T cells to recognize antigens displayed by antigen presenting cells in the context of class I MHC molecules. CD8 functions either as a homodimer composed of two CD8 α chains, or as a heterodimer com-

posed of one CD8 α chain and one CD8 β chain. Both CD8 α and CD8 β are members of the immunoglobulin superfamily. Both CD8 α and CD8 β chains comprise one unique immunoglobulin-like domain constituting the extracellular part of the protein, which is anchored to the cell membrane through a long glycosylated peptide chain. The two chains constituting the CD8 dimer (i.e., heterodimer CD8 α /CD8 β or homodimer CD8 α /CD8 α) are associated through a disulfide bond.

[0092] In the NCBI databases (<https://www.ncbi.nlm.nih.gov>), the reference human CD8 α gene sequence corresponds to NCBI Gene ID: 925, as updated on Aug. 5, 2022. The human CD8 α (hCD8 α) gene consists of 10 exons on chromosome 2p11.2, and encodes a 235 amino acid protein for isoform 1 and a 198 amino acid protein for isoform 2. Human CD8 α isoform 1 refers to the protein referenced as NP_001139345.1, NP_001759.3 or NP_001369627.1 in the NCBI databases, while isoform 2 refers to the protein referenced as NP_741969.1. Alternative names for CD8 α include “T-cell surface glycoprotein CD8 alpha chain”, “CD8 antigen”, “alpha polypeptide”, “Leu2 T-lymphocyte antigen”, “OKT8 T-cell antigen”, “T cell co-receptor”, “T-cell antigen Leu2”, “T-lymphocyte differentiation antigen T8”, “T8 T-cell antigen”, “p32”, “Leu-2” and “Leu2” as non-limiting examples. Herein, the expressions “CD8 α ” and “CD8” are used indifferently.

[0093] In the NCBI databases, the reference human CD8 β gene sequence corresponds to NCBI Gene ID: 926, as updated on Aug. 5, 2022. The human CD8 β gene consists of 8 exons on chromosome 2p11.2, and encodes 6 isoforms by alternative splicing: a 246 amino acid protein for isoform 1, a 243 amino acid protein for isoform 2, a 221 amino acid protein for isoform 3, a 213 amino acid protein for isoform 4, a 210 amino acid protein for isoform 5 and a 198 amino acid protein for isoform 6. Human CD8 β isoform 1 refers to the protein referenced as XP_011531466.1 in the NCBI databases, isoform 2 refers to the protein referenced as NP_757362.1, isoform 3 refers to the protein referenced as NP_742099.1, isoform 4 refers to the protein referenced as NP_742100.1, isoform 5 refers to the protein referenced as NP_004922.1, and isoform 6 refers to the protein referenced as NP_001171571.1. Alternatives names for CD8 β include “LY3”, “P37”, “LEU2”, “LYT3”, “CD8B1”, “T-cell surface glycoprotein CD8 beta chain”, “CD8 antigen”, “beta polypeptide 1”, “T lymphocyte surface glycoprotein beta chain” as non-limiting examples.

[0094] “CDR” or “complementarity determining region” means the non-contiguous antigen combining sites found within the variable region of both the heavy chain and the light chain. The precise amino acid sequence boundaries of a given CDR can be determined using any of a number of well-known schemes, including those described by Kabat et al. (1991), “Sequences of Proteins of Immunological Interest” 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (“Kabat” numbering scheme); Al-Lazikani et al., (1997) JMB 273,927-948 (“Chothia” numbering scheme); or a combination thereof such as the AbM definition which is a compromise between the two used by Oxford Molecular's AbM antibody modelling software. More recently, a universal numbering system has been developed and widely adopted, ImMunoGeneTics (IMGT) Information System® (Lefranc et al., Nucleic Acids Res. 27: 209-212 1999). IMGT® is an integrated information system specializing in immunoglobulins (Ig), T cell receptors

(TCR) and major histocompatibility complex (MHC) of human and other vertebrates. Herein, the CDRs are referred to in terms of both the amino acid sequence and the location within the light or heavy chain (e.g., V_H -CDR1, V_H -CDR2, V_H -CDR3, V_L -CDR1, V_L -CDR2, V_L -CDR3). As the “location” of the CDRs within the structure of the immunoglobulin variable region is conserved between species and present in structures called loops, by using numbering systems that align variable region sequences according to structural features, CDR and framework amino acid residues may be readily identified. This information can be used in grafting and replacement of CDRs from immunoglobulins of one species into an acceptor framework from, typically, a human antibody. Correspondence between the Kabat numbering and the IMGT® unique numbering system is also well-known to one skilled in the art (e.g., Lefranc et al., supra).

[0095] “Epitope” refers to a specific arrangement of amino acids located on a protein or proteins to which an antibody, or an antigen-binding fragment thereof, specifically binds. Epitopes often consist of a chemically active surface grouping of molecules such as amino acids or sugar side chains, and have specific three-dimensional structural characteristics as well as specific charge characteristics. Epitopes can be linear (or sequential) or conformational, i.e., involving two or more sequences of amino acids in various regions of the antigen that may not necessarily be contiguous.

[0096] “Fc domain,” “Fc portion,” and “Fc region” may be used interchangeably and refer to a C-terminal fragment of an antibody heavy chain, e.g., from about amino acid (aa) 230 to about aa 450 of human gamma heavy chain or its counterpart sequence in other types of antibody heavy chains (e.g., α , δ , ϵ and μ for human antibodies), or a naturally occurring allotype thereof.

[0097] “Framework region” or “FR region” or “non-CDR region” includes the amino acid residues that are part of the variable region, but are not part of the CDRs (e.g., using the Kabat definition of CDRs, the Chothia definition of CDRs, or the IMGT® numbering definition of CDRs). Therefore, a variable region framework ranges from about 100 to 120 amino acids in length but includes only those amino acids outside of the CDRs.

[0098] For the specific example of a heavy chain variable region (VH) and for the CDRs as defined by Kabat or Chothia:

[0099] FR1 may correspond to the domain of the variable region encompassing amino acids 1-25 according to Chothia/AbM’s definition, or 5 amino acid residues later according to Kabat’s definition;

[0100] FR2 may correspond to the domain of the variable region encompassing amino acids 36-49;

[0101] FR3 may correspond to the domain of the variable region encompassing amino acids 67-98; and

[0102] FR4 may correspond to the domain of the variable region from amino acid 104-110 to the end of the variable region.

The framework regions for the light chain are similarly separated by each of the CDRs of the light chain variable region (VL).

In naturally occurring antibodies, the six CDRs present on each monomeric antibody are short, non-contiguous sequences of amino acids that are specifically positioned to form the antigen-binding site as the antibody assumes its three-dimensional configuration in an aqueous environment. As indicated above, the remainders of the heavy and light

variable regions show less inter-molecular variability in amino acid sequence and correspond to the framework regions. The framework regions largely adopt a j-sheet conformation and the CDRs form loops which connect, and in some cases form part of, the j-sheet structure. Thus, these framework regions act to form a scaffold for positioning the six CDRs in correct orientation by inter-chain, non-covalent interactions. The antigen-binding site formed by the positioned CDRs defines a surface complementary to the epitope on the immunoreactive antigen. This complementary surface promotes the non-covalent binding of the antibody to the immunoreactive antigen epitope.

[0103] “Heavy chain region” includes amino acid sequences derived from the constant domains of an immunoglobulin heavy chain. A protein comprising a heavy chain region comprises at least one of a CH1 domain, a hinge region (e.g., upper, middle, and/or lower hinge domain), a CH2 domain, a CH3 domain, or a variant or fragment thereof. In some embodiments, the antibody, or the antigen-binding fragment thereof, as described herein may comprise the Fc region of an immunoglobulin heavy chain (e.g., a hinge portion, a CH2 domain, and a CH3 domain). In some embodiments, the antibody, or the antigen-binding fragment thereof, as described herein lacks at least a region of a constant domain (e.g., all or part of a CH2 domain). In some embodiments, at least one, and preferably all, of the constant domains are derived from a human immunoglobulin heavy chain. For example, in some embodiments, the heavy chain region comprises a fully human hinge domain. In some embodiments, the heavy chain region comprises a fully human Fc region (e.g., hinge, CH2 and CH3 domain from a human immunoglobulin). In some embodiments, the constituent constant domains of the heavy chain region are from different immunoglobulin molecules. For example, a heavy chain region of a protein may comprise a CH2 domain derived from an IgG1 molecule and a hinge region derived from an IgG3 or IgG4 molecule. In some embodiments, the constant domains are chimeric domains comprising regions of different immunoglobulin molecules. For example, a hinge may comprise a first region from an IgG1 molecule and a second region from an IgG3 or IgG4 molecule. In some embodiments, the constant domains of the heavy chain region may be modified such that they vary in amino acid sequence from the naturally occurring (wild-type) immunoglobulin molecule. That is, the antibody, or the antigen-binding fragment thereof, as described herein may comprise alterations or modifications to one or more of the heavy chain constant domains (CH1, hinge, CH2 or CH3) and/or to the light chain constant domain (CL). Exemplary modifications include additions, deletions or substitutions of one or more amino acids in one or more domains.

[0104] “Hinge region” includes the region of a heavy chain molecule that joins the CH1 domain to the CH2 domain. This hinge region comprises approximately 25 amino acid residues and is flexible, thus allowing the two N-terminal antigen-binding regions to move independently. Hinge regions can be subdivided into three distinct domains: upper, middle, and lower hinge domains.

[0105] “Identity” or “identical”, when used in the present invention in a relationship between the sequences of two or more polypeptides or of two or more nucleic acids, refers to the degree of sequence relatedness between polypeptides or nucleic acids (respectively), as determined by the number of matches between strings of two or more amino acid residues

or of two or more nucleotides, respectively. “Identity” measures the percent of identical matches between the smaller of two or more sequences with gap alignments (if any) addressed by a particular mathematical model or computer program (i.e., “algorithms”). Identity of related polypeptides or nucleic acid sequences can be readily calculated by known methods. Such methods include, but are not limited to, those described in Computational Molecular Biology, Lesk, A. M., ed., Oxford University Press, New York, 1988; Biocomputing: Informatics and Genome Projects, Smith, D. W., ed., Academic Press, New York, 1993; Computer Analysis of Sequence Data, Part 1, Griffin, A. M., and Griffin, H. G., eds., Humana Press, New Jersey, 1994; Sequence Analysis in Molecular Biology, von Heinje, G., Academic Press, 1987; Sequence Analysis Primer, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo et al., SIAM J. Applied Math. 48, 1073 (1988). Preferred methods for determining identity are designed to give the largest match between the sequences tested. Methods of determining identity are described in publicly available computer programs. Preferred computer program methods for determining identity between two sequences include the GCG program package, including GAP (Devereux et al., Nucleic Acids Res. 1984 Jan. 11; 12(1 Pt 1):387-95; Genetics Computer Group, University of Wisconsin, Madison, Wis.), BLASTP, BLASTN, and FASTA (Altschul et al., J. Mol. Biol. 215, 403-410 (1990)). The BLASTX program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (BLAST Manual, Altschul et al. NCB/NLM/NIH Bethesda, Md. 20894; Altschul et al., J. Mol. Biol. 215, 403-410 (1990)). The well-known Smith Waterman algorithm may also be used to determine identity.

[0106] “Isolated” or “non-naturally occurring” with reference to a biological component (such as an antibody or a nucleic acid) refers to a biological component altered or removed from the natural state. For example, an antibody or a nucleic acid naturally present in a living animal is not “isolated” but the same antibody or nucleic acid partially or completely separated from the coexisting materials of its natural state is “isolated”. An isolated antibody or nucleic acid can exist in substantially purified form, or can exist in a non-native environment such as, for example, a host cell. Typically, a preparation of isolated antibody or nucleic acid contains the antibody or nucleic acid at least about 80% pure, at least about 85% pure, at least about 90% pure, at least about 95% pure, greater than 95% pure, greater than about 96% pure, greater than about 97% pure, greater than about 98% pure, or greater than about 99% pure. Nucleic acids and proteins, such as antibodies, that are “non-naturally occurring” or have been “isolated” include nucleic acids and proteins purified by standard purification methods. The term also embraces nucleic acids and proteins prepared by recombinant expression in a host cell as well as chemically synthesized nucleic acids.

[0107] “Monoclonal antibody (mAb)” refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprised in the population are identical except for possible naturally occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. Furthermore, in contrast to polyclonal antibody preparations that include different antibodies directed against different determinants (epitopes),

each monoclonal antibody is directed against a single determinant (epitope) on the antigen. In addition to their specificity, the monoclonal antibodies are advantageous in that they may be synthesized uncontaminated by other antibodies. The modifier “monoclonal” is not to be construed as requiring production of the antibody by any particular method. For example, a monoclonal antibody, or an antigen-binding fragment thereof, as described herein may be prepared by the well-known hybridoma methodology, or may be made using recombinant DNA methods in bacterial, eukaryotic, animal, or plant cells. The “monoclonal antibodies” may also be isolated from phage antibody libraries using techniques commonly known in the field.

[0108] As used herein, the term “nucleic acid” or “polynucleotide” refers to a polymer of nucleotides covalently linked by phosphodiester bonds, such as deoxyribonucleic acids (DNA) or ribonucleic acids (RNA), in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids containing known analogues of natural nucleotides that have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions), alleles, orthologs, SNPs, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues.

[0109] “Pharmaceutically acceptable excipient” or “pharmaceutically acceptable carrier” refers to an excipient or carrier that does not produce an adverse, allergic or other untoward reaction when administered to a mammal, preferably a human. It includes any and all solvents, such as, for example, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents. A pharmaceutically acceptable excipient or carrier refers to a non-toxic solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. For human administration, preparations should meet sterility, pyrogenicity, general safety and purity standards as required by the regulatory offices such as the FDA (US Food and Drug Administration) or EMA (European Medicines Agency).

[0110] “Single-chain Fv”, also abbreviated as “sFv” or “scFv”, refers to a fusion protein comprising at least one antibody fragment comprising a variable region of a light chain (VL) and at least one antibody fragment comprising a variable region of a heavy chain (V_H), wherein the light and heavy chain variable regions are contiguously linked, e.g., via a synthetic linker, such as a short flexible polypeptide linker, and capable of being expressed as a single chain polypeptide, and wherein the scFv retains the specificity of the intact antibody from which it is derived.

[0111] “Subject” refers to a warm-blooded animal, more preferably a mammal. The term “mammal” refers here to any mammal, including humans. Preferably, the mammal is a primate, more preferably a human. In some embodiments, a subject may be a “patient”, who/which is awaiting the receipt of, or is receiving medical care or was/is/will be the object of a medical procedure, or is monitored for the development of the targeted disease or condition. In some embodiments, the subject is an adult (for example a subject

above the age of 18). In some embodiments, the subject is a child (for example a subject below the age of 18). In some embodiments, the subject is a male. In some embodiments, the subject is a female.

[0112] “Variable”, “variable region” or “variable domain” refer to the fact that certain regions of the variable domains V_H and V_L differ extensively in sequence among antibodies and are used in the binding and specificity of each particular antibody for its target antigen. However, the variability is not evenly distributed throughout the variable domains of antibodies. It is concentrated in three segments called “hypervariable loops” in each of the V_L domain and the V_H domain which form part of the antigen-binding site. The 6 hypervariable loops may each comprise part of a CDR, as defined hereinabove.

[0113] A first object of the present invention relates to an isolated antibody, or to an antigen-binding fragment thereof, which specifically binds to cluster of differentiation 8 alpha (CD8 α). A first object of the present invention thus relates to an isolated anti-CD8 α antibody, or to an antigen-binding fragment thereof.

[0114] As used herein, an antibody or antigen-binding fragment thereof is said to be “specific for”, “immunospecific” or to “specifically bind” its cognate antigen (e.g., CD8 α) if it reacts at a detectable level with said antigen. The binding properties of an antibody, or an antigen-binding fragment thereof, to its cognate antigen (or cells or tissues expressing said antigen) may generally be determined and assessed using immunodetection methods well-known in the art including, for example, enzyme-linked immunosorbent assay (ELISA), immunofluorescence-based assays, such as immuno-histochemistry (IHC) and/or fluorescence-activated cell sorting (FACS); or by surface plasmon resonance (SPR).

[0115] The affinity of an antibody, or an antigen-binding fragment thereof, for its cognate antigen (e.g., CD8 α) can be readily determined using conventional techniques, for example, those described by Scatchard, 1949. *Ann NY Acad Sci.* 51:660-672. Affinity of an antibody, or an antigen-binding fragment thereof, for its cognate antigen is commonly expressed as an equilibrium dissociation constant (K_D). Thus, in some embodiments, the isolated anti-CD8 α antibody, or an antigen-binding fragment thereof, is said to be “immunospecific”, “specific for” or to “specifically bind” to its cognate antigen CD8 α if it reacts at a detectable level with said antigen with a K_D of less than or equal to 10^{-6} M.

[0116] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein specifically binds to human CD8 α (hCD8 α or huCD8 α). In other words, in some embodiments, the isolated antibody, or the antigen-binding fragment thereof, is an anti-hCD8 α antibody, or anti-hCD8 α antigen-binding fragment thereof.

[0117] The binding of an antibody, or an antigen-binding fragment thereof, for its cognate antigen may also be assessed through the determination of the concentration of antibody, or antigen-binding fragment thereof, that gives half-maximal binding (i.e., EC50). For example, the EC50 of an antibody, or an antigen-binding fragment thereof, may be determined with an enzyme-linked immunosorbent assay (ELISA), in particular with a dose-response ELISA. In some embodiments, the isolated anti-CD8 α antibody, or an antigen-binding fragment thereof, binds CD8 α , preferably hCD8 α , with an EC50 of less than 10 nM, preferably less

than 5 nM, more preferable less than 2.5 nM, in particular as determined with dose-response ELISA.

[0118] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein specifically binds human CD8 α having an amino acid sequence as set forth in SEQ ID NO: 30, corresponding to NCBI accession number NP_001759.3, last modified on Jun. 12, 2022.

[0119] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, specifically binds to the extracellular domain of CD8 α .

[0120] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein specifically binds to human CD8 α isoform 1 and isoform 2. In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein specifically binds to human CD8 α isoform 1 and isoform 2 both secreted form and tethered secreted form.

[0121] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein specifically binds to CD8 α , in particular to human CD8 α , present at the surface of leukocytes. In some embodiments, leukocytes encompass CD8+ T lymphocytes (also known as CD8+ T cells), CD8 α -expressing dendritic cells (also referred to as CD8+ dendritic cells) and CD8 α -expressing Natural Killer (NK) cells (also referred to as CD8+NK cells).

[0122] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is able to deplete CD8 α -expressing cells (which may also be referred to as CD8+ cells). Thus, in some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is a depleting antibody, that is to say a CD8 α -expressing cell depleting antibody (which may also be referred to as a CD8+ cell depleting antibody). The CD8 α -expressing cells may be CD8 α -expressing leukocytes, in particular CD8 α -expressing T cells (also referred to as CD8+ T cells), CD8 α -expressing NK cells (also referred to as CD8+NK cells) and/or CD8 α -expressing dendritic cells (also referred to as CD8+ dendritic cells). The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may thus be able to deplete CD8+ T cells and/or CD8+NK cells. Accordingly, in some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is a CD8 α -expressing T cell depleting antibody, also referred to as a CD8+ T cell depleting antibody.

[0123] The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may be able to induce the in vitro depletion of CD8 α -expressing cells in a sample upon contact with said sample. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may be able to induce the in vivo depletion of CD8 α -expressing cells in an organism or subject upon administration to said organism or subject.

[0124] As used herein, the terms “a CD8 α -expressing cell depleting antibody” or “a CD8+ cell depleting antibody” refer to an antibody which binds to CD8 α (i.e., CD8) expressed at the surface of CD8 α -expressing cells (i.e., CD8+ cells) and which mediate the destruction or depletion of said cells upon binding to CD8 α (i.e., CD8).

[0125] As used herein “deplete”, “depleting”, “depletion” with respect to cells expressing CD8 α as described herein refer to a measurable decrease or reduction in the number of CD8 α -expressing cells in a sample, in an organism or in a

subject. For example, depletion with respect to cells expressing CD8 α as described herein may refer to a measurable decrease or reduction in the number of circulating CD8 α -expressing cells, and/or tissue-infiltrating CD8 α -expressing cells, and/or CD8 α -expressing cells present in the lymph nodes and/or in the spleen in an organism or in a subject.

[0126] The decrease or reduction may be at least about 10%, e.g., at least about 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, or more, in particular with reference to the initial number of CD8 α -expressing cells in the sample, organism or subject. As used herein, “initial number of CD8 α -expressing cells” (or “number of CD8 α -expressing cells at baseline”) refers to the number of CD8 α -expressing cells in a sample, organism or subject before depletion induced by the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein (i.e., before administration of the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein). In some embodiments, the initial number of CD8 α -expressing cells in an organism or subject is a normal number of CD8 α -expressing cells, that is to say a number of CD8 α -expressing cells within the range considered normal or standard (i.e., normal range) for the organism or subject. For example, a normal number of CD8+ T cells (in particular circulating CD8+ T cells) in healthy human adults ranges from 150 to 1000 cells/mm³. A normal mean number of CD8+ T cells (in particular circulating CD8+ T cells) in healthy human adults may thus be about 550 cells/ μ L. In some embodiments, the decrease or reduction may be at least about 10%, e.g., at least about 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99%, or more, with reference to a number of CD8 α -expressing cells in the organism or subject within the normal range for said organism or subject.

[0127] In some embodiments, the terms “deplete”, “depleting”, “depletion” with respect to cells expressing CD8 α as described herein refer to a decrease or reduction in the number of CD8 α -expressing cells in a sample, organism or subject to an amount below detectable limits.

[0128] As indicated above, the depletion of CD8 α -expressing cells induced by the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may be a complete depletion. By “complete depletion” it is meant that, after depletion induced by the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, CD8 α -expressing cells are decreased by at least about 90%, preferably at least about 95%, 96%, 97%, 98%, or 99%, in particular with reference to the initial number of CD8 α -expressing cells in the sample, organism or subject, preferably for at least about 1, 2, 3, 4, 5, 6, or 7 day(s).

[0129] The depletion of CD8 α -expressing cells induced by the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may be a transient depletion. By “transient depletion” it is meant that, depletion of the CD8 α -expressing cells induced by the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is followed by a repopulation of CD8 α -expressing cells, preferably starting about 2, 3, 4, 5, 6, 7 or 8 week(s) after the initial depletion.

[0130] As used herein, “repopulation” means that after CD8 α -expressing cell depletion, the number of CD8 α -expressing cells in a sample, in an organism or in a subject

increases. The increase may be sufficient to reach at least about 10%, e.g., at least about 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%, preferably at least about 55%, 60%, 65%, 70%, or 75%, more preferably at least about 80%, 85%, 90%, or 95%, or more, in particular with reference to the initial number of CD8 α -expressing cells in the sample, organism or subject. In some embodiments, the increase may be sufficient to reach at least about 10%, e.g., at least about 15%, 20%, 25%, 30%, 35%, 40%, 45%, or 50%, preferably at least about 55%, 60%, 65%, 70%, or 75%, more preferably at least about 80%, 85%, 90%, or 95%, or more, with reference to a number of CD8 α -expressing cells in the organism or subject within the normal range for said organism or subject.

[0131] The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, may deplete CD8 α -expressing cells as described herein by binding to CD8 α and by activating antibody-dependent cell mediated cytotoxicity (ADCC), complement dependent cytotoxicity (CDC) and/or antibody dependent cellular phagocytosis (ADCP); and/or by inhibiting proliferation of CD8 α -expressing cells; and/or by inducing CD8 α -expressing cells cell death (for example via apoptosis).

[0132] The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, may be able to deplete CD8 α -expressing cells as described herein through antibody-dependent cell mediated cytotoxicity.

[0133] As used herein the term “antibody-dependent cell mediated cytotoxicity” or “ADCC” refers to a form of cytotoxicity in which antibodies bound onto Fc receptors (FcRs) present on certain cytotoxic cells (e.g., NK cells, neutrophils, monocytes and macrophages) enable these cytotoxic effector cells to bind specifically to an antigen-bearing target cell (e.g., a CD8 α -expressing cell as described herein) and subsequently to kill the target cell. To assess the ADCC activity of a molecule of interest, an in vitro ADCC assay, such as that described in U.S. Pat. No. 5,500,362 or U.S. Pat. No. 5,821,337, may be performed.

[0134] The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, may be able to deplete CD8 α -expressing cells as described herein through complement dependent cytotoxicity.

[0135] As used herein, the term “complement dependent cytotoxicity” or “CDC” refers to the lysis of a target cell in the presence of complement. Activation of the classical complement pathway is initiated by the binding of the first component of the complement system to antibodies which are bound to their cognate antigen (e.g., CD8 α). To assess complement activation, a CDC assay such as that described in Gazzano-Santoro et al., 1997. J Immunol Methods. 202 (2):163-71, may be performed.

[0136] The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, may be able to deplete CD8 α -expressing cells as described herein through antibody dependent phagocytosis.

[0137] As used herein the term “antibody dependent cellular phagocytosis” or “ADCP” refers to a form of phagocytosis in which antibodies bound onto Fc receptors (FcRs) present on certain phagocytic cells (e.g., monocytes, macrophages, neutrophils, dendritic cells, and mast cells) enable these phagocytic cells to bind specifically to an antigen-bearing target cell (e.g., a CD8 α -expressing cell as described herein) and subsequently to phagocyte the target cell.

[0138] Methods for assessing whether a compound, such as an antibody or antigen-binding fragment thereof, is able to deplete CD8 α -expressing cells are well-known in the art and comprise, for example, the assays described hereinafter in the example section.

[0139] Assays for evaluating whether an anti-CD8 α antibody, or antigen-binding fragment thereof, is depleting CD8 α -expressing cells include the detection and quantification by flow cytometry of hCD8 α -expressing cells in transgenic mice expressing the hCD8 α transgene after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, an anti-CD8 α antibody, or antigen-binding fragment thereof, able to decrease the number of hCD8 α -expressing cells, in particular as compared to the baseline number of hCD8 α -expressing cells (i.e., the number of hCD8 α -expressing cells before injection of the anti-CD8 α antibody, or antigen-binding fragment thereof), is a CD8 α -expressing cell depleting antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, 3, 4, or 5 hour(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, 3, 4, 5, 6 or 7 day(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, or 3 week(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof.

[0140] Assays for evaluating whether an anti-CD8 α antibody, or antigen-binding fragment thereof, is depleting CD8 α -expressing cells also include the detection and quantification by flow cytometry of CD8 α -expressing cells in non-human primates, for example in cynomolgus monkeys (*Macaca fascicularis*) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, an anti-CD8 α antibody, or antigen-binding fragment thereof, able to decrease the number of CD8 α -expressing cells as compared to the baseline number of hCD8 α -expressing cells (i.e., the number of hCD8 α -expressing cells before injection of the anti-CD8 α antibody, or antigen-binding fragment thereof), is a CD8 α -expressing cell depleting antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 hour(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, or 65 day(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the decrease in the number of hCD8 α -expressing cells is assessed 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 week(s) after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof.

[0141] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is able to limit necrosis after myocardial infarction. In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is able to reduce the infarct size after myocardial infarction. In some embodiments, the isolated anti-CD8 α antibody, or

antigen-binding fragment thereof, as described herein is able to improve cardiac function after myocardial infarction. In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is able to improve the left ventricular ejection fraction after myocardial infarction.

[0142] Methods for assessing whether a compound, such as an antibody or antigen-binding fragment thereof, is able to limit necrosis, reduce the infarct size, improve cardiac function, and/or improve the left ventricular ejection fraction after myocardial infarction are well-known in the art and comprise, for example, the assays described hereinafter in the example section.

[0143] Assays for evaluating whether an anti-CD8 α antibody, or antigen-binding fragment thereof, is able to limit necrosis after myocardial infarction include measuring the percentage of necrosis on heart tissue slices from transgenic mice expressing the hCD8 α transgene after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the percentage of necrosis is assessed 3 days after myocardial infarction. In some embodiments, heart tissue slices are stained with 2,3,5-triphenyltetrazolium chloride (TTC).

[0144] Assays for evaluating whether an anti-CD8 α antibody, or antigen-binding fragment thereof, is able to reduce the infarct size after myocardial infarction include calculating the infarct size by measuring the total tissue area and the infarcted tissue area on heart tissue slices from transgenic mice expressing the hCD8 α transgene after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, the infarct size is assessed 3 days after myocardial infarction. In some embodiments, heart tissue slices are stained with 2,3,5-triphenyltetrazolium chloride (TTC).

[0145] Assays for evaluating whether an anti-CD8 α antibody, or antigen-binding fragment thereof, is able to improve cardiac function and/or improve the left ventricular ejection fraction after myocardial infarction include calculating left ventricular end-diastolic volume (EDV) and end-systolic volume (ESV) in transgenic mice expressing the hCD8 α transgene after injection of the anti-CD8 α antibody, or antigen-binding fragment thereof. In some embodiments, cardiac function is assessed 21 days after myocardial infarction. In some embodiments, left ventricular end-diastolic volume (EDV) and end-systolic volume (ESV) are calculated 21 days after myocardial infarction. In some embodiments, cardiac function is recorded using an imaging system adapted for mouse cardiovascular imaging.

[0146] In some embodiments, “isolated”, as in “isolated antibody or isolated antigen-binding fragment thereof”, refers to an antibody, or antigen-binding fragment thereof, that is substantially free of other proteins or antibodies having different antigenic specificities. Thus, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is substantially free of proteins or antibodies, or antigen-binding fragments thereof, that specifically bind antigens other than CD8 α . The isolated antibody, or the antigen-binding fragment thereof, that specifically binds CD8 α as described herein may, however, have cross-reactivity to other related antigens, such as CD8 α from other genera or species.

[0147] Moreover, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein may be substantially free of other cellular material and/or chemi-

cals, in particular those that would interfere with any uses (such as any therapeutic uses) of the antibody, or antigen-binding fragment thereof, including without limitation, enzymes, hormones, and other proteinaceous or non-proteinaceous components.

[0148] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, as described herein is purified.

[0149] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is purified to obtain a purity greater than about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 98%, or 99% by weight of antibody or antigen-binding fragment, preferably greater than about 95%, 96%, 97%, 98%, or 99% by weight of antibody or antigen-binding fragment. In some embodiments, the purity is determined by analytical size exclusion chromatography (SEC), such as size exclusion high-performance liquid chromatography (SE HPLC).

[0150] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, specifically binds to human CD8 α and to at least one ortholog of human CD8 α . Thus, in some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, binds to human CD8 α and to at least one CD8 α from another genus or species. In other words, in some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, displays cross-reactivity (cross-reacts). In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, binds to human CD8 α and to cynomolgus monkey CD8 α (in particular to CD8 α of *Macaca fascicularis*).

[0151] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, does not bind to mouse CD8 α (in particular to CD8 α of *Mus musculus*).

[0152] In some embodiments, the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, does not exhibit any off-target binding.

[0153] In some embodiments, the antigen-binding fragment of the anti-CD8 α antibody is a molecule selected from the group comprising or consisting of a single-chain antibody or scFv, a dimeric single-chain antibody or di-scFv, a Fv, a Fab, a Fab', a Fab'-SH, and a F(ab')₂.

[0154] Antigen-binding fragments of antibodies can be obtained using standard methods. For instance, Fab or F(ab')₂ fragments may be produced by protease digestion of the isolated antibodies, according to conventional techniques. Alternatively, antigen-binding fragments of antibodies, such as Fab fragments, may be expressed as recombinant proteins.

[0155] In some embodiments, the isolated anti-CD8 α antibody is a whole antibody. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, is selected from the group comprising or consisting of a Fc silent antibody or antigen-binding fragment thereof (i.e., an antibody or antigen-binding fragment thereof comprising a Fc silent), an antibody or antigen-binding fragment thereof with an engineered Fc such as a defucosylated Fc (defucosylated antibody), and a bispecific antibody or antigen-binding fragment thereof. In some embodiments, the antigen-binding fragment of the anti-CD8 α antibody is a diabody, a triabody, or a tetraabody.

[0156] In some embodiments, the anti-CD8 α antibody, or the antigen-binding fragment thereof, is from the IgG class.

The anti-CD8 α antibody, or antigen-binding fragment thereof, may thus be from the IgG1, IgG2, IgG3, or IgG4 subclass.

[0157] In some embodiments, the anti-CD8 α antibody, or the antigen-binding fragment thereof, is from the IgG1 subclass, preferably from the human IgG1 subclass. In some embodiments, the anti-CD8 α antibody, or the antigen-binding fragment thereof, is thus an IgG1 antibody, preferably a human IgG1 antibody or a chimeric human IgG1 antibody.

[0158] In some embodiments, the anti-CD8 α antibody, or the antigen-binding fragment thereof, is from the IgG2 subclass, preferably from the human IgG2 subclass. In some embodiments, the anti-CD8 α antibody, or the antigen-binding fragment thereof, is thus an IgG2 antibody, preferably a human IgG2 antibody or a chimeric human IgG2 antibody. In particular, the IgG2 antibody may be a IgG2b antibody.

[0159] As used herein, a "chimeric antibody" refers to an antibody or antigen-binding fragment thereof comprising a first amino acid sequence linked to a second amino acid sequence with which it is not naturally linked in nature. The amino acid sequences may normally exist in separate proteins that are brought together in the chimeric (or fusion) protein or may normally exist in the same protein but are placed in a new arrangement in the chimeric (or fusion) protein. A chimeric protein may be created, for example, by chemical synthesis, or by creating and translating a polynucleotide in which the peptide regions are encoded in the desired relationship. The term "chimeric antibody" comprises antibodies and antigen-binding fragment thereof in which:

[0160] (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the variable region is linked to a constant region of a different or altered class, effector function and/or species; or

[0161] (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region, or portion thereof, having a different or altered antigen specificity; or with corresponding sequences from another species or from another antibody class or subclass.

[0162] In the present invention, unless otherwise specified, the position of the complementary-determining regions (CDRs) is determined using the Kabat nomenclature.

[0163] According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein comprises a variable region of the heavy chain (also referred to as heavy chain variable region or VH) which comprises at least one, preferably at least two, more preferably the three following CDRs:

[0164] V_H-CDR1: NX₁X₂MN, wherein

[0165] X₁ is N or Y,

[0166] X₂ is D or A or Y;

[0167] V_H-CDR2: X₃ISGSSX₄YIX₅YADFVKG (SEQ ID NO: 1), wherein

[0168] X₃ is D or S,

[0169] X₄ is S or R,

[0170] X₅ is D or G or Y; and/or

[0171] V_H-CDR3: SSX₆X₇X₈X₉YX₁₀X₁₁X₁₂X₁₃MDV (SEQ ID NO: 2), wherein

[0172] X₆ is N or no amino acid,

[0173] X₇ is Y or no amino acid,

[0174] X₈ is Y or G or no amino acid,

[0175] X₉ is D or S or no amino acid,

[0176] X₁₀ is S or N or F,

[0177] X_1 is A or G,

[0178] X_{12} is S or D or N,

[0179] X_{13} is A or G.

[0180] Examples of V_H -CDR1 having an amino acid sequence corresponding to NX_1X_2MN as described above include, without being limited to: NNAMN (SEQ ID NO: 6), NYDMN (SEQ ID NO: 12), and NYYMN (SEQ ID NO: 18). Thus, in some embodiments, the V_H -CDR1 having an amino acid sequence as set forth in sequence NX_1X_2MN is selected from the group comprising or consisting of SEQ ID NO: 6, SEQ ID NO: 12, and SEQ ID NO: 18.

[0181] Examples of V_H -CDR2 having an amino acid sequence as set forth in SEQ ID NO: 1 as described above include, without being limited to, DISGSSRYIGYADFVK (SEQ ID NO: 7), DISGSSSYIDYADFVK (SEQ ID NO: 13), and SISGSSRYIYYADFVK (SEQ ID NO: 19). Thus, in some embodiments, the V_H -CDR2 having an amino acid sequence as set forth in SEQ ID NO: 1 is selected from the group comprising or consisting of SEQ ID NO: 7, SEQ ID NO: 13, and SEQ ID NO: 19.

[0182] Examples of V_H -CDR3 having an amino acid sequence as set forth in SEQ ID NO: 2 as described above include, without being limited to, SSYYSGSGMDV (SEQ ID NO: 8), SSYYSGSGMDV (SEQ ID NO: 14), and SSGSYFGNAMDV (SEQ ID NO: 20). Thus, in some embodiments, the V_H -CDR3 having an amino acid sequence as set forth in SEQ ID NO: 2 is selected from the group comprising or consisting of SEQ ID NO: 8, SEQ ID NO: 14, and SEQ ID NO: 20.

[0183] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

[0184] a V_H -CDR1 selected from the group comprising or consisting of NNAMN (SEQ ID NO: 6), NYDMN (SEQ ID NO: 12), and NYYMN (SEQ ID NO: 18);

[0185] a V_H -CDR2 selected from the group comprising or consisting of DISGSSRYIGYADFVK (SEQ ID NO: 7), DISGSSSYIDYADFVK (SEQ ID NO: 13), and SISGSSRYIYYADFVK (SEQ ID NO: 19); and/or

[0186] a V_H -CDR3 selected from the group comprising or consisting of SSYYDYDYNADAMDV (SEQ ID NO: 8), SSYYSGSGMDV (SEQ ID NO: 14), and SSGSYFGNAMDV (SEQ ID NO: 20).

[0187] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

- V_H -CDR1: (SEQ ID NO: 6)
NNAMN;

- V_H -CDR2: (SEQ ID NO: 7)
DISGSSRYIGYADFVK;
and/or

- V_H -CDR3: (SEQ ID NO: 8)
SSYYDYDYNADAMDV.

[0188] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a

V_H which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

- V_H -CDR1: (SEQ ID NO: 12)
NYDMN;

- V_H -CDR2: (SEQ ID NO: 13)
DISGSSSYIDYADFVK;
and/or

- V_H -CDR3: (SEQ ID NO: 14)
SSYYSGSGMDV.

[0189] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

V_H -CDR1: (SEQ ID NO: 18)
NYYMN;

 V_H -CDR2: (SEQ ID NO: 19)
SISGSSRYIYYADFVK;
and/or

 V_H -CDR3: (SEQ ID NO: 20)
SSGSYFGNAMDV

[0190] In some embodiments, V_H -CDR1, V_H -CDR2 and/or V_H -CDR3 have an amino acid sequence as set forth in any one of SEQ ID NOs 6-8, 12-14, and 18-20, as described above with 1, 2, or 3, or more amino acid(s) being substituted by a different amino acid. For example, in some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H -CDR1 having an amino acid sequence as set forth in SEQ ID NO: 6, SEQ ID NO: 12, or in SEQ ID NO: 18 with 1, 2, or 3, or more amino acid(s) being substituted by a different amino acid. In some embodiments, V_H -CDR1, V_H -CDR2 and/or V_H -CDR3 as described above have an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the corresponding amino acid sequence as set forth in any one of SEQ ID NOs 6-8, 12-14, and 18-20. For example, in some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H -CDR1 having an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence as set forth in SEQ ID NO: 6, SEQ ID NO: 12, or in SEQ ID NO: 18.

[0191] According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a variable region of the light chain (also referred to as light chain variable region or VL) which comprises at least one, preferably at least two, more preferably the three following CDRs:

[0192] VL-CDR1: AGTSSDVGGX₁₄X₁₅X₁₆VS (SEQ ID NO: 3), wherein

[0193] X_{14} is G or N or Y,

[0194] X_{15} is S or Y,

[0195] X_{16} is S or Y;

[0196] VL-CDR2: X₁₇DSX₁₈RPS (SEQ ID NO: 4), wherein

[0197] X₁₇ is Q or S or Y,

[0198] X₁₈ is Y or S; and/or

[0199] VL-CDR3: SSX₁₉TX₂₀YSTRV (SEQ ID NO: 5), wherein

[0200] X₁₉ is Y or D,

[0201] X₂₀ is Y or Q or S.

[0202] Examples of V_L-CDR1 having an amino acid sequence as set forth in SEQ ID NO: 3 as described above include, without being limited to, AGTSSDVGGNSYVS (SEQ ID NO: 9), AGTSSDVGGGSSVS (SEQ ID NO: 15), and AGTSSDVGGYYSVS (SEQ ID NO: 21). Thus, in some embodiments, the V_L-CDR1 having an amino acid sequence as set forth in SEQ ID NO: 3 is selected from the group comprising or consisting of SEQ ID NO: 9, SEQ ID NO: 15, and SEQ ID NO: 21.

[0203] Examples of V_L-CDR2 having an amino acid sequence as set forth in SEQ ID NO: 4 as described above include, without being limited to, SDSSRPS (SEQ ID NO: 10), QDSYRPS (SEQ ID NO: 16), and YDSSRPS (SEQ ID NO: 22). Thus, in some embodiments, the V_L-CDR2 having an amino acid sequence as set forth in SEQ ID NO: 4 is selected from the group comprising or consisting of SEQ ID NO: 10, SEQ ID NO: 16, and SEQ ID NO: 22.

[0204] Examples of V_L-CDR3 having an amino acid sequence as set forth in SEQ ID NO: 5 as described above include, without being limited to, SSYTQYSTRV (SEQ ID NO: 11), SSYTYSTRV (SEQ ID NO: 17), and SSDTSYSTRV (SEQ ID NO: 23). Thus, in some embodiments, the V_L-CDR3 having an amino acid sequence as set forth in SEQ ID NO: 5 is selected from the group comprising or consisting of SEQ ID NO: 11, SEQ ID NO: 17, and SEQ ID NO: 23.

[0205] In some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a VL which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

[0206] a V_L-CDR1 selected from the group comprising or consisting of AGTSSDVGGNSYVS (SEQ ID NO: 9), AGTSSDVGGGSSVS (SEQ ID NO: 15), and AGTSSDVGGYYSVS (SEQ ID NO: 21);

[0207] a V_L-CDR2 selected from the group comprising or consisting of SDSSRPS (SEQ ID NO: 10), QDSYRPS (SEQ ID NO: 16), and YDSSRPS (SEQ ID NO: 22); and/or

[0208] a V_L-CDR3 selected from the group comprising or consisting of SSYTQYSTRV (SEQ ID NO: 11), SSYTYSTRV (SEQ ID NO: 17), and SSDTSYSTRV (SEQ ID NO: 23).

[0209] In some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a VL which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

V_L-CDR1: (SEQ ID NO: 9)
AGTSSDVGGNSYVS;
V_L-CDR2: (SEQ ID NO: 10)
SDSSRPS;

-continued
and/or

V_L-CDR3: (SEQ ID NO: 11)

SSYTQYSTRV.

[0210] In some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a VL which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

V_L-CDR1: (SEQ ID NO: 15)

AGTSSDVGGGSSVS;

V_L-CDR2: (SEQ ID NO: 16)

QDSYRPS;
and/or

V_L-CDR3: (SEQ ID NO: 17)

SSYTYSTRV

[0211] In some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a VL which comprises at least one (e.g., 1, 2 or 3) of the following CDRs, and preferably the three following CDRs:

V_L-CDR1: (SEQ ID NO: 21)

AGTSSDVGGYYSVS;

V_L-CDR2: (SEQ ID NO: 22)

YDSSRPS;
and/or

V_L-CDR3: (SEQ ID NO: 23)

SSDTSYSTRV.

[0212] In some embodiments, V_L-CDR1, V_L-CDR2 and/or V_L-CDR3 have an amino acid sequence as set forth in any one of SEQ ID NOs 9-11, 15-17, and 21-23 as described above with 1, 2, 3 or more amino acid(s) being substituted by a different amino acid. For example, in some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a V_L-CDR1 having an amino acid sequence as set forth in SEQ ID NO: 9, SEQ ID NO: 15, or in SEQ ID NO: 21 with 1, 2, or 3, or more amino acid(s) being substituted by a different amino acid. In some embodiments, V_L-CDR1, V_L-CDR2 and/or V_L-CDR3 as described above have an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the corresponding amino acid sequence as set forth in any one of SEQ ID NOs 9-11, 15-17, and 21-23. For example, in some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises a V_L-CDR1 having an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence as set forth in SEQ ID NO: 9, SEQ ID NO: 15, or in SEQ ID NO: 21.

[0213] In some embodiments, the isolated anti-CD8α antibody, or the antigen-binding fragment thereof, comprises: a variable region of the heavy chain (VH) which comprises at least one, preferably at least two, more preferably the three following CDRs:

- [0214] V_H -CDR1: NX_1X_2MN , wherein
- [0215] X_1 is N or Y,
- [0216] X_2 is D or A or Y;
- [0217] V_H -CDR2: $X_3ISGSSX_4YIX_5YADFVKG$ (SEQ ID NO: 1), wherein
- [0218] X_3 is D or S,
- [0219] X_4 is S or R,
- [0220] X_5 is D or G or Y; and/or
- [0221] V_H -CDR3: $SSX_6X_7X_8X_9YX_{10}X_{11}X_{12}X_{13}MDV$ (SEQ ID NO: 2), wherein
- [0222] X_6 is N or no amino acid,
- [0223] X_7 is Y or no amino acid,
- [0224] X_8 is Y or G or no amino acid,
- [0225] X_9 is D or S or no amino acid,
- [0226] X_{10} is S or N or F,
- [0227] X_{11} is A or G,
- [0228] X_{12} is S or D or N,
- [0229] X_{13} is A or G; and
- a variable region of the light chain (VL) which comprises at least one, preferably at least two, more preferably the three following CDRs:
- [0230] VL-CDR1: $AGTSSDVGGX_{14}X_{15}X_{16}VS$ (SEQ ID NO: 3), wherein
- [0231] X_{14} is G or N or Y,
- [0232] X_{15} is S or Y,
- [0233] X_{16} is S or Y;
- [0234] VL-CDR2: $X_{17}DSX_{18}RPS$ (SEQ ID NO: 4), wherein
- [0235] X_{17} is Q or S or Y,
- [0236] X_{18} is Y or S; and/or
- [0237] VL-CDR3: $SSX_{19}TX_{20}YSTRV$ (SEQ ID NO: 5), wherein
- [0238] X_{19} is Y or D,
- [0239] X_{20} is Y or Q or S.
- [0240] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises: a V_H which comprises the three following CDRs:
- [0241] V_H -CDR1: NX_1X_2MN , wherein
- [0242] X_1 is N or Y,
- [0243] X_2 is D or A or Y;
- [0244] V_H -CDR2: $X_3ISGSSX_4YIX_5YADFVKG$ (SEQ ID NO: 1), wherein
- [0245] X_3 is D or S,
- [0246] X_4 is S or R,
- [0247] X_5 is D or G or Y; and
- [0248] V_H -CDR3: $SSX_6X_7X_8X_9YX_{10}X_{11}X_{12}X_{13}MDV$ (SEQ ID NO: 2), wherein
- [0249] X_6 is N or no amino acid,
- [0250] X_7 is Y or no amino acid,
- [0251] X_8 is Y or G or no amino acid,
- [0252] X_9 is D or S or no amino acid,
- [0253] X_{10} is S or N or F,
- [0254] X_{11} is A or G,
- [0255] X_{12} is S or D or N,
- [0256] X_{13} is A or G; and
- a VL which comprises the three following CDRs:
- [0257] VL-CDR1: $AGTSSDVGGX_{14}X_{15}X_{16}VS$ (SEQ ID NO: 3), wherein
- [0258] X_{14} is G or N or Y,
- [0259] X_{15} is S or Y,
- [0260] X_{16} is S or Y;

- [0261] VL-CDR2: $X_{17}DSX_{18}RPS$ (SEQ ID NO: 4), wherein
- [0262] X_{17} is Q or S or Y,
- [0263] X_{18} is Y or S,
- [0264] VL-CDR3: $SSX_{19}TX_{20}YSTRV$ (SEQ ID NO: 5), wherein
- [0265] X_{19} is Y or D,
- [0266] X_{20} is Y or Q or S.
- [0267] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises: a V_H which comprises the three following CDRs:
- [0268] a V_H -CDR1 selected from the group comprising or consisting of NNAMN (SEQ ID NO: 6), NYDMN (SEQ ID NO: 12), and NYYMN (SEQ ID NO: 18);
- [0269] a V_H -CDR2 selected from the group comprising or consisting of DISGSSRYIGYADFVKG (SEQ ID NO: 7), DISGSSSYIDYADFVKG (SEQ ID NO: 13), and SISGSSRYIYYADFVKG (SEQ ID NO: 19); and
- [0270] a V_H -CDR3 selected from the group comprising or consisting of SSNYDYDNADAMDV (SEQ ID NO: 8), SSYYSGSGMDV (SEQ ID NO: 14), and SSGSYFGNAMDV (SEQ ID NO: 20); and
- a VL which comprises the three following CDRs:
- [0271] a V_L -CDR1 selected from the group comprising or consisting of AGTSSDVGGNSYVS (SEQ ID NO: 9), AGTSSDVGGGSSVS (SEQ ID NO: 15), and AGTSSDVGGYYVS (SEQ ID NO: 21);
- [0272] a V_L -CDR2 selected from the group comprising or consisting of SDSSRPS (SEQ ID NO: 10), QDSYRPS (SEQ ID NO: 16), and YDSSRPS (SEQ ID NO: 22); and
- [0273] a V_L -CDR3 selected from the group comprising or consisting of SSYTQYSTRV (SEQ ID NO: 11), SSYTYSTRV (SEQ ID NO: 17), and SSDTSYSTRV (SEQ ID NO: 23).

[0274] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises the six following CDRs:

V_H -CDR1:	(SEQ ID NO: 6)
NNAMN,	
V_H -CDR2:	(SEQ ID NO: 7)
DISGSSRYIGYADFVKG,	
V_H -CDR3:	(SEQ ID NO: 8)
SSNYDYDNADAMDV,	
V_L -CDR1:	(SEQ ID NO: 9)
AGTSSDVGGNSYVS,	
V_L -CDR2:	(SEQ ID NO: 10)
SDSSRPS,	
and	
V_L -CDR3:	(SEQ ID NO: 11);
SSYTQYSTRV,	
or	
V_H -CDR1:	(SEQ ID NO: 12)
NYDMN,	

-continued

V_H -CDR2: (SEQ ID NO: 13)
 DISGSSSYIDYADFVKG,
 V_H -CDR3: (SEQ ID NO: 14)
 SSYYSGSGMDV,
 V_L -CDR1: (SEQ ID NO: 15)
 AGTSSDVGGGSSVS,
 V_L -CDR2: (SEQ ID NO: 16)
 QDSYRPS,
 and
 V_L -CDR3: (SEQ ID NO: 17);
 SSYTYSTRV,
 or
 V_H -CDR1: (SEQ ID NO: 18)
 NYDMN,
 V_H -CDR2: (SEQ ID NO: 19)
 SISGSSRYIYADFVKG,
 V_H -CDR3: (SEQ ID NO: 20)
 SSGSYFGNAMDV,
 V_L -CDR1: (SEQ ID NO: 21)
 AGTSSDVGGYYSVS,
 V_L -CDR2: (SEQ ID NO: 22)
 YDSSRPS,
 and
 V_L -CDR3: (SEQ ID NO: 23)
 SSDTSYSTRV.

[0275] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

a V_H which comprises the three following CDRs:

V_H -CDR3: (SEQ ID NO: 8)
 SSNYDYNDAMDV,
 V_H -CDR1: (SEQ ID NO: 6)
 NNAMN,
 V_H -CDR2: (SEQ ID NO: 7)
 DISGSSRYIGYADFVKG,
 and

and

a V_L which comprises the three following CDRs:

V_L -CDR1: (SEQ ID NO: 9)
 AGTSSDVGGNSYVS,

-continued

V_L -CDR2: (SEQ ID NO: 10)
 SDSSRPS,
 and
 V_L -CDR3: (SEQ ID NO: 11)
 SSYTYSTRV

[0276] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs having an amino acid sequence as set forth in SEQ ID NOs: 6-11 as described above with 1, 2, 3 or more amino acid(s) being substituted by a different amino acid. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs as described above having an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the corresponding amino acid sequence as set forth in SEQ ID NOs: 6-11.

[0277] An example of anti-CD8 α antibody comprising a V_H comprising V_H -CDR1, V_H -CDR2 and V_H -CDR3 as set forth in SEQ ID NOs: 6, 7 and 8, respectively, and a V_L comprising V_L -CDR1, V_L -CDR2 and V_L -CDR3 as set forth in SEQ ID NOs: 9, 10 and 11, respectively, is hMP08-R3-F08.

[0278] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises: a V_H which comprises the three following CDRs:

V_H -CDR3: (SEQ ID NO: 14)
 SSYYSGSGMDV;
 V_H -CDR1: (SEQ ID NO: 12)
 NYDMN,
 V_H -CDR2: (SEQ ID NO: 13)
 DISGSSSYIDYADFVKG,
 and

and a V_L which comprises the three following CDRs:

V_L -CDR1: (SEQ ID NO: 15)
 AGTSSDVGGGSSVS,
 V_L -CDR2: (SEQ ID NO: 16)
 QDSYRPS,
 and
 V_L -CDR3: (SEQ ID NO: 17)
 SSYTYSTRV

[0279] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs having an amino acid sequence as set forth in SEQ ID NOs: 12-17 as described above with 1, 2, 3 or more amino acid(s) being substituted by a different amino acid. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs as described above having an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the corresponding amino acid sequence as set forth in SEQ ID NOs: 12-17.

[0280] An example of anti-CD8 α antibody comprising a V_H comprising V_H -CDR1, V_H -CDR2 and V_H -CDR3 as set forth in SEQ ID NOs: 12, 13 and 14, respectively, and a VL comprising V_L -CDR1, V_L -CDR2 and V_L -CDR3 as set forth in SEQ ID NOs: 15, 16 and 17, respectively, is hMP08-R3-C11.

[0281] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises: a V_H which comprises the three following CDRs:

V_H -CDR1: (SEQ ID NO: 18)
 NYYYMN,
 V_H -CDR2: (SEQ ID NO: 19)
 SISGSSRYIYYADFVKG,
 and
 V_H -CDR3: (SEQ ID NO: 20)
 SSGSYFGNAMDV;

and a VL which comprises the three following CDRs:

V_L -CDR1: (SEQ ID NO: 21)
 AGTSSDVGGYYSVS,
 V_L -CDR2: (SEQ ID NO: 22)
 YDSSRPS,
 and
 V_L -CDR3: (SEQ ID NO: 23)
 SSDTSYSTRV

[0282] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs having an amino acid sequence as set forth in SEQ ID NOs: 18-23 as described above with 1, 2, 3 or more amino acid(s) being substituted by a different amino acid. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises CDRs as described above having an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the corresponding amino acid sequence as set forth in SEQ ID NOs: 18-23.

[0283] An example of anti-CD8 α antibody comprising a V_H comprising V_H -CDR1, V_H -CDR2 and V_H -CDR3 as set forth in SEQ ID NOs: 18, 19 and 20, respectively, and a VL comprising V_L -CDR1, V_L -CDR2 and V_L -CDR3 as set forth in SEQ ID NOs: 21, 22 and 23, respectively, is hMP09-R3-D03.

[0284] According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a variable region of the heavy chain (V_H) comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28.

[0285] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 24, SEQ ID

NO: 26, and SEQ ID NO: 28, with 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more amino acid(s) substituted by a different amino acid. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 24, SEQ ID NO: 26, and SEQ ID NO: 28, with 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more amino acid(s) substituted by a different amino acid, wherein said amino acid substitution(s) do(es) not occur in any of the three CDRs (V_H -CDR1, V_H -CDR2, and V_H -CDR3).

[0286] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising or consisting of an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising three CDRs (V_H -CDR1, V_H -CDR2, and V_H -CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28.

[0287] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a V_H comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 24, SEQ ID NO: 26, and SEQ ID NO: 28.

[0288] According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a variable region of the light chain (VL) comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29 and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0289] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, and SEQ ID NO: 29, with 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more amino acid(s) substituted by a different amino acid. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, and SEQ ID NO: 29, with 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16,

17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35 or more amino acid(s) substituted by a different amino acid, wherein said amino acid substitution(s) do(es) not occur in any of the three CDRs (V_L —CDR1, V_L —CDR2, and V_L —CDR3).

[0290] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising or consisting of an amino acid sequence that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29. In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising three CDRs (V_L —CDR1, V_L —CDR2, and V_L —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0291] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, and SEQ ID NO: 29.

[0292] According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0293] a variable region of the heavy chain (V_H) comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28; and

[0294] a variable region of the light chain (VL) comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29 and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0295] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0296] a V_H comprising three CDRs (V_H —CDR1, V_H —CDR2, and V_H —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28; and

[0297] a VL comprising three CDRs (V_L —CDR1, V_L —CDR2, and V_L —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%,

85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0298] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0299] a V_H comprising or consisting of a sequence as set forth in SEQ ID NO: 24 or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24; and

[0300] a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0301] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0302] a V_H comprising three CDRs (V_H —CDR1, V_H —CDR2, and V_H —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 24; and

[0303] a VL comprising three CDRs (V_L —CDR1, V_L —CDR2, and V_L —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0304] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0305] a V_H comprising or consisting of a sequence as set forth in SEQ ID NO: 26 or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 26; and

[0306] a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29 and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

[0307] In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:

[0308] a V_H comprising three CDRs (V_H —CDR1, V_H —CDR2, and V_H —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 26; and

[0309] a VL comprising three CDRs (V_L —CDR1, V_L —CDR2, and V_L —CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework

- regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.
- [0310]** In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0311]** a V_H comprising or consisting of a sequence as set forth in SEQ ID NO: 28 or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 28; and
- [0312]** a VL comprising or consisting of a sequence selected from the group comprising or consisting of SEQ ID NO: 25, SEQ ID NO: 27, SEQ ID NO: 29, and sequences having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.
- [0313]** In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0314]** a V_H comprising three CDRs (V_H -CDR1, V_H -CDR2, and V_H -CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 28; and
- [0315]** a VL comprising three CDRs (V_L -CDR1, V_L -CDR2, and V_L -CDR3) as described above and an amino acid sequence of the framework regions (i.e., the non-CDR regions) that shares at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with the amino acid sequence of the framework regions (i.e., the non-CDR regions) of SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.
- [0316]** According to some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0317]** a V_H comprising or consisting of SEQ ID NO: 24, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24, and a VL comprising or consisting of SEQ ID NO: 25, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25; or
- [0318]** a V_H comprising or consisting of SEQ ID NO: 26, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 26, and a VL comprising or consisting of SEQ ID NO: 27, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 27; or
- [0319]** a V_H comprising or consisting of SEQ ID NO: 28, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 28, and a VL comprising or consisting of SEQ ID NO: 29, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 29.
- [0320]** In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0321]** a V_H comprising or consisting of SEQ ID NO: 24, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 24; and

- [0322]** a VL comprising or consisting of SEQ ID NO: 25, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 25.
- [0323]** An example of anti-CD8 α antibody comprising a V_H as set forth in SEQ ID NO: 24, and a VL as set forth in SEQ ID NO: 25, is hMP08-R3-F08.
- [0324]** In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0325]** a V_H comprising or consisting of SEQ ID NO: 26, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 26; and
- [0326]** a VL comprising or consisting of SEQ ID NO: 27, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 27.
- [0327]** An example of anti-CD8 α antibody comprising a V_H as set forth in SEQ ID NO: 26, and a VL as set forth in SEQ ID NO: 27, is hMP08-R3-C11.
- [0328]** In some embodiments, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, comprises:
- [0329]** a V_H comprising or consisting of SEQ ID NO: 28, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 28; and
- [0330]** a VL comprising or consisting of SEQ ID NO: 29, or a sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 29.
- [0331]** An example of anti-CD8 α antibody comprising a V_H as set forth in SEQ ID NO: 28, and a VL as set forth in SEQ ID NO: 29, is hMP09-R3-D03.
- [0332]** Another object of the present invention is a fusion protein comprising the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein.
- [0333]** Another object of the present invention is a nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein. Another object of the present invention is a nucleic acid encoding the fusion protein as described herein.
- [0334]** In some embodiments, the nucleic acid is an isolated nucleic acid. In some embodiments, an “isolated nucleic acid” refers to a nucleic acid that is substantially separated from other nucleic acid sequences as well as proteins or complexes such as ribosomes and polymerases, which naturally accompany a native sequence. The term embraces a nucleic acid sequence that has been removed from its naturally occurring environment, and includes recombinant or cloned nucleic acid, chemically synthesized analogues, and analogues biologically synthesized by heterologous systems. A substantially pure nucleic acid includes isolated forms of the nucleic acid. This refers to the nucleic acid as originally isolated and does not exclude nucleic acid sequences later added to the isolated nucleic acid by the hand of man.
- [0335]** In some embodiments, the isolated nucleic acid is purified.
- [0336]** In some embodiments, the isolated nucleic acid is purified to (i) greater than about 80%, 85%, 90%, 91%, 92%, 93%, 94%, or 95%, or more by weight of nucleic acid as determined by absorbance methods or fluorescence methods (such as, for example, by measuring the ratio of absorbance at 260 and 280 nm (A₂₆₀/A₂₈₀)), and most preferably to

greater than about 96%, 97%, 98%, or 99% by weight; or (ii) homogeneity as shown by agarose gel electrophoresis and using an intercalating agent such as, for example, ethidium bromide, SYBR Green, GelGreen.

[0337] In some embodiments, the nucleic acid encodes at least a heavy chain variable region (V_H) and/or a light chain variable region (VL) of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein. In some embodiments, the nucleic acid encodes at least a V_H and a VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein on separate nucleic acid molecules or on the same nucleic acid molecule. The nucleic acid may encode variable and constant regions of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein. The nucleic acid may encode heavy and light chains of the anti-CD8 α antibody, or the antigen-binding fragment thereof, on separate nucleic acid molecules or on the same nucleic acid molecule.

[0338] According to some embodiments, the nucleic acid comprises or consists of a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described above.

[0339] According to some embodiments, the nucleic acid comprises or consists of a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described above.

[0340] According to some embodiments, the nucleic acid comprises or consists of:

[0341] a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described above; and/or

[0342] a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described above.

[0343] According to some embodiments, the nucleic acid comprises or consists of:

[0344] a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being any one of SEQ ID NOs: 31, 33, or 35, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NOs: 31, 33, or 35; and/or

[0345] a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being any one of SEQ ID NOs: 32, 34, or 36, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NOs: 32, 34, or 36.

[0346] According to some embodiments, the nucleic acid comprises or consists of:

[0347] a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 31, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 31, and/or a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 32, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 32; or

[0348] a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 33, or any sequence

sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 33, and/or a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 34, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 34; or

[0349] a sequence encoding the V_H of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 35, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 35, and/or a sequence encoding the VL of the anti-CD8 α antibody, or the antigen-binding fragment thereof, said sequence being SEQ ID NO: 36, or any sequence sharing at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%, or more identity with SEQ ID NO: 36.

[0350] Typically, the nucleic acid described herein is a DNA (deoxyribonucleic acid) or RNA (ribonucleic acid) molecule, which may be included in any suitable vector.

[0351] Another object of the present invention is a vector, such as, for example, an expression vector or a cloning vector, comprising a nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein.

[0352] The terms “vector”, “cloning vector” and “expression vector” mean the vehicle by which a nucleic acid sequence (e.g., a DNA molecule) encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein can be introduced into a host cell, so as to transform or transfect the host cell and promote expression (e.g., transcription and/or translation) of the introduced nucleic acid sequence.

[0353] In some embodiments, the vector, cloning vector, or expression vector, is a plasmid, a cosmid, an episome, an artificial chromosome, a phage, or a viral vector.

[0354] Examples of plasmids include replicating plasmids comprising an origin of replication, or integrative plasmids, such as for instance pUC, pcDNA, and pBR.

[0355] Examples of viral vector include adenoviral, adeno-associated virus (AAV), retroviral, herpes virus vectors. Such viral vectors may be produced by techniques known in the art, such as by transfecting packaging cells or by transient transfection with helper plasmids or viruses.

[0356] Any expression vector for animal cells can be used, so long as a nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein can be inserted and expressed.

[0357] The vector as described herein may comprise regulatory elements, such as a promoter, an enhancer, and/or a terminator, for example to cause or direct expression of the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein upon introduction within a host cell or upon administration to a subject. Examples of promoters and enhancers used in expression vectors for animal cells include, but are not limited to, early promoter and enhancer of SV40, LTR promoter and enhancer of Moloney mouse leukemia virus, and promoter and enhancer of immunoglobulin H chain.

[0358] In some embodiments, the vector as described herein comprises a nucleic acid sequence encoding the variable region of the heavy chain (V_H) of the anti-CD8 α

antibody, or the antigen-binding fragment thereof, as described above operably linked to regulatory elements. In some embodiments, the vector as described above comprises a sequence encoding the variable region of the light chain (VL) of the anti-CD8 α antibody, or the antigen-binding fragment thereof, as described above operably linked to regulatory elements.

[0359] In some embodiments, the vector as described above is monocistronic. By “monocistronic”, it is meant that a single nucleic acid is expressed with a single vector. In some embodiments, the vector as described above is polycistronic. By “polycistronic”, it is meant that at least two or more nucleic acid sequences are expressed with a single vector.

[0360] Another object of the invention is a host cell comprising the vector as described herein. In some embodiments, the host cell is an isolated host cell. Said host cell may be used for the recombinant production of the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein.

[0361] Host cells may be prokaryote cells, or eukaryote cells, such as, for example, yeast or mammalian cells.

[0362] It should be noted that, with regards to animal and human cells, the term “host cell” generally refers to a cell of a cultured cell line. Animals and human beings into whom a vector encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein, has been introduced are explicitly excluded from the definition of a “host cell”.

[0363] Another object of the present invention is a method of producing, and optionally purifying, the isolated anti-CD8 α antibody, or the antigen-binding fragment thereof, as described herein.

[0364] In some embodiments, the method comprises:

[0365] introducing in vitro or ex vivo a nucleic acid or a vector as described herein into a host cell (that is to say transforming or transfecting a host cell with a nucleic acid or a vector as described herein);

[0366] culturing in vitro or ex vivo host cells transformed or transfected with the nucleic acid or with the vector, under conditions suitable for expression of the anti-CD8 α antibody, or the antigen-binding fragment thereof;

[0367] optionally, selecting the host cells which express and/or secrete the anti-CD8 α antibody, or the antigen-binding fragment thereof; and

[0368] recovering the expressed and/or secreted anti-CD8 α antibody, or the expressed and/or secreted antigen-binding fragment thereof.

[0369] Such a method is well-known in the art and can be used for large scale production of antibodies or antigen-binding fragments thereof, including monoclonal antibodies intended for in vitro, ex vivo and/or in vivo uses, such as therapeutic uses.

[0370] In some embodiments, the recovered anti-CD8 α antibody, or the recovered antigen-binding fragment thereof, is further purified. Methods for purifying an antibody, or an antigen-binding fragment thereof, are well-known in the art and include, without limitation, use of an anti-CH1 antibody, protein A-Sepharose, gel electrophoresis, chromatography, in particular affinity chromatography.

[0371] Another object of the present invention is a composition comprising, consisting essentially of, or consisting of:

[0372] at least one anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein;

[0373] at least one fusion protein as described herein,

[0374] at least one nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein;

[0375] at least one vector comprising such a nucleic acid molecule; or

[0376] at least one host cell comprising such a vector.

[0377] As used herein, “consisting essentially of”, with reference to a composition, means that the at least one anti-CD8 α antibody, or the antigen-binding fragment thereof, fusion protein nucleic acid, vector, or host cell is the only active agent, therapeutic agent, or agent with a biologic activity within said composition.

[0378] Another object of the present invention is a pharmaceutical composition comprising, consisting essentially of, or consisting of:

[0379] at least one anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein;

[0380] at least one fusion protein as described herein,

[0381] at least one nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein;

[0382] at least one vector comprising such a nucleic acid molecule; or

[0383] at least one host cell comprising such a vector;

[0384] and at least one pharmaceutically acceptable excipient.

[0385] Pharmaceutically acceptable excipients that may be used in the pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances (for example sodium carboxymethyl-cellulose), polyethylene glycol, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat.

[0386] Another object of the present invention is a medicament comprising, consisting essentially of, or consisting of:

[0387] at least one anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein;

[0388] at least one fusion protein as described herein,

[0389] at least one nucleic acid encoding the anti-CD8 α antibody, or the antigen-binding fragment thereof, or the fusion protein as described herein;

[0390] at least one vector comprising such a nucleic acid molecule; or

[0391] at least one host cell comprising such a vector.

[0392] In some embodiments, the composition, pharmaceutical composition, or medicament as described herein are formulated for administration to a subject.

[0393] The composition, pharmaceutical composition, or medicament as described herein may be administered or formulated to be administered systemically, orally, parenterally, by injection, topically, by inhalation spray, rectally, nasally, or via an implanted reservoir. The term administration used herein includes subcutaneous, intravenous, intra-

muscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques.

[0394] In some embodiments, the composition, pharmaceutical composition, or medicament as described herein are formulated for administration by injection, such as for example subcutaneous injection, or by infusion, such as for example intravenous infusion. Examples of forms adapted for injection include, but are not limited to, solutions, such as, for example, sterile aqueous solutions, gels, dispersions, emulsions, suspensions, solid forms suitable for using to prepare solutions or suspensions upon the addition of a liquid prior to use, such as, for example, powder, liposomal forms and the like.

[0395] In some embodiments, regimens or dosages used for administration of the composition, pharmaceutical composition, or medicament can be adapted as function of various parameters, and in particular as a function of the mode of administration used, of the relevant pathology, or of the desired duration of treatment. For example, it is well within the skill of the art to start with a dose at a level lower than those required to achieve the desired therapeutic effect and to gradually increase the dose of anti-CD8 α antibody, an antigen-binding fragment thereof, or fusion protein as described herein until the desired effect is achieved. The dose of anti-CD8 α antibody, an antigen-binding fragment thereof, or fusion protein as described herein may be varied over a wide range from 0.01 to 10 000 mg per adult per day. A composition, pharmaceutical composition or medicament typically may for example comprise from about 0.01 mg to about 10 000 mg, preferably from about 0.1 mg to about 5000 mg of anti-CD8 α antibody, an antigen-binding fragment thereof, or fusion protein as described herein.

[0396] Another object of the present invention is an anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein, for use as a medicament. Another object of the present invention is a fusion protein as described herein for use as a medicament.

[0397] Another object of the present invention is a nucleic acid encoding an anti-CD8 α antibody, or an antigen-binding fragment thereof, or a fusion protein as described herein, or a vector comprising such a nucleic acid molecule, or a host cell comprising such a vector, for use as a medicament.

[0398] Another object of the present invention is an anti-CD8 α antibody, or antigen binding fragment thereof, as described herein for use in the treatment of a CD8- related disease.

[0399] As used herein "CD8- related disease" refers to a disease, disorder or condition caused by or potentialized by an increased proportion of cells expressing CD8, in particular cells expressing CD8 α , in a subject and/or by an increased level of expression of CD8, in particular CD8 α , in cells of the subject.

[0400] Another object of the present invention is a fusion protein as described herein for use in the treatment of a CD8- related disease.

[0401] Another object of the present invention is a nucleic acid encoding an anti-CD8 α antibody, or antigen binding fragment thereof, or a fusion protein as described herein, or

a vector comprising such a nucleic acid molecule, or a host cell comprising such a vector, for use in the treatment of a CD8- related disease.

[0402] In some embodiments, the CD8- related disease is a cardiovascular disease. In some embodiments the cardiovascular disease is myocardial infarction or acute myocardial infarction.

[0403] Examples of cardiovascular diseases include, but are not limited to, myocardial infarction, acute myocardial infarction, post-myocardial infarction cardiac remodeling, post-myocardial infarction heart failure, stroke, coronary artery disease (i.e., atherosclerosis), acute coronary syndrome, myocarditis, and atrial fibrillation.

[0404] In some embodiments, the cardiovascular disease is selected from myocardial infarction, acute myocardial infarction, post-myocardial infarction cardiac remodeling, post-myocardial infarction heart failure, stroke, coronary artery disease, acute coronary syndrome, myocarditis, and atrial fibrillation.

[0405] Another object of the present invention is a method for treating a disease as described herein in a subject in need thereof, said method comprising administering to the subject at least one of:

[0406] an anti-CD8 α antibody, or an antigen binding fragment thereof, as described herein;

[0407] a fusion protein as described herein;

[0408] a nucleic acid as described herein encoding an anti-CD8 α antibody, an antigen-binding fragment thereof, or a fusion protein;

[0409] a vector as described herein comprising such a nucleic acid molecule; or

[0410] a host cell as described herein comprising such a vector.

[0411] As mentioned above, in some embodiments, the disease to be treated is a CD8-related disease. In some embodiments, the disease to be treated is a cardiovascular disease, in particular myocardial infarction or acute myocardial infarction. In some embodiments, the cardiovascular disease is selected from myocardial infarction, acute myocardial infarction, post-myocardial infarction cardiac remodeling, post-myocardial infarction heart failure, stroke, coronary artery disease, acute coronary syndrome, myocarditis, and atrial fibrillation.

[0412] In some embodiments, the method comprises administering a therapeutically effective dose of the anti-CD8 α antibody, or the antigen binding fragment thereof, as described herein; the fusion protein as described herein; the nucleic acid encoding as described herein an anti-CD8 α antibody, an antigen-binding fragment thereof, or a fusion protein; the vector as described herein comprising such a nucleic acid molecule; or the host cell as described herein comprising such a vector.

[0413] A therapeutically effective dose may for example correspond to a dose ranging from 0.0001 mg/kg to about 100 mg/kg of body weight per day, preferably from 0.001 mg/kg to about 50 mg/kg of body weight per day. It will be appreciated that these doses are exemplary and that an optimal dosage can be adapted taking into account the affinity and tolerability of the anti-CD8 α antibody, antigen-binding fragment, or fusion protein in the composition, pharmaceutical composition, or medicament.

[0414] The specific therapeutically effective dose level for any particular patient may depend upon a variety of factors

including the disease being treated and the severity of the disease; activity of the isolated anti-CD8 α antibody or antigen-binding fragment thereof, fusion protein, nucleic acid, expression vector, composition, pharmaceutical composition or medicament employed; the age, body weight, general health, sex and diet of the subject; the time of administration, route of administration, and rate of excretion of the specific anti-CD8 α antibody or antigen-binding fragment thereof, fusion protein, nucleic acid, expression vector, composition, pharmaceutical composition or medicament employed; the duration of the treatment; drugs used in combination or coincidental with the specific isolated anti-CD8 α antibody or antigen-binding fragment thereof, fusion protein, nucleic acid, expression vector, composition, pharmaceutical composition or medicament employed; and like factors well-known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than those required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved. The total dose required for each treatment may be administered by multiple doses or in a single dose.

[0415] Another object of the present invention is a pharmaceutical composition for treating or for use in the treatment of a disease as described herein in a subject in need thereof, wherein said pharmaceutical composition comprises at least one of:

[0416] an anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein;

[0417] a fusion protein as described herein;

[0418] a nucleic acid as described herein encoding an anti-CD8 α antibody, an antigen-binding fragment thereof, or a fusion protein;

[0419] a vector as described herein comprising such a nucleic acid molecule; or

[0420] a host cell as described herein comprising such a vector;

and a least one pharmaceutically acceptable excipient.

[0421] Another object of the present invention is the use of at least one of:

[0422] an anti-CD8 α antibody, or an antigen-binding fragment thereof, as described herein;

[0423] a fusion protein as described herein;

[0424] a nucleic acid as described herein encoding an anti-CD8 α antibody, an antigen-binding fragment thereof, or a fusion protein;

[0425] a vector as described herein comprising such a nucleic acid molecule; or

[0426] a host cell as described herein comprising such a vector;

in the manufacture of a medicament for the treatment of a disease as described herein in a subject in need thereof.

[0427] As mentioned above, in some embodiments, the disease to be treated is a CD8- related disease. In some embodiments, the disease to be treated is a cardiovascular disease, in particular myocardial infarction or acute myocardial infarction. In some embodiments, the cardiovascular disease is selected from myocardial infarction, acute myocardial infarction, post-myocardial infarction cardiac remodeling, post-myocardial infarction heart failure, stroke,

coronary artery disease, acute coronary syndrome, myocarditis, and atrial fibrillation.

[0428] The present invention further relates to the use of the isolated anti-CD8 α antibody, or antigen binding fragment thereof, or fusion protein as described herein for depleting CD8 α -expressing cells in a sample, and to an in vitro method for depleting CD8 α -expressing cells in a sample, said method comprising a step of contacting said sample with the isolated anti-CD8 α antibody, or antigen binding fragment thereof, or fusion protein as described herein.

[0429] As used herein, a “sample” refers to any biological material obtained via suitable methods known to the person skilled in the art from a subject. The sample may be collected in a clinically acceptable manner, e.g., in a way that cells, nucleic acids (such as DNA and RNA), proteins and/or extracellular vesicles are preserved. A “sample” may be a body tissue and/or a bodily fluid, preferably a bodily fluid. Examples of bodily fluids include, but are not limited to, blood, plasma, serum, lymph, ascetic fluid, cystic fluid, urine, bile, nipple exudate, vomitus, breast milk, tears, wound drainage, feces, vaginal secretions, synovial fluid, bronchoalveolar lavage fluid, sputum, amniotic fluid, peritoneal fluid, cerebrospinal fluid, pleural fluid, pericardial fluid, semen, saliva, sweat and alveolar macrophages, tissue lysates, biopsies and extracts prepared from diseased tissues.

[0430] In some embodiments, the sample is previously taken or retrieved from a subject, i.e., the methods as described herein do not comprise an active step of recovering a sample from the subject. Consequently, according to some embodiments, the methods as described herein are non-invasive methods, i.e., the methods of the invention are in vitro methods.

[0431] The invention further relates to the use of the isolated anti-CD8 α antibody, or antigen binding fragment thereof, or fusion protein as described herein for detecting or quantifying CD8 α in a sample, cell, tissue, organ, organism or subject; and to methods for detecting and/or quantifying CD8 α in a sample, cell, tissue, organ, organism, or subject comprising contacting said sample, cell, tissue, or organ or administering said organism or subject with the isolated anti-CD8 α antibody, or antigen binding fragment thereof, or fusion protein as described herein.

[0432] The uses and methods for detecting and/or quantifying CD8 α may be in vitro or in vivo.

[0433] In some embodiments, the isolated anti-CD8 α antibody, or antigen binding fragment thereof, as described herein is labeled for detection or diagnostic purposes.

[0434] Assays suitable for detecting or quantifying CD8 α using the isolated anti-CD8 α antibody, or antigen binding fragment thereof, or fusion protein as described herein are well-known in the art, and include ELISA, sandwich ELISA, RIA, FACS, tissue immunohistochemistry, Western-blot and immunoprecipitation.

TABLE OF SEQUENCES		
SEQ ID	Sequence	
NO	function	Sequence
	Consensus	NX_1X_2MN
	VH-CDR1	X_1 is N or Y X_2 is D or A or Y
1	Consensus	$X_3ISGSSX_4YIX_5YADFVKG$
	VH-CDR2	X_3 is D or S X_4 is S or R X_5 is D or G or Y
2	Consensus	$SSX_6X_7X_8X_9YX_{10}X_{11}X_{12}X_{13}MDV$
	VH-CDR3	X_6 is N or no amino acid X_7 is Y or no amino acid X_8 is Y or G or no amino acid X_9 is D or S or no amino acid X_{10} is S or N or F X_{11} is A or G X_{12} is S or D or N X_{13} is A or G
3	Consensus	$AGTSSDVGGX_{14}X_{15}X_{16}VS$
	VL-CDR1	X_{14} is G or N or Y X_{15} is S or Y X_{16} is S or Y
4	Consensus	$X_{17}DSX_{18}RPS$
	VL-CDR2	X_{17} is Q or S or Y X_{18} is Y or S
5	Consensus	$SSX_{19}TX_{20}YSTRV$
	VL-CDR3	X_{19} is Y or D X_{20} is Y or Q or S
6	hMP08-R3- F08-VH- CDR1	NNAMN
7	hMP08-R3- F08-VH- CDR2	DISGSSRYIGYADFVKG
8	hMP08-R3- F08-VH- CDR3	SSNYDYDNADAMDV
9	hMP08-R3- F08-VL- CDR1	AGTSSDVGGNSYVS
10	hMP08-R3- F08-VL- CDR2	SDSSSRPS
11	hMP08-R3- F08-VL- CDR3	SSYTQYSTRV
12	hMP08-R3- C11-VH- CDR1	NYDMN
13	hMP08-R3- C11-VH- CDR2	DISGSSSYIDYADFVKG
14	hMP08-R3- C11-VH- CDR3	SSYYSGSGMDV
15	hMP08-R3- C11-VL- CDR1	AGTSSDVGGGSSVS

- continued

TABLE OF SEQUENCES		
SEQ ID NO	Sequence function	Sequence
16	hMP08-R3- C11-VL- CDR2	QDSYRPS
17	hMP08-R3- C11-VL- CDR3	SSYTYSTRV
18	hMP09-R3- D03-VH- CDR1	NYYMN
19	hMP09-R3- D03-VH- CDR2	SISGSSRYIYYADFVKG
20	hMP09-R3- D03-VH- CDR3	SSGSYFGNAMDV
21	hMP09-R3- D03-VL- CDR1	AGTSSDVGGYYSVS
22	hMP09-R3- D03-VL- CDR2	YDSSRPS
23	hMP09-R3- D03-VL- CDR3	SSDTSYSTRV
24	hMP08-R3- F08-VH	EVOLVESGGS L V K P G G S L R L S C A A S G F T F S N N A M N W V R Q A P G K G L E W I S D I S G S S R Y I G Y A D F V K G R F T I S R D N A K N S L Y L Q M N S L R A E D T A V Y Y C V R S S N Y D Y N A D A M D V W G R G T L V T V S S
25	hMP08-R3- F08-VL	Q S V L T Q P A S V S G S P G Q S I T I S C A G T S S D V G G N S Y V S W Y Q Q H P G K A P K L M I Y D S S R P S G V S N R F S G S K S G N T A S L T I S G L Q A E D E A D Y Y C S S Y T Q Y S T R V F G G G T K L A V L
26	hMP08-R3- C11-VH	EVOLVESGGS L V K P G G S L R L S C A A S G F T F S N Y D M N W V R Q A P G K G L E W I S D I S G S S Y I D Y A D F V K G R F T I S R D N A K N S L Y L Q M N S L R A E D T A V Y Y C V R S S Y Y S G S G M D V W G R G T L V T V S S
27	hMP08-R3- C11-VL	Q S V L T Q P A S V S G S P G Q S I T I S C A G T S S D V G G G S V S W Y Q Q H P G K A P K L M I Y Q D S Y R P S G V S N R F S G S K S G N T A S L T I S G L Q A E D E A D Y Y C S S Y T Y S T R V F G G G T K L A V L
28	hMP09-R3- D03-VH	EVOLVESGGS L V K P G G S L R L S C A A S G F T F S N Y Y M N W V R Q A P G K G L E W I S S I S G S S R Y I Y Y A D F V K G R F T I S R D N A K N S L Y L Q M N S L R A E D T A V Y Y C V R S S G S Y F G N A M D V W G R G T L V T V S S
29	hMP09-R3- D03-VL	Q S V L T Q P A S V S G S P G Q S I T I S C A G T S S D V G G Y S V S W Y Q Q H P G K A P K L M I Y D S S R P S G V S N R F S G S K S G N T A S L T I S G L Q A E D E A D Y Y C S S D T S Y S T R V F G G G T K L A V L
30	Human CD8a (NCBI Reference Sequence: NP_001759.3)	M A L P V T A L L L P L A L L L H A A R P S Q F R V S P L D R T W N L G E T V E L K C Q V L L S N P T S G C S W L F Q P R G A A A S P T F L L Y L S Q N K P K A A E G L D T Q R F S G K R L G D T F V L T L S D F R E N E G Y Y F C S A L S N S I M Y F S H F V P V F L P A K P T T T P A P R P P T P A P T I A S Q P L S R P E A C R P A A G G A V H T R G L D F A C D I Y I W A P L A G T C G V L L L S L V I T L Y C N H R N R R R V C K C P R P V V K S G D K P S L S A R Y V
31	hMP08-R3- F08-VH nucleic acid	G C G G C C G C A A A C T A C A A G A C A G A C T T G C A A A A G A A G G C A T G C A C A G C T C A G C A C T G C T C T G T T G C C T G G T C C T C T G A C T G G G G T G A G G C C G A G G T G C A G C T G G T G G A G A G C G G G G G C A G C C T G G T G A A G C C A G G G G A A G C C T G A G G C T G A G C T G T G C C G C A T C C G G T T T A C A T T T T C C A C A C G C C A T G A A C T G G G T G A G C A G G C A C C G G C A A A G G C T G G A G T G G A T T T

- continued

TABLE OF SEQUENCES		
SEQ ID NO	Sequence function	Sequence
		CAGACATTTCCGGCAGTAGCAGATACATCGGCTACGCCGA CTTCGTGAAGGGCAGGTTTCAATCAGCAGAGACAACGC TAAGAACAGCCTGTACCTGCAGATGAACCTCCCTGAGGGCC GAGGACACAGCCGTGTACTACTGCGTGAGGAGCTCCAAC TACTATGACTACAACGCCGACGCTATGGATGTGTGGGAA GGGGCACCTGGTGACCGTGAGCTCCGCTAGCACCAAGG GCCCATCGGTCTCCCCCTGGCACCTCTCTCAAGAGCAC CTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGTCAAGGA CTACTTCCCCGAACCGGTGACGGTGTCTGTGGAACCTCAGGC GCCCTGACCAGCGCGGTGCACACTTCCCGGCTGTCCTAC AGTCCTCAGGACTCTACTCCCTCAGCAGCGTGGTGACCGT GCCCTCCAGCAGCTTGGGCACCCAGACCTACATCTGCAAC GTGAATCACAAGCCAGCAACACCAAGGTGGACAAGAAA GTTGAGCCCAATCTTGTGACAAAACCTCACACATGCCAC CGTGCCAGCAGCTGAACCTCTGGGGGGACCGTCAGTCTT CCTCTTCCCCCAAAACCAAGGACACCTCATGATCTCC CGGACCCCGAGGTCACATGCGTGGTGGTGAGCTGAGC CACGAAGACCTGAGGTCAAGTCAACTGGTACGTGGAC GCGGTGGAGGTGCATAATGCCAAGACAAGCCGCGGGAG GAGCAGTACAACAGCAGTACCGTGTGGTCAAGCTCTCA CCGTCTGCACAGGACTGGCTGAATGGCAAGGAGTACA AGTGCAAGGTCTCCAACAAGCCCTCCAGCCCCATCGA GAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACC ACAGGTGTACACCTGCCCCCATCCCGGAGGAGATGAC CAAGAACCAGGTGAGCTGACCTGCCTGGTCAAGGCTTC TATCCCGAGCATCGCCGTGGAGTGGGAGAGCAATGGG CAGCCGAGACAACCTACAAGACCACGCTCCCGTGCTG GACTCCGACGGCTCCTTCTCTCTACAGCAAGCTCACCG TGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCAT GCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGA GAAGAGCCTCTCCCTGTCTCGGGTAAATGATTCTAGA
32	hMP08-R3- F08-VL nucleic acid	GCGGCCGCAAACTACAAGACAGACTTGCAAAAGAAGGCA TGCACAGCTCAGCACTGCTCTGTTGCCTGGTCTCTGACT GGGGTGAGGGCCAGTCCGTGCTGACCCAGCCCGCTCCG TGTCCGAAGCCAGGACAGTCCATCACCATCTCTGCGC CGGCACCTCTCCGACGTGGGAGGAACAGCTACGTGTCC TGGTATCAACAGCACCCCGCAAGCACCCAAACTGATG ATCTACTCCGACAGTTCCTCGCCCTCCGCGCTGTCCAACA GGTTCAGCGGCTCCAAGAGCGGCAACCGCTCTCTGAC TATCTCCGGCTGCAGGCTGAAGACGAAGCCGACTACTAC TGCTCTCTTACACCCAGTCTTCCACCCGCTGTTCGGCG GCGGCACCAAACTGGCCGTGCTGGGTGAGCCCAAGGCCA ACCCCACTGTCACTCTGTTCCCGCCCTCTCTGAGGAGCTC CAAGCCAACAAGGCCACACTAGTGTGTCTGATCAGTGACT TCTACCCGGGAGCTGTGACAGTGGCTGGAAGGCAGATG GCAGCCCCGTCAAGGCGGGAGTGGAGACCACCAACCCT CCAACAGAGCAACAACAAGTACGCGGCGAGCAGCTACC TGAGCCTGACGCCGAGCAGTGAAGTCCACAGAAGCT ACAGCTGCCAGGTACGCATGAAGGAGCACCGTGGAGA AGACAGTGGCCCTACAGAATGTTCTATGATTCTAGA
33	hMP08-R3- C11-VH nucleic acid	GCGGCCGCAAACTACAAGACAGACTTGCAAAAGAAGGCA TGCACAGCTCAGCACTGCTCTGTTGCCTGGTCTCTGACT GGGGTGAGGGCCAGGTGCAGCTGGTGGAGAGCGGGGC AGCCTGGTGAAGCCAGGGGGAAGCTGAGGCTGAGCTGT GCCGCATCCGGTTTACATTTTCCAACCTACGACATGAACT GGGTGAGGCAGGCCCCCGCAAGGACTGGAGTGGATT CAGACATCTCCGGCAGCAGTTCCTACATCGACTACGCTGA TTTTGTGAAGGGCAGGTTTCAATCTCCCGGACAAACGCT AAGAACTCCCTGTACCTGCAGATGAACAGTCTGAGGGCC GAGGACACCGCTGTGTACTACTGCGTGAGGTCTCTACT ATAGTGGCAGCGCATGGACGTGTGGGCGGGGAACCC TGGTGACCGTGTCTCTGCTAGCACCAAGGGCCATCGGT CTTCCTTGGCACCTCTCTCAAGAGCACTCTGGGGGC ACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCG AACCGGTGACCGTGTCTGGAACCTCAGGCGCCCTGACCA GCGGCGTGACACCTTCCCGGCTGTCTACAGTCTCTCAGG ACTCTACTCCCTCAGCAGCGTGGTACCGTGCCCTCCAGC AGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCACA

- continued

TABLE OF SEQUENCES		
SEQ ID	Sequence	
NO	function	Sequence
		AGCCAGCAACACCAAGGTGGACAAGAAAGTTGAGCCCA AATCTTGTGACAAACTCACACATGCCACCGTGCCAGC ACCTGAACTCCTGGGGGACCGTCAGTCTTCTCTTCCCC CCAAAACCAAGGACACCTCATGATCTCCCGGACCCCG AGGTCACATGCGTGGTGGTGGACGTGAGCCACGAAGACC CTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGG TGCATAATGCCAAGACAAGCCGCGGGAGGAGCAGTACA ACAGCACGTACCGTGTGGTCAGCGTCTCACCCTCCTGCA CCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGT CTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCATC TCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGTAC ACCTTGCCCCCATCCCGGGAGGAGATGACCAAGAACAG GTCAGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCG ACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGA ACAACACTACAAGACCAGCCTCCCGTGTGGACTCCGACGG CTCCTTCTTCTCTACAGCAAGCTCACCCTGGACAAGAGC AGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGATGC ATGAGGCTCTGCACAACCACTACACGCAGAAGAGCCTCTC CCTGTCTCCGGTAAATGATTCTAGA
34	hMP08-R3- C11-VL nucleic acid	GCGGCCGCAAACTACAAGACAGACTTGCAAAGAAGGCA TGCACAGCTCAGCACTGCTCTGTTGCCCTGGTCTCTGACT GGGGTGAGGGCCAGTCCGTGCTGACCCAGCCCGCTCCG TGTCCGGAAGCCAGGACAGTCCATCACCATCTCCTGCGC CGGCACCTCCTCCGACGTGGGAGGAGGAAGCTCCGTGAG CTGGTATCAACAGCACCCCGGCAAAGCACCCAACTGAT GATCTACCAGGACAGCTACAGGCCCTCCGGCGTGTCCAAC AGGTTACGCGGCTCCAAGAGCGGAACACAGCCTCTCTG ACCATCTCCGGCTGCAGGCTGAGGACGAAGCCGACTACT ACTGCTCATCTACACTTACTATTCCACCAGGGTGTTCGG AGGCGGAACAAACTGGCCGTGCTGGGTGAGCCCAAGGC CAACCCCACTGTCACTCTGTTCCTCCCGCTCCTCTGAGGAG CTCCAAGCCAACAAGGCCACACTAGTGTGTCTGATCAGTG ACTTCTACCCGGGAGCTGTGACAGTGGCCTGGAAGGCAG ATGGCAGCCCCGTCAAGGCGGGAGTGGAGACCACCAAC CCTCCAACAGAGCAACAACAGTACGCGGCCAGCAGCT ACCTGAGCCTGACGCCCAGCAGTGGAAAGTCCCACAGAA GCTACAGCTGCCAGGTACGCTATGAAGGGAGCACCGTGG AGAAGACAGTGGCCCCACAGAATGTTATGATTCTAGA
35	hMP09-R3- D03-VH nucleic acid	GCGGCCGCAAACTACAAGACAGACTTGCAAAGAAGGCA TGCACAGCTCAGCACTGCTCTGTTGCCCTGGTCTCTGACT GGGGTGAGGGCCGAGGTGCAGCTGGTGGAGAGCGGGGC AGCCTGGTGAAGCCAGGGGGAAGCCTGAGGCTGAGCTGT GCCGCATCCGGGTTTACATTTTCCAACACTATATGAACT GGGTGAGGCAGCCCCCTGGCAAGGGCCTGGAGTGGAATTT CATCTATCTCCGGCAGTAGTCGATACATCTACTACGCTGA CTTTGTGAAGGGCCGTTTACCATCTCCAGGGACAACGCT AAGAAGTCCCTGTATCTGCAGATGAACCTCCCTGCGCGCCG AGGACACCGCGTGTACTACTGCGTGCCTCCTCCGGCAG CTACTTCGGAAATGCTATGGATGTGTGGGGGAGGGGGAC CCTGGTGACCGTGTCTCCGCTAGCACAAGGGCCCATCG GTCTTCCCCCTGGCACCTCCTCCAAGAGCACCTCTGGGG GCACAGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCC CGAACCGGTGACGGTGTCTGGAACTCAGGCGCCCTGAC CAGCGGCGTGACACCTTCCCGGCTGTCTACAGTCTCTCA GGACTCTACTCCCTCAGCAGCGTGGTACCGTGCCCTCCA GCAGCTTGGGCACCCAGACCTACATCTGCAACGTGAATCA CAAGCCCAGCAACCAAGGTGGACAAGAAAGTTGAGCC

- continued

TABLE OF SEQUENCES		
SEQ ID NO	Sequence function	Sequence
		CAAATCTTGTGACAAACTCACACATGCCACCGTGCCCA GCACCTGAACTCCTGGGGGACCGTCAGTCTTCTCTTCC CCCCCAAACCAAGGACACCCCTCATGATCTCCCGACCCC CGAGGTCACATGCGTGGTGGTGGACGTAGCCACGAAGA CCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGA GGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGCAGTA CAACAGCACGTACCGTGTGGTACGCGTCTCACCGTCTG CACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGAAG GTCTCCAACAAAGCCCTCCAGCCCCATCGAGAAAACCA TCTCCAAGCCAAAGGGCAGCCCCGAGAACCACAGGTGT ACACCCCTGCCCCATCCCGGGAGGAGATGACCAAGAACC AGGTACAGCTGACCTGCCTGGTCAAGGCTTCTATCCAG CGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGA GAACAACACTACAAGACCACGCTCCCGTGTGGACTCCGAC GGCTCCTTCTCTCTACAGCAAGCTCACCGTGGACAAGA GCAGGTGGCAGCAGGGGAACGTCTTCTCATGCTCCGTGAT GCATGAGGCTCTGCACAACCACTACACGCAGAAGGCCT CTCCCTGTCTCCGGTAATGATTCTAGA
36	hMP09-R3- D03-VL nucleic acid	GCGGCCGCAAACTACAAGACAGACTTGCAAAAGAAGGCA TGCACAGCTCAGCACTGCTCTGTTGCCTGGTCTCTGACT GGGGTGAGGGCCCACTCCGTGCTGACCCAGCCCGCCTCG TGTCCGGAAGCCAGGACAGTCCATCACCATCTCTGCGC CGGCACCTCCTCCGACGTGGGAGGATACTCTCCGTGAGC TGGTATCAACAGCACCCCGGCAAGCCCCAACTGATG ATCTACTACGACAGTCCCGGCCCTCCGGCGTGTCTAACA GGTTCCTCCGGTCCAAGAGCGGCAACCCGCGAGCTGAC CATCTCCGGCTGACAGCAGAAGACGAGGCCGACTACTA CTGCTCCTCCGACACCTCTACAGCACCAGGGTGTTCGGC GGTGGCACCAGCTGGCCGTGCTGGGTGAGCCCAAGGCC AACCCCACTGTCACTCTGTTCCCGCCCTCTCTGAGGAGC TCCAAGCCAACAAGGCCACACTAGTGTGTCTGATCAGTGA CTCTTACCCGGGAGCTGTGACAGTGGCTGGAAGGCAGAT GGCAGCCCCGTCAAGGCGGAGTGGAGACCACCAACCC TCCAACAGAGCAACAACAAGTACGCGGCCAGCAGCTAC CTGAGCCTGACGCCCGAGCAGTGGAAAGTCCCACAGAAGC TACAGCTGCCAGGTACGCATGAAGGAGCACCGTGGAG AAGACAGTGGCCCTACAGAATGTTCTGATTCTAGA

EXAMPLES

[0435] The present invention is further illustrated by the following examples.

TABLE 1

Sequences of the antibodies of the examples.			
	MP08-R3-F08	MP08-R3-C11	MP09-R3-D03
V _H -CDR1	SEQ ID NO: 6	SEQ ID NO: 12	SEQ ID NO: 18
V _H -CDR2	SEQ ID NO: 7	SEQ ID NO: 13	SEQ ID NO: 19
V _H -CDR3	SEQ ID NO: 8	SEQ ID NO: 14	SEQ ID NO: 20
V _L -CDR1	SEQ ID NO: 9	SEQ ID NO: 15	SEQ ID NO: 21
V _L -CDR2	SEQ ID NO: 10	SEQ ID NO: 16	SEQ ID NO: 22
V _L -CDR3	SEQ ID NO: 11	SEQ ID NO: 17	SEQ ID NO: 23
VH sequence	SEQ ID NO: 24	SEQ ID NO: 26	SEQ ID NO: 28
VL sequence	SEQ ID NO: 25	SEQ ID NO: 27	SEQ ID NO: 29

Example 1: Antibody production

Materials and Methods

[0436] Three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) were produced in both human IgG1 lambda and rat IgG2b kappa formats and purified. For

IgG expression, the V_H and VL chain fragments were cloned into an IgG expression system encoding heavy and light chains on separate pcDNA3.4 plasmids. Constructs were sequenced to validate correct insertion of V_H and VL sequences.

[0437] Plasmids were transiently transfected into ExpiCHO cells and IgG containing supernatants were filtered (0.22 μm) and purified on protein A columns. Antibodies that did not reach 98% purity went through a second purification step by preparative SEC (size exclusion chromatography). A limulus amebocyte lysate test was conducted to measure endotoxin levels in purified IgGs.

Results

[0438] The summary of the antibody production, purification and quality control is presented in Table 2.

[0439] All three antibody clones displayed high expression yields both in the human IgG1 format, with high amounts recovered after protein A purification (up to 400 mg/L culture). In the rat IgG2b format, while clone MP08-R3-F08 expressed to a similar high yield, clones MP08-R3-C11 and MP09-R3-D03 expressed to a moderate yield, with 6.3 and 7.5 mg of purified IgG recovered respectively from 100 mL cultures.

[0440] The six antibodies were analyzed by SDS-PAGE: 1.3 g of each antibody clone were loaded per well in reducing and non-reducing conditions. All six antibodies showed a standard migration profile with one band at 50 kDa and another band at 25 kDa, corresponding to the heavy and light chains respectively (data not shown).

was observed on mouse CD8 α (FIG. 1B for human IgG1 clones and FIG. 2B for rat IgG2b clones).

[0446] Binding specificity tested on human CD4 showed that none of the three antibody clones bound to this protein, neither in human IgG1 nor in rat IgG2b format (FIG. 1C for human IgG1 clones and FIG. 2C for rat IgG2b clones).

TABLE 2

Summary of production purification and quality control of the 3 antibody clones in both human IgG1 and rat IgG2b formats.											
Antibody name		Purity							Yield (mg) after endotoxin removal		
		Yield (mg)		Analytical SEC after protein A purification		Analytical SEC after preparative SEC					
		after protein A purification	SDS-PAGE	SEC-214 nm	SEC-280 nm	SEC-214 nm	SEC-280 nm	Yield (mg) after SEC	Delivered quantity (mg)	Endotoxin level (EU/mg)	
Human	MP08-R3-F08	34.7	>95%	73.7%	74.7%	100%	100%	16.8	11.06	11.1	<1
IgG1	MP08-R3-C11	46.4	>95%	91.4%	91.5%	99.3%	99.4%	27.3	24.96	25	<1
lambda	MP09-R3-D03	22.4	>95%	98.6%	98.7%	/	/	/	/	22.4	<1
Rat	MP08-R3-F08	16.4	>95%	53.1%	53.4%	99.4%	99%	4.1	/	4.1	<1
IgG2b	MP08-R3-C11	6.3	>95%	77.3%	77.6%	99.4%	99%	0.5	/	0.5	<1
kappa	MP09-R3-D03	7.5	>95%	82.4%	83%	99.5%	99.5%	0.2	/	0.2	>1, <5

Example 2: Antibody Binding to Recombinant CD8 α Protein

Materials and Methods

[0441] Five g/mL of streptavidin were coated on Nunc Maxisorp 96-well plates (ThermoFisher #442404) at 100 μ L/well and incubated 1 hour at room temperature. Biotinylated human CD8 α (Fisher Scientific #17196181) and mouse CD8 α (Sino biological #50389-M08H-B) were added at 0.5 μ g/mL in 100 μ L/well and incubated overnight at 4° C. Human CD4 (Sino biological #10400-H08H) was coated passively at 1 μ g/mL in 100 μ L/well and incubated overnight at 4° C. Plates were then blocked using 200 μ L of PBS containing 4% skimmed milk for 1 hour at room temperature.

[0442] The three antibody clones (in both human IgG1 lambda and rat IgG2b kappa formats) were diluted in milk-PBS and tested at different concentrations (75, 25, 8.3, 2.8, 0.93, 0.31, 0.103, 0.034, 0.011, 0.0038, 0.0013 and 0.00042 μ g/mL). 100 μ L/well were incubated 1 hour at room temperature.

[0443] Human IgGs were detected using an anti-hFab-HRP (Sigma #A0293, 1:4000 dilution), while rat IgGs were detected using an anti-rat-HRP (Sigma #A9037, 1:10 000 dilution), at 100 μ L/well and incubated 1 hour at room temperature.

[0444] TMB ((3,3',5,5' tetramethylbenzidine) Thermo Scientific #12617087) was added for a 3-minute incubation. Reaction was stopped with H₂SO₄ and absorbance at 450 nm was read on a microplate reader (Pherastar FS, BMG Labtech).

Results

[0445] All three tested antibody clones were capable of binding to human CD8 α , with EC50 ranging from 0.28 to 2.0 nM for clones in the human IgG1 format (FIG. 1A and Table 3) and ranging from 0.20 to 0.99 nM for clones in the rat IgG2b format (FIG. 2A and Table 3). No cross-reactivity

[0447] These results demonstrate that the three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) possess strong binding towards human CD8 α , as well as a good specificity towards human CD8 α . Additionally, none of the three antibody clones exhibit binding towards human CD4 or cross-reactivity towards mouse CD8 α .

TABLE 3

Summary of the EC50, CI 95% and Rsquared of the three antibody clones in the human IgG1 and rat IgG2b formats.					
		MP08-R3-F08	MP08-R3-C11	MP09-R3-D03	
Human IgG1	EC50 (nM)	0.28	2.4	2.0	
	CI 95% (nM)	0.26 to 0.30	1.8 to 3.1	1.5 to 2.8	
	Rsquared	0.28	2.4	2.0	
Rat IgG2	EC50 (nM)	0.20	0.48	0.99	
	CI 95% (nM)	0.18 to 0.22	0.36 to 0.66	0.69 to 1.4	
	Rsquared	0.999	0.993	0.990	

Example 3: Antibody Binding to CD8+ and Non-Binding to CD8- Cells

Materials and Methods

[0448] 200,000 HPB-ALL cells (expressing CD8 α and referred to as CD8+) or 1299 cells (not expressing CD8 α and referred to as CD8-) were incubated with primary antibodies: mouse anti-CD8 α (Sino biological #10980-MM28, 1:50 dilution), or with one of the three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11, and MP09-R3-D03) in the human IgG1 or rat IgG2b format, at a concentration of 10, 3.3, 1.1 or 0.37 μ g/mL. HPB-ALL is a cell line of human lymphocytes derived from T cell leukemia. 1299 (or H1299) is a cell line of human epithelial-like cells derived from non-small cell lung carcinoma.

[0449] Human antibodies were detected using an anti-human F(ab')₂-AF647 antibody (Jackson #109-605-006,

1:100 dilution), rat antibodies with an anti-rat-FITC antibody (Sigma #F1763, 1:320 dilution) and the anti-human CD8 α mouse antibody with an anti-mouse-FITC antibody (Sigma #12-506, 1:100 dilution). Results were acquired on a CytoFLEX cytometer (Beckman Coulter) and analysed using FlowJo.

Results

[0450] Cells were validated by flow cytometry using control experiments (data not shown). Background signal from secondary antibodies was determined on both cell lines (data not shown). An irrelevant human IgG1 termed “13R4” showed no binding on either of the cell lines (FIG. 3D for HPB-ALL cells and FIG. 4D for H1299 cells). CD8 α expression was also assessed using a commercial anti-CD8 α antibody, showing a strong signal on HPB-ALL cells (CD8+ cells) and no signal on H1299 cells (CD8- cells), as expected (FIG. 3E for HPB-ALL cells and FIG. 4E for H1299 cells).

[0451] Antibody binding to both cell lines was assessed by flow cytometry. All three antibody clones in the human IgG1 format bound to HPB-ALL CD8+ cells (FIGS. 3A-C showing the 3.3 μ g/mL concentration), while no binding was observed on H1299 CD8- cells (FIGS. 4A-C showing the 3.3 μ g/mL concentration).

[0452] At the lowest concentration (0.37 μ g/mL), clones MP08-R3-C11 and MP08-R3-F08 showed the strongest signals (data not shown), suggesting higher affinities.

[0453] Similar results were obtained with the rat IgG2b format, with clones MP08-R3-C11 and MP08-R3-F08 giving the strongest signals at 0.37 μ g/mL (data not shown).

[0454] These results confirm the results obtained on recombinant human CD8 α protein and demonstrate that the three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) possess a strong binding and good specificity towards human CD8 α .

Example 4: Antibody Binding to Primary CD8+ T Cells

Materials and Methods

[0455] Human PBMCs (peripheral blood mononuclear cells) were prepared from a buffy coat using density gradient. Buffy coat was diluted 1/2 in PBS-SVF 2% and 30 mL were loaded onto 15 mL of Lymphoprep (Fisher Scientific #17171036). Tubes were centrifuged 20 min at 2000 rpm with no brake. PBMC were taken at the lymphoprep/plasma interface, washed with PBS-SVF and centrifuged 5 min at 1200 rpm. CD8+ T cells isolation was performed using CD8 α + T Cell Isolation Kit (Miltenyi Biotec #130-104-075) according to manufacturer's instructions.

[0456] Cells were incubated with primary antibodies: mouse anti-CD8 α (Sino biological #10980-MM28, 1:50 dilution), or with one of the three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11, and MP09-R3-D03) in the human IgG1 or rat IgG2b format, at a concentration of 10, 3.3, 1.1 or 0.37 μ g/mL. Human antibodies were detected using an anti-human F(ab')₂-AF647 antibody (Jackson #109-605-006, 1:100 dilution), rat antibodies with anti-rat-FITC antibody (Sigma #F1763, 1:320 dilution) and the anti-human CD8 α mouse antibody with an anti-mouse-FITC antibody (Sigma #12-506, 1:100 dilution). Results

were acquired on a CytoFLEX cytometer (Beckman Coulter) and analyzed using FlowJo.

Results

[0457] Cells were validated by flow cytometry using control experiments (data not shown). Background signal from secondary antibodies was determined (data not shown). An irrelevant human IgG1 termed “13R4” showed no binding to fresh CD8+ T cells (FIG. 5D). CD8 α expression was also assessed using a commercial anti-CD8 α antibody, showing a strong signal on purified primary CD8+ T cells (FIG. 5E).

[0458] Antibody binding to fresh primary CD8+ T cells was assessed by flow cytometry, showing binding of all three antibody clones in the human IgG1 format (FIGS. 5A-C).

[0459] As with the HPB-ALL CD8+ cell line, at the lowest concentration (0.37 μ g/mL), clones MP08-R3-C11 and MP08-R3-F08 showed again the strongest signals (data not shown), suggesting a higher affinity for these two clones.

[0460] Similar results were obtained in rat IgG2b format with clones MP08-R3-C11 and MP08-R3-F08 giving the strongest signals at 0.37 μ g/mL (data not shown).

[0461] These results confirm the results obtained on recombinant human CD8 α protein and on CD8+/CD8- cell lines and demonstrate that the three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) exhibits good binding for human CD8 α .

Example 5: Human CD8 Depletion In Vivo

Materials and Methods

Mice

[0462] B6; SJL-Tg(CD8 α CD8b)57Scr/J mice were used (Jackson Laboratory strain #003202, LaFace et al. Human CD8 transgene regulation of HLA recognition by murine T cells. *J Exp Med.* 1995 Nov. 1; 182(5):1315-25); and later designed as hCD8 mice. These hCD8 mice express a transgene containing the human CD8 (hCD8) alpha and beta chains expressed under the control of the murine p56^{lck} proximal promoter.

[0463] Preliminary experiments confirmed the absence of expression of human CD8 α (i.e., hCD8 α) in control C57B16 mice (named control WT mice, FIG. 6A) and strong human CD8 α expression in transgenic mice expressing hCD8 α transgene (FIG. 6B). However, this hCD8 α expression is not restricted to CD8 T cells.

[0464] Mice were maintained under identical standard conditions including housing, regular care, and normal chow diet in the same animal facility throughout the duration of the experiments (12 hours light/dark cycle with ad libitum access to food and water). Mice were aged 9 to 12 weeks at the beginning of experiments. Only males were used except for the hCD8 100 μ g depletion group where half were females because of mice availability.

Antibody Injection

[0465] Upon isoflurane anesthesia, antibodies in the human IgG1 format (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) were injected intravenously through the retro-orbital sinus with a 30-gauge insulin syringe. Mice were returned to their home cage after complete recovery from anesthesia.

Blood Sampling

[0466] At the required timepoints, blood was sampled from the submandibular vein. Upon isoflurane anesthesia, the left submandibular vein was punctured using a 25-gauge needle. A few drops of blood corresponding approximately to 100 μ L were collected in a collection tube containing 5 μ L of heparin. After sample collection, bleeding was stopped by application of a sterile gauge. After complete recovery from the anesthesia, mice were returned to their home cage.

Flow Cytometry

[0467] For each sample, 50 μ L of blood were incubated with 10 μ L of FACS antibody solution for 30 minutes at room temperature.

[0468] In blood samples, erythrocytes were lysed using BD FACS lysis solution (BD Biosciences) after staining. Acquisition was performed on a LSR Fortessa cytometer, and further analysis performed using FlowJO v10.8. Forward scatter (FSC) and side scatter (SSC) parameters were used to gate live cells excluding red blood cells, debris, and cell aggregates. Cells were later selected based upon CD45 expression. For hCD8 transgenic mice, the expression level of hCD8 α was assessed in total CD45+ cells; with control mice experiments, the expression of mCD8 α was assessed from the CD45+CD11b-CD3+ cells.

Data Representation and Normalization

[0469] The number of hCD8 α + cells was evaluated within the total CD45+ cell population and then normalized to the average ratio of hCD8 α + in control non-depleted mice in order to assess the repopulation of the hCD8 α + cells.

Results

[0470] In order to evaluate the efficiency and duration of depletion of the hCD8 α -expressing cells (hCD8+ cells), the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) were injected into mice at 3 different doses (50, 100 and 200 μ g), with 3 mice per group. Immediate confirmation of depletion was performed by flow cytometry analysis 1 hour after antibody injection. Follow-up of hCD8 α cell repopulation was done weekly for the 2 low doses; and at day 7, 14 and 42 for the high dose. In mice from the MP08-R3-C11 50 μ g group one animal was excluded because of incorrect antibody injection.

[0471] Depletion of hCD8 α -expressing cells was complete 1 hour after injection for all clones at 200 μ g, while it was almost complete for the 50 μ g and 100 μ g doses (FIGS. 7A-C and 8A-C). At day 7 after injection, hCD8 α depletion was total for all clones and doses (FIGS. 7A-C and 8A-C).

[0472] Later timepoints showed progressive hCD8 α + repopulation starting with the 50 μ g dose, with the threshold of 50% repopulation attained by all clones at 50 μ g by day 28 with very similar evolution (FIG. 8C). For higher dose, only clones MP08-R3-C11 (FIG. 7B) and MP09-R3-D03 (FIG. 7C) reached 50% repopulation at 100 μ g and only MP09-R3-D03 reached 50% repopulation at 200 μ g by day 42. The repopulation within 100 μ g and 200 μ g groups follow very similar trends for clones MP08-R3-F08 (FIG. 7A) and MP09-R3-D03 (FIG. 7C).

[0473] In order to investigate the cross-reactivity of the three human CD8 α antibody clones with mouse CD8 α , each

of the three antibody clones were injected in control C57B16 mice and the evolution of the blood CD8 T cell population was evaluated.

[0474] As shown on the flow cytometry plots (FIG. 9A-D) and on the quantification (FIG. 9E), none of the three anti-human CD8 α antibody clone was able to deplete mouse CD8 T cells in control mice, neither 1 hour after injection nor 7 days after. These results demonstrate that all three antibody clones do not cross-react with mouse CD8 α .

CONCLUSION

[0475] These results show that all three antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) are capable of strongly binding to human CD8 α recombinant protein, but also human CD8 α expressed at the surface of CD8 α -expressing cell lines and at the surface of primary CD8+ T cells. Additionally, all three antibody clones possess good specificity for human CD8 α , with no binding to human CD4 and no cross-reactivity for mouse CD8 α .

[0476] Furthermore, all three antibody clones are capable of depleting CD8 α -expressing cells in vivo. Also, all three clones possess good specificity for human CD8 α , with no cross-reactivity for mouse CD8 α in vivo.

[0477] All clones induce a complete or almost complete depletion of CD8 α -expressing cells within 1 hour after injection in human CD8 α -expressing mice, depending on the dose. In addition, depletion is complete for all clones and doses for at least 7 days. After day 7 post-depletion, repopulation of hCD8 α -expressing cells begins progressively. At the 50 μ g dose, repopulation reaches 50% by day 28 with all clones, while the 50% repopulation is reached only by day 42 with clones MP08-R3-C11 and MP09-R3-D03 at dose 100 μ g, and only with clone MP09-R3-D03 at dose 200 μ g.

[0478] Thus, depletion of CD8 α -expressing cells by the anti-CD8 α antibodies of the invention is fast, almost complete if not complete, and transient. Indeed, CD8 α -expressing cell repopulation after depletion is controlled and depends upon the dose of antibody used.

[0479] Taken together, these results indicate that the anti-CD8 α antibodies as described herein could be valuable therapeutic tools for transiently depleting CD8 α -expressing cells in a controlled manner.

Example 6: Antibody Specificity

Materials and Methods

[0480] The Retrogenix Cell Microarray Technology platform was used to screen for specific off-target binding interactions of the anti-CD8 α antibody clone MP08-R3-F08.

Pre-Screen

[0481] The level of background binding of the anti-CD8 α antibody clone MP08-R3-F08 to fixed untransfected HEK293 cells, and to cells overexpressing CD8 α (isoform 1, isoform 2 secreted or tethered secreted) was evaluated. Binding to untransfected cells and target-expressing cells was assessed using an AlexaFluor 647 labelled anti-human IgG Fc detection antibody, followed by fluorescence imaging.

Library Screen

[0482] The level of binding of the anti-CD8 α antibody clone MP08-R3-F08 was screened using fixed human HEK293 cells, individually over-expressing 6105 full-length human plasma membrane proteins, human secreted proteins, and cell surface-tethered human secreted proteins, as well as additional 400 human heterodimers.

[0483] Binding detection was performed using the same AlexaFluor 647 labelled anti-human IgG Fc detection antibody as above, followed by fluorescence imaging. The anti-CD8 α antibody clone MP08-R3-F08 was tested against 2 replicates. Fluorescent images were analyzed and quantitated using Image Quant software.

[0484] A protein interaction was defined as a duplicate spot showing a raised signal compared to background levels. Interactions were classified as strong, medium, weak or very weak depending on the intensity of the duplicate spots.

Confirmation Screen

[0485] All proteins identified as interacting with the anti-CD8 α antibody clone MP08-R3-F08 at the Library screen step were re-expressed in HEK293 cells. The level of binding of the anti-CD8 α antibody clone MP08-R3-F08 was assessed on both fixed and non-fixed cells in the same manner than for the library screen using fluorescence imaging, and compared to Rituximab (anti-CD20 antibody) used as control antibody. The goal was to determine which interactions were repeatable and specific to the anti-CD8 α antibody clone MP08-R3-F08.

Results

Pre-Screen

[0486] The anti-CD8 α antibody clone MP08-R3-F08 showed low levels of background binding to fixed untransfected HEK293 cells at all three tested concentrations (i.e., 2, 5 and 20 μ g/mL). The anti-CD8 α antibody clone MP08-R3-F08 exhibited binding towards over-expressed CD8 α (isoform 2 both secreted or tethered secreted) at all 3 concentrations, while binding towards over-expressed CD8 α (isoform 1) was observed only at 5 and 20 μ g/mL. Based on these results, further screenings were performed at 20 μ g/mL on fixed cells.

Library Screen

[0487] After screening 6105 full-length human plasma membrane proteins, human secreted proteins, and cell surface-tethered human secreted proteins, as well as additional 400 human heterodimers over-expressed in HEK293 cells using 20 μ g/mL anti-CD8 α antibody clone MP08-R3-F08, 23 interactions were identified and further analyzed.

Confirmation Screen

[0488] For confirmation screen, the 23 proteins identified as interacting with anti-CD8 α antibody clone MP08-R3-F08 at the step of library screen, as well as CD8 α (isoform 1) and two control receptors (CD20 and EGFR) were over-expressed in HEK293 cells. Cells were used fixed or non-fixed and interaction with the anti-CD8 α antibody clone MP08-R3-F08 was evaluated and compared with the control antibody (i.e., Rituximab).

[0489] On fixed cells microarrays, 22 out of the 23 proteins identified as interacting with the anti-CD8 α antibody clone MP08-R3-F08 were reproducibly observed as interacting with anti-CD8 α antibody clone MP08-R3-F08 (FIG. 10A).

[0490] The exception was FCGR3B, which showed a very weak intensity in the library screen, and lack of reproducibility into the confirmation screen indicates that the interaction was not real. Seventeen out of the 23 identified proteins were bound by both the anti-CD8 α antibody clone MP08-R3-F08 and control antibody, and were therefore classified as non-specific interactions (FIGS. 10A-B). Those included Fc gamma receptors, which were presumably Fc-domain mediated, and IGHG3, which is bound directly by the detection antibody.

[0491] Additional three out of the 23 proteins, as well as CD8 α (isoform 1) appeared to be specific to the anti-CD8 α antibody clone MP08-R3-F08. However, signal intensities were so weak and close to background levels that they were categorized as non-significant.

[0492] After excluding the non-reproducible, non-specific, and non-significant interactions, only two remained that were significant and specific to the anti-CD8 α antibody clone MP08-R3-F08: CD8 α isoform 2 both secreted or tethered secreted forms. On non-fixed cells, the interactions with CD8 α isoform 2 both secreted or tethered secreted forms of the anti-CD8 α antibody clone MP08-R3-F08 was confirmed (FIG. 10C).

[0493] As a conclusion, anti-CD8 α antibody clone MP08-R3-F08 showed a significant specific interaction with CD8 α (isoform 2 both secreted or tethered secreted) on both fixed and live cell microarrays. No other specific interactions were identified, indicating high specificity of the anti-CD8 α antibody clone MP08-R3-F08 for its target CD8 α isoform 2.

Example 7: CD8 Depletion in Non-Human Primates

Materials and Methods

Animals

[0494] Four naïve healthy male Cynomolgus Monkeys aged 2 to 4 years old were used for this study. Animals were housed in stainless steel cages, one monkey per cage. The room temperature was maintained between 2° and 29° C. with a relative humidity of 40-70%. Fluorescent illumination was on 12-hour light (08:00-20:00) and 12-hour dark cycle. Monkeys had illimited access to water and food.

Antibody Injection

[0495] Each monkey was intravenously infused through the saphenous vein with either (1) 20 mg/kg bodyweight of anti-CD8 α antibody clone MP08-R3-F08, (2) 20 mg/kg bodyweight of anti-CD8 α antibody clone MP08-R3-C11, (3) 20 mg/kg bodyweight of anti-CD8 α antibody clone MP09-R3-D3, or (4) 7.5 mg/kg bodyweight of anti-CD8 α antibody clone MP08-R3-F08. Antibody infusion was completed within 30 minutes.

Sample Collection

[0496] Whole blood was collected into potassium (K2) EDTA tubes from each animal at day -1, day 0 (1h), day 0 (6h), day 0 (24h), day 5, day 10, day 15 and day 22

post-injection for the three monkeys which received the 20 mg/kg dose of antibody and at day -1, day 0 (3*h*), day 0 (6*h*), day 0 (24*h*), day 7, day 14, day 28, day 48 and day 62 post-injection for the monkey which received the 7.5 mg/kg dose of antibody.

Flow Cytometry

[0497] From whole blood samples, erythrocytes were lysed using BD FACS lysis solution (BD Biosciences cat #555898). After lysis, pelleted cells were washed twice with PBS 1% FBS. Cells were resuspended in PBS 1% FBS and incubated with FcR blocking buffer for 10 minutes, before staining for 1 h at 4° C. with antibodies as detailed in Table 4 below. After staining, cells were washed twice with PBS 1% FBS, fixed with fixation buffer for 10 minutes, washed again and resuspended in PBS before acquisition on a flow cytometer. Forward scatter (FSC) and side scatter (SSC) parameters were used to gate live cells excluding red blood cells, debris, and cell aggregates. Cells were later selected based upon CD45 expression. CD8+ cells were identified as CD45+CD11b-CD3+CD4-CD8+ cells.

TABLE 4

Antibodies used for flow cytometry in non-human primate experiments.		
Antibody/Dye	Cat. #	Vendor
CD45-BV510	563530	BD
CD11b-APC	17-0118-42	ebioscience
CD20-FITC	555622	BD
CD3-BV605	562994	BD
CD4-BB700	566479	BD
CD8- BV786	563823	BD
CD16-R718	566969	BD
CD14-BV650	563419	BD
CD56-PE	556647	BD
CD66b-BV421	562940	BD
Live/Dead- APC-CY7	565388	BD

Data Representation

[0498] The number of CD8α+ cells was evaluated within the total CD45+ cell population and represented as a percentage of the CD45+ cells.

Results

[0499] In order to evaluate the efficiency of depletion of the CD8α-expressing cells, the 3 antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) were injected into monkeys at a dose of 20 mg/kg bodyweight, with one monkey per group. Immediate confirmation of depletion was performed by flow cytometry analysis 1 hour and 6 hours after antibody injection. Follow-up of CD8α-expressing cell depletion was done at 24 hours, day 5, day 10, day 15 and day 22 after antibody injection.

[0500] Depletion of CD8α-expressing cells was complete 1 hour after injection for clone MP08-R3-F08, and the percentage of CD8+ cells remained at 0% until day 22 (FIG. 10A). Antibody clones MP08-R3-C11 and MP09-R3-D03 induced an important decrease of CD8α-expressing cells, which lasted until day 22 (FIG. 10A). However, the depletion of CD8α-expressing cells induced by antibody clones MP08-R3-C11 and MP09-R3-D03 was not total.

[0501] Since antibody clone MP08-R3-F08 induced the most efficient depletion of the CD8α-expressing cells, another dose was tested. In order to evaluate the efficiency of depletion of the CD8α-expressing cells, antibody clone MP08-R3-F08 was injected into one additional monkey at a dose of 7.5 mg/kg bodyweight. Immediate confirmation of depletion was performed by flow cytometry analysis 1 hour and 6 hours after antibody injection. Follow-up of CD8α-expressing cell depletion was done at 24 hours, day 7, day 14, day 28, day 48 and day 62 after antibody injection.

[0502] Even at the lower dose of 7.5 mg/kg bodyweight, antibody clone MP08-R3-F08 induced a complete depletion of CD8α-expressing cells within 1 hour after injection. Additionally, the percentage of CD8+ cells remained at 0% until day 14 post injection. From day 14, CD8α-expressing cells started re-increasing very progressively and reached 2.5% by day 62 (FIG. 10B).

Example 8: Effect of the Anti-CD8α Antibody in a Mouse Model of Myocardial Infarction

Materials and Methods

Mice

[0503] B6; SJL-Tg(CD8aCD8b)57Scr/J were used (Jackson Laboratory strain #003202); and later designed as hCD8 mice, as described in Example 5 above.

[0504] Mice were maintained under identical standard conditions including housing, regular care, and normal chow diet in the same animal facility throughout the duration of the experiments (12 h light/dark cycle with ad libitum access to food and water). Mice were aged 9 to 15 weeks at the beginning of experiments. Both males and females were used.

Myocardial Infarction

[0505] Myocardial infarction (MI) was induced by permanent ligation of the left anterior descending artery in its proximal third, as previously described (Zouggari et al., Nature medicine, 2013). Mice were anesthetized by peritoneal injection of ketamine (100 mg per kg body weight; Ketamine 1000, Vibrac) and Xylazine (10 mg per kg body weight; Xylazine 2%, Bayer), intubated and ventilated with air using a small animal respirator (SAR 830/AP Ventilator, CWE Incorporated, Ardmore, USA). Body temperature was maintained to 37° C. using a homeothermic blanket system (Harvard Apparatus). The chest wall was shaved and a thoracotomy was performed in the fourth left intercostal space. The left ventricle was localized and the pericardial sac was opened. The left anterior descending artery was permanently ligated using a 7/0 PVDF monofilament suture (Peters surgical, France) 5 mm below the site of its emergence from the left atrium. Permanent whitening of the myocardium below the ligation was considered indicative of a successful coronary occlusion. The thorax and the skin were closed with 6/0 PVDF monofilament sutures (Peters surgical, France), with local application of lidocaine (Lidocaine 20 mg/mL, Aguettant). The endotracheal tube was removed once spontaneous respiration resumed, and mice were placed in a cage on a warm pad maintained at 37° C. until completely vigil. To reduce pain, buprenorphine (0.1 mg/kg) was injected before, 6 hours after the onset of surgery and twice a day for 48 to 72 hours.

Antibody Injection

[0506] One hour after the end of surgery, mice were injected intravenously with anti-CD8 α antibody clone MP08-R3-F08 or sterile saline solution. Using a 29 Gauge insulin syringe, 100 μ L of a solution containing either 100 g of anti-CD8 α antibody clone MP08-R3-F08 or saline solution was slowly injected in the retro-orbital sinus over 10 seconds. Mice were then placed in a cage on a warm pad maintained at 37° C. and returned to their home cage after complete recovery from anesthesia.

Organ Harvesting

[0507] Mice were euthanized using 150 mg/kg Euthasol in 250 μ L of saline solution injected intraperitoneally upon isoflurane sedation. Blood from the vena cava was collected using a 27-gauge needle. Spleen and heart are weighed and rinsed in PBS for further analysis.

Cell Suspension Preparation and Flow Cytometry

[0508] Mice were sacrificed at day 3 or 21 post-MI. Peripheral blood was drawn via inferior vena using a syringe with a 27 Gauge needle primed with heparin solution. For blood staining, erythrocytes were lysed using BD FACS lysis solution (BD Biosciences). Spleens were surgically removed and dissociated, obtaining a single cell suspension that was filtered through 40 m nylon mesh (BD Biosciences). Cell suspensions were centrifuged at 400 g for 15 min at 4° C. Red blood cells were removed using a red blood lysis buffer (Sigma-Aldrich) and cells were washed with PBS and resuspended in 5 mL PBS supplemented with 3% FBS.

[0509] General characteristics of antibodies are summarized in Table 5 below. Surface marker staining was performed by incubating the cell suspension (50 μ L blood or 50 μ L of splenocytes suspension) with 10 μ L of the antibody cocktail for 30 minutes at 4° C. before washing with PBS. Forward scatter (FSC) and side scatter (SSC) parameters were used to gate live cells excluding red blood cells, debris, and cell aggregates. Cells were analyzed using a BD LSR Fortessa flow cytometer (BD Biosciences) and data were analyzed using FlowJo v10.9 software (BD Biosciences).

[0510] The expression level of hCD8 α was assessed in total CD45+ cells.

TABLE 5

Antibodies used for flow cytometry in mouse model of MI experiments					
Marker	Dye	Manufacturer	Clone	Reference	Dilution
CD45	PerCP	BD Bioscience	30-F11	557235	1:100
CD11b	BV605	BD Bioscience	M1/70	563015	1:100
CD3	PerCP-Cy5,5	eBioscience	145-2C11	45-0031-82	1:100
B220	V500	BD Bioscience	RA3-6B2	561226	1:100
CD4	Efluor450	eBioscience	RM4-5	48-0042-82	1:100
mCD8 α	AF700	BD Bioscience	53-6.7	557959	1:100
hCD8 α	FITC	eBioscience	HIT8 α	11-0089-42	1:100
NK1.1	APC	eBioscience	PK136	25-5941-82	1:100
Ly6C	APC-Cy7	BD Bioscience	AL-21	560596	1:100
Ly6G	PE	BD Bioscience	1A8	551461	1:100

Infarct Size Evaluation at Day 3 by TTC Staining

[0511] To evaluate infarct size at an early time point (3 days after MI), 2,3,5-triphenyltetrazolium chloride (TTC, Sigma Aldrich) was used. Hearts were harvested and sectioned into four 1.5-2 mm thick slices from the apex, in a semi-frozen state. The slices were then incubated for 40 min in PBS—1% TTC solution at 37° C. Sections were then rinsed twice 5 min with PBS at room temperature. Slices were photographed and infarct size was calculated using ImageJ blindly by measuring the total tissue area and the infarcted tissue area on each tissue slice.

Echocardiography

[0512] Cardiac function was assessed 21 days after MI. Chest wall was shaved using depilatory cream. Upon isoflurane anesthesia (2.5 % Isoflurane & 2 μ L/min oxygen for induction; 2% isoflurane & 1 μ L/min oxygen for echography), cardiac function was recorded using a VEVO 2100 Imaging System and a MS400 transducer appropriate for mouse cardiovascular imaging (18-38 MHz) (VisualSonics, Canada). Data were analyzed using the cardiac package in VEVO-Lab (VisualSonics, Canada). Left ventricular end-diastolic volume (EDV) and end-systolic volume (ESV) were calculated on cine-loops using the 2-dimensional area-length method, allowing the calculation of left ventricular ejection fraction (EF) as $EF (\%) = [(EDV - ESV) / EDV] \times 100$.

Histopathological Analyses

[0513] Cardiac remodeling following MI was assessed at day 21. Hearts were harvested, rinsed in PBS; atria and right ventricle were excised and the remaining left ventricle (LV) was embedded in Tissue-TEK O.C.T. compound (Sakura) and frozen in liquid nitrogen-cooled isopentane and stored at -80° C. Hearts were cut along their length into 8 m thick cardiac muscle cryosections (CM 3050S, Leica). Serial sections were mounted on microscope slides, each section being spaced from the next one by 500 m. In this way, 8 cuts allowed entire LV tissue analysis. Masson's trichrome and Sirius Red staining were performed for infarct size and interstitial fibrosis evaluation, respectively. Infarct size was calculated as a percentage of infarct area to total LV area. The collagen volume fraction was calculated as the ratio of the total area of interstitial fibrosis to the total cardiac area in the entire visual field of the section.

Results

[0514] In order to evaluate the effect of the anti-CD8 α antibody of the invention on myocardial infarction, the anti-CD8 α antibody clone MP08-R3-F08 was injected into transgenic mice expressing human CD8 α one hour after induction of myocardial infarction.

[0515] First, depletion of hCD8 α cells in the blood and spleen of hCD8 α -expressing transgenic mice which received either saline solution (control) or the anti-CD8 α antibody clone MP08-R3-F08 was assessed, 3 days after induction of myocardial infarction and injection of saline solution or anti-CD8 α antibody clone MP08-R3-F08.

[0516] As shown on FIG. 12A-C and FIG. 13A-C depletion of CD45+hCD8 α + cells is total in the blood of mice treated with the anti-CD8 α antibody clone MP08-R3-F08 (FIGS. 12B & C) as compared to control mice (FIGS. 12A & C), while there is more than 90% depletion in the spleen

of mice treated with the anti-CD8 α antibody clone MP08-R3-F08 (FIGS. 13B & C) as compared to control mice (FIGS. 13A & C).

[0517] In addition, 3 days after induction of myocardial infarction and injection of saline solution or anti-CD8 α antibody clone MP08-R3-F08, hearts were harvested from the mice, sectioned and stained with 2,3,5-triphenyltetrazolium chloride (TTC) in order to measure the necrosis induced after myocardial infarction.

[0518] As shown of FIG. 14, there is a significant 20% decrease in necrosis area in transgenic mice treated with the anti-CD8 α antibody clone MP08-R3-F08, as compared to control transgenic mice treated with saline solution.

Conclusion

[0519] These results show that the anti-CD8 α antibody clone MP08-R3-F08 possess a strong specificity for its target CD8 α isoform 2 (both secreted form and tethered secreted form). Additionally, the anti-CD8 α antibody clone MP08-R3-F08 does not exhibit any off-target binding.

[0520] Furthermore, all three anti-CD8 α antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) are

capable of depleting CD8 α -expressing cells in vivo in non-human primates. All anti-CD8 α antibody clones (i.e., MP08-R3-F08, MP08-R3-C11 and MP09-R3-D03) induce a strong or even complete depletion of CD8 α -expressing cells within 1 hour after antibody injection in non-human primates. Moreover, depletion is total for the anti-CD8 α antibody clone MP08-R3-F08 for at least 14 days. 14 days post-depletion, repopulation of the CD8 α -expressing cells begins slowly and progressively at the 7.5 mg/kg body-weight dose, and reaches only 2.5% at day 62 post-injection. Thus, depletion of CD8 α -expressing cells by the anti-CD8 α antibodies of the invention is fast, strong if not complete, and transient. Repopulation of CD8 α -expressing cells after depletion is slow and progressive.

[0521] Moreover, the anti-CD8 α antibody clone MP08-R3-F08 is able to limit the necrosis area induced after myocardial infarction in hCD8 α -expressing transgenic mice.

[0522] Taken together, these results indicate that the anti-CD8 α antibodies as described herein could be valuable therapeutic tools for transiently depleting CD8 α -expressing cells in a controlled manner and for the treatment of cardiovascular diseases such as myocardial infarction.

SEQUENCE LISTING

```
Sequence total quantity: 36
SEQ ID NO: 1          moltype = AA  length = 17
FEATURE              Location/Qualifiers
source                1..17
                     mol_type = protein
                     organism = synthetic construct
VARIANT              1
                     note = X is D or S
VARIANT              7
                     note = X is S or R
VARIANT              10
                     note = X is D or G or Y
SEQUENCE: 1
XISGSSXYIX YADFKVG                                     17

SEQ ID NO: 2          moltype = AA  length = 14
FEATURE              Location/Qualifiers
source                1..14
                     mol_type = protein
                     organism = synthetic construct
VARIANT              3
                     note = X is N or no amino acid
VARIANT              4
                     note = X is Y or no amino acid
VARIANT              5
                     note = X is Y or G or no amino acid
VARIANT              6
                     note = X is D or S or no amino acid
VARIANT              8
                     note = X is S or N or F
VARIANT              9
                     note = X is A or G
VARIANT              10
                     note = X is S or D or N
VARIANT              11
                     note = X is A or G
SEQUENCE: 2
SSXXXXYXXX XMDV                                         14

SEQ ID NO: 3          moltype = AA  length = 14
FEATURE              Location/Qualifiers
source                1..14
                     mol_type = protein
                     organism = synthetic construct
VARIANT              10
```

-continued

VARIANT	note = X is G or N or Y 11	
VARIANT	note = X is S or Y 12	
SEQUENCE: 3 AGTSSDVGGX XXVS	note = X is S or Y	14
SEQ ID NO: 4 FEATURE source	moltype = AA length = 7 Location/Qualifiers 1..7 mol_type = protein organism = synthetic construct	
VARIANT	1	
VARIANT	note = X is Q or S or Y 4	
SEQUENCE: 4 XDSXRPS	note = X is Y or S	7
SEQ ID NO: 5 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein organism = synthetic construct	
VARIANT	3	
VARIANT	note = X is Y or D 5	
SEQUENCE: 5 SSXTXYSTRV	note = X is Y or Q or S	10
SEQ ID NO: 6 FEATURE source	moltype = AA length = 5 Location/Qualifiers 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 6 NNAMN		5
SEQ ID NO: 7 FEATURE source	moltype = AA length = 17 Location/Qualifiers 1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 7 DISGSSRYIG YADFKVG		17
SEQ ID NO: 8 FEATURE source	moltype = AA length = 14 Location/Qualifiers 1..14 mol_type = protein organism = synthetic construct	
SEQUENCE: 8 SSNYDYDNAD AMDV		14
SEQ ID NO: 9 FEATURE source	moltype = AA length = 14 Location/Qualifiers 1..14 mol_type = protein organism = synthetic construct	
SEQUENCE: 9 AGTSSDVGGN SYVS		14
SEQ ID NO: 10 FEATURE source	moltype = AA length = 7 Location/Qualifiers 1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 10 SDSSRPS		7
SEQ ID NO: 11 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein	

-continued

SEQUENCE: 11 SSYTQYSTRV	organism = synthetic construct	10
SEQ ID NO: 12 FEATURE source	moltype = AA length = 5 Location/Qualifiers 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 12 NYDMN		5
SEQ ID NO: 13 FEATURE source	moltype = AA length = 17 Location/Qualifiers 1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 13 DISGSSSYID YADFVKG		17
SEQ ID NO: 14 FEATURE source	moltype = AA length = 11 Location/Qualifiers 1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 14 SSYYSGSGMD V		11
SEQ ID NO: 15 FEATURE source	moltype = AA length = 14 Location/Qualifiers 1..14 mol_type = protein organism = synthetic construct	
SEQUENCE: 15 AGTSSDVGGG SSVS		14
SEQ ID NO: 16 FEATURE source	moltype = AA length = 7 Location/Qualifiers 1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 16 QDSYRPS		7
SEQ ID NO: 17 FEATURE source	moltype = AA length = 10 Location/Qualifiers 1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 17 SSYTYSTRV		10
SEQ ID NO: 18 FEATURE source	moltype = AA length = 5 Location/Qualifiers 1..5 mol_type = protein organism = synthetic construct	
SEQUENCE: 18 NYVMN		5
SEQ ID NO: 19 FEATURE source	moltype = AA length = 17 Location/Qualifiers 1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 19 SISGSSRYIY YADFVKG		17
SEQ ID NO: 20 FEATURE source	moltype = AA length = 12 Location/Qualifiers 1..12 mol_type = protein organism = synthetic construct	
SEQUENCE: 20 SSGSYFGNAM DV		12

-continued

SEQ ID NO: 21	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		
AGTSSDVGGY YSVS		14
SEQ ID NO: 22	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 22		
YDSSRPS		7
SEQ ID NO: 23	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		
SSDTSYSTRV		10
SEQ ID NO: 24	moltype = AA length = 123	
FEATURE	Location/Qualifiers	
source	1..123	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 24		
EVQLVESGGG LVKPGGSLRL SCAASGFTFS NNAMNWRQA PGKGLEWISD ISGSSRYIGY		60
ADFKVGRFTI SRDPAKNSLY LQMSLRAED TAVYYCVRSS NYDYDADAM DVWGRGTLVT		120
VSS		123
SEQ ID NO: 25	moltype = AA length = 110	
FEATURE	Location/Qualifiers	
source	1..110	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 25		
QSVLTQPASV SGSPGQSITI SCAGTSSDVG GNSYVSWYQQ HPGKAPKLMI YSDSSRPSGV		60
SNRFSGSKSG NTASLTISGL QAEDADYYC SSYTQYSTRV FGGGTKLAVL		110
SEQ ID NO: 26	moltype = AA length = 120	
FEATURE	Location/Qualifiers	
source	1..120	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 26		
EVQLVESGGG LVKPGGSLRL SCAASGFTFS NYDMNWRQA PGKGLEWISD ISGSSSYIDY		60
ADFKVGRFTI SRDPAKNSLY LQMSLRAED TAVYYCVRSS YSGSGMDVW GRGTLVTSS		120
SEQ ID NO: 27	moltype = AA length = 110	
FEATURE	Location/Qualifiers	
source	1..110	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 27		
QSVLTQPASV SGSPGQSITI SCAGTSSDVG GGSSVSWYQQ HPGKAPKLMI YQDSYRPSGV		60
SNRFSGSKSG NTASLTISGL QAEDADYYC SSYTYSTRV FGGGTKLAVL		110
SEQ ID NO: 28	moltype = AA length = 121	
FEATURE	Location/Qualifiers	
source	1..121	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 28		
EVQLVESGGG LVKPGGSLRL SCAASGFTFS NYDMNWRQA PGKGLEWISD ISGSSRYIY		60
ADFKVGRFTI SRDPAKNSLY LQMSLRAED TAVYYCVRSS GSYFGNAMDV WGRGTLVTSS		120
S		121
SEQ ID NO: 29	moltype = AA length = 110	
FEATURE	Location/Qualifiers	
source	1..110	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 29
 QSVLTQPASV SGSPGQSITI SCAGTSSDVG GYYSVSWYQQ HPGKAPKLMI YYDSSRPSGV 60
 SNRFSGSKSG NTASLTISGL QAEDEADYYC SSSTSYSYTRV FGGGTKLAVL 110

SEQ ID NO: 30 moltype = AA length = 235
 FEATURE Location/Qualifiers
 source 1..235
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 30
 MALPVTALLL PLALLLHAAR PSQFRVSPLD RTWNLGETVE LKCQVLLSNP TSGCSWLFQP 60
 RGAAASPTFL LYLSQNKPKA AEGLDTQRFS GKRLGDTFVL TLSDFRRENE GYYFCSALSN 120
 SIMYFSHFVP VFLPAKPTTT PAPRPPTPAP TIASQPLSLR PEACRPAAGG AVHTRGLDFA 180
 CDIYIWAPLA GTCGVLLLSL VITLYCNHRN RRRVKCKPRP VVKSGDKPSL SARYV 235

SEQ ID NO: 31 moltype = DNA length = 1461
 FEATURE Location/Qualifiers
 source 1..1461
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 31
 gcggccgcaa actacaagac agacttgcaa aagaaggcat gcacagctca gcaactgctct 60
 gttgcctggt cctcctgact ggggtgaggg ccgaggtgca gctggtggag agcgggggca 120
 gcctgggtgaa gccaggggga agcctgaggg tgagctgtgc cgcacccggg ttacatcttt 180
 ccaacaacgc catgaactgg gtgaggcagg cacccggaag agggctggag tggatttcag 240
 acatttcagg cagtagcaga tacatcggct acgcccagct cgtgaagggc aggttcacaa 300
 tcagcagaga caacgctaag aacagcctgt acctgcagat gaactccctg agggccgagg 360
 acacagccgt gtactactgc gtgaggagct ccaactacta tgactacaac gccgacgcta 420
 tggatgtgtg gggaaagggc accctgggtga ccgtgagctc cgctagcacc aaggggcccat 480
 cggctctccc cctggcacc cctctcaaga gcacctctgg gggcacagcg gccctgggct 540
 gctcgggtcaa ggactacttc ccgaaccggg tgacggtgtc gtggaactca ggcgcctga 600
 ccagcggcgt gcacaccttc ccggctgttc tacagtcttc aggaactctac tccctcagca 660
 gcgtgggtgac cgtgccctcc agcagcttgg gcacccagac ctacatctgc aacgtgaatc 720
 acaagcccgag caacaccaag gtggacaaga aagttgagcc caaatcttgt gacaaaactc 780
 acacatgccc accgtgccc gcacctgaac tcttgggggg accgtcagtc tctctcttcc 840
 ccccaaaacc caaggacacc ctcatgatct cccggacccc cgaggtcaca tgcgtgggtg 900
 tggacgtgag ccacgaagac cctgaggtca agttcaactg gtacgtggac ggcgtggagg 960
 tgcataatgc caagacaaag ccgaggagg agcagtaaca cagcagctac cgtgtgtgca 1020
 gcgtcctcac cgtcctgcac caggactggc tgaatggcaa ggagtagaag tgcagggtct 1080
 ccaacaaagc cctcccagcc cccatcgaga aaacctctc caaagcaca gggcagcccc 1140
 gagaaccaca ggtgtacacc ctgcccccat cccggggagg gatgaccaag aaccagggtc 1200
 gcctgacctg cctgggtcaa ggcttctatc ccagcgacat cgccgtggag tgggagagca 1260
 atgggcagcc ggagaacca cgcctcccg gctggactcc gacggctcct 1320
 tcttctctca cagaagctc accgtggaca agagcaggtg gcagcagggg aacgtcttct 1380
 catgctcgt gatcatgag gctctgcaca accactacac gcagaagagc ctctccctgt 1440
 ctccgggtaa atgattctag a 1461

SEQ ID NO: 32 moltype = DNA length = 750
 FEATURE Location/Qualifiers
 source 1..750
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 32
 gcggccgcaa actacaagac agacttgcaa aagaaggcat gcacagctca gcaactgctct 60
 gttgcctggt cctcctgact ggggtgaggg ccagtcctgt gctgaccag cccgctccg 120
 tgtccggaag cccaggacag tccatacca tctcctgcgc cggcacctcc tccgacgtgg 180
 gaggaacacg ctacgtgtcc ttgtatcaac agcaccgccg caaagcacc aaactgatga 240
 tctactccga cagttcccgc cctccggcg tgtccaacag gttcagcggc tccaagagcg 300
 gcaacaccgc ctccctgact atctccggcc tgcagggtga agacgaagcc gactactact 360
 gctcctccta caccagctac tccaccgcg tgttcggcgg cggcacaaa ctggccgtgc 420
 tgggtcagcc caaggccaac ccaactgtca ctctgttccc gccctcctct gaggaagctc 480
 aagccaacaa ggccacacta gtgtgtctga tcagtgaact ctaccgggga gctgtgacag 540
 tggcctggaa ggcagatggc agccccgtca aggggggagt ggagaccacc aaacctcca 600
 aacagagcaa caacaagtac gcggccagca gctacctgag cctgacgccc gagcagtgga 660
 agtcccacag aagctacagc tgccagggtc cgcaggaagg gagcaccgtg gagaagacag 720
 tggccctcac agaattgtca tgattctaga 750

SEQ ID NO: 33 moltype = DNA length = 1452
 FEATURE Location/Qualifiers
 source 1..1452
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 33
 gcggccgcaa actacaagac agacttgcaa aagaaggcat gcacagctca gcaactgctct 60
 gttgcctggt cctcctgact ggggtgaggg ccgaggtgca gctggtggag agcgggggca 120
 gcctgggtgaa gccaggggga agcctgaggg tgagctgtgc cgcacccggg ttacatcttt 180

-continued

```

ccaactacga catgaactgg gtgaggcagg cccccggcaa aggactggag tggatttcag 240
acatctccgg cagcagttcc tacatcgact acgctgattt tgtgaagggc aggttcacaa 300
tctcccggga caacgctaag aactccctgt acctgcagat gaacagctcg agggccgagg 360
acaccgctgt gtactactgc gtgaggtcct cctactatag tggcagcggc atggacgtgt 420
ggggccgggg aaccctgggt accgtgtcct ctgctagcac caagggccca tcggtcttcc 480
ccctggcacc ctctccaag agcacctctg ggggcacagc ggccctgggc tgctgtgtca 540
aggactactt ccccgcaacc gtgacgggtg cgtggaactc aggcgcctcg accagcggcg 600
tgcacacctt cccggctgtc ctacagtcct caggactcta ctccctcagc agcgtggtga 660
ccgtgccctc cagcagcttg ggcacccaga cctacatctg caacgtgaat cacaagccca 720
gcaacaccaa ggtggacaag aaagttaggc ccaaatcttg tgacaaaact cacacatgcc 780
caccgtgccc agcacctgaa ctctcggggg gaccgtcagt cttctcttcc cccccaaac 840
ccaaggacac cctcatgatc tcccggaccc ccgaggtcac atgctgtgtg gtggacgtga 900
gccacgaaga ccttgagggtc aagttcaact ggtacgtgga cggcgtggag gtgcataatg 960
ccaagacaaa gcccggggag gagcagtaca acagcacgta ccgtgtgtgc agcgtcttca 1020
ccgtcctgca ccaggactgg ctgaatggca aggagtacaa gtgcaaggtc tccaacaaag 1080
ccctccagc ccccatcgag aaaaccatct ccaaagccaa agggcagccc cgagaaccac 1140
agggtgtcac cctgccccca tcccgggagg agatgaccaa gaaccaggtc agcgtgacct 1200
gcctggtcaa aggtttctat ccacgcgaca tcgcccgtga gtgggagagc aatgggcagc 1260
cggagaacaa ctacaagacc acgcctcccg tgctggactc cgacggctcc ttcttctct 1320
acagcaagct caccgtggac aagagcaggt ggcagcaggg gaacgtcttc tcatgctccg 1380
tgatgcatga ggctctgcac aaccactaca cgcagaagag cctctccctg tctccgggtg 1440
aatgattcta ga 1452

```

```

SEQ ID NO: 34      moltype = DNA length = 750
FEATURE
source             Location/Qualifiers
                   1..750
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 34
gcgcccgcaa actacaagac agacttgcaa aagaaggcat gcacagctca gactgctct 60
gttgcttggt cctcctgact ggggtgaggg cccagtcctg gctgacccag cccgcctccg 120
tgtccggaag cccaggacag tccatcacca tctcctgcgc cggcacctcc tccgacgtgg 180
gaggaggaag ctccgtgagc ttgtatcaac agcaccccgg caaagcacc aaactgatga 240
tctaccagga ctgtacagc cctccggcg gtccaacag gttcagcggc tccaagagcg 300
gaaacacagc ctctctgacc atctccggcc tgcaggctga ggacgaagcc gactactact 360
gtcatcctca cacttactat tccaccaggg tgttcggagg cggaacaaaa ctggccgtgc 420
tgggtcagcc caaggccaac cccactgtca ctctgttccc gccctcctct gaggaagctcc 480
aagccaacaa ggccacacta gtgtgtctga tcaagtactt ctaccgggga gctgtgacag 540
tggcttgaaa ggcagatggc agccccgtca agggcggagt ggagaccacc aaaccctcca 600
aacagagcaa caacaagacc gcggccagca gctacctgag cctgacgccc gagcagtgga 660
agtcccacag aagctacagc tgcagggtca gcctgaagg gagcacctg gagaaagacag 720
tggcccttac agaattgtca tgattctaga 750

```

```

SEQ ID NO: 35      moltype = DNA length = 1455
FEATURE
source             Location/Qualifiers
                   1..1455
                   mol_type = other DNA
                   organism = synthetic construct

```

```

SEQUENCE: 35
gcgcccgcaa actacaagac agacttgcaa aagaaggcat gcacagctca gactgctct 60
gttgcttggt cctcctgact ggggtgaggg ccgagggtgca gctggtggag agcgggggca 120
gctcggtgaa gccaggggga agcctgaggg tgagctgtgc cgcacccggg ttacatattt 180
ccaaetacta ttgaactgg gtgaggcagg cccctggcaa gggcctggag tggatttcat 240
ctatctccgg cagtagtcta tacatctact acgctgactt tgtgaagggc cgttccacca 300
tctccaggga caacgctaag aactccctgt atctgcagat gaactccctg cgcgccgagg 360
acaccgctgt gtactactgc gtgcgctcct ccggcagcta cttcggaat gctatggatg 420
tgtgggggag ggggaccctg gtgaccgtgt cctccgctag caccaagggc ccacgtgct 480
tcccctggc accctcctcc aagagcacct ctgggggcac agcggccctg ggctgcttg 540
tcaaggacta cttccccgaa ccgggtgacgg tgtcgtggaa ctcaggcgcc ctgaccagcg 600
gcgtgcacac cttcccggtc gtccctacagt cctcaggact ctactccctc agcagcgtgg 660
tgaccgtgcc ctccagcagc ttgggcaccc agacctacat ctgcaacgtg aatcacaagc 720
ccagcaacac caaggtggac aagaaaagtg agcccaaatc ttgtgacaaa actcacacat 780
gcccaccgtg cccagcacct gaactcctgg ggggaccgtc agtcttctcc tccccccaa 840
aaccacaagg acccctcatg atctcccggg cccccgaggt cacatgcgtg gtggtggagc 900
tgagccacga agaccctgag gtcaagttca actggtacgt ggacggcgtg gaggtgcata 960
atgccaaagc aaagccgcgg gaggagcagt acaacagcac gtaccgtgtg gtcagcgtcc 1020
tcaccgtcct gcaccaggac tggctgaatg gcaaggagta caagtgaag gtctccaaca 1080
aagccctccc agccccctac gagaaaacca tctccaaagc caaagggcag ccccgagaac 1140
cacagggtga caccctgccc ccatcccggg aggagatgac caagaaccag gtcagcctga 1200
cctgcttggt caaaggcttc tatcccagcg acatcgccgt ggagtgggag agcaatgggc 1260
agccggagaa caactacaag accacgctc ccgtgctgga ctccgacggc tcttcttcc 1320
tctacagcaa gctcaccgtg gacaagagca ggtggcagca ggggaacgtc ttctcatgct 1380
ccgtgatgca tgaggctctg cacaaccact acacgcagaa gagcctctcc ctgtctccgg 1440
gtaaatgatt ctaga 1455

```

```

SEQ ID NO: 36      moltype = DNA length = 750

```

-continued

FEATURE	Location/Qualifiers
source	1..750
	mol_type = other DNA
	organism = synthetic construct

SEQUENCE: 36

gcggccgcaa	actacaagac	agacttgcaa	aagaaggcat	gcacagctca	gcactgctct	60
gttgccctggt	cctcctgact	gggggtgagg	cccagtcctg	gctgacccag	cccgcctccg	120
tgcccggaag	cccaggacag	tccatcacca	tctcctgcgc	cggcacctcc	tccgacgtgg	180
gaggatacta	ctccgtgagc	tggtatcaac	agcaccgccg	caaagccccc	aaactgatga	240
tctactacga	cagctcccg	ccctccggcg	tgtctaacag	gttctccggc	tccaagagcg	300
gcaacaccgc	cagcctgacc	atctccggcc	tgcaggcaga	agacgaggcc	gactactact	360
gctcctccga	cacctctac	agcaccagg	tggtccggcg	tggcaccaag	ctggccgtgc	420
tgggtcagcc	caaggccaac	cccactgtca	ctctgttccc	gccctcctct	gaggagctcc	480
aagccaacaa	ggccacacta	gtgtgtctga	tcagtgaact	ctaccgggga	gctgtgacag	540
tggcctgga	ggcagatggc	agccccgtca	aggcgggagt	ggagaccacc	aaaccctcca	600
aacagagcaa	caacaagtac	gcggccagca	gctacctgag	cctgacgccc	gagcagtgga	660
agtccacag	aagctacagc	tgccagggtca	cgcattgaagg	gagcaccgtg	gagaagacag	720
tggcccttac	agaatgttca	tgattctaga				750

1. An isolated anti-CD8 α antibody, or an antigen-binding fragment thereof, wherein:

- a) the variable region of the heavy chain (V_H) of said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

V_H -CDR1: NX_1X_2MN , wherein

X_1 is N or Y,

X_2 is D or A or Y,

V_H -CDR2: $X_3ISGSSX_4YIX_5YADFVK$ (SEQ ID NO: 1), wherein

X_3 is D or S,

X_4 is S or R,

X_5 is D or G or Y,

V_H -CDR3: $SSX_6X_7X_8X_9YX_{10}X_{11}X_{12}X_{13}MDV$ (SEQ ID NO: 2), wherein

X_6 is N or no amino acid,

X_7 is Y or no amino acid,

X_8 is Y or G or no amino acid,

X_9 is D or S or no amino acid,

X_{10} is S or N or F,

X_{11} is A or G,

X_{12} is S or D or N,

X_{13} is A or G; and

- b) the variable region of the light chain (V_L) of said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

V_L -CDR1: $AGTSSDVGGX_{14}X_{15}X_{16}VS$ (SEQ ID NO: 3), wherein

X_{14} is G or N or Y,

X_{15} is S or Y,

X_{16} is S or Y,

V_L -CDR2: $X_{17}DSX_{18}RPS$ (SEQ ID NO: 4), wherein

X_{17} is Q or S or Y,

X_{18} is Y or S,

V_L -CDR3: $SSX_{19}TX_{20}YSTRV$ (SEQ ID NO: 5), wherein

X_{19} is Y or D,

X_{20} is Y or Q or S.

2. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein:

- a) the V_H of said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

V_H -CDR1:

(SEQ ID NO: 6)

NNAMN,

(SEQ ID NO: 12)

NYDMN,

or

(SEQ ID NO: 18)

NYYMN;

V_H -CDR2:

(SEQ ID NO: 7)

DISGSSRYIGYADFVK,

(SEQ ID NO: 13)

DISGSSSYIDYADFVK,

or

(SEQ ID NO: 19)

SISGSSRYIYADFVK;

V_H -CDR3:

(SEQ ID NO: 8)

SSNYDYNDAMDV,

(SEQ ID NO: 14)

SSYYSYSGMDV,

or

(SEQ ID NO: 20)

SSGSYFGNAMDV;

- b) the V_L of said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises the three following CDRs:

V_L -CDR1:

(SEQ ID NO: 9)

AGTSSDVGGNSYVS,

(SEQ ID NO: 15)

AGTSSDVGGGSSVS,

or

(SEQ ID NO: 21)

AGTSSDVGGYYSVS;

-continued

V_L -CDR2: (SEQ ID NO: 10)
 SDSSRPS,
 (SEQ ID NO: 16)
 QDSYRPS,
 or
 (SEQ ID NO: 22)
 YDSSRPS
 V_L -CDR3: (SEQ ID NO: 11)
 SSYTQYSTRV,
 (SEQ ID NO: 17)
 SSYTYSTRV,
 or
 (SEQ ID NO: 23)
 SSDTSYSTRV.

3. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises the six following CDRs:

V_H -CDR1: (SEQ ID NO: 6)
 NNAMN,
 V_H -CDR2: (SEQ ID NO: 7)
 DISGSSRYIGYADFVKG,
 V_H -CDR3: (SEQ ID NO: 8)
 SSNYDYDADAMDV,
 V_L -CDR1: (SEQ ID NO: 9)
 AGTSSDVGGNSYVS,
 V_L -CDR2: (SEQ ID NO: 10)
 SDSSRPS,
 and
 V_L -CDR3: (SEQ ID NO: 11)
 SSYTQYSTRV;
 or
 V_H -CDR1: (SEQ ID NO: 12)
 NYDMN,
 V_H -CDR2: (SEQ ID NO: 13)
 DISGSSSYIDYADFVKG,
 V_H -CDR3: (SEQ ID NO: 14)
 SSYYSGSGMDV,
 V_L -CDR1: (SEQ ID NO: 15)
 AGTSSDVGGSSVS,
 V_L -CDR2: (SEQ ID NO: 16)
 QDSYRPS,

-continued
and

V_L -CDR3: (SEQ ID NO: 17)
 SSYTYSTRV;
 or
 V_H -CDR1: (SEQ ID NO: 18)
 NYYMN,
 V_H -CDR2: (SEQ ID NO: 19)
 SIGSSRYIYADFVKG,
 V_H -CDR3: (SEQ ID NO: 20)
 SSGSYFGNAMDV,
 V_L -CDR1: (SEQ ID NO: 21)
 AGTSSDVGGYYSVS,
 V_L -CDR2: (SEQ ID NO: 22)
 YDSSRPS,
 and
 V_L -CDR3: (SEQ ID NO: 23)
 SSDTSYSTRV.

4. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises a variable region of the heavy chain (V_H) comprising a sequence as set forth in SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28, or a sequence having at least 80% identity with SEQ ID NO: 24, SEQ ID NO: 26, or SEQ ID NO: 28.

5. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises a variable region of the light chain (V_L) comprising a sequence as set forth in SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29, or a sequence having at least 80% identity with SEQ ID NO: 25, SEQ ID NO: 27, or SEQ ID NO: 29.

6. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, comprises:

a V_H comprising a sequence as set forth in SEQ ID NO: 24, or a sequence having at least 80% identity with SEQ ID NO: 24, and a V_L comprising a sequence as set forth in SEQ ID NO: 25, or a sequence having at least 80% identity with SEQ ID NO: 25; or

a V_H comprising a sequence as set forth in SEQ ID NO: 26, or a sequence having at least 80% identity with SEQ ID NO: 26, and a V_L comprising a sequence as set forth in SEQ ID NO: 27, or a sequence having at least 80% identity with SEQ ID NO: 27; or

a V_H comprising a sequence as set forth in SEQ ID NO: 28, or a sequence having at least 80% identity with SEQ ID NO: 28, and a V_L comprising a sequence as set forth in SEQ ID NO: 29, or a sequence having at least 80% identity with SEQ ID NO: 29.

7. The isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, wherein said isolated anti-CD8 α antibody, or antigen-binding fragment thereof, is a CD8 α -expressing cell depleting antibody.

8. A fusion protein comprising the anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1.

9. A nucleic acid molecule encoding the anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1.

10. A pharmaceutical composition comprising the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1, and at least one pharmaceutically acceptable excipient.

11. A method for treating a CD8- related disease in a subject in need thereof, said method comprising administering to the subject at least one isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1.

12. The method according to claim 11, wherein said CD8-related disease is a cardiovascular disease.

13. An in vitro method for depleting CD8 α -expressing cells in a sample, comprising a step of contacting said sample with the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1.

14. An in vitro method for detecting and/or quantifying CD8 α in a sample, cell, tissue or organ, comprising contacting said sample, cell, tissue or organ with the isolated anti-CD8 α antibody, or antigen-binding fragment thereof, according to claim 1.

* * * * *