

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12409180
Kind Code	B2
Date of Patent	September 09, 2025
Inventor(s)	Spargo; Peter Lionel et al.

Formulation production process

Abstract

The present invention relates to a process for producing a sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles, wherein the process comprises: (a) heating ensifentrine particles at a temperature of from 100° C. to 220° C. to obtain sterile ensifentrine particles; and (b) combining the sterile ensifentrine particles with a sterile liquid vehicle to produce the sterile liquid pharmaceutical composition suitable for administration by inhalation. A process for producing an ampule comprising the sterile liquid pharmaceutical composition is also described.

Inventors: Spargo; Peter Lionel (Canterbury, GB), Haywood; Phillip A. (Buntingford, GB), French; Edward James (Canterbury, GB)

Applicant: Verona Pharma PLC (Cardiff, GB)

Family ID: 80934600

Assignee: Verona Pharma PLC (Cardiff, GB)

Appl. No.: 18/946234

Filed: November 13, 2024

Prior Publication Data

Document Identifier	Publication Date
US 20250064814 A1	Feb. 27, 2025

Foreign Application Priority Data

GB	2202297	Feb. 21, 2022
----	---------	---------------

Related U.S. Application Data

Publication Classification

Int. Cl.: A61K31/519 (20060101); A61K9/00 (20060101); A61K9/16 (20060101)

U.S. Cl.:

CPC A61K31/519 (20130101); A61K9/0075 (20130101); A61K9/1682 (20130101);

Field of Classification Search

CPC: C07D (217/00); A61K (31/475); A61K (31/519)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4482556	12/1983	Lal et al.	N/A	N/A
5378818	12/1994	Mayer et al.	N/A	N/A
5985878	12/1998	Stokbroekx et al.	N/A	N/A
6794391	12/2003	Oxford et al.	N/A	N/A
7105663	12/2005	Oxford et al.	N/A	N/A
7378424	12/2007	Oxford et al.	N/A	N/A
8221772	12/2011	Johnson et al.	N/A	N/A
8242127	12/2011	Oxford et al.	N/A	N/A
9062047	12/2014	Walker et al.	N/A	N/A
9700558	12/2016	Walker et al.	N/A	N/A
9717732	12/2016	Walker et al.	N/A	N/A
9956171	12/2017	Spargo	N/A	A61P 35/00
10463665	12/2018	Spargo et al.	N/A	N/A
10471063	12/2018	Walker et al.	N/A	N/A
10710998	12/2019	Spargo	N/A	N/A
10864213	12/2019	Abbott-Banner et al.	N/A	N/A
10945950	12/2020	Spargo	N/A	A61P 11/08
11491155	12/2021	Spargo et al.	N/A	N/A
11759467	12/2022	Abbott-Banner et al.	N/A	N/A
12168011	12/2023	Spargo	N/A	A61K 9/10
12194045	12/2024	Spargo et al.	N/A	N/A
12251384	12/2024	Spargo et al.	N/A	N/A
2003/0036542	12/2002	Oxford et al.	N/A	N/A

2003/0229108	12/2002	De Belin et al.	N/A	N/A
2004/0076668	12/2003	Berchielli et al.	N/A	N/A
2004/0171828	12/2003	Oxford et al.	N/A	N/A
2004/0176353	12/2003	Oxford et al.	N/A	N/A
2004/0247628	12/2003	Lintz et al.	N/A	N/A
2005/0054655	12/2004	Beaulieu et al.	N/A	N/A
2005/0186276	12/2004	Berchielli et al.	N/A	N/A
2005/0261271	12/2004	Feng et al.	N/A	N/A
2007/0197489	12/2006	Karlsson	514/174	A61P 11/06
2008/0003283	12/2007	Feng et al.	N/A	N/A
2008/0108807	12/2007	Feng et al.	N/A	N/A
2008/0108808	12/2007	Feng et al.	N/A	N/A
2008/0113993	12/2007	De Belin et al.	N/A	N/A
2008/0125437	12/2007	Dong et al.	N/A	N/A
2008/0142759	12/2007	Pays	N/A	N/A
2008/0161562	12/2007	Feng et al.	N/A	N/A
2008/0177064	12/2007	Feng et al.	N/A	N/A
2008/0188501	12/2007	Feng et al.	N/A	N/A
2008/0199410	12/2007	Johnson et al.	N/A	N/A
2008/0206163	12/2007	Oxford et al.	N/A	N/A
2008/0254127	12/2007	Watanabe et al.	N/A	N/A
2009/0012059	12/2008	Feng et al.	N/A	N/A
2009/0075939	12/2008	He et al.	N/A	N/A
2010/0113413	12/2009	Dong et al.	N/A	N/A
2010/0204471	12/2009	Dong et al.	N/A	N/A
2011/0076276	12/2010	Guo et al.	N/A	N/A
2011/0190261	12/2010	Dong et al.	N/A	N/A
2012/0251594	12/2011	Longest et al.	N/A	N/A
2012/0302533	12/2011	Oxford et al.	N/A	N/A
2013/0078256	12/2012	Guo et al.	N/A	N/A
2013/0158021	12/2012	Dong et al.	N/A	N/A
2013/0225616	12/2012	Walker et al.	N/A	N/A
2014/0242174	12/2013	Walker	N/A	N/A
2016/0000790	12/2015	Walker et al.	N/A	N/A
2016/0008363	12/2015	Walker et al.	N/A	N/A
2017/0112839	12/2016	Abbott-Banner et al.	N/A	N/A
2017/0239178	12/2016	Spargo et al.	N/A	N/A
2017/0266190	12/2016	Walker et al.	N/A	N/A
2018/0021337	12/2017	Spargo et al.	N/A	N/A
2018/0369139	12/2017	Spargo et al.	N/A	N/A
2019/0330206	12/2018	Spargo	N/A	N/A
2020/0016158	12/2019	Spargo et al.	N/A	N/A
2021/0106585	12/2020	Abbott-Banner et al.	N/A	N/A
2021/0379053	12/2020	Spargo et al.	N/A	N/A
2022/0265549	12/2021	Spargo et al.	N/A	N/A
2023/0112220	12/2022	Spargo et al.	N/A	N/A
2024/0165023	12/2023	Spargo et al.	N/A	N/A

2024/0165054	12/2023	Rheault et al.	N/A	N/A
2024/0165055	12/2023	Rheault et al.	N/A	N/A
2024/0165117	12/2023	Spargo et al.	N/A	N/A
2024/0173327	12/2023	Rickard et al.	N/A	N/A
2024/0382489	12/2023	Spargo et al.	N/A	N/A
2025/0000866	12/2024	Spargo et al.	N/A	N/A
2025/0064814	12/2024	Spargo et al.	N/A	N/A
2025/0064815	12/2024	Spargo et al.	N/A	N/A
2025/0073240	12/2024	Rheault et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
527997	12/1982	AU	N/A
2847693	12/1979	DE	N/A
4240981	12/1993	DE	N/A
19612194	12/1996	DE	N/A
19820947	12/1997	DE	N/A
19821263	12/1997	DE	N/A
19726241	12/1997	DE	N/A
0601401	12/1993	EP	N/A
0876366	12/1997	EP	N/A
0983334	12/1999	EP	N/A
1305026	12/2002	EP	N/A
1418896	12/2003	EP	N/A
1438019	12/2003	EP	N/A
1519711	12/2004	EP	N/A
1586571	12/2004	EP	N/A
1651611	12/2005	EP	N/A
1825467	12/2006	EP	N/A
1892269	12/2007	EP	N/A
1993540	12/2007	EP	N/A
2063903	12/2008	EP	N/A
2167476	12/2009	EP	N/A
2001213867	12/2000	JP	N/A
3675274	12/2004	JP	N/A
2005263780	12/2004	JP	N/A
101361145	12/2013	KR	N/A
WO-9726258	12/1996	WO	N/A
WO-9736039	12/1996	WO	N/A
WO-9845293	12/1997	WO	N/A
WO-9851772	12/1997	WO	N/A
WO-9901607	12/1998	WO	N/A
WO-0058308	12/1999	WO	N/A
WO-0058309	12/1999	WO	N/A
WO-0193841	12/2000	WO	N/A
WO-02/17894	12/2001	WO	N/A
WO-03000343	12/2002	WO	N/A
WO-03035030	12/2002	WO	N/A
WO-03037262	12/2002	WO	N/A

WO-2004004684	12/2003	WO	N/A
WO-2005007092	12/2004	WO	N/A
WO-2005007138	12/2004	WO	N/A
WO-2005021510	12/2004	WO	N/A
WO-2005095381	12/2004	WO	N/A
WO-2006061398	12/2005	WO	N/A
WO-2008023249	12/2007	WO	N/A
WO-2008036293	12/2007	WO	N/A
WO-2008140553	12/2007	WO	N/A
WO-2009005674	12/2008	WO	N/A
WO-2012020016	12/2011	WO	A61K 31/519
WO-2012037782	12/2011	WO	N/A
WO-2012051426	12/2011	WO	N/A
WO-2013/034909	12/2012	WO	N/A
WO-2013/034910	12/2012	WO	N/A
WO-2013063468	12/2012	WO	N/A
WO-2013092791	12/2012	WO	N/A
WO-2013118855	12/2012	WO	N/A
WO-2013128283	12/2012	WO	N/A
WO-2014/037726	12/2013	WO	N/A
WO-2014/037727	12/2013	WO	N/A
WO-2014140647	12/2013	WO	A61K 31/137
WO-2014140648	12/2013	WO	N/A
WO-2015173551	12/2014	WO	N/A
WO-2016042313	12/2015	WO	A61P 43/00
WO-2016128742	12/2015	WO	N/A
WO-2018020249	12/2017	WO	N/A
WO-2020074894	12/2019	WO	N/A
WO-2021028679	12/2020	WO	N/A
WO-2021/150268	12/2020	WO	N/A
WO-2021171034	12/2020	WO	N/A
WO-2022/261417	12/2021	WO	N/A
WO-2023/076205	12/2022	WO	N/A
WO-2023/156794	12/2022	WO	N/A
WO-2024/033624	12/2023	WO	N/A
WO-2024/033625	12/2023	WO	N/A
WO-2024/033626	12/2023	WO	N/A
WO-2024/033627	12/2023	WO	N/A
WO-2024/084212	12/2023	WO	N/A
WO-2025/003643	12/2024	WO	N/A

OTHER PUBLICATIONS

Akers (Sterile Drug Products: Formulation, Packaging, Manufacture and Quality. Published 2010) (Year: 2010). cited by examiner

Bjermer, L. et al., Efficacy and safety of a first-in-class inhaled PDE3/4 inhibitor (ensifentrine) vs salbutamol in asthma, Pulmonary Pharmacology & Therapeutics, vol. 58, (2019). cited by applicant

Blasko, G. et al., Pyrimido[1,6-a]pyrido[3,4-b]indoles as new platelet inhibiting agents, European Journal of Medicinal Chemistry, vol. 21, 2 (1986):91-5. cited by applicant

Chen, E.H. et al., Modifications of primaquine as antimalarials. 1. 5-Phenoxy derivatives of primaquine, *Journal of Medicinal Chemistry*, vol. 20, 8 (1977):1107-9. cited by applicant

Dong-Sheng, M. et al., Synthesis of Daimuron, *College of Chemistry and Chemical Engineering, Huaxue Yu Nianhe*, vol. 6, (2003):293-295. cited by applicant

Franciosi, L.G. et al., Efficacy and safety of RPL554, a dual PDE3 and PDE4 inhibitor, in healthy volunteers and in patients with asthma or chronic obstructive pulmonary disease: findings from four clinical trials, *The Lancet. Respiratory Medicine*, vol. 1, 9(2013):714-727. cited by applicant

Frank, A.W., Synthesis of some carbonyl derivatives of tris(aminomethyl)phosphine oxide, *Phosphorus and Sulfur and the Related Elements*, vol. 22, 3(1985):265-76. cited by applicant

Houlihan, W.J. et al., Synthesis and proton-NMR spectra, *Journal of Heterocyclic Chemistry*, vol. 19, 6(1982):1453-6. cited by applicant

Jansen, M., Derivatives of some nuclear methoxylated B-phenylethylamines, vol. 50, (1931):617-37. cited by applicant

Kienzle, F. et al., Synthesis of 6,7-dihydro-2H-pyrimido[6, 1-a]isoquinolin-4(3H)-ones and analogous compounds and their activity as blood platelet aggregation inhibitors, *Helvetica Chimica Acta*, vol. 69, 7(1986): 1671-80. cited by applicant

Kijima, I, et al., Synthesis and reactivities of triisocyanatoantimony, Japan, *Nippon Kagaku Kaishi*, vol. 12, (1986):1754-57. cited by applicant

Lai, B. et al., Synthesis of 2-substituted-6,7-dihydro-4H-pyrimido[6,1-a]thieno[2,3-c]- and [3,2-c]pyridin-4-ones, *Heterocycles*, vol. 24, 7(1986):1977-85. cited by applicant

Lal, B. et al., Trequinsin, a potent new antihypertensive vasodilator in the series of 2-(arylimino)-3-alkyl-9,10-dimethoxy-3,4,6,7-tetrahydro-2H-pyrimido[6,1-a]isoquinolin-4-ones, *Journal of Medicinal Chemistry*, vol. 27, 11(1984):1470-80. cited by applicant

Mannich, C. et al., *Berichte der Deutschen Chemischen Gesellschaft*, vol. 43, (1910):189-97. cited by applicant

No Author, European Medicines Agency, Guideline on the sterilisation of the medicinal product, active substance, excipient and primary container, Apr. 11, 2016. cited by applicant

No Author, European Medicines Agency, Guideline on the sterilisation of the medicinal product, active substance, excipient and primary container, Mar. 6, 2019. cited by applicant

Pai, N.R. et al., Synthesis of novel analogs 3,4-dihydro-1H-quinolin-2-one derivatives as typical Antidepressant, Sedative and anti-Parkinson agents, *Heterocyclic Letters*, vol. 2, 1 (2012):117-128. cited by applicant

Pfleiderer, V.W. et al., Synthesis of 9-substituted xanthines, *Justus Liebigs Annalen der Chemie*, vol. 631, (1960):168-74. cited by applicant

Ratel, M. et al., Imidazolium-Based Ionic Liquid Surfaces for Biosensing, *Analytical Chemistry*, vol. 85, 12 (2013):5770-5777. cited by applicant

Tran, P. et al., Structure-activity relationship of human glutaminyl cyclase inhibitors having an N-(5-methy-1H-imidazol-1-yl)propyl thiourea template, *Bioorganic & Medicinal Chemistry*, vol. 21, 13 (2013):3821-3830. cited by applicant

USP 1229.8 Dry Heat Sterilization 2018. cited by applicant

SPAN 20 (sorbitan monolaurate product page) (Year: 2024). cited by applicant

TWEEN 20 (polyoxyethylene sorbitan monolaurate product page) (Year: 2024). cited by applicant

Agusti et al., Global Initiative for chronic Obstructive Lung Disease 2023 Report: GOLD Executive Summary, *European Respiratory Journal Task Force Report*, 2023, 61:2300239, 1-26. cited by applicant

Anoro Ellipta Prescribing Information, 2013, revised Jun. 2023, 37 pages. cited by applicant

Bateman et al., Aclidinium bromide and formoterol fumarate as a fixed-dose combination in COPD: pooled analysis of symptoms and exacerbations from two six-month, multicentre, randomised studies (ACLIFORM and AUGMENT), *Respiratory Research*, 2015, 16(1):92, 1-13. cited by applicant

Bevespi Aerosphere Prescribing Information, 2016, revised Mar. 2023, 12 pages. cited by applicant

Breztri Aerosphere Prescribing Information, 2020, revised Jul. 2020, 19 pages. cited by applicant

De la Loge et al., Relationship Between FEV1 and Patient-Reported Outcomes Changes: Results of a Meta-Analysis of Randomized Trials in Stable COPD, *Chronic Obstr. Pulm. Dis.*, Mar. 15, 2016, 3(2):519-538. cited by applicant

Donohue, James F., Minimal clinically important differences in COPD lung function, *Chronic Obstr. Pulm. Dis.*, Mar. 2005, 2(1):111-124. cited by applicant

Duaklir Pressair Prescribing Information, 2019, revised Mar. 2019, 33 pages. cited by applicant

E-RS (EXACT-Respiratory Symptoms), User Manual (Version 3.0), Oct. 2014, 26 pages. cited by applicant

Eneje et al., A Single Dose of a Novel Nebulised PDE3/4 Inhibitor RPL554 Increased FEV1 in Patients with Cystic Fibrosis and Demonstrated a Reproducible Pharmacokinetic Profile, *Pediatric Pulmonology*, 2018, 53(Suppl. S2)S228, Abstract 215, 1 page. cited by applicant

Ferguson et al. Triple therapy with budesonide/glycopyrrolate/formoterol fumarate with co-suspension delivery technology versus dual therapies in chronic obstructive pulmonary disease (KRONOS): a double-blind, parallel-group, multicentre, phase 3 randomised controlled trial, *Lancet Respir. Med.*, 2018, 6:747-758. cited by applicant

Jones et al., Minimally Clinically Important Differences in Pharmacological Trials, *American Journal of Respiratory and Critical Care Medicine*, Feb. 1, 2014, 189(3):250-255. cited by applicant

Jones, Paul W., St. George's Respiratory Questionnaire: MCID, *Chronic Obstr. Pulm. Dis.*, 2005, 2(1):75-79. cited by applicant

Kumar et al., "Basic Concepts of Sterilization Techniques," *Research Journal of Pharmacology and Pharmacodynamics*, Oct.-Dec. 2021, 13(4):155-161. cited by applicant

Lonhala Magnair Prescribing Information, 1961, revised Dec. 2017, 64 pages. cited by applicant

Mahler et al., The MCID of the Transition Dyspnea Index is a Total Score of One Unit, *Chronic Obstr. Pulm. Dis.*, 2005, 2(1):99-103. cited by applicant

Miravittles M et al., Pharmacological strategies to reduce exacerbation risk in COPD: a narrative review, *Respiratory Research*, 2016, 17(1):112, 1-15. cited by applicant

Newman et al., Low Oral Bioavailability of RPL554, a First-in-Class Dual PDE3/4 Inhibitor, Demonstrates that Its Nebulized, Inhaled Formulation Is Appropriate for Delivering Optimal Pulmonary Dose, *Am. J. Respir. Crit. Care Med.*, 2018; 197:A3022, Abstract, 3 pages. cited by applicant

Rheault et al., The effect of fluconazole on the pharmacokinetics of ensifentrine in healthy individuals, *European Respiratory Journal*, 2021; 58:(Suppl. 65):PA2137, Abstract, 4 pages. cited by applicant

Rickard et al., The Dual Phosphodiesterase (PDE) 3 and 4 Inhibitor Ensifentrine Does Not Prolong QT Interval in Healthy Volunteers, *Am. J. Respir. Crit. Care Med.*, 2022, 205:A5605, Abstract, 1 page. cited by applicant

Singh et al., "The short term bronchodilator effects of the dual PDE3 and PDE4 inhibitor RPL554 in COPD," *European Respiratory Journal*, 2018; in press (<https://doi.org/10.1183/13993003.01074-2018>), 33 pages. cited by applicant

Singh et al., A Phase I, Randomised, Double Blind, Placebo Controlled, Study to Assess the Safety, Tolerability and Pharmacokinetics of Multiple Inhaled Doses of RPL554 Administered by Nebuliser to Healthy Male Subjects and Stable COPD Patients, *Am. J. Respir. Crit. Care Med.*, 2016, 193:A6838, Abstract, 3 pages. cited by applicant

Singh et al., A Phase I, Randomised, Double Blind, Placebo Controlled, Study to Assess the Safety, Tolerability and Pharmacokinetics of Single Inhaled Doses of the Dual Phosphodiesterase 3/4 (PDE3/4) Inhibitor RPL554 Administered by Nebuliser to Healthy Male Subjects, *Am. J. Respir. Crit. Care Med.*, 2016, 193:A6839, Abstract, 3 pages. cited by applicant

Stiolto Respimat Prescribing Information, 2015, revised Jan. 2025, 21 pages. cited by applicant

Striverdi Respimat Medical Review, Center for Drug Evaluation and Research, NDA 203108, 2012, 385 pages. cited by applicant

Striverdi Respimat Prescribing Information, 2014, revised Jan. 2025, 18 pages. cited by applicant

SYMBICORT Prescribing Information, 2006, revised Feb. 2009, 73 pages. cited by applicant

Terry et al., “Inhalation Therapy for Stable COPD: 20 Years of GOLD Reports,” *Advances in Therapy*, Apr. 3, 2020, 37:1812-1828. cited by applicant

Trujillo et al., Decrease in exacerbations during the coronavirus disease 2019 pandemic in a cohort of veterans with COPD, *Chronic Obstr. Pulm. Dis.*, 2021, 8(4): 572-579. cited by applicant

Utibron Neohaler Prescribing Information, 2015, revised May 2019, 28 pages. cited by applicant

Verona Pharma, Clinical Study Protocol, Version 1.0, May 28, 2018, 66 pages. cited by applicant

U.S. Appl. No. 18/825,447, filed Sep. 5, 2024. cited by applicant

U.S. Appl. No. 18/949,105, filed Nov. 15, 2024. cited by applicant

Abbott-Banner, K.H. et al., Dual PDE3/4 and PDE4 inhibitors: Novel treatments for COPD and other inflammatory airway diseases, *Basic & Clinical Pharmacology & Toxicology*, vol. 114, (2014):365-376. cited by applicant

Albert, RK et al., Azithromycin for prevention of exacerbations of Copd. *N. Engl. J. Med.* 2011; 365(8):689-98. cited by applicant

Anzueto, A. et al., Ensifentrine, a novel phosphodiesterase 3 and 4 inhibitor for the treatment of chronic obstructive pulmonary disease: randomized, double-blind, placebo-controlled, multicenter phase III trials (the ENHANCE Trials). *Am J Respir Crit Care Med.* 2023;208(4):406-416. cited by applicant

Anzueto, A. et al., Effect of Ensifentrine, a Novel PDE3 and PDE4 Inhibitor, on Lung Function, Symptoms and Exacerbations in Patients with COPD: The ENHANCE Trials (PowerPoint slides), *ATS 2023*, Washington DC May 19-24, 2023. cited by applicant

Anzueto, A. et al., Treatment with Ensifentrine, a Dual PDE3 and PDE4 Inhibitor, Significantly Reduced Exacerbation Rate and Risk in Subjects with COPD: Sub-group Results from the Phase 3 Trial, *ENHANCE-2* (Poster at *ATS 2023—May 19-24, 2023*). cited by applicant

Anzueto, A. et al., Treatment with ensifentrine, a dual PDE3 and PDE4 inhibitor, significantly reduced exacerbation rate and risk in subjects with COPD: sub-group results from the phase 3 trial, *enhance-2*, *Am J Respir Crit Care Med*, (2023):207:A4998. cited by applicant

Anzueto, A., Impact of exacerbations on COPD. *Eur. Respir. Rev.* 2010; 19(116):113-8. cited by applicant

Au, DH et al., The effects of smoking cessation on the risk of chronic obstructive pulmonary disease exacerbations. *J Gen Intern Med.* 2009;24(4):457-63. cited by applicant

Bafadhel, M et al., Predictors of exacerbation risk and response to budesonide in patients with chronic obstructive pulmonary disease : a post-hoc analysis of three randomised trials. *Lancet Respir Med* 2018 ; 6(2) : 117-26. cited by applicant

Barjaktarevic, I. et al., Ensifentrine, a Novel Dual Phosphodiesterase (PDE) 3 and 4 Inhibitor, Significantly Reduces Annualized Exacerbations and Delays the Time to First Exacerbation in COPD: Pooled Sub-group Analyses of ENHANCE-1 and ENHANCE-2 Phase 3 Trials (Poster at *ATS 2023—May 19-24, 2023*). cited by applicant

Barjaktarevic, I. et al., Ensifentrine, a novel dual phosphodiesterase (PDE) 3 and 4 inhibitor, significantly reduces annualized exacerbations and delays the time to first exacerbation in COPD: pooled sub-group analyses of enhance-1 and enhance-2 phase 3 trials, *Am J Respir Crit Care Med*, (2023):207:A5008. cited by applicant

Brat, K et al., Prognostic properties of the GOLD 2023 classification system. *Int J Chron Obstruct Pulmon Dis.* 2023; 18:661-667. cited by applicant

Breo USPI 2013. cited by applicant

Cazzola et al., “Ensifentrine (RPL554): an investigational PDE3/4 inhibitor for the treatment of

COPD,” Expert Opinion on Investigational Drugs, Sep. 1, 2019, 28(10):827-833. cited by applicant

Cazzola, M. et al., Ensifentrine (RPL554): and inhaled ‘bifunctional’ dual PDE3/4 inhibitor for the treatment of asthma and chronic obstructive pulmonary disease, *Pharmaceutical Patent Analyst*, vol. 7, 6(2018):249-257. cited by applicant

Daliresp (roflumilast). Package insert. AstraZeneca; 2020. cited by applicant

Daliresp USPI 2011. cited by applicant

Developing innovative therapies for the treatment of respiratory diseases, Nov. 2023 (presentation from Verona website). cited by applicant

Dose Ranging Study of RPL554 in Chronic Obstructive Pulmonary Disease (COPD) Patients, *ClinicalTrials.gov*, (2019). <https://www.clinicaltrials.gov/study/NCT03443414>

<https://www.clinicaltrials.gov/study/NCT03443414>. cited by applicant

Dransfield, MT et al., Acute exacerbations and lung function loss in smokers with and without chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2017;195(3):324-330. cited by applicant

Duffy, SP et al., Chronic obstructive pulmonary disease: evaluation and management. *Med Clin North Am*. 2019; 103(3):453-461. cited by applicant

ENHANCE-1 Phase 3 data—Dec. 2022 (presentation from Verona website). cited by applicant

ENHANCE-2 Phase 3 data—Aug. 2022 (presentation from Verona website). cited by applicant

Experimental report filed with response dated Dec. 13, 2021 to first office action on Chinese Application No. 202010090019.5, including translation of experimental report. cited by applicant

Experimental report filed with response dated Jan. 17, 2022 on Korean Patent Application No. 10-2017-7006862, including translation of experimental report. cited by applicant

Experimental report filed with response dated Jul. 4, 2022 to second office action on Chinese Application No. 202010090019.5, including translation of experimental report. cited by applicant

Ferguson, G.T. et al., A Dose-Ranging Study of the Novel Inhaled Dual PDE 3 and 4 Inhibitor Ensifentrine in Patients with COPD Receiving Maintenance Tiotropium Therapy, *International Journal of Chronic Obstructive Pulmonary Disease*, vol. 16, (2021): 1137-1148, with Supplement to Manuscript, 32 pages. cited by applicant

Ferguson, G.T. et al., A Dose-Ranging Study of the Novel Inhaled Dual PDE 3 and 4 Inhibitor Ensifentrine in Patients with COPD Receiving Maintenance Tiotropium Therapy, *International Journal of Chronic Obstructive Pulmonary Disease*, vol. 16, (2021):1137-1148. cited by applicant

Global Initiative for Chronic Obstructive Lung Disease, 2021 Report, 2021, 54 pages. cited by applicant

Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease 2022 report. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Accessed Jun. 13, 2024. https://goldcopd.org/wp-content/uploads/2021/12/GOLD-REPORT-2022-v1.1-22Nov2021_WMV.pdf. cited by applicant

Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease 2023 report. Global Initiative for Chronic Obstructive Lung Disease (GOLD). cited by applicant

Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease 2024 report. Global Initiative for Chronic Obstructive Lung Disease (GOLD). Accessed Apr. 16, 2024. <https://goldcopd.org/2024-gold-report/>. cited by applicant

Hankinson, J.L. et al., Spirometric reference values from a sample of the General U.S. Population, *Am J Respir Crit Care Med*, vol. 159, (1999):179-187. cited by applicant

Harries, TH et al., Blood eosinophil count, a marker of inhaled corticosteroid effectiveness in preventing COPD exacerbations in post-hoc RCT and observational studies: systematic review and meta-analysis. *Respir Res*. Jan. 3, 2020;21(1):3. cited by applicant

Hurst, JR et al., Prognostic risk factors for moderate-to-severe exacerbations in patients with chronic obstructive pulmonary disease: a systematic literature review. *Respir Res*. 2022;23(1):213.

cited by applicant

Iheanacho, I. et al., Economic burden of chronic obstructive pulmonary disease (COPD): a systematic literature review. *Int J Chron Obstruct Pulmon Dis.* 2020;15:439-460. cited by applicant

International Search Report on PCT/GB2023/052082 dated Oct. 20, 2023. cited by applicant

International Search Report on PCT/GB2023/052083 dated Oct. 31, 2023. cited by applicant

International Search Report on PCT/GB2023/052084 dated Oct. 23, 2023. cited by applicant

International Search Report on PCT/GB2023/052085 dated Nov. 15, 2023. cited by applicant

Kew, KM et al., Inhaled steroids and risk of pneumonia for chronic obstructive pulmonary disease. *Cochrane Database Syst. Rev.* 2014;2014(3):Cd010115. cited by applicant

Labiris, N.R. et al., Pulmonary drug delivery. Part II: The role of inhalant delivery devices and drug formulations in therapeutic effectiveness of aerosolized medications, *Br J Clin Pharmacol.*, vol. 56, (2003):600-612. cited by applicant

Le Brun, P.P.H. et al., A review of the technical aspects of drug nebulization, *Pharmacy World and Science*, vol. 22, 3(2000):75-81. cited by applicant

Lipson, DA et al., FULFIL trial: once-daily triple therapy for patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2017;196(4):438-446. cited by applicant

Lipson, DA et al., Once-daily single-inhaler triple versus dual therapy in patients with COPD. *N Engl J Med.* 2018; 378(18):1671-1680. cited by applicant

Mahler, DA et al., Effect of ensifentrine on dyspnea in patients with moderate-to-severe chronic obstructive pulmonary disease: pooled analysis of the ENHANCE trials. *Expert Rev Respir Med.* Aug. 2024;18(8):645-654. doi: 10.1080/17476348.2024.2389960. Epub Aug. 8, 2024. PMID: 39106052. cited by applicant

Mahler, DA et al., Ensifentrine, A Novel, Selective Inhibitor of PDE3 and PDE4, Improved Dyspnea in Subjects With Symptomatic, Moderate-to-Severe COPD Over 24 Weeks (Abstract and Poster), A27. Emerging Treatments and Therapeutic Strategies in Copd: Results of Clinical Trials and Observational Studies / Poster Discussion Session / Sunday May 19/09:15 AM / San Diego Convention Center, Room 33A-C. *American Journal of Respiratory and Critical Care Medicine* 2024;209:A1205. cited by applicant

Mannino, D et al., Treatment patterns for chronic obstructive pulmonary disease (COPD) in the United States: results from an observational cross-sectional physician and patient survey. *Int J Chron Obstruct Pulmon Dis.* 2022; 17:749-761. cited by applicant

Mantero, M. et al., Acute exacerbations of COPD: risk factors for failure and relapse, *International Journal of COPD*, vol. 12, (2017):2687-2693. cited by applicant

Martinez, FJ et al., Determinants of response to roflumilast in severe chronic obstructive pulmonary disease. Pooled analysis of two randomized trials. *Am J Respir Crit Care Med.* Nov. 15, 2018;198(10):1268-1278. cited by applicant

Matera, M.G. et al., Prospects for COPD treatment, *Current Opinion In Pharmacology*, vol. 56, (2021):74-84. cited by applicant

Maurer et al., "Late Breaking Abstract—RPL554, a first-in-class dual PDE3/4 inhibitor, causes significant bronchodilation and symptom relief; a Phase 2B COPD study," *European Respiratory Journal*, 2018, 52:OA1940, 1-5. cited by applicant

Milara, J et al., Oxidative stress-induced glucocorticoid resistance is prevented by dual PDE3/PDE4 inhibition in human alveolar macrophages. *Clin Exp Allergy.* 2011;41(4):535-46. cited by applicant

Miller, M.R. et al., Standardisation of spirometry, *Eur Respir J.*, vol. 26, (2005):319-338. cited by applicant

Miravittles, M et al., A pooled analysis of mortality in patients with COPD receiving dual bronchodilation with and without additional inhaled corticosteroid. *Int J Chron Obstruct Pulmon Dis.* 2022;17:545-558. cited by applicant

Negewo et al., "Peripheral blood eosinophils: a surrogate marker for airway eosinophilia in stable

COPD,” International Journal of COPD, 2016, 11:1495-1504. cited by applicant

Nici, L et al., Pharmacologic management of chronic obstructive pulmonary disease. an official American Thoracic Society clinical practice guideline. Am J Respir Crit Care Med. 2020;201(9):e56-e69. cited by applicant

Pascoe, S et al., Blood eosinophils and treatment response with triple and dual combination therapy in chronic obstructive pulmonary disease: analysis of the IMPACT trial. Lancet Respir Med 2019;7:745-756. cited by applicant

Pasquale, MK et al., Impact of exacerbations on health care cost and resource utilization in chronic obstructive pulmonary disease patients with chronic bronchitis from a predominantly Medicare population. Int J Chron Obstruct Pulmon Dis. 2012; 7:757-64. cited by applicant

Pavord, ID et al., Blood eosinophil count and pneumonia risk in patients with chronic obstructive pulmonary disease: a patient-level meta-analysis. Lancet Respir Med. 2016;4(9):731-741. cited by applicant

Preparing to launch the first novel MOA in COPD in 10 years, Oct. 18, 2023 (presentation from Verona website). cited by applicant

Rabe, KF et al., Anti-inflammatory effects of roflumilast in chronic obstructive pulmonary disease (ROBERT): a 16-week, randomised, placebo-controlled trial. Lancet Respir Med. Nov. 2018;6(11):827-836. doi: 10.1016/S2213-2600(18)30331-X. Epub Sep. 14, 2018. Erratum in: Lancet Respir Med. Oct. 12, 2018;: PMID: 30224319. cited by applicant

Rabe, KF et al., Triple inhaled therapy at two glucocorticoid doses in moderate-to-very-severe Copd. N Engl J Med. 2020; 383(1):35-48. cited by applicant

Record History | ver. 42: Jul. 11, 2022 | NCT04535986 | ClinicalTrials.gov. cited by applicant

Record History | ver. 43: May 13, 2022 | NCT04542057 | ClinicalTrials.gov. cited by applicant

Rheault, T. et al., Effect of ensifentrine on lung function and health-related quality of life (QoL) by COPD severity: analysis from a phase 2b dose ranging study, American Journal of Respiratory and Critical Care Medicine, (2021):203:A2254. cited by applicant

Roca, J. et al., References values for forced spirometry, Eur Respir J., vol. 11, (1998):1354-1362. cited by applicant

Rothnie, KJ et al., Natural history of chronic obstructive pulmonary disease exacerbations in a general practice-based population with chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 2018; 198(4):464-471. cited by applicant

Schmidt, DT et al., The effect of selective and non-selective phosphodiesterase inhibitors on allergen- and leukotriene C(4)-induced contractions in passively sensitized human airways. Br J Pharmacol. 2000;131(8):1607-18. cited by applicant

Sciurba, F.C. et al., Ensifentrine, a novel dual phosphodiesterase (PDE) 3 and 4 inhibitor, improves lung function, symptoms, quality of life and reduces exacerbation rate and risk in patients with COPD: results from replicate phase 3 trials, Am J Respir Crit Care Med, (2023):207, poster. cited by applicant

Sciurba, F.C. et al., Ensifentrine, a novel dual phosphodiesterase (PDE) 3 and 4 inhibitor, improves lung function, symptoms, quality of life and reduces exacerbation rate and risk in patients with COPD: results from replicate phase 3 trials, Am J Respir Crit Care Med, (2023):207:A5005. cited by applicant

Sciurba, FC et al., Dual Phosphodiesterase 3 and 4 Inhibitor Ensifentrine Reduces Exacerbation Rate and Risk in Patients With Moderate to Severe COPD. Chest. Aug. 27, 2024:S0012-3692(24)04937-7. doi: 10.1016/j.chest.2024.07.168. Epub ahead of print. PMID: 39197510. cited by applicant

Sciurba, FC et al., Ensifentrine, A Novel, Selective Inhibitor of PDE3 and PDE4, Reduced Moderate/Severe Exacerbation Rate and Risk in Subjects With COPD Regardless of Baseline Blood Eosinophils (Abstract and Poster), B52. Evidence for Therapeutic Strategies in Copd: From Established To Emerging/ Thematic Poster Session / Monday, May 20/09:15 AM-04:15 PM / San

Diego Convention Center, Area F. American Journal of Respiratory and Critical Care Medicine 2024;209:A3820. cited by applicant

Singh et al., “Blood Eosinophil Counts in Chronic Obstructive Pulmonary Disease: A Biomarker of Inhaled Corticosteroid Effects,” Tuberculosis and Respiratory Disease, 2020, 83:185-194. cited by applicant

Singh, D. et al., A dose-ranging study of the inhaled dual phosphodiesterase 3 and 4 inhibitor ensifentrine in COPD, Respiratory Research, vol. 21, (2020). cited by applicant

Singh, D. et al., Efficacy and safety of CHF6001, a novel inhaled PDE4 inhibitor in COPD: the PIONEER study, Respir Res., vol. 21, (2020):246. cited by applicant

Singh, D. et al., The short-term bronchodilator effects of the dual phosphodiesterase 3 and 4 inhibitor RPL554 in COPD, Eur Respir J., vol. 52, (2018):1801074. cited by applicant

Sonnex, K et al., Impact of smoking status on the efficacy of inhaled corticosteroids in chronic obstructive pulmonary disease: a systematic review. BMJ Open. 2020; 10(4):e037509. cited by applicant

Spiriva Respimat USPI (US product insert), 2014. cited by applicant

Suissa, S, Number needed to treat in COPD: exacerbations versus pneumonias. Thorax. 2013;68(6):540-3. cited by applicant

Suissa, Samy, Single-inhaler triple versus dual bronchodilator therapy for Gold E and other exacerbating patients with COPD: Real-world comparative effectiveness and safety, European Respiratory Journal 2023. cited by applicant

Symbicort USPI 2006. cited by applicant

Tashkin DP, et al., Concomitant inhaled corticosteroid use and the risk of pneumonia in COPD: a matched-subgroup post hoc analysis of the UPLIFT trial. Respir Res. 2018; 19(1):196. cited by applicant

Taylor, N.P., Verona sets sights on Philb after CPOD drug comes through early trial, Fierce Biotech article, (2015). cited by applicant

Tudorza USPI 2012. cited by applicant

Turner, MJ et al., The dual phosphodiesterase 3 and 4 inhibitor RPL554 stimulates CFTR and ciliary beating in primary cultures of bronchial epithelia. Am J Physiol Lung Cell Mol Physiol. 2015;310(1):L59-70. cited by applicant

Verona Pharma Announces Analyses Demonstrating Ensifentrine Reduced Exacerbation Rates Across Subgroups in Phase 3 ENHANCE-2 Trial for COPD, Oct. 14, 2022 (press release from Verona website). cited by applicant

Verona Pharma, Clinical Study Protocol, Version 5.0, Apr. 30, 2021 (Year: 2021). cited by applicant

Vogelmeier, CF et al., Evaluation of exacerbations and blood eosinophils in UK and US COPD populations. Respir Res. 2019;20(1): 178. cited by applicant

Watz, H et al., Symptom improvement following treatment with the inhaled dual phosphodiesterase 3 and 4 inhibitor ensifentrine in patients with moderate to severe COPD—A Detailed Analysis. Int J Chron Obstruct Pulmon Dis. 2020;15:2199-2206. cited by applicant

Zafari, Z. et al., Projecting long-term health and economic burden of COPD in the United States. Chest. 2021; 159(4):1400-1410. cited by applicant

Zafari, Z. et al., The projected economic and health burden of sub-optimal asthma control in Canada. Respiratory Medicine. 2018; 138:7-12. cited by applicant

Zuo, H. et al., Phosphodiesterases as therapeutic targets for respiratory diseases, Pharmacology & Therapeutics, vol. 197, (2019):225-242. cited by applicant

Background/Summary

CROSS REFERENCE (1) This application is a continuation of U.S. application Ser. No. 18/422,922, filed Jan. 25, 2024, which is a continuation of International Patent Application No. PCT/GB2023/050372, filed Feb. 20, 2023, which claims the benefit of GB Application No. 2202297.4, filed Feb. 21, 2022, each of which is incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

(1) The present invention relates to a process for producing a sterile formulation comprising an active pharmaceutical ingredient.

BACKGROUND OF THE INVENTION

(2) Certain pharmaceutical products must meet stringent sterility standards before they can be administered to patients. In view of the importance of sterility for pharmaceutical products, a large number of different processes, protocols and techniques have been developed for sterilisation of pharmaceutical products. These include steam sterilisation, dry heat sterilisation, ionising radiation sterilisation (including x-ray and gamma-ray sterilisation), gas sterilisation, sterile filtration and aseptic processing. The selection of a sterilisation protocol for a new pharmaceutical product is a complex process which relies on evaluation of the various available methods and their suitability for the specific product.

(3) Ensifentrine (N-(2-((2E)-9,10-dimethoxy-4-oxo-2-[(2,4,6-trimethylphenyl)imino]-6,7-dihydro-2H-pyrimido [6,1-a]isoquinolin-3(4H)-yl)ethyl) urea; also known as RPL554) is a dual PDE3/PDE4 inhibitor and is described in WO 00/58308 A1. Ensifentrine has both bronchodilatory and anti-inflammatory activity and is useful in the treatment of respiratory disorders such as asthma and chronic obstructive pulmonary disease (COPD). The structure of ensifentrine is shown below.

(4) ##STR00001##

(5) Ensifentrine may be formulated as a suspension of particles in a diluent as described in WO 2016/042313 A1.

(6) No suitable method for sterilising a liquid pharmaceutical composition comprising ensifentrine particles has been described. There is a need to develop a process for producing a sterilised ensifentrine suspension.

SUMMARY OF THE INVENTION

(7) The inventors have found that, of all the available sterilisation techniques, dry heat sterilization of particles of active agent followed by aseptic compounding is particularly well-suited to formulation of ensifentrine as a suspension. Other techniques such as terminal sterilisation or dry gamma-ray sterilisation were found to cause degradation of the formulations.

(8) The invention accordingly provides a process for producing a sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles, wherein the process comprises: (a) heating ensifentrine particles at a temperature of from 100° C. to 220° C. to obtain sterile ensifentrine particles; and (b) combining the sterile ensifentrine particles with a sterile liquid vehicle to produce the sterile liquid pharmaceutical composition suitable for administration by inhalation.

(9) The invention further provides a process for producing an ampule comprising a sterile liquid pharmaceutical composition suitable for administration by inhalation, the process comprising: (i) producing a sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles by the process defined herein; and (ii) filling an ampule with the sterile liquid pharmaceutical composition suitable for administration by inhalation.

Description

DETAILED DESCRIPTION OF THE INVENTION

- (1) The process of the invention produces a sterile liquid pharmaceutical composition.
- (2) The sterile liquid pharmaceutical composition is suitable for administration by inhalation. The sterile liquid pharmaceutical composition comprises sterile ensifentrine particles.
- (3) The term “sterile” refers to a composition which is essentially free of viable microbes. As such, a sterile composition is typically a composition which is substantially free of viable bacteria or fungi. The term “sterile” is well known in the art.
- (4) The sterile liquid pharmaceutical composition may have a sterility assurance level (SAL) of less than or equal to 10^{-3} , less than or equal to 10^{-6} , or less than or equal to 10^{-9} . For instance, an SAL of 10^{-6} means that the probability that a final product is non-sterile is $1/10^6$. The sterile liquid pharmaceutical composition may have a total bioburden limit of less than or equal to 10 CFU/mL, or less than or equal to 1 CFU/mL. “CFU” is a colony forming unit. The bioburden may be as measured using a plate count method as described in USP 31 <61>, for instance the pour-plate method.
- (5) Typically, the sterility of the liquid pharmaceutical composition is as determined in accordance with USP <71> or Ph Eur 2.6.1. The liquid pharmaceutical composition typically meets the acceptance criteria as defined in USP <71> or Ph Eur 2.6.1.
- (6) The pharmaceutical composition is a liquid pharmaceutical composition and, as such, is liquid under ambient conditions (for instance at temperatures from 10 to 40° C.).
- (7) The sterile liquid pharmaceutical composition comprises sterile ensifentrine particles. The ensifentrine particles comprise ensifentrine (i.e. ensifentrine free base) or a pharmaceutically acceptable salt thereof. Typically, the ensifentrine particles comprise ensifentrine free base. The ensifentrine particles typically comprise at least 90.0 wt % of ensifentrine or a pharmaceutically acceptable salt thereof, preferably at least 95.0 wt %. The ensifentrine particles may consist essentially of ensifentrine or a pharmaceutically acceptable salt thereof, or may consist of ensifentrine or a pharmaceutically acceptable salt thereof. For instance, the ensifentrine particles may consist of ensifentrine free base.
- (8) A composition which consists essentially of a component comprises only that component and other components which do not materially affect the essential characteristics of the component of which the composition essentially consists. A composition consisting essentially of a component typically comprises at least 99.5 wt % of that component relative to the total weight of the composition.
- (9) The sterile ensifentrine particles in the sterile liquid pharmaceutical composition may be suspended in the sterile liquid vehicle. The sterile liquid pharmaceutical composition accordingly typically comprises a suspension of the ensifentrine particles. It may also be the case that some or all of the ensifentrine particles in the sterile liquid pharmaceutical composition have settled to the bottom of a receptacle containing the sterile liquid pharmaceutical composition, for instance after storage for a period of time. The ensifentrine particles may be re-suspended in any suitable way, for instance by agitation of the sterile liquid pharmaceutical composition.
- (10) Step (a) of the process typically comprises heating the ensifentrine particles at a temperature of from 120° C. to 200° C. The process may comprise heating the ensifentrine particles at a temperature of from 140° C. to 180° C., for instance from 150° C. to 170° C. or from 155° C. to 165° C.
- (11) The ensifentrine particles may be heated for any period of time suitable for producing sterile ensifentrine particles. The process typically comprises heating the ensifentrine particles at the temperature for a time of from 10 minutes to 24 hours. The process may comprise heating the ensifentrine particles at the temperature for a time of from 30 minutes to 360 minutes, for instance from 45 minutes to 160 minutes or from 60 minutes to 140 minutes.
- (12) The process preferably comprises heating the ensifentrine particles at a temperature of from

145° C. to 175° C. for a time of from 45 minutes to 160 minutes. For instance, the process may comprise heating the ensifentrine particles at a temperature of from 155° C. to 165° C. for a time of from 110 minutes to 130 minutes.

(13) Heating the ensifentrine particles at a temperature in a certain range for a certain time means that the temperature of the ensifentrine particles is held at a temperature within that range for the total time. The ensifentrine particles are typically heated for the relevant time in a single heating step, but the heating may alternatively be conducted in two or more separate heating steps. For instance, heating at a temperature of from 140° C. to 180° C. for 60 minutes may comprising a single heating step for a duration of 60 minutes or two separate heating steps each having a duration of 30 minutes. Typically the ensifentrine particles are heated in a single heating step at the temperature for the time.

(14) Step (a) is typically a dry heat sterilisation process. The ensifentrine particles are typically in the form of a dry powder. The process accordingly may comprise heating a dry powder comprising ensifentrine particles at a temperature of from 100° C. to 220° C. to obtain a sterile dry powder comprising ensifentrine particles. Typically, the ensifentrine particles are subjected to dry-heat treatment on their own. The process may comprise heating a dry powder of ensifentrine particles (i.e. a dry powder consisting essentially of ensifentrine particles) at a temperature of from 100° C. to 220° C. to obtain a sterile dry powder of ensifentrine particles. A dry powder is typically a powder having a water content of less than 5.0 wt %, less than 1.0 wt % or less than 0.1 wt %.

(15) The ensifentrine particles typically have a particle size distribution with a Dv50 of from 0.5 µm to 5.0 µm. The ensifentrine particles preferably have a Dv50 of from 1.0 µm to 2.0 µm. Typically, the Dv10 of the ensifentrine particles is from 0.2 µm to 1.0 µm and the Dv90 of the ensifentrine particles is from 2.5 µm to 6.0 µm. For instance, the Dv10 of the ensifentrine particles may be from 0.4 µm to 0.6 µm and the Dv90 of the ensifentrine particles may be from 2.8 µm to 4.2 µm.

(16) Particle sizes are described herein by reference to the Dv50 value, which is the median particle size for a volume distribution. Thus, half the volume of the particles have diameters of less than the Dv50 value and half the volume of the particles have diameters of greater than the Dv50 value. This is a well-known manner in which to describe particle size distributions. The parameters of Dv10 and Dv90 may also be used to characterise a particle size distribution of a sample. 10% of the volume of particles have a diameter of less than the Dv10 value. 90% of the volume of the particles have a diameter of less than the Dv90 value.

(17) The technique used to measure the Dv50 (and Dv10 and Dv90) values as stated herein is typically laser diffraction. The particle size distribution of the ensifentrine particles may be as measured by laser diffraction using a wet powder dispersion system. For instance, the particle size distribution can be measured by laser diffraction using a Malvern Spraytec in conjunction with a wet dispersion cell. Typically, the instrument parameters for the Malvern Spraytec are as follows: particle—standard opaque particle; refractive index Particle—1.50; refractive index (imaginary)—0.50; density of particle—1.00; refractive index of dispersant—1.33; controller unit—1000 RPM; measurement type—timed; initial sampling time—30 s; obscuration—20%-30%; dispersant—1% Polysorbate 20 in deionised water.

(18) The ensifentrine particles may be produced by any pharmaceutically acceptable size reduction process or particle size controlled production process. For instance, the particles may be produced by spray-drying a solution of ensifentrine, by controlled crystallisation, or by size reduction of a solid form of ensifentrine, for example by air jet milling, mechanical micronisation or media milling.

(19) The process further comprises combining the sterile ensifentrine particles with a sterile liquid vehicle to produce the sterile liquid pharmaceutical composition suitable for administration by inhalation. Combining the sterile ensifentrine particles with a sterile liquid vehicle typically comprises mixing the sterile ensifentrine particles with a sterile liquid vehicle.

(20) The sterile liquid vehicle typically comprises one or more diluents. The diluent may be any suitable liquid diluent. For instance, the sterile liquid vehicle typically comprises a diluent which is water. The sterile liquid vehicle may comprise one or more additional diluents or the sterile liquid vehicle may comprise a single diluent. The sterile liquid vehicle may comprise a single diluent which is water.

(21) A sterile liquid vehicle comprising water may be used to produce a sterile liquid pharmaceutical composition suitable for use in a nebulizer. The sterile liquid vehicle typically further comprises one or more additional excipients selected from surfactants, buffers and tonicity adjusters.

(22) The sterile liquid vehicle typically further comprises a tonicity adjuster. Examples of tonicity adjusters include sodium chloride, potassium chloride, glucose, glycerine and mannitol. Preferably, the tonicity adjuster is sodium chloride.

(23) The concentration of the tonicity adjuster in the sterile liquid vehicle is typically greater than or equal to 1.0 mg/mL (for instance from 1.0 to 50.0 mg/mL). Preferably, the tonicity adjuster concentration is from 4.0 to 20.0 mg/ml or from 6.0 to 12.0 mg/mL.

(24) The sterile liquid vehicle typically further comprises one or more surfactants. The one or more surfactant may comprise a non-ionic surfactant, an anionic surfactant, a cationic surfactant, a zwitterionic surfactant or a mixture thereof. Typically, the one or more surfactants comprise a non-ionic surfactant.

(25) Examples of surfactants include lecithin, oleic acid, polyoxyethylene glycol alkyl ethers (for instance PEG 300, PEG 600, PEG 1000, Brij 30, Brij 35, Brij 56, Brij 76 and Brij 97), polypropylene glycol (for instance PPG 2000), glucoside alkyl ethers, polyoxyethylene glycol octylphenol ethers, polyoxyethylene glycol alkylphenol ethers, glycerol alkyl esters, polyoxyethylene glycol sorbitan alkyl esters (polysorbates, for instance polysorbate 20, polysorbate 40, polysorbate 60 and polysorbate 80), sorbitan alkyl esters (for instance sorbitan monolaurate (Span® 20), sorbitan monooleate (Span® 80) and sorbitan trioleate (Span® 85)), cocamide MEA, cocamide DEA, dodecyldimethylamine oxide, block copolymers of polyethylene glycol and polypropylene glycol (poloxamers), block copolymers of polyethylene glycol and polypropylene oxide (for instance Pluronic surfactants), polyvinyl pyrrolidone K25, polyvinyl alcohol, oligolactic acid, sodium dioctyl sulfosuccinate and polyethoxylated tallow amine (POEA).

(26) Preferably, the one or more surfactants comprise a polysorbate and/or a sorbitan alkyl ester. The one or more surfactants may for instance comprise polysorbate 20 (polyoxyethylene (20) sorbitan monolaurate), polysorbate 40 (polyoxyethylene (20) sorbitan monopalmitate), polysorbate 60 (polyoxyethylene (20) sorbitan monostearate) or polysorbate 80 (polyoxyethylene (20) sorbitan monooleate). The one or more surfactants may for instance comprise sorbitan monolaurate (Span® 20), sorbitan monooleate (Span® 80) or sorbitan trioleate (Span® 85). Preferably, the sterile liquid vehicle comprises polysorbate 20 and/or sorbitan monolaurate (Span® 20).

(27) The sterile liquid vehicle may comprise two or more surfactants, or the sterile liquid vehicle may comprise a single surfactant. For instance the composition may comprise a single surfactant which is a polysorbate, for instance polysorbate 20. The sterile liquid vehicle may comprise two or more surfactants. For instance, the sterile liquid vehicle may comprise polysorbate 20 and sorbitan monolaurate (Span® 20).

(28) The total concentration of the one or more surfactants in the sterile liquid vehicle is typically from 0.01 to 2.0 mg/mL. Preferably, the total surfactant concentration is from 0.1 to 1.0 mg/mL. For instance, the total surfactant concentration in the sterile liquid vehicle may be from 0.25 to 0.75 mg/mL. The sterile liquid vehicle may for instance comprise a polysorbate at a concentration of from 0.1 to 1.0 mg/mL and optionally a sorbitan alkyl ester at a concentration of from 0.01 to 0.1 mg/mL.

(29) The sterile liquid vehicle typically further comprises one or more buffers. The buffer can be used to control the pH of the liquid pharmaceutical composition. A buffer typically comprises a

weak acid and its conjugate base. Examples of buffers include a citrate buffer, a phosphate buffer, an acetate buffer, and a bicarbonate buffer.

(30) Preferably, the buffer is a phosphate buffer. For instance, the sterile liquid vehicle may comprise sodium dihydrogen phosphate dihydrate and/or disodium phosphate dihydrate.

(31) The pH of the sterile liquid vehicle is typically from 6.0 to 7.5, for instance from 6.2 to 7.2. The pH of the sterile liquid vehicle may be from 6.5 to 6.9. The pH of the sterile liquid vehicle is typically the pH as measured at a temperature of 20° C. The pH of the sterile liquid vehicle may be measured by any suitable technique. For instance, the pH may be as measured using a potentiometric pH meter.

(32) The total concentration of the one or more buffers in the sterile liquid vehicle is typically from 0.1 to 20.0 mg/mL. Preferably the buffer concentration is from 1.0 to 2.0 mg/mL. The concentration of the buffer includes both the acid and conjugate base components of the buffer.

(33) Typically, the sterile liquid vehicle comprises: water, one or more tonicity adjusters, one or more buffers, and one or more surfactants. The sterile liquid vehicle may for instance comprise: water, sodium chloride, sodium dihydrogen phosphate dihydrate, disodium phosphate dihydrate, polysorbate 20 and sorbitan laurate.

(34) The process typically further comprises producing the sterile liquid vehicle. The sterile liquid vehicle may be produced by sterilising a liquid vehicle, for instance by aseptic filtration, heat treatment or gamma-ray treatment of a liquid vehicle. The sterile liquid vehicle is typically produced by aseptic filtration of a liquid vehicle. Aseptic filtration typically comprises filtering a liquid vehicle through a filter having a nominal pore size of no greater than 0.5 µm, or no greater than 0.3 µm (for instance around 0.2 µm).

(35) The process of the invention produces a sterile liquid pharmaceutical composition through aseptic processing and compounding. As such, the process typically does not further comprise an additional step of terminal sterilisation (which has been found to cause the presence of impurities in the composition). For instance, the process typically does not further comprise heating the sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles at a temperature of 100° C. or greater.

(36) The concentration of ensifentrine particles in the sterile liquid pharmaceutical composition produced by the process can be any suitable concentration, for instance from 0.01 to 400 mg/mL. Typically, the concentration of ensifentrine particles is from 0.1 to 5.0 mg/mL. Preferably, the concentration of ensifentrine particles is from 0.1 to 2.5 mg/mL. For instance, the concentration of ensifentrine particles may be from 0.15 to 0.5 mg/mL or from 1.0 to 2.0 mg/mL.

(37) For instance, the sterile liquid pharmaceutical composition may comprise: water; particles consisting of ensifentrine free base at a concentration of from 0.1 to 20 mg/mL; one or more tonicity adjusters at a total concentration of from 1.0 to 15 mg/mL; one or more buffers at a total concentration of from 0.1 to 4 mg/ml; and one or more surfactants at a total concentration of from 0.05 to 3 mg/mL.

(38) The sterile liquid pharmaceutical composition may comprise: water; particles consisting of ensifentrine free base at a concentration of from 0.5 to 6 mg/mL; sodium chloride at a concentration of from 5 to 12 mg/ml; sodium dihydrogen phosphate dihydrate at a concentration of from 0.3 to 2 mg/mL; disodium phosphate dihydrate at a concentration of from 0.3 to 2 mg/mL; polysorbate 20 at a concentration of from 0.1 to 1.5 mg/ml; and sorbitan laurate at a concentration of from 0.01 to 0.5 mg/mL.

(39) The invention also provides a process for producing an ampule comprising a sterile liquid pharmaceutical composition suitable for administration by inhalation. The process comprises: (i) producing a sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles by a process as defined herein; and (ii) filling an ampule with the sterile liquid pharmaceutical composition suitable for administration by inhalation. The ampule may be a glass ampule or a plastic ampule. The ampule is typically a polyethylene plastic ampule.

The ampule may be individually overwrapped in an aluminium foil pouch after filling.

(40) The ampule may be a blow-fill-seal ampule. For instance, the process may be a process for producing an ampule using blow-fill-seal technology, which ampule comprises a sterile liquid pharmaceutical composition suitable for administration by inhalation, the process comprising: (i) producing a sterile liquid pharmaceutical composition suitable for administration by inhalation comprising ensifentrine particles by the process defined herein; and (ii) producing a blow-fill-seal ampule filled with the sterile liquid pharmaceutical composition suitable for administration by inhalation.

EXAMPLES

Methods

(41) Assay testing was performed in duplicate with the key high performance liquid chromatography (HPLC) parameters shown in Table 1.

(42) TABLE-US-00001 TABLE 1 Assay parameters Mobile phase Acetonitrile:water:TFA 45:55:0.1 Column Waters X-Bridge phenyl, 3.5 μ m, 150 $\times$ 4.6 mm Column temperature 40° C. Flow rate 1.5 mL/min Injection volume 10 μ L Detection UV@ 254 nm Runtime 6 minutes Autosampler Ambient temperature Sample and standard 0.1 mg/mL concentration

(43) Impurities determination was performed in duplicate with the key high performance liquid chromatography (HPLC) parameters shown in Table 2.

(44) TABLE-US-00002 TABLE 2 Impurities determination parameters Mobile phase A - Purified water/Acetonitrile/TFA (95/5/0.1) B - Acetonitrile/Water/TFA (95/5/0.1) Column Waters X-Bridge phenyl, 3.5 μ m, 150 $\times$ 4.6 mm Column temperature 30° C. Autosampler Ambient temperature Flow rate 1.0 mL/min Injection volume 10 μ L Detection UV@ 254 nm Gradient Time % A % B 0 100 0 2 100 0 15 0 100 25 0 100 27 100 0 37 100 0 Diluent Acetonitrile:water (50:50) Sample concentration 0.2 mg/mL Standard concentration 2 μ g/mL

Products Tested

(45) The effect of different sterilisation techniques were assessed on: (i) suspension formulations comprising ensifentrine particles (terminal sterilisation); and (ii) ensifentrine particles prior to formulation as a suspension (aseptic compounding and filling).

(46) Three aqueous suspensions having the formulations shown in Tables 3 to 5 were prepared.

(47) TABLE-US-00003 TABLE 3 Variant 1a (0.4 mg/ml, pH 6.7) Constituent Concentration (mg/ml) Micronised ensifentrine 0.4 Polysorbate 20 0.5 Span[®] 20 (sorbitan monolaureate) 0.05 Monosodium phosphate anhydrous 5.72 Dibasic sodium phosphate anhydrous 6.80 Sodium chloride 4.80 Water N/A

(48) TABLE-US-00004 TABLE 4 Variant 2a (0.4 mg/ml, pH 5.9) Constituent Concentration (mg/ml) Micronised ensifentrine 0.4 Polysorbate 80 0.5 Citric acid monohydrate 1.68 Trisodium citrate dihydrate 27.06 Sodium chloride 9.0 Water N/A

(49) TABLE-US-00005 TABLE 5 Variant 3a (0.4 mg/mL, pH 7.1) Constituent Concentration (mg/ml) Micronised ensifentrine 0.4 Polysorbate 20 0.5 Sodium chloride 9.0 NaOH N/A Water N/A

(50) The unformulated ensifentrine particles were assessed as a dry powder.

Sterilisation Techniques

(51) The sterilisation techniques shown in Table 6 were assessed.

(52) TABLE-US-00006 TABLE 6 sterilisation techniques Technique Method A Terminal sterilisation of the aqueous suspension formulations by heat treatment: each of variants 1a, 2a and 3a of the suspension formulation were heat treated in an autoclave at 121° C. for 15 minutes. B Terminal sterilisation of the aqueous suspension formulations by gamma irradiation: each of variants 1a, 2a and 3a of the suspension formulation were gamma irradiated at 25kGy-30kGy. C Dry heat sterilisation of micronised ensifentrine: a dry powder of the micronised ensifentrine was heat treated in an open glass vial in an oven at 150° C. for 150 minutes. D Dry heat sterilisation of micronised ensifentrine: a dry powder of the micronised ensifentrine was heat treated in an open

glass vial in an oven at 160° C. for 120 minutes. E Dry heat sterilisation of micronised ensifentrine: a dry powder of the micronised ensifentrine was heat treated in an open glass vial in an oven at 170° C. for 60 minutes. F Dry gamma irradiation of micronised ensifentrine: a dry powder of the micronised ensifentrine was gamma irradiated at 25kGy-30kGy.

(53) The sterilised micronised ensifentrine obtained from techniques C to F may be subsequently combined with an aseptic vehicle to obtain a suspension formulation.

Results

(54) The assays and related substances of the products obtained by each of sterilisation techniques A to F were assessed by the HPLC methods described above. The results are summarised in Table 7.

(55) TABLE-US-00007 TABLE 7 assessment of products Technique Assessment A: terminal heat No change in appearance of the suspension formulations was treatment observed. Significant impurities were however observed for variants 1a and 2a (RTT 0.79, 2.5% nominal) and variant 3a (RRT 0.87, 5.8% nominal). The assay results were correspondingly lower compared to untreated samples, with variant 1a dropping by 5%, variant 2a by 2.5% and variant 3a by approximately 8%. B: terminal gamma Upon irradiation, a brown discolouration of the glass was apparent. irradiation Variant 3a was found to have converted to a pale yellow solution. The related substances analysis showed clear degradation to a large number of impurities totalling nearly 7% nominal for variant 2a and 2.6% nominal for variant 3a. Variant 1a showed no change in total related substances when compared to the untreated sample, but there was a drop in assay suggesting product loss or degradation that was not visible in the chromatogram. A large drop in assay was observed for all variants. C: dry heat No significant changes observed. treatment (150° C. for 150 minutes) D: dry heat No significant changes observed. treatment (160° C. for 120 minutes) E: dry heat No significant changes observed. treatment: (170° C. for 60 minutes) F: dry gamma An increase in total impurities of 0.5% w/w was observed with a irradiation decrease in assay of around 1 to 1.5% w/w.

Conclusion

(56) The results in Table 7 indicated that terminal heat treatment (technique A), terminal gamma irradiation (technique B) and dry gamma irradiation (technique F) caused some degradation when used for preparation of a sterilised suspension of ensifentrine particles.

(57) It was found, however, that ensifentrine particles were highly resistant to sterilisation by dry heat treatment, with no significant changes in purity or assay following treatment at temperatures of from 150° C. to 170° C. Dry heat treatment at 160° C. for at least 120 minutes is preferred. The sterilised ensifentrine particles obtained by dry heat treatment can be combined with the suspension vehicle in aseptic compounding and processing to obtain a sterile suspension formulation comprising ensifentrine.

Claims

1. A sterile liquid pharmaceutical composition comprising sterile ensifentrine particles suitable for administration by inhalation, wherein the composition is produced by a method comprising: (a) dry heating ensifentrine particles as a dry powder at a temperature of from 145° C. to 175° C. for 45 minutes to 160 minutes to obtain the sterile ensifentrine particles, and (b) combining the sterile ensifentrine particles with a sterile liquid vehicle to produce the sterile liquid pharmaceutical composition suitable for administration by inhalation; wherein the dry heating produces less than a 0.5% w/w increase in total impurities.
2. The composition of claim 1, wherein the dry heating is performed at a temperature of 155° C. to 165° C. for 110 minutes to 130 minutes.
3. The composition of claim 1, wherein the sterile ensifentrine particles comprise at least 95% by weight of ensifentrine or a pharmaceutically acceptable salt thereof relative to a total weight of the

sterile ensifentrine particles.

4. The composition of claim 1, wherein the sterile ensifentrine particles comprise at least 95% by weight of ensifentrine free base relative to a total weight of the sterile ensifentrine particles.
 5. The composition of claim 1, wherein the sterile ensifentrine particles have a particle size distribution with a Dv10 of from 0.1 μm to 1.0 μm .
 6. The composition of claim 1, wherein the sterile ensifentrine particles have a particle size distribution with a Dv50 of from 0.5 μm to 5.0 μm .
 7. The composition of claim 1, wherein the sterile ensifentrine particles have a particle size distribution with a Dv90 of from 2.5 μm to 6 μm .
 8. The composition of claim 1, wherein the method further comprises subjecting a liquid vehicle to aseptic filtration to produce the sterile liquid vehicle.
 9. The composition of claim 1, wherein the sterile liquid vehicle comprises a diluent and optionally one or more additional excipients selected from surfactants, buffers, and tonicity adjusters.
 10. The composition of claim 9 comprising (a) sterile ensifentrine particles at a concentration of from 0.5 mg/mL to 6 mg/mL; (b) one or more surfactants at a concentration of from 0.01 mg/mL to 2 mg/mL; (c) a buffer at a concentration of from 0.6 mg/mL to 4 mg/mL; (d) a tonicity adjuster at a concentration of from 5 mg/mL to 12 mg/mL.
 11. The composition of claim 10, wherein the sterile liquid vehicle comprises water, sodium chloride, sodium dihydrogen phosphate dihydrate, disodium hydrogen phosphate dihydrate, polysorbate 20, or sorbitan laurate.
 12. The composition of claim 11, comprising (a) sterile ensifentrine particles at a concentration of from 0.5 mg/mL to 6 mg/mL; (b) polysorbate 20 at a concentration of from 0.1 mg/mL to 1.5 mg/mL; (c) sorbitan monolaurate at a concentration of from 0.01 mg/mL to 0.5 mg/mL; (d) sodium dihydrogen phosphate dihydrate at a concentration of from 0.3 mg/mL to 2 mg/mL; (e) disodium hydrogen phosphate dihydrate at a concentration of from 0.3 mg/mL to 2 mg/mL; and (f) sodium chloride at a concentration of from 5 mg/mL to 12 mg/mL, wherein a total weight of ensifentrine in the sterile liquid pharmaceutical composition is from 2.7 to 3.3 mg.
 13. The composition of claim 1, wherein the sterile liquid pharmaceutical composition comprises the sterile ensifentrine particles at a concentration of from 0.1 mg/mL to 5 mg/mL.
 14. The composition of claim 13, wherein the sterile liquid pharmaceutical composition comprises the sterile ensifentrine particles at a concentration of from 1 mg/mL to 2 mg/mL.
-