



(12) **United States Patent**
Higuchi et al.

(10) **Patent No.: US 12,410,483 B2**
(45) **Date of Patent: *Sep. 9, 2025**

(54) **METHODS OF DETECTING BLADDER
CANCER**

(71) Applicant: **Cepheid**, Sunnyvale, CA (US)

(72) Inventors: **Russell Higuchi**, Alameda, CA (US);
Stacey Ellen Wallace, Sunnyvale, CA
(US); **Edwin Wei-Lung Lai**, Menlo
Park, CA (US)

(73) Assignee: **Cepheid**, Sunnyvale, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **18/485,615**

(22) Filed: **Oct. 12, 2023**

(65) **Prior Publication Data**

US 2024/0175089 A1 May 30, 2024

Related U.S. Application Data

(60) Continuation of application No. 16/272,978, filed on
Feb. 11, 2019, now abandoned, which is a division of
application No. 14/394,352, filed as application No.
PCT/US2013/037334 on Apr. 19, 2013, now Pat. No.
10,202,653.

(60) Provisional application No. 61/770,803, filed on Feb.
28, 2013, provisional application No. 61/636,194,
filed on Apr. 20, 2012.

(51) **Int. Cl.**
C12Q 1/6886 (2018.01)

(52) **U.S. Cl.**
CPC **C12Q 1/6886** (2013.01); **C12Q 2600/158**
(2013.01); **C12Q 2600/16** (2013.01); **C12Q**
2600/166 (2013.01)

(58) **Field of Classification Search**
None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,155,018 A 10/1992 Gillespie et al.
5,958,349 A 9/1999 Petersen et al.
6,403,037 B1 6/2002 Chang et al.
6,440,725 B1 8/2002 Pourahmadi et al.
6,783,736 B1 8/2004 Taylor et al.
6,818,185 B1 11/2004 Petersen et al.
7,351,538 B2 4/2008 Fuchs et al.
7,402,422 B2 7/2008 Fuchs et al.
8,114,588 B2 2/2012 Yoshiki et al.
2004/0076955 A1 4/2004 Mack
2009/0282496 A1 11/2009 Chang
2010/0086932 A1 4/2010 Asensio et al.

FOREIGN PATENT DOCUMENTS

EP 2138848 A1 * 12/2009 C12Q 1/6886
JP 20085322477 A 8/2008
WO 2011079191 A1 6/2011
WO 2012067899 A2 5/2012
WO 2014018926 A1 1/2014
WO 2014118334 A1 8/2014
WO 2014138396 A1 9/2014

OTHER PUBLICATIONS

Stevenson et. al. 2008 The use of Armored RNA as a multi-purpose
internal control for RT-PCR. Journal of Virological Methods
150 :73-76 (Year: 2008).*

Kalendar et. al. 2009. FastPCR software for PCR primer and probe
design and repeat search. Genomes and Genomics 3 (Special Issue
1), 1-14 (Year: 2009).*

NCBI Reference Sequence: NM_000756.2, 2011, *Homo sapiens*
corticotropin releasing hormone (CRH), mRNA (Year: 2011).*

NCBI Reference Sequence: NM_001127598.1. 2011. *Homo sapiens*
insulin-like growth factor 2 (somatomedin A) (IGF2), transcript
variant 3, mRNA (Year: 2011).*

GenBank: BC031559.1. 2006. *Homo sapiens* keratin 20, mRNA
(cDNA clone MGC:35423 IMAGE:5189289), complete cds (Year:
2006).*

GenBank: BC007320.2. 2006. *Homo sapiens* annexin A10, mRNA
(cDNA clone MGC:1303 IMAGE:2988009 (Year: 2006).*

Mengual et. al. 2010. Gene Expression Signature in Urine for
Diagnosing and Assessing Aggressiveness of Bladder Urothelial
Carcinoma. Clin Cancer Res; 16(9); 2624-3 (Year: 2010).*

Holyoake et al. Development of a Multiplex RNAUrine Test for the
Detection and Stratification of Transitional Cell Carcinoma of the
Bladder. 2008 Clin Cancer Res 2008; 14(3), 742-749 (Year: 2008).*

Aguino de Muro, "Probe Design, Production, and Applications",
Medical Biomethods Handbook, Humana Press, pp. 13-23 (2005).

Blalock et al. "Gene Expression Analysis of Urine Sediment:
Evaluation for Potential Noninvasive Markers of Interstitial Cystitis/
Bladder Pain Syndrome," The Journal of Urology, 187:725-732
(2012).

Brabender et al. "The molecular signature of normal squamous
esophageal epithelium identifies the presence of a field effect and
can discriminate between patients with Barrett's esophagus and
patients with Barrett's-associated adenocarcinoma," Cancer Epidemiol
Biomarkers Prev. 14:2113-2117 (2005).

Buchumensky et al. "Cytokeratin 20: a new marker for early
detection of bladder cell carcinoma" J Urol. 160(6 Pt 1):1971-4
(1998).

Chen et al. "Identification of potential bladder cancer markers in
urine by abundant-protein depletion coupled with quantitative
proteomics", Journal of Proteomics, 85, pp. 28-43 (2013).

(Continued)

Primary Examiner — Jennifer Dunston

Assistant Examiner — Tiffany Nicole Grooms

(74) *Attorney, Agent, or Firm* — McAndrews, Held &
Malloy, Ltd.

(57) **ABSTRACT**

Compositions and methods for detecting bladder cancer are
provided. In some embodiments, methods of monitoring
recurrence of bladder cancer are provided. In some embodi-
ments, the methods comprise detecting a set of markers
consisting of CRH, IGF2, KRT20, and ANXA10.

21 Claims, No Drawings

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

- Christoph et al. "Urinary cytokeratin 20 mRNA expression has the potential to predict recurrence in superficial transitional cell carcinoma of the bladder" *Cancer Lett.* 245(1-2):121-6. 2007 (Epub 2006).
- Damrauer et al. "Intrinsic subtypes of high-grade bladder cancer reflect the hallmarks of breast cancer biology," *PNAS.* 111:3110-3115 (2014).
- DeGraff et al. "Loss of the urothelial differentiation marker FOXA1 is associated with high grade, late stage bladder cancer and increased tumor proliferation," *PLoS One.* 7(5):e36669 (2012).
- Eissa et al. "Comparison of CD44 and cytokeratin 20 mRNA in voided urine samples as diagnostic tools for bladder cancer," *Clin Biochem.* 41(16-17):1335-41 (2008).
- Eissa et al. "Comparison of cytokeratin 20 RNA and angiogenin in voided urine samples as diagnostic tools for bladder carcinoma," *Clin Biochem.* 37(9):803-10 (2004).
- Eissa et al. "The clinical relevance of urine-based markers for diagnosis of bladder cancer," *Med Oncol.* 28(2):513-8 (2011).
- Elsamman et al. "Differences in gene expression between noninvasive and invasive transitional cell carcinoma of the human bladder using complementary deoxyribonucleic acid microarray: preliminary results," *Urol Oncol.* 24(2):109-15 (2006).
- European Patent Office, "Communication pursuant to Rule 114(2) EPC", Third Party Observations to EP2839028 A1, issued in European Patent Application No. 13718488.3, mailed Sep. 16, 2016, 10 pages.
- Gallagher et al. "Recurrence of urothelial carcinoma of the bladder: a role for insulin-like growth factor-II loss of imprinting and cytoplasmic E-cadherin immunolocalization," *Clin Cancer Res.* 14(21):6829-38 (2008).
- GenBank Accession No. NM_000044, 11 pages. 2014.
- GenBank Accession No. NM_000756, 4 pages. 2014.
- GenBank Accession No. NM_001011645, 7 pages. 2014.
- GenBank Accession No. NM_006952, 4 pages. 2014.
- GenBank Accession No. NM_007193, 4 pages. 2014.
- GenBank Accession No. NM_007313, 7 pages. 2014.
- GenBank Accession No. NM_019010, 5 pages. 2014.
- GenBank: BC007320.2. 2006. *Homo sapiens* annexin A10, mRNA (cDNA clone MGC:1303 IMAGE:2988009), complete cds.
- GenBank: BC031559.1. 2006. *Homo sapiens* keratin 20, mRNA (cDNA clone MGC:35423 IMAGE:5189289), complete cds.
- Ginzinger, D. "Gene quantification using real-time quantitative PCR: An emerging technology hits the mainstream," *Experimental Hematology*, 30:503-512 (2002).
- Guo et al. "Quantitative detection of cytokeratin 20 mRNA in urine samples as diagnostic tools for bladder cancer by real-time PCR," *Exp Oncol.* 31(1):43-7 (2009).
- Holyoake et al. "Development of a Multiplex RNA Urine Test for the Detection and Stratification of Transitional Cell Carcinoma of the Bladder," *Clinical Cancer Research*, vol. 14, No. 1, pp. 742-749 (2008).
- International Search Report and Written Opinion of the International Searching Authority for PCT/US2013/037334, mailed Jun. 19, 2013, 13 pages.
- International Search Report and Written Opinion of the International Searching Authority PCT/US2014/021199, mailed Jun. 10, 2014, 20 pages.
- Ishii et al. Bladder cancer discovered by ovarian metastasis: cytokeratin expression is useful when making differential diagnosis. *Int J Urol.* 12(1):104-7 (2005).
- Izquierdo et al. "Molecular characterization of upper urinary tract tumours," *BJU Int.* 106(6):868-72. 2010 (Epub 2009).
- Jiang et al. "Cytokeratin 7 and cytokeratin 20 in primary urinary bladder carcinoma and matched lymph node metastasis," *Arch Pathol Lab Med.* 125(7):921-3 (2001).
- Kalendar et al. "FastPCR Software for PCR Primer and Probe Design and Repeat Search," *Genes, Genomes and Genomics* 3 (Special Issue 1), 1-14) and NM000756.2 (NCBI Reference Sequence: NM_000756.2, 2011", 14 pages (2009).
- Kawahito et al. "Corticotropin releasing hormone in colonic mucosa in patients with ulcerative colitis. *Gut.*" 37(4):544-51 (1995).
- Kim et al. "Decreased expression of annexin A10 in gastric cancer and its overexpression in tumor cell growth suppression," *Oncol Rep.* 24(3):607-12 (2010).
- Kim et al. "Reduced expression and homozygous deletion of annexin A10 in gastric carcinoma. *Int J Cancer.*" 125(8):1842-50 (2009).
- Langner et al. "Prognostic value of keratin subtyping in transitional cell carcinoma of the upper urinary tract," *Virchows Arch.* 445(5):442-8 (2004).
- Lassmann et al. "Identification of occult tumor cells in node negative lymph nodes of colorectal cancer patients by cytokeratin 20 gene and protein expression," *Int J Colorectal Dis.* 19(2):87-94. 2004 (Epub 2003).
- Lee et al. "Seven-Color, Homogeneous Detection of Six PCR Products," *BioTechniques.* 27:342-349 (1999).
- Liu S. et al. "Down-regulation of annexin A10 in hepatocellular carcinoma is associated with vascular invasion, early recurrence, and poor prognosis in synergy with p53 mutation," *Am J Pathol.* 160(5):1831-7 (2002).
- Liu, D. "Choice of endogenous control for gene expression in nonsmall cell lung cancer," *European Respiratory Journal.* 26:1002-1008 (2005).
- Livingstone, "IGF2 and cancer", *Endocrine Related Cancer*, vol. 20, No. 6, pp. R321-R339 (2013).
- Lobban et al. "Uroplakin Gene Expression by Normal and Neoplastic Human Urothelium," *Am J Pathol.* 153:1957-1967 (1998).
- Lopez-Beltran et al. "Invasive micropapillary urothelial carcinoma of the bladder," *Hum Pathol.* 41(8):1159-64 (2010).
- Lu et al. "MicroRNA expression profiles classify human cancers," *Nature.* 435:834-838 (2005).
- Lu S. et al. "Expression and prognostic significance of gastric-specific annexin A10 in diffuse- and intestinal-type gastric carcinoma," *J Gastroenterol Hepatol.* 26(1):90-7 (2011).
- Maniatis et al. "Amplification of cDNA Generated by Reverse Transcription of mRNA", *Molecular Cloning—A Laboratory Manual*, 2nd Edition, (1998).
- Marin-Aguilera et al. "Utility of urothelial mRNA markers in blood for staging and monitoring bladder cancer. *Urology.*" 79(1):240.e9-15. 2012 (Epub 2011).
- Markou et al. "Molecular Characterization of Circulating Tumor Cells in Breast Cancer by a Liquid Bead Array Hybridization Assay," *Clinical Chemistry*, vol. 57:3, pp. 421-430 (2011).
- McNeill et al. "Evaluation and validation of candidate endogenous control genes for real-time quantitative PCR studies of breast cancer", *BMC Molecular Biology*, vol. 8, No. 107, 13 pages (2007).
- Mengual et al. "DNA Microarray Expression Profiling of Bladder Cancer Allows Identification of Noninvasive Diagnostic Markers," *The Journal of Urology*, 182:741-748 (2009).
- Mengual et al. "Gene expression signature in urine for diagnosing and assessing aggressiveness of bladder urothelial carcinoma," *Clin Cancer Res.* 16(9):2624-33 (2010).
- Mengual et al. "Partially degraded RNA from bladder washing is a suitable sample for studying gene expression profiles in bladder cancer," *Eur Urol.* 50(6):1347-55; Editorial Comment 1355-6 (2006).
- Mengual et al. "Validation Study of a Noninvasive Urine Test for Diagnosis and Prognosis Assessment of Bladder Cancer: Evidence for Improved Models", *The Journal of Urology*, vol. 191, pp. 261-269 (2014).
- Mengual et al. Supplementary data to *Clinical Cancer Research*, vol. 16, 10 pages (2010).
- Minas et al. "Intratumoral CRH modulates immuno-escape of ovarian cancer cells through FasL regulation" *Br J Cancer.* 97(5):637-45 (2007).
- Miyamoto et al. "GATA binding protein 3 is down-regulated in bladder cancer yet strong expression is an independent predictor of poor prognosis in invasive tumor," *Human Pathology.* 43:2033-2040 (2012).
- Moon et al. "Benzylidihydroxyoctenone, a novel anticancer agent, induces apoptosis via mitochondrial-mediated pathway in androgen-

(56)

References Cited**OTHER PUBLICATIONS**

sensitive LNCaP prostate cancer cells," *Bioorganic & Medicinal Chemistry Letters*. 19:742-744 (2009).

Munksgaard et al. "Low ANXA10 expression is associated with disease aggressiveness in bladder cancer," *Br J Cancer*. 105(9):1379-87 (2011).

National Cancer Institute "What You Need To Know About: Bladder Cancer" (NIH Publication No. 10-1559, published Aug. 2010). NCBI Reference Sequence: NM_001127598.1, *Homo sapiens* insulin-like growth factor 2 (somatomedin A) (IGF2), transcript variant 3, mRNA), 2011.

NCBI Reference Sequence: NM-000756.2, *Homo sapiens* corticotropin releasing hormone (CRH), mRNA, 2011.

Olsburgh et al. "Uroplakin gene expression in normal human tissues and locally advanced bladder cancer," *J Pathol*. 199:41-49 (2003).

O'Sullivan et al. "A Multigene Urine Test for the Detection and Stratification of Bladder Cancer in Patients Presenting with Hematuria," *The Journal of Urology*, vol. 188, pp. 741-747 (2012).

Ozdemir et al. "IMP3 expression in urothelial carcinomas of the urinary bladder," *Türk Patoloji Derg*. 27(1):31-7 (2011).

Parker et al. "Current and Emerging Bladder Cancer Urinary Biomarkers," *The Scientific World Journal*, 11, pp. 1103-1112 (2011).

Pignot et al. "Hedgehog pathway activation in human transitional cell carcinoma of the bladder," *Br J Cancer*, 106(6):1177-86 (2012).

Plesea et al. "Detection and morphological profile of early stages bladder carcinoma: preliminary study," *Virchows Arch*. 459(Suppl 1):S296 (2011).

Pu et al. "The value of combined use of survivin, cytokeratin 20 and mucin 7 mRNA for bladder cancer detection in voided urine," *J Cancer Res Clin Oncol*. 134(6):659-65. 2008 (Epub 2007).

Qiagen, "Using endogenous control genes in real-time RT-PCR", <https://www.qiagen.com/us/spotlight-pages/newsletters-and-magazines/articles/endogenous-controls> (2016).

Qian et al. "Characteristics of hepatic igf-ii expression and monitored levels of circulating igf-ii mRNA in metastasis of hepatocellular carcinoma," *Am J Clin Pathol*. 134(5):799-806 (2010).

Raica et al. "Cytokeratin 20, 34betaE12 and overexpression of HER-2/neu in urine cytology as predictors of recurrences in superficial urothelial carcinoma," *Rom J Morphol Embryol*. 46(1):11-5 (2005).

Retz et al. "Mucin 7 and cytokeratin 20 as new diagnostic urinary markers for bladder tumor," *J Urol*. 169(1):86-9 (2003).

Ribal et al. "Molecular staging of bladder cancer with RT-PCR assay for CK20 in peripheral blood, bone marrow and lymph nodes: comparison with standard histological staging," *Anticancer Res*. 26(1A):411-9 (2006).

Rosser et al. "Bladder Cancer-Associated Gene Expression Signatures Identified by Profiling of Exfoliated Urothelia. *Cancer Epidemiology, Biomarkers, & Prevention*, 18(2):444-453 (2009).

Sambrook et al. "Molecular Cloning: A Laboratory Manual," 3rd Edition, Cold Spring Harbor Laboratory Press, New York, pp. 8.1-8.126 (2001).

Sanchez-Carbajo et al. "Gene discovery in bladder cancer progression using cDNA microarrays," *Am J Pathol*. 163(2):505-16 (2003).

Schmittgen et al. "Analyzing real-time PCR data by the comparative Ct Method" *Nature Protocols* 2(6), pp. 1101-1108 (2008).

Sharma et al. "Cancer-Testis Antigens: Expression and Correlation with Survival in Human Urothelial Carcinoma," *Clinical Cancer Research*, vol. 12, No. 18, pp. 5442-5447 (2006).

Shulman et al. "Evaluation of Four Different Systems for Extraction of RNA from Stool Suspensions Using MS-2 Coliphage as an Exogenous Control for RT-PCR Inhibition," *PLoS One*. 7(7):e39455 (2012).

Siracusano et al. "The simultaneous use of telomerase, cytokeratin 20 and CD4 for bladder cancer detection in urine," *Eur Urol*. 47(3):327-33. 2005 (Epub 2004).

Soong et al. "Quantitative reverse transcription-polymerase chain reaction detection of cytokeratin 20 in noncolorectal lymph nodes," *Clin Cancer Res*. 7(11):3423-9 (2001).

Stevenson et al. "The use of Armored RNA as a multi-purpose internal control for RT-PCR", *Journal of Virological Methods* 150 :73-76 (2008).

Taqman Array Gene information , 18 pages (2016).

Taqman Array Protocol, Publication No. 4391016 Rev F. total of 53 pages (2011).

Varallyay et al. "MicroRNA detection by northern blotting using locked nucleic acid probes," *Nature Protocols*. 3:190-196 (2008).

Varga et al. "Methylation of a CpG Island with the Uroplakin Ib promoter: A possible mechanism for loss of uroplakin Ib expression in bladder carcinoma," *Neoplasia*. 6(2): 128-135 (2004).

Vrooman et al. "Urinary Markers in Bladder Cancer", *European Urology* 53, pp. 909-916 (2008).

Walkerpeach et al. "Ribonuclease-resistant RNA controls (Armored RNA) for reverse transcription-PCR, branched DNA, and genotyping assays for hepatitis C virus," *Clin Chem*. 45:2079-2085 (1999).

Wallace et al. "Development of a 90-Minute Integrated Noninvasive Urinary Assay for Bladder Cancer Detection," *The Journal of Urology*, vol. 199, Issue 3, pp. 655-662 (2018).

Watson et al. "Urinary insulin-like growth factor 2 identifies the presence of urothelial carcinoma of the bladder," *BJU Int*. 103(5):694-7. 2009 (Epub 2008).

Weber et al. "Detection of disseminated medullary thyroid carcinoma cells in cervical lymph nodes by cytokeratin 20 reverse transcription-polymerase chain reaction," *World J Surg*. 26(2):148-52. 2002 (Epub 2001).

Whitehead et al. "Variation in tissue-specific gene expression among natural populations," *Genome Biology*, 6:R13 (2005).

Wilkinson, M. "A rapid and convenient method for isolation of nuclear, cytoplasmic and total cellular RNA," *Nucleic Acids Res*. 16:10934 (1988).

Wilkinson, M. "A rapid and convenient method for isolation of nuclear, cytoplasmic and total cellular RNA," *Nucleic Acids Res*. 16:10933 (1988).

Williams et al. "Androgen receptor expression in genitourinary neoplasms," *Modern Pathology* 26(S2):258A Feb. 2013, and 102nd Annual Meeting of the United States and Canadian Academy of Pathology (USCAP); Baltimore MD, USA (2013).

Wu et al. "Uroplakin II as a promising marker for molecular diagnosis of nodal metastases from bladder cancer: comparison with cytokeratin 20," *J Urol*. 174(6):2138-42, Editorial Comment 2142-3 (2005).

Ye et al. "CK20 and Ki-67 as significant prognostic factors in human bladder carcinoma," *Clin Exp Med*. 10(3):153-8 (2010).

Zheng et al. "Dihydrotestosterone upregulates the expression of epidermal growth factor receptor and ERBB2 in androgen receptor-positive bladder cancer cells. *Endocrine Related Cancer*," 18:451-464 (2011).

* cited by examiner

1

METHODS OF DETECTING BLADDER CANCER

1. CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a continuation of U.S. application Ser. No. 16/272,978, filed Feb. 11, 2019, which is a divisional of U.S. application Ser. No. 14/394,352, filed Oct. 14, 2014, which is a 371 application of International Application No. PCT/US2013/037334, filed Apr. 19, 2013, which claims the benefit of U.S. Provisional Application No. 61/636,194, filed Apr. 20, 2012 and of U.S. Provisional Application No. 61/770,803 filed Feb. 28, 2013, all of which are incorporated herein by reference in their entireties.

2. SEQUENCE LISTING

The present application is filed with a Sequence Listing in electronic format. The Sequence Listing is provided as a file entitled "01148-0006-02US_Sequence_Listing_ST26" created on Sep. 16, 2023, which is 69,632 bytes in size. The information in the electronic format of the sequence listing is incorporated herein by reference in its entirety.

3. FIELD OF THE INVENTION

Compositions and methods for detecting bladder cancer are provided. In particular, bladder cancer markers and panels of markers useful in the detection of bladder cancer are provided.

4. BACKGROUND

386,000 cases of bladder cancer are diagnosed globally each year, including 70,500 cases per year in the United States. The incidence of bladder cancer is three times higher in men than in women. The highest incidence and prevalence are found in the European Union, North America, North Africa, and the Middle East.

Smoking is the greatest risk factor for bladder cancer. Additional risk factors include chemical exposure, chemotherapy (such as Cytoxan), radiation treatment, and chronic bladder infection.

Bladder tumors include papillary tumors, which are urothelial carcinomas that grow narrow, finger-like projections; and nonpapillary (sessile) tumors, such as carcinoma-in-situ, which are less common but have a high risk of becoming invasive.

Symptoms of bladder cancer can include abdominal pain, blood in the urine, bone pain or tenderness, fatigue, painful urination, frequent urination, urinary urgency, incontinence, and weight loss. Diagnosis is generally based on imaging, urinalysis, and/or biopsy.

The prognosis for bladder cancer depends on the stage of cancer at diagnosis. The prognosis for early tumors is favorable, while the prognosis for advanced tumors is poor. Long-term follow up is recommended to detect cancer recurrence, which occurs in up to 70% of bladder cancers. For the first two years, cystoscopy and urine cytology are recommended every 3 to 4 months, and then at longer intervals in subsequent years, often for the patient's lifetime. These methods are invasive and costly, making bladder cancer one of the most expensive cancers to treat from diagnosis until death.

Existing non-invasive diagnostic tests include ImmunoCyt™ (Scimedx, Denville, NJ) and UroVysion® (Abbott

2

Molecular, Abbott Park, IL). ImmunoCyt™ is a cytology assay that uses a cocktail of three monoclonal antibodies labeled with fluorescent markers to detect certain cellular markers of bladder cancer in exfoliated cells isolated from urine samples. ImmunoCyt™ is used in conjunction with standard urine cytology to improve cytology's sensitivity at detecting tumor cells. UroVysion® is also a cytology-based assay, which detects aneuploidy in certain chromosomes via fluorescent in situ hybridization (FISH). Determination of the results is conducted by enumerating signals through microscopic examination of the nucleus of cells in urine.

Improved methods for early detection of bladder cancer are needed. In particular, an accurate urine-based diagnostic test that does not rely on cytology could reduce the need for costly and invasive cystoscopy and labor-intensive and potentially subjective cytology assays.

5. SUMMARY

Compositions and methods for detecting bladder cancer are provided. In particular, bladder cancer markers and panels of markers useful in the detection of bladder cancer are provided. In some embodiments, the levels of CRH, IGF2, KRT20 and ANXA10 mRNA are measured, for example, by quantitative RT-PCR, and the results can be used to determine whether or not a subject has bladder cancer. In some embodiments, the levels of CRH, IGF2, KRT20, and ANXA10 mRNA are normalized to an endogenous control. In some embodiments, the endogenous control is ABL mRNA. In some embodiments, an endogenous control is selected that is expected to be expressed at similar levels in bladder urothelial cells from subjects with and without bladder cancer. In some embodiments, the sample is a urine sample. In some embodiments, the present methods are used to monitor subjects with a history of bladder cancer for tumor recurrence. In some embodiments, the subject has been treated with Bacillus Calmette-Guerin (BCG) within the past three months. In some embodiments, the present methods are used to detect bladder cancer in subjects with no history of bladder cancer. In some such embodiments, the subjects have symptoms of bladder cancer. Nonlimiting exemplary symptoms of bladder cancer include abdominal pain, blood in the urine, bone pain or tenderness, fatigue, painful urination, frequent urination, urinary urgency, incontinence, and weight loss.

In some embodiments, methods for detecting the presence of bladder cancer in a subject are provided. In some embodiments, a method comprises detecting the levels of each marker of a set of bladder cancer markers in a sample from the subject, wherein the set of bladder cancer markers consists of corticotrophin releasing hormone (CRH), insulin-like growth factor 2 (IGF2), keratin 20 (KRT20) and annexin 10 (ANXA10). In some embodiments, detection of an elevated level of at least one marker indicates the presence of bladder cancer in the subject. In some embodiments, a method further comprises detecting an endogenous control. In some embodiments, the endogenous control is selected from ABL, GUSB, GAPDH, TUBB, and UPK1. In some embodiments, the endogenous control is ABL. In some embodiments, a method comprises detecting an exogenous control. In some embodiments, the exogenous control is an RNA. In some such embodiments, the exogenous control is an Armored RNA®.

In some embodiments, detecting comprises RT-PCR. In some embodiments, detecting comprises quantitative RT-PCR. In some embodiments, a method comprises comparing a Ct value or a Δ Ct value to a threshold Ct value or Δ Ct

4

[illegible]

5

9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 39, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

In some embodiments, a method further comprises contacting RNA from the sample with an endogenous control primer pair. In some such embodiments, the endogenous control primer pair is for detecting an endogenous control is selected from ABL, GUSB, GAPDH, TUBB, and UPK1. In some embodiments, the endogenous control primer pair is for detecting ABL. In some embodiments, the endogenous control primer pair comprises a first primer comprising SEQ ID NO: 8 and a second primer comprising SEQ ID NO: 9, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the endogenous control primer pair comprises a first primer comprising SEQ ID NO: 41 and a second primer comprising SEQ ID NO: 42, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the endogenous control primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 8 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 9, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the endogenous control primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 41 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 42, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

In some embodiments, a method comprises contacting RNA from the sample with an exogenous control primer pair. In some such embodiments, the exogenous control primer pair is for detecting an exogenous RNA. In some embodiments, the exogenous control primer pair comprises a first primer comprising SEQ ID NO: 23 and a second primer comprising SEQ ID NO: 24, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the exogenous control primer pair comprises a first primer comprising SEQ ID NO: 44 and a second primer comprising SEQ ID NO: 45, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the exogenous control primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 23 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 24, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the exogenous control primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 44 and a second primer

6

comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 45, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

In some embodiments, the method comprises forming a set of bladder cancer marker amplicons, wherein the set of bladder cancer marker amplicons consists of a CRH amplicon, an IGF2 amplicon, a KRT20 amplicon, and an ANXA10 amplicon, and contacting the bladder cancer marker amplicons with a set of bladder cancer marker probes, wherein the set of bladder cancer marker probes consists of a first probe for detecting the CRH amplicon, a second probe for detecting the IGF2 amplicon, a third probe for detecting the KRT20 amplicon, and a fourth probe for detecting the ANXA10 amplicon. In some embodiments, the first probe comprises SEQ ID NO: 21 or SEQ ID NO: 37, wherein the first probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second probe comprises SEQ ID NO: 34 or SEQ ID NO: 18, wherein the second probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third probe comprises SEQ ID NO: 15 or SEQ ID NO: 31, wherein the third probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth probe comprises SEQ ID NO: 28 or SEQ ID NO: 40, wherein the fourth probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the first probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 21 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 37, wherein the first probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 34 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 18, wherein the second probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 15 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 31, wherein the third probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 28 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 40, wherein the fourth probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, each bladder cancer marker probe comprises a dye, and wherein each dye is detectably different from the other three labels. In some embodiments, each bladder cancer marker probe comprises a fluorescent dye and a quencher molecule.

In some embodiments, a method comprises forming an endogenous control amplicon, and contacting the endogenous control amplicon with an endogenous control probe. In some embodiments, the endogenous control probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 10, 11, 12, or 43, wherein the endogenous control probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the endogenous control probe comprises a dye that is detectably different from the dyes of the bladder cancer marker probes.

In some embodiments, a method comprises forming an exogenous control amplicon, and contacting the exogenous control amplicon with an exogenous control probe. In some embodiments, the exogenous control probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 25 or 46, wherein the exogenous control probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the exogenous control probe comprises a dye that is detectably different from the dyes of the bladder cancer marker probes and the endogenous control probe.

In some embodiments, the set of bladder cancer markers are detected in a single multiplex reaction.

In some embodiments, the sample comprises urothelial cells. In some embodiments, the sample is selected from a urine sample and a bladder washing sample. In some embodiments, the subject has a history of bladder cancer. In some embodiments, the subject is being monitored for recurrence of bladder cancer.

In some embodiments, compositions are provided. In some embodiments, a composition comprises a set of bladder cancer marker primer pairs, wherein the set of bladder cancer marker primer pairs consists of a first primer pair for detecting CRH, a second primer pair for detecting IGF2, a third primer pair for detecting KRT20, and a fourth primer pair for detecting ANXA10. In some embodiments, the first primer pair comprises a first primer comprising SEQ ID NO: 19 and a second primer comprising SEQ ID NO: 20, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the first primer pair comprises a first primer comprising SEQ ID NO: 35 and a second primer comprising SEQ ID NO: 36, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second primer pair comprises a first primer comprising SEQ ID NO: 16 and a second primer comprising SEQ ID NO: 17, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second primer pair comprises a first primer comprising SEQ ID NO: 32 and a second primer comprising SEQ ID NO: 33, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third primer pair comprises a first primer comprising SEQ ID NO: 13 and a second primer comprising SEQ ID NO: 14, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third primer pair comprises a first primer comprising SEQ ID NO: 29 and a second primer comprising SEQ ID NO: 30, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth primer pair comprises a first primer comprising SEQ ID NO: 26 and a second primer comprising SEQ ID NO: 27, wherein

each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth primer pair comprises a first primer comprising SEQ ID NO: 38 and a second primer comprising SEQ ID NO: 39, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth primer pair comprises a first primer comprising SEQ ID NO: 48 and a second primer comprising SEQ ID NO: 39, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

In some embodiments, a composition comprises a set of bladder cancer marker primer pairs, wherein the set of bladder cancer marker primer pairs consists of a first primer pair for detecting CRH, a second primer pair for detecting IGF2, a third primer pair for detecting KRT20, and a fourth primer pair for detecting ANXA10. In some embodiments, the first primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 19 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 20, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the first primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 35 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 36, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 16 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 17, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 32 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 33, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 13 and a second primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 14, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third primer pair comprises a first primer comprising at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 29 and a second primer comprising at least 8, at least 9, at

In some embodiments, a composition further comprises a set of bladder cancer marker probes, wherein the set of bladder cancer marker probes consists of a first probe for detecting a CRH amplicon, a second probe for detecting an IGF2 amplicon, a third probe for detecting a KRT20 amplicon, and a fourth probe for detecting an ANXA10 amplicon. In some embodiments, the first probe comprises SEQ ID NO: 21 or SEQ ID NO: 37, wherein the first probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second probe comprises SEQ ID NO: 34 or SEQ ID NO: 18, wherein the second probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the third probe comprises SEQ ID NO: 15 or SEQ ID NO: 31, wherein the third probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the fourth probe comprises SEQ ID NO: 28 or SEQ ID NO: 40, wherein the fourth probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the first probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 21 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 37, wherein the first probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long. In some embodiments, the second probe comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 34 or at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, or at least 15, at least 16, at least 17, or at least 18 nucleotides of SEQ ID NO: 18, wherein the second probe is less than 50, less than 45, less

control probe comprises a dye that is detectably different from the dyes of the bladder cancer marker probes.

In some embodiments, a composition is a lyophilized composition. In some embodiments, the composition is a solution. In some embodiments, the composition further comprises urothelial cells. In some embodiments, the urothelial cells are from a urine sample.

Further embodiments and details of the inventions are described below.

6. DETAILED DESCRIPTION

6.1. Definitions

To facilitate an understanding of the present invention, a number of terms and phrases are defined below:

As used herein, the terms “detect”, “detecting” or “detection” may describe either the general act of discovering or discerning or the specific observation of a detectably labeled composition.

As used herein, the term “detectably different” refers to a set of labels (such as dyes) that can be detected and distinguished simultaneously.

As used herein, the terms “patient” and “subject” are used interchangeably to refer to a human. In some embodiments, the methods described herein may be used on samples from non-human animals.

As used herein, “bladder cancer” is a tumor, such as a transitional cell carcinoma, arising from the lining of the bladder, and includes low grade and high grade bladder cancers, as well as metastatic bladder cancer. “Low grade bladder cancer” refers to superficial tumors that project into the interior of the bladder cavity. Low grade bladder cancers have a high rate of recurrence. “High grade bladder cancer” refers to a fast-growing and/or invasive tumor that invades the bladder wall. High grade bladder cancers have the potential to spread (i.e., metastasize) to other areas of the body. “Metastatic bladder cancer” refers to invasive bladder cancer that has spread to one or more locations in the body beyond the bladder.

As used herein, the terms “oligonucleotide,” “polynucleotide,” “nucleic acid molecule,” and the like, refer to nucleic acid-containing molecules, including but not limited to, DNA or RNA. The term encompasses sequences that include any of the known base analogs of DNA and RNA including, but not limited to, 4-acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxymethyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethyl-aminomethyluracil, dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudouracil, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

As used herein, the term “oligonucleotide,” refers to a single-stranded polynucleotide having fewer than 500 nucleotides. In some embodiments, an oligonucleotide is 8 to 200, 8 to 100, 12 to 200, 12 to 100, 12 to 75, or 12 to 50

nucleotides long. Oligonucleotides may be referred to by their length, for example, a 24 residue oligonucleotide may be referred to as a “24-mer.”

As used herein, the term “complementary” to a target RNA (or target region thereof), and the percentage of “complementarity” of the probe sequence to that of the target RNA sequence is the percentage “identity” to the sequence of target RNA or to the reverse complement of the sequence of the target RNA. In determining the degree of “complementarity” between probes used in the compositions described herein (or regions thereof) and a target RNA, such as those disclosed herein, the degree of “complementarity” is expressed as the percentage identity between the sequence of the probe (or region thereof) and sequence of the target RNA or the reverse complement of the sequence of the target RNA that best aligns therewith. The percentage is calculated by counting the number of aligned bases that are identical as between the 2 sequences, dividing by the total number of contiguous nucleotides in the probe, and multiplying by 100. When the term “complementary” is used, the subject oligonucleotide is at least 90% complementary to the target molecule, unless indicated otherwise. In some embodiments, the subject oligonucleotide is at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to the target molecule.

A “primer” or “probe” as used herein, refers to an oligonucleotide that comprises a region that is complementary to a sequence of at least 8 contiguous nucleotides of a target nucleic acid molecule, such as an mRNA or a DNA reverse-transcribed from an mRNA. In some embodiments, a primer or probe comprises a region that is complementary to a sequence of at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of a target molecule. When a primer or probe comprises a region that is “complementary to at least x contiguous nucleotides of a target molecule,” the primer or probe is at least 95% complementary to at least x contiguous nucleotides of the target molecule. In some embodiments, the primer or probe is at least 96%, at least 97%, at least 98%, at least 99%, or 100% complementary to the target molecule.

A “sample,” as used herein, includes urine samples (including samples derived from urine samples), and other types of human samples. As used herein, urine samples include, but are not limited to, whole urine, a sample comprising cells from a urine sample, a sample comprising the cell pellet isolated by centrifugation of a urine sample, a sample comprising cells isolated by filtration of a urine sample, and the like. In some embodiments, a urine sample comprises a preservative, such as a preservative that causes damage, such as lysis, of red and/or white blood cells. In some embodiments, a sample is a human sample other than a urine sample, such as a tissue sample (including a bladder tissue and/or bladder tumor sample), a blood sample (including whole blood, serum, plasma, etc.), etc. In some embodiments, a sample is a bladder washing sample.

As used herein, “corticotrophin releasing hormone” or “CRH” refers to an mRNA that encodes CRH, as well as the CRH protein. In some embodiments, CRH is human CRH. Nonlimiting exemplary human CRH mRNA sequences are found at GenBank Accession No. NM_000756, and at SEQ ID NO: 1.

As used herein, “insulin-like growth factor 2” or “IGF2” refers to an mRNA that encodes IGF2, as well as the IGF2

protein. In some embodiments, IGF2 is human IGF2. Non-limiting exemplary human IGF2 mRNA sequences are shown in SEQ ID NOs: 2 to 4.

As used herein, “keratin 20” or “KRT20” refers to an mRNA that encodes KRT20, as well as the KRT20 protein. In some embodiments, KRT20 is human KRT20. Non-limiting exemplary human KRT20 mRNA sequences are found at GenBank Accession No. NM_019010, and at SEQ ID NO: 5.

As used herein, “annexin A10” or “ANXA10” refers to an mRNA that encodes ANXA10, as well as the ANXA10 protein. In some embodiments, ANXA10 is human ANXA10. Non-limiting exemplary human ANXA10 mRNA sequences are found at GenBank Accession No. NM_007193, and at SEQ ID NO: 6.

An “endogenous control,” as used herein refers to a moiety that is naturally present in the sample to be used for detection, and which can be used to normalize the levels of the bladder cancer markers described herein (including, but not limited to, CRH, IGF2, KRT20 and ANXA10). Thus, an endogenous control is typically a moiety that is present at similar levels from cell to cell, and at similar levels in cells from subjects with bladder cancer and cells from subjects without bladder cancer. In some embodiments, an endogenous control is an RNA (such as an mRNA, tRNA, ribosomal RNA, etc.). Non-limiting exemplary endogenous controls include ABL mRNA, GUSB mRNA, GAPDH mRNA, TUBB mRNA, and UPK1a mRNA. Non-limiting exemplary human ABL mRNA sequences are found at GenBank Accession No. NM_007313, and at SEQ ID NO: 7. In some embodiments, an endogenous control is selected that can be detected in the same manner as the bladder cancer markers are detected and, in some embodiments, simultaneously with the bladder cancer markers.

An “exogenous control,” as used herein, refers to a moiety that is added to a sample to be used for detection. An exogenous control is typically selected that is not expected to be present in the sample to be used for detection, or is present at very low levels in the sample such that the amount of the moiety naturally present in the sample is either undetectable or is detectable at a much lower level than the amount added to the sample as an exogenous control. In some embodiments, an exogenous control comprises a nucleotide sequence that is not expected to be present in the sample type used for detection of the bladder cancer markers. In some embodiments, an exogenous control comprises a nucleotide sequence that is not known to be present in the species from whom the sample is taken. In some embodiments, an exogenous control comprises a nucleotide sequence from a different species than the subject from whom the sample was taken. In some embodiments, an exogenous control comprises a nucleotide sequence that is not known to be present in any species. In some embodiments, an exogenous control is selected that can be detected in the same manner as the bladder cancer markers are detected and, in some embodiments, simultaneously with the bladder cancer markers. In some embodiments, an exogenous control is an RNA. In some embodiments, an RNA is an Armored RNA®, which comprises RNA packaged in a bacteriophage protective coat. See, e.g., Walker-Peach et al., (*Clin. Chem.* 45:12: 2079-2085 (1999)).

In the sequences herein, “U” and “T” are used interchangeably, such that both letters indicate a uracil or thymine at that position. One skilled in the art will understand from the context and/or intended use whether a uracil or thymine is intended and/or should be used at that position in the sequence. For example, one skilled in the art would

understand that native RNA molecules typically include uracil, while native DNA molecules typically include thymine. Thus, where an RNA sequence includes “T”, one skilled in the art would understand that that position in the native RNA is likely a uracil.

In the present disclosure, “a sequence selected from” encompasses both “one sequence selected from” and “one or more sequences selected from.” Thus, when “a sequence selected from” is used, it is to be understood that one, or more than one, of the listed sequences may be chosen.

In the present disclosure, a method that comprises detecting a “a set of bladder cancer markers consisting of . . .” involves detection of only the bladder cancer markers of the set, and not any further bladder cancer markers. The method may comprise additional components or steps, however, such as detecting endogenous and/or exogenous controls. Similarly, a method or composition that comprises “a set of bladder cancer marker primer pairs” and/or “a set of bladder cancer marker probes” can include primer pairs and/or probes for only the bladder cancer markers of the set, and not for any other bladder cancer markers. The method or composition may comprise additional components, however, such as one or more endogenous control primer pairs and/or one or more exogenous control primer pairs.

6.2. Detecting Bladder Cancer

The present inventors have developed an assay for detecting bladder cancer that involves detecting just four markers, CRH, IGF2, KRT20 and ANXA10 (plus one or two controls). The presently described assays have several advantages over existing and previously described diagnostics for bladder cancer. For example, the present assays do not rely on cytology, which can be costly, and requires trained cytologists for accurate interpretation of results. Instead, the present assays rely on the polymerase chain reaction (PCR), and can be carried out in a substantially automated manner, for example, using the GeneXpert® system (Cepheid, Sunnyvale, CA). Mengual et al., *Clin Cancer Res* (2010) 16: 2624-2633, recently described a “12+2 gene expression signature,” which detects 12 to 14 genes in an array format. Surprisingly, the present inventors have found that a much smaller signature, of just four markers (plus one or two controls), is sufficient to provide equivalent or better sensitivity and/or specificity over existing diagnostic assays for bladder cancer, including the Mengual et al. assay. Further, because the present assays use just four markers (plus one or two controls), they can be carried out in a single reaction mixture, using 4 to 6 detectably different dyes, such as fluorescent dyes. The present assays can be completed in under 3 hours, and in some embodiments, under 2 hours, using an automated system, for example, the GeneXpert® system. Existing tests can require several days for a laboratory to complete and send results. In addition, the present assay can be carried out on much smaller volumes of urine (in some embodiments, 5 ml or less). Thus, the present assays, which rely on PCR of just four markers (plus one or two controls) rather than cytology, allows for a fast, one-pot reaction for diagnosis of bladder cancer, which in many instances can be carried out at the point of care using an automated system such as GeneXpert®.

6.2.1. General Methods

Compositions and methods for detecting bladder cancer are provided. In some embodiments, compositions and methods for detecting low grade bladder cancer are pro-

vided. In some embodiments, compositions and methods of detecting high grade bladder cancer are provided. In some embodiments, compositions and methods for monitoring the recurrence of bladder cancer are provided.

In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10, and at least one endogenous control. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10, and at least one endogenous control. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10, and at least one exogenous control. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10, and at least one exogenous control. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10, at least one endogenous control, and at least one exogenous control. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10, at least one endogenous control, and at least one exogenous control.

In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10 mRNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA, and at least one endogenous control RNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10 mRNA, and at least one endogenous control RNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA, and at least one exogenous control RNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10 mRNA, and at least one exogenous control RNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA, at least one endogenous control RNA, and at least one exogenous control RNA. In some embodiments, a method of detecting bladder cancer comprises detecting the levels of a set of markers consisting of CRH, IGF2, KRT20 and ANXA10 mRNA, at least one endogenous control RNA, and at least one exogenous control RNA.

In the present disclosure, the term "target RNA" is used for convenience to refer to CRH, IGF2, KRT20 and ANXA10 mRNAs and also to other target RNAs, such as exogenous and/or endogenous control RNAs. Thus, it is to be understood that when a discussion is presented in terms of a target RNA, that discussion is specifically intended to encompass CRH, IGF2, KRT20 and ANXA10 mRNAs, and/or other target RNAs.

In some embodiments, the level of one or more target RNAs is detected in a urine sample. In some embodiments, the level of one or more target RNAs is determined in a urine

sample that has been preserved in a manner that causes damage, such as lysis, to red blood cells and/or white blood cells. In some embodiments, the level of one or more target RNAs is detected in urothelial cells isolated from a urine sample, either with or without preservative treatment. In some embodiments, the urothelial cells are isolated by filtration.

In some embodiments, detection of an elevated level of one or more target RNAs selected from CRH, IGF2, KRT20 and ANXA10 in a sample from a subject indicates the presence of bladder cancer in the subject. In some embodiments, the detecting is done quantitatively. In other embodiments, the detecting is done qualitatively. In some embodiments, detecting a target RNA comprises forming a complex comprising a polynucleotide and a nucleic acid selected from a target RNA, a DNA amplicon of a target RNA, and a complement of a target RNA. In some embodiments, detecting a target RNA comprises RT-PCR. In some embodiments, detecting a target RNA comprises quantitative RT-PCT. In some embodiments, the level of the target RNA is compared to a normal or control level of the target RNA.

In some embodiments, the levels of target RNAs, such as CRH, IGF2, KRT20 and ANXA10 mRNA, can be measured in samples collected at one or more times from a patient to monitor the status or progression of bladder cancer in the patient. In some embodiments, a patient with a history of bladder cancer, such as a history of low grade bladder cancer or a history of high grade bladder cancer, is monitored by detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA at regular or semi-regular intervals. In some such embodiments, the patient is monitored by detecting the levels of the target RNAs at least once per month, at least once every two months, at least once every three months, at least once every four months, at least once every five months, at least once every six months, at least once every nine months, at least once per year, or at least once every two years.

In some embodiments, a sample to be tested is a urine sample (such as a voided urine sample), or is derived from a urine sample. In some embodiments, a preservative is added to the urine sample, for example, to damage (e.g., lyse) red and/or white blood cells present in the urine sample. By damaging or lysing red and/or white blood cells prior to isolation of urothelial cells, contamination by the red and/or white blood cells can be reduced. In some embodiments, the urine sample is centrifuged to concentrate the urothelial cells. In some embodiments, the urine sample is filtered to isolate the urothelial cells from other urine and preservative materials. In some such embodiments, the filter is part of a GeneXpert cartridge (Cepheid, Sunnyvale, CA).

In some embodiments, less than 5 ml, less than 4 ml, less than 3 ml, or less than 2 ml of urine are used in the present methods. In some embodiments, the urine sample is analyzed without a centrifugation step. Thus, in some embodiments, the present methods are carried out in the absence of centrifugation. In some embodiments, a larger volume of urine may be used, and in some such embodiments, a centrifugation step may be used to concentrate the urothelial cells prior to analysis.

In some embodiments, the sample to be tested is another bodily fluid, such as blood, sputum, mucus, saliva, semen, etc. In some embodiments, a sample to be tested is a blood sample. In some embodiments, the blood sample is whole blood. In some embodiments, the blood sample is a sample of blood cells. In some embodiments, the blood sample is plasma. In some embodiments, the blood sample is serum.

The clinical sample to be tested is, in some embodiments, fresh (i.e., never frozen). In other embodiments, the sample is a frozen specimen. In some embodiments, the sample is a tissue sample, such as a formalin-fixed paraffin embedded sample. In some embodiments, the sample is a liquid cytology sample.

In some embodiments, the methods described herein are used for early detection of bladder cancer in a sample of urothelial cells, such as those obtained from voided urine. In some embodiments, the methods described herein are used for monitoring for recurrence of bladder cancer using a sample of urothelial cells, such as those obtained from voided urine.

In some embodiments, the sample to be tested is obtained from an individual who has one or more of the following risk factors: history of smoking, hematuria, history of bladder or other cancers, and exposure to known carcinogens such as benzene. In some embodiments, the sample is obtained from an individual who has diagnostic signs or clinical symptoms that may be associated with bladder cancer, such as blood in the urine, frequent urination, urinary urgency, incontinence, difficulty urinating, abdominal pain, unexplained weight loss and/or loss of appetite. In some embodiments, the sample to be tested is obtained from an individual who has previously been diagnosed with low grade or high grade bladder cancer. In some such embodiments, the individual is monitored for recurrence of bladder cancer.

Bladder cancer can be divided into stages, which indicate the growth pattern of the primary tumor. Table A shows the stages of bladder cancer according to the American Joint Committee on Cancer (AJCC). The stages shown cover only the "T" portion of the "TNM" system. The "T" portion refers to the primary tumor, while "N" refers to spread of the cancer to the lymph nodes, and "M" refers to whether the cancer has metastasized to distant sites.

TABLE A

Bladder cancer stages	
Stage	description
T0	No evidence of primary tumor
Ta	Non-invasive papillary carcinoma
Tis/ CIS	Non-invasive flat carcinoma (carcinoma in situ)
T1	Tumor has grown from the lining of the bladder into the connective tissue, but has not grown into the muscle layer of the bladder
T2	Tumor has grown into the muscle layer
T2a	Tumor has grown into the inner half of the muscle layer
T2b	Tumor has grown into the outer half of the muscle layer
T3	Tumor has grown through the muscle layer of the bladder and into the fatty tissue that surrounds it
T3a	Tumor's spread to fatty tissue can only be seen under a microscope
T3b	Tumor's spread to fatty tissue can be seen on imaging tests or can be seen or felt by surgeon
T4	Tumor has spread beyond the fatty tissue to nearby organs or structures
T4a	Tumor has spread to the stroma of the prostate, or to the uterus and/or vagina
T4b	Tumor has spread to the pelvic wall or abdominal wall

In some embodiments, methods described herein can be used for routine screening of healthy individuals with no risk factors. In some embodiments, methods described herein are used to screen asymptomatic individuals having one or more of the above-described risk factors.

In some embodiments, the methods described herein can be used to detect low grade bladder cancer. In some embodiments, the methods described herein can be used to detect high grade bladder cancer. In some embodiments, the meth-

ods described herein detect at least 15%, at least 17%, at least 20%, at least 22%, at least 25%, at least 27%, at least 30%, at least 31%, at least 32%, at least 33%, at least 34%, or at least 35% of low grade bladder cancers. In some embodiments, the methods described herein detect at least 85%, at least 87%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% of high grade bladder cancers.

In some embodiments, the methods described herein can be used to assess the effectiveness of a treatment for bladder cancer in a patient. In some embodiments, target RNA levels, such as the levels of CRH, IGF2, KRT20 or ANXA10 mRNA, are determined at various times during the treatment, and are compared to target RNA levels from an archival sample taken from the patient before the beginning of treatment. In some embodiments, target RNA levels are compared to target RNA levels from an archival normal sample taken from the patient. Ideally, target RNA levels in the normal sample evidence no aberrant changes in target RNA levels.

In some embodiments, use of the levels of CRH, IGF2, KRT20 and ANXA10 mRNA for monitoring recurrence of bladder cancer is provided. In some embodiments, an elevated level of one or more, two or more, three or more, or all four mRNAs indicates that bladder cancer has recurred in the patient.

In any of the embodiments described herein, RNA levels may be detected concurrently or simultaneously in the same or separate assay reactions. In some embodiments, RNA levels are detected at different times, e.g., in serial assay reactions.

In some embodiments, a method comprises detecting the level of CRH, IGF2, KRT20 and ANXA10 mRNA in a sample from a subject, wherein detection of a level of any one of the four mRNAs that is greater than a normal level of the RNA indicates the presence of bladder cancer in the subject. In some embodiments, detection of elevated levels of two, three, or four bladder cancer marker mRNAs indicates the presence of bladder cancer in the subject. In some embodiments, detection of elevated levels of at least two, at least three, or all four of the bladder cancer marker mRNAs indicates a greater risk of high grade bladder cancer.

In some embodiments, a method of facilitating diagnosis of bladder cancer in a subject is provided. Such methods comprise detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA in a sample from the subject. In some embodiments, information concerning the levels of CRH, IGF2, KRT20 and/or ANXA10 mRNA in the sample from the subject is communicated to a medical practitioner. A "medical practitioner," as used herein, refers to an individual or entity that diagnoses and/or treats patients, such as a hospital, a clinic, a physician's office, a physician, a nurse, or an agent of any of the aforementioned entities and individuals. In some embodiments, detecting the levels of CRH, IGF2, KRT20 and ANXA10 mRNA is carried out at a laboratory that has received the subject's sample from the medical practitioner or agent of the medical practitioner. The laboratory carries out the detection by any method, including those described herein, and then communicates the results to the medical practitioner. A result is "communicated," as used herein, when it is provided by any means to the medical practitioner. In some embodiments, such communication may be oral or written, may be by telephone, in person, by e-mail, by mail or other courier, or may be made by directly depositing the information into, e.g., a database accessible by the medical practitioner, including databases

not controlled by the medical practitioner. In some embodiments, the information is maintained in electronic form. In some embodiments, the information can be stored in a memory or other computer readable medium, such as RAM, ROM, EEPROM, flash memory, computer chips, digital video discs (DVD), compact discs (CDs), hard disk drives (HDD), magnetic tape, etc.

In some embodiments, methods of detecting the presence bladder cancer are provided. In some embodiments, methods of diagnosing bladder cancer are provided. In some embodiments, the method comprises obtaining a sample from a subject and providing the sample to a laboratory for detection of levels of CRH, IGF2, KRT20 and ANXA10 mRNA in the sample. In some embodiments, the method further comprises receiving a communication from the laboratory that indicates the level of at least one RNA selected from CRH, IGF2, KRT20 and ANXA10 mRNA in the sample. In some embodiments, bladder cancer is present if the level of any one of the four mRNAs is greater than a normal or control level of the mRNA. A "laboratory," as used herein, is any facility that detects the levels of CRH, IGF2, KRT20 and ANXA10 mRNA in a sample by any method, including the methods described herein, and communicates the level to a medical practitioner. In some embodiments, a laboratory is under the control of a medical practitioner. In some embodiments, a laboratory is not under the control of the medical practitioner.

When a laboratory communicates the level of at least one RNA selected from CRH, IGF2, KRT20 and ANXA10 to a medical practitioner, in some embodiments, the laboratory communicates a numerical value representing the level of the RNA in the sample, with or without providing a numerical value for a normal level. In some embodiments, the laboratory communicates the level of the RNA by providing a qualitative value, such as "high," "low," "elevated," "decreased," "positive" (such as "CRH positive" or "CRH and KRT20 positive"), etc. In some embodiments, the laboratory communicates a suggested diagnosis, such as "bladder cancer positive" or "positive for cancer," and the like; or simply "cancer positive" or "cancer negative."

As used herein, when a method relates to detecting bladder cancer, determining the presence of bladder cancer, monitoring for bladder cancer, and/or diagnosing bladder cancer, the method includes activities in which the steps of the method are carried out, but the result is negative for the presence of bladder cancer. That is, detecting, determining, monitoring, and diagnosing bladder cancer include instances of carrying out the methods that result in either positive or negative results.

In some embodiments, more than one RNA is detected simultaneously in a single reaction. In some embodiments, CRH, IGF2, KRT20 and ANXA10 mRNAs are detected simultaneously in a single reaction. In some embodiments, CRH, IGF2, KRT20 and ANXA10 mRNAs and at least one endogenous control and/or at least one exogenous control are detected simultaneously in a single reaction. In some embodiments, CRH, IGF2, KRT20 and ANXA10 mRNAs, and endogenous control, and an exogenous control are detected simultaneously in a single reaction.

6.2.2. Exemplary Controls

In some embodiments, a normal level (a "control") of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, can be determined as an average level or range that is characteristic of normal urothelial cells or other reference material, against which the level measured in the sample can

be compared. The determined average or range of a target RNA in normal subjects can be used as a benchmark for detecting above-normal levels of the target RNA that are indicative of bladder cancer. In some embodiments, normal levels of a target RNA can be determined using individual or pooled RNA-containing samples from one or more individuals, such as from normal urothelial cells isolated from urine of healthy individuals.

In some embodiments, determining a normal level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, comprises detecting a complex comprising a polynucleotide for detection hybridized to a nucleic acid selected from a target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA. That is, in some embodiments, a normal level can be determined by detecting a DNA amplicon of the target RNA, or a complement of the target RNA rather than the target RNA itself. In some embodiments, a normal level of such a complex is determined and used as a control. The normal level of the complex, in some embodiments, correlates to the normal level of the target RNA. Thus, when a normal level of a target is discussed herein, that level can, in some embodiments, be determined by detecting such a complex.

In some embodiments, a normal level of a target RNA is, or has been, determined by the same method as the level of the target RNA from a patient sample. In some such embodiments, the method is RT-PCR (such as real-time RT-PCR, quantitative RT-PCR, etc.).

In some embodiments, a control comprises RNA from cells of a single individual, e.g., from normal urothelial cells isolated from urine of a healthy individual. In some embodiments, a control comprises RNA from blood, such as whole blood or serum, of a single individual. In some embodiments, a control comprises RNA from a pool of cells from multiple individuals. In some embodiments, a control comprises RNA from a pool of urine from multiple individuals. In some embodiments, a control comprises commercially-available human RNA (see, for example, Ambion). In some embodiments, a normal level or normal range has already been predetermined prior to testing a sample for an elevated level.

In some embodiments, the normal level of a target RNA can be determined from one or more continuous cell lines, typically cell lines previously shown to have levels of RNAs that approximate the levels in normal urothelial cells.

In some embodiments, quantitation of target RNA levels requires assumptions to be made about the total RNA per cell and the extent of sample loss during sample preparation. In order to correct for differences between different samples or between samples that are prepared under different conditions, the quantities of target RNAs in some embodiments are normalized to the levels of at least one endogenous control and/or at least one exogenous control.

In some embodiments, a control RNA is an endogenous control RNA. An endogenous control RNA may be any RNA suitable for the purpose, for example, RNAs that are present at approximately constant levels from cell to cell and in urothelial cells from both bladder cancer and non-bladder cancer patients. Nonlimiting exemplary endogenous control RNAs include ABL, GUSB, GAPDH, TUBB, and UPK1a. In some embodiments, one endogenous control is used for normalization. In some embodiments, more than one endogenous control is used for normalization.

In some embodiments, the level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, is normalized to an endogenous control RNA. Normalization may comprise, for example, determination of the difference of the level of

the target RNA to the level of the endogenous control RNA. In some such embodiments, the level of the RNAs are represented by a Ct value obtained from quantitative PCR. In some such embodiments, the difference is expressed as ΔCt . ΔCt may be calculated as $Ct[\text{target RNA}] - Ct[\text{endogenous control}]$ or $Ct[\text{endogenous control}] - Ct[\text{target RNA}]$. In certain embodiments, $\Delta Ct = Ct[\text{endogenous control}] - Ct[\text{marker}]$. In some embodiments, a threshold ΔCt value is set, above or below which bladder cancer is indicated. In some such embodiments, the ΔCt threshold is set as the ΔCt value below which 95% of normal samples are correctly characterized. In some such embodiments, a ΔCt value that is higher than the threshold ΔCt value is indicative of bladder cancer.

In some embodiments, linear discriminant analysis (LDA) is used, for example, to combine two or more of the markers into a single combined scale. In some such embodiments, a single threshold value is used for the markers included in the LDA.

In some embodiments, a control RNA is an exogenous control RNA. In some such embodiments, the exogenous control RNA is an Armored RNA®, which is protected by a bacteriophage coat. An exogenous control RNA may, in some embodiments, be used to determine if the detection assay reaction has failed, and therefore the results are not meaningful. For example, if an exogenous control RNA is not amplified in the assay reaction, then a negative result for the target RNAs is likely not meaningful because the levels reflect the reaction failing rather than the target RNA levels being low. Reaction failure can occur for any number of reasons, including, but not limited to, the presence of a reaction inhibitor in the sample (an “inhibitory sample”), compromised reagents, the presence of an RNase, etc. An exogenous RNA control may be added at any stage of the sample collection and analysis. For example, in some embodiments, the exogenous control RNA is added to the sample at the time preservative is added, is added to the sample when it is received by the diagnostic laboratory, is added to the sample immediately prior to analysis, or is added to the sample during analysis (as a nonlimiting example, during or after lysis of the urothelial cells but before addition of the amplification reagents).

In some embodiments, the level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, is compared to a reference level, e.g., from a confirmed bladder cancer. In some such embodiments, a similar level of a target RNA relative to the reference sample indicates bladder cancer.

In some embodiments, a level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, that is at least about 20%, 30%, 40%, 50%, 60%, 70%, 80%, or 90% greater than a normal level of the respective target RNA indicates the presence of bladder cancer. In some embodiments, a level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, that is at least about two-fold, at least about 3-fold, at least about 4-fold, at least about 5-fold, at least about 6-fold, at least about 7-fold, at least about 8-fold, at least about 9-fold, or at least about 10-fold greater than a normal level of the respective target RNA indicates the presence of bladder cancer.

In some embodiments, a control level of a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, is determined contemporaneously, such as in the same assay or batch of assays, as the level of the target RNA in a sample. In some embodiments, a control level of a target RNA is not determined contemporaneously as the level of the target

RNA in a sample. In some such embodiments, the control level has been determined previously.

In some embodiments, the level of an endogenous control and/or an exogenous control is determined contemporaneously, such as in the same assay or batch of assays, as the level of the target RNA in a sample. In some embodiments, an assay comprises reagents for determining the levels of CRH, IGF2, KRT20 and ANXA10 mRNA, and an endogenous control simultaneously in the same assay reaction. In some embodiments, an assay comprises reagents for determining the levels of CRH, IGF2, KRT20 and ANXA10 mRNA, and an exogenous control simultaneously in the same assay reaction. In some embodiments, an assay comprises reagents for determining the levels of CRH, IGF2, KRT20, ANXA10 mRNA, an endogenous control, and an exogenous control simultaneously in the same assay reaction. In some such embodiments, for example, an assay reaction comprises primer sets for amplifying each of CRH, IGF2, KRT20 and ANXA10 mRNAs, a primer set for amplifying an endogenous control and/or a primer set for amplifying an exogenous control, and detectably different labeled probes for detecting the amplification products (such as, for example, TaqMan® probes with detectably different dyes for each different amplicon to be detected).

In some embodiments, the level of a target RNA is not compared to a control level, for example, when it is known that the target RNA is present at very low levels, or not at all, in normal cells. In such embodiments, detection of a high level of the target RNA in a sample is indicative of bladder cancer.

6.2.3. Exemplary Sample Preparation

6.2.3.1. Exemplary Urine Preservatives

In some embodiments, a preservative is added to the urine sample. In some embodiments, the preservative is added within one hour, two hours, three hours, or six hours of the time the urine sample was collected (e.g., voided). In some embodiments, a preservative is added to the urine sample within one hour, two hours, three hours, or six hours before the sample is analyzed by the methods described herein.

In some embodiments, a preservative causes damage, such as lysis, of red blood cells and/or white blood cells, but does not damage urothelial cells. Red blood cells and/or white blood cells may be present in the urine as a result of a tumor and/or infection. In some such embodiments, adding the preservative allows for improved enrichment of the urothelial cells, for example by filtration. In some embodiments, a preservative lowers the pH of the urine sample and improves solubility of urine salts. In some such embodiments, the preservative facilitates passage of the salts through a filter in a filtration step. A desirable pH of preserved urine to be passed through a filter is between about 2.5 and 4. In some embodiments, a desirable pH of preserved urine is between about 2.7 and 3.7. In some embodiments, a desirable pH of preserved urine is between about 3 and 3.5. In some embodiments, a desirable pH of preserved urine is about 3.2.

In some embodiments a preservative is added such that the urine/preservative sample comprises 0.875M to 2.625M guanidine hydrochloride, 0.25% to 0.75% N-acetyl-L-cysteine, 6.25 to 18.75 mM sodium citrate, and 0.625% to 1.875% Tween-20, and has a pH of 3 to 3.5. In some embodiments a preservative is added such that the urine/preservative sample comprises about 1.75 M guanidine

hydrochloride, about 0.5% N-acetyl-L-cysteine, about 12.5 mM sodium citrate, and about 1.25% Tween-20, and has a pH of about 3.2.

A nonlimiting exemplary commercial preservative is PreservCyt (Hologic, Bedford, MA).

6.2.3.2. Exemplary Cell Enrichment

In some embodiments, urothelial cells are enriched by centrifugation. In some such embodiments, the cell pellet is resuspended in the supernatant and/or a preservative. Resuspension of the cell pellet can be used to adjust the concentration of cells in solution. The resuspended cell pellet may be used (for example, with lysis) in the methods described herein, or may be subject to an additional enrichment step, such as filtration.

In some embodiments, urothelial cells are enriched by filtration. Nonlimiting exemplary filter pore sizes that may be suitable for capturing urothelial cells include 0.8 μm , 2 μm , 8 μm , and 10 μm . In some embodiments, a filter pore size is selected that allows pass-through of red blood cells and/or white blood cells, while retaining most urothelial cells. In some embodiments, a filter is located within a GeneXpert cartridge designed for carrying out a bladder cancer diagnostic assay described herein.

6.2.3.3. Exemplary mRNA Preparation

Target RNA can be prepared by any appropriate method. Total RNA can be isolated by any method, including, but not limited to, the protocols set forth in Wilkinson, M. (1988) Nucl. Acids Res. 16(22): 10,933; and Wilkinson, M. (1988) Nucl. Acids Res. 16(22): 10934, or by using commercially-available kits or reagents, such as the TRIzol® reagent (Invitrogen), Total RNA Extraction Kit (iNtRON Biotechnology), Total RNA Purification Kit (Norgen Biotek Corp.), RNAqueous™ (Ambion), MagMAX™ (Ambion), RecoverAll™ (Ambion), RNeasy (Qiagen), etc.

In some embodiments, RNA levels are measured in a sample in which RNA has not first been purified from the cells. In some such embodiments, the cells are subject to a lysis step to release the RNA. Nonlimiting exemplary lysis methods include sonication (for example, for 2-15 seconds, 8-18 μm at 36 kHz); chemical lysis, for example, using a detergent; and various commercially available lysis reagents (such as RNeasy lysis buffer, Qiagen). In some embodiments, RNA levels are measured in a sample in which RNA has been isolated.

In some embodiments, RNA is modified before a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, is detected. In some embodiments, all of the RNA in the sample is modified. In some embodiments, just the particular target RNAs to be analyzed are modified, e.g., in a sequence-specific manner. In some embodiments, RNA is reverse transcribed. In some such embodiments, RNA is reverse transcribed using MMLV reverse transcriptase. Nonlimiting exemplary conditions for reverse transcribing RNA using MMLV reverse transcriptase include incubation from 5 to 20 minutes at 40° C. to 50° C.

When a target RNA is reverse transcribed, a DNA complement of the target RNA is formed. In some embodiments, the complement of a target RNA is detected rather than a target RNA itself (or a DNA copy of the RNA itself). Thus, when the methods discussed herein indicate that a target RNA is detected, or the level of a target RNA is determined, such detection or determination may be carried out on a complement of a target RNA instead of, or in addition to, the target

RNA itself. In some embodiments, when the complement of a target RNA is detected rather than the target RNA, a polynucleotide for detection is used that is complementary to the complement of the target RNA. In some such embodiments, a polynucleotide for detection comprises at least a portion that is identical in sequence to the target RNA, although it may contain thymidine in place of uridine, and/or comprise other modified nucleotides.

6.2.4. Exemplary Analytical Methods

As described above, methods are presented for detecting bladder cancer. The methods comprise detecting a panel of bladder cancer markers consisting of CRH, IGF2, KRT20 and ANXA10, and optionally including at least one endogenous control and/or at least one exogenous control. In some embodiments, detection of an elevated level of one of the four bladder cancer markers indicates the presence of bladder cancer. In some embodiments, detection of an elevated level of two, three, or all four of the bladder cancer markers indicates the presence of bladder cancer. In some embodiments, the bladder cancer is low grade bladder cancer. In some embodiments, the bladder cancer is high grade bladder cancer. In some embodiments, the bladder cancer is a recurrence of bladder cancer in a patient with a history of bladder cancer.

Any analytical procedure capable of permitting specific and quantifiable (or semi-quantifiable) detection of a target RNA, such as CRH, IGF2, KRT20 and ANXA10 mRNAs, may be used in the methods herein presented. Such analytical procedures include, but are not limited to, RT-PCR methods, and other methods known to those skilled in the art.

In some embodiments, the method of detecting a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA, comprises amplifying cDNA complementary to the target RNA. Such amplification can be accomplished by any method. Exemplary methods include, but are not limited to, real time PCR, endpoint PCR, and amplification using T7 polymerase from a T7 promoter annealed to a cDNA, such as provided by the SenseAmp Plus™ Kit available at Implen, Germany.

When a target RNA or a cDNA complementary to a target RNA is amplified, in some embodiments, a DNA amplicon of the target RNA is formed. A DNA amplicon may be single stranded or double-stranded. In some embodiments, when a DNA amplicon is single-stranded, the sequence of the DNA amplicon is related to the target RNA in either the sense or antisense orientation. In some embodiments, a DNA amplicon of a target RNA is detected rather than the target RNA itself. Thus, when the methods discussed herein indicate that a target RNA is detected, or the level of a target RNA is determined, such detection or determination may be carried out on a DNA amplicon of the target RNA instead of, or in addition to, the target RNA itself. In some embodiments, when the DNA amplicon of the target RNA is detected rather than the target RNA, a polynucleotide for detection is used that is complementary to the complement of the target RNA. In some embodiments, when the DNA amplicon of the target RNA is detected rather than the target RNA, a polynucleotide for detection is used that is complementary to the target RNA. Further, in some embodiments, multiple polynucleotides for detection may be used, and some polynucleotides may be complementary to the target RNA and some polynucleotides may be complementary to the complement of the target RNA.

In some embodiments, the method of detecting one or more target RNAs, such as CRH, IGF2, KRT20 or ANXA10 mRNA, comprises RT-PCR, as described below. In some embodiments, detecting one or more target RNAs comprises real-time monitoring of an RT-PCR reaction, which can be accomplished by any method. Such methods include, but are not limited to, the use of TaqMan®, Molecular beacon, or Scorpion probes (i.e., energy transfer (ET) probes, such as FRET probes) and the use of intercalating dyes, such as SYBR green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc.

Nonlimiting exemplary conditions for amplifying cDNA that has been reverse transcribed from the target RNAs are as follows. An exemplary cycle comprises an initial denaturation at 90° C. to 100° C. for 2 to 5 minutes, followed by cycling that comprises denaturation at 90° C. to 100° C. for 1 to 10 seconds, annealing at 60° C. to 70° C. for 10 to 30 seconds, and extension at 60° C. to 75° C. for 10 to 40 seconds. In some embodiments, for the first cycle following the initial denaturation step, the cycle denaturation step is omitted. In some embodiments, Taq polymerase is used for amplification. In some embodiments, the cycle is carried out at least 10 times, at least 15 times, at least 20 times, at least 25 times, at least 30 times, at least 35 times, or at least 45 times. In some such embodiments, Taq is used with a hot start function. In some embodiments, the amplification reaction occurs in a GeneXpert cartridge, and amplification of the four bladder cancer marker target RNAs occurs in the same reaction. In some embodiments, detection of CRH, IGF2, KRT20 and ANXA10 mRNAs occurs in less than 3 hours, less than 2.5 hours, or less than 2 hours, from initial denaturation through the last extension.

In some embodiments, detection of a target RNA comprises forming a complex comprising a polynucleotide that is complementary to a target RNA or to a complement thereof, and a nucleic acid selected from the target RNA, a DNA amplicon of the target RNA, and a complement of the target RNA. Thus, in some embodiments, the polynucleotide forms a complex with a target RNA. In some embodiments, the polynucleotide forms a complex with a complement of the target RNA, such as a cDNA that has been reverse transcribed from the target RNA. In some embodiments, the polynucleotide forms a complex with a DNA amplicon of the target RNA. When a double-stranded DNA amplicon is part of a complex, as used herein, the complex may comprise one or both strands of the DNA amplicon. Thus, in some embodiments, a complex comprises only one strand of the DNA amplicon. In some embodiments, a complex is a triplex and comprises the polynucleotide and both strands of the DNA amplicon. In some embodiments, the complex is formed by hybridization between the polynucleotide and the target RNA, complement of the target RNA, or DNA amplicon of the target RNA. The polynucleotide, in some embodiments, is a primer or probe.

In some embodiments, a method comprises detecting the complex. In some embodiments, the complex does not have to be associated at the time of detection. That is, in some embodiments, a complex is formed, the complex is then dissociated or destroyed in some manner, and components from the complex are detected. An example of such a system is a TaqMan® assay. In some embodiments, when the polynucleotide is a primer, detection of the complex may comprise amplification of the target RNA, a complement of the target RNA, or a DNA amplicon of a target RNA.

In some embodiments the analytical method used for detecting at least one target RNA in the methods set forth herein includes real-time quantitative RT-PCR. In some

embodiments, the analytical method used for detecting at least one target RNA includes the use of a TaqMan® probe. The assay uses energy transfer (“ET”), such as fluorescence resonance energy transfer (“FRET”), to detect and quantitate the synthesized PCR product. Typically, the TaqMan® probe comprises a fluorescent dye molecule coupled to the 5'-end and a quencher molecule coupled to the 3'-end, such that the dye and the quencher are in close proximity, allowing the quencher to suppress the fluorescence signal of the dye via FRET. When the polymerase replicates the chimeric amplicon template to which the TaqMan® probe is bound, the 5'-nuclease of the polymerase cleaves the probe, decoupling the dye and the quencher so that the dye signal (such as fluorescence) is detected. Signal (such as fluorescence) increases with each RT-PCR cycle proportionally to the amount of probe that is cleaved.

In some embodiments, quantitation of the results of real-time RT-PCR assays is done by constructing a standard curve from a nucleic acid of known concentration and then extrapolating quantitative information for target RNAs of unknown concentration. In some embodiments, the nucleic acid used for generating a standard curve is an RNA (for example, an endogenous control, or an exogenous control). In some embodiments, the nucleic acid used for generating a standard curve is a purified double-stranded plasmid DNA or a single-stranded DNA generated in vitro.

In some embodiments, where the amplification efficiencies of the target nucleic acids and the endogenous reference are approximately equal, quantitation is accomplished by the comparative Ct (cycle threshold, e.g., the number of PCR cycles required for the fluorescence signal to rise above background) method. Ct values are inversely proportional to the amount of nucleic acid target in a sample. In some embodiments, Ct values of a target RNA can be compared with a control or calibrator, such as an exogenous control RNA. In some embodiments, the Ct values of the exogenous control and the target RNA are normalized to an appropriate endogenous control. Nonlimiting exemplary endogenous controls are discussed herein.

In some embodiments, a threshold Ct (or a “cutoff Ct”) value for a target RNA, below which bladder cancer is indicated, has previously been determined. In such embodiments, a control sample may not be assayed concurrently with the test sample. In some embodiments, as discussed herein, a ΔCt threshold value is determined, above which bladder cancer is indicated, has previously been determined.

In addition to the TaqMan® assays, other real-time RT-PCR chemistries useful for detecting and quantitating PCR products in the methods presented herein include, but are not limited to, Molecular Beacons, Scorpion probes and intercalating dyes, such as SYBR Green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc., which are discussed below.

In various embodiments, real-time RT-PCR detection is utilized to detect, in a single multiplex reaction, all four bladder cancer markers of the panel described herein, and optionally, at least one endogenous control and/or at least one exogenous control. In some multiplex embodiments, a plurality of probes, such as TaqMan® probes, each specific for a different RNA target, is used. In some embodiments, each target RNA-specific probe is spectrally distinguishable from the other probes used in the same multiplex reaction.

In some embodiments, quantitation of real-time RT PCR products is accomplished using a dye that binds to double-stranded DNA products, such as SYBR Green, EvaGreen, thiazole orange, YO-PRO, TO-PRO, etc. In some embodiments, the assay is the QuantiTect SYBR Green PCR assay from Qiagen. In this assay, total RNA is first isolated from

a sample. Total RNA is subsequently poly-adenylated at the 3'-end and reverse transcribed using a universal primer with poly-dT at the 5'-end. In some embodiments, a single reverse transcription reaction is sufficient to assay multiple target RNAs. Real-time RT-PCR is then accomplished using target RNA-specific primers and an miScript Universal Primer, which comprises a poly-dT sequence at the 5'-end. SYBR Green dye binds non-specifically to double-stranded DNA and upon excitation, emits light. In some embodiments, buffer conditions that promote highly-specific annealing of primers to the PCR template (e.g., available in the QuantiTect SYBR Green PCR Kit from Qiagen) can be used to avoid the formation of non-specific DNA duplexes and primer dimers that will bind SYBR Green and negatively affect quantitation. Thus, as PCR product accumulates, the signal from SYBR Green increases, allowing quantitation of specific products.

Real-time RT-PCR is performed using any RT-PCR instrumentation available in the art. Typically, instrumentation used in real-time RT-PCR data collection and analysis comprises a thermal cycler, optics for fluorescence excitation and emission collection, and optionally a computer and data acquisition and analysis software.

In some embodiments, the analytical method used in the methods described herein is a DASL® (cDNA-mediated Annealing, Selection, Extension, and Ligation) Assay. In some embodiments, total RNA is isolated from a sample to be analyzed by any method. Total RNA may then be polyadenylated (>18 A residues are added to the 3'-ends of the RNAs in the reaction mixture). The RNA is reverse transcribed using a biotin-labeled DNA primer that comprises from the 5' to the 3' end, a sequence that includes a PCR primer site and a poly-dT region that binds to the poly-dA tail of the sample RNA. The resulting biotinylated cDNA transcripts are then hybridized to a solid support via a biotin-streptavidin interaction and contacted with one or more target RNA-specific polynucleotides. The target RNA-specific polynucleotides comprise, from the 5'-end to the 3'-end, a region comprising a PCR primer site, region comprising an address sequence, and a target RNA-specific sequence.

In some DASL® embodiments, the target RNA-specific sequence comprises at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19 contiguous nucleotides having a sequence that is the same as, or complementary to, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19 contiguous nucleotides of a bladder cancer marker target RNA, an endogenous control RNA, or an exogenous control RNA.

After hybridization, the target RNA-specific polynucleotide is extended, and the extended products are then eluted from the immobilized cDNA array. A second PCR reaction using a fluorescently-labeled universal primer generates a fluorescently-labeled DNA comprising the target RNA-specific sequence. The labeled PCR products are then hybridized to a microbead array for detection and quantitation.

In some embodiments, the analytical method used for detecting and quantifying the levels of the at least one target RNA in the methods described herein is a bead-based flow cytometric assay. See Lu J. et al. (2005) *Nature* 435:834-838, which is incorporated herein by reference in its entirety. An example of a bead-based flow cytometric assay is the xMAP® technology of Luminex, Inc. (See luminexcorp.com/technology/index.html). In some embodiments, total RNA is isolated from a sample and is then labeled with

biotin. The labeled RNA is then hybridized to target RNA-specific capture probes (e.g., FlexmiR™ products sold by Luminex, Inc. at luminexcorp.com/products/assays/index.html) that are covalently bound to microbeads, each of which is labeled with 2 dyes having different fluorescence intensities. A streptavidin-bound reporter molecule (e.g., streptavidin-phycoerythrin, also known as "SAPE") is attached to the captured target RNA and the unique signal of each bead is read using flow cytometry. In some embodiments, the RNA sample is first polyadenylated, and is subsequently labeled with a biotinylated 3DNA™ dendrimer (i.e., a multiple-arm DNA with numerous biotin molecules bound thereto), using a bridging polynucleotide that is complementary to the 3'-end of the poly-dA tail of the sample RNA and to the 5'-end of the polynucleotide attached to the biotinylated dendrimer. The streptavidin-bound reporter molecule is then attached to the biotinylated dendrimer before analysis by flow cytometry. In some embodiments, biotin-labeled RNA is first exposed to SAPE, and the RNA/SAPE complex is subsequently exposed to an anti-phycoerythrin antibody attached to a DNA dendrimer, which can be bound to as many as 900 biotin molecules. This allows multiple SAPE molecules to bind to the biotinylated dendrimer through the biotin-streptavidin interaction, thus increasing the signal from the assay.

In some embodiments, the analytical method used for detecting and quantifying the levels of the at least one target RNA in the methods described herein is by gel electrophoresis and detection with labeled probes (e.g., probes labeled with a radioactive or chemiluminescent label), such as by Northern blotting. In some embodiments, total RNA is isolated from the sample, and then is size-separated by SDS polyacrylamide gel electrophoresis. The separated RNA is then blotted onto a membrane and hybridized to radiolabeled complementary probes. In some embodiments, exemplary probes contain one or more affinity-enhancing nucleotide analogs as discussed below, such as locked nucleic acid ("LNA") analogs, which contain a bicyclic sugar moiety instead of deoxyribose or ribose sugars. See, e.g., Várallyay, E. et al. (2008) *Nature Protocols* 3(2):190-196, which is incorporated herein by reference in its entirety.

In some embodiments, detection and quantification of one or more target RNAs is accomplished using microfluidic devices and single-molecule detection. In some embodiments, target RNAs in a sample of isolated total RNA are hybridized to two probes, one which is complementary to nucleic acids at the 5'-end of the target RNA and the second which is complementary to the 3'-end of the target RNA. Each probe comprises, in some embodiments, one or more affinity-enhancing nucleotide analogs, such as LNA nucleotide analogs and each is labeled with a different fluorescent dye having different fluorescence emission spectra (i.e., detectably different dyes). The sample is then flowed through a microfluidic capillary in which multiple lasers excite the fluorescent probes, such that a unique coincident burst of photons identifies a particular target RNA, and the number of particular unique coincident bursts of photons can be counted to quantify the amount of the target RNA in the sample. In some alternative embodiments, a target RNA-specific probe can be labeled with 3 or more distinct labels selected from, e.g., fluorophores, electron spin labels, etc., and then hybridized to an RNA sample.

Optionally, the sample RNA is modified before hybridization. The target RNA/probe duplex is then passed through channels in a microfluidic device and that comprise detectors that record the unique signal of the 3 labels. In this way, individual molecules are detected by their unique signal and

counted. See U.S. Pat. Nos. 7,402,422 and 7,351,538 to Fuchs et al., U.S. Genomics, Inc., each of which is incorporated herein by reference in its entirety.

6.2.5. Exemplary Automation and Systems

In some embodiments, gene expression is detected using an automated sample handling and/or analysis platform. In some embodiments, commercially available automated analysis platforms are utilized. For example, in some embodiments, the GeneXpert® system (Cepheid, Sunnyvale, CA) is utilized.

The present invention is illustrated for use with the GeneXpert system. Exemplary sample preparation and analysis methods are described below. However, the present invention is not limited to a particular detection method or analysis platform. One of skill in the art recognizes that any number of platforms and methods may be utilized.

The GeneXpert® utilizes a self-contained, single use cartridge. Sample extraction, amplification, and detection may all carried out within this self-contained "laboratory in a cartridge." (See e.g., U.S. Pat. Nos. 5,958,349, 6,403,037, 6,440,725, 6,783,736, 6,818,185; each of which is herein incorporated by reference in its entirety.)

Components of the cartridge include, but are not limited to, processing chambers containing reagents, filters, and capture technologies useful to extract, purify, and amplify target nucleic acids. A valve enables fluid transfer from chamber to chamber and contain nucleic acids lysis and filtration components. An optical window enables real-time optical detection. A reaction tube enables very rapid thermal cycling.

In some embodiments, the GenXpert® system includes a plurality of modules for scalability. Each module includes a plurality of cartridges, along with sample handling and analysis components.

In some embodiments, the GeneXpert® sample preparation method utilizes filtration in order to capture and concentrate cells from urine. In some embodiments, a filter pore size of 0.8 μm is utilized. This size facilitates capture of all cells in urine. In other embodiments, pore sizes of 0.5 to 10 μm , 0.5 to 5 μm , 0.8 to 10 μm , 0.8 to 5 μm , 0.8 to 2 μm , 2 to 5 μm , 2 to 10 μm , 2 to 8 μm , 5 to 8 μm , or 5 to 10 μm are utilized. Certain filters (such as 5 μm , 8 μm , and 10 μm) allow the removal of most red and white blood cells from the sample while capturing the larger urothelial cells, which are the assay target cells. In some embodiments, this sample preparation method improves assay specificity by removing white blood cells that may be present due to infection or inflammation. In some instances, sample preparation methods such as centrifugation of whole urine followed by RNA isolation from the urine pellet do not allow for removal of white blood cells. In some embodiments, the efficiency of cell capture by filtration is higher compared to centrifugation, and may provide more consistent results.

After the cells from the urine are captured on the filter, in some embodiments, they are washed and then lysed using sonication (2-15 seconds, 8-16 μm at 36 kHz). The cell lysate is then collected and used to reconstitute the RT-PCR reagents, which are present in the cartridge as lyophilized particles.

In some embodiments, RT-PCR is used to amplify and analyze the presence or expression levels of the bladder cancer markers. In some embodiments, the reverse transcription uses MMLV RT enzyme and an incubation of 5 to 20 minutes at 40° C. to 50° C. In some embodiments, the PCR uses Taq polymerase with hot start function, such as AptaTaq

(Roche). In some embodiments, the initial denaturation is at 90° C. to 100° C. for 2 to 5 minutes; the cycling denaturation temperature is 90° C. to 100° C. for 1 to 10 seconds; the cycling anneal temperature is 60° C. to 70° C. for 10 to 30 seconds; and the cycling extend temperature is 60° C. to 75° C. for 10 to 40 seconds; and up to 50 cycles are performed.

The present invention is not limited to particular primer and/or probe sequences. Exemplary amplification primers and detection probes are described in the Examples.

In some embodiments, an off-line centrifugation is used to improve assay results with samples with low cellular content. The sample, with or without the preservative added, is centrifuged and the supernatant removed. The pellet is then resuspended in a smaller volume of either supernatant or the preservative. The resuspended pellet is then added to a GeneXpert® cartridge as previously described.

6.2.6. Exemplary Data Analysis

In some embodiments, a computer-based analysis program is used to translate the raw data generated by the detection assay (e.g., the expression level of the bladder cancer markers described herein) into data of predictive value for a clinician. The clinician can access the predictive data using any suitable means. Thus, in some embodiments, the present invention provides the further benefit that the clinician, who is not likely to be trained in genetics or molecular biology, need not understand the raw data. The data is presented directly to the clinician in its most useful form. The clinician is then able to immediately utilize the information in order to optimize the care of the subject.

The present invention contemplates any method capable of receiving, processing, and transmitting the information to and from laboratories conducting the assays, information provides, medical personal, and subjects. For example, in some embodiments of the present invention, a sample (e.g., a biopsy or a serum or urine sample) is obtained from a subject and submitted to a profiling service (e.g., clinical lab at a medical facility, genomic profiling business, etc.), located in any part of the world (e.g., in a country different than the country where the subject resides or where the information is ultimately used) to generate raw data. Where the sample comprises a tissue or other biological sample, the subject may visit a medical center to have the sample obtained and sent to the profiling center, or subjects may collect the sample themselves (e.g., a urine sample) and directly send it to a profiling center. Where the sample comprises previously determined biological information, the information may be directly sent to the profiling service by the subject (e.g., an information card containing the information may be scanned by a computer and the data transmitted to a computer of the profiling center using an electronic communication systems). Once received by the profiling service, the sample is processed and a profile is produced (i.e., expression data), specific for the diagnostic or prognostic information desired for the subject.

The profile data is then prepared in a format suitable for interpretation by a treating clinician. For example, rather than providing raw expression data, the prepared format may represent a diagnosis or risk assessment (e.g., expression level of the bladder cancer markers described herein or diagnosis of bladder cancer) for the subject, along with recommendations for particular treatment options. The data may be displayed to the clinician by any suitable method. For example, in some embodiments, the profiling service

generates a report that can be printed for the clinician (e.g., at the point of care) or displayed to the clinician on a computer monitor.

In some embodiments, the information is first analyzed at the point of care or at a regional facility. The raw data is then sent to a central processing facility for further analysis and/or to convert the raw data to information useful for a clinician or patient. The central processing facility provides the advantage of privacy (all data is stored in a central facility with uniform security protocols), speed, and uniformity of data analysis. The central processing facility can then control the fate of the data following treatment of the subject. For example, using an electronic communication system, the central facility can provide data to the clinician, the subject, or researchers.

In some embodiments, the subject is able to directly access the data using the electronic communication system. The subject may choose further intervention or counseling based on the results. In some embodiments, the data is used for research use. For example, the data may be used to further optimize the inclusion or elimination of markers as useful indicators of a particular condition or stage of disease or as a companion diagnostic to determine a treatment course of action.

6.2.7. Exemplary Polynucleotides

In some embodiments, polynucleotides are provided. In some embodiments, synthetic polynucleotides are provided. Synthetic polynucleotides, as used herein, refer to polynucleotides that have been synthesized in vitro either chemically or enzymatically. Chemical synthesis of polynucleotides includes, but is not limited to, synthesis using polynucleotide synthesizers, such as OligoPilot (GE Healthcare), ABI 3900 DNA Synthesizer (Applied Biosystems), and the like. Enzymatic synthesis includes, but is not limited, to producing polynucleotides by enzymatic amplification, e.g., PCR. A polynucleotide may comprise one or more nucleotide analogs (i.e., modified nucleotides) discussed herein.

In some embodiments, a polynucleotide is provided that comprises a region that is identical to, or complementary to, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of a sequence selected from CRH, IGF2, KRT20, and ANXA10 mRNA. In some embodiments, a polynucleotide is provided that comprises a region that is identical to, or complementary to, a span of 6 to 100, 8 to 100, 8 to 75, 8 to 50, 8 to 40, or 8 to 30 contiguous nucleotides of a sequence selected from CRH, IGF2, KRT20, and ANXA10 mRNA. Nonlimiting exemplary polynucleotides are shown in Tables 1 and 6.

In various embodiments, a polynucleotide comprises fewer than 500, fewer than 300, fewer than 200, fewer than 150, fewer than 100, fewer than 75, fewer than 50, fewer than 40, or fewer than 30 nucleotides. In various embodiments, a polynucleotide is between 6 and 200, between 8 and 200, between 8 and 150, between 8 and 100, between 8 and 75, between 8 and 50, between 8 and 40, or between 8 and 30 nucleotides long.

In some embodiments, the polynucleotide is a primer. In some embodiments, the primer is labeled with a detectable moiety. In some embodiments, a primer is not labeled. A primer, as used herein, is a polynucleotide that is capable of specifically hybridizing to a target RNA or to a cDNA

reverse transcribed from the target RNA or to an amplicon that has been amplified from a target RNA or a cDNA (collectively referred to as "template"), and, in the presence of the template, a polymerase and suitable buffers and reagents, can be extended to form a primer extension product.

In some embodiments, the polynucleotide is a probe. In some embodiments, the probe is labeled with a detectable moiety. A detectable moiety, as used herein, includes both directly detectable moieties, such as fluorescent dyes, and indirectly detectable moieties, such as members of binding pairs. When the detectable moiety is a member of a binding pair, in some embodiments, the probe can be detectable by incubating the probe with a detectable label bound to the second member of the binding pair. In some embodiments, a probe is not labeled, such as when a probe is a capture probe, e.g., on a microarray or bead. In some embodiments, a probe is not extendable, e.g., by a polymerase. In other embodiments, a probe is extendable.

In some embodiments, the polynucleotide is a FRET probe that in some embodiments is labeled at the 5'-end with a fluorescent dye (donor) and at the 3'-end with a quencher (acceptor), a chemical group that absorbs (i.e., suppresses) fluorescence emission from the dye when the groups are in close proximity (i.e., attached to the same probe). In other embodiments, the dye and quencher are not at the ends of the FRET probe. Thus, in some embodiments, the emission spectrum of the dye should overlap considerably with the absorption spectrum of the quencher.

6.2.7.1. Exemplary Polynucleotide Modifications

In some embodiments, the methods of detecting at least one target RNA described herein employ one or more polynucleotides that have been modified, such as polynucleotides comprising one or more affinity-enhancing nucleotide analogs. Modified polynucleotides useful in the methods described herein include primers for reverse transcription, PCR amplification primers, and probes. In some embodiments, the incorporation of affinity-enhancing nucleotides increases the binding affinity and specificity of a polynucleotide for its target nucleic acid as compared to polynucleotides that contain only deoxyribonucleotides, and allows for the use of shorter polynucleotides or for shorter regions of complementarity between the polynucleotide and the target nucleic acid.

In some embodiments, affinity-enhancing nucleotide analogs include nucleotides comprising one or more base modifications, sugar modifications and/or backbone modifications.

In some embodiments, modified bases for use in affinity-enhancing nucleotide analogs include 5-methylcytosine, isocytosine, pseudoisocytosine, 5-bromouracil, 5-propynyluracil, 6-aminopurine, 2-aminopurine, inosine, diaminopurine, 2-chloro-6-aminopurine, xanthine and hypoxanthine.

In some embodiments, affinity-enhancing nucleotide analogs include nucleotides having modified sugars such as 2'-substituted sugars, such as 2'-O-alkyl-ribose sugars, 2'-amino-deoxyribose sugars, 2'-fluoro-deoxyribose sugars, 2'-fluoro-arabinose sugars, and 2'-O-methoxyethyl-ribose (2'MOE) sugars. In some embodiments, modified sugars are arabinose sugars, or d-arabino-hexitol sugars.

In some embodiments, affinity-enhancing nucleotide analogs include backbone modifications such as the use of peptide nucleic acids (PNA; e.g., an oligomer including nucleobases linked together by an amino acid backbone). Other backbone modifications include phosphorothioate

linkages, phosphodiester modified nucleic acids, combinations of phosphodiester and phosphorothioate nucleic acid, methylphosphonate, alkylphosphonates, phosphate esters, alkylphosphonothioates, phosphoramidates, carbamates, carbonates, phosphate triesters, acetamides, carboxymethyl esters, methylphosphorothioate, phosphorodithioate, p-ethoxy, and combinations thereof.

In some embodiments, a polynucleotide includes at least one affinity-enhancing nucleotide analog that has a modified base, at least nucleotide (which may be the same nucleotide) that has a modified sugar, and/or at least one internucleotide linkage that is non-naturally occurring.

In some embodiments, an affinity-enhancing nucleotide analog contains a locked nucleic acid ("LNA") sugar, which is a bicyclic sugar. In some embodiments, a polynucleotide for use in the methods described herein comprises one or more nucleotides having an LNA sugar. In some embodiments, a polynucleotide contains one or more regions consisting of nucleotides with LNA sugars. In other embodiments, a polynucleotide contains nucleotides with LNA sugars interspersed with deoxyribonucleotides. See, e.g., Frieden, M. et al. (2008) *Curr. Pharm. Des.* 14(11):1138-1142.

6.2.7.2. Exemplary Primers

In some embodiments, a primer is provided. In some embodiments, a primer is identical to, or complementary to, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of a sequence selected from CRH, IGF2, KRT20, and ANXA10 mRNA. In some embodiments, a primer is provided that comprises a region that is identical to, or complementary to, a span of 6 to 100, 8 to 100, 8 to 75, 8 to 50, 8 to 40, or 8 to 30 contiguous nucleotides of a sequence selected from CRH, IGF2, KRT20, and ANXA10 mRNA. Nonlimiting exemplary primers are shown in Tables 1 and 6. In some embodiments, a primer may also comprise portions or regions that are not identical or complementary to the target RNA. In some embodiments, a region of a primer that is identical or complementary to a target RNA is contiguous, such that any region of a primer that is not identical or complementary to the target RNA does not disrupt the identical or complementary region.

In some embodiments, a primer comprises a portion that is identically present in a target RNA. In some such embodiments, a primer that comprises a region that is identically present in the target RNA is capable of selectively hybridizing to a cDNA that has been reverse transcribed from the RNA, or to an amplicon that has been produced by amplification of the target RNA or cDNA. In some embodiments, the primer is complementary to a sufficient portion of the cDNA or amplicon such that it selectively hybridizes to the cDNA or amplicon under the conditions of the particular assay being used.

As used herein, "selectively hybridize" means that a polynucleotide, such as a primer or probe, will hybridize to a particular nucleic acid in a sample with at least 5-fold greater affinity than it will hybridize to another nucleic acid present in the same sample that has a different nucleotide sequence in the hybridizing region. Exemplary hybridization conditions are discussed herein, for example, in the context of a reverse transcription reaction or a PCR amplification reaction. In some embodiments, a polynucleotide will

hybridize to a particular nucleic acid in a sample with at least 10-fold greater affinity than it will hybridize to another nucleic acid present in the same sample that has a different nucleotide sequence in the hybridizing region.

In some embodiments, a primer is used to reverse transcribe a target RNA, for example, as discussed herein. In some embodiments, a primer is used to amplify a target RNA or a cDNA reverse transcribed therefrom. Such amplification, in some embodiments, is quantitative PCR, for example, as discussed herein. In some embodiments, a primer comprises a detectable moiety.

In some embodiments, primer pairs are provided. Such primer pairs are designed to amplify a portion of a target mRNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA, or an endogenous control RNA, or an exogenous control RNA. In some embodiments, a primer pair is designed to produce an amplicon that is 50 to 1500 nucleotides long, 50 to 1000 nucleotides long, 50 to 750 nucleotides long, 50 to 500 nucleotides long, 50 to 400 nucleotides long, 50 to 300 nucleotides long, 50 to 200 nucleotides long, 50 to 150 nucleotides long, or 50 to 100 nucleotides long. Nonlimiting exemplary primer pairs are shown in Tables 1 and 6. In some embodiments, a primer pair is designed that spans an intron in the genomic sequence so that the mRNA, without the intron, is more preferably amplified than the genomic sequence. By "spans an intron" is meant that one primer of the primer pair is complementary to a sequence in the mRNA or a cDNA reverse transcribed from the mRNA that is at least partially located 5' to an intron in the genomic sequence and one primer of the primer pair is complementary to a sequence in the mRNA or a cDNA reverse transcribed from the mRNA that is at least partially located 3' to the same intron in the genomic sequence. In some embodiments, one primer of the primer pair is complementary to a sequence in the mRNA or a cDNA reverse transcribed from the mRNA that is located 5' to an intron in the genomic sequence and one primer of the primer pair is complementary to a sequence in the mRNA or a cDNA reverse transcribed from the mRNA that is located 3' to the same intron in the genomic sequence. In some embodiments, one of the primers in the primer pair may be complementary to a sequence in the mRNA or a cDNA reverse transcribed from the mRNA that is spliced together when the intron is removed such that the contiguous complementary sequence is not found in the genomic sequence. A primer pair comprising such a primer is still considered to span an intron.

6.2.7.3. Exemplary Probes

In various embodiments, methods of detecting the presence of bladder cancer comprise hybridizing nucleic acids of a sample with a probe. In some embodiments, the probe comprises a portion that is complementary to a target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA. In some embodiments, the probe comprises a portion that is identically present in the target RNA. In some such embodiments, a probe that is complementary to a target RNA is complementary to a sufficient portion of the target RNA such that it selectively hybridizes to the target RNA under the conditions of the particular assay being used. In some embodiments, a probe that is complementary to a target RNA comprises a region that is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of the target RNA, such as

CRH, IGF2, KRT20 or ANXA10 mRNA. Nonlimiting exemplary probes are shown in Tables 1 and 6. A probe that is complementary to a target RNA may also comprise portions or regions that are not complementary to the target RNA. In some embodiments, a region of a probe that is complementary to a target RNA is contiguous, such that any region of a probe that is not complementary to the target RNA does not disrupt the complementary region.

In some embodiments, the probe comprises a portion that is identically present in the target RNA, such as CRH, IGF2, KRT20 or ANXA10 mRNA. In some such embodiments, a probe that comprises a region that is identically present in the target RNA is capable of selectively hybridizing to a cDNA that has been reverse transcribed from the RNA, or to an amplicon that has been produced by amplification of the target RNA or cDNA. In some embodiments, the probe is complementary to a sufficient portion of the cDNA or amplicon such that it selectively hybridizes to the cDNA or amplicon under the conditions of the particular assay being used. In some embodiments, a probe that is complementary to a cDNA or amplicon comprises a region that is complementary to at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, at least 25, at least 26, at least 27, at least 28, at least 29, or at least 30 contiguous nucleotides of the cDNA or amplicon. A probe that is complementary to a cDNA or amplicon may also comprise portions or regions that are not complementary to the cDNA or amplicon. In some embodiments, a region of a probe that is complementary to a cDNA or amplicon is contiguous, such that any region of a probe that is not complementary to the cDNA or amplicon does not disrupt the complementary region.

In some embodiments, the method of detectably quantifying one or more target RNAs comprises: (a) reverse transcribing a target RNA to produce a cDNA that is complementary to the target RNA; (b) amplifying the cDNA from (a); and (c) detecting the amount of a target RNA using real time RT-PCR and a detection probe (which may be simultaneous with the amplification step (b)).

As described above, in some embodiments, real time RT-PCR detection may be performed using a FRET probe, which includes, but is not limited to, a TaqMan® probe, a Molecular beacon probe and a Scorpion probe. In some embodiments, the real time RT-PCR detection and quantification is performed with a TaqMan® probe, i.e., a linear probe that typically has a fluorescent dye covalently bound at one end of the DNA and a quencher molecule covalently bound at the other end of the DNA. The FRET probe comprises a sequence that is complementary to a region of the cDNA such that, when the FRET probe is hybridized to the cDNA, the dye fluorescence is quenched, and when the probe is digested during amplification of the cDNA, the dye is released from the probe and produces a fluorescence signal. In such embodiments, the amount of target RNA in the sample is proportional to the amount of fluorescence measured during cDNA amplification.

The TaqMan® probe typically comprises a region of contiguous nucleotides having a sequence that is complementary to a region of a target RNA or its complementary cDNA that is reverse transcribed from the target RNA template (i.e., the sequence of the probe region is complementary to or identically present in the target RNA to be detected) such that the probe is specifically hybridizable to the resulting PCR amplicon. In some embodiments, the probe comprises a region of at least 6 contiguous nucleotides having a sequence that is fully complementary to or iden-

tically present in a region of a cDNA that has been reverse transcribed from a target RNA template, such as comprising a region of at least 8 contiguous nucleotides, at least 10 contiguous nucleotides, at least 12 contiguous nucleotides, at least 14 contiguous nucleotides, or at least 16 contiguous nucleotides having a sequence that is complementary to or identically present in a region of a cDNA reverse transcribed from a target RNA to be detected.

In some embodiments, the region of the cDNA that has a sequence that is complementary to the TaqMan® probe sequence is at or near the center of the cDNA molecule. In some embodiments, there are independently at least 2 nucleotides, such as at least 3 nucleotides, such as at least 4 nucleotides, such as at least 5 nucleotides of the cDNA at the 5'-end and at the 3'-end of the region of complementarity.

In some embodiments, Molecular Beacons can be used to detect and quantitate PCR products. Like TaqMan® probes, Molecular Beacons use FRET to detect and quantitate a PCR product via a probe having a fluorescent dye and a quencher attached at the ends of the probe. Unlike TaqMan® probes, Molecular Beacons remain intact during the PCR cycles. Molecular Beacon probes form a stem-loop structure when free in solution, thereby allowing the dye and quencher to be in close enough proximity to cause fluorescence quenching. When the Molecular Beacon hybridizes to a target, the stem-loop structure is abolished so that the dye and the quencher become separated in space and the dye fluoresces. Molecular Beacons are available, e.g., from Gene Link™ (see genelink.com/newsite/products/mbintro.asp).

In some embodiments, Scorpion probes can be used as both sequence-specific primers and for PCR product detection and quantitation. Like Molecular Beacons, Scorpion probes form a stem-loop structure when not hybridized to a target nucleic acid. However, unlike Molecular Beacons, a Scorpion probe achieves both sequence-specific priming and PCR product detection. A fluorescent dye molecule is attached to the 5'-end of the Scorpion probe, and a quencher is attached to the 3'-end. The 3' portion of the probe is complementary to the extension product of the PCR primer, and this complementary portion is linked to the 5'-end of the probe by a non-amplifiable moiety. After the Scorpion primer is extended, the target-specific sequence of the probe binds to its complement within the extended amplicon, thus opening up the stem-loop structure and allowing the dye on the 5'-end to fluoresce and generate a signal. Scorpion probes are available from, e.g., Premier Biosoft International (see premierbiosoft.com/tech_notes/Scorpion.html).

In some embodiments, labels that can be used on the FRET probes include colorimetric and fluorescent dyes such as Alexa Fluor dyes, BODIPY dyes, such as BODIPY FL; Cascade Blue; Cascade Yellow; coumarin and its derivatives, such as 7-amino-4-methylcoumarin, aminocoumarin and hydroxycoumarin; cyanine dyes, such as Cy3 and Cy5; eosins and erythrosins; fluorescein and its derivatives, such as fluorescein isothiocyanate; macrocyclic chelates of lanthanide ions, such as Quantum Dye™; Marina Blue; Oregon Green; rhodamine dyes, such as rhodamine red, tetramethylrhodamine and rhodamine 6G; Texas Red; fluorescent energy transfer dyes, such as thiazole orange-ethidium heterodimer; and, TOTAB.

Specific examples of dyes include, but are not limited to, those identified above and the following: Alexa Fluor 350, Alexa Fluor 405, Alexa Fluor 430, Alexa Fluor 488, Alexa Fluor 500, Alexa Fluor 514, Alexa Fluor 532, Alexa Fluor 546, Alexa Fluor 555, Alexa Fluor 568, Alexa Fluor 594, Alexa Fluor 610, Alexa Fluor 633, Alexa Fluor 647, Alexa Fluor 660, Alexa Fluor 680, Alexa Fluor 700, and, Alexa

Fl

uor 750; amine-reactive BODIPY dyes, such as BODIPY 493/503, BODIPY 530/550, BODIPY 558/568, BODIPY 564/570, BODIPY 576/589, BODIPY 581/591, BODIPY 630/650, BODIPY 650/655, BODIPY FL, BODIPY R6G, BODIPY TMR, and, BODIPY-TR; Cy3, Cy5, 6-FAM, Fluorescein Isothiocyanate, HEX, 6-JOE, Oregon Green 488, Oregon Green 500, Oregon Green 514, Pacific Blue, REG, Rhodamine Green, Rhodamine Red, Renographin, ROX, SYPRO, TAMRA, 2',4',5',7'-Tetrabromosulfonefluorescein, and TET.

Examples of dye/quencher pairs (i.e., donor/acceptor pairs) include, but are not limited to, fluorescein/tetramethylrhodamine; IAEDANS/fluorescein; EDANS/dabcyl; fluorescein/fluorescein; BODIPY FL/BODIPY FL; fluorescein/QSY 7 or QSY 9 dyes. When the donor and acceptor are the same, FRET may be detected, in some embodiments, by fluorescence depolarization. Certain specific examples of dye/quencher pairs (i.e., donor/acceptor pairs) include, but are not limited to, Alexa Fluor 350/Alexa Fluor 488; Alexa Fluor 488/Alexa Fluor 546; Alexa Fluor 488/Alexa Fluor 555; Alexa Fluor 488/Alexa Fluor 568; Alexa Fluor 488/Alexa Fluor 594; Alexa Fluor 488/Alexa Fluor 647; Alexa Fluor 546/Alexa Fluor 568; Alexa Fluor 546/Alexa Fluor 594; Alexa Fluor 546/Alexa Fluor 647; Alexa Fluor 555/Alexa Fluor 594; Alexa Fluor 555/Alexa Fluor 647; Alexa Fluor 568/Alexa Fluor 647; Alexa Fluor 594/Alexa Fluor 647; Alexa Fluor 350/QSY35; Alexa Fluor 350/dabcyl; Alexa Fluor 488/QSY 35; Alexa Fluor 488/dabcyl; Alexa Fluor 488/QSY 7 or QSY 9; Alexa Fluor 555/QSY 7 or QSY 9; Alexa Fluor 568/QSY 7 or QSY 9; Alexa Fluor 568/QSY 21; Alexa Fluor 594/QSY 21; and Alexa Fluor 647/QSY 21. In some instances, the same quencher may be used for multiple dyes, for example, a broad spectrum quencher, such as an Iowa Black® quencher (Integrated DNA Technologies, Coralville, IA) or a Black Hole Quencher™ (BHQ™; Sigma-Aldrich, St. Louis, MO).

In some embodiments, for example, in a multiplex reaction in which two or more moieties (such as amplicons) are detected simultaneously, each probe comprises a detectably different dye such that the dyes may be distinguished when detected simultaneously in the same reaction. One skilled in the art can select a set of detectably different dyes for use in a multiplex reaction.

Specific examples of fluorescently labeled ribonucleotides useful in the preparation of RT-PCR probes for use in some embodiments of the methods described herein are available from Molecular Probes (Invitrogen), and these include, Alexa Fluor 488-5-UTP, Fluorescein-12-UTP, BODIPY FL-14-UTP, BODIPY TMR-14-UTP, Tetramethylrhodamine-6-UTP, Alexa Fluor 546-14-UTP, Texas Red-5-UTP, and BODIPY TR-14-UTP. Other fluorescent ribonucleotides are available from Amersham Biosciences (GE Healthcare), such as Cy3-UTP and Cy5-UTP.

Examples of fluorescently labeled deoxyribonucleotides useful in the preparation of RT-PCR probes for use in the methods described herein include Dinitrophenyl (DNP)-1'-dUTP, Cascade Blue-7-dUTP, Alexa Fluor 488-5-dUTP, Fluorescein-12-dUTP, Oregon Green 488-5-dUTP, BODIPY FL-14-dUTP, Rhodamine Green-5-dUTP, Alexa Fluor 532-5-dUTP, BODIPY TMR-14-dUTP, Tetramethylrhodamine-6-dUTP, Alexa Fluor 546-14-dUTP, Alexa Fluor 568-5-dUTP, Texas Red-12-dUTP, Texas Red-5-dUTP, BODIPY TR-14-dUTP, Alexa Fluor 594-5-dUTP, BODIPY 630/650-14-dUTP, BODIPY 650/665-14-dUTP; Alexa Fluor 488-7-OBEA-dCTP, Alexa Fluor 546-16-OBEA-dCTP, Alexa Fluor 594-7-OBEA-dCTP, Alexa Fluor 647-

12-OBEA-dCTP. Fluorescently labeled nucleotides are commercially available and can be purchased from, e.g., Invitrogen.

In some embodiments, dyes and other moieties, such as quenchers, are introduced into polynucleotide used in the methods described herein, such as FRET probes, via modified nucleotides. A "modified nucleotide" refers to a nucleotide that has been chemically modified, but still functions as a nucleotide. In some embodiments, the modified nucleotide has a chemical moiety, such as a dye or quencher, covalently attached, and can be introduced into a polynucleotide, for example, by way of solid phase synthesis of the polynucleotide. In other embodiments, the modified nucleotide includes one or more reactive groups that can react with a dye or quencher before, during, or after incorporation of the modified nucleotide into the nucleic acid. In specific embodiments, the modified nucleotide is an amine-modified nucleotide, i.e., a nucleotide that has been modified to have a reactive amine group. In some embodiments, the modified nucleotide comprises a modified base moiety, such as uridine, adenosine, guanosine, and/or cytosine. In specific embodiments, the amine-modified nucleotide is selected from 5-(3-aminoallyl)-UTP; 8-[(4-amino)butyl]-amino-ATP and 8-[(6-amino)butyl]-amino-ATP; N6-(4-amino)butyl-ATP, N6-(6-amino)butyl-ATP, N4-[2,2-oxy-bis(ethylamine)]-CTP; N6-(6-Amino)hexyl-ATP; 8-[(6-Amino)hexyl]-amino-ATP; 5-propargylamino-CTP, 5-propargylamino-UTP. In some embodiments, nucleotides with different nucleobase moieties are similarly modified, for example, 5-(3-aminoallyl)-GTP instead of 5-(3-aminoallyl)-UTP. Many amine modified nucleotides are commercially available from, e.g., Applied Biosystems, Sigma, Jena Bioscience and TriLink.

Exemplary detectable moieties also include, but are not limited to, members of binding pairs. In some such embodiments, a first member of a binding pair is linked to a polynucleotide. The second member of the binding pair is linked to a detectable label, such as a fluorescent label. When the polynucleotide linked to the first member of the binding pair is incubated with the second member of the binding pair linked to the detectable label, the first and second members of the binding pair associate and the polynucleotide can be detected. Exemplary binding pairs include, but are not limited to, biotin and streptavidin, antibodies and antigens, etc.

In some embodiments, multiple target RNAs are detected in a single multiplex reaction. In some such embodiments, each probe that is targeted to a unique cDNA is spectrally distinguishable when released from the probe. Thus, each target RNA is detected by a unique fluorescence signal.

One skilled in the art can select a suitable detection method for a selected assay, e.g., a real-time RT-PCR assay. The selected detection method need not be a method described above, and may be any method.

6.3. Exemplary Compositions and Kits

In another aspect, compositions are provided. In some embodiments, compositions are provided for use in the methods described herein.

In some embodiments, compositions are provided that comprise at least one target RNA-specific primer. The term "target RNA-specific primer" encompasses primers that have a region of contiguous nucleotides having a sequence that is (i) identically present in a target RNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA, or (ii) complementary to the sequence of a region of contiguous nucleotides found

in a target RNA, such CRH, IGF2, KRT20, or ANXA10 mRNA. In some embodiments, a composition is provided that comprises at least one pair of target RNA-specific primers. The term “pair of target RNA-specific primers” encompasses pairs of primers that are suitable for amplifying a defined region of a target RNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA. A pair of target RNA-specific primers typically comprises a first primer that comprises a sequence that is identical to the sequence of a region of a target RNA (although the primer will typically comprise DNA or modified nucleosides rather than RNA) and a second primer that comprises a sequence that is complementary to a region of a target RNA. A pair of primers is typically suitable for amplifying a region of a target mRNA that is 50 to 1500 nucleotides long, 50 to 1000 nucleotides long, 50 to 750 nucleotides long, 50 to 500 nucleotides long, 50 to 400 nucleotides long, 50 to 300 nucleotides long, 50 to 200 nucleotides long, 50 to 150 nucleotides long, or 50 to 100 nucleotides long. Nonlimiting exemplary primers, and pairs of primers, are shown in Tables 1 and 6.

In some embodiments, a composition comprises four pairs of target RNA-specific primers, one pair for amplifying each of CRH, IGF2, KRT20, and ANXA10 mRNA. In some embodiments, a composition additionally comprises a pair of target RNA-specific primers for amplifying an endogenous control RNA and/or one pair of target RNA-specific primers for amplifying an exogenous control RNA.

In some embodiments, a composition comprises at least one target RNA-specific probe. The term “target RNA-specific probe” encompasses probes that have a region of contiguous nucleotides having a sequence that is (i) identically present in a target RNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA, or (ii) complementary to the sequence of a region of contiguous nucleotides found in a target RNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA. Nonlimiting exemplary target-specific probes are shown in Tables 1 and 6.

In some embodiments, a composition (including a composition described above that comprises four or more pairs of target RNA-specific primers) comprises four probes, one probe for detecting each of CRH, IGF2, KRT20, and ANXA10 mRNA. In some embodiments, a composition additionally comprises a probe for detecting an endogenous control RNA and/or a probe for detecting an exogenous control RNA.

In some embodiments, a composition is an aqueous composition. In some embodiments, the aqueous composition comprises a buffering component, such as phosphate, tris, HEPES, etc., and/or additional components, as discussed below. In some embodiments, a composition is dry, for example, lyophilized, and suitable for reconstitution by addition of fluid. A dry composition may include one or more buffering components and/or additional components.

In some embodiments, a composition further comprises one or more additional components. Additional components include, but are not limited to, salts, such as NaCl, KCl, and MgCl₂; polymerases, including thermostable polymerases such as Taq; dNTPs; reverse transcriptases, such as MMLV reverse transcriptase; RNase inhibitors; bovine serum albumin (BSA) and the like; reducing agents, such as β-mercaptoethanol; EDTA and the like; etc. One skilled in the art can select suitable composition components depending on the intended use of the composition.

In some embodiments, compositions are provided that comprise at least one polynucleotide for detecting at least one target RNA. In some embodiments, the polynucleotide is used as a primer for a reverse transcriptase reaction. In

some embodiments, the polynucleotide is used as a primer for amplification. In some embodiments, the polynucleotide is used as a primer for RT-PCR. In some embodiments, the polynucleotide is used as a probe for detecting at least one target RNA. In some embodiments, the polynucleotide is detectably labeled. In some embodiments, the polynucleotide is a FRET probe. In some embodiments, the polynucleotide is a TaqMan® probe, a Molecular Beacon, or a Scorpion probe.

In some embodiments, a composition comprises at least one FRET probe having a sequence that is identically present in, or complementary to a region of, CRH, IGF2, KRT20, or ANXA10 mRNA. In some embodiments, a FRET probe is labeled with a donor/acceptor pair such that when the probe is digested during the PCR reaction, it produces a unique fluorescence emission that is associated with a specific target RNA. In some embodiments, when a composition comprises multiple FRET probes, each probe is labeled with a different donor/acceptor pair such that when the probe is digested during the PCR reaction, each one produces a unique fluorescence emission that is associated with a specific probe sequence and/or target RNA. In some embodiments, the sequence of the FRET probe is complementary to a target region of a target RNA. In other embodiments, the FRET probe has a sequence that comprises one or more base mismatches when compared to the sequence of the best-aligned target region of a target RNA.

In some embodiments, a composition comprises a FRET probe consisting of at least 8, at least 9, at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 nucleotides, wherein at least a portion of the sequence is identically present in, or complementary to a region of, CRH, IGF2, KRT20, or ANXA10 mRNA. In some embodiments, at least 8, at least 9, at least 10, at least 11, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 nucleotides of the FRET probe are identically present in, or complementary to a region of, CRH, IGF2, KRT20, or ANXA10 mRNA. In some embodiments, the FRET probe has a sequence with one, two or three base mismatches when compared to the sequence or complement of small CRH, IGF2, KRT20, or ANXA10 mRNA.

In some embodiments, a kit comprises a polynucleotide discussed above. In some embodiments, a kit comprises at least one primer and/or probe discussed above. In some embodiments, a kit comprises at least one polymerase, such as a thermostable polymerase. In some embodiments, a kit comprises dNTPs. In some embodiments, kits for use in the real time RT-PCR methods described herein comprise one or more target RNA-specific FRET probes and/or one or more primers for reverse transcription of target RNAs and/or one or more primers for amplification of target RNAs or cDNAs reverse transcribed therefrom.

In some embodiments, one or more of the primers and/or probes is “linear”. A “linear” primer refers to a polynucleotide that is a single stranded molecule, and typically does not comprise a short region of, for example, at least 3, 4 or 5 contiguous nucleotides, which are complementary to another region within the same polynucleotide such that the primer forms an internal duplex. In some embodiments, the primers for use in reverse transcription comprise a region of at least 4, such as at least 5, such as at least 6, such as at least 7 or more contiguous nucleotides at the 3'-end that has a sequence that is complementary to region of at least 4, such

as at least 5, such as at least 6, such as at least 7 or more contiguous nucleotides at the 5'-end of a target RNA.

In some embodiments, a kit comprises one or more pairs of linear primers (a "forward primer" and a "reverse primer") for amplification of a cDNA reverse transcribed from a target RNA, such as CRH, IGF2, KRT20, or ANXA10 mRNA. Accordingly, in some embodiments, a first primer comprises a region of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides having a sequence that is identical to the sequence of a region of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides at a first location in the mRNA. Furthermore, in some embodiments, a second primer comprises a region of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides having a sequence that is complementary to the sequence of a region of at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 16, at least 17, at least 18, at least 19, at least 20, at least 21, at least 22, at least 23, at least 24, or at least 25 contiguous nucleotides at a second location in the mRNA, such that a PCR reaction using the two primers results in an amplicon extending from the first location of the mRNA to the second location of the mRNA.

In some embodiments, the kit comprises at least two, at least three, or at least four sets of primers, each of which is for amplification of a cDNA that is reverse transcribed from a different target RNA, including CRH, IGF2, KRT20, and ANXA10 mRNA. In some embodiments, the kit further comprises at least one set of primers for amplifying a control RNA, such as an endogenous control and/or an exogenous control.

In some embodiments, probes and/or primers for use in the compositions described herein comprise deoxyribo-

nucleotides. In some embodiments, probes and/or primers for use in the compositions described herein comprise deoxyribonucleotides and one or more nucleotide analogs, such as LNA analogs or other duplex-stabilizing nucleotide analogs described above. In some embodiments, probes and/or primers for use in the compositions described herein comprise all nucleotide analogs. In some embodiments, the probes and/or primers comprise one or more duplex-stabilizing nucleotide analogs, such as LNA analogs, in the region of complementarity.

In some embodiments, the kits for use in real time RT-PCR methods described herein further comprise reagents for use in the reverse transcription and amplification reactions. In some embodiments, the kits comprise enzymes such as reverse transcriptase, and a heat stable DNA polymerase, such as Taq polymerase. In some embodiments, the kits further comprise deoxyribonucleotide triphosphates (dNTP) for use in reverse transcription and amplification. In further embodiments, the kits comprise buffers optimized for specific hybridization of the probes and primers.

The following examples are for illustration purposes only, and are not meant to be limiting in any way.

7. EXAMPLES

7.1. Example 1: Detection of High Grade and Low Grade Bladder Cancer

More than 30 mRNA markers and 20 microRNA markers were evaluated in both bladder tissue and urine samples to determine the most accurate panel for detection of bladder cancer. Based on results from over 200 urine samples using eight markers and the GeneXpert system (Cepheid, Sunnyvale, CA), a panel consisting of CRH, IGF2, KRT20, and ANXA10 mRNA markers (with or without at least one endogenous and/or at least one exogenous control) was selected.

Various reaction compositions were designed for use in the GeneXpert® system, for detecting various combinations of CRH, IGF2, KRT20, and ANXA10 mRNA. Table 1 shows the sequences of the primers and probes used to detect each of the target RNAs by quantitative RT-PCR in the various reaction compositions.

TABLE 1

Primer and probe sequences					
oligo name	target	sequence	SEQ ID NO	Reagent ("TSR")	Formul.
ABLa3a4 PrmrFwd4	ABL	GATCAACACTGCTTCTGATGGCAA	8	CL3, CL4, CL1	
ABLa3a4 PrmrRev1	ABL	CCACCGTTGAATGATGATGAACCAA	9	CL3, CL4, CL1	
ABLa3a4 Probe 1	ABL	F4-CCTCCGAGAGCCGCTTCAAC-Q4	10	CL3	
ABL probe F6	ABL	F6-CCTCCGAGAGCCGCTTCAAC-Q6	11	CL4	
ABL probe F1	ABL	F1-CCTCCGAGAGCCGC(T-dabsyl) TCAAC-Q1	12	CL1	
KRT20 For	KRT20	TTGAAGAGCTGCGAAGTCAGAT	13	CL3, CL4	
KRT20 Rev	KRT20	TGAAGTCCTCAGCAGCCAGTT	14	CL3, CL4	
KRT20 Probe (F3)	KRT20	F3-TCAACTGCAAAATGCTCGGTGTGTCC-Q3	15	CL3, CL4	
IGF2 For_4	IGF2	CGCGGCTTCTACTTTCAGCAG	16	CL3, CL4	
IGF2 Rev_4	IGF2	GCGGAAACAGCACTCCTCAA	17	CL3, CL4	
IGF2 Probe_2	IGF2	F5-TGTGAGCCGTCGAGCCGTG-Q5	18	CL3, CL4	

TABLE 1-continued

Primer and probe sequences					
oligo name	target	sequence	SEQ ID NO	Reagent	Formul.
CRH_For	CRH	ACCCGGCTCACCTGCGAA	19	CL3, CL4, CL1	
CRH_Rev	CRH	GGACTCCCGCGGACACAA	20	CL3, CL4, CL1	
CRH_probe 3	CRH	F2-TCCTGGGAAGCGAGTGCCCTAA-Q2	21	CL3, CL1	
CRH_probe_F1	CRH	F1-CCTGGGAAGCGAG(T-Dabsyl)GCCCTAA-Q1	22	CL4	
Armored RNA [®] Fwd	exogenous control	GGCTATTCTCTCTTGGCAGAT	23	CL1	
Armored RNA [®] Rev	exogenous control	TGCTTGAGCTCCAGTCCCTAAG	24	CL1	
Armored RNA [®] _Probe	exogenous control	F6-AGCCGAGAAGGCGGAGTCTGGC-Q6	25	CL1	
ANXA10-FW	ANXA10	GTGAAACAAGTTTATGCAATCGATCAA	26	CL1	
ANXA10-RV3	ANXA10	GATTGAAATTGGGAGCTGGGAA	27	CL1	
ANXA10-F3	ANXA10	F3-TCATCCCTGAGGTTAACAATTACCATCAA-Q3	28	CL1	

F1 through F6 are detectably different dyes that can be detected and distinguished simultaneously in a multiplex reaction, and Q1 to Q6 are quenchers (in the present example, Q2, Q4, Q5, and Q6 are the same quencher).

The final primer and probe compositions of three different reaction compositions are shown in Table 2.

For each sample to be tested, 5 mL of voided urine was added to 5 mL preservative (3.5M guanidine HCl, 1% N-acetyl-L-cysteine, 25 mM sodium citrate, and 2.5% Tween-20, pH 3.2), preferably within 1 hour of sample collection. The preserved samples were transported on ice and stored at 4° C. Clinical information for each sample was

TABLE 2

Primers and probes in TSR CL3, CL4, and CL1					
Target	Label	Purpose	Final conc. forw. primer	Final conc. rev primer	Final conc. probe
TSR CL3					
ABL	F4	Normalization (endogenous control)	400 nM	400 nM	150 nM
KRT20	F3	Bladder cancer marker	400 nM	400 nM	75 nM
IGF2	F5	Bladder cancer marker	400 nM	400 nM	200 nM
CRH	F2	Bladder cancer marker	400 nM	400 nM	200 nM
TSR CL4					
ABL	F6	Normalization (endogenous control)	400 nM	400 nM	400 nM
KRT20	F3	Bladder cancer marker	400 nM	400 nM	75 nM
IGF2	F5	Bladder cancer marker	400 nM	400 nM	300 nM
CRH	F1	Bladder cancer marker	400 nM	400 nM	600 nM
TSR CL1					
ABL	F1	Normalization (endogenous control)	400 nM	400 nM	600 nM
Armored RNA [®]	F6	Exogenous control	400 nM	400 nM	400 nM
ANXA10	F3	Bladder cancer marker	400 nM	400 nM	75 nM
CRH	F2	Bladder cancer marker	400 nM	400 nM	300 nM

Each reaction contained 50-90 mM KCl, 3-5 mM MgCl₂, 400-825 μM dNTPs, 20 mM Tris, pH 8.5, 0.01% sodium azide, and 0.9 units/μl of RNase inhibitor. MMLV reverse transcriptase (0.375 units/μl) and AptAq (0.25 units/μl; Roche) were used for reverse transcription and amplification, respectively. TSR CL1 included an Armored RNA[®] exogenous control (SEQ ID NO: 47; Asuragen, Austin, TX).

provided by the collection sites. The number of red blood cells per millilitre was determined by microscopic evaluation.

Prior to use, the preserved urine was inverted three times to mix. 1.2 mL of preserved urine was loaded into a GeneXpert cartridge for analysis. The cartridge contained a 0.8 μm filter to capture urothelial cells. The captured cells were washed and lysed using sonication (2-15 seconds, 8-16

µm at 36 kHz) within the cartridge. The lysate was then used to reconstitute the reagents used for real-time RT-PCR (described above). The reaction cycle used was: 10 minutes at 45° C., followed by 2 minutes at 95° C., and then 45 cycles of (a) 5 seconds at 95° C., 20 seconds at 60° C., and 20 seconds at 72° C., using a GeneXpert® cartridge in a GeneXpert® system. Delta Ct (ΔCt) was calculated as Ct (ABL)–Ct (marker). The ΔCt cutoff was set as the ΔCt that gave at least 95% specificity with samples from patients not expected to have bladder cancer (data not shown). A ΔCt above the ΔCt cutoff for any one of the markers was

considered a positive result, indicative of the presence of bladder cancer.

Some samples were also tested using UroVysion® (Abbott Laboratories, Abbott Park, IL). The results of that experiment are shown in Table 3 (high grade bladder cancer) and Table 4 (low grade bladder cancer). ΔCts above the threshold, indicating a positive result, are highlighted. Each of the three TSR lots, CL3, CL4, and CL1, detected 100% of high grade bladder cancer samples, as did UroVysion.

For low grade bladder cancer, the detection rate was 37% (7/19), compared to only 16% (3/19) for UroVysion®.

TABLE 3

Detection of high grade bladder cancer															
						history of	TSR lot CL3			TSR lot CL4			CIC TSR lot CL1		
Sample ID		stage	grade	UroVysion® Result	cytology	bladder cancer	CRH -10	KRT20 2.2	IGF2 -1	KRT20 4	IGF2 0.5	CRH -5	ANXA10 -0.5	CRH -3	GX result
67001	pTa	high	positive	suspicious	yes		-20	4.9	-20	4.8	1.1	-20	-1.6	-3.6	positive
67006	pT1	high	positive	positive	no		-0.7	4.4	-6.2	5.7	-2.8	-0.6	-20	2.2	positive
67009	pT2	high	positive	positive	no		-20	4.9	-0.4	4.9	1.5	-20	1.1	-20	positive
75211	pT2	high	positive	negative	no		-0.8	1.4	-4.1						positive
75216	pTa, CIC	high	positive	suspicious	yes		-3.5	2	-0.8	0.9	-0.6	-3.4			positive
75218	pTa	high	positive	negative	no		0.9	3.8	-0.9	3.8	-1.6	0.5			positive
75245	pTa	high	positive	NA	no		20	1.6	6.7	0.4	6.2	20	1.9	8.2	positive
75247	CIS	high	positive	atypical	no		-1.4	3.4	2.7	3.6	3.7	-3.8	-0.2	0.4	positive
75248	pT1/ CIS	high	positive	atypical	no		-20	4.3	0	4.8	1	-20	0.4	-5.1	positive
75249	pT1	high	positive	negative	no		-20	3.3	2.1	3.4	2.5	-20	3.7	-20	positive
75258	CIS, pTa	high	positive	atypical	no					4.2	-3.7	0.6	-2	1.9	positive
75246			positive	positive	no		-20	4.3	5.3	4.4	5.9	9.1	-1	3.4	positive

TABLE 4

Detection of low grade bladder cancer															
Sample ID		stage	grade	UroVysion ® Result	cytology	history of bladder cancer	TSR lot CL3			TSR lot CL4			CIC TSR lot CL1		
							CRH -10	KRT20 2.2	IGF2 -1	KRT20 4	IGF2 0.5	CRH -5	ANXA10 -0.5	CRH -3	GX result
67002	pTa	low	negative	atypical	yes	-20	3.9	-3.8	3.3	-1.5	-20	-1.7	-20	positive	
67003	pTa	low	negative	negative	no	-20	-20	-3.2	-20	-20	-20	-20	-20	negative	
67004	pTa	low	negative	negative	no	-2	4.7	-2.3	5.2	0.1	-1.7	2.4	2.2	positive	
67010	pTa	low	positive	atypical	no	-20	-5.3	-8.8	-3.8	-6.7	-20	-5	-20	negative	
67011	pTa	low	positive	atypical	no	-1.3	4.6	1.9	4	2.4	-2	-1.3	1	positive	
67018	pTa	low	negative	atypical	yes	-20	-20	-4.4	-20	-20	-20	-20	-20	negative	
67050	pTa	low	negative	atypical	yes	-20	-0.4	3.4	1.9	3.3	-20	-0.5	-20	positive	
67100	pTa	low	borderline	atypical	yes				-2.9	-2.7	-20	-20	-20	negative	
75161	pTa	low	negative	atypical	yes	-20	-3	-4.7	-3.6	-1.3	-20	-5.7	-20	negative	
75183	pTa	low	positive	negative	yes	-1.5	3.1	-2	3.4	-0.5	0.7	-4.1	2.3	positive	
75184	pTa	low	inconclusive	negative	yes	-20	-0.6	1.3	-1.5	1.8	-20	-20	-20	positive	
75185	pTa	low	negative	negative	no	-20	3	5	1.3	4.7	-20	1.2	-4.2	positive	
75191	pTa	low	negative	negative	yes	-20	-0.4	-3.4	1.1	-0.5	-20	-4.1	-20	negative	
75202	pTa	low	negative	negative	yes	-20	0.3	-2.3	-0.4	-1.5	-20	-2.4	-20	negative	
75236	pTa	low	negative	negative	no	-20	-5.7	-7.9	-20	-8.9	-20	-3.2	-20	negative	
75251	pTa	low	negative	negative	yes	-20	-3.6	-5.1	-1	-4.4	-20	-20	-20	negative	
75257	pTa	low	negative	negative	no				-2.1	-0.9	-20	-20	-20	negative	
75265	pTa	low	negative	negative	yes				1.4	-20	-20	-20	-20	negative	

47

A summary of the sensitivity for high grade bladder cancer and low grade bladder cancer, and the specificity in patients with a low risk of bladder cancer, is shown in Table 5 for each of the individual markers tested in Tables 3 and 4.

TABLE 5

Summary of sensitivity and specificity of individual markers			
Marker	Sensitivity, high grade bladder cancer	Sensitivity, low grade bladder cancer	Specificity, low risk of bladder cancer
CRH	15/30 (50%)	9/53 (17%)	220/221 (99%)
KRT20	11/21 (52%)	6/34 (18%)	144/145 (99%)
IGF2	13/21 (62%)	8/34 (24%)	144/145 (99%)

48

TABLE 5-continued

Summary of sensitivity and specificity of individual markers			
Marker	Sensitivity, high grade bladder cancer	Sensitivity, low grade bladder cancer	Specificity, low risk of bladder cancer
ANXA10	5/9 (56%)	2/19 (11%)	74/76 (97%)
4 marker combo	12/12 (100%)	7/19 (37%)	83/88 (94%)

Exemplary alternative primers and probes for detecting the four markers, KRT20, IGF2, CRH, and ANXA10, are shown in Table 6. Table 6 also shows an exemplary set of primers and probes for detecting an exogenous control and an endogenous control, ABL. The dyes and quenchers Q1 to Q6 may be the same. One skilled in the art could select a suitable set, for example, a set of detectably different dyes for use in a multiplex assay. The predicted amplicon length for each set of primers is also shown, as well as the length of any intervening intron(s) between the primer sites on the genomic copy of the target.

TABLE 6

Primer and probe sequences						
name	5' mod	sequence	3' mod	SEQ ID NO	amplicon length (bp)	intron length (bp)
KRT20 For_3		CGACTACAGTGCATATTACAGACAA		29	113	2142
KRT20 Rev_2		CAGCAGCCAGTTTAGCATTATCAA		30		
KRT20 Probe	F1	TCAACTGCAAAA (T-dabsyl) GCT CGGTGTGTCC	Q1	31		
IGF2 For_5		GGACCGCGGCTTCTACTTCA		32	95	1701
IGF2 Rev_5		CCAGGTCACAGCTGCGGAA		33		
IGF2 Probe_2_F4	F4	TGTGAGCCGTGCGAGCCGTG	Q4	34		
CRH_For 4		TGCGAAGCGCCTGGGAAGC		35	66	801
CRH_Rev		GGACTCCCGCGGACACAA		36		
CRH_probe_F2	F2	TGCCCCTAACATGCGGCTGCC	Q2	37		
ANXA10_For_3		TCAGCGCTGCAATGCACAA		38	122	22, 947
ANXA10_For_4		CTGCAATGCACAAAGGATGA		48	117	22, 947
ANXA10_Rev_5		GGCCAGCCATCACATCTTTGAA		39		
ANXA10_Probe_3	F3	TAGAGCATGTATGGCCGGACCT	Q3	40		
ABLa3a4 PrmrFwd4		GATCAACACTGCTTCTGATGGCAA		41	92	7666
ABLa3a4 PrmrRev1		CCACCGTTGAATGATGATGAACCAA		42		
ABL Probe F5	F5	CCTCCGAGAGCCGCTTCAAC	Q5	43		
Armored RNA ® Fwd		GGCTATTCTCCTCTGGCAGAT		44	101	NA
Armored RNA ® Rev		TGCTTGAGCTCCAGTCCCTAAG		45		
Armored RNA ® Probe	F6	AGCCGAGAAGCGGAGTCTGGC	Q6	46		

7.2. Example 2: Assay Sensitivity for Detecting Bladder Cancer Using Marker Panel KRT20, IGF2, CRH, and ANXA10 in a Larger Cohort

Urine samples were collected from subjects at seven different sites. Eligibility criteria for inclusion in the study included:

- 18 years or older;
- Documented informed consent as required by the reviewing IRB or HREC, and a signed Experimental Subjects Bill of Rights for patients in California;
- At least one of the following criteria:
 - A history or recurrence of bladder cancer;
 - A referral for cystoscopy evaluation due to micro- or gross-hematuria in urine;
 - A referral for urology evaluation, but no previous history of bladder cancer or clinical evidence of bladder cancer;
- Consent to provide at least 15 ml voided urine in addition to that required for standard of care;
- Consent to allow pathology results for any biopsy specimens taken during cystoscopy procedure and other medical records to be reported.

Exclusion criteria included only under 18 years of age and first voided urine. Patients currently or previously treated with Bacillus Calmette-Guerin (BCG) and patients currently or previously treated with intravesical therapy or transurethral resection of bladder or radiation therapy for bladder

filter to capture urothelial cells. The captured cells were washed and lysed using sonication (2-15 seconds, 8-16 μ m at 36 kHz) within the cartridge. The lysate was then used to reconstitute the reagents used for real-time RT-PCR (described above). The reaction cycle used was: 10 minutes at 45° C., followed by 2 minutes at 95° C., and then 45 cycles of (a) 5 seconds at 95° C., 20 seconds at 60° C., and 20 seconds at 72° C., using a GeneXpert® cartridge in a GeneXpert® system. For ANXA10, KRT20 and IGF2, delta Ct (Δ Ct) was calculated as Ct (ABL)–Ct (marker). The Δ Ct cutoff was set as the Δ Ct that gave high (>90%) specificity with samples from patients not expected to have bladder cancer (data not shown). A Δ Ct above the Δ Ct cutoff for any one of the markers was considered a positive result, indicative of the presence of bladder cancer. For CRH, Ct values were used instead of Δ Ct to determine positivity for the CRH marker. A CRH Ct value <45 was considered a positive result, indicative of the presence of bladder cancer. In addition to the four bladder cancer markers (KRT20, IGF2, CRH, and ANXA10), the GeneXpert® bladder cancer assay included two controls: primers and probe for detecting ABL mRNA in the samples, and primers and probe for detecting an Armored RNA® exogenous control RNA.

In the first analysis, 132 samples collected from patients who had positive cystoscopy results for bladder cancer were tested with the GeneXpert® bladder cancer assay. Sixty of those samples had also been tested using UroVysion®. Table 7 shows the results for those 132 samples.

TABLE 7

Assay sensitivity by bladder cancer stage and grade								
	Xpert Bladder				UroVysion			
	Invalid/			Sensitivity	inconclusive			Sensitivity
	POS	NEG	Error**		POS	NEG		
Stage:								
All	94	35	3	72.9%	29	26	5	52.7%
Ta, Grade Low	29	23	2	55.8%	5	18	4	21.7%
Ta, Grade High	20	2		90.9%	4	6	1	40.0%
T1	13	2		86.7%	6	1		85.7%
T2	11	2		84.6%	4	0		100.0%
T3	3	0		100.0%	1	0		100.0%
T4	2	0		100.0%				
CIS	11	1		91.7%	7	0		100.0%
UNK	5	5	1	50.0%	2	1		66.7%
Grade:								
All	94	35	3	72.9%	29	26	5	52.7%
Low Grade	33	28	2	54.1%	6	19	4	24.0%
High Grade	61	7	1	89.7%	23	7	1	76.7%

**Two invalid results were due to low ABL Ct, suggesting a low number of cells in the sample, and one of the invalid results was due to poor sample quality.

cancer were eligible for the study. In addition, repeat enrollment during the course of the study was also permitted.

Two of the collection sites provided the results of UroVysion® analysis on the urine samples. For each sample to be tested 15 mL of voided urine was added to 15 mL of preservative (3.5M guanidine HCl, 1% N-acetyl-L-cysteine, 25 mM sodium citrate, and 2.5% Tween-20, pH 3.2), preferably within 1 hour of sample collection. The preserved samples were transported on ice and stored at 4° C. Clinical information for each sample was provided by the collection sites.

Prior to use, the preserved urine was inverted three times to mix. 4 mL of preserved urine was loaded into a GeneXpert cartridge for analysis. The cartridge contained a 0.8 μ m

As shown in Table 7, for these samples, the GeneXpert® bladder cancer assay had a sensitivity of 54.1% for low grade bladder cancer and a sensitivity of 89.7% for high grade bladder cancer. In contrast, UroVysion® had a sensitivity of just 24% for low grade bladder cancer and a sensitivity of 76.6% for high grade bladder cancer. Further, the GeneXpert® bladder cancer assay was able to detect all grades and stages of bladder cancer.

The same data set was then divided according to three patient groups: (A) patients with a history of bladder cancer who were currently being monitored for recurrence of bladder cancer, (B) patients who had been treated with Bacillus Calmette-Guerin (BCG) within the three months prior to sample collection, and (C) patients who were

symptomatic for bladder cancer and had no prior history of bladder cancer. The results for those patient groups are shown in Table 8.

TABLE 8

Assay sensitivity by patient population								
Grade:	Xpert Bladder*				UroVysion			
	POS	NEG	Invalid/Error**	Sensitivity	POS	NEG	inconclusive	Sensitivity
Monitoring (Population A)								
All	50	20	1	71.4%	11	12		47.8%
Low Grade	21	18		53.8%	2	11	1	15.4%
High Grade	29	2	1	93.5%	9	1		90.0%
Treated with BCG in last 3 months (Population B)								
All	4	2		66.7%	1			100.0%
Low Grade								
High Grade	4	2		66.7%	1			100.0%
Symptomatic (Population C)								
All	40	13	2	75.5%	17	14	4	54.8%
Low Grade	12	10	2	54.5%	4	8	3	33.3%
High Grade	28	3		90.3%	13	6	1	68.4%

**Two invalid results were due to low ABL Ct, suggesting a low number of cells in the sample, and one of the invalid results was due to poor sample quality.

25

As shown in Table 8, the GeneXpert® bladder cancer assay had a similar sensitivity for low grade and high grade bladder cancer in patients being monitored for bladder cancer and in patients who were symptomatic of bladder cancer as in the patient group as a whole (see Table 7). In patients who had been treated with BCG within the last three months, the GeneXpert® bladder cancer assay had a sensitivity of 66.7%, although the sample size was too small (6 samples) to draw any conclusions from that result.

In order to have a direct comparison of the GeneXpert® bladder cancer assay and UroVysion®, a dataset was selected that included only samples that had been tested with both assays. Table 9 shows the results for that dataset, with the patients separated into two groups: (A&B) patients with a history of bladder cancer who were currently being monitored for recurrence of bladder cancer, combined with patients who had been treated with Bacillus Calmette-Guerin (BCG) within the three months prior to sample collection (these groups were combined because only one sample from a BCG-treated patient had been tested with both assays), and (C) patients who were symptomatic for bladder cancer and had no prior history of bladder cancer.

TABLE 9

Assay sensitivity for samples tested with both GeneXpert ® and UroVysion ®								
Grade:	Xpert Bladder				UroVysion			
	POS	NEG	Invalid/Error**	Sensitivity	POS	NEG	inconclusive	Sensitivity
Monitoring and BCG treated (Populations A&B)								
All	18	7	1	72.0%	12	13	1	48.0%
Low Grade	6	7	1	46.2%	2	12		14.3%
High Grade	12	0		100.0%	10	1	1	90.9%
Symptomatic (Population C)								
All	26	8	1	76.5%	17	14	4	54.8%
Low Grade	7	7	1	50.0%	4	8	3	33.3%
High Grade	19	1		95.0%	13	6	1	68.4%

As shown in FIG. 9, for those samples that have been tested with both GeneXpert® bladder cancer assay and UroVysion®, the GeneXpert® bladder cancer assay showed

greater sensitivity than UroVysion® for detecting low grade and high grade cancer in both patient groups.

Next, the data set was divided into samples that had been archived, meaning they were tested more than one week after collection (the samples ranged from 8 days old up to nine months old), and samples that were fresh, meaning they were tested within one week of collection. The results of that analysis are shown in Table 10.

TABLE 10

Assay sensitivity in archived and fresh samples					
Grade:	Xpert Bladder				
	n	POS	NEG	Invalid/Error**	Sensitivity
Archived Samples					
All	89	61	25	3	70.9%
Low Grade	46	23	21	2	52.3%
High Grade	43	38	4	1	90.5%

25

30

35

40

45

53

TABLE 10-continued

Assay sensitivity in archived and fresh samples					
Grade:	Xpert Bladder				Sensitivity
	n	POS	NEG	Invalid/Error**	
Fresh Samples					
All	43	33	10		76.7%
Low Grade	17	10	7		58.8%
High Grade	26	23	3		88.5%

**Two invalid results were due to low ABL Ct, suggesting a low number of cells in the sample, and one of the invalid results was due to poor sample quality.

As shown in Table 10, the GeneXpert® bladder cancer assay had a similar sensitivity for detecting low grade and high grade bladder cancer in fresh and archived samples.

7.3. Example 3: Assay Specificity for Detecting Bladder Cancer Using Marker Panel KRT20, IGF2, CRH, and ANXA10 in a Larger Cohort

In addition to the samples from patients with positive cystoscopy results for bladder cancer, urine samples were collected at the seven sites from patients with negative cystoscopy results for bladder cancer, but who were being monitored for recurrence of bladder cancer, had received BCG within the three months prior to sample collection, and who appeared to be symptomatic for bladder cancer but had no prior history of bladder cancer. In addition, urine samples were collected from patients with urology referrals for other suspected conditions, such as kidney stones. Finally, urine samples were collected from healthy individuals. Urine samples were preserved and analyzed using the GeneXpert® bladder cancer assay as described in Example 2.

For the samples from patients with negative cystoscopy results for bladder cancer, the assay results were divided according to the three patient groups: (A) patients with a history of bladder cancer who were currently being monitored for recurrence of bladder cancer, (B) patients who had been treated with Bacillus Calmette-Guerin (BCG) within the three months prior to sample collection, and (C) patients who were symptomatic for bladder cancer and had no prior

54

history of bladder cancer. The results for those patient groups are shown in Table 11.

TABLE 11

Assay specificity in cystoscopy negative patients by population Xpert Bladder				
	POS	NEG	Invalid/Error	Specificity
Monitoring (Population A)				
5	55	156	17	73.9%
BCG Treated (Population B)				
10	1	6	0	85.7%
Symptomatic (Population C)				
15	16	86	10	84.3%

As shown in Table 11, the GeneXpert® bladder cancer assay had a specificity of 73.9%, 85.7%, and 84.3% for patient groups (A), (B), and (C), respectively.

Next, the specificity of the GeneXpert® bladder cancer assay in patients who were suspected of having other urological conditions, but not bladder cancer, was determined. Seventy patient samples were collected in this category. The results with the GeneXpert® bladder cancer assay were 14 positives, 50 negatives, and 6 invalid results. The specificity of the GeneXpert® bladder cancer assay for this patient population was therefore 78.1%.

Finally, the specificity of the GeneXpert® bladder cancer assay in healthy individuals was determined. Fifty-five samples were collected in this category. The results were 4 positives and 51 negatives, indicating a specificity of 92.7% for this subject category.

All publications, patents, patent applications and other documents cited in this application are hereby incorporated by reference in their entireties for all purposes to the same extent as if each individual publication, patent, patent application or other document were individually indicated to be incorporated by reference for all purposes.

While various specific embodiments have been illustrated and described, it will be appreciated that changes can be made without departing from the spirit and scope of the invention(s).

TABLE OF CERTAIN SEQUENCES

SEQ ID NO	Description	Sequence
1	Human CRH mRNA	TCGTTCCCTTG GCAGGGCCCT ATGATTTATG CAGGAGCAGA GGCAGCACGC AATCGAGCTG TCAAGAGAGC GTCAGCTTAT TAGGCAAATG CTGCGTGGTT TTTGAAGAGG GTCGACACTA TAAATCCCA CTCCAGGCTC TGGAGTGGAG AAACTCAGAG ACCAAGTCCA TTGAGAGACT GAGGGGAAAG AGAGGAGAGA AAGAAAAAGA GAGTGGGAAC AGTAAAGAGA AAGGAAGACA ACCTCCAGAG AAAGCCCCCG GAGACGTCTC TCTGCAGAGA GCGGCAGCA CCGGCTCAC CTGCGAAGCG CCTGGGAAGC GAGTGCCCTT AACATGCGGC TGCCGCTGCT TGTGTCCGCG GGAGTCCTGC TGGTGGCTCT CCGCCCTGC CCGCATGCA GGGCGCTCCT GAGCCGCGGG CCGTCCCGG GAGCTCGGA GGCCTCGAG CACCCCTAGC CCTTGGATT CTTCAGCCG CCGCCGAGT CCGAGCAGCC CCAGCAGCCG CAGGCTCGGC CGGTCTGCT CCGCATGGGA GAGGAGTACT TCCTCCGCTT GGGGAACCTC AACAGAGCC CCGCCGCTCC CCTTTCGCCC GCCTCCTCGC TCCTCGCCGG AGGCAGCGGC AGCCGCCCTT CGCCGGAACA GGCGACCGCC AACTTTTTC GCGTGTGCT GCAGCAGCTG CTGCTGCCTC GGCGCTCGCT CGACAGCCCC GCGCTCTCG CCGAGCGCGG CGCTAGGAAT GCCCTCGCG GCCACAGGA GGCACCGGAG AGAGAAAGGC GGTCCGAGGA GCCTCCATC TCCCTGGATC TCACCTTCCA CCTCCTCCG GAAGTCTTGG AAATGCCAG GGCCGAGCAG TTAGCACAGC AAGCTCACAG CACAGGAAA TTCATGGAGA TTATTGGGAA ATAAACGGT GCGTTTGGCC AAAAAGAATC TGCAATTAGC ACAAAAAAAA TTTAAAAAAA TACAGTATTC TGTACCATAG CGCTGCTCTT ATGCCATTG TTTATTTTTA TATAGCTTGA AACATAGAGG

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		GAGAGAGGGA	GAGAGCCTAT	ACCCCTTACT	TAGCATGCAC
		CACGTGCAGC	AGCAACACAA	TGTTATTTCGT	TTTGTCTACG
		GTTTCCAGGT	GTTTATAGTG	GTGTTTAAAA	GAGAATGTAG
		AAACGTTTGT	TTTGAAAAAG	CAGACAGAAG	TCACTCAATT
		TGGTCTGAGC	CAAAGAGAAT	GCCATTCTCT	TGGGTGGGTA
		TGTAAGCTCT	TTGAAACAAC	TTTCTCTTGT	AAACGTTTCA
		ATCTTTCCAG	TCCTTGGTCA	GTTTGGTTGT	GTAAGAGAAT
		TATATTTTTA	ATAAAGTTG	CAAAGGTAAT	CATG
2	Human IGF2 mRNA, transcript variant 2	CCGCTAATGT	ACCATGCCCT	GGTGCTGGAA	AGTGCCTGAG
		CAGCGGCCTC	AGCACTACCA	AGTTGGCACA	AAGCTCCCCA
		GGCTCAGGGA	AACGAGTGGA	GGGGATGAGG	AGGTGAGGGG
		ATTTCAAGTTG	GCATTTGAGC	AGGTGCCATG	CTCAGCGGAG
		CCCATCTGTA	GGGGCCGTAT	TAACATGCAC	ACTCTAAAAG
		TTCTCCAGCC	TCAGCTTTGT	CCCTCTCCTC	CTCCACGTCA
		AGGGTCTGGA	CGCCACAGCC	AGGGCACCCT	CTGCTTTGGT
		AATATTGGCC	AGGCCCGCGG	ATCATCGTCC	AGGCAGTTTC
		CTTGGGCACC	AGTGACTCCC	CGGTCTCTTT	TATCCACTGT
		CGGGGACTGC	GCAGGGACTA	GAGTACAGGG	GCCGAAGAGT
		CTTGTTGGG	AGGAGGTGGA	TTCCAGCCCC	CAGCCCCAGG
		GCTGCCAGCT	CAGCCCCCTG	CCCAGCCTGC	CCCACAGCCT
		AGGCCAGAGA	GCCCAGTCCT	GAGGTGAGCT	GCTGTGCCT
		CGACCCAGC	GCTCCCAGAA	CTGAGGCTGG	CAGCCAGCCC
		CCCAACTGCG	AGGCAGAGAG	ACACCAATGG	GAATCCCAAT
		ATGCTGGTGC	TTCTCACCTT	CTTGGCCTTC	GCCTCGTGCT
		TTACCGCCCC	AGTGAGACCC	TGTGCGGCGG	GGAGCTGGTG
		AGTTCTGCTG	TGGGGACCGC	GGCTTCTACT	TCAGCAGGCC
		GTGAGCCGTC	GCAGCCGTGG	CATCGTTGAG	GAGTGCTGTT
		TGACCTGGCC	CTCCTGGAGA	CGTACTGTGC	TACCCCGGCC
		GGGACGTGTC	GACCCCTCCG	ACCGTGCTTC	CGGACAACTT
		CCCGTGGCA	AGTTCTTCCA	ATATGACACC	TGGAAGCAGT
		CCTGCGCAGG	GGCCTGCCTG	CCCTCCTGCG	TGCCCCGCCG
		TCGCGCAAGGA	GCTCGAGGCG	TTCAGGGAGG	CCAAACGTCA
		ATTGCTCTAC	CCACCCAAGA	CCCCGCCCAC	GGGGGCGCCC
		GGCCAGCAAT	CGGAAGTGAG	CAAACTGCC	GCAAGTCTGC
		CACCATCTTG	CAGCCTCCTC	CTGACCACGG	ACGTTTCCAT
		CCCGAAATC	TCTCGGTTCC	ACGTCCCCCT	GGGGCTTCTC
		CCCGTGCCC	CGCCTCCCCG	AAACAGGCTA	CTCTCCTCGG
		CGGGCTGAGG	AAGCAGACGA	GCATCTTCAA	ACATGTACAA
		CTTTAAACAC	CCTTCACATA	CCCTCCCCCC	AAATTATCCC
		CACACATAAA	AAATCAAAAC	ATTAACTAA	CCCCCTTCCC
		AACAACCCCTC	TTAAACTAA	TTGGCTTTT	AGAAACACCC
		CAGAAATTGG	CTTTAAAAAA	AACAACCACC	AAAAAAATC
		AAAAAAAAG	TATTAATAAC	GAATTGGCTG	AGAAACAATT
		AGGAATTGG	CACTCCCCAC	CCCCCTCTTT	CTCTTCTCCC
		AGTCAAATTG	GCCTGGACTT	GAGTCCCTGA	ACCAGCAAAG
		GACCCAGAA	ATCACAGGTG	GGCACGTCGC	TGCTACCGCC
		TCACGGGAAT	TTTCAGGGTA	AACCTGGCAT	CCGAAAATAG
		GACTGGCTCC	TCACTCCCTT	TTCCATCACT	AAAAATCACA
		AGGGACCCAG	TAAGACCAAA	GGAGGGGAGG	ACAGAGCATG
		TCCATGCAAA	TGAAATGTAA	TTGGCAGCAG	CCTCACCCCC
		TCTCAATTCC	CATCCTAAAA	AGCACTCATA	CTTTATGCAT
		ACACACACAC	AACACACAGC	ACACGCATGA	ACACAGCACA
		CAGCACACAC	ACAAACGCAC	AGCACACACA	GCACACAGAT
		CACACACACA	AACGCACAGC	ACACACACGC	ACACACATGC
		CACAAACGCA	CGGCACACAC	ACGCACACAC	ATGCACACAC
		CAAAACGCACA	GCACACACAA	ACGCACAGCA	CACACGCACA
		ACACGAGCAC	ACAGCACACA	AACGCACAGC	ACACGCACAC
		ACAGCACACA	CACTAGCACA	CAGCACACAC	ACAAAGACAC
		TGCACACACA	GCACACACAC	GCGAACACAG	CACACACGAA
		ACAGCACACA	CACAAACACA	GCACACACAT	GCACACAGCA
		CACGACACAC	ATGAACACAG	CACACAGCAC	ACACATGCAC
		CACGCATGCA	CAGCACACAT	GAACACAGCA	CACACACAAA
		ACACATGCAC	ACACAGCACA	CACACTCATG	CGCAGCACAT
		AGCTCACAGC	ACACAAACAC	GCAGCACACA	CGTTGCACAC
		ACCTGCACAC	ACACATGCGC	ACACACACGC	ACACCCCCAC
		GAAAAACAATA	AGCATATCTA	AGCAACTACG	ATATCTGTAT
		AAAGTCCCGC	TAAGATTCTC	CAATGTTTTT	ATGGTCTGAG
		GTTCCCATCT	CCACTGCCCC	TCGGCCCTGT	CTGTGCCCTG
		GAGGGGGCTC	AGATGGTGCG	GCCTGAGTGT	GCGGCCGCGC
		TACACCCGTA	GGGTGGGCGG	GGTGTGTCCC	AGGCCTAATT
		ACCATGACAG	AGATGCCCTT	GTGAGGCTGG	CCTCCTTGGC
		ACGGCCCCCG	CAGCGTGAGC	CACGATGCTC	CCCATACCCC

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		GATACACCTT	ACTTACTGTG	TGTTGGCCCA	GCCAGAGTGA
		TGGCCACATT	GGAGATGGCG	GTAGCTGAGC	AGACATGCCC
		CCTGACTCCC	TGGTGTGCTC	CTGGAAGGAA	GATCTTGGGG
		CGGAGCACAC	CTAGGGATCA	TCTTTGCCCG	TCTCTGGGG
		AAATGTGGAG	TCCTCGGGGG	CCGTGCACTG	ATGCGGGGAG
		CTGGCGGTTG	GAGGGGTGGG	TGGGGGGCAG	TGGGGGCTGG
		TTCTGGGGTA	GGAAGTGGTC	CCGGGAGATT	TTGGATGGAA
		GATTGACAGC	AGACTTGCAG	AATTACATAG	AGAAATTAGG
		TTTCATGTCA	ATTGATCTAT	TCCCCCTCTT	TGTTTCTTGG
		CTTTTTTTTT	TTTTTTTGT	TTTTTTTAC	CCCTCCTTAG
		TCAGAAACCA	AATTAAACCC	CCCCCCCATG	TAACAGGGGG
		AAGCAAGAAC	GCACGAAGCC	AGCCTGGAGA	CCACCACGTC
		CCATTTATCG	CCCTGATTGG	ATTTTGT	TCATCTGTCC
		GGTTGAGTTG	AGGGTGGAGC	CTCCTGGGGG	GCACTGGCCA
		TTGGAGAAGT	CAGAGGGGAG	TGGAGAAGGC	CACGTGCTCG
		GGGGACAGTG	GCTGGTCCCC	AGAAGTCTTG	AGGGCGGAGG
		GCAGGGTCTC	CTCAGGTGTC	AGGAGGGTGC	TCGGAGGCCA
		CTCCTGGCTG	GCCTGAGGCT	GGCCGGAGGG	GAAGGGGCTA
		AAACAGAGGG	TTCCATCAGG	CTGGGGCAGG	GTGGCCGCTT
		TGAGGAACCC	TCCCCCTCTC	CTCGGTGACA	TCTTGCCCGC
		CCTGCCTTGT	CTCCAGGAGG	TCCGAAGCTC	TGTGGGACCT
		AGGTGGGGTG	AGGCCGGGGA	GTAGGGAGGT	CAGGCGGGTC
		GAGCAGGAGA	GCTGCCAGGT	CTGCCCATCG	ACCAGGTTGC
		GAGCCCACGG	GTCTGGTGAT	GCCATAGCAG	CCACCACCGC
		GCTGCGCAG	GGACTCGGCC	TCTGGGAGGT	TTACCTCGCC
		CCCCCAGCTC	AGCCCCCCTG	CACGCAGCCC	GACTAGCAGT
		GAGGCTTCTG	GGTCTGGTGG	ACGGGGCTGG	CATGACCCCG
		ATGCCAGTCC	GCCTCAGTCG	CAGAGGGTCC	CTCGCAAGC
		TGGGCCATTG	GGAACATTGG	ACAGAAGCCC	AAAGAGCCAA
		TTGTGGAACC	CACATTGGCC	TGAGATCCAA	AACGCTTCGA
		ATTACCTGCC	CATTCTGTCG	GACACCCACC	CACCCAGTGT
		CTCGCCGAGG	TGGGTGTTCC	CGGGGGCACT	TGCCGACCAG
		CCCCAGGTTT	GCAGCTCTCC	CCTGGGCCAC	TAACCATCCT
		GCCTGTCTGA	CCTCCGTGCC	TAGTCGTGGC	TCTCCATCTT
		CGTGTCCCCA	ATGTCTTCAG	TGGGGGGCCC	CCTCTTGGGT
		GCCATCACCT	GAAGACCCCC	ACGCCAAACA	CTGAATGTCA
		CCGCCTCGGT	CCACCTTGCG	GCCCCGTGTT	GACTCAACTC
		ACGCTAATAT	TTCCGGCAAA	ATCCCATGCT	TGGGTTTTGT
		GTAACGCTTG	CAATCCCAAT	AAAGCATTAA	AAGTCATGAA
		AAAAAA			AAAAAA
3	Human IGF2 mRNA, transcript variant 2	CGCCTGTCCC	CCTCCCGAGG	CCCGGGCTCG	CGACGGCAGA
		GGCCCAAAAC	GAGCTGGGCG	CCCGCGGTCC	GGGTGCAGCC
		CCCCCAGTCA	CCGCCCTCCC	CGGCCCTCG	ACGTGGCGCC
		CTTCTCTGTG	CTCCCGCGCC	CCCTCTTGGC	GTCTGGCCCC
		CTTTCTCCCG	CAACCTTCCC	TTGCTCTCCT	CCCGTCCCCC
		GCCTCCGACT	CCCTCCCCCC	CTACGCCCCG	CCCTCTCGCC
		CAAAAGTGGAT	TAATTACACG	CTTTCTGTTT	CTCTCCGTGC
		CGCTGTGCGC	CTGCCCGCCT	CTCGCTGTCC	TCTCTCCCCC
		TTCCGGCCCC	CCCTTTCACG	TTCACTCTGT	CTCTCCCACT
		CCCTCTATCC	TTGATACAAC	AGCTGACCTC	ATTTCCCGAT
		CCCCGAAAAG	TACAACATCT	GGCCCCGCC	AGCCCGAAGA
		TCCCTGGACA	ATCAGACGAA	TTCTCCCCCC	CCCCCAAAA
		CCCCCGCTCT	GCCCCGTGCG	ACATTCGGCC	CCCGCGACTC
		GCGCTGGCAG	AGGAGTGTCC	GGCAGGAGGG	CCAACGCCCG
		TGCGACACGC	AGCAGGGAGG	TGGGCGGCAG	CGTCGCCGGC
		CAATGGGAAT	CCCAATGGGG	AAGTCGATGC	TGGTGCTTCT
		GCCTTCGCCT	CGTGCTGCAT	TGCTGCTTAC	CGCCCCAGTG
		CGGCGGGGAG	CTGGTGGACA	CCCTCCAGTT	CGTCTGTGGG
		TCTACTTTCAG	CAGGCCCGCA	AGCCGTGTGA	GCCGTGCGAG
		GTTGAGGAGT	GCTGTTTCCG	CAGCTGTGAC	CTGGCCCTCC
		CTGTGCTACC	CCCGCCAAGT	CCGAGAGGGA	CGTGTGACCC
		TGCTTCCGGA	CAACTTCCCC	AGATACCCCG	TGGGCAAGTT
		GACACCTTGA	AGCAGTCCAC	CCAGCGCCTG	CGCAGGGGCC
		CCTGCGTGCC	CGCCGGGGTC	ACGTGCTCGC	CAAGGAGCTC
		GGGAGGCCAA	ACGTACCCGT	CCCCGTGATT	CTCTACCCAC
		GCCACGCGGG	GCGCCCCCCC	AGAGATGGCC	AGCAATCGGA
		ACTGCCCAA	GTCTGCAGCC	CGGCGCCACC	ATCCTGCAGC
		CCACGGACGT	TTCCATCAGG	TTCCATCCCC	AAAATCTCTC
		CCCCCTGGGG	CTTCTCCTGA	CCAGTCCCC	GTGCCCCGCC
		AGGCTACTCT	CCTCGGCCCC	CTCCATCGGG	CTGAGGAAGC
		CTTCAAACAT	GTACAAAATC	GATTGGCTTT	AAACACCCTT
		CCCCCAAAAT	TATCCCCAAT	TATCCCCACA	CATAAAAAAT
					CAAAACATTA

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		AACTAACCCC	CTTCCCCCCC	CCCCACAACA	ACCCTCTTAA
		CTTTTGTAGAA	ACACCCACACA	AAAGCTCAGA	AATTGGCTTT
		ACCACCAAAA	AAAATCAATT	GGCTAAAAAA	AAAAAGTATT
		TGGCTGAGAA	ACAATTGGCA	AAATAAAGGA	ATTTGGCACT
		CTCTTTCTCT	TCTCCCTTGG	ACTTTGAGTC	AAATTGGCCT
		CCCTGAACCA	GCAAGAGAGAA	AAGAAGGACC	CCAGAAATCA
		CGTCGCTGCT	ACCGCCATCT	CCCTTCTCAC	GGGAATTTTC
		GGCCATCCGA	AAATAGCAAC	AACCCAGACT	GGCTCCTCAC
		ATCACTAAAA	ATCACAGAGC	AGTCAGAGGG	ACCCAGTAAG
		GGGAGGACAG	AGCATGAAAA	CCAAAATCCA	TGCAATGAA
		CACGACCCTC	ACCCCCAAAT	CTTACATCTC	AATTCCCATC
		CTCATACTTT	ATGCATCCCC	GCAGCTACAC	ACACACAACA
		GCATGAACAC	AGCACACACA	CGAGCACAGC	ACACACAAA
		CACACAGCAC	ACAGATGAGC	ACACAGCACA	CACACAAAAG
		ACACGCACAC	ACATGCACAC	ACAGCACACA	AACGCACGGC
		ACACACATGC	ACACACAGCA	CACACACAAA	CGCACAGCAC
		ACAGCACACA	CGCACACACA	GCACACACAC	GAGCACACAG
		CACAGCACAC	GCACACACAT	GCACACACAG	CACACACACT
		ACACACACAA	AGACACAGCA	CACACATGCA	CACACAGCAC
		ACACAGCACA	CACGAACACA	GCACACACAG	CACACACACA
		ACACATGCAC	ACAGCACACG	CACACACAGC	ACACACATGA
		CAGCACACAC	ATGCACACAC	AGCACACACG	CATGCACAGC
		ACAGCACACA	CACAAACACA	CAGCACACAC	ATGCACACAC
		CTCATGCGCA	GCACATACAT	GAACACAGCT	CACAGCACAC
		CACACACGTT	GCACACGCAA	GCACCCACCT	GCACACACAC
		ACACGCACAC	CCCCACAAAA	TTGGATGAAA	ACAATAAGCA
		ACTACGATAT	CTGTATGGAT	CAGGCCAAAG	TCCCGCTAAG
		GTTTTTCATGG	TCTGAGCCCC	GCTCCTGTTC	CCATCTCCAC
		CCCTGTCTGT	GCCCTGCCTC	TCAGAGGAGG	GGGCTCAGAT
		GAGTGTGCGG	CCGGCGGCAT	TTGGGATACA	CCCGTAGGGT
		TGTCCAGGC	CTAATTCCAT	CTTTCCACCA	TGACAGAGAT
		GGCTGGCCTC	CTTGGCGCCT	GTCCCCACGG	CCCCCGCAGC
		ATGCTCCCCA	TACCCACACC	ATTCCCAGATA	CACCTTACTT
		GGCCCAGCCA	GAGTGAGGAA	GGAGTTTGGC	CACATTGGAG
		CTGAGCAGAC	ATGCCCCAC	GAGTAGCCTG	ACTCCCTGGT
		AAGGAAGATC	TTGGGGACCC	CCCCACCGGA	GCACACCTAG
		TGCCCCTCTC	CTGGGGACCC	CCCAAGAAAT	GTGGAGTCCT
		GCACTGATGC	GGGGAGTGTG	GGAAGTCTGG	CGGTTGGAGG
		GGGCAGTGGG	GGCTGGGCGG	GGGGAGTTCT	GGGGTAGGAA
		GAGATTTTGG	ATGGAAGT	CAGGAGGATT	GACAGCAGAC
		ACATAGAGAA	ATTAGGAACC	CCCAAATTTT	ATGTCAATTG
		CCTCTTTGTT	TCTTGGGGCA	TTTTTCCTTT	TTTTTTTTTT
		TTTTACCCCT	CCTTAGCTTT	ATGCGCTCAG	AAACCAAATT
		CCCATGTAAC	AGGGGGGCAG	TGACAAAAGC	AAGAACGCAC
		TGGAGACCAC	CACGTCTCTC	CCCCCGCCAT	TTATCGCCCT
		TGTTTTTCAT	CTGTCCCTGT	TGCTTGGGTT	GAGTTGAGGG
		TGGGGGGCAC	TGGCCACTGA	GCCCCCTTGG	AGAAGTCAGA
		GGAAGCCACT	GTCCGGCCTG	GCTTCTGGGG	ACAGTGGCTG
		GTCTGAGGG	CGGAGGGGGG	GTTTGGGCAG	GGTCTCCTCA
		GGGTGCTCGG	AGGCCACAGG	AGGGGGCTCC	TGGCTGGCCT
		GGAGGGGAAG	GGGCTAGCAG	GTGTGTAAC	AGAGGGTTCC
		GGCAGGGTGG	CCGCCTTCCG	CACACTTGAG	GAACCTTCCC
		GTGACATCTT	GCCCCCCCCCT	CAGCACCCCTG	CCTTGTCTCC
		AAGTCTGTGT	GGACCTCTTG	GGGGCAAGGT	GGGGTAGGGC
		GGAGGTCAGG	CGGGTCTGAG	CCACAGAGC	AGGAGAGCTG
		CCATCGACCA	GGTTGCTTGG	GCCCCGGAGC	CCACGGGTCT
		TAGCAGCCAC	CACCGCGCG	CCTAGGGCTG	CGGCAGGGAC
		GGAGGTTTAC	CTCGCCCCCA	CTTGTGCCCC	CAGCTCAGCC
		CAGCCCCACT	AGCAGTCTAG	AGGCCTGAGG	CTTCTGGGTC
		GGCTGGCATG	ACCCCGGGGG	TCGTCCATGC	CAGTCCGCCT
		GGGTCCCTCG	GCAAGCGCCC	TGTGAGTGGG	CCATTGCGAA
		AAGCCCAAAG	AGCCAAATTG	TCACAATTGT	GGAACCCACA
		ATCCAAACG	CTTCGAGGCA	CCCCAAATTA	CCTGCCCAT
		CCCACCCACC	CAGTGTATATA	TTCTGCCTCG	CCGGAGTGGG
		GGCACTTGCC	GACCAGCCCC	TTGCGTCCCC	AGGTTTGCAG
		GGCCACTAAC	CATCCTGGCC	CGGGCTGCCT	GTCTGACCTC
		CGTGGCTCTC	CATCTTGTCT	CCTCCCCGTG	TCCCCAATGT
		GGGCCCCCTC	TTGGGTCCCC	TCCTCTGCCA	TCACCTGAAG
		CAAACACTGA	ATGTCACCTG	TGCCCTGCCG	CTCGGTCCAC
		GTGTTTGACT	CAACTCAACT	CCTTTAACGC	TAATATTTCC
		CATGCTTGGG	TTTTGTCTTT	AACCTTGTA	CGCTTGCAAT
		CATTAAAGT	CATGAAAAAA	AAAAAAAATA	AA

-continued

TABLE OF CERTAIN SEQUENCES						
SEQ ID NO	Description	Sequence				
4	Human IGF2 mRNA, transcript variant 3	GGCCGCGCGC	CCTCAGGACG	TGGACAGGGA	GGGCTTCCCC	GTGTCCAGGA
		AAGCGACCGG	GCATTGCCCC	CAGTCTCCCC	CAAATTTGGG	CATTGTCCCC
		GGGTCTTCCA	ACGGACTGGG	CGTTGCTCCC	GGACACTGAG	GACTGGCCCC
		GGGGTCTCGC	TCACCTTCAG	CAGCGTCCAC	CGCCTGCCAC	AGAGCGTTCC
		ATCGCTCGCT	GCCTGAGCTC	CTGGTGCGCC	CGCGGACGCA	GCCTCCAGCT
		TCGCGGAGAT	GGTTTCCCCA	GACCCCAAAA	TTATCGTGGT	GGCCCCCGAG
		ACCGAACTCG	CGTCTATGCA	AGTCCAACGC	ACTGAGGACG	GGGTAACCAT
		TATCCAGATA	TTTTGGGTGG	GCCGCAAAGG	CGAGCTACTT	AGACGCACCC
		CGGTGAGCTC	GGCCATGCAG	ACACCAATGG	GAATCCCAAT	GGGGAAGTCG
		ATGCTGGTGC	TTCTCACCTT	CTTGGCCTTC	GCCTCGTGCT	GCATTGTCTC
		TTACCGCCCC	AGTGAGACCC	TGTGCGGCGG	GGAGCTGGTG	GACACCCCTC
		AGTTCTGTCT	TGGGGACCGC	GGCTTCTACT	TCAGCAGGCC	CGCAAGCCGT
		GTGAGCCGTC	GCAGCCGTGG	CATCGTTGAG	GAGTGCTGTT	TCCGCAGCTG
		TGACCTGGCC	CTCCTGGAGA	CGTACTGTGC	TACCCCGGCC	AAGTCCGAGA
		GGGACGTGTC	GACCCCTCCG	ACCGTGCTTC	CGGACAACTT	CCCCAGATAC
		CCCGTGGGCA	AGTTCTTCCA	ATATGACACC	TGGAAGCAGT	CCACCCAGCG
		CCTGCGCAGG	GGCCTGCCTG	CCCTCCTGCG	TGCCCCGCCG	GGTCACGTGC
		TCGCCAAGGA	GCTCGAGGCG	TTCAGGGAGG	CCAAACGTCA	CCGTCCCCTG
		ATTGCTCTAC	CCACCCAAGA	CCCCGCCAC	GGGGGCGCCC	CCCCAGAGAT
		GGCCAGCAAT	CGGAAGTGAG	CAAACTGCC	GCAAGTCTGC	AGCCCGGCGC
		CACCATCTGT	CAGCCTCCTC	CTGACCACGG	ACGTTTCCAT	CAGGTTCCAT
		CCCGAAATC	TCTCGTTCC	ACGTCCCCCT	GGGGCTTCTC	CTGACCCAGT
		CCCCGTGCCC	CGCCTCCCCG	AAACAGGCTA	CTCTCCTCGG	CCCCCTCCAT
		CGGGCTGAGG	AAGCACAGCA	GCATCTTCAA	ACATGTACAA	AATCGATTGG
		CTTTAAACAC	CCTTCACATA	CCCTCCCCCC	AAATTATCCC	CAATTATCCC
		CACACATAAA	AAATCAAAAC	ATTAACATAA	CCCCCTTCCC	CCCCCCCCAC
		AACAACCCCT	TTAAACTAA	TTGGCTTTT	AGAAACACCC	CACAAAAGCT
		CAGAAATTGG	CTTTAAAAAA	AACAACCACC	AAAAAAATC	AATTGGCTAA
		AAAAAAAAG	TATTAATAAC	GAATTGGCTG	AGAAACAATT	GGCAAAATAA
		AGGAATTGG	CACTCCCCAC	CCCCCTCTTT	CTCTTCTCCC	TTGGACTTTG
		AGTCAATTG	GCCTGGACTT	GAGTCCCTGA	ACCAGCAAAG	AGAAAAGAAG
		GACCCAGAA	ATCACAGGTG	GGCACGTCGC	TGCTACCGCC	ATCTCCCTTC
		TCACGGGAAT	TTTCAGGGTA	AACTGGCCAT	CCGAAAATAG	CAACAACCCA
		CAGTGGCTCC	TCACTCCCTT	TTCCATCACT	AAAAATCACA	GAGCAGTCAG
		AGGGACCCAG	TAAGACCAAA	GGAGGGGAGG	ACAGAGCATG	AAAACCAAAA
		TCCATGCAAA	TGAAATGTAA	TTGGCAGCAC	CCTCACCCCC	AAATCTTACA
		TCTCAATTCC	CATCCTAAAA	AGCACTCATA	CTTTATGCAT	CCCCGAGCT
		ACACACACAC	AACACACAGC	ACACGCATGA	ACACAGCACA	CACACGAGCA
		CAGCACACAC	ACAAACGCAC	AGCACACACA	GCACACAGAT	GAGCACACAG
		CACACACACA	AACGCACAGC	ACACACACGC	ACACACATGC	ACACACAGCA
		CACAAACGCA	CGGCACACAC	ACGCACACAC	ATGCACACAC	AGCACACACA
		CAAAACGACA	GCACACACAA	ACGCACAGCA	CACACGCACA	CACAGCACAC
		ACACGAGCAC	ACAGCACACA	AACGCACAGC	ACACGCACAC	ACATGCACAC
		ACAGCACACA	CACTAGCACA	CAGCACACAC	ACAAAGACAC	AGCACACACA
		TGCACACACA	GCACACACAC	GCGAACACAG	CACACAGCAA	CACAGCACAC
		ACAGCACACA	CACAAACACA	GCACACACAT	GCACACAGCA	CACGCACACA
		CAGCACACAC	ATGAACACAG	CACACAGCAC	ACACATGCAC	ACACAGCACA
		CACGCATGCA	CAGCACACAT	GAACACAGCA	CACACACAAA	CACACAGCAC
		ACACATGCAC	ACACAGCACA	CACACTCATG	CGCAGCACAT	ACATGAACAC
		AGCTCACAGC	ACACAAACAC	GCAGCACACA	CGTTGCACAC	GCAAGCACCC
		ACCTGCACAC	ACACATGCGC	ACACACACGC	ACACCCCCAC	AAAATTGGAT
		GAAAACAATA	AGCATATCTA	AGCAACTACG	ATATCTGTAT	GGATCAGGCC
		AAAGTCCCGC	TAAGATTCTC	CAATGTTTTT	ATGGTCTGAG	CCCCGCTCCT
		GTTCCCATCT	CCACTGCCCC	TCGGCCCTGT	CTGTGCCCTG	CCTCTCAGAG
		GAGGGGGCTC	AGATGGTGCG	GCCTGAGTGT	GCGGCCGGCG	GCATTTGGGA
		TACACCCGTA	GGGTGGGCGG	GGTGTGTCCC	AGGCCTAATT	CCATCTTTCC
		ACCATGACAG	AGATGCCCTT	GTGAGGCTGG	CCTCCTTGGC	GCCTGTCCCC
		ACGGCCCCCG	CAGCGTGAGC	CACGATGCTC	CCCATACCCC	ACCCATTCCC
		GATACACCTT	ACTTACTGTG	TGTTGGCCCA	GCCAGAGTGA	GGAAGGAGTT
		TGGCCACATT	GGAGATGGCG	GTAGCTGAGC	AGACATGCCC	CCACGAGTAG
		CCTGACTCCC	TGGTGTGCTC	CTGGAAGGAA	GATCTTGGGG	ACCCCCCAC
		CGGAGCACAC	CTAGGGATCA	TCTTTGCCCG	TCTCCTGGGG	ACCCCCCAAG
		AAATGTGGAG	TCCTCGGGGG	CCGTGCATGT	ATGCGGGGAG	TGTGGGAAAT
		CTGGCGGTTG	GAGGGGTGGG	TGGGGGGCAG	TGGGGGCTGG	GCGGGGGGAG
		TTCTGGGGTA	GGAAGTGCTC	CCGGGAGATT	TTGGATGGAA	AAGTCAGGAG
		GATTGACAGC	AGACTTGACG	AATTACATAG	AGAAATTAGG	AACCCCCAAA
		TTTCATGTCA	ATTGATCTAT	TCCCCCTCTT	TGTTTCTTGG	GGCATTTTTC
		CTTTTTTTTT	TTTTTTTGTT	TTTTTTTAC	CCCTCCTTAG	CTTTATGCGC
		TCAGAAACCA	AATTAAACCC	CCCCCCCATG	TAACAGGGGG	GCAGTGACAA
		AAGCAAGAAC	GCACGAAGCC	AGCCTGGAGA	CCACCACGTC	CTGCCCCCG
		CCATTTATCG	CCCTGATTGG	ATTTTGTGTT	TCATCTGTCC	CTGTTGCTTG
		GGTTGAGTTG	AGGGTGGAGC	CTCCTGGGGG	GCACGTGCCA	CTGAGCCCC
		TTGGAGAAGT	CAGAGGGGAG	TGGAGAAGGC	CACTGTCCGG	CCTGGCTTCT

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		GGGGACAGTG GCTGGTCCCC AGAAGTCCTG AGGGCGGAGG GGGGGGTTGG			
		GCAGGGTCTC CTCAGGTGTC AGGAGGGTGC TCGGAGGCCA CAGGAGGGGG			
		CTCCTGGCTG GCCTGAGGCT GGCCGGAGGG GAAGGGGCTA GCAGGTGTGT			
		AAACAGAGGG TTCCATCAGG CTGGGGCAGG GTGGCCGCTT TCCGCACACT			
		TGAGGAACCC TCCCCCTCTCC CTCGGTGACA TCTTGCCCGC CCCTCAGCAC			
		CCTGCCTTGT CTCCAGGAGG TCCGAAGCTC TGTGGGACCT CTGGGGGGCA			
		AGGTGGGGTG AGGCCGGGGA GTAGGGAGGT CAGGCGGGTC TGAGCCCA			
		GAGCAGGAGA GCTGCCAGGT CTGCCATCG ACCAGGTTGC TTGGGCCCCG			
		GAGCCACGG GTCTGGTGAT GCCATAGCAG CCACCACCGC GGCCTTAGG			
		GCTGCGCAG GGACTCGGCC TCTGGGAGGT TTACCTCGCC CCCACTGTG			
		CCCCAGCTC AGCCCCCTG CACGCAGCCC GACTAGCAGT CTAGAGGCCT			
		GAGGCTTCTG GGTCTGGTG ACGGGGCTGG CATGACCCCG GGGGTGCTCC			
		ATGCCAGTCC GCCTCAGTCG CAGAGGGTCC CTCGGCAAGC GCCCTGTAG			
		TGGGCCATTG GGAACATTGG ACAGAAGCCC AAAGAGCCAA ATTGTCACAA			
		TTGTGGAACC CACATTGGCC TGAGATCCAA AACGCTTCGA GGCACCCCAA			
		ATTACCTGCC CATTCTGTCAG GACACCCACC CACCCAGTGT TATATTCTGC			
		CTCGCCGGAG TGGGTGTTCC CGGGGGCACT TGCCGACCAG CCCCTTGCGT			
		CCCCAGTTT GCAGCTCTCC CTTGGGCCAC TAACCATCCT GCGCCGGGCT			
		GCCTGTCTGA CCTCCGTGCC TAGTCGTGGC TCTCCATCTT GTCTCTCC			
		CGTGTCCCCA ATGTCTTCAG TGGGGGGCCC CCTCTTGGGT CCCCTCCTCT			
		GCCATCACCT GAAGACCCCC ACGCCAAACA CTGAATGTCA CCTGTGCCTG			
		CCGCTCGGT CCACCTTGGC GCCCGTGTTC GACTCACTC AACTCCTTTA			
		ACGCTAATAT TTCCGGCAAA ATCCCATGCT TGGGTTTTGT CTTTAACCTT			
		GTAACGCTTG CAATCCCAAT AAAGCATTAA AAGTCATGAA AAAAAAAAAA			
		AAAAAA			
5	Human KRT20 mRNA	GAGACACACT CTGCCCCAAC CATCCTGAAG CTACAGGTGC TCCCTCCTGG			
		AATCTCCAAT GGATTTCACT CGCAGAAGCT TCCACAGAAG CCTGAGCTCC			
		TCCTTGCAAG CCCCTGTAGT CAGTACAGTG GGCATGCAGC GCCTCGGGAC			
		GACACCCAGC GTTTATGGGG GTGCTGGAGG CCGGGGCATC CGCATCTCCA			
		ACTCCAGACA CACGGTGAAC TATGGGAGCG ATCTCACAGG CGCGGGGAC			
		CTGTTTGTG GCAATGAGAA AATGGCCATG CAGAACCTAA ATGACCGTCT			
		AGCGAGTAC CTAGAAAAGG TCGGACCCCT GGAGCAGTCC AACTCCAAAC			
		TTGAAGTGCA AATCAAGCAG TGGTACGAAA CCAACGCCCC GAGGGCTGGT			
		CGCGACTACA GTGCATATTA CAGACAAATT GAAGAGCTGC GAAGTCAGAT			
		TAAGGATGCT CAACTGCAAA ATGCTCGGTG TGTCTGCAA ATTGATAATG			
		CTAAACTGGC TGCTGAGGAC TTCAGACTGA AGTATGAGAC TGAGAGAGGA			
		ATACGTCTAA CAGTGAAGC TGATCTCCAA GGCCTGAATA AGGTCTTTGA			
		TGACCTAACC CTACATAAAA CAGATTGGA GATTCAAATT GAAGAACTGA			
		ATAAAGACCT AGCTCTCCTC AAAAAGGAGC ATCAGGAGGA AGTCGATGGC			
		CTACACAAGC ATCTGGGCAA CACTGTCAAT GTGGAGGTTG ATGCTGCTCC			
		AGGCCTGAAC CTTGGCGTCA TCATGAATGA AATGAGGCAG AAGTATGAAG			
		TCATGGCCCA GAAGAACCTT CAAGAGGCCA AAGAACAGTT TGAGAGACAG			
		ACTGCAGTTC TGCAGCAACA GGTACAGTG AATACTGAAG AATTAAGG			
		AACTGAGGTT CAACTAACGG AGCTGAGAC CACCTCCAG AGCCTTGAGA			
		TAGAACTCCA GTCCATCTC AGCATGAAAG AGTCTTTGGA GCACACTCTA			
		GAGGAGACCA AGGCCCGTTA CAGCAGCCAG TTAGCCAAAC TCCAGTCGCT			
		GTTGAGCTCT CTGGAGGCC AACTGATGCA GATTCCGAGT AACATGGAAC			
		GCCAGAACAA CGAATACCAT ATCCTTCTTG ACATAAAGAC TCGACTTGAA			
		CAGGAAATTG CTACTTACCG CCGCTTCTG GAAGGAGAAG ACGTAAAAAC			
		TACAGAATAT CAGTTAAGCA CCCTGGAAGA GAGAGATATA AAGAAAACCA			
		GGAAGATTAA GACAGTCGTG CAAGAAGTAG TGGATGGCAA GGTCTGTCTA			
		TCTGAAGTCA AAGAGGTGGA AGAAAATATC TAAATAGCTA CCAGAAGGAG			
		ATGCTGCTGA GGTTTTGAAA GAAATTTGGC TATAATCTTA TCTTTGCTCC			
		CTGCAAGAAA TCAGCCATAA GAAAGCACTA TTAATACTCT GCAGTGATTA			
		GAAGGGGTGG GGTGGCGGGA ATCCTATTTA TCAGACTCTG TAATTGAATA			
		TAAATGTTTT ACTCAGAGGA GCTGCAAATT GCCTGCAAAA ATGAAATCCA			
		GTGAGCACTA GAATATTTAA AACATCATTA CTGCATCTT TATCATGAAG			
		CACATCAATT ACAAGCTGTA GACCACCTAA TATCAATTTG TAGGTAATGT			
		TCCTGAAAT TGCAATACAT TTCAATTATA CTAAACCTCA CAAAGTAGAG			
		GAATCCATGT AAATTGCAAA TAAACCACTT TCTAATTTT TCTGTTTCT			
		GAATTGTAAA ACCCCCTTGG GGAGTCCCTG GTTTCTTATT GAGCCAATTT			
		CTGGG			
6	Human ANXA10 mRNA	ATCCAGATTT GCTTTTACAT TTTCTGCCT GAGTCTGAGG TGAACAGTGA			
		ACATATTTAC ATTTGATTTA ACAGTGAACC TTAATTCCTT CTGGCTTCAC			
		AGTGAAACAA GTTTATGCAA TCGATCAAAT ATTTTCATCC CTGAGGTTAA			
		CAATTACCAT CAAAATGTTT TGTGGAGACT ATGTGCAAGG AACCATCTTC			
		CCAGCTCCCA ATTTCAATCC CATAATGGAT GCCCAAATGC TAGGAGGAGC			
		ACTCCAAGGA TTTGACTGTG ACAAAGACAT GCTGATCAAC ATCTGACTC			
		AGCGCTGCAA TGCACAAAGG ATGATGATTG CAGAGGCATA CCAGAGCATG			
		TATGGCCGGG ACCTGATTGG GGATATGAGG GAGCAGCTTT CGGATCACTT			
		CAAAGATGTG ATGGCTGGCC TCATGTACCC ACCACCACTG TATGATGCTC			

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		ATGAGCTCTG	GCATGCCATG	AAGGGAGTAG	GCACTGATGA
		ATTGAAATAC	TAGCTTCAAG	AACAAATGGA	GAAATTTTCC
		AGCCTACTGC	TTGCAATACA	GCAATAACCT	CCAAGAGGAC
		AGACCTCAGG	ACACTTCAGA	GATACTCTCA	TGAACCTGGT
		AGAGAGGAAG	GATATACAGA	CCCTGCGATG	GCTGCTCAGG
		CCTATGGGAA	GCCTGTCAGC	AGAAGACGGG	GGAGCACAAA
		AAATGATCCT	GTGCAACAAG	AGCTACCAGC	AGCTGCGGCT
		GAATTTCAAA	ATATTTCTGG	GCAAGATATG	GTAGATGCCA
		TTATGATGGA	TACTTTTCAGG	AGCTGCTGGT	TGCAATTGTT
		GAGACAAACC	AGCCTATTTT	GCTTATAGAT	TATATAGTGC
		TTTGGTTTCC	ATAATAAAAC	TGTAATCAGG	ATTCTCATTG
		AATAGACCTG	CTGACCATAA	GGAAACGATA	CAAAGAGCGA
		CCCTATTTCA	TGATATCAGA	AATTTTGCTT	CAGGGCATT
		CTGCTTGCCA	TCTGTGCTGG	TGATGCTGAG	GACTACTAAA
		CTTGAGATAC	TGTGCACTCC	TCTTTCTAGA	CACTTCCAAA
		CTCACAAATT	TGTACTGTTT	ATGGCACTAT	TAACAAAAT
		ATTTTCTCTT	CTATCTTTGA	AATTATTCTA	AGCCAAAGAA
		GAAAGTATAT	GATACTGAAT	TTGCCTACTA	TCCTGAATTT
		TAATCAGCAA	TTAAATAAAT	TGTGCATGAT	GGAATAATAG
		TTGGAATAGA	TTTTATTATA	ATGTGAACCA	TCAACAACCT
					ACAACAA
7	Human ABL mRNA	GGTTGGTGAC	TTCCACAGGA	AAAGTTCTGG	AGGAGTAGCC
		AGCGTTTCCT	TTATGTGTGA	GAATTGAAAT	GACTAGCATT
		TTGAGCATCC	CCTGTGAATA	TTTCTGTTTA	GGTTTTTCTT
		AATTGTTATT	CAGCCCGTTT	AAAACAAATC	AAGAAACTTT
		TGCAATTACA	TGAAATTGAT	AACCGCGAAA	ATAATTGGAA
		CAAGTGTCAA	CCTAAAAAAA	GTGCTTCCTT	TTGTTATGGA
		CTGTGATTGA	CTTCAATTGC	TGACTTGTGG	AGATGCAGCG
		CCCACGTATA	TGCCATTTC	CTCTACGCTC	GCTGACCGTT
		TTGAACCTTC	TTCTGGAAAG	GGGTACCTAT	TATTACTTTA
		GCCTGGAAAA	GTACTTGGGG	ACCAAAGAAG	GCCAAGCTTG
		ATTTTATCAA	AGGAGCAGGG	AAGAAGGAAT	CATCGAGGCA
		CACTGCAATG	TTTTTGTGGA	ACATGAAGCC	CTTCAGCGGC
		TGACTTTGAG	CCTCAGGCTC	TGAGTGAAGC	CGCTCGTTGG
		AAAACCTTCT	CGCTGGACCC	AGTGAATAATG	ACCCCAACCT
		CTGTATGATT	TTGTGGCCAG	TGGAGATAAC	ACTCTAAGCA
		TGAAAAGCTC	CGGGTCTTAG	GCTATAATCA	CAATGGGGAA
		CCCAAACCAA	AAATGGCCAA	GGCTGGGTCC	CAAGCAACTA
		GTCAACAGTC	TGGAGAAACA	CTCCTGGTAC	CATGGGCTTG
		TGCCGCTGAG	TATCTGCTGA	GCAGCGGGAT	CAATGGCAGC
		GTGAGAGTGA	GAGCAGTCCT	GGCCAGAGGT	CCATCTCGCT
		GGGAGGGTGT	ACCATTACAG	GATCAACACT	GCTTCTGATG
		CGTCTCTCTC	GAGAGCCGCT	TCAACACCCT	GGCCGAGTTG
		ATTCAACGGT	GGCCGACGGG	CTCATCACCA	CGCTCCATTA
		AAGCGCAACA	AGCCCACTGT	CTATGGTGTG	TCCCCCAACT
		GGAGATGGAA	CGCACGGACA	TCACCATGAA	GCACAAGCTG
		AGTACGGGGA	GGTGTAACGAG	GGCGTGTGGA	AGAAATACAG
		CGCCTGAAGA	CCTTGAAGGA	GGACACCATG	GAGGTGGAAG
		AGAAGCTGCA	GTCAATGAAAG	AGATCAAACA	CCCTAACCTG
		TTGGGGTCTG	CACCCGGGAG	CCCCCGTTCT	ATATCATCAC
		ACCTACGGGA	ACCTCTTGGA	CTACCTGAGG	GAGTGCACCC
		GAACGCCGTG	GTGCTGCTGT	ACATGGCCAC	TCAGATCTCG
		AGTACCTGGA	GAAGAAAAAC	TTCATCCACA	GAGATCTTGC
		TGCCTGGTAG	GGGAGAACCA	CTTGGTGAAG	GTAGCTGATT
		CAGGTTGATG	ACAGGGGACA	CCTACACAGC	CCATGCTGGA
		CCATCAAATG	GACTGCACCC	GAGAGCCTGG	CCTACAACAA
		AAGTCCGACG	TCTGGGCATT	TGGAGTATTG	CTTTGGGAAA
		TGGCATGTCC	CCTTACCCGG	GAATTGACCT	GTCCAGGTG
		TAGAGAAGGA	CTACCGCATG	GAGCGCCCAG	AAGGCTGCCC
		CTTTGCTGAA	TGCGAGCATG	TTGGCAGTGG	AATCCCTCTG
		TCTCAGACGA	AGTGGAAAAAG	GAGCTGGGGA	AACAAGGCGT
		GTGAGTACCT	TGCTGCAGGC	CCCAGAGCTG	CCCACCAAGA
		CAGGAGAGCT	GCAGAGCACA	GAGACACCAC	TGACGTGCCT
		ACTCCAAGGG	CCAGGGAGAG	AGCGATCCTC	TGGACCATGA
		TCTCCATTGC	TCCCTCGAAA	AGAGCGAGGT	CCCCCGGAGG
		TGAAGATGAG	CGCCTTCTCC	CCAAAGACAA	AAAGACCAAC
		CCTTGATCAA	GAAGAAGAAG	AAGACAGCCC	CAACCCCTCC
		AGCTCCTTCC	GGGAGATGGA	CGGCCAGCCG	GAGCGCAGAG
		GGAAGAGGGC	CGAGACATCA	GCAACGGGGC	ACTGGCTTTC
		ACACAGCTGA	CCCAGCCAAG	TCCCCAAAGC	CCAGCAATGG
		CCCAATGGAG	CCCTCCGGGA	GTCCGGGGGC	TCAGGCTTCC
		CCTGTGGAAG	AAGTCCAGCA	CGCTGACCAG	CAGCCGCCTA
					GCCACCGGCG

-continued

TABLE OF CERTAIN SEQUENCES					
SEQ ID NO	Description	Sequence			
		AGGAGGAGGG	CGGTGGCAGC	TCCAGCAAGC	GCTTCCTGCG
		GCCTCCTGCG	TTCCCATAGG	GGCCAAGGAC	ACGGAGTGGA
		GCTGCCTCGG	GACTTGCACT	CCACGGGAAG	ACAGTTTGAC
		TTGGAGGGCA	CAAAAGTGAG	AAGCCGGCTC	TGCCTCGGAA
		GAGAACAGGT	CTGACCAGGT	GACCCGAGGC	ACAGTAACGC
		GCTGGTGAAA	AAGAATGAGG	AAGCTGCTGA	TGAGGTCTTC
		TGGAGTCCAG	CCCGGGCTCC	AGCCCGCCCA	ACCTGACTCC
		CGGCGGCAGG	TCACCGTGGC	CCCTGCCTCG	GGCCTCCCCC
		AGCTGGAAAG	GGCAGTGCCCT	TAGGGACCCC	TGCTGCAGCT
		CCCCCACCAG	CAAAGCAGGC	TCAGGTGCAC	CAGGGGCAC
		CCCGCCGAGG	AGTCCAGAGT	GAGGAGGCAC	AAGCACTCCT
		AGGGAGGGAC	AAGGGGAAAT	TGTCCAGGCT	CAAACCTGCC
		CACCAGCAGC	CTCTGCAGGG	AAGGCTGGAG	GAAAGCCCTC
		AGCCAGGAGG	CGGCCGGGGA	GGCAGTCCTG	GGCGCAAAGA
		GAGTCTGGTT	GATGCTGTGA	ACAGTGACGC	TGCCAAGCCC
		GAGAGGGCCT	CAAAAAGCCC	GTGCTCCCGG	CCACTCCAAA
		GCCAAGCCGT	CGGGGACCCC	CATCAGCCCA	GCCCCCGTTC
		GCCATCAGCA	TCCTCGGCC	TGGCAGGGGA	CCAGCCGTCT
		TCATCCCTCT	CATATCAACC	CGAGTGTCTC	TTCGGAAAAC
		CCAGAGCGGA	TCGCCAGCGG	CGCCATCACC	AAGGGCGTGG
		CACCGAGGCG	CTGTGCCTCG	CCATCTCTAG	GAACCTCCGAG
		GCCACAGCGC	AGTGTGGAG	GCCGGCAAAA	ACCTCTACAC
		AGCTATGTGG	ATTCCATCCA	GCAAATGAGG	AACAAGTTTG
		GGCCATCAAC	AAACTGGAGA	ATAATCTCCG	GGAGCTTCAG
		CGACAGCAGG	CAGTGGTCCA	GCGGCCACTC	AGGACTTCAG
		AGTTCGGTGA	AGGAAATCAG	TGACATAGTG	CAGAGGTAGC
		GGTCAGGTGT	CAGGCCCGCT	GGAGCTGCCT	GCAGCACATG
		CATACCCGTG	ACAGTGCTG	ACAAGGGACT	AGTGAGTCAG
		CAGGAGCTCT	GCGCCAGGCA	GAGCTGAGGG	CCCTGTGGAG
		CTACCTACGT	TTGCACCCGC	TGCCCTCCCG	CACCTTCCTC
		CGTCTCTGTC	CTCGAATTTT	ATCTGTGGAG	TTCTTGCTCC
		GTCCGGCATGC	CAGGACCCGC	CAGCCCCGCT	CCCACCTAGT
		GAGCTCTCCA	GGCCAGGTGG	GAACGGCTGA	TGTGGACTGT
		TTTTTCTCTC	TGGAGCCCCT	CCTCCCCCGG	CTGGGCCTCC
		TCTCCAAGAA	TGGAAGCCTG	AACTGAGGCC	TTGTGTGTCA
		CTGCACTCCC	TGGCCTTGCC	CGTCGTGTGC	TGAAGACATG
		CGCATTTCGG	GAAGGGCATG	CACGGGCATG	CACACGGCTG
		CCTCTGCTGC	TGCCCCGGGT	GGGGTGCAC	CGCCATTTC
		GACAGCTCTT	GATTTGGGTG	GAAAACAGGG	TGCTAAAGCC
		TGGGTCTTGG	GCAGGTGGGA	GCTGAAAAGG	ATCGAGGCAT
		CTTTCCATCT	GTCCACATCC	CCAGAGCCCA	GCTCTTGCTC
		TGCACTGTGA	ATCCTGGCAA	GAAAGCTTGA	GTCTCAAGGG
		CTGTCACTGC	CGACATCCCT	CCCCCAGCAG	AATGGAGGCA
		GAGGCAGTGG	CTAGTGGGGT	GAACAGCTGG	TGCCAAATAG
		GGCCCCAGGA	GGTCTGCAAG	GGCCCAGAGT	GAACCGTCCT
		TGGGTGCCCT	GAAAGGGCCC	TTCCCCCTCC	CCACTCCTCT
		AGATTCTTAC	AAGGCCCTTT	CCTTTGGAAC	AAGACAGCCT
		GAGTCTTTGA	AGCATTTCAA	AGCCCTGCCT	CTGTGTAGCC
		AGAATAGAGC	TGCCACTGGG	CACCTGCGCA	CAGGTGGGAG
		GGCCAGTCCT	GGTCTTGGCT	GCACTCTTGA	ACTGGGCGAA
		AATTACCGTG	AGTGACATAG	CCTCATGTTC	TGTGGGGGTC
		GTTAGGAAAA	CCACAAACGG	AGCCCCTGAA	AGCCTCACGT
		GCACGCCTGC	CATCTTCTCC	CCGAGGCTGC	CCCAGGCCGG
		CGGGGGCTGT	GACTCTGGGC	AGGGACCCGG	GGTCTCTCTG
		AGCAGCTAAC	TCCGAGAGCA	GTGGGCAGGT	GGCCGCCCTC
		GCCGGGAGAA	GCCACCTTCC	CACCCCTTCA	TACCGCCTCG
		CTCGCACAGG	CCCTAGCTTT	ACGCTCATCA	CCTAAACTTG
		TTCTGATAGA	AATGGTTTCC	TCTGGATCGT	TTTATGCGGT
		CATCACCTCT	TTGCCCCCGA	CGGCTGTGAC	GCAGCCGGAG
		GTCACCGACA	GCGGCCCTGA	AGACAGAGCA	AAGCGCCAC
		CGACTGCCTG	TCTCCATGAG	GTACTGGTCC	CTTCCTTTTG
		GTGCCACTAT	ATTTTACACG	TATCTCTTGG	TATGCATCTT
		TCTTTTCTAA	GTGGCGTGTG	CATAGCGTCC	TGCCCTGCCC
		CTGTGGTGGC	TCCCCCTCTG	CTTCTCGGGG	TCCAGTGCAT
		TATATGATTG	TCTGTGGTTT	TTTTTGAATC	CAAATCTGTC
		TTTTTTAAAT	AAATCAGTGT	TTACATTAGA	A
8	Armored RNA [®] sequence	GAUGCUCACU	UCAUCUAUGG	UUACCCUGGG	ACUUUUACAC
		AGCAUCAUCC	UCUGCAUGGU	CAGGUCAUGG	AUCGGCAUCC
		GGGAUUUAGG	CAUCUGCAGU	CUUACUGCUC	AUCGGCUGAU
		UAUCCCCCAU	CCAAGCAAGC	UUGUGAUCCU	CCGCCAUUUA
		AUAACAUUUA	GGACUUAAAG	CUAUGCAAUU	AUACCUUUGU
		CAAGACCUAA	UAUUUUUUUU	UCAUCAUUAA	UGCCUUUUUGA
					UGGAUCAGGC

-continued

TABLE OF CERTAIN SEQUENCES

SEQ ID NO	Description	Sequence
		AACCAUUUUAU AAUAUGUUC ACCAGCCGAA GUCAGUAGUG AUUGGGUGGU
		UCCUGGCUUG GGAUCAUGCC GCUGCAGAGG CUAUUCUCCU CUUGGCAGAU
		UGUCUGUAGC CGAGAAGGCG GAGUCUGGCA AUGAUC AUGC AUACAGUGUA
		CGACAGCCUU AGGGACUGGA GCUCAAGCAG UGUUCCUCA ACCAGUCACA

SEQUENCE LISTING

Sequence total quantity: 48

SEQ ID NO: 1 moltype = DNA length = 1434

FEATURE Location/Qualifiers

source 1..1434

mol_type = genomic DNA

organism = Homo sapiens

SEQUENCE: 1

```

tcgttccttg gcagggccct atgatttatg caggagcaga ggcagcacgc aatcgagctg 60
tcaagagagc gtcagcttat taggcaaatg ctgcgtgggt tttgaagagg gtcgacacta 120
taaaatccca ctccaggctc tggagtggag aaactcagag accaagtcca ttgagagact 180
gaggggaaag agaggagaga aagaaaaaga gagtgggaac agtaaagaga aaggaagaca 240
acctccagag aaagcccccg gagacgtctc tctgcagaga ggcggcagca cccggctcac 300
ctgcgaagcg cctgggaagc gagtgcctc aacatgcggc tgcgctgct tgtgtccgcg 360
ggagtctctc tgggtgctct cctgcctctc ccgccatgca ggcgctcct gagcccgagg 420
ccggtcccg gacgtcggca ggcgcgcag caccctcagc ccttggattt ctccagccg 480
ccgcccagct ccgagcagcc ccagcagcc caggctcggc cggctcctgct ccgcatggga 540
gaggagtact tctcccgctt ggggaacctc aacaagagcc cggccgctcc cctttcgccc 600
gcctcctcgc tctcgcgcgg aggcagcggc agcccgccct cgcgggaaca ggcgacggc 660
aaactttttc gcgtgttgct gcagcagctg ctgctgcctc ggcgctcgtc cgacagcccc 720
gcggctctcg cggagcgcgg cgctaggaat gccctcggcg gccaccagga ggcaccggag 780
agagaaagcg ggtccgagga gctcccatc tccctggatc tcaccttcca cctcctccgg 840
gaagtcttgg aaatggccag gcccgagcag ttagcacagc aagctcacag caacaggaaa 900
ctcatggaga ttattgggaa ataaaacggg gcgtttggcc aaaaagaatc tgcatttagc 960
acaaaaaaaa tttaaaaaaa tacagtattc tgtaccatag cgctgctctt atgccatttg 1020
ttatttttta tatagcttga aacatagagg gagagaggga gagagcctat accccttact 1080
tagcatgcac aaagtgtatt cacgtgcagc agcaacacaa tgttattcgt tttgtctacg 1140
tttagtttcc gtttcagagt gtttatagtg gtgttttaaa gagaatgtag acctgtgaga 1200
aaacgttttg tttgaaaaag cagacagaag tcaactcaatt gtttttgggt tggctcgagc 1260
caaagagaat gccattctct tgggtgggta agactaaatc tgtaagctct ttgaaacaac 1320
tttctcttgt aaacgtttca gtaataaaac atctttccag tccttggtca gtttgggtgt 1380
gtaagagaat gttgaatact tatattttta ataaaagttg caaaggtaat catg 1434

```

SEQ ID NO: 2 moltype = DNA length = 5156

FEATURE Location/Qualifiers

source 1..5156

mol_type = genomic DNA

organism = Homo sapiens

SEQUENCE: 2

```

ccgctaattg accatgcctt ggtgctggaa agtgctgag ccagctgccc cagcggcctc 60
agcactacca agttggcaca aagctcccca aattcggagg ggctcaggga aacgagtgga 120
ggggatgagg aggtgagggg taaacccatc atttcagttg gcatttgagc aggtgccatg 180
ctcagcggag atgaggctct cccatctgta ggggcggtat taacatgcac actctaaaag 240
tgccctctgt tttccagacc tcagctttgt cctctctctc ctccaagctc acctggccag 300
agggctctga cgcacagacc agggcaccct ctgctttggt ggtgactgct aatattggcc 360
agggcggcgg atcatcgtcc aggcagtttc ggcagagagc cttgggcacc agtgactccc 420
cggtcctctt tatccactgc ccaggagctg cggggactgc gcagggacta gactacaggg 480
gccgaagagt caccaccgag cttgtgtggg aggaggtgga ttccagcccc cagccccagg 540
gctctgaatc gctgccagct cagccccctg cccagcctgc cccacagcct gagccccagc 600
agggcagaga gccagctctt gaggtgagct gctgtggcct gtggcccagg cgacccccagc 660
gctcccagaa ctgaggcttg cagccagccc cagcctcagc cccaactcgc aggcagagag 720
acaccaatgg gaatcccaat ggggaagtcg atgctggtgc ttctcaactt cttggccctc 780
gcctcgtgct cgcattgtcg ttaccgcccc agtgagaccc tgtgcggcgg ggagctgggtg 840
gacacccctc agttcgtctg tggggaccgc ggcttctact tcagcaggcc cgcaagccgt 900
gtgagccgtc gcagccgttg catcgttgag gactgtgtgt tccgcagctg tgacctggcc 960
ctcctggaga ctcctgctgc taacccccgc aagtcagaga gggacgtgtc gacccctccg 1020
accgtgcttc cggacaactt cccagatata cccgtgggca agttcttcca atatgacacc 1080
tggaagcagt ccccccagcg cctgcgcagg ggcctgcctg cctcctcgcg tgcggccggg 1140
ggtcacgtgc tcgccaagga gctcgaggcg ttcaggaggg ccaaaagctc ccgtccccctg 1200
attgctctac ccaccaaga ccccgccccc gggggcgccc cccagagatg ggccagcaat 1260
cggaaagtga caaaactgcc gcaagtctgc agcccgccgc caccatcctg cagcctcctc 1320
ctgaccacgg acgtttccat cagggtccat cccgaaaatc tctcggttcc acgtccccct 1380
ggggcttctc ctgaccagct ccccgctccc cgctcccccg aaacaggcta ctctcctcgg 1440

```

-continued

```

ccccctccat cgggctgagg aagcacagca gcattctcaa acatgtacaa aatcgattgg 1500
ctttaaacac ccttcacata cctccccccc aaattatccc caattatccc cacacataaa 1560
aatcaaaaac attaaactaa cccctctccc cccccccac aacaaccctc ttaaaactaa 1620
ttggcttttt agaaacaccc caaaaaagct cagaaattgg ctttaaaaaa aacaaccacc 1680
aaaaaaaatc aattggctaa aaaaaaaaag tattaaaaac gaattggctg agaaacaatt 1740
ggcaaaataa aggaatttgg cactccccac cccctctctt ctctctctcc ttggactttg 1800
agtcaaatgg gctcggactt gagtccctga accagcaaaag agaaaagaag gaccccagaa 1860
atcacagggt ggacgctgcg tgctacggcc atctcccttc tcacgggaat ttccagggtg 1920
aactggccat ccgaaaatag caacaaccca gactggctcc tcactccctt ttccatcact 1980
aaaaatcaca gacgagtcag agggacccag taagacaaa ggaggggagg acagagcatg 2040
aaaacccaaa tccatgcaaa tgaaatgtaa ttggcacgac cctcaccccc aaatcttaca 2100
tctcaattcc catcctaaaa agcactcata ctttatgcat ccccgagct acacacacac 2160
aacacacagc acacgcatga acacagcaca cacacgagca cagcacacac acaaacgcac 2220
agcacacaca cagcacagat gacgacacag cagcacacac aacgcacagc acacacagc 2280
acacacatgc acacacagca cacaacgcga cggcacacac acgcacacac atgcacacac 2340
agcacacaca caaacgcaca gcacacacaa acgcacagca cacacgcaca cacagcacac 2400
acacgagcac acagcacaca aacgcacagc acatgcacac acagcacaca 2460
cactagcaca cagcacacac acaaaagcac agcacacaca tgcacacaca gcacacacac 2520
gcgaacacag cacacacgaa cacagcacac acagcacaca cacaacaca gcacacacat 2580
gcacacagca cagcacacaa cagcacacac atgaacacag cacacagcac acacatgcac 2640
acacagcaca cagcagtcga cagcacacat gaacacagca cacacacaaa cacacagcac 2700
acacatgcac acacagcaca cacactcatg cgcagcacat acatgaacac agctcacagc 2760
acacaaacac gcacagcaca cgttgacac gcaagcaccc acctgcacac acacatgcgc 2820
acacacagcg acacccccac aaaattggat gaaaacaata agcatatcta agcaactacg 2880
atatctgtat ggatcaggcc aaagtcccg taagattctc caatgttttc atggtctgag 2940
ccccgctcct gttccccctt ccaactgccc tcggccctgt ctgtgccctg cctctcagag 3000
gagggggctc agatgggtgc gcttgagtgt gcggccggcg gcatttggga tacaccctga 3060
gggtggcgcg ggtgtgtccc aggcctaatt ccatctttcc accatgacag agatgccctt 3120
gtgaggctgg cctccttggc gctgtcccc acggcccccg cagcgtgagc cagcatgctc 3180
cccatacccc acccattccc gatcacctt acttactgtg tgttggccca gccagagtga 3240
ggaaggagtt tggccacatt ggagatggcg gtagctgagc agacatgccc ccacgagtag 3300
cctgactccc ctggtgtgtc ctgggaaggaa gatcttgggg accccccac cggagcacac 3360
ctagggatca tctttgcccc tctcctgggg accccccaa aaatgtggag tctcggggg 3420
ccgtgcactg atgcggggag tgtgggaagt ctggcggttg gaggggtggg tggggggcag 3480
tgggggctgg cgggggggag ttctggggta ggaagtggtc cggggagatt ttggatggaa 3540
aagtccaggag gatgacagc agacttgacg aattacatag agaaattagg aacccccaaa 3600
tttcatgtca attgatctat tccccctctt tgtttcttgg ggcatttttc cttttttttt 3660
tttttttgtt ttttttttac cctccttag ctttatgcgc tcagaaacca aattaaaccc 3720
ccccccatg taacaggggg gcagtgacaa aagcaagaac gcacgaagcc agcctggaga 3780
ccaccacgtc ctgcccccg ccatttatcg cctgattgg attttgttt tcactgttcc 3840
ctgttgcttg ggttgagtgg aggggtggag cctcggggg gcactggcca ctgagcccc 3900
ttggagaagt cagaggggag tggagaaggc cactgtccgg cctggcttct ggggacagt 3960
gctggtcccc agaagctctg agggcgaggg ggggggttgg gcagggtctc ctacaggtgc 4020
aggaggggtc tcggaggcca caggaggggg ctcctggctg gcctgaggct ggccggaggg 4080
gaaggggcta gcagggtgtg aaacagaggg ttccatcagg ctggggcagg gtggccgct 4140
tcgcacactc tgagggaacc tccccctccc ctcggtgaca tcttgcgcg cctcagcac 4200
cctgccttgt ctccaggagg tccgaagctc tgtgggacct cttgggggca aggtgggggtg 4260
aggccgggga gtaggggagt caggcggtgc tgagccaca gagcaggaga gctgccaggt 4320
ctgcccacgc accaggttgc ttgggccccg gagccacgg gtctggtgat gccatagcag 4380
ccaccaccgc ggcgcctagg gctcgggag ggaactcgcc tctgggaggt ttacctcgcc 4440
cccacttgtg cccccagctc agccccctg cagcgagccc gactagcagt ctgagggcct 4500
gaggctcttg ggtcctggtg acggggctgg catgaccccg ggggtcgtcc atgccagtcc 4560
gectcagtcg cagaggggtc ctcggaagc gccctgtgag tgggcccatt ggaacattgg 4620
acagaagccc aaagagccaa attgtcacia ttgtggaacc cacattggcc tgagatccaa 4680
aacgcttcga ggcaccccaa attactgcc cattcgtcag gacaccaccc caccagtggt 4740
tatattctgc ctgcgggag tgggtgttcc cgggggacct tgccgaccag ccccttgctg 4800
ccccaggttt gcagctctcc cctggggccc taacctcct ggcccgggct gctgtctga 4860
cctcgtgccc tagtcgtggc tctccatctt gtctcctccc cgtgtcccca atgtcttcag 4920
tggggggccc cctcttgggt cccctcctct gccatcacct gaagaccccc acgcccacaa 4980
ctgaatgtca cctgtgcctg ccgcctcggt ccaccttgcg gcccggtttt gactcaactc 5040
aactccttta acgctaatat ttccggcaaa atcccatgct tgggttttgt ctttaacctt 5100
gtaacgcttg caatcccaat aaagcattaa aagtcatgaa aaaaaaaaaa aaaaaa 5156

```

```

SEQ ID NO: 3      moltype = DNA length = 5182
FEATURE
source            Location/Qualifiers
                  1..5182
                  mol_type = genomic DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 3
cgctgtcccc cctcccgagg cccgggctcg cgacggcaga gggctccgtc ggcccaaac 60
gagctggggc cccgctgagcc ggggtgcagcc tccactccgc ccccagtcac ccgcctcccc 120
cgcccccctc acgtggcgcc cttccctccg cttctctgtg ctcccgccgc cctctttggc 180
gtctggcccc gggccccgct ctttctcccc caacctcccc tctgctcccc cccgtcccc 240
ccagctccta gcctccgaact cctccccccc ctacgccccg cctctcgccc ttcccgaaac 300
caaatgggat taattacacg ctttctgttt ctctcgtgct tgttctctcc cgtgtgctgc 360
ctgcccgcct ctegtgttcc ttctcccccc tcgcccctctc ttggccccc ccttttcacg 420
ttcactctgt ctctcccatc atctctgccc cctctatccc ttgatacaac agctgacctc 480
atttcccgat accttttccc cccgaaaaag tacaacatct ggcccgcgcc agcccgaaag 540
cagcccgctc tccctggaca atcagacgaa ttctcccccc cccccaaaaa aaaagccatc 600

```

-continued

```

ccccgcgtct gcccgcgtgc acattcggcc cccgcgactc ggcagagcgg gcgttggcag 660
aggagtgtcc ggcaggagg ccaacgccc ctgttcggtt tgcgacacgc agcaggagg 720
tggtcgccag cgtcgccggtc ttccagacac caatgggaat cccaatgggg aagtcgatgc 780
tggtgcttct cacttcttg gcttcgcct cgtgctgcac tgcgtgttac cgcgccagt 840
agaccctgtg cggcggggag ctggtggaca cctccagtt cgtctgtggg gaccgaggct 900
tctacttcag caggcccgcga agcgtgtga gccgtcgag ccgtggcatc gttgaggagt 960
gctgtttccg cagctgtgac ctggccctcc tggagacgta ctgtgtacc ccgcgaagt 1020
ccgagaggga cgtgtcgacc cctccgaccg tgcttcggga caacttccc agatacccc 1080
tggtcaagtt cttccaatat gacacctgga agcagtcac ccagcgccgt cgcaggggcc 1140
tgctgcctct cctgcgtgct cgccgggggtc acgtgctgc caaggagctc gaggcgttca 1200
gggaggccaa acgtccacgt cccctgattg ctctaccac ccaagacccc gccacgggg 1260
gcccccccc agagatggcc agcaatcgga agtgagcaaa actgcccga gtctgcagcc 1320
cggcgccacc atctgcagc ctctcctga ccacggagct tccatcagg tccatccc 1380
aaaaatcttc cgtgtccagc cccctgggg ctctcctga cccagtcacc gtgcccccc 1440
tccccgaac aggtactctc cctcgccccc ctccatcggt ctgaggagg acagcagcat 1500
cttcaaacat gtacaaaatc gattggcttt aaacaccct cacaaccct ccccccagt 1560
tatccccaat tctccacaca cataaaaaat caaaaacatta aactaacccc cttcccccc 1620
ccccacacaa accctcttaa aactaattgg ctttttagaa acaccccaca aaagtcaga 1680
aattggcttt aaaaaaaaca accaccaaaa aaaatcaatt gggtaaaaaa aaaaagtatt 1740
aaaaacgaat tggctgagaa acaattggca aaataaaggga atttggcact ccccccccc 1800
ctctttctct tctcccttgg actttgagtc aaattggcct ggacttgagt ccttgaacca 1860
gcaaagagaa aagaaggacc ccagaaatca caggtgggca cgtcgtgct accgcatct 1920
ccttctcacc gggaattttc agggtaaaact ggccatccga aaatagcaac aaccagact 1980
ggctcctcac tcccttttcc atcactaaaa atcacagagc agtcagagg acccagtaag 2040
accaaaggag gggaggacag agcatgaaaa ccaaaatcca tgcaaatgaa atgtaattgg 2100
cacgacctc acccccaaat ctacatctc aatccccac ctaaaaagca ctcatcttt 2160
atgcatcccc gcagctacac acacacaaca cacagcacac gcatgaacac agcacacaca 2220
cgagcacagc acacacacaa acgcacagca cacacagcac acagatgagc acacagcaca 2280
cacacaaaag cacagcacac acacgcacac acatgcacac acagcacaca aacgcacggc 2340
acacacacgc acacacatgc acacacagca cacacacaaa cgcacagcac acacaaaagc 2400
acagcacaca cgcacacaca gcacacacac gagcacacag cacacaaaag cacagcacac 2460
gcacacacat gcacacacag cacacacact agcacacagc acacacacaa agcacagca 2520
cacacatgca cacacagcac acacacgcga acacagcaca caggaacaca gcacacagc 2580
cacacacaca aacacagcac acacatgcac acagcacagc cacacacagc acacacatga 2640
acacagcaca cagcacacac atgcacacac agcacacacg catgcacagc acacatgaac 2700
acagcacaca cacaacacaca cagcacacac atgcacacac agcacacaca ctcatgcga 2760
gcacatacat gaacacagct cagacacac aaacacgcag cacacacgtt gcacacgca 2820
gcacccacct gcacacacac atgcgcacac acacgcacac cccacaaaaa ttggatgaaa 2880
acaataagca tatctaagca actacgatat ctgtatggat caggccaaaag tcccgctaag 2940
attctccaat gttttcttgg tctgagcccc gctcctgttc ccatctccac tgccccctcg 3000
cctgtctgt gcccctgcctc tcagaggagg gggctcagat ggtgcggcct gagtgtgagg 3060
cggcgggcat ttgggataca cccgtagggt gggcggggtg tgtcccaggc ctaattccat 3120
ctttccacca tagacagat gccctgtgga ggtggcctc gtcgccacgg 3180
cccccgagc gtgagccagc atgctcccc taacccccc attcccgata caccctactt 3240
actgtgtgtt ggcgccgcca gagttaggaa ggagtttggc cacattggag atggcggtag 3300
ctgagcagac atgcccccat gagtgcctg actccctggt gtgctcctgg aagggaagatc 3360
ttggggaccc cccaccgga gcacacctag ggatcatctt tgcccgtctc ctggggaccc 3420
cccaagaat gtggagtcct cgggggcccgt gcatgagtc ggggagtggt ggaagtctgg 3480
cggttggagg ggtgggtggg gggcagtggt ggtggggcgg ggggagttct ggggtaggaa 3540
gtggtcccg gtgatatttg atggaaaagt caggaggatt gacagcagc ttgcagaatt 3600
acatagagaa attaggaacc cccaaatttc atgtcaattg atctattccc cctcttgtt 3660
tcttggggca tttttctctt tttttttttt ttttaacctt ccttagcttt 3720
atgcgtctag aaaccaaatt aaaccccccc ccatgtaac agggggggcag tgacaaaagc 3780
aagaacgcac gaagccagcc tggagaccac cacgtcctgc ccccccgcct tatcgccc 3840
gattggattt ggtttttcat ctgtccctgt tgcttgggtt gatttgaggg tggagcctcc 3900
tggtggggcag tggccactga gcccccttgg agaagtcaga ggggagtgga gaaggccact 3960
gtccggcctg gcttctgggg acagtggctg gtcccagaaa gtccagagg cggagggggg 4020
ggttggggcag ggtctcctca ggtgtcagga ggtgtcctcg aggcacagg agggggctcc 4080
tggttggcct gaggtgggac ggagggggaag gggttagcag gtgtgtaaac agagggttcc 4140
atcaggctgg ggcagggtgg ccgccttccg cacacttgag gaacccctcc ctctccctcg 4200
gtgacatctt gcccgccctc cagcacccctg ccttgtctcc aggaggtccg aagctctgtg 4260
ggacctcttg ggggcaaggt ggggtgaggg cggggagtag ggaggtcagg cgggtctgag 4320
cccacagagc agggagagctg ccaggctctgc ccatcgacca ggtgtcttgg gcccgggagc 4380
ccacgggtct ggtgatgcca tagcagccac caccgcgcg cctagggtcg cggcagggac 4440
tcggcctctg gggaggttac ctgcgcccc cttgtgcccc cagctcagcc cccctgcagc 4500
cagcccgact agcagctag aggcctgagg cttctgggtc ctgttgacgg ggttggcatg 4560
accccggggg tcttcctatgc cagtccgctc cagtgcaga gggtcctctg gcaagcgccc 4620
tgtgagtggt ccattcgcaa cattggacag aagcccaaa agccaaaattg tcaaatgtt 4680
ggaaaccaca ttgcccgtgag atccaaaacg cttcgaggca ccccaaatca cctgccatt 4740
cgtcaggaca cccaccacc cagtgttata ttctgcctcg ccggagtggt tgttccccgg 4800
ggcacttgcc gaccagcccc ttgcgtcccc aggtttgcag ctctccctcg ggcactaac 4860
catcctggcc cgggtgctct gtctgacctc cgtgcctagt cgtggctctc catcttgtct 4920
ctctccctgt tccccaatgt ctcagtggtg gggccccctc ttgggtcccc tctcttgcca 4980
tcacctgaag acccccacgc caaacactga atgtcacctg tgccctgccc ctcggtccac 5040
cttgcggccc gtgtttgact caactcaact cctttaacgc taatatttcc ggcaaaatcc 5100
catgcttggt ttttgtctt aaacttgtaa cgcttgcaat cccaataaag cattaaaagt 5160
catgaaaaaa aaaaaaaaaa aa 5182

```


-continued

FEATURE	Location/Qualifiers					
source	1..4856					
	mol_type = genomic DNA					
	organism = Homo sapiens					
SEQUENCE: 4						
ggcgcgcgcg	cctcaggacg	tggacaggga	gggcttcccc	gtgtccagga	aagcgaccgg	60
gcattgcccc	cagtctcccc	caaatttggg	cattgtcccc	gggtcttcca	acggactggg	120
cgttgctccc	ggacactgag	gactggcccc	ggggctctgc	tcaccttcag	cagcgtccac	180
cgctgccac	agagcgcttg	atcgctcgct	gcctgagctc	ctgggtgcgc	cgcggaacga	240
gcctccagct	tcgcggagat	ggtttcccca	gacccccaaa	ttatcgtggt	ggcccccgag	300
accgaactcg	cgctatgca	agtccaacgc	actgaggacg	gggtaacccat	tatccagata	360
ttttgggtgg	gcccgaagg	cgagctactt	agacgcaccc	cggtgagctc	ggccatgcag	420
acaccaatgg	gaatcccaat	ggggaagtgc	atgctgggtc	ttctcacctt	cttggccttc	480
gcctcggtgc	gcattgtgtc	ttaccgcccc	agtgcgaccc	tggtgcggcg	ggagctgggtg	540
gacacccctc	agttcgtctg	tggggaccgc	ggcttctact	tcagcaggcc	cgcaagccgt	600
gtgagccgtc	gcagccgtgg	catcggtgag	gagtgctgtt	tcgcagctgt	tgacctggcc	660
ctcctggaga	tcgccaagga	tacccccgcc	aagtccgaga	gggacgtgtc	gacccctccg	720
accgtgcttc	cggacaactt	ccccagatag	cccggtgggc	agttcttcca	atatgacacc	780
tggagcagct	cccccagcg	cctgcgcagg	ggcctgcctg	ccctcctgcg	tgccccgcgg	840
ggtcacgtgc	tcgccaagga	gctcgaggcg	ttcagggagg	ccaaacgtca	ccgtccccctg	900
attgtctctac	cccccgaaga	ccccgcccac	gggggcgcgc	ccccagagat	ggccagcaat	960
cggaagtggag	caaaaactgc	gcaagtctgc	agcccggcgc	caccatcctg	cagcctctctc	1020
ctgaccacgg	acgtttccat	cagggttccat	cccgaataac	tctcggttcc	acgtccccct	1080
ggggcttctc	ctgaccagct	ccccgtgccc	cgctcccccg	aaacaggcta	ctctctctcg	1140
ccccctccat	cgggctgagg	aagcacagca	gcattctcaa	acatgtacaa	aatcgattgg	1200
ctttaaacac	ccttcacata	ccctcccccc	aaattatccc	caattatccc	cacacataaa	1260
aatcaaaaac	attaaactaa	cccccttccc	ccccccccac	aacaaccttc	ttaaaactaa	1320
ttggcttttt	agaaaacacc	caaaaaagct	cagaaattgg	ctttaaaaaa	aacaaccacc	1380
aaaaaaaaac	aattggctaa	aaaaaaaaag	tattaaaaac	gaattggctg	agaaacaatt	1440
ggcaaaataa	aggaatttgg	cactccccac	ccccctcttt	ctcttctccc	ttggactttg	1500
agtcaaaattg	gcctggactt	gagtcctcga	accagcaaa	agaaaagaag	gaccccgtaa	1560
atcacagggtg	ggcacgtcgc	tgctaccgcc	atctcccttc	tcacgggaat	tttcagggta	1620
aactggccat	cggaaaatag	caacaaccca	gactggctcc	tcactccctt	ttccatcact	1680
aaaaatcaca	gcagcgtcag	agggaccag	taagaccaaa	ggagggggagg	acagagcatg	1740
aaaacccaaa	tcctatgcaa	tgaaatgtaa	ttggcacgac	cctcaccccc	aaatcttaca	1800
tctcaattcc	catcctaaaa	agcactcata	ctttatgcac	ccccgcagct	acacacacac	1860
aaacacacgc	caacgcagca	acacagcaca	cacacgagca	cagcacacac	acaaacgcac	1920
agcacacaca	gcacacagat	gagcacacag	cacacacaca	aacgcacagc	acacacacgc	1980
acacacatgc	acacacagca	cacaaacgca	cggcacacac	acgcacacac	atgcacacac	2040
agcacacaca	caaacgcaca	gcacacacaa	acgcacagca	cacacgcaca	cacagcacac	2100
acacgagcac	caacgcacaca	aacgcacagc	acacgcacac	acatgcacac	acagcacaca	2160
cactagcac	cagcacacac	acaaagacac	agcacacaca	tgcacacaca	gcacacacac	2220
gcgaacacag	cacacacgaa	cacagcacac	acagcacaca	cacaaacaca	gcacacacat	2280
gcacacagca	cacgcacaca	cagcacacac	atgaacacag	cacacagcac	acacatgcac	2340
acacagcaca	cacgcgatgca	cagcacacat	gaacacagca	cacacacaaa	cacacagcac	2400
acacatgcac	acacagcaca	cacactcatg	cgcagcacat	acatgaacac	agctcacagc	2460
acacaaacac	gcagcacaca	cgttgccac	gcaagcaccc	acctgcacac	acacatgcgc	2520
acacacacgc	acacccccac	aaaattggat	gaaaacaata	agcatatcta	agcaactacg	2580
atatctgtat	ggatcaggcc	aaagtccccg	taagattctc	caatgttttc	atggctcgag	2640
ccccgctcct	gttccccatc	ccactgcccc	tcggccctgt	ctgtgccctg	cctctcagag	2700
gagggggctc	agatgggtgc	gcctgagtg	cgggccgcgc	gcatttggga	tacaccctga	2760
gggtggggcg	ggtgtgtccc	agggcctaatt	ccatcttttc	accatgacag	agatgccctt	2820
gtgagggctg	cctccttgcc	gcctgtcccc	acggcccccg	cagcgtgagc	cacgatgctc	2880
cccatacccc	accattcccc	gatacacctt	acttactgtg	tgttggccca	gccagagtga	2940
ggaaggagtt	tggccacatt	ggagatggcg	gtagctgagc	agacatgccc	ccacgagtga	3000
cctgactccc	ttggtgtgct	ctggaaggaa	gatcttgggg	acccccccac	cggagcacac	3060
ctagggtaca	tctttgcccc	tctcctgggg	accccccaag	aaatgtggag	tctcgggggg	3120
ccgtgcactg	atgcggggag	tgtgggaagt	ctggcgggtg	gaggggtggg	tggggggcag	3180
tgggggctgg	gcggggggag	ttctggggta	ggaagtggtc	ccgggagatt	ttggatggaa	3240
aagtgcaggag	gattgacagc	agacttgacg	aattacatag	agaaattagg	aacccccaaa	3300
tttcatgtca	attgatctat	tccccctctt	tgtttcttgg	ggcatttttc	cttttttttt	3360
tttttttgtt	ttttttttac	ccctccttag	ctttatgcgc	tcagaaacca	aattaaaccc	3420
ccccccatg	taaacggggg	gcagtgacaa	aagcaagaac	gcacgaagcc	agcctggaga	3480
ccaccacgtc	ctgccccccg	ccatttatcg	ccctgatagg	attttgtttt	tcactcgtcc	3540
ctgttgcttg	gggtgagttg	aggggtggagc	ctcctggggg	gcactggcca	ctgagcccc 3600	
ttggagaagt	cagaggggag	tggaagaagg	cactgtccgg	cctggctctc	ggggacagtg 3660	
gctggtcccc	agaagtcctg	agggcggagg	gggggggttg	gcagggtctc	ctcaggtgtc 3720	
aggaggggtg	tcggaaggca	caggaggggg	ctcctggctg	gcctgaggct	ggccggagggt 3780	
gaaggggcta	gcagggtgtg	aaacagaggg	ttccatcagg	ctggggcagg	gtggccgcct 3840	
tccgcacact	tgaggaaccc	tccccctctc	ctcggtgaca	tcttgcccgc	ccctcagcac 3900	
cctgccttgt	ctccaggagg	tcggaagctc	tgtgggaact	cttgggggca	aggtgggggtg 3960	
aggccgggga	gtagggaggt	caggcggggtc	tgagcccaca	gagcaggaga	gctgccaggt 4020	
ctgccccatg	accaggttgc	ttggggcccc	gagccccagg	gtctggtgat	gccatagcag 4080	
ccaccaccgc	ggcgccctagg	gctcgggcag	ggactcggcc	tctgggaggt	ttacctgcgc 4140	
cccacttgtg	ccccccagctc	agccccctgt	cacgcagccc	gactagcagt	ctagaggcct 4200	
gaggcttctg	ggtcctgggtg	acggggctgg	catgaccccc	ggggctcgtc	atgccagtc 4260	
gcctcagtcg	cagaggggtc	ctcggaagc	gccctgtgag	tgggccattc	ggaacattgg 4320	
acagaagccc	aaagagccaa	attgtcacaa	tttgtgaacc	cacatitggc	tgagatccaa 4380	
aacgcttcga	ggcaccccaa	attacctgcc	cattcgtcag	gacaccacc	caccagtggt 4440	

-continued

tatattctgc	ctcgccggag	tgggtgttcc	cgggggcact	tgccgaccag	ccccctgcgt	4500
ccccaggttt	gcagctctcc	cctggggccac	taaccatcct	ggcccgggct	gcctgtctga	4560
cctccgtgcc	tagctgtggc	tctccatctt	gtctcctccc	cgtgtcccca	atgtcttcag	4620
tggggggccc	cctcttgggt	ccccctctct	gccatcacct	gaagaccccc	acgccaacaa	4680
ctgaatgtca	cctgtgcctg	cgcctcgggt	ccaccttgcg	gcccgtgttt	gactcaactc	4740
aactccttta	acgctaatat	ttccggcaaa	atcccatgct	tgggttttgt	ctttaacctt	4800
gtaacgcttg	caatcccaat	aaagcattaa	aagtcatgaa	aaaaaaaaaa	aaaaaa	4856

SEQ ID NO: 5 moltype = DNA length = 1805
 FEATURE Location/Qualifiers
 source 1..1805
 mol_type = genomic DNA
 organism = Homo sapiens

SEQUENCE: 5

gagacacact	ctgccccaac	catcctgaag	ctacaggtgc	tcctcctcgt	aatctccaat	60
ggatttcagt	cgcagaagct	tcacagaag	cctgagctcc	tcctgcagg	ccccgtagt	120
cagtacagtg	ggcatgcagc	gccctgggac	gacacccagc	gtttatgggg	gtgctggagg	180
cggggggcatc	cgcctctcca	actccagaca	cacggtgaac	tatggggagc	atctcacagg	240
cggcggggac	ctgtttgttg	gcaatgagaa	aatggccatg	cagaacctaa	atgaccgtct	300
agcgagctac	ctagaaaagg	tgcggacctt	ggagcagtc	aactccaaac	ttgaagtga	360
aatcaagcag	tggtagcaaa	ccaacgcccc	gagggctggt	cgcgactaca	gtgcatatta	420
cagacaaatt	gaagagctgc	gaagtcagat	taaggatgct	caactgcaaa	atgctcggtg	480
tgtcctgcaa	attgataatt	ctaaactggc	tgctgaggac	ttcagactga	agtatgagac	540
tgagagagga	atacgtctaa	cagtggaaag	tgatctccaa	ggcctgaata	aggtctttga	600
tgacctaacc	ctacataaaa	cagatttggg	gattcaaat	gaagaactga	ataaagacct	660
agctctctct	aaaaaggagc	atcaggagga	agtcgatggc	ctacacaagc	atctggggca	720
cactgtcaat	gtggagggtg	atgctgctcc	aggcctgaac	cttggcgta	tcataaatga	780
aatgaggcag	aagtatgaag	tcatggccca	gaagaacctt	caagaggcca	aagaacagtt	840
tgagagacag	actgagcttc	tgagcaaca	ggtcacagt	aatactgaag	aattaaaagg	900
aactgaggtt	caactaacgg	agctgagacg	cacctcccag	agccttgaga	tagaactcca	960
gtcccatctc	agcatgaaag	agtctttgga	gcacactcta	gaggagacca	aggcccggtt	1020
cagcagccag	ttagccaaac	tcagtcgct	gttgagctct	ctggaggccc	aactgatgca	1080
gattcggagt	aacatggaac	gccagaacaa	cgaataccat	atcctctctg	acataaagac	1140
tcgacttgaa	caggaaattg	ctacttaccg	ccgccttctg	gaaggagaa	acgtaaaaac	1200
tacagaatat	cagttaagca	ccttggaaga	gagagatata	aagaaaacca	ggaagattaa	1260
gacagtctgt	caagaagtag	tgatggcaca	ggtcgtgtca	tctgaagtca	aagagggtga	1320
agaaaatatc	taaatagcta	ccagaaggag	atgctgctga	ggtttttgaa	gaaatttggc	1380
tataatctta	tctttgtctc	ctgcaagaaa	tcagccataa	gaaagcacta	tttaactctt	1440
gcagtgtatta	gaaggggtgg	ggtggcgggg	atcctattta	tcagactctg	taattgaata	1500
taaatgtttt	actcagagga	gctgc aaatt	gcctgcaaaa	atgaaatcca	gtgagcacta	1560
gaatatattaa	aacatcatta	ctgccatctt	tatcatgaag	cacatcaatt	acaagctgta	1620
gaccacctaa	tatcaatttg	taggtaatgt	tcctgaaaaa	tgcaatacat	ttcaattata	1680
ctaaacctca	caagtagag	gaatccatgt	aaattgcaaa	taaacacctt	tctaattttt	1740
tcctgtttct	gaattgtaaa	accccccttg	ggagtcctct	gtttcttatt	gagccaattt	1800
ctggg						1805

SEQ ID NO: 6 moltype = DNA length = 1447
 FEATURE Location/Qualifiers
 source 1..1447
 mol_type = genomic DNA
 organism = Homo sapiens

SEQUENCE: 6

atccagattt	gcttttacat	tttcttgct	gagctctgag	tgaacagtga	acatatattac	60
atttgattta	acagtgaacc	ttaattcttt	ctggtctcac	agtgaacaaa	gtttatgcaa	120
tcgatcaaat	attttcatcc	ctgaggttaa	caattaccat	caaaatgttt	tgtggagact	180
atgtgcaagg	aacctatctc	ccagctccca	atttcaatcc	cataatggat	gccccaaatgc	240
taggaggagc	actccaagga	tttgactgtg	acaaagacat	gctgatcaac	attctgactc	300
agcgctgcaa	tgacacaaag	atgatgattg	cagaggcata	ccagagcatg	tatggccggg	360
acctgatttg	ggatgatgag	gagcagcttt	cggatcactt	caaagatgtg	atggctggcc	420
tcagtgtacc	accaccactg	tatgatgtct	atgagctctg	gcctgcatg	aagggagtag	480
gcactgtatg	gaattgcctc	attgaaatac	tagcttcaag	aacaaatgga	gaaattttcc	540
agatgcgaga	agcctactgc	ttgcaataca	gcaataacct	ccaagaggac	atttattcag	600
agacctcagg	acacttcaga	gatactctca	tgaacttggt	ccaggggacc	agagagggaag	660
gatatacaga	ccctgcgatg	gctgctcagg	atgcaatggt	cctatgggaa	gcctgtcagc	720
agaagacggg	ggagcacaaa	accatgctgc	aaatgatctc	gtgcaacaag	agctaccagc	780
agctgcggct	gggtttccag	gaatttcaaa	atatctctgg	gcaagatag	gtagatgcca	840
ttaatgaatg	ttatgatgga	tactttcagg	agctgctggt	tgcaattgtt	ctctgtgttc	900
gagacaaacc	agcctatttt	gcttatagat	tatatagtgc	aattcatgac	tttgggtttcc	960
ataataaaa	tgtaatcagg	attctcattg	ccagaagtga	aatagacctg	ctgaccataa	1020
ggaaacgata	caaagagcga	tatggaaaat	ccctatttca	tgatatcaga	aatttttgctt	1080
cagggcatta	taagaaagca	ctgcttgcca	tctgtgctgg	tgatgctgag	gactactaaa	1140
atgaagagga	cttgaggtac	tgtgcactcc	tctttctaga	cacttccaaa	tagagatttt	1200
ctcacaattt	tgtactgttc	atggcactat	taacaaaact	atacaatcat	attttctctt	1260
ctatctttga	aattttcta	agccaaagaa	aactatgaat	gaaagtatat	gatactgaat	1320
ttgcctacta	tcctgaattt	gcctactatc	taatcagcaa	ttaaataaat	tgtgcatgat	1380
ggaataatag	aaaaattgca	ttggaataga	ttttatttaa	atgtgaacca	tcaacaacct	1440
acaacaa						1447

SEQ ID NO: 7 moltype = DNA length = 5881

-continued

FEATURE	Location/Qualifiers				
source	1..5881				
	mol_type = genomic DNA				
	organism = Homo sapiens				
SEQUENCE: 7					
ggttggtgac	ttccacagga	aaagtctctg	aggagtagcc	aaagaccatc	agcgtttcct 60
ttatgtgtga	gaattgaaat	gactagcatt	attgaccctt	ttcagcatcc	cctgtgaata 120
ttctgtttaa	ggtttttctt	ctgaaaaaga	aattgttatt	cagcccgttt	aaaacaaatc 180
aagaactttt	tgggtaacat	tgcaattaca	tgaattgat	aaccgcgaaa	ataattggaa 240
ctcctgcttg	caagtgtcaa	ccataaaaaa	gtgcttcctt	ttgttatgga	agatgtcttt 300
ctgtgattga	cttcaattgc	tgacttgtgg	agatgcagcg	aatgtgaaat	cccacgtata 360
tgccatttcc	ctctacgcct	gctgaccgtt	ctggaagatc	ttgaaccctc	ttctggaaag 420
gggtacctat	tattacttta	tggggcagca	gcctggaaaa	gtacttgggg	accaaagaag 480
gccaagcttg	cctgocctgc	attttatcaa	aggagcaggg	aagaaggaaat	catcgaggca 540
tgggggtcca	cactgcaatg	tttttgtgga	acatgaagcc	cttcagcgcc	cagtagcatc 600
tgactttgag	cctcagggtc	tgagtgaagc	cgctcggttg	aactccaagg	aaaaccttct 660
cgctggagcc	agtgaaaaat	accccaacct	tttcgttgca	ctgtatgatt	tttgggccag 720
tggagataac	actctaagca	taactaaagg	tgaaaagctc	cggtctctag	gctataatca 780
caatggggaa	tggtgtgaag	cccaaaccaa	aaatggccaa	ggctgggtcc	caagcaacta 840
catcacgcca	gtcaacagtc	tggagaaaac	ctcctggtag	catgggcctg	tgtcccgcaa 900
tgccgctgag	tatctgtgta	gcagcgggat	caatggcagc	ttcttggtgc	gtgagagtga 960
gagcagtcct	ggcagaggtg	ccatctcgct	gagatacgaa	gggaggggtg	accattacag 1020
gatcaaacct	gctctgtgat	gcaagctcta	cgctctctcc	gagagccgct	tcaacacctc 1080
ggccgagttg	gttcatcatc	attcaacggt	ggccgacggg	ctcatcacca	cgctccatta 1140
tcacgcccc	aagcgcaaca	agccactgt	ctatggtgtg	tccccaaact	acgacaaagt 1200
ggagatggaa	gcacccgaca	tcacatgaa	gcacaagctg	ggcggggggc	agtcacggga 1260
ggtgtacgag	ggcgtgtgga	agaaatacag	cctgacgggt	ggcgtgaaga	ccttgaagga 1320
ggacaccatg	gaggtggaag	agttcttgaa	agaagctgca	gtcatgaaag	agatcaaaac 1380
ccctaaccctg	gtgcagctcc	ttggggctcg	cacccgggag	cccccgctct	atatcatcac 1440
tgagttcatg	acctacggga	acctcctgga	ctacctgagg	gagtgcaacc	ggcaggaggt 1500
gaacgccgtg	gtgctgctgt	acatggccac	tcagatctcg	tcagccatgg	agtacctgga 1560
gaagaaaaac	ttcatccaca	gagatcttgc	tgcccgaaac	tgectggtag	gggagaacca 1620
cttggtgaag	gtagctgatt	ttggcctgag	caggttgatg	acaggggaca	cctacacagc 1680
ccatgctgga	gccaagtctc	ccatcaaatg	gactgcaccc	gagagcctgg	cctacaacaa 1740
gttctccatc	aagtcgcagc	tctgggcatt	tggaagtatt	ctttgggaaa	ttgtacctta 1800
tggtcatgtc	ccttaccggt	gaattgacct	gtcccagggt	tatgagctgc	tagagaagga 1860
ctaccgcatt	gagcgcccag	aaggctgccc	agagaaggtc	tatgaactca	tgcgagcatt 1920
ttggcagttg	aatccctctg	accggccctc	ctttgtctga	atccaccaag	cctttgaaac 1980
aatgttccag	gaatccagta	tctcagacga	agtggaaaaa	gagctgggga	aacaaggcgt 2040
ccgtggggct	gtgagtaact	tgctgcaggc	cccagagctg	cccaccaaga	cgaggacctc 2100
caggagagct	gcagagcaca	gagacaccac	tgacgtgcct	gagatgcctc	actccaaggg 2160
caggggagag	agcgatccct	tggaccatga	gcctgcccgt	tctccattgc	tcctccgaaa 2220
agagcgaggt	ccccggagg	gcggcctgaa	tgaagatgag	cgctctctcc	ccaaagacaa 2280
aaagaccaac	ttgttcagcg	ccttgatcaa	gaagaagaag	aagacagccc	caacccctcc 2340
caaacgcagc	agctccttcc	gggagatgga	cgccagcccg	gagcgacagc	gggcccggga 2400
ggaagagggc	gcagacatga	gcaacggggc	actggcttcc	acccctctgg	acacagctga 2460
cccagccaa	tcctccaaag	ccagcaatgg	ggctggggct	cccaatggag	ccctccggga 2520
gtccgggggc	tcaggcttcc	ggctctccca	cctgtggaag	aagtcacaga	cgctgaccag 2580
agccgcctta	gcccggcg	aggaggagg	cggtggcagc	tcacgcaagc	gcttctcgcg 2640
ctcttgctcc	gcccctctgc	ttcccatgg	ggccaaaggc	acggagtggg	ggtcagtcac 2700
gctgcctcgg	gacttgacgt	ccacgggaag	acagtttgac	tcgtccacat	ttggagggca 2760
caaaagttag	aagccggctc	tgctcggaa	gagggcaggg	gagaacaggt	ctgaccaggt 2820
gacccgaggg	acagttaacg	ctccccccag	gctggtgaaa	aagaatgagg	aagctgctga 2880
tgaggtcttc	aaagacatca	tggagtccag	ccggggctcc	agcccgccca	acctgactcc 2940
aaaaacccctc	cgccggcagg	tcacctgggc	ccctgcctcg	ggcctccccc	acaagggaaga 3000
agctggaaag	ggcagtgctc	tagggacccc	tgctgcagct	gagccagtga	ccccaccag 3060
caaaagcagg	tcagggtcac	cagggggcac	cagcaagggc	cccgcagagg	agtcacaggt 3120
gaggaggcac	aagcactcct	ctgagtcgcc	agggaaggac	aaggggaaat	tgtccaggct 3180
caaacctgcc	ccgcgcgcgc	caccagcagc	ctctgcaggg	aaggctggag	gaaagccctc 3240
gcagagcccg	agccaggagg	cgccggggga	ggcagtcctg	ggcgcaaaag	caaaagccac 3300
gagtcctggt	gatgctgtga	acagtgaacg	tgccaagccc	agccagcccg	gagagggcct 3360
caaaaagccc	gtgctcccg	ccactccaaa	gccacagtc	gccaaagcgt	cggggacccc 3420
catcagccca	gcccccgctc	cctccacgtt	gccatcagca	tcctcgcccc	tgccagggga 3480
ccagccgtct	tcacccgctc	catatccctc	catatcaacc	cgagtgtctc	ttcggaatac 3540
ccgccagcct	ccagagcgga	tcgccagcgg	cgccatcacc	aaggggctgg	tcctggacag 3600
caccgaggcg	ctgtgcctcg	ccatctctag	gaactccgag	cagatggcca	gccacagcgc 3660
agtgtcggag	gcccggcaaaa	acctctacac	gttctgcgtg	agctatgtgg	attccatcca 3720
gcaaatgagg	aacaagtttg	ccttccgaga	ggccatcaac	aaactggaga	ataatctccg 3780
ggagcttcag	atctgcccgg	cgacagcagg	cagtgggtcca	gcggccactc	aggacttcag 3840
caagctcttc	agttcggtga	aggaaatcag	tgacatagtg	cagaggtagc	agcagtcagg 3900
ggtcaggtgt	caggcccgct	ggagctgcct	gcagcacatg	cggtctcgcc	cataccgctg 3960
acagtggctg	acaagggact	agtgagtcag	caccttggcc	caggagctct	gcgccaggca 4020
gagctgaggg	ccctgtggag	tcagctctta	ctacctacgt	ttgcaccgcc	tgccctcccg 4080
caacctctct	ctccccgcct	cgtctctgtc	ctcgaatttt	atctgtggag	ttcctgctcc 4140
gtggactgca	gtcggcatgc	caggacccgc	cagcccccgt	cccacctagt	gccccagact 4200
gagctctcca	ggccaggtgg	gaacggctga	tgtggactgt	ctttttcatt	tttttctctc 4260
tggagccctc	cctcccccg	ctgggcctcc	ttcttccact	tctccaagaa	tggaagcctg 4320
aaactgagcc	ttgtgtgtca	ggccctctgc	ctgcactccc	tgcccttgcc	cgctggtgtc 4380
tgaagacatg	tttcaagaac	cgcatttcgg	gaagggcatt	cacgggcatt	cacacggctg 4440

-continued

```

gtcactctgc cctctgtgc tgcgggggt ggggtgact cgccatttcc tcacgtgcag 4500
gacagctctt gatttgggtg gaaaacaggg tgctaaagcc aaccagcctt tgggtcctgg 4560
gcaggtggga gctgaaaagg atcgaggcat ggggcatgtc ctttccatct gtccacatcc 4620
ccagagccca gctcttgctc tcttgtgacg tgcactgtga atcctggcaa gaaagcttga 4680
gtctcaaggg tggcaggtca ctgtcactgc cgacatccct ccccagcag aatggaggca 4740
ggggacaagg gaggcagtggt ctagtgggggt gaacagctgg tgccaaatag ccccagactg 4800
ggcccaggca ggtctgcaag ggcccagagt gaaccgtcct ttcacacatc tgggtgccct 4860
gaaagggccc ttccctccc ccactcctct aagacaaagt agattcttac aaggcccttt 4920
cctttggaac aagacagcct tcacttttct gaggttctga agcatttcaa agccctgcct 4980
ctgtgtagcc gcctgagag agaataagac tgccactggg cactgcgca caggtgggag 5040
gaaagggcct ggccagtcct ggtcctggct gcactcttga actgggcgaa tgtcttattt 5100
aattaccgtg agtgacatag cctcatgttc tgtgggggtc atcagggagg gttaggaaaa 5160
ccacaaacgg agccctgaa agcctcacgt atttcacaga gcacgcctgc catcttctcc 5220
ccgagggctgc cccaggccgg agcccagata cgggggctgt gactctgggc agggacccgg 5280
ggtctcctgg accttgacag agcagctaac tccgagagca gtgggcaggt gcccgccct 5340
gaggettccac gccgggagaa gccaccttcc cacccttca taccgcctcg tgccagcagc 5400
ctcgacacag ccctagcttt acgctcatca cctaaacttg tactttattt ttctgataga 5460
aatggtttcc tctggatcgt tttatgcggt tcttacagca catcacctct ttgccccga 5520
cggctgtgac gcagccggag ggaggcacta gtcaccgaca gcggccttga agacagagca 5580
aagcgcccac ccaggctccc cgactgcctg tctccatgag gtactgttcc cttccttttg 5640
ttaacgtgat gtgccactat attttacacg tatctcttgg tatgcactct ttatagacgc 5700
tcttttctaa gtggcgtgtg catagcgtcc tgccctgccc cctcgggggc ctgtggtggc 5760
tccccctctg cttctcgggg tccagtgcac tttgtttctg tatatgattc tctgtgggtt 5820
tttttgaatc caaatctgtc ctctgtagta ttttttaaat aaatcagtgt ttacattaga 5880
a 5881

```

```

SEQ ID NO: 8      moltype = DNA length = 24
FEATURE          Location/Qualifiers
source           1..24
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 8
gatcaacact gctctgatg gcaa 24

```

```

SEQ ID NO: 9      moltype = DNA length = 25
FEATURE          Location/Qualifiers
source           1..25
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 9
ccaccgttga atgatgatga accaa 25

```

```

SEQ ID NO: 10     moltype = DNA length = 20
FEATURE          Location/Qualifiers
source           1..20
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 10
cctccgagag ccgcttcaac 20

```

```

SEQ ID NO: 11     moltype = DNA length = 20
FEATURE          Location/Qualifiers
source           1..20
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 11
cctccgagag ccgcttcaac 20

```

```

SEQ ID NO: 12     moltype = DNA length = 20
FEATURE          Location/Qualifiers
source           1..20
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 12
cctccgagag ccgcttcaac 20

```

```

SEQ ID NO: 13     moltype = DNA length = 22
FEATURE          Location/Qualifiers
source           1..22
                 mol_type = other DNA
                 organism = synthetic construct

```

```

SEQUENCE: 13
ttgaagagct gcgaagtcag at 22

```

```

SEQ ID NO: 14     moltype = DNA length = 21
FEATURE          Location/Qualifiers
source           1..21
                 mol_type = other DNA
                 organism = synthetic construct

```

-continued

SEQUENCE: 14		
tgaagtcctc agcagccagt t		21
SEQ ID NO: 15	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 15		
tcaactgcaa aatgctcggg gtgtcc		26
SEQ ID NO: 16	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 16		
cgcggtcttct acttcagcag		20
SEQ ID NO: 17	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 17		
gaggaaacag cactctctaa		20
SEQ ID NO: 18	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 18		
tgtgagccgt cgcagccgtg		20
SEQ ID NO: 19	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 19		
acccgggtca cctgcgaa		18
SEQ ID NO: 20	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 20		
ggactccgc ggacacaa		18
SEQ ID NO: 21	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 21		
tcctgggaag cgagtgcctt taa		23
SEQ ID NO: 22	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 22		
cctgggaagc gagtgccct aa		22
SEQ ID NO: 23	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 23		
ggctattctc ctcttgag at		22
SEQ ID NO: 24	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	

-continued

SEQUENCE: 24	organism = synthetic construct	
tgcttgagct ccagtccta ag		22
SEQ ID NO: 25	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 25		
agccgagaag gcggagctctg gc		22
SEQ ID NO: 26	moltype = DNA length = 27	
FEATURE	Location/Qualifiers	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 26		
gtgaaacaag tttatgcaat cgatcaa		27
SEQ ID NO: 27	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 27		
gattgaaatt gggagctggg aa		22
SEQ ID NO: 28	moltype = DNA length = 29	
FEATURE	Location/Qualifiers	
source	1..29	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 28		
tcatccctga ggtaacaat taccatcaa		29
SEQ ID NO: 29	moltype = DNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 29		
cgactacagt gcatattaca gacaa		25
SEQ ID NO: 30	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 30		
cagcagccag tttagcatta tcaa		24
SEQ ID NO: 31	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
source	1..26	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 31		
tcaactgcaa aatgctcggt gtgtcc		26
SEQ ID NO: 32	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 32		
ggaccgcggc ttctacttca		20
SEQ ID NO: 33	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 33		
ccaggtcaca gctgoggaa		19
SEQ ID NO: 34	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	

-continued

SEQUENCE: 34	mol_type = other DNA	
tgtgagccgt cgcagccgtg	organism = synthetic construct	20
SEQ ID NO: 35	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 35		
tcgaagcgc ctgggaagc		19
SEQ ID NO: 36	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 36		
ggactccgc ggacacaa		18
SEQ ID NO: 37	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 37		
tgcccctaac atgcggctgc c		21
SEQ ID NO: 38	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 38		
tcagcgctgc aatgcacaa		19
SEQ ID NO: 39	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 39		
ggccagccat cacatctttg aa		22
SEQ ID NO: 40	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 40		
tagagcatgt atggccggga cct		23
SEQ ID NO: 41	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 41		
gatcaacact gcttctgatg gcaa		24
SEQ ID NO: 42	moltype = DNA length = 25	
FEATURE	Location/Qualifiers	
source	1..25	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 42		
ccaccgttga atgatgatga accaa		25
SEQ ID NO: 43	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 43		
cctccgagag ccgcttcaac		20
SEQ ID NO: 44	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	

-continued

```

source                1..22
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 44
ggctattctc ctcttgccag at                                22

SEQ ID NO: 45         moltype = DNA length = 22
FEATURE              Location/Qualifiers
source                1..22
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 45
tgcttgagct ccagtcacct a g                                22

SEQ ID NO: 46         moltype = DNA length = 22
FEATURE              Location/Qualifiers
source                1..22
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 46
agccgagaag gcggagctcg gc                                22

SEQ ID NO: 47         moltype = RNA length = 500
FEATURE              Location/Qualifiers
source                1..500
                      mol_type = other RNA
                      organism = synthetic construct

SEQUENCE: 47
gatgtcact tcctctatgg ttaccctggg acttttacac caacagaact agcatcatcc 60
ctgcatggg caggtcatgg atcgccatcc tgacagtctc gggaattagg catctgcagt 120
cttactgctc atcggtgat gatgtgctg taatcccat ccaagcaagc ttgtgatcct 180
ccgccattat cccaaatggg ataacattta ggacttaaa ctagcaatt atcaccttgt 240
ttttcaacag caagacctaa tattttcttt tcatcattaa tgccttttga tggatcaggc 300
aaccatttat aaatatgttc accagccgaa gtcagtagtg attgggtggt tcctggcttg 360
ggatcatgcc gctgcagagg ctattctctc cttggcagat tgtctgtagc cgagaaggcg 420
gagtctggca atgatgatgc atacagtgtg cgacagcctt agggactgga gctcaagcag 480
tgttctctca accagtcaca                                     500

SEQ ID NO: 48         moltype = DNA length = 20
FEATURE              Location/Qualifiers
source                1..20
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 48
ctgcaatgca caaaggatga                                     20

```

40

What is claimed is:

1. A method for detecting bladder cancer markers in a subject comprising detecting the levels of each bladder cancer marker amplicon of a set of bladder cancer markers in a urine sample or bladder washing sample from the subject, wherein the set of bladder cancer markers consists of corticotrophin releasing hormone (CRH), insulin-like growth factor 2 (IGF2), keratin 20 (KRT20) and annexin 10 (ANXA10), by contacting RNA from the sample with a set of primers, conducting one or more polymerase chain reactions (PCR) and detecting a set of bladder cancer marker amplicons that is produced by the PCR.

2. The method of claim 1, wherein the method further comprises detecting an endogenous control and/or exogenous control.

3. The method of claim 2, wherein the endogenous control is selected from ABL, GUSB, GAPDH, TUBB, and UPK1.

4. The method of claim 2, wherein the exogenous control is an RNA.

5. The method of claim 2, wherein the method comprises comparing a Ct value or a Δ Ct value to a threshold Ct value or Δ Ct value, wherein Δ Ct is the Ct value for the control minus the Ct value for the marker.

6. The method of claim 1, wherein the detecting comprises RT-PCR.

7. The method of claim 1, wherein the detecting comprises a RT-PCR reaction that takes less than 2 hours from an initial denaturation step through a final extension step.

8. The method of claim 1, wherein the set of primers comprises a first and second primer for detecting CRH comprising:

a) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 19 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 20, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long; or

b) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 35 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 36, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

9. The method of claim 1, wherein the set of primers comprises a first and second primer for detecting IGF2 comprising:

a) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 16 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 17, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long; or

65

91

- b) a first primer comprising at least 12 contiguous nucleotides of at least 12 contiguous nucleotides of SEQ ID NO: 32 and a second primer comprising SEQ ID NO: 33, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

10. The method of claim 1, wherein the set of primers comprises a first and second primer for detecting KRT20 comprising:

- a) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 13 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 14, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long; or
 b) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 29 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 30, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

11. The method of claim 1, wherein the set of primers comprises a first and second primer for detecting ANXA10 comprising:

- a) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 26 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 27, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long; or
 b) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 38 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 39, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long; or
 c) a first primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 48 and a second primer comprising at least 12 contiguous nucleotides of SEQ ID NO: 39, wherein each primer is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

12. The method of claim 1, wherein the set of bladder cancer marker amplicons comprises a CRH amplicon, an IGF2 amplicon, a KRT20 amplicon, and an ANXA10 amplicon, and wherein the method comprises contacting the

92

bladder cancer marker amplicons with a set of bladder cancer marker probes, wherein the set of bladder cancer marker probes comprises a probe for detecting the CRH amplicon, a probe for detecting the IGF2 amplicon, a probe for detecting the KRT20 amplicon, and a probe for detecting the ANXA10 amplicon.

13. The method of claim 12, wherein the probe for detecting CRH comprises at least 12 contiguous nucleotides of SEQ ID NO: 21, SEQ ID NO: 22, or SEQ ID NO: 37, wherein the probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

14. The method of claim 12, wherein the probe for detecting IGF2 comprises at least 12 contiguous nucleotides of SEQ ID NO: 34 or at least 12 SEQ ID NO: 18, wherein the probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

15. The method of claim 12, wherein the probe for detecting KRT20 comprises at least 12 contiguous nucleotides of SEQ ID NO: 15 or SEQ ID NO: 31, wherein the probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

16. The method of claim 12, wherein the fourth probe for detecting ANXA10 comprises at least 12 contiguous nucleotides of SEQ ID NO: 28 or SEQ ID NO: 40, wherein the probe is less than 50, less than 45, less than 40, less than 35, or less than 30 nucleotides long.

17. The method of claim 12, wherein each bladder cancer marker probe comprises a dye, and wherein each dye is detectably different from the other dyes.

18. The method of claim 17, wherein each probe comprises a quencher molecule.

19. The method of claim 12, wherein the method further comprises forming an endogenous control amplicon, and contacting the endogenous control amplicon with an endogenous control probe, and/or forming an exogenous control amplicon, and contacting the exogenous control amplicon with an exogenous control probe, wherein each probe comprises a dye, and wherein each dye is detectably different from the other dyes.

20. The method of claim 1, wherein the set of bladder cancer markers are detected in a single multiplex reaction.

21. The method of claim 1, wherein the subject has at least one symptom of bladder cancer and/or has a history of bladder cancer.

* * * * *