



US012405265B2

(12) **United States Patent**
Brafman et al.(10) **Patent No.: US 12,405,265 B2**(45) **Date of Patent: Sep. 2, 2025**(54) **TRANSIENT REPORTERS AND METHODS FOR BASE EDITING ENRICHMENT**(71) Applicant: **ARIZONA BOARD OF REGENTS ON BEHALF OF ARIZONA STATE UNIVERSITY**, Scottsdale, AZ (US)(72) Inventors: **David Brafman**, Phoenix, AZ (US); **Xiao Wang**, Chandler, AZ (US); **Nicholas Brookhouser**, Tempe, AZ (US); **Stefan Tekel**, Tempe, AZ (US); **Kylie Standage-Beier**, Phoenix, AZ (US)(73) Assignee: **ARIZONA BOARD OF REGENTS ON BEHALF OF ARIZONA STATE UNIVERSITY**, Scottsdale, AZ (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 212 days.

(21) Appl. No.: **17/347,360**(22) Filed: **Jun. 14, 2021**(65) **Prior Publication Data**

US 2021/0389303 A1 Dec. 16, 2021

Related U.S. Application Data

(60) Provisional application No. 63/038,220, filed on Jun. 12, 2020.

(51) **Int. Cl.****G01N 33/50** (2006.01)**C07K 14/435** (2006.01)**C12N 9/22** (2006.01)**C12N 15/11** (2006.01)**C12N 15/90** (2006.01)(52) **U.S. Cl.**CPC ... **G01N 33/5073** (2013.01); **C07K 14/43595** (2013.01); **C12N 9/22** (2013.01); **C12N 15/11** (2013.01); **C12N 15/907** (2013.01); **C12N 2310/20** (2017.05); **C12N 2800/80** (2013.01)(58) **Field of Classification Search**CPC **G01N 33/5073**; **C07K 14/43595**; **C12N 9/22**; **C12N 15/11**; **C12N 15/907**; **C12N 2310/20**; **C12N 2800/80**; **C12N 15/90**
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**8,987,257 B2 * 3/2015 Radetich A61P 25/28
544/118

2020/0032251 A1 * 1/2020 McLaughlin C12N 15/907

OTHER PUBLICATIONSGauthier, Nicholas P., et al. "Cell-selective labeling using amino acid precursors for proteomic studies of multicellular environments." *nature methods* 10.8 (2013): 768-773. (Year: 2013).*

Synthetic construct mCherry-P2A-wild type lysine racemase (mCherry-P2A-wild-type-lyr) gene, complete cds GenBank: KC962563.1 (Year: 2013).*

Standage-Beier, Kylie, et al. "A transient reporter for editing enrichment (TREE) in human cells." *Nucleic acids research* 47.19 (2019): e120-e120. (Year: 2019).*Anzalone et al., Search-and-replace genome editing without double-strand breaks or donor DNA, *Nature*, 2019, 576:149-157.Billon et al., CRISPR-mediated base editing enables efficient disruption of eukaryotic genes through induction of STOP codons, *Molecular Cell*, 2017, 67:1068-1079.Boettcher et al., Choosing the Right Tool for the Job: RNAi, TALEN, or CRISPR, *Molecular Cell*, 2015, 58:575-585.Brookhouser et al., BIG-TREE: Base-Edited Isogenic hPSC Line Generation Using a Transient Reporter for Editing Enrichment, *Stem Cell Reports*, 2020, 14:184-191.Brookhouser et al., May I Cut in? Gene Editing Approaches in Human Induced Pluripotent Stem Cells, *Cells*, 2017, vol. 6, pp. 1-21.Carlson-Stevermer et al., Genome editing in human pluripotent stem cells, *Methods Mol. Biol.*, 2017, 1590:165-174.Chapman et al., Playing the end game: DNA double-strand break repair pathway choice, *Molecular Cell*, 2012, 47:497-510.Chen et al., Enhanced proofreading governs CRISPR-Cas9 targeting accuracy, *Nature*, 2017, 550:407-410.Cheng et al., Expanding C-T base editing toolkit with diversified cytidine deaminases, *Nature Communications*, 2019, 10:3612, pp. 1-10.Coelho et al., BE-FLARE: a fluorescent reporter of base editing activity reveals editing characteristics of APOBEC3A and APOBEC3B, *BMC Biol.*, 2018, 16:150, pp. 1-11.Dai et al., The genes associated with early-onset Alzheimer's disease, *Oncotarget*, 2018, 9:15132-15143.Eggenchwil et al., Improved bi-allelic modification of a transcriptionally silent locus in patient-derived iPSC by Cas9 nickase, *Sci Rep*, 2016, 6:38198, pp. 1-14.Gaudelli et al., Programmable base editing of A,T to G,C in genomic DNA without DNA cleavage, *Nature*, 2017, 551:464-471.Gaudelli et al., Directed evolution of adenine base editors with increased activity and therapeutic application, *Nature Biotechnology*, 2020, 38(7):892-900.Glaser et al., GFP to BFP Conversion: A Versatile Assay for the Quantification of CRISPR/Cas9-mediated Genome Editing, *Molecular Therapy—Nucleic Acids*, 2016, 5:e334, 4 pages.Grunewald et al., A dual-deaminase CRISPR base editor enables concurrent adenine and cytosine editing, *Nature Biotechnology*, 2020, 38(7):861-864.Gurumurthy et al., Creation of CRISPR-based germline-genome-engineered mice without ex vivo handling of zygotes by i-GONAD, *Nature Protocols*, 2019, 14(8):2452-2482.

(Continued)

Primary Examiner — Kara D Johnson*Assistant Examiner* — Constantina E Stavrou(74) *Attorney, Agent, or Firm* — Quarles & Brady, LLP(57) **ABSTRACT**

Provided herein are compositions and methods for real-time identification and isolation of base-edited cell populations. Also provided herein are methods for producing enriched isogenic lines of genetically modified cells, including base-edited human pluripotent stem cells. In particular, provided herein are methods utilizing transient expression of reporter proteins, the detectable signal of which is altered following base editing. Using the transient reporter with a base editor permits enrichment of isogenic populations of base-edited cells.

1 Claim, 57 Drawing Sheets**Specification includes a Sequence Listing.**

(56)

References Cited**OTHER PUBLICATIONS**

- Haapaniemi et al., CRISPR-Cas9 genome editing induces a p53-mediated DNA damage response. *Nat. Med.*, 2018, 24:927-930.
- Hauser et al., Impact of apolipoprotein E on Alzheimer's disease, *Curr. Alzheimer Res.*, 2013, 10:809-817.
- He et al., Knock-in of large reporter genes in human cells via CRISPR/Cas9-induced homology-dependent and independent DNA repair, *Nucleic Acids Res.*, 2016, 44:e85, 14 pages.
- Hockemeyer et al., Induced Pluripotent Stem Cells Meet Genome Editing, *Cell Stem Cell*, 2016, 18:573-586.
- Hsu et al., Development and applications of CRISPR-Cas9 for genome engineering, *Cell*, 2014, 157:1262-1278.
- Hu et al., Evolved Cas9 variants with broad PAM compatibility and high DNA specificity, *Nature*, 2018, 556:57-63.
- Huang et al., Developing ABEmax-NG with Precise Targeting and Expanded Editing Scope to Model Pathogenic Splice Site Mutations In Vivo, *iScience*, 2019, 15:640-648.
- Huang et al., Circularly permuted and PAM-modified Cas9 variants broaden the targeting scope of base editors, *Nat. Biotechnol.*, 2019, 37:626-631.
- Ihry et al., Genome-scale CRISPR screens identify human pluripotency-specific genes, *Cell Rep.*, 2019, 27:616-630.
- Ihry et al., p53 inhibits CRISPR-Cas9 engineering in human pluripotent stem cells, *Nat. Med.*, 2018, 24:939-946.
- Katti et al., GO: a functional reporter system to identify and enrich base editing activity, *Nucleic Acids Res.*, 2020, 48:2841-2852.
- Khera et al., Genome-wide polygenic scores for common diseases identify individuals with risk equivalent to monogenic mutations, *Nat. Genet.*, 2018, 50:1219-1224.
- Kim et al., Genome-wide target specificity of CRISPR RNA-guided adenine base editors, *Nature Biotechnology*, 2019, 37:430-435.
- Kim et al., CRISPR/Cpf1-mediated DNA-free plant genome editing, *Nature Communications*, 2017, 8:14406, pp. 1-7.
- Kim et al., Highly efficient RNA-guided base editing in mouse embryos, *Nature Biotechnology*, 2017, 35:435-437.
- Kim et al., Increasing the genome-targeting scope and precision of base editing with engineered Cas9-cytidine deaminase fusions, *Nature Biotechnology*, 2017, 35:371-376.
- Kluesner et al., EditR: A Method to Quantify Base Editing from Sanger Sequencing, *CRISPR Journal*, 2018, 1:239-250.
- Kobayashi et al., Pathogenic variant burden in the ExAC database: an empirical approach to evaluating population data for clinical variant interpretation, *Genome Med.*, 2017, 9:13, pp. 1-14.
- Koblan et al., Improving cytidine and adenine base editors by expression optimization and ancestral reconstruction, *Nat. Biotechnol.*, 2018, 36:843-846.
- Komor et al., Editing the genome without double-stranded DNA breaks, *ACS Chem. Biol.*, 2018, 13:383-388.
- Komor et al., Programmable editing of a target base in genomic DNA without double-stranded DNA cleavage, *Nature*, 2016, 533:420-424.
- Komor et al., Improved base excision repair inhibition and bacteriophage Mu Gam protein yields C:G-to-T:A base editors with higher efficiency and product purity, *Science Advances*, 2017, 3:eao4774, pp. 1-9.
- Kosicki et al., Repair of double-strand breaks induced by CRISPR-Cas9 leads to large deletions and complex rearrangements, *Nat. Biotechnol.*, 2018, 36:765-771.
- Kwart et al., Precise and efficient scarless genome editing in stem cells using CORRECT, *Nat. Protoc.*, 2017, 12:329-354.
- Landrum et al., ClinVar: improving access to variant interpretations and supporting evidence, *Nucleic Acids Res.*, 2018, 46:D1062-D1067.
- Lee et al., CRISPR-pass: Gene rescue of nonsense mutations using adenine base editors, *Molecular Therapy*, 2019, 27(8):1364-1371.
- Lee et al., Simultaneous targeting of linked loci in mouse embryos using base editing, *Scientific Reports*, 2019, 9:1662, pp. 1-8.
- Levy et al., Cytosine and adenine base editing of the brain, liver, retina, heart and skeletal muscle of mice via adeno-associated viruses, *Nature Biomedical Engineering*, 2020, 4(1):97-110.
- Li et al., CRISPR-Cas9-mediated base-editing screening in mice identifies DND1 amino acids that are critical for primordial germ cell development, *Nature Cell Biology*, 2018, 20(11):1315-1325.
- Li et al., Base editing with a Cpf1-cytidine deaminase fusion, *Nature Biotechnology*, 2018, 36(4):324-327.
- Lin et al., Prime genome editing in rice and wheat, *Nature Biotechnology*, 2020, 38(5):582-585.
- Lino et al., Delivering CRISPR: a review of the challenges and approaches, *Drug Deliv.*, 2018, 25:1234-1257.
- Liu et al., Delivery strategies of the CRISPR-Cas9 geneediting system for therapeutic applications, *J Control Release*, 2017, 266:17-26.
- Liu et al., Efficient generation of mouse models of human diseases via ABE- and BE-mediated base editing, *Nat Commun*, 2018, 9:2338, pp. 1-8.
- Liu et al., Highly efficient base editing with expanded targeting scope using SpCas9-NG in rabbits, *FASEB J.*, 2020, 34:588-596.
- Logue et al., Two rare AKAP9 variants are associated with Alzheimer's disease in African Americans, *Alzheimers Dement*, 2014, 10:609-618.
- Ma et al., Targeted AID-mediated mutagenesis (TAM) enables efficient genomic diversification in mammalian cells, *Nature Methods*, 2016, 13(12):1029-1035.
- Martin et al., A panel of eGFP reporters for single base editing by APOBEC-Cas9 editosome complexes, *Sci Rep*, 2019, 9:497, pp. 1-8.
- Moon et al., Recent advances in the CRISPR genome editing tool set, *Exp. Mol. Med.*, 2019, 51:130, pp. 1-11.
- Nishida et al., Targeted nucleotide editing using hybrid prokaryotic and vertebrate adaptive immune systems, *Science*, 2016, 353(6305):1248, aaf8729-1 thru aaf8729-8.
- Ortmann et al., Variability of human pluripotent stem cell lines, *Curr. Opin. Genet. Dev.*, 2017, 46:179-185.
- Panyam et al., Biodegradable nanoparticles for drug and gene delivery to cells and tissue, *Advanced Drug Delivery Reviews*, 2012, 64:61-71.
- Paquet et al., Efficient introduction of specific homozygous and heterozygous mutations using CRISPR/Cas9, *Nature*, 2016, 533:125-129.
- Park et al., Disease-specific induced pluripotent stem cells, *Cell*, 2008, 134:877-886.
- Pau, A table of senile forms of cataract. Clinical picture, prognosis, biochemistry and etiology, *Klin Monbl Augenheilkd*, 1988, 193:550-553.
- Rees et al., Base editing: precision chemistry on the genome and transcriptome of living cells, *Nat Rev Genet*, 2018, 19:770-788.
- Rees et al., Development of hRad51-Cas9 nickase fusions that mediate HDR without double-stranded breaks, *Nat Commun*, 2019, 10:2212, pp. 1-12.
- Rees et al., Improving the DNA specificity and applicability of base editing through protein engineering and protein delivery, *Nature Communications*, 2017, 8:15790, pp. 1-10.
- Ryu et al., Adenine base editing in mouse embryos and an adult mouse model of Duchenne muscular dystrophy, *Nature Biotechnology*, 2018, 36(6):536-539.
- Sakata et al., Base editors for simultaneous introduction of C-to-T and A-to-G mutations, *Nature Biotechnology*, 2020, 38(7):865-869.
- Shimatani et al., Targeted base editing in rice and tomato using a CRISPR-Cas9 cytidine deaminase fusion, *Nature Biotechnology*, 2017, 35(5):441-443.
- Si et al., Impact of single nucleotide polymorphisms on P450 oxidoreductase and peroxisome proliferator-activated receptor alpha on tacrolimus pharmacokinetics in renal transplant recipients, *Pharmacogenomics Journal*, 2019, 19:42-52.
- Slaymaker et al., Rationally engineered Cas9 nucleases with improved specificity, *Science*, 2016, 351:84-88.
- Sod-Moriah et al., Long term effects of dibromochloropropane (DBCP) on male rats' reproductive system, *Andrologia*, 1988, 20:60-66.

(56)

References Cited**OTHER PUBLICATIONS**

St Martin et al., A fluorescent reporter for quantification and enrichment of DNA editing by APOBEC-Cas9 or cleavage by Cas9 in living cells, *Nucleic Acids Res.*, 2018, 46:e84, 10 pages.

Standage-Beier et al., A transient reporter for editing enrichment (TREE) in human cells, *Nucleic Acids Res.*, 2019, 47:e120, pp. 1-13.

Stemmer et al., CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target Prediction Tool, *PLoS ONE*, 2015, 10:e0124633, 11 pages.

Takayama et al., Highly efficient biallelic genome editing of human ES/IPS cells using a CRISPR/Cas9 or TALEN system, *Nucleic Acids Res.*, 2017, 45:5198-5207.

Tan et al., Engineering of high-precision base editors for site-specific single nucleotide replacement, *Nat. Commun.*, 2019, 10:439, pp. 1-10.

Thuronyi et al., Continuous evolution of base editors with expanded target compatibility and improved activity, *Nat. Biotechnol.*, 2019, 37:1070-1079.

Tong et al., Highly efficient DSB-free base editing for streptomycetes with CRISPR-BEST, *Proceedings of the National Academy of Sciences*, 2019, 116(41):20366-20375.

Traxler et al., A genome-editing strategy to treat B-hemoglobinopathies that recapitulates a mutation associated with a benign genetic condition, *Nat. Med.*, 2016, 22:987-990.

Van Rhee et al., Genetic correlations of polygenic disease traits: from theory to practice, *Nat. Rev. Genet.*, 2019, 20:567-581.

Wang et al., BEON: A Functional Fluorescence Reporter for Quantification and Enrichment of Adenine Base-Editing Activity, *Mol. Ther.*, 2020, 28(7):1696-1705.

Wang et al., Efficient Gene Silencing by Adenine Base Editor-Mediated Start Codon Mutation, *Mol. Ther.*, 2020, 28:431-440.

Wang et al., Expanding targeting scope, editing window, and base transition capability of base editing in *Corynebacterium glutamicum*, *Biotechnology and Bioengineering*, 2019, 116(11):3016-3029.

Xie et al., Efficient base editing for multiple genes and loci in pigs using base editors, *Nature Communications*, 2019, 10:2852, pp. 1-13.

Yang et al., Methods and applications of CRISPR/Cas system for genome editing in stem cells, *Cell Regen*, 2019, 8:33-41.

Yang et al., Engineering and optimising deaminase fusions for genome editing, *Nature Communications*, 2016, 7:13330, pp. 1-11.

Yu et al., Small Molecules Enhance CRISPR Genome Editing in Pluripotent Stem Cells, *Cell Stem Cell*, 2015, 16:142-147.

Zafra et al., Optimized base editors enable efficient editing in cells, organoids and mice, *Nature Biotechnology*, 2018, 36(9):888-893.

Zhang et al., Simultaneous zygotic inactivation of multiple genes in mouse through CRISPR/Cas9-mediated base editing, *Development*, 2018, 145, pp. 1-11.


Zhang et al., Dual base editor catalyzes both cytosine and adenine base conversions in human cells, *Nature Biotechnology*, 2020, 38(7):856-860.

Zheng et al., Highly efficient base editing in bacteria using a Cas9-cytidine deaminase fusion, *Commun Biol*, 2018, 1:32, pp. 1-6.


Zong et al., Precise base editing in rice, wheat and maize with a Cas9-cytidine deaminase fusion, *Nature Biotechnology*, 2017, 35(5):438-440.

* cited by examiner


FIGS. 1A-1F




FIGS. 2A-2F




FIGS. 4A-4C




FIGS. 5A-5D



FIGS. 6A-6I



FIGS. 7A-7D



FIGS. 8A-8C







FIG. 9

U6 Promoter
sg(BG) guide
sg(BG) hairpin
U6 Promoter
sg(NT) guide
sg(NT) hairpin


SEQ ID NO: 262

>pDI.sgRNA.sg(BG).sg(NT) 3492 bp
TCGCGCGTTTCGGTGATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCACAGCTTGCTGTGTA
GCGGATGCCGGGAGCAGACAAAGCCCGTCAGGGCGCGTCAGCGGGTGTTGGCGGGTGTCGGGGCTGGCTTAA
CTATGCGGCATCAGAGCAGATTGTACTGAGAGTGCACCATATGCGGTGTGAAATACCGCACAGATGCGTAAGGA
GAAAATACCGCATCAGGCGCCATTGCGCATTCAGGCTGCGCAACTGTGGGAAGGGCGATCGGTGCGGGCCCTC
TTGCGTATTACGCCAGCTGGCGAAAGGGGGATGTGCTGCAAGGCGATTAAAGTTGGGTAACGCCAGGGTTTTCC
CAGTCACGACGTTGTAAAACGACGGCCAGTGAATTCGAGGGCCCTATTTCCCATGATTCTTCATATTTGCATATA
CCTATACAAAGCTCTTAGAGAGATAATTTGCAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATACGTGAC
GTAGAAAGTAATAATTTCTTGGGTAGTTTTCAGATTTTAAATTAATGTTTAAATGGACTATCATATGCTTACCGTA
ACTTGAAGATATTTGATTTCTTGGCTTTATATATCTTTGCGAAAGGAGGAAACACCGACCCCAAGGCGGTGCGAGT
CTTGTTTTAGAGCTAGAAAATAGCAAGTTAAATAAGGCTAGTCCGTTTATCAACTTGAAAAAGTGGCACCAGCTCG
GTGCTTTTGTGTTTAGAGCTAGAAATAGCAAGTTAAATAAGGCTAGTCCGTTTATAGCGCGTCCGCCAATTCTG
CAGACAGAGAGGGCCCTATTTCCCATGATTCTTCATATTTGCATATACGATACAGGCTCTTAGADAGATAATTG
GAATTAATTTGACTGTAAACACAAAGATATTAGTACAAAATACGTGACGTAGAAAGTAATAATTTCTTGGGTAGTTT
GCAGTTTAAAAATTATGTTTAAATGGACTATCATATGCTTACCGTAAGTTGAAAGTATTTGATTTCTTGGCTTT
ATATATCTTGTGGAAAGGACGAAACACCGGGCTCTCGAGAGAGACCTGTTTATAGAGCTAGAAAATAGCAAGTTAA
TAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGCTCGGTGCTTTTGTGTTTAGAGCTAGAAAATAGCAA
GTTAAATAAGGCTAGTCCGTTTATAGCGCGTGCGCCAATTCTGCAGACAAAAGCTTGGCGTAATCATGGTCAT
AGCTGTTTCTGTGTGAAATTGTTATCCGCTCACAAATCCACACAACATACGAGCCGGAAGCATAAAGTGTAAG
CCTGGGTGCGCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCCGCTTCCAGTCGGGAAAC
CTGTGCTGCCAGCTGCATTAATGAATCGGCCAACGCGCGGGGAGAGGCGSTTTGCGTATTGGGCGCTCTTCCG
CTTCTCGCTCACTGACTCGCTGCGCTCGGTGCTTGGGTGCGGCGAGCGGTATCAGCTCACTCAAAGCGGT
AATACGGTTATCCACAGAATCAGGGGATAACGCAGGAAGAAGATGTGAGCAAAAGGCCAGCAAAAGGCCAGG
AACCGTAAAAAGGCCCGCTTGTGCGCTTTTCCATAGGCTCCGCCCCCTGACGAGCATCACAAAAATCGAGC
CTCAAGTCAGAGGAGTGGCGAAACCCGACAGGACTATAAAGATACCAAGGCGTTTCCCGCTGGAAGCTCCCTCGT
CGCTCTCTGTTCCGACCCCTGCCGCTTACCGGATACCTGTCCGCTTTCTCCCTTCCGGAAGCGTGGCGCTTTC
TCATAGCTCAGCTGTAGGTATCTCAGTTGCGGTGTAGTGTGTTGCTCCAAGCTGGGCTGTGTGCACGAACCCC
CCGTTCCAGCCCGACCGCTGCGCCTTATCCGTAAGTATCGTCTTGAATCCACCGGTAAGACACGACTTATCG
CCACTGGCAGCAGCTAGTAAGTAGTTGCGCAGTTAAGATTTGCGCAACGTTGTTGCCATTGCTACAGGCTTGAAGT
GGTGGCCTAACTACGCTACACTAGAAGAACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTTACCTTCGGA
AAAAGAGTTGGTAGCTCTTGATCCGCGCAAAACAAACCACCGCTGGTAGCGGTGGTTTTTTGTTTGCAAGCAGCA
GATTACGCGCAGAAAAAAGGATCTCAAGAAGATCCTTTGATCTTTCTACGGGGTCTGACGCTCAGTGGAAACGA
AAACTCACGTTAAGGGATTTTGGTCATGAGATTACAAAAAGGATCTTCACTAGATCCTTTTAAATTAATAAATGAA
GTTTTAAATCAATCTAAAGTATATAGTAAGTAACTGGTCTGACAGTTACCAATGCTTAATCAGTGAGGCACCTAT
CTCAGCGATCTGTCTATTTGTTTCATCCATAGTTGCGTACTCGCGCTCGTGTAGATAACTACGATACGGGAGGG
CTTACCATCTGGCCCCAGTGCTGCAATGATACCGCGAGACCCACGCTCACCAGCTCCAGATTATCAGCAATAA
ACGAGCCAGCCGGAAGGGCGGAGCGCAGAGTGGTCTGCAACTTTATCCGCTCCATCCAGTCTATTAATTGT
TGCCGGGAAGCTAGAGTAAGTAGTTGCGCAGTTAAGATTTGCGCAACGTTGTTGCCATTGCTACAGGCTATCGT
GGTGTACGCTCGTCTGTTTGGTATGGCTTCACTCAGCTCCGTTTCCCAACGATCAAGGCGAGTTACATGATCCC
CCATGTTGTGCAAAAAAGCGGTTAGCTCCTTGGGTCTCCGATCGTTGTGCAAGTAAGTTGGCCGCAAGTGTAT
CACTCATGGTTATGGCAGCACTGCATAATTTCTTACTGTCTATGCCATCCGTAAGATGCTTTTCTGTGACTGGTGA
GTACTCAACCAAGTCTGAGAAATAGTGTATGCGGCGACCGAGTTGCTCTTGGCCGGCGTCAACCGGATTA
ATACCGCGCCACATAGCAGAACTTTAAAGTGCTCATTCATTGAAAAACGTTCTTGGGGCGGAAAACTCTCAAGGA
TCTTACCGCTGTTGAGATCCAGTTCGATGTAACCCACTCGTGCACCCAACGATCTTACGATCTTTTACTTTTAC
CAGCGTTTCTGGGTGAGCAAAAAACAGGAAGGCAAAATGCCGCAAAAAAGGGAATAAGGGCGACACGGAAATGT
TGAATGCTCATACTTCTCTTTTCAATATTATTGAAGCATTTATCAGGGTTATTGTCTCATGAGCGGATACATATT
TGAATGTATTAGAAAAATAAAATAAGGGGTTCCGCGCAATTTCCCGGAAAAAGTGCCACCTGACGCTTAAGA
AACCATTATTATCATGACATTAACCTATAAAAAATAGGCGTATCAGGAGGCCCTTTCGTC


FIGS. 13A-13E




FIGS. 14A-14B




FIGS. 15A-15C



FIGS. 16A-16D



FIGS. 17A-17C



FIGS. 18A-18B




FIG. 19




FIG. 20




FIG. 21




FIG. 23

[illegible]

FIG. 24




FIG. 26




FIG. 27

Untransfected Site-1

a

	Editing window	% of Reads		Editing window	% of Reads
SEQ ID NO: 263	WT	99.7	SEQ ID NO: 279	GGCCAGACTGAGCAGGTGATGG	5.2
SEQ ID NO: 264	18	0.1	SEQ ID NO: 280	GGTCCAGACTGAGCAGGTGATGG	0.0
SEQ ID NO: 265	17	0.1	SEQ ID NO: 281	GGCTCAGACTGAGCAGGTGATGG	0.1
SEQ ID NO: 266	16	0.1	SEQ ID NO: 282	GGCTTAGACTGAGCAGGTGATGG	0.3
SEQ ID NO: 267	12	0.0	SEQ ID NO: 283	GGCCAGATTGAGCAGGTGATGG	0.0
SEQ ID NO: 268	18,17	0.0	SEQ ID NO: 284	GGTTCAGACTGAGCAGGTGATGG	0.0
SEQ ID NO: 269	18,16	0.0	SEQ ID NO: 285	GGTCTAGACTGAGCAGGTGATGG	0.0
SEQ ID NO: 270	18,12	0.0	SEQ ID NO: 286	GGTCCAGATTGAGCAGGTGATGG	0.0
SEQ ID NO: 271	17,16	0.0	SEQ ID NO: 287	GGCTTAGACTGAGCAGGTGATGG	85.2
SEQ ID NO: 272	17,12	0.0	SEQ ID NO: 288	GGCTCAGATTGAGCAGGTGATGG	0.1
SEQ ID NO: 273	16,12	0.0	SEQ ID NO: 289	GGCCTAGATTGAGCAGGTGATGG	0.2
SEQ ID NO: 274	18,17,16	0.0	SEQ ID NO: 290	GGTTTAGACTGAGCAGGTGATGG	6.2
SEQ ID NO: 275	17,16,12	0.0	SEQ ID NO: 291	GGCTTAGATTGAGCAGGTGATGG	2.4
SEQ ID NO: 276	18,16,12	0.0	SEQ ID NO: 292	GGTCTAGATTGAGCAGGTGATGG	0.0
SEQ ID NO: 277	18,17,12	0.0	SEQ ID NO: 293	GGTTCAGATTGAGCAGGTGATGG	0.0
SEQ ID NO: 278	18,17,16,12	0.0	SEQ ID NO: 294	GGTTTAGATTGAGCAGGTGATGG	0.1

TREE:Site-1

Untransfected APOE

b

	Editing window	% of Reads		Editing window	% of Reads
SEQ ID NO: 295	WT	93.4	SEQ ID NO: 303	GAAGCGCTGGCAGGTGATCCAGG	84.2
SEQ ID NO: 296	16	2.5	SEQ ID NO: 304	GAAGTGCTTGGCAGGTGATCCAGG	5.2
SEQ ID NO: 297	14	0.1	SEQ ID NO: 305	GAAGCGTCTGGCAGGTGATCCAGG	1.1
SEQ ID NO: 298	13	0.1	SEQ ID NO: 306	GAAGCGCTTGGCAGGTGATCCAGG	1.0
SEQ ID NO: 299	16,14	1.1	SEQ ID NO: 307	GAAGTGCTTGGCAGGTGATCCAGG	1.2
SEQ ID NO: 300	16,13	0.4	SEQ ID NO: 308	GAAGTGCTTGGCAGGTGATCCAGG	1.0
SEQ ID NO: 301	14,13	0.5	SEQ ID NO: 309	GAAGCGTCTGGCAGGTGATCCAGG	7.0
SEQ ID NO: 302	16,14,13	1.9	SEQ ID NO: 310	GAAGTGTTTGGCAGGTGATCCAGG	19.2

TREE:APOE

FIG. 29




FIG. 30



FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37





FIG. 39



U6 Promoter
Sg2x Stop
sgRNA hairpin
U6 Promoter
Sg(NT) guide
sgRNA hairpin
pUC19 Backbone

GGGTAAATCATGCTCATAGGCTTCTCTGTGTGTAATTTGTTATCCGGTCAACAATTCACACAAC
 ATATCCAGGCGCGGAAGCATAAAGTGTAAAGCTGGGTGGCTATGAGTAGAGCTAACTACCAT
 AATTGGTGGGCTCACTGCCCGCTTCAGTCGGGAACCTGTGGTGGAGCTGCACTGATTAAT
 GAATCGGCGCAACCGCGGCGAGAGCGGTTTGGCTATTGGGGGCTCTTCGCTCTCTCGC
 TCACTAGCTCGCTGGCTCGGTGTGGGTGCGCGAGCGGTATCAGTCACTCAAAGGCT
 GGTAAATACGGTTATCCACAGATCAAGGGAATACCGAGGAAGACATGTGACGAAGAAGCG
 CAGCAAAAAGCGGAGGAACGGTAAAGAGGCGGCTTCTGGGCTTTTCCATAGGCTCCGCGC
 CCGCTGACGAGCATCAAAATCGAGCTCAAGCTAGAGGTGGCAACCGACGACAGCT
 ATAAAGATCAACAGGGGTTCGCCGTGGAGCTCCCTCGGCGCTCTCTGTCCGACCCGT
 CCGCTTACCAGTACCTGTGCGCTTCTCCCTTGGGAAGCTGGGCTTCTCATAGCTC
 AAGCGTGTAGGTATCTCAAGTTGGGTAGTGTCTGGCTCAAGCTGGGCTGTGGAGCGAA
 CCGCGCTTACGCGGACCGCTCGCGTATCCGGTAACTATCTGTGTAGTCAACAACCGG
 TAAGACACGCACTATCGCATCGCAAGCGCACTGGTAAACAGGATTAGCAAGAGCGAGTAT
 TTGAGCGGCTGCTACAGATGTTCTGAAGTGTGGCTAACTACGGCTACACTAGAAGACATCA
 GTATTGTTGTTATCGCTCTGCTGTAAGCGAGTATACCTTGGCAAAAAGAGTTGGTAGCTCTTG
 ATTCGCGAACAACCAACCGCGGTGTGACGGTGGTTTTTTTGTTCAGCAAGCAAGATTACGG
 CGCAAAAAGAAGATCTCAAGAAGATCCTTGATCTTTCTAAGGGGCTTGACCTCAGTAGTC
 CAGCAAAACTCAAGTTAGAGGATTTGGCTATGATGATTTCAAAGAGCTTCACTCTAGTATG
 GTTTTAAATTAATAATGAAGTTTAAATCAATCAAGATATATGAGTAAGCTTGTCTGACA
 GTTTACCAATGGTAACTAGGAGCGACCTATCCAGCATCTGTCTATTCTCATCCATAG
 TTGGCTGACTCCCGCTGTGTAGTAACTACGATAGCGGAGGCTTACCATCTGGCCCCAG
 TTGTGCAATGATACCGCGAGCCAGCGCTCCACCGCTCCAGATTATCAGGAATCAACAACG
 TTGCGAGGAAGGCGCGAGCGGAAGTGGCTGTCACATTTATCGCGCTCAATCCAGTCTA
 TTATAATTGTCGCGGAAGCTAGATGAAGTGGTGGCAGTAAAGTTTGGCGCAACGTTGTT
 GGCATCTGTACAGGATCGTGGTGTGACGCTGTGGTGTGGTATCGGCTTCACTCAGCTCCG
 GTTCTCAACATCAAGCGGCTATCATATCCCGATCCCGATTTGTGCAAAAAGCGTTAGTCTG
 TTGGCTGCTCGATGGTTGACAGAATGAAGTTGGCGGAGTGTATCACTCATGGTTATGGC
 AGCATGCAATATCTCTTACTGTCATGCCATCGGTAAAGTCTTCTGTGCTAGTGGTGATGTA
 CTCACCAAGTCTATGAGATAGTGTATCGCGGACCGAGTGGCTCTTGGCGCGGTCA
 ATATTCAGGGAATACCGCGCATAGCAGCTATAAAGTGGCTCATCATGAGAAACAGGCTTC
 TTGTGGGTGAACAACCTCGAGGATCTACCGGTGTGGAGTCCAGTGTGAATGAACCCACTC
 GTGCACCAACTGATCTCAGCATCTTTACTTCAACAGCGGTTTCTGGGTGAGCAAAAATCA
 GGAAGCAAAAATTCGCGCAAAAAGGATTAAGCGGACGACGGAATTTGTAATCTCATAC
 TCTCTGCTTTTCAATATTGAAGCATTTATCAGGGTATTGTCTATGAGCGGATACATATT
 TGAATGTATTAGAAAAATAACAATAGGGGTTCCGCGCACATTTCCCGGAAAGTGGCAC
 TCGACGTCTAGAAAACCATTTATTATCATGACATTAACTATAAAAATAGCGGTATCAACGAGGCG
 CTTCTGCTG

FIG. 40




FIG. 40 (CONTINUED)

FIG. 41 (CONTINUED)

FIG. 41 (CONTINUED)

FIG. 42 (CONTINUED)

FIG. 43 (CONTINUED)

FIG. 44 (CONTINUED)

FIG. 45 (Continued)

BFP coding sequence

Protospacer

Histidine 66 "CAC" CodonPAM "CGG"

SEQ ID NO: 258

>BFP reporter coding sequence

ATGGTGAAGCAAGGCGAGGAGCTGTTCACCGGGTGTGCCATCCTGGTCGAGCTGGACGGCAGC
TAAACGGCCACAAGTTCAGCGTGTCCGGGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCCCT
GAAGTTCATCTGCACCAACCGGCAAGCTGCCCCGTGCCCTGGCCCCACCCCTGACCCACCTGACCCACG
GCGTGCAGTGCTTCGGCCGTACCCCGACCAATGAGCAGCAGACTTCTTCAAGTCCGCCATGCC
GAAGGCTACGTCCAGGAGCGCACCATCTTCTTCAAGGACGACGGCACTACAAGACCCGCCGAGG
TGAAGTTCGAGGGCGACACCCCTGGTGAAACCGCATCGAGCTGAAGGCAATCGACTTCAAGGAGACGG
CAACATCCTGGGGCACAAGCTGGAGTACAATAACAGCCACAACGTCTATATCATGGCCGACAAG
CAGAAGAACGGCATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCG
CCGACCACTACCAAGACAACCCCATCGGGCAGCGCCCGTGTCTGCTGCCCGACAAACCACTACCTG
AGCACCCAGTCCGCCCTGAGCAAAAGACCCCAACGAGAAAGCGCGATCACATGGTCTGCTGGAGTTCTGT
GACCGCGCCGGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

FIG. 45 (Continued)

SEQ ID NO: 259

>XMAS 1xStop Reporter Coding Sequence
 ATGGTGAGCAAGGGCGAGGAGGATAACATGGGCATCATCAAGGAGTTCAATGGCTTCAAGGTGCACA
 TGGAGGGCTCCGTGAACGGCCACGAGTTCGAGATCGAGGGCGAGGGCGAGGGCCGCCCTACGAGGG
 CACCCAGACCGCCAAAGCTGAAGGTGACCAAGGTGGCCCCCTGCCCCCTTGGCTGGGACATCTCTGTCCC
 CTCAGTTCAATGTACGGCTCCAAAGGCTACGTGAAGCACCCCGCCGACATCCCCGACTACTTGAAGCTG
 TCCTTCCCGAGGGCTTCAAGTGGGAGCCGCTGATGAATCTTGAAGACCGCGGCGCTGGTGACCCGTGAC
 CCAGGACTCTCTCCGAGGACGGCGAGTTCTATACAAAGTGAAGCTGGCGGCAACCAACTTCCCT
 CCGACGGCCCCGTAAATGCAGAGAAGAACCATGGGCTGGGAGGCTTCTCCGAGCGGATGTACCCCGA
 GGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCTGAAGGACGGCGGCACTACGACGCT
 GAGGTCAAGACCACCTACAAGGCCAAGAAGCCGTGCAAGTGGCCGGGCTTACAAGTCAACATCA
 AGTTGGACATCACCTCCACAAACGAGGACTACACCATCGTGGACACAGTACGAACGCCGCGAGGGCCG
 CCACCTCACCGGGCGCATGGACGAGCTGTACAGCCCCGGGAGGGCAGAGGAGTCTTAAATGTC
 GGTGACGTGAGGAGAGATCCCGGCCCTACTAGTTGATGGGGTGGTTCAAGAGGGGCATGCCGTGAGCA
 AAGGCGAGGAGCTGTTCACCGGGGTGGTGGCCATCTCTGGTGGAGCTGGACGGCGACGTAAACGGCCA
 CAAATTACGCTGTCCGGGAGGGCGAGGGGATGCCACCTACGGCAAGCTGACCTTGAAGTTCACT
 GCACCAACCGCAAGCTGCCCGTCCCTGGCCCCACCTCTGTGACCACTTGAACCTTACGGGTGCAATGC
 TTCAGCGCTACCCCGACCATGAAGCAGCAGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGT
 CCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCTGGCGCGAGGTGAAGTTCCAG
 GCGCACACCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGG
 GGCACAAGCTGGAGTACAACACTACAAGCCACAAGTCTATATCATGGCCGACAAGCAGAGAACGG
 CATCAAGGTGAACCTTCAAGATCCGCCACAACATCGAGGACGGCGAGGTGACGCTGGCGACCACTAC
 AGCAGAACACCCCATCGGGACCGCCCGTGTCTGCCCCACAACCACTACTCTAAGCACCCCAATCC
 GCCCTGAGCAAAAGACCCCAACGAGAAAGCCGATCATCATGCTCTCTGGAGTTCTGTGACCGCGCCG
 GGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

>XMAS 1xStop Reporter Protospacer and PAM sequence
 GTTGAATGGGGTGGTTCAAGGAGGG

SEQ ID NO: 315

FIG. 45 (Continued)


SEQ ID NO: 260

>XMAS 1xStop Reporter Coding Sequence
 ATGTTGAGCAAGGGCGAGGAGGATAACATGGCCATCATCAAGGAGTTATGCGCTTCAAGGTGCACA
 TGGAGGCTCCGTGAACGGCCACGAGTTGAGATCGAGGCGAGGGCGAGGGCCGCCCTACGAGGG
 CACCAGACCGCCAAAGCTGAAGGTGACCAAGGTGGCCCTCCCTTCGCTTGGACATCCTGTGTCC
 CTCAGTTTCATGTACGGCTCCAAAGGCTACGTGAAGCACCCCGCGACATCCCGACTACTTGAAGCTG
 TCCTTCCCGAGGGCTTCAAGTGGAGCGCGTGATGAACCTTGGAGGACGGCGGTGGTGACCGTGAC
 CCAGGACTCCTCCCTGACGAGCGGAGTTTCTACAAAGGTGAAGCTGGCGGCAACCACTTCCCT
 CCGACGGCCCGGTAAATGCAGAAAGACCAATGGGTGGAGGCTCTCCGAGCGGATGTACCCCGA
 GGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCTGAGGACGGCGGCCACTACGACGCT
 GAGGTCAAGACCACTACAAGGCCAAGAAGCCCGTGCAGCTGCCCGGCCCTACAACGTCAACATCA
 AGTTGGACATCACTCCCAACAAGAGGACTACACCATCGTGGAAACAGTACGAACGCCCGAGGGCCG
 CCACTCCACCGCGGCAATGACGAGCTGTACAAGCCCGGAGAGGGGAGAGGAAATGTTCTAAACATGC
 GGTGACGTGGAGGAGAAATCCCGCCCTACTAGTTGATGAGGGGTTCAGGAGGGGCATGCGTGAGCA
 AGGGCGAGGAGCTGTTCAACCGGGTGTGCCCATCTTGGTCGAGCTGGACGGCGACGTAAACGGCCA
 CAAGTTACGCGTGTCCGCGAGGGCGAGGGCGATGCCACCTACGGCAAGCTGACCTGAAGTTCACT
 GCACCAACGGCAAGCTGCCCGTGCCTGGCCACCCCTCGTGACCACCTGACCTACGGCGTGCAGTGC
 TTCAGCCGCTACCCGACCACATGAAGCAGCACGACTTCTTCAAGTCCGCCATGCCCGAAGGCTACGT
 CCAGGAGCGCACCATCTTCTTCAAGGACGACGGCAACTACAAGACCCCGCGCGAGGTGAAGTTCGAG
 GCGGACACCTGGTGAACCGCATCGAGCTGAAGGGCATCGACTTCAAGGAGGACGGCAACATCCTGG
 GGCACAAGCTGGAGTACAACTACACAGCCACAACGTCTATATCATGGCCGACAAGCAGAGAACGG
 CATCAAGGTGAACTTCAAGATCCGCCACAACATCGAGGACGGCAGCGTGCAGCTCGCCGACCACTACC
 AGCAGAACACCCCATCGCGACGGCCCGTGTCTGCTGCCCGACAACCACTACTGAGCACCAGTCC
 GCCCTGAGCAAGACCCCAACGAGAAAGCGGATCACATGGTCTCTGTGAGTTCTGTGACCCGCCCG
 GGATCACTCTCGGCATGGACGAGCTGTACAAGTAA

>XMAS 2xStop Reporter Protospacer and PAM sequence
 GTTGATGAGGTGTTTCAGGAGGG

SEQ ID NO: 261

FIG. 46



XMAS 1x stop Translation prior to editing: SEQ ID NO: 316
MVSKEEDNMAIIEFMRFKVHMEGVSNGHEFEIEGEGEGRPYEGTAKLVTKGGPLFAWDILSPQFMYGSKAYVKHPADIPDYLKLSFPEG
FKWERVMNFEDGGVTVTQDSSLQDGEFIYKVKLRGTNFPSDGPVMQKKTMGWEASSERMYPEDGALKGEIKQRLKLDGGHYDAEVKTTYKA
KKPVQLPGAYNVNIKDITSHNEDYTIVEQYERAEGRHSTGGMDLEYKPREGRGSLTTCGDVEENPGPTS-
WGGSGGACVSKGEELFTGVWPILVELDGVNNGHKFSVSGEGEGDATYGKLTLCFICTTGKLPVPWPTLVTTLTYGVCFSRYPDHMKQHDFKSA
MPEGVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYVNSHNVIYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQ
QNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLVLEFVTAAGITLGMDELYK-

XMAS1x stop Translation after editing: SEQ ID NO: 317
MVSKEEDNMAIIEFMRFKVHMEGVSNGHEFEIEGEGEGRPYEGTAKLVTKGGPLFAWDILSPQFMYGSKAYVKHPADIPDYLKLSFPEG
FKWERVMNFEDGGVTVTQDSSLQDGEFIYKVKLRGTNFPSDGPVMQKKTMGWEASSERMYPEDGALKGEIKQRLKLDGGHYDAEVKTTYKA
KKPVQLPGAYNVNIKDITSHNEDYTIVEQYERAEGRHSTGGMDLEYKPREGRGSLTTCGDVEENPGPTS-
WGGSGGACVSKGEELFTGVWPILVELDGVNNGHKFSVSGEGEGDATYGKLTLCFICTTGKLPVPWPTLVTTLTYGVCFSRYPDHMKQHDFKSA
MPEGVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYVNSHNVIYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQ
QNTPIGDGPVLLPDNHYLSTQSALS KDPNEKRDHMLVLEFVTAAGITLGMDELYK-

Figure 46 (continued)

XMAS 2x stop Translation no editing ; SEQ ID NO: 318

MVSKGEEDNMAIIEFMRFKVHMEGVSNGHEFEIEGEGEGRPYEGTQTAKLVTKGGPLPFAWDILSPQFMYGSKAYVVKHPADIPDYLKL
SFPEGFKWERVMNFEDGGVTVTQDSSLQDGEFIYKVKLRGTNFPSDGPMQKKTMGWEASSERMYPEDGALKGEIKQRLKLDGGHY
DAEVKTTYKAKKPVQLPGAYNVNIKLDITSHNEDYTIQEYERAEGRHSTGGMDELYKPREGRGSLTCGDVEENPGPTS--
GGSGGACVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTTLYGVQCFSRYPDHIKQHDFF
KSAMPEGVYQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNNSHNVYIMADKQKNGIKVNFKIRHNIEDGSV
QLADHYQQNTPIGDPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK-

XMAS 2x stop After editing SEQ ID NO: 319

MVSKGEEDNMAIIEFMRFKVHMEGVSNGHEFEIEGEGEGRPYEGTQTAKLVTKGGPLPFAWDILSPQFMYGSKAYVVKHPADIPDYLKL
SFPEGFKWERVMNFEDGGVTVTQDSSLQDGEFIYKVKLRGTNFPSDGPMQKKTMGWEASSERMYPEDGALKGEIKQRLKLDGGHY
DAEVKTTYKAKKPVQLPGAYNVNIKLDITSHNEDYTIQEYERAEGRHSTGGMDELYKPREGRGSLTCGDVEENPGPTS--
VSKGEELFTGVVPILVELDGDVNGHKFSVSGEGEGDATYGKLTCLKFICTTGKLPVPWPTLVTTLYGVQCFSRYPDHIKQHDFFKSAMPEG
YVQERTIFFKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNNSHNVYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQ
QNTPIGDPVLLPDNHYLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK-

1

TRANSIENT REPORTERS AND METHODS FOR BASE EDITING ENRICHMENT

CROSS-REFERENCE TO RELATED PATENT APPLICATIONS

The present application claims the benefit of priority under 35 U.S.C. § 119(e) to U.S. Provisional Application No. 63/038,220, filed on Jun. 12, 2020, the content of which is incorporated herein by reference in its entirety.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH

This invention was made with government support under R01 GM106081, R01 GM121698, R01 GM131405 and R21 AG056706 awarded by the National Institutes of Health. The government has certain rights in the invention.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 8, 2024, is named 112624_01257_M20-204L_SL.txt and is 171,515 bytes in size.

BACKGROUND

The rapid advancement of CRISPR/Cas-based technologies has allowed for the modification (i.e., deletion, mutation and insertion) of human cells at precise genomic locations. For applications in which precise editing of a single nucleotide is desired, the CRISPR/Cas machinery can be used to introduce site-specific double-stranded breaks (DSB) followed by homology-directed repair (HDR) using an exogenous DNA template. However, HDR is inefficient in mammalian cells, especially in recalcitrant cells such as human pluripotent stem cells (hPSCs), and repair of DSB is predominantly achieved through non-homologous end joining (NHEJ). In addition, NHEJ results in insertion or deletion of nucleotides (indels), resulting in undesired disruption (e.g. frameshift mutations, premature stop codons, deletion) of the targeted genes.

As an alternative to standard gene editing approaches that require a DSB, several groups have reported the development of deaminase base editors that do not rely on HDR to introduce single nucleotide genomic changes. Broadly speaking, these base editors consist of a fusion of three components—a DOA nickase Cas endonuclease, cytidine deaminase (APOBEC1), and a DNA uracil glycosylase inhibitor (UGI). This complex is capable of converting cytosine to thymine (or adenine to guanine on the complementary strand) without the need for a DSB and homology repair template. Overall, genome modification through the use of base editors has been shown to result in formation of fewer indels when compared to HDR-based methods.

Despite the advantages that deaminase base editors offer, identification and isolation of cell populations that have been successfully edited remains challenging. Specifically, there is no readily detectable phenotype to distinguish edited from unedited cells. In turn, isolation of edited cell populations requires single cell isolation followed by downstream sequencing verification. Some progress has been made to help enrich for edited cells, such as co-transfecting plasmids with a fluorescent reporter and using flow cytometry to

2

isolate reporter-positive cells. Similarly, base editors fused to fluorescent proteins have been used to enrich for edited cell populations. However, these techniques are only reporters of transfection (RoT) and do not report on base editing activity within a cell population. Accordingly, there remains a need in the art for materials and efficient methods for selecting and enriching for base-edited human cells. There also remains a need in the art for efficient methods for producing isogenic populations of base edited human cells, particularly human pluripotent stem cell populations having targeted genetic modifications.

BRIEF SUMMARY OF THE DISCLOSURE

In a first aspect, a polynucleotide encoding one or more reporter polypeptides, the polynucleotide including a PAM site adjacent to a base that when the base is edited a change in a function or characteristic of the one or more reporter polypeptides occurs. The polynucleotide may encode at least one reporter polypeptide with at least 90% sequence identity to SEQ ID NO: 2, wherein the polynucleotide encodes histidine at amino acid at position number 66 relative to SEQ ID NO: 1, and encodes glycine at amino acid position number 72 relative to SEQ ID NO: 1. Alternatively, the polynucleotide may encode a reporter polypeptide with at least 90% sequence identity to one of SEQ ID NO: 316 or 318. In one alternative, the polynucleotide comprises a polynucleotide selected from the group consisting of SEQ ID NO: 258, 259 and 260.

In a second aspect, provided herein is a kit. The kit can comprise or consist essentially of a first nucleic acid sequence encoding one or more reporter proteins, wherein the first nucleic acid includes a PAM site adjacent to a base that when edited causes a change in a function or characteristic of the one or more reporter proteins; a second nucleic acid sequence encoding a first sgRNA adjacent to a protospacer adjacent motif (PAM), wherein the first sgRNA comprises a protospacer sequence and is complementary to a portion of the nucleic acid sequence encoding one or more reporter proteins; a third nucleic acid sequence encoding a second sgRNA adjacent to a protospacer adjacent motif (PAM), wherein the sgRNA comprises a protospacer sequence and is complementary to a portion of a gene of interest to be base edited or comprises a cloning site to allow insertion of a complementary portion of a gene of interest to be base edited; and a fourth nucleic acid sequence encoding a base editor. The base editor can be selected from a cytidine deaminase base editor, an adenine base editor, Cas9-mediated adenosine base editor, and a prime editor. One or more of the first, second, third, or fourth nucleic acids can be provided in one or more vectors. The vector can be an episomal vector. The reporter protein can be a fluorescent protein or a variant thereof, luciferase or a variant thereof, β -galactosidase (lacZ), chloramphenyl acetyltransferase (CAT), β -glucuronidase (GUS), secretory alkaline phosphatase (SEAP), a survival selection protein, or a reporter protein that directly or indirectly produces or catalyzes a colorimetric reaction. The fluorescent protein can be a green fluorescent protein (GFP), a blue fluorescent protein (BFP), red fluorescent protein (RFP), luciferase, or mCherry, or a variant thereof. The fluorescent protein can be a BFP variant comprising a histidine at amino acid position 66 (numbered relative to SEQ ID NO:1). Alternatively, the reporter protein may be a fusion protein of two fluorescent proteins linked via a linker including at least one stop codon and a PAM site.

The fourth nucleic acid sequence encoding a base editor can be a vector comprising a base editor operably linked to a heterologous promoter.

In another aspect, provided herein is a method for selecting a base edited cell. The method can comprise or consist essentially of introducing into a cell a first nucleic acid sequence encoding one or more reporter proteins, a second nucleic acid sequence encoding a first sgRNA adjacent to a protospacer adjacent motif (PAM), wherein the first sgRNA comprises a protospacer sequence and is complementary to a portion of the nucleic acid sequence encoding one or more reporter proteins; a third nucleic acid encoding a second sgRNA adjacent to a protospacer adjacent motif (PAM), wherein the second sgRNA comprises a protospacer adjacent sequence and is complementary to a portion of a gene of interest to be base edited; and a fourth nucleic acid sequence encoding a base editor, wherein the first nucleic acid includes a PAM site adjacent to a base that when edited causes a change in a function or characteristic of the one or more reporter proteins and wherein the change in function or characteristic results in a detectable signal; culturing the cell for about 48 hours to about 72 hours under conditions sufficient for expression of proteins encoded by the first, second, third and fourth nucleic acid sequences; sorting cells based on the presence or absence of a detectable signal, wherein a change in the detectable signal indicates that the base editor caused a base-to-base conversion or other genetic modification in the first nucleic acid sequence; and selecting cells exhibiting the changed detectable signal from the sorted cells, thereby selecting base edited cells. The base editor can be selected from a cytidine deaminase base editor, an adenine base editor, Cas9-mediated adenosine base editor, and a prime editor. One or more of the first, second, and third nucleic acids can be provided in a vector. The vector can be an episomal vector. The reporter protein can be a fluorescent protein or a variant thereof, luciferase or a variant thereof, β -galactosidase (lacZ), chloramphenyl acetyltransferase (CAT), β -glucuronidase (GUS), secretory alkaline phosphatase (SEAP), a survival selection protein, or a reporter protein that directly or indirectly produces or catalyzes a colorimetric reaction. The fluorescent protein can be a green fluorescent protein (GFP), a blue fluorescent protein (BFP), red fluorescent protein (RFP), luciferase, mCherry, or a variant or combination thereof. The fluorescent protein can be a BFP variant comprising a histidine at amino acid position 66 (numbered relative to SEQ ID NO:1). The cell can be a human cell. The human cell can be a human pluripotent stem cell. The human pluripotent stem cell can be a human induced pluripotent stem cell obtained from a somatic cell of a human subject having a disease-associated single nucleotide polymorphism. Selecting can be performed using flow cytometry. Sorting can be performed using a fluorescence activated cell sorter (FACS).

The foregoing and other advantages of the invention will appear from the following description. In the description, reference is made to the accompanying drawings which form a part hereof, and in which there is shown by way of illustration a preferred embodiment of the invention. Such embodiment does not necessarily represent the full scope of the invention, however, and reference is made therefore to the claims and herein for interpreting the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1F demonstrate conversion of BFP to GFP enables detection of base-editing activity in cells. (A) A

mutant BFP was designed to convert to GFP upon a C-to-T nucleotide conversion. The protospacer sequence (underlined black) for the sgRNA, sg (BG), targeting the 'CAC' codon (underlined blue) resulting in a C-to-T conversion to 'TAC' (underlined green) and the corresponding amino acid change of histidine (blue) to tyrosine (green) at the 66th amino acid position in BFP. A PAM (underlined red) was placed in the position to orient the base editing window (underlined orange) around the C nucleotide (red) to facilitate BFP_{H66} to GFP_{Y66} conversion. All alternative C-to-T conversions in the editing window resulted in silent mutations of the coding sequence. (B) The BFP mutant was cloned into a vector, pEF-BFP, with a human EF1 α promoter driving expression. Targeting pEF-BFP with a cytidine deaminase base editor results in a C-to-T conversion causing a shift in the fluorescent emission spectra from BFP to GFP. (C) Representative fluorescent microscopy images of HEK293 cells transfected with pEF-BFP, pCMV-BE4-Gam and sg(NT) (top row) or sg(BG) (bottom row). (D) Representative flow cytometry plots of HEK293 cells transfected with pEF-BFP, pCMV-BE4-Gam and sg(NT) (top) or sg(BG) (bottom). Y-axis is a non-fluorescent control channel. (E) Schematic for isolation and detection of editing of episomal DNA after transfection. (F) Representative Sanger sequencing chromatogram of amplicons of episomal DNA isolated from HEK293 cells transfected with pEF-BFP, pCMV-BE4-Gam and sg(BG). The presence of T-nucleotide (red trace) at the target nucleotide (red asterisk) demonstrates the C-to-T base conversion responsible for the amino acid change of histidine to tyrosine at the 66th amino acid position and subsequent shift of the BFP emission spectra of the resultant protein to a GFP variant.

FIGS. 2A-2F demonstrate BFP-to-GFP conversion reports on base-editing at a chromosomal locus. (A) A pEF-BFP-PuroR vector was integrated into the C1ORF228 locus using homology-independent targeted integration to generate the HEK293-BFP cell line. (B) Schematic for plasmid or RNP base editing optimization using the HEK293-BFP line. (C) Representative fluorescent microscopy images of HEK293-BFP cells transfected with 600 ng pCMV-BE4-Gam and 200 ng sg(NT) (top row) or sg(BG) (bottom row). Scale bar=200 μ m. (D) Editing efficiencies (GFP-positive cells) of HEK293-BFP cells transfected with various amounts of pCMV-BE4-Gam and ratios with the sg(BG) vector. n=3, * P <0.05. (E) Representative fluorescent microscopy images of HEK293-BFP cells transfected with BE3-sg(BG) or -sg(NT) RNP complexes. (F) Schematic for RNP complex generation and transfection. BE3 was overexpressed, purified, complexed and validated in vitro, and transfected. Editing efficiencies (GFP-positive cells) of HEK293-BFP cells transfected with RNP complexes using various delivery reagents. n=3, * P <0.05, ** P <0.01.

FIGS. 3A-3E demonstrate enrichment of base-edited cell populations using TREE. (A) Plasmid map of pDT-sgRNA vector that contains sg(BG) and sg(TS). Expression for both sgRNA cassettes is driven by separate U6 promoters (orange arrows). The BbsI restriction sites allow for direct restriction enzyme-based cloning of new target sites. (B) Schematic for enrichment of edited cells using TREE. HEK293 cells are co-transfected with pEF-BFP, pCMV-BE4-Gam and pDT-sgRNA vectors. After 48 h post-transfection, flow cytometry is used to sort cell populations into GFP-positive and -negative fractions. (C) Schematic for enrichment of edited cells using reporter of transfection (RoT). HEK293 cells are co-transfected with pEF-GFP, pCMV-BE4-Gam and sg(TS) vectors. After 48 h post-transfection, flow cytometry is used

to sort cell populations into GFP-positive and -negative fractions. (D) Representative flow cytometry plots of (i) untransfected HEK293 cells and (ii) HEK293 cells transfected with pEF-BFP only as well as HEK293 cells in which TREE was applied targeting (iii) Site-1, (iv) Site-2 and (v) Site-3. (E) Quantification of base editing efficiency at Site-1, Site-2 and Site-3 in GFP-positive, GFP-negative and unsorted cell populations isolated using TREE- or RoT-based enrichment strategies. $n=3$; $*=P<0.05$, $**=P<0.01$.

FIGS. 4A-4C demonstrate TREE enables efficient multiplex base editing. (A) Plasmid map of pMT-sgRNA vector that contains sg(BG) in addition to sgRNA for multiple target sites. Expression for all sgRNA cassettes is driven by separate U6 promoters (orange arrows). The HindIII restriction site allows for additional sgRNAs for target sites to be cloned in through restriction enzyme-based cloning. (B) Quantification of multiplex base editing efficiency at Site-1, Site-2 and Site-3 in GFP-positive, GFP-negative and unsorted cell populations using TREE- or RoT-based enrichment strategies. $n=3$; $*=P<0.05$, $**=P<0.01$. (C) Clonal analysis of editing at multiple genomic loci using TREE. 40 GFP-positive clones were isolated via single-cell sorting. Editing was detected via PCR and Sanger sequencing. Blank icon indicates no editing observed, half-red icon indicates heterozygous C and T at the target site, and solid red icon indicates homozygous T edits at the genomic site.

FIGS. 5A-5D demonstrate highly efficient editing in human pluripotent stem cells (hPSCs) using TREE (A) Quantification of base editing efficiency (percentage GFP-positive cells) when hPSCs were co-transfected with pEF-BFP, sg(BG) and various base editing vectors. $n=3$; $*=P<0.05$, $**=P<0.01$. (B) Schematic for enrichment of edited hPSC using TREE. HPSCs were co-transfected with pEF-BFP, pEF-BE4-Gam/pEF-AncBE4 and pDT-sgRNA vectors. 48 h post-transfection, flow cytometry was used to sort cell populations into GFP-positive and -negative fractions. (C) Representative flow cytometry plots of (i) untransfected hPSCs cells and (ii) hPSCs transfected with pEF-BFP only as well as hPSCs cells in which TREE was applied targeting Site-1 utilizing (iii) pEF-BE4-Gam or (iv) pEF-AncBE4. (D) Quantification of base editing efficiency at Site-1 in GFP-positive, GFP-negative and unsorted cell populations isolated using TREE- or RoT-based enrichment strategies in which pEF-BE4-Gam or pEF-AncBE4 was employed. $n=3$; $*=P<0.05$, $**=P<0.01$.

FIGS. 6A-6I demonstrate efficient generation of isogenic hPSC lines using BIG-TREE. (A) Schematic for generation of clonal isogenic hPSC lines using BIG-TREE. HPSCs are co-transfected with pEF-BFP, pEF-AncBE4max, and pDT-sgRNA plasmid vectors. Forty-eight hours post transfection, FACS is used to isolate single GFP-positive cells into 96-well plates. Cells are subsequently expanded, and target clones are identified by Sanger sequencing of the target loci. (B) Schematic of vectors used for BIG-TREE- and RoT-based generation of clonal hPSC lines in which the APOE (158R) locus has been targeted. (C) Schematic of the APOE (158R) target locus in exon 4 of the APOE gene. Successful base editing of the APOE(158R) locus would result in a C-to-T conversion causing a change in the amino acid position at 158 from an arginine (APOE3) to a cysteine (APOE2). Representative Sanger sequences of the APOE (158R) locus of unedited parental hPSC lines as well as clonal hPSC lines that have been edited at the APOE(158R) are shown. Each line shown is representative of clones obtained from three independent parental hPSC populations (hPSC lines 1-3) with different genetic backgrounds. (D) Distribution of genotypes in clonal hPSCs derived from

hPSC line 1 that was targeted at the APOE(158R) locus using BIG-TREE- or RoT-based methods. (E) Distribution of genotypes in clonal hPSCs derived from hPSC lines 2 and 3 that were generated via BIG-TREE-based targeting at the APOE(158R) locus. (F) Karyotype analysis of representative clones edited at the APOE(158R) locus. (G) Phase contrast images of representative clones edited at the APOE (158R) locus. (H) Immunofluorescence staining of representative clones edited at the APOE(158R) locus for pluripotency markers NANOG, OCT4, and SOX2. (I) Alpha fetoprotein (AFP), smooth muscle actin (SMA), and beta-III tubulin (TUJ1) immunofluorescence staining of representative clones edited at the APOE(158R) locus that had been subject to tri-lineage differentiation.

FIGS. 7A-7D demonstrate BIG-TREE-Based Gene Knockout of APOE in hPSCs. (A) Schematic of the APOE (39Q) locus in exon 3 of the APOE gene. Successful base editing of the APOE(39Q) locus would result in a C-to-T conversion causing a change in the amino acid at position 39 from a glutamine to a premature stop codon. (B) Representative Sanger sequencing of the APOE(39Q) locus in unedited wild-type cells (Q39/Q39; left panel) as well as hPSC clones in which a heterozygous (Q39/X39; middle panel) or homozygous (X39/X39; right panel) stop codon has been introduced. (C) Distribution of genotypes in clonal hPSCs that were generated via BIG-TREE targeting the APOE(39Q) locus. (D) Measurement of ApoE secretion in the condition medium of wild-type (Q39/Q39) and homozygous edited (X39/X39) hPSCs ($n=3$ independent experiments, $p<0.05$, two-tailed Student's t test).

FIGS. 8A-8C demonstrate simultaneous base editing of multiple loci in hPSCs using BIG-TREE. (A) Schematic of plasmid vectors used for BIG-TREE-based generation of clonal hPSC lines in which multiple loci have been simultaneously targeted. The pMT-sgRNA vector contains sg(BG) in addition to sgRNA for multiple target sites (S1, genomic site 1; S2, genomic site 2; S3, genomic site 3). (B) Representative Sanger sequencing chromatographs of the site 1, site 2, and site 3 loci in clonal hPSCs that have been generated via BIG-TREE multiplexed base editing. (C) Distribution of genotypes in clonal hPSCs that were generated via BIG-TREE multiplexed base editing.

FIG. 9 presents pDT-sgRNA sequence.

FIG. 10 is a schematic illustration of vector pDT-sgRNA. FIG. 11 is a schematic illustration of vector pEF-AncBE4max.

FIG. 12 is a schematic illustration of vector pEF-BE4Gam. Additional vector map illustrations are provided in FIGS. 39-44.

FIGS. 13A-13E illustrate a fluorescent reporter system for real-time measurement of adenosine base editing activity. (a) The XMAS-TREE reporter vector consists of a human EF1 α promoter driving expression of an mCherry cassette followed by a stop codon (TGA) then a GFP cassette. Targeting pEF-XMAS with an adenine base editor and sg(XMAS) will result in an A-to-G conversion, enabling expression of the downstream GFP reporter. (b) Two versions of pEF-XMAS-TREE plasmid were designed, one with a single stop codon (XMAS-1 $\times$ Stop) and another with two stop codons (XMAS-2 $\times$ Stop), preceding the coding sequence for GFP. The protospacer sequence (underlined black) for the sgRNA, sg(XMAS), targeting the TGA codon (underlined red) resulting in an A-to-G conversion to TGG and the corresponding amino acid change to tryptophan. The PAM sequence (underlined red) was placed to position the base editing window (underlined orange) around the target nucleotides. FIG. 13B discloses SEQ ID NOS 322, 200, 323,

201, 324, 202, and 324, respectively, in order of appearance. (c) Representative fluorescence microscopy images of HEK293 cells transfected with pEF-XMAS-1xStop (left panels) or pEF-XMAS-2xStop (right panels), pCMV-ABEmax, and sg(NT) (top panels) or sg(XMAS) (bottom panels). (d) Flow cytometry and (e) fluorescence microscopy analysis of HEK293 cells at various time points after transfection with pEF-XMAS-1xStop (left panels) or pEF-XMAS-2xStop (right panels), pCMVABEmax, and sg(XMAS).

FIGS. 14A-14B demonstrate identification and enrichment of base-edited cell populations using XMAS-TREE. (a) Schematic for identification and enrichment of adenosine base edited cells using XMAS-TREE. Cells are transfected with pEF-XMAS, pCMV-ABEmax, and pST-sgRNA vectors. Posttransfection, flow cytometry is used to sort cell populations into reporter-positive and -negative populations based upon mCherry and GFP expression levels. (b) Quantification of base editing efficiency at various genomic loci in unsorted (white bar), mCherry-negative/GFP-negative (grey bar), mCherry-positive/GFP-negative (red bar), and mCherry-positive/GFP-positive (orange bar) isolated cells using XMAS-TREE-based methods. Statistical comparisons were made between cells positive for the transfection reporter but not the editing reporter (i.e., mCherry-positive/GFP-negative) and cells positive for both (i.e., mCherry-positive/GFP-positive); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

FIGS. 15A-15C demonstrate that XMAS-TREE enables highly efficient multiplex adenosine base editing. (a) Cells were transfected with pEF-XMAS, pCMV-ABEmax, and a pMT-sgRNA that simultaneously targeted Site-1/Site-3/ Site-4 or Site-5/HBG1/HBG2. Base editing was quantified in unsorted as well as reporter-positive and -negative cell populations. Statistical comparisons were made between cells positive for the transfection reporter but not the editing reporter (i.e., mCherry-positive/GFP-negative) and cells positive for both (i.e., mCherry-positive/GFP-positive); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. (b) Schematic for employing XMAS-TREE for generation of clonal lines that have been simultaneously edited at multiple loci. HEK293 cells are co-transfected with pEF-XMAS, pCMV-ABEmax, and pMT-sgRNA. At 48 hours post-transfection, single mCherry-positive/GFP-positive cells are sorted into 96-well plates. After expansion, target clones are identified by Sanger sequencing at the target sites. (c) Analysis of clonal editing efficiency at multiple independent genomic sites using XMAS-TREE. A total of 30 clones were examined at each locus. White box indicates no editing observed at a specified locus, half-filled box indicates mono-allelic targeting at the genomic site, and full box indicates bi-allelic editing at the target locus.

FIGS. 16A-16D demonstrate that XMAS-TREE can be employed for the highly efficient base editing of human pluripotent stem cells (hPSCs). (a) Representative fluorescence microscopy images of hPSCs transfected with pEF-XMAS-1xStop (top panels) or pEF-XMAS-2xStop (bottom panels), pEFABEmax, and sg(XMAS). (b) Flow cytometry and (c) fluorescence microscopy analysis of hPSCs at various time points after transfection with pEF-XMAS-1xStop (top panels) or pEF-XMAS-2xStop (bottom panels), pEF-ABEmax, and sg(XMAS). (d) Quantification of base editing efficiency at various genomic loci in unsorted (white bar), mCherry-negative/GFP-negative (grey bar), mCherry-positive/GFP-negative (red bar), and mCherry-positive/GFP-positive (orange bar) isolated hPSCs using XMAS-TREE-based methods. Statistical comparisons were made between hPSCs positive for the transfection reporter but not the

editing reporter (i.e., mCherry-positive/GFP-negative) and cells positive for both (i.e., mCherry-positive/GFP-positive); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

FIGS. 17A-17C demonstrate highly efficient generation of clonal isogenic hPSC lines using XMAS-TREE. (a) Schematic for enrichment of adenosine base-edited cells using XMAS-TREE and reporter of transfection (RoT) based approaches. (b) Quantification of relative base editing in mCherry-positive cells isolated using RoT and mCherry-positive/GFP-positive cells isolated using XMASTREE. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$ (c) Analysis of clonal editing efficiency in hPSCs that were targeted at the Site-3 locus using XMAS-TREE- or RoT-based methods.

FIGS. 18A-18B demonstrate simultaneous adenosine base editing of multiple target sites in hPSCs using XMAS-TREE. (a) HPSCs were co-transfected with pEF-XMAS, pEF-ABEmax, and a pMT-sgRNA that simultaneously targeted Site-5/HBG1/HBG2. Flow cytometry was used to sort reporter-positive and -negative cell populations and base-editing was quantified at target loci. Statistical comparisons were made between cells positive for the transfection reporter but not the editing reporter (i.e., mCherry-positive/GFP-negative) and cells positive for both (i.e., mCherry-positive/GFP-positive); * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$. (b) Quantification of relative base editing in mCherry-positive and mCherry-positive/GFP-positive cells isolated using RoT and XMAS-TREE, respectively.

FIG. 19 shows Sanger sequencing chromatographs of Site-1, Site-2, and Site-3 of GFP-positive, GFP-negative, and unsorted cell populations isolated with TREE- and RoT-based approaches.

FIG. 20 demonstrates that TREE allows for base editing of refractory APOE(R158) locus in HEK293 cells. (a) HEK293 cells were transfected with pEF-GFP, pCMV-BE4-Gam, and sg(TS). Comparison of transfection efficiency (percentage of GFP-positive cells) and editing efficiency (percentage of C-to-T conversion at target nucleotide) in unsorted cell populations at Site-1, Site-2, Site-2, and APOE (R158) locus. (b) Representative Sanger sequencing chromatographs of APOE(R158) locus in GFP-positive, GFP-negative, and unsorted cell populations isolated with RoT-based methods. FIG. 20B discloses SEQ ID NOS 325 and 210, respectively, in order of appearance. (c) Representative flow cytometry plot of HEK293 cells in which TREE was applied targeting the APOE(R158) locus. (d) HEK293 cells were transfected with pEF-BFP, pCMV-BE4-Gam, and pDT-sgRNA. Comparison of transfection efficiency (percentage of BFP-positive cells) and editing efficiency (percentage of C-to-T conversion at target nucleotide) in unsorted cell populations at Site-1, Site-2, Site-3, and APOE (R158) locus. (e) Representative Sanger sequencing chromatographs of APOE(R158) locus in GFP-positive, GFP-negative, and unsorted cell populations isolated with TREE-based methods. FIG. 20E discloses SEQ ID NOS 325 and 211, respectively, in order of appearance.

FIG. 21 demonstrates that TREE fluorescent output in HEK293 cells is transient. (a) HEK293 cells were transfected with pEF-BFP, pCMV-BE4-Gam, and pDT-sgRNA and GFP-positive cells were isolated by flow cytometry. Replated GFP-positive cells were analyzed by fluorescent microscopy and flow cytometry at various time points post-sorting. (b) Representative fluorescent microscopy images of cells prior to cell sorting (D-1, Pre-sort) and various time points (D0, D7, D10) after sorting. (c) Representative flow cytometry plots of (i) untransfected HEK293

cells, (ii) pEF-GFP transfected HEK293 cells, and (iii) TREE-enriched GFP-positive HEK293 cells 10 days after sorting.

FIG. 22 shows an Analysis of multiplexed edited HEK293 cells using TREE- and RoT-based methods. (a) Representative flow cytometry plot of HEK293 cells in which multiplex TREE was applied simultaneously targeting Site-1, Site-2, and Site-3. (b) Representative Sanger sequencing chromatographs of the Site-1, Site-2, and Site-3 loci in GFP-positive, GFP-negative, and unsorted cell populations isolated with TREE multiplex-based methods. (c) Comparison of base editing efficiencies at Site-1, Site-2, and Site-3 in GFP-positive, GFP-negative, and unsorted cell populations using TREE-based methods to target these sites individually or in a multiplexed manner. $n=3$; N.S.=not significant.

FIG. 23 shows an analysis of off-target sites in multiplexed edited HEK293 cells using TREE- and RoT-based methods. GFP-positive cell populations isolated from TREE and RoT approaches were PCR-amplified and subjected to Sanger sequencing on the top predicted off-target loci for the sgRNA sequences for sg(BG) and genomic Sites 1-3. The C nucleotides in red text are potential Cs that can undergo C-to-T conversion within the editing window in the protospacer. The numbers below each C are quantification of the percentage of Cs of the Sanger sequence chromatograms using EditR.

FIG. 24 depicts the identification of exclusive targeting events in clonal population in edited HEK293 cells using TREE. Representative Sanger sequencing chromatographs of clonal cell populations that contain edits exclusively at the target C and not any other Cs within the editing window.

FIG. 25 demonstrates that TREE allows for base editing in hPSCs. (a) Representative flow cytometry plots in which TREE was employed in hPSCs utilizing (i) untransfected (ii) pCMV-BE4-Gam or (iii) pCMV-AncBE4. (b) Editing efficiency (percentage GFP-positive cells) of targeting in hPSCs line with various amounts of pEF-AncBE4 plasmid and ratios with the sg(BG) vector. $n=3$, $*=p<0.05$. (c) Representative Sanger sequencing chromatographs of Site-1 in GFP-positive, GFP-negative, and unsorted cell populations isolated with TREE- and RoT-based methods in which pEF-BE4-Gam or pEF-AncBE4 was utilized. (d) Representative flow cytometry plot of hPSCs in which TREE was applied targeting the APOE(R158) locus. (e) Representative Sanger sequencing chromatographs of APOE(R158) locus in GFP-positive, GFP-negative, and unsorted cell populations isolated with TREE-based methods.

FIG. 26 shows that TREE fluorescent output in hPSCs is transient. Representative flow cytometry plots of (i) untransfected hPSCs, (ii) TREE-enriched GFP-positive hPSCs 0 days (iii) 14 days after sorting.

FIG. 27 demonstrates a Next generation sequencing (NGS) analysis of allelic outcomes at target sites in hPSCs. NGS analysis for the target site when TREE was applied to edit Site-1 or the APOE(R158) in hPSCs. The number to left of the allelic outcome indicates the position upstream (5') relative to the PAM. Abbreviation: WT=wild-type unedited locus.

FIG. 28 depicts an Analysis of bystander editing in BIG-TREE edited hPSC clones. (a) Schematic of editing window and nucleotide position for target sites associated with FIG. 1 (APOE158R) and FIG. 2 (Genomic Sites 1-3). Red indicates target C within the editing window. Blue indicates bystander C within the editing window. (b) Distribution of bystander edits at the APOE(158R) locus in

clonal hPSCs. (c) Distribution of bystander edits at genomic Sites 1-3 in clonal hPSCs that were generated via multiplexed base editing.

FIG. 29 shows Off-target analysis of hPSC clones generated via BIG-TREE. Representative clonal lines were subjected to Sanger sequencing on the top predicted off-target loci for the sgRNA sequences for (a) sg(BG) and (b) sg(APOE158R).

FIG. 30 shows an Analysis of FAD-related genotypes in clonal hPSC lines generated from hPSC lines 2 and 3. Representative clonal lines derived from hPSC lines 2 and 3 were subject to Sanger sequencing to confirm that they retained the APP V717I or PSEN1 A246E FAD-related mutations after editing at the APOE(158R) locus.

FIG. 31 shows Validation of BIG-TREE-based gene knock-out (KO) using a HEK293-mCherry line. (a) Schematic of using BIG-TREE to introduce stop codons into the genomically integrated mCherry cassette. Successful targeting with sg(mCh1), sg(mCh2), or sg(mCh3) will result in a C-to-T conversion causing a change in the amino acid at position 47, 114, or 168, respectively, from a glutamine (Q) to a premature stop codon (X). (b) Representative phase contrast (top panels) and fluorescent (bottom panels) images of HEK293-mCherry cells that had been targeted with a control non-targeting sgRNA [sg(NT)] or a sgRNA targeting mCherry [sg(mCh1), sg(mCh2), or sg(mCh3)]. (c) Representative flow cytometry plots of HEK293-mCherry cells that been targeted with a control non-targeting sgRNA [sg(NT)] or a sgRNA targeting mCherry [sg(mCh1), sg(mCh2), or sg(mCh3)]. (d) Sanger sequencing of sg(mCh1-3) protospacer sequences with codon translations indicated. Sequencing indicates conversion from target 'CAG' (Q) to 'TAG' (X). (e) Correlation between observed mCherry loss by flow cytometry and percent conversion of the target C to T within Sanger sequencing reads.

FIG. 32 shows that transfection efficiency is not predictive of editing efficiency. HEK293 cells were transfected with pEF-mCherry, pCMV-ABEmax, and sg(TS). Comparison of transfection efficiency (percentage of mCherry-positive cells) and editing efficiency (percentage of A-to-G conversion at target nucleotides) in unsorted cell populations targeted at various genomic loci.

FIG. 33 shows flow cytometry-based characterization of XMAS-TREE reporter. Representative flow cytometry plots of HEK293 cells transfected with pEF-XMAS-1xStop or pEF-XMAS-2xStop, pCMV-ABEmax, and sg(NT) or sg(XMAS).

FIG. 34 demonstrates a comparison of XMAS-TREE editing efficiency in individual- or multiplexed-targeted genomic sites. Quantification of base editing efficiencies at targeted loci mCherry-positive/GFP-positive cell populations using XMAS-TREE-based targeting in a single or multiplexed manner. N.S.=not significant, $*=p<0.05$.

FIG. 35 shows off-target analysis of HEK293 clones generated using XMAS-TREE-based methods. Representative clonal lines were analyzed by Sanger sequencing at the top predicted off-target loci for the sgRNA sequences for (a) sg(XMAS), (b) sg(Site-1), (c) sg(Site-3), and (d) sg(Site-4).

FIG. 36 shows the characterization of XMAS-TREE reporter in hPSCs. Representative flow cytometry of hPSCs transfected pEF-XMAS-1xStop or pEF-XMAS-2xStop, pEF-ABEmax, and sg(NT) or sg(XMAS).

FIG. 37 demonstrates Representative Sanger sequencing chromatographs of edited hPSCs enriched using XMAS-TREE and RoT-based approaches. Sanger sequencing chromatographs of Site-3 and PSEN1 of unsorted hPSCs as well

11

as mCherry-positive and mCherry-positive/GFP-positive cells isolated using RoT-based and XMAS-TREE strategies, respectively.

FIG. 38 shows the distribution of genotypes in clonal hPSCs generated using XMAS-TREE-based methods. Analysis of clonal editing efficiency in hPSCs that were targeted at the PSEN1 locus.

FIG. 39 shows a vector map and sequence for pDT-sgRNA-1xStop.

FIG. 40 shows a vector map and sequence for pDT-sgRNA-2xStop.

FIG. 41 shows a vector map and sequence for pEF-XMAS-1xStop.

FIG. 42 shows a vector map and sequence for pEF-XMAS-2xStop.

FIG. 43 shows a vector map and sequence for pEF-ABE7.10.

FIG. 44 shows a vector map and sequence for pEF-ABEmax.

FIG. 45 shows sequences for mCherry-1xStop-GFP and mCherry-2xStop-GFP and XMAS 1xStop reporter sequences, XMAS 2xStop reporter sequences. FIG. 45 discloses SEQ ID NOS 311, 320, 312 and 321, respectively, in order of appearance.

FIG. 46 shows sequences for GFP and mCherry and the sequences of the XMAS 1x stop and XMAS 2x stop prior to and after editing. FIG. 46 discloses SEQ ID NOS 313-316, 326, 317-318, 327, and 319, respectively, in order of appearance.

DETAILED DESCRIPTION

The compositions and methods provided herein are based at least in part on the Inventors' development of real-time, fluorescent-based methods for identification and isolation of base-edited cell populations. In particular, the disclosure provides a transient reporter for editing enrichment (TREE) to efficiently select and isolate base-edited cells from non-edited cells. As described herein, TREE takes advantage of a detectable change in a reporter protein signal as a direct reporter of base editing activity within a cell. Compared to conventional cell enrichment strategies that employ reporters of transfection (RoT), TREE significantly improved the editing efficiency at multiple independent loci, with efficiencies approaching 80%. Using these methods, it is possible to target multiple separate loci for gene editing and to identify and separate base edited cells from non-edited cells, all without homology directed repair (HDR). The combination of TREE with these base-editing methods yield isogenic genetically modified human pluripotent stem cell lines, with single-nucleotide editing efficiencies of >80% across multiple hPSC lines. Also described herein are methods that are particularly advantageous for efficient generation of loss-of-function and gain-of-function hPSC lines via introduction of premature stop codons or other genetic modifications, and for multiplex editing of hPSCs at several independent loci. These methods allow for the precise and efficient base editing of hPSCs for use in developmental biology, disease modeling, drug screening, and cell-based therapies.

Accordingly, in a first aspect, a polynucleotide encoding one or more reporter protein. The polynucleotide includes a PAM site adjacent to a base that when edited causes a change in a function or characteristic of the one or more reporter proteins. In one embodiment, the polynucleotide encodes a reporter protein with at least 90% sequence similarity to SEQ ID NO: 2, wherein the polynucleotide encodes histidine at amino acid at position number 66, relative to SEQ ID

12

NO: 1 is provided, and encodes glycine at amino acid position number 72 relative to SEQ ID NO: 1. Alternatively, a reporter polypeptide may include two fluorescent proteins with at least 90% sequence identity to one of SEQ ID NO: 316 or 318. In another alternative, the polynucleotide comprises a polynucleotide selected from the group consisting of SEQ ID NO: 258, 259 and 260. The polypeptides may include a polypeptide having at least 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity to any one of the polypeptides provided herein. The polynucleotides encoding the polypeptides provided herein may include polynucleotides having at least 50%, 60%, 70%, 80%, 85%, 90%, 95%, 98%, 99% sequence identity to the polynucleotides provided herein. The polynucleotides encoding the polypeptides provided herein may be operably linked to a heterologous promoter. The polynucleotides may be included on constructs or in a vector, such as an expression vector, to allow for expression of the polypeptides in a cell.

The constructs and vectors provided herein may be prepared by methods available to those of skill in the art. Notably each of the constructs claimed are recombinant molecules and as such do not occur in nature. Generally, the nomenclature used herein and the laboratory procedures utilized in the present invention include molecular, biochemical, and recombinant DNA techniques that are well known and commonly employed in the art. Standard techniques available to those skilled in the art may be used for cloning, DNA and RNA isolation, amplification and purification. Such techniques are thoroughly explained in the literature.

The constructs provided herein may include a promoter operably linked to any one of the polynucleotides described herein. As used herein, the terms "heterologous promoter," "promoter," "promoter region," or "promoter sequence" refer generally to transcriptional regulatory regions of a gene, which may be found at the 5' or 3' side of the polynucleotides described herein, or within the coding region of the polynucleotides, or within introns in the polynucleotides. Typically, a promoter is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. The typical 5' promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. Within the promoter sequence is a transcription initiation site (conveniently defined by mapping with nuclease S1), as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase.

In some embodiments, the disclosed polynucleotides are operably connected to the promoter. As used herein, a polynucleotide is "operably connected" or "operably linked" when it is placed into a functional relationship with a second polynucleotide sequence. For instance, a promoter is operably linked to a polynucleotide if the promoter is connected to the polynucleotide such that it may effect transcription of the polynucleotides. Heterologous promoters useful in the practice of the present invention include, but are not limited to, constitutive, inducible, temporally-regulated, developmentally regulated, chemically regulated, tissue-preferred and tissue-specific promoters. The heterologous promoter may be a plant, animal, bacterial, fungal, or synthetic promoter.

Vectors including any of the constructs or polynucleotides described herein are provided. The term "vector" is intended to refer to a polynucleotide capable of transporting another

polynucleotide to which it has been linked. In some embodiments, the vector may be a “plasmid,” which refers to a circular double-stranded DNA loop into which additional DNA segments may be ligated. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome, such as some viral vectors or transposons, such as lentiviral vectors. Vectors may carry genetic elements, such as those that confer resistance to certain drugs or chemicals.

Cells including any of the polynucleotides, constructs, or vectors described herein are provided. Suitable “cells” that may be used in accordance with the present invention include eukaryotic or prokaryotic cells. Suitable eukaryotic cells include, without limitation, plant cells, fungal cells, and animal cells. Suitable prokaryotic cells include, without limitation, gram-negative and gram-positive bacterial species.

In a second aspect provided herein, is a kit comprising nucleic acid sequences that encode components having base editing activity when introduced into a cell as well as components that, when expressed in the cell, function as a transient reporter of successful base editing. As described herein, the transient reporter-containing compositions and methods of their use enable enrichment of base edited cells.

In some cases, the kit comprises a one or more vectors comprising nucleic acid sequences that encode elements that perform base editing and the transient reporter of base editing enrichment. In some cases, the composition comprises one or more vectors comprising a first nucleic acid encoding one or more reporter proteins, wherein the first nucleic acid includes a PAM site adjacent to a base that when edited causes a change in a function or characteristic of the one or more reporter proteins, a second nucleic acid sequence encoding a base editing targeting cassette (preferably comprising one or more sgRNAs and protospacer adjacent motifs (PAMs) directed to the nucleic acid sequence that encodes the reporter protein), a third nucleic acid sequence encoding a base editing targeting cassette (preferably comprising one or more sgRNAs and protospacer adjacent motifs (PAMs) directed to a nucleic acid sequence of interest), and a fourth nucleic acid sequence encoding a base editor,

Any appropriate base editor capable of single nucleotide modifications without a need for double stranded DNA breaks can be used. In some cases, the base editor is a cytidine deaminase base editor (CBE) or adenine base editor (ABE). CBEs and ABEs install C•G-to-T•A and A•T-to-G•C transitions, respectively, and have been successfully used in many cell types including mammalian cells and plant cells. Cytidine deaminase base editors convert cytidine to uridine within a small editing window near the protospacer adjacent motif (PAM) site. Uridine is subsequently converted to thymidine through base excision repair, creating a C-to-T conversion (or G-to-A on the opposite strand). ABEs can convert adenine into inosine through deamination in any ABE protospacer (e.g. NGG, NG, and more) adjacent motif. The editing window will vary based on the type of base editor. For instance, the editing window for ABE is typically 12 to 17 nucleotides upstream of the ABE protospacer adjacent motif.

In some cases, the base editor is a prime editor. Prime editors are engineered Cas9 nickase-reverse transcriptase (RT) fusion proteins. When used in combination with prime

editing guide RNAs (pegRNAs) that encode the desired edit, prime editors can edit bases in plant and animal cells without donor DNA or double-strand breaks. Unlike CBEs and ABE, prime editors are able to introduce point mutations, insertions, deletions, and all 12 possible base-to-base conversions. See Anzalone et al., i 576:149-157 (2019).

In some cases, the base editor is a Cas9-mediated adenosine base editor (referred to herein as “XMAS”). As described in Example 3, use of a Cas9-mediated adenosine base editor according to the methods of this disclosure introduces an A-to-G conversion. When a Cas9-mediated adenosine base editor used, for example, to target a TGA stop codon located between coding sequence for two different reporter proteins, the A-to-G conversion changes the TGA stop codon to TGG. Accordingly, expression of both reporter proteins allows for the real-time detection of adenosine base editing.

In some cases, the base editor is a dual adenine and cytosine base editor (e.g. CRISPR-Cas9-based synchronous programmable adenine and cytosine editor (SPACE), a codon-optimized fusion of cytosine deaminase PmCDA1, adenosine deaminase TadaA, and a Cas9 nickase (Target-ACEmax), or a fusion of both adenine and cytosine deaminases with a Cas9 nickase (A&C-BEmax). When a dual editor is used, both C-T and A-G conversions happen simultaneously without the need for two base editor constructs.

In some cases, the base editor is any enzyme capable of modifying nucleobases is fused to a catalytically inactivated or impaired zinc finger nuclease (ZFN), or a transcription activator like effector (TALE). A ZF or TALE is designed to bind to a specific region of DNA, eliminating the potentially narrow editing window found with CRISPR systems.

In the context of the present disclosure, the following abbreviations for the commonly occurring nucleic acid bases are used. “A” refers to adenosine, “C” refers to cytosine, “G” refers to guanosine, “T” refers to thymidine, and “U” refers to uridine.

The reporter protein can be any detectable protein (e.g., detectable by fluorescence, color, bioluminescence indirect reporter system) for which a single base-to-base conversion or, in some cases, an insertion or deletion resulting in an observable change in the detectable protein. For example, the reporter protein can comprise a single nucleotide variation relative to the wild-type reporter protein. When a base-to-base conversion or other mutation is introduced into sequence encoding the reporter protein by base editing, the expressed reporter protein exhibits a detectable change (e.g., change in emitted fluorescent, change in color, change by way of indirect reporter) relative to the non-edited reporter protein. Examples of reporter proteins appropriate for this disclosure include, without limitation, fluorescent proteins (e.g., green fluorescent protein (GFP) and variants thereof, blue fluorescent protein (BFP) and variants thereof, red fluorescent protein (RFP) and variants thereof), luciferase and variants thereof, mCherry), β -galactosidase (lacZ), chloramphenicol acetyltransferase (CAT), β -glucuronidase (GUS), secretory alkaline phosphatase (SEAP), survival selection genes such as but not limited to antibiotic resistance, auxotrophies, flux redirection, toxin pumps, biosensors, reporter proteins that directly or indirectly produce or catalyze a colorimetric reaction, and those set forth in Table 1 below. In some cases, the reporter protein can be a blue fluorescent protein (BFP) variant that comprises a single nucleotide variation relative to the wild-type reporter protein. Referring to FIGS. 1A-1F, a blue fluorescent protein (BFP) variant can comprise a histidine at amino acid position 66 (numbered relative to SEQ ID NO:1) where the

histidine is encoded by a C-A-C codon. In some cases, the reporter protein is a mutated, inactive form of luciferase. In such cases, it is advantageous to design a nucleic acid encoding the mutant luciferase such that a single base-to-base conversion or other genetic modification will introduce a corrective edit, resulting in expression of an active luciferase enzyme. In other embodiments the reporter protein is a fusion protein of two fluorescent proteins linked via a linker including at least one stop codon and a PAM site positioned to allow targeting and editing of the stop codon into a codon coding for an amino acid. Base editing allows for translation of the second reporter in the fusion protein and expression of the second reporter. The linker is positioned between the two reporter proteins such that expression of the reporter construct comprising the fusion protein in a cell results in expression of only the first reporter protein in the fusion protein. Once the polynucleotide encoding the fusion protein is edited, then both the first and second reporter will be detected. In this case the first reporter may be used as a measure of transfection efficiency and the second may be used to monitor and select for gene edited cells.

TABLE 1

Exemplary Reporter Proteins and Base Edits			
Gene Name:	Edit:	Function:	Notes:
Blue Fluorescent Protein (BFP)	C-to-T (CBE) H66Y	Change emission spectra from 448 nm to 508 nm	TREE
Green Fluorescent Protein (GFP)	A-to-G (ABE) Y66H	Change emission spectra from 508 nm to 448 nm	TREE reaction run in reverse
Activation of fluorescent Proteins	A-to-G (ABE) Stop-to-W	Removal of stop codon in coding sequence	XMAS-TREE
Introduction of premature stop codons	C-to-T (CBE) Q-to-Stop	Addition of premature stop codons	BIG-TREE
Activation of antibiotic resistance cassette	A-to-G (ABE) Stop-to-W	Removal of stop codon in coding sequence	
Activation of auxotrophic selection marker	A-to-G (ABE) Stop-to-W	Removal of stop codon in coding sequence	
Activation of gene function via editing splice junctions	CBE or ABE	Editing to restore functional splice junction in reporter	

In some cases, the nucleic acid sequence that encodes a base editing cassette comprises a sgRNA adjacent to (e.g., located 5' to) a protospacer adjacent motif (PAM), where the sgRNA comprises a protospacer sequence and targets a portion of the nucleic acid sequence encoding the reporter protein for base editing. As used herein, the term "base editing cassette" refers to an expression cassette or framework comprising nucleic acid sequence encoding a RNA oligonucleotide containing, in some cases, 18-22 base pairs (the single guide RNA or "sgRNA") and PAM sequence. As used herein, a "single guide RNA" (sgRNA) is nucleotide sequence that is complementary to at least a portion of a target nucleic acid to be genetically modified by a base

editor. Generally, a sgRNA comprises a nucleotide sequence that is partially or wholly complementary to a target sequence (such as a target genomic sequence or sequence in an expression vector) and comprises a target base pair. A sgRNA target site also comprises a PAM located immediately downstream from the target site. In some cases, the PAM is a *S. pyogenes* Cas9 PAM 'NGG.' The PAM sequence may vary if other Cas9 variants are used. For some embodiments, the sgRNA preferably comprises a sequence of at least 10 contiguous nucleotides, and often a sequence of 18-22 contiguous nucleotides or more. In some embodiments, a sgRNA molecule can be from 20 to 300 or more bases in length, or more. In certain embodiments, a sgRNA molecule can be from 20 to 300 bases in length, or 20 to 120 bases, or 30 to 50 bases, or 39 to 46 bases. In some cases, no sgRNA is needed.

As used herein, the term "encoding" refers to the inherent property of specific sequences of nucleotides in a polynucleotide, such as a gene, a cDNA, or an mRNA, to serve as templates for synthesis of other polymers and macromolecules in biological processes having either a defined sequence of nucleotides (i.e., rRNA, tRNA and mRNA) or a defined sequence of amino acids and the biological properties resulting therefrom. Thus, a gene encodes a protein if transcription and translation of mRNA corresponding to that gene produces the protein in a cell or other biological system. Both the coding strand, the nucleotide sequence of which is identical to the mRNA sequence and is usually provided in sequence listings, and the non-coding strand, used as the template for transcription of a gene or cDNA, can be referred to as encoding the protein or other product of that gene or cDNA.

As used herein, the term "target site" or "target sequence" refers to a genomic nucleic acid sequence that defines a portion of a nucleic acid to which biological molecules involved in base editing may specifically bind under conditions sufficient for binding to occur.

Preferably, nucleic acid sequences and vectors comprising sgRNAs are designed to allow for the facile cloning of new target sites via restriction enzyme digestion and ligation of oligonucleotides that target the desired genomic sequence. Depending on the PAM specificity and editing window configured in the base editing cassettes, the methods can be used in conjunction with any base editors or base editor variants. For instance, the PAM sequence and edit distance can be modified to match the editing specificity and window of the new base editor. Such modifications are straightforward to achieve with the BFP vector using TREE, or the stop codon between RFP and GFP using BIG-TREE.

By way of example, the sgRNA can be designed for editing of a nucleic acid sequence encoding a BFP variant as the reporter protein. As described in the Examples that follow, the sgRNA is designed to bring a cytidine deaminase base editor to the BFP variant-encoding nucleic acid sequence for base editing. Referring to FIG. 1A, the protospacer sequence (underlined black) for the sgRNA, sg(BG), targeting the 'CAC' codon (underlined blue) resulting in a C-to-T conversion to 'TAC' (underlined green) and the corresponding amino acid change of histidine (blue) to tyrosine (green) at the 66th amino acid position in BFP. A PAM (underlined red) was placed in the position to orient the base editing window (underlined orange) around the C nucleotide (red) to facilitate BFPH66 to GFPY66 conversion.

In some cases, the amino acid sequence for wild-type BFP is

(SEQ ID NO: 1)
 MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGDATYGKLTCLKFICT
 TGKLPVPWPTLVTTLTLYGVQCFSRYPDHMKQHDFFKSAMPEGYVQERTIF
 FKDDGNYKTRAEVKFEGDTLVNRIELKGIDEDKGNILGHKLEYNNSHN
 VYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNH
 YLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK.

In some cases, the amino acid sequence for variant BFP is

(SEQ ID NO: 2)
 MVSKGEELFTGVVPILVELDGDVNGHKFSVSGEGDATYGKLTCLKFICT
 TGKLPVPWPTLVTTLTHGVQCFGRYPDHMKQHDFFKSAMPEGYVQERTIF
 FKDDGNYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNNSHN
 VYIMADKQKNGIKVNFKIRHNIEDGSVQLADHYQQNTPIGDGPVLLPDNH
 YLSTQSALS KDPNEKRDHMLLEFVTAAGITLGMDELYK*.

In SEQ ID NO:2, the histidine at residue position 66 is underlined. The S-to-G modification to introduce the PAM site is double underlined.

Preferably, nucleic acid sequence encoding the reporter protein is operably linked to a promoter that drives expression of the reporter protein upon introduction into a cell. A promoter, generally, is a region of nucleic acid that initiates transcription of a nucleic acid encoding a product. A promoter may be located upstream (e.g., 0 bp to -100 bp, -30 bp, -75 bp, or -90 bp) from the transcriptional start site of a nucleic acid encoding a product, or a transcription start site may be located within a promoter. A promoter may have a length of 100-1000 nucleotide base pairs, or 50-2000 nucleotide base pairs. In some embodiments, promoters have a length of at least 2 kilobases (e.g., 2-5 kb, 2-4 kb, or 2-3 kb). The term "expression" as used herein is defined as the transcription and/or translation of a particular nucleotide sequence driven by its promoter. As used herein, the term "operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame. As exemplified in FIG. 1B, the promoter can be human EF1a promoter, which is a strong constitutively active promoter, however other promoters can be used. In some cases the promoter is an inducible promoter or a cell-type specific promoter. Examples of base editing in different organisms and plus DOI citations for each example are set forth in Table 2.

TABLE 2

Examples of Base Editing in Different Organisms		
	C-T	A-G
<u>bacteria</u>		
<i>E. coli</i>	10.1038/ncomms13330, 10.1038/s42003-018-0035-5	
<i>B. melitensis</i>	10.1038/s42003-018-0035-5	
<i>Streptomyces</i>	10.1073/pnas.1913493116	
<i>C. glutamicum</i>	10.1002/bit.27121	
<u>yeast</u>		
<i>S. cerevisiae</i>	10.1126/science.aaf8729	
<u>plants</u>		
rice, wheat, maize	10.1038/nbt.3811	
rice and tomato	10.1038/nbt.3833	
rice and wheat	10.1038/s41587-020-0455-x	
<u>mammalian</u>		
mouse embryo	10.1038/s41598-018-33533-5,	
mice	10.1038/s41556-018-0202-4,	
	10.1038/nbt.3816	
	10.1038/nbt.4194	
	10.1038/nature26155,	
	10.1038/s41587-019-0134-y,	
	10.1016/j.molcel.2017.08.008,	
	10.1038/nbt.4102,	
	10.1038/s41467-019-11562-6,	10.1038/s41587-019-0050-1,
	10.1038/ncomms15790,	10.1038/nbt.4172,
	10.1038/nbt.3803,	10.1038/nature24644,
	10.1038/nature17946,	10.1016/j.ymthe.2019.05.013,
human cells	10.1038/nmeth.4027	10.1038/s41587-020-0491-6
mouse embryo		10.1038/s41467-018-04768-7
Pig	10.1038/s41467-019-10421-8	
mice, mouse zygote, mammalian		10.1016/j.ymthe.2019.11.022
mice and mice embryo		10.1038/nbt.4148
mice organs	10.1038/s41551-019-0501-5	10.1038/s41551-019-0501-5
dual base editing	10.1038/s41587-020-0527-y,	
	10.1038/s41587-020-0509-0,	
	10.1038/s41587-020-0535-y	

As used herein, the term “vector” is intended to mean a nucleic acid molecule capable of transporting another nucleic acid. By way of example, appropriate vectors for the compositions and methods of this disclosure include episomal vectors, viral vectors (e.g., retrovirus, adenovirus, baculovirus), plasmids, RNA vectors, or linear or circular DNA or RNA molecules which may comprise or consist of a chromosomal, non-chromosomal, semi-synthetic, or synthetic nucleic acid. Large numbers of suitable vectors are known to those of skill in the art and commercially available. Preferred vectors are episomal vectors, which are capable of autonomous replication due to the presence of an origin of replication. Preferably, vectors of the base editing compositions described herein are episomal vectors, meaning they are capable of autonomous replication due to the presence of an origin of replication. While the nucleic acid sequences can be provided on separate vectors, it will be understood that it is possible to configure a single or pair of expression vectors comprising the base editing elements described herein.

Methods

In another aspect, provided herein are methods for genome engineering (e.g., for altering or manipulating the expression of one or more genes or one or more gene products) in cells, in vitro, in vivo, or ex vivo. In particular, the methods provided herein are useful for targeted base editing or base correction in any cell.

In some cases, the methods comprise multiplex editing at multiple loci. In such cases, the methods described herein can be performed using a vector comprising dual-targeting sgRNAs. In such cases, a first nucleic acid sequence encodes a sgRNA for base editing in the reporter protein, and a second nucleic acid sequence encodes a sgRNA for a genomic target site. In some cases, methods described herein can be performed using a vector comprising a multiplex-targeting sgRNAs. In such cases, a first nucleic acid sequence encodes a sgRNA for base editing in the reporter protein, and a second, third, fourth, etc. nucleic acid sequence encodes a sgRNA for a genomic target site. When introduced into a cell with nucleic acid sequences encoding the base editor and reporter protein, it is possible to base edit two or more genomic target sites and also promote a base-to-base conversion or other genetic modification that yields a detectable change in fluorescent emission or color of the transient reporter protein. The Examples demonstrate successful base editing at two or more genomic target sites and plus introduction of a C-to-T conversion in nucleic acid sequence encoding a BFP variant, thus resulting in a shift in fluorescent emission spectra from BFP to GFP as described herein. In this manner, base-edited cells can be sorted and selected based on fluorescence emission spectra or other detectable signals. Accordingly, also provided herein are engineered cells that have been genetically modified according to these methods.

Although human pluripotent stem cells are exemplified herein, it will be understood by practitioners in the art that the base editing compositions and methods can be used with other cell types, including a variety of human cell types and cells of other types of animals. In some cases, the cell is a mammalian cell. Preferably, the mammalian cell is a human cell. Cells appropriate for use in the methods of this disclosure include, without limitation, pluripotent stem cells, multipotent cells, dissociated organs or organoids, terminally differentiated cells, immune cells, hematopoietic stem progenitor cells (HSPCs) (e.g., umbilical cord blood HSPCs), and fibroblasts. In addition to mammalian cells, it will be understood by practitioners in the art that the base

editing compositions and methods can be used with other cell types such as bacteria cells, yeast cells, plant cells, and other single celled organisms.

As used herein, the term “pluripotent stem cell” (hPSC) means a cell capable of continued self-renewal and capable, under appropriate conditions, of differentiating into cells of all three germ layers. hPSCs exhibit a gene expression profile that includes SOX2⁺ and OCT4⁺. Examples of human PSCs (hPSCs) include human embryonic stem cells (hESCs) and human induced pluripotent stem cells (hiPSCs). As used herein, “iPS cells” or “iPSCs” refer to cells that are substantially genetically identical to their respective differentiated somatic cell of origin and display characteristics similar to higher potency cells, such as ES cells, as described herein. The cells can be obtained by reprogramming non-pluripotent (e.g., multipotent or somatic) cells. In some cases, the modified cells are human pluripotent stem cells such as human embryonic stem cells or induced pluripotent stem cells. In some cases, the modified cells are human embryonic stem cells isolated from human embryonic tissues. In other cases, the modified cells are cells isolated from human blastocysts and then modified. In some cases, the modified cells are human placental or umbilical cord stem cells.

Induced pluripotent stem cells exhibit morphological properties (e.g., round shape, large nucleoli and scant cytoplasm) and growth properties (e.g., doubling time of about seventeen to eighteen hours) akin to ESCs. In addition, iPS cells express pluripotent cell-specific markers (e.g., Oct-4, SSEA-3, SSEA-4, Tra-1-60 or Tra-1-81, but not SSEA-1). Induced pluripotent stem cells, however, are not immediately derived from embryos. As used herein, “not immediately derived from embryos” means that the starting cell type for producing iPS cells is a non-pluripotent cell, such as a multipotent cell or terminally differentiated cell, such as somatic cells obtained from a post-natal individual.

Subject-specific somatic cells for reprogramming into induced pluripotent stem cells can be obtained or isolated from a target tissue of interest by biopsy or other tissue sampling methods. In some cases, subject-specific cells are manipulated in vitro prior to use in a method of this disclosure. For example, subject-specific cells can be expanded, differentiated, genetically modified, contacted to polypeptides, nucleic acids, or other factors, cryo-preserved, or otherwise modified prior to use according to the methods of this disclosure.

For the methods described herein, gene editing systems or components thereof (e.g., a vector encoding a base editor protein, a gRNA) are introduced into a cell (e.g., a human pluripotent stem cell). As used herein, the term “introducing” encompasses a variety of methods of introducing DNA into a cell, either in vitro or in vivo, such methods including transformation, transduction, transfection (e.g., electroporation), nucleofection (an electroporation-based transfection method which enables transfer of nucleic acids such as DNA and RNA into cells by applying a specific voltage and reagents) and infection. Where the introducing involves electroporation (e.g., nucleofection), a polynucleotide (e.g., a plasmid, a single stranded DNA, a minicircle DNA, RNA) is electroporated into a target cell. Vectors are useful for introducing DNA encoding molecules into cells. Any appropriate delivery vector can be used with the methods described herein. For example, delivery vectors include exosomes, viruses (viral vectors), and viral particles. Preferably, the delivery vector is a viral vector, such as a lentiviral or baculo- or preferably adeno-viral/adeno-associated viral (AAV) vectors, but other non-viral means of delivery are

known (such as yeast systems, microvesicles, gene guns/ means of attaching vectors to gold nanoparticles). Other methods of introducing a nucleic acid into a host cell are known in the art, and any known method can be used to introduce a nucleic acid (e.g., vector or expression construct) into a cell for the methods provided herein. Suitable methods include, without limitation, viral or bacteriophage infection, transfection, conjugation, protoplast fusion, lipofection, electroporation, calcium phosphate precipitation, polyethyleneimine (PEI)-mediated transfection, DEAE-dextran mediated transfection, liposome-mediated transfection, particle gun technology, calcium phosphate precipitation, direct micro injection, nanoparticle-mediated nucleic acid delivery (see, e.g., Panyam et al., *Adv. Drug Deliv. Rev.*), and the like. Delivery of components may also include use of ribonucleo- proteins complexed with sgRNA alone, or in combination with nucleic acid delivery.

A cell is “genome edited” or “genetically modified” if the cell includes a modification to its genome compared to a non-genome edited cell of the same type. In some cases, a non-genome edited cell is a wild-type cell. As used herein, the terms “genetically modified” and “genetically engineered” are used interchangeably and refer to a prokaryotic or eukaryotic cell that includes an exogenous polynucleotide, regardless of the method used for insertion. In some cases, a cell has been modified to comprise a non-naturally occurring nucleic acid molecule that has been created or modified by the hand of man (e.g., using recombinant DNA technology) or is derived from such a molecule (e.g., by transcription, translation, etc.). A cell that contains an exogenous, recombinant, synthetic, and/or otherwise modified polynucleotide is considered to be an engineered or “genome edited” cell. Genetically editing or modifying a cell refers to modifying cellular nucleic acid within a cell, including genetic modifications to endogenous and/or exogenous nucleic acids within the cell. Genetic modifications can comprise deletions, insertions, integrations of exogenous DNA, gene correction and/or gene mutation.

The term “substantially pure cell composition of genetically modified cells” as used herein refers to a cell composition comprising at least 70%, more preferentially at least 90%, most preferentially at least 95% of genetically modified cells in the cell composition obtained by methods of this disclosure. The terms “purified” or “enriched” cell populations are used interchangeably herein, and refer to cell populations, in vitro or ex vivo, that contain a higher proportion of a specified cell type or cells having a specified characteristic than are found in vivo (e.g., in a tissue).

As used herein, the term “isogenic” refers to cells or organisms that are genetically related, or having the same or closely similar genotypes, such as cells of a cell line. For example, cells of a clonal population of cells are isogenic to each other. In some cases, a first population of human pluripotent stem cells can have a wild-type, genetically unmodified genome, and a second population of pluripotent stem cells can be isogenic to the first population except that they have been genetically modified (which term as used herein includes progeny of modified cells) to comprise a particular genetic modification. Individual cells of the second population may be isogenic to each other if obtained by clonal expansion of a single genetically modified cell.

In some cases, cells into which nucleic acid sequences encoding a base editor, reporter protein, and base editing cassette have been introduced and then cultured for about 48 to about 72 hours are sorted using any sorting technique capable of detecting expression of green fluorescent protein and, optionally, other cell markers. Methods and techniques

for assessing the expression and/or levels of cell markers are known in the art. Antibodies and reagents for detection of such markers are well known in the art, and readily available. Assays and methods for detecting such markers include, but are not limited to, flow cytometry, including intracellular flow cytometry, ELISA, ELISPOT, cytometric bead array or other multiplex methods, Western Blot and other immunoaffinity-based methods. In preferred embodiments, cells genetically modified according to the methods of this disclosure are detected and sorted by fluorescence-based flow cytometry. As used herein, the term “flow cytometry” refers to a cell analysis technique that detects and measures physical and chemical characteristics of a population of cells or particles in a rapidly flowing fluid stream as they pass in front of a viewing aperture. The term “flow cytometry” encompasses fluorescence-based flow cytometry in which, generally, lasers are used to detect and count cells based on fluorescence emission of fluorophores associated with the cells. In some cases, flow cytometry is performed using a specialized flow cytometer known as a fluorescence activated cell sorter (FACS). Cell sorters like FACS use fluidics and fluorescence components similar to those in flow cytometers, but are able to divert a specific population from within a heterogeneous sample into a separate tube, typically based on specified fluorescence characteristics.

Any appropriate technique can be used to as an additional means to confirm that base editing has occurred. For example, Sanger sequencing or next generation sequencing (NGS) can be used to detect C-to-T conversions.

As used herein, the terms “complementary” or “complementarity” are used in reference to “polynucleotides” and “oligonucleotides” (which are interchangeable terms that refer to a sequence of nucleotides) related by the base-pairing rules. For example, the sequence “5'-C-A-G-T,” is complementary to the sequence “5'-A-C-T-G.” Complementarity can be “partial” or “total.” “Partial” complementarity is where one or more nucleic acid bases is not matched according to the base pairing rules. “Total” or “complete” complementarity between nucleic acids is where each and every nucleic acid base is matched with another base under the base pairing rules.

In another aspect, provided herein is a cell culture composition comprising the isogenic line of genetically modified human pluripotent stem cells produced according to the methods of this disclosure and a chemically defined culture medium. The term “chemically defined culture medium” or “chemically defined medium,” as used herein, means that the molecular identity, chemical structure, and quantity of each medium ingredient is definitively known. The term “ingredient,” as used herein, refers to a component the molecular identity and quantity of which is known. In some cases, a chemically defined medium is made xeno-free, and incorporates human proteins, which can be produced using recombinant technology or derived from placenta or other human tissues, in lieu of animal-derived proteins. In some embodiments, all proteins added to the medium are recombinant human proteins.

In some cases, the method is performed using human induced pluripotent stem cells obtained by reprogramming a somatic cell of a human subject, such as a human subject that has a genetic disorder caused by a single nucleotide polymorphism. In such cases, the resulting base-edited human cells are autologous to the human subject and, in some cases, can be administered back to the subject (e.g., cell therapy).

In another aspect, provided herein is an isogenic population of human cells genetically modified ex vivo to comprise a base-to-base conversion or other genetic modification. In

some cases, the base-to-base conversion corrects or suppress a disease-associated single nucleotide polymorphism (SNP) or modifies an disease-associated isoform of a particular protein. As demonstrated in the Examples, the base editing-transient reporter enrichment methods of this disclosure were used to modify a nucleotide sequence encoding a APOE2 isoform of human APOE protein, such that the modified cell did not express any other isoform of the APOE protein. In this example, the human cells were induced pluripotent stem cells obtained from a somatic cell of a human subject having Familial Alzheimer's disease (FAD). Although correction of a SNP associated with FAD is exemplified herein, the methods of this disclosure can be used to correct introduce disease-correcting or disease-suppressing base edits in human cells provided that there is an appropriate PAM in the correct location downstream of the target SNP.

The terms "nucleic acid" and "nucleic acid molecule," as used herein, refer to a compound comprising a nucleobase and an acidic moiety, e.g., a nucleoside, a nucleotide, or a polymer of nucleotides. Nucleic acids generally refer to polymers comprising nucleotides or nucleotide analogs joined together through backbone linkages such as but not limited to phosphodiester bonds. Nucleic acids include deoxyribonucleic acids (DNA) and ribonucleic acids (RNA) such as messenger RNA (mRNA), transfer RNA (tRNA), etc. Typically, polymeric nucleic acids, e.g., nucleic acid molecules comprising three or more nucleotides are linear molecules, in which adjacent nucleotides are linked to each other via a phosphodiester linkage. In some embodiments, "nucleic acid" refers to individual nucleic acid residues (e.g. nucleotides and/or nucleosides). In some embodiments, "nucleic acid" refers to an oligonucleotide chain comprising three or more individual nucleotide residues. As used herein, the terms "oligonucleotide" and "polynucleotide" can be used interchangeably to refer to a polymer of nucleotides (e.g., a string of at least three nucleotides). In some embodiments, "nucleic acid" encompasses RNA as well as single and/or double-stranded DNA. Nucleic acids may be naturally occurring, for example, in the context of a genome, a transcript, an mRNA, tRNA, rRNA, siRNA, snRNA, a plasmid, cosmid, chromosome, chromatid, or other naturally occurring nucleic acid molecule. On the other hand, a nucleic acid molecule may be a non-naturally occurring molecule, e.g., a recombinant DNA or RNA, an artificial chromosome, an engineered genome, or fragment thereof, or a synthetic DNA, RNA, DNA/RNA hybrid, or include non-naturally occurring nucleotides or nucleosides. Furthermore, the terms "nucleic acid," "DNA," "RNA," and/or similar terms include nucleic acid analogs, i.e. analogs having other than a phosphodiester backbone. Nucleic acids can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, etc. Where appropriate, e.g., in the case of chemically synthesized molecules, nucleic acids can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, and backbone modifications. A nucleic acid sequence is presented in the 5' to 3' direction unless otherwise indicated. In some embodiments, a nucleic acid is or comprises natural nucleosides (e.g. adenosine, thymidine, guanosine, cytidine, uridine, deoxyadenosine, deoxythymidine, deoxyguanosine, and deoxycytidine); nucleoside analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoado-

sine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (e.g., methylated bases); intercalated bases; modified sugars (e.g., 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (e.g., phosphorothioates and 5'-N-phosphoramidite linkages).

Nucleic acids and/or other constructs (including cell populations) described in this disclosure may be isolated. As used herein, "isolated" means altered or removed from the natural state. For example, a nucleic acid or a peptide naturally present in a living animal is not "isolated," but the same nucleic acid or peptide partially or completely separated from the coexisting materials of its natural state is "isolated." An isolated nucleic acid or protein can exist in substantially purified form, or can exist in a non-native environment such as, for example, a host cell.

The terms "protein," "peptide," and "polypeptide" are used interchangeably herein and refer to a polymer of amino acid residues linked together by peptide (amide) bonds. The terms refer to a protein, peptide, or polypeptide of any size, structure, or function. Typically, a protein, peptide, or polypeptide will be at least three amino acids long. A protein, peptide, or polypeptide may refer to an individual protein or a collection of proteins. One or more of the amino acids in a protein, peptide, or polypeptide may be modified, for example, by the addition of a chemical entity such as a carbohydrate group, a hydroxyl group, a phosphate group, a farnesyl group, an isofarnesyl group, a fatty acid group, a linker for conjugation, functionalization, or other modification, etc. A protein, peptide, or polypeptide may also be a single molecule or may be a multi-molecular complex. A protein, peptide, or polypeptide may be just a fragment of a naturally occurring protein or peptide. A protein, peptide, or polypeptide may be naturally occurring, recombinant, or synthetic, or any combination thereof. A protein may comprise different domains, for example, a nucleic acid binding domain and a nucleic acid cleavage domain. In some embodiments, a protein comprises a proteinaceous part, e.g., an amino acid sequence constituting a nucleic acid binding domain.

Nucleic acids, proteins, and/or other compositions (e.g., cell population) described herein may be purified. As used herein, "purified" means separate from the majority of other compounds or entities, and encompasses partially purified or substantially purified. Purity may be denoted by a weight by weight measure and may be determined using a variety of analytical techniques such as but not limited to mass spectrometry, HPLC, etc.

Articles of Manufacture

In another aspect, provided herein is an article of manufacture such as a kit comprising a composition comprising expression vectors to perform base editing and editing enrichment using TREE as described herein. In certain embodiments, the kit comprises a plurality of vectors to achieve targeted base editing and for expression of the transient reporter (TREE). In some cases, the kit further comprises reagents and other materials useful for introducing vectors into cells and for culturing modified cells according to the methods. In some cases, the kit further comprises instructions for performing the methods of this disclosure.

In another aspect, provided herein is a kit for generating substantially pure populations of base-edited cells including, as a non-limiting example, base-edited human pluripotent stem cells. In exemplary embodiments, the kit comprises one or more of (i) a culture medium suitable for maintaining cells in vitro or ex vivo; (ii) base editing vectors as described

herein; (iii) reagents for introduction of the base editing vectors into cells (e.g., transfection reagents, transduction reagents, electroporation reagents), and (iv) instructions describing a method for generating substantially pure populations of base-edited cells as described herein, the method employing one or more of the culture medium, the base editing vectors, and the reagents for introducing vectors into the cells. In some cases, the kit further comprises reagents and/or materials for flow cytometry and cell sorting using FACS.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as though set forth in their entirety in the present application.

In interpreting this disclosure, all terms should be interpreted in the broadest possible manner consistent with the context. It is understood that certain adaptations of the invention described in this disclosure are a matter of routine optimization for those skilled in the art, and can be implemented without departing from the spirit of the invention, or the scope of the appended claims.

So that the compositions and methods provided herein may more readily be understood, certain terms are defined:

As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. Any reference to “or” herein is intended to encompass “and/or” unless otherwise stated.

The phrase “and/or,” as used herein in the specification and in the claims, should be understood to mean “either or both” of the elements so conjoined, i.e., elements that are conjunctively present in some cases and disjunctively present in other cases. Multiple elements listed with “and/or” should be construed in the same fashion, i.e., “one or more” of the elements so conjoined. Other elements may optionally be present other than the elements specifically identified by the “and/or” clause, whether related or unrelated to those elements specifically identified. Thus, as a non-limiting example, a reference to “A and/or B”, when used in conjunction with open-ended language such as “comprising” can refer, in one embodiment, to A only (optionally including elements other than B); in another embodiment, to B only (optionally including elements other than A); in yet another embodiment, to both A and B (optionally including other elements); etc.

The terms “comprising”, “comprises” and “comprised of” as used herein are synonymous with “including”, “includes” or “containing”, “contains”, and are inclusive or open-ended and do not exclude additional, non-recited members, elements, or method steps. The phraseology and terminology used herein is for the purpose of description and should not be regarded as limiting. The use of “including,” “comprising,” “having,” “containing,” “involving,” and variations thereof, is meant to encompass the items listed thereafter and additional items. Embodiments referenced as “comprising” certain elements are also contemplated as “consisting essentially of” and “consisting of” those elements. Use of ordinal terms such as “first,” “second,” “third,” etc., in the claims to modify a claim element does not by itself connote any priority, precedence, or order of one claim element over another or the temporal order in which acts of a method are performed. Ordinal terms are used merely as labels to distinguish one claim element having a certain name from another element having a same name (but for use of the ordinal term), to distinguish the claim elements.

The terms “about” and “approximately” shall generally mean an acceptable degree of error for the quantity mea-

sured given the nature or precision of the measurements. “About” as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass variations of $\pm 20\%$ or $\pm 10\%$, more preferably $\pm 5\%$, even more preferably $\pm 1\%$, and still more preferably $\pm 0.1\%$ from the specified value, as such variations are appropriate to perform the disclosed methods. Alternatively, and particularly in biological systems, the terms “about” and “approximately” may mean values that are within an order of magnitude, preferably within 5-fold and more preferably within 2-fold of a given value. Numerical quantities given herein are approximate unless stated otherwise, meaning that the term “about” or “approximately” can be inferred when not expressly stated.

Unless otherwise defined, all technical terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. As used herein and in the claims, the singular forms “a,” “an,” and “the” include the singular and the plural reference unless the context clearly indicates otherwise. Thus, for example, a reference to “an agent” includes a single agent and a plurality of such agents. Any reference to “or” herein is intended to encompass “and/or” unless otherwise stated.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as though set forth in their entirety in the present application.

EXAMPLES

Example 1—Transient Reporter for Editing Enrichment (TREE)

Current approaches to identify cell populations that have been modified with deaminase base editing technologies are inefficient and rely on downstream sequencing techniques. In this example, a blue fluorescent protein (BFP) that converts to green fluorescent protein (GFP) upon a C-to-T substitution was used as an assay to report directly on base editing activity within a cell. Using this assay, various base editing transfection parameters and delivery strategies were developed. Moreover, this assay was used in conjunction with flow cytometry to develop a transient reporter for editing enrichment (TREE) to efficiently purify base-edited cell populations. Compared to conventional cell enrichment strategies that employ reporters of transfection (RoT), TREE significantly improved the editing efficiency at multiple independent loci, with efficiencies approaching 80%. In addition, the BFP-to-GFP conversion assay was employed to develop base editor vector design for human pluripotent stem cells (hPSCs), a cell type that is resistant to genome editing and in which modification via base editors has not been previously reported. At last, using these optimized vectors in the context of TREE allowed for the highly efficient editing of hPSCs. It is envisioned that TREE is a readily adoptable method to facilitate base editing applications in synthetic biology, disease modeling, and regenerative medicine.

Materials and Methods

Plasmid construction: Unless otherwise noted, all molecular cloning polymerase chain reactions (PCR) were performed using Phusion® High-Fidelity DNA polymerase (New England Biolabs, Ipswich, MA, USA) using the using the manufacturer’s recommended protocols. All restriction enzyme (New England Biolabs) digests were performed

according to the manufacturer's instructions. Ligation reactions were performed with T4 DNA Ligase (New England Biolabs) according to the manufacturer's instructions. PCR primers and oligonucleotides were synthesized by Integrated DNA Technologies (Coralville, IA, USA). All PCR products and intermediate plasmid products were confirmed via Sanger sequencing (DNASU Sequencing Core Facility and Genewiz).

For construction of the pEF-BFP plasmid, we utilized PCR to add the H-66 and protospacer adjacent motif (PAM) site mutations into a GFP cassette (Addgene #11154). PCR products containing these mutations were digested with SapI/EcoRI and SapI/NotI and ligated into a EcoRI/NotI digested EF1 α expression vector (Addgene #11154).

For construction of the pDT-sgRNA vector, sgRNAs were synthesized as pairs of oligonucleotides (Table 4). Subsequently, 5' phosphates were added to each oligonucleotide pair by incubating 1 μ g oligonucleotide in 50 μ l reactions containing 1 $\times$ T4 DNA Ligase Buffer (New England Biolabs) and 10 units of T4 Polynucleotide Kinase at 37 $^{\circ}$ C. overnight. Oligonucleotides were then duplexed by heating the kinase reactions to 90 $^{\circ}$ C. on an aluminum heating block for 5 minutes followed by slowly returning the reaction to room temperature over 1 hour. Following duplexing, guides were cloned into a modified pSB1C3 vector containing a U6 promoter, inverted BbsI restriction enzyme digestion sites, and a *Streptococcus pyogenes* recognized sgRNA hairpin. For construction of pMT-sgRNA, pairs of sgRNAs (Table 4) were PCR amplified with primers adding EcoRI/SapI restriction enzyme digestion sites or SapI/XbaI restriction enzyme digestion sites. Purified PCR products were then digested with the respective restriction enzymes and ligated into EcoRI/XbaI digested pUC19 vector (Addgene #50005). The resultant vector contained pairs of sgRNA expression cassettes. To add additional sgRNA expression cassettes, pairs of sgRNAs were PCR amplified with primers that add HindIII/SapI or SapI/HindIII restriction enzyme digestion sites. These products were then digested with HindIII/SapI and ligated into HindIII digested and dephosphorylated pDT-sgRNA vector.

For insertion of the EF1 α promoter into the pCMV-BE4-Gam (Addgene #100806) and pCMV-AncBE4max (Addgene #112094), EF1 α was PCR amplified from an EF1 α expression vector (Addgene #11154) adding SpeI/NotI restriction enzyme digestion sites. After purification and digestion, these PCR products were ligated into SpeI/NotI digested and dephosphorylated pCMV-BE4-Gam or pCMVAncBE4max vectors.

Cell Culture: All media component were purchased from ThermoFisher Scientific (Waltham, MA, USA) unless indicated otherwise. HEK293 cells were cultured on poly-L-ornithine (4 μ g/ml; Sigma Aldrich, St Louis MO, USA) coated plates in the following media: 1 $\times$ high glucose Dulbecco's modified Eagle's medium (DMEM), 10% (v/v) fetal bovine serum, 1% (v/v) L-glutamine penicillin/streptomycin. Culture medium changed was every other day and cells were passaged with Accutase (ThermoFisher) every 5 days. HPSCs were cultured on 12-well tissue culture plates coated with MatrigelTM (BD Biosciences, San Jose, CA, USA) in Essential 8TM Medium (E8) (ThermoFisher). HPSCs were cultured in mTESR1 medium (STEM CELL Technologies). Culture medium was changed every day and cells were passaged with Accutase every 4-5 days. After passaging, the medium was supplemented with 5 μ M Rho kinase inhibitor (ROCKi; Y-27632 [BioGems, Westlake Village, CA, USA]) for 24 hours to aid in single cell survival.

Isolation of Episomal DNA: After 48 hours following transfection, HEK293 cells were dissociated from the tissue plates with Accutase, washed twice with phosphate-buffered saline and resuspended in RNase-A containing solution. Cells were then lysed via alkaline lysis and the resultant debris was precipitated via centrifugation at $1.2 \times 10^4 \times g$ for 10 min. Supernatant DNA was isolated by column DNA purification using the manufacture recommended protocol (Sigma-Aldrich: NA0160).

Generation of HEK293-BFP line: The HEK293T-BFP cell line was generated via homology independent targeted integration (HITI). Briefly, the BFP coding sequence was PCR amplified with primers adding EcoRI restriction enzyme digestion sites. The resultant PCR product was EcoRI digested, phosphorylated and ligated into an EcoRI/SmaI digested vector containing an EF1 α promoter, puromycin resistance cassette and HITI protospacer sequence (pEF-BFP-PuroR). The pEFBFP-PuroR vector was co-transfected in HEK293s with pX330 (Addgene #42230) and a custom sgRNA vector (pHSG(C1ORF228)-1C3) targeting the C1ORF228 locus. Transfections were conducted in a 24-well plate with 300 ng pX330, 400 ng pEF-BFP-PuroR, 50 ng sgRNA vector, 1.5 μ l Lipofectamine 3000 (ThermoFisher Scientific) and 1 μ l P3000 transfection reagent. Cells were passaged at 72 hours post-transfection into a single well of a 6-well plate and selected with 0.5 μ g/ml puromycin for 2 weeks.

Results

BFP-to-GFP Conversion Allows for Detection of Base-Editing Activity

To establish that BFP to GFP conversion could be used as the basis for an assay to detect genomic base editing, we utilized a BFP mutant that converts to a GFP upon a C-to-T nucleotide conversion (FIG. 1A). Briefly, this BFP mutant (BFP^{H66}) contains a histidine at the 66th amino acid position encoded by a 'CAC' codon. The C-to-T conversion of that codon to 'TAC' or 'TAT' will result in an amino acid change from a histidine to a tyrosine. In turn, this amino acid change will cause a shift of the emission spectra of the resultant protein generating a GFP variant (GFP^{T66}). Because the optimal nucleotide base editing window is typically 12-18 nt upstream from the PAM, we also placed a *S. pyogenes* Cas9 PAM 'NGG' in a position that would enable base editing to occur at the target 'CAC' codon. To verify the utility of this fluorescent protein to report on base editing activity, we cloned the BFP coding sequence into a vector with a human EF1 α promoter to drive expression (pEF-BFP; FIG. 1B). In addition, we designed a sgRNA vector [sg(BG)] that would direct the base editing machinery to the target 'CAC' codon resulting in a C-to-T conversion and the subsequent amino acid change of histidine to tyrosine at the 66th amino acid position (FIG. 1A). HEK293 cells were co-transfected with pEF-BFP, a base editing vector (pCMV-BE4-Gam) and sg(BG) or a control non-targeting sgRNA [sg(NT)]. Fluorescent microscopy (FIG. 1C) and flow cytometry (FIG. 1D) revealed that targeting pEF-BFP with sg(BG) resulted in the generation of BFP/GFP double positive cells. However, targeting pEF-BFP with sg(NT) did not result in the generation of any BFP/GFP positive cells. To confirm GFP expression was a consequence of direct editing of the target codon in pEF-BFP, we implemented a strategy to isolate and detect editing of episomal DNA after transfection (v 1E). Sanger sequencing of isolated pEF-BFP DNA established that editing had occurred at the target 'CAC' codon in pEF-BFP resulting in a change to 'TAC' or 'TAT' reflected

in the GFP emission (FIG. 1F). Overall, these results confirm that the GFP-to-BFP conversion corresponds to C-to-T conversion at targeted base editing sites.

Next, we wanted to establish that the BFP-to-GFP conversion would correlate with base-editing efficiency at a chromosomal locus. To that end, we employed a HEK293 cell line (herein referred to as HEK293-BFP) in which BFP^{H66} was stably integrated into a known genomic location (C1ORF228; FIG. 2A). We then used this line to enable the analysis of the efficiency of base editing genomic loci (FIG. 2B). To first assess plasmid-based base editing, we co-transfected pCMV-BE4-Gam and sg(BG) plasmid DNA in HEK293-BFP cells. Targeting with sg(BG), but not sg(NT), resulted in generation of detectable GFP+ cells, indicating successful base editing at the targeted genomic loci (FIG. 2C). Moreover, we were able to use this assay to systematically evaluate genomic base editing efficiencies using a range of pCMV-BE4-Gam plasmid amounts at varying ratios with the sg(BG) vector (FIG. 2D). This analysis revealed that base editing plasmid concentration and base editor to sgRNA ratios could enhance genomic base editing efficiencies approximately 2-fold. Because ribonucleoprotein (RNP) complex-based strategies have been previously shown as an attractive alternative to plasmid-based Cas9 genome engineering, we also utilized BFP-to-GFP conversion as an assay to optimize RNP-driven base editing. As such, we generated RNPs through the in vitro complexing of purified base editing protein with sg(BG) or sg(NT) (FIG. 2E). Our initial analysis revealed that RNP delivery using the same transfection reagent that was used for plasmid delivery of the base editor (i.e., LipofectamineTM 3000) did not result in substantial BFP-to-GFP conversion (FIG. 2F). In turn, we utilized BFP-to-GFP to evaluate various commercially available transfection reagents to optimize RNP delivery for base editing applications. From this analysis, we were able to determine that LipofectamineTM 2000 allowed for a >4-fold increase in genomic base editing efficiency compared to other commercially available reagents such as LipofectamineTM and CRISPRMAX (FIG. 2F). Despite this, RNP-driven delivery was about 4-fold less efficient in genomic base editing compared to plasmid delivery. Thus, for the remainder of this example we proceeded with plasmid delivery of base editors. Nonetheless, this collective data demonstrates that BFP-to-GFP conversion correlates to base editing efficiency at genomic loci. Moreover, this approach allows for the facile and systematic optimization of base editing in human cells using plasmid- and RNP-based approaches.

Development of Transient Reporter for Editing Enrichment (TREE) to Identify and Efficiently Isolate Base-Edited Cell Populations

Conventional base editing approaches that use reporters of transfection (herein abbreviated as RoT) only report on the efficiency of plasmid delivery to a cell but not directly on the efficiency of base editing within these cells. As such, it was hypothesized that we BFP-to-GFP conversion, which directly correlates to base editing activity within a cell, could be employed as a TREE to allow for the identification and enrichment of cells in which targeted genomic base editing had occurred. To facilitate this, we engineered a dual-targeting sgRNA (pDT-sgRNA) vector that contains both sg(BG) and a sgRNA for a genomic target site [sg(TS)] (FIG. 3A). Moreover, the pDT-sgRNA vector was designed to allow for the facile cloning of new target sites via BbsI restriction enzyme digestion and ligation of sg(TS) oligonucleotides (FIG. 3A). Accordingly, we designed pDT-sgRNA vectors with sequences targeting three genomic

locations (Sites 1-3). To utilize TREE for enrichment of cells that have been edited at specific loci, we co-transfected these pDT-sgRNA vectors with pEF-BFP and pCMV-BE4-Gam into HEK293 cells using the optimized base editing parameters identified using the BFP-to-GFP conversion assay (FIG. 3B). Flow cytometry was then used to isolate GFP-positive and GFP-negative cells. For comparison, we used a conventional RoT as a strategy to enrich for edited cell populations (FIG. 3C). Specifically, after co-transfecting HEK293 cells with pEF-GFP and sg(TS) plasmids, we used flow cytometry to sort for GFP-positive and -negative cell populations. Flow cytometry analysis of cells in which TREE was applied confirmed the presence of BFP and GFP-positive cell populations indicative of active base editing (FIG. 3D). Importantly, in these cell populations there was also a significant percentage of cells that were BFP-positive but GFP-negative, suggesting that isolating cell populations exclusively based upon a reporter of transfection would significantly limit the enrichment of edited cells. To confirm this, we performed Sanger sequencing of the targeted genomic sites in GFP-positive, GFP-negative and unsorted cell populations isolated from TREE and RoT approaches (FIG. 3E and FIG. 19). As expected, GFP-positive cells isolated using both TREE- and RoT-based strategies were enriched for edited cells when compared to GFP-negative and unsorted cell populations. We found that base editing efficiency at these three target loci in HEK293 cells using RoT-based approaches was similar to those reported previously (Table 11). Importantly, this analysis also revealed across all three targeted sites that GFP-positive cells isolated via TREE had a statistically significant higher frequency of base editing than GFP-positive cells isolated using traditional RoT approaches.

Because of the success of targeting these loci, it was investigated if TREE could be utilized to target additional genomic sites that display very low editing efficiency when traditional RoT approaches are applied. One such example is the APOE locus, a well-established risk factor associated with altered probability of sporadic Alzheimer's disease onset. Human APOE has three major isoforms, ApoE2, ApoE3 and ApoE4, which differ by two amino acid substitutions at positions 112 and 158 in exon 4-ApoE2 (Cys112, Cys158), ApoE3 (Cys112, Arg158), ApoE4 (Arg112, Arg158). Attempts to use base editing to convert ApoE3 to ApoE2 by targeting the APOE(R158) locus revealed undetectable levels of editing in unsorted cell populations despite similar transfection efficiencies when other genomic sites (Sites 1-3) were targeted (FIG. 20). In addition, our attempts to use RoT-based methods in HEK293 cells to convert ApoE3 to ApoE2 by targeting the APOE(R158) locus revealed very low levels of editing in GFP+ isolated cells (FIG. 20B), further establishing the APOE(R158) locus as recalcitrant to genomic editing. Then, TREE-based methods were used to edit this same loci in HEK293 cells by co-transfecting pEF-BFP, pCMV-BE4-Gam and pDT-sgRNA with a sg(TS) targeting the APOE(R158) locus. As expected, flow cytometry analysis demonstrated that the transfection efficiency when TREE was used to target the APOE(R158) locus was similar to when TREE was used to target other genomic sites (FIG. 20C). In addition, despite these similarities in transfection efficiencies, there was no detectable editing in the unsorted cell populations using TREE to target the APOE(R158) locus, thereby confirming the difficulty in editing this genomic location (FIG. 20D). However, unlike in GFP-positive isolated using RoT methods, GFP-positive cells purified using TREE methods displayed a high level of base editing at the APOE(R158) locus

(FIG. 20E). Together, these results demonstrate that TREE can not only provide for a higher level of enrichment of base-edited cell populations compared to conventional RoT strategies but also can allow for isolation of base-edited cells at genomic loci that were not previously achievable with traditional RoT approaches.

At last, we wanted to confirm that the fluorescent signal associated with cells isolated by TREE was transient. To that end, we measured the long-term fluorescence of GFP-positive cells purified after TREE-based editing (FIG. 21A). Notably, analysis of these cells by fluorescent microscopy (FIG. 21B) and flow cytometry (FIG. 21C) revealed no long-term detectable GFP signal, verifying that the TREE fluorescent output is indeed transient in nature. Multiplex Base-Editing Using TREE

It was further investigated whether TREE could be utilized in conjunction with multiplexed genome engineering strategies. To accomplish this, we generated a multi-targeted vector (pMT-sgRNA) that contains sg(BG) as well as sgRNA for genomic targets Sites 1-3 (FIG. 4A). In a similar manner to when TREE was employed to target a single locus, we utilized TREE to simultaneously target multiple genomic sites by co-transfecting HEK293 cells with pMT-sgRNA, pEF-BFP and pCMV-BE4-Gam. In parallel, we used a RoT-based approach by co-transfecting HEK293 cells with pMT-sgRNA, pEF-GFP and pCMV-BE4-Gam. After 48 h, GFP-negative and GFP-positive cells were isolated using flow cytometry (FIG. 22A). Along similar lines to single locus targeting, Sanger sequencing of the multiplex targeted genomic sites in GFP-positive cell populations isolated from TREE and RoT approaches revealed that TREE allowed for statistically significant higher frequency of base editing than RoT approaches (FIG. 4B and FIG. 22B). Importantly, this analysis revealed that there was no statistically significant difference in editing efficiency when TREE was used to target these sites individually or a multiplexed manner (FIG. 22C). Finally, we wanted to determine if TREE increased the likelihood of C-to-T conversions at off-target loci. Therefore, in GFP-positive cell populations isolated from TREE and RoT approaches we PCR-amplified and Sanger sequenced the top predicted off-target loci for the sgRNA sequences used for multiplexed editing. Overall, quantification of the Sanger chromatographs by EditR revealed no observable C-to-T conversions at these off-target loci in either GFP-positive cells isolated with TREE- or RoT-based strategies when compared to that of untransfected cells (FIG. 23).

Sanger sequencing that was performed on bulk sorted GFP-positive cells suggested that multiplex editing in conjunction with TREE could result in multiplexed editing in the same cell. To confirm that this indeed occurred, we again used our multi-targeting vector (pMT-sgRNA) in conjunction with TREE to simultaneously target genomic Sites 1-3 in HEK293 cells. We then sorted single GFP-positive cells into a 96-well plate. After expansion, Sanger sequencing of the multiplexed genomic sites was performed on a total of 40 clones. This analysis revealed that 36 out of the 40 clones had base editing at more than one genomic site (FIG. 4C). Remarkably, this analysis revealed that almost 80% of the isolated clones (31 out of 40) had biallelic conversions at all three genomic loci.

One of the caveats of all base-editing approaches, regardless if RoT- or TREE-based enrichment strategies are employed, is that base editors can potentially edit non-target Cs that are located in an 6 nt window (termed the editing window) within the protospacer. As a consequence, this could potentially limit the application of base editing

approaches in which conversion of non-target Cs result in a non-silent mutation or other phenotypic changes. To that end, we wanted to determine if any of our clones contained edits exclusively at the target C and not any other Cs within the editing window. Indeed, we identified a number of clones in which at genomic Site 2 and Site 3 modification only occurred at the target C (FIG. 24). Interestingly, we did not identify any clones in which at genomic Site 1 such exclusive modification of the target C occurred. We speculate that because another C occurs immediately adjacent to this target C, that such exclusive modification will require the use of recently published site-specific editors that allow for single nucleotide changes free from off-targeting conversions within the editing window.

TREE Allows for Highly Efficient Editing in Human Pluripotent Stem Cells (hPSCs)

Single base pair modification in hPSCs via CRISPR/Cas9-induced DSB followed by HDR suffers from low efficiencies. In addition, genomic modification of hPSCs using deaminase-based DNA base editor has yet to be reported. Therefore, we wanted to investigate if TREE could be utilized to efficiently edit specific loci in hPSCs. Hence, we co-transfected pEF-BFP and pCMV-BE4-Gam into hPSCs using a transfection reagent (Lipofectamine™ Stem) that had been previously used by others for the efficient delivery of Cas9-related plasmids to hPSCs. Surprisingly, we did not observe many GFP-positive cells in these cell populations (FIG. 5A and FIG. 25A). As such, we performed similar experiments in which we employed a more recently published, higher efficiency base editor, AncBE4max (herein referred to AncBE4). Briefly, AncBE4 is an improved version of BE4 that has been codon optimized for expression and contains an ancestral reconstructed deaminase to increase base editing efficiency at target loci. Nonetheless, similar to when BE4-Gam was utilized, we observed very few GFP-positive cells when AncBE4 was used (FIG. 25A). Because previous reports have suggested that the CMV promoter is inefficient for transgene expression in pluripotent stem cells, we replaced the promoter driving base editor expression with EF1a. When hPSCs were co-transfected with pEF-BE4-Gam or pEF-AncBE4 as well as pEF-BFP and sg(BG), a significant number of GFP-positive cells were observed (FIG. 5A). Using the pEF-AncBE4 vector, we also developed editing efficiency in hPSCs by using a range of base editor amount at varying ratio with sg(BG) (FIG. 25B). Similar to our experiments with HEK293 cells, this analysis revealed that base editing efficiencies were significantly affected by these parameters. Interestingly, the most optimal parameters in hPSCs differed from those identified in HEK293 cells (FIG. 2D) highlighting the utility of this assay to evaluate these variables. Using these base editing vector designs, we applied TREE to target a genomic loci in hPSCs by co-transfecting pEF-BE4-Gam/pEF-AncBE4, pEF-BFP and pMT-sgRNA (with a sg(TS) targeting site 1) (FIG. 5B). In turn, flow cytometry was used to isolate GFP-positive and -negative cell populations (FIG. 5C). Subsequently, Sanger sequencing was performed on the targeted genomic site in GFP-positive, GFP-negative and unsorted cell populations isolated from TREE and RoT approaches in which pEF-BE4-Gam and pEF-AncBE4 was used (FIG. 25C). This analysis demonstrated that GFP-positive hPSCs isolated via TREE had a statistically significant higher frequency of base editing than GFP-positive hPSCs isolated using traditional RoT approaches (FIG. 5D). In addition, TREE employed with the pEF-AncBE4 vector allowed for the efficient modification of the difficult to edit APOE(R158) locus (FIG. 7D-S7E).

Similar to our work with HEK293 cells, we wanted to confirm that the fluorescent output of TREE was transient in nature. In that regard, we measured the fluorescence of GFP-positive hPSCs isolated after TREE-based editing. Flow cytometry analysis revealed that after 2 weeks of culture there was no detectable GFP signal (FIG. 26), demonstrating that the fluorescent signal associated with hPSCs purified by TREE was transient.

Collectively, although this data demonstrates that TREE can be utilized for the efficient base editing of hPSCs, one of the caveats of all base editing approaches is the C-to-T conversion of non-target Cs within the editing window. Indeed, the Sanger sequencing analysis of GFP-positive populations isolated from TREE revealed editing of such non-target Cs when either Site 1 (FIG. 25C) or the APOE (R158)(FIG. 25E) locus was targeted in hPSCs. As such, to determine whether TREE allowed allelic outcomes in which targeting only occurred at the desired C, we performed NGS of PCR amplicons of Site 1 and APOE(R158) in GFP-positive cells purified using TREE. This analysis revealed at both these loci a very modest number of allelic outcomes in which base editing occurred exclusively at the target C, free from con-founding C-to-T conversions at other sites within the targeting window (FIG. 27). Instead, the most common editing outcome was one in which the majority of the Cs in the editing window were converted to Ts (FIG. 27). This suggests that for future applications which require a higher percentage of allelic outcomes where editing occurs only at the target C the use of recently published base editors that have a narrower editing window will be required. Nonetheless, this collective data demonstrates the broad utility of TREE to allow for the efficient editing in hPSCs.

Discussion

Since the first deaminase base editor was developed by Komor et al., multiple additional base-editing technologies have been rapidly developed with various endonucleases, deaminases, targeting windows and PAM specificities. Application of these emerging base editors to new cell types requires a slow, iterative process in which various base editing parameters are tested and editing efficiency is assessed through downstream sequencing methods. Additionally, as demonstrated herein, transfection efficiency does not precisely correlate with editing efficiency, so reporters of transfection do not provide accurate information about the efficacy of various base editing strategies. Here, it is described how BFP-to-GFP conversion and TREE can be utilized to rapidly optimize various factors that influence base editing efficiency, including base editor plasmid concentration and design as well as base editor to sgRNA ratios. In fact, these data show that these parameters are cell line-specific, demonstrating the advantage of TREE to allow for the high-throughput evaluation of base editing approaches. In the future, TREE may be used in the context of high-throughput screening to identify small molecules to further enhance base editing efficiency in a manner similar to that which has been previously achieved with CRISPR-mediated HDR approaches.

It has been shown that CRISPR/Cas9 genome engineering is compatible with a variety of delivery methods (e.g., lipid-mediated transfection, electroporation) and expression systems (e.g., plasmid DNA, Cas9-gRNA ribonucleo-protein complexes [RNP]), each with advantages and disadvantages that have been reviewed extensively elsewhere. In this example, we employed lipid-based delivery reagents that have been previously employed by others for the

CRISPR/Cas9-based editing of HEK293 cells (Lipofectamine 3000) and hPSCs (Lipofectamine Stem). Given TREE's ease of use and readily detectable fluorescent output we anticipate that TREE can be employed with whatever transfection method that is preferred by the end user. For instance, the data presented here demonstrated that the TREE base editing assay was compatible with both plasmid and RNP approaches. Although we observed that the overall genomic base editing efficiency of RNP-based expression was lower than that of lipid-based expression, we provide proof-of-principle that TREE can be employed in future applications where the advantages of RNPs are desirable.

One potential limitation of the use of the plasmid DNA expression systems in the context of TREE approaches is random integration of all or part of the plasmid DNA into the genome of targeted cells. It should be noted that it has been reported by others that the stable integration of circular plasmid DNA into the host genome is infrequent, especially for cells such as hPSCs where it has been reported on the order of 1 per 1×10^5 cells. Indeed, as it relates to potential integration of the pEF-BFP plasmid, we demonstrate that the fluorescent output of TREE is transient in both HEK293 cells and hPSCs, suggesting that this plasmid does not integrate into the genome. As it relates to the integration of the base editing and sgRNA plasmids, it has been shown by others in CRISPR/Cas9 genome engineering that the Cas9 and sgRNA plasmids can be integrated at on- and off-target sites. However, we speculate that because base editors do not introduce DSBs the integration of these plasmids into the genome would be infrequent. In fact, we did not observe any integration of these plasmids when Sanger sequencing or NGS was performed at the on- or off-target sites. Moving forward, undesirable insertions of plasmid DNA sequences at target sites can be detected using PCR-based methods followed by Sanger sequencing or NGS of the resultant amplicons. On the other hand, similar insertions at off-target or random genomic sites are difficult to detect and will require the use of more comprehensive techniques such as whole genome sequencing.

Human cell models are critical for elucidating the mechanisms of disease progression as well as identifying and testing potential therapeutic interventions. Because a high percentage of human diseases are due to single nucleotide poly-morphisms (SNPs), base editors can allow for the precise engineering of in vitro models of human disease. Here we provide proof-of-principle that TREE can be employed to edit disease-relevant loci. Specifically, we demonstrate that TREE enables for the enrichment of cells that had been edited at the APOE(R158) locus, a gene associated with altered risk of Alzheimer's disease onset. Notably, conventional RoT-based methods did not allow for significant enrichment of edited cells at this same refractory locus. In addition, because many human diseases are multi-genetic disorders that are a result of complex gene interactions, we also investigated the ability of TREE to be utilized in multiplexed genome engineering applications. By using a multi-targeted vector, we demonstrated that compared to RoT-based methods TREE resulted in a significantly higher level of cells enriched for simultaneous editing at multiple independent loci. In fact, we demonstrated that through analysis of single cell clones that 90% of the clones had simultaneous base editing at more than one genomic site and almost 80% of the clones had biallelic conversions at all three targeted loci. In this vein, TREE provides a highly efficient method for generating cell-based models of multi-genetic diseases.

Many immortalized cell lines, such as HEK293s, are aneuploid with unknown mutations and dosage at key disease-relevant genes. Alternatively, hPSCs, which have a normal euploid karyotype and the potential to differentiate into all cell types of the mature adult body, represent an attractive alternative to immortalized cell lines for disease modeling and drug screening applications. In particular, the ability to use gene editing technologies to generate isogenic hPSC lines that differ only with respect to disease mutations has great potential as it relates to precisely defining genotype to phenotype relationships. The RNA-guided CRISPR-Cas9 system has the potential to allow for precise genetic modifications in hPSCs through the introduction of site-specific DSBs. Although previous reports demonstrate that introduction of DSB via CRISPR/Cas9 significantly improves the ability to obtain knock out cell lines from hPSCs by the NHEJ pathway, single base modification using CRISPR/Cas9-induced DSB followed by HDR is extremely inefficient (1-2% of sequenced colonies in which one allele is targeted and <1% where both alleles are targeted. Recently, it has been reported that co-delivery of Cas9, sgRNA, and a puromycin selection cassette followed by transient puromycin selection can increase the HDR-mediated genome engineering in hPSCs. However, these strategies rely on the introduction of DSBs, which in pluripotent stem cells can lead to large deletions and complex chromosomal rearrangements, significant cytotoxicity and increased acquisition of p53 mutations. In addition, it has been shown that the use of antibiotic selection, even in a transient manner, may lead to the selection of hPSCs, with chromosomal abnormalities. Yet, to our knowledge, base editors, which do not have these same limitations as CRISPR/Cas9-induced DSB followed by HDR, have not previously been used with hPSCs. In fact, our initial attempts to apply base editors in the context of both RoT- and TREE-based approaches with hPSCs did not allow for observable modification of target loci. Instead, by replacing the standard CMV promoter in the base editing plasmids with an EF1a promoter, we were able to achieve modification of genomic sites using both RoT- and TREE-centered methods. However, TREE allowed for significantly higher enrichment of edited hPSCs when compared to RoT isolation strategies. We contend that the use of TREE with hPSCs will significantly advance the use of these cells in disease modeling, drug screening, and cell-based therapies.

Despite their tremendous potential in a variety of downstream applications, base editing approaches have a few of caveats that should be noted, regardless of whether RoT- or TREE-based enrichment strategies are employed. First, as is the case with all Cas9-directed genome editing approaches, is the potential for genome modification at off-target loci. In this work, GFP-positive cells isolated via TREE did not display untargeted C-to-T conversions at the off-target genomic loci examined. Recently, it has been reported that base editors can induce site-specific inosine formation on RNA. Accordingly, in the future, the effect of TREE-based approaches on unwanted RNA modifications should be examined. Another limitation of base editing methods is modification of additional C nucleotides that are in close proximity to the target C. In fact, some base editors can cause C-to-T conversions at any Cs in up to a 9-nt window within the protospacer. Such C-to-T modifications could be especially problematic if they result in amino acid alterations during translation, induce epigenetic changes or cause other phenotypic changes in targeted cells. To that end, through clonal isolation and next generation sequencing (NGS) analysis we identified that such exclusive modifications of the target C were achieved in both edited HEK293

cells or hPSCs that were enriched using TREE-based methods. It should be noted, though, that at genomic Site-1, where a C lies adjacent to the target C, allelic outcomes in which modification only occurred at the target C were rare events. Moving forward, modified base editors that have a narrow editing window could be easily employed with TREE to target such genomic loci that contain multiple Cs in close proximity to the target C.

In summary, these data demonstrate that TREE allows not only for the optimization of base editing strategies in the context of a variety of cell types and genomic locations but also the enrichment of cell populations to be utilized in variety of downstream applications. In particular, with the rate at which the genome editing field has been progressing over the past few years, TREE is a readily adoptable method that will expedite and improve tractability of single-nucleotide genome engineering methods.

Example 2—Producing Base-Edited Isogenic Cell Lines Using a Transient Reporter for Editing Enrichment (BIG-TREE)

Current CRISPR-targeted single-nucleotide modifications and subsequent isogenic cell line generation in human pluripotent stem cells (hPSCs) require the introduction of deleterious double-stranded DNA breaks followed by inefficient homology-directed repair (HDR). This section describes the development of Cas9 deaminase base-editing technologies to co-target genomic loci and an episomal reporter to enable single-nucleotide genomic changes in hPSCs without HDR. Using this method, a base-edited isogenic hPSC line was generated using a transient reporter for editing enrichment (BIG-TREE) which allows for single-nucleotide editing efficiencies of >80% across multiple hPSC lines. Also described herein is use of BIG-TREE for efficient generation of loss-of-function hPSC lines via introduction of premature stop codons. Finally, BIG-TREE achieves efficient multiplex editing of hPSCs at several independent loci. These methods allow for the precise and efficient base editing of hPSCs for use in developmental biology, disease modeling, drug screening, and cell-based therapies.

Materials and Methods

Cells and Culture Conditions: Cell lines, media compositions, and conditions for culture of hPSC and HEK293 are listed in the Supplemental Experimental Procedures (End of Example 2).

Plasmid Construction: All plasmids were constructed using conventional restriction enzyme-based molecular cloning techniques. For construction of the sgRNA plasmids, the sgRNA sequences listed in Table 12 were used. Additional details for molecular cloning and plasmid construction are provided in the Supplemental Experimental Procedures (End of Example 2).

```
sgRNA Sequences in TREE: Guide and Cas9 Handle
>sg (BG)
                                     (SEQ ID NO: 3)
GACCCACGGCGTGCAGTGCTTGTGTTTAGAGCTAGAAATAGCAAGTTAAAA
TAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGAGTCGGTGCTTT
TTTGTTTT
```

37

-continued

>sg (NT) (SEQ ID NO: 4)
GGGTCTTCGAGAAGACCTGTTTGTAGAGCTAGAAATAGCAAGTTAAATAA
GGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTTTT
GTTTT

>sg (Site-1) (SEQ ID NO: 5)
GGCCAGACTGAGCACGTGAGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (Site-2) (SEQ ID NO: 6)
GAACACAAAGCATAGACTGCGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (Site-3) (SEQ ID NO: 7)
GGCACTGCGGCTGGAGGTGGGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (APOE R158) (SEQ ID NO: 8)
GAAGCGCTGGCAGTGTACCGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (C10RF228) (SEQ ID NO: 9)
GTGCTGTTAGCACCTGGAAAGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
TAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTT
TTGTTTT

>sg (APOE Q39X) (SEQ ID NO: 10)
GTGGCAGAGCGGCCAGCGCTGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (mCh1) (SEQ ID NO: 11)
GCACCAGACCGCCAGCTGAGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
TAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTT
TTGTTTT

>sg (mCh2) (SEQ ID NO: 12)
GACCCAGACTCCTCCCTGCGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
AAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTTT
TTGTTTT

>sg (mCh3) (SEQ ID NO: 13)
GCAAGCAGAGGCTGAAGCTGAGTTTGTAGAGCTAGAAATAGCAAGTTAAAT
TAAGGCTAGTCCGTTATCAACTTGAAAAAGTGGCACCAGTCGGTGCTTT
TTGTTTT

38

hPSC Base Editing, Clonal Isolation, and Characterization: Methods for transfection of hPSCs, clonal isolation, and characterization via tri-lineage differentiation are described in the Supplemental Experimental Procedures (End of Example 2).

Genotyping and Sequence Analysis at Off- and On-Target Analysis: Genomic DNA was prepared from expanded clones using the DNeasy kit (QIAGEN). PCR was performed with the primers listed in Table 5 using the methods described in the Supplemental Experimental Procedures (End of Example 2).

Karyotype Analysis: For each cell line, cytogenetic analysis was performed (Cell Line Genetics) on 20 metaphase cells using standard protocols for G-banding.

Immunofluorescence: Detailed protocols for immunofluorescence and antibodies used are provided in the Supplemental Experimental Procedures (End of Example 2).

HEK293 Transfections: Methods for transfection of HEK293s and Sanger sequencing of resultant populations are described in the Supplemental Experimental Procedures (End of Example 2).

Fluorescence Microscopy: Fluorescent imaging was performed on a Nikon Ti-Eclipse inverted microscope using the filters and acquisition settings described in the Supplemental Experimental Procedures (End of Example 2).

Flow Cytometry: Cells were dissociated with Accutase for 10 minutes at 37° C., triturated, and passed through a 40 mm cell strainer. Cells were then washed twice with flow cytometry buffer (BD Biosciences) and resuspended at a maximum concentration of 5×10⁶ cells per 100 mL. Flow cytometry analysis was performed on an Attune NxT (Thermo Fisher Scientific). Flow cytometry files were analyzed using FlowJo (FlowJo LLC, Ashland, OR, USA).

Apolipoprotein E ELISA: Cells were seeded in a 6-well plate at a density of 3×10⁵ cells per well. Medium was changed every 24 hours (h). On day 3, 24-h conditioned medium was collected, and ApoE levels in the medium were measured with the Human APOE (AD2) ELISA Kit (Thermo Scientific).

Statistical Analysis: Unless otherwise noted, all data are displayed as means±SD.

Results

Highly Efficient Generation of Clonal Isogenic hPSC Lines Using BIG-TREE

It was previously demonstrated that the efficiency at which a base editor is delivered to a cell does not precisely correlate with editing efficiency at a genomic locus. To overcome this limitation, an assay was developed, termed transient reporter for editing enrichment (TREE) (Standage-Beier et al., 2019). TREE utilizes a BFP variant that converts to a GFP upon a C-to-T nucleotide change (Standage-Beier et al., 2019). More specifically, this BFP mutant contains a histidine at the 66th amino acid position encoded by a “CAC” codon. The C-to-T conversion of that codon to a “TAC” or “TAT” will cause an amino acid change from a histidine to a tyrosine as well as a shift in the emission spectra of the modified protein resulting in a GFP variant. Thus, co-transfection of cells with this BFP construct (pEF-BFP), a base editor (pEF-AncBE4max), and a single guide RNA (sgRNA) targeting the “CAC” codon, sg(BG), will result in a BFP-to-GFP conversion in which the base editor machinery is present and actively functioning. In addition, we found that this BFP-to-GFP conversion was highly predictive of the likelihood of base editing at genomic loci

within the same cell that had been transfected with a sgRNA for a genomic target site, sg(TS).

This section describes efforts to extend this work to develop a rapid and efficient resource that uses TREE as the basis for the generation of clonal isogenic hPSC lines, termed base-edited isogenic hPSC line generation using a transient reporter for editing enrichment (BIG-TREE) (FIG. 6A). As proof-of-principle, we aimed to edit the APOE locus, a risk factor associated with altered probability of sporadic Alzheimer disease (AD) onset (Hauser and Ryan, 2013). Human APOE has three common isoforms that differ from each other by two amino acids at position 112 and 158 (APOE2=Cys112, Cys158; APOE3=Cys112, Arg158; APOE4=Arg112, Arg158). To this end, we transfected a non-demented control hPSC line (herein referred to as hPSC line 1) that has an APOE3/E3 genotype with pEF-BFP, pEF-AncBE4max, and a dual-targeting sgRNA (pDT-sgRNA) vector that contains both sg(BG) and a sgRNA for the APOE(R158) locus (FIG. 6B, top). Consequently, successful targeting of the APOE(R158) locus would result in C-to-T conversion that would cause a change from an APOE3 genotype (R158) to an APOE2 genotype (C158) (FIG. 6C). At 48 hours post transfection, fluorescent activated cell sorting (FACS) was used to sort single GFP-positive cells into 96-well plates. Clonal lines were then passaged and expanded over the course of 18 days prior to detailed analysis. First, genomic DNA was isolated from ten clones and the target region of the APOE locus, APOE (R158), was subject to Sanger sequencing after PCR amplification (FIG. 6C). Remarkably, this analysis revealed that 90% of the clones isolated had been edited, with seven of the clones having a homozygous and two of the clones having a heterozygous edit at the APOE(R158) locus (FIG. 6D). For comparison, we used a more conventional reporter of transfection (RoT) approach in which this same hPSC line was transfected with a plasmid in which a GFP and the AncBE4max base editor are driven by the same promoter, connected by a P2A post-translational self-cleavage peptide tag (pEFAncBE4max-P2A-GFP), as well as the same sgRNA for the APOE locus (FIG. 6B, bottom). In a manner analogous to that described for the BIG-TREE-based approach, single GFP-positive cells were then sorted into 96-well plates, expanded, and subject to Sanger sequencing. Analysis of ten clonal lines revealed this traditional RoT-based approach was significantly less efficient with only a single clone displaying a heterozygous edit at the target APOE(R158) locus (FIG. 6D). Given the large variability that exists between individual hPSC lines (Ortmann and Vallier, 2017), we wanted to determine the robustness of BIG-TREE to efficiently generate isogenic pairs in other independent hPSC lines. In this vein, we employed BIG-TREE to target the APOE (R158 locus) in two hPSC lines derived from patients with familial AD (FAD) (herein referred to hPSC line 2 and hPSC 3). Analysis of single cell clones by Sanger sequencing (FIG. 6C) revealed that across all three hPSC lines tested, over 80% (33/41 clones examined) had an edit at the APOE(R158) locus, and greater than 50% of those edits were homozygous in nature (FIG. 6E). Importantly, we did not observe the presence of indels at the target site in any of clones examined. Finally, one of the limitations of base editor techniques, regardless if BIG-TREE strategies are employed, is that base editors can induce changes in the protospacer at a C other than the target C within the editing window-termed bystander editing (FIG. 21A). Indeed, with respect to generating isogenic lines at the APOE(R158) locus, editing at these bystander Cs was a common occurrence (FIG. 21B). In fact, only one of the

clones analyzed (line 2, clone 5) had a heterozygous edit exclusively at the target C and no other Cs within the editing window. However, it should be noted that these bystander edits did not alter the amino acid sequence.

We performed detailed phenotypic analysis on representative biallelic edited clones from each hPSC line. Overall, these clones had a normal euploid karyotype (FIG. 6F), characteristic hPSC morphology (FIG. 1G), high expression of key pluripotency markers (FIG. 1H), and demonstrated tri-lineage differentiation potential (FIG. 6I). In addition, off-target analysis was performed at the top predicted sites for sg(BG) as well as the sgRNA used to target the APOE (R158) locus. At all of the off-target sites analyzed, we did not observe any C-to-T conversions at these off-target loci (FIG. 29). Furthermore, indels were not identified at any of the off-target sites in clones analyzed. Finally, Sanger sequencing revealed that the AD-related mutations in the hPSC clones derived from the FAD lines were retained in the edited clones (FIG. 30). Taken together, this analysis reveals that TREE can be employed for the highly efficient generation of isogenic hPSCs across multiple independent cell lines.

BIG-TREE can be Utilized for the Engineering of Gene Knockout hPSC Lines

To date, engineering of hPSC loss-of-function lines using CRISPR-based approaches has involved the generation of Cas9-mediated DSBs followed by non-homologous end joining (NHEJ), which typically results in a frameshift mutation and introduction of a downstream premature stop codon. Because of the aforementioned caveats associated with such DSB-driven approaches, we wanted to determine if BIG-TREE could be utilized to generate gene knockout hPSC lines without the introduction of DSBs. Because base editors have not been utilized previously to generate loss-of-function in hPSCs, we first wanted to establish this proof-of-principle in HEK293 cells. First, to validate base editor targeted introduction of premature stop codons, we designed a series of sgRNAs targeting an mCherry cassette in an HEK293T line, which would lead to conversion of a "CAG" codon encoding for glutamine to a "TAG" stop codon (FIG. 31A). We observed loss of mCherry expression via fluorescent microscopy and flow cytometry when targeting with sgRNAs (FIGS. 31B and 31C). In addition, we confirmed the targeted addition of stop codons by Sanger sequencing (FIG. 31D). Finally, this analysis revealed that loss of mCherry fluorescent signal was a direct consequence of introduction of a premature stop codon introduced into the genomically integrated mCherry cassette (FIG. 31E). Next, we sought to employ BIG-TREE to introduce premature stop codons in hPSCs at a disease relevant locus. To this end, we transfected hPSC line 1 with pEF-BFP, pEF-AncBE4max, and a dual-targeting sgRNA (pDT-sgRNA) vector that contained both sg(BG) and a sgRNA for the glutamine residue at amino acid position 39 in exon 3 of the APOE locus. Successful targeting would result in conversion of the glutamine encoding "CAA" codon to a premature "TAA" stop codon (FIG. 7A). Similar to as previously described, we isolated clonal cell lines established from single GFP-positive sorted cells. Analysis of these clones by Sanger sequencing (FIG. 7B) revealed that more than 80% of the clones had a stop codon introduced at the target site with greater than 50% of the edited clones displaying a biallelic modification (FIG. 7C). Importantly, none of the clones analyzed had indels at the same target site. Lastly, to demonstrate that introduction of a premature stop codon in exon 3 results in functional loss of APOE, we measured the amount of APOE in the conditioned media secreted by

unedited and edited cells using ELISA. Compared with the unedited wild-type (Q39/Q39) cells that secreted robust amounts of APOE, cells in which a premature stop codon had been introduced into both alleles (X39/X39) did not secrete any detectable levels of APOE (FIG. 7D). Collectively, these data show that BIG-TREE enables efficient generation of loss-of-function hPSC lines through the introduction of premature stop codons.

BIG-TREE Enables High-Frequency, Multiplex Base Editing in hPSCs

Finally, we wanted to determine if BIG-TREE could be utilized with multiplexed genome modification methods to establish hPSC lines that had been simultaneously edited at multiple genomic locations. Accordingly, a multi-targeting vector (pMT-sgRNA) that contains sg(BG) as well as sgRNAs for three independent genomic target sites (FIG. 8A) was used. Analogous to when BIG-TREE was used to target a single genomic location, TREE was employed to simultaneously target multiple loci by co-transfecting hPSC line 1 with pMT-sgRNA, pEF-BFP, and pEF-AncBE4-max. Sanger sequencing was then performed on the multiplex targeted genomic sites in clonal hPSC lines derived from single GFP-positive cells (FIG. 8B). Along similar lines to when BIG-TREE was used to target a single genomic locus, Sanger sequencing revealed that more than 80% of clones had been targeted at all three sites with all clones displaying biallelic edits (FIG. 8C). Moreover, indels were not identified in any of the clones across all three target sites. Lastly, examination of potential bystander edits within the editing window (FIG. 28A) revealed a number of clones in which at genomic site 2 and site 3 modification only occurred at the target C and not any other Cs within the editing window (FIG. 28C). Specifically, of the ten clones that had homozygous edits at the target C at all three sites, two clones were free from bystander edits at both sites 2 and 3 (clones 1 and 2) and five clones were free from bystander edits at site 3 only (clones 3-7). However, it should be noted that we did not identify any clones in which at genomic site 1 such exclusive modification of the target C occurred. We speculate that because another C occurs immediately adjacent to this target C, that such exclusive modification is likely a rare event that will require site-specific base editors that allow for single-nucleotide changes free from bystander editing at adjacent nucleotides.

Discussion

In summary, we establish that BIG-TREE is a fast and efficient protocol for the generation of clonal isogenic hPSC lines with homozygous and heterozygous single base pair edits. Because the number of diseases that are a consequence of single point mutations, as well as the growing number of genomic variants of uncertain significance that have been identified through large-scale sequencing efforts, the ability to rapidly engineer isogenic hPSC lines will have a significant impact on the establishment of in vitro models to assess pathogenic risk and dissect disease-causing mechanisms. In addition, in this example, we demonstrate that BIG-TREE can be employed to generate effective loss-of-function cell lines through the introduction of premature stop codons. Currently, most CRISPR/Cas9-based approaches to generate gene knockouts involve the introduction of deleterious DSBs followed by NHEJ-mediated repair that results in frameshift and loss of gene function. As we describe in this example, the ability to rapidly generate gene knockouts without the need for DSBs will have important implications for the use of hPSCs to elucidate the function of specific

genes in development and disease. Lastly, we establish that BIG-TREE can allow for the generation of clonal hPSC lines that have been simultaneously edited at multiple independent loci, an important consideration given that many diseases are polygenetic in nature. By comparison, conventional CRISPR/Cas9-based approaches are too inefficient in hPSCs to employ multiplexing editing strategies.

Since the first base editors were engineered, numerous additional base editors with targeting windows, editing efficiencies, PAM specificities, and deaminases have been generated. In the context of BIG-TREE, we employed AncBE4max, which displays a relatively high editing efficiency with low off-target activity. However, one of the limitations of AncBE4max is that it can induce C-to-T conversions at bystander Cs within the editing window. Although bystander editing was a common occurrence in our clonal populations, we did observe clones with exclusive modifications of the target C. More specifically, when generating isogenic lines edited at the APOE(R158) locus, we only isolated one clone that had a monoallelic edit exclusively at the target C. Nonetheless, all of the bystander edits that we observed at the APOE(R158) locus did not impact the amino acid sequence, mitigating the impact on the downstream application of these hPSC lines. With regard to the multiplex editing, we did observe several clones that were free from bystander edits at genomic sites 2 and 3. However, at genomic site 1, where a C is present in the base pair position directly next to the target C, we did not isolate any clones where modification only occurred at the target C. In the future, given the ease of use, we anticipate that utilizing BIG-TREE with these other base editor variants with a narrow editing window will be easily achieved. In this regard, the end-user can select to employ such base editors with a more stringent editing window if editing at a bystander C is not tolerable (e.g., results in changes in the amino acid coding sequence).

In general, there are several enabling aspects to the methods presented in the example that will allow for the facile adoption by a broad set of researchers. First, the high editing frequencies do not require the screening of large numbers of clones to identify those with the desired modification. Moreover, we demonstrate that BIG-TREE is robust, as it allows for the efficient editing of multiple loci and across several independent hPSC lines. Because of these efficiencies, clonal lines can be identified, expanded, and characterized in the course of a few weeks. Along similar lines, the high efficiency of BIG-TREE allows for the biallelic or multiplexed targeting without the need for sequential re-targeting. In addition, BIG-TREE is compatible with off-the-shelf chemical transfection reagents and does not require the cloning of complex viral constructs or the use of specialized cell transfection systems. In fact, all sgRNA vectors were designed to allow for the facile cloning of new target sites via BbsI restriction enzyme digestion and ligation of oligonucleotides that target the desired genomic sequence. Lastly, BIG-TREE offers the flexibility to be used in conjunction with other base editor variants that have altered PAM specificities and editing windows. For instance, the PAM sequence and edit distance can be modified to match the editing specificity and window of the new base editor. Such modifications are straightforward to achieve with the BFP vector using TREE, or the stop codon between RFP and GFP using BIG-TREE. In this manner, BIG-TREE is a readily adoptable method that will enhance and accelerate the use of base-editing approaches in hPSCs.

Supplemental Experimental Procedures

Human iPSC and HEK293 culture. HPSCs were maintained in mTeSR1 medium (Stemcell Technologies) on

feeder-free Matrigel (Corning)-coated plates. Subculture was performed every 3 days using Accutase (Life Technologies) in mTeSR1 medium supplemented with 5 μ M Y-27632 (Tocris). Control and AD-patient hPSCs were generated from dermal fibroblasts as previously described (Park et al., 2008).

TABLE 3

HPSC lines described in the example			
Cell Line	Disease Status	MMSE	Mutation
hPSC Line 1	Non-demented control	30	n/a
hPSC Line 2	Familial AD	n/a	APP V717I
hPSC Line 3	Familial AD	n/a	PSEN1 A246E

mCherry expressing HEK293 line was generated using lentiviral integration of a constitutively expressing mCherry transgene as previously described (Standage-Beier et al., 2019). HEK293 cells were cultured on poly-L-ornithine (4 μ g/mL; Sigma Aldrich, St. Louis MO, USA) coated plates in the following media: 1 $\times$ high glucose DMEM, 10% (v/v) fetal bovine serum, 1% (v/v) L-glutamine penicillin/streptomycin. Culture medium was changed every other day and cells were passaged with Accutase every 5 days.

Plasmid construction. Unless otherwise noted, for all molecular cloning PCRs were performed using Phusion High-Fidelity DNA polymerase (New England Biolabs, Ipswich, MA, USA) using the manufacturer's recommended protocols. All restriction enzyme (New England Biolabs) digests were performed according to the manufacturer's instructions. Ligation reactions were performed with T4 DNA ligase (New England Biolabs) according to the manufacturer's instructions. PCR primers and oligonucleotides were synthesized by Integrated DNA Technologies (Coralville, IA, USA). All PCR products and intermediate plasmid products were confirmed via Sanger sequencing (DNASU Sequencing Core Facility). Complete plasmid sequences will be made available upon request.

For construction of the pEF-BFP plasmid, we utilized PCR to add the H-66 and PAM site mutations into a GFP cassette (Addgene #11154). PCR products containing these mutations were digested with SapI/EcoRI and SapI/NotI and ligated into an EcoRI/NotI digested EF1 α expression vector (Addgene #11154).

For construction of the pDT-sgRNA vector, sgRNAs were synthesized as pairs of oligonucleotides (Table 12). Subsequently, 5' phosphates were added to each oligonucleotide pair by incubating 1 μ g oligo nucleotide in 50 μ L reactions containing 1 $\times$ T4 DNA Ligase Buffer (New England Biolabs) and 10 units of T4 Polynucleotide Kinase at 37° C. overnight. Oligonucleotides were then duplexed by heating the kinase reactions to 90° C. on an aluminum heating block for 5 minutes followed by slowly returning the reaction to room temperature over 1 hour. Following duplexing, guides were cloned into a modified pSB1C3 vector containing a U6 promoter, inverted BbsI restriction enzyme digestion sites, and an *S. pyogenes* sgRNA hairpin. For construction of pMT-sgRNA, pairs of sgRNAs (Table 12) were PCR amplified with primers adding EcoRI/SapI restriction enzyme digestion sites or SapI/XbaI restriction enzyme digestion sites. Purified PCR products were then digested with the respective restriction enzymes and ligated into EcoRI/XbaI digested pUC19 vector (Addgene #50005). The resultant vector contained pairs of sgRNA expression cassettes. To add additional sgRNA expression cassettes, pairs of sgRNAs were PCR amplified with primers that add HindIII/SapI or

SapI/HindIII restriction enzyme digestion sites. These products were then digested with HindIII/SapI and ligated into HindIII digested and dephosphorylated pDT-sgRNA vector.

For insertion of the EF1 α promoter into pCMV-AncBE4max (Addgene #112094), EF1 α was PCR amplified from an EF1 α expression vector (Addgene #11154) adding SpeI/NotI restriction enzyme digestion sites. After purification and digestion, these PCR products were ligated into SpeI/NotI digested and dephosphorylated pCMV-AncBE4max vector.

hPSC base editing and clonal isolation. hPSCs were passaged onto Matrigel-coated 12-well plates with 5 μ M Y-27632. Media was changed, and transfection were performed 24 hours after passage. 900 ng base editor (pEF1 α -AncBE4max), 300 ng sgRNA, and 300 ng pEF1 α -BFP was transfected per well using 4 μ L Lipofectamine Stem transfection reagent (Life Technologies). Media was changed 24 hours post-transfection. Cells were dissociated using Accutase 48 hours post-transfection and passed through a 0.45 μ m filter. Single GFP-positive hPSCs were FACS sorted into 96-well Matrigel coated plates in mTeSR1 supplemented with CloneR (Stemcell Technologies), plates were immediately centrifuged at 100 $\times$ g for 1 minute and incubated at 37° C. Media was changed 48 hours post-sort with fresh mTeSR1 supplemented with CloneR. 96 hours post-sort, media was changed to mTeSR1 without supplement and clonal hPSC colonies were expanded with fresh media changes daily until ready for subculture.

Genotyping and Sequence Analysis.

Clones were amplified with the primers listed in Table 13 to determine genotype following base editing. Genomic DNA was prepared from expanded clones using the DNeasy kit (Qiagen) and PCR products were generated with Phusion High-Fidelity Polymerase (New England Biolabs). Amplicons were purified using the QIAquick PCR purification kit (Qiagen) according to manufacturer's instructions prior to Sanger sequencing (Genewiz). For multiplex clones, hPSCs were directly added to a 50 μ L master mix consisting of 1 $\times$ Phire Hot Start II DNA Polymerase (ThermoFisher), 1 μ M forward primer, and 1 μ M reverse primer. PCR was performed using the following conditions: 98° C. for 5 minutes, followed by 40 cycles at 99° C. for 5 seconds, 56° C. for 5 seconds, and 72° C. for 20 seconds, followed by a final 5 min 72° C. extension. All products sizes were confirmed on a 1% agarose gel prior to Sanger sequencing.

HEK293 transfections. HEK293 cells stably expressing mCherry were transfected in 24 well tissue culture plates at 40% confluence with the following reagents per well: 300 ng pEF1 α -AncBE4max, 100 ng sgRNA vector or sg(NT),

0.75 μ L Lipofectamine 3000 Transfection Reagent (ThermoFisher), and 1 μ L P3000 reagent (Thermo Fisher). Flow cytometry was performed at 7 days post-transfection to evaluate loss of mCherry expression. Genomic DNA was isolated and mCherry was PCR amplified before Sanger sequencing to determine editing efficiency.

Immunofluorescence. Cultures were gently washed twice with PBS prior to fixation. Cultures were then fixed for 15 min at room temperature (RT) with BD Cytofix Fixation Buffer (BD Biosciences). The cultures were then washed twice with PBS and permeabilized with BD Phosflow Perm Buffer III (BD Biosciences) for 30 min at 40° C. Cultures were then washed twice with PBS. Primary antibodies were incubated overnight at 40° C and then washed twice with PBS at room temperature. Secondary antibodies were incubated at RT for 1 hr. Nucleic acids were stained for DNA with Hoechst 33342 (2 μ g/mL; Life Technologies) for 10 min at RT and then washed twice with PBS. Antibodies used are as

follows at the following concentrations: NANOG (ThermoFisher Scientific; Cat #PA1-097, RRID:AB_2539867; 1:500), OCT4 (ThermoFisher Scientific; Cat #PA5-27438, RRID:AB_2544914; 1:500), SOX2

(ThermoFisher Scientific; Cat #PA1-094, RRID: AB_2539862; 1:500), AFP (Santa Cruz Biotechnology; Cat #sc-15375, RRID:AB_2223935; 1:50), SMA (Santa Cruz Biotechnology; Cat #sc-53015, RRID:AB_628683; 1:50), TUJ1

(Fitzgerald; Cat #10R-T136A, RRID:AB_1289248; 1:1000), Alexa 488 donkey anti-mouse (ThermoFisher Scientific; Cat #A-21206, RRID:AB_2535792; 1:500), and Alexa 488 donkey anti-rabbit (ThermoFisher Scientific; Cat #A-21202, RRID:AB_141607; 1:500).

Tri-lineage differentiation of edited hPSCs. hPSCs were harvested using Accutase and plated on ultra-low attachment plates in mTeSR1 medium. The following day, media was changed to differentiation medium (DM; DMEM/F12, 20% FBS, 1% Pen/Strep). After 5 days, embryoid bodies were plated on Matrigel-coated plates and cultured with DM. After 21 days in DM, cells were fixed, permeabilized, and stained for germ layer markers.

Fluorescence microscopy. All imaging was performed on a Nikon Ti-Eclipse inverted microscope with an LED-based Lumencor SOLA SE Light Engine using a Semrock band pass filter. GFP was visualized with an excitation at 472 nm and emission at 520 nm. BFP was visualized with the DAPI fluorescence channel with excitation at 395 nm and emission at 460 nm. mCherry was visualized with an excitation of 562 nm and emission at 641/75 nm.

Flow cytometry. Cells were dissociated with Accutase for 10 min at 37° C., triturated, and passed through a 40 µm cell strainer. Cells were then washed twice with flow cytometry buffer (BD Biosciences) and resuspended at a maximum concentration of 5×10⁶ cells per 100 µL. Flow cytometry analysis was performed on an Attune NxT (Thermo Fisher Scientific). Flow cytometry files were analyzed using with FlowJo (FlowJo LLC, Ashland, OR, USA).

Off-target analysis. For the data presented in FIG. 28, analysis was performed for the top three off-target loci for sg(BG) and sg(APOE-R158) predicted in silico via CCTop using default parameters for *S. pyogenes* Cas9 against human genome reference sequence hg38 (Stemmer et al., 2015). Determination of base editing at these off-target sites was performed in a similar manner to that at on-target sites. The PCR primers used to analyze these off-target sites are presented in Table 13.

Quantification of editing in mCherry expressing HEK293 cells. Sanger sequencing of PCRs from genomic DNA from mCherry HEK cells treated with or without base editor and sgRNA were analyzed using EditR (Kluesner et al., 2018). For forward sequencing reactions, the “sgRNA Sequence” was the same as the protospacer. For reverse sequencing reads, the “sgRNA Sequence” was the reverse complement of the protospacer. The 5' and 3' start are the corresponding nucleotide number (starting at 1 for the first nucleotide of the sequencing read) 100 bp upstream and downstream of the protospacer, respectively.

Apolipoprotein E (APOE) ELISA. Cells were seeded in a 6 well plate at a density of 3×10⁵ cells per well. Media was changed every 24 hours. On day 3, 24-hour conditioned media was collected and ApoE levels in the medium were measured with the Human APOE (AD2) ELISA Kit (Thermo Scientific).

Statistical analysis. Unless otherwise noted, all data are displayed as mean ± standard deviation (S.D).

SUPPLEMENTAL REFERENCES

- Kluesner, M. G., Nedveck, D. A., Lahr, W. S., Garbe, J. R., Abrahante, J. E., Webber, B. R., and Moriarity, B. S. (2018). EditR: A Method to Quantify Base Editing from Sanger Sequencing. *CRISPR J.* 1, 239-250.
- 10 Park, I.-H., Arora, N., Huo, H., Maherali, N., Ahfeldt, T., Shimamura, A., Lensch, M. W., Cowan, C., Hochedlinger, K., and Daley, G. Q. (2008). Disease-specific induced pluripotent stem cells. *Cell* 134, 877-886.
- 15 Stemmer, M., Thumberger, T., Del Sol Keyer, M., Wittbrodt, J., and Mateo, J. L. (2015). CCTop: An Intuitive, Flexible and Reliable CRISPR/Cas9 Target Prediction Tool. *PLoS One* 10, e0124633.

Example 3—CasMAs (XMAS)-TREE: A Cas9-Mediated Adenosine Transient Reporter for Editing Enrichment

Adenine base editors (ABE) enable single nucleotide modifications without the need for harmful double stranded DNA breaks (DSBs) induced by conventional CRISPR/Cas9-based approaches. However, most approaches that employ ABEs require inefficient downstream technologies to identify targeted cell populations. This example demonstrates development and characterization of a fluorescence-based method, entitled Cas9-mediated adenosine transient reporter for editing enrichment (CasMAs-TREE; herein abbreviated XMAS-TREE), to facilitate the real-time identification of base-edited cell populations. In particular, this section demonstrates use of XMAS-TREE to detect ABE activity. These studies also demonstrate that, at several independent loci, XMAS-TREE can be used to rapidly identify and purify modified cell populations. In addition, this section demonstrates that XMAS-TREE can be used in concert with multiplex editing schemes to efficiently edit several independent loci. In addition, XMAS-TREE can be used to edit human pluripotent stem cells (hPSCs), a cell type refractory to traditional gene editing approaches. In particular, XMAS-TREE allows for the efficient generation of clonal isogenic hPSCs at loci not editable using typical reporter of transfection (RoT)-based enrichment techniques. Collectively, XMAS-TREE is an easily implemented method that will greatly facilitate the use of ABEs in downstream basic biomedical science and translational applications.

Results

Development of a Fluorescent Reporter for Cas9-Mediated Adenosine Base Editing

As we have previously shown with cytosine base editors (CBEs), conventional approaches that use reporters of transfection, such as co-transfection or co-expression with a fluorescent protein (herein abbreviated as RoT) only report on the efficiency of plasmid delivery to a cell but not directly on the efficiency of base editing within these cells. To determine if the same was true with adenosine base editing approaches, HEK293 cells were transfected with a reporter plasmid (mCherry), an adenine base editor (ABEmax; pCMV-ABEmax), and a sgRNA for a genomic target site [sg(TS)]. This analysis revealed no correlation between transfection efficiency (percentage of mCherry-positive cells) and editing efficiency (percentage of A-to-G conver-

sion at target nucleotide) (FIG. 32). To that end, we sought to leverage our experience developing fluorescent reporters of editing activity to enable XMAS-TREE. To establish a fluorescent assay to detect ABE activity within a cell, we engineered a construct encoding a mCherry fluorescent protein followed by a stop codon (TGA) immediately preceding the coding sequence for a green fluorescent protein (GFP). Consequently, the A-to-G conversion of that codon to 'TGG' (encoding tryptophan) will enable translational read-through and expression of GFP. To determine the utility of this fluorescent-based construct to report on ABE activity, a vector was assembled with a human EF1 α promoter to drive expression of the fluorescent reporters (pEF-XMAS; FIG. 13A). In addition, we engineered two versions of this vector, one with a single stop codon (pEF-XMAS-1xStop) and another with two stop codons (pEF-XMAS-2xStop; FIG. 13B). It was speculated that A-to-G conversion of two stop codons within the editing window would provide a higher degree of stringency with respect to reporting on base editing activity within a cell. In addition, we designed a sgRNA vector [sg(XMAS)] that would direct the ABE to the target 'TGA' resulting in an A-to-G conversion and allow for subsequent translation of the downstream GFP cassette. Next, HEK293 cells were co-transfected with pEF-XMAS, pCMV-ABEmax, and sg(XMAS) or a control non-targeting sgRNA [sg(NT)]. Fluorescence microscopy (FIG. 13C) and flow cytometry (FIG. 13D) revealed that targeting pEF-XMAS with sg(XMAS) resulted in the generation of mCherry/GFP double positive cells, suggesting A-to-G base editing in the target codons allowing for GFP expression. Conversely, targeting pEF-XMAS-1xStop or pEF-XMAS-2xStop with sg(NT) did not result in the generation of any GFP positive cells (FIG. 33). Despite similarities in transfection efficiency between pEF-XMAS-1xStop and pEF-XMAS-2xStop (as measured by percentage of mCherry-positive cells), the percentage of GFP-positive cells was significantly lower in sg(XMAS) targeted cells transfected with pEF-XMAS-2xStop, suggesting that a higher level of base editing activity was necessary for the activation of GFP expression with the 2xStop plasmid. Interestingly, a significant percentage of cells that were mCherry-positive were not GFP-positive, verifying that the reporter of transfection (mCherry) does not report on base editing activity within a cell (FIG. 13D). Finally, we wanted to demonstrate that the fluorescent output associated with the XMAS-TREE reporter was transient. As such, the long-term fluorescence of cells transfected with pEF-XMAS and targeted with sg(XMAS) was measured. Indeed, analysis of these cells by flow cytometry (FIG. 13D) and fluorescence microscopy (FIG. 13E) revealed no long-term detectable fluorescent signal, confirming that the XMAS-TREE fluorescent output was transient. Collectively, this data establishes that editing of the XMAS-TREE plasmid provides a transient fluorescent reporter for base editing activity within a cell.

XMAS-TREE Allows for the Identification and Isolation of Base-Edited Cell Populations

Next, we wanted to demonstrate the utility of XMAS-TREE for the identification and isolation of cells in which targeted genomic adenosine base editing had occurred. To facilitate this, a dual-targeting sgRNA (pDT-sgRNA) vector that contains both sg(XMAS) and a guide matching an endogenous target site, sg(TS) was designed. Additionally, the pDT-sgRNA vector was designed to allow for the straightforward cloning of new target sites via BbsI restriction enzyme digestion and ligation of sg(TS) oligonucleotides. We designed pDT-sgRNA vectors with sequences targeting five genomic loci (Sites 1-5) as well as the pro-

motor of the γ -globin genes HBG1 and HBG2. To utilize XMAS-TREE for enrichment of cells that have been edited at a specific genomic location, we co-transfected these pDT-sgRNA vectors with pEF-XMAS-1xStop or 2xStop and pCMV-ABE into HEK293 cells (FIG. 14A). Flow cytometry was then used to isolate reporter positive cell populations and Sanger sequencing was performed on the targeted genomic sites in isolated populations (FIG. 14A). As expected, mCherry-positive/GFP-positive cells were enriched for edited cells when compared to double-negative cell populations (FIG. 14B). Importantly the transfection marker mCherry-positive population had significantly reduced editing compared to the editing reporter positive GFP-positive population. This demonstrates the benefit of utilizing a real-time reporter of base editing. (FIG. 14B). Finally, comparison of mCherry-positive/GFP-positive cells isolated using the 2xStop versus the 1xStop vector revealed that use of the 2xStop vector led to increased editing efficiencies, especially at loci (i.e., HBG1, HBG2) that were more resistant to editing. This suggests that at more difficult to edit loci, the 2xStop plasmid might provide a higher level of stringency necessary to enrich for edited cell populations. Overall, these results confirm that XMAS-TREE could be used to identify and enrich for adenosine base edited cell populations at a variety of genomic target sites.

XMAS-TREE Enables Efficient Multiplex Base Editing at Genomic Loci

XMAS-TREE was further evaluated to determine if it could be used for multiplexed genome editing. To that end, we generated a multi-targeting vector (pMT-sgRNA) that contains sg(TREE) as well as sgRNAs for multiple genomic targets. More specifically, we generated two pMT-sgRNA vectors—one that would target Site-1/Site-3/Site-4 and another that would simultaneously edit Site-5/HBG1/HBG2. We employed XMAS-TREE to simultaneously target multiple genomic sites by co-transfecting HEK293 cells with pMT-sgRNA, pEF-XMAS, and pCMV-ABEmax. Reporter-positive and -negative cells were isolated by flow cytometry and analyzed by Sanger sequencing at the targeted loci. Consistent with single locus targeting, mCherry-positive/GFP-positive cells displayed a significantly higher frequency of base editing at the target sites than editing levels that were observed in unsorted, mCherry-negative/GFP-negative, and mCherry-positive/GFP-negative cell populations (FIG. 15A). Importantly, there was no significant reduction in editing efficiency when XMAS-TREE was used to target these sites individually or a multiplexed fashion (FIG. 34).

Initial analysis of bulk sorted mCherry-positive/GFP-positive cells suggested that multiplexed editing with XMAS-TREE resulted in a large percentage of cells that had been simultaneously edited at multiple loci. To verify this observation, XMAS-TREE was used for the clonal isolation of base edited populations (FIG. 15B). Briefly, we co-transfected HEK293 cells with pEF-XMAS, pCMV-ABEmax, and a pMT-sgRNA designed to simultaneously target genomic Site-1/Site-3/Site-4. Single GFP-positive cells were sorted into a 96-well plate and expanded prior to analysis. Genomic DNA was isolated from clonal populations and the multiplexed genomic sites were subject to Sanger sequencing after PCR amplification. Remarkably, this analysis revealed that greater than 90% of the clones isolated had been edited, with 26 out of the 30 clones having biallelic conversions at all three genomic loci (FIG. 15C). In addition, we did not observe indels in any of the clones at these target sites. Lastly, we wanted to determine if XMAS-TREE increased A-to-G conversion at off-target loci. There-

fore, in several clones that had biallelic edits at all three target sites, we performed off-target analysis at the top predicted sites for sg(XMAS) as well as the sgRNAs used to target Site-1/Site-3/Site-4. At all of the off-target sites analyzed, we did not observe substantial A-to-G edits at these off-target loci (FIG. 35). In addition, indels were not observed at any of the off-target sites in the clones analyzed. Collectively, these results demonstrate the broad utility of XMAS-TREE to allow for the highly efficient, simultaneous editing of multiple independent loci.

Highly Efficient Editing of Human Pluripotent Stem Cells (hPSCs) Using XMAS-TREE

Traditional CRISPR-based approaches to modify single base pairs in hPSCs suffer from extremely low efficiencies. Therefore, we wanted to determine if XMAS-TREE could be utilized to efficiently mediate A-to-G conversions at specific loci in hPSCs. To confirm that the XMAS reporter was functioning in hPSCs, we transfected hPSCs with pEF-XMAS1xStop/2xStop, pEF-ABEmax, and sg(XMAS) or sg(NT). Similar to our experiments with HEK293 cells, fluorescence microscopy (FIG. 16A) and flow cytometry (FIG. 16B) with sg(XMAS), but not with sg(NT) (FIG. 36), resulted in the generation of mCherry-positive/GFP-positive cells, indicative of adenosine base editing of the pEF-XMAS reporter plasmid. Additionally, this analysis revealed that the proportion of cells that were positive for the base editing reporter (GFP) relative to the transfection reporter (mCherry) were markedly reduced in hPSCs, consistent with reports that hPSCs are recalcitrant to genomic modification. In this vein, these results suggest that purifying hPSC populations solely with a reporter of transfection (mCherry) would significantly dilute out cells with targeted genomic base edits. In addition, the level of base editing of the 2xStop plasmid was significantly lower than that observed with the 1xStop, suggesting that the 2xStop plasmid provides a higher degree of stringency in identifying base edited populations in hPSCs. Finally, flow cytometry (FIG. 16B) and fluorescence analysis (FIG. 16C) demonstrated that there was no detectable mCherry or GFP signal after 2 weeks of culture, confirming that the fluorescent signal associated with the XMAS-TREE reporter was transient in hPSCs.

Since we established pEF-XMAS reports on functional base editing in hPSCs, we wanted to determine if XMAS-TREE could be employed to enrich for cells with single-base pair edits at target loci in hPSCs. In this regard, we co-transfected hPSCs with pEF-XMAS1xStop/2xStop and pEF-ABEmax along with a pDT-sgRNA targeting genomic Site-1 or single base pair changes in AKAP9 and PSEN1 that have been previously associated with increased risks of developing Alzheimer's disease (AD). In turn, flow cytometry was used to purify reporter-positive and -negative cell populations and Sanger sequencing was performed on the targeted genomic locations in isolated populations. This analysis demonstrated that mCherry-positive/GFP-positive cells displayed a statistically significant increase in editing efficiency at the target loci when compared to other populations analyzed (FIG. 16D). In fact, in the more difficult to edit loci, AKAP9 and PSEN1, editing was virtually absent in populations not positive for our base editing reporter (GFP). Furthermore, mCherry-positive/GFP-positive cells isolated using the 2xStop plasmid allowed for greater level of enrichment. Together, these results demonstrated that XMAS-TREE can be used for the isolation of base-edited hPSC populations.

XMAS-TREE Enables Highly Efficient Generation of Clonal Isogenic hPSC Lines

We next wanted to compare the editing efficiency enabled by XMAS-TREE compared to conventional reporters of transfection (RoT). Accordingly, we co-transfected hPSCs with a reporter plasmid (pEF-mCherry), an adenine base editor (pEF-ABEmax), and a sgRNA for various genomic target sites [sg(TS)] (FIG. 17A). Flow cytometry was then used to sort mCherry-positive cell populations (RoT) and Sanger sequencing was performed on the targeted genomic sites. This analysis revealed that across all targeted sites that mCherry-positive/GFP-positive cells isolated using XMAS-TREE had a significantly higher frequency of base editing than mCherry-positive cells isolated using traditional RoT approaches (FIG. 17B). In fact, several targeted loci (i.e. Site-3, PSEN) displayed undetectable levels of editing when traditional RoT approaches were applied (FIG. 37). We then wanted to directly compare the efficiency by which XMAS-TREE and RoT-based methods could be utilized to generate clonal isogenic lines modified at these difficult to edit sites. To this end, we transfected hPSCs with pEF-XMAS, pEF-ABEmax, and pDT-sgRNA containing a sgRNA to target genomic Site-3. Single mCherry-positive/GFP-positive cells were sorted into 96-well plates, expanded, and subject to Sanger sequencing. Of the 10 clones analyzed, 80% had a homozygous A-to-G edit at the genomic Site-3 locus (FIG. 17C). Importantly, indels were not identified in any of the clones at the target site. For comparison to a more conventional RoT approach to generate isogenic lines, this same hPSC line was transfected with a plasmid in which the base-editor (ABEmax) was co-transfected with a pEFmCherry vector as well as the same sgRNA for the Site-3 locus. After 48 hours post-transfection, single GFP-positive cells were sorted into 96-well plates. Clonal lines were then passaged, expanded, and subjected to Sanger sequencing at the targeted locus. Notably, analysis of 10 clonal lines revealed that this RoT-based approach did not result in generation of a single isogenic clone at the target site (FIG. 17C). This ability of XMAS-TREE to generate isogenic clonal lines at sites that did not display significant editing in bulk RoT approaches was also confirmed at the PSEN locus (FIG. 38). In sum, these results demonstrate that XMAS-TREE can not only provide for a higher level of enrichment of base-edited cell populations compared to RoT approaches, but also can allow for the generation of isogenic lines at genomic loci that are not achievable with conventional RoT methods.

Multiplex Editing of hPSCs Using XMAS-TREE

Lastly, we wanted to establish that XMAS-TREE could allow for multiplexed genome modification in hPSCs. hPSCs were co-transfected with pEF-XMAS, pEF-ABEmax, and a pMT-sgRNA with sgRNAs targeting Site-5, HBG1, and HBG2. Similar to our results obtained with HEK293 cells, mCherry-positive/GFP-positive cells had a statistically significant higher level of base editing at all three target sites when compared to those in unsorted, mCherry-negative/GFP-negative, and mCherry-positive/GFP-negative cell populations (FIG. 18A). In addition, direct comparison of multiplex editing using XMAS-TREE and RoT approaches demonstrated that XMAS-TREE allowed for a statistically significant higher level of base editing than by RoT-based methods (FIG. 18B). Altogether, this data demonstrates that XMAS-TREE enables efficient simultaneous editing of multiple loci in hPSCs.

Discussion

Together, CBEs and ABEs have the potential ability to modify up to 60% of the disease-causing point mutations.

That said, BEs can be used in the context of cellular models of human disease models to establish genotype-to-phenotype relationship associated with genetic risk factors, investigate disease mechanisms, and test therapeutic strategies. In our previous work, we describe the development of a transient reporter for editing enrichment (TREE) as a fluorescence-based assay to report on cytosine base editing (CBE) activity within a single cell. In this work, we develop an analogous reporter system, Cas9-mediated adenosine transient reporter for editing enrichment (CasMAS-TREE; XMAS-TREE) that allows for the real-time detection of adenosine base editing for the identification and enrichment of base-edited cell populations. Notably, at several loci, XMAS-TREE allows for the targeted gene editing at efficiencies approaching 90%. As part of these efforts, we also utilized XMAS-TREE to enrich for cells that have been edited at several disease-relevant loci including those associated with sickle-cell anemia (i.e., HBG1, HBG2) and Alzheimer's disease (i.e., AKAP9, PSEN1). In addition, we demonstrate that XMAS-TREE can be used in the context of multiplex genome engineering strategies to facilitate simultaneous A-to-G (or T-to-C) conversions at several independent loci at the same efficiencies when single loci were targeted. Critically, the ability of XMAS-TREE to generate clonal lines that had been simultaneously edited at multiple loci will enable the facile generation of cell-based models of polygenetic diseases. Finally, we establish that the same XMAS-TREE-based methods can be applied in human pluripotent stem cells (hPSCs), a cell population in which gene editing technologies, including base editors and multiplex genome modification, have been challenging to implement. In particular, we show that XMAS-TREE can facilitate the establishment of isogenic hPSC lines at loci that were not able to be modified using well-accepted reporter of transfection (RoT) methods. In fact, we show that at certain target sites that XMAS-TREE can allow for derivation of isogenic clonal populations with biallelic modification with 80% efficiency. Notably, all targeted clones were free from indels at all on-target sites. The clonal targeting efficiencies that we observe with XMAS-TREE in hPSCs are significantly higher than those previously reported with other CRISPR/Cas9-based methods, which are often in the single digits at most loci. In addition, the inefficiencies associated with these well-established methods make it difficult to achieve homozygous or multiplexed editing in hPSCs.

We speculate that XMAS-TREE can be utilized in other applications not described in this example. For example, several groups have reported the generation of additional ABEs with non-NGG PAM specificities, narrower targeting windows, and reduced by-product formation 34-36. Accordingly, future application of XMAS-TREE with these next-generation ABE variants will be straightforward. In addition, we anticipate that XMAS-TREE can be applied to induce alterations in target gene expression. More specifically, we previously described how CBEs can be used with other TREE-based strategies to generate gene knockout lines without the introduction of DSBs through in-frame conversion of 'CAG' codon encoding for glutamine to a 'TAG' premature stop codon. However, these approaches do not allow targeting for all genes and can be limited by the propensity of CBEs to induce genome-wide Cas9-independent off-target mutations. As an alternative, Wang and colleagues recently described an ABE-mediated strategy to induce gene knockout through modification of the ATG start codon to ACG or GTG. Moving forward, XMAS-TREE can be utilized with such strategies to enrich for cell populations with targeted gene knockouts.

In summary, there are several features of XMAS-TREE based methods that will enable extensive use by the research community. First, XMAS-TREE only requires the use of common lipid-based reagents for cell transduction. We envision that XMAS-TREE is compatible with other DNA delivery systems (i.e., electroporation) or expression methods (i.e., ribonucleoprotein complexes [RNP]) that have been utilized in other CRISPR/Cas9- and BE-based genome engineering strategies. In the future, XMAS-TREE associated plasmids can also be easily cloned into nonintegrating viral vectors to facilitate the development of in vivo gene editing methods 2. Second, we have designed the sgRNA vectors to allow for the simple restriction enzyme-based cloning of new target sites. In this regard, we show that XMAS-TREE can allow for the highly efficient editing of a diverse set of loci across multiple cell lines. In the future, XMAS-TREE can be easily utilized in other animal, primary, or immortalized cell types. In addition, because of the high editing efficiencies associated with XMAS-TREE, establishment of clonal lines with the targeted base pair edit does not require the screening of hundreds of clones, which is typical of other methods. Finally, we demonstrate that the use of XMAS-1xStop and -2xStop plasmids allows the end-user to balance the need for cell yield versus editing stringency. Specifically, the XMAS-1xStop plasmid provides for a higher degree of cell yield compared to the XMAS-2xStop plasmid while allowing for enrichment of editing cells at levels higher than conventional approaches. Alternatively, the XMAS-2xStop plasmid enables a higher degree enrichment at the target loci, especially at difficult to edit genomic locations. Collectively, these enabling features of XMAS-TREE will significantly enhance the use of ABE-based technologies in a variety of contexts and cell populations.

It should be noted that the above description, attached figures and their descriptions are intended to be illustrative and not limiting. Many themes and variations of this disclosure will be suggested to one skilled in this art, in light of the disclosure. All such themes and variations are within the contemplation hereof. For instance, while this invention has been described in conjunction with the various exemplary embodiments outlined above, various alternatives, modifications, variations, improvements, and/or substantial equivalents, whether known or that rare or may be presently unforeseen, may become apparent to those having at least ordinary skill in the art. Various changes may be made without departing from the spirit and scope of the invention. Therefore, the invention is intended to embrace all known or later-developed alternatives, modifications, variations, improvements, and/or substantial equivalents of these exemplary embodiments.

TABLE 4

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
Site-1	GGCCCAGACTGAGCAGCTG A (SEQ ID NO: 14)
Site-2	GAACACAAAGCATAGACTG C (SEQ ID NO: 15)

53

TABLE 4-continued

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
Site-3	GGCACTGCGGCTGGAGGTG G (SEQ ID NO: 16)
APOE(R158)	GAAGCGCCTGGCAGTGTAC C (SEQ ID NO: 17)

54

TABLE 4-continued

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
BFP(H66Y)	GACCCACGGCGTGCAGTGCT T (SEQ ID NO: 18)
C1ORF228	GTGCTGTTAGCACCTGGAA A (SEQ ID NO: 19)

TABLE 5

List of primers used in this example to amplify on-target sites.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-1	ATGTGGGCTGCCTAGAAAGG (SEQ ID NO: 20)	CCCAGCCAAACTTGTCACCC (SEQ ID NO: 21)
Site-2	CCAGCCCCATCTGTCAAAC (SEQ ID NO: 22)	TGAATGGATTCTTGGAAACAATGA (SEQ ID NO: 23)
Site-3	TGGTCTTCTTTCCCTCCCTGCCCTCC (SEQ ID NO: 24)	GGCCTGGAGGCGGGGCTCAGAGA (SEQ ID NO: 25)
APOE(R158)	GGACGAGACCATGAAGGAGTTGAAGGC (SEQ ID NO: 26)	CCACCTGCTCCTTCACCTCGTCCAG (SEQ ID NO: 27)

TABLE 6

Parameters for EditR analysis.				
Target Site	Sequencing Direction	Protospacer	5' bound	3' bound
Site 1	Forward	GGCCCAGACTGAGCAGGTGA (SEQ ID NO: 28)	GGCCTGGGTCAA (SEQ ID NO: 29)	TTCTTTCTCTG (SEQ ID NO: 30)
	Reverse	TCACGTGCTCAGTCTGGGCC (SEQ ID NO: 31)	GAGGAAAGGAAGCCCTGCT (SEQ ID NO: 32)	CAGGCCAGGGCTGGA (SEQ ID NO: 33)
Site-2	Forward	GAACACAAAGCATAGACTGC (SEQ ID NO: 34)	CCCGCTGGCCCTGT (SEQ ID NO: 35)	TCAGGCTGGCCCGC (SEQ ID NO: 36)
	Reverse	GCAGTCTATGCTTTGTGTTC (SEQ ID NO: 37)	CCAGCCCGCTGGCCCTGTA (SEQ ID NO: 38)	AGCTATTGAGGCT (SEQ ID NO: 39)
Site 3	Forward	GTGGCACTGCGGCTGGAGGT (SEQ ID NO: 40)	GATGACAGGCAGGGGCA (SEQ ID NO: 41)	CAGCACCAGA (SEQ ID NO: 42)
	Reverse	ACCTCCAGCCGAGTGCC (SEQ ID NO: 43)	CCGCGGTGCCCTGCCT (SEQ ID NO: 44)	AAGCGGAGACTCTGGTGC (SEQ ID NO: 45)
APOE(R158)	Forward	GAAGCGCCTGGCAGTGTACC (SEQ ID NO: 46)	CTGCGCAAGCTGCG (SEQ ID NO: 47)	TCGGCGCCCTCGCG (SEQ ID NO: 48)
	Reverse	GGTACACTGCCAGGCGCTTC (SEQ ID NO: 49)	GGATGGCGCTGA (SEQ ID NO: 50)	GCCTCGCCTCCACC (SEQ ID NO: 51)

TABLE 7

PCR conditions for each target site analyzed by Sanger sequencing.						
Target	Initial denature time and temperature	Denature time and temperature	Annealing time and temperature 40 cycles	Extension time and temperature	Final extension time and temperature	
Site-1	98° C., 45 seconds	98° C., 10 seconds	54° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes	
Site-2	98° C., 45 seconds	98° C., 10 seconds	56° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes	

TABLE 7-continued

PCR conditions for each target site analyzed by Sanger sequencing.					
Target	Initial denature time and temperature	Denature time and temperature	Annealing time and temperature 40 cycles	Extension time and temperature	Final extension time and temperature
Site-3	98° C., 45 seconds	98° C., 10 seconds	56° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes
APOE(R158)	98° C., 45 seconds	98° C., 10 seconds	62° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes

TABLE 8

List of primers used in this example to amplify off-target sites. Abbreviations: BG-OT = Off-targets associated with sg(BG), Site 1-OT = Off-targets associated with sg(Site-1), Site2-OT = Off-targets associated with sg(Site-2), Site3-OT = Off-targets associated with sg(Site-3).		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
BG-OT1	GATGCGCTTCCGGAAGACC (SEQ ID NO: 52)	GCTTCTTGAGCTTCTCAGCG (SEQ ID NO: 53)
BG-OT2	GGTAGCATGTTACGGCACCAG (SEQ ID NO: 54)	CATCCCTAGTACCGAATCCCATATAGC (SEQ ID NO: 55)
BG-OT3	CATCCTCCACCTAAGCCTTTCAA (SEQ ID NO: 56)	TTGAGTTAATAGCATTATAACAATTTCCACA (SEQ ID NO: 57)
BG-OT4	ACTCCTTACAACCGGAAGGCAAAC (SEQ ID NO: 58)	TGGACGTGGTGAAGCCCGTGGTG (SEQ ID NO: 59)
BG-OT5	TAGGTCTCTAGGGGCCTCTG (SEQ ID NO: 60)	AGGCTGCCCAACAGCCCCACT (SEQ ID NO: 61)
Site1-OT1	TCCCCTGTTGACCTGGAGAA (SEQ ID NO: 62)	CACTGTACTTGCCCTGACCA (SEQ ID NO: 63)
Site1-OT2	TGAGATGTGGGAGAGGG (SEQ ID NO: 64)	TTGGTGTGACAGGGAGCAA (SEQ ID NO: 65)
Site1-OT3	GTCCAAAGGCCAAGAACCT (SEQ ID NO: 66)	TGAGAGGGAACAGAAGGGCT (SEQ ID NO: 67)
Site1-OT4	GCTCATCTTAATCTGCTCAGCC (SEQ ID NO: 68)	TCCTAGCACTTTGGAAGGTCG (SEQ ID NO: 69)
Site1-OT5	AAAGGAGCAGCTCTTCCTGG (SEQ ID NO: 70)	GTCTGCACCATCTCCACAA (SEQ ID NO: 71)
Site2-OT1	GTGTGGAGAGTGAGTAAGCCA (SEQ ID NO: 72)	ACGGTAGGATGATTTTCAGGCA (SEQ ID NO: 73)
Site2-OT2	TTTTTTGGTACTCGAGTGTTATTCAG (SEQ ID NO: 74)	CACAAAGCAGTGTAGCTCAGG (SEQ ID NO: 75)
Site3-OT1	GGCATGGCTTCTGAGACTCA (SEQ ID NO: 76)	CCCCTTGCACTCCCTGTCTTT (SEQ ID NO: 77)
Site3-OT2	GAAGAGGCTGCCCATGAGAG (SEQ ID NO: 78)	TTTGGCAATGGAGGCATTGG (SEQ ID NO: 79)
Site3-OT3	GGTCTGAGGCTCGAATCCTG (SEQ ID NO: 80)	CTGTGGCCTCCATATCCCTG (SEQ ID NO: 81)
Site3-OT4	TTTCCACCAGAACTCAGCCC (SEQ ID NO: 82)	CCTCGGTTCTCCACAACAC (SEQ ID NO: 83)
Site3-OT5	GCAGGGGAGGATAAAGCAG (SEQ ID NO: 84)	CACGGGAAGGACAGGAGAAG (SEQ ID NO: 85)

TABLE 9

List of primers used in this example for NGS analysis.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-1	ATGTGGGCTGCCTAGAAAGG (SEQ ID NO: 86)	CCCAGCCAACTTGTCACCC (SEQ ID NO: 87)
APOE(R158)	GGACGAGACCATGAAGGAGTTGAAGGC (SEQ ID NO: 88)	CCACCTGCTCCTTCACCTCGTCCAG (SEQ ID NO: 89)

TABLE 10

PCR conditions for each target site subjected to NGS analysis.					
Target	Initial denature time and temperature	Denature time and temperature	Annealing time and temperature 40 cycles	Extension time and temperature	Final extension time and temperature
Site-1	98° C., 45 seconds	98° C., 10 seconds,	54° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes
APOE(R158)	98° C., 45 seconds	98° C., 10 seconds,	62° C., 5 seconds	72° C., 20 seconds	72° C., 10 minutes

TABLE 11

Comparison of editing efficiency using RoT-based approaches at the same target loci in this example, Komar et al., and Koblan et al.									
Figure 3E Standage-Beier et al. Reporter of Transfection				Figure 5C Komar et.al Sci Adv. 2017 Aug 30;3(8) No Reporter			Figure 1C Koblan et. al Nat Biotechnol. 2018 October;36(9):843-846 Reporter of Transfection		
	Unsorted	Reporter–	Reporter+	Unsorted	Reporter–	Reporter+	Unsorted	Reporter–	Reporter+
Site-1 (HEK 3)	21.3 ± 2.9	3.3 ± 2.8	40.7 ± 7.0	~45	N/A	N/A	~38	N/A	~55
Site-2 (HEK 2)	36.6 ± 3.8	13.3 ± 5.9	49.7 ± 5.1	~35	N/A	N/A	~20	N/A	~38
Site-3 (HEK 4)	24.0 ± 6.6	7.6 ± 5.0	45.3 ± 1.5	~45	N/A	N/A	~25	N/A	~40

TABLE 12

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
BFP (H66Y)	GACCCACGGCGTGCACTGCT T (SEQ ID NO: 90)
Site-1	GGCCCAGACTGAGCACGTGA (SEQ ID NO: 91)
Site-2	GAACACAAAGCATAGACTGC (SEQ ID NO: 92)
Site-3	GGCACTGCGGCTGGAGGTGG (SEQ ID NO: 93)
APOE (R158C)	GAAGCGCCTGGCAGTGATCC (SEQ ID NO: 94)

TABLE 12-continued

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
APOE (Q39X)	GTGGCAGAGCGGCCAGCGCT (SEQ ID NO: 95)
mCh1	GCACCCAGACCGCCAAGCTG A (SEQ ID NO: 96)
mCh2	GACCCAGGACTCCTCCCTGC (SEQ ID NO: 97)
mCh3	GCAAGCAGAGGCTGAAGCTG A (SEQ ID NO: 98)
Non-target (NT)	GGGTCTTCAGAAAGACCT (SEQ ID NO: 99)

TABLE 13

List of primer sequences used in this example for on- and off-target sites. Abbreviations: BG-OT = Off-targets associated with sg(BG), APOE(R158C) = Off-targets associated with sg(APOE ^{R158C})		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-1	ATGTGGGCTGCCTAGAAAGG (SEQ ID NO: 100)	CCCAGCCAACTTGTCACCC (SEQ ID NO: 101)

TABLE 13-continued

List of primer sequences used in this example for on- and off-target sites.
Abbreviations: BG-OT = Off-targets associated with sg(BG), APOE(R158C) = Off-targets associated with sg(APOE^{R158C})

Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-2	CCAGCCCCATCTGTCAAAC (SEQ ID NO: 102)	TGAATGGATTCTTGGAAACAATGA (SEQ ID NO: 103)
Site-3	TGGTCTTTCTTCCCTCCCTGCCCTC (SEQ ID NO: 104)	GGCCTGGAGGCGGGGCTCAGAGA (SEQ ID NO: 105)
APOE(R158C)	GGACGAGACCATGAAGGAGTTGAAGG (SEQ ID NO: 106)	CCACCTGCTCCTTCACCTCGTCCAG (SEQ ID NO: 107)
APOE(Q39X)	TCAGAAGGACCTGACCCACCT (SEQ ID NO: 108)	ATGAAACCTGGACCTGGGAGGTATA (SEQ ID NO: 109)
mCherry	AGCTGTGACCGCGCCTACG (SEQ ID NO: 110)	GGGATTCTCCTCCACGTCAC (SEQ ID NO: 111)
BG-OT1	GATGCGCTTCCGGAAGACC (SEQ ID NO: 112)	GCTTCTTGAGCTTCTCAGCG (SEQ ID NO: 113)
BG-OT2	GGTAGCATGTTCCAGCACCAG (SEQ ID NO: 114)	CATCCCTAGTACCGAATCCCATATAGC (SEQ ID NO: 115)
BG-OT3	CATCCTCCACCTAAGCCTTTCAA (SEQ ID NO: 116)	TTGAGTTAATAGCATTATAACAATTTC CACA (SEQ ID NO: 117)
APOE(R158C) - OT1	GATACACCATAAAGGGTTTGACTG (SEQ ID NO: 118)	ACCATTTCCTCCCAATTCTACTC (SEQ ID NO: 119)
APOE(R158C) - OT2	CATCTGCATTGGCTTGAAACATC (SEQ ID NO: 120)	TTACAAAAGTGCTAAATGATGCACAT (SEQ ID NO: 121)
APOE(R158C) - OT3	ACTCAGTAAAGCTCCTCTTCAAC (SEQ ID NO: 122)	TTTGTCTTAGGTCCACTGGGC (SEQ ID NO: 123)

35

TABLE 14

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
XMAS-1xStop	GTTGATGGGGTGGTTCAGGA (SEQ ID NO: 124)
XMAS-2xStop	GTTGATGAGGTGGTTCAGGA (SEQ ID NO: 125)
Site-1	GAACACAAAGCATAGACTGC (SEQ ID NO: 126)
Site-2	GAGTATGAGGCATAGACTGC (SEQ ID NO: 127)
Site-3	GATGAGATAATGATGAGTCA (SEQ ID NO: 128)
Site-4	GGATTGACCCAGGCCAGGGC (SEQ ID NO: 129)

40

45

50

TABLE 14-continued

List of sgRNA sequences used in this example.	
Site	Sequence (5'→3')
Site-5	GTAGAAAAAGTATAGACTGC (SEQ ID NO: 130)
HBG1	GCTTGACCAATAGCCTTGACA (SEQ ID NO: 131)
HBG2	GATATTGCTTGGAGATAGTG (SEQ ID NO: 132)
AKAP9	GAATAAGTTGAAGAAAAAG (SEQ ID NO: 133)
PSEN	GCACAGAAGATACCGAGACTG (SEQ ID NO: 134)
Non-target (NT)	GGGTCTTCGAGAAGACCT (SEQ ID NO: 135)

TABLE 15

List of primer sequences used in this example.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-1	TCCTTGGAAACAATGATAACAAGAC (SEQ ID NO: 136)	CCAGCCCCATCTGTCAAAC (SEQ ID NO: 137)
Site-2	GCTTATATTCTAGGGAGACAGACAT (SEQ ID NO: 138)	ACCTGAGGTCAGAAGTTGAGA (SEQ ID NO: 139)

TABLE 15-continued

List of primer sequences used in this example.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
Site-3	GTCTGAGGTCACACAGTGGG (SEQ ID NO: 140)	AGAGCAGGGACCACATCTAC (SEQ ID NO: 141)
Site-4	GCCAAACTTGTCAACCGTA (SEQ ID NO: 142)	ATGTGGGCTGCCTAGAAAGG (SEQ ID NO: 143)
Site-5	TCCATTTATATGAAATGTTGAGAAAAG GCAAAT (SEQ ID NO: 144)	GTAACATATGCTCTCTGATTCTCC TATTAGC (SEQ ID NO: 145)
HBG1	CCTACCTTCCCAGGGTTT (SEQ ID NO: 146)	AAGAAGTCCTGGTATCTTCTATG (SEQ ID NO: 147)
HBG2	TCAGACGTTCAGAAAGCGAG (SEQ ID NO: 148)	GACAAGAAGGTGAAAAACGGCTG (SEQ ID NO: 149)
AKAP9	GATTCAAAGCATACCAGAGAATAGT (SEQ ID NO: 150)	TCAAACATAGTATGCATTTCAACAAC (SEQ ID NO: 151)
PSEN1	GAGTGTAAGCTGTTTTTCTCAGGTT (SEQ ID NO: 152)	GAATACCCAACCATAAGAAGAACAG (SEQ ID NO: 153)

TABLE 16

Phire PCR conditions for each target site analyzed by Sanger sequencing.					
Target	Initial denature time and temperature	Denature time and temperature	Annealing time and temperature 40 cycles	Extension time and temperature	Final extension time and temperature
Site-1	98 C. 5 min	98 C. 5 sec	58 C. 5 sec	72 C. 30 sec	72 C. 5 min
Site-2	98 C. 5 min	98 C. 5 sec	62 C. 5 sec	72 C. 30 sec	72 C. 5 min
Site-3	98 C. 5 min	98 C. 5 sec	56.8 C. 5 sec	72 C. 30 sec	72 C. 5 min
Site-4	98 C. 5 min	98 C. 5 sec	61.3 C. 5 sec	72 C. 30 sec	72 C. 5 min
Site-5	98 C. 5 min	98 C. 5 sec	65 C. 5 sec	72 C. 30 sec	72 C. 5 min
HBG1	98 C. 5 min	98 C. 5 sec	59.2 C. 5 sec	72 C. 30 sec	72 C. 5 min
HBG2	98 C. 5 min	98 C. 5 sec	59 C. 5 sec	72 C. 30 sec	72 C. 5 min
AKAP9	98 C. 5 min	98 C. 5 sec	64 C. 5 sec	72 C. 30 sec	72 C. 5 min
PSEN1	98 C. 5 min	98 C. 5 sec	63 C. 5 sec	72 C. 30 sec	72 C. 5 min

40

TABLE 17

Parameters for EditR analysis.			
Target Site	Protospacer	5' bound	3' bound
Site-1	GAACACAAAGCATAGAC TGC (SEQ ID NO: 154)	50	120
Site-2	GAGTATGAGGCATAGAC TGC (SEQ ID NO: 155)	140	200
Site-3	GATGAGATAATGATGAG TCA (SEQ ID NO: 156)	100	180
Site-4	GGATTGACCCAGGCCA GGGC (SEQ ID NO: 157)	80	160
Site-5	GCAGTCTATACTTTTTC TAC (SEQ ID NO: 158)	40	120

TABLE 17-continued

Parameters for EditR analysis.			
Target Site	Protospacer	5' bound	3' bound
HBG1	CTTGACCAATAGCCTTG ACA (SEQ ID NO: 159)	160	260
HBG2	ATATTGCATTGAGATA GTG (SEQ ID NO: 160)	120	220
AKAP9	GAAAATAGTTGAAGAAA AAG (SEQ ID NO: 161)	200	300
PSEN1	CACAGAAGATACCGAGA CTG (SEQ ID NO: 162)	120	200

TABLE 18

List of primers used in this example to amplify off-target sites.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
XMAS-OT1	CAGCATTATCCATTGTGCGCA (SEQ ID NO: 163)	TGGAGACAGCGAGTCTACAGC (SEQ ID NO: 164)

TABLE 18-continued

List of primers used in this example to amplify off-target sites.		
Primer	Forward Sequence (5'→3')	Reverse Sequence (5'→3')
XTAS-OT2	TAACACCATTATAGCTGAAGTGGGG (SEQ ID NO: 165)	TGAGTTACACACAAGCCAGTTAAATTC (SEQ ID NO: 166)
XTAS-OT3	AGGGAGTGGACATGAGGCGA (SEQ ID NO: 167)	CCCAAGAGGAAGTCCCAAGG (SEQ ID NO: 168)
Site-1-OT1	CCTTGGGAAGAGAAGGGGTC (SEQ ID NO: 169)	GAGATACCGGAAGCTTTGATGTAAGA (SEQ ID NO: 170)
Site-1-OT2	CTTGGGGAGAAAGGTCCAGG (SEQ ID NO: 171)	CAAGCTTTTCTCTGGGATGTAAAA (SEQ ID NO: 172)
Site-1-OT3	CTGGCAAGCTGTTCTCACATG (SEQ ID NO: 173)	GAGGCTGAGGCAGGAGTATG (SEQ ID NO: 174)
Site-3-OT1	GTTTTTCAGTAGAAGAGTATATAATACATA AT (SEQ ID NO: 175)	ATATTCTCAGCCTAGGCCTG (SEQ ID NO: 176)
Site-3-OT2	TGTTGGACATGGGTGCCTTATT (SEQ ID NO: 177)	TTCACCCCTCTCTGGATGGCG (SEQ ID NO: 178)
Site-3-OT3	GCAGGAGGAGGCAAGTGAAG (SEQ ID NO: 179)	CAGAGAAATAACACTCTGGCAGCTG (SEQ ID NO: 180)
Site-4-OT1	CAGCATTATACGCAGTATTGTTATTG (SEQ ID NO: 181)	TCATTTTCGTGTTGTGCTTTTACTTAA A (SEQ ID NO: 182)
Site-4-OT2	GTGAGCAGTAAACTTAATTGTTGATACA ATAAATC (SEQ ID NO: 183)	CTTTTAGAATGAAAGTGTGCATCTTAGTA AAGAAA (SEQ ID NO: 184)
Site-4-OT3	GTTCTCTACTGATTCTCAGCAGG (SEQ ID NO: 185)	CACAAAAGGGATAAATGCTCTATCCATT (SEQ ID NO: 186)

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 327

<210> SEQ ID NO 1

<211> LENGTH: 239

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 1

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
1 5 10 15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20 25 30

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35 40 45

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50 55 60

Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg Tyr Pro Asp His Met Lys
65 70 75 80

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85 90 95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100 105 110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115 120 125

-continued

```

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
 130                      135                      140

Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145                      150                      155                      160

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
                      165                      170                      175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
                      180                      185                      190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
                      195                      200                      205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210                      215                      220

Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
225                      230                      235

```

```

<210> SEQ ID NO 2
<211> LENGTH: 239
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
                        polypeptide

```

```

<400> SEQUENCE: 2

```

```

Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly Val Val Pro Ile Leu
 1                      5                      10                      15

Val Glu Leu Asp Gly Asp Val Asn Gly His Lys Phe Ser Val Ser Gly
20                      25                      30

Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile
35                      40                      45

Cys Thr Thr Gly Lys Leu Pro Val Pro Trp Pro Thr Leu Val Thr Thr
50                      55                      60

Leu Thr His Gly Val Gln Cys Phe Gly Arg Tyr Pro Asp His Met Lys
65                      70                      75                      80

Gln His Asp Phe Phe Lys Ser Ala Met Pro Glu Gly Tyr Val Gln Glu
85                      90                      95

Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu
100                     105                     110

Val Lys Phe Glu Gly Asp Thr Leu Val Asn Arg Ile Glu Leu Lys Gly
115                     120                     125

Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu Gly His Lys Leu Glu Tyr
130                      135                      140

Asn Tyr Asn Ser His Asn Val Tyr Ile Met Ala Asp Lys Gln Lys Asn
145                      150                      155                      160

Gly Ile Lys Val Asn Phe Lys Ile Arg His Asn Ile Glu Asp Gly Ser
165                      170                      175

Val Gln Leu Ala Asp His Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly
180                      185                      190

Pro Val Leu Leu Pro Asp Asn His Tyr Leu Ser Thr Gln Ser Ala Leu
195                      200                      205

Ser Lys Asp Pro Asn Glu Lys Arg Asp His Met Val Leu Leu Glu Phe
210                      215                      220

Val Thr Ala Ala Gly Ile Thr Leu Gly Met Asp Glu Leu Tyr Lys
225                      230                      235

```

-continued

<210> SEQ ID NO 3
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 3

gacccacggc gtgcagtgt tgttttagag ctgaaatag caagttaaaa taaggctagt 60
ccgttatcaa cttgaaaaag tggcaccgag tcggtgcttt ttgtttt 108

<210> SEQ ID NO 4
<211> LENGTH: 105
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 4

gggtcttcga gaagacctgt ttttagagta gaaatagcaa gttaaaataa ggctagtccg 60
ttatcaactt gaaaaagtgg caccgagtcg gtgctttttt gtttt 105

<210> SEQ ID NO 5
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 5

ggcccagact gagcacgtga gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 6
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 6

gaacacaaag catagactgc gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 7
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 7

ggcactgcgg ctggagggtgg gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 8
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 8

gaagcgctg gcagtgtacc gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 9
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 9

gtgctgtag caccctggaa agtttttagag ctgaaatag caagttaaaa taaggctagt 60
ccggttatcaa cttgaaaaag tggcaccgag tcggtgcttt tttgtttt 108

<210> SEQ ID NO 10
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 10

gtggcagagc ggccagcgct gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 11
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 11

gcaccagac cgccaagctg agtttttagag ctgaaatag caagttaaaa taaggctagt 60
ccggttatcaa cttgaaaaag tggcaccgag tcggtgcttt tttgtttt 108

<210> SEQ ID NO 12
<211> LENGTH: 107
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 12

gaccaggac tcctccctgc gtttttagagc tagaaatagc aagttaaaat aaggctagtc 60
cgttatcaac ttgaaaaagt ggcaccgagt cggtgctttt ttgtttt 107

<210> SEQ ID NO 13
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

-continued

<400> SEQUENCE: 13

gcaagcagag gctgaagctg agtttttagag ctagaaatag caagttaaaa taaggctagt 60

ccgttatcaa cttgaaaaag tggcacgag tcggtgcttt tttgtttt 108

<210> SEQ ID NO 14

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 14

ggcccagact gagcacgtga 20

<210> SEQ ID NO 15

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 15

gaacacaaag catagactgc 20

<210> SEQ ID NO 16

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 16

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 17

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 17

gaagcgctg gcagtgtacc 20

<210> SEQ ID NO 18

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 18

gaccacggc gtgcagtgt t 21

<210> SEQ ID NO 19

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 19

gtgctgttag caccctggaa a

21

<210> SEQ ID NO 20

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 20

atgtgggctg cctagaaagg

20

<210> SEQ ID NO 21

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 21

cccagccaaa cttgtcaacc

20

<210> SEQ ID NO 22

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 22

ccagcccat ctgtcaaaact

20

<210> SEQ ID NO 23

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 23

tgaatggatt ccttggaac aatga

25

<210> SEQ ID NO 24

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 24

tggctcttctt tccctcccc tgccctcc

28

<210> SEQ ID NO 25

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 25

ggcctggagg cgggggctca gaga

24

<210> SEQ ID NO 26

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 26

ggacgagacc atgaaggagt tgaaggc

27

<210> SEQ ID NO 27

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 27

ccacctgctc cttcacctcg tccag

25

<210> SEQ ID NO 28

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 28

ggcccagact gagcacgtga

20

<210> SEQ ID NO 29

<211> LENGTH: 12

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 29

ggcctgggctc aa

12

<210> SEQ ID NO 30

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 30

ttcctttcct ctg

13

<210> SEQ ID NO 31

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 31

-continued

tcacgtgctc agtctgggcc 20

<210> SEQ ID NO 32
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 32

gaggaaagga agccctgct 19

<210> SEQ ID NO 33
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 33

caggccaggg ctgga 15

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 34

gaacacaaaag catagactgc 20

<210> SEQ ID NO 35
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 35

cccgtggcc ctgt 14

<210> SEQ ID NO 36
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 36

tcaggctggc ccgc 14

<210> SEQ ID NO 37
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 37

-continued

gcagtcctatg ctttgtgttc 20

<210> SEQ ID NO 38
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 38

ccagcccgcgt ggccctgta 19

<210> SEQ ID NO 39
<211> LENGTH: 13
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 39

agctattcag gct 13

<210> SEQ ID NO 40
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 40

gtggcactgc ggctggaggt 20

<210> SEQ ID NO 41
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 41

gatgacaggc aggggca 17

<210> SEQ ID NO 42
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 42

cagcaccaga 10

<210> SEQ ID NO 43
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 43

acctccagcc gcagtgcc 18

-continued

<210> SEQ ID NO 44
<211> LENGTH: 17
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 44

ccgcggtgcc cctgcct 17

<210> SEQ ID NO 45
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 45

aagcggagac tctggtgc 18

<210> SEQ ID NO 46
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 46

gaagcgcctg gcagtgtacc 20

<210> SEQ ID NO 47
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 47

ctgcgcaagc tgcg 14

<210> SEQ ID NO 48
<211> LENGTH: 14
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 48

tcggcgccct cgcg 14

<210> SEQ ID NO 49
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 49

ggtacactgc caggcgcttc 20

-continued

<210> SEQ ID NO 50
<211> LENGTH: 12
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 50

ggatggcgct ga 12

<210> SEQ ID NO 51
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 51

gcctcgctc ccacc 15

<210> SEQ ID NO 52
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 52

gatgcgcttc cggaagacc 19

<210> SEQ ID NO 53
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 53

gcttcttgag cttctcagcg 20

<210> SEQ ID NO 54
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 54

ggtagcatgt tcaggcacca g 21

<210> SEQ ID NO 55
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 55

catccctagt accgaatccc atatagc 27

-continued

<210> SEQ ID NO 56
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 56

catcctccca cctaagcctt tcaa 24

<210> SEQ ID NO 57
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 57

ttgagttaat agcattataa caatttcac a 31

<210> SEQ ID NO 58
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 58

actccttaca accggaaggc aaac 24

<210> SEQ ID NO 59
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 59

tggacgtggt gaagcccgtg gtg 23

<210> SEQ ID NO 60
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 60

taggtctcta gggggcctct g 21

<210> SEQ ID NO 61
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 61

aggctgcca acagccccac t 21

<210> SEQ ID NO 62

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 62

tccccctgttg acctggagaa 20

<210> SEQ ID NO 63
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 63

cactgtactt gccctgacca 20

<210> SEQ ID NO 64
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 64

tgagatgtgg gcagaagg 19

<210> SEQ ID NO 65
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 65

ttggtgttga cagggagcaa 20

<210> SEQ ID NO 66
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 66

gtccaaaggc ccaagaacct 20

<210> SEQ ID NO 67
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 67

tgagaggga cagaaggct 20

<210> SEQ ID NO 68
<211> LENGTH: 22

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 68

gctcatctta atctgctcag cc 22

<210> SEQ ID NO 69
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 69

tcctagcact ttggaaggtc g 21

<210> SEQ ID NO 70
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 70

aaaggagcag ctcttctctgg 20

<210> SEQ ID NO 71
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 71

gtctgcacca tctcccacaa 20

<210> SEQ ID NO 72
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 72

gtgtggagag tgagtaagcc a 21

<210> SEQ ID NO 73
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 73

acggtaggat gatttcaggc a 21

<210> SEQ ID NO 74
<211> LENGTH: 26
<212> TYPE: DNA

-continued

```

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 74

ttttttggta ctcgagtgtt attcag                                26

<210> SEQ ID NO 75
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 75

cacaaagcag tgtagctcag g                                    21

<210> SEQ ID NO 76
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 76

ggcatggctt ctgagactca                                    20

<210> SEQ ID NO 77
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 77

cccccttgca cccctgtett t                                    21

<210> SEQ ID NO 78
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 78

gaagaggctg cccatgagag                                    20

<210> SEQ ID NO 79
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

<400> SEQUENCE: 79

tttgcaatg gaggcattgg                                    20

<210> SEQ ID NO 80
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 80

ggtctgaggc tcgaatcctg 20

<210> SEQ ID NO 81
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 81

ctgtggcctc catatccctg 20

<210> SEQ ID NO 82
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 82

tttccaccag aactcagccc 20

<210> SEQ ID NO 83
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 83

cctcggttcc tccacaacac 20

<210> SEQ ID NO 84
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 84

gcaggggagg gataaagcag 20

<210> SEQ ID NO 85
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 85

cacgggaagg acaggagaag 20

<210> SEQ ID NO 86
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 86

atgtgggctg cctagaaagg 20

<210> SEQ ID NO 87

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 87

cccagccaaa cttgtcaacc 20

<210> SEQ ID NO 88

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 88

ggacgagacc atgaaggagt tgaaggc 27

<210> SEQ ID NO 89

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 89

ccacctgctc cttcacctcg tccag 25

<210> SEQ ID NO 90

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 90

gacccacggc gtgcagtgt t 21

<210> SEQ ID NO 91

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 91

ggcccagact gagcacgtga 20

<210> SEQ ID NO 92

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 92

gaacacaaaag catagactgc 20

<210> SEQ ID NO 93
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 93

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 94
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 94

gaagcgcctg gcagtgtacc 20

<210> SEQ ID NO 95
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 95

gtggcagagc ggccagcgt 20

<210> SEQ ID NO 96
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 96

gcacccagac cgccaagctg a 21

<210> SEQ ID NO 97
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

<400> SEQUENCE: 97

gaccaggac tcttcctgc 20

<210> SEQ ID NO 98
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 oligonucleotide

-continued

<400> SEQUENCE: 98

gcaagcagag gctgaagctg a

21

<210> SEQ ID NO 99

<211> LENGTH: 18

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 99

gggtcttcga gaagacct

18

<210> SEQ ID NO 100

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 100

atgtgggctg cctagaaagg

20

<210> SEQ ID NO 101

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 101

cccagccaaa cttgtcaacc

20

<210> SEQ ID NO 102

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 102

ccagcccat ctgtcaaaact

20

<210> SEQ ID NO 103

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 103

tgaatggatt ccttggaac aatga

25

<210> SEQ ID NO 104

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

-continued

<400> SEQUENCE: 104

tggtcttctt tccctcccc tgccctcc

28

<210> SEQ ID NO 105

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 105

ggcctggagg cgggggctca gaga

24

<210> SEQ ID NO 106

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 106

ggacgagacc atgaaggagt tgaaggc

27

<210> SEQ ID NO 107

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 107

ccacctgctc cttcacctcg tccag

25

<210> SEQ ID NO 108

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 108

tcagaaggac cctgaccac ct

22

<210> SEQ ID NO 109

<211> LENGTH: 26

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 109

atgaaacctg gacctgggga ggtata

26

<210> SEQ ID NO 110

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 110

-continued

agctgtgacc ggcgcctacg 20

<210> SEQ ID NO 111
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 111

gggattctcc tccacgtcac 20

<210> SEQ ID NO 112
<211> LENGTH: 19
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 112

gatgcgcttc cggaagacc 19

<210> SEQ ID NO 113
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 113

gcttcttgag cttctcagcg 20

<210> SEQ ID NO 114
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 114

ggtagcatgt tcaggcacca g 21

<210> SEQ ID NO 115
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 115

catccctagt accgaatccc atatagc 27

<210> SEQ ID NO 116
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 116

-continued

catcctccca cctaagcctt tcaa 24

<210> SEQ ID NO 117
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 117

ttgagttaat agcattataa caatttcac a 31

<210> SEQ ID NO 118
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 118

gatacaccat aaaggggttt gactg 25

<210> SEQ ID NO 119
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 119

accatttccc cccaattcta ctc 23

<210> SEQ ID NO 120
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 120

catctgcatt ggcttgaaac atc 23

<210> SEQ ID NO 121
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 121

ttacaaaagt gctaaatgat gcacat 26

<210> SEQ ID NO 122
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 122

actcagtaaa gctcctcttc aac 23

-continued

<210> SEQ ID NO 123
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 123

ttttgcttag gtccactggg c 21

<210> SEQ ID NO 124
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 124

gttgatgggg tggttcagga 20

<210> SEQ ID NO 125
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 125

gttgatgagg tggttcagga 20

<210> SEQ ID NO 126
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 126

gaacacaaag catagactgc 20

<210> SEQ ID NO 127
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 127

gagtatgagg catagactgc 20

<210> SEQ ID NO 128
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 128

gatgagataa tgatgagtca 20

-continued

<210> SEQ ID NO 129
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 129

ggattgaccc aggccagggc 20

<210> SEQ ID NO 130
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 130

gtagaaaaag tatagactgc 20

<210> SEQ ID NO 131
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 131

gcttgaccaa tagccttgac a 21

<210> SEQ ID NO 132
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 132

gatatttgca ttgagatagt g 21

<210> SEQ ID NO 133
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 133

gaaaatagtt gaagaaaaag 20

<210> SEQ ID NO 134
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 134

gcacagaaga taccgagact g 21

-continued

<210> SEQ ID NO 135
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 135

gggtcttcga gaagacct 18

<210> SEQ ID NO 136
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 136

tccttggaac caatgataac aagac 25

<210> SEQ ID NO 137
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 137

ccagccccat ctgtcaaac 20

<210> SEQ ID NO 138
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 138

gcttatattc tagggagaca gacat 25

<210> SEQ ID NO 139
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 139

acctgaggtc agaagtttga ga 22

<210> SEQ ID NO 140
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 140

gtctgaggtc acacagtggg 20

<210> SEQ ID NO 141

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 141

agagcaggga ccacatctac 20

<210> SEQ ID NO 142
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 142

gccaaacttg tcaaccagta 20

<210> SEQ ID NO 143
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 143

atgtgggctg cctagaaagg 20

<210> SEQ ID NO 144
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 144

tccatttata tgaaatgttc agaaaaggca aat 33

<210> SEQ ID NO 145
<211> LENGTH: 32
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 145

gtaactatat gctctctgat tctcctatta gc 32

<210> SEQ ID NO 146
<211> LENGTH: 18
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 146

cctaccttcc cagggttt 18

<210> SEQ ID NO 147
<211> LENGTH: 23

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 147

aagaagtcct ggtatcttct atg 23

<210> SEQ ID NO 148
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 148

tcagacgttc cagaagcgag 20

<210> SEQ ID NO 149
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 149

gacaagaagg tgaaaaacgg ctg 23

<210> SEQ ID NO 150
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 150

gattcaaagc ataccagaga atagt 25

<210> SEQ ID NO 151
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 151

tcaaactagt atgcatttca acaac 25

<210> SEQ ID NO 152
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 152

gagtgtagct gtttttctca gggt 24

<210> SEQ ID NO 153
<211> LENGTH: 25
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 153

gaatacccaa ccataagaag aacag 25

<210> SEQ ID NO 154
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 154

gaacacaaaag catagactgc 20

<210> SEQ ID NO 155
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 155

gagtatgagg catagactgc 20

<210> SEQ ID NO 156
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 156

gatgagataa tgatgagtca 20

<210> SEQ ID NO 157
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 157

ggattgaccc aggccagggc 20

<210> SEQ ID NO 158
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 158

gcagtctata ctttttctac 20

<210> SEQ ID NO 159
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 159

cttgaccaat agccttgaca 20

<210> SEQ ID NO 160
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 160

atatttgcac tgagatagtg 20

<210> SEQ ID NO 161
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 161

gaaaatagtt gaagaaaaag 20

<210> SEQ ID NO 162
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 162

cacagaagat accgagactg 20

<210> SEQ ID NO 163
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 163

cagcattatc catttgctgc ca 22

<210> SEQ ID NO 164
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
primer

<400> SEQUENCE: 164

tggagacagc gagtctacag c 21

<210> SEQ ID NO 165
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 165

taacaccatt atagctgaag tgggg 25

<210> SEQ ID NO 166
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 166

tgagttacac acaagccagt taaattc 27

<210> SEQ ID NO 167
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 167

agggagtgga catgaggcga 20

<210> SEQ ID NO 168
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 168

cccaagagga agtcccaagg 20

<210> SEQ ID NO 169
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 169

ccttggaag agaaggggtc 20

<210> SEQ ID NO 170
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 170

gagataccgg aagctttgat gtaaga 26

<210> SEQ ID NO 171
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

```

    primer

<400> SEQUENCE: 171

cttggggaga aaggtccagg                20

<210> SEQ ID NO 172
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 172

caagcttttc ctctgggat gtaaaa          26

<210> SEQ ID NO 173
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 173

ctggcaagct gttctcacat g              21

<210> SEQ ID NO 174
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 174

gaggctgagg caggagtatg                20

<210> SEQ ID NO 175
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 175

gttttcagta gaagagtata taatacataa t  31

<210> SEQ ID NO 176
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

<400> SEQUENCE: 176

atattctcag cctaggcctg                20

<210> SEQ ID NO 177
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    primer

```

-continued

<400> SEQUENCE: 177

tggtggacat gggcgctta tt

22

<210> SEQ ID NO 178

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 178

ttcacctct ctggatggcg

20

<210> SEQ ID NO 179

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 179

gcaggaggag gcagtgaaag

20

<210> SEQ ID NO 180

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 180

cagagaaata acactctggc agctg

25

<210> SEQ ID NO 181

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 181

cagcatttat cagcagtat tgttattg

28

<210> SEQ ID NO 182

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 182

tcatttcgtg ttgtgcttta tcacttaaaa

30

<210> SEQ ID NO 183

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

-continued

<400> SEQUENCE: 183

gtgagcagta aacttaattg ttgatacaat aaatc	35
--	----

<210> SEQ ID NO 184

<211> LENGTH: 35

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 184

cttttagaat gaaagtgtgc atcttagtaa agaaa	35
--	----

<210> SEQ ID NO 185

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 185

gttcctcact gattctcagc agg	23
---------------------------	----

<210> SEQ ID NO 186

<211> LENGTH: 29

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 186

cacaaaaggg ataaatgctc tatccattt	29
---------------------------------	----

<210> SEQ ID NO 187

<211> LENGTH: 3439

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 187

tgcgcgcttt cggtgatgac ggtgaaaacc tctgacacat gcagctcccc gagacgggtca	60
--	----

cagcttgctc gtaagcggat gccgggagca gacaagcccc tcagggcgcg tcagcgggtg	120
---	-----

ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc	180
---	-----

accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc	240
---	-----

attcgccatt caggtcgcgc aactgttggg aagggcgata ggtgcgggcc tcttcgctat	300
---	-----

tacgccagct ggcgaaaagg ggatgtgctg caaggcgatt aagttgggta acgccagggt	360
---	-----

tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cgagggccta tttcccatga	420
---	-----

ttccttcata tttgcatata cgatacaagg ctgttagaga gataattgga attaatttga	480
---	-----

ctgtaaacac aaagatatta gtacaaaata cgtgacgtag aaagtaataa tttcttgggt	540
---	-----

agtttgagct tttaaaatta tgttttaaaa tggactatca tatgcttacc gtaacttgaa	600
---	-----

agtatttcga tttcttggtt ttatatatct tgtggaaagg acgaaacacc gttgatgggg	660
---	-----

tgggttcagga gtttttagagc tagaaatagc aagttaaaat aaggctagtc cgttatcaac	720
---	-----

ttgaaaaagt ggcaccgagc cggtgctttt ttgttttaga gctagaaata gcaagttaaa	780
---	-----

-continued

ataaggctag	tccgttttta	gcgcgtgcgc	caattctgca	gacagagagg	gcctatttcc	840
catgattcct	tcatatttgc	atatacgata	caaggctgtt	agagagataa	ttggaattaa	900
tttgactgta	aacacaaaga	tattagtaca	aaatacgtga	cgtagaaagt	aataatttct	960
tgggtagttt	gcagttttta	aattatgttt	taaaatggac	tatcatatgc	ttaccgtaac	1020
ttgaaagtat	ttcgatttct	tggttttata	tatcttgtgg	aaaggacgaa	acaccgggtc	1080
ttcgagaaga	cctgttttag	agctagaaat	agcaagttaa	aataaggcta	gtccgttatc	1140
aaactgaaaa	agtgccaccg	agtcggtgct	ttttttctag	agtcgacctg	caggcatgca	1200
agcttgccgt	aatcatgggc	atagctgttt	cctgtgtgaa	attgttatcc	gctcacaatt	1260
ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	ggggtgccta	atgagtgcgc	1320
taactcacat	taattgcgtt	gcgctcactg	cccgttttcc	agtcgggaaa	cctgtcgtgc	1380
cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcgtat	tgggcgtctc	1440
tccgcttctc	cgtcactga	ctcgtgcgc	tccgtcgttc	ggctgcggcg	agcggtatca	1500
gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataacgc	aggaaagaac	1560
atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgtt	gctggcggtt	1620
ttccataggc	tccgcccccc	tgacgagcat	cacaaaaatc	gacgctcaag	tcagaggtag	1680
cgaaccgccg	caggactata	aagataccag	gcgtttcccc	ctggaagctc	cctcgtgcgc	1740
tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	ttcgggaagc	1800
gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cgggtgtagg	cgttcgtccc	1860
aaagctgggt	gtgtgcacga	acccccggtt	cagcccgacc	gctgcgcctt	atccggtaac	1920
tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	agccactggc	1980
aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	gtggtggcct	2040
aactacggct	acactagaag	aacagtatct	ggtatctgcg	ctctgctgaa	gccagttacc	2100
ttcggaaaaa	gagttgtag	ctcttgatcc	ggcaaaaaaa	ccaccgctgg	tagcggtagg	2160
ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	agatcctttg	2220
atctttttcta	cgggggtctga	cgtcagtggt	aacgaaaaact	cacgttaagg	gatttttggtc	2280
atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	aagttttaaa	2340
tcaatctaaa	gtatatatga	gtaaaccttg	tctgacagtt	accaatgctt	aatcagtgag	2400
gcacctatct	cagcgatctg	tctatttctg	tcatccatag	ttgcctgact	ccccgtcgtg	2460
tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	gataccgcga	2520
gaccacgcct	caccggctcc	agattttatca	gcaataaaac	agccagccgg	aagggccgag	2580
cgcagaagtg	gtcctgcaac	tttatccgcc	tccatccagt	ctattaattg	ttgccgggaa	2640
gctagagtaa	gtagttcgcc	agttaatagt	ttgcgcaacg	ttgttgccat	tgctacaggc	2700
atcgtgggtg	cacgctcgtc	gtttggtatg	gcttcattca	gctccgggtc	ccaacgatca	2760
aggcgagtta	catgatcccc	catgttgtgc	aaaaaagcgg	ttagctcctt	cggtcctccg	2820
atcgttgtca	gaagtaagtt	ggccgcagtg	ttatcactca	tggttatggc	agcactgcat	2880
aattctctta	ctgtcatgcc	atccgtaaga	tgtttttctg	tgactggtga	gtactcaacc	2940
aagtcattct	gagaatagtg	tatgcggcga	ccgagttgct	cttgcccggc	gtcaatacgg	3000
gataataccg	cgccacatag	cagaacttta	aaagtgtcca	tcattggaaa	acgttcttcg	3060
gggcgaaaaa	tctcaaggat	cttaccgctg	ttgagatcca	gttcgatgta	accaactcgt	3120

-continued

```
gcacccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg agcaaaaaca 3180
ggaaggcaaa atgccgcaaa aaaggggaata agggcgacac ggaaatgttg aatactcata 3240
ctcttccttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat gagcggatac 3300
atatttgaat gtatttagaa aaataaacia ataggggttc cgcgacatt tccccgaaaa 3360
gtgccacctg acgtctaaga aaccattatt atcatgacat taacctataa aaataggcgt 3420
atcacgaggc cctttcgtc 3439
```

<210> SEQ ID NO 188

<211> LENGTH: 3439

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 188

```
tcgcgcggtt cggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacgggtca 60
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 120
ttggcgggtg tcggggcttg cttactatg cggcatcaga gcagattgta ctgagagtgc 180
accatatgcg gtgtgaaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc 240
attcgccatt caggctgcgc aactgttggg aagggcgatc ggtgcgggccc tcttcgctat 300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt 360
tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cgagggccta tttcccatga 420
ttccttcata tttgcatata cgatacaagg ctgttagaga gataattgga attaatttga 480
ctgtaaacac aaagatatta gtacaaaata cgtgacgtag aaagtaataa tttcttgggt 540
agtttgagct tttaaaatta tgttttaaaa tggactatca tatgcttacc gtaacttgaa 600
agtatttcga tttcttggct ttatatatct tgtggaaaagg acgaaacacc gttgatgagg 660
tgggttcagga gtttttagagc tagaaatagc aagttaaaat aaggctagtc cgttatcaac 720
ttgaaaaagt ggcaccgagt cggtgctttt ttgttttaga gctagaaaata gcaagttaaa 780
ataaggctag tccgttttta gcgcgtgcgc caattctgca gacagagagg gcctatttcc 840
catgattcct tcatatttgc atatacgata caaggctgtt agagagataa ttggaattaa 900
tttgactgta aacacaaaga tattagtaca aaatacgtga cgtagaaagt aataatttct 960
tgggtagttt gcagttttta aattatgttt taaaatggac tatcatatgc ttaccgtaac 1020
ttgaaagtat ttcgatttct tggttttata tatcttgtgg aaaggacgaa acaccgggtc 1080
ttcgagaaga cctgttttag agctagaaat agcaagttaa aataaggcta gtccgttatc 1140
aacttgaaaa agtggcaccg agtcggtgct ttttttctag agtcgacctg caggcatgca 1200
agcttggcgt aatcatggtc atagctgttt cctgtgtgaa attgttatcc gctcacaatt 1260
ccacacaaca tacgagccgg aagcataaag tgtaaaacct ggggtgccta atgagtgcgc 1320
taactcacat taattgcgtt gcgctcactg cccgctttcc agtcgggaaa cctgtcgtgc 1380
cagctgcatt aatgaatcgg ccaacgcgcg gggagaggcg gtttgcttat tgggcgctct 1440
tccgcttctc cgctcactga ctgcgtgcgc tcggtcggtc ggctgcggcg agcgggtatca 1500
gctcactcaa aggcggtaat acggttatcc acagaatcag gggataacgc aggaaagaac 1560
atgtgagcaa aaggccagca aaaggccagg aaccgtaaaa aggcgcgctt gctggcggtt 1620
ttocatagcc tccgcccccc tgacgagcat cacaaaaatc gacgctcaag tcagagggtg 1680
```

-continued

cgaaacccga caggactata aagataccag gcgtttcccc ctggaagctc cctcgtgcgc	1740
tctcctgttc cgaccctgcc gcttacccga tacctgtccg cctttctccc ttcgggaagc	1800
gtggcgcttt ctcatactc acgctgtagg tatctcagtt cgggtgtaggt cgttcgctcc	1860
aagctgggct gtgtgcacga acccccgtt cagcccgacc gctgcgcctt atccggtaac	1920
tatcgtcttg agtccaaccc ggtaagacac gacttatcgc cactggcagc agccactggt	1980
aacaggatta gcagagcgag gtatgtaggc ggtgctacag agttcttgaa gtgggtggcct	2040
aactacggct aactagaag aacagtattt ggtatctgcg ctctgctgaa gccagttacc	2100
ttcggaaaaa gagttgtag ctcttgatcc ggcaaaaaa ccaccgctgg tagcggtggt	2160
ttttttgttt gcaagcagca gattacgcgc agaaaaaag gatctcaaga agatcctttg	2220
atcttttcta cggggctctga cgctcagtg aacgaaaact cactgtaagg gattttggtc	2280
atgagattat caaaaaggat cttcacctag atccttttaa attaaaaatg aagttttaaa	2340
tcaatctaaa gtatatatga gtaaaacttg tctgacagtt accaatgctt aatcagtgag	2400
gcacctatct cagcgatctg tctatttctg tcatccatag ttgcctgact ccccgctgtg	2460
tagataacta cgataccgga gggcctacca tctggcccca gtgctgcaat gataccgcga	2520
gaccacgct caccggctcc agatttatca gcaataaacc agccagccgg aagggccgag	2580
cgcagaagtg gtcttgaac ttatccgcc tccatccagt ctattaattg ttgccgggaa	2640
gctagagtaa gtagtccgc agttaatagt ttgcgcaacg ttgttgccat tgctacaggc	2700
atcgtggtgt cagctcgtc gtttggtatg gcttcattca gctccggttc ccaacgatca	2760
aggcgagtta catgatcccc catgttggtc aaaaaagcgg ttagctcctt cggctcctcg	2820
atcgttgta gaagtaagtt ggcgcagtg ttatcactca tggttatggc agcaactgcat	2880
aattctctta ctgtcatgcc atccgtaaga tgcttttctg tgactggtga gtactcaacc	2940
aagtcattct gagaatagtg tatgcggcga ccgagttgct cttgccggc gtcaatacgg	3000
gataataccg cgccacatag cagaacttta aaagtgtca tcattggaaa acgtttctcg	3060
ggggcaaaa tctcaaggat cttaccgctg ttgagatcca gttcgatga acccaactgt	3120
gcaccaact gatcttcagc atcttttact ttcaccagcg tttctgggtg agcaaaaaca	3180
ggaaggcaaa atgcgcgaaa aaagggaata agggcgacac ggaaatgttg aatactcata	3240
ctcttctttt ttcaatatta ttgaagcatt tatcagggtt attgtctcat gagcggatac	3300
atatttgaat gtatttagaa aaataaaca ataggggttc cgcgcacatt tccccgaaa	3360
gtgccacctg acgtctaaga aaccattatt atcatgacat taacctataa aaataggcgt	3420
atcacgaggc cctttcgtc	3439

<210> SEQ ID NO 189

<211> LENGTH: 5849

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 189

gtcgacattg attattgact agatcatcgc gtgaggctcc ggtgccgctc agtgggcaga	60
gcgcacatcg cccacagtcc ccgagaagtt ggggggaggg gtcggcaatt gaaccggtgc	120
ctagagaagg tggcgcgggg taaactggga aagtgatgtc gtgtactggc tccgcctttt	180
tcccagggtt ggggggagaac cgtatataag tgcagtagtc gccgtgaacg ttctttttcg	240

-continued

caacgggttt gccgccagaa cacaggttaag tgccgtgtgt ggttcccgcg ggcctggcct	300
ctttacgggt tatggccctt gcgtgccttg aattacttcc acgcccctgg ctgcagtacg	360
tgattcttga tcccagagctt cgggttgga gtgggtggga gagttcgagg ccttgcgctt	420
aaggagcccc ttcgcctcgt gcttgagttg aggccctggc tgggcgctgg ggccgcgcg	480
tgcaaatctg gtggcacctt cgcgcctgtc tcgctgcttt cgataagtct ctageccattt	540
aaaatttttg atgacctgct gcgacgcttt ttttctggca agatagtctt gtaaagtcgg	600
gccaaagtct gcacactggg atttcggttt ttggggccgc gggcggcgac ggggcccgtg	660
cgtcccagcg cacatgttcg gcgaggcggg gcctgcgagc gcggccaccg agaatcggac	720
gggggtagtc tcaagctggc cggcctgctc tggtgccctg cctcgcgcgc cctgtatcgc	780
ccccgccctg ggccggcaagg ctggcccgtt cggcaccagt tgcgtgagcg gaaagatggc	840
cgttcccgcg ccctgctgca gggagctcaa aatggaggac gcggcgctcg ggagagcggg	900
cgggtgagtc acccacacaa aggaaaaggg cctttccgtc ctcagccgtc gcttcatgtg	960
actccacgga gtaccgggcg ccgtccaggc acctcgatta gttctcgagc ttttgagta	1020
cgtcgtcttt aggttggggg gaggggtttt atgcgatgga gtttcccac actgagtggg	1080
tgagactga agttaggcca gcttggcact tgatgtaatt ctccctggaa tttgcccttt	1140
ttgagtttgg atcttggttc attctcaagc ctcagacagt ggttcaaagt tttttcttc	1200
catttcaggt gtcgtgagga attctgcagt cgacgggtacc gcctacgcta gcgctaccgg	1260
tcgccaccat ggtgagcaag ggcgaggagg ataacatggc catcatcaag gagttcatgc	1320
gcttcaaggt gcacatggag ggctccgtga acggccacga gttcgagatc gagggcgagg	1380
gcgagggccg cccctacgag ggcaccaga ccgccaagct gaagggtgacc aagggtggcc	1440
ccctgccctt cgctgggac atcctgtccc ctcagttcat gtacggtccc aaggcctacg	1500
tgaagcacc cgccgacatc cccgactact tgaagctgtc cttcccagag ggcttcaagt	1560
gggagcgcgt gatgaacttc gaggaaggcg gcgtgggtgac cgtgaccag gactcctccc	1620
tcgaggacgg cgagttcatc tacaagggtga agctgcgcgg caccaacttc ccctccgacg	1680
gcccgcgaat gcagaagaag accatgggct gggaggcctc ctccgagcgg atgtaccccg	1740
aggacggcgc cctgaagggc gagatcaagc agaggctgaa gctgaaggac ggccggccact	1800
acgacgctga ggtcaagacc acctacaagg ccaagaagcc cgtgcagctg cccggcgctt	1860
acaacgtcaa catcaagttg gacatcacct cccacaacga ggactacacc atcgtggaac	1920
agtacgaacg cgccgagggc cgccactcca ccggcggcat ggacgagctg tacaagcccc	1980
gggagggcag aggaagtctt ctaacatcg gtgacgtgga ggagaatccc ggccctacta	2040
gttgatgggg tggttcagga ggggcatgct tgagcaaggc cgaggagctg ttcaccgggg	2100
tgggtcccat cctggctgag ctggacggcg acgtaaacgg ccacaagtcc agcgtgtccg	2160
gcgagggcga gggcgatgcc acctacggca agctgaccct gaagtctatc tgcaccaccg	2220
gcaagctgcc cgtgccctgg cccaccctcg tgaccacct gacctacggc gtgcagtgtc	2280
tcagccgcta ccccgaccac atgaagcagc acgacttctt caagtccgcc atgccgaag	2340
gtacgtcca ggagcgacc atcttcttca aggacgacgg caactacaag acccgcgccg	2400
aggtgaagtt cgagggcgac accctgggtga accgcatcga gctgaagggc atcgacttca	2460
aggaggacgg caacatcctg gggcacaagc tggagtacaa ctacaacagc cacaacgtct	2520
atatcatggc cgacaagcag aagaacggca tcaagggtgaa cttcaagatc cgccacaaca	2580
tcgaggacgg cagcgtgcag ctgcgcgacc actaccagca gaacaccccc atcggcgacg	2640

-continued

gccccgtgct gctgcccgcac aaccactacc tgagcaccca gtcgcacctg agcaaagacc	2700
ccaacgagaa gcgcgatcac atggctcctgc tggagttcgt gaccgcgcgc gggatcactc	2760
tcggcatgga cgagctgtac aagtaaagcg gccgcactcc tcagggtgcag gctgcctatc	2820
agaagggtggt ggctgggtgtg gccaatgccc tggctcacia ataccactga gatctttttc	2880
cctctgccaa aaattatggg gacatcatga agccccttga gcatctgact tctggctaata	2940
aaaggaaatt tattttcatt gcaatagtgt gttggaattt tttgtgtctc tcaactcgaa	3000
ggacatatgg gagggcaaat catttaaac atcagaatga gtatttggtt tagagtttgg	3060
caacatatgc catatgctgg ctgccatgaa caaagggtggc tataaagagg tcatcagtat	3120
atgaaacagc cccctgtgtt ccattcctta ttccatagaa aagccttgac ttgaggttag	3180
atttttttta tattttgttt tgtgttattt tttcttttaa catccctaaa attttcctta	3240
catgttttac tagccagatt ttctctctc tcctgactac tcccagtcac agctgtccct	3300
cttctcttat gaagatccct cgacctgcag cccaagcttg gcgtaatcat ggtcatagct	3360
gtttcctgtg tgaattgtt atccgctcac aattccacac aacatacgag ccggaagcat	3420
aaagtgtaaa gcctgggggtg cctaatgagt gagctaactc acattaattg cgttgcgctc	3480
actgcccgtt ttccagtcgg gaaacctgtc gtgccagcgg atccgcactc caattagtca	3540
gcaaccatag tcccgcacct aactccgcc atcccgcctc taactccgc cagttccgcc	3600
cattctccgc cccatggctg actaattttt tttatttatg cagaggccga ggccgcctcg	3660
gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg cttttgcaaa	3720
aagctaactt gtttattgca gcttataatg gttacaaata aagcaatagc atcacaatt	3780
tcacaaataa agcatTTTTT tcaactgcatt ctagtgtggg tttgtccaaa ctcatcaatg	3840
tatcttatca tgtctggatc cgtctgatta atgaatcggc caacgcgcgg ggagaggcgg	3900
tttgctattt gggcgctctt ccgcttctc gctcactgac tcgctgcgct cggctgctcg	3960
gctgcggcga gcggtatcag ctcaactcaa ggcggttaata cggttatcca cagaatcagg	4020
ggataacgca ggaaagaaca tgtgagcaaa agggccagca aaggccagga accgtaaaaa	4080
ggccgcgttg ctggcgTTTT tccataggct ccgccccct gacgagcatc acaaaaatcg	4140
acgctcaagt cagagggtggc gaaacccgac aggactataa agataccagg cgtttccccc	4200
tggaagctcc ctgctgcgct ctctctgttc gacctgccc cttaccggat acctgtccgc	4260
ctttctccct tcgggaagcg tggcgcttcc tcaatgtcca cgctgtaggt atctcagttc	4320
ggtgtaggtc gttcgtcca agctgggctg tgtgcacgaa cccccgttc agcccgaccg	4380
ctgcgcctta tccggttaact atcgtcttga gtccaacccg gtaagacacg acttatcgcc	4440
actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg gtgctacaga	4500
gttcttgaag tgggtggccta actacggcta cactagaagg acagtatttg gtatctgcgc	4560
tctgtgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg gaaaacaaac	4620
caccgctggt agcgggtggt tttttgttg caagcagcag attacgcgca gaaaaaaagg	4680
atctcaagaa gatcctttga tctttctac ggggtctgac gctcagtgga acgaaaaactc	4740
acgttaagggt attttgttca tgagattatc aaaaaggatc ttcacctaga tctttttaa	4800
ttaaaaatga agttttaaat caatctaaag tatatatgag taaacttgggt ctgacagtta	4860
ccaatgctta atcagtgagg cacctatctc agcgatctgt ctatttcggt catccatagt	4920
tgccctgactc cccgctgtgt agataactac gatacgggag ggcttaccat ctggccccag	4980

-continued

tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag caataaacca	5040
gccagccgga agggccgagc gcagaagtgg tcttgcaact ttatccgect ccatccagtc	5100
tattaattgt tgccgggaag ctgagtaag tagttcgcca gttaatagtt tgcgcaacgt	5160
tgttgccatt gctacaggca tctgtgtgtc acgctcgctg tttggtatgg ctteattcag	5220
ctccggttcc caacgatcaa ggcgagttac atgatcccc atgttgtgca aaaaagcgg	5280
tagctccttc ggtcctccga tctgtgtcag aagtaagttg gccgcagtgt tatcactcat	5340
ggttatggca gcaactgcata attctcttac tgtcatgcca tccgtaagat gcttttctgt	5400
gactggtagg tactcaacca agtcattctg agaatagttg atgcggcgac cgagttgctc	5460
ttgccggcgg tcaatacggg ataataccgc gccacatagc agaactttaa aagtgtctat	5520
cattggaaaa cgttcttcgg ggcgaaaact ctcaaggatc ttaccgctgt tgagatccag	5580
ttcgatgtaa cccactcgtg caccacaactg atcttcagca tcttttactt tcaccagcgt	5640
ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa aagggaataa gggcgacacg	5700
gaaatgttga atactcatac tcttctttt tcaatattat tgaagcattt atcagggtta	5760
ttgtctcatg agcggataca tatttgaatg tatttagaaa aataacaaa taggggttcc	5820
gcgcacattt ccccgaaaag tgccacctg	5849

<210> SEQ ID NO 190

<211> LENGTH: 5849

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 190

gtcgacattg attattgact agatcatcgc gtgaggctcc ggtgcccgtc agtgggcaga	60
gcgcacatcg cccacagtcc ccgagaagtt ggggggaggg gtcggcaatt gaaccggtgc	120
ctagagaagg tggcgcgagg taaactggga aagtgatgtc gtgtactggc tccgcctttt	180
tcccagggtt gggggagaac cgtatataag tgcagtagtc gccgtgaacg ttctttttcg	240
caacgggttt gccgccagaa cacaggtaag tgccgtgtgt ggttcccgcg ggccctggcct	300
ctttacgggt tatggccctt gcgtgccttg aattaactcc acgcccctgg ctgcagtacg	360
tgattcttga tcccagactt cgggttgga gtgggtggga gagttcgagg ccttgcgctt	420
aaggagcccc ttgcctcgt gcttgagttg aggcctggcc tgggcgctgg ggccgcgcg	480
tgcgaaatctg gtggcacctt cgcgcctgtc tcgctgcttt cgataagtct ctagccattt	540
aaaatttttg atgacctgct gcgacgcttt ttttctggca agatagtctt gtaaatgcgg	600
gccaaagatct gcacactggt atttcggttt ttggggccgc gggcggcgac ggggcccgtg	660
cgtcccagcg cacatgttcg gcgaggcggg gcctgcgagc gcggccaccg agaatcggac	720
gggggtagtc tcaagctggc cggcctgctc tgggtgcctgg cctcgcgcg ccgtgtatcg	780
ccccgccctg ggcggcaagg ctggcccggg cggcaccagt tgctgtagcg gaaagatggc	840
cgttccccgg ccctgctgca gggagctcaa aatggaggac gcggcgctcg ggagagcggg	900
cgggtgagtc acccacacaa aggaaaaggg cctttccgtc ctacgccgtc gcttcatgtg	960
actccacgga gtaccggggc ccgtccaggc acctcgatta gttctcgagc ttttgagta	1020
cgtcgtcttt aggttggggg gaggggtttt atgcgatgga gtttcccac actgagtggg	1080
tggagactga agttaggcca gcttggcact tgatgtaatt ctccctggaa tttgcccttt	1140

-continued

ttgagtttgg atcttgggttc attctcaagc ctcagacagt ggttcaaagt ttttttcttc	1200
catttcaggt gtcgtgagga attctgcagt cgacggtacc gcctacgcta gcgctaccgg	1260
tcgccaccat ggtgagcaag ggcgaggagg ataacatggc catcatcaag gagttcatgc	1320
gcttcaaggt gcacatggag ggctccgtga acggccacga gtctgagatc gagggcgagg	1380
gcgagggccg cccctacgag ggcacccaga ccgccaagct gaagggtgacc aagggtggcc	1440
ccctgccctt cgctgggac atcctgtccc ctcagttcat gtacggctcc aaggcctacg	1500
tgaagcacc cgccgacatc cccgactact tgaagctgtc cttccccgag ggcttcaagt	1560
gggagcgct gatgaacttc gaggacggcg gcgtggtgac cgtgaccag gactcctccc	1620
tcgaggacgg cgagttcatc tacaagggtga agctgcgcgg caccaacttc cctccgacg	1680
gccccgtaat gcagaagaag accatgggct gggaggcctc ctccgagcgg atgtaccccg	1740
aggacggcgc cctgaagggc gagatcaagc agaggctgaa gctgaaggac ggcgccact	1800
acgacgctga ggtcaagacc acctacaagg ccaagaagcc cgtgcagctg cccggcgct	1860
acaacgtcaa catcaagttg gacatcacct cccacaacga ggactacacc atcgtggaac	1920
agtacgaac cgccgagggc cgccactcca ccggcggcat ggacgagctg tacaagcccc	1980
gggagggcag aggaagtctt ctaacatgcg gtgacgtgga ggagaatccc ggccctacta	2040
gttgatgagg tggttcagga ggggcatgcg tgagcaaggc cgaggagctg ttcaccgggg	2100
tgggtcccat cctggtcgag ctggacggcg acgtaaacgg ccacaagttc agcgtgtccg	2160
gcgagggcga gggcgatgcc acctacggca agctgaccct gaagttcatc tgcaccaccg	2220
gcaagctgcc cgtgccctgg cccaccctcg tgaccaccct gacctacggc gtgcagtgtc	2280
tcagccgcta ccccgaccac atgaagcagc acgacttctt caagtccgcc atgccgaag	2340
gtacgtcca ggagcgacac atcttcttca aggacgacgg caactacaag acccgcgccg	2400
aggatgaatt cgagggcgac accctgggtga accgcacga gctgaagggc atcgacttca	2460
aggaggacgg caacatcctg gggcacaagc tggagtacaa ctacaacagc cacaacgtct	2520
atatcatggc cgacaagcag aagaacggca tcaagggtgaa cttcaagatc cgccacaaca	2580
tcgaggacgg cagcgtgcag ctgcgcgacc actaccagca gaacaccccc atcggcgacg	2640
gccccgtgct gctgcccgac aaccactacc tgagcaccca gtccgccctg agcaaagacc	2700
ccaacgagaa cgcgatcac atgggtcctgc tggagttcgt gaccgcgcgc gggatcactc	2760
tcggcatgga cgagctgtac aagtaaagcg gccgcactcc tcagggtgcag gctgcctatc	2820
agaagggtgtt ggctgggtgt gccaatgccc tgggtcacaa ataccactga gatctttttc	2880
cctctgcca aaattatggg gacatcatga agccccctga gcatctgact tctggctaat	2940
aaaggaaatt tattttcatt gcaatagtgt gttggaattt tttgtgtctc tactcggaa	3000
ggacatatgg gagggcaaat catttaaac atcagaatga gtatttgggt tagagtttgg	3060
caacatatgc catatgctgg ctgccatgaa caaagggtggc tataaagagg tcatcagtat	3120
atgaaacagc cccctgctgt ccattcctta ttccatagaa aagccttgac ttgaggttag	3180
atttttttta tattttgttt tgtgttattt ttttcttta catccctaaa attttcctta	3240
catgttttac tagccagatt tttcctctc tctgactac tcccagtc atgtgtccct	3300
cttctcttat gaagatccct cgacctgcag cccaagcttg gcgtaatcat ggtcatagct	3360
gtttcctgtg tgaaattgtt atccgctcac aattccacac aacatacgag ccggaagcat	3420
aaagtgtaaa gcctgggggtg cctaataagt gagctaactc acattaattg cgttgcgctc	3480
actgcccgtc ttccagtcgg gaaacctgtc gtgccagcgg atccgcatct caattagtca	3540

-continued

gcaaccatag tcccgccect aactccgccc atcccgccecc taactccgcc cagttccgcc	3600
cattctccgc cccatggctg actaattttt tttatttatg cagaggccga ggccgcctcg	3660
gcctctgagc tattccagaa gtagtgagga ggcttttttg gaggcctagg cttttgcaaa	3720
aagctaactt gtttattgca gcttataatg gttacaaata aagcaatagc atcacaaatt	3780
tcacaaataa agcatttttt tcaactgcatt ctagtgtggg tttgtccaaa ctcatcaatg	3840
tatcttatca tgtctggatc cgtctgatta atgaatcgcc caacgcgcgg ggagaggcgg	3900
tttgcgattt gggcgctctt ccgcttcttc gctcaactgac tcgctgcgct cggtcggtcg	3960
gctgcggcga gcggtatcag ctcaactcaa ggcgtaata cggttatcca cagaatcagg	4020
ggataacgca ggaagaaca tgtgagcaaa aggcagcaa aaggccagga accgtaaaaa	4080
ggccgcgctg ctggcgtttt tccataggct ccgccccctc gacgagcgc aaaaaatcg	4140
acgctcaagt cagagggtgg gaaaccgac aggactataa agataccagg cgtttccccc	4200
tggaagctcc ctctgctgct ctctctgttc gacctgccc cttaccggat acctgtccgc	4260
ctttctccct tcgggaagcg tggcgctttc tcaatgctca cgtgttaggt atctcagttc	4320
ggtgtaggtc gttcgctcca agctgggctg tgtgcacgaa cccccgttc agcccgaccg	4380
ctgcgcctta tccggttaact atcgtcttga gtccaaccgc gtaagacacg acttatcgcc	4440
actggcagca gccactggta acaggattag cagagcgagg tatgtaggcg gtgctacaga	4500
gttcttgaag tgggtggccta actacggcta cactagaagg acagtatttg gtatctgcgc	4560
tctgctgaag ccagttacct tcggaaaaag agttggtagc tcttgatccg gaaacaaac	4620
caccgctggt agcgggtggt tttttgttg caagcagcag attacgcgca gaaaaaagg	4680
atctcaagaa gatcctttga tctttttctac ggggtctgac gctcagtgga acgaaaaactc	4740
acgttaaggg attttggtca tgagattatc aaaaaggatc ttcacctaga tctttttaa	4800
ttaaaaaatga agttttaaat caatctaaag tatatatgag taaacttggt ctgacagtta	4860
ccaatgctta atcagtgagg cacctatctc agcgatctgt ctatttcgtt catccatagt	4920
tgccctgactc ccgctcggtg agataactac gatacgggag ggcttaccat ctggccccag	4980
tgctgcaatg ataccgcgag acccacgctc accggctcca gatttatcag caataaacca	5040
gccagccgga agggccgagc gcagaagtgg tcctgcaact ttatccgctc ccatccagtc	5100
tattaattgt tgccgggaag ctgagtaga tagttcgcca gttaatatgt tgcgcaacgt	5160
tgttgccatt gctacaggca tcgtggtgtc acgctcgctg tttggtaggt cttcattcag	5220
ctccggttcc caacgatcaa ggcgagttac atgatcccc atgttgtgca aaaaagcgg	5280
tagctccttc ggtcctccga tcgttgctcag aagtaagttg gccgcagtgt tatcactcat	5340
ggttatggca gcaactgcata attctcttac tgtcatgcca tccgtaagat gctttttctgt	5400
gactggtgag tactcaacca agtcattctg agaatagtgt atgcggcgac cgagttgctc	5460
ttgcccgcg tcaatacggg ataataccgc gccacatagc agaactttaa aagtgtctcat	5520
cattggaaaa cgttcttcgg ggcgaaaaact ctcaaggatc ttaccgctgt tgagatccag	5580
ttcgatgtaa cccactcgtg caccacaactg atcttcagca tcttttactt tcaccagcgt	5640
ttctgggtga gcaaaaacag gaaggcaaaa tgccgcaaaa aagggaataa gggcgacacg	5700
gaaatgttga atactcatat tcttcctttt tcaatattat tgaagcattt atcagggtta	5760
ttgtctcatg agcggataca tatttgaatg tatttagaaa aataaacaac taggggttcc	5820
gcgcacattt cccgaaaaag tgccacctg	5849

-continued

```

<210> SEQ ID NO 191
<211> LENGTH: 9319
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polynucleotide

<400> SEQUENCE: 191

gctgcttcgc gatgtacggg ccagatatac gcgttgacat tgattattga ctagtcgtga      60
ggctccgggtg cccgtcagtg ggcagagcgc acatcgccca cagtcccgga gaagttgggg      120
ggaggggtcgc gcaattgaac cgggtgcctag agaaggtggc gcggggtaaa ctgggaaagt      180
gatgtcgtgt actggctccg cctttttccc gaggggtggg gagaaccgta tataagtgca      240
gtagtcgcgc tgaacgttct ttttcgcaac gggtttgccg ccagaacaca ggtaagtgcc      300
gtgtgtggtt cccgcggggc tggcctcttt acgggttatg gcccttgctg gccttgaatt      360
acttccacgc ccctggctgc agtacgtgat tcttgatccc gagcttcggg ttggaagtgg      420
gtgggagagt tcgaggcctt gcgcttaagg agccccctcg cctcgtgctt gagttgaggc      480
ctggcctggg cgtcggggcc gccgcgtgcg aatctggtgg caccttcgcg cctgtctcgc      540
tgctttcgat aagtctctag ccatttaaaa tttttgatga cctgctgcga cgcttttttt      600
ctggcaagat agtcttgtaa atgcggggcca agatctgcac actggtattt cggttttttg      660
ggccgcgggc ggcgacgggg cccgtgcgtc ccagcgcaca tggtcggcga ggcggggcct      720
gcgagcgcgc ccacgcagaa tcggacgggg gtagtctcaa gctggccggc ctgctctggt      780
gcctggcctc gcgcgcgcgt gtatcgcccc gccctgggcg gcaaggctgg cccggtcggc      840
accagttcgc tgagcggaaa gatggcgcgt tcccggccct gctgcaggga gctcaaatg      900
gaggacgcgc cgctcgggag agcgggcggg tgagtcaccc acacaaagga aaagggcctt      960
tccgtcctca gccgtcgtt catgtgactc cacggagtac cgggcgcgt ccaggcacct      1020
cgattagtto tcgagctttt ggagtagctc gtcttttagt tggggggagg ggttttatgc      1080
gatggagttt cccacactg agtgggtgga gactgaagtt aggccagctt ggcacttgat      1140
gtaattctcc ttggaatttg ccttttttga gtttggtatc tggttcattc tcaagcctca      1200
gacagtgtt caaagttttt ttcttccatt tcaggtgtcg tgaggcggc gctaatacga      1260
ctcactatag ggagagccgc caccatgtcc gaagtcgagt tttccatga gtactggatg      1320
agacacgcat tgactctcgc aaagagggct tgggatgaac gcgaggtgcc cgtgggggca      1380
gtactcgtgc ataacaatcg cgtaatcggc gaaggttggga ataggccgat cggacgccac      1440
gacccactg cacatgcgga aatcatggcc cttcgacagg gagggcttgt gatgcagaat      1500
tatcgactta tcgatgcgc gctgtacgtc acgcttgaac cttgcgtaat gtgcgcggga      1560
gctatgattc actccgcac tggacgagtt gtattcgggtg cccgcgacgc caagacgggt      1620
gccgcaggtt cactgatgga cgtgctgcat caccaggca tgaaccaccg ggtagaaatc      1680
acagaaggca tattggcgga cgaatgtgcg gcgctgttgt ccgacttttt tcgcatgcgg      1740
aggcaggaga tcaaggccca gaaaaagca caatcctcta ctgactctgg tggttcttct      1800
ggtggttcta gcggcagcga gactcccggt acctcagagt ccgccacacc cgaaagttct      1860
ggtggttctt ctggtggttc ttccgaagtc gagttttccc atgagtactg gatgagacac      1920
gcattgactc tcgcaaagag ggctcgagat gaacgcgagg tgcccgtggg ggcagtactc      1980
gtgctcaaca atcgcgtaat cggcgaaggt tggaataggg caatcgact ccacgacccc      2040

```

-continued

actgcacatg	cggaaatcat	ggcccttcga	caggaggagg	ttgtgatgca	gaattatcga	2100
cttatcgatg	cgacgctgta	cgtcacgttt	gaaccttgcg	taatgtgcgc	gggagctatg	2160
attcactccc	gcattggacg	agttgtattc	ggtgttcgca	acgccaagac	gggtgccgca	2220
ggttcactga	tggacgtgct	gcattacca	ggcatgaacc	accgggtaga	aatcacagaa	2280
ggcatattgg	cggacgaatg	tgcggcgctg	ttgtgttact	tttttcgcat	gccaggcgag	2340
gtctttaacg	cccagaaaaa	agcacaatcc	tctactgact	ctggtggttc	ttctggtggt	2400
tctagcggca	gcgagactcc	cgggacctca	gagtcgcgca	caccgaaaag	ttctggtggt	2460
tcttctggtg	gttctgataa	aaagtattct	attggtttag	ccatcggcac	taattccgtt	2520
ggatgggctg	tcataacoga	tgaatacaaa	gtaccttcaa	agaaatttaa	ggtgtgtggg	2580
aacacagacc	gtcattcgat	taaaaagaat	cttatcggtg	ccctcctatt	cgatagtggc	2640
gaaacggcag	aggcgactcg	cctgaaacga	accgctcgga	gaagggtatac	acgtcgcaag	2700
aaccgaatat	gttacttaca	agaaattttt	agcaatgaga	tggccaaagt	tgacgattct	2760
ttctttcacc	gtttggaaga	gtccttcctt	gtcgaagagg	acaagaaaca	tgaacggcac	2820
cccatccttg	gaaacatagt	agatgaggtg	gcatatcatg	aaaagtaccc	aacgatttat	2880
cacctcagaa	aaaagctagt	tgactcaact	gataaagcgg	acctgagggt	aatctacttg	2940
gctcttgccc	atatgataaa	gttcggtggg	cactttctca	ttgagggtga	tctaaatccg	3000
gacaactcgg	atgtcgacaa	actgttcac	cagttagtag	aaacctataa	tcagttgttt	3060
gaagagaacc	ctataaatgc	aagtggcgtg	gatgcgaagg	ctattcttag	cggccgctc	3120
tctaaatccc	gacggctaga	aaacctgatc	gcacaattac	ccgagagaaa	gaaaaatggg	3180
ttgttcggta	accttatagc	gctctcacta	ggcctgacac	caaattttaa	gtcgaacttc	3240
gacttagctg	aagatgcaa	attgcagctt	agtaaggaca	cgtagatga	cgatctcgac	3300
aatctactgg	cacaaattgg	agatcagtat	gcggacttat	ttttggctgc	caaaaacctt	3360
agcgatgcaa	tcctcctatc	tgacatactg	agagttaata	ctgagattac	caaggcgccg	3420
ttatccgctt	caatgatcaa	aagggtacgat	gaacatcacc	aagacttgac	acttctcaag	3480
gccttagtcc	gtcagcaact	gcctgagaaa	tataaggaaa	tattctttga	tcagtcgaaa	3540
aacgggtacg	caggttatat	tgacggcgga	gcgagtcgaag	aggaattcta	caagtttatc	3600
aaacctatat	tagagaagat	ggatgggacg	gaagagttgc	ttgtaaaact	caatcgcgaa	3660
gatctactgc	gaaagcagcg	gactttcgac	aacggtagca	ttccacatca	aatccactta	3720
ggcgaattgc	atgtataact	tagaaggcag	gaggattttt	atccgttcct	caaagacaat	3780
cgtgaaaaga	ttgagaaaat	cctaaccttt	cgcatacctt	actatgtggg	acctctggcc	3840
cgagggaact	ctcggttcgc	atggatgaca	agaaagtccg	aagaaacgat	tactccatgg	3900
aattttgagg	aagttgtcga	taaagtgctg	tcagctcaat	cgttcatcga	gaggatgacc	3960
aactttgaca	agaatttacc	gaacgaaaaa	gtattgccta	agcacagttt	actttacgag	4020
tatttcacag	tgtacaatga	actcacgaaa	gttaagtatg	tactgagggg	catgcgtaaa	4080
ccgcctttc	taagcggaga	acagaagaaa	gcaatagtag	atctgttatt	caagaccaac	4140
cgcaaagtga	cagttaagca	attgaaagag	gactacttta	agaaaattga	atgcttcgat	4200
tctgtcgaga	tctccggggt	agaagatcga	tttaatgcgt	cacttggtac	gtatcatgac	4260
ctcctaaaga	taattaaaga	taaggacttc	ctggataacg	aagagaatga	agatatctta	4320
gaagatatag	tgttgactct	tacctctttt	gaagatcggg	aatgattga	ggaaagacta	4380
aaaacatacg	ctcacctggt	cgacgataag	gttatgaaac	agttaagag	gcgtcgctat	4440

-continued

acgggctggg gacgattgtc gcggaaactt atcaacggga taagagacaa gcaaagtgg	4500
aaaactattc tcgattttct aaagagcgac ggcttcgcca ataggaactt tatgcagctg	4560
atccatgatg actctttaac cttcaaagag gatatacaaa aggcacaggt ttccggacaa	4620
ggggactcat tgcaogaaca tattgcgaat cttgctggtt cgccagccat caaaaagggc	4680
atactccaga cagtcaaagt agtggatgag ctagttaagg tcatgggacg tcacaaaccg	4740
gaaaacattg taatcgagat ggcacgcgaa aatcaaacga ctcagaaggg gcaaaaaaac	4800
agtcgagagc ggatgaagag aatagaagag ggtattaaag aactgggcag ccagatctta	4860
aaggagcatc ctgtggaaaa tacccaattg cagaacgaga aactttacct ctattacct	4920
caaaatggaa gggacatgta tgttgatcag gaactggaca taaaccgttt atctgattac	4980
gacgtcgatc acattgtacc ccaatccttt ttgaaggacg attcaatcga caataaagt	5040
cttacacgct cggataagaa ccgagggaaa agtgacaatg ttccaagcga ggaagtcgta	5100
aagaaaatga agaactattg gcggcagctc ctaaatgcga aactgataac gcaaagaaag	5160
ttcgataact taactaaagc tgagaggggt ggcttgtctg aacttgacaa ggccggattt	5220
attaacgctc agctcgtgga aaccgcgcaa atcacaagc atgttgaca gatactagat	5280
tcccgaatga atacgaaata cgacgagaa c gataagctga ttccgggaagt caaagtaatc	5340
actttaaagt caaaattggt gtcggacttc agaaaggatt ttcaattcta taaagttagg	5400
gagataaata actaccacca tgcgcacgac gcttatctta atgccgtcgt agggaccgca	5460
ctcattaaga aatacccgaa gctagaaagt gagtttgtgt atggtgatta caaagtatat	5520
gacgtccgta agatgatcgc gaaaagcgaa caggagatag gcaaggctac agccaaatac	5580
ttcttttatt ctaacattat gaatttcttt aagacggaaa tcaactctggc aaacggagag	5640
atacgcaaac gacctttaat tgaaccaat ggggagacag gtgaaatcgt atgggataag	5700
ggccgggact tcgcgacggt gagaaaagtt ttgtccatgc cccaagtcaa catagtaaa	5760
aaaactgagg tgcagaccgg agggttttca aaggaatcga ttcttccaaa aaggaatagt	5820
gataagctca tcgctcgtaa aaaggactgg gacccgaaaa agtacgggtg cttcgatagc	5880
cctacagttg cctattctgt cctagtagtg gcaaaagtg agaagggaaa atccaagaaa	5940
ctgaagtcag tcaaagaatt attggggata acgattatgg agcgtcgtc tttgaaaag	6000
aaccccatcg acttccttga ggcgaaaggt tacaaggag taaaaagga tctcataatt	6060
aaactacca agtatagtct gtttgagtta gaaaatggc gaaaacggat gttggctagc	6120
gcgggagagc ttcaaaaggg gaacgaactc gcactaccgt ctaaatacgt gaatttctg	6180
tatttagcgt ccattacga gaagtgtgaa ggttcacctg aagataacga acagaagcaa	6240
ctttttgttg agcagcacia acattatctc gacgaaatca tagagcaaat ttcggaattc	6300
agtaagagag tcactcctagc tgatgccaat ctggacaaag tattaagcgc atacaacaag	6360
cacagggata aaccatcacg tgagcaggcg gaaaatatta tccatttgtt tactcttacc	6420
aaactcggcg ctccagcgc attcaagtat ttgacacaa cgatagatcg caaacgatac	6480
acttctacca aggaggtgct agacgcgaca ctgattcacc aatccatcac gggattatat	6540
gaaactcgga tagatttctc acagcttggg ggtgactctg gtggttctcc caagaagaag	6600
aggaaagtct aaccggtcat catcaccatc accattgagt ttaaacccgc tgatcagcct	6660
cgactgtgcc ttctagtctc cagccatctg ttgtttgccc ctccccctg ccttctctga	6720
ccctggaagg tgccactccc actgtccttt cctaataaaa tgaggaaatt gcatcgcat	6780

-continued

gtctgagtag	gtgtcattct	attctggggg	gtggggtggg	gcaggacagc	aagggggagg	6840
attggaaga	caatagcagg	catgctgggg	atgcggtggg	ctctatggct	tctgaggcgg	6900
aaagaaccag	ctggggctcg	ataccgtcga	cctctagcta	gagcttggcg	taatcatggt	6960
catagctgtt	tcctgtgtga	aattgttata	cgtcacaaat	tccacacaac	atacgagccg	7020
gaagcataaa	gtgtaaagcc	taggggtgcct	aatgagttag	ctaactcaca	ttaattgcgt	7080
tgcgctcact	gcccgttttc	cagtcgggaa	acctgtcgtg	ccagctgcac	taatgaatcg	7140
gccaacgcgc	ggggagaggc	ggtttgcgta	ttgggcgctc	ttccgcttcc	tcgctcactg	7200
actcgctcgc	ctcggtcggt	cggtcgcggc	gagcggtatc	agctcactca	aaggcggtaa	7260
tacggttatc	cacagaatca	ggggataacg	caggaaagaa	catgtgagca	aaaggccagc	7320
aaaaggccag	gaaccgtaaa	aaggccgcgt	tgtggcggtt	tttccatagg	ctccgcccc	7380
ctgacgagca	tcacaaaaat	cgacgctcaa	gtcagagggt	gcgaaccccg	acaggactat	7440
aaagatacca	ggcggtttccc	cctggaagct	ccctcggtcg	ctctcctggt	ccgacctgc	7500
cgttaccgg	atacctgtcc	gcctttctcc	cttcgggaag	cgtggcgctt	tctcatagct	7560
cacgctgtag	gtatctcagt	tcgggttagg	tcgttcgctc	caagctgggc	tgtgtgcacg	7620
aacccccgt	tcagcccgac	cgtgcgcct	tatccggtaa	ctatcgtctt	gagtcacaac	7680
cggtaagaca	cgacttatcg	ccactggcag	cagccactgg	taacaggatt	agcagagcga	7740
ggtatgtagg	cgggtctaca	gagttcttga	agtgggtggc	taactacggc	tacactagaa	7800
gaacagtatt	tggtatctgc	gctctgctga	agccagttac	cttcggaaaa	agagttggta	7860
gctcttgatc	cggcaaacaa	accaccgctg	gtagcgggtg	tttttttgtt	tgcaagcagc	7920
agattacgcg	cagaaaaaaa	ggatctcaag	aagatccttt	gatcttttct	acggggtctg	7980
acgctcagtg	gaacgaaaaa	tcacgttaag	ggattttggt	catgagatta	tcaaaaagga	8040
tcttcaccta	gatcctttta	aattaaaaat	gaagttttta	atcaatctaa	agtatatatg	8100
agtaaacttg	gtctgacagt	taccaatgct	taatcagtg	ggcacctatc	tcagcgatct	8160
gtctatttcg	ttcatccata	gttgctgac	tccccgtcgt	gtagataact	acgatacggg	8220
agggcttacc	atctggcccc	agtgtgcaa	tgataccgcg	agacccacgc	tcaccggctc	8280
cagatttata	agcaataaac	cagccagccg	gaaggggcga	gcgcagaagt	ggtcctgcaa	8340
ctttatccgc	ctccatccag	tctattaatt	gttgccggga	agctagagta	agtagttcgc	8400
cagttaatag	tttgcgcaac	gttggtgcca	ttgctacagg	catcgtgggtg	tcacgctcgt	8460
cgtttggtat	ggcttcattc	agtcccggtt	cccaacgata	aaggcgagtt	acatgatccc	8520
ccatgttgtg	caaaaaagcg	gttagctcct	tcggctcctc	gatcgttgtc	agaagtaagt	8580
tggccgcagt	gttatcactc	atgggtatgg	cagcaactgca	taattctctt	actgtcatgc	8640
catccgtaag	atgcttttct	gtgactgggtg	agtactcaac	caagtcattc	tgagaatagt	8700
gtatgcggcg	accgagttgc	tcttgcccg	cgtcaatacg	ggataatacc	gcgccacata	8760
gcagaacttt	aaaagtgtc	atcattggaa	aacgttcttc	ggggcgaaaa	ctctcaagga	8820
tcttaaccgt	gttgagatcc	agttcgatgt	aacccactcg	tgaccccaac	tgatcttcag	8880
catcttttac	tttcaccagc	gtttctgggt	gagcaaaaaa	aggaaggcaa	aatgccgcaa	8940
aaaagggaat	aagggcgaca	cggaaatgtt	gaatactcat	actcttcctt	tttcaatatt	9000
attgaagcat	ttatcagggt	tattgtctca	tgagcggata	catatttgaa	tgtatttaga	9060
aaaataaaca	aataggggtt	cgcgcacat	ttccccgaaa	agtgccacct	gacgtcgacg	9120
gatcgggaga	tcgatctccc	gatccctag	ggtcgactct	cagtacaatc	tgctctgatg	9180

-continued

```

ccgcatagtt aagccagtat ctgctccctg cttgtgtgtt ggaggtcgct gagtagtgcg 9240
cgagcaaaat ttaagctaca acaaggcaag gcttgaccga caattgcatg aagaatctgc 9300
ttagggtagt gcgttttgc 9319

```

```

<210> SEQ ID NO 192
<211> LENGTH: 9403
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
                        polynucleotide

```

```

<400> SEQUENCE: 192

```

```

gctgcttcgc gatgtacggg ccagatatac gcgttgacat tgattattga ctagtctgta 60
ggctccgggtg cccgtcagtg ggcagagcgc acatcgccca cagtocccga gaagtggggg 120
ggaggggtcg gcaattgaac cgggtgcctag agaagggtggc gcggggtaaa ctgggaaagt 180
gatgtcgtgt actggctccg cctttttccc gaggggtggg gagaaaccga tataagtga 240
gtagtcgccg tgaacgttct ttttcgcaac ggggttgccg ccagaacaca ggtaagtgcc 300
gtgtgtgtgt cccgcggggc tggcctcttt acgggttatg gcccttgctg gcttgaatt 360
acttccacgc ccctggctgc agtacgtgat tcttgatccc gagcttcggg ttggaagtgg 420
gtgggagagt tcgaggcctt gcgcttaagg agcccttcg cctcgtgctt gagttgaggc 480
ctggcctggg cgtgggggccc gccgcgtgcg aatctggtgg caccttcgcg cctgtctcgc 540
tgctttcgat aagtctctag ccatttaaaa tttttgatga cctgctgcga cgcttttttt 600
ctggcaagat agtctttaa atcggggcca agatctgcac actgggtattt cggttttttg 660
ggccgcggggc ggcgacgggg cccgtgcgtc ccagcgcaca tgttcggcga ggccggggcct 720
gcgagcgcgg ccaccagaaa tcggacgggg gtagtctcaa gctggccggc ctgctctggt 780
gcctggcctc gcgcgcgcgt gtatcgcccc gccctgggcg gcaaggctgg ccgggtcggc 840
accagttgct tgagcggaaa gatggccgct tcccggccct gctgcaggga gctcaaaatg 900
gaggacgcgg cgtcggggag agcggggggg tagtcaccc acacaaagga aaagggcctt 960
tccgtcctca gccgtcgtt catgtgactc cacggagtac cgggcgcgt ccaggcacct 1020
cgattagtct tcgagctttt ggagtacgtc gtcttttagt tggggggagg ggttttatgc 1080
gatggagttt cccacactg agtgggtgga gactgaagtt aggccagctt ggcacttgat 1140
gtaattctcc ttggaatttg ccttttttga gtttggtatc tggttcatc tcaagcctca 1200
gacagtgttt caaagttttt ttcttccatt tcaggtgtcg tgaggcggcc gctaatacga 1260
ctcactatag ggagagccgc caccatgaaa cggacagccg acggaagcga gttcagtgca 1320
ccaaagaaga agcggaaaagt ccttgaagtc gagtttagcc acgagtattg gatgaggcac 1380
gcactgaccc tggcaaaagc agcatgggat gaaagagaag tcccgtggg cgcctgctg 1440
gtgcacaaca atagagtgat cggagaggga tggaaacagg caatcggccg ccacgaccct 1500
accgcacacg cagagatcat ggcactgagg cagggaggcc tggtoatgca gaattaccgc 1560
ctgatcgatg ccaccctgta tgtgacactg gagccatgct tgatgtgcgc aggagcaatg 1620
atccacagca ggatcggaag agtgggtgtc ggagcacggg acgccaagac cggcgagca 1680
ggctccctga tggatgtgct gcaccacccc ggcataaacc accgggtgga gatcacagag 1740
ggaatcctgg cagacgagtg cgcgcctctg ctgagcgatt tctttagaat gcggagacag 1800
gagatcaagg ccagaagaa ggcacagagc tccaccgact ctggaggatc tagcggagga 1860

```

-continued

tcctctggaa gcgagacacc aggcacaagc gagtccgcca caccagagag ctccggcggc	1920
tcctccggag gatcctctga ggtggagttt tcccacgagt actggatgag acatgccctg	1980
accctggcca agagggcacg cgatgagagg gaggtgcctg tgggagccgt gctggtgctg	2040
aacaatagag tgatcggcga gggctggaac agagccatcg gcctgcacga cccaacagcc	2100
catgccgaaa ttatggccct gagacagggc ggcttggtca tgcagaacta cagactgatt	2160
gacgccacco tgtacgtgac attcagacct tgcgtgatgt gcgcggcgcg catgatccac	2220
tctaggatcg gcccggtggt gtttgcgctg aggaacgcaa aaaccggcgc cgcaggctcc	2280
ctgatggacg tgctgcacta ccccgcatg aatcacccgc tcgaaattac cgagggaatc	2340
ctggcagatg aatgtgccgc cctgctgtgc tatttcttcc ggatgcctag acaggtgttc	2400
aatgctcaga agaaggccca gagctccacc gactccggag gatctagcgg aggtccctct	2460
ggctctgaga cacctggcac aagcgagagc gcaacacctg aaagcagcgg gggcagcagc	2520
ggggggtcag acaagaagta cagcatcggc ctggccatcg gcaccaactc tgtgggctgg	2580
gccgtgatca ccgacgagta caaggtgcc agcaagaaat tcaaggtgct gggcaacacc	2640
gaccggcaca gcatcaagaa gaacctgatc ggagccctgc tgttcgacag cggcgaaaca	2700
gccgaggcca cccggctgaa gagaaccgcc agaagaagat acaccagacg gaagaaccgg	2760
atctgctatc tgcaagagat cttcagcaac gagatggcca aggtggacga cagcttcttc	2820
cacagactgg aagagtcctt cctggtggaa gaggataaga agcacgagcg gcaccccatc	2880
ttcggcaaca tcgtggacga ggtggcctac cacgagaagt accccaccat ctaccacctg	2940
agaaagaaac tgggtggacg caccgacaag gccgacctgc ggctgatcta tctggccctg	3000
gcccatcatg tcaagtcccg gggccacttc ctgatcgagg gcgacctgaa ccccgacaa	3060
agcgacgtgg acaagctgtt catccagctg gtgcagacct acaaccagct gttcgaggaa	3120
aaccccatca acgcccagcg cgtggacgcc aaggccatcc tgtctgccag actgagcaag	3180
agcagacggc tggaaaatct gatcgcccag ctgcccggcg agaagaagaa tggcctgttc	3240
ggaaacctga ttgcctgag cctgggcctg acccccaact tcaagagcaa cttcgacctg	3300
gccgaggatg ccaaactgca gctgagcaag gacacctacg acgacgacct ggacaacctg	3360
ctggcccaga tcggcgacca gtacgcccac ctgtttctgg ccgccaagaa cctgtccgac	3420
gccatcctgc tgagcgacat cctgagagtg aacaccgaga tcaccaaggc cccctgagc	3480
gcctctatga tcaagagata cgacgagcac caccaggacc tgacctgct gaaagctctc	3540
gtgcggcagc agctgcctga gaagtacaaa gagattttct tcgaccagag caagaacggc	3600
tacgccggct acattgacgg cggagccagc caggaagagt tctacaagtt catcaagccc	3660
atcctggaaa agatggacgg caccgaggaa ctgctcgtga agctgaacag agaggacctg	3720
ctgcggaagc agcggacctt cgacaacggc agcatcccc accagatcca cctgggagag	3780
ctgcacgcca ttctcggcg gcaggaagat ttttacctat tcctgaagga caaccgggaa	3840
aagatcgaga agatcctgac cttccgcatc cctactacg tgggccctct ggccagggga	3900
aacagcagat tcgcctggat gaccagaaa agcaggagaaa ccatcacccc ctggaacttc	3960
gaggaagtgg tggacaaggc cgcttccgcc cagagcttca tcgagcggat gaccaacttc	4020
gataagaacc tgcccaacga gaaggtgctg cccaagcaca gcctgctgta cgagtacttc	4080
accgtgtata acgagctgac caaagtgaat tacgtgaccg agggaaatgag aaagcccgcc	4140
ttcctgagcg gcgagcagaa aaaggccatc gtggacctgc tgttcaagac caaccggaaa	4200

-continued

gtgaccgtga agcagctgaa agaggactac ttcaagaaaa tctagtgctt cgactccgtg	4260
gaaatctccg gcgtggaaga tctgttcaac gctccctgg gcacatacca cgatctgctg	4320
aaaattatca aggacaagga cttcctggac aatgaggaaa acgaggacat tctggaagat	4380
atcgtgctga cctgacact gtttgaggac agagagatga tctaggaacg gctgaaaacc	4440
tatgcccacc tgttcgacga caaagtgatg aagcagctga agcggcggag atacaccggc	4500
tggggcaggc tgagccgga gctgatcaac ggcacccggg acaagcagtc cggcaagaca	4560
atcctggatt tcctgaagtc cgacggcttc gccaacagaa acttcatgca gctgatccac	4620
gacgacagcc tgaccttaa agaggacatc cagaaagccc aggtgtccgg ccaggggcat	4680
agcctgcacg agcacattgc caatctggcc ggcagccccg ccattaagaa gggcatcctg	4740
cagacagtga aggtggtgga cgagctcgtg aaagtgatgg gccggcacia gcccgagaac	4800
atcgtgatcg aaatggccag agagaaccag accaccaga agggacagaa gaacagccgc	4860
gagagaatga agcggatcga agagggtatc aaagagctgg gcagccagat cctgaaagaa	4920
caccccgtag aaaacaccca gctgcagaa gagaagctgt acctgtacta cctgcagaat	4980
ggcggggata tgtacgtgga ccaggaaactg gacatcaacc ggctgtccga ctacgatgtg	5040
gaccatatcg tgctcagag ctttctgaag gacgactcca tgcacaacia ggtgtgacc	5100
agaagcgaca agaaccgggg caagagcgac aacgtgcctt ccgaagaggt cgtgaagaag	5160
atgaagaact actggcgga gctgtgtaac gccaaagtga ttaccagag aaagtctgac	5220
aatctgacca aggcgagag aggcggcctg agcgaactgg ataaggccgg cttcatcaag	5280
agacagctgg tggaaacccg gcagatcaca aagcacgtgg cacagatcct ggactcccgg	5340
atgaacacta agtacgacga gaatgacaag ctgatccggg aagtgaagat gatcacccctg	5400
aagtccaagc tgggtgtccga tttccggaag gatttccagt tttacaaagt gcgcgagatc	5460
aacaactacc accacgcccc cgacgcctac ctgaacgccc tctgtgggaa cgccctgatc	5520
aaaaagtacc ctaagctgga aagcgagttc gtgtacggcg actacaaggt gtacgacgtg	5580
cgggaagtga tgcgaagag cgagcaggaa atcggcaagg ctaccgcaa gtactttctc	5640
tacagcaaca tcatgaactt tttcaagacc gagattaccc tggccaacgg cgagatccgg	5700
aagcggcctc tgatcgagac aaacggcgaa accggggaga tctgtgtgga taagggccgg	5760
gattttgccg ccgtgcggaa agtgtgagc atgccccaa tgaatatcgt gaaaaagacc	5820
gaggtgcaga caggcggtt cagcaagag tctatcctgc ccaagaggaa cagcgataag	5880
ctgatcgcca gaaagaagga ctgggaccct aagaagtacg gcggtctga cagccccacc	5940
gtggcctatt ctgtgctggt ggtggccaaa gtggaaaagg gcaagtccaa gaaactgaag	6000
agtgtgaaag agctgctggg gatcaccatc atggaaagaa gcagcttcca gaagaatccc	6060
atcgactttc tggaaagcaa gggctacaaa gaagtgaaaa aggacctgat catcaagctg	6120
cctaagtact ccctgttcca gctggaaaac ggcggaaga gaatgtggc ctctgccggc	6180
gaactgcaga agggaaacga actggccctg ccctccaaat atgtgaactt cctgtacctg	6240
gccagccact atgagaagct gaagggtcc cccgaggata atgagcagaa acagctgttt	6300
gtggaacagc acaagcacta cctggacgag atcatcgagc agatcagcga gttctccaag	6360
agagtgatcc tggccgagc taatctggac aaagtgtgtt ccgcctacaa caagcaccgg	6420
gataagccca tcagagagca ggcgagaaat atcatccacc tgtttacctt gaccaatctg	6480
ggagccctg ccgccttcaa gtactttgac accaccatcg accggaagag gtacaccagc	6540
accaaagagg tgctggagc caccctgatc caccagagca tcaccggcct gtacgagaca	6600

-continued

cggtatcgacc	tgtctcagct	gggaggtgac	tctggcggct	caaaaagaac	cgccgacggc	6660
agcgaattcg	agcccaagaa	gaagaggaaa	gtetaaccgg	tcatacatcac	catcaccatt	6720
gagtttaaac	cgcgtgatca	gcctcgactg	tgcccttctag	ttgccagcca	tctgttgttt	6780
gccccctccc	cgtgccttcc	ttgacctg	aagggtgccac	tcccactgtc	ctttcctaata	6840
aaaatgagga	aattgcatcg	cattgtctga	gtaggtgtca	ttctattctg	gggggtgggg	6900
tggggcagga	cagcaagggg	gaggattggg	aagacaatag	caggcatgct	ggggatgcgg	6960
tgggctctat	ggcttctgag	gcggaaagaa	ccagctgggg	ctcgataccg	tcgacctcta	7020
gctagagctt	ggcgtaatca	tggtcatagc	tgtttctctg	gtgaaattgt	tatccgctca	7080
caattccaca	caacatacga	gccggaagca	taaagtgtaa	agcctagggt	gcctaatagag	7140
tgagctaact	cacattaatt	gcgttgcgct	cactgcccgc	tttccagtcg	ggaacacctgt	7200
cgtgccagct	gcattaatga	atcggccaac	gcgcggggag	aggcggtttg	cgtattgggc	7260
gctcttccgc	ttcctcgctc	actgactcgc	tcgcctcggg	cggtcggctg	cggcgagcgg	7320
tatcagctca	ctcaaaggcg	gtaatacggg	tatccacaga	atcaggggat	aacgcaggaa	7380
agaacatgtg	agcaaaaggc	cagcaaaagg	ccaggaaccg	taaaaaggcc	gcgttgctgg	7440
cgtttttcca	taggctccgc	ccccctgacg	agcatcaca	aaatcgacgc	tcaagtcaga	7500
ggtggcgaaa	cccacagga	ctataaagat	accaggcggt	tccccctgga	agtcacctcg	7560
tgcgctctcc	tgttccgacc	ctgccgctta	ccggatacct	gtccgccttt	ctcccttcgg	7620
gaagcgtggc	gctttctcat	agctcacgct	gtaggtatct	cagttcgggtg	taggtcggtc	7680
gtccaagct	gggctgtgtg	cacgaacccc	cggttcagcc	cgaccgctgc	gccttatccg	7740
gtaactatcg	tcttgagtcc	aacccggtaa	gacacgactt	atcgccactg	gcagcagcca	7800
ctggtaacag	gattagcaga	gcgaggtatg	taggcgggtc	tacagagtcc	ttgaagtggg	7860
ggcctaacta	cggtacact	agaagaacag	tatttggtat	ctgcgctctg	ctgaagccag	7920
ttaccttcgg	aaaaagagtt	ggtagctctt	gatccggcaa	acaaccacc	gctggtagcg	7980
gtgggttttt	tgtttgcaag	cagcagatta	cgcgcagaaa	aaaaggatct	caagaagatc	8040
ctttgatctt	ttctacgggg	tctgacgctc	agtggaaacga	aaactcacgt	taagggattt	8100
tggatcatgag	attatcaaaa	aggatcttca	cctagatcct	tttaaattaa	aatgaagtt	8160
ttaaatcaat	ctaaagtata	tatgagtaaa	cttggtctga	cagttacca	tgcttaataca	8220
gtgaggcacc	tatctcagcg	atctgtctat	ttcgttcatc	catagttgcc	tgactccccg	8280
tcgtgtagat	aactacgata	cgggagggct	taccatctgg	cccagtgct	gcaatgatac	8340
cgcgagaccc	acgctcaccc	gctccagatt	tatcagcaat	aaaccagcca	gccggaaggg	8400
ccgagcgcag	aagtggtcct	gcaactttat	ccgcctccat	ccagtctatt	aattgttgcc	8460
gggaagctag	agtaagtagt	tcgccagtta	atagtttgcg	caacgttggt	gccattgcta	8520
caggcatcgt	ggtgtcacgc	tcgtcgtttg	gtatggcttc	attcagctcc	ggttcccaac	8580
gatcaaggcg	agttacatga	tcccccatgt	tgtagcaaaa	agcgggttagc	tccttcgggtc	8640
ctccgatcgt	tgtagaaggt	aagttggccg	cagtggtatc	actcatgggt	atggcagcac	8700
tgcataatcc	tcttactgtc	atgccatccg	taagatgctt	ttctgtgact	ggtgagtact	8760
caaccaagtc	attctgagaa	tagtgtatgc	ggcgaccgag	ttgctcttgc	ccggcgtcaa	8820
tacgggataa	taccgcgcga	catagcagaa	ctttaaaagt	gctcatcatt	ggaaaacgtt	8880
cttcggggcg	aaaactctca	aggatcttac	cgtgttgtag	atccagttcg	atgtaaccca	8940

-continued

```

ctcgtgcacc caactgatct tcagcatctt ttactttcac cagcgtttct gggtagagcaa 9000
aacaggaag gcaaaatgcc gcaaaaaagg gaataagggc gacacggaaa tgttgaatac 9060
tcatactctt cctttttcaa tattattgaa gcatttatca gggttattgt ctcagtagcg 9120
gatacatatt tgaatgtatt tagaaaaata aacaaatagg ggttcgcgc acatttcccc 9180
gaaaagtgcc acctgacgtc gacggatcgg gagatcgatc tcccgatccc ctagggtcga 9240
ctctcagtac aatctgctct gatgccgcat agttaagcca gtatctgctc cctgcttggtg 9300
tgttggaggt cgctgagtag tgcgcgagca aaatttaagc tacaacaagg caaggcttga 9360
cgcacaattg catgaagaat ctgcttaggg ttaggcgttt tgc 9403

```

```

<210> SEQ ID NO 193
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      peptide

```

```

<400> SEQUENCE: 193

```

```

Leu Thr His Gly Val Gln Cys Phe Gly Arg
1           5           10

```

```

<210> SEQ ID NO 194
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 194

```

```

ctgacccacg gcgtgcagtg cttcggccgc 30

```

```

<210> SEQ ID NO 195
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      peptide

```

```

<400> SEQUENCE: 195

```

```

Leu Thr Tyr Gly Val Gln Cys Phe Gly Arg
1           5           10

```

```

<210> SEQ ID NO 196
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 196

```

```

ctgacctacg gcgtgcagtg cttcggccgc 30

```

```

<210> SEQ ID NO 197
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

-continued

<400> SEQUENCE: 197

cctgacccac ggcgtgcagt gcttcggcc

29

<210> SEQ ID NO 198

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 198

cgaaacaccg ggtcttcgag aagacctgtt ttagag

36

<210> SEQ ID NO 199

<211> LENGTH: 36

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 199

ctctaaaaca ggtcttctcg aagaccgggt gtttcg

36

<210> SEQ ID NO 200

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 200

agttgatggg gtggttcagg agggg

25

<210> SEQ ID NO 201

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 201

agttgatgag gtggttcagg agggg

25

<210> SEQ ID NO 202

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 202

agttggtggg gtggttcagg agggg

25

<210> SEQ ID NO 203

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 203

-continued

agttggtggg gtggttcagg agggg 25

<210> SEQ ID NO 204
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 204

ggcccagact gagcacgtga 20

<210> SEQ ID NO 205
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 205

ggcccagact gagcacgtga 20

<210> SEQ ID NO 206
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 206

gaacacaaaag catagactgc 20

<210> SEQ ID NO 207
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 207

gaacacaaaag catagactgc 20

<210> SEQ ID NO 208
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 208

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 209
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 209

-continued

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 210
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 210

gaagcgctg gcagtgtacc 20

<210> SEQ ID NO 211
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 211

gaagcgctg gcagtgtacc 20

<210> SEQ ID NO 212
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 212

ggcccagact gagcacgtga 20

<210> SEQ ID NO 213
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 213

ggcccagact gagcacgtga 20

<210> SEQ ID NO 214
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 214

gaacacaaag catagactgc 20

<210> SEQ ID NO 215
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 215

gaacacaaag catagactgc 20

-continued

<210> SEQ ID NO 216
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 216

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 217
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 217

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 218
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 218

ccctacatcg tgcagtgctt 20

<210> SEQ ID NO 219
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 219

ccccaagtag tgcagtgctt 20

<210> SEQ ID NO 220
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 220

aaccaagatg tgcagtgctt 20

<210> SEQ ID NO 221
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 221

aaccagcgcc tgcagtgctt 20

-continued

<210> SEQ ID NO 222
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 222

ccccatggct tgctgtgctt 20

<210> SEQ ID NO 223
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 223

caccagact gagcacgtgc 20

<210> SEQ ID NO 224
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 224

gacacagacc gggcacgtga 20

<210> SEQ ID NO 225
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 225

agctcagact gagcaagtga 20

<210> SEQ ID NO 226
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 226

agaccagact gagcaagaga 20

<210> SEQ ID NO 227
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 227

gagccagaat gagcacgtga 20

-continued

<210> SEQ ID NO 228
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 228

gaacacaatg catagattgc 20

<210> SEQ ID NO 229
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 229

gcagtctatg ctttatgttt 20

<210> SEQ ID NO 230
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 230

tgcactgcgg ccggaggagg 20

<210> SEQ ID NO 231
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 231

ggctctgcgg ctggaggggg 20

<210> SEQ ID NO 232
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 232

ggcacgacgg ctggaggtgg 20

<210> SEQ ID NO 233
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 233

ggcatcacgg ctggaggtgg 20

<210> SEQ ID NO 234

-continued

<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 234

ggcgctgcgg cgggaggtgg 20

<210> SEQ ID NO 235
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 235

ggcccagact gagcacgtga 20

<210> SEQ ID NO 236
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 236

ggcccagact gagcacgtga 20

<210> SEQ ID NO 237
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 237

gaagcgcctg gcagtgtacc 20

<210> SEQ ID NO 238
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 238

gaagcgcctg gcagtgtacc 20

<210> SEQ ID NO 239
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 239

ggcccagact gagcacgtga 20

<210> SEQ ID NO 240
<211> LENGTH: 20

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 240

gaacacaaag catagactgc 20

<210> SEQ ID NO 241
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 241

ggcactgcgg ctggaggtgg 20

<210> SEQ ID NO 242
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 242

gtgatcatca tcacc 15

<210> SEQ ID NO 243
<211> LENGTH: 15
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 243

tggactgagt ggctc 15

<210> SEQ ID NO 244
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 244

gtagaggagg tggttcagga 20

<210> SEQ ID NO 245
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 245

ttagagcagg tggttcagga 20

<210> SEQ ID NO 246
<211> LENGTH: 20
<212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 246

caatacaaaag gatagactgc 20

<210> SEQ ID NO 247
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 247

ggagagagag catagactgc 20

<210> SEQ ID NO 248
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 248

taagacacaa tgatgagtca 20

<210> SEQ ID NO 249
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 249

ctggagatat tgatgagtca 20

<210> SEQ ID NO 250
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 250

aggttgcccc aggccagggc 20

<210> SEQ ID NO 251
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 251

cagtggaccc aggccagggc 20

<210> SEQ ID NO 252
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 252

tttgatgagg aggttcagga 20

<210> SEQ ID NO 253
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 253

ccaaacaaaa catagactgc 20

<210> SEQ ID NO 254
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 254

ggtgagagaa agatgagtca 20

<210> SEQ ID NO 255
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 255

ttgtggaccc aggccagggc 20

<210> SEQ ID NO 256
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 256

gatgagataa tgatgagtca 20

<210> SEQ ID NO 257
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 257

cacagaagat accgagactg 20

<210> SEQ ID NO 258
<211> LENGTH: 720
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 258

atggtgagca agggcgagga gctgttcacc ggggtggtgc ccatcctggt cgagctggac	60
ggcgacgtaa acggccacaa gtccagcgtg tccggcgagg gcgagggcga tgccacctac	120
ggcaagctga ccttgaagtt catctgcacc accggcaagc tgcccgtgcc ctggcccacc	180
ctcgtgacca ccttgaccca cggcgtgcag tgcttcggcc gctaccccga ccacatgaag	240
cagcacgact tcttcaagtc cgccatgccc gaaggctacg tccaggagcg caccatcttc	300
ttcaaggacg acggcaacta caagaccgc gccaggtga agttcgaggg cgacacctg	360
gtgaaccgca tcgagctgaa gggcatcgac ttcaaggagg acggcaacat cctggggcac	420
aaagtggagt acaactacaa cagccacaac gtctatatca tggccgacaa gcagaagaac	480
ggcatcaagg tgaacttcaa gatccgccac aacatcgagg acggcagcgt gcagctcgcc	540
gaccactacc agcagaacac ccccatcggc gacggccccg tgctgctgcc cgacaaccac	600
tacctgagca ccagtcgcg cctgagcaaa gaccccaacg agaagcgcga tcacatggtc	660
ctgctggagt tcgtgaccgc cgccgggatc actctcgga tggacgagct gtacaagtaa	720

<210> SEQ ID NO 259

<211> LENGTH: 1518

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polynucleotide

<400> SEQUENCE: 259

atggtgagca agggcgagga ggataacatg gccatcatca aggagttcat gcgcttcaag	60
gtgcacatgg agggctccgt gaacggccac gaggctcgaga tcgagggcga gggcgagggc	120
cggccctacg agggcaccca gaccgccaag ctgaaggatga ccaaggggtg cccctgccc	180
ttcgccctggg acatcctgtc cctcagttc atgtacggct ccaaggccta cgtgaagcac	240
cccgccgaca tccccgacta cttgaagctg tccttccccg agggcttcaa gtgggagcgc	300
gtgatgaact tcgaggacgg cggcgtggtg accgtgaccc aggactcctc cctgcaggac	360
ggcgagtcca tctacaaggt gaagctgcgc ggcaccaact tccccccga cggccccgta	420
atgcagaaga agaccatggg ctgggaggcc tcctccgagc ggatgtaccc cgaggacggc	480
gcccctgaagg gcgagatcaa gcagaggctg aagctgaagg acggcggcca ctacgacgct	540
gaggtaaga ccacctacaa ggccaagaag cccgtgcagc tgcccggcgc ctacaacgtc	600
aacatcaagt tggacatcac ctcccacaac gaggactaca ccatcgtgga acagtaacga	660
cgcgccgagg gccgccatc caccggcggc atggacgagc tgtacaagcc ccgggagggc	720
agagggaatc ttctaacatg cggtagcgtg gaggagaatc ccggccctac tagttgatgg	780
ggtggttcag gaggggcatg cgtgagcaag ggcgaggagc tggtcaccgg ggtggtgcc	840
atcctggctg agctggacgg cgacgtaaac ggccacaagt tcagcgtgtc cggcgagggc	900
gagggcgatg ccacctacgg caagctgacc ctgaagttca tctgcaccac cggcaagctg	960
cccgtgcctt ggcccaccct cgtgaccacc ctgacctacg gcgtgcagtg ctteagccgc	1020
taccocgacc acatgaagca gcacgacttc ttcaagtccg ccatgccoga aggtacgtc	1080
caggagcgca ccatcttctt caaggacgac ggcaactaca agaccgcgc cgaggtgaag	1140
ttcgagggcg acaccctggt gaaccgcac gagctgaagg gcacgactt caaggaggac	1200

-continued

```

ggcaacatcc tggggcacaa gctggagtag aactacaaca gccacaacgt ctatatcatg 1260
gccgacaagc agaagaacgg catcaagggt aacttcaaga tccgccacaa catcgaggac 1320
ggcagcgtgc agctcgccga ccactaccag cagaacaccc ccacggcga cggccccgtg 1380
ctgctgcccc acaaccacta cctgagcacc cagtccgccc tgagcaaaga ccccaacgag 1440
aagcgcgatc acatgggtcct gctggaggtc gtgaccgccc cggggatcac tctcgcatg 1500
gacgagctgt acaagtaa 1518

```

```

<210> SEQ ID NO 260
<211> LENGTH: 1518
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

```

```

<400> SEQUENCE: 260

```

```

atggtgagca agggcgagga ggataacatg gccatcatca aggagttcat gcgcttcaag 60
gtgcacatgg agggctccgt gaacggccac gaggtcgaga tcgagggcga gggcgagggc 120
cgccctacg agggcaccca gaccgccaag ctgaagggtg ccaagggtgg cccctgccc 180
ttcgctggg acatcctgtc ccctcagttc atgtacggct ccaaggccta cgtgaagcac 240
ccgcccgaca tccccgacta cttgaagctg tccttccccg agggcttcaa gtgggagcgc 300
gtgatgaact tcgaggacgg cggcgtggtg accgtgaccc aggactcctc cctgcaggac 360
ggcgagttca tctacaaggt gaagctgcgc ggcaccaact tcccctccga cggccccgta 420
atgcagaaga agaccatggg ctgggaggcc tcctccgagc ggatgtaccc caggagcggc 480
gcccctgaagg gcgagatcaa gcagaggctg aagctgaagg acggcgggca ctacgacgct 540
gaggtcaaga ccacctacaa ggccaagaag cccgtgcagc tgcccggcgc ctacaacgtc 600
aacatcaagt tggacatcac ctcccacaac gaggactaca ccacgtgga acagtacgaa 660
cgcgcccagg gccgcactc caccggcggc atggacgagc tgtacaagcc cgggagggc 720
agaggaagtc ttctaacatg cggtagcgtg gaggagaatc ccggccctac tagttgatga 780
ggtggttcag gaggggcatg cgtgagcaag ggcgaggagc tgttcaccgg ggtggtgccc 840
atcctggtcg agctggacgg cgacgtaaac ggcacaaagt tcagcgtgtc cggcgagggc 900
gagggcgatg ccacctacgg caagctgacc ctgaagttca tctgcaccac cggcaagctg 960
cccgtgcctt ggcccaccct cgtgaccacc ctgacctacg gcgtgcagtg ctteagccgc 1020
taccgccgac acatgaagca gcacgacttc ttcaagtccg ccacgcccga aggtacgtc 1080
caggagcgca ccattttctt caaggacgac ggcaactaca agaccgcgc cgagggtgaag 1140
ttcgagggcg acaccctggt gaaccgcatc gagctgaagg gcacgactt caaggaggac 1200
ggcaacatcc tggggcacaa gctggagtag aactacaaca gccacaacgt ctatatcatg 1260
gccgacaagc agaagaacgg catcaagggt aacttcaaga tccgccacaa catcgaggac 1320
ggcagcgtgc agctcgccga ccactaccag cagaacaccc ccacggcga cggccccgtg 1380
ctgctgcccc acaaccacta cctgagcacc cagtccgccc tgagcaaaga ccccaacgag 1440
aagcgcgatc acatgggtcct gctggaggtc gtgaccgccc cggggatcac tctcgcatg 1500
gacgagctgt acaagtaa 1518

```

```

<210> SEQ ID NO 261
<211> LENGTH: 23

```

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        oligonucleotide

<400> SEQUENCE: 261

gttgatgagg tggttcagga ggg                23

<210> SEQ ID NO 262
<211> LENGTH: 3490
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        polynucleotide

<400> SEQUENCE: 262

tcgcgcgttt cggatgatgac ggtgaaaacc tctgacacat gcagctcccc gagacgggtca    60
cagcttgctc gtaagcggat gccgggagca gacaagcccc tcagggcgcg tcagcgggtg    120
ttggcgggtg tcggggcttg cttaactatg cggcatcaga gcagattgta ctgagagtgc    180
accatattcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgcc    240
attcgccatt caggctgcgc aactgttggg aagggcgacg ggtgcgggcc tcttcgctat    300
tacgccagct ggcgaaaggg ggatgtgctg caaggcgatt aagttgggta acgccagggt    360
tttcccagtc acgacgttgt aaaacgacgg ccagtgaatt cgagggccta tttcccatga    420
ttccttcata tttcatata cgtacaagg ctgtagaga gataattgga attaatttga    480
ctgtaaacac aaagatatta gtacaaaata cgtgacgtag aaagtaataa tttcttgggt    540
agtttgcagt tttaaaatta tgttttaaaa tggactatca tatgcttacc gtaacttgaa    600
agtatttcga tttcttggtt ttatatatct tgtggaaaagg acgaaacacc gacccacggc    660
gtgcagtgtc tgttttagag ctagaaatag caagttaaaa taaggctagt ccgttatcaa    720
cttgaaaaag tggcaccgag tcggtgcttt tttgttttag agctagaaat agcaagttaa    780
aataaggcta gtcggttttt agcgcgtgcg ccaattctgc agacagagag ggcctatttc    840
ccatgattcc ttcatttttg catatacgat acaaggctgt tagagagata attggaatta    900
atttgactgt aaacacaaag atattagtag aaaatacgtg acgtagaaag taataatttc    960
ttgggtagtt tgcagtttta aaattatgtt ttaaaatgga ctatcatatg cttaccgtaa   1020
cttgaaagta tttcgatttc ttggctttat atatcttgty gaaaggacga aacaccgggt   1080
cttcgagaag acctgtttta gagctagaaa tagcaagtta aaataaggct agtccgttat   1140
caactgaaa aagtggcacc gagtcggtgc ttttttggtt tagagctaga aatagcaagt   1200
taaaataagg ctagtccgtt tttagcgcgt gcgccaattc tgcagacaaa aagcttggcg   1260
taatcatggt catagctgtt tcctgtgtga aattgttatc cgctcacaat tccacacaa   1320
atacgagccg gaagcataaa gtgtaaagcc tggggtgcct aatgagttag ctaactcaca   1380
ttaattgcgt tgcgctcact gcccgcttcc cagtcgggaa acctgtcgtg ccagctgcat   1440
taatgaatcg gccaacgcgc ggggagaggc ggtttgcgta ttgggcgctc ttccgcttcc   1500
tcgctcactg actcgtgcg ctcggtcgtt cggctgcggc gagcgggtatc agctcactca   1560
aaggcggtaa tacggttatc cacagaatca ggggataacg caggaaagaa catgtgagca   1620
aaaggccagc aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggtt tttccatagg   1680
ctccgcccc ctgacgagca tcacaaaaat cgacgctcaa gtcagagggtg gcgaaacccg   1740

```

-continued

```

acaggactat aaagatacca ggcgtttccc cctggaagct cctcgtgcg ctctcctgtt 1800
cgcaccctgc cgcttacccg atacctgtcc gcctttctcc ctccgggaag cgtggcgctt 1860
tctcatagct cagcgtgtag gtatctcagt tcggtgtagg tcgttcgctc caagctgggc 1920
tgtgtgcacg aacccccctg tcagcccgac cgtcgccct tatccggtaa ctatcgtctt 1980
gagtccaacc cggtaagaca cgacttatcg ccactggcag cagccactgg taacaggatt 2040
agcagagcga ggtatgtagg cggtgctaca gaggcttga agtggtggcc taactacggc 2100
tacactagaa gaacagtatt tggatatctgc gctctgtga agccagttac ctccgaaaa 2160
agagttggta gctcttgatc cggcaacaa accaccgctg gtagcggtag ttttttgtt 2220
tgcaagcagc agattacgag cagaaaaaaa ggatctcaag aagatccttt gatcttttct 2280
acggggtctg acgctcagtg gaacgaaaaac tcacgttaag ggattttggt catgagatta 2340
tcaaaaagga tcttcaccta gatcctttta aataaaaaat gaagttttaa atcaatctaa 2400
agtatatatg agtaaacttg gctcgacagt taccaatgct taatcagtga ggcacctatc 2460
tcagcgatct gtctatttcg ttcattcata gttgcctgac tcccgcctgt gtagataact 2520
acgatacggg agggcctacc atctggcccc agtgctgcaa tgataccgag agaccacgc 2580
tcaccggctc cagatttata agcaataaac cagccagccg gaagggccga gcgcagaagt 2640
ggctctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga agctagagta 2700
agtagttcgc cagttaatag tttgcgaac gttgttgcca ttgctacagg catcgtggtg 2760
tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgac aaggcgagtt 2820
acatgatccc ccatgttggtg caaaaaagcg gttagctcct tcggtcctcc gatcgttgtc 2880
agaagtaagt tggccgcagt gttatcactc atgggttatgg cagcactgca taattctctt 2940
actgtcatgc catcogtaag atgcttttct gtgactgggt agtactcaac caagtcattc 3000
tgagaatagt gtatgcggcg accgagttgc tcttgcccg cgtaatacg ggataatacc 3060
gcgccacata gcagaacttt aaaagtgtc atcattggaa aacgtttctt cggggcgaaaa 3120
ctctcaagga tcttaccgct gttgagatcc agttcgatgt aacccactcg tgcacccaac 3180
tgatcttcag catcttttac tttaccagc gtttctgggt gagcaaaaac aggaaggcaa 3240
aatgccgcaa aaaaggaat aagggcgaca cggaaatgtt gaatactcat actcttctt 3300
tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata catatttgaa 3360
tgtatttaga aaaataaaca aataggggtt ccgcgcacat tccccgaaa agtgccacct 3420
gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg tatcagcagg 3480
ccctttcgtc 3490

```

```

<210> SEQ ID NO 263
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

```

```

<400> SEQUENCE: 263

```

```

ggcccagact gagcacgtga tgg

```

23

```

<210> SEQ ID NO 264
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 264

ggtccagact gagcacgtga tgg 23

<210> SEQ ID NO 265
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 265

ggctcagact gagcacgtga tgg 23

<210> SEQ ID NO 266
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 266

ggcctagact gagcacgtga tgg 23

<210> SEQ ID NO 267
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 267

ggcccagatt gagcacgtga tgg 23

<210> SEQ ID NO 268
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 268

ggttcagact gagcacgtga tgg 23

<210> SEQ ID NO 269
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 269

ggtctagact gagcacgtga tgg 23

<210> SEQ ID NO 270
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

oligonucleotide

<400> SEQUENCE: 270

ggtccagatt gagcacgtga tgg 23

<210> SEQ ID NO 271
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 271

ggcttagact gagcacgtga tgg 23

<210> SEQ ID NO 272
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 272

ggctcagatt gagcacgtga tgg 23

<210> SEQ ID NO 273
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 273

ggcctagatt gagcacgtga tgg 23

<210> SEQ ID NO 274
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 274

ggtttagact gagcacgtga tgg 23

<210> SEQ ID NO 275
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 275

ggcttagatt gagcacgtga tgg 23

<210> SEQ ID NO 276
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 276

ggtctagatt gagcacgtga tgg 23

<210> SEQ ID NO 277

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 277

ggttcagatt gagcacgtga tgg 23

<210> SEQ ID NO 278

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 278

ggttttagatt gagcacgtga tgg 23

<210> SEQ ID NO 279

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 279

ggcccagact gagcacgtga tgg 23

<210> SEQ ID NO 280

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 280

ggtccagact gagcacgtga tgg 23

<210> SEQ ID NO 281

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 281

ggctcagact gagcacgtga tgg 23

<210> SEQ ID NO 282

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

-continued

<400> SEQUENCE: 282

ggcctagact gagcacgtga tgg 23

<210> SEQ ID NO 283

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 283

ggcccagatt gagcacgtga tgg 23

<210> SEQ ID NO 284

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 284

ggttcagact gagcacgtga tgg 23

<210> SEQ ID NO 285

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 285

ggtctagact gagcacgtga tgg 23

<210> SEQ ID NO 286

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 286

ggtccagatt gagcacgtga tgg 23

<210> SEQ ID NO 287

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 287

ggcttagact gagcacgtga tgg 23

<210> SEQ ID NO 288

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 288

-continued

ggctcagatt gagcacgtga tgg 23

<210> SEQ ID NO 289
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 289

ggcctagatt gagcacgtga tgg 23

<210> SEQ ID NO 290
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 290

ggttttagact gagcacgtga tgg 23

<210> SEQ ID NO 291
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 291

ggcttagatt gagcacgtga tgg 23

<210> SEQ ID NO 292
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 292

ggtctagatt gagcacgtga tgg 23

<210> SEQ ID NO 293
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 293

ggttcagatt gagcacgtga tgg 23

<210> SEQ ID NO 294
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 294

-continued

ggttttagatt gaggcagtgga tgg 23

<210> SEQ ID NO 295
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 295

gaagcgccctg gcagtggtacc agg 23

<210> SEQ ID NO 296
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 296

gaagtgcctg gcagtggtacc agg 23

<210> SEQ ID NO 297
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 297

gaagcgtctg gcagtggtacc agg 23

<210> SEQ ID NO 298
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 298

gaagcgcttg gcagtggtacc agg 23

<210> SEQ ID NO 299
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 299

gaagtgtctg gcagtggtacc agg 23

<210> SEQ ID NO 300
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 300

gaagtgtctg gcagtggtacc agg 23

-continued

<210> SEQ ID NO 301
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 301

gaagcgtttg gcagtggtacc agg 23

<210> SEQ ID NO 302
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 302

gaagtgtttg gcagtggtacc agg 23

<210> SEQ ID NO 303
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 303

gaagcgcttg gcagtggtacc agg 23

<210> SEQ ID NO 304
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 304

gaagtgcctg gcagtggtacc agg 23

<210> SEQ ID NO 305
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 305

gaagcgcttg gcagtggtacc agg 23

<210> SEQ ID NO 306
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
oligonucleotide

<400> SEQUENCE: 306

gaagcgcttg gcagtggtacc agg 23

-continued

```

<210> SEQ ID NO 307
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 307

```

```

gaagtgtctg gcagtgtacc agg                                     23

```

```

<210> SEQ ID NO 308
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 308

```

```

gaagtgcttg gcagtgtacc agg                                     23

```

```

<210> SEQ ID NO 309
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 309

```

```

gaagcgtttg gcagtgtacc agg                                     23

```

```

<210> SEQ ID NO 310
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      oligonucleotide

```

```

<400> SEQUENCE: 310

```

```

gaagtgtttg gcagtgtacc agg                                     23

```

```

<210> SEQ ID NO 311
<211> LENGTH: 257
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
      polypeptide

```

```

<400> SEQUENCE: 311

```

```

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe
 1             5             10             15

```

```

Met Arg Phe Lys Val His Met Glu Gly Ser Val Asn Gly His Glu Phe
 20             25             30

```

```

Glu Ile Glu Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly Thr Gln Thr
 35             40             45

```

```

Ala Lys Leu Lys Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Trp Asp
 50             55             60

```

```

Ile Leu Ser Pro Gln Phe Met Tyr Gly Ser Lys Ala Tyr Val Lys His
 65             70             75             80

```

```

Pro Ala Asp Ile Pro Asp Tyr Leu Lys Leu Ser Phe Pro Glu Gly Phe

```

-continued

85					90					95					
Lys	Trp	Glu	Arg	Val	Met	Asn	Phe	Glu	Asp	Gly	Gly	Val	Val	Thr	Val
			100					105					110		
Thr	Gln	Asp	Ser	Ser	Leu	Gln	Asp	Gly	Glu	Phe	Ile	Tyr	Lys	Val	Lys
		115					120					125			
Leu	Arg	Gly	Thr	Asn	Phe	Pro	Ser	Asp	Gly	Pro	Val	Met	Lys	Lys	Thr
	130					135					140				
Met	Gly	Trp	Glu	Ala	Ser	Ser	Glu	Arg	Met	Tyr	Pro	Glu	Asp	Gly	Ala
145						150					155				160
Leu	Lys	Gly	Glu	Ile	Lys	Gln	Arg	Leu	Lys	Leu	Lys	Asp	Gly	Gly	His
			165					170					175		
Tyr	Asp	Ala	Glu	Val	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Pro	Val	Gln
		180						185					190		
Leu	Pro	Gly	Ala	Tyr	Asn	Val	Asn	Ile	Lys	Leu	Asp	Ile	Thr	Ser	His
		195					200					205			
Asn	Glu	Asp	Tyr	Thr	Ile	Val	Glu	Gln	Tyr	Glu	Arg	Ala	Glu	Gly	Arg
	210					215					220				
His	Ser	Thr	Gly	Gly	Met	Asp	Glu	Leu	Tyr	Lys	Pro	Arg	Glu	Gly	Arg
225						230					235				240
Gly	Ser	Leu	Leu	Thr	Cys	Gly	Asp	Val	Glu	Glu	Asn	Pro	Gly	Pro	Thr
			245						250				255		

Ser

<210> SEQ ID NO 312
 <211> LENGTH: 257
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 312

Met	Val	Ser	Lys	Gly	Glu	Glu	Asp	Asn	Met	Ala	Ile	Ile	Lys	Glu	Phe
1			5					10					15		
Met	Arg	Phe	Lys	Val	His	Met	Glu	Gly	Ser	Val	Asn	Gly	His	Glu	Phe
		20					25					30			
Glu	Ile	Glu	Gly	Glu	Gly	Glu	Gly	Arg	Pro	Tyr	Glu	Gly	Thr	Gln	Thr
	35					40					45				
Ala	Lys	Leu	Lys	Val	Thr	Lys	Gly	Gly	Pro	Leu	Pro	Phe	Ala	Trp	Asp
	50					55				60					
Ile	Leu	Ser	Pro	Gln	Phe	Met	Tyr	Gly	Ser	Lys	Ala	Tyr	Val	Lys	His
65				70				75						80	
Pro	Ala	Asp	Ile	Pro	Asp	Tyr	Leu	Lys	Leu	Ser	Phe	Pro	Glu	Gly	Phe
		85					90					95			
Lys	Trp	Glu	Arg	Val	Met	Asn	Phe	Glu	Asp	Gly	Gly	Val	Val	Thr	Val
		100					105					110			
Thr	Gln	Asp	Ser	Ser	Leu	Gln	Asp	Gly	Glu	Phe	Ile	Tyr	Lys	Val	Lys
	115					120					125				
Leu	Arg	Gly	Thr	Asn	Phe	Pro	Ser	Asp	Gly	Pro	Val	Met	Lys	Lys	Thr
	130					135					140				
Met	Gly	Trp	Glu	Ala	Ser	Ser	Glu	Arg	Met	Tyr	Pro	Glu	Asp	Gly	Ala
145					150					155					160
Leu	Lys	Gly	Glu	Ile	Lys	Gln	Arg	Leu	Lys	Leu	Lys	Asp	Gly	Gly	His
			165					170					175		
Tyr	Asp	Ala	Glu	Val	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Pro	Val	Gln

-continued

180	185	190
Leu Pro Gly Ala Tyr Asn Val	Asn Ile Lys Leu Asp Ile Thr Ser His	
195	200	205
Asn Glu Asp Tyr Thr Ile Val	Glu Gln Tyr Glu Arg Ala Glu Gly Arg	
210	215	220
His Ser Thr Gly Gly Met Asp	Glu Leu Tyr Lys Pro Arg Glu Gly Arg	
225	230	235
Gly Ser Leu Leu Thr Cys Gly	Asp Val Glu Glu Asn Pro Gly Pro Thr	
245	250	255

Ser

<210> SEQ ID NO 313

<211> LENGTH: 240

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 313

Met Arg Glu Gln Gly Arg Gly	Ala Val His Arg Gly Gly	Ala His Pro
1	5	10

Gly Arg Ala Gly Arg Arg Arg	Lys Arg Pro Gln Val Gln Arg Val Arg
20	25

Arg Gly Arg Gly Arg Cys His	Leu Arg Gln Ala Asp Pro Glu Val His
35	40

Leu His His Arg Gln Ala Ala	Arg Ala Leu Ala His Pro Arg Asp His
50	55

Pro Asp Leu Arg Arg Ala Val	Leu Gln Pro Leu Pro Arg Pro His Glu
65	70

Ala Ala Arg Leu Leu Gln Val	Arg His Ala Arg Arg Leu Arg Pro Gly
85	90

Ala His His Leu Leu Gln Gly	Arg Arg Gln Leu Gln Asp Pro Arg Arg
100	105

Gly Glu Val Arg Gly Arg His	Pro Gly Glu Pro His Arg Ala Glu Gly
115	120

His Arg Leu Gln Gly Gly Arg	Gln His Pro Gly Ala Gln Ala Gly Val
130	135

Gln Leu Gln Gln Pro Gln Arg	Leu Tyr His Gly Arg Gln Ala Glu Glu
145	150

Arg His Gln Gly Glu Leu Gln	Asp Pro Pro Gln His Arg Gly Arg Gln
165	170

Arg Ala Ala Arg Arg Pro Leu	Pro Ala Glu His Pro His Arg Arg Arg
180	185

Pro Arg Ala Ala Ala Arg Gln	Pro Leu Pro Glu His Pro Val Arg Pro
195	200

Glu Gln Arg Pro Gln Arg Glu	Ala Arg Ser His Gly Pro Ala Gly Val
210	215

Arg Asp Arg Arg Arg Asp His	Ser Arg His Gly Arg Ala Val Gln Val
225	230

<210> SEQ ID NO 314

<211> LENGTH: 236

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic

-continued

polypeptide

<400> SEQUENCE: 314

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe
 1 5 10 15

Met Arg Phe Lys Val His Met Glu Gly Ser Val Asn Gly His Glu Phe
 20 25 30

Glu Ile Glu Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly Thr Gln Thr
 35 40 45

Ala Lys Leu Lys Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Trp Asp
 50 55 60

Ile Leu Ser Pro Gln Phe Met Tyr Gly Ser Lys Ala Tyr Val Lys His
 65 70 75 80

Pro Ala Asp Ile Pro Asp Tyr Leu Lys Leu Ser Phe Pro Glu Gly Phe
 85 90 95

Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val Thr Val
 100 105 110

Thr Gln Asp Ser Ser Leu Gln Asp Gly Glu Phe Ile Tyr Lys Val Lys
 115 120 125

Leu Arg Gly Thr Asn Phe Pro Ser Asp Gly Pro Val Met Gln Lys Lys
 130 135 140

Thr Met Gly Trp Glu Ala Ser Ser Glu Arg Met Tyr Pro Glu Asp Gly
 145 150 155 160

Ala Leu Lys Gly Glu Ile Lys Gln Arg Leu Lys Leu Lys Asp Gly Gly
 165 170 175

His Tyr Asp Ala Glu Val Lys Thr Thr Tyr Lys Ala Lys Lys Pro Val
 180 185 190

Gln Leu Pro Gly Ala Tyr Asn Val Asn Ile Lys Leu Asp Ile Thr Ser
 195 200 205

His Asn Glu Asp Tyr Thr Ile Val Glu Gln Tyr Glu Arg Ala Glu Gly
 210 215 220

Arg His Ser Thr Gly Gly Met Asp Glu Leu Tyr Lys
 225 230 235

<210> SEQ ID NO 315

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic oligonucleotide

<400> SEQUENCE: 315

gttgatgggg tggttcagga ggg

23

<210> SEQ ID NO 316

<211> LENGTH: 504

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 316

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe
 1 5 10 15

Met Arg Phe Lys Val His Met Glu Gly Ser Val Asn Gly His Glu Phe
 20 25 30

-continued

Glu	Ile	Glu	Gly	Glu	Gly	Glu	Gly	Arg	Pro	Tyr	Glu	Gly	Thr	Gln	Thr
	35						40					45			
Ala	Lys	Leu	Lys	Val	Thr	Lys	Gly	Gly	Pro	Leu	Pro	Phe	Ala	Trp	Asp
	50					55					60				
Ile	Leu	Ser	Pro	Gln	Phe	Met	Tyr	Gly	Ser	Lys	Ala	Tyr	Val	Lys	His
65				70					75					80	
Pro	Ala	Asp	Ile	Pro	Asp	Tyr	Leu	Lys	Leu	Ser	Phe	Pro	Glu	Gly	Phe
			85					90					95		
Lys	Trp	Glu	Arg	Val	Met	Asn	Phe	Glu	Asp	Gly	Gly	Val	Val	Thr	Val
			100					105				110			
Thr	Gln	Asp	Ser	Ser	Leu	Gln	Asp	Gly	Glu	Phe	Ile	Tyr	Lys	Val	Lys
	115					120					125				
Leu	Arg	Gly	Thr	Asn	Phe	Pro	Ser	Asp	Gly	Pro	Val	Met	Gln	Lys	Lys
	130			135							140				
Thr	Met	Gly	Trp	Glu	Ala	Ser	Ser	Glu	Arg	Met	Tyr	Pro	Glu	Asp	Gly
145				150					155						160
Ala	Leu	Lys	Gly	Glu	Ile	Lys	Gln	Arg	Leu	Lys	Leu	Lys	Asp	Gly	Gly
			165					170						175	
His	Tyr	Asp	Ala	Glu	Val	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Pro	Val
	180						185					190			
Gln	Leu	Pro	Gly	Ala	Tyr	Asn	Val	Asn	Ile	Lys	Leu	Asp	Ile	Thr	Ser
	195					200						205			
His	Asn	Glu	Asp	Tyr	Thr	Ile	Val	Glu	Gln	Tyr	Glu	Arg	Ala	Glu	Gly
	210				215						220				
Arg	His	Ser	Thr	Gly	Gly	Met	Asp	Glu	Leu	Tyr	Lys	Pro	Arg	Glu	Gly
225				230					235					240	
Arg	Gly	Ser	Leu	Leu	Thr	Cys	Gly	Asp	Val	Glu	Glu	Asn	Pro	Gly	Pro
			245					250						255	
Thr	Ser	Trp	Gly	Gly	Ser	Gly	Gly	Ala	Cys	Val	Ser	Lys	Gly	Glu	Glu
			260				265					270			
Leu	Phe	Thr	Gly	Val	Val	Pro	Ile	Leu	Val	Glu	Leu	Asp	Gly	Asp	Val
	275					280					285				
Asn	Gly	His	Lys	Phe	Ser	Val	Ser	Gly	Glu	Gly	Glu	Gly	Asp	Ala	Thr
	290				295						300				
Tyr	Gly	Lys	Leu	Thr	Leu	Lys	Phe	Ile	Cys	Thr	Thr	Gly	Lys	Leu	Pro
305				310					315					320	
Val	Pro	Trp	Pro	Thr	Leu	Val	Thr	Thr	Leu	Thr	Tyr	Gly	Val	Gln	Cys
			325					330						335	
Phe	Ser	Arg	Tyr	Pro	Asp	His	Met	Lys	Gln	His	Asp	Phe	Phe	Lys	Ser
		340					345					350			
Ala	Met	Pro	Glu	Gly	Tyr	Val	Gln	Glu	Arg	Thr	Ile	Phe	Phe	Lys	Asp
	355					360					365				
Asp	Gly	Asn	Tyr	Lys	Thr	Arg	Ala	Glu	Val	Lys	Phe	Glu	Gly	Asp	Thr
	370				375						380				
Leu	Val	Asn	Arg	Ile	Glu	Leu	Lys	Gly	Ile	Asp	Phe	Lys	Glu	Asp	Gly
385				390					395					400	
Asn	Ile	Leu	Gly	His	Lys	Leu	Glu	Tyr	Asn	Tyr	Asn	Ser	His	Asn	Val
		405						410					415		
Tyr	Ile	Met	Ala	Asp	Lys	Gln	Lys	Asn	Gly	Ile	Lys	Val	Asn	Phe	Lys
	420						425					430			
Ile	Arg	His	Asn	Ile	Glu	Asp	Gly	Ser	Val	Gln	Leu	Ala	Asp	His	Tyr
	435					440					445				
Gln	Gln	Asn	Thr	Pro	Ile	Gly	Asp	Gly	Pro	Val	Leu	Leu	Pro	Asp	Asn

-continued

450	455	460
His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys		
465	470	475 480
Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr		
	485	490 495
Leu Gly Met Asp Glu Leu Tyr Lys		
	500	

<210> SEQ ID NO 317
 <211> LENGTH: 505
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 317

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe		
1	5	10 15
Met Arg Phe Lys Val His Met Glu Gly Ser Val Asn Gly His Glu Phe		
	20	25 30
Glu Ile Glu Gly Glu Gly Glu Gly Arg Pro Tyr Glu Gly Thr Gln Thr		
	35	40 45
Ala Lys Leu Lys Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Trp Asp		
	50	55 60
Ile Leu Ser Pro Gln Phe Met Tyr Gly Ser Lys Ala Tyr Val Lys His		
65	70	75 80
Pro Ala Asp Ile Pro Asp Tyr Leu Lys Leu Ser Phe Pro Glu Gly Phe		
	85	90 95
Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val Thr Val		
	100	105 110
Thr Gln Asp Ser Ser Leu Gln Asp Gly Glu Phe Ile Tyr Lys Val Lys		
	115	120 125
Leu Arg Gly Thr Asn Phe Pro Ser Asp Gly Pro Val Met Gln Lys Lys		
	130	135 140
Thr Met Gly Trp Glu Ala Ser Ser Glu Arg Met Tyr Pro Glu Asp Gly		
145	150	155 160
Ala Leu Lys Gly Glu Ile Lys Gln Arg Leu Lys Leu Lys Asp Gly Gly		
	165	170 175
His Tyr Asp Ala Glu Val Lys Thr Thr Tyr Lys Ala Lys Lys Pro Val		
	180	185 190
Gln Leu Pro Gly Ala Tyr Asn Val Asn Ile Lys Leu Asp Ile Thr Ser		
	195	200 205
His Asn Glu Asp Tyr Thr Ile Val Glu Gln Tyr Glu Arg Ala Glu Gly		
	210	215 220
Arg His Ser Thr Gly Gly Met Asp Glu Leu Tyr Lys Pro Arg Glu Gly		
225	230	235 240
Arg Gly Ser Leu Leu Thr Cys Gly Asp Val Glu Glu Asn Pro Gly Pro		
	245	250 255
Thr Ser Trp Trp Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu		
	260	265 270
Glu Leu Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp		
	275	280 285
Val Asn Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala		
	290	295 300

-continued

```

Thr Tyr Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu
305                310                315                320

Pro Val Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln
                325                330                335

Cys Phe Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys
                340                345                350

Ser Ala Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys
                355                360                365

Asp Asp Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp
370                375                380

Thr Leu Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp
385                390                395                400

Gly Asn Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn
                405                410                415

Val Tyr Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe
                420                425                430

Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His
                435                440                445

Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp
450                455                460

Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu
465                470                475                480

Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile
                485                490                495

Thr Leu Gly Met Asp Glu Leu Tyr Lys
500                505

```

```

<210> SEQ ID NO 318
<211> LENGTH: 503
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
                        polypeptide

```

```

<400> SEQUENCE: 318

```

```

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe
1          5          10          15

Met Arg Phe Lys Val His Met Glu Gly Ser Val Asn Gly His Glu Phe
20        25        30

Glu Ile Glu Gly Glu Gly Glu Arg Pro Tyr Glu Gly Thr Gln Thr
35        40        45

Ala Lys Leu Lys Val Thr Lys Gly Gly Pro Leu Pro Phe Ala Trp Asp
50        55        60

Ile Leu Ser Pro Gln Phe Met Tyr Gly Ser Lys Ala Tyr Val Lys His
65        70        75        80

Pro Ala Asp Ile Pro Asp Tyr Leu Lys Leu Ser Phe Pro Glu Gly Phe
85        90        95

Lys Trp Glu Arg Val Met Asn Phe Glu Asp Gly Gly Val Val Thr Val
100       105       110

Thr Gln Asp Ser Ser Leu Gln Asp Gly Glu Phe Ile Tyr Lys Val Lys
115       120       125

Leu Arg Gly Thr Asn Phe Pro Ser Asp Gly Pro Val Met Gln Lys Lys
130       135       140

Thr Met Gly Trp Glu Ala Ser Ser Glu Arg Met Tyr Pro Glu Asp Gly
145       150       155       160

```

-continued

Ala Leu Lys Gly Glu Ile Lys Gln Arg Leu Lys Leu Lys Asp Gly Gly
 165 170 175
 His Tyr Asp Ala Glu Val Lys Thr Thr Tyr Lys Ala Lys Lys Pro Val
 180 185 190
 Gln Leu Pro Gly Ala Tyr Asn Val Asn Ile Lys Leu Asp Ile Thr Ser
 195 200 205
 His Asn Glu Asp Tyr Thr Ile Val Glu Gln Tyr Glu Arg Ala Glu Gly
 210 215 220
 Arg His Ser Thr Gly Gly Met Asp Glu Leu Tyr Lys Pro Arg Glu Gly
 225 230 235 240
 Arg Gly Ser Leu Leu Thr Cys Gly Asp Val Glu Glu Asn Pro Gly Pro
 245 250 255
 Thr Ser Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu Glu Leu
 260 265 270
 Phe Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn
 275 280 285
 Gly His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr
 290 295 300
 Gly Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val
 305 310 315 320
 Pro Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe
 325 330 335
 Ser Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala
 340 345 350
 Met Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp
 355 360 365
 Gly Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu
 370 375 380
 Val Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn
 385 390 395 400
 Ile Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr
 405 410 415
 Ile Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile
 420 425 430
 Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln
 435 440 445
 Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His
 450 455 460
 Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg
 465 470 475 480
 Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu
 485 490 495
 Gly Met Asp Glu Leu Tyr Lys
 500

<210> SEQ ID NO 319

<211> LENGTH: 505

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 319

Met Val Ser Lys Gly Glu Glu Asp Asn Met Ala Ile Ile Lys Glu Phe

-continued

1	5	10	15
Met Arg Phe	Lys Val His Met	Glu Gly Ser Val Asn Gly His	Glu Phe
	20	25	30
Glu Ile Glu	Gly Glu Gly Glu	Gly Arg Pro Tyr Glu Gly Thr	Gln Thr
	35	40	45
Ala Lys Leu	Lys Val Thr Lys	Gly Gly Pro Leu Pro Phe Ala Trp	Asp
	50	55	60
Ile Leu Ser	Pro Gln Phe Met	Tyr Gly Ser Lys Ala Tyr Val Lys	His
	65	70	80
Pro Ala Asp	Ile Pro Asp Tyr	Leu Lys Leu Ser Phe Pro Glu Gly	Phe
	85	90	95
Lys Trp Glu	Arg Val Met Asn Phe	Glu Asp Gly Gly Val Val Thr	Val
	100	105	110
Thr Gln Asp	Ser Ser Leu Gln Asp	Gly Glu Phe Ile Tyr Lys Val	Lys
	115	120	125
Leu Arg Gly	Thr Asn Phe Pro Ser	Asp Gly Pro Val Met Gln Lys	Lys
	130	135	140
Thr Met Gly	Trp Glu Ala Ser Ser	Glu Arg Met Tyr Pro Glu Asp	Gly
	145	150	160
Ala Leu Lys	Gly Glu Ile Lys Gln	Arg Leu Lys Leu Lys Asp	Gly Gly
	165	170	175
His Tyr Asp	Ala Glu Val Lys Thr	Thr Tyr Lys Ala Lys Lys	Pro Val
	180	185	190
Gln Leu Pro	Gly Ala Tyr Asn Val	Asn Ile Lys Leu Asp Ile Thr	Ser
	195	200	205
His Asn Glu	Asp Tyr Thr Ile Val	Glu Gln Tyr Glu Arg Ala Glu	Gly
	210	215	220
Arg His Ser	Thr Gly Gly Met Asp	Glu Leu Tyr Lys Pro Arg Glu	Gly
	225	230	240
Arg Gly Ser	Leu Leu Thr Cys Gly	Asp Val Glu Glu Asn Pro Gly	Pro
	245	250	255
Thr Ser Trp	Trp Gly Gly Ser Gly	Gly Ala Cys Val Ser Lys Gly	Glu
	260	265	270
Glu Leu Phe	Thr Gly Val Val Pro	Ile Leu Val Glu Leu Asp Gly	Asp
	275	280	285
Val Asn Gly	His Lys Phe Ser Val	Ser Gly Glu Gly Glu Gly Asp	Ala
	290	295	300
Thr Tyr Gly	Lys Leu Thr Leu Lys	Phe Ile Cys Thr Thr Gly Lys	Leu
	305	310	320
Pro Val Pro	Trp Pro Thr Leu Val	Thr Thr Leu Thr Tyr Gly Val	Gln
	325	330	335
Cys Phe Ser	Arg Tyr Pro Asp His	Met Lys Gln His Asp Phe Phe	Lys
	340	345	350
Ser Ala Met	Pro Glu Gly Tyr Val	Gln Glu Arg Thr Ile Phe Phe	Lys
	355	360	365
Asp Asp Gly	Asn Tyr Lys Thr Arg	Ala Glu Val Lys Phe Glu Gly	Asp
	370	375	380
Thr Leu Val	Asn Arg Ile Glu Leu	Lys Gly Ile Asp Phe Lys Glu	Asp
	385	390	400
Gly Asn Ile	Leu Gly His Lys Leu	Glu Tyr Asn Tyr Asn Ser His	Asn
	405	410	415
Val Tyr Ile	Met Ala Asp Lys Gln	Lys Asn Gly Ile Lys Val Asn	Phe
	420	425	430

-continued

Lys Ile Arg His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His
435 440 445

Tyr Gln Gln Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp
450 455 460

Asn His Tyr Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu
465 470 475 480

Lys Arg Asp His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile
485 490 495

Thr Leu Gly Met Asp Glu Leu Tyr Lys
500 505

<210> SEQ ID NO 320

<211> LENGTH: 246

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 320

Trp Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu Glu Leu Phe
1 5 10 15

Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly
20 25 30

His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly
35 40 45

Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro
50 55 60

Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser
65 70 75 80

Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met
85 90 95

Pro Glu Gly Tyr Tyr Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly
100 105 110

Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val
115 120 125

Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile
130 135 140

Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile
145 150 155 160

Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg
165 170 175

His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln
180 185 190

Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr
195 200 205

Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp
210 215 220

His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly
225 230 235 240

Met Asp Glu Leu Tyr Lys
245

<210> SEQ ID NO 321

<211> LENGTH: 245

<212> TYPE: PRT

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 321

Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu Glu Leu Phe Thr
 1 5 10 15
 Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His
 20 25 30
 Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys
 35 40 45
 Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp
 50 55 60
 Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg
 65 70 75 80
 Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro
 85 90 95
 Glu Gly Tyr Tyr Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn
 100 105 110
 Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn
 115 120 125
 Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu
 130 135 140
 Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met
 145 150 155 160
 Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His
 165 170 175
 Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn
 180 185 190
 Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu
 195 200 205
 Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His
 210 215 220
 Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met
 225 230 235 240
 Asp Glu Leu Tyr Lys
 245

<210> SEQ ID NO 322
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 322

Trp Gly Gly Ser Gly Gly
 1 5

<210> SEQ ID NO 323
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 323

-continued

Gly Gly Ser Gly Gly
1 5

<210> SEQ ID NO 324
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 324

Ser Trp Trp Gly Gly Ser Gly Gly
1 5

<210> SEQ ID NO 325
<211> LENGTH: 6
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic peptide

<400> SEQUENCE: 325

Lys Arg Leu Ala Val Tyr
1 5

<210> SEQ ID NO 326
<211> LENGTH: 246
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic polypeptide

<400> SEQUENCE: 326

Trp Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu Glu Leu Phe
1 5 10 15

Thr Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly
20 25 30

His Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly
35 40 45

Lys Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro
50 55 60

Trp Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser
65 70 75 80

Arg Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met
85 90 95

Pro Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly
100 105 110

Asn Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val
115 120 125

Asn Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile
130 135 140

Leu Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile
145 150 155 160

Met Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg
165 170 175

-continued

```

His Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln
    180                      185                      190

Asn Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr
    195                      200                      205

Leu Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp
    210                      215                      220

His Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly
    225                      230                      235                      240

Met Asp Glu Leu Tyr Lys
    245

<210> SEQ ID NO 327
<211> LENGTH: 245
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
    polypeptide

<400> SEQUENCE: 327

Gly Gly Ser Gly Gly Ala Cys Val Ser Lys Gly Glu Glu Leu Phe Thr
1      5      10      15

Gly Val Val Pro Ile Leu Val Glu Leu Asp Gly Asp Val Asn Gly His
    20      25      30

Lys Phe Ser Val Ser Gly Glu Gly Glu Gly Asp Ala Thr Tyr Gly Lys
    35      40      45

Leu Thr Leu Lys Phe Ile Cys Thr Thr Gly Lys Leu Pro Val Pro Trp
    50      55      60

Pro Thr Leu Val Thr Thr Leu Thr Tyr Gly Val Gln Cys Phe Ser Arg
    65      70      75      80

Tyr Pro Asp His Met Lys Gln His Asp Phe Phe Lys Ser Ala Met Pro
    85      90      95

Glu Gly Tyr Val Gln Glu Arg Thr Ile Phe Phe Lys Asp Asp Gly Asn
    100     105     110

Tyr Lys Thr Arg Ala Glu Val Lys Phe Glu Gly Asp Thr Leu Val Asn
    115     120     125

Arg Ile Glu Leu Lys Gly Ile Asp Phe Lys Glu Asp Gly Asn Ile Leu
    130     135     140

Gly His Lys Leu Glu Tyr Asn Tyr Asn Ser His Asn Val Tyr Ile Met
    145     150     155     160

Ala Asp Lys Gln Lys Asn Gly Ile Lys Val Asn Phe Lys Ile Arg His
    165     170     175

Asn Ile Glu Asp Gly Ser Val Gln Leu Ala Asp His Tyr Gln Gln Asn
    180     185     190

Thr Pro Ile Gly Asp Gly Pro Val Leu Leu Pro Asp Asn His Tyr Leu
    195     200     205

Ser Thr Gln Ser Ala Leu Ser Lys Asp Pro Asn Glu Lys Arg Asp His
    210     215     220

Met Val Leu Leu Glu Phe Val Thr Ala Ala Gly Ile Thr Leu Gly Met
    225     230     235     240

Asp Glu Leu Tyr Lys
    245

```

We claim:

1. A polynucleotide encoding one or more reporter polypeptides, the polynucleotide including a PAM site adjacent

to a base that when edited causes a change in a function or characteristic of the one or more reporter polypeptides, wherein the polynucleotide comprises a sequence selected

231

from the group consisting of: SEQ ID NO: 258; SEQ ID NO:259; and SEQ ID NO: 260; and wherein the one or more reporter polypeptides are selected from:

- a reporter polypeptide comprising SEQ ID NO: 2, wherein the polypeptide comprises histidine at amino acid position number 67 and glycine at amino acid position number 73;
- a reporter polypeptide comprising SEQ ID NO: 316; and
- a reporter polypeptide comprising SEQ ID NO: 318.

* * * * *

10

232