

(19) **United States**
(12) **Patent Application Publication**
Young
(10) **Pub. No.: US 2025/0276053 A1**
(43) **Pub. Date: Sep. 4, 2025**

(54) **ROTAVIRUS VACCINES**

(71) Applicant: **VST LLC dba MEDGENE**,
Brookings, SD (US)

(72) Inventor: **Alan John Young**, Brookings, SD (US)

(73) Assignee: **VST LLC dba MEDGENE**,
Brookings, SD (US)

(21) Appl. No.: **18/593,328**

(22) Filed: **Mar. 1, 2024**

Publication Classification

(51) **Int. Cl.**
A61K 39/12 (2006.01)
A61K 39/00 (2006.01)
C07K 14/005 (2006.01)

(52) **U.S. Cl.**


CPC **A61K 39/12** (2013.01); **C07K 14/005**
(2013.01); **A61K 2039/54** (2013.01); **A61K**
2039/541 (2013.01); **A61K 2039/545**
(2013.01); **A61K 2039/55566** (2013.01); **A61K**
2039/575 (2013.01); **C12N 2720/12322**
(2013.01); **C12N 2720/12334** (2013.01); **C12N**
2720/12351 (2013.01)

(57) **ABSTRACT**


The present invention describes immunogenic compositions containing immunogenic polypeptides of Rotavirus (RV), including immunogenic compositions containing antigens other than RV antigens and antigens that may be used in immunization against pathogens that cause diarrheal diseases. Methods of eliciting an immune response with the immunogenic compositions as disclosed and methods of treating an RV infection are also described.

Specification includes a Sequence Listing.

Average Scour Scores - Medgene vs. Competitor



Average Scour Scores of Litters that Scoured - Medgene vs. Competitor



Average Scour Scores - Medgene vs. Competitor




FIG. 1a

Average Scour Scores of Litters that
Scoured - Medgene vs. Competitor




FIG. 1b

Medgene Scour Scores by Parity




FIG. 2a

Competitor Scour Scores by Parity




FIG. 2b

ROTAVIRUS VACCINES

BACKGROUND OF THE INVENTION

Field of Invention

[0001] The present invention pertains generally to compositions that elicit immune responses against Rotavirus infection. In particular, the invention relates to immunogenic compositions (e.g., vaccines) comprising immunogenic polypeptides of Rotavirus A and C. Immunogenic compositions, in addition, may contain antigens other than Rotavirus antigens. Methods of eliciting an immune response with the immunogenic compositions as disclosed herein and methods of treating a Rotavirus infection are also described.

Background Information

[0002] Swine Rotaviruses are a prevalent and economically devastating disease that continue to plague the swine industry. Current vaccines necessarily rely on outdated strains that may not cover the full breadth of Rotaviruses impacting producers today.

[0003] Rotaviruses (RVs) are endemic in swine populations, and all swine herds certainly have a history of RV infection and circulation. Rotavirus A (RVA) and C (RVC) are the most common among all RV species reported in swine. RVA was considered most prevalent and pathogenic in swine; however, RVC has been emerging as a significant cause of enteritis in newborn piglets. Newborn piglets are highly susceptible to RV infection at birth.

[0004] Rotaviruses (RVs) are double-stranded RNA viruses belonging to the Rotavirus genus in the Reoviridae family. The RV genome is approximately 18,500 bp in size and consists of 11 segments of dsRNA encoding six structural proteins (VP1-VP4, VP6 and VP7) and five non-structural proteins (NSP 1-NSP5/6). The inner capsid of the virion comprises VP1, VP2, and VP3 segments. The VP6 forms the middle layer of the capsid, while the outer capsid is composed of the VP7 and VP4 proteins. VP7, a glycoprotein with a molecular weight of 37 kDa, constitutes 30% of the virus protein, and forms the smooth external surface of the outer shell. The minor component of the outer shell, VP4, is present as a series of spikes that project outward from the VP7 shell. VP4 is non-glycosylated, has a molecular weight of 88 kDa, and constitutes 1.5% of the virus protein. The VP4 gets proteolytically cleaved into VP5* and VP8* segments. The VP8* forms the head of the VP4 protein which helps in host attachment and infectivity. VP4 has been implicated in several important functions, including cell attachment and penetration, hemagglutination, neutralization, host range, and virulence. RVs are unique since the NSP4 produces an enterotoxin, which contributes to viral pathogenesis.

[0005] RV species are classified based on sequencing of the VP6 gene. A binary classification system is used to address vast Rotavirus diversity on the basis of sequencing of G (VP7) and P types (VP4). The dual (G/P) typing system has been extended to a complete genome classification system based on nucleotide sequencing of all 11 RV segments with nucleotide percent identity cut-off values set for each segment. In this system, VP7-VP4-VP6-VP1-VP2-VP3-NSP1-NSP2-NSP3-NSP4-NSP5/6 RV genes are designated as Gx-P[x]-Ix-Rx-Cx-Mx-Ax-Nx-Tx-Ex-Hx.

[0006] Currently, nine RV species (RVA-RVD and RVF-J) have been classified by the International Committee on Taxonomy of Viruses. However, only species A, B, C, E, and H have been reported from swine. RVs are ubiquitous in nature, and mixed infections involving several RV strains appear more common as pigs grow older. Once infected, piglets may exhibit clinical or subclinical symptoms and eventual recovery in most cases. However, neonatal and suckling piglets without an established adaptive immune system are worst affected. RVA is the most characterized species among RVs due to its wide host range, high prevalence, and pathogenicity.

[0007] RV transmission is through fecal-oral route, and the piglets become infected with RV shed from sows and other piglets. The target sites of RV replication are the mature, non-dividing enterocytes in the small intestine. Enterocytes contain the enterokinase enzyme which is necessary for activating trypsin, which activates RVs and facilitates viral entry into the cells. The virus particularly affects the middle and the tip of the villi causing destruction and eventually resulting in villous atrophy. The extent of villous atrophy induced by RV is lesser compared to other enteric viral pathogens of pigs such as transmissible gastroenteritis virus (TGEV) or porcine epidemic diarrhea virus (PEDV). The normal villus-height/crypt-depth ratio of the intestinal villi is approximately 7:1. TGEV reduces this ratio to 1:1, whereas RV infection slightly changes it to 5:1. The severity of RV infection depends on the length of the villi and the percentage of enterocytes that are affected. Eventually, the mature columnar epithelial cells on the villi are replaced by immature cuboidal enterocytes that are unable to produce digestive enzymes and have lost their absorptive capabilities.

[0008] RVA has been reported from swine population globally. In the US, RVA infection is most common in pigs 21-55 days old, with slightly less neonatal cases. However, RVA is still a major cause of neonatal diarrhea in piglets worldwide. RVA prevalence ranging from 9.4% to 81.1% have been reported from swine populations in the US. To date, 42 G genotypes and 58 P genotypes have been identified within RVAs. Of these, 12 G genotypes (G1-G6, G8-G12, and G26) and 16 P genotypes (P[1], P[5]-P[8], P[11], P[13]-P[14], P[19], P[23], P[26]-P[27], P[32] and P[34]) have been reported from swine populations. A study reported G5 (71.43%) as the most prevalent RVA genotype in the US followed by G4 (8.19%), G3 (3.57%), G9 (2.31%), G11 (1.89%), G10 (1.26%), and G1 (1.05%). Prevalence of other genotypes (G2, G6, and G10) was less than 1%. Among P genotypes, P[7] was the most common genotype (77.22%) followed by P[6](12.07%), whereas other P types individually constituted less than 1% of the reported RVA genotypes.

[0009] RVC has been emerging as a significant cause of enteritis in neonatal piglets. Porcine RVC was first identified in 1980 and considered as an enteric pathogen with a moderate prevalence rate between 4 and 31%. In the US, RVC is a major cause of diarrhea in neonatal pigs, particularly in piglets younger than 3 days old. In the US, 51.1% of porcine intestinal samples collected during 2009-2011 were positive for RVC. A study reported an overall RVC prevalence of 19.5% in diarrheic and non-diarrheic piglets collected from swine farms located in Ohio, US. The prevalence of RVC was 23.5% among nursing piglets compared to only 8.5% in weaned piglets. Another study from the US detected RVC in 46% samples of porcine origin (feces, fecal

swabs, intestinal, or lung tissues) collected during 2009-2011 in the US and Canada. Of these, RVC was detected in 16% of samples from very young pigs (<3 days old) and 21% of samples from young pigs (4-20 days old). Interestingly, 34% of RVC positive samples were negative for RVA/RVB, and the highest percentage of single RVC infections was in very young (78%, <3 days) and young pigs (65%, 4-22 days) compared to 6-39% in older age groups. Interestingly, single infections of RVC are more common in 0-3 days old piglets, with co-infection with other RV species being more prevalent post-weaning.

[0010] In swine, 15 G genotypes (G12, G13, G8, G6, G5, G14, G9, G1, G17, G15, G7, G10, G3, G18, G16), and 16 P genotypes (P[1], P[5]-P[9], and P[12]-P[21]) of RVC have been identified. The G6 genotype (70%) was the dominant RVC genotype followed by G5 (17%), G1 (12%), and G9 (1%). A study reported higher fecal prevalence (76.1%) of RVC from healthy and diarrheic piglets in the US. A recent study reported presence of RVC in piglets less than 1-week-old in Australian swine herds. Importantly, single RVC genotypes (either G5 or G6) were detected from neonatal piglets; however, older piglets (5-11 weeks) harbored multiple genotypes of RVC (G1 and G3). It is evident that RVC infections are more prevalent among neonatal piglets than weaned piglets, but the reason(s) are not completely understood. While not being bound by theory, likely reasons include lack of an RVC vaccine for use in swine, insufficient maternal RVC antibodies in colostrum or low minimum infectious dose of RVC required for infecting piglets compared to other swine enteric viruses.

[0011] As Rotaviruses exist in multiple genotypes (generally denoted P and G types) due to reassortment of viral genes, a more flexible and temporally responsive vaccine platform is required to match the needs of producers with modern vaccine technology.

SUMMARY OF THE INVENTION

[0012] The present invention provides immunogenic compositions comprising Rotavirus antigens, in particular as a part of subunit vaccines.

[0013] In embodiments, methods for producing Rotavirus-derived immunogenic polypeptides and/or peptides may be mixed or co-expressed with adjuvants are disclosed. Immunogenic compositions may include one or more polypeptides and/or adjuvants as described herein. For example, immunogenic compositions may comprise other antigens that may be used in immunization against pathogens that cause other diseases, such as antigens derived from non-Rotavirus pathogens.

[0014] In embodiments, a process for producing a polypeptide is disclosed including the step of culturing a host cell transformed with a nucleic acid as described herein under conditions which induce polypeptide expression. In a related aspect, a Rotavirus protein may be expressed by recombinant technology and used to develop an immunogenic composition comprising a recombinant antigenic subunit, where such expressed polypeptide is generated using baculovirus/insect cell methodology.

[0015] In one aspect, a process for producing nucleic acid is disclosed, where the nucleic acid encoding a Rotavirus derived protein or polypeptide is prepared (at least in part) by chemical synthesis. In a related aspect, the process includes amplifying nucleic acids using a primer-based amplification method (e.g., PCR).

[0016] In another aspect, a process for producing a protein complex is disclosed, including administering Rotavirus derived polypeptides, or fragments thereof, to a subject. In a related aspect, the process includes admixing Rotavirus-derived polypeptides with a pharmaceutically acceptable carrier or diluent, including an adjuvant. In a related aspect, the adjuvant is a Water/Oil/Water adjuvant. In a further related aspect, the composition may include the set of polypeptides as set forth in SEQ ID NOs:2 and 46 or SEQ ID NOs:2, 4, 6, 8, 10, 12, 14, 16, and 18.

[0017] In embodiments, a method of eliciting an immunological response in a subject is disclosed including administering a composition of the instant disclosure. In a related aspect, the method further includes administering an adjuvant. In a further related aspect, the method includes administering the immunogenic composition to the subject via topical, parenteral or mucosal route.

[0018] In one aspect, the administration may be multiple administrations, where a first immunogenic composition and a second immunogenic composition are the same. In another aspect, the first immunogenic composition and the second immunogenic composition are different.

[0019] In one aspect, administration is performed two or more times.

[0020] In embodiments, a method for treating an infection by a Rotavirus is disclosed including administering to a subject in need thereof a therapeutically effective amount of an immunogenic composition as described herein. In a related aspect, the Rotavirus is of the G5/P[7] genotype for RVA and G6 genotype for RVC. In a further related aspect, the subject is a neonatal piglet.

[0021] In one aspect, multiple therapeutically effective doses of the immunogenic composition are administered to a subject.

[0022] In a related aspect, the method includes mucosally administering a therapeutically effective amount of a first immunogenic composition comprising the set of Rotavirus antigens and topically or parenterally administering a therapeutically effective amount of a second immunogenic composition comprising the set of Rotavirus antigens.

[0023] In one aspect, multiple therapeutically effective doses of the immunogenic composition are administered to a subject. In another aspect, an immunogenic composition comprises a separate, non-Rotavirus antigen.

[0024] These and other embodiments of the instant subject matter as disclosed will readily occur to those of skill in the art in view of the instant disclosure.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] FIGS. 1 *a, b* illustrate the results from the vaccination between MEDGENE™'s (Applicants') vaccine and an RNA-based vaccine in swine.

[0026] FIGS. 2 *a, b* illustrate the Parity results from the vaccination between MEDGENE™'s (Applicants') vaccine and an RNA-based vaccine in swine.

DETAILED DESCRIPTION OF THE INVENTION

[0027] Before the present composition, methods, and methodologies are described, it is to be understood that this invention is not limited to particular compositions, methods, and experimental conditions described, as such compositions, methods, and conditions may vary. It is also to be

understood that the terminology used herein is for purposes of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only in the appended claims.

[0028] As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural references unless the context clearly dictates otherwise. Thus, for example, references to “a nucleic acid” includes one or more nucleic acids, and/or compositions of the type described herein which will become apparent to those persons skilled in the art upon reading this disclosure and so forth.

[0029] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Any methods and materials similar or equivalent to those described herein may be used in the practice or testing of the invention, as it will be understood that modifications and variations are encompassed within the spirit and scope of the instant disclosure.

[0030] As used herein, “about,” “approximately,” “substantially” and “significantly” will be understood by a person of ordinary skill in the art and will vary in some extent depending on the context in which they are used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, “about” and “approximately” will mean plus or minus <10% of particular term and “substantially” and “significantly” will mean plus or minus >10% of the particular term. In embodiments, compositions may “contain,” “comprise” or “consist essentially of” a particular component or group of components, where the skilled artisan would understand the latter to mean the scope of the claim is limited to the specified materials or steps “and those that do not materially affect the basic and novel characteristic(s)” of the claimed invention.

[0031] We have shown that a baculovirus-produced, rotavirus specific bivalent vaccine elicited high titers of neutralizing antibodies against the target Rotavirus A. Mice were immunized with several putative subunit vaccines, and blood collected to analyze immunoreactivity. Mice immunized with a bivalent vaccine developed neutralizing titers exceeding 1:640 whereas control mice failed to develop any titers. Interesting, the inclusion of additional protein targets did not increase vaccine effectiveness, and in fact a trivalent vaccine reduced apparent neutralization titers to 1:320. Based upon this encouraging data, we have continued to examine the specificity of the immune response in vaccination trials. Specifically, mice were vaccinated with baculovirus-produced proteins whose sequences were derived from circulating strains of protein Rotavirus. Sera obtained from these vaccinated animals was then used in standard serum neutralization (SN) assays as a correlate of expected protection. While these experiments are ongoing, we have established critical epitopes within the target proteins as evidenced by effective cross-neutralization in vitro, combined with in silico analysis of predicted protein structure. Based upon the data obtained, it is clear that cross-protection against significantly divergent Rotavirus A isolates can be generated, and these predictive algorithms are currently being tested directly in swine. Given new regulatory pathways, this work opens the door to the availability of targeted vaccines to protect against currently circulating strains of Rotavirus A, and the potential for development of vaccines

against other strains of Rotaviruses (B and C) that can be easily produced using analogous technology.

[0032] As used herein, the terms “Rotavirus” and “Rotavirus-like virus” refer to members of the genus Rotavirus of the family Reoviridae. The term Rotavirus includes strains in all genogroups of the virus. Currently, nine species of Rotavirus referred to as follows: A, B, C, D, E, F, G, H, I, and J. A large number of Rotaviruses isolates have been partially or completely sequenced. The term Rotavirus also includes isolates not characterized at the time of filing.

[0033] The terms “polypeptide” and “protein” refer to a polymer of amino acid residues and are not limited to a minimum length of the product. Thus, peptides, oligopeptides, dimers, multimers, and the like, are included within the definition. Both full-length proteins and fragments thereof are encompassed by the definition. The terms also include post-expression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation and the like. Furthermore, for purposes of the present disclosure, a “polypeptide” refers to a protein which includes modifications, such as deletions, additions and substitutions (generally conservative in nature), to the native sequence, so long as the protein maintains the desired activity. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts which produce the proteins or errors due to PCR amplification.

[0034] “Substantially purified” generally refers to isolation of a substance (compound, polynucleotide, protein, polypeptide, polypeptide composition) such that the substance comprises the majority percent of the sample in which it resides. Typically, in a sample, a substantially purified component comprises about 50%, about 80%-85%, or about 90-95% of the sample. Techniques for purifying polynucleotides and polypeptides of interest are well-known in the art and include, for example, ion-exchange chromatography, affinity chromatography and sedimentation according to density.

[0035] By “isolated” is meant, when referring to a polypeptide, that the indicated molecule is separate and discrete from the whole organism or cell with which the molecule is found in nature or is present in the substantial absence of other biological macro-molecules of the same type. The term “isolated” with respect to a polynucleotide is a nucleic acid molecule devoid, in whole or part, of sequences normally associated with it in nature; or a sequence, as it exists in nature, but having heterologous sequences in association therewith; or a molecule disassociated from the chromosome.

[0036] As used herein, the terms “label” and “detectable label” refer to a molecule capable of detection, including, but not limited to, radioactive isotopes, fluorescers, chemiluminescers, enzymes, enzyme substrates, enzyme cofactors, enzyme inhibitors, chromophores, dyes, metal ions, metal sols, ligands (e.g., biotin or haptens) and the like. The term “fluorescer” refers to a substance or a portion thereof which is capable of exhibiting fluorescence in the detectable range. Particular examples of labels which may be used include fluorescein, rhodamine, dansyl, umbelliferone, Texas red, luminol, acradimium esters, NADPH and α - β -galactosidase.

[0037] “Homology” refers to the percent identity between two polynucleotide or two polypeptide moieties. Two nucleic acid, or two polypeptide sequences are “substan-

tially homologous” to each other when the sequences exhibit at least about 50% sequence identity, at least about 75% sequence identity, at least about 80%-85% sequence identity, at least about 90% sequence identity, and at least about 95%-98% sequence identity over a defined length of the molecules. As used herein, substantially homologous also refers to sequences showing complete identity to the specified sequence.

[0038] In general, “identity” refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Percent identity may be determined by a direct comparison of the sequence information between two molecules by aligning the sequences, counting the exact number of matches between the two aligned sequences, dividing by the length of the shorter sequence, and multiplying the result by 100. Readily available computer programs may be used to aid in the analysis, such as ALIGN, Dayhoff, M. O. in *Atlas of Protein Sequence and Structure* M. O. Dayhoff, ed., 5 Suppl. 3:353-358, National biomedical Research Foundation, Washington, D.C., which adapts the local homology algorithm of Smith and Waterman *Advances in Appl. Math.* 2:482-489, 1981 for peptide analysis. Programs for determining nucleotide sequence identity are available in the Wisconsin Sequence Analysis Package, Version 8 (available from Genetics Computer Group, Madison, Wis.) for example, the BESTFIT, FASTA and GAP programs, which also rely on the Smith and Waterman algorithm. These programs are readily utilized with the default parameters recommended by the manufacturer and described in the Wisconsin Sequence Analysis Package referred to above. For example, percent identity of a particular nucleotide sequence to a reference sequence may be determined using the homology algorithm of Smith and Waterman with a default scoring table and a gap penalty of six nucleotide positions.

[0039] Another method of establishing percent identity in the context of the present disclosure is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm may be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the “Match” value reflects “sequence identity.” Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example, BLASTN and BLASTP may be used using the following default parameters: genetic code=standard; filter=none; strand=both; cutoff=60; expect=10; Matrix=BLOSUM62; Descriptions=50 sequences; sort by=HIGH SCORE; Databases=non-redundant, GenBank+EMBL+DDBJ+PDB+GenBank CDS translations+Swiss protein+Spupdate+PIR. Details of these programs are readily available.

[0040] Alternatively, homology may be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease (s), and size determination of the digested fragments. DNA sequences that are substantially homologous may be identified in a Southern hybridization experiment under, for

example, stringent conditions, as defined for that particular system. Defining appropriate hybridization conditions is within the skill of the art.

[0041] “Recombinant” as used herein to describe a nucleic acid molecule means a polynucleotide of genomic, cDNA, viral, semisynthetic, or synthetic origin which, by virtue of its origin or manipulation, is not associated with all or a portion of the polynucleotide with which it is associated in nature. The term “recombinant” as used with respect to a protein or polypeptide means a polypeptide produced by expression of a recombinant polynucleotide. In general, the gene of interest is cloned and then expressed in transformed organisms, as described further below. The host organism expresses the foreign gene to produce the protein under expression conditions.

[0042] The term “transformation” refers to the insertion of an exogenous polynucleotide into a host cell, irrespective of the method used for the insertion. For example, direct uptake, transduction or f-mating are included. The exogenous polynucleotide may be maintained as a non-integrated vector, for example, a plasmid, or alternatively, may be integrated into the host genome.

[0043] “Recombinant host cells,” “host cells,” “cells,” “cell lines,” “cell cultures,” and other such terms denoting microorganisms or higher eukaryotic cell lines cultured as unicellular entities refer to cells which may be, or have been, used as recipients for recombinant vector or other transferred DNA, and include the original progeny of the original cell which has been transfected.

[0044] A “coding sequence” or a sequence which “encodes” a selected polypeptide, is a nucleic acid molecule which is transcribed (in the case of DNA) and translated (in the case of mRNA) into a polypeptide in vivo when placed under the control of appropriate regulatory sequences (or “control elements”). The boundaries of the coding sequence may be determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A coding sequence may include, but is not limited to, cDNA from viral, prokaryotic or eukaryotic mRNA, genomic DNA sequences from viral or prokaryotic DNA, and even synthetic DNA sequences. A transcription termination sequence may be located 3' to the coding sequence.

[0045] Typical “control elements,” include, but are not limited to, transcription promoters, transcription enhancer elements, transcription termination signals, polyadenylation sequences (located 3' to the translation stop codon), sequences for optimization of initiation of translation (located 5' to the coding sequence), and translation termination sequences.

[0046] The term “nucleic acid” includes DNA and RNA, and also their analogues, such as those containing modified backbones (e.g., phosphorothioates, and the like), and also peptide nucleic acids (PNA), and the like. The present disclosure provides nucleic acids comprising sequences complementary to those described above (e.g., for antisense or probing purposes).

[0047] “Operably linked” refers to an arrangement of elements wherein the components so described are configured so as to perform their usual function. Thus, a given promoter operably linked to a coding sequence is capable of effecting the expression of the coding sequence when the proper enzymes are present. The promoter need not be contiguous with the coding sequence, so long as it functions to direct the expression thereof. Thus, for example, inter-

vening untranslated yet transcribed sequences may be present between the promoter sequence and the coding sequence and the promoter sequence may still be considered “operably linked” to the coding sequence.

[0048] “Encoded by” refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 to 5 amino acids, at least 8 to 10 amino acids, and at least 15 to 20 amino acids from a polypeptide encoded by the nucleic acid sequence.

[0049] “Expression cassette” or “expression construct” refers to an assembly which is capable of directing the expression of the sequence(s) or gene(s) of interest. An expression cassette generally includes control elements, as described above, such as a promoter which is operably linked to (so as to direct transcription of) the sequence(s) or gene(s) of interest, and often includes a polyadenylation sequence as well. In embodiments, the expression cassette described herein may be contained within a plasmid construct. In addition to the components of the expression cassette, the plasmid construct may also include, one or more selectable markers, a signal which allows the plasmid construct to exist as single-stranded DNA (e.g., a M13 origin of replication), at least one multiple cloning site, and a “mammalian” origin of replication (e.g., a SV40 or adenovirus origin of replication).

[0050] “Purified polynucleotide” refers to a polynucleotide of interest or fragment thereof which is essentially free, e.g., contains less than about 50%, less than about 70%, and less than about at least 90%, of the protein with which the polynucleotide is naturally associated. Techniques for purifying polynucleotides of interest are well-known in the art and include, for example, disruption of the cell containing the polynucleotide with a chaotropic agent and separation of the polynucleotide(s) and proteins by ion-exchange chromatography, affinity chromatography and sedimentation according to density.

[0051] The term “transfection” is used to refer to the uptake of foreign DNA by a cell. A cell has been “transfected” when exogenous DNA has been introduced inside the cell membrane. A number of transfection techniques are generally known in the art. Such techniques may be used to introduce one or more exogenous DNA moieties into suitable host cells. The term refers to both stable and transient uptake of the genetic material, and includes uptake of peptide- or antibody-linked DNAs.

[0052] A “vector” is capable of transferring nucleic acid sequences to target cells (e.g., viral vectors, non-viral vectors, particulate carriers, and liposomes). Typically, “vector construct,” “expression vector,” and “gene transfer vector,” mean any nucleic acid construct capable of directing the expression of a nucleic acid of interest and which may transfer nucleic acid sequences to target cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

[0053] By “fragment” is intended a molecule consisting of only a part of the intact full-length sequence and structure. A fragment of a polypeptide may include a C-terminal deletion, an N-terminal deletion, and/or an internal deletion of the native polypeptide. A fragment of a polypeptide will generally include at least about 5-10 contiguous amino acid residues of the full-length molecule, at least about 15-25 contiguous amino acid residues of the full-length molecule, and at least about 20-50 or more contiguous amino acid

residues of the full-length molecule, or any integer between 5 amino acids and the number of amino acids in the full-length sequence, provided that the fragment in question retains the ability to elicit the desired biological response. A fragment of a nucleic acid may include a 5'-deletion, a 3'-deletion, and/or an internal deletion of a nucleic acid. Nucleic acid fragments will generally include at least about 5-1000 contiguous nucleotide bases of the full-length molecule and may include at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides of the full-length molecule, or any integer between 5 nucleotides and the number of nucleotides in the full-length sequence. Such fragments may be useful in hybridization, amplification, production of immunogenic fragments, or nucleic acid immunization.

[0054] By “immunogenic fragment” is meant a fragment of an immunogen which includes one or more epitopes and thus may modulate an immune response or may act as an adjuvant for a co-administered antigen. Such fragments may be identified using any number of epitope mapping techniques, well known in the art. For example, linear epitopes may be determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the peptides with antibodies while the peptides are still attached to the supports. Such techniques are known in the art and described in, e.g., U.S. Pat. No. 4,708,871, incorporated herein by reference in its entirety. Similarly, conformational epitopes are readily identified by determining spatial conformation of amino acids such as by, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance.

[0055] Immunogenic fragments, for purposes of the present disclosure, will usually be at least about 2 amino acids in length, about 5 amino acids in length, and at least about 10 to about 15 amino acids in length. There is no critical upper limit to the length of the fragment, which could comprise nearly the full-length of the protein sequence, or even a fusion protein comprising two or more epitopes.

[0056] As used herein, the term “epitope” generally refers to the site on an antigen which is recognized by a T-cell receptor and/or an antibody. In embodiments, it is a short peptide derived from or as part of a protein antigen. However, the term is also intended to include peptides with glycopeptides and carbohydrate epitopes. Several different epitopes may be carried by a single antigenic molecule. The term “epitope” also includes modified sequences of amino acids or carbohydrates which stimulate responses which recognize the whole organism. It is advantageous if the selected epitope is an epitope of an infectious agent, which agent causes the infectious disease.

[0057] The epitope may be generated from knowledge of the amino acid and corresponding DNA sequences of the peptide or polypeptide, as well as from the nature of particular amino acids (e.g., size, charge, and the like) and the codon dictionary, without undue experimentation. Some guidelines in determining whether a protein will stimulate a response, include: Peptide length—the peptide is about 8 or 9 amino acids long to fit into the MHC class I complex and about 13-25 amino acids long to fit into a class II MHC complex. This length is a minimum for the peptide to bind to the MHC complex. In one aspect, the peptides may be longer than these lengths because cells may cut peptides. The peptide may contain an appropriate anchor motif which

will enable it to bind to the various class I or class II molecules with high enough specificity to generate an immune response. This may be done, without undue experimentation, by comparing the sequence of the protein of interest with published structures of peptides associated with the MHC molecules. Thus, the skilled artisan may ascertain an epitope of interest by comparing the protein sequence with sequences listed in the protein database.

[0058] As used herein, the term “T cell epitope” refers generally to those features of a peptide structure which are capable of inducing a T cell response and a “B cell epitope” refers generally to those features of a peptide structure which are capable of inducing a B cell response.

[0059] An “immunological response” to an antigen or composition is the development in a subject of a humoral and/or a cellular immune response to an antigen present in the composition of interest. For purposes of the present disclosure, a “humoral immune response” refers to an immune response mediated by antibody molecules, while a “cellular immune response” is one mediated by T-lymphocytes and/or other white blood cells. One important aspect of cellular immunity involves an antigen-specific response by cytolytic T-cells (“CTL”s). CTLs have specificity for peptide antigens that are presented in association with proteins encoded by the major histocompatibility complex (MHC) and expressed on the surfaces of cells. CTLs help induce and promote the destruction of intracellular microbes, or the lysis of cells infected with such microbes. Another aspect of cellular immunity involves an antigen-specific response by helper T-cells. Helper T-cells act to help stimulate the function, and focus the activity of, nonspecific effector cells against cells displaying peptide antigens in association with MHC molecules on their surface. A “cellular immune response” also refers to the production of cytokines, chemokines and other such molecules produced by activated T-cells and/or other white blood cells, including those derived from CD4+ and CD8+ T-cells.

[0060] A composition or vaccine that elicits a cellular immune response may serve to sensitize a vertebrate subject by the presentation of antigen in association with MHC molecules at the cell surface. The cell-mediated immune response is directed at, or near, cells presenting antigen at their surface. In addition, antigen-specific T-lymphocytes may be generated to allow for the future protection of an immunized host.

[0061] The ability of a particular antigen to stimulate a cell-mediated immunological response may be determined by a number of assays, such as by lymphoproliferation (lymphocyte activation) assays, CTL cytotoxic cell assays, or by assaying for T-lymphocytes specific for the antigen in a sensitized subject. Such assays are well known in the art. Recent methods of measuring cell-mediated immune response include measurement of intracellular cytokines or cytokine secretion by T-cell populations, or by measurement of epitope specific T-cells.

[0062] Thus, an immunological response as used herein may be one that stimulates the production of antibodies (e.g., neutralizing antibodies that block bacterial toxins and pathogens such as viruses entering cells and replicating by binding to toxins and pathogens, typically protecting cells from infection and destruction). The antigen of interest may also elicit production of CTLs. Hence, an immunological response may include one or more of the following effects: the production of antibodies by B-cells; and/or the activation

of suppressor T-cells and/or memory/effector T-cells directed specifically to an antigen or antigens present in the composition or vaccine of interest. These responses may serve to neutralize infectivity, and/or mediate antibody-complement, or antibody dependent cell cytotoxicity (ADCC) to provide protection to an immunized host. Such responses may be determined using standard immunoassays and neutralization assays, well known in the art. The innate immune system of mammals also recognizes and responds to molecular features of pathogenic organisms via activation of Toll-like receptors and similar receptor molecules on immune cells. Upon activation of the innate immune system, various non-adaptive immune response cells are activated to, e.g., produce various cytokines, lymphokines and chemokines. Cells activated by an innate immune response include immature and mature Dendritic cells of the monocyte and plasmacytoid lineage (MDC, PDC), as well as gamma, delta, alpha and beta T cells and B cells and the like. Thus, the present disclosure also contemplates an immune response wherein the immune response involves both an innate and adaptive response.

[0063] An “immunogenic composition” is a composition that comprises an antigenic molecule where administration of the composition to a subject results in the development in the subject of a humoral and/or a cellular immune response to the antigenic molecule of interest.

[0064] The terms “immunogenic” protein or polypeptide refer to an amino acid sequence which elicits an immunological response as described above. An “immunogenic” protein or polypeptide, as used herein, includes the full-length sequence of the protein in question, including the precursor and mature forms, analogs thereof, or immunogenic fragments thereof.

[0065] “Gene transfer” or “gene delivery” refers to methods or systems for reliably inserting DNA or RNA of interest into a host cell. Such methods may result in transient expression of non-integrated transferred DNA, extrachromosomal replication and expression of transferred replicons (e.g., episomes), or integration of transferred genetic material into the genomic DNA of host cells. Gene delivery expression vectors include, but are not limited to, vectors derived from bacterial plasmid vectors, viral vectors, non-viral vectors, alphaviruses, pox viruses and vaccinia viruses. When used for immunization, such gene delivery expression vectors may be referred to as vaccines or vaccine vectors.

[0066] The term “derived from” is used herein to identify the original source of a molecule but is not meant to limit the method by which the molecule is made which may be, for example, by chemical synthesis or recombinant means.

[0067] Generally, a viral polypeptide is “derived from” a particular polypeptide of a virus (viral polypeptide) if it is (i) encoded by an open reading frame of a polynucleotide of that virus (viral polynucleotide), or (ii) displays sequence identity to polypeptides of that virus as described above.

[0068] A polynucleotide “derived from” a designated sequence refers to a polynucleotide sequence which comprises a contiguous sequence of approximately at least about 6 nucleotides, at least about 8 nucleotides, at least about 10-12 nucleotides, and at least about 15-20 nucleotides corresponding, i.e., identical or complementary to, a region of the designated nucleotide sequence. The derived polynucleotide will not necessarily be derived physically from the nucleotide sequence of interest, but may be generated in any manner, including, but not limited to, chemical synthe-

sis, replication, reverse transcription or transcription, which is based on the information provided by the sequence of bases in the region(s) from which the polynucleotide is derived. As such, it may represent either a sense or an antisense orientation of the original polynucleotide.

[0069] A Rotavirus polynucleotide, oligonucleotide, nucleic acid, protein, polypeptide, or peptide, as defined above, is a molecule derived from a Rotavirus, respectively, including, without limitation, any of the various isolates of Rotavirus. The molecule need not be physically derived from the particular isolate in question, but may be synthetically or recombinantly produced.

[0070] The Rotavirus genome consists of eleven segments of ds RNA encoding six structural proteins VP1-VP4, VP6 and VP7 and five non-structural proteins NSP1-NSP5/6.

[0071] Nucleic acid and protein sequences for a number of Rotavirus isolates are also known. Representative Rotavirus sequences are presented in SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17 and 45. Additional representative sequences, including sequences of Rotavirus, and their encoded polypeptides from Rotavirus isolates are listed in the National Center for Biotechnology Information (NCBI) database. See, for example, but not limited to, GenBank entries: BAA03855.1; BAF80156.1, CAA67568.1, BAF59645.1, X99126.1, AB275386.1, D16352.1 and AB008292.1, all of which sequences (as entered by the date of filing of this application) are herein incorporated by reference.

[0072] As used herein, the terms “capsid protein” or “capsid polypeptide” or “outer capsid” in reference to a Rotavirus refer to VP4 or VP7 comprising a sequence homologous or identical to the capsid polypeptide of a Rotavirus VP4 or VP7, and include sequences displaying at least about 80-100% sequence identity thereto, including any percent identity within these ranges, such as 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% or greater, but less than 100% sequence identity thereto.

[0073] As used herein, the terms “minor structural protein” or “minor structural polypeptide” or “inner or middle capsid” in reference to a Rotavirus refer to a polypeptide comprising VP1, VP2, VP3, and VP6 sequence homologous or identical to the polypeptide encoded by the open reading frame following the coding region for the capsid protein in the Rotavirus genome, and include sequences displaying at least about 80-100% sequence identity thereto, including any percent identity within these ranges, such as 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% or greater, but less than 100% sequence identity thereto.

[0074] An “antigen” refers to a molecule containing one or more epitopes (either linear, conformational or both) that will stimulate a host’s immune-system to make a humoral and/or cellular antigen-specific response. The term is used interchangeably with the term “immunogen.” Normally, a B-cell epitope will include at least about 5 amino acids but may be as small as 3-4 amino acids. A T-cell epitope, such as a CTL epitope, will include at least about 7-9 amino acids, and a helper T-cell epitope at least about 12-20 amino acids. Normally, an epitope will include between about 7 and 15 amino acids, such as 9, 10, 12 or 15 amino acids. The term “antigen” denotes both subunit antigens (i.e., antigens which are separate and discrete from a whole organism with which the antigen is associated in nature), as well as, killed, attenuated or inactivated bacteria, viruses, fungi, parasites or other microbes. Antibodies such as anti-idiotypic antibodies,

or fragments thereof, and synthetic peptide mimotopes, which may mimic an antigen or antigenic determinant, are also captured under the definition of antigen as used herein. Similarly, an oligonucleotide or polynucleotide which expresses an antigen or antigenic determinant *in vivo*, such as in gene therapy and DNA immunization applications, is also included in the definition of antigen herein.

[0075] The term “antibody” encompasses polyclonal and monoclonal antibody preparations, as well as preparations including hybrid antibodies, altered antibodies, chimeric antibodies and, humanized antibodies, as well as: hybrid (chimeric) antibody molecules and any functional fragments obtained from such molecules, wherein such fragments retain specific-binding properties of the parent antibody molecule.

[0076] The terms “hybridize” and “hybridization” refer to the formation of complexes between nucleotide sequences which are sufficiently complementary to form complexes via Watson-Crick base pairing. Where a primer “hybridizes” with target (template), such complexes (or hybrids) are sufficiently stable to serve the priming function required by, e.g., the DNA polymerase to initiate DNA synthesis.

[0077] As used herein, a “biological sample” refers to a sample of tissue or fluid isolated from a subject, including but not limited to, for example, blood, plasma, serum, fecal matter, urine, bone marrow, bile, spinal fluid, lymph fluid, samples of the skin, external secretions of the skin, respiratory, intestinal, and genitourinary tracts, tears, saliva, milk, blood cells, organs, biopsies and also samples of *in vitro* cell culture constituents including but not limited to conditioned media resulting from the growth of cells and tissues in culture medium, e.g., recombinant cells, and cell components. In particular, Rotavirus may be obtained from biological samples including, but not limited to, oral or fecal matter.

[0078] By “subject” is meant any member of the family Suidae, genus *sus*, and more specifically, *Sus domesticus*. The term does not denote a particular age. Thus, both adult and newborn individuals are intended to be covered, as well as females.

[0079] The terms “variant,” “analog” and “mutein” refer to biologically active derivatives of the reference molecule that retain desired activity, such as antigenic activity in inducing an immune response against Rotavirus. In general, the terms “variant” and “analog” refer to compounds having a native polypeptide sequence and structure with one or more amino acid additions, substitutions (generally conservative in nature) and/or deletions, relative to the native molecule, so long as the modifications do not destroy biological activity and which are “substantially homologous” to the reference molecule as defined below. In general, the amino acid sequences of such analogs will have a high degree of sequence homology to the reference sequence, e.g., amino acid sequence homology of more than 50%, generally more than 60%-70%, even more particularly 80%-85% or more, such as at least 90%-95% or more, when the two sequences are aligned. Often, the analogs will include the same number of amino acids but will include substitutions, as explained herein. The term “mutein” further includes polypeptides having one or more amino acid-like molecules including but not limited to compounds comprising only amino and/or imino molecules, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, and the like), polypeptides

with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring (e.g., synthetic), cyclized, branched molecules and the like. The term also includes molecules comprising one or more N-substituted glycine residues (a “peptoid”) and other synthetic amino acids or peptides. (See, e.g., U.S. Pat. Nos. 5,831,005; 5,877,278; and 5,977,301). In embodiments, the analog or mutein has at least the same antigenic activity as the native molecule. Methods for making polypeptide analogs and muteins are known in the art and are described further below.

[0080] As explained above, analogs generally include substitutions that are conservative in nature, i.e., those substitutions that take place within a family of amino acids that are related in their side chains. Specifically, amino acids are generally divided into four families: (1) acidic—aspartate and glutamate; (2) basic—lysine, arginine, histidine; (3) non-polar—alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar—glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. Phenylalanine, tryptophan, and tyrosine are sometimes classified as aromatic amino acids. For example, it is reasonably predictable that an isolated replacement of leucine with isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar conservative replacement of an amino acid with a structurally related amino acid, will not have a major effect on the biological activity. For example, the polypeptide of interest may include up to about 5-10 conservative or non-conservative amino acid substitutions, or even up to about 15-25 conservative or non-conservative amino acid substitutions, or any integer between 5-25, so long as the desired function of the molecule remains intact. One of skill in the art may readily determine regions of the molecule of interest that may tolerate change by reference to Hopp/Woods and Kyte-Doolittle plots, well known in the art.

[0081] The term “multiple epitope fusion antigen” or “multiple epitope fusion protein” as used herein intends a polypeptide in which multiple Rotavirus antigens are part of a single, continuous chain of amino acids, which chain does not occur in nature. The Rotavirus antigens may be connected directly to each other by peptide bonds or may be separated by intervening amino acid sequences. The fusion antigens may also contain sequences exogenous to the Rotavirus. Moreover, the sequences present may be from multiple genotypes and/or isolates of Rotavirus.

[0082] By “therapeutically effective amount” in the context of the immunogenic compositions is meant an amount of an immunogen (e.g., immunogenic polypeptide, fusion protein, polyprotein, or nucleic acid encoding an antigen) which will induce an immunological response, either for antibody production or for treatment or prevention of Rotavirus infection. Such a response will generally result in the development in the subject of an antibody-mediated and/or a secretory or cellular immune response to the composition. Usually, such a response includes but is not limited to one or more of the following effects; the production of antibodies from any of the immunological classes, such as immunoglobulins A, D, E, G or M; the proliferation of B and T lymphocytes; the provision of activation, growth and differentiation signals to immunological cells; expansion of helper T cell, suppressor T cell, and/or cytotoxic T cell and/or γ , δ -T cell populations.

[0083] For purposes of the present disclosure, an “effective amount” of an adjuvant will be that amount which enhances an immunological response to a co-administered antigen or nucleic acid encoding an antigen.

[0084] As used herein, “treatment” refers to any of (i) the prevention of infection or reinfection, as in a traditional vaccine, (ii) the reduction or elimination of symptoms, and (iii) the substantial or complete elimination of the pathogen in question. Treatment may be effected prophylactically (prior to infection) or therapeutically (following infection).

[0085] Before describing the present disclosure in detail, it is to be understood that the practice of the present disclosure will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, recombinant DNA techniques and immunology all of which are within the ordinary skill of the art. Such techniques are explained fully in the literature. Although a number of methods and materials similar or equivalent to those described herein may be used in the practice of the present invention as claimed, the materials and methods are described herein.

[0086] The present disclosure includes compositions and methods for immunizing a subject against Rotavirus infection. The instant disclosure provides immunogenic compositions comprising nucleic acids encoding capsid proteins and/or other immunogenic polypeptides from one or more strains of Rotavirus, compositions comprising immunogenic polypeptides derived from one or more strains of Rotavirus. Immunogenic polypeptides to be used in the practice of the instant subject matter may include Rotavirus-derived polypeptides, including multiple epitope fusion antigens, and/or Rotavirus-encoded polyproteins. In addition, immunogenic compositions may comprise one or more adjuvants or nucleic acids encoding adjuvants, wherein immunogenic polypeptides are mixed or co-expressed with adjuvants. Immunogenic compositions may also comprise additional antigens other than Rotavirus antigens, such as antigens that may be used in immunization against pathogens that cause diarrheal diseases.

[0087] In order to further an understanding of the subject matter as disclosed, a more detailed discussion is provided below regarding the production of nucleic acids and polypeptides for use in immunogenic compositions and methods of using such compositions in the treatment or prevention of Rotavirus infection.

Structural Polypeptides, Nonstructural Polypeptides, and Polyproteins

[0088] The immunogenic compositions described herein may comprise one or more polypeptides derived from one or more genotypes and/or isolates of Rotavirus. Polypeptides that may be used in the practice of the subject matter as disclosed herein include structural proteins, nonstructural proteins, and polyproteins. Such polypeptides may be full-length proteins or variants or immunogenic fragments thereof capable of eliciting an immune response to a Rotavirus.

[0089] The polypeptides in immunogenic compositions may be encoded by any region of a Rotavirus genome. Multiple polypeptides may be included in immunogenic compositions. Such compositions may comprise polypeptides from the same Rotavirus isolate or from different strains and isolates, including isolates having any of the various species of Rotavirus (e.g., Rotavirus A, B, C, D, E,

F, G, H, I or J), to provide increased protection against a broad range of Rotavirus species. Multiple viral strains of Rotavirus are known (or not), and multiple polypeptides comprising epitopes derived from any of these strains may be used in immunogenic compositions.

[0090] The antigens used in the immunogenic compositions of the present disclosure may be present in the composition as individual separate polypeptides. Generally, the recombinant proteins of the present disclosure are expressed as a GST-fusion protein and/or a His-tagged fusion protein.

Multi-epitope Fusion Proteins

[0091] The immunogenic compositions described herein may also comprise multiple epitope fusion proteins. Such fusion proteins include multiple epitopes derived from two or more viral polypeptides of one or more genotypes and/or isolates of Rotavirus. Multiple epitope fusion proteins offer two principal advantages: first, a polypeptide that may be unstable or poorly expressed on its own may be assisted by adding a suitable hybrid partner that overcomes the problem; second, commercial manufacture is simplified as only one expression and purification need be employed in order to produce two polypeptides which are both antigenically useful.

[0092] Multi-epitope fusion proteins may contain one or more of the various domains of Rotavirus full-length proteins VP4 and/or VP7 (also referred to herein as Rotavirus major structural proteins) and/or VP1, VP2, VP3, and/or VP6 (also referred to herein as a Rotavirus minor structural protein); and NSP1-NSP5/6 (also referred to herein as a Rotavirus non-structural protein), or fragments thereof, derived from one or more Rotavirus isolates. The polypeptides in fusion proteins may be derived from the same Rotavirus isolate or from different strains and isolates, including isolates having any of the various Rotavirus species, to provide increased protection against a broad range of Rotavirus species. Multiple viral strains of Rotavirus are known, and epitopes derived from any of these strains may be used in a fusion protein.

[0093] It is well known that any given species of organism varies from one individual organism to another and further that a given organism such as a virus may have a number of different strains. For example, as explained above, Rotavirus includes at least nine species (A-J). Each strain includes a number of antigenic determinants that are in homologous regions present in all species of Rotavirus but are slightly different from one viral strain to another. Thus, a multiple epitope fusion antigen may include multiple polypeptides from different viral strains of Rotavirus, each comprising a particular homologous region but each having a different form of an antigenic determinant. In general, antigenic determinants may have a high degree of homology in terms of amino acid sequence, which degree of homology is generally 30% or more, 40% or more, when aligned.

[0094] Multiple epitope fusion antigens may be represented by the formula $\text{NH}_2\text{-A}\{-\text{X-L}\}_n\text{-B-COOH}$, wherein: X is an amino acid sequence of a Rotavirus antigen or a fragment thereof; L is an optional linker amino acid sequence; A is an optional N-terminal amino acid sequence; B is an optional C-terminal amino acid sequence; and n is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.

[0095] If an —X— moiety has a leader peptide sequence in its wild-type form, this may be included or omitted in the multiple epitope fusion antigen. In some embodiments, the

leader peptides will be deleted except for that of the —X— moiety located at the N-terminus of the hybrid protein i.e., the leader peptide of X_1 will be retained, but the leader peptides of $X_2 \dots X_n$ will be omitted. This is equivalent to deleting all leader peptides and using the leader peptide of X_1 as moiety -A-.

[0096] For each n instances of (—X-L—), linker amino acid sequence -L- may be present or absent. For instance, when $n=2$ the hybrid may be $\text{NH}_2\text{-X}_1\text{-L}_1\text{-X}_2\text{-L}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-X}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-L}_1\text{-X}_2\text{-COOH}$, $\text{NH}_2\text{-X}_1\text{-X}_2\text{-L}_2\text{-COOH}$, and the like. Linker amino acid sequence(s)-L- will typically be short, e.g., 20 or fewer amino acids (i.e., 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include short peptide sequences which facilitate cloning, poly-glycine linkers (Gly, where $n=2, 3, 4, 5, 6, 7, 8, 9, 10$ or more), and histidine tags (His_n where $n=3, 4, 5, 6, 7, 8, 9, 10$ or more). Other suitable linker amino acid sequences will be apparent to those skilled in the art. A useful linker is GSGGGG, with the Gly-Ser dipeptide being formed from a BamHI restriction site, which aids cloning and manipulation, and the (Gly)₄ tetrapeptide being a typical poly-glycine linker. In addition, protease substrate sequences may also be added (e.g., TEV protease: ENLYFQG).

[0097] -A- is an optional N-terminal amino acid sequence. This will typically be short, e.g., 40 or fewer amino acids (i.e., 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include leader sequences to direct protein trafficking or short peptide sequences which facilitate cloning or purification (e.g., a histidine tag His_n where $n=3, 4, 5, 6, 7, 8, 9, 10$ or more). Other suitable N-terminal amino acid sequences will be apparent to those skilled in the art. If X_1 lacks its own N-terminus methionine, -A- is an oligopeptide (e.g., with 1, 2, 3, 4, 5, 6, 7 or 8 amino acids) which provides a N-terminus methionine.

[0098] —B— is an optional C-terminal amino acid sequence. This will typically be short, e.g., 40 or fewer amino acids (i.e., 40, 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include sequences to direct protein trafficking, short peptide sequences which facilitate cloning or purification (e.g., His_n where $n=3, 4, 5, 6, 7, 8, 9, 10$ or more), or sequences which enhance protein stability. Other suitable C-terminal amino acid sequences will be apparent to those skilled in the art, including that such His_n sequences may be removed when a TEV protease substrate sequence precedes it (e.g., ENLYFQGH_n).

[0099] The individual antigens of the immunogenic composition within the multiple epitope fusion antigen (individual —X— moieties) may be from one or more strains or from one or more M types. Where $n=2$, for instance, X_2 may be from the same strain or type as X_1 or from a different strain or type. Where $n=3$, the strains might be (i) $X_1=X_2=X_3$, (ii) $X_1=X_2$ not equal to X_3 , (iii) X_1 not equal to $X_2=X_3$, (iv) X_1 not equal to X_2 not equal to X_3 , or (v) $X_1=X_3$ not equal to X_2 , and the like.

[0100] Where multiple epitope fusion antigens are used, the individual antigens within the fusion protein (i.e., individual —X— moieties) may be from one or more strains. Where $n=2$, for instance, X_2 may be from the same strain as X_1 or from a different strain. Where $n=3$, the strains might be (i) $X_1=X_2=X_3$ (ii) $X_1=X_2$ not equal to X_3 (iii) X_1 not equal

to $X_2=X_3$ (iv) X_1 not equal to X_2 not equal to X_3 or (v) $X_1=X_3$ not equal to X_2 , and the like.

[0101] Accordingly, in embodiments, antigenic determinants from different Rotavirus species may be present. Representative multiepitope fusion proteins for use in the present disclosure, comprising polypeptides derived from Rotavirus isolates, are discussed below. However, it is to be understood that multiepitope fusion proteins comprising other epitopes derived from Rotavirus genomes or multiepitope fusion proteins comprising different arrangements of epitopes will also find use in immunogenic compositions as disclosed.

[0102] In certain embodiments, the fusion protein comprises one or more capsid and/or minor structural polypeptides from one or more isolates of Rotavirus.

[0103] In another embodiment, the fusion protein comprises Rotavirus polypeptides from more than one Rotavirus strain.

[0104] In all fusions described herein, the viral regions need not be in the order in which they occur naturally. Moreover, each of the regions may be derived from the same or different Rotavirus isolates. The various Rotavirus polypeptides present in the various fusions described above may either be full-length polypeptides or portions thereof.

[0105] If desired, the fusion proteins, or the individual components of these proteins, also may contain other amino acid sequences, such as amino acid linkers or signal sequences, as well as ligands useful in protein purification, such as glutathione-S-transferase and staphylococcal protein A.

Nucleic Acids

[0106] Nucleic acids for use as disclosed herein, for example, in polypeptide production, may be derived from any of the various regions of a Rotavirus genome. Representative sequences from Rotavirus isolates are listed herein. Thus, nucleic acids for use as disclosed herein include those derived from one or more sequences from any pathogenic Rotavirus species or isolate.

[0107] Representative sequences from Rotavirus include SEQ ID NOs:1, 3, 5, 7, 9, 11, 13, 15, 17 and 45.

[0108] Any of these sequences, as well as fragments and variants thereof that may be used in nucleic acid immunization to elicit an immune response to a Rotavirus will find use in the present methods. Thus, the present disclosure provides variants of the above sequences displaying at least about 80-100% sequence identity thereto, including any percent identity within these ranges, such as 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99% or greater, but less than 100% sequence identity thereto. The present disclosure also provides polynucleotides encoding immunogenic fragments of a Rotavirus polypeptide derived from any of the above sequences or a variant thereof. Polynucleotides may also comprise coding sequences for polypeptides which occur naturally or may be artificial sequences which do not occur in nature.

[0109] Polynucleotides may contain less than an entire Rotavirus genome, or alternatively may include the sequence of an entire viral genomic RNA. For example, polynucleotides may comprise one or more sequences from the Rotavirus genome. Polynucleotides may also comprise the entire viral genomic Rotavirus or less than the entire viral genomic Rotavirus from multiple species and/or isolates of Rotavirus.

[0110] In embodiments, polynucleotides comprise an Rotavirus sequence coding for the full-length polyprotein of a Rotavirus. In other embodiments, polynucleotides comprise one or more portions of the Rotavirus RNA sequence of a Rotavirus.

[0111] In another example, a polynucleotide may comprise a nucleotide sequence encoding a portion of a Rotavirus polyprotein.

[0112] In embodiments, the polynucleotides comprise sequences encoding one or more capsid proteins of a Rotavirus. For example, polynucleotides may comprise one or more sequences coding for structural proteins (e.g., VP4 and VP7) of a Rotavirus. In certain embodiments, polynucleotides comprise sequences coding for at least two capsid proteins from multiple species and/or isolates of Rotavirus.

[0113] In embodiments, polynucleotides comprise one or more Rotavirus sequences coding for the capsid proteins of one or more isolates of Rotavirus. In certain embodiments, polynucleotides comprise one or more sequences coding for the capsid proteins of one or more isolates of Rotavirus.

[0114] In embodiments, the present disclosure provides polynucleotides encoding a multiepitope fusion protein as described herein. Multiepitope fusion proteins may comprise sequences from one or more species and/or isolates of Rotavirus.

[0115] Nucleic acids according to the instant disclosure may be prepared in many ways (e.g., by chemical synthesis, from genomic or cDNA libraries, from the organism itself, etc.) and may take various forms (e.g., single stranded, double stranded, vectors, probes, and the like). In embodiments, nucleic acids are prepared in substantially pure form (i.e., substantially free from other host cell or non-host cell nucleic acids).

[0116] For example, nucleic acids may be obtained by screening cDNA and/or genomic libraries from cells infected with virus, or by deriving the gene from a vector known to include the same. For example, polynucleotides of interest may be isolated from a genomic library derived from viral RNA, present in, for example, hair or blood samples from an infected individual. Alternatively, Rotavirus nucleic acids may be isolated from infected mammals or from biological samples collected from infected individuals. An amplification method such as PCR may be used to amplify polynucleotides from either Rotavirus genomic RNA or cDNA encoding therefor. Alternatively, polynucleotides may be synthesized in the laboratory, for example, using an automatic synthesizer. The nucleotide sequence may be designed with the appropriate codons for the particular amino acid sequence desired. In general, one will select preferred codons for the intended host in which the sequence will be expressed. The complete sequence of the polynucleotide of interest may be assembled from overlapping oligonucleotides prepared by standard methods and assembled into a complete coding sequence. The polynucleotides may be RNA or single- or double-stranded DNA. In embodiments, the polynucleotides are isolated free of other components, such as proteins and lipids.

[0117] Thus, particular nucleotide sequences may be obtained from vectors harboring the desired sequences or synthesized completely or in part using various oligonucleotide synthesis techniques known in the art, such as site-directed mutagenesis and polymerase chain reaction (PCR) techniques where appropriate. In particular, one method of obtaining nucleotide sequences encoding the desired

sequences is by annealing complementary sets of overlapping synthetic oligonucleotides produced in a conventional, automated polynucleotide synthesizer, followed by ligation with an appropriate DNA ligase and amplification of the ligated nucleotide sequence via PCR. Primer sequences may include, but are not limited to, SEQ ID NOs:19-44 and 47-52.

Production of Immunogenic Polypeptides

[0118] Polypeptides described herein may be prepared in any suitable manner (e.g., recombinant expression, purification from cell culture, chemical synthesis, and the like) and in various forms (e.g., native, fusions, non-glycosylated, lipidated, and the like). Such polypeptides include naturally-occurring polypeptides, recombinantly produced polypeptides, synthetically produced polypeptides, or polypeptides produced by a combination of these methods. Means for preparing such polypeptides are well understood in the art. Polypeptides are prepared in substantially pure form (i.e., substantially free from other host cell or non-host cell proteins).

[0119] Polypeptides may be conveniently synthesized chemically, by any of several techniques that are known to those skilled in the peptide art. In general, these methods employ the sequential addition of one or more amino acids to a growing peptide chain. Normally, either the amino or carboxyl group of the first amino acid is protected by a suitable protecting group. The protected or derivatized amino acid may then be either attached to an inert solid support or utilized in solution by adding the next amino acid in the sequence having the complementary (amino or carboxyl) group suitably protected, under conditions that allow for the formation of an amide linkage. The protecting group is then removed from the newly added amino acid residue and the next amino acid (suitably protected) is then added, and so forth. After the desired amino acids have been linked in the proper sequence, any remaining protecting groups (and any solid support, if solid phase synthesis techniques are used) are removed sequentially or concurrently, to render the final polypeptide. By simple modification of this general procedure, it is possible to add more than one amino acid at a time to a growing chain, for example, by coupling (under conditions which do not racemize chiral centers) a protected tripeptide with a properly protected dipeptide to form, after deprotection, a pentapeptide.

[0120] Typical protecting groups include t-butyloxycarbonyl (Boc), 9-fluorenylmethoxycarbonyl (Fmoc) benzyloxycarbonyl (Cbz); p-toluenesulfonyl (Tx); 2,4-dinitrophenyl; benzyl (Bzl); biphenylisopropylloxycarboxy-carbonyl, t-amylloxycarbonyl, isobornylloxycarbonyl, o-bromobenzyloxycarbonyl, cyclohexyl, isopropyl, acetyl, o-nitrophenylsulfonyl and the like. Typical solid supports are cross-linked polymeric supports. These may include divinylbenzene cross-linked-styrene-based polymers, for example, divinylbenzene-hydroxymethylstyrene copolymers, divinylbenzene-chloromethylstyrene copolymers and divinylbenzene-benzhydrylaminopolystyrene copolymers.

[0121] The polypeptides of the present disclosure may also be chemically prepared by other methods such as by the method of simultaneous multiple peptide synthesis.

[0122] Alternatively, the above-described immunogenic polypeptides, polyproteins, and multiepitope fusion proteins may be produced recombinantly. Once coding sequences for the desired proteins have been isolated or synthesized, they

may be cloned into any suitable vector or replicon for expression. Numerous cloning vectors are known to those of skill in the art, and the selection of an appropriate cloning vector is a matter of choice. A variety of bacterial, yeast, plant, mammalian and insect expression systems are available in the art and any such expression system may be used. Optionally, a polynucleotide encoding these proteins may be translated in a cell-free translation system. Such methods are well known in the art.

[0123] Examples of recombinant DNA vectors for cloning and host cells which they may transform include the bacteriophage λ (*E. coli*), pBR322 (*E. coli*), pACYC177 (*E. coli*), pKT230 (gram-negative bacteria), pGV1106 (gram-negative bacteria), pLAFR1 (gram-negative bacteria), pME290 (non-*E. coli* gram-negative bacteria), pHV14 (*E. coli* and *Bacillus subtilis*), pBD9 (*Bacillus*), pIJ61 (*Streptomyces*), pUC6 (*Streptomyces*), YIp5 (*Saccharomyces*), YCp19 (*Saccharomyces*) and bovine papilloma virus (mammalian cells).

[0124] Insect cell expression systems, such as baculovirus systems, may also be used and are known to those of skill in the art and described in, e.g., Summers and Smith, Texas Agricultural Experiment Station Bulletin No. 1555 (1987). Materials and methods for baculovirus/insect cell expression systems are commercially available in kit form from, inter alia, Invitrogen, San Diego, CA ("MaxBac" kit).

[0125] Plant expression systems may also be used to produce the immunogenic proteins. Generally, such systems use virus-based vectors to transfect plant cells with heterologous genes.

[0126] Viral systems, such as a vaccinia based infection/transfection system, will also find use with the subject matter as disclosed herein. In this system, cells are first transfected in vitro with a vaccinia virus recombinant that encodes the bacteriophage T7 RNA polymerase. This polymerase displays exquisite specificity in that it only transcribes templates bearing T7 promoters. Following infection, cells are transfected with the DNA of interest, driven by a T7 promoter. The polymerase expressed in the cytoplasm from the vaccinia virus recombinant transcribes the transfected DNA into RNA which is then translated into protein by the host translational machinery. The method provides for high level, transient, cytoplasmic production of large quantities of RNA and its translation product(s).

[0127] The gene may be placed under the control of a promoter, ribosome binding site (for bacterial expression) and, optionally, an operator (collectively referred to herein as "control" elements), so that the DNA sequence encoding the desired immunogenic polypeptide is transcribed into RNA in the host cell transformed by a vector containing this expression construction. The coding sequence may or may not contain a signal peptide or leader sequence. With the present subject matter as disclosed herein, both the naturally occurring signal peptides or heterologous sequences may be used. Leader sequences may be removed by the host in post-translational processing. See, e.g., U.S. Pat. Nos. 4,431,739; 4,425,437; 4,338,397, each herein incorporated by reference in their entireties. Such sequences include, but are not limited to, the tpa leader, as well as the honey bee mellitin signal sequence.

[0128] Other regulatory sequences may also be desirable which allow for regulation of expression of the protein sequences relative to the growth of the host cell. Such regulatory sequences are known to those of skill in the art, and examples include those which cause the expression of a

gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Other types of regulatory elements may also be present in the vector, for example, enhancer sequences.

[0129] The control sequences and other regulatory sequences may be ligated to the coding sequence prior to insertion into a vector. Alternatively, the coding sequence may be cloned directly into an expression vector which already contains the control sequences and an appropriate restriction site.

[0130] In embodiments, it may be necessary to modify the coding sequence so that it may be attached to the control sequences with the appropriate orientation; i.e., to maintain the proper reading frame. It may also be desirable to produce mutants or analogs of the immunogenic polypeptides. Mutants or analogs may be prepared by the deletion of a portion of the sequence encoding the protein, by insertion of a sequence, and/or by substitution of one or more nucleotides within the sequence. Techniques for modifying nucleotide sequences, such as site-directed mutagenesis, are well known to those skilled in the art.

[0131] The expression vector is then used to transform an appropriate host cell. A number of mammalian cell lines are known in the art and include immortalized cell lines available from the American Type Culture Collection (ATCC), such as, but not limited to, Chinese hamster ovary (CHO) cells, HeLa cells, baby hamster kidney (BHK) cells, monkey kidney cells (COS), human hepatocellular carcinoma cells (e.g., Hep G2), as well as others. Similarly, bacterial hosts such as *E. coli*, *Bacillus subtilis*, and *Streptococcus* spp., will find use with the present expression constructs. Yeast hosts useful with the subject matter as disclosed include, inter alia, *Saccharomyces cerevisiae*, *Candida albicans*, *Candida maltosa*, *Hansenula polymorpha*, *Kluyveromyces fragilis*, *Kluyveromyces lactis*, *Pichia guilliermondii*, *Pichia pastoris*, *Schizosaccharomyces pombe* and *Yarrowia lipolytica*. Insect cells for use with baculovirus expression vectors include, inter alia, *Aedes aegypti*, *Autographa californica*, *Bombyx mori*, *Drosophila melanogaster*, *Spodoptera frugiperda*, and *Trichoplusia ni*.

[0132] Depending on the expression system and host selected, the proteins as disclosed herein are produced by growing host cells transformed by an expression vector described above under conditions whereby the protein of interest is expressed. The selection of the appropriate growth conditions is within the skill of the art. The cells are then disrupted, using chemical, physical or mechanical means, which lyse the cells yet keep the Rotavirus immunogenic polypeptides substantially intact. Intracellular proteins may also be obtained by removing components from the cell wall or membrane, e.g., by the use of detergents or organic solvents, such that leakage of the immunogenic polypeptides occurs.

[0133] For example, methods of disrupting cells for use with the subject matter as disclosed herein include but are not limited to: sonication or ultrasonication; agitation; liquid or solid extrusion; heat treatment; freeze-thaw; desiccation; explosive decompression; osmotic shock; treatment with lytic enzymes including proteases such as trypsin, neuraminidase and lysozyme; alkali treatment; and the use of detergents and solvents such as bile salts, sodium dodecylsulphate, Triton, NP40 and CHAPS. The particular technique used to disrupt the cells is largely a matter of

choice and will depend on the cell type in which the polypeptide is expressed, culture conditions and any pre-treatment used.

[0134] Following disruption of the cells, cellular debris is removed, generally by centrifugation, and the intracellularly produced Rotavirus immunogenic polypeptides are further purified, using standard purification techniques such as but not limited to, column chromatography, ion-exchange chromatography, size-exclusion chromatography, electrophoresis, HPLC, immunoabsorbent techniques, affinity chromatography, immunoprecipitation, and the like.

[0135] For example, one method for obtaining the intracellular Rotavirus immunogenic polypeptides as disclosed herein involves affinity purification, such as by immunoaffinity chromatography using specific antibodies. The choice of a suitable affinity resin is within the skill in the art. After affinity purification, immunogenic polypeptides may be further purified using conventional techniques well known in the art, such as by any of the techniques described above.

[0136] It may be desirable to produce multiple polypeptides simultaneously (e.g., structural and/or nonstructural proteins from one or more viral strains or viral polypeptides in combination with polypeptide adjuvants). Production of two or more different polypeptides may readily be accomplished by e.g., co-transfecting host cells with constructs encoding the different polypeptides. Co-transfection may be accomplished either in trans or cis, i.e., by using separate vectors or by using a single vector encoding the polypeptides. If a single vector is used, expression of the polypeptides may be driven by a single set of control elements or, alternatively, the sequences coding for the polypeptides may be present on the vector in individual expression cassettes, regulated by individual control elements.

[0137] The polypeptides described herein may be attached to a solid support. The solid supports which may be used in the practice with the subject matter as disclosed herein include substrates such as nitrocellulose (e.g., in membrane or microtiter well form); polyvinylchloride (e.g., sheets or microtiter wells); polystyrene latex (e.g., beads or microtiter plates); polyvinylidene fluoride; diazotized paper; nylon membranes; activated beads, magnetically responsive beads, and the like.

[0138] Typically, a solid support is first reacted with a solid phase component (e.g., one or more Rotavirus antigens) under suitable binding conditions such that the component is sufficiently immobilized to the support. Sometimes, immobilization of the antigen to the support may be enhanced by first coupling the antigen to a protein with better binding properties. Suitable coupling proteins include, but are not limited to, macromolecules such as serum albumins including bovine serum albumin (BSA), keyhole limpet hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, and other proteins well known to those skilled in the art. Other molecules that may be used to bind the antigens to the support include polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and the like. Such molecules and methods of coupling these molecules to the antigens, are well known to those of ordinary skill in the art.

[0139] If desired, polypeptides may be labeled using conventional techniques. Suitable labels include fluorophores, chromophores, radioactive atoms (particularly ^{32}P and ^{125}I), electron-dense reagents, enzymes, and ligands having specific binding partners. Enzymes are typically detected by

their activity. For example, horseradish peroxidase is usually detected by its ability to convert 3,3',5,5'-tetramethylbenzidine (TMB) to a blue pigment, quantifiable with a spectrophotometer. "Specific binding partner" refers to a protein capable of binding a ligand molecule with high specificity, as for example in the case of an antigen and a monoclonal antibody specific therefor. Other specific binding partners include biotin and avidin or streptavidin, IgG and protein A, and the numerous receptor-ligand couples known in the art. A single label or a combination of labels may be used in the as disclosed herein.

[0140] Once formulated, the compositions as disclosed herein may be administered directly to the subject (e.g., as described above) or, alternatively, delivered ex vivo, to cells derived from the subject, using methods such as those described above.

Immunogenic Compositions

[0141] The present disclosure also provides compositions comprising one or more of the immunogenic polypeptides and/or polyproteins multi-epitope fusion proteins described herein. Different polypeptides, polyproteins, and multiple epitope fusion proteins may be mixed together in a single formulation. Within such combinations, an antigen of the immunogenic composition may be present in more than one polypeptide, or multiple epitope polypeptide, or polyprotein.

[0142] The immunogenic compositions may comprise a mixture of polypeptides, which in turn may be delivered using the same or different vehicles. Antigens may be administered individually or in combination, in e.g., prophylactic (i.e., to prevent infection) or therapeutic (to treat infection) immunogenic compositions. The immunogenic composition may be given more than once (e.g., a "prime" administration followed by one or more "boosts") to achieve the desired effects. The same composition may be administered in one or more priming and one or more boosting steps. Alternatively, different compositions may be used for priming and boosting.

[0143] The immunogenic compositions will generally include one or more "pharmaceutically acceptable excipients or vehicles" such as water, saline, glycerol, ethanol, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.

[0144] Immunogenic compositions will typically, in addition to the components mentioned above, comprise one or more "pharmaceutically acceptable carriers." These include any carrier which does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers typically are large, slowly metabolized macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and lipid aggregates (such as oil droplets or liposomes). Such carriers are well known to those of ordinary skill in the art. A composition may also contain a diluent, such as water, saline, glycerol, and the like. Additionally, an auxiliary substance, such as a wetting or emulsifying agent, pH buffering substance, and the like, may be present. A thorough discussion of pharmaceutically acceptable components is available in Gennaro (2000) Remington: The Science and Practice of Pharmacy. 20th ed., ISBN: 0683306472.

[0145] Pharmaceutically acceptable salts may also be used in compositions as disclosed herein, for example, mineral

salts such as hydrochlorides, hydrobromides, phosphates, or sulfates, as well as salts of organic acids such as acetates, propionates, malonates, or benzoates. Especially useful protein substrates are serum albumins, keyhole limpet hemocyanin, immunoglobulin molecules, thyroglobulin, ovalbumin, *tetanus* toxoid, and other proteins well known to those of skill in the art. Compositions as disclosed may also contain liquids or excipients, such as water, saline, glycerol, dextrose, ethanol, or the like, singly or in combination, as well as substances such as wetting agents, emulsifying agents, or pH buffering agents. Antigens may also be adsorbed to, entrapped within or otherwise associated with liposomes and particulate carriers such as PLG.

[0146] Antigens may be conjugated to a carrier protein in order to enhance immunogenicity. This is particularly useful in compositions in which a saccharide or carbohydrate antigen is used.

[0147] Carrier proteins may include, but are not limited to, bacterial toxins or toxoids, such as diphtheria or *tetanus* toxoids. The CRM₁₉₇ diphtheria toxoid may be used. Other carrier polypeptides include the *N. meningitidis* outer membrane protein (EP-A-0372501), synthetic peptides (EP-A-0378881 and EP-A-0427347), heat shock proteins (WO 93/17712 and WO 94/03208), *pertussis* proteins (WO 98/58668 and EP-A-0471177), protein D from *H. influenzae* (WO 00/56360), cytokines (WO 91/01146), lymphokines, hormones, growth factors, toxin A or B from *C. difficile* (WO 00/61761), iron-uptake proteins, such as transferring (WO 01/72337), etc. Where a mixture comprises capsular saccharide from both serigraphs A and C, it may be that the ratio (w/w) of MenA saccharide:MenC saccharide is greater than 1 (e.g., 2:1, 3:1, 4:1, 5:1, 10:1 or higher). Different saccharides may be conjugated to the same or different type of carrier protein. Any suitable conjugation reaction may be used, with any suitable linker where necessary.

[0148] Immunogenic compositions, including vaccines as disclosed may be administered in conjunction with other immunoregulatory agents. For example, a vaccine as disclosed herein may include an adjuvant. Adjuvants include, but are not limited to, one or more of the following types of adjuvants described below.

Mineral Containing Compositions

[0149] Mineral containing compositions suitable for use as adjuvants disclosed herein include mineral salts, such as aluminum salts and calcium salts. Salts as disclosed herein includes mineral salts such as hydroxides (e.g., oxyhydroxides), phosphates (e.g., hydroxyphosphates, orthophosphates), sulfates, and the like, or mixtures of different mineral compounds (e.g., a mixture of a phosphate and a hydroxide adjuvant, optionally with an excess of the phosphate), with the compounds taking any suitable form (e.g., gel, crystalline, amorphous, and the like). The mineral containing compositions may also be formulated as a particle of metal salt (WO00/23105).

[0150] Aluminum salts may be included in vaccines such that the dose of Al³⁺ is between 0.2 and 1.0 mg per dose.

[0151] In embodiments, the aluminum based adjuvant for use as disclosed is alum (aluminum potassium sulfate (AlK(SO₄)₂)), or an alum derivative, such as that formed in-situ by mixing an antigen in phosphate buffer with alum, followed by titration and precipitation with a base such as ammonium hydroxide or sodium hydroxide.

[0152] Another aluminum-based adjuvant for use in vaccine formulations of the present invention is aluminum hydroxide adjuvant ($\text{Al}(\text{OH})_3$) or crystalline aluminum oxyhydroxide (AlOOH), which is an excellent adsorbant, having a surface area of approximately $500 \text{ m}^2/\text{g}$. Alternatively, aluminum phosphate adjuvant (AlPO_4) or aluminum hydroxyphosphate, which contains phosphate groups in place of some or all of the hydroxyl groups of aluminum hydroxide adjuvant is provided. In embodiments, aluminum phosphate adjuvants provided herein are amorphous and soluble in acidic, basic and neutral media.

[0153] In embodiments, the adjuvant as disclosed herein comprises both aluminum phosphate and aluminum hydroxide. In one aspect, the adjuvant has a greater amount of aluminum phosphate than aluminum hydroxide, such as a ratio of 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1 or greater than 9:1, by weight aluminum phosphate to aluminum hydroxide. More particular still, aluminum salts in the vaccine are present at 0.4 to 1.0 mg per vaccine dose, or 0.4 to 0.8 mg per vaccine dose, or 0.5 to 0.7 mg per vaccine dose, or about 0.6 mg per vaccine dose.

[0154] Generally, the aluminum-based adjuvant(s), or ratio of multiple aluminum-based adjuvants, such as aluminum phosphate to aluminum hydroxide is selected by optimization of electrostatic attraction between molecules such that the antigen carries an opposite charge as the adjuvant at the desired pH. For example, aluminum phosphate adjuvant ($\text{iep}=4$) adsorbs lysozyme, but not albumin at pH 7.4. Should albumin be the target, aluminum hydroxide adjuvant would be selected (i.e., 11.4). Alternatively, pretreatment of aluminum hydroxide with phosphate lowers its isoelectric point, making it a preferred adjuvant for more basic antigens.

Oil-Emulsions

[0155] Oil-emulsion compositions suitable for use as adjuvants may include squalene-water emulsions, such as MF59 (5% Squalene, 0.5% TWEEN 80™, and 0.5% SPAN 85™, formulated into submicron particles using a microfluidizer). See WO90/14837. MF59 is used as the adjuvant in the FLUAD™ influenza virus trivalent subunit vaccine.

[0156] Particularly adjuvants for use in the compositions are submicron oil-in-water emulsions. Submicron oil-in-water emulsions for use herein may be squalene/water emulsions optionally containing varying amounts of MTP-PE, such as a submicron oil-in-water emulsion containing 4-5% w/v squalene, 0.25-1.0% w/v TWEEN80™ (polyoxyethylenesorbitan monooleate), and/or 0.25-1.0% SPAN85™ (sorbitan trioleate), and, optionally, N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(β , β -dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), for example, the submicron oil-in-water emulsion known as “MF59” (International Publication No. WO90/14837; U.S. Pat. Nos. 6,299,884 and 6,451,325.) MF59 contains 4-5% w/v Squalene (e.g., 4.3%), 0.25-0.5% w/v TWEEN80™, and 0.5% w/v SPAN85™ and optionally contains various amounts of MTP-PE, formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton, Mass.). For example, MTP-PE may be present in an amount of about 0-500 g/dose, 0-250 g/dose and 0-100 g/dose. As used herein, the term “MF59-0” refers to the above submicron oil-in-water emulsion lacking MTP-PE, while the term MF59-MTP denotes a formulation that contains MTP-PE. For instance, “MF59-100” contains 100 μg MTP-PE per

dose, and so on. MF69, another submicron oil-in-water emulsion for use herein, contains 4.3% w/v squalene, 0.25% w/v TWEEN80™, and 0.75% w/v SPAN85™ and optionally MTP-PE. Yet another submicron oil-in-water emulsion is MF75, also known as SAF, containing 10% squalene, 0.4% TWEEN80™, 5% pluronic-blocked polymer L121, and thr-MDP, also microfluidized into a submicron emulsion. MF75-MTP denotes an MF75 formulation that includes MTP, such as from 100-400 g MTP-PE per dose. [0157] Submicron oil-in-water emulsions, methods of making the same and immunostimulating agents, such as muramyl peptides, for use in the compositions, are described in detail in International Publication No. WO90/14837 and U.S. Pat. Nos. 6,299,884 and 6,451,325.

[0158] Complete Freund's adjuvant (CFA) and incomplete Freund's adjuvant (IFA) may also be used as adjuvants.

Saponin Formulations

[0159] Saponin formulations, may also be used as adjuvants. Saponins are a heterologous group of sterol glycosides and triterpenoid glycosides that are found in the bark, leaves, stems, roots and even flowers of a wide range of plant species. Saponins isolated from the bark of the *Quilalaia saponaria* Molina tree have been widely studied as adjuvants. Saponins may also be commercially obtained from *Smilax ornata* (sarsaparilla), *Gypsophilla paniculata* (brides veil), and *Saponaria officinalis* (soap root). Saponin adjuvant formulations include purified formulations, such as QS21, as well as lipid formulations, such as ISCOMs.

[0160] Saponin compositions have been purified using High Performance Thin Layer Chromatography (HP-TLC) and Reversed Phase High Performance Liquid Chromatography (RP-HPLC). Specific purified fractions using these techniques have been identified, including QS7, QS17, QS18, QS21, QH-A, QH-B and QH-C. In embodiments, the saponin is QS21. A method of production of QS21 is disclosed in U.S. Pat. No. 5,057,540. Saponin formulations may also comprise a sterol, such as cholesterol (see WO96/33739).

[0161] Combinations of saponins and cholesterol may be used to form unique particles called Immunostimulating Complexes (ISCOMs). ISCOMs typically also include a phospholipid such as phosphatidylethanolamine or phosphatidylcholine. Any known saponin may be used in ISCOMs. In embodiments, the ISCOM includes one or more of Quil A, QHA and QHC. ISCOMs are further described in EP0109942, WO96/11711 and WO96/33739. Optionally, the ISCOMS may be devoid of (an) additional detergent(s). See WO00/07621.

Bacterial or Microbial Derivatives

[0162] Adjuvants suitable for use as disclosed herein include bacterial or microbial derivatives such as:

(1) Non-Toxic Derivatives of Enterobacterial Lipopolysaccharide (LPs)

[0163] Such derivatives include Monophosphoryl lipid A (MPL) and 3-O-deacylated MPL (3dMPL). 3dMPL is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains. One “small particle” form of 3 De-O-acylated monophosphoryl lipid A is disclosed in EP 0 689 454. Such “small particles” of 3dMPL are small enough to be sterile filtered through a 0.22 micron membrane (see

EP 0 689 454). Other non-toxic LPS derivatives include monophosphoryl lipid A mimics, such as aminoalkyl glucosaminide phosphate derivatives e.g., RC-529.

(2) Lipid A Derivatives

[0164] Lipid A derivatives include derivatives of lipid A from *Escherichia coli* such as OM-174.

(3) Immunostimulatory Oligonucleotides

[0165] Immunostimulatory oligonucleotides suitable for use as adjuvants may include nucleotide sequences containing a CpG motif (a sequence containing an unmethylated cytosine followed by guanosine and linked by a phosphate bond). Bacterial double stranded RNA or oligonucleotides containing palindromic or poly(dG) sequences have also been shown to be immunostimulatory.

[0166] The CpG's may include nucleotide modifications/analogs such as phosphorothioate modifications and may be double-stranded or single-stranded. Optionally, the guanosine may be replaced with an analog such as 2'-deoxy-7-deazaguanosine.

[0167] The CpG sequence may be directed to TLR9, such as the motif GTCGTT or TTCGTT. The CpG sequence may be specific for inducing a Th1 immune response, such as a CpG-A ODN, or it may be more specific for inducing a B cell response, such a CpG-B ODN. In embodiments, the CpG is a CpG-A ODN.

[0168] In embodiments, the CpG oligonucleotide may be constructed so that the 5' end is accessible for receptor recognition. Optionally, two CpG oligonucleotide sequences may be attached at their 3' ends to form "immunomers."

(4) ADP-Ribosylating Toxins and Detoxified Derivatives Thereof.

[0169] Bacterial ADP-ribosylating toxins and detoxified derivatives thereof may be used as adjuvants. In embodiments, the protein may be derived from *E. coli* (i.e., *E. coli* heat labile enterotoxin "LT", cholera ("CT"), or *pertussis* ("PT"). The use of detoxified ADP-ribosylating toxins as mucosal adjuvants is described in WO95/17211 and as parenteral adjuvants in WO98/42375. In embodiments, the adjuvant is a detoxified LT mutant such as LT-K63, LT-R72, and LTR192G.

Bioadhesives and Mucoadhesives

[0170] Bioadhesives and mucoadhesives may also be used as adjuvants. Suitable bioadhesives include esterified hyaluronic acid microspheres or mucoadhesives such as cross-linked derivatives of polyacrylic acid, polyvinyl alcohol, polyvinyl pyrrolidone, polysaccharides and carboxymethylcellulose. Chitosan and derivatives thereof may also be used as adjuvants. E.g., WO99/27960.

Muramyl Peptides

[0171] Examples of muramyl peptides suitable for use as adjuvants include N-acetyl-muramyl-L-threonine-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-1-alanyl-d-isoglutamine (nor-MDP), and N-acetylmuramyl-1-alanyl-d-isoglutaminyl-1-alanine-2-(1'-2'-dipalmitoyl-s-n-glycero-3-hydroxyphosphoryloxy)-ethylamine MTP-PE).

Imidazoquinoline Compounds.

[0172] Examples of imidazoquinoline compounds suitable for use as adjuvants include Imiquimod and its analogues (see, e.g., U.S. Pat. Nos. 4,689,338, 5,389,640, 5,268,376, 4,929,624, 5,266,575, 5,352,784, 5,494,916, 5,482,936, 5,346,905, 5,395,937, 5,238,944, and 5,525,612).

Thiosemicarbazone Compounds.

[0173] Examples of thiosemicarbazone compounds, as well as methods of formulating, manufacturing, and screening for compounds all suitable for use as adjuvants include those described in WO04/60308. The thiosemicarbazones are particularly effective in the stimulation of human peripheral blood mononuclear cells for the production of cytokines, such as TNF- α .

Tryptanthrin Compounds.

[0174] Examples of tryptanthrin compounds, as well as methods of formulating, manufacturing, and screening for compounds all suitable for use as adjuvants as disclosed herein include those described in WO04/64759. The tryptanthrin compounds are particularly effective in the stimulation of human peripheral blood mononuclear cells for the production of cytokines, such as TNF- α .

[0175] Combinations of aspects of one or more of the adjuvants identified above may be applied to the compositions as disclosed herein. For example, the following adjuvant compositions may be used:

[0176] (1) a saponin and an oil-in-water emulsion (WO99/11241); (2) a saponin (e.g., QS21)+a non-toxic LPS derivative (e.g., 3dMPL) (see WO94/00153); (3) a saponin (e.g., QS21)+a non-toxic LPS derivative (e.g., 3dMPL)+a cholesterol; (4) a saponin (e.g., QS21)+3dMPL+IL-12 (optionally+a sterol) (WO98/57659); (5) combinations of 3dMPL with, for example, QS21 and/or oil-in-water emulsions (See European patent applications 0835318, 0735898 and 0761231); (6) SAF, containing 10% Squalene, 0.4% TWEEN80™, 5% pluronic-block polymer L121, and thr-MDP, either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion. (7) RIBI™ adjuvant system (RAS), (Ribi Immunochem) containing 2% Squalene, 0.2% TWEEN80™, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), MPL+CWS (DETOX™); and (8) one or more mineral salts (such as an aluminum salt)+a non-toxic derivative of LPS (such as 3dPML). (9) one or more mineral salts (such as an aluminum salt) and one or more immunostimulatory oligonucleotides (such as a nucleotide sequence including a CpG motif) and one or more detoxified ADP-ribosylating toxins (such as LT-K63 and LT-R72), (10) inulin and inulin acetate formulations (see, e.g., WO 2013/110050, herein incorporated in its entirety).

Additional Antigens

[0177] Compositions of the as disclosed herein optionally may comprise one or more additional polypeptide antigens which are not derived from Rotavirus proteins. Such antigens include bacterial, viral, or parasitic antigens.

[0178] In some embodiments, Rotavirus antigens are combined with one or more antigens including, but not limited to, antigens derived from a bacteria or virus, such as Orthomyxovirus (influenza), Pneumovirus (RSV), Paramyxovirus (PIV and Mumps), Morbillivirus (measles), Togavirus (Rubella), Enterovirus (polio), HBV, Coronavirus (SARS), and Varicella-zoster virus (VZV), Epstein Barr virus (EBV), *Streptococcus pneumoniae*, *Neisseria meningitides*, *Streptococcus pyogenes* (Group A *Streptococcus*), *Moraxella catarrhalis*, *Bordetella pertussis*, *Staphylococcus aureus*, *Clostridium tetani* (*Tetanus*), *Corynebacterium diphtheriae* (Diphtheria), *Haemophilus influenzae* B (Hib), *Pseudomonas aeruginosa*, *Streptococcus agalactiae* (Group B *Streptococcus*), and *E. coli*.

[0179] In other embodiments, Rotavirus antigens are combined with one or more antigens including, but not limited to, *Neisseria meningitides*, *Streptococcus pneumoniae*, *Streptococcus pyogenes* (Group A *Streptococcus*), *Moraxella catarrhalis*, *Bordetella pertussis*, *Staphylococcus aureus*, *Staphylococcus epidermis*, *Clostridium tetani* (*Tetanus*), *Corynebacterium diphtheriae* (Diphtheria), *Haemophilus influenzae* B (Hib), *Pseudomonas aeruginosa*, *Legionella pneumophila*, *Streptococcus agalactiae* (Group B *Streptococcus*), *Enterococcus faecalis*, *Helicobacter pylori*, *Chlamydia pneumoniae*, Orthomyxovirus (influenza), Pneumovirus (RSV), Paramyxovirus (PIV and Mumps), Morbillivirus (measles), Togavirus (Rubella), Enterovirus (polio), HBV, Coronavirus (SARS), Varicella-zoster virus (VZV), Epstein Barr virus (EBV), Cytomegalovirus (CMV).

[0180] In other embodiments, Rotavirus antigens are combined with one or more antigens which are useful in a vaccine designed to protect individuals against pathogens that cause diarrheal diseases. Such antigens include, but are not limited to, rotavirus, *Shigella* spp., enterotoxigenic *Escherichia coli* (ETEC), *Vibrio cholerae*, and *Campylobacter jejuni* antigens. In embodiments, one or more Norovirus antigens may be derived from Norwalk virus, Snow Mountain virus, and/or Hawaii virus are combined with a rotavirus antigen in an immunogenic composition.

[0181] Antigens which may find use with the present compositions include, but are not limited to, one or more of the following antigens set forth below, or antigens derived from one or more of the pathogens set forth below:

Bacterial Antigens

[0182] Suitable Bacterial antigens as disclosed herein include proteins, polysaccharides, lipopolysaccharides, and outer membrane vesicles which may be isolated, purified or derived from a bacteria. In addition, bacterial antigens may include bacterial lysates and inactivated bacteria formulations. Bacteria antigens may be produced by recombinant expression. Bacterial antigens include epitopes which may be exposed on the surface of the bacteria during at least one stage of its life cycle. Bacterial antigens may be conserved across multiple serotypes. Bacterial antigens include antigens derived from one or more of the bacteria set forth below as well as the specific antigens examples identified below.

[0183] *Neisseria meningitides*: *Meningitides* antigens may include proteins (such as those identified in References 1-7), saccharides (including a polysaccharide, oligosaccharide or lipopolysaccharide), or outer-membrane vesicles purified or derived from *N. meningitides* serogroup such as A, C, W135, Y, and/or B. *Meningitides* protein antigens may be selected

from adhesions, autotransporters, toxins, Fe acquisition proteins, and membrane associated proteins (e.g., integral outer membrane protein).

[0184] *Streptococcus pneumoniae*: *Streptococcus pneumoniae* antigens may include a saccharide (including a polysaccharide or an oligosaccharide) and/or protein from *Streptococcus pneumoniae*. Saccharide antigens may be selected from serotypes 1, 2, 3, 4, 5, 6B, 7F, 8, 9N, 9V, 10A, 11A, 12F, 14, 15B, 17F, 18C, 19A, 19F, 20, 22F, 23F, and 33F. Protein antigens may be selected from a protein identified in WO 98/18931, WO 98/18930, U.S. Pat. Nos. 6,699,703, 6,800,744, WO 97/43303, and WO 97/37026. *Streptococcus pneumoniae* proteins may be selected from the Poly Histidine Triad family (PhtX), the Choline Binding Protein family (CbpX), CbpX truncates, LytX family, LytX truncates, CbpX truncate-LytX truncate chimeric proteins, pneumolysin (Ply), PspA, PsaA, Sp128, Sp101, Sp130, Sp125 or Sp133.

[0185] *Streptococcus pyogenes* (Group A *Streptococcus*): Group A *Streptococcus* antigens may include a protein identified in WO 02/34771 or WO 2005/032582 (including GAS 40), fusions of fragments of GAS M proteins (including those described in WO 02/094851), fibronectin binding protein (Sfb1), Streptococcal heme-associated protein (Shp), and Streptolysin S (SagA).

[0186] *Moraxella catarrhalis*: *Moraxella* antigens include antigens identified in WO 02/18595 and WO 99/58562, outer membrane protein antigens (HMW-OMP), C-antigen, and/or LPS.

[0187] *Bordetella pertussis*: *Pertussis* antigens include *pertussis* holotoxin (PT) and filamentous haemagglutinin (FHA) from *B. pertussis*, optionally also combination with pertactin and/or agglutinogens 2 and 3 antigen.

[0188] *Staphylococcus aureus*: *Staph aureus* antigens include *S. aureus* type 5 and 8 capsular polysaccharides optionally conjugated to nontoxic recombinant *Pseudomonas aeruginosa* exotoxin A, such as STAPHVAX™, or antigens derived from surface proteins, invasins (leukocidin, kinases, hyaluronidase), surface factors that inhibit phagocytic engulfment (capsule, Protein A), carotenoids, catalase production, Protein A, coagulase, clotting factor, and/or membrane-damaging toxins (optionally detoxified) that lyse eukaryotic cell membranes (hemolysins, leukotoxin, leukocidin).

[0189] *Staphylococcus epidermis*: *S. epidermidis* antigens include slime-associated antigen (SAA).

[0190] *Clostridium tetani* (*Tetanus*): *Tetanus* antigens include *tetanus* toxoid (TT), may be used as a carrier protein in conjunction/conjugated with the compositions of the present disclosure.

[0191] *Corynebacterium diphtheriae* (Diphtheria): Diphtheria antigens include diphtheria toxin, including detoxified, such as CRM₁₉₇. Additionally, antigens capable of modulating, inhibiting or associated with ADP ribosylation are contemplated for combination/co-administration/conjugation with the compositions of the present disclosure. The diphtheria toxoids may be used as carrier proteins.

[0192] *Haemophilus influenzae* B (Hib): Hib antigens include a Hib saccharide antigen.

[0193] *Pseudomonas aeruginosa*: *Pseudomonas* antigens include endotoxin A, Wzz protein, *P. aeruginosa* LPS, more particularly LPS isolated from PAO1 (05 serotype), and/or Outer Membrane Proteins, including Outer Membrane Proteins F (OprF).

[0194] *Legionella pneumophila*. Bacterial antigens may be derived from *Legionella pneumophila*.

[0195] *Streptococcus agalactiae* (Group B *Streptococcus*): Group B *Streptococcus* antigens include a protein or saccharide antigen identified in WO 02/34771, WO 03/093306, WO 04/041157, or WO 2005/002619 (including proteins GBS 80, GBS 104, GBS 276 and GBS 322, and including saccharide antigens derived from serotypes Ia, Ib, Ia/c, II, III, IV, V, VI, VII and VIII).

[0196] *Neisseria gonorrhoeae*: *Gonorrhoeae* antigens include Por (or porin) protein, such as PorB, a transferring binding protein, such as TbpA and TbpB, a opacity protein (such as Opa), a reduction-modifiable protein (Rmp), and outer membrane vesicle (OMV) preparations (see e.g., WO99/24578, WO99/36544, WO99/57280, WO02/079243).

[0197] *Chlamydia trachomatis*: *Chlamydia trachomatis* antigens include antigens derived from serotypes A, B, Ba and C (agents of trachoma, a cause of blindness), serotypes L₁, L₂ & L₃ (associated with Lymphogranuloma venereum), and serotypes, D-K. *Chlamydia trachomatis* antigens may also include an antigen identified in WO 00/37494, WO 03/049762, WO 03/068811, or WO 05/002619, including PepA (CT045), LcrE (CT089), ArtJ (CT381), DnaK (CT396), CT398, OmpH-like (CT242), L7/L12 (CT316), OmcA (CT444), AtosS (CT467), CT547, Eno (CT587), HrtA (CT823), and MurG (CT761).

[0198] *Treponema pallidum* (Syphilis): Syphilis antigens include TmpA antigen.

[0199] *Haemophilus ducreyi* (causing chancroid): *Ducreyi* antigens include outer membrane protein (DsrA).

[0200] *Enterococcus faecalis* or *Enterococcus faecium*: Antigens include a trisaccharide repeat or other *Enterococcus* derived antigens provided in U.S. Pat. No. 6,756,361.

[0201] *Helicobacter pylori*: *H. pylori* antigens include Cag, Vac, Nap, HopX, HopY and/or urease antigen.

[0202] *Staphylococcus saprophyticus*: Antigens include the 160 kDa hemagglutinin of *S. saprophyticus* antigen.

[0203] *Yersinia enterocolitica* Antigens include LPS.

[0204] *E. coli*: *E. coli* antigens may be derived from enterotoxigenic *E. coli* (ETEC), enteroaggregative *E. coli* (EAaggEC), diffusely adhering *E. coli* (DAEC), enteropathogenic *E. coli* (EPEC), and/or enterohemorrhagic *E. coli* (EHEC).

[0205] *Bacillus anthracis* (anthrax): *B. anthracis* antigens are optionally detoxified and may be selected from A-components (lethal factor (LF) and edema factor (EF)), both of which may share a common B-component known as protective antigen (PA).

[0206] *Yersinia pestis* (plague): Plague antigens include F1 capsular antigen.

[0207] *Mycobacterium tuberculosis*: Tuberculosis antigens include lipoproteins, LPS, BCG antigens, a fusion protein of antigen 85B (Ag85B) and/or ESAT-6 optionally formulated in cationic lipid vesicles, *Mycobacterium tuberculosis* (Mtb) isocitrate dehydrogenase associated antigens, and/or MP751 antigens.

[0208] *Rickettsia*: Antigens include outer membrane proteins, including the outer membrane protein A and/or B (OmpB).

[0209] *Listeria monocytogenes*. Bacterial antigens may be derived from *Listeria monocytogenes*.

[0210] *Chlamydia pneumoniae*: Antigens include those identified in WO 02/02606.

[0211] *Vibrio cholerae*: Antigens include proteinase antigens, LPS, particularly lipopolysaccharides of *Vibrio cholerae* II, 01 Inaba O-specific polysaccharides, *V. cholera* 0139, antigens of IEM108 vaccine, and/or Zonula occludens toxin (Zot).

[0212] *Salmonella typhi* (typhoid fever): Antigens include capsular polysaccharides, including conjugates (Vi, i.e., vax-TyVi).

[0213] *Borrelia burgdorferi* (Lyme disease): Antigens include lipoproteins (such as OspA, OspB, Osp C and Osp D), other surface proteins such as OspE-related proteins (Erps), decorin-binding proteins (such as DbpA), and antigenically variable VI proteins, such as antigens associated with P39 and P13 VlsE Antigenic Variation Protein.

[0214] *Porphyromonas gingivalis*: Antigens include *P. gingivalis* outer membrane protein (OMP).

[0215] *Klebsiella*: Antigens include an OMP, including OMP A, or a polysaccharide optionally conjugated to *tetanus* toxoid.

[0216] Further bacterial antigens of the instant disclosure may be capsular antigens, polysaccharide antigens or protein antigens of any of the above. Further bacterial antigens may also include an outer membrane vesicle (OMV) preparation. Additionally, antigens include live, attenuated, and/or purified versions of any of the aforementioned bacteria. The antigens of the present disclosure may be derived from gram-negative or gram-positive bacteria. The antigens of the present disclosure may be derived from aerobic or anaerobic bacteria.

[0217] Additionally, any of the above bacterial-derived saccharides (polysaccharides, LPS, LOS or oligosaccharides) may be conjugated to another agent or antigen, such as a carrier protein (for example CRM₁₉₇). Such conjugation may be direct conjugation effected by reductive amination of carbonyl moieties on the saccharide to amino groups on the protein, as provided in U.S. Pat. No. 5,360,897. Alternatively, the saccharides may be conjugated through a linker, such as, with succinamide or other linkages.

Viral Antigens

[0218] Viral antigens suitable for use in the compositions as disclosed include purified subunit formulations, viral proteins which may be isolated, purified or derived from a virus, and Virus Like Particles (VLPs). Viral antigens may be derived from viruses propagated on cell culture or other substrate. Alternatively, viral antigens may be expressed recombinantly. Viral antigens include epitopes which are exposed on the surface of the virus during at least one stage of its life cycle. Viral antigens may be conserved across multiple serotypes or isolates. Viral antigens include antigens derived from one or more of the viruses set forth below as well as the specific antigens examples identified below.

[0219] Orthomyxovirus: Viral antigens may be derived from an Orthomyxovirus, such as Influenza A, B and C. Orthomyxovirus antigens may be selected from one or more of the viral proteins, including hemagglutinin (HA), neuraminidase (NA), nucleoprotein (NP), matrix protein (M1), membrane protein (M2), one or more of the transcriptase components (PB1, PB2 and PA). In embodiments, antigens include HA and NA.

[0220] Influenza antigens may be derived from interpanemic (annual) flu strains. Alternatively, influenza antigens may be derived from strains with the potential to cause pandemic a pandemic outbreak (i.e., influenza strains with

new haemagglutinin compared to the haemagglutinin in currently circulating strains, or influenza strains which are pathogenic in avian subjects and have the potential to be transmitted horizontally in the human population, or influenza strains which are pathogenic to humans).

[0221] Paramyxoviridae viruses: Viral antigens may be derived from Paramyxoviridae viruses, such as Pneumoviruses (RSV), Paramyxoviruses (PIV) and Morbilliviruses (Measles).

[0222] Pneumovirus: Viral antigens may be derived from a Pneumovirus, such as Respiratory syncytial virus (RSV), Bovine respiratory syncytial virus, Pneumonia virus of mice, and Turkey rhinotracheitis virus. In embodiments, the Pneumovirus is RSV. Pneumovirus antigens may be selected from one or more of the following proteins, including surface proteins Fusion (F), Glycoprotein (G) and Small Hydrophobic protein (SH), matrix proteins M and M2, nucleocapsid proteins N, P and L and nonstructural proteins NS1 and NS2. Pneumovirus antigens may include F, G and M. Pneumovirus antigens may also be formulated in or derived from chimeric viruses. For example, chimeric RSV/PIV viruses may comprise components of both RSV and PIV.

[0223] Paramyxovirus: Viral antigens may be derived from a Paramyxovirus, such as Parainfluenza virus types 1-4 (PIV), Mumps, Sendai viruses, Simian virus 5, Bovine parainfluenza virus and Newcastle disease virus. In embodiments, the Paramyxovirus is PIV or Mumps. Paramyxovirus antigens may be selected from one or more of the following proteins: Hemagglutinin-Neuraminidase (HN), Fusion proteins F1 and F2, Nucleoprotein (NP), Phosphoprotein (P), Large protein (L), and Matrix protein (M). Paramyxovirus proteins may include HN, F1 and F2. Paramyxovirus antigens may also be formulated in or derived from chimeric viruses. For example, chimeric RSV/PIV viruses may comprise components of both RSV and PIV. Commercially available mumps vaccines include live attenuated mumps virus, in either a monovalent form or in combination with measles and rubella vaccines (MMR).

[0224] Morbillivirus: Viral antigens may be derived from a Morbillivirus, such as Measles. Morbillivirus antigens may be selected from one or more of the following proteins: hemagglutinin (H), Glycoprotein (G), Fusion factor (F), Large protein (L), Nucleoprotein (NP), Polymerase phosphoprotein (P), and Matrix (M). Commercially available measles vaccines include live attenuated measles virus, typically in combination with mumps and rubella (MMR).

[0225] Picornavirus: Viral antigens may be derived from Picornaviruses, such as Enteroviruses, Rhinoviruses, Hepatitis virus, Cardioviruses and Aphthoviruses. Antigens derived from Enteroviruses, such as Poliovirus are may be used.

[0226] Enterovirus: Viral antigens may be derived from an Enterovirus, such as Poliovirus types 1, 2 or 3, Coxsackie A virus types 1 to 22 and 24, Coxsackie B virus types 1 to 6, Echovirus (ECHO) virus types 1 to 9, 11 to 27 and 29 to 34 and Enterovirus 68 to 71. In embodiments, the Enterovirus may be poliovirus. Enterovirus antigens may include one or more of the following Capsid proteins VP1, VP2, VP3 and VP4. Commercially available polio vaccines include Inactivated Polio Vaccine (IPV) and Oral poliovirus vaccine (OPV).

[0227] Heparnavirus: Viral antigens may be derived from an Heparnavirus, such as Hepatitis A virus (HAV). Commercially available HAV vaccines include inactivated HAV vaccine.

[0228] Togavirus: Viral antigens may be derived from a Togavirus, such as a Rubivirus, an Alphavirus, or an Arterivirus. Antigens derived from Rubivirus, such as Rubella virus, may be used. Togavirus antigens may be selected from E1, E2, E3, C, NSP-1, NSPO-2, NSP-3 or NSP-4. Togavirus antigens include E1, E2 or E3. Commercially available Rubella vaccines include a live cold-adapted virus, typically in combination with mumps and measles vaccines (MMR).

[0229] Flavivirus: Viral antigens may be derived from a Flavivirus, such as Tick-borne encephalitis (TBE), Dengue (types 1, 2, 3 or 4), Yellow Fever, Japanese encephalitis, West Nile encephalitis, St. Louis encephalitis, Russian spring-summer encephalitis, Powassan encephalitis. Flavivirus antigens may be selected from PrM, M, C, E, NS-1, NS-2a, NS2b, NS3, NS4a, NS4b, and NS5. Flavivirus antigens may include PrM, M and E. Commercially available TBE vaccine include inactivated virus vaccines.

[0230] Pestivirus: Viral antigens may be derived from a Pestivirus, such as Bovine viral diarrhea (BVDV), Classical swine fever (CSFV) or Border disease (BDV).

[0231] Hepadnavirus: Viral antigens may be derived from a Hepadnavirus, such as Hepatitis B virus. Hepadnavirus antigens may be selected from surface antigens (L, M and S), core antigens (HBc, HBe). Commercially available HBV vaccines include subunit vaccines comprising the surface antigen S protein.

[0232] Hepatitis C virus: Viral antigens may be derived from a Hepatitis C virus (HCV). HCV antigens may be selected from one or more of E1, E2, E1/E2, NS3/45 polyprotein, NS 3/45-core polyprotein, core, and/or peptides from the nonstructural regions.

[0233] Rhabdovirus: Viral antigens may be derived from a Rhabdovirus, such as a Lyssavirus (Rabies virus) and Vesiculovirus (VSV). Rhabdovirus antigens may be selected from glycoprotein (G), nucleoprotein (N), large protein (L), nonstructural proteins (NS). Commercially available Rabies virus vaccine comprise killed virus grown on human diploid cells or fetal rhesus lung cells.

[0234] Caliciviridae; Viral antigens may be derived from Caliciviridae, such as Norwalk virus, and Norwalk-like Viruses, such as Hawaii Virus and Snow Mountain Virus.

[0235] Coronavirus: Viral antigens may be derived from a Coronavirus, SARS, Human respiratory coronavirus, Avian infectious bronchitis (IBV), Mouse hepatitis virus (MHV), and Porcine transmissible gastroenteritis virus (TGEV). Coronavirus antigens may be selected from spike (S), envelope (E), matrix (M), nucleocapsid (N), and Hemagglutinin-esterase glycoprotein (HE). In embodiments, the Coronavirus antigen is derived from a SARS virus. SARS viral antigens are described in WO 04/92360;

[0236] Retrovirus: Viral antigens may be derived from a Retrovirus, such as an Oncovirus, a Lentivirus or a Spumavirus. Oncovirus antigens may be derived from HTLV-1, HTLV-2 or HTLV-5. Lentivirus antigens may be derived from HIV-1 or HIV-2. Retrovirus antigens may be selected from gag, pol, env, tax, tat, rex, rev, nef, vif, vpr, and vpr. HIV antigens may be selected from gag (p24gag and p55gag), env (gp160 and gp41), pol, tat, nef, rev vpr, miniproteins, (e.g., p55 gag and gp140v delete). HIV anti-

gens may be derived from one or more of the following strains: HIV_{IIIb}, HIV_{SF2}, HIV_{LAV}, HIV_{LAI}, HIV_{MN}, HIV-1_{CM235}, HIV-1_{US4}.

[0237] Reovirus: Viral antigens may be derived from a Reovirus, such as an Orthoreovirus, a Rotavirus, an Orbivirus, or a Coltivirus. Reovirus antigens may be selected from structural proteins $\lambda 1$, $\lambda 2$, $\lambda 3$, $\mu 1$, $\mu 2$, $\sigma 1$, $\sigma 2$, or $\sigma 3$, or nonstructural proteins σNS , μNS , or $\sigma 1s$. Reovirus antigens may be derived from a Rotavirus. Rotavirus antigens may be selected from VP1, VP2, VP3, VP4 (or the cleaved product VP5 and VP8), NSP 1, VP6, NSP3, NSP2, VP7, NSP4, or NSP5. Rotavirus antigens may include VP4 (or the cleaved product VP5 and VP8), and VP7. See, e.g., WO 2005/021033, WO 2003/072716, WO 2002/11540, WO 2001/12797, WO 01/08495, WO 00/26380, WO 02/036172; herein incorporated by reference in their entireties.

[0238] Parvovirus: Viral antigens may be derived from a Parvovirus, such as Parvovirus B19. Parvovirus antigens may be selected from VP-1, VP-2, VP-3, NS-1 and NS-2. In embodiments, the Parvovirus antigen is capsid protein VP-2.

[0239] Delta hepatitis virus (HDV): Viral antigens may be derived HDV, particularly δ -antigen from HDV (see, e.g., U.S. Pat. No. 5,378,814).

[0240] Hepatitis E virus (HEV): Viral antigens may be derived from HEV.

[0241] Hepatitis G virus (HGV): Viral antigens may be derived from HGV.

[0242] Human Herpesvirus: Viral antigens may be derived from a Human Herpesvirus, such as Herpes Simplex Viruses (HSV), Varicella-zoster virus (VZV), Epstein-Barr virus (EBV), Cytomegalovirus (CMV), Human Herpesvirus 6 (HHV6), Human Herpesvirus 7 (HHV7), and Human Herpesvirus 8 (HHV8). Human Herpesvirus antigens may be selected from immediate early proteins (α), early proteins (β), and late proteins (γ). HSV antigens may be derived from HSV-1 or HSV-2 strains. HSV antigens may be selected from glycoproteins gB, gC, gD and gH, fusion protein (gB), or immune escape proteins (gC, gE, or gI). VZV antigens may be selected from core, nucleocapsid, tegument, or envelope proteins. A live attenuated VZV vaccine is commercially available. EBV antigens may be selected from early antigen (EA) proteins, viral capsid antigen (VCA), and glycoproteins of the membrane antigen (MA). CMV antigens may be selected from capsid proteins, envelope glycoproteins (such as gB and gH), and tegument proteins.

[0243] Papovaviruses: Antigens may be derived from Papovaviruses, such as Papillomaviruses and Polyomaviruses. Papillomaviruses include HPV serotypes 1, 2, 4, 5, 6, 8, 11, 13, 16, 18, 31, 33, 35, 39, 41, 42, 47, 51, 57, 58, 63 and 65. In embodiments, HPV antigens are derived from serotypes 6, 11, 16 or 18. HPV antigens may include capsid proteins (L1) and (L2), or E1-E7, or fusions thereof. Polyomavirus viruses include BK virus and JK virus. Polyomavirus antigens may be selected from VP1, VP2 or VP3.

[0244] Circovirus: Antigens may be derived from Circoviruses, such as Porcine circovirus (PCV) 1, PCV 2, PCV 3, and PCV 4.

Fungal Antigens

[0245] Suitable fungal antigens may be derived from one or more of the fungi set forth below.

[0246] Fungal antigens may be derived from Dermato-phytes, including: *Epidermophyton floccosum*, *Microsporum audouini*, *Microsporum canis*, *Microsporum distortum*,

Microsporum equinum, *Microsporum gypsum*, *Microsporum nanum*, *Trichophyton concentricum*, *Trichophyton equinum*, *Trichophyton gallinae*, *Trichophyton gypseum*, *Trichophyton megnini*, *Trichophyton mentagrophytes*, *Trichophyton quinckeanum*, *Trichophyton rubrum*, *Trichophyton schoenleini*, *Trichophyton tonsurans*, *Trichophyton verrucosum*, *T. verrucosum* var. *album*, var. *discoides*, var. *ochraceum*, *Trichophyton violaceum*, and/or *Trichophyton faviforme*.

[0247] Fungal pathogens may be derived from *Aspergillus fumigatus*, *Aspergillus flavus*, *Aspergillus niger*, *Aspergillus nidulans*, *Aspergillus terreus*, *Aspergillus sydowii*, *Aspergillus flavatus*, *Aspergillus glaucus*, *Blastoschizomyces capitatus*, *Candida albicans*, *Candida enolase*, *Candida tropicalis*, *Candida glabrata*, *Candida krusei*, *Candida parapsilosis*, *Candida stellatoidea*, *Candida kusei*, *Candida parakwsei*, *Candida lusitanae*, *Candida pseudotropicalis*, *Candida guilliermondi*, *Cladosporium carrionii*, *Coccidioides immitis*, *Blastomyces dermatidis*, *Cryptococcus neoformans*, *Geotrichum clavatum*, *Histoplasma capsulatum*, *Klebsiella pneumoniae*, *Paracoccidioides brasiliensis*, *Pneumocystis carinii*, *Pythium insidiosum*, *Pityrosporum ovale*, *Sacharomyces cerevisiae*, *Saccharomyces boulardii*, *Saccharomyces pombe*, *Scedosporium apiospermum*, *Sporothrix schenckii*, *Trichosporon beigelii*, *Toxoplasma gondii*, *Penicillium marneffei*, *Malassezia* spp., *Fonsecaea* spp., *Wangiella* spp., *Sporothrix* spp., *Basidiobolus* spp., *Conidiobolus* spp., *Rhizopus* spp., *Mucor* spp., *Absidia* spp., *Mortierella* spp., *Cunninghamella* spp., *Saksenaea* spp., *Alternaria* spp., *Curvularia* spp., *Helminthosporium* spp., *Fusarium* spp., *Aspergillus* spp., *Penicillium* spp., *Monoilinia* spp., *Rhizoctonia* spp., *Paeclomyces* spp., *Pithomyces* spp., and *Cladosporium* spp.

[0248] Processes for producing a fungal antigens are well known in the art (see U.S. Pat. No. 6,333,164). In one method, a solubilized fraction extracted and separated from an insoluble fraction obtainable from fungal cells of which cell wall has been substantially removed or at least partially removed, characterized in that the process comprises the steps of: obtaining living fungal cells; obtaining fungal cells of which cell wall has been substantially removed or at least partially removed; bursting the fungal cells of which cell wall has been substantially removed or at least partially removed; obtaining an insoluble fraction; and extracting and separating a solubilized fraction from the insoluble fraction.

Respiratory Antigens

[0249] The compositions of the as disclosed herein may include one or more antigens derived from a pathogen which causes respiratory disease. For example, respiratory antigens may be derived from a respiratory virus such as Orthomyxoviruses (influenza), Pneumovirus (RSV), Paramyxovirus (PIV), Morbillivirus (measles), Togavirus (Rubella), VZV, and Coronavirus (SARS). Respiratory antigens may be derived from a bacteria which causes respiratory disease, such as *Streptococcus pneumoniae*, *Pseudomonas aeruginosa*, *Bordetella pertussis*, *Mycobacterium tuberculosis*, *Mycoplasma pneumoniae*, *Chlamydia pneumoniae*, *Bacillus anthracis*, and *Moraxella catarrhalis*. Examples of specific antigens derived from these pathogens are described above.

[0250] The immunogenic compositions as disclosed herein may be prepared in various forms. For example, the compositions may be prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable for solution

in, or suspension in, liquid vehicles prior to injection may also be prepared (e.g., a lyophilized composition or a spray-freeze dried composition). The composition may be prepared for topical administration e.g., as an ointment, cream or powder. The composition may be prepared for oral administration e.g., as a tablet or capsule or as a spray. The composition may be prepared for pulmonary administration e.g., as an inhaler, using a fine powder or a spray. The composition may be prepared as a suppository or pessary. The composition may be prepared for nasal, aural or ocular administration e.g., as drops. Preparation of such pharmaceutical compositions is within the general skill of the art. See, e.g., Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pa., 18th edition, 1990.

[0251] The composition may be in kit form, designed such that a combined composition is reconstituted just prior to administration to a patient. Such kits may comprise one or more Rotavirus antigens or nucleic acids encoding such antigens in liquid form, and any of the additional antigens and adjuvants as described herein.

[0252] Immunogenic compositions comprising polypeptide antigens as disclosed are vaccine compositions. The pH of such compositions is between 6 and 8, about 7. The pH may be maintained by the use of a buffer. The composition may be sterile and/or pyrogen-free. The composition may be isotonic with respect to the subject. Vaccines according to the instant disclosure may be used either prophylactically or therapeutically, but will typically be prophylactic and may be used to treat animals (including farm, game, companion and laboratory mammals).

[0253] Immunogenic compositions used as vaccines comprise an immunologically effective amount of antigen(s) and/or nucleic acids encoding antigen(s), as well as any other components, as needed. By "immunologically effective amount," it is meant that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment or prevention. This amount varies depending upon the health and physical condition of the individual to be treated, age, the taxonomic group of individual to be treated (e.g., swine, cattle, and the like), the capacity of the individual's immune system to synthesize antibodies, the degree of protection desired, the formulation of the vaccine, the treating veterinarian's assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that may be determined through routine trials.

Administration

[0254] Compositions of as disclosed herein will generally be administered directly to a subject. Direct delivery may be accomplished by parenteral injection (e.g., subcutaneously, intraperitoneally, intravenously, intramuscularly, or to the interstitial space of a tissue), or mucosally, such as by rectal, oral (e.g., tablet, spray), vaginal, topical, transdermal (see, e.g., WO99/27961) or transcutaneous (see e.g., WO02/074244 and WO02/064162), intranasal (see, e.g., WO03/028760), ocular, aural, pulmonary or other mucosal administration. Immunogenic compositions may also be administered topically by direct transfer to the surface of the skin. Topical administration may be accomplished without utilizing any devices, or by contacting naked skin with the immunogenic composition utilizing a bandage or a bandage-like device (see, e.g., U.S. Pat. No. 6,348,450).

[0255] In embodiments, the mode of administration may be parenteral, mucosal or a combination of mucosal and parenteral immunizations. In one aspect, the mode of administration is parenteral, mucosal or a combination of mucosal and parenteral immunizations in a total of 1-2 vaccinations 1-3 weeks apart. In one aspect, the route of administration includes but is not limited to oral delivery, intra-muscular delivery and a combination of oral and intra-muscular delivery.

[0256] It has already been demonstrated that mucosal and systemic immune responses to antigens, such as *Helicobacter pylori* antigens may be enhanced through mucosal priming followed by systemic boosting immunizations. In embodiments, the method for treating an infection by a Rotavirus, comprises mucosally administering to a subject in need thereof a first immunogenic composition comprising Rotavirus antigens followed by parenterally administering a therapeutically effective amount of a second immunogenic composition comprising Rotavirus antigens.

[0257] The immunogenic composition may be used to elicit systemic and/or mucosal immunity, to elicit an enhanced systemic and/or mucosal immunity.

[0258] In embodiments, the immune response is characterized by the induction of a serum IgG and/or intestinal IgA immune response.

[0259] As noted above, prime-boost methods may be employed where one or more gene delivery vectors and/or polypeptide antigens are delivered in a "priming" step and, subsequently, one or more second gene delivery vectors and/or polypeptide antigens are delivered in a "boosting" step. In certain embodiments, priming and boosting with one or more gene delivery vectors or polypeptide antigens described herein is followed by additional boosting with one or more polypeptide-containing compositions (e.g., polypeptides comprising Rotavirus antigens).

[0260] In any method involving co-administration, the various compositions may be delivered in any order. Thus, in embodiments including delivery of multiple different compositions or molecules, the nucleic acids need not be all delivered before the polypeptides. For example, the priming step may include delivery of one or more polypeptides and the boosting comprises delivery of one or more nucleic acids and/or one or more polypeptides. Multiple polypeptide administrations may be followed by multiple nucleic acid administrations or polypeptide and nucleic acid administrations may be performed in any order. Thus, one or more of the gene delivery vectors described herein and one or more of the polypeptides described herein may be co-administered in any order and via any administration route. Therefore, any combination of polynucleotides and polypeptides described herein may be used to elicit an immune reaction.

Dosage Regime

[0261] Dosage treatment may be according to a single dose schedule or a multiple dose schedule. Multiple doses may be used in a primary immunization schedule and/or in a booster immunization schedule. In a multiple dose schedule, the various doses may be given by the same or different routes, e.g., a parenteral prime and mucosal boost, a mucosal prime and parenteral boost, and the like.

[0262] In embodiments, the dosage regime enhances the avidity of the antibody response leading to antibodies with a neutralizing characteristic. An in-vitro neutralization assay may be used to test for neutralizing antibodies.

[0263] There is a strong case for a correlation between serum antibody levels and protection from disease caused by Rotavirus.

Tests to Determine the Efficacy of an Immune Response

[0264] One way of assessing efficacy of therapeutic treatment involves monitoring infection after administration of a composition of the as disclosed. One way of assessing efficacy of prophylactic treatment involves monitoring immune responses against the antigens in the compositions of the as disclosed after administration of the composition.

[0265] Another way of assessing the immunogenicity of the component proteins of the immunogenic compositions of the present disclosure is to express the proteins recombinantly and to screen patient sera or mucosal secretions by immunoblot. A positive reaction between the protein and the patient serum indicates that the patient has previously mounted an immune response to the protein in question—that is, the protein is an immunogen. This method may also be used to identify immunodominant proteins and/or epitopes.

[0266] Another way of checking efficacy of therapeutic treatment involves monitoring infection after administration of the compositions of the present disclosure. One way of checking efficacy of prophylactic treatment involves monitoring immune responses both systemically (such as monitoring the level of IgG1 and IgG2a production) and mucosally (such as monitoring the level of IgA production) against the antigens in the compositions of the present disclosure after administration of the composition. Typically, serum specific antibody responses are determined post-immunization but pre-challenge whereas mucosal specific antibody body responses are determined post-immunization and post-challenge.

[0267] The immunogenic compositions of the present disclosure may be evaluated in in vitro and in vivo animal models prior to host. Particularly useful mouse models include those in which intraperitoneal immunization is followed by either intraperitoneal challenge or intranasal challenge.

[0268] The efficacy of immunogenic compositions of the present disclosure may also be determined in vivo by challenging animal models of infection, e.g., guinea pigs or mice or rhesus macaques, with the immunogenic compositions. The immunogenic compositions may or may not be derived from the same strains as the challenge strains. In embodiments, the immunogenic compositions may be derivable from the same strains as the challenge strains.

[0269] In vivo efficacy models include but are not limited to: (i) A murine infection model using human strains; (ii) a murine disease model which is a murine model using a mouse-adapted strain, such as strains which are particularly virulent in mice and (iii) a primate model using human isolates. A human challenge model, which is supported by the NIH and Center for Disease Control (CDC) is also available.

[0270] The immune response may be one or both of a TH1 immune response and a TH2 response. The immune response may be an improved or an enhanced or an altered immune response. The immune response may be one or both of a systemic and a mucosal immune response. In embodiments, the immune response is an enhanced systemic and/or mucosal response.

[0271] An enhanced systemic and/or mucosal immunity is reflected in an enhanced TH1 and/or TH2 immune response. In embodiments, the enhanced immune response includes an increase in the production of IgG1 and/or IgG2a and/or IgA. In embodiments, the mucosal immune response is a TH2 immune response. In one aspect, the mucosal immune response includes an increase in the production of IgA.

[0272] Activated TH2 cells enhance antibody production and are therefore of value in responding to extracellular infections. Activated TH2 cells may secrete one or more of IL-4, IL-5, IL-6, and IL-10. A TH2 immune response may result in the production of IgG1, IgE, IgA and memory B cells for future protection.

[0273] A TH2 immune response may include one or more of an increase in one or more of the cytokines associated with a TH2 immune response (such as IL-4, IL-5, IL-6 and IL-10), or an increase in the production of IgG1, IgE, IgA and memory B cells. In embodiments, the enhanced TH2 immune response will include an increase in IgG1 production.

[0274] A TH1 immune response may include one or more of an increase in CTLs, an increase in one or more of the cytokines associated with a TH1 immune response (such as IL-2, IFN γ , and TNF β), an increase in activated macrophages, an increase in NK activity, or an increase in the production of IgG2a. In embodiments, the enhanced TH1 immune response will include an increase in IgG2a production.

[0275] Immunogenic compositions of the present disclosure, in particular, immunogenic composition comprising one or more antigens of the present disclosure may be used either alone or in combination with other antigens optionally with an immunoregulatory agent capable of eliciting a Th1 and/or Th2 response.

[0276] The immunogenic composition of the present disclosure may also comprise one or more immunoregulatory agents, such as a mineral salt, such as an aluminum salt and an oligonucleotide containing a CpG motif. In embodiments, the immunogenic composition includes both an aluminum salt and an oligonucleotide containing a CpG motif. Alternatively, the immunogenic composition includes an ADP ribosylating toxin, such as a detoxified ADP ribosylating toxin and an oligonucleotide containing a CpG motif. In one aspect, the one or more immunoregulatory agents include an adjuvant. The adjuvant may be selected from one or more of the group consisting of a TH1 adjuvant and TH2 adjuvant, further discussed above.

[0277] The immunogenic compositions of the present composition may elicit both a cell mediated immune response as well as a humoral immune response in order to effectively address an infection. This immune response may induce long lasting (e.g., neutralizing) antibodies and a cell mediated immunity that may quickly respond upon exposure to one or more infectious antigens. By way of example, evidence of neutralizing antibodies in a subject's blood samples is considered as a surrogate parameter for protection since their formation is of decisive importance for virus elimination in TBE infections.

Use of the Immunogenic Compositions as Medicaments

[0278] The instant disclosure also provides a composition for use as a medicament. The medicament may be able to raise an immune response in a mammal (i.e., it is an immunogenic composition) and may be a vaccine. The

present disclosure also provides the use of the instant compositions in the manufacture of a medicament for raising an immune response in a mammal. The medicament may be a vaccine. In embodiments, the vaccine is used to prevent and/or treat an intestinal infection such as gastroenteritis, including acute gastroenteritis. The gastroenteritis may result from an imbalance in ion and/or water transfer resulting in both watery diarrhea and/or intestinal peristalsis and/or motility (vomiting).

[0279] The instant disclosure provides methods for inducing or increasing an immune response using the compositions described above. The immune response may be protective and may induce antibodies and/or cell-mediated immunity (including systemic and mucosal immunity). Immune responses include booster responses.

[0280] The present disclosure also provides a method for raising an immune response in a mammal comprising the step of administering an effective amount of a composition of the instant disclosure. The immune response may be protective and may involve antibodies and/or cell-mediated immunity. In embodiments, the immune response includes one or both of a TH1 immune response and a TH2 immune response. The method may raise a booster response.

Kits

[0281] The present disclosure also provides kits comprising one or more containers of compositions as described herein. Compositions may be in liquid form or may be lyophilized, as may individual antigens. Suitable containers for the compositions include, for example, bottles, vials, syringes, and test tubes. Containers may be formed from a variety of materials, including glass or plastic. A container may have a sterile access port (for example, the container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle).

[0282] The kit may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution, or dextrose solution. It may also contain other materials useful to the end-user, including other pharmaceutically acceptable formulating solutions such as buffers, diluents, filters, needles, and syringes or other delivery device. The kit may further include a third component comprising an adjuvant.

[0283] The kit may also comprise a package insert containing written instructions for methods of inducing immunity or for treating infections. The package insert may be an unapproved draft package insert or may be a package insert approved by the Food and Drug Administration (FDA) or other regulatory body.

[0284] In embodiments, a delivery device is pre-filled with the immunogenic compositions as disclosed herein.

Methods of Producing Rotavirus Specific Antibodies

[0285] The Rotavirus polypeptides described herein may be used to produce Rotavirus-specific polyclonal and monoclonal antibodies that specifically bind to/are selective for Rotavirus antigens, respectively. Polyclonal antibodies may be produced by administering Rotavirus polypeptides to a mammal, such as a mouse, a rabbit, a goat, or a horse. Serum from the immunized animal is collected and the antibodies are purified from the plasma by, for example, precipitation with ammonium sulfate, followed by chromatography,

including affinity chromatography. Techniques for producing and processing polyclonal antisera are known in the art.

[0286] Monoclonal antibodies directed against Rotavirus-specific epitopes present in the polypeptides may also be readily produced. Normal B cells from a mammal, such as a mouse, immunized with Rotavirus polypeptides, may be fused with, for example, HAT-sensitive mouse myeloma cells to produce hybridomas. Hybridomas producing Rotavirus-specific antibodies may be identified using RIA or ELISA and isolated by cloning in semi-solid agar or by limiting dilution. Clones producing Rotavirus-specific antibodies are isolated by another round of screening.

[0287] Antibodies, i.e., monoclonal and antibodies from polyclonal sera (polyclonal), which are directed against Rotavirus epitopes, are particularly useful for detecting the presence of Rotavirus antigens in a sample, such as a serum sample from a Rotavirus-infected subject. An immunoassay for Rotavirus antigens may utilize one antibody or several antibodies. An immunoassay for Rotavirus antigens may use, for example, a monoclonal antibody directed towards Rotavirus epitopes, a combination of monoclonal antibodies directed towards epitopes of Rotavirus polypeptides, monoclonal antibodies directed towards epitopes of different Rotavirus polypeptides, polyclonal antibodies directed towards the same Rotavirus antigens, polyclonal antibodies directed towards different Rotavirus antigens, or a combination of monoclonal and polyclonal antibodies. Immunoassay protocols may be based, for example, upon competition, direct reaction, or sandwich type assays using, for example, labeled antibody. The labels may be, for example, fluorescent, chemiluminescent, or radioactive.

[0288] The polyclonal or monoclonal antibodies may further be used to isolate Rotavirus particles or antigens by immunoaffinity columns. The antibodies may be affixed to a solid support by, for example, adsorption or by covalent linkage so that the antibodies retain their immunoselective activity. Optionally, spacer groups may be included so that the antigen binding site of the antibody remains accessible. The immobilized antibodies may then be used to bind Rotavirus particles or antigens from a biological sample, such as blood or plasma. The bound Rotavirus particles or antigens are recovered from the column matrix by, for example, a change in pH.

[0289] All patent literature cited in the instant disclosure is incorporated by reference in their entireties herein.

EXAMPLES

Example 1. Baculovirus Protein Subunit Production

[0290] The Recombinant Baculovirus protein expression system was based on nucleic acid sequences for targeted Rotavirus viral proteins (i.e., SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16, 18 and 46). The final sequences were optimized for expression in in-house *Spodoptera frugiperda* insect cells (Sf9) to ensure that appropriate restriction endonuclease sites were present at the termination of the sequences.

[0291] A pBacPAK8 cloning vector (Clontech Laboratories, Inc., Mountain View, CA) was used to prepare plasmid vectors containing the target sequences. The plasmid vectors contain flanking sequences homologous to the linear BestBac 2.0 Baculovirus vector (Expression Systems, Davis, CA), such that when the plasmid containing the various VP4/VP7 inserts were co-transfected into Sf9 cells with the linear BestBac 2.0 virus Baculovirus backbone (Expression

Systems, Davis, CA), homologous recombination exchanges the insert for the polyhedrin gene of the Baculovirus. The resulting Baculovirus containing the Rotavirus viral sequences expressed under control of the polyhedrin promoter were then harvested. Cells and virus were grown in culture media obtained from Expression Systems (Davis, CA, Media ES 99-300) formulated without animal origin ingredients. Gentamicin solution is added to a final concentration of 10 µg/ml from purchased stock solution (Thermo Fisher Scientific, Waltham, MA, Gibco Cat #15710). At final harvest, infected cultures were centrifuged to remove the cells and the supernatants collected. The supernatants were processed through 0.2-micron sterile disposable filter. The premaster cultures were titrated to determine final concentrations.

[0292] Sf9 cells was scaled up to production quantities utilizing glass or sterile disposable plastic vessel volumes. Upon reaching the final cell culture volume required for production, virus infection occurs in the same vessel as the final passage of cells. Culture mixing was achieved through shaking/rocking of the container or utilizing low shear type impeller design. Mixing speed and intensity was adjusted to maintain cells in suspension without creating excess shear or foaming which will cause cell disruption.

[0293] Viral fluids were inactivated with beta-propiolactone (BPL) at a final concentration of 0.2-0.3%. Prior to inactivation, the pH of the disrupted fluids were adjusted to 7.5-8.0 using 2-ION NaOH as base or 10-38% HCl or 10% nitric acid as acid. The disrupted fluids were allowed to warm to room temperature for 1-18 hours prior to the addition of BPL. BPL was added at the concentration specified above, with mixing. After the addition of BPL, the viral fluids were transferred to an inactivation container utilizing a “bottom to bottom” transfer process to ensure that all fluids come into contact with BPL. The disrupted fluids were incubated at 17-27° C. for 18-48 hours with agitation. After the inactivation process was complete, the pH was adjusted to 7.0-7.5 with acid or base as mentioned above. The inactivated virus fluids were stored at 2-8° C. until further processing. The antigens were prepared with Water/Oil/Water (WOW) adjuvants.

Example 2. Evaluation of Rotavirus Antigens Based Vaccine in Mice

[0294] Experimental prescription platform vaccines were produced according to the above outline of production containing either one or two relevant proteins from each isolate (encoding SEQ ID NO:18, VP7 RVA and SEQ ID NO:46, VP4 RVA).

either homologous or heterologous Rotavirus isolates. In addition, viral sequences corresponding to the native virus, or the adapted sequence used in recombinant baculovirus were analyzed for predicted secondary structure and immune epitopes. In a final series of experiments, test vaccines used in mouse studies were administered to feeder pigs, and sera collected for analysis of cross-neutralization with homologous and heterologous virus.

[0296] A field isolate was identified and successfully cultured (termed Rotavirus A1). Based upon viral sequences derived from the genome, baculoviruses were produced independently, each of which expressed a single target viral protein from the homologous virus Rotavirus A1. These proteins were prepared according to the above and formulated either individually or in combination vaccine products that were then administered to separate groups of mice. Serum harvested from these mice was then tested for neutralization of homologous virus, and the results presented in Table 1.

TABLE 1

Serum neutralization titers to Rotavirus A1 following vaccination with various protein targets.	
Protein Target	SN Titer
VP4	1280
	160
VP7	40
	40
VP4 + VP7	1280
	640
VP4 + VP6 + VP7	320
No vaccine	<20

Example 3. Cross-Neutralization of Mouse Sera Against Homologous and Heterologous Virus

[0297] Based upon the results obtained in Example 2, a second study was carried-out to examine the relative importance of sequence conservation of the target proteins in providing protection in mice to either heterologous or homologous virus. Briefly, mice were vaccinated with each of two proteins individually or with a combination of both proteins, and the result sera analyzed for cross-neutralization. The original virus designation, as well as the recombinant baculovirus construct designation are presented in Table 2 for one experiment.

TABLE 2

Protein Sequence Homology between the native viral protein and test baculovirus.						
Baculovirus	Rotavirus Isolate					
	6-091920	3-080820	1-073020	8-100620	7-100620	4-090920
Subunit Protein						
SEQ ID NO: 46	82.3%	82.0%	77.9%	82.8%	82.3%	77.8%
SEQ ID NO: 2	78.8%	78.5%	95.4%	81.9%	N/A	NA

[0295] Mice were vaccinated twice at 3-week intervals with an appropriately reduced dose of the swine vaccine, and serum harvested 28-35 days post second vaccination. Sera was then tested in standard virus neutralization assays using

[0298] Based upon the serological results obtained in vitro, the relative importance of each protein and the importance of sequence homology to the target virus was compared, with the results expressed in Table 3.

TABLE 3

Serum neutralization titers following vaccination with either a combination or single-protein vaccine.						
Vaccine	Rotavirus Isolate					
Subunit Protein	6-091920	3-080820	1-073020	8-100620	7-100620	4-090920
SEQ ID NOs: 46 and 2	Neg	Neg	1:525	Neg	1:390	1:431
SEQ ID NO: 46	Neg	Neg	1:593	Neg	1:403	1:403
SEQ ID NO: 2	Neg	Neg	1:80	Neg	Neg	Neg
Sham	Neg	Neg	1:48	Neg	Neg	Neg

[0299] It was evident that one of the two selected viral proteins appeared to provide the majority of protection observed, and that the sequence homology between the target viral protein and the vaccine construct was not highly predictable.

Example 3. Field Evaluation of Rotavirus Antigens Based Vaccine

[0300] The vaccine study was designed to incorporate general field evaluation to demonstrate Rotavirus A & C vaccine reduction in clinical signs of infection. The primary outcome being to increase economical swine production by decreasing Rotavirus A & C impact.

[0301] Four thousand (4000) doses was considered the minimum number of doses deemed necessary to accurately evaluate if the vaccine is reducing the incidence of Rotavirus. The Rotavirus antigens were manufactured as described above with the Rotavirus A & C genes of interest included (SEQ ID NOs: 1, 3, 5, 7, 9, 11, 13, 15, and 17) to form the combination-antigen vaccine (MEDGENE™s/Applicants' vaccine). A 21-day withdrawal period before slaughter was used.

TABLE 4

Trial Background Data.	
Species/Breed/Strain	Porcine Line 03
Sex	Mixed
Age at Day 0	At least 21 days of age
Number of Doses Used	4000
Housing	Pigs were housed according to farm standards (SOPs)
Feed and Water	Animals were fed age-appropriate feed according to SOPs. Water was available ad libitum.
Vaccination	All sows and gilts received the same vaccine, given as a 2 mL intramuscular dose. Vaccine was used according to label instructions, under the direction of the veterinarian.
Summary and Analysis	Performance parameters were recorded weekly.
Parameters	Evaluation was at 16 weeks.
	Success is litters treated/week reduced from 10-20 to <10 litters/week. Success is an improvement in pre-wean mortality from 20% to <15%.
Farrowing Piglet	(i) Executed when observes three consecutive weeks of litter diarrhea incidence in the vaccination time-period that is greater than the baseline litter diarrhea incidence.
Diagnostic Workup	(ii) Executed at discretion. (iii) Tissue submissions for pathology was defined as 3-5 piglets with clinical diarrhea.

TABLE 4-continued

Trial Background Data.	
The enteric workup objective was to determine diarrhea etiology.	

[0302] The Trial started with breeding group 10 and data was collected past breeding group 21. Each sow's litter's score was based on a score card (not shown). The scores were as follows:

[0303] Score 0

[0304] Clean dry pigs

[0305] No loose stools

[0306] Score 1

[0307] Majority of litter at Score 0

[0308] 1-2 with loose stool

[0309] Score 2

[0310] Less than about half of litter with loose stools

[0311] Score 3

[0312] Half of litter with loose stool

[0313] Evidence of loose stool on all pigs

[0314] Predominantly dry

[0315] Few to most pigs with red anus and visibly wet

[0316] Score 4

[0317] More than half of litter with loose stool

[0318] Score 5

[0319] Entire litter with loose stool

[0320] Wet and chilled pigs

[0321] Loss in body condition

[0322] Scores were collected 4 days post farrowing. Groups 10-15 & 18 were mixed groups, Groups 16-17 were all competitor (SEQUIVITY™, Merck Animal Health, Rahway, NJ; "RNA-based vaccine") vaccine, Groups 19-21 were all MEDGENE™ (Applicants') vaccine.

TABLE 5

Trial Parity Results.	
Breeding Groups	4
Sows on Trial	1109
	MEDGENE™ Vaccine - 565
	RNA-based Vaccine - 544
Parity* Breakdown	MEDGENE™ Vaccine
	P1 - 173
	P2 - 64
	P3 - 89
	P4 - 78
	P5+ - 166
	RNA-based Vaccine
	P1 - 106
	P2 - 87

TABLE 5-continued

Trial Parity Results.	
	P3 - 99
	P4 - 81
	P5+ - 166

*Parity - is a swine age/reproduction measurement. P1 - the female has given birth (farrowed) to her first litter of pigs, Parity 2 - female has farrowed her 2nd litter of pigs, P3 - third and so on.

[0323] MEDGENE™ vaccine contained a combination of peptides as set forth in SEQ ID NOs: 2 (VP7 RVA), 4 (VP4 RVA), 6 (VP4 RVA), 8 (VP7 RVC), 10 (VP4 RVC), 12 (VP4 RVA), 14 (VP4 RVC), 16 (VP7 RVC) and 18 (VP7 RVC).
[0324] FIGS. 1 and 2 show the results of the trials. As may be seen in FIG. 1, the MEDGENE™ vaccine has lower

scores, and thus, is more effective than the RNA-based vaccine. Further, from FIG. 2, it may be seen that (though not statistically significant) there is a more positive trend for Parity for the MEDGENE™ vaccine vs. the RNA-based vaccine.

[0325] Although the invention has been described with reference to the above examples, it will be understood that modifications and variations are encompassed within the spirit and scope of the invention. Accordingly, the invention is limited only by the following claims.

[0326] The sequence listing submitted on 2024 Mar. 1 has the file name “Rotavirus.xml”, was created on 2024 Mar. 1, and has a file size of 81 KB, which sequence listing is incorporated herein by reference.

SEQUENCE LISTING

Sequence total quantity: 52

SEQ ID NO: 1 moltype = DNA length = 1058
 FEATURE Location/Qualifiers
 source 1..1058
 mol_type = other DNA
 organism = Rotavirus A

SEQUENCE: 1
 accgagctcg aattcatgta tgggaatcgaa tacacaaccg tccttacctt ccttatatct 60
 ataactccttc tcaattatat attgaagtct ctgacgagca ccatggattt tatcatctac 120
 agattcctcc tgcttattgc gatagtgagt ccatttggta agacgcaaaa ttacgggaatc 180
 aacctcccaa taacaggctc tatggatacg gcatacgcaa actcatcaca acaagaaacg 240
 tttcttactt ctacattgtg cctttactac ccgactgagg ctagcactca gattggcgat 300
 gctgagtggg agaacacact ttcccaactc tttctcacta aaggctggcc gactgggtcg 360
 gtatatattta aggagtacac aaatatcgct tccttttcca tcgatccaca actgtactgc 420
 gattacaacg tagttttgat gaaatatgac tcaaccctgg aattggacat gtcggagctt 480
 gcggacetta ttctcaacga gtggctgtgt aatccgatgg acatcacatt gtactattat 540
 cagcagacgg acgaagcgaa caaatggatt agcatgggtc agtcttgac tggttaaagt 600
 tgcccaactca atacacaaac tctgggcata ggatgcacca ctacaatac tgcgacgttc 660
 gaagaagtag caagagatga aaaatttgga atcacgatg tcgtcgatgg agtcaacccat 720
 aaactggacg ttactaccaa tacatgtacg ataagaaatt gcaaaaaatt gggaccgaga 780
 gaaaaatgtt ccataataca agttggtggt agtgaggtcc ttgatgttac tgctgacccc 840
 actaccgctc cgcagacaga gcgcatgatg agggtaaaact ggaagaaatg gtggcaagtt 900
 ttttatactg tcgtagatta cataaaccag atcgtccaag tcatgagtaa aagatctctg 960
 tcaactgaact ctgcccgctt ttattatagg gtgaaggcg aaacttgta ctttcaaggc 1020
 catcaccatc accatcacta ggccggccgt taattaat 1058

SEQ ID NO: 2 moltype = AA length = 1023
 FEATURE Location/Qualifiers
 source 1..1023
 mol_type = protein
 organism = Rotavirus A

SEQUENCE: 2
 METTYRGLYI LEGLUTYRTH RTHRVALLEU THRPHELEUI LESERILEIL ELEULEUASN 60
 TYRILELEUL YSSERLEUTH RSERTHRMET ASPPHEILEI LETYRARGPH ELEULEULEU 120
 ILEALAILLEV ALSERPROPH EVALYSTHR GLNASNTYRG LYILEASNLE UPROILETHR 180
 GLYSERMETA SPTHRALTATY RALAASNSER SERGLNGLNG LUTHRPHELE UTHRSERTHR 240
 LEUCYSLEUT YRTYRPROTH RGLUALASER THRGLNILEG LYASPALAGL UTRPLYSASN 300
 THRLEUSERG LNLEUPHELE UTHRLYSGLY TRPPROTHRG LYSERVALTY RPHELYSGLU 360
 TYRTHRASNI LEALASERPH ESERILEASP PROGLNLEUT YRCYSASPTY RASNVALVAL 420
 LEUMETLYST YRASPSERTH RLEUGLULEU ASPMETSERG LULEUALAAS PLEUIELELU 480
 ASNGLUTRPL EUCYSASNPR OMETASPILE THRLEUTYRT YRTYRGLNGL NTHRASPGLU 540
 ALAASNLYST RPILERERME TGLYGLNSER CYSTRHVALL YSVALCYSPR OLEUASNTHR 600
 GLNTHRLEUG LYILEGLYCY STHRTHRTHR ASNTHRALAT HRPHEGLUGL UVALALAARG 660
 ASPLGLULYS EUVALILETH RASPVALVAL ASPGLYVALA SNHISLYSLE UASPVALTHR 720
 THRASNTHRC YSTHRILEAR GASNCYSLYS LYSLEUGLYP ROARGGLUAS NVALALAILL 780
 ILEGLNVALG LYGLYSERGL UVALLEUASP VALTHRALAA SPPROTHRTH RALAPROGLN 840
 THRGLUARGM ETMETARGVA LASNTRPLYS LYSTRPTRPG LNVALPHETY RTHRVALVAL 900
 ASPTYRILEA SNGLNILEVA LGLNVALMET SERLYSARGS ERARGSERLE UASNSEALA 960
 ALAPHETYRT YRARGVALLY SGLYGLUASN LEUTYRPHEG LNGLYHISHI SHISHISHIS 1020
 HIS 1023

SEQ ID NO: 3 moltype = DNA length = 2408
 FEATURE Location/Qualifiers
 source 1..2408
 mol_type = other DNA

-continued

organism = Rotavirus A

SEQUENCE: 3

accgagctcg	aattcatggc	gagcttgatt	taccgacaat	tacttacgaa	ctcctacact	60
gtggatctgt	cggacgaaat	tcagattatt	gggagcgaga	aggcccaaaa	tgtgaccatc	120
aatccggggc	catttgctca	aactgggtat	gcacccgta	actggggacc	aggtgagacc	180
aatgatagta	cgaccgtaga	gccggtgctt	gacggaccct	atcagccgac	cacgtttaac	240
cctcctgtaa	gctattggat	cctcttatca	ccgtcgaacg	ctggcgctgt	cgtagagggt	300
acaaacaact	cagatagatg	gctcgcaacc	attttaatcg	agccaaatgt	gacttctgag	360
tcgcgacacat	atacactctt	cggccgcgaa	gagcagatca	ccgtggagaa	tgtcagtagc	420
acgaagtgga	aattttaga	cctcgcgaa	accgacataa	atggaacgtt	cactcaacac	480
ggggcactgc	tctcggtatc	taagctttac	ggggttatga	agttttctgg	cagattgtac	540
acgtataatg	gagagactcc	caacgcccact	accggttatt	acaccaccac	caactacgat	600
actgtaaata	tgacatcaca	ttgcgatttc	tacatcatcc	cccgtagtga	agagaatgcc	660
tgtacaaact	acataaataa	tggtctacca	ccgatccaga	ataccagaaa	cgctgttcca	720
gttagcctca	cttctcgctc	tattgtacat	acacgagcac	aggcaaacga	ggacattgtg	780
gttagtaaaa	cagcctgtgt	gaaagaaatg	cagtataacc	gcgatattac	gatccgtttc	840
aagttcgcta	acgctattat	taagagtgga	ggcctggggg	acaagtgggt	tgagataaag	900
tttaagccag	ccaactacca	gtacaactac	accogtgatg	gtgaggttat	aactgcccac	960
acgacctgct	cggtaaatgg	gataaacgac	ttcagttaca	acggcggtag	cctacactacg	1020
gattttgtat	ttagtagata	cgaggctcatt	aaggagaaca	gttacgtgta	cgtagactac	1080
tggtgatgaca	gcccaagcatt	ccgcaacatg	gtctatgtgc	gtagtcttgc	ggcgaatcta	1140
aactcagtgta	cctgtacttg	cggaactat	gactttcgtc	ttccggtggg	ggcttggcca	1200
gtaatgactg	gaggggctgt	cagctctacg	cctgtgtggg	tcacactgag	cactcaattt	1260
acggattttg	tatcactaaa	cagcctacga	tttaggttct	ccctgtctgt	agaggagccg	1320
ccctttgcaa	tagccggaac	gcgagttagc	ggactctacg	ggctccctgc	tgccaaccct	1380
aataacggta	aggactacta	tgagattctg	ggtagggttt	ccctaactctc	actagtctct	1440
agtaacgacg	actatcagac	cccagataatg	aacagcgtca	ccgtgaggca	ggatttagaa	1500
agacaaactca	acgaactctg	acaggagtct	aattctctga	gtcaagaaat	agcgatgagt	1560
cagttgatag	atctggcact	gctaccattg	gatattgtct	caatgttttc	tggtataaag	1620
tcgacaatag	atgcggcgaa	aagcatggcg	acctcggtta	tgaagaagt	caagcgcagt	1680
aatctagcta	gcagactttc	gacgcttact	gactcactaa	gcgacgcagc	ctcttcgata	1740
tcacgttaacc	catccatacg	cagcgtcgga	tcctcagtc	gtgtgtggac	ggacatctcc	1800
acacaggcga	ctgatattat	tgataccttt	aaatcagtg	gtacccaaac	gtcgacaatt	1860
tcaagaagat	tacgaactta	ggaaattact	acccaaacag	aaggtatgaa	ttttgatgat	1920
atcagcgcg	cagtactgaa	aacccaaatc	gataaatctg	ttcaaatctc	acccaaacaca	1980
ctaccggaca	tcgttacaga	agcaagtga	aagttcatcc	ctaactcgta	ataccgtgta	2040
attaaagatg	acgaggttct	ggaggctggg	actgacggac	gtttcttcgc	ctataaagta	2100
gatacctttg	acgaaatacc	tttcgatgtc	cagaagtgtg	cagatcttgt	gacggatagt	2160
cctgtcataa	gtgcaatcat	agattttaaa	acattgaaaa	atctgaatga	caactatggc	2220
atcacaagg	aaaagccctt	taatctgttg	agatccgacc	caagggtgct	aagggagtgc	2280
atcaatcaag	aaaatccgat	aatacgtaat	cgcattgaac	agcttatact	acaatgtaga	2340
cttaaggcg	aaaacttgta	ctttcaaggc	catcaccatc	accatcacta	ggcgccgct	2400
taattaat						2408

SEQ ID NO: 4 moltype = AA length = 2373

FEATURE Location/Qualifiers

source 1..2373

 mol_type = protein

 organism = Rotavirus A

SEQUENCE: 4

METALASERL	EUILEYRAR	GGLNLEULEU	THRASNERT	YRTHRVALAS	PLEUSERASP	60
GLUILEGLNI	LEILEGLYSE	RGLULYSALA	GLNASNVALT	HRILEASNPR	OGLYPROPHE	120
ALAGLNTHRG	LYTYRALAPR	OVALASNTRP	GLYPROGLYG	LUTHRASNAS	PSERTHRTHR	180
VALGLUPROV	ALLEUASPGL	YPROTYRGLN	PROTHRTHRP	HEASNPROPR	OVALSERTYR	240
TRPILELEUL	EUSERPROSE	RASNALAGLY	VALVALVALG	LUGLYTHRAS	NASNSEASP	300
ARGTRPLEUA	LATHRILELE	UILEGLUPRO	ASNVALTHRS	ERGLUSERAR	GTHRTYRTHR	360
LEUPHEGLYA	RGGLNGLUGL	NILETHRVAL	GLUASNVALS	ERTHRTHRLY	STRPLYSPE	420
VALASPLEUA	LALYSTHRAS	PILEASNGLY	THRPETHRG	LNHISGLYPR	OLEULEUSER	480
ASPTHRLYSL	EUTYRGLYVA	LMETLYSPHE	SERGLYARGL	EUTYRTHRTY	RASNGLYGLU	540
THRPROASNA	LATHRTHRGL	YTYRTYRTHR	THRTHRASNT	YRASPTHRVA	LASNMTTHR	600
SERHISCYSA	SPPHETRYIL	EILEPROARG	SERGLUGLUA	SNALACYSTH	RASNTYRILE	660
ASNASNGLYL	EUPROPROIL	EGLNASNTHR	ARGASNVALV	ALPROVALSE	RLEUTHRSE	720
ARGSERILEV	ALHISTHRAR	GALAGLNALA	ASNGLUASPI	LEVALVALSE	RLYSTHRSE	780
LEUTRPLYSG	LUMETGLNLY	SARNARGASP	ILETHRILEA	RGPHELYSPH	EALASNALA	840
ILEILELYSS	ERGLYGLYLE	UGLYTYRLYS	TRPSERGLUI	LESERPHELY	SPROALAASN	900
TYRGLNLYRA	SNTRYRTHAR	GASPGLYGLU	VALILETHRA	LAHISTHRTH	RCYSSSERVAL	960
ASNGLYILEA	SNASPPHESE	RTYRASNGLY	GLYSERLEUP	ROTHRASPPH	EVALILESER	1020
ARGTYRGLUV	ALILELYSGL	UASNSERTYR	VALTYRVALA	SPTYRTRPAS	PASPSEGLN	1080
ALAPHEARGA	SNMETVALTY	RVALARGSER	LEUALAALAA	SNLEUASNSE	RVALTHRCYS	1140
THRGLYGLYA	SNTRYASPPH	EARGLEUPRO	VALGLYALAT	RPPROVALME	TTHRGLYGLY	1200
ALAVALSERL	EUARGPROAL	AGLYVALTHR	LEUSERTHRG	LNPETHTRAS	PPHEVALSER	1260
LEUASNSERL	EUARGPHEAR	GPHESEERLEU	SERVALGLUG	LUPROPROPH	EALALEALA	1320
ARGTHRARGV	ALSERGLYLE	UTYRGLYLEU	PROALAALAA	SNPROASNAS	NGLYLYSASP	1380
TYRTYRGLUI	LELEUGLYAR	GPHESEERLEU	ILESERLEUV	ALPROSERAS	NASPASPTYR	1440
GLNTHRPROI	LEMETASNSE	RVALTHRVAL	ARGGLNASPL	EUGLUARGGL	NLEUASNGLU	1500
LEUARGGLNG	LUPHEASNSE	RLEUSERGLN	GLUILEALAM	ETSERGLNLE	UILEASPLEU	1560

-continued

ALALEULEUP	ROLEUASPM	TPHERMET	PHESERGLYI	LELYSSERTH	RILEASPALA	1620
ALALYSSERM	ETALATHRSE	RVALMETLYS	LYSPHELYSA	RGSERASNLE	UALASERSE	1680
ILESERTHRL	EUTHRASPE	RLEUSERASP	ALAALASERS	ERILESERAR	GASNPROSER	1740
ILEARGSERV	ALGLYERSE	RVALSERVAL	TRPTHRASPI	LESERTHRL	NALATHRASP	1800
ILESERASPT	HRPHELYSSE	RVALSERTHR	GLNTHRSERT	HRILESERAR	GARGLEUARG	1860
LEULYSGLUI	LETHRTHRL	NTHRGLUGLY	METASNPHEA	SPASPILESE	RALAALAVAL	1920
LEULYSTRHL	YSILEASPLY	SSERVALGLN	ILESERPROA	SNTHRLEUPR	OASPILEVAL	1980
THRGLUALAS	ERGLULYSPH	EILEPROASN	ARGTHRTYRA	RGVALILELY	SASPASPGLU	2040
VALLEUGLUA	LAGLYTHRAS	PGLYARGPHE	PHEALATYRL	YSVALASPTH	RPHEASPGLU	2100
ILEPROPHEA	SPVALGLNLY	SPHEALAASP	LEUVALTHRA	SPSERPROVA	LILESERALA	2160
ILEILEASPP	HELYSTRHLE	ULYSASNLEU	ASNASPASNT	YRGLYILETH	RARGGLUGLN	2220
ALAPHEASNL	EULEUARGSE	RASPPROARG	VALLEUARGG	LUPHEILEAS	NGLNGLUASN	2280
PROILEILEA	RGASNARGIL	EGLUGLNLEU	ILELEUGLNC	YSARGLEULY	SGLYGLUASN	2340
LEUTYRPHEG	LNGLYHISHI	SHISHISHIS	HIS			2373

SEQ ID NO: 5 moltype = DNA length = 1076
 FEATURE Location/Qualifiers
 source 1..1076
 mol_type = other DNA
 organism = Rotavirus A

SEQUENCE: 5

accgagctcg	aattcatggt	ttgcaccacc	ctctacacgg	tctgcgtgat	cctgtgtatt	60
ctcttcattt	acatgtgtct	tttccgtaag	atgttccacc	tcattacgga	taccgtaata	120
gtacgcataa	taatctccaa	tgtgtgtggc	tggactcaag	gtcaaatggt	catcgacaat	180
atgcattacg	agggaatat	cgagacgatt	atcaatgcta	cgaatccatt	taacgtagaa	240
agtctttgca	tatatctccc	aaaagcgggt	gttggaaagtc	agggaccogg	ccaaatcaac	300
ggatacctga	acgacggaaa	ctacgccccaa	accatagcta	cactgttcga	gactaagggc	360
ttccctaagg	gaagtgtggt	gctgaagact	tacgcaaaga	caagtgactt	tatctcttcg	420
gtagagatga	cgtgttctga	taacatagtc	atcatccggg	atgatccgga	taagtctgag	480
gcaatcgaa	agatagcaga	gtggatattg	aatgtatgga	agtgtgatga	tatggatctt	540
aacatataca	cgtacgaaca	aatagggaata	gacaaccttt	gggctgcatt	tggcgacgac	600
tgtgacatta	gcgtttgccc	gctcgatacg	accatgaatg	gcataggatg	ccttcgggcg	660
agtacagaaa	cgtacgaggt	ggatctccaa	aacaccgagt	tggccctgat	taatgtcgta	720
gataacgtaa	aacataggat	acagatgaac	acaggcagat	gcaaacctaa	aaattgcgtc	780
aaggggtgag	cacgcctgaa	cacggctata	atccgcata	ccaagtctag	tagtttcgac	840
aatagtctca	gtccaatgaa	caacggacag	actacgagat	cgttcaaaat	caacgccccaa	900
aaatggtgga	cgtctctcta	tacaatcatc	gactatatta	atcagattgt	acaaacgatg	960
acaccgaggg	cgcgcgccat	ttatccagag	ggttggatgc	tccgctatgc	gaaggcgcaa	1020
aacttgctact	ttcaaggcca	tcaccatcac	catcactagg	cggccgctta	attaat	1076

SEQ ID NO: 6 moltype = AA length = 1041
 FEATURE Location/Qualifiers
 source 1..1041
 mol_type = protein
 organism = Rotavirus A

SEQUENCE: 6

METVALCYST	HRTHRLEUTY	RTHRVALCYS	VALILELEUC	YSILELEUPH	EILETYRMET	60
LEULEUPHEA	RGLYSMETPH	EHISLEUILE	THRASPTHR	ALILEVALAL	AILEILEILE	120
SERASNYSV	ALGLYTRPTH	RGLNGLYGLN	METPHEILEA	SPASNMTTHI	STYRGLUGLY	180
ASNILEGLUT	HRILEILEAS	NALATHRASN	PROPHEASNV	ALGLUSERLE	UCYSILETYR	240
PHEPROLYSA	LAVALVALGL	YSERGLNGLY	PROGLYGLNI	LEASNGLYTY	RLEUASNASP	300
GLYASNTYRA	LAGLNTHRL	EALATHRLEU	PHEGLUTHRL	YSGLYPHEPR	OLYSGLYSER	360
VALVALLEUL	YSTHRTYRAL	ALYSTHRSER	ASPPHEILES	ERSERVALGL	UMETHTRCYS	420
SERTYRASNI	LEVALILEIL	EPROASPASP	PROASPLYSS	ERGLUALAIL	EGLUGLNILE	480
ALAGLUTRPI	LELEUASNVA	LTRPLYSCYS	ASPASPMETA	SPLEUASNIL	ETYRTHRTYR	540
GLUGLNILEG	LYILEASPAS	NLEUTRPALA	ALAPHEGLYA	SPASPCYSAS	PILESERVAL	600
CYSPROLEUA	SPTHRTHRME	TASNGLYILE	GLYCYSLLEUP	ROALASERTH	RGLUTHRTYR	660
GLUVALVALS	ERASNASNT	RGLULEUALA	LEUILEASN	ALVALASPAS	NVALLYSHIS	720
ARGILEGLNM	ETASNTHRGL	YARGCYSLYS	LEULYSASNC	YSVALLYSGL	YGLUALAARG	780
LEUASNTHRA	LAILEILEAR	GILESERLYS	SERSERSERP	HEASPASNSE	RLEUSERPRO	840
METASNASNG	LYGLNTHRTH	RARGSERPHE	LYSILEASNA	LALYSLYSTR	PTRPTHRILE	900
PHETYRTHRI	LEILEASPTY	RILEASNTHR	ILEVALGLNT	HRMETTHRPR	OARGHISARG	960
ALAILETYRP	ROGLUGLYTR	PMETLEUARG	TYRALALYS	LYGLUASNLE	UTYRPHEGLN	1020
GLYHISHISH	ISHISHISHI	S				1041

SEQ ID NO: 7 moltype = DNA length = 1076
 FEATURE Location/Qualifiers
 source 1..1076
 mol_type = other DNA
 organism = Rotavirus C

SEQUENCE: 7

accgagctcg	aattcatggt	gtgtacaacc	ctgtatacgg	tttgcgttat	tttgtgcatt	60
ttgttcattt	acatgctttt	gttccgcaaa	atgttccacc	tcatacagga	tgctacaatc	120
ataacccttg	tgatatactaa	tgttattgga	tggtcacaa	gtcagatggt	tacagacgac	180
atctattaca	acggaacgt	cgaactgtg	gtgaactcga	ctgatccatt	tgacgtgaga	240
agtctttgca	tttactttcc	taacgcgctc	gtcggatcgc	agggacctgg	aaaaactgat	300

-continued

```

ggatacctca acgatggttaa ttatgcccag actattgccg cactcttcga aacgaagggt 360
tttccccgtg gatcgatcgt tctcaagaca tatacgaaag tctcagactt cataaattcg 420
gtggaaatga cgtgctcgta taacatcgta attataccgg actcgttgac caacagcgaa 480
tcaatcgaga ggattgcaga gtggatcctt aatggtttggc gttgtgacga catgaatcct 540
gatatctaca catacgaaca gacaggtatc gacaacttgt gggctgcgtt cggtagcgac 600
tgcgacatca gcgtatgtcc gttggacaca accatgaacg gcctcggctg tagtccagca 660
tcaacagaaa cgtatgaagt gctttccaac gatactcagc tcgcccctgt taacgttgtg 720
gacaacatta aacatcgcat tcagatgaat accgcaactt gtaaaactaa aaactgcac 780
aaaggagaa cagactctaa taccgcgctg attagaatct caacatccag ctccctcgac 840
aactcgctga gcccgctcaa tgacggccaa acaacaaggt cttttaagat aaacgcgaag 900
aagtgggtga cgattttcta cacgatcata gactacatca ataccatag acagactatg 960
gcgcccagac acagagcgat atatectgag ggttggtatgc tgcgttatgc gaaggcgaa 1020
aactgtact ttcaaggcca tcaccatcac catcactagg cggccgctta attaat 1076

```

SEQ ID NO: 8 moltype = AA length = 1041

FEATURE Location/Qualifiers

source 1..1041

mol_type = protein

organism = Rotavirus C

SEQUENCE: 8

```

METVALCYST HRTHRLEUTY RTHRVALCYS VALILELEUC YSILELEUPH EILETYRMET 60
LEULEUPHEA RGLYSMETPH EHISLEUILE THRASPALAT HRILEILETH RLEUVALILE 120
SERASNCYSI LEGLYTRPSE RGLNGLYGLN METPHETHRA SPASPILETY RTYRASNGLY 180
ASNVALGLUT HRVALVALAS NSERTHRASP PROPHEASPV ALARGSERLE UCYSILEITYR 240
PHEPROASNA LAVALVALGL YSERGLNGLY PROGLYLYST HRASPGLYTY RLEUASNASP 300
GLYASNTYRA LAGLNTHRIL EALAALALEU PHEGLUTHRL YSGLYPHEPR OARGGLYSER 360
ILEVALLEUL YSTHRTYRTH RLYSVALSER ASPPHEILEA SNSERVALGL UMETTHRCYS 420
SERTYRASNI LEVALILEIL EPROASPSPER LEUTHRASNS ERGLUSERIL EGLUARGILE 480
ALAGLUTRPI LELEUASNVA LTRPARGCYS ASPASPMETA SNLEUASPIL ETYRTHRTYR 540
GLUGLNTHRG LYILEASPAS NLEUTRPALA ALAPHEGLYS ERASPCYSAS PILESERVAL 600
CYSPROLEUA SPTHRTHRME TASNGLYILE GLYCYSSERP ROALASERTH RGLUTHRTYR 660
GLUVALLEUS ERASNASPTH RGLNLEUALA LEULEUASN V ALVALASPAS NILELYSHIS 720
ARGILEGLNM ETASNTHRAL ATHRCYSLYS LEULYSASNC YSILELYSGL YGLUALAARG 780
LEUASNTHRA LALEUILEAR GILESERTHR SERSERSERP HEASPASNSE RLEUSERPRO 840
LEUASNASPG LYGLNTHRTH RARGSERPHE LYSILEASNA LALYSLYSTR PTRPTHRILE 900
PHETYRTHRI LEILEASPTY RILEASNTHR ILEVALGLNT HRMETALAPR OARGHISARG 960
ALAILETYRP ROGLUGLYTR PMTLEUARG TYRALALYSG LYGLUASNLE UTYRPHEGLN 1020
GLYHISHISH ISHISHISHI S 1041

```

SEQ ID NO: 9 moltype = DNA length = 2294

FEATURE Location/Qualifiers

source 1..2294

mol_type = other DNA

organism = Rotavirus C

SEQUENCE: 9

```

accgagctcg aattcatggc atcaagcttg taccagcaat tgatttcgca gaattattac 60
tctaccggca acgagatact tctcgataag caaacgacac gcacaactaa ggattttgta 120
gaagcaggaa attatactta cgtcgaactg ccacctacgg aatggggcgc aggcctcgaca 180
tttgagtcga cgtttaaaag ttctaataa acaggtcctc atagcaatgc cgttattgag 240
tggaataacc tcataaattc ggatatctgg gtattgtacc agaagccact cgacatcacc 300
gttccaatcc gtttgctcaa gcacggcccc gagaattacg gaaatgtagc ggcgtttgag 360
ttgtggtagc gtaaggcggg tcatactggt actagcattt actactccgc gattagtaac 420
ccaaataaca cgggttacgtt gacttctgat agtttgggtc ttttctggaa cgaaggtcaa 480
actacccttg ataaaaagac tgttaacttt aattggaata tgggcggaat ttgggtccgt 540
cctagttagg gaacgagagt agatatctgt atgtcgaaac tggataaacac agatgatgca 600
tcatttaatt ggattcagtg gaaacacgag ttcccagatg cctcaaattc agctaacgct 660
aacatgtatg tagaatacta ccttgcatcc agcgatccat atcacgagt gaaagagctc 720
caacgcccta cagtgacgac catcaatatg caaatgatga aagcagtgaa cggtgacttg 780
atcaatgtag acgaggttat ctccaaagac tccctttggc aagagatgag ctatgttagg 840
gatataactc tcgagtgtaa gatcgagctc gaagtattta aaggcggtgg ctggggctat 900
gattacacaa gcctacacct caaaaccgta aatcacacat acagttacaa ccgcgcggcg 960
gagaatgtga cgcgccacgt gacagtgtcc ttcaataaca tgaagggaac ctccctacgga 1020
ggctcacttc ctacggattt caagataaac aggtttgacg tgatggatat cgacacctat 1080
atgtatattg actattggga cgactccgat atgttcaaga acatgggttta tgtacgcgat 1140
tcaggggcag atgtgaacgg ctctctttac tataaagaga tgtcctattc gtttagaatc 1200
cccgttggtg tacacccaag gataaatagt agtggatcgc agttcacgta cgaaaagtgc 1260
atgttgagtc aacagttcac cgaccagttg gcactgaaca gcatgaggtt tgtattttcg 1320
gttacatctg cgaaacggctg gtttatgacc tcaggcggac ccaatacagag gcgcatagct 1380
tcaggcactg gcttcgcata tagtgatggc cacacttcga gaacagtcgg tacactcagt 1440
ttcatctcgc tcattcccttc taatcccaac taccagacgc ccatagctag ctccctcgag 1500
gtcaggatgg accctgacgc caagattaac gatctgagga acgatttcaa ccagctcgcg 1560
agctccgtcg cactcggaga cattatgtct ctggctatgt cacctctcac tttttctaat 1620
cttctggagt cgttacggcg aataaactagt agtgtgaaag atgttgccgc caacgtaagt 1680
aaaaaatttc gtaacactaa gatgtttaag aaagccacaa aacctaaata taaggagtgc 1740
tcatttggtg atttgcttga ggatgtaacg aatgtgacca agaactccaa taccatgaac 1800
tttgacgata tcacgtctgc cgtgatgata agtacgacaa acaagctgca attgactgac 1860

```

-continued

gtagacaccc	tgtcagaat	tgtcgtagg	tctgcagatg	actttatacc	gtcccgtagc	1920
taccgcatga	tgcagaataa	cacgggtgat	gaagcgacac	catcacgtgt	gttttcttac	1980
gatcttgcaa	ccatgcagca	gagagacttc	gacattgata	agtttatcca	gcttgcttca	2040
aagtcaccgc	taatttcagc	gategttgat	ttcgccacac	tgaaggccat	gagggagacg	2100
tatggaacat	cggtcgaagt	gattcataag	cttgtggcct	cggattctcc	aacaatattg	2160
tctttcatca	ataataacaa	tccacttatc	aggaacagga	tcgaagaact	gctcagacag	2220
tgccgcatta	agggcgaaaa	cttgtacttt	caaggccatc	accatcacca	tcactaggcg	2280
gccgcttaat	taat					2294

SEQ ID NO: 10 moltype = AA length = 2259
 FEATURE Location/Qualifiers
 source 1..2259
 mol_type = protein
 organism = Rotavirus C

SEQUENCE: 10

METALASERS	ERLEUTYRGL	NGLNLEUILE	SERGLNASNT	YRTYRSERTH	RGLYASNGLU	60
ILELEULEUA	SPLYSGLNTH	RTHRARGTHR	THRLYSASPP	HEVALGLUAL	AGLYASNTYR	120
THRTYRALAG	LNLEUPROPR	OTHRGLUTRP	GLYALAGLYS	ERTHRPHEGL	USERTHRPHE	180
LYSSERSERA	SNILETHRGL	YPROHISSE	ASNALAVALI	LEGLUTRPAS	NASNLEUILE	240
ASNSERASPI	LETROPVALLE	UTYRGLNLYS	PROLEUASPI	LETHRVALPR	OILEARGLEU	300
LEULYSHISG	LYPROGLUAS	NTYRGLYASN	VALALAALAP	HEGLULEUTR	PTYRGLYLYS	360
ALAGLYHIHT	HRVALTHRSE	RILETYRTYR	SERALAILES	ERASNPROAS	NASNTHRVAL	420
THRLEUTHRS	ERASPSEERLE	UVALLEUPHE	TRPASNGLUG	LYGLNTHRTH	RLEUASPLYLS	480
LYSTHRVALA	SNPHEASNTR	PASNMTGLY	GLYILELEUV	ALARGPROSE	RARGGLYTHR	540
ARGVALASPI	LECYSMETSE	RASNMTASP	ASNTHRASPA	SPALASERPH	EASNTRPILE	600
GLNTRPLYSH	ISGLUPHEPR	OARGSERSER	ASNSERALAA	SNVALASNME	TTYRVALGLU	660
TYRTYRLEUA	LASERSERAS	PPROTyrHis	GLULEULYSG	LULEUGLNAR	GPROTHRVAL	720
THRTHRILEA	SNMETGLNME	TMETLYSALA	VALASNGLYA	SPLUULEEAS	NVALASPLGU	780
VALILESERL	YSASPSEERLE	UTRPGLNGLU	METARGTYRV	ALARGASPIL	ETHRLEUGLU	840
CYSLYSILEG	LUSERGLUVA	LILELYSGLY	GLYGLYTRPG	LYTYRASPTY	RTHRSEERILE	900
THRPHELYST	HRVALSNHNI	STHRTYRSE	TYRASNARGA	LAGLYGLUAS	NVALTHRALA	960
HISVALTHRV	ALSERPHEAS	NASNMTLYS	GLUARGSERT	YRGLYGLYSE	RLEUPROTHR	1020
ASPPHELYSI	LEASNARGPH	EASPVALMET	ASPILEASPT	HRTYRMETTY	RILEASPTYR	1080
TRPASPASPS	ERASPMETPH	ELYSASNMET	VALTYRVALA	RGASPLEUAR	GALAASPVVAL	1140
ASNGLYPHEP	HETYRTYRLY	SGLUMETSER	TYRSERPHEA	RGILEPROVA	LGLYVALHIS	1200
PROARGILEA	SNSERSERGL	YILEGLUPHE	THRTYRGLUL	YSCYSMETLE	USERGLNGLN	1260
PHETHRASPG	LNLEUALALE	UASNSEEMET	ARGPHEVALP	HEARGVALTH	RSERALAASN	1320
GLYTRPPHEM	ETTHRSEERGL	YGLYPROASN	THRARGARGI	LEALASERGL	YTHRGLYPHE	1380
ALATYRSERA	SPGLYHISTH	RSEERARGTHR	VALGLYTHRL	EUSERPHEIL	ESERLEUILE	1440
PROSERASNP	ROASNTYRGL	NTHRPROILE	ALASERSERS	ERTHRVALAR	GMETASPLEU	1500
ASPARGLYSI	LEASNASPLE	UARGASNASP	PHEASNGLNL	EUALASERSE	RVALALALEU	1560
GLYASPILEM	ETSERLEUAL	AMETSERPRO	LEUTHRPHE	ERASNLEULE	UGLUSERVAL	1620
PROALAILET	HRSERSERVA	LLYSASPVAL	ALAALAASNV	ALMETLYSLY	SPHEARGASN	1680
THRLYSMETP	HELYSLYSAL	ATHRLYSPRO	LYSTYRLYSG	LUPHEILEIL	EGLYASPLEU	1740
LEUGLUASPV	ALTHRASNVA	LTHRLYSASN	SERASNTHRM	ETASNPHAS	PASPILETHR	1800
SERALAVALM	ETILESERTH	RTHRASNLYS	LEUGLNLEUT	HRASPVALAS	PTHREUSER	1860
GLUILEVALV	ALARGSERAL	AASPASPPHE	ILEPROSERA	RGSERTYRAR	GMETILEGLU	1920
ASNASNTHRV	ALTYRGLUAL	ATHRPROSER	ARGVALPHES	ERTYRASPLE	UALATHRMET	1980
GLNGLNARGA	SPYHEASPI	EASPLYSPHE	ILEGLNLEUA	LASERLYSSE	RPROVALILE	2040
SERALAILEV	ALASPPHEAL	ATHRLEULYS	ALAMETARGG	LUTHRTYRGL	YTHRSERVAL	2100
GLUVALILIEH	ISLYSLUVA	LALASERASP	SERPROTHRI	LELEUSERPH	EILEASNASN	2160
ASNASNPROL	EUILEARGAS	NARGILEGLU	GLULEULEUA	RGGLNLCYSAR	GILELYSGLY	2220
GLUASNLEUT	YRPHEGLNGL	YHISHISHIS	HISHISHIS			2259

SEQ ID NO: 11 moltype = DNA length = 2414
 FEATURE Location/Qualifiers
 source 1..2414
 mol_type = other DNA
 organism = Rotavirus A

SEQUENCE: 11

accgagctcg	aattcatggc	aagccttatt	tacagacagc	tgctgacgaa	cagctacaca	60
atcgatctta	gtgacgaat	tgaagaaatc	ggatcgctca	aatcacagga	cgtcacaata	120
aaccttgccc	catttgctca	gacaggttat	gcgcctgtga	gctggggccc	tgccgagacg	180
aacgattcaa	cgaccatcga	gccggttctg	gatggcccat	atcgcccgac	cacctttaac	240
ccccgcatag	aatactggac	gctgtttgcy	cctaataaca	aaggagtcgt	tgccgaactt	300
acgaacaacg	cggatatctg	gttgccgcat	atccttatag	agccaaacgc	ttcaaggagg	360
attcgcactt	ataccatctt	cggtaacacg	gtcaatttga	cggtcgagaa	tgatccccaa	420
actaagtggg	aatttgctga	ttccgctcgc	cgctcgcaac	acgacactta	tgctcctgaa	480
ggaactcttc	ttagtgatag	taaattgcag	gccgccatga	aatatgggtg	taggctgttt	540
actttcatag	gtgatactcc	ttccgcagcg	cctcaggaat	acggttacga	gactaataac	600
tattcggcgc	ttgaggttaa	gtcattttgc	gacttctaca	tgattccccg	cttgccctaga	660
gaggtgtgca	gaaactatat	caaccaagga	cttcgcgcct	tacagaatac	ccgcaacggt	720
gtaccgcttg	cattgtcggc	tcgttcggtt	atagcgcagc	gtgtttcagt	taatgaggac	780
atagtgatat	cgaagacatc	cctctggaaa	gagatgcagt	acaataggga	tataaccatc	840
aggttcaagt	tcgctaacca	aataatcaaa	tcgggcggac	tggtttataa	atgggcgcag	900
atatcatttc	gtcctgcaaa	ctatcaatat	acctatattc	gtgacggcga	ggaggttaac	960

-continued

gcgcacacta	cctgttcggt	gaacggcgta	aacaatttct	cctacaacgg	cggttccctc	1020
ccaacggact	tcgtcatatc	gaggtatgag	gttctcaaag	agaactcggt	cgtttacatc	1080
gattactggg	atgatatgca	agcctttcgt	aacatgggtg	acgtcaggtc	actggcaacg	1140
gacctcaact	cgatcacatg	ctcgggaggg	agttataatt	ttgcgctccc	acttgccaac	1200
ttcccagtg	tgctctggcg	tgccgtgtct	ctccacccga	gcggtgtgac	gctcagcacg	1260
cagtttacag	actacgtgag	cctcaatagt	ctgaggttcc	gctttagact	ggcggttgaa	1320
gagccaccgt	tctccattac	gagaacaaga	gtttctcgct	tgtacggact	tccagctgtc	1380
aaccggaata	acaacaagga	tttttatgaa	atggcgggaa	gattctcatt	gatctctctt	1440
gtaccatcaa	atgacgatta	ccaaactccg	ataatgaata	gcgtcacagt	tagacaagat	1500
ttggaaggcg	agttggggaga	acttcgtgat	gaatttaacg	ccctctcaca	gcaaatcgca	1560
atgagccagt	tgatagatct	tgacttgctt	ccgctggaca	tggtctctat	gtttagcggg	1620
atcaagggca	cgatagacat	cgcaaaaagt	atggcgacaa	aggtcatgaa	gaagtccgt	1680
aagtccaatc	tcgcaaatag	ctattccacg	cttacggaat	cgttgtcaga	tgccgcgtcg	1740
agcatcagta	gaggtctccac	aatcccgagt	atcggtctta	gcgcttcggc	gtggaccgag	1800
gtaagtacaa	cgatagacga	cacaacaaat	gcaacgaaca	gcatagcaac	tcagacagca	1860
actatctcca	agaggtctcg	cttgaaagaa	atatcaactc	aaaccgacgg	aatgaatttc	1920
gatgatattc	ctgctgtctg	gctgaagact	aaaatagaca	agtcaactca	aatcgcgccc	1980
aacacactcg	ctgacatcgt	tacagaggcc	tccgaaaagt	ttataccaaa	tagggcttat	2040
cgcgtcatcg	acaatgacga	ggtattcgaa	gcgggcacgg	acggacgctt	ttcgctctac	2100
agagtcgaaa	acttcgagga	gataccggtc	gatgtacaga	agttcgcgga	ccttgtagcc	2160
gattcaccag	ttatttcagc	aataatagac	tttaaaactc	tgaagaacct	caacgataat	2220
tatggcatca	ccaagcaaca	ggcgtacaac	cttcttcgta	gcgacccgag	ggctcctcagg	2280
gagttcataa	accaggaaaa	cccgataatc	cgcaaccgta	tcgaatcttt	gattatgcaa	2340
tgagggttga	agggcgaaaa	ctgtgacttt	caaggccatc	accatcacca	tcactaggcg	2400
gccgcttaat	taat					2414

SEQ ID NO: 12 moltype = AA length = 2379
FEATURE Location/Qualifiers
source 1..2379
 mol_type = protein
 organism = Rotavirus A

SEQUENCE: 12

METALASERL	EUILETYRAR	GGLNLEULEU	THRASNSSERT	YRTHRELEAS	PLEUSERASP	60
GLUILEGLUG	LUILEGLYSE	RLEULYSSER	GLNASPVALT	HRILEASNPR	OGLYPROPHE	120
ALAGLNTHRG	LYTYRALAPR	OVALSERTRP	GLYPROGLYG	LUTHRASNAS	PSERTHRTHR	180
ILEGLUPROV	ALLUASPLG	YPROTYRARG	PROTHRTHRP	HEASNPROPR	OILEGLUTYR	240
TRPTHRLEUP	HEALAPROAS	NASNLYSGLY	VALVALALAG	LULEUTHRAS	NASNLAASP	300
ILETRPLEUA	LAILEILELE	UILEGLUPRO	ASNALASERA	RGGLUILEAR	GTHRTYRTHR	360
ILEPHEGLYG	LNGLNVALAS	NLEUTHRVAL	GLUASNVALS	ERGLNTHRLY	STRPLYSPE	420
VALASPPHEA	RGARGARGSE	RGLNHHISASP	THRTYRVALL	EUGLUGLYTH	RLEULEUSER	480
ASPTHRLYSL	EUGLNALAL	AMETLYSTYR	GLYALAARGL	EUPHETHRPH	EILEGLYASP	540
THRPROSERA	LAALAPROGL	NGLUTYRGLY	TYRGLUTHRA	SNASNTYRSE	RALAVALGLU	600
VALLYSSERP	HECYSASPPH	ETYRMETILE	PROARGLEUP	ROARGGLUVA	LCYSARGASN	660
TYRILEASNG	LNGLYLEUPR	OPROILEGLN	ASNTHRARGA	SNVALVALPR	OVALALALEU	720
SERALAARGS	ERVALILEAL	AGLNARGVAL	SERVALASNG	LUASPILEVA	LILESERLYS	780
THRSEERLEUT	RPLYSGULVE	TGLNTYRASN	ARGASPILET	HRILEARGPH	ELYSPEAL	840
ASNGLNILEI	LELYSSERGL	YGLYLEUGLY	TYRLYSTRPA	LAGLUILESE	RPHEARGPRO	900
ALASANTYRG	LNTYRTHRTY	RILEARGASP	GLYGLUGLUV	ALTHRALAHI	STHRTHRCYS	960
SERVALASNG	LYVALASNA	NPHESERTYR	ASNGLYGLYS	ERLEUPROTH	RASPPHEVAL	1020
ILESERARGT	YRGLUVALLE	ULYSGLUASN	SERPHEVALT	YRILEASPTY	RTRPASPASP	1080
SERGLNALAP	HEARGASNME	TVALTYRVAL	ARGSERLEUA	LATHRASPLE	UASNSEIRILE	1140
THRCYSSERG	LYGLYSERTY	RASNPHLEAL	LEUPROLEUG	LYASNPHEP	OVALMETSER	1200
GLYGLYALAV	ALSERLEUHI	SROSERGLY	VALTHRLEUS	ERTHRGLNPH	ETHRASPTYR	1260
VALSERLEUA	SNSEIRLEUA	GPHEARGPHE	ARGLEUALAV	ALGLUGLUP	OPROPHESES	1320
ILETHRARGT	HRARGVASE	RARGLEUTYR	GLYLEUPROA	LAVALASNPR	OASNASNASN	1380
LYSASPPHET	YRGLUMETAL	AGLYARGPHE	SERLEUILES	ERLEUVALPR	OSERASNASP	1440
ASPTYRGLNT	HRPROILEME	TASNSEIRVAL	THRVALARGG	LNASPLEUGL	UARGGLNLEU	1500
GLYGLULEUA	RGASPLGUPH	EASNALALEU	SERGLNGLNI	LEALAMETSE	RGLNLEUILE	1560
ASPLEUALAL	EULEUPROLE	UASPMETPHE	SERMETPHES	ERGLYILELY	SGLYTHRILE	1620
ASPILEALAL	YSSERMETAL	ATHRLYSVAL	METLYSLYSP	HEARGLYSSE	RASNLEUALA	1680
ASNSEIRILES	ERTHRLUTH	RGLUSERLEU	SERASPALAA	LASERSERIL	ESERARGGLY	1740
SERTHRIELA	RGSERILEGL	YSERSEIRALA	SERLATRPT	HRGLUVALSE	RTHRTHRIEL	1800
ALAASPTHRT	HRASNALATH	RASNSEIRILE	ALATHRGLNT	HRALATHRIL	ESERLYSARG	1860
LEUARGLEUL	YSGULILESE	RTHRGLNTHR	ASPGLYMETA	SNPHEASPAS	PILESERALA	1920
ALAVALLEUL	YSTRHLYSIL	EASPLYSSER	THRGLNILEA	LAPROASNT	RLEUPROASP	1980
ILEVALTHRG	LUALASERGL	ULYSPEHILE	PROASNARGA	LATYRARGVA	LILEASPASN	2040
ASPLUVALP	HEGLUALAGL	YTHRASPLGY	ARGPHEPHEA	LATYRARGVA	LGLUTHRPHE	2100
GLUGLUILEP	ROPHEASPPA	LGLNLYSPHE	ALAASPLEUV	ALTHRASPS	RPROVALILE	2160
SERALAILEI	LEASPPHELY	STHRLLEULYS	ASNLEUASNA	SPASNTYRGL	YILETHRLYS	2220
GLNGLNALAT	YRASNLEULE	UARGSERASP	PROARGVALL	EUARGGLUPH	EILEASNGLN	2280
GLUASNPROI	LEILEARGAS	NARGILEGLU	SERLEUILEM	ETGLNCSYAR	GLEULYSGLY	2340
GLUASNLEUT	YRPHEGLNGL	YHISHISHIS	HISHISHIS			2379

SEQ ID NO: 13 moltype = DNA length = 2279
FEATURE Location/Qualifiers
source 1..2279

-continued

```

mol_type = other DNA
organism = Rotavirus C

SEQUENCE: 13
accgagctcg aattcatggc tagcagcctt taccatcagc tcattttctca gaattattac 60
agtcacagaa accgagtcct taaagatttg caaactacta aaacaacagt cgactacgta 120
gacgctggca actacacata ttctcagctt ccgcccacca aatgggggtgc tggcgccact 180
tttgaatctt ttttttcggc cgccgaaatc accggccctc ataccaatag ggttatcgaa 240
tggaagaacc ttcttaatag caatcagtggt ctcctttttcc ctaaaccaac agacactgtc 300
aagctcttga agcatggtcc tcagacgtat gacaatactt tggctgcatg cgagctttgg 360
tatggaaagg ctaatactac agtgacgtct gaacattact cgtctctgtc agactcttgg 420
gtcaacgtta acagcgattc actcgctact ttctggaaacg ctggcgccac cacattcaac 480
aaaaaggtcg tcaatttcgc ctggaacatg ggaggtatca tgataaaacc ttcatcgcaa 540
cttctcgtc tcgaaattta tatggccaac atgaacaatt tcaatggcga taacttcaat 600
tgggaggagt ggcatttcac ccttcctagg tctaagcaa ctatcaatat atatattgac 660
tattacctgg cttccagtgga ccggtataat cagcttaagg agctccaaca gcccacgatt 720
actacgttcg agatgcgcac gatgcttctg aatagagatg gtcaaatcaa cgtaaatgaa 780
gtagtcagca aagatttcac ctggcagga atgcgctacg cgctgacat cacactccaa 840
tgcaagatcg agtcgaggt ggtgaaaggt ggcggttggg gttatgacta tacatctatc 900
aggttcaaga ctgttaacca tacgtacagt tacactagag cgggcgaaaa cgtaaatgca 960
catgtcacga tatcgttcaa taacttgaaa gagagaagct acggaggttc gcttccgacg 1020
gatttcaaga taggtaagtt cgatataata gatgtagata cgatatatga cattgactac 1080
gggagtgatt ccgagatcct caagaacatg gtatacgtaa gagatctcac cgcaaatatt 1140
ggtagcttct tccattatag agagtagagc tattatttcc aaataccggt cggagttcat 1200
ctgtgtatcc acagcagtggt gtgcaggttc atgtacgaaa ggtgcctttt gtctcagcaa 1260
tttacggatc aagttgcttt gaactcgatg aggttcattt tcaaaagtaac ggcggctaac 1320
ggctgggttc tgacgagcgg aaatattaat actcgtagga tagcgagcgg cacaggcttt 1380
gcttacgcca atggttacgt gtcacagacc gtgggcaaca tcagcttcat tagccttatt 1440
ccctcgaaac ctaattatca gacacccatc gctccagct ccacgctccg tatggacttg 1500
gagcgttaaga ttaatgatct taggaatgac tttaaaccaac tggcatcttc agtggtcttc 1560
ggagatatcc tgtcgcttgc cactcgccg ctcacatttg ctaacctgtt ggagtcagtg 1620
cctgcagtag ctgactccgt gaaggatgtg gcggcgaatg taatgaaaag gttcagaaat 1680
actaaaatgt ttaaacgtgc cgcaaaaccc aaatataaag agttcatcat aggtgatctt 1740
cttgaggatg taaccaatgt gacccgtaat cccaactcta tgaatttcga tgatatacag 1800
agtgcocatg tagtttccac tacaaacaaa cttcaattga cggatgtgga caccctttct 1860
gagatcgttg ccaggctctg tgatgacttc attccgaata ggtcctaccg tatgatcgaa 1920
aacggagttg tccatgaagc aacacaaaat ggtgtttttt cgtatgatct tgtaactctg 1980
cagcaacggt cttcgacatg ggataaattt cttcaacttg ctagaagtc acctgttatt 2040
tcggccatag ttgatttcgc gacccctaag gcaatgaggg acacgtatgg catcggtaca 2100
gatgtacttt acaaatgtgt tgtaagtgat acgccaaccc ttctgtcatt tattaataac 2160
aacaatccac tcattccgaa taggatcgaa ggactgtcga ggcaatgtag aattaagggc 2220
gaaaacttgt actttcaagg ccatcaccat caccatcact aggcggcgcc ttaattaat 2279

```

```

SEQ ID NO: 14      moltype = AA length = 2244
FEATURE            Location/Qualifiers
source              1..2244
                    mol_type = protein
                    organism = Rotavirus C

```

```

SEQUENCE: 14
METALASERS ERLEUTYRHI SGLNLEUILE SERGLNASNT YRTYRSERTH RGLYASNGLU 60
ILELEULYSA SPLEUGLNTH RTHRLYSTHR THRVALASPT YRVALASPAL AGLYASNTYR 120
THRTYRSERG LNLUPROPR OTHRLYSTRP GLYALAGLYA LATHRPEHGL USERPHEPHE 180
SERALAALAG LUILETHRLG YPROHISTHR ASNARGVALI LEGLUTRPLY SASNLEULEU 240
ASNSEASNSG LNTRPLEULE UPHEPROLYS PROTHRASPT HRVALLYSLE ULEULYSHIS 300
GLYPROGLNT HRTYRASPAS NTHRLEUALA ALACYSGLUL EUTRPTYRGL YLYSALAASN 360
THRTHRVALT HRSERGUHI STYRSERSE LEUSERASPS ERTRPVVALS NVALASNSER 420
ASPSERLEUV ALLEUPHETR PASNALAGLY GLYTHRTRHP HEASNLYSLY SVALVALASN 480
PHEALATRPA SNMETGLYGL YILEMETILE LYSPROSERS ERGLNLEUPR OARGLEUGLU 540
ILETYRMETA LAASNMETAS NASNPHEASN GLYASPASNP HEASNTRPGL UGLUTRPHIS 600
PHETHRLEUP ROARGSERAS NALATHRILE ASNILETYRI LEASPTYRTY RLEUALASER 660
SERASPPROT YRASNGLNLE ULYSGLULEU GLNGLNPROT HRILETHRTH RPHEGLUMET 720
ARGMETMETL EULEUASNRL GASPGLYGLN ILEASNVALA SNGLUVALVA LSERLYSASP 780
SERLEUTRPG LNLUMETAR GTYRALAARG ASPILETHRL EUGLNCYSLY SILEGLUSER 840
GLUVALVALL YSGLYGLYGL YTRPGLYTYR ASPTYRTHRS ERILEARGPV ELYSTHRVAL 900
ASNHISTHRT YRSERTYRTH RARGALAGLY GLUASNVALA SNALAHISVA LTHRILESER 960
PHEASNASNL EULYSGLUAR GSERTYRGLY GLYSERLEUP ROTHRASPPH ELYSILEGLY 1020
LYSPHEASPI LEILEASPVA LASPTHRTYR METTYRILEA SPTYRTRPAS PASPSERGLU 1080
ILEPHELYSA SNMETVALTY RVALARGASP LEUTHRALAA SNILEGLYAS PPHEPHEHIS 1140
TYRARGGLUM ETSERTYRTY RPHEGLNILE PROVALGLYV ALHISPROGL YILEHISSE 1200
SERGLYVALA RGPHEMETTY RGLUARGCYS LEULEUSERG LNGLNPHETH RASPGLNVAL 1260
ALALEUASNS ERMETARGPH EILEPHELYS VALTHRALAA LAASNGLYTR PPHEMETTHR 1320
SERGLYASNI LEASNTHRAR GARGILEALA SERGLYTHRG LYPHEALATY RALAASNGLY 1380
TYRVALSERG LNTHRVALGL YASNILESER PHEILESERL EUILEPROSE RASNPROASN 1440
TYRGLNTHRP ROILEALASE RSERSERTHR VALARGMETA SPLEUGLUAR GLYSILEASN 1500
ASPLEUARGA SNASPPHEAS NGLNLEUALA SERSERVALA LALEUGLYAS PILELEUSER 1560
LEUALATHRS ERPROLEUTH RPHEALAASN LEULEUGLUS ERVALPROAL AVALALASER 1620
SERVALLYSA SPVALALAAL AASNVALMET LYSARGPHEA RGASNTHRLY SMETPHELYS 1680

```

-continued

ARGALALAL	YSPROLYSTY	RLYSLGLUPHE	ILEILEGLYA	SPLEULEUGL	UASPVALTHR	1740
ASNVALTHRA	RGASNPROAS	NSERMETASN	PHEASPASPI	LETHRSEAL	AMETMETILE	1800
SERTHRTHRA	SNLYSLEUGL	NLEUTHRASP	VALASPTHRL	EUSERGLUIL	EVALALAARG	1860
SERALAASPA	SPPHEILEPR	OASNARGSER	TYRARGMETI	LEGLUASNGL	YVALVALHIS	1920
GLUALATHRG	LNASNGLYVA	LPHESERTYR	ASPLEUVALT	HRLEUGLNGL	NARGSERPHE	1980
ASPMETASPL	YSPHELEUGL	NLEUALASER	LYSSERPROV	ALILESERAL	AILEVALASP	2040
PHEALATHRL	BULYSALAME	TARGASPTHR	TYRGLYLEG	LYTHRASPVA	LLEUTYRLYS	2100
LEUVALVALS	ERASPTHPR	OTHRLEULEU	SERPHEILEA	SNASNASNAS	NPROLEUILE	2160
ARGASNARGI	LEGLUGLYLE	ULEUARGGLN	CYSARGILEL	YSGLYGLUAS	NLEUTYRPHE	2220
GLNGLYHISH	ISHISHISHI	SHIS				2244

SEQ ID NO: 15 moltype = DNA length = 2276
 FEATURE Location/Qualifiers
 source 1..2276
 mol_type = other DNA
 organism = Rotavirus C

SEQUENCE: 15

accgagctcg	aattcatggc	aagttcattg	tataagcaac	tgataagcca	aaactattat	60
agtacgggca	acgacatcct	gctcgatcgc	cagacgaacg	agacaactgt	cgattacgtt	120
gatatttgta	attactcgta	cacccaactg	ccccctacta	gttgggtg	gggcatgacg	180
ttcaagtcgg	ccttcaacgc	cgaagagatt	accggtccta	acactgggtga	tatagatttg	240
aataacctca	caaacgccc	cggttggatt	ctctatgaca	aacctacctc	aacgaagcgc	300
ttgcttaaac	tgggtccga	cggttacgat	aacgtatatg	cggttttcga	gctctggtat	360
ggaaaagcta	ataccatcgt	agagtcctac	tattatgcct	ctgcgcaaaa	ctccgagaaa	420
accgctacgg	ccagatga	ctctctcgta	ctgttcttta	atgttggata	tacaggtcct	480
actaaacaga	tcgtgaaatt	caactggaac	atggggaggaa	ttttgggttag	gccgacagca	540
gacggccgtg	tggatgtctg	ctggcgggac	atgaatgact	ttaactcaga	caatttcaat	600
tggaagcct	ggaagcgctc	ttcccgcgt	agtaatatca	atatgtatac	agaatactat	660
ctcgcaaatg	ttgatccgta	taaccagctg	aaggctctca	atcagctcac	cgctaaaaat	720
atagagctta	gaatgatgaa	agcaacaaag	caacagggac	caattaatgt	ggatgaggtc	780
gtttccaaag	attcattgtg	gcaggagatg	aggtatgcta	gagatatcac	acttaagtgt	840
aaagtcgaat	cgacatcat	taaaggcgcc	ggttggggct	atgactatac	aagcatcacg	900
ttcaaaactg	tgaactacgt	gtacaataac	accogttctg	gagaggaggt	caatgccac	960
gtgacaattt	cattcaacaa	catgaagaa	cgcagctatg	gcggttcact	tccactgac	1020
ttcaagatca	agagattcga	cggttattgat	aacgatacgt	atatgtatat	cgattattgg	1080
gacgactcag	atatatttaa	gaatatgggt	tacgtacgcg	acctgagggc	aaacatgggt	1140
gactttttct	attatagtga	gatgtggcac	agatttccaa	ttccagtagg	aaccaccca	1200
ggcctgcatt	ccagcgcgct	ccaatttaca	tacgaaaaat	gtttgttgag	ccaacagttc	1260
acagatatgg	ttcggtgaa	ctcgatgcgc	ttcattttcc	gcgttacctc	gtccaacggt	1320
tggtatatca	cttcgggag	tcgaataacg	cgcaaatag	cttcgggaac	cggatcgcg	1380
tactcaaacg	gctacataag	tgagactgtc	ggaaacatta	gctttatctc	attgattcct	1440
tctaattccta	attatcagac	accgattgca	tcgtctcga	cagtaaggat	ggatcttgac	1500
cgtaaaatta	acgactccg	caatgacttt	aacaatctgg	ctagctctgt	cgctctggga	1560
gacataatgt	cattggcaat	gtcacccctg	acatttgcga	acctgcttga	gtcggtccca	1620
cgcataaact	gtagcgctca	agatgtcgca	gccaaactga	tgaagaagt	taggaatact	1680
aaagatgttca	agcgtgccac	aaagacgaag	tacaaggaat	acatcatcgg	cgacctctcg	1740
gaagacgtga	caaatgtcac	gaggggatga	aacaagatga	atttcgacga	tatcacttcg	1800
gcgatgatgg	tgtcagcgac	caataaactg	caactgactg	atattgatac	atttagtgaa	1860
attgtagtga	ggtcggaaga	cgactttata	ccaaacaggt	cttacaggat	gatcgagaat	1920
ggtatcgtac	atgaggtcac	cccaactaga	gtcttctctt	atgaccttaa	aacctcgcaa	1980
caaacccgact	tcgatattgga	caaattcatg	caattggcct	ccgagtcctc	ggtcataatc	2040
gcaataatag	actttgcaac	gctgaaggcc	atgcgtgaca	cttatgggat	atcgactgat	2100
ataatgtaca	aacttatagc	aagcgactcg	cctactattc	tgtcttttat	caacaacaat	2160
aacctctcta	taagaaatag	aatcgagggt	ttgcttcgtc	aatgcagaat	taagggcgaa	2220
aacttgact	ttcaaggcca	tcaccatcac	catcactagg	cggcgcgcta	attaat	2276

SEQ ID NO: 16 moltype = AA length = 2241
 FEATURE Location/Qualifiers
 source 1..2241
 mol_type = protein
 organism = Rotavirus C

SEQUENCE: 16

METALASERS	ERLEUTYRLY	SGLNLEUILE	SERGLNASNT	YRTYRSERTH	RGLYASNASP	60
ILELEULEUA	SPARGGLNTH	RASNGLUTHR	THRVALASPT	YRVALASPI	EGLYASNTYR	120
SERTYRTHRG	LNLEUPROPR	OTHRSETRP	GLYALAGLYM	ETTHRPHEL	SSERALAPHE	180
ASNALAGLUG	LUILETHRGL	YPROASNTHR	GLYASPILEA	SPLEUASNAS	NLEUTHRASN	240
ALAASNGLYT	RPILELEUTY	RASPLYSPRO	THRSERTHRL	YSARGLEULE	ULYSLEUGLY	300
PROASPART	YRASPAENVA	LTYRALAALA	PHEGLULEUT	RPTYRGLYLY	SALAASNTHR	360
ILEVALGLUS	ERILETYRTY	RALASERALA	GLNASNSERG	LULYSTHRL	ATHRVALGLN	420
TYRASPSERL	EUVALLEUPH	EPHEASNVAL	GLTYRTHRG	LYLEUTHRLY	SGLNILEVAL	480
LYSPHEASNT	RPASNMTGL	YGLYILELEU	VALARGPROT	HRALAASPL	YARGVALASP	540
VALCYSMETA	LAASPMETAS	NASPPHEASN	SERASPASNP	HEASNTRPGL	UALATRPLYS	600
ARGSERPHEP	ROARGSERAS	NILEASNMT	TYRTHRGLUT	YRTYRLEUAL	AASNVALASP	660
PROTYRASNG	LNLEULYSAL	NLEUASNGLN	LEUTHRALAL	YSASNILEGL	ULEUARGMET	720
METLYSALAT	HRLYSGNLGL	NGLYPROILE	ASNVALASPG	LUVALVALSE	RLYSASPSER	780
LEUTRPGLNG	LUMETARGTY	RALAARGASP	ILETHRLEUL	YSCSYLSYL	EGLUSERASP	840

-continued

ILEILELYSG	LYGLYGLYTR	PGLYTYRASP	TYRTHRSERI	LETHRPHELY	STHRVALASN	900
TYRVALTYRL	YSTYRTHRAR	GSERGYGLU	GLUVALASNA	LAHISVALTH	RILESERPHE	960
ASNASNMTL	YSGLUARGSE	RTYRGYGLY	SERLEUPROT	HRASPPHELY	SILELYSARG	1020
PHEASPVALI	LEASPASNAS	PTHRTYRMET	TYRILEASPT	YRTRPASPAS	PSERASPILE	1080
PHELYSASN	ETVALTYRVA	LARGASPLEU	ARGALAASN	ETGLYASPPH	EPHETYRTYR	1140
SERGLUMETT	RPHISARGPH	EPROILEPRO	VALGLYTHRH	ISPROGLYLE	UHISSERSE	1200
GLYVALGLNP	HEHTRTYRGL	ULYSCYSLEU	LEUSERGLNG	LNPHEHTRAS	PMETVALALA	1260
LEUASNSERM	ETARGPHEIL	EPHEARGVAL	THRSERSERA	SNGLYTRPTY	RILETHRSE	1320
GLYGLYPROA	SNTHRARGLY	SVALALASER	GLYTHRGLYP	HEALATYRSE	RASNGLYTYR	1380
ILESERGLUT	HRARGVLYPH	NILESERPHE	ILESERLEUI	LEPROSERAS	NPROASNTYR	1440
GLNTHRPROI	LEALASERSE	RSETRHRVAL	ARGMETASPL	EUASPARGLY	SILEASNASP	1500
LEUARGASNA	SPPEASNAS	NLEUALASER	SERVALALAL	EUGLYASPII	EMETSERLEU	1560
ALAMETSERP	ROLEUTHRPH	EALAASNLEU	LEUGLUSERV	ALPROALAIL	ETHRSESRSE	1620
VALLYSASPV	ALALAALAAS	NVALMETLYS	LYSPHEARGA	SNTHRLYSME	TPHELYSARG	1680
ALATHRLYST	HRLYSTYRLY	SGLUTYRILE	ILEGLYASPL	EULEUGLUAS	PVALTHRASN	1740
VALTHRARGA	SPVALASNPH	SMETASNPH	ASPASPILET	HRSERALAME	TMETVALSER	1800
ALATHRASNL	YSLEUGLNLE	UTHRASPILE	ASPTHRPHE	ERGLUILEVA	LVALARGSER	1860
ALAASPASPP	HEILEPROAS	NARGSERTYR	ARGMETILEG	LUASNGLYIL	EVALHISGLU	1920
ALATHRPROT	HRARGVLYPH	ESERTYRASP	LEULYSTHRL	EUGLNLNTH	RASPPHEAS	1980
METASPLYSP	HEMETGLNLE	UALASERGLU	SERPROVALI	LESERALAIL	EILEASPPHE	2040
ALATHRLEUL	YSALAMETAR	GASPTHRTYR	GLYILESERT	HRASPILEME	TTYRLYSLEU	2100
ILEALASERA	SPSERPROTH	RILELEUSER	PHEILEASNA	SNASNANPR	OLEUILEARG	2160
ASNARGILEG	LUGLYLEULE	UARGGLNCYS	ARGILELYSG	LYGLUASNLE	UTYRPHEGLN	2220
GLYHISHISH	ISHISHISHI	S				2241

SEQ ID NO: 17 moltype = DNA length = 1088
 FEATURE Location/Qualifiers
 source 1..1088
 mol_type = other DNA
 organism = Rotavirus C

SEQUENCE: 17

accgagctcg	aattcatggt	atgtacaacc	ctctacacaa	tttgtgcaat	attgtgcatt	60
ctccttatct	atacactggt	tctcagaaag	atgttccatt	tgcttactga	cgcccttgta	120
gttattctta	tcatatctgc	gtgtgcccga	agcagtaagt	gccagctctt	taccaatgac	180
atcagttacg	acggcaacat	tgaatctgtg	atcaatacaa	ctaaccggtt	taatgttgag	240
agccctttga	tatactttcc	aaacgcccga	ataggttcac	caggaccggg	taaacgagat	300
ggacttctca	acgataataa	ctacgcccag	acacttgcaa	cacttttcga	aacaaaggga	360
tttcctaagg	gatcagtgat	attcaagacc	tataccaaaa	cgagcgactt	cattaactct	420
gttgagatga	cgtgcagtta	taatttggtt	ataattccag	atgagccaaa	aaacagtgaa	480
gcaatagagc	agatagccga	atggattttg	aatgtctggc	aatgcgatga	catgaatctt	540
gacatctata	catcagcaaa	aatcaataaa	gataaacctt	gggcagcggt	tggtgaagat	600
tgtgatattt	cggtatgtcc	cgttgatact	acaatgaatg	gaatcggtcg	ctctccggct	660
agtagcggaa	cgtatgaagt	cggtgcgaac	gataccacgc	ttgctttgat	agatgtagtt	720
gataatgtca	agtatagaat	acaaatgtcc	agtagactga	atcagcttca	atgtaagctg	780
aagaattcgc	ttaaaggcga	accaagggtg	aatactgcaa	ttatccgtat	tagtagctcc	840
tccttctttg	ataattccct	ttctccgctg	tcaaacggcc	aaaccacgag	agcgttcaag	900
atcaatgcta	aaaagtgggt	gaaaaatttc	tacactataa	tcgattacat	caaaccttgc	960
gttcagacga	tgaccccaag	gcacgtgtgc	gtctatccct	aggggttgat	gcttcgctac	1020
gctaagggag	aaaacttgta	ctttcaaggc	catcaccatc	accatcacta	ggcggccgct	1080
taattaat						1088

SEQ ID NO: 18 moltype = AA length = 1053
 FEATURE Location/Qualifiers
 source 1..1053
 mol_type = protein
 organism = Rotavirus C

SEQUENCE: 18

METVALCYST	HRTHRLEUTY	RTHRILECYS	ALAILELEUC	YSILELEULE	UILEYRTHR	60
LEUPHELEUA	RGLYSMETPH	EHISLEULEU	THRASPALAL	EUVALVALIL	ELEUILEILE	120
SERALACYSA	LAGLYSERSE	RASNGLYGLN	LEUPHETHRA	SNASPILESE	RTYRASPGLY	180
ASNILEGLUS	ERVALILEAS	NTHRTHRASN	PROPHEASNV	ALGLUSERLE	UCYSILETYR	240
PHEPROASNA	LAALAILEGL	YSERPROGLY	PROGLYLYSA	LAASPGLYLE	ULEUASNASP	300
ASNASNTRYA	LAGLNTHRLE	UALATHRLEU	PHEGLUTHRL	YSGLYPHEPR	OLYSGLYSER	360
VALILEPHEL	YSTHRTYRTH	RLYSTRHSE	ASPPHEILEA	SNSERVALGL	UMETTHRCYS	420
SERTYRASN	EUVALILEIL	EPROASPLU	PROLYSASNS	ERGLUALAIL	EGLUGLNILE	480
ALAGLUTRPI	LELEUASNVA	LTRPGLNCYS	ASPASPMETA	SNLEUASPII	ETYRTHRTYR	540
GLUGLNILEA	SNLYSASPAS	NLEUTRPALA	ALAPHEGLYG	LUASPCYSAS	PILESERVAL	600
CYSPROLEUA	SPTHRTHRME	TASNGLYILE	GLYCYSSERP	ROALASERTH	RGLUTHRTYR	660
GLUVALVALS	ERASNASPTH	RGLNLEUALA	LEUILEASPV	ALVALASPAS	NVALLYSTYR	720
ARGILEGLNM	ETSESRERTH	RLEUASNGLN	LEUGLNCYSL	YSLEULYSAS	NCYSVALLYS	780
GLYGLUPROA	RGLEUASNTH	RALAILEILE	ARGILESERT	HRSESRERSE	RPHEASPASN	840
SERLEUSERP	ROLEUSERAS	NGLYGLNTHR	THRARGALAP	HELYSILEAS	NALALYSLYS	900
TRPTRPLYSI	LEPHETYRTH	RILEILEASP	TYRILEASNT	HRLEUVALGL	NTHRMETTHR	960
PROARGHISA	RGALAVALTY	RPROGLUGLY	TRPMETLEUA	RGTYRALALY	SGLYGLUASN	1020
LEUTYRPHEG	LNGLYHISHI	SHISHISHIS	HIS			1053

-continued

SEQ ID NO: 19	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 19		
gtctgcgagc agttgttt		18
SEQ ID NO: 20	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 20		
cgctctaaca taccacccta aa		22
SEQ ID NO: 21	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 21		
ggatcgggtgg tagaccatta ttt		23
SEQ ID NO: 22	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 22		
caacgccact accggttatt a		21
SEQ ID NO: 23	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 23		
cgaagttcgt tgagttgtct ttc		23
SEQ ID NO: 24	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 24		
cctgctgccca accctaataa		20
SEQ ID NO: 25	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 25		
cgtggtatgc aaggtagtat t		21
SEQ ID NO: 26	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 26		
gtaaggcggg tcatactgtt ac		22
SEQ ID NO: 27	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 27		
aggagcggttc cttcatgtta tt		22
SEQ ID NO: 28	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	

-continued

SEQUENCE: 28	organism = Synthetic construct	
tggaagaga tgcgtatg		19
SEQ ID NO: 29	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 29		
tggtcacatt cgttacatcc tc		22
SEQ ID NO: 30	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
source	1..23	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 30		
catctcgctc atcccttcta atc		23
SEQ ID NO: 31	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 31		
cgttgcgggt attctgtata gg		22
SEQ ID NO: 32	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 32		
atggtgctag gctgtttact t		21
SEQ ID NO: 33	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 33		
atgtctatcg tgccottgat ac		22
SEQ ID NO: 34	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 34		
ccagctgtca acccgaataa		20
SEQ ID NO: 35	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 35		
gtcactggaa gccaggtaat ag		22
SEQ ID NO: 36	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
source	1..22	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 36		
cgattcactc gtactcttct gg		22
SEQ ID NO: 37	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
source	1..20	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 37		
ggcgatgggt gtctgataat		20

-continued

SEQ ID NO: 38	moltype = DNA	length = 22	
FEATURE	Location/Qualifiers		
source	1..22		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 38			
tgtcagggttc atgtacgaaa gg			22
SEQ ID NO: 39	moltype = DNA	length = 21	
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 39			
agggtcgcgga aatcaactat g			21
SEQ ID NO: 40	moltype = DNA	length = 22	
FEATURE	Location/Qualifiers		
source	1..22		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 40			
gaccgcgtaat cccaactcta tg			22
SEQ ID NO: 41	moltype = DNA	length = 22	
FEATURE	Location/Qualifiers		
source	1..22		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 41			
gagagccttc agctgggttat ac			22
SEQ ID NO: 42	moltype = DNA	length = 22	
FEATURE	Location/Qualifiers		
source	1..22		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 42			
ggcggacatg aatgacttta ac			22
SEQ ID NO: 43	moltype = DNA	length = 21	
FEATURE	Location/Qualifiers		
source	1..21		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 43			
gcgacatctt tgacgctact a			21
SEQ ID NO: 44	moltype = DNA	length = 23	
FEATURE	Location/Qualifiers		
source	1..23		
	mol_type = other DNA		
	organism = Synthetic construct		
SEQUENCE: 44			
agtgagactg tcggaacat tag			23
SEQ ID NO: 45	moltype = DNA	length = 2407	
FEATURE	Location/Qualifiers		
source	1..2407		
	mol_type = other DNA		
	organism = Rotavirus A		
SEQUENCE: 45			
accgagctcg aattcatggc ttcgcttata tacaggcaac tgctcagcaa ttcgtatacc			60
gttgatctgt ctgacgagat tcaaattatc ggctcggaga agacacaaaa cgtcaccata			120
aaccccgggac cgtttcgcgca gcacaggttat gcacctgtca attgggggacc ggggtgagaca			180
aacgactcaa caactgtcga gccagctcttg gacggcccggt atcaaccgac gacctttaat			240
ccaccggttt cgtattggat tcttctttcg cctagtaagt ccggtgtagt cgtcgaaggc			300
actaataact cagaccgttg gctcgcaact atattgatag aaccggaacgt aatttccgaa			360
actcgcacat acacattggt tggcagacag gagcaaatga cggtggaaaa tgtatcaaca			420
accaagtggga aattcgtcga ccttgccaaa accgatatca atggaaacgtt caccgcagcat			480
ggtcgcgttt tgtcggcacac caaattgtac ggagtaatga aatttagtggt tcgtctgtac			540
acttataatg gagaaacacc caatgctacc acaggatatt atactaccac caattatgac			600
accgtgaaca tgaccagcta ttgtgacctc tatatcatac cgcgcctctga agagtgcagt			660
tgtactagtt atattaataa tggcctgccc ccaattcaga atacgcgtga tgttgttcca			720
gtgagtttga cgtctcgcct catagtcocat actagggcgc aagcaaatga agatattgtg			780
gtcagcaaaa catctctgtg qaataaqaatg caatacaata qqatataaac qatacgtttc			840

-continued

```

aaatttgcga acgtataaat aaaaagcggg ggactggggt acaagtggag cgagatttct 900
ttcaagccgg ccaactatca gtacaattat acgagagacg gtgagggtcat cactgcccat 960
actacctgta gtgtcaatgg tattaatgat tttagttaca atggagggtc cttgccgacc 1020
gatttttgta tcagcaggta tgaagtaata aaagaaaata gttatgtgta tgtggactat 1080
tgggatgact cacaggcttt cagaaacatg gtctatgtcc gtagtttggc tgccaatctc 1140
aattctgtta catgcacagg tggcaattat gatttttagac tgcccgtggg tgccctggcct 1200
gttatgactg gaggcgctgt cctactgcgt ccggccggcg ttacgctctc cacgcagttt 1260
acggattttg taagcttgaa ctccttgaga tttagggtca gcttgcgggt ggaagagcct 1320
ccttttacaa ttacacgcac taggggtgtcc ggactgtatg gtctcccggc ggcaaatcct 1380
aacaacggaa aggattacta cgagataact ggtcgcttta gcctcatcag cctcgtgccg 1440
agcaacgacg attatcgac gcctatcatg aactctgtga ccgtgcgtca agaccttgaa 1500
cgccagctta acgaattgcg ccaagaattt aattcattga gtcaagaagt tgctatgagc 1560
cagcttatcg acctggcgct gcttctcttc gatatgttct ctatgttcag cggataaaag 1620
tcgaccatcg acgcccgcga atctatggcc acttcagtta tgaagaagt taaacgctcg 1680
aacctcgcat cttctattag caccctcaca gattctctct cagacgctgc atccagtatt 1740
tcgaggaatc cgagtataag aagtgtctga tcttctgtgt cggtttggac agatataagt 1800
acccaagcga cgatactttc cgatactttc agaagtgtgt caaccagag gtcaacaatt 1860
tcgcgcaggc tccgtctgaa ggagataacg actcaaaactg aaggcatgaa ttttgacgat 1920
atttcagccg ctgtgcttaa gacaaaaatc gataagtcgg tgcaaatctt gcccaatcag 1980
ttgccggaca ttgttacgga ggcttccgag aaattttatt cgaatcgcac gtatcgtgtc 2040
attaagcagc atgaggtcct tgaggctgga accgacggcc gcttctttgc atacaaggct 2100
gacactttcg atgaataacc attcgacgtc caaaagttcg ccgacctgt tactgacagc 2160
cctgtgataa gcgcgacatc tgactttaag acgcttaaaa atcttaacga caactacggt 2220
ataactaggg aacaggcttt caacctgctg cgttccgac ctcgcgtctt gcgtgagttt 2280
attaatcagg agaaccatc tattcgcaac cgcacgagc agctcattct ccagtgcagg 2340
cttaaggcgc aaaacttgta ctttcaaggc catcaccatc accatcacta ggcgcccgct 2400
taattaa 2407

```

```

SEQ ID NO: 46      moltype = AA length = 2373
FEATURE           Location/Qualifiers
source            1..2373
                  mol_type = protein
                  organism = Rotavirus A

```

```

SEQUENCE: 46
METALASERL EUILETYRAR GGLNLEULEU SERASNSERT YRTHRVALAS PLEUSERASP 60
GLUILEGLNI LEILEGLYSE RGLULYSTHR GLNASNVALT HRILEASNPR OGLYPROPHE 120
ALAGLNTHRG LYTYRALAPR OVALASNTRP GLYPROGLYG LUTHRASNAS PSERTHRTHR 180
VALGLUPROV ALLEUASPGL YPROTYRGLN PROTHRTHRP HEASNPROPR OVALSERTYR 240
TRPILELEUL EUSERPROSE RASNALAGLY VALVALVALG LUGLYTHRAS NASNSERASP 300
ARGTRPLEUA LATHRILELE UILEGLUPRO ASNVALILES ERGLUSERAR GTHRTRYTHR 360
LEUPHEGLYA RGGLNGLUGL NILETHRVAL GLUASNVALS ERTHRTHRLY STRPLYSPE 420
VALASPLEUA LALYSTHRAS PILEASNGLY THRPHETHRG LNHISGLYPR OLEULEUSER 480
ASPTHRLYSL EUTYRGLYVA PMETLYSPHE SERGLYARGL EUTYRTHRTY RASNGLYGLU 540
THRPROASNA LATHRTHRGL YTYRTYRTHR THRTHRASNT YRASPTHRAV LASNMETTHR 600
SERTYRCYSA SPPEHYRIL EILEPROARG SERGLUGLUS ERALACYSTH RSERTYRILE 660
ASNASNGLYL EUPROPROAL EGLNASNTHR ARGASNVALV ALPROVALSE RLEUTHRSER 720
ARGSERILEV ALHISTHRAR GALAGLNALA ASNGLUASPI LEVALVALSE RLYSTHRSER 780
LEUTRPLYSG LUMETGLNTY RASNARGASP ILETHRILEA RGPHELYSPH EALAASNALA 840
ILEILELYSS ERGLYGLYLE UGLYTYRLYS TRPSEGLUI LESERPHELY SPROALAASN 900
TYRGLNTYRA SNTYRTHRAR GASPGLYGLU VALILETHRA LAHISTHRTH RCYSSERVAL 960
ASNGLYILEA SNASPPHESE RTYRASNGLY GLYSERLEUP ROTHRASPPH EVALILESER 1020
ARGTYRGLUV ALILELYSGL UASNSERTYR VALTYRVALA SPTYRTRPAS PASPSEGLN 1080
ALAPHEARGA SNMETVALTY RVALARGSER LEUALAALAA SNLEUASNSE RVALTHRCYS 1140
THRGLYGLYA SNTYRASPPH EARGLEUPRO VALGLYALAT RPPROVALME TTHRGLYGLY 1200
ALAVALSERL EUARGPROAL AGLYVALTHR LEUSERTHRG LNPHEHTRAS PPHEVALSER 1260
LEUASNSERL EUARGPHEAR GPHESEERLEU SERVALGLUG LUPROPROPH ETHRILETHR 1320
ARGTHRARGV ALSERGLYLE UTYRGLYLEU PROALAALAA SNPROASNAS NGLYLYSASP 1380
TYRTYRGLUI LELEUGLYA GPHESEERLEU ILESEERLEUV ALPROSERAS NASPASPTYR 1440
GLNTHRPROI LEMETASNSE RVALTHRVAL ARGGLNASPL EUGLUARGGL NLEUASNGLU 1500
LEUARGGLNG LUPHEASNSE RLEUSERGLN GLUVALALAM ETSEERGLNLE UILEASPLEU 1560
ALALEULEUP ROLEUASPMET PHESEERMET PHESEERGLYI LELYSSERTH RILEASPALA 1620
ALALYSSERM ETALATHRSE RVALMETLYS LYSPHELYSA RGSERASNLE UALASERSER 1680
ILESEERTHL EUTHRASPSR RLEUSERASP ALAALASERS ERILESEERAS GASNPROSER 1740
ILEARGSERA LAGLYSERSE RVALSERVAL TRPTHRASPI LESERTHRGL NALATHRASP 1800
ILESERASPT HRPHEARGSE RVALSERTHR GLNTHRSERT HRILESEERAR GARGLEUARG 1860
LEULYSGLUI LETHRTHRGL NTHRGLUGLY METASNPHEA SPASPILESE RALAALAVAL 1920
LEULYSTHRL YSILEASPLY SSERVALGLN ILESERPROA SNTHRLEUPR OASPILEVAL 1980
THRGLUALAS THRGLUALAS EILEPROASN ARGTHRTYRA RGVALILELY SASPASPLU 2040
VALLEUGLUA LAGLYTHRAS PGLYARGPHE PHEALATYRL YSVALASPTH RPHEASPGLU 2100
ILEPROPHEA SPVALGLNLY SPHEALAASP LEUVALTHRA SPSEERPROVA LILESERALA 2160
ILEILEASP HEGLYTHRLE ULYSASNLEU ASNASPASNT YRGLYILETH RARGGLUGLN 2220
ALAPHEASNL EULEUARGSE RASPPROARG VALLEUARGG LUPHEILEAS NGLNGLUASN 2280
PROILEILEA RGASNARGIL EGLUGLNLEU ILELEUGLNC YSARGLEULY SGLYGLUASN 2340
LEUTYRPHEG LNGLYHISHI SHISHISHIS HIS 2373

```

```

SEQ ID NO: 47      moltype = DNA length = 21

```

-continued

FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 47		
gagccaacgg tctgagttat t		21
SEQ ID NO: 48	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 48		
tcaaccgacg acctttaac c		21
SEQ ID NO: 49	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
source	1..21	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 49		
aatctcgctc cacttgtaac c		21
SEQ ID NO: 50	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
source	1..18	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 50		
tagggcgcaa gcaaatga		18
SEQ ID NO: 51	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
source	1..24	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 51		
cgactttata ccgctgaaca taga		24
SEQ ID NO: 52	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
source	1..19	
	mol_type = other DNA	
	organism = Synthetic construct	
SEQUENCE: 52		
gccgagcaac gacgattat		19

We claim herein:

1. A method for producing Rotavirus-derived immunogenic polypeptides and/or peptides, wherein optionally, said immunogenic polypeptides and/or peptides are mixed or co-expressed with one or more adjuvants.

2. The method of claim 1, comprising culturing a host cell transformed with a nucleic acid under conditions which induce expression of said polypeptides and/or peptides.

3. The method of claim 2, wherein the immunogenic polypeptides and/or peptides comprise a recombinant sub-unit vaccine, and wherein such expressed polypeptides and/or peptides are generated using baculovirus/insect cell methodology.

4. The method of claim 1, wherein the polypeptides and/or peptides comprise the amino acid sequence as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16 and 18.

5. The method of claim 2, wherein the nucleic acid encoding the Rotavirus derived immunogenic polypeptides and/or peptides is prepared by chemical synthesis.

6. The method of claim 5, wherein the nucleic acid encoding the Rotavirus derived immunogenic polypeptides and/or peptides is generated using a primer-based amplification method.

7. The method of claim 6, wherein the primer-based amplification method is PCR.

8. An immunogenic composition comprising the polypeptides as set forth in SEQ ID NOs: 2 and 46 or SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16 and 18.

9. A method of eliciting an immunological response in a subject comprising administering a composition as set forth in claim 8.

10. The method of claim 9, further comprising administering an adjuvant.

11. The method of claim 10, wherein administering said immunogenic composition to said subject is via topical, parenteral or mucosal administration.

12. The method of claim 9, wherein said administration is by multiple administrations.

13. The method of claim 12, wherein a first immunogenic composition and a second immunogenic composition are the same.

14. The method of claim 12, wherein a first immunogenic composition and the second immunogenic composition are different.

15. A method for treating an infection by a Rotavirus comprising administering to a subject in need thereof a therapeutically effective amount of an immunogenic composition of claim **8**.

16. A composition comprising the combination of polypeptides as set forth in SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, 16 and 18.

17. The composition of claim **16**, where said composition is formulated as a vaccine.

18. The composition of claim **17**, wherein the composition further comprises an adjuvant.

19. The composition of claim **18**, wherein the adjuvant is a water/oil/water adjuvant.

20. The composition of claim **16**, wherein the composition elicits neutralizing antibodies at a single dose.

* * * * *