


(43) **Pub. Date:** **Sep. 4, 2025**

Specification includes a Sequence Listing.






FIG. 1




FIG. 1 Continued

1A




FIG. 2A

1B




FIG. 2B

1C





FIG. 2C

ID





FIG. 2D

1E





FIG. 2E

2B




FIG. 3B

2C




FIG. 3C

2D







FIG. 3D

23



3A



3B




FIG. 4B

3C




FIG. 4C

3D






FIG. 4D


3E



[illegible]

FIG. 5A

4B



4C




FIG. 5C

4D






FIG. 5D



5A




FIG. 6A

5B






FIG. 6B

5C



5D






FIG. 6D


$$\text{CH}_2\text{Cl}_2$$

53

34

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525

 $\text{fl}(\text{ppm})$

FIG. 6E

6A




FIG. 7A

6B






FIG. 7B

6C



6D




FIG. 7D

6E




FIG. 7E

7A





FIG. 8A

7B



7C




FIG. 8C

7D




FIG. 8D

FIG. 8E

FIG. 8E

Chemical Structure: The structure shows a polymer repeat unit with a backbone containing a nitrogen atom (N) and a carbonyl group (C=O). The side chain includes a thioether linkage (-S-) and an ether linkage (-O-). Protons are labeled as follows: 'a' for the hydroxyl group, 'b' for the methylene group adjacent to the hydroxyl, 'c' for the methylene group adjacent to the thioether, 'd' for the methylene group adjacent to the ether, 'e' for the methylene group adjacent to the nitrogen, 'f' for the methylene group adjacent to the carbonyl, 'g' for the methine group, 'h' for the methylene group adjacent to the carbonyl, 'i' for the carbonyl group, and 'j' for the methylene group adjacent to the thioether.

1H NMR Spectrum: The spectrum is recorded in DMSO-d₆. The x-axis represents the chemical shift in ppm, ranging from 0 to 10. The peaks are assigned as follows: a broad peak at 1.2-1.6 ppm (a+c+d+e), a sharp peak at 2.4 ppm (f), a sharp peak at 2.8 ppm (h), a sharp peak at 3.0 ppm (j), a sharp peak at 3.3 ppm (b+j), a sharp peak at 3.6 ppm (b), a sharp peak at 4.2 ppm (g+i), and a sharp peak at 4.8 ppm (a). A solvent peak for DMSO is visible at 2.5 ppm.

FIG. 9A

8B




FIG. 9B


Chemical Structure: CCCCC(=O)OCCOCCCCNCCCCC(=O)OCCOCCCCNCCCCC(=O)OCCOCCCCN (simplified representation of the repeating unit). Protons are labeled: a (HO), b (CH2), c (CH2), d (CH2), e (CH2), f (CH2), g (N-CH2), h (CH2), i (CH2), j (CH2).


1H NMR Spectrum (DMSO-d₆):

- CDCl₃:** 7.26 ppm (triplet, solvent)
- DMSO:** 2.50 ppm (singlet, solvent)
- a:** 4.5-5.0 ppm (broad multiplet, OH)
- b:** 3.7 ppm (multiplet, CH₂)
- c+i:** 4.1 ppm (multiplet, CH₂)
- d+e+j:** 1.4-1.8 ppm (multiplet, CH₂)
- f+h:** 2.1-2.3 ppm (multiplet, CH₂)

FIG. 9C

8D





9A




FIG. 10A

9B






FIG. 10B

9C



9D



9E





FIG. 10E

10A



10B




FIG. 11B

10C




FIG. 11C

10D




FIG. 11D

10E






FIG. 11E



11B





FIG. 12B

11C



11D





FIG. 12D

11E



PEG-1A




FIG. 13A

PEG-1B




FIG. 13B

PEG-2A






FIG. 14A



PEG-3A





FIG. 15A

PEG-3B



PEG-4A




FIG. 16A

PEG-4B




FIG. 16B




FIG. 17A




FIG. 17B




FIG. 17C




FIG. 18




FIG. 19




FIG. 20

POLYMER-BASED NUCLEIC ACID MOLECULE DELIVERY VEHICLE HAVING IONIZATION MOIETY

INCORPORATION OF SEQUENCE LISTING

[0001] This application contains a sequence listing submitted in Computer Readable Form (CRF). The CRF file containing the sequence listing entitled “PK6387466-SequenceListing.xml”, which was created on May 14, 2025, and is 3,687 bytes in size. The information in the sequence listing is incorporated herein by reference in its entirety.

TECHNICAL FIELD

[0002] The present invention relates to a polymer-based nucleic acid molecule delivery vehicle having an ionizable moiety, and more specifically, to a polymer based on poly beta-aminoester or poly beta-aminoacrylate. Furthermore, the polymer according to the present invention can provide high mRNA delivery efficiency through pH sensitivity by attaching PEG to both ends to synthesize a triple copolymer.

BACKGROUND ART

[0003] mRNA is a substance that carries genetic information for producing proteins in the human body. Through mRNA delivery, it is possible to induce the production of proteins that are originally present in the human body, as well as the production of proteins that do not exist in the body. Since the outbreak of COVID-19, the development of mRNA delivery vehicles has been actively progressing as mRNA vaccines that activate immune responses through mRNA delivery have been utilized. Several mRNA delivery vehicles based on lipid nanoparticles have been developed, but they have problems due to rapid diffusion to organs other than the target site (off-target effect) and high immunogenicity, which can cause various side effects such as myocarditis and pericarditis. In addition, since vaccines are inoculated up to 2 to 4 times, the possibility of exposure to these side effects increases. Therefore, it is necessary to develop an mRNA delivery vehicle with high mRNA expression efficiency, low off-target effect, relatively low immunogenicity, and low side effects.

[0004] In order to improve the disadvantages of the above mRNA delivery vehicle, the present researchers studied mRNA delivery using polymers. As a result, a novel polymer that has pH sensitivity and has high mRNA delivery efficiency in vivo was newly discovered. Based on the above, the present researchers completed the present invention.

DETAILED DESCRIPTION OF INVENTION

Technical Problem


[0005] An object of the present invention is to provide a nucleic acid molecule delivery vehicle based on a polymer having pH sensitivity for high nucleic acid molecule delivery efficiency and low side effects.

[0006] One aspect of the object of the present invention is to provide a mRNA delivery vehicle based on a polymer having a relatively long duration of expression in order to overcome the short-term expression of mRNA in general.

[0007] Another object of the present invention is to provide a novel polymer for providing a nucleic acid molecule delivery vehicle based on a polymer having pH sensitivity.

Solution to Problem

[0008] In order to achieve the above object, the present invention provides a polymer suitable for intracellular nucleic acid delivery, comprising a repeating unit of Formula 1 below:



[0009] in the formula,

[0010] R1 is a bond or a C1-6 alkylene,

[0011] R2 and R3 are each independently (i) hydrogen, (ii) C1-6 alkyl which may be substituted with alkylamine or hydroxy, (iii) a 5- or 6-membered carbocycle or heterocycle, or (iv) C1-6 alkyl substituted with a 5- or 6-membered carbocycle or heterocycle,

[0012] said R2 and R3 may be linked together to form a C1-6 alkylene,


[0013] said 5- or 6-membered carbocycle or heterocycle may be substituted with C1-3 alkyl or C1-3 alkoxyacyl,

[0014] X is a nitrogen atom or a bond, and when X is a bond, said R3 is absent,

[0015] R4 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and


[0016] — may represent a single bond or a double bond.

[0017] The above repeating unit may be n (an integer between 1 and 500), and in this case, Formula 1 may be represented as follows.



[0018] (the definitions of the groups are as defined in relation to Formula 1 above)

[0019] The present invention provides a polymer having a repeating unit of Formula 2 below:




[0020] in the formula,

[0021] R5 is a C1-6 alkylene,

[0022] R6 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and

[0023] --- may represent a single bond or a double bond.

[0024] The above repeating unit may be n (an integer between 1 and 500), and in this case, Formula 2 may be represented as follows.












[0025] (the definitions of the groups are as defined in relation to Formula 2 above)

[0026] In one embodiment, the polymer may be a polymer in which the repeating unit of Formula 1 is formed by a reaction between any one of the monomers of Formulas 3 to 13 below and any one of the monomers of Formulas 14 to 23 below.


Formula	Structure
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	

-continued











Formula	Structure
15	
16	
17	
18	
19	
20	
21	
22	
23	

[0027] In one embodiment, the polymer may be a polymer in which the repeating unit of Formula 2 is formed by a

reaction between a monomer of Formula 24 below and any one of the monomers of Formulas 14 to 23 below.

Formula	Structure
14	






-continued

Formula	Structure
15	
16	
17	
18	
19	
20	
21	
22	
23	
24	

[0028] The present invention provides a polymer comprising a repeating unit selected from the group consisting of repeating units set forth in the table below.

Manufacturing Example No.	Chemical structural formula
1 1A	
2 1B	
3 1C	
4 1D	
5 1E	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
6	2A	
7	2B	
8	2C	
9	2D	
10	2E	






-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
11	3A	
12	3B	
13	3C	




-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
14	3D	
15	3E	
16	4A	





-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
20	4E	
21	5A	
22	5B	
23	5C	
24	5D	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
25	5E	
26	6A	
27	6B	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
31	7A	
32	7B	
33	7C	
34	7D	

-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
35	7E	
36	8A	
37	8B	




-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
38	8C	
39	8D	
40	8E	

-continued

Man- ufac- tur- ing Ex- am- ple No.	Chemical structural formula
41 9A	
42 9B	
43 9C	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
44	9D	
45	9E	
46	10A	






-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
47	10B	
48	100	





-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
51	11A	
52	11B	
53	11C	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
58	1H	
59	1I	
60	1J	
61	2F	
62	2G	

-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
63	2H	
64	2I	
65	2J	
66	3F	

-continued

Man- ufac- turing Ex- am- ple No.	Code No.	Chemical structural formula
67	3G	
68	3H	
69	3I	






-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
70	3J	
71	4F	
72	4G	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
73	4H	
74	4I	
75	4J	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
76	5F	
77	5G	
78	5H	
79	5I	
80	5J	

-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
81	6F	
82	6G	
83	6H	



-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
84	6I	
85	6J	
86	7F	



-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
91	8F	
92	8G	
93	8H	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
102	10G	
103	10H	





-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
104	10I	
105	10J	

-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
109	11I	
110	11J	
111	PEG- 1A	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
112	PEG- 1B	
113	PEG- 1C	
114	PEG- 1D	
115	PEG- 1E	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code	Chemical structural formula
116	PEG- 2A	
117	PEG- 2B	
118	PEG- 2C	
119	PEG- 2D	
120	PEG- 2E	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
121	PEG- 3A	
122	PEG- 3B	
123	PEG- 3C	

-continued

Man- ufac- turing Ex- am- ple No.	Code	Chemical structural formula
124	PEG- 3D	
125	PEG- 3E	
126	PEG- 4A	





-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
127	PEG- 4B	
128	PEG- 4C	
129	PEG- 4D	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
130	PEG- 4E	
131	PEG- 5A	
132	PEG- 5B	
133	PEG- 5C	
134	PEG- 5D	





-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
135	PEG- 5E	
136	PEG- 6A	
137	PEG- 6B	
138	PEG- 6C	



-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
139	PEG- 6D	
140	PEG- 6E	
141	PEG- 7A	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
142	PEG- 7B	
143	PEG- 7C	
144	PEG- 7D	
145	PEG- 7E	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
146	PEG- 8A	
147	PEG- 8B	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
148	PEG-8C	
149	PEG-8D	
150	PEG-8E	



-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
151	PEG-9A	
152	PEG-9B	
153	PEG-9C	

-continued

Man- ufac- turing Ex- am- ple No.	Code	Chemical structural formula
154	PEG- 9D	
155	PEG- 9E	
156	PEG- 10A	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
157	PEG- 10B	
158	PEG- 10C	




-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
159	PEG- 10D	The chemical structure of PEG-10D consists of a central poly(ethylene glycol) (PEG) chain, represented by a wavy line with a subscript 'n'. This central chain is flanked by two identical side chains. Each side chain starts with a methyl group attached to a carbon atom, which is also bonded to an oxygen atom. This oxygen atom is part of an ester linkage to a 4-(4-methylpiperazin-1-yl)butanoate group. The side chain is terminated by a methyl group attached to a carbon atom, which is also bonded to an oxygen atom. The entire structure is enclosed in brackets with a subscript 'n'.
160	PEG- 10E	The chemical structure of PEG-10E consists of a central poly(ethylene glycol) (PEG) chain, represented by a wavy line with a subscript 'n'. This central chain is flanked by two identical side chains. Each side chain starts with a methyl group attached to a carbon atom, which is also bonded to an oxygen atom. This oxygen atom is part of an ester linkage to a 4-(4-methylpiperazin-1-yl)butanoate group. The side chain is terminated by a methyl group attached to a carbon atom, which is also bonded to an oxygen atom. The entire structure is enclosed in brackets with a subscript 'n'.





-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
161	PEG- 11A	
162	PEG- 11B	
163	PEG- 11C	

-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
164	PEG- 11D	
165	PEG- 11E	
166	PEG- 1F	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
167	PEG- 1G	
168	PEG- 1H	
169	PEG- 1I	
170	PEG- 1J	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
171	PEG- 2F	
172	PEG- 2G	
173	PEG- 2H	
174	PEG- 2I	
175	PEG- 2J	

-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
176	PEG- 3F	
177	PEG- 3G	
178	PEG- 3H	






-continued

Man- ufac- turing Ex- am- ple No.	Code	Chemical structural formula
179	PEG- 3I	
180	PEG- 3J	
181	PEG- 4F	





-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
182	PEG- 4G	
183	PEG- 4H	
184	PEG- 4I	





-continued

Man- ufac- tur- ing Ex- am- ple No.	Code	Chemical structural formula
185	PEG- 4J	
186	PEG- 5F	
187	PEG- 5G	
188	PEG- 5H	
189	PEG- 5I	

-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
190	PEG- 5J	
191	PEG- 6F	
192	PEG- 6G	
193	PEG- 6H	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
194	PEG- 6f	
195	PEG- 6j	
196	PEG- 7F	
197	PEG- 7G	



-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
198	PEG- 7H	
199	PEG- 7I	
200	PEG- 7I	



-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
203	PEG- 8H	
204	PEG- 8I	
205	PEG- 8J	



-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
209	PEG- 91	
210	PEG- 91	




-continued

Man- ufac- turing Ex- am- ple No.	Code	Chemical structural formula
211	PEG- 10F	
212	PEG- 10G	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
213	PEG- 10H	
214	PEG- 10I	

-continued

Man- ufac- turing Ex- am- ple No.	Code No.	Chemical structural formula
215	PEG-10J	
216	PEG-11F	
217	PEG-11G	

-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
218	PEG- 11H	
219	PEG- 11I	
220	PEG- 11J	

-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
221	b3A	
222	b3B	
223	b3C	
224	b3D	









-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula
225	b3E	
226	b4A	
227	b4B	




-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
228	b4C	
229	b4D	
230	b4E	
231	b7A	




-continued

Man- ufac- tur- ing Ex- am- ple	Code No.	Chemical structural formula	
232	b7B		
233	b7C		
234	b7D		
235	b7E		




-continued

Man- ufac- turing Ex- am- ple No.	Code No.	Chemical structural formula
236	b8A	
237	b8B	
238	b8C	

-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
245	b9E	
246	b10A	
247	b10B	



-continued

Man- ufac- turing Ex- am- ple No.	Code No.	Chemical structural formula
248	b10C	
249	b10D	
250	b10E	



-continued

Man- ufac- tur- ing Ex- am- ple No.	Code No.	Chemical structural formula
251	b11A	
252	b11B	
253	b11C	

-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
257	b12B	
258	b12C	

-continued

Man- ufac- turing Ex- am- ple	Code No.	Chemical structural formula
259	b12D	
260	b12E	


[0029] (in the above formulas, n is an integer between 1 and 500)

[0041] (the definitions of the groups are as defined in relation to Formula 26 above)

[0042] When the polymer according to the present invention comprises a structure of Formula 25 above, the polymer may comprise a structure of Formula 27 below.

[0053] The present invention provides a polymeric nanoparticle in which the polymer and a nucleic acid molecule form a complex. The polymeric nanoparticle according to the present invention preferably does not include a lipid nanoparticle (LNP).

[Formula 27]



[0043] in the formula,

[0044] R5 is a C1-6 alkylene,

[0045] R6 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and


[0046] ----- may represent a single bond or a double bond.

[0047] The above repeating unit may be n (an integer between 1 and 500), and in this case, Formula 27 may be represented as follows.

[0054] The nucleic acid molecule that can be included in the polymeric nanoparticle according to the present invention may be RNA. The RNA is preferably mRNA.

[0055] The nucleic acid molecule that can be included in the polymeric nanoparticle according to the present invention may be DNA.


[0056] The polymeric nanoparticle according to the present invention may be one in which the nucleic acid molecule is isolated in a pH-dependent manner and delivered into cells in the body.



[0048] (the definitions of the groups are as defined in relation to Formula 27 above)

[0049] In one embodiment, one or both ends of the polymer may be linked to a PEG-containing moiety having a structure of Formula 28 below:

[Formula 28]



[0050] in the formula,

[0051] ===== may represent a double bond or a triple bond, and

[0052] n may be an integer in the range of 1 to 500.

[0057] The polymeric nanoparticle according to the present invention may be one in which when delivered into the body, the expression of the nucleic acid molecule is maintained for a longer period of time compared to when the same nucleic acid molecule is delivered into the body in the form of a non-polymeric lipid nanoparticle.

[0058] The present invention provides a method for delivering a nucleic acid molecule into a cell using the polymeric nanoparticle. The nucleic acid molecule delivery method according to the present invention preferably does not use a lipid nanoparticle (LNP).

[0059] The polymeric nanoparticle according to the present invention may be delivered by parenteral, intramuscular, subcutaneous, or intravenous administration.

[0060] The present invention provides a pharmaceutical composition comprising the polymeric nanoparticle. The pharmaceutical composition according to the present invention preferably does not include a lipid nanoparticle (LNP).

[0061] The pharmaceutical composition comprising the polymeric nanoparticle according to the present invention may be a gene therapeutic agent.

[0062] The pharmaceutical composition comprising the polymeric nanoparticle according to the present invention may be a vaccine.

Effects of Invention

[0063] Since the nucleic acid molecule delivery vehicle based on a polymer having pH sensitivity according to the present invention has a high nucleic acid molecule delivery efficiency, it can be used as an excellent delivery vehicle. In particular, the nucleic acid molecule delivery vehicle according to the present invention can be utilized as a therapeutic gene delivery vehicle and an mRNA vaccine.

[0064] In addition, depending on the length of the monomer and polymer, the nucleic acid molecule can be delivered to a desired site, and the nucleic acid molecule can be released in response to a specific stimulus through an additional stimulus-sensitive linker.

[0065] In addition, the nucleic acid molecule delivery vehicle according to the present invention can be degraded after releasing the nucleic acid molecule within the cell, so it has low side effects.

BRIEF DESCRIPTION OF DRAWINGS

[0066] FIG. 1 is a schematic diagram of the synthesis of the polymer delivery vehicle according to the present invention and a schematic diagram of a polymeric nanoparticle (PNP) comprising the polymer delivery vehicle and mRNA.

[0067] FIGS. 2A to 2E are graphs showing the ¹H-NMR results of polymers 1A to 1E.

[0068] FIGS. 3A to 3E are graphs showing the ¹H-NMR results of polymers 2A to 2E.

[0069] FIGS. 4A to 4E are graphs showing the ¹H-NMR results of polymers 3A to 3E.

[0070] FIGS. 5A to 5E are graphs showing the ¹H-NMR results of polymers 4A to 4E.

[0071] FIGS. 6A to 6E are graphs showing the ¹H-NMR results of polymers 5A to 5E.

[0072] FIGS. 7A to 7E are graphs showing the ¹H-NMR results of polymers 6A to 6E.

[0073] FIGS. 8A to 8E are graphs showing the ¹H-NMR results of polymers 7A to 7E.

[0074] FIGS. 9A to 9E are graphs showing the ¹H-NMR results of polymers 8A to 8E.

[0075] FIGS. 10A to 10E are graphs showing the ¹H-NMR results of polymers 9A to 9E.

[0076] FIGS. 11A to 11E are graphs showing the ¹H-NMR results of polymers 10A to 10E.

[0077] FIGS. 12A to 12E are graphs showing the ¹H-NMR results of polymers 11A to 11E.

[0078] FIGS. 13A and 13B are graphs showing the ¹H-NMR results of polymers PEG-1A and PEG-1B.

[0079] FIGS. 14A and 14B are graphs showing the ¹H-NMR results of polymers PEG-2A and PEG-2B.

[0080] FIGS. 15A and 15B are graphs showing the ¹H-NMR results of polymers PEG-3A and PEG-3B.

[0081] FIGS. 16A and 16B are graphs showing the ¹H-NMR results of polymers PEG-4A and PEG-4B.

[0082] FIGS. 17A to 17C are photographs showing the electrophoresis results of the polymeric nanoparticles (PNP) according to the present invention.

[0083] FIG. 18 is a graph showing the zeta potential of the polymeric nanoparticles (PNP) according to the present invention.

[0084] FIG. 19 is a photograph showing the mRNA delivery of the polymeric nanoparticles (PNP) according to the present invention in vivo.

[0085] FIG. 20 is a graph showing the expression duration of mRNA delivered to the polymeric nanoparticles (PNP) according to the present invention.

BEST MODE FOR CARRYING OUT THE INVENTION

[0086] The manufacturing examples and embodiments of the present invention will be described in detail so that those of ordinary skill in the art to which the present invention belongs can easily practice the present invention. However, the present invention may be implemented in various forms and is not limited to the manufacturing examples and embodiments described herein.

[0087] Throughout the present specification, when a certain part “includes” a certain component, it means that other components may be further included, rather than excluding other components, unless otherwise specifically stated.

[0088] Throughout the present specification, “polymer”, “macromolecule,” and “polymer” are used interchangeably with each other with the same meaning.

[0089] The term “repeating unit” used in describing “polymer” in the present specification refers to a structure in which the same structure appears twice or more repeatedly throughout the length of the polymer, and the polymer may not necessarily be composed of only the repeating unit. For example, the polymer may further comprise branched structures and/or end-capping agents in addition to the given repeating units.

[0090] As used herein, the term “substituted”, for example “substituted alkyl,” means that one or more hydrogen atoms of the alkyl are each independently replaced by a non-hydrogen substituent. The substituent may include, but is not limited to, any of the substituents described herein, for example, halogen, hydroxyl, alkyl, hydroxyalkyl, haloalkyl, cyanoalkyl, alkoxyalkyl, carbonyl (e.g., carboxyl, alkoxy-carbonyl, formyl, or acyl), thiocarbonyl (e.g., thioester, thioacetate, or thioformate), alkoxy, phosphoryl, phosphate, phosphonate, phosphinate, amino, amido, amidine, imine, cyano, nitro, azido, sulfhydryl, alkylthio, sulfate, sulfonate, sulfamoyl, sulfonamido, sulfonyl, heterocyclyl, aralkyl, aryl, or heteroaryl. The substituted moiety on the hydrocarbon chain may itself be substituted, if desired.

[0091] In the present invention, an “alkyl group” is a hydrocarbon having substituted or unsubstituted primary, secondary, tertiary and/or quaternary carbon atoms, and includes a saturated aliphatic group which may be straight-chain, branched, cyclic, or a combination thereof. For example, the alkyl group may have 1 to 20 carbon atoms (i.e., a C1-C20 alkyl), 1 to 10 carbon atoms (i.e., a C1-C10 alkyl), or 1 to 6 carbon atoms (i.e., a C1-C6 alkyl). Examples of suitable alkyl groups may include, but are not limited to, methyl (Me, —CH₃), ethyl (Et, —CH₂CH₃), 1-propyl (n-Pr, n-propyl, —CH₂CH₂CH₃), 2-propyl (i-Pr, i-propyl, —CH(CH₃)₂), 1-butyl (n-Bu, n-butyl, —CH₂CH₂CH₂CH₃), 2-methyl-1-propyl (i-Bu, i-butyl, —CH₂CH(CH₃)CH₂CH₃), 2-butyl (s-Bu, s-butyl, —CH(CH₃)CH₂CH₃), 2-methyl-2-propyl (t-Bu, t-butyl, —C(CH₃)₃), 1-pentyl (n-pentyl, —CH₂CH₂CH₂CH₂CH₃), 2-pentyl

(—CH(CH₃)CH₂CH₂CH₃), 3-pentyl (—CH(CH₂CH₃)₂), 2-methyl-2-butyl (—C(CH₃)₂CH₂CH₃), 3-methyl-2-butyl (—CH(CH₃)CH(CH₃)₂), 3-methyl-1-butyl (—CH₂CH₂CH(CH₃)₂), 2-methyl-1-butyl (—CH₂CH(CH₃)CH₂CH₃), 1-hexyl (—CH₂CH₂CH₂CH₂CH₂CH₃), 2-hexyl (—CH(CH₃)CH₂CH₂CH₂CH₃), 3-hexyl (—CH(CH₂CH₃)CH₂CH₂CH₃), 2-methyl-2-pentyl (—C(CH₃)₂CH₂CH₂CH₃), 3-methyl-2-pentyl (—CH(CH₃)CH(CH₃)CH₂CH₃), 4-methyl-2-pentyl (—CH(CH₃)CH₂CH(CH₃)₂), 3-methyl-3-pentyl (—C(CH₃)(CH₂CH₃)₂), 2-methyl-3-pentyl (—CH(CH₂CH₃)CH(CH₃)₂), 2,3-dimethyl-2-butyl (—C(CH₃)₂CH(CH₃)₂), 3,3-dimethyl-2-butyl (—CH(CH₃)C(CH₃)₃), and octyl (—(CH₂)₇CH₃).

[0092] Moreover, as used throughout the specification, examples and claims, the term “alkyl” is intended to include both unsubstituted and substituted alkyl groups, the latter of which refers to an alkyl moiety having a substituent that replaces a hydrogen on one or more carbons of the hydrocarbon backbone, which includes a haloalkyl group such as trifluoromethyl.

[0093] As used herein, the term “alkylene” refers to a saturated hydrocarbon group having two valences, which may be branched, straight-chain, cyclic, or a combination thereof, and which is derived by removing two hydrogen atoms from the same carbon atom or two different carbon atoms of a parent alkane. For example, an alkylene group may have 1 to 20 carbon atoms, 1 to 10 carbon atoms, or 1 to 6 carbon atoms. Examples of suitable alkylene groups may include, but are not limited to, methylene (—CH₂—) and 1,2-ethylene (—CH₂—CH₂—).

[0094] As used herein, the term “heterocycle” refers to a substituted or unsubstituted, monovalent or divalent, saturated or partially saturated non-aromatic ring structure, preferably a 3- to 10-membered ring, more preferably a 3- to 7-membered ring, wherein the ring structure comprises one or more heteroatoms, preferably 1 to 4 heteroatoms, and more preferably 1 to 2 heteroatoms. The terms “heterocyclyl,” “heterocycle,” “heterocyclic,” and “heterocycloalkyl” also include polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjacent rings, wherein at least one of the rings is heterocyclic, for example, the other cyclic rings may be cycloalkyl, cycloalkenyl, cycloalkynyl, aryl, heteroaryl, and/or heterocyclyl. The bicyclic and polycyclic heterocyclic ring systems may be fused, bridged, or spiro ring systems. The substituted heterocycles include heterocyclic rings substituted with any of the substituents disclosed herein, including, for example, a carbonyl group. The heterocyclyl groups include, for example, piperidine, piperazine, pyrrolidine, morpholine, lactones, lactams, and the like. Additionally exemplary heterocycles may include, but are not limited to, dihydropyridyl, dihydroindolyl, tetrahydropyridyl (piperidyl), tetrahydrothiophenyl, sulfur-oxidized tetrahydrothiophenyl, indolenyl, piperidinyl, 4-piperidinyl, pyrrolidinyl, 2-pyrrolidinyl, pyrrolinyl, tetrahydrofuranlyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, decahydroquinolinyl, octahydroisoquinolinyl, 6H-1,2,5-thiadiazinyl, 2H,6H-1,5,2-dithiazinyl, pyranlyl, chromenyl, xanthenyl, phenoxathinyl, 2H-pyrrolyl, 3H-indolyl, 4H-quinoliziny, phthalazinyl, naphthyridinyl, quinoxalinyl, quinazolinyl, cinnolinyl, pteridinyl, 4aH-carbazolyl, carbazolyl, β-carbolinyl, phenanthridinyl, acridinyl, phenanthrolinyl, phenazinyl, phenothiazinyl, furazanyl, phenoxazinyl, isochromanyl, chromanyl, imidazolidinyl,

imidazolinyl, pyrazolidinyl, pyrazolinyl, piperazinyl, quinuclidinyl, morpholinyl, and oxazolidinyl (each of which may be substituted or unsubstituted).

[0095] As used herein, the term “carbocycle” refers to a non-aromatic saturated or unsaturated, monovalent or divalent ring which may be monocyclic, bicyclic, or polycyclic and in which each of the ring atoms is carbon. Cycloalkyl groups may have from 3 to 7 carbon atoms as a monocycle, from 7 to 12 carbon atoms as a bicycle, and up to about 20 carbon atoms as a polycycle. Monocyclic cycloalkyls have from 3 to 7 ring atoms, more typically 5 or 6 ring atoms. Bicyclic cycloalkyls may have from 7 to 12 ring atoms and may be fused ring systems, spirocyclic ring systems, or bridged ring systems. In exemplary cycloalkyl groups, the atoms may be arranged in a bicyclo[4,5], [5,5], [5,6], or [6,6] system. In certain embodiments, the cycloalkyl contains from 3 to 20 atoms, or from 3 to 10 atoms, or more preferably from 3 to 7 atoms. Examples of cycloalkyls may include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, and the like. Unless otherwise specified, cycloalkyl may be substituted by one or more substituents described herein.

[0096] As used herein, the term “alkoxyacyl” refers to a group of —C(=O)OR. R is alkyl as defined herein, which may be substituted or unsubstituted.

[0097] — as used in the chemical structural formulas herein represents a single bond or a double bond, according to the convention used in the art.

[0098] ~ as used in the chemical structural formulas herein represents a bond connected to another unit, according to the convention used in the art.

[0099] As used herein, the term “amine” refers to —NH₂, in which the hydrogen atom is unsubstituted or substituted with a substituent such as alkyl or aryl, wherein the substituent such as alkyl or aryl that substitutes the hydrogen atom is as defined herein, and may be substituted or unsubstituted.

[0100] As used herein, the term “amine monomer” is a monomer containing an amine functional group, and preferably, may be selected from the group consisting of amine monomers described in Table 1 below.

TABLE 1





No.	Structural formula
1	
2	
3	
4	

TABLE 1-continued

No.	Structural formula
5	
6	
7	
8	
9	

TABLE 1-continued

No.	Structural formula
10	
11	
12	

[0101] As used herein, the term “acrylate monomer” is a monomer in the form of an ester derivative of acrylic acid, and may be selected from the group consisting of acrylate monomers described in Table 2 below.

TABLE 2





No.	Structural formula
A	
B	
C	
D	
E	

[0102] As used herein, the term “alkynoate monomer” is a monomer in the form of an ester derivative of alkynoic acid, and may be selected from the group consisting of alkynoate monomers described in Table 3 below.

TABLE 3

No.	Structural formula
F	

TABLE 3-continued


No.	Structural formula
G	
H	
I	
J	

[0103] The polymer according to the present invention is characterized by including an amine monomer and an acrylate or alkynoate monomer together, and the two monomers are alternately connected (see schematic diagram of FIG. 1).

[0104] The polymer according to the present invention is characterized by having an ionizable moiety, where the “ionizable moiety” means a part of the polymer that is ionized depending on the pH. For example, it may be a tertiary amine present in the polymer chemical structure, and those of ordinary skill in the art are well aware of which moiety can be ionized depending on the pH.


[0105] Hereinafter, the present invention will be described in more detail through the examples, but the following examples are for illustrative purposes only and are not intended to limit the scope of the present invention.

Manufacturing Example 1




[0106] Acrylate monomer A (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. Dichloromethane (CH_2Cl_2 , DCM) was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 1A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 2



[0107] Acrylate monomer B (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 1B having the ionizable moiety of the present invention was obtained.


Manufacturing Example 3






[0108] Acrylate monomer C (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 1C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 4



[0109] Acrylate monomer D (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 1D having the ionizable moiety of the present invention was obtained.


Manufacturing Example 5



[0110] Acrylate monomer E (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 2500 mg/ml. Thereafter, the reaction


was carried out at 50° C. for 48 hours with reflux, and then polymer 1E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 6



[0111] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 2 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. Dimethylsulfoxide (DMSO) was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 2A having the ionizable moiety of the present invention was obtained.

Manufacturing Example 7




[0112] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 2 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at

90° C. for 24 hours, and then polymer 2B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 8


Preparation of polymer 2C



[0113] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 2 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 2C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 9


Preparation of polymer 2D



[0114] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 2 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 2D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 10


Preparation of polymer 2E



[0115] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 2 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 2E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 11


Preparation of polymer 3A




[0116] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 3 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 3A having the ionizable moiety of the present invention was obtained.

Manufacturing Example 12

Preparation of polymer 3B







[0118] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 3 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 3C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 14


[0117] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 3 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 3B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 13




[0119] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 3 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 3D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 15




[0120] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 3 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 3E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 16



[0121] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 4 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 4A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 17



[0122] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 4 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration


was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 4B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 18



[0123] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 4 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 4C having the ionizable moiety of the present invention was obtained.


Manufacturing Example 19



[0124] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 4 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 4D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 20


Preparation of polymer 4A



[0125] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 4 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 4E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 21

Preparation of polymer 5A





[0126] Acrylate monomer A (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction

was carried out at 50° C. for 48 hours with reflux, and then polymer 5A having the ionizable moiety of the present invention was obtained.

Manufacturing Example 22


Preparation of polymer 5B





[0127] Acrylate monomer B (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 5B having the ionizable moiety of the present invention was obtained.


Manufacturing Example 23



[0128] Acrylate monomer C (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction

was carried out at 50° C. for 48 hours with reflux, and then polymer 5C having the ionizable moiety of the present invention was obtained.


Manufacturing Example 24



[0129] Acrylate monomer D (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 50° C. for 48 hours with reflux, and then polymer 5D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 25

Preparation of polymer 5E




[0130] Acrylate monomer E (1000 mg) was placed in a 2-neck round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the acrylate monomer was 1.04. DCM was added to the flask so that the final concentration was 1000 mg/ml. Thereafter, the reaction

was carried out at 50° C. for 48 hours with reflux, and then polymer 5E having the ionizable moiety of the present invention was obtained.


Manufacturing Example 26

Preparation of polymer 6A



[0131] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 6 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 6A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 27



[0132] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 6 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at


90° C. for 24 hours, and then polymer 6B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 28




[0133] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 6 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 6C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 29




[0134] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 6 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 6D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 30




[0135] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 6 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 6E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 31




-continued



[0136] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 7 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 7A having the ionizable moiety of the present invention was obtained.

Manufacturing Example 32



Preparation of polymer 7B




[0137] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 7 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 7B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 33

Preparation of polymer 7C




-continued



[0138] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 7 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 7C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 34


Preparation of polymer 7D



[0139] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 7 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 7D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 35

Preparation of polymer 7E




[0140] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 7 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 7E having the ionizable moiety of the present invention was obtained.

that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 8A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 37

Preparation of polymer 8B



Manufacturing Example 36


Preparation of polymer 8A




[0142] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 8 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 8B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 38

Preparation of polymer 8C




[0141] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 8 was added to the vial so



[0143] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 8 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 8C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 39


Preparation of polymer 8D



[0144] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 8 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 8D having the ionizable moiety of the present invention was obtained.


Manufacturing Example 40

Preparation of polymer 8E




[0145] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 8 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 8E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 41



[0146] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 9 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 9A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 42



[0147] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 9 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at


90° C. for 24 hours, and then polymer 9B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 43




[0148] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 9 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 9C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 44



[0149] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 9 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 9D having the ionizable moiety of the present invention was obtained.



Manufacturing Example 45

Preparation of polymer 9E

[0150] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 9 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 9E having the ionizable moiety of the present invention was obtained.


Manufacturing Example 46

Preparation of Polymer 10A




[0151] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 10 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 10A having the ionizable moiety of the present invention was obtained.

Manufacturing Example 47

Preparation of polymer 10B

[0152] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 10 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 10B having the ionizable moiety of the present invention was obtained.


Manufacturing Example 48



[0153] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 10 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration


was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 10C having the ionizable moiety of the present invention was obtained.

Manufacturing Example 49





[0154] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 10 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 10D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 50




[0155] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 10 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 10E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 51




[0156] Acrylate monomer A (1000 mg) was placed in a 20 ml vial, and then amine monomer 11 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 11A having the ionizable moiety of the present invention was obtained.


Manufacturing Example 52



[0157] Acrylate monomer B (1000 mg) was placed in a 20 ml vial, and then amine monomer 11 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 11B having the ionizable moiety of the present invention was obtained.

Manufacturing Example 53







[0158] Acrylate monomer C (1000 mg) was placed in a 20 ml vial, and then amine monomer 11 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 11C having the ionizable moiety of the present invention was obtained.

so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 1000 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 11D having the ionizable moiety of the present invention was obtained.

Manufacturing Example 55




Manufacturing Example 54




[0160] Acrylate monomer E (1000 mg) was placed in a 20 ml vial, and then amine monomer 11 was added to the vial so that the molar ratio with the acrylate monomer was 1.04. DMSO was added to the vial so that the final concentration was 2500 mg/ml. Thereafter, the reaction was carried out at 90° C. for 24 hours, and then polymer 11E having the ionizable moiety of the present invention was obtained.

Manufacturing Example 56




[0159] Acrylate monomer D (1000 mg) was placed in a 20 ml vial, and then amine monomer 11 was added to the vial



[0161] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the acrylate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 1F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 57



Preparation of polymer 1G



[0162] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 1G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 58


Preparation of polymer 1H



[0163] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 1H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 59


Preparation of polymer 1I



[0164] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 1 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 1I having the ionizable moiety of the present invention was obtained.


Manufacturing Example 60

Preparation of polymer 1J




[0165] monomer 1 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 1J having the ionizable moiety of the present invention was obtained.

Manufacturing Example 61



[0166] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 2 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 2F having the ionizable moiety of the present invention was obtained.


Manufacturing Example 62



[0167] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 2 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room


temperature for 6 hours, and then polymer 2G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 63




[0168] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 2 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 2H having the ionizable moiety of the present invention was obtained.


Manufacturing Example 64



[0169] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 2 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 2I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 65






[0170] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 2 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 2J having the ionizable moiety of the present invention was obtained.

Manufacturing Example 66


Preparation of polymer 3F



[0171] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 3 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 3F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 67

Preparation of polymer 3G




[0172] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 3 was added to the flask so that the molar ratio with the alkynoate

monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 3G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 68


Preparation of polymer 3H



[0173] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 3 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 3H having the ionizable moiety of the present invention was obtained.


Manufacturing Example 69

Preparation of polymer 3I




[0174] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 3 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 3I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 70


Preparation of polymer 3J

[0175] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 3 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 3J having the ionizable moiety of the present invention was obtained.

-continued




Manufacturing Example 71

Preparation of polymer 4F


[0177] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 4 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 4G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 73

Preparation of polymer 4H



[0176] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 4 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 4F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 72

Preparation of polymer 4G


[0178] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 4 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 4H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 74




[0179] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 4 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 4I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 75



[0180] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 4 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 4J having the ionizable moiety of the present invention was obtained.


Manufacturing Example 76



[0181] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 5F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 77


Preparation of polymer 5G



[0182] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 5G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 78

Preparation of polymer 5H





[0183] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room

temperature for 6 hours, and then polymer 5H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 79

Preparation of polymer 5I






[0184] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 5I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 80

Preparation of polymer 5J




[0185] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 5 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room

temperature for 6 hours, and then polymer 5J having the ionizable moiety of the present invention was obtained.


Manufacturing Example 81

Preparation of polymer 6F



[0186] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 6 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 6F having the ionizable moiety of the present invention was obtained.


Manufacturing Example 82



[0187] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 6 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room

temperature for 6 hours, and then polymer 6G having the ionizable moiety of the present invention was obtained.


Manufacturing Example 83



[0188] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 6 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 6H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 84


Preparation of polymer 6I



[0189] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 6 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 6I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 85


Preparation of polymer 6J




[0190] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 6 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 6J having the ionizable moiety of the present invention was obtained.

Manufacturing Example 86


Preparation of polymer 7F







[0191] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 7 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 7F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 87




[0192] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 7 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 7G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 88




[0193] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 7 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 7H having the ionizable moiety of the present invention was obtained.


Manufacturing Example 89



[0194] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 7 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 7I having the ionizable moiety of the present invention was obtained.


Manufacturing Example 90





[0195] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 7 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 7J having the ionizable moiety of the present invention was obtained.


Manufacturing Example 91




[0196] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 8 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room

temperature for 6 hours, and then polymer 8F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 92




-continued



[0197] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 8 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 8G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 93

Preparation of polymer 8H




[0198] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 8 was added to the flask so that the molar ratio with the alkynoate

monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 8H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 94


Preparation of polymer 8I



[0199] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 8 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 8I having the ionizable moiety of the present invention was obtained.



Manufacturing Example 95

Preparation of polymer 8J




[0203] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 9 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 9H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 99




[0204] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 9 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 9I having the ionizable moiety of the present invention was obtained.


Manufacturing Example 100



[0205] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 9 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 9J having the ionizable moiety of the present invention was obtained.

Manufacturing Example 101






[0206] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 10 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 10F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 102

Preparation of polymer 10G





[0207] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 10 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room

temperature for 6 hours, and then polymer 10G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 103



Preparation of polymer 10H






[0208] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 10 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 10H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 104




[0209] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 10 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 10I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 105




[0210] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 10 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 10J having the ionizable moiety of the present invention was obtained.

Manufacturing Example 106




[0211] Alkynoate monomer F (500 mg) was placed in a round bottom flask, and then amine monomer 11 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 11F having the ionizable moiety of the present invention was obtained.

Manufacturing Example 107




[0212] Alkynoate monomer G (500 mg) was placed in a round bottom flask, and then amine monomer 11 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 11G having the ionizable moiety of the present invention was obtained.

Manufacturing Example 108




[0213] Alkynoate monomer H (500 mg) was placed in a round bottom flask, and then amine monomer 11 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 11H having the ionizable moiety of the present invention was obtained.

Manufacturing Example 109




[0214] Alkynoate monomer I (500 mg) was placed in a round bottom flask, and then amine monomer 11 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 11I having the ionizable moiety of the present invention was obtained.

Manufacturing Example 110



[0215] Alkynoate monomer J (500 mg) was placed in a round bottom flask, and then amine monomer 11 was added to the flask so that the molar ratio with the alkynoate monomer was 1.04, and 4 mL of dichloromethane was added to the flask. Thereafter, the reaction was carried out at room temperature for 6 hours, and then polymer 11J having the ionizable moiety of the present invention was obtained.


Manufacturing Example 111




[0216] Polymer 1A prepared in the same manner as in Manufacturing Example 1 above and 648.8 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried

out at 50°C. for 24 hours with reflux, and then PEG-1A was obtained.

Manufacturing Example 112




-continued



[0217] Polymer 1B prepared in the same manner as in Manufacturing Example 2 above and 752.3 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 50° C. for 24 hours with reflux, and then PEG-1B was obtained.

Manufacturing Example 113

Preparation of polymer PEG-1C




[0218] Polymer 1C prepared in the same manner as in Manufacturing Example 3 above and 858.7 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried

out at 50° C. for 24 hours with reflux, and then PEG-1C was obtained.

Manufacturing Example 114


Preparation of polymer PEG-1D



[0219] Polymer 1D prepared in the same manner as in Manufacturing Example 4 above and 858.7 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 50° C. for 24 hours with reflux, and then PEG-1D was obtained.

Manufacturing Example 115

Preparation of polymer PEG-1E




[0220] Polymer 1E prepared in the same manner as in Manufacturing Example 5 above and 563.0 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 2500 mg/ml and added to a flask. The reaction was carried

out at 50° C. for 24 hours with reflux, and then PEG-1E was obtained.

Manufacturing Example 116


Preparation of polymer PEG-2A




[0221] Polymer 2A prepared in the same manner as in Manufacturing Example 6 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-2A was obtained.

Manufacturing Example 117

Preparation of polymer PEG-2B






[0222] Polymer 2B prepared in the same manner as in Manufacturing Example 7 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-2B was obtained.

Manufacturing Example 118


Preparation of polymer PEG-2C



[0223] Polymer 2C prepared in the same manner as in Manufacturing Example 8 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-2C was obtained.

Manufacturing Example 119

Preparation of polymer PEG-2D





[0224] Polymer 2D prepared in the same manner as in Manufacturing Example 9 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000

mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-2D was obtained.

Manufacturing Example 120


Preparation of polymer PEG-2E






[0225] Polymer 2E prepared in the same manner as in Manufacturing Example 10 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-2E was obtained.


Manufacturing Example 121



[0226] Polymer 3A prepared in the same manner as in Manufacturing Example 11 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-3A was obtained.

Manufacturing Example 122






[0227] Polymer 3B prepared in the same manner as in Manufacturing Example 12 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-3B was obtained.

Manufacturing Example 123


Preparation of polymer PEG-3C




[0228] Polymer 3C prepared in the same manner as in Manufacturing Example 13 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-3C was obtained.

Manufacturing Example 124

Preparation of polymer PEG-3D






[0229] Polymer 3D prepared in the same manner as in Manufacturing Example 14 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-3D was obtained.

Manufacturing Example 125


Preparation of polymer PEG-3E




[0230] Polymer 3E prepared in the same manner as in Manufacturing Example 15 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-3E was obtained.

Manufacturing Example 126

Preparation of polymer PEG-4A






[0231] Polymer 4A prepared in the same manner as in Manufacturing Example 16 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-4A was obtained.

Manufacturing Example 127


Preparation of polymer PEG-4B




[0232] Polymer 4B prepared in the same manner as in Manufacturing Example 17 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-4B was obtained.

Manufacturing Example 128

Preparation of polymer PEG-4C






[0233] Polymer 4C prepared in the same manner as in Manufacturing Example 18 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-4C was obtained.

Manufacturing Example 129


Preparation of polymer PEG-4D




[0234] Polymer 4D prepared in the same manner as in Manufacturing Example 19 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-4D was obtained.

Manufacturing Example 130

Preparation of polymer PEG-4E




-continued



[0235] Polymer 4E prepared in the same manner as in Manufacturing Example 20 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-4E was obtained.

Manufacturing Example 131

Preparation of polymer PEG-5A




[0236] Polymer 5A prepared in the same manner as in Manufacturing Example 21 above and 648.8 mg of PEG-acrylate were dissolved in CH₂Cl₂ to a concentration of 1000 mg/ml and added to a flask. The reaction was carried

out at 50° C. for 24 hours with reflux, and then PEG-5A was obtained.

Manufacturing Example 132


Preparation of polymer PEG-5B



[0237] Polymer 5B prepared in the same manner as in Manufacturing Example 22 above and 752.3 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 50° C. for 24 hours with reflux, and then PEG-5B was obtained.

Manufacturing Example 133


Preparation of polymer PEG-5C



[0238] Polymer 5C prepared in the same manner as in Manufacturing Example 23 above and 858.7 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 50° C. for 24 hours with reflux, and then PEG-5C was obtained.

Manufacturing Example 134

Preparation of polymer PEG-5D




[0239] Polymer 5D prepared in the same manner as in Manufacturing Example 24 above and 858.7 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 1000 mg/ml and added to a flask. The reaction was carried


out at 50° C. for 24 hours with reflux, and then PEG-5D was obtained.

Manufacturing Example 135

Preparation of polymer PEG-5E



-continued




[0240] Polymer 5E prepared in the same manner as in Manufacturing Example 25 above and 563.0 mg of PEG-acrylate were dissolved in CH_2Cl_2 to a concentration of 2500 mg/ml and added to a flask. The reaction was carried

out at 50° C. for 24 hours with reflux, and then PEG-5E was obtained.

Manufacturing Example 136


Preparation of polymer PEG-6A




[0241] Polymer 6A prepared in the same manner as in Manufacturing Example 26 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-6A was obtained.

Manufacturing Example 137

Preparation of polymer PEG-6B




-continued




[0242] Polymer 6B prepared in the same manner as in Manufacturing Example 27 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-6B was obtained.

Manufacturing Example 138


Preparation of polymer PEG-6C

[0243] Polymer 6C prepared in the same manner as in Manufacturing Example 28 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-6C was obtained.

Manufacturing Example 139

Preparation of polymer PEG-6D


-continued



[0244] Polymer 6D prepared in the same manner as in Manufacturing Example 29 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-6D was obtained.


Manufacturing Example 140

Preparation of polymer PEG-6E




[0245] Polymer 6E prepared in the same manner as in Manufacturing Example 30 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-6E was obtained.

Manufacturing Example 141




[0246] Polymer 7A prepared in the same manner as in Manufacturing Example 31 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-7A was obtained.

Manufacturing Example 142




[0247] Polymer 7B prepared in the same manner as in Manufacturing Example 32 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-7B was obtained.

Manufacturing Example 143




[0248] Polymer 7C prepared in the same manner as in Manufacturing Example 33 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-7C was obtained.

Manufacturing Example 144




[0249] Polymer 7D prepared in the same manner as in Manufacturing Example 34 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-7D was obtained.

Manufacturing Example 145



[0250] Polymer 7E prepared in the same manner as in Manufacturing Example 35 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-7E was obtained.


Manufacturing Example 146



[0251] Polymer 8A prepared in the same manner as in Manufacturing Example 36 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-8A was obtained.

Manufacturing Example 147


Preparation of polymer PEG-8B




[0252] Polymer 8B prepared in the same manner as in Manufacturing Example 37 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-8B was obtained.

Manufacturing Example 148

Preparation of polymer PEG-8C



-continued



[0253] Polymer 8C prepared in the same manner as in Manufacturing Example 38 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-8C was obtained.

Manufacturing Example 149



mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-8D was obtained.

Manufacturing Example 150

Preparation of Polymer PEG-8E

[0255] Polymer 8E prepared in the same manner as in Manufacturing Example 40 above and 563.0 mg of PEG-

Preparation of polymer PEG-8D




[0254] Polymer 8D prepared in the same manner as in Manufacturing Example 39 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000

acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-8E was obtained.

Manufacturing Example 151


Preparation of polymer PEG-9A



[0256] Polymer 9A prepared in the same manner as in Manufacturing Example 41 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-9A was obtained.


Manufacturing Example 152

Preparation of polymer PEG-9B




[0257] Polymer 9B prepared in the same manner as in Manufacturing Example 42 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-9B was obtained.

Manufacturing Example 153




[0258] Polymer 9C prepared in the same manner as in Manufacturing Example 43 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-9C was obtained.

Manufacturing Example 154




[0259] Polymer 9D prepared in the same manner as in Manufacturing Example 44 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-9D was obtained.

Manufacturing Example 155



[0260] Polymer 9E prepared in the same manner as in Manufacturing Example 45 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-9E was obtained.


Manufacturing Example 156



[0261] Polymer 10A prepared in the same manner as in Manufacturing Example 46 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-10A was obtained.

Manufacturing Example 157


Preparation of polymer PEG-10B




[0262] Polymer 10B prepared in the same manner as in Manufacturing Example 47 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-10B was obtained.

Manufacturing Example 158

Preparation of polymer PEG-10C




-continued



[0263] Polymer 10C prepared in the same manner as in Manufacturing Example 48 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-10C was obtained.

Manufacturing Example 159


Preparation of polymer PEG-10D



[0264] Polymer 10D prepared in the same manner as in Manufacturing Example 49 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-10D was obtained.

Manufacturing Example 160


Preparation of polymer PEG-10E




[0265] Polymer 10E prepared in the same manner as in Manufacturing Example 50 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-10E was obtained.

Manufacturing Example 161

Preparation of polymer PEG-11A






[0266] Polymer 11A prepared in the same manner as in Manufacturing Example 51 above and 648.8 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-11A was obtained.

Manufacturing Example 162


Preparation of polymer PEG-11B




[0267] Polymer 11B prepared in the same manner as in Manufacturing Example 52 above and 752.3 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-11B was obtained.

Manufacturing Example 163

Preparation of polymer PEG-11C






[0268] Polymer 11C prepared in the same manner as in Manufacturing Example 53 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-11C was obtained.

Manufacturing Example 164


Preparation of polymer PEG-11D




[0269] Polymer 11D prepared in the same manner as in Manufacturing Example 54 above and 858.7 mg of PEG-acrylate were dissolved in DMSO to a concentration of 1000 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-11D was obtained.

Manufacturing Example 165

Preparation of polymer PEG-11E






[0270] Polymer 11E prepared in the same manner as in Manufacturing Example 55 above and 563.0 mg of PEG-acrylate were dissolved in DMSO to a concentration of 2500 mg/ml and added to a flask. The reaction was carried out at 90° C. for 24 hours, and then PEG-11E was obtained.

Manufacturing Example 166


Preparation of polymer PEG-1F



[0271] Polymer 1F prepared in the same manner as in Manufacturing Example 56 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH₂Cl₂ and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-1F was obtained.


Manufacturing Example 167

Preparation of polymer PEG-1G




[0272] Polymer 1G prepared in the same manner as in Manufacturing Example 57 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-1G was obtained.

Manufacturing Example 168



[0273] Polymer 1H prepared in the same manner as in Manufacturing Example 58 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-1H was obtained.


Manufacturing Example 169



[0274] Polymer 1I prepared in the same manner as in Manufacturing Example 59 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-1I was obtained.

Manufacturing Example 170


Preparation of polymer PEG-1J



[0275] Polymer 1J prepared in the same manner as in Manufacturing Example 60 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-1J was obtained.

Manufacturing Example 171


Preparation of polymer PEG-2F



[0276] Polymer 2F prepared in the same manner as in Manufacturing Example 61 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-2F was obtained.

Manufacturing Example 172


Preparation of polymer PEG-2G



[0277] Polymer 2G prepared in the same manner as in Manufacturing Example 62 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-2G was obtained.

Manufacturing Example 173


Preparation of polymer PEG-2H



[0278] Polymer 2H prepared in the same manner as in Manufacturing Example 63 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-2H was obtained.

Manufacturing Example 174


Preparation of polymer PEG-2I



[0279] Polymer 2I prepared in the same manner as in Manufacturing Example 64 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-2I was obtained.

Manufacturing Example 175


Preparation of polymer PEG-2J



[0280] Polymer 2J prepared in the same manner as in Manufacturing Example 65 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-2J was obtained.

Manufacturing Example 176


Preparation of polymer PEG-3F



[0281] Polymer 3F prepared in the same manner as in Manufacturing Example 66 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-3F was obtained.

Manufacturing Example 177


Preparation of polymer PEG-3G



[0282] Polymer 3G prepared in the same manner as in Manufacturing Example 67 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-3G was obtained.

Manufacturing Example 178


Preparation of polymer PEG-3H



[0283] Polymer 3H prepared in the same manner as in Manufacturing Example 68 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-3H was obtained.


Manufacturing Example 179

Preparation of polymer PEG-3I




[0284] Polymer 3I prepared in the same manner as in Manufacturing Example 69 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-3I was obtained.

Manufacturing Example 180




[0285] Polymer 3J prepared in the same manner as in Manufacturing Example 70 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-3J was obtained.

Manufacturing Example 181




[0286] Polymer 4F prepared in the same manner as in Manufacturing Example 71 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-4F was obtained.

Manufacturing Example 182



[0287] Polymer 4G prepared in the same manner as in Manufacturing Example 72 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-4G was obtained.


Manufacturing Example 183



[0288] Polymer 4H prepared in the same manner as in Manufacturing Example 73 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-4H was obtained.

Manufacturing Example 184


Preparation of polymer PEG-4I



[0289] Polymer 4I prepared in the same manner as in Manufacturing Example 74 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-4I was obtained.

Manufacturing Example 185


Preparation of polymer PEG-4J



[0290] Polymer 4J prepared in the same manner as in Manufacturing Example 75 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-4J was obtained.

Manufacturing Example 186


Preparation of polymer PEG-5F



[0291] Polymer 5F prepared in the same manner as in Manufacturing Example 76 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-5F was obtained.

Manufacturing Example 187


Preparation of polymer PEG-5G



[0292] Polymer 5G prepared in the same manner as in Manufacturing Example 77 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-5G was obtained.

Manufacturing Example 188


Preparation of polymer PEG-5H



[0293] Polymer 5H prepared in the same manner as in Manufacturing Example 78 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-5H was obtained.

Manufacturing Example 189


Preparation of polymer PEG-5I



[0294] Polymer 5I prepared in the same manner as in Manufacturing Example 79 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-5I was obtained.

Manufacturing Example 190


Preparation of polymer PEG-5J




[0295] Polymer 5J prepared in the same manner as in Manufacturing Example 80 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-5J was obtained.

Manufacturing Example 191

Preparation of polymer PEG-6F




-continued




[0296] Polymer 6F prepared in the same manner as in Manufacturing Example 81 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-6F was obtained.

Manufacturing Example 192


Preparation of polymer PEG-6G

[0297] Polymer 6G prepared in the same manner as in Manufacturing Example 82 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-6G was obtained.

Manufacturing Example 193

Preparation of polymer PEG-6H


-continued



[0298] Polymer 6H prepared in the same manner as in Manufacturing Example 83 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-6H was obtained.

Manufacturing Example 194

Preparation of polymer PEG-6I





[0299] Polymer 6I prepared in the same manner as in Manufacturing Example 84 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to

a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-6I was obtained.

[0300] Manufacturing Example 195


Preparation of polymer PEG-6J






[0301] Polymer 6J prepared in the same manner as in Manufacturing Example 85 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-6J was obtained.

Manufacturing Example 196




[0302] Polymer 7F prepared in the same manner as in Manufacturing Example 86 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-7F was obtained.

Manufacturing Example 197




[0303] Polymer 7G prepared in the same manner as in Manufacturing Example 87 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-7G was obtained.

Manufacturing Example 198




[0304] Polymer 7H prepared in the same manner as in Manufacturing Example 88 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-7H was obtained.

Manufacturing Example 199




[0305] Polymer 7I prepared in the same manner as in Manufacturing Example 89 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-7I was obtained.

Manufacturing Example 200



[0306] Polymer 7J prepared in the same manner as in Manufacturing Example 90 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-7J was obtained.


Manufacturing Example 201



[0307] Polymer 8F prepared in the same manner as in Manufacturing Example 91 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-8F was obtained.

Manufacturing Example 202


Preparation of polymer PEG-8G




[0308] Polymer 8G prepared in the same manner as in Manufacturing Example 92 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-8G was obtained.

Manufacturing Example 203

Preparation of polymer PEG-8H




-continued



[0309] Polymer 8H prepared in the same manner as in Manufacturing Example 93 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-8H was obtained.


Manufacturing Example 204

Preparation of polymer PEG-8I

[0310] Polymer 8I prepared in the same manner as in Manufacturing Example 94 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-8I was obtained.

Manufacturing Example 205


Preparation of polymer PEG-8J



[0311] Polymer 8J prepared in the same manner as in Manufacturing Example 95 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-8J was obtained.

Manufacturing Example 206


Preparation of polymer PEG-9F



[0312] Polymer 9F prepared in the same manner as in Manufacturing Example 96 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-9F was obtained.

Manufacturing Example 207


Preparation of polymer PEG-9G



[0313] Polymer 9G prepared in the same manner as in Manufacturing Example 97 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-9G was obtained.

Manufacturing Example 208


Preparation of polymer PEG-9H



[0314] Polymer 9H prepared in the same manner as in Manufacturing Example 98 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-9H was obtained.

Manufacturing Example 209


Preparation of polymer PEG-9I



[0315] Polymer 9I prepared in the same manner as in Manufacturing Example 99 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-9I was obtained.

Manufacturing Example 210


Preparation of polymer PEG-9J



[0316] Polymer 9J prepared in the same manner as in Manufacturing Example 100 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-9J was obtained.

Manufacturing Example 211


Preparation of polymer PEG-10F




[0317] Polymer 10F prepared in the same manner as in Manufacturing Example 101 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-10F was obtained.

Manufacturing Example 212

Preparation of polymer PEG-10G




-continued



[0318] Polymer 10G prepared in the same manner as in Manufacturing Example 102 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-10G was obtained.


Manufacturing Example 213

Preparation of polymer PEG-10H

[0319] Polymer 10H prepared in the same manner as in Manufacturing Example 103 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-10H was obtained.

Manufacturing Example 214


Preparation of polymer PEG-10I




[0320] Polymer 10I prepared in the same manner as in Manufacturing Example 104 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-10I was obtained.

Manufacturing Example 215

Preparation polymer PEG-10J






[0321] Polymer 10J prepared in the same manner as in Manufacturing Example 105 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-10J was obtained.

Manufacturing Example 216


Preparation of polymer PEG-11F




[0322] Polymer 11F prepared in the same manner as in Manufacturing Example 106 above and 329.5 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-11F was obtained.

Manufacturing Example 217

Preparation of polymer PEG-11G





-continued




[0323] Polymer 11G prepared in the same manner as in Manufacturing Example 107 above and 383.0 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-11G was obtained.

Manufacturing Example 218

Preparation of polymer PEG-11H




Preparation of polymer PEG-11I



[0325] Polymer 11I prepared in the same manner as in Manufacturing Example 109 above and 438.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-11I was obtained.

Manufacturing Example 220


Preparation of polymer PEG-11J



[0326] Polymer 11J prepared in the same manner as in Manufacturing Example 110 above and 285.3 mg of PEG-alkynoate were dissolved in 4 mL of CH_2Cl_2 and added to a flask. The reaction was carried out at room temperature for 24 hours, and then PEG-11J was obtained.

Manufacturing Example 221

Preparation of polymer b3A





[0327] Amine monomer 3, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, dimethylformamide (N,N-dimethylformamide, DMF) was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150

mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b3A was obtained. — used in the reaction scheme in the present specification is a notation that is well known to those of ordinary skill in the art, indicating that a multi-step reaction occurs and that intermediates are omitted.

Manufacturing Example 222

Preparation of polymer b3B






[0328] Amine monomer 3, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b3B was obtained.


Manufacturing Example 223

[0329] Amine monomer 3, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b3C was obtained.

Preparation of polymer b3C




Manufacturing Example 224

Preparation of polymer b3D


[0330] Amine monomer 3, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90°C for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b3D was obtained.

Manufacturing Example 225

Preparation of polymer b3E


Manufacturing Example 227

Preparation of polymer b4B


[0333] Amine monomer 4, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b4B was obtained.

Manufacturing Example 228

Preparation of polymer b4C

-continued




[0334] Amine monomer 4, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b4C was obtained.


Manufacturing Example 229

[0335] Amine monomer 4, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b4D was obtained.

Preparation of polymer b4D




Manufacturing Example 230


Preparation of polymer b4E

[0336] Amine monomer 4, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b4E was obtained.

Manufacturing Example 231


Preparation of polymer b7A



[0337] Amine monomer 7, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b7A was obtained.

Manufacturing Example 232

Preparation of polymer b7B




[0338] Amine monomer 7, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an


end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b7B was obtained.

Manufacturing Example 233

Preparation of polymer b7C



-continued




[0339] Amine monomer 7, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b7C was obtained.


Manufacturing Example 234

[0340] Amine monomer 7, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b7D was obtained.

Preparation of polymer b7D




Manufacturing Example 235

Preparation of polymer b7E


[0341] Amine monomer 7, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b7E was obtained.

Manufacturing Example 236

Preparation of polymer b8A

-continued




[0342] Amine monomer 8, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b8A was obtained.

Manufacturing Example 237


[0343] Amine monomer 8, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b8B was obtained.

Preparation of polymer b8B



Manufacturing Example 238

Preparation of polymer b8C




[0344] Amine monomer 8, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an


end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b8C was obtained.

Manufacturing Example 239

Preparation of polymer b8D




Manufacturing Example 241

Preparation of polymer b9A


[0347] Amine monomer 9, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b9A was obtained.

Manufacturing Example 242

Preparation of polymer b7B

-continued




[0348] Amine monomer 9, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b9B was obtained.

Manufacturing Example 243


[0349] Amine monomer 9, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b9C was obtained.

Preparation of polymer b9C



Manufacturing Example 244

Preparation of polymer b9D





[0350] Amine monomer 9, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an

end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b9D was obtained.

Manufacturing Example 245


Preparation of polymer b9E





[0351] Amine monomer 9, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b9E was obtained.


Manufacturing Example 246



[0352] Amine monomer 10, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b10A was obtained.

Manufacturing Example 247

Preparation of polymer b10B




[0353] Amine monomer 10, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b10B was obtained.

Manufacturing Example 248


[0354] Amine monomer 10, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b10C was obtained.

Manufacturing Example 249



[0355] Amine monomer 10, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b10D was obtained.

Manufacturing Example 250




[0356] Amine monomer 10, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b10E was obtained.

Manufacturing Example 251

[0357] Amine monomer 11, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b11A was obtained.

Manufacturing Example 252


Preparation of polymer b11B



[0358] Amine monomer 11, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b11B was obtained.

Manufacturing Example 253


Preparation of polymer b11C



[0359] Amine monomer 11, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b11C was obtained.


Manufacturing Example 254

Preparation of polymer b11D



[0360] Amine monomer 11, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b11D was obtained.


Manufacturing Example 255



[0361] Amine monomer 11, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b11E was obtained.

Manufacturing Example 256


Preparation of polymer b12A



[0362] Amine monomer 12, acrylate monomer A, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b12A was obtained.


Manufacturing Example 257

Preparation of polymer b12B.



[0363] Amine monomer 12, acrylate monomer B, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b12B was obtained.

Manufacturing Example 258




[0364] Amine monomer 12, acrylate monomer C, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b12C was obtained.

Manufacturing Example 259

[0365] Amine monomer 12, acrylate monomer D, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b12D was obtained.

Manufacturing Example 260



[0366] Amine monomer 12, acrylate monomer E, and branched dendrimer were added to a round bottom flask in a molar ratio of 0.4:1:0.2. Thereafter, DMF was added to the flask so that the concentration was 150 mg/ml, and the reaction was carried out at 90° C. for 48 hours. Thereafter, the mixture was cooled to room temperature, and then an end-capping agent was dissolved in DMF to a concentration of 150 mg/ml and added in a molar ratio of 2, and the reaction was carried out for 24 hours at room temperature, and then b12E was obtained.

Example 1

Preparation of Polymeric Nanoparticles (PNP)

[0367] mRNA was added to the polymer prepared by any one of Manufacturing Examples 1 to 260 above, and polymeric nanoparticles (PNP), which are nucleic acid delivery vehicles of the present invention, were prepared by the following method.

[0368] The polymer prepared by any one of Manufacturing Example 1 to 260 above was diluted in a pH 5.0, 25 mM sodium acetate buffer at a ratio for each polymer. Thereafter, 6 μ l of the polymer solution and 6 μ l of the mRNA solution having a concentration of 1 mg/ml were mixed and then stabilized at 4° C. for 30 minutes. Thereafter, the mixture was diluted with 38 μ l of PBS to make a total volume of 50 μ l, and used in the following test examples.

Test Example 1

Analysis of Synthesized Polymer

[0369] The polymers obtained by the methods of Manufacturing Example 1 to 260 above were analyzed by the following methods.

1.1 ¹H-NMR

[0370] ¹H-NMR analysis was performed on the synthesized polymer using deuterated chloroform (CDCl₃) solvent, and the results are shown in FIGS. 2A to 16B. As shown in FIGS. 2A to 16B, it was confirmed that all polymers were prepared normally.

1.2 Hydrophobic Gel Permeable Chromatography

[0371] The solvent was removed from the synthesized polymer, and then analysis was performed using hydrophobic gel permeable chromatography (hydrophobic GPC) by dissolving the polymer in tetrahydrofuran (THF) solvent at a concentration of 1 to 3 mg/ml.

Test Example 2

Analysis of Polymeric Nanoparticle (PNP) Delivery Vehicle

[0372] The polymeric nanoparticle (PNP) prepared in the same manner as in Example 1 above was analyzed by the following method.

2.1 Agarose Gel Electrophoresis

[0373] Agarose gel electrophoresis was performed on the polymers having pH sensitivity by adjusting the N/P ratio (the ratio of amine; N contained in the polymer and the ratio of phosphate; P contained in mRNA) of mRNA and polymer. The pH of the 25 mM NaOAc buffer was set to 5.0, and

5 μ l of the solution in which the polymer was dissolved according to the N/P ratio was mixed with 5 μ l of mRNA at a concentration of 200 ng/5 μ l, and then gently vortexed. Thereafter, the mixture was stabilized at 4° C. for 30 minutes, loaded onto a 1 w/v % agarose gel containing GelRed, and electrophoresed at 100 V for 10 minutes. As a result of the experiment, as shown in FIGS. 17A to 17C, when a complex is formed at different N/P (Polymer/mRNA) ratios for each polymer, it can be confirmed that the mRNA band is not formed in the gel but is caught in the inlet. For example, the 6B polymer completely formed a complex starting from an N/P ratio of 4.

2.2 Zeta Potential and Dynamic Light Scattering Measurement

[0374] For the polymers having pH sensitivity, the N/P ratio for each polymer was adjusted, and 6 μ l of the solution dissolved in a pH 5.0, 25 mM NaOAc buffer was mixed with 6 μ l of luciferase mRNA (FLuc) of SEQ ID NO: 1 at a concentration of 1 mg/ml, and then gently vortexed. Thereafter, the mixture was stabilized at 4° C. for 30 minutes, and 38 μ l of PBS was added to make a final volume of 50 μ l.

[0375] The zeta potential and size of the polymer-mRNA complex were measured by diluting the complex solution described above 10 times in distilled water and using a zeta potential meter and a dynamic light scattering (DLS) device. As shown in FIG. 18, it was confirmed that the zeta potential changed depending on the pH. Through this, it was confirmed that the mRNA complex was formed due to a positive charge at low pH, and it was confirmed that it could have low toxicity and high mRNA delivery efficiency by changing to neutral in the pH in the body.

Test Example 3

Confirmation of mRNA Expression In Vivo of Polymeric Nanoparticles (PNP)

[0376] The polymeric nanoparticles prepared according to Example 1 above were injected intramuscularly into animals to confirm the expression of mRNA.

3.1 Confirmation of In Vivo mRNA Expression

[0377] A representative polymer was synthesized according to the method described in the above manufacturing example, and polymeric nanoparticles were prepared according to Manufacturing Example 261, and mRNA expression level screening was performed. The method for forming the polymer-mRNA complex used for in vivo injection to confirm the mRNA expression is as follows. The weight ratio of each polymer having pH sensitivity and a micelle structure was adjusted, and 6 μ l of the solution dissolved in a pH 5.0, 25 mM NaOAc buffer was mixed with 6 μ l of FLuc of SEQ ID NO: 1 at a concentration of 1 mg/ml, and then gently vortexed. Thereafter, the mixture was stabilized at 4° C. for 30 minutes, and 38 μ l of PBS was added to make a final volume of 50 μ l. Regarding the preparation of lipid nanoparticles (LNPs) used as a comparative group, the following preparation method was used. The molar ratio of ionizable lipid (SM-102):DSPC:cholesterol:PEG-lipid was set to 50:10:38.5:1.5 and dissolved in ethanol. Thereafter, the mRNA diluted in a 25 mM, pH 5.0 sodium acetate buffer was mixed with the mixture dissolved in ethanol so that the volume ratio was 3:1 and the N/P ratio was 5.67. Thereafter, Centrifugation was performed at least three times at 1000 g for 5 minutes with PBS added using a 100 kDa Amicon. Finally, the mRNA was diluted with PBS so that the concentration was 6 μ g/50 μ l.

TABLE 4

SEQ ID NO	RNA sequence
1	ATGGAGGACGCCAAGAACATCAAGAAGGGCCCCGCCCTTCTACCCCTGGAGG ACGGCACCGCGCGAGCAGCTGCACAAGGCCATGAAGCGGTACGCCCTGGTGC CCGGCACCATCGCCTTCAACGACGCCACATCGAGGTGGACATCACCTACGCCGA GTACTTCGAGATGAGCGTGCCTGGCCGAGGCCATGAAGCGGTACGCCCTGAAC ACCAACCACCGGATCGTGGTGTGCAGCGAGAAGAGCCTGCAGTTCTTCATGCCCG TGCTGGGCGCCTGTTTCATCGCGGTGGCCGTGGCCCCGCCCAACGACATCTACAA CGAGCGGGAGGTGCTGAACAGCATGGGCATCAGCCAGCCACCGTGGTGTCTGTG AGCAAGAAGGGCTGCAGAAGATCCTGAACGTGCAGAAGAAGCTGCCCATCATC CAGAAGATCATCATCATGGACAGCAAGACCGACTACCAGGGCTTCCAGAGCATGT ACACCTTCGTGACCAGCCACCTGCCCCCGGCTTCAACGAGTACGACTTCGTGCC GAGAGCTTCGACCGGACAAAGCCATCGCCCTGATCATGAACAGCAGCGGCAGC ACCGGCTTGCCTCAAGGGCGTGGCCCTGCCCCACCGGACCGCCTGCGTGGGTTCA GCCACGCCCGGACCCCATCTTCGGCAACAGATCATCCCGACACCGCCATCCT GAGCGTGGTGGCTTCCACCACGGCTTCGGCATGTTACCCACCTGGGCTACCTGA TCTGCGGCTTCCGGGTGGTGTGATGTACCGGTTTCAGGAGGAGCTGTCTCTGCG GAGCCTGCAGACTACAAGATCCAGAGCGCCTGCTGGTGCACCCCTGTTCAGC TTCTTCGCCAAGAGCACCTGATCGACAAGTACGACCTGAGCAACCTGCACGAGA TCGCCAGCGCGCGGCCGCCCTGAGCAAGGAGGTGGGCGAGGCGTGGCCAAGC GGTTCACCTGCCCGCATCCGCGAGGGCTACGGCTGACCGAGACCAACAGCGC CATCCTGATCACCCCCGAGGGCGACGACAAGCCGCGCGCGTGGGCAAGGTGGT GCCCTTCTTCGAGGCCAAGGTGGTGGACCTGGACACCGGCAAGACCTGGGCGTG AACCAGCGGGCGAGCTGTGCGTGGCGGCCCATGATCATGAGCGGTACGTG AACAAACCCGAGGCCACCAACGCCCTGATCGACAAGGACGGCTGGTGCACAGC GCGGACATCGCCTACTGGGACGAGGACGAGCACTTCTTCATCGTGGACCGGCTGA AGAGCCTGATCAAGTACAAGGGCTACAGGTGGCCCCCGCGAGCTGGAGAGCA TCCTGTGTCAGCACCCCAACATCTTCGACGCGCGGTGGCCGGCTGCCCGACGA CGAGCGCGCGAGCTGCCCGCGCGGTGGTGTGCTGGAGCACGGCAAGACCAT GACCGAGAGGAGATCGTGGACTACGTGGCCAGCCAGGTGACCAACGCCAAGAA GCTGCGGGGCGCGGTGGTGTCTGTCGACGAGGTGCCCAAGGGCTGACCGGCAA GCTGGACGCCCGGAAGATCCGGGAGATCCTGATCAAGGCCAAGAAGGGCGGCAA GATCGCCGTGTA

[0378] In order to confirm in vivo mRNA expression using the composition above, luciferase mRNA (FLuc) of SEQ ID NO: 1 was injected into the gastrocnemius muscle of the mouse hind leg at a concentration of 6 µg/50 µl. Thereafter, mRNA expression was confirmed after 24 hours. 200 µl of luciferin was injected intraperitoneally into each individual at a concentration of 15 mg/ml. 10 minutes after luciferin injection, mRNA expression was confirmed with an exposure time of 15 seconds.

[0379] As a result, as shown in Table 5 below, when delivering mRNA using the polymers according to the present invention, it was confirmed that mRNA expression was equivalent or higher than that of free mRNA (in the present specification, free mRNA refers to mRNA without a delivery vehicle), and it was confirmed that among them, 22 polymers showed higher mRNA expression than lipid nanoparticle (LNP) delivery vehicles. In addition, experiments using representative branched polymers also showed a high mRNA expression rate compared to free mRNA. Therefore, it can be seen that the delivery vehicle of the present invention can be used as an mRNA delivery vehicle.

TABLE 5

Manufacturing Example	Item No.	mRNA expression efficiency
1	1A	3.61
3	1C	9.56
4	1D	6.39
5	1E	5.39
6	2A	2.02
9	2D	8.37

TABLE 5-continued

Manufacturing Example	Item No.	mRNA expression efficiency
10	2E	27.79
11	3A	6.58
13	3C	17.81
14	3D	1.07
15	3E	10.19
16	4A	1.76
17	4B	23.72
18	4C	58.63
19	4D	38.72
20	4E	16.00
21	5A	2.24
27	6B	41.22
29	6D	21.58
30	6E	67.60
31	7A	36.43
32	7B	37.83
33	7C	11.28
34	7D	62.72
35	7E	6.29
36	8A	29.36
37	8B	78.62
38	8C	333.17
39	8D	50.90
40	8E	32.18
41	9A	2.12
42	9B	27.33
43	9C	92.24
44	9D	160.44
45	9E	18.38
50	10E	2.54
51	11A	36.90
53	11C	6.14
55	11E	2.43

TABLE 5-continued

Manufacturing Example	Item No.	mRNA expression efficiency
222	b3B	13.47
223	b3C	1.45
225	b3E	1.24
227	b4B	658.79
228	b4C	100.52
229	b4D	115.61
235	b7E	18.36
238	b8C	218.05
239	b8D	24.59
242	b9B	45.64
243	b9C	3.80
244	b9D	105.24
247	b10B	72.51
248	b10C	18.37
249	b10D	19.68
250	b10E	182.42
252	b11B	867.09
253	b11C	67.04
255	b12E	10.02
256	b12A	98.05
258	b12C	80.08
256	b12D	10.26
Free mRNA	comparative group	1
Lipid nanoparticle	comparative group	12.27

Test Example 4

Confirmation of In Vivo mRNA Expression of Polymeric Nanoparticles (PNP) Comprising PEG

[0380] The mRNA expression efficiency using polymeric nanoparticles (PNP) comprising PEG at both ends was confirmed.

[0381] The polymeric nanoparticles prepared according to Example 1 above were injected intramuscularly into animals to confirm the expression rate of mRNA.

[0382] Eight polymers (PEG-1A, PEG-1B, PEG-2A, PEG-2B, PEG-3A, PEG-3B, PEG-4A, PEG-4B) prepared in the same manner as in the above manufacturing examples was screened for the in vivo mRNA expression efficiency. The method for forming the polymer-mRNA complex used for in vivo injection to confirm the mRNA expression is as follows. The weight ratio of each polymer was adjusted, and 6 μ l of the solution dissolved in a pH 5.0, 25 mM NaOAc buffer was mixed with 6 μ l of luciferase mRNA (FLuc) of SEQ ID NO: 1 at a concentration of 1 mg/ml, and then

gently vortexed. Thereafter, the mixture was stabilized at 4° C. for 30 minutes, and 38 μ l of PBS was added to make a final volume of 50 μ l.

[0383] In order to confirm in vivo mRNA expression using the composition above, it was injected into the gastrocnemius muscle of the mouse hind leg. Thereafter, mRNA expression was confirmed 24 hours after intramuscular injection. 200 μ l of luciferin was injected intraperitoneally into each individual at a concentration of 15 mg/ml. 10 minutes after luciferin injection, mRNA expression was confirmed with an exposure time of 15 seconds.

[0384] As a result, as shown in FIG. 19, PEG was included at both ends of the polymers prepared in Manufacturing Examples 111 to 220 above, and at this time, the mRNA expression of representative polymers was confirmed. As a result, it was confirmed that the expression rate was equal to or higher than that of free mRNA. In addition, it was confirmed that 1B prepared by Manufacturing Example 2 had poor solubility, but PEG-1B, which had PEG attached to both ends, had improved solubility and mRNA expression was significantly increased. Therefore, it can be seen that the delivery vehicle of the present invention can be effectively used as an mRNA delivery vehicle.

Test Example 5

Confirmation of In Vivo Expression Retention Time of Polymeric Nanoparticles (PNP) Comprising PEG

[0385] In order to confirm the expression retention time of mRNA, polymer-mRNA complexes were prepared using representative polymers of the manufacturing examples in the same manner as in Test Example 3, and then injected into the hind legs of mice, and measured for up to 20 days. The results are shown in FIG. 20.

[0386] As shown in FIG. 20, it was confirmed that in the case of lipid nanoparticle-mRNA complex (LNP-mRNA), expression reached its peak at 4 hours and then dropped below the baseline value at about 48 hours, whereas in the case of all polymeric nanoparticle-mRNA complexes used in the test, the expression time was maintained longer than that of lipid nanoparticle-mRNA complexes. In particular, it was confirmed that in the case of 6B and 8C polymer complexes, mRNA expression was maintained for more than 10 days, which means that the administration frequency can be drastically reduced compared to the existing method in the development of a gene therapeutic agent.

SEQUENCE LISTING

```

Sequence total quantity: 1
SEQ ID NO: 1          moltype = RNA  length = 1653
FEATURE              Location/Qualifiers
source                1..1653
                     mol_type = mRNA
                     organism = unidentified

SEQUENCE: 1
atggaggacg ccaagaacat caagaagggc cccgccccct tctacccccct ggaggacggc 60
accgccggcg agcagctgca caaggccatg aagcggtagc ccctgggtgc cggcaccatc 120
gccttcacccg acgcccacat cgagggtggac atcacctacg ccgagtaactt cgagatgagc 180
gtgcgggttg cggaggccat gaagcggtag gccctgaaca ccaaccaccg gatcgtggtg 240
tgcagcgaga acagcctgca gttcttcatg cccgtgctgg gcgccctgtt catcggcgtg 300
gccgtggccc cggccaaaga catctacaac gagcgggagc tgctgaacag catgggcate 360
agccagccca ccgtggtggt cgtgagcaag aagggcctgc agaagatcct gaacgtgcag 420
aagaagctgc ccatcatcca gaagatcatc atcatggaca gcaagaccga ctaccagggc 480
ttccagagca tgtacacett cgtgaccagc caccctgccc cgggcttcaa cgagtacgac 540
ttcgtgcccc agagcttcga cgggacaag accatcgccc tgatcatgaa cagcagcgcc 600

```


-continued

```

agcaccggcc tgcaccaagg cgtggccctg ccccaaccga ccgcctgcgt gcggttcage 660
cacgcccggg accccatctt cggcaaccag atcatcccg acacggccat cctgagcgtg 720
gtgcccttcc accacggctt cggcatgttc accaccctgg gctacctgat ctgcggcttc 780
cgggtggtgc tgatgtaccg gttcgaggag gagctgttcc tgcggagcct gcaggactac 840
aagatccaga gcgcctgtct ggtgcccacc ctgttcagct tcttcgcaa gagcacctg 900
atcgacaagt acgacctgag caacctgcac gagatcgcca gcggcggcgc cccctgagc 960
aaggaggtgg gcgaggccgt ggccaagcgg ttccacctgc ccggcatccg gcagggtac 1020
ggcctgaccg agaccaccag cgccatctct atcaccccg agggcgacga caagcccggc 1080
gcccgtggga aggtggtgcc cttcttcgag gccaaagggt tggacctgga caccggcaag 1140
accctgggcy tgaaccagcg gggcgagctg tgcgtgcggg gcccatgat catgagcggc 1200
tacgtgaaca acccgagggc caccacgccc ctgatcgaca aggacggctg gctgcacagc 1260
ggcgacatcg cctactggga cgaggacgag cactcttcca tcgtggaccg gctgaagagc 1320
ctgatcaagt acaagggcta ccagggtggc ccgcccgcgc tggagagcat cctgtgcag 1380
caccacaaca tcttcgacgc cggcgtggcc ggccctgccc acgacgacgc cggcgagctg 1440
cccgccggcc tgggtggtgt ggagcacggc aagaccatga ccgagaagga gatcgtggac 1500
tacgtggcca gccagggtgac caccgccaag aagctgcggg gcggcgtggt gttcgtggac 1560
gaggtgcccc agggcctgac cggcaagctg gacgcccga agatccggga gatcctgatc 1620
aaggccaaga agggcgggca gatcgccgtg tga 1653

```

1. A polymer suitable for intracellular nucleic acid delivery, comprising a repeating unit of Formula 1 below:



in the formula,

R1 is a bond or a C1-6 alkylene,

R2 and R3 are each independently (i) hydrogen, (ii) C1-6 alkyl which may be substituted with alkylamine or hydroxy, (iii) a 5- or 6-membered carbocycle or heterocycle, or (iv) C1-6 alkyl substituted with a 5- or 6-membered carbocycle or heterocycle,

said R2 and R3 may be linked together to form a C1-6 alkylene,


said 5- or 6-membered carbocycle or heterocycle may be substituted with C1-3 alkyl or C1-3 alkoxyacyl,

X is a nitrogen atom or a bond, and when X is a bond, said R3 is absent,

R4 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and

----- represents a single bond or a double bond.

2. A polymer suitable for intracellular nucleic acid delivery, comprising a repeating unit of Formula 2 below:



in the formula,

R5 is a C1-6 alkylene,














R6 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and

----- represents a single bond or a double bond.





3. The polymer according to claim 1, wherein the repeating unit of Formula 1 is formed by a reaction between any one of the monomers of Formulas 3 to 13 below and any one of the monomers of Formulas 14 to 23 below:

Formula	Structure
3	
4	
5	
6	

-continued







Formula	Structure
7	
8	
9	
10	
11	
12	
13	
14	
15	
16	
17	
18	
19	

-continued






Formula	Structure
20	
21	
22	
23	

4. The polymer according to claim 2, wherein the repeating unit of Formula 2 is formed by a reaction between a





monomer of Formula 24 below and any one of the monomers of Formulas 14 to 23 below:

Formula	Structure
14	
15	
16	
17	
18	
19	

-continued









Formula	Structure
20	
21	
22	
23	
24	

5. A polymer comprising a repeating unit selected from the group consisting of repeating units of the chemical structures set forth in the table below:

Chemical structural formula










-continued

Chemical structural formula










-continued

Chemical structural formula









-continued

Chemical structural formula









-continued

Chemical structural formula








-continued

Chemical structural formula








-continued

Chemical structural formula







-continued

Chemical structural formula








-continued

Chemical structural formula












-continued

Chemical structural formula









-continued

Chemical structural formula









-continued

Chemical structural formula










-continued

Chemical structural formula









-continued

Chemical structural formula








-continued

Chemical structural formula








-continued

Chemical structural formula







-continued

Chemical structural formula







-continued

Chemical structural formula









-continued

Chemical structural formula




-continued

Chemical structural formula








-continued

Chemical structural formula










-continued

Chemical structural formula










-continued

Chemical structural formula









-continued

Chemical structural formula








-continued

Chemical structural formula








-continued

Chemical structural formula







-continued

Chemical structural formula








-continued

Chemical structural formula









-continued

Chemical structural formula




-continued

Chemical structural formula











-continued

Chemical structural formula









-continued

Chemical structural formula




-continued

Chemical structural formula








-continued

Chemical structural formula







-continued

Chemical structural formula




-continued

Chemical structural formula




-continued

Chemical structural formula




-continued

Chemical structural formula







-continued

Chemical structural formula








-continued

Chemical structural formula






-continued

Chemical structural formula



-continued

Chemical structural formula




-continued

The figure displays four chemical structures of poly(2,2,5-trimethyl-1,3-dioxane-5-carboxylic acid) (PTMDC) and its derivatives. The structures are arranged in two rows, each showing a repeating unit of the polymer chain and a corresponding side chain.







- Top Row:** The repeating unit is shown with a side chain consisting of a 4-aminobutyl group linked to a 4-aminobutyl group via an amide bond. The side chain is labeled with a subscript n .
- Bottom Row:** The repeating unit is shown with a side chain consisting of a 4-aminobutyl group linked to a 4-aminobutyl group via an amide bond. The side chain is labeled with a subscript n .

Chemical structural formula










-continued

Chemical structural formula







-continued

Chemical structural formula









-continued

Chemical structural formula




-continued

Chemical structural formula




-continued

Chemical structural formula




Ⓜ indicates text missing or illegible when filed

6. The polymer according to claim 1, wherein the polymer comprises a branched monomer of Formula 25 below:



[Formula 25]

7. The polymer according to claim 1, wherein the polymer comprises a structure of Formula 26 below:



[Formula 26]

hydroxy, (iii) a 5- or 6-membered carbocycle or heterocycle, or (iv) C1-6 alkyl substituted with a 5- or 6-membered carbocycle or heterocycle,

said R2 and R3 may be linked together to form a C1-6 alkylene,

said 5- or 6-membered carbocycle or heterocycle may be substituted with C1-3 alkyl or C1-3 alkoxyacyl,

X is a nitrogen atom or a bond, and when X is a bond, said R3 is absent,

in the formula,


R1 is a bond or a C1-6 alkylene,

R2 and R3 are each independently (i) hydrogen, (ii) C1-6 alkyl which may be substituted with alkylamine or

R4 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and

----- represents a single bond or a double bond.

8. The polymer according to claim 2, wherein the polymer comprises a structure of Formula 27 below:




in the formula,

R5 is a C1-6 alkylene,

R6 is a linear or branched C1-15 alkylene, 1 to 5 carbon atoms of said alkylene may be replaced by oxygen or sulfur, and

— represents a single bond or a double bond.

9. The polymer according to claim 1, wherein one or both ends of the polymer are linked to a PEG-containing moiety comprising a structure of Formula 28 below:



in the formula,

== represents a double bond or a triple bond, and

n is an integer in the range of 1 to 500.

10. A polymeric nanoparticle in which the polymer according to claim 1 and a nucleic acid molecule form a complex.

11. The polymeric nanoparticle according to claim 10, wherein the nucleic acid molecule is RNA.

12. The polymeric nanoparticle according to claim 10, wherein the nucleic acid molecule is DNA.

13. The polymeric nanoparticle according to claim 10, wherein the nucleic acid molecule is isolated in a pH-dependent manner and delivered into cells in the body.

14. The polymeric nanoparticle according to claim 11, wherein when delivered into the body, the expression of the nucleic acid molecule is maintained for a longer period of time compared to when the same nucleic acid molecule is delivered into the body in the form of a non-polymeric lipid nanoparticle.

15. A method for delivering a nucleic acid molecule into a cell using the polymeric nanoparticle according to claim 10.

16. The method according to claim 15, wherein the polymeric nanoparticle is delivered by parenteral, intramuscular, subcutaneous, or intravenous administration.

17. A pharmaceutical composition comprising the polymeric nanoparticle according to claim 10.

18. The pharmaceutical composition according to claim 17, wherein the pharmaceutical composition is a gene therapeutic agent.

19. The pharmaceutical composition according to claim 17, wherein the pharmaceutical composition is a vaccine.

* * * * *