



US012416005B2

(12) **United States Patent**
Cigan et al.(10) **Patent No.:** **US 12,416,005 B2**(45) **Date of Patent:** ***Sep. 16, 2025**(54) **PIG WITH A GENETICALLY MODIFIED
CD163 GENE RESISTANT TO PRRSV**11,208,659 B2 * 12/2021 Cigan A01K 67/0276
11,535,850 B2 * 12/2022 Cigan A01K 67/0275
2014/0068797 A1 3/2014 Doudna et al.(71) Applicant: **Genus PLC**, DeForest, WI (US)

FOREIGN PATENT DOCUMENTS

(72) Inventors: **Andrew Mark Cigan**, Madison, WI (US); **Jonathan Edward Lightner**, Jefferson, WI (US); **Matthew Scott Culbertson**, Wilmington, NC (US); **William Thomas Christianson**, Hendersonville, TN (US); **Benjamin Beaton**, DeForest, WI (US); **Brian Burger**, Madison, WI (US); **Dylan Barnes**, Madison, WI (US); **Matthew Campbell**, Madison, WI (US)WO WO2017023337 2/2017
WO WO 2018073237 * 4/2018
WO WO2019203807 10/2019

OTHER PUBLICATIONS

(73) Assignee: **Genus plc**, DeForest, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 238 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **17/515,139**(22) Filed: **Oct. 29, 2021**(65) **Prior Publication Data**

US 2022/0042017 A1 Feb. 10, 2022

Related U.S. Application Data

(63) Continuation of application No. 17/307,369, filed on May 4, 2021, now Pat. No. 11,208,659.

(60) Provisional application No. 63/021,370, filed on May 7, 2020, provisional application No. 63/020,128, filed on May 5, 2020.

(51) **Int. Cl.****A01K 67/027** (2024.01)
A01K 67/0275 (2024.01)
C12N 9/22 (2006.01)
C12N 15/113 (2010.01)
C12N 15/877 (2010.01)(52) **U.S. Cl.**CPC **C12N 15/113** (2013.01); **A01K 67/0275** (2013.01); **C12N 9/22** (2013.01); **C12N 15/877** (2013.01); **A01K 2217/052** (2013.01); **A01K 2227/108** (2013.01); **C12N 2310/20** (2017.05); **C12N 2320/34** (2013.01)(58) **Field of Classification Search**USPC 800/17
See application file for complete search history.(56) **References Cited**

U.S. PATENT DOCUMENTS

4,873,191 A 10/1989 Wagner et al.
9,820,475 B2 11/2017 Prather et al.
10,091,975 B2 10/2018 Prather et al.
10,827,730 B2 11/2020 Prather et al.Yang, Dongshan, et al., Generation of PPARy mono-allelic knock-out pigs via zinc-finger nucleases and nuclear transfer cloning, *Cell Research*, 2011, 21:979-982.Yoon, In J., et al., Persistent and contact infection in nursery pigs experimentally infected with reproductive and respiratory syndrome (PPRS) virus, *Swine Health and Production*, 1993, 1(4):5-8.Yoshioka, Kojn, et al., Birth of Piglets Derived from Porcine Zygotes Cultured in a Chemically Defined Medium, *Biol of Reprod*, 2002, 66:112-119.Zhang, Qingzhan and Yoo, Dongwan, PRRS virus receptors and their role for pathogenesis, *Veterinary Microbiology*, 2015, 177:229-241.Zhao, Jianguo, et al., Histone Deacetylase Inhibitors Improve In Vitro and In Vivo Development Competence of Somatic Cell Nuclear Transfer Porcine Embryos, *Cellular Reprogramming*, 2010, 12(1):75-83.Zhao, Jianguo, et al., Significant Improvement in Cloning Efficiency of an Inbred Miniature Pig by Histone Deacetylase Inhibitor Treatment after Somatic Cell Nuclear Transfer, *Biology of Reproduction*, 2009, 81:525-530.Abeydeera, L.R. and Day, B.N., Fertilization and subsequent development in vitro of pig oocytes inseminated in a modified tris-buffered medium with frozen-thawed ejaculated spermatozoa, *Biol. Reprod.*, 1997, 57:729-734.Agung et al., In vitro fertilization and development of porcine oocytes matured in follicular fluid, *J Reprod Dev.*, 2013, 59:103-106.Alkan, F., et al., CRISPR-Cas9 off-targeting assessment with nucleic acid duplex energy parameters, *Genome Biol.*, 2018, 19:177.Altschul, S.P., et al., Basic local alignment search tool, *J. Mol. Biol.*, 1990, 215:403-410.Appeltant et al., Porcine oocyte maturation in vitro: role of cAMP and oocyte-secreted factors—A practical approach, *J Reprod Dev.*, 2016, 62:439-449.

(Continued)

Primary Examiner — Michael C Wilson(74) *Attorney, Agent, or Firm* — Daniel M. Lorentzen;
Christopher J. Arnold(57) **ABSTRACT**The present disclosure relates to methods and compositions useful for prevention of porcine reproductive and respiratory syndrome virus (PRRSv) in animals, including animals of the species *Sus scrofa*. The present teachings relate to swine wherein at least one allele of a CD163 gene has been inactivated, and to specific methods and nucleic acid sequences used in gene editing to inactivate the CD163 gene. Swine wherein both alleles of the CD163 gene are inactivated are resistant to porcine reproductive and respiratory syndrome virus (PRRSv). Elite lines comprising homozygous CD163-edited genes retain their superior properties.**3 Claims, No Drawings****Specification includes a Sequence Listing.**

(56)

References Cited

OTHER PUBLICATIONS

- Bae, S., et al., Cas-OFFinder: a fast and versatile algorithm that searches for potential off-target sites of Cas9 RNA-guided endonucleases, *Bioinformatics*, 2014, 30:1473-1475.
- Cameron, P., et al., Mapping porcine landscape of CRISPR-Cas9 cleavage, *Nat Methods* 2017, 14:600-606.
- Chenna, R., et al., Multiple sequence alignment with the Clustal series of programs, *Nucleic Acids Research*, 2003, 31:3497-3500.
- Fowler et al., The production of pig preimplantation embryos in vitro: Current progress and future prospects, *Reprod Biol.*, 2018, 18:203-211.
- Fonfara et al., Phylogeny of Cas9 determines functional exchangeability of dual-RNA and Cas9 among orthologous type II CRISPR-Cas systems, *Nucleic Acids Research*, 2014, 42:2577-2590.
- Gnirke, A., et al., Solution hybrid selection with ultra-long oligonucleotides for massively parallel targeted sequencing, *Nat. Biotechnol.*, 2009, 27:182-189.
- Guo, Chunhe, et al., Highly Efficient Generation of Pigs Harboring a Partial Deletion of the CD163 SRCR5 Domain, Which are Fully Resistant to Porcine Reproductive and Respiratory Syndrome Virus 2 Infection, *Frontiers in immunology*, 2019, 10(1846):1-14.
- Haft, Daniel H., et al., A guild of 45 CRISPR-associated (Cas) protein families and multiple CRISPR/Cas subtypes exist in prokaryotic genomes, *PLoS Comput. Bio.*, 2005, 1(6):474-483.
- Jinek, M., et al., A programmable dual RNA-guided DNA endonuclease in adaptive bacterial immunity, *Science*, 2012, 337:816-821.
- Joung et al., TALENs: a widely applicable technology for targeted genome editing, *Nat. Rev. Mol. Cell. Biol.*, 2013, 14:49-55.
- Kang, Soo Ji, et al., Comparison of Seven Commercial TaqMan Master Mixes and Two Real-Time PCR Platforms Regarding the Rapid Detection of Porcine DNA, *Food Sci Anim Resour*, 2021, 41(1):85-94.
- Karvelis et al., Rapid characterization of CRISPR-Cas9 protospacer adjacent motif sequence elements, *Genome Biology*, 2015, 16:253.
- Larkin Ma et al., Clustal W and Clustal X version 2.0, *Bioinformatics*, 2007, 23:2947-2948.
- Lavitrano et al., Efficient production by sperm-mediated gene transfer of human decay accelerating factor (hDAF) transgenic pigs for xenotransplantation, *Proc. Natl. Acad. Sci. USA*, 2002, 99:14230-14236.
- Lavitrano et al., Sperm-mediated gene transfer, *Reprod. Fert. Develop.*, 2006, 18:19-23.
- Lo, Transformation by iontophoretic microinjection of DNA: multiple integrations without tandem insertions, *Mol. Cell. Biol.*, 1983, 3:1803-1814.
- Onofre, Gabriela et al., Scavenger receptor CD163 and its biological functions, *Acta Medica*, 2009, 52:57-61.
- Ran, F. A., et al., In vivo genome editing using *Staphylococcus aureus* Cas9, *Nature*, 2015, 520:186-191.
- Sapranaukas et al., The *Streptococcus thermophilus* CRISPR/Cas system provides immunity in *Escherichia coli*, *Nucleic Acids Research*, 2011, 39:9275-9282.
- Shah et al., Protospacer recognition motifs: mixed identities and functional diversity, *RNA Biol.*, 2013, 10:891-899.
- Suzuki, C., et al., Effects of amino acid supplements and replacement of polyvinyl alcohol with bovine serum albumin in porcine zygote medium, *Reprod. Fert. Dev.*, 2006 18:789-795.
- Svitashev et al., Genome editing in maize directed by CRISPR-Cas9 ribonucleoprotein complexes, *Nature Communications*, 2016, 7:113274.
- Thompson et al., Germ line transmission and expression of a corrected HPRT gene produced by gene targeting in embryonic stem cells, *Cell*, 1989, 56:313-321.
- Thompson, J.D. and Clustal, W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice et al., *Nucleic Acids Research*, 1994, 22:4673-4680.
- Tsai, S.Q., et al., GUIDE-seg enables genome-wide profiling of off-target cleavage by CRISPR-Cas nucleases, *Nat. Biotechnol.* 2015, 33:187-197.
- Tsai, S.Q., et al., CIRCLE-seq: a highly sensitive in vitro screen for genome-wide CRISPR-Cas9 nuclease off-targets, *Nat. Methods*, 2017, 14:607-614.
- Van Der Putten et al., Efficient insertion of genes into the mouse germ line via retroviral vectors, *Proc. Natl. Acad. Sci. USA*, 1985, 82:6148-1652.
- Wakayama et al., Full-term development of mice from enucleated oocytes injected with cumulus cell nuclei, *Nature*, 1998, 394:369-374.
- Wilmut et al., Viable offspring derived from fetal and adult mammalian cells, *Nature*, 1997, 385:810-813.
- Yoshioka, K., et al., Birth of piglets derived from porcine zygotes cultured in a chemically defined medium, *Biol. Reprod.*, 2002, 60:112-119.
- Yoshioka, K., Defined System for In Vitro Production of Porcine Embryos Using a Single Basic Medium, *J. Reprod. Dev.* 2008, 54:208-213.
- Yoshioka, K., J., Development and application of a chemically defined medium for the in vitro production of porcine embryos, *Reprod. Dev.*, 2011, 57:9-16.
- Zetsche et al., Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system, *Cell*, 2015, 163:759-771.
- Zhang et al., Off-target Effects in CRISPR/Cas9-mediated Genome Engineering, *Mol Ther Nucleic Acids*, 2015, 4:e264.
- Zhao, C., et al., CRISPR-offinder: a CRISPR guide RNA design and off-target searching tool for user-defined protospacer adjacent motif, *Int. J. Biol. Sci.*, 2017, 13:1470-1478.
- Merck Sharp and Dohme Corp., Normal Rectal Temperature Ranges, 2009, pp. I.
- Miao, Yi-hang, et al., Centrosome Abnormalities During Porcine Oocyte Aging, *Environmental and Molecular Mutagenesis*, 2009, 50:666-671.
- Misinzo, Gerald, M., et al., Involvement of proteases in porcine reproductive and respiratory syndrome virus uncoating upon internalization in primary macrophages, *Vet. Res.*, 2008, 39:55-69.
- Molitor, T.W., et al., Immunity to PRRSV: Double-edged sword, *Veterinary Microbiology*, 1997, 55:265-276.
- Morgan, S.B., et al., Pathology and Virus Distribution in the Lung and Lymphoid Tissues of Pigs Experimentally Inoculated with Three Distinct Type 1 PRRS Virus Isolates of Varying Pathogenicity, *Transboundary and Emerging Diseases*, 2014, 63(2016):285-295.
- Murtaugh, M.P., et al., Comparison of the structural protein coding sequences of the VR-2332 and Lelystad virus strains of the PRRS virus, *Arch Virol*, 1995, 140:1451-1460.
- Nath, Deepa, et al., The Amino-terminal Immunoglobulin-like Domain of Sialoadhesin Contains the Sialic Acid Binding Site, *J Biol Chem*, 1995, 270(44):26184-26191.
- Nauwynck, H.J., et al., Entry of porcine reproductive and respiratory syndrome virus into porcine alveolar macrophages via receptor-mediated endocytosis, *J General Virology*, 1999, 80:297-305.
- Nelson, Chris J., et al., Porcine Reproductive and Respiratory Syndrome Virus Comparison: Divergent Evolution on Two Continents, *J Virology*, 1999, 73(1):270-280.
- Neumann, Eric J., et al., Assessment of the economic impact of porcine reproductive and respiratory syndrome on swine production in the United States, *JAVMA*, 2005, 227(3):385-392.
- Nielsen, H.S., et al., Generation of an Infectious Clone of VR-2332, a Highly Virulent North American-Type Isolate of Porcine Reproductive and Respiratory Syndrome Virus, *J Virology*, 2003, 77(6):3702-3711.
- Nielsen, Marianne Jensby, et al., The macrophage scavenger receptor CD163: endocytic properties of cytoplasmic tail variants, *J Leukocyte Biology*, 2006, 79:837-845.
- Niu, Yaya, et al., Generation of Gene-Modified Cynomolgus Monkeys via Cas9/RNA-Mediated Gene Targeting in One-Cell Embryos, *Cell*, 2014, 156:836-843.
- Nussbaum, Diana J. and Prather, Randall S., Differential Effects of Protein Synthesis Inhibitors on Porcine Oocyte Activation, *Mol Reprod and Develop*, 1995, 41:70-75.
- Oetke, Cornelia, et al., Sialoadhesin-Deficient Mice Exhibit Subtle Changes in B- and T-Cell Populations and Reduced Immunoglobulin M Levels, *Mol and Cell Biol*, 2006, 26(4):1549-1557.

(56)

References Cited**OTHER PUBLICATIONS**

- Park, Kwang-Wook, et al., Developmental Potential of Porcine Nuclear Transfer Embryos Derived from Transgenic Fetal Fibroblasts Infected with the Gene for the Green Fluorescent Protein: Comparison of Different Pasion/Activation Conditions, *Biol of Reprod*, 2001, 65:1681-1685.
- Park, Kwang-Wook, et al., Production fo Nuclear Transfer-Derived Swine That Express the Enhanced Green Fluorescent Protein, *Animal Biotech*, 2001, 12(2), 173-181.
- Patience, John F. and Thacker, Phil A., *Nutrition of the Breeding Herd*, Prairie Swine Centre, 1989, pp. 149-171.
- Patton, John B., et al., Modulation of CD163 receptor expression and replication of porcine reproductive and respiratory syndrome virus in porcine macrophages, *Virus Research*, 2009, 140:161-171.
- Plagemann, Peter O.W., *Lactage Dehydrogenase-Elevating Virus and Related Viruses*, Fields Virology, 3rd Ed., 1996, Ch. 36, pp. 1105-1120.
- Popescu, Luca, et al., Genetically edited pigs lacking CD163 show no resistance following infection with the African swine fever virus isolate, Georgia Jan. 2007, *Virology*, 2017, 501:102-106.
- Prather, Randall S., et al., Genetic engineering alveolar macrophages for host resistance to PRRSV, *Veterinary Microbiology*, 2017, 209:124-129.
- Prather, Randall S., et al., Knockout of maternal CD163 protects fetuses from infection with porcine reproductive and respiratory syndrome virus (PRRSV), *Nature Scientific Reports*, 2017, 7(13371):1-5.
- Prather, R.S., et al., Genetic Engineering of Pigs for PRRSV Resistance, *Animal Sciences*, Univ Missouri-Columbia, College of Ag Food and Natural Resources, Animal Reproductive Biology Group, Slides, pp. 1-43.
- Prather, Randall S., et al., An Intact Staloadhesin (Sn/SIGLEC1/CD169) Is Not Required for Attachment/Internalization of the Porcine Reproductive and Respiratory Syndrome Virus, *J Virology*, 2013, 87(17):9538-9546.
- Prather, R.S., Genetic Engineering of Pigs for PRRSV resistance, *Animal Sciences*, Univ Missouri-Columbia, College of Ag Food and Natural Resources, Animal Reproductive Biology Group, Slides, pp. 1-55.
- Prather, R.S., et al., Genetic engineering the pig to better understand PRRSV infection, 2014 North American PRRS Symposium, 2014, p. 33.
- Prather, R.S., Genetic Engineering for Profitable Port Production: Is Resistance to PRRSV Possible?, *Animal Sciences*, Univ Missouri-Columbia, College of Ag Food and Natural Resources, Animal Reproductive Biology Group, Slides, pp. 1-70.
- Prather, Randall S., et al. CRISPR/Cas9-Mediated Genetic Engineering: Is CD163 an Entry Mediator for PRRSV Infection?, *International Plant and Animal Genome Conference XXIII*, Abstract, Jan. 9-15, 2015, 1 page.
- Provost, Chantale, et al., Identification of a new cell line permissive to porcine reproductive and respiratory syndrome virus infection and replication which is phenotypically distinct from MARC-145 cell line, *Virology Journal*, 2012, 9(267):1-14.
- Reed, L.J. and Muench, H., A Simple Method of Estimating Fifty Per Cent Endpoints, *American Journal of Hygiene*, 1938, 27(3):493-497.
- Rezaee, Ramin and Abdollahi, Mohammad, The Importance of Transistability in Drug Discovery, *Expert Opinion on Drug Discovery*, 2017, 12(3):237-239.
- Ritter, M., et al., The Scavenger Receptor CD163: Regulation, Promoter Structure and Genomic Organization, *Pathobiology*, 1999, 67:257-261.
- Ritter, Mirko, et al., Genomic Organization and Chromosomal Localization of the Human CD163 (M130) Gene: A Member of the Scavenger Receptor Cysteine-Rich Superfamily, *Biochem and Biophysical Research Communications*, 1999, 260(2):466-474.
- Robl, J.M., et al., Nuclear Transplantation in Bovine Embryos, *J Anim Sci.*, 1987, 64:642-647.
- Ropp, Susan L., et al., Characterization of Emerging European-Like Porcine Reproductive and Respiratory Syndrome Virus Isolates in the United States, *J Virology*, 2004, 78(7):3684-3703.
- Ross, Jason W., et al., Optimization of square-wave electroporation for transfection of porcine fetal fibroblasts, *Transgenic Res*, 2010, 19(4):611-620.
- Rowland, Raymond R.R., et al., Control of porcine reproductive and respiratory syndrome (PRRS) through genetic improvements in disease resistance and tolerance, *Frontiers in Genetics*, 2012, 3(260):1-6.
- Rowland, Raymond R.R., et al., The Evolution of Porcine Reproductive and Respiratory Syndrome Virus: Quasispecies and Emergence of a Virus Subpopulation during Infection of Pigs with VR-2332, *Virology*, 1999, 259:262-266.
- Saeidnia, Soodabeh, et al., From in vitro Experiments to in vivo and Clinical Studies; Pros and Cons, *Current Drug Discovery Technologies*, 2015, 12(4):218-224.
- Sanchez, Carmen, et al., The Porcine 2A10 Antigen Is Homologous to Human CD163 and Related to Macrophage Differentiation, *J Immunology*, 19 162:5230-5237.
- Sanchez-Torres, C., et al., Expression of porcine CD163 on monocytes/macrophages correlates with permissiveness to African swine fever infection, *Arch Virol*, 2003, 148:2307-2323.
- Sangamo BioSciences, Efficient Generation of Transgenic Pigs Using Zinc Finger Nuclease (ZFN) Technology, Demonstration of ZEN Technology for Efficient Creation of Animals as a Source of Organs for Transplantation into Humans, Press Release, 2011, p. 1.
- Schaer, Christian A., et al., Constitutive Endocytosis of CD163 Mediates Hemoglobin-Heme Uptake and Determines the Noninflammatory and Protective Transcriptional Response of Macrophages to Hemoglobin, *Circ Res*, 2006, 99:943-950.
- Schaer, Dominik J., et al., CD163 is the macrophage scavenger receptor for native and chemically modified hemoglobins in the absence of haptoglobin, *Blood*, 2006, 107(1):373-380.
- Schaer, DJ, et al., Hemophagocytic macrophages constitute a major compartment of hemne oxygenase expression in sepsis, *Eur J Haematol*, 2006, 77:432-436.
- Schurgers, Evelien, et al., Discrepancy between the in vitro and in vivo effects of murine mesenchymal stem cells on T-cell proliferation and collagen-induced arthritis, *Arthritis Research & Therapy*, 2010, 12(R31):3-11.
- Aigner, Bernhard, et al., Transgenic pigs as models for translational biomedical research, *J Mol Med*, 2010, 88:653-664.
- Albina, E., et al., Immune response and persistence of the porcine reproductive and respiratory syndrome virus in infected pigs and far units, *The Veterinary Record*, May 28, 1994, 134:567-573.
- Allende, R., et al., North American and European porcine reproductive and respiratory syndrome viruses differ in non-structural protein coding regions, *J General Virology*, 1999, 80:307-315.
- Allende, R., et al., Porcine Reproductive and Respiratory Syndrome Virus: Deseption of Persistence in Individual Pigs upon Experimental Infection, *J Virology*, Nov. 2000, 74(22):10834-10837.
- American Society for Cell Biology, "Small details between 'in vivo' and 'in vitro' studies make for big differences." *ScienceDaily*, Dec. 13, 2010.
- Andreyev, V.G., et al., Genetic variation and phylogenetic relationships of 22 poreine reproductive and respiratory syndrome virus (PRRSV) field strains based on sequence analysis of open reading frame 5, *Arch Virol* 1997, 142:993-1001.
- ATCC CRL-3216, 293T, Organism: *Homo sapiens*, human, Tissue: embryonic kidney.
- Bauer, B.K., et al., l Arginine Supplementation in vitro Increases Porcine Embryo Development and Affects mRNA Transcript Expression, Reproduction, Fertility and Development, Dec. 7, 2010, 23(1):107.
- Beaton, Benjamin P. and Wells, Kevin D., Compound Transgenics: Recombinase-Mediated Gene Stacking, *Transgenic Animal Technology*, pp. 565-678.
- Benfield, David A., et al., Characterization of swine infertility and respiratory syndrome (SIRS) virus (isolate ATCC VR-2332), *J Vet Diagn Invest*, 1992 4:127-133.
- Benfield, David A., et al., Pathogenesis and persistence of PRRS, Allen D. Leman Swine Conference, 1998, pp. 169-171.

(56)

References Cited**OTHER PUBLICATIONS**

- Berg, H., 200-Biological Implications of Electric Field Effects Part V. Fusion of Blastomeres and Blastocysts of Mouns Embryos, *Bioelectrochemistry and Bioenergetics*, 1982, 9:223-228.
- Boddicker, Nicholas J., et al., Genome-wide association and genomic prediction for host response to porcine reproductive and respiratory syndrome virus infection, *Genetics Selection Evolution*, 2014, 46(18):1-14.
- Bookstein, Robert, et al., Promoter deletion and loss of retinoblastoma gene expression in human prostate carcinoma, *Proc. Natl. Acad. Sci. USA*, 1990, 87:7762-7766.
- Borg, Natalie A., et al., CD1d-lipid-antigen recognition by the semi-invariant NKT T-cell receptor, *Nature*, Jul. 5, 2007, 448:44-49.
- Brinster, Ralph L., et al., Factors affecting the efficiency of introducing foreign DNA into mice by microinjecting eggs, *Proc. Natl. Acad. Sci.*, Jul. 1985, 82:4438-4442.
- Burkard, Christine, et al., Pigs Lacking the Scavenger Receptor Cystine-Rich Domain 5 of CD163 Are Resistant to Porcine Reproductive and Respiratory Syndrome Virus 1 Infection, *J. Virology*, 2018, 92(16):1-17.
- Burkard, Christine, et al., Precision engineering for PRRSV resistance in pigs: Macrophages from genome edited pigs lacking CD163 SRCR5 domain are fully resistant to both PRRSV genotypes while maintaining biological function, *PLOS Pathogens*, Feb. 23, 2017, 13(2):1-28.
- Burlac, C., et al., Identification of human preformed antibody targets in GTKO pigs, *Xenotransplantation* 2012, 19:92-101.
- Calvert, Jay G., et al., CD163 Expression Confers Susceptibility to Porcine Reproductive and Respiratory Syndrome Viruses, *J of Virology*, Jul. 2007, 81(14):7371-7379.
- Carter, D. Bart, et al., Phenotyping of Transgenic Cloned Piglets, *Cloning and Stem Cells*, 2002, 4(2):131-145.
- Chin, Kun-Kuet and Shwu-Pen Chang, The -104G nucleotide of the human CYP21 gene is important for CYP21 transcription activity and protein interaction, *Nucleic Acids Research*, 1998, 26(8):1959-1964.
- Christopher-Hennings, Jane, et al., Persistence of porcine reproductive and respiratory syndrome virus in serum and semen of adult boars, *J Vet Diagn Invest*, 1995, 7:456-464.
- Christopher-Hennings, J, et al., Effects of a modified-live virus vaccine against porcine reproductive and respiratory syndrome in boars, *AJVR*, 1997, 58(1):40-45.
- Ciotti, M., et al., Coding defect and a TATA box mutation at the bilirubin UDP-glucuronosyltransferase gene cause Crigler-Najjar type 1 disease, *Biochimica et Biophysica Acta*, 1998, 1407:40-S0.
- Cong, Le, et al., Multiplex Genome Engineering Using CRISPR/Cas Systems, *Science*, Feb. 15, 2013, 339(6121):819-823.
- Cooper, David K C. Modifying the sugar icing on the transplantation cake, *Glycobiology*, 2016, 26(6):571-581.
- Crocker, Paul R. and Gordon, Siamon, Properties and Distribution of a Lectin-Like Hemagglutinin Differentially Expressed by Murine Stromal Tissue Macrophages, *J Exp Med*, The Rockefeller Univ. Press, Dec. 1986, 164:1862-1875.
- Crocker, Pant R., et al., Molecular analysis of sialoside binding to sialoadhesin by NMR and site-directed mutagenesis, *Biochem J*, 1999, 341:355-361.
- Dai, Yifan, et al., Targeted disruption of the $\alpha 1, 3$ -galactosyltransferase gene in cloned pigs, *Nature*, Mar. 2002, 20:251-255.
- Das, Phani B., et al., The Minor Envelope Glycoproteins GP2a and GP4 of Porcine Reproductive and Respiratory Syndrome Virus Interact with the Receptor CD163, *J Virology*, Feb. 2010, 84(4):1731-1740.
- Dee, S. A. and Molitor, T. W., Elimination of porcine reproductive and respiratory syndrome virus using a test and removal process, *Veterinary Record*, 1998, 143:474-476.
- Delputte, P.L., et al., Effect of virus-specific antibodies on attachment, internalization and infection of porcine reproductive and respiratory syndrome virus in primary macrophages, *Vet Immunology and Immunopathology*, 2004, 102:179-188.
- Delputte, P.L., et al., Involvement of the Matrix Protein in Attachment of Porcine Reproductive and Respiratory Syndrome Virus to a Heparinlike Receptor on Porcine Alveolar Macrophages, *J Virology*, May 2002, 76(9) 4312-4320.
- Delputte, P.L., et al., Poreine Arterivirus Infection of Alveolar Macrophages Is Mediated by Sialic Acid on the Virus, *J Virology*, Aug. 2004, 78(15):8094-8101.
- Delputte, P.L., et al., Porcine Arterivirus Attachment to the Macrophage-Specific Receptor Sialoadhesin Is Dependent on the Sialic Acid-Binding Activity of the N-Terminal Immunoglobulin Domain of Sialoadhesin, *J Virology*, Sep. 2007, 81(17):9546-9550.
- Delputte, P.L., et al., Porcine Sialoadhesin (CD169/Siglec-1) Is an Endocytic Receptor that Allows Targeted Delivery of Toxins and Antigens to Macrophages, *PLoS ONE*, Feb. 2011, 6(2):1-12.
- Etzerodt, Anders, et al., Plasma Clearance of Hemoglobin and Haptoglobin in Mice and Effect of CD163 Gene Targeting Disruption, *Antioxidants & Redox Signaling*, 2013, 18(17):2254-2263.
- Etzerodt, Andres and Moestrup, Soren K., CD163 and Inflammation: Biological, Diagnostic, and Therapeutic Aspects, *Antioxidants & Redox Signaling*, 2013, 18(17):2352-2363.
- Fabrick, Babs O., et al., The macrophage scavenger receptor CD163, *Immunobiology*, 2005, 210:153-160.
- Fabrick, Babs O., et al., The macrophage scavenger receptor CD163 functions as an innate immune sensor for bacteria, *Blood Journal*, Jan. 22, 2009, 113(4):887-892.
- Fisher, D. and Goodall, A.H., Membrane fusion by viruses and chemical agents, *Techniques in Cellular Physiology*, 1981, P115:1-36.
- Fridman, AL and Tainsky, MA, Critical pathways in cellular senescence and immortalization revealed by gene expression profiling, *Oncogene*, 2008, 27:5975-5987.
- Gaj, Thomas, et al., ZEN, TALEN and CRISPR/Cas-based methods for genome engineering, *Trends Biotechnol*, 2013, 31(7):397-405.
- Galili, Uri, Xenotransplantation and ABO incompatible transplantation: The similarities they share, *Transfusion and Apheresis Science*, 2006, 35:45-58.
- Gaudreault, N., et al., Factors affecting the permissiveness of porcine alveolar macrophages for porcine reproductive and respiratory syndrome virus, *Arch Virol*, 2009, 154:133-136.
- Gerrits, Roger J., et al., Perspectives for artificial insemination and genomics to improve global swine populations, *Theriogenology*, 2005, 63:283-299.
- Goryenin, Igor, et al., Applications of Whole Cell and Large Pathway Mathematical Models in the Pharmaceutical Industry, Chapter 12, *Metabolic Engineering in the Post Genomic Era*, Horizon Bioscience, 2004, p. 344.
- Graham, Christopher F., The Fusion of Cells with One- and Two-Cell Mouse Embryos, *Wistar Inst Symp Monogr.*, 1969, 9:19-35.
- Graversen, Jonas H., et al., Targeting the Hemoglobin Scavenger receptor CD163 in Macrophages Highly Increases the Anti-Inflammatory Potency of Dexamethasone, *Molecular Therapy*, Aug. 2012, 20(8):1550-1558.
- Groenen, Martien A. M., et al., Analyses of pig genomes provide insight into porcine demography and evolution, *Nature*, Nov. 15, 2012, 491(7424):393-398.
- Hai, Tang et al., One-step generation of knockout pigs by zygote injection of CRISPR/Cas System, *Cell Research*, 2014, 34:372-375.
- Halbur, P.O., et al., Comparison of the Pathogenicity of Two US Porcine Reproductive and Respiratory Syndrome Virus Isolates with that of the Lelystad Virus, *Vet Pathol*, 1995, 32:648-660.
- Hammer, Robert E., et al., Production of transgenic rabbits, sheep and pigs by microinjection, *Nature*, 1985, 315:680-683.
- Hao, Y.H., et al., Production of endothelial nitric oxide synthase (eNOS) over-expressing piglets, *Transgenic Res*, 2006, 15:739-750.
- Hauschild, Janet, et al., Efficient generation of a biallelic knockout in pigs using zinc-finger nucleases, *PNAS*, 2011, 108(36):1-6.
- Holtkamp, Derald J., et al., Assessment of the economic impact of porcine reproductive and respiratory syndrome virus on United States port producers, *J Swine Health and Production*, 2013, 21(2):72-84.
- Huang, Y.W., et al., Porcine DC-SIGN: Moleenlat cloning, gene structure, tissue distribution and binding characteristics, *Dev and Comparative Immunology*, 2009, 33:464-480.

(56)

References Cited

OTHER PUBLICATIONS

- Hwang, Woong Y, et al., Efficient genome editing in zebrafish using a CRISPR-Cas system, *Nature Biotechnology*, 2013, 31(3):227-229.
- Hyder, Salvoan M., et al., Identification of an Estrogen Response Element in the 3'-Flanking Region of the Murine *c-fos* Protooncogene, *J Biological Chemistry*, 1992, 267(25):18047-18054.
- Im, Gi-Sun, et al., In vitro development of preimplantation porcine nuclear transfer embryos cultured in different media and gas atmospheres, *Theriogenology*, 2004, 61:1125-1135.
- Jeney, Viktona, et al., Pro-oxidant and cytotoxic effects of circulating heme, 2002, 100(3):879-887.
- Kefpaber, Kerry K., Reproductive Failure of Unknown Etiology, 1989, 1(2):1-10.
- Kim, Hyongbum and Kim, Jin-Soo, A guide to genome engineering with programmable nucleases, *Nature Reviews Genetic*, 2014, 15:321-334.
- Kim, H.S., et al., Enhanced replication of porcine reproductive and respiratory syndrome (PRRS) virus in a homogeneous subpopulation of MA-104 cell line, *Arch Vitol*, 1993, 133:477-483.
- Kim, Jeong-Ki, et al., Defining the Cellulat Target(s) of Porcine Reproductive and Respiratory Syndrome Virus Blocking Monoclonal Antibody 7G10, *J Virology*, 2006, 80(2):689-696.
- Kleinjan, Dirk-Jan and Coutinho, Pedro, Cis-ruption mechanisms: disruption of cis-regulatory control as a cause of human genetic disease, *Briefings Functional Genomics and Proteomics*, 2009, 8(4):317-332.
- Kolston, Paul J., Comparing in vitro, in situ, and in vivo experimental data in a three-dimensional model of mammalian cochlear mechanics, *Proc Natl Acad Sci*, 1999, 96:3676-3681.
- Kristiansen, Mette, et al., Identification of the haemoglobin scavenger receptor, *Nature*, 2001, 409: 198-201.
- Kwon, Deug-Nam, et al., Production of biallelic CMP-Neu5Ac hydroxylase knock-out pigs, *Scientific Reports*, 2013, 3(1981)1-10.
- Ladinig, Andrea, et al., Pathogenicity of three type 2 porcine reproductive and respiratory syndrome virus strains in experimentally inoculated pregnant gilts, *Virus Research*, 2015, 203:34-35.
- Lager, Kelly M., et al., Evaluation of protective immunity in gilts inoculated with the NADC-8 isolate of porcine reproductive and respiratory syndrome virus (PRRSV) and challenge-exposed with an antigenically distinct PRRSV isolate, *AJVR*, 1998, 60(8):1022-1027.
- Lai, Liangxue, et al., Generation of cloned transgenic pigs rich in omega-3 fatty acids, *Nature Biotechnology*, 2006, 24(4):435-436.
- Lai, Liangxue, et al., Production of α -1,3-Galactosyltransferase Knockout Pigs by Nuclear Transfer Cloning, *Science*, 2002, 295:1089-1092.
- Lai, Liangxue, and Prather, Randall S., Creating genetically modified pigs by using nuclear transfer, *Reprod Biol and Endocrinology*, 2003, 1(82):1-6.
- Lai, Liangxue and Prather, Randall S., Production of Cloned Pigs by Using Somatic Dells as Donors, *Cloning and Stem Cells*, 2003, 5(4):233-241.
- Lai, Liangxue and Prather, Randall S., A method for Producing Cloned Pigs by Using Somatic Cells as Donors, *Methods in Molecular Biology, Germ Cell Protocols*, 2004, vol. 2, Ch. 9, pp. 149-163.
- Lawson, Stevewn R., et al., Porcine reproductive and respiratory syndrome virus infection of gnotobiotic pigs: sites of virus replication and co-localization with MAC-387 staining at 21 days post-infection, *Virus Research*, 1997, 51:105-113.
- Lee, Kiho, et al., Engraftment of human iPS cells and allogeneic porcine cells into pigs with inactivated RAG2 and accompanying severe combined immunodeficiency, *PNAS*, 2014, 111(20):7260-7265.
- Lee, Kiho, et al., Piglets Produced From Cloned Blastocysts Cultured In Vitro With GM-CSF, *Mol Reprod, Dev*, 2013, 80(2):145-154.
- Li, Dali, et al., Heritable gene targeting in the mouse and rat using a CRISPR-Cas system, *Nature Biotechnology*, 2013, 31(8):681-683.
- Li, Rongfeng, et al., Cloned Transgenic Swine Via In Vitro Production and Cryopreservation, *Biology of Reprod*, 2006, 75:226-230.
- Li, Ying and Jaiswal, Anil K., Regulation of Human NAD(P)H:Quinone Oxidoreductase Gene, *J Biol Chem*, 1992, 267(21):15097-15104.
- Lillico, Simon G., et al., Live pigs produced from genome edited zygotes, *Scientific Reports*, 2013, 3:2847-2850.
- Loudianos, O., et al., Molecular characterization of Wilson disease in the Sardinian population—evidence of a founder effect, *Hum Matat*, 1999, 14(4):294-303.
- Lunney, Joan K. and Chen, Hongbo, Genetic control of host resistance to porcine reproductive and respiratory syndrome virus (PRRSV) infection, *Virus Research*, 2010, 154:161-169.
- MacHat, Zoltan, et al., Complete Activation of Porcine Oocytes Induced by the Sulfhydryl Reagent, Thimerosal, *Biol of Reprod*, 1997, 57:1123-1127.
- MacHat, Zoltan, et al., Development of Early Porcine Embryos In Vitro and In Vivo, *Biol of Reprod*, 1998, 59:451-455.
- Madsen, Mette, et al., Molecular Characterization of the Haptoglobin Hemoglobin Receptor CD163, *J Biol Chem*, 2004, 279(49):51561-51567.
- Mansour, Suzanne L., et al., Disruption of the proto-oncogene *int-2* in mouse embryo-derived stem cells. a general strategy for targeting mutations to non-selectable genes, *Nature*, 1988, 336:348-352.
- Mayes, M.A., et al., Parthenogenic Activation of Pig Oocytes by Protein Kinase Inhibition, *Biol of Reprod*, 1995, 53:270-275.
- McGrath, James and Solter, Davor, Nuclear Transplantation in the Mouse Embryo by Microsurgery and Cell Fusion, *Science*, 1983, 220(4603):1300-1302.
- Meng, X.J., Heterogeneity of porcine reproductive and respiratory syndrome virus: implications for current vaccine efficacy and future vaccine development, *Veterinary Microbiology*, 2000, 74:309-329.
- Meng, Xiang-Jin, et al., Sequence comparison of open reading frames 2 to 5 of low and high virulence United States isolates of porcine reproductive and respiratory syndrome virus, *J Gen Virology*, 1995, 76:3181-3188.
- Mengeling, William L., et al., Identification and clinical assessment of suspected vaccine-related field strains of porcine reproductive and respiratory syndrome virus, *AJVR*, 1999, 60(3): 334-340.
- Shanmukhappa, Kumar, et al., Role of CD151, A tetraspantin, in porcine reproductive and respiratory syndrome virus infection, *Virology Journal*, 2007, 4:62.
- Shimozawa, Nobuhiro, et al., Abnormalities in Cloned Mice Are Not Transmitted to the Progeny, *Genesis*, 2002, 35:203-207.
- Snuder, Eric J., and Meulenberg, Janneke J.M., The molecular biology of arteriviruses, *J Gen Virology*, 1998, 79:961-979.
- Soares, Mignel, P., Bach, Fritz H., Heme oxygenase-1: from biology to therapeutic potential, *Trends in Molecular Medicine*, 2009, 15(2):50-58.
- Stein, Michael, et al., Interleukin 4 Potently Enhances Murine Macrophage Mannose Receptor Activity: A Marker of Alternative Immmmmmologie Macrophage Activation, *J Exp Med*, 1992, 176:287-292.
- Stephen, Sam L., et al., Scavenger Receptors and Their Potential as Therapeutic Targets in the Treatment of Cardiovascular Disease, *International Journal of Hypertension*, 2010, pp. 1-21.
- Stephenson, Rachel J., et al., Multiplex Serology for Common Viral Infections in Feral Pigs (*Sus scrofa*) in Hawaii between 2007 and 2010, *J. Wildlife Diseases*, 2015, 51(1):239-243.
- Suarez, Paloma, et al., Open Reading Frame 5 of Porcine Reproductive and Respiratory Syndrome Virus as a Cause of Virus-Induced Apoptosis, *J Virology*, 1996, 70(5):2876-2882.
- Sulahian, Timothy H., et al., Human Monocytes Express CD163, Which is Unregulated by IL-10 and Identical to p155. Cytokine, 2000, 12(9):1312-1321.
- Sur, Jung-Hyang, et al., In Vivo Detection of Poreme Reproductive and Respiratory Syndrome Virus RNA by In Situ Hybridization at Different Times Postinfection, *J Clinical Microbiology*, 1996, 34(9):2280-2286.

(56)

References Cited**OTHER PUBLICATIONS**

- Sur, Jung-Hiyang, et al., Porcine Reproductive and Respiratory Syndrome Virus Replicates in Testis Germ Cells, Alters Spermatogenesis, and Induces Germ Cell Death by Apoptosis, *J Virology*, 1997, 71(12):9170-9179.
- Tait-Burkard, Christine, et al., Livestock: 2.0—genome editing for fitter, healthier, and more productive farmed animals, *Genome Biology*, 2018, 19(204):1-11.
- Terns, Michael P. and Terns, Rebecca M., CRISPR-Based Adaptive Immune Systems, *Curr Opin Microbiol*, 2011, 14(3):321-327.
- Trible, Benjamin R., et al., Recognition of the Different Structural Forms of the Capsid Protein Determines the Outcome following Infection with Porcine Circovirus Type 2, *J Virology*, 2012, 86(24):13508-13514.
- Vadori, M. and Cozzi, E., The immunological barriers to xenotransplantation, *Tissue Antigens*, 2015, 86:239-253.
- Van Breedam, Wander, et al., Porcine reproductive and respiratory syndrome virus entry into the porcine macrophage, *J General Virology*, 2010, 91:1659-1667.
- Van Breedam, Wander, et al., The M/GP5 Glycoprotein Complex of Porcine Reproductive and Respiratory Syndrome Virus Binds the Sialoadhesin Receptor in a Sialic Acid-Dependent Manner, *PLoS Pathogens*, 2010, 6(1):1-11.
- Van Den Heuvel, Michel M., et al., Regulation of CD163 on human macrophages: cross-linking of CD163 induces signaling and activation, *J Leukocyte Biology*, 1999, 66(5):858-866.
- Van Den Hoff, Maurice J.B., et al., Electroporation in 'intracellular' buffer increases cell survival, *Nucleic Acids Research*, 1992, 20(11):2902.
- Vanderheiden, Nathalie, et al., Involvement of Sialoadhesin in Entry of Porcine Reproductive and Respiratory Syndrome Virus into Porcine Alveolar Macrophages, 2003, *J Virology*, 77(15):8207-8215.
- Van Gorp, Hanne, et al., Scavenger receptor CD163, a Jack-of-all-trades and potential target for cell-directed therapy, 2010, 47:1650-1660.
- Van Gorp, Hanne, et al., Identification of the CD163 Protein Domains Involved in Infection of the Porcine Reproductive and Respiratory Syndrome Virus, 2010, *J Virology*, 84(6):3101-3105.
- Van Gorp, Hanne, et al., Sialoadhesin and CD163 join forces during entry of the porcine reproductive and respiratory syndrome virus, *J General Virology*, 2008, 89:2943-2953.
- Vinson, Mary, et al., Characterization of the Sialic Acid-binding Site in Sialoadhesin by Site-directed Mutagenesis, *J Biological Chemistry*, 1996, 271(16):9267-9272.
- Walters, Eric M. and Prather, Randall S., Advancing Swine Models for Human Health and Diseases, *Missouri Medicine*, 2013, 110(3):212-215.
- Wang, Haoyi, et al., One-Step Generation of Mice Carrying Mutations in Multiple Genes by CRISPR/Cas-Mediated Genome Engineering, *Cell*, 2013, 153(4):910-918.
- Watanabe, Masahito, et al., Knockout of exogenous EGFP gene in porcine somatic cells using zinc-finger nucleases, *Biochem and Biophysical Research Communications*, 2010, 402:14-18.
- Welch, Siao-Kun and Calvert, Jay G., A brief review of CD163 and its role in PRRSV infection, *Virus Research*, 2010, 154:98-103.
- Wells, Kevin D., et al., Replacement of Porcine CD163 Scavenger Receptor Cysteine-Rich Domain 5 with a CD163-Like Homolog Confers Resistance of Pigs to Genotype 1 but Not Genotype 2 Porcine Reproductive and Respiratory Syndrome Virus, *J Virology*, 2017, 91(2):1-11.
- Wensvoort, O., et al., Mystery swine disease in the Netherlands: The isolation of Lelystad virus, *Veterinary Quarterly*, 1991, 13(3):121-130.
- Whitworth, Kristin M., et al., Gene-edited pigs are protected from porcine reproductive and respiratory syndrome virus, *Nature Biotechnology*, 2016, 34(1):20-22.
- Whitworth, K. M., et al., "Disruption the Sialoadhesin and CD163 Genes to Create Pigs Resistant to PRRSV Infectivity," Abstract, Swine in Biomedical Research in Chicago, Illinois, Jul. 17-19, 2011, S1-25, pp. 39-40.
- Whitworth, Kristin M., et al., Method of Oocyte Activation Affects Cloning Efficiency in Pigs, *Mol Reprod Dev*, 2002, 76:1.11.
- Whitworth, Kristin M., et al., Activation Method Does Not Alter Abnormal Placental Gene Expression and Development in Cloned Pigs, *Mol Reprod Dev*, 2010, 77:1016-1030.
- Whitworth, K. M., et al., "Pigs Resistant to PRRSV Infectivity." Poster Presented at Swine in Biomedical Research in Chicago, Illinois, Jul. 18, 2011, 1 page.
- Whitworth, Kristin M., et al., Scriptaid Corrects Gene Expression of a Few Aberrantly Reprogrammed Transcripts in Nuclear Transfer Pig Blastocyst Stage Embryos, *Cellular Reprogramming*, 2011, 13(3):191-204.
- Whitworth, Kristin M. and Prather, Randall S., Gene editing as applied to prevention of reproductive porcine reproductive and respiratory syndrome, *Mol Reprod Dev*, 2017, 33.
- Whitworth, Kristin M., et al., Use of the CRISPR/Cas9 System to Produce Genetically Engineered Pigs from In Vitro-Derived Oocytes and Embryos, *Biol of Reproduction*, 2014, 91(3):78, 1-13.
- Whitworth, K. M., et al., "Use of the CRISPR/Cas9 System to Produce Pigs With a Genetically Modified CD163 Gene by Using Somatic Cell Nuclear Transfer of In Vitro Derived Oocytes," Abstract, Swine in Biomedical Research Conference, Raleigh, North Carolina, Jul. 20-22, 2014, p. 103.
- Whitworth, Kristin M., et al., Use of the CRISPR/Cas9 system to produce pigs with a genetically modified CD163 gene by somatic cell nuclear transfer of in vitro derived oocytes, Division of Animal Science, University of Missouri, Columbia, Poster, p. 1.
- Whitworth, K.M., et al., Gene editing of CD163 protects pigs from PRRSV infectivity, Abstracts from the UC Davis Transgenic Animal Research Conference XI, Aug. 13-17, 2017, *Transgenic Res*, 2018, 27:473-474.
- Whyte, Jeffrey J., et al., Gene Targeting With Zinc Finger Nucleases to Produce Cloned eGFP Knockout Pigs, *Mol Reprod Dev*, 2011, 78(1):1-3.
- Whyte, Jeffrey and Prather, Randall S., Genetic modifications of pigs for medicine and agriculture, *Mol Reprod Dev*, 2011, 78(10-11):879-891.
- Wiedenheft, Blake, et al., RNA-guided genetic silencing systems in bacteria and archaea, *Nature*, 2012, 482:331-338.
- Wills, R. W., et al., Porcine reproductive and respiratory syndrome virus: Persistent infection, *Veterinary Microbiology* 1997, 55:231-240.
- Winckler, C. and Willen, S., The Reliability and Repeatability of a Lameness Scoring System for Use as an Indicator of Welfare in Dairy Cattle, *Acta Agric Scand Sect A, Animal Sci Suppl*, 2001, 30:103-107.
- Wissink, E.H.J., et al., Identification of porcine alveolar macrophage glycoproteins involved in infection of porcine respiratory and reproductive syndrome virus, *Arch Virol*, 2003, 148:177-187.

* cited by examiner

1

PIG WITH A GENETICALLY MODIFIED CD163 GENE RESISTANT TO PRRSV

CROSS REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of and priority to U.S. Non-Provisional application Ser. No. 17/307,369, filed May 4, 2021. Ser. No. 17/307,369 claims priority to U.S. Provisional Application 63/020,128 filed on May 5, 2020 and to U.S. Provisional Application 63/021,370 filed on May 7, 2020. Each application is hereby incorporated by reference, each in their entirety.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

The official copy of the sequence listing is submitted electronically via EFS-Web as an ASCII-formatted sequence listing with a file named RD-12-2020-US2-SEQLIST, created on Oct. 27, 2021, and having a size of 137,066 bytes is filed concurrently with the specification. The sequence listing contained in this ASCII-formatted document is part of the specification and is herein incorporated by reference in its entirety.

FIELD

The present disclosure relates to methods for improving the health of porcine species. In particular, the present disclosure relates to methods of protecting *Sus scrofa* animals and elite lines from infection by porcine reproductive and respiratory syndrome virus (PRRSv) through targeted polynucleotide edits of CD163 to prevent infection and generate resistant animals, herds, and cell lines.

BACKGROUND

Viral infections are a major source of morbidity and mortality in the livestock industry. In particular, Porcine Reproductive and Respiratory Syndrome (PRRS) is a pan-zootic infectious disease of pigs, causing major economic losses to the world-wide pig industry. PRRS manifests in pigs of all ages but primarily causes late-term abortions and stillbirths in sows and respiratory disease in piglets. The causative agent of the disease is the positive-strand RNA PRRS virus (PRRSv). PRRS is the most economically important disease of domestic swine in North America, Europe and Asia, costing producers in North America more than \$600 million annually.

Currently, there are no effective treatment programs for acute PRRS. As a result, when incidence of PRRS is detected on a farm, depopulation, sufficient cleaning/disinfection, and proper disposal of the carcasses must be used to eliminate the virus. In more extreme cases, whole herd depopulation-repopulation has been documented as an effective method of eliminating the PRRSv from endemically infected herds; however, this method results in significant loss.

Vaccines for PRRSv do exist; however, these vaccines have been unable to control the disease largely due to the genetic diversity within the structural proteins of the virus. Consequently, prevention of infection is currently the best control measure. As a prophylactic measure, farms in a country or zone where PRRSv exists must use stringent control measures, involving an assessment of the health

2

status of replacement gilts and boars, as well as 45-60 days of isolation and acclimatization for incoming stock.

In recent years, more attention has been given to the role CD163 may play in the occurrence of PRRS. Despite the significant heterogeneity in strains of PRRSv, strains of PRRSv all share a tropism for CD163-positive cells. Although CD163 is a virus receptor, the CD163 scavenger receptor is also involved in the adhesion of monocytes to endothelial cells. Functions and a detailed description of CD163 are provided in Onofre, Gabriela et al., *ACTA MEDICA*, 2009, 52, 57-61.

CD163 is a 130 kDa type 1 membrane protein considered to be a fusion receptor for the PRRS virus; it is mapped to chromosome 5 in pigs. The basic transcript encodes for a protein of 1076 amino acids. There are five reported isoforms of CD163; three of the isoforms display different splicing forms of their cytoplasmic domains. Generally, however, the genomic molecule sequence of CD163 comprises 17 exons coding for a peptide signal sequence, nine scavenger receptor cysteine-rich (SRCR) domains, two proline serine threonine (PST) linker domains, a cytoplasmic domain, and a short cytoplasmic tail. CD163 has been described as the receptor for PRRSv. Domain 5 (SRCR5) of the protein is the interaction site for the virus. Exon 7 of CD163 encodes the SRCR domain 5 (SRCR5) that serves as an interaction site for the PRRSv in vitro. Burkard (Burkard, C., *PLoS Pathog.* 2017, 23, 13, e1006206.) demonstrated that removal of CD163 exon 7 confers PRRSv resistance to porcine macrophages. The guides used in that work (set forth as targeting sequences including the PAM in SEQ ID NOs: 272 and 273), however, may lack sufficient activity and specificity for gene editing as part of a commercial breeding program. Further work by Whitworth and colleagues included creating a 123 bp deletion in Exon 7 using guides as set forth (including the PAM) in SEQ ID NO: 354 and SEQ ID NO: 211 (Whitworth, K. M., *Biol. Reprod.*, 2014, 91, 1-13). Whitworth et al. (Whitworth, K. M., *Nature Biotechnology*, 2016, 34, 20-22) reported the preparation of PRRSv resistant pigs by knocking out the function of CD163.

Genome editing includes altering the genome by deleting, inserting, or substituting specific nucleic acid sequences. The alteration can be gene- or location-specific. Genome editing can use site-directed nucleases, such as Cas proteins and their cognate polynucleotides.

Clustered regularly interspaced short palindromic repeats (CRISPR) and CRISPR-associated proteins (Cas) constitute the CRISPR-Cas system.

Cas9 is an exemplary Type II CRISPR Cas protein. Cas9 is an endonuclease that can be programmed by the tracrRNA/crRNA to cleave, in a site-specific manner, a DNA target sequence using two distinct endonuclease domains (HNH and RuvC/RNase H-like domains) (see U.S. Patent Application Publication No. 2014-0068797, published 6 Mar. 2014; see also Jinek, M., et al., *Science*, 337:816-821 (2012) and Karvelis et al. *Genome Biology* (2015) 16:253.

The foregoing CD163 edits, while demonstrating some indication of efficacy against PRRSv, cannot be made as precisely and effectively as the edits disclosed herein. There is a need to improve the health of a porcine herd by editing the CD163 gene using guides for improved editing activity and reduced unintended edits, while conferring resistance to PRRSv.

SUMMARY

The present specification provides for and includes edited CD163 genes that confer PRRSv resistance on pigs comprising the edited gene.

In some embodiments, the present teachings provide for and include a CD163 gene edited to confer PRRSV resistance in *Sus scrofa* wherein the edit excises exon 7 and the edited gene can comprise a repaired genomic sequence as set forth in any one of SEQ ID NOs: 426-458 and 520-555. In some configurations, the edited gene can comprise a repaired genomic sequence selected from the group consisting of SEQ ID NOs: 426-458. In various configurations, the repaired gene sequence can comprise SEQ ID NO: 426, SEQ ID NO: 427, SEQ ID NO: 428, SEQ ID NO: 429, SEQ ID NO: 430, SEQ ID NO: 431, SEQ ID NO: 432, SEQ ID NO: 433, SEQ ID NO: 434, SEQ ID NO: 435, SEQ ID NO: 436, SEQ ID NO: 437, SEQ ID NO: 438, SEQ ID NO: 439, SEQ ID NO: 440, SEQ ID NO: 441, SEQ ID NO: 442, SEQ ID NO: 443, SEQ ID NO: 444, SEQ ID NO: 445, SEQ ID NO: 446, SEQ ID NO: 447, SEQ ID NO: 448, SEQ ID NO: 449, SEQ ID NO: 450, SEQ ID NO: 451, SEQ ID NO: 452, SEQ ID NO: 453, SEQ ID NO: 454, SEQ ID NO: 455, SEQ ID NO: 456, SEQ ID NO: 457, SEQ ID NO: 458, SEQ ID NO: 520, SEQ ID NO: 521, SEQ ID NO: 522, SEQ ID NO: 523, SEQ ID NO: 524, SEQ ID NO: 525, SEQ ID NO: 526, SEQ ID NO: 527, SEQ ID NO: 528, SEQ ID NO: 529, SEQ ID NO: 530, SEQ ID NO: 531, SEQ ID NO: 532, SEQ ID NO: 533, SEQ ID NO: 534, SEQ ID NO: 535, SEQ ID NO: 536, SEQ ID NO: 537, SEQ ID NO: 538, SEQ ID NO: 539, SEQ ID NO: 540, SEQ ID NO: 541, SEQ ID NO: 542, SEQ ID NO: 543, SEQ ID NO: 544, SEQ ID NO: 545, SEQ ID NO: 546, SEQ ID NO: 547, SEQ ID NO: 548, SEQ ID NO: 549, SEQ ID NO: 550, SEQ ID NO: 551, SEQ ID NO: 552, SEQ ID NO: 553, SEQ ID NO: 554, or SEQ ID NO: 555. In various configurations, the edit can be created using a pair of gRNAs, wherein the pair of gRNAs can be SEQ ID NOs: 229 and 256, SEQ ID NOs: 230 and 256, SEQ ID NOs: 231 and 256, SEQ ID NOs: 237 and 256, SEQ ID NOs: 241 and 256, SEQ ID NOs: 229 and 258, SEQ ID NOs: 230 and 258, SEQ ID NOs: 231 and 258, SEQ ID NOs: 237 and 258, SEQ ID NOs: 241 and 258, SEQ ID NOs: 229 and 261, SEQ ID NOs: 230 and 261, SEQ ID NOs: 231 and 261, SEQ ID NOs: 237 and 261, SEQ ID NOs: 241 and 261, SEQ ID NOs: 219 and 256, SEQ ID NOs: 221 and 256, SEQ ID NOs: 224 and 256, SEQ ID NOs: 227 and 256, SEQ ID NOs: 219 and 258, SEQ ID NOs: 221 and 258, SEQ ID NOs: 224 and 258, SEQ ID NOs: 227 and 258, SEQ ID NOs: 219 and 261, SEQ ID NOs: 221 and 261, SEQ ID NOs: 224 and 261, SEQ ID NOs: 227 and 261, SEQ ID NOs: 249 and 256, SEQ ID NOs: 250 and 256, SEQ ID NOs: 249 and 258, SEQ ID NOs: 250 and 258, SEQ ID NOs: 249 and 261, or SEQ ID NOs: 250 and 261. In various configurations, the edit can be created using a pair of gRNAs, wherein the pair of targeting regions can be SEQ ID NOs: 229 and 256, SEQ ID NOs: 230 and 256, SEQ ID NOs: 231 and 256, SEQ ID NOs: 241 and 256, SEQ ID NOs: 229 and 258, SEQ ID NOs: 231 and 258, SEQ ID NOs: 241 and 258, SEQ ID NOs: 219 and 256, SEQ ID NOs: 221 and 256, SEQ ID NOs: 224 and 256, SEQ ID NOs: 227 and 256, SEQ ID NOs: 227 and 258, SEQ ID NOs: 221 and 261, SEQ ID NOs: 249 and 256, SEQ ID NOs: 250 and 256, SEQ ID NOs: 249 and 258, or SEQ ID NOs: 249 and 261.

In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 229 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 230 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 231 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 237 and 256. In various configurations, the edit can be created using

a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 241 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 229 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 230 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of targeting regions can be SEQ ID NOs: 231 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 237 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 241 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 229 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 230 and 261. In various configurations, the edit can be a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 231 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 237 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 241 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 219 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 221 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 224 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 227 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 219 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 221 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 224 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 227 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 219 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 224 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 227 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 249 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 250 and 256. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 249 and 258. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 249 and 261. In various configurations, the edit can be created using a pair of gRNAs wherein the pair of gRNAs can be SEQ ID NOs: 250 and 261. In various configurations, the repaired genomic sequence can comprise

SEQ ID NO: 453 and the edit can be created using sequences set forth in SEQ ID NOs: 249 and 256.

In various configurations, the instant disclosure provides for and includes a *Sus scrofa* cell that can comprise the CD163 gene of the present teachings. In some configurations, the present teachings further provide for and include a cell line that can comprise a plurality of the *Sus scrofa* cell. In some configurations, the cell line can be a fibroblast cell line. In various configurations, the cell, plurality of cells, or cell line can be derived from PIC line 2, PIC line 3, PIC line 15, PIC line 19, PIC line 27, PIC line 62, or PIC line 65. The present teachings further provide for an embryo, piglet, or mature adult that can comprise a plurality of the cell.

The present disclosure also provides for a CD163 gene edited to confer PRRSV resistance in *Sus scrofa* wherein the edit creates a stop codon resulting in a predicted exon 7 amino acid sequence that can be selected from the group consisting of 506-517. In various configurations, the amino acid sequence can be set forth in SEQ ID NO: 506, SEQ ID NO: 507, SEQ ID NO: 508, SEQ ID NO: 509, SEQ ID NO: 510, SEQ ID NO: 511, SEQ ID NO: 512, SEQ ID NO: 513, SEQ ID NO: 514, SEQ ID NO: 515, SEQ ID NO: 516, or SEQ ID NO: 517. In various configurations, the predicted amino acid sequence of exon 7 can be set forth in SEQ ID NO: 513. In some configurations, the edit can be created using gRNAs selected from the group consisting of SEQ ID NOs: 351 and 365, SEQ ID NOs: 351 and 387, SEQ ID NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 348 and 395, SEQ ID NOs: 352 and 365, SEQ ID NOs: 352 and 387, SEQ ID NOs: 352 and 399, SEQ ID NOs: 353 and 365, SEQ ID NOs: 353 and 387, SEQ ID NOs: 353 and 399, SEQ ID NOs: 354 and 390, SEQ ID NOs: 354 and 388, SEQ ID NOs: 354 and 395, SEQ ID NOs: 358 and 361, SEQ ID NOs: 358 and 362, SEQ ID NOs: 358 and 368, SEQ ID NOs: 358 and 384, SEQ ID NOs: 358 and 394, SEQ ID NOs: 358 and 399, SEQ ID NOs: 359 and 390, SEQ ID NOs: 359 and 388, SEQ ID NOs: 359 and 395, SEQ ID NOs: 360 and 368, SEQ ID NOs: 360 and 384, SEQ ID NOs: 360 and 389, SEQ ID NOs: 360 and 394, SEQ ID NOs: 360 and 397, SEQ ID NOs: 361 and 365, SEQ ID NOs: 361 and 387, SEQ ID NOs: 362 and 390, SEQ ID NOs: 362 and 388, SEQ ID NOs: 362 and 395, SEQ ID NOs: 364 and 365, SEQ ID NOs: 364 and 387, SEQ ID NOs: 364 and 399, SEQ ID NOs: 365 and 368, SEQ ID NOs: 365 and 384, SEQ ID NOs: 365 and 389, SEQ ID NOs: 365 and 394, SEQ ID NOs: 365 and 397, SEQ ID NOs: 366 and 368, SEQ ID NOs: 366 and 384, SEQ ID NOs: 366 and 389, SEQ ID NOs: 366 and 394, and SEQ ID NOs: 366 and 397. In various configurations, the edit can be created using gRNAs selected from the group consisting of SEQ ID NOs: 351 and 365, SEQ ID NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 354 and 390, SEQ ID NOs: 358 and 394, SEQ ID NOs: 362 and 390, and SEQ ID NOs: 366 and 394. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 351 and 365. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 351 and 387. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 348 and 390. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 348 and 388. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 348 and 395. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 352 and 365. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 352 and 387. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 352 and 399. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 353 and 365. In various

configurations, the gRNAs used to create the edit can be SEQ ID NOs: 353 and 387. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 353 and 399. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 354 and 390. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 354 and 388. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 354 and 395. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 361. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 362. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 368. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 384. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 394. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 358 and 399. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 359 and 390. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 359 and 388. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 359 and 395. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 360 and 368. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 360 and 384. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 360 and 389. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 360 and 394. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 360 and 397. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 361 and 365. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 361 and 387. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 362 and 390. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 362 and 388. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 362 and 395. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 364 and 365. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 364 and 387. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 364 and 399. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 365 and 368. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 365 and 384. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 365 and 389. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 365 and 397. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 366 and 368. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 366 and 384. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 366 and 394. In various configurations, the gRNAs used to create the edit can be SEQ ID NOs: 366 and 397.

In various configurations, the edited CD163 gene can have a nucleic acid sequence selected from the group consisting of SEQ ID NO: 459-504. In various configurations, the repaired gene can have a nucleic acid sequence set forth in SEQ ID NO: 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490,

491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, or 504. In various configurations, the repaired gene can have a nucleic acid sequence as set forth in SEQ ID NO: 489. In various configurations the predicted amino acid sequence of exon 7 can be as set forth in SEQ ID NO: 513. In various configurations, the predicted amino acid sequence of the gene can be SEQ ID NO: 513, the repaired gene can have a nucleic acid sequence set forth in SEQ ID NO: 489, and the edit can be created using sequences set forth in SEQ ID NOs: 362 and 390.

The present disclosure provides for and includes a *Sus scrofa* cell that can comprise the CD163 gene edited to comprise an exogenous stop codon as described supra. The present disclosure also provides for a cell line that can comprise a plurality of the cell comprising the CD163 gene edited to comprise an exogenous stop codon. In some configurations, the cell line can be a fibroblast cell line. In various configurations, the cell can be derived from PIC line 2, PIC line 3, PIC line 15, PIC line 19, PIC line 27, PIC line 62, or PIC line 65. The present disclosure also provides for and includes an embryo, piglet, or mature adult comprising a plurality of the cell that can comprise the CD163 gene edited to comprise an exogenous stop codon.

In various embodiments, the present teachings provide for and include a pair of gRNAs for editing a *Sus scrofa* CD163 gene having a sequence as set forth in SEQ ID NOs: 229 and 256, SEQ ID NOs: 230 and 256, SEQ ID NOs: 231 and 256, SEQ ID NOs: 237 and 256, SEQ ID NOs: 241 and 256, SEQ ID NOs: 229 and 258, SEQ ID NOs: 230 and 258, SEQ ID NOs: 231 and 258, SEQ ID NOs: 237 and 258, SEQ ID NOs: 241 and 258, SEQ ID NOs: 229 and 261, SEQ ID NOs: 230 and 261, SEQ ID NOs: 231 and 261, SEQ ID NOs: 237 and 261, SEQ ID NOs: 241 and 261, SEQ ID NOs: 219 and 256, SEQ ID NOs: 221 and 256, SEQ ID NOs: 224 and 256, SEQ ID NOs: 227 and 256, SEQ ID NOs: 219 and 258, SEQ ID NOs: 221 and 258, SEQ ID NOs: 224 and 258, SEQ ID NOs: 227 and 258, SEQ ID NOs: 219 and 261, SEQ ID NOs: 221 and 261, SEQ ID NOs: 224 and 261, SEQ ID NOs: 227 and 261, SEQ ID NOs: 249 and 256, SEQ ID NOs: 250 and 256, SEQ ID NOs: 249 and 258, SEQ ID NOs: 250 and 258, SEQ ID NOs: 249 and 261, SEQ ID NOs: 250 and 261, SEQ ID NOs: 351 and 365, SEQ ID NOs: 351 and 387, SEQ NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 348 and 395, SEQ ID NOs: 352 and 365, SEQ ID NOs: 352 and 387, SEQ ID NOs: 352 and 399, SEQ ID NOs: 353 and 365, SEQ ID NOs: 353 and 387, SEQ ID NOs: 353 and 399, SEQ ID NOs: 354 and 390, SEQ ID NOs: 354 and 388, SEQ ID NOs: 354 and 395, SEQ ID NOs: 358 and 361, SEQ ID NOs: 358 and 362, SEQ ID NOs: 358 and 368, SEQ ID NOs: 358 and 384, SEQ ID NOs: 358 and 394, SEQ ID NOs: 358 and 399, SEQ ID NOs: 359 and 390, SEQ ID NOs: 359 and 388, SEQ ID NOs: 359 and 395, SEQ ID NOs: 360 and 368, SEQ ID NOs: 360 and 384, SEQ ID NOs: 360 and 389, SEQ ID NOs: 360 and 394, SEQ ID NOs: 360 and 397, SEQ ID NOs: 361 and 365, SEQ ID NOs: 361 and 387, SEQ ID NOs: 362 and 390, SEQ ID NOs: 362 and 388, SEQ ID NOs: 362 and 395, SEQ ID NOs: 364 and 365, SEQ ID NOs: 364 and 387, SEQ ID NOs: 364 and 399, SEQ ID NOs: 365 and 368, SEQ ID NOs: 365 and 384, SEQ ID NOs: 365 and 389, SEQ ID NOs: 365 and 394, SEQ ID NOs: 365 and 397, SEQ ID NOs: 366 and 368, SEQ ID NOs: 366 and 384, SEQ ID NOs: 366 and 389, SEQ ID NOs: 366 and 394, or SEQ ID NOs: 366 and 397. In some configurations, the pair of gRNAs can have a sequence as set forth in SEQ ID NOs: 229 and 256, SEQ ID NOs: 230 and 256, SEQ ID NOs: 231 and 256, SEQ ID NOs: 241 and 256, SEQ ID NOs: 229 and 258, SEQ ID NOs: 231 and 258, SEQ ID NOs: 241 and 258, SEQ ID NOs: 219

and 256, SEQ ID NOs: 221 and 256, SEQ ID NOs: 224 and 256, SEQ ID NOs: 227 and 256, SEQ ID NOs: 227 and 258, SEQ ID NOs: 221 and 261, SEQ ID NOs: 249 and 256, SEQ ID NOs: 250 and 256, SEQ ID NOs: 249 and 258, SEQ ID NOs: 249 and 261, SEQ ID NOs: 351 and 365, SEQ ID NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 354 and 390, SEQ ID NOs: 358 and 394, SEQ ID NOs: 362 and 390, or SEQ ID NOs: 366 and 394. In various configurations, the pair of gRNAs can have a sequence as set forth in SEQ ID NOs: 249 and 256. In various configurations, the pair of gRNAs can have a sequence as set forth in SEQ ID NOs: 362 and 390.

The present disclosure provides for and includes a CRISPR complex for editing a CD163 gene of *Sus scrofa* comprising a pair of gRNAs of the present teachings.

The present disclosure provides for and includes a method for editing a CD163 gene of *Sus scrofa* comprising using a CRISPR-CAS complex comprising a pair of gRNAs of the present teachings.

The present disclosure provides for and includes a method for preparing a PRSSV resistant *Sus scrofa* cell by using a CRISPR-CAS complex comprising the pair of gRNAs of the present teachings.

The present disclosure provides for and includes a method of producing PRSSV resistant *Sus scrofa* animals, comprising: a) editing a CD163 gene of a *Sus scrofa* cell or a plurality of *Sus scrofa* cells using a CRISPR complex comprising a pair of gRNAs of the present teachings; and b) producing an animal from the cell or plurality of cells.

The present disclosure provides for and includes the use of a cell line according to the present teachings in producing PRSSV resistant animals.

The present disclosure also provides for and includes an embryo, piglet, or mature adult comprising a plurality of the cell according to the present teachings.

In various embodiments the present disclosure provides for and includes a method of determining the presence or absence of an edited sequence having 90% or 95% identity with a sequence set forth in SEQ ID NO: 453 comprising performing real time PCR with a) differentially labeled probes of sequences set forth in SEQ ID NO: 564 and SEQ ID NO: 558 or 561; b) a primer pair set forth in SEQ ID NOs: 562 and 563; and c) a primer pair set forth in SEQ ID NOs: 556 and 557 or SEQ ID NOs: 559 and 560. In some configurations, the edited sequence can have 100% identity with the sequence set forth in SEQ ID NO: 453. In some configurations, the edited sequence can have 90% identity with the sequence set forth in SEQ ID NO: 453. In some configurations, the edited sequence can have 95% identity with the sequence set forth in SEQ ID NO: 453.

In various embodiments, the present teachings provide for and include a PCR primer selected from the group consisting of SEQ ID NO: 556, SEQ ID NO: 557, SEQ ID NO: 559, SEQ ID NO: 560, SEQ ID NO: 562, and SEQ ID NO: 563. In some configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 556. In various configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 557. In various configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 559. In various configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 560. In various configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 562. In various configurations, the PCR primer can have a sequence as set forth in SEQ ID NO: 563.

In some embodiments, the present disclosure provides for and includes a real time PCR probe selected from the group consisting of SEQ ID NO: 558, SEQ ID NO: 561, and SEQ

ID NO: 564. In some configurations, the probe has a sequence as set forth in SEQ ID NO: 558. In various configurations, the probe has a sequence as set forth in SEQ ID NO: 561. In various configurations, the probe has a sequence as set forth in SEQ ID NO: 564.

In various embodiments, the present teachings provide for and include the use of PCR primers set forth in a) SEQ ID NO: 556 and 557 and SEQ ID NO: 562 and 563 or b) SEQ ID NO: 559 and 560 and SEQ ID NO: 562 and 563 to determine the presence or absence of an edited genome sequence set forth in SEQ ID NO: 453.

In various embodiments, the present teachings provide for and include the use of PCR probes set forth in a) SEQ ID NOs: 558 and 564 or b) SEQ ID NOs: 561 and 564 to determine the presence or absence of an edited genome sequence set forth in SEQ ID NO: 453.

In various embodiments, the present teachings provide for and include a method of creating a PRRSV resistant pig comprising editing the pig's genome to comprise a genomic sequence as set forth in SEQ ID NO: 453. In some configurations, the editing the pig's genome can comprise administering gRNAs having sequences set forth in SEQ ID NOs: 249 and 256. In various configurations, the administering comprises injecting pre-formed RNP complexes comprising the gRNAs and a CAS protein into a zygote, embryo, or MII oocyte. In various configurations, the pig is a PICTTM line 2 (Pig Improvement Company, Ltd, Basingstoke, UK), PICTTM line 3, PICTTM line 15, PICTTM line 19, PICTTM line 27, PICTTM line 62, or PICTTM line 65 pig.

In various embodiments, the present disclosure provides for and includes, a *Sus scrofa* animal comprising an edited gene that confers PRRSV resistance in *Sus scrofa* wherein the edit excises exon 7 and the edited gene comprises a repaired genomic sequence set forth in SEQ ID NO: 453. In some configurations, the edit can be made with guide RNAs (gRNAs) having sequences set forth in SEQ ID NOs: 249 and 256. In various configurations, the animal can be an edited animal of PICTTM line 2, PICTTM line 3, PICTTM line 15, PICTTM line 19, PICTTM line 27, PICTTM line 62, or PICTTM line 65. In various configurations, the present disclosure provides for and includes a cell prepared from the animal of the present teachings. In various configurations, the present disclosure provides for and includes a cell line prepared from the cell according to the present teachings. In some configurations, the cell line can be a fibroblast cell line.

In some embodiments, the present teachings provide for and include a CD163 gene edited to confer PRRSV resistance in *Sus scrofa* wherein the edit excises exon 7 and the edited gene comprises a repaired genomic sequence set forth in SEQ ID NO: 453. In some configurations, the edit is created using sequences set forth in SEQ ID NOs: 249 and 256. In various configurations, the present disclosure provides for a *Sus scrofa* cell comprising the CD163 gene according to the present teachings. In some configurations, the present disclosure provides for a cell line comprising a plurality of the cell according to the present teachings. In some configurations, the cell line can be a fibroblast cell line. In various configurations, the cell can be isolated from PIC line 2, PIC line 3, PIC line 15, PIC line 19, PIC line 27, PIC line 62, or PIC line 65.

In various embodiments, the present disclosure provides for and includes a pair of gRNAs for editing a *Sus scrofa* CD163 gene comprising the guide sequences set forth in SEQ ID NOs: 249 and 256.

In various embodiments, the present teachings provide for and include a method of creating a PRRSV resistant pig comprising editing the pig's genome to comprise a genomic

sequence as set forth in SEQ ID NO: 453. In some configurations, editing the pig's genome can comprise administering gRNAs having sequences set forth in SEQ ID NOs: 249 and 256. In some configurations, the administering can comprise injecting pre-formed RNP complexes comprising the gRNAs and a CAS protein into a zygote, embryo, or MII oocyte. In various configurations, the pig can be a PICTTM line 2, PICTTM line 3, PICTTM line 15, PICTTM line 19, PICTTM line 27, PICTTM line 62, or PICTTM line 65 pig.

DETAILED DESCRIPTION

The aspects of the present teachings include, but are not limited to, particular methods of improving the health of a porcine species by targeted inactivation of CD163, which can vary and are understood by skilled artisans. It is further to be understood that all terminology used herein is for the purpose of describing particular aspects only, and is not intended to be limiting in any manner or scope. For example, as used in this specification and the appended claims, the singular forms "a," "an," and "the" can include plural referents unless the content clearly indicates otherwise. Further, all units, prefixes, and symbols may be denoted in their SI accepted forms. Numeric ranges recited within the specification are inclusive of the numbers defining the range and include each integer within the defined range.

So that the present application may be more readily understood, certain terms are first defined. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which aspects of the present application pertain. Many methods and materials similar, modified, or equivalent to those described herein can be used in the practice of the aspects of the present application without undue experimentation, the preferred materials and methods are described herein. In describing and claiming the aspects of the present application, the following terminology will be used in accordance with the definitions set out below.

As used herein, "animal cell" means a cell, including, but not limited to, a somatic cell, culture cell, gamete cell, blood cell, zygote, and embryonic cell. These animal cells can be reproductive or non-reproductive cells. As used herein, cells may be isolated from an animal or embryo and maintained in tissue culture. Also included are mixed cultures that can comprise gene edited cells of the present specification and a non-gene edited support cell or feeder cell.

As used herein, the terms "gene edited," "genetically edited," and "genome-edited," refer to the use of homing technology with naturally occurring or artificially engineered endonucleases, often referred to as "homing endonucleases," or "targeting endonucleases." "Genome-editing" and "gene editing," refer to altering the genome by deleting, inserting, or substituting specific nucleic acid sequences. The altering can be gene or location specific, but need not be altering the sequence of a gene per se. Genome editing can use endonucleases such as the CRISPR system to cut a nucleic acid, thereby generating a site for the alteration. Other endonucleases are available and are suitable for use; however, off-site cutting and specificity can be significant problems. In systems like CRISPR and others, the nuclease can be directed to the target site by complexing with a polynucleotide, herein called a "target sequence," to introduce a site specific DSB. Not to be limited by theory, the DSB can then be repaired by endogenous non-homologous end joining (NHEJ) machinery.

A number of endonucleases are known that are suitable for, and have been adapted to, gene editing. Gene editing

methods are known in the art, including Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) systems (e.g., the CRISPR/Cas9 system), Transcription Activator-Like Effector Nucleases (TALENs), and Gene editing nucleases including Zinc Finger Nucleases (ZFNs). Gene edited animals can be distinguished from transgenic animals by the incorporation of exogenous DNA sequences, particularly sequences having identity with sequences from foreign species in the latter. Here, the terms include, and provide for, small deletions and insertions that do not introduce more than 10 nucleotides. The terms also encompass progeny animals such as those created by sexual crosses or asexual propagation from the initial gene edited animal.

As used herein, the term “repair template” or “repair sequences” refers to a polynucleotide introduced into a cell undergoing DSB repair at nuclease targeted locations in the genome to guide the repair and provide accurate and selective editing. Repair templates can be used to selectively change or delete sequences at a locus having a DSB and generally comprise a 5' genome hybridizing region and a 3' genome hybridizing region. In an aspect, the 5' genome hybridizing and 3' genome hybridizing regions can share at least 80% homology. In an aspect, the 5' genome hybridizing and 3' genome hybridizing regions can share identity. Further descriptions of the repair templates of the present disclosure are provided below. For deletion templates, a contiguous region of the genome can be separated by the deletion of one or more nucleotides. Targeted editing of individual bases can be prepared using repair templates comprising a contiguous region of a genome and having one or more base changes, including addition of a stop codon. The percentage identity of a repair template to a contiguous region of the genome can be between 80% and 100% using an ungapped alignment. As provided herein, a core region of identity within a repair template is provided that may be flanked by a flanking homology region sharing between 80% and 100% homology to the targeted chromosomal region. Not to be limited by theory, the core region identity can increase the fidelity of the repair process while extended regions of homology on either side increase the efficiency and allow for genotypic variation among cells targeted for editing.

As used herein, the term “wild type” or “WT” refers to a phenotype, genotype, or gene that predominates in a natural population of pigs or line of pigs. When used to compare phenotypes and genotypes of gene edited cells and animals of specific lines, the term “wild type” refers to non-gene edited cells and animals of the same line. In an aspect, the present disclosure provides for comparison of edited pigs to non-edited pigs of a similar breed having a similar genetic background. In an aspect, the present disclosure provides for comparison of edited pigs to non-edited pigs of the same breeding line.

As used herein, the term “knock-out” refers to the disruption of gene function by reduction or elimination of its expression. Knock-outs may be generated through the creation of double-strand breaks (DSBs) in the chromosome which can then be repaired using either non-homologous end joining (NHEJ) or homologous recombination of DNA repair templates or targeting vectors by homology-directed repair (HDR). Not to be limited by theory, HDR knock-outs may also be prepared by microhomology-mediated end joining providing a repair template to insert, delete, or edit genomic sequences. Knock-outs may also be generated through replacement vectors, or hit-and-run vectors, or random insertion of a gene trap vector resulting in complete, partial, or conditional loss of gene function.

References herein to a deletion in a nucleotide sequence spanning a range are inclusive of all nucleotides in the listed range. For example, a 5 base pair deletion from nucleotide “a” to nucleotide “e” means that each of nucleotides “a,” “b,” “c,” “d,” and “e” have been deleted (where “b,” “c,” and “d” are between “a” and “e”).

As used herein, the term “edit” includes alterations in the nucleotide sequence of a polynucleotide, such as, for example, a gene, coding DNA sequence (CDS), or non-coding DNA sequence, compared to the wild-type sequence. The term “edits” may include insertions, deletions, splice-donor site edits, point-edits, and the like.

As used herein, the term “clustered regularly interspaced short palindromic repeats” or “CRISPR” refers to a gene editing system utilizing a CRISPR segment of genetic material, and the RNA segments and enzymes it produces, to identify and modify specific DNA sequences in the genome of other organisms. CRISPR systems include Type I, Type II, and Type III CRISPR systems. As used herein, the term “CRISPR associated protein” or “Cas” refers to a protein family that can be strictly associated with CRISPR elements and always occurs near a repeat cluster of CRISPR segments. For example, Cas proteins can include, but are not limited to, Cas9 family member proteins, Cas6 family member proteins (e.g., Csy4 and Cas6), and Cas5 family member proteins. Examples of Cas protein families and methods of identifying the same have been disclosed in Haft, Daniel H., et al., *PLoS Comput. Bio.*, 2005, 1, e60.

While not being limited by any particular scientific theory, a CRISPR nuclease can form a complex with a guide RNA (gRNA), which hybridizes with a complementary target nucleic acid molecule, thereby guiding the CRISPR nuclease to the target nucleic acid molecule. The crRNA comprises a repeat sequence and a spacer sequence which can be complementary to a specific protospacer sequence in an invading pathogen. It is the spacer sequence that can be designed to be complementary to target sequences in a eukaryotic genome. CRISPR nucleases can associate with their respective crRNAs in their active forms. The present specification provides for, and includes a crRNA that can comprise an RNA spacer comprising the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. As used herein, each of SEQ ID NOs: 22 to 271 and 347 to 425 provides for, and includes, the corresponding RNA sequence substituting uridine for the thymidine and ribose for deoxy-ribose. Also, as used herein, the SEQ ID NOs include the PAM (see below) which is present in the genome sequence targeted. Skilled artisans will recognize that the gRNA targeting this sequence would not include the PAM, and therefore would recognize that a gRNA having a sequence set forth in these SEQ ID NOs would only include the first 20 nucleotides, thus excluding the PAM.

Some CRISPR nucleases, such as CasX and Cas9, can require another non-coding RNA component, referred to as a trans-activating crRNA (tracrRNA), to have functional activity. A crRNA comprising a spacer sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425 can be covalently linked to the 5' end of a tracrRNA into one nucleic acid molecule in what is herein referred to as a “single guide RNA” (sgRNA), as described in Jinek, et al., *Science*, 337, 2012, 816-821. As used herein, the tracrRNA is also referred to as “guide RNA backbone” or “the backbone” sequence. As used herein, gRNA includes both a single guide RNA or separate molecules comprising the spacer sequence in the crRNA for use with a separate tracrRNA. The gRNA can

guide the active Cas complex to a target site complementary to the spacer sequence in the crRNA, where the Cas nuclease can cleave the target site. The crRNA can have a region of complementarity to a potential DNA target sequence and a second region that can form base-pair hydrogen bonds with the tracrRNA to form a secondary structure, typically to form at least one stem structure. The region of complementarity to the DNA target sequence can be the spacer or spacer sequence. The tracrRNA and a crRNA can interact through a number of base-pair hydrogen bonds to form secondary RNA structures. Complex formation between tracrRNA/crRNA and Cas9 protein can result in a conformational change of the Cas9 protein that facilitates binding to DNA, endonuclease activities of the Cas9 protein, and crRNA-guided site-specific DNA cleavage by the endonuclease Cas9 (Svitashev et al., Nature Communications, 2016, 7, 113274). In practice, the 20 nucleotides and the guide RNA backbone can be DNA (that can be expressed from a promoter to be transcribed in vivo (in cells) or in vitro (via T7 polymerase) to form an RNA guide or the guide RNA backbone can be chemically synthesized dual (crRNA and trRNA) guide or single guide RNA.

For a Cas9 protein/tracrRNA/crRNA complex to cleave a double-stranded DNA target sequence, the DNA target sequence can be adjacent to a cognate protospacer adjacent motif (PAM). By designing a crRNA to have an appropriate spacer sequence, the complex can be targeted to cleave at a locus of interest, e.g., a locus at which sequence modification is desired.

A variety of Type II CRISPR-Cas system crRNA and tracrRNA sequences, and associated predicted secondary structures are known in the art (see, e.g., Ran, F. A., et al., Nature, 2015, 520, 186-191; Fonfara et al., Nucleic Acids Research, 2014, 42, 2577-2590). The Type II CRISPR-Cas system of Ran et al., is provided herein in an aspect.

The spacer of Type II CRISPR-Cas systems can hybridize to a nucleic acid target sequence that is located 5' or 3' of a PAM, depending upon the Cas protein to be used. A PAM can vary depending upon the Cas polypeptide to be used. For example, if Cas9 from *S. pyogenes* is used, the PAM can be a sequence in the nucleic acid target sequence that comprises the sequence 5'-NRR-3', wherein R can be either A or G, N is any nucleotide, and N is immediately 3' of the nucleic acid target sequence targeted by the nucleic acid target binding sequence. Preferably, the PAM of *S. pyogenes* comprises 5'-NGG-3'. In another example, if Cas9 from *S. thermophilus* CRISPR3 (Sther CR3) is used, the PAM can be a sequence in the nucleic acid target sequence that comprises the sequence 5'-nGGnG-3'. (See Sapranaukas et al., Nucleic Acids Research, 2011, 39, 9275-9282).

Other Cas proteins recognize other PAMs, and one of skill in the art can determine the PAM for any particular Cas protein. A growing number of CAS proteins and systems are known in the art that are suitable for use with the targeting sequences of the present specification. See Shah et al., RNA Biol., 2013, 10, 891-899. A representative sample of CAS systems and their PAM sequences is provided in Table 1 below.

TABLE 1

Exemplary CRISPR/Cas Systems		
Species/Variant of Cas9	PAM Sequence	
<i>Streptococcus pyogenes</i> (SP); SpCas9	3' NGG	Wu et al., Nat Biotechnol., 2014, 32, 670-676
SpCas9 D1135E variant	3' NGG (reduced NAG binding)	Nishimasu et al., Science, 2018, 361, 1259-1262.
SpCas9 VRER variant	3' NGCG	Nishimasu et al. (2018)
SpCas9 EQR variant	3' NGAG	Nishimasu et al. (2018)
SpCas9 VQR variant	3' NGAN or NGNG	Nishimasu et al. (2018)
<i>Staphylococcus aureus</i> (SA); SaCas9	3' NNGRRT or NNGRR (N)	Kleinstiver et al., Nature, 2015, 523, 481-485
<i>Acidaminococcus</i> sp. (AsCpf1) and <i>Lachnospiraceae</i> bacterium (LbCpf1)	5' TTTV	Fagerlund, R., et al., 2015, Genome Biology, 16, 251
AsCpf1 RR variant	5' TYCV	Nishimasu et al., Mol Cell., 2017, 67, 139-147
LbCpf1 RR variant	5' TYCV	Nishimasu et al. (2017)
AsCpf1 RVR variant	5' TATV	Nishimasu et al. (2017)

TABLE 1-continued

Exemplary CRISPR/Cas Systems		
Species/Variant of Cas9	PAM Sequence	
<i>Neisseria meningitidis</i> (NM)	3' NNNNGAT	Hou et al., 2013, Proc. Natl. Acad. Sci. USA, 110, 15644-156449
<i>Treponema denticola</i> (TD)	3' NAAAC	Sun et al., Biotechnol J., 2018, 13, e1700588
<i>Campylobacter jejuni</i> (Cj)	NNNRYAC	Kim, et al., Nat Commun., 2017, 8, 14500.
<i>Streptococcus thermophilus</i> CR1 (St)	NNAGAAW	Toth et al., 2016, Biol Direct., 11, 46
<i>Streptococcus thermophilus</i> CR3	nGGnG	Müller et al., Mol Ther., 2016, 24, 636-644

Methods that rely on any CRISPR/CAS system can have a PAM sequence of NRR (e.g., NGG and NAA) as provided below for the *S. pyogenes* sequences or can have a PAM sequence of NGGNG as provided for the *S. thermophilus* sequences. Table 3 and the sequence listing provide the targeting sequences and chromosomal locations. It will also be appreciated by those of skill in the art that known CAS systems having different PAM requirements can be engineered to recognize and utilize the target sequences disclosed herein. Examples of modifications are presented in Table 1.

The terms “CRISPR/CasN” or “CRISPR/CasN system” refer to a programmable nuclease system for genetic editing that includes a CasN (e.g., Cas2, Cas5, Cas6, Cas9, etc.) protein, or derivative thereof, and one or more non-coding RNAs that can provide the function of a CRISPR RNA (crRNA) and trans-activating RNA (tracrRNA) for the CasN. The crRNA and tracrRNA can be used individually or can be combined to produce a “guide RNA” (gRNA). The crRNA or gRNA can provide a sequence that is complementary to the genomic target.

The term “Cpf1” or “CAS12” refers to another programmable RNA-guided endonuclease of a class 2 CRISPR-Cas system, described and used for gene editing purposes (Zetsche et al., *Cell*, 163:759-771, 2015). This system can use a non-specific endonuclease unit from the Cpf1 protein family, with a specificity of cleavage conferred by a single crRNA (lacking tracrRNA). Similar to Cas9, the Cpf1 coding sequence can be fused to UTR sequences described herein to improve its stability, and thus the efficiency of the resulting gene editing method.

As used herein, the terms “transcription activator-like effector nucleases” or “TALENs” refer to nucleases engineered to enable the targeted alteration of a given DNA sequence. TALENs can comprise a non-specific DNA-cleaving nuclease fused to a TALE DNA-binding domain that can be engineered to allow targeted gene editing. A “TALE DNA-binding domain” or “TALE” can be a polypeptide comprising one or more TALE repeat domains/units. The repeat domains can be involved in binding of the TALE to its cognate target DNA sequence. A single “repeat unit” (also referred to as a “repeat”) can be 33-35 amino acids in length and exhibits at least some sequence homology with other TALE repeat sequences within a naturally occurring

TALE protein. A “designed” DNA binding protein can be a protein not occurring in nature whose design/composition results principally from rational criteria. TALENs are discussed and disclosed in Joung et al., *Nat. Rev. Mol. Cell. Biol.*, 2013, 14, 49-55. A “selected TALE” can be a protein not found in nature whose production results primarily from an empirical process such as phage display, interaction trap, or hybrid selection. The present specification provides for, and includes, the use of TALENs to the pigs described herein.

“Resistance” or “disease resistance” refers to the extent to which an organism can defend itself from and/or withstand the attack of a pathogen and remain unaffected. An organism may demonstrate complete resistance, meaning it remains virtually unaffected by a pathogen. Alternatively, an organism may demonstrate partial resistance, wherein the extent to which the pathogen affects the organisms can be less than a comparable organism with no resistance. Resistance may stem from a particular characteristic of the organism, allowing the organism to avoid the outcome of organism-pathogen interactions. Resistance can be demonstrated by the extent to which an organism can avoid the disease symptoms associated with, or reduce the incidence/severity of clinical signs, or reduce the clinical symptoms associated with a pathogen.

The terms “increased resistance” and “reduced susceptibility” refer to a statistically significant reduction of the incidence and/or severity of clinical signs or clinical symptoms which are associated with infection by a given pathogen. For example, “increased resistance” or “reduced susceptibility” refer to a statistically significant reduction of the incidence and/or severity of clinical signs or clinical symptoms which are associated with infection by PRRSV in an animal comprising a deleted or inactivated chromosomal sequence in a CD163 gene protein as compared to a control animal having an unmodified chromosomal sequence. The term “statistically significant reduction of clinical symptoms” means, but is not limited to, the frequency in the incidence of at least one clinical symptom in the modified group of subjects and, in some aspects, clinical symptoms may be statistically reduced by at least 80% lower than in the non-modified control group after the challenge with the infectious agent.

As used herein, the terms “reduction of the incidence and/or severity of clinical signs” or “reduction of clinical

symptoms” mean reducing the number of infected subjects in a group, reducing or eliminating the number of subjects exhibiting clinical signs of infection, or reducing the severity of any clinical signs that are present in one or more subjects, in comparison to wild-type infection in an otherwise similar genetic background. For example, these terms encompass any clinical signs of infection, lung pathology, viremia, antibody production, reduction of pathogen load, pathogen shedding, reduction in pathogen transmission, or reduction of any clinical sign symptomatic of PRRS when compared to an otherwise similar background. Comparisons of clinical signs between non-edited CD163 pigs and CD163 edited pigs can be individually or between herds. In an aspect, an individual pig does not present clinical signs. In an aspect, a herd of CD163 edited pigs present reduced clinical signs. In an aspect, the size of the herd can be at least 100 animals. In an aspect, CD163 edited pigs of the present specification can have reduced clinical signs of reproductive syndrome. In an aspect, a herd of CD163 edited pigs can have a reduced number of premature farrowings, reduced numbers of stillborn or mummified piglets, reduced numbers of PRRSv-positive piglets, or reductions in delays to return to service of sow. In an aspect, CD163 edited pigs of the present specification can have reduced clinical signs in sows and gilts including, but not limited to, reduced anorexia, reduced fever, reduced lethargy, reduced pneumonia, reduced agalactica, and reduced subcutaneous and hind limb edema. In an aspect, gilts and sows that are CD163 edited pigs of the present specification can have reduced red/blue discoloration of the ears and vulva. In an aspect, CD163 edited sows can exhibit reduced delays to return to estrus after weaning. The present specification provides for, and includes, reduced deaths among a herd of CD163 edited gilts and sows. Also included and provided for by the present specification are reduced clinical signs in finishing pigs. In an aspect, a herd of CD163 edited finishing pigs can have reduced respiratory clinical signs selected from the group consisting of fever, sneezing, hyperpnoea, dyspnea, coughing, pneumonia, lethargy, periocular edema, and oculo-nasal discharge. Preferably these clinical signs can be reduced in one or more animals of the present teachings by at least 10% in comparison to subjects not having a modification in the CD163 gene and having a similar background and that become infected. In an aspect, clinical signs can be reduced in subjects of the present teachings by at least 80%. In another aspect, clinical signs can be reduced in pigs of the present teachings by at least 85%. In a further aspect, clinical signs can be reduced in pigs of the present teachings by at least 90%. In yet another aspect, clinical signs can be reduced in pigs of the present teachings by at least 95%. In aspects according to the present disclosure, clinical signs can be reduced by 80% to 100% relative to non-edited CD163 pigs.

The term “breeding” as used herein refers to a process comprising the selection of superior male and superior female animals use for creation of the next generation of offspring. This process further comprises the union of male and female gametes so that fertilization occurs. Such a union may be brought about by mating (copulation) or by in vitro or in vivo artificial methods. Such artificial methods can include, but are not limited to, artificial insemination, surgical assisted artificial insemination, in vitro fertilization, intracytoplasmic sperm injection, zona drilling, in vitro culture of fertilized oocytes, ovary transfer, and ovary splitting. The term “breeding” as used herein also can include

transferring of a fertilized oocyte into the reproductive tract of a female animal in order to allow for more offspring from a particular elite female.

As used herein, “reference sequence” is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length promoter sequence, or the complete promoter sequence. In an aspect, the reference sequence can be the Sscrofa11.1 reference genome (GenBank accession: GCA_000003025.6).

The terms “percent identity” or “percent identical” as used herein in reference to two or more nucleotide or protein sequences is calculated by (i) comparing two optimally aligned sequences (nucleotide or protein) over a window of comparison, (ii) determining the number of positions at which the identical nucleic acid base (for nucleotide sequences) or amino acid residue (for proteins) occurs in both sequences to yield the number of matched positions, (iii) dividing the number of matched positions by the total number of positions in the window of comparison, and then (iv) multiplying this quotient by 100% to yield the percent identity. If the “percent identity” is being calculated in relation to a reference sequence without a particular comparison window being specified, then the percent identity can be determined by dividing the number of matched positions over the region of alignment by the total length of the reference sequence. Accordingly, for purposes of the present application, when two sequences (query and subject) are optimally aligned (with allowance for gaps in their alignment), the “percent identity” for the query sequence can be equal to the number of identical positions between the two sequences divided by the total number of positions in the query sequence over its length (or a comparison window), which is then multiplied by 100%. When percentage of sequence identity is used in reference to proteins, it is recognized that residue positions which are not identical often can differ by conservative amino acid substitutions, where amino acid residues can be substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity can be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have “sequence similarity” or “similarity.”

For optimal alignment of sequences to calculate their percent identity, various pair-wise or multiple sequence alignment algorithms and programs are known in the art, such as ClustalW or Basic Local Alignment Search Tool (BLAST®, National Library of Medicine, Bethesda, MD), etc., that can be used to compare the sequence identity or similarity between two or more nucleotide or protein sequences. Although other alignment and comparison methods are known in the art, the alignment and percent identity between two sequences (including the percent identity ranges described above) can be as determined by the ClustalW algorithm, see, e.g., Chenna, R., et al., *Nucleic Acids Research*, 2003, 31, 3497-3500; Thompson, J. D., et al., *Nucleic Acids Research*, 1994, 22, 4673-4680; Larkin M A et al., *Bioinformatics*, 2007, 23, 2947-2948; and Altschul, S. F., et al., *J. Mol. Biol.*, 1990, 215, 403-410.

Identity to a sequence used herein can be expressed in terms of a percent identity between two sequences as determined according to alignment of the two sequences. The present specification provides for repair template sequences that can have at least 80% identity to a contiguous

region of a genome. In an aspect, the repair template can have a 5' region and a 3' region sharing at least 80% identity to a contiguous region of the genome and a core region having identity to the chromosome sequences flanking the intended gene edit site.

The present specification provides for, and includes, but is not limited to, target sequences that can be 100% identical to the target sequences selected from the group consisting of SEQ ID NOs: 22 to 271 and 347 to 425. These target sequences can comprise a PAM sequence such as those listed on Table 1. Accordingly, when incorporated into a guide RNA, in an aspect the spacer region can share 100% identity to a sequence selected from the group of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. Also included are spacer sequences that can have from 15 to 20 nucleotides of each of SEQ ID NOs: 22 to 271 or from 15 to 20 nucleotides of each of SEQ ID NOs: 347 to 425. (The remainder of the sequence can comprise the PAM sequence which can be present in the genome, but not the gRNA molecules.) Also included and provided for are guide RNA spacer sequences that can have one or more mismatches to the target sequence. In an aspect, the mismatch can be at the distal end of the sequence target sequence to ensure that identity at the endonuclease cleavage site is maintained. Typically, the mismatches can occur at the 5' end of the target sequence relative to the nuclease site at the 3' end. In an aspect, a target sequence, or the spacer sequences of a guide RNA prepared therefrom, may have a single mismatch. In another aspect, a target sequence, or the spacer sequences of a guide RNA prepared therefrom, may have two (2) mismatches. In another aspect, a sequence, or the spacer sequences of a guide RNA prepared therefrom, may have three (3) mismatches. In another aspect, a sequence, or the spacer sequences of a guide RNA prepared therefrom, that may have less than four (4) mismatches are included. In some aspects, the mismatch regions can be limited to terminal nucleotides distal to the PAM sequences and cleavage site.

In an aspect, the target sequences and the spacer sequence of the gRNA guide can have at least 90% identity to a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. In a further aspect, the RNA guide can have at least 90% identity and 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 95% identity. In a further aspect, the RNA guide can have at least 95% identity or 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 96% identity. In a further aspect, the RNA guide can have at least 96% identity or 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 97% identity. In a further aspect, the RNA guide can have at least 97% identity or 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 98% identity. In a further aspect, the RNA guide can have at least 98% identity or 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 99% identity. In a further aspect, the RNA guide can have at least 99% identity or 100% identity at 15 nucleotides at the 3' end of the sequence. In an aspect, the target sequences and the RNA guide can have at least 100% identity. Also included and provided for by the present specification are spacer

sequences that can comprise the first 15 nucleotides each of SEQ ID NOs: 22 to 271 or the first 15 nucleotides each of SEQ ID NOs: 347 to 425. In an aspect, the spacer sequences can comprise the first 16 nucleotides in each of SEQ ID NOs: 22 to 271 or the first 16 nucleotides in each of SEQ ID NOs: 347 to 425. In another aspect, the spacer sequences can comprise the first 17 nucleotides in each of SEQ ID NOs: 22 to 271 or the first 17 nucleotides in each of SEQ ID NOs: 347 to 425. In an aspect, the spacer sequences can comprise the first 18 nucleotides in each of SEQ ID NOs: 22 to 271 or the first 18 nucleotides in each of SEQ ID NOs: 347 to 425. In yet another aspect, the spacer sequences can comprise the first 19 nucleotides in each of SEQ ID NOs: 22 to 271 or the first 19 nucleotides in each of SEQ ID NOs: 347 to 425.

The preferred meaning of "fertilization" as used herein to refer to pig breeding encompasses any technique that produces a viable embryo. Fertilization can include both insemination of female pigs, and in vitro or ex vivo fertilization. There are three commonly available ways to inseminate a sow, namely traditional artificial insemination (AI), intrauterine insemination (IUI), and deep intrauterine insemination (DIUI). These techniques all rely on providing a dose of semen, typically fresh and unfrozen, for insemination. In vitro fertilization (IVF) can be the harvesting of unfertilized oocytes(s) and the subsequent fertilization of those oocytes with semen in vitro (i.e., in the laboratory) instead of in vivo (i.e., in the live animal; insemination discussed above) as in standard ET. The fertilized oocytes(s) or embryo(s) from the oocyte donor can then be transferred into another female (embryo recipient). Embryo transfer (ET) can be the harvesting of fertilized oocytes(s) or embryo(s) from one female (embryo donor) and transfer of those embryo(s) into another female (embryo recipient) whose reproductive status can be synchronized with that of the donor.

As described in Cameron et. al. (*Nat Methods*, 2017, 14, 600-606), the CRISPR-Cas9 system can be used for genome editing in both basic research and biotechnology. The application of CRISPR and related technologies to gene or genome editing can be accompanied by unwanted off-target cleavage activity and possible pathogenic and other negative phenotypic consequences. To minimize off-target cleavage, a variety of genome-wide experimental methods have been recently developed but some methods are potentially biased due to cellular context or inefficiencies in recovering relevant cleavage sites. One approach, the SITE-SEQ® assay (Caribou Biosciences, Inc., Berkeley CA) disclosed in Cameron et. al. enables one to comprehensively list Cas9 cleavage sites in a sample genome, then probe those sites for cellular off-target editing in follow-up experiments through (1) extraction and purification of high-molecular weight genomic DNA, (2) execution of Cas9 Ribonucleoprotein (RNP) cleavage in vitro, (3) fragmentation, adapter ligation, and affinity purification to enrich for Cas9 cleaved fragments, and (4) amplification and indexing of SITE-SEQ® libraries for ILLUMINA® (ILLUMINA®, San Diego, CA) sequencing. (See Cameron et al., *Nat Methods*, 2017, 14, 600-606.) Various other methods of discovering and reducing the number of potential off-target edits are known in the genome editing arts. While such bioinformatics tools can be invaluable for identifying suitable target sequences, the error rate is generally higher than desired and the approach can be limited when the system lacks extensive sequence data. Further, the fidelity of DSB break repair can vary, thus confounding the ability to predict and prevent unwanted off target effects. Further testing of in silico selected target

sequences can identify select sequences that have higher efficiencies and fewer off target cuts. Testing in cell culture systems combined with high-throughput sequencing methods can be used to identify poor performing target constructs.

For example, the type II CRISPR system, which is derived from *Streptococcus pyogenes* (*S. pyogenes*), can be reconstituted in mammalian cells using Cas9, a specificity-determining CRISPR RNA (crRNA) comprising a backbone sequence and a sequence selected from the group consisting of the first twenty nucleotides of each of SEQ ID NOs: 22 to 271 and the first twenty nucleotides of each of SEQ ID NOs: 347 to 425, and an auxiliary trans-activating RNA (tracrRNA). The term “off target effect” broadly refers to any impact distinct from and not intended as a result of the on-target treatment or procedure. Examples of off-target edits can include double strand breaks at unintended locations that lead to DNA insertions of unintended nucleotides or repair templates, deletions, or rearrangements. For some systems, for example the *S. pyogenes* and *S. thermophilus* CRISPR system, the crRNA and tracrRNA duplexes can be fused to generate a single-guide RNA (sgRNA). The first 20 nucleotides of the sgRNA can be complementary to the target DNA sequence and can be the spacer region, and those 20 nucleotides can be followed at the 3' end by a protospacer adjacent motif (PAM) in the genome (but not the guide). In an aspect, the first 20 nucleotides of the sgRNA and PAM sequence can be a sequence selected from the group consisting of SEQ ID NOs: 22 to 271 and 347 to 425. Accordingly, as provided below at Table 2 and SEQ ID NOs: 22 to 271 and 347 to 425, specific targeting of the Cas nuclease from either *S. pyogenes* or *S. thermophilus* can be accomplished combining the crRNAs with a backbone sequence. Although the 20-nucleotide guide sequence plus PAM sequence (e.g., 23 to 25 nucleotides) of the sgRNA can provide tightly controlled targeting and cleavage, it has been discovered that off-target cleavage activity can still occur on DNA sequences with between 3-5 base pair mismatches in the PAM-distal part of the sgRNA-guiding sequence. Further, different types of guide RNA structures actually affect the cleavage precision, increasing or decreasing cleave on off-target sites. Various techniques, as well as a further description of off-target cleavage, are reviewed in Zhang et al., *Mol Ther Nucleic Acids*, 2015, 4, e264. The mechanisms and effects of off-target cleavage are still poorly understood, meaning it can be difficult to predict and to compensate for its effects. However, the consequences can be severe; off-target cleavage can ultimately lead to genomic instability and disrupt the functionality of otherwise normal genes.

The present specification provides for, and includes, pigs that can have inactivating edits in both alleles of the CD163 gene and that can be resistant to infection with PRRSV. The CD163 pigs disclosed herein further do not comprise new sequences or polypeptides, nor do they comprise non-native amino acids resulting from frameshifts or missense mutations.

The gene edits disclosed herein allow for the protection of CD163-positive fetuses (e.g., fetuses that have one or two wild-type CD163 alleles). CD163-positive fetuses can be protected from PRRSV infection while in utero so long as the dam possesses inactivating edits in both alleles of her CD163 genes (PCT/US2018/027944). For example, dams having inactivating edits in both alleles of the CD163 gene can be mated with males having two wild-type CD163 alleles, and the resulting heterozygous fetuses will be protected from PRRSV infection in utero.

In an aspect, pigs having inactivating edits in one allele of CD163 can be generated using the methods of the present specification. These pigs having heterozygous CD163 alleles (one edited, one wild-type) can be bred with other pigs also having heterozygous CD163 alleles or homozygous edited CD163 alleles, and offspring from this breeding having homozygous edited CD163 alleles can be selected for resistance to infection with PRRSV. The present disclosure also provides for, and includes, porcine animals and populations, and methods for creating or improving porcine animals and populations, in which the animals can be homozygous for one or more particular genetic markers or alleles. In various aspects, the present disclosure provides for, and includes, animals, and methods for creating or improving crossing porcine animals and populations, by generating animals that can be heterozygous for one or more particular genetic markers or alleles, and crossing said animals with other animals heterozygous for one or more of the particular genetic markers or alleles to produce animals that can be homozygous for the one or more particular genetic markers or alleles. In some aspects, multiple rounds of crossing and/or back-crossing may be required to obtain homozygosity for each of the particular genetic markers and alleles. In other aspects, a single cross may be sufficient to obtain homozygosity.

Various techniques known in the art can be used to inactivate genes to make knock-out animals and/or to introduce edited genes into animals to produce founder animals and to make animal lines, in which the knockout or nucleic acid construct can be integrated into the genome. Such techniques can include, without limitation, pronuclear microinjection (U.S. Pat. No. 4,873,191), retrovirus mediated gene transfer into germ lines (Van der Putten et al., *Proc. Natl. Acad. Sci. USA*, 1985, 82, 6148-1652), gene targeting into embryonic stem cells (Thompson et al., *Cell*, 1989, 56, 313-321), electroporation of embryos (Lo, *Mol. Cell. Biol.*, 1983, 3, 1803-1814), sperm-mediated gene transfer (Lavitrano et al., *Proc. Natl. Acad. Sci. USA*, 2002, 99, 14230-14235; Lavitrano et al., *Reprod. Fert. Develop.*, 2006, 18, 19-23), and in vitro transformation of somatic cells, such as cumulus or mammary cells, or adult, fetal, or embryonic stem cells, followed by nuclear transplantation (Wilmut et al., *Nature*, 1997, 385, 810-813 and Wakayama et al., *Nature*, 1998, 394, 369-374). Pronuclear microinjection, sperm mediated gene transfer, and somatic cell nuclear transfer can be other useful techniques. An animal can be genomically edited in that all of its cells have the edit, including its germ line cells. When methods are used that produce an animal that is mosaic in its modification, the animals may be mosaic and can be further inbred and progeny that are genomically edited can be produced and selected using standard methods. Cloning, for instance, may be used to make a mosaic animal if its cells are edited at the blastocyst stage, or genomic modification can take place when a single-cell is edited. In an aspect, an inactivated knock-out edit can be homozygous.

In an embryo/zygote microinjection aspect, a nucleic acid construct, mRNA, protein, polynucleotides, or combinations thereof, can be introduced into a fertilized egg. In an aspect, one or two cell fertilized eggs can be used. One and two cell fertilized eggs can provide a visible nuclear structure containing the genetic material from the sperm head and the egg within the protoplasm. In an aspect, pronuclear staged fertilized eggs can be obtained in vitro or in vivo (i.e., surgically recovered from the oviduct of donor animals). In another aspect, in vitro fertilized eggs can be produced, for example, from collected swine ovaries by follicles aspirated

using methods known in the art. See, e.g., Agung et al., *J Reprod Dev.*, 2013, 59, 103-106 and Appeltant et al., *J Reprod Dev.*, 2016, 62, 439-449. In an aspect, mature oocytes can be provided for use in the in vitro fertilization methods. In another aspect, mature oocytes can be injected with the CRISPR/Cas gene editing system of the present specification.

The present specification further provides for in vitro fertilization of mature oocytes. In an aspect, the oocytes can be matured in vitro as provided above. In another aspect, mature oocytes can be collected from gilts. In vitro fertilization is performed according to established methods. See Appeltant et al., *J Reprod Dev.*, 2011, 57, 9-16, and Fowler et al., *Reprod Biol.*, 2018, 18, 203-211.

Zygotes or embryos can be obtained for germline editing by artificial insemination and flushing. The collected zygotes or embryos can then be edited using the methods provided herein. In an aspect, linearized nucleic acid constructs, mRNA, proteins, polynucleotides or combinations thereof can be injected or otherwise introduced, for example, by electroporation, into one of the pronuclei or into the cytoplasm of a zygote or embryo post-fertilization, or into a gamete cell pre-fertilization. In an aspect, a pre-formed RNP complex comprising the Cas nuclease protein and a guide RNA comprising a backbone of SEQ ID NO: 19 and a targeting sequence selected from any one of SEQ ID NOs: 22 to 271 and 347 to 425 can be prepared and injected into the embryo, zygote, or oocyte. In another aspect, a pre-formed RNP complex comprising the Cas nuclease and a guide RNA comprising a backbone of SEQ ID NO: 19 and a spacer sequence selected from an RNA sequence of a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 can be prepared and injected into the embryo, zygote, or oocyte, together with a repair template comprising SEQ ID NOs: 1 to 13 listed in Table 6. Also included, and provided for, by the present specification are the combinations of spacer sequences and repair templates that can have the sequences of SEQ ID NOs: 1 to 13 as recited in Table 6. In an aspect, the injected zygotes or embryos can be transferred to a recipient female (e.g., into the oviducts of a recipient female) and allowed to develop in the recipient female to produce the transgenic or gene edited animals. In an aspect, the methods further provide for in vitro or in vivo fertilized zygotes or embryos that can be centrifuged at 15,000×g for 5 minutes to sediment lipids allowing visualization of the pronucleus. The zygotes or embryos can be injected with using an EPPENDORF® FEMTOJET® (EPPENDORF® AG, Germany) injector and can be cultured until blastocyst formation. Rates of embryo cleavage and blastocyst formation and quality can be recorded.

In an aspect, the CRISPR/Cas editing system and gRNA can be provided as a nucleic acid construct. In another aspect, the Cas nuclease may be provided as an mRNA together with a guide RNA comprising a backbone sequence and a targeting sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. In another aspect, the Cas nuclease may be provided as an mRNA together with a guide RNA comprising a backbone sequence and a targeting sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 together with a repair template listed in Table 6 and selected from the group consisting of SEQ ID NOs: 1 to 13. In a further aspect, the gene targeting complex can be provided by microinjection of a transcribable DNA encoding a Cas nuclease and a guide RNA comprising a

backbone sequence and a targeting sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. In aspects according to the present specification, the backbone sequence can be a sequence of SEQ ID NO: 19. Also included and provided for by the present specification are the combinations of targeting sequences and repair templates that can have one of the sequences recited in SEQ ID NOs: 1 to 13. It would be understood by persons of skill in the art that various combinations of nuclease, gRNA, and optionally repair template sequences, can be introduced into oocytes, zygotes, blastula, and embryos to achieve the goals of the present methods. It will be further understood that other backbone sequences can be used in conjunction with the targeting sequences of SEQ ID NOs: 22 to 271 and SEQ ID NOs: 347 to 425 for use with other Cas nucleases. Any Cas nuclease having a PAM sequence of NGG or NGGNG can be suitable for the preparation of a backbone sequence for combination with the targeting sequences of the present specification. In various aspects, the desired edit can have a final genomic sequence of SEQ ID NOs: 426 to 505. In some configurations, the desired edit lacks exon 7 and can have a final nucleotide sequence as set forth in SEQ ID NOs: 426 to 458. In various configurations, the final genomic sequence can include an exogenous stop codon and can have a final genomic sequence as set forth in SEQ ID NOs: 459-504. In various configurations, these partial CD163 genes can have an exon 7 amino acid sequence as set forth in SEQ ID NOs: 506 to 517.

In an aspect, the injected zygotes or embryos can be transferred to a recipient female (e.g., into the oviducts of a recipient female) and allowed to develop in the recipient female to produce the transgenic or gene edited animals. In an aspect, the methods further provide for in vitro or in vivo fertilized zygotes or embryos that can be centrifuged at 15,000×g for 5 minutes to sediment lipids allowing visualization of the pronucleus. The zygotes or embryos can be injected using, for example, an EPPENDORF® FEMTOJET® injector and can be cultured until blastocyst formation. Rates of embryo cleavage and blastocyst formation and quality can be recorded.

The present specification provides for, and includes, a recipient sow having one or more embryos that can be injected with a CRISPR/Cas/gRNA combination according to the present disclosure. In an aspect, a pre-formed RNP complex can be provided comprising the Cas nuclease and a guide RNA comprising a backbone of SEQ ID NO: 19 and a spacer sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. In various aspects, two pre-formed RNP complexes can be provided comprising the Cas nuclease and a guide RNA comprising a backbone of SEQ ID NO: 19 and each comprising a different spacer sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. Also provided for, and included, are recipient sows that can have one or more embryos injected with a pre-formed RNP complex that can comprise the Cas nuclease and a guide RNA comprising a targeting sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347-425. In an aspect, the pre-formed RNP complex can be prepared and injected into the embryo, zygote, or oocyte, together with a repair

template listed in Table 6 (SEQ ID NOs: 1 to 13). In an aspect, a recipient sow can be a sow of a different line than the donor embryos.

Embryos or zygotes can be surgically transferred into uteri of synchronous recipients. Typically, 100-200 (or 150-200) embryos or zygotes can be deposited into the ampulla-isthmus junction of the oviduct using a catheter. After surgery, real-time ultrasound examination of pregnancy can be performed. In an aspect, the present specification can include a recipient sow having 1 to 100, but more typically 30 to 60 CRISPR/Cas/gRNA combination treated embryos wherein said embryos can comprise a gene edited CD163 gene comprising a sequence of SEQ ID NOs: 1 to 18 or 426 to 505. In an aspect, the transferred embryos can comprise a mosaic of edited cells. Also included are CD163 gene edited embryos that can be non-mosaic.

Methods of improving the health of a livestock animal or herd of livestock can comprise modifying a target sequence in the genome of an animal cell to form a truncated CD163 polypeptide. Also provided for, and included, are improved animals that can have truncated CD163 polypeptides. In some aspects, the predicted truncated polypeptides do not include non-native amino acids. Without being bound by theory, the truncated gene product can be rapidly digested in vivo by cellular proteases. In an aspect of the present specification, the truncated CD163 can result in the complete elimination of the protein. In a further aspect, the truncated CD163 can be non-detectable in vivo. In an aspect, the truncated CD163 can be non-detectable by immunofluorescence labelling and FACS analysis. In yet another aspect, the truncated CD163 can be non-detectable when using any of the expression analysis methods provided in the specification, including those provided in Example 5. In yet another aspect, the truncated CD163 can be non-detectable when using any of the expression analysis methods provided in the specification, including those provided in Example 8.

In another aspect, a gene edited CD163 pig can comprise a truncated CD163 gene with no more than 25 non-native amino acids. Not to be limited by theory, truncated CD163 proteins can be destabilized and targeted for degradation within the cell. Accordingly, CD163 protein sequences and CD163 polypeptides comprising non-native amino acids can be non-detectable. In another aspect, the predicted truncated CD163 protein can comprise no more than 211 amino acids of the native protein. In an aspect, the truncated CD163 protein can be predicted to comprise no more than 144 amino acids of the native protein. In an aspect, the truncated CD163 protein can be predicted to comprise no more than 133 amino acids of the native protein. In yet another aspect, the truncated CD163 protein can be predicted to comprise no more than 129 amino acids of the native protein. In an aspect, the truncated CD163 protein can be predicted to comprise no more than 113 amino acids of the native protein. In yet another aspect, the truncated CD163 protein can be predicted to comprise no more than 108 amino acids of the native protein. In a further aspect, the truncated CD163 protein can be predicted to comprise no more than 93 amino acids of the native protein. In yet another aspect, the truncated CD163 protein can be predicted to comprise no more than 74 amino acids of the native protein. As provided herein, truncated CD163 proteins can comprise between 32 to 211 amino acids of native CD163 polypeptide sequences and no more than 25 non-native amino acids and can be undetectable in cells and cell extracts.

In other aspects, altered CD163 protein can have a truncation with a single amino acid substitution. In some

aspects, the altered CD163 protein can have a predicted amino acid sequence of no more than 1,010 amino acids.

The present specification provides for, and includes, gene edited CD163 pigs that can have truncations of the CD163 protein (e.g., amino acids 1 to 40 of CD163 reference sequence NP 999141). Not to be limited by theory, it is believed that truncations within the signal peptide (also known as the signal sequence) can result in a failure of the protein to translocate to the cellular membrane and the polypeptide is subsequently degraded within the cell. Accordingly, CD163 proteins having signal peptide truncations can result in CD163 null animals that are resistant to infection by PRRSV. In an aspect, the gene edited CD163 animals having proteins truncated in the signal sequence can comprise no more than 25 amino acids of non-native amino acid sequences. In an aspect, the gene edited pigs can comprise no more than the first 39 amino acids of the native CD163 protein. In an aspect, the gene edited pigs can comprise no more than the first 39 amino acids of the native CD163 protein and no more than 15 non-native amino acids. Also included, and provided for, by the present specification are gene edited pigs that can be predicted to comprise no more than the first 38 amino acids of the native CD163 protein. In an aspect, the gene edited pigs can comprise no more than the first 38 amino acids of the native CD163 protein and no more than 15 non-native amino acids. Also included, and provided for, by the present specification are gene edited pigs that can be predicted to comprise no more than the first 36 amino acids of the native CD163 protein. In an aspect, the gene edited pigs can comprise no more than the first 36 amino acids of the native CD163 protein and no more than 15 non-native amino acids. Truncated proteins can be non-detectable using methods known to those of skill in the art.

In an aspect, the gene edited pigs can be predicted to comprise no more than the first 34 amino acids of the native CD163 protein and no more than 15 non-native amino acids. In an aspect, the gene edited pigs can be predicted to comprise no more than the first 32 amino acids of the native CD163 protein and no more than 15 non-native amino acids. Truncated proteins can be non-detectable using methods known to those of skill in the art.

Further, the present disclosure provides for and includes truncations of the CD163 protein that can include deletion of exon 7 (SEQ ID NOs: 426-458), sequences where an exogenous stop codon is introduced into exon 7 (SEQ ID NO: 459-504). These truncations can include CD163 proteins comprising no more than 1,010 amino acids.

Breeding techniques can be used to create animals that are homozygous for the inactivated gene from the initial homozygous or heterozygous founder or other heterozygous animals. Homozygous animals can be generated. Gene edited pigs described herein can be bred with other edited or wild-type pigs of interest to ultimately generate pigs that are homozygous or heterozygous for the edited gene. In an aspect, the homozygous animal can be an animal of any one of the lines generated by crosses with PIC™ elite porcine lines 2, 3, 15, 19, 27, 62, 65, and combinations thereof. In an aspect, the homozygous animal can be a hybrid animal prepared by a cross between animals of Line 2 and Line 3.

Gene editing has been used to address various diseases, including PRRS, in swine. While there have been several gene editing technologies developed over the past 15 years, generally these platforms can be designed to introduce a double-stranded break at a specific region of a genome. The introduced break can then be repaired by the cell's own machinery.

One repair pathway, non-homologous end-joining (NHEJ), is evolutionarily conserved throughout all kingdoms of life and is the predominant double-strand break repair pathway in mammalian cells. This repair process can result in the random insertion or deletion of nucleotides across the cut site. As a practical matter, this incomplete fidelity can be used to advantage by providing in trans a DNA repair template. In an aspect, a DNA repair template carrying an alternative allele of the targeted site (for example, a single base polymorphism or a single or multiple base insertion or deletion) can be co-delivered with the gene editing reagents, and through the use of the cell's homologous recombination machinery, the DNA repair template can direct the repair to generate a new allele. In aspects according to the present specification, DNA repair templates can be designed and selected based on the location of high efficiency of CRISPR/CAS cleavage (e.g., high editing frequency from Table 3) and the ability to introduce deletions that can result in truncated proteins that comprise only wild type sequences. Repair templates can increase the efficiency and accuracy of gene editing. DNA repair templates suitable for the methods of the present application can include SEQ ID NOs: 1 to 13. In an aspect, the DNA repair templates can be paired with a guide sequence as described below in Table 6. Unlike transgenesis, genome editing according to the present specification does not result in the introduction of foreign DNA sequences into the genome.

However, as these exogenously added DNA repair templates have the possibility to randomly integrate in the genome, it is also advantageous to identify and use guide pairs that result in the deletion of DNA sequences such that the joined ends can result in the generation of an in-frame translational stop codon across the joined ends; when the cut sites of two guides are repaired by NHEJ in an end-to-end manner, this new DNA sequence, when transcribed into mRNA and translated into protein could terminate the production of the CD163 protein. Although function of this terminated CD163 protein can be lost, often, but not always, premature termination of protein synthesis can result in an unstable polypeptide that can be degraded and not detectable by standard methods. Guide pairs that cut exon 7 of the CD163 gene such that the joined ends form an exogenous stop codon can include SEQ ID NOs: 351 and 365, SEQ ID NOs: 351 and 387, SEQ ID NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 348 and 395, SEQ ID NOs: 352 and 365, SEQ ID NOs: 352 and 387, SEQ ID NOs: 352 and 399, SEQ ID NOs: 353 and 365, SEQ ID NOs: 353 and 387, SEQ ID NOs: 353 and 399, SEQ ID NOs: 354 and 390, SEQ ID NOs: 354 and 388, SEQ ID NOs: 354 and 395, SEQ ID NOs: 358 and 361, SEQ ID NOs: 358 and 362, SEQ ID NOs: 358 and 368, SEQ ID NOs: 358 and 384, SEQ ID NOs: 358 and 394, SEQ ID NOs: 358 and 399, SEQ ID NOs: 359 and 390, SEQ ID NOs: 359 and 388, SEQ ID NOs: 359 and 395, SEQ ID NOs: 360 and 368, SEQ ID NOs: 360 and 384, SEQ ID NOs: 360 and 389, SEQ ID NOs: 360 and 394, SEQ ID NOs: 360 and 397, SEQ ID NOs: 361 and 365, SEQ ID NOs: 361 and 387, SEQ ID NOs: 362 and 390, SEQ ID NOs: 362 and 388, SEQ ID NOs: 362 and 395, SEQ ID NOs: 364 and 365, SEQ ID NOs: 364 and 387, SEQ ID NOs: 364 and 399, SEQ ID NOs: 365 and 368, SEQ ID NOs: 365 and 384, SEQ ID NOs: 365 and 389, SEQ ID NOs: 365 and 394, SEQ ID NOs: 365 and 397, SEQ ID NOs: 366 and 368, SEQ ID NOs: 366 and 384, SEQ ID NOs: 366 and 389, SEQ ID NOs: 366 and 394, and SEQ ID NOs: 366 and 397.

In one aspect, the present teachings can involve a simple, precise and reproducible single CD163 loss of function edit in elite pigs such that the edit occurs early in the CD163

gene. A single guide RNA-Cas protein combination and DNA repair template can be selected for directing a specific gene edit in elite pigs, based on several considerations and approaches: efficient cutting near the 5' end of the CD163 gene in tissue culture, bioinformatic review identifying guides with few mismatches in the pig genome, high specificity of on-target cutting as determined by biochemical prescreen, and the ability to target specific gene edits at an intended site in pig embryo-like cells. Examples of specific gene edits of CD163 are discussed in detail below. In an aspect, a specific gene edit according to the present specification can include a CD163 gene comprising a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426 to 505. In an aspect, cells comprising an edited CD163 gene can comprise at least one allele having a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426 to 505. In an aspect, cells comprising two gene edited alleles of the CD163 gene can comprise, at both alleles, a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426 to 505. In an aspect, both gene edited CD163 genes can comprise the same sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426-505 (e.g., two alleles of SEQ ID NO: 1 or two alleles of SEQ ID NO: 2, etc.). In an aspect, the genome of a cell can comprise gene edited CD163 genes comprising one each of a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426-505 (e.g., an allele of SEQ ID NO: 1 with an allele of SEQ ID NO: 2, and all combinations thereof).

Also provided for, and included, are mixtures of cells that can comprise CD163 edited cells and non-gene edited cells. In an aspect, the mixture can be an embryo. In another aspect, the mixture can be a cell culture. In a further aspect, the mixture of cells does not comprise a reproductive cell. In an aspect, the present specification can provide for, and includes, a tissue culture of CD163 edited cells and methods to prepare such cultures. Cultures according to the present specification can include mixtures of CD163 edited cells (e.g., cells having an allele of SEQ ID NO: 1 with cells having an allele of SEQ ID NO: 2, cells having an allele of SEQ ID NO: 1 with cells having an allele of SEQ ID NO: 3, and all combinations thereof). In an aspect, the tissue culture of non-reproductive cells can include cells comprising a single CD163 edit. In a further aspect, cells of a single CD163 edit in culture may comprise one or both edited alleles.

Specific examples of endonuclease and guide RNA backbone sequences that can be used in the products and methods of the present teachings are listed in Table 2.

TABLE 2

Endonuclease and Guide Backbone Sequences	
Sequence Identity	SEQ ID NO:
Guide RNA Backbone (DNA sequence shown)	19
<i>Streptococcus pyogenes</i> endonuclease protein sequence	20
<i>Streptococcus thermophilus</i> CR3 endonuclease protein sequence	21

Elite porcine nucleus lines can be sequenced and aligned against a public reference for the CD163 gene. These data can be used for the development of gene editing reagents for the PRRS-resistance project. These sequences can be scanned for the presence of conserved RNA-guided

CRISPR-Cas9 recognition sites which can consist of the 3 nucleotide or 5 nucleotide motif, nGG (AGG, CGG, TGG, GGG), nGGnG (AGGTG, CGGTG, TGGTG, GGGTG, AGGGG, CGGGG, TGGGG, GGGGG, AGGAG, CGGAG, TGGAG, GGGAG, AGGCG, CGGCG, TGGCG, GGGCG), respectively. This motif, called the PAM sequence, can be located adjacent to, and 3' of, a 20 nucleotide spacer sequence which can be used for base-pairing with the RNA. Suitable sites can be identified, and crRNA guide or single guide sequences can be prepared, and are presented below in Table 3.

While not limited to any particular theory, when complexed, the guideRNA-Cas9 protein can recognize a DNA site for cleavage, which can then be repaired by cellular components by either non-homologous end joining (random repair) or by a DNA template repair pathway (homology directed repair, HDR).

Guide RNAs (gRNA) can be generated across exons 1 to exons 7 of CD163. Edits in the virus binding domain (Domain 5) of the CD163 protein can inhibit the ability of the virus to bind to the protein, thus preventing uptake of the virus into pig lung macrophages. When the edit in CD163 is homozygous in pigs, the animals can be resistant to PRRS infection. The present specification provides for, and includes, gRNA sequences including the crRNA sequences listed in Table 3 that can be selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. SEQ ID NOs: 22 to 271 and 347 to 425 are shown as DNA sequences. The corresponding RNA counterpart, and the presence of the PAM motif which is generally not included in gRNAs, would be apparent to a person of skill in the art. Also provided herein, are RNA sequences that can comprise sequences selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. SEQ ID NOs: 22 to 271 and 347 to 425 can be combined with the gRNA backbone sequence and expressed as RNA for use in the preparation of a gRNA and then mixed with the Cas nuclease to prepare an active ribonucleotide protein complex (sgRNP). The sgRNP complex can then be injected into a mature oocyte, a zygote, or early embryo, either alone or together with a repair template selected from the group consisting of SEQ ID NOs: 1 to 13. After cleavage and template directed repair, the CD163 gene can be edited to incorporate the repair template sequences, and the edited genome of a CD163 gene edited cell can comprise a sequence selected from the group consisting of SEQ ID NOs: 1 to 13.

In one aspect, the present disclosure can include a simple, precise, and reproducible single CD163 loss of function edit in elite pigs such that the edit occurs early in the CD163 gene (exons 1-7). In an aspect, a pig can have a CD163 gene comprising a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 and 426 to 505. Also included and provided for is a pig that can comprise SEQ ID NOs: 1 to 18 and 425 to 505 in a genomic region comprising the markers of Table 8. Also included and provided for is a pig that can comprise SEQ ID NO: 2 in a genomic region comprising the markers of Table 8. Also included and provided for is a pig that can comprise SEQ ID NO: 426-458 in a genomic region comprising the markers of Table 8. Also included and provided for is a pig that can comprise SEQ ID NO: 459-504 in a genomic region comprising the markers of Table 8. Using porcine embryonic fibroblast cell lines derived from these elite pigs, 250 guide RNA and Cas9 protein combinations can be tested to identify gene editing pairs that efficiently cut and generate edits in the CD163 gene. Selected specific guides and endonuclease protein combi-

nations found are listed in Table 3, along with the location of the target, their endonuclease activity (average editing frequency), and the position on CD163 of the edit. Greater average editing frequency combinations are preferred because they have more effective editing activity. Combinations having an average editing frequency greater than 1 are desired. In an aspect, combinations can have an average editing frequency greater than 5. In another aspect, combinations can have an average editing frequency greater than 10. In yet another aspect, combinations can have an average editing frequency greater than 15. The present disclosure provides for, and includes, combinations that can have an average editing frequency greater than 20. In an aspect, combinations can have an average editing frequency greater than 25. In yet another aspect, combinations can have an average editing frequency greater than 30.

As shown in Table 3, the average editing frequency varied greatly between target sequences. Moreover, sequences targeting overlapping sequences resulted in different editing frequencies depending on the Cas protein source. For example, SEQ ID NOs: 24 and 25 targeted overlapping locations and had a 10-fold difference in editing frequency. At this site, with these sequences, *S. pyogenes* provided a nearly a 20-fold higher editing frequency. In contrast, RNPs incorporating SEQ ID NOs: 33 and 34 resulted in indistinguishable average editing frequencies between the Cas proteins. While RNPs incorporating a Cas9 nuclease from *S. pyogenes* can provide superior average editing frequencies, that is not always the case. As shown for SEQ ID NOs: 60 and 61, RNPs incorporating the Cas9 protein from *S. thermophilus* resulted in superior editing frequencies (e.g., 3.9 vs. 22.0). Accordingly, the present specification provides for efficient guide RNA sequences or efficient targeting sequences for targeting CD163 using the CRISPR/Cas system. In an aspect, RNPs can comprise a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425.

All position designations herein refer to the Sscrofa11.1 reference genome (GenBank accession: GCA_000003025.6).

Target site sequences SEQ ID NOs: 22 to 271 and 347 to 425 can be editing target sequences within the porcine CD163 gene. In each target site sequence, the first 20 nucleotides can correspond to the guide RNA spacer, and the remaining 3 or 5 nucleotides can be the PAM sequences for *S. pyogenes* or *S. thermophilus*, respectively. The first 20 bases of SEQ ID NOs: 22 to 271 and 347 to 425 correspond to the DNA equivalent guide RNA sequences, with the appropriate ribonuclease bases substituted for deoxyribonuclease bases, although the DNA bases can be used instead of RNA bases as guides. Guide RNA sequences can be paired with *S. pyogenes* or *S. thermophilus* endonucleases to direct DNA breaks in the porcine genome in either coding or non-coding regions of the CD163 gene, which can result in DNA deletions and/or DNA nucleotide insertions. Depending on the locations of the target sites as disclosed in Table 3, pairs of guide/endonuclease combinations can be selected to delete regions of the CD163 gene between the pairs of guide/endonuclease combinations.

Target site sequences SEQ ID NOs: 212 to 271 and 347 to 425 are targeting sequences that can be used to design guides for deletion of exon 7 of the CD163 gene. Pairing of intron 6 targeting guides SEQ ID NOs: 212 to 253, and 262 to 271 with intron 7 targeting guides SEQ ID NOs: 254 to 261 using *S. pyogenes* can cause two double strand breaks on the CD163 gene, resulting in excision of exon 7. Pairing of targeting guides SEQ ID NOs: 249 and 256 also can result

in excision of Exon 7. Pairing of some guides can result in the introduction of a stop codon in Exon 7, including SEQ ID NOs: 351 and 365, SEQ ID NOs: 351 and 387, SEQ ID NOs: 348 and 390, SEQ ID NOs: 348 and 388, SEQ ID NOs: 348 and 498, SEQ ID NOs: 352 and 365, SEQ ID NOs: 352 and 387, SEQ ID NOs: 352 and 399, SEQ ID NOs: 353 and 365, SEQ ID NOs: 353 and 387, SEQ ID NOs: 353 and 399, SEQ ID NOs: 354 and 390, SEQ ID NOs: 354 and 388, SEQ ID NOs: 354 and 395, SEQ ID NOs: 358 and 361, SEQ ID NOs: 358 and 362, SEQ ID NOs: 358 and 368, SEQ ID NOs: 358 and 384, SEQ ID NOs: 358 and 394, SEQ ID NOs: 358 and 399, SEQ ID NOs: 359 and 390, SEQ ID NOs: 359 and 388, SEQ ID NOs: 359 and 395, SEQ ID NOs: 360 and 368, SEQ ID NOs: 360 and 384, SEQ ID NOs: 360 and 389, SEQ ID NOs: 360 and 394, SEQ ID NOs: 360 and 397, SEQ ID NOs: 361 and 365, SEQ ID NOs: 361 and 387, SEQ ID NOs: 362 and 390, SEQ ID NOs: 362 and 388, SEQ ID NOs: 362 and 395, SEQ ID NOs: 364 and 365, SEQ ID NOs: 364 and 387, SEQ ID NOs: 364 and 399, SEQ ID NOs: 365 and 368, SEQ ID NOs: 365 and 384, SEQ ID NOs: 365 and 389, SEQ ID NOs: 365 and 394, SEQ ID NOs: 365 and 397, SEQ ID NOs: 366 and 368, SEQ ID NOs: 366 and 384, SEQ ID NOs: 366 and 389, SEQ ID NOs: 366 and 394, or SEQ ID NOs: 366 and 397.

For each targeting sequence, Table 3 lists its SEQ ID No., the species of CAS9 nuclease homing arm used, the location of the targeting sequence on the *Sus scrofa* genome including the PAM sequence, the editing efficiency as measured by the average fraction of edits for a particular guide in fetal fibroblast cell assays, and the exon on CD163 targeted.

TABLE 3

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
22	<i>S. pyogenes</i>	chr5: 63300192-63300214	12.9	Exon 1/15
23	<i>S. pyogenes</i>	chr5: 63300222-63300244	1.4	Exon 1/15
24	<i>S. pyogenes</i>	chr5: 63300236-63300258	5.8	Exon 1/15
25	<i>S. thermophilus</i>	chr5: 63300236-63300260	0.3	Exon 1/15
26	<i>S. pyogenes</i>	chr5: 63300250-63300272	8.6	Exon 1/15
27	<i>S. pyogenes</i>	chr5: 63300251-63300273	0.1	Exon 1/15
28	<i>S. pyogenes</i>	chr5: 63300275-63300297	25.7	Exon 1/15
29	<i>S. pyogenes</i>	chr5: 63300288-63300310	24.9	Exon 1/15
30	<i>S. pyogenes</i>	chr5: 63300293-63300315	12.8	Exon 1/15
31	<i>S. pyogenes</i>	chr5: 63300305-63300327	0.4	Exon 1/15
32	<i>S. pyogenes</i>	chr5: 63300308-63300330	3.4	Exon 1/15
33	<i>S. thermophilus</i>	chr5: 63300308-63300332	3.7	Exon 1/15
34	<i>S. pyogenes</i>	chr5: 63300327-63300349	5.3	Exon 1/15
35	<i>S. pyogenes</i>	chr5: 63300336-63300358	3.5	Exon 1/15
36	<i>S. pyogenes</i>	chr5: 63301950-63301972	14.4	Exon 2/15
37	<i>S. pyogenes</i>	chr5: 63301951-63301973	16.3	Exon 2/15
38	<i>S. pyogenes</i>	chr5: 63301962-63301984	13.6	Exon 2/15
39	<i>S. pyogenes</i>	chr5: 63301981-63302003	3.6	Exon 2/15
40	<i>S. pyogenes</i>	chr5: 63301995-63302017	21.8	Exon 2/15
41	<i>S. pyogenes</i>	chr5: 63301997-63302019	8.2	Exon 2/15
42	<i>S. thermophilus</i>	chr5: 63301997-63302021	10.1	Exon 2/15
43	<i>S. pyogenes</i>	chr5: 63303121-63303143	16.4	Exon 3/15
44	<i>S. thermophilus</i>	chr5: 63303121-63303145	16.1	Exon 3/15
45	<i>S. pyogenes</i>	chr5: 63303129-63303151	31.4	Exon 3/15
46	<i>S. pyogenes</i>	chr5: 63303136-63303158	22.8	Exon 3/15
47	<i>S. pyogenes</i>	chr5: 63303137-63303159	42.9	Exon 3/15
48	<i>S. thermophilus</i>	chr5: 63303137-63303161	10.7	Exon 3/15
49	<i>S. pyogenes</i>	chr5: 63303140-63303162	36.7	Exon 3/15
50	<i>S. thermophilus</i>	chr5: 63303140-63303164	32.6	Exon 3/15
51	<i>S. pyogenes</i>	chr5: 63303158-63303180	21.1	Exon 3/15
52	<i>S. pyogenes</i>	chr5: 63303166-63303188	23.6	Exon 3/15
53	<i>S. thermophilus</i>	chr5: 63303166-63303190	22.1	Exon 3/15
54	<i>S. pyogenes</i>	chr5: 63303169-63303191	40.0	Exon 3/15
55	<i>S. thermophilus</i>	chr5: 63303169-63303193	19.5	Exon 3/15
56	<i>S. pyogenes</i>	chr5: 63303181-63303203	9.9	Exon 3/15

TABLE 3-continued

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
57	<i>S. thermophilus</i>	chr5: 63303181-63303205	10.4	Exon 3/15
58	<i>S. pyogenes</i>	chr5: 63303184-63303206	20.6	Exon 3/15
59	<i>S. thermophilus</i>	chr5: 63303184-63303208	2.8	Exon 3/15
60	<i>S. pyogenes</i>	chr5: 63303189-63303211	3.9	Exon 3/15
61	<i>S. thermophilus</i>	chr5: 63303189-63303213	22.0	Exon 3/15
62	<i>S. pyogenes</i>	chr5: 63303190-63303212	18.1	Exon 3/15
63	<i>S. pyogenes</i>	chr5: 63303191-63303213	23.5	Exon 3/15
64	<i>S. pyogenes</i>	chr5: 63303209-63303231	23.6	Exon 3/15
65	<i>S. pyogenes</i>	chr5: 63303213-63303235	4.1	Exon 3/15
66	<i>S. pyogenes</i>	chr5: 63303214-63303236	16.2	Exon 3/15
67	<i>S. pyogenes</i>	chr5: 63303220-63303242	14.6	Exon 3/15
68	<i>S. pyogenes</i>	chr5: 63303226-63303248	15.4	Exon 3/15
69	<i>S. pyogenes</i>	chr5: 63303243-63303265	21.6	Exon 3/15
70	<i>S. pyogenes</i>	chr5: 63303250-63303272	1.5	Exon 3/15
71	<i>S. pyogenes</i>	chr5: 63303251-63303273	30.3	Exon 3/15
72	<i>S. pyogenes</i>	chr5: 63303278-63303300	20.1	Exon 3/15
73	<i>S. pyogenes</i>	chr5: 63303278-63303300	1.5	Exon 3/15
74	<i>S. pyogenes</i>	chr5: 63303282-63303304	23.9	Exon 3/15
75	<i>S. pyogenes</i>	chr5: 63303283-63303305	20.8	Exon 3/15
76	<i>S. pyogenes</i>	chr5: 63303294-63303316	33.8	Exon 3/15
77	<i>S. pyogenes</i>	chr5: 63303299-63303321	8.2	Exon 3/15
78	<i>S. pyogenes</i>	chr5: 63303305-63303327	6.1	Exon 3/15
79	<i>S. pyogenes</i>	chr5: 63303315-63303337	25.2	Exon 3/15
80	<i>S. pyogenes</i>	chr5: 63303319-63303341	28.7	Exon 3/15
81	<i>S. pyogenes</i>	chr5: 63303338-63303360	4.5	Exon 3/15
82	<i>S. pyogenes</i>	chr5: 63303339-63303361	2.8	Exon 3/15
83	<i>S. pyogenes</i>	chr5: 63303357-63303379	18.3	Exon 3/15
84	<i>S. pyogenes</i>	chr5: 63303358-63303380	26.6	Exon 3/15
85	<i>S. pyogenes</i>	chr5: 63303374-63303396	16.1	Exon 3/15
86	<i>S. pyogenes</i>	chr5: 63303378-63303400	35.4	Exon 3/15
87	<i>S. thermophilus</i>	chr5: 63303378-63303402	40.0	Exon 3/15
88	<i>S. pyogenes</i>	chr5: 63303379-63303401	24.4	Exon 3/15
89	<i>S. pyogenes</i>	chr5: 63303380-63303402	25.6	Exon 3/15
90	<i>S. pyogenes</i>	chr5: 63303406-63303428	28.1	Exon 3/15
91	<i>S. pyogenes</i>	chr5: 63303413-63303435	13.7	Exon 3/15
92	<i>S. thermophilus</i>	chr5: 63303413-63303437	19.0	Exon 3/15
93	<i>S. thermophilus</i>	chr5: 63303419-63303443	15.2	Exon 3/15
94	<i>S. pyogenes</i>	chr5: 63303421-63303443	16.5	Exon 3/15
95	<i>S. pyogenes</i>	chr5: 63303428-63303450	35.8	Exon 3/15
96	<i>S. pyogenes</i>	chr5: 63303441-63303463	3.4	Exon 3/15
97	<i>S. pyogenes</i>	chr5: 63306087-63306109	19.8	Exon 4/15
98	<i>S. pyogenes</i>	chr5: 63306091-63306113	4.5	Exon 4/15
99	<i>S. thermophilus</i>	chr5: 63306091-63306115	3.1	Exon 4/15
100	<i>S. pyogenes</i>	chr5: 63306098-63306120	29.9	Exon 4/15
101	<i>S. thermophilus</i>	chr5: 63306098-63306122	31.5	Exon 4/15
102	<i>S. pyogenes</i>	chr5: 63306101-63306123	29.2	Exon 4/15
103	<i>S. pyogenes</i>	chr5: 63306108-63306130	6.9	Exon 4/15
104	<i>S. thermophilus</i>	chr5: 63306108-63306132	14.7	Exon 4/15
105	<i>S. pyogenes</i>	chr5: 63306116-63306138	21.6	Exon 4/15
106	<i>S. pyogenes</i>	chr5: 63306127-63306149	26.4	Exon 4/15
107	<i>S. pyogenes</i>	chr5: 63306140-63306162	6.9	Exon 4/15
108	<i>S. pyogenes</i>	chr5: 63306144-63306166	35.2	Exon 4/15
109	<i>S. thermophilus</i>	chr5: 63306144-63306168	3.3	Exon 4/15
110	<i>S. pyogenes</i>	chr5: 63306147-63306169	8.8	Exon 4/15
111	<i>S. thermophilus</i>	chr5: 63306147-63306171	3.2	Exon 4/15
112	<i>S. pyogenes</i>	chr5: 63306148-63306170	5.2	Exon 4/15
113	<i>S. pyogenes</i>	chr5: 63306149-63306171	28.0	Exon 4/15
114	<i>S. pyogenes</i>	chr5: 63306193-63306215	9.3	Exon 4/15
115	<i>S. pyogenes</i>	chr5: 63306236-63306258	14.2	Exon 4/15
116	<i>S. pyogenes</i>	chr5: 63306251-63306273	26.3	Exon 4/15
117	<i>S. thermophilus</i>	chr5: 63306251-63306275	4.4	Exon 4/15
118	<i>S. pyogenes</i>	chr5: 63306257-63306279	1.3	Exon 4/15
119	<i>S. pyogenes</i>	chr5: 63306263-63306285	7.8	Exon 4/15
120	<i>S. pyogenes</i>	chr5: 63306273-63306295	35.0	Exon 4/15
121	<i>S. pyogenes</i>	chr5: 63306287-63306309	13.4	Exon 4/15
122	<i>S. pyogenes</i>	chr5: 63306296-63306318	2.0	Exon 4/15
123	<i>S. pyogenes</i>	chr5: 63306315-63306337	31.2	Exon 4/15
124	<i>S. pyogenes</i>	chr5: 63306332-63306354	10.1	Exon 4/15
125	<i>S. pyogenes</i>	chr5: 63306336-63306358	31.1	Exon 4/15
126	<i>S. thermophilus</i>	chr5: 63306336-63306360	52.8	Exon 4/15
127	<i>S. pyogenes</i>	chr5: 63306337-63306359	30.3	Exon 4/15
128	<i>S. pyogenes</i>	chr5: 63306338-63306360	43.2	Exon 4/15
129	<i>S. pyogenes</i>	chr5: 63306364-63306386	2.5	Exon 4/15
130	<i>S. pyogenes</i>	chr5: 63306371-63306393	0.2	Exon 4/15

33

TABLE 3-continued

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
131	<i>S. thermophilus</i>	chr5: 63306371-63306395	0.3	Exon 4/15
132	<i>S. pyogenes</i>	chr5: 63309028-63309050	4.5	Exon 5/15
133	<i>S. pyogenes</i>	chr5: 63309034-63309056	5.1	Exon 5/15
134	<i>S. pyogenes</i>	chr5: 63309035-63309057	41.8	Exon 5/15
135	<i>S. thermophilus</i>	chr5: 63309035-63309059	21.9	Exon 5/15
136	<i>S. pyogenes</i>	chr5: 63309053-63309075	15.1	Exon 5/15
137	<i>S. pyogenes</i>	chr5: 63309061-63309083	0.2	Exon 5/15
138	<i>S. pyogenes</i>	chr5: 63309077-63309099	15.7	Exon 5/15
139	<i>S. thermophilus</i>	chr5: 63309077-63309101	11.2	Exon 5/15
140	<i>S. pyogenes</i>	chr5: 63309084-63309106	10.1	Exon 5/15
141	<i>S. thermophilus</i>	chr5: 63309084-63309108	23.8	Exon 5/15
142	<i>S. pyogenes</i>	chr5: 63309085-63309107	11.3	Exon 5/15
143	<i>S. pyogenes</i>	chr5: 63309086-63309108	23.4	Exon 5/15
144	<i>S. pyogenes</i>	chr5: 63309094-63309116	13.8	Exon 5/15
145	<i>S. pyogenes</i>	chr5: 63309104-63309126	7.8	Exon 5/15
146	<i>S. pyogenes</i>	chr5: 63309108-63309130	1.4	Exon 5/15
147	<i>S. pyogenes</i>	chr5: 63309109-63309131	7.0	Exon 5/15
148	<i>S. pyogenes</i>	chr5: 63309130-63309152	24.8	Exon 5/15
149	<i>S. pyogenes</i>	chr5: 63309144-63309166	20.8	Exon 5/15
150	<i>S. pyogenes</i>	chr5: 63309145-63309167	0.1	Exon 5/15
151	<i>S. pyogenes</i>	chr5: 63309146-63309168	37.4	Exon 5/15
152	<i>S. pyogenes</i>	chr5: 63309173-63309195	2.5	Exon 5/15
153	<i>S. pyogenes</i>	chr5: 63309173-63309195	2.3	Exon 5/15
154	<i>S. pyogenes</i>	chr5: 63309189-63309211	11.5	Exon 5/15
155	<i>S. pyogenes</i>	chr5: 63309193-63309215	2.7	Exon 5/15
156	<i>S. pyogenes</i>	chr5: 63309194-63309216	10.4	Exon 5/15
157	<i>S. pyogenes</i>	chr5: 63309200-63309222	3.8	Exon 5/15
158	<i>S. thermophilus</i>	chr5: 63309205-63309229	19.0	Exon 5/15
159	<i>S. pyogenes</i>	chr5: 63309207-63309229	21.0	Exon 5/15
160	<i>S. pyogenes</i>	chr5: 63309210-63309232	19.0	Exon 5/15
161	<i>S. pyogenes</i>	chr5: 63309233-63309255	8.3	Exon 5/15
162	<i>S. pyogenes</i>	chr5: 63309250-63309272	13.6	Exon 5/15
163	<i>S. pyogenes</i>	chr5: 63309252-63309274	23.0	Exon 5/15
164	<i>S. pyogenes</i>	chr5: 63309273-63309295	2.3	Exon 5/15
165	<i>S. thermophilus</i>	chr5: 63309273-63309297	34.4	Exon 5/15
166	<i>S. pyogenes</i>	chr5: 63309274-63309296	15.1	Exon 5/15
167	<i>S. pyogenes</i>	chr5: 63309275-63309297	19.4	Exon 5/15
168	<i>S. thermophilus</i>	chr5: 63309284-63309308	14.4	Exon 5/15
169	<i>S. pyogenes</i>	chr5: 63309286-63309308	7.0	Exon 5/15
170	<i>S. pyogenes</i>	chr5: 63309308-63309330	0.5	Exon 5/15
171	<i>S. thermophilus</i>	chr5: 63309308-63309332	2.4	Exon 5/15
172	<i>S. pyogenes</i>	chr5: 63309323-63309345	16.0	Exon 5/15
173	<i>S. pyogenes</i>	chr5: 63309841-63309863	0.1	Exon 6/15
174	<i>S. pyogenes</i>	chr5: 63309857-63309879	8.4	Exon 6/15
175	<i>S. thermophilus</i>	chr5: 63309857-63309881	8.0	Exon 6/15
176	<i>S. pyogenes</i>	chr5: 63309860-63309882	40.3	Exon 6/15
177	<i>S. thermophilus</i>	chr5: 63309860-63309884	32.3	Exon 6/15
178	<i>S. pyogenes</i>	chr5: 63309863-63309885	37.2	Exon 6/15
179	<i>S. pyogenes</i>	chr5: 63309886-63309908	4.7	Exon 6/15
180	<i>S. thermophilus</i>	chr5: 63309886-63309910	3.5	Exon 6/15
181	<i>S. pyogenes</i>	chr5: 63309889-63309911	38.1	Exon 6/15
182	<i>S. pyogenes</i>	chr5: 63309889-63309911	34.1	Exon 6/15
183	<i>S. thermophilus</i>	chr5: 63309889-63309913	17.5	Exon 6/15
184	<i>S. pyogenes</i>	chr5: 63309892-63309914	4.9	Exon 6/15
185	<i>S. pyogenes</i>	chr5: 63309907-63309929	13.5	Exon 6/15
186	<i>S. pyogenes</i>	chr5: 63309911-63309933	5.5	Exon 6/15
187	<i>S. pyogenes</i>	chr5: 63309933-63309955	0.7	Exon 6/15
188	<i>S. thermophilus</i>	chr5: 63309933-63309957	10.0	Exon 6/15
189	<i>S. pyogenes</i>	chr5: 63309934-63309956	6.4	Exon 6/15
190	<i>S. pyogenes</i>	chr5: 63309935-63309957	18.7	Exon 6/15
191	<i>S. pyogenes</i>	chr5: 63309955-63309977	12.0	Exon 6/15
192	<i>S. pyogenes</i>	chr5: 63309963-63309985	5.6	Exon 6/15
193	<i>S. pyogenes</i>	chr5: 63309970-63309992	0.5	Exon 6/15
194	<i>S. pyogenes</i>	chr5: 63309971-63309993	12.5	Exon 6/15
195	<i>S. pyogenes</i>	chr5: 63309977-63309999	5.3	Exon 6/15
196	<i>S. pyogenes</i>	chr5: 63310021-63310043	2.2	Exon 6/15
197	<i>S. pyogenes</i>	chr5: 63310035-63310057	10.3	Exon 6/15
198	<i>S. pyogenes</i>	chr5: 63310038-63310060	0.2	Exon 6/15
199	<i>S. pyogenes</i>	chr5: 63310058-63310080	8.2	Exon 6/15
200	<i>S. pyogenes</i>	chr5: 63310077-63310099	6.2	Exon 6/15
201	<i>S. pyogenes</i>	chr5: 63310078-63310100	8.6	Exon 6/15
202	<i>S. pyogenes</i>	chr5: 63310092-63310114	2.7	Exon 6/15
203	<i>S. pyogenes</i>	chr5: 63310098-63310120	11.1	Exon 6/15
204	<i>S. thermophilus</i>	chr5: 63310098-63310122	8.0	Exon 6/15

34

TABLE 3-continued

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
205	<i>S. pyogenes</i>	chr5: 63310099-63310121	12.2	Exon 6/15
206	<i>S. pyogenes</i>	chr5: 63310100-63310122	21.1	Exon 6/15
207	<i>S. thermophilus</i>	chr5: 63310100-63310124	0.1	Exon 6/15
208	<i>S. pyogenes</i>	chr5: 63310103-63310125	3.3	Exon 6/15
209	<i>S. pyogenes</i>	chr5: 63310152-63310174	4.8	Exon 6/15
210	<i>S. pyogenes</i>	chr5: 63323061-63323083	32.8	Exon 7/15
211	<i>S. pyogenes</i>	chr5: 63323147-63323169	17.0	Exon 7/15
212	<i>S. pyogenes</i>	chr5: 63322548-63322570	13.6	Intron 6
213	<i>S. pyogenes</i>	chr5: 63322549-63322571	1.7	Intron 6
214	<i>S. pyogenes</i>	chr5: 63322566-63322588	15.6	Intron 6
215	<i>S. pyogenes</i>	chr5: 63322594-63322616	0.0	Intron 6
216	<i>S. pyogenes</i>	chr5: 63322597-63322619	0.0	Intron 6
217	<i>S. pyogenes</i>	chr5: 63322646-63322668	21.1	Intron 6
218	<i>S. pyogenes</i>	chr5: 63322647-63322669	12.6	Intron 6
219	<i>S. pyogenes</i>	chr5: 63322681-63322703	44.3	Intron 6
220	<i>S. pyogenes</i>	chr5: 63322683-63322705	3.9	Intron 6
221	<i>S. pyogenes</i>	chr5: 63322693-63322715	33.9	Intron 6
222	<i>S. pyogenes</i>	chr5: 63322694-63322716	26.1	Intron 6
223	<i>S. pyogenes</i>	chr5: 63322714-63322736	5.7	Intron 6
224	<i>S. pyogenes</i>	chr5: 63322731-63322753	42.8	Intron 6
225	<i>S. pyogenes</i>	chr5: 63322756-63322778	5.8	Intron 6
226	<i>S. pyogenes</i>	chr5: 63322757-63322779	21.2	Intron 6
227	<i>S. pyogenes</i>	chr5: 63322770-63322792	36.1	Intron 6
228	<i>S. pyogenes</i>	chr5: 63322799-63322821	29.9	Intron 6
229	<i>S. pyogenes</i>	chr5: 63322800-63322822	43.2	Intron 6
230	<i>S. pyogenes</i>	chr5: 63322809-63322831	33.3	Intron 6
231	<i>S. pyogenes</i>	chr5: 63322810-63322832	46.8	Intron 6
232	<i>S. pyogenes</i>	chr5: 63322834-63322856	7.8	Intron 6
233	<i>S. pyogenes</i>	chr5: 63322835-63322857	18.3	Intron 6
234	<i>S. pyogenes</i>	chr5: 63322839-63322861	18.8	Intron 6
235	<i>S. pyogenes</i>	chr5: 63322839-63322861	13.2	Intron 6
236	<i>S. pyogenes</i>	chr5: 63322840-63322862	2.8	Intron 6
237	<i>S. pyogenes</i>	chr5: 63322845-63322867	55.3	Intron 6
238	<i>S. pyogenes</i>	chr5: 63322848-63322870	27.2	Intron 6
239	<i>S. pyogenes</i>	chr5: 63322852-63322874	24.4	Intron 6
240	<i>S. pyogenes</i>	chr5: 63322859-63322881	22.2	Intron 6
241	<i>S. pyogenes</i>	chr5: 63322875-63322897	25.3	Intron 6
242	<i>S. pyogenes</i>	chr5: 63322887-63322909	4.2	Intron 6
243	<i>S. pyogenes</i>	chr5: 63322888-63322910	3.5	Intron 6
244	<i>S. pyogenes</i>	chr5: 63322891-63322913	44.3	Intron 6
245	<i>S. pyogenes</i>	chr5: 63322900-63322922	58.2	Intron 6
246	<i>S. pyogenes</i>	chr5: 63322906-63322928	3.7	Intron 6
247	<i>S. pyogenes</i>	chr5: 63322926-63322948	38.1	Intron 6
248	<i>S. pyogenes</i>	chr5: 63322927-63322949	0.5	Intron 6
249	<i>S. pyogenes</i>	chr5: 63322947-63322969	31.7	Intron 6
250	<i>S. pyogenes</i>	chr5: 63322957-63322979	50.0	Intron 6
251	<i>S. pyogenes</i>	chr5: 63322957-63322979	11.0	Intron 6
252	<i>S. pyogenes</i>	chr5: 63322991-63323013	4.5	Intron 6
253	<i>S. pyogenes</i>	chr5: 63322992-63323014	0.5	Intron 6
254	<i>S. pyogenes</i>	chr5: 63323338-63323360	19.8	Intron 7
255	<i>S. pyogenes</i>	chr5: 63323339-63323361	8.1	Intron 7
256	<i>S. pyogenes</i>	chr5: 63323361-63323383	21.1	Intron 7
257	<i>S. pyogenes</i>	chr5: 63323362-63323384	30.4	Intron 7
258	<i>S. pyogenes</i>	chr5: 63323362-63323384	5.6	Intron 7
259	<i>S. pyogenes</i>	chr5: 63323363-63323385	2.8	Intron 7
260	<i>S. pyogenes</i>	chr5: 63323367-63323389	23.2	Intron 7
261	<i>S. pyogenes</i>	chr5: 63323368-63323390	26.2	Intron 7
262	<i>S. thermophilus</i>	chr5: 63322644-63322668	Activity not tested	Intron 6
263	<i>S. thermophilus</i>	chr5: 63322647-63322671	Activity not tested	Intron 6
264	<i>S. thermophilus</i>	chr5: 63322678-63322702	Activity not tested	Intron 6
265	<i>S. thermophilus</i>	chr5: 63322681-63322705	Activity not tested	Intron 6
266	<i>S. thermophilus</i>	chr5: 63322755-63322779	Activity not tested	Intron 6
267	<i>S. thermophilus</i>	chr5: 63322807-63322831	Activity not tested	Intron 6
268	<i>S. thermophilus</i>	chr5: 63322845-63322869	Activity not tested	Intron 6
269	<i>S. thermophilus</i>	chr5: 63322850-63322874	Activity not tested	Intron 6

35

TABLE 3-continued

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
270	<i>S. thermophilus</i>	chr5: 63322955-63322979	Activity not tested	Intron 6
271	<i>S. thermophilus</i>	chr5: 63322989-63323013	Activity not tested	Intron 6
347	<i>S. pyogenes</i>	chr5: 63323002-63323024	2.1	Exon 7
348	<i>S. pyogenes</i>	chr5: 63323011-63323033	31.9	Exon 7
349	<i>S. pyogenes</i>	chr5: 63323015-63323037	7.7	Exon 7
350	<i>S. pyogenes</i>	chr5: 63323017-63323039	56.5	Exon 7
351	<i>S. pyogenes</i>	chr5: 63323018-63323040	74.3	Exon 7
352	<i>S. pyogenes</i>	chr5: 63323019-63323041	52.3	Exon 7
353	<i>S. pyogenes</i>	chr5: 63323022-63323044	44.0	Exon 7
354	<i>S. pyogenes</i>	chr5: 63323023-63323045	61.6	Exon 7
355	<i>S. pyogenes</i>	chr5: 63323024-63323046	49.4	Exon 7
356	<i>S. pyogenes</i>	chr5: 63323028-63323050	3.1	Exon 7
357	<i>S. pyogenes</i>	chr5: 63323029-63323051	0.8	Exon 7
358	<i>S. pyogenes</i>	chr5: 63323040-63323062	42.4	Exon 7
359	<i>S. pyogenes</i>	chr5: 63323052-63323074	66.7	Exon 7
360	<i>S. pyogenes</i>	chr5: 63323053-63323075	13.4	Exon 7
361	<i>S. pyogenes</i>	chr5: 63323061-63323083	77.9	Exon 7
362	<i>S. pyogenes</i>	chr5: 63323071-63323093	60.8	Exon 7
363	<i>S. pyogenes</i>	chr5: 63323072-63323094	70.6	Exon 7
364	<i>S. pyogenes</i>	chr5: 63323073-63323095	75.2	Exon 7
365	<i>S. pyogenes</i>	chr5: 63323098-63323120	64.4	Exon 7
366	<i>S. pyogenes</i>	chr5: 63323102-63323124	55.0	Exon 7
367	<i>S. pyogenes</i>	chr5: 63323105-63323127	51.2	Exon 7
368	<i>S. pyogenes</i>	chr5: 63323108-63323130	58.3	Exon 7
369	<i>S. pyogenes</i>	chr5: 63323125-63323147	14.5	Exon 7
370	<i>S. pyogenes</i>	chr5: 63323126-63323148	22.6	Exon 7
371	<i>S. pyogenes</i>	chr5: 63323131-63323153	49.9	Exon 7
372	<i>S. pyogenes</i>	chr5: 63323139-63323161	68.9	Exon 7
373	<i>S. pyogenes</i>	chr5: 63323147-63323169	57.4	Exon 7
374	<i>S. pyogenes</i>	chr5: 63323159-63323181	12.9	Exon 7
375	<i>S. pyogenes</i>	chr5: 63323160-63323182	19.7	Exon 7
376	<i>S. pyogenes</i>	chr5: 63323161-63323183	33.6	Exon 7
377	<i>S. pyogenes</i>	chr5: 63323162-63323184	65.3	Exon 7
378	<i>S. pyogenes</i>	chr5: 63323163-63323185	51.4	Exon 7
379	<i>S. pyogenes</i>	chr5: 63323173-63323195	57.6	Exon 7
380	<i>S. pyogenes</i>	chr5: 63323174-63323196	52.1	Exon 7
381	<i>S. pyogenes</i>	chr5: 63323175-63323197	48.2	Exon 7
382	<i>S. pyogenes</i>	chr5: 63323177-63323199	33.7	Exon 7
383	<i>S. pyogenes</i>	chr5: 63323181-63323203	18.3	Exon 7
384	<i>S. pyogenes</i>	chr5: 63323187-63323209	57.0	Exon 7
385	<i>S. pyogenes</i>	chr5: 63323197-63323219	33.3	Exon 7
386	<i>S. pyogenes</i>	chr5: 63323198-63323220	58.3	Exon 7
387	<i>S. pyogenes</i>	chr5: 63323219-63323241	20.7	Exon 7
388	<i>S. pyogenes</i>	chr5: 63323220-63323242	42.2	Exon 7
389	<i>S. pyogenes</i>	chr5: 63323221-63323243	43.4	Exon 7
390	<i>S. pyogenes</i>	chr5: 63323231-63323253	21.8	Exon 7
391	<i>S. pyogenes</i>	chr5: 63323251-63323273	46.8	Exon 7
392	<i>S. pyogenes</i>	chr5: 63323252-63323274	42.6	Exon 7
393	<i>S. pyogenes</i>	chr5: 63323255-63323277	22.8	Exon 7
394	<i>S. pyogenes</i>	chr5: 63323267-63323289	85.4	Exon 7
395	<i>S. pyogenes</i>	chr5: 63323268-63323290	41.1	Exon 7
396	<i>S. pyogenes</i>	chr5: 63323268-63323290	53.6	Exon 7
397	<i>S. pyogenes</i>	chr5: 63323269-63323291	32.1	Exon 7
398	<i>S. pyogenes</i>	chr5: 63323277-63323299	16.6	Exon 7
399	<i>S. pyogenes</i>	chr5: 63323278-63323300	39.0	Exon 7
400	<i>S. pyogenes</i>	chr5: 63323279-63323301	11.7	Exon 7
401	<i>S. pyogenes</i>	chr5: 63323282-63323304	48.9	Exon 7
402	<i>S. pyogenes</i>	chr5: 63323283-63323305	n/a	Exon 7
403	<i>S. pyogenes</i>	chr5: 63323287-63323309	11.6	Exon 7
404	<i>S. pyogenes</i>	chr5: 63323288-63323310	69.3	Exon 7
405	<i>S. pyogenes</i>	chr5: 63323295-63323317	46.0	Exon 7
406	<i>S. pyogenes</i>	chr5: 63323300-63323322	54.2	Exon 7
407	<i>S. thermophilus</i>	chr5: 63323019-63323043	25.5	Exon 7
408	<i>S. thermophilus</i>	chr5: 63323022-63323046	24.7	Exon 7
409	<i>S. thermophilus</i>	chr5: 63323061-63323085	37.8	Exon 7
410	<i>S. thermophilus</i>	chr5: 63323071-63323095	30.4	Exon 7
411	<i>S. thermophilus</i>	chr5: 63323096-63323120	33.4	Exon 7
412	<i>S. thermophilus</i>	chr5: 63323102-63323126	19.8	Exon 7
413	<i>S. thermophilus</i>	chr5: 63323105-63323129	0.6	Exon 7
414	<i>S. thermophilus</i>	chr5: 63323159-63323183	14.8	Exon 7
415	<i>S. thermophilus</i>	chr5: 63323160-63323184	7.8	Exon 7
416	<i>S. thermophilus</i>	chr5: 63323161-63323185	3.6	Exon 7

36

TABLE 3-continued

List of target sequences and editing activities in porcine fetal fibroblasts				
SEQ ID NO:	Cas9 Nuclease	Location of target	Average edited fraction	Position on CD163
417	<i>S. thermophilus</i>	chr5: 63323163-63323187	7.9	Exon 7
418	<i>S. thermophilus</i>	chr5: 63323175-63323199	33.8	Exon 7
419	<i>S. thermophilus</i>	chr5: 63323175-63323199	11.5	Exon 7
420	<i>S. thermophilus</i>	chr5: 63323219-63323243	8.1	Exon 7
421	<i>S. thermophilus</i>	chr5: 63323253-63323277	32.6	Exon 7
422	<i>S. thermophilus</i>	chr5: 63323275-63323299	0.2	Exon 7
423	<i>S. thermophilus</i>	chr5: 63323277-63323301	12.3	Exon 7
424	<i>S. thermophilus</i>	chr5: 63323280-63323304	26.0	Exon 7
425	<i>S. thermophilus</i>	chr5: 63323295-63323319	30.5	Exon 7

Efficient guideRNA-Cas9 pairs which cut sites early in the CD163 gene (exons 1-4) were selected and further screened for potential off-target binding within the pig genome.

In some instances, the use of guide RNA and endonuclease has been observed to result in cleavage of DNA at unintended locations in the genome. Because the repair process for dsDNA breaks can be random, off-target cleavage events can result in undesirable changes to coding or regulatory regions in the genome. Therefore, in instances where a number of single or paired guides can perform an intended edit with similar editing frequencies, it is advantageous to consider off-target cleavage in choosing a guide or guide pair for editing experiments.

A number of computational and biochemical approaches for elucidating off-targets have been developed. Computational approaches can include, but are not limited to, Cas-OFFinder (Bae, S., et al., Bioinformatics, 2014, 30, 1473-1475.), CRISPR-offender (Zhao, C., et al., Int. J. Biol. Sci., 2017, 13, 1470-1478), and CRISPR-OFF (Alkan, F., et al., Genome Biol., 2018, 19, 177). Other computational approaches are readily available to those skilled in the art. Biochemical approaches can include, but are not limited to, GUIDE-Seq (Tsai, S. Q., et al., Nat Biotechnol. 2015, 33, 187-197), SITE-SEQ® (Cameron, P., et al., Nat Methods 2017, 14, 600-606), and CIRCLE-seq (Tsai, S. Q., et al., Nat. Methods, 2017, 14, 607-614). Other biochemical approaches are readily available to those skilled in the art. While computational methods are relatively fast and inexpensive compared to biochemical ones, biochemical approaches have been shown to be superior in identifying validated off-target edits.

A subset of guideRNAs that have demonstrated a high frequency of intended edits in porcine fetal fibroblasts, as described above, were assayed for specificity using SITE-SEQ®. Using naked gDNA and RNP editing reagents in vitro, SITE-SEQ® can provide a list of potential cleavage sites. Off-target cleavage in a cellular environment can be more complicated than that simulated biochemically in vitro. Factors such as effective RNP concentration and target availability due to chromatin state, among other factors, can contribute to the number of off-target edits realized in porcine cells or in edited pigs. Fortunately, biochemical methods such as SITE-SEQ® can provide researchers a list of sites to interrogate. These sites can be interrogated by methods that include, but are not limited to, TOPO cloning and sequencing of TOPO clones, ILLUMINA® amplicon sequencing, Nanopore sequencing, other NGS sequencing methods, and sequence capture (Gnirke, A., et al., Nat. Biotechnol., 2009, 27, 182-189).

Screening was performed for biochemically-identified off-target sites in edited porcine fibroblasts and subsequently in injected embryos. Guides with validated off-target edits were de-prioritized for use in generating edited pigs. Animals generated using guides with known off-targets can be interrogated for the presence of off-target edits using the strategies outlined above. Animals that do contain off-target edits can either be removed from the breeding program or the off-target edits can be removed via breeding. This screening can include bioinformatic methods, such as BLAST® searching, to identify sequences in the *Sus scrofa* genome that contain 1-5 mismatches with the guideRNA, that could therefore allow for off-target binding. The number of potential off-target binding sites in the genome when allowing these mismatches in the computer algorithm is detailed below in Table 4.

TABLE 4

Mismatch Detection					
SITE	LOCATION-Spy coordinates	SEQ ID		MISMATCHES ≤ 5	
		SEQ ID Spy	SEQ ID Stherm	Spy	Stherm C3
1	chr5: 63300236-63300258	19	20	920	142
2	chr5: 63300308-63300330	32	33	625	51
3	chr5: 63301997-63302019	41	42	506	37
4	chr5: 63303121-63303143	43	44	1625	201
5	chr5: 63303137-63303159	47	48	474	32
6	chr5: 63303140-63303162	49	50	548	49
7	chr5: 63303166-63303188	52	53	692	92
8	chr5: 63303169-63303191	54	55	782	94
9	chr5: 63303181-63303203	56	57	1346	200
10	chr5: 63303184-63303206	58	59	1873	382
11	chr5: 63303189-63303211	60	61	1642	240
12	chr5: 63303378-63303400	86	87	588	53
13	chr5: 63303413-63303435	91	92	545	50
14	chr5: 63306108-63306130	103	104	480	49
15	chr5: 63306144-63306166	108	109	344	21
16	chr5: 63306147-63306169	110	111	451	44
17	chr5: 63306251-63306273	116	117	1343	65
18	chr5: 63306336-63306358	125	126	718	66
19	chr5: 63306371-63306393	130	131	1310	131
20	chr5: 63309035-63309057	134	135	690	83
21	chr5: 63309077-63309099	138	139	949	95
22	chr5: 63309084-63309106	140	141	1234	99
23	chr5: 63309273-63309295	164	165	418	47
24	chr5: 63309308-63309330	170	171	747	95
25	chr5: 63309857-63309879	174	175	795	47
26	chr5: 63309860-63309882	176	177	736	94
27	chr5: 63309886-63309908	179	180	1829	124
28	chr5: 63309889-63309911	182	183	1575	94
29	chr5: 63309933-63309955	187	188	871	95
30	chr5: 63310098-63310120	203	204	533	73
31	chr5: 63310100-63310122	206	207	537	53

One of the factors that contributes to cutting at these off-target sites can be whether the gene editing components are delivered as a DNA vector or as an RNA-protein complex; sites that may have as few as 1 or 2 base mismatches can be faithfully discriminated from the intended target site when the Cas9 and guide RNA can be delivered as a single guide ribonucleotide protein complex (sgRNP) rather than delivered on a DNA vector.

A second consideration to maximize specificity of a Cas9 guide-RNA reagent can be to prescreen guide RNA-protein pairs using in vitro biochemical methods. Several laboratories have published methods to screen for off-target edit sites. These biochemical approaches can identify potential off-target Cas9 cleavage sites in purified genomic DNA. Using these assays, genomic DNA can be digested with a range of sgRNP concentrations, from limiting to saturating,

thus permitting the recovery of both high- and low-cleavage-sensitivity off-target sites. See, e.g., Cameron et al., "SITE-SEQ®: A Genome-wide Method to Measure Cas9 Cleavage," Protocol Exchange (2017).

The off-target sites identified by off-target screening can be used to guide careful and comprehensive examination of possible off-target sites in cells, measuring both editing frequency and functional cellular consequence. Several selected guide RNA-Cas protein pairs can be screened, and the screening can demonstrate: 1) efficient cutting nearer to the 5' end of the CD163 gene, and 2) few mismatch sequences as determined by bioinformatic methods.

Using this guide selection criteria, several guide RNA-Cas protein pairs can be selected for cutting activity in porcine parthenotes. Porcine oocytes can be doubled using an electric current, injected with guide RNA-Cas protein, allowed to develop for 7 days, then harvested and DNA sequenced across the intended target site. Several guide RNA-Cas protein pairs can be identified to confer edits at a high frequency in parthenotes and are selected for further development.

The present specification provides for, and includes, methods for identifying and selecting optimal target sites for CRISPR/Cas mediated cleavage and gene editing. In an aspect, the method can comprise identifying a target region of a genome for editing; identifying all 20 nucleotide sequences in the target region; performing a bioinformatic screen to identify and remove sequences that match non-target sites in the pig genome that contain 1 to 4 mismatches and have a suitable PAM sequence based on the Cas protein; preparing CRISPR/Cas RNP complexes comprising a guide RNA backbone and target sequences, introducing the CRISPR/Cas RNP complexes into a porcine cell in culture, and determining the average editing frequency of the guide RNA/Cas combination. In an aspect, determining the average editing frequency of the guide RNA/Cas combination can comprise amplifying, by PCR, a region surrounding the target site, performing amplicon deep sequencing, and comparing it to untreated cells. In an aspect, the method provides for selecting preferred target sequences that can have an average editing frequency of at least 15. Also provided for are methods for identifying and selecting optimal target sites for CRISPR/Cas mediated cleavage and gene editing wherein the guide RNA and Cas proteins can be provided as part of an expression vector or vectors. Suitable cells are known to persons of skill in the art and can include, but are not limited to, primary fetal fibroblasts of an intended porcine breeding line.

Multiple repair templates can be designed to generate in-frame stop codons. The repair templates listed, when used with appropriate guide RNAs in an endonuclease system, can generate an in-frame termination codon (TAA, TGA, TAG) after repair. Different lengths of repair templates can be used, having sequence identity on either side of the edit site. In an aspect, a repair template can share at least 15 nucleotides on either side of the CRISPR/Cas endonuclease cleavage site (e.g., 15-15 nucleotides). In other aspects, repair templates can share more than 100 nucleotides on either side of the edit site (e.g., >100~>100). Repair templates can be single, double, or staggered strands of complementary DNA using overhangs on the ends. In addition, these templates are not limited to DNA but also could be RNA or modified nucleotides (inosine, for example) or a mixture of these bases and have 5' and/or 3' ends protected from degradation. Because there are exonucleases that cleave nucleotides from the ends, protecting the ends with modified bases can prevent exonuclease digestion in the cell

of both DNA and RNA. Not to be limited by theory, increasing the length of the region of identity on both sides of the edit site can increase the efficiency and specificity of the repair process (e.g., homology arms). Included and provided for by the present specification are repair templates that can have 100% identity over at least 25 nucleotides on both sides of the edit site. This core identity region ensures that the desired edit (for example, deletions for SEQ ID NOs: 1 to 13) is accurate and efficient. In an aspect, the core identity region can be at least 40 nucleotides flanking the edit site. In yet another aspect, the core identity region can be at least 50 nucleotides on both sides of the edit site. In an aspect, the flanking core region can comprise 60 nucleotides of 100% identity. Also included in an aspect are repair templates that can have 70 identical nucleotides to the chromosomal region flanking the edit site. In an aspect, the core identity region can comprise 75 nucleotides of identity to the chromosomal region flanking the edit site. In an aspect, the core identity region can comprise 25 to 40 flanking nucleotides. In a further aspect, the core identity region can comprise 40 to 75 flanking nucleotides.

In some aspects, the core identity region can be further flanked by additional regions of homology to the target site (the "flanking homology regions"). As provided herein, in an aspect, the flanking homology regions can comprise 100% identity to the target site. As examples, SEQ ID NOs: 1 to 13 shared 100% homology to the CD163 region on either side of the targeted edit site as found in lines 2, 3, 15, 19, 27, 62, or 65 (e.g., 100% homology in the core region and the flanking homology regions). Also included are repair templates that can comprise SEQ ID NOs: 1 to 13, wherein the core identity region can comprise 100% identity to 25 nucleotides of the genome on either side of the desired edit (e.g., bases 50 to 100) and can be flanked by at least 80% identity to the genome on either side of the core region (e.g., nucleotides 1 to 49 and 101 to 150 of SEQ ID NOs: 1 to 18). In another aspect, the core region can have 100% homology and the flanking homology regions can share 85% homology. In a further aspect, the core region can have 100% homology and the flanking homology regions can share 90% homology. In yet another aspect, the core region can have 100% homology and the flanking homology regions can share 95% homology. Also included in an aspect are repair templates that can have a core region that has 100% homology and the flanking homology regions share 97% homology. In some aspects, the flanking homology regions can share 99% homology. Not to be limited by theory, it is thought that increasing the length of the flanking homology regions can introduce the desired CD163 edits into related animals without further modifying the genome. That is, specific repair sequences incorporating polymorphisms or changes in the CD163 genome of other pigs are specifically included and provided for.

The present specification includes, and provides for, flanking homology regions that can have greater than 50 nucleotides on each side of the edit site. In an aspect, the flanking homology regions can have greater than 75 nucleotides on each side of the edit site. In an aspect, the flanking homology regions can have greater than 100 nucleotides on each side of the edit site. Also included are flanking homology regions that can have greater than 200 nucleotides on each side of the edit site. In aspects, the flanking homology regions can have between 30 and 1000 nucleotides on each side of the edit site.

The present specification includes, and provides for, additional modifications at the 5' and 3' ends of repair template. Repair templates are not limited to DNA but also could be

RNA or modified nucleotides (inosine, for example) or a mixture of these bases and can have 5' and or 3' ends protected from degradation. Because there are exonucleases that cleave nucleotides from the ends, protecting the ends with modified bases can prevent exonuclease digestion in the cell of both DNA and RNA. Accordingly, as provided herein, the 5' and 3' ends of a repair template comprising a flanking homology region can be modified to prevent degradation.

The present specification provides for, and includes, ribonucleoprotein (RNP) complexes that can comprise a guide polynucleotide and a Cas protein. In an aspect, the Cas protein can be a *S. thermophilus* Cas9. In another aspect, the Cas protein can be Cas9 from *S. pyogenes*. Other suitable Cas proteins are known in the art and exemplary Cas systems are described above in Table 1. The selection of suitable Cas proteins depends on the required PAM sequence combination, and methods of identifying Cas proteins and modifying PAM sequences are known. Similarly, Cas systems differ in their requirements for guide RNA backbone sequences (e.g., tracrRNA sequences). In an aspect, the RNP complex can comprise a Cas protein and a guide nucleotide having at least 98% sequence identity to an RNA sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. In another aspect, the RNP complex can comprise a guide RNA comprising a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. In an aspect, the RNP complex can comprise a guide polynucleotide having 99% identity to a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. In an aspect, the guide polynucleotide of the RNP complex can comprise a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and 347 to 425. As provided herein, the RNP complex can comprise the sequences of any one of SEQ ID NOs: 22 to 271 and 347 to 425 combined with an RNA backbone as part of an sgRNA. In an aspect, the RNP complex can be pre-formed prior to injection into a target cell or can be injected or introduced separately.

Also provided for, and included, in the present specification are isolated guide RNAs that can comprise a spacer selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 and the first 20 nucleotides of each of SEQ ID NOs: 347 to 425. In an aspect, the Cas protein can be a protein comprising SEQ ID NOs: 20 or 21.

The present specification also includes, and provides for, DNA vectors that can encode guide RNAs for the preparation of CRISPR/Cas RNPs. In general, a vector encoding a guide RNA can comprise a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 or SEQ ID NOs: 347 to 425 and a guide RNA backbone arranged in cis and as part of a single transcription unit. Upon expression in a suitable cell, the DNA vector can produce an sgRNA of the present specification. Suitable expression vector backbones, including promoters, selectable markers and replication origins, are well known to persons of skill in the art. In practice, the 20 nucleotides and the guide RNA backbone can be DNA (when expressed from a promoter to be transcribed in vivo (in cells) or in vitro (via T7 polymerase) to form an RNA guide or the backbone can be chemically synthesized dual (crRNA and trRNA) guide or single guide RNA.

Specific guideRNAs can be paired to excise parts of the CD163 gene by using paired guideRNAs to delete sections

of the gene located between those guides. Exemplary pairs of guideRNAs (listed as DNA sequences) are provided in Table 5. Guide pairs without an exon 7 amino acid sequence listed remove the entire exon. The complete amino acid sequence for these deletions is set forth in SEQ ID NO: 553. All guides in Table 5 can create the desired sequence by excising the DNA in between the two guides and NHEJ repair between the cut sites without the use of a DNA repair template. The guide pairs that result in the amino acid sequences set forth in SEQ ID NOs. 506-517 can introduce an exogenous stop codon across the cut ends of the DNA introduced by the guide.

TABLE 5

Exemplary Guide RNA Pairs						
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Repaired Sequence SEQ ID	Exon 7AA SEQ ID
229	63322816	256	63323377	561	426	N/A
230	63322814	256	63323377	562	427	N/A
231	63322826	256	63323377	551	428	N/A
237	63322861	256	63323377	516	429	N/A
241	63322891	256	63323377	486	430	N/A
229	63322816	258	63323378	562	431	N/A
230	63322814	258	63323378	563	432	N/A
231	63322826	258	63323378	552	433	N/A
237	63322861	258	63323378	517	434	N/A
241	63322891	258	63323378	487	435	N/A
229	63322816	261	63323373	558	436	N/A
230	63322814	261	63323373	559	437	N/A
231	63322826	261	63323373	548	438	N/A
237	63322861	261	63323373	513	439	N/A
241	63322891	261	63323373	483	440	N/A
219	63322697	256	63323377	680	441	N/A
221	63322709	256	63323377	668	442	N/A
224	63322747	256	63323377	630	443	N/A
227	63322786	256	63323377	591	444	N/A
219	63322697	258	63323378	681	445	N/A
221	63322709	258	63323378	669	446	N/A
224	63322747	258	63323378	631	447	N/A
227	63322786	258	63323378	592	448	N/A
219	63322697	261	63323373	677	449	N/A
221	63322709	261	63323373	665	450	N/A
224	63322747	261	63323373	627	451	N/A
227	63322786	261	63323373	588	452	N/A
249	63322963	256	63323377	414	453	N/A
250	63322973	256	63323377	404	454	N/A
249	63322963	258	63323378	415	455	N/A
250	63322973	258	63323378	405	456	N/A
249	63322963	261	63323373	411	457	N/A
250	63322973	261	63323373	401	458	N/A
351	63323023	365	63323103	80	459	506
351	63323023	387	63323235	212	460	506
348	63323027	390	63323236	209	461	507
348	63323027	388	63323236	209	462	507
348	63323027	395	63323284	257	463	507
352	63323035	365	63323103	68	464	508
352	63323035	387	63323235	200	465	508
352	63323035	399	63323283	248	466	508
353	63323038	365	63323103	65	467	509
353	63323038	387	63323235	197	468	509
353	63323038	399	63323283	245	469	509
354	63323039	390	63323236	197	470	509
354	63323039	388	63323236	197	471	509
354	63323039	395	63323284	245	472	509
358	63323056	361	63323077	21	473	510
358	63323056	362	63323087	31	474	510
358	63323056	368	63323124	68	475	510
358	63323056	384	63323203	147	476	510
358	63323056	394	63323272	216	477	510
358	63323056	399	63323283	227	478	511
359	63323057	390	63323236	179	479	511
359	63323057	388	63323236	179	480	511
359	63323057	395	63323284	227	481	511
360	63323058	368	63323124	66	482	511

TABLE 5-continued

Exemplary Guide RNA Pairs						
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Repaired Sequence SEQ ID	Exon 7AA SEQ ID
360	63323058	384	63323203	145	483	511
360	63323058	389	63323237	179	484	511
360	63323058	394	63323272	214	485	511
360	63323058	397	63323285	227	486	511
361	63323077	365	63323103	26	487	512
361	63323077	387	63323235	158	488	512
362	63323087	390	63323236	149	489	513
362	63323087	388	63323236	149	490	512
362	63323087	395	63323284	197	491	513
364	63323089	365	63323103	14	492	514
364	63323089	387	63323235	146	493	514
364	63323089	399	63323283	194	494	514
365	63323103	368	63323124	21	495	515
365	63323103	384	63323203	100	496	516
365	63323103	389	63323237	134	497	516
365	63323103	394	63323272	169	498	516
365	63323103	397	63323285	182	499	516
366	63323118	368	63323124	6	500	517
366	63323118	384	63323203	85	501	517
366	63323118	389	63323237	119	502	517
366	63323118	394	63323272	154	503	517
366	63323118	397	63323285	167	504	517
354	63323039	211	63323163	123	505	518

In aspects, the guide RNA-Cas protein pairs can further comprise a DNA repair template. Exemplary guides and repair templates creating stop codons are listed in Table 6. These and other guides can be paired with other repair templates to generate in-frame stop codons or perturb, interfere, or eliminate splicing of exons, as examples of disrupting CD163 mRNA translation or processing. Other examples can include, but are not limited to, the elimination of the start ATG codon, or programming repair outcomes when using paired guide RNAs where a repair template could promote a single repair outcome over random non-homologous end joining when paired nucleases remove an exon (e.g., deletion of exon 7 corresponding to domain 5) in CD163. Without being limited by theory, analysis reveals consistent reduction of bioinformatic mismatches with inclusion of the extra nG in the PAM (nGGnG) as well as reduced off-target cutting in vitro and in vivo. Reduction of mismatches can include those from an alignment perspective (whether the DNA sequence has mismatches with the RNA guide sequence). See Table 4. A mismatch may not be an off-target edit (cutting or presence of indel, in vitro or in vivo, respectively), but an off-target edit is likely due to a mismatch. In the most preferred aspects, the methods disclosed herein produce no off-target edits in pigs, and the pigs disclosed herein have no off-target edits in their genomes.

TABLE 6

Guide RNAs and repair templates for editing Exon 2 of CD163							
Region Chr5 Location of guideRNAs binding site	63301998- 02017	63301997- 02016	63303213- 03232	63303283- 03302	63303315- 03334	63303315- 03334	63303338- 03357
Repair template	1	2	3	4	5	6	7
SEQ ID NO							
Guide RNA	274	275	276	277	278	279	280
spacer (DNA)							
SEQ ID NO							
Guide RNA	287	288	289	290	291	292	293
spacer (RNA)							
SEQ ID NO							
WT Region	300	301	302	303	304	305	306
SEQ ID NO.							
Trans-lation of WT region	313	314	315	316	317	318	319
Translation of	326	327	328	329	330	331	332
Deletion							
Base deletion	chr5:	chr5:	chr5:	chr5:	chr5:	chr5:	chr5:
coordinates	63301999- 63302005	63301999- 63302005	63303234- 63303235	63303304- 63303307	63303336- 63303336	63303337- 63303337	63303351- 63303356
Deleted bases (#)	CTTGGTC (7)	CTTGGTC (7)	GG (2)	GGCT (4)	G (1)	G (1)	CTTGTC (6)
Region Chr5 Location of guideRNAs binding site	63303379- 03398	63303379- 03398	63306148- 06167	63306263- 06282	63306364- 06383	63323147- 23166	
Repair template	8	9	10	11	12	13	
SEQ ID NO							
Guide RNA	281	282	283	284	285	286	
spacer (DNA)							
SEQ ID NO							
Guide RNA	294	295	296	297	298	299	
spacer (RNA)							
SEQ ID NO							
WT Region	307	308	309	310	311	312	
SEQ ID NO.							
Trans-lation of WT region	320	321	322	323	324	325	
Translation of	333	334	335	336	337	338	
Deletion							
Base deletion	chr5:	chr5:	chr5:	chr5:	chr5:	chr5:	
coordinates	63303399- 63303401	63303400- 63303402	63306156- 63306158	63306282- 63306284	63306373- 63306377	63323159- 63323162	
Deleted bases (#)	GGG (3)	GGG (3)	TTC (3)	CTC (3)	CGATC (5)	CGGC (4)	

The stochastic natural cellular repair process of non-homologous end joining (NHEJ) repair sometimes involves inserting or deleting single or multiple nucleotides at the double-strand break. As a consequence of this repair process, if a double-strand break occurs in the coding region of a gene, a shift in the translational reading frame of the encoded mRNA can result until an in-frame stop codon, terminating protein translation. Although the translation of this naturally, accidentally mutated gene can produce a shortened protein relative to the protein product of an unmodified version, the amino acid sequence of this newly encoded polypeptide would be unique, possibly reducing or even improving the fitness of cells within the target organism. In an aspect, to preclude the creation of frame shifting mutations and the translation of this set of undesirable polypeptides, edited pigs can be screened for formation of an in-frame translational stop codon at or near the endonuclease cut site that is a consequence of a separate, naturally occurring NHEJ repair.

The excision of genomic DNA sequence using two intronic guideRNAs in conjunction with an endonuclease can be accomplished by NHEJ repairs that include, but are

45

not limited to, the end-to-end joining of nuclease cut sites. Because introns are non-coding, NHEJ repair outcomes that include indels around the nuclease cut sites can accomplish the excision of the intended region of DNA which can include intron and/or exon sequences. Because some NHEJ repair outcomes occur more frequently than others, it is advantageous to consider repair outcome frequency when choosing a guideRNA pair for use in gene editing experiments. SEQ ID NO: 520-555 illustrate repair outcomes of exon 7 excisions observed in blastocysts from guide pairs as set forth in Table 7. The designed repair outcome for these guide pairs is listed in Table 5.

TABLE 7

Observed Repair Outcomes in Blastocysts		
SEQ ID NO: (5')	SEQ ID NO: (3')	Repair outcome SEQ ID
249	261	520
249	261	521
249	261	522

60

65

TABLE 7-continued

Observed Repair Outcomes in Blastocysts		
SEQ ID NO: (5')	SEQ ID NO: (3')	Repair outcome SEQ ID
249	261	523
249	261	524
249	261	525
249	261	526
249	261	527
249	256	528
249	256	529
249	256	530
249	256	531
249	256	532
249	256	533
249	256	534
249	256	535
241	258	536
241	258	537
241	258	538
241	258	539
241	258	540
241	258	541
241	258	542
241	258	543
241	258	544
221	261	545
221	261	546
221	261	547
221	261	548
221	261	549
229	256	550
229	256	551
229	256	552
229	256	553
229	256	554
229	256	555

In another aspect, during gene editing, in-frame stop codons that do not result in the addition of new amino acids can be created by including a DNA repair template together with the guide RNA-Cas protein pair. In an aspect, a DNA repair template can be a dsDNA. In another aspect, the DNA repair template can be a ssDNA. Co-introduction of a double- or single-stranded DNA repair template can be used to either delete or insert DNA nucleotides to form an in-frame translational stop codon (TAA, TGA, TAG) at or near the double-strand break site initiated by the endonuclease. DNA repair templates can further comprise polynucleotide modification templates containing several nucleotide changes in comparison to the native sequence, which can directly edit the target DNA sequence and can be co-transfected with the endonuclease editing reagents to generate edited CD163 genes with in-frame stop codons. In an aspect, the encoded proteins from these gene edited CD163 genes, when transcribed into mRNA and then translated into protein, can only synthesize a shortened and non-functional form of the CD163 polypeptide. Not to be limited by theory, DNA repair templates having longer regions of sequence homology can be more efficient. In certain embodiments, DNA repair templates can contain regions of sequence identity (homology arms) within the DNA repair template which can flank the sequence change and can range from fewer than 50 nucleotides to greater than 1000 nucleotides.

Several aspects of Cas9-protein/guide RNA combinations together with DNA repair templates are shown in Table 6. As provided in Table 6, several Cas9-protein/guide RNA combinations can be paired with DNA repair templates having, but not being limited to, 75 bases of sequence homology on each side of the targeted deletion (50 core homology bases

and flanking homology regions of 50 on either side). In an aspect, the protein-guide RNA complexes and template combinations can be transfected into cells. Without being limited by theory, repair of the double-strand break using the DNA repair template can direct the formation of an in-frame translational stop codon as the result of deletion of single or multiple nucleotides at or adjacent to the break site. Animals and cells obtained from this method demonstrate that endonuclease-directed double-strand breaks at the porcine CD163 gene can be repaired using a co-introduced DNA repair template. As provided herein, repair templates having a sequence of SEQ ID NOs: 1 to 13 can introduce an in-frame translational stop codon in Exon 2, thereby producing a shortened and non-functional CD163 protein.

Use of a DNA repair template according to the present specification is not limited to abolishing the function of CD163. The present specification further includes, and provides for, a repair template that can direct the removal or addition of nucleotides to the gene. In an aspect, the stability or half-life modulation of the encoded CD163 mRNA can be modulated by editing according to the present methods. In yet another aspect, DNA sequences which encode amino acids of the mature protein responsible for binding PRRS virus can be removed or replaced. In an aspect, CD163 expression and or activity can be reduced by at least 90% but not abolished. In an aspect of the present teachings, using the spacer sequences described with other repair templates would be contemplated by those skilled in the art to allow for the formation of an in frame stop codon by the removal of bases and the introduction of bases or a combination thereof.

Included, and provided for, by the present specification are pigs derived from elite porcine lines comprising edited CD163 genes. In aspects, the elite porcine lines can be PICT[™] Line 15, PICT[™] Line 17, PICT[™] Line 27, PICT[™] Line 65, PICT[™] Line 14, PICT[™] Line 62, PIC337, PIC800, PIC280, PIC327, PIC408, PICT[™] 399, PIC410, PIC415, PIC359, PIC380, PIC837, PIC260, PIC265, PIC210, PICT[™] Line 2, PICT[™] Line 3, PICT[™] Line 4, PICT[™] Line 5, PICT[™] Line 18, PICT[™] Line 19, PICT[™] Line 92, PIC95, PICT[™] CAMBOROUGH® (Pig Improvement Company, Limited, Basingstoke, UK), PIC1070, PICT[™] CAMBOROUGH® 40, PICT[™] CAMBOROUGH® 22, PIC1050, PICT[™] CAMBOROUGH® 29, PICT[™] CAMBOROUGH® 48, or PICT[™] CAMBOROUGH® x54. In various aspects, the elite porcine lines can be PICT[™] elite porcine lines 2, 3, 15, 19, 27, 62, or 65. In another aspect, the pigs can comprise edited CD163 genes derived from elite porcine lines. In aspects, the pigs can comprise an edited CD163 in a CD163 genomic region having the genotypes shown in Table 8 to Table 14. Table 8 to Table 14 present each position on chromosome 5 in the vicinity of the CD163 gene in which a single nucleotide polymorphism exists.

Each table presents alleles that are homozygous in that line and provides a distinguishing genetic signature of the line, as well as its unique genome edited region. All the genetic signatures are based on the Sscrofa11.1 reference genome (GenBank accession: GCA_000003025.6). Accordingly, the transmission of the edited CD163 line can be followed, and animals comprising an edited CD163 gene obtained from the line can be identified. Pigs of other lines are heterozygous or opposite genotype at multiple alleles when compared to the lines of Table 8 to Table 14. In this way, the combination of genetic signatures in each line can be used to distinguish between pigs belonging to a particular line and pigs not belonging to a particular line. In an aspect, the genetic signatures provide for methods to breed and

47

track the CD163 edited genomes from generation to generation. In an aspect, progeny generations that comprise the CD163 edited genomes having a genetic signature according to Table 8 to Table 14 can be prepared.

TABLE 8

Line 2 Genetic Signatures	
Position	Genotype
60306428	C/C
60306527	C/C
60308030	C/C
60320580	C/C
60322529	A/A
60324162	T/T
60327987	C/C
60328009	T/T
60335421	T/T
60338654	G/G
60344946	A/A
60345163	C/C
60345689	G/G
60345715	G/G
60345722	A/A
60345749	G/G
60345775	A/A
60345825	A/A
60346409	G/G
60346607	A/A
60346641	G/G
60346691	T/T
60346734	A/A
60347297	G/G
60350295	C/C
60350343	G/G
60350448	C/C
60350470	T/T
60350475	C/C
60350564	A/A
60350571	A/A
60350572	A/A
60350911	C/C
60351055	G/G
60351604	C/C
60351855	T/T
60351857	C/C
60351972	G/G
60352165	C/C
60352923	C/C
60353408	A/A
60353562	T/T
60353576	C/C
60353659	T/T
60353721	G/G
60354428	G/G
60354925	C/C
60355415	G/G
60355420	T/T
60355448	G/G
60355529	A/A
60355530	A/A
60355774	C/C
60356277	C/C
60356351	C/C
60356575	C/C
60356578	T/T
60356861	G/G
60356885	T/T
60356898	T/T
60356914	C/C
60357001	T/T
60357014	T/T
60358409	C/C
60358449	G/G
60358469	T/T
60358475	T/T
60358551	A/A
60358568	G/G
60358641	T/T

48

TABLE 8-continued

Line 2 Genetic Signatures	
Position	Genotype
60358704	T/T
60358774	T/T
60368913	C/C
60368916	A/A
60368921	T/T
60368940	G/G
60368963	C/C
60416097	G/G
60543479	C/C
60614797	G/G
60614964	G/G
60920926	G/G
61479206	C/C
61479353	T/T
61653091	C/C
61783688	A/A
62091204	A/A
62416522	A/A
62476273	C/C
62687883	C/C
63044336	A/A
63546615	C/C
63553656	G/G
63558087	G/G
63640722	G/G
63792071	T/T
64009226	C/C
64459105	T/T
64460101	A/A
64527707	A/A
64577968	C/C
64943306	T/T
65242696	T/T
65242725	A/A
65242729	A/A
65242736	G/G
65249484	T/T
65260283	T/T
65269510	C/C
65273886	T/T
65276133	C/C
65277053	T/T
65277156	C/C
65277320	G/G
65280282	A/A
65282683	C/C
65283434	C/C
65283970	C/C
65284829	C/C
65285100	C/C
65285902	G/G
65285914	T/T
65286400	C/C
65294497	T/T
65383805	C/C
65405986	C/C
65410118	C/C
65410147	G/G
65410198	C/C
65410435	C/C
65410440	T/T
65410441	C/C
65410443	C/C
65410447	C/C
65411265	C/C
65411431	G/G
65411524	C/C
65411698	G/G
65411831	C/C
65411848	G/G
65411854	T/T
65411966	C/C
65412729	G/G
65413215	G/G
65413759	C/C
65417199	A/A

49

TABLE 8-continued

Line 2 Genetic Signatures	
Position	Genotype
65417256	G/G
65417261	A/A
65417273	C/C
65417287	G/G
65417716	G/G
65417797	C/C
65417800	C/C
65419169	G/G
65419410	G/G
65420017	T/T
65420184	G/G
65420415	A/A
65420591	G/G
65420680	C/C
65420681	A/A
65420693	G/G
65420947	G/G
65420949	T/T
65421133	G/G
65421195	G/G
65421255	G/G
65421365	G/G
65421421	C/C
65422005	G/G
65422008	G/G
65422057	T/T
65422141	C/C
65422701	C/C
65422725	C/C
65425730	G/G
65601310	C/C
65602328	C/C
65711602	C/C
65761990	C/C
66293766	G/G
66296268	A/A

TABLE 9

Line 3 Genetic Signatures	
Position	Genotype
60414864	T/T
60415813	C/C
60415854	G/G
60572371	A/A
60663464	C/C
60663499	A/A
61016527	G/G
61442674	T/T
61870484	C/C
61989476	G/G
62510031	G/G
62602514	T/T
63044336	A/A
63345577	G/G
63452493	A/A
63669675	G/G
63754659	T/T
63810627	C/C
63810632	G/G
63810707	T/T
63810733	G/G
63810819	C/C
63810857	C/C
63810861	G/G
64010555	T/T
64010599	A/A
64452338	T/T
64455982	C/C
64457532	T/T
64458163	T/T

50

TABLE 9-continued

Line 3 Genetic Signatures	
Position	Genotype
64458711	G/G
64555715	G/G
64561148	G/G
64561209	G/G
64561659	G/G
64561828	T/T
64561873	G/G
64561935	C/C
64562021	C/C
64562771	C/C
64563462	G/G
64682243	C/C
64682929	G/G
64685072	T/T
64692673	G/G
64692905	T/T
64693058	T/T
64705228	G/G
64720067	A/A
64722426	T/T
64722539	C/C
64726936	G/G
64729340	C/C
64730596	G/G
64736752	G/G
64761650	G/G
64767369	G/G
64768411	G/G
64769978	T/T
64769989	C/C
64770307	G/G
64773053	A/A
64774093	G/G
64775512	G/G
64775565	C/C
64780089	C/C
64780152	C/C
64780196	C/C
64783169	G/G
64783774	C/C
64783896	G/G
64784000	T/T
64784139	C/C
64784235	T/T
64784410	A/A
64784653	A/A
64784725	C/C
64784844	A/A
64784875	T/T
64785001	A/A
64785078	C/C
64813032	G/G
64819974	C/C
64893099	G/G
64899922	A/A
64934758	C/C
65348500	G/G
65546268	A/A
65546327	G/G
65841270	T/T
65858531	G/G
65979134	G/G
65985520	C/C
66035868	A/A
66042409	C/C
66042542	C/C
66042551	C/C
66042557	A/A
66042563	G/G
66155827	A/A
66155859	A/A
66155863	T/T
66156160	A/A
66156183	G/G
66156533	G/G
66167582	C/C

51

TABLE 9-continued

Line 3 Genetic Signatures	
Position	Genotype
66167809	C/C
66167823	T/T
66167877	A/A
66167888	C/C
66167905	A/A
66168135	C/C
66168261	C/C
66168311	G/G
66168615	G/G
66168688	G/G
66168742	G/G
66168880	G/G
66168947	T/T
66168952	A/A
66168993	A/A

TABLE 10

Line 15 Genetic Signatures	
Position	Genotype
60391943	G/G
60392234	T/T
60394009	A/A
60414796	T/T
60601475	A/A
60630898	A/A
60630910	A/A
60786421	G/G
60787517	G/G
60792231	G/G
60795724	C/C
60915104	C/C
61016227	A/A
61086145	T/T
61275101	C/C
61396087	A/A
61434042	G/G
61515967	T/T
61759785	C/C
61759821	C/C
61865586	A/A
61867498	C/C
62192420	G/G
62195570	A/A
62196101	C/C
62442547	G/G
62509196	T/T
62675268	G/G
62843969	G/G
62852334	A/A
63025151	A/A
63025152	C/C
63168616	A/A
63266682	T/T
63367266	G/G
63424311	T/T
63535058	T/T
63653754	G/G
63655538	T/T
63669242	A/A
63669944	C/C
63669946	G/G
63900332	A/A
64138082	C/C
64171653	C/C
64471845	G/G
64472363	A/A
64472504	T/T
64472616	T/T
64523343	A/A
64536599	G/G

52

TABLE 10-continued

Line 15 Genetic Signatures	
Position	Genotype
64537982	T/T
64893205	A/A
64916422	G/G
65039734	C/C
65175496	C/C
65177506	G/G
65294016	C/C
65571397	A/A
65573608	T/T
65576707	C/C
65984917	G/G
65984984	T/T
65985058	T/T
65985392	C/C
66042472	C/C
66099282	A/A
66119075	T/T

TABLE 11

Line 19 Genetic Signatures	
Position	Genotype
60320100	A/A
60320144	G/G
60320580	C/C
60507836	C/C
60512716	G/G
60564418	A/A
60666662	T/T
60683738	T/T
60782197	T/T
60782278	T/T
60782642	A/A
60782683	G/G
60782756	G/G
60782997	G/G
60783031	C/C
60783198	G/G
61131519	G/G
61131736	C/C
61189029	C/C
61359517	A/A
61390748	C/C
61465215	T/T
61567329	C/C
61567357	A/A
61567365	A/A
61567410	C/C
61567494	C/C
61567594	G/G
61975156	G/G
61975367	G/G
61975382	A/A
61975388	C/C
62363513	T/T
62367647	G/G
62378299	G/G
62610431	T/T
62952267	T/T
62965473	T/T
63108711	G/G
63109224	T/T
63174233	T/T
63174257	A/A
63393845	G/G
63393849	T/T
63393851	T/T
63860532	A/A
63861247	A/A
63862368	T/T
64138082	C/C

53

TABLE 11-continued

Line 19 Genetic Signatures	
Position	Genotype
64171653	C/C
64181448	G/G
64402245	T/T
64455172	G/G
64514012	C/C
64514013	A/A
64522169	G/G
64522173	T/T
64522177	T/T
64522178	C/C
64794635	G/G
65031621	C/C
65031630	C/C
65031684	T/T
65222383	T/T
65222385	A/A
65242693	C/C
65497229	G/G
65499233	C/C
65502398	G/G
65562917	C/C
65566225	T/T
65567066	T/T
65894449	C/C
65904537	T/T
65959573	A/A
65998062	C/C
66158553	A/A
66158950	C/C
66159103	A/A
66159212	G/G

TABLE 12

Line 27 Genetic Signatures	
Position	Genotype
60347934	A/A
60356984	C/C
60692943	C/C
60693260	G/G
60693769	G/G
60938885	T/T
60960728	A/A
61436306	G/G
61464114	C/C
61468375	G/G
61736415	A/A
61736429	G/G
61757961	G/G
61820110	A/A
61822513	T/T
61858837	G/G
62159353	T/T
62313847	C/C
62315464	A/A
62570257	C/C
62570793	A/A
62571313	T/T
62917536	G/G
62917597	T/T
62917618	C/C
62917629	T/T
62917780	G/G
62917866	T/T
62917889	T/T
62918448	G/G
62918456	A/A
62918465	A/A
62918655	T/T
63042154	C/C
63043261	G/G

54

TABLE 12-continued

Line 27 Genetic Signatures	
Position	Genotype
63046692	G/G
63433705	G/G
63490990	A/A
63626018	A/A
63626423	T/T
63626826	C/C
63630149	A/A
63631098	C/C
63977777	A/A
63977821	G/G
64460204	C/C
64582243	C/C
64582246	G/G
64582272	A/A
64600735	G/G
65056396	A/A
65056671	T/T
65332987	A/A
65335837	C/C
65390844	A/A
65404042	A/A
65404045	A/A
65404063	C/C
65705261	G/G
65740018	T/T
65740030	A/A
66235840	G/G

TABLE 13

Line 62 Genetic Signatures	
Position	Genotype
60336383	A/A
60343172	A/A
60345254	G/G
60447765	G/G
60452143	C/C
60452167	T/T
60462769	T/T
60653700	T/T
60654939	A/A
60654943	A/A
61085770	A/A
61108831	G/G
61117357	A/A
61234908	C/C
61331270	A/A
61388824	G/G
61755575	C/C
61759095	C/C
61760230	C/C
61927492	T/T
61927506	A/A
62006220	T/T
62115641	T/T
62169134	T/T
62368918	C/C
62471593	C/C
62474001	A/A
62474006	C/C
62474365	A/A
62480042	A/A
62480261	G/G
62480699	A/A
62480892	A/A
62481902	G/G
62482914	T/T
62483893	C/C
62483897	T/T
62847072	G/G
62874766	G/G

55

TABLE 13-continued

Line 62 Genetic Signatures	
Position	Genotype
62897778	A/A
63100290	C/C
63219344	G/G
63361759	G/G
63388357	C/C
63442314	G/G
63590333	T/T
63670724	G/G
63714834	A/A
63792005	T/T
63985848	A/A
63985923	G/G
64131088	C/C
64471400	C/C
64483848	G/G
64484177	T/T
64689271	A/A
64758235	C/C
64790188	C/C
64895383	C/C
64991645	A/A
65045805	T/T
65146186	C/C
65368620	C/C
65432698	C/C
65542716	A/A
65637507	G/G
65771614	C/C
65771615	G/G
65772162	T/T
66033248	T/T
66033476	C/C
66033484	C/C

TABLE 14

Line 65 Genetic Signatures	
Position	Genotype
60322300	C/C
60411747	A/A
60412758	G/G
60412775	A/A
60412826	C/C
60413294	G/G
60459719	A/A
60462179	C/C
60492928	A/A
60492938	T/T
60666662	T/T
60722708	C/C
60770183	G/G
60770197	C/C
60777431	C/C
60781592	T/T
60782197	T/T
60782278	T/T
60782642	A/A
60782683	G/G
60782756	G/G
60782997	G/G
60783006	G/G
60783031	C/C
60783198	G/G
60791719	T/T
60900261	G/G
60911551	A/A
60912669	A/A
61223574	A/A
61486607	C/C
61756324	G/G
61832527	C/C

56

TABLE 14-continued

Line 65 Genetic Signatures	
Position	Genotype
61895566	C/C
61895833	A/A
62354225	C/C
62364072	G/G
62383525	G/G
62399452	T/T
62404734	A/A
62432982	C/C
62841388	G/G
62931496	T/T
63156406	C/C
63156417	T/T
63266198	A/A
63383925	A/A
63392495	T/T
63549670	C/C
63593491	G/G
63609938	A/A
63641998	C/C
63678793	G/G
63784770	G/G
63810897	C/C
63810903	C/C
63810920	G/G
63810922	T/T
63810928	G/G
63872757	C/C
63944636	G/G
63944765	C/C
64138639	C/C
64194054	G/G
64242768	T/T
64270989	G/G
64753488	G/G
64758888	A/A
64758900	C/C
64759492	T/T
64795760	T/T
64797970	A/A
64798690	G/G
64860395	G/G
64860396	C/C
64862131	C/C
65371202	T/T
65371472	G/G
65371901	G/G
65371970	C/C
65372072	A/A
65409493	A/A
65437347	G/G
65573362	T/T
65601655	G/G
65747069	T/T
65843843	A/A
65849424	T/T
66248756	A/A

Elite PIC™ lines 2, 3, 15, 19, 27, 62 and 65 are lines selected for superior commercial phenotypes. In an aspect, the CD163 gene edited cells and animals can be free of deleterious alleles that are present in wild populations and in many commercial herds. In aspects, CD163 gene edited cells and animals can be free of one or more of the deleterious alleles selected from the group consisting of epetheliogenesis imperfecta, melanotic skin tumors, dermatosis vegetans, abnormal mammae, shortened vertebral column, kinky tail, rudimentary tail, Hairlessness, Hairlessness (2), Woolly hair, Hydrocephalus, Tassels, Legless, Three-legged, Syndactyly, Polydactyly, Pulawska factor, Heterochromia iridis, Congenital tremor A III, Congenital tremor A IV, Congenital ataxia, Hind leg paralysis, Bentleg, Thickleg, Malignant hyperthermia, Hemophilia (von Willebrand's disease), Leu-

kemia, Hemolytic disease, edema, Acute respiratory distress ("barker"), Rickets, 25 Renal hypoplasia, Renal cysts, Uterus aplasia, Porcine Stress Syndrome (PSS), halothane (HAL), Dipped Shoulder (Humpy Back, Kinky Back, Kyphosis), Hyperostosis, Mammary Hypoplasia, and Undeveloped Udder. As provided herein, the improved methods of preparing CD163 gene edited animals can avoid introducing new mutations at the deleterious loci or generating new amorphic, hypomorphic, hypermorphic, neomorphic, antimorphic mutations at non-target sites. These latter changes can be particularly undesirable in elite lines as they can interfere with genes involved in desirable traits that can be controlled by multiple loci in a continuous, quantitative way in a population. Many such Quantitative Trait Loci (QTL) are known and can be typically characterized by a bell-shaped curve the trait value can be plotted against the number of observed animals. Such polygenic inheritance of traits can be common among traits recognized as commercially important such as, but not limited to, backfat, average daily feed intake, lifetime daily gain, and loin depth.

Similarly, traits associated with productivity can be multigenic and controlled by multiple QTLs. These traits were typically measured by visual inspection but methods now can include ultrasound to measure backfat thickness (bfp), loin depth (ldp) and intramuscular fat (uip). As provided in Table 15, elite lines can have desirable phenotypic traits including high backfat and loin depth while having high lifetime daily gain. Similarly, the sows of the elite lines can be fecund and have large litters, few still-born, and sufficient teats to wean and nurse the piglets.

TABLE 15

Desirable Phenotypic Traits			
LINE	TRT	average	standard deviation
2	Backfat, mm	7.8	1.81
3	Backfat, mm	9.44	2.77
15	Backfat, mm	8.16	2.03
65	Backfat, mm	7.45	1.88
2	Average Daily Feed Intake, kg	1.99	0.24
3	Average Daily Feed Intake, kg	2.13	0.25
15	Average Daily Feed Intake, kg	2.22	0.24
65	Average Daily Feed Intake, kg	2.2	0.27
2	Lifetime Daily Gain, grams/day	683.83	73.53
3	Lifetime Daily Gain, grams/day	704.23	80.15
15	Lifetime Daily Gain, grams/day	755.61	69.32
65	Lifetime Daily Gain, grams/day	800.6	84.48
2	Loin depth, mm	65.31	7.09
3	Loin depth, mm	62.61	7.14
15	Loin depth, mm	67.17	7.15
65	Loin depth, mm	81.2	8
2	Total Born per litter	13.71	3.19
3	Total Born per litter	15.06	3.41
15	Total Born per litter	10.39	2.71
65	Total Born per litter	10.31	2.52
2	Still born per litter	1.19	1.43
3	Still born per litter	1.25	1.50
15	Still born per litter	1.22	1.37
65	Still born per litter	0.77	1.16
2	Teat number	15.43	1.27
3	Teat number	15.44	1.29

The present specification provides for, and includes, gene edited pigs of selected elite lines that can be homozygous for CD163 knockout edits (CD163^{-/-}). In an aspect, the line can be selected from the group consisting of PICTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, Line 65, and progeny thereof comprising the edited CD163 genes described herein. Gene edited lines 2, 3, 15, 19, 27, 62, and 65 can comprise CD163 genomic regions as provided at Table 8 to

Table 14, and can be readily distinguished from each other, from unimproved lines, and from other elite lines. In aspects according to the present specification, the heterozygous and homozygous pigs of the present specification can be free of off-site mutations. The present specification provides for, and includes, pigs and cells that can have edited CD163 genes comprising SEQ ID NOs: 1 to 18 and 426 to 505 and that can share a genetic signature comprising at least 90% of the genotypic markers of Table 8 to Table 14. The present specification provides for, and includes, pigs and cells that can have edited CD163 genes comprising SEQ ID NO: 2 and sharing a genetic signature comprising at least 90% of the genotypic markers of Table 8 to Table 14. In an aspect, the genetic signature of a CD163 edited pig or cell can share a genetic signature comprising at least 95% of the genotypic markers of Table 8 to Table 14. Also included are CD163 edited pigs or cells that can share a genetic signature comprising at least 97% of the genotypic markers of Table 8 to Table 14. In an aspect, the CD163 edited pigs or cells can share a genetic signature comprising at least 98% of the genotypic markers of Table 8 to Table 14. Another aspect provides for CD163 edited pigs or cells that can share a genetic signature comprising at least 99% of the genotypic markers of Table 8 to Table 14.

As provided herein, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, can retain desirable commercial traits and can be free of deleterious off-target mutations and can comprise an edited CD163 gene comprising any of SEQ ID NOs: 1 to 18 and 426-505. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, can comprise at least 90% of the loin depth of the non-edited pig line and can comprise an edited CD163 gene comprising any of SEQ ID NOs: 1 to 18 and 426-505. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, can comprise at least 90% of the lifetime daily gain of the non-edited pig line. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, can comprise at least 90% of the average daily feed intake of the non-edited pig line. The present specification provides for, and includes, pigs and cells that can have edited CD163 genes comprising SEQ ID NOs: 1 to 18 and 426 to 505 and can share a genetic signature comprising at least 90% of the genotypic markers of Table 8 to Table 14.

As provided herein, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, can retain desirable reproductive traits. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, comprising a gene-edited CD163^{-/-} of SEQ ID NOs: 1 to 18 or 426 to 505 can comprise at least 90% of the total born per litter of the non-edited pig line. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, comprising a gene-edited CD163^{-/-} of SEQ ID NOs: 1 to 18 or 426 to 505 can comprise no more than 110% of the number of still born piglets compared to the non-edited pig line. In aspects of the present disclosure, the gene-edited CD163^{-/-} animals and cells of Lines 2, 3, 15, 19, 27, 62, and 65, comprising a gene-edited CD163^{-/-} of SEQ ID NOs: 1 to 18 or 426 to 505 can comprise at least 90% of the average number of teats compared to the non-edited pig line.

In an aspect, the gene edited CD163^{-/-} animals and cells of Line 2 having a gene-edited CD163^{-/-} of SEQ ID NOs: 1 to 18 or 426 to 505 can have backfat that is at least 90% of the amount found in unedited Line 2 animals. In an aspect,

In an aspect, the gene edited CD163^{-/-} animals and cells of Line 2 having a gene-edited CD163^{-/-} of SEQ ID NOS:1 to 18 or 426 to 505 can have a number of total born per litter that is at least 90% of the amount found in unedited Line 2 animals. In an aspect, gene edited CD163^{-/-} animals and cells of Line 2 can have a number of total born per litter that is at least 95% of the unedited Line 2 animal. In an aspect,

The present specification provides for and includes CD163 edited animals comprising the germplasm of PICTTM Line 15, PICTTM Line 17, PICTTM Line 27, PICTTM Line 65, PICTTM Line 14, PICTTM Line 62, PIC337, PIC800, PIC280,

PIC327, PIC408, PICT[™] 399, PIC410, PIC415, PIC359, PIC380, PIC837, PIC260, PIC265, PIC210, PICT[™] Line 2, PICT[™] Line 3, PICT[™] Line 4, PICT[™] Line 5, PICT[™] Line 18, PICT[™] Line 19, PICT[™] Line 92, PIC95, PICT[™] CAMBOROUGH®, PIC1070, PICT[™] CAMBOROUGH® 40, PICT[™] CAMBOROUGH® 22, PIC1050, PICT[™] CAMBOROUGH® 29, PICT[™] CAMBOROUGH® 48, or PICT[™] CAMBOROUGH® ×54. As used herein, the term germplasm includes an intact genome present in cells or nuclei and comprising chromosomes. The term germplasm may include any gamete, germ cell, or any somatic cell from which an animal can be cloned. The edited germplasm can comprise an edit having an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The edited germplasm can comprise an edit having an edited genomic sequence of any one of SEQ ID NOs: 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, or 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 15. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 17. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 27. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 65. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 14. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 62. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC337. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC800. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC280. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC327. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC408. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] 399. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC410. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC415. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC359. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC380. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC837. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC260. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC265. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC210. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 2. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 3. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 4. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 5. The edited

germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 18. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 19. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 92. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC95. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH®. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC1070. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH® 40. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH® 22. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PIC1050. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH® 29. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH® 48. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] CAMBOROUGH® ×54.

The present specification provides for and includes CD163 edited animals comprising the germplasm of PICT[™] Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65. The edited germplasm can comprise an edit having an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The edited germplasm can comprise an edit having an edited genomic sequence of any one of SEQ ID NOs: 426, 427, 428, 429, 430, 431, 432, 433, 434, 435, 436, 437, 438, 439, 440, 441, 442, 443, 444, 445, 446, 447, 448, 449, 450, 451, 452, 453, 454, 455, 456, 457, 458, 459, 460, 461, 462, 463, 464, 465, 466, 467, 468, 469, 470, 471, 472, 473, 474, 475, 476, 477, 478, 479, 480, 481, 482, 483, 484, 485, 486, 487, 488, 489, 490, 491, 492, 493, 494, 495, 496, 497, 498, 499, 500, 501, 502, 503, 504, or 505. The edited germplasm can comprise an edit having an edited genomic sequence as set forth in SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 2 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 3 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 15 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 19 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 27 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 62 with a CD163 edited sequence comprising SEQ ID NO: 453. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 65 with a CD163 edited sequence comprising SEQ ID NO: 453.

The edited germplasm can comprise an edit having an edited genomic sequence as set forth in SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICT[™] Line 2 with a

CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 3 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 15 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 19 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 27 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 62 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 65 with a CD163 edited sequence comprising SEQ ID NO: 489. The edited germplasm can have a predicted exon 7 amino acid sequence from any one of SEQ ID NOs: 506-517. The edited germplasm can have a predicted exon 7 amino acid sequence as set forth in SEQ ID NO: 513.

The edited germplasm can comprise an edit having an edited genomic sequence as set forth in SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 2 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 3 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 15 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 19 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 27 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 62 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can comprise a genome that is 80%, 85%, 90%, 95% similar or identical to PICTTM Line 65 with a CD163 edited sequence comprising SEQ ID NO: 505. The edited germplasm can have a predicted exon 7 amino acid sequence as set forth in SEQ ID NO: 518.

The present specification also provides for and includes cells of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65. In some embodiments, a cell of PICTTM Line 2 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 2 can comprise an editing genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 3 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 3 can comprise an editing genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 15 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 15 can comprise an edited genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 19 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 19 can comprise an editing

genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 27 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 27 can comprise an editing genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 62 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 62 can comprise an editing genomic sequence as set forth in SEQ ID NO: 453. In some embodiments, a cell of PICTTM Line 65 can comprise an edited genomic sequence of any one of SEQ ID NOs: 426 to 505. The cell of PICTTM Line 65 can comprise an editing genomic sequence as set forth in SEQ ID NO: 453. The edited germplasm can have a predicted exon 7 amino acid sequence as set forth in SEQ ID NO: 513.

The present specification provides for, and includes, hybrid animals that can comprise the CD163 gene edits characterized by SEQ ID NOs: 1 to 18 and 426 to 505. In some configurations, gene edit can comprise SEQ ID NO: 453. In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® (PICTTM UK Limited, Basingstoke, UK) line. CAMBOROUGH® pigs are hybrids that can be prepared by a cross between Line 2 and Line 3. In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® line having an edited CD163 gene of SEQ ID NO: 2. In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® line having an edited CD163 gene of SEQ ID NO: 426-458. In various aspects, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® line having an edited CD163 gene of SEQ ID NO: 453. In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® line having an edited CD163 gene of SEQ ID NO: 459-504. In various aspects, the hybrid animal can be a CD163^{-/-} hybrid animal of the CAMBOROUGH® line having an edited CD163 gene sequence of SEQ ID NO: 489. CAMBOROUGH® hybrid pigs are pigs that have large litters with uniform and vigorous piglets. CAMBOROUGH® hybrid pigs have a long productive life and have a low mortality. CAMBOROUGH® CD163^{-/-} hybrid pigs retain these desirable commercial traits. In an aspect, CAMBOROUGH® CD163^{-/-} hybrid pigs have litter sizes that are indistinguishable from non-gene edited CAMBOROUGH® hybrid pigs. In aspects according to the present specification, the heterozygous and homozygous pigs of the present specification are free of off-site mutations.

The present specification further provides, and includes, methods for preparing CD163 gene edited hybrid animals. In an aspect, a first parent can comprise a CD163 gene edited boar, gilt, or sow of one of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65 for crossing to a second parent. In an aspect, the genome of a gene edited first parent can comprise a sequence selected from the group consisting of SEQ ID NOs: 1 to 18 or 426 to 505. In some aspects, the genome can comprise SEQ ID NO: 453. In various aspects, the genome can comprise SEQ ID NO: 489. In various aspects, the genome can comprise SEQ ID NO: 505. In an aspect, the method can comprise a second parent selected from the group consisting of a CD163 gene edited boar, gilt, or sow of one of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65. Also provided for, and included, by the present specification are methods of preparing CD163 edited animals that can comprise crossing a progeny of any one of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65.

In an aspect, the present specification provides for heterozygous pigs from lines 2, 3, 15, and 65 (Table 8, Table 9, Table 10, and Table 14, respectively) that can have at least one copy of the CD163 gene successfully edited in a knock-out edit. These edited pigs can exhibit a healthy phenotype with no noticeable deleterious effects from the edit. In another aspect, the specification provides for pigs from lines 19, 27, and 62 (Table 11, Table 12, and Table 13, respectively) that can be edited using the methods disclosed herein to generate edited pigs with healthy phenotypes. The heterozygous pigs can be crossed with non-edited animals of the corresponding line to produce F1 heterozygous pigs. In an embodiment, heterozygous pigs of lines 2, 3, 15, and 65 can be crossed to a second heterozygous pig of lines 2, 3, 15, and 65, and homozygous CD163 edited pigs can be produced in Mendelian proportions. Notably, the gene edits can be unique and identifiable using SEQ ID NOs: 1 to 18 and 426 to 505 and share genetic signatures comprising at least 90% of the genotypic markers of Table 8 to Table 14, thereby enabling the detection and breeding of the CD163 gene edited genomes in any progeny generation. In an aspect, the genetic signatures can share 95% or more of the genotypic markers of Table 8 to Table 14. In another aspect, the genetic signatures share 97% or more of the genotypic markers of Table 8 to Table 14. Also included are genetic signatures that can share 98% or more of the genotypic markers of Table 8 to Table 14. Importantly, the linkage of the CD163 edits to the genomic regions identifiable using the genotypic markers of Table 8 to Table 14 can enable the preparation of any progeny animal having the desired edits and the tracking of the edited regions in any number of progeny generations. In aspects according to the present specification, the heterozygous and homozygous pigs of the present specification can be free of off-site mutations.

The present specification provides for, and includes, an embryo or zygote that can be obtained from an elite line of pigs. In an aspect, the embryo or zygote can be obtained from an elite porcine line selected from the group consisting of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, or Line 65. In an aspect, the embryo or zygote can be a frozen embryo or zygote. In another aspect, the embryo or zygote can be a frozen blastocyst. As provided herein, the embryos can be prepared from in vitro matured oocytes collected from estrus synchronized gilts. The surrounding cumulus cells can be removed from the in vitro matured oocytes and incubated with washed boar spermatozoa and incubated. After incubation, presumptive zygotes can be microinjected with an RNP mixture comprising the CRISPR-Cas endonuclease and guide RNA combinations comprising the first 20 nucleotides of SEQ ID NOs: 22 to 271 or 347 to 425 listed in Table 3. Injected embryos can be transferred to a surrogate female at the 1 to 4 cell stage. In an aspect, the RNP mixture can further include repair templates listed in Table 6 (SEQ ID NOs: 1 to 13). In an aspect, injected zygotes can be surgically implanted into the oviducts of estrus synchronized, un-mated surrogate females by a mid-line laparotomy under general anesthesia (each surrogate receives 40-60 injected embryos).

The present specification provides for, and includes, gene edited pigs of selected elite lines that can be CD163^{-/-}. In an aspect, the line can be PICTTM Line 15, PICTTM Line 17, PICTTM Line 27, PICTTM Line 65, PICTTM Line 14, PICTTM Line 62, PIC337, PIC800, PIC280, PIC327, PIC408, PICTTM 399, PIC410, PIC415, PIC359, PIC380, PIC837, PIC260, PIC265, PIC210, PICTTM Line 2, PICTTM Line 3, PICTTM Line 4, PICTTM Line 5, PICTTM Line 18, PICTTM Line 19, PICTTM Line 92, PIC95, PICTTM CAMBOROUGH®, PIC1070, PICTTM

CAMBOROUGH® 40, PICTTM CAMBOROUGH® 22, PIC1050, PICTTM CAMBOROUGH® 29, PICTTM CAMBOROUGH® 48, or PICTTM CAMBOROUGH® x54. In an aspect, the line can be selected from the group consisting of PICTTM Line 2, Line 3, Line 15, Line 19, Line 27, Line 62, Line 65, and progeny thereof comprising the edited CD163 genes described herein. Gene edited lines 2, 3, 15, 19, 27, 62, and 65 comprise CD163 genomic regions as provided above at Table 8 to Table 14, and can be readily distinguished from each other, from unimproved lines, and from other elite lines. Similarly, progeny of lines 2, 3, 15, 19, 27, 62, and 65 comprising the CD163^{-/-} genomic regions as provided in Table 8 to Table 14 can be identified. Accordingly, the present specification provides for progeny pigs that can have a CD163^{-/-} genomic region.

The present specification provides for and includes hybrid porcine lines comprising an edited CD163 gene of the present teachings. In some aspects, the hybrid porcine line can be produced by crossing an edited PICTTM line with at least one other edited PICTTM line. In some aspects, the porcine line can be produced by serial crosses to introduce germplasm from three or more porcine lines. In an aspect, the line can be PICTTM Line 15, PICTTM Line 17, PICTTM Line 27, PICTTM Line 65, PICTTM Line 14, PICTTM Line 62, PIC337, PIC800, PIC280, PIC327, PIC408, PICTTM 399, PIC410, PIC415, PIC359, PIC380, PIC837, PIC260, PIC265, PIC210, PICTTM Line 2, PICTTM Line 3, PICTTM Line 4, PICTTM Line 5, PICTTM Line 18, PICTTM Line 19, PICTTM Line 92, PIC95, PICTTM CAMBOROUGH®, PIC1070, PICTTM CAMBOROUGH® 40, PICTTM CAMBOROUGH® 22, PIC1050, PICTTM CAMBOROUGH® 29, PICTTM CAMBOROUGH® 48, or PICTTM CAMBOROUGH® x54.

In various aspects, PICTTM Line 65 is sold under the trade name PIC337. In various aspects, PICTTM line 62 is sold under the tradename PIC408. In various aspects, hybrid pigs made by crossing PICTTM lines 15 and 17 are sold under the tradenames PIC800 or PIC280. In various aspects, PICTTM Line 27 is sold under the tradename PIC327. In various aspects, hybrids created from crossing PICTTM Line 65 and PICTTM Line 62 is sold under the tradenames PIC399, PIC410 or PIC415. In various aspects, hybrids created from crossing PICTTM Line 65 and Pic Line 27 are sold under the tradename PIC359. In various aspects, hybrids prepared from crossing PICTTM Line 800 pigs (which is a hybrid of PICTTM Line 15 and PICTTM Line 17) to PICTTM Line 65 pigs are sold under the tradenames PIC380 or PIC837. In various aspects, PICTTM Line 14 is sold under the trade name PIC260. In various aspects, hybrids created from crossing PICTTM Line 14 and PICTTM Line 65 are sold under the tradename PIC265. In various aspects, hybrids created by crossing PICTTM Line 2 and PICTTM Line 3 are sold under the tradenames PIC210, PICTTM CAMBOROUGH®, and PIC1050. In various aspects, hybrids of PICTTM Line 3 and PICTTM Line 92 are sold under the tradename PIC95. In various configurations, hybrids made from crossing PICTTM Line 19 and PICTTM Line 3 are sold under the tradename PIC1070. In various aspects, hybrids created by crossing PICTTM Line 18 and PICTTM Line 3 are sold under the tradename PICTTM CAMBOROUGH® 40. In various aspects, hybrids created from crossing PICTTM Line 19 and PIC1050 (which is itself a hybrid of PICTTM lines 2 and 3) are sold under the tradename PICTTM CAMBOROUGH® 22. In various aspects, hybrids created from crossing PICTTM Line 2 and PIC1070 (which is itself a hybrid of PICTTM lines 19 and 3) are sold under the tradename PICTTM CAMBOROUGH® 29. In various aspects, hybrids created from crossing PICTTM Line 18 and PIC1050 (which is itself a hybrid of PICTTM lines 2 and 3) are sold under the

tradename PICTTM CAMBOROUGH® 48. In various aspects, hybrids created from crossing PICTTM Line 4 and PICTTM Line 5 are sold under the tradename PICTTM CAMBOROUGH® x54. The present teachings provide for and include pigs of any of the forgoing lines or hybrids comprising a CD163 edit of the present teachings.

In various aspects, the present teachings provide for an include pigs comprising an edited CD163 gene comprising an edited sequence set forth in SEQ ID NOs: 453, 489, or 505 wherein the pig can be a pig of PICTTM Line 15, PICTTM Line 17, PICTTM Line 27, PICTTM Line 65, PICTTM Line 14, PICTTM Line 62, PICTTM Line 2, PICTTM Line 3, PICTTM Line 4, PICTTM Line 5, PICTTM Line 18, PICTTM Line 19, PICTTM Line 92, or a hybrid of two or more of these lines. In various aspects, the hybrid pig can be a cross of PICTTM Line 15 and PICTTM Line 17, PICTTM Line 65 and PICTTM Line 62, PICTTM Line 65 and PICTTM Line 27, a serial hybrid of PICTTM Line 15 and PICTTM Line 17, wherein the hybrid offspring is then crossed to PICTTM Line 65, PICTTM Line 14 and PICTTM Line 65, PICTTM Line 2 and PICTTM Line 3, PICTTM Line 3 and PICTTM Line 92, PICTTM Line 19 and PICTTM Line 3, PICTTM Line 18 and PICTTM Line 3, a serial hybrid between a hybrid pig of PICTTM Line 2 and PICTTM Line 3 and a pig of PICTTM Line 19, a serial hybrid between a hybrid pig of PICTTM Line 19 and PICTTM Line 3 and a pig of PICTTM Line 2, a serial hybrid between a hybrid of PICTTM Line 2 and PICTTM Line 3 crossed to a pig of PICTTM Line 18, or a hybrid of PICTTM Line 4 and PICTTM Line 5.

In various aspects, the hybrid line comprising an edited CD163 gene can be produced by crossing two or more of lines 2, 3, 15, 19, 27, 62, or 65. In some aspects, the hybrid line comprising an edited CD163 gene can be a CAMBOROUGH® line. CAMBOROUGH® pigs are hybrids that can be prepared by a cross between Line 2 and Line 3. In various aspects, the hybrid line can comprise a PICTTM 837 hybrid line comprising an edited CD163 gene. PICTTM 837 pigs are hybrids that can be prepared by crossing PICTTM Line 800 pigs to PICTTM Line 65 pigs.

The present specification provides for, and includes, hybrid CD163^{-/-} progeny lines comprising one CD163^{-/-} genomic region that can be obtained from lines 2, 3, 15, 19, 27, 62, or 65, and a second CD163^{-/-} genomic region that can be obtained from a different line selected from lines 2, 3, 15, 19, 27, 62, or 65. In an aspect, a hybrid CD163^{-/-} progeny line can comprise a CD163^{-/-} allele in a genomic region according to Table 8 and a CD163^{-/-} allele in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise a CD163^{-/-} allele in a genomic region according to Table 8 and a CD163^{-/-} allele in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise a CD163^{-/-} allele in a genomic region according to Table 9 and a CD163^{-/-} allele in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise a CD163^{-/-} allele in a genomic region according to Table 9 and a CD163^{-/-} allele in a genomic region according to Table 14.

The present specification provides for, and includes, hybrid animals that can comprise the CD163 gene edits characterized by SEQ ID NOs: 1 to 18 and 426 to 505. In an aspect, the hybrid animals comprising the CD163 gene edits can be characterized by SEQ ID NO: 2. In an aspect, the hybrid animals comprising the CD163 gene edits can be characterized by SEQ ID NO: 426-458. In an aspect, the hybrid animals comprising the CD163 gene edits can be characterized by SEQ ID NO: 453. In an aspect, the hybrid animals comprising the CD163 gene edits can be character-

ized by SEQ ID NO: 459-504. In an aspect, the hybrid animals comprising the CD163 gene edits can be characterized by SEQ ID NO: 489. In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animals of the CAMBOROUGH® line. CAMBOROUGH® pigs are hybrids that can be prepared by a cross between Line 2 and Line 3.

In various aspects, the hybrid pig can comprise SEQ ID NO: 2 in a genomic region according to Table 8 and SEQ ID NO: 2 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 453 in a genomic region according to Table 8 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 489 in a genomic region according to Table 8 and SEQ ID NO: 489 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 2 in a genomic region according to Table 8 and SEQ ID NO: 2 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 453 in a genomic region according to Table 8 and SEQ ID NO: 453 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 489 in a genomic region according to Table 8 and SEQ ID NO: 489 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 2 in a genomic region according to Table 9 and SEQ ID NO: 2 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 453 in a genomic region according to Table 9 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 489 in a genomic region according to Table 9 and SEQ ID NO: 489 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 2 in a genomic region according to Table 8 and SEQ ID NO: 2 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 453 in a genomic region according to Table 8 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 489 in a genomic region according to Table 8 and SEQ ID NO: 489 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 2 in a genomic region according to Table 9 and SEQ ID NO: 2 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 453 in a genomic region according to Table 9 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 489 in a genomic region according to Table 9 and SEQ ID NO: 489 in a genomic region according to Table 10. In aspects according to the present specification, the heterozygous and homozygous pigs of the present specification can be free of off-site mutations.

In an aspect, the hybrid animal can be a CD163^{-/-} hybrid animals of the CAMBOROUGH® line. CAMBOROUGH® pigs are hybrids that can be prepared by a cross between Line 2 and Line 3. In an aspect, the hybrid pig can comprise

SEQ ID NO: 426 to 505 in a genomic region according to Table 8 and SEQ ID NO: 426-505 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 453 in a genomic region according to Table 8 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, the hybrid pig can comprise SEQ ID NO: 489 in a genomic region according to Table 8 and SEQ ID NO: 489 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 426 to 505 in a genomic region according to Table 8 and SEQ ID NO: 426 to 505 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 453 in a genomic region according to Table 8 and SEQ ID NO: 489 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 489 in a genomic region according to Table 8 and SEQ ID NO: 453 in a genomic region according to Table 14. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 426 to 505 in a genomic region according to Table 9 and SEQ ID NO: 426 to 505 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 453 in a genomic region according to Table 9 and SEQ ID NO: 453 in a genomic region according to Table 10. In an aspect, a hybrid CD163^{-/-} progeny line can comprise SEQ ID NO: 489 in a genomic region according to Table 9 and SEQ ID NO: 489 in a genomic region according to Table 10.

In various aspects, an edited pig of PICTTM line 2, PICTTM line 3, PICTTM line 15, PICTTM line 19, PICTTM line 27, PICTTM line 62, or PICTTM line 65 can have a CD163 gene comprising a predicted amino acid sequence as set forth in SEQ ID NO: 518. In various aspects, the edited CD163 gene can have a 123 bp deletion as set forth in SEQ ID NO: 505. In various configurations, this deletion can have an Exon 7 Amino Acid sequence of AHRKPRLV - - - TVVSLGGAHFGEGSGQI-WAEFQCEGHESHLSCPVA PRPDGTCSHSRDVGVC CS (SEQ ID NO: 518). In various aspects, a pig having an edited CD163 gene comprising an amino acid sequence set forth in SEQ ID NO: 518 can be a hybrid offspring between two PICTTM lines. In various aspects, the pig can have an edited CD163 Exon 7 sequence comprising a nucleotide sequence set forth in SEQ ID NO: 505. In various aspects, the pig can be an offspring of a cross between PICTTM line 2 and PICTTM line 3. In aspects according to the present specification, the heterozygous and homozygous pigs of the present specification can be free of off-site mutations.

Importantly, the CD163 edited pigs and cells of the present specification retain their desirable commercial phenotypes. The edited pigs can exhibit a healthy phenotype with no noticeable acute deleterious effects from the edit. In an aspect, the pigs of lines 2, 3, 15, 19, 27, 62, and 65, can retain the commercially desirable phenotypes as provided in Table 15.

Methods for improving the health of existing herds of livestock can comprise modifying the CD163 gene locus using the methods described above. In an aspect, the method can comprise introducing into a pig cell an endonuclease or a polynucleotide encoding said endonuclease, and a guide polynucleotide comprising a sequence selected from the group consisting of the first 20 nucleotides of each of SEQ ID NOs: 22 to 271 or the first 20 nucleotides of each of 347 to 425, incubating the cell under conditions that permit the endonuclease to act upon the DNA at, or near, the target sequence and thereby induce recombination, homology-directed repair, or non-homologous end joining at or near the target site, identifying at least one cell having a modification

at said target sequence, and producing an animal from an animal cell. In aspects of the present specification, the endonuclease can be an RNP complex of a guide RNA and a Cas protein from *S. thermophilus*. In another aspect, the endonuclease can be an RNP complex of a guide RNA and a Cas protein from *S. pyogenes*.

In an aspect, the method can further comprise providing to the cell a repair guide comprising a sequence selected from the group consisting of nucleotides 50 to 100 of SEQ ID NOs: 1 to 13. In another aspect, a repair guide can comprise a sequence selected from the group consisting of nucleotides 50 to 100 of SEQ ID NOs: 1 to 13 and can further comprise 85% homology to nucleotides 1 to 49 and 101 to 150 of SEQ ID NOs: 1 to 13. In an aspect, the repair guide can comprise a sequence selected from the group consisting of SEQ ID NOs: 1 to 13. As provided herein, the method for improving the health of existing herds provides for the maintenance of desirable commercial phenotypes as discussed above. In an aspect, the desirable commercial phenotypes can be at least 90% of the phenotypes observed in herds of non-edited pigs having a similar genetic background.

The CD163 gene edited locus can be introduced into the herd by conventional breeding or by methods incorporating artificial insemination. To prepare homozygous animals, crosses between parents having at least one gene edited CD163 locus comprising a sequence of SEQ ID NOs: 1 to 18 or 426 to 505 can be performed and homozygous progeny (present at a 1:4 Mendelian ratio) can be selected. As provided herein, a combination of parents and one gene edited CD163 locus can be suitable for improving the herd. Further breeding of animals having the CD163 gene edited loci can improve the health of the herd until all the animals of the herd comprise a CD163 gene edited locus as described. Notably, the health of the herd can improve significantly well before the herd has been bred fully to a CD163 gene edited herd. Specifically, and as discussed above, pig fetuses in a PRRSV resistant gene edited CD163 sow are themselves protected from PRRSV. Further, one of ordinary skill in the art would know that, as the numbers of homozygous CD163 edited animals increase, suitable pig vectors for PRRSV transmission decrease (e.g., herd immunity develops). Accordingly, the present methods can provide for improving a herd by introducing herd immunity.

One useful method of detecting the desired edit is to use real-time PCR. PCR primers flanking the region of interest and a probe that specifically anneals to the region of interest. The probe is labelled with both a fluorophore and a quencher. In the PCR reaction, the primers and probe hybridize in a sequence-dependent manner to the complementary DNA strand of the region of interest. Because the probe is intact, the fluorophore and quencher are in close proximity and the quencher absorbs fluorescence emitted by the fluorophore. The polymerase extends from the primers and begins DNA synthesis. When the polymerase reaches the probe, the exonuclease activity of the polymerase cleaves the hybridized probe. As a result of cleavage, the fluorophore is separated from the quencher and fluoresces. This fluorescence is detected by the real time instrument. These steps are repeated for each PCR cycle and allow detection of specific products.

In the instant application, three separate sets of primers and probes were designed. The first set of primers (SEQ ID NO: 556 and 557) flanked the unedited genomic sequence comprising SEQ ID NO: 249 and a probe (SEQ ID NO: 558) which binds to the unedited genomic DNA in between the primers. The second set of primers (SEQ ID NO: 559 and

560) flanked the unedited genomic sequence comprising SEQ ID NO: 256, and a probe (SEQ ID NO: 561) binds the unedited between the primers. The final set of primers (SEQ ID NO: 562 and 563) flanked the desired Exon 7 deletion edit created by excision of the sequence between the cut sites of SEQ ID NO: 249 and SEQ ID NO: 256. A probe (SEQ ID NO: 564) was designed to bind the desired edit in between these primers. A commercial real-time PCR kit was then used to probe various animals for the desired edit. A variety of commercial real-time PCR kits exist including, such as, but without limitation, PRIMETIME® from IDT, TAQMAN® (Roche Molecular Systems, Inc, Pleasanton, CA) from Applied Biosystems, and various kits from Qiagen and Bio-Rad. Skilled persons will recognize that any such kit can be used with the primers and methods of the present teachings to achieve like results.

EXAMPLES

Example 1

This Example illustrates target site selection of a porcine CD163 gene knockout in pig cells.

A *Streptococcus* Cas9/gRNA RNA-directed DNA endonuclease was used to generate DNA sequence insertions, deletions, and combinations thereof, in the porcine CD163 gene (Sscrofa11.1, GenBank accession: GCA000003025.6), whereby guide RNAs and protein combinations were delivered singly or as pairs, such that sequence changes in the CD163 gene reduce or abolish the function, stability, or expression of the CD163 messenger RNA or protein. The sequence of the CD163 gene from 120 nucleotides upstream (chr5:63300192) of the translational start site to 59 nucleotides downstream of CD163 exon 7 (chr5:63323390), was screened for the guide RNA binding sites having either an adjacent nGG or nGGnG PAM sequence required for cutting by the Cas9 proteins derived from *Streptococcus pyogenes* or *Streptococcus thermophilus* CRISPR3 (*S. thermophilus* CR3), respectively, the sequences of which are in the sequence listing filed herewith, as listed in Table 2. The DNA sequences for the target sites for editing, the locations of the target sites on the CD163 gene, and the editing activity (measured as described in Example 3) are provided in Table 3. Guide RNA molecules had the same sequence as the target, with corresponding RNA nucleotides, without the PAM sequence (nGG for *S. pyogenes* and nGGnG for *S. thermophilus*). Guide polynucleotide molecules could also consist of DNA bases or mixtures of DNA and RNA bases. The target site sequences are listed in the sequence listing filed herewith as SEQ ID NOs: 22 to 271 and 347 to 425. The target site sequences shown are conserved across pig germplasm and screened by DNA sequencing.

Example 2

This example illustrates nucleofection for the delivery of guide RNA/Cas9 endonuclease to porcine fetal fibroblasts.

To test the DNA cutting activity in living cells to produce an edited porcine CD163 allele, the CRISPR-Cas endonuclease and guide RNA targeting the sequences listed in Table 3 were nucleofected into porcine fetal fibroblast cells. Porcine fetal fibroblast (PFF) cells lines were prepared from 28-35 day-old fetuses and 3.2 µg of Cas9 protein (*S. pyogenes* or *S. thermophilus*) and 2.2 µg of in vitro transcribed single guide RNA were combined in water to a total volume of 2.23 µl, then were nucleofected into PFF cells using a Lonza electroporator. In preparation for nucleofection, PFF

cells were harvested using TrypLE express (recombinant Trypsin), upon which the culture medium was removed from cells, washed 1× with HBSS or DPBS, and incubated for 3-5 minutes at 38.5° C. in the presence of TrypLE. Cells were then harvested with complete medium. Cells were pelleted via centrifugation (300×g for 5 minutes at room temperature), supernatant was discarded, then the cells were resuspended in 10 mL PBS to obtain single cell suspension counting cells using trypan blue staining.

The appropriate amount of cells was pelleted by centrifugation (300×g for 5 minutes at room temperature), the supernatant was discarded, and the cells were resuspended in nucleofection buffer P3 at a final concentration of 7.5×10⁶ cells/ml. 20 µl of the cell suspension were added to each well of a nucleofection plate containing the RNP mixture using a multichannel pipette, then mixed gently to resuspend the cells. The RNP/cell mixture was transferred in the nucleofection plate, nucleofected with program CM138 (supplied by the manufacturer). 80 µl of warm Embryonic Fibroblast Medium, EFM, (Dulbecco's Modified Eagle's Medium (DMEM) containing 2.77 mM glucose, 1.99 mM L-glutamine, and 0.5 mM sodium pyruvate, supplemented with 100 µM 2-Mercaptoethanol, 1× Eagle's minimum essential medium non-essential amino acids (MEM NEAA), 100 µg/mL Penicillin-Streptomycin, and 12% Fetal Bovine Serum were added to each well after nucleofection. The suspensions were mixed gently by pipetting, then 100 µl were transferred to a 12 well plate containing 900 µl of EFM pre-incubated at 38.5° C. The plate was then incubated at 38.5° C., 5% CO₂ for 48 hours. Forty-eight hours post nucleofection, genomic DNA was prepared from transfected and control PFF cells, 15 µl of QUICKEXTRACT™ DNA Extraction Solution (Lucigen, Madison, WI) were added to pelleted cells then lysed by incubating for 10 mins at 37° C., for 8 mins at 65° C., for 5 mins at 95° C., then lysate was held at 4° C. until used for DNA sequencing.

Example 3

This example illustrates the editing frequency of guide RNA/Cas9 combinations directed against porcine CD163.

Nucleotide sequence changes were introduced into the porcine CD163 gene by delivering Cas9 protein complexed with guide RNAs to fetal fibroblast cells as described in Example 2.

To evaluate DNA double strand cleavage at a porcine CD163 genomic target site mediated by the guide RNA/Cas endonuclease system, a region of approximately 250 bp genomic DNA surrounding the target site was amplified by PCR and the PCR product was then examined by amplicon deep sequencing for the presence of edits. After transfection in triplicate, PFF genomic DNA was extracted as described in Example 2. The region surrounding the intended target site was PCR amplified with NEB Q5 Polymerase, adding sequences necessary for amplicon-specific barcodes and ILLUMINA® (ILLUMINA®, San Diego, CA) sequencing using tailed primers through two rounds of PCR. The resulting PCR amplifications were deep sequenced on an ILLUMINA® MISEQ® Personal Sequencer (ILLUMINA®, San Diego, CA). The resulting reads were examined for the presence of edits at the expected site of cleavage by comparison to control experiments where the Cas9 protein and guide RNA were omitted from the transfection or by comparison to the reference genome. To calculate the frequency of NHEJ edits for a target site, Cas9 protein, guide RNA combination, the total number of edited reads (amplicon sequences containing insertions or deletions when com-

pared to the DNA sequences from control treatments or reference genome) were divided by total read number (wild-type plus edited reads) of an appropriate length containing a perfect match to the barcode and forward primer. Total read counts averaged approximately 7000 per sample and NHEJ activity is expressed as the average (n=3) edited fraction in Table 3. As shown in Table 3, from 0 to 58.2% of the reads contained edits and the average editing frequency across all combinations was approximately 16%. This example demonstrates that, in fetal fibroblast cells, the porcine CD163 gene nucleotide sequence was edited through the stimulation of double-strand breaks mediated by transfecting Cas9 protein from either *S. pyogenes* or *S. thermophilus* complexed with various guide RNAs.

Example 4

This example illustrates generation of an in-frame stop codon using DNA repair template.

DNA repair templates were used to introduce stop codons in the porcine CD163 gene when co-delivered with Cas9 protein and guide RNA. The endonuclease-guide complexes and repair templates disclosed in Table 5 were generated and used to edit CD163 genes in blastocysts. The base deletions were verified by the methods of Example 5.

Example 5

This example illustrates the molecular characterization of edited animal genomes.

A tissue sample was taken from an animal whose genome was been edited according to the examples herein. Tail, ear notch, or blood samples were suitable tissue types. The tissue sample was frozen at -20° C. within 1 hour of sampling to preserve integrity of the DNA in the tissue sample.

DNA was extracted from tissue samples after proteinase K digestion in lysis buffer. Characterization was performed on two different sequence platforms, short sequence reads using the ILLUMINA® platform and long sequence reads on an Oxford NANOPORE™ platform (Oxford NANOPORE™ Technologies, Oxford, UK).

For short sequence reads, two-step PCR was used to amplify and sequence the region of interest. The first step was a locus-specific PCR which amplified the locus of interest from the DNA sample using a combined locus-specific primer with a vendor-specific primer. The second step attached the sequencing index and adaptor sequences to the amplicon from the first step so that sequencing could occur.

The locus-specific primers for the first step PCR were chosen so that they amplified a region <300 bp such that ILLUMINA® paired-end sequencing reads could span the amplified fragment. Multiple amplicons were preferred to provide redundancy should deletions or naturally occurring point mutations prevent primers from correctly binding. Sequence data for the amplicon was generated using an ILLUMINA® sequencing platform (MISEQ®, ILLUMINA®, San Diego, CA). Sequence reads are analyzed to characterize the outcome of the editing process.

For long sequence reads, two-step PCR was used to amplify and sequence the region of interest. The first step is a locus-specific PCR which amplified the locus of interest from the DNA sample using a combined locus-specific primer with a vendor-specific adapter. The second step PCR attached the sequencing index to the amplicon from the first-step PCR so that the DNA was ready for preparing a

sequencing library. The step 2 PCR products underwent a set of chemical reactions from a vendor kit to polish the ends of the DNA and ligate on the adapter containing the motor protein to allow access to the pores for DNA strand-based sequencing.

The locus specific primers for the first step PCR range were designed to amplify different regions of the CD163 gene and amplified regions differed in length. Normalized DNA is then mixed with vendor supplied loading buffer and is loaded onto the NANOPORE™ flowcell.

Long sequence reads, while having lower per base accuracy than short reads, are very useful for observing the long range context of the sequence around the target site.

Example 6

This example illustrates methods of making pigs having edited CD163 genes conferring PRRSV resistance.

Porcine oocytes were isolated, fertilized, and then the resulting zygotes are edited to generate gene edited pigs.

CD163 RNP complexes were microinjected into the cytoplasm of in vivo or in vitro fertilized porcine one-cell zygotes. These zygotes were then incubated to generate edited multicellular embryos and transferred to surrogate gilts via standard methods to birth gene edited pigs. To prepare embryo donors and surrogates, pubertal gilts from PIC™ Line 2, Line 3, Line 15, and Line 65 were subjected to estrus synchronization by treatment with 0.22% altrenogest solution (20-36 mg/animal) for 14 days. Follicular growth was induced by the administration of PMSG 36 hours following the last dose of Matrix, and ovulation was induced by the administration of hCG 82 hours after PMSG administration. To generate in vivo fertilized zygotes, females in standing heat were then artificially inseminated (AI) with boar semen from the corresponding PIC™ line. In vivo derived zygotes were recovered surgically 12-24 hours after AI by retrograde flushing the oviduct with sterile TL-HEPES medium supplemented with 0.3% BSA (w/v). Fertilized zygotes were subjected to a single 2-50 picoliter (pl) cytoplasmic injection of Cas9 protein and guide RNA complex (25-50 ng/μl and 12.5-35 ng/μl) targeting CD163 and cultured in PZM5 medium (Yoshioka, K., et al., Biol. Reprod., 2002, 60: 112-119; Suzuki, C., et al., Reprod. Fert. Dev., 2006 18, 789-795; Yoshioka, K., J. Reprod. Dev. 2008, 54, 208-213). Injected zygotes were surgically implanted into the oviducts of estrus synchronized, unmated surrogate females by a mid-line laparotomy under general anesthesia (each surrogate received 20-60 injected embryos).

In vitro fertilized embryos for gene editing were derived from non-fertilized PIC™ oocytes. Immature oocytes from estrus synchronized PIC™ gilts were collected from medium size (3-6 mm) follicles. Oocytes with evenly dark cytoplasm and intact surrounding cumulus cells were then selected for maturation. Cumulus oocyte complexes were placed in a well containing 500 μl of maturation medium, TCM-199 (Invitrogen) with 3.05 mM glucose, 0.91 mM sodium pyruvate, 0.57 mM cysteine, 10 ng/ml EGF, 0.5 μg/ml luteinizing hormone (LH), 0.5 μg/ml FSH, 10 ng/ml gentamicin (Sigma), and 10% follicular fluid for 42-44 h at 38.5° C. and 5% CO₂, in humidified air. At the end of the maturation, the surrounding cumulus cells were removed from the oocytes by vortexing for 3 min in the presence of 0.1% hyaluronidase. Then, in vitro matured oocytes were placed in 100 μl droplets of IVF medium (modified Tris-buffered medium containing 113.1 mM NaCl, 3 mM KCl, 7.5 mM CaCl₂, 11 mM glucose, 20 mM Tris, 2 mM

caffeine, 5 mM sodium pyruvate, and 2 mg/ml bovine serum albumin (BSA)) in groups of 25-30 oocytes and were fertilized according to established protocol (Abeydeera, *Biol. Reprod.*, 57:729-734, 1997) using fresh extended boar semen. One ml of extended semen was mixed with Dulbecco's Phosphate Buffered Saline (DPBS) containing 1 mg/ml BSA to a final volume of 10 ml and centrifuged at 1000×g, 25° C. for 4 minutes, and spermatozoa were washed in DPBS three times. After the final wash, spermatozoa were re-suspended in mTBM medium and added to oocytes at a final concentration of 1×10⁵ spermatozoa/ml, and co-incubated for 4-5 h at 38.5° C. and 5% CO₂. Presumptive zygotes were microinjected 5 hours post IVF and transferred to a surrogate female after 18-42 hours (1-4 cell stage). Each surrogate receives 20-60 injected embryos. Pregnancies were confirmed by lack of return to estrus (21 days) and ultrasound at 28 days post embryo transfer.

To establish the frequency of Cas9-guide RNA targeted gene editing in porcine embryos, uninjected control zygotes and injected surplus zygotes generated by in vitro fertilization were cultivated in PZM3 or PZM5 medium at 38.5° C. for 5-7 days. Blastocysts were harvested at day 7 post cultivation and the genomic DNA isolated for next generation sequencing.

Example 7

This example illustrates the generation and characterization of gene edited pigs.

Animals from PICTM lines 2, 3, 15, and 65 were edited by the methods described in Example 6. Successful edits were confirmed using the methods of Example 5. Fibroblast cell lines were grown from collagenase treated ear notch samples extracted from the edited animals and deposited with the American Type Culture Collection (ATCC®). The ATCC® has an address of 10801 University Boulevard, Manassas, VA 20110-2209. A representative sample of CD163 edited PICTM Line 2 was deposited with the ATCC on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125814. A representative sample of CD163 edited PICTM Line 3 was deposited with the ATCC on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125815. A representative sample of CD163 edited PICTM Line 15 was deposited with the ATCC on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125816. A representative sample of CD163 edited PICTM Line 65 was deposited with the ATCC® on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125813. Each deposit was made according to the Budapest Treaty. Representative animals from each line were confirmed to have heterozygous edits as specified in Table 16.

TABLE 16

Verified Gene Edited Animals			
Line number	CD163 Allele 1	CD163 Allele 2	ATCC Deposit No.
2	Wild type sequence	Deleted nucleotides chr5: 63301999-63302005	PTA-125814
3	Wild type sequence	Deleted nucleotides chr5: 63301999-63302005	PTA-125815
15	Wild type sequence	Deleted nucleotides chr5: 63301999-63302005	PTA-125816
65	Wild type sequence	Deleted nucleotides chr5: 63301999-63302005	PTA-125813

Each of the animals in Table 16 presented a healthy phenotype with deleted nucleotides 63301999-63302005 (in exon 2) from chromosome 5.

Additional cell lines were grown from collagenase treated ear notch samples from unedited animals from PICTM lines 19, 27, and 62, were deposited with the ATCC. A representative sample of PICTM Line 19 was deposited with the ATCC® on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125811. A representative sample of PICTM Line 27 was deposited with the ATCC on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125907. A representative sample of PICTM Line 62 was deposited with the ATCC on Apr. 3, 2019 and assigned ATCC® Patent Deposit Number PTA-125812. Each deposit was made according to the Budapest Treaty. Using conventional cloning methods, animals are generated from the cell lines deposited as PTA-125811, PTA-125812, and PTA-125907 and edited using the methods of Example 6 in order to generate edited lines.

Single nucleotide polymorphisms (SNPs) in the vicinity of the CD163 gene were analyzed for each of the deposited PICTM lines (2, 3, 15, 19, 27, 62, and 65) and SNP profiles of each of the lines that are capable of distinguishing each line are selected. The starting dataset for defining the line signatures was a collection of 330 whole genome sequenced animals from the 7 deposited PICTM lines. A 6 Mb region of Chromosome 5 centered on the CD163 gene was extracted for the signatures, and variation within and between the lines was examined for each nucleotide in this region in order to identify a relatively small number of SNPs that together formed a signature for the line.

To be a candidate for inclusion in the signature of a given line, the following criteria were imposed on each chromosomal position: sequence coverage had to exist for 90% of animals in the line; the above had to be true for at least 5 of the 7 lines; and all animals with data in the target line must have had the same homozygous genotype. For each of the other lines, a genotype frequency for this genotype was calculated across all the sequenced animals for that line. A cutoff was imposed of at least 30% of the difference between the highest and lowest per-line genotype frequency of the other 6 lines (i.e., there must be a spread of genotype frequencies within the 6 lines).

A combination of metrics for discriminating power and even distribution were used to select a subset of genotypes that could define each line. Table 8 to Table 14 provide the positions on chromosome 5 for the SNPs for which the homozygous allele was fixed in each porcine line. These sets of homozygous alleles distinguish each porcine line from the other lines. The genotype listed indicates which allele is homozygous at each position, as indicated in Table 8 to Table 14.

Example 8

This example compares differing levels of PRRSV resistance of immune cells isolated from wild type and gene edited pigs.

CD163 surface expression analysis was conducted on Monocyte Derived Macrophages (MoMØs) recovered from edited and wild type animals, each with and without edits to the CD163 gene according to the methods described in the previous examples. The edits tested are presented in Table 17. Four edits comprise deletions of various sizes as shown, and the fifth comprises a 2 base pair insertion; all edits in Table 15 are in exon 2. All deletions or insertions result in a truncated CD163 polypeptide. CD163 expression was

assessed by immunofluorescence labelling and FACS analysis using clone 2A10/11, a mouse anti-pig CD163 monoclonal antibody. CD163 edited cells, MoMØs, lacked a functional epitope on the surface of the cell as evident from cell surface expression analysis.

TABLE 17

CD163 gene edits tested for protein expression					
Region Chr5 Location of guideRNAs binding site	63301997- 02016	63301997- 02016	63301997- 02016	63301997- 02016	63301997- 02016
SEQ ID NO Guide RNA spacer (DNA)	14 42	15 42	16 42	17 42	18 42
SEQ ID NO Guide RNA spacer (RNA)	288	288	288	288	288
SEQ ID NO Translation of Deletion	339	340	341	342	343
Base deletion or insertion coordinates	chr5: 6330201 2-63302013	chr5: 6330200 9-63302041	chr5: 6330201 2-63302013	chr5: 6330200 5-63302020	chr5: 6330201 1-63302020
Deleted or inserted bases (#)	TC (2 bp deletion)	SEQ ID NO: 344; (33 bp deletion)	TC (2 bp insertion)	SEQ ID NO: 345; (16 bp deletion)	SEQ ID NO: 346; (10 bp deletion)

to transfection. Editing frequency for guide pairs was determined as described in Example 3. The frequencies of NHEJ-mediated repairs whereby the endonuclease cut sites are brought together resulting in the deletion of CD163 exon 7 are shown in Table 18.

To test the two base pair deletion edit to CD163 in homozygous edited cells for PRRSV viral infection, MoMØs were infected with PRRSV type 1 and type 2. Twenty-four (24) hours post infectivity, in a microscope field of view, cells were counted to determine the number of cells which contain replicating PRRSV. CD163 homozygous edited MoMØ cells were found to be not permissive to both PRRSV types 1 and 2.

Example 9

This example demonstrates the use of two guides for removal of exon 7 from *S. scrofa* CD163 in porcine fibroblasts.

In order to remove CD163 exon 7 DNA sequences which encode SRCR5 of the mature CD163 polypeptide, DNA sequences located in the intronic regions 450 bp upstream and 59 bp downstream of CD163 exon 7 were examined for *Streptococcus pyogenes* (NGG) and *Streptococcus thermophilus* (NGGNG) Cas9 protein and guideRNA recognition sites. 48 sites were identified within the 450 bp of intron 6 and 10 sites were identified within a 59 bp region of intron 7. The ability of these 58 sites to bind to Cas9 and guideRNAs to direct gene editing was first tested using single guides in porcine fetal fibroblasts. A subset of these single guideRNA-Cas9 proteins—the guideRNAs which generated a high frequency of edits across the spacer recognition site—were further tested as guide pairs for their ability to remove CD163 exon 7. Guides were introduced to porcine fetal fibroblasts by nucleofection as described in Example 2: each guide was prepared as an RNP and then the two sets of complexes were combined in a total volume of 2.23 µl prior

TABLE 18

End-to-end repair frequencies using paired guideRNAs for CD163 exon 7 deletion in porcine fetal fibroblasts					
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Desired repair outcome (%)
229	63322816	256	63323377	561	23.3
230	63322814	256	63323377	562	24.7
231	63322826	256	63323377	551	15.3
237	63322861	256	63323377	516	20.7
241	63322891	256	63323377	486	20.0
229	63322816	258	63323378	562	5.3
230	63322814	258	63323378	563	7.0
231	63322826	258	63323378	552	4.3
237	63322861	258	63323378	517	15.7
241	63322891	258	63323378	487	5.7
229	63322816	261	63323373	558	15.0
230	63322814	261	63323373	559	18.0
231	63322826	261	63323373	548	0.0
237	63322861	261	63323373	513	12.0
241	63322891	261	63323373	483	14.0
219	63322697	256	63323377	680	28.0
221	63322709	256	63323377	668	33.7
224	63322747	256	63323377	630	27.0
227	63322786	256	63323377	591	22.7
219	63322697	258	63323378	681	7.7
221	63322709	258	63323378	669	13.0
224	63322747	258	63323378	631	7.3
227	63322786	258	63323378	592	6.3
219	63322697	261	63323373	677	14.5
221	63322709	261	63323373	665	20.7
224	63322747	261	63323373	627	14.7
227	63322786	261	63323373	588	11.3
249	63322963	256	63323377	414	38.0
250	63322973	256	63323377	404	36.7
249	63322963	258	63323378	415	20.3
250	63322973	258	63323378	405	14.7
249	63322963	261	63323373	411	17.7
250	63322973	261	63323373	401	11.7

79

The excision guides had a wide range of editing frequencies for the desired edit.

Example 10

This example illustrates the excision of Exon 7 in porcine blastocysts using dual guide RNAs.

A subset of guides screened in porcine fetal fibroblasts were additionally tested for their ability to remove CD163 exon 7 in porcine blastocysts. The subset of guides to be tested in porcine embryos was chosen based on a combination of their efficacy in generating exon 7 deletions in porcine fibroblasts and low number of off-target edits for each guide in the pair (see Detailed Description). Edited porcine embryos were generated as described above. Briefly, oocytes recovered from slaughterhouse ovaries were in vitro fertilized as described in Example 6. The sgRNP solution was injected into the cytoplasm of presumptive zygotes at 4-5 hours post-fertilization by using a single pulse from a FEMTOJET® 4i microinjector (Eppendorf; Hamburg, Germany) with settings at pi=200 hPa, ti=0.25 s, pc=15 hPa. Glass capillary pipettes (Sutter Instrument, Navato, CA, USA) with an outer diameter of 1.2 mm and an inner diameter of 0.94 mm were pulled to a very fine point of <0.5. Microinjection was performed in TL-Hepes (ABT360, LLC) supplemented with 3 mg/ml BSA (Proliant) on the heated stage of an inverted microscope equipped with Narishige (Narishige International USA, Amityville, NY) micromanipulators. Following injections, presumptive zygotes were cultured for 7 days in PZM5 (Cosmo Bio, Co LTD, Tokyo, Japan) in an incubator environment of 5% CO₂, 5% O₂, 90% N₂. Editing frequency of blastocysts was determined as described in Example 3. The frequencies of end-to-end NHEJ repairs resulting in the deletion of CD163 exon 7 are shown in Table 19.

TABLE 19

End-to-end repair frequencies using paired guideRNAs for in CD163 exon 7 deletion in porcine embryos					
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Desired repair outcome (%)
229	63322816	256	63323377	561	38.0
230	63322814	256	63323377	562	21.0
231	63322826	256	63323377	551	24.0
241	63322891	256	63323377	486	29.0
229	63322816	258	63323378	562	7.0
231	63322826	258	63323378	552	12.0
241	63322891	258	63323378	487	20.0
219	63322697	256	63323377	680	35.0
221	63322709	256	63323377	668	36.0
224	63322747	256	63323377	630	24.0
227	63322786	256	63323377	591	0.0
227	63322786	258	63323378	592	0.0
221	63322709	261	63323373	665	15.0
249	63322963	256	63323377	414	44.0
250	63322973	256	63323377	404	14.0
249	63322963	258	63323378	415	7.0
249	63322963	261	63323373	411	53.0

This example demonstrates that a number of guide pairs can be used to delete CD163 exon 7, but the efficiency can vary greatly between guide pairs and cell types.

Example 11

This example demonstrates the use of two guides to introduce a premature stop codon into exon 7 of *S. scrofa* CD163.

80

Guides within exon 7 of CD163 were screened by bioinformatic methods for their ability to generate an in-frame stop codon when the cuts generated by those guides are ligated together during NHEJ. Bioinformatic predictions were tested in porcine fetal fibroblasts as described in Example 2: guides were complexed separately before being combined in a total volume of 2.23 µl prior to transfection. Editing frequency was determined as described in Example 3. The frequencies of end-to-end NHEJ repairs resulting in the introduction of a premature stop codon in exon 7 of CD163 are shown in Table 20.

TABLE 20

End-to-end repair frequencies for introduction of a stop codon in CD163 exon 7 deletion in porcine fetal fibroblasts					
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Desired repair outcome (%)
351	63323023	365	63323103	80	50.5
351	63323023	387	63323235	212	1.0
348	63323027	390	63323236	209	53.5
348	63323027	388	63323236	209	50.0
348	63323027	395	63323284	257	31.0
352	63323035	365	63323103	68	29.0
352	63323035	387	63323235	200	12.5
352	63323035	399	63323283	248	21.0
353	63323038	365	63323103	65	21.0
353	63323038	387	63323235	197	6.0
353	63323038	399	63323283	245	11.0
354	63323039	390	63323236	197	55.5
354	63323039	388	63323236	197	18.0
354	63323039	395	63323284	245	17.0
358	63323056	361	63323077	21	0.0
358	63323056	362	63323087	31	4.0
358	63323056	368	63323124	68	37.0
358	63323056	384	63323203	147	34.0
358	63323056	394	63323272	216	56.0
358	63323056	399	63323283	227	34.0
359	63323057	390	63323236	179	13.0
359	63323057	388	63323236	179	3.0
359	63323057	395	63323284	227	4.0
360	63323058	368	63323124	66	7.5
360	63323058	384	63323203	145	6.0
360	63323058	389	63323237	179	0.0
360	63323058	394	63323272	214	16.5
360	63323058	397	63323285	227	0.0
361	63323077	365	63323103	26	42.0
361	63323077	387	63323235	158	0.0
362	63323087	390	63323236	149	45.0
362	63323087	388	63323236	149	17.5
362	63323087	395	63323284	197	22.5
364	63323089	365	63323103	14	0.0
364	63323089	387	63323235	146	14.5
364	63323089	399	63323283	194	35.5
365	63323103	368	63323124	21	0.0
365	63323103	384	63323203	100	17.0
365	63323103	389	63323237	134	0.0
365	63323103	394	63323272	169	32.5
365	63323103	397	63323285	182	1.0
366	63323118	368	63323124	6	0.0
366	63323118	384	63323203	85	29.0
366	63323118	389	63323237	119	14.0
366	63323118	394	63323272	154	48.5
366	63323118	397	63323285	167	10.0
354	63323039	211	63323163	123	29.0

This example illustrates that guide pairs designed to introduce a stop codon have a wide variety of editing efficiencies in porcine fibroblasts.

Example 12

This example illustrates the editing efficiency of guides introducing stop codons in porcine blastocysts.

81

A subset of guides screened in porcine fetal fibroblasts were additionally tested for their ability to introduce a premature stop codon in exon 7 of CD163 in porcine embryos. The subset of guides to be tested in porcine embryos was chosen based on a combination of their efficacy in introducing a premature stop codon in exon 7 of CD163 in porcine fibroblasts and the lack of observed off-targets for each guide in the pair as described supra. Editing frequency was determined as described in Example 3. The frequencies of end-to-end NHEJ repairs resulting in the introduction of a premature stop codon in exon 7 of CD163 are shown in Table 21.

TABLE 21

End-to-end repair frequencies for introduction of a stop codon in CD163 exon 7 deletion in porcine blastocysts					
SEQ ID NO: (5')	Cut site (5')	SEQ ID NO: (3')	Cut site (3')	Deletion size (bp)	Desired repair outcome (%)
351	63323023	365	63323103	80	39.6
348	63323027	390	63323236	209	31
348	63323027	388	63323236	209	35.4
354	63323039	390	63323236	197	30.4
358	63323056	394	63323272	216	29.2
362	63323087	390	63323236	149	31
366	63323118	394	63323272	154	33
354	63323039	211	63323163	123	27.4

This example demonstrates that guide pairs that can be used to introduce a premature stop codon into exon 7 of CD163 have a wide variety of editing efficiencies.

Example 13

This example illustrates variable repair outcomes for NHEJ repair.

A subset of guideRNAs designed to delete CD163 exon 7 were tested for their ability to delete the exon 7 coding and flanking regions in porcine blastocysts, as described in Example 9. Editing frequency of blastocysts was determined as described in Example 3. In this example, a subset of the DNA sequences observed in porcine blastocysts in vivo as the result of NHEJ-mediated repair are shown for five guideRNA pairs in Table 7. It is known that simple nucleotide deletions also occur, as well as more complex NHEJ-mediated repair DNA sequences which contain deletions, insertions, rearrangements, inversions, and any combination thereof. Without being limited by theory, these varied repair outcomes also occur as a result of endonuclease cutting of DNA using single and paired guides in porcine blastocysts. Table 7 also shows that the frequencies of the observed DNA sequence associated with each repair outcome vary between guideRNA pairs. In some, but not all, instances the frequency of DNA sequence associated with the end-to-end joining of cut sites of the paired guideRNA was the most highly represented repair outcome. For some guideRNA pairs there was a single predominant DNA repair outcome, while for other guideRNA pairs there were DNA repair outcomes that occurred with equal frequency. Most of the DNA repair outcomes shown for these five guideRNA pairs resulted in the deletion of CD163 exon 7 DNA sequences corresponding to the SRCR 5 domain. Therefore, the decision for which guideRNA pair to use for generation of edited pigs depends largely on the tolerance for multiple alleles in a population.

This example demonstrates that the NHEJ-mediated repair outcomes that result when two intronic guideRNAs

82

are used to delete genomic DNA vary between guideRNA pairs, not only in the manner in which DNA breaks are resolved but in the frequency of these resolutions. It is advantageous to screen guideRNA pairs in fibroblasts or embryos to observe DNA repair outcomes of single or dual guideRNAs for the generation of edited animals.

Example 14

This example illustrates a real time PCR assay for identifying the presence of the spacer sequence set forth in SEQ ID NO: 249 and/or the desired Exon 7 excision edit in cells.

Two sets of primers and two probes were designed for this assay. One set of primers were designed to flank the spacer sequence set forth in SEQ ID NO: 249 in the unedited genome. The sequences of these primers are set forth in SEQ ID NOs: 556 and 557. A probe of sequence SEQ ID NO: 558 and labeled with the HEX fluorescent moiety was designed to anneal to the unedited genome between the PCR primers. The other set of primers, having sequences set forth in SEQ ID NOs: 562 and 563, were designed to flank the desired edit sequence. A probe, having a sequence set forth in SEQ ID NO: 564 and labeled with the FAM fluorescent moiety, was designed to anneal to nucleotides spanning the joining region of the edit. Real time PCR was performed using both primer sets and with genomic DNA extracted from tail and/or ear samples of pigs that had known allelic status (wild type, homozygous, or heterozygous). 5 μ l of 2 $\times$ PRIME-TIME[®] Master Mix (Integrated DNA Technologies, Coralville, IA) was mixed with 0.5 μ l of each primer (10 μ M), 0.5 μ l of each probe (2.5 μ M), and 2 μ l of genomic DNA. PCR was performed with 3 minutes initial denaturing at 95° C., then 35 cycles of: 95° C. for 15 seconds, 64° C. for 30 seconds, and 72° C. for 30 seconds. A final elongation was performed at 72° C. for 2 minutes, and then the reaction was held in the cyclor at 4° C. Fluorescence was measured and charted and, as expected, the homozygotes were close to the y axis (representing the FAM moiety wavelength), the heterozygotes grouped near the center of the chart, and the wild type pigs grouped close to the X axis (representing the HEX moiety wavelength). The assay was therefore successful in detecting the edit based on the spacer sequence set forth in SEQ ID NO: 249.

Example 15

This example illustrates a real time PCR assay for identifying the presence of the spacer sequence set forth in SEQ ID NO: 256 and/or the desired Exon 7 excision edit in cells.

Two sets of primers and two probes were designed for this assay. One set of primers flanked the spacer sequence set forth in SEQ ID NO: 256. The sequence of these primers is set forth in SEQ ID NOs: 559 and 560. A probe, having a sequence set forth in SEQ ID NO: 561 and labeled with the HEX fluorescent moiety, was designed to anneal to the unedited version of the spacer sequence. The other set of primers and probe are designed to target the desired edit and are described in Example 14 (SEQ ID NOs: 562-564.) Real time PCR was performed as described in Example 14, but with the instant primers and probes. Fluorescence was charted and, as expected, the homozygotes were close to the y axis (representing the FAM moiety), the heterozygotes grouped near the center of the chart, and the wild type pigs grouped close to the X axis (representing the HEX moiety). The assay was therefore successful in detecting the edit based on the spacer sequence set forth in SEQ ID NO: 256.

83

Example 16

This example illustrates a comparison of CD163 CRISPR-CAS gRNA activity in cells between previously published guide pairs and guide pairs of the present teachings.

Each pair of guides was tested in porcine fibroblasts as described in Example 2. Each pair was further tested in porcine blastocysts as described in Example 10. The results are shown below in Table 22.

TABLE 22

Comparative Activity in Porcine Cells			
guide pair SEQ ID NOs:	Source	% desired repair fibroblasts	% desired repair blastocysts
272 and 273	Burkard 2017	6	20
249 and 256	Present Disclosure	38	44
354 and 211	Whitworth 2014	17	n.d.
362 and 390	Present Disclosure	45	31

These data illustrate that the guides of the present disclosure provide at least a two-fold improvement in the percentage of cells that have the desired edit relative to the guides that were previously disclosed in the literature. Therefore, the guides of the present disclosure are more efficient than the guides previously disclosed in the literature.

Example 17

This example illustrates CD163 edited pigs challenged with PRRSV Type I.

Pigs from PICTTM Line 2 were edited with guides as set forth in SEQ ID NOs: 249 and 256 as described in Example 6. Edits were confirmed as described in Example 5. Edited pigs were then crossbred to create pigs that were homozygous for the edit. These homozygous edited pigs were inoculated with 3 ml of PRSSv Type I (SD13-15) having 10⁴ to 10⁵ virions (4-5 log TCID₅₀). 1.5 ml was administered intramuscularly with a 21 gauge needle. The remaining 1.5 ml was administered intranasally. Serum samples were obtained on Day 0 (prior to inoculation on that day), Day 3, Day 5, Day 10, Day 14, and Day 21. Realtime PCR to determine the presence of virus in the serum samples using TETRACORE® EZ-PRRSV MPX 4.0 Master Mix and Enzyme with ROX (TETRACORE®, Rockville, MD) according to manufacturer directions. The real time PCR EU adjusted counts are shown in Table 23. The counts have been inverted using standard methods known in the art to make the data more intuitive—higher numbers indicate more virus detected.

TABLE 23

Realtime PCR EU Adjusted Counts for Type I PRRSV Challenge							
Animal	CD163	Day 0	Day 3	Day 5	Day 10	Day 14	Day 21
2	WT	0	19.8	22.6	20	19.6	18.3
4	WT	0	17	17.7	18.6	16.8	17.5
5	WT	0	18.9	21.5	17.3	14.2	13.6
6	WT	0	15.8	19.7	15.6	14.5	13.4
17	WT	0	17.1	17.8	18.5	18	15.8
18	WT	0	15.8	17.9	17	16.8	15
20	WT	0	17.2	20.3	18.8	17	12.8
3	Edit	0	0	0	0	0	0
7	Edit	0	0	0	0	0	0

84

TABLE 23-continued

Realtime PCR EU Adjusted Counts for Type I PRRSV Challenge							
Animal	CD163	Day 0	Day 3	Day 5	Day 10	Day 14	Day 21
14	Edit	0	0	0	0	0	0
19	Edit	0	0	0	0	0	0

No PRRSV was detected in the serum of the edited animals throughout the experiment.

The serum samples were also subjected to ELISA using the IDEXX PRRS X3 antibody test kit; the test was performed by an accredited Veterinary Diagnostic Laboratory. The results are shown as a Sample:Positive ratio. Ratios greater than or equal to 0.40 are considered positive. The ratios for each sample are shown in Table 24.

TABLE 24

ELISA S/P Results for Type I PRRSV Challenge							
Animal	CD163	Day 0	Day 3	Day 5	Day 10	Day 14	Day 21
2	WT	0.0	0.0	0.0	0.5	0.6	1.4
4	WT	0.0	0.0	0.0	0.1	0.2	0.8
5	WT	0.0	0.0	0.0	0.9	1.3	1.8
6	WT	0.0	0.0	0.0	1.3	1.3	1.9
17	WT	0.0	0.0	0.0	0.2	0.2	1.1
18	WT	0.0	0.0	0.0	0.3	0.3	1.4
20	WT	0.0	0.0	0.0	0.3	0.4	1.2
3	Edit	0.0	0.0	0.0	0.0	0.0	0.0
7	Edit	0.0	0.0	0.0	0.0	0.0	0.0
14	Edit	0.0	0.0	0.0	0.0	0.0	0.0
19	Edit	0.0	0.0	0.0	0.0	0.0	0.0

The edited pigs do not have any positive ratios. In contrast, by day 21, all of the wild type pigs have positive ratios. This further illustrates that there is no PRRSV circulating in the edited pigs' serum.

This example illustrates that pigs edited with guides of SEQ ID NOs: 249 and 256 are resistant to PRRSV Type I virus infection.

Example 18

This example illustrates CD163 edited pigs challenged with PRRSV Type II.

Pigs from PICTTM Lines 2 and 3 were edited with guides as set forth in SEQ ID NOs: 249 and 256 as described in Example 6. Edits were confirmed as described in Example 5. Edited pigs were then crossbred to create pigs that were homozygous for the edit. These homozygous edited pigs were then inoculated with 3 ml of PRRSV Type II (NVSL 97-7895) having 10⁴ to 10⁵ virions (4-5 log TCID₅₀). 1.5 ml was administered intramuscularly with a 21 gauge needle. The remaining 1.5 ml was administered intranasally. Serum samples were obtained on Day 0 (prior to inoculation on that day), Day 3, Day 5, Day 10, Day 14, and Day 21. Realtime PCR to determine the presence of virus in the serum samples using TETRACORE® EZ-PRRSV MPX 4.0 Master Mix and Enzyme with ROX according to manufacturer directions. The NA adjusted counts for real time PCR are shown in

Table 25. The numbers were inverted to make them more intuitive—the higher the count, the more virus is present.

85

TABLE 25

Realtime PCR NA Adjusted Counts for Type II PRRSv Challenge							
Animal	CD163	Day 0	Day 3	Day 5	Day 10	Day 14	Day 21
21	WT	0.0	18.4	22.0	21.9	18.0	16.1
23	WT	0.0	16.4	22.0	21.8	20.0	19.7
25	WT	N/A	17.2	21.2	21.9	19.2	19.8
30	WT	0.0	16.2	17.8	21.4	21.1	17.6
33	WT	0.0	18.5	N/A	21.8	18.5	15.3
35	WT	N/A	17.0	22.9	21.8	20.5	20.9
36	WT	0.0	16.7	21.4	21.9	19.9	17.4
38	WT	0.0	16.0	18.1	20.2	20.9	9.4
24	Edit	N/A	1.9	0.0	0.0	0.0	0.0
37	Edit	0.0	0.0	0.0	0.0	0.0	0.0

The edited pigs have very little to no virus counts relative to the wild type pigs.

The serum samples were also subjected to ELISA using the IDEXX PRRS X3 antibody test kit; the test was performed by an accredited Veterinary Diagnostic Laboratory. The results are shown as a Sample:Positive ratio. Ratios greater than or equal to 0.40 are considered positive. The ratios for each sample are shown in Table 26.

86

TABLE 26

ELISA S/P Ratios for Type II PRRSv Challenge							
Animal	Edit	Day 0	Day 3	Day 5	Day 10	Day 14	Day 21
21	WT	0.0	0.0	0.0	1.7	2.0	1.9
23	WT	0.0	0.0	0.0	1.2	1.7	1.9
25	WT	N/A	0.0	0.0	1.2	1.6	1.7
30	WT	0.0	0.0	0.0	0.4	0.6	0.6
33	WT	0.0	0.0	N/A	0.7	0.6	0.7
35	WT	N/A	0.0	0.0	1.3	1.3	1.4
36	WT	0.0	0.0	0.0	1.2	1.5	1.8
38	WT	0.0	0.0	0.0	0.5	0.9	1.0
24	Edit	N/A	0.2	0.1	0.2	0.1	0.1
37	Edit	0.0	0.0	0.0	0.0	0.0	0.0

The edited pigs do not have any positive ratios; in contrast, by day 10, all of the wild type pigs have positive ratios. Taken together, these data illustrate that there is no virus circulating in the edited pigs' blood.

This example illustrates that the pigs with a CD163 gene edited with SEQ ID NOS: 249 and 256 are resistant to PRRSv Type II infection.

The contents of each of the foregoing references and applications are incorporated herein by reference in their entireties. Having described the present disclosure in detail, it will be apparent that modifications and variations are possible without departing from the scope of the teachings defined in the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 564

<210> SEQ ID NO 1

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 1

tttcgcagac tttagaagat gttctgcccc ttttaagttcc ttcacttttg ctgtagtcgc 60

tggtctcagt gcttgactag ttctcttggt gactactttg acaaatttac ttgtaacctc 120

gcccactgtg acaagaaaca ctgaaaagca 150

<210> SEQ ID NO 2

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 2

tttcgcagac tttagaagat gttctgcccc ttttaagttcc ttcacttttg ctgtagtcgc 60

tggtctcagt gcttgactag ttctcttggt gactactttg acaaatttac ttgtaacctc 120

gcccactgtg acaagaaaca ctgaaaagca 150

<210> SEQ ID NO 3

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 3

gtggtgaaaa caagtgtctt ggaagagtgg aggtgaaagt gcaggaggag tggggaactg 60

tgtgtaataa tggtgtacat ggatgtggtc tctgttgttt gtaggcagct gggatgtcca 120

actgctatca aagccactgg atgggctaata 150

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 4

tggtctgggac atggatgtgg tctctgttgt ttgtaggcag ctgggatgac caactgctat	60
caaagccact ggatgaatgt tagtcaggt tctggacgca tttggatgga tcatgtttct	120
tgtcgaggga atgagtcagc tctctgggac	150

<210> SEQ ID NO 5
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 5

gtaggcagct gggatgtcca actgctatca aagccactgg atgggctaatt ttagtgagcag	60
gttctggacg catttgatgg atcatgtttc ttgtcgaggg aatgagtcag ctctctggga	120
ctgcaaacat gatggatggg gaaagcataa	150

<210> SEQ ID NO 6
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 6

taggcagctg ggatgtccaa ctgctatcaa agccactgga tgggctaatt ttagtgagcag	60
ttctggacgc atttgatgga tcatgtttct ttgtcgagga atgagtcagc tctctgggac	120
tgcaaacatg atggatgggg aaagcataac	150

<210> SEQ ID NO 7
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 7

gtccaactgc tatcaaagcc actggatggg ctaatttttag tgcaggttct ggacgcattt	60
ggatggatca tgtttgaggg aatgagtcag ctctctggga ctgcaaacat gatggatggg	120
gaaagcataa ctgtactcac caacaggatg	150

<210> SEQ ID NO 8
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 8

ctggacgcat ttggatggat catgtttctt gtcgaggga tgagtcagct ctctgggact	60
gcaaacatga tggatgaaag cataactgta ctcaccaaca ggatgctgga gtaacctgct	120
caggtgaagac atacacaaat aagtcaagcc	150

<210> SEQ ID NO 9
 <211> LENGTH: 150
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

 <400> SEQUENCE: 9

tggaacgatt tggatggatc atgtttcttg tcgagggaat gagtcagctc tctgggactg	60
---	----

-continued

 caaacatgat ggatgaaagc ataactgtac tcaccaacag gatgctggag taacctgctc 120

aggtaagaca tacacaaata agtcaagcct 150

<210> SEQ ID NO 10

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 10

ctctagatgg atctgattta gagatgaggc tggatgaatgg aggaaacggg tgcttaggaa 60

gaatagaagt caaataagga cgggtggggaa cagtgtgtga tgataacttc aacataaatc 120

atgcttctgt gggttgtaaa caacttgaat 150

<210> SEQ ID NO 11

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 11

cttctgtggt ttgtaaacaa cttgaatgtg gaagtgtgt cagtttctct gggtcagcta 60

attttgaga aggttgacca atctggttg atgatctgt atgcaatgga aatgagtcag 120

ctctctggaa ctgcaaacat gaaggatggg 150

<210> SEQ ID NO 12

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 12

tgatgatctt gtatgcaatg gaaatgagtc agctctctgg aactgcaaac atgaaggatg 60

gggaaagcac aattgatgct gaggatgctg gagtgatttg cttaagtaag gactgacctg 120

gggttgttct gttctccatg agagggcaaa 150

<210> SEQ ID NO 13

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 13

agacacgtgg ggcacgtct gtgattctga cttctctctg gaggcggcca gcgtgctgtg 60

cagggaacta cagtgactgt gggttccctc ctggggggag ctcacttttg agaaggaagt 120

ggacagatct gggctgaaga attccagtgt 150

<210> SEQ ID NO 14

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 14

agaagatgtt ctgcccattt aagttccttc acttttctg tagtgcgtgt tctcagtgcc 60

tgcttggtca ctagtcttg gtgagtactt tgacaaattt acttgtaacc tagccactg 120

tgacaagaaa cactgaaaag caaataattc 150

<210> SEQ ID NO 15

<211> LENGTH: 150

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 15

gactttagaa gatgttctgc ccatttaagt tccttcactt ttgctgtagt cgtgtttctc	60
agtgccctgct tggtcacttg taacctagcc cactgtgaca agaaacactg aaaagcaaat	120
aattctcctg aagtctagat agcatctaaa	150

<210> SEQ ID NO 16

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 16

gaagatgttc tgcccattta agttccttca cttttgctgt agtcgctgtt ctcaagtgcct	60
gcttggtcac tagttctctc ttggtgagta ctttgacaaa ttacttgta acctagccca	120
ctgtgacaag aaacactgaa aagcaataaa	150

<210> SEQ ID NO 17

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 17

tttcgcagac tttagaagat gttctgccca tttaagttcc ttcacttttg ctgtagtcgc	60
tgttctcagt gctgtcttgg tgagtacttt gacaaattta cttgtaacct agcccactgt	120
gacaagaaac actgaaaagc aaataattct	150

<210> SEQ ID NO 18

<211> LENGTH: 150

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 18

tagaagatgt tctgccatt taagttcctt cacttttgcg gtagtcgctg ttctcagtgc	60
ctgcttggtc actaggagta ctttgacaaa ttacttgta acctagccca ctgtgacaag	120
aaacactgaa aagcaataaa ttctctgaa	150

<210> SEQ ID NO 19

<211> LENGTH: 82

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 19

gttttagagc tagaaatagc aagttaaaat aaggctagtc cggtatcaac ttgaaaaagt	60
ggcaccgagt cgggtgctttt tt	82

<210> SEQ ID NO 20

<211> LENGTH: 1380

<212> TYPE: PRT

<213> ORGANISM: *Streptococcus pyogenes*

<400> SEQUENCE: 20

Gly Met Asp Lys Lys Tyr Ser Ile Gly Leu Asp Ile Gly Thr Asn Ser	
1 5 10 15	

Val Gly Trp Ala Val Ile Thr Asp Glu Tyr Lys Val Pro Ser Lys Lys	
20 25 30	

Phe Lys Val Leu Gly Asn Thr Asp Arg His Ser Ile Lys Lys Asn Leu	
---	--

-continued

35	40	45
Ile Gly Ala Leu Leu Phe Asp Ser Gly Glu Thr Ala Glu Ala Thr Arg		
50	55	60
Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Lys Asn Arg Ile		
65	70	75
Cys Tyr Leu Gln Glu Ile Phe Ser Asn Glu Met Ala Lys Val Asp Asp		
85	90	95
Ser Phe Phe His Arg Leu Glu Glu Ser Phe Leu Val Glu Glu Asp Lys		
100	105	110
Lys His Glu Arg His Pro Ile Phe Gly Asn Ile Val Asp Glu Val Ala		
115	120	125
Tyr His Glu Lys Tyr Pro Thr Ile Tyr His Leu Arg Lys Lys Leu Val		
130	135	140
Asp Ser Thr Asp Lys Ala Asp Leu Arg Leu Ile Tyr Leu Ala Leu Ala		
145	150	155
His Met Ile Lys Phe Arg Gly His Phe Leu Ile Glu Gly Asp Leu Asn		
165	170	175
Pro Asp Asn Ser Asp Val Asp Lys Leu Phe Ile Gln Leu Val Gln Thr		
180	185	190
Tyr Asn Gln Leu Phe Glu Glu Asn Pro Ile Asn Ala Ser Gly Val Asp		
195	200	205
Ala Lys Ala Ile Leu Ser Ala Arg Leu Ser Lys Ser Arg Arg Leu Glu		
210	215	220
Asn Leu Ile Ala Gln Leu Pro Gly Glu Lys Lys Asn Gly Leu Phe Gly		
225	230	235
Asn Leu Ile Ala Leu Ser Leu Gly Leu Thr Pro Asn Phe Lys Ser Asn		
245	250	255
Phe Asp Leu Ala Glu Asp Ala Lys Leu Gln Leu Ser Lys Asp Thr Tyr		
260	265	270
Asp Asp Asp Leu Asp Asn Leu Leu Ala Gln Ile Gly Asp Gln Tyr Ala		
275	280	285
Asp Leu Phe Leu Ala Ala Lys Asn Leu Ser Asp Ala Ile Leu Leu Ser		
290	295	300
Asp Ile Leu Arg Val Asn Thr Glu Ile Thr Lys Ala Pro Leu Ser Ala		
305	310	315
Ser Met Ile Lys Arg Tyr Asp Glu His His Gln Asp Leu Thr Leu Leu		
325	330	335
Lys Ala Leu Val Arg Gln Gln Leu Pro Glu Lys Tyr Lys Glu Ile Phe		
340	345	350
Phe Asp Gln Ser Lys Asn Gly Tyr Ala Gly Tyr Ile Asp Gly Gly Ala		
355	360	365
Ser Gln Glu Glu Phe Tyr Lys Phe Ile Lys Pro Ile Leu Glu Lys Met		
370	375	380
Asp Gly Thr Glu Glu Leu Leu Val Lys Leu Asn Arg Glu Asp Leu Leu		
385	390	395
Arg Lys Gln Arg Thr Phe Asp Asn Gly Ser Ile Pro His Gln Ile His		
405	410	415
Leu Gly Glu Leu His Ala Ile Leu Arg Arg Gln Glu Asp Phe Tyr Pro		
420	425	430
Phe Leu Lys Asp Asn Arg Glu Lys Ile Glu Lys Ile Leu Thr Phe Arg		
435	440	445
Ile Pro Tyr Tyr Val Gly Pro Leu Ala Arg Gly Asn Ser Arg Phe Ala		
450	455	460

Trp	Met	Thr	Arg	Lys	Ser	Glu	Glu	Thr	Ile	Thr	Pro	Trp	Asn	Phe	Glu
465					470					475					480
Glu	Val	Val	Asp	Lys	Gly	Ala	Ser	Ala	Gln	Ser	Phe	Ile	Glu	Arg	Met
			485						490					495	
Thr	Asn	Phe	Asp	Lys	Asn	Leu	Pro	Asn	Glu	Lys	Val	Leu	Pro	Lys	His
			500					505					510		
Ser	Leu	Leu	Tyr	Glu	Tyr	Phe	Thr	Val	Tyr	Asn	Glu	Leu	Thr	Lys	Val
		515					520					525			
Lys	Tyr	Val	Thr	Glu	Gly	Met	Arg	Lys	Pro	Ala	Phe	Leu	Ser	Gly	Glu
	530					535					540				
Gln	Lys	Lys	Ala	Ile	Val	Asp	Leu	Leu	Phe	Lys	Thr	Asn	Arg	Lys	Val
545					550					555					560
Thr	Val	Lys	Gln	Leu	Lys	Glu	Asp	Tyr	Phe	Lys	Lys	Ile	Glu	Cys	Phe
				565					570					575	
Asp	Ser	Val	Glu	Ile	Ser	Gly	Val	Glu	Asp	Arg	Phe	Asn	Ala	Ser	Leu
			580					585					590		
Gly	Thr	Tyr	His	Asp	Leu	Leu	Lys	Ile	Ile	Lys	Asp	Lys	Asp	Phe	Leu
		595					600					605			
Asp	Asn	Glu	Glu	Asn	Glu	Asp	Ile	Leu	Glu	Asp	Ile	Val	Leu	Thr	Leu
						615					620				
Thr	Leu	Phe	Glu	Asp	Arg	Glu	Met	Ile	Glu	Glu	Arg	Leu	Lys	Thr	Tyr
625					630					635					640
Ala	His	Leu	Phe	Asp	Asp	Lys	Val	Met	Lys	Gln	Leu	Lys	Arg	Arg	Arg
				645					650					655	
Tyr	Thr	Gly	Trp	Gly	Arg	Leu	Ser	Arg	Lys	Leu	Ile	Asn	Gly	Ile	Arg
			660					665					670		
Asp	Lys	Gln	Ser	Gly	Lys	Thr	Ile	Leu	Asp	Phe	Leu	Lys	Ser	Asp	Gly
		675					680					685			
Phe	Ala	Asn	Arg	Asn	Phe	Met	Gln	Leu	Ile	His	Asp	Asp	Ser	Leu	Thr
						695					700				
Phe	Lys	Glu	Asp	Ile	Gln	Lys	Ala	Gln	Val	Ser	Gly	Gln	Gly	Asp	Ser
705					710					715					720
Leu	His	Glu	His	Ile	Ala	Asn	Leu	Ala	Gly	Ser	Pro	Ala	Ile	Lys	Lys
				725					730					735	
Gly	Ile	Leu	Gln	Thr	Val	Lys	Val	Val	Asp	Glu	Leu	Val	Lys	Val	Met
			740					745					750		
Gly	Arg	His	Lys	Pro	Glu	Asn	Ile	Val	Ile	Glu	Met	Ala	Arg	Glu	Asn
		755					760					765			
Gln	Thr	Thr	Gln	Lys	Gly	Gln	Lys	Asn	Ser	Arg	Glu	Arg	Met	Lys	Arg
					775						780				
Ile	Glu	Glu	Gly	Ile	Lys	Glu	Leu	Gly	Ser	Gln	Ile	Leu	Lys	Glu	His
785					790					795					800
Pro	Val	Glu	Asn	Thr	Gln	Leu	Gln	Asn	Glu	Lys	Leu	Tyr	Leu	Tyr	Tyr
				805					810					815	
Leu	Gln	Asn	Gly	Arg											

Lys	Asn	Tyr	Trp	Arg	Gln	Leu	Leu	Asn	Ala	Lys	Leu	Ile	Thr	Gln	Arg
				885					890				895		
Lys	Phe	Asp	Asn	Leu	Thr	Lys	Ala	Glu	Arg	Gly	Gly	Leu	Ser	Glu	Leu
				900					905				910		
Asp	Lys	Ala	Gly	Phe	Ile	Lys	Arg	Gln	Leu	Val	Glu	Thr	Arg	Gln	Ile
				915					920				925		
Thr	Lys	His	Val	Ala	Gln	Ile	Leu	Asp	Ser	Arg	Met	Asn	Thr	Lys	Tyr
				930					935				940		
Asp	Glu	Asn	Asp	Lys	Leu	Ile	Arg	Glu	Val	Lys	Val	Ile	Thr	Leu	Lys
				945									950		
Ser	Lys	Leu	Val	Ser	Asp	Phe	Arg	Lys	Asp	Phe	Gln	Phe	Tyr	Lys	Val
				965									970		
Arg	Glu	Ile	Asn	Asn	Tyr	His	His	Ala	His	Asp	Ala	Tyr	Leu	Asn	Ala
				980									985		
Val	Val	Gly	Thr	Ala	Leu	Ile	Lys	Lys	Tyr	Pro	Lys	Leu	Glu	Ser	Glu
				995									1000		
Phe	Val	Tyr	Gly	Asp	Tyr	Lys	Val	Tyr	Asp	Val	Arg	Lys	Met	Ile	
				1010									1015		
Ala	Lys	Ser	Glu	Gln	Glu	Ile	Gly	Lys	Ala	Thr	Ala	Lys	Tyr	Phe	
				1025									1030		
Phe	Tyr	Ser	Asn	Ile	Met	Asn	Phe	Phe	Lys	Thr	Glu	Ile	Thr	Leu	
				1040									1045		
Ala	Asn	Gly	Glu	Ile	Arg	Lys	Arg	Pro	Leu	Ile	Glu	Thr	Asn	Gly	
				1055									1060		
Glu	Thr	Gly	Glu	Ile	Val	Trp	Asp	Lys	Gly	Arg	Asp	Phe	Ala	Thr	
				1070									1075		
Val	Arg	Lys	Val	Leu	Ser	Met	Pro	Gln	Val	Asn	Ile	Val	Lys	Lys	
				1085									1090		
Thr	Glu	Val	Gln	Thr	Gly	Gly	Phe	Ser	Lys	Glu	Ser	Ile	Leu	Pro	
				1100									1105		
Lys	Arg	Asn	Ser	Asp	Lys	Leu	Ile	Ala	Arg	Lys	Lys	Asp	Trp	Asp	
				1115									1120		
Pro	Lys	Lys	Tyr	Gly	Gly	Phe	Asp	Ser	Pro	Thr	Val	Ala	Tyr	Ser	
				1130									1135		
Val	Leu	Val	Val	Ala	Lys	Val	Glu	Lys	Gly	Lys	Ser	Lys	Lys	Leu	
				1145									1150		
Lys	Ser	Val	Lys	Glu	Leu	Leu	Gly	Ile	Thr	Ile	Met	Glu	Arg	Ser	
				1160									1165		
Ser	Phe	Glu	Lys	Asn	Pro	Ile	Asp	Phe	Leu	Glu	Ala	Lys	Gly	Tyr	
				1175									1180		
Lys	Glu	Val	Lys	Lys	Asp	Leu	Ile	Ile	Lys	Leu	Pro	Lys	Tyr	Ser	
				1190									1195		
Leu	Phe	Glu	Leu	Glu	Asn	Gly	Arg	Lys	Arg	Met	Leu	Ala	Ser	Ala	
				1205									1210		
Gly	Glu	Leu	Gln	Lys	Gly	Asn	Glu	Leu	Ala	Leu	Pro	Ser	Lys	Tyr	
				1220									1225		
Val	Asn	Phe	Leu	Tyr	Leu	Ala	Ser	His	Tyr	Glu	Lys	Leu	Lys	Gly	
				1235									1240		
Ser	Pro	Glu	Asp	Asn	Glu	Gln	Lys	Gln	Leu	Phe	Val	Glu	Gln	His	
				1250									1255		
Lys	His	Tyr	Leu	Asp	Glu	Ile	Ile	Glu	Gln	Ile	Ser	Glu	Phe	Ser	
				1265									1270		
Lys	Arg	Val	Ile	Leu	Ala	Asp	Ala	Asn	Leu	Asp	Lys	Val	Leu	Ser	
				1280									1285		

-continued

1280	1285	1290
Ala Tyr Asn Lys His Arg Asp Lys Pro Ile Arg Glu Gln Ala Glu		
1295	1300	1305
Asn Ile Ile His Leu Phe Thr Leu Thr Asn Leu Gly Ala Pro Ala		
1310	1315	1320
Ala Phe Lys Tyr Phe Asp Thr Thr Ile Asp Arg Lys Arg Tyr Thr		
1325	1330	1335
Ser Thr Lys Glu Val Leu Asp Ala Thr Leu Ile His Gln Ser Ile		
1340	1345	1350
Thr Gly Leu Tyr Glu Thr Arg Ile Asp Leu Ser Gln Leu Gly Gly		
1355	1360	1365
Asp Gly Ser Gly Ser Pro Lys Lys Lys Arg Lys Val		
1370	1375	1380

<210> SEQ ID NO 21

<211> LENGTH: 1400

<212> TYPE: PRT

<213> ORGANISM: Streptococcus thermophilus

<400> SEQUENCE: 21

Ser Asn Ala Met Thr Lys Pro Tyr Ser Ile Gly Leu Asp Ile Gly Thr		
1	5	10
Asn Ser Val Gly Trp Ala Val Ile Thr Asp Asn Tyr Lys Val Pro Ser		
20	25	30
Lys Lys Met Lys Val Leu Gly Asn Thr Ser Lys Lys Tyr Ile Lys Lys		
35	40	45
Asn Leu Leu Gly Val Leu Leu Phe Asp Ser Gly Ile Thr Ala Glu Gly		
50	55	60
Arg Arg Leu Lys Arg Thr Ala Arg Arg Arg Tyr Thr Arg Arg Arg Asn		
65	70	75
Arg Ile Leu Tyr Leu Gln Glu Ile Phe Ser Thr Glu Met Ala Thr Leu		
85	90	95
Asp Asp Ala Phe Phe Gln Arg Leu Asp Asp Ser Phe Leu Val Pro Asp		
100	105	110
Asp Lys Arg Asp Ser Lys Tyr Pro Ile Phe Gly Asn Leu Val Glu Glu		
115	120	125
Lys Val Tyr His Asp Glu Phe Pro Thr Ile Tyr His Leu Arg Lys Tyr		
130	135	140
Leu Ala Asp Ser Thr Lys Lys Ala Asp Leu Arg Leu Val Tyr Leu Ala		
145	150	155
Leu Ala His Met Ile Lys Tyr Arg Gly His Phe Leu Ile Glu Gly Glu		
165	170	175
Phe Asn Ser Lys Asn Asn Asp Ile Gln Lys Asn Phe Gln Asp Phe Leu		
180	185	190
Asp Thr Tyr Asn Ala Ile Phe Glu Ser Asp Leu Ser Leu Glu Asn Ser		
195	200	205
Lys Gln Leu Glu Glu Ile Val Lys Asp Lys Ile Ser Lys Leu Glu Lys		
210	215	220
Lys Asp Arg Ile Leu Lys Leu Phe Pro Gly Glu Lys Asn Ser Gly Ile		
225	230	235
Phe Ser Glu Phe Leu Lys Leu Ile Val Gly Asn Gln Ala Asp Phe Arg		
245	250	255
Lys Cys Phe Asn Leu Asp Glu Lys Ala Ser Leu His Phe Ser Lys Glu		
260	265	270

Ser	Tyr	Asp	Glu	Asp	Leu	Glu	Thr	Leu	Leu	Gly	Tyr	Ile	Gly	Asp	Asp
		275					280					285			
Tyr	Ser	Asp	Val	Phe	Leu	Lys	Ala	Lys	Lys	Leu	Tyr	Asp	Ala	Ile	Leu
		290					295					300			
Leu	Ser	Gly	Phe	Leu	Thr	Val	Thr	Asp	Asn	Glu	Thr	Glu	Ala	Pro	Leu
		305					310					315			
Ser	Ser	Ala	Met	Ile	Lys	Arg	Tyr	Asn	Glu	His	Lys	Glu	Asp	Leu	Ala
				325							330			335	
Leu	Leu	Lys	Glu	Tyr	Ile	Arg	Asn	Ile	Ser	Leu	Lys	Thr	Tyr	Asn	Glu
				340											350
Val	Phe	Lys	Asp	Asp	Thr	Lys	Asn	Gly	Tyr	Ala	Gly	Tyr	Ile	Asp	Gly
				355											365
Lys	Thr	Asn	Gln	Glu	Asp	Phe	Tyr	Val	Tyr	Leu	Lys	Asn	Leu	Leu	Ala
				370											380
Glu	Phe	Glu	Gly	Ala	Asp	Tyr	Phe	Leu	Glu	Lys	Ile	Asp	Arg	Glu	Asp
						390									400
Phe	Leu	Arg	Lys	Gln	Arg	Thr	Phe	Asp	Asn	Gly	Ser	Ile	Pro	Tyr	Gln
						405									415
Ile	His	Leu	Gln	Glu	Met	Arg	Ala	Ile	Leu	Asp	Lys	Gln	Ala	Lys	Phe
						420									430
Tyr	Pro	Phe	Leu	Ala	Lys	Asn	Lys	Glu	Arg	Ile	Glu	Lys	Ile	Leu	Thr
						435									445
Phe	Arg	Ile	Pro	Tyr	Tyr	Val	Gly	Pro	Leu	Ala	Arg	Gly	Asn	Ser	Asp
						450									460
Phe	Ala	Trp	Ser	Ile	Arg	Lys	Arg	Asn	Glu	Lys	Ile	Thr	Pro	Trp	Asn
						470									480
Phe	Glu	Asp	Val	Ile	Asp	Lys	Glu	Ser	Ser	Ala	Glu	Ala	Phe	Ile	Asn
						485									495
Arg	Met	Thr	Ser	Phe	Asp	Leu	Tyr	Leu	Pro	Glu	Glu	Lys	Val	Leu	Pro
						500									510
Lys	His	Ser	Leu	Leu	Tyr	Glu	Thr	Phe	Asn	Val	Tyr	Asn	Glu	Leu	Thr
						515									525
Lys	Val	Arg	Phe	Ile	Ala	Glu	Ser	Met	Arg	Asp	Tyr	Gln	Phe	Leu	Asp
						530									540
Ser	Lys	Gln	Lys	Lys	Asp	Ile	Val	Arg	Leu	Tyr	Phe	Lys	Asp	Lys	Arg
						550									560
Lys	Val	Thr	Asp	Lys	Asp	Ile	Ile	Glu	Tyr	Leu	His	Ala	Ile	Tyr	Gly
						565									575
Tyr	Asp	Gly	Ile	Glu	Leu	Lys	Gly	Ile	Glu	Lys	Gln	Phe	Asn	Ser	Ser
						580									590
Leu	Ser	Thr	Tyr	His	Asp	Leu	Leu	Asn	Ile	Ile	Asn	Asp	Lys	Glu	Phe
						595									605
Leu	Asp	Asp	Ser	Ser	Asn	Glu	Ala	Ile	Ile	Glu	Glu	Ile	Ile	His	Thr
						610									620
Leu	Thr	Ile	Phe	Glu	Asp	Arg	Glu	Met	Ile	Lys	Gln	Arg	Leu	Ser	Lys
						630									640
Phe	Glu	Asn	Ile	Phe	Asp	Lys	Ser	Val	Leu	Lys	Lys	Leu	Ser		

-continued

690	695	700
Ser Phe Lys Lys Lys Ile Gln Lys Ala Gln Ile Ile Gly Asp Glu Asp 705 710 715 720		
Lys Gly Asn Ile Lys Glu Val Val Lys Ser Leu Pro Gly Ser Pro Ala 725 730 735		
Ile Lys Lys Gly Ile Leu Gln Ser Ile Lys Ile Val Asp Glu Leu Val 740 745 750		
Lys Val Met Gly Gly Arg Lys Pro Glu Ser Ile Val Val Glu Met Ala 755 760 765		
Arg Glu Asn Gln Tyr Thr Asn Gln Gly Lys Ser Asn Ser Gln Gln Arg 770 775 780		
Leu Lys Arg Leu Glu Lys Ser Leu Lys Glu Leu Gly Ser Lys Ile Leu 785 790 795 800		
Lys Glu Asn Ile Pro Ala Lys Leu Ser Lys Ile Asp Asn Asn Ala Leu 805 810 815		
Gln Asn Asp Arg Leu Tyr Leu Tyr Tyr Leu Gln Asn Gly Lys Asp Met 820 825 830		
Tyr Thr Gly Asp Asp Leu Asp Ile Asp Arg Leu Ser Asn Tyr Asp Ile 835 840 845		
Asp His Ile Ile Pro Gln Ala Phe Leu Lys Asp Asn Ser Ile Asp Asn 850 855 860		
Lys Val Leu Val Ser Ser Ala Ser Asn Arg Gly Lys Ser Asp Asp Phe 865 870 875 880		
Pro Ser Leu Glu Val Val Lys Lys Arg Lys Thr Phe Trp Tyr Gln Leu 885 890 895		
Leu Lys Ser Lys Leu Ile Ser Gln Arg Lys Phe Asp Asn Leu Thr Lys 900 905 910		
Ala Glu Arg Gly Gly Leu Leu Pro Glu Asp Lys Ala Gly Phe Ile Gln 915 920 925		
Arg Gln Leu Val Glu Thr Arg Gln Ile Thr Lys His Val Ala Arg Leu 930 935 940		
Leu Asp Glu Lys Phe Asn Asn Lys Lys Asp Glu Asn Asn Arg Ala Val 945 950 955 960		
Arg Thr Val Lys Ile Ile Thr Leu Lys Ser Thr Leu Val Ser Gln Phe 965 970 975		
Arg Lys Asp Phe Glu Leu Tyr Lys Val Arg Glu Ile Asn Asp Phe His 980 985 990		
His Ala His Asp Ala Tyr Leu Asn Ala Val Ile Ala Ser Ala Leu Leu 995 1000 1005		
Lys Lys Tyr Pro Lys Leu Glu Pro Glu Phe Val Tyr Gly Asp Tyr 1010 1015 1020		
Pro Lys Tyr Asn Ser Phe Arg Glu Arg Lys Ser Ala Thr Glu Lys 1025 1030 1035		
Val Tyr Phe Tyr Ser Asn Ile Met Asn Ile Phe Lys Lys Ser Ile 1040 1045 1050		
Ser Leu Ala Asp Gly Arg Val Ile Glu Arg Pro Leu Ile Glu Val 1055 1060 1065		
Asn Glu Glu Thr Gly Glu Ser Val Trp Asn Lys Glu Ser Asp Leu 1070 1075 1080		
Ala Thr Val Arg Arg Val Leu Ser Tyr Pro Gln Val Asn Val Val 1085 1090 1095		
Lys Lys Val Glu Glu Gln Asn His Gly Leu Asp Arg Gly Lys Pro 1100 1105 1110		

-continued

Lys Gly Leu Phe Asn Ala Asn Leu Ser Ser Lys Pro Lys Pro Asn
 1115 1120 1125
 Ser Asn Glu Asn Leu Val Gly Ala Lys Glu Tyr Leu Asp Pro Lys
 1130 1135 1140
 Lys Tyr Gly Gly Tyr Ala Gly Ile Ser Asn Ser Phe Ala Val Leu
 1145 1150 1155
 Val Lys Gly Thr Ile Glu Lys Gly Ala Lys Lys Lys Ile Thr Asn
 1160 1165 1170
 Val Leu Glu Phe Gln Gly Ile Ser Ile Leu Asp Arg Ile Asn Tyr
 1175 1180 1185
 Arg Lys Asp Lys Leu Asn Phe Leu Leu Glu Lys Gly Tyr Lys Asp
 1190 1195 1200
 Ile Glu Leu Ile Ile Glu Leu Pro Lys Tyr Ser Leu Phe Glu Leu
 1205 1210 1215
 Ser Asp Gly Ser Arg Arg Met Leu Ala Ser Ile Leu Ser Thr Asn
 1220 1225 1230
 Asn Lys Arg Gly Glu Ile His Lys Gly Asn Gln Ile Phe Leu Ser
 1235 1240 1245
 Gln Lys Phe Val Lys Leu Leu Tyr His Ala Lys Arg Ile Ser Asn
 1250 1255 1260
 Thr Ile Asn Glu Asn His Arg Lys Tyr Val Glu Asn His Lys Lys
 1265 1270 1275
 Glu Phe Glu Glu Leu Phe Tyr Tyr Ile Leu Glu Phe Asn Glu Asn
 1280 1285 1290
 Tyr Val Gly Ala Lys Lys Asn Gly Lys Leu Leu Asn Ser Ala Phe
 1295 1300 1305
 Gln Ser Trp Gln Asn His Ser Ile Asp Glu Leu Cys Ser Ser Phe
 1310 1315 1320
 Ile Gly Pro Thr Gly Ser Glu Arg Lys Gly Leu Phe Glu Leu Thr
 1325 1330 1335
 Ser Arg Gly Ser Ala Ala Asp Phe Glu Phe Leu Gly Val Lys Ile
 1340 1345 1350
 Pro Arg Tyr Arg Asp Tyr Thr Pro Ser Ser Leu Leu Lys Asp Ala
 1355 1360 1365
 Thr Leu Ile His Gln Ser Val Thr Gly Leu Tyr Glu Thr Arg Ile
 1370 1375 1380
 Asp Leu Ala Lys Leu Gly Glu Gly Gly Ser Pro Lys Lys Lys Arg
 1385 1390 1395
 Lys Val
 1400

<210> SEQ ID NO 22
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 22

gaagcctttc tgtatttttg tgg

23

<210> SEQ ID NO 23
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 23

-continued

atgattttta gaattcttag tgg 23

<210> SEQ ID NO 24
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 24

tcttagtggt tctcttcttc agg 23

<210> SEQ ID NO 25
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 25

tcttagtggt tctcttcttc aggag 25

<210> SEQ ID NO 26
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 26

ttcttcagga gaacatttct agg 23

<210> SEQ ID NO 27
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 27

tcttcaggag aacatttcta ggg 23

<210> SEQ ID NO 28
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 28

ataatacaag aagatttaaa tgg 23

<210> SEQ ID NO 29
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 29

atttaaattg cataaaacct tgg 23

<210> SEQ ID NO 30
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 30

aatggcataa aaccttgga tgg 23

<210> SEQ ID NO 31
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 31

-continued

ttctgagttt gtccattcca agg	23
 <210> SEQ ID NO 32 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 32	
tggaatggac aaactcagaa tgg	23
 <210> SEQ ID NO 33 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 33	
tggaatggac aaactcagaa tgggtg	25
 <210> SEQ ID NO 34 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 34	
atggtgtctac atgaaaaactc tgg	23
 <210> SEQ ID NO 35 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 35	
catgaaaact ctggatctgc agg	23
 <210> SEQ ID NO 36 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 36	
aaaagtgaag gaacttaaat ggg	23
 <210> SEQ ID NO 37 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 37	
caaaagtga ggaacttaaa tgg	23
 <210> SEQ ID NO 38 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 38	
agcgactaca gcaaaagtga agg	23
 <210> SEQ ID NO 39 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa	

-continued

<400> SEQUENCE: 39

cgctgttctc agtgctgct tgg 23

<210> SEQ ID NO 40

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 40

aagagaacta gtgaccaagc agg 23

<210> SEQ ID NO 41

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 41

tgcttggtca ctagttctct tgg 23

<210> SEQ ID NO 42

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 42

tgcttggtca ctagttctct tggtg 25

<210> SEQ ID NO 43

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 43

tgttacagga ggaaaagaca agg 23

<210> SEQ ID NO 44

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 44

tgttacagga ggaaaagaca aggag 25

<210> SEQ ID NO 45

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 45

gaggaaaaga caaggagctg agg 23

<210> SEQ ID NO 46

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 46

agacaaggag ctgaggctaa cgg 23

<210> SEQ ID NO 47

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 47

gacaaggagc tgaggctaac ggg 23

<210> SEQ ID NO 48

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 48

gacaaggagc tgaggctaac gggtg 25

<210> SEQ ID NO 49

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 49

aaggagctga ggctaacggg tgg 23

<210> SEQ ID NO 50

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 50

aaggagctga ggctaacggg tggtg 25

<210> SEQ ID NO 51

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 51

ggtggtgaaa acaagtgtc tgg 23

<210> SEQ ID NO 52

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 52

aaacaagtgc tctggaagag tgg 23

<210> SEQ ID NO 53

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 53

aaacaagtgc tctggaagag tggag 25

<210> SEQ ID NO 54

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 54

caagtgtctt ggaagagtgg agg 23

<210> SEQ ID NO 55

<211> LENGTH: 25

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 55

caagtgcctct ggaagagtgg aggtg 25

<210> SEQ ID NO 56

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 56

aagagtggag gtgaaagtgc agg 23

<210> SEQ ID NO 57

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 57

aagagtggag gtgaaagtgc aggag 25

<210> SEQ ID NO 58

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 58

agtgagggtg aaagtgcagg agg 23

<210> SEQ ID NO 59

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 59

agtgagggtg aaagtgcagg aggag 25

<210> SEQ ID NO 60

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 60

aggtgaaagt gcaggaggag tgg 23

<210> SEQ ID NO 61

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 61

aggtgaaagt gcaggaggag tgggg 25

<210> SEQ ID NO 62

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 62

ggtgaaagtg caggaggagt ggg 23

<210> SEQ ID NO 63

<211> LENGTH: 23

-continued

<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 63

gtgaaagtgc aggaggagtg ggg 23

<210> SEQ ID NO 64
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 64

tggggaactg tgtgtaataa tgg 23

<210> SEQ ID NO 65
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 65

gaactgtgtg taataatggc tgg 23

<210> SEQ ID NO 66
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 66

aactgtgtgt aataatggct ggg 23

<210> SEQ ID NO 67
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 67

gtgtaataat ggctgggaca tgg 23

<210> SEQ ID NO 68
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 68

taatggctgg gacatggatg tgg 23

<210> SEQ ID NO 69
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 69

atgtgggtctc tgttgtttgt agg 23

<210> SEQ ID NO 70
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 70

ctctgttggt ttaggcagc tgg 23

<210> SEQ ID NO 71

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 71

tctgttggtt gtaggcagct ggg 23

<210> SEQ ID NO 72
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 72

ccagtggctt tgatagcagt tgg 23

<210> SEQ ID NO 73
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 73

ccaactgcta tcaaagccac tgg 23

<210> SEQ ID NO 74
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 74

ctgctatcaa agccactgga tgg 23

<210> SEQ ID NO 75
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 75

tgctatcaaa gccactggat ggg 23

<210> SEQ ID NO 76
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 76

actaaaatta gcccatccag tgg 23

<210> SEQ ID NO 77
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 77

ggatgggcta attttagtgc agg 23

<210> SEQ ID NO 78
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 78

gctaatttta gtgcagggttc tgg 23

-continued

<210> SEQ ID NO 79
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 79

gtgcagggttc tggacgcatt tgg 23

<210> SEQ ID NO 80
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 80

agggttctgga cgcatttgga tgg 23

<210> SEQ ID NO 81
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 81

atggatcatg tttcttgctg agg 23

<210> SEQ ID NO 82
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 82

tggatcatgt ttcttgctga ggg 23

<210> SEQ ID NO 83
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 83

gagggaatga gtcagctctc tgg 23

<210> SEQ ID NO 84
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 84

agggaatgag tcagctctct ggg 23

<210> SEQ ID NO 85
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 85

ctctgggact gcaaacatga tgg 23

<210> SEQ ID NO 86
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 86

gggactgcaa acatgatgga tgg 23

-continued

<210> SEQ ID NO 87
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 87

gggactgcaa acatgatgga tgggg 25

<210> SEQ ID NO 88
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 88

ggactgcaaa catgatggat ggg 23

<210> SEQ ID NO 89
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 89

gactgcaaac atgatggatg ggg 23

<210> SEQ ID NO 90
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 90

gcataactgt actcaccaac agg 23

<210> SEQ ID NO 91
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 91

tgtactcacc aacaggatgc tgg 23

<210> SEQ ID NO 92
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 92

tgtactcacc aacaggatgc tggag 25

<210> SEQ ID NO 93
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 93

agggttactcc agcatcctgt tggtg 25

<210> SEQ ID NO 94
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 94

agggttactcc agcatcctgt tgg 23

-continued

<210> SEQ ID NO 95
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 95

gatgctggag taacctgctc agg 23

<210> SEQ ID NO 96
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 96

tgtgtatgtc ttacctgagc agg 23

<210> SEQ ID NO 97
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 97

atggatctga tttagagatg agg 23

<210> SEQ ID NO 98
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 98

atctgattta gagatgaggc tgg 23

<210> SEQ ID NO 99
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 99

atctgattta gagatgaggc tggtg 25

<210> SEQ ID NO 100
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 100

ttagagatga ggctgggtgaa tgg 23

<210> SEQ ID NO 101
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 101

ttagagatga ggctgggtgaa tggag 25

<210> SEQ ID NO 102
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 102

-continued

gagatgaggc tggatgaatgg agg 23

<210> SEQ ID NO 103
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 103

ggctgggtgaa tggaggaaac cgg 23

<210> SEQ ID NO 104
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 104

ggctgggtgaa tggaggaaac cggtg 25

<210> SEQ ID NO 105
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 105

aatggaggaa accggtgctt agg 23

<210> SEQ ID NO 106
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 106

cttctattct tcctaagcac cgg 23

<210> SEQ ID NO 107
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 107

agaatagaag tcaaatttca agg 23

<210> SEQ ID NO 108
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 108

tagaagtcaa atttcaagga cgg 23

<210> SEQ ID NO 109
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 109

tagaagtcaa atttcaagga cggtg 25

<210> SEQ ID NO 110
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 110

-continued

aagtcaaatt tcaaggacgg tgg	23
 <210> SEQ ID NO 111 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 111	
aagtcaaatt tcaaggacgg tgggg	25
 <210> SEQ ID NO 112 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 112	
agtcaaattt caaggacggt ggg	23
 <210> SEQ ID NO 113 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 113	
gtcaaatttc aaggacggtg ggg	23
 <210> SEQ ID NO 114 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 114	
caacataaat catgcttctg tgg	23
 <210> SEQ ID NO 115 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 115	
ggaagtgtg tcagtttctc tgg	23
 <210> SEQ ID NO 116 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 116	
ttctctgggt cagctaattt tgg	23
 <210> SEQ ID NO 117 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 117	
ttctctgggt cagctaattt tggag	25
 <210> SEQ ID NO 118 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa	

-continued

<400> SEQUENCE: 118

ggttcagcta attttggaaggagg 23

<210> SEQ ID NO 119

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 119

gctaattttg gagaagggttc tgg 23

<210> SEQ ID NO 120

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 120

gagaagggttc tggaccaatc tgg 23

<210> SEQ ID NO 121

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 121

acaagatcat caaaccagat tgg 23

<210> SEQ ID NO 122

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 122

tttgatgatc ttgtatgcaa tgg 23

<210> SEQ ID NO 123

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 123

atggaaatga gtcagctctc tgg 23

<210> SEQ ID NO 124

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 124

ctctggaact gcaaacatga agg 23

<210> SEQ ID NO 125

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 125

ggaactgcaa acatgaagga tgg 23

<210> SEQ ID NO 126

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 126

ggaactgcaa acatgaagga tgggg

25

<210> SEQ ID NO 127

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 127

gaactgcaaa catgaaggat ggg

23

<210> SEQ ID NO 128

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 128

aactgcaaac atgaaggatg ggg

23

<210> SEQ ID NO 129

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 129

gcacaattgt gatcatgctg agg

23

<210> SEQ ID NO 130

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 130

tgtgatcatg ctgaggatgc tgg

23

<210> SEQ ID NO 131

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 131

tgtgatcatg ctgaggatgc tggag

25

<210> SEQ ID NO 132

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 132

agcagacctg aaactgagag tgg

23

<210> SEQ ID NO 133

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 133

catctaccac tctcagtttc agg

23

<210> SEQ ID NO 134

<211> LENGTH: 23

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 134

ctgaaactga gagtggtaga tgg 23

<210> SEQ ID NO 135
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 135

ctgaaactga gagtggtaga tggag 25

<210> SEQ ID NO 136
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 136

gatggagtca ctgaatgttc agg 23

<210> SEQ ID NO 137
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 137

cactgaatgt tcaggaagat tgg 23

<210> SEQ ID NO 138
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 138

agattggaag tgaaattcca agg 23

<210> SEQ ID NO 139
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 139

agattggaag tgaaattcca aggag 25

<210> SEQ ID NO 140
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 140

aagtgaatt ccaaggagaa tgg 23

<210> SEQ ID NO 141
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 141

aagtgaatt ccaaggagaa tgggg 25

<210> SEQ ID NO 142
<211> LENGTH: 23

-continued

<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 142

agtgaaattc caaggagaat ggg 23

<210> SEQ ID NO 143
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 143

gtgaaattcc aaggagaatg ggg 23

<210> SEQ ID NO 144
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 144

agattgttcc ccattctect tgg 23

<210> SEQ ID NO 145
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 145

tggggaacaa tctgtgatga tgg 23

<210> SEQ ID NO 146
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 146

gaacaatctg tgatgatggc tgg 23

<210> SEQ ID NO 147
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 147

aacaatctgt gatgatggct ggg 23

<210> SEQ ID NO 148
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 148

ggatagtgat gatgccgtg tgg 23

<210> SEQ ID NO 149
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 149

cagttgctta catgccacag cgg 23

<210> SEQ ID NO 150

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 150

cgctgtggca tgtaagcaac tgg 23

<210> SEQ ID NO 151
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 151

gctgtggcat gtaagcaact ggg 23

<210> SEQ ID NO 152
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 152

ccaatggcag tgacagcagt tgg 23

<210> SEQ ID NO 153
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 153

ccaactgctg tcaactgccat tgg 23

<210> SEQ ID NO 154
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 154

actggcggtta actcgaccaa tgg 23

<210> SEQ ID NO 155
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 155

tggtcgagtt aacgccagtg agg 23

<210> SEQ ID NO 156
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 156

ggtcgagtta acgccagtga ggg 23

<210> SEQ ID NO 157
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 157

gttaacgcca gtgagggaaac tgg 23

-continued

<210> SEQ ID NO 158
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 158

aatgtgtcca gttccctcac tggcg 25

<210> SEQ ID NO 159
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 159

aatgtgtcca gttccctcac tgg 23

<210> SEQ ID NO 160
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 160

gtgaggggaac tggacacatt tgg 23

<210> SEQ ID NO 161
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 161

cttgacagtg tttcttgcca tgg 23

<210> SEQ ID NO 162
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 162

agagagcaga ctcgtgtcca tgg 23

<210> SEQ ID NO 163
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 163

atggacacga gtctgtcttc tgg 23

<210> SEQ ID NO 164
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 164

ggcagtgtag acaccatgaa tgg 23

<210> SEQ ID NO 165
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 165

ggcagtgtag acaccatgaa tgggg 25

-continued

<210> SEQ ID NO 166
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 166

gcagtgtaga caccatgaat ggg 23

<210> SEQ ID NO 167
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 167

cagtgtagac accatgaatg ggg 23

<210> SEQ ID NO 168
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 168

aataatgctt tccccattca tgggtg 25

<210> SEQ ID NO 169
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 169

aataatgctt tccccattca tgg 23

<210> SEQ ID NO 170
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 170

tgcaatcata atgaagatgc tgg 23

<210> SEQ ID NO 171
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 171

tgcaatcata atgaagatgc tgggtg 25

<210> SEQ ID NO 172
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 172

gatgctggtg tgacatgttc tgg 23

<210> SEQ ID NO 173
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 173

ttttgcagat ggatcagatc tgg 23

-continued

<210> SEQ ID NO 174
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 174

gatctggaac tgagacttaa agg 23

<210> SEQ ID NO 175
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 175

gatctggaac tgagacttaa aggtg 25

<210> SEQ ID NO 176
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 176

ctggaactga gacttaaagg tgg 23

<210> SEQ ID NO 177
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 177

ctggaactga gacttaaagg tggag 25

<210> SEQ ID NO 178
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 178

gaactgagac ttaaagggtgg agg 23

<210> SEQ ID NO 179
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 179

cagccactgt gctgggacag tgg 23

<210> SEQ ID NO 180
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 180

cagccactgt gctgggacag tggag 25

<210> SEQ ID NO 181
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 181

-continued

cctccactgt ccagcacag tgg 23

<210> SEQ ID NO 182
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 182

ccactgtgct gggacagtgg agg 23

<210> SEQ ID NO 183
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 183

ccactgtgct gggacagtgg aggtg 25

<210> SEQ ID NO 184
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 184

ctgtgctggg acagtggagg tgg 23

<210> SEQ ID NO 185
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 185

ggaggtggaa attcagaaac tgg 23

<210> SEQ ID NO 186
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 186

gtggaaattc agaaactggt agg 23

<210> SEQ ID NO 187
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 187

gaaaagtgtg tgatagaagc tgg 23

<210> SEQ ID NO 188
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 188

gaaaagtgtg tgatagaagc tgggg 25

<210> SEQ ID NO 189
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 189

-continued

aaaagtgtgt gatagaagct ggg	23
 <210> SEQ ID NO 190 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 190	
aaagtgtgtg atagaagctg ggg	23
 <210> SEQ ID NO 191 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 191	
gggactgaaa gaagctgatg tgg	23
 <210> SEQ ID NO 192 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 192	
aagaagctga tgtggtttgc agg	23
 <210> SEQ ID NO 193 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 193	
tgatgtggtt tgcaggcagc tgg	23
 <210> SEQ ID NO 194 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 194	
gatgtggttt gcaggcagct ggg	23
 <210> SEQ ID NO 195 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 195	
gtttgcaggc agctgggatg tgg	23
 <210> SEQ ID NO 196 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 196	
tcaagtttat tccaaaacca agg	23
 <210> SEQ ID NO 197 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa	

-continued

<400> SEQUENCE: 197

aaaccaaggc aacaaacaca tgg 23

<210> SEQ ID NO 198

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 198

cagccatgtg tttgttgct tgg 23

<210> SEQ ID NO 199

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 199

ctgtttgtaa gcagctgtaa tgg 23

<210> SEQ ID NO 200

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 200

atggaaatga aacttctctt tgg 23

<210> SEQ ID NO 201

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 201

tggaaatgaa acttctcttt ggg 23

<210> SEQ ID NO 202

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 202

ctctttggga ctgcaagaat tgg 23

<210> SEQ ID NO 203

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 203

gggactgcaa gaattggcag tgg 23

<210> SEQ ID NO 204

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 204

gggactgcaa gaattggcag tgggg 25

<210> SEQ ID NO 205

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 205

ggactgcaag aattggcagt ggg 23

<210> SEQ ID NO 206

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 206

gactgcaaga attggcagtg ggg 23

<210> SEQ ID NO 207

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 207

gactgcaaga attggcagtg ggggtg 25

<210> SEQ ID NO 208

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 208

tgcaagaatt ggcagtgggg tgg 23

<210> SEQ ID NO 209

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 209

cttacctgag caggtaattt tgg 23

<210> SEQ ID NO 210

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 210

ggtcgtgttg aagtacaaca tgg 23

<210> SEQ ID NO 211

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 211

ggaactacag tgcggcactg tgg 23

<210> SEQ ID NO 212

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 212

agaggcaatt caatttactt ggg 23

<210> SEQ ID NO 213

<211> LENGTH: 23

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 213

gagaggcaat tcaatttact tgg 23

<210> SEQ ID NO 214

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 214

gtttctttta acagactgag agg 23

<210> SEQ ID NO 215

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 215

ggttaagagt acaatcatca agg 23

<210> SEQ ID NO 216

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 216

tgatgattgt actcttaacc tgg 23

<210> SEQ ID NO 217

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 217

cagaaataaa gcagaagaca tgg 23

<210> SEQ ID NO 218

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 218

catgtcttct gctttatttc tgg 23

<210> SEQ ID NO 219

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 219

ctccagatta cagtaaatgg agg 23

<210> SEQ ID NO 220

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 220

gtctctcatt tactgtaate tgg 23

<210> SEQ ID NO 221

<211> LENGTH: 23

-continued

<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 221

gtaaatggag gactgagtat agg 23

<210> SEQ ID NO 222
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 222

taaatggagg actgagtata ggg 23

<210> SEQ ID NO 223
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 223

gggctaataaa gtagagagaa tgg 23

<210> SEQ ID NO 224
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 224

gaatggatgc atattatctg tgg 23

<210> SEQ ID NO 225
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 225

tccaatgtga tgaatgaagt agg 23

<210> SEQ ID NO 226
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 226

gcctacttca ttcacacat tgg 23

<210> SEQ ID NO 227
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 227

tgaagtaggc aaatactcaa agg 23

<210> SEQ ID NO 228
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 228

aaagcatgct ccaagaatta tgg 23

<210> SEQ ID NO 229

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 229

aagcatgctc caagaattat ggg 23

<210> SEQ ID NO 230
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 230

cttctggaac ccataattct tgg 23

<210> SEQ ID NO 231
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 231

caagaattat gggttccaga agg 23

<210> SEQ ID NO 232
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 232

aaagtcccag aattgtctcc agg 23

<210> SEQ ID NO 233
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 233

aagtcccaga attgtctcca ggg 23

<210> SEQ ID NO 234
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 234

ccttcctgg agacaattct ggg 23

<210> SEQ ID NO 235
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 235

cccagaattg tctccaggga agg 23

<210> SEQ ID NO 236
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 236

tccttcctg gagacaattc tgg 23

-continued

<210> SEQ ID NO 237
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 237

attgtctcca gggaaggaca ggg 23

<210> SEQ ID NO 238
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 238

gtctccaggg aaggacaggg agg 23

<210> SEQ ID NO 239
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 239

tagacctccc tgccttccc tgg 23

<210> SEQ ID NO 240
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 240

aggacaggga ggtctagaat cgg 23

<210> SEQ ID NO 241
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 241

gaatcggcta agccactgt agg 23

<210> SEQ ID NO 242
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 242

ttggtttttc tgcctacagt ggg 23

<210> SEQ ID NO 243
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 243

cttggttttt ctgcctacag tgg 23

<210> SEQ ID NO 244
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 244

ctgtaggcag aaaaaccaag agg 23

-continued

<210> SEQ ID NO 245
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 245

gaaaaaccaa gaggcataaa tgg 23

<210> SEQ ID NO 246
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 246

gggaagccat tcatgcctct tgg 23

<210> SEQ ID NO 247
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 247

agagagttaa aagtgagaaa ggg 23

<210> SEQ ID NO 248
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 248

cagagagtga aaagtgagaa agg 23

<210> SEQ ID NO 249
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 249

ctggcttact cctatcatga agg 23

<210> SEQ ID NO 250
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 250

cctatcatga agggaaatat tgg 23

<210> SEQ ID NO 251
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 251

ccaatatattt ccttcataat agg 23

<210> SEQ ID NO 252
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 252

aaaaaatagc atttcggtga ggg 23

-continued

<210> SEQ ID NO 253
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 253

gaaaaaatag catttcggtg agg 23

<210> SEQ ID NO 254
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 254

acaaagtga caccattccct ggg 23

<210> SEQ ID NO 255
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 255

aacaaagtga acacattccc tgg 23

<210> SEQ ID NO 256
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 256

tcccatgccca tgaagagggt agg 23

<210> SEQ ID NO 257
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 257

ccctaccctc ttcattggcat ggg 23

<210> SEQ ID NO 258
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 258

cccatgccat gaagagggtg ggg 23

<210> SEQ ID NO 259
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 259

accctaccct cttcatggca tgg 23

<210> SEQ ID NO 260
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 260

-continued

gccatgaaga gggtagggtt agg 23

<210> SEQ ID NO 261
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 261

acctaaccct accctcttca tgg 23

<210> SEQ ID NO 262
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 262

cagaaataaa gcagaagaca tggag 25

<210> SEQ ID NO 263
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 263

catgtcttct gctttatttc tggtg 25

<210> SEQ ID NO 264
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 264

tgactccaga ttacagtaaa tggag 25

<210> SEQ ID NO 265
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 265

gtcctccatt tactgtaac tggag 25

<210> SEQ ID NO 266
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 266

gcctacttca ttcacacat tggag 25

<210> SEQ ID NO 267
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 267

cttctggaac ccataattct tggag 25

<210> SEQ ID NO 268
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 268

-continued

attgtctcca gggaaggaca gggag	25
 <210> SEQ ID NO 269 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 269	
tagacctccc tgctcttccc tggag	25
 <210> SEQ ID NO 270 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 270	
ccaatatttt ccttcgatgat aggag	25
 <210> SEQ ID NO 271 <211> LENGTH: 25 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 271	
aaaaaatagc atttcggtga gggag	25
 <210> SEQ ID NO 272 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 272	
gaatcggcta agccactgt agg	23
 <210> SEQ ID NO 273 <211> LENGTH: 23 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 273	
cccatgccat gaagagggtg ggg	23
 <210> SEQ ID NO 274 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 274	
aagagaacta gtgaccaagc	20
 <210> SEQ ID NO 275 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Sus scrofa <400> SEQUENCE: 275	
tgettgtgca ctagtctct	20
 <210> SEQ ID NO 276 <211> LENGTH: 20 <212> TYPE: DNA <213> ORGANISM: Sus scrofa	

-continued

<400> SEQUENCE: 276

gaactgtgtg taataatggc 20

<210> SEQ ID NO 277

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 277

tgctatcaaa gccactggat 20

<210> SEQ ID NO 278

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 278

gtgcaggttc tggacgcatt 20

<210> SEQ ID NO 279

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 279

gtgcaggttc tggacgcatt 20

<210> SEQ ID NO 280

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 280

atggatcatg tttcttgctg 20

<210> SEQ ID NO 281

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 281

ggactgcaaa catgatggat 20

<210> SEQ ID NO 282

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 282

ggactgcaaa catgatggat 20

<210> SEQ ID NO 283

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 283

agtcaaattt caaggacggt 20

<210> SEQ ID NO 284

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 284

gctaattttg gagaagggttc

20

<210> SEQ ID NO 285

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 285

gcacaattgt gatcatgctg

20

<210> SEQ ID NO 286

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 286

ggaactacag tgcggcactg

20

<210> SEQ ID NO 287

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 287

aagagaacua gugaccaagc

20

<210> SEQ ID NO 288

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 288

ugcuugguca cuaguucucu

20

<210> SEQ ID NO 289

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 289

gaacugugug uaauaauggc

20

<210> SEQ ID NO 290

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 290

ugcuaucaaa gccacuggau

20

<210> SEQ ID NO 291

<211> LENGTH: 20

<212> TYPE: RNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 291

gugcagguuc uggacgcauu

20

<210> SEQ ID NO 292

<211> LENGTH: 20

<212> TYPE: RNA

-continued

<213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 292
 gugcagguuc uggaogcauu 20

 <210> SEQ ID NO 293
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 293
 auggaucaug uuucuugucg 20

 <210> SEQ ID NO 294
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 294
 ggacugcaaa caugauggau 20

 <210> SEQ ID NO 295
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 295
 ggacugcaaa caugauggau 20

 <210> SEQ ID NO 296
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 296
 agucaaaauu caaggacggu 20

 <210> SEQ ID NO 297
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 297
 gcuaauuuug gagaagguuc 20

 <210> SEQ ID NO 298
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 298
 gcacaaugu gaucaugcug 20

 <210> SEQ ID NO 299
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: *Sus scrofa*
 <400> SEQUENCE: 299
 ggaacuacag ugcggcacug 20

 <210> SEQ ID NO 300
 <211> LENGTH: 157

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 300

tttcgcagac tttagaagat gttctgccca tttaagttcc ttcacttttg ctgtagtcgc      60
tgttctcagt gcctgcttgg tcactagttc tcttggtgag tactttgaca aatttacttg      120
taacctagcc cactgtgaca agaaacactg aaaagca                                157

<210> SEQ ID NO 301
<211> LENGTH: 157
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 301

tttcgcagac tttagaagat gttctgccca tttaagttcc ttcacttttg ctgtagtcgc      60
tgttctcagt gcctgcttgg tcactagttc tcttggtgag tactttgaca aatttacttg      120
taacctagcc cactgtgaca agaaacactg aaaagca                                157

<210> SEQ ID NO 302
<211> LENGTH: 152
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 302

gtggtgaaaa caagtgtctt ggaagagtgg aggtgaaagt gcaggaggag tggggaactg      60
tgtgtaataa tggctgggac atggatgtgg tctctgttgt ttgtaggcag ctgggatgtc      120
caactgctat caaagccact ggatgggcta at                                152

<210> SEQ ID NO 303
<211> LENGTH: 154
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 303

tggctgggac atggatgtgg tctctgttgt ttgtaggcag ctgggatgtc caactgctat      60
caaagccact ggatgggcta attttagtgc aggttctgga cgcatttgga tggatcatgt      120
ttcttgtcga gggaaatgagt cagctctctg ggac                                154

<210> SEQ ID NO 304
<211> LENGTH: 151
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 304

gtaggcagct gggatgtcca actgctatca aagccactgg atgggctaatt ttagtgacag      60
gttctggacg catttgatg gatcatgttt cttgtcgagg gaatgagtc gctctctggg      120
actgcaaaca tgatggatgg ggaagcata a                                151

<210> SEQ ID NO 305
<211> LENGTH: 151
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 305

gtaggcagct gggatgtcca actgctatca aagccactgg atgggctaatt ttagtgacag      60
gttctggacg catttgatg gatcatgttt cttgtcgagg gaatgagtc gctctctggg      120

```

-continued

actgcaaaca tgatggatgg ggaaagcata a 151

<210> SEQ ID NO 306
 <211> LENGTH: 156
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 306

gtccaactgc tatcaaagcc actggatggg ctaatttttag tgcaggttct ggacgcattt 60
 ggatggatca tgtttcttgt cgaggggaatg agtcagctct ctgggactgc aaacatgatg 120
 gatggggaaa gcataactgt actcaccaac aggatg 156

<210> SEQ ID NO 307
 <211> LENGTH: 153
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 307

ctggacgcat ttggatggat catgtttctt gtcgaggga tgagtcagct ctctgggact 60
 gcaaacatga tggatgggga aagcataact gtactcacca acaggatgct ggagtaacct 120
 gctcaggtaa gacatacaca aataagtcaa gcc 153

<210> SEQ ID NO 308
 <211> LENGTH: 153
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 308

ctggacgcat ttggatggat catgtttctt gtcgaggga tgagtcagct ctctgggact 60
 gcaaacatga tggatgggga aagcataact gtactcacca acaggatgct ggagtaacct 120
 gctcaggtaa gacatacaca aataagtcaa gcc 153

<210> SEQ ID NO 309
 <211> LENGTH: 153
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 309

ctctagatgg atctgattta gagatgaggc tggatgaatgg aggaaccgg tgetttaggaa 60
 gaatagaagt caaatattcaa ggacggtggg gaacagtgtg tgatgataac ttcaacataa 120
 atcatgcttc tgtggtttgt aaacaacttg aat 153

<210> SEQ ID NO 310
 <211> LENGTH: 153
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 310

cttctgtggt ttgtaaacaa cttgaatgtg gaagtgtgt cagtttctct gggtcagcta 60
 attttgaga aggttctgga ccaatctggt ttgatgatct tgtatgcaat ggaaatgagt 120
 cagctctctg gaactgcaaa catgaaggat ggg 153

<210> SEQ ID NO 311
 <211> LENGTH: 155
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 311

-continued

```

tgatgatctt gtatgcaatg gaaatgagtc agctctctgg aactgcaaac atgaaggatg    60
gggaaagcac aattgcatc atgctgagga tgetggagtg atttgettaa gtaaggactg    120
acctgggttt gttctgttct ccatgagagg gcaaaa                                155

```

```

<210> SEQ ID NO 312
<211> LENGTH: 154
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 312

```

```

agacacgtgg ggcaccgtct gtgattctga cttctctctg gaggcggcca gcgtgctgtg    60
cagggaaacta cagtgcggca ctgtggtttc cctcctgggg ggagctcact ttggagaagg    120
aagtggacag atctgggctg aagaattcca gtgt                                154

```

```

<210> SEQ ID NO 313
<211> LENGTH: 28
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 313

```

```

Phe Arg Arg Cys Ser Ala His Leu Ser Ser Phe Thr Phe Ala Val Val
1         5             10             15
Ala Val Leu Ser Ala Cys Leu Val Thr Ser Ser Leu
                20             25

```

```

<210> SEQ ID NO 314
<211> LENGTH: 28
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 314

```

```

Phe Arg Arg Cys Ser Ala His Leu Ser Ser Phe Thr Phe Ala Val Val
1         5             10             15
Ala Val Leu Ser Ala Cys Leu Val Thr Ser Ser Leu
                20             25

```

```

<210> SEQ ID NO 315
<211> LENGTH: 50
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 315

```

```

Gly Glu Asn Lys Cys Ser Gly Arg Val Glu Val Lys Val Gln Glu Glu
1         5             10             15
Trp Gly Thr Val Cys Asn Asn Gly Trp Asp Met Asp Val Val Ser Val
                20             25             30
Val Cys Arg Gln Leu Gly Cys Pro Thr Ala Ile Lys Ala Thr Gly Trp
                35             40             45
Ala Asn
50

```

```

<210> SEQ ID NO 316
<211> LENGTH: 51
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 316

```

```

Gly Trp Asp Met Asp Val Val Ser Val Val Cys Arg Gln Leu Gly Cys
1         5             10             15

```

-continued

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
20 25 30

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
35 40 45

Leu Trp Asp
50

<210> SEQ ID NO 317

<211> LENGTH: 44

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 317

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
1 5 10 15

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
20 25 30

Leu Trp Asp Cys Lys His Asp Gly Trp Gly Lys His
35 40

<210> SEQ ID NO 318

<211> LENGTH: 44

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 318

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
1 5 10 15

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
20 25 30

Leu Trp Asp Cys Lys His Asp Gly Trp Gly Lys His
35 40

<210> SEQ ID NO 319

<211> LENGTH: 51

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 319

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
1 5 10 15

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
20 25 30

Leu Trp Asp Cys Lys His Asp Gly Trp Gly Lys His Asn Cys Thr His
35 40 45

Gln Gln Asp
50

<210> SEQ ID NO 320

<211> LENGTH: 35

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 320

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
1 5 10 15

Leu Trp Asp Cys Lys His Asp Gly Trp Gly Lys His Asn Cys Thr His
20 25 30

Gln Gln Asp

-continued

35

<210> SEQ ID NO 321
 <211> LENGTH: 35
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 321

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
 1 5 10 15
 Leu Trp Asp Cys Lys His Asp Gly Trp Gly Lys His Asn Cys Thr His
 20 25 30
 Gln Gln Asp
 35

<210> SEQ ID NO 322
 <211> LENGTH: 48
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 322

Gly Ser Asp Leu Glu Met Arg Leu Val Asn Gly Gly Asn Arg Cys Leu
 1 5 10 15
 Gly Arg Ile Glu Val Lys Phe Gln Gly Arg Trp Gly Thr Val Cys Asp
 20 25 30
 Asp Asn Phe Asn Ile Asn His Ala Ser Val Val Cys Lys Gln Leu Glu
 35 40 45

<210> SEQ ID NO 323
 <211> LENGTH: 50
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 323

Ser Val Val Cys Lys Gln Leu Glu Cys Gly Ser Ala Val Ser Phe Ser
 1 5 10 15
 Gly Ser Ala Asn Phe Gly Glu Gly Ser Gly Pro Ile Trp Phe Asp Asp
 20 25 30
 Leu Val Cys Asn Gly Asn Glu Ser Ala Leu Trp Asn Cys Lys His Glu
 35 40 45
 Gly Trp
 50

<210> SEQ ID NO 324
 <211> LENGTH: 36
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 324

Asp Asp Leu Val Cys Asn Gly Asn Glu Ser Ala Leu Trp Asn Cys Lys
 1 5 10 15
 His Glu Gly Trp Gly Lys His Asn Cys Asp His Ala Glu Asp Ala Gly
 20 25 30
 Val Ile Cys Leu
 35

<210> SEQ ID NO 325
 <211> LENGTH: 51
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 325

Asp Thr Trp Gly Thr Val Cys Asp Ser Asp Phe Ser Leu Glu Ala Ala
 1 5 10 15
 Ser Val Leu Cys Arg Glu Leu Gln Cys Gly Thr Val Val Ser Leu Leu
 20 25 30
 Gly Gly Ala His Phe Gly Glu Gly Ser Gly Gln Ile Trp Ala Glu Glu
 35 40 45
 Phe Gln Cys
 50

<210> SEQ ID NO 326

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 326

Phe Arg Arg Cys Ser Ala His Leu Ser Ser Phe Thr Phe Ala Val Val
 1 5 10 15
 Ala Val Leu Ser Ala
 20

<210> SEQ ID NO 327

<211> LENGTH: 21

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 327

Phe Arg Arg Cys Ser Ala His Leu Ser Ser Phe Thr Phe Ala Val Val
 1 5 10 15
 Ala Val Leu Ser Ala
 20

<210> SEQ ID NO 328

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 328

Gly Glu Asn Lys Cys Ser Gly Arg Val Glu Val Lys Val Gln Glu Glu
 1 5 10 15
 Trp Gly Thr Val Cys Asn Asn Gly
 20

<210> SEQ ID NO 329

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 329

Gly Trp Asp Met Asp Val Val Ser Val Val Cys Arg Gln Leu Gly Cys
 1 5 10 15
 Pro Thr Ala Ile Lys Ala Thr Gly
 20

<210> SEQ ID NO 330

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 330

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser

-continued

1 5 10 15

Gly Arg Ile

<210> SEQ ID NO 331
 <211> LENGTH: 19
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 331

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
 1 5 10 15

Gly Arg Ile

<210> SEQ ID NO 332
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 332

Pro Thr Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser
 1 5 10 15

Gly Arg Ile Trp Met Asp His Val
 20

<210> SEQ ID NO 333
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 333

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
 1 5 10 15

Leu Trp Asp Cys Lys His Asp Gly
 20

<210> SEQ ID NO 334
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 334

Gly Arg Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala
 1 5 10 15

Leu Trp Asp Cys Lys His Asp Gly
 20

<210> SEQ ID NO 335
 <211> LENGTH: 22
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 335

Gly Ser Asp Leu Glu Met Arg Leu Val Asn Gly Gly Asn Arg Cys Leu
 1 5 10 15

Gly Arg Ile Glu Val Lys
 20

<210> SEQ ID NO 336
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 336

Ser Val Val Cys Lys Gln Leu Glu Cys Gly Ser Ala Val Ser Phe Ser
1 5 10 15
Gly Ser Ala Asn Phe Gly Glu Gly
20

<210> SEQ ID NO 337

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 337

Asp Asp Leu Val Cys Asn Gly Asn Glu Ser Ala Leu Trp Asn Cys Lys
1 5 10 15
His Glu Gly Trp Gly Lys His Asn
20

<210> SEQ ID NO 338

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 338

Asp Thr Trp Gly Thr Val Cys Asp Ser Asp Phe Ser Leu Glu Ala Ala
1 5 10 15
Ser Val Leu Cys Arg Glu Leu Gln
20

<210> SEQ ID NO 339

<211> LENGTH: 24

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 339

Val Ala Val Leu Ser Ala Cys Leu Val Thr Ser Ser Trp Arg Lys Arg
1 5 10 15
Gln Gly Ala Glu Ala Asn Gly Trp
20

<210> SEQ ID NO 340

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 340

Val Ala Val Leu Ser Ala Cys Leu Val Thr Cys Asn Leu Ala His Cys
1 5 10 15
Asp Lys Lys His
20

<210> SEQ ID NO 341

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 341

Val Ala Val Leu Ser Ala Cys Leu Val Thr Ser Ser Leu Leu Glu Glu
1 5 10 15
Lys Thr Arg Ser
20

-continued

<210> SEQ ID NO 342

<211> LENGTH: 14

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 342

Val Ala Val Leu Ser Ala Cys Leu Glu Glu Lys Thr Arg Ser
 1 5 10

<210> SEQ ID NO 343

<211> LENGTH: 20

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 343

Val Ala Val Leu Ser Ala Cys Leu Val Thr Arg Ser Thr Leu Thr Asn
 1 5 10 15

Leu Leu Val Thr
 20

<210> SEQ ID NO 344

<211> LENGTH: 33

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 344

agttctcttg gtgagtactt tgacaaattt act 33

<210> SEQ ID NO 345

<211> LENGTH: 16

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 345

cactagttct cttggt 16

<210> SEQ ID NO 346

<211> LENGTH: 10

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 346

ttctcttggt 10

<210> SEQ ID NO 347

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 347

atgctatttt ttcagccac agg 23

<210> SEQ ID NO 348

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 348

tttcagccca caggaaacc agg 23

<210> SEQ ID NO 349

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 349

agccacagga aaaccaggc tgg 23

<210> SEQ ID NO 350

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 350

aaccagcctg ggtttcctgt ggg 23

<210> SEQ ID NO 351

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 351

caaccagcct ggtttcctgt tgg 23

<210> SEQ ID NO 352

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 352

cacaggaaac ccaggctggt tgg 23

<210> SEQ ID NO 353

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 353

aggaaaccca ggtggttgg agg 23

<210> SEQ ID NO 354

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 354

ggaaaccag gctggttggg ggg 23

<210> SEQ ID NO 355

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 355

gaaaccagg ctggttggag ggg 23

<210> SEQ ID NO 356

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 356

atgtccctc caaccagcct ggg 23

<210> SEQ ID NO 357

<211> LENGTH: 23

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 357

aatgtccct ccaaccagcc tgg 23

<210> SEQ ID NO 358

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 358

ggaggggaca ttccctgctc tgg 23

<210> SEQ ID NO 359

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 359

acttcaacac gaccagagca ggg 23

<210> SEQ ID NO 360

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 360

tacttcaaca cgaccagagc agg 23

<210> SEQ ID NO 361

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 361

ggtcgtgttg aagtacaaca tgg 23

<210> SEQ ID NO 362

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 362

aagtacaaca tggagacacg tgg 23

<210> SEQ ID NO 363

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 363

agtacaacat ggagacacgt ggg 23

<210> SEQ ID NO 364

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 364

gtacaacatg gagacacgtg ggg 23

<210> SEQ ID NO 365

<211> LENGTH: 23

-continued

<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 365

agagaagtca gaatcacaga cgg 23

<210> SEQ ID NO 366
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 366

ctgtgattct gacttctctc tgg 23

<210> SEQ ID NO 367
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 367

tgattctgac ttctctctgg agg 23

<210> SEQ ID NO 368
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 368

ttctgacttc tctctggagg cgg 23

<210> SEQ ID NO 369
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 369

aggcggccag cgtgctgtgc agg 23

<210> SEQ ID NO 370
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 370

ggcggccagc gtgctgtgca ggg 23

<210> SEQ ID NO 371
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 371

tagttccctg cacagcacgc tgg 23

<210> SEQ ID NO 372
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 372

ctgtgcaggg aactacagtg cgg 23

<210> SEQ ID NO 373

-continued

<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 373

ggaactacag tgcggcactg tgg 23

<210> SEQ ID NO 374
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 374

eggcactgtg gtttccctcc tgg 23

<210> SEQ ID NO 375
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 375

ggcactgtgg tttccctcct ggg 23

<210> SEQ ID NO 376
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 376

gcactgtggt ttccctcctg ggg 23

<210> SEQ ID NO 377
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 377

cactgtgggt tccctcctgg ggg 23

<210> SEQ ID NO 378
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 378

actgtgggtt ccctcctggg ggg 23

<210> SEQ ID NO 379
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 379

aaagtgagct cccccagga ggg 23

<210> SEQ ID NO 380
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 380

caaagtgagc tccccccagg agg 23

-continued

<210> SEQ ID NO 381
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 381

ctcctggggg gagctcactt tgg 23

<210> SEQ ID NO 382
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 382

ctccaaagtg agctccccc agg 23

<210> SEQ ID NO 383
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 383

gggggagctc actttggaga agg 23

<210> SEQ ID NO 384
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 384

gctcactttg gagaaggaag tgg 23

<210> SEQ ID NO 385
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 385

gagaaggaag tggacagatc tgg 23

<210> SEQ ID NO 386
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 386

agaaggaagt ggacagatct ggg 23

<210> SEQ ID NO 387
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 387

ggctgaagaa ttccagtgtg agg 23

<210> SEQ ID NO 388
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 388

gctgaagaat tccagtgtga ggg 23

-continued

<210> SEQ ID NO 389
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 389

ctgaagaatt ccagtgtgag ggg 23

<210> SEQ ID NO 390
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 390

gggactcgtg cccctcacac tgg 23

<210> SEQ ID NO 391
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 391

tactgggcag agtgaaaggt ggg 23

<210> SEQ ID NO 392
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 392

ctactgggca gaggaaagg tgg 23

<210> SEQ ID NO 393
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 393

gtgctactgg gcagagtgaagg 23

<210> SEQ ID NO 394
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 394

cgtcaggcg ggggtgctact ggg 23

<210> SEQ ID NO 395
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 395

ccagtagcac cccgccctga cgg 23

<210> SEQ ID NO 396
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 396

ccgtcaggc ggggtgctac tgg 23

-continued

<210> SEQ ID NO 397
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 397

cagtagcacc ccgccctgac ggg 23

<210> SEQ ID NO 398
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 398

ctacatgtcc cgtcaggcg ggg 23

<210> SEQ ID NO 399
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 399

gctacatgtc ccgtcagggc ggg 23

<210> SEQ ID NO 400
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 400

ggctacatgt cccgtcaggg cgg 23

<210> SEQ ID NO 401
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 401

tgtggctaca tgtcccgta ggg 23

<210> SEQ ID NO 402
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 402

ctgtgggtac atgtcccgta agg 23

<210> SEQ ID NO 403
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 403

acgggacatg tagccacagc agg 23

<210> SEQ ID NO 404
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 404

-continued

cgggacatgt agccacagca ggg 23

<210> SEQ ID NO 405
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 405

tgtagccaca gcagggacgt cgg 23

<210> SEQ ID NO 406
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 406

ctacgccgac gtcctgtgtg tgg 23

<210> SEQ ID NO 407
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 407

cacaggaaac ccagggtggt tggag 25

<210> SEQ ID NO 408
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 408

aggaaaccca ggctggttgg agggg 25

<210> SEQ ID NO 409
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 409

ggtcgtgttg aagtacaaca tggag 25

<210> SEQ ID NO 410
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 410

aagtacaaca tggagacacg tgggg 25

<210> SEQ ID NO 411
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 411

agagaagtca gaatcacaga cgggtg 25

<210> SEQ ID NO 412
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 412

-continued

ctgtgattct gacttctctc tggag 25

<210> SEQ ID NO 413
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 413

tgattctgac ttctctctgg aggcg 25

<210> SEQ ID NO 414
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 414

cggcactgtg gtttcctcc tgggg 25

<210> SEQ ID NO 415
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 415

ggcactgtgg tttccctect ggggg 25

<210> SEQ ID NO 416
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 416

gcactgtggt ttcctctctg ggggg 25

<210> SEQ ID NO 417
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 417

actgtggttt ccctctctggg gggag 25

<210> SEQ ID NO 418
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 418

ctctggggg gagctcactt tggag 25

<210> SEQ ID NO 419
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 419

ctccaaagtg agctccccc aggag 25

<210> SEQ ID NO 420
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 420

ggctgaagaa ttccagtgtg agggg 25

<210> SEQ ID NO 421
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 421

gtgctactgg gcagagtga aggtg 25

<210> SEQ ID NO 422
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 422

ctacatgtcc cgtcagggcg gggcg 25

<210> SEQ ID NO 423
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 423

ggctacatgt cccgtcaggg cgggg 25

<210> SEQ ID NO 424
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 424

tgtggctaca tgtcccgta gggcg 25

<210> SEQ ID NO 425
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 425

tgtagccaca gcagggacgt cggcg 25

<210> SEQ ID NO 426
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 426

gaatgaagta ggcaaatact caaaggaaag agaaagcatg ctccaagaat ggtaggggta 60

ggtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 427
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 427

atgaatgaag taggcaaata ctcaaaggaa agagaaagca tgctccaaga ggtaggggta 60

ggtagtcaca gacatctttt taaagccctg tctccttcca 100

-continued

```

<210> SEQ ID NO 428
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 428

ggcaaatact caaaggaaag agaaagcatg ctccaagaat tatgggttcc ggtaggggta      60
ggtagtcaca gacatctttt taaagccctg tctccttcca                             100

<210> SEQ ID NO 429
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 429

agaattatgg gttccagaag gcaaagtccc agaattgtct ccagggaagg ggtaggggta      60
ggtagtcaca gacatctttt taaagccctg tctccttcca                             100

<210> SEQ ID NO 430
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 430

agaattgtct ccagggaagg acagggaggt ctagaatcgg ctaagcccac ggtaggggta      60
ggtagtcaca gacatctttt taaagccctg tctccttcca                             100

<210> SEQ ID NO 431
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 431

gaatgaagta ggcaaatact caaaggaaag agaaagcatg ctccaagaat gtaggggttag      60
gtagtcacag acatcttttt aaagccctgt ctccttccag                             100

<210> SEQ ID NO 432
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 432

atgaatgaag taggcaaata ctcaaaggaa agagaaagca tgctccaaga gtaggggttag      60
gtagtcacag acatcttttt aaagccctgt ctccttccag                             100

<210> SEQ ID NO 433
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 433

ggcaaatact caaaggaaag agaaagcatg ctccaagaat tatgggttcc gtaggggttag      60
gtagtcacag acatcttttt aaagccctgt ctccttccag                             100

<210> SEQ ID NO 434
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 434

```

-continued

```

agaattatgg gttccagaag gcaaagtccc agaattgtct ccaggggaagg gtagggtagg      60
gtagtcacag acatcttttt aaagccctgt ctccttcag                               100

```

```

<210> SEQ ID NO 435
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 435

```

```

agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac gtagggtagg      60
gtagtcacag acatcttttt aaagccctgt ctccttcag                               100

```

```

<210> SEQ ID NO 436
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 436

```

```

gaatgaagta ggcaataact caaaggaaag agaaagcatg ctccaagaat agagggtagg      60
gttaggtagt cacagacatc tttttaaagc cctgtctcct                               100

```

```

<210> SEQ ID NO 437
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 437

```

```

atgaatgaag taggcaaata ctcaaaggaa agagaaagca tgctccaaga agagggtagg      60
gttaggtagt cacagacatc tttttaaagc cctgtctcct                               100

```

```

<210> SEQ ID NO 438
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 438

```

```

ggcaataact caaaggaaag agaaagcatg ctccaagaat tatgggttcc agagggtagg      60
gttaggtagt cacagacatc tttttaaagc cctgtctcct                               100

```

```

<210> SEQ ID NO 439
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 439

```

```

agaattatgg gttccagaag gcaaagtccc agaattgtct ccaggggaagg agagggtagg      60
gttaggtagt cacagacatc tttttaaagc cctgtctcct                               100

```

```

<210> SEQ ID NO 440
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

```

```

<400> SEQUENCE: 440

```

```

agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac agagggtagg      60
gttaggtagt cacagacatc tttttaaagc cctgtctcct                               100

```

```

<210> SEQ ID NO 441
<211> LENGTH: 100

```

-continued

<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 441

atgtcttctg ctttatttct ggtgtgcctt tgactccaga ttacagtaaa ggtaggggta 60
ggtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 442
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 442

ttatttctgg tgtgccttgg actccagatt acagtaaatg gaggactgag ggtaggggta 60
ggtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 443
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 443

tggaggactg agtatagggc taaaaagtag agagaatgga tgcattattat ggtaggggta 60
ggtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 444
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 444

atgcattatta tctgtgtgct ccaatgtgat gaatgaagta ggcaaatact ggtaggggta 60
ggtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 445
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 445

atgtcttctg ctttatttct ggtgtgcctt tgactccaga ttacagtaaa gtaggggtag 60
gtagtcacag acatcttttt aaagccctgt ctccttccag 100

<210> SEQ ID NO 446
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 446

ttatttctgg tgtgccttgg actccagatt acagtaaatg gaggactgag gtaggggtag 60
gtagtcacag acatcttttt aaagccctgt ctccttccag 100

<210> SEQ ID NO 447
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 447

tggaggactg agtatagggc taaaaagtag agagaatgga tgcattattat gtaggggtag 60

-continued

gtagtcacag acatcttttt aaagccctgt ctcttccag	100
--	-----

<210> SEQ ID NO 448
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 448

atgcatatta tctgtggtct ccaatgtgat gaatgaagta ggcaaatact gtagggtagg	60
---	----

gtagtcacag acatcttttt aaagccctgt ctcttccag	100
--	-----

<210> SEQ ID NO 449
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 449

atgtcttctg ctttatttct ggtgtgcctt tgactccaga ttacagtaaa agagggtagg	60
---	----

gttaggtagt cacagacatc tttttaaagc cctgtctcct	100
---	-----

<210> SEQ ID NO 450
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 450

ttatttctgg tgtgcctttg actccagatt acagtaaagtg gaggactgag agagggtagg	60
--	----

gttaggtagt cacagacatc tttttaaagc cctgtctcct	100
---	-----

<210> SEQ ID NO 451
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 451

tggaggactg agtatagggc taaaaagtag agagaatgga tgcattatag agagggtagg	60
---	----

gttaggtagt cacagacatc tttttaaagc cctgtctcct	100
---	-----

<210> SEQ ID NO 452
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 452

atgcatatta tctgtggtct ccaatgtgat gaatgaagta ggcaaatact agagggtagg	60
---	----

gttaggtagt cacagacatc tttttaaagc cctgtctcct	100
---	-----

<210> SEQ ID NO 453
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 453

catgaatggc ttccctttct cacttttcac tctctggctt actcctatca ggtaggggta	60
---	----

ggtagtcaca gacatctttt taaagccctg tctccttcca	100
---	-----

<210> SEQ ID NO 454
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 454

ttccctttct cacttttcac tctctggctt actcctatca tgaaggaaaa gtaggggta 60
gtagtcaca gacatctttt taaagccctg tctccttcca 100

<210> SEQ ID NO 455

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 455

catgaatggc ttccctttct cacttttcac tctctggctt actcctatca gtaggggtag 60
gtagtcacag acatcttttt aaagccctgt ctccttccag 100

<210> SEQ ID NO 456

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 456

ttccctttct cacttttcac tctctggctt actcctatca tgaaggaaaa gtaggggtag 60
gtagtcacag acatcttttt aaagccctgt ctccttccag 100

<210> SEQ ID NO 457

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 457

catgaatggc ttccctttct cacttttcac tctctggctt actcctatca agagggtagg 60
gttaggtagt cacagacatc tttttaaaag cctgtctcct 100

<210> SEQ ID NO 458

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 458

ttccctttct cacttttcac tctctggctt actcctatca tgaaggaaaa agagggtagg 60
gttaggtagt cacagacatc tttttaaaag cctgtctcct 100

<210> SEQ ID NO 459

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 459

tattggaatc atattctccc tcaccgaaat gctatttttt cagcccacag gtgattctga 60
cttctctctg gagggcgcca gcgtgctgtg cagggaacta 100

<210> SEQ ID NO 460

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 460

tattggaatc atattctccc tcaccgaaat gctatttttt cagcccacag gtgaggggca 60
cgagtccac ctttcactct gccagtagc accccgcctt 100

-continued

<210> SEQ ID NO 461
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 461

ggaatcatat tctccctcac cgaaatgcta ttttttcagc ccacaggaaa tgaggggcac 60
gagtcccacc tttcactctg cccagtagca ccccgcctg 100

<210> SEQ ID NO 462
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 462

ggaatcatat tctccctcac cgaaatgcta ttttttcagc ccacaggaaa tgaggggcac 60
gagtcccacc tttcactctg cccagtagca ccccgcctg 100

<210> SEQ ID NO 463
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 463

ggaatcatat tctccctcac cgaaatgcta ttttttcagc ccacaggaaa tgacgggaca 60
tgtagccaca gcagggaagt cggcgtagtc tgctcaagt 100

<210> SEQ ID NO 464
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 464

attctccctc accgaaatgc tattttttca gcccacagga aaccagggt gtgattctga 60
cttctctctg gagggggcca gcgtgctgtg cagggaacta 100

<210> SEQ ID NO 465
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 465

attctccctc accgaaatgc tattttttca gcccacagga aaccagggt gtgaggggca 60
cgagtccac ctttcaactct gccagtagc acccgcct 100

<210> SEQ ID NO 466
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 466

attctccctc accgaaatgc tattttttca gcccacagga aaccagggt ctgacgggac 60
atgtagccac agcagggaagc tcggcgtagt ctgctcaagt 100

<210> SEQ ID NO 467
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 467

-continued

ctccctcacc gaaatgctat tttttcagcc cacaggaaac ccaggctggt gtgattctga 60

cttctctctg gagggcgcca gcgtgctgtg cagggaacta 100

<210> SEQ ID NO 468

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 468

ctccctcacc gaaatgctat tttttcagcc cacaggaaac ccaggctggt gtgaggggca 60

cgagtccacc ctttactctt gccagtagc accccgcctt 100

<210> SEQ ID NO 469

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 469

ctccctcacc gaaatgctat tttttcagcc cacaggaaac ccaggctggt ctgacgggac 60

atgtagccac agcaggggacg tcggcgtagt ctgctcaagt 100

<210> SEQ ID NO 470

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 470

tccctcaccg aaatgctatt ttttcagccc acaggaaacc caggctgggt tgagggggcac 60

gagtccacc ttttactctg cccagtagca ccccgccctg 100

<210> SEQ ID NO 471

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 471

tccctcaccg aaatgctatt ttttcagccc acaggaaacc caggctgggt tgagggggcac 60

gagtccacc ttttactctg cccagtagca ccccgccctg 100

<210> SEQ ID NO 472

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 472

tccctcaccg aaatgctatt ttttcagccc acaggaaacc caggctgggt tgacgggaca 60

tgtagccaca gcaggggacgt cggcgtagtc tgctcaagtg 100

<210> SEQ ID NO 473

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 473

attttttcag cccacaggaa acccaggctg gttggagggg acattccctg acatggagac 60

acgtggggca ccgtctgtga ttctgacttc tctctggagg 100

<210> SEQ ID NO 474

-continued

<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 474

atTTTTtcag cccacaggaa acccaggctg gttggagggg acattccctg acgtggggca 60
ccgtctgtga ttctgacttc tctctggagg cggccagcgt 100

<210> SEQ ID NO 475
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 475

atTTTTtcag cccacaggaa acccaggctg gttggagggg acattccctg aggcggccag 60
cgtgctgtgc agggaactac agtgccggcac tgtggtttcc 100

<210> SEQ ID NO 476
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 476

atTTTTtcag cccacaggaa acccaggctg gttggagggg acattccctg aagtggacag 60
atctgggctg aagaattcca gtgtgagggg cacgagtccc 100

<210> SEQ ID NO 477
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 477

atTTTTtcag cccacaggaa acccaggctg gttggagggg acattccctg agcaccgccg 60
cctgacggga catgtagcca cagcaggggac gtcggcgtag 100

<210> SEQ ID NO 478
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 478

atTTTTtcag cccacaggaa acccaggctg gttggagggg acattccctg ctgacgggac 60
atgtagccac agcagggacg tcggcgtagt ctgctcaagt 100

<210> SEQ ID NO 479
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 479

ttttttcagc ccacaggaaa cccaggctgg ttggagggga cattccctgc tgagggggcac 60
gagtcccacc tttactctg cccagtagca ccccgcctg 100

<210> SEQ ID NO 480
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 480

ttttttcagc ccacaggaaa cccaggctgg ttggagggga cattccctgc tgagggggcac 60

-continued

gagtcccacc ttctactctg ccagtagca ccccgccctg 100

<210> SEQ ID NO 481
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 481

ttttttcagc ccacaggaac ccaggctgg ttgagggga cattccctgc tgacgggaca 60

tgtagccaca gcagggacgt cggcgtagtc tgctcaagtg 100

<210> SEQ ID NO 482
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 482

tttttcagcc cacaggaac ccaggctgg ttgaggggac attccctgct aggcggccag 60

cgtgctgtgc aggaactac agtgcggcac tgtggtttcc 100

<210> SEQ ID NO 483
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 483

tttttcagcc cacaggaac ccaggctgg ttgaggggac attccctgct aagtgacag 60

atctgggctg aagaattcca gtgtgagggg cagcagtc 100

<210> SEQ ID NO 484
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 484

tttttcagcc cacaggaac ccaggctgg ttgaggggac attccctgct gaggggcacg 60

agtcccacct ttctactctgc ccagtagcac cccgccctga 100

<210> SEQ ID NO 485
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 485

tttttcagcc cacaggaac ccaggctgg ttgaggggac attccctgct agcaccgcc 60

cctgacggga catgtagcca cagcagggac gtcggcgtag 100

<210> SEQ ID NO 486
<211> LENGTH: 100
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 486

tttttcagcc cacaggaac ccaggctgg ttgaggggac attccctgct gacgggacat 60

gtagccacag cagggacgtc ggcgtagtct gctcaagtga 100

<210> SEQ ID NO 487
<211> LENGTH: 100
<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 487

cccaggctgg ttggagggga cattccctgc tctggctgtg ttgaagtaca gtgattctga 60

cttctctctg gaggcggcca gcgtgctgtg cagggaacta 100

<210> SEQ ID NO 488

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 488

cccaggctgg ttggagggga cattccctgc tctggctgtg ttgaagtaca gtgaggggca 60

cgagtccac ctttactct gccagtagc accccgcct 100

<210> SEQ ID NO 489

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 489

ttggagggga cattccctgc tctggctgtg ttgaagtaca acatggagac tgaggggac 60

gagtccacc tttactctg ccagtagca ccccgccctg 100

<210> SEQ ID NO 490

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 490

ttggagggga cattccctgc tctggctgtg ttgaagtaca acatggagac tgaggggac 60

gagtccacc tttactctg ccagtagca ccccgccctg 100

<210> SEQ ID NO 491

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 491

ttggagggga cattccctgc tctggctgtg ttgaagtaca acatggagac tgacgggaca 60

tgtagccaca gcagggaagt cggcgtagtc tgctcaagtg 100

<210> SEQ ID NO 492

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 492

ggaggggaca ttccctgctc tggctgtgtt gaagtacaac atggagacac gtgattctga 60

cttctctctg gaggcggcca gcgtgctgtg cagggaacta 100

<210> SEQ ID NO 493

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 493

ggaggggaca ttccctgctc tggctgtgtt gaagtacaac atggagacac gtgaggggca 60

cgagtccac ctttactct gccagtagc accccgcct 100

-continued

<210> SEQ ID NO 494

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 494

```
ggaggggaca ttccctgctc tggctgtgtt gaagtacaac atggagacac ctgacgggac    60
atgtagccac agcaggggacg tcggcgtagt ctgctcaagt    100
```

<210> SEQ ID NO 495

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 495

```
ctgctctggt cgtgttgaag tacaacatgg agacacgtgg ggcaccgtct aggcggccag    60
cgtgctgtgc agggaactac agtgcggcac tgtggtttcc    100
```

<210> SEQ ID NO 496

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 496

```
ctgctctggt cgtgttgaag tacaacatgg agacacgtgg ggcaccgtct aagtggacag    60
atctgggctg aagaattcca gtgtgagggg cacgagtcct    100
```

<210> SEQ ID NO 497

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 497

```
ctgctctggt cgtgttgaag tacaacatgg agacacgtgg ggcaccgtct gaggggcacg    60
agtccacact ttcactctgc ccagtagcac cccgccctga    100
```

<210> SEQ ID NO 498

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 498

```
ctgctctggt cgtgttgaag tacaacatgg agacacgtgg ggcaccgtct agcaccgccg    60
cctgacggga catgtagcca cagcagggac gtcggcgtag    100
```

<210> SEQ ID NO 499

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 499

```
ctgctctggt cgtgttgaag tacaacatgg agacacgtgg ggcaccgtct gacgggacat    60
gtagccacag cagggacgtc ggcgtagtct gctcaagtga    100
```

<210> SEQ ID NO 500

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 500

tgaagtacaa catggagaca cgtggggcac cgtctgtgat tctgacttct aggcggccag 60

cgtgctgtgc agggaactac agtgcggcac tgtggtttcc 100

<210> SEQ ID NO 501

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 501

tgaagtacaa catggagaca cgtggggcac cgtctgtgat tctgacttct aagtggacag 60

atctgggctg aagaattcca gtgtgagggg cagcagtc 100

<210> SEQ ID NO 502

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 502

tgaagtacaa catggagaca cgtggggcac cgtctgtgat tctgacttct gaggggcacg 60

agtccacct ttcactctgc ccagtagcac cccgccctga 100

<210> SEQ ID NO 503

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 503

tgaagtacaa catggagaca cgtggggcac cgtctgtgat tctgacttct agcaccgccg 60

cctgacggga catgtagcca cagcagggac gtcggcgtag 100

<210> SEQ ID NO 504

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 504

tgaagtacaa catggagaca cgtggggcac cgtctgtgat tctgacttct gacgggacat 60

gtagccacag cagggacgtc ggcgtagtct gctcaagtga 100

<210> SEQ ID NO 505

<211> LENGTH: 100

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 505

tccttcaccg aaatgctatt ttttcagccc acaggaaacc caggctgggt ctgtggtttc 60

cctcctgggg ggagctcact ttggagaagg aagtggacag 100

<210> SEQ ID NO 506

<211> LENGTH: 3

<212> TYPE: PRT

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 506

Ala His Arg

1

<210> SEQ ID NO 507

-continued

<211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 507

Ala His Arg Lys
 1

<210> SEQ ID NO 508
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 508

Ala His Arg Lys Pro Arg Leu
 1 5

<210> SEQ ID NO 509
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 509

Ala His Arg Lys Pro Arg Leu Val
 1 5

<210> SEQ ID NO 510
 <211> LENGTH: 13
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 510

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro
 1 5 10

<210> SEQ ID NO 511
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 511

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys
 1 5 10

<210> SEQ ID NO 512
 <211> LENGTH: 21
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 512

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15

Arg Val Glu Val Gln
 20

<210> SEQ ID NO 513
 <211> LENGTH: 24
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 513

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15

Arg Val Glu Val Gln His Gly Asp

-continued

20

<210> SEQ ID NO 514
 <211> LENGTH: 25
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 514

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15
 Arg Val Glu Val Gln His Gly Asp Thr
 20 25

<210> SEQ ID NO 515
 <211> LENGTH: 29
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 515

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15
 Arg Val Glu Val Gln His Gly Asp Thr Trp Gly Thr Val
 20 25

<210> SEQ ID NO 516
 <211> LENGTH: 29
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 516

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15
 Arg Val Glu Val Gln His Gly Asp Thr Trp Gly Thr Val
 20 25

<210> SEQ ID NO 517
 <211> LENGTH: 34
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 517

Ala His Arg Lys Pro Arg Leu Val Gly Gly Asp Ile Pro Cys Ser Gly
 1 5 10 15
 Arg Val Glu Val Gln His Gly Asp Thr Trp Gly Thr Val Cys Asp Ser
 20 25 30

Asp Phe

<210> SEQ ID NO 518
 <211> LENGTH: 64
 <212> TYPE: PRT
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 518

Ala His Arg Lys Pro Arg Leu Val Thr Val Val Ser Leu Leu Gly Gly
 1 5 10 15
 Ala His Phe Gly Glu Gly Ser Gly Gln Ile Trp Ala Glu Glu Phe Gln
 20 25 30
 Cys Glu Gly His Glu Ser His Leu Ser Leu Cys Pro Val Ala Pro Arg
 35 40 45
 Pro Asp Gly Thr Cys Ser His Ser Arg Asp Val Gly Val Val Cys Ser
 50 55 60

-continued

```

<210> SEQ ID NO 519
<211> LENGTH: 1010
<212> TYPE: PRT
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 519

Met Asp Lys Leu Arg Met Val Leu His Glu Asn Ser Gly Ser Ala Asp
 1              5              10              15

Phe Arg Arg Cys Ser Ala His Leu Ser Ser Phe Thr Phe Ala Val Val
      20              25              30

Ala Val Leu Ser Ala Cys Leu Val Thr Ser Ser Leu Gly Gly Lys Asp
      35              40              45

Lys Glu Leu Arg Leu Thr Gly Gly Glu Asn Lys Cys Ser Gly Arg Val
      50              55              60

Glu Val Lys Val Gln Glu Glu Trp Gly Thr Val Cys Asn Asn Gly Trp
      65              70              75              80

Asp Met Asp Val Val Ser Val Val Cys Arg Gln Leu Gly Cys Pro Thr
      85              90              95

Ala Ile Lys Ala Thr Gly Trp Ala Asn Phe Ser Ala Gly Ser Gly Arg
      100              105              110

Ile Trp Met Asp His Val Ser Cys Arg Gly Asn Glu Ser Ala Leu Trp
      115              120              125

Asp Cys Lys His Asp Gly Trp Gly Lys His Asn Cys Thr His Gln Gln
      130              135              140

Asp Ala Gly Val Thr Cys Ser Asp Gly Ser Asp Leu Glu Met Arg Leu
      145              150              155              160

Val Asn Gly Gly Asn Arg Cys Leu Gly Arg Ile Glu Val Lys Phe Gln
      165              170              175

Gly Arg Trp Gly Thr Val Cys Asp Asp Asn Phe Asn Ile Asn His Ala
      180              185              190

Ser Val Val Cys Lys Gln Leu Glu Cys Gly Ser Ala Val Ser Phe Ser
      195              200              205

Gly Ser Ala Asn Phe Gly Glu Gly Ser Gly Pro Ile Trp Phe Asp Asp
      210              215              220

Leu Val Cys Asn Gly Asn Glu Ser Ala Leu Trp Asn Cys Lys His Glu
      225              230              235              240

Gly Trp Gly Lys His Asn Cys Asp His Ala Glu Asp Ala Gly Val Ile
      245              250              255

Cys Leu Asn Gly Ala Asp Leu Lys Leu Arg Val Val Asp Gly Val Thr
      260              265              270

Glu Cys Ser Gly Arg Leu Glu Val Lys Phe Gln Gly Glu Trp Gly Thr
      275              280              285

Ile Cys Asp Asp Gly Trp Asp Ser Asp Asp Ala Ala Val Ala Cys Lys
      290              295              300

Gln Leu Gly Cys Pro Thr Ala Val Thr Ala Ile Gly Arg Val Asn Ala
      305              310              315              320

Ser Glu Gly Thr Gly His Ile Trp Leu Asp Ser Val Ser Cys His Gly
      325              330              335

His Glu Ser Ala Leu Trp Gln Cys Arg His His Glu Trp Gly Lys His
      340              345              350

Tyr Cys Asn His Asn Glu Asp Ala Gly Val Thr Cys Ser Asp Gly Ser
      355              360              365

Asp Leu Glu Leu Arg Leu Lys Gly Gly Gly Ser His Cys Ala Gly Thr

```


-continued

370	375	380
Val Glu Val Glu Ile Gln Lys Leu Val Gly Lys Val Cys Asp Arg Ser 385 390 395 400		
Trp Gly Leu Lys Glu Ala Asp Val Val Cys Arg Gln Leu Gly Cys Gly 405 410 415		
Ser Ala Leu Lys Thr Ser Tyr Gln Val Tyr Ser Lys Thr Lys Ala Thr 420 425 430		
Asn Thr Trp Leu Phe Val Ser Ser Cys Asn Gly Asn Glu Thr Ser Leu 435 440 445		
Trp Asp Cys Lys Asn Trp Gln Trp Gly Gly Leu Ser Cys Asp His Tyr 450 455 460		
Asp Glu Ala Lys Ile Thr Cys Ser Gly Tyr Thr Gln Ile Arg Leu Val 465 470 475 480		
Asn Gly Lys Thr Pro Cys Glu Gly Arg Val Glu Leu Asn Ile Leu Gly 485 490 495		
Ser Trp Gly Ser Leu Cys Asn Ser His Trp Asp Met Glu Asp Ala His 500 505 510		
Val Leu Cys Gln Gln Leu Lys Cys Gly Val Ala Leu Ser Ile Pro Arg 515 520 525		
Gly Ala Pro Phe Gly Lys Gly Ser Glu Gln Val Trp Arg His Met Phe 530 535 540		
His Cys Thr Gly Thr Glu Lys His Met Gly Asp Cys Ser Val Thr Ala 545 550 555 560		
Leu Gly Ala Ser Leu Cys Ser Ser Gly Gln Val Ala Ser Val Ile Cys 565 570 575		
Ser Gly Asn Gln Ser Gln Thr Leu Ser Pro Cys Asn Ser Ser Ser Ser 580 585 590		
Asp Pro Ser Ser Ser Ile Ile Ser Glu Glu Asn Gly Val Ala Cys Ile 595 600 605		
Gly Ser Gly Gln Leu Arg Leu Val Asp Gly Gly Gly Arg Cys Ala Gly 610 615 620		
Arg Val Glu Val Tyr His Glu Gly Ser Trp Gly Thr Ile Cys Asp Asp 625 630 635 640		
Ser Trp Asp Leu Asn Asp Ala His Val Val Cys Lys Gln Leu Ser Cys 645 650 655		
Gly Trp Ala Ile Asn Ala Thr Gly Ser Ala His Phe Gly Glu Gly Thr 660 665 670		
Gly Pro Ile Trp Leu Asp Glu Ile Asn Cys Asn Gly Lys Glu Ser His 675 680 685		
Ile Trp Gln Cys His Ser His Gly Trp Gly Arg His Asn Cys Arg His 690 695 700		
Lys Glu Asp Ala Gly Val Ile Cys Ser Glu Phe Met Ser Leu Arg Leu 705 710 715 720		
Ile Ser Glu Asn Ser Arg Glu Thr Cys Ala Gly Arg Leu Glu Val Phe 725 730 735		
Tyr Asn Gly Ala Trp Gly Ser Val Gly Arg Asn Ser Met Ser Pro Ala 740 745 750		
Thr Val Gly Val Val Cys Arg Gln Leu Gly Cys Ala Asp Arg Gly Asp 755 760 765		
Ile Ser Pro Ala Ser Ser Asp Lys Thr Val Ser Arg His Met Trp Val 770 775 780		
Asp Asn Val Gln Cys Pro Lys Gly Pro Asp Thr Leu Trp Gln Cys Pro 785 790 795 800		

-continued

Ser Ser Pro Trp Lys Lys Arg Leu Ala Ser Pro Ser Glu Glu Thr Trp
 805 810 815
 Ile Thr Cys Ala Asn Lys Ile Arg Leu Gln Glu Gly Asn Thr Asn Cys
 820 825 830
 Ser Gly Arg Val Glu Ile Trp Tyr Gly Gly Ser Trp Gly Thr Val Cys
 835 840 845
 Asp Asp Ser Trp Asp Leu Glu Asp Ala Gln Val Val Cys Arg Gln Leu
 850 855 860
 Gly Cys Gly Ser Ala Leu Glu Ala Gly Lys Glu Ala Ala Phe Gly Gln
 865 870 875 880
 Gly Thr Gly Pro Ile Trp Leu Asn Glu Val Lys Cys Lys Gly Asn Glu
 885 890 895
 Thr Ser Leu Trp Asp Cys Pro Ala Arg Ser Trp Gly His Ser Asp Cys
 900 905 910
 Gly His Lys Glu Asp Ala Ala Val Thr Cys Ser Glu Ile Ala Lys Ser
 915 920 925
 Arg Glu Ser Leu His Ala Thr Gly Arg Ser Ser Phe Val Ala Leu Ala
 930 935 940
 Ile Phe Gly Val Ile Leu Leu Ala Cys Leu Ile Ala Phe Leu Ile Trp
 945 950 955 960
 Thr Gln Lys Arg Arg Gln Arg Gln Arg Leu Ser Val Phe Ser Gly Gly
 965 970 975
 Glu Asn Ser Val His Gln Ile Gln Tyr Arg Glu Met Asn Ser Cys Leu
 980 985 990
 Lys Ala Asp Glu Thr Asp Met Leu Asn Pro Ser Gly Asp His Ser Glu
 995 1000 1005
 Val Gln
 1010

<210> SEQ ID NO 520
 <211> LENGTH: 101
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 520

ctcaacttttc actctctggc ttactcctat caagagggta gggtaggta gtcacagaca 60
 tctttttaaa gccctgtctc cttccaggat acacacaaat c 101

<210> SEQ ID NO 521
 <211> LENGTH: 100
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 521

ctcaacttttc actctctggc ttactcctat cagagggttag ggttaggtag tcacagacat 60
 ctttttaaag ccctgtctcc ttccaggata cacacaaatc 100

<210> SEQ ID NO 522
 <211> LENGTH: 98
 <212> TYPE: DNA
 <213> ORGANISM: Sus scrofa

<400> SEQUENCE: 522

ctcaacttttc actctctggc ttactcctat cagggtaggg ttaggtagtc acagacatct 60
 ttttaaagcc ctgtctcctt ccaggataca cacaaatc 98

-continued

<210> SEQ ID NO 523
<211> LENGTH: 108
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 523

ctcacttttc actctctggc ttactcctat catggcatgg gagggtaggg ttaggtagtc 60
acagacatct ttttaaagcc ctgtctcctt ccaggatata cacaatc 108

<210> SEQ ID NO 524
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 524

ctcacttttc aggggttaggt agtcacagac atctttttaa agccctgtct ccttccagga 60
tacacacaaa tc 72

<210> SEQ ID NO 525
<211> LENGTH: 90
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 525

ctcacttttc actctctggc ttactcctag ggtaggtag tcacagacat ctttttaaag 60
ccctgtctcc ttccaggata cacacaaatc 90

<210> SEQ ID NO 526
<211> LENGTH: 75
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 526

ctcacttttc actctctggc ttactcccca gacatctttt taaagccctg tctccttcca 60
ggatacacac aaatc 75

<210> SEQ ID NO 527
<211> LENGTH: 96
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 527

ctcacttttc actctctggc ttactccaga ggtaggggtt aggtagtcac agacatcttt 60
ttaaagccct gtctccttcc aggatacaca caaatc 96

<210> SEQ ID NO 528
<211> LENGTH: 97
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 528

ctcacttttc actctctggc ttactcctat caggtagggt taggtagtca cagacatctt 60
tttaaagccc tgtctccttc caggatacac acaaatc 97

<210> SEQ ID NO 529
<211> LENGTH: 90
<212> TYPE: DNA
<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 529

-continued

ctcacttttc actctctggc ttactcctag ggtaggtag tcacagacat ctttttaaag 60

ccctgtctcc ttccaggata cacacaaatc 90

<210> SEQ ID NO 530

<211> LENGTH: 99

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 530

ctcacttttc actctctggc ttactcctat caagggtagg gttaggtagt cacagacatc 60

tttttaaagc cctgtctcct tccaggatac acacaaatc 99

<210> SEQ ID NO 531

<211> LENGTH: 93

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 531

ctcacttttc actctctggc ttactcctat cagggttagg tagtcacaga catcttttta 60

aagccctgtc tccttcagg atacacacaa atc 93

<210> SEQ ID NO 532

<211> LENGTH: 92

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 532

ctcacttttc actctctggc ttactcgggt aggggttagg agtcacagac atcttttta 60

agccctgtct ccttcagga tacacacaaa tc 92

<210> SEQ ID NO 533

<211> LENGTH: 98

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 533

ctcacttttc actctctggc ttactcctat caaggtaggg ttaggtagtc acagacatct 60

ttttaaagcc ctgtctcctt ccaggatata cacaaatc 98

<210> SEQ ID NO 534

<211> LENGTH: 99

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 534

ctcacttttc actctctggc ttactcctat cacaggtagg gttaggtagt cacagacatc 60

tttttaaagc cctgtctcct tccaggatac acacaaatc 99

<210> SEQ ID NO 535

<211> LENGTH: 95

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 535

ctcacttttc actctctggc ttactcctat ggtaggggta ggtagtcaca gacatctttt 60

taaagccctg tctccttcca ggatacacac aaatc 95

<210> SEQ ID NO 536

-continued

```

<211> LENGTH: 125
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 536
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac    60
ggtagggtta ggtagtcaca gacatctttt taaagccctg tctccttcca ggatacacac    120
aaatc                                           125

<210> SEQ ID NO 537
<211> LENGTH: 124
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 537
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac    60
gtagggttag gtagtcacag acatcttttt aaagccctgt tctcttcag gatacacaca    120
aatc                                           124

<210> SEQ ID NO 538
<211> LENGTH: 123
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 538
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccat    60
tagggtagg tagtcacaga catcttttta aagccctgtc tccttcagg atacacacaa    120
atc                                           123

<210> SEQ ID NO 539
<211> LENGTH: 114
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 539
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctagggttag    60
gtagtcacag acatcttttt aaagccctgt tctcttcag gatacacaca aatc          114

<210> SEQ ID NO 540
<211> LENGTH: 123
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 540
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac    60
tagggtagg tagtcacaga catcttttta aagccctgtc tccttcagg atacacacaa    120
atc                                           123

<210> SEQ ID NO 541
<211> LENGTH: 116
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 541
gcaaagtccc agaattgtct ccaggggaagg acagggaggt ctagaatcgg ctaagcccac    60
aggtagtcac agacatcttt ttaaagccct gtctccttcc aggatacaca caaatc       116

```

-continued

<210> SEQ ID NO 542

<211> LENGTH: 124

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 542

gcaaagtccc agaattgtct ccaggaagg acaggagggt ctagaatcgg ctaagcccag 60

gtagggttag gtagtcacag acatcttttt aaagccctgt ctcttccag gatacacaca 120

aatc 124

<210> SEQ ID NO 543

<211> LENGTH: 122

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 543

gcaaagtccc agaattgtct ccaggaagg acaggagggt ctagaatcgg ctaagcccat 60

ggggtaggt agtcacagac atctttttaa agccctgtct ccttccagga tacacacaaa 120

tc 122

<210> SEQ ID NO 544

<211> LENGTH: 113

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 544

gcaaagtccc agaattgtct ccaggaagg acaggagggt ctagaatcgg tagggtaggt 60

tagtcacaga catcttttta aagccctgtc tccttccagg atacacacaa atc 113

<210> SEQ ID NO 545

<211> LENGTH: 113

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 545

atctctggtg tgcctttgac tccagattac agtaaatgga ggactgaggg tagggtaggt 60

tagtcacaga catcttttta aagccctgtc tccttccagg atacacacaa atc 113

<210> SEQ ID NO 546

<211> LENGTH: 117

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 546

atctctggtg tgcctttgac tccagattac agtaaatgga ggactgagag agggtagggt 60

taggtagtca cagacatctt ttaaagccc tgtctccttc caggatacac acaaatc 117

<210> SEQ ID NO 547

<211> LENGTH: 106

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 547

atctctggtg tgcctttgac tccagattac agtaaatgga gggtagggtt aggtagtcac 60

agacatcttt ttaaagccct gtctccttc aggatacaca caaatc 106

<210> SEQ ID NO 548

<211> LENGTH: 112

<212> TYPE: DNA

-continued

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 548

atttctggtg tgcctttgac tccagattac agtaaatgga ggactgaggg tgggttaggt 60

agtcacagac atctttttaa agccctgtct ccttccagga tacacacaaa tc 112

<210> SEQ ID NO 549

<211> LENGTH: 117

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 549

atttctggtg tgcctttgac tccagattac agtaaatgga ggactgagag aggatagggt 60

taggtagtca cagacatctt tttaaagccc tgtctccttc caggatacac acaaatac 117

<210> SEQ ID NO 550

<211> LENGTH: 90

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 550

gaaagagaaa gcatgctcca agaatggtag ggtaggtag tcacagacat ctttttaaag 60

ccctgtctcc ttccaggata cacacaaatac 90

<210> SEQ ID NO 551

<211> LENGTH: 89

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 551

gaaagagaaa gcatgctcca agaaggtagg gttaggtagt cacagacatc tttttaaagc 60

cctgtctcct tccaggatac acacaaatac 89

<210> SEQ ID NO 552

<211> LENGTH: 91

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 552

gaaagagaaa gcatgctcca agaatgggta gggtaggta gtcacagaca tctttttaa 60

gcctgtcttc cttccaggat acacacaaat c 91

<210> SEQ ID NO 553

<211> LENGTH: 87

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 553

gaaagagaaa gcatgctcca agagtagggt taggtagtca cagacatctt tttaaagccc 60

tgtctccttc caggatacac acaaatac 87

<210> SEQ ID NO 554

<211> LENGTH: 97

<212> TYPE: DNA

<213> ORGANISM: *Sus scrofa*

<400> SEQUENCE: 554

gaaagagaaa gcatgctcca agaatccaag aggtagggt taggtagtca cagacatctt 60

tttaaagccc tgtctccttc caggatacac acaaatac 97

-continued

<210> SEQ ID NO 555
<211> LENGTH: 92
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 555

gaaagagaaa gcatgctcca agaatagggt agggtaggt agtcacagac atctttttaa 60
agccctgtct ccttcagga tacacacaaa tc 92

<210> SEQ ID NO 556
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 556

ctaagccac tgtaggcaga a 21

<210> SEQ ID NO 557
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 557

ctgggtttcc tgtgggctga a 21

<210> SEQ ID NO 558
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 558

tggcttactc ctatcatgaa ggaa 24

<210> SEQ ID NO 559
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 559

atgtagccac agcagggacg t 21

<210> SEQ ID NO 560
<211> LENGTH: 22
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 560

ccaagcggat ttgtgtgtat cc 22

<210> SEQ ID NO 561
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 561

ttcccatgcc atgaagaggg tagggtta 28

<210> SEQ ID NO 562
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Sus scrofa

-continued

<400> SEQUENCE: 562

gctaagccca ctgtaggcag a

21

<210> SEQ ID NO 563

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 563

gcggatttgt gtgtatcctg g

21

<210> SEQ ID NO 564

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Sus scrofa

<400> SEQUENCE: 564

tggettactc ctatcaggta gggt

24

What is claimed is:

1. A method of screening comprising:

- i) breeding a genetically edited pig whose genome comprises a CD163 gene that has a homozygous deletion of exon 7 and comprises the nucleic acid sequence of SEQ ID NO: 453, wherein the pig exhibits resistance to porcine reproductive and respiratory syndrome virus (PRRSV) such that offspring are obtained;
- ii) screening the genomes of the offspring for the presence of a CD163 gene that has a homozygous deletion of

exon 7 and comprises the nucleic acid sequence of SEQ ID NO: 453; and

iii) screening the offspring for resistance to PRRSV.

2. The method of claim 1, wherein the screening is performed using polymerase chain reaction (PCR) and primers comprising the nucleic acid sequences of SEQ ID NO: 556 and 557.

3. The method of claim 1, wherein the screening is performed using polymerase chain reaction (PCR) and primers comprising the nucleic acid sequences of SEQ ID NO: 562 and 563.

* * * * *