



US 20250281502A1

(19) **United States**

(12) **Patent Application Publication**
DUTRIEUX et al.

(10) **Pub. No.: US 2025/0281502 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **NEW DRUG APPLICATION**

Publication Classification

(71) Applicant: **Centre National de la Recherche Scientifique, PARIS (FR)**

(51) **Int. Cl.**

A61K 31/5517 (2006.01)

A61K 31/4045 (2006.01)

A61K 31/4709 (2006.01)

A61P 35/00 (2006.01)

(72) Inventors: **Laure DUTRIEUX, Montpellier (FR); Jérôme MOREAUX, Montpellier (FR); Philippe PASERO, Montpellier (FR); Raphaël RODRIGUEZ, Paris (FR); Michel COGNÉ, Rennes (FR)**

(52) **U.S. Cl.**

CPC A61K 31/5517 (2013.01); A61K 31/4045 (2013.01); A61K 31/4709 (2013.01); A61P 35/00 (2018.01)

(21) Appl. No.: **18/575,733**

(22) PCT Filed: **Jul. 1, 2022**

(86) PCT No.: **PCT/EP2022/068304**

§ 371 (c)(1),

(2) Date: **Dec. 29, 2023**

(57)

ABSTRACT

A method is disclosed for treating an individual afflicted by a multiple myeloma by a composition comprising at least one G-quadruplex (G4) stabilizer. Also disclosed is a composition comprising at least one G-quadruplex (G4) stabilizer for its use in a method for treating an individual afflicted by a multiple myeloma.


(30) **Foreign Application Priority Data**

Jul. 1, 2021 (EP) 21305916.5

Jul. 27, 2021 (EP) 21306047.8

Jul. 27, 2021 (FR) FR2108147

Specification includes a Sequence Listing.






Fig. 1

Fig. 2A




Fig. 2B




Fig. 2C




Fig. 2D




Fig. 2E




Fig. 3A




Fig. 3B




Fig. 3C




Fig. 3D




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 4D




Fig. 5A




Fig. 5B




Fig. 6A




Fig. 6B




Fig. 6C




Fig. 7A




Fig. 7B




Fig. 8A




Fig. 8B




Fig. 9A





Fig. 9B






Fig. 10




Fig. 11




Fig. 12

Fig. 13A




Fig. 13B




Fig. 14A





Fig. 14B






Fig. 15

Fig. 16A




Fig. 16B




Fig. 16C




Fig. 17A




Fig. 17B




Fig. 17C




Fig. 18A




Fig. 18B




Fig. 18C




Fig. 19A




Fig. 19B





Fig. 19C



NEW DRUG APPLICATION

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application is a filing under 35 U.S.C. 371 as the National Stage of International Application No. PCT/EP2022/068304, filed Jul. 1, 2022, entitled “NEW DRUG APPLICATION”, which claims priority to European Application No. EP 21305916.5 filed with the European Patent Office on Jul. 1, 2021, and entitled “NEW DRUG APPLICATION”; European Application No. EP 21306047.8 filed with the European Patent Office on Jul. 27, 2021, and entitled “NEW DRUG APPLICATION”; and French Application No. FR 2108147 filed with the French Patent Office on Jul. 27, 2021, and entitled “NEW DRUG APPLICATION”, all of which are incorporated herein by reference in their entirety for all purposes.

INCORPORATION BY REFERENCE OF MATERIAL IN XML FILE

[0002] This application incorporates by reference the Sequence Listing contained in the following XML file submitted Dec. 29, 2023:

[0003] File name: 4692-15000 Sequence listing.xml; filed on Dec. 29, 2023; and having a file size of 30.7 KB.

[0004] The information in the Sequence Listing is incorporated herein in its entirety for all purposes.

[0005] The invention relates to a new drug application.

[0006] Multiple myeloma (MM) is the second most frequent hematological malignancy, characterized by the accumulation of malignant plasma cells (PCs) within the bone marrow. More than 5000 new cases are identified in France each year, and multiple myeloma represents 2% of cancer-related cell death.

[0007] To date, there is no definitive treatment for this pathology and a majority of patients will invariably relapse. Oncogenic transformation in MM is thought to occur within the secondary lymphoid organs. Malignant PCs present a high rate of somatic mutations, suggesting that the oncogenic event occurs after the end of the somatic hypermutation (SHM) process, which physiologically takes place in the germinal centers (GCs). However, the precise molecular events leading to myelomagenesis remain obscure. Broad genomic instability is frequent in MM and not restricted to the IgH locus, but IgH S regions are exquisitely involved in various translocations (notably with MAF, MAFB, CCND1, CCND3, WWOX, FGFR3/MMSET . . .) which thus stand as late “post-GC” events. These events can yield fused transcripts, the frequency of which is clearly correlated with a bad prognosis. A better understanding of early myelomagenesis and myeloma progression will help define therapeutic areas of interest.

[0008] The invention intends to obviate this lack in the art.

[0009] The object of the invention is to provide a new efficient drug for treating multiple myeloma.

[0010] Thus, the invention relates to a composition comprising at least one G-quadruplex (G4) stabilizer for its use in a method for treating an individual afflicted by a multiple myeloma.

[0011] In the invention, the term “individual” refers to a mammal individual, preferably a human individual.

[0012] The inventors unexpectedly discovered that a patient with MM can be treated by a therapeutic composition

targeting the transcription/replication conflicts (TRCs) resolution cell machinery, and more particularly by stabilising the G4 structures.

[0013] TRCs occur at R-loop structures during the replication stage of a cell. R-loops are three-stranded nucleic acid structures, formed by the annealing of an RNA moiety with double-stranded DNA constituting an RNA:DNA hybrid. These structures are physiologically enriched near promoters and transcription termination sites, and are involved in immunoglobulin (Ig) class switch recombination (CSR), transcription initiation and termination, and telomere elongation. Unscheduled R-loop formation interferes with replication fork progression and increases the collision rate between the replication and transcription machineries, known as transcription/replication conflicts (TRCs).

[0014] G-quadruplex (G4) are four-stranded secondary DNA structures, constituted of at least two stacked guanine tetrads stabilized by Hoogsteen hydrogen bonds and cations, forming a planar complex (G-quartet). These G-quartets are stabilized by a central counterion, typically K⁺, and stack upon each other forming stable structures. These highly stable non-canonical structures are present at telomeres, at the promoter of many genes, and at replication origins. G4s can be formed in the displaced DNA strand of a R-loop in order to stabilize it.

[0015] G4 stabilizers are compounds that avoid the G4 structures to untie, with the result that G4 stabilizers impede with the resolution of R-loop structures and trigger the occurrence of TRCs.

[0016] Unexpectedly, the inventors identified that G4 stabilizers are able to induce apoptosis or to inhibit cell cycle of primary cells from individuals afflicted by a multiple myeloma.

[0017] In particular, the at least one G4 stabilizer is selected in a group consisting of Quarfloxin, Pidnarulex, MM41 (4,9-Bis((3-(4-methylpiperazin-1-yl)propyl)amino)-2,7-bis(3-morpholinopropyl)benzo[Imn][3,8]phenanthroline-1,3,6,8(2H,7H)-tetraone), Telomestatin, BMSG-SH-3 (2,7-Bis-[5-(4-methyl-piperazin-1-yl)-pentyl]-4,9-bis-(3-(4-methyl-piperazin-1-yl)-propylamino)-benzo[Imn][3,8]phenanthroline-1,3,6,8-tetraone), BRACO-19 (N,N'-(9-(4-(Dimethylamino)phenylamino)acridine-3,6-diyl)bis(3-(pyrrolidin-1-yl)propanamide) hydrochloride), CM03 ((2,7-bis(3-morpholinopropyl)-4-((2-(pyrrolidin-1-yl)ethyl)amino)benzo[Imn][3,8]phenanthroline-1,3,6,8(2H,7H)-tetraone), PDP, Pyridostatin, carboxyPyridostatin, PhenDC3 (3,3'-[1,10-Phenanthroline-2,9-diylbis(carbonylimino)]bis[1-methylquinolinium]1,1,1-trifluoromethanesulfonate (1:2)), AQ1, TMPyP4 (Meso-Tetra (N-methyl-4-pyridyl) porphine tetra tosylate), RHPS4 (3,11-Difluoro-6,8,13-trimethylquino[4,3,2-kl]acridinium methylsulfate), 360A(2-N,6-N-bis(1-methylquinolin-1-ium-3-yl)pyridine-2,6-dicarboxamide), FG (bis-guanylylhydrazone derivative of diimidazo [1,2-a:1,2-c]pyrimidine), 20A, Emetine and in particular combinations thereof. In particular, the G4 stabilizer is Pyridostatin.

[0018] PDP corresponds to compound 7 in Emanuela Ruggiero and Sara N Richter, *Nucleic Acids Research*, Volume 46, Issue 7, 20 Apr. 2018, Pages 3270-3283.

[0019] AQ1 is described on FIG. 2 in Eleonora Zorzan et al., *Oncotarget*, 2016, Vol. 7, No. 16.

[0020] 20A corresponds to compound 3 in N. M. Smith et al., *Organic & Biomolecular Chemistry*, 2011, issue 17.

[0021] Advantageously, the at least one G4 stabilizer is associated with a pharmaceutical acceptable vehicle.

[0022] An “acceptable pharmaceutical vehicle” refers in the invention to any carrier, emulsion or excipient that does not impede with the therapeutic effect of the composition nor harm the health of the individual. It is within the skills of a physician to determine the said acceptable pharmaceutical vehicle.

[0023] It also is within the skills of a physician to determine the specific therapeutically effective dosage regimen, as this dosage regimen will be dependent upon a variety of factors including, but not limited to: the severity of the multiple myeloma; the age; the body weight; general health; the sex; the diet; the time course of administration; the route of administration; the duration of the treatment; the drugs that are concomitantly administered in combination with the pharmaceutical composition within the scope of the present invention.

[0024] In some embodiments, the dosage regimen of said at least one G4 stabilizer ranges from about 0.0001 mg to about 1,000 mg per adult, per day. Preferably, the individual is administered with an amount of about 0.0001, 0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 7.5, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 150, 250, 500 and 750 mg of said at least one G4 stabilizer in order to adjust the dosage regimen that is the most suitable to a particular individual in need of the treatment.

[0025] A pharmaceutical composition within the scope of the present invention may contain from about 0.01 mg to about 500 mg of said at least one G4 stabilizer, preferably from about 1 mg to about 100 mg of said at least one G4 stabilizer.

[0026] In a preferred embodiment, an effective amount of at least one G4 stabilizer is routinely administered to an individual in need thereof, at a dosage regimen from about 0.0002 mg/kg to about 20 mg/kg of body weight per day, in particular from about 0.001 mg/kg to 7 mg/kg of body weight per day.

[0027] The optimal amount of said at least one G4 stabilizer to be comprised in a pharmaceutical dosage unit according to the invention may be easily adapted by the one skilled in the art using routine known protocols or methods.

[0028] Said composition may be administered by any suitable route, i.e. including, but not limited to, an oral, sublingual, subcutaneous, intraperitoneal, intramuscular, intravenous, intrathecal and rectal administration.

[0029] In one embodiment of the invention, the composition further comprises a drug commonly used for treating multiple myeloma, and possibly to which a resistance occurs.

[0030] In the invention, “a drug commonly used for treating multiple myeloma” refers to anticancer drugs or compounds.

[0031] Resistance to a drug, regarding MM, means that said drug is not able to affect survival and/or proliferation of the cells that constitute MM (induce apoptosis and/or necrosis and inhibit cell proliferation). If a resistance occurs, it means that the malignant cells of the MM were initially sensitive to the drug, but further to the treatment, or during the treatment, mutations may occur in some cells, such that the target of the drug is not any more sensitive to the drug. Therefore, the cells become insensitive to the drug and a resistance appears, i.e. the tumor grows from the resistant cells.

[0032] In particular, the composition further comprises at least one histone deacetylase inhibitor.

[0033] The inventors unexpectedly discovered that stabilizing G4 structures potentializes the histone deacetylase inhibitors cytotoxicity on multiple myeloma cells. The inventors have identified that the combination between at least one G4 stabilizer and at least one histone deacetylase inhibitor is able to induce apoptosis or to inhibit cell cycle of primary cells from individuals afflicted by a multiple myeloma with a synergetic effect.

[0034] In the invention, the term “histone deacetylase inhibitor” or “HDACi” refers to histone deacetylase inhibitor that can be grouped in four classes: hydroxamates (panobinostat (LBH-589), trichostatin-A (TSA), vorinostat (SAHA), belinostat (PXD101), NVP-LAQ824 and givinostat (ITF2357)), cyclic peptide (romidepsin (depsipeptide)), aliphatic acids (valproic acid (VPA) and sodium phenylbutyrate) and benzamides (MS-275, MGCD0103). HDACi are characterized as class I-specific HDACs inhibitors (MGCD0103, romidepsin and MS-275) or as pan-HDAC inhibitors, denoting activity against both classes I and II HDACis (TSA, panobinostat, vorinostat and belinostat).

[0035] In particular, the at least one histone deacetylase inhibitor is selected in a group consisting of Panobinostat, trichostatin-A, vorinostat, belinostat, NVP-LAQ824 (Dacinostat), givinostat, romidepsin, valproic acid, sodium phenylbutyrate, MS-275 (N-(2-aminophenyl)-4-[N-(pyridine-3-yl-methoxy-carbonyl) aminomethyl]benzamide), MGCD0103 (Mocetinostat), and in particular combinations thereof. Particularly, the histone deacetylase inhibitor is Panobinostat.

[0036] Advantageously, the at least one G4 stabilizer and at least one histone deacetylase inhibitor are used simultaneously, separately, or sequentially.

[0037] By a simultaneous use, it is meant in the invention that all the compounds are injected or administered to an individual at the same time. Separately use means that the compounds are provided in a separate formulation but are injected or administered at the same time. Sequentially means that the compounds are delivered to the individual separately over the time.

[0038] Advantageously, the at least one G4 stabilizer and at least one histone deacetylase inhibitor are associated with a pharmaceutical acceptable vehicle. The pharmaceutical acceptable vehicle is as defined above.

[0039] It is within the skills of a physician to determine the specific therapeutically effective dosage regimen, as this dosage regimen will be dependent upon a variety of factors including, but not limited to: the severity of the multiple myeloma; the age; the body weight; general health; the sex; the diet; the time course of administration; the route of administration; the duration of the treatment; the drugs that are concomitantly administered in combination with the pharmaceutical composition within the scope of the present invention.

[0040] In some embodiments, the dosage regimen of each of said at least one G4 stabilizer and at least one histone deacetylase inhibitor ranges from about 0.0001 mg to about 1,000 mg per adult, per day. Preferably, the individual is administered with an amount of about 0.0001, 0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 7.5, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 150, 250, 500 and 750 mg of each of said at least one G4 stabilizer and at least one histone

deacetylase inhibitor in order to adjust the dosage regimen that is the most suitable to a particular individual in need of the treatment.

[0041] A pharmaceutical composition within the scope of the present invention may contain from about 0.01 mg to about 500 mg of each of the said at least one G4 stabilizer and at least one histone deacetylase inhibitor, preferably from about 1 mg to about 100 mg of said at least one G4 stabilizer and at least one histone deacetylase inhibitor.

[0042] In a preferred embodiment, an effective amount of each of the said at least one G4 stabilizer and at least one histone deacetylase inhibitor is routinely administered to an individual in need thereof, at a dosage regimen from about 0.0002 mg/kg to about 20 mg/kg of body weight per day, in particular from about 0.001 mg/kg to 7 mg/kg of body weight per day.

[0043] The optimal amount of the said at least one G4 stabilizer and at least one histone deacetylase inhibitor to be comprised in a pharmaceutical dosage unit according to the invention may be easily adapted by the one skilled in the art using routine known protocols or methods.

[0044] Said composition may be administered by any suitable route, i.e. including, but not limited to, an oral, sublingual, subcutaneous, intraperitoneal, intramuscular, intravenous, intrathecal and rectal administration.

[0045] In particular, the composition further comprises at least one bromodomain and extraterminal (BET) proteins inhibitor.

[0046] The inventors unexpectedly discovered that stabilizing G4 structures potentializes the BET proteins inhibitors cytotoxicity on multiple myeloma cells. The inventors have identified that the combination between at least one G4 stabilizer and at least one histone BET proteins inhibitor/deacetylase inhibitor is able to induce apoptosis or to inhibit cell cycle of primary cells from individuals afflicted by a multiple myeloma with a synergetic effect.

[0047] BET proteins inhibitors are a class of drugs that reversibly bind the bromodomains of BET proteins BRD2, BRD3, BRD4, and BRDT, and prevent protein-protein interaction between BET proteins and acetylated histones and transcription factors.

[0048] Particularly, the at least one BET proteins inhibitor is selected in the group consisting of RVX-208 (2-[4-(2-hydroxyethoxy)-3,5-dimethylphenyl]-5,7-dimethoxy-4-(3H)-quinazolinone), I-BET-762 ((4S)-6-(4-Chlorophenyl)-N-ethyl-8-methoxy-1-methyl-4H-[1,2,4]triazolo[4,3-a][1,4]benzodiazepine-4-acetamide), OTX015 ((6S)-4-(4-Chlorophenyl)-N-(4-hydroxyphenyl)-2,3,9-trimethyl-6H-thieno[3,2-f][1,2,4]triazolo[4,3-a][1,4]diazepine-6-acetamide), CPI-0610 (Pelabresib), GSK525762 (Molibresib), ABBV-075 (Mivebresib), FT-1101 ((2S)-5-(Cyclobutyl)-3,4-dihydro-2-methyl-6-[1-(4-piperidinyl)-1H-pyrazol-4-yl]-1(2H)-quinolinyl]cyclopropylmethanone), INCB057643 (2,2,4-trimethyl-8-(6-methyl-7-oxo-1H-pyrrolo[2,3-c]pyridin-4-yl)-6-methylsulfonyl-1,4-benzoxazin-3-one), ZEN003694, GSK2820151, CC-90010 (Trotabresib), PLX51107 (4-[6-(3,5-dimethyl-1,2-oxazol-4-yl)-1-[(1S)-1-pyridin-2-ylethyl]pyrrolo[3,2-b]pyridin-3-yl]benzoic acid), ABBV-744 (N-ethyl-4-[2-(4-fluoro-2,6-dimethylphenoxy)-5-(2-hydroxypropan-2-yl)phenyl]-6-methyl-7-oxo-1H-pyrrolo[2,3-c]pyridine-2-carboxamide), BAY1238097 ((4S)-7,8-dimethoxy-N,4-dimethyl-1-[4-(4-methylpiperazin-1-yl)phenyl]-4,5-dihydro-2,3-benzodiazepine-3-carboxamide), B1894999 (6-[1-benzyl-6-(4-meth-

ylpiperazin-1-yl)benzimidazol-2-yl]-N,3-dimethyl-[1,2,4]triazolo[4,3-a]pyrazin-8-amine), BMS-986158 (2-[3-(3,5-dimethyltriazol-4-yl)-5-[(S)-oxan-4-yl(phenyl)methyl]pyrido[3,2-b]indol-7-yl]propan-2-ol), GS-5829 ([2-cyclopropyl-6-(3,5-dimethyl-1,2-oxazol-4-yl)-1H-benzimidazol-4-yl]-dipyridin-2-ylmethanol), INCB054328, R06870810 (2-[(9S)-7-(4-chlorophenyl)-4,5,13-trimethyl-3-thia-1,8,11,12-tetrazatricyclo[8.3.0.0.2,6]trideca-2(6),4,7,10,12-pentaen-9-yl]-N-[3-(4-methylpiperazin-1-yl)propyl]acetamide), and in particular combinations thereof.

[0049] In particular, the at least one BET proteins inhibitor is selected in the group consisting of RVX-208, I-BET-762, OTX015, CPI-0610, GSK525762, ABBV-075, FT-1101, INCB057643, CC-90010, PLX51107, ABBV-744, BAY1238097, B1894999, BMS-986158, GS-5829, R06870810, and combinations thereof.

[0050] Advantageously, the at least one G4 stabilizer and at least one BET proteins inhibitor/histone deacetylase inhibitor are used simultaneously, separately, or sequentially.

[0051] Advantageously, the at least one G4 stabilizer and at least one BET proteins inhibitor are associated with a pharmaceutical acceptable vehicle. The pharmaceutical acceptable vehicle is as defined above.

[0052] It is within the skills of a physician to determine the specific therapeutically effective dosage regimen, as this dosage regimen will be dependent upon a variety of factors including, but not limited to: the severity of the multiple myeloma; the age; the body weight; general health; the sex; the diet; the time course of administration; the route of administration; the duration of the treatment; the drugs that are concomitantly administered in combination with the pharmaceutical composition within the scope of the present invention.

[0053] In some embodiments, the dosage regimen of each of said at least one G4 stabilizer and at least one BET proteins inhibitor ranges from about 0.0001 mg to about 1,000 mg per adult, per day. Preferably, the individual is administered with an amount of about 0.0001, 0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 7.5, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 150, 250, 500 and 750 mg of each of said at least one G4 stabilizer and at least one BET proteins inhibitor/histone deacetylase inhibitor in order to adjust the dosage regimen that is the most suitable to a particular individual in need of the treatment.

[0054] A pharmaceutical composition within the scope of the present invention may contain from about 0.01 mg to about 500 mg of each of the said at least one G4 stabilizer and at least one BET proteins inhibitor/histone deacetylase inhibitor, preferably from about 1 mg to about 100 mg of said at least one G4 stabilizer and at least one BET proteins inhibitor/histone deacetylase inhibitor.

[0055] In a preferred embodiment, an effective amount of each of the said at least one G4 stabilizer and at least one BET proteins inhibitor is routinely administered to an individual in need thereof, at a dosage regimen from about 0.0002 mg/kg to about 20 mg/kg of body weight per day, in particular from about 0.001 mg/kg to 7 mg/kg of body weight per day.

[0056] The optimal amount of the said at least one G4 stabilizer and at least one BET proteins inhibitor to be comprised in a pharmaceutical dosage unit according to the invention may be easily adapted by the one skilled in the art using routine known protocols or methods.

[0057] Said composition may be administered by any suitable route, i.e. including, but not limited to, an oral, sublingual, subcutaneous, intraperitoneal, intramuscular, intravenous, intrathecal and rectal administration.

[0058] In particular, the composition further comprises at least one nitrogen mustard.

[0059] The inventors unexpectedly discovered that stabilizing G4 structures potentializes the nitrogen mustards cytotoxicity on multiple myeloma cells. The inventors have identified that the combination between at least one G4 stabilizer and at least one nitrogen mustard is able to induce apoptosis or to inhibit cell cycle of primary cells from individuals afflicted by a multiple myeloma with a synergistic effect.

[0060] Nitrogen mustards nonspecific DNA alkylating agents. They are cytotoxic organic compounds with the chloroethylamine ($\text{Cl}(\text{CH}_2)_2\text{NR}_2$) functional group which form cyclic aminium ions (aziridinium rings) by intramolecular displacement of the chloride by the amine nitrogen. This aziridinium group then alkylates DNA once it is attacked by the N-7 nucleophilic centre on the guanine base.

[0061] Particularly, the at least one nitrogen mustard is selected in the group consisting of Chlormethine, Chlorambucil, Melphalan, Cyclophosphamide, Ifosfamide, Estramustine, Prednimustine, Bendamustine, Melphalan flufenamide (Melflufen) and combinations thereof. In particular, the nitrogen mustard is Melphalan.

[0062] Advantageously, the at least one G4 stabilizer and at least one nitrogen mustard inhibitor are used simultaneously, separately, or sequentially.

[0063] Advantageously, the at least one G4 stabilizer and at least one nitrogen mustard are associated with a pharmaceutical acceptable vehicle. The pharmaceutical acceptable vehicle is as defined above.

[0064] It is within the skills of a physician to determine the specific therapeutically effective dosage regimen, as this dosage regimen will be dependent upon a variety of factors including, but not limited to: the severity of the multiple myeloma; the age; the body weight; general health; the sex; the diet; the time course of administration; the route of administration; the duration of the treatment; the drugs that are concomitantly administered in combination with the pharmaceutical composition within the scope of the present invention.

[0065] In some embodiments, the dosage regimen of each of said at least one G4 stabilizer and at least one nitrogen mustard ranges from about 0.0001 mg to about 1,000 mg per adult, per day.

[0066] Preferably, the individual is administered with an amount of about 0.0001, 0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 7.5, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 150, 250, 500 and 750 mg of each of said at least one G4 stabilizer and at least one nitrogen mustard in order to adjust the dosage regimen that is the most suitable to a particular individual in need of the treatment.

[0067] A pharmaceutical composition within the scope of the present invention may contain from about 0.01 mg to about 500 mg of each of the said at least one G4 stabilizer and at least one nitrogen mustard, preferably from about 1 mg to about 100 mg of said at least one G4 stabilizer and at least one nitrogen mustard.

[0068] The optimal amount of the said at least one G4 stabilizer and at least one nitrogen mustard to be comprised in a pharmaceutical dosage unit according to the invention

may be easily adapted by the one skilled in the art using routine known protocols or methods.

[0069] Said composition may be administered by any suitable route, i.e. including, but not limited to, an oral, sublingual, subcutaneous, intraperitoneal, intramuscular, intravenous, intrathecal and rectal administration.

[0070] In another embodiment, the individual has been prognosed with a poor outcome.

[0071] The term “outcome” refers to the survival, the relapse or the death of the individual. The outcome may relate to disease-free survival (DFS), progression-free survival (PFS), event free survival (EFS) or overall survival (OS).

[0072] Illustratively, a “poor outcome” may refer to a disease relapse or death of the individual. The disease relapse of multiple myeloma may be defined as an increase in circulating monoclonal peak, an increase in medullary plasmacytosis and the return of one or more clinical evidence (hypercalcemia, renal failure, anaemia and bone tissue injuries). In particular, a “poor outcome” may refer to an overall survival after 1000 days below 80%, after 1500 days below 60% or even after 2500 days below 40%. Particularly, a “poor outcome” may refer to an event free survival after 1000 days below 50%. A “poor outcome” may also refer to a median overall survival of the individual around 1600 days or 55 months. Oppositely, a “good outcome” may refer to survival of the individual, with or without relapse episode. In particular, a “good outcome” may refer to an overall survival after 1000 days over 80%, after 1500 days over 80% or even after 2500 days over 60%. Particularly, a “good outcome” may refer to an event free survival after 1000 days over 50%.

[0073] A “good outcome” may also refer to a median overall survival over 2500 days.

[0074] The term “overall survival” refers to the length of time from either the date of diagnosis or the beginning of treatment for a disease, such as cancer, that patients diagnosed with the disease are still alive.

[0075] The term “event free survival” refers to the length of time after primary treatment for a cancer during which the patient remains free of certain complications or events that the treatment was intended to prevent or delay. These events may include the return of the cancer or the onset of certain symptoms, such as bone pain from cancer that has spread to the bone.

[0076] The term “disease-free survival” refers to the length of time after primary treatment for a cancer during which the patient survives without any signs or symptoms of that cancer.

[0077] The term “progression-free survival” refers to the length of time during and after the treatment of a disease, such as cancer, that a patient lives with the disease but it does not get worse.

[0078] In particular, the poor outcome of the individual is in vitro determined by carrying out the following steps:

[0079] a) measuring in a biological sample from said individual, the expression level of n genes, wherein n is an integer from 3 to 9 and wherein the n genes are selected from a group of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;

[0080] b) calculating a score TRC_{score} according to the following formula

$$TRC_{score} = \sum_{i=1}^n \beta_i \times C_i$$

[0081] wherein β_i represents the regression β coefficient reference value of one of the n genes selected at step a), wherein $C_i=1$ if the expression level of the said gene is higher than an expression level of reference and $C_i=-1$ if the expression level of the said gene is lower than or equal to ELR_i ,

[0082] c) prognosing that

[0083] said individual with a score TCR_{score} higher than a reference value TRC_{ref} is likely to have a poor outcome, or

[0084] said individual with a score TCR_{score} lower than a reference value TRC_{ref} is likely to have a good outcome.

[0085] In particular, the poor outcome of the individual is in vitro determined by carrying out the following steps:

[0086] a) measuring in a biological sample from said individual, the expression level of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;

[0087] b) calculating a score TRC_{score} according to the following formula

$$TRC_{score} = \sum_{i=1}^9 \beta_i \times C_i$$

[0088] wherein β_i represents the regression β coefficient reference value for the gene of nucleic acid sequence SEQ ID NO: i ,

[0089] wherein $C_i=1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is higher than an expression level of reference and $C_i=-1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is lower than or equal to ELR_i ,

[0090] c) prognosing that

[0091] said individual with a score TCR_{score} higher than a reference value TRC_{ref} is likely to have a poor outcome, or

[0092] said individual with a score TCR_{score} lower than a reference value TRC_{ref} is likely to have a good outcome,

[0093] wherein the reference value TRC_{ref} is of -0.39535 .

Step a)

[0094] The 466 different combinations of n genes from the group of 9 genes of the invention are listed in Table 2 below.

TABLE 2

Comb.	Genes Symbol		
1	BRIP1	DDX1	DDX23
2	BRIP1	DDX1	EXOSC5
3	BRIP1	DDX1	FANCD2
4	BRIP1	DDX1	HNRNPU
5	BRIP1	DDX1	PRMT5
6	BRIP1	DDX1	SRPK2
7	BRIP1	DDX1	XRN2
8	BRIP1	DDX23	EXOSC5
9	BRIP1	DDX23	FANCD2
10	BRIP1	DDX23	HNRNPU
11	BRIP1	DDX23	PRMT5
12	BRIP1	DDX23	SRPK2
13	BRIP1	DDX23	XRN2
14	BRIP1	EXOSC5	FANCD2
15	BRIP1	EXOSC5	HNRNPU
16	BRIP1	EXOSC5	PRMT5
17	BRIP1	EXOSC5	SRPK2
18	BRIP1	EXOSC5	XRN2
19	BRIP1	FANCD2	HNRNPU
20	BRIP1	FANCD2	PRMT5
21	BRIP1	FANCD2	SRPK2
22	BRIP1	FANCD2	XRN2
23	BRIP1	HNRNPU	PRMT5
24	BRIP1	HNRNPU	SRPK2
25	BRIP1	HNRNPU	XRN2
26	BRIP1	PRMT5	SRPK2
27	BRIP1	PRMT5	XRN2
28	BRIP1	SRPK2	XRN2
29	DDX1	DDX23	EXOSC5
30	DDX1	DDX23	FANCD2
31	DDX1	DDX23	HNRNPU
32	DDX1	DDX23	PRMT5
33	DDX1	DDX23	SRPK2
34	DDX1	DDX23	XRN2
35	DDX1	EXOSC5	FANCD2
36	DDX1	EXOSC5	HNRNPU
37	DDX1	EXOSC5	PRMT5
38	DDX1	EXOSC5	SRPK2

TABLE 2-continued

Comb.	Genes Symbol		
39	DDX1	EXOSC5	XRN2
40	DDX1	FANCD2	HNRNPU
41	DDX1	FANCD2	PRMT5
42	DDX1	FANCD2	SRPK2
43	DDX1	FANCD2	XRN2
44	DDX1	HNRNPU	PRMT5
45	DDX1	HNRNPU	SRPK2
46	DDX1	HNRNPU	XRN2
47	DDX1	PRMT5	SRPK2
48	DDX1	PRMT5	XRN2
49	DDX1	SRPK2	XRN2
50	DDX23	EXOSC5	FANCD2
51	DDX23	EXOSC5	HNRNPU
52	DDX23	EXOSC5	PRMT5
53	DDX23	EXOSC5	SRPK2
54	DDX23	EXOSC5	XRN2
55	DDX23	FANCD2	HNRNPU
56	DDX23	FANCD2	PRMT5
57	DDX23	FANCD2	SRPK2
58	DDX23	FANCD2	XRN2
59	DDX23	HNRNPU	PRMT5
60	DDX23	HNRNPU	SRPK2
61	DDX23	HNRNPU	XRN2
62	DDX23	PRMT5	SRPK2
63	DDX23	PRMT5	XRN2
64	DDX23	SRPK2	XRN2
65	EXOSC5	FANCD2	HNRNPU
66	EXOSC5	FANCD2	PRMT5
67	EXOSC5	FANCD2	SRPK2
68	EXOSC5	FANCD2	XRN2
69	EXOSC5	HNRNPU	PRMT5
70	EXOSC5	HNRNPU	SRPK2
71	EXOSC5	HNRNPU	XRN2
72	EXOSC5	PRMT5	SRPK2
73	EXOSC5	PRMT5	XRN2
74	EXOSC5	SRPK2	XRN2
75	FANCD2	HNRNPU	PRMT5
76	FANCD2	HNRNPU	SRPK2
77	FANCD2	HNRNPU	XRN2
78	FANCD2	PRMT5	SRPK2
79	FANCD2	PRMT5	XRN2
80	FANCD2	SRPK2	XRN2
81	HNRNPU	PRMT5	SRPK2
82	HNRNPU	PRMT5	XRN2
83	HNRNPU	SRPK2	XRN2
84	PRMT5	SRPK2	XRN2
85	BRIP1	DDX1	DDX23 EXOSC5
86	BRIP1	DDX1	DDX23 FANCD2
87	BRIP1	DDX1	DDX23 HNRNPU
88	BRIP1	DDX1	DDX23 PRMT5
89	BRIP1	DDX1	DDX23 SRPK2
90	BRIP1	DDX1	DDX23 XRN2
91	BRIP1	DDX1	EXOSC5 FANCD2
92	BRIP1	DDX1	EXOSC5 HNRNPU
93	BRIP1	DDX1	EXOSC5 PRMT5
94	BRIP1	DDX1	EXOSC5 SRPK2
95	BRIP1	DDX1	EXOSC5 XRN2
96	BRIP1	DDX1	FANCD2 HNRNPU
97	BRIP1	DDX1	FANCD2 PRMT5
98	BRIP1	DDX1	FANCD2 SRPK2
99	BRIP1	DDX1	FANCD2 XRN2
100	BRIP1	DDX1	HNRNPU PRMT5
101	BRIP1	DDX1	HNRNPU SRPK2
102	BRIP1	DDX1	HNRNPU XRN2
103	BRIP1	DDX1	PRMT5 SRPK2
104	BRIP1	DDX1	PRMT5 XRN2
105	BRIP1	DDX1	SRPK2 XRN2
106	BRIP1	DDX23	EXOSC5 FANCD2
107	BRIP1	DDX23	EXOSC5 HNRNPU
108	BRIP1	DDX23	EXOSC5 PRMT5
109	BRIP1	DDX23	EXOSC5 SRPK2
110	BRIP1	DDX23	EXOSC5 XRN2
111	BRIP1	DDX23	FANCD2 HNRNPU
112	BRIP1	DDX23	FANCD2 PRMT5
113	BRIP1	DDX23	FANCD2 SRPK2

TABLE 2-continued

Comb.	Genes Symbol			
114	BRIP1	DDX23	FANCD2	XRN2
115	BRIP1	DDX23	HNRNPU	PRMT5
116	BRIP1	DDX23	HNRNPU	SRPK2
117	BRIP1	DDX23	HNRNPU	XRN2
118	BRIP1	DDX23	PRMT5	SRPK2
119	BRIP1	DDX23	PRMT5	XRN2
120	BRIP1	DDX23	SRPK2	XRN2
121	BRIP1	EXOSC5	FANCD2	HNRNPU
122	BRIP1	EXOSC5	FANCD2	PRMT5
123	BRIP1	EXOSC5	FANCD2	SRPK2
124	BRIP1	EXOSC5	FANCD2	XRN2
125	BRIP1	EXOSC5	HNRNPU	PRMT5
126	BRIP1	EXOSC5	HNRNPU	SRPK2
127	BRIP1	EXOSC5	HNRNPU	XRN2
128	BRIP1	EXOSC5	PRMT5	SRPK2
129	BRIP1	EXOSC5	PRMT5	XRN2
130	BRIP1	EXOSC5	SRPK2	XRN2
131	BRIP1	FANCD2	HNRNPU	PRMT5
132	BRIP1	FANCD2	HNRNPU	SRPK2
133	BRIP1	FANCD2	HNRNPU	XRN2
134	BRIP1	FANCD2	PRMT5	SRPK2
135	BRIP1	FANCD2	PRMT5	XRN2
136	BRIP1	FANCD2	SRPK2	XRN2
137	BRIP1	HNRNPU	PRMT5	SRPK2
138	BRIP1	HNRNPU	PRMT5	XRN2
139	BRIP1	HNRNPU	SRPK2	XRN2
140	BRIP1	PRMT5	SRPK2	XRN2
141	DDX1	DDX23	EXOSC5	FANCD2
142	DDX1	DDX23	EXOSC5	HNRNPU
143	DDX1	DDX23	EXOSC5	PRMT5
144	DDX1	DDX23	EXOSC5	SRPK2
145	DDX1	DDX23	EXOSC5	XRN2
146	DDX1	DDX23	FANCD2	HNRNPU
147	DDX1	DDX23	FANCD2	PRMT5
148	DDX1	DDX23	FANCD2	SRPK2
149	DDX1	DDX23	FANCD2	XRN2
150	DDX1	DDX23	HNRNPU	PRMT5
151	DDX1	DDX23	HNRNPU	SRPK2
152	DDX1	DDX23	HNRNPU	XRN2
153	DDX1	DDX23	PRMT5	SRPK2
154	DDX1	DDX23	PRMT5	XRN2
155	DDX1	DDX23	SRPK2	XRN2
156	DDX1	EXOSC5	FANCD2	HNRNPU
157	DDX1	EXOSC5	FANCD2	PRMT5
158	DDX1	EXOSC5	FANCD2	SRPK2
159	DDX1	EXOSC5	FANCD2	XRN2
160	DDX1	EXOSC5	HNRNPU	PRMT5
161	DDX1	EXOSC5	HNRNPU	SRPK2
162	DDX1	EXOSC5	HNRNPU	XRN2
163	DDX1	EXOSC5	PRMT5	SRPK2
164	DDX1	EXOSC5	PRMT5	XRN2
165	DDX1	EXOSC5	SRPK2	XRN2
166	DDX1	FANCD2	HNRNPU	PRMT5
167	DDX1	FANCD2	HNRNPU	SRPK2
168	DDX1	FANCD2	HNRNPU	XRN2
169	DDX1	FANCD2	PRMT5	SRPK2
170	DDX1	FANCD2	PRMT5	XRN2
171	DDX1	FANCD2	SRPK2	XRN2
172	DDX1	HNRNPU	PRMT5	SRPK2
173	DDX1	HNRNPU	PRMT5	XRN2
174	DDX1	HNRNPU	SRPK2	XRN2
175	DDX1	PRMT5	SRPK2	XRN2
176	DDX23	EXOSC5	FANCD2	HNRNPU
177	DDX23	EXOSC5	FANCD2	PRMT5
178	DDX23	EXOSC5	FANCD2	SRPK2
179	DDX23	EXOSC5	FANCD2	XRN2
180	DDX23	EXOSC5	HNRNPU	PRMT5
181	DDX23	EXOSC5	HNRNPU	SRPK2
182	DDX23	EXOSC5	HNRNPU	XRN2
183	DDX23	EXOSC5	PRMT5	SRPK2
184	DDX23	EXOSC5	PRMT5	XRN2
185	DDX23	EXOSC5	SRPK2	XRN2
186	DDX23	FANCD2	HNRNPU	PRMT5
187	DDX23	FANCD2	HNRNPU	SRPK2
188	DDX23	FANCD2	HNRNPU	XRN2

TABLE 2-continued

Comb.	Genes Symbol				
189	DDX23	FANCD2	PRMT5	SRPK2	
190	DDX23	FANCD2	PRMT5	XRN2	
191	DDX23	FANCD2	SRPK2	XRN2	
192	DDX23	HNRNPU	PRMT5	SRPK2	
193	DDX23	HNRNPU	PRMT5	XRN2	
194	DDX23	HNRNPU	SRPK2	XRN2	
195	DDX23	PRMT5	SRPK2	XRN2	
196	EXOSC5	FANCD2	HNRNPU	PRMT5	
197	EXOSC5	FANCD2	HNRNPU	SRPK2	
198	EXOSC5	FANCD2	HNRNPU	XRN2	
199	EXOSC5	FANCD2	PRMT5	SRPK2	
200	EXOSC5	FANCD2	PRMT5	XRN2	
201	EXOSC5	FANCD2	SRPK2	XRN2	
202	EXOSC5	HNRNPU	PRMT5	SRPK2	
203	EXOSC5	HNRNPU	PRMT5	XRN2	
204	EXOSC5	HNRNPU	SRPK2	XRN2	
205	EXOSC5	PRMT5	SRPK2	XRN2	
206	FANCD2	HNRNPU	PRMT5	SRPK2	
207	FANCD2	HNRNPU	PRMT5	XRN2	
208	FANCD2	HNRNPU	SRPK2	XRN2	
209	FANCD2	PRMT5	SRPK2	XRN2	
210	HNRNPU	PRMT5	SRPK2	XRN2	
211	BRIP1	DDX1	DDX23	EXOSC5	FANCD2
212	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU
213	BRIP1	DDX1	DDX23	EXOSC5	PRMT5
214	BRIP1	DDX1	DDX23	EXOSC5	SRPK2
215	BRIP1	DDX1	DDX23	EXOSC5	XRN2
216	BRIP1	DDX1	DDX23	FANCD2	HNRNPU
217	BRIP1	DDX1	DDX23	FANCD2	PRMT5
218	BRIP1	DDX1	DDX23	FANCD2	SRPK2
219	BRIP1	DDX1	DDX23	FANCD2	XRN2
220	BRIP1	DDX1	DDX23	HNRNPU	PRMT5
221	BRIP1	DDX1	DDX23	HNRNPU	SRPK2
222	BRIP1	DDX1	DDX23	HNRNPU	XRN2
223	BRIP1	DDX1	DDX23	PRMT5	SRPK2
224	BRIP1	DDX1	DDX23	PRMT5	XRN2
225	BRIP1	DDX1	DDX23	SRPK2	XRN2
226	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU
227	BRIP1	DDX1	EXOSC5	FANCD2	PRMT5
228	BRIP1	DDX1	EXOSC5	FANCD2	SRPK2
229	BRIP1	DDX1	EXOSC5	FANCD2	XRN2
230	BRIP1	DDX1	EXOSC5	HNRNPU	PRMT5
231	BRIP1	DDX1	EXOSC5	HNRNPU	SRPK2
232	BRIP1	DDX1	EXOSC5	HNRNPU	XRN2
233	BRIP1	DDX1	EXOSC5	PRMT5	SRPK2
234	BRIP1	DDX1	EXOSC5	PRMT5	XRN2
235	BRIP1	DDX1	EXOSC5	SRPK2	XRN2
236	BRIP1	DDX1	FANCD2	HNRNPU	PRMT5
237	BRIP1	DDX1	FANCD2	HNRNPU	SRPK2
238	BRIP1	DDX1	FANCD2	HNRNPU	XRN2
239	BRIP1	DDX1	FANCD2	PRMT5	SRPK2
240	BRIP1	DDX1	FANCD2	PRMT5	XRN2
241	BRIP1	DDX1	FANCD2	SRPK2	XRN2
242	BRIP1	DDX1	HNRNPU	PRMT5	SRPK2
243	BRIP1	DDX1	HNRNPU	PRMT5	XRN2
244	BRIP1	DDX1	HNRNPU	SRPK2	XRN2
245	BRIP1	DDX1	PRMT5	SRPK2	XRN2
246	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU
247	BRIP1	DDX23	EXOSC5	FANCD2	PRMT5
248	BRIP1	DDX23	EXOSC5	FANCD2	SRPK2
249	BRIP1	DDX23	EXOSC5	FANCD2	XRN2
250	BRIP1	DDX23	EXOSC5	HNRNPU	PRMT5
251	BRIP1	DDX23	EXOSC5	HNRNPU	SRPK2
252	BRIP1	DDX23	EXOSC5	HNRNPU	XRN2
253	BRIP1	DDX23	EXOSC5	PRMT5	SRPK2
254	BRIP1	DDX23	EXOSC5	PRMT5	XRN2
255	BRIP1	DDX23	EXOSC5	SRPK2	XRN2
256	BRIP1	DDX23	FANCD2	HNRNPU	PRMT5
257	BRIP1	DDX23	FANCD2	HNRNPU	SRPK2
258	BRIP1	DDX23	FANCD2	HNRNPU	XRN2
259	BRIP1	DDX23	FANCD2	PRMT5	SRPK2
260	BRIP1	DDX23	FANCD2	PRMT5	XRN2
261	BRIP1	DDX23	FANCD2	SRPK2	XRN2
262	BRIP1	DDX23	HNRNPU	PRMT5	SRPK2
263	BRIP1	DDX23	HNRNPU	PRMT5	XRN2

TABLE 2-continued

Comb.	Genes Symbol				
264	BRIP1	DDX23	HNRNPU	SRPK2	XRN2
265	BRIP1	DDX23	PRMT5	SRPK2	XRN2
266	BRIP1	EXOSC5	FANCD2	HNRNPU	PRMT5
267	BRIP1	EXOSC5	FANCD2	HNRNPU	SRPK2
268	BRIP1	EXOSC5	FANCD2	HNRNPU	XRN2
269	BRIP1	EXOSC5	FANCD2	PRMT5	SRPK2
270	BRIP1	EXOSC5	FANCD2	PRMT5	XRN2
271	BRIP1	EXOSC5	FANCD2	SRPK2	XRN2
272	BRIP1	EXOSC5	HNRNPU	PRMT5	SRPK2
273	BRIP1	EXOSC5	HNRNPU	PRMT5	XRN2
274	BRIP1	EXOSC5	HNRNPU	SRPK2	XRN2
275	BRIP1	EXOSC5	PRMT5	SRPK2	XRN2
276	BRIP1	FANCD2	HNRNPU	PRMT5	SRPK2
277	BRIP1	FANCD2	HNRNPU	PRMT5	XRN2
278	BRIP1	FANCD2	HNRNPU	SRPK2	XRN2
279	BRIP1	FANCD2	PRMT5	SRPK2	XRN2
280	BRIP1	HNRNPU	PRMT5	SRPK2	XRN2
281	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU
282	DDX1	DDX23	EXOSC5	FANCD2	PRMT5
283	DDX1	DDX23	EXOSC5	FANCD2	SRPK2
284	DDX1	DDX23	EXOSC5	FANCD2	XRN2
285	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5
286	DDX1	DDX23	EXOSC5	HNRNPU	SRPK2
287	DDX1	DDX23	EXOSC5	HNRNPU	XRN2
288	DDX1	DDX23	EXOSC5	PRMT5	SRPK2
289	DDX1	DDX23	EXOSC5	PRMT5	XRN2
290	DDX1	DDX23	EXOSC5	SRPK2	XRN2
291	DDX1	DDX23	FANCD2	HNRNPU	PRMT5
292	DDX1	DDX23	FANCD2	HNRNPU	SRPK2
293	DDX1	DDX23	FANCD2	HNRNPU	XRN2
294	DDX1	DDX23	FANCD2	PRMT5	SRPK2
295	DDX1	DDX23	FANCD2	PRMT5	XRN2
296	DDX1	DDX23	FANCD2	SRPK2	XRN2
297	DDX1	DDX23	HNRNPU	PRMT5	SRPK2
298	DDX1	DDX23	HNRNPU	PRMT5	XRN2
299	DDX1	DDX23	HNRNPU	SRPK2	XRN2
300	DDX1	DDX23	PRMT5	SRPK2	XRN2
301	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5
302	DDX1	EXOSC5	FANCD2	HNRNPU	SRPK2
303	DDX1	EXOSC5	FANCD2	HNRNPU	XRN2
304	DDX1	EXOSC5	FANCD2	PRMT5	SRPK2
305	DDX1	EXOSC5	FANCD2	PRMT5	XRN2
306	DDX1	EXOSC5	FANCD2	SRPK2	XRN2
307	DDX1	EXOSC5	HNRNPU	PRMT5	SRPK2
308	DDX1	EXOSC5	HNRNPU	PRMT5	XRN2
309	DDX1	EXOSC5	HNRNPU	SRPK2	XRN2
310	DDX1	EXOSC5	PRMT5	SRPK2	XRN2
311	DDX1	FANCD2	HNRNPU	PRMT5	SRPK2
312	DDX1	FANCD2	HNRNPU	PRMT5	XRN2
313	DDX1	FANCD2	HNRNPU	SRPK2	XRN2
314	DDX1	FANCD2	PRMT5	SRPK2	XRN2
315	DDX1	HNRNPU	PRMT5	SRPK2	XRN2
316	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5
317	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2
318	DDX23	EXOSC5	FANCD2	HNRNPU	XRN2
319	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2
320	DDX23	EXOSC5	FANCD2	PRMT5	XRN2
321	DDX23	EXOSC5	FANCD2	SRPK2	XRN2
322	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2
323	DDX23	EXOSC5	HNRNPU	PRMT5	XRN2
324	DDX23	EXOSC5	HNRNPU	SRPK2	XRN2
325	DDX23	EXOSC5	PRMT5	SRPK2	XRN2
326	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2
327	DDX23	FANCD2	HNRNPU	PRMT5	XRN2
328	DDX23	FANCD2	HNRNPU	SRPK2	XRN2
329	DDX23	FANCD2	PRMT5	SRPK2	XRN2
330	DDX23	HNRNPU	PRMT5	SRPK2	XRN2
331	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
332	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
333	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
334	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
335	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2
336	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
337	BRIP1	DDX1	DDX23	EXOSC5	FANCD2 HNRNPU
338	BRIP1	DDX1	DDX23	EXOSC5	FANCD2 PRMT5

TABLE 2-continued

Comb.	Genes Symbol					
339	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	SRPK2
340	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	XRN2
341	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5
342	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	SRPK2
343	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	XRN2
344	BRIP1	DDX1	DDX23	EXOSC5	PRMT5	SRPK2
345	BRIP1	DDX1	DDX23	EXOSC5	PRMT5	XRN2
346	BRIP1	DDX1	DDX23	EXOSC5	SRPK2	XRN2
347	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	PRMT5
348	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	SRPK2
349	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	XRN2
350	BRIP1	DDX1	DDX23	FANCD2	PRMT5	SRPK2
351	BRIP1	DDX1	DDX23	FANCD2	PRMT5	XRN2
352	BRIP1	DDX1	DDX23	FANCD2	SRPK2	XRN2
353	BRIP1	DDX1	DDX23	HNRNPU	PRMT5	SRPK2
354	BRIP1	DDX1	DDX23	HNRNPU	PRMT5	XRN2
355	BRIP1	DDX1	DDX23	HNRNPU	SRPK2	XRN2
356	BRIP1	DDX1	DDX23	PRMT5	SRPK2	XRN2
357	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5
358	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	SRPK2
359	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	XRN2
360	BRIP1	DDX1	EXOSC5	FANCD2	PRMT5	SRPK2
361	BRIP1	DDX1	EXOSC5	FANCD2	PRMT5	XRN2
362	BRIP1	DDX1	EXOSC5	FANCD2	SRPK2	XRN2
363	BRIP1	DDX1	EXOSC5	HNRNPU	PRMT5	SRPK2
364	BRIP1	DDX1	EXOSC5	HNRNPU	PRMT5	XRN2
365	BRIP1	DDX1	EXOSC5	HNRNPU	SRPK2	XRN2
366	BRIP1	DDX1	EXOSC5	PRMT5	SRPK2	XRN2
367	BRIP1	DDX1	FANCD2	HNRNPU	PRMT5	SRPK2
368	BRIP1	DDX1	FANCD2	HNRNPU	PRMT5	XRN2
369	BRIP1	DDX1	FANCD2	HNRNPU	SRPK2	XRN2
370	BRIP1	DDX1	FANCD2	PRMT5	SRPK2	XRN2
371	BRIP1	DDX1	HNRNPU	PRMT5	SRPK2	XRN2
372	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5
373	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2
374	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	XRN2
375	BRIP1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2
376	BRIP1	DDX23	EXOSC5	FANCD2	PRMT5	XRN2
377	BRIP1	DDX23	EXOSC5	FANCD2	SRPK2	XRN2
378	BRIP1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2
379	BRIP1	DDX23	EXOSC5	HNRNPU	PRMT5	XRN2
380	BRIP1	DDX23	EXOSC5	HNRNPU	SRPK2	XRN2
381	BRIP1	DDX23	EXOSC5	PRMT5	SRPK2	XRN2
382	BRIP1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2
383	BRIP1	DDX23	FANCD2	HNRNPU	PRMT5	XRN2
384	BRIP1	DDX23	FANCD2	HNRNPU	SRPK2	XRN2
385	BRIP1	DDX23	FANCD2	PRMT5	SRPK2	XRN2
386	BRIP1	DDX23	HNRNPU	PRMT5	SRPK2	XRN2
387	BRIP1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
388	BRIP1	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
389	BRIP1	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
390	BRIP1	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
391	BRIP1	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2
392	BRIP1	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
393	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5
394	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2
395	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	XRN2
396	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2
397	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	XRN2
398	DDX1	DDX23	EXOSC5	FANCD2	SRPK2	XRN2
399	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2
400	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	XRN2
401	DDX1	DDX23	EXOSC5	HNRNPU	SRPK2	XRN2
402	DDX1	DDX23	EXOSC5	PRMT5	SRPK2	XRN2
403	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2
404	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	XRN2
405	DDX1	DDX23	FANCD2	HNRNPU	SRPK2	XRN2
406	DDX1	DDX23	FANCD2	PRMT5	SRPK2	XRN2
407	DDX1	DDX23	HNRNPU	PRMT5	SRPK2	XRN2
408	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
409	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
410	DDX1	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
411	DDX1	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
412	DDX1	EXOSC5	HARNPU	PRMT5	SRPK2	XRN2
413	DDX1	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2

TABLE 2-continued

Comb.	Genes Symbol						
414	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2	
415	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2	
416	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2	
417	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2	
418	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2	
419	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2	
420	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2	
421	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5
422	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2
423	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	XRN2
424	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2
425	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	XRN2
426	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	SRPK2	XRN2
427	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2
428	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	XRN2
429	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	SRPK2	XRN2
430	BRIP1	DDX1	DDX23	EXOSC5	PRMT5	SRPK2	XRN2
431	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2
432	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	XRN2
433	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	SRPK2	XRN2
434	BRIP1	DDX1	DDX23	FANCD2	PRMT5	SRPK2	XRN2
435	BRIP1	DDX1	DDX23	HNRNPU	PRMT5	SRPK2	XRN2
436	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
437	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
438	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
439	BRIP1	DDX1	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
440	BRIP1	DDX1	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2
441	BRIP1	DDX1	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
442	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
443	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
444	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
445	BRIP1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
446	BRIP1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2
447	BRIP1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
448	BRIP1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
449	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2
450	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	XRN2
451	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2	XRN2
452	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2	XRN2
453	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2	XRN2
454	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
455	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
456	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2	XRN2
457	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5 SRPK2
458	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5 XRN2
459	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	SRPK2 XRN2
460	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	PRMT5	SRPK2 XRN2
461	BRIP1	DDX1	DDX23	EXOSC5	HNRNPU	PRMT5	SRPK2 XRN2
462	BRIP1	DDX1	DDX23	FANCD2	HNRNPU	PRMT5	SRPK2 XRN2
463	BRIP1	DDX1	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2 XRN2
464	BRIP1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2 XRN2
465	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5	SRPK2 XRN2
466	BRIP1	DDX1	DDX23	EXOSC5	FANCD2	HNRNPU	PRMT5 SRPK2 XRN2

[0095] In the invention, a “biological sample” refers to a biological sample obtained, reached, collected or isolated from an individual, in vivo or in situ. Such samples may be, but not limited to, organs, tissues, fractions thereof and cells isolated from an individual. For example, suitable biological samples include but are not limited to a cell culture, a cell line, a tissue biopsy such as a bone marrow aspirate, a biological fluid such as a blood, pleural effusion or a serum sample, and the like. An advantageous biological sample includes but is not limited to a blood sample, a tissue biopsy, including a bone marrow aspirate. The biological sample as defined in the invention may be a crude sample, or may be purified to various degrees prior to storage, processing, or measurement.

[0096] The expression level of the n genes is measured by well-known protocol in the art. These methods are for

instance, DNA-CHIPS containing probesets of said n genes, so that an expression level can be determined for each of said n genes. Other methods can be used, such that quantitative PCR strategy by using specific couples of primers for each of said n genes, with either a specific Taqman probe for each of said 9 genes, or SYBR® compounds.

[0097] Advantageously, the expression level can be evaluated by measuring the expression level of mRNA for each of the n genes. This measurement may be carried out by using the well-known techniques available in the art. In this case, mRNA may be extracted, for example using lytic enzymes or chemical solutions or extracted by commercially available nucleic-acid-binding resins following the manufacturer’s instructions. Extracted mRNA may be subsequently detected by hybridization, such as Northern blot, and/or amplification, such as quantitative or semi-quantitative RT-

PCR. Other methods of amplification include ligase chain reaction (LCR), transcription-mediated amplification (TMA), strand displacement amplification (SDA) and nucleic acid sequence-based amplification (NASBA).

[0098] Advantageously, the level of mRNA expression for each of the n genes may be measured by the mean of quantification of the cDNA synthesized from said mRNA, as a template, by one reverse transcriptase. Methods for determining the quantity of mRNA by microarrays or by RNA sequencing may also be used.

[0099] In certain embodiments, complexes between the double-stranded nucleic acids resulting from amplification and fluorescent SYBR® molecules may be obtained and then the fluorescence signal generated by the SYBR® molecules complexed with the said amplified nucleic acids may be measured.

[0100] The determination of the expression level of said n genes could be to carry out by a northern blot analysis, but due to the low efficiency of such a method, the skilled person will prefer the quantitative methods to obtain a more precise expression level of said n genes.

[0101] The group of 9 genes of the invention with the corresponding probe set and CDS (or one of the CDS if the gene expression different variants) are represented in Table 1.

TABLE 1

Probe set	Gene symbole	CDS SEQ ID	β coefficient	Expression level of reference (ELR)
235609_at	BRIP1	SEQ ID NO:1	0.25527251	192
201241_at	DDX1	SEQ ID NO:2	0.2787536	3182
201440_at	DDX23	SEQ ID NO:3	0.23044892	1015
218481_at	EXOSC5	SEQ ID NO:4	0.25527251	1412
242560_at	FANCD2	SEQ ID NO:5	0.25527251	288
200594_x_at	HNRNPU	SEQ ID NO:6	0.32221929	11244
217786_at	PRMT5	SEQ ID NO:7	0.23044892	1616
203182_s_at	SRPK2	SEQ ID NO:8	0.32221929	966
223002_s_at	XRN2	SEQ ID NO:9	0.32221929	767

Step b)

[0102] The regression β coefficient reference values may be easily determined by the skilled man in the art for each gene of nucleic acid sequence SEQ ID NO: i using a Cox model. The Cox model is based on a modelling approach to the analysis of survival data. The purpose of the model is to simultaneously explore the effects of several variables on survival. The Cox model is a well-recognised statistical technique for analysing survival data. When it is used to analyse the survival of patients in a clinical trial, the model allows us to isolate the effects of treatment from the effects of other variables. The logrank test cannot be used to explore (and adjust for) the effects of several variables, such as age and disease duration, known to affect survival. Adjustment for variables that are known to affect survival may improve the precision with which the inventors can estimate the treatment effect. The regression method introduced by Cox is used to investigate several variables at a time. It is also known as proportional hazards regression analysis. Briefly,

the procedure models or regresses the survival times (or more specifically, the so-called hazard function) on the explanatory variables. The hazard function is the probability that an individual will experience an event (for example, death) within a small-time interval, given that the individual has survived up to the beginning of the interval. It can therefore be interpreted as the risk of dying at time t . The quantity $h_0(t)$ is the baseline or underlying hazard function and corresponds to the probability of dying (or reaching an event) when all the explanatory variables are zero. The baseline hazard function is analogous to the intercept in ordinary regression (since $\exp(0)=1$). The regression coefficient β gives the proportional change that can be expected in the hazard, related to changes in the explanatory variables. The coefficient β is estimated by a statistical method called maximum likelihood. In survival analysis, the hazard ratio (HR) (Hazard Ratio= $\exp(\beta)$) is the ratio of the hazard rates corresponding to the conditions described by two sets of explanatory variables. For example, in a drug study, the treated population may die at twice the rate per unit time as the control population. The hazard ratio would be 2, indicating higher hazard of death from the treatment.

[0103] In one embodiment, the regression β coefficient reference values are described in Table 1.

[0104] In the abovementioned formula, $C_i=1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is higher than an expression level of reference ELR $_i$ or $C_i=-1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is lower than or equal to ELR $_i$, wherein “ i ” is as defined above for each DNA repair pathway group.

[0105] Expression level of reference ELR $_i$ may consist of “cut-off” values. A cut-off value is a value of expression of a gene that allows to separate the individuals according to their outcome (good or bad) for a given gene. If the measured expression value of the gene of an individual is higher than the cut-off value, the individual has a good outcome and vice-versa. The cut-off values may be obtained using Maxstat algorithm.

[0106] For example, each reference cut-off value ELR $_i$ for each gene may be determined by carrying out a method comprising the steps of:

- [0107]** a) providing a collection of samples from patients suffering from acute myeloid leukemia;
- [0108]** b) determining the expression level of the relevant gene for each sample contained in the collection provided at step a);
- [0109]** c) ranking the samples according to said expression level;
- [0110]** d) classifying said samples in pairs of subsets of increasing, respectively decreasing, number of members ranked according to their expression level;
- [0111]** e) providing, for each sample provided at step a), information relating to the actual clinical outcome for the corresponding cancer patient (i.e. the duration of the disease-free survival (DFS), or the event free survival (EFS), or the overall survival (OS) or both);
- [0112]** f) for each pair of subsets of tumour tissue samples, obtaining a Kaplan Meier percentage of survival curve;
- [0113]** g) for each pair of subsets of tumour tissue samples calculating the statistical significance (p value) between both subsets;

[0114] h) selecting as reference value ELR for the expression level, the value of expression level for which the p value is the smallest.

[0115] For example, the expression level of a gene has been assessed for 100 samples of 100 patients. The 100 samples are ranked according to the expression level of the gene. Sample 1 has the highest expression level and sample 100 has the lowest expression level. A first grouping provides two subsets: on one side sample Nr 1 and on the other side the 99 other samples. The next grouping provides on one side samples 1 and 2 and on the other side the 98 remaining samples etc., until the last grouping: on one side samples 1 to 99 and on the other side sample Nr 100. According to the information relating to the actual clinical outcome for the corresponding cancer patient, Kaplan Meier curves are prepared for each of the 99 groups of two subsets. Also, for each of the 99 groups, the p value between both subsets was calculated. The reference value ELR_i is then selected such as the discrimination based on the criterion of the minimum p value is the strongest. In other terms, the expression level corresponding to the boundary between both subsets for which the p value is minimum is considered as the reference value. It should be noted that according to the experiments made by the inventors, the reference value ELR_i is not necessarily the median value of expression levels.

[0116] In one embodiment, the ELR_i are described in Table 1.

Step c)

[0117] In case said individual has a score TRC_{score} higher than the reference value TRC_{ref} , said individual is likely to have an overall survival after 1000 days below 80%. By “below 80%”, it is meant in the invention 79%, 78%, 77%, 76%, 75%, 74%, 73%, 72%, 71%, 70%, 69%, 68%, 67%, 66%, 65%, 64%, 63%, 62%, 61%, 60%, 59%, 58%, 57%, 56%, 55%, 54%, 53%, 52%, 51%, 50%, 49%, 48%, 47%, 46%, 45%, 44%, 43%, 42%, 41%, 40%, 39%, 38%, 37%, 36%, 35%, 34%, 33%, 32%, 31%, 30%, 29%, 28%, 27%, 26%, 25%, 24%, 23%, 22%, 21%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% and 0%. In particular, the said individual is likely to have an overall survival after 1000 days below after 1500 days below 60%. Particularly, the said individual is likely to have an overall survival after 2500 days below 40%. Alternatively or additionally, the said individual is likely to have an event free survival after 1000 days below 50%.

[0118] In case said individual has a score TRC_{score} lower than the reference value TRC_{ref} , the said individual is likely to have an overall survival after 1000 days equal or over 80%. By “equal or over 80%”, it is meant in the invention 80%, 79%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and 100%. In particular, the said individual is likely to have an overall survival after 1500 days over 80%. Particularly, the said individual is likely to have an overall survival after 2500 days equal or over 60%. By “equal or over 60%” it is meant in the invention 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and 100%. Alternatively or additionally, the said individual is likely to have an event free

survival after 1000 days equal or over 50%. By “equal or over 50%” it is meant in the invention 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% and 100%.

[0119] The invention also relates to a method for treating an individual afflicted by a multiple myeloma, the method comprising administering an effective amount of a composition comprising a, i.e., at least one, G-quadruplex (G4) stabilizer and/or a, i.e., at least one histone deacetylase (HDAC) inhibitor and/or a, i.e., at least one bromodomain and extraterminal (BET) proteins inhibitor and/or a, i.e., at least one, nitrogen mustard.

[0120] The composition may comprise

[0121] solely a, i.e., at least one G4 stabilizer,

[0122] solely a, i.e., at least one HDAC inhibitor,

[0123] solely a, i.e., at least one BET proteins inhibitor,

[0124] solely at least one nitrogen mustard,

[0125] a, i.e., at least one G4 stabilizer and a, i.e., at least one HDAC inhibitor,

[0126] a, i.e., at least one G4 stabilizer and a, i.e., at least one BET proteins inhibitor,

[0127] at least one G4 stabilizer and at least one nitrogen mustard,

[0128] at least one G4 stabilizer and at least one HDAC inhibitor and at least one nitrogen mustard,

[0129] at least one G4 stabilizer and at least one BET proteins inhibitor and at least one nitrogen mustard,

[0130] a, i.e., at least one HDAC inhibitor and a, i.e., at least one BET protein inhibitor,

[0131] a, i.e., at least one G4 stabilizer and a, i.e., at least one HDAC inhibitor and a, i.e., at least one BET proteins inhibitor, or

[0132] at least one G4 stabilizer and at least one HDAC inhibitor and at least one BET protein inhibitor and at least one nitrogen mustard.

[0133] The term “effective amount”, also referred to herein as “dosage”, refers to an amount of the composition that is suitable to treat a multiple myeloma in an individual.

[0134] In particular, the said method comprises the following steps:

[0135] a) identifying the patient able to respond to a composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor and/or at least one nitrogen mustard by the following sub steps:

[0136] i) measuring in a biological sample from said individual, the expression level of n genes, wherein n is an integer from 3 to 9 and wherein the n genes are selected from a group of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;

[0137] ii) calculating a score TRC_{score} according to the following formula

$$TRC_{score} = \sum_{i=1}^n \beta_i \times C_i$$

- [0138] wherein β_i represents the regression β coefficient reference value of one of the n genes selected at step a), wherein $C_i=1$ if the expression level of the said gene is higher than an expression level of reference and $C_i=-1$ if the expression level of the said gene is lower than or equal to ELR_i ,
- [0139] iii) confirming that the said individual has a score TRC_{score} higher than a reference value TRC_{ref} ,
- [0140] iv) concluding that the individual is able to respond a composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor,
- [0141] b) administering to the individual the composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor.
- [0142] In particular, the said method comprises the following steps:
- [0143] a) identifying the patient able to respond to a composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor and/or at least one nitrogen mustard by the following sub steps:
- [0144] i) measuring in a biological sample from said individual, the expression level of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;
- [0145] ii) calculating a score TRC_{score} according to the following formula
- $$TRC_{score} = \sum_{i=1}^9 \beta_i \times C_i$$
- [0146] wherein β_i represents the regression β coefficient reference value for the gene of nucleic acid sequence SEQ ID NO: i ,
- [0147] wherein $C_i=1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is higher than an expression level of reference $C_i=-1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO: i is lower than or equal to ELR_i ,
- [0148] iii) confirming that the said individual has a score TRC_{score} higher than a reference value TRC_{ref} wherein the reference value TRC_{ref} is of -0.39535 ,
- [0149] iv) concluding that the individual is able to respond a composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor,
- [0150] b) administering to the individual the composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor.
- [0151] The invention also relates to the use of a composition comprising at least one HDACi and at least one BET inhibitor for its use in the treatment of an individual afflicted by a multiple myeloma.
- [0152] Advantageously, the at least one HDACi and at least one at least one BET inhibitor are used simultaneously, separately, or sequentially.
- [0153] Advantageously, the at least one HDACi and at least one BET proteins inhibitor are associated with a pharmaceutical acceptable vehicle. The pharmaceutical acceptable vehicle is as defined above.
- [0154] It is within the skills of a physician to determine the specific therapeutically effective dosage regimen, as this dosage regimen will be dependent upon a variety of factors including, but not limited to: the severity of the multiple myeloma; the age; the body weight; general health; the sex; the diet; the time course of administration; the route of administration; the duration of the treatment; the drugs that are concomitantly administered in combination with the pharmaceutical composition within the scope of the present invention.
- [0155] In some embodiments, the dosage regimen of each of said at least one HDACi and at least one BET proteins inhibitor ranges from about 0.0001 mg to about 1,000 mg per adult, per day. Preferably, the individual is administered with an amount of about 0.0001, 0.0005, 0.001, 0.005, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 7.5, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 15 100, 250, 500 and 750 mg of each of said at least one G4 stabilizer and at least one histone deacetylase inhibitor in order to adjust the dosage regimen that is the most suitable to a particular individual in need of the treatment.
- [0156] A pharmaceutical composition within the scope of the present invention may contain from about 0.01 mg to about 500 mg of each of the said at least one G4 stabilizer and at least one histone deacetylase inhibitor, preferably from about 1 mg to about 100 mg of said at least one G4 stabilizer and at least one histone deacetylase inhibitor.
- [0157] In a preferred embodiment, an effective amount of each of the said at least one HDACi and at least one BET proteins inhibitor is routinely administered to an individual in need thereof, at a dosage regimen from about 0.0002 mg/kg to about 20 mg/kg of body weight per day, in particular from about 0.001 mg/kg to 7 mg/kg of body weight per day.
- [0158] The optimal amount of the said at least one HDACi and at least one BET proteins inhibitor to be comprised in a pharmaceutical dosage unit according to the invention may be easily adapted by the one skilled in the art using routine known protocols or methods.
- [0159] Said composition may be administered by any suitable route, i.e. including, but not limited to, an oral, sublingual, buccal, subcutaneous, transdermal, topical, intraperitoneal, intramuscular, intravenous, subdermal, intrathecal and intranasal and rectal administration.
- [0160] The invention further relates to a pharmaceutical composition comprising at least one G-quadruplex (G4) stabilizer and at least one histone deacetylase inhibitor and/or at least one bromodomain and extraterminal (BET) proteins inhibitor and/or at least one nitrogen mustard.
- [0161] The invention also relates to a pharmaceutical composition comprising at least one histone deacetylase inhibitor and at least one bromodomain and extraterminal (BET) proteins inhibitor.

[0162] The invention further relates to a method for prognosing the outcome of an individual afflicted by a multiple myeloma, wherein said method comprises the following steps:

- [0163] a) measuring in a biological sample from said individual, the expression level of n genes, wherein n is an integer from 3 to 9 and wherein the n genes are selected from a group of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;
- [0164] b) calculating a score TRC_{score} according to the following formula

$$TRC_{score} = \sum_{i=1}^n \beta_i \times C_i$$

[0165] wherein β_i represents the regression β coefficient reference value of one of the n genes selected at step a), wherein $C_i=1$ if the expression level of the said gene is higher than an expression level of reference and $C_i=-1$ if the expression level of the said gene is lower than or equal to ELRi,

[0166] c) prognosing that

[0167] said individual with a score TRC_{score} higher than a reference value TRC_{ref} is likely to have a bad outcome, or

[0168] said individual with a score TRC_{score} lower than a reference value TRC_{ref} is likely to have a good outcome.

[0169] The different combinations of n genes from the group of 9 genes of the invention are listed in Table 2 above.

[0170] In particular, n equal to 9 and the reference value TRC_{ref} is of -0.39535 .

[0171] The invention also relates to a composition comprising at least one G-quadruplex (G4) stabilizer and/or at least one histone deacetylase inhibitor and/or at least one at least one bromodomain and extraterminal (BET) proteins inhibitor and/or at least one nitrogen mustard for its use in a method for treating an individual afflicted by multiple myeloma, wherein the individual has been prognosed with a poor outcome according to the above-mentioned method.

LEGENDS TO THE FIGURES

[0172] FIG. 1 represents the expression of R-loops and TRC resolution genes for the different stage of Memory B cell (MBC) to plasma cell (PC) differentiation. The genes significantly overexpressed in PrePBs compared to MBCs, PBs and PCs were determined with a SAM (Significance Analysis of Microarrays) multiclass analysis (False discovery rate (FDR)=0), identifying 41 unique genes. An unsupervised hierarchical clustering was completed and the normalized expression value for each gene is indicated in shade of grey, with black representing high expression and white representing low expression. MBC1 to 5 represent 5 different samples of memory B cells. PrePB1 to 5 represent 5 different samples of preplasmablasts. PB1 to 5 represents 5 different samples of plasmablasts, PC1 to 5 represent 5 different samples of Plasma cells.

[0173] FIG. 2 represents Kaplan Meier curves showing the percentage of survival of the patients of the training cohort vs time (days) based on different genes' expression. In all graphs, curve 1 represents low gene expression results and curve 2 represents high expression results. The gene expres-

sion prognostic value was determined using Maxstat R algorithm. FIG. 1A represents the results obtained for BRIP1 expression, wherein for curve 1 $n=92$, for curve 2 $n=253$, p -value=0.0012 and hazard ratio (HR)=1.8. FIG. 1B represents the results obtained for DDX1 expression, wherein for curve 1 $n=264$, for curve 2 $n=81$, p -value=0.0011 and HR=1.9. FIG. 1C represents the results obtained for DDX23, wherein for curve 1 $n=304$, for curve 2 $n=41$, p -value=0.044 and HR=1.7. FIG. 1D represents the results obtained for EXOSC5 expression, wherein for curve 1 $n=259$, for curve 2 $n=86$, p -value=0.0019 and HR=1.8. FIG. 1E represents the results obtained for FANCD2 expression, wherein for curve 1 $n=168$, for curve 2 $n=177$, p -value=0.0037 and HR=1.8.

[0174] FIG. 3 represents Kaplan Meier curves showing the percentage of survival of the patients of the training cohort vs time (days) based on different genes' expression. In all graphs, curve 1 represents low gene expression results and curve 2 represents high expression results. The gene expression prognostic value was determined using Maxstat R algorithm. FIG. 3A represents the results obtained for HNRNPU expression, wherein for curve 1 $n=304$, for curve 2 $n=41$, p -value=0.0014 and HR=2.1. FIG. 3B represents the results obtained for PRMT5 expression, wherein for curve 1 $n=268$, for curve 2 $n=77$, p -value=0.013 and HR=1.7. FIG. 3C represents the results obtained for SRPK2 expression, wherein for curve 1 $n=87$, for curve 2 $n=258$, p -value=0.0031 and HR=2.1. FIG. 3D represents the results obtained for XRN2 expression, wherein for curve 1 $n=298$, for curve 2 $n=47$, p -value=0.0014 and HR=2.1.

[0175] FIGS. 4A and 4C represents Kaplan Meier curves showing the percentage of event free survival of the patients of the training cohort (FIG. 4A) or of the validation cohort (FIG. 4C) vs time (days) based on the TRC_{score} . FIGS. 4B and 4D represents Kaplan Meier curves showing the percentage of survival of the patients of the training cohort (FIG. 4A) or of the validation cohort (FIG. 4C) vs time (days) based on the TRC_{score} . Patients of the different cohort were split in two groups: high risk patients ($n=119$ for the training cohort; $n=71$ for the validation cohort) and low risk patients ($n=226$ for the training cohort; $n=135$ for the validation cohort). High risk patients have a TRC_{score} higher than TRC_{ref} and low risk patients have a TRC_{score} lower or equal to TRC_{ref} . In all graphs, curve 1 represents the results for high-risk patients, and curve 2 represents the results for low risk patients. The parameters to compute the TRCs score of patients in the HM cohort and the proportions delineating the two prognostic groups were those defined with the TT2 cohort.

[0176] FIG. 5A represents the IC50 of Panobinostat (nM) as function of the TRC_{score} of 11 Huma Myeloma Cell Lines (HMCLs): JJN3, OPM2, RPM18266, LP1, AMO1, SKMM2, XG1, XG2, XG6, XG7, XG12, XG19, XG20 and XG21. The represented best-fit line has a $r=-0.55$ with a p -value=0.05. FIG. 5B represents tumor cell viability as function of the TRC_{score} of 12 primary myeloma cells. The represented best-fit line has a $r=-0.67$ with a p -value lower than 0.01. Mononuclear cells from tumour samples of 12 patients with MM were cultured for 4 days in the presence of IL6 (2 ng/mL) with or without increasing concentrations (0 nM, 0.6 nM, 1.25 nM, 2.5 nM, 5 nM, 10 nM and 20 nM) of Panobinostat. At day 4 of culture, the count of viable MM cells was determined using CD138 and CD38 staining by flow cytometry.

[0177] FIG. 6A represents the IC₅₀ of Pyridostatin (nM) for 11 HMCLs: a is OPM2; b is XG12; c is XG21; d is XG19; e is RPM18266; f is LP1; g is XG6; h is JJN3; i is XG2; j is XG7; k: L363; l is XG1; m is AMO1; n is SKMM2 and o is XG20. IC₅₀ of Pyridostatin $\pm$ SD are representative of 3 independent experiments. NA: not reached. FIG. 6B represents the Pyridostatin toxicity on bone marrow cells (x-axis is the Pyridostatin concentration in μ M; y-axis is the cell count in percentage of the control (0 μ M of Pyridostatin)). The toxicity of Pyridostatin was assessed on primary bone marrow cells from one MM patient (dark grey bars), co-cultured with normal bone marrow microenvironment (light grey). The toxicity on MM cells and normal bone marrow cells was assessed by flow cytometry using CD138 and CD28 marker cells. FIG. 6C represents the Pyridostatin toxicity on bone marrow cells (x-axis is the Pyridostatin concentration in μ M; y-axis is the cell count in percentage of the control (0 μ M of Pyridostatin)). The toxicity of Pyridostatin was assessed on primary bone marrow cells from five MM patient (black bars) and co-cultured with normal bone marrow microenvironment (grey bars). The toxicity on MM cells and normal bone marrow cells was assessed by flow cytometry using CD138 and CD28 marker cells. Data are mean values of five independent experiments. P-value: * <0.05 ; ** <0.01 ; *** <0.001 using a student t-test for pairs.

[0178] FIGS. 7A and 7B represent Western blots. Doxy stands for doxycycline. PDS stands for Pyridostatin (concentration in μ M). XG7 cell line was transduced with a doxycycline-inducible lentivirus containing RNase H gene (XG7-RH). XG7-RH was cultured with or without doxycycline in absence or presence of Pyridostatin during 5 h or 24 h. Western blot membranes were stained with anti-Phospho-Chk2, anti-Phospho-P53, anti-P53, anti-PARP, anti-myc (FIG. 7A) and anti-p21, anti-p16, anti-p27 and anti- γ H2AX (FIG. 7B). Tubulin protein level was used as control.

[0179] FIGS. 8A and 8B represent the IC₅₀ of a bromodomain and extraterminal proteins inhibitor (FIG. 8A: I-BET-762; FIG. 8B: RVX-208) as function of the TRC_{score} of 11 Human Myeloma Cell Lines (HMCLs): JJN3, OPM2, RPM18266, LP1, AMO1, SKMM2, XG1, XG2, XG6, XG7, XG12, XG19, XG20 and XG21. For FIG. 8A, the represented best-fit line has a $r=-0.66$ with a p-value <0.05 . For FIG. 8B, the represented best-fit line has a $r=-0.88$ with a p-value <0.05 . HMCLs with high TRC score are significantly more sensitive to I-BET-762 and RVX-208 bromodomain and extraterminal proteins inhibitors treatment compared to cell lines with low TRC score.

[0180] FIGS. 9A and 9B represent cell viability of two HMCLs (9A: JJN3; 9B: XG7) treated with increasing concentration of Pyridostatin (x-axis in μ M) and Panobinostat (y-axis in μ M) for 4 days. The synergistic combination is represented in shade of grey (arbitrary unit), with black representing high synergism and white representing high antagonism.

[0181] FIG. 10 represents cell viability of XG2 cell line treated with increasing concentration of Pyridostatin (x-axis in μ M) and Panobinostat (y-axis in μ M) for 4 days. The synergistic combination is represented in shade of grey (arbitrary unit), with black representing high synergism and white representing high antagonism.

[0182] FIG. 11 represents the toxicity of Pyridostatin alone, Panobinostat alone and the combination of Pyridostatin and Panobinostat on bone marrow cells. X-axis is the concentration in μ M of the drug: 0: control; A: Pyridostatin

at 1.1; B: Panobinostat at 2.5; C: Pyridostatin at 1.1 and Panobinostat at 2.5. Y-axis is the cell count in percentage of the control (0 μ M of drug)). The toxicity was assessed on primary bone marrow cells from one MM patient cultured alone (black bars) and co-cultured with normal bone marrow microenvironment (grey bars). The toxicity on MM cells and normal bone marrow cells was assessed by flow cytometry using CD138 and CD28 marker cells.

[0183] FIG. 12 represents cell viability of XG7 cell line treated with increasing concentration of Pyridostatin (x-axis in μ M) and I-BET-762 (y-axis in μ M) for 4 days. The synergistic combination is represented in shade of grey (arbitrary unit), with black representing high synergism and white representing high antagonism.

[0184] FIG. 13 represents cell viability of XG1 cell line (A) and XG2 cell line (B) treated with increasing concentration of Pyridostatin (x-axis in μ M) and Melphan (y-axis in μ M) for 4 days. The synergistic combination is represented in shade of grey (arbitrary unit), with black representing high synergism and white representing high antagonism.

[0185] FIG. 14 represents cell viability of XG2 cell line (A) and XG2 Melphalan-resistant (MeIR) cell line (B) treated with increasing concentration of Pyridostatin (x-axis in μ M) and Melphan (y-axis in μ M) for 4 days. The synergistic combination is represented in shade of grey (arbitrary unit), with black representing high synergism and white representing high antagonism.

[0186] FIG. 15 represents Melphalan IC₅₀ on XG2 (A) and XG2-MeIR (B) cell lines. Grey bars relate to the respective myeloma cells treated with increasing concentration of Melphan (y-axis in μ M). For the XG2 cell line, Melphalan IC₅₀ was 0.6325 μ M. For the XG2 cell line, Melphalan IC₅₀ was 0.35 μ M. Black bars relate to the respective myeloma cells treated with increasing concentration of Melphalan (y-axis in μ M) and Pyridostatin at 1.25 μ M for 4 days. For the XG2 cell line, Melphalan IC₅₀ was 1.4875 μ M. For the XG2 cell line, Melphalan IC₅₀ was 0.6325 μ M. P-value: ** <0.01 ; *** <0.001 using a student t-test for pairs. "ns" stands for non-significant.

[0187] FIG. 16 relates to DDX23 depletion in XG7 cells. Two groups of XG7 cells were transduced with a doxycycline-inducible lentivirus containing an shRNA targeting DDX23 giving XG7-shDDX23-1 group cell and XG7-shDDX23-2 group cell. Each group cell was cultured without or with doxycycline for 24 or 48 hours. Protein detection was assayed using western blot analysis (A). Membranes were stained with anti-PARP, anti-DDX23, anti- γ H2AX and anti-tubulin. Tubulin protein level was used as control for assaying DDX23 protein level (B). Y-axis on FIG. 17B represents the ratio of DDX23 level on tubulin level. DDX23 mRNA level was assayed by RT-qPCR (C). On FIGS. 16B and 16C, black bars relate to XG7-shDDX23-1 group cell, and black bar relate to XG7-shDDX23-2 group cell. P-value: * <0.05 ; ** <0.01 ; *** <0.001 using a student t-test for pairs. "ns" stands for non-significant.

[0188] FIG. 17 relates to DDX1 depletion in XG7 cells. Two groups of XG7 cells were transduced with a doxycycline-inducible lentivirus containing an shRNA targeting DDX1 giving XG7-shDDX1-4 group cell and XG7-shDDX1-5 group cell. Each group cell was cultured without or with doxycycline for 24 or 48 hours. Protein detection was assayed using western blot analysis (A). Membranes were stained with anti-PARP, anti-DDX23, anti- γ H2AX and anti-tubulin. Tubulin protein level was used as control for

assaying DDX23 protein level (B). Y-axis on FIG. 17B represents the ratio of DDX1 level on tubulin level. DDX1 mRNA level was assayed by RT-qPCR (C). On FIGS. 16B and 16C, grey bars relate to XG7-shDDX1-4 group cell, and black bar relate to XG7-shDDX1-5 group cell. P-value: * <0.05 ; ** <0.01 ; *** <0.001 using a student t-test for pairs. “ns” stands for non-significant.

[0189] FIG. 18 represents cell viability of XG7 cells depleted for DDX23. XG7-shDDX23-1 and XG7-shDDX23-2 group cells were exposed to doxycycline and cell viability was analyzed by trypan blue assay. Results are those of three independent experiments. FIG. 18A is a diagram representing XG7-shDDX23-1 group cell viability after 2, 3 or 6 days of culture without (control in grey bars) or with (black bars) doxycycline at 1 $\mu\text{g/ml}$. Y-axis represents the ratio of cells relatively to the control. FIG. 18B is a diagram representing XG7-shDDX23-2 group cell viability after 2, 3 or 6 days of culture without (control in grey bars) or with (black bars) doxycycline at 1 $\mu\text{g/ml}$. Y-axis represents the ratio of cells relatively to the control. FIG. 18C represents the number of cells after 0, 2, 3 or 6 days of culture without or with doxycycline at 1 $\mu\text{g/ml}$ (1: XG7-shDDX23-1 without doxycycline; 2: XG7-shDDX23-1 with doxycycline; 3: XG7-shDDX23-2 without doxycycline; 4: XG7-shDDX23-2 with doxycycline). Cell count was analyzed by trypan blue assay. Results are mean of three independent experiments.

[0190] FIG. 19 represents cell viability of XG7 cells depleted for DDX1. XG7-shDDX1-4 and XG7-shDDX1-5 group cells were exposed to doxycycline and cell viability was analyzed by trypan blue assay. Results are those of three independent experiments. FIG. 19A is a diagram representing XG7-shDDX1-4 group cell viability after 2, 3 or 6 days of culture without (control in grey bars) or with (black bars) doxycycline at 1 $\mu\text{g/ml}$. Y-axis represents the ratio of cells relatively to the control. FIG. 19B is a diagram representing XG7-shDDX1-5 group cell viability after 2, 3 or 6 days of culture without (control in grey bars) or with (black bars) doxycycline at 1 $\mu\text{g/ml}$. Y-axis represents the ratio of cells relatively to the control. FIG. 19C represents the number of cells after 0, 2, 3 or 6 days of culture without or with doxycycline at 1 $\mu\text{g/ml}$ (1: XG7-shDDX1-4 without doxycycline; 2: XG7-shDDX1-4 with doxycycline; 3: XG7-shDDX1-5 without doxycycline; 4: XG7-shDDX1-5 with doxycycline). Cell count was analyzed by trypan blue assay. Results are mean of three independent experiments.

EXAMPLE

1. Patients and Method

Cell Lines

[0191] JJN3, OPM2, RPM18266, LP1, AMO1, SKMM2 human myeloma cell lines (HMCLs) were obtained from ATCC (Molsheim, France) and were cultured in RPMI 1640 medium, supplemented with 10% fetal calf serum (Biochrom, Berlin, Germany) and 2 mM L-Glutamine. The interleukin-6-dependent cell lines XG1, XG2, XG6, XG7, XG12, XG19, XG20, XG21 were obtained as previously described and maintained in the presence of 2 ng/mL recombinant IL-6 (R&D Systems, Oxon, UK), 10% fetal calf serum and 2 mM L-Glutamine.

Gene Expression Data

[0192] Affymetrix data from two independent cohorts of previously untreated patients with MM were used. These data are publicly available through the ArrayExpress database (E-MTAB-372). The training cohort (TT2 cohort) included 345 patients with MM from the University of Arkansas for Medical Sciences (UAMS, Little Rock, AR, USA). These data can be accessed at the online Gene Expression Omnibus (GSE2658). The validation cohort consisted of 206 patients with MM and was called Heidelberg-Montpellier (HM) cohort. This cohort also included five BMPC samples from healthy donors. Samples were obtained after signature of a written informed consent form in accordance with the Declaration of Helsinki and after approval by the Ethics Committees of Montpellier (DC-2008-417) and Heidelberg. After Ficoll-density gradient centrifugation, plasma cells were purified using anti-CD138 MACS microbeads (Miltenyi Biotech, Bergisch Gladbach, Germany). The clinical characteristics of the two cohorts have been previously described (Moreaux, J. et al, *Mol. Cancer Ther.* 11, 2685-2692 (2012) and Hose, D. et al., *Haematologica* 96, 87-95 (2011).

[0193] Affymetrix data from another cohort of patient treated with Daratumumab, the Daratumumab cohort, was also used. The clinical characteristics of this cohort have been described in Pochard et al., Integrative approach to find predictive markers of response to daratumumab in multiple myeloma, EHA Library, 2021.

Selection of Prognostic Genes

[0194] To establish gene expression (GE)-based risk scores, the inventors selected probe sets whose expression values were significantly associated with overall survival, using MaxStat R function and Benjamini Hochberg multiple testing correction (adjusted p-value <0.05).

Building Gene Expression-Based Risk Score ($\text{TCR}_{\text{Score}}$) on Training Cohort

[0195] The GE-based risk score was built as the sum of the beta coefficients weighted by +1 or -1 according to the patient signal above or below/equal the probe set MaxStat value. Patients from the training cohort were ranked according to increased prognostic score and for a given score value X, the difference in survival of patients with a prognostic score $\leq X$ or $> X$ was computed using MaxStat analysis. Survival analyses were assessed using Kaplan-Meier method, and survival curves were compared using log-rank test.

[0196] The build TCRscore was validated with the validation cohort, using the cutoff values determined for the training cohort. Survival analyses were assessed using Kaplan-Meier method, and survival curves were compared using log-rank test.

Patient Samples

[0197] All primary samples were collected with the approval of the institutional research board from Montpellier University hospital (DC2008-417) and in accordance with the Declaration of Helsinki. Patient's written informed consent was obtained. Mononuclear cells were isolated from total BM using Ficoll density gradient centrifugation (Nycomed, Zurich, Switzerland) and cultured in the presence of

1 ng/mL IL-6. Four days after treatment with increased concentrations of panobinostat or pyridostatin, the percentage CD138+ viable plasma cells and CD138- viable non-myeloma cells was determined by flow cytometry.

Compounds

[0198] Panobinostat, pyridostatin and I-BET-762 were obtained from Sellekchem (Munich, Germany). For in vitro studies, pyridostatin was dissolved in water, panobinostat and I-BET-762 in dimethyl sulfoxide (DMSO). Aliquots were stored at -20° C.

Cell Count and Cell Viability

[0199] Cells in culture were stained using trypan blue solution (Sigma-Aldrich, St. Louis MO, USA) and viable cells were counted under a light microscope. The CellTiter-Glo Luminescent Viability assay (Promega, Leiden, The Netherlands) was used to measure the effect on cell viability according to manufacturer's instructions, and the 50% inhibition (IC50) was determined using GraphPad Prism software (<http://www.graphpad.com/scientific-software/prism/>).

Drug Combination Study

[0200] The interaction between pyridostatin and panobinostat, pyridostatin and I-BET-762 and pyridostatin and Melphan was investigated with a concentration matrix test, in which increasing concentration of each single drug were assessed with all possible combinations of the other drugs. Cell viability was measured using ATP quantification to obtain the viability matrix. For each combination, the percentage of expected growing cells in case of effect independence was calculated according to the Bliss equation: $fuC = fuA \cdot fuB$, where fuC is the expected fraction of cells unaffected by the drug combination in the case of effect independence, and fuA and fuB are the fractions of cells unaffected by treatment A and B, respectively. The difference between the fraction of living cells in the cytotoxicity test and the fuC value was considered as an estimation of the interaction effect, with positive values indicating synergism and negative values antagonism.

Western Blot Analysis Western blot analysis was performed as previously described (Viziteu, E. et al., *Leukemia* 31, 2104-2113, 2017). Tubulin was used as loading control. All antibodies used were purchased from Cell Signaling, except for p21, p27, Myc, and phospho-Histone H2A.X. Myc antibody was obtained from Santa Cruz biotechnology (Texas, USA), phospho-Histone H2A.X antibody from Merck Millipore (Sigma-Aldrich), and p27 antibody was from Sigma-Aldrich.

Statistical Analysis

[0201] Statistical analysis was performed using GenomicScape web tool (<http://www.genomicscape.com>). A Mann-Whitney U test and 1-way analysis of variance was used to compare 2 groups or multiple groups, respectively. GEP data were normalized with MAS5 algorithm and analyzed with GenomicScape and R and Bioconductor programs. Probe sets were selected for prognostic significance using the Maxstat R function and the Benjamini Hochberg multiple testing correction. The difference in OS was assayed with a log-rank test, and survival curves were plotted using the Kaplan-Meier method. The prognostic value of BRIP1, DDX1, DDX23, EXOSC5, FANCD2, HNRNP, PRMT5,

SRPK2 and XRN2 was computed using a maximally selected rank test from the R package maxstat, which allowed us to determine the optimal cut point for continuous variables.

Inducible Depletion of DDX1 and DDX23

[0202] XG7 cells were transduced with shDDX1 or shDDX23 lentiviruses and stable transduced cells (XG7-shDDX23-1; XG7-shDDX23-2; XG7shDDX1-4; XG7-shDDX1-5) were obtained by adding 10 µg/ml puromycin. The expression of the shRNAs was induced by adding Doxycycline (1 µg/ml) 24 h after plating the cells.

RT-qPCR

[0203] Total RNA was extracted using the RNeasy Kit (Qiagen) and reverse transcribed with the Reverse Transcription Kit (Qiagen). The measurement of gene expression was performed using the Roche LC480 Sequence Detection System. For each primer, serial dilutions of standard cDNA were amplified to create a standard curve, and values of unknown samples were estimated relative to this standard curve in order to assess PCR efficiency. Ct values were obtained for rRNA 18S and the respective genes of interest during the log phase of the cycle. Gene expression was normalized to that of rRNA 18s ($\Delta Ct = Ct \text{ gene of interest} - Ct \text{ 18s}$) and compared with the values obtained for a known positive control using the following formula: $100/2^{\Delta\Delta Ct}$ where $\Delta\Delta Ct = \Delta Ct \text{ unknown} - \Delta Ct \text{ positive control}$.

2. Results

[0204] Multiple myeloma is clinically and biologically heterogeneous with several genetic alterations proposed as driving events in myelomagenesis, which are associated with growth advantage and cell cycle deregulation. Malignant PCs continue to produce elevated levels of immunoglobulins, underlying that transcription is highly active in those cells. Like non-transformed plasma cells, myeloma cells contain a well-developed endoplasmic reticulum (ER) and Golgi complex tailored to produce and secrete large amounts of immunoglobulins, underlying their characteristic morphology.

[0205] Besides, during normal PC differentiation, MBCs differentiate into pre-plasmablasts (prePBs), plasmablasts (PBs), early PCs, and finally, long-lived PCs that produce high Ig amounts. the pre-plasmablastic stage is associated with high proliferation following B cell activation (50% of cells in S-phase) and the start of Ig secretion. This led the inventors to hypothesis that conflicts between replication and transcription should be tightly managed at this specific stage to avoid tumorigenesis. Then, the inventors investigate the TRCs levels in malignant PCs.

Probe Sets Selection

[0206] The inventors first aimed to identify genes involved in TRCs resolution significantly overexpressed in prePBs during B to PC differentiation. A list set of 83 genes involved in TRCs resolution was defined using the review of the literature. In that purpose, the inventors used their own in vitro model of B to PC differentiation. Indeed, the inventors have shown that PC generation can be modeled using multi-step culture systems to reproduce the sequential cell differentiation occurring in the different organs/tissues in vivo. In this model, memory B cells (MBCs) differentiate

into pre-plasmablasts (prePBs), plasmablasts (PBs), early PCs and, finally, into long-lived PCs (LLPCs) (Jourdan M, Caraux A, Caron G, Robert N, Fiol G, Reme T, et al., *J Immunol* 2011; 187:3931-41; Jourdan M, Caraux A, De Vos J, Fiol G, Larroque M, Cognot C, et al., *Blood* 2009; 114:5173-81 and Jourdan M, Cren M, Robert N, Bollore K, Fest T, Duperray C, et al., *Leukemia* 2014, August; 28(8): 1647-56). Using Significance Analysis of Microarrays (SAM) multi-class analysis, the inventors found 41 genes significantly overexpressed in prePBs compared to MBCs, PBs and PCs with a false discovery rate <5% (FIG. 1). These results confirmed the hypothesis of the inventors that the TRCs resolution is highly relevant in prePBs.

[0207] Considering the potent role of TRCs resolution in MM cell adaptation to replication stress, the inventors sought to identify the TRCs resolution factors that are associated with a poor outcome in MM. They first compared the Affymetrix gene expression profiles of purified normal bone marrow PCs (n=5) and purified MM cells from newly diagnosed patients (n=206) using the GenomicScape web-tool (Kassambara, A. et al., *PLoS Comput. Biol.* 11, (2015) showed that 13 TRC resolution factors were significantly overexpressed in MM cell samples (fold change 2, false discovery rate <5%; 1000 permutations) (Data not shown).

[0208] Then, the inventors analysed the publicly available Affymetrix gene expression profiles (Gene Expression Omnibus, GSE2658) of the training cohort with the MaxStat R function and found out a high expression of 9 of these 13 TRC resolution factors was associated with poor MM outcome (FIGS. 2 and 3). This result suggested to the inventors that increasing TRCs could be a specific therapeutic strategy to kill malignant plasma cells. In that objective, the inventors gathered the prognostic value of these 9 genes within a GEP-based TRC resolution score (TRC_{score}). High TRC_{score} has been identified associated with a poor outcome (EFS and OS) in two independent cohorts (training and validation cohorts) of newly diagnosed MM patients treated by high dose therapy and autologous stem cell transplantation (FIG. 4).

HDAC Inhibitors' Sensitivity

[0209] By investigating the link between the TRC score and the response of MM cells to drugs using their own collection of human myeloma cell lines (HMCL), the inventors identified that HMCLs with high TRC score values are significantly more sensitive to Panobinostat histone deacetylase (HDAC) inhibitor, currently used in MM treatment at relapse (FIG. 5A). Without being bound by any theory, the inventors suggest that this result could be explained by that during chromatin opening, acetylation promotes R-loop formation that will then constitute obstacles to replication fork progression.

[0210] Furthermore, the inventors validated these results using primary MM cells from patients (n=12) co-cultured with their bone marrow microenvironment (FIG. 5B). A high TRC score value is associated with a higher toxicity of Panobinostat in primary MM cells. This underlines that the TRC score allows the identification of a subgroup of MM patients with a poor outcome that could benefit from Panobinostat HDACi treatment.

G-Quadruplex (G4) Stabilizers' Sensitivity in MM Cell Lines

[0211] The inventors then investigated the therapeutic interest of G4 stabilizers to kill MM cells. Treatment with

the G4 stabilizer Pyridostatin (PDS) was found to be associated with significant toxicity in 10 HMCLs with an IC50 ≤ 2 μ M whereas 5 cell lines demonstrated higher resistance to PDS (FIG. 6A). PDS inhibits MM cell growth and induces apoptosis with significant accumulation in G2/M phase of the cell cycle. Importantly, experiments confirmed the toxicity of PDS on primary MM cells of patients cocultured with their bone marrow microenvironment (FIG. 6B). Interestingly, the inventors confirmed that toxicity of PDS on primary malignant PCs of MM patients, without any significant toxicity on the co-cultured normal cells from the bone marrow microenvironment (FIG. 6C).

RNase H Expression in MM Cell Line

[0212] Inducible RNase H expression in MM cell line resulted in a significant reduction of DNA damage response after PDS treatment with a decreased phosphorylation of Chk2, p53 and a decrease of p21 and p16 expression (FIG. 7). Since RNase H specifically degrades RNA:DNA hybrids, this result supports the view that spontaneous replication stress and genomic instability are caused by R-loops in MM cells.

Bet Proteins Inhibitors' Sensitivity in MM Cell Lines

[0213] The inventors also found a correlation between HMCLs TRC score and the response to two Bromodomain and Extra-Terminal motif (BET) proteins inhibitors, I-BET-762 and RVX-208 (FIGS. 8A and 8B). Without being bound by any theory, the inventors suggest that this result could be explained by that BET proteins inhibition may increase R-loop formation and DNA damage.

[0214] Overall, these results demonstrate the ability of the score TRC_{score} to identify interesting drug combinations using PDS in MM cells.

Drugs Combination with G4 Stabilizer

[0215] The inventors investigated the sensitivity of HMCLs for different drugs combination with G4 stabilizer: HDAC inhibitors and BET proteins inhibitors.

[0216] The combination of PDS and Panobinostat was found to be a synergistic effect in three HMCLs (FIGS. 9 and 10). Moreover, it was showed the absence of too much toxicity of this combination on primary samples from MM patients (FIG. 11).

[0217] Furthermore, the synergistic effect of the combination of PDS with I-BET-762 was confirmed for one cell line (FIG. 12).

[0218] The combination of PDS and Melphalan was found to show a synergistic effect in two HMCLs (FIG. 13). These results underline that PDS can increase the DNA damage induced by Melphalan.

[0219] At that stage the inventors hypothesize that PDS could be used as a strategy to increase MM patients' response to Melphalan. This is demonstrated in FIGS. 14 and 15 where PDS is shown to re-sensitize the resistant cells (XG2-MelR) to Melphalan.

Ddx1 and Ddx23 Depletions

[0220] The inventors searched to validate the pathophysiological roles of two genes of the TRC score, namely DDX1 and DDX23. To that purpose, the inventors transduced XG7 cells with inducible shRNAs targeting either DDX1 or

DDX23. The inventors validated the depletion of the two proteins by western-blot (FIGS. 16A, B and 17A, B) and RT-qPCR (FIGS. 16C and 17C). The inventors found that the depletion of DDX1 affected cell viability and decreased cell proliferation (FIG. 19) whereas DDX23 depletion had a mild effect on cell viability and proliferation (FIG. 18).

[0221] Moreover, the inventors found that DDX1 and DDX23 depletions led to an increased phosphorylation of γ H2AX which points out spontaneous DNA damage formation (FIGS. 16A and 17A). Together, these results show that DDX1 and DDX23 are critical for MM cells survival and in TRCs prevention.

SEQUENCE LISTING

Sequence total quantity: 9
 SEQ ID NO: 1 moltype = DNA length = 3750
 FEATURE Location/Qualifiers
 source 1..3750
 mol_type = genomic DNA
 organism = Homo sapiens

SEQUENCE: 1

atgtcttcaa	tgtggtctga	atatacaatt	gggtgggtga	agatttactt	tccttataaa	60
gcttaccggt	cacagcttgc	tatgatgaat	tctattctca	gaggattaaa	cagcaagcaa	120
cattgtttgt	tggagagtcc	cacaggaagt	ggaaaaagct	tagccttact	ttgttctgct	180
ttagcatggc	aacaatctct	tagtgggaaa	ccagcagatg	agggcgtaag	tgaaaaagct	240
gaagtacaat	tgtcatgttg	ttgtgcatgc	cattcaaagg	attttacaaa	caatgacatg	300
aaccaaggaa	cttcacgtca	tttcaactat	ccaagcacac	caccttctga	aagaaatggc	360
acttcaccaa	cttgtcaaga	ctccctgaa	aaaaccactc	tggctgcaaa	gttatctgct	420
aagaacacagg	cattccatata	cagagatgaa	aatgatgatt	ttcaagtaga	gaagaaaaga	480
attcgaccct	tagaaactac	acagcagatt	agaaaaagct	attgctttgg	aacagaagta	540
cacaatttgg	atgcaaaagt	tgattcagga	aagactgtaa	aactcaactc	tccactggaa	600
aagataaaact	ctctttcggc	acagaaaacc	cctggccact	gttctagggt	ctgttggtct	660
actaaacaag	gaacacgtca	agagtcacgt	aataccatta	agaaggatca	tacagggaaa	720
tccaagatac	ccaaaaatata	ttttgggaca	cgcacacaca	agcagattgc	tcagattact	780
agagagctcc	ggagcagcgc	atattcaggg	gttccaatga	ctattcttct	cagcagggat	840
catacttggt	tccatcctga	ggtagtcggt	aacttcaaca	gaaatgagaa	gtgcatggaa	900
ttgctagatg	ggaaaaacgg	aaaactcctg	tatttttatc	atggagttca	taaaattagt	960
gatcagacaca	cattacagac	tttccaaggg	atgtgcaaa	cctgggatata	agaagaactt	1020
gtcagcctgg	ggaagaaact	aaaggcctgt	ccatattaca	cagcccagaa	actaatacaa	1080
gatgctgaca	tcataatttg	tccctacaac	tatcttctag	atgcacaaa	aagggaaaagt	1140
atggatttaa	ctctgaaaag	acaggtttgt	attttagatg	aagctcataa	catcgaggag	1200
tgtgctcggg	aatcagcaag	ttacagtgtg	acagaagttc	agcttcggtt	tgctcgggat	1260
gaactagata	gtatggtcaa	caataatata	aggaagaaag	atcatgaacc	cctacgagat	1320
gtgtgctgta	gcctcattaa	ttggttagaa	gcaaacgctg	aatatcttgt	agaaagagat	1380
tatgaatcag	cttgtaaaat	atggagtgtg	aatgaaatgc	tcttaacttt	acacaaaatg	1440
ggtatcacca	ctgctacttt	teccattttg	cagggcacatt	tttctgctgt	tcctcaaaaa	1500
gaggaaaaaa	ttccaccaat	ttatggtaaa	gaggaggcaa	gagaagtacc	tggtattagt	1560
gcatacaact	aaataatgct	taaaggactt	tttatggtac	ttgactatct	ttttaggcaa	1620
aatagcagat	ttgcagatga	ttataaaatt	gcgattcaac	agacttactc	ctggacaaat	1680
cagattgata	tttcagacaa	aaatgggttg	ttggttctac	caaaaaataa	gaaacgttca	1740
cgacagaaaa	ctgcagttca	tgtgctaaac	ttttggtgct	taaatccagc	tggtggcctt	1800
tcagatatta	atggcagaag	tcagaccatt	gttttgacat	ctggtagact	atcaccaagt	1860
aaatcctttt	cgtcagaact	tttggtttaca	tttactatcc	agctggaggc	taatacatac	1920
attaaaaaatt	cacaggtttg	ggttggtacc	attgggtcag	gccccagggg	tcggaatctc	1980
tgtgctacct	tcagaaatac	tgaaacattt	gagttccaag	atgaagtggg	agcacttttg	2040
ttatctgtgt	ccagactcgt	gagccaagga	attttgtggt	tcttgccatc	ttacaagtta	2100
ttagaaaaat	taaaagaacg	ttggctctct	actggtttat	ggcataatct	ggagttggtg	2160
aagacagtca	ttgtagaacc	acagggagga	gaaaaaacaa	attttgatga	attactgcag	2220
gtgtactatg	acgcaatcaa	atacaaaagg	gagaaagatg	gagctctcct	ggtagcagtt	2280
tgtcgtggta	aagttagtga	gggtctggat	ttctcagatg	acaatgcccg	tgctgtcata	2340
acaataggaa	ttccttttcc	aaatgtgaaa	gatctacagg	ttgaactaaa	acgacaatac	2400
aatgaccaca	attcaaaatt	gagaggtctt	ctacctggcc	gtcagtggtg	tgaaattcaa	2460
gcatacaggg	ccttaaacca	ggccttggtg	agatgtatta	gacacagaaa	tgattgggga	2520
gctcttattc	tagtggatga	tcgctttagg	aataacccaa	gtcgctatat	atctggactt	2580
tctaaatggg	tacggcagca	gattcagcac	cattcaacct	ttgaaagtgc	actggaatcc	2640
ttggctgaat	tttccaaaaa	gcatacaaaa	gttcttaatt	tatccataaa	ggacagaacc	2700
aataacacag	acaatgagtc	tacacttgaa	gtgacctctt	taaaagtacg	tacctcacct	2760
tatttactgg	aagcagcaag	tcactctatc	ccagaaaatt	ttgtggaaga	tgaagcaaa	2820
atatgtgtct	aggaactaca	gtgtcctaaa	attattacca	aaaattcacc	ttaccaagt	2880
agcattatct	ccagaaagga	gaaaaatgat	ccagtatctc	tggaagaagc	agggaaaagc	2940
gaaaaaattg	tgatttccag	atccacaagc	ccaactttca	acaaaacaa	aaagagagtt	3000
agctggtcaa	gcttttaatt	tttgggacag	tattttactg	gtaaaatacc	gaaggcaaca	3060
cctgagctcg	ggctcatcga	gaatagtgc	tctagtctct	cccgtttcaa	aacagagaag	3120
atggaaaagt	aaactgtttt	gcccttctact	gataaatgtg	aatcctcaaa	tctgacagta	3180
aacacatcgt	ttggatcatg	ccctcaatca	gaaaccatta	tttcatcatt	aaagattgat	3240
gccaccctta	ctagaaaaaa	tcattctgaa	catccgctct	gttctgaaga	agccctggat	3300
ccagacattg	aattgtctct	agtaagtga	gaagataaac	agtccacttc	aaatagagat	3360
tttgaaacag	aagcagaaga	tgaatctatc	tattttacac	ctgaacttta	tgatcctgaa	3420
gatcacagat	aagaaaaaaa	tgacctagct	gaaactgata	gaggaaatag	attggctaac	3480
aattcagatt	gcatttttag	taaaagacct	tttgaaatta	gaactataaa	agaagtagat	3540

-continued

tcagccagag	aagtgaagc	tgaggattgc	atagatacaa	agttgaatgg	aattctgcat	3600
attgaagaaa	gtaaaattga	tgacattgat	ggtaatgtaa	aaacaacttg	gataaatgaa	3660
ctggaaactgg	gaaaaactca	tgaatagaa	ataaagaact	ttaaaccatc	tccttccaaa	3720
aataaaggca	tgtttcctgg	ttttaagtaa				3750

SEQ ID NO: 2 moltype = DNA length = 2223
 FEATURE Location/Qualifiers
 source 1..2223
 mol_type = genomic DNA
 organism = Homo sapiens

SEQUENCE: 2

atggcggcct	tctccgagat	gggtgtaatg	cctgagattg	cacaagctgt	ggaagagatg	60
gattggctcc	tcccaactga	tatccaggct	gaatctatcc	cattgatctt	aggaggagggt	120
gatgtactta	tggctgcaga	aacaggaagt	ggcaaaactg	gtgcttttag	tattccagtt	180
atccagatag	tttatgaaac	tctgaaagac	caacagggaag	gcaaaaaagg	aaaaacaaca	240
atataaactg	gtgcttcagt	gctgaacaaa	tggcagatga	accatataga	cagaggatct	300
gcttttgcaa	tgggtgcaga	tggtctttgt	tgtcaaaagca	gagaagtaaa	ggaatggcat	360
gggtgtagag	ctactaaagg	attaatgaaa	gggaaacact	actatgaagt	atcctgtcat	420
gaccaagggt	tatgcagggt	cggtgggtct	accatgcagg	cctctttgga	cctaggtact	480
gacaagtttg	tatttggtct	tggtggaaaca	ggaaagaaat	cccataacaa	acaatttgat	540
aattatggag	aggaattcac	tatgcatgat	accattggat	gttacctgga	tatagataag	600
ggacatgtca	agttctccaa	aaatggaaaa	gatcttggtc	tggcatttga	aataccacca	660
catatgaaaa	accaagcctc	ctttcctgcc	tgtgttttga	agaatgctga	actgaaattt	720
aacttcgggtg	aagagggaatt	taagtttcca	ccaaaagatg	gctttgttgc	tctttccaaag	780
gcaccggatg	ttacatttgt	caaatcacag	cactcaggta	atgcacagggt	gacacaaaca	840
aagtttctcc	gcaatgtctc	gaaagctctc	attgttgaaac	cttcccggga	gttagctgaa	900
caaaccttga	acaacatcaa	gcagtttaag	aaatacattg	ataatcctaa	attaaggagg	960
cttctgataa	tggagggttg	tgcagcccgg	gatcagctct	ctgttttggga	aaatggagta	1020
gatatagttg	taggtactct	gggaagacta	gatgacttgg	tgtcaactgg	aaagctgaac	1080
ttatctcaag	ttagattcct	ggctcctggat	gaagctgatg	ggcttctttc	tcaaggttat	1140
tctgatttta	taaataggat	gcacaatcag	attcctcagg	ttacctctga	tggaaaaaga	1200
cttcagggtga	ttgtttgtct	tgccactttg	cattctttcg	atgtaaagaa	actgtccgag	1260
aagataatgc	attttctcat	atgggttgac	ttaaaaggag	aagactctgt	tccagatact	1320
gtacaccatg	ttgttgtccc	gtaaatcccc	aaaactgaca	gactctggga	aaggcttgga	1380
aagagccaca	ttagaactga	tgatgtacat	gcaaaaagata	acacaagacc	tgggtgctaat	1440
agtccagaga	tgtggtctga	agctattaaa	atcctgaaag	gggagtagtc	tgtccgggca	1500
atcaaggaaac	ataagatgga	tcaagcaatt	atcttctgta	gaaccaaaat	tgactgtgat	1560
aacttggagc	agttactttat	acaacaagga	ggaggacctg	ataaaaaagg	acaccagttc	1620
tcatgtgttt	gtcttcatgg	tgacagaaag	cctcatgaga	gaaagcaaaa	cttggaaga	1680
tttaagaaag	gagatgtaa	attcttgatt	tgcacagatg	tagctgctag	aggaattgat	1740
atccacgggtg	ttccttatgt	tataaatgtc	actctgcccg	atgaaaagca	aaactacgta	1800
catcgaaattg	gcagagtagg	aagagctgaa	aggatgggtc	tggcaatttc	cctgggtgca	1860
acagaaaaag	aaaaggtttg	gtaccatgta	tgtagcagcc	gtggaaaaag	gtgttataac	1920
acaagactca	aggaagatgg	aggctgtacc	atatggtaca	acgagatgca	gttactatct	1980
gagatagaag	aacacctgaa	ctgtaccatt	tctcaggttg	agccggatat	aaaggtacca	2040
gtggatgaat	ttgatgggaa	agttacctac	ggtcagaaaa	gggctgctgg	tgggtggaagc	2100
tataaaggcc	atgtggatat	tttggcacct	actgttcaag	agttggctgc	ccttggaag	2160
gaggcgacga	catctttcct	gcacatctggc	taccttctca	accagctgtt	cagaaccttc	2220
tga						2223

SEQ ID NO: 3 moltype = DNA length = 2463
 FEATURE Location/Qualifiers
 source 1..2463
 mol_type = genomic DNA
 organism = Homo sapiens

SEQUENCE: 3

atggcaggag	agctgggtga	caaaaaggac	cgtgatgcac	caccttccaa	ggaggaaagg	60
aagcgatcac	ggactcctga	cagagagcgg	gatagagacc	gggaccggaa	gtcttcccca	120
tctaagata	gaaagcggca	tcttcaagg	gatagacgtc	gaggaggcag	ccgttctcgc	180
tctcgttccc	gttccaaatc	tgacagaaag	gaacgacggc	acaaagaacg	agaacgagat	240
aaggagcggg	atcggataaa	caaggaccca	gatcgagaca	aggatgggca	cagacgggac	300
aaggaccgta	aacgatccag	cttatctcct	ggtcgaggaa	aagacttta	atctcggga	360
gacagagact	ctaagaagga	tgaagaggat	gaacatgggt	ataagaagcc	taaggcccag	420
ccattatccc	tggaggagct	tctggccaag	aaaaaggctg	aggaagaagc	tgaggctaag	480
cccaagttcc	tctctaaagc	agaacgagag	gctgaagctc	taaagcgacg	gcagcaggag	540
gtggaagagc	ggcagaggat	gcttgaagaa	gagagggaag	aaaggaaaac	gttccaagac	600
ttgggcagga	agatgttgga	agatcctcag	gaacgggaac	gtcgggaacg	caggagagg	660
atggaaacggg	agaccaatgg	aaatgaggat	gaggaaaggc	ggcagaagat	ccgggaagag	720
aaggataaga	gcaaggaact	gcatgccatt	aaggagcgtt	acctgggtgg	catcaaaaag	780
cggcgccgaa	cgagacatct	caatgaccgg	aaatttggtt	ttgagtggga	tgcactgag	840
gacacatcca	ttgactacaa	ccccctgtac	aaagaacggc	accaggtgca	gttgtaggg	900
cgaggcttca	ttgcaggcat	tgacctcaag	cagcagaagc	gagagcagtc	acgtttctat	960
ggagacctaa	tggagaagag	gcgaaccttg	gaagaaaagg	agcaggagga	ggcaagactc	1020
cgcaaaacttc	gtaagaagga	agccaagcag	cgtgggatg	atcgtcattg	gtctcagaaa	1080
aagttagatg	agatgacgga	cagggaactg	cggatcttcc	gtgaggacta	cagcatcacc	1140
accaaagggtg	gcaagatccc	caatcccatc	cgatcctgga	aagactcttc	tctgccccca	1200

-continued

```

cacatcttgg aggtcattga taagtgtggc tacaaggaac caacacctat acagcgtcag 1260
gcaattccca ttgggctaca gaatcgtgac atcattgggtg tggctgagac tggcagtggc 1320
aagacagcag ccttccctcat cctctcgtct gtctggatca ccacacttcc caaaattgac 1380
aggatcgaag agtcagacca aggcccttat gccatcatcc tggctcccac ccgtgagttg 1440
gctcaacaga ttgaggaaga gaccatcaag tttgggaaac cgctaggtat ccgcaactgtg 1500
gctgtcattg gtggcatctc cagagaagac cagggcttca ggctgcgcac gggttgtgag 1560
attgtgattg ctacccctgg gcgtttgatt gatgtgctgg agaaccgcta cctgggtgctg 1620
agccgctgta cctatgtggg tctggatgag gcagatagga tgattgacat gggctttgag 1680
ccagatgtcc agaagatcct ggagcacatg cctgtcagca accagaagcc agacacggat 1740
gaggtcagag accctgagaa gatgctggcc aactttgagt cgggaaaaca taagtaccgc 1800
caaacagtca tgttcacggc caccatgccc ccagcgggtg agcgtctggc caggagctat 1860
cttcggcgac ctgctgtggg gtacattggc tccgcaggca agcccatga gcgtgtggaa 1920
cagaaggtct tccctatgtc agagtcagaa aagaggaaaa agctgctggc aatcttggag 1980
caaggtcttg accaccctat cattattttt gtcaaccaga agaagggtcg cgacgtgtg 2040
gccaatccc tggagaagat ggggtacaat gcttgacac tgcacggtag aaaagccag 2100
gagcagcgag agtttgcgtt gtccaacctc aaggctgggg ccaaggatat tttgtggct 2160
acagatgtgg ctggtcgtgg tattgacatc caagatgtgt ctatgggtgt caactatgat 2220
atggccaaaa atattgaaga ttacatccac cgcattggcc gcacgggacg agcaggcaag 2280
agtggggtgg ccataccctt cctcacaaaa gaggactctg ctgtgttcta cgagctgaag 2340
caagctatcc tggaaagccc agtgtcttcc tgtcccccg aactagccaa ccaccagat 2400
gcccagcata agccaggcac catcctcacc aagaagcgcc gggaagagac catcttggc 2460
tga 2463

```

```

SEQ ID NO: 4      moltype = DNA length = 708
FEATURE          Location/Qualifiers
source            1..708
                  mol_type = genomic DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 4
atggaggagg agacgcatac tgacgccaaa atccgtgctg aaaatggaac aggggtccagc 60
cctcgggggtc ctgggtcgac cctccggcac tttgcctgcg aacagaacct gctgtcgcg 120
ccagatggct ctgcttcctt cctgcaaggt gacacctctg tccctggcggg tgtgtacggg 180
cgggccgagg tgaaggtcag caaagagatt ttcaacaagg ccacactcga agtgatcctg 240
agggcgaaga ttgggtgtgc tgggtgttgc gagaagagcc gggagcggct gatcaggaac 300
acgtgcgagg cgggtggtgt gggcacgttg cacccccga cctccatcac cgtgggtgctg 360
caggttgtca gcgatgcggg ctctctcctg gcctgttgtc tgaatgcgcg ctgcatggca 420
ttggtggtat caggtgtgct catgcgggct ctcttctgtg gggctgcctg cgccctggac 480
tctgatggga cctcgtgctt ggatcctaca tccaagcaag aaaaggaggc ccgggcagtc 540
ctgacctttg ccctggacag cgtggaacgg aagctgtcta tgtccagcac caagggggtc 600
tactcagaca ctgagctcca gcagtgcctg gctgcggccc aggcggcttc gcaacacgtc 660
ttccgtttct accgggaatc gctgcagagg cgttactcca agagctga 708

```

```

SEQ ID NO: 5      moltype = DNA length = 4416
FEATURE          Location/Qualifiers
source            1..4416
                  mol_type = genomic DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 5
atgggtttcca aaagaagact gtcaaaatct gaggataaag agagcctgac agaagatgcc 60
tccaaaacca ggaagcaacc actttccaaa aagacaaaaga aatctcatat tgctaataa 120
gttgaaagaa atgacagcat ctttgtaaag cttcttaaga tatcaggaaat tattcttaaa 180
acgggagaga gtcagaatca actagctgtg gatcaaatag ctttccaaaa gaagctcttt 240
cagaccctga ggagacaccc ttctatccc aaaataatag aagaatttgt tagtgacctg 300
gagctctaca ttgagatga agacagtttc aggaactgcc ttttgtcttg tgagcgtctg 360
caggatgagg aagccagtat ggggtgcattc tattctaaga gtctcatcaa actgcttctg 420
gggattgaca tactgcagcc tgccattatc aaaaccttat ttgagaagtt gccagaatat 480
ttttttgaaa acaagaacag tgatgaaatc aacataacct gactcattgt cagtcaacta 540
aaatggcttg acagagttgt ggatggcaag gacctcacca ccaagatcat gcagctgatc 600
agtattgtct cagagaacct gcagcatgac atcatcacca gcctacctga gatcctaggg 660
gattcccagc acgctgatgt ggggaaagaa ctcagtgacc tactgataga gaatacttca 720
ctcactgtcc caatcctgga tctcctttca agcctccgac ttgaccctaaa cttcctattg 780
aaggttcgcc agttgggtgat ggataagttg tctgtctatta gattggagga ttacacctgtg 840
ataataaagt tcattcttca ttccgtaaca gccatggata cacttgaggt aatttctgag 900
cttcgggaga agttggatct gcagcattgt gttttgccat cacgggtaca ggcttcccaa 960
gtaaagttag aaagttaaagg acgagcaagt tctcaggaa atcaagaaag cagcgtcag 1020
agctgtatta ttctctctt tgatgtaata aagtcagcta ttagatatga gaaaaccatt 1080
tcagaagcct ggattaaggc aattgaaaac actgcctcag tatctgaaca caaggtgttt 1140
gacctgggtg tgcttttcat catctatagc accaatactc agacaaagaa gtacattgac 1200
aggggtgctaa gaaataagat tcgatcaggc tgcattcaag aacagctgct ccagagtaca 1260
ttctctgttc attacttagt ccttaaggat atgtgttcat ccattctgtc gctggctcag 1320
agtttgcctc actctctaga ccagagtata atttcatttg gcagctctct atacaatat 1380
gcatttaagt tttttgacac gtactgccag caggaagtgg ttggtgcctt agtgacctat 1440
atctgcagtg ggaatgaagc tgaagttagt actgccttag atgtccttct agagttggta 1500
gtgttaaac catctgctat gatgatgaat gctgtctttg taaagggcat tttagattat 1560
ctggataaca tatccctca gcaaatacga aaactcttct atgttctcag cacactggca 1620
tttagcaaac agaatgaagc cagcagccac atccaggatg acatgcactt ggtgataaga 1680

```

-continued

```

aagcagctct ctagaccgtt attcaagtac aagctcattg ggattattgg tgetgtgacc 1740
atggctggca tcatggcggc agacagaagt gaatcaccta gtttgaccca agagagagcc 1800
aacctgagcg atgagcagtg cacacaggtg acctccttgt tgcagtttgt tcattcctgc 1860
agtggagcagt ctcctcaggg cctcgcactt tactatgatg aatttgccaa cctgatccaa 1920
catgaaaaagc tggatccaaa agccctggaa tgggttgggc ataccatctg taatgatttc 1980
caggatgcct tcgtagtgga ctcctgtgtt gttccggaag gtgactttcc atttccctgtg 2040
aaagcactgt acggactgga agaatacgac actcaggatg ggattgccat aaacctctcg 2100
ccgtctgtgt tttctcagga ctttgcaaaa gatgggggtc cggtgacctc acaggaatca 2160
ggccaaaaat tgggtgtctc gctgtgcctg gctccgtatt tccggttact gagactttgt 2220
gtggagagac agcataacgg aaacttgagg gagattgatg gtctactaga ttgtcctata 2280
ttcctaactg acctggagcc tggagagaag ttggagtcca tgtctgctaa agagcgttca 2340
ttcatgtgtt ctctcatatt tcttactctc aactgggtcc gagagattgt aaatgccttc 2400
tgccagggaaa catcacctga gatgaagggg aagggtgtca ctcggttaaa gcacattgta 2460
gaattgcaaaa taactcctga aaagtacttg gcagtcaccc cagactatgt cctcctctt 2520
ggaaactttg atgtggaaac tttagatata acacctcata ctgttactgc tatttcagca 2580
aaaatcgaaa agaaaggaaa aatagaaaag aaacaaaaaa cagatggcag caagacatcc 2640
tcctctgaca cactttcaga agagaaaaat tcagaatgtg acctacgcc atctcataga 2700
ggccagctaa acaaggagtt cacagggaag gaagaaaaga catcattgtt actacataat 2760
tccatgctt tttccgaga gctggacatt gaggtcttct ctattctaca ttgtggactt 2820
gtgacgaagt tcactctaga tactgaaatg cacactgaag ctacagaagt tgtgcaactt 2880
ggggccctcg agctgctttt cttgctggaa gatctctccc agaagctgga gagtatgctg 2940
acacctccta ttgcccagg agtccccttt ctcaagaaca aaggaagccg gaatatgga 3000
ttctcacatc tccaacagag atctgcccga gaaattgttc attgtgtttt tcaactgctg 3060
acccaatgt gtaaccacct ggagaacatt cacaactatt ttcagtgttt agctgctgag 3120
aatcacggtg tagttgatgg accaggagtg aaagttcagg agtaccacat aatgtcttcc 3180
tgctatcaga ggctgctgca gatttttcat gggctttttg cttggagtgg attttctcaa 3240
cctgaaaatc agaatttact gtattcagcc ctccatgtcc ttagtagccg actgaaacag 3300
ggagaacaca gccagccttt ggaggaaacta ctacgccaga gcgtccatta cttgcagaat 3360
ttccatcaaa gcattcccag tttccagtgt gctctttatc tcactcagact tttgatgggt 3420
attttgagga aatcaacagc ttctgctcag aacaagaaaa aaattgtctc ccttgccaga 3480
caattcctct gtcgggtgtg gccaaagtgg gataaagaga agagcaacat ctctaagac 3540
cagctccatg cctcgtctct tatctacctg gagcacacag agagcattct gaaggccata 3600
gaggagattg ctggtgttgg tgtcccagaa ctgatcaact ctctaaaga tgcattctcc 3660
tccacattcc ctacactgac caggcatact tttgtgtttt tcttcctgtg gatgatggct 3720
gaactagaga agacgggtgaa aaaaattgag cctggcacag cagcagactc gcagcagatt 3780
catgaagaga aactcctcta ctggaacatg gctgttcgag acttcagtat cctcatcaac 3840
ttgataaagg tatttgatag tcatcctgtt ctgcatgtat gtttgaagta tggcgctctc 3900
tttgtggaag catttctgaa gcaatgtatg ccgctcctag acttcagttt tagaaaaaac 3960
cgggaagatg ttctgagcct actggaacc ttccagttgg acacaaggct gcttcacac 4020
ctgtgtgggc attccaagat tcaccaggac acgagactca cccaacatgt gcctctgctc 4080
aaaaagacc ttggaactttt agtttcgaga gtcaaaagta tgcctactct caacaattgt 4140
agagaggctt tctggctggg caatctaaaa aaccgggact tgcagggtga agagattaa 4200
tccaaaaatt cccaggagag cagacagat gagagtgagg atgacatgtc atcccaggcc 4260
tccaagagca aagccactga ggtatctcta caaaaccac cagagtctgg cactgatgg 4320
tgcatttgt taattgttct aagttgggtg agcagaactt tgcctactta tgtttattgt 4380
caaatgcttc tatgccatt tccattccct ccataa 4416

```

SEQ ID NO: 6 moltype = DNA length = 2478

FEATURE Location/Qualifiers

source 1..2478

mol_type = genomic DNA

organism = Homo sapiens

SEQUENCE: 6

```

atgagttcct cgctgtttaa tgtaaaaaag ctgaagggtg cggagctgaa agaggagctc 60
aagaagcgac gcctttctga caagggtctc aaggccgagc tcatggagcg actccaggct 120
gcgctggacg cagaggaggg cgggggcccgc cccgccatgg agcccgggaa cggcagccta 180
gacctgggcg gggattccgc tgggcgctcg ggagcaggcc tcgagcagga gcccgcggcc 240
ggcgccgatg aagaggagga ggaagaggag aaggaatctc cgctctggac 300
ggcgaccaga tggagctagg agaggagaa cggggccggc gggcgggcga ctggggcccg 360
atggaggagg aggaggccgc ctcggaagac gagaacggcg acgatcaggg ttccaggaa 420
ggggaagatg agctcgggga cgaagaggaa ggcgcgggcg acgagaacgg gcacggggag 480
cagcagcctc aaccgccggc gacgcagcag caacagcccc aacagcagcg cggggccggc 540
aaggaggccg cggggaagag cagcggcccc acctcgctgt tcgcggtgac ggtggcgccg 600
cccggggcga ggcagggccg gcagcaggcg ggaggtaaga agaaggcgga aggcggcgga 660
ggcgcggtc gcccgggggt tccggcgggc ggggacggca aaacagaaca gaaaggcgga 720
gataaaaaga ggggtgttaa aagaccacga gaagatcatg gccgtggata ttttgagtac 780
attgaagaga acaagtatag cagagccaaa tctcctcagc cacctgttga agaagaagat 840
gaacacttgc atgacacagt ggtttgtctt gatacttata attgtgatct acattttaaa 900
atatcaagag atcgtctcag tgcttcttcc cttacaatgg agagttttgc ttttctttgg 960
gctggaggaa gagcatccta ttgtgtgtca aaaggcaaa gtgtttttga gatgaaggtt 1020
acagagaaga tccagttatg gcatttatat acaaaagata ttgacatata tgaagttcgt 1080
attggctggt cactaactac aagtggaatg ttacttgggt aagaagaatt ttcttatggg 1140
tattctctaa aaggaaataa aacatgcaac tgtgagactg aagattatgg aaaaaagttt 1200
gatgaaaatg atgtgattac atgttttgc taaactttgaa gtgatgaagt agaactctcg 1260
tatgctaaga atggacaaga tcttggcgtt gccttcaaaa tcagtaagga agttcttgct 1320
ggacggccac tgttcccgcg tgttctctgc cacaactgtg cagttgaatt taatttttgt 1380

```

-continued

```

cagaaggaaa agccatattt tccaatacct gaagagtata ctttcatcca gaacgtcccc 1440
ttagaggatc gagttagagg accaaagggg cctgaagaga agaaagattg tgaagttgtg 1500
atgatgattg gcttgccagg agctggaaaa actacctggg ttactaaaca tgcagcagaa 1560
aatccaggga aataatacat tcttggcaca aatactatta tggataagat gatggtggca 1620
ggttttaaga agcaaatggc agatactgga aaactgaaca cactgttgca gagagcccc 1680
cagtgtcttg ggaaatttat tgagattgct gcccgaaaaga agcgaaattt tattctggat 1740
cagacaaatg tgtctgctgc tgcccagagg agaaaaatgt gcctgtttgc aggtctccag 1800
cgaaaagctg ttgtagtttg cccaaaagat gaagactata agcaaaagaa acagaagaaa 1860
gcagaagttag aggggaaga cctaccagaa catgctggctc tcaaaatgaa aggaaacttt 1920
acccctccag aggttagctga gtgctttgat gaaataacct atgttgaaat tcagaaggaa 1980
gaagcccaaa aactcttgga gcaatataag gaagaaagca aaaaggctct tccaccagaa 2040
aagaacacaga acactggctc aaagaaaagc aataaaaata agagtggcaa gaaccagttt 2100
aacagagggtg gtggccatag aggacgtgga ggattcaata tgcgtgggtg aaatttcaga 2160
ggaggagccc ctgggaatcg tggcgatat ataggagg gcaacatgcc acagagaggt 2220
ggtggcggtg gaggaagtgg tggaaatcgcc tatccatacc ctctgcccc tgttttctct 2280
ggcgtggtga gttactcaaa agaggggaac tacaaacagag gtggaatgcc caacagaggg 2340
aactacaacc agaactctag aggacgagga aacaatcgtg gctacaaaaa tcaatctcag 2400
ggctacaacc agtggcagca ggttcaattc tggggtcaga agccatggag tcagcattat 2460
caccaaggat attattga                               2478

```

```

SEQ ID NO: 7      moltype = DNA length = 1914
FEATURE
source            Location/Qualifiers
                  1..1914
                  mol_type = genomic DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 7
atggcgccga tggcggtcgg ggggtgctggt gggagcccgcg tgtccagcgg gagggacctg 60
aattgcgtcc ccgaaatagc tgacacacta ggggctgtgg ccaagcaggg gtttgatttc 120
ctctgcattc ctgtcttcca tccgcgtttc aagagggagt tcattcagga acctgctaag 180
aatccggccc gtccccagac acgatcagac ctactgctgt caggaaaggga ctggaatacg 240
ctaattgtgg gaaagctttc tccatggatt cgtccagact caaaagtgga gaagattcgc 300
aggaactccg agcgggccat gttacaggag ctgaattttg tggcatattt ggttcttcca 360
gctttcctgc tgccccttaa tcaggaagat aacaccaacc tggccagagt ttgaccaac 420
cacatccaca ctggccatca ctcttccatg ttctggatgc gggtaacctt ggtggcacca 480
gaggacctga gagatgatat aattgagaat gcaccaacta cacacacaga ggagatagat 540
ggggaggaga aaacgtggat gtggtggcac aacttccgga ctttgtgtga ctatagtaag 600
aggattgcag tggctcttga aattggggct gacctcccat ctaatcatgt cattgatcgc 660
tggcttgggg agcccatcaa agcagccatt ctccccacta gcattttcct gaccaataag 720
aagggttttc ctgttcttcc taagatgcac cagagggtca tcttccggct cctcaagttg 780
gaggtgcagt tcatcatcac aggcaccaac caccactcag agaaggagtt ctgctcctac 840
ctccaatacc tggaaatactt aagccagaac cgtcctccac ctaatgccta tgaactcttt 900
gccaagggtc atgaagacta tctgcagtc ccgcttcagc cactgatgga caatctggaa 960
tctcagacat ctgaagtgtt tgaaaaggag cccatcaaat actctcagta ccagcaggcc 1020
atctataaat gtctgctaga ccgagtacca gaagaggaga aggataccaa tgtccaggta 1080
ctgatggtgc tgggagcagg acggggaccc ctgggtgaacg cttccctgcg ggcagccaa 1140
caggccgacc ggcggtataa gctgtatgct gtggagaaaa acccaaatgc cgtggtgacg 1200
ctagagaact ggcagtttga agaattggga agccaagtga ccgtagtctc atcagacatg 1260
agggaaatggg tggctccaga gaaagcagac atcattgtca gtgagctctt gggctcattt 1320
gctgacaatg aattgtcgcc tgagtgcctg gatggagccc agcacttctt aaaagatgat 1380
ggtgtgagca tcccggggga gtacacttcc tttctggctc ccatctcttc ctccaagctg 1440
tacaatgagg tccagacctg tagggagaag gaccgtgacc ctgagggcca gtttgagatg 1500
ccttatgtgg taccgtgcat caacttccac cagctctctg caccacagcc ctgtttcacc 1560
ttcagccatc ccaacagaga tctatgatt gacaacaacc gctattgcac cttggaattt 1620
cctgtggagg tgaacacagt actacatggc tttgcggctc actttgagac tgtgctttat 1680
caggacatca cctctgagat ccgtccagag actcactctc ctgggatggt ctcagtgttt 1740
cccatctctt tccctattaa gcagcccata acggtacgtg aaggccaaac catctgtgtg 1800
cgtttctggc gatgcagcaa ttccaagaag gtgtgggtatg agtgggctgt gacagcacca 1860
gtctgttctg ctattcataa cccacaggcc cgctcatata ccattggcct ctag 1914

```

```

SEQ ID NO: 8      moltype = DNA length = 2100
FEATURE
source            Location/Qualifiers
                  1..2100
                  mol_type = genomic DNA
                  organism = Homo sapiens

```

```

SEQUENCE: 8
atgagctccc ggaaagtgtc ggccattcag gcccgaaaag ggaggccgaa aagagagaaa 60
catccgaaaa agccggagcc tcaacagaaa gctcctttag ttccctctcc tccaccgcca 120
ccaccaccac caccgccacc ttggccagac cccacacccc cggagccaga ggaggagatc 180
ctgggatcag atgatgagga gcaagaggac cctgcccact actgcaaaag tggatatcat 240
ccagtgaaaa ttggagacct ctccaatggc cggtatcatg ttattagaaa gcttggatgg 300
gggcacttct ctactgtctg gctgtgctgg gatatgcagg ggaaaagatt tgttgcaatg 360
aaagtgttaa aaagtgcaca gcaattatag gagacagcct tggatgaaat aaaattgtct 420
aaatgtgttc gagaaagtga tcccagtgac ccaacaaaag acatgggtgtt ccagctcatt 480
gacgacttca agatttcagg catgaatggg atacatgtct gcatggtctt cgaagtactt 540
ggccaccatc tccccaagtg gatcatcaaa tccaactatc aaggcctccc agtacgttgt 600
gtgaagagta tcattcgaca ggtccttcaa gggttagatt acttacacag taagtgaag 660

```

-continued

```

atcattcata ctgacataaa gccggaaaat atcttgatgt gtgtggatga tgcataatgt 720
agaagaatgg cagctgaggg cactgagtggt cagaaagcag gtgctcctcc tccttcaggg 780
tctgcagtga gtacgggtcc acagcagaaa cctataggaa aaatatctaa aaacaaaaag 840
aaaaaactga aaaagaaaca gaagaggcag gctgagttat tggagaagcg cctgcaggag 900
atagaagaat tggagcgaga agctgaaaag aaaataatag aagaaaacat cacctcagct 960
gcaccttcca atgaccagga tggcgaatac tgcccagagg tgaaactaaa aacaacagga 1020
ttagaggagg cggtcgaggg agagactgca aaggacaatg gtgaagctga ggaccaggaa 1080
gagaaagaag atgctgagaa agaaaacatt gaaaaagatg aagatgatgt agatcaggaa 1140
cttgcgaaaca tagacctac gtgtagataa tcacctaaaa ccaatggcca tattgagaat 1200
ggcccatctc cactggagca gcaactggac gatgaagatg atgatgaaga agactgcca 1260
aatcctgagg aatataatct tgatgagcca aatgcagaaa gtgattacac atatagcagc 1320
tcctatgaac aattcaatgg tgaattgcca aatggacgac ataaaattcc cgagtccagc 1380
ttcccagagt tttccacctc gttgttctct ggatccttag aacctgtggc ctgaggctct 1440
gtgctttctg acttactgag caagaggaga gcagtcctac ccatgacaga 1500
agcagaacgg tttcagctcc cagtactggg gatttgccaa aagcaaaaac ccgggcagct 1560
gacttggtgg tgaatccctt ggatccgagg aatgcagata aaattagagt aaaaattgct 1620
gacctgggaa atgcttgggt ggtgcataaa cacttcacgg aagacatcca gacgcgtcag 1680
taccgctcca tagagggttt aataggagcg ggttacagca cccctgcgga catctggagc 1740
acggcgtgta tggcctgaca gctggcaacg ggagattatt tgttgaacc acattctggg 1800
gaagactatt ccagagagca agaccacata gccacatca tagagctgct aggcagtatt 1860
ccaaggcact ttgctctatc tggaaaatat tctcgggaat tcttcaatcg cagaggagaa 1920
ctgcgacaca tccaacagct gaagccctgg agcctctttg atgtacttgt ggaaaagtat 1980
ggctggcccc atgaagatgc tgcacagttt acagatttcc tgatcccgat gttagaaatg 2040
gttcagaaaa aacgagcctc agctggcgaa tgccttcggc atccttggtt gaattcttag 2100

```

```

SEQ ID NO: 9      moltype = DNA length = 3087
FEATURE          Location/Qualifiers
source           1..3087
                 mol_type = genomic DNA
                 organism = Homo sapiens

```

```

SEQUENCE: 9
atggggtccg ccgcctgccc ccggggagcc ttgcccggagc tcgcgcctcg ctgccagcct 60
cgcgagcagt cgcagcccca caccgatgg gacgcgggct gtgggattca gcaccccggg 120
ggcgaggaat tcaggacctc ccgcggggca agggcctata gggttccgaa ctgcaggag 180
ggtcgctcct cccctactcg ctttttccc gcaccggaag gcccgcgcca ctgctttgtt 240
tcctctccag accgcgcatt ttgggtctcg gaagagggtc agaggctgtt gttgagcaat 300
gcatgccagc caaaagaatg caatgggtga aagattccag ttgatgccag taaacctaat 360
ccaaatgatg tggagtgtga taactctgat ttggatatga atggaatcat ccatccctgt 420
actcatctcg aagacaaacc agcacaaaaa aatgaagatg aaatgatggt tgcaattttt 480
gagtacattg acagactttt cagtattgta agaccaagaa gacttctcta catggcaata 540
gatggagtgg caccacgtgc taaaatgaac cagcagcgtt caaggagggt caggggcatca 600
aaagaaggaa tggagcagc agtcgagaag cagcgagatc ggggaagaaa attggcaaaa 660
ggtggctttc ttctctccaga agaaataaaa gaaagatttg acagcaactg tattacacca 720
ggaaactgaat tcctggacaa tcttgctaaa tgccttcgct attacatagc tgatcgttta 780
aataatgacc ctgggtggaa aaatttgaca gttattttat ctgatgctag tgctcctggt 840
gaaggagaac ataaaatcat ggattacatt agaaggcaaa gagccagcc taacctgac 900
ccaaatactc atcattgttt attgtggagc gatgetgatc tcattatgct tggccttgcc 960
acacatgaac cgaactttac cattattaga gaagaattca aacaaacaa gcccaaacca 1020
tgttggtctt gtaatcagtt tggacatgag gtcaaaagatt gtgaagggtt gccaaagaaa 1080
aagaagggaa agcatgatga acttgccgat agtcttcctt gtgcagaagg agagtttatc 1140
tcccttcggc ttaatgttct tcgtgagtat ttggaaaag aactcacaat gccagcccta 1200
ccattccatc ttgatgttga gaggagcatt gatgaactgg ttttcatggt cttctttgtg 1260
ggaaatgact tcctccctca ttggccatcg ttagagatta gggaaaaatgc aattgaccgt 1320
ttggttaaca tatacaaaaa tgtggtacac aaaactgggg gttaccttac agaaagtggg 1380
tatgtcaatc tgcaaaagat acagatgatc atgttagcag ttggtgaagt tgaggatagc 1440
atttttaaaa agagaaagga tgatgaggac agttttagaa gacgacagaa agaaaaaaga 1500
aagagaatga agagagatca accagctttc actcctagtg gaatattaac tcctcatgcc 1560
ttgggttcaa gaaattccac aggtttctca gttagccagta atccgagaca agcagcctat 1620
gaaatgagga tgcagaataa ctctagtcct tcgatatctc ctaatacagag ttccacatct 1680
gatggctccc cgtctccatt aggaggaatt aagcgaaaaa cagaagacag tgacagtga 1740
cctgagccag aggataatgt cagggtatgg gaagctggct ggaagcagcg gtactacaag 1800
aacaatttgg atgtggatgc agctgatgag aaattccgtc ggaaagtgtt gcagtcgtac 1860
gttgaaagac tttgctgggt tcttagatat tattaccagg gctgtgcttc ctggaagtgg 1920
tattatccat ttcattatgc accatttgct tcagactttg aaggcattgc agacatgcca 1980
tctgattttg agaagggtac gaaccggtt aaaccactag aacaacttat gggggtattt 2040
ccagctgcaa gtggttaatt tctacctcca tcatggcgga agctcatgag tgatcctgat 2100
tctagtataa ttgacttcta tcctgaagat tttgctattg atttgaatgg gaagaaatat 2160
gcatggcaag gtgttgctct ctggccattc gtggatgagc gaaggctacg agctgcccta 2220
gaagagggat acccagacct cactccagaa gagaccagaa gaaacagcct tggaggtgat 2280
gtcttatttg tggggaaaca tcaccactc catgacttca ttttagagct gtaccagaca 2340
ggttccacag agccagtggg ggtaccccc gaactatgtc atgggattca aggaaagtgt 2400
tctttggatg aagaagccat tcttccagat caaatagtat gttctcctgt tcctatgtta 2460
agggatctga cacagaacac tgtagtcaat attaatttta aagaccaca gtttctgtaa 2520
gattacattt ttaaagctgt aatgcttcca ggagcaagaa agccagcagc agtactgaaa 2580
cctagtgact tggaaaatc cagcaatgga cggcagtgga agcctcagct tggctttaac 2640
cgtgaccgga ggctgtgca cctggatcag gcagccttca ggactttggg ccatgtgatg 2700

```

-continued

ccaagaggct	caggaactgg	catttacagc	aatgctgcac	caccacctgt	gaattaccag	2760
ggaaacttat	acaggccgct	tttgagagga	caagcccaga	ttccaaaact	tatgtcaa	2820
atgaggcccc	aggatttcctg	gcgaggctcct	cctccccttt	tcacagcaga	aaggtttgac	2880
agaggcggtg	gggctgaacc	tctgctccca	tggaaccgga	tgctgcaa	ccagaatgca	2940
gccttccagc	caaaccagta	ccagatgcta	gctgggcctg	gtgggtatcc	acccagacga	3000
gatgatcgtg	gagggagaca	gggatatccc	agagaaggaa	ggaaataccc	tttgccacca	3060
ccctcaggaa	gatacaattg	gaattaa				3087

1. A composition comprising at least one G-quadruplex (G4) stabilizer for its use in a method for treating an individual afflicted by a multiple myeloma.

2. The composition according to claim 1, wherein the at least one G4 stabilizer is selected in a group consisting of Quarfloxin, Pidnarulex, MM41, Telomestatin, BMSG-SH-3, BRACO-19, CMO3, PDP, Pyridostatin, carboxyPyridostatin, PhenDC3, AQ1, TMPyP4, RHPS4, 360A, FG, 20A, Emetine and combinations thereof, preferably the G4 stabilizer is Pyridostatin.

3. The composition according to claim 1, wherein the composition further comprises at least one histone deacetylase inhibitor.

4. The composition according to claim 3, wherein the at least one histone deacetylase inhibitor is selected in a group consisting of Panobinostat, trichostatin-A, vorinostat, belinostat, NVP-LAQ824, givinostat, romidepsin, valproic acid, sodium phenylbutyrate, MS-275, MGCD0103 and combinations thereof, preferably the histone deacetylase inhibitor is Panobinostat.

5. The composition according to claim 1, wherein the composition further comprises at least one bromodomain and extraterminal (BET) proteins inhibitor.

6. The composition according to claim 5, wherein the at least one BET proteins inhibitor is selected in the group consisting of RVX-208, I-BET-762, OTX015, CPI-0610, GSK525762, ABBV-075, FT-1101, INCB057643, ZEN003694, GSK2820151, CC-90010, PLX51107, ABBV-744, BAY1238097, B1894999, BMS-986158, GS-5829, INCB054328, RO6870810 and combinations thereof.

7. The composition according to claim 1, wherein the composition further comprises at least one nitrogen mustard.

8. The composition according to claim 7, wherein the nitrogen mustard is selected in the group consisting of Chlormethine, Chlorambucil, Melphalan, Cyclophosph-

amide, Ifosfamide, Estramustine, Prednimustine, Bendamustine, Melphalan flufenamide and combinations thereof.

9. The composition according to claim 1, wherein the individual has been prognosed with a poor outcome.

10. The composition according to claim 9, wherein the poor outcome of the individual is in vitro determined by carrying out the following steps:

- measuring in a biological sample from said individual, the expression level of 9 genes consisting of the nucleic acid sequences SEQ ID NO:1 to SEQ ID NO:9;
- calculating a score TRC_{score} according to the following formula

$$TRC_{score} = \sum_{i=1}^9 \beta_i \times C_i$$

wherein β_i represents the regression β coefficient reference value for the gene of nucleic acid sequence SEQ ID NO:i,

wherein $C_i=1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO:i is higher than an expression level of reference $C_i=-1$ if the expression level of the gene of nucleic acid sequence SEQ ID NO:i is lower than or equal to ELRi,

- prognosing that said individual with a score TRC_{score} higher than a reference value TRC_{ref} is likely to have a bad outcome, or said individual with a score TRC_{score} lower than a reference value TRC_{ref} is likely to have a good outcome,

wherein the reference value TRC_{ref} is of -0.39535 .

* * * * *