



US 20250283872A1

(19) **United States**

(12) **Patent Application Publication**
Bredt et al.

(10) **Pub. No.: US 2025/0283872 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **EXPRESSION SYSTEMS FOR THE
ALPHA2ALPHA5BETA2 NICOTINIC
ACETYLCHOLINE RECEPTOR AND
METHODS OF USE THEREOF**

(71) Applicant: **Janssen Pharmaceutica NV, Beerse
(BE)**

(72) Inventors: **David S Bredt, La Jolla, CA (US);
Shenyan Gu, San Diego, CA (US);
Min Lei O'Carroll, San Diego, CA
(US)**

(21) Appl. No.: **18/288,024**

(22) PCT Filed: **Apr. 22, 2022**

(86) PCT No.: **PCT/EP2022/060744**

§ 371 (c)(1),

(2) Date: **Oct. 23, 2023**

Related U.S. Application Data

(60) Provisional application No. 63/178,813, filed on Apr.
23, 2021.

Publication Classification

(51) **Int. Cl.**

G01N 33/50 (2006.01)

C07K 14/705 (2006.01)

C12N 9/10 (2006.01)

G01N 33/94 (2006.01)

(52) **U.S. Cl.**

CPC ... G01N 33/5041 (2013.01); **C07K 14/70571**

(2013.01); **C12N 9/1029** (2013.01); **G01N**

33/944 (2013.01); **C12Y 203/01006** (2013.01);


C12Y 203/01057 (2013.01)

(57)

ABSTRACT

Disclosed herein are isolated recombinant cells for the
expression of $\alpha 2 \alpha 5 \beta 2$ nicotinic acetylcholine receptor
(nAChR) and methods of use thereof.

Specification includes a Sequence Listing.






Figure 1




Figure 2

EXPRESSION SYSTEMS FOR THE ALPHA2ALPHA5BETA2 NICOTINIC ACETYLCHOLINE RECEPTOR AND METHODS OF USE THEREOF

CROSS REFERENCE TO RELATED APPLICATION

[0001] This application claims the benefit of U.S. Provisional Application 63/178,813, filed on Apr. 23, 2021, which is incorporated by reference herein in its entirety.

FIELD OF INVENTION

[0002] This invention relates to isolated recombinant cells for the expression of $\alpha 2\alpha 5\beta 2$ nicotinic acetylcholine receptor (nAChR) and methods of use thereof.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[0003] The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 28, 2022, is named PRD4128WOPCT1_SL.txt and is 24,576 bytes in size.

BACKGROUND OF INVENTION

[0004] Nicotinic acetylcholine receptors (nAChRs) are implicated in several neurological and psychiatric conditions including nicotine addiction, pain, and psychotic disorders (Gotti et al., Neuronal nicotinic receptors: from structure to pathology, Prog Neurobiol. 2004 December; 74 (6): 363-96). Nicotine addiction associated with smoking is the major cause of lung cancer. Genome-wide association studies show that a nucleotide polymorphism (SNP) rs16969968, which encodes a change (D398N) in sequence of the $\alpha 5$ subunit of the nAChR increases susceptibility to nicotine addiction. The habenulo-interpeduncular pathway which is specifically enriched in the $\alpha 5$ nAChR is a key neurocircuit controlling nicotine intake. For these reasons, the $\alpha 5$ nAChR has been proposed as a valuable drug target (Fowler et al., Habenular nicotinic receptors containing the $\alpha 5^*$ subunit controls nicotine intake, Nature. 2011 Mar. 31; 471 (7340): 597-601; Maskos, The nicotinic receptor alpha5 coding polymorphism rs16969968 as a major target in disease: Functional dissection and remaining challenges, Journal of Neurochemistry. 2020:154-241-250). However, identification of compounds that modulate $\alpha 5$ -containing nAChRs (such as $\alpha 2\alpha 5\beta 2$ nAChR) has not been possible, as this receptor is not functionally expressed in recombinant cell lines typically used for drug discovery.

[0005] There still is a need to develop a cell line that expresses $\alpha 5$ subunit of nAChR robustly and can be used for drug discovery.

BRIEF SUMMARY OF THE INVENTION

[0006] Provided herein are isolated recombinant cells comprising: a) a heterologous nucleic acid encoding an $\alpha 5$ subunit of nicotinic acetylcholine receptor (nAChR) and b) a heterologous nucleic acid encoding a chaperone protein selected from transmembrane inner ear expressed protein TMIE and protein FAM163B.

[0007] In one embodiment of the isolated recombinant cells provided above, the cells further comprise: c) a hetero-

ologous nucleic acid encoding an $\alpha 2$ subunit of nAChR; d) a heterologous nucleic acid encoding a $\beta 2$ subunit of nAChR; e) a heterologous nucleic acid encoding diamine acetyltransferase 1 (SAT1); and f) a heterologous nucleic acid encoding choline O-acetyltransferase (CHAT), wherein, the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of nAChR forms a $\alpha 2\alpha 5\beta 2$ nAChR.

[0008] In a further embodiment of the isolated recombinant cells, the recombinant cells are mammalian cells.

[0009] In a yet further embodiment of the isolated recombinant cells, the mammalian cells are selected from the group consisting of a human embryonic kidney 293T (HEK293T) cell, a HEK293F cell, a HeLa cell, a Chinese hamster ovary (CHO) cell, a NIH 3T3 cell, a MCF-7 cell, a Hep G2 cell, a baby hamster kidney (BHK) cell, and a Cos7 cell.

[0010] In a yet further embodiment of the isolated recombinant cells, the $\alpha 2$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 1.

[0011] In a yet further embodiment of the isolated recombinant cells, the $\alpha 5$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 2.

[0012] In a yet further embodiment of the isolated recombinant cells, the $\beta 2$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 3.

[0013] In a yet further embodiment of the isolated recombinant cells, the TMIE comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 4.

[0014] In a yet further embodiment of the isolated recombinant cells, the FAM163B comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 5.

[0015] In a yet further embodiment of the isolated recombinant cells, the SAT1 comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 6.

[0016] In a yet further embodiment of the isolated recombinant cells, the CHAT comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 7.

[0017] Further provided herein are methods for identifying agonists, antagonists, or positive allosteric modulators of $\alpha 5$ containing nAChR, the method comprising: a) contacting the isolated recombinant cell of any one of claims 1-11 with an agent and b) determining the activity of the $\alpha 5$ containing nAChR of the isolated recombinant cell, wherein the agent is identified as an agonist or positive allosteric modulator (PAM) if the agent enhances the activity of the $\alpha 5$ containing nAChR and the agent is identified as an antagonist if the agent decreases the activity of the $\alpha 5$ containing nAChR as compared to the activity of the $\alpha 5$ containing nAChR when the isolated recombinant cell was not contacted with the agent.

[0018] Yet further provided herein are methods for identifying agonists, antagonists, or positive allosteric modulators of $\alpha 2\alpha 5\beta 2$ nAChR, the method comprising: a) contacting the isolated recombinant cell of any one of claims 2-11 with an agent and b) determining the activity of the $\alpha 2\alpha 5\beta 2$ nAChR of the isolated recombinant cell, wherein the agent is identified as an agonist or positive allosteric modulator

(PAM) if the agent enhances the activity of the $\alpha 2\alpha 5\beta 2$ nAChR and the agent is identified as an antagonist if the agent decreases the activity of the $\alpha 2\alpha 5\beta 2$ nAChR as compared to the activity of the $\alpha 2\alpha 5\beta 2$ nAChR when the isolated recombinant cell was not contacted with the agent.

[0019] In one embodiment of the method, step b) comprises determining calcium flux of the isolated recombinant cell, wherein the agent is identified as an agonist if the agent enhances the calcium flux as compared to the calcium flux when the isolated recombinant cell was not contacted with the agent.

[0020] In a further embodiment of the method, step b) comprises determining calcium flux and nicotine-evoked calcium flux of the isolated recombinant cell, wherein the agent is identified as a PAM if the agent does not enhance calcium flux and enhances the nicotine-evoked calcium flux as compared to the calcium flux and nicotine-evoked calcium flux when the isolated recombinant cell was not contacted with the agent

[0021] In a yet further embodiment of the method, step b) comprises determining nicotine-evoked calcium flux of the isolated recombinant cell, wherein the agent is identified as an antagonist if the agent decreases the nicotine-evoked calcium flux as compared to the nicotine-evoked calcium flux when the isolated recombinant cell was not contacted with the agent.

[0022] In a yet further embodiment of the method, the isolated recombinant cell is incubated at about 25° C.-35° C. for about 20-50 hours prior to being contacted with the agent.

[0023] In a yet further embodiment of the method, the agent is a small molecule or peptide.

[0024] Yet further provided herein are kits comprising (i) the isolated recombinant cells as provided above and (ii) instructions for use.

BRIEF DESCRIPTION OF THE DRAWINGS

[0025] FIG. 1 is a graph showing functional expression of $\alpha 2\beta 2$ nAChR and $\alpha 2\alpha 5\beta 2$ nAChR when co-expressed with different chaperone proteins.

[0026] FIG. 2 is concentration response curves for nicotine-evoked activation of $\alpha 2\beta 2$ nAChR and $\alpha 2\alpha 5\beta 2$ nAChR when co-expressed with different chaperone proteins.

DETAILED DESCRIPTION OF THE INVENTION

[0027] Various publications, articles and patents are cited or described in the background and throughout the specification; each of these references is herein incorporated by reference in its entirety. Discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is for the purpose of providing context for the invention. Such discussion is not an admission that any or all of these matters form part of the prior art with respect to any inventions disclosed or claimed.

[0028] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood to one of ordinary skill in the art to which this invention pertains. Otherwise, certain terms used herein have the meanings as set forth in the specification.

[0029] It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural reference unless the context clearly dictates otherwise.

[0030] Unless otherwise stated, any numerical values, such as a concentration or a concentration range described herein, are to be understood as being modified in all instances by the term “about.” Thus, a numerical value typically includes $\pm 10\%$ of the recited value. For example, a concentration of 1 mg/ml includes 0.9 mg/ml to 1.1 mg/ml. Likewise, a concentration range of 1% to 10% (w/v) includes 0.9% (w/v) to 11% (w/v). As used herein, the use of a numerical range expressly includes all possible sub-ranges, all individual numerical values within that range, including integers within such ranges and fractions of the values unless the context clearly indicates otherwise.

[0031] Unless otherwise indicated, the term “at least” preceding a series of elements is to be understood to refer to every element in the series. Those skilled in the art will recognize or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the invention.

[0032] As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains” or “containing,” or any other variation thereof, will be understood to imply the inclusion of a stated integer or group of integers but not the exclusion of any other integer or group of integers and are intended to be non-exclusive or open-ended. For example, a composition, a mixture, a process, a method, an article, or an apparatus that comprises a list of elements is not necessarily limited to only those elements but can include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, “or” refers to an inclusive or and not to an exclusive or. For example, a condition A or B is satisfied by any one of the following: A is true (or present) and B is false (or not present), A is false (or not present) and B is true (or present), and both A and B are true (or present).

[0033] As used herein, the conjunctive term “and/or” between multiple recited elements is understood as encompassing both individual and combined options. For instance, where two elements are conjoined by “and/or,” a first option refers to the applicability of the first element without the second. A second option refers to the applicability of the second element without the first. A third option refers to the applicability of the first and second elements together. Any one of these options is understood to fall within the meaning, and therefore satisfy the requirement of the term “and/or” as used herein. Concurrent applicability of more than one of the options is also understood to fall within the meaning, and therefore satisfy the requirement of the term “and/or.”

[0034] As used herein, the term “consists of,” or variations such as “consist of” or “consisting of,” as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, but that no additional integer or group of integers can be added to the specified method, structure, or composition.

[0035] As used herein, the term “consists essentially of,” or variations such as “consist essentially of” or “consisting essentially of,” as used throughout the specification and claims, indicate the inclusion of any recited integer or group of integers, and the optional inclusion of any recited integer

or group of integers that do not materially change the basic or novel properties of the specified method, structure or composition. See M.P.E.P. § 2 111.03.

[0036] It should also be understood that the terms “about,” “approximately,” “generally,” “substantially,” and like terms, used herein when referring to a dimension or characteristic of a component of the preferred invention, indicate that the described dimension/characteristic is not a strict boundary or parameter and does not exclude minor variations therefrom that are functionally the same or similar, as would be understood by one having ordinary skill in the art. At a minimum, such references that include a numerical parameter would include variations that, using mathematical and industrial principles accepted in the art (e.g., rounding, measurement or other systematic errors, manufacturing tolerances, etc.), would not vary the least significant digit.

[0037] The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same, when compared and aligned for maximum correspondence, as measured using one of the following sequence comparison algorithms or by visual inspection. For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

[0038] Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI), or by visual inspection (see generally, *Current Protocols in Molecular Biology*, F. M. Ausubel et al., eds., *Current Protocols*, a joint venture between Greene Publishing Associates, Inc. and John Wiley & Sons, Inc., (1995 Supplement) (Ausubel)).

[0039] Examples of algorithms that are suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1990) *J. Mol. Biol.* 215:403-410 and Altschul et al. (1997) *Nucleic Acids Res.* 25:3389-3402, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length *W* in the query sequence, which either match or satisfy some positive-valued threshold score *T* when aligned with a word of the same length in a database sequence. *T* is referred to as the neighborhood word score threshold (Altschul et al, *supra*). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are then extended in

both directions along each sequence for as far as the cumulative alignment score can be increased.

[0040] Cumulative scores are calculated using, for nucleotide sequences, the parameters *M* (reward score for a pair of matching residues; always >0) and *N* (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity *X* from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters *W*, *T*, and *X* determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (*W*) of 11, an expectation (*E*) of 10, *M*=5, *N*=-4, and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength (*W*) of 3, an expectation (*E*) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1992)).

[0041] In addition to calculating percent sequence identity, the BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin & Altschul, *Proc. Natl. Acad. Sci. USA* 90:5873-5877 (1993)). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (*P(N)*), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.1, more preferably less than about 0.01, and most preferably less than about 0.001.

[0042] A further indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules hybridize to each other under stringent conditions.

[0043] As used herein, the term “polynucleotide,” synonymously referred to as “nucleic acid molecule,” “nucleotides” or “nucleic acids,” refers to any polyribonucleotide or polydeoxyribonucleotide, which can be unmodified RNA or DNA or modified RNA or DNA. “Polynucleotides” include, without limitation single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that can be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, “polynucleotide” refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term polynucleotide also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. “Modified” bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications can be made to DNA and RNA; thus, “poly-

nucleotide” embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. “Polynucleotide” also embraces relatively short nucleic acid chains, often referred to as oligonucleotides.

[0044] As used herein, the terms “peptide,” “polypeptide,” or “protein” can refer to a molecule comprised of amino acids and can be recognized as a protein by those of skill in the art. The conventional one-letter or three-letter code for amino acid residues is used herein. The terms “peptide,” “polypeptide,” and “protein” can be used interchangeably herein to refer to polymers of amino acids of any length. The polymer can be linear or branched, it can comprise modified amino acids, and it can be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art.

[0045] The peptide sequences described herein are written according to the usual convention whereby the N-terminal region of the peptide is on the left and the C-terminal region is on the right. Although isomeric forms of the amino acids are known, it is the L-form of the amino acid that is represented unless otherwise expressly indicated.

Isolated Recombinant Cells and Methods of Making Recombinant Cells

[0046] The inventions disclosed herein are based, at least in part, on the unexpected finding that co-expressing certain chaperone proteins with $\alpha 5$ containing nicotinic acetylcholine receptors (nAChR) (such as $\alpha 2\alpha 5\beta 2$ nAChR) in cells generated cells that highly express $\alpha 5$ containing nAChR, making the cells useful for drug discovery. Provided herein are methods of making recombinant cells expressing $\alpha 5$ subunit of nAChR and isolated recombinant cells for the expression of $\alpha 5$ subunit of nAChR. Also provided herein are methods of making recombinant cells expressing $\alpha 2\alpha 5\beta 2$ nAChR and isolated recombinant cells for the expression of $\alpha 2\alpha 5\beta 2$ nAChR.

[0047] As used herein, the terms “ $\alpha 2\alpha 5\beta 2$ nicotinic acetylcholine receptor”, “ $\alpha 2\alpha 5\beta 2$ nAChR”, “ $\alpha 2\alpha 5\beta 2$ nicotinic acetylcholine receptor”, and “ $\alpha 2\alpha 5\beta 2$ nAChR” are used interchangeably and refer to the $\alpha 2\alpha 5\beta 2$ nicotinic acetylcholine receptor protein, preferably the human $\alpha 2\alpha 5\beta 2$ nAChR, which is a member of a protein family of cholinergic receptors.

[0048] $\alpha 2\alpha 5\beta 2$ nAChR is a ligand-gated ion channel composed of $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits. The $\alpha 2$ subunit is encoded by the gene CHRNA2 (NM_000742), the $\alpha 5$ subunit is encoded by the gene CHRNA5 (NM_000745), and the $\beta 2$ subunit is encoded by the gene CHRN2 (NM_000748). When expressed together, the subunits co-assemble to form an $\alpha 2\alpha 5\beta 2$ nAChR, that in chick is specifically enriched in the optic lobe and in mammal in the habenulo-interpeduncular system (see e.g., Balestra et al., Chick Optic Lobe Contains a Developmentally Regulated $\alpha 2\alpha 5\beta 2$ Nicotinic Receptor Subtype, *MOL* 58:300-311, 2000/1/841787 and Salas et al., “Nicotinic Receptors in the

Habenulo-Interpeduncular System Are Necessary for Nicotine Withdrawal in Mice, *J. Neurosci.* 29:3014-18, 2009).

[0049] “Recombinant cells” refers to one or more individual cells as well as to a recombinant cell line in which the cells are heterologously expressing protein(s). As used herein, “heterologous expression” of a protein in a cell refers to modifying the cell to express the protein by introducing an exogenous nucleic acid into the cell, e.g., an exogenous nucleic acid that encodes the protein to be expressed. A “heterologous nucleic acid” refers to a nucleic acid exogenous to a cell that is introduced into the cell. In some embodiments, the heterologous nucleic acid is DNA. In some embodiments, the heterologous nucleic acid is RNA. Heterologous expression of a protein in a cell can be achieved using a variety of methods. For example, an expression vector comprising a nucleic acid encoding the protein that is operably linked to a nucleic acid encoding a promoter capable of driving expression of the protein (e.g., a constitutive promoter) may be introduced into the cell.

[0050] The term “expression” as used herein, refers to the biosynthesis of a gene product. The term encompasses the transcription of a gene into RNA. The term also encompasses translation of RNA into one or more polypeptides, and further encompasses all naturally occurring post-transcriptional and post-translational modifications.

[0051] In a general aspect, the invention relates to methods of making or generating cells expressing $\alpha 5$ containing nAChR (such as an $\alpha 2\alpha 5\beta 2$ nAChR) that are useful for drug discovery. In one embodiment, the method comprises introducing into a cell a nucleic acid encoding $\alpha 5$ subunit of nAChR and a nucleic acid encoding a chaperone protein selected from transmembrane inner ear expressed protein TMIE and protein FAM163B. The nucleic acid(s) encoding the $\alpha 5$ subunit of nAChR and the chaperone protein (i.e., TMIE or FAM163B) can be in an expression vector (e.g., in a single expression vector or in separate expression vectors). In some embodiments, the nucleic acid encoding the $\alpha 5$ subunit of nAChR and the chaperone protein (i.e., TMIE or FAM163B) is operably linked to a promoter capable of driving expression of the respective protein. In some other embodiments, each of the nucleic acids encoding the $\alpha 5$ subunit of nAChR and the chaperone protein (i.e., TMIE or FAM163B) is operably linked to a promoter capable of driving expression of the respective protein. In some embodiments, the promoter is a constitutive promoter.

[0052] In one aspect, the invention relates to methods of making or generating cells expressing $\alpha 2\alpha 5\beta 2$ nAChR that are useful for drug discovery. In one embodiment, the method comprises introducing into a cell a nucleic acid encoding $\alpha 2$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding $\alpha 5$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding $\beta 2$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding Diamine acetyltransferase 1 (SAT1), a nucleic acid encoding choline O-acetyltransferase (CHAT), and one or both of a nucleic acid encoding TMIE and a nucleic acid encoding FAM163B. The nucleic acid(s) encoding the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits $\alpha 2\alpha 5\beta 2$ nAChR, SAT1, CHAT, and one selected from TMIE and FAM163B can be in an expression vector (e.g., in a single expression vector or in separate expression vectors). In some embodiments, the nucleic acid encoding the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of $\alpha 2\alpha 5\beta 2$ nAChR, SAT1, CHAT, and one or both of TMIE and FAM163B is operably linked to a promoter capable of driving expression of the respective protein. In some other embodiments, each

of the nucleic acids encoding the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of $\alpha 2\alpha 5\beta 2$ nAChR, SAT1, CHAT, and one or both of TMIE and FAM163B is operably linked to a promoter capable of driving expression of the respective protein. In some embodiments, the promoter is a constitutive promoter.

[0053] In another aspect, the method comprises introducing into a cell a nucleic acid encoding $\alpha 2$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding $\alpha 5$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding $\beta 2$ subunit of $\alpha 2\alpha 5\beta 2$ nAChR, a nucleic acid encoding SAT1, a nucleic acid encoding CHAT, and one or both of a nucleic acid encoding TMIE and a nucleic acid encoding FAM163B, wherein the cell generated by the method expresses $\alpha 2\alpha 5\beta 2$ nAChR at an increased level compared to the same cell without the nucleic acids encoding the $\alpha 2$, $\alpha 5$, $\beta 2$ subunits of $\alpha 2\alpha 5\beta 2$ nAChR, and the nucleic acids encoding SAT1, CHAT, TMIE or FAM163B.

[0054] In another aspect, the invention relates to cells genetically modified to express TMIE or FAM163B, wherein the genetically modified cell expresses the at least one protein at an increased level relative to the expression of the same protein in the unmodified cell under the same (or substantially the same) conditions. In one embodiment, the cells are genetically modified to express SAT1, CHAT, and one or both selected from TMIE and FAM163B, wherein genetically modified cell expresses SAT1, CHAT, and one or both selected from TMIE and FAM163B at an increased level relative to the expression of SAT1, CHAT, and one or both selected from TMIE and FAM163B, respectively, in the unmodified cell under the same (or substantially the same) conditions.

[0055] In another aspect, the invention relates to cells genetically modified to express $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of $\alpha 2\alpha 5\beta 2$ nAChR, SAT1, CHAT, and TMIE or FAM163B, wherein the genetically modified cell expresses these proteins protein at an increased level relative to the expression of the same protein in the unmodified cell under the same (or substantially the same) conditions.

[0056] In another aspect, the invention relates to isolated recombinant cells comprising at least one expression vector selected from the group consisting of an expression vector comprising a nucleic acid sequence encoding SAT1, an expression vector comprising a nucleic acid sequence encoding CHAT, and an expression vector comprising a nucleic acid sequence encoding TMIE or FAM163B. In one embodiment, the invention relates to isolated recombinant cells comprising at least one expression vector selected from the group consisting of an expression vector comprising a nucleic acid sequence encoding SAT1, an expression vector comprising a nucleic acid sequence encoding CHAT, an expression vector comprising a nucleic acid sequence encoding TMIE, and an expression vector comprising a nucleic acid sequence encoding FAM163B.

[0057] In one aspect, the invention relates to isolated recombinant cells comprising a heterologous nucleic acid encoding an $\alpha 5$ subunit of nAChR and a heterologous nucleic acid encoding a chaperone protein selected from transmembrane inner ear expressed protein (TMIE) and

protein FAM163B. In one embodiment, the heterologous nucleic acids are introduced into the recombinant cells in the form of expression vectors. In one embodiment, the isolated recombinant cells disclosed herein comprises an expression vector comprising the nucleic acid encoding the $\alpha 5$ subunit of nAChR and an expression vector comprising the nucleic acid encoding the chaperone protein selected from TMIE and FAM163B. In one embodiment, the isolated recombinant cells disclosed herein comprises an expression vector comprising the nucleic acid encoding the $\alpha 5$ subunit of nAChR, an expression vector comprising the nucleic acid encoding TMIE, and an expression vector comprising the nucleic acid encoding FAM163B.

[0058] In a further aspect, the invention relates to isolated recombinant cells comprising a heterologous nucleic acid encoding an $\alpha 2$ subunit of nAChR, a heterologous nucleic acid encoding an $\alpha 5$ subunit of nAChR, a heterologous nucleic acid encoding $\beta 2$ subunit of nAChR, a heterologous nucleic acid encoding diamine acetyltransferase 1 (SAT1), a heterologous nucleic acid encoding choline O-acetyltransferase (CHAT), and a heterologous nucleic acid encoding TMIE and/or FAM163B, wherein the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits forms an $\alpha 2\alpha 5\beta 2$ nAChR. In one embodiment, the heterologous nucleic acids are introduced into the recombinant cells in the form of expression vectors. In one embodiment, the isolated recombinant cells disclosed herein comprises an expression vector comprising the nucleic acid encoding the $\alpha 2$ subunit of nAChR, an expression vector comprising the nucleic acid encoding the $\alpha 5$ subunit of nAChR, an expression vector comprising the nucleic acid encoding the $\beta 2$ subunit of nAChR, an expression vector comprising the nucleic acid encoding SAT1, an expression vector comprising the nucleic acid encoding CHAT, and an expression vector comprising the nucleic acid encoding TMIE or FAM163B. In one embodiment, the isolated recombinant cells disclosed herein comprises an expression vector comprising the nucleic acid encoding the $\alpha 2$ subunit of nAChR, an expression vector comprising the nucleic acid encoding the $\alpha 5$ subunit of nAChR, an expression vector comprising the nucleic acid encoding the $\beta 2$ subunit of nAChR, an expression vector comprising the nucleic acid encoding SAT1, an expression vector comprising the nucleic acid encoding CHAT, an expression vector comprising the nucleic acid encoding TMIE, and an expression vector comprising the nucleic acid encoding FAM163B.

[0059] In any one of the embodiments described herein, the nucleic acid(s) encoding any one or more of the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits nAChR, SAT1, CHAT, TMIE, and FAM163B can be in a single expression vector or in separate expression vectors.

[0060] In one embodiment, the $\alpha 2$ subunit of nAChR comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 1.

SEQ ID NO: 1:
MGPSCPVLSTKLSLWLLLTTPAGGEEAKRPPPRAPGDLSSPSPTAL

PQGGSHTEDETLFKHLFRGYNRWARPVPNTSDVVIVFRGLSLIAQLDIV

DEKNQMMTTNVWLKQEWSDYKLRWNPTDFGNITSLRVPSEMIWIPDIVL

YNNADGEFAVTHMTKAHLFSTGTVHWVPPAIYKSSCSIDVTFFPFDDQN

-continued

CKMKFGSWTYDKAKIDLEQMEQTVDLKDYWESGEWAIVNATGTYNKKY
DCCAETYPDVTYAFVIRRLPLFYTNLIIPCLLISCLTVLVFYLPSDCG
EKITLCISVLLSLTVFLLLITEIIPSTSLVIPLIGEYLLFTMIFVTLSTI
VITVFVLNVHHRSPSTHTMPHWVRGALLGCVPRWLLMNRPPPPVELCHP
LRLKLSPSYHWLESNVDAEEREVVVEEDRWACAGHVAPSVGTLCSHGH
LHSGASGPKAEALLQEGELLLSPHMQKALEGVHYIADHLRSEDADSSVK
EDWKYVAMVIDRIFLWLFIIIVCFGLTIGLFLPPFLAGMI

[0061] In one embodiment, the $\alpha 5$ subunit of nAChR comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 2.

SEQ ID NO: 2
MAARGSPRALRLLLLVLQVAGRCGLAGAAGGAQRGLSEPSIAKHEDS
LLKDLFQDYERWVRPVEHLNDKIKIKFGLAISQLVDVDEKNQMLMTNVW
LKQEWIDVKLRWNPDDYGGIKVIRVPSDSVWTPDIVLFDNADGRFEGTS
TKTVIRYNGTVTWTTPANYKSSCTIDVTFFPDLQNCMKFGSWTYDGS
QVDIILEDQDVKRDFDNGEWEIVSATGSKGNRTDSCCWYPYVTYSFV
IKRLPLFYTLFLIIPICIGLSFLTIVLVFYLPSNEGEKICLCTSVLVSLTV
FLLVIEEIIIPSSSKVIPLIGEYLVFTMIFVTLSIMVTVFAINIHRSSS
THNAMAPLVRKIFLHTLPKLLCMRSHVDYRFTQKEETESGSGPKSSRNT
LEAALDSIRYITRHMKEVDREVVEDWKFIQVLDRLMFLWTFVSVIV
GSLGLFVPVIYKWNILIPVHIGNANK

[0062] In one embodiment, the $\beta 2$ subunit of nAChR comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 3.

SEQ ID NO: 3
MARRCGPVALLLGFGLLRLCSGVWGTDTTEERLVEHLDPSTYNKLRPA
TNGSELVTQVMVSLAQ LISVHEREQIMTTNVWLTQEWEDYRLTWKPEE
FDNMKKVRLPSKHIWLPDVLVLYNNADGMYEVSFYSSNAVVSVDGSIWFLP
PAIYKSACKIEVKHFFPDQONCTMKFRSWTYDRTEIDLVLKSEVASLDD
FTPSGEWDIVLPGRRNENPDDSTYVDITYDFIIRKPLFYTNLIIPC
VLITSLAILVFYLPSPDCGKMTLCISVLLALTIVFLLLIKIVPPTSLDV
PLVGKYLMTMVLVTFISIVTSCVVLNVHHRSPTHHTMAPWVKVVFLEKL
PALLFMQQPRHHCARQLRLRRRQREREGAGALFFREAPGADSCTCFVN
RASVQGLAGAFGAEPAPVAGPGRSGEPCCGLREAVDGVRFIADHMRSE
DDDQSVSEDWKYVAMVIDRLFLWIFVFCVFTIGMFLQPLFQNYTTT
FLHSDHSAAPSSK

[0063] In one embodiment, the TMIE comprises an amino acid sequence with at least 85%, or at least 90%, or at least

95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 4. Or, the TMIE comprises an amino acid sequence with at least 60%, at least 70%, at least 80%, at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 4 and has protein chaperone property, which includes the property of enhancing the expression of the $\alpha 5$ subunit of nAChR.

SEQ ID NO: 4
MAGWPGAGPLCVLGGALGVCLAGVAGQLVEPSTAPPKPKPPPLTKETV
VFWDMRLWHVVGIFSLFVLSIIITLCCVFNCRVPRTRKEIEARYLQRKA
AKMYTDKLETVPPLNELTEVPGEDKKKKKKKKKSDVTVAIKVEEDEKN
EAKKKKGEK

[0064] In one embodiment, the FAM163B comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 5. Or, the FAM163B comprises an amino acid sequence with at least 60%, at least 70%, at least 80%, at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 5 and has protein chaperone property, which includes the property of enhancing the expression of the $\alpha 5$ subunit of nAChR.

SEQ ID NO: 5
MTAGTVVITGGILATVILLCI IAVLCYCRLQYCYCKKDESEDEEEDP
AVHSHLPPLHSNRNLVLTNGPALYPTASTSFSQKSPQARALCRSCSHCE
PPTFFLQEPPEEEDVLNGGERVLYKSVSQEDVELPPGGFGGLQALNPN
RLSAMREAFARSRSISTDV

[0065] In one embodiment, the SAT1 comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 6. Or, the SAT1 comprises an amino acid sequence with at least 60%, at least 70%, at least 80%, at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 6 and has protein chaperone property, which includes the property of enhancing the expression of the $\alpha 2$ and $\beta 2$ subunits of nAChR.

SEQ ID NO: 6
MAKFVIRPATAADCSDILRLIKELAKYEYMEEQVILTEKDLLEDGFGHEH
PFYHCLVAEVPKEHWTPEGHSIVGFAMYFYFTDPWIGKLLYLEDFVMS
DYRFGIGSEILKNLSQVAMRCRCSSMHFLVAEWNEPSINFYKRRGASD
LSSEEGWRLFKIDKEYLLKMATEE

[0066] In one embodiment, the CHAT comprises an amino acid sequence with at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 7. Or, the CHAT comprises an amino acid sequence with at least 60%, at least 70%, at least 80%, at least 85%, or at least 90%, or at least 95%, such as 95%, 96%, 97%, 98%, 99%, or 100% identity to the amino acid sequence of SEQ ID NO: 7 and has protein

chaperone property, which includes the property of enhancing the expression of the $\alpha 2$ and $\beta 2$ Subunits of nAChR.

SEQ ID NO: 7

```

MAAKTPSSEESGLPKLPVPPLQQTTLATYLQCMRHLVSEEQFRKSQAIIVQ
QFGAPGGLGETLQQKLLERQEKANTANVWSEYWLNDMYLNNRLALPVNSSP
AVIFARQHFPGETDDQLRFAASLISGVLSYKALLDSHSIPTDCAKGQLSG
QPLCMKQYYGLFSSYRLPGHTQDTLVAQNSSIMPEPEHVIVACCNQFFV
LDVVINFRRLSEGDLFTQLRKIVKMASNEDERLPPIGLLTSDGRSEWAE
ARTVLVKDSTNRDSLDMIERICLVCLDAPGGVELSDTHRALQLLHGGG
YSKNGANRWYDKSLQFVVGRDTCGVVCEHSPFDGIVLVQCTEHLKHM
TQSSRKILIRADSVSELPAPRRRLRWKCSPEIQGHLASSAEKLQRIVKNLD
FIVYKFDNYGKTFIKKQKSPDAFIQVALQLAFYRLHRRLVPTYESASI
RRFQEGRVNIRSATPEALFVRAVTDHKAAPVASEKLLLLKDAIRAQT
AYTVMAITGMAIDNHLALRELARAMCKELPEMFMDETYLMNRFVLST
SQVPTTTEMFCCYGPVVPNGYGACYNPQPETILFCISSFHSCKETSSSK
FAKAVEESLIDMRDLCSLLPPTESKPLATKEKATRPSQGHQP

```

[0067] Any suitable means for introducing heterologous nucleic acid into a cell can be used herein to prepare the recombinant cells disclosed herein, such as DNA transfection (e.g., via a DNA vector) and RNA transduction. In one embodiment, the heterologous nucleic acid to be introduced into cells to generate the recombinant cells are prepared by using a vector, preferably an expression vector. The term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double-stranded DNA loop into which additional DNA segments can be inserted. Another type of vector is a viral vector wherein additional DNA segments can be inserted. Expression vectors are those vectors capable of directing the expression of genes to which they are operably linked. The expression vectors used herein comprise a nucleic acid encoding a protein sequence in a form suitable for expression of the nucleic acid in a host cell. Thus, the expression vectors can include one or more regulatory sequences, such as a promoter, selected on the basis of the host cells to be used for expression, operably linked to the nucleic acid sequence to be expressed. When used in reference to a expression vector, “operably linked” is intended to mean that the nucleotide sequence of interest is linked to the regulatory sequence(s) in a manner allowing for expression of the nucleotide sequence (e.g., in an in vitro transcription/translation system or in a host cell when the vector is introduced into the host cell). It will be appreciated by those of ordinary skill in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed and the level of expression of protein desired as well as the intended use of the vector.

[0068] Any vector known to those skilled in the art in view of the present disclosure can be used, such as a plasmid, a cosmid, a phage vector, or a viral vector. In one embodiment, the vector is an expression vector such as a plasmid. The vector can include any element to establish a conventional function of an expression vector, for example, a promoter, ribosome binding element, terminator, enhancer, selection

marker, and origin of replication. The promoter can be a constitutive, inducible or repressible promoter. A number of expression vectors capable of delivering nucleic acids to a cell are known in the art and can be used herein. Conventional cloning techniques or artificial gene synthesis can be used to generate an expression vector according to embodiments of the invention.

[0069] Any cell known to those skilled in the art in view of the present disclosure can be used for recombinant expression of $\alpha 2$, $\alpha 5$, $\beta 2$ subunits of nAChR, SAT1, CHAT, TMIE, and FAM163B. In one embodiment, the recombinant cell is a mammalian cell. Suitable the mammalian cells may be selected from human embryonic kidney 293T (HEK293T) cell, HEK293F cells, Hela cells, Chinese hamster ovary (CHO) cells, NIH 3T3 cells, MCF-7 cells, Hep G2 cells, baby hamster kidney (BHK) cells, and Cos7 cells.

Methods of Identifying Agonists, Antagonists, or Positive Allosteric Modulators of $\alpha 2\alpha 5\beta 2$ nAChR

[0070] Further provided herein are methods of identifying agonists, antagonists, or positive allosteric modulators (PAMs) of an $\alpha 2\alpha 5\beta 2$ nAChR. PAMs are compounds that bind at sites on the protein surface other than the active sites, and therefore change the conformation of the protein binding sites.

[0071] In one embodiment, the method comprises culturing the isolated recombinant cells disclosed herein under conditions where the recombinant cells grow, contacting the recombinant cells with an agent, and determining if the agent is an agonist, antagonist, or PAM of the $\alpha 2\alpha 5\beta 2$ nAChR, wherein an agonist or PAM enhances the activity of the $\alpha 2\alpha 5\beta 2$ nAChR and an antagonist decreases the activity of the $\alpha 2\alpha 5\beta 2$ nAChR as compared to the activity of the $\alpha 2\alpha 5\beta 2$ nAChR in a recombinant cell that was not contacted with an agent. Agonists, as used herein, refer to molecules/compounds/peptides that serve to enhance the function of the $\alpha 2\alpha 5\beta 2$ nAChR. PAMs, as used herein, refer to molecules/compounds/peptides that enhance the effect of $\alpha 2\alpha 5\beta 2$ nAChR's response to a ligand without directly activating the receptor. As used herein, the term “enhance”, “enhanced”, “increase”, or “increased”, when used with respect to $\alpha 2\alpha 5\beta 2$ nAChR activity refers to an increase in the signaling through the receptor, relative to the corresponding signaling observed in a cell in which an agonist or PAM is not administered. Antagonists, as used herein, refer to molecules/compounds/peptides that serve to block, decrease, or dampen the function of the $\alpha 2\alpha 5\beta 2$ nAChR. In particular embodiments, the agent is a small molecule or peptide.

[0072] In one embodiment, FLIPR assay is used to identify agonists of $\alpha 2\alpha 5\beta 2$ nAChR mediated calcium flux. In this assay, the recombinant cells are incubated with a calcium sensitive dye (such as Ca5), exposed to a test compound, and calcium flux is imaged by FLIPR^{TETRA}.

[0073] In one embodiment, FLIPR assay is used to identify antagonists of nicotine-evoked $\alpha 2\alpha 5\beta 2$ nAChR mediated calcium flux. In this assay, the recombinant cells are incubated with a calcium sensitive dye (such as Ca5). Using a double addition protocol, the recombinant cells are exposed to test compounds during the first duration, and to nicotine (e.g., at EC₈₀ (1 μ M)) during the second duration. Thereafter, the calcium flux is imaged by FLIPR^{TETRA}. As shown in FIG. 2, antagonists of $\alpha 2\alpha 5\beta 2$ nAChR reduce calcium flux.

[0074] In one embodiment, FLIPR assay is used to identify compounds that positively modulate or potentiate nicotine-evoked $\alpha 2\alpha 5\beta 2$ nAChR mediated calcium flux. In this assay, the recombinant cells are incubated with a calcium sensitive dye (such as Ca5). Using a double addition protocol, the recombinant cells are exposed to test compounds during the first duration, and to nicotine (e.g., at EC_{80} (1 μ M)) during the second duration. Thereafter, the calcium flux is imaged by FLIPR^{TETRA}. If a test compound enhances nicotine-evoked $\alpha 2\alpha 5\beta 2$ nAChR mediated calcium flux, yet does not enhances calcium flux as an agonist (determined as above), the test compound is termed a positive allosteric modulator (PAM) or potentiator of $\alpha 2\alpha 5\beta 2$ nAChR.

[0075] In a preferred embodiment, the cells are incubated at about 25-35° C. for about 20-50 hr prior to the FLIPR assay or other assay for measuring $\alpha 2\alpha 5\beta 2$ nAChR activity. In some embodiments, the cells are incubated at about 25° C., about 26° C., about 27° C., about 28° C., about 29° C., about 30° C., about 31° C., about 32° C., about 33° C., about 34° C., or about 35° C., for about 20 hr, about 25 hr, about 30 hr, about 35 hr, about 40 hr, about 45 hr, or about 50 hr prior to the assay.

Expression System and Kits

[0076] Further provided herein are expression systems and kits comprising the isolated recombinant cells. The expression systems and kits may further include instructions for use

Example

[0077] It will be appreciated by those skilled in the art that changes could be made to the embodiments described above without departing from the broad inventive concept thereof. It is understood, therefore, that this invention is not limited to the particular embodiments disclosed, but it is intended to cover modifications within the spirit and scope of the present invention as defined by the present description.

Material

Name	Company	Cat. No
FreeStyle 293 Expression Media	Gibco	12338018
OptiPro SFM	Gibco	12309019
FreeStyle MAX Reagent	Invitrogen	16447100
Recovery Cell Culture Freezing Media	Gibco	12648010
DMEM-high glucose	Sigma	D5671-500ML
Defined FBS	HyClone	SH30070.03
Sodium Pyruvate	HyClone	SH30239.01
Glutamax	HyClone	35050061
DMSO	Sigma Aldrich	41648
Ca5 dye	Molecular Devices	R8186
Nicotine	Tocris	3546
384-well pp plates	Greiner	781280
384-well poly-D-lysine plates	Corning	354663
Custom HBSS	HyClone	SH3A4475.01
Calcium Chloride	Amresco	E506-500ML
Magnesium Chloride	Amresco	E525-500ML
Freezing Vial	Nalgene	5000-0020

Cell Line

[0078] FreeStyle 293F cell line (ThermoFisher Scientific Cat #R79007)

Vector Constructs

[0079] pcDNA3.1- $\alpha 2$
[0080] pcDNA3.1- $\alpha 5$
[0081] pcDNA3.1- $\beta 2$
[0082] pcDNA3.1-SAT1
[0083] pcDNA3.1-CHAT
[0084] pcDNA3.1-TMIE
[0085] pcDNA3.1-FAM163B

Culturing Media

[0086] FreeStyle 293 Expression Medium

Transfection Reagents

[0087] FreeStyle MAX Reagent

Planting Media

[0088] DMEM
[0089] 10% FBS
[0090] 1% pen-strep

Compound Dilution/Ca5 Buffer.

[0091] HEPES-buffered saline solution (500 ml) supplemented with 1 mM Mg^{2+} and 2 mM Ca^{2+}

Calcium 5 (Ca5) Dye (Molecular Devices)

[0092] Calcium dye diluted at 25 $\times$ concentration in Hanks Buffered Salt Solution. Dye diluted to 1 $\times$ in HEPES assay buffer with 2 mM $CaCl_2$) and 1 mM $MgCl_2$.

Methods and Procedures

Day 1 Cell Preparation

[0093] Prepare a cell suspension of 1×10^6 cells per ml by centrifuging cells and resuspending in fresh, pre-warmed media;
[0094] With 1900 ml of cells shaking, prepare the following transfection mix in 50 ml of OptiPro SFM medium;
[0095] Add 2.5 ml FreeStyle MAX reagent to 47.5 ml of OptiPro SFM to an empty 50 ml tube;
[0096] In a second 50 ml tube, add 2.5 mg total DNA ($\alpha 2$, $\alpha 5$, $\beta 2$, SAT1, CHAT, and TMIE or FAM163B at a ratio of 3:3:3:3:1:2) in OptiPro SFM to a final volume of 50 ml and vortex gently;
[0097] Transfer the diluted DNA to the FreeStyle MAX reagent and gently mix the solution immediately and incubate for 10 minutes at room temperature;
[0098] Transfer the 100 ml FreeStyle MAX/DNA transfection mix to the cells and homogenize the culture;
[0099] Incubate 1 hour at 37° C. with 8% CO_2 ;
[0100] Pour cells into sterile 500 ml tubes;
[0101] Spin at 1000 rpm for 7 minutes;
[0102] Remove supernatant; and
[0103] Resuspend Cells in freezing media, aliquot into cryotube and frozen in -80° C.

Day 2 Plate Cells

- [0104] Ethanol vials to decontaminate;
 [0105] Pour cells into 50 ml conical tube;
 [0106] Slowly add 10 ml warmed plating media drop wise;
 [0107] Pipet up and down gently a few times to dislodge pellet if it exists;
 [0108] Spin at 1000 RPM for 7 min, aspirate supernatant, and resuspend in 25 ml plating;
 [0109] Seed 35,000 cells/well in 35 μ l (1.0×10^6 cells/ml); and
 [0110] Incubate plates for 24 hours at 37° C. Then transfer the plate to 30° C. with 5% CO₂ in a humidified atmosphere for 24-48 hours.

Day 3 FLIPR Assay

- [0111] Wash plate with plate washer (4 washes $\times$ 100 μ l/wash) using compound dilution buffer, leaving 25 μ l in each well;
 [0112] Add 25 μ l of 2 $\times$ Ca5 dye to each well;
 [0113] Incubate at room temperature for 1 hour;
 [0114] Wash plate with plate washer (4 washes $\times$ 100 μ l/wash) using compound dilution buffer, leaving 50 μ l in each well;
 [0115] Transfer plates to FLIPR for compound addition.
 Functional Expression of $\alpha 2\beta 2$ nAChR and $\alpha 2\alpha 5\beta 2$ nAChR
 [0116] Human $\alpha 2$ and $\beta 2$ subunits of $\alpha 2\beta 2$ nAChR were co-transfected with specified combinations of cDNAs (vector alone; SAT1+CHAT; SAT1+CHAT+TMIE, or SAT1+CHAT+FAM163B) in HEK293T cells and incubated at 37° C. overnight followed by 30° C. for 24-48 hours. The transfected cells were incubated for one hour at room temperature with Ca5 dye followed by stimulation with E_{max}

nicotine (33 μ M). Nicotine-evoked Ca²⁺ signal of the transfected are graphed in FIG. 1 (left). As shown, co-transfection with SAT1 and CHAT enhances $\alpha 2\beta 2$ nAChR function (i.e., nicotine-evoked calcium flux), while co-transfection with TMIE or FAM163B does not affect $\alpha 2\beta 2$ nAChR function.

[0117] Separately, human $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of $\alpha 2\alpha 5\beta 2$ nAChR were co-transfected with specified combinations of cDNAs (vector alone; SAT1+CHAT; SAT1+CHAT+TMIE, or SAT1+CHAT+FAM163B) in HEK293T cells and incubated at 37° C. overnight followed by 30° C. for 24-48 hours. The transfected cells were incubated for one hour at room temperature with Ca5 dye followed by stimulation with E_{max} nicotine (33 μ M). Nicotine-evoked Ca²⁺ signal of the transfected are graphed in FIG. 1 (right). As shown, TMIE or FAM163B co-transfection (in the presence of SAT1 and CHAT) dramatically enhances $\alpha 2\beta 2\alpha 5$ nAChR function (i.e., nicotine-evoked calcium flux).

Concentration Response Curve for Antagonist on Nicotine-Evoked Activity $\alpha 2\beta 2$ nAChR and $\alpha 2\alpha 5\beta 2$ nAChR

[0118] Human nAChR subunits $\alpha 2$ and $\beta 2$ subunits with or without $\alpha 5$ were co-transfected with specified combinations of cDNAs (SAT1+CHAT; SAT1+CHAT+TMIE, or SAT1+CHAT+FAM163B) in HEK293T cells. Following incubation at 37° C. overnight cells were incubated at 30° C. for an additional 24-48 hours. Cells were then incubated with Ca5 dye for one hour at room temperature and stimulated with various of concentrations of nicotine. Nicotine-evoked calcium flux was recorded using a FLIPR^{TETRA} imager. The FLIPR signals were averaged and plotted with percentage to maximum responses per each transfection. As shown in FIG. 2, co-transfection with FAM163B or TMIE left shifted the nicotine potency on both $\alpha 2\beta 2$ nAChR and $\alpha 2\beta 2\alpha 5$ nAChR.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 529

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```
Met Gly Pro Ser Cys Pro Val Phe Leu Ser Phe Thr Lys Leu Ser Leu
1              5              10              15

Trp Trp Leu Leu Leu Thr Pro Ala Gly Gly Glu Glu Ala Lys Arg Pro
20              25              30

Pro Pro Arg Ala Pro Gly Asp Pro Leu Ser Ser Pro Ser Pro Thr Ala
35              40              45

Leu Pro Gln Gly Gly Ser His Thr Glu Thr Glu Asp Arg Leu Phe Lys
50              55              60

His Leu Phe Arg Gly Tyr Asn Arg Trp Ala Arg Pro Val Pro Asn Thr
65              70              75              80

Ser Asp Val Val Ile Val Arg Phe Gly Leu Ser Ile Ala Gln Leu Ile
85              90              95

Asp Val Asp Glu Lys Asn Gln Met Met Thr Thr Asn Val Trp Leu Lys
100             105             110

Gln Glu Trp Ser Asp Tyr Lys Leu Arg Trp Asn Pro Thr Asp Phe Gly
115             120             125
```

-continued

Asn	Ile	Thr	Ser	Leu	Arg	Val	Pro	Ser	Glu	Met	Ile	Trp	Ile	Pro	Asp
130						135					140				
Ile	Val	Leu	Tyr	Asn	Asn	Ala	Asp	Gly	Glu	Phe	Ala	Val	Thr	His	Met
145				150					155						160
Thr	Lys	Ala	His	Leu	Phe	Ser	Thr	Gly	Thr	Val	His	Trp	Val	Pro	Pro
			165					170						175	
Ala	Ile	Tyr	Lys	Ser	Ser	Cys	Ser	Ile	Asp	Val	Thr	Phe	Phe	Pro	Phe
			180					185					190		
Asp	Gln	Gln	Asn	Cys	Lys	Met	Lys	Phe	Gly	Ser	Trp	Thr	Tyr	Asp	Lys
	195					200						205			
Ala	Lys	Ile	Asp	Leu	Glu	Gln	Met	Glu	Gln	Thr	Val	Asp	Leu	Lys	Asp
	210					215					220				
Tyr	Trp	Glu	Ser	Gly	Glu	Trp	Ala	Ile	Val	Asn	Ala	Thr	Gly	Thr	Tyr
225				230					235						240
Asn	Ser	Lys	Lys	Tyr	Asp	Cys	Cys	Ala	Glu	Ile	Tyr	Pro	Asp	Val	Thr
			245						250					255	
Tyr	Ala	Phe	Val	Ile	Arg	Arg	Leu	Pro	Leu	Phe	Tyr	Thr	Ile	Asn	Leu
		260					265						270		
Ile	Ile	Pro	Cys	Leu	Leu	Ile	Ser	Cys	Leu	Thr	Val	Leu	Val	Phe	Tyr
		275					280					285			
Leu	Pro	Ser	Asp	Cys	Gly	Glu	Lys	Ile	Thr	Leu	Cys	Ile	Ser	Val	Leu
	290					295					300				
Leu	Ser	Leu	Thr	Val	Phe	Leu	Leu	Leu	Ile	Thr	Glu	Ile	Ile	Pro	Ser
305				310						315					320
Thr	Ser	Leu	Val	Ile	Pro	Leu	Ile	Gly	Glu	Tyr	Leu	Leu	Phe	Thr	Met
			325						330					335	
Ile	Phe	Val	Thr	Leu	Ser	Ile	Val	Ile	Thr	Val	Phe	Val	Leu	Asn	Val
		340					345						350		
His	His	Arg	Ser	Pro	Ser	Thr	His	Thr	Met	Pro	His	Trp	Val	Arg	Gly
		355				360						365			
Ala	Leu	Leu	Gly	Cys	Val	Pro	Arg	Trp	Leu	Leu	Met	Asn	Arg	Pro	Pro
	370				375						380				
Pro	Pro	Val	Glu	Leu	Cys	His	Pro	Leu	Arg	Leu	Lys	Leu	Ser	Pro	Ser
385				390						395					400
Tyr	His	Trp	Leu	Glu	Ser	Asn	Val	Asp	Ala	Glu	Glu	Arg	Glu	Val	Val
			405						410					415	
Val	Glu	Glu	Glu	Asp	Arg	Trp	Ala	Cys	Ala	Gly	His	Val	Ala	Pro	Ser
		420					425					430			
Val	Gly	Thr	Leu	Cys	Ser	His	Gly	His	Leu	His	Ser	Gly	Ala	Ser	Gly
	435						440					445			
Pro	Lys	Ala	Glu	Ala	Leu	Leu	Gln	Glu	Gly	Glu	Leu	Leu	Leu	Ser	Pro
	450					455					460				
His	Met	Gln	Lys	Ala	Leu	Glu	Gly	Val	His	Tyr	Ile	Ala	Asp	His	Leu
465				470						475					480
Arg	Ser	Glu	Asp	Ala	Asp	Ser	Ser	Val	Lys	Glu	Asp	Trp	Lys	Tyr	Val
			485						490					495	
Ala	Met	Val	Ile	Asp	Arg	Ile	Phe	Leu	Trp	Leu	Phe	Ile	Ile	Val	Cys
		500					505						510		
Phe	Leu	Gly	Thr	Ile	Gly	Leu	Phe	Leu	Pro	Pro	Phe	Leu	Ala	Gly	Met
	515					520						525			

-continued

Ile

<210> SEQ ID NO 2

<211> LENGTH: 468

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Ala Ala Arg Gly Ser Gly Pro Arg Ala Leu Arg Leu Leu Leu Leu
1 5 10 15
Val Gln Leu Val Ala Gly Arg Cys Gly Leu Ala Gly Ala Ala Gly Gly
20 25 30
Ala Gln Arg Gly Leu Ser Glu Pro Ser Ser Ile Ala Lys His Glu Asp
35 40 45
Ser Leu Leu Lys Asp Leu Phe Gln Asp Tyr Glu Arg Trp Val Arg Pro
50 55 60
Val Glu His Leu Asn Asp Lys Ile Lys Ile Lys Phe Gly Leu Ala Ile
65 70 75 80
Ser Gln Leu Val Asp Val Asp Glu Lys Asn Gln Leu Met Thr Thr Asn
85 90 95
Val Trp Leu Lys Gln Glu Trp Ile Asp Val Lys Leu Arg Trp Asn Pro
100 105 110
Asp Asp Tyr Gly Gly Ile Lys Val Ile Arg Val Pro Ser Asp Ser Val
115 120 125
Trp Thr Pro Asp Ile Val Leu Phe Asp Asn Ala Asp Gly Arg Phe Glu
130 135 140
Gly Thr Ser Thr Lys Thr Val Ile Arg Tyr Asn Gly Thr Val Thr Trp
145 150 155 160
Thr Pro Pro Ala Asn Tyr Lys Ser Ser Cys Thr Ile Asp Val Thr Phe
165 170 175
Phe Pro Phe Asp Leu Gln Asn Cys Ser Met Lys Phe Gly Ser Trp Thr
180 185 190
Tyr Asp Gly Ser Gln Val Asp Ile Ile Leu Glu Asp Gln Asp Val Asp
195 200 205
Lys Arg Asp Phe Phe Asp Asn Gly Glu Trp Glu Ile Val Ser Ala Thr
210 215 220
Gly Ser Lys Gly Asn Arg Thr Asp Ser Cys Cys Trp Tyr Pro Tyr Val
225 230 235 240
Thr Tyr Ser Phe Val Ile Lys Arg Leu Pro Leu Phe Tyr Thr Leu Phe
245 250 255
Leu Ile Ile Pro Cys Ile Gly Leu Ser Phe Leu Thr Val Leu Val Phe
260 265 270
Tyr Leu Pro Ser Asn Glu Gly Glu Lys Ile Cys Leu Cys Thr Ser Val
275 280 285
Leu Val Ser Leu Thr Val Phe Leu Leu Val Ile Glu Glu Ile Ile Pro
290 295 300
Ser Ser Ser Lys Val Ile Pro Leu Ile Gly Glu Tyr Leu Val Phe Thr
305 310 315 320
Met Ile Phe Val Thr Leu Ser Ile Met Val Thr Val Phe Ala Ile Asn
325 330 335
Ile His His Arg Ser Ser Ser Thr His Asn Ala Met Ala Pro Leu Val
340 345 350

-continued

Arg	Lys	Ile	Phe	Leu	His	Thr	Leu	Pro	Lys	Leu	Leu	Cys	Met	Arg	Ser
		355					360					365			
His	Val	Asp	Arg	Tyr	Phe	Thr	Gln	Lys	Glu	Glu	Thr	Glu	Ser	Gly	Ser
	370					375					380				
Gly	Pro	Lys	Ser	Ser	Arg	Asn	Thr	Leu	Glu	Ala	Ala	Leu	Asp	Ser	Ile
385					390					395					400
Arg	Tyr	Ile	Thr	Arg	His	Ile	Met	Lys	Glu	Asn	Asp	Val	Arg	Glu	Val
			405						410					415	
Val	Glu	Asp	Trp	Lys	Phe	Ile	Ala	Gln	Val	Leu	Asp	Arg	Met	Phe	Leu
		420						425				430			
Trp	Thr	Phe	Leu	Phe	Val	Ser	Ile	Val	Gly	Ser	Leu	Gly	Leu	Phe	Val
	435						440					445			
Pro	Val	Ile	Tyr	Lys	Trp	Ala	Asn	Ile	Leu	Ile	Pro	Val	His	Ile	Gly
	450					455					460				
Asn	Ala	Asn	Lys												
465															

<210> SEQ ID NO 3
 <211> LENGTH: 502
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Met	Ala	Arg	Arg	Cys	Gly	Pro	Val	Ala	Leu	Leu	Leu	Gly	Phe	Gly	Leu
1			5					10					15		
Leu	Arg	Leu	Cys	Ser	Gly	Val	Trp	Gly	Thr	Asp	Thr	Glu	Glu	Arg	Leu
		20					25					30			
Val	Glu	His	Leu	Leu	Asp	Pro	Ser	Arg	Tyr	Asn	Lys	Leu	Ile	Arg	Pro
	35					40					45				
Ala	Thr	Asn	Gly	Ser	Glu	Leu	Val	Thr	Val	Gln	Leu	Met	Val	Ser	Leu
	50				55					60					
Ala	Gln	Leu	Ile	Ser	Val	His	Glu	Arg	Glu	Gln	Ile	Met	Thr	Thr	Asn
65				70					75						80
Val	Trp	Leu	Thr	Gln	Glu	Trp	Glu	Asp	Tyr	Arg	Leu	Thr	Trp	Lys	Pro
		85						90					95		
Glu	Glu	Phe	Asp	Asn	Met	Lys	Lys	Val	Arg	Leu	Pro	Ser	Lys	His	Ile
	100						105					110			
Trp	Leu	Pro	Asp	Val	Val	Leu	Tyr	Asn	Asn	Ala	Asp	Gly	Met	Tyr	Glu
	115					120					125				
Val	Ser	Phe	Tyr	Ser	Asn	Ala	Val	Val	Ser	Tyr	Asp	Gly	Ser	Ile	Phe
	130				135						140				
Trp	Leu	Pro	Pro	Ala	Ile	Tyr	Lys	Ser	Ala	Cys	Lys	Ile	Glu	Val	Lys
145				150					155						160
His	Phe	Pro	Phe	Asp	Gln	Gln	Asn	Cys	Thr	Met	Lys	Phe	Arg	Ser	Trp
			165					170						175	
Thr	Tyr	Asp	Arg	Thr	Glu	Ile	Asp	Leu	Val	Leu	Lys	Ser	Glu	Val	Ala
		180					185						190		
Ser	Leu	Asp	Asp	Phe	Thr	Pro	Ser	Gly	Glu	Trp	Asp	Ile	Val	Ala	Leu
	195					200					205				
Pro	Gly	Arg	Arg	Asn	Glu	Asn	Pro	Asp	Asp	Ser	Thr	Tyr	Val	Asp	Ile
	210				215						220				
Thr	Tyr	Asp	Phe	Ile	Ile	Arg	Arg	Lys	Pro	Leu	Phe	Tyr	Thr	Ile	Asn
225				230						235					240

```

<210> SEQ ID NO 4
<211> LENGTH: 156
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Ala Gly Trp Pro Gly Ala Gly Pro Leu Cys Val Leu Gly Gly Ala
1          5          10          15
Ala Leu Gly Val Cys Leu Ala Gly Val Ala Gly Gln Leu Val Glu Pro
          20          25          30
Ser Thr Ala Pro Pro Lys Pro Lys Pro Pro Pro Leu Thr Lys Glu Thr
          35          40          45
Val Val Phe Trp Asp Met Arg Leu Trp His Val Val Gly Ile Phe Ser
          50          55          60
Leu Phe Val Leu Ser Ile Ile Ile Thr Leu Cys Cys Val Phe Asn Cys
65          70          75          80
Arg Val Pro Arg Thr Arg Lys Glu Ile Glu Ala Arg Tyr Leu Gln Arg

```

```
<210> SEQ ID NO 5
<211> LENGTH: 166
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

[illegible]

```
<210> SEQ ID NO 6
<211> LENGTH: 171
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens
```

Met	Ala	Lys	Phe	Val	Ile	Arg	Pro	Ala	Thr	Ala	Ala	Asp	Cys	Ser	Asp
1				5					10					15	
Ile	Leu	Arg	Leu	Ile	Lys	Glu	Leu	Ala	Lys	Tyr	Glu	Tyr	Met	Glu	Glu
			20					25					30		
Gln	Val	Ile	Leu	Thr	Glu	Lys	Asp	Leu	Leu	Glu	Asp	Gly	Phe	Gly	Glu
			35				40					45			
His	Pro	Phe	Tyr	His	Cys	Leu	Val	Ala	Glu	Val	Pro	Lys	Glu	His	Trp
	50					55					60				
Thr	Pro	Glu	Gly	His	Ser	Ile	Val	Gly	Phe	Ala	Met	Tyr	Tyr	Phe	Thr

-continued

65	70	75	80
Tyr Asp Pro Trp Ile Gly Lys Leu Leu Tyr Leu Glu Asp Phe Phe Val			
	85	90	95
Met Ser Asp Tyr Arg Gly Phe Gly Ile Gly Ser Glu Ile Leu Lys Asn			
	100	105	110
Leu Ser Gln Val Ala Met Arg Cys Arg Cys Ser Ser Met His Phe Leu			
	115	120	125
Val Ala Glu Trp Asn Glu Pro Ser Ile Asn Phe Tyr Lys Arg Arg Gly			
	130	135	140
Ala Ser Asp Leu Ser Ser Glu Glu Gly Trp Arg Leu Phe Lys Ile Asp			
	145	150	155
Lys Glu Tyr Leu Leu Lys Met Ala Thr Glu Glu			
	165	170	

<210> SEQ ID NO 7

<211> LENGTH: 630

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Met Ala Ala Lys Thr Pro Ser Ser Glu Glu Ser Gly Leu Pro Lys Leu			
1	5	10	15
Pro Val Pro Pro Leu Gln Gln Thr Leu Ala Thr Tyr Leu Gln Cys Met			
	20	25	30
Arg His Leu Val Ser Glu Glu Gln Phe Arg Lys Ser Gln Ala Ile Val			
	35	40	45
Gln Gln Phe Gly Ala Pro Gly Gly Leu Gly Glu Thr Leu Gln Gln Lys			
	50	55	60
Leu Leu Glu Arg Gln Glu Lys Thr Ala Asn Trp Val Ser Glu Tyr Trp			
	65	70	75
Leu Asn Asp Met Tyr Leu Asn Asn Arg Leu Ala Leu Pro Val Asn Ser			
	85	90	95
Ser Pro Ala Val Ile Phe Ala Arg Gln His Phe Pro Gly Thr Asp Asp			
	100	105	110
Gln Leu Arg Phe Ala Ala Ser Leu Ile Ser Gly Val Leu Ser Tyr Lys			
	115	120	125
Ala Leu Leu Asp Ser His Ser Ile Pro Thr Asp Cys Ala Lys Gly Gln			
	130	135	140
Leu Ser Gly Gln Pro Leu Cys Met Lys Gln Tyr Tyr Gly Leu Phe Ser			
	145	150	155
Ser Tyr Arg Leu Pro Gly His Thr Gln Asp Thr Leu Val Ala Gln Asn			
	165	170	175
Ser Ser Ile Met Pro Glu Pro Glu His Val Ile Val Ala Cys Cys Asn			
	180	185	190
Gln Phe Phe Val Leu Asp Val Val Ile Asn Phe Arg Arg Leu Ser Glu			
	195	200	205
Gly Asp Leu Phe Thr Gln Leu Arg Lys Ile Val Lys Met Ala Ser Asn			
	210	215	220
Glu Asp Glu Arg Leu Pro Pro Ile Gly Leu Leu Thr Ser Asp Gly Arg			
	225	230	235
Ser Glu Trp Ala Glu Ala Arg Thr Val Leu Val Lys Asp Ser Thr Asn			
	245	250	255

-continued

Arg	Asp	Ser	Leu	Asp	Met	Ile	Glu	Arg	Cys	Ile	Cys	Leu	Val	Cys	Leu	260	265	270	
Asp	Ala	Pro	Gly	Gly	Val	Glu	Leu	Ser	Asp	Thr	His	Arg	Ala	Leu	Gln	275	280	285	
Leu	Leu	His	Gly	Gly	Gly	Tyr	Ser	Lys	Asn	Gly	Ala	Asn	Arg	Trp	Tyr	290	295	300	
Asp	Lys	Ser	Leu	Gln	Phe	Val	Val	Gly	Arg	Asp	Gly	Thr	Cys	Gly	Val	305	310	315	320
Val	Cys	Glu	His	Ser	Pro	Phe	Asp	Gly	Ile	Val	Leu	Val	Gln	Cys	Thr	325	330	335	
Glu	His	Leu	Leu	Lys	His	Met	Thr	Gln	Ser	Ser	Arg	Lys	Leu	Ile	Arg	340	345	350	
Ala	Asp	Ser	Val	Ser	Glu	Leu	Pro	Ala	Pro	Arg	Arg	Leu	Arg	Trp	Lys	355	360	365	
Cys	Ser	Pro	Glu	Ile	Gln	Gly	His	Leu	Ala	Ser	Ser	Ala	Glu	Lys	Leu	370	375	380	
Gln	Arg	Ile	Val	Lys	Asn	Leu	Asp	Phe	Ile	Val	Tyr	Lys	Phe	Asp	Asn	385	390	395	400
Tyr	Gly	Lys	Thr	Phe	Ile	Lys	Lys	Gln	Lys	Cys	Ser	Pro	Asp	Ala	Phe	405	410	415	
Ile	Gln	Val	Ala	Leu	Gln	Leu	Ala	Phe	Tyr	Arg	Leu	His	Arg	Arg	Leu	420	425	430	
Val	Pro	Thr	Tyr	Glu	Ser	Ala	Ser	Ile	Arg	Arg	Phe	Gln	Glu	Gly	Arg	435	440	445	
Val	Asp	Asn	Ile	Arg	Ser	Ala	Thr	Pro	Glu	Ala	Leu	Ala	Phe	Val	Arg	450	455	460	
Ala	Val	Thr	Asp	His	Lys	Ala	Ala	Val	Pro	Ala	Ser	Glu	Lys	Leu	Leu	465	470	475	480
Leu	Leu	Lys	Asp	Ala	Ile	Arg	Ala	Gln	Thr	Ala	Tyr	Thr	Val	Met	Ala	485	490	495	
Ile	Thr	Gly	Met	Ala	Ile	Asp	Asn	His	Leu	Leu	Ala	Leu	Arg	Glu	Leu	500	505	510	
Ala	Arg	Ala	Met	Cys	Lys	Glu	Leu	Pro	Glu	Met	Phe	Met	Asp	Glu	Thr	515	520	525	
Tyr	Leu	Met	Ser	Asn	Arg	Phe	Val	Leu	Ser	Thr	Ser	Gln	Val	Pro	Thr	530	535	540	
Thr	Thr	Glu	Met	Phe	Cys	Cys	Tyr	Gly	Pro	Val	Val	Pro	Asn	Gly	Tyr	545	550	555	560
Gly	Ala	Cys	Tyr	Asn	Pro	Gln	Pro	Glu	Thr	Ile	Leu	Phe	Cys	Ile	Ser	565	570	575	
Ser	Phe	His	Ser	Cys	Lys	Glu	Thr	Ser	Ser	Ser	Lys	Phe	Ala	Lys	Ala	580	585	590	
Val	Glu	Glu	Ser	Leu	Ile	Asp	Met	Arg	Asp	Leu	Cys	Ser	Leu	Leu	Pro	595	600	605	
Pro	Thr	Glu	Ser	Lys	Pro	Leu	Ala	Thr	Lys	Glu	Lys	Ala	Thr	Arg	Pro	610	615	620	
Ser	Gln	Gly	His	Gln	Pro											625	630		

What is claimed is:

1. An isolated recombinant cell comprising:
 - a) a heterologous nucleic acid encoding an $\alpha 5$ subunit of nicotinic acetylcholine receptor (nAChR); and
 - b) a heterologous nucleic acid encoding a chaperone protein selected from transmembrane inner ear expressed protein TMIE and protein FAM163B.
2. The isolated recombinant cell of claim 1, further comprising:
 - c) a heterologous nucleic acid encoding an $\alpha 2$ subunit of nAChR;
 - d) a heterologous nucleic acid encoding a $\beta 2$ subunit of nAChR;
 - e) a heterologous nucleic acid encoding diamine acetyltransferase 1 (SAT1); and
 - f) a heterologous nucleic acid encoding choline O-acetyltransferase (CHAT),
 wherein, the $\alpha 2$, $\alpha 5$, and $\beta 2$ subunits of nAChR forms a $\alpha 2\alpha 5\beta 2$ nAChR.
3. The isolated recombinant cell of claim 1, wherein the recombinant cell is a mammalian cell.
4. The isolated recombinant cell of claim 3, wherein the mammalian cell is selected from the group consisting of a human embryonic kidney 293T (HEK293T) cell, a HEK293F cell, a HeLa cell, a Chinese hamster ovary (CHO) cell, a NIH 3T3 cell, a MCF-7 cell, a Hep G2 cell, a baby hamster kidney (BHK) cell, and a Cos7 cell.
5. The isolated recombinant cell of claim 1, wherein the $\alpha 2$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 1.
6. The isolated recombinant cell of claim 1, wherein the $\alpha 5$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 2.
7. The isolated recombinant cell of claim 1, wherein the $\beta 2$ subunit of the $\alpha 2\alpha 5\beta 2$ nAChR comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 3.
8. The isolated recombinant cell of claim 1, wherein the TMIE comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 4.
9. The isolated recombinant cell of claim 1, wherein the FAM163B comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 5.
10. The isolated recombinant cell of claim 2, wherein the SAT1 comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 6.
11. The isolated recombinant cell of claim 2, wherein the CHAT comprises an amino acid sequence with at least 95% identity to the amino acid sequence of SEQ ID NO: 7.
12. A method for identifying agonists, antagonists, or positive allosteric modulators of $\alpha 5$ containing nAChR, the method comprising:

- a) contacting the isolated recombinant cell of claim 1 with an agent; and
- b) determining the activity of the $\alpha 5$ containing nAChR of the isolated recombinant cell, wherein the agent is identified as an agonist or positive allosteric modulator (PAM) if the agent enhances the activity of the $\alpha 5$ containing nAChR and the agent is identified as an antagonist if the agent decreases the activity of the $\alpha 5$ containing nAChR as compared to the activity of the $\alpha 5$ containing nAChR when the isolated recombinant cell was not contacted with the agent.

13. A method for identifying agonists, antagonists, or positive allosteric modulators of $\alpha 2\alpha 5\beta 2$ nAChR, the method comprising:

- a) contacting the isolated recombinant cell of claim 2 with an agent; and
- b) determining the activity of the $\alpha 2\alpha 5\beta 2$ nAChR of the isolated recombinant cell, wherein the agent is identified as an agonist or positive allosteric modulator (PAM) if the agent enhances the activity of the $\alpha 2\alpha 5\beta 2$ nAChR and the agent is identified as an antagonist if the agent decreases the activity of the $\alpha 2\alpha 5\beta 2$ nAChR as compared to the activity of the $\alpha 2\alpha 5\beta 2$ nAChR when the isolated recombinant cell was not contacted with the agent.

14. The method of claim 13, wherein step b) comprises determining calcium flux of the isolated recombinant cell, wherein the agent is identified as an agonist if the agent enhances the calcium flux as compared to the calcium flux when the isolated recombinant cell was not contacted with the agent.

15. The method of claim 13, wherein step b) comprises determining calcium flux and nicotine-evoked calcium flux of the isolated recombinant cell, wherein the agent is identified as a PAM if the agent does not enhance calcium flux and enhances the nicotine-evoked calcium flux as compared to the calcium flux and nicotine-evoked calcium flux when the isolated recombinant cell was not contacted with the agent.

16. The method of claim 13, wherein step b) comprises determining nicotine-evoked calcium flux of the isolated recombinant cell, wherein the agent is identified as an antagonist if the agent decreases the nicotine-evoked calcium flux as compared to the nicotine-evoked calcium flux when the isolated recombinant cell was not contacted with the agent.

17. The method of claim 13, wherein the isolated recombinant cell is incubated at about 25° C.-35° C. for about 20-50 hours prior to being contacted with the agent.

18. The method of claim 13, wherein the agent is a small molecule or peptide.

19. A kit comprising (i) the isolated recombinant cell of claim 1, and (ii) instructions for use.

* * * * *