



US012410253B2

(12) **United States Patent**
Burman et al.

(10) **Patent No.:** **US 12,410,253 B2**

(45) **Date of Patent:** **Sep. 9, 2025**

(54) **COMPOSITIONS COMPRISING ANTI-NRP2 ANTIBODIES**

(71) Applicant: **aTyr Pharma, Inc.**, San Diego, CA (US)

(72) Inventors: **Luke Burman**, San Diego, CA (US); **Yeeting Chong**, San Diego, CA (US); **Leslie Ann Greene**, San Diego, CA (US); **David King**, Encinitas, CA (US); **Zhiwen Xu**, La Jolla, CA (US); **Ryan Andrew Adams**, San Diego, CA (US)

(73) Assignee: **aTyr Pharma, Inc.**, San Diego, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **18/477,113**

(22) Filed: **Sep. 28, 2023**

(65) **Prior Publication Data**

US 2024/0092917 A1 Mar. 21, 2024

Related U.S. Application Data

(62) Division of application No. 17/062,100, filed on Oct. 2, 2020, now Pat. No. 11,807,687.

(Continued)

(51) **Int. Cl.**

A61K 39/395 (2006.01)

A61P 35/04 (2006.01)

(Continued)

(52) **U.S. Cl.**

CPC **C07K 16/2863** (2013.01); **A61K 39/3955** (2013.01); **A61P 35/04** (2018.01);

(Continued)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,428,965 B1 8/2002 Ginty et al.
7,736,655 B2 6/2010 Klagsbrun et al.
(Continued)

FOREIGN PATENT DOCUMENTS

CN 1726226 A 1/2006
CN 101326196 A 12/2008
(Continued)

OTHER PUBLICATIONS

Herold et al., Determinants of the assembly and function of antibody variable domains, Scientific Reports, 7:12276, DOI: 10.1038/s41598-017-12519-9, 17 pages, Sep. 2017.*

(Continued)

Primary Examiner — Claire Kaufman

(74) *Attorney, Agent, or Firm* — COOLEY LLP


(57) **ABSTRACT**

Provided are affinity matured and humanized antibodies and antigen-binding fragments thereof that specifically bind to human neuropilin-2 (NRP2) polypeptides, including those that modulate binding interactions between human NRP2 and at least one NRP2 ligand, and which thereby modulate subsequent NRP2-mediated downstream signaling events, including related therapeutic compositions and methods for modulating NRP2 activity and treating diseases such as NRP2-associated diseases.

15 Claims, 23 Drawing Sheets

Specification includes a Sequence Listing.

Inhibition of Sema3F Signaling



Related U.S. Application Data

(60) Provisional application No. 63/024,960, filed on May 14, 2020, provisional application No. 62/910,042, filed on Oct. 3, 2019.

(51) **Int. Cl.**

C07K 16/28 (2006.01)

A61K 39/00 (2006.01)

(52) **U.S. Cl.**

CPC . **A61K 39/001103** (2018.08); **A61K 2039/505** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/33** (2013.01); **C07K 2317/52** (2013.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

8,318,163 B2 11/2012 Appleton et al.
8,466,264 B2 6/2013 Appleton et al.
8,529,905 B2 9/2013 Klagsbrun et al.
8,648,173 B2 2/2014 Wu et al.
8,722,043 B2 5/2014 Borg et al.
8,920,805 B2 12/2014 Wu et al.
8,956,622 B2 2/2015 Bagnard et al.
11,505,610 B2 11/2022 Burman et al.
11,767,520 B2 9/2023 Mendlein et al.
11,807,687 B2 11/2023 Burman et al.
12,065,495 B2 8/2024 Burman et al.
2002/0127227 A1 9/2002 Holmes et al.
2003/0045691 A1 3/2003 Ono et al.
2003/0088075 A1 5/2003 Shitara et al.
2003/0113324 A1 6/2003 Alitalo et al.
2004/0005647 A1 1/2004 Denardo et al.
2004/0028685 A1 2/2004 Kinch et al.
2004/0038339 A1 2/2004 Kufer et al.
2004/0214766 A1 10/2004 Alitalo et al.
2005/0064518 A1 3/2005 Albone et al.
2006/0280747 A1 12/2006 Fuh et al.
2007/0189962 A1 8/2007 Pastan et al.
2009/0196850 A1 8/2009 Romagne et al.
2010/0130904 A1 5/2010 Kirber
2010/0172921 A1 7/2010 Wu et al.
2010/0310573 A1 12/2010 Nakagawa et al.
2011/0064670 A1 3/2011 Gogineni et al.
2011/0091377 A1 4/2011 Alani et al.
2011/0091384 A1 4/2011 Alani et al.
2011/0243932 A1 10/2011 Barrett et al.
2012/0027779 A1 2/2012 Borg et al.
2012/0282637 A1 11/2012 Huber et al.
2013/0131319 A1 5/2013 Igawa et al.
2013/0266517 A1 10/2013 Saldanha et al.
2014/0193420 A1 7/2014 Aburatani et al.
2014/0234312 A1 8/2014 Wu et al.
2015/0030600 A1 1/2015 Marks et al.
2015/0218267 A1 8/2015 Brodeur et al.
2015/0307589 A1 10/2015 Bagnard et al.
2016/0000873 A1 1/2016 Long et al.
2016/0017041 A1 1/2016 Violette et al.
2016/0046711 A1 2/2016 Bialucha et al.
2016/0108113 A1 4/2016 Ayalon et al.
2016/0108131 A1 4/2016 Berne et al.
2016/0312300 A1 10/2016 Alani et al.
2017/0100456 A1 4/2017 Briscoe et al.
2018/0193454 A1 7/2018 Rothstein et al.
2018/0282402 A1 10/2018 Adams et al.
2019/0309076 A1 10/2019 Burman et al.
2021/0163606 A1 6/2021 Burman et al.
2023/0312730 A1 10/2023 Burman et al.
2024/0417476 A1 12/2024 Burman et al.
2025/0074990 A1 3/2025 Burman et al.

FOREIGN PATENT DOCUMENTS

CN 101743253 A 6/2010
CN 101754771 A 6/2010

CN 102597776 A 7/2012
CN 102667484 A 9/2012
CN 103459422 A 12/2013
CN 104203268 A 12/2014
CN 105407891 A 3/2016
EP 1968565 B1 11/2015
EP 3024460 B1 7/2020
JP 2010527350 A 8/2010
JP 2012500832 A 1/2012
JP 2013507641 A 3/2013
WO WO-9929729 A2 6/1999
WO WO-9929858 A1 6/1999
WO WO-9930157 A2 6/1999
WO WO-2004035537 A2 4/2004
WO WO-2007020075 A1 2/2007
WO WO-2007056470 A2 5/2007
WO WO-2007095749 A1 8/2007
WO WO-2008143665 A1 11/2008
WO WO-2008143666 A2 11/2008
WO WO-2009043933 A1 4/2009
WO WO-2010065940 A1 6/2010
WO WO-2010102251 A2 9/2010
WO WO-2010119704 A1 10/2010
WO WO-2011032013 A1 3/2011
WO WO-2011047033 A2 4/2011
WO WO-2012069557 A1 5/2012
WO WO-2013106765 A1 7/2013
WO WO-2013116742 A1 8/2013
WO WO-2014105810 A1 7/2014
WO WO-2015136471 A1 9/2015
WO WO-2017205377 A2 11/2017
WO WO-2018102589 A2 6/2018
WO WO-2018195302 A1 10/2018
WO WO-2019075472 A1 4/2019
WO WO-2019086351 A1 5/2019
WO WO-2019147973 A1 8/2019
WO WO-2019195770 A1 10/2019
WO WO-2020200196 A1 10/2020
WO WO-2021058548 A1 4/2021
WO WO-2021067761 A1 4/2021
WO WO-2021202590 A1 10/2021
WO WO-2023063208 A1 4/2023
WO WO-2023076998 A1 5/2023
WO WO-2023245117 A2 12/2023

OTHER PUBLICATIONS

Sela-Culang et al., The structural basis of antibody-antigen recognition, *Front. Immunol.* 4:302, 13 pages, Oct. 2013.*
Qiu et al., Engineering an anti-CD52 antibody for enhanced deamidation stability, *mAbs*, 11(7):1266-1275, 2019.*
Alto, L. T. et al., "Semaphorins and their signaling mechanisms," *Methods. Mol. Biol.* 1493:1-25 (2017).
Anonymous, "aTyr Pharma Presents Preclinical Research Characterizing Effects of ATYR2810 in Highly Aggressive Tumor Subtypes at the 2022 AACR Annual Meeting : BioSpace," Apr. 11, 2022, 3 pages, Retrieved from the Internet: URL:<https://www.biospace.com/article/releases/atyr-pharma-presents-preclinical-research-characterizing-effects-of-tyr2810-in-highly-aggressive-tumor-subtypes-at-the-2022-aacr-annual-meeting/>.
Appleton, B. A. et al., "Structural studies of neuropilin/antibody complexes provide insights into semaphorin and VEGF binding," *The EMBO Journal*, 26(23):4902-4912 (Nov. 2007).
Aung, N. Y. et al., "Specific neuropilins expression in alveolar macrophages among tissue-specific macrophages," *PLoS One* 11(2): pp. e0147358 (2016).
Bannerman, P. et al., "Peripheral nerve regeneration is delayed in neuropilin 2-deficient mice," *J. Neurosci. Res.* 86(14):3163-3169 (2008).
Bielenberg, D. R. et al., "Increased smooth muscle contractility in mice deficient for neuropilin 2," *Am. J. Path.* 181(2):548-559 (2012).
Bielenberg, D. R. et al., "Neuropilins in neoplasms: Expression, regulation, and function," *Exp. Cell. Res.* 312:584-593 (2006).
Binch, A. L. A. et al., "Class 3 semaphorins expression and association with innervation and angiogenesis within the degenerate human intervertebral disc," *Oncotarget* 6(21):18338-18354 (2015).

(56)

References Cited**OTHER PUBLICATIONS**

- Bogan and Thorn, "Anatomy of hot spots in protein interfaces", *J Mol Biol.* Jul. 3, 1998; 280(1): 1-9.
- Boscolo, E. et al., "AKT hyper-phosphorylation associated with PI3K mutations in lymphatic endothelial cells from a patient with lymphatic malformation," *Angiogenesis* 18(2):151-162 (2015).
- Cao, Y. et al., "Neuropilin-2 promotes extravasation and metastasis by interacting with endothelial 5 integrin," *Cancer Res.* 73(14):4579-4590 (2013).
- Casset, F., et al., "A Peptide Mimetic of an Anti-cd4 Monoclonal Antibody by Rational Design," *Biochemical and Biophysical Research Communications*, Jul. 2003, vol. 307 (1), pp. 198-205.
- Caunt, M. et al., "Blocking neuropilin-2 function inhibits tumor cell metastasis," *Cancer Cell*, vol. 13, No. 4, Apr. 2008, pp. 331-342.
- Chen, L. et al., "131I-labeled monoclonal antibody targeting neuropilin receptor type-2 for tumor SPECT imaging," *Int. J. Oncol.* 50:649-659 (2017).
- Chen, Y., et al., "Selection and Analysis of an Optimized Anti-vegfr Antibody: Crystal Structure of an Affinity-matured Fab in Complex With Antigen," *Journal of Molecular Biology*, Nov. 1999, vol. 293 (4), pp. 865-881.
- Chong, Y. E. et al., "ATYR2810, a Neuropilin-2 antibody, selectively blocks the NRP2/VEGFR signaling axis and sensitizes aggressive cancers to chemotherapy," Jan. 2021, Retrieved from the Internet: URL: https://atyrpharma.com/wp-content/uploads/2021/05/2021-Keystone_CSC_poster_vS.pdf [retrieved on Dec. 6, 2021], 1 page.
- Corliss, B. A. et al., "Macrophages: An inflammatory link between angiogenesis and lymphangiogenesis," *Microcirculation* 23(2):95-121 (2016).
- De Pascalis et al., "Grafting of 'Abbreviated' Complementarity-Determining Regions Containing Specificity-Determining Residues Essential for Ligand Contact to Engineer a Less Immunogenic Humanized Monoclonal Antibody," *The Journal of Immunology*, Sep. 2002, vol. 169, No. 6, pp. 3076-3084.
- Dong, X. et al., "Elevated expression of neuropilin-2 associated with unfavorable prognosis in hepatocellular carcinoma," *OncoTargets and Therapy* 10:3827-3833 (2017).
- Drabkin, H. A. et al., "A triad of NRP2, DLX and p53 proteins in lung cancer metastasis," *Oncotarget* 8(57):96464-96465 (2017).
- Dutta, S. et al., "Neuropilin-2 regulates endosomal maturation and EGFR trafficking to support cancer cell pathobiology," *Cancer Res.* 76(2):418-428 (2016).
- Elaimy, A. L. et al., "VEGF-neuropilin-2 signaling promotes stem-like traits in breast cancer cells by TAZ-mediated repression of the Rac GAP 2-chimaerin," *Sci. Signal* 11, pp. eaao6897 (2018).
- Elpek, G. O., "Neuropilins and liver," *World J. Gastroent.* 21(23):7065-7073 (2015).
- Extended European Search Report for European Application No. 19781711.7, mailed Dec. 16, 2021, 11 pages.
- Fassold, A. et al., "Soluble neuropilin-2, a nerve repellent receptor, is increased in rheumatoid arthritis synovium and aggravates sympathetic fiber repulsion and arthritis," *Arthritis & Rheum.* 60(10):2892-2901 (2009).
- Forsyth et al., "Deep mutational scanning of an antibody against epidermal growth factor receptor using mammalian cell display and massively parallel pyrosequencing," *mAbs.* Jul.-Aug. 2013; 5(4): 523-32. Epub May 29, 2013.
- Fujita, H. et al., "Expressions of neuropilin-1, neuropilin-2 and semaphorin 3A Mrna in the rat brain after middle cerebral artery occlusion," *Brain Research* 914(1-2):1-14 (2001).
- Fusco, F. et al., "Progressive bladder remodeling due to bladder outlet obstruction: a systematic review of morphological and molecular evidences in humans," *BMC Urology* 18:1-11 (2018).
- Gemmill, R. M. et al., "The neuropilin 2 isoform NRP2b uniquely supports TGF-mediated progression in lung cancer," *Sci. Signal.* 10, pp. eaag0528 (2017).
- Goel, H. L. et al., "Enhancing integrin function by VEGF/neuropilin signaling," *Cell Adhesion & Migration*, 6(6):554-560 (2012).
- Goel, H. L. et al., "GLI1 regulates a novel neuropilin-2/61 integrin based autocrine pathway that contributes to breast cancer initiation," *EMBO Mol. Med.* 5:488-508 (2013).
- Goel, H. L. et al., "VEGF/Neuropilin-2 regulation of Bmi-1 and consequent repression of IGF-1R define a novel mechanism of aggressive prostate cancer," *Cancer Discov.* 2(10):906-921 (2012).
- Gogineni, A. et al., "Inhibition of VEGF-C modulates distal lymphatic remodeling and secondary metastasis," *PloS One* 8(7):pp. e68755 (2013).
- Grandclement, C. et al., "Neuropilin-2 Expression Promotes TGF-1-Mediated Epithelial to Mesenchymal Transition in Colorectal Cancer Cells," *PLoS One* 6(7):pp. e20444 (2011).
- Guan, J., et al., "The U4 Antibody Epitope on Human Papillomavirus 16 Identified by Cryo-electron Microscopy" pp. 12108-12117. *Journal of Virology*, vol. 89, No. 23, Sep. 23, 2015, abstract; Genbank supplement, pp. 1-3; DOI: 10.1128/JVI.02020-15.
- Guo et al., "Protein tolerance to random amino acid change," *Proc Natl Acad Sci USA.* Jun. 22, 2004; 101(25): 9205-10. Epub Jun. 14, 2004.
- Guo, H-F. et al., "Neuropilin functions as an essential cell surface receptor," *J. Biol. Chem.* 290(49):29120-29126 (2015).
- Harding, J. et al., "Lymphangiogenesis is induced by mycobacterial granulomas via vascular endothelial growth factor receptor-3 and supports systemic T-Cell responses against mycobacterial antigen," *Am. J. Path.* 185(2):432-445 (2015).
- Henno, A. et al., "Altered expression of angiogenesis and lymphangiogenesis markers in the uninvolved skin of plaque-type psoriasis," *Br. J. Derm.*, 160:581-590 (2009).
- Hey-Cunningham, A. J. et al., "Dysregulation of vascular endothelial growth factors and their neuropilin receptors in the eutopic endometrium of women with endometriosis," *Repro. Sci.*, Nov. 2013, vol. 20, No. 11, pp. 1382-1389.
- Huang, Z-L. et al., "Establishment of a bead-based duplex assay for the simultaneous quantitative detection of Neuropilin-1 and Neuropilin-2 using xMAP technology and its clinical application," *J. Clin. Lab. Anal.*, 2019, 6 pages.
- Immormino, R. M. et al., "Neuropilin-2 regulates airway inflammatory responses to inhaled lipopolysaccharide," *Am J Physiol Lung Cell Mol Physiol.*, 315(2):L202-L211 (Aug. 2018).
- International Search Report and Written Opinion for International Application No. PCT/US2019/026128, mailed Aug. 12, 2019, 12 pages.
- International Search Report and Written Opinion for International Application No. PCT/US2020/054017, mailed Feb. 25, 2021, 13 pages.
- International Search Report and Written Opinion for International Application No. PCT/US2022/078780 dated Mar. 15, 2023, 18 pages.
- International Search Report and Written Opinion for PCT Application No. PCT/US2023/063208 dated Jul. 18, 2023, 11 pages.
- International Search Report and Written Opinion for PCT/US2021/021148 Jul. 28, 2021, 12 pages.
- Karkkainen, M. J. et al., "A model for gene therapy of human hereditary lymphedema," *PNAS* 98(22):12677-12682 (2001).
- Karpanen, T. et al., "Functional interaction of VEGF-C and VEGF-D with neuropilin receptors," *The FASEB J.* 20(9):1462-1472 (2007).
- Li, J., et al., "Repertoire diversification in mice with an IgH-locus targeted transgene for the 1-3 rearranged VH domain of a physiologically selected anti-ssDNA antibody," *Molecular Immunology*, Mar. 5, 2005, vol. 42, No. 12, pp. 1475-1484.
- Lippow, et al., "Computational design of antibody-affinity improvement beyond in vivo maturation," *Nat Biotechnol.* (Oct. 2007); 25(10): 1171-1176. Epub Sep. 23, 2007.
- Liu, B. et al., "Identification of prognostic biomarkers by combined mRNA and miRNA expression microarray analysis in pancreatic cancer," *Trans. Onc.* 11(3):700-714 (2018).
- Lu, Y. et al., "Identification of circulating neuropilin-1 and dose-dependent elevation following anti-neuropilin-1 antibody administration," *mAbs*, 1(4):364-369 (2009).
- MacCallum, R.M., et al., "Antibody-Antigen Interactions: Contact Analysis and Binding Site Topography," *Journal of Molecular Biology*, Oct. 11, 1996, vol. 262, No. 5, pp. 732-745.

(56)

References Cited

OTHER PUBLICATIONS

- Makinen, T. et al., "Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble VEGF receptor-3," *Nature Med.* 7(2):199-205 (2001).
- Matthews, B. W., "Structural and genetic analysis of protein stability," *Annu. Rev. Biochem.*, 62:139-160 (1993).
- Mendes-Da-Cruz, D. A. et al., "Semaphorin 3F and Neuropilin-2 Control the Migration of Human T-Cell Precursors," *PloS One* 9(7), 7 pages (2014).
- Mercurio, A. M., "VEGF/Neuropilin signaling in cancer stem cells," *Int. J. Mol. Sci.*, 20(3):490 (2019).
- Meyer, L. A. T. et al., "Current drug design to target the semaphorin/neuropilin/plexin complexes," *Cell Adhes. & Migration* 10(6):700-708 (2016).
- Miaskowski, C. et al., "Lymphatic and angiogenic candidate genes predict the development of secondary lymphedema following breast cancer surgery," *PloS One* 8(4), e60164, 20 pages (2013).
- Moran, T. P. et al., "Neuropilin-2 is a negative regulator of allergic airway inflammation," *J. Allergy Clin. Immunol.*, AB180 Abstracts, 568 (Feb. 2018).
- Mucka, P. et al., "Inflammation and lymphedema are exacerbated and prolonged by neuropilin 2 deficiency," *Am. J. Path.* 186(11):2803-2812 (2016).
- Nakayama, H. et al., "Regulation of mTOR signalling by semaphorin 3F-neuropilin 2 interactions in vitro and in vivo," *Scientific Reports* 5, 11789, 14 pages (2015).
- Napolitano, V. et al., "Neuropilins controlling cancer therapy responsiveness," *Int. J. Mol. Sci.*, 20(8):2049 (2019).
- Nasarre, P. et al., "Semaphorin SEMA3F and VEGF have opposing effects on cell attachment and spreading," *Neoplasia* 5(1):83-92 (2003).
- Nasarre, P. et al., "The emerging role of class-3 semaphorins and their neuropilin receptors in oncology," *Oncotargets and Therapy* 7:1663-1687 (2014).
- Neufeld, G. et al., "Semaphorins in angiogenesis and tumor progression," *Cold Spring Harbor Perspect. in Med.* 2, 13 pages, a006718 (2012).
- NG and Henikoff, "Predicting the Effects of Amino Acid Substitutions on Protein Function," *Annu. Rev. Genomics Hum. Genet.* (2006); 7: 61-80.
- NG, T. et al., "Neuropilin 2 signaling is involved in cell positioning of adult-born neurons through glycogen synthase kinase-3," *J. Biol. Chem.* 291(48):25088-25095 (2016).
- Niland, S. et al., "Neuropilins in the context of tumor vasculature," *Int. J. Mol. Sci.*, 20(3):639 (2019).
- Ogata, F. et al., "Excess lymphangiogenesis cooperatively induced by macrophages and CD4+ T cells drives the pathogenesis of lymphedema," *J. Invest. Derm.* 136:706-714 (2016).
- Papadopolou, K. et al., "Genotyping and mRNA profiling reveal actionable molecular targets in biliary tract cancers," *Am. J. Cancer Res.* 8(1):2-15 (2018).
- Parker, M. W. et al., "Furin processing of semaphorin 3F determines its anti-angiogenic activity by regulating direct binding and competition for neuropilin," *Biochemistry* 49(19):4068-4075 (2010).
- Parker, M. W. et al., "Microplate-based screening for small molecule inhibitors of Neuropilin-2/VEGF-C interactions," *Anal. Biochem.* 453:4-6 (May 2014).
- Parker, M. W. et al., "Structural basis for VEGF-C binding to neuropilin-2 and sequestration by a soluble splice form," *Structure*, 23(4):677-687 (2015).
- Partial European Search Report for European Application No. EP20870841.2, dated Oct. 5, 2023, 15 pages.
- Pavlovic, H., "Soluble VEGFR2: an anti-lymphatic variant of VEGF receptors," *Ann. N.Y. Acad. Sci.* 1207(Suppl 1):E7-E15 (2010).
- Pellet-Many, C. et al., "Neuropilins: structure, function and role in disease," *Biochemical Journal*, vol. 411, No. 2, Apr. 2008, pp. 211-226.
- Prud'homme, G. J. et al., "Neuropilins are multifunctional coreceptors involved in tumor initiation, growth, metastasis and immunity," *Oncotarget* 3(9):921-939 (2012).
- Raaben, M. et al., "NRP2 and CD63 are host factors for Lujo virus cell entry," *Cell Host Microbe* 22(5):688-695.e5 (2017).
- Rey-Gallardo, A. et al., "Polysialylated neuropilin-2 enhances human dendritic cell migration through the basic C-terminal region of CCL21," *Glycobiology* 20(9):1139-1146 (2010).
- Rizzolio, S. et al., "Downregulating neuropilin-2 triggers a novel mechanism enabling EGFR-dependent resistance to oncogene-targeted therapies," *Cancer. Res.* 78(4):1058-1068 (2017).
- Rollenhagen, M. et al., "Polysialic acid on neuropilin-2 is exclusively synthesized by the polysialyltransferase ST8SialV and attached to mucin-type O-glycans located between the b2 and c domain," *The J. Biol. Chem.* 288(32):22880-22892 (2013).
- Roy, S. et al., "Multifaceted role of neuropilins in the immune system: Potential targets for immunotherapy," *Front. Immunol.*, Oct. 10, 2017, vol. 8, p. 1228.
- Rudikoff, S., et al., "Single amino acid substitution altering antigen-binding specificity," *Proceedings of the National Academy of Sciences*, 1982; 79(6): 1979-1983.
- Saban, M. R. et al., "VEGF signaling mediates bladder neuroplasticity and inflammation in response to BCG," *BMC Physiology*, 11(16):1-20 (2011).
- Saban, R., "Angiogenic factors, bladder neuroplasticity and interstitial cystitis-new pathobiological insights," *Trans. Androl. Urol.* 4(5):555-562 (2015).
- Sainz-Jaspeado, M. et al., "Cytokines regulating lymphangiogenesis," *Curr. Opin. Immunol.* 53:58-63 (2018).
- Samuel, S., et al., "Neuropilin-2 Mediated [beta]-Catenin Signaling and Survival in Human Gastro-Intestinal Cancer Cell Lines," *PLOS One*, vol. 6, No. 10, Oct. 20, 2011, 11 pages, e23208, DOI: 10.1371/journal.pone.0023208.
- Sarabipour, S. et al., "VEGF-A121a binding to Neuropilins—A concept revisited," *Cell Adhes. & Migration* 12(3):204-214 (2018).
- Schellenburg, S. et al., "Role of neuropilin-2 in the immune system," *Molecular Immunology*, 90:239-244 (2017).
- Schwarz, Q. et al., "Neuropilin, you gotta let me know, Should I stay or should I go?" *Cell Adhesion & Migration* 4(1):61-66 (2010).
- Simeon et al., "In vitro-engineered non-antibody protein therapeutics," *Protein Cell* 2018, 9(1):3-14, doi: 10.1007/s13238-017-0386-6.
- Stanton, M. J. et al., "Angiogenic growth factor axis in autophagy regulation," *Autophagy* 9(5):789-790 (2013).
- Stine, M. J. et al., "Integration of genotypic and phenotypic screening reveals molecular mediators of melanoma-stromal interaction," *Cancer. Res.* 71(7):2433-2444 (Apr. 2011).
- Sulpice, E. et al., "Neuropilin-1 and neuropilin-2 act as coreceptors, potentiating proangiogenic activity," *Blood* 111(4):2036-2045 (2008).
- Tu, D.-G. et al., "Promotion of metastasis of thyroid cancer cells via NRP-2 mediated induction," *Oncol. Lett.* 12:4224-4230 (2016).
- Uni Prot Database, P01857, IGHG1_Human, Retrieved online: <url: <https://www.uniprot.org/uniprotkb/P01857/entry#structure>, Retrieved on Jan. 12, 2023, pp. 1, 2 and 11, 2023.
- UniProtKB Accession No. A0A1C0YHR0, Uncharacterized protein, Nov. 2, 2016 [online] 1-3; [Retrieved on May 15, 2023]. Retrieved from the internet: URL: <https://www.uniprot.org/uniprot/A0A1C0YHR0> Entire document; amino acid sequence residues 21-31, 4 pages.
- Vaahomeri, K. et al., "Lymphangiogenesis guidance by paracrine and pericellular factors," *Genes & Dev* 31:1615-1634 (2017).
- Vasquez, E. et al., "Deletion of neuropilin 2 enhances detrusor contractility following bladder outlet obstruction," *JCI Insight* 2(3):e90617, 10 pages (2017).
- Verlinden, L. et al., "Nrp2 deficiency leads to trabecular bone loss and is accompanied by enhanced osteoclast and reduced osteoblast numbers," *Bone* 55:465-475 (2013).
- Wang, J. et al., "NRP-2 in tumor lymphangiogenesis and lymphatic metastasis," *Cancer. Lett.* 418:176-184 (2018).
- Werneburg, S. et al., "Polysialylation and lipopolysaccharide-induced shedding of E-selectin ligand-1 and neuropilin-2 by microglia and THP1 macrophages," *GLIA* 64:1314-1330 (2016).

(56)

References Cited

OTHER PUBLICATIONS

Wild, J. R. L. et al., "Neuropilins: expression and roles in the epithelium," *Int. J. Exp. Path.* 93:81-103 (2012).

Wittmann, P. et al., "Neuropilin-2 induced by transforming growth factor-augments migration of hepatocellular carcinoma cells, *BMC Cancer*, 2015, vol. 15, No. 1, pp. 1-8.

Wu, H., et al., "Humanization of a murine monoclonal antibody by simultaneous optimization of framework and CDR residues", *Journal of Molecular Biology* (1999); 294(1): 151-162.

Xu, Y. et al., "Neuropilin-2 mediates VEGF-C-induced lymphatic sprouting together with VEGFR3," *J. Cell. Biol.* 188(1):115-130 (2010).

Yang, F. et al., "Understanding lymphangiogenesis in knockout models, the cornea, and ocular diseases for the development of therapeutic interventions," *Surv. Ophthalmol.*, 61(3):272-296 (May-Jun. 2016).

Yang, Y. et al., "Preparation, Purification, and Identification of a Monoclonal Antibody Against NRP2 b1b2 Domain," *Monoclonal Antibodies in Immunodiagnosis and Immunotherapy*, vol. 34, No. 5, Oct. 2015, pp. 354-359.

Yasuoka, H. et al., "Neuropilin-2 expression in breast cancer: correlation with lymph node metastasis, poor prognosis, and regulation of CXCR4 expression," *BMC Cancer*, Dec. 2009, vol. 9, pp. 1-7.

Yelland, T. et al., "Crystal structure of the neuropilin-1 MAM domain: completing the neuropilin-1 ectodomain picture," *Structure* 24(11):2008-2015 (2016).

Yuan, L. et al., "Abnormal lymphatic vessel development in neuropilin 2 mutant mice," *Development* 129:4797-4806 (2002).

Zhu, H. et al., "VEGF-C inhibition reverses resistance of bladder cancer cells to cisplatin via upregulating maspin," *Mol. Med. Reports*, 2015, vol. 12, pp. 3163-3169.

Extended European Search Report for European Application No. EP20870841.2, mailed on Jan. 11, 2024, 12 pages.

International Preliminary Report on Patentability for International Application No. PCT/US2022/078780, mailed Apr. 30, 2024, 13 pages.

International Preliminary Report on Patentability for International Application No. PCT/US2023/063208, mailed Sep. 6, 2024 7 pages.

International Preliminary Report on Patentability for International Application No. PCT/US2023/068511, mailed Dec. 26, 2024, 8 pages.



International Search Report and Written Opinion for International Application No. PCT/US2023/068511, mailed on Jan. 10, 2024, 12 pages.

Lambrinos, G. et al., "Neuropilin 2 is a novel regulator of distal colon contractility," *Am. J. Pathol.*, Nov. 2022, vol. 192, No. 11, pp. 1592-1603.

Pan et al., "Molecular mechanisms for tumor resistance to chemotherapy," *Clinical and Experimental Pharmacology and Physiology*, Aug. 2016, vol. 43, No. 8, pp. 723-737.

Salahuddin et al., "Pulmonary Eosinophilia," NIH Bookshelf, last updated May 22, 2023, in: StatPearls, Treasure Island (FL): StatPearls Publishing, 8 pages, at url: <https://www.ncbi.nlm.nih.gov/books/NBK470600/>.

* cited by examiner

*FIG. 1A**FIG. 1B*




FIG. 2A




FIG. 2B




FIG. 3







FIG. 4




FIGs. 5A-5B




FIGs. 6A-6B




FIGs. 7A-7B





FIGs. 8A-8B



FIGs. 9A-9B



FIGs. 10A-10B



FIGs. 11A-11B






FIG. 12

13A TNBC MDA-MB-231 NPR2-high cells



13B TNBC BT549 NPR2-high cells



FIGs. 13A-13B




FIG. 14




FIG. 15




FIG. 16




FIG. 17





FIG. 18


19A

Tumor Growth Curves



19B


Terminal Tumor Weights



FIGs. 19A-19B


20A

Number of lung metastases



20B

Serum AB Conc. vs # Lung Metastases



FIGs. 20A-20B




FIG. 21






FIG. 22



FIGs. 23A-23B

1

**COMPOSITIONS COMPRISING ANTI-NRP2
ANTIBODIES****CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a divisional application of U.S. application Ser. No. 17/062,100, filed Oct. 2, 2020, now U.S. Pat. No. 11,807,687, issued Nov. 7, 2023, which claims the benefit under 35 U.S.C. § 119 (e) to U.S. Provisional Application No. 63/024,960, filed May 14, 2020; and U.S. Provisional Application No. 62/910,042, filed Oct. 3, 2019, each of which is incorporated by reference in its entirety.

**STATEMENT REGARDING THE SEQUENCE
LISTING**

The Sequence Listing XML associated with this application is provided in XML file format and is hereby incorporated by reference into the specification. The name of the XML file containing the Sequence Listing XML is ATYR_136_03US_ST26.xml. The XML file is about 211,947 bytes, was created on Mar. 12, 2025, and is being submitted electronically via USPTO Patent Center.

BACKGROUND**Technical Field**

Embodiments of the present disclosure relate to affinity matured and humanized antibodies and antigen-binding fragments thereof that specifically bind to human neuropilin-2 (NRP2) polypeptides, including those that modulate binding interactions between human NRP2 and at least one NRP2 ligand, and which thereby modulate subsequent NRP2-mediated downstream signaling events, including related therapeutic compositions and methods for modulating NRP2 activity and treating diseases such as NRP2-associated diseases.

Description of the Related Art

Recent research developments suggest that tRNA synthetases play important roles in cellular responses beyond their well-characterized role in protein synthesis. In particular, there is a growing recognition that tRNA synthetases participate in a range of previously-unrecognized roles in responding to cellular stress and tissue homeostasis, in both intracellular and extracellular environments.

Significant progress has been made in elucidating the role of extracellular HARS derived proteins, including the identification of a putative cellular receptor, neuropilin-2 (NRP2 or NRP-2). Interactions of HARS with NRP2 appear to be mediated by the N-terminal region of HARS, and can lead to important changes in the cellular function of NRP2.

Accordingly, the current discovery of this new pathway of regulation represents a previously unknown mechanism, which acts as a central regulator of cellular processes, including, for example, axonal guidance, endocytosis, cell migration, proliferation, survival, apoptosis, lymphangiogenesis, cellular differentiation, and cell attachment with direct relevance to cancer initiation, growth, metastasis, and chemoresistance, as well as muscular, vascular, neuronal, bone, and immune homeostasis. The deregulation of any of these processes may lead to a spectrum of diseases, which may be addressed by the development of anti-NRP2 anti-

2

bodies that selectively target the neuropilin-2 axis. The present disclosure provides such antibodies and related embodiments.

BRIEF SUMMARY

Embodiments of the present disclosure include antibodies or antigen-binding fragments thereof that specifically bind to a human neuropilin-2 (NRP2) polypeptide (anti-NRP2 antibody).

Certain embodiments include a therapeutic composition, comprising at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin-2 (NRP2) polypeptide (anti-NRP2 antibody), wherein the at least one antibody or antigen-binding fragment thereof comprises: a heavy chain variable region (V_H) sequence that comprises complementary determining region V_H CDR1, V_H CDR2, and V_H CDR3 sequences selected from Table A1 or Table A3 and variants thereof which specifically bind to the human NRP2 polypeptide; and a light chain variable region (V_L) sequence that comprises complementary determining region V_L CDR1, V_L CDR2, and V_L CDR3 sequences selected from Table A1 or Table A3 and variants thereof which specifically bind to the human NRP2 polypeptide.

In some embodiments:

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 1-3, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 4-6, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 7-9, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 10-12, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 13-15, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 16-18, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 19-21, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 22-24, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 25-27, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 28-30, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 34-36, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 34-39, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 40-42, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 57-59, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 60-62, respectively, including variants thereof; or

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 63-65, respectively, and the

3

V_LCDR1, V_LCDR2, and V_LCDR3 sequences comprise SEQ ID NOs: 66-68, respectively, including variants thereof.

In some embodiments, the V_H sequence is at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from Table A2, optionally wherein the V_H sequence has 1, 2, 3, 4, or 5 alterations in the framework regions. In some embodiments, the V_L sequence is at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from Table A2, optionally wherein the V_L sequence has 1, 2, 3, 4, or 5 alterations in the framework regions.

In some embodiments:

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 43, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 44;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 45, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 46;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 47, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 48;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 49, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 50;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 51, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 52;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 53, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 54;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 55, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 56;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 69, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 70;

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 71, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 72; or

the V_H sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 73, and the V_L sequence comprises a sequence at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to SEQ ID NO: 74.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a full-length human NRP2 polypeptide or a human NRP2 polypeptide selected from Table N1, optionally with an affinity of about 10 pM to about 500 pM or to about 50 nM, or about,

4

at least about, or no more than about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900 pM, 1 nM, 10 nM, 25 nM, or 50 nM, or optionally with an affinity that ranges from about 10 pM to about 500 pM, about 10 pM to about 400 pM, about 10 pM to about 300 pM, about 10 pM to about 200 pM, about 10 pM to about 100 pM, about 10 pM to about 50 pM, or about 20 pM to about 500 pM, about 20 pM to about 400 pM, about 20 pM to about 300 pM, about 20 pM to about 200 pM, about 20 pM to about 100 pM, about 20 pM to about 50 pM, or about 30 pM to about 500 pM, about 30 pM to about 400 pM, about 30 pM to about 300 pM, about 30 pM to about 200 pM, about 30 pM to about 100 pM, about 30 pM to about 50 pM, or about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 1 nM to about 5 nM, about 5 nM to about 10 nM, about 10 nM to 25 nM, or about 25 nM to about 50 nM, optionally wherein the at least one antibody or antigen-binding fragment thereof specifically binds to the human NRP2 polypeptide in its native form but does not substantially bind to the human NRP2 polypeptide in its denatured form.

In some embodiments, the at least one antibody or antigen-binding fragment thereof that specifically binds to at least one epitope in a neuropilin domain selected from one or more of the neuropilin b1 domain, neuropilin a1 domain, neuropilin a2 domain, neuropilin b2 domain, neuropilin c domain, neuropilin a1/a2 combined domain, neuropilin b1/b2 combined domain, neuropilin a2/b1 combined domain, neuropilin b2/c combined domain, neuropilin a2/b1/b2 combined domain, neuropilin a2/b1/b2/c combined domain, neuropilin a1/a2/b1 combined domain, neuropilin a1/a2/b1/b2 combined domain, neuropilin a1/a2/b1/b2/c combined domain, and the neuropilin b1/b2/c combined domain, optionally with an affinity of about 10 pM to about 500 pM or to about 50 nM, or about, at least about, or no more than about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900 pM, 1 nM, 10 nM, 25 nM, or 50 nM, or optionally with an affinity that ranges from about 10 pM to about 500 pM, about 10 pM to about 400 pM, about 10 pM to about 300 pM, about 10 pM to about 200 pM, about 10 pM to about 100 pM, about 10 pM to about 50 pM, or about 20 pM to about 500 pM, about 20 pM to about 400 pM, about 20 pM to about 300 pM, about 20 pM to about 200 pM, about 20 pM to about 100 pM, about 20 pM to about 50 pM, or about 30 pM to about 500 pM, about 30 pM to about 400 pM, about 30 pM to about 300 pM, about 30 pM to about 200 pM, about 30 pM to about 100 pM, about 30 pM to about 50 pM, or about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 1 nM to about 5 nM, about 5 nM to about 10 nM, about 10 nM to 25 nM, or about 25 nM to about 50 nM.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin a1 domain, the neuropilin a2 domain, and/or the neuropilin a1a2 combined domain, including adjacent linker regions, optionally at about residues;

(neuropilin a1 domain) 20-148, 30-141, 40-141, 50-141, 60-141, 70-141, 80-141, 90-141, 100-141, 110-141,

5

120-141, 130-141; 20-130, 20-120, 20-110, 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, or 20-30 as defined by a human NRP2 precursor sequence (see Table N1);

(neuropilin a2 domain) 142-280, 150-265, 160-265, 170-265, 180-265, 190-265, 200-265, 210-265, 220-265, 230-265, 240-265, 250-265, 260-265, 141-270, 141-260, 141-250, 141-240, 141-230, 141-220, 141-210, 141-200, 141-190, 141-180, 141-170, 141-160, 141-150, 200-250, 210-250, 220-250, 230-250, 200-240, 210-240, 220-240, 230-240, 227-247, 228-247, 229-247, 230-247, 231-247, 232-247, 233-247, 234-247, 235-247, 236-247; 227-246, 227-245, 227-244, 227-243, 227-242, 227-241, 227-240, 227-239, 227-238; 235-240, 236-239, 236-238, or residue 237 as defined by a human NRP2 precursor sequence (see Table N1); or

(combined a1a2 domain) 20-280, 30-280, 40-280, 50-280, 60-280, 70-280, 80-280, 90-280, 100-280, 110-280, 120-280, 130-280, 140-280, 150-280, 160-280, 170-280, 180-280, 190-280, 200-280, 210-280, 220-280, 230-280, 240-280, 260-280, 270-280, 20-270, 20-260, 20-250, 20-240, 20-230, 20-220, 20-210, 20-200, 20-190, 20-180, 20-170, 20-160, 20-150, 20-140, 20-130, 20-120, 20-110, 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, or 20-30 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin b1 domain, the neuropilin b2 domain, and/or the neuropilin b1/b2 combined domain, including adjacent linker regions, optionally at about residues;

(neuropilin b1 domain) 299-420, 266-426, 280-426, 290-426, 300-426, 310-426, 320-426, 330-426, 340-426, 350-426, 360-426, 370-426, 380-426, 390-426, 400-426, 410-426, 420-426, 280-420, 280-410, 280-400, 280-390, 280-380, 280-370, 280-360, 280-350, 280-340, 280-330, 280-320, 280-310, 280-300, or 280-290 as defined by a human NRP2 precursor sequence (see Table N1), optionally wherein the epitope is a discontinuous epitope that comprises one, two, or three of residues 299Y, 354N, and/or 416S as defined by the human NRP2 precursor sequence;

(neuropilin b2 domain) 438-591, 450-591, 460-591, 470-591, 480-591, 490-591, 500-591, 510-591, 520-591, 530-591, 540-591, 550-591, 560-591, 570-591, 580-591, 438-590, 438-580, 438-570, 438-560, 438-550, 438-540, 438-530, 438-520, 438-510, 438-500, 438-490, 438-480, 438-470, 438-460, or 438-450 as defined by a human NRP2 precursor sequence (see Table N1); or

(neuropilin b1/b2 combined domain) 266-591, 276-591, 286-591, 296-591, 306-591, 316-591, 326-591, 336-591, 346-591, 356-591, 366-591, 376-591, 386-591, 396-591, 406-591, 416-591, 426-591, 436-591, 446-591, 456-591, 466-591, 476-591, 486-591, 498-591, 508-591, 518-591, 528-591, 538-591, 548-591, 558-591, 568-591, 578-591, 588-591, 266-581, 266-571, 266-561, 266-551, 266-541, 266-531, 266-521, 266-511, 266-501, 266-491, 266-481, 266-471, 266-461, 266-451, 266-441, 266-431, 266-421, 266-411, 266-401, 266-391, 266-381, 266-371, 266-361, 266-351, 266-341, 266-331, 266-321, 266-311, 266-301, 266-291, 266-281, or 266-271 as defined by a human NRP2 precursor sequence (see Table N1).

6

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin a2/b1 combined domain and/or the neuropilin b2c combined domain, including adjacent linker regions, optionally at about residues;

(neuropilin a2b1 combined domain) 149-437, 159-426, 169-426, 179-426, 189-426, 199-426, 209-426, 219-426, 229-426, 239-426, 249-426, 259-426, 269-426, 279-426, 289-426, 299-426, 309-426, 319-426, 329-426, 339-426, 349-426, 359-426, 369-426, 379-426, 389-426, 399-426, 409-426, 419-426, 149-436, 149-426, 149-416, 149-406, 149-396, 149-386, 149-376, 149-366, 149-356, 149-346, 149-336, 149-326, 149-316, 149-306, 149-296, 149-286, 149-276, 149-266, 149-256, 149-246, 149-236, 149-226, 149-216, 149-206, 149-196, 146-186, 146-176, 146-166, or 146-155 as defined by a human NRP2 precursor sequence (see Table N1); or

(neuropilin b2c combined domain) 438-794, 448-794, 458-794, 468-794, 478-794, 487-794, 497-794, 507-794, 517-794, 527-794, 537-794, 547-794, 557-794, 567-794, 587-794, 597-794, 607-794, 617-794, 627-794, 637-794, 647-794, 657-794, 667-794, 677-794, 687-794, 697-794, 707-794, 717-794, 727-794, 737-794, 747-794, 757-794, 767-794, 777-794, 787-794, 427-794, 438-784, 438-774, 438-764, 438-754, 438-744, 438-734, 438-728, 438-714, 438-704, 438-694, 438-684, 438-674, 438-664, 438-654, 438-644, 438-634, 438-624, 438-614, 438-604, 438-596, 438-586, 438-576, 438-566, 438-556, 438-546, 438-536, 438-526, 438-516, 438-506, 438-494, 438-484, 438-474, 438-464, 438-454, 438-444 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin c domain, including adjacent linker regions, optionally at about residues 591-794, 600-794, 610-794, 620-794, 630-794, 640-794, 650-794, 660-794, 670-794, 680-794, 690-794, 700-794, 710-794, 720-794, 730-794, 740-794, 750-794, 760-794, 770-794, 780-794, 790-794, 591-790, 591-780, 591-770, 591-760, 591-750, 591-740, 591-730, 591-720, 591-710, 591-700, 591-690, 591-680, 591-670, 591-660, 591-650, 591-640, 591-630, 591-620, 591-610, or 591-600 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin b1/b2c combined domain, including adjacent linker regions, optionally at about residues 276-794, 286-794, 296-794, 306-794, 316-794, 326-794, 336-794, 346-794, 356-794, 366-794, 376-794, 387-794, 396-794, 406-794, 416-794, 426-794, 436-794, 446-794, 456-794, 466-794, 476-794, 486-794, 496-794, 506-794, 516-794, 526-794, 536-794, 546-794, 556-794, 566-794, 576-794, 586-794, 596-794, 606-794, 616-794, 626-794, 636-794, 646-794, 656-794, 666-794, 676-794, 686-794, 696-794, 706-794, 716-794, 726-794, 736-794, 746-794, 756-794, 766-794, 776-794, 786-794, 266-794, 276-784, 276-774, 276-764, 276-754, 276-744, 276-734, 276-724, 276-714, 276-704, 276-694, 276-684, 276-674, 276-664, 276-654, 276-644, 276-634, 276-624, 276-614, 276-604, 276-594, 276-584, 276-574, 276-564, 276-554, 276-544, 276-534, 276-524, 276-514, 276-504, 276-594, 276-584, 276-574, 276-564, 276-554, 276-544, 276-534, 276-524, 276-514, 276-504, or 276-496 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the juxtamembrane domain (see Table N1), optionally selected from one or more of the juxtamembrane domain of NRP2a (variant 1), the juxtamembrane domain of NRP2a (variant 2), the juxtamembrane domain of NRP2a (variant 3), the juxtamembrane domain of NRP2b (variant 4), and the juxtamembrane domain of NRP2b (variant 5), including combinations thereof.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a conformational epitope composed of two or more discontinuous epitope regions, optionally a conformational epitope comprising or consisting of:

- (a) a first epitope region within the a1 domain, and second epitope region within the a2 domain of the human NRP2 polypeptide;
- (b) a first epitope region within the a1 domain, and second epitope region within the b1 domain of the human NRP2 polypeptide;
- (c) a first epitope region within the a1 domain, and second epitope region within the b2 domain of the human NRP2 polypeptide;
- (d) a first epitope region within the a1 domain, and second epitope region within the c domain of the human NRP2 polypeptide;
- (e) a first epitope region within the a1 domain, and second epitope region within the juxtamembrane domain of the human NRP2 polypeptide selected from variant 1, 2, 3, 4 and 5;
- (f) a first epitope region within the a2 domain, and second epitope region within the b1 domain of the human NRP2 polypeptide;
- (g) a first epitope region within the a2 domain, and second epitope region within the b2 domain of the human NRP2 polypeptide;
- (h) a first epitope region within the a2 domain, and second epitope region within the c domain of the human NRP2 polypeptide;
- (i) a first epitope region within the a2 domain, and second epitope region within the juxtamembrane domain of the human NRP2 polypeptide selected from variant 1, 2, 3, 4 and 5;
- (j) a first epitope region within the b1 domain, and second epitope region within the b2 domain of the human NRP2 polypeptide;
- (k) a first epitope region within the b1 domain, and second epitope region within the c domain of the human NRP2 polypeptide;
- (l) a first epitope region within the b1 domain, and second epitope region within the juxtamembrane domain of the human NRP2 polypeptide selected from variant 1, 2, 3, 4 and 5;
- (m) a first epitope region within the b2 domain, and second epitope region within the c domain of the human NRP2 polypeptide;
- (n) a first epitope region within the b2 domain, and second epitope region within the juxtamembrane domain of the human NRP2 polypeptide selected from variant 1, 2, 3, 4 and 5; or
- (o) a first epitope region within the c domain, and second epitope region within the juxtamembrane domain of the human NRP2 polypeptide selected from variant 1, 2, 3, 4 and 5.

In some embodiments, the at least one antibody or antigen-binding fragment thereof modulates binding of the human NRP2 polypeptide to at least one NRP2 ligand

(optionally an NRP2 ligand selected from Table N2 or Table N3 and/or a human histidyl-tRNA synthetase (HRS) polypeptide selected from Table H1, optionally a HRS splice variant selected from one or more of SV9 (HRS (1-60)), SV11 (HRS (1-60)+(399-509)) and SV14 (HRS (1-100)+(399-509)).

In some embodiments, the at least one antibody or antigen-binding fragment thereof is a blocking antibody which inhibits about or at least about 80-100% of the theoretical maximal binding between the human NRP2 polypeptide and the least one NRP2 ligand, after pre-incubation with the human NRP2 polypeptide in a stoichiometrically equivalent amount, optionally about or at least about 80, 85, 90, 95, or 100% of the theoretical maximal binding. In some embodiments, the at least one antibody or antigen-binding fragment thereof is a partial blocking antibody which inhibits about or at least about 20-80% of the theoretical maximal binding between the human NRP2 polypeptide and the at least one NRP2 ligand, after pre-incubation with the human NRP2 polypeptide in a stoichiometrically equivalent amount, optionally about or at least about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, or 80% of the theoretical maximal binding. In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an HRS polypeptide-interacting region of the NRP2 polypeptide, and mimics or agonizes one or more signaling activities of the HRS polypeptide binding to the NRP2 polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an HRS polypeptide-interacting region of the NRP2 polypeptide, and modulates binding/signaling activity between the NRP2 polypeptide and at least one NRP2 ligand. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand.

In some embodiments, the at least one antibody or antigen-binding fragment thereof agonizes or enhances the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand. In some embodiments, the at least one NRP2 ligand is selected from:

- a VEGF selected from one or more of VEGF-A145, VEGF-A165, VEGF-C, VEGF-D and PIGF-2;
- a VEGF receptor (VEGFR) selected from VEGFR2 and VEGFR3;
- a semaphorin selected from one or more of SEMA3-A, SEMA-3B, SEMA-3C, SEMA-3D SEMA-3F, and SEMA-3G;
- a plexin selected from one or more of plexin A1, A2, A3, A4, and D1;
- a growth factor selected from one or more of fibroblast growth factor (FGF), hepatocyte growth factor (HGF), and platelet derived growth factor (PDGF);
- a growth factor receptor selected from one or more of a fibroblast growth factor receptor (FGFR), a hepatocyte growth factor receptor (HGFR), and a platelet derived growth factor receptor (PDGFR);
- a galectin or a galectin receptor
- a transcription factor selected from FAC1 and bromoprotein PHD finger transcription factor;
- an adaptor protein selected from one or more of GIPC1, GIPC2 and GIPC3;
- an integrin selected from Table N3, optionally one or more of $\alpha_1\beta_1$, $\alpha_1\beta_3$, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_v\beta_6$, $\alpha_v\beta_8$, $\alpha_6\beta_1$, and $\alpha_6\beta_4$;
- a transforming growth factor beta selected from one or more of TGF β 1, TGF β 2, TGF β 3, and their corresponding TGF β receptors; and

an HRS polypeptide selected from Table H1, optionally an HRS splice variant selected from one or more of HisRS^{N1}, HisRS^{N2}, HisRS^{N3}, HisRS^{N4}(SV9), HisRS^{N5}, HisRS^{C1}, HisRS^{C2}, HisRS^{C3}, HisRS^{C4}, HisRS^{C5}, HisRS^{C6}, HisRS^{C7}, HisRS^{C8}(SV11), and HisRS^{C9}(SV14).

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and VEGFR2 or VEGFR3 or VEGF-C. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a HRS polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and VEGFR2 or VEGFR3 or VEGF-C. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and VEGFR3 without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and VEGFR3 or VEGF-C without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a different ligand, optionally an HRS polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor without substantially modulating the ligand binding of semaphorin 3 to NRP2. In some embodiments, the plexin receptor is selected from plexin A1, A2, A3, A4, and D1. In some embodiments, the semaphorin is selected from semaphorin 3B, 3C, 3D, 3F, and 3G.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 contiguous amino acids within the human NRP2 a2 domain, wherein the at least one antibody or antigen-binding fragment thereof selectively inhibits receptor dimerization between NRP2 and plexin A1 without substantially inhibiting dimerization between NRP2 and FLT4 (VEGFR3). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope within amino acids 232-242 of a human NRP2 precursor (see Table N1). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a discontinuous epitope comprised within amino acids 299-416 the human NRP2 b1 domain, wherein the at least one antibody or antigen-binding fragment thereof selectively inhibits receptor dimerization between NRP2 and FLT4 (VEGFR3) and KDR (VEGFR2) without substantially inhibiting dimerization between NRP2 and plexin A1.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 contiguous amino acids within the human NRP2 b2 domain, wherein the at least one antibody

or antigen-binding fragment thereof inhibits receptor dimerization between NRP2 and FLT4 (VEGFR3) and inhibits dimerization between NRP2 and plexin A1. In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 contiguous amino acids within the human NRP2 c domain, wherein the at least one antibody or antigen-binding fragment thereof inhibits receptor dimerization between NRP2 and plexin A1 and partially inhibits dimerization between NRP2 and FLT4 (VEGFR3).

In some embodiments, the at least one antibody or antigen-binding fragment thereof has an affinity (K_d or EC₅₀) for each of (i) a human NRP2 polypeptide and (ii) the corresponding region of a cynomolgus monkey NRP2 polypeptide, wherein the affinity for (i) and (ii) is within the range of about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 0.4 to about 1.2 nM, about 0.9 to about 5.5 nM, about 0.9 to about 5 nM, or about 1 nM to about 10 nM. In some embodiments, the at least one antibody or antigen-binding fragment thereof has an affinity (K_d or EC₅₀) for each of (i) a human NRP2 polypeptide and (ii) the corresponding region of a murine NRP2 polypeptide, wherein the affinity for (i) and (ii) is within the range of about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, or about 1 nM to about 10 nM.

In some embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to the NRP2a isoform (optionally variants 1, 2, and/or 3 of Table N1) of NRP2, and does not substantially bind to the NRP2b isoform (optionally variants 4 and/or 5 of Table N1). In some embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to the NRP2b isoform (optionally variants 4 and/or 5 of Table N1), and does not substantially bind to the NRP2a isoform (optionally variants 1, 2, and/or 3 of Table N1). In some embodiments, the at least one antibody or antigen-binding fragment thereof reduces the homo- or hetero-dimerization between NRP2 polypeptides, optionally by about or at least about 20-100% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, or 100%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptides in a substantially stoichiometrically equivalent amount, optionally in the presence of an NRP2 ligand.

In some embodiments, the at least one antibody or antigen-binding fragment thereof enhances the homo- or hetero-dimerization between NRP2 polypeptides, optionally by about or at least about 20-100% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, or 100%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptides in a substantially stoichiometrically equivalent amount, optionally in the presence of an NRP2 ligand. In some embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to a human NRP2 polypeptide (see Table N1) relative to a murine NRP2 polypeptide, optionally where its affinity for the human NRP2 polypeptide is significantly stronger than its affinity for the murine NRP2 polypeptide, optionally by about or at least about 2, 5, 10, 20, 30, 40, 50, 100, 500, or 1000-fold or more. In some embodiments, the at least one antibody or antigen-binding fragment thereof binds to the human NRP2

11

polypeptide and does not substantially bind to the murine NRP2 polypeptide, optionally wherein the murine NRP2 polypeptide is a *Mus musculus* NRP2 polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof binds to an epitope in the b1 domain that comprises residues 299Y, 354N, and 416S, as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof comprises an IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), or IgM Fc domain, optionally a human Fc domain, or a hybrid and/or variant thereof. In some embodiments, the at least one antibody or antigen-binding fragment thereof comprises an IgG Fc domain with high effector function in humans, optionally an IgG1 or IgG3 Fc domain. In some embodiments, the at least one antibody or antigen-binding fragment thereof comprises an IgG Fc domain with low effector function in humans, optionally an IgG2 or IgG4 Fc domain. In some embodiments, the at least one antibody or antigen-binding fragment thereof comprises an IgG1 or IgG4 Fc domain, optionally selected from Table F1. In some embodiments, the at least one antibody or antigen-binding fragment thereof comprises a modified IgG1 or IgG4 Fc domain which has altered binding to FcRn, optionally wherein the modified IgG1 or IgG4 Fc domain comprises any one or more of YD (M252Y/T256D), DQ (T256D/T307Q), DW (T256D/T307W), YTE (M252Y/S254T/T256E), AAA (T307A/E380A/N434A), LS (M428L/N434S), M252Y, T256D/E, K288D/N, T307Q/W, E380C, N434FY, and/or Y436H/N/W mutations (EU numbering), including combinations thereof.

In some embodiments, the at least one antibody or antigen-binding fragment thereof is a monoclonal antibody. In some embodiments, the at least one antibody or antigen-binding fragment thereof is a humanized antibody. In some embodiments, the at least one antibody or antigen-binding fragment thereof is an Fv fragment, a single chain Fv (scFv) polypeptide, an adnectin, an anticalin, an aptamer, an avimer, a camelid antibody, a designed ankyrin repeat protein (DARPin), a minibody, a nanobody, or a unibody.

In some embodiments, the therapeutic composition has a purity of at least about 80%, 85%, 90%, 95%, 98%, or 99% on a protein basis with respect to the at least one antibody or antigen-binding fragment, and is substantially aggregate-free. In some embodiments, the therapeutic composition is substantially endotoxin-free. In some embodiments, the therapeutic composition is a sterile, injectable solution, optionally suitable for intravenous, intramuscular, subcutaneous, or intraperitoneal administration.

In some embodiments, the therapeutic composition further comprises at least one additional agent selected from one or more of a cancer immunotherapy agent, a chemotherapeutic agent, a hormonal therapeutic agent, and a kinase inhibitor. In some embodiments, the cancer immunotherapy agent is selected from one or more of an immune checkpoint modulatory agent, a cancer vaccine, an oncolytic virus, a cytokine, and a cell-based immunotherapies. In some embodiments, the immune checkpoint modulatory agent is a polypeptide, optionally an antibody or antigen-binding fragment thereof or a ligand, or a small molecule. In some embodiments, the immune checkpoint modulatory agent comprises

- (a) an antagonist of a inhibitory immune checkpoint molecule; or
- (b) an agonist of a stimulatory immune checkpoint molecule,

12

optionally wherein the immune checkpoint modulatory agent specifically binds to the immune checkpoint molecule.

In some embodiments, the inhibitory immune checkpoint molecule is selected from one or more of Programmed Death-Ligand 1 (PD-L1), Programmed Death 1 (PD-1), Programmed Death-Ligand 2 (PD-L2), Cytotoxic T-Lymphocyte-Associated protein 4 (CTLA-4), Indoleamine 2,3-dioxygenase (IDO), tryptophan 2,3-dioxygenase (TDO), T-cell Immunoglobulin domain and Mucin domain 3 (TIM-3), Lymphocyte Activation Gene-3 (LAG-3), V-domain Ig suppressor of T cell activation (VISTA), B and T Lymphocyte Attenuator (BTLA), CD160, Herpes Virus Entry Mediator (HVEM), and T-cell immunoreceptor with Ig and ITIM domains (TIGIT).

In some embodiments:

the antagonist is a PD-L1 and/or PD-L2 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, atezolizumab (MPDL3280A), avelumab (MSB0010718C), and durvalumab (MEDI4736);

the antagonist is a PD-1 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, nivolumab, pembrolizumab, MK-3475, AMP-224, AMP-514, PDR001, and pidilizumab;

the antagonist is a CTLA-4 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, ipilimumab, and tremelimumab;

the antagonist is an IDO antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, indoximod (NLG-8189), 1-methyl-tryptophan (1MT), β -Carboline (norharmaline; 9H-pyrido[3,4-b]indole), rosmarinic acid, and epacadostat;

the antagonist is a TDO antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, 680C91, and LM10;

the antagonist is a TIM-3 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

the antagonist is a LAG-3 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, and BMS-986016;

the antagonist is a VISTA antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

the antagonist is a BTLA, CD160, and/or HVEM antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

and/or the antagonist is a TIGIT antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto.

In some embodiments, the stimulatory immune checkpoint molecule is selected from one or more of OX40, CD40, Glucocorticoid-Induced TNFR Family Related Gene (GITR), CD137 (4-1BB), CD27, CD28, CD226, and Herpes Virus Entry Mediator (HVEM).

13

In some embodiments:

the agonist is an OX40 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, OX86, Fc-OX40L, and GSK3174998;

the agonist is a CD40 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, CP-870,893, dacetuzumab, Chi Lob 7/4, ADC-1013, and rhCD40L;

the agonist is a GITR agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, INCAGN01876, DTA-1, and MEDI1873;

the agonist is a CD137 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, utomilumab, and 4-1BB ligand;

the agonist is a CD27 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, varlilumab, and CDX-1127 (1F5);

the agonist is a CD28 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, and TAB08; and/or

the agonist is an HVEM agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto.

In some embodiments, the cancer vaccine is selected from one or more of Oncophage, a human papillomavirus HPV vaccine optionally Gardasil or Cervarix, a hepatitis B vaccine optionally Engerix-B, Recombivax HB, or Twinrix, and sipuleucel-T (Provenge), or comprises a cancer antigen selected from one or more of human Her2/neu, Her1/EGF receptor (EGFR), Her3, A33 antigen, B7H3, CD5, CD19, CD20, CD22, CD23 (IgE Receptor), MAGE-3, C242 antigen, 5T4, IL-6, IL-13, vascular endothelial growth factor VEGF (e.g., VEGF-A) VEGFR-1, VEGFR-2, CD30, CD33, CD37, CD40, CD44, CD51, CD52, CD56, CD74, CD80, CD152, CD200, CD221, CCR4, HLA-DR, CTLA-4, NPC-1C, tenascin, vimentin, insulin-like growth factor 1 receptor (IGF-1R), alpha-fetoprotein, insulin-like growth factor 1 (IGF-1), carbonic anhydrase 9 (CA-IX), carcinoembryonic antigen (CEA), guanylyl cyclase C, NY-ESO-1, p53, survivin, integrin α v β 3, integrin α 5 β 1, folate receptor 1, transmembrane glycoprotein NMB, fibroblast activation protein alpha (FAP), glycoprotein 75, TAG-72, MUC1, MUC16 (or CA-125), phosphatidylserine, prostate-specific membrane antigen (PSMA), NR-LU-13 antigen, TRAIL-R1, tumor necrosis factor receptor superfamily member 10b (TNFRSF10B or TRAIL-R2), SLAM family member 7 (SLAMF7), EGP40 pancarcinoma antigen, B-cell activating factor (BAFF), platelet-derived growth factor receptor, glycoprotein EpCAM (17-1A), Programmed Death-1, protein disulfide isomerase (PDI), Phosphatase of Regenerating Liver 3 (PRL-3), prostatic acid phosphatase, Lewis-Y antigen, GD2 (a disialoganglioside expressed on tumors of neuroectodermal origin), glypican-3 (GPC3), and mesothelin.

In some embodiments, the oncolytic virus selected from one or more of talimogene laherparepvec (T-VEC), coxsackievirus A21 (CAVATAK™), Oncorine (H101), pelareorep (REOLYSIN®), Seneca Valley virus (NTX-010), Sen-

14

ecavirus SVV-001, ColoAd1, SEPREHVIR (HSV-1716), CGTG-102 (Ad5/3-D24-GMCSF), GL-ONC1, MV-NIS, and DNX-2401.

In some embodiments, the cytokine selected from one or more of interferon (IFN)- α , IL-2, IL-12, IL-7, IL-21, and Granulocyte-macrophage colony-stimulating factor (GM-CSF).

In some embodiments, the cell-based immunotherapy agent comprises cancer antigen-specific T-cells, optionally ex vivo-derived T-cells. In some embodiments, the cancer antigen-specific T-cells are selected from one or more of chimeric antigen receptor (CAR)-modified T-cells, and T-cell Receptor (TCR)-modified T-cells, tumor infiltrating lymphocytes (TILs), and peptide-induced T-cells.

In some embodiments, the at least one chemotherapeutic agent is selected from one or more of an alkylating agent, an anti-metabolite, a cytotoxic antibiotic, a topoisomerase inhibitor (type I or type II), and an anti-microtubule agent.

In some embodiments:

the alkylating agent is selected from one or more of nitrogen mustards (optionally mechlorethamine, cyclophosphamide, mustine, melphalan, chlorambucil, ifosfamide, and busulfan), nitrosoureas (optionally N-Nitroso-N-methylurea (MNU), carmustine (BCNU), lomustine (CCNU), semustine (MeCCNU), fotemustine, and streptozotocin), tetrazines (optionally dacarbazine, mitozolomide, and temozolomide), aziridines (optionally thiotepe, mytomycin, and diaziquone (AZQ)), cisplatins and derivatives thereof (optionally carboplatin and oxaliplatin), and non-classical alkylating agents (optionally procarbazine and hexamethylmelamine);

the anti-metabolite is selected from one or more of anti-folates (optionally methotrexate and pemetrexed), fluoropyrimidines (optionally 5-fluorouracil and capecitabine), deoxynucleoside analogues (optionally ancitabine, enocitabine, cytarabine, gemcitabine, decitabine, azacitidine, fludarabine, nelarabine, cladribine, clofarabine, fludarabine, and pentostatin), and thiopurines (optionally thioguanine and mercaptopurine);

the cytotoxic antibiotic is selected from one or more of anthracyclines (optionally doxorubicin, daunorubicin, epirubicin, idarubicin, pirarubicin, aclarubicin, and mitoxantrone), bleomycins, mitomycin C, mitoxantrone, and actinomycin;

the topoisomerase inhibitor is selected from one or more of camptothecin, irinotecan, topotecan, etoposide, doxorubicin, mitoxantrone, teniposide, novobiocin, merbarone, and aclarubicin; and/or

the anti-microtubule agent is selected from one or more of taxanes (optionally paclitaxel and docetaxel) and vinca alkaloids (optionally vinblastine, vincristine, vindesine, vinorelbine).

In some embodiments, the at least one hormonal therapeutic agent is a hormonal agonist or a hormonal antagonist. In some embodiments, the hormonal agonist is selected from one or more of a progestogen (progestin), a corticosteroid (optionally prednisolone, methylprednisolone, or dexamethasone), insulin like growth factors, VEGF derived angiogenic and lymphangiogenic factors (optionally VEGF-A, VEGF-A145, VEGF-A165, VEGF-C, VEGF-D, PIGF-2), fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor (PDGF), transforming growth factor (TGF)-beta, an androgen, an estrogen, and a somatostatin analog. In some embodiments, the hormonal antagonist is selected from one or more of a hormone synthesis inhibitor, optionally an aromatase inhibi-

tor or a gonadotropin-releasing hormone (GnRH) or an analog thereof, and a hormone receptor antagonist, optionally a selective estrogen receptor modulator (SERM) or an anti-androgen, or an antibody directed against a hormonal receptor, optionally cixutumumab, dalotuzumab, figitumumab, ganitumab, istiratutumab, robatumumab, alacizumab pegol, bevacizumab, icrucumab, ramucirumab, fresolimumab, metelimumab, naxitamab, cetuximab, depatuxizumab mafodotin, futuximab, imgatuzumab, laprituximab emtansine, matuzumab, modotuximab, necitumumab, nimotuzumab, panitumumab, tomuzotuximab, zalutumumab, aprutumab ixadotin, bemarituzumab, olaratumab, or tovetumab.

In some embodiments, the kinase inhibitor is selected from one or more of adavosertib, afanitib, aflibercept, axitinib, bevacizumab, bosutinib, cabozantinib, cetuximab, cobimetinib, crizotinib, dasatinib, entrectinib, erdafitinib, erlotinib, fostamitinib, gefitinib, ibrutinib, imatinib, lapatinib, lenvatinib, mubritinib, nilotinib, panitumumab, pazopanib, pegaptanib, ponatinib, ranibizumab, regorafenib, ruxolitinib, sorafenib, sunitinib, SU6656, tofacitinib, trastuzumab, vandetanib, and vemurafenib.

Also included are methods of treating a disease or condition in a subject in need thereof, comprising administering to the subject a therapeutic composition comprising at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin-2 (NRP2) polypeptide, wherein the at least one antibody or antigen-binding fragment thereof modulates (e.g., interferes with) binding of the human NRP2 polypeptide to a human histidyl-tRNA synthetase (HRS) polypeptide, optionally as a therapeutic composition described herein.

In some embodiments, the disease or condition is an NRP2-associated disease or condition. In some embodiments, the disease or condition is selected from one or more of cancer and diseases and pathways associated with cancer, including cancer cell growth, initiation, migration, adhesion, invasion, chemoresistance, and/or metastasis; diseases associated with inflammation, autoimmunity, and related inflammatory diseases, including diseases associated with inappropriate immune cell activation or migration such as Graft versus host disease (GVHD); diseases associated with lymphatic development, lymphangioma, lymphangiogenesis, and lymphatic damage, including, for example, edema, lymphedema, secondary lymphedema, inappropriate fat absorption and deposition, excess fat deposition, and vascular permeability; diseases associated with infections, including latent infections; diseases associated with allergic disorders/diseases, allergic responses, including, for example, chronic obstructive pulmonary disorder (COPD), neutrophilic asthma, antineutrophil cytoplasmic antibody (ANCA)-associated systemic vasculitis, systemic lupus erythematosus, rheumatoid arthritis, inflammasome-related diseases, and skin-related neutrophil-mediated diseases such as pyoderma gangrenosum; diseases associated with granulomatous inflammatory diseases, including sarcoidosis and granulomas; diseases associated with fibrosis including fibrotic diseases, fibrosis, endothelial to mesenchymal transition (EMT), and wound healing; diseases associated with inappropriate smooth muscle contractility, smooth muscle compensation and decompensation, and inappropriate vascular smooth muscle cell migration and adhesion; diseases associated with inappropriate autophagy, phagocytosis, and efferocytosis; diseases associated with inappropriate migratory cell movement; diseases associated with neuronal dis-

eases, peripheral nervous system remodeling, and pain perception; and diseases associated with bone development and bone remodeling.

In some embodiments, the disease is a cancer, optionally wherein the cancer expresses or overexpresses NRP2, optionally wherein the cancer displays NRP2-dependent growth, NRP2-dependent adhesion, NRP2-dependent migration, and/or NRP2-dependent invasion. In some embodiments, the cancer expresses or overexpresses NRP2 but does not substantially express neuropilin-1 (NRP1). Certain methods are directed to reducing or preventing re-emergence of a cancer in a subject in need thereof, wherein administration of the therapeutic composition enables generation of an immune memory to the cancer. In some embodiments, the subject has lymphedema.

Certain embodiments comprise administering to the subject at least one additional agent selected from one or more of a cancer immunotherapy agent, a chemotherapeutic agent, a hormonal therapeutic agent, and a kinase inhibitor. In some embodiments, the at least one anti-NRP2 antibody or antigen-binding fragment thereof and the at least one agent are administered separately, as separate compositions. In some embodiments, the at least one anti-NRP2 antibody and the at least one agent are administered together as part of the same therapeutic composition, optionally as a therapeutic composition described herein. In some embodiments, the cancer immunotherapy agent is selected from one or more of an immune checkpoint modulatory agent, a cancer vaccine, an oncolytic virus, a cytokine, and a cell-based immunotherapies.

In some embodiments, the immune checkpoint modulatory agent is a polypeptide, optionally an antibody or antigen-binding fragment thereof or a ligand, or a small molecule. In some embodiments, the immune checkpoint modulatory agent comprises

- (a) an antagonist of an inhibitory immune checkpoint molecule; or
 - (b) an agonist of a stimulatory immune checkpoint molecule.
- optionally, wherein the immune checkpoint modulatory agent specifically binds to the immune checkpoint molecule.

In some embodiments, the inhibitory immune checkpoint molecule is selected from one or more of Programmed Death-Ligand 1 (PD-L1), Programmed Death 1 (PD-1), Programmed Death-Ligand 2 (PD-L2), Cytotoxic T-Lymphocyte-Associated protein 4 (CTLA-4), Indoleamine 2,3-dioxygenase (IDO), tryptophan 2,3-dioxygenase (TDO), T-cell Immunoglobulin domain and Mucin domain 3 (TIM-3), Lymphocyte Activation Gene-3 (LAG-3), V-domain Ig suppressor of T cell activation (VISTA), B and T Lymphocyte Attenuator (BTLA), CD160, Herpes Virus Entry Mediator (HVEM), and T-cell immunoreceptor with Ig and ITIM domains (TIGIT).

In some embodiments:

- the antagonist is a PD-L1 and/or PD-L2 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, atezolizumab (MPDL3280A), avelumab (MSB0010718C), and durvalumab (MEDI4736), optionally wherein the cancer is selected from one or more of colorectal cancer, melanoma, breast cancer, non-small-cell lung carcinoma, bladder cancer, and renal cell carcinoma;
- the antagonist is a PD-1 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds

thereto, nivolumab, pembrolizumab, MK-3475, AMP-224, AMP-514PDR001, and pidilizumab, optionally wherein the PD-1 antagonist is nivolumab and the cancer is optionally selected from one or more of Hodgkin's lymphoma, melanoma, non-small cell lung cancer, hepatocellular carcinoma, renal cell carcinoma, and ovarian cancer;

the PD-1 antagonist is pembrolizumab and the cancer is optionally selected from one or more of melanoma, non-small cell lung cancer, small cell lung cancer, head and neck cancer, and urothelial cancer;

the antagonist is a CTLA-4 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, ipilimumab, tremelimumab, optionally wherein the cancer is selected from one or more of melanoma, prostate cancer, lung cancer, and bladder cancer;

the antagonist is an IDO antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, indoximod (NLG-8189), 1-methyl-tryptophan (1MT), β -Carboline (norharmane; 9H-pyrido[3,4-b]indole), rosmarinic acid, and epacadostat, and wherein the cancer is optionally selected from one or more of metastatic breast cancer and brain cancer optionally glioblastoma multiforme, glioma, gliosarcoma or malignant brain tumor;

the antagonist is a TDO antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, 680C91, and LM10;

the antagonist is a TIM-3 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

the antagonist is a LAG-3 antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto, and BMS-986016;

the antagonist is a VISTA antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

the antagonist is a BTLA, CD160, and/or HVEM antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto;

the antagonist is a TIGIT antagonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule that specifically binds thereto.

In some embodiments, the stimulatory immune checkpoint molecule is selected from one or more of OX40, CD40, Glucocorticoid-Induced TNFR Family Related Gene (GITR), CD137 (4-1BB), CD27, CD28, CD226, and Herpes Virus Entry Mediator (HVEM).

In some embodiments:

the agonist is an OX40 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, OX86, Fc-OX40L, and GSK3174998;

the agonist is a CD40 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, CP-870,893, dacetuzumab, Chi Lob 7/4, ADC-1013, and rhCD40L, and wherein the cancer is optionally

selected from one or more of melanoma, pancreatic carcinoma, mesothelioma, and hematological cancers optionally lymphoma such as Non-Hodgkin's lymphoma;

the agonist is a GITR agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, INCAGN01876, DTA-1, and MEDI1873;

the agonist is a CD137 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, utomilumab, and 4-1BB ligand;

the agonist is a CD27 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, varilumab, and CDX-1127 (1F5);

the agonist is a CD28 agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto, and TAB08; and/or

the agonist is an HVEM agonist optionally selected from one or more of an antibody or antigen-binding fragment or small molecule or ligand that specifically binds thereto.

In some embodiments, the cancer vaccine is selected from one or more of Oncophage, a human papillomavirus HPV vaccine optionally Gardasil or Cervarix, a hepatitis B vaccine optionally Engerix-B, Recombivax HB, or Twinrix, and sipuleucel-T (Provenge), or comprises a cancer antigen selected from one or more of human Her2/neu, Her1/EGF receptor (EGFR), Her3, A33 antigen, β_7 H3, CD5, CD19, CD20, CD22, CD23 (IgE Receptor), MAGE-3, C242 antigen, 5T4, IL-6, IL-13, vascular endothelial growth factor VEGF (e.g., VEGF-A) VEGFR-1, VEGFR-2, CD30, CD33, CD37, CD40, CD44, CD51, CD52, CD56, CD74, CD80, CD152, CD200, CD221, CCR4, HLA-DR, CTLA-4, NPC-1C, tenascin, vimentin, insulin-like growth factor 1 receptor (IGF-1R), alpha-fetoprotein, insulin-like growth factor 1 (IGF-1), carbonic anhydrase 9 (CA-IX), carcinoembryonic antigen (CEA), guanylyl cyclase C, NY-ESO-1, p53, survivin, integrin $\alpha v \beta 3$, integrin $\alpha 5 \beta 1$, folate receptor 1, transmembrane glycoprotein NMB, fibroblast activation protein alpha (FAP), glycoprotein 75, TAG-72, MUC1, MUC16 (or CA-125), phosphatidylserine, prostate-specific membrane antigen (PMSA), NR-LU-13 antigen, TRAIL-R1, tumor necrosis factor receptor superfamily member 10b (TNFRSF10B or TRAIL-R2), SLAM family member 7 (SLAMF7), EGP40 pancarcinoma antigen, B-cell activating factor (BAFF), platelet-derived growth factor receptor, glycoprotein EpCAM (17-1A), Programmed Death-1, protein disulfide isomerase (PDI), Phosphatase of Regenerating Liver 3 (PRL-3), prostatic acid phosphatase, Lewis-Y antigen, GD2 (a disialoganglioside expressed on tumors of neuroectodermal origin), glypican-3 (GPC3), and mesothelin, optionally wherein the subject has or is at risk for having a cancer that comprises the corresponding cancer antigen.

In some embodiments, the oncolytic virus selected from one or more of talimogene laherparepvec (T-VEC), coxsackievirus A21 (CAVATAKTM), Oncorine (H101), pelareo-rep (REOLYSIN[®]), Seneca Valley virus (NTX-010), Senecavirus SVV-001, ColoAd1, SEPREHVIR (HSV-1716), CGTG-102 (Ad5/3-D24-GMCSF), GL-ONC1, MV-NIS, and DNX-2401.

In some embodiments, the cytokine selected from one or more of interferon (IFN)- α , IL-2, IL-12, IL-7, IL-21, and Granulocyte-macrophage colony-stimulating factor (GM-CSF).

In some embodiments, the cell-based immunotherapy agent comprises cancer antigen-specific T-cells, optionally ex vivo-derived T-cells. In some embodiments, the cancer antigen-specific T-cells are selected from one or more of chimeric antigen receptor (CAR)-modified T-cells, and T-cell Receptor (TCR)-modified T-cells, tumor infiltrating lymphocytes (TILs), and peptide-induced T-cells.

In some embodiments, the at least one chemotherapeutic agent is selected from one or more of an alkylating agent, an anti-metabolite, a cytotoxic antibiotic, a topoisomerase inhibitor (type I or type II), and an anti-microtubule agent.

In some embodiments:

the alkylating agent is selected from one or more of nitrogen mustards (optionally mechlorethamine, cyclophosphamide, mustine, melphalan, chlorambucil, ifosfamide, and busulfan), nitrosoureas (optionally N-Nitroso-N-methylurea (MNU), carmustine (BCNU), lomustine (CCNU), semustine (MeCCNU), fotemustine, and streptozotocin), tetrazines (optionally dacarbazine, mitozolomide, and temozolomide), aziridines (optionally thiotepa, mytomycin, and diaziquone (AZQ)), cisplatin and derivatives thereof (optionally carboplatin and oxaliplatin), and non-classical alkylating agents (optionally procarbazine and hexamethylmelamine);

the anti-metabolite is selected from one or more of anti-folates (optionally methotrexate and pemetrexed), fluoropyrimidines (optionally 5-fluorouracil and capecitabine), deoxynucleoside analogues (optionally ancitabine, enocitabine, cytarabine, gemcitabine, decitabine, azacitidine, fludarabine, nelarabine, cladribine, clofarabine, fludarabine, and pentostatin), and thiopurines (optionally thioguanine and mercaptopurine);

the cytotoxic antibiotic is selected from one or more of anthracyclines (optionally doxorubicin, daunorubicin, epirubicin, idarubicin, pirarubicin, aclarubicin, and mitoxantrone), bleomycins, mitomycin C, mitoxantrone, and actinomycin;

the topoisomerase inhibitor is selected from one or more of camptothecin, irinotecan, topotecan, etoposide, doxorubicin, mitoxantrone, teniposide, novobiocin, merbarone, and aclarubicin; and/or

the anti-microtubule agent is selected from one or more of taxanes (optionally paclitaxel and docetaxel) and vinca alkaloids (optionally vinblastine, vincristine, vindesine, vinorelbine).

In some embodiments, the at least one hormonal therapeutic agent is a hormonal agonist or a hormonal antagonist. In some embodiments, the hormonal agonist is selected from one or more of a progestogen (progesterone), a corticosteroid (optionally prednisolone, methylprednisolone, or dexamethasone), insulin like growth factors, VEGF derived angiogenic and lymphangiogenic factors (optionally VEGF-A, VEGF-A145, VEGF-A165, VEGF-C, VEGF-D, PIGF-2), fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor (PDGF), transforming growth factor (TGF)-beta, an androgen, an estrogen, and a somatostatin analog. In some embodiments, the hormonal antagonist is selected from one or more of a hormone synthesis inhibitor, optionally an aromatase inhibitor or a gonadotropin-releasing hormone (GnRH) or an analog thereof, and a hormone receptor antagonist, optionally a selective estrogen receptor modulator (SERM) or an anti-androgen, or an antibody directed against a hormonal receptor, optionally cixutumumab, dalotuzumab, figitumumab, ganitumab, istiratumab, robatumumab, alacizumab pegol, bevacizumab, icrucumab, ramucirumab, fresolim-

umab, metelimumab, naxitamab, cetuximab, depatuxizumab mafodotin, futuximab, imgatuzumab, laprituximab emtansine, matuzumab, modotuximab, necitumumab, nimotuzumab, panitumumab, tomuzotuximab, zalutumumab, aprutumab ixadotin, bemarituzumab, olaratumab, or tovetumab.

In some embodiments, the kinase inhibitor is selected from one or more of adavosertib, afanitib, aflibercept, axitinib, bevacizumab, bosutinib, cabozantinib, cetuximab, cobimetinib, crizotinib, dasatinib, entrectinib, erdafitinib, erlotinib, fostamitinib, gefitinib, ibrutinib, imatinib, lapatinib, lenvatinib, mubritinib, nilotinib, panitumumab, pazopanib, pegaptanib, ponatinib, ranibizumab, regorafenib, ruxolitinib, sorafenib, sunitinib, SU6656, tofacitinib, trastuzumab, vandetanib, and vemurafenib.

In some embodiments, the cancer is a primary cancer. In some embodiments, the cancer is a metastatic cancer, optionally a metastatic cancer that expresses NRP2a and/or NRP2b. In some embodiments, the cancer is selected from one or more of melanoma (e.g., metastatic melanoma), pancreatic cancer, bone cancer, prostate cancer, small cell lung cancer, non-small cell lung cancer (NSCLC), mesothelioma, leukemia (e.g., lymphocytic leukemia, chronic myelogenous leukemia, acute myeloid leukemia, relapsed acute myeloid leukemia), lymphoma, hepatoma (hepatocellular carcinoma), sarcoma, B-cell malignancy, breast cancer, ovarian cancer, colorectal cancer, glioma, glioblastoma multiforme, meningioma, pituitary adenoma, vestibular schwannoma, primary CNS lymphoma, primitive neuroectodermal tumor (medulloblastoma), kidney cancer (e.g., renal cell carcinoma), bladder cancer, uterine cancer, esophageal cancer, brain cancer, head and neck cancers, cervical cancer, testicular cancer, thyroid cancer, and stomach cancer. In some embodiments, the metastatic cancer is selected from one or more of:

- (a) a bladder cancer which has metastasized to the bone, liver, and/or lungs;
- (b) a breast cancer which has metastasized to the bone, brain, liver, and/or lungs;
- (c) a colorectal cancer which has metastasized to the liver, lungs, and/or peritoneum;
- (d) a kidney cancer which has metastasized to the adrenal glands, bone, brain, liver, and/or lungs;
- (e) a lung cancer which has metastasized to the adrenal glands, bone, brain, liver, and/or other lung sites;
- (f) a melanoma which has metastasized to the bone, brain, liver, lung, and/or skin/muscle;
- (g) a ovarian cancer which has metastasized to the liver, lung, and/or peritoneum;
- (h) a pancreatic cancer which has metastasized to the liver, lung, and/or peritoneum;
- (i) a prostate cancer which has metastasized to the adrenal glands, bone, liver, and/or lungs;
- (j) a stomach cancer which has metastasized to the liver, lung, and/or peritoneum;
- (l) a thyroid cancer which has metastasized to the bone, liver, and/or lungs; and
- (m) a uterine cancer which has metastasized to the bone, liver, lung, peritoneum, and/or vagina.

In some embodiments, the subject has, and/or is selected for treatment based on having, increased circulating or serum levels of at least one NRP2 ligand (optionally an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1), either bound or free, relative to the levels of a healthy control or matched control standard or population of subject(s), optionally about or at least about 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700,

21

800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 3000, 4000, or 5000 pM of the at least one NRP2 ligand, or about or at least about 30-100, 40-100, 50-100, 30-2000, 40-2000, 50-2000, 60-2000, 70-2000, 80-2000, 90-2000, 100-2000, 200-2000, 300-2000, 400-2000, 500-2000, 600-2000, 700-2000, 800-2000, 900-2000, 1000-2000, 2000-3000, 3000-4000, or 4000-5000 pM of the at least one NRP2 ligand.

In some embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of at least one NRP2 ligand (optionally an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1) and/or a coding mRNA thereof relative to a healthy control or matched control standard or population of subject(s), optionally a cancer which has increased levels or expression of the at least one NRP2 ligand and/or a coding mRNA thereof relative to a non-cancerous control cell or tissue, optionally relative to a non-cancerous cell or tissue of the same type as the cancer, optionally wherein the HRS polypeptide is a splice variant selected from HisRS^{N1}, HisRS^{N2}, HisRS^{N3}, HisRS^{N4}, HisRS^{N5}, HisRS^{C1}, HisRS^{C2}, HisRS^{C3}, HisRS^{C4}, HisRS^{C5}, HisRS^{C6}, HisRS^{C7}, HisRS^{C8}, and HisRS^{C9}.

In some embodiments, the subject has, and/or is selected for treatment based on having, increased circulating or serum levels of a soluble neuropilin 2 (NRP2) polypeptide (optionally selected from Table N1), either bound or free, relative to the levels of a healthy control or matched control standard or population of subject(s), optionally circulating or serum levels of about or at least about 10, 20, 30, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 3000, 4000, 5000 pM of the soluble NRP2 polypeptide, or optionally circulating or serum levels about 30-50, 50-100, 100-2000, 200-2000, 300-2000, 400-2000, 500-2000, 600-2000, 700-2000, 800-2000, 900-2000, 1000-2000, 2000-3000, 3000-4000, 4000-5000 pM of the soluble NRP2 polypeptide.

In some embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of an NRP2 polypeptide (optionally selected from Table N1) and/or a coding mRNA thereof relative to a healthy control or matched control standard or population of subject(s), optionally a cancer which has increased levels or expression of an NRP2 polypeptide (optionally selected from Table N1) and/or a coding mRNA thereof relative to a non-cancerous control cell or tissue, optionally relative to a non-cancerous cell or tissue of the same type as the cancer.

In some embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of NRP2a and/or NRP2b, or an altered ratio of NRP2a: NRP2b expression, relative to a healthy control or matched control standard or population of subject(s). In some embodiments, the levels of NRP2b are increased by about or at least about 10%, 20%, 30%, 40%, 50%, 100%, 200%, 300%, 400%, 500%, 600%, 700%, 800%, 900%, 1000% compared to a healthy control or matched control standard or population of subject(s). In some embodiments, the healthy control or matched control standard or population of subject(s) comprises average ranges for age-matched samples of cancerous or non-cancerous cells or tissue of the same type as the cancer, which comprise specific characteristics such as drug resistance, metastatic potential, aggressiveness, genetic signature (optionally p53 mutation(s), PTEN deletion, IGF1R expression), and/or expression patterns. In some embodiments, the sub-

22

ject has, and/or is selected for treatment based on having, increased circulating levels of HRS: NRP2 complexes relative to a healthy or matched control standard or population of subject(s).

Certain embodiments comprise administering the at least one anti-NRP2 antibody in an amount and at a frequency sufficient to achieve an average, sustained serum or circulating levels of a soluble NRP2 polypeptide of about or less than about 500 pM, 400 pM, 300 pM, 200 pM, 100 pM, 50 pM, 40 pM, 30 pM, 20 pM, or 10 pM. Certain embodiments comprise administering the at least one anti-NRP2 antibody in an amount and at a frequency sufficient to achieve a reduction in the circulating levels of HRS: NRP2 complexes, optionally a reduction of about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 95, 99, or 100%.

In some embodiments, the at least one anti-NRP2 antibody enhances the immune response to the cancer by about, or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more, relative to a control. In some embodiments, the at least one anti-NRP2 antibody reduces the rate of in vitro growth of the cancer by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some embodiments, the at least one anti-NRP2 antibody reduces the in vitro adhesiveness of the cancer to a substrate by about or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control, optionally wherein the substrate comprises laminin.

In some embodiments, the at least one anti-NRP2 antibody reduces the invasiveness of the cancer by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some embodiments, the at least one anti-NRP2 antibody inhibits the rate of migration or motility of the cancer by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some embodiments, the at least one anti-NRP2 antibody inhibits the rate of autophagy or endosome maturation (optionally endosome acidification) of the cancer or associated immune cells by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some embodiments, the at least one anti-NRP2 antibody enhances the susceptibility of the cancer to an additional agent selected from one or more of a chemotherapeutic agent, hormonal therapeutic agent, and kinase inhibitor by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to the additional agent alone. In some embodiments, the at least one anti-NRP2 antibody enhances an anti-tumor and/or immunostimulatory activity of the cancer immunotherapy agent by about, or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more, relative to the cancer immunotherapy agent alone.

Certain embodiments comprise administering the at least one anti-NRP2 antibody in an amount and at a frequency sufficient to achieve a steady state concentration, or average circulating concentration, of the at least one anti-NRP2 antibody of between about 1 nM and about 1 μ M, between

about 1 nM and about 100 nM, between about 1 nM and about 10 nM, or between about 1 nM and about 3 μ M.

Also included are patient care kits, comprising:

- (a) at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin-2 (NRP2) polypeptide; and optionally
- (b) at least one additional agent selected from a cancer immunotherapy agent, a chemotherapeutic agent, a hormonal therapeutic agent, and a kinase inhibitor.

In some embodiments, (a) and (b) are in separate therapeutic compositions. In some embodiments, (a) and (b) are in the same therapeutic composition. In some embodiments, the at least one chemotherapeutic agent is selected from one or more of an alkylating agent, an anti-metabolite, a cytotoxic antibiotic, a topoisomerase inhibitor (type I or type II), and an anti-microtubule agent.

In some embodiments:

the alkylating agent is selected from one or more of nitrogen mustards (optionally mechlorethamine, cyclophosphamide, mustine, melphalan, chlorambucil, ifosfamide, and busulfan), nitrosoureas (optionally N-Nitroso-N-methylurea (MNU), carmustine (BCNU), lomustine (CCNU), semustine (MeCCNU), fotemustine, and streptozotocin), tetrazines (optionally dacarbazine, mitozolomide, and temozolomide), aziridines (optionally thiotepea, mytomycin, and diaziquone (AZQ)), cisplatin and derivatives thereof (optionally carboplatin and oxaliplatin), and non-classical alkylating agents (optionally procarbazine and hexamethylmelamine);

the anti-metabolite is selected from one or more of anti-folates (optionally methotrexate and pemetrexed), fluoropyrimidines (optionally 5-fluorouracil and capecitabine), deoxynucleoside analogues (optionally ancitabine, enocitabine, cytarabine, gemcitabine, decitabine, azacitidine, fludarabine, nelarabine, cladribine, clofarabine, fludarabine, and pentostatin), and thiopurines (optionally thioguanine and mercaptopurine);

the cytotoxic antibiotic is selected from one or more of anthracyclines (optionally doxorubicin, daunorubicin, epirubicin, idarubicin, pirarubicin, aclarubicin, and mitoxantrone), bleomycins, mitomycin C, mitoxantrone, and actinomycin;

the topoisomerase inhibitor is selected from one or more of camptothecin, irinotecan, topotecan, etoposide, doxorubicin, mitoxantrone, teniposide, novobiocin, merbarone, and aclarubicin; and/or

the anti-microtubule agent is selected from one or more of taxanes (optionally paclitaxel and docetaxel) and vinca alkaloids (optionally vinblastine, vincristine, vindesine, vinorelbine).

In some embodiments, the at least one hormonal therapeutic agent is a hormonal agonist or a hormonal antagonist. In some embodiments, the hormonal agonist is selected from one or more of a progestogen (progesterone), a corticosteroid (optionally prednisolone, methylprednisolone, or dexamethasone), insulin like growth factors, VEGF derived angiogenic and lymphangiogenic factors (optionally VEGF-A, VEGF-A145, VEGF-A165, VEGF-C, VEGF-D, PlGF-2), fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor (PDGF), transforming growth factor (TGF)- β , an androgen, an estrogen, and a somatostatin analog. In some embodiments, the hormonal antagonist is selected from one or more of a hormone synthesis inhibitor, optionally an aromatase inhibitor or a gonadotropin-releasing hormone (GnRH) or an analog thereof, and a hormone receptor antagonist, option-

ally a selective estrogen receptor modulator (SERM) or an anti-androgen, or an antibody directed against a hormonal receptor, optionally cixutumumab, dalotuzumab, figitumumab, ganitumab, istratuzumab, robatumumab, alacizumab pegol, bevacizumab, icrucumab, ramucirumab, fresolimumab, metelimumab, naxitamab, cetuximab, depatuxizumab mafodotin, futuximab, imgatuzumab, laprituximab emtansine, matuzumab, modotuximab, necitumumab, nimotuzumab, panitumumab, tomuzotuximab, zalutumumab, aprutumab ixadotin, bemarituzumab, olaratumab, or tovetumab.

In some embodiments, the kinase inhibitor is selected from one or more of adavosertib, afanitib, aflibercept, axitinib, bevacizumab, bosutinib, cabozantinib, cetuximab, cobimetinib, crizotinib, dasatinib, entrectinib, erdafitinib, erlotinib, fostamitinib, gefitinib, ibrutinib, imatinib, lapatinib, lenvatinib, mubritinib, nilotinib, panitumumab, pazopanib, pegaptanib, ponatinib, ranibizumab, regorafenib, ruxolitinib, sorafenib, sunitinib, SU6656, tofacitinib, trastuzumab, vandetanib, and vemurafenib.

Also included are bioassay systems, comprising a substantially pure anti-NRP2 antibody or antigen-binding fragment thereof, optionally as defined herein, and a host cell line that expresses a human NRP2 polypeptide on the cell surface. In some embodiments, the NRP2 polypeptide is labeled with a detectable label. In some embodiments, the anti-NRP2 antibody is labeled with a detectable label. In some embodiments, the NRP2 polypeptide is functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator of biological activity of the NRP2 polypeptide. In some embodiments, the NRP2 polypeptide is selected from Table N1. Certain bioassay systems comprise at least one NRP2 ligand (optionally an NRP2 ligand selected from Table N2 or Table N3 and/or a human histidyl-tRNA synthetase (HRS) polypeptide selected from Table H1), optionally wherein the host cell expresses the at least one NRP2 ligand. In some embodiments, the HRS polypeptide is selected from Table H1, optionally wherein the HRS polypeptide comprises a HRS splice variant, optionally selected from HisRS^{N1}, HisRS^{N2}, HisRS^{N3}, HisRS^{N4}, HisRS^{N5}, HisRS^{C1}, HisRS^{C2}, HisRS^{C3}, HisRS^{C4}, HisRS^{C5}, HisRS^{C6}, HisRS^{C7}, HisRS^{C8}, and HisRS^{C9}. In some embodiments, the at least one NRP2 ligand is selected from Table N2 or Table N3.

Also included are detection systems, comprising a cell that expresses a human neuropilin 2 (NRP2) polypeptide, at least one NRP2 ligand (optionally a recombinant NRP2 ligand selected from Table N2 or Table N3 and/or a human histidyl-tRNA synthetase (HRS) polypeptide selected from Table H1), and a human or humanized anti-NRP2 antibody or antigen-binding fragment thereof, optionally as defined herein, which modulates the interaction between the NRP2 polypeptide and the at least one NRP2 ligand. In some embodiments, the anti-NRP2 antibody is labeled with a detectable label. In some embodiments, the NRP2 polypeptide is selected from Table N1. In some embodiments, the HRS polypeptide comprises a HRS splice variant selected from Table H1, optionally selected from HisRS^{N1}, HisRS^{N2}, HisRS^{N3}, HisRS^{N4}, HisRS^{N5}, HisRS^{C1}, HisRS^{C2}, HisRS^{C3}, HisRS^{C4}, HisRS^{C5}, HisRS^{C6}, HisRS^{C7}, HisRS^{C8}, and HisRS^{C9}. In some embodiments, the at least one NRP2 ligand is selected from Table N2 or Table N3. In some embodiments, the NRP2 polypeptide and/or the at least one NRP2 ligand is/are functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator of biological activity of the NRP2 polypeptide or the at least one NRP2 ligand.

Also included are diagnostic systems, comprising a cell that comprises a neuropilin 2 (NRP2) polypeptide, and at least one NRP2 ligand that specifically binds to the NRP2 polypeptide (optionally an NRP2 ligand selected from Table N2 or Table N3 and/or a human histidyl-tRNA synthetase (HRS) polypeptide selected from Table H1), wherein the cell comprises an indicator molecule that indicates a change in the levels or activity of the NRP2 polypeptide in response to interaction with the at least one NRP2 ligand.

Also included are cellular compositions, comprising an engineered population of cells in which at least one cell comprises one or more polynucleotides encoding a human or humanized anti-NRP2 antibody or antigen-binding fragment thereof, as defined herein, wherein the cells are capable of growing in a serum-free medium.

Also included are cellular growth devices, comprising a human or humanized anti-NRP2 antibody or antigen-binding fragment thereof, as defined herein, an engineered population of cells in which at least one cell comprises one or more polynucleotides encoding said anti-NRP2 antibody or antigen-binding fragment thereof, at least about 10 liters of a serum-free growth medium, and a sterile container.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B illustrate the general domain structure of neuropilins (1A) and exemplary neuropilin co-receptor functions (1B).

FIGS. 2A-2B illustrate the domain structure of NRP2 isoforms and exemplary NRP2 ligand binding domains. All NRP2 isoforms are identical through the MAM domain, after this, is a short possible retained intron that codes for GENFK (SEQ ID NO: 153). There are then two possible splicing frames (forms a and b), these code different juxtamembrane domains, transmembrane helixes, and cytoplasmic domains. Within the first there is an alternative splice acceptor that removes 17 amino acids (variant 3, or form c). Each form has different spacing from the membrane, which may affect co-receptor specificity. The A/C and B forms, also have different transmembrane domains, where the A/C forms contain a dimerization motif (GXXXG).

FIG. 3 shows the binding of anti-NRP2 antibodies to clonal Expi293 cells expressing human NRP2. The four anti-NRP2 antibodies, and isotype control antibody (human IgG4) were added to human NRP2 expressing cells at the concentrations shown, and cell binding determined by FACS analysis as described in the examples.

FIG. 4 shows the binding of anti-NRP2 antibodies to clonal Expi293 cells expressing cynomolgus monkey NRP2. The four anti-NRP2 antibodies, and isotype control antibody (human IgG4) were added to cynomolgus monkey NRP2 expressing cells at the concentrations shown, and cell binding determined by FACS analysis as described in the examples.

FIGS. 5A-5B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with KDR in the presence and absence of VEGF-A. The indicated antibodies were assessed in a receptor dimerization assay as described in the examples. 5A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 5B shows the net effect at 80 minutes

FIGS. 6A-6B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with FLT4 in the presence and absence of VEGF-C. The indicated antibodies were assessed

in a receptor dimerization assay as described in the examples. 6A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 6B shows the net effect at 80 minutes

FIGS. 7A-7B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with PLXNA1 in the presence and absence of SEMA 3F. The indicated antibodies were assessed in a receptor dimerization assay as described in the examples. 7A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 7B shows the net effect at 80 minutes

FIGS. 8A-8B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with KDR in the presence and absence of VEGF-A. The indicated antibodies were assessed in a receptor dimerization assay as described in the examples. 8A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 8B shows the net effect at 80 minutes.

FIGS. 9A-9B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with FLT4 in the presence and absence of VEGF-C. The indicated antibodies were assessed in a receptor dimerization assay as described in the examples. 9A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 9B shows the net effect at 80 minutes.

FIGS. 10A-10B show the effects of anti-NRP2 antibodies on NRP2 receptor dimerization with PLXN A1 in the presence and absence of SEMA 3F. The indicated antibodies were assessed in a receptor dimerization assay as described in the examples. 10A shows normalized responses for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. 10B shows the net effect at 80 minutes

FIGS. 11A-11B show the measurement of antibody binding to Expi293 cells over-expressing NRP2. 11A shows an example of MFI profiles stained with mouse/human reactive control antibody where untransfected cell histogram is shown in light grey, while cells over-expressing mouse NRP2 I383V variant are shown in dark grey. Gating for NRP2 over-expressing cells is shown. 11B shows a chart of MFI staining of aNRP2-10v10 at 10 nM of human and mouse wild type NRP2, and mouse NRP2 receptors containing individual mutations representing human residues.

FIG. 12 shows the structure of the human NRP2 b1 domain from PBD file 2QQK. A ribbon diagram showing amino acids 280-426 is displayed. Side chains of residues 299Y, 354N, 416S, and 319T are highlighted in white. Distances between the alpha carbon of 299Y, 354N, and 416S in angstroms are shown alongside a dotted line.

FIGS. 13A-13B shows the inhibitory effect of the anti-NRP2 antibody on anchorage-independent growth and sensitivity to chemotherapeutic agents in triple negative breast cancer cells (TNBC) cells in 3D soft agar colony formation

assays. The fluorescence readings of the colony formation assay are shown as dot plots with mean $\pm$ SEM, for TNBC cells MDA-MB-231 (13A) or BT549 (13B) treated with a-NRP2-10v10 vs the isotype control hIgG4 at 100 nM in combination with chemo drugs-Cisplatin or 5-FU. The statistical significance is indicated by asterisks (* <0.05 , ** <0.01 , **** <0.0001 by Student's t test). In both cases, the anti-tumor effects of the anti-NRP2 antibody are significantly more pronounced in combination with cisplatin or 5-FU.

FIG. 14 shows the inhibitory effect of the anti-NRP2 antibody in combination with chemotherapeutic agents and a VEGF-A antibody drug (Bevacizumab) on anchorage-independent growth of TNBC MDA-MB-231 cells in 3D methylcellulose colony formation assays. The luminescence readings of the colony formation assay are shown as dot plots with mean $\pm$ SEM, for TNBC cells MDA-MB-231 treated with aNRP2-10v10 vs the isotype control hIgG4 at 100 nM in combination with chemo drugs-Cisplatin (at 2 doses) or Bevacizumab (at 100 nM). The statistical significance is indicated by asterisks (** $p<0.01$, *** $p<0.001$, **** $p<0.0001$ by Student's t test).

FIG. 15 shows the inhibition of human lymphatic endothelial cell (HLEC) migration toward vascular endothelial growth factor C (VEGF-C) by antibodies aNRP2-10v10 and aNRP2-11v7. HLECs were inhibited by both NRP2 antibodies to levels consistent with positive control antibodies (α KDR or 3C5) directed against VEGFR2 or VEGFR3 respectively.

FIG. 16 shows that treatment of cells with the antibodies aNRP2-14v10 (NRP2_14) or aNRP2-11v7 (NRP2_11) blocked the reduction in phospho-Akt levels upon Semaphorin 3F treatment in U251 glioblastoma cells. Treatment of U251 cells with Semaphorin 3F results in an approximately 50% decrease in intracellular phospho-Akt levels that can be blocked by both NRP2_11 and NRP2_14, but is not blocked by VEGF-C blocking antibody aNRP2-10v10 or by mIgG1 and hIgG4 control antibodies.

FIG. 17 shows a schematic of Semaphorin 3F signaling via NRP2. Treatment with Semaphorin 3F results in blocking PI3K activity and thereby reducing phospho-Akt levels.

FIG. 18 shows that antibodies aNRP2-14v10 and aNRP2-11v7, but not aNRP2-10v10, block semaphorin 3F mediated inhibition of AKT phosphorylation. Treatment of U251 cells with Semaphorin 3F resulted in an approximately 50% decrease in intracellular phospho-Akt levels that can be blocked by both the semaphorin blocking antibodies aNRP2-11v7 and aNRP2-14v10, but is not blocked by the VEGF-C blocking antibody aNRP2-10v10 or with mIgG1 and hIgG4 control antibodies.

FIGS. 19A-19B show that the mouse surrogate VEGF Blocking antibody aNRP2-28 inhibits Tumor Growth in a Murine Melanoma Model (B16.F10). Animals bearing B16-F10 tumors and treated with three doses of the mouse surrogate antibody aNRP2-28, which recognizes mouse NRP2 (and is a functional surrogate for anti-human NRP2 antibody aNRP2-10v10). FIG. 19A shows that treatment with the mouse surrogate antibody aNRP2-28 (grey triangles), showed tumor growth inhibition compared to the IgG control group (black squares), and Bevacizumab (black triangles) reaching statistical significance ($p<0.05$) on Day 14, 16, and 19. FIG. 19B shows terminal tumor weights for each treatment group.

FIG. 20A-20B show the anti-metastatic effects of VEGF-blocker (aNRP2-28) in a model of spontaneous metastasis (4T1). FIG. 20A shows that animals bearing 4T1 tumors and treated with mouse surrogate antibody aNRP2-28 showed a

reduction in metastatic nodules in the lung at termination, reaching statistical significance ($p<0.05$) compared to the IgG control (black bar graph). FIG. 20B shows serum antibody concentration versus the lung of lung metastases, demonstrating a linear correlation between terminal antibody concentration and number of metastatic nodules.

FIG. 21 shows the synergistic effect of VEGF-Blockers (aNRP2-10v10/aNRP2-28) in combination with cisplatin in the TNBC Xenograft Model (MDA-MB-231). The addition of pooled anti-NRP2 antibodies (black symbols) to the cisplatin treatment regimen increased the tumor-inhibitory effect of cisplatin starting on day 40 post inoculation, reaching statistical significance on day 60 compared to the control animals (grey symbols).

FIG. 22 shows that the VEGF Blocker (aNRP2-10v10) enhances activity of the chemotherapeutic drug 5-FU in the NSCLC Xenograft model (A549). The combination of 5-FU and aNRP2-10 (black symbols) performed better than 5-FU with control IgG (grey symbols), evidencing that targeting NRP2 increases the efficacy of the chemotherapeutic drug 5-FU in an art-accepted model of NSCLC. A statistically significant difference was reached on day 51 post cell inoculation.

FIGS. 23A-23B show that the VEGF Blocker (aNRP2-28) inhibits neo-lymphangiogenesis in an animal model of corneal injury. FIG. 23A shows representative image scans of Lyve-1 (lymphangiogenic marker) stained corneas of the IgG control and aNRP2-28 group. Plotting the mean values of % Lyve-1+ area for each treatment group. FIG. 23B shows a clear trend towards reduced sprouting of lymphatic vessels in the aNRP2-10 treatment groups as compared to the IgG control group.

DETAILED DESCRIPTION

Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which the disclosure belongs. Although any methods, materials, compositions, reagents, cells, similar or equivalent similar or equivalent to those described herein can be used in the practice or testing of the subject matter of the present disclosure, preferred methods and materials are described. All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

Standard techniques may be used for recombinant DNA, oligonucleotide synthesis, and tissue culture and transformation (e.g., electroporation, lipofection). Enzymatic reactions and purification techniques may be performed according to manufacturer's specifications or as commonly accomplished in the art or as described herein. These and related techniques and procedures may be generally performed according to conventional methods well known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification. Unless specific definitions are provided, the nomenclature utilized in connection with, and the laboratory procedures and techniques of, molecular biology, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein

are those well-known and commonly used in the art. Standard techniques may be used for recombinant technology, molecular biological, microbiological, chemical syntheses, chemical analyses, pharmaceutical preparation, formulation, and delivery, and treatment of patients.

For the purposes of the present disclosure, the following terms are defined below.

The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” includes “one element”, “one or more elements” and/or “at least one element”.

By “about” is meant a quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length that varies by as much as 30, 25, 20, 15, 10, 9, 8, 7, 6, 5, 4, 3, 2 or 1% to a reference quantity, level, value, number, frequency, percentage, dimension, size, amount, weight or length.

The term “antigen” refers to a molecule or a portion of a molecule capable of being bound by a selective binding agent, such as an antibody, and additionally capable of being used in an animal to produce antibodies capable of binding to an epitope of that antigen. An antigen may have one or more epitopes. As used herein, the term “antigen” includes substances that are capable, under appropriate conditions, of inducing an immune response to the substance and of reacting with the products of the immune response. For example, an antigen can be recognized by antibodies (humoral immune response) or sensitized T-lymphocytes (T helper or cell-mediated immune response), or both. Antigens can be soluble substances, such as toxins and foreign proteins, or particulates, such as bacteria and tissue cells; however, only the portion of the protein or polysaccharide molecule known as the antigenic determinant (epitopes) combines with the antibody or a specific receptor on a lymphocyte. More broadly, the term “antigen” includes any substance to which an antibody binds, or for which antibodies are desired, regardless of whether the substance is immunogenic. For such antigens, antibodies can be identified by recombinant methods, independently of any immune response.

An “antagonist” refers to biological structure or chemical agent that interferes with or otherwise reduces the physiological action of another agent or molecule. In some instances, the antagonist specifically binds to the other agent or molecule. Included are full and partial antagonists.

An “agonist” refers to biological structure or chemical agent that increases or enhances the physiological action of another agent or molecule. In some instances, the agonist specifically binds to the other agent or molecule. Included are full and partial agonists.

The term “anergy” refers to the functional inactivation of a T-cell, or B-cell response to re-stimulation by antigen.

As used herein, the term “amino acid” is intended to mean both naturally occurring and non-naturally occurring amino acids as well as amino acid analogs and mimetics. Naturally-occurring amino acids include the 20 (L)-amino acids utilized during protein biosynthesis as well as others such as 4-hydroxyproline, hydroxylysine, desmosine, isodesmosine, homocysteine, citrulline and ornithine, for example. Non-naturally occurring amino acids include, for example, (D)-amino acids, norleucine, norvaline, p-fluorophenylalanine, ethionine and the like, which are known to a person skilled in the art. Amino acid analogs include modified forms of naturally and non-naturally occurring amino acids. Such modifications can include, for example, substitution or replacement of chemical groups and moieties on the amino

acid or by derivatization of the amino acid. Amino acid mimetics include, for example, organic structures which exhibit functionally similar properties such as charge and charge spacing characteristic of the reference amino acid.

For example, an organic structure which mimics arginine (Arg or R) would have a positive charge moiety located in similar molecular space and having the same degree of mobility as the ϵ -amino group of the side chain of the naturally occurring Arg amino acid. Mimetics also include constrained structures so as to maintain optimal spacing and charge interactions of the amino acid or of the amino acid functional groups. Those skilled in the art know or can determine what structures constitute functionally equivalent amino acid analogs and amino acid mimetics.

As used herein, the term “antibody” encompasses not only intact polyclonal or monoclonal antibodies, but also fragments thereof (such as dAb, Fab, Fab', F(ab')₂, Fv), single chain (ScFv), synthetic variants thereof, naturally occurring variants, fusion proteins comprising an antibody portion with an antigen-binding fragment of the required specificity, humanized antibodies, chimeric antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen-binding site or fragment (epitope recognition site) of the required specificity. Certain features and characteristics of antibodies (and antigen-binding fragments thereof) are described in greater detail herein.

An antibody or antigen-binding fragment can be of essentially any type. As is well known in the art, an antibody is an immunoglobulin molecule capable of specific binding to a target, such as an immune checkpoint molecule, through at least one epitope recognition site, located in the variable region of the immunoglobulin molecule.

The term “antigen-binding fragment” as used herein refers to a polypeptide fragment that contains at least one CDR of an immunoglobulin heavy and/or light chain that binds to the antigen of interest. In this regard, an antigen-binding fragment of the herein described antibodies may comprise 1, 2, 3, 4, 5, or all 6 CDRs of a V_H and V_L sequence from antibodies that bind to a target molecule.

The binding properties of antibodies and antigen-binding fragments thereof can be quantified using methods well known in the art (see Davies et al., Annual Rev. Biochem. 59:439-473, 1990). In some embodiments, an antibody or antigen-binding fragment thereof specifically binds to a target molecule, for example, an NRP2 polypeptide or an epitope or complex thereof, with an equilibrium dissociation constant that is about or ranges from about $\leq 10^{-7}$ M to about 10^{-8} M. In some embodiments, the equilibrium dissociation constant is about or ranges from about $\leq 10^{-9}$ M to about $\leq 10^{-10}$ M. In certain illustrative embodiments, an antibody or antigen-binding fragment thereof has an affinity (K_d or EC₅₀) for a target molecule (to which it specifically binds) of about, at least about, or less than about, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM.

A molecule such as a polypeptide or antibody is said to exhibit “specific binding” or “preferential binding” if it reacts or associates more frequently, more rapidly, with greater duration and/or with greater affinity with a particular cell, substance, or particular epitope than it does with alternative cells or substances, or epitopes. An antibody “specifically binds” or “preferentially binds” to a target molecule or epitope if it binds with greater affinity, avidity, more readily, and/or with greater duration than it binds to other substances or epitopes, for example, by a statistically significant amount. Typically one member of the pair of

molecules that exhibit specific binding has an area on its surface, or a cavity, which specifically binds to and is therefore complementary to a particular spatial and/or polar organization of the other member of the pair of molecules. Thus, the members of the pair have the property of binding specifically to each other. For instance, an antibody that specifically or preferentially binds to a specific epitope is an antibody that binds that specific epitope with greater affinity, avidity, more readily, and/or with greater duration than it binds to other epitopes. It is also understood by reading this definition that, for example, an antibody (or moiety or epitope) that specifically or preferentially binds to a first target may or may not specifically or preferentially bind to a second target. The term is also applicable where, for example, an antibody is specific for a particular epitope which is carried by a number of antigens, in which case the specific binding member carrying the antigen-binding fragment or domain will be able to bind to the various antigens carrying the epitope; for example, it may be cross reactive to a number of different forms of a target antigen from multiple species that share a common epitope.

Immunological binding generally refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific, for example by way of illustration and not limitation, as a result of electrostatic, ionic, hydrophilic and/or hydrophobic attractions or repulsion, steric forces, hydrogen bonding, van der Waals forces, and other interactions. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (Kd) of the interaction, wherein a smaller Kd represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and on geometric parameters that equally influence the rate in both directions. Thus, both the "on rate constant" (Kon) and the "off rate constant" (Koff) can be determined by calculation of the concentrations and the actual rates of association and dissociation. The ratio of Koff/Kon enables cancellation of all parameters not related to affinity, and is thus equal to the dissociation constant Kd. As used herein, the term "affinity" includes the equilibrium constant for the reversible binding of two agents and is expressed as Kd or EC₅₀. Affinity of a binding protein to a ligand such as affinity of an antibody for an epitope can be, for example, from about 100 nanomolar (nM) to about 0.1 nM, from about 100 nM to about 1 picomolar (pM), or from about 100 nM to about 1 femtomolar (fM). As used herein, the term "avidity" refers to the resistance of a complex of two or more agents to dissociation after dilution. In some embodiments, affinity is expressed in terms of the half maximal effective concentration (EC₅₀), which refers to the concentration of an agent, such as an antibody, or an anti-NRP2 antibody, as disclosed herein, which induces a response halfway between the baseline and maximum after a specified exposure time. The EC₅₀ is commonly used as a measure of an antibody's potency.

Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. See, e.g., Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. Monoclonal antibodies specific for a polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Also

included are methods that utilize transgenic animals such as mice to express human antibodies. See, e.g., Neuberger et al., *Nature Biotechnology* 14:826, 1996; Lonberg et al., *Handbook of Experimental Pharmacology* 113:49-101, 1994; and Lonberg et al., *Internal Review of Immunology* 13:65-93, 1995. Particular examples include the VELOCIMMUNE® platform by REGENEREX® (see, e.g., U.S. Pat. No. 6,596,541).

Antibodies can also be generated or identified by the use of phage display or yeast display libraries (see, e.g., U.S. Pat. No. 7,244,592; Chao et al., *Nature Protocols* 1:755-768, 2006). Non-limiting examples of available libraries include cloned or synthetic libraries, such as the Human Combinatorial Antibody Library (HuCAL), in which the structural diversity of the human antibody repertoire is represented by seven heavy chain and seven light chain variable region genes. The combination of these genes gives rise to 49 frameworks in the master library. By superimposing highly variable genetic cassettes (CDRs=complementarity determining regions) on these frameworks, the vast human antibody repertoire can be reproduced. Also included are human libraries designed with human-donor-sourced fragments encoding a light-chain variable region, a heavy-chain CDR-3, synthetic DNA encoding diversity in heavy-chain CDR-1, and synthetic DNA encoding diversity in heavy-chain CDR-2. Other libraries suitable for use will be apparent to persons skilled in the art.

In certain embodiments, antibodies and antigen-binding fragments thereof as described herein include a heavy chain and a light chain CDR set, respectively interposed between a heavy chain and a light chain framework region (FR) set which provide support to the CDRs and define the spatial relationship of the CDRs relative to each other. As used herein, the term "CDR set" refers to the three hypervariable regions of a heavy or light chain V region. Proceeding from the N-terminus of a heavy or light chain, these regions are denoted as "CDR1," "CDR2," and "CDR3" respectively. An antigen-binding site, therefore, includes six CDRs, comprising the CDR set from each of a heavy and a light chain V region. A polypeptide comprising a single CDR, (e.g., a CDR1, CDR2 or CDR3) is referred to herein as a "molecular recognition unit." Crystallographic analysis of a number of antigen-antibody complexes has demonstrated that the amino acid residues of CDRs form extensive contact with bound antigen, wherein the most extensive antigen contact is with the heavy chain CDR3. Thus, the molecular recognition units are primarily responsible for the specificity of an antigen-binding site.

As used herein, the term "FR set" refers to the four flanking amino acid sequences which frame the CDRs of a CDR set of a heavy or light chain V region. Some FR residues may contact bound antigen; however, FRs are primarily responsible for folding the V region into the antigen-binding site, particularly the FR residues directly adjacent to the CDRs. Within FRs, certain amino residues and certain structural features are very highly conserved. In this regard, all V region sequences contain an internal disulfide loop of around 90 amino acid residues. When the V regions fold into a binding-site, the CDRs are displayed as projecting loop motifs which form an antigen-binding surface. It is generally recognized that there are conserved structural regions of FRs which influence the folded shape of the CDR loops into certain "canonical" structures-regardless of the precise CDR amino acid sequence. Further, certain FR

residues are known to participate in non-covalent interdomain contacts which stabilize the interaction of the antibody heavy and light chains.

The structures and locations of immunoglobulin variable domains may be determined by reference to Kabat, E. A. et al., Sequences of Proteins of Immunological Interest. 4th Edition. US Department of Health and Human Services. 1987, and updates thereof.

Also include are "monoclonal" antibodies, which refer to a homogeneous antibody population wherein the monoclonal antibody is comprised of amino acids (naturally occurring and non-naturally occurring) that are involved in the selective binding of an epitope. Monoclonal antibodies are highly specific, being directed against a single epitope. The term "monoclonal antibody" encompasses not only intact monoclonal antibodies and full-length monoclonal antibodies, but also fragments thereof (such as Fab, Fab', F(ab')₂, Fv), single chain (scFv), variants thereof, fusion proteins comprising an antigen-binding portion, humanized monoclonal antibodies, chimeric monoclonal antibodies, and any other modified configuration of the immunoglobulin molecule that comprises an antigen-binding fragment (epitope recognition site) of the required specificity and the ability to bind to an epitope. It is not intended to be limited as regards the source of the antibody or the manner in which it is made (e.g., by hybridoma, phage selection, recombinant expression, transgenic animals). The term includes whole immunoglobulins as well as the fragments etc. described above under the definition of "antibody."

The proteolytic enzyme papain preferentially cleaves IgG molecules to yield several fragments, two of which (the F(ab) fragments) each comprise a covalent heterodimer that includes an intact antigen-binding site. The enzyme pepsin is able to cleave IgG molecules to provide several fragments, including the F(ab')₂ fragment which comprises both antigen-binding sites. An Fv fragment for use according to certain embodiments can be produced by preferential proteolytic cleavage of an IgM, and on rare occasions of an IgG or IgA immunoglobulin molecule. Fv fragments are, however, more commonly derived using recombinant techniques known in the art. The Fv fragment includes a non-covalent VH:VL heterodimer including an antigen-binding site which retains much of the antigen recognition and binding capabilities of the native antibody molecule. See Inbar et al., PNAS USA. 69:2659-2662, 1972; Hochman et al., Biochem. 15:2706-2710, 1976; and Ehrlich et al., Biochem. 19:4091-4096, 1980.

In certain embodiments, single chain Fv (scFv) antibodies are contemplated. For example, Kappa bodies (III et al., Prot. Eng. 10:949-57, 1997); minibodies (Martin et al., EMBO J 13:5305-9, 1994); diabodies (Holliger et al., PNAS 90:6444-8, 1993); or Janusins (Traunecker et al., EMBO J 10:3655-59, 1991; and Traunecker et al., Int. J. Cancer Suppl. 7:51-52, 1992), may be prepared using standard molecular biology techniques following the teachings of the present application with regard to selecting antibodies having the desired specificity.

A single chain Fv (scFv) polypeptide is a covalently linked VH:VL heterodimer which is expressed from a gene fusion including VH- and VL-encoding genes linked by a peptide-encoding linker. Huston et al. (PNAS USA. 85 (16): 5879-5883, 1988). A number of methods have been described to discern chemical structures for converting the naturally aggregated—but chemically separated-light and heavy polypeptide chains from an antibody V region into an scFv molecule which will fold into a three dimensional structure substantially similar to the structure of an antigen-

binding site. See, e.g., U.S. Pat. Nos. 5,091,513 and 5,132,405, to Huston et al.; and U.S. Pat. No. 4,946,778, to Ladner et al.

In certain embodiments, the antibodies or antigen-binding fragments described herein are in the form of a "diabody." Diabodies are multimers of polypeptides, each polypeptide comprising a first domain comprising a binding region of an immunoglobulin light chain and a second domain comprising a binding region of an immunoglobulin heavy chain, the two domains being linked (e.g., by a peptide linker) but unable to associate with each other to form an antigen-binding site: antigen-binding sites are formed by the association of the first domain of one polypeptide within the multimer with the second domain of another polypeptide within the multimer (WO94/13804). A dAb fragment of an antibody consists of a VH domain (Ward et al., Nature 341:544-546, 1989). Diabodies and other multivalent or multispecific fragments can be constructed, for example, by gene fusion (see WO94/13804; and Holliger et al., PNAS USA. 90:6444-6448, 1993)).

Minibodies comprising a scFv joined to a CH3 domain are also included (see Hu et al., Cancer Res. 56:3055-3061, 1996). See also Ward et al., Nature. 341:544-546, 1989; Bird et al., Science. 242:423-426, 1988; Huston et al., PNAS USA. 85:5879-5883, 1988; PCT/US92/09965; WO94/13804; and Reiter et al., Nature Biotech. 14:1239-1245, 1996.

Where bispecific antibodies are to be used, these may be conventional bispecific antibodies, which can be manufactured in a variety of ways (Holliger and Winter, Current Opinion Biotechnol. 4:446-449, 1993), e.g., prepared chemically or from hybrid hybridomas, or may be any of the bispecific antibody fragments mentioned above. Diabodies and scFv can be constructed without an Fc region, using only variable domains, potentially reducing the effects of anti-idiotypic reaction.

Bispecific diabodies, as opposed to bispecific whole antibodies, may also be particularly useful because they can be readily constructed and expressed in *E. coli*. Diabodies (and many other polypeptides such as antibody fragments) of appropriate binding specificities can be readily selected using phage display (WO94/13804) from libraries. If one arm of the diabody is to be kept constant, for instance, with a specificity directed against antigen X, then a library can be made where the other arm is varied and an antibody of appropriate specificity selected. Bispecific whole antibodies may be made by knobs-into-holes engineering (Ridgeway et al., Protein Eng., 9:616-621, 1996).

In certain embodiments, the antibodies or antigen-binding fragments described herein are in the form of a UniBody®. A UniBody® is an IgG4 antibody with the hinge region removed (see GenMab Utrecht, The Netherlands; see also, e.g., US20090226421). This antibody technology creates a stable, smaller antibody format with an anticipated longer therapeutic window than current small antibody formats. IgG4 antibodies are considered inert and thus do not interact with the immune system. Fully human IgG4 antibodies may be modified by eliminating the hinge region of the antibody to obtain half-molecule fragments having distinct stability properties relative to the corresponding intact IgG4 (GenMab, Utrecht). Halving the IgG4 molecule leaves only one area on the UniBody® that can bind to cognate antigens (e.g., disease targets) and the UniBody® therefore binds univalently to only one site on target cells. For certain cancer cell surface antigens, this univalent binding may not stimulate the cancer cells to grow as may be seen using bivalent antibodies having the same antigen specificity, and hence

UniBody® technology may afford treatment options for some types of cancer that may be refractory to treatment with conventional antibodies. The small size of the UniBody® can be a great benefit when treating some forms of cancer, allowing for better distribution of the molecule over larger solid tumors and potentially increasing efficacy.

In certain embodiments, the antibodies and antigen-binding fragments described herein are in the form of a nanobody. Minibodies are encoded by single genes and are efficiently produced in almost all prokaryotic and eukaryotic hosts, for example, *E. coli* (see U.S. Pat. No. 6,765,087), molds (for example *Aspergillus* or *Trichoderma*) and yeast (for example *Saccharomyces*, *Kluyvermyces*, *Hansenula* or *Pichia* (see U.S. Pat. No. 6,838,254). The production process is scalable and multi-kilogram quantities of nanobodies have been produced. Nanobodies may be formulated as a ready-to-use solution having a long shelf life. The Nanoclone method (see WO 06/079372) is a proprietary method for generating Nanobodies against a desired target, based on automated high-throughput selection of B-cells.

In some embodiments, the antibodies or antigen-binding fragments described herein are in the form of an aptamer (see, e.g., Ellington et al., *Nature*. 346, 818-22, 1990; and Tuerk et al., *Science*. 249, 505-10, 1990, incorporated by reference). Examples of aptamers included nucleic acid aptamers (e.g., DNA aptamers, RNA aptamers) and peptide aptamers. Nucleic acid aptamers refer generally to nucleic acid species that have been engineered through repeated rounds of in vitro selection or equivalent method, such as SELEX (systematic evolution of ligands by exponential enrichment), to bind to various molecular targets such as small molecules, proteins, nucleic acids, and even cells, tissues and organisms. See, e.g., U.S. Pat. Nos. 6,376,190; and 6,387,620, incorporated by reference.

Peptide aptamers typically include a variable peptide loop attached at both ends to a protein scaffold, a double structural constraint that typically increases the binding affinity of the peptide aptamer to levels comparable to that of an antibody's (e.g., in the nanomolar range). In certain embodiments, the variable loop length may be composed of about 10-20 amino acids (including all integers in between), and the scaffold may include any protein that has good solubility and compacity properties. Certain exemplary embodiments utilize the bacterial protein Thioredoxin-A as a scaffold protein, the variable loop being inserted within the reducing active site (-Cys-Gly-Pro-Cys-loop in the wild protein), with the two cysteines lateral chains being able to form a disulfide bridge. Methods for identifying peptide aptamers are described, for example, in U.S. Application No. 2003/0108532, incorporated by reference. Peptide aptamer selection can be performed using different systems known in the art, including the yeast two-hybrid system.

In some embodiments, the antibodies or antigen-binding fragments described herein are in the form of an avimer. Avimers refer to multimeric binding proteins or peptides engineered using in vitro exon shuffling and phage display. Multiple binding domains are linked, resulting in greater affinity and specificity compared to single epitope immunoglobulin domains. See, e.g., Silverman et al., *Nature Biotechnology*. 23:1556-1561, 2005; U.S. Pat. No. 7,166,697; and U.S. Application Nos. 2004/0175756, 2005/0048512, 2005/0053973, 2005/0089932 and 2005/0221384, incorporated by reference.

In some embodiments, the antibodies or antigen-binding fragments described herein are in the form of an adnectin. Adnectins refer to a class of targeted biologics derived from human fibronectin, an abundant extracellular protein that

naturally binds to other proteins. See, e.g., U.S. Application Nos. 2007/0082365; 2008/0139791; and 2008/0220049, incorporated by reference. Adnectins typically consists of a natural fibronectin backbone, as well as the multiple targeting domains of a specific portion of human fibronectin. The targeting domains can be engineered to enable an adnectin to specifically recognize an NRP2 polypeptide or an epitope thereof.

In some embodiments, the antibodies or antigen-binding fragments described herein are in the form of an anticalin. Anticalins refer to a class of antibody mimetics that are typically synthesized from human lipocalins, a family of binding proteins with a hypervariable loop region supported by a structurally rigid framework. See, e.g., U.S. Application No. 2006/0058510. Anticalins typically have a size of about 20 kDa. Anticalins can be characterized by a barrel structure formed by eight antiparallel β -strands (a stable β -barrel scaffold) that are pairwise connected by four peptide loops and an attached α -helix. In certain aspects, conformational deviations to achieve specific binding are made in the hypervariable loop region(s). See, e.g., Skerra, *FEBS J.* 275:2677-83, 2008, incorporated by reference.

In some embodiments, the antibodies or antigen-binding fragments described herein are in the form of a designed ankyrin repeat protein (DARPin). DARPins include a class of non-immunoglobulin proteins that can offer advantages over antibodies for target binding in drug discovery and drug development. Among other uses, DARPins are ideally suited for in vivo imaging or delivery of toxins or other therapeutic payloads because of their favorable molecular properties, including small size and high stability. The low-cost production in bacteria and the rapid generation of many target-specific DARPins make the DARPin approach useful for drug discovery. Additionally, DARPins can be easily generated in multispecific formats, offering the potential to target an effector DARPin to a specific organ or to target multiple receptors with one molecule composed of several DARPins. See, e.g., Stumpp et al., *Curr Opin Drug Discov Devel.* 10:153-159, 2007; U.S. Application No. 2009/0082274; and PCT/EP2001/10454, incorporated by reference.

Also included are heavy chain dimers, such as antibodies from camelids and sharks. Camelid and shark antibodies comprise a homodimeric pair of two chains of V-like and C-like domains (neither has a light chain). Since the VH region of a heavy chain dimer IgG in a camelid does not have to make hydrophobic interactions with a light chain, the region in the heavy chain that normally contacts a light chain is changed to hydrophilic amino acid residues in a camelid. VH domains of heavy-chain dimer IgGs are called VHH domains. Shark Ig-NARs comprise a homodimer of one variable domain (termed a V-NAR domain) and five C-like constant domains (C-NAR domains).

In camelids, the diversity of antibody repertoire is determined by the complementary determining regions (CDR) 1, 2, and 3 in the VH or VHH regions. The CDR3 in the camel VHH region is characterized by its relatively long length averaging 16 amino acids (Muyldermans et al., 1994, *Protein Engineering* 7 (9): 1129). This is in contrast to CDR3 regions of antibodies of many other species. For example, the CDR3 of mouse VH has an average of 9 amino acids. Libraries of camelid-derived antibody variable regions, which maintain the in vivo diversity of the variable regions of a camelid, can be made by, for example, the methods disclosed in U.S. patent application No. 20050037421, published Feb. 17, 2005

In certain embodiments, the antibodies or antigen-binding fragments thereof are humanized. These embodiments refer to a chimeric molecule, generally prepared using recombinant techniques, having an antigen-binding site derived from an immunoglobulin from a non-human species and the remaining immunoglobulin structure of the molecule based upon the structure and/or sequence of a human immunoglobulin. The antigen-binding site may comprise either complete variable domains fused onto constant domains or only the CDRs grafted onto appropriate framework regions in the variable domains. Epitope binding sites may be wild type or modified by one or more amino acid substitutions. This eliminates the constant region as an immunogen in human individuals, but the possibility of an immune response to the foreign variable region remains (LoBuglio et al., PNAS USA 86:4220-4224, 1989; Queen et al., PNAS USA. 86:10029-10033, 1988; Riechmann et al., Nature. 332:323-327, 1988). Illustrative methods for humanization of antibodies include the methods described in U.S. Pat. No. 7,462,697.

Another approach focuses not only on providing human-derived constant regions, but modifying the variable regions as well so as to reshape them as closely as possible to human form. It is known that the variable regions of both heavy and light chains contain three complementarity-determining regions (CDRs) which vary in response to the epitopes in question and determine binding capability, flanked by four framework regions (FRs) which are relatively conserved in a given species and which putatively provide a scaffolding for the CDRs. When nonhuman antibodies are prepared with respect to a particular epitope, the variable regions can be "reshaped" or "humanized" by grafting CDRs derived from nonhuman antibody on the FRs present in the human antibody to be modified. Application of this approach to various antibodies has been reported by Sato et al., Cancer Res. 53:851-856, 1993; Riechmann et al., Nature 332:323-327, 1988; Verhoeven et al., Science 239:1534-1536, 1988; Kettleborough et al., Protein Engineering. 4:773-3783, 1991; Maeda et al., Human Antibodies Hybridoma 2:124-134, 1991; Gorman et al., PNAS USA. 88:4181-4185, 1991; Tempest et al., Bio/Technology 9:266-271, 1991; Co et al., PNAS USA. 88:2869-2873, 1991; Carter et al., PNAS USA. 89:4285-4289, 1992; and Co et al., J Immunol. 148:1149-1154, 1992. In some embodiments, humanized antibodies preserve all CDR sequences (for example, a humanized mouse antibody which contains all six CDRs from the mouse antibodies). In other embodiments, humanized antibodies have one or more CDRs (one, two, three, four, five, six) which are altered with respect to the original antibody, which are also termed one or more CDRs "derived from" one or more CDRs from the original antibody.

In certain embodiments, the antibodies are "chimeric" antibodies. In this regard, a chimeric antibody is comprised of an antigen-binding fragment of an antibody operably linked or otherwise fused to a heterologous Fc portion of a different antibody. In certain embodiments, the Fc domain or heterologous Fc domain is of human origin. In certain embodiments, the Fc domain or heterologous Fc domain is of mouse origin. In other embodiments, the heterologous Fc domain may be from a different Ig class from the parent antibody, including IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and IgG4), and IgM. In further embodiments, the heterologous Fc domain may be comprised of CH2 and CH3 domains from one or more of the different Ig classes. As noted above with regard to humanized antibodies, the antigen-binding fragment of a chimeric antibody may comprise

only one or more of the CDRs of the antibodies described herein (e.g., 1, 2, 3, 4, 5, or 6 CDRs of the antibodies described herein), or may comprise an entire variable domain (VL, VH or both).

As used herein, a subject "at risk" of developing a disease, or adverse reaction may or may not have detectable disease, or symptoms of disease, and may or may not have displayed detectable disease or symptoms of disease prior to the treatment methods described herein. "At risk" denotes that a subject has one or more risk factors, which are measurable parameters that correlate with development of a disease, as described herein and known in the art. A subject having one or more of these risk factors has a higher probability of developing disease, or an adverse reaction than a subject without one or more of these risk factor(s).

"Biocompatible" refers to materials or compounds which are generally not injurious to biological functions of a cell or subject and which will not result in any degree of unacceptable toxicity, including allergenic and disease states.

The term "binding" refers to a direct association between two molecules, due to, for example, covalent, electrostatic, hydrophobic, and ionic and/or hydrogen-bond interactions, including interactions such as salt bridges and water bridges.

The term "chemoresistance" refers to the change in therapeutic sensitivity of a cancer cell population over time following exposure to chemotherapy, including resistance to at least one of a cancer immunotherapy agent, a chemotherapeutic agent, a hormonal therapeutic agent, and/or a kinase inhibitor. Eventually, chemoresistance leads to the relapse and/or metastasis, of the cancer, and challenges the improvement of clinical outcome for the cancer patients. It remains the main obstacle to long term successful cancer therapy. For example, approximately 30 percent of women diagnosed with early-stage breast cancer ultimately develop resistance and eventually progress to metastatic breast cancer. The molecular mechanisms of chemoresistance include the induction of transporter pumps, oncogenes, tumor suppressor genes, mitochondrial alteration, DNA repair, autophagy, epithelial-mesenchymal transition (EMT), cancer stemness, and exosome production. These processes may operate via distinct mechanisms, alone or in combination with each other, but ultimately coordinate to prevent cell death in response to a specific targeted chemotherapeutic agent. For example, such processes provide alternative pro-growth signals and/or eliminate or otherwise reduce apoptotic pathways. Accordingly, agents that reduce chemoresistance could find utility in the treatment or reduction of chemoresistant cancers.

By "coding sequence" is meant any nucleic acid sequence that contributes to the code for the polypeptide product of a gene. By contrast, the term "non-coding sequence" refers to any nucleic acid sequence that does not directly contribute to the code for the polypeptide product of a gene.

Throughout this disclosure, unless the context requires otherwise, the words "comprise," "comprises," and "comprising" will be understood to imply the inclusion of a stated step or element or group of steps or elements but not the exclusion of any other step or element or group of steps or elements.

By "consisting of" is meant including, and limited to, whatever follows the phrase "consisting of." Thus, the phrase "consisting of" indicates that the listed elements are required or mandatory, and that no other elements may be present. By "consisting essentially of" is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed

elements. Thus, the phrase “consisting essentially of” indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be present depending upon whether or not they materially affect the activity or action of the listed elements.

The term “effector function”, or “ADCC effector function” in the context of antibodies refers to the ability of that antibody to engage with other arms of the immune system, including for example, the activation of the classical complement pathway, or through engagement of Fc receptors. Complement dependent pathways are primarily driven by the interaction of C1q with the C1 complex with clustered antibody Fc domains. Antibody dependent cellular cytotoxicity (ADCC), is primarily driven by the interaction of Fc receptors (FcRs) on the surface of effector cells (natural killer cells, macrophages, monocytes and eosinophils) which bind to the Fc region of an IgG which itself is bound to a target cell. Fc receptors (FcRs) are key immune regulatory receptors connecting the antibody mediated (humoral) immune response to cellular effector functions. Receptors for all classes of immunoglobulins have been identified, including FcγR (IgG), FcεRI (IgE), FcαRI (IgA), FcμR (IgM) and FcγRI (IgD). There are at least three classes of receptors for human IgG found on leukocytes: CD64 (FcγRI), CD32 (FcγRIIa, FcγRIIb and FcγRIIc) and CD16 (FcγRIIIa and FcγRIIIb). FcγRI is classed as a high affinity receptor (nanomolar range KD) while FcγRII and FcγRIII are low to intermediate affinity (micromolar range KD). Upon Fc binding a signaling pathway is triggered which results in the secretion of various substances, such as lytic enzymes, perforin, granzymes and tumour necrosis factor, which mediate in the destruction of the target cell. The level of ADCC effector function varies for human IgG subtypes. Although this is dependent on the allotype and specific FcγR, in simple terms ADCC effector function is “high” for human IgG1 and IgG3, and “low” for IgG2 and IgG4.

The term “endotoxin free” or “substantially endotoxin free” relates generally to compositions, solvents, and/or vessels that contain at most trace amounts (e.g., amounts having no clinically adverse physiological effects to a subject) of endotoxin, and preferably undetectable amounts of endotoxin. Endotoxins are toxins associated with certain micro-organisms, such as bacteria, typically gram-negative bacteria, although endotoxins may be found in gram-positive bacteria, such as *Listeria monocytogenes*. The most prevalent endotoxins are lipopolysaccharides (LPS) or lipooligo-saccharides (LOS) found in the outer membrane of various Gram-negative bacteria, and which represent a central pathogenic feature in the ability of these bacteria to cause disease. Small amounts of endotoxin in humans may produce fever, a lowering of the blood pressure, and activation of inflammation and coagulation, among other adverse physiological effects.

Therefore, in pharmaceutical production, it is often desirable to remove most or all traces of endotoxin from drug products and/or drug containers, because even small amounts may cause adverse effects in humans. A depyrogenation oven may be used for this purpose, as temperatures in excess of 300° C. are typically required to break down most endotoxins. For instance, based on primary packaging material such as syringes or vials, the combination of a glass temperature of 250° C. and a holding time of 30 minutes is often sufficient to achieve a 3 log reduction in endotoxin levels. Other methods of removing endotoxins are contemplated, including, for example, chromatography and filtration methods, as described herein and known in the art.

Endotoxins can be detected using routine techniques known in the art. For example, the Limulus Amoebocyte Lysate assay, which utilizes blood from the horseshoe crab, is a very sensitive assay for detecting presence of endotoxin.

In this test, very low levels of LPS can cause detectable coagulation of the limulus lysate due a powerful enzymatic cascade that amplifies this reaction. Endotoxins can also be quantitated by enzyme-linked immunosorbent assay (ELISA). To be substantially endotoxin free, endotoxin levels may be less than about 0.001, 0.005, 0.01, 0.02, 0.03, 0.04, 0.05, 0.06, 0.08, 0.09, 0.1, 0.5, 1.0, 1.5, 2, 2.5, 3, 4, 5, 6, 7, 8, 9, or 10 EU/mg of active compound. Typically, 1 ng lipopolysaccharide (LPS) corresponds to about 1-10 EU.

The term “epitope” includes any determinant, preferably a polypeptide determinant, capable of specific binding to an immunoglobulin or T-cell receptor. An epitope includes a region of an antigen that is bound by an antibody. In certain embodiments, epitope determinants include chemically active surface groupings of molecules such as amino acids, sugar side chains, phosphoryl or sulfonyl, and may in certain embodiments have specific three-dimensional structural characteristics, and/or specific charge characteristics. Epitopes can be contiguous or non-contiguous in relation to the primary structure of the antigen, for example, an NRP2 polypeptide. In particular embodiments, an epitope comprises, consists, or consists essentially of about, at least about, or no more than about 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, or 20 contiguous amino acids (i.e., a linear epitope) or non-contiguous amino acids (i.e., conformational epitope) of a reference sequence (see, e.g., Table N1) or target molecule described herein.

An “epitope” includes that portion of an antigen or other macromolecule capable of forming a binding interaction that interacts with the variable region binding pocket of a binding protein. Such binding interaction can be manifested as an intermolecular contact with one or more amino acid residues of a CDR. Antigen binding can involve a CDR3 or a CDR3 pair. An epitope can be a linear peptide sequence (i.e., “continuous”) or can be composed of noncontiguous amino acid sequences (i.e., “conformational” or “discontinuous”). A binding protein can recognize one or more amino acid sequences; therefore an epitope can define more than one distinct amino acid sequence. Epitopes recognized by binding protein can be determined by peptide mapping and sequence analysis techniques well known to one of skill in the art. A “cryptic epitope” or a “cryptic binding site” is an epitope or binding site of a protein sequence that is not exposed or substantially protected from recognition within an unmodified polypeptide, but is capable of being recognized by a binding protein of a denatured or proteolyzed polypeptide. Amino acid sequences that are not exposed, or are only partially exposed, in the unmodified polypeptide structure are potential cryptic epitopes. If an epitope is not exposed, or only partially exposed, then it is likely that it is buried within the interior of the polypeptide. Candidate cryptic epitopes can be identified, for example, by examining the three-dimensional structure of an unmodified polypeptide.

The term “half maximal effective concentration” or “EC₅₀” refers to the concentration of an agent (e.g., antibody) as described herein at which it induces a response halfway between the baseline and maximum after some specified exposure time; the EC₅₀ of a graded dose response curve therefore represents the concentration of a compound at which 50% of its maximal effect is observed. EC₅₀ also represents the plasma concentration required for obtaining 50% of a maximum effect in vivo. Similarly, the “EC₉₀”

refers to the concentration of an agent or composition at which 90% of its maximal effect is observed. The “EC₉₀” can be calculated from the “EC₅₀” and the Hill slope, or it can be determined from the data directly, using routine knowledge in the art. In some embodiments, the EC₅₀ of an agent (e.g., antibody) is less than about 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, 200 or 500 nM. In some embodiments, an agent will have an EC₅₀ value of about 1 nM or less.

“Immune response” means any immunological response originating from immune system, including responses from the cellular and humeral, innate and adaptive immune systems. Exemplary cellular immune cells include for example, lymphocytes, macrophages, T cells, B cells, NK cells, neutrophils, eosinophils, dendritic cells, mast cells, monocytes, and all subsets thereof. Cellular responses include for example, effector function, cytokine release, phagocytosis, efferocytosis, translocation, trafficking, proliferation, differentiation, activation, repression, cell-cell interactions, apoptosis, etc. Humeral responses include for example IgG, IgM, IgA, IgE, responses and their corresponding effector functions.

The “half-life” of an agent such as an antibody can refer to the time it takes for the agent to lose half of its pharmacologic, physiologic, or other activity, relative to such activity at the time of administration into the serum or tissue of an organism, or relative to any other defined time-point. “Half-life” can also refer to the time it takes for the amount or concentration of an agent to be reduced by half of a starting amount administered into the serum or tissue of an organism, relative to such amount or concentration at the time of administration into the serum or tissue of an organism, or relative to any other defined time-point. The half-life can be measured in serum and/or any one or more selected tissues.

The terms “modulating” and “altering” include “increasing,” “enhancing” or “stimulating,” as well as “decreasing” or “reducing,” typically in a statistically significant or a physiologically significant amount or degree relative to a control. An “increased,” “stimulated” or “enhanced” amount is typically a “statistically significant” amount, and may include an increase that is 1.1, 1.2, 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more times (e.g., 500, 1000 times) (including all integers and ranges in between e.g., 1.5, 1.6, 1.7, 1.8, etc.) the amount produced by no composition (e.g., the absence of agent) or a control composition. A “decreased” or “reduced” amount is typically a “statistically significant” amount, and may include a 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, 16%, 17%, 18%, 19%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% decrease (including all integers and ranges in between) in the amount produced by no composition (e.g., the absence of an agent) or a control composition. Examples of comparisons and “statistically significant” amounts are described herein.

The term “migratory cells” refers to cells that are capable of movement from one place to another in response to a stimulus. Exemplary migratory cells include immune cells such as monocytes, Natural Killer (NK) cells, dendritic cells (immature or mature), subsets of dendritic cells including myeloid, plasmacytoid (also called lymphoid) and Langerhans cells, macrophages such as histiocytes, tissue resident macrophages such as Kupffer’s cells, microglia cells in the CNS, alveolar macrophages, and peritoneal macrophages, macrophage subtypes such as M0, M1, Mox, M2a, M2b, and

M2c macrophages, neutrophils, eosinophils, mast cells, basophils, B cells including plasma B cells, memory B cells, B-1 cells, and B-2 cells, CD45RO (naive T) cells, CD45RA (memory T) cells, CD4 Helper T Cells including Th1, Th2, and Tr1/Th3 cells, CD8 Cytotoxic T Cells, Regulatory T Cells, Gamma Delta T Cells, and thymocytes. Additional examples of migratory cells include fibroblasts, fibrocytes, tumor cells, and stem cells. The term “cell migration” refers to the movement of migratory cells, and the term “modulation of cell migration” refers to the modulation of the movement of any such migratory cells.

The terms “polypeptide,” “protein” and “peptide” are used interchangeably and mean a polymer of amino acids not limited to any particular length. The term “enzyme” includes polypeptide or protein catalysts. The terms include modifications such as myristoylation, sulfation, glycosylation, phosphorylation and addition or deletion of signal sequences. The terms “polypeptide” or “protein” means one or more chains of amino acids, wherein each chain comprises amino acids covalently linked by peptide bonds, and wherein said polypeptide or protein can comprise a plurality of chains non-covalently and/or covalently linked together by peptide bonds, having the sequence of native proteins, that is, proteins produced by naturally-occurring and specifically non-recombinant cells, or genetically-engineered or recombinant cells, and comprise molecules having the amino acid sequence of the native protein, or molecules having deletions from, additions to, and/or substitutions of one or more amino acids of the native sequence. In certain embodiments, the polypeptide is a “recombinant” polypeptide, produced by recombinant cell that comprises one or more recombinant DNA molecules, which are typically made of heterologous polynucleotide sequences or combinations of polynucleotide sequences that would not otherwise be found in the cell.

The term “polynucleotide” and “nucleic acid” includes mRNA, RNA, cRNA, cDNA, and DNA. The term typically refers to polymeric form of nucleotides of at least 10 bases in length, either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The terms “isolated DNA” and “isolated polynucleotide” and “isolated nucleic acid” refer to a molecule that has been isolated free of total genomic DNA of a particular species. Therefore, an isolated DNA segment encoding a polypeptide refers to a DNA segment that contains one or more coding sequences yet is substantially isolated away from, or purified free from, total genomic DNA of the species from which the DNA segment is obtained. Also included are non-coding polynucleotides (e.g., primers, probes, oligonucleotides), which do not encode a polypeptide. Also included are recombinant vectors, including, for example, expression vectors, viral vectors, plasmids, cosmids, phagemids, phage, viruses, and the like.

Additional coding or non-coding sequences may, but need not, be present within a polynucleotide described herein, and a polynucleotide may, but need not, be linked to other molecules and/or support materials. Hence, a polynucleotide or expressible polynucleotides, regardless of the length of the coding sequence itself, may be combined with other sequences, for example, expression control sequences.

“Expression control sequences” include regulatory sequences of nucleic acids, or the corresponding amino acids, such as promoters, leaders, enhancers, introns, recognition motifs for RNA, or DNA binding proteins, polyadenylation signals, terminators, internal ribosome entry sites (IRES), secretion signals, subcellular localization sig-

nals, and the like, which have the ability to affect the transcription or translation, or subcellular, or cellular location of a coding sequence in a host cell. Exemplary expression control sequences are described in Goeddel; Gene Expression Technology: Methods in Enzymology 185, Academic Press, San Diego, Calif. (1990).

A “promoter” is a DNA regulatory region capable of binding RNA polymerase in a cell and initiating transcription of a downstream (3' direction) coding sequence. As used herein, the promoter sequence is bounded at its 3' terminus by the transcription initiation site and extends upstream (5' direction) to include the minimum number of bases or elements necessary to initiate transcription at levels detectable above background. A transcription initiation site (conveniently defined by mapping with nuclease S1) can be found within a promoter sequence, as well as protein binding domains (consensus sequences) responsible for the binding of RNA polymerase. Eukaryotic promoters can often, but not always, contain “TATA” boxes and “CAT” boxes. Prokaryotic promoters contain Shine-Dalgarno sequences in addition to the -10 and -35 consensus sequences.

A large number of promoters, including constitutive, inducible and repressible promoters, from a variety of different sources are well known in the art. Representative sources include for example, viral, mammalian, insect, plant, yeast, and bacterial cell types), and suitable promoters from these sources are readily available, or can be made synthetically, based on sequences publicly available on line or, for example, from depositories such as the ATCC as well as other commercial or individual sources. Promoters can be unidirectional (i.e., initiate transcription in one direction) or bi-directional (i.e., initiate transcription in either a 3' or 5' direction). Non-limiting examples of promoters include, for example, the T7 bacterial expression system, pBAD (araA) bacterial expression system, the cytomegalovirus (CMV) promoter, the SV40 promoter, the RSV promoter. Inducible promoters include the Tet system, (U.S. Pat. Nos. 5,464,758 and 5,814,618), the Ecdysone inducible system (No et al., Proc. Natl. Acad. Sci. (1996) 93 (8): 3346-3351; the T-REx™ system (Invitrogen Carlsbad, CA), LacSwitch® (Stratagene, (San Diego, CA) and the Cre-ERT tamoxifen inducible recombinase system (Indra et al. Nuc. Acid. Res. (1999) 27 (22): 4324-4327; Nuc. Acid. Res. (2000) 28 (23): e99; U.S. Pat. No. 7,112,715; and Kramer & Fussenegger, Methods Mol. Biol. (2005) 308:123-144) or any promoter known in the art suitable for expression in the desired cells.

An “expressible polynucleotide” includes a cDNA, RNA, mRNA or other polynucleotide that comprises at least one coding sequence and optionally at least one expression control sequence, for example, a transcriptional and/or translational regulatory element, and which can express an encoded polypeptide upon introduction into a cell, for example, a cell in a subject.

Various viral vectors that can be utilized to deliver an expressible polynucleotide include adenoviral vectors, herpes virus vectors, vaccinia virus vectors, adeno-associated virus (AAV) vectors, and retroviral vectors. In some instances, the retroviral vector is a derivative of a murine or avian retrovirus, or is a lentiviral vector. Examples of retroviral vectors in which a single foreign gene can be inserted include, but are not limited to: Moloney murine leukemia virus (MoMuLV), Harvey murine sarcoma virus (HaMuSV), murine mammary tumor virus (MuMTV), SIV, BIV, HIV and Rous Sarcoma Virus (RSV). A number of additional retroviral vectors can incorporate multiple genes. All of these vectors can transfer or incorporate a gene for a selectable marker so that transduced cells can be identified

and generated. By inserting a polypeptide sequence of interest into the viral vector, along with another gene that encodes the ligand for a receptor on a specific target cell, for example, the vector may be made target specific. Retroviral vectors can be made target specific by inserting, for example, a polynucleotide encoding a protein. Illustrative targeting may be accomplished by using an antibody to target the retroviral vector. Those of skill in the art will know of, or can readily ascertain without undue experimentation, specific polynucleotide sequences which can be inserted into the retroviral genome to allow target specific delivery of the retroviral vector.

In particular embodiments, the expressible polynucleotide is a modified RNA or modified mRNA polynucleotide, for example, a non-naturally occurring RNA analog. In certain embodiments, the modified RNA or mRNA polypeptide comprises one or more modified or non-natural bases, for example, a nucleotide base other than adenine (A), guanine (G), cytosine (C), thymine (T), and/or uracil (U). In some embodiments, the modified mRNA comprises one or more modified or non-natural internucleotide linkages. Expressible RNA polynucleotides for delivering an encoded therapeutic polypeptide are described, for example, in Kormann et al., Nat Biotechnol. 29:154-7, 2011; and U.S. Application Nos. 2015/0111248; 2014/0243399; 2014/0147454; and 2013/0245104, which are incorporated by reference in their entireties.

The term “isolated” polypeptide or protein referred to herein means that a subject protein (1) is free of at least some other proteins with which it would typically be found in nature, (2) is essentially free of other proteins from the same source, e.g., from the same species, (3) is expressed by a cell from a different species, (4) has been separated from at least about 50 percent of polynucleotides, lipids, carbohydrates, or other materials with which it is associated in nature, (5) is not associated (by covalent or non-covalent interaction) with portions of a protein with which the “isolated protein” is associated in nature, (6) is operably associated (by covalent or non-covalent interaction) with a polypeptide with which it is not associated in nature, or (7) does not occur in nature. Such an isolated protein can be encoded by genomic DNA, cDNA, mRNA or other RNA, or may be of synthetic origin, or any combination thereof. In certain embodiments, the isolated protein is substantially free from proteins or polypeptides or other contaminants that are found in its natural environment that would interfere with its use (therapeutic, diagnostic, prophylactic, research or otherwise).

In certain embodiments, the “purity” of any given agent (e.g., polypeptide such as an antibody) in a composition may be defined. For instance, certain compositions may comprise an agent such as a polypeptide agent that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% pure on a protein basis or a weight-weight basis, including all decimals and ranges in between, as measured, for example and by no means limiting, by high performance liquid chromatography (HPLC), a well-known form of column chromatography used frequently in biochemistry and analytical chemistry to separate, identify, and quantify compounds.

A “lipid nanoparticle” or “solid lipid nanoparticle” refers to one or more spherical nanoparticles with an average diameter of between about 10 to about 1000 nanometers, and which comprise a solid lipid core matrix that can solubilize lipophilic molecules. The lipid core is stabilized by surfactants (e.g., emulsifiers), and can comprise one or more of triglycerides (e.g., tristearin), diglycerides (e.g., glycerol behenate), monoglycerides (e.g., glycerol monostearate),

fatty acids (e.g., stearic acid), steroids (e.g., cholesterol), and waxes (e.g., cetyl palmitate), including combinations thereof. Lipid nanoparticles are described, for example, in Petrilli et al., *Curr Pharm Biotechnol.* 15:847-55, 2014; and U.S. Pat. Nos. 6,217,912; 6,881,421; 7,402,573; 7,404,969; 7,550,441; 7,727,969; 8,003,621; 8,691,750; 8,871,509; 9,017,726; 9,173,853; 9,220,779; 9,227,917; and 9,278,130, which are incorporated by reference in their entireties. Certain compositions described herein are formulated with one or more lipid nanoparticles.

The terms or “Neuropilin 2-associated disease” or “NRP2-associated disease” refer to diseases and conditions in which NRP2 activity, expression, and/or spatial distribution plays a role in the pathophysiology of that disease or condition. In some instances, NRP2 associated diseases are modulated by the anti-NRP2 antibodies of the present disclosure by altering the interaction of NRP2 with at least one NRP2 ligand to impact NRP2 activity, signaling, expression, and/or spatial distribution. Exemplary NRP2-associated diseases and conditions include without limitation, cancer and diseases or pathologies associated with cancer including cancer cell growth, cancer initiation, cancer migration, cancer cell adhesion, invasion, chemoresistance, and metastasis. Also included are diseases associated with inflammation and autoimmunity, and related inflammatory diseases, including disease associated with inappropriate immune cell activation or migration such as graft versus host disease (GVHD). Additional examples include diseases associated with lymphatic development, lymphangiogenesis, and lymphatic damage, including edema, lymphedema, secondary lymphedema, inappropriate fat absorption and deposition, excess fat deposition, and vascular permeability. Also included are diseases associated with infections including latent infections, and diseases associated with allergic disorders/diseases and allergic responses, including chronic obstructive pulmonary disorder (COPD), neutrophilic asthma, antineutrophil cytoplasmic antibody (ANCA)-associated systemic vasculitis, systemic lupus erythematosus, rheumatoid arthritis, inflammasome-related disease(s), and skin-related neutrophil-mediated disease(s) such as pyoderma gangrenosum. Additional examples include diseases associated with granulomatous inflammatory diseases including sarcoidosis and granulomas, and fibrotic diseases including endometriosis, fibrosis, endothelial to mesenchymal transition (EMT), and wound healing, among others. Also included are diseases associated with inappropriate smooth muscle contractility, smooth muscle compensation and decompensation, vascular smooth muscle cell migration and/or adhesion, and diseases associated with inappropriate autophagy, phagocytosis, and efferocytosis. Also included are diseases associated with inappropriate migratory cell movement, as described herein. Additional examples include neuronal diseases, including diseases associated with peripheral nervous system remodeling and pain perception. Also included are diseases associated with bone development and/or bone remodeling. Typically, the term “inappropriate” refers to an activity or characteristic that associates with or causes a pathology or disease state.

The term “reference sequence” refers generally to a nucleic acid coding sequence, or amino acid sequence, to which another sequence is being compared. All polypeptide and polynucleotide sequences described herein are included as references sequences, including those described by name and those described in the Tables and the Sequence Listing.

Certain embodiments include biologically active “variants” and “fragments” of the polypeptides (e.g., antibodies) described herein, and the polynucleotides that encode the

same. “Variants” contain one or more substitutions, additions, deletions, and/or insertions relative to a reference polypeptide or polynucleotide (see, e.g., the Tables and the Sequence Listing). A variant polypeptide or polynucleotide comprises an amino acid or nucleotide sequence with at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity or similarity or homology to a reference sequence, as described herein, and substantially retains the activity of that reference sequence. Also included are sequences that consist of or differ from a reference sequences by the addition, deletion, insertion, or substitution of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150 or more amino acids or nucleotides and which substantially retain the activity of that reference sequence. In certain embodiments, the additions or deletions include C-terminal and/or N-terminal additions and/or deletions.

The terms “sequence identity” or, for example, comprising a “sequence 50% identical to,” as used herein, refer to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, Wis., USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., *Nucl. Acids Res.* 25:3389, 1997.

The term “solubility” refers to the property of an agent (e.g., antibody) provided herein to dissolve in a liquid solvent and form a homogeneous solution. Solubility is typically expressed as a concentration, either by mass of solute per unit volume of solvent (g of solute per kg of solvent, g per dL (100 mL), mg/mL, etc.), molarity, molality, mole fraction or other similar descriptions of concentration. The maximum equilibrium amount of solute that can dissolve per amount of solvent is the solubility of that solute in that solvent under the specified conditions, including temperature, pressure, pH, and the nature of the solvent. In certain embodiments, solubility is measured at physiological pH, or other pH, for example, at pH 5.0, pH 6.0, pH 7.0, pH 7.4, pH 7.6, pH 7.8, or pH 8.0 (e.g., about pH 5-8). In certain embodiments, solubility is measured in water or a physiological buffer such as PBS or NaCl (with or without NaPO₄). In specific embodiments, solubility is measured at relatively lower pH (e.g., pH 6.0) and relatively higher salt (e.g., 500 mM NaCl and 10 mM NaPO₄). In certain embodiments, solubility is measured in a biological fluid (solvent) such as blood or serum. In certain embodiments, the temperature can be about room temperature (e.g., about 20, 21,

47

22, 23, 24, 25° C.) or about body temperature (37° C.). In certain embodiments, an agent has a solubility of at least about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 70, 80, 90 or 100 mg/ml at room temperature or at 37° C.

A “subject” or a “subject in need thereof” or a “patient” or a “patient in need thereof” includes a mammalian subject such as a human subject.

“Substantially” or “essentially” means nearly totally or completely, for instance, 95%, 96%, 97%, 98%, 99% or greater of some given quantity.

By “statistically significant,” it is meant that the result was unlikely to have occurred by chance. Statistical significance can be determined by any method known in the art. Commonly used measures of significance include the p-value, which is the frequency or probability with which the observed event would occur, if the null hypothesis were true. If the obtained p-value is smaller than the significance level, then the null hypothesis is rejected. In simple cases, the significance level is defined at a p-value of 0.05 or less.

“Therapeutic response” refers to improvement of symptoms (whether or not sustained) based on administration of one or more therapeutic agents.

As used herein, the terms “therapeutically effective amount,” “therapeutic dose,” “prophylactically effective amount,” or “diagnostically effective amount” is the amount of an agent (e.g., anti-NRP2 antibody, immunotherapy agent) needed to elicit the desired biological response following administration.

As used herein, “treatment” of a subject (e.g., a mammal, such as a human) or a cell is any type of intervention used in an attempt to alter the natural course of the individual or cell. Treatment includes, but is not limited to, administration of a pharmaceutical composition, and may be performed either prophylactically or subsequent to the initiation of a pathologic event or contact with an etiologic agent. Also included are “prophylactic” treatments, which can be directed to reducing the rate of progression of the disease or condition being treated, delaying the onset of that disease or condition, or reducing the severity of its onset. “Treatment” or “prophylaxis” does not necessarily indicate complete eradication, cure, or prevention of the disease or condition, or associated symptoms thereof.

The term “wild-type” refers to a gene or gene product (e.g., a polypeptide) that is most frequently observed in a population and is thus arbitrarily designed the “normal” or “wild-type” form of the gene.

Each embodiment in this specification is to be applied to every other embodiment unless expressly stated otherwise. Anti-NRP2 Antibodies

Certain embodiments include antibodies, and antigen-binding fragments thereof, which specifically bind to a human neuropilin 2 (NRP2) polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof modulates (e.g., interferes with) binding of the human NRP2 polypeptide to at least one NRP2 ligand, such as a human histidyl-tRNA synthetase (HRS) polypeptide or other NRP2 ligand.

In certain embodiments, an antibody or antigen-binding fragment thereof is characterized by or comprises a heavy chain variable region (V_H) sequence that comprises complementary determining region V_H CDR1, V_H CDR2, and V_H CDR3 sequences, and a light chain variable region (V_L) sequence that comprises complementary determining region V_L CDR1, V_L CDR2, and V_L CDR3 sequences. Exemplary V_H , V_H CDR1, V_H CDR2, V_H CDR3, V_L , V_L CDR1,

48

V_L CDR2, and V_L CDR3 sequences are provided in Table A1, Table A2, and Table A3 below.

TABLE A1

Exemplary CDR Sequences		
Description	Sequence	SEQ ID NO:
aNRP2-1v3		
V_H CDR1	GYTFTSYWMH	1
V_H CDR2	AIYPGNSDTSYNQQFKG	2
V_H CDR3	RGGGYFDY	3
V_L CDR1	KASQNVGAAVA	4
V_L CDR2	SASNRYT	5
V_L CDR3	QQYSSYPPLT	6
aNRP2-2v4		
V_H CDR1	GYTFTSYWMH	7
V_H CDR2	VIHPNSASTFYNERFKT	8
V_H CDR3	PGTVRRSDY	9
V_L CDR1	RSSQNIHVHSTGMTYLE	10
V_L CDR2	KVSNRFS	11
V_L CDR3	FQGSHPVPT	12
aNRP2-10v5		
V_H CDR1	GFNIKDYIYH	13
V_H CDR2	RIDVEDDETKYAPKFQG	14
V_H CDR3	PIYGAREAWFAY	15
V_L CDR1	TASSSVSSSYLH	16
V_L CDR2	RTSNLAS	17
V_L CDR3	HQYRSPPPT	18
aNRP2-10v10		
V_H CDR1	GFNIKDYIVH	19
V_H CDR2	RIDVEDDETKYAPKFQG	20
V_H CDR3	PIYGAREAWFAY	21
V_L CDR1	TASSSVSSSYLH	22
V_L CDR2	RTSNLAS	23
V_L CDR3	HQYRSPPPT	24
aNRP2-10v13		
V_H CDR1	GFNIKDYIVH	57
V_H CDR2	RIDVHDETKYAPKFQG	58
V_H CDR3	PIYGAREAWFAY	59
V_L CDR1	TASSSVSSSYLH	60
V_L CDR2	RTSNLAS	61
V_L CDR3	HQYRSPPPT	62
aNRP2-11v7		
V_H CDR1	GYTFTSFGIS	25
V_H CDR2	EIYPRSGNTYYNENFKG	26
V_H CDR3	SSGYGSTPPFY	27
V_L CDR1	RASQDISNYLN	28
V_L CDR2	YTSRLHS	29
V_L CDR3	QQGNTLPWT	30
aNRP2-14v9		
V_H CDR1	GFSLTSYGVH	31
V_H CDR2	LIWGGSTDYSPAFIS	32
V_H CDR3	NSYSSGYAMDY	33
V_L CDR1	KASQNVGHAVA	34
V_L CDR2	SASNRYT	35
V_L CDR3	QQYSRPPYT	36
aNRP2-14v10		
V_H CDR1	GFSLTSYGVH	37
V_H CDR2	LIWGGSTDYSPAFIS	38
V_H CDR3	NSYSSGYAMDY	39
V_L CDR1	KASQNVGTAVA	40
V_L CDR2	SASNRYT	41
V_L CDR3	QQRSRPPYT	42

49

TABLE A1-continued

Exemplary CDR Sequences		
Description	Sequence	SEQ ID NO:
aNRP2-28v2/4		
V _H CDR1	GFSLSTYSIS	63
V _H CDR2	IIGDAGGIIYATWAKS	64

50

TABLE A1-continued

Exemplary CDR Sequences		
Description	Sequence	SEQ ID NO:
V _H CDR3	DGTAFDI	65
V _L CDR1	QASQSIYSKLG	66
V _L CDR2	RASTLAS	67
V _L CDR3	QQDYSYINVDNI	68

TABLE A2

Exemplary Polypeptide Sequences		
Description	Sequence	SEQ ID NO:
aNRP2-1v3		
Heavy chain variable region (V _H)	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYWMHWVRQAPGQGLEWM GAIYPGNSDTSYNQQFKGRVTMTTRDTSTSTVMELSSLRSEDTAVYYC ARRGGGYFDYWGGGTLTVSS	43
Light chain variable region (V _L)	DIQLTQSPSFLSASVGDRTVITCKASQNVGA AVAWYQQKPGKAPKLLI YSASNRYTGVPSPRFSGSGSGTEFTLTISSLQPEDFATYYCQQYSSYP LTFGGGTKVEIK	44
aNRP2-2v4		
Heavy chain variable region (V _H)	QVQLVQSGAEVKKPGASVKVSKASGYTFTSYWMHWVRQAPGQGLEWM GVIHPNSASTFYNERFKTRATMTVDRSSSTAYMELSSLRSEDTAVYYC SRPGTVRRSDYWGGGTTTVSS	45
Light chain variable region (V _L)	DVVMTQSPSLSPVTLGQPASISCRSSQNI VHSTGNTYLEWYQQRPQGS PRLLIYKVSNRFSGVPDRFSGSGSGTDFTLKISRVEADVGVYCFQG SHVPWTFGGGTKVEIK	46
aNRP2-10v5		
Heavy chain variable region (V _H)	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYIIHWVQQAPGKGLEWM GRIDVEDDETKYAPKFQGRVTITADTSTD TAYMELSSLRSEDTAVYYC ATPIYGSREAWFAYWGQGT LTVSS	47
Light chain variable region (V _L)	DIQMTQSPSSLSASVGDRTVITCTASSSVSSSYLHWYQQKPGKAPKLL IYRTSNLASGVPSPRFSGSGSGTDFTLTISSLQPEDFATYYCHQYYRSP PTFGGGTKVEIK	48
aNRP2-10v10		
Heavy chain variable region (V _H)	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYIVHWVQQAPGKGLEWM GRIDVEDDETKYAPKFQGRVTITADTSTD TAYMELSSLRSEDTAVYYC ATPIYGAREAWFAYWGQGT LTVSS	49
Light chain variable region (V _L)	DIQMTQSPSSLSASVGDRTVITCTASSSVSSSYLHWYQQKPGKAPKLL IYRTSNLASGVPSPRFSGSGSGTDFTLTISSLQPEDFATYYCHQYYRSP PTFGGGTKVEIK	50
aNRP2-10v13		
Heavy chain variable region (V _H)	EVQLVQSGAEVKKPGATVKISCKVSGFNIKDYIVHWVQQAPGKGLEWM GRIDVHDETKYAPKFQGRVTITADTSTD TAYMELSSLRSEDTAVYYC ATPIYGAREAWFAYWGQGT LTVSS	69
Light chain variable region (V _L)	DIQMTQSPSSLSASVGDRTVITCTASSSVSSSYLHWYQQKPGKAPKLL IYRTSNLASGVPSPRFSGSGSGTDFTLTISSLQPEDFATYYCHQYYRSP PTFGGGTKVEIK	70
aNRP2-11v7		
Heavy chain variable region (V _H)	QVQLVQSGAEVKKPGASVKVSKASGYTFTSPGISWVRQAPGQGLEWI GEIYPRSGNTYYNENFKGRATMTADKSTSTAYMELRSLRSDDTAVYYC ARSSGYGSTPPPYWGQGT LTVSS	51
Light chain variable region (V _L)	DIQMTQSPSSLSASVGDRTVITCRASQDISNYLNWYQQKPGKAVKLLI YYTSRLHSGVPSRFSGSGSGTDYFTFTISSLQPEDATYYCQQGNTLPW TFGGGTKVEIK	52

TABLE A2-continued

Exemplary Polypeptide Sequences		
Description	Sequence	SEQ ID NO:
aNRP2-14v9		
Heavy chain variable region (V_H)	QLQLVESGGGVVQPGRSLRLSCTVSGFSLTSYGVHWRQAPGKGLEWV GLIWSGGSTDYSPAFISRFTISEDNSKSTVYLQMNSLRAEDTAVYFCA RNSYSSGGYYAMDYWGQGTTVTVSS	53
Light chain variable region (V_L)	DIQLTQSPSFLSASVGDRTVITCKASQNVGHAVAWYQQKPGKAPKLLI YSASNRYTGVPSPRFGSGSGTEFTLTISLQPEDFATYYCQQYSRYPP YTPGGGTKVEIK	54
aNRP2-14v10		
Heavy chain variable region (V_H)	QLQLVESGGGVVQPGRSLRLSCTVSGFSLTSYGVHWRQAPGKGLEWV GLIWSGGSTDYSPAFISRFTISEDNSKSTVYLQMNSLRAEDTAVYFCA RNSYSSGGYYAMDYWGQGTTVTVSS	55
Light chain variable region (V_L)	DIQLTQSPSFLSASVGDRTVITCKASQNVGTAVAWYQQKPGKAPKLLI YSASNRYTGVPSPRFGSGSGTEFTLTISLQPEDFATYYCQQRSRYPP YTPGGGTKVEIK	56
aNRP2-28v2		
Heavy chain variable region (V_H)	QSVKESEGGFLFKPTDRTLTLCTVSGFSLSTYSISWVRQAPGNLEWIG IIGDAGGIYATWAKSRSTITRSTALNTVTLKMTGLTAADTATYFCAR DGTAFDIWGPGLTVTVSS	71
Light chain variable region (V_L)	AYDMTQTPASVEVVVGTVTIKCQASQSIYSKLGWYQQKPGQPPKLLI YRASTLASGVSSRFKSGSGTEYTLTISGVQCDAAATYYCQQDYSYIN VDNIFGGGTEVVVK	72
aNRP2-28v4		
Heavy chain variable region (V_H)	QSVKESEGGFLFKPTDRTLTLCTVSGFSLSTYSISWVRQAPGNLEWIG IIGDAGGIYATWAKSRSTITRSTALNTVTLKMTGLTAADTATYFCAR DGTAFDIWGPGLTVTVSS	73
Light chain variable region (V_L)	AYDMTQTPASVEVVVGTVTIKCQASQSIYSKLGWYQQKPGQPPKLLI YRASTLASGVSSRFKSGSGTEYTLTISGVQADDAATYYCQQDYSYIN VDNIFGGGTEVVVK	74

Thus, in certain embodiments, an antibody or antigen-binding fragment thereof comprises

- a heavy chain variable region (V_H) sequence that comprises complementary determining region V_H CDR1, V_H CDR2, and V_H CDR3 sequences selected from Table A1 and variants thereof which specifically bind to a human NRP2 polypeptide (selected, for example, from Table N1); and
- a light chain variable region (V_L) sequence that comprises complementary determining region V_L CDR1, V_L CDR2, and V_L CDR3 sequences selected from Table A1 and variants thereof which specifically bind to the human NRP2 polypeptide (selected, for example, from Table N1).

In certain embodiments, the CDR sequences are as follows:

- the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 1-3, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 4-6, respectively, including variants thereof;
- the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 7-9, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 10-12, respectively, including variants thereof;
- the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 13-15, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 16-18, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 19-21, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 22-24, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 25-27, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 28-30, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 34-36, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 34-39, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 40-42, respectively, including variants thereof;

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 57-59, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 60-62, respectively, including variants thereof; or

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 63-65, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 66-68, respectively, including variants thereof.

53

Also included are variants thereof, including affinity matured variants, which bind to human NRP2, for example, variants having 1, 2, 3, 4, 5, or 6 alterations in one or more of the CDR regions, for example, one or more the V_HCDR1, V_HCDR2, V_HCDR3, V_LCDR1, V_LCDR2, and/or V_LCDR3 sequences described herein. Exemplary “alterations” include amino acid substitutions, additions, and deletions.

In certain embodiments, the V_H sequence is at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from Table A2, including, for example, wherein the V_H sequence has 1, 2, 3, 4, or 5 alterations in one or more framework regions.

In some embodiments, the V_L sequence is at least 80, 85, 90, 95, 97, 98, 99, or 100% identical to a sequence selected from Table A2, including, for example, wherein the V_L sequence has 1, 2, 3, 4, or 5 alterations in one or more framework regions.

In some embodiments, the V_H and V_L sequences of an antibody or antigen-binding fragment are as follows:

- the V_H sequence comprises SEQ ID NO: 43, and the V_L sequence comprises SEQ ID NO: 44;
- the V_H sequence comprises SEQ ID NO: 45, and the V_L sequence comprises SEQ ID NO: 46;

54

- the V_H sequence comprises SEQ ID NO: 47, and the V_L sequence comprises SEQ ID NO: 48;
- the V_H sequence comprises SEQ ID NO: 49, and the V_L sequence comprises SEQ ID NO: 50;
- the V_H sequence comprises SEQ ID NO: 51, and the V_L sequence comprises SEQ ID NO: 52;
- the V_H sequence comprises SEQ ID NO: 53, and the V_L sequence comprises SEQ ID NO: 54;
- the V_H sequence comprises SEQ ID NO: 55, and the V_L sequence comprises SEQ ID NO: 56;
- the V_H sequence comprises SEQ ID NO: 69, and the V_L sequence comprises SEQ ID NO: 70;
- the V_H sequence comprises SEQ ID NO: 71, and the V_L sequence comprises SEQ ID NO: 72; or
- the V_H sequence comprises SEQ ID NO: 73, and the V_L sequence comprises SEQ ID NO: 74.

Also included are variants thereof, for example, variants having 1, 2, 3, 4, or 5 alterations in one or more framework regions. Exemplary “alterations” include amino acid substitutions, additions, and deletions.

Table A3 below summarizes exemplary redundant antibody CDRs, including exemplary consensus CDR sequences, and provides amino acid code for the same.

TABLE A3

Redundant antibody CDRs			
Antibody Chain	CDR1	CDR2	CDR3
aNRP2-10 V _H	GFNX ₂₂ KDX ₄₂ X ₁₄ X ₂₂ H (SEQ ID NO: 75)	RIDX ₄₃ EDDETKYAPKFQG (SEQ ID NO: 76)	X ₁ X ₂ X ₃ X ₄ X ₅ RX ₆ X ₇ X ₈ X ₉ X ₁₀ X ₁₁ (SEQ ID NO: 77)
aNRP2-10 V _L	TASSSVSSSYLH (SEQ ID NO: 16)	X ₂₁ TSNLAS (SEQ ID NO: 78)	X ₁₂ X ₁₃ X ₁₄ X ₁₅ X ₁₆ X ₁₇ X ₁₈ X ₁₉ X ₂₀ (SEQ ID NO: 79)
aNRP2-11 V _H	GYTFTSFGIS (SEQ ID NO: 25)	EIYPRSGNTYYNENFKG (SEQ ID NO: 26)	X ₂₃ X ₂₄ GX ₂₅ YX ₂₆ STPX ₂₇ X ₂₈ X ₂₉ (SEQ ID NO: 80)
aNRP2-11 V _L	RASQDISNYLN (SEQ ID NO: 28)	YTSRLHS (SEQ ID NO: 29)	QQGNTLPWT (SEQ ID NO: 30)
aNRP2-14 V _H	GFSLTSYGVH (SEQ ID NO: 31)	LIWSGGSTDYSPAFIS (SEQ ID NO: 32)	NX ₃₀ X ₃₁ X ₃₂ X ₃₃ GYXX ₃₄ X ₄₁ DX ₃₅ (SEQ ID NO: 81)
aNRP2-14 V _L	KASQNVGX ₄₀ AVA (SEQ ID NO: 82)	SASNRYT (SEQ ID NO: 5)	QQX ₃₆ X ₃₇ X ₃₈ X ₃₉ PPYT (SEQ ID NO: 83)

Redundant Amino acid code

X ₁	A, E, F, G, H, K, L, N, P, T, Y
X ₂	D, E, H, I, K, L, P, Q, R, S, T, V, Y
X ₃	A, F, G, H, I, K, L, P, Q, R, S, T, V, Y
X ₄	A, F, G, S, T, Y
X ₅	A, F, G, H, I, K, N, Q, R, S, T, V, Y
X ₆	A, E, P, Q, S, T
X ₇	A, I, L, N, Q, S, V
X ₈	F, W, Y
X ₉	F, L
X ₁₀	A, D, F, G, H, K, L, N, P, R, S, T, Y
X ₁₁	A, D, E, F, G, H, I, K, L, N, R, S, T, V, Y
X ₁₂	D, F, G, H, I, K, L, N, R, S, T, V
X ₁₃	A, D, E, G, Q, S, T
X ₁₄	F, Y
X ₁₅	F, G, H, N, Y
X ₁₆	H, N, Q, R, S, T, Y
X ₁₇	A, F, H, I, L, N, P, S, T, V
X ₁₈	A, D, F, G, H, K, L, P, S, T, V, Y
X ₁₉	A, G, L, P, S, T
X ₂₀	A, D, E, F, G, H, I, K, L, N, Q, R, S, T, V, Y
X ₂₁	R, S
X ₂₂	I, V
X ₂₃	S, T
X ₂₄	G, K, L, S, T, V
X ₂₅	L, Y

TABLE A3-continued

Redundant antibody CDRs			
Antibody Chain	CDR1	CDR2	CDR3
X ₂₆		A, G, S	
X ₂₇		A, F, H, L, N, Q, S, T, Y	
X ₂₈		A, I, P, T, V	
X ₂₉	A, E, G, H, I, K, L, N, Q, R, S, T, Y		
X ₃₀		F, G, I, S, T, V, Y	
X ₃₁		H, F, Y	
X ₃₂		A, F, G, S	
X ₃₃		F, S, Y	
X ₃₄		A, G, H, S, P, T	
X ₃₅	D, G, H, I, A, V, L, R, S, Y		
X ₃₆		F, H, K, R, Y	
X ₃₇		A, I, L, N, R, S, T, V	
X ₃₈	F, G, I, K, L, N, R, S, T, V, Y		
X ₃₉		A, F, G, I, Q, R, S, T, Y	
X ₄₀		A, G, H, N, Q, R, S, T	
X ₄₁		M, K	
X ₄₂		S, Y	
X ₄₃		V, P	

Thus, in certain embodiments, the at least one anti-NRP2 antibody or antigen-binding fragment thereof comprises a CDR sequence, for example, a CDR1 consensus sequence, from Table A3.

Neuropilin-2 is a cell surface receptor protein that modulates a broad range of cellular functions through its roles as an essential cell surface receptor and co-receptor for a variety of ligands (see, e.g., Guo and Vander Kooi, *J. Cell. Biol.* 290 No 49:29120-29126, 2015). For instance, it functions during epithelial to mesenchymal transition (EMT), for example, by promoting TGF- β 1-mediated EMT in colorectal and other cancer cells (see, e.g., Grandclement et al., *PLOS ONE* 6 (7) e20444, 2011), and by mediating EMT or endo-EMT in fibroblasts, myofibroblasts, and endothelial cells to promote fibrosis formation (see, e.g., Pardali et al., *Int. J. Mol. Sci.* 18:2157, 2017).

Neuropilin-2 expression promotes lymphangiogenesis (see, e.g., Doci et al., *Cancer Res.* 75:2937-2948, 2015). Single nucleotide polymorphisms (SNPs) in NRP2 are associated with lymphedema (see, e.g., Miaskowski et al., *PLOS ONE* 8 (4) e60164, 2013). NRP2 also regulates smooth muscle contractility (see, e.g., Bielenberg et al., *Amer. J. Path.* 181:548-559, 2012), regulates autophagy, for example, in cancer (see, e.g., Stanton et al., *Cancer Res.* 73:160-171, 2013), contributes to tumor initiation, survival, and metastasis (see, e.g., Goel et al., *EMBO Mol. Med.* 5:488-508, 2013; and Samuel et al., *PLOS ONE* 6 (10) e23208, 2011), and regulates immune cell activation and migration (see, e.g., Mendes-da-Cruz et al., *PLOS ONE* 9 (7) e103405, 2014). Neuropilins are also multifunctional co-receptors involved in tumor initiation, growth, metastasis and immunity (see, e.g., Prud'homme et al., *Oncotarget* 3:921-939, 2012).

Neuropilin-2 is expressed in various cells of the immune system, including lymphoid cells such as B and T cells, and myeloid cells such as basophils, eosinophils, monocytes, dendritic cells, neutrophils, and macrophages, including tissue-specific macrophages, for example, alveolar macrophages. It is also expressed in endothelial and epithelial cells in the lung and other tissues, and in muscle cells [see, e.g., Bielenberg et al., *Amer. J. Path.* 181:548-559, 2012; Aung, et al., *PLOS ONE* 11 (2) e0147358, 2016; Schellenburg et al., *Mol. Imm.* 90:239-244, 2017; and Wild et al., *Int. J. Exp. Path.* 93:81-103, 2012).

Neuropilin-2 also plays a key role in endosome development, for example, by regulating late endosomal maturation, an important aspect of phagocytosis and efferocytosis, which respectively contribute to clearance of infections and apoptotic cells (See, e.g., Diaz-Vera et al., *J. Cell. Sci.* 130:697-711, 2017; Dutta et al., *Cancer Res.* 76:418-428, 2016).

Neuropilin-2 is known to be a key player in the pathophysiology of many diseases (e.g., "NRP2-associated diseases") and interacts with a broad array of soluble ligands including semaphorin 3F, VEGF-C and D, and TGF- β (see, for example, Table N2 and Table N3), and an array of cellular receptors and co-factors (see, for example, FIGS. 1A-1B and FIG. 2). NRP2 is also polysialated on dendritic cells, and actively interacts with the chemokine CCL21 to mediate immune cell migration, and for which single nucleotide polymorphisms associated with ILD and RA have been described (see, e.g., Rey-Gallardo et al., *Glycobiology* 20:1139-1146, 2010; Stahl et al., *Nat. Genet.* 42:508-514, 2013; and Miller et al., *Arthritis Rheum.* 65:3239-3247). Additionally, soluble, circulating forms of NRP-2 are known (see, e.g., Parker et al., *Structure* 23 (4) 677-687, 2015), and internal studies have confirmed the existence of circulating complexes of HRS polypeptides and NRP-2 polypeptides in circulation. Accordingly, given the central role played by NRP2 in pathophysiology in a broad range of diseases, it is evident that interactions between NRP2 and NRP2 ligand(s) (for example, NRP2 ligands from Table N2 and Table N3), and the modulation of those interactions with antibodies against NRP2 to selectively change the corresponding biological activities, provides broad potential for the treatment of diseases, including NRP2 associated diseases.

NRP2 is a single transmembrane receptor with a predominant extracellular region containing two CUB domains (a1/a2 combined domain), two Factor V/VIII homology domains (b1/b2 combined domain), a MAM domain (c domain) (see FIGS. 1A-1B), and a short juxtamembrane region that connects the c domain to the transmembrane domain (which traverses the plasma membrane). The a1a2 combined domain interacts with sema region of the semaphorins, and the b1 domain interacts with the semaphorin PSI and Ig-like domains. NRP2 has a higher affinity for SEMA3F and 3G; in contrast, SEMAs 3A, 3B and 3E preferentially interact with NRP1. Both NRP1 and NRP2

have similar affinity for SEMA 3C. The b1b2 combined domain interacts with several growth factors containing heparin-binding domains, including VEGF C & D, placenta growth factor (PIGF)-2, fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor (PDGF), and transforming growth factor (TGF)-beta (see, for example, Prud'homme et al., *Oncotarget*. 3:921-939, 2012). NRP2 also interacts with various growth factor-specific receptors, and interactions with these receptors occur independently of binding to SEMAs. In this context, integrins and growth factor receptors like VEGF receptor, TGF-beta receptor, c-Met, EGFR, FGFR, PDGFR, have been shown to interact with NRPs and in general appear to increase the affinity of each ligand for its receptor and to modulate down stream signaling. The c domain (Mam) domain does not appear to be directly required for ligand binding, but may impact ligand specificity, receptor signaling, and NRP2 dimerization. The juxtamembrane region differs significantly between the NRP2a and NRP2b isoforms, and may also impact ligand specificity, dimerization, and signaling.

Accordingly, anti-NRP2 antibodies and antigen-binding fragments thereof that bind to the a1 and/or a2 domains of NRP2 have the potential to selectively modulate semaphorin binding. Likewise, anti-NRP2 antibodies and antigen-binding fragments thereof that bind to the b1 domain have the potential to modulating both semaphorin and VEGF and growth factor binding, and anti-NRP2 antibodies that bind to the b2 domain have the potential to selectively modulate VEGF and growth factor binding. Antibodies and antigen-binding fragments thereof that bind to the c domain might not directly impact NRP2 ligand binding, but have the potential to modulate NRP2 downstream signaling, for example, by modulating (e.g., promoting or enhancing) NRP2 receptor dimerization.

Anti-NRP2 antibodies and antigen binding fragments thereof that promote NRP2 receptor homodimerization could modulate NRP2 receptor activity, and provide agonistic or antagonistic antibodies depending on the nature of the binding site. Such antibodies and antigen binding fragments thereof could modulate (e.g., enhance) the activities of NRP2 ligands, even if they do not directly modulate ligand binding. Additional diversity in the functional effects of specific anti-NRP2 antibodies may be expected based on their binding mode, and as a result of steric effects, which may indirectly impact ligand binding.

NRP2 can form homodimers as well as heterodimers, and is heavily glycosylated. NRP2 has different splice variants which are between about 551 and 926 amino acids long. Two major variants for NRP2 are categorized as NRP2a and NRP2b. These differ in their intracellular C terminal part (FIGS. 1A-1B) in which for NRP2a, the c-terminal domain comprises 42 amino acids and a PDZ-binding domain with the C-terminal SEA amino acid sequence. By contrast, NRP2b comprises a 46 amino acid C terminal domain which shares about 11% sequence homolog with the intracellular, juxtamembrane, and transmembrane sequences of NRP2a. Between the MAM domain and the transmembrane domain, additional splicing can occur and 5 additional amino acids (GENFK; SEQ ID NO: 153) can be added to either the NRP2a, or NRP2b forms—these variants are named based on the number additional amino acids added through alternative splicing. Thus the two variants of NRP2a are named NRP2a (17) (or variant 1) and NRP2a (22) (or variant 2), and the two transmembrane variants for NRP2b are named NRP2b (0) (or variant 4) and NRP2b (5) (or variant 5). Additionally, a soluble form called sNRP2b (or variant 6) can be generated. Exemplary NRP2 polypeptide sequences are provided in Table N1 below, including the mature (after cleavage of the N-terminal signal peptide) and precursor forms of the various isoforms of NRP2.

TABLE N1

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
Human full length NRP2	1-931	MDMFLTWVFLALYFSRHQVRGQPDPPCGRLNSKDAGYI	84
Variant 1 precursor		TSPGYPDYPSHQNCWEIVYAPEPNQKIVLNFNPHFETEK	
NRP2a (22)		HDCKYDFIEIRDGDSADLLGKHCGNIAPPTIISSGSM	
		YIRFTSDYARQGAGFSLRYEIFKTGSEDCSKNFTSPNGTI	
		ESPGFPPEKYPHNLDCFTTILAKPKMEIILQLIFDLEHDP	
		LQVGEGDCKYDWLDIWDGI PHVGLIGKYCGTKTPSELRS	
		STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP	
		LGMESGRIANEQISASSTYS DGRWTPQQSRLHGDDNGWTP	
		NLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVKS	
		YKLEVSTNGEDWMVYRHGKNHKVFQANNDATVVLNKLHA	
		PLLTRFVRIRPQTWHSIALRLLELFGCRVTDAPCSNMLGM	
		LSGLIADSQISASSTQEYLWSPSAARLVSSRSGWFPRIPQ	
		AQPGEEWLQVDLGTPTKTVKGVI IQGARGGDSITAVEARAF	
		VRKFVSYSLNGKDWEYIQDPRTQPKLFEGNMHYDTPDI	
		RRFDP I PAQYVRVYPERWSPAGIMRLEVLGCDWTD SKPT	
		VETLGPTVKSEETTPYPTEEATECGENCSEFDDKDLQL	
		PSGFNCNFDLEPCGWMDHAKWLRTTWASSSSPNDRTF	
		PDDRNFLRLQSDS QREGQYARLISPPVHLPRSPVCMFQY	
		QATGGRGVALQVREASQESKLLWVIREDDQGEWKHGRI I	
		LPSYDMEYQIVFEGVIGKGRSGEIAIDDIRISTDVPLENC	
		MEPISAFAGENFKVDIPEIHEREGYEDEIDDEYEVDSNS	
		SSATSGSGAPSTDKEKSWLYTLDPILITIIAMSSLGVLLG	
		ATCAGLLLYCTCSYSGLSSRSCTTLENYFELYDGLKHKV	
		KMNHQKCCSEA	

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
Human NRP2 Variant 2 precursor NRP2a (17)	1-926	MDMFPLTWVFLALYFSRHQVRGQDPDPPCGRLNSKDAGYI TSPGYPDYPSHQNCEWIVYAPEPNQKIVLNFNPHFEIEK HDCKYDFIEIRDGDESEADLLGKHCGNIAPPTIISSGSML YIKFTSDYARQGAGFSLRYEIFKTGSDEDCSKNFTSPNGTI ESPGFPEKYPHNLDCTFTILAKPKMEIILQFLIFDLEHDP LQVGEGDCKYDWLDIWDGIPHVGPLIGKYCGTKTPSELRS STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP LGMESGRIANEQISASSTYSDGRWTPQQSRLHGDDNGWTP NLDSNKEYLQVDLRFLTMLTAIATQGAI SRETQNGYYVKS YKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHA PLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCSNMLGM LSGLIADSQISASSTQEYLWSPSAARLVSSRSGWFPRIPQ AQPGEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAF VRKFVSYSLNGKDWEYIQDPRTQQPKLFEGNMHYDTPDI RRFDPIPAQYVRVYPERWSPAGIGMRLEVLGCDWTD SKPT VETLGPTVKSEETTPYPTEEEATECGENC SFEDDKDLQL PSGFNCNFDLEEPCGWMDHAKWLRTWASSSSPNDRTF PDDRNFLRLQSDSQREGQYARLISPPVHLPRSPVCMEFQY QATGGRGVALQVVREASQESKLLWVIREDDQGEWKHGRII LPSYDMEYQIVFEGVIGKGRSGEIAIDDIRISTDVPLENC MEPISAFAVDIPEIHEREGYEDEIDDEYEDWNSNSSATS SGGAPSTDKEKSWLYTLDPILITIIAMSSLGVLGATCAG LLLYCTCSYSGLSSRSCTTLENYNFELYDGLKHKVKMNHQ KCCSEA	85
Human NRP2 Variant 3 precursor NRP2a (0)	1-909	MDMFPLTWVFLALYFSRHQVRGQDPDPPCGRLNSKDAGYI TSPGYPDYPSHQNCEWIVYAPEPNQKIVLNFNPHFEIEK HDCKYDFIEIRDGDESEADLLGKHCGNIAPPTIISSGSML YIKFTSDYARQGAGFSLRYEIFKTGSDEDCSKNFTSPNGTI ESPGFPEKYPHNLDCTFTILAKPKMEIILQFLIFDLEHDP LQVGEGDCKYDWLDIWDGIPHVGPLIGKYCGTKTPSELRS STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP LGMESGRIANEQISASSTYSDGRWTPQQSRLHGDDNGWTP NLDSNKEYLQVDLRFLTMLTAIATQGAI SRETQNGYYVKS YKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHA PLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCSNMLGM LSGLIADSQISASSTQEYLWSPSAARLVSSRSGWFPRIPQ AQPGEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAF VRKFVSYSLNGKDWEYIQDPRTQQPKLFEGNMHYDTPDI RRFDPIPAQYVRVYPERWSPAGIGMRLEVLGCDWTD SKPT VETLGPTVKSEETTPYPTEEEATECGENC SFEDDKDLQL PSGFNCNFDLEEPCGWMDHAKWLRTWASSSSPNDRTF PDDRNFLRLQSDSQREGQYARLISPPVHLPRSPVCMEFQY QATGGRGVALQVVREASQESKLLWVIREDDQGEWKHGRII LPSYDMEYQIVFEGVIGKGRSGEIAIDDIRISTDVPLENC MEPISAFADYEYEDWNSNSSATSGGAPSTDKEKSWLYTL DPILITIIAMSSLGVLGATCAGLLYCTCSYSGLSSRSC TTLENYNFELYDGLKHKVKMNHQKCCSEA	86
Human NRP2 Variant 4 precursor NRP2b (5)	1-906	MDMFPLTWVFLALYFSRHQVRGQDPDPPCGRLNSKDAGYI TSPGYPDYPSHQNCEWIVYAPEPNQKIVLNFNPHFEIEK HDCKYDFIEIRDGDESEADLLGKHCGNIAPPTIISSGSML YIKFTSDYARQGAGFSLRYEIFKTGSDEDCSKNFTSPNGTI ESPGFPEKYPHNLDCTFTILAKPKMEIILQFLIFDLEHDP LQVGEGDCKYDWLDIWDGIPHVGPLIGKYCGTKTPSELRS STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP LGMESGRIANEQISASSTYSDGRWTPQQSRLHGDDNGWTP NLDSNKEYLQVDLRFLTMLTAIATQGAI SRETQNGYYVKS YKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHA PLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCSNMLGM LSGLIADSQISASSTQEYLWSPSAARLVSSRSGWFPRIPQ AQPGEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAF VRKFVSYSLNGKDWEYIQDPRTQQPKLFEGNMHYDTPDI RRFDPIPAQYVRVYPERWSPAGIGMRLEVLGCDWTD SKPT VETLGPTVKSEETTPYPTEEEATECGENC SFEDDKDLQL PSGFNCNFDLEEPCGWMDHAKWLRTWASSSSPNDRTF PDDRNFLRLQSDSQREGQYARLISPPVHLPRSPVCMEFQY QATGGRGVALQVVREASQESKLLWVIREDDQGEWKHGRII LPSYDMEYQIVFEGVIGKGRSGEIAIDDIRISTDVPLENC MEPISAFAGENFKGGTLLPGTEPTVDTPMQPIPAYWYV MAGGAVLVLSVALALVLHYHFRYAAKKTDSITYKTS HYTNGAPLAVEPTLTIKLEQDRGSHC	87

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
Human NRP2 Variant 5 precursor NRP2b (0)	1-901	MDMFPLTWVFLALYFSRHQVRGQDPDPPCGRLNSKDAGYI TSPGYPDYPSHQNCWEIVYAPEPNQKIVLNFNPHFEIEK HDCKYDFIEIRDGDEESADLLGKHCGNIAPPTIISSGSML YIKFTSDYARQGAGFSLRYEIKFTGSEDCSKNFTSPNGTI ESPGFPEKYPHNLDCTFTILAKPKMEIILQFLIFDLEHDP LQVGECDCKYDWLDIWDGIPHVGLIGKYCGTKTPSELRS STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP LGMESGRIANEQISASSTYSDGRWTPQQSRLHGDDNGWTP NLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVVK YKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHA PLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCSNMLGM LSGLIADSQISASSTQEYLWSPSAARLVSSRSWGFPRIPQ AQPGEELQVDLGTPTVKGVIIQGARGGDSITAVEARAF VRKFKVSYSLNGKDWEYIQDPRTQQPKLFEKNMHYDTPDI RRFDPIPAQYVVRVYPERWSPAGIMRLEVLGCDWTDKPT VETLGPVTKSEETTPYPTREEATECGENCSEDDKDLQL PSGFNCNFDLEPCGWMYDHAKWLRITWASSSSPNDRTF PDDRNFLRLQSDSQREGQYARLISPPVHLPRSPVCMFQY QATGGRGVALQVVREASQESKLWVIREDDQGEWKHGRII LPSYDMEYQIVFEGVIGKRSGEIAIDDIRISTDVPLENC MEPISAFAGGTLPGTEPTVDTVPMQPIPAYWYVMAAGG AVLVLVVALALVLYHHRFRYAAKKTDHSTYKTSHTYTN APLAVEPTLTIKLEQDRGSHC	88
Human NRP2 Variant 6 precursor S9NRP2b Soluble NRP2	1-555	MDMFPLTWVFLALYFSRHQVRGQDPDPPCGRLNSKDAGYI TSPGYPDYPSHQNCWEIVYAPEPNQKIVLNFNPHFEIEK HDCKYDFIEIRDGDEESADLLGKHCGNIAPPTIISSGSML YIKFTSDYARQGAGFSLRYEIKFTGSEDCSKNFTSPNGTI ESPGFPEKYPHNLDCTFTILAKPKMEIILQFLIFDLEHDP LQVGECDCKYDWLDIWDGIPHVGLIGKYCGTKTPSELRS STGILSLTFHTDMAVAKDGFSAARYLVHQEPLNFQCNVP LGMESGRIANEQISASSTYSDGRWTPQQSRLHGDDNGWTP NLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVVK YKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHA PLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCSNMLGM LSGLIADSQISASSTQEYLWSPSAARLVSSRSWGFPRIPQ AQPGEELQVDLGTPTVKGVIIQGARGGDSITAVEARAF VRKFKVSYSLNGKDWEYIQDPRTQQPKVGCWRPL	89
Human NRP2 Variant 2 NRP2a (17) (mature)	23-926	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCWEIVYAP EPNQKIVLNFNPHFEIEKHCKYDFIEIRDGDEESADLLG KHCGNIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIK FTGSEDCSKNFTSPNGTIESPGFPEKYPHNLDCTFTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSDG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTREVRIRPQTWHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSQISASSTQEYLWSP SAARLVSSRSWGFPRIPQAQPGEELQVDLGTPTVKGVII IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEKNMHYDTPDIRFDPIPAQYVVRVYPERWSPAG IMRLEVLGCDWTDKPTVETLGPVTKSEETTPYPTREE ATECGENCSEDDKDLQLPSGFNCNFDLEPCGWMYDHA KWLRTWASSSSPNDRTFPDDRNFLRLQSDSQREGQYARL ISPPVHLPRSPVCMFQYQATGGRGVALQVVREASQESKL LWVIREDDQGEWKHGRIILPSYDMEYQIVFEGVIGKRS GEIAIDDIRISTDVPLENCMEPISAFVADIPEIHEREGYED EIDDEYVDWSNSSSATSGSGAPSTDKEKSWLYTLDPI LTIAMSLGLVLLGATCAGLLLYCTCSYGLSSRSCTTLEN YNFELYDGLKHKVKMNHQKCCSEA	90
NRP2 splice variant 5 NRP2b (0) (mature)	23-901	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCWEIVYAP EPNQKIVLNFNPHFEIEKHCKYDFIEIRDGDEESADLLG KHCGNIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIK FTGSEDCSKNFTSPNGTIESPGFPEKYPHNLDCTFTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSDG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTREVRIRPQTWHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSQISASSTQEYLWSP	91

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
		SAARLVSSRSWFPRIIPAQPGEEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLLEVLCDDWTDKPTVETLGTPTVKEETTTPTPTTEE ATECGENCSEDDKDLQPLSGFNCNFDLEPCGWMYDHA KWLRTTWASSSSPNDRTPDDRNLRLQSDSQREGQYARL ISPPVHLPRSPVCMFQYQATGGRGVALQVVREASQESKL LWVIREDDQGGWKHGRIILPSYDMEYQIVFEGVIGKGRSG EIAIDDIRISTDVPLENCMEPISAFAGGTLPGTEPTVDT VPMQPIPAYWYVMAAGGAVLVLSVALALVLHYHFRFYA AKKTDHSITYKTSHTNGAPLAVEPTLTIKLEQDRGSHC	
Soluble NRP2 S9Nrp-2b (mature)	23-555	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPHFEIEKHDCKYDFIEIRDGDSEADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSAR RYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYG RWTQQSRLHGGDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVYSYKLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTRFVIRPQTWHSIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSWFPRIIPAQPGEEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKVGCWRPL	92
NRP2 a1 domain	28-141	CGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAPEPNQK IVLNFNPHFEIEKHDCKYDFIEIRDGDSEADLLGKHCN IAPPTIISSGSMLYIKFTSDYARQAGFSLRYEIF	93
NRP2 a2 domain	149-265	CSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAKPKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV YCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSARYY	94
NRP2 b1 domain	280-426	PLGMESGRIANEQISASSTYSYGRTWTQQSRLHGGDNGWTPNLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVYSYKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHAPLLTRFVIRPQTWHSIALRLLELFG	95
NRP2 b2 domain	438-591	LGMLSGLIADSIASSTQEYLWSPSAARLVSSRSWFPRIIPAQPGEEWLQVDLGTPTVKGVIIQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAGIGMRLLEVLCDDWTDKPTVETLGTPTVKEETTTPTPTTEE	96
NRP2 c domain	641-794	PSGFNCNFDLEPCGWMYDHAHWLRTTWASSSSPNDRTPDDRNLRLQSDSQREGQYARLISPPVHLPRSPVCMFQYQATGGRGVALQVVREASQESKLWVIREDDQGGWKHGRIILPSYDMEYQIVFEGVIGKGRSGEIAIDDIRISTD	97
NRP2 a1a2 combined domains	23-265	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPHFEIEKHDCKYDFIEIRDGDSEADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSARYY	98
NRP2 a2b1 combined domains	149-426	CSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAKPKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV YCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSARYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYGRTWTQQSRLHGGDNGWTPNLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVYSYKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVLNKLHAPLLTRFVIRPQTWHSIALRLLELFG	99
NRP2 a1a2b1 combined domains	23-426	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPHFEIEKHDCKYDFIEIRDGDSEADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSARYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYGRTWTQQSRLHGGDNGWTPNLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNGYVYSYKLEVSTNGEDWMVYRHGKNHK	100

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
		VFQANNDATEVVLNKLHAPLLTRFVRIRPQTHSGIALRL ELFG	
NRP2 a1a2b1b2 combined domains	23-595	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCWEIVYAP EPNQKIVLNFNPHFEIEKHDCYDFIEIRDGDESADLLG KHCGNIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIF KTGSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYYL VHQEPL ENFQCNVPLGMESGRIANEQISASSTYSDG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTRFVRIRPQTHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSGWFPRI PQAQPGEEWLQVDLGTPTVKGV I IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLEVLGCDWT	101
NRP2 a2b1b2 combined domains	145-595	GSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAKPK MEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHVGP LIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYL VHQEPL ENFQCNVPLGMESGRIANEQISASSTYSDGRW TPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAIAT QGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHKV QANNDATEVVLNKLHAPLLTRFVRIRPQTHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSGWFPRI PQAQPGEEWLQVDLGTPTVKGV I IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLEVLGCDWT	102
NRP2 b1b2 combined domains	276-595	QCNVPLGMESGRIANEQISASSTYSDGRWTPQQSRLHGDD NGWTPNLDSNKEYLQVDLRFLTMLTAIATQGAISRETQNG YVVKSYKLEVSTNGEDWMVYRHGKNHKVFQANNDATEVVL NKLHAPLLTRFVRIRPQTHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSGWFPRI PQAQPGEEWLQVDLGTPTVKGV I IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLEVLGCDWT	103
NRP2 v2-Fc fusion protein	23-855	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCWEIVYAP EPNQKIVLNFNPHFEIEKHDCYDFIEIRDGDESADLLG KHCGNIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIF KTGSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYYL VHQEPL ENFQCNVPLGMESGRIANEQISASSTYSDG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTRFVRIRPQTHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSGWFPRI PQAQPGEEWLQVDLGTPTVKGV I IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLEVLGCDWTDKPTVETLGTPTVKS EETTPYPT EEE ATECGENCSFEDDKDLQLPSGFNCNFDLEPCGWMYDHA KWLRTTWASSSSPNDRFPDDRNLRLQSDSQREGQYARL ISPPVHLPRSPVCMFQYQATGGRGVALQVVRASQESKL LWVIREQDQGEWKHGRIILPSYDMEYQIVFEGVIGKGRSG EIAIDDIRISDTPLENCMEPI SAFAVDIPEIHEREGYED EIDDEYEVDWSNSSATSGSGAPSTDKEKSWLYDKTHTCP PCPAPELLGGPSVFLFPPKPKDTLMI SRTPEVTCVVVDVS HEDPEVKFNWYVDGVVHNAKTPREEQYNSTYRVVSVLT VLHQDWLNGKEYCKVSNKALPAPIEKTISKAKGQPREPQ VYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP ENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQQGNVFSQCSV MHEALHNHYTQKSLSLSPGK	104
NRP2 a2b1b2-Fc	145-595	GSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAKPK MEIILQFLIFDLEHDPQVGECDCKYDWLDIWDGIPHVGP LIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYL VHQEPL ENFQCNVPLGMESGRIANEQISASSTYSDGRW TPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAIAT QGAISRETQNGYVVKSYKLEVSTNGEDWMVYRHGKNHKV	105

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
		QANNDATEVVLNKLHAPLLTRFVRIRPQTWHSGIALRLEL FGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSPSA ARLVSSRSGWFPRIPQAQPGEEWLQVDLGTPTVKGVIIQ GARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPRTQ QPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAGIG MRLEVLGCDWTDKTHTCPPCAPELLGGPSVFLFPPKPKD TLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALP APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLV KGFYPSDIAVEWESNGQPENNYKTTPPVLDSDGSFFLYSK LTVDKSRWQQGNVFSQVMHEALHNHYTQKSLSLSPGK	
NRP2 b2c combined domains	438-794	LGMLSGLIADSIASSTQEYLWSPSAARLVSSRSGWFPR IPQAQPGEEWLQVDLGTPTVKGVIIQGARGGDSITAVEA RAFVRKFKVSYSLNGKDWEYIQDPRTQPKLFEGNMHYDTP DIRRFDPIPAQYVRVYPERWSPAGIMRLEVLGCDWTD SKPTVETLGPTVKSEETTPYPTTEEEATECGENCSEDDKD LQLPSGFNCNFDLFLEPCGWMYDHAKWLRTTWASSSSPND RTFPDDRNFRLRLQSDSQREGQYARLISPPVHLPRSPVCME FQYQATGGRGVALQVVREASQESKLLWVIREDDQGGWKHG RIILPSYDMEYQIVFEGVIGKRSGEIAIDDIRISTD	106
NRP2 b1b2c combined domains	276-794	QCNVPLGMESGRIANEQISASSTYSRGRWTPQQSRLHGDD NGWTPNLDNKEYLQVDLRLTMTLTAIATQGAISRETQNG YYVKSYLEVSTNGEDWMVYRHGKNHKVFOANNDATEVVL NKLHAPLLTRFVRIRPQTWHSGIALRLELFGCRVTDAPCS NMLGMLSGLIADSIASSTQEYLWSPSAARLVSSRSGWF PRIPQAQPGEEWLQVDLGTPTVKGVIIQGARGGDSITAVEA RAFVRKFKVSYSLNGKDWEYIQDPRTQPKLFEGNMHYDTP DIRRFDPIPAQYVRVYPERWSPAGIMRLEVLGCDWTD SKPTVETLGPTVKSEETTPYPTTEEEATECGENCSEDDKD LQLPSGFNCNFDLFLEPCGWMYDHAKWLRTTWASSSSPND RTFPDDRNFRLRLQSDSQREGQYARLISPPVHLPRSPVCME FQYQATGGRGVALQVVREASQESKLLWVIREDDQGGWKHG RIILPSYDMEYQIVFEGVIGKRSGEIAIDDIRISTD	107
NRP2 a2b1b2c combined domains	149-802	CSKNFTSPNGTIESPGFPEKYPHNLDCTFTILAKPKMEII LQFLIFDLEHDPQVGEQDCKYDWDLDWDGIPHVGPLIGK YCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSAARYLVH QEPLNFQCNVPLGMESGRIANEQISASSTYSRGRWTPQQ SRLHGDDNGWTPNLDNKEYLQVDLRLTMTLTAIATQGAIS RETQNGYYVKSYLEVSTNGEDWMVYRHGKNHKVFOANN DATEVVLNKLHAPLLTRFVRIRPQTWHSGIALRLELFGCR VTDAPCSNMLGMLSGLIADSIASSTQEYLWSPSAARLV SSRSGWFPRIPQAQPGEEWLQVDLGTPTVKGVIIQGARG GDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPRTQPKL FEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAGIMRLE VLGCDWTDKPTVETLGPTVKSEETTPYPTTEEEATECGE NCSFEDDKDLQLPSGFNCNFDLFLEPCGWMYDHAKWLRT TWASSSSPNDRTFPDDRNFRLRLQSDSQREGQYARLISPPV HLPRSPVCMEFQYQATGGRGVALQVVREASQESKLLWVIRE DDQGGWKHGRRIILPSYDMEYQIVFEGVIGKRSGEIAIDD IRISTDVPLENCME	108
NRP2 a1a2b1b2c combined domains	23-802	QPDPPCGRLNSKDAGYITSPGYPQDYPSHQNCWEIVYAP EPNQKIVLNFNPHFEIEKHDCYDFIEIRDGDESADLLG KHCGNIAPPTIISSGSMLYIKFTSDYARQAGFSLRYEIF KTGSEDCSKNFTSPNGTIESPGFPEKYPHNLDCTFTILAK PKMEIILQFLIFDLEHDPQVGEQDCKYDWDLDWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSRGR WTPQQSRLHGDDNGWTPNLDNKEYLQVDLRLTMTLTAI ATQGAISRETQNGYYVKSYLEVSTNGEDWMVYRHGKNHK VFQANNDATEVVLNKLHAPLLTRFVRIRPQTWHSGIALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSGWFPRIPQAQPGEEWLQVDLGTPTVKGVII QGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPRT QPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAGI MRLEVLGCDWTDKPTVETLGPTVKSEETTPYPTTEEEATE CGENCSEDDKDQLQLPSGFNCNFDLFLEPCGWMYDHAK WLRTTWASSSSPNDRTFPDDRNFRLRLQSDSQREGQYARL ISPPVHLPRSPVCMEFQYQATGGRGVALQVVREASQESKL LWVIREDDQGGWKHGRRIILPSYDMEYQIVFEGVIGKRS GEIAIDDIRISTDVPLENCME	109

TABLE N1-continued

Exemplary Human NRP2 Polypeptides			
Name	Residues	Sequence	SEQ ID NO:
NRP2a a1a2b1b2c combined domains + juxtamembrane	23-859	QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPFHEIEKHDCKYDFIEIRDGDSSESADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVTNGEDWMVYRHGKNHK VFQANNDATFVVLNKLHAPLLTRFVRIRPQTWHSGLALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSWGFPRIPQAQPGEEWLQVDLGTPTKTVKGI IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLVGLGCDWTDKPTVETLGTPTVKSSEETTPYPTTEE ATECGENCSFEDDKDLQLPSGFNCNFDLEPCGWMYDHA KWLRTTWASSSSPNDRTPDDRNLRLQSDSQREGQYARL ISPPVHLPRSPVCMFQYQATGGRGVALQVVRASQESKL LWVIREQGGGEWKHGRIILPSYDMEYQIVFEGVIGKGRSG EIAIDDIRISTDVPLENCMEPISAFAVDIPEIHEREGYED EIDDEYEVDWSNSSSATSGSGAPSTDKEKSWLYTLDP	110
NRP2b a1a2b1b2c combined domains + juxtamembrane		QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPFHEIEKHDCKYDFIEIRDGDSSESADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVTNGEDWMVYRHGKNHK VFQANNDATFVVLNKLHAPLLTRFVRIRPQTWHSGLALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP SAARLVSSRSWGFPRIPQAQPGEEWLQVDLGTPTKTVKGI IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR TQQPKLFEGNMHYDTPDIRRFDPIPAQYVRVYPERWSPAG IGMRLVGLGCDWTDKPTVETLGTPTVKSSEETTPYPTTEE ATECGENCSFEDDKDLQLPSGFNCNFDLEPCGWMYDHA KWLRTTWASSSSPNDRTPDDRNLRLQSDSQREGQYARL ISPPVHLPRSPVCMFQYQATGGRGVALQVVRASQESKL LWVIREQGGGEWKHGRIILPSYDMEYQIVFEGVIGKGRSG EIAIDDIRISTDVPLENCMEPISAFAGGTLPLGTEPTVDT VPMQPIPAY	111
NRP2a Juxtamembrane Variant 1	803-864	PISAFAGENFKVDIPEIHEREGYEDEIDDEYEVDWSNSSS ATSGSGAPSTDKEKSWLYTLDP	112
NRP2a Juxtamembrane Variant 2	803-859	PISAFAVDIPEIHEREGYEDEIDDEYEVDWSNSSSATSGS GAPSTDKEKSWLYTLDP	113
NRP2a Juxtamembrane Variant 3	803-842	PISAFADYEVDWSNSSSATSGSGAPSTDKEKSWLYTLDP	114
NRP2b Juxtamembrane Variant 4	803-837	PISAFAGENFKGGTLLPGTEPTVDTVPMQPIPAYW	115
NRP2b Juxtamembrane Variant 5	803-832	PISAFAGGTLLPGTEPTVDTVPMQPIPAYW	116
NRP2 (23-595) -Fc		QPDPPCGRLNSKDAGYITSPGYPDYPSHQNCEWIVYAP EPNQKIVLNFNPFHEIEKHDCKYDFIEIRDGDSSESADLLG KHCNIIAPPTIISSGSMLYIKFTSDYARQGAGFSLRYEIF KTGSSEDCSKNFTSPNGTIESPGFPEKYPHNLDCFTTILAK PKMEIILQFLIFDLEHDPQVGECDKYDWLDIWDGIPHV GPLIGKYCGTKTPSELRSSTGILSLTFHTDMAVAKDGFSA RYYLVHQEPLNFQCNVPLGMESGRIANEQISASSTYSYG RWTPQQSRLHGDDNGWTPNLDSNKEYLQVDLRFLTMLTAI ATQGAISRETQNGYVVKSYKLEVTNGEDWMVYRHGKNHK VFQANNDATFVVLNKLHAPLLTRFVRIRPQTWHSGLALRL ELFGCRVTDAPCSNMLGMLSGLIADSIASSTQEYLWSP	117

TABLE N1-continued

Exemplary Human NRP2 Polypeptides		
Name	Residues Sequence	SEQ ID NO:
	SAARLVSSRSRGWFPRIQPAQPGEEWLQVDLGTPTKTVKGI	
	IQGARGGDSITAVEARAFVRKFKVSYSLNGKDWEYIQDPR	
	TQQPKLFEKNMHYDTPDIRFDPIPAQYVRVYPERWSPAG	
	IGMRLEVLGCDWTDKTHTCPPCAPPELLGGPSVFLFPPKP	
	KDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNA	
	KTKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKA	
	LPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTC	
	LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLY	
	SKLTVDKSRWQQGNVFCSCVMHEALHNHYTQKSLSLSPGK	

15

In certain embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a full-length human NRP2 polypeptide or a human NRP2 polypeptide selected from Table N1. In some embodiments, the antibody or antigen-binding fragment thereof binds to the human NRP2 polypeptide with an affinity of about 10 pM to about 500 pM or to about 50 nM, or about, at least about, or no more than about 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900 pM, 1 nM, 10 nM, 25 nM, or 50 nM, or optionally with an affinity that ranges from about 10 pM to about 500 pM, about 10 pM to about 400 pM, about 10 pM to about 300 pM, about 10 pM to about 200 pM, about 10 pM to about 100 pM, about 10 pM to about 50 pM, or about 20 pM to about 500 pM, about 20 pM to about 400 pM, about 20 pM to about 300 pM, about 20 pM to about 200 pM, about 20 pM to about 100 pM, about 20 pM to about 50 pM, or about 30 pM to about 500 pM, about 30 pM to about 400 pM, about 30 pM to about 300 pM, about 30 pM to about 200 pM, about 30 pM to about 100 pM, about 30 pM to about 50 pM, or about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 1 nM to about 5 nM, about 5 nM to about 10 nM, about 10 nM to 25 nM, or about 25 nM to about 50 nM.

In some embodiments, the at least one antibody or antigen-binding fragment thereof that specifically binds to at least one epitope in at least one neuropilin domain. Exemplary neuropilin domains include one or more of the neuropilin a1 domain, neuropilin a2 domain, neuropilin b1 domain, neuropilin b2 domain, neuropilin c domain, neuropilin a1/a2 combined domain, neuropilin b1/b2 combined domain, neuropilin a2/b1 combined domain, neuropilin b2/c combined domain, neuropilin a2/b1/b2 combined domain, neuropilin a2/b1/b2/c combined domain, neuropilin a1/a2/b1 combined domain, neuropilin a1/a2/b1/b2 combined domain, neuropilin a1/a2/b1/b2/c combined domain, a1/a2/b1/b2/c/juxtamembrane combined domain, and the neuropilin b1/b2/c combined domain (see Table N1 for residues of the domains). In specific embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin b1 domain, the neuropilin b2 domain, and/or the neuropilin b1/b2 combined domain (see Table N1). In particular embodiments, the antibody or antigen-binding fragment thereof binds to the at least one domain (or at least one epitope therein) with an affinity of about 10 pM to about 500 pM or to about 50 nM, or about, at least about, or no more than about 10, 20, 30, 40,

50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 300, 400, 500, 600, 700, 800, 900 pM, 1 nM, 10 nM, 25 nM, or 50 nM, or optionally with an affinity that ranges from about 10 pM to about 500 pM, about 10 pM to about 400 pM, about 10 pM to about 300 pM, about 10 pM to about 200 pM, about 10 pM to about 100 pM, about 10 pM to about 50 pM, or about 20 pM to about 500 pM, about 20 pM to about 400 pM, about 20 pM to about 300 pM, about 20 pM to about 200 pM, about 20 pM to about 100 pM, about 20 pM to about 50 pM, or about 30 pM to about 500 pM, about 30 pM to about 400 pM, about 30 pM to about 300 pM, about 30 pM to about 200 pM, about 30 pM to about 100 pM, about 30 pM to about 50 pM, or about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 1 nM to about 5 nM, about 5 nM to about 10 nM, about 10 nM to 25 nM, or about 25 nM to about 50 nM.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin a1 domain, the neuropilin a2 domain, and/or the neuropilin a1a2 combined domain, including adjacent linker regions, for example, at about residues (neuropilin a1 domain) 20-148, 30-141, 40-141, 50-141, 60-141, 70-141, 80-141, 90-141, 100-141, 110-141, 120-141, 130-141; 20-130, 20-120, 20-110, 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, or 20-30 as defined by a human NRP2 precursor sequence (see Table N1); or, for example, at about residues (neuropilin a2 domain) 142-280, 150-265, 160-265, 170-265, 180-265, 190-265, 200-265, 210-265, 220-265, 230-265, 240-265, 250-265, 260-265, 141-270, 141-260, 141-250, 141-240, 141-230, 141-220, 141-210, 141-200, 141-190, 141-180, 141-170, 141-160, 141-150, 200-250, 210-250, 220-250, 230-250, 200-240, 210-240, 220-240, 230-240, 227-247, 228-247, 229-247, 230-247, 231-247, 232-247, 233-247, 234-247, 235-247, 236-247; 227-246, 227-245, 227-244, 227-243, 227-242, 227-241, 227-240, 227-239, 227-238; 235-240, 236-239, 236-238, or residue 237 as defined by a human NRP2 precursor sequence (see Table N1); or, for example, at about residues (combined a1a2 domain) 20-280, 30-280, 40-280, 50-280, 60-280, 70-280, 80-280, 90-280, 100-280, 110-280, 120-280, 130-280, 140-280, 150-280, 160-280, 170-280, 180-280, 190-280, 200-280, 210-280, 220-280, 230-280, 240-280, 260-280, 270-280, 20-270, 20-260, 20-250, 20-240, 20-230, 20-220, 20-210, 20-200, 20-190, 20-180, 20-170, 20-160, 20-150, 20-140, 20-130, 20-120, 20-110, 20-100, 20-90, 20-80, 20-70, 20-60, 20-50, 20-40, or 20-30 as defined by a human NRP2 precursor sequence (see Table N1).

In particular embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin b1 domain, the neuropilin b2 domain, and/or the neuropilin b1/b2 combined domain, including adjacent linker regions, for example, at about residues (neuropilin b1 domain) 266-426, 280-426, 290-426, 299-420, 300-426, 310-426, 320-426, 330-426, 340-426, 350-426, 360-426, 370-426, 380-426, 390-426, 400-426, 410-426, 420-426, 280-420, 280-410, 280-400, 280-390, 280-380, 280-370, 280-360, 280-350, 280-340, 280-330, 280-320, 280-310, 280-300, or 280-290 as defined by a human NRP2 precursor sequence (see Table N1), including a discontinuous epitope that comprises one, two, or three of residues Y299, N354, and/or S416 as defined by the human NRP2 precursor sequence; (neuropilin b2 domain) 438-591, 450-591, 460-591, 470-591, 480-591, 490-591, 500-591, 510-591, 520-591, 530-591, 540-591, 550-591, 560-591, 570-591, 580-591, 438-590, 438-580, 438-570, 438-560, 438-550, 438-540, 438-530, 438-520, 438-510, 438-500, 438-490, 438-480, 438-470, 438-460, 438-450 as defined by a human NRP2 precursor sequence (see Table N1); or (neuropilin b1/b2 combined domain) 266-591, 276-591, 286-591, 296-591, 306-591, 316-591, 326-591, 336-591, 346-591, 356-591, 366-591, 376-591, 386-591, 396-591, 406-591, 416-591, 426-591, 436-591, 446-591, 456-591, 466-591, 476-591, 486-591, 498-591, 508-591, 518-591, 528-591, 538-591, 548-591, 558-591, 568-591, 578-591, 588-591, 266-581, 266-571, 266-561, 266-551, 266-541, 266-531, 266-521, 266-511, 266-501, 266-491, 266-481, 266-471, 266-461, 266-451, 266-441, 266-431, 266-421, 266-411, 266-401, 266-391, 266-381, 266-371, 266-361, 266-351, 266-341, 266-331, 266-321, 266-311, 266-301, 266-291, 266-281, or 266-271 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin a2/b1 combined domain and/or the neuropilin b2c combined domain, including adjacent linker regions, for example, at about residues (neuropilin a2b1 combined domain) 149-437, 159-426, 169-426, 179-426, 189-426, 199-426, 209-426, 219-426, 229-426, 239-426, 249-426, 259-426, 269-426, 279-426, 289-426, 299-426, 309-426, 319-426, 329-426, 339-426, 349-426, 359-426, 369-426, 379-426, 389-426, 399-426, 409-426, 419-426, 149-436, 149-426, 149-416, 149-406, 149-396, 149-386, 149-376, 149-366, 149-356, 149-346, 149-336, 149-326, 149-316, 149-306, 149-296, 149-286, 149-276, 149-266, 149-256, 149-246, 149-236, 149-226, 149-216, 149-206, 149-196, 146-186, 146-176, 146-166, or 146-155 as defined by a human NRP2 precursor sequence (see Table N1); or, for example, at about residues (neuropilin b2c combined domain) 438-794, 448-794, 458-794, 468-794, 478-794, 487-794, 497-794, 507-794, 517-794, 527-794, 537-794, 547-794, 557-794, 567-794, 587-794, 597-794, 607-794, 617-794, 627-794, 637-794, 647-794, 657-794, 667-794, 677-794, 687-794, 697-794, 707-794, 717-794, 727-794, 737-794, 747-794, 757-794, 767-794, 777-794, 787-794, 427-794, 438-784, 438-774, 438-764, 438-754, 438-744, 438-734, 438-728, 438-714, 438-704, 438-694, 438-684, 438-674, 438-664, 438-654, 438-644, 438-634, 438-624, 438-614, 438-604, 438-596, 438-586, 438-576, 438-566, 438-556, 438-546, 438-536, 438-526, 438-516, 438-506, 438-494, 438-484, 438-474, 438-464, 438-454, 438-444 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least

one epitope in the neuropilin c domain, including adjacent linker regions, for example, at about residues 591-794, 600-794, 610-794, 620-794, 630-794, 640-794, 650-794, 660-794, 670-794, 680-794, 690-794, 700-794, 710-794, 720-794, 730-794, 740-794, 750-794, 760-794, 770-794, 780-794, 790-794, 591-790, 591-780, 591-770, 591-760, 591-750, 591-740, 591-730, 591-720, 591-710, 591-700, 591-690, 591-680, 591-670, 591-660, 591-650, 591-640, 591-630, 591-620, 591-610, or 591-600 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin b1/b2c combined domain, including adjacent linker regions, for example, at about residues 276-794, 286-794, 296-794, 306-794, 316-794, 326-794, 336-794, 346-794, 356-794, 366-794, 376-794, 387-794, 396-794, 406-794, 416-794, 426-794, 436-794, 446-794, 456-794, 466-794, 476-794, 486-794, 496-794, 506-794, 516-794, 526-794, 536-794, 546-794, 556-794, 566-794, 576-794, 586-794, 596-794, 606-794, 616-794, 626-794, 636-794, 646-794, 656-794, 666-794, 676-794, 686-794, 696-794, 706-794, 716-794, 726-794, 736-794, 746-794, 756-794, 766-794, 776-794, 786-794, 266-794, 276-784, 276-774, 276-764, 276-754, 276-744, 276-734, 276-724, 276-714, 276-704, 276-694, 276-684, 276-674, 276-664, 276-654, 276-644, 276-634, 276-624, 276-614, 276-604, 276-594, 276-584, 276-574, 276-564, 276-554, 276-544, 276-534, 276-524, 276-514, 276-504, 276-496 as defined by a human NRP2 precursor sequence (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin C/juxtamembrane combined domain.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin juxtamembrane domain of NRP2a (variant 1). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin juxtamembrane domain of NRP2a (variant 2). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin juxtamembrane domain of NRP2a (variant 3). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin juxtamembrane domain of NRP2b (variant 4). In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to at least one epitope in the neuropilin juxtamembrane domain of NRP2b (variant 5).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a conformational epitope composed of two or more discontinuous epitope regions. In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to a conformational epitope comprising or consisting of:

- (a) a first epitope region within the a1 domain, and second epitope region within the a2 domain of the human NRP2 polypeptide;
- (b) a first epitope region within the a1 domain, and second epitope region within the b1 domain of the human NRP2 polypeptide;

75

- (c) a first epitope region within the a1 domain, and second epitope region within the b2 domain of the human NPR2 polypeptide;
- (d) a first epitope region within the a1 domain, and second epitope region within the c domain of the human NPR2 polypeptide;
- (e) a first epitope region within the a1 domain, and second epitope region within the juxtamembrane domain of the human NPR2 polypeptide selected from variant 1, 2, 3, 4 and 5;
- (f) a first epitope region within the a2 domain, and second epitope region within the b1 domain of the human NPR2 polypeptide;
- (g) a first epitope region within the a2 domain, and second epitope region within the b2 domain of the human NPR2 polypeptide;
- (h) a first epitope region within the a2 domain, and second epitope region within the c domain of the human NPR2 polypeptide;
- (i) a first epitope region within the a2 domain, and second epitope region within the juxtamembrane domain of the human NPR2 polypeptide selected from variant 1, 2, 3, 4 and 5;
- (j) a first epitope region within the b1 domain, and second epitope region within the b2 domain of the human NPR2 polypeptide;
- (k) a first epitope region within the b1 domain, and second epitope region within the c domain of the human NPR2 polypeptide;
- (l) a first epitope region within the b1 domain, and second epitope region within the juxtamembrane domain of the human NPR2 polypeptide selected from variant 1, 2, 3, 4 and 5;

76

- (m) a first epitope region within the b2 domain, and second epitope region within the c domain of the human NPR2 polypeptide;
- (n) a first epitope region within the b2 domain, and second epitope region within the juxtamembrane domain of the human NPR2 polypeptide selected from variant 1, 2, 3, 4 and 5; or
- (o) a first epitope region within the c domain, and second epitope region within the juxtamembrane domain of the human NPR2 polypeptide selected from variant 1, 2, 3, 4 and 5.

In some embodiments, the antibody or antigen-binding fragment thereof specifically binds to at least one epitope within a region of a human NRP2 polypeptide that binds to or interacts with at least one “NRP2 ligand”, including any molecule that interacts with or binds reversibly to human NRP2, including any one or more variants of human NRP2. General examples of “NRP2 ligands” include polypeptides such as HRS polypeptides, soluble ligands, receptors (e.g., cell surface receptors), including growth factors, growth factor receptors, and others, and specific examples of NRP2 ligands are detailed herein. In some embodiments, the at least one antibody or antigen-binding fragment thereof modulates (e.g., antagonizes, interferes with, agonizes, enhances) binding of the human NRP2 polypeptide to at least one “NRP2 ligand”.

As noted above, in certain embodiments the at least one NRP2 ligand is an HRS polypeptide. Thus, in certain embodiments, an antibody or antigen-binding fragment thereof specifically binds to at least one epitope within a region of a human NRP2 polypeptide that binds to or interacts with at least one human HRS polypeptide, and thereby modulates binding of the human NRP2 polypeptide to the human HRS polypeptide. Exemplary HRS polypeptides are provided in Table H1 below.

TABLE H1

Exemplary Human HRS polypeptides			
Name	Residues	Sequence	SEQ ID NO:
FL cytosolic wild type	1-509	MAERAALEELVKLGGERVRLKQKASAELEIEEVAKLLKL KAQLGPDESKQKFLKTPKGTRDYSRQMAVREKVFVDVIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKLIYDLKDQGGG LLSLRYDLTVPFARYLAMNKL TNIKRYHIAKVYRRDNPAMT RGRYREFYQCFDIAGNFDPMIPDAECLKIMCEILSSLQIG DFLVKVNDRILDGMFAICGVSDSKFRTICSSVDKLDKVS EEVKNEVMGEGKLAPEVADRIGDYVQQHGGVSLVEQLLQDP KLSQNKQALEGLGDLKLLFEYLTFLGIDDKISFDLSLARGL DYTGVIYEAVLLQTPAQAGEEPLGVGSVAAGGRYDGLVGM FDPKGRKVPVGLSIGVERIFSIQRLAELEEKIRTTETQ VLVSAQKLLLEERLKLVSLEWDAGIKAEALLYKKNPKLLNQ LQYCEEAGIPLVAIIGEQELKDGVIKLRSVTSREEVDVRRE DLVEEIKRRTGQPLCIC	118
HisRS1 ^{N1}	1-141	MAERAALEELVKLGGERVRLKQKASAELEIEEVAKLLKL KAQLGPDESKQKFLKTPKGTRDYSRQMAVREKVFVDVIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKLIYDLKDQGGG LLSLRYDLTVPFARYLAM	119
HisRS1 ^{N2}	1-408	MAERAALEELVKLGGERVRLKQKASAELEIEEVAKLLKL KAQLGPDESKQKFLKTPKGTRDYSRQMAVREKVFVDVIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKLIYDLKDQGGG LLSLRYDLTVPFARYLAMNKL TNIKRYHIAKVYRRDNPAMT RGRYREFYQCFDIAGNFDPMIPDAECLKIMCEILSSLQIG DFLVKVNDRILDGMFAICGVSDSKFRTICSSVDKLDKVS EEVKNEVMGEGKLAPEVADRIGDYVQQHGGVSLVEQLLQDP KLSQNKQALEGLGDLKLLFEYLTFLGIDDKISFDLSLARGL DYTGVIYEAVLLQTPAQAGEEPLGVGSVAAGGRYDGLVGM FDPKGRKVPVGLSIGVERIFSIQRLAELEEKIRTTETQ	120

TABLE H1-continued

Exemplary Human HRS polypeptides			
Name	Residues	Sequence	SEQ ID NO:
HisRS1 ^{N3}	1-113	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKGRDYS PRQMAVREK VFDVIIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKL	121
HisRS1 ^{N4}	1-60	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPK	122
HisRS1 ^{N5}	1-243 + 27aa	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKGRDYS PRQMAVREK VFDVIIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKLIYDLKDQGG LLSLRYDLTVPPARYLAMNKL TNIKRYHIAKVYRRDNPAMT RGRYREFYQCDFDIAGNFDPMIPDAECLKIMCEILSSLQIG DFLVKVNDRRI LDGMFAICGVSDSKFRTICSSVDKLDKVG Y PWWNSCSRILNYPKTSRPRAWET	123
HisRS1 ^{C1}	405-509	RTTETQVLVASAQKKLLEERLKL VSELWDAGI KAE LLYKKN PKLLNQLQYCEEAGIPLVAI IGEQELKDGVIKLRSVTSREE VDVRREDLV E EIKRRTGQPLCIC	124
HisRS1 ^{C2}	1-60 + 175-509	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKDFDIAGNFDPMIPDAECLKIMC EILSSLQIGDFLVKVNDRRI LDGMFAICGVSDSKFRTICSS VDKLDKVSWE EVKNEMVGEKGLAPEVADRIGDYVQHGGS LVEQLLDQPKLSQNKQALEGLGDLKLLFEYLT LFGIDDKIS FDLSLARGLDYYTGVI YEAVLLQTPAQAGEEPLGVGSVAAG GRYDGLVGMFDPKGRKVP CVGLSIGVERIF SIVEQRLEALE EKIRTTETQVLVASAQKKLLEERLKL VSELWDAGI KAE LLY KKNPKLLNQLQYCEEAGIPLVAI IGEQELKDGVIKLRSVTS REEVDVRREDLV E EIKRRTGQPLCIC	125
HisRS1 ^{C3}	1-60 + 211-509	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKVNDRRI LDGMFAICGVSDSKFR TICSSVDKLDKVSWE EVKNEMVGEKGLAPEVADRIGDYVQ HGGVSLVEQLLDQPKLSQNKQALEGLGDLKLLFEYLT LFGI DDKISFDLSLARGLDYYTGVI YEAVLLQTPAQAGEEPLGVG SVAAGGRYDGLVGMFDPKGRKVP CVGLSIGVERIF SIVEQR LEALEEKIRTTETQVLVASAQKKLLEERLKL VSELWDAGI K AE LLYKKNPKLLNQLQYCEEAGIPLVAI IGEQELKDGVIKLR SVTSREEVDVRREDLV E EIKRRTGQPLCIC	126
HisRS1 ^{C4}	1-100 + 211-509	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKGRDYS PRQMAVREK VFDVIIR CFKRHGAEVIDTPVFELKVNDRRI LDGMFAICGVSDSKFRT ICSSVDKLDKVSWE EVKNEMVGEKGLAPEVADRIGDYVQ HGGVSLVEQLLDQPKLSQNKQALEGLGDLKLLFEYLT LFGID DKISFDLSLARGLDYYTGVI YEAVLLQTPAQAGEEPLGVGS VAAGGRYDGLVGMFDPKGRKVP CVGLSIGVERIF SIVEQR LEALEEKIRTTETQVLVASAQKKLLEERLKL VSELWDAGI K AE LLYKKNPKLLNQLQYCEEAGIPLVAI IGEQELKDGVIKLR SVTSREEVDVRREDLV E EIKRRTGQPLCIC	127
HisRS1 ^{C5}	1-174 + 211-509	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKGRDYS PRQMAVREK VFDVIIR CFKRHGAEVIDTPVFELKETLMGKYGEDSKLIYDLKDQGG LLSLRYDLTVPPARYLAMNKL TNIKRYHIAKVYRRDNPAMT RGRYREFYQCVNDRRI LDGMFAICGVSDSKFRTICSSVDKL DKVSWE EVKNEMVGEKGLAPEVADRIGDYVQHGGSVLEQ LLQDPKLSQNKQALEGLGDLKLLFEYLT LFGIDDKISFDLS LARGLDYYTGVI YEAVLLQTPAQAGEEPLGVGSVAAGGRYD GLVGMFDPKGRKVP CVGLSIGVERIF SIVEQRLEALEEKIR TTETQVLVASAQKKLLEERLKL VSELWDAGI KAE LLYKKNP KLLNQLQYCEEAGIPLVAI IGEQELKDGVIKLRSVTSREEV DVRREDLV E EIKRRTGQPLCIC	128
HisRS1 ^{C6}	1-60 + 101-509	MAERAAL EELVKLQGERVRLKQQKASAE LIEEEVAKLLKL KAQLGPDES KQKFV LKTPKETLMGKYGEDSKLIYDLKDQGG ELLSRYDLTVPPARYLAMNKL TNIKRYHIAKVYRRDNPAM TRGRYREFYQCDFDIAGNFDPMIPDAECLKIMCEILSSLQI GDFLVKVNDRRI LDGMFAICGVSDSKFRTICSSVDKLDKVS WEEVKNEMVGEKGLAPEVADRIGDYVQHGGSVLEQLLQD PKLSQNKQALEGLGDLKLLFEYLT LFGIDDKISFDLSLARG LDYYTGVI YEAVLLQTPAQAGEEPLGVGSVAAGGRYDGLV GMFDPKGRKVP CVGLSIGVERIF SIVEQRLEALEEKIRTTET QVLVASAQKKLLEERLKL VSELWDAGI KAE LLYKKNPKLLN	129

TABLE H1-continued

Exemplary Human HRS polypeptides			
Name	Residues	Sequence	SEQ ID NO:
		QLQYCEEAGIPLVAIIGEQLKDGVIKLRSVTSREEVDVRR EDLVEEIKRRTGQPLCIC	
HisRS1 ^{C7}	1-100 + 175-509	MAERAAL EELVKLQGERVRGLKQQKASAE LIEEEVAKLLKL KAQLGPDESKQKFV LKTPKGTRDYS PRQMAVREKVF DVIIR CFKRHGAEVIDTPVFELKDFDIAGNFDPMIPDAECLKIMCE ILSSLQIGDFLVKVNDRRILDGMFAICGVSDSKFRTICSSV DKLDKVSWE EVKNE MVGEKGLAPEVADRI GDYVQQHGGVSL VEQLLQDPKLSQNKQALEGLGDLKLLFEYLT LFGIDDKISF DLSLARGLDYYTGV IYEAVLLQTPAQAGEEPLGVGSVAAGG RYDGLVGMFDPKGRKVP CVGLSIGVERIF SIVEQRLEALEE KIRTTETQVLVSAQKKLLEERLKL VSELWDAGIKAELLYK KNPKLLNQLQYCEEAGIPLVAIIGEQLKDGVIKLRSVTSR EEVDVRREDLVEEIKRRTGQPLCIC	130
HisRS1 ^{C8}	1-60 + 399-509	MAERAAL EELVKLQGERVRGLKQQKASAE LIEEEVAKLLKL KAQLGPDESKQKFV LKTPKALEEKIRTTETQVLVSAQKKL LEERLKL VSELWDAGIKAELLYKKNPKLLNQLQYCEEAGIP LVAIIGEQLKDGVIKLRSVTSREEVDVRREDLVEEIKRRT GQPLCIC	131
HisRS1 ^{C9}	1-100 + 399-509	MAERAAL EELVKLQGERVRGLKQQKASAE LIEEEVAKLLKL KAQLGPDESKQKFV LKTPKGTRDYS PRQMAVREKVF DVIIR CFKRHGAEVIDTPVFELKALEEKIRTTETQVLVSAQKKLL EERLKL VSELWDAGIKAELLYKKNPKLLNQLQYCEEAGIPL VAIIGEQLKDGVIKLRSVTSREEVDVRREDLVEEIKRRTG QPLCIC	132
HisRS1 ^{C10}	369-509	MFDPKGRKVP CVGLSIGVERIF SIVEQRLEALEEKIRTTET QVLVSAQKKLLEERLKL VSELWDAGIKAELLYKKNPKLLN QLQYCEEAGIPLVAIIGEQLKDGVIKLRSVTSREEVDVRR EDLVEEIKRRTGQPLCIC	133
HisRS1 ^{I1}	191-333	CLKIMCEILSSLQIGDFLVKVNDRRILDGMFAICGVSDSKF RTICSSVDKLDKVSWE EVKNE MVGEKGLAPEVADRI GDYVQ QHGGVSLVEQLLQDPKLSQNKQALEGLGDLKLLFEYLT LFG IDDKISFDLSLARGLDYYTG	134
FL mito. wild type	1-506	MPLLGLLPRRAWASLLS QLLRPPCASCTGAVRC SQVAEAV LTSQLKAHQEKPNFI IKT PKGTRDLSPQH MVVREKILD LVI SCFKRHGAKGMDTPAFELKETL TEKYGEDSGLMYDLKDQGG ELLSLR YDLTVPFARYLAMNKVKMKRYHVGVWRRESPTI VQGRYREFCQCFDIAGQFDPMPIDAECLKIMCEILSGLQL GDFLIKVNDRI VDGMFVCGVPESKFRAICSSIDKLDKMA WKDVRHEMVVKGLAPEVADRI GDYVQCHGGVSLVEQMFQD PRLSQNKQALEGLGDLKLLFEYLT LFGIADKISFDLSLARG LDYYTGV IYEAVLLQTPAQAGEEPLNVGSVAAGGRYDGLVG MFDPKGHKVP CVGLSIGVERIF YIVEQRMKTKEKVRTTET QVFVATPQKNFLQERLKL IAE LWDSGIKAEMLYKNNPKLLT QLHYCESTGIPLVV IIGEQLKEGV I KIRSVASREEVAIKR ENFVAEIQR LSES	135
	152-398	HVGK VWRRESPTIVQGRYREFCQCFDIAGQFDPMPIDAE LKIMCEILSGLQLGDFLIKVNDRI VDGMFVCGVPESKFR AICSSIDKLDKMAWKDVRHEMVVKGLAPEVADRI GDYVQC HGGVSLVEQMFQDPRLSQNKQALEGLGDLKLLFEYLT LFGI ADKISFDLSLARGLDYYTGV IYEAVLLQTPAQAGEEPLNVG SVAAGGRYDGLVGMFDPKGHKVP CVGLSIGVERIF YIVEQR M	136
	294-372	QALEGLGDLKLLFEYLT LFGIDDKISFDLSLARGLDYYTGV IYEAVLLQTPAQAGEEPLGVGSVAAGGRYDGLVGMFDP	137
Amino- acylation domain and anticodon binding domain	54-509	FVLKTPKGTRDYS PRQMAVREKVF DVIIRCFKRHGAEVIDT PVFELKETLMGYGEDSKLIYDLKQGGELLSLRYDLTVPF ARYLAMNKL TNIKRYHIAKVYRRDNPAMTRGRYREFYQCF DIAGNFDPMIPDAECLKIMCEILSSLQIGDFLVKVNDRRIL DGMFAICGVSDSKFRTICSSVDKLDKVSWE EVKNE MVGEK GLAPEVADRI GDYVQQHGGVSLVEQLLQDPKLSQNKQALEGL GDLKLLFEYLT LFGIDDKISFDLSLARGLDYYTGV IYEAVL LQTPAQAGEEPLGVGSVAAGGRYDGLVGMFDPKGRKVP CVG LSIGVERIF SIVEQRLEALEEKIRTTETQVLVSAQKKLLE ERLKL VSELWDAGIKAELLYKKNPKLLNQLQYCEEAGIPLV AIIIGEQLKDGVIKLRSVTSREEVDVRREDLVEEIKRRTGQ PLCIC	138

TABLE H1-continued

Exemplary Human HRS polypeptides			
Name	Residues	Sequence	SEQ ID NO:
Amino-acylation domain	54-398	FVLKTPKGTRDYSPRQMAVREKVFVDVIRCFKRHGAEVIDT PVFELKETLMGKYGEDSKLIYDLKDQGGELLRLRYDLTVPF ARYLAMNKLTKRYHIAKVYRRDNPAMTRGRYREFYQCDF DIAGNFDPMIPDAECLKIMCEILSSLQIGDFLVKVNDRIIL DGMFAICGVSDSKFRTICSSVDKLDKVSWEVKNEMVGEKG LAPEVADRIGDYVQQHGGVSLVEQLLQDPKLSQNKQALEGL GDLKLLFEYLTFLFGIDDKISFDLSLARGLDYYTGVIYEAVL LQTPAQAGEEPLGVGSVAAGGRYDGLVGMFDPKGRKVPVCG LSIGVERIFSIVEQRLE	139
Amino-acylation (core) domain	61-398	GTRDYSPRQMAVREKVFVDVIRCFKRHGAEVIDTPVFELKE TLMGKYGEDSKLIYDLKDQGGELLRLRYDLTVPFARYLAMN KLTNIAKVYRRDNPAMTRGRYREFYQCDFDIAGNFD PMIPDAECLKIMCEILSSLQIGDFLVKVNDRIILDGMFAIC GVSDSKFRTICSSVDKLDKVSWEVKNEMVGEKGLAPEVAD RIGDYVQQHGGVSLVEQLLQDPKLSQNKQALEGLGDLKLLF EYLTFLFGIDDKISFDLSLARGLDYYTGVIYEAVLLQTPAQ GEEPLGVGSVAAGGRYDGLVGMFDPKGRKVPVCGLSIGVER IFSIVEQRLE	140
Anticodon binding domain	399-509	ALEEKIRTTETQVLVASAQKKLLEERLKLSELWDAGIKAE LLYKKNPKLLNQLQYCEEAGIPLVAIIGEQLKDGVIKLRS VTSREEVDVRRREDLVVEIKRRTGQPLCIC	141
Anticodon binding (core) domain	406-501	TTETQVLVASAQKKLLEERLKLSELWDAGIKAE LLYKKNP KLLNQLQYCEEAGIPLVAIIGEQLKDGVIKLRSVTSREEV DVRREDLVVEIKRR	142
HRS WHEP consensus		X_A -L- X_B -Q-G-X-X-V-R-X-L-K-X-X-K-A- X_C -V-X- X-L-L-X-L-K- X_D Where: X is any amino acid X_A is 0-50 amino acids X_B is about 5-7 amino acids, preferably 6 amino acids X_C is about 7-9 amino acids, preferably 8 amino acids X_D is 0-50 amino acids	143

Thus, in certain embodiments, the at least one NRP2 ligand is selected from Table H1, and the anti-NRP2 antibody or antigen-binding fragment thereof modulates (e.g., interferes with) binding of a human NRP2 polypeptide (for example, a human NRP2 polypeptide selected from Table N1) to a human HRS polypeptide selected from Table H1. In some embodiments, the anti-NRP2 antibody or antigen-binding fragment specifically binds to an HRS polypeptide-interacting region of the NRP2 polypeptide, and in some instances mimics one or more signaling activities of the HRS polypeptide binding to the NRP2 polypeptide, for example, as an agonist antibody. An "HRS polypeptide-interacting region" includes a region or domain of a human NRP2 polypeptide that interacts with a region or domain of human HRS polypeptide, for example, at a ligand binding site for a different NRP2 ligand (examples of which are provided herein), a dimerization domain, a protein-protein interaction domain, or at a site which is allosterically sensitive within a NRP2 polypeptide to modulate the activity of the NRP2 polypeptide.

In certain embodiments, an antibody or antigen-binding fragment thereof is a "blocking antibody", which fully or substantially inhibits the binding between a human NRP2

polypeptide (selected, for example, from Table N1) and an NRP2 ligand such as a human HRS polypeptide (selected, for example, from Table H1) or other NRP2 ligand (for example, selected from Table N2 or Table N3). In some embodiments, a "blocking antibody" inhibits about or at least about 80-100% (e.g., 80, 85, 90, 95, or 100%) of the theoretical maximal binding between the NRP2 polypeptide and the NRP2 ligand (for example, HRS polypeptide) after pre-incubation of the "blocking antibody" with the NRP2 polypeptide in a substantially stoichiometrically equivalent amount. As used herein, a "stoichiometrically equivalent amount" refers to a situation where the number of moles of one substance (e.g., anti-NRP2 antibody) is equivalent or substantially equivalent to the number of moles at least one other substance (e.g., NRP2 polypeptide) in a given equation or reaction.

In certain embodiments, an antibody or antigen-binding fragment thereof is a "partial-blocking antibody", which at least partially but not fully inhibits the binding between a human NRP2 polypeptide (selected, for example, from Table N1) and an NRP2 ligand such as a human HRS polypeptide (selected, for example, from Table H1) or other NRP2 ligand (for example, selected from Table N2 or Table

83

N3). In some embodiments, a “partial-blocking antibody” inhibits about or at least about 20-80% (e.g., 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, or 80%) of the theoretical maximal binding between the NRP2 polypeptide and the NRP2 ligand (for example, HRS polypeptide) after pre-incubation of the “partial-blocking antibody” with the NRP2 polypeptide in a stoichiometric amount.

In specific embodiments, the at least one antibody or antigen-binding fragment thereof specifically inhibits or otherwise reduces the binding between a human NRP2 polypeptide and a HRS polypeptide splice variant selected from Table H1, for example, a HRS splice variant selected from one or more of HisRS^{N1}, HisRS^{N2}, HisRS^{N3}, HisRS^{N4} (SV9), HisRS^{N5}, HisRS^{C1}, HisRS^{C2}, HisRS^{C3}, HisRS^{C4}, HisRS^{C5}, HisRS^{C6}, HisRS^{C7}, HisRS^{C8} (SV11), and HisRS^{C9} (SV14).

As noted above, NRP2 interacts with multiple NRP2 ligands other than HRS, which mediate downstream signaling events. Additional examples of NRP2 ligands are provided in Table N2 and Table N3 below.

TABLE N2

Exemplary Neuropilin Ligands		
Ligand	NRP1	NRP2
VEGF-A121	+	
VEGF-A145		+
VEGF-A165	+	+

84

TABLE N2-continued

Exemplary Neuropilin Ligands		
Ligand	NRP1	NRP2
VEGF-B167	+	
VEGF-C	+	+
VEGF-D	+	+
VEGF-E	+	
10 PIGF-2	+	+
VEGFR	+R1 and R2	+R1, R2, R3
Heparin	+	+
Sema3A	+	
Sema3B, C, D, F, and G	+	+
15 Plexins A1, A2, A3, A4, D1	+	+
GIPC1, 2, and 3	+	+
TGF- β 1, β 2, and β 3 receptors, and LAP	+	+
TbRI and TbRII	+	+
FGF-1, 2, 4, and 7	+	+
20 FGF receptor 1	+	+
Hepatocyte growth factor receptors		+
Integrins (see Table N3)	+	+
Fibronectin	+	
Galectin-1 and Galectin Receptors	+	+
25 Li-CAM	+	+
Glat-1	+	
HRS polypeptides (see Table H1)		+

TABLE N3

Vertebrate integrins as NRP2 ligands			
Name	Synonyms	Distribution	Ligands
$\alpha_1\beta_1$	VLA-1	Many	Collagens, laminins
$\alpha_2\beta_1$	VLA-2	Many	Collagens, laminins
$\alpha_3\beta_1$	VLA-3	Many	Laminin-5
$\alpha_4\beta_1$	VLA-4	Hematopoietic cells	Fibronectin, VCAM-1
$\alpha_5\beta_1$	VLA-5;	widespread	fibronectin and proteinases
	fibronectin receptor		
$\alpha_6\beta_1$	VLA-6;	widespread	laminins
	laminin receptor		
$\alpha_7\beta_1$		muscle, glioma	laminins
$\alpha_L\beta_2$	LFA-1	T-lymphocytes	ICAM-1, ICAM-2
$\alpha_M\beta_2$	Mac-1, CR3	Neutrophils and monocytes	Serum proteins, ICAM-1
$\alpha_{IIb}\beta_3$	Fibrinogen receptor;	Platelets	fibrinogen, fibronectin ^[24]
	gpIIbIIIa		
$\alpha_v\beta_1$		ocular melanoma; neurological tumors	vitronectin; fibrinogen
$\alpha_v\beta_3$	vitronectin receptor	activated endothelial cells, melanoma, glioblastoma	vitronectin, fibronectin, fibrinogen, osteopontin, Cyr61, thyroxine, TETRAC
$\alpha_v\beta_5$		widespread, esp. fibroblasts, epithelial cells	vitronectin and adenovirus
$\alpha_v\beta_6$		proliferating epithelia, esp. lung and mammary gland	fibronectin; T6E β 1 + 3
$\alpha_v\beta_8$		neural tissue; peripheral nerve	fibronectin; T6E β 1 + 3
$\alpha_6\beta_4$		Epithelial cells	Laminin

Thus, in certain embodiments, the at least one NRP2 ligand is selected from Table N2 and/or Table N3.

For example, in some aspects, the at least one NRP2 ligand is a VEGF (vascular endothelial growth factor) ligand selected from VEGF-A145, VEGF-A165, VEGF-C, VEGF-D, and PlGF-2. VEGF-VEGFR2/3-NRP2 interactions are associated with promoting cell migration, cell growth, cell survival, and cell attachment, and also with lymphangiogenesis, increasing vascular permeability, activating integrin signaling, promoting vesicular trafficking and internalization, and slowing cellular differentiation. Accordingly, anti-NRP2 antibodies which modulate VEGF related NRP2 ligands would be expected find utility in modulating one or more of these pathways.

In certain aspects, the at least one NRP2 ligand is a semaphorin selected from one or more of SEMA-3B, SEMA-3C, SEMA-3D, SEMA-3F, and SEMA-3B, or a plexin receptor selected from one or more of plexins A1, A2, A3, A4, and D1. SEMAs typically antagonize the effects of VEGF-C, through there is a close dynamic interplay between VEGF and Sema signaling pathways. SEMAs typically function in the immune system to control cell movement, cell migration, cell-cell communication, and cell activation. SEMA Plexin-NRP2 interactions are associated with inhibiting cell migration, inhibiting cell growth, promoting apoptosis, inhibiting cell attachment, inhibiting integrin signaling, promoting cellular differentiation, inhibiting lymphangiogenesis, reducing vascular permeability, promoting microtubule destabilization, mediating the collapse of actin cytoskeleton & cell contraction including growth cone collapse and actomyosin contraction, and preventing neuronal cell spreading and inhibiting axon outgrowth. Accordingly, anti-NRP2 antibodies which modulate SEMA-related NRP2 ligands would be expected find utility in modulating one or more of these pathways.

In some aspects, the at least one NRP2 ligand is an integrin selected from one or more of $\alpha V\beta 1$, $\alpha V\beta 3$, $\alpha V\beta 5$, $\alpha V\beta 6$, $\alpha V\beta 8$, $\alpha 6\beta 1$ and $\alpha 6\beta 4$. Integrin-NRP2 interactions are generally associated with increased cell adhesion, cell growth, cancer growth and invasiveness. Accordingly anti-NRP2 antibodies which modulate integrin related NRP2 ligands would be expected find utility in modulating one or more of these pathways.

In some aspects, the at least one NRP2 ligand is selected from TGF β 1, TGF β 2, TGF β 3, and their corresponding TGF β receptors. TGF- β signaling is strongly involved in the regulation of EMT in cancer, and also in fibrosis development (see, for example, Gemmill et al., Sci. Signal. 10 eaag0528, 2017). NRP2b expression is preferentially upregulated by TGF- β signaling in abnormal lungs, and shows little or no expression in normal lung. NRP2b expression enhances migration, invasion, metastasis, chemoresistance, and tumorsphere formation, and also enhances acquired EGFR inhibitor resistance associated with EMT in cancer cells. Accordingly, anti-NRP2 antibodies which modulate TGF- β related NRP2 ligands would be expected find utility in modulating one or more of these pathways, and find utility in the treatment of cancer chemoresistance. In certain embodiments, an anti-NRP2 antibody or antigen-binding fragment thereof modulates binding/signaling activity between an NRP2 polypeptide and at least one of the NRP2 ligands from Table N2 and/or Table N3, for example, by specifically binding to an NRP2 ligand-interacting region of the NRP2 polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to the NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table N1) of NRP2,

and does not substantially bind to the NRP2b isoform (e.g., variants 4 and/or 5 of Table N1) of NRP2. In some embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to the NRP2b isoform (e.g., variants 4 and/or 5 of Table N1), and does not substantially bind to the NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table N1) of NRP2.

In some instances, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand. For example, in some embodiments, the anti-NRP2 antibody antagonizes or reduces the theoretical maximal binding/signaling between the NRP2 polypeptide and the NRP2 ligand by about or at least about 20-100% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, or 100%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptide/ligand in a substantially stoichiometrically equivalent amount.

In some instances, the at least one antibody or antigen-binding fragment thereof reduces or inhibits the dimerization between two NRP2 polypeptides. For example, in some embodiments, the anti-NRP2 antibody antagonizes or reduces the theoretical maximal dimerization between two NRP2 polypeptides by about or at least about 20-100% (e.g., about 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 90, or 100%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptides in a substantially stoichiometrically equivalent amount.

In some instances, the at least one antibody or antigen-binding fragment thereof agonizes or enhances the dimerization between two NRP2 polypeptides. For instance, in some embodiments, the anti-NRP2 antibody agonizes or enhances the basal dimerization state of two NRP2 polypeptides by about or at least about 20%-500% (e.g., about 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptides in a substantially stoichiometrically equivalent amount.

In some instances, the at least one antibody or antigen-binding fragment thereof agonizes or enhances the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand. For instance, in some embodiments, the anti-NRP2 antibody agonizes or enhances the theoretical maximal binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand by about or at least about 20%-500% (e.g., about 20, 30, 40, 50, 60, 70, 80, 90, 100, 200, 300, 400 or 500%) after pre-incubation of the anti-NRP2 antibody with the NRP2 polypeptide in a substantially stoichiometrically equivalent amount.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of semaphorins to, or via, the NRP2 polypeptide. In some aspects, such antibodies do not substantially block the interaction of VEGF-C or related NRP2 ligands. In some aspects, such antibodies are agonistic antibodies with respect to semaphorin signaling. In some aspects, such antibodies are antagonistic antibodies with respect to semaphorin signaling.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of VEGF-C or related NRP2 ligands to, or via, the NRP2 polypeptide. In some aspects, such antibodies do not substantially block the interaction of semaphorins. In some embodiments, such antibodies selectively modulate both the binding of VEGF-C or related NRP2 ligands and semaphorins to the NRP2 polypeptide. In some embodiments, such antibodies are agonistic antibodies

with respect to VEGF-C signaling. In some aspects, such antibodies are antagonistic antibodies with respect to VEGF-C signaling.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of integrins or related NRP2 ligands to the NRP2 polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of TGF β 1, TGF β 2, TGF β 3, or their corresponding TGF β receptors to the NRP2 polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor, and/or their corresponding receptors to the NRP2 polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and VEGFR2, VEGFR3, and/or VEGF-C.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a HRS polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a HRS polypeptide, and without substantially modulating the binding/signaling activity between the NRP2 polypeptide and VEGFR2, VEGFR3, and/or VEGF-C.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and VEGFR2 and/or VEGFR3 without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and VEGFR2, VEGFR3, and/or VEGF-C without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a HRS polypeptide.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and a plexin receptor without substantially modulating the ligand binding of semaphorin 3 to NRP2.

In some embodiments, the plexin receptor is selected from plexin A1, A2, A3, A4, and D1. In some embodiments, the semaphorin is selected from semaphorin 3B, 3C, 3D, 3F, and 3G.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 amino acids within the human NRP2 a2 domain, wherein the at least one antibody or antigen-binding fragment thereof selectively inhibits receptor dimerization between NRP2 and plexin A1 without substantially inhibiting dimerization between NRP2 and FLT4 (VEGFR3). In some embodiments, the at least one antibody or antigen-

binding fragment thereof specifically binds to an epitope within amino acids 232-242 of a human NRP2 precursor (see Table N1).

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 amino acids within the human NRP2 b1 domain, wherein the at least one antibody or antigen-binding fragment thereof selectively inhibits receptor dimerization between NRP2 and FLT4 (VEGFR3) without substantially inhibiting dimerization between NRP2 and plexin A1.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 amino acids within the human NRP2 b2 domain, wherein the at least one antibody or antigen-binding fragment thereof inhibits receptor dimerization between NRP2 and FLT4 (VEGFR3) and inhibits dimerization between NRP2 and plexin A1.

In some embodiments, the at least one antibody or antigen-binding fragment thereof specifically binds to an epitope of at least 5 amino acids within the human NRP2 c domain, wherein the at least one antibody or antigen-binding fragment thereof inhibits receptor dimerization between NRP2 and plexin A1 and partially inhibits dimerization between NRP2 and FLT4 (VEGFR3).

In some embodiments, the at least one antibody or antigen-binding fragment thereof has an affinity (K_d or EC_{50}) for each of (i) a human NRP2 polypeptide and (ii) the corresponding region of a cynomolgus monkey NRP2 polypeptide (see, for example, UniProt G7PL91), wherein the affinity for (i) and (ii) is within the range of about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, about 0.4 to about 1.2 nM, about 0.9 to about 5.5 nM, about 0.9 to about 5 nM, or about 1 nM to about 10 nM.

In some embodiments, the at least one antibody or antigen-binding fragment thereof has an affinity (K_d or EC_{50}) for each of (i) a human NRP2 polypeptide and (ii) the corresponding region of a murine NRP2 polypeptide, wherein the affinity for (i) and (ii) is within the range of about 20 pM to about 200 pM, about 30 pM to about 300 pM, about 40 pM to about 400 pM, about 50 pM to about 500 pM, about 60 pM to about 600 pM, about 70 pM to about 700 pM, about 80 pM to about 800 pM, about 90 pM to about 900 pM, about 100 pM to about 1 nM, or about 1 nM to about 10 nM.

In certain embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to a human NRP2 polypeptide (see Table N1) relative to a murine NRP2 polypeptide, for instance, where its affinity for a human NRP2 polypeptide is significantly stronger than its affinity for a murine NRP2 polypeptide, for example, by about or at least about 2, 5, 10, 20, 30, 40, 50, 100, 500, or 1000-fold or more. In particular embodiments, the at least one antibody or antigen-binding fragment thereof binds selectively to a human NRP2 polypeptide and does not substantially bind to a murine NRP2 polypeptide. Certain exemplary murine NRP2 polypeptides include the *Mus musculus* NRP2 polypeptide (see, for example, UniProt 035375).

Merely for illustrative purposes, the binding interactions between a human NRP2 polypeptide and an NRP2 ligand can be detected and quantified using a variety of routine methods, including biacore assays (for example, with appropriately tagged soluble reagents, bound to a sensor chip),

FACS analyses with cells expressing a NRP2 polypeptide on the cell surface (either native, or recombinant), immunoassays, fluorescence staining assays, ELISA assays, and microcalorimetry approaches such as ITC (Isothermal Titration calorimetry).

In certain embodiments, an antibody or antigen-binding fragment thereof comprises variant or otherwise modified Fc region(s), including those having altered properties or biological activities relative to wild-type Fc region(s). Examples of modified Fc regions include those having mutated sequences, for instance, by substitution, insertion, deletion, or truncation of one or more amino acids relative to a wild-type sequence, hybrid Fc polypeptides composed of domains from different immunoglobulin classes/subclasses, Fc polypeptides having altered glycosylation/sialylation patterns, and Fc polypeptides that are modified or derivatized, for example, by biotinylation (see, e.g., US Application No. 2010/0209424), phosphorylation, sulfation, etc., or any combination of the foregoing. Such modifications can be employed to alter (e.g., increase, decrease) the binding properties of the Fc region to one or more particular FcRs (e.g., FcγRI, FcγRIIa, FcγRIIb, FcγRIIc, FcγRIIIa, FcγRIIIb, FcRn), its pharmacokinetic properties (e.g., stability or half-life, bioavailability, tissue distribution, volume of distribution, concentration, elimination rate constant, elimination rate, area under the curve (AUC), clearance, C_{max} , t_{max} , C_{min} , fluctuation), its immunogenicity, its complement fixation or activation, and/or the CDC/ADCC/ADCP-related activities of the Fc region, among other properties described herein, relative to a corresponding wild-type Fc sequence of an antibody or antigen-binding fragment thereof. Included are modified Fc regions of human and/or mouse origin.

Also included are antibodies or antigen-binding fragments thereof that comprise hybrid Fc regions, for example, Fc regions that comprise a combination of Fc domains (e.g., hinge, CH₂, CH₃, CH₄) from immunoglobulins of different species (e.g., human, mouse), different Ig classes, and/or different Ig subclasses. General examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of CH₂/CH₃ domains: IgA1/IgA1, IgA1/IgA2, IgA1/IgD, IgA1/IgE, IgA1/IgG1, IgA1/IgG2, IgA1/IgG3, IgA1/IgG4, IgA1/IgM, IgA2/IgA1, IgA2/IgA2, IgA2/IgD, IgA2/IgE, IgA2/IgG1, IgA2/IgG2, IgA2/IgG3, IgA2/IgG4, IgA2/IgM, IgD/IgA1, IgD/IgA2, IgD/IgD, IgD/IgE, IgD/IgG1, IgD/IgG2, IgD/IgG3, IgD/IgG4, IgD/IgM, IgE/IgA1, IgE/IgA2, IgE/IgD, IgE/IgE, IgE/IgG1, IgE/IgG2, IgE/IgG3, IgE/IgG4, IgE/IgM, IgG1/IgA1, IgG1/IgA2, IgG1/IgD, IgG1/IgE, IgG1/IgG1, IgG1/IgG2, IgG1/IgG3, IgG1/IgG4, IgG1/IgM, IgG2/IgA1, IgG2/IgA2, IgG2/IgD, IgG2/IgE, IgG2/IgG1, IgG2/IgG2, IgG2/IgG3, IgG2/IgG4, IgG2/IgM, IgG3/IgA1, IgG3/IgA2, IgG3/IgD, IgG3/IgE, IgG3/IgG1, IgG3/IgG2, IgG3/IgG3, IgG3/IgG4, IgG3/IgM, IgG4/IgA1, IgG4/IgA2, IgG4/IgD, IgG4/IgE, IgG4/IgG1, IgG4/IgG2, IgG4/IgG3, IgG4/IgG4, IgG4/IgM, IgM/IgA1, IgM/IgA2, IgM/IgD, IgM/IgE, IgM/IgG1, IgM/IgG2, IgM/IgG3, IgM/IgG4, IgM/IgM (or fragments or variants thereof), and optionally include a hinge from one or more of IgA1, IgA2, IgD, IgG1, IgG2, IgG3, or IgG4, and/or a CH₄ domain from IgE and/or IgM. In specific embodiments, the hinge, CH₂, CH₃, and CH₄ domains are from human Ig.

Additional examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of CH₂/CH₄ domains: IgA1/IgE, IgA2/IgE, IgD/IgE, IgE/IgE, IgG1/IgE, IgG2/IgE, IgG3/IgE, IgG4/IgE, IgM/IgE, IgA1/IgM, IgA2/IgM, IgD/IgM, IgE/IgM, IgG1/IgM, IgG2/IgM, IgG3/IgM, IgG4/IgM, IgM/IgM (or

fragments or variants thereof), and optionally include a hinge from one or more of IgA1, IgA2, IgD, IgG1, IgG2, IgG3, IgG4, and/or a CH₃ domain from one or more of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4, or IgM. In specific embodiments, the hinge, CH₂, CH₃, and CH₄ domains are from human Ig.

Certain examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of CH₃/CH₄ domains: IgA1/IgE, IgA2/IgE, IgD/IgE, IgE/IgE, IgG1/IgE, IgG2/IgE, IgG3/IgE, IgG4/IgE, IgM/IgE, IgA1/IgM, IgA2/IgM, IgD/IgM, IgE/IgM, IgG1/IgM, IgG2/IgM, IgG3/IgM, IgG4/IgM, IgM/IgM (or fragments or variants thereof), and optionally include a hinge from one or more of IgA1, IgA2, IgD, IgG1, IgG2, IgG3, IgG4, and/or a CH₂ domain from one or more of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4, or IgM. In specific embodiments, the hinge, CH₂, CH₃, and CH₄ domains are from human Ig.

Particular examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of hinge/CH₂ domains: IgA1/IgA1, IgA1/IgA2, IgA1/IgD, IgA1/IgE, IgA1/IgG1, IgA1/IgG2, IgA1/IgG3, IgA1/IgG4, IgA1/IgM, IgA2/IgA1, IgA2/IgA2, IgA2/IgD, IgA2/IgE, IgA2/IgG1, IgA2/IgG2, IgA2/IgG3, IgA2/IgG4, IgA2/IgM, IgD/IgA1, IgD/IgA2, IgD/IgD, IgD/IgE, IgD/IgG1, IgD/IgG2, IgD/IgG3, IgD/IgG4, IgD/IgM, IgG1/IgA1, IgG1/IgA2, IgG1/IgD, IgG1/IgE, IgG1/IgG1, IgG1/IgG2, IgG1/IgG3, IgG1/IgG4, IgG1/IgM, IgG2/IgA1, IgG2/IgA2, IgG2/IgD, IgG2/IgE, IgG2/IgG1, IgG2/IgG2, IgG2/IgG3, IgG2/IgG4, IgG2/IgM, IgG3/IgA1, IgG3/IgA2, IgG3/IgD, IgG3/IgE, IgG3/IgG1, IgG3/IgG2, IgG3/IgG3, IgG3/IgG4, IgG3/IgM, IgG4/IgA1, IgG4/IgA2, IgG4/IgD, IgG4/IgE, IgG4/IgG1, IgG4/IgG2, IgG4/IgG3, IgG4/IgG4, IgG4/IgM (or fragments or variants thereof), and optionally include a CH₃ domain from one or more of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4, or IgM, and/or a CH₄ domain from IgE and/or IgM. In specific embodiments, the hinge, CH₂, CH₃, and CH₄ domains are from human Ig.

Certain examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of hinge/CH₃ domains: IgA1/IgA1, IgA1/IgA2, IgA1/IgD, IgA1/IgE, IgA1/IgG1, IgA1/IgG2, IgA1/IgG3, IgA1/IgG4, IgA1/IgM, IgA2/IgA1, IgA2/IgA2, IgA2/IgD, IgA2/IgE, IgA2/IgG1, IgA2/IgG2, IgA2/IgG3, IgA2/IgG4, IgA2/IgM, IgD/IgA1, IgD/IgA2, IgD/IgD, IgD/IgE, IgD/IgG1, IgD/IgG2, IgD/IgG3, IgD/IgG4, IgD/IgM, IgG1/IgA1, IgG1/IgA2, IgG1/IgD, IgG1/IgE, IgG1/IgG1, IgG1/IgG2, IgG1/IgG3, IgG1/IgG4, IgG1/IgM, IgG2/IgA1, IgG2/IgA2, IgG2/IgD, IgG2/IgE, IgG2/IgG1, IgG2/IgG2, IgG2/IgG3, IgG2/IgG4, IgG2/IgM, IgG3/IgA1, IgG3/IgA2, IgG3/IgD, IgG3/IgE, IgG3/IgG1, IgG3/IgG2, IgG3/IgG3, IgG3/IgG4, IgG3/IgM, IgG4/IgA1, IgG4/IgA2, IgG4/IgD, IgG4/IgE, IgG4/IgG1, IgG4/IgG2, IgG4/IgG3, IgG4/IgG4, IgG4/IgM (or fragments or variants thereof), and optionally include a CH₂ domain from one or more of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4, or IgM, and/or a CH₄ domain from IgE and/or IgM. In specific embodiments, the hinge, CH₂, CH₃, and CH₄ domains are from human Ig.

Some examples include hybrid Fc regions that comprise, consist of, or consist essentially of the following combination of hinge/CH₄ domains: IgA1/IgE, IgA1/IgM, IgA2/IgE, IgA2/IgM, IgD/IgE, IgD/IgM, IgG1/IgE, IgG1/IgM, IgG2/IgE, IgG2/IgM, IgG3/IgE, IgG3/IgM, IgG4/IgE, IgG4/IgM (or fragments or variants thereof), and optionally include a CH₂ domain from one or more of IgA1, IgA2, IgD, IgE,

IgG1, IgG2, IgG3, IgG4, or IgM, and/or a CH₃ domain from one or more of IgA1, IgA2, IgD, IgE, IgG1, IgG2, IgG3, IgG4, or IgM.

Specific examples of hybrid Fc regions can be found, for example, in WO 2008/147143, which are derived from combinations of IgG subclasses or combinations of human IgD and IgG.

Also included are antibodies or antigen-binding fragments thereof having derivatized or otherwise modified Fc regions. In certain aspects, the Fc region may be modified by phosphorylation, sulfation, acylation, glycosylation, methylation, farnesylation, acetylation, amidation, and the like, for instance, relative to a wild-type or naturally-occurring Fc region. In certain embodiments, the Fc region may comprise wild-type or native glycosylation patterns, or alternatively, it may comprise increased glycosylation relative to a native form, decreased glycosylation relative to a native form, or it may be entirely deglycosylated. As one example of a modified Fc glycoform, decreased glycosylation of an Fc region reduces binding to the C1q region of the first complement component C1, a decrease in ADCC-related activity, and/or a decrease in CDC-related activity. Certain embodiments thus employ a deglycosylated or aglycosylated Fc region. See, e.g., WO 2005/047337 for the production of exemplary aglycosylated Fc regions. Another example of an Fc region glycoform can be generated by substituting the Q295 position with a cysteine residue (see, e.g., U.S. Application No. 2010/0080794), according to the Kabat et al. numbering system. Certain embodiments may include Fc regions where about 80-100% of the glycoprotein in Fc region comprises a mature core carbohydrate structure that lacks fructose (see, e.g., U.S. Application No. 2010/0255013). Some embodiments may include Fc regions that are optimized by substitution or deletion to reduce the level of fucosylation, for instance, to increase affinity for FcγRI, FcγRIa, or FcγRIIIa, and/or to improve phagocytosis by FcγRIIIa-expressing cells (see U.S. Application Nos. 2010/0249382 and 2007/0148170).

As another example of a modified Fc glycoform, an Fc region of an antibody or antigen-binding fragment thereof may comprise oligomannose-type N-glycans, and optionally have one or more of the following: increased ADCC effector activity, increased binding affinity for FcγRIIIA (and certain other FcRs), similar or increased binding specificity for the target of the NRP2 polypeptide, similar or higher binding affinity for the target of the NRP2 polypeptide, and/or similar or lower binding affinity for mannose receptor, relative to a corresponding Fc region that contains complex-type N-glycans (see, e.g., U.S. Application No. 2007/0092521 and U.S. Pat. No. 7,700,321). As another example, enhanced affinity of Fc regions for FcγRs has been achieved using engineered glycoforms generated by expression of antibodies in engineered or variant cell lines (see, e.g., Umana et al., *Nat Biotechnol.* 17:176-180, 1999; Davies et al., *Biotechnol Bioeng.* 74:288-294, 2001; Shields et al., *J Biol Chem.* 277:26733-26740, 2002; Shinkawa et al., 2003, *J Biol Chem.* 278:3466-3473, 2003; and U.S. Application No. 2007/0111281). Certain Fc region glycoforms comprise an increased proportion of N-glycoside bond type complex sugar chains, which do not have the 1-position of fucose bound to the 6-position of N-acetylglucosamine at the reducing end of the sugar chain (see, e.g., U.S. Application No. 2010/0092997). Particular embodiments may include IgG Fc region that is glycosylated with at least one galactose moiety connected to a respective terminal sialic acid moiety by an α-2,6 linkage, optionally where the Fc region has a higher anti-inflammatory activity relative to a correspond-

ing, wild-type Fc region (see U.S. Application No. 2008/0206246). Certain of these and related altered glycosylation approaches have generated substantial enhancements of the capacity of Fc regions to selectively bind FcRs such as FcγRIII, to mediate ADCC, and to alter other properties of Fc regions, as described herein.

Certain variant, fragment, hybrid, or otherwise modified Fc regions of an antibody or antigen-binding fragment thereof may have altered binding to one or more FcRs, and/or corresponding changes to effector function, relative to a corresponding, wild-type Fc sequence (e.g., same species, same Ig class, same Ig subclass). For instance, such Fc regions may have increased binding to one or more of Fcγ receptors, Fcα receptors, Fcε receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In other embodiments, variant, fragment, hybrid, or modified Fc regions may have decreased binding to one or more of Fcγ receptors, Fcα receptors, Fcε receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. Specific FcRs are described elsewhere herein.

In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to increase binding to one or more of Fcγ receptors, Fcα receptors, Fcα receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an IgG1 or IgG3 Fc domain, comprising one or more mutations to increase binding to one or more of Fcγ receptors, Fcα receptors, Fcα receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to increase effector function. In some embodiments the at least one antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function.

In some embodiments, an antibody is blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the blocking antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function. In some embodiments, an antibody is a partial-blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the a partial-blocking antibody comprises an Fc domain selected from a human IgG1 and IgG3, comprising one or more mutations to increase effector function. In some embodiments, an antibody is a non-blocking antibody that comprises an Fc domain with high effector activity. In some embodiments, the non-blocking antibody comprises an Fc domain selected from a human IgG1 or IgG3, comprising one or more mutations to increase effector function.

In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to decrease binding to one or more of Fcγ receptors, Fcα receptors, Fcα receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an IgG1 or IgG3 Fc domain, comprising one or more mutations to decrease binding to one or more of Fcγ receptors, Fcα receptors, Fcε receptors, and/or the neonatal Fc receptor, relative to a corresponding, wild-type Fc sequence. In some embodiments, an antibody comprises an Fc domain, comprising one or more mutations to decrease effector function. In some embodiments, an antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function.

In some embodiments, an antibody is a blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function. In some embodiments, an antibody is a partial-blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the partial-blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function. In some embodiments, an antibody is a non-blocking antibody comprising an Fc domain with low effector activity. In some embodiments, the non-blocking antibody comprises an Fc domain selected from a human IgG2 and IgG4, comprising one or more mutations to decrease effector function.

Specific examples of Fc variants having altered (e.g., increased, decreased) effector function/FcR binding can be found, for example, in U.S. Pat. Nos. 5,624,821 and 7,425,619; U.S. Application Nos. 2009/0017023, 2009/0010921, and 2010/0203046; and WO 2000/42072 and WO 2004/016750. Certain examples include human Fc regions having one or more substitutions at position 298, 333, and/or 334, for example, S298A, E333A, and/or K334A (based on the numbering of the EU index of Kabat et al.), which have been shown to increase binding to the activating receptor FcγRIIIa and reduce binding to the inhibitory receptor FcγRIIb. These mutations can be combined to obtain double and triple mutation variants that have further improvements in binding to FcRs. Certain embodiments include a S298A/E333A/K334A triple mutant, which has increased binding to FcγRIIIa, decreased binding to FcγRIIb, and increased ADCC (see, e.g., Shields et al., *J Biol Chem.* 276:6591-6604, 2001; and Presta et al., *Biochem Soc Trans.* 30:487-490, 2002). See also engineered Fc glycoforms that have increased binding to FcRs, as disclosed in Umana et al., *supra*; and U.S. Pat. No. 7,662,925. Some embodiments include Fc regions that comprise one or more substitutions selected from 434S, 252Y/428L, 252Y/434S, and 428L/434S (see U.S. Application Nos. 2009/0163699 and 20060173170), based on the EU index of Kabat et al.

Certain variant, fragment, hybrid, or modified Fc regions may have altered effector functions, relative to a corresponding, wild-type Fc sequence. For example, such Fc regions may have increased complement fixation or activation, increased C1q binding affinity, increased CDC-related activity, increased ADCC-related activity, and/or increased ADCC-related activity, relative to a corresponding, wild-type Fc sequence. In other embodiments, such Fc regions may have decreased complement fixation or activation, decreased C1q binding affinity, decreased CDC-related activity, decreased ADCC-related activity, and/or decreased ADCC-related activity, relative to a corresponding, wild-type Fc sequence. As merely one illustrative example, an Fc region may comprise a deletion or substitution in a complement-binding site, such as a C1q-binding site, and/or a deletion or substitution in an ADCC site. Examples of such deletions/substitutions are described, for example, in U.S. Pat. No. 7,030,226. Many Fc effector functions, such as ADCC, can be assayed according to routine techniques in the art. (see, e.g., Zuckerman et al., *CRC Crit Rev Microbiol.* 7:1-26, 1978). Useful effector cells for such assays includes, but are not limited to, natural killer (NK) cells, macrophages, and other peripheral blood mononuclear cells (PBMC). Alternatively, or additionally, certain Fc effector functions may be assessed in vivo, for example, by employing an animal model described in Clynes et al. *PNAS.* 95:652-656, 1998.

Certain variant hybrid, or modified Fc regions may have altered stability or half-life relative to a corresponding, wild-type Fc sequence. In certain embodiments, such Fc regions may have increased half-life relative to a corresponding, wild-type Fc sequence. In other embodiments, variant hybrid, or modified Fc regions may have decreased half-life relative to a corresponding, wild-type Fc sequence. Half-life can be measured in vitro (e.g., under physiological conditions) or in vivo, according to routine techniques in the art, such as radiolabeling, ELISA, or other methods. In vivo measurements of stability or half-life can be measured in one or more bodily fluids, including blood, serum, plasma, urine, or cerebrospinal fluid, or a given tissue, such as the liver, kidneys, muscle, central nervous system tissues, bone, etc.

As one example, modifications to an Fc region that alter its ability to bind the FcRn can alter its half-life in vivo, or other properties. In some embodiments, a modified Fc domain, for example, a modified IgG1 or IgG4 Fc domain, comprises at least one mutation to alter FcRn binding as described, for example, by Zalevsky et al. (*Nature Biotechnology.* 28 (2): 157-159, 2010) or Mackness et al. (*mAbs.* 11 (7): 1276-1288, 2019). In specific embodiments, a modified IgG1 or IgG3 Fc domain comprises any one or more of YD (M252Y/T256D), DQ (T256D/T307Q), DW (T256D/T307W), YTE (M252Y/S254T/T256E), AAA (T307A/E380A/N434A), LS (M428L/N434S), M252Y, T256D/E, K288D/N, T307Q/W, E380C, N434FY, or Y436H/N/W mutations (EU numbering), including combinations thereof. In some embodiments, a modified IgG1 or IgG3 Fc domain comprises any one or more of M252Y, T256D/E, T307Q/W, and/or N434F/Y mutations (EU numbering), including combinations thereof. Assays for measuring the in vivo pharmacokinetic properties (e.g., in vivo mean elimination half-life) and non-limiting examples of Fc modifications that alter its binding to the FcRn are described, for example, in U.S. Pat. Nos. 7,217,797 and 7,732,570; and U.S. Application Nos. US 2010/0143254 and 2010/0143254.

Additional non-limiting examples of modifications to alter stability or half-life include substitutions/deletions at one or more of amino acid residues selected from 251-256, 285-290, and 308-314 in the CH₂ domain, and 385-389 and 428-436 in the CH₃ domain, according to the numbering system of Kabat et al. See U.S. Application No. 2003/0190311. Specific examples include substitution with leucine at position 251, substitution with tyrosine, tryptophan or phenylalanine at position 252, substitution with threonine or serine at position 254, substitution with arginine at position 255, substitution with glutamine, arginine, serine, threonine, or glutamate at position 256, substitution with threonine at position 308, substitution with proline at position 309, substitution with serine at position 311, substitution with aspartate at position 312, substitution with leucine at position 314, substitution with arginine, aspartate or serine at position 385, substitution with threonine or proline at position 386, substitution with arginine or proline at position 387, substitution with proline, asparagine or serine at position 389, substitution with methionine or threonine at position 428, substitution with tyrosine or phenylalanine at position 434, substitution with histidine, arginine, lysine or serine at position 433, and/or substitution with histidine, tyrosine, arginine or threonine at position 436, including any combination thereof. Such modifications optionally increase affinity of the Fc region for the FcRn and thereby increase half-life, relative to a corresponding, wild-type Fc region.

Certain variant hybrid, or modified Fc regions may have altered solubility relative to a corresponding, wild-type Fc

sequence. In certain embodiments, such Fc regions may have increased solubility relative to a corresponding, wild-type Fc sequence. In other embodiments, variant hybrid, or modified Fc regions may have decreased solubility relative to a corresponding, wild-type Fc sequence. Solubility can be measured, for example, in vitro (e.g., under physiological conditions) according to routine techniques in the art. Exemplary solubility measurements are described elsewhere herein.

Additional examples of variants include IgG Fc regions having conservative or non-conservative substitutions (as described elsewhere herein) at one or more of positions 250, 314, or 428 of the heavy chain, or in any combination thereof, such as at positions 250 and 428, or at positions 250 and 314, or at positions 314 and 428, or at positions 250, 314, and 428 (see, e.g., U.S. Application No. 2011/0183412). In specific embodiments, the residue at position 250 is substituted with glutamic acid or glutamine, and/or the residue at position 428 is substituted with leucine or phenylalanine. As another illustrative example of an IgG Fc variant, any one or more of the amino acid residues at positions 214 to 238, 297 to 299, 318 to 322, and/or 327 to 331 may be used as a suitable target for modification (e.g., conservative or non-conservative substitution, deletion). In particular embodiments, the IgG Fc variant CH₂ domain contains amino acid substitutions at positions 228, 234, 235, and/or 331 (e.g., human IgG4 with Ser228Pro and Leu235Ala mutations) to attenuate the effector functions of the Fc region (see U.S. Pat. No. 7,030,226). Here, the numbering of the residues in the heavy chain is that of the EU index (see Kabat et al., "Sequences of Proteins of Immunological Interest," 5th Ed., National Institutes of Health, Bethesda, Md. (1991)). Certain of these and related embodiments have altered (e.g., increased, decreased) FcRn binding and/or serum half-life, optionally without reduced effector functions such as ADCC or CDC-related activities.

Additional examples include variant Fc regions that comprise one or more amino acid substitutions at positions 279, 341, 343 or 373 of a wild-type Fc region, or any combination thereof (see, e.g., U.S. Application No. 2007/0224188). The wild-type amino acid residues at these positions for human IgG are valine (279), glycine (341), proline (343) and tyrosine (373). The substitution(s) can be conservative or non-conservative, or can include non-naturally occurring amino acids or mimetics, as described herein. Alone or in combination with these substitutions, certain embodiments may also employ a variant Fc region that comprises at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more amino acid substitutions selected from the following: 235G, 235R, 236F, 236R, 236Y, 237K, 237N, 237R, 238E, 238G, 238H, 238I, 238L, 238V, 238W, 238Y, 244L, 245R, 247A, 247D, 247E, 247F, 247M, 247N, 247Q, 247R, 247S, 247T, 247W, 247Y, 248F, 248P, 248Q, 248W, 249L, 249M, 249N, 249P, 249Y, 251H, 251I, 251W, 254D, 254E, 254F, 254G, 254H, 254I, 254K, 254L, 254M, 254N, 254P, 254Q, 254R, 254V, 254W, 254Y, 255K, 255N, 256H, 256I, 256K, 256L, 256V, 256W, 256Y, 257A, 257I, 257M, 257N, 257S, 258D, 260S, 262L, 264S, 265K, 265S, 267H, 267I, 267K, 268K, 269N, 269Q, 271T, 272H, 272K, 272L, 272R, 279A, 279D, 279F, 279G, 279H, 279I, 279K, 279L, 279M, 279N, 279Q, 279R, 279S, 279T, 279W, 279Y, 280T, 283F, 283G, 283H, 283I, 283K, 283L, 283M, 283P, 283R, 283T, 283W, 283Y, 285N, 286F, 288N, 288P, 292E, 292F, 292G, 292I, 292L, 293S, 293V, 301W, 304E, 307E, 307M, 312P, 315F, 315K, 315L, 315P, 315R, 316F, 316K, 317P, 317T, 318N, 318P, 318T, 332F, 332G, 332L, 332M, 332S, 332V, 332W, 339D, 339E, 339F, 339G, 339H, 339I, 339K, 339L, 339M, 339N, 339Q, 339R, 339S,

339W, 339Y, 341D, 341E, 341F, 341H, 341I, 341K, 341L, 341M, 341N, 341P, 341Q, 341R, 341S, 341T, 341V, 341W, 341Y, 343A, 343D, 343E, 343F, 343G, 343H, 343I, 343K, 343L, 343M, 343N, 343Q, 343R, 343S, 343T, 343V, 343W, 343Y, 373D, 373E, 373F, 373G, 373H, 373I, 373K, 373L, 373M, 373N, 373Q, 373R, 373S, 373T, 373V, 373W, 375R, 376E, 376F, 376G, 376H, 376I, 376L, 376M, 376N, 376P, 376Q, 376R, 376S, 376T, 376V, 376W, 376Y, 377G, 377K, 377P, 378N, 379N, 379Q, 379S, 379T, 380D, 380N, 380S, 380T, 382D, 382F, 382H, 382I, 382K, 382L, 382M, 382N, 382P, 382Q, 382R, 382S, 382T, 382V, 382W, 382Y, 385E, 385P, 386K, 423N, 424H, 424M, 424V, 426D, 426L, 427N, 429A, 429F, 429M, 430A, 430D, 430F, 430G, 430H, 430I, 430K, 430L, 430M, 430N, 430P, 430Q, 430R, 430S, 430T, 430V, 430W, 430Y, 431H, 431K, 431P, 432R, 432S, 438G, 438K, 438L, 438T, 438W, 439E, 439H, 439Q, 440D, 440E, 440F, 440G, 440H, 440I, 440K, 440L, 440M, 440Q, 440T, 440V or 442K. As above, the numbering of the residues in the heavy chain is that of the EU index (see Kabat et al., supra). Such variant Fc regions typically confer an altered effector function or altered serum half-life upon the antibody to which the variant Fc region is operably attached. Preferably the altered effector function is an increase in ADCC, a decrease in ADCC, an increase in CDC, a decrease in CDC, an increase in C1q binding affinity, a decrease in C1q binding affinity, an increase in FcR (preferably FcRn) binding affinity or a decrease in FcR (preferably FcRn) binding affinity as compared to a corresponding Fc region that lacks such amino acid substitution(s).

Additional examples include variant Fc regions that comprise an amino acid substitution at one or more of position(s) 221, 222, 224, 227, 228, 230, 231, 223, 233, 234, 235, 236, 237, 238, 239, 240, 241, 243, 244, 245, 246, 247, 249, 250, 258, 262, 263, 264, 265, 266, 267, 268, 269, 270, 271, 272, 273, 274, 275, 276, 278, 280, 281, 283, 285, 286, 288, 290, 291, 293, 294, 295, 296, 297, 298, 299, 300, 302, 313, 317, 318, 320, 322, 323, 324, 325, 326, 327, 328, 329, 330, 331, 332, 333, 334, 335 336 and/or 428 (see, e.g., U.S. Pat. No. 7,662,925). In specific embodiments, the variant Fc region comprises at least one amino acid substitution selected from the group consisting of: P230A, E233D, L234E, L234Y, L234I, L235D, L235S, L235Y, L235I, S239D, S239E, S239N, S239Q, S239T, V240I, V240M, F243L, V264I, V264T, V264Y, V266I, E272Y, K274T, K274E, K274R, K274L, K274Y, F275W, N276L, Y278T, V302I, E318R, S324D, S324I, S324V, N325T, K326I, K326T, L328M, L328I, L328Q, L328D, L328V, L328T, A330Y, A330L, A330I, I332D, I332E, I332N, I332Q, T335D, T335R, and T335Y. In other specific embodiments, the variant Fc region comprises at least one amino acid substitution selected from the group consisting of: V264I, F243L/V264I, L328M, I332E, L328M/I332E, V264I/I332E, S298A/I332E, S239E/I332E, S239Q/I332E, S239E, A330Y, I332D, L328I/I332E, L328Q/I332E, V264T, V240I, V266I, S239D, S239D/I332D, S239D/I332E, S239D/I332N, S239D/I332Q, S239E/I332D, S239E/I332N, S239E/I332Q, S239N/I332D, S239N/I332E, S239Q/I332D, A330Y/I332E, V264I/A330Y/I332E, A330L/I332E, V264I/A330L/I332E, L234E, L234Y, L234I, L235D, L235S, L235Y, L235I, S239T, V240M, V264Y, A330I, N325T, L328D/I332E, L328V/I332E, L328T/I332E, L328I/I332E, S239E/V264I/I332E, S239Q/V264I/I332E, S239E/V264I/A330Y/I332E, S239D/A330Y/I332E, S239N/A330Y/I332E, S239D/A330L/I332E, S239N/A330L/I332E, V264I/S298A/I332E, S239D/S298A/I332E, S239N/S298A/I332E, S239D/V264I/I332E, S239D/V264I/S298A/I332E, S239D/V264I/A330L/I332E, S239D/I332E/A330I, P230A, P230A/E233D/I332E,

E272Y, K274T, K274E, K274R, K274L, K274Y, F275W, N276L, Y278T, V302I, E318R, S324D, S324I, S324V, K326I, K326T, T335D, T335R, T335Y, V240I/V266I, S239D/A330Y/I332E/L234I, S239D/A330Y/I332E/L235D, S239D/A330Y/I332E/V240I, S239D/A330Y/I332E/V264T, S239D/A330Y/I332E/K326E, and S239D/A330Y/I332E/K326T. In more specific embodiments, the variant Fc region comprises a series of substitutions selected from the group consisting of: N297D/I332E, F241Y/F243Y/V262T/V264T/N297D/I332E, S239D/N297D/I332E, S239E/N297D/I332E, S239D/D265Y/N297D/I332E, S239D/D265H/N297D/I332E, V264E/N297D/I332E, Y296N/N297D/I332E, N297D/A330Y/I332E, S239D/D265V/N297D/I332E, S239D/D265I/N297D/I332E, and N297D/S298A/A330Y/I332E. In specific embodiments, the variant Fc region comprises an amino acid substitution at position 332 (using the numbering of the EU index, Kabat et al., supra). Examples of substitutions include 332A, 332D, 332E, 332F, 332G, 332H, 332K, 332L, 332M, 332N, 332P, 332Q, 332R, 332S, 332T, 332V, 332W and 332Y. The numbering of the residues in the Fc region is that of the EU index of Kabat et al. Among other properties described herein, such variant Fc regions may have increased affinity for an FcγR, increased stability, and/or increased solubility, relative to a corresponding, wild-type Fc region. Further examples include variant Fc regions that comprise one or more of the following amino acid substitutions: 224N/Y, 225A, 228L, 230S, 239P, 240A, 241L, 243S/L/G/H/I, 244L, 246E, 247L/A, 252T, 254T/P, 258K, 261Y, 265V, 266A, 267G/N, 268N, 269K/G, 273A, 276D, 278H, 279M, 280N, 283G, 285R, 288R, 289A, 290E, 291L, 292Q, 297D, 299A, 300H, 301C, 304G, 305A, 306I/F, 311R, 312N, 315D/K/S, 320R, 322E, 323A, 324T, 325S, 326E/R, 332T, 333D/G, 335I, 338R, 339T, 340Q, 341E, 342R, 344Q, 347R, 351S, 352A, 354A, 355W, 356G, 358T, 361D/Y, 362L, 364C, 365Q/P, 370R, 372L, 377V, 378T, 383N, 389S, 390D, 391C, 393A, 394A, 399G, 404S, 408G, 409R, 411I, 412A, 414M, 421S, 422I, 426F/P, 428T, 430K, 431S, 432P, 433P, 438L, 439E/R, 440G, 441F, 442T, 445R, 446A, 447E, optionally where the variant has altered recognition of an Fc ligand and/or altered effector function compared with a parent Fc polypeptide, and wherein the numbering of the residues is that of the EU index as in Kabat et al. Specific examples of these and related embodiments include variant Fc regions that comprise or consist of the following sets of substitutions: (1) N276D, R292Q, V305A, I377V, T394A, V412A

and K439E; (2) P244L, K246E, D399G and K409R; (3) S304G, K320R, S324T, K326E and M358T; (4) F243S, P247L, D265V, V266A, S383N and T411I; (5) H224N, F243L, T393A and H433P; (6) V240A, S267G, G341E and E356G; (7) M252T, P291L, P352A, R355W, N390D, S408G, S426F and A431S; (8) P228L, T289A, L365Q, N389S and S440G; (9) F241L, V273A, K340Q and L441F; (10) F241L, T299A, I332T and M428T; (11) E269K, Y300H, Q342R, V422I and G446A; (12) T225A, R301C, S304G, D312N, N315D, L351S and N421S; (13) S254T, L306I, K326R and Q362L; (14) H224Y, P230S, V323A, E333D, K338R and S364C; (15) T335I, K414M and P445R; (16) T335I and K414M; (17) P247A, E258K, D280N, K288R, N297D, T299A, K322E, Q342R, S354A and L365P; (18) H268N, V279M, A339T, N361D and S426P; (19) C261Y, K290E, L306F, Q311R, E333G and Q438L; (20) E283G, N315K, E333G, R344Q, L365P and S442T; (21) Q347R, N361Y and K439R; (22) S239P, S254P, S267N, H285R, N315S, F372L, A378T, N390D, Y391C, F404S, E430K, L432P and K447E; and (23) E269G, Y278H, N325S and K370R, wherein the numbering of the residues is that of the EU index as in Kabat et al. (see, e.g., U.S. Application No. 2010/0184959).

Variant Fc regions can also have one or more mutated hinge regions, as described, for example, in U.S. Application No. 2003/0118592. For instance, one or more cysteines in a hinge region can be deleted or substituted with a different amino acid. The mutated hinge region can comprise no cysteine residues, or it can comprise 1, 2, or 3 fewer cysteine residues than a corresponding, wild-type hinge region. In some embodiments, an Fc region having a mutated hinge region of this type exhibits a reduced ability to dimerize, relative to a wild-type Ig hinge region.

In particular embodiments, the Fc region comprises, consists, or consists essentially of the Fc from human IgG1 or IgG4 (see, e.g., Allberse and Schuurman, *Immunology*, 105:9-19, 2002), or a fragment or variant thereof. Table F1 below provides exemplary sequences (CH1, hinge (underlined), CH2, and CH3 regions) from human IgG1 and IgG4. Examples of variant IgG4 sequences that can be employed are described, for example, in Peters et al., *JBC*, 287:24525-24533, 2012, and include substitutions at C227, C230, C127 (e.g., C127S), and C131 (e.g., C131S). Other variants that can be used include a L445P substitution in IgG4 (denoted as IgG4-2) or a D356E and L358M substitution in IgG1, (denoted as IgG1m(zf)).

TABLE F1

Exemplary IgG4 Fc Sequences		
Name	Sequence	SEQ ID NO:
Wild-type IgG4	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQKTYTCNVDPKPSNTKVDKRVES KYGGPPCPSCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSDQED PEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVHLQDNLNGKEYK CKVSNKGLPSSIEKTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVK GFYPDSIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEG NVFSCSVMHREALHNYTQKSLSLSLGK	144
IgG4-M265Y, S267T, T269E (YTE, Kabat numbering)	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQKTYTCNVDPKPSNTKVDKRVES KYGGPPCPSCPAPEFLGGPSVFLFPPKPKDTLYITREPEVTCVVVDVSDQED PEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVHLQDNLNGKEYK CKVSNKGLPSSIEKTIISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVK GFYPDSIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEG NVFSCSVMHREALHNYTQKSLSLSLGK	145

TABLE F1-continued

Exemplary IgG4 Fc Sequences		
Name	Sequence	SEQ ID NO:
IgG4-S241P (Kabat numbering)	ASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALTSGV HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYTCNVDPKPSNTKVDKRVES KYGPCCPCCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVSQED PEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVHLQDNLNGKEYK CKVSNKGLPSSIEKTIKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVK GFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRWQEG NVFSCSVMHEALHNHYTQKSLSLGLK	146
IgG1m(za) GenBank: AH007035.2	ASTKGPSVFPLAPSSKSTSGGTAAALGCLVKDYFPEPVTVSWNSGALTSGV HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEP KSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVDVS HEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVHLQDNLNGK EYCKKVSNAKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCL LVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSRW QQGNVFCSCVMHEALHNHYTQKSLSLSPGK	147
Kappa Km3	RTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSG NSQESVTEQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC	148

As noted above, antibodies having altered Fc regions typically have altered (e.g., improved, increased, decreased) pharmacokinetic properties relative to corresponding wild-type Fc region. Examples of pharmacokinetic properties include stability or half-life, bioavailability (the fraction of a drug that is absorbed), tissue distribution, volume of distribution (apparent volume in which a drug is distributed immediately after it has been injected intravenously and equilibrated between plasma and the surrounding tissues), concentration (initial or steady-state concentration of drug in plasma), elimination rate constant (rate at which drugs are removed from the body), elimination rate (rate of infusion required to balance elimination), area under the curve (AUC or exposure; integral of the concentration-time curve, after a single dose or in steady state), clearance (volume of plasma cleared of the drug per unit time), C_{max} (peak plasma concentration of a drug after oral administration), t_{max} (time to reach C_{max}), C_{min} (lowest concentration that a drug reaches before the next dose is administered), and fluctuation (peak trough fluctuation within one dosing interval at steady state).

In particular embodiments, an antibody or antigen-binding fragment thereof has a biological half life at about pH 7.4, at about a physiological pH, at about 25° C. or room temperature, and/or at about 37° C. or human body temperature (e.g., in vivo, in serum, in a given tissue, in a given species such as rat, mouse, monkey, or human), of about or at least about 30 minutes, about 1 hour, about 2 hour, about 3 hours, about 4 hours, about 5 hours, about 6 hours, about 12 hours, about 18 hours, about 20 hours, about 24 hours, about 30 hours, about 36 hours, about 40 hours, about 48 hours, about 50 hours, about 60 hours, about 70 hours, about 72 hours, about 80 hours, about 84 hours, about 90 hours, about 96 hours, about 120 hours, or about 144 hours or more, or about 1 week, or about 2 weeks, or about 3 weeks, or about 4 weeks, or about 5 weeks, or about 6 weeks or more, or any intervening half-life, including all ranges in between.

In some embodiments, an antibody or antigen-binding fragment thereof has a T_m of about or at least about 60, 62, 64, 66, 68, 70, 72, 74, or 75° C. In some embodiments, an antibody or antigen-binding fragment thereof has a T_m of about 60° C. or greater.

In some embodiments, an antibody or antigen-binding fragment thereof conjugated to one or more cytotoxic or chemotherapeutic agents. General examples of cytotoxic or chemotherapeutic agents include, without limitation, alkylating agents, anti-metabolites, anthracyclines, anti-tumor antibiotics, platinum, type I topoisomerase inhibitors, type II topoisomerase inhibitors, vinca alkaloids, and taxanes. Specific examples of cytotoxic or chemotherapeutic agents include, without limitation, cyclophosphamide, cilengitide, lomustine (CCNU), melphalan, procarbazine, carmustine (BCNU), enzastaurin, busulfan, daunorubicin, doxorubicin, gefitinib, erlotinib, idarubicin, temozolomide, epirubicin, mitoxantrone, bleomycin, cisplatin, carboplatin, oxaliplatin, camptothecins, irinotecan, topotecan, amsacrine, etoposide, etoposide phosphate, teniposide, temsirolimus, everolimus, vincristine, vinblastine, vinorelbine, vindesine, CT52923, paclitaxel, imatinib, dasatinib, sorafenib, pazopanib, sunitinib, vatalanib, gefitinib, erlotinib, AEE-788, dichloroacetate, tamoxifen, fasudil, SB-681323, semaxanib, donepezil, galantamine, memantine, rivastigmine, tacrine, rasagiline, naltrexone, lubiprostone, safinamide, istradefylline, pimavanterin, pitolisant, isradipine, pridopidine (ACR16), tetrabenazine, bexarotene, glatirimer acetate, fingolimod, and mitoxantrone, including pharmaceutically acceptable salts and acids thereof. Further examples of cytotoxic or chemotherapeutic agents include alkylating agents such as thiopeta, cyclophosphamide (CYTOXANTM); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as aclacinomysins, actinomycin, anthramycin, azaserine, bleomycins, cactinomycin, calicheamicin, carubicin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin, epirubicin, esorubicin, idarubicin,

marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, pottfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as calusterone, dromostanolone propionate, epitioestanol, mepitioestane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amсарine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; mitoguanzone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK; razoxane; sizofiran; spirogermanium; tenoazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thioteпа; taxoids, e.g., paclitaxel (TAXOL®, Bristol-Myers Squibb Oncology, Princeton, N.J.) and doxetaxel (TAXOTERER®, Rhne-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide; daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylomithine (DMFO); retinoic acid derivatives such as Targretin™ (bexarotene), Panretin™ (alitretinoin); ONTAK™ (denileukin diftiox); esperamicins; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

The antibodies or antigen-binding fragments thereof can be used in any of the compositions, methods, and/or kits described herein, and combined with one or more of the immunotherapy agents described herein.

Additional Therapeutic Agents and Compositions

Immunotherapy Agents. Certain embodiments employ one or more cancer immunotherapy agents. In certain instances, an immunotherapy agent modulates the immune response of a subject, for example, to increase or maintain a cancer-related or cancer-specific immune response, and thereby results in increased immune cell inhibition or reduction of cancer cells. Exemplary immunotherapy agents include polypeptides, for example, antibodies and antigen-binding fragments thereof, ligands, and small peptides, and mixtures thereof. Also include as immunotherapy agents are small molecules, cells (e.g., immune cells such as T-cells), various cancer vaccines, gene therapy or other polynucleotide-based agents, including viral agents such as oncolytic viruses, and others known in the art. Thus, in certain embodiments, the cancer immunotherapy agent is selected from one or more of immune checkpoint modulatory agents, cancer vaccines, oncolytic viruses, cytokines, and a cell-based immunotherapies.

In certain embodiments, the cancer immunotherapy agent is an immune checkpoint modulatory agent. Particular examples include "antagonists" of one or more inhibitory immune checkpoint molecules, and "agonists" of one or more stimulatory immune checkpoint molecules. Generally,

immune checkpoint molecules are components of the immune system that either turn up a signal (co-stimulatory molecules) or turn down a signal, the targeting of which has therapeutic potential in cancer because cancer cells can perturb the natural function of immune checkpoint molecules (see, e.g., Sharma and Allison, *Science*. 348:56-61, 2015; Topalian et al., *Cancer Cell*. 27:450-461, 2015; Pardoll, *Nature Reviews Cancer*. 12:252-264, 2012). In some embodiments, the immune checkpoint modulatory agent (e.g., antagonist, agonist) "binds" or "specifically binds" to the one or more immune checkpoint molecules, as described herein.

In particular embodiments, the immune checkpoint modulatory agent is a polypeptide or peptide. The terms "peptide" and "polypeptide" are used interchangeably herein, however, in certain instances, the term "peptide" can refer to shorter polypeptides, for example, polypeptides that consist of about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, or 50 amino acids, including all integers and ranges (e.g., 5-10, 8-12, 10-15) in between. Polypeptides and peptides can be composed of naturally-occurring amino acids and/or non-naturally occurring amino acids, as described herein.

Antibodies are also included as polypeptides. Thus, in some embodiments, the immune checkpoint modulatory polypeptide agent is an antibody or "antigen-binding fragment thereof", as described elsewhere herein.

In some embodiments, the agent is or comprises a "ligand," for example, a natural ligand, of the immune checkpoint molecule. A "ligand" refers generally to a substance or molecule that forms a complex with a target molecule (e.g., biomolecule) to serve a biological purpose, and includes a "protein ligand," which generally produces a signal by binding to a site on a target molecule or target protein. Thus, certain agents are protein ligands that, in nature, bind to an immune checkpoint molecule and produce a signal. Also included are "modified ligands," for example, protein ligands that are fused to a pharmacokinetic modifier, for example, an Fc region derived from an immunoglobulin.

The binding properties of polypeptides can be quantified using methods well known in the art (see Davies et al., *Annual Rev. Biochem.* 59:439-473, 1990). In some embodiments, a polypeptide specifically binds to a target molecule, for example, an immune checkpoint molecule or an epitope thereof, with an equilibrium dissociation constant that is about or ranges from about $\leq 10^{-7}$ to about 10^{-8} M. In some embodiments, the equilibrium dissociation constant is about or ranges from about $\leq 10^{-9}$ M to about $\leq 10^{-10}$ M. In certain illustrative embodiments, the polypeptide has an affinity (K_d or EC_{50}) for a target described herein (to which it specifically binds) of about, at least about, or less than about, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM.

In some embodiments, the agent is a "small molecule," which refers to an organic compound that is of synthetic or biological origin (biomolecule), but is typically not a polymer. Organic compounds refer to a large class of chemical compounds whose molecules contain carbon, typically excluding those that contain only carbonates, simple oxides of carbon, or cyanides. A "biomolecule" refers generally to an organic molecule that is produced by a living organism, including large polymeric molecules (biopolymers) such as peptides, polysaccharides, and nucleic acids as well, and small molecules such as primary secondary metabolites, lipids, phospholipids, glycolipids, sterols, glycerolipids, vitamins, and hormones. A "polymer" refers generally to a

large molecule or macromolecule composed of repeating structural units, which are typically connected by covalent chemical bond.

In certain embodiments, a small molecule has a molecular weight of about or less than about 1000-2000 Daltons, typically between about 300 and 700 Daltons, and including about or less than about 50, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000 or 2000 Daltons.

Certain small molecules can have the "specific binding" characteristics described for herein polypeptides such as antibodies. For instance, in some embodiments a small molecule specifically binds to a target, for example, an immune checkpoint molecule, with a binding affinity (K_d or EC_{50}) of about, at least about, or less than about, 0.01, 0.05, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 40, or 50 nM.

In some embodiments, the immune checkpoint modulatory agent is an antagonist or inhibitor of one or more inhibitory immune checkpoint molecules. Exemplary inhibitory immune checkpoint molecules include Programmed Death-Ligand 1 (PD-L1), Programmed Death-Ligand 2 (PD-L2), Programmed Death 1 (PD-1), Cytotoxic T-Lymphocyte-Associated protein 4 (CTLA-4), Indoleamine 2,3-dioxygenase (IDO), tryptophan 2,3-dioxygenase (TDO), T-cell Immunoglobulin domain and Mucin domain 3 (TIM-3), Lymphocyte Activation Gene-3 (LAG-3), V-domain Ig suppressor of T cell activation (VISTA), B and T Lymphocyte Attenuator (BTLA), CD160, and T-cell immunoreceptor with Ig and ITIM domains (TIGIT).

In certain embodiments, the agent is a PD-1 (receptor) antagonist or inhibitor, the targeting of which has been shown to restore immune function in the tumor environment (see, e.g., Phillips et al., *Int Immunol.* 27:39-46, 2015). PD-1 is a cell surface receptor that belongs to the immunoglobulin superfamily and is expressed on T cells and pro-B cells. PD-1 interacts with two ligands, PD-L1 and PD-L2. PD-1 functions as an inhibitory immune checkpoint molecule, for example, by reducing or preventing the activation of T-cells, which in turn reduces autoimmunity and promotes self-tolerance. The inhibitory effect of PD-1 is accomplished at least in part through a dual mechanism of promoting apoptosis in antigen specific T-cells in lymph nodes while also reducing apoptosis in regulatory T cells (suppressor T cells). Some examples of PD-1 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to PD-1 and reduces one or more of its immune-suppressive activities, for example, its downstream signaling or its interaction with PD-L1. Specific examples of PD-1 antagonists or inhibitors include the antibodies nivolumab, pembrolizumab, PDR001, MK-3475, AMP-224, AMP-514, and pidilizumab, and antigen-binding fragments thereof (see, e.g., U.S. Pat. Nos. 8,008,449; 8,993,731; 9,073,994; 9,084,776; 9,102,727; 9,102,728; 9,181,342; 9,217,034; 9,387,247; 9,492,539; 9,492,540; and U.S. Application Nos. 2012/0039906; 2015/0203579).

In some embodiments, the agent is a PD-L1 antagonist or inhibitor. As noted above, PD-L1 is one of the natural ligands for the PD-1 receptor. General examples of PD-L1 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to PD-L1 and reduces one or more of its immune-suppressive activities, for example, its binding to the PD-1 receptor. Specific examples of PD-L1 antagonists include the antibodies atezolizumab (MPDL3280A), avelumab (MSB0010718C), and durvalumab (MEDI4736), and anti-

gen-binding fragments thereof (see, e.g., U.S. Pat. Nos. 9,102,725; 9,393,301; 9,402,899; 9,439,962).

In some embodiments, the agent is a PD-L2 antagonist or inhibitor. As noted above, PD-L2 is one of the natural ligands for the PD-1 receptor. General examples of PD-L2 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to PD-L2 and reduces one or more of its immune-suppressive activities, for example, its binding to the PD-1 receptor.

In some embodiments, the agent is a CTLA-4 antagonist or inhibitor. CTLA4 or CTLA-4 (cytotoxic T-lymphocyte-associated protein 4), also known as CD152 (cluster of differentiation 152), is a protein receptor that functions as an inhibitory immune checkpoint molecule, for example, by transmitting inhibitory signals to T-cells when it is bound to CD80 or CD86 on the surface of antigen-presenting cells. General examples CTLA-4 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to CTLA-4. Particular examples include the antibodies ipilimumab and tremelimumab, and antigen-binding fragments thereof. At least some of the activity of ipilimumab is believed to be mediated by antibody-dependent cell-mediated cytotoxicity (ADCC) killing of suppressor Tregs that express CTLA-4.

In some embodiments, the agent is an IDO antagonist or inhibitor, or a TDO antagonist or inhibitor. IDO and TDO are tryptophan catabolic enzymes with immune-inhibitory properties. For example, IDO is known to suppress T-cells and NK cells, generate and activate Tregs and myeloid-derived suppressor cells, and promote tumor angiogenesis. General examples of IDO and TDO antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to IDO or TDO (see, e.g., Platten et al., *Front Immunol.* 5:673, 2014) and reduces or inhibits one or more immune-suppressive activities. Specific examples of IDO antagonists or inhibitors include indoximod (NLG-8189), 1-methyl-tryptophan (1MT), β -Carboline (norharmane; 9H-pyrido[3,4-b]indole), rosmarinic acid, and epacadostat (see, e.g., Sheridan, *Nature Biotechnology.* 33:321-322, 2015). Specific examples of TDO antagonists or inhibitors include 680C91 and LM10 (see, e.g., Pilotte et al., *PNAS USA.* 109:2497-2502, 2012).

In some embodiments, the agent is a TIM-3 antagonist or inhibitor. T-cell Immunoglobulin domain and Mucin domain 3 (TIM-3) is expressed on activated human CD4⁺ T-cells and regulates Th1 and Th17 cytokines. TIM-3 also acts as a negative regulator of Th1/Tc1 function by triggering cell death upon interaction with its ligand, galectin-9. TIM-3 contributes to the suppressive tumor microenvironment and its overexpression is associated with poor prognosis in a variety of cancers (see, e.g., Li et al., *Acta Oncol.* 54:1706-13, 2015). General examples of TIM-3 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to TIM-3 and reduces or inhibits one or more of its immune-suppressive activities.

In some embodiments, the agent is a LAG-3 antagonist or inhibitor. Lymphocyte Activation Gene-3 (LAG-3) is expressed on activated T-cells, natural killer cells, B-cells and plasmacytoid dendritic cells. It negatively regulates cellular proliferation, activation, and homeostasis of T-cells, in a similar fashion to CTLA-4 and PD-1 (see, e.g., Workman and Vignali, *European Journal of Immun.* 33:970-9, 2003; and Workman et al., *Journal of Immun.* 172:5450-5, 2004), and has been reported to play a role in Treg suppressive function (see, e.g., Huang et al., *Immunity.* 21:503-13, 2004). LAG3 also maintains CD8⁺ T-cells in a tolerogenic state and combines with PD-1 to maintain CD8 T-cell

exhaustion. General examples of LAG-3 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to LAG-3 and inhibits one or more of its immune-suppressive activities. Specific examples include the antibody BMS-986016, and antigen-binding fragments thereof.

In some embodiments, the agent is a VISTA antagonist or inhibitor. V-domain Ig suppressor of T cell activation (VISTA) is primarily expressed on hematopoietic cells and is an inhibitory immune checkpoint regulator that suppresses T-cell activation, induces Foxp3 expression, and is highly expressed within the tumor microenvironment where it suppresses anti-tumor T cell responses (see, e.g., Lines et al., *Cancer Res.* 74:1924-32, 2014). General examples of VISTA antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to VISTA and reduces one or more of its immune-suppressive activities.

In some embodiments, the agent is a BTLA antagonist or inhibitor. B- and T-lymphocyte attenuator (BTLA; CD272) expression is induced during activation of T-cells, and it inhibits T-cells via interaction with tumor necrosis family receptors (TNF-R) and β_7 family of cell surface receptors. BTLA is a ligand for tumor necrosis factor (receptor) superfamily, member 14 (TNFRSF14), also known as herpes virus entry mediator (HVEM). BTLA-HVEM complexes negatively regulate T-cell immune responses, for example, by inhibiting the function of human CD8+ cancer-specific T-cells (see, e.g., Derré et al., *J Clin Invest* 120:157-67, 2009). General examples of BTLA antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to BTLA-4 and reduce one or more of its immune-suppressive activities.

In some embodiments, the agent is an HVEM antagonist or inhibitor, for example, an antagonist or inhibitor that specifically binds to HVEM and interferes with its interaction with BTLA or CD160. General examples of HVEM antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to HVEM, optionally reduces the HVEM/BTLA and/or HVEM/CD160 interaction, and thereby reduces one or more of the immune-suppressive activities of HVEM.

In some embodiments, the agent is a CD160 antagonist or inhibitor, for example, an antagonist or inhibitor that specifically binds to CD160 and interferes with its interaction with HVEM. General examples of CD160 antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to CD160, optionally reduces the CD160/HVEM interaction, and thereby reduces or inhibits one or more of its immune-suppressive activities.

In some embodiments, the agent is a TIGIT antagonist or inhibitor. T cell Ig and ITIM domain (TIGIT) is a co-inhibitory receptor that is found on the surface of a variety of lymphoid cells, and suppresses antitumor immunity, for example, via Tregs (Kurtulus et al., *J Clin Invest.* 125:4053-4062, 2015). General examples of TIGIT antagonists or inhibitors include an antibody or antigen-binding fragment or small molecule that specifically binds to TIGIT and reduce one or more of its immune-suppressive activities (see, e.g., Johnston et al., *Cancer Cell.* 26:923-37, 2014).

In certain embodiments, the immune checkpoint modulatory agent is an agonist of one or more stimulatory immune checkpoint molecules. Exemplary stimulatory immune checkpoint molecules include OX40, CD40, Glucocorti-

coid-Induced TNFR Family Related Gene (GITR), CD137 (4-1BB), CD27, CD28, CD226, and Herpes Virus Entry Mediator (HVEM).

In some embodiments, the agent is an OX40 agonist. OX40 (CD134) promotes the expansion of effector and memory T cells, and suppresses the differentiation and activity of T-regulatory cells (see, e.g., Croft et al., *Immunol Rev.* 229:173-91, 2009). Its ligand is OX40L (CD252). Since OX40 signaling influences both T-cell activation and survival, it plays a key role in the initiation of an anti-tumor immune response in the lymph node and in the maintenance of the anti-tumor immune response in the tumor microenvironment. General examples of OX40 agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to OX40 and increases one or more of its immunostimulatory activities. Specific examples include OX86, OX-40L, Fc-OX40L, GSK3174998, MEDI0562 (a humanized OX40 agonist), MEDI6469 (murine OX4 agonist), and MEDI6383 (an OX40 agonist), and antigen-binding fragments thereof.

In some embodiments, the agent is a CD40 agonist. CD40 is expressed on antigen-presenting cells (APC) and some malignancies. Its ligand is CD40L (CD154). On APC, ligation results in upregulation of costimulatory molecules, potentially bypassing the need for T-cell assistance in an antitumor immune response. CD40 agonist therapy plays an important role in APC maturation and their migration from the tumor to the lymph nodes, resulting in elevated antigen presentation and T cell activation. Anti-CD40 agonist antibodies produce substantial responses and durable anticancer immunity in animal models, an effect mediated at least in part by cytotoxic T-cells (see, e.g., Johnson et al. *Clin Cancer Res.* 21:1321-1328, 2015; and Vonderheide and Glennie, *Clin Cancer Res.* 19:1035-43, 2013). General examples of CD40 agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to CD40 and increases one or more of its immunostimulatory activities. Specific examples include CP-870,893, dacetuzumab, Chi Lob 7/4, ADC-1013, CD40L, rhCD40L, and antigen-binding fragments thereof.

In some embodiments, the agent is a GITR agonist. Glucocorticoid-Induced TNFR family Related gene (GITR) increases T cell expansion, inhibits the suppressive activity of Tregs, and extends the survival of T-effector cells. GITR agonists have been shown to promote an anti-tumor response through loss of Treg lineage stability (see, e.g., Schaer et al., *Cancer Immunol Res.* 1:320-31, 2013). These diverse mechanisms show that GITR plays an important role in initiating the immune response in the lymph nodes and in maintaining the immune response in the tumor tissue. Its ligand is GITRL. General examples of GITR agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to GITR and increases one or more of its immunostimulatory activities. Specific examples include GITRL, INCAGN01876, DTA-1, MEDI1873, and antigen-binding fragments thereof.

In some embodiments, the agent is a CD137 agonist. CD137 (4-1BB) is a member of the tumor necrosis factor (TNF) receptor family, and crosslinking of CD137 enhances T-cell proliferation, IL-2 secretion, survival, and cytolytic activity. CD137-mediated signaling also protects T-cells such as CD8+ T-cells from activation-induced cell death. General examples of CD137 agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to CD137 and increases one or more of its immunostimulatory activities. Specific examples include the CD137 (or 4-1BB) ligand (see, e.g., Shao and Schwarz, J

Leukoc Biol. 89:21-9, 2011) and the antibody utomilumab, including antigen-binding fragments thereof.

In some embodiments, the agent is a CD27 agonist. Stimulation of CD27 increases antigen-specific expansion of naïve T cells and contributes to T-cell memory and long-term maintenance of T-cell immunity. Its ligand is CD70. The targeting of human CD27 with an agonist antibody stimulates T-cell activation and antitumor immunity (see, e.g., Thomas et al., *Oncoimmunology*. 2014; 3: e27255. doi: 10.4161/onci.27255; and He et al., *J Immunol*. 191:4174-83, 2013). General examples of CD27 agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to CD27 and increases one or more of its immunostimulatory activities. Specific examples include CD70 and the antibodies varilumab and CDX-1127 (IF5), including antigen-binding fragments thereof.

In some embodiments, the agent is a CD28 agonist. CD28 is constitutively expressed CD4+ T cells some CD8+ T cells. Its ligands include CD80 and CD86, and its stimulation increases T-cell expansion. General examples of CD28 agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to CD28 and increases one or more of its immunostimulatory activities. Specific examples include CD80, CD86, the antibody TAB08, and antigen-binding fragments thereof.

In some embodiments, the agent is CD226 agonist. CD226 is a stimulating receptor that shares ligands with TIGIT, and opposite to TIGIT, engagement of CD226 enhances T-cell activation (see, e.g., Kurtulus et al., *J Clin Invest*. 125:4053-4062, 2015; Bottino et al., *J Exp Med*. 1984:557-567, 2003; and Tahara-Hanaoka et al., *Int Immunol*. 16:533-538, 2004). General examples of CD226 agonists include an antibody or antigen-binding fragment or small molecule or ligand (e.g., CD112, CD155) that specifically binds to CD226 and increases one or more of its immunostimulatory activities.

In some embodiments, the agent is an HVEM agonist. Herpesvirus entry mediator (HVEM), also known as tumor necrosis factor receptor superfamily member 14 (TNFRSF14), is a human cell surface receptor of the TNF-receptor superfamily. HVEM is found on a variety of cells including T-cells, APCs, and other immune cells. Unlike other receptors, HVEM is expressed at high levels on resting T-cells and down-regulated upon activation. It has been shown that HVEM signaling plays a crucial role in the early phases of T-cell activation and during the expansion of tumor-specific lymphocyte populations in the lymph nodes. General examples of HVEM agonists include an antibody or antigen-binding fragment or small molecule or ligand that specifically binds to HVEM and increases one or more of its immunostimulatory activities.

In certain embodiments, the cancer immunotherapy agent is a cancer vaccine. Exemplary cancer vaccines include Oncophage, human papillomavirus HPV vaccines such as Gardasil or Cervarix, hepatitis B vaccines such as Engerix-B, Recombivax HB, or Twinrix, and sipuleucel-T (Provenge). In some embodiments, the cancer vaccine comprises or utilizes one or more cancer antigens, or cancer-associate d antigens. Exemplary cancer antigens include, without limitation, human Her2/neu, Her1/EGF receptor (EGFR), Her3, A33 antigen, β_7 H3, CD5, CD19, CD20, CD22, CD23 (IgE Receptor), MAGE-3, C242 antigen, 5T4, IL-6, IL-13, vascular endothelial growth factor VEGF (e.g., VEGF-A) VEGFR-1, VEGFR-2, VEGFR-3, NRP2, CD30, CD33, CD37, CD40, CD44, CD51, CD52, CD56, CD74, CD80, CD152, CD200, CD221, CCR4, HLA-DR, CTLA-4, NPC-1C, tenascin, vimentin, insulin-like growth factor 1 receptor

(IGF-1R), alpha-fetoprotein, insulin-like growth factor 1 (IGF-1), carbonic anhydrase 9 (CA-IX), carcinoembryonic antigen (CEA), guanylyl cyclase C, NY-ESO-1, p53, survivin, integrin $\alpha\beta$ 3, integrin α 5 β 1, folate receptor 1, transmembrane glycoprotein NMB, fibroblast activation protein alpha (FAP), glycoprotein 75, TAG-72, MUC1, MUC16 (or CA-125), phosphatidylserine, prostate-specific membrane antigen (PSMA), NR-LU-13 antigen, TRAIL-R1, tumor necrosis factor receptor superfamily member 10b (TNFRSF10B or TRAIL-R2), SLAM family member 7 (SLAMF7), EGP40 pancarcinoma antigen, B-cell activating factor (BAFF), platelet-derived growth factor receptor, glycoprotein EpCAM (17-1A), Programmed Death-1, protein disulfide isomerase (PDI), Phosphatase of Regenerating Liver 3 (PRL-3), prostatic acid phosphatase, Lewis-Y antigen, GD2 (a disialoganglioside expressed on tumors of neuroectodermal origin), glypican-3 (GPC3), and mesothelin.

In certain embodiments, the cancer immunotherapy agent is an oncolytic virus. An oncolytic virus is a virus that preferentially infects and kills cancer cells. Included are naturally-occurring and man-made or engineered oncolytic viruses. Most oncolytic viruses are engineered for tumor selectivity, although there are naturally-occurring examples such as Reovirus and the SVV-001 Seneca Valley virus. General examples of oncolytic viruses include VSV, Poliovirus, Reovirus, Senecavirus, and RIGVIR, and engineered versions thereof. Non-limiting examples of oncolytic viruses include herpes simplex virus (HSV) and engineered version thereof, talimogene laherparepvec (T-VEC), coxsackievirus A21 (CAVATAK™), Oncorine (H101), pelareorep (REOLYSIN®), Seneca Valley virus (NTX-010), Senecavirus SVV-001, ColoAd1, SEPREHVIR (HSV-1716), CGTG-102 (Ad5/3-D24-GMCSF), GL-ONC1, MV-NIS, and DNX-2401, among others.

In certain embodiments, the cancer immunotherapy agent is a cytokine. Exemplary cytokines include interferon (IFN)- α , IL-2, IL-12, IL-7, IL-21, and Granulocyte-macrophage colony-stimulating factor (GM-CSF).

In certain embodiments, the cancer immunotherapy agent is cell-based immunotherapy, for example, a T-cell based adoptive immunotherapy. In some embodiments, the cell-based immunotherapy comprises cancer antigen-specific T-cells, optionally ex vivo-derived T-cells. In some embodiments, the cancer antigen-specific T-cells are selected from one or more of chimeric antigen receptor (CAR)-modified T-cells, and T-cell Receptor (TCR)-modified T-cells, tumor infiltrating lymphocytes (TILs), and peptide-induced T-cells. In specific embodiments, the CAR-modified T-cell is targeted against CD-19 (see, e.g., Maude et al., *Blood*. 125: 4017-4023, 2015).

In certain instances, the cancer to be treated associates with the cancer antigen, that is, the cancer antigen-specific T-cells are targeted against or enriched for at least one antigen that is known to associate with the cancer to be treated. In some embodiments, the cancer antigen is selected from one or more of CD19, human Her2/neu, Her1/EGF receptor (EGFR), Her3, A33 antigen, β_7 H3, CD5, CD20, CD22, CD23 (IgE Receptor), MAGE-3, C242 antigen, 5T4, IL-6, IL-13, vascular endothelial growth factor VEGF (e.g., VEGF-A) VEGFR-1, VEGFR-2, CD30, CD33, CD37, CD40, CD44, CD51, CD52, CD56, CD74, CD80, CD152, CD200, CD221, CCR4, HLA-DR, CTLA-4, NPC-1C, tenascin, vimentin, insulin-like growth factor 1 receptor (IGF-1R), alpha-fetoprotein, insulin-like growth factor 1 (IGF-1), carbonic anhydrase 9 (CA-IX), carcinoembryonic antigen (CEA), guanylyl cyclase C, NY-ESO-1, p53, survivin, inte-

grin $\alpha v\beta 3$, integrin $\alpha 5\beta 1$, folate receptor 1, transmembrane glycoprotein NMB, fibroblast activation protein alpha (FAP), glycoprotein 75, TAG-72, MUC1, MUC16 (or CA-125), phosphatidylserine, prostate-specific membrane antigen (PMSA), NR-LU-13 antigen, TRAIL-R1, tumor necrosis factor receptor superfamily member 10b (TNFRSF10B or TRAIL-R2), SLAM family member 7 (SLAMF7), EGP40 pancarcinoma antigen, B-cell activating factor (BAFF), platelet-derived growth factor receptor, glycoprotein EpCAM (17-1A), Programmed Death-1, protein disulfide isomerase (PDI), Phosphatase of Regenerating Liver 3 (PRL-3), prostatic acid phosphatase, Lewis-Y antigen, GD2 (a disialoganglioside expressed on tumors of neuroectodermal origin), glypican-3 (GPC3), and mesothelin.

Additional exemplary cancer antigens include 5T4, 707-AP, 9D7, AFP, AlbZIP HPG1, alpha-5-beta-1-integrin, alpha-5-beta-6-integrin, alpha-actinin-4/m, alpha-methylacyl-coenzyme A racemase, ART-4, ARTC1/m, B7H4, BAGE-1, BCL-2, bcr/abl, beta-catenin/m, BING-4, BRCA1/m, BRCA2/m, CA 15-3/CA 27-29, CA 19-9, CA72-4, CA125, calreticulin, CAMEL, CASP-8/m, cathepsin B, cathepsin L, CDC27/m, CDK4/m, CDKN2A/m, CEA, CLCA2, CML28, CML66, COA-1/m, coactosin-like protein, collagen XXIII, COX-2, CT-9/BRD6, Cten, cyclin β_1 , cyclin D1, cyp-B, CYPB1, DAM-10, DAM-6, DEK-CAN, EFTUD2/m, EGFR, ELF2/m, EMMPRIN, EpCam, EphA2, EphA3, ErbB3, ETV6-AML1, EZH2, FGF-5, FN, Frau-1, G250, GAGE-1, GAGE-2, GAGE-3, GAGE-4, GAGE-5, GAGE-6, GAGE7b, GAGE-8, GDEP, GnT-V, gp100, GPC3, GPNMB/m, HAGE, HAST-2, hepsin, Her2/neu, HERV-K-MEL, HLA-A*0201-R1 7I, HLA-A1 1/m, HLA-A2/m, HNE, homeobox NKX3.1, HOM-TES-14/SCP-1, HOM-TES-85, HPV-E6, HPV-E7, HSP70-2M, HST-2, hTERT, ICE, IGF-1 R, IL-13Ra2, IL-2R, IL-5, immature laminin receptor, kallikrein-2, kallikrein-4, Ki67, KIAA0205, KIAA0205/m, KK-LC-1, K-Ras/m, LAGE-A1, LDLR-FUT, MAGE-A1, MAGE-A2, MAGE-A3, MAGE-A4, MAGE-A6, MAGE-A9, MAGE-A10, MAGE-A12, MAGE-B1, MAGE- β_2 , MAGE-B3, MAGE-B4, MAGE-B5, MAGE-B6, MAGE-B10, MAGE-B1 6, MAGE-B1 7, MAGE-C1, MAGE-C2, MAGE-C3, MAGE-D1, MAGE-D2, MAGE-D4, MAGE-E1, MAGE-E2, MAGE-F1, MAGE-H1, MAGE-L2, mammaglobin A, MART-1/melan-A, MART-2, MART-2/m, matrix protein 22, MCI R, M-CSF, ME1/m, mesothelin, MG50/PXDn, MMP1 1, MN/CA IX-antigen, MRP-3, MUC-1, MUC-2, MUM-1/m, MUM-2/m, MUM-3/m, myosin class I/m, NA88-A, N-acetylglucosaminyltransferase-V, Neo-PAP, Neo-PAP/m, NFYC/m, NGEF, NMP22, NPM/ALK, N-Ras/m, NSE, NY-ESO-B, NY-ESO-1, OA1, OFA-iLRP, OGT, OGT/m, OS-9, OS-9/m, osteocalcin, osteopontin, pi 5, p190 minor bcr-abl, p53, p53/m, PAGE-4, PAI-1, PAI-2, PAP, PART-1, PATE, PDEF, Pim-1 Kinase, Pin-1, Pml/PARalpha, POTE, PRAME, PRDX5/m, prostein, proteinase-3, PSA, PSCA, PSGR, PSM, PSMA, PTPRK/m, RAGE-1, RBAF600/m, RHAMM/CD1 68, RU1, RU2, S-100, SAGE, SART-1, SART-2, SART-3, SCC, SIRT2/m, Sp1 7, SSX-1, SSX-2/HOM-MEL-40, SSX-4, STAMP-1, STEAP-1, survivin, survivin-2B, SYT-SSX-1, SYT-SSX-2, TA-90, TAG-72, TARP, TEL-AML1, TGF-beta, TGFbetaRII, TGM-4, TPI/m, TRAG-3, TRG, TRP-1, TRP-2/6b, TRP/INT2, TRP-p8, tyrosinase, UPA, VEGFR1, VEGFR-2/FLK-1, and WT1. Certain preferred antigens include p53, CA125, EGFR, Her2/neu, hTERT, PAP, MAGE-A1, MAGE-A3, Mesothelin, MUC-1, GP100,

MART-1, Tyrosinase, PSA, PSCA, PSMA, STEAP-1, Ras, CEA and WT1, and more preferably PAP, MAGE-A3, WT1, and MUC-1.

In some embodiments the antigen is selected from MAGE-A1 (e.g., MAGE-A1 according to accession number M77481), MAGE-A2, MAGE-A3, MAGE-A6 (e.g., MAGE-A6 according to accession number NM_005363), MAGE-C1, MAGE-C2, melan-A (e.g., melan-A according to accession number NM_00551 1), GP100 (e.g., GP100 according to accession number M77348), tyrosinase (e.g., tyrosinase according to accession number NM_000372), survivin (e.g., survivin according to accession number AF077350), CEA (e.g., CEA according to accession number NM_004363), Her-2/neu (e.g., Her-2/neu according to accession number M1 1 730), WT1 (e.g., WT1 according to accession number NM_000378), PRAME (e.g., PRAME according to accession number NM_0061 15), EGFR1 (epidermal growth factor receptor 1) (e.g., EGFR1 (epidermal growth factor receptor 1) according to accession number AF288738), MUC1, mucin-1 (e.g., mucin-1 according to accession number NM_002456), SEC61 G (e.g., SEC61 G according to accession number NM_014302), hTERT (e.g., hTERT accession number NM_198253), 5T4 (e.g., 5T4 according to accession number NM_006670), TRP-2 (e.g., TRP-2 according to accession number NM_001 922), STEAP1 (Six-transmembrane epithelial antigen of prostate 1), PSCA, PSA, PSMA, etc.

In some embodiments, the cancer antigen is selected from PCA, PSA, PSMA, STEAP, and optionally MUC-1, including fragments, variants, and derivatives thereof. In some embodiments, the cancer antigen selected from NY-ESO-1, MAGE-C1, MAGE-C2, survivin, 5T4, and optionally MUC-1, including fragments, variants, and derivatives thereof.

In some instances, cancer antigens encompass idiotypic antigens associated with a cancer or tumor disease, particularly lymphoma or a lymphoma associated disease, for example, wherein the idiotypic antigen is an immunoglobulin idiotype of a lymphoid blood cell or a T cell receptor idiotype of a lymphoid blood cell.

In some instances, the cancer antigen-specific T-cells are selected from one or more of chimeric antigen receptor (CAR)-modified T-cells (e.g., targeted against a cancer antigen), and T-cell Receptor (TCR)-modified T-cells, tumor infiltrating lymphocytes (TILs), and peptide-induced T-cells.

The skilled artisan will appreciate that the various cancer immunotherapy agents described herein can be combined with any one or more of the various anti-NRP2 antibodies (including antigen-binding fragments thereof) described herein, and used according to any one or more of the methods or compositions described herein.

Chemotherapeutic Agents. Certain embodiments employ one or more chemotherapeutic agents, for example, small molecule chemotherapeutic agents. Non-limiting examples of chemotherapeutic agents include alkylating agents, anti-metabolites, cytotoxic antibiotics, topoisomerase inhibitors (type I or type II), an anti-microtubule agents, among others.

Examples of alkylating agents include nitrogen mustards (e.g., mechlorethamine, cyclophosphamide, mustine, melphalan, chlorambucil, ifosfamide, and busulfan), nitrosoureas (e.g., N-Nitroso-N-methylurea (MNU), carmustine (BCNU), lomustine (CCNU), semustine (MeCCNU), fotemustine, and streptozotocin), tetrazines (e.g., dacarbazine, mitozolomide, and temozolomide), aziridines (e.g., thiotepa, mytomycin, and diaziquone (AZQ)), cisplatin and derivatives thereof (e.g., carboplatin and oxa-

liplatin), and non-classical alkylating agents (optionally procarbazine and hexamethylmelamine).

Examples of anti-metabolites include anti-folates (e.g., methotrexate and pemetrexed), fluoropyrimidines (e.g., 5-fluorouracil and capecitabine), deoxynucleoside analogues (e.g., ancitabine, enocitabine, cytarabine, gemcitabine, decitabine, azacitidine, fludarabine, nelarabine, cladribine, clofarabine, fludarabine, and pentostatin), and thiopurines (e.g., thioguanine and mercaptopurine);

Examples of cytotoxic antibiotics include anthracyclines (e.g., doxorubicin, daunorubicin, epirubicin, idarubicin, pirarubicin, aclarubicin, and mitoxantrone), bleomycins, mitomycin C, mitoxantrone, and actinomycin. Examples of topoisomerase inhibitors include camptothecin, irinotecan, topotecan, etoposide, doxorubicin, mitoxantrone, teniposide, novobiocin, merbarone, and aclarubicin.

Examples of anti-microtubule agents include taxanes (e.g., paclitaxel and docetaxel) and vinca alkaloids (e.g., vinblastine, vincristine, vindesine, vinorelbine).

The skilled artisan will appreciate that the various chemotherapeutic agents described herein can be combined with any one or more of the various anti-NRP2 antibodies (including antigen-binding fragments thereof) described herein, and used according to any one or more of the methods or compositions described herein.

Hormonal Therapeutic Agents. Certain embodiments employ at least one hormonal therapeutic agent. General examples of hormonal therapeutic agents include hormonal agonists and hormonal antagonists. Particular examples of hormonal agonists include progestogen (progestin), corticosteroids (e.g., prednisolone, methylprednisolone, dexamethasone), insulin like growth factors, VEGF derived angiogenic and lymphangiogenic factors (e.g., VEGF-A, VEGF-A145, VEGF-A165, VEGF-C, VEGF-D, PlGF-2), fibroblast growth factor (FGF), galectin, hepatocyte growth factor (HGF), platelet derived growth factor (PDGF), transforming growth factor (TGF)-beta, androgens, estrogens, and somatostatin analogs. Examples of hormonal antagonists include hormone synthesis inhibitors such as aromatase inhibitors and gonadotropin-releasing hormone (GnRH) s agonists (e.g., leuprolide, goserelin, triptorelin, histrelin) including analogs thereof. Also included are hormone receptor antagonist such as selective estrogen receptor modulators (SERMs; e.g., tamoxifen, raloxifene, toremifene) and anti-androgens (e.g., flutamide, bicalutamide, nilutamide).

Also included are hormonal pathway inhibitors such as antibodies directed against hormonal receptors. Examples include inhibitors of the IGF receptor (e.g., IGF-IR1) such as cixutumumab, dalotuzumab, figitumumab, ganitumab, istratutumab, and robatumumab; inhibitors of the vascular endothelial growth factor receptors 1, 2 or 3 (VEGFR1, VEGFR2 or VEGFR3) such as alacizumab pegol, bevacizumab, icrucumab, ramucirumab; inhibitors of the TGF-beta receptors R1, R2, and R3 such as fresolimumab and metelimumab; inhibitors of c-Met such as naxitamab; inhibitors of the EGF receptor such as cetuximab, depatuxizumab mafodotin, futuximab, imgatuzumab, laprituximab emtansine, matuzumab, modotuximab, necitumumab, nimo-tuzumab, panitumumab, tomuzotuximab, and zalutumumab; inhibitors of the FGF receptor such as aprutumab ixadotin and bemarituzumab; and inhibitors of the PDGF receptor such as olaratumab and tovetumab.

The skilled artisan will appreciate that the various hormonal therapeutic agents described herein can be combined with any one or more of the various anti-NRP2 antibodies (including antigen-binding fragments thereof) described

herein, and used according to any one or more of the methods or compositions described herein.

Kinase Inhibitors. Certain embodiments employ at least one kinase inhibitor, including tyrosine kinase inhibitors. Examples of kinase inhibitors include, without limitation, 5 adavosertib, afanitib, aflibercept, axitinib, bevacizumab, bosutinib, cabozantinib, cetuximab, cobimetinib, crizotinib, dasatinib, entrectinib, erdafitinib, erlotinib, fostamitinib, gefitinib, ibrutinib, imatinib, lapatinib, lenvatinib, mubritinib, nilotinib, panitumumab, pazopanib, pegaptanib, ponatinib, ranibizumab, regorafenib, ruxolitinib, sorafenib, sunitinib, SU6656, tofacitinib, trastuzumab, vandetanib, and vemurafenib. Exemplary PI3 kinase inhibitors include 10 alpelisib, buparlisib, copanlisib, CUDC-907, dactolisib, duvelisib, GNE-477, idelasib, IPI-549, LY294002, ME-401, perifosine, PI-103, pictilisib, PWT33597, RP6503, taselisib, umbralisib, voxalisib, wortmannin, and XL147

The skilled artisan will appreciate that the various kinase inhibitors described herein can be combined with any one or more of the various anti-NRP2 antibodies (including antigen-binding fragments thereof) described herein, and used according to any one or more of the methods or compositions described herein.

Methods of Use and Therapeutic Compositions

Embodiments of the present disclosure relate in part to the discovery that human histidyl-tRNA synthetase (HRS) polypeptides have unexpected biological properties which are relevant to treating a broad range of diseases and conditions, and that certain of these properties relate to the interactions 30 between HRS and human neuropilin 2 (NRP2). Accordingly, antibodies directed against human NRP2, which interfere with the binding between NRP2 and NRP2 ligands and/or modulate NRP2 receptor signaling can be used as standalone therapies in the treatment of diseases, including NRP2-associated diseases, or in combination with other therapeutic agents as described herein.

Certain embodiments therefore include methods of treating, ameliorating the symptoms of, and/or reducing the progression of, a disease or condition in a subject in need thereof, comprising administering to the subject at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin-2 (NRP2) polypeptide. In some instances, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 40 ligand. In some instances, the at least one antibody or antigen-binding fragment thereof mimics or otherwise enhances one or more signaling activities of the NRP2/NRP2 ligand interaction, for example, by acting as an agonist antibody.

In some embodiments the anti-NRP2 antibody binds selectively to the NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table N1) of NRP2, and does not substantially bind to the NRP2b isoform (e.g., variants 4 and/or 5 of Table N1). In some embodiments the anti-NRP2 antibody binds selectively to the NRP2b isoform (e.g., variants 4 and/or 5 of Table N1), and does not substantially bind to the NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table N1). Such binding characteristics can apply to any one or more of the 50 embodiments described herein.

In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of VEGF-C or related NRP2 ligands to or via an NRP2 polypeptide. In some aspects, such antibodies or antigen-binding fragment thereof do not substantially block the interaction of semaphorins to an NRP2 polypeptide. In some aspects, such antibodies or antigen-

binding fragment thereof are antagonistic antibodies with respect to VEGF-C signaling via an NRP2 polypeptide. In some embodiments, such antibodies are agonistic antibodies with respect to semaphorin 3F signaling via an NRP2 polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between an NRP2 polypeptide and VEGFR2 or VEGFR3 without substantially modulating the binding/signaling activity between an NRP2 polypeptide and a plexin receptor and/or a semaphorin.

Some embodiments comprise administering at least one anti-NRP2 antibody to a subject in an amount and at a frequency sufficient to achieve an average, sustained blood plasma concentration of soluble NRP2 of about or less than about 500 pM, 400 pM, 300 pM, 200 pM, 100 pM, 50 pM, 40 pM, 30 pM, 20 pM, or 10 pM. Certain embodiments comprise administering at least one anti-NRP2 antibody in an amount and at a frequency sufficient to achieve a reduction in the circulating levels of HRS: NRP2 complexes, for example, a reduction of about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 95, 99, or 100%. Some embodiments include administering the at least one anti-NRP2 antibody in an amount and at a frequency sufficient to achieve a steady state concentration, or average circulating concentration, of the at least one anti-NRP2 antibody of between about 1 nM and about 1 μ M, between about 1 nM and about 100 nM, between about 1 nM and about 10 nM, or between about 1 nM and about 3 μ M.

In certain embodiments, the disease is or condition is an NRP2-associated disease or condition. In some embodiments, the NRP2-associated disease or condition is selected from one or more of cancer and diseases and pathways associated with cancer, including cancer cell growth, initiation, migration, adhesion, invasion, chemoresistance, and/or metastasis; diseases associated with inflammation, autoimmunity, and related inflammatory diseases, including diseases associated with inappropriate immune cell activation or migration such as Graft versus host disease (GVHD); diseases associated with lymphatic development, lymphangiogenesis, and lymphatic damage, including, for example, edema, lymphedema, secondary lymphedema, inappropriate fat absorption and deposition, excess fat deposition, and vascular permeability; diseases associated with infections, including latent infections; diseases associated with allergic disorders/diseases, allergic responses, including, for example, chronic obstructive pulmonary disorder (COPD), neutrophilic asthma, antineutrophil cytoplasmic antibody (ANCA)-associated systemic vasculitis, systemic lupus erythematosus, rheumatoid arthritis, inflammasome-related diseases, and skin-related neutrophil-mediated diseases such as pyoderma gangrenosum; diseases associated with granulomatous inflammatory diseases, including sarcoidosis and granulomas; diseases associated with fibrosis including fibrotic diseases, fibrosis, endothelial to mesenchymal transition (EMT), and wound healing; diseases associated with inappropriate smooth muscle contractility, smooth muscle compensation and decompensation, and inappropriate vascular smooth muscle cell migration and adhesion; diseases associated with inappropriate autophagy, phagocytosis, and efferocytosis; diseases associated with neuronal diseases, peripheral nervous system remodeling, and pain perception; and diseases associated with bone development and bone remodeling.

In some embodiments, the disease is a cancer. Here, upregulation of NRP2 expression is associated with tumorigenesis, and in particular tumor metastasis and development of chemoresistance, and is correlated with more aggressive

disease in several tumor types. Moreover, the semaphorin/plexin/neuropilin signaling axis influences many of the hallmarks of cancer (see, for example, Franzolin and Tamagnone *Int. J. Mol. Sci.* 20, 377; doi: 10.3390/ijms20020377, 2019); Nasarre et al., *OncoTargets and Therapy* 2014;7 1663-1687; Neufeld et al., *Cold Spring Harb Perspect Med.* 2: a006718, 2012). For example, Semaphorin 3F (SEMA3F), which was initially identified as tumor suppressor gene in lung cancer, and is consistently downregulated in highly metastatic tumor cells, including prostate, bladder, and melanoma cells in vitro and in vivo. Consistent with these studies, increased expression of NRP2 in prostate cancer cells is also induced by the loss of the tumor suppressor gene phosphatase and tensin homolog (PTEN), and its expression correlates with Gleason grade (see, for example, Zhao et al., *Thoracic Cancer* 8:203-213, 2017). Additionally, P53 mutations upregulate NRP2 expression via suppression of DLX2 transcription leading to increased cell mobility. About 50% of human tumors and cancers contain a mutation in the p53 tumor suppressor gene, with the vast majority of these mutations occurring in the DNA binding domain, and such is linked to poorer prognosis (see, for example, Drabkin et al., *Oncotarget* 8 (No 57) 96464-96465, 2017). Additionally, TGF- β signaling is involved in the expression of NRP2b, and up-regulation of EMT in cancer, which may explain why in advanced tumors, higher production of TGF- β is positively associated with tumor aggressiveness and poor prognosis (see, for example, Malfettone et al., *Cancer Lett.* 392:39-50, 2017).

Certain embodiments thus include methods of treating ameliorating the symptoms of, or inhibiting the progression of, a cancer in a subject in need thereof, comprising administering to the subject at least one antibody or antigen-binding fragment thereof that specifically binds to a human NRP2 polypeptide (an anti-NRP2 antibody), and which modulates (e.g., interferes with) binding of the human NRP2 polypeptide to a NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or a human HRS polypeptide from Table H1). Certain embodiments include reducing or preventing the re-emergence of a cancer in a subject in need thereof, for example, a metastatic cancer and/or a chemoresistant cancer, wherein administration of the therapeutic composition enables generation of an immune memory to the cancer.

For the treatment of cancer, in some instances, an anti-NRP2 antibody enhances the immune response to the cancer by about, or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more, relative to an untreated control. Exemplary immune responses include increasing or enhancing immune cell invasion of a solid tumor, and increasing the biological activity against the cancer. In certain embodiments, an anti-NRP2 antibody enhances an adaptive immune response to the cancer, and in some embodiments, an anti-NRP2 antibody enhances an innate immune response to the cancer. In some instances, an anti-NRP2 antibody directly or indirectly enhances a T-cell-mediated response to the cancer. In some instances, an anti-NRP2 antibody enhances a B-cell-mediated or antibody-mediated response to the cancer. In some instances, an anti-NRP2 antibody modulates a macrophage responses to the cancer. In some instances, an anti-NRP2 antibody modulates immune cell, or cancer autophagy. In some instances, an anti-NRP2 antibody modulates immune cell phagocytosis. In some instances, an anti-NRP2 antibody modulates cancer cell apoptosis. In some instances, an anti-NRP2 antibody modulates immune cell efferocytosis and/or cancer cell autophagy.

In some embodiments, an anti-NRP2 antibody enhances macrophage responses to the cancer. In some embodiments, an anti-NRP2 inhibits macrophage responses to the cancer. In some embodiments of the anti-NRP2 antibody, the antibody enhances autophagy. In some embodiments, an anti-NRP2 inhibits autophagy. In some embodiments, an anti-NRP2 enhances phagocytosis. In some embodiments, an anti-NRP2 inhibits phagocytosis. In some embodiments, an anti-NRP2 enhances apoptosis. In some embodiments of the anti-NRP2 antibody, the antibody inhibits apoptosis. In some embodiments, an anti-NRP2 antibody enhances effecrocytosis. In some embodiments, an anti-NRP2 inhibits effecrocytosis.

In some instances, an anti-NRP2 antibody reduces cancer initiation, cancer cell migration, adhesion, or cancer cell metastasis by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some instances, an anti-NRP2 antibody reduces cancer mediated lymphangiogenesis by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In some embodiments, the at least one anti-NRP2 antibody reduces the rate of in vitro growth of the cancer (for example, cancer cells isolated from a biopsy or other sample grown in vitro) by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In some embodiments, the at least one anti-NRP2 antibody reduces the adhesiveness of the cancer (for example, cancer cells isolated from a biopsy or other sample grown in vitro) to a substrate by about or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control. In some instances, the substrate comprises laminin.

In some embodiments, the at least one anti-NRP2 antibody reduces the invasiveness of the cancer (for example, cancer cells isolated from a biopsy or other sample grown in vitro) by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In some embodiments, the at least one anti-NRP2 antibody inhibits the rate of migration or motility of the cancer or a migratory cell (for example, cancer or immune cells

isolated from a biopsy or other sample grown in vitro) by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In some embodiments, the at least one anti-NRP2 antibody inhibits the rate of autophagy or endosome maturation (for example, endosome acidification) of the cancer or associated immune cells by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to an untreated control.

In some embodiments, the at least one anti-NRP2 antibody enhances the susceptibility of the cancer to an additional agent (for example, chemotherapeutic agent, hormonal therapeutic agent, and or kinase inhibitor) by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more relative to the additional agent alone. In some embodiments, the at least one anti-NRP2 antibody enhances an anti-tumor and/or immunostimulatory activity of a cancer immunotherapy agent by about or at least about 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more, relative to the cancer immunotherapy agent alone.

Also include are combination therapies for treating cancers, including methods of treating ameliorating the symptoms of, or inhibiting the progression of, a cancer in a subject in need thereof, comprising administering to the subject at least one antibody or antigen-binding fragment thereof that specifically binds to a human NRP2 polypeptide (an anti-NRP2 antibody) in combination with at least one additional agent, for example, a cancer immunotherapy agent, a chemotherapeutic agent, a hormonal therapeutic agent, and/or a kinase inhibitor. Exemplary cancer immunotherapy agents, chemotherapeutic agents, hormonal therapeutic agents, and kinase inhibitors are described elsewhere herein.

In some instances, an anti-NRP2 antibody and the at least one additional agent are administered separately, for example, in separate therapeutic compositions and at the same or different times. In some embodiments, an anti-NRP2 antibody and the at least one additional agent are administered as part of the same therapeutic composition, at the same time.

Particular methods employ one or more anti-NRP2 antibodies, or antigen-binding fragments thereof, as part of (i.e., in addition to) a combination therapy regimen. Exemplary combination regimens are provided in Table M1 below.

TABLE M1

Combination Therapy Regimens		
Cancer Type	Agents	Acronym
Breast Cancer	Cyclophosphamide, methotrexate, 5-fluorouracil, vinorelbine	CMF
	Doxorubicin, cyclophosphamide	AC
Hodgkin's lymphoma	Docetaxel, doxorubicin, cyclophosphamide	TAC
	Doxorubicin, bleomycin, vinblastine, dacarbazine	ABVD
	Mustine, vincristine, procarbazine, prednisolone	MOPP
Non-Hodgkin's lymphoma	Cyclophosphamide, doxorubicin, vincristine, prednisolone	CHOP
Germ cell tumor	Bleomycin, etoposide, cisplatin	BEP
Stomach cancer	Epirubicin, cisplatin, 5-fluorouracil	ECF
	Epirubicin, cisplatin, capecitabine	ECX
Bladder cancer	Methotrexate, vincristine, doxorubicin, cisplatin	MVAC
Lung cancer	Cyclophosphamide, doxorubicin, vincristine, vinorelbine	CAV

TABLE M1-continued

Combination Therapy Regimens		
Cancer Type	Agents	Acronym
Colorectal cancer	5-fluorouracil, folinic acid, oxaliplatin	FOLFOX
Pancreatic Cancer	Leucovorin, fluorouracil, irinotecan (Camptosar), oxaliplatin	FOLFIRINOX
	Gemcitabine, nabpaclitaxel	ABRAXANE

In some embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with at least one additional agent) increase median survival time of a subject by 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks, 15 weeks, 20 weeks, 25 weeks, 30 weeks, 40 weeks, or longer. In certain embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with at least one additional agent) increase median survival time of a subject by 1 year, 2 years, 3 years, or longer. In some embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with cancer immunotherapy agent) increase progression-free survival by 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 7 weeks, 8 weeks, 9 weeks, 10 weeks or longer. In certain embodiments, the methods or therapeutic compositions described herein increase progression-free survival by 1 year, 2 years, 3 years, or longer.

In certain embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with at least one additional agent) are sufficient to result in tumor regression, as indicated by a statistically significant decrease in the amount of viable tumor, for example, at least a 10%, 20%, 30%, 40%, 50% or greater decrease in tumor mass, or by altered (e.g., decreased with statistical significance) scan dimensions. In certain embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with at least one additional agent) are sufficient to result in stable disease. In certain embodiments, the methods and therapeutic compositions described herein (for example, anti-NRP2 antibody, alone or in combination with cancer immunotherapy agent) are sufficient to result in clinically relevant reduction in symptoms of a particular disease indication known to the skilled clinician.

In some embodiments, an anti-NRP2 antibody increases, complements, or otherwise enhances the anti-tumor and/or immunostimulatory activity of the cancer immunotherapy agent, relative to the cancer immunotherapy agent alone. In some embodiments, an anti-NRP2 antibody enhances the anti-tumor and/or immunostimulatory activity of the cancer immunotherapy agent by about, or at least about, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 2000% or more, relative to the cancer immunotherapy agent alone.

The methods and therapeutic compositions described herein can be used in the treatment of any variety of cancers or tumors. In some embodiments, the cancer is a primary cancer, i.e., a cancer growing at the anatomical site where tumor progression began and yielded a cancerous mass. In some embodiments, the cancer is a secondary or metastatic cancer, i.e., a cancer which has spread from the primary site or tissue of origin into one or more different sites or tissues. In some embodiments, the cancer expresses or overex-

presses NRP2. In some embodiments, the subject or patient has a cancer selected from one or more of melanoma (e.g., metastatic melanoma), an epithelial or epithelial-derived tumor, pancreatic cancer, bone cancer, prostate cancer, small cell lung cancer, non-small cell lung cancer (NSCLC), mesothelioma, leukemia (e.g., lymphocytic leukemia, chronic myelogenous leukemia, acute myeloid leukemia, relapsed acute myeloid leukemia), lymphoma, hepatoma (hepatocellular carcinoma or HCC), sarcoma, B-cell malignancy, breast cancer (for example, estrogen receptor positive (ER+), estrogen receptor negative (ER-), Her2 positive (Her2+), Her2 negative (Her2-), or a combination thereof, e.g., ER+/Her2+, ER+/Her2-, ER-/Her2+, or ER-/Her2-; or "triple negative" breast cancer which is estrogen receptor-negative, progesterone receptor-negative, and HER2-negative), ovarian cancer, colorectal cancer, glioma (e.g., astrocytoma, oligodendroglioma, ependymoma, or a choroid plexus papilloma), glioblastoma multiforme (e.g., giant cell glioblastoma or a gliosarcoma), meningioma, pituitary adenoma, vestibular schwannoma, primary CNS lymphoma, primitive neuroectodermal tumor (medulloblastoma), kidney cancer (e.g., renal cell carcinoma), bladder cancer, uterine cancer, esophageal cancer, brain cancer, head and neck cancers, cervical cancer, testicular cancer, thyroid cancer, stomach cancer, virus-induced tumors such as, for example, papilloma virus-induced carcinomas (e.g., cervical carcinoma, cervical cancer), adenocarcinomas, herpes virus-induced tumors (e.g., Burkitt's lymphoma, EBV-induced B-cell lymphoma), hepatitis B-induced tumors (hepatocellular carcinomas), HTLV-1-induced and HTLV-2-induced lymphomas, acoustic neuroma, lung cancers (e.g., lung carcinoma, bronchial carcinoma), small-cell lung carcinomas, pharyngeal cancer, anal carcinoma, glioblastoma, rectal carcinoma, lymphangioma, astrocytoma, brain tumors, retinoblastoma, basalioma, brain metastases, medulloblastomas, vaginal cancer, pancreatic cancer, testicular cancer, Hodgkin's syndrome, meningiomas, Schneeberger disease, hypophysis tumor, Mycosis fungoides, carcinoids, neurinoma, spinalioma, Burkitt's lymphoma, laryngeal cancer, renal cancer, thymoma, corpus carcinoma, bone cancer, non-Hodgkin's lymphomas, urethral cancer, CUP syndrome, head/neck tumors, oligodendroglioma, vulval cancer, intestinal cancer, colon carcinoma, oesophageal cancer (e.g., oesophageal carcinoma), wart involvement, tumors of the small intestine, craniopharyngeomas, ovarian carcinoma, genital tumors, ovarian cancer (e.g., ovarian carcinoma), pancreatic cancer (e.g., pancreatic carcinoma), endometrial carcinoma, liver metastases, penile cancer, tongue cancer, gall bladder cancer, leukaemia, plasmocytoma, and lid tumor.

In some embodiments, as noted above, the cancer or tumor is a metastatic cancer, for example, a metastatic cancer that expresses NRP2a and/or NRP2b. Further to the above cancers, exemplary metastatic cancers include, without limitation, bladder cancers which have metastasized to the bone, liver, and/or lungs; breast cancers which have

metastasized to the bone, brain, liver, and/or lungs; colorectal cancers which have metastasized to the liver, lungs, and/or peritoneum; kidney cancers which have metastasized to the adrenal glands, bone, brain, liver, and/or lungs; lung cancers which have metastasized to the adrenal glands, bone, brain, liver, and/or other lung sites; melanomas which have metastasized to the bone, brain, liver, lung, and/or skin/muscle; ovarian cancers which have metastasized to the liver, lung, and/or peritoneum; pancreatic cancers which have metastasized to the liver, lung, and/or peritoneum; prostate cancers which have metastasized to the adrenal glands, bone, liver, and/or lungs; stomach cancers which have metastasized to the liver, lung, and/or peritoneum; thyroid cancers which have metastasized to the bone, liver, and/or lungs; and uterine cancers which have metastasized to the bone, liver, lung, peritoneum, and/or vagina; among others.

In some embodiments, for example, where the cancer immunotherapy agent is a PD-1 or PD-L1 antagonist or inhibitor, the subject has one or more biomarkers (e.g., increased PD-1 or PD-L1 levels in cells such as cancer cells or cancer-specific CTLs) that make the suitable for PD-1 or PD-L1 inhibitor therapy. For instance, in some embodiments, the subject has increased fractions of programmed cell death 1 high/cytotoxic T lymphocyte-associated protein 4 high (e.g., PD-1^{hi}CTLA-4^{hi}) cells within a tumor-infiltrating CD8⁺ T cell subset (see, e.g., Daud et al., *J Clin Invest*. 126:3447-3452, 2016). As another example, in some embodiments, the subject has increased levels of Bim (B cell lymphoma 2-interacting (Bcl2-interacting) mediator) in circulating tumor-reactive (e.g., PD-1⁺CD11a^{hi}CD8⁺) T cells, and optionally has metastatic melanoma (see, e.g., Dronca et al., *JCI Insight*. May 5; 1 (6): e86014, 2016).

Certain specific combinations include an anti-NRP2 antibody and a PD-L1 antagonist or inhibitor, for example, atezolizumab (MPDL3280A), avelumab (MSB0010718C), and durvalumab (MEDI4736), for treating a cancer selected from one or more of colorectal cancer, melanoma, breast cancer, non-small-cell lung carcinoma, bladder cancer, and renal cell carcinoma.

Some specific combinations include an anti-NRP2 antibody and a PD-1 antagonist, for example, nivolumab, for treating a cancer selected from one or more of Hodgkin's lymphoma, melanoma, non-small cell lung cancer, hepatocellular carcinoma, renal cell carcinoma, and ovarian cancer.

Particular specific combinations include an anti-NRP2 antibody and a PD-1 antagonist, for example, pembrolizumab, for treating a cancer selected from one or more of melanoma, non-small cell lung cancer, small cell lung cancer, head and neck cancer, and urothelial cancer.

Certain specific combinations include an anti-NRP2 antibody and a CTLA-4 antagonist, for example, ipilimumab and tremelimumab, for treating a cancer selected from one or more of melanoma, prostate cancer, lung cancer, and bladder cancer.

Some specific combinations include an anti-NRP2 antibody and an IDO antagonist, for example, indoximod (NLG-8189), 1-methyl-tryptophan (1MT), β -Carboline (norharmane; 9H-pyrido[3,4-b]indole), rosmarinic acid, or epacadostat, for treating a cancer selected from one or more of metastatic breast cancer and brain cancer optionally Glioblastoma Multiforme, glioma, gliosarcoma or malignant brain tumor.

Certain specific combinations include an anti-NRP2 antibody and the cytokine INF- α for treating melanoma, Kaposi sarcoma, and hematologic cancers. Also included is the

combination of an anti-NRP2 antibody and IL-2 (e.g., Aldesleukin) for treating metastatic kidney cancer or metastatic melanoma.

Some specific combinations include an anti-NRP2 antibody and a T-cell based adoptive immunotherapy, for example, comprising CAR-modified T-cells targeted against CD-19, for treating hematological cancers such as acute lymphoblastic leukemia (ALL), chronic lymphocytic leukemia (CLL), and B-cell neoplasms (see, e.g., Maude et al., 2015, *supra*; Lorentzen and Straten, *Scand J Immunol*. 82:307-19, 2015; and Ramos et al., *Cancer J*. 20:112-118, 2014).

The methods for treating cancers can be combined with other therapeutic modalities. For example, a combination therapy described herein can be administered to a subject before, during, or after other therapeutic interventions, including symptomatic care, radiotherapy, surgery, transplantation, hormone therapy, photodynamic therapy, antibiotic therapy, or any combination thereof. Symptomatic care includes administration of corticosteroids, to reduce cerebral edema, headaches, cognitive dysfunction, and emesis, and administration of anti-convulsants, to reduce seizures. Radiotherapy includes whole-brain irradiation, fractionated radiotherapy, and radiosurgery, such as stereotactic radiosurgery, which can be further combined with traditional surgery.

Certain embodiments include the use of the anti-NRP2 antibodies described herein to modulate lymphangiogenesis, and treat related conditions such as lymphedema or tumor metastasis. The lymphatic system consists of networks of interconnected capillaries, collecting vessels and lymph nodes that absorb, collect and transport the fluid and protein filtered from the blood vascular system. This system provides a critical homeostatic function: in humans, lymphatic vessels return >4 liters of fluid and a substantial amount of protein per day back into the great veins of the neck.

Lymphatic vascular dysfunction (lymphedema) results in the accumulation of excess fluid (edema) in the interstitium. Although lymphedema is typically not life-threatening, it has serious health consequences, including pain, immobility, fibrosis, inflammation, adipose tissue accumulation, and tissue damage. Because the lymphatic system is also a critical component of immune responses, lymphedema is typically accompanied by an increased risk of infection and other immune system problems.

Lymphangiogenesis is the formation of new lymphatic vessels from preexisting lymphatic vessels and is associated with diverse pathological conditions including metastatic dissemination, graft rejection (e.g., cornea, kidney and heart), type 2 diabetes, obesity, hypertension, and lymphedema (See, e.g. Alitalo et al. *Nature* 438:946-953, 2005; Karaman et al. *J Clin Invest* 124:922-928, 2014; Kim et al., *J Clin Invest* 124:936-942, 2014; Maby-E1 Hajjami et al., *Histochem Cell Biol* 130:1063-107, 2008; Machnik et al., *Nat Med* 15:545-552; Mortimer et al., 2014. *J Clin Invest* 124:915-921; Skobe et al., 2009. *Nat Med* 15:993-994).

Lymphatic vessel invasion in and around a primary tumor compared to invasion of blood vessels is a prognostic marker of the aggressiveness of various types of cancers. Growth of lymphatic vessels is also involved in graft rejection (Dietrich, T., et al., *J Immunol* 184:535-539, 2010, Hall et al., *Arch Otolaryngol Head Neck Surg* 129:716-719, 2003.; Maula et al., *Cancer Res* 63:1920-1926, 2003; Miyata et al., *J Urol* 176:348-353, 2006; Saad et al., *Mod Pathol* 19:1317-1323, 2006; Schoppmann et al., *Ann Surg* 240:306-312, 2004; Zeng et al., *Prostate* 65:222-230, 2005).

Despite the well-established significance of lymphatics in the pathogenesis of numerous diseases, there has been little progress in the development of anti-lymphangiogenic agents compared to the abundance of anti-angiogenic agents that have entered clinical trials. Therefore, the development of additional lymphangiogenesis inhibitors is of interest for the treatment of a range of conditions, including lymphedema and cancer metastasis.

Anti-lymphangiogenic agents are useful, for example, for treatment of debilitating diseases of the eye, where the growth of lymphatic vessels is the major reason of corneal graft rejection and a major contributor to neovascularization associated with age related macular degeneration (Dietrich, T., et al., *J Immunol* 184:535-539, 2010). In particular, penetrating keratoplasty is the most common form of solid tissue transplantation, with approximately 40,000 corneal transplantations performed each year in the United States. The success rate of penetrating keratoplasty is as high as 90% for uncomplicated first grafts performed in avascular low-risk beds. However, the rejection rate of the corneal grafts placed in high-risk vascularized host beds is extremely high (70% to 90%). Thus the development of safe and targeted new regimens to inhibit lymphangiogenesis are needed to promote graft survival and inhibit neovascularization.

Anti-lymphangiogenesis drugs are useful also for treatment of dry eye disease. Significant upregulation of pro-lymphangiogenic factors (e.g., VEGF-C, VEGF-D, and VEGFR-3) and selective growth of lymphatic vessels without concurrent growth of blood vessels has been demonstrated in corneas with dry eye disease (Goyal et al., *Arch Ophthalmol* 128:819-824, 2010). Dry eye disease is an immune-mediated disorder affecting about 5 million Americans. It severely impacts the vision-related quality of life and the symptoms can be debilitating. The current therapeutic options for dry eye disease are limited, mostly palliative, and expensive. Therefore, development of lymphangiogenesis inhibitors is of therapeutic value for treatment of dry eye disease.

It is currently accepted that metastases are responsible for the vast majority (90%) of deaths from solid tumors (Gupta and Massague, *Cell* 127, 679-695, 2006). The complex process of metastasis involves a series of distinct steps including detachment of tumor cells from the primary tumor, intravasation of tumor cells into lymphatic or blood vessels, and extravasation and growth of tumor cells in secondary sites. Analysis of regional lymph nodes in many tumor types suggests that the lymphatic vasculature is an important route for the dissemination of human cancers. Furthermore, in almost all carcinomas, the presence of tumor cells in lymph nodes is the most important adverse prognostic factor. While it was previously thought that such metastases exclusively involved passage of malignant cells along pre-existing lymphatic vessels near tumors, recent experimental studies and clinicopathological reports (See, e.g., Achen et al., *Br J Cancer* 94, 1355-1360, 2006 and Nathanson, *Cancer* 98, 413-423, 2003) suggest that lymphangiogenesis can be induced by solid tumors and can promote tumor spread. These and other recent studies suggest targeting lymphatics and lymphangiogenesis may be a useful therapeutic strategy to restrict the development of cancer metastasis, which would have a significant benefit for many patients.

Accordingly, there is a need for methods and compositions that inhibit the activities of pro-lymphangiogenic factors and for methods that prevent or treat graft rejection, dry-eye disease tumor metastasis, lymphedema, and inflammatory conditions.

Given the role of NRP2 in these processes, anti-NRP2 antibodies or antigen-binding fragments thereof represent powerful new tools for the development of both pro and anti-lymphangiogenic therapies. Such differential effects may be mediated for example, via the uses of different anti-NRP2 compositions, differential NRP2a or NRP2b selectivity, differential dosing, differential treatment duration, or the appropriate use of additional co-factors, such as, for example, VEGF-C, or Semaphorin 3F and/or 3G.

Accordingly, in some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of VEGF-C or related NRP2 ligands to or via an NRP2 polypeptide. In some aspects, such antibodies or antigen-binding fragment thereof do not substantially block the interaction of semaphorins to an NRP2 polypeptide. In some embodiments, the at least one antibody or antigen-binding fragment thereof selectively modulates the binding and/or signaling of Semaphorin 3F or related NRP2 ligands to or via an NRP2 polypeptide. In some aspects, such antibodies do not substantially block the interaction of VEGF-C to an NRP2 polypeptide. In some embodiments, such antibodies selectively modulate both the binding of VEGF-C or related NRP2 ligands and semaphorins to an NRP2 polypeptide.

In some embodiments, such antibodies are agonistic antibodies with respect to VEGF-C signaling. In some aspects, such antibodies are antagonistic antibodies with respect to VEGF-C signaling. In some embodiments, such antibodies are agonistic antibodies with respect to semaphorin 3F signaling. In some aspects, such antibodies are antagonistic antibodies with respect to semaphorin 3F signaling.

In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between an NRP2 polypeptide and VEGFR2 or VEGFR3 without substantially modulating the binding/signaling activity between the NRP2 polypeptide and a plexin receptor and/or a semaphorin. In some embodiments, the at least one antibody or antigen-binding fragment thereof antagonizes the binding/signaling activity between an NRP2 polypeptide and a plexin receptor and/or a semaphorin without substantially modulating the binding/signaling activity between the NRP2 polypeptide and VEGFR2, VEGFR3, and/or VEGF-C.

Certain embodiments thus include methods of treating ameliorating the symptoms of, or inhibiting the progression of, lymphangiogenesis, or neovascularization in a subject in need thereof, comprising administering to the subject at least one antibody or antigen-binding fragment thereof that specifically binds to a human NRP2 polypeptide, and which modulates (e.g., interferes with) binding of the human NRP2 polypeptide to VEGF-C, VEGFR1, VEGFR2, and/or VEGFR3.

Some embodiments include the use of the anti-NRP2 antibodies described herein to modulate smooth muscle contractility, and treat related conditions. Reduced smooth muscle (SM) contractility in the bladder can stem from numerous etiologies including partial obstruction secondary to benign prostatic hyperplasia (BPH), posterior urethral valves, diabetes mellitus, multiple sclerosis, spinal cord injury, or idiopathic causes. (See, e.g., Drake et al., *Nat Rev Urol.* 11 (8): 454-464, 2014). In conditions such as BPH or posterior urethral valves, the bladder contracts against an obstructed outlet. The initial response is adaptive, involving a compensatory phase of SM hypertrophy that enables increased force generation to overcome the increased outlet resistance. When the demand outstrips the adaptive capability of the bladder, contractile performance becomes less

efficient, residual volumes increase, and the bladder remodels, ultimately leading to a loss of detrusor contractility as the bladder decompensates. (See, e.g., Zderic et al., *J Cell Mol Med.* 16 (2): 203-217, 2012). The prevalence of underactive detrusor function is reported to be as high as 48% in adults (Osman et al., *Eur Urol*; 65 (2): 389-398, 2014). Furthermore, existing pharmacological treatments for restoration of SM contraction such as muscarinic agonists or cholinesterase inhibitors have shown limited efficacy and adverse effects (Barendrecht et al., *BJU Int.* 99 (4): 749-752, 2007).

Recent studies have identified bladder smooth muscle as a major site of Nrp2 expression, demonstrated the inhibition of RhoA and cytoskeletal stiffness, and observed increased contractility of bladder SM strips from mice with ubiquitous or smooth muscle-specific deletion of NRP2 in vivo, when compared with tissues from NRP2-intact littermate controls (See, e.g., Bielenberg et al., *Am. J. Pathol.* 181 548-559, 2012; and Vasquez et al., *JCI Insight* 2 (3) e90617, 2017).

Collectively, these findings suggest that downregulation of NRP2 is an important component of the compensatory response to obstruction in experimental animals and in humans with obstruction, and suggest that Nrp2 may represent a novel pharmacological target for maintenance or restoration of detrusor contractility in the decompensated bladder.

Furthermore, recent studies have shown that targeting NRP2 in bladders undergoing decompensation has the potential to restore contractility in spite of ongoing obstruction. (Vasquez et al., *JCI Insight* 2 (3) e90617, 2017). These findings argue that the NRP2 axis represents a potentially novel pharmacologic target for restoration of SM contractility in partial bladder outlet obstruction syndromes, and provide an important therapeutic opportunity for the development of antibody-based modulators of NRP2 function.

Pharmacological management of diminished detrusor contractility to date has focused on stimulation of parasympathetic activity to enhance bladder contractility and reduction of outflow resistance to facilitate bladder emptying (Chancellor et al., *Urology* 72 (5) 966-967, 2008). However, analysis of 10 randomized clinical trials of parasympathomimetic drugs in patients with poorly contractile bladders, revealed either a worsening of symptoms or a lack of significant improvement (Barendrecht et al., *BJU Int.* 99 (4) 749-752, 2007). The increase in contractility following NRP2 deletion in the decompensating bladder suggests that NRP2 may be a useful target to mitigate reduced detrusor contractility under conditions of chronic obstruction.

Given the role of NRP2 in this process, the anti-NRP2 antibodies described herein could provide new therapies to modulate smooth muscle contractility, including for example, the treatment of reduced smooth muscle (SM) contractility in the bladder, and more specifically syndromes associated with partial bladder outlet obstruction syndromes.

Certain embodiments thus include methods of treating ameliorating the symptoms of, or inhibiting the progression of partial bladder outlet obstruction syndromes in a subject in need thereof, comprising administering to the subject at least one antibody or antigen-binding fragment thereof that specifically binds to a human NRP2 polypeptide (an anti-NRP2 antibody), and which modulates (e.g., interferes with) binding of the human NRP2 polypeptide to a NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3).

Certain embodiments also include methods of modulating (e.g., increasing, reducing) smooth muscle contractility in a subject in need thereof, comprising administering to the subject an anti-NRP2 antibody, or antigen-binding fragment

thereof, or therapeutic composition provided herein. Certain embodiments include treating, ameliorating the symptoms of, and/or reducing the progression of, reduced smooth muscle contractility in a subject in need thereof, comprising administering to the subject an anti-NRP2 antibody or antigen-binding fragment thereof, or therapeutic composition provided herein.

Methods for identifying subjects with one or more of the diseases or conditions described herein are known in the art. In some embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1) and/or a coding mRNA thereof relative to a healthy control. For instance, in some embodiments, the levels of the at least one NRP2 ligand in the diseases subject, cells, or tissue are about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 1000 or more times the levels of the at least one NRP2 ligand in a healthy control. In some embodiments, the subject has, and/or is selected for treatment based on having, a cancer which has increased levels or expression of at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1) and/or a coding mRNA thereof relative to a non-cancerous control cell or tissue. For instance, in some embodiments, the levels of the at least one NRP2 ligand in the cancer cells or tissue are about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100, 1000 or more times the levels of the NRP2 ligand in a non-cancerous control or standard. Thus, certain embodiments include methods of selecting a subject for treatment, comprising (i) detecting increased expression levels of at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1) and/or coding mRNA in the subject relative to a control or reference, and (ii) administering to the subject a therapeutic composition comprising at least one anti-NRP2-antibody or antigen-binding fragment thereof, as described herein. In particular embodiments, the HRS polypeptide is a splice variant of full-length HRS. In some embodiments, the HRS splice variant is selected from one or more of HisRSN1, HisRSN2, HisRSN3, HisRSN4, HisRSN5, HisRSC1, HisRSC2, HisRSC3, HisRSC4, HisRSC5, HisRSC6, HisRSC7, HisRSC8, and HisRSC9.

In some embodiments, the subject has, and/or is selected for treatment based on having, increased circulating or serum levels of a soluble neuropilin 2 (NRP2) polypeptide (selected, for example, from Table N1), either bound to an HRS polypeptide or free, relative to the levels of a healthy or matched control population of subject(s). For instance, in certain embodiments, the circulating or serum levels are about or at least about 10, 20, 30, 50, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, 1800, 1900, 2000, 3000, 4000, 5000 pM of the soluble NRP2 polypeptide, or the circulating or serum levels are about 30-50, 50-100, 100-2000, 200-2000, 300-2000, 400-2000, 500-2000, 600-2000, 700-2000, 800-2000, 900-2000, 1000-2000, 2000-3000, 3000-4000, 4000-5000 pM of the soluble NRP2 polypeptide.

In certain embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of an NRP2 polypeptide (optionally selected from Table N1) and/or a coding mRNA thereof relative to a healthy control (e.g., an NRP2-associated disease). In some embodiments, the NRP2 polypeptide is an NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table

N1) or a fragment thereof. In some embodiments, the NRP2 polypeptide is an NRP2b isoform (e.g., variants 4 and/or 5 of Table N1) or a fragment thereof.

For example, in certain embodiments, the levels of the NRP2 polypeptide in the diseased subject, cells, or tissue are about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more times the levels of NRP2 polypeptide in a healthy control. In some embodiments, the subject has, and/or is selected for treatment based on having, a cancer which has increased levels or expression of a NRP2 polypeptide (selected, for example, from Table N1) and/or a coding mRNA thereof relative to a control cell or tissue, optionally relative to a non-cancerous cell or tissue of the same type as the cancer. For instance, in some embodiments, the levels of the NRP2 polypeptide in the cancer cells or tissue are about or at least about 1.5, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 30, 40, 50, 60, 70, 80, 90, 100 or more times the levels of NRP2 polypeptide in a non-cancerous control or standard. Some embodiments thus include methods of selecting a subject for treatment, comprising (i) detecting increased expression levels of a NRP2 polypeptide and/or a coding mRNA thereof in the subject relative to a control or reference, and (ii) administering to the subject a therapeutic composition comprising at least one anti-NRP2-antibody or antigen-binding fragment thereof, as described herein.

In some embodiments, the subject has, and/or is selected for treatment based on having, a disease associated with increased levels or expression of NRP2a (e.g., variants 1, 2, and/or 3 of Table N1) and/or NRP2b (e.g., variants 4 and/or 5 of Table N1), or an altered ratio of NRP2a: NRP2b expression, relative to a healthy control or matched control standard or population of subject(s). In some embodiments, the subject has significantly higher expression or levels of NRP2b relative to a healthy control or matched control standard or population of subject(s). In some embodiments, the levels of NRP2b are increased by about or at least about 10%, 20%, 30%, 40%, 50%, 100%, 200%, 300%, 400%, 500%, 600%, 700%, 800%, 900%, 1000% compared to a healthy control or matched control standard or population of subject(s). In some embodiments, the subject has, and/or is selected for treatment based on having, increased circulating levels of HRS: NRP2 complexes relative to a healthy or matched control standard or population of subject(s). Certain embodiments therefore include methods of selecting a subject for cancer treatment, comprising (i) detecting increased expression levels of HRS: NRP2 complexes in the subject relative to a control or reference, and (ii) administering to the subject a therapeutic composition comprising at least one anti-NRP2-antibody or antigen-binding fragment thereof, as described herein.

In some embodiments, the healthy control or matched control standard or population of subject(s) comprises average ranges for age-matched samples of diseased or non-diseased cells or tissue of the same type, which comprise specific characteristics such as drug resistance, metastatic potential, aggressiveness, genetic signature (e.g., p53 mutations, PTEN deletion, IGF1R expression), and/or expression patterns.

For in vivo use, as noted above, for the treatment of human or non-human mammalian disease or testing, the agents described herein are generally incorporated into one or more therapeutic or pharmaceutical compositions prior to administration, including veterinary therapeutic compositions.

Thus, certain embodiments relate to therapeutic compositions that comprise at least one antibody or antigen-

binding fragment thereof that specifically binds to a human NRP2 polypeptide, as described herein. In some instances, a therapeutic or pharmaceutical composition comprises one or more of the agents described herein in combination with a pharmaceutically- or physiologically-acceptable carrier or excipient. Certain therapeutic compositions further comprise at least one cancer immunotherapy agent, as described herein.

Some therapeutic compositions comprise (and certain methods utilize) only one anti-NRP2 antibody or antigen-binding fragment thereof. Certain therapeutic compositions comprise (and certain methods utilize) a mixture of at least two, three, four, or five different anti-NRP2 antibodies or antigen-binding fragments thereof.

For instance, certain therapeutic compositions comprise at least two anti-NRP2 antibodies, including a first antibody or antigen-binding fragment thereof that specifically binds to at least one first epitope of a human NRP2 polypeptide, and a second antibody or antigen-binding fragment thereof that specifically binds to at least one second epitope of a human NRP2 polypeptide, wherein the at least one first epitope differs from the at least one second epitope. In some embodiments, the first and the second antibody or antigen-binding fragment thereof specifically and non-competitively bind to the same domain of the NRP2 polypeptide. In some embodiments, the first anti-NRP2 antibody or antigen-binding fragment thereof binds selectively to a first epitope that is specific to the NRP2a isoform (e.g., variants 1, 2, and/or 3 of Table N1), and the second anti-NRP2 antibody or antigen-binding fragment thereof binds selectively to a second epitope which is specific to the NRP2b isoform (e.g., variants 4 and/or 5 of Table N1).

In some embodiments, the first and the second antibody or antigen-binding fragment thereof specifically and non-competitively bind to different domains of the NRP2 polypeptide.

In some embodiments, the first antibody antagonizes the binding/signaling activity between the NRP2 polypeptide and the at least one NRP2 ligand. In certain embodiments, the second antibody or antigen-binding fragment thereof agonizes or enhances the binding/signaling activity between the NRP2 polypeptide and at least one NRP2 ligand.

In some embodiments, the first and the second antibody or antigen-binding fragments thereof are both blocking antibodies, for example, for at least two different NRP2 ligands. In some embodiments, the first and the second antibody or antigen-binding fragments thereof are both partial-blocking antibodies, for example, for at least two different NRP2 ligands. In some instances, the first and the second antibodies or antigen-binding fragments thereof are both non-blocking antibodies, for example, with respect to at least two different NRP2 ligands.

In some instances, the first antibody or antigen-binding fragment thereof is a blocking antibody and the second antibody or antigen-binding fragment thereof is a partial-blocking antibody. In certain instances, the first antibody or antigen-binding fragment thereof is a blocking antibody and the second antibody or antigen-binding fragment thereof is a non-blocking antibody.

In some embodiments, the first and the second antibodies or antigen-binding fragments thereof both comprise an IgG Fc domain with high effector function in humans, for example, an IgG1 or IgG3 Fc domain. In some embodiments, the first and the second antibodies or antigen-binding fragments thereof comprise an IgG Fc domain with low effector function in humans, for example, an IgG2 or IgG4 Fc domain.

In some instances, the first antibody or antigen-binding fragment thereof comprises an IgG Fc domain with high effector function in humans, for example, an IgG1 or IgG3 Fc domain, and the second antibody or antigen-binding fragment thereof comprises an IgG Fc domain with low effector function in humans, for example, an IgG2 or IgG4 Fc domain.

In particular embodiments, the therapeutic composition comprising the agents such as antibodies or other polypeptide agents (e.g., anti-NRP2 antibodies) is substantially pure on a protein basis or a weight-weight basis, for example, the composition has a purity of at least about 80%, 85%, 90%, 95%, 98%, or 99% on a protein basis or a weight-weight basis.

In some embodiments, the antibodies (e.g., anti-NRP2 antibodies) or other polypeptide agents provided herein do not form aggregates, have a desired solubility, and/or have an immunogenicity profile that is suitable for use in humans, as described herein and known in the art. Thus, in some embodiments, the therapeutic composition comprising a polypeptide agent (for example, an antibody such as an anti-NRP2 antibody) is substantially aggregate-free. For example, certain compositions comprise less than about 10% (on a protein basis) high molecular weight aggregated proteins, or less than about 5% high molecular weight aggregated proteins, or less than about 4% high molecular weight aggregated proteins, or less than about 3% high molecular weight aggregated proteins, or less than about 2% high molecular weight aggregated proteins, or less than about 1% high molecular weight aggregated proteins. Some compositions comprise a polypeptide agent (e.g., an antibody such as an anti-NRP2 antibody) that is at least about 50%, about 60%, about 70%, about 80%, about 90% or about 95% monodisperse with respect to its apparent molecular mass.

In some embodiments, polypeptide agents such as antibodies (e.g., anti-NRP2 antibodies) are concentrated to about or at least about 0.1 mg/ml, 0.2 mg/ml, 0.3 mg/ml, 0.4 mg/ml, 0.5 mg/ml, 0.6, 0.7, 0.8, 0.9, 1 mg/ml, 2 mg/ml, 3 mg/ml, 4 mg/ml, 5 mg/ml, 6 mg/ml, 7 mg/ml, 8 mg/ml, 9 mg/ml, 10 mg/ml, 11, 12, 13, 14 or 15 mg/ml and are formulated for biotherapeutic uses.

To prepare a therapeutic or pharmaceutical composition, an effective or desired amount of one or more agents is mixed with any pharmaceutical carrier(s) or excipient known to those skilled in the art to be suitable for the particular agent and/or mode of administration. A pharmaceutical carrier may be liquid, semi-liquid or solid. Solutions or suspensions used for parenteral, intradermal, intraocular, subcutaneous, direct instillation into the bladder, or topical application may include, for example, a sterile diluent (such as water), saline solution (e.g., phosphate buffered saline; PBS), fixed oil, polyethylene glycol, glycerin, propylene glycol or other synthetic solvent; antimicrobial agents (such as benzyl alcohol and methyl parabens); antioxidants (such as ascorbic acid and sodium bisulfite) and chelating agents (such as ethylenediaminetetraacetic acid (EDTA)); buffers (such as acetates, citrates and phosphates). If administered intravenously (e.g., by IV infusion), suitable carriers include physiological saline or phosphate buffered saline (PBS), and solutions containing thickening and solubilizing agents, such as glucose, polyethylene glycol, polypropylene glycol and mixtures thereof.

Administration of agents described herein, in pure form or in an appropriate therapeutic or pharmaceutical composition, can be carried out via any of the accepted modes of administration of agents for serving similar utilities. The

therapeutic or pharmaceutical compositions can be prepared by combining an agent-containing composition with an appropriate physiologically acceptable carrier, diluent or excipient, and may be formulated into preparations in solid, semi-solid, liquid or gaseous forms, such as tablets, capsules, powders, granules, ointments, solutions, suppositories, injections, inhalants, gels, microspheres, and aerosols. In addition, other pharmaceutically active ingredients (including other small molecules as described elsewhere herein) and/or suitable excipients such as salts, buffers and stabilizers may, but need not, be present within the composition.

Administration may be achieved by a variety of different routes, including oral, parenteral, nasal, intravenous, intraocular, intradermal, intramuscular, subcutaneous, installation into the bladder, or topical. Preferred modes of administration depend upon the nature of the condition to be treated or prevented. Particular embodiments include administration by IV infusion.

Carriers can include, for example, pharmaceutically- or physiologically-acceptable carriers, excipients, or stabilizers that are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically-acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as polysorbate 20 (TWEEN™), polyethylene glycol (PEG), and poloxamers (PLURON-ICS™), and the like.

In some embodiments, one or more agents can be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nanoparticles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences, 16th edition, Oslo, A., Ed., (1980). The particle(s) or liposomes may further comprise other therapeutic or diagnostic agents.

The precise dosage and duration of treatment is a function of the disease being treated and may be determined empirically using known testing protocols or by testing the compositions in model systems known in the art and extrapolating therefrom. Controlled clinical trials may also be performed. Dosages may also vary with the severity of the condition to be alleviated. A pharmaceutical composition is generally formulated and administered to exert a therapeutically useful effect while minimizing undesirable side effects. The composition may be administered one time, or may be divided into a number of smaller doses to be administered at intervals of time. For any particular subject, specific dosage regimens may be adjusted over time according to the individual need.

Typical routes of administering these and related therapeutic or pharmaceutical compositions thus include, without limitation, oral, topical, transdermal, inhalation, parenteral,

sublingual, buccal, ocular, rectal, vaginal, and intranasal. The term parenteral as used herein includes subcutaneous injections, intravenous, instillation into the bladder, intramuscular, intrasternal injection or infusion techniques. Therapeutic or pharmaceutical compositions according to certain embodiments of the present disclosure are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a subject or patient. Compositions that will be administered to a subject or patient may take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a herein described agent in aerosol form may hold a plurality of dosage units. Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see Remington: The Science and Practice of Pharmacy, 20th Edition (Philadelphia College of Pharmacy and Science, 2000). The composition to be administered will typically contain a therapeutically effective amount of an agent described herein, for treatment of a disease or condition of interest.

A therapeutic or pharmaceutical composition may be in the form of a solid or liquid. In one embodiment, the carrier(s) are particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral oil, injectable liquid or an aerosol, which is useful in, for example, inhalatory administration. When intended for oral administration, the pharmaceutical composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid. Certain embodiments include sterile, injectable solutions.

As a solid composition for oral administration, the pharmaceutical composition may be formulated into a powder, granule, gel, compressed tablet, pill, capsule, chewing gum, wafer or the like. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, gum tragacanth or gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin; a flavoring agent such as peppermint, methyl salicylate or orange flavoring; and a coloring agent. When the pharmaceutical composition is in the form of a capsule, for example, a gelatin capsule, it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol or oil.

The therapeutic or pharmaceutical composition may be in the form of a liquid, for example, an elixir, syrup, solution, gel, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration, preferred compositions contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

The liquid therapeutic or pharmaceutical compositions, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium

chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium, polyethylene glycols, glycerin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid therapeutic or pharmaceutical composition intended for either parenteral, intraocular, or oral administration should contain an amount of an agent such that a suitable dosage will be obtained. Typically, this amount is at least 0.01% of the agent of interest in the composition. When intended for oral administration, this amount may be varied to be between 0.1 and about 70% of the weight of the composition. Certain oral therapeutic or pharmaceutical compositions contain between about 4% and about 75% of the agent of interest. In certain embodiments, therapeutic or pharmaceutical compositions and preparations are prepared so that a parenteral dosage unit contains between 0.01 to 10% by weight of the agent of interest prior to dilution.

The therapeutic or pharmaceutical compositions may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, bee wax, mineral oil, diluents such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a therapeutic or pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device.

The therapeutic or pharmaceutical compositions may be intended for rectal administration, in the form, for example, of a suppository, which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter, and polyethylene glycol.

The therapeutic or pharmaceutical composition may include various materials, which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials that form the coating shell are typically inert, and may be selected from, for example, sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule. The therapeutic or pharmaceutical compositions in solid or liquid form may include a component that binds to agent and thereby assists in the delivery of the compound. Suitable components that may act in this capacity include monoclonal or polyclonal antibodies, one or more proteins or a liposome.

The therapeutic or pharmaceutical composition may consist essentially of dosage units that can be administered as an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or by a suitable pump system that dispenses the active ingredients. Aerosols may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol

includes the necessary container, activators, valves, subcontainers, and the like, which together may form a kit. One of ordinary skill in the art, without undue experimentation may determine preferred aerosols.

The compositions described herein may be prepared with carriers that protect the agents against rapid elimination from the body, such as time release formulations or coatings. Such carriers include controlled release formulations, such as, but not limited to, implants and microencapsulated delivery systems, and biodegradable, biocompatible polymers, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, polyorthoesters, polylactic acid and others known to those of ordinary skill in the art.

The therapeutic or pharmaceutical compositions may be prepared by methodology well known in the pharmaceutical art. For example, a therapeutic or pharmaceutical composition intended to be administered by injection may comprise one or more of salts, buffers and/or stabilizers, with sterile, distilled water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the agent so as to facilitate dissolution or homogeneous suspension of the agent in the aqueous delivery system.

The therapeutic or pharmaceutical compositions may be administered in a therapeutically effective amount, which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the subject; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disorder or condition; and the subject undergoing therapy. In some instances, a therapeutically effective daily dose is (for a 70 kg mammal) from about 0.001 mg/kg (i.e., ~ 0.07 mg) to about 100 mg/kg (i.e., ~ 7.0 g); preferably a therapeutically effective dose is (for a 70 kg mammal) from about 0.01 mg/kg (i.e., ~ 0.7 mg) to about 50 mg/kg (i.e., ~ 3.5 g); more preferably a therapeutically effective dose is (for a 70 kg mammal) from about 1 mg/kg (i.e., ~ 70 mg) to about 25 mg/kg (i.e., ~ 1.75 g). In some embodiments, the therapeutically effective dose is administered on a weekly, bi-weekly, or monthly basis. In specific embodiments, the therapeutically effective dose is administered on a weekly, bi-weekly, or monthly basis, for example, at a dose of about 1-10 or 1-5 mg/kg, or about 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10 mg/kg.

The combination therapies described herein may include administration of a single pharmaceutical dosage formulation, which contains an anti-NRP2 antibody and an additional therapeutic agent (e.g., immunotherapy agent, chemotherapeutic agent, hormonal therapeutic agent, kinase inhibitor), as well as administration of compositions comprising an anti-NRP2 antibody and an additional therapeutic agent in its own separate pharmaceutical dosage formulation. For example, an anti-NRP2 antibody as described herein and additional therapeutic agent can be administered to the subject together in a single oral dosage composition such as a tablet or capsule, or each agent administered in separate oral dosage formulations. Similarly, an anti-NRP2 antibody as described herein and additional therapeutic agent can be administered to the subject together in a single parenteral dosage composition such as in a saline solution or other physiologically acceptable solution, or each agent administered in separate parenteral dosage formulations. As another example, for cell-based therapies, an anti-NRP2 antibody can be mixed with the cells prior to administration, administered as part of a separate composition, or both.

Where separate dosage formulations are used, the compositions can be administered at essentially the same time, i.e., concurrently, or at separately staggered times, i.e., sequentially and in any order; combination therapy is understood to include all these regimens.

Also included are patient care kits, comprising (a) at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin 2 (NRP2) polypeptide (an anti-NRP2 antibody), as described herein; and optionally (b) at least one additional therapeutic agent (e.g., immunotherapy agent, chemotherapeutic agent, hormonal therapeutic agent, kinase inhibitor). In certain kits, (a) and (b) are in separate therapeutic compositions. In some kits, (a) and (b) are in the same therapeutic composition.

The kits herein may also include a one or more additional therapeutic agents or other components suitable or desired for the indication being treated, or for the desired diagnostic application. The kits herein can also include one or more syringes or other components necessary or desired to facilitate an intended mode of delivery (e.g., stents, implantable depots, etc.).

In some embodiments, a patient care kit contains separate containers, dividers, or compartments for the composition(s) and informational material(s). For example, the composition(s) can be contained in a bottle, vial, or syringe, and the informational material(s) can be contained in association with the container. In some embodiments, the separate elements of the kit are contained within a single, undivided container. For example, the composition is contained in a bottle, vial or syringe that has attached thereto the informational material in the form of a label. In some embodiments, the kit includes a plurality (e.g., a pack) of individual containers, each containing one or more unit dosage forms (e.g., a dosage form described herein) of an anti-NRP2 antibody and optionally at least one additional therapeutic agent. For example, the kit includes a plurality of syringes, ampules, foil packets, or blister packs, each containing a single unit dose of an anti-NRP2 antibody and optionally at least one additional therapeutic agent. The containers of the kits can be air tight, waterproof (e.g., impermeable to changes in moisture or evaporation), and/or light-tight.

The patient care kit optionally includes a device suitable for administration of the composition, e.g., a syringe, inhalant, dropper (e.g., eye dropper), swab (e.g., a cotton swab or wooden swab), or any such delivery device. In some embodiments, the device is an implantable device that dispenses metered doses of the agent(s). Also included are methods of providing a kit, e.g., by combining the components described herein.

Bioassays and Analytical Assays for Drug Release Assays and Product Specifications, Diagnostics, and Reagents

Also included are bioassays that relate to anti-NRP2 antibodies and related agents such as therapeutic and diagnostic reagents. Examples include bioassays and analytical assays that measure purity, biological activity, affinity, solubility, pH, endotoxin levels, among others, many of which are described herein. Also included are assays that establish dose response curves and/or provide one or more bases for comparison between different batches of antibody. Batch comparisons can be based on any one or more of chemical characterization, biological characterization, and clinical characterization. Also included are methods of evaluating the potency, stability, pharmacokinetics, and immunogenicity of a selected antibody. Among other uses, these and other

methods can be used for lot releasing testing of biologic or chemical agents, including anti-NRP2 antibodies, described herein.

Certain embodiments include the use of bioaffinity assays. Such assays can be used to assess the binding affinity, for example, between an anti-NRP2 antibody and at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1), including its ability to interfere with the interaction between a human NRP2 polypeptide and the at least one NRP2 ligand, or other cellular binding partner. Certain exemplary binding affinity assays may utilize ELISA assays, and other immunoassays as described herein and known in the art. Certain assays utilize high-performance receptor binding chromatography (see, e.g., Roswall et al., *Biologicals* 24:25-39, 1996). Other exemplary binding affinity assays may utilize surface plasmon resonance (SPR)-based technologies. Examples include BIAcore technologies, certain of which integrate SPR technology with a microfluidics system to monitor molecular interactions in real time at concentrations ranging from pM to mM. Also included are KINEXA™ assays, which provide accurate measurements of binding specificity, binding affinity, and binding kinetics/rate constants.

Certain embodiments relate to immunoassays for evaluating or optimizing the immunogenicity of anti-NRP2 antibodies. Examples include ex vivo human cellular assays and in vitro immuno-enzymatic assays to provide useful information on the immunogenic potential of a therapeutic protein. Ex vivo cell-response assays can be used, for example, to reproduce the cellular co-operation between antigen-presenting cells (APCs) and T-cells, and thereby measure T-cells activation after contact with a protein of interest. Certain in vitro enzymatic assays may utilize a collection of recombinant HLA-DR molecules that cover a significant portion of a relevant human population, and may include automated immuno-enzymatic assays for testing the binding of peptides (stemming from the fragmentation of the therapeutic protein) with the HLA-DR molecules. Also included are methods of reducing the immunogenicity of a selected protein, such as by using these and related methods to identify and then remove or alter one or more T-cell epitopes from an anti-NRP2 antibody.

Also included are biological release assays (e.g., cell-based assays) for measuring parameters such as specific biological activities, including non-canonical biological activities, and cytotoxicity. Certain specific biological assays include, for example, cell-based assays that utilize a cellular binding partner (e.g., cell-surface receptor (for example a NRP2 polypeptide and/or at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3) presented on the cell surface), which is either endogenously, or recombinantly expressed on the cell surface), which is functionally coupled to a readout, such as a fluorescent or luminescent indicator of NRP2 or NRP2 ligand binding, or functional activity, as described herein.

For instance, specific embodiments include a cell that either endogenously or recombinantly expresses a human NRP2 polypeptide on the cell surface, which allows assessment of the ability of anti-NRP2 antibody to bind NRP2. In some embodiments, the anti-NRP2 antibody and/or the NRP2 polypeptide is/are functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator to measure the binding and/or biological activity of the NRP2 polypeptide. In some embodiments, the cell also expresses at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table

H1), wherein the at least one NRP2 ligand is coupled to a readout or indicator, such as a fluorescent or luminescent indicator of binding and/or biological activity of the at least one NRP2 ligand.

Also included are in vivo biological assays to characterize the pharmacokinetics of an anti-NRP2 antibody, typically utilizing engineered, or wild type mice, rat, monkey or other mammal (see, e.g., Lee et al., *The Journal of Pharmacology* 281:1431-1439, 1997). Examples of cytotoxicity-based biological assays include release assays (e.g., chromium or europium release assays to measure apoptosis; see, e.g., von Zons et al., *Clin Diagn Lab Immunol* 4:202-207, 1997), among others, which can assess the cytotoxicity anti-NRP2 antibodies, whether for establishing dose response curves, batch testing, or other properties related to approval by various regulatory agencies, such as the Food and Drug Administration (FDA).

Also included are assays for evaluating the effects of an anti-NRP2 antibody on immune cells. Examples include an assay system, comprising an activated population of T-cells and at least one anti-NRP2 antibody, wherein the at least one anti-NRP2 antibody interferes with binding of NRP2 to at least one NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1).

Certain embodiments include an assay system, comprising a single monoclonal anti-NRP2 antibody and at least one human NRP2 polypeptide, wherein the anti-NRP2 antibody binds to the NRP2 polypeptide. In some instances, the at least one antibody comprises an IgG4 Fc domain.

Also included are testing material(s), comprising a purified NRP2 polypeptide, wherein said purified NRP2 polypeptide is bound to a solid substrate in a manner that enables antibody binding detection.

Such assays and materials can be used, for example, to develop a dose response curve for a selected anti-NRP2 antibody, and/or to compare the dose response curve of different batches of proteins or other agents. A dose-response curve is an X-Y graph that relates the magnitude of a stressor to the response of a receptor, such as an NRP2-NRP2 ligand (for example, an NRP2 ligand from Table N2 or Table N3 and/or an HRS polypeptide from Table H1) interaction; the response may be a physiological or biochemical response, such as a non-canonical biological activity in a cell in vitro or in a cell or tissue in vivo, a therapeutically effective amount as measured in vivo (e.g., as measured by EC₅₀), or death, whether measured in vitro or in vivo (e.g., cell death, organismal death). Death is usually indicated as an LD₅₀, a statistically-derived dose that is lethal to 50% of a modeled population, though it can be indicated by LC₀₁ (lethal dose for 1% of the animal test population), LC100 (lethal dose for 100% of the animal test population), or LC_{LO} (lowest dose causing lethality). Almost any desired effect or endpoint can be characterized in this manner.

The measured dose of a response curve is typically plotted on the X axis and the response is plotted on the Y axis. More typically, the logarithm of the dose is plotted on the X axis, most often generating a sigmoidal curve with the steepest portion in the middle. The No Observable Effect Level (NOEL) refers to the lowest experimental dose for which no measurable effect is observed, and the threshold dose refers to the first point along the graph that indicates a response above zero. As a general rule, stronger drugs generate steeper dose response curves. For many drugs, the desired effects are found at doses slightly greater than the threshold dose, often because lower doses are relatively ineffective and higher doses lead to undesired side effects. For in vivo

generated dose response curves, a curve can be characterized by values such as $\mu\text{g/kg}$, mg/kg , or g/kg of body-weight, if desired.

For batch comparisons, it can be useful to calculate the coefficient of variation (CV) between different dose response curves of different batches (e.g., between different batches of anti-NRP2 antibody), in part because the CV allows comparison between data sets with different units or different means. For instance, in certain exemplary embodiments, two or three or more different batches of anti-NRP2 antibodies or other agents have a CV between them of less than about 30%, 20%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, or 1% for a 4, 5, 6, 7, or 8 point dose curve. In certain embodiments, the dose response curve is measured in a cell-based assay, and its readout relates to an increase or a decrease in a selected activity of an anti-NRP2 antibody. In certain embodiments, the dose response curve is measured in a cell release assay or animal model (e.g., mouse model), and its readout relates to cell death or animal death. Other variations will be apparent to persons skilled in the art.

Expression and Purification Systems

Certain embodiments include methods and related compositions for expressing and purifying an anti-NRP2 antibody or other polypeptide-based agent described herein. Such recombinant anti-NRP2 antibodies can be conveniently prepared using standard protocols as described for example in Sambrook, et al., (1989, supra), in particular Sections 16 and 17; Ausubel et al., (1994, supra), in particular Chapters 10 and 16; and Coligan et al., Current Protocols in Protein Science (John Wiley & Sons, Inc. 1995-1997), in particular Chapters 1, 5 and 6. As one general example, anti-NRP2 antibodies may be prepared by a procedure including one or more of the steps of: (a) preparing a construct comprising a polynucleotide sequences that encode an anti-NRP2 antibody heavy and light chain and that are operably linked to a regulatory element; (b) introducing the constructs into a host cell; (c) culturing the host cell to express an anti-NRP2 antibody; and (d) isolating an anti-NRP2 antibody from the host cell.

Anti-NRP2 antibody polynucleotides are described elsewhere herein. In order to express a desired polypeptide, a nucleotide sequence encoding an anti-NRP2 antibody, or a functional equivalent, may be inserted into appropriate expression vector, i.e., a vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding a polypeptide of interest and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described in Sambrook et al., Molecular Cloning, A Laboratory Manual (1989), and Ausubel et al., Current Protocols in Molecular Biology (1989).

A variety of expression vector/host systems are known and may be utilized to contain and express polynucleotide sequences. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with virus expression vectors (e.g., baculovirus); plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors

(e.g., Ti or pBR322 plasmids); or animal cell systems, including mammalian cell and more specifically human cell systems.

The “control elements” or “regulatory sequences” present in an expression vector are those non-translated regions of the vector—enhancers, promoters, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements may vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, may be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the PBLUESCRIPT phagemid (Stratagene, La Jolla, Calif.) or PSORT1 plasmid (Gibco BRL, Gaithersburg, Md.) and the like may be used. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are generally preferred. If it is necessary to generate a cell line that contains multiple copies of the sequence encoding a polypeptide, vectors based on SV40 or EBV may be advantageously used with an appropriate selectable marker.

In bacterial systems, a number of expression vectors may be selected depending upon the use intended for the expressed polypeptide. For example, when large quantities are needed, vectors which direct high level expression of fusion proteins that are readily purified may be used. Such vectors include, but are not limited to, the multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene), in which the sequence encoding the polypeptide of interest may be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced; pIN vectors (Van Heeke & Schuster, J. Biol. Chem. 264:5503-5509 (1989)); and the like. pGEX Vectors (Promega, Madison, Wis.) may also be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems may be designed to include heparin, thrombin, or factor XA protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

Certain embodiments may employ *E. coli*-based expression systems (see, e.g., Structural Genomics Consortium et al., Nature Methods. 5:135-146, 2008). These and related embodiments may rely partially or totally on ligation-independent cloning (LIC) to produce a suitable expression vector. In specific embodiments, protein expression may be controlled by a T7 RNA polymerase (e.g., pET vector series). These and related embodiments may utilize the expression host strain BL21 (DE3), a λ DE3 lysogen of BL21 that supports T7-mediated expression and is deficient in *lon* and *ompT* proteases for improved target protein stability. Also included are expression host strains carrying plasmids encoding tRNAs rarely used in *E. coli*, such as ROSETTA™ (DE3) and Rosetta 2 (DE3) strains. Cell lysis and sample handling may also be improved using reagents sold under the trademarks BENZONASE® nuclease and BUGBUSTER® Protein Extraction Reagent. For cell culture, auto-inducing media can improve the efficiency of many expression systems, including high-throughput expression systems. Media of this type (e.g., OVERNIGHT EXPRESS™ Autoinduction System) gradually elicit protein expression through metabolic shift without the addition of

artificial inducing agents such as IPTG. Particular embodiments employ hexahistidine tags (such as those sold under the trademark HIS. TAG® fusions), followed by immobilized metal affinity chromatography (IMAC) purification, or related techniques. In certain aspects, however, clinical grade proteins can be isolated from *E. coli* inclusion bodies, without or without the use of affinity tags (see, e.g., Shimp et al., Protein Expr Purif. 50:58-67, 2006). As a further example, certain embodiments may employ a cold-shock induced *E. coli* high-yield production system, because over-expression of proteins in *Escherichia coli* at low temperature improves their solubility and stability (see, e.g., Qing et al., Nature Biotechnology. 22:877-882, 2004).

Also included are high-density bacterial fermentation systems. For example, high cell density cultivation of *Ralstonia eutropha* allows protein production at cell densities of over 150 g/L, and the expression of recombinant proteins at titers exceeding 10 g/L.

In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH may be used. For reviews, see Ausubel et al. (supra) and Grant et al., Methods Enzymol. 153:516-544 (1987). Also included are *Pichia* pandoris expression systems (see, e.g., Li et al., Nature Biotechnology. 24, 210-215, 2006; and Hamilton et al., Science, 301:1244, 2003). Certain embodiments include yeast systems that are engineered to selectively glycosylate proteins, including yeast that have humanized N-glycosylation pathways, among others (see, e.g., Hamilton et al., Science. 313:1441-1443, 2006; Wildt et al., Nature Reviews Microbiol. 3:119-28, 2005; and Gerngross et al., Nature-Biotechnology. 22:1409-1414, 2004; U.S. Pat. Nos. 7,629, 163; 7,326,681; and 7,029,872). Merely by way of example, recombinant yeast cultures can be grown in Fernbach Flasks or 15L, 50L, 100L, and 200L fermentors, among others.

In cases where plant expression vectors are used, the expression of sequences encoding polypeptides may be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV may be used alone or in combination with the omega leader sequence from TMV (Takamatsu, EMBO J. 6:307-311 (1987)). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used (Coruzzi et al., EMBO J. 3:1671-1680 (1984); Broglie et al., Science 224:838-843 (1984); and Winter et al., Results Probl. Cell Differ. 17:85-105 (1991)). These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (see, e.g., Hobbs in McGraw Hill, Yearbook of Science and Technology, pp. 191-196 (1992)).

An insect system may also be used to express a polypeptide of interest. For example, in one such system, *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* cells. The sequences encoding the polypeptide may be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of the polypeptide-encoding sequence will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses may then be used to infect, for example, *S. frugiperda* cells or *Trichoplusia* cells in which the polypeptide of interest may be expressed (Engelhard et al., Proc. Natl. Acad. Sci. U.S.A. 91:3224-3227 (1994)). Also included are baculovirus expression systems, including those that utilize SF9, SF21,

and *T. ni* cells (see, e.g., Murphy and Piwnica-Worms, Curr Protoc Protein Sci. Chapter 5: Unit5.4, 2001). Insect systems can provide post-translation modifications that are similar to mammalian systems.

In mammalian host cells, a number of viral-based expression systems are generally available. For example, in cases where an adenovirus is used as an expression vector, sequences encoding a polypeptide of interest may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain a viable virus which is capable of expressing the polypeptide in infected host cells (Logan & Shenk, Proc. Natl. Acad. Sci. U.S.A. 81:3655-3659 (1984)). In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells.

Examples of useful mammalian host cell lines include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells sub-cloned for growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TR1 cells (Mather et al., Annals N.Y. Acad. Sci. 383:44-68 (1982)); MRC 5 cells; FS4 cells; and a human hepatoma line (Hep G2). Other useful mammalian host cell lines include Chinese hamster ovary (CHO) cells, including DHFR-CHO cells (Urlaub et al., PNAS USA 77:4216 (1980)); and myeloma cell lines such as NSO and Sp2/0. For a review of certain mammalian host cell lines suitable for antibody production, see, e.g., Yazaki and Wu, Methods in Molecular Biology, Vol. 248 (B. K. C Lo, ed., Humana Press, Totowa, N.J., 2003), pp. 255-268. Certain preferred mammalian cell expression systems include CHO and HEK293-cell based expression systems. Mammalian expression systems can utilize attached cell lines, for example, in T-flasks, roller bottles, or cell factories, or suspension cultures, for example, in 1 L and 5 L spinners, 5 L, 14 L, 40 L, 100 L and 200 L stir tank bioreactors, or 20/50 L and 100/200 L WAVE bioreactors, among others known in the art.

Also included is the cell-free expression of proteins. These and related embodiments typically utilize purified RNA polymerase, ribosomes, tRNA and ribonucleotides; these reagents may be produced by extraction from cells or from a cell-based expression system.

Specific initiation signals may also be used to achieve more efficient translation of sequences encoding a polypeptide of interest. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding the polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a portion thereof, is inserted, exogenous translational control signals including the ATG initiation codon should be provided. Furthermore, the initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons may be of various origins, both natural and

synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used, such as those described in the literature (Scharf. et al., *Results Probl. Cell Differ.* 20:125-162 (1994)).

In addition, a host cell strain may be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, post-translational modifications such as acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to facilitate correct insertion, folding and/or function. Different host cells such as yeast, CHO, HeLa, MDCK, HEK293, and W138, in addition to bacterial cells, which have or even lack specific cellular machinery and characteristic mechanisms for such post-translational activities, may be chosen to ensure the correct modification and processing of the foreign protein.

For long-term, high-yield production of recombinant proteins, stable expression is generally preferred. For example, cell lines which stably express a polynucleotide of interest may be transformed using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1-2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be proliferated using tissue culture techniques appropriate to the cell type. Transient production, such as by transient transfection or infection, can also be employed. Exemplary mammalian expression systems that are suitable for transient production include HEK293 and CHO-based systems.

Any number of selection systems may be used to recover transformed or transduced cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11:223-232 (1977)) and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22:817-823 (1990)) genes which can be employed in tk- or aprt-cells, respectively. Also, antimetabolite, antibiotic or herbicide resistance can be used as the basis for selection; for example, dhfr which confers resistance to methotrexate (Wigler et al., *PNAS USA* 77:3567-70 (1980)); npt, which confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin et al., *J. Mol. Biol.* 150:1-14 (1981)); and als or pat, which confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murry, *supra*). Additional selectable genes have been described, for example, trpB, which allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci. U.S.A.* 85:8047-51 (1988)). The use of visible markers has gained popularity with such markers as green fluorescent protein (GFP) and other fluorescent proteins (e.g., RFP, YFP), anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, being widely used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system (see, e.g., Rhodes et al., *Methods Mol. Biol.* 55:121-131 (1995)).

Also included are high-throughput protein production systems, or micro-production systems. Certain aspects may

utilize, for example, hexa-histidine fusion tags for protein expression and purification on metal chelate-modified slide surfaces or MagneHis Ni-Particles (see, e.g., Kwon et al., *BMC Biotechnol.* 9:72, 2009; and Lin et al., *Methods Mol Biol.* 498:129-41, 2009)). Also included are high-throughput cell-free protein expression systems (see, e.g., Sitaraman et al., *Methods Mol Biol.* 498:229-44, 2009). These and related embodiments can be used, for example, to generate microarrays of anti-NRP2 antibodies which can then be used for screening libraries to identify antibodies and antigen-binding domains that interact with the NRP2 polypeptide(s) of interest.

A variety of protocols for detecting and measuring the expression of polynucleotide-encoded products, using binding agents or antibodies such as polyclonal or monoclonal antibodies specific for the product, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), western immunoblots, radioimmunoassays (RIA), and fluorescence activated cell sorting (FACS). These and other assays are described, among other places, in Hampton et al., *Serological Methods, a Laboratory Manual* (1990) and Maddox et al., *J. Exp. Med.* 158:1211-1216 (1983).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides include oligolabeling, nick translation, end-labeling or PCR amplification using a labeled nucleotide. Alternatively, the sequences, or any portions thereof may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits. Suitable reporter molecules or labels, which may be used include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with a polynucleotide sequence of interest may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. Certain specific embodiments utilize serum free cell expression systems. Examples include HEK293 cells and CHO cells that can grown on serum free medium (see, e.g., Rosser et al., *Protein Expr. Purif.* 40:237-43, 2005; and U.S. Pat. No. 6,210,922).

An antibody, or antigen-binding fragment thereof, produced by a recombinant cell may be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides may be designed to contain signal sequences which direct secretion of the encoded polypeptide through a prokaryotic or eukaryotic cell membrane. Other recombinant constructions may be used to join sequences encoding a polypeptide of interest to nucleotide sequence encoding a polypeptide domain which will facilitate purification and/or detection of soluble proteins. Examples of such domains include cleavable and non-cleavable affinity purification and epitope tags such as avidin, FLAG tags, poly-histidine tags (e.g., 6 $\times$ His), cMyc tags, V5-tags, glutathione S-transferase (GST) tags, and others.

The protein produced by a recombinant cell can be purified and characterized according to a variety of techniques known in the art. Exemplary systems for performing

protein purification and analyzing protein purity include fast protein liquid chromatography (FPLC) (e.g., AKTA and Bio-Rad FPLC systems), high-pressure liquid chromatography (HPLC) (e.g., Beckman and Waters HPLC). Exemplary chemistries for purification include ion exchange chromatography (e.g., Q, S), size exclusion chromatography, salt gradients, affinity purification (e.g., Ni, Co, FLAG, maltose, glutathione, protein A/G), gel filtration, reverse-phase, ceramic HYPERD[®] ion exchange chromatography, and hydrophobic interaction columns (HIC), among others known in the art. Also included are analytical methods such as SDS-PAGE (e.g., coomassie, silver stain), immunoblot, Bradford, and ELISA, which may be utilized during any step of the production or purification process, typically to measure the purity of the protein composition.

Also included are methods of concentrating anti-NRP2 antibodies and antigen-binding fragments thereof, and composition comprising concentrated soluble proteins. In different aspects such concentrated solutions of anti-NRP2 antibodies may comprise proteins at a concentration of about 5 mg/ml; or about 8 mg/ml; or about 10 mg/ml; about 15 mg/ml; or about 20 mg/mL.

In some aspects, such compositions may be substantially monodisperse, meaning that an at least one anti-NRP2 antibody exists primarily (i.e., at least about 90%, or greater) in one apparent molecular weight form when assessed for example, by size exclusion chromatography, dynamic light scattering, or analytical ultracentrifugation.

In some aspects, such compositions have a purity (on a protein basis) of at least about 90%, or in some aspects at least about 95% purity, or in some embodiments, at least 98% purity. Purity may be determined via any routine analytical method as known in the art.

In some aspects, such compositions have a high molecular weight aggregate content of less than about 10%, compared to the total amount of protein present, or in some embodiments such compositions have a high molecular weight aggregate content of less than about 5%, or in some aspects such compositions have a high molecular weight aggregate content of less than about 3%, or in some embodiments a high molecular weight aggregate content of less than about 1%. High molecular weight aggregate content may be determined via a variety of analytical techniques including for example, by size exclusion chromatography, dynamic light scattering, or analytical ultracentrifugation.

Examples of concentration approaches contemplated herein include lyophilization, which is typically employed when the solution contains few soluble components other than the protein of interest. Lyophilization is often performed after HPLC run, and can remove most or all volatile components from the mixture. Also included are ultrafiltration techniques, which typically employ one or more selective permeable membranes to concentrate a protein solution. The membrane allows water and small molecules to pass through and retains the protein; the solution can be forced against the membrane by mechanical pump, gas pressure, or centrifugation, among other techniques.

In certain embodiments, the reagents, anti-NRP2 antibodies, or related agents have a purity of at least about 90%, as measured according to routine techniques in the art. In certain embodiments, such as diagnostic compositions or certain therapeutic compositions, an anti-NRP2 antibody composition has a purity of at least about 95%. In specific embodiments, such as therapeutic or pharmaceutical compositions, an anti-NRP2 antibody composition has a purity of at least about 97% or 98% or 99%. In other embodiments, such as when being used as reference or research reagents,

anti-NRP2 antibodies can be of lesser purity, and may have a purity of at least about 50%, 60%, 70%, or 80%. Purity can be measured overall or in relation to selected components, such as other proteins, e.g., purity on a protein basis.

Purified anti-NRP2 antibodies can also be characterized according to their biological characteristics. Binding affinity and binding kinetics can be measured according to a variety of techniques known in the art, such as Biacore[®] and related technologies that utilize surface plasmon resonance (SPR), an optical phenomenon that enables detection of unlabeled interactants in real time. SPR-based biosensors can be used in determination of active concentration, screening and characterization in terms of both affinity and kinetics. The presence or levels of one or more canonical or non-canonical biological activities can be measured according to cell-based assays, including those that utilize a cellular binding partner of a selected anti-NRP2 antibody, which is functionally coupled to a readout or indicator, such as a fluorescent or luminescent indicator of biological activity, as described herein.

In certain embodiments, as noted above, an anti-NRP2 antibody composition is substantially endotoxin free, including, for example, about 95% endotoxin free, preferably about 99% endotoxin free, and more preferably about 99.99% endotoxin free. The presence of endotoxins can be detected according to routine techniques in the art, as described herein. In specific embodiments, an anti-NRP2 antibody composition is made from a eukaryotic cell such as a mammalian or human cell in substantially serum free media. In certain embodiments, as noted herein, an anti-NRP2 antibody composition has an endotoxin content of less than about 10 EU/mg of anti-NRP2 antibody, or less than about 5 EU/mg of anti-NRP2 antibody, less than about 3 EU/mg of anti-NRP2 antibody, or less than about 1 EU/mg of anti-NRP2 antibody.

In certain embodiments, an anti-NRP2 antibody composition comprises less than about 10% wt/wt high molecular weight aggregates, or less than about 5% wt/wt high molecular weight aggregates, or less than about 2% wt/wt high molecular weight aggregates, or less than about or less than about 1% wt/wt high molecular weight aggregates.

Also included are protein-based analytical assays and methods, which can be used to assess, for example, protein purity, size, solubility, and degree of aggregation, among other characteristics. Protein purity can be assessed a number of ways. For instance, purity can be assessed based on primary structure, higher order structure, size, charge, hydrophobicity, and glycosylation. Examples of methods for assessing primary structure include N- and C-terminal sequencing and peptide-mapping (see, e.g., Allen et al., *Biologicals*. 24:255-275, 1996)). Examples of methods for assessing higher order structure include circular dichroism (see, e.g., Kelly et al., *Biochim Biophys Acta*. 1751:119-139, 2005), fluorescent spectroscopy (see, e.g., Meagher et al., *J. Biol. Chem.* 273:23283-89, 1998), FT-IR, amide hydrogen-deuterium exchange kinetics, differential scanning calorimetry, NMR spectroscopy, immunoreactivity with conformationally sensitive antibodies. Higher order structure can also be assessed as a function of a variety of parameters such as pH, temperature, or added salts. Examples of methods for assessing protein characteristics such as size include analytical ultracentrifugation and size exclusion HPLC (SEC-HPLC), and exemplary methods for measuring charge include ion-exchange chromatography and isoelectric focusing. Hydrophobicity can be assessed, for example, by reverse-phase HPLC and hydrophobic interaction chromatography HPLC. Glycosylation can affect pharmacokinetics

(e.g., clearance), conformation or stability, receptor binding, and protein function, and can be assessed, for example, by mass spectrometry and nuclear magnetic resonance (NMR) spectroscopy.

As noted above, certain embodiments include the use of SEC-HPLC to assess protein characteristics such as purity, size (e.g., size homogeneity) or degree of aggregation, and/or to purify proteins, among other uses. SEC, also including gel-filtration chromatography (GFC) and gel-permeation chromatography (GPC), refers to a chromatographic method in which molecules in solution are separated in a porous material based on their size, or more specifically their hydrodynamic volume, diffusion coefficient, and/or surface properties. The process is generally used to separate biological molecules, and to determine molecular weights and molecular weight distributions of polymers. Typically, a biological or protein sample (such as a protein extract produced according to the protein expression methods provided herein and known in the art) is loaded into a selected size-exclusion column with a defined stationary phase (the porous material), preferably a phase that does not interact with the proteins in the sample. In certain aspects, the stationary phase is composed of inert particles packed into a dense three-dimensional matrix within a glass or steel column. The mobile phase can be pure water, an aqueous buffer, an organic solvent, or a mixture thereof. The stationary-phase particles typically have small pores and/or channels which only allow molecules below a certain size to enter. Large particles are therefore excluded from these pores and channels, and their limited interaction with the stationary phase leads them to elute as a "totally-excluded" peak at the beginning of the experiment. Smaller molecules, which can fit into the pores, are removed from the flowing mobile phase, and the time they spend immobilized in the stationary-phase pores depends, in part, on how far into the pores they penetrate. Their removal from the mobile phase flow causes them to take longer to elute from the column and results in a separation between the particles based on differences in their size. A given size exclusion column has a range of molecular weights that can be separated. Overall, molecules larger than the upper limit will not be trapped by the stationary phase, molecules smaller than the lower limit will completely enter the solid phase and elute as a single band, and molecules within the range will elute at different rates, defined by their properties such as hydrodynamic volume. For examples of these methods in practice with pharmaceutical proteins, see Bruner et al., *Journal of Pharmaceutical and Biomedical Analysis*. 15:1929-1935, 1997.

Protein purity for clinical applications is also discussed, for example, by Anicetti et al. (*Trends in Biotechnology*. 7:342-349, 1989). More recent techniques for analyzing protein purity include, without limitation, the LabChip GXII, an automated platform for rapid analysis of proteins and nucleic acids, which provides high throughput analysis of titer, sizing, and purity analysis of proteins. In certain non-limiting embodiments, clinical grade proteins such as protein fragments and antibodies can be obtained by utilizing a combination of chromatographic materials in at least two orthogonal steps, among other methods (see, e.g., *Therapeutic Proteins: Methods and Protocols*. Vol. 308, Eds., Smales and James, Humana Press Inc., 2005). Typically, protein agents (e.g., anti-NRP2 antibodies, and antigen-binding fragments) are substantially endotoxin-free, as measured according to techniques known in the art and described herein.

Protein solubility assays are also included. Such assays can be utilized, for example, to determine optimal growth

and purification conditions for recombinant production, to optimize the choice of buffer(s), and to optimize the choice of anti-NRP2 antibodies or variants thereof. Solubility or aggregation can be evaluated according to a variety of parameters, including temperature, pH, salts, and the presence or absence of other additives. Examples of solubility screening assays include, without limitation, microplate-based methods of measuring protein solubility using turbidity or other measure as an end point, high-throughput assays for analysis of the solubility of purified recombinant proteins (see, e.g., Stenvall et al., *Biochim Biophys Acta*. 1752:6-10, 2005), assays that use structural complementation of a genetic marker protein to monitor and measure protein folding and solubility in vivo (see, e.g., Wigley et al., *Nature Biotechnology*. 19:131-136, 2001), and electrochemical screening of recombinant protein solubility in *Escherichia coli* using scanning electrochemical microscopy (SECM) (see, e.g., Nagamine et al., *Biotechnology and Bioengineering*. 96:1008-1013, 2006), among others. Anti-NRP2 antibodies with increased solubility (or reduced aggregation) can be identified or selected for according to routine techniques in the art, including simple in vivo assays for protein solubility (see, e.g., Maxwell et al., *Protein Sci*. 8:1908-11, 1999).

Protein solubility and aggregation can also be measured by dynamic light scattering techniques. Aggregation is a general term that encompasses several types of interactions or characteristics, including soluble/insoluble, covalent/non-covalent, reversible/irreversible, and native/denatured interactions and characteristics. For protein therapeutics, the presence of aggregates is typically considered undesirable because of the concern that aggregates may cause an immunogenic reaction (e.g., small aggregates), or may cause adverse events on administration (e.g., particulates). Dynamic light scattering refers to a technique that can be used to determine the size distribution profile of small particles in suspension or polymers such as proteins in solution. This technique, also referred to as photon correlation spectroscopy (PCS) or quasi-elastic light scattering (QELS), uses scattered light to measure the rate of diffusion of the protein particles. Fluctuations of the scattering intensity can be observed due to the Brownian motion of the molecules and particles in solution. This motion data can be conventionally processed to derive a size distribution for the sample, wherein the size is given by the Stokes radius or hydrodynamic radius of the protein particle. The hydrodynamic size depends on both mass and shape (conformation). Dynamic scattering can detect the presence of very small amounts of aggregated protein (<0.01% by weight), even in samples that contain a large range of masses. It can also be used to compare the stability of different formulations, including, for example, applications that rely on real-time monitoring of changes at elevated temperatures. Accordingly, certain embodiments include the use of dynamic light scattering to analyze the solubility and/or presence of aggregates in a sample that contains an anti-NRP2 antibody of the present disclosure.

Although the foregoing embodiments have been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to one of ordinary skill in the art in light of the teachings of this disclosure that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims. The following examples are provided by way of illustration only and not by way of limitation. Those of skill in the art will readily

recognize a variety of noncritical parameters that could be changed or modified to yield essentially similar results.

EXAMPLES

Reference Sequences			
Protein	Species	NCBI Reference Sequence	Region and mutations
NRP2v1	Human	NP_957718.1	
NRP2v2	Human	NP_003863.2	
NRP2v3	Human	NP_958436.1	
NRP2v4	Human	NP_061004.3	
NRP2v5	Human	NP_957719.1	
NRP2v2	crab-eating macaque	XP_005574085.1	
NRP2v2	Mouse	NP_001070872.1	
SEMA3F	Human	NP_004177.3	19-779 R583A R586A
VEGF-C	Human	NP_005420.1	112-227
SEMA3F	Mouse	NP_001298080.1	19-779 R583A R586A
VEGF-C	Mouse	NP_033532	108-223
FLT4	Human	NP_891555	
PLXNA1	Human	NP_115618.3	L839P
KDR	Human	NP_002244.1	
VEGF-A	Human	NP_001165097.1	27-191

Example 1

Characterization of Humanized, Affinity Matured Anti-Neuropilin 2 Antibodies

Recombinant, humanized anti-NRP2 antibodies were generated from a panel of murine monoclonal antibodies targeting different domains of human NRP2. Humanized antibodies were affinity matured to produce highly selective, potent antibodies with sub nanomolar affinities in the context of primarily human IgG constant domain to limit the potential for immunogenicity in humans. Initial assessments of binding affinity as measured by surface plasmon resonance (SPR), and flow cytometry (FACS) binding to cell lines expressing recombinant NRP2, is presented in Table E1.

TABLE E1

Antibodies				
Antibody	Isotype	Target domain	Affinity (nM)	EC50 (nM)
aNRP2-1v3	hIgG4	b2	17	n.d.
aNRP2-2v4	hIgG4	c	2.7	n.d.
aNRP2-10v5	hIgG4	b1	0.20	n.d.
aNRP2-10v10	hIgG4	b1	0.029	0.09
aNRP2-10v13	hIgG4-YTE	b1	0.358	0.65
aNRP2-11v7	hIgG4	b2	0.95	6.3
aNRP2-14v9	hIgG4	a2	0.15	1.2
aNRP2-14v10	hIgG4	a2	0.17	47.5
aNRP2-28v2/4	mIgG1(D265A)	b1	0.0599	1.58

Humanization and affinity maturation. Humanized murine monoclonal antibodies were prepared via grafting of murine CDRs onto human IgG4 with the selective replacement of murine specific amino acid substitutions compared to the corresponding human sequences when necessary to stabilize the antibody, and screening to confirm binding to NRP2 polypeptides. Humanized antibodies were affinity matured via the systematic mutation of their CDR sequences, and the most advantageous recombination of the higher affinity combinations identified were fully characterized. Recombi-

nant antibodies were purified from conditioned medium starting at 2 weeks of culture by flowing over a Protein A affinity column, eluting and storing in Phosphate Buffered Saline (1xPBS), pH 7.4. Each lot was tested for protein concentration, purity, and endotoxin level. Purity by SDS-PAGE was routinely >90%.

Anti-NRP2 antibody binding and affinity measurements. Surface plasmon resonance (SPR) methods were used to demonstrate binding of anti-NRP2 antibodies to human NRP2 antigen and to measure binding affinities as summarized in Table E1. SPR experiments were conducted on a Bio-Rad ProteOn XPR36 Protein Interaction Array instrument. Biotinylated anti-human antibody (ThermoFisher Cat #7103302100) was immobilized on ProteOn NLC sensor chips. Anti-NRP2 antibodies were subsequently flowed over and captured by the anti-human antibody. Human NRP2 antigen protein was flowed over the captured antibody at varying concentrations (150, 50, 16.67, 5.56, 1.85 nM). The sensor chip surface was regenerated between each analyte run to remove anti-NRP2 antibodies and NRP2 protein. Data was double referenced against a surface with no anti-NRP2 antibody captured (immobilized anti-human antibody only) and a buffer only blank. Affinity constants were derived by globally fitting sensograms to a Langmuir (1:1) interaction model in the ProteOn manager software. For each anti-NRP2 antibody, data from the multiple NRP2 concentrations was fit as a single data set with dissociation rate constant (k_d), association rate constant (k_a) and R_{max} values as global parameters. The reported binding affinity is the equilibrium dissociation constant (KD) calculated from k_d/k_a .

Running buffer: 50 mM HEPES, 300 mM NaCl, 5 mM CaCl_2 , 0.005% Tween-20, pH 7.4

Immobilized antibody: CaptureSelect Human Fab-kappa Kinetics Biotin Conjugate, ThermoFisher Cat #7103302100

Antigen: Human NRP2 (aa23-855) with C-terminal Avi-, Myc- and His-tags

Regeneration buffer: 10 mM glycine pH 2.0

Purification of plasmid DNA. Plasmid containing human NRP2 were purchased from Origene. To generate a large stock of plasmids and purify the DNA, plasmid was transformed into chemically competent *E. coli* cells (One Shot TOP10) according to the manufacturer's instructions. The transformed bacteria were grown in the recommended liquid medium (Difco Dehydrated Miler Luria-Bertani medium powder resuspended in water) containing kanamycin monosulfate (50 g/mL) as an antibiotic. The plasmid DNA was then purified using the QIAGEN DNA maxi prep kit according to the kit's instructions. DNA concentration and purity were measured on a spectrophotometer (Nanodrop 2000). An A260/A280 absorbance ratio between 1.8 and 2.0 was required for transfection.

Generation of Expi293-hNRP2 clonal cells stably over-expressing human NRP2. A plasmid (Origene Technologies Cat #RC220706) encoding the human NRP2 variant 2 transcript NM_003872 (hNRP2) fused to a Myc-DDK tag was purchased. The vector was PCR amplified using Q5 polymerase (New England Biolabs Cat #M0491) with the following primer pairs:

(SEQ ID NO: 149)
5' -TGAGGATGACAAAGATTTGCAGCT-3'

(SEQ ID NO: 150)
5' -ACCGCGCGCGCGCTTTATGCCTCGGAGCAGCACTT-3'

147

-continued

(SEQ ID NO: 151)
 5' -AGTGCCAAGCAAGCAACTCAA-3'

(SEQ ID NO: 152)
 5' -AAGTGTGCTCCGAGGCATAAACGGCCGCGGT-3'

The resulting PCR products were then fused, cut with MfeI/AgeI (New England Biolabs Cat #R3589, R3552), and ligated into a vector fragment of RC220706 cut with the same enzymes. This vector, containing an untagged human NRP2 transcript, was then linearized and re-suspended in 10 mM Tris-0.1 mM EDTA. Suspension Expi293 cells, (a HEK293 based transient expression system which is maintained in suspension cultures from ThermoFisher; Cat #A14527), were grown in expression medium (ThermoFisher Cat #A1435101) at 37° C. and 8% CO₂. The linearized plasmid described above was transfected into Expi293 cells using an SF Cell Line 4D-Nucleofector® X Kit L (Lonza Cat #V4XC-2012) and standard protocol T-030 for suspension HEK293 cells. Cells were allowed to recover in static culture for 17 hours, transferred to suspension and recovered an additional 72 hours, and then were selected with 200-350 µg/mL G418 in 50 µg increments (ThermoFisher Cat #10131035). Cell densities and viabilities were monitored for a period of 3 weeks, with fresh media/antibiotic replacement every 2-3 days.

To select human NRP2 over-expressing clonal cells, cells were suspended into 96-well plates by limited dilution. Single colonies were transferred to falcon tubes for further expansion under maintained selection pressure. Clonal cells that over-express human NRP2 were validated by a-NRP2 staining followed with flow cytometry analysis.

Binding of anti-NRP2 antibodies to Expi293 cells expressing human NRP2 (hNRP2) clonal cells. Expi293-hNRP2 clonal cells were collected by centrifugation at 300 g for 5 minutes, and washed twice by DPBS (with calcium and magnesium, ThermoFisher Cat #14040133). Washed cells were added to a 96-well V-bottom plate (ThermoFisher Cat #1424572) at 100,000 cells/well in 50 µL of DPBS added with Zombie Violet viability stain (Biolegend Cat #423114, diluted at 1:500). From this step on, cells were protected from light. Cells were kept at room temperature for 10 min and pelleted down at 300 g for 5 minutes at 4° C. Supernatants containing Zombie Violet were discarded, and without washing, 30 µL of antibodies diluted in the flow wash buffer (FWB, DPBS plus 2% FBS and 0.1% sodium azide to inhibit receptor internalization) were added to the cells. For anti-NRP2 antibody staining on Expi293-hNRP2 clonal cells, in-house human anti-NRP2 antibodies were tested at final concentrations of 0.006-100 nM at 4-fold dilutions. An isotype control human IgG4 antibody (Biolegend Cat #403701) was used instead of the anti-NRP2 antibody to demonstrate the specificity of the staining. Binding was allowed to proceed on ice for 40-60 minutes. Cells were then pelleted at 300 g for 5 minutes at 4° C., supernatants removed, and cells washed twice by adding 150 µL of FWB and centrifuging the cells again in the same conditions. To detect a-NRP2 binding on the cell surface, Cy3-conjugated goat anti-human IgG Fc (Jackson ImmunoResearch Cat #109165098) was added to the cells at a final concentration of 3 µg/mL in 30 µL of FWB. After 30-40 minutes of incubation on ice, cells were pelleted at 300 g for 5 minutes at 4° C. and supernatants removed, and the cells were washed twice as described previously. Cells were then resuspended in FWB before acquisition on the Cytotflex. Gains were set based on staining controls. Cells were collected with typically more than 10,000 events. Cells were

148

then analyzed on FlowJo analysis software by excluding dead cells and gating singlets. The statistical analysis was performed using GraphPad Prism. A four-parameter variable-slope curve was fitted to the data ([agonist] vs. response) using non-linear regression, and the EC₅₀ and r² for each curve was determined.

The EC₅₀ of four anti-NRP2 antibodies directed against different domains of NRP2, were measured for binding to Expi293-hNRP2 clonal cells (see Table E1 and FIG. 3). They showed specific binding to NRP2 with EC₅₀s in the subnanomolar or nanomolar range. By contrast, no binding was observed by the isotype control human IgG4 antibody.

Example 2

Binding of Anti-Human Neuropilin 2 Antibodies to Cynomolgus Monkey Nrp2

Cross reactivity of the anti-NRP2 antibodies to cynomolgus monkey NRP2 was assessed by surface plasmon resonance (SPR) on recombinant proteins, and flow cytometry (FACS) binding to NRP2 expressed on the surface of HEK 293 cells. Establishing cross reactivity between cynomolgus monkey and human NRP2 is an important therapeutic development consideration to ensure that potential therapeutic candidates can be readily assessed for toxicity in animal studies; particularly if there is little or no cross reactivity of the antibodies between human and rodent NRP2.

Plasmid DNA. Plasmids containing human or cynomolgus monkey NRP2 were purchased from Origene. Plasmid propagation and purification was performed as described for the human NRP2 plasmid.

Expi293 propagation and transient transfection. Expi293 cells, a HEK293 transient expression system maintained in suspension cultures, were used to express cynomolgus monkey NRP2. Cells were grown in 60 mL Expi293 medium within 250 mL vented suspension culture flasks. The propagation of cells was carried out in a Multitron Cell incubator at 37° C., with 8% CO₂, 80% humidity, and shaking at 225 rpm. During the maintenance and expansion phase, Expi293 cells were split twice a week at 0.3×10⁶ cells/mL in order to keep the density within the optimal range for transfection and the viability high. Cell densities and viabilities were determined using a cell counter (Cedex HiRes Cell Analyzer).

The day prior to DNA transfection, the Expi293 cells were seeded at a density of 2.0×10⁶ cells/mL in order to maintain high viability (>95%) and low density (not to exceed 3-5×10⁶ cells/mL), thereby providing cells with fresh nutrients and avoiding transfection impediments caused by any secreted substances.

Prior to DNA transfection, Expi293 cells were counted and reseeded at 2.5×10⁶ cells/mL in 50 mL shake flasks (TPP TubeSpin bioreactor tubes). The transfection procedure was performed using the Expifectamine kit. The manufacturer's protocol was adapted for a transfection in a total volume of 5 mL instead of 30 mL (i.e., all volumes were divided by a factor of 6). Five µg of each plasmid DNA was diluted in Opti-MEM reduced serum medium, complexed with Expifectamine transfection reagent, and transfected into Expi293 cells. The transfected pools were cultured with shaking at 225 rpm to ensure complete suspension of the cells within the 50 mL bioreactors. Following the manufacturer's protocol, Expifectamine kit enhancers were added after 16-18 hours, and the cultures analyzed by flow cytometry 2 days post-transfection.

Binding of a-NRP2 to cynomolgus monkey NRP2-transfected Expi293 cells. Cynomolgus monkey (cyno) NRP2-

transfected Expi293 cells were collected by centrifugation at 300 g for 5 minutes, and washed twice by DPBS (with calcium and magnesium, ThermoFisher Cat #14040133). Washed cells were added to a 96-well V-bottom plate (ThermoFisher Cat #1424572) at 100,000 cells/well in 50 μ L of DPBS added with Zombie Violet viability stain (Biolegend Cat #423114, diluted at 1:500). From this step on, cells were protected from light. Cells were kept at room temperature for 10 min and pelleted down at 300 g for 5 minutes at 4° C. Supernatants containing Zombie Violet were discarded, and without washing, 30 μ L of antibodies diluted in the flow wash buffer (FWB, DPBS plus 2% FBS and 0.1% sodium azide to inhibit receptor internalization) were added to the cells. For a-NRP2 staining on Cyno NRP2-transfected Expi293 cells, in-house human anti-NRP2 antibodies were tested at final concentrations of 0.006-100 nM at 4-fold dilutions. For each test, NRP2 over-expression on the cell surface was validated by a-NRP2 staining (10 μ g/mL R&D #AF567 a-NRP2) in separate wells. Binding was allowed to proceed on ice for 40-60 minutes. Cells were then pelleted at 300 g for 5 minutes at 4° C., supernatants removed, and cells washed twice by adding 150 μ L of FWB and centrifuging the cells again in the same conditions. To detect a-NRP2 binding on the cell surface, Cy3-conjugated goat anti-human IgG Fc or AF647-conjugated donkey anti-goat IgG (Jackson ImmunoResearch Cat #109165098 & 705605147) was added to the cells at a final concentration of 3 μ g/mL in 30 μ L of FWB. After 30-40 minutes of incubation on ice, cells were pelleted at 300 g for 5 minutes at 4° C. and supernatants removed, and the cells were washed twice as described previously. Cells were then resuspended in FWB before acquisition on the Cytoflex. Gains were set based on staining controls. Cells were collected with typically more than 10,000 events. Cells were then analyzed on FlowJo analysis software by excluding dead cells and gating singlets.

As additional controls, the following conditions were tested: 1) An isotype control human IgG4 antibody (BioLegend Cat #403701) was used instead of the anti-NRP2 antibody to demonstrate the specificity of the staining; and 2) Staining with R&D #AF567 a-NRP2 was also determined using mock-transfected Expi293 cells to demonstrate over-expression of Cyno NRP2 on Expi293 cells. The statistical analysis was performed using GraphPad Prism. A four-parameter variable-slope curve was fitted to the data ([agonist] vs. response) using non-linear regression, and the EC₅₀ and r² for each curve was determined.

The EC₅₀ of three anti-NRP2 antibodies, each against a distinct domain of NRP2, were measured for binding to Cyno NRP2-transfected Expi293 cells (see Table E2 and FIG. 4). They all showed specific binding to NRP2 with EC₅₀s in the nanomolar range. By contrast, no binding was observed by the isotype control human IgG4 antibody.

Anti-NRP2 antibody binding and affinity measurements. Surface plasmon resonance (SPR) methods were used to demonstrate binding of anti-NRP2 antibodies to cynomolgus NRP2 antigen and to measure binding affinities as summarized in Table E2. SPR experiments were conducted on a Bio-Rad ProteOn XPR36 Protein Interaction Array instrument. Biotinylated anti-human antibody (ThermoFisher Cat #7103302100) was immobilized on ProteOn NLC sensor chips. Anti-NRP2 antibodies were subsequently flowed over and captured by the anti-human antibody. Cynomolgus NRP2 antigen protein was flowed over the captured antibody at varying concentrations (150, 50, 16.67, 5.56, 1.85 nM). The sensor chip surface was regenerated between each analyte run to remove anti-NRP2 antibodies

and NRP2 protein. Data was double referenced against a surface with no anti-NRP2 antibody captured (immobilized anti-human antibody only) and a buffer only blank. Affinity constants were derived by globally fitting sensograms to a bivalent analyte interaction model in the ProteOn manager software. For each anti-NRP2 antibody, data from the multiple NRP2 concentrations was fit as a single data set with dissociation rate constant (k_d), association rate constant (k_a) and R_{max} values as global parameters. The reported binding affinity is the equilibrium dissociation constant (KD) calculated from k_d/k_a .

Running buffer: 50 mM HEPES, 300 mM NaCl, 5 mM CaCl₂, 0.005% Tween-20, pH 7.4

Immobilized antibody: CaptureSelect Human Fab-kappa Kinetics Biotin Conjugate, ThermoFisher Cat #7103302100

Antigen: Cynomolgus NRP2 (aa23-854) with C-terminal Fc-tag

Regeneration buffer: 10 mM glycine pH 2.0

TABLE E2

Comparison of binding affinity to cynomolgus monkey NRP2			
Antibody	Domain Specificity	EC50 (nM)	Affinity (nM)
aNRP2-14v9	a2	2.8	0.017
aNRP2-10v10	b1	2.9	0.032
aNRP2-11v7	b2	5.7	12
Isotype control antibody	n.d.	n.d.	n.d.

The results from these studies demonstrate that each of the tested antibodies showed comparable binding affinities to human and cyno NRP2, which affinities that were all low nanomolar and within about 5-6 fold of each other between species.

Example 3

Characterization of Anti-Nrp2 Antibodies on Receptor NRP2 Dimerization

To further extend the assessment of the biological activity of the anti-NRP2 antibodies, their activity was assessed in a receptor dimerization assay. In brief, vectors encoding a split luciferase pBiT1.1 and pBiT2.1 were obtained from Promega corporation (Madison, Wisconsin). The complete extracellular domain and transmembrane helices of NRP2a (version 2), NRP2b (version 5) (See table N1), FLT4 (VEGFR3), KDR (VEGFR2), and plexin A1 (PLXNA1) were cloned into the vectors and screened for optimal orientation, receptor density, and ligand induced dimerization, following established methods. Expi293 cells (Fisher) were transfected at 1 million cells/mL ~20 hours prior to the assay with NRP2 and a co-receptor at equal mass amounts. Cells were counted, and 100,000 live cells were plated in a well of a white luminometer plate in Live Cell Imaging Solution (Fisher). Nano-Glo assay substrate (Promega) was added, and the plate was read on a Glomax 96 luminometer at room temperature to establish baseline luminescence. Antibodies were added to the well at 100 nM and the plate was again read to establish a new baseline and monitor the effects of the antibodies on spontaneous receptor dimerization. Depending on the receptor pair, typically 200 nM SEMA3F-p95 (aTyr Pharma), 2 nM VEGF-C(R&D Systems, Minnesota), or 2 nM VEGF165 (R&D Systems, Minnesota) were then added and the plate was again read to measure dimerization of receptors. Normalized responses

for each individual well were calculated to the time point prior to addition of ligand, then normalized to no-antibody/no-ligand (baseline signal). Replicates were then processed to give average response and standard deviation. Results are summarized in Table E3 below, and shown graphically in FIGS. 3-10.

TABLE E3

Receptor Dimerization blocking summary:				
Antibody	Domain specificity	Ability to Block NRP2/PLXNA1	Ability to Block NRP2/FLT4(VEGFR3)	Ability to Block NRP2/KDR(VEGFR2)
aNRP2-14v9	a2	Very strong (80-100% inhibition)	None (Does not substantially block)	None (Does not substantially block)
aNRP2-10v10	b1	Weak/None (Does not substantially block)	Very strong (80-100% inhibition)	Completely (80-100% inhibition)
aNRP2-11v7	b2	Enhances receptor complex formation (greater than 20%)	Partial blocking (20-80% inhibition)	Enhances receptor complex formation (greater than 20%)
NRP2 ^B	b1	Enhances receptor complex formation (greater than 20%)	None (Does not substantially block)	None (Does not substantially block)

The results demonstrate that all of the tested antibodies show differential functional activity in these assays. Surprisingly specific antibodies show extremely specific and non-obvious functional differentiation at the level of receptor dimerization, demonstrating that specific epitopes have unique functional roles in this receptor system.

For example, antibody aNRP2-14v9, which binds to the a2 domain of NRP2, was demonstrated to exhibit extremely potent inhibition of NRP2-plexin heterodimerization, while having no significant impact on NRP2-VEGFR3 heterodimerization. These characteristics suggest that this antibody would potentially inhibit semaphorin 3 signaling through NRP2, without significantly inhibiting the signaling through VEGFR2 or 3. Such characteristics therefore make this antibody well suited for applications where the selective modulation of semaphorin signaling is required. For example, for modulating effects of semaphorin signaling on axonal growth avoidance, lymphatic sprouting, cell migration, smooth muscle contractility, immune cell activation and exhaustion, as more generally modulating cell growth and proliferation.

Additionally, antibody aNRP2-10v10, which binds to the b1 domain of NRP2, shows the ability to potentially inhibit NRP2-VEGFR3 heterodimerization, without significantly inhibiting NRP2-plexin heterodimerization. These characteristics suggest that this antibody would potentially inhibit VEGF signaling through NRP2, without significantly inhibiting the signaling through the plexin receptor/semaphorin 3 pathways. Such characteristics therefore make this antibody well suited for applications where the selective modulation of VEGF signaling is required. For example, for modulating effects of VEGF on lymphangiogenesis, neovascularization, angiogenesis, cancer cell growth, cancer initiation, cancer migration, cancer cell adhesion, invasion, chemoresistance development and metastasis. Importantly this antibody does not significantly impair semaphorin 3 signaling, making it uniquely suited for applications where endogenous semaphorin signaling may be advantageous—for example in any of the applications listed above.

By contrast antibody aNRP2-11v7, which binds to the b2 domain of NRP2, shows the ability to inhibit the heterodimerization of NRP2 to FLT4, but increased the ligand induced dimerization signal for both Plexin and KDR. Suggesting that it stabilizes heterodimeric and higher

order complexes of NRP2. These characteristics suggest that this antibody could have unique context dependent signaling effects to inhibit FLT4 signaling while promoting signaling through alternative NRP2 dependent pathways.

Most surprisingly, in relation to antibody aNRP2-10v10, the previously described NRP2B antibody (Caunt et al.,

Cancer Cell. 13:331-342, 2008), shows no ability to inhibit ligand induced dimerization for any of the tested receptor pairs, while enhancing the ligand induced dimerization of the NRP2/PLXNA1 receptor pair in this assay system. These differential effects of the NRP2B suggests that it binds to a unique epitope of NRP2 compared to aNRP2-10v10, and has functionally distinct effects.

Example 4

Epitope Mapping of aNRP2-10v10 Antibody

Human NRP2a (version 2; see Table N1) and mouse NRP2v2 cDNAs were obtained from Origene. Because we have previously demonstrated that the aNRP2-10v10 antibody binds specifically to the b1 domain of human NRP2, but not mouse NRP2, mutations in the b1 domain of mouse NRP2 sequence (F299Y, K354N, 1383V, M400A, 1407V, and L416S) were made in isolation to restore the amino acids which differ between the human and mouse proteins in this region. Human and mouse NRP2, expression vectors, along with expression vectors encoding all 6 mutants were transfected into Expi293 cells at 1 million cells/mL, by standard techniques. Approximately 40 hours post transfection cells were co-stained with aNRP2-10v10, and a species agnostic anti-NRP2 a1 specific binding antibody. Cells were gated on live, single, NRP2 expressing cells (a1 binder positive), as shown in FIG. 11A, and the degree of aNRP2-10v10 staining was quantified as mean fluorescent intensity (MFI), as shown in FIG. 11B.

The mutations F299Y, K354N, and L416S restored partial, but incomplete, binding activity of aNRP2-10v10 to mouse NRP2, while mutation of the remaining residues, 1383V, M400A, and 1407V showed no improvement in binding (see FIG. 11B). All substitutions that improve binding of aNRP2-10v10 to mouse NRP2 map on a shared surface of the b1 domain in close proximity in the published human NRP2 PDB structure 2QOK. The three residues map 15.7, 8.9, and 14.3 angstroms apart in the structure, and form a shared epitope. Y299, N354 and S416 also surround the binding pocket ("C-wall") for the C-terminus of VEGF-C, and are shown surrounding T319 which forms the base of the binding pocket (see FIG. 12), where previous work has shown a mutation T319R, blocks binding of VEGF-C (Parker 2015).

153

Example 5

Inhibition of Anchorage-Independent Growth and Sensitization of Tumor Cells to Chemotherapy by aNRP2-10v10

Anchorage-independent growth is the ability of transformed cells to grow independently of a solid surface, and is a hallmark of carcinogenesis. The soft agar colony formation assay is a well-established method for characterizing this capability in vitro (J Vis Exp. 2014; (92): 51998). Human triple negative breast cancer (TNBC) cell lines MDA-MB-231 (HTB-26™) and BT549 (HTB122™) were obtained from ATCC and employed in this assay.

MDA-MB-231 was cultured in 1×RPMI media containing 10% FBS and 1% P/S. BT549 was cultured in 1×RPMI media containing 10% FBS, 0.023 U/mL insulin and 1% P/S. Cells were recovered and expanded from frozen stocks for 2-3 passages. The NRP2-high population was sorted out on FACS equipment following cell surface staining by a fluorophore-conjugated anti-NRP2 antibody. The colony formation assay was performed using the soft agar colony formation fluorometric assay kit obtained from Cell Biolab (#CBA-130-CB). Briefly, the sorted NRP2-high cells were suspended as single cell suspensions, and seeded at 1-1.5 K cells/96-well in the 3D soft agar matrix. After overnight incubation to stabilize the cells, the treatment reagents were spiked into the culture media. The chemo drug Cisplatin or 5-FU was added at a final concentration of IC30 or 1/2 IC30, based on a preliminary test that determined their inhibition curves on MDA-MB-231 cells, in combination with the antibodies a-NRP2-10v10 or its isotype control hIgG4 at 100 nM. Reagents were renewed by spiking-in every 3 days. After 6 or 9 days of treatment, the soft agar matrix was dissolved and the formed cell colonies were quantified using CyQuant dye diluted in PBS (provided in the Cell Biolab kit), which produced a fluorescence signal proportional to viable cells.

The NRP2-high population of TNBC cells were sorted for use in the colony formation assay because NRP2 is expressed preferentially on breast cancer stem cells (CSCs) and VEGF/NRP2 signaling was shown to be important for the genesis of TNBCs and tumor initiation (EMBO Mol Med. 2013 5 (4): 488-508). Compared to the hIgG4 control, the a-NRP2-10v10 antibody in combination with either Cisplatin or 5-FU showed a significant inhibition of colony formation by MDA-MB-231 cells (FIG. 13A). Similar effect was observed for the BT549 cells at certain doses of Cisplatin or 5-FU. (FIG. 13B). In a separate test with aNRP2-10v10 treatment alone, no obvious effect in the colony formation assay was observed with these cells. The results suggest that aNRP2-10v10 sensitized the TNBC CSC-like cells to chemo drug treatment and has the potential to be developed in a combination therapy in combating cancer.

Example 6

Inhibition of Anchorage Independent Growth and Sensitization to Cisplatin and Bevacizumab Therapy by aNRP2-10V10

The immunostaining of TNBC MDA-MB-231 cells revealed that >90% of these cells exhibit a cancer stem cell (CSC)-like phenotype (CD44+/CD24-), therefore we further performed the colony formation assay using parental MDA-MB-231 cells without pre-sorting. The matrix for 3D colony formation assay was changed to methylcellulose which is more solid, better mimics the extracellular matrix in vivo, and shows better signal-to-background than soft

154

agar. Other experimental details are essentially the same as described in the previous example.

The results demonstrated that the anti-NRP2 antibody aNRP2-10v10 in combination with either the chemo drug Cisplatin or the anti-VEGF-A drug Bevacizumab, showed a significantly enhanced inhibition of colony formation by MDA-MB-231 cells, compared to either treatment with the isotype control hIgG4 antibody (see FIG. 14). aNRP2-10v10 thus significantly augments the effectiveness of both cisplatin and Bevacizumab in treating triple negative breast cancer.

Example 7

Inhibition of Cell Migration of Human Lymphatic Endothelial Cell (HLEC) Migration Toward Vascular Endothelial Growth Factor C (VEGF-C) by Antibodies ANRP2-10V10 and ANRP2-11V7

Migration assays were performed using a modified Boyden Chamber with 8 µm pore size membrane, 16-well CIM plate (ACEA Biosciences.) Both sides of the membrane were coated with 5 µg/ml Fibronectin (Sigma) for 30 minutes at 37° C., 5% CO₂. 160 µl of 191.5 pM VEGF-C or assay medium (0.1% BSA, ECM) was added to the lower chamber, the upper well attached, 50 µl assay medium added to the upper chamber before replacing the lid, and the plate was incubated for 1 hour at 37° C., 5% CO₂ before baseline measurement was taken. HLECs were harvested, washed, and re-suspended in assay medium at 100,000 cells/well with aNRP2 antibody or control proteins and incubated for 30 minutes at 37° C., 5% CO₂. Cells were added to the plate and incubated at room temperature for 30 minutes. Plate was inserted into xCELLigence RTCA DP (ACEA Biosciences) instrument, and migration was measured real-time every 5 minutes over a 24 hour period (2 wells per condition).

The results demonstrated that the anti-NRP2, VEGF-C blocking antibodies, aNRP2-10v10, and aNRP2-11v7 were able to effectively block the migration of HLEC cells towards VEGF-C in a dose dependent manner (see FIG. 15). HLECs were inhibited by both NRP2 antibodies to levels consistent with positive control antibodies (αKDR or 3C5) directed against VEGFR2 or VEGFR3 respectively.

Example 8

Blocking of Semaphorin 3F Mediated Inhibition of Akt Phosphorylated by Antibodies ANRP2-14V10 and ANRP2-11V7, but not ANRP2-10V10

As shown schematically in FIG. 17, the treatment of cells with Sema3F should specifically block PI3K activity via NRP2 mediated interaction with plexin A1, thereby resulting in reduced phospho-Akt levels. To directly test the ability of the anti-NRP2 antibodies to block this effect, lysates were prepared following standard protocols (MesoScale Discovery, Phospho (Ser473)/Total Akt Assay Whole Cell Lysate Kit #K15100D). Serum starved and confluent U251 cells (Sigma, Cat. 09063001) were treated with 10 nM rapamycin (Sigma, Cat. 553211, 30 minutes), and blocking antibodies or controls were added (10 µg/mL, 30 minutes), followed by 25 nM Sema3F or serum free media for 1 hour in triplicate. Lysate protein concentrations were determined by BCA assay and samples were equalized to 0.4 mg/mL. Phospho and Total AKT were measured following the manufacturer's instructions and percent phosphorylation was calculated by dividing phosphorylated AKT relative units by total AKT relative units. Blocking of Sema3F for each antibody or

155

control was determined by comparing percent phosphorylation of Sema3F treated cells to percent phosphorylation of non-Sema3F treated cells.

Treatment of U251 cells with Sema3F resulted (see FIG. 16) in an approximately 50% decrease in intracellular phospho-Akt levels that can be blocked by both the semaphorin blocking antibodies aNRP2-11v7 and aNRP2-14v10, but is not blocked by the VEGF-C blocking antibody aNRP2-10v10 or with mIgG1 and hIgG4 control antibodies.

Example 9

Evaluation of Anti-Tumor Activity of Test Compound ANRP2-28 on B16-F10 Mouse Melanoma in C57BL/6 Mice

Studies were performed to investigate the in vivo anti-cancer potential of antibody aNRP2-10v10 using a surrogate antibody, aNRP2-28, (which recognizes mouse NRP2 at a similar epitope region as the antibody aNRP2-10v10 (which does not cross react with mouse NRP2) in a syngeneic mouse model using B16-F10 cells (a mouse melanoma cancer model) prepared as described below.

Animals and Animal husbandry. All mice were ear tagged for identification purpose. Upon arrival, animals were examined to ensure that they were healthy. The animals were housed in autoclaved solid floor polycarbonate cages. Housing and sanitation were performed according to industry standards. All animal handling was performed in a laminar flow hood located in a clean room. In all experiments, euthanasia criteria were set for an upper limit on subcutaneous tumor volume (usually $>2,000 \text{ mm}^3$) and body weight loss (usually $>20\%$ body weight loss for an individual during the study). Animal work was performed at either Comparative Biosciences, Inc., 786 Lucerne Dr., Sunnyvale, CA 94085 or at aTyr Pharma, 3545 John Hopkins Court, Suite 250, San Diego, CA 92121. The test antibody dosing regimen was initiated once the mean tumor volume reached $50\text{-}100 \text{ mm}^3$. Tumor measurements were recorded two times a week until study termination, as described below.

Tumor monitoring. Once palpable, sub-cutaneous tumors were measured three times a week with a digital caliper. Tumor volumes were calculated using formula:

$$\text{Tumor Volume} = \text{length} \times \text{width} \times \text{width} \times 1/2$$

Statistical analysis. Data are expressed as mean $\pm$ SEM. In experiments in which animals were euthanized due to tumor burden or body weight loss, the terminal tumor volume was carried forward for statistical analysis. In cases where an animal was found dead but did not have a large tumor (cause of mortality unknown), the animal's data were removed prior to statistical analysis. Significance of difference over time was tested with a 2-way repeated measures ANOVA followed by Dunnett's post-hoc test. Group comparisons were conducted using 1-way ANOVA (parametric or Kruskal Wallis, as noted in the figure legends). A p value <0.05 was considered significant.

Cell culture and implantation. Mouse B16-F10 cell line was purchased from ATCC. The cells were cultured in 75 cm^2 flasks containing DMEM media supplemented with 10% fetal bovine serum (FBS) and incubated at 37°C . in humidified atmosphere of $5\% \text{ CO}_2$. As cells reached 90% confluence, cultures were expanded to 175 cm^2 flasks until sufficient cells are available for injection. 10,000 cancer cells in PBS with 20% matrigel were subcutaneously into the right flank of each mouse. The day on which the tumor was implanted is designated as Day 0.

156

Treatment Regimen. Thirty (30) C57BL/6 mice (Charles River, female, 6-8 wks old) were used in this study. The animals were assigned to 3 study groups of 10 mice randomly, and housed as described in the Materials and Methods. The antibody dosing regimen is shown in Table E4 below; in brief animals received intraperitoneal injections of the control IgG antibody, positive control antibody (Bevacizumab), and anti-NRP2 antibody (as described in the Materials and Methods), administered according to the protocol below (Table E4).

TABLE E4

GROUP TREATMENTS					
Group	#Mice	Test Materials	Dose (mg/kg)	Route of Admin. (ROA)	Frequency
1	10	Control IgG	25	IP	BIW
2	10	aNRP2-28	25	IP	BIW
3	10	Bevacizumab	5	IP	BIW

* IP = Intraperitoneal injection

Summary of results. Animals bearing B16-F10 tumors and treated with three doses of the mouse surrogate antibody aNRP2-28, which recognizes mouse NRP2 (and is a functional surrogate for anti-human NRP2 antibody aNRP2-10v10). Treatment with aNRP2-28 resulted in tumor growth inhibition compare to the IgG control group, reaching statistical significance ($p < 0.05$) on Day 14, 16 and 19 (FIG. 19A). Accordingly, aNRP2-28 treated mice had the lowest terminal tumor weights (FIG. 19B). In addition, the anti-NRP2 antibody had more potent anti-tumor activity in this study than the VEGF-a blocking antibody Bevacizumab that was chosen as the positive control. There was no evidence of toxicity from animal body weight measurements and observations during the study (data not shown). These results demonstrate that anti-NRP2 antibodies have clear potential for therapeutic impact on cancer growth in this murine melanoma model system.

Example 10

Evaluation of Antimetastatic Activity of Test Compound ANRP2-28 in a Model of Spontaneous Metastasis

Studies were performed to investigate the in vivo anti-metastatic potential of antibody aNRP2-10v10 using a surrogate antibody, aNRP2-28, which recognizes mouse NRP2 at a similar epitope region as antibody, aNRP2-10v10 (which does not cross react with mouse NRP2) in 4T1 cells (a melanoma cancer model) prepared as described below. The test antibody dosing regimen was initiated one day before cell injection and animal weights and tumor measurements were recorded three times a week until study termination.

Mouse 4T1 cell line was purchased from ATCC. The cells were cultured in 75 cm^2 flasks containing RPMI-1640 media supplemented with 10% FBS and incubated at 37°C . in humidified atmosphere of $5\% \text{ CO}_2$. As cells reached 90% confluence, cultures were expanded to 150 cm^2 flasks until sufficient cells are available for injection. 50,000 cancer cells in PBS were implanted subcutaneously into the right flank of each mouse.

Treatment Regimen. Twenty (20) Balb/c mice (Charles River, female, 5-6 wks old) were used in this study. The animals were assigned to 3 study groups of 10 mice randomly, and housed as described in the Materials and Meth-

157

ods. The antibody dosing regimen is shown in Table E5 below; in brief animals received intraperitoneal injections of 10 mg/kg of the control IgG antibody, and aNRP2-28 (as described in the Materials and Methods), administered according to the protocol below (Table E5); starting one day before cancer cell implantation until termination on day 23).

TABLE E5

GROUP TREATMENTS					
Group	#Mice	Test Materials	Dose (mg/kg)	Route of Admin. (ROA)	Frequency
1	10	Control IgG	10	IP	days -1, 2, 6, 9, 13, 15, 20, 23
2	10	aNRP2-28	10	IP	days -1, 2, 6, 9, 13, 15, 20, 23

* IP = Intraperitoneal injection

Summary of results. Animals bearing 4T1 tumors and treated with mouse surrogate antibody aNRP2-28, which recognizes mouse NRP2 (and is a functional surrogate for anti-human NRP2 antibody aNRP2-10v10) throughout the study showed a reduction in metastatic nodules in the lung at termination, reaching statistical significance ($p \leq 0.05$) (FIG. 20A). Accordingly, there is a linear correlation between terminal antibody concentration and number of metastatic nodules (FIG. 20B). There was no evidence of toxicity from animal body weight measurements and observations during the study (data not shown). These results demonstrate that anti-NRP2 antibodies have clear potential for prophylactic, and potentially therapeutic impact on development of distant metastasis in this model system.

Example 11

Evaluation of Anti-Tumor Activity of Test Compounds in Combination with Cisplatin in a Xenograft Model of Triple-Negative Breast Cancer

Studies were performed to investigate the in vivo anticancer potential of the combination of the mouse surrogate antibody aNRP2-28, which recognizes mouse NRP2 (to block endogenous mouse NRP2) in combination with the anti-human NRP2 antibody aNRP2-10v10 (to block NRP2 on the human MDA-MB-231 cells) in combination with cisplatin in a human xenograft mouse model using MDA-MB-231 cells (Triple Negative Breast Cancer, TNBC cancer model). The dosing regimen was initiated when the mean tumor volume of all study animals reached 50-100 mm³.

Treatment Regimen. Twenty (20) NSG mice (Jackson Laboratory, female, 5-6 wks old) were used in this study. The animals were assigned to two study groups of 10 mice randomly. The MDA-MB-231 cell line was prepared for injection as described in the Materials and Methods. The dosing regimen is shown in Table E6 below; In brief animals received injections of 25 mg/kg of each the mouse surrogate antibody aNRP2-28 and aNRP2-10v10 or IgG control, which were administered to mice intraperitoneally according to the protocol below (Table E6).

Human MDA-MB-231 cell line was purchased from ATCC. The cells were cultured in 75 cm² flasks containing DMEM media supplemented with 10% fetal bovine serum (FBS) and incubated at 37° C. in humidified atmosphere of 5% CO₂. As cells reached 90% confluence, cultures were expanded to 175 cm² flasks until sufficient cells are available for injection. 1.25×10⁶ cancer cells in PBS with 20% matri-

158

gel were injected orthotopically into the mammary fat pad. The day on which the tumor was implanted is designated as Day 0.

TABLE E6

GROUP TREATMENTS					
Group	#Mice	Materials	Dose (mg/kg)	ROA	Frequency
1	10	Control IgG/ Cisplatin	25/1-5	IP	day 19- termination
2	10	aNRP2-10/ aNRP2-28/ Cisplatin	25/25/ 1-5	IP	day 19- termination

* IP = Intraperitoneal injection

Summary of results. There was significant loss in body weight observed in both groups likely due to cisplatin-associated toxicity. Addition of the anti-NRP2 antibodies to the cisplatin treatment regimen increased the tumor-inhibitory effect of cisplatin starting on day 40 post inoculation, reaching statistical significance on day 60 (FIG. 21). These results demonstrate that targeting NRP2 in conjunction with cisplatin treatment enhances the chemotherapy efficacy in this experimental model of TNBC.

Example 12

Evaluation of Anti-Tumor Activity of aNRP2-10 in Combination with Fluorouracil in a Xenograft Model of Non-Small Cell Lung Cancer

This study aimed to investigate the in vivo anticancer potential of aNRP2-10v10 in combination with fluorouracil (5-FU) in a human xenograft mouse model using A549 cells (NSCLC cancer model). The dosing regimen was initiated when the mean tumor volume of all study animals reached 50-100 mm³ on day 16.

Treatment Regimen. Twenty (20) Nu/J mice (Jackson Laboratory, female, 6-8 wks old) were used in this study. The animals were assigned to two study groups of 10 mice randomly. The A549 cell line was prepared for injection as described in the Materials and Methods. The dosing regimen is shown in Table E7 below; In brief animals received injections of 12.5 mg/kg of aNRP2-10 or IgG control, which were administered to mice intraperitoneally twice a week according to the protocol below (Table E7). Both groups also received 2.5 mg/kg of 5-FU three times a week.

Human A549 cell line was purchased from ATCC. The cells were thawed and cultured in 75 cm² flasks containing DMEM media supplemented with 10% FBS and incubated at 37° C. in humidified atmosphere of 5% CO₂. As cells reached 90% confluence, cultures were expanded to 150 cm² flasks until sufficient cells are available for injection. 2×10⁶ cancer cells in PBS with 20% matrigel were subcutaneously implanted into right flank of each mouse. The day on which the tumor was implanted is designated as Day 0.

TABLE E7

GROUP TREATMENTS					
Group	#Mice	Materials	Dose (mg/kg)	ROA	Frequency
1	10	Control IgG/5-FU	12.5/2.5	IP	BIW/TIW from day 16 on
2	10	aNRP2- 10/5-FU	12.5/2.5	IP	BIW/TIW from day 16 on

* IP = Intraperitoneal injection

During the in-life phase, animals were monitored with daily clinical observations and weekly body weights. Additionally, ocular assessments of injured eyes were scored and photos taken on post-injury Day 7 and Day 14. All animals were euthanized on Day 14. Injured eyes from all animals were removed and dissected. Whole-mounted corneas were fixed, stained with LYVE-1, and analyzed immunohistochemically to examine the efficacy of Test Articles.

onstrate that aNRP2-28 can inhibit neolymphangiogenesis in an experimental model of corneal injury. FIG. 23A shows representative image scans of Lyve-1 stained corneas of the IgG control and aNRP2-28 group. Plotting the mean values of % Lyve-1+ area for each treatment group. FIG. 23B shows a clear trend towards reduced sprouting of lymphatic vessels in the aNRP2-10 treatment groups as compared to the IgG control group. The observed effect was close to statistical significance ($p=0.07$). These data show that systemic administration of anti-NRP2 antibodies, such as aNRP2-28 and its anti-human surrogate shows pharmacologic activity in an experimental model of neovascularization.

All publications, patent applications, and issued patents cited in this specification are herein incorporated by reference as if each individual publication, patent application, or issued patent were specifically and individually indicated to be incorporated by reference.

All publications, patent applications, and issued patents cited in this specification are herein incorporated by reference as if each individual publication, patent application, or 25 issued patent were specifically and individually indicated to be incorporated by reference.

Study Design & Treatment Regimen. The study consisted of 40 female C57BL/6 mice separated into four groups. Prior to study, animals were randomized into four groups, 10 mice per group. On Day 0, neovascularization (NV) of the cornea was induced from alkali-burn injury in the eyes of animals under anesthesia. Animals were treated intraperitoneally with control IgG (Group 1), aNRP2-28 (Group 2) and Bevacizumab (Group 3) on post-injury Day 0, 3, 7 and 10.

Sequence total quantity: 153		
SEQ ID NO: 1	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable heavy CDR1 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 1		
GYTFTSYWMH		10
SEQ ID NO: 2	moltype = AA length = 17	
FEATURE	Location/Qualifiers	
REGION	1..17	
	note = Variable heavy CDR2 sequence	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 2		
AIYPGNSDTS YNQQFKG		17
SEQ ID NO: 3	moltype = AA length = 8	
FEATURE	Location/Qualifiers	
REGION	1..8	
	note = Variable heavy CDR3 sequence	
source	1..8	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 3		
RGGGYFDY		8
SEQ ID NO: 4	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Variable light CDR1 sequence	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 4		
KASQNVGA AV A		11
SEQ ID NO: 5	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	

-continued

source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 5 SASNRYT		7
SEQ ID NO: 6 FEATURE REGION	moltype = AA length = 10 Location/Qualifiers 1..10 note = Variable light CDR3 sequence	
source	1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 6 QQYSSYPLLT		10
SEQ ID NO: 7 FEATURE REGION	moltype = AA length = 10 Location/Qualifiers 1..10 note = Variable heavy CDR1 sequence	
source	1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 7 GYTFTSYWMH		10
SEQ ID NO: 8 FEATURE REGION	moltype = AA length = 17 Location/Qualifiers 1..17 note = Variable heavy CDR2 sequence	
source	1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 8 VIHPNSASTF YNERFKT		17
SEQ ID NO: 9 FEATURE REGION	moltype = AA length = 9 Location/Qualifiers 1..9 note = Variable heavy CDR3 sequence	
source	1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 9 PGTVRRSDY		9
SEQ ID NO: 10 FEATURE REGION	moltype = AA length = 16 Location/Qualifiers 1..16 note = Variable light CDR1 sequence	
source	1..16 mol_type = protein organism = synthetic construct	
SEQUENCE: 10 RSSQNIVHST GNTYLE		16
SEQ ID NO: 11 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Variable light CDR2 sequence	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 11 KVSNRFS		7
SEQ ID NO: 12 FEATURE REGION	moltype = AA length = 9 Location/Qualifiers 1..9 note = Variable light CDR3 sequence	
source	1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 12 FQGSHPWT		9
SEQ ID NO: 13 FEATURE REGION	moltype = AA length = 10 Location/Qualifiers 1..10	

-continued

source	note = Variable heavy CDR1 sequence 1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 13 GFNIKDYIHH		10
SEQ ID NO: 14 FEATURE REGION	moltype = AA length = 17 Location/Qualifiers 1..17 note = Variable heavy CDR2 sequence	
source	1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 14 RIDVEDDETK YAPKFQG		17
SEQ ID NO: 15 FEATURE REGION	moltype = AA length = 12 Location/Qualifiers 1..12 note = Variable heavy CDR3 sequence	
source	1..12 mol_type = protein organism = synthetic construct	
SEQUENCE: 15 PIYGSREAWF AY		12
SEQ ID NO: 16 FEATURE REGION	moltype = AA length = 12 Location/Qualifiers 1..12 note = Variable light CDR1 sequence	
source	1..12 mol_type = protein organism = synthetic construct	
SEQUENCE: 16 TASSSVSSSY LH		12
SEQ ID NO: 17 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Variable light CDR2 sequence	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 17 RTSNLAS		7
SEQ ID NO: 18 FEATURE REGION	moltype = AA length = 9 Location/Qualifiers 1..9 note = Variable light CDR3 sequence	
source	1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 18 HQYYRSPPT		9
SEQ ID NO: 19 FEATURE REGION	moltype = AA length = 10 Location/Qualifiers 1..10 note = Variable heavy CDR1 sequence	
source	1..10 mol_type = protein organism = synthetic construct	
SEQUENCE: 19 GFNIKDYVHH		10
SEQ ID NO: 20 FEATURE REGION	moltype = AA length = 17 Location/Qualifiers 1..17 note = Variable heavy CDR2 sequence	
source	1..17 mol_type = protein organism = synthetic construct	
SEQUENCE: 20 RIDVEDDETK YAPKFQG		17
SEQ ID NO: 21 FEATURE	moltype = AA length = 12 Location/Qualifiers	

-continued

REGION	1..12	
	note = Variable heavy CDR3 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 21		
PIYGAREAWF AY		12
SEQ ID NO: 22	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable light CDR1 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 22		
TASSSVSSSY LH		12
SEQ ID NO: 23	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 23		
RTSNLAS		7
SEQ ID NO: 24	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = Variable light CDR3 sequence	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 24		
HQYYRSPPT		9
SEQ ID NO: 25	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable heavy CDR1 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 25		
GYTFTSFGIS		10
SEQ ID NO: 26	moltype = AA length = 17	
FEATURE	Location/Qualifiers	
REGION	1..17	
	note = Variable heavy CDR2 sequence	
source	1..17	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 26		
EIYPRSGNTY YNENFKG		17
SEQ ID NO: 27	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable heavy CDR3 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 27		
SSGYYGSTPF PY		12
SEQ ID NO: 28	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Variable light CDR1 sequence	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 28		
RASQDISNYL N		11
SEQ ID NO: 29	moltype = AA length = 7	

-continued

FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 29		
YTSRLHS		7
SEQ ID NO: 30	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = Variable light CDR3 sequence	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 30		
QQGNTLPWT		9
SEQ ID NO: 31	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable heavy CDR1 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 31		
GPSLTSYGVH		10
SEQ ID NO: 32	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Variable heavy CDR2 sequence	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 32		
LIWSGGSTDY SPAFIS		16
SEQ ID NO: 33	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable heavy CDR3 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 33		
NSYSSGGYYAM DY		12
SEQ ID NO: 34	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Variable light CDR1 sequence	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 34		
KASQNVGHAV A		11
SEQ ID NO: 35	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 35		
SASNRYT		7
SEQ ID NO: 36	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable light CDR3 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 36		
QQYSRYPPYT		10

-continued

SEQ ID NO: 37	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable heavy CDR1 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 37		
GFSLTSYGVH		10
SEQ ID NO: 38	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Variable heavy CDR2 sequence	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 38		
LIWSGGSTDY SPAFIS		16
SEQ ID NO: 39	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable heavy CDR3 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 39		
NSYSSGYIAM DY		12
SEQ ID NO: 40	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Variable light CDR1 sequence	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 40		
KASQNVGTAV A		11
SEQ ID NO: 41	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 41		
SASNRYT		7
SEQ ID NO: 42	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable light CDR3 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 42		
QQRSRYPPTYT		10
SEQ ID NO: 43	moltype = AA length = 117	
FEATURE	Location/Qualifiers	
REGION	1..117	
	note = Variable heavy chain region	
source	1..117	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 43		
QVQLVQSGAE VKKPGASVKV SCKASGYTFT SYWMHWVRQA PGQGLEWMGA IYPGNSDTSY		60
NQQPKGRVTM TRDTSTSTVY MELSSLRSED TAVYYCARRG GGYFDYWQGQ TLVTVSS		117
SEQ ID NO: 44	moltype = AA length = 108	
FEATURE	Location/Qualifiers	
REGION	1..108	
	note = Light chain variable region	
source	1..108	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 44		

-continued

DIQLTQSPSF LSASVGDRVT ITCKASQNVG AAVAWYQQKP GKAPKLLIYS ASNRYTGVPS 60
 RFSGSGSGTE FTLTISSLQP EDFATYYCQQ YSSYPLLTFG GGTKVEIK 108

SEQ ID NO: 45 moltype = AA length = 118
 FEATURE Location/Qualifiers
 REGION 1..118
 note = Variable heavy chain region
 source 1..118
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 45
 QVQLVQSGAE VKKPGASVKV SCKASGYTFT SYWMHWVRQA PGQGLEWMGV IHPNSASTFY 60
 NERFKTRATM TVDRSSSTAY MELSSLRSED TAVYYCSRPG TVRRSDYWGQ GTTDTVSS 118

SEQ ID NO: 46 moltype = AA length = 112
 FEATURE Location/Qualifiers
 REGION 1..112
 note = Light chain variable region
 source 1..112
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 46
 DVVMTQSPLS LPVTLGQPAS ISCRSSQNIV HSTGNTYLEW YQRRPGQSPR LLIYKVSNRF 60
 SGVPRDRFSSG GSGTDFTLKI SRVEAEDVGV YYCFQGSHPV WTFGGGTKE IK 112

SEQ ID NO: 47 moltype = AA length = 121
 FEATURE Location/Qualifiers
 REGION 1..121
 note = Variable heavy chain region
 source 1..121
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 47
 EVQLVQSGAE VKKPGATVKI SCKVSGFNIK DYYIHVVQQA PGKGLEWMGR IDVEDDETKY 60
 APKFQGRVTI TADTSTDYAY MELSSLRSED TAVYYCATPI YGSREAWFAY WGQGTLVTVS 120
 S 121

SEQ ID NO: 48 moltype = AA length = 108
 FEATURE Location/Qualifiers
 REGION 1..108
 note = Light chain variable region
 source 1..108
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 48
 DIQMTQSPSS LSASVGDRVT ITCTASSSVS SSYLHWYQQK PGKAPKLLIY RTSNLASGVP 60
 SRFSGSGSGT DFTLTISLQ PEDFATYYCH QYYRSPPTFG GGTKVEIK 108

SEQ ID NO: 49 moltype = AA length = 121
 FEATURE Location/Qualifiers
 REGION 1..121
 note = Variable heavy chain region
 source 1..121
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 49
 EVQLVQSGAE VKKPGATVKI SCKVSGFNIK DYYVHWVQQA PGKGLEWMGR IDVEDDETKY 60
 APKFQGRVTI TADTSTDYAY MELSSLRSED TAVYYCATPI YGAREAWFAY WGQGTLVTVS 120
 S 121

SEQ ID NO: 50 moltype = AA length = 108
 FEATURE Location/Qualifiers
 REGION 1..108
 note = Light chain variable region
 source 1..108
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 50
 DIQMTQSPSS LSASVGDRVT ITCTASSSVS SSYLHWYQQK PGKAPKLLIY RTSNLASGVP 60
 SRFSGSGSGT DFTLTISLQ PEDFATYYCH QYYRSPPTFG GGTKVEIK 108

SEQ ID NO: 51 moltype = AA length = 121
 FEATURE Location/Qualifiers
 REGION 1..121
 note = Variable heavy chain region
 source 1..121
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 51

-continued

QVQLVQSGAE VKKPGASVKV SCKASGYTFT SFGISWVRQA PGQGLEWIGE IYPRSGNTYY 60
 NENFKGRATM TADKSTSTAY MELRSLRSDD TAVYYCARSS GYYGSTPPFY WGQGTLVTVS 120
 S 121

SEQ ID NO: 52 moltype = AA length = 107
 FEATURE Location/Qualifiers
 REGION 1..107
 note = Light chain variable region
 source 1..107
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 52
 DIQMTQSPSS LSASVGDRVT ITCRASQDIS NYLNWYQQKP GKAVKLLIYY TSRLHSGVPS 60
 RFGSGSGSTD YTFTISLQP EDIATYYCQQ GNTLPWTFGG GTKVEIK 107

SEQ ID NO: 53 moltype = AA length = 120
 FEATURE Location/Qualifiers
 REGION 1..120
 note = Variable heavy chain region
 source 1..120
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 53
 QLQLVESGGG VVQPGRLRL SCTVSGFSLT SYGVHWVRQA PGKGLEWVGL IWSGGSTDYS 60
 PAFISRFTIS EDNSKSTVYL QMNSLRAEDT AVYFCARNSY SSGYYAMDYWGQTTVTVSS 120

SEQ ID NO: 54 moltype = AA length = 108
 FEATURE Location/Qualifiers
 REGION 1..108
 note = Light chain variable region
 source 1..108
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 54
 DIQLTQSPSF LSASVGDRVT ITCKASQNVG HAVAWYQQKP GKAPKLLIYS ASNRYTGVPVS 60
 RFGSGSGSTE FTLTISLQP EDFATYYCQQ YSRYPPTYFG GGTKVEIK 108

SEQ ID NO: 55 moltype = AA length = 120
 FEATURE Location/Qualifiers
 REGION 1..120
 note = Variable heavy chain region
 source 1..120
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 55
 QLQLVESGGG VVQPGRLRL SCTVSGFSLT SYGVHWVRQA PGKGLEWVGL IWSGGSTDYS 60
 PAFISRFTIS EDNSKSTVYL QMNSLRAEDT AVYFCARNSY SSGYYAMDYWGQTTVTVSS 120

SEQ ID NO: 56 moltype = AA length = 108
 FEATURE Location/Qualifiers
 REGION 1..108
 note = Light chain variable region
 source 1..108
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 56
 DIQLTQSPSF LSASVGDRVT ITCKASQNVG TAVAWYQQKP GKAPKLLIYS ASNRYTGVPVS 60
 RFGSGSGSTE FTLTISLQP EDFATYYCQQ RSRYPPTYFG GGTKVEIK 108

SEQ ID NO: 57 moltype = AA length = 10
 FEATURE Location/Qualifiers
 REGION 1..10
 note = Variable heavy CDR1 sequence
 source 1..10
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 57
 GFNIKDYVH 10

SEQ ID NO: 58 moltype = AA length = 17
 FEATURE Location/Qualifiers
 REGION 1..17
 note = Variable heavy CDR2 sequence
 source 1..17
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 58
 RIDVHDETK YAPKFQG 17

-continued

SEQ ID NO: 59	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable heavy CDR3 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 59		
PIYGAREAWF AY		12
SEQ ID NO: 60	moltype = AA length = 12	
FEATURE	Location/Qualifiers	
REGION	1..12	
	note = Variable light CDR1 sequence	
source	1..12	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 60		
TASSSVSSSY LH		12
SEQ ID NO: 61	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable light CDR2 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 61		
RTSNLAS		7
SEQ ID NO: 62	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = Variable light CDR3 sequence	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 62		
HQYRSPPT		9
SEQ ID NO: 63	moltype = AA length = 10	
FEATURE	Location/Qualifiers	
REGION	1..10	
	note = Variable heavy CDR1 sequence	
source	1..10	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 63		
GFSLSTYSIS		10
SEQ ID NO: 64	moltype = AA length = 16	
FEATURE	Location/Qualifiers	
REGION	1..16	
	note = Variable heavy CDR2 sequence	
source	1..16	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 64		
IIGDAGGIY ATWAKS		16
SEQ ID NO: 65	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Variable heavy CDR3 sequence	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 65		
DGTAFDI		7
SEQ ID NO: 66	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Variable light CDR1 sequence	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 66		
QASQSIYSKL G		11

-continued

SEQ ID NO: 67	moltype = AA	length = 7	
FEATURE	Location/Qualifiers		
REGION	1..7		
source	note = Variable light CDR2 sequence		
	1..7		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 67			
RASTLAS	7		
SEQ ID NO: 68	moltype = AA	length = 12	
FEATURE	Location/Qualifiers		
REGION	1..12		
source	note = Variable light CDR3 sequence		
	1..12		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 68			
QQDYSYINVD NI	12		
SEQ ID NO: 69	moltype = AA	length = 121	
FEATURE	Location/Qualifiers		
REGION	1..121		
source	note = Variable heavy chain region		
	1..121		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 69			
EVQLVQSGAE VKKPGATVKI	SCKVSGFNIK	DYYVHWVQQA	PGKGLEWMGR IDVHDDETKY 60
APKFGQGRVTI TADTSTDYAY	MELSSLRSED	TAVYYCATPI	YGAREAWFAY WGQGTLVTVS 120
S			121
SEQ ID NO: 70	moltype = AA	length = 108	
FEATURE	Location/Qualifiers		
REGION	1..108		
source	note = Light chain variable region		
	1..108		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 70			
DIQMTQSPSS LSASVGDRVT	ITCTASSSVS	SSYLHWYQQK	PGKAPKLLIY RTSNLASGVP 60
SRFSGSGSGT DFTLTISLQ	PEDFATYYCH	QYYRSPPTFG	GGTKVEIK 108
SEQ ID NO: 71	moltype = AA	length = 114	
FEATURE	Location/Qualifiers		
REGION	1..114		
source	note = Variable heavy chain region		
	1..114		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 71			
QSVKESEGL FKPTDTLTLT	CTVSGFSLST	YSISWVRQAP	GNGLEWIGII GDAGGIYAT 60
WAKRSRSTITR STALNTVTLK	MTGLTAADTA	TYFCARDGTA	FDIWGPGLTV TVSS 114
SEQ ID NO: 72	moltype = AA	length = 110	
FEATURE	Location/Qualifiers		
REGION	1..110		
source	note = Light chain variable region		
	1..110		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 72			
AYDMTQTPAS VEVVVGTVT	IKQASQSIY	SKLGWYQQKP	GQPPKLLIYR ASTLASGVSS 60
RFKGSGSGTE YTLTISGVQC	DDAATYYCQ	DYSYINVNDNI	FGGGTEVVVK 110
SEQ ID NO: 73	moltype = AA	length = 114	
FEATURE	Location/Qualifiers		
REGION	1..114		
source	note = Variable heavy chain region		
	1..114		
	mol_type = protein		
	organism = synthetic construct		
SEQUENCE: 73			
QSVKESEGL FKPTDTLTLT	CTVSGFSLST	YSISWVRQAP	GNGLEWIGII GDAGGIYAT 60
WAKRSRSTITR STALNTVTLK	MTGLTAADTA	TYFCARDGTA	FDIWGPGLTV TVSS 114
SEQ ID NO: 74	moltype = AA	length = 110	
FEATURE	Location/Qualifiers		

-continued

```

REGION                1..110
                        note = Light chain variable region
source                1..110
                        mol_type = protein
                        organism = synthetic construct

SEQUENCE: 74
AYDMTQTPAS VEVVVGTVT IKQASQSIY SKLGWYQQKP GQPPKLLIYR ASTLASGVSS 60
RFKGSQSGTE YTLTISGVQA DDAATYYCQQ DYSYINVDNI FGGGTEVVVK      110

SEQ ID NO: 75          moltype = AA length = 10
FEATURE               Location/Qualifiers
REGION                1..10
                        note = consensus heavy chain CDR1
VARIANT               4
                        note = Xaa = Ile or Val
VARIANT               7
                        note = Xaa = Ser or Tyr
VARIANT               8
                        note = Xaa = Phe or Tyr
VARIANT               9
                        note = Xaa = Ile or Val
source                1..10
                        mol_type = protein
                        organism = synthetic construct

SEQUENCE: 75
GFNXKDXXXH                                10

SEQ ID NO: 76          moltype = AA length = 17
FEATURE               Location/Qualifiers
REGION                1..17
                        note = Consensus heavy chain CDR2
VARIANT               4
                        note = Xaa = Val or Pro
source                1..17
                        mol_type = protein
                        organism = synthetic construct

SEQUENCE: 76
RIDXEDDETK YAPKFQG                                17

SEQ ID NO: 77          moltype = length =
SEQUENCE: 77
000

SEQ ID NO: 78          moltype = AA length = 7
FEATURE               Location/Qualifiers
REGION                1..7
                        note = Consensus light chain CDR2
VARIANT               1
                        note = Xaa = Arg or Ser
source                1..7
                        mol_type = protein
                        organism = synthetic construct

SEQUENCE: 78
XTSNLAS                                7

SEQ ID NO: 79          moltype = length =
SEQUENCE: 79
000

SEQ ID NO: 80          moltype = AA length = 12
FEATURE               Location/Qualifiers
REGION                1..12
                        note = Consensus heavy chain CDR3
VARIANT               1
                        note = Xaa = Ser or Thr
VARIANT               2
                        note = Xaa = Gly, Lys, Leu, Ser, Thr or Val
VARIANT               4
                        note = Xaa = Leu or Tyr
VARIANT               6
                        note = Xaa = Ala, Gly or Ser
VARIANT               10
                        note = Xaa = Ala, Phe, His, Leu, Asn, Gln, Ser, Thr or Tyr
VARIANT               11
                        note = Xaa = Ala, Ile, Pro, Thr or Val
VARIANT               12
                        note = Xaa = Ala, Glu, Gly, His, Ile, Lys, Leu, Asn, Gln,
                        Arg, Ser, Thr or Tyr
source                1..12

```

-continued

```

mol_type = protein
organism = synthetic construct

SEQUENCE: 80
XXGXYSSTPX XX                                     12

SEQ ID NO: 81
FEATURE
REGION
VARIANT
VARIANT
VARIANT
VARIANT
VARIANT
VARIANT
VARIANT
source
mol_type = AA length = 12
Location/Qualifiers
1..12
note = Consensus heavy chain CDR3
2
note = Xaa = Phe, Gly, Ile, Ser, Thr, Val or tyr
3
note = Xaa = His, Phe or Tyr
4
note = Xaa = Ala, Phe, Gly or Ser
5
note = Xaa = Phe, Ser or Tyr
9
note = Xaa = Ala, Gly, His, Ser, Pro or Thr
10
note = Xaa = Met or Lys
12
note = Xaa = Asp, Gly, His, Ile, Ala, Val, Leu, Arg, Ser or
Tyr
1..12
mol_type = protein
organism = synthetic construct

SEQUENCE: 81
NXXXXGYXX DX                                     12

SEQ ID NO: 82
FEATURE
REGION
VARIANT
source
mol_type = AA length = 11
Location/Qualifiers
1..11
note = Consensus light chain CDR1
8
note = Xaa = Ala, Gly, His, Asn, Gln, Arg, Ser or Thr
1..11
mol_type = protein
organism = synthetic construct

SEQUENCE: 82
KASQNVGXAV A                                     11

SEQ ID NO: 83
FEATURE
REGION
VARIANT
VARIANT
VARIANT
VARIANT
VARIANT
source
mol_type = AA length = 10
Location/Qualifiers
1..10
note = Consensus light chain CDR3
3
note = Xaa = Phe, His, Lys, Arg or tyr
4
note = Xaa = Ala, Ile, Leu, Asn, Arg, Ser, Thr or Val
5
note = Xaa = Phe, Gly, Ile, Lys, Leu, Asn, Arg, Ser, Thr,
Val or Tyr
6
note = Xaa = Ala, Phe, Gly, Ile, Gln, Arg, Ser, Thr or Tyr
1..10
mol_type = protein
organism = synthetic construct

SEQUENCE: 83
QXXXXXPYT                                         10

SEQ ID NO: 84
FEATURE
source
mol_type = AA length = 931
Location/Qualifiers
1..931
mol_type = protein
organism = Homo sapiens

SEQUENCE: 84
MDMFLPLTWVF LALYFSRHQV RGQPDPPCGG RLNSKDAGYI TSPGYPDQYP SHQNCWEIVY 60
APEPNQKIVL NFNPHFEIEK HDCKYDFIEI RDGDSSESADL LGKHCGNIAP PTIISGSMML 120
YIRFTSDYAR QGAGFSLRYE IFKTGSEDCS KNFTSPNGTI ESPGFPEKYP HNLDCFTTIL 180
AKPKMEIILQ FLIFDLEHDP LQVGECDCKY DWLDIWDGIP HVGPLIGKYC GTKTPSELRS 240
STGILSLTFH TDMAVAKDGF SARYYLVHQE PLENFQCNPV LGMESGRIAN EQISASSTYS 300
DGRWTPQQSR LHGDDNGWTP NLDSNKEYLQ VDLRFLTMLT AIATQGAISR ETQNGYYVKS 360
YKLEVSTNGE DWMVYRHGKN HKVFQANND A TEVVLNKLHA PLLTRFVRIR PQTWHSGIAL 420
RLELFGCRVT DAPCSNMLGM LSGLIADSI SASSTQEYLW SPSAARLVSS RSGWFPRIPQ 480
AQPGEELQV DLGTPKTVKG VIIQGARGGD SITAVEARAF VRKFKVSYSL NGKDWEYIQD 540
PRTQQPKLFE GNMHYDTPDI RRFDPIPAQY VRVYPERWSP AGIGMRLEVL GCDWTDKPT 600
VETLGPTVKS EETTPPYTE EEATECGENC SFEDDKDLQL PSGFNCNFD LEEP CGWMYD 660
HAKWLRTTWA SSSSPNDRTE PDDRNFLRLQ SDSQREGQYA RLISPPVHLP RSPVCMFQY 720

```

QATGGRGVAL	QVVRASQES	KLLWVIREQD	GGEWKHGRII	LPSYDMEYQI	VFEGVIGKGR	780
SGETAIDDIR	ISTDVPLENC	MEPISAFAGE	NFKVDIPEIH	EREGYDEEDL	DEYEDVWSNS	800
SSATSSGSGA	STDKEKSWL	TLDPILITII	AMSSLGVLLG	ATCAGLLLYC	TCSYSGLSSR	840
SCTTLENYNF	ELYDGLKHWY	KMNHKCKCSE	A			931

```
SEQ ID NO: 85      moltype = AA  length = 926
FEATURE            Location/Qualifiers
source             1..926
                  mol_type = protein
                  organism = Homo sapiens
```

SEQUENCE: 85		Organism = Homo sapiens		
MDMPFLTWVF	LALYFSRHQV	RGGQDPDPCG	RLNSKDAYI	TSPGYPLQDYP SHQNC EWIV 60
APBNQKIVL	LNPNHFIEIK	HDCKYDFIEI	RDGDESADL	LGKHCGNIA PPTIISG SML 120
YIKFTSDYAR	QGAGFSLRYE	IFKTGSEDCS	KNFTSPNGTI	ESPGFPEKYP HNL DCTFTIL 180
AKPKMEIILQ	FLI PDLHDKP	LQVGEGHQCE	DWLDIWDGIP	HVGLSILGKYC GTKTPSELRS 240
STGLSLTLFH	TDMAVAKDGF	SARYLVHCKY	PLENQCNPVP	LMGESGRIAN EGI SASSTYS 300
DGRVTPQOSR	LHGDDNGWTP	NLDSNKEYLK	VDLRFLTMLT	AIA TQGAISR ETQNGYVVK 360
YKLEVSTNGE	DWMVYRHGKN	HKVFQAMNDA	TEVVNLKQLH	PLLTRFVRIR PQTWHS GIAL 420
RLEFLGCRVT	DAPCSNMLMG	LSGLIADSPQ	SASSVTEYLW	SPSAARLVSS RQGWFPRI PQ 480
AQPGEEWLQV	DLGTPTKYTK	VIIQAGARGD	SITAVEARAF	VRKFVKSYSL NGKDWEYIQD 540
PRTQQPKLFE	GNMHYDTPDI	RRFDPIPAQY	VRVYPERWSP	AGIGMRLEVL GCDTWESKPT 600
VETLGLPTVKS	EBETTPPYTE	EEATECGECS	SFEDDKLQL	PSGFNCNPDF LEEPCGWMDY 660
HAKWLVTTWA	SSSSPNDRPT	PPDRNFLRLQ	SDSQREGQYA	RLTSIPVHLV RSPVCMFEQY 720
QATGGRGVAL	QVVRASEGS	KLKLVIREQD	GEWKHKGRII	LPSYDMEYQI VFEVGIGKGR 780
SGEIAIDDIR	ISTDVPLENC	MEPISAFAVD	IPEIHEREGY	EDEIDDEYEV DWSNSSATS 840
GSASPSTDKS	KSWLYTLDP I	LITTIAMSSL	GVLLGATCAG	LLLYCTCSYS GLSSRSCTLL 900
ENYNFELYDQ	LKHKKVMNHO	KCCSEA		

```
SEQ ID NO: 86      moltype = AA  length = 909
FEATURE            Location/Qualifiers
source             1..909
                  mol_type = protein
                  organism = Homo sapiens
```

SEQUENCE: 86	SEQUENCE: 87	SEQUENCE: 88	SEQUENCE: 89
MDMPFLTWVF	LALYFSRHQV	RQGPDPCCGG	RLNSKDAGYI
APENQKIVL	LNPNHFIEIK	HDCKYDFIEI	RDGDESADL
YIKFTSDYAR	QQAGFSLRYE	IFKTSGEDCS	KNFTSPNGTI
AKPKMEIILQ	FLIADLEHPD	LQVGEGEDCK	DWLDIWDGPI
STGLSLTFPH	TDMAVAKDFG	SARYLVHQEJ	PLENFQCNVP
DGRWTPQOSR	LHGDDNGWTP	NLDSNKEYLK	VDLRFLTMLT
YKLEVSTNGE	DWMVYRHGKN	HKVFQAMNDA	TEVVNLKQLH
RLELFGCRVT	DAPCSNMLMG	LSGLIADSPS	SAASARLVSS
AQPGEEWLQV	DLGTPKTYKG	VITQGARGGD	SITAVEARAF
PRTQQPKLFE	GNMHYDTPDI	RFFDPIPAQY	VRVYPERWSP
VETLGLPTVKS	BETTTPTYPE	EEATECEGCS	SPEDDKLQL
HAKWRLTTWA	SSSSPNDRFT	PPDRNFLRLQ	SDSQREGGYA
QATGGRGVAL	QVVRASEGS	KLWLVIREQD	GEWKKHORIA
SGEIAIDDIR	ISTDVPLENC	MEPISAFADE	YEVDSNSSSS
DLPIITIAM	SSLGVLLGAT	CAGLLLYCTC	SYSLGSSRSC
NHOKCCSEA			

```
SEQ ID NO: 87      moltype = AA  length = 906
FEATURE            Location/Qualifiers
source             1..906
                  mol_type = protein
                  organism = Homo sapiens
```

[illegible]

```
SEQ ID NO: 88      moltype = AA  length = 901
FEATURE            Location/Qualifiers
source             1..901
                  mol_type = protein
                  organism = Homo sapiens
```

-continued

SEQUENCE: 88

MDMFPLTWVF	LALYFSRHQV	RGQDPDPPCGG	RLNSKDAGYI	TSPGYPDYYP	SHQNCWEIVY	60
APEPNQKIVL	NFNPHFEIEK	HDCKYDFIEI	RDGDSSESADL	LGKHCIGNIAP	PTIISSGSML	120
YIKFTSDYAR	QGAGFSLRYE	IFKTGSEDCS	KNFTSPNGTI	ESPGFPEKYP	HNLDCFTTIL	180
AKPKMEIILQ	FLIFDLEHDP	LQVGECDCKY	DWLDIWDGIP	HVGPLIGKYC	GTKTPSELRS	240
STGILSLTFH	TDMAVAKDGF	SARYYLVHQE	PLENFQCNVP	LGMESEGRIAN	EQISASSTYS	300
DGRWTPQQSR	LHGDDNGWTP	NLDSNKEYLQ	VDLRFLLMTL	AIATQGAISR	ETQNGYVVK	360
YKLEVSTNGE	DWMVYRHGKN	HKVFQANMDA	TEVVLNKLHA	PLLTRFVRIR	PQTHSGIAL	420
RLELPGCRVT	DAPCSNMLGM	LSGLIADSI	SASSTQEYLW	SPSAARLVSS	RSWGFPRIPQ	480
AQPGEEWLQV	DLGTPKTVKG	VIIQGARGGD	SITAVEARAF	VRKFKVSYSL	NGKDEYIQD	540
PRTQQPKLFE	GNMHYDTPDI	RRFDPIPAQY	VRVYPERWSP	AGIGMRLEVL	GCDWTDKSKPT	600
VETLGPTVKS	EETTPYPTPE	EEATECGENC	SFEDDKDLQL	PSGFNCNFD	LEEPCGWMYD	660
HAKWLRTTWA	SSSPNDRTF	PDDRNFLRLQ	SDSQREGQYA	RLISPPVHLR	RSPVCMFQY	720
QATGGRGVAL	QVVREASQES	KLLWVIREDO	GGEWKHGRII	LPSYDMEYQI	VFEGVIGKGR	780
SGEIAIDDIR	ISTDVPLENC	MEPISAFAGG	TLLPGTEPTV	DTVPMQPIPA	YWYYVMAAGG	840
AVLVLVSVSL	ALVLHYHRFR	YAAKKTDSHI	TYKTSHTYNG	APLAVEPTLT	IKLEQDRGSH	900
C						901

SEQ ID NO: 89 moltype = AA length = 555
 FEATURE Location/Qualifiers
 source 1..555
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 89

MDMFPLTWVF	LALYFSRHQV	RGQDPDPPCGG	RLNSKDAGYI	TSPGYPDYYP	SHQNCWEIVY	60
APEPNQKIVL	NFNPHFEIEK	HDCKYDFIEI	RDGDSSESADL	LGKHCIGNIAP	PTIISSGSML	120
YIKFTSDYAR	QGAGFSLRYE	IFKTGSEDCS	KNFTSPNGTI	ESPGFPEKYP	HNLDCFTTIL	180
AKPKMEIILQ	FLIFDLEHDP	LQVGECDCKY	DWLDIWDGIP	HVGPLIGKYC	GTKTPSELRS	240
STGILSLTFH	TDMAVAKDGF	SARYYLVHQE	PLENFQCNVP	LGMESEGRIAN	EQISASSTYS	300
DGRWTPQQSR	LHGDDNGWTP	NLDSNKEYLQ	VDLRFLLMTL	AIATQGAISR	ETQNGYVVK	360
YKLEVSTNGE	DWMVYRHGKN	HKVFQANMDA	TEVVLNKLHA	PLLTRFVRIR	PQTHSGIAL	420
RLELPGCRVT	DAPCSNMLGM	LSGLIADSI	SASSTQEYLW	SPSAARLVSS	RSWGFPRIPQ	480
AQPGEEWLQV	DLGTPKTVKG	VIIQGARGGD	SITAVEARAF	VRKFKVSYSL	NGKDEYIQD	540
PRTQQPKVGC	SWRPL					555

SEQ ID NO: 90 moltype = AA length = 904
 FEATURE Location/Qualifiers
 source 1..904
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 90

QPDPPCGGRL	NSKDAGYITS	PGYPQDYPSH	QNCWEIVYAP	EPNQKIVLNF	NPHFEIEKHD	60
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT	IISSGSMLYI	KFTSDYARQG	AGFSLRYEIF	120
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ	180
VGECDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	240
RYLVHQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL	300
DSNKEYLQVD	LRFLTMLTAI	ATQGAISR	QNGYVVKSYK	LEVSTNGEDW	MVYRHGKNHK	360
VFQANNDAT	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	PCSNMLGMLS	420
GLIADSI	SSTQEYLWSP	SAARLVSSRS	GWFPRIPQAQ	PGEWLQVDL	GTPKTVKGVI	480
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKLFEGN	MHYDTPDIRR	540
FDPIPAQYVR	VYPERWSPAG	IGMRLEVLGC	DWTDKSKPTE	TLGPTVKSEE	TTTPYPTTEE	600
ATECGENC	EDDKDLQLPS	GFNCNFDLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTFPD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFYQQA	TGGRGVALQV	VREASQESKL	720
LWVIREDDQ	EWKHGRIILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780
PISAFADVTP	BIHEREGYED	EIDDEYEVWD	SNSSSATS	GAPSTDKEKS	WLYTLDPILI	840
TIAMSSSLGV	LLGATCAGLL	LYCTCSYSG	SSRSCTTLEN	YNFELYDGLK	HKVKMNHQKC	900
CSEA						904

SEQ ID NO: 91 moltype = AA length = 879
 FEATURE Location/Qualifiers
 source 1..879
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 91

QPDPPCGGRL	NSKDAGYITS	PGYPQDYPSH	QNCWEIVYAP	EPNQKIVLNF	NPHFEIEKHD	60
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT	IISSGSMLYI	KFTSDYARQG	AGFSLRYEIF	120
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ	180
VGECDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	240
RYLVHQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL	300
DSNKEYLQVD	LRFLTMLTAI	ATQGAISR	QNGYVVKSYK	LEVSTNGEDW	MVYRHGKNHK	360
VFQANNDAT	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	PCSNMLGMLS	420
GLIADSI	SSTQEYLWSP	SAARLVSSRS	GWFPRIPQAQ	PGEWLQVDL	GTPKTVKGVI	480
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKLFEGN	MHYDTPDIRR	540
FDPIPAQYVR	VYPERWSPAG	IGMRLEVLGC	DWTDKSKPTE	TLGPTVKSEE	TTTPYPTTEE	600
ATECGENC	EDDKDLQLPS	GFNCNFDLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTFPD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFYQQA	TGGRGVALQV	VREASQESKL	720
LWVIREDDQ	EWKHGRIILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780
PISAFAGGTL	LPTEPTVDT	VPMQPIPAYW	YVMAAGGAV	LVLVSVLAL	VLHYHRFRYA	840
AKKTDHSITY	KTSHYTINGAP	LAVEPTLTIK	LEQDRGSHC			879

-continued

SEQ ID NO: 92 moltype = AA length = 533
 FEATURE Location/Qualifiers
 source 1..533
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 92

QDPDPCGGRL	NSKDAGYITS	PGYPQDYPSH	QNCWEIVYAP	EPNQKIVLNF	NPHFEIEKHD	60
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT	IISGSMLEYI	KFTSDYARQG	AGFSLRYEIF	120
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPQL	180
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFSA	240
RYYLHVQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYS DG	RWTPQQSRLH	GDDNGWTPNL	300
DSNKEYLQVD	LRFLTMLTAI	ATQGAISRET	QNGYVVKSYK	LEVSTNGEDW	MVVRHGKNHK	360
VFQANNDAT	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	PCSNMLGMLS	420
GLIADSQISA	SSTQEYLWSP	SAARLVSSRS	GWFPRIQQAQ	PGEEWLQVDL	GTPKTVKGV	480
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKVGC	CSW RPL	533

SEQ ID NO: 93 moltype = AA length = 114
 FEATURE Location/Qualifiers
 source 1..114
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 93

CGRLNSKDA	GYITSPGYPO	DYPHQNCWE	IVYAPEPNQK	IVLNFNPHFE	IEKHDCKYDF	60
IEIRDGDS	ADLLGKHCGN	IAPPTIISG	SMLYIKFTSD	YARQGAGFSL	RYEI	114

SEQ ID NO: 94 moltype = AA length = 117
 FEATURE Location/Qualifiers
 source 1..117
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 94

CSKNFTSPNG	TIESPGFPEK	YPHNLDCTFT	ILAKPKMEII	LQFLIFDLEH	DPLQVGEGDC	60
KYDWLDIWDG	IPHVGPLIGK	YCGTKTPSEL	RSSTGILSLT	FHTDMAVAKD	GFSARYY	117

SEQ ID NO: 95 moltype = AA length = 147
 FEATURE Location/Qualifiers
 source 1..147
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 95

PLGMESGRIA	NEQISASSTY	SDGRWTPQQS	RLHGDDNGWT	PNLDSNKEYL	QVDLRFLTML	60
TAIATQGAIS	RETQNGYVVK	SYKLEVSTNG	EDWMVYRHGK	NHKVFQANND	ATEVVNLKLH	120
APLLTRFVRI	RPQTWHSGLA	LRLELFG				147

SEQ ID NO: 96 moltype = AA length = 154
 FEATURE Location/Qualifiers
 source 1..154
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 96

LGMLSGLIAD	SQISASSTQE	YLWSPSAARL	VSSRSGWFPR	IPQAQPGEEW	LQVDLGTPTK	60
VKGVIIQGAR	GGDSITAVEA	RAFVRKFKVS	YSLNGKDWEY	IQDPRTQQPK	LFEGNMHYDT	120
PDIRRFDPIP	AQYVRVYPER	WSPAGIGMRL	EVLG			154

SEQ ID NO: 97 moltype = AA length = 154
 FEATURE Location/Qualifiers
 source 1..154
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 97

PSGFNCNPDF	LEEPCGMYD	HAKWLRTTWA	SSSSPNDRTF	PDDRNFLRLQ	SDSQREGQYA	60
RLISPPVHLP	RSPVCMFQY	QATGGRGVAL	QVVREASQES	KLLWVIREQD	GGEWKHGRII	120
LPSYDMEYQI	VFEGVIGKGR	SGEIAIDDIR	ISTD			154

SEQ ID NO: 98 moltype = AA length = 243
 FEATURE Location/Qualifiers
 source 1..243
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 98

QDPDPCGGRL	NSKDAGYITS	PGYPQDYPSH	QNCWEIVYAP	EPNQKIVLNF	NPHFEIEKHD	60
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT	IISGSMLEYI	KFTSDYARQG	AGFSLRYEIF	120
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPQL	180
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFSA	240
RYY						243

SEQ ID NO: 99 moltype = AA length = 278
 FEATURE Location/Qualifiers

-continued

```

source                1..278
                      mol_type = protein
                      organism = Homo sapiens

SEQUENCE: 99
CSKNFTSPNG TIESPGFPEK YPHNLDCTFT ILAKPKMEII LQFLIFDLEH DPLQVGEGDC 60
KYDWLDIWDG IPHVGPLIGK YCGTKTPSEL RSSTGILSLT FHTDMAVAKD GFSARYLVH 120
QEPLNFQCN VPLGMESGRI ANEQISASST YSDGRWTPQQ SRLHGDDNGW TPNLDSNKEY 180
LQVDLRFLTM LTAIATQGA SRETQNGYYV KSYKLEVSTN GEDWMVYRHG KNHKVFQANN 240
DATEVVLNKL HAPLLTRFVR IRPQTWHSGL ALRLELFG 278

SEQ ID NO: 100        moltype = AA length = 404
FEATURE              Location/Qualifiers
source                1..404
                      mol_type = protein
                      organism = Homo sapiens

SEQUENCE: 100
QPDPPCGGRL NSKDAYGITS PGYPQDYPSH QNCEWIVYAP EPNQKIVLNF NPHFEIEKHD 60
CKYDFIEIRD GDSADLLG KHCGNIAPPT IISGSMYI KFTSDYARQG AGFSLRYEIF 120
KTGSEDCSKN FTSPNGTIES PGFPEKYPHN LDCTFTILAK PKMEIILQFL IFDLEHDPLQ 180
VGEGDCKYDW LDIWDGIPHV GPLIGKYCGT KTPSELRSST GILSLTFHTD MAVAKDGFS 240
RYYLVHQEPL ENFQCNVPLG MESGRIANEQ ISASSTYSDG RWTPQQSRLH GDDNGWTPNL 300
DSNKEYLQVD LRFLTMLTAI ATQGAISRET QNGYVVKSYK LEVSTNGEDW MVYRHGKNHK 360
VFQANNDATE VVLNKLHAPL LTRFVRIRPQ TWHSGIALRL ELFG 404

SEQ ID NO: 101        moltype = AA length = 573
FEATURE              Location/Qualifiers
source                1..573
                      mol_type = protein
                      organism = Homo sapiens

SEQUENCE: 101
QPDPPCGGRL NSKDAYGITS PGYPQDYPSH QNCEWIVYAP EPNQKIVLNF NPHFEIEKHD 60
CKYDFIEIRD GDSADLLG KHCGNIAPPT IISGSMYI KFTSDYARQG AGFSLRYEIF 120
KTGSEDCSKN FTSPNGTIES PGFPEKYPHN LDCTFTILAK PKMEIILQFL IFDLEHDPLQ 180
VGEGDCKYDW LDIWDGIPHV GPLIGKYCGT KTPSELRSST GILSLTFHTD MAVAKDGFS 240
RYYLVHQEPL ENFQCNVPLG MESGRIANEQ ISASSTYSDG RWTPQQSRLH GDDNGWTPNL 300
DSNKEYLQVD LRFLTMLTAI ATQGAISRET QNGYVVKSYK LEVSTNGEDW MVYRHGKNHK 360
VFQANNDATE VVLNKLHAPL LTRFVRIRPQ TWHSGIALRL ELFGCRVTD PCSNMLGMLS 420
GLIADSQISA SSTQEYLWSP SAARLVSSRS GWFPRIPQAQ PGEWLQVDLT GTPKTVKGVI 480
IQARGGDSI TAVEARAFVR KFKVSYSLNG KDWEYIQDPR TQPKLFEGNMH MYDTPDIRR 540
FDPIPAQYVR VYPERWSPAG IGMRLVLGCDW DWT 573

SEQ ID NO: 102        moltype = AA length = 451
FEATURE              Location/Qualifiers
source                1..451
                      mol_type = protein
                      organism = Homo sapiens

SEQUENCE: 102
GSEDCSKNFT SPNGTIESPG FPEKYPHNLD CTFTILAKPK MEIILQFLIF DLEHDPLQVG 60
EGDCKYDWLD IWDGIPHVGP LIGKYCGTK PSELRSSTGI LSLTFHTDMA VAKDGFSARY 120
YLVHQEPLEN FQCNVPLGME SGRIANEQIS ASSTYSDGRW TPQQSRLHGD DNGWTPNLDS 180
NKEYLQVDLR FLTMLTAIAT QGAISRETQN GYVVKSYKLE VSTNGEDWMV YRHGKNHKVF 240
QANNDATEV LNKHLAPLLT RFVRIRPQTH HSGIALRLEL FGCRVTDAPC SNMLGMLSG 300
IADSQISASS TQEYLWSPSA ARLVSSRSRW FPRIPQAQPG EEWLQVDLGT PKTVKGVI 360
GARGGDSITA VEARAFVRKF KVSYSNLGKD WEYIQDPRTO QPKLFEGNMH YDTPDIRRFD 420
PIPAQYVRVY PERWSPAGIG MRLEVLGCDW T 451

SEQ ID NO: 103        moltype = AA length = 320
FEATURE              Location/Qualifiers
source                1..320
                      mol_type = protein
                      organism = Homo sapiens

SEQUENCE: 103
QCNVPLGMES GRIANEQISA SSTYSDGRWT PQQSRLHGDD NGWTPNLDSN KEYLQVDLRF 60
LTMLTAIATQ GAISRETQNG YVVKSYKLEV STNGEDWMVY RHGKNHKVFQ ANNDATEVVL 120
NKLHAPLLTR FVRIRPQTH SGIALRLELFG GCRVTDAPCS NMLGMLSGLI ADSQISASST 180
QEYLWSPSAA RLVSSRSRGW PRIPQAQPG EWLQVDLGT KTVKGVIQG ARGGDSITAV 240
EARAFVRKFK VSYSLNGKDW EYIQDPRTOQ PKLFEGNMHY DTPDIRRFD IPQYVRVYP 300
ERWSPAGIGM RLEVLGCDWT 320

SEQ ID NO: 104        moltype = AA length = 1060
FEATURE              Location/Qualifiers
REGION              1..1060
                      note = NRP2 - Fc fusion Construct
source              1..1060
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 104
QPDPPCGGRL NSKDAYGITS PGYPQDYPSH QNCEWIVYAP EPNQKIVLNF NPHFEIEKHD 60
CKYDFIEIRD GDSADLLG KHCGNIAPPT IISGSMYI KFTSDYARQG AGFSLRYEIF 120

```

-continued

KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ	180
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	240
RYLVHQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL	300
DSNKEYLQVD	LRFLTMLTAI	ATQGAISRET	QNGYVVKSYK	LEVSTNGEDW	MVYRHGKNHK	360
VQANNDATE	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	PCSNMLGMLS	420
GLIADSQISA	SSTQEYLWSP	SAARLVSSRS	GWFPRIPOAQ	PGEWLQVDL	GTPKTVKGV	480
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQPKLFEGN	MHYDTPDIR	540
FPIPAQYVR	VYPERWSPAG	IGMRLEVLC	DWTDCKPTVE	TLGPTVKSEE	TTTPYPTTEE	600
ATECGENCFS	EDDKDLQLPS	GFNCNFDLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTPFD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFQYQA	TGGRGVALQV	VREASQESKL	720
LWVIREDDQG	EWKHGRIILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780
PISAPAVDIP	EIHEREGYED	EIDDEYEVDW	SNSSSATSGS	GAPSTDKEKS	WLYDKTHTCP	840
PCPAPELLGG	PSVFLFPPKP	KDTLMISRTP	EVTGVVDVVS	HEDPEVKFNW	YVDGVEVHNA	900
KTKPREEQYN	STYRVVSVLT	VLHQDWLNGK	EYKCKVSNKA	LPAPIEKTIS	KAKGQPREPQ	960
VYTLPPSRDE	LTKNQVSLTC	LVKGFYPSDI	AVEWESNGQP	ENNYKTTTPV	LSDSGSFFLY	1020
SKLTVDKSRW	QQGNVFCSCV	MHEALHNHYT	QKSLSLSPGK			1060

SEQ ID NO: 105 moltype = AA length = 678
 FEATURE Location/Qualifiers
 REGION 1..678
 note = NRP2 - Fc fusion construct
 source 1..678
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 105

GSEDCSKNPT	SPNGTIESPG	FPEKYPHNLD	CTFTILAKPK	MEIILQFLIF	DLEHDPLQVG	60
EGDCKYDWLD	IWDGIPHVGP	LIGKYCGTKT	PSELRSSTGI	LSLTFHTDMA	VAKDGFSA	120
YLVHQEPLN	QCNVPLGME	SGRIANEQIS	ASSTYSDGRW	TPQQSRLHGD	DNGWTPNLDS	180
NKEYLQVDLR	FLTMLTAIAT	QGAISRETQN	GYVVKSYKLE	VSTNGEDWMV	YRHGKNHKVF	240
QANNDATEV	VVLNKLHAPLLT	RFVRIRPQTW	HSGIALRLEL	FGCRVTDAPC	SNMLGMLSGL	300
IADSQISASS	TQEYLWSPSA	ARLVSSRSGW	FPRIPOAQPG	EEWLQVDLGT	PKTVKGVIIQ	360
GARGGDSITA	VEARAFVRKFP	KVSYSLNGKD	WEYIQDPRTQ	QPKLFEGNMH	YDTPDIRRFD	420
PPIPAQYVRVY	PERWSPAGIG	MRLEVLCGDW	TDKTHTCPPC	PAPELLGGPS	VFLFPPKPKD	480
TLMISRTPEV	TCVVVDVSHE	DPEVKFNWYV	DGVEVHNAKT	KPREEQYNST	YRVVSVLTVL	540
HQDWLNGKEY	KCKVSNKALP	APIEKTISKA	KGQPREPQVY	TLPPSRDEL	KNQVSLTCLV	600
KGFYPSDIAV	EWESNGQFEN	NYKTTTPVLD	SDGSFFLYSK	LTVDKSRWQQ	GNVFCSCVMH	660
EALHNHYTQK	SLSLSPGK					678

SEQ ID NO: 106 moltype = AA length = 357
 FEATURE Location/Qualifiers
 source 1..357
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 106

LGMLSGLIAD	SQISASSTQE	YLWSPSAARL	VSSRSGWFPR	IPQAQPGEEW	LQVDLGTPTK	60
VKGVIQGAR	GGDSITAVEA	RAFVRKFKVS	YSLNGKDWEY	IQDPRTQOPK	LFEGNMHYDT	120
PDIRRFDPIP	AQYVRVYPER	WSPAGIGMRL	EVLGCDWTD	KPTVETLGPT	VKSEETTPY	180
PTEEATECG	ENCSFEDDKD	LQLPSGFNCN	FDLEEPCCGW	MYDHAKWLRT	TWASSSSPND	240
RTFPDDRNL	RLQSDSQREG	QYARLISPPV	HLPRSPVCME	FQYQATGGRG	VALQVREAS	300
QESKLLWVIR	EDQGGGEWKH	RIILPSYDME	YQIVFEGVIG	KGRSGEIAID	DIRISTD	357

SEQ ID NO: 107 moltype = AA length = 519
 FEATURE Location/Qualifiers
 source 1..519
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 107

QCNVPLGMES	GRIANEQISA	SSTYSDGRWT	PQQSRLHGDD	NGWTPNLDSN	KEYLQVDLRF	60
LTMLTAIATQ	GAISRETQNG	YVVKSYKLEV	STNGEDWMVY	RHGKNHKVFQ	ANNDATEVVL	120
NKLHAPLLTR	FVRIRPQTW	SGIALRLELF	GCRVTDAPCS	NMLGMLSGLI	ADSQISASST	180
QEYLWSPSAA	RLVSSRSWGF	PRIPQAQPG	EWLQVDLGT	KTVKGVIIQ	ARGGDSITAV	240
EARAFVRKFP	VSYSNLNGKD	EYIQDPRTQQ	PKLFEGNMHY	DTPDIRRFD	IPAQYVRVYP	300
ERWSPAGIGM	RLEVLCGDWT	DSKPTVETLG	PTVKSEETTT	PYPTEEEATE	CGENCSEFDD	360
KDLQLPSGFN	CNFDLFLEPC	GWMYDHAKWL	RTTWASSSSP	NDRTFPDDRN	FLRLQSDSQR	420
EGQYARLISP	PVHLPRSPVC	MEFQYQATGG	RGVALQVVRE	ASQESKLLWV	IREDDQGGWK	480
HGRIILPSYD	MEYQIVFEGV	IGKRSGEIA	IDDIRISTD			519

SEQ ID NO: 108 moltype = AA length = 654
 FEATURE Location/Qualifiers
 source 1..654
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 108

CSKNFTSPNG	TIESPGFPEK	YPHNLDCTFT	ILAKPKMEII	LQFLIFDLEH	DPLQVGEGDC	60
KYDWLDIWDG	IPHVGLIGK	YCGTKTPSEL	RSSTGILSLT	FHTDMAVAKD	GFSARYLVH	120
QEPLENFQCN	VPLGMESGRI	ANEQISASST	YSDGRWTPQQ	SRLHGDDNGW	TPNLDSNKEY	180
LQVDLRLFTM	LTAIATQGA	SRETQNGYV	KSYKLEVSTN	GEDWMVYRHG	KNHKVFQANN	240
DATEVVLNKL	HAPLLTRFVR	IRPQTWHSGL	ALRLLELFGC	VTDAPCSNML	GMLSGLIADS	300
QISASSTQEY	LWSPSAARLV	SSRSGWFPR	PQAQPGEEWL	QVDLGTPTKV	KGVIIQGAR	360

-continued

GDSITAVEAR	AFVRKFKVSY	SLNGKDWEYI	QDPRTQQPKL	PEGNMHYDTP	DIRRFDPIPA	420
QYVRVYPERW	SPAGIGMRLE	VLGCDWTDK	PTVETLGPV	KSEETTTYP	TEEEATECGE	480
NCSFEDDKDL	QLPSGFNCNF	DFLEPCGWM	YDHAKWLRTT	WASSSSPNDR	TFPDDRNFLR	540
LQSDSQREGQ	YARLISPPVH	LPRSPVCMF	QYQATGGRGV	ALQVVREASQ	ESKLLWVIRE	600
DQGGWKHGR	IILPSYDMEY	QIVFEGVIGK	GRSGEIAIDD	IRISTDVPLE	NCME	654

SEQ ID NO: 109 moltype = AA length = 780
 FEATURE Location/Qualifiers
 source 1..780
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 109

QDPDPCGGRL	NSKDAYGITS	PGYPQDYP	SH	QNCEWIVYAP	EPNQKIVLNF	NPHFEIEKHD	60	
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT		IISSGSMLYI	KFTSDYARQG	AGFSLRYEIF	120	
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ		180	
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	A	240	
RYYLHVQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL		300	
DSNKEYLQVD	LRFLTMLTAI	ATQGAISRET	QNGYYVKS	YK	LEVSTNGEDW	MVYRHGKNHK	360	
VPQANNDATE	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	A	PCSNNMLGMLS	420	
GLIADSQISA	SSTQEYLWSP	SAARLVSSRS	GWFPRIQQAQ	PGEEWLQVDL	GTPKTVKGVI		480	
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKLFEGN	MHYDTPDIRR		540	
FDPIPAQYVR	VYPERWSPAG	IGMRLEVLGC	DWTDKPTVE	TLGPTVKSEE	TTTTPYPTTEE		600	
ATECGENC	SF	EDDKDLQLPS	GFNCNFD	FLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTPFD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFYQQA	TGGRGVALQV	VREASQESKL		720	
LWVIRE	DQGG	EWKHGRILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780	

SEQ ID NO: 110 moltype = AA length = 837
 FEATURE Location/Qualifiers
 source 1..837
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 110

QDPDPCGGRL	NSKDAYGITS	PGYPQDYP	SH	QNCEWIVYAP	EPNQKIVLNF	NPHFEIEKHD	60	
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT		IISSGSMLYI	KFTSDYARQG	AGFSLRYEIF	120	
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ		180	
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	A	240	
RYYLHVQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL		300	
DSNKEYLQVD	LRFLTMLTAI	ATQGAISRET	QNGYYVKS	YK	LEVSTNGEDW	MVYRHGKNHK	360	
VPQANNDATE	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	A	PCSNNMLGMLS	420	
GLIADSQISA	SSTQEYLWSP	SAARLVSSRS	GWFPRIQQAQ	PGEEWLQVDL	GTPKTVKGVI		480	
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKLFEGN	MHYDTPDIRR		540	
FDPIPAQYVR	VYPERWSPAG	IGMRLEVLGC	DWTDKPTVE	TLGPTVKSEE	TTTTPYPTTEE		600	
ATECGENC	SF	EDDKDLQLPS	GFNCNFD	FLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTPFD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFYQQA	TGGRGVALQV	VREASQESKL		720	
LWVIRE	DQGG	EWKHGRILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780	
PISAF	AVDIP	BIHEREGYED	EIDDEYEVDW	SNSSSATSGS	GAPSTDKEKS	WLYTLDP	837	

SEQ ID NO: 111 moltype = AA length = 809
 FEATURE Location/Qualifiers
 source 1..809
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 111

QDPDPCGGRL	NSKDAYGITS	PGYPQDYP	SH	QNCEWIVYAP	EPNQKIVLNF	NPHFEIEKHD	60	
CKYDFIEIRD	GDSESADLLG	KHCGNIAPPT		IISSGSMLYI	KFTSDYARQG	AGFSLRYEIF	120	
KTGSEDCSKN	FTSPNGTIES	PGFPEKYPHN	LDCTFTILAK	PKMEIILQFL	IFDLEHDPLQ		180	
VGEGDCKYDW	LDIWDGIPHV	GPLIGKYCGT	KTPSELRSST	GILSLTFHTD	MAVAKDGFS	A	240	
RYYLHVQEPL	ENFQCNVPLG	MESGRIANEQ	ISASSTYSDG	RWTPQQSRLH	GDDNGWTPNL		300	
DSNKEYLQVD	LRFLTMLTAI	ATQGAISRET	QNGYYVKS	YK	LEVSTNGEDW	MVYRHGKNHK	360	
VPQANNDATE	VVLNKLHAPL	LTRFVRIRPQ	TWHSGLALRL	ELFGCRVTD	A	PCSNNMLGMLS	420	
GLIADSQISA	SSTQEYLWSP	SAARLVSSRS	GWFPRIQQAQ	PGEEWLQVDL	GTPKTVKGVI		480	
IQGARGGDSI	TAVEARAFVR	KFKVSYSLNG	KDWEYIQDPR	TQQPKLFEGN	MHYDTPDIRR		540	
FDPIPAQYVR	VYPERWSPAG	IGMRLEVLGC	DWTDKPTVE	TLGPTVKSEE	TTTTPYPTTEE		600	
ATECGENC	SF	EDDKDLQLPS	GFNCNFD	FLE	EPCGWMYDHA	KWLRTTWASS	SSPNDRTPFD	660
DRNFLRLQSD	SQREGQYARL	ISPPVHLPRS	PVCMEFYQQA	TGGRGVALQV	VREASQESKL		720	
LWVIRE	DQGG	EWKHGRILP	SYDMEYQIVF	EGVIGKGRSG	EIAIDDIRIS	TDVPLENCME	780	
PISAF	AGGT	LPGTEPTVDT	VPMQIPAY				809	

SEQ ID NO: 112 moltype = AA length = 62
 FEATURE Location/Qualifiers
 source 1..62
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 112

PISAF	AGENF	KVDIPEIHER	EGYEDEIDDE	YEVDWSN	SSSS	ATSGSGAPST	DKEKSWLYTL	60
DP								62

SEQ ID NO: 113 moltype = AA length = 57
 FEATURE Location/Qualifiers

-continued

source 1..57
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 113
 PISAFVDIP EIHEREGYED EIDDEYEVDW SNSSSATSGS GAPSTDKEKS WLYTLDP 57

SEQ ID NO: 114 moltype = AA length = 40
 FEATURE Location/Qualifiers
 source 1..40
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 114
 PISAFADYE VDWNSSSAT SGSGAPSTDK EKSWLYTLDP 40

SEQ ID NO: 115 moltype = AA length = 35
 FEATURE Location/Qualifiers
 source 1..35
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 115
 PISAFAGENF KGGTLLPGTE PTVDTVPMQP IPAYW 35

SEQ ID NO: 116 moltype = AA length = 30
 FEATURE Location/Qualifiers
 source 1..30
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 116
 PISAFAGGTL LPGTEPTVDT VPMQPIPAYW 30

SEQ ID NO: 117 moltype = AA length = 800
 FEATURE Location/Qualifiers
 REGION 1..800
 note = NRP2 - Fc fusion construct
 source 1..800
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 117
 QPDPPCGGRL NSKDAGYITS PGYPQDYP SH QNCEWIVYAP EPNQKIVLNF NPHFEIEKHD 60
 CKYDFIEIRD GDSESADLLG KHCGNIAPPT IISGSMLYI KFTSDYARQG AGFSLRYEIF 120
 KTGSSEDCSKN FTSPNGTIES PGFPEKYPHN LDCTFTILAK PKMEIILQFL IFDLEHDPLQ 180
 VEGEDCKYDW LDIWDGIPHV GPLIGKYCGT KTPSELRSST GILSLTFHTD MAVAKDGFSA 240
 RYYLVHQEPL ENFQCNVPLG MESGRIANEQ ISASSTYSDG RWTPQQSRLH GDDNGWTPNL 300
 DSNKEYLQVD LRFLTMLTAI ATQGAISRET QNGYVVKSYK LEVSTNGEDW MVYRHGKQNHK 360
 VFQANNDAT VVLNKLHAPL LTRFVRIRPQ TWHSGIALRL ELFGCRVTD PCSNMLGMLS 420
 GLIADSQISA SSTQEYLWSP SAARLVSSRS GWFPRIPOAQ PGEWLVQVDL GTPKTVKGVI 480
 IQGARGGDSI TAVEARAFVR KFKVSYSLNG KDWEYIQDPR TQQPKLFEGN MHYDTPDIRR 540
 FDPIPAQYVR VYPERWSPAG IGMRLVLGCG DWTDKTHTCP PCPAPELLGG PSVFLFPPPK 600
 KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN STYRVVSVLT 660
 VLHQDWLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE LTKNQVSLTC 720
 LVKGFYPSDI AVEWESNGQP ENNYKTPPV LDSDGSFFLY SKLTVDKSRW QQGNVFSCSV 780
 MHEALHNHYT QKSLSLSPGK 800

SEQ ID NO: 118 moltype = AA length = 509
 FEATURE Location/Qualifiers
 source 1..509
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 118
 MAERAALEEL VKLQGERVRG LKQKASAE IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 GTRDYSPRQM AVREKVFVDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ 120
 GGELLSLRYP LTVPFARYLA MNKLTNIKRY HIAKVYRRDN PAMTRGRYRE FYQCDFDIAG 180
 NFDPMIPDAE CLKIMCEILS SLQIGDFLVK VNDRLILDGM FAICGVSDSK FRTICSSVDK 240
 LDKVSWEEVK NEMVGEKGLA PEVADRIGDY VQQHGGVSLV EQLLQDPKLS QNKQALEGLG 300
 DLKLLFEYLT LFGIDDKISF DLSLARGLDY YTGVIYEAVL LQTPAQAGEE PLGVGSVAAG 360
 GRYDGLVGMF DPKGRKVPV GLSIGVERIF SIVEQRLEAL EEKIRTETQ VLVASQKKL 420
 LEERLKLVE LWDAGIKAE LYKKNPKLLN QLQYCEEAGI PLVAIIGQE LKDGVIKLR 480
 VTSREEVDVR REDLVEIKR RTGQPLCIC 509

SEQ ID NO: 119 moltype = AA length = 141
 FEATURE Location/Qualifiers
 source 1..141
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 119
 MAERAALEEL VKLQGERVRG LKQKASAE IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 GTRDYSPRQM AVREKVFVDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ 120
 GGELLSLRYP LTVPFARYLA M 141

-continued

SEQ ID NO: 120 moltype = AA length = 408
 FEATURE Location/Qualifiers
 source 1..408
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 120
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 GTRDYSRQPM AVREKVFVDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ 120
 GGELLSLRYD LTVPFARYLA MNKLNTIKRY HIAKVYRRDN PAMTRGRYRE FYQCDFDIAG 180
 NFDPMIPDAE CLKIMCEILS SLQIGDFLVK VNDRLDGM FAICGVSDSK FRTICSSVDK 240
 LDKVSWEDEVK NEMVGEKGLA PEVADRIGDY VQQHGGVSLV EQLLQDPKLS QNKQALEGLG 300
 DLKLLFEYLT LFGIDDKISF DLSLARGLDY YTGVIYEAVL LQTPAQAGEE PLGVGSVAAG 360
 GRYDGLVGMF DPKGRKVPCV GLSIGVERIF SIVEQRLEAL EEKIRTTE 408

SEQ ID NO: 121 moltype = AA length = 113
 FEATURE Location/Qualifiers
 source 1..113
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 121
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 GTRDYSRQPM AVREKVFVDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKL 113

SEQ ID NO: 122 moltype = AA length = 60
 FEATURE Location/Qualifiers
 source 1..60
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 122
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60

SEQ ID NO: 123 moltype = AA length = 270
 FEATURE Location/Qualifiers
 REGION 1..270
 note = HRS construct
 source 1..270
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 123
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 GTRDYSRQPM AVREKVFVDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ 120
 GGELLSLRYD LTVPFARYLA MNKLNTIKRY HIAKVYRRDN PAMTRGRYRE FYQCDFDIAG 180
 NFDPMIPDAE CLKIMCEILS SLQIGDFLVK VNDRLDGM FAICGVSDSK FRTICSSVDK 240
 LDKVGYPWWN SCSRLNYPK TSRPRAWET 270

SEQ ID NO: 124 moltype = AA length = 105
 FEATURE Location/Qualifiers
 source 1..105
 mol_type = protein
 organism = Homo sapiens

SEQUENCE: 124
 RTTETQVLVA SAQKKLEER LKLVSELWDA GIKAELLYKK NPKLLNQLQY CEEAGIPLVA 60
 IIGEQLKDG VIKLRVTSR EEVDVRREDL VEEIKRRTGQ PLCIC 105

SEQ ID NO: 125 moltype = AA length = 395
 FEATURE Location/Qualifiers
 REGION 1..395
 note = human HRS construct
 source 1..395
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 125
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
 DFDIAGNFDIP MIPDAECLKI MCEILSSLQI GDFLVKVNDR RILDGMFAIC GVSDSKERTI 120
 CSSVDKLDKV SWEVKNEMV GEKGLAPEVA DRIGDYVQQH GGVSLVEQLL QDPKLSQNKQ 180
 ALEGLDGLL LFEYLTFLGI DDKISFDLSL ARGLDYYTGV IYEAULLQTP AQAGEEPLGV 240
 GSVAAGGRYD GLVGMFDPKG RKVPCVGLSI GVERIFSIVE QRLEALEEKI RTTETQVLVA 300
 SAQKKLEER LKLVSELWDA GIKAELLYKK NPKLLNQLQY CEEAGIPLVA IIGEQLKDG 360
 VIKLRVTSR EEVDVRREDL VEEIKRRTGQ PLCIC 395

SEQ ID NO: 126 moltype = AA length = 359
 FEATURE Location/Qualifiers
 REGION 1..359
 note = human HRS construct
 source 1..359
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 126
 MAERAALEEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60

-continued

VNDRRILDGM	FAICGVSDSK	FRTICSSVDK	LDKVSWEVK	NEMVGEKGLA	PEVADRIGDY	120
VQQHGGVSLV	EQLLQDPKLS	QNKQALEGLG	DLKLLFEYLT	LFGIDDKISF	DLSLARGLDY	180
YTGVIYEAVL	LQTPAQAGEE	PLGVGSVAAG	GRYDGLVGMF	DPKGRKVPVCV	GLSIGVERIF	240
SIVEQRLEAL	EEKIRTTETQ	VLVASAQKKL	LEERLKLVS	LWDAGIKAE	LYKKNPKLLN	300
QLQYCEEAGI	PLVAIIGEQE	LKDGVIKLS	VTSREEVDVR	REDLVEEIKR	RTGQPLCIC	359

SEQ ID NO: 127 moltype = AA length = 399
 FEATURE Location/Qualifiers
 REGION 1..399
 note = human HRS construct
 source 1..399
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 127

MAERAALDEL	VKLQGERVRG	LKQKASAE	IEEEVAKLLK	LKAQLGPDES	KQKFVLKTPK	60
GTRDYSRPM	AVREKVPDVI	IRCFKRHGA	VIDTPVFELK	VNDRRILDGM	FAICGVSDSK	120
FRTICSSVDK	LDKVSWEVK	NEMVGEKGLA	PEVADRIGDY	VQQHGGVSLV	EQLLQDPKLS	180
QNKQALEGLG	DLKLLFEYLT	LFGIDDKISF	DLSLARGLDY	YTGVIYEAVL	LQTPAQAGEE	240
PLGVGSVAAG	GRYDGLVGMF	DPKGRKVPVCV	GLSIGVERIF	SIVEQRLEAL	EEKIRTTETQ	300
VLVASAQKKL	LEERLKLVS	LWDAGIKAE	LYKKNPKLLN	QLQYCEEAGI	PLVAIIGEQE	360
LKDGVIKLS	VTSREEVDVR	REDLVEEIKR	RTGQPLCIC			399

SEQ ID NO: 128 moltype = AA length = 473
 FEATURE Location/Qualifiers
 REGION 1..473
 note = human HRS construct
 source 1..473
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 128

MAERAALDEL	VKLQGERVRG	LKQKASAE	IEEEVAKLLK	LKAQLGPDES	KQKFVLKTPK	60
GTRDYSRPM	AVREKVPDVI	IRCFKRHGA	VIDTPVFELK	ETLMGKYGED	SKLIYDLKDQ	120
GGELLSRLYD	LTVPFARYLA	MNKLNTIKRY	HIKAVYRRDN	PAMTRGRYRE	FYQCVNDRRI	180
LDGMFAICGV	SDSKERTICS	SVDKLDKVS	EEVKNEVMGE	KGLAPEVADR	IGDYVQQHGG	240
VSLVEQLLQD	PKLSQNKQAL	EGLGDLKLLF	EYLTFLGIDD	KISFDLSLAR	GLDYYTGVYI	300
EAVLLQTPAQ	AGEEPLGVGS	VAAGGRYDGL	VGMFDPKGRK	VPCVGLSIGV	ERIFSIVEQR	360
LEALEEKIRT	TETQVLVASA	QKKLLEERLK	LVSELWDAGI	KAELLYKKNP	KLLNQLQYCE	420
EAGIPLVAII	GEQELKDGVI	KLSVTSREE	VDVREDLVE	EIKRRTGQPL	CIC	473

SEQ ID NO: 129 moltype = AA length = 469
 FEATURE Location/Qualifiers
 REGION 1..469
 note = human HRS construct
 source 1..469
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 129

MAERAALDEL	VKLQGERVRG	LKQKASAE	IEEEVAKLLK	LKAQLGPDES	KQKFVLKTPK	60
ETLMGKYGED	SKLIYDLKDQ	GGELLSRLYD	LTVPFARYLA	MNKLNTIKRY	HIKAVYRRDN	120
PAMTRGRYRE	FYQCDFDIAG	NFDPMPIDAE	CLKIMCEILS	SLQIGDFLVK	VNDRRILDGM	180
FAICGVSDSK	FRTICSSVDK	LDKVSWEVK	NEMVGEKGLA	PEVADRIGDY	VQQHGGVSLV	240
EQLLQDPKLS	QNKQALEGLG	DLKLLFEYLT	LFGIDDKISF	DLSLARGLDY	YTGVIYEAVL	300
LQTPAQAGEE	PLGVGSVAAG	GRYDGLVGMF	DPKGRKVPVCV	GLSIGVERIF	SIVEQRLEAL	360
EEKIRTTETQ	VLVASAQKKL	LEERLKLVS	LWDAGIKAE	LYKKNPKLLN	QLQYCEEAGI	420
PLVAIIGEQE	LKDGVIKLS	VTSREEVDVR	REDLVEEIKR	RTGQPLCIC		469

SEQ ID NO: 130 moltype = AA length = 435
 FEATURE Location/Qualifiers
 REGION 1..435
 note = human HRS construct
 source 1..435
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 130

MAERAALDEL	VKLQGERVRG	LKQKASAE	IEEEVAKLLK	LKAQLGPDES	KQKFVLKTPK	60
GTRDYSRPM	AVREKVPDVI	IRCFKRHGA	VIDTPVFELK	DFDIAGNFD	MIPDAECLKI	120
MCEILSSLQI	GDIFVKNDR	RILDGMFAIC	GVSDSKFRTI	CSSVDKLDKV	SWEVKNEVM	180
GEKGLAPEVA	DRIGDYVQQH	GGVSLVEQLL	QDPKLSQNKQ	ALEGLGDLKL	LFEYTLFGI	240
DDKISFDLSL	ARGLDYYTGV	IYEAVALQTP	AQAGEEPLGV	GSAAGGRYD	GLVGMFDPKG	300
RKVPCVGLSI	GVERIFSIVE	QRLEALEEKI	RTTETQVLVA	SAQKKLLEER	LKLVSSELWDA	360
GKAEALLYKK	NPKLLNQLQY	CEEAGIPLVA	IIGEQELKDG	VIKLSVTSR	EEVDVRRDEL	420
VEEIKRRTGQ	PLCIC					435

SEQ ID NO: 131 moltype = AA length = 171
 FEATURE Location/Qualifiers
 REGION 1..171
 note = human HRS construct
 source 1..171
 mol_type = protein

-continued

```

                organism = synthetic construct
SEQUENCE: 131
MAERAAL EEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
ALEEKIR TTE TQVLVASAQK KLL EERLKL V SELWDAGIKA ELLYKKNPKL LNQLQYCEEA 120
GIPLVAI IGE QELKDGVIKL RSVTSREEVD VRREDLVEEI KRRTGQPLCI C 171

SEQ ID NO: 132      moltype = AA length = 211
FEATURE            Location/Qualifiers
REGION            1..211
                  note = human HRS construct
source            1..211
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 132
MAERAAL EEL VKLQGERVRG LKQKASAEI IEEVAKLLK LKAQLGPDES KQKFVLKTPK 60
GTRDYS PRQM AVREKVFDVI IRCFKRHGAE VIDTPVFELK ALEEKIR TTE TQVLVASAQK 120
KLL EERLKL V SELWDAGIKA ELLYKKNPKL LNQLQYCEEA GIPLVAI IGE QELKDGVIKL 180
RSVTSREEVD VRREDLVEEI KRRTGQPLCI C 211

SEQ ID NO: 133      moltype = AA length = 141
FEATURE            Location/Qualifiers
source            1..141
                  mol_type = protein
                  organism = Homo sapiens

SEQUENCE: 133
MFDPKGR KVP CVGLSIGVER IFSIVEQRLE ALEEKIR TTE TQVLVASAQK KLL EERLKL V 60
SELWDAG IKA ELLYKKNPKL LNQLQYCEEA GIPLVAI IGE QELKDGVIKL RSVTSREEVD 120
VRREDLVE EI KRRTGQPLCI C 141

SEQ ID NO: 134      moltype = AA length = 143
FEATURE            Location/Qualifiers
source            1..143
                  mol_type = protein
                  organism = Homo sapiens

SEQUENCE: 134
CLKIMCE ILS SLQIGDFLVK VNDRRILDGM FAICGVSDSK FRTICSSVDK LDKVSWEEVK 60
NEMVGEK GLA PEVADRIGDY VQCHGGVSLV EQLLQDPKLS QNKQALEGLG DLKLLFEYLT 120
LFGIDDK ISF DLSLARGLDY YTG 143

SEQ ID NO: 135      moltype = AA length = 506
FEATURE            Location/Qualifiers
source            1..506
                  mol_type = protein
                  organism = Homo sapiens

SEQUENCE: 135
MPLLGLL PRR AWASLLSQLL RPPCASCTGA VRCQSQVAEA VLTSQLKAHQ EKPNIKI TP 60
KGTRDL SPQH MUVREKILD L VISCFKRHGA KGMDTPAFEL KETLTEKYGE DSGLMYDLKD 120
QGGE LLSLR Y DLTVPFARYL AMNKVKMKR YHVGVWRRE SPTIVQGRYR EPCQCDFDIA 180
GQFDPMI PDA ECLKIMCEI L SGLQLGDFLI KVNDRIIDVG MFAVCGVPES KFRAICSSID 240
KLDKMAWK DV RHEMVVKGL APEVADRIGD YVQCHGGVSL VEQMFQDPRL SQNKQALEGL 300
GDLKLLF EYL TLFGIADKIS FDLSLARGLD YTGVIYEAV LLQTPTQAGE EPLNVGSVA 360
GGRYDGL VGM FDPKGHKVPC VGLSIGVERI FYIVEQRMKT KGEKVRTTET QVFVATPQKN 420
FLQERLKL IA ELWDSGIKAE MLYKNNPKLL TQLHYCESTG IPLVVIIEGQ ELKEGVIKIR 480
SVASREEVA I KRENFVAEI Q KRLSES 506

SEQ ID NO: 136      moltype = AA length = 247
FEATURE            Location/Qualifiers
source            1..247
                  mol_type = protein
                  organism = Homo sapiens

SEQUENCE: 136
HVGKVWR RES PTIVQGRYRE FCQCDFDIAG QFDPMIPDAE CLKIMCEILS GLQLGDFLIK 60
VNDRIIDV GM FAVCGVPESK FRAICSSIDK LDKMAWKDVR HEMVVKKGLA PEVADRIGDY 120
VQCHGGVSL V EQMFQDPRLS QNKQALEGLG DLKLLFEYLT LFGIADKISF DLSLARGLDY 180
YTGVIYEAV L LQTPTQAGEE PLNVGSVAAG GRYDGLVGMF DPKGHKVPCV GLSIGVERIF 240
YIVEQRM 247

SEQ ID NO: 137      moltype = AA length = 79
FEATURE            Location/Qualifiers
source            1..79
                  mol_type = protein
                  organism = Homo sapiens

SEQUENCE: 137
QALEGLG DLK LLFEYLT LFG IDDKISFDLS LARGLDYYTG VIYEAVLLQT PAQAGEEPLG 60
VGSVAAGGR Y DGLVGMFDP 79

SEQ ID NO: 138      moltype = AA length = 456
FEATURE            Location/Qualifiers
source            1..456

```

-continued

```

mol_type = protein
organism = Homo sapiens

SEQUENCE: 138
FVLKTPKGTR DYSRQMAVR EKVPDVIIRC FKRHGAEVID TPVPFELKETL MGKYGEDSKL 60
IYDLKDQGGG LLSRLYDLTV PFARYLAMNK LTNIKRYHIA KVVRRDNPAM TRGRYREFYQ 120
CDFDIAGNFD PMIPDAECLK IMCEILSSLQ IGDPLVKVND RRILDGMFAI CGVSDSKFRT 180
ICSSVDKLDK VSWEEVKNEV VGEKGLAPEV ADRIGDYVQQ HGGVSLVEQL LQDPKLSQNK 240
QALEGLGLDK LLFEYLTFLG IDDKISFDLS LARGLDYYTG VIYEAVLLQT PAQAGEEPLG 300
VGSVAAGGRY DGLVGMFDPK GRKVPCVGLS IGVVERIFSIV EQRLEALEEK IRTTETQVLV 360
ASAQKLLLEE RLKLVSELWD AGIKAELLYK KNPKLLNQLQ YCEEAGIPLV AIIGEQLKLD 420
GVIKLRSVTS REEVDVRRED LVBEIKRRTG QPLCIC 456

SEQ ID NO: 139      moltype = AA length = 345
FEATURE            Location/Qualifiers
source              1..345
                    mol_type = protein
                    organism = Homo sapiens

SEQUENCE: 139
FVLKTPKGTR DYSRQMAVR EKVPDVIIRC FKRHGAEVID TPVPFELKETL MGKYGEDSKL 60
IYDLKDQGGG LLSRLYDLTV PFARYLAMNK LTNIKRYHIA KVVRRDNPAM TRGRYREFYQ 120
CDFDIAGNFD PMIPDAECLK IMCEILSSLQ IGDPLVKVND RRILDGMFAI CGVSDSKFRT 180
ICSSVDKLDK VSWEEVKNEV VGEKGLAPEV ADRIGDYVQQ HGGVSLVEQL LQDPKLSQNK 240
QALEGLGLDK LLFEYLTFLG IDDKISFDLS LARGLDYYTG VIYEAVLLQT PAQAGEEPLG 300
VGSVAAGGRY DGLVGMFDPK GRKVPCVGLS IGVVERIFSIV EQRLE 345

SEQ ID NO: 140      moltype = AA length = 338
FEATURE            Location/Qualifiers
source              1..338
                    mol_type = protein
                    organism = Homo sapiens

SEQUENCE: 140
GTRDYSRQPM AVREKVPDVI IRCFKRHGAE VIDTPVFELK ETLMGKYGED SKLIYDLKDQ 60
GGELLSRLYD LTVPFARYLA MNKLTNIKRY HIAKVYRRDN PAMTRGRYRE FYQCDFDIAG 120
NFDPMIPDAE CLKIMCEILS SLQIGDFLVK VNDRRILDGM FAICGVSDSK FRTICSSVDK 180
LDKVSWEVKV NEMVGEKGLA PEVADRIGDY VQQHGGVSLV EQLLQDPKLS QNKQALEGLG 240
DKLLFEYLT LFGIDDKISF DLSLARGLDY YTGVIYEAVL LQTPAQAGEE PLGVGSVAAG 300
GRYDGLVGMP DPKGRKVPCV GLSIGVERIF SIVEQRLE 338

SEQ ID NO: 141      moltype = AA length = 111
FEATURE            Location/Qualifiers
source              1..111
                    mol_type = protein
                    organism = Homo sapiens

SEQUENCE: 141
ALEEKIRTTT TQVLVASAQK KLEEERLKL VSELWDAGIKA ELLYKKNPKL LNQLQYCEEA 60
GIPLVAIIGE QELKDGVIKL RSVTSREEVD VRREDLVEEI KRRTGQPLCI C 111

SEQ ID NO: 142      moltype = AA length = 96
FEATURE            Location/Qualifiers
source              1..96
                    mol_type = protein
                    organism = Homo sapiens

SEQUENCE: 142
TTETQVLVAS AQKLLLEERL KLVSELWDAG IKAELLYKKN PKLLNQLQYC EEAGIPLVAI 60
IGEQLKDGVI IKLRSVTSRE EVDVRREDLV EEIKRR 96

SEQ ID NO: 143      moltype = AA length = 138
FEATURE            Location/Qualifiers
REGION              1..138
                    note = Consensus HRS WHEP sequence
VARIANT              1..50
                    note = Xaa is any amino acid or absent
VARIANT              52..56
                    note = Xaa can be any naturally occurring amino acid
VARIANT              57..58
                    note = Xaa is any amino acid or absent
VARIANT              61..62
                    note = Xaa is any amino acid
VARIANT              65
                    note = Xaa is any amino acid
VARIANT              68..69
                    note = Xaa is any amino acid
VARIANT              72..78
                    note = Xaa can be any naturally occurring amino acid
VARIANT              79..80
                    note = Xaa is any amino acid or absent
VARIANT              82..83
                    note = Xaa is any amino acid
VARIANT              86

```

-continued

VARIANT note = Xaa is any amino acid
89..138
note = Xaa is any amino acid or absent
source 1..138
mol_type = protein
organism = synthetic construct

SEQUENCE: 143
XXXXXXXXXX XXXXXXXXXXXX XXXXXXXXXXXX XXXXXXXXXXXX LXXXXXXXXQ 60
XXVRLKXXX XXXXXXXXXXXX VXXLLXLKXX XXXXXXXXXXXX XXXXXXXXXXXX 120
XXXXXXXXXX XXXXXXXX 138

SEQ ID NO: 144 moltype = AA length = 327
FEATURE Location/Qualifiers
source 1..327
mol_type = protein
organism = Homo sapiens

SEQUENCE: 144
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 60
GLYSLSSVVT VPSSSLGTTK YTCNVDHKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV 120
FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY 180
RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK 240
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG 300
NVFSCSVME ALHNHYTQKS LSLSLGK 327

SEQ ID NO: 145 moltype = AA length = 327
FEATURE Location/Qualifiers
REGION 1..327
note = Humanized IgG4
source 1..327
mol_type = protein
organism = synthetic construct

SEQUENCE: 145
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 60
GLYSLSSVVT VPSSSLGTTK YTCNVDHKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV 120
FLFPPKPKDT LYITREPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY 180
RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK 240
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG 300
NVFSCSVME ALHNHYTQKS LSLSLGK 327

SEQ ID NO: 146 moltype = AA length = 327
FEATURE Location/Qualifiers
REGION 1..327
note = humanized IgG4
source 1..327
mol_type = protein
organism = synthetic construct

SEQUENCE: 146
ASTKGPSVFP LAPCSRSTSE STAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 60
GLYSLSSVVT VPSSSLGTTK YTCNVDHKPS NTKVDKRVES KYGPPCPSCP APEFLGGPSV 120
FLFPPKPKDT LMISRTPEVT CVVVDVSQED PEVQFNWYVD GVEVHNAKTK PREEQFNSTY 180
RVVSVLTVLH QDWLNGKEYK CKVSNKGLPS SIEKTISKAK GQPREPQVYT LPPSQEEMTK 240
NQVSLTCLVK GFYPSDIAVE WESNGQPENN YKTTTPVLDS DGSFFLYSRL TVDKSRWQEG 300
NVFSCSVME ALHNHYTQKS LSLSLGK 327

SEQ ID NO: 147 moltype = AA length = 330
FEATURE Location/Qualifiers
source 1..330
mol_type = protein
organism = Homo sapiens

SEQUENCE: 147
ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 60
GLYSLSSVVT VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKTHTCP PCPAPELLGG 120
PSVFLFPPKP KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN 180
STYRVVSVLT VLNQDNLNGK EYKCKVSNKA LPAPIEKTIS KAKGQPREPQ VYTLPPSRDE 240
LTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSGGSFFLY SKLTVDKSRW 300
QQGNVFSV MHEALHNHYT QKSLSLSPGK 330

SEQ ID NO: 148 moltype = AA length = 107
FEATURE Location/Qualifiers
source 1..107
mol_type = protein
organism = Homo sapiens

SEQUENCE: 148
RTVAAPSVFI FPPSDEQLKS GTASVVCLLN NFYPREAKVQ WKVDNALQSG NSQESVTEQD 60
SKDSTYLSL TLTLKADYE KHKVYACEVT HQGLSSPVTK SFNRGEC 107

SEQ ID NO: 149 moltype = DNA length = 24
FEATURE Location/Qualifiers
misc_feature 1..24

-continued

```

source          note = primer
                1..24
                mol_type = other DNA
                organism = synthetic construct

SEQUENCE: 149
tgaggatgac aaagatttgc agct                                24

SEQ ID NO: 150      moltype = DNA  length = 36
FEATURE            Location/Qualifiers
misc_feature        1..36
                    note = primer
source              1..36
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 150
accgcggccg gccgtttatg cctcggagca gcactt                    36

SEQ ID NO: 151      moltype = DNA  length = 22
FEATURE            Location/Qualifiers
misc_feature        1..22
                    note = primer
source              1..22
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 151
agtgccaagc aagcaactca aa                                    22

SEQ ID NO: 152      moltype = DNA  length = 36
FEATURE            Location/Qualifiers
misc_feature        1..36
                    note = primer
source              1..36
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 152
aagtgtctgt ccgaggcata aacggccggc cgcggt                    36

SEQ ID NO: 153      moltype = AA  length = 5
FEATURE            Location/Qualifiers
source              1..5
                    mol_type = protein
                    organism = Homo sapiens

SEQUENCE: 153
GENFK                                                        5

```

The invention claimed is:

1. A therapeutic composition, comprising a pharmaceutically-acceptable carrier and at least one antibody or antigen-binding fragment thereof that specifically binds to a human neuropilin-2 (NRP2) polypeptide at an epitope in the neuropilin a2 domain of human NRP2, wherein the at least one antibody or antigen-binding fragment thereof comprises:

a heavy chain variable region (V_H) sequence that comprises complementary determining region V_H CDR1, V_H CDR2, and V_H CDR3 sequences; and a light chain variable region (V_L) sequence that comprises complementary determining region V_L CDR1, V_L CDR2, and V_L CDR3 sequences, wherein:

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 34-36, respectively; or

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 37-39, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 40-42, respectively,

including variants thereof that differ by substitution of one amino acid in V_H CDR3, V_L CDR1 and/or V_L CDR3, wherein the V_H CDR3 sequence comprises SEQ ID NO: 81, the V_L CDR1 sequence comprises SEQ ID NO: 82, and the V_L CDR3 sequence comprises SEQ ID NO: 83, wherein X5 of SEQ ID NO: 83 is F, G, I, K, L, N, R, T, V, or Y.

2. The therapeutic composition of claim 1, wherein: the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 31-33, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 34-36, respectively; or

the V_H CDR1, V_H CDR2, and V_H CDR3 sequences comprise SEQ ID NOs: 37-39, respectively, and the V_L CDR1, V_L CDR2, and V_L CDR3 sequences comprise SEQ ID NOs: 40-42, respectively.

3. The therapeutic composition of claim 2, wherein: the V_H sequence comprises a sequence at least 95% identical to a SEQ ID NO: 53, and the V_L sequence comprises a sequence at least 95% identical to SEQ ID NO: 54; or

the V_H sequence comprises a sequence at least 95% identical to SEQ ID NO: 55, and the V_L sequence comprises a sequence at least 95% identical to SEQ ID NO: 56.

4. The therapeutic composition of claim 3, wherein: the V_H sequence comprises SEQ ID NO: 53, and the V_L sequence comprises SEQ ID NO: 54; or the V_H sequence comprises SEQ ID NO: 55, and the V_L sequence comprises SEQ ID NO: 56.

5. The therapeutic composition of claim 1, wherein the at least one antibody or antigen-binding fragment thereof comprises an IgA (including subclasses IgA1 and IgA2), IgD, IgE, IgG (including subclasses IgG1, IgG2, IgG3, and

209

IgG4), or IgM Fc domain, or variant thereof, optionally wherein the Fc domain is a human Fc domain.

6. The therapeutic composition of claim 5, wherein the at least one antibody or antigen-binding fragment thereof comprises an IgG1 or IgG3 Fc domain.

7. The therapeutic composition of claim 5, wherein the at least one antibody or antigen-binding fragment thereof comprises an IgG2 or IgG4 Fc domain.

8. The therapeutic composition of claim 5, wherein the at least one antibody or antigen-binding fragment thereof comprises an IgG1 or IgG4 Fc domain, optionally selected from SEQ ID NOs: 144-147.

9. The therapeutic composition of claim 1, wherein the at least one antibody or antigen-binding fragment thereof comprises a modified IgG1 or IgG4 Fc domain which has altered binding to FcRn, optionally wherein the modified IgG1 or IgG4 Fc domain comprises any one or more of YD (M252Y/T256D), DQ (T256D/T307Q), DW (T256D/T307W), YTE (M252Y/S254T/T256E), AAA (T307A/E380A/N434A), LS (M428L/N434S), M252Y, T256D/E, K288D/N, T307Q/W, E380C, N434F/Y, and/or Y436H/N/W mutations (EU numbering).

210

10. The therapeutic composition of claim 1, wherein the at least one antibody or antigen-binding fragment thereof is a monoclonal antibody and/or a humanized antibody.

11. The therapeutic composition of claim 1, wherein the at least one antibody or antigen-binding fragment thereof is an Fv fragment, a single chain Fv (scFv) polypeptide, or a minibody.

12. The therapeutic composition of claim 1, wherein the composition has a purity of at least about 80% on a protein basis with respect to the at least one antibody or antigen-binding fragment, and is substantially aggregate-free.

13. The therapeutic composition of claim 12, wherein the therapeutic composition is substantially endotoxin-free.

14. The therapeutic composition of claim 1, wherein the therapeutic composition is a sterile, injectable solution, optionally suitable for intravenous, intramuscular, subcutaneous, or intraperitoneal administration.

15. The therapeutic composition of claim 1, which is a concentrated lyophilized composition.

* * * * *