

(19) **United States**

(12) **Patent Application Publication**
Maslow et al.

(10) **Pub. No.: US 2025/0276051 A1**
(43) **Pub. Date: Sep. 4, 2025**

(54) **POWASSAN VIRAL ANTIGENS AND RELATED COMPOSITIONS, AND USES THEREOF TO VACCINATE AND TREAT PATIENTS**

(71) Applicant: **GeneOne Life Science, Inc.**, Seoul (KR)

(72) Inventors: **Joel Maslow**, Seoul (KR); **Moonsup Jeong**, Seoul (KR); **Christine Roberts**, Seoul (KR); **Youngran Cho**, Seoul (KR); **Young Park**, Seoul (KR)

(73) Assignee: **GeneOne Life Science, Inc.**, Seoul (KR)

(21) Appl. No.: **17/786,310**

(22) PCT Filed: **Dec. 18, 2020**

(86) PCT No.: **PCT/US2020/066233**

§ 371 (c)(1),

(2) Date: **Jun. 16, 2022**

Related U.S. Application Data

(60) Provisional application No. 62/949,821, filed on Dec. 18, 2019.

Publication Classification

(51) **Int. Cl.**

A61K 39/12 (2006.01)

A61K 39/00 (2006.01)

A61P 31/14 (2006.01)

C07K 14/005 (2006.01)

C12N 7/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/12** (2013.01); **A61P 31/14**

(2018.01); **C07K 14/005** (2013.01); **C12N 7/00**

(2013.01); **A61K 2039/53** (2013.01); **A61K**

2039/54 (2013.01); **A61K 2039/575** (2013.01);

C12N 2770/24122 (2013.01); **C12N**


2770/24134 (2013.01)

(57)


ABSTRACT

Provided herein is an immunogenic composition comprising a synthetic consensus antigen to POWV. Also disclosed herein is a method of treating an infection by POWV in a subject in need thereof, by administering the immunogenic composition to the subject.

Specification includes a Sequence Listing.



ELISA analysis measuring binding antibody against POWV prME antigen in immunized mice



Lane 1	DMEM (10% FBS, 1% P/S)
Lane 2	pVAX1
Lane 3	pVAX1-IgE leader-prME
Lane 4	pVAX1-IgE leader-prM signal (20 aa)-prME
Lane 5	pVAX1-IgE leader-portion of prM signal (12 aa)-prME
Lane 6	pVAX1-Met-prM signal (20 aa)-prME
Lane 7	pVAX1-JEV signal-prME

Fig. 1. Western Blot analysis of POWV-prM protein expressed from transfected cells




Fig. 2. ELISA analysis measuring binding antibody against POWV prME antigen in immunized mice




Fig. 3. Comparison of POWV specific binding antibody level between groups at each




Fig. 4. Endpoint titers for POWV-E specific antibodies

POWASSAN VIRAL ANTIGENS AND RELATED COMPOSITIONS, AND USES THEREOF TO VACCINATE AND TREAT PATIENTS

TECHNICAL FIELD

[0001] The present invention relates to compositions comprising an immunological Powassan virus antigen, compositions comprising a recombinant nucleic acid sequence for generating one or more synthetic antigens, and functional fragments thereof, and combinations thereof. The compositions of the invention provide improved methods for inducing immune responses, and for prophylactically and/or therapeutically immunizing individuals against Powassan virus.

BACKGROUND

[0002] Vaccines help the body fight disease by training the immune system to recognize and destroy harmful substances and diseased cells. Vaccines can be largely grouped into two types, preventive and treatment vaccines. Preventive vaccines are given to healthy people to prevent the development of specific diseases, while treatment vaccines, also referred to as immunotherapies, are given to a person who has been diagnosed with disease.

[0003] Nucleic acid based vaccines work by inducing expression of a small fraction of genes within the host's cells, which in turn, enhance the ability of the host's immune system to eliminate the infecting pathogen. As such, clinical response of a viral vaccine can depend on the ability of the vaccine to obtain a high-level of immunogenicity and have sustained long-term expression.

[0004] The Powassan virus (POWV) is a flavivirus. The Powassan virus has been noted as the only tick-borne flavivirus in North America with human pathogenicity. Powassan virus is also found in the warm climate across Eurasia, where it is part of the tick-borne encephalitis virus-complex. Powassan encephalitis is severe, and neurologic sequelae are common. There are currently no medications or vaccines to treat or prevent the POWV. People affected by Powassan virus generally show symptoms 1 to 3 weeks after infection. The initial symptoms include fever, headache, nausea, occasional confusion, and weakness. Neurologic complications of Powassan virus infection may include: seizures, aphasia, cranial nerve palsies, paresis and altered mental status. About 10% of POWV encephalitis cases are fatal and half the survivors have permanent symptoms that affect their brain.

[0005] Therefore, there remains a need to discover novel compositions and methods to prevent POWV infection and for those affected, novel compositions and methods to treat POWV infections.

SUMMARY

[0006] Aspects of the present invention are directed to immunological compositions comprising a nucleotide sequence encoding a consensus POWV antigen described herein. The consensus POWV antigen comprises an amino acid sequence that is prME, and can include a signal sequence such as JEV signal, prM signal or a portion of prM signal. The nucleotide sequence can include a leader sequence such as an IgE leader.

[0007] Some aspects of the present invention are directed to methods of treating or preventing POWV infection in a subject in need thereof by administering the immunological compositions provided herein.

BRIEF DESCRIPTION OF DRAWINGS

[0008] FIG. 1 shows an image of a western blot of POWV-prM protein expression from transfected cells.

[0009] FIG. 2 shows graphs of ELISA analysis that illustrate measurement of antibody binding against POWV prME antigen from sera from the various construct (pVAX1, pVAX1-IgE leader-prME, pVAX1-IgE leader-portion of prM signal-prME, pVAX1-prM signal-prME, or pVAX1-JEV signal-prME) immunized mice.

[0010] FIG. 3 shows graphs of ELISA analysis that compare antibody binding against POWV prME between the various mouse sera (pVAX1, pVAX1-IgE leader-prME, pVAX1-IgE leader-portion of prM signal-prME, pVAX1-prM signal-prME, or pVAX1-JEV signal-prME transfected mice).

[0011] FIG. 4 shows graphs that provide endpoint titers for POWV-E specific antibodies for each of the constructs ((pVAX1, pVAX1-IgE leader-prME, pVAX1-IgE leader-portion of prM signal-prME, pVAX1-prM signal-prME, or pVAX1-JEV signal-prME).

DETAILED DESCRIPTION OF EMBODIMENTS

Definitions

[0012] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art. In case of conflict, the present document, including definitions, will control. Preferred methods and materials are described below, although methods and materials similar or equivalent to those described herein can be used in practice or testing of the present invention. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. The materials, methods, and examples disclosed herein are illustrative only and not intended to be limiting.

[0013] The terms “comprise(s),” “include(s),” “having,” “has,” “can,” “contain(s),” and variants thereof, as used herein, are intended to be open-ended transitional phrases, terms, or words that do not preclude the possibility of additional acts or structures. The singular forms “a,” “and” and “the” include plural references unless the context clearly dictates otherwise. The present disclosure also contemplates other embodiments “comprising,” “consisting of” and “consisting essentially of,” the embodiments or elements presented herein, whether explicitly set forth or not.

[0014] “Adjuvant” as used herein means any molecule added to the immunogenic composition described herein to enhance the immunogenicity of the antigen.

[0015] “Antigen” as used herein means molecules that can be recognized by the immune system, and more specifically are bound by antibodies or a cell surface receptor on immune cells such as T-cells. Preferably, Antigens invoke an immune response, such as inducing a humoral or cell-mediated immune response. Antigens are usually proteins, peptides (amino acid chains) and polysaccharides (chains of monosaccharides/simple sugars) but lipids and nucleic acids become antigens only when combined with proteins and

polysaccharides. The antigen may originate from within the body ("self-antigen") or from the external environment ("non-self").

[0016] "Coding sequence" or "encoding nucleic acid" as used herein means the nucleic acids (RNA or DNA molecule) that comprise a nucleotide sequence which encodes a protein. The coding sequence can further include initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of an individual or mammal to which the nucleic acid is administered.

[0017] "Complement" or "complementary" as used herein means Watson-Crick (e.g., A-T/U and C-G) or Hoogsteen base pairing between nucleotides or nucleotide analogs of nucleic acid molecules.

[0018] "Consensus" or "Consensus Sequence" as used herein may mean a nucleic acid sequence, or corresponding polypeptide sequence, constructed based on analysis of an alignment of multiple subtypes of a particular antigen. The sequence may be used to induce broad immunity against multiple subtypes, serotypes, or strains of a particular pathogen.

[0019] As used herein, the term "expressible form" refers to gene constructs that contain the necessary regulatory elements operably linked to a coding sequence that encodes a target protein, such that when present in the cell of the individual, the coding sequence will be expressed.

[0020] "Fragment" as used herein means a nucleotide sequence or a portion thereof that encodes a polypeptide capable of eliciting an immune response in a mammal. The fragments can be DNA or RNA fragments selected from at least one of the various nucleotide sequences that encode protein fragments set forth below.

[0021] "Immunogenic fragment" with respect to polypeptide sequences means a polypeptide capable of eliciting an immune response in a mammal that cross reacts with a full length endogenous antigen. Fragments of consensus proteins can comprise at least 80%, at least 90% or at least 95% of a consensus protein. In some embodiments, fragments of consensus proteins can comprise at least 520 amino acids or more, at least 530 amino acids or more, at least 540 amino acids or more, at least 550 amino acids or more, at least 560 amino acids or more, at least 570 amino acids or more, at least 580 amino acids or more, at least 590 amino acids or more, at least 600 amino acids or more, at least 610 amino acids or more, at least 620 amino acids or more, at least 630 amino acids or more, at least 640 amino acids or more, at least 650 amino acids or more, or at least 660 amino acids or more of a consensus protein.

[0022] As used herein, the term "genetic construct" refers to the DNA or RNA molecules that comprise a nucleotide sequence which encodes a protein. The coding sequence includes initiation and termination signals operably linked to regulatory elements including a promoter and polyadenylation signal capable of directing expression in the cells of the individual to whom the nucleic acid molecule is administered. As used herein, the term "expressible form" refers to gene constructs that contain the necessary regulatory elements operable linked to a coding sequence that encodes a protein such that when present in the cell of the individual, the coding sequence will be expressed.

[0023] "Identical" or "identity" as used herein in the context of two or more nucleic acids or polypeptide sequences, means that the sequences have a specified per-

centage of residues that are the same over a specified region. The percentage can be calculated by optimally aligning the two sequences, comparing the two sequences over the specified region, determining the number of positions at which the identical residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the specified region, and multiplying the result by 100 to yield the percentage of sequence identity. In cases where the two sequences are of different lengths or the alignment produces one or more staggered ends and the specified region of comparison includes only a single sequence, the residues of single sequence are included in the denominator but not the numerator of the calculation. When comparing DNA and RNA, thymine (T) and uracil (U) can be considered equivalent. Identity can be performed manually or by using a computer sequence algorithm such as BLAST or BLAST 2.0

[0024] "Immune response" as used herein means the activation of a host's immune system, e.g., that of a mammal, in response to the introduction of antigen. The immune response can be in the form of a cellular or humoral response, or both.

[0025] "Nucleic acid" or "oligonucleotide" or "polynucleotide" as used herein means at least two nucleotides covalently linked together. The depiction of a single strand also defines the sequence of the complementary strand. Thus, a nucleic acid also encompasses the complementary strand of a depicted single strand. Many variants of a nucleic acid can be used for the same purpose as a given nucleic acid. Thus, a nucleic acid also encompasses substantially identical nucleic acids and complements thereof. A single strand provides a probe that can hybridize to a target sequence under stringent hybridization conditions. Thus, a nucleic acid also encompasses a probe that hybridizes under stringent hybridization conditions.

[0026] Nucleic acids can be single stranded or double stranded, or can contain portions of both double stranded and single stranded sequence. The nucleic acid can be DNA, both genomic and cDNA, RNA, or a hybrid, where the nucleic acid can contain combinations of deoxyribo- and ribo-nucleotides, and combinations of bases including uracil, adenine, thymine, cytosine, guanine, inosine, xanthine hypoxanthine, isocytosine and isoguanine. Nucleic acids can be obtained by chemical synthesis methods or by recombinant methods.

[0027] "Operably linked" as used herein means that expression of a gene is under the control of a promoter with which it is spatially connected. A promoter can be positioned 5' (upstream) or 3' (downstream) of a gene under its control. The distance between the promoter and a gene can be approximately the same as the distance between that promoter and the gene it controls in the gene from which the promoter is derived. As is known in the art, variation in this distance can be accommodated without loss of promoter function.

[0028] A "peptide," "protein," or "polypeptide" as used herein can mean a linked sequence of amino acids and can be natural, synthetic, or a modification or combination of natural and synthetic.

[0029] "Promoter" as used herein means a synthetic or naturally-derived molecule which is capable of conferring, activating or enhancing expression of a nucleic acid in a cell. A promoter can comprise one or more specific transcriptional regulatory sequences to further enhance expression

and/or to alter the spatial expression and/or temporal expression of same. A promoter can also comprise distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A promoter can be derived from sources including viral, bacterial, fungal, plants, insects, and animals. A promoter can regulate the expression of a gene component constitutively or differentially with respect to cell, the tissue or organ in which expression occurs or, with respect to the developmental stage at which expression occurs, or in response to external stimuli such as physiological stresses, pathogens, metal ions, or inducing agents. Representative examples of promoters include the bacteriophage T7 promoter, bacteriophage T3 promoter, SP6 promoter, lac operator-promoter, tac promoter, SV40 late promoter, SV40 early promoter, RSV-LTR promoter, CMV IE promoter, SV40 early promoter or SV40 late promoter and the CMV IE promoter.

[0030] “Signal peptide” and “leader sequence” are used interchangeably herein and refer to an amino acid sequence that can be linked at the amino terminus of a tumor microenvironment protein set forth herein. Signal peptides/leader sequences typically direct localization of a protein. Signal peptides/leader sequences used herein preferably facilitate secretion of the protein from the cell in which it is produced. Signal peptides/leader sequences are often cleaved from the remainder of the protein, often referred to as the mature protein, upon secretion from the cell. Signal peptides/leader sequences are linked at the N terminus of the protein.

[0031] “Subject” as used herein can mean a mammal that is capable of being administered the immunogenic compositions described herein. The mammal can be, for example, a human, chimpanzee, dog, cat, horse, cow, rabbit, groundhog, squirrel, mouse, rat, or other rodents.

[0032] “Substantially identical” as used herein can mean that a first and second amino acid sequence are at least 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% over a region of 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 300, 400, 500, 600 or more amino acids. Substantially identical can also mean that a first nucleotide sequence and a second nucleotide sequence are at least 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% over a region of 60, 65, 70, 75, 80, 85, 90, 95, 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100 or more nucleotides.

[0033] “Treatment” or “treating,” as used herein can mean protecting of a subject from a disease through means of preventing, suppressing, repressing, or completely eliminating the disease. In one embodiment, preventing the disease involves administering an immunogenic composition of the present invention to a subject prior to onset of the disease. In one embodiment, preventing the disease involves administering an immunogenic composition of the present invention to a subject following a treatment so as to prevent reoccurrence or further progression of the disease. Suppressing the disease involves administering an immunogenic composition of the present invention to a subject after induction of the disease but before its clinical appearance. Repressing the disease involves administering an immunogenic composition of the present invention to a subject after clinical appearance of the disease.

[0034] “Variant” used herein with respect to a nucleic acid means (i) a portion or fragment of a referenced nucleotide sequence; (ii) the complement of a referenced nucleotide sequence or portion thereof; (iii) a nucleic acid that is substantially identical to a referenced nucleic acid or the complement thereof; or (iv) a nucleic acid that hybridizes under stringent conditions to the referenced nucleic acid, complement thereof, or a sequences substantially identical thereto.

[0035] Variant can further be defined as a peptide or polypeptide that differs in amino acid sequence by the insertion, deletion, or conservative substitution of amino acids, but retain at least one biological activity. Representative examples of “biological activity” include the ability to be bound by a specific antibody or to promote an immune response. Variant can also mean a protein with an amino acid sequence that is substantially identical to a referenced protein with an amino acid sequence that retains at least one biological activity. A conservative substitution of an amino acid, i.e., replacing an amino acid with a different amino acid of similar properties (e.g., hydrophilicity, degree and distribution of charged regions) is recognized in the art as typically involving a minor change. These minor changes can be identified, in part, by considering the hydropathic index of amino acids, as understood in the art. Kyte et al., J. Mol. Biol. 157:105-132 (1982). The hydropathic index of an amino acid is based on a consideration of its hydrophobicity and charge. It is known in the art that amino acids of similar hydropathic indexes can be substituted and still retain protein function. In one aspect, amino acids having hydropathic indexes of ± 2 are substituted. The hydrophilicity of amino acids can also be used to reveal substitutions that would result in proteins retaining biological function. A consideration of the hydrophilicity of amino acids in the context of a peptide permits calculation of the greatest local average hydrophilicity of that peptide, a useful measure that has been reported to correlate well with antigenicity and immunogenicity. Substitution of amino acids having similar hydrophilicity values can result in peptides retaining biological activity, for example immunogenicity, as is understood in the art. Substitutions can be performed with amino acids having hydrophilicity values within ± 2 of each other. Both the hydrophobicity index and the hydrophilicity value of amino acids are influenced by the particular side chain of that amino acid. Consistent with that observation, amino acid substitutions that are compatible with biological function are understood to depend on the relative similarity of the amino acids, and particularly the side chains of those amino acids, as revealed by the hydrophobicity, hydrophilicity, charge, size, and other properties.

[0036] A variant may be a nucleotide sequence that is substantially identical over the full length of the full gene sequence or a fragment thereof. The nucleotide sequence may be 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical over the full length of the gene sequence or a fragment thereof. A variant may be an amino acid sequence that is substantially identical over the full length of the amino acid sequence or fragment thereof. The amino acid sequence may be 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% identical over the full length of the amino acid sequence or a fragment thereof.

[0037] “Vector” as used herein means a nucleic acid sequence containing an origin of replication. A vector can be a viral vector, bacteriophage, adeno-associated virus (AAV), bacterial artificial chromosome or yeast artificial chromosome. A vector can be a DNA or RNA vector. A vector can be an extrachromosomal vector, and preferably, is a DNA plasmid or mRNA.

[0038] For the recitation of numeric ranges herein, each intervening number there between with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the number 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9, and 7.0 are explicitly contemplated.

Description

[0039] Aspects of the present invention are directed to immunological compositions comprising a nucleotide sequence encoding a consensus POWV antigen described herein. The consensus POWV antigen comprises an amino acid sequence that is prME, and can include a signal sequence such as JEV signal, prM signal or a portion of prM signal. The nucleotide sequence can include a leader sequence such as an IgE leader.

[0040] Some aspects of the present invention are directed to methods of treating or preventing POWV infection in a subject in need thereof by administering the immunological compositions provided herein.

[0041] In one embodiment, the POWV antigen comprises an antigen selected from the group consisting of a POWV-prME. In some embodiments, there are nucleotide expression constructs comprising prME plus an N-terminal signal sequence, which can be a JEV signal or a prM signal, or fragments thereof. In other embodiments, one or more of the aforementioned nucleotides is also comprised of a N-terminal leader sequence, which in some embodiments can be an IgE leader sequence.

[0042] The present disclosure provides an optimized consensus sequence encoding a POWV antigen. In one embodiment, the POWV antigen encoded by the optimized consensus sequence is capable of eliciting an immune response in a mammal. In one embodiment, the POWV antigen encoded by the optimized consensus sequence can comprise an epitope(s) that makes it particularly effective as an immunogen against which an immune response can be induced.

[0043] The consensus sequence can be derived from two or more native POWV nucleic acid sequences. Modifications to the consensus sequence can include codon optimization, RNA optimization, addition of a Kozak sequence for increased translation initiation, and/or the addition of an immunoglobulin or other leader sequence to increase immunogenicity. The encoded POWV antigen can comprise a signal peptide such as an immunoglobulin signal peptide, for example, but not limited to, an immunoglobulin E (IgE) or immunoglobulin (IgG) signal peptide. The encoded antigen by the POWV sequence can be designed to elicit stronger cellular and/or humoral immune responses than a corresponding native antigen.

[0044] In one embodiment, the encoded POWV antigen is operably linked to one or more regulatory elements. In one embodiment, a regulatory element is a leader sequence. In one embodiment, the POWVDNA sequence operably linked to an IgE leader encoding sequence is set forth in SEQ ID

NO:6. In one embodiment, the optimized consensus-encoded POWV antigen operably linked to an IgE leader sequence.

[0045] In some embodiment, the POWV antigen is an immunogenic domain of one or more of the functional proteins. In some instances, the POWV antigen is the D3 domain of the envelope protein (E). In some instances, the nucleotide constructs of the invention include the D3 domain of the E protein repeated in the open reading frame, and in particular repeated three times, thereby expressing three D3 domains.

[0046] In one embodiment, a regulatory element is a start codon. Therefore, in one embodiment, the invention relates to a nucleic acid sequence as set forth in SEQ ID NO:1 (pVAX1-IgE leader-prME), SEQ ID NO:2 (pVAX1-IgE leader-prMsignal-prME), SEQ ID NO:3 (pVAX1-IgE leader-portion of prMsignal-prME), SEQ ID NO:4 (pVAX1-prM signal-prME), or SEQ ID NO:5 (pVAX1-JEV signal-prME). In one embodiment, the invention relates to an amino acid sequence as set forth in SEQ ID NO:7 prM consensus or SEQ ID NO:8 E consensus, or a fragment or homolog thereof, operably linked to an amino acid encoded by a start codon (e.g., a Methionine) at the N-terminus.

[0047] In one embodiment, a regulatory element is at least one stop codon. Therefore, in one embodiment, the invention relates to a nucleic acid sequence as set forth in SEQ ID NO:9 (consensus prME) or a fragment or homolog thereof, operably linked to a nucleotide sequence comprising at least one stop codon at the 3' terminus. In one embodiment, the nucleotide sequence is operably linked to two stop codons to increase the efficiency of translational termination.

[0048] In one embodiment, the optimized consensus sequence encoding a POWV antigen can encode a peptide having the amino acid sequence set forth in SEQ ID NO:7 or SEQ ID NO: 8. In one embodiment, the optimized consensus sequence can have the nucleotide sequence set forth in SEQ ID NO: 9, which is a part of the inserts in constructs SEQ ID NO: 1, 2, 3, 4, or 5. In some embodiments, the sequence can be the nucleotide sequence having at least about 96%, 97%, 98%, 99% or 100% identity over an entire length of the nucleotide sequence set forth in SEQ ID NO:9. In other embodiments, sequence can be the nucleotide sequence that encodes the amino acid sequence having at least about 96%, 97%, 98%, 99%, or 100% identity over an entire length of the amino acid sequence set forth in SEQ ID NO:7 or SEQ ID NO:8. In some embodiments, the optimized consensus POWV antigen can be encoded by an RNA that is a transcript from a DNA sequence having at least about 96%, 97%, 98%, 99% or 100% identity over an entire length of the nucleic acid sequence set forth in the SEQ ID NO:9. In some embodiments, the optimized consensus POWV antigen can be encoded by an RNA that encodes an amino acid sequence having at least about 96%, 97%, 98%, 99% or 100% identity over an entire length of the amino acid sequence set forth in SEQ ID NO:7 or SEQ ID NO:8.

[0049] In some embodiments, the antigen can have an amino acid sequence having at least about 96%, 97%, 98%, 99%, or 100% identity over an entire length of the amino acid sequence set forth in SEQ ID NO:7 or SEQ ID NO:8.

[0050] Immunogenic fragments of SEQ ID NO:7 or SEQ ID NO:8 can be provided. Immunogenic fragments can comprise at least at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least

99% of the full length of SEQ ID NO:7 or SEQ ID NO:8. In some embodiments, immunogenic fragments include a leader sequence, such as for example an immunoglobulin leader, such as the IgE leader. In some embodiments, immunogenic fragments are free of a leader sequence.

[0051] In one embodiment, the nucleic acid sequence comprises an RNA sequence encoding a consensus POWV immunogen sequence described herein. For example, nucleic acids may comprise an RNA sequence encoding one or more of SEQ ID NO:7 or SEQ ID NO:8, a variant thereof, a fragment thereof or any combination thereof.

[0052] Immunogenic fragments of proteins with amino acid sequences homologous to immunogenic fragments of SEQ ID NO:7 or SEQ ID NO:8 can be provided. Such immunogenic fragments can comprise at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% of proteins that are 95% homologous to SEQ ID NO:7 or SEQ ID NO:8. Some embodiments relate to immunogenic fragments that have 90% homology to the immunogenic fragments of consensus protein sequences herein. Some embodiments relate to immunogenic fragments that have 97% homology to the immunogenic fragments of consensus protein sequences herein. Some embodiments relate to immunogenic fragments that have 98% homology to the immunogenic fragments of consensus protein sequences herein. Some embodiments relate to immunogenic fragments that have 99% homology to the immunogenic fragments of consensus protein sequences herein. In some embodiments, immunogenic fragments include a leader sequence, such as for example an immunoglobulin leader, such as the IgE leader. In some embodiments, immunogenic fragments are free of a leader sequence.

[0053] Some embodiments relate to immunogenic fragments of SEQ ID NO:9. Immunogenic fragments can be at least 80%, at least 85%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98% or at least 99% of the full length of SEQ ID NO:9. Immunogenic fragments can be at least 96%, at least 97% at least 98% or at least 99% homologous to fragments of SEQ ID NO:9. In some embodiments, immunogenic fragments include sequences that encode a leader sequence, such as for example an immunoglobulin leader, such as the IgE leader. In some embodiments, fragments are free of coding sequences that encode a leader sequence.

Immunogenic Composition

[0054] Provided herein are immunogenic compositions, such as vaccines, comprising an antigenic sequence, an encoded antigen, a fragment thereof, a variant thereof, or a combination thereof.

[0055] The immunogenic composition can be a DNA vaccine, an RNA vaccine, a peptide vaccine, or a combination vaccine. The vaccine can include an optimized consensus nucleotide sequence encoding an antigen. The nucleotide sequence can be DNA, RNA, mRNA, cDNA, a variant thereof, a fragment thereof, or a combination thereof. The nucleotide sequence can also include additional sequences that encode linker, leader, or tag sequences that are linked to the antigen by a peptide bond. The peptide vaccine can include an antigen, a variant thereof, a fragment thereof, or a combination thereof. The combination DNA and peptide vaccine can include the above described optimized consensus nucleotide sequence and the encoded antigen.

[0056] The vaccine of the present invention can have features required of effective vaccines such as being safe so that the vaccine itself does not cause illness or death; being protective against illness; inducing neutralizing antibody; inducing protective T cell responses; and providing ease of administration, few side effects, and biological stability.

Immune Response

[0057] The immunogenic composition can induce an immune response in the subject administered the composition.

[0058] The immunogenic composition can further induce an immune response when administered to different tissues such as the muscle or skin. The immunogenic composition can further induce an immune response when administered via electroporation, or injection, or subcutaneously, or intramuscularly.

Fragments

[0059] In one embodiment, the immunogenic fragment is an immunogenic fragment of a full length antigen of the invention. As used herein, an immunogenic fragment is a fragment of a full length nucleic acid or amino acid sequence that can induce an immune response significantly similar to that of the full length sequence. In one embodiment, an immunogenic fragment comprises an immunogenic epitope of a full length sequence. In one embodiment, the immunogenic fragment induces an immune response at least about 0.7-fold, at least about 0.8-fold, at least about 0.9-fold, at least about 1.0-fold, at least about 1.1-fold, at least about 1.2-fold, at least about 1.3-fold, at least about 1.4-fold, at least about 1.5-fold, at least about 2.0-fold or greater than 2.0-fold as compared to the full length sequence.

[0060] The immunogenic fragment can induce a humoral immune response in the subject administered the immunogenic fragment. The humoral immune response can be induced in the subject administered the immunogenic fragment by about 1.5-fold to about 16-fold, about 2-fold to about 12-fold, or about 3-fold to about 10-fold. The humoral immune response can be induced in the subject administered the immunogenic fragment by at least about 1.5-fold, at least about 2.0-fold, at least about 2.5-fold, at least about 3.0-fold, at least about 3.5-fold, at least about 4.0-fold, at least about 4.5-fold, at least about 5.0-fold, at least about 5.5-fold, at least about 6.0-fold, at least about 6.5-fold, at least about 7.0-fold, at least about 7.5-fold, at least about 8.0-fold, at least about 8.5-fold, at least about 9.0-fold, at least about 9.5-fold, at least about 10.0-fold, at least about 10.5-fold, at least about 11.0-fold, at least about 11.5-fold, at least about 12.0-fold, at least about 12.5-fold, at least about 13.0-fold, at least about 13.5-fold, at least about 14.0-fold, at least about 14.5-fold, at least about 15.0-fold, at least about 15.5-fold, or at least about 16.0-fold as compared to a subject not administered immunogenic fragment.

[0061] The humoral immune response induced by the immunogenic fragment can include an increased level of IgG antibodies associated with the subject administered the immunogenic fragment as compared to a subject not administered the immunogenic fragment. The level of IgG antibody associated with the subject administered the immunogenic fragment can be increased by about 1.5-fold to about 16-fold, about 2-fold to about 12-fold, or about 3-fold to about 10-fold as compared to the subject not administered

the immunogenic fragment. The level of IgG antibody associated with the subject administered the immunogenic fragment can be increased by at least about 1.5-fold, at least about 2.0-fold, at least about 2.5-fold, at least about 3.0-fold, at least about 3.5-fold, at least about 4.0-fold, at least about 4.5-fold, at least about 5.0-fold, at least about 5.5-fold, at least about 6.0-fold, at least about 6.5-fold, at least about 7.0-fold, at least about 7.5-fold, at least about 8.0-fold, at least about 8.5-fold, at least about 9.0-fold, at least about 9.5-fold, at least about 10.0-fold, at least about 10.5-fold, at least about 11.0-fold, at least about 11.5-fold, at least about 12.0-fold, at least about 12.5-fold, at least about 13.0-fold, at least about 13.5-fold, at least about 14.0-fold, at least about 14.5-fold, at least about 15.0-fold, at least about 15.5-fold, or at least about 16.0-fold as compared to a subject not administered the immunogenic fragment.

[0062] The induced cellular immune response can include an increased CD8⁺ T cell response associated with the subject administered the immunogenic fragment as compared to the subject not administered the immunogenic fragment. The CD8⁺ T cell response associated with the subject administered the immunogenic fragment can be increased by about 2-fold to about 30-fold, about 3-fold to about 25-fold, or about 4-fold to about 20-fold as compared to the subject not administered the immunogenic fragment. The CD8⁺ T cell response associated with the subject administered the immunogenic fragment can be increased by at least about 1.5-fold, at least about 2.0-fold, at least about 3.0-fold, at least about 4.0-fold, at least about 5.0-fold, at least about 6.0-fold, at least about 6.5-fold, at least about 7.0-fold, at least about 7.5-fold, at least about 8.0-fold, at least about 8.5-fold, at least about 9.0-fold, at least about 9.5-fold, at least about 10.0-fold, at least about 10.5-fold, at least about 11.0-fold, at least about 11.5-fold, at least about 12.0-fold, at least about 12.5-fold, at least about 13.0-fold, at least about 13.5-fold, at least about 14.0-fold, at least about 14.5-fold, at least about 15.0-fold, at least about 16.0-fold, at least about 17.0-fold, at least about 18.0-fold, at least about 19.0-fold, at least about 20.0-fold, at least about 21.0-fold, at least about 22.0-fold, at least about 23.0-fold, at least about 24.0-fold, at least about 25.0-fold, at least about 26.0-fold, at least about 27.0-fold, at least about 28.0-fold, at least about 29.0-fold, or at least about 30.0-fold as compared to a subject not administered the immunogenic fragment.

[0063] The induced cellular immune response can include an increased frequency of CD107a/IFN γ /T-bet triple-positive CD8 T cells that are reactive against the native antigen. The frequency of CD107a/IFN γ /T-bet triple-positive CD8 T cells associated with the subject administered the immunogenic fragment can be increased by at least about 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold, 12-fold, 13-fold, 14-fold, 15-fold, 16-fold, 17-fold, 18-fold, 19-fold, or 20-fold as compared to a subject not administered the immunogenic fragment.

[0064] The induced cellular immune response can include an increased frequency of CD107a/IFN γ double-positive CD8 T cells that are reactive against the native antigen. The frequency of CD107a/IFN γ double-positive CD8 T cells associated with the subject administered the immunogenic fragment can be increased by at least about 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold, 12-fold, 13-fold, or 14-fold as compared to a subject not administered the immunogenic.

[0065] The cellular immune response induced by the immunogenic fragment can include eliciting a CD4⁺ T cell response. The elicited CD4⁺ T cell response can be reactive with the native antigen genetically related to the optimized consensus antigen. The elicited CD4⁺ T cell response can be polyfunctional. The induced cellular immune response can include eliciting a CD4⁺ T cell response, in which the CD4⁺ T cells produce IFN- γ , TNF- α , IL-2, or a combination of IFN- γ and TNF- α .

[0066] The induced cellular immune response can include an increased frequency of CD4⁺ T cells that produce IFN- γ . The frequency of CD4⁺ IFN- γ ⁺ T cells associated with the subject administered the immunogenic fragment can be increased by at least about 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold, 12-fold, 13-fold, 14-fold, 15-fold, 16-fold, 17-fold, 18-fold, 19-fold, or 20-fold as compared to a subject not administered the immunogenic fragment.

[0067] The induced cellular immune response can include an increased frequency of CD4⁺ T cells that produce TNF- α . The frequency of CD4⁺ TNF- α ⁺ T cells associated with the subject administered the immunogenic fragment can be increased by at least about 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 7-fold, 8-fold, 9-fold, 10-fold, 11-fold, 12-fold, 13-fold, 14-fold, 15-fold, 16-fold, 17-fold, 18-fold, 19-fold, 20-fold, 21-fold, or 22-fold as compared to a subject not administered the immunogenic fragment.

[0068] The induced cellular immune response can include an increased frequency of CD4⁺ T cells that produce both IFN- γ and TNF- α . The frequency of CD4⁺IFN- γ ⁺TNF- α ⁺ associated with the subject administered the immunogenic fragment can be increased by at least about 2-fold, 2.5-fold, 3.0-fold, 3.5-fold, 4.0-fold, 4.5-fold, 5.0-fold, 5.5-fold, 6.0-fold, 6.5-fold, 7.0-fold, 7.5-fold, 8.0-fold, 8.5-fold, 9.0-fold, 9.5-fold, 10.0-fold, 10.5-fold, 11.0-fold, 11.5-fold, 12.0-fold, 12.5-fold, 13.0-fold, 13.5-fold, 14.0-fold, 14.5-fold, 15.0-fold, 15.5-fold, 16.0-fold, 16.5-fold, 17.0-fold, 17.5-fold, 18.0-fold, 18.5-fold, 19.0-fold, 19.5-fold, 20.0-fold, 21-fold, 22-fold, 23-fold, 24-fold, 25-fold, 26-fold, 27-fold, 28-fold, 29-fold, 30-fold, 31-fold, 32-fold, 33-fold, 34-fold, or 35-fold as compared to a subject not administered the immunogenic fragment.

[0069] The immunogenic fragment of the present invention can have features required of effective vaccines such as being safe so the vaccine itself does not cause illness or death; is protective against illness resulting from exposure to live pathogens such as viruses or bacteria; induces neutralizing antibody to prevent infection of cells; induces protective T cells against intracellular pathogens; and provides ease of administration, few side effects, biological stability, and low cost per dose.

[0070] The immunogenic fragment can further induce an immune response when administered to different tissues such as the muscle or skin. The immunogenic fragment can further induce an immune response when administered via electroporation, or injection, or subcutaneously, or intramuscularly.

Vector

[0071] The nucleotide construct described above can be placed in one or more vectors. The one or more vectors can contain an origin of replication. The one or more vectors can be a plasmid, RNA, viral, including vesicular stomatitis virus, measles virus, modified vaccinia Ankara virus, adeno-

virus, AAV, bacteriophage, bacterial artificial chromosome or yeast artificial chromosome. The one or more vectors can be either a self-replication extra chromosomal vector.

[0072] Vectors include, but are not limited to, plasmids, expression vectors, recombinant viruses, any form of recombinant “naked DNA” vector, and the like. A “vector” comprises a nucleic acid which can infect, transfect, transiently or permanently transduce a cell. It will be recognized that a vector can be a naked nucleic acid, or a nucleic acid complexed with protein or lipid. The vector optionally comprises viral or bacterial nucleic acids and/or proteins, and/or membranes (e.g., a cell membrane, a viral lipid envelope, etc.). Vectors include, but are not limited to replicons (e.g., RNA replicons, bacteriophages) to which fragments of DNA may be attached and become replicated. Vectors thus include, but are not limited to RNA, autonomous self-replicating circular or linear DNA or RNA (e.g., plasmids, viruses, and the like, see, e.g., U.S. Pat. No. 5,217,879), and include both the expression and non-expression plasmids. Where a recombinant microorganism or cell culture is described as hosting an “expression vector” that may be extra-chromosomal circular or linear DNA. Where a vector is being maintained by a host cell, the vector may either be stably replicated by the cells during mitosis as an autonomous structure, or is incorporated within the host’s genome.

[0073] The one or more vectors can be an expression construct, which is generally a plasmid that is used to introduce a specific gene into a target cell. Once the expression vector is inside the cell, the protein that is encoded by the gene is produced by the cellular-transcription and translation machinery ribosomal complexes. The plasmid is frequently engineered to contain regulatory sequences that act as enhancer and promoter regions and lead to efficient transcription of the gene carried on the expression vector.

[0074] The vectors may have expression signals such as a strong promoter, a strong termination codon, adjustment of the distance between the promoter and the cloned gene, and the insertion of a transcription termination sequence and a PTIS (portable translation initiation sequence).

Expression Vectors

[0075] The vector can be a circular plasmid or a linear nucleic acid. The circular plasmid and linear nucleic acid are capable of directing expression of a particular nucleotide sequence in an appropriate subject cell. The vector can have a promoter operably linked to the antigen-encoding nucleotide sequence, which may be operably linked to termination signals. The vector can also contain sequences required for proper translation of the nucleotide sequence. The vector comprising the nucleotide sequence of interest may be chimeric, meaning that at least one of its components is heterologous with respect to at least one of its other components. The expression of the nucleotide sequence in the expression cassette may be under the control of a constitutive promoter or of an inducible promoter, which initiates transcription only when the host cell is exposed to some particular external stimulus. In the case of a multicellular organism, the promoter can also be specific to a particular tissue or organ or stage of development.

Plasmid

[0076] The one or more vectors can be a plasmid. The plasmid may be useful for transfecting cells with the recom-

binant nucleic acid construct. The plasmid may be useful for introducing the recombinant nucleic acid construct into the subject. The plasmid may also comprise a regulatory sequence, which may be well suited for gene expression in a cell into which the plasmid is administered.

[0077] The plasmid may also comprise a mammalian origin of replication in order to maintain the plasmid extra-chromosomally and produce multiple copies of the plasmid in a cell. The plasmid may be pVAX1, pCEP4 or pREP4 from Invitrogen (San Diego, CA), which may comprise the Epstein Barr virus origin of replication and nuclear antigen EBNA-1 coding region, which may produce high copy episomal replication without integration. The backbone of the plasmid may be pAV0242. The plasmid may be a replication defective adenovirus type 5 (Ad5) plasmid.

[0078] The plasmid can also be pIDV or pIDV-II.

[0079] The plasmid may be pSE420 (Invitrogen, San Diego, Calif.), which may be used for protein production in *Escherichia coli* (*E.coli*). The plasmid may also be pYES2 (Invitrogen, San Diego, Calif.), which may be used for protein production in *Saccharomyces cerevisiae* strains of yeast. The plasmid may also be of the MAXBAC™ complete baculovirus expression system (Invitrogen, San Diego, Calif.), which may be used for protein production in insect cells. The plasmid may also be pcDNA1 or pcDNA3 (Invitrogen, San Diego, Calif.), which may be used for protein production in mammalian cells such as Chinese hamster ovary (CHO) cells.

RNA

[0080] In one embodiment, the nucleic acid is an RNA molecule. In one embodiment, the RNA molecule is transcribed from a DNA sequence described herein. For example, in some embodiments, the RNA molecule is encoded by a DNA sequence at least 90% homologous to SEQ ID NO:9, or a variant thereof or a fragment thereof. In another embodiment, the nucleotide sequence comprises an RNA sequence transcribed by a DNA sequence encoding a polypeptide sequence at least 90% homologous to one of SEQ ID NO:7 or SEQ ID NO:8 or a variant thereof or a fragment thereof. Accordingly, in one embodiment, the invention provides an RNA molecule encoding one or more of the POWV antigens. The RNA may be plus-stranded. Accordingly, in some embodiments, the RNA molecule can be translated by cells without needing any intervening replication steps such as reverse transcription. A RNA molecule useful with the invention may have a 5' cap (e.g. a 7-methylguanosine). This cap can enhance in vivo translation of the RNA. The 5' nucleotide of a RNA molecule useful with the invention may have a 5' triphosphate group. In a capped RNA this may be linked to a 7-methylguanosine via a 5'-to-5' bridge. A RNA molecule may have a 3' poly-A tail. It may also include a poly-A polymerase recognition sequence (e.g. AAUAAA) near its 3' end. A RNA molecule useful with the invention may be single-stranded. A RNA molecule useful with the invention may comprise synthetic RNA. In some embodiments, the RNA molecule is a naked RNA molecule. In one embodiment, the RNA molecule is comprised within a vector.

[0081] In one embodiment, the RNA has 5' and 3' UTRs. In one embodiment, the 5' UTR is between zero and 3000 nucleotides in length. The length of 5' and 3' UTR sequences to be added to the coding region can be altered by different methods, including, but not limited to, designing primers for

PCR that anneal to different regions of the UTRs. Using this approach, one of ordinary skill in the art can modify the 5' and 3' UTR lengths required to achieve optimal translation efficiency following transfection of the transcribed RNA.

[0082] The 5' and 3' UTRs can be the naturally occurring, endogenous 5' and 3' UTRs for the gene of interest. Alternatively, UTR sequences that are not endogenous to the gene of interest can be added by incorporating the UTR sequences into the forward and reverse primers or by any other modifications of the template. The use of UTR sequences that are not endogenous to the gene of interest can be useful for modifying the stability and/or translation efficiency of the RNA. For example, it is known that AU-rich elements in 3' UTR sequences can decrease the stability of RNA. Therefore, 3' UTRs can be selected or designed to increase the stability of the transcribed RNA based on properties of UTRs that are well known in the art.

[0083] In one embodiment, the 5' UTR can contain the Kozak sequence of the endogenous gene. Alternatively, when a 5'UTR that is not endogenous to the gene of interest is being added by PCR as described above, a consensus Kozak sequence can be redesigned by adding the 5' UTR sequence. Kozak sequences can increase the efficiency of translation of some RNA transcripts, but does not appear to be required for all RNAs to enable efficient translation. The requirement for Kozak sequences for many RNAs is known in the art. In other embodiments, the 5' UTR can be derived from an RNA virus whose RNA genome is stable in cells. In other embodiments, various nucleotide analogues can be used in the 3' or 5' UTR to impede exonuclease degradation of the RNA.

[0084] In one embodiment, the RNA has both a cap on the 5' end and a 3' poly(A) tail which determine ribosome binding, initiation of translation and stability of RNA in the cell.

[0085] In one embodiment, the RNA is a nucleoside-modified RNA. Nucleoside-modified RNA have particular advantages over non-modified RNA, including for example, increased stability, low or absent innate immunogenicity, and enhanced translation.

Circular and Linear Vectors

[0086] The vector may be a circular plasmid, which may transform a target cell and exist extrachromosomally (e.g., autonomous replicating plasmid with an origin of replication).

[0087] The vector can be pVAX, pcDNA3.0, or provax, or any other expression vector capable of expressing DNA encoding the antigen and enabling a cell to translate the sequence to an antigen that is recognized by the immune system.

[0088] Also provided herein is a linear nucleic acid immunogenic composition, or linear expression cassette ("LEC"), that is capable of being efficiently delivered to a subject and expressing one or more desired antigens. The LEC may be any linear DNA devoid of any phosphate backbone. The DNA may encode one or more antigens. The LEC may contain a promoter, an intron, a stop codon, and/or a polyadenylation signal. The expression of the antigen may be controlled by the promoter. The LEC may not contain any antibiotic resistance genes and/or a phosphate backbone. The LEC may not contain other nucleotide sequences unrelated to the desired antigen gene expression.

[0089] The LEC may be derived from any plasmid capable of being linearized. The plasmid may be capable of expressing the antigen. The plasmid can be pNP (Puerto Rico/34) or pM2 (New Caledonia/99). The plasmid may be pVAX, pVAX1 pcDNA3.0, pIDV, pIDV-II, or provax, or any other expression vector capable of expressing DNA encoding the antigen and enabling a cell to translate the sequence to an antigen that is recognized by the immune system.

[0090] The LEC can be pcrM2. The LEC can be perNP, pcrNP and pcrMR can be derived from pNP (Puerto Rico/34) and pM2 (New Caledonia/99), respectively.

Promoter, Intron, Stop Codon, and Polyadenylation Signal

[0091] The vector may have a promoter. A promoter may be any promoter that is capable of driving gene expression and regulating expression of the isolated nucleic acid. Such a promoter is a cis-acting sequence element required for transcription via a DNA dependent RNA polymerase, which transcribes the antigen sequence described herein. Selection of the promoter used to direct expression of a heterologous nucleic acid depends on the particular application. The promoter may be positioned about the same distance from the transcription start in the vector as it is from the transcription start site in its natural setting. However, variation in this distance may be accommodated without loss of promoter function.

[0092] The promoter may be operably linked to the nucleotide sequence encoding the antigen and signals required for efficient polyadenylation of the transcript, ribosome binding sites, and translation termination. The promoter may be a CMV promoter, SV40 early promoter, SV40 later promoter, metallothionein promoter, murine mammary tumor virus promoter, CAG promoter, Rous sarcoma virus promoter, polyhedrin promoter, or another promoter shown effective for expression in eukaryotic cells.

[0093] The vector may include an enhancer and an intron with functional splice donor and acceptor sites. The vector may contain a transcription termination region downstream of the structural gene to provide for efficient termination. The termination region may be obtained from the same gene as the promoter sequence or may be obtained from different genes.

Viral Vectors

[0094] In one embodiment, viral vectors are provided herein which are capable of delivering a nucleic acid of the invention to a cell. The expression vector may be provided to a cell in the form of a viral vector. Viral vector technology is well known in the art and is described, for example, in Sambrook et al. (2001), and in Ausubel et al. (1997), and in other virology and molecular biology manuals. Viruses, which are useful as vectors include, but are not limited to, adenoviruses, adeno-associated viruses, measles virus, vesicular stomatitis virus, modified vaccinia Ankara virus. In general, a suitable vector contains an origin of replication functional in at least one organism, a promoter sequence, convenient restriction endonuclease sites, and one or more selectable markers. (See, e.g., WO 01/96584; WO 01/29058; and U.S. Pat. No. 6,326,193.

Multiple Vectors

[0095] The immunogenic composition may comprise a plurality of copies of a single nucleic acid molecule such a

single plasmid, or a plurality of copies of two or more different nucleic acid molecules such as two or more different plasmids. For example an immunogenic composition may comprise plurality of two, three, four, five, six, seven, eight, nine or ten or more different nucleic acid molecules. Such compositions may comprise plurality of two, three, four, five, six, or more different plasmids.

[0096] Immunogenic compositions may comprise nucleic acid molecules, such as plasmids, that collectively contain coding sequence for a POWV antigen. Immunogenic compositions may comprise nucleic acid molecules, such as plasmids, that collectively contain coding sequence for multiple antigens. In one embodiment, the antigens are a POWV antigen and one or more additional adjuvants. Immunogenic compositions may comprise nucleic acid molecules, such as plasmids, that collectively contain coding sequence for one or more antigen and one or more adjuvant.

Methods

[0097] Provided herein are methods of treating, protecting against, and/or preventing a POWV associated disease in a subject in need thereof by administering one or more immunogenic composition described herein to the subject. Administration of the immunogenic composition to the subject can induce or elicit an immune response in the subject.

[0098] Provided herein is a method for delivering the immunogenic composition for providing genetic constructs and proteins of the consensus antigen which comprise epitopes that make them particular effective against POWV. The method of delivering the immunogenic composition or vaccination may be provided to induce a therapeutic and prophylactic immune response. The vaccination process may generate in the mammal an immune response against POWV. The immunogenic composition may be delivered to an individual to modulate the activity of the mammal's immune system and enhance the immune response. The delivery of the immunogenic composition may be the transfection of the consensus antigen as a nucleic acid molecule that is expressed in the cell and delivered to the surface of the cell upon which the immune system recognized and induces a cellular, humoral, or cellular and humoral response. The delivery of the immunogenic composition may be used to induce or elicit an immune response in mammals against POWV by administering to the mammals the immunogenic composition as discussed above.

[0099] Upon delivery of the immunogenic composition and plasmid into the cells of the mammal, the transfected cells will express and secrete consensus antigens for each of the plasmids injected from the immunogenic composition. These proteins will be recognized as foreign by the immune system and antibodies will be made against them. These antibodies will be maintained by the immune system and allow for an effective response.

[0100] The immunogenic composition may be administered to a mammal to elicit an immune response in a mammal. The mammal may be human, primate, non-human primate, cow, cattle, sheep, goat, antelope, water buffalo, bison, bovids, deer, hedgehogs, rabbits, mice, rats, groundhogs, squirrels, other rodents, and chicken.

[0101] The induced immune response can include an induced humoral immune response and/or an induced cellular immune response. The humoral immune response can be induced by about 1.5-fold to about 16-fold, about 2-fold to about 12-fold, or about 3-fold to about 10-fold. The

induced cellular immune response can include a CD8⁺ T cell response, which is induced by about 2-fold to about 30-fold, about 3-fold to about 25-fold, or about 4-fold to about 20-fold.

[0102] The immunogenic composition dose can be between 1 µg to 10 mg active component/kg body weight/time, and can be 20 µg to 10 mg component/kg body weight/time. The immunogenic composition can be administered every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, or 31 days. The number of immunogenic composition doses for effective treatment can be 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10.

[0103] The immunogenic composition can be formulated in accordance with standard techniques well known to those skilled in the pharmaceutical art. Such compositions can be administered in dosages and by techniques well known to those skilled in the medical arts taking into consideration such factors as the age, sex, weight, and condition of the particular subject, and the route of administration.

[0104] The immunogenic composition can be administered prophylactically or therapeutically. In prophylactic administration, the immunogenic compositions can be administered in an amount sufficient to induce an immune response. In therapeutic applications, the immunogenic compositions are administered to a subject in need thereof in an amount sufficient to elicit a therapeutic effect. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the particular composition of the immunogenic composition regimen administered, the manner of administration, the stage and severity of the disease, the general state of health of the subject, and the judgment of the prescribing physician.

[0105] The immunogenic composition can be administered by methods well known in the art as described in Donnelly et al. (Ann. Rev. Immunol. 15:617-648 (1997)); Felgner et al. (U.S. Pat. No. 5,580,859, issued Dec. 3, 1996); Felgner (U.S. Pat. No. 5,703,055, issued Dec. 30, 1997), and Carson et al. (U.S. Pat. No. 5,679,647, issued Oct. 21, 1997), the contents of all of which are incorporated herein by reference in their entirety. The nucleic acid of the immunogenic composition can be complexed to particles or beads that can be administered to an individual, for example, using a vaccine gun. One skilled in the art would know that the choice of a pharmaceutically acceptable carrier, including a physiologically acceptable compound, depends, for example, on the route of administration of the expression vector.

[0106] The immunogenic composition can be delivered via a variety of routes. Typical delivery routes include parenteral administration, e.g., intradermal, intraepidermal, intramuscular or subcutaneous delivery. Other routes include oral administration, intranasal, and intravaginal routes. For the nucleic acid of the immunogenic composition in particular, the immunogenic composition can be delivered to the interstitial spaces of tissues of an individual (Felgner et al., U.S. Pat. Nos. 5,580,859 and 5,703,055, the contents of all of which are incorporated herein by reference in their entirety). The immunogenic composition can also be administered to muscle, or can be administered via intradermal or subcutaneous injections, or transdermally, such as by iontophoresis. Epidermal administration of the immunogenic composition can also be employed. Epidermal administration can involve mechanically or chemically irritating the

outermost layer of epidermis to stimulate an immune response to the irritant (Carson et al., U.S. Pat. No. 5,679,647, the contents of which are incorporated herein by reference in its entirety).

[0107] The immunogenic composition can also be formulated for administration via the nasal passages. Formulations suitable for nasal administration, wherein the carrier is a solid, can include a coarse powder having a particle size, for example, in the range of about 10 to about 500 microns which is administered in the manner in which snuff is taken, i.e., by rapid inhalation through the nasal passage from a container of the powder held close up to the nose. The formulation can be a nasal spray, nasal drops, or by aerosol administration by nebulizer. The formulation can include aqueous or oily solutions of the immunogenic composition.

[0108] The immunogenic composition can be a liquid preparation such as a suspension, syrup or elixir. The immunogenic composition can also be a preparation for parenteral, subcutaneous, intradermal, intramuscular or intravenous administration (e.g., injectable administration), such as a sterile suspension or emulsion.

[0109] The immunogenic composition can be incorporated into liposomes, microspheres or other polymer matrices (Felgner et al., U.S. Pat. No. 5,703,055; Gregoriadis, *Liposome Technology*, Vols. I to III (2nd ed. 1993), the contents of which are incorporated herein by reference in their entirety). Liposomes can consist of phospholipids or other lipids, and can be nontoxic, physiologically acceptable and metabolizable carriers that are relatively simple to make and administer.

Combination Treatments

[0110] The immunogenic composition may be administered in combination with other proteins and/or genes encoding CCL20, α -interferon, γ -interferon, platelet derived growth factor (PDGF), TNF α , TNF β , GM-CSF, epidermal growth factor (EGF), cutaneous T cell-attracting chemokine (CTACK), epithelial thymus-expressed chemokine (TECK), mucosae-associated epithelial chemokine (MEC), IL-12, IL-15 including IL-15 having the signal sequence deleted and optionally including the different signal peptide such as the IgE signal peptide, MHC, CD80, CD86, IL-28, IL-1, IL-2, IL-4, IL-5, IL-6, IL-10, IL-18, MCP-1, MIP-1a, MIP-1B, IL-8, RANTES, L-selectin, P-selectin, E-selectin, CD34, GlyCAM-1, MadCAM-1, LFA-1, VLA-1, Mac-1, p150.95, PECAM, ICAM-1, ICAM-2, ICAM-3, CD2, LFA-3, M-CSF, G-CSF, mutant forms of IL-18, CD40, CD40L, vascular growth factor, fibroblast growth factor, IL-7, nerve growth factor, vascular endothelial growth factor, Fas, TNF receptor, Flt, Apo-1, p55, WSL-1, DR3, TRAMP, Apo-3, AIR, LARD, NGRF, DR4, DR5, KILLER, TRAIL-R2, TRICK2, DR6, Caspase ICE, Fos, c-jun, Sp-1, Ap-1, Ap-2, p38, p65Rel, MyD88, IRAK, TRAF6, I κ B, Inactive NIK, SAP K, SAP-1, JNK, interferon response genes, NFKB, Bax, TRAIL, TRAILrec, TRAILrecDRCS, TRAIL-R3, TRAIL-R4, RANK, RANK LIGAND, OX40, OX40 LIGAND, NKG2D, MICA, MICB, NKG2A, NKG2B, NKG2C, NKG2E, NKG2F, TAP1, TAP2 and functional fragments thereof or combinations thereof. In some embodiments, the immunogenic composition is administered in combination with one or more of the following nucleic acid molecules and/or proteins: nucleic acid molecules selected from the group consisting of nucleic acid molecules comprising coding sequence that encode one or more of CCL20,

IL-12, IL-15, IL-28, CTACK, TECK, MEC and RANTES or functional fragments thereof, and proteins selected from the group consisting of: CCL20, IL-12 protein, IL-15 protein, IL-28 protein, CTACK protein, TECK protein, MEC protein or RANTES protein or functional fragments thereof.

Routes of Administration

[0111] The vaccine or pharmaceutical composition can be administered by different routes including orally, parenterally, sublingually, transdermally, rectally, transmucosally, topically, via inhalation, via buccal administration, intrapleurally, intravenous, intraarterial, intraperitoneal, subcutaneous, intramuscular, intranasal, intrathecal, intradermal, intraepidermal, and intraarticular or combinations thereof. For veterinary use, the composition can be administered as a suitably acceptable formulation in accordance with normal veterinary practice. The veterinarian can readily determine the dosing regimen and route of administration that is most appropriate for a particular animal. The vaccine can be administered by traditional syringes, needleless injection devices, microneedle device, "microprojectile bombardment gene guns", or other physical methods such as electroporation ("EP"), "hydrodynamic method", or ultrasound.

[0112] The vector of the vaccine can be administering to the mammal by several well-known technologies including DNA injection (also referred to as DNA vaccination) with and without in vivo electroporation, liposome mediated, nanoparticle facilitated, recombinant vectors such as recombinant adenovirus, recombinant adenovirus associated virus and recombinant vaccinia. The one or more POWV constructs can be administered via DNA injection, and can optionally be accompanied by methods to enhance cellular uptake that may include magnetoporation, ultrasound, heat, electroporation, physical methods such as tattooing.

[0113] The immunogenic composition may be administered by different routes including orally, parenterally, sublingually, transdermally, rectally, transmucosally, topically, via inhalation, via buccal administration, intrapleurally, intravenous, intraarterial, intraperitoneal, subcutaneous, intramuscular, intranasal, intrathecal, and intraarticular or combinations thereof.

[0114] The immunogenic composition may be administered by traditional syringes, needleless injection devices, "microprojectile bombardment gene guns", aerosol bombardment devices, or other physical methods such as electroporation ("EP"), "hydrodynamic method", or ultrasound. Electroporation apparatuses capable of performing the in vivo electroporation for transfecting in situ cells with one or more of the powassan constructs described herein include any commercially available electroporation devices. In addition, other publicly describe electroporation devices may be utilized upon configuring the device to perform the desired electroporation to cause reversible member pores to form and maintain the viability of the cells. The electroporation devices can include such devices described and made or utilized by the following companies: Inovio Pharmaceuticals (Collectra), Collectis (Easy Vax), Ichor (Trigrig), IGEA (CliniPorator), BTX Harvard, among others.

[0115] The plasmid of the immunogenic composition may be delivered to the mammal by several well-known technologies including DNA or RNA injection (also referred to as nucleic acid vaccination) with and without in vivo electroporation, liposome mediated, nanoparticle facilitated,

recombinant vectors such as recombinant adenovirus, recombinant adenovirus associated virus and recombinant vaccinia.

Generation of Antigens In Vitro and Ex Vivo

[0116] In one embodiment, the optimized consensus POWV antigen is generated in vitro or ex vivo. For example, in one embodiment, a nucleic acid encoding an optimized consensus POWV antigen can be introduced and expressed in an in vitro or ex vivo cell.

[0117] Methods of introducing and expressing genes into a cell are known in the art. In the context of an expression vector, the vector can be readily introduced into a host cell, e.g., mammalian, bacterial, yeast, or insect cell by any method in the art. For example, the expression vector can be transferred into a host cell by physical, chemical, or biological means.

[0118] Physical methods for introducing a polynucleotide into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Methods for producing cells comprising vectors and/or exogenous nucleic acids are well-known in the art. See, for example, Sambrook et al. (2012, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, New York). Preferred methods for the introduction of a polynucleotide into a host cell are calcium phosphate transfection, lipid nanoparticles, and silica nanoparticles.

[0119] Biological methods for introducing a polynucleotide of interest into a host cell include the use of DNA and RNA vectors. Viral vectors, and especially retroviral vectors, have become the most widely used method for inserting genes into mammalian, e.g., human cells. Other viral vectors can be derived from lentivirus, poxviruses, herpes simplex virus I, adenoviruses and adeno-associated viruses, and the like. See, for example, U.S. Pat. Nos. 5,350,674 and 5,585,362.

[0120] Chemical means for introducing a polynucleotide into a host cell include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. An exemplary colloidal system for use as a delivery vehicle in vitro and in vivo is a liposome (e.g., an artificial membrane vesicle).

[0121] In the case where a non-viral delivery system is utilized, an exemplary delivery vehicle is a liposome. The use of lipid formulations is contemplated for the introduction of the nucleic acids into a host cell (in vitro, ex vivo or in vivo). In another aspect, the nucleic acid may be associated with a lipid. The nucleic acid associated with a lipid may be encapsulated in the aqueous interior of a liposome, interspersed within the lipid bilayer of a liposome, attached to a liposome via a linking molecule that is associated with both the liposome and the oligonucleotide, entrapped in a liposome, complexed with a liposome, dispersed in a solution containing a lipid, mixed with a lipid, combined with a lipid, contained as a suspension in a lipid, contained or complexed with a micelle, or otherwise associated with a lipid. Lipid, lipid/DNA or lipid/expression vector associated compositions are not limited to any particular structure in solution. For example, they may be present in a bilayer structure, as micelles, or with a “collapsed” structure. They may also simply be interspersed in a solution, possibly

forming aggregates that are not uniform in size or shape. Lipids are fatty substances which may be naturally occurring or synthetic lipids. For example, lipids include the fatty droplets that naturally occur in the cytoplasm as well as the class of compounds which contain long-chain aliphatic hydrocarbons and their derivatives, such as fatty acids, alcohols, amines, amino alcohols, and aldehydes.

EXPERIMENTAL EXAMPLES

[0122] The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

[0123] Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the present invention and practice the claimed methods. The following working examples therefore, specifically point out the preferred embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

EXAMPLES

Example 1 Origin of Plasmid Backbone

[0124] pVAX1 (Invitrogen V260-20) was purchased from Thermo Fisher Scientific and a master cell bank was established in *Escherichia coli* strain DH10B-TIR. The purified plasmid DNA derived from the master cell bank was delivered to GenScript for constructing five powassan virus (POWV) DNA vaccines described, below.

Example 2 Construction of Powassan virus (POWV) DNA Vaccines

[0125] Five POWV DNA candidate vaccines were designed and constructed with the same antigen sequence, which encodes the POWV pre-membranemembrane, and envelope (prME) genes with different signal peptide sequences and with and without leader sequence.

[0126] The consensus prME protein coding sequence was generated from 22 different POWV prM protein sequences and 37 different POWV E protein sequences using MegAlign software (DNASTAR, Madison, WI). The signal peptide sequence of the five constructs contained either (1) an IgE leader sequence alone, (2) an IgE leader sequence plus the 20 amino acid consensus prM signal sequence, (3) an IgE leader plus the first 12 amino acids of the prM signal sequence, (4) the 20 amino acid consensus prM signal sequence, and (5) a JEV signal sequence, respectively. The prM signal sequence was generated as a consensus sequence from 22 different POWV prM signal peptide sequences.

[0127] The Genbank accession numbers used for generating the consensus envelope (E) sequences are: AAL32141.1, AAL32142.1, AAL32143.1, AAL32144.1, AAL32145.1, AAL32146.1, AAL32147.1, AAL32148.1, AAL32149.1, AAL32150.1, AAL32151.1, AAL32152.1, AAL32153.1, AAL32154.1, AAL32169.1, AAK61347.1, ACD67873.1, ACD88752.1, ACF05267.1, ADK37752.1, ADK37753.1, ADK37754.1, ADK37755.1, ADK37756.1, ADK37757.1,

AEN83681.1, AEN83682.1, AIG95651.1, ALP82430.1, ALP82431.1, AAA02739.1, AMY50382.1, AMZ80134.1, AMQ49162.1, AMQ49164.1, AMQ49165.1, AMQ49163.1
[0128] The Genbank accession numbers used for generating the consensus pre-membrane (prM) and prM signal sequences are: AAL32169.1, AAK61347.1, ACD67873.1, ACD88752.1, ACF05267.1, ADK37752.1, ADK37753.1, ADK37754.1, ADK37755.1, ADK37756.1, ADK37757.1, AEN83681.1, AEN83682.1, AIG95651.1, ALP82430.1, ALP82431.1, AAA02739.1, AMY50382.1, AMQ49162.1, AMQ49164.1, AMQ49165.1, AMQ49163.1

[0129] Once the consensus prME sequence was obtained, each signal peptide sequence (with and without leader sequence) of the five constructs was added to the N-terminal to facilitate expression and mRNA export, and further modified with upstream Kozak sequence. This sequence was then codon optimized and synthesized by GenScript. The synthesized insert was subcloned into the pVAX1 at the BamHI and XhoI sites, creating constructs:

- [0130]** Candidate 1 pVAX1-IgE-prME (SEQ ID NO:1)
- [0131]** Candidate 2 pVAX1-IgE-prMsignal-prME (SEQ ID NO:2)
- [0132]** Candidate 3 pVAX1-IgE-portion of prMsignal-prME (SEQ ID NO:3)
- [0133]** Candidate 4 pVAX1-prMsignal-prME (SEQ ID NO:4)
- [0134]** Candidate 5 pVAX1-JEVsignal-prME (SEQ ID NO:5)

Example 3 Detection of Expressed POWV prM Protein in Transfected Cells

[0135] To assess expression of the vaccine candidates, HEK293T cells were transfected with POWV DNA vaccines, respectively, using X-fect reagent (Clontech, Takara Korea Biomedical, Inc., Seoul Korea) and the cell supernatants were collected 48 hours after transfection. Total protein (20 µg) was subjected to a SDS-PAGE under reducing condition and analyzed by western blot using POWV prM polyclonal antibody (GeneTex Inc., Irvine, CA). The Western blot data indicated that four of the five POWV DNA vaccines expressed POWV-prM protein having approx. 18 KDa. One vaccine candidate containing two complete signal sequences, the IgE-leader sequence plus prM signal sequence, did not show evidence of prM expression (FIG. 1.).

Example 4. Induction of POWV Specific Binding Antibody in Mice by the POWV DNA Vaccination

[0136] POWV prME antigen specific binding antibody in sera from immunized C57BL/6 mice was measured by

ELISA using recombinant POWV-E (rPOWV-E) (Bioclone Inc., Sandiego, CA) as a coating antigen and the ability of the POWV vaccines to induce antibody immune response was evaluated between groups. Mice in each group were immunized with 25 µg of POWV DNA vaccine by intramuscular (i.m.) electroporation for 3 times 2 weeks apart. The sera were collected at day 0 (1 day before the first injection), day 21 (1 week after the second injection), day 35 (1 week after the third injection), and day 60 (approx. 4 weeks after the third injection).

[0137] POWV specific antibody levels were significantly elevated in all groups on day 21, 1 week after the second immunization, except pVAX1 control group. However, in pVAX1-JEV signal-prME group, POWV specific humoral responses were not further boosted by third immunization (FIG. 2.).

[0138] The data sorted by timepoint was shown that binding antibody against rPOWV-E was markedly increased in pVAX1-IgE leader-portion of prM signal-prME group on day 35, 1 week after the third immunization, and maintained for 4 more weeks thereafter. This result suggests that POWV vaccine having the IgE leader and the portion of prM signal sequence together as signal peptide sequence is more potent than other groups in inducing POWV-E specific binding antibodies (FIG. 3.).

[0139] The endpoint titer was determined as the highest dilution of sera meeting the cut-off value which is absorbance above four times the mean of naïve (preimmunization) and 0.15. The cut-off value applied in this study is more stringent than the 'mean +2 or 3 SD' or the '2 or 3 mean' reading that generally used in ELISA.

[0140] As shown in FIG. 4., the group immunized with POWV vaccine having IgE leader-portion of prM signal sequence was distinguished from other groups in terms of high endpoint titers on day 35 and 60. And interestingly, two groups, pVAX1-IgE leader-prME and pVAX1-IgE leader-portion of prM signal-prME, represented time dependent increase of humoral responses and the high endpoint titers was on day 60 (FIG. 4.).

[0141] It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents.

[0142] Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, compositions, formulations, or methods of use of the invention, may be made without departing from the spirit and scope thereof.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1

<211> LENGTH: 4997

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pVAX1-IgE-prME

<400> SEQUENCE: 1

-continued

gctgcttcgc gatgtacggg ccagatatat gcgttgacat tgattattga ctagttatta	60
atagtaatca attacggggg cattagtcca tagcccatat atggagttcc gcgttacata	120
acttacggta aatggcccg ctaggtgacc gcccaacgac ccccgcccat tgacgtcaat	180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga	240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtaogcc	300
ccctattgac gtcaatgacg gtaaatggcc cgcctggcat tatgccagat acatgacctt	360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgctatta ccatggtgat	420
gcggttttgg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag	480
tctccacccc attgacgtca atgggagttt gttttggcac caaaatcaac gggactttcc	540
aaaatgtcgt aacaactccg ccccatgac gcaaatgggc ggtaggcgtg tacgggtgga	600
ggtctatata agcagagctc tctggctaac tagagaaccc actgcttact ggcttatcga	660
aattaatacg actcactata gggagaccca agctggctag cgtttaaaact taagcttggg	720
accgagctcg gatccgccac catggactgg acctggatcc tgttcctggg ggcagcagca	780
acaagggtgc acagcaccac aatccacagg gaccgcgagg gctacatggt catgagggca	840
agcgaagag atgcagcatc ccagggtgcgc gtgcagaacg gaacctgcgt gatcctggcc	900
acagacatgg gcgagtgggt cgaggattcc atcacctact cttgcgtgac aatcgaccag	960
gaggaggagc ctgtggacgt ggattgcttt tgtaggggag tggatcgctg gaagctggag	1020
tatggccggt gtggcagaca ggcaggcagc cggggcaacg ggagcgtggt catcccaacc	1080
cacgcacaga aggacatggt ggggaaggga cagcagatggc tgaaggcgga caacatcagg	1140
gatcacgtga cccgcgtgga gggctggatg tggagaata agctgctgac agcagcaatc	1200
gtggccctgg catggctgat ggtggactct tggatggccc gggtagacct gatcctgctg	1260
gcccctgagc tgggacctgt gtacgcaacc aggtgcacac acctggagaa ccgcgatttc	1320
gtgaccggca cacagggcac cacaagagtg tccctgggtc tggagctggg aggatgcgtg	1380
accatcacag cagagggcaa gccaaagcatc gacgtgtggc tggaggatat ctttcaggag	1440
tcccagcag agacccggga gtattgctg cagcccaagc tgaccaatac aaaggtggag	1500
gccagatgac ccaccacagg cctgcccaca ctgccagagg agcaccaggc caacatggtg	1560
tgcaagcggg accagtccga tagaggctgg ggcaatcact gtggcttctt tggcaagggc	1620
tctatcgtgg cctgcgcca gttcgagtgt gaggaggcca agaaggccgt gggccacgtg	1680
tacgactcta ccaagatcac atatgtggtg aagggtggag cacacaccgg cgattacctg	1740
gcagccaacg agacaaacag caataggaag tccgcccagt ttaccgtggc cagcgagaaa	1800
gtgatcctgc gcctgggcca ctatggcgac gtgagcctga cctgcaaggt ggcaagcgga	1860
atcgacgtgg cacagacagt ggtcatgtct ctgggcagct ccaaggatca cctgcctagc	1920
gcctggcagg tgcacaggga ctgggtcgag gatctggccc tgccatggaa gcacaaggac	1980
aaccaggatt ggaattccgt ggagaagctg gtggagtctg gaccacctca cgagtgaa	2040
atggacgtgt ttaatctggg cgatcagacc gccgtgctgc tgaagtccct ggcaggagt	2100
cctctggcat ctgtggaggg ccagaagtac cacctgaagt ctggacacgt gacctgcgac	2160
gtgggactgg agaagctgaa gctgaagggc accacatata gcatgtcgca taaggccaag	2220
tttaagtgga agagggtgcc agtggactcc ggacacgata ccgtggtcat ggaggtgtct	2280

-continued

tacacaggca ggcacaagcc atgtagaatc cccgtgcggg cagtggcaca cggagtgcc	2340
gccgtgaacg tggccatgct gatcaccct aaccaacca tcgagacaaa tggcggcggc	2400
ttcatcgaga tgcagctgcc acccggcgac aatatcatct atgtgggcga tctgagccag	2460
cagtgggtcc agaagggtc caccatcgcc cggatgttg agaagacacg gagaggcctg	2520
gagagactgt ccgtgggtgg agagcacgca tgggattttg gctctgtggg aggcgtgctg	2580
tctagcgtgg gcaaggcaat ccacaccgtg ctgggaggcg ccttcaatac actgtttggc	2640
ggcgtgggct tcatacccaa gatgctgctg ggagtggccc tgggtgggct gggcctgaac	2700
gcccggatc ctaccatgag catgacattt ctggccgtgg gcgcctgac cctgatgatg	2760
acaatgggcg tgggcgccta atgactcgag tctagaggcg ccgtttaaac ccgctgatca	2820
gcctcgactg tgccttctag ttgccagcca tctgttgttt gcccctccc cgtgccttcc	2880
ttgacctgg aaggtgccac tcccactgtc ctttcctaataaaatgagga aattgcatcg	2940
cattgtctga gtaggtgtca ttctattctg gggggtgggg tggggcagga cagcaagggg	3000
gaggattggg aagacaatag caggcatgct ggggatgcgg tgggctctat ggttctact	3060
gggcggtttt atggacagca agcgaaccgg aattgccagc tggggcgccc tctggttaagg	3120
ttgggaagcc ctgcaaagta aactggatgg ctttcttgcc gccaggatc tgatggcgca	3180
ggggatcaag ctctgatcaa gagacaggat gaggatcgtt tcgcatgatt gaacaagatg	3240
gattgcacgc aggttctccg gccgcttggg tggagaggct attcggctat gactgggcac	3300
aacagacaat cggctgtctc gatgccggcg tgttcggct gtcagcgag gggcgcccg	3360
ttctttttgt caagaccgac ctgtccggtg cctgaatga actgcaagac gaggcagcgc	3420
ggctatcgtg gctggccacg acgggcgttc cttgcgcagc tgtgctcgac gttgtcactg	3480
aagcgggaag ggaactgctg ctattggcg aagtgccggg gcaggatctc ctgtcatctc	3540
acctgtctcc tgcgagaaa gtatccatca tggctgatgc aatgcggcgg ctgcatacgc	3600
ttgatccgcg tacctgccc ttcgaccacc aagcgaacaa tcgcatcgag cgagcacgta	3660
ctcgatgga agccggtctt gtcgatcagg atgatctgga cgaagagcat caggggctcg	3720
cgcacgcga actgttcgcc aggtcaagg cgagcatgcc cgacggcgag gatctcgtcg	3780
tgacctatgg cgatgcctgc ttgccgaata tcatggtgga aaatggccgc ttttctggat	3840
tcacgactg tggccgctg ggtgtggcgg accgctatca ggacatagcg ttggctaccc	3900
gtgatattgc tgaagagctt ggcggcgaat gggctgaccg cttcctcgtg ctttacggta	3960
tcgcccgtcc cgattcgag cgcacgcct tctatgcct tcttgacgag ttcttctgaa	4020
ttattaacgc ttacaatttc ctgatcgggt attttctcct tacgcatctg tgcggatttt	4080
cacaccgcat caggtggcac ttttcgggga aatgtgcgcg gaacctctat ttgtttattt	4140
ttctaatac attcaaatat gtatccgctc atgagacaat aacctgata aatgcttcaa	4200
taatagcacg tgctaaaact tcatttttaa tttaaaagga tctaggtgaa gatccttttt	4260
gataatctca tgacaaaaat ccctaacgt gagttttcgt tccactgagc gtcagacccc	4320
gtagaaaaga tcaaggatc ttcttgagat ccttttttcc tgcgcgtaat ctgctgcttg	4380
caaacaaaaa aaccaccgct accagcgtg gtttgttgc cggatcaaga gctaccaact	4440
ctttttccga aggttaactg cttcagcaga gcgcagatac caaatactgt tcttctagt	4500
tagccgtagt taggccacca cttcaagaac tctgtagcac cgcctacata cctcgctctg	4560

-continued

```

ctaatacctgt taccagtggc tgctgccagt ggcgataagt cgtgtcttac cgggttggac 4620
tcaagacgat agttaccgga taaggcgcag cggtcgggct gaacggggggg ttcgtgcaca 4680
cagcccagct tggagcgaaac gacctacacc gaactgagat acctacagcg tgagctatga 4740
gaaagcgcca cgcttccga agggagaaaag gcggacaggt atccggtaaag cggcagggtc 4800
ggaacaggag agcgcacgag ggagcttcca ggggaaacg cctggatatct ttatagtcct 4860
gtcgggttct gccacctctg acttgagcgt cgatttttgt gatgctcgtc aggggggcg 4920
agcctatgga aaaacgccag caacgcggcc tttttacggt tcctggcctt ttgctggcct 4980
tttctcaca tgttctt 4997

```

```

<210> SEQ ID NO 2
<211> LENGTH: 5060
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pVAX1-IgE-prM signal-prME

```

```

<400> SEQUENCE: 2

```

```

gctgcttcgc gatgtacggg ccagatatac gcgttgacat tgattattga ctagttatta 60
atagtaatac attacggggt cattagttca tagcccatat atggagttcc gcgttacata 120
acttacggta aatggccgcg ctgggtgacc gcccaacgac ccccgcccat tgacgtcaat 180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga 240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtacgcc 300
ccctattgac gtcaatgacg gtaaatggcc cgctggcat tatgccagat acatgacctt 360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgtatta ccatggtgat 420
gcggttttgg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag 480
tctccacccc attgacgtca atgggagttt gttttggcac caaaatcaac gggactttcc 540
aaaatgtcgt aacaactccg ccccatgac gcaaatgggc ggtaggcgtg tacggtgga 600
ggtctatata agcagagctc tctggctaac tagagaaccc actgcttact ggcttatcga 660
aattaatacg actcactata gggagaccca agctggctag cgtttaaact taagcttgg 720
accgagctcg gatccgccac catggactgg acctggattc tgttctggtg gcagcagca 780
acacgggtgc acagcatgag cggcgtggac tggacctgga tcttctgac aatggccctg 840
atgaccatgg ccatggccac cacaatccac agggaccgcg agggctacat ggtcatgagg 900
gcaagcggaa gagatgcagc atcccagggt cgggtgcaga acggaacatg cgtgatcctg 960
gccaccgaca tgggcgagtg gtgcgaggat tccatcacat actcttgcgt gaccatcgac 1020
caggaggagg agcctgtgga cgtggattgc tttttagagg gcgtggatcg cgtgaagctg 1080
gagtatggcc ggtgtggcag acaggcaggc agccggggca agcggagcgt ggtcatccca 1140
acccacgccc agaaggacat ggtgggaaga ggacacgcat ggctgaaggg cgacaacatc 1200
agggatcacg tgacacgcgt ggagggtggt atgtggaaga ataagctgct gaccgcagca 1260
atcgtggccc tggcctggct gatggtggac tcttggatgg ccagggtgac cgtgatcctg 1320
ctggccctga gcctgggacc cgtgtacgca accagggtgca cacacctgga gaaccggat 1380
ttcgtgaccg gaacacaggg aaccacacgc gtgtccctgg tgctggagct gggaggatgc 1440
gtgaccatca cagcagaggg caagccaagc atcgacgtgt ggctggagga tatctttcag 1500

```

-continued

gagtccccg	cggagacacg	ggagtattgc	ctgcacgcca	agctgaccaa	tacaaagggtg	1560
gaggccagat	gtccaaccac	aggacctgcc	accctgccag	aggagcacca	ggccaacatg	1620
gtgtgcaagc	gggaccagtc	cgatagaggc	tggggcaatc	actgtggctt	ctttggcaag	1680
ggctctatcg	tggcctgcgc	caagttcgag	tgtgaggagg	ccaagaaggc	cgtggggccac	1740
gtgtacgact	ctaccaagat	cacatatgtg	gtgaagggtg	agccccacac	aggcgattac	1800
ctggccgcca	acgagaccaa	cagcaatagg	aagtccgccc	agtttacgtg	ggccagcgag	1860
aaagtgatcc	tgcgcctggg	cgactatggc	gacgtgagcc	tgacatgcaa	ggtggcaagc	1920
ggaatcgacg	tggcacagac	cgtggtcatg	tctctgggca	gctccaagga	tcacctgcct	1980
agcgcttgcc	agggtcacag	ggactgggtc	gaggatctgg	ccctgccatg	gaagcacaag	2040
gacaaccagg	attggaattc	cgtggagaag	ctgggtgagt	tcggaccacc	tcacgcagtg	2100
aagatggacg	tgtttaatct	ggcgcatcag	acagccgtgc	tgctgaagtc	cctggcagga	2160
gtgcctctgg	catctgtgga	gggccagaag	taccacctga	agtcctggaca	cgtgacctgc	2220
gacgtgggac	tggagaagct	gaagctgaag	ggcaccacat	atagcatgtg	cgataaggcc	2280
aagttaagt	ggaagagggt	gccagtggac	tccggacacg	atacagtggg	catggagggtg	2340
tcttacaccg	gcagcgacaa	gccatgtaga	atccccgtgc	gggcagtggc	acacggagtg	2400
cccgccgtga	acgtggccat	gctgatcacc	cctaacccaa	ccatcgagac	aaatggcggc	2460
ggcttcacgc	agatgcagct	gccaccgggc	gacaatatca	tctatgtggg	cgatctgagc	2520
cagcagtggt	tccagaaggg	ctccacaatc	ggccggatgt	ttgagaagac	ccggagaggc	2580
ctggagagac	tgtcogtggt	gggagagcac	gcattgggatt	ttggctctgt	gggaggcgtg	2640
ctgtctagcg	tgggcaaggc	catccacaca	gtgctgggcg	gcgccttcaa	tacctgttt	2700
ggcggcgtgg	gcttcacccc	caagatgctg	ctgggagtg	ccctgggtgtg	gctgggcctg	2760
aacgcaagga	atcctaccat	gagcatgaca	tttctggccg	tgggcgcctt	gacctgatg	2820
atgacaatgg	gcgtgggcgc	ctaagtactc	gagtcctagag	ggcccgttta	aaccgcgtga	2880
tcagcctcga	ctgtgccttc	tagttgccag	ccatctgttg	tttgcccttc	ccccgtgcct	2940
tccttgaccc	tgggaagggtg	cactcccact	gtcctttcct	aataaaatga	ggaaattgca	3000
tcgcattgtc	tgagtaggtg	tcattctatt	ctgggggggtg	gggtggggca	ggacagcaag	3060
ggggaggatt	gggaagacaa	tagcaggcat	gctggggatg	cggtgggctc	tatggcttct	3120
actgggcggg	tttatggaca	gcaagcgaac	cgggaattgcc	agctggggcg	ccctctggta	3180
agggtgggaa	gccctgcaaa	gtaaactgga	tggctttctt	gccgccaagg	atctgatggc	3240
gcaggggatc	aagctctgat	caagagacag	gatgaggatc	gtttcgcatg	attgaacaag	3300
atggattgca	cgcaggttct	ccggccgctt	gggtggagag	gctattcggc	tatgactggg	3360
cacaacagac	aatcggtgc	tctgatgccg	ccgtgttccg	gctgtcagcg	caggggcgcc	3420
cggttctttt	tgtcaagacc	gacctgtccg	gtgccctgaa	tgaactgcaa	gacgaggcag	3480
cgcggctatc	gtggctggcc	acgacgggcg	ttccttgccg	agctgtgctc	gacgttgtca	3540
ctgaagcggg	aagggactgg	ctgctattgg	gcgaagtgcc	ggggcaggat	ctcctgtcat	3600
ctcaccttgc	tcctgccgag	aaagtatcca	tcattggctga	tgcaatgcgg	cggctgcata	3660
cgccttgatcc	ggctacctgc	ccattcgacc	accaagcgaa	acatcgcatc	gagcgagcac	3720
gtactcggat	ggaagccggg	cttgtcgatc	aggatgatct	ggacgaagag	catcaggggc	3780

-continued

```

tcgcgccagc cgaactgttc gccaggctca aggcgagcat gcccgacggc gaggatctcg 3840
tcgtgaccca tggcgatgcc tgcttgccga atatcatggg ggaaaatggc cgcttttctg 3900
gattcatcga ctgtggcccg ctgggtgtgg cggaccgcta tcaggacata gcgttggtta 3960
cccgatgat tctgaagag ctggcgggc aatgggtga ccgcttcctc gtgctttacg 4020
gtatcgccgc tcccatttcg cagcgcatcg ccttctatcg ccttcttgac gagttcttct 4080
gaattattaa cgcttacaat ttctgatgc ggtattttct ccttacgcac ctgtgcggta 4140
tttcacaccg catcagggtg cacttttcgg ggaaatgtgc gcggaacccc tatttggtta 4200
tttttctaaa tacattcaaa tatgatccg ctcatgagac aataaccctg ataaatgctt 4260
caataatagc acgtgctaaa acttcatttt taatttaaaa ggatctaggt gaagatcctt 4320
tttgataatc tcatgaccaa aatcccttaa cgtgagtttt cgttccactg agcgtcagac 4380
cccgtagaaa agatcaaagg atcttcttga gatccttttt ttctgcgcgt aatctgctgc 4440
ttgcaacaaa aaaaaccacc gctaccagcg gtggtttgtt tgccggatca agagctacca 4500
actcttttct cgaaggtaac tggcttcagc agagcgagc taccaaatac tgttcttcta 4560
gtgtagccgt agttaggcca ccacttcaag aactctgtag caccgcctac atacctcgt 4620
ctgctaatac tgttaccagt ggtgctgcc agtggcgata agtcgtgtc ttaccgggtg 4680
gactcaagac gatagttacc ggataaggcg cagcggctcg gctgaacggg gggttcgtgc 4740
acacagccca gcttgagcgc aacgacctac accgaactga gatacctaca gcgtgagcta 4800
tgagaaagcg ccacgcttcc cgaagggaga aaggcggaca ggtatccggt aagcggcagg 4860
gtcgaacacg gagagcgcac gagggagctt ccagggggaa acgcctggta tctttatagt 4920
cctgtcgggt ttcgccacct ctgacttgag cgtcgatttt tgtgatgtc gtcagggggg 4980
cggagcctat ggaaaaacgc cagcaacgcg gcctttttac ggttcctggc cttttgctgg 5040
ccttttgctc acatgtttct 5060

```

```

<210> SEQ ID NO 3
<211> LENGTH: 5033
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pVAX1-IgE-portion of prM signal-prME
<400> SEQUENCE: 3

```

```

gctgcttcgc gatgtacggg ccagatatca gcgttgacat tgattattga ctagttatta 60
atagtaatca attacggggg cattagttca tagcccatat atggagtacc gcgttacata 120
acttacggta aatggcccg ctagctgacc gcccaacgac ccccgcccat tgacgtcaat 180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga 240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtaacgc 300
ccctattgac gtcaatgacg gtaaatggcc gccttggcat tatgccagc acatgacctt 360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgctatta ccatggtgat 420
gcgggttttg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag 480
tctccacccc attgacgtca atgggagttt gttttggcac caaaatcaac gggactttcc 540
aaaatgtcgt aacaactccg cccattgac gcaaatgggc ggtaggcgtg tacggtggga 600
ggtctatata agcagagctc tctggctaac tagagaaccc actgcttact ggcttatcga 660

```

-continued

aattaatacgc actcactata gggagaccca agctggctag cgtttaaaact taagcttggc	720
accgagctcg gatccgccac catggactgg acctggatc tgttctctgg ggcagcagca	780
acacgggtgc acagctttct gaccatggcc ctgatgacaa tggccatggc caccacaatc	840
cacagggacc gcgagggeta catggctcatg agggcaagcg gaagagatgc agcatcccag	900
gtgagagtgc agaacggcac ctgcgtgatc ctggccacag acatgggcga gtggtgcgag	960
gattccatca cctactcttg cgtgacaatc gaccaggagg aggagcctgt ggacgtggat	1020
tgttctctgta gggcgctgga tgcgtgaag ctggagtatg gccggtgtgg cagacaggca	1080
ggcagccggg gcaagcggag cgtggtcatc ccaaccacg cacagaagga catggtggga	1140
aggggacacg catggctgaa gggcgacaac atcagggatc acgtgacccg cgtggagggc	1200
tggatgtgga agaataagct gctgacagca gcaatcgtgg ccctggcatg gctgatggtg	1260
gactcttgga tggccagggt gaccgtgatc ctgctggccc tgagcctggg acccgtgtac	1320
gcaaccaggt gcacacacct ggagaaccgc gatttctgta ccggaacaca gggaaaccaca	1380
cgcgtgtccc tgggtctgga gctgggagga tgcgtgacca tcacagcaga gggcaagcca	1440
agcatcgacg tgtggctgga ggatatcttt caggagtccc cagcagagac ccgggagtat	1500
tgccctgcacg ccaagctgac caatacaaag gtggaggcca gatgtccac cacaggccct	1560
gccacactgc cagaggagca ccaggccaac atgggtgtgca agcgggacca gtccgataga	1620
ggctggggca atcactgtgg cttctttggc aagggtctta tcgtggcctg cgccaagtgc	1680
gagtgtgagg agggcaagaa ggcctgggc cactgttacg actctacaa gatcacatat	1740
gtggtgaagg tggagccaca caccggcgat tacctggcag ccaacgagac aaacagcaat	1800
aggaagtccg ccagctttac cgtggccagc gagaaagtga tcctgcgcct gggcgactat	1860
ggcgacgtga gcctgacctg caagggtgga agcggaatcg acgtggcaca gacagtggtc	1920
atgtctctgg gcagctccaa ggatcacctg cctagcgctt ggcagggtgca cagggactgg	1980
ttcgaggatc tggccctgcc atggaagcac aaggacaacc aggatggaa ttccgtggag	2040
aagctgggtg agttcggacc acctcacgca gtgaagatgg acgtgtttaa tctgggcat	2100
cagaccgccc tgctgtgaa gtccctggca ggagtgcctc tggcatctgt ggagggccag	2160
aagtaccacc tgaagtctgg acacgtgacc tgcgacgtgg gactggagaa gctgaagctg	2220
aagggcacca catatagcat gtgcgataag gccaaagtta agtgggaagag ggtgccagt	2280
gactccggac acgataccgt ggtcatggag gtgtcttaca caggcagcga caagccatgt	2340
agaatccccg tgcgggcagt ggcacacgga gtgcccgcg tgaacgtggc catgctgatc	2400
acccctaacc caaccatcga gacaaatggc ggcggcttca tcgagatgca gctgccaccc	2460
ggcgacaata tcctctatgt gggcgatctg agccagcagt ggttcagaa gggctccacc	2520
atcgcccgga tgtttgagaa gacacggaga ggcctggaga gactgtccgt ggtgggagag	2580
cacgcatggg attttgctc tgtgggaggc gtgctgtcta gcgtgggcaa ggcaatccac	2640
accgtgctgg gagggcctt caatacactg tttggcgcg tgggcttcat cccaagatg	2700
ctgctgggag tggccctggg gtggctgggc ctgaacgccc ggaatcctac catgagcatg	2760
acatttctgg ccgtgggcgc cctgacctg atgatgacaa tggcgctggg cgctaataga	2820
ctcgagtcta gagggccctg ttaaacccgc tgatcagcct cgactgtgcc ttctagtgtc	2880
cagccatctg ttgtttgccc ctccccctg ccttccttga ccctggaagg tgccactccc	2940

-continued

```

actgtccttt cctaataaaa tgaggaaatt gcatcgatt gtctgagtag gtgtcattct 3000
attctggggg gtggggtggg gcaggacagc aagggggagg attgggaaga caatagcagg 3060
catgctgggg atgcgggtggg ctctatggct tctactgggc gggttttatgg acagcaagcg 3120
aaccggaatt gccagctggg gcgccctctg gtaagggtgg gaagccctgc aaagtaaact 3180
ggatggcttt cttgccgcca aggatctgat ggccgagggg atcaagctct gatcaagaga 3240
caggatgagg atcgtttcgc atgattgaac aagatggatt gcacgcagggt tctccggccg 3300
cttgggtgga gaggtatctt ggctatgact gggcacaaca gacaatcggc tgctctgatg 3360
ccgccgtgtt ccggtgtgca gcgcaggggc gcccggttct ttttgtcaag accgacctgt 3420
ccggtgccct gaatagaact caagacgagg cagcgcggct atcgtggctg gccacgacgg 3480
gcgttccttg cgcagctgtg ctgcagcttg tcaactgaagc gggaaggagc tggctgctat 3540
tgggcgaagt gccggggcag gatctcctgt catctcacct tgctcctgcc gagaaagtat 3600
ccatcatggc tgatgcaatg cggcggctgc atacgcttga tccggctacc tgccattcgc 3660
accaccaagc gaaacatgc atcgagcgag cagctactcg gatggaagc ggtcttgtcg 3720
atcaggatga tctggacgaa gagcatcagg ggctcgcgcc agccgaactg ttcgccaggc 3780
tcaaggcgag catgccgac ggcgaggatc tcgtcgtgac ccatggcgat gcctgcttgc 3840
cgaatatcat ggtggaaaat ggcgcctttt ctggattcat cgactgtggc cggctgggtg 3900
tggcggaccg ctatcaggac atagcgttgg ctaccgtga tattgctgaa gagcttggcg 3960
gcgaatgggc tgaccgcttc ctgctgcttt acggtatcgc cgctcccgat tcgcagcgca 4020
tcgccttcta tcgccttctt gacgagttct tctgaattat taacgcttac aatttctgta 4080
tgcggtattt tctccttacg catctgtgcg gtatttcaca ccgcatcagg tggcactttt 4140
cggggaaaatg tgccgggaac cctatattgt ttatttttct aaatacatc aaatatgtat 4200
ccgctcatga gacaataacc ctgataaatg cttcaataat agcacgtgct aaaaattcat 4260
ttttaattta aaaggatcta ggtgaagatc ctttttgata atctcatgac caaaatccct 4320
taacgtgagt tttcgttcca ctgagcgtca gaccccgtag aaaagatcaa aggatcttct 4380
tgagatcctt tttttctgcg cgtaactctgc tgcttgcaaa caaaaaaac accgctacca 4440
gcggtgggtt gtttgccgga tcaagagcta ccaactcttt ttccgaagggt aactggcttc 4500
agcagagcgc agataccaaa tactgttctt ctagtgtagc cgtagttagg ccaccacttc 4560
aagaactctg tagcaccgcc tacatacttc gctctgctaa tcctgttacc agtggtgct 4620
gccagtggcg ataagtcgtg tcttaccggg ttggaactca gacgatagtt accggataag 4680
gcgcagcggc cgggctgaac ggggggttcg tgcacacagc ccagcttgga gcgaacgacc 4740
tacaccgaac tgagatacct acagcgtgag ctatgagaaa gcgccacgct tcccgaagg 4800
agaaaggcgg acaggtatcc ggtaagcggc agggtcggaa caggagagcg cacgaggag 4860
cttccagggg gaaacgcctg gtatctttat agtcctgtcg gggttcgcca cctctgactt 4920
gagcgtcgat ttttgtgatg ctgcgcaggg gggcggagcc tatggaaaaa cgccagcaac 4980
gcggcctttt tacggttcct ggccttttgc tggccttttg ctcacatggt ctt 5033

```

<210> SEQ ID NO 4

<211> LENGTH: 5006

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: pVAX1-prM signal-prME

<400> SEQUENCE: 4

gctgcttcgc gatgtacggg ccagatatac gcgttgacat tgattattga ctagttatta	60
atagtaatca attacggggg cattagttca tagcccatat atggagttcc gcgttacata	120
acttacggta aatggcccg ctaggtgacc gcccaacgac ccccgcccat tgacgtcaat	180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga	240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtacgcc	300
ccctattgac gtcaatgacg gtaaatggcc cgctggcat tatgccagt acatgacctt	360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgctatta ccatggtgat	420
gcggttttgg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag	480
tctccacccc attgacgtca atgggagttt gttttggcac caaaatcaac gggactttcc	540
aaaatgtcgt aacaactccg ccccatgac gcaaatgggc ggtaggcgtg tacggtggga	600
ggtctatata agcagagctc tctggctaac tagagaacct actgcttact ggcttatcga	660
aattaatacg actcactata gggagacca agctggctag cgtttaaact taagcttggg	720
accgagctcg gatccgccac catgagcggc gtggactgga cctggatctt cctgacaatg	780
gcccgtgatg ccatggccat gggccaccaca atccacaggg accgcgaggg ctacatggtc	840
atgagggcaa gcggaagaga tgcagcatcc caggtgcggg tgcagaacgg aacatgcgtg	900
atcctggcca ccgacatggg cgagtgggtg gaggattcca tcacatactc ttgcgtgacc	960
atcgaccagg agggaggagc tgtggacgtg gattgctttt gtaggggcgt ggatcgctg	1020
aagctggagt atggccgggtg tggcagacag gcaggcagcc ggggcaagcg gagcgtggtc	1080
atcccaaccc acgcccagaa ggacatgggt ggaagaggac acgcatggct gaaggcgcgac	1140
aacatcaggg atcacgtgac acgcgtggag ggctggatgt ggaagaataa gctgctgacc	1200
gcagcaatcg tggccctggc ctggctgatg gtggactctt ggatggccag ggtgaccgtg	1260
atcctgctgg ccctgagcct gggaccctg tacgcaacca ggtgcacaca cctggagaac	1320
cgcgatttcc tgaccggaac acaggggaac acacgcgtg ccctgggtgct ggagctggga	1380
ggatgcgtga ccatcacagc agagggcaag ccaagcatcg acgtgtggct ggaggatatc	1440
tttcaggagt ccccgccga gacacgggag tattgcctgc acgccaagct gaccaataca	1500
aaggtggagg ccagatgtcc aaccacagga cctgccaccc tgccagagga gcaccaggcc	1560
aacatgggtg gcaagcggga ccagtcgat agaggtctgg gcaatcactg tggtttcttt	1620
ggcaagggct ctatcgtggc ctgcgccaaag ttcgagtgtg aggaggccaa gaaggccgtg	1680
ggccacgtgt acgactctac caagatcaca tatgtggtga aggtggagcc ccacacaggc	1740
gattacctgg ccgccaacga gaccaacagc aataggaagt ccgcccagtt taccgtggcc	1800
agcgagaaa tgatcctgcg cctgggcgac tatggcgacg tgagcctgac atgcaagggtg	1860
gcaagcggaa tcgacgtggc acagacgtg gtcattgtctc tgggcagctc caaggatcac	1920
ctgcctagcg cctggcaggt gcacagggac tggttcgagg atctggccct gccatggaag	1980
cacaaggaca accaggattg gaattccgtg gagaagctgg tggagttcgg accacctcac	2040
gcagtgaaga tggacgtgtt taatctgggc gatcagacag ccgtgctgct gaagtccctg	2100
gcaggagtgc ctctggcatc tgtggagggc cagaagtacc acctgaagtc tggacacgtg	2160

-continued

acctgcgacg	tgggactgga	gaagctgaag	ctgaaggcca	ccacatatag	catgtgcgat	2220
aaggccaagt	ttaagtggaa	gaggggtcca	gtggactccg	gacacgatac	agtggtcatg	2280
gaggtgtctt	acaccgcgag	cgacaagcca	tgtagaatcc	ccgtgcgggc	agtggcacac	2340
ggagtgcctg	ccgtgaacgt	ggccatgctg	atcaccccta	acccaacct	cgagacaaat	2400
ggcggcggct	tcctcgagat	gcagctgcca	cccgcgaca	atatcatcta	tgtggcgcat	2460
ctgagccagc	agtggttcca	gaagggtccc	acaatcgccc	ggatgtttga	gaagaccggg	2520
agaggcctgg	agagactgtc	cgtggtggga	gagcacgcat	gggattttgg	ctctgtggga	2580
ggcgtgctgt	ctagcgtggg	caaggccatc	cacacagtgc	tgggcggcgc	cttcaatacc	2640
ctgtttggcg	gcgtgggctt	catccccaag	atgctgctgg	gagtggccct	ggtgtggctg	2700
ggcctgaacg	caaggaatcc	taccatgagc	atgacatttc	tggccgtggg	cgccctgacc	2760
ctgatgatga	caatgggcgt	gggcgcctaa	tgactcgagt	ctagagggcc	cgtttaaac	2820
cgctgatcag	cctcgactgt	gccttctagt	tgccagccat	ctgtgtttg	ccctccccc	2880
gtgccttcct	tgacctgga	aggtgccact	cccactgtcc	tttcctaata	aaatgaggaa	2940
attgcatcgc	attgtctgag	taggtgtcat	tctattctgg	ggggtgggg	ggggcaggac	3000
agcaaggggg	aggattggga	agacaatagc	aggcatgctg	gggatgcggt	gggctctatg	3060
gcttctactg	ggcggtttta	tggacagcaa	gcgaaccgga	attgccagct	ggggcgccct	3120
ctggtaaagt	tgggaagccc	tgcaaagtaa	actggatggc	tttcttgccg	ccaaggatct	3180
gatggcgcag	gggatcaagc	tctgatcaag	agacaggatg	aggatcgttt	cgcatgattg	3240
aacaagatgg	attgcacgca	ggttctccgg	ccgcttgggt	ggagaggcta	ttcggtatg	3300
actgggcaca	acagacaatc	ggctgctctg	atgccgccgt	gttcggctg	tcagcgcagg	3360
ggcgcgccgt	tctttttgtc	aagaccgacc	tgccgggtgc	cctgaatgaa	ctgcaagacg	3420
aggcagcgcg	gctatcgtgg	ctggccacga	cgggcgttcc	ttgcgcagct	gtgctcgacg	3480
ttgtcactga	agcgggaagg	gactggctgc	tattgggcga	agtgcggggg	caggatctcc	3540
tgatcatctc	ccttgctcct	gccagaaaag	tatccatcat	ggctgatgca	atgcggcggc	3600
tgcatacgct	tgatccggct	acctgccc	tcgaccacca	agcgaaacat	cgcatcgagc	3660
gagcacgtac	tcggatggaa	gccggtcttg	tcgatcagga	tgatctggac	gaagagcatc	3720
aggggctcgc	gccagccgaa	ctgttcgcca	ggctcaaggc	gagcatgccc	gacggcgagg	3780
atctcgtcgt	gacccatggc	gatgcctgct	tgccgaatat	catggtggaa	aatggccgct	3840
tttctggatt	catcgactgt	ggcggctggg	gtgtggcgga	ccgctatcag	gacatagcgt	3900
tggctacccg	tgatattgct	gaagagcttg	gcggcgaatg	ggctgaccgc	ttcctcgtgc	3960
tttacgggat	cgccgtctcc	gatttcgcagc	gcacgcctt	ctatgcctt	cttgacgagt	4020
tcttctgaat	tattaacgct	tacaatttcc	tgatgcggta	ttttctcctt	acgcactctgt	4080
gcggtatttc	acaccgcatc	aggtggcact	tttcggggaa	atgtgcgcgg	aaccctatt	4140
tgtttatttt	tctaaataca	ttcaaata	tatccgctca	tgagacaata	accctgataa	4200
atgcttcaat	aatagcacgt	gctaaaactt	cattttta	ttaaaaggat	ctagggtgaag	4260
atcctttttg	ataatctcat	gacccaaa	ccttaacgtg	agttttcgtt	ccactgagcg	4320
tcagaccccg	tagaaaagat	caaaggatct	tcttgagatc	ctttttttct	gcgcgtaatc	4380
tgctgcttgc	aaacaaaaaa	accaccgcta	ccagcgggtg	tttgtttgcc	ggatcaagag	4440

-continued

```

ctaccaactc tttttccgaa ggtaactggc ttcagcagag cgcagatacc aaatactgtt 4500
cttctagtggt agccgtagtt aggccaccac ttcaagaact ctgtagcacc gcctacatac 4560
ctcgctctgc taatcctgtt accagtggtt gctgccagtg gcgataagtc gtgtcttacc 4620
gggttggtgact caagacgata gttaccggat aaggcgcagc ggtcgggctg aacggggggt 4680
tcgtgcacac agcccagctt ggagcgaacg acctacaccg aactgagata cctacagcgt 4740
gagctatgag aaagcgccac gcttcccga gggagaaagg cggacaggta tccggttaagc 4800
ggcaggggtcg gaacaggaga gcgcacgagg gagcttcag ggggaaacgc ctggtatctt 4860
tatagtcctg tcgggtttcg ccacctctga cttgagcgtc gatttttgtg atgctcgtca 4920
ggggggcgga gcctatggaa aaacgcccagc aacgcggcct ttttacggtt cctggccttt 4980
tgctggcctt ttgtcacat gttctt 5006

```

```

<210> SEQ ID NO 5
<211> LENGTH: 5015
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pVAX1-JEV signal-prME

```

```

<400> SEQUENCE: 5

```

```

gctgcttcgc gatgtacggg ccagatatat gcgttgacat tgattattga ctagtattata 60
atagtaatca attacgggggt cattagtcca tagcccatat atggagttcc gcgttacata 120
acttacggta aatggcccg cctggctgacc gcccaacgac ccccgcccat tgacgtcaat 180
aatgacgtat gttcccatag taacgccaat agggactttc cattgacgtc aatgggtgga 240
gtatttacgg taaactgccc acttggcagt acatcaagtg tatcatatgc caagtaagcc 300
ccctattgac gtcaatgacg gtaaatggcc cgcctggcat tatgccagat acatgacctt 360
atgggacttt cctacttggc agtacatcta cgtattagtc atcgctatta ccatggtgat 420
gcgggttttg cagtacatca atgggcgtgg atagcggttt gactcacggg gatttccaag 480
tctccacccc attgacgtca atgggagttt gttttggcac caaaatcaac gggactttcc 540
aaaatgtcgt aacaactccg ccccatggac gcaaatgggc ggtaggcgtg tacgggtgga 600
ggtctatata agcagagctc tctggctaac tagagaaccc actgcttact ggcttatcga 660
aattaatacg actcactata gggagaccca agctggctag cgtttaaact taagcttggt 720
accgagctcg gatccgccac catgggcaag cggtcgcgcg gctctatcat gtggctggcc 780
agcctggcag tggtcacgcg atgcgcagga gcaaccacaa tccacaggga ccgcgagggc 840
tacatggcca tgagggcacg tggaagagat gcagcaagcc aggtgagagt gcagaacggc 900
acctgcgtga tcctggccac agacatgggc gagtggtgcg aggatagcat cacctactcc 960
tgcgtagcaa tcgaccagga ggaggagcct gtggacgtgg attgcttctg taggggcgtg 1020
gatcgcgtga agctggagta tggccggtgc ggcagacagg caggcagccg gggcaagcgg 1080
agcgtggcca tcccaaccca cgcacagaag gacatggtgg gaaggggaca cgcattggctg 1140
aaggcgacac acatcaggga tcacgtgacc cgcgtggagg gctggatgtg gaagaataag 1200
ctgctgacag cagcaatcgt ggccttggca tggctgatgg tggactcctg gatggccagg 1260
gtgaccgtga tcctgctggc cctgtctctg ggaccctgtg acgcaaccag gtgcacacac 1320
ctggagaacc gcgatttcgt gaccggaaca cagggaacca cacgcgtgtc cctggtgctg 1380

```

-continued

gagctgggag gatgogtgac catcacagca gagggcaagc catctatcga cgtgtggctg	1440
gaggatatct ttcaggagag cccagcagag acccgggagt attgectgca cgccaagctg	1500
accaatacaa aggtggaggc cagatgtccc accacaggcc ctgccacact gccagaggag	1560
caccaggcca acatgggtgt caagcgggac cagagcgata gaggtgggg caatcactgt	1620
ggcttctttg gcaagggctc catcgtggcc tgcgccaaat tcgagtgtga ggaggccaag	1680
aaggccgtgg gccacgtgta cgactccacc aagatcacat atgtggtgaa ggtggagcca	1740
cacaccggcg attacctggc agccaacgag acaaactcta ataggaagag cgcccagttt	1800
accgtggcct ctgagaaagt gatcctgcgc ctgggcgact atggcgacgt gagcctgacc	1860
tgcaaggtgg cctctggcat cgacgtggcc cagacagtgg tcatgtccct gggcagctcc	1920
aaggatcacc tgccttctgc ctggcaggtg cacagggact ggctcgagga tctggccctg	1980
ccatggaagc acaaggacaa ccaggattgg aatagcgtgg agaagctggt ggagttcgga	2040
ccacctcacg cagtgaagat ggacgtgttt aatctgggcg atcagaccgc cgtgctgctg	2100
aagagcctgg caggagtgcc tctggcatcc gtggagggcc agaagtacca cctgaagtcc	2160
ggacacgtga cctgcgacgt gggactggag aagctgaagc tgaagggcac cacatattct	2220
atgtgcgata aggccaaagt taagtgaag aggggtgccag tggacagcgg acacgatacc	2280
gtggtcatgg aggtgtccta cacaggctct gacaagccat gtagaatccc cgtgcgggca	2340
gtggcacacg gagtgccgc cgtgaacgtg gccatgctga tccccctaa cccaaccatc	2400
gagacaaatg cggcgcgctt catcgagatg cagctgccac ccggcgacaa tatcatctat	2460
gtggcgcatc tgtctcagca gtggttcag aagggcagca ccatcgcccg gatgtttgag	2520
aagacacgga gaggcctgga gagactgagc gtggtgggag agcacgcatg ggattttggc	2580
tccgtgggag cgtgctgtgc tagcgtgggc aaggcaatcc acaccgtgct gggaggcgcc	2640
ttcaatacac tgtttggcgg cgtgggcttc atccccaga tgetgctggg agtggccctg	2700
gtgtggctgg gcctgaacgc ccggaatcct accatgagca tgacatttct ggcctggggc	2760
gccctgaccc tgatgatgac aatgggcgtg ggcgccta atgactcgagtc tagagggccc	2820
gtttaaaccc gctgatcagc ctgcactgtg cttctagtt gccagccatc tgttgtttgc	2880
ccctcccccg tgccttcctt gacctggaa ggtgccactc cactgtcct ttctaataa	2940
aatgaggaaa ttgcatcgca ttgtctgagt aggtgtcatt ctattctggg ggggtgggtg	3000
gggcaggaca gcaaggggga ggattgggaa gacaatagca ggcattgctg ggatgcggtg	3060
ggctctatgg cttctactgg cgggttttat ggacagcaag cgaaccggaa ttgccagctg	3120
gggcgccctc tggttaagggt ggaagccct gcaaagtaaa ctggatggct ttcttgccgc	3180
caaggatctg atggcgagg ggatcaagct ctgatcaaga gacaggatga ggatcgtttc	3240
gcatgattga acaagatgga ttgcacgcag gttctccggc cgcttgggtg gagaggctat	3300
tgggtatga ctgggcacaa cagacaatcg gctgctctga tgccgctgtg ttccgctgt	3360
cagcgcaggg cgcgccggtt ctttttgtca agaccgacct gtccggtgcc ctgaatgaac	3420
tgcaagacga ggcagcgagg ctatcgtggc tggccacgac gggcggttcc tgcgcagctg	3480
tgctcgacgt tgtcactgaa gcgggaaggg actggtgct attgggcgaa gtgccggggc	3540
aggatctcct gtcattctac cttgctcctg ccgagaaagt atccatcatg gctgatgcaa	3600
tgccggcgct gcatacgtt gatccggcta cctgcccatt cgaccaccaa gcgaaacatc	3660

-continued

```

gcatcgagcg agcacgtact cggatggaag cgggtcttgt cgatcaggat gatctggacg 3720
aagagcatca ggggctcgcg ccagccgaac tggtcgccag gctcaaggcg agcatgcccg 3780
acggcgagga tctcgctgtg acccatggcg atgectgtct gccgaatc atggtggaaa 3840
atggccgctt ttctggttc atcgactgtg gccggtggg tgtggcgga cgtatcagg 3900
acatagcgtt ggctaccctg gatattgtg aagagcttg cggcgaatgg gctgaccgt 3960
tcctcgctgt ttacgggtat gccgctcccg attcgacg catcgcttc tatcgcttc 4020
ttgacgagtt cttctgaatt attaacgctt acaatttcct gatgcggtat ttctcctta 4080
cgcatctgtg cggtatattca caccgcatca ggtggcactt ttcggggaaa tgtgcgga 4140
acccctattt gtttattttt ctaaatatcat tcaaatatgt atccgctcat gagacaataa 4200
ccctgataaa tgcttcaata atagcagctg ctaaaacttc atttttaatt taaaaggatc 4260
taggtgaaga tcctttttga taatctcatg accaaaatcc cttaacgtga gttttcgttc 4320
cactgagcgt cagacccctg agaaaagatc aaaggatctt cttgagatcc ttttttctg 4380
cgcgtaatct gctgcttgca aacaaaaaaa ccaccgtac cagcgtggtt ttgtttgccg 4440
gatcaagagc taccaactct ttttcgaag gtaactggct tcagcagagc gcagatacca 4500
aatactgttc ttctagtgtg gccgtagtta gccaccact tcaagaactc tgtagaccg 4560
ctacataacc tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg 4620
tgtcttaccg gggtggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga 4680
acgggggggtt cgtgcacaca gccagcttg gagcgaacga cctacaccga actgagatac 4740
ctacagcgtg agctatgaga aagcgccacg cttccgaag ggagaaaggc ggacaggat 4800
ccggtaaagc gcagggtcgg aacaggagag cgcacgaggg agcttccagg gggaaacgcc 4860
tggtatcttt atagtctgt cgggtttcgc cactctgac ttgagcgtcg atttttgtga 4920
tgctcgtcag gggggcgag cctatggaaa aacgccagca acgcgccctt tttacggttc 4980
ctggcctttt gctggccttt tgctcacatg ttctt 5015

```

```

<210> SEQ ID NO 6
<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: IgE leader

```

```

<400> SEQUENCE: 6

```

```

atggactgga cctggattct gttcctgggt gcagcagcaa cacgggtgca cagc 54

```

```

<210> SEQ ID NO 7
<211> LENGTH: 164
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PrM consensus sequence

```

```

<400> SEQUENCE: 7

```

```

Thr Thr Ile His Arg Asp Arg Glu Gly Tyr Met Val Met Arg Ala Ser
1           5           10           15
Gly Arg Asp Ala Ala Ser Gln Val Arg Val Gln Asn Gly Thr Cys Val
20           25           30
Ile Leu Ala Thr Asp Met Gly Glu Trp Cys Glu Asp Ser Ile Thr Tyr

```

-continued

35	40	45
Ser Cys Val Thr Ile Asp	Gln Glu Glu Glu Pro	Val Asp Val Asp Cys
50	55	60
Phe Cys Arg Gly Val Asp	Arg Val Lys Leu Glu	Tyr Gly Arg Cys Gly
65	70	75 80
Arg Gln Ala Gly Ser Arg	Gly Lys Arg Ser Val	Val Ile Pro Thr His
85	90	95
Ala Gln Lys Asp Met Val	Gly Arg Gly His Ala	Trp Leu Lys Gly Asp
100	105	110
Asn Ile Arg Asp His Val	Thr Arg Val Glu Gly	Trp Met Trp Lys Asn
115	120	125
Lys Leu Leu Thr Ala Ala	Ile Val Ala Leu Ala	Trp Leu Met Val Asp
130	135	140
Ser Trp Met Ala Arg Val	Thr Val Ile Leu Leu	Ala Leu Ser Leu Gly
145	150	155 160
Pro Val Tyr Ala		

<210> SEQ ID NO 8
 <211> LENGTH: 497
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: E consensus sequence
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (430)..(430)
 <223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<400> SEQUENCE: 8

Thr Arg Cys Thr His Leu	Glu Asn Arg Asp Phe Val	Thr Gly Thr Gln
1	5 10	15
Gly Thr Thr Arg Val Ser	Leu Val Leu Glu Leu Gly	Gly Cys Val Thr
20	25	30
Ile Thr Ala Glu Gly Lys	Pro Ser Ile Asp Val Trp	Leu Glu Asp Ile
35	40	45
Phe Gln Glu Ser Pro Ala	Glu Thr Arg Glu Tyr Cys	Leu His Ala Lys
50	55	60
Leu Thr Asn Thr Lys Val	Glu Ala Arg Cys Pro Thr	Thr Gly Pro Ala
65	70	75 80
Thr Leu Pro Glu Glu His	Gln Ala Asn Met Val Cys	Lys Arg Asp Gln
85	90	95
Ser Asp Arg Gly Trp Gly	Asn His Cys Gly Phe Phe	Gly Lys Gly Ser
100	105	110
Ile Val Ala Cys Ala Lys	Phe Glu Cys Glu Glu Ala	Lys Lys Ala Val
115	120	125
Gly His Val Tyr Asp Ser	Thr Lys Ile Thr Tyr Val	Val Lys Val Glu
130	135	140
Pro His Thr Gly Asp Tyr	Leu Ala Ala Asn Glu Thr	Asn Ser Asn Arg
145	150	155 160
Lys Ser Ala Gln Phe Thr	Val Ala Ser Glu Lys Val	Ile Leu Arg Leu
165	170	175
Gly Asp Tyr Gly Asp Val	Ser Leu Thr Cys Lys Val	Ala Ser Gly Ile
180	185	190
Asp Val Ala Gln Thr Val Val Met Ser Leu Gly Ser Ser Lys Asp His		

-continued

195					200					205					
Leu	Pro	Ser	Ala	Trp	Gln	Val	His	Arg	Asp	Trp	Phe	Glu	Asp	Leu	Ala
210						215					220				
Leu	Pro	Trp	Lys	His	Lys	Asp	Asn	Gln	Asp	Trp	Asn	Ser	Val	Glu	Lys
225					230					235				240	
Leu	Val	Glu	Phe	Gly	Pro	Pro	His	Ala	Val	Lys	Met	Asp	Val	Phe	Asn
				245					250					255	
Leu	Gly	Asp	Gln	Thr	Ala	Val	Leu	Leu	Lys	Ser	Leu	Ala	Gly	Val	Pro
			260					265					270		
Leu	Ala	Ser	Val	Glu	Gly	Gln	Lys	Tyr	His	Leu	Lys	Ser	Gly	His	Val
		275					280					285			
Thr	Cys	Asp	Val	Gly	Leu	Glu	Lys	Leu	Lys	Leu	Lys	Gly	Thr	Thr	Tyr
290						295					300				
Ser	Met	Cys	Asp	Lys	Ala	Lys	Phe	Lys	Trp	Lys	Arg	Val	Pro	Val	Asp
305					310					315					320
Ser	Gly	His	Asp	Thr	Val	Val	Met	Glu	Val	Ser	Tyr	Thr	Gly	Ser	Asp
				325					330					335	
Lys	Pro	Cys	Arg	Ile	Pro	Val	Arg	Ala	Val	Ala	His	Gly	Val	Pro	Ala
			340					345					350		
Val	Asn	Val	Ala	Met	Leu	Ile	Thr	Pro	Asn	Pro	Thr	Ile	Glu	Thr	Asn
		355					360					365			
Gly	Gly	Gly	Phe	Ile	Glu	Met	Gln	Leu	Pro	Pro	Gly	Asp	Asn	Ile	Ile
370						375					380				
Tyr	Val	Gly	Asp	Leu	Ser	Gln	Gln	Trp	Phe	Gln	Lys	Gly	Ser	Thr	Ile
385					390					395					400
Gly	Arg	Met	Phe	Glu	Lys	Thr	Arg	Arg	Gly	Leu	Glu	Arg	Leu	Ser	Val
				405					410					415	
Val	Gly	Glu	His	Ala	Trp	Asp	Phe	Gly	Ser	Val	Gly	Gly	Xaa	Leu	Ser
			420					425					430		
Ser	Val	Gly	Lys	Ala	Ile	His	Thr	Val	Leu	Gly	Gly	Ala	Phe	Asn	Thr
		435					440					445			
Leu	Phe	Gly	Gly	Val	Gly	Phe	Ile	Pro	Lys	Met	Leu	Leu	Gly	Val	Ala
450						455					460				
Leu	Val	Trp	Leu	Gly	Leu	Asn	Ala	Arg	Asn	Pro	Thr	Met	Ser	Met	Thr
465					470					475					480
Phe	Leu	Ala	Val	Gly	Ala	Leu	Thr	Leu	Met	Met	Thr	Met	Gly	Val	Gly
			485						490					495	

Ala

<210> SEQ ID NO 9

<211> LENGTH: 1989

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: prME consensus

<400> SEQUENCE: 9

accacaatcc acagggaccg cgagggctac atggctcatga gggcaagcgg aagagatgca 60

gcattcccagg tgcgcgtgca gaacggaacc tgcgtgatcc tggccacaga catgggagag 120

tgggtgcgagg attccatcac ctactcttgc gtagacaatcg accaggagga ggagcctgtg 180

gacgtggatt gctttttagt gggcgtggat cgcgtgaagc tggagtatgg ccggtgtggc 240

-continued

agacaggcag	gcagccgggg	caagcggagc	gtggtcatcc	caaccacgc	acagaaggac	300
atggtgggaa	ggggacacgc	atggctgaag	ggcgacaaca	tcagggatca	cgtgaccgc	360
gtggagggct	ggatgtggaa	gaataagctg	ctgacagcag	caatcgtggc	cctggcatgg	420
ctgatggtgg	actcttgat	ggcccggtg	accgtgatcc	tgctggccct	gagcctggga	480
cccggtacg	caaccaggtg	cacacacctg	gagaaccgcg	atttcgtgac	cggcacacag	540
ggcaccacaa	gagtgtccct	ggtgctggag	ctgggaggat	gcgtgaccat	cacagcagag	600
ggcaagccaa	gcacgcagct	gtggctggag	gatatcttcc	aggagtcccc	agcagagacc	660
cgggagtatt	gcctgcacgc	caagctgacc	aatacaaaag	tggaggccag	atgtcccacc	720
acaggccctg	ccacactgcc	agaggagcac	caggccaaca	tggtgtgcaa	gcgggaccag	780
tccgatagag	gctggggcaa	tcaactgtgc	ttctttggca	agggctctat	cgtggcctgc	840
gccaaagtcc	agtgtgagga	ggccaagaag	gccgtgggcc	acgtgtacga	ctctaccaag	900
atcacatatg	tggtgaaggt	ggagccacac	accggcgatt	acctggcagc	caacgagaca	960
aacagcaata	ggaagtccgc	ccagtttacc	gtggccagcg	agaaagtgat	cctgcgcctg	1020
ggcgactatg	gcgacgtgag	cctgacctgc	aaggtggcaa	gcggaatcga	cgtggcacag	1080
acagtgggtc	tgtctctggg	cagctccaag	gatcacctgc	ctagcgcctg	gcaggtgcac	1140
agggactggt	tcgaggatct	ggccctgcc	tggaaacaca	aggacaacca	ggattggaat	1200
tccgtggaga	agctgggtga	gttcggacca	cctcacgcag	tgaagatgga	cgtgtttaat	1260
ctgggcgatc	agaccgcctg	gctgctgaag	tccctggcag	gagtgcctct	ggcatctgtg	1320
gagggccaga	agtaccacct	gaagtctgga	cacgtgacct	gcgacgtggg	actggagaag	1380
ctgaagctga	agggcaccac	atatagcatg	tgcgataaag	ccaagttaa	gtggaagagg	1440
gtgccagtgg	actccggaca	cgataccgtg	gtcatggagg	tgtcttacac	aggcagcgac	1500
aagccatgta	gaatccccgt	gcgggcagtg	gcacacggag	tgcccgccgt	gaacgtggcc	1560
atgctgatca	cccctaacc	aaccatcgag	acaaatggcg	gcggcttcat	cgagatgcag	1620
ctgccacccg	gcgacaatat	catctatgtg	ggcgatctga	gccagcagtg	gttcagaaag	1680
ggctccacca	tcggccggat	gtttgagaag	acacggagag	gcctggagag	actgtccgtg	1740
gtgggagagc	acgcatggga	ttttggtct	gtgggaggcg	tgctgtctag	cgtgggcaag	1800
gcaatccaca	ccgtgctggg	aggcgccctc	aatacactgt	ttggcgccgt	gggcttcac	1860
cccaagatgc	tgctgggagt	ggccctggtg	tggtggggcc	tgaacgccc	gaatcctacc	1920
atgagcatga	catttctggc	cgtgggcgcc	ctgacctga	tgatgacaat	gggcgtgggc	1980
gcctaataga						1989

What is claimed is:

1. An immunological composition comprising a nucleotide sequence encoding a consensus POWV antigen, wherein the consensus POWV antigen comprises an amino acid sequence selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, an amino acid sequence that is 90% identical or greater to SEQ ID NO:2, and an amino acid sequence that is 90% identical or greater to SEQ ID NO:4.

2. The immunological composition of claim 1, wherein the nucleotide sequence comprises a nucleotide sequence selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:3, a nucleotide sequence that is 90% identical or

greater to SEQ ID NO:1, and a nucleotide sequence that is 90% identical or greater to SEQ ID NO:3.

3. The immunogenic composition of claim 1, wherein the nucleic acid molecule is selected from the group consisting of a DNA molecule and an RNA molecule.

4. The immunological composition of claim 1, wherein the nucleotide sequence comprises one or more plasmids.

5. The immunological composition of claim 1, further comprising a nucleotide sequence encoding an adjuvant.

6. The immunogenic composition of claim 1, wherein a nucleotide sequence encoding the peptide is operably linked

to at least one regulatory sequence selected from the group consisting of a start codon, an IgE leader sequence and a stop codon.

7. The immunogenic composition of claim 1, wherein the nucleic acid molecule comprises an expression vector.

8. The immunogenic composition of claim 1, wherein the nucleic acid molecule is incorporated into a viral particle.

9. The immunogenic composition of claim 1, further comprising a pharmaceutically acceptable excipient.

10. A method of treating or preventing POWV infection in a subject in need thereof, the method comprising administering the immunological composition of claim 1 to the subject.

11. The method of claim 12, wherein the administering step comprises electroporation, aerosol administration, gene-gun administration, microneedle injection, or syringe needle injection.

12. A method of inducing an immune response to POWV in a subject in need thereof, the method comprising administering an immunogenic composition of claim 1 to the subject.

13. The method of claim 12, wherein the administering step comprises electroporation.

14. A nucleic acid molecule comprising one or more nucleotide sequences selected from the group consisting of: SEQ ID NO:1, SEQ ID NO:3, a nucleotide sequence that is 90% identical or greater to SEQ ID NO:1, and a nucleotide sequence that is 90% identical or greater to SEQ ID NO:3.

15. A protein comprising one or more amino acid sequences selected from the group consisting of: SEQ ID NO:2, SEQ ID NO:4, an amino acid sequence that is 90% identical or greater to SEQ ID NO:2, and an amino acid sequence that is 90% identical or greater to SEQ ID NO:4.

* * * * *