



US 20250277006A1

(19) **United States**(12) **Patent Application Publication**
Ander et al.(10) **Pub. No.: US 2025/0277006 A1**(43) **Pub. Date: Sep. 4, 2025**(54) **FC BINDING POLYPEPTIDES****Publication Classification**(71) Applicant: **Cytiva Bioprocess R&D AB**, Uppsala (SE)(51) **Int. Cl.****C07K 14/31** (2006.01)**C07K 1/22** (2006.01)(72) Inventors: **Mats Arvid Ander**, Uppsala (SE);
Daniel Ivansson, Uppsala (SE);
Andreas Lars, **Magnus Jonsson**,
Uppsala (SE)(52) **U.S. Cl.**CPC **C07K 14/31** (2013.01); **C07K 1/22**
(2013.01); **C07K 2319/30** (2013.01)(21) Appl. No.: **18/689,737**

(57)

ABSTRACT(22) PCT Filed: **Sep. 23, 2022**(86) PCT No.: **PCT/EP2022/076493**

§ 371 (c)(1),

(2) Date: **Mar. 6, 2024**

The present disclosure relates to a class of engineered polypeptides having a binding affinity for the Fc region of immunoglobulins while exhibiting a significantly reduced binding affinity to the VH3 region of immunoglobulins. The present disclosure also relates to methods for isolating an immunoglobulin using said polypeptides as well as to related products, such as separation matrices.

(30) **Foreign Application Priority Data**

Sep. 24, 2021 (GB) 2113626.2


Specification includes a Sequence Listing.

Fig. 1A




Fig. 1B




Fig. 2




Fig. 3




Fig. 4




Fig. 5





Fig. 6






Fig. 8




Fig. 9




Fig. 10A

denotation	SEQ ID NO:	Amino acid sequence
ZGE85	1	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSGSKaILAEAKKSNDaQAAPK
ZGE86	2	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSGSKaILAEAKKINDaQAAPK
ZGE87	3	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQSLKDDPSRSKaILAEAKKVNDaQAAPK
ZGE88	4	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSRSKaILAEAKKLNDaQAAPK
ZGE89	5	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQQLKDDPSRSKaILAEAKKINDaQAAPK
ZGE90	6	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQALKDDPSRSKaILAEAKKINDaQAAPK
ZGE91	7	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQSLKDDPSGSKaILAEAKKVNDaQAAPK
ZGE92	8	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQTLKDDPSESKaILAEAKKRNDaQAAPK
ZGE93	9	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSESKaILAEAKKRNDaQAAPK
ZGE94	10	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQALKDDPSDSKaILAEAKKLNDaQAAPK
ZGE95	11	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQALKDDPSGSKaILAEAKKRNDaQAAPK
ZGE96	12	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQELKDDPSGSKaILAEAKKLNDaQAAPK
ZGE97	13	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQHLKDDPSKSKaILAEAKKLNDaQAAPK
ZGE98	14	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSDSKaILAEAKKVNDaQAAPK
ZGE99	15	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQGLKDDPSQSKaILAEAKKINDaQAAPK
ZGE100	16	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQQLKDDPSDSKaILAEAKKINDaQAAPK
	17	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQALKDDPSKSKaILAEAKKLNDaQAAPK
	18	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQALKDDPSRSKaILAEAKKLNDaQAAPK
	19	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQDLKDDPSKSKaILAEAKKINDaQAAPK
	20	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQDLKDDPSNSKaILAEAKKSNDaQAAPK
	21	VDKFDKEAQEAfYEILHLPNLTEEQRNafIQDLKDDPSRSKaILAEAKKVNDaQAAPK

Fig. 10B

Denotation	SEQ ID NO:	Amino acid sequence
	22	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQELKDDPSRSKAILAEAKKINDAQAPK
	23	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQGLKDDPSDSKAILAEAKKRND AQAPK
	24	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQGLKDDPSNSKAILAEAKKINDAQAPK
	25	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQGLKDDPSRSKAILAEAKKINDAQAPK
	26	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQGLKDDPSRSKAILAEAKKRND AQAPK
	27	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQHLKDDPSKSKAILAEAKKINDAQAPK
	28	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQHLKDDPSQSKAILAEAKKSND AQAPK
	29	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQHLKDDPSRSKAILAEAKKINDAQAPK
	30	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQHLKDDPSSSKAILAEAKKRND AQAPK
	31	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQHLKDDPSSSKAILAEAKKVND AQAPK
	32	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQKLKDDPSESKAILAEAKKINDAQAPK
	33	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQKLKDDPSKSKAILAEAKKINDAQAPK
	34	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQKLKDDPSRSKAILAEAKKINDAQAPK
	35	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQNLKDDPSHSKAILAEAKKSND AQAPK
	36	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSDSKAILAEAKKRND AQAPK
	37	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSHSKAILAEAKKRND AQAPK
	38	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSHSKAILAEAKKSND AQAPK
	39	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSKSKAILAEAKKINDAQAPK
	40	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSKSKAILAEAKKSND AQAPK
	41	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSQSKAILAEAKKRND AQAPK
	42	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQPLKDDPSSSKAILAEAKKINDAQAPK
	43	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQQLKDDPSDSKAILAEAKKVND AQAPK
	44	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQQLKDDPSNSKAILAEAKKSND AQAPK
	45	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQQLKDDPSQSKAILAEAKKINDAQAPK
	46	VDKFDKEAQEAQAFYEILHLPNLTEEQRNAFIQQLKDDPSRSKAILAEAKKVND AQAPK

Fig. 10C

Denotation	SEQ ID NO:	Amino acid sequence
	47	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQRLKDDPSQSKAILAEAKKRNDAAQAPK
	48	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQRLKDDPSQSKAILAEAKKSNDAAQAPK
	49	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQRLKDDPSRSKAILAEAKKINDAAQAPK
	50	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQRLKDDPSRSKAILAEAKKLNDAAQAPK
	51	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQSLKDDPSRSKAILAEAKKRNDAAQAPK
	52	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQSLKDDPSRSKAILAEAKKSNDAAQAPK
	53	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQTLKDDPSKSKAILAEAKKINDAAQAPK
	54	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQTLKDDPSQSKAILAEAKKINDAAQAPK
	55	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQTLKDDPSRSKAILAEAKKRNDAAQAPK
	56	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQTLKDDPSSSKAILAEAKKRNDAAQAPK
	57	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQTLKDDPSKSKAILAEAKKGNDAQAPK
	58	VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQSLKDDPSVSKAILAEAKKLNDAAQAPK
Z	59	VDNKFNKEQQNAFAFYEILHLPNLNEEQRNAFIQSLKDDPSQSANLLAEAKKLNDAAQAPK
C	60	ADNKFNKEQQNAFAFYEILHLPNLTEEQRNAFIQSLKDDPSVSKEILAEAKKLNDAAQAPK
B	61	ADNKFNKEQQNAFAFYEILHLPNLNEEQRNAFIQSLKDDPSQSANLLAEAKKLNDAAQAPK
A	62	ADNNFNKEQQNAFAFYEILNMPNLNEEQRNAFIQSLKDDPSQSANLLAEAKKLNESQAPK
D	63	ADAQONKFNKDQSAFYEILNMPNLNEEQRNAFIQSLKDDPSQSTNVLGEAKKLNESQAPK
E	64	AQHDEAQONAFYQVILNMPNLNADQRNGFIQSLKDDPSQSANVLGEAQKLNDQSAPK
ZGE90-6	203	(VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQALKDDPSRSKAILAEAKKINDAAQAPK) εC
ZGE91-6	204	(VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQSLKDDPSGSKAILAEAKKVNDAAQAPK) εC
ZGE96-6	205	(VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQELKDDPSGSKAILAEAKKLNDAAQAPK) εC
	206	(VDAKFDKEAQEAFAFYEILHLPNLTEEQRNAFIQSLKDDPSVSKAILAEAKKLNDAAQAPK) εC
	208	VDNKFNKEQQNAFAFYEILHLPNLNEEQRNAFIQGLKDDPSGSANLLAEAKKSNDAAQAPK
	209	VDNKFNKEQQNAFAFYEILHLPNLNEEQRNAFIQGLKDDPSGSANLLAEAKKINDAAQAPK
	210	VDNKFNKEQQNAFAFYEILHLPNLNEEQRNAFIQSLKDDPSRSANLLAEAKKVNDAAQAPK

Fig. 10D

Denotation	SEQ ID NO:	Amino acid sequence
	211	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSRSANLLAEAKKLNDAAQAPK
	212	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQQLKDDPSRSANLLAEAKKINDAAQAPK
	213	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQALKDDPSRSANLLAEAKKINDAAQAPK
	214	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQSLKDDPSGSANLLAEAKKVNDAAQAPK
	215	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQTLKDDPESANLLAEAKKRNDAAQAPK
	216	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSESANLLAEAKKRNDAAQAPK
	217	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQALKDDPSDSANLLAEAKKLNDAAQAPK
	218	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQALKDDPSGSANLLAEAKKRNDAAQAPK
	219	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIOELKDDPSGSANLLAEAKKLNDAAQAPK
	220	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIOHLKDDPSKSANLLAEAKKLNDAAQAPK
	221	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSDSANLLAEAKKVNDAAQAPK
	222	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSQSANLLAEAKKINDAAQAPK
	223	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQQLKDDPSDSANLLAEAKKINDAAQAPK
	224	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQALKDDPSKSANLLAEAKKLNDAAQAPK
	225	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQALKDDPSRSANLLAEAKKLNDAAQAPK
	226	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQDLKDDPSKSANLLAEAKKINDAAQAPK
	227	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQDLKDDPSNSANLLAEAKKSNDAAQAPK
	228	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQDLKDDPSRSANLLAEAKKVNDAAQAPK
	229	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIOELKDDPSRSANLLAEAKKLNDAAQAPK
	230	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSDSANLLAEAKKRNDAAQAPK
	231	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPESNSANLLAEAKKINDAAQAPK
	232	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSRSANLLAEAKKINDAAQAPK
	233	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIQGLKDDPSRSANLLAEAKKRNDAAQAPK
	234	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIOHLKDDPSKSANLLAEAKKINDAAQAPK
	235	VDNKFNKEQQNAFYIEILHLPNLNEEQRNAFIOHLKDDPSQSANLLAEAKKSNDAAQAPK

Fig. 10E

Denotation	SEQ ID No:	Amino acid sequence
	236	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSRSANLLAEAKKINDAQAPK
	237	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSSSANLLAEAKKRNDAAQAPK
	238	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSSSANLLAEAKKVNDAAQAPK
	239	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSESANLLAEAKKINDAQAPK
	240	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSKSANLLAEAKKLNDAAQAPK
	241	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQHLKDDPSRSANLLAEAKKINDAQAPK
	242	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQNLKDDPSHSANLLAEAKKSNDAAQAPK
	243	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSDSANLLAEAKKRNDAAQAPK
	244	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSHSANLLAEAKKRNDAAQAPK
	245	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSHSANLLAEAKKSNDAAQAPK
	246	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSKSANLLAEAKKLNDAAQAPK
	247	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSKSANLLAEAKKSNDAAQAPK
	248	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSQSANLLAEAKKRNDAAQAPK
	249	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQPLKDDPSSSANLLAEAKKINDAAQAPK
	250	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQQLKDDPSDSANLLAEAKKVNDAAQAPK
	251	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQQLKDDPSNSANLLAEAKKSNDAAQAPK
	252	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQQLKDDPSQSANLLAEAKKINDAAQAPK
	253	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQQLKDDPSRSANLLAEAKKVNDAAQAPK
	254	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQRLKDDPSQSANLLAEAKKRNDAAQAPK
	255	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQRLKDDPSQSANLLAEAKKSNDAAQAPK
	256	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQRLKDDPSRSANLLAEAKKINDAAQAPK
	257	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQRLKDDPSRSANLLAEAKKLNDAAQAPK
	258	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQSLKDDPSRSANLLAEAKKRNDAAQAPK
	259	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQSLKDDPSRSANLLAEAKKSNDAAQAPK
	260	VDNKFNKEQQNAFYEILHLPNLNEEQRNAFIQTLKDDPSKSANLLAEAKKINDAAQAPK

Fig. 10F

Denotation	SEQ ID NO:	Amino acid sequence
	261	VDNKFNKEQQNAFYEIILHLPNLNLEEQRNAFIQTLKDDPSQSANLLAEAKKINDAQAPK
	262	VDNKFNKEQQNAFYEIILHLPNLNLEEQRNAFIQTLKDDPSRSANLLAEAKKRNDQAQPK
	263	VDNKFNKEQQNAFYEIILHLPNLNLEEQRNAFIQTLKDDPSSSANLLAEAKKRNDQAQPK
	264	VDNKFNKEQQNAFYEIILHLPNLNLEEQRNAFIQTLKDDPSKSANLLAEAKKGNDAQAPK

Fig. 11A




Fig. 11B




Fig. 11C




Fig. 11D




Fig. 12A




Fig. 12B




Fig. 12C





Fig. 12D



FC BINDING POLYPEPTIDES

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of PCT/EP2022/076493, filed on Sep. 23, 2022, which claims the priority benefit of GB Application No. 2113626.2 filed on Sep. 24, 2021, the entire contents of which are incorporated herein by reference.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted electronically in ST.26 XML format and is hereby incorporated by reference in its entirety. Said ST.26 XML copy, created on Sep. 15, 2022, is named Sequence_Listing.xml and is 580,000 bytes in size.

FIELD OF THE INVENTION

[0003] The present disclosure relates to a class of engineered polypeptides having a binding affinity for the Fc region of immunoglobulins while exhibiting a significantly reduced binding affinity to the VH3 region of immunoglobulins. The present disclosure also relates to methods for isolating an immunoglobulin using said polypeptides as well as to related products.

BACKGROUND

[0004] Immunoglobulins represent the most prevalent biopharmaceutical products in either manufacture or development worldwide. The high commercial demand for and hence value of this particular therapeutic market has led to the emphasis being placed on pharmaceutical companies to maximize the productivity of their respective mAb manufacturing processes whilst controlling the associated costs.

[0005] Affinity chromatography is used in most cases, as one of the key steps in the purification of these immunoglobulin molecules, such as monoclonal or polyclonal antibodies. A particularly interesting class of affinity reagents is proteins capable of specific binding to invariable parts of an immunoglobulin molecule, such interaction being independent of the antigen-binding specificity of the antibody. Such reagents can be widely used for affinity chromatography recovery of immunoglobulins from different samples, such as but not limited to serum or plasma preparations or cell culture derived feed stocks. An example of such a protein is staphylococcal protein A (SpA), containing domains capable of binding to the Fc and Fab portions of IgG immunoglobulins from different species. These domains are commonly denoted as the E-, D-, A-, B- and C-domains.

[0006] SpA-based proteins have due to their high affinity and selectivity found a widespread use in the field of biotechnology, e.g. as ligands in affinity chromatography for capture and purification of antibodies as well as for detection or quantification. At present, SpA-based affinity medium is probably the most widely used affinity medium for isolation of monoclonal antibodies and their fragments from different samples including industrial cell culture supernatants. Accordingly, various matrices comprising protein A or protein A-derived ligands are commercially available, for example, in the form of MabSelect™ SuRe, MabSelect™ SuRe LX, MabSelect™ Prisma protein A and HiScreen Fibro™ Prisma from Cytiva, Uppsala, Sweden.

[0007] Certain Protein A and Protein A-derived ligands have binding affinity for both the Fc part of an antibody and for some VH domains of antibodies, in particular VH3. As a result, co-purification of product-related impurities such as half-antibodies and truncated variants may occur and require elution schemes which are complex and/or not sufficiently mild. When using these ligands in library applications aiming at finding novel binders against antibody domains, having inherent VH domain interactions can be problematic.

SUMMARY OF THE INVENTION

[0008] It is an object of the present disclosure to provide Fc binding SpA-derived ligands which exhibit no, or significantly reduced affinity for the VH3 region of immunoglobulins. Such ligands could for example be used in methods for isolating an immunoglobulin and/or fragments thereof.

[0009] It is an object of the present disclosure to provide Fc binding SpA-derived ligands allowing for efficient isolation of immunoglobulins and/or fragments thereof while alleviating the abovementioned and other drawbacks of the prior art.

[0010] It is one object to provide Fc binding SpA-derived ligands enabling mild elution of purified antibodies.

[0011] It is a further object to provide Fc binding SpA-derived ligands which decreases or prevents co-purification of product-related impurities such as half-antibodies and truncated variants.

[0012] It is also a further object to provide Fc binding SpA-derived ligands which avoid competing interactions, e.g. to simplify identification of altered immunoglobulin variants showing changed sensitivity to pH-elution.

[0013] It is one object to provide Fc binding SpA-derived ligands which avoid competing interactions and therefore are useful in platform solutions for purification of all Fc containing products.

[0014] It is one object to provide Fc binding SpA-derived ligands which avoid competing binding interactions from VH3 and thus exhibit increased binding capacity for Fc containing products, in other words Fc binding SpA-derived ligands which are capable of binding more Fc containing products.

[0015] These and other objects which are evident to the skilled person from the present disclosure are met by different aspects of the invention as claimed in the appended claims and as generally disclosed herein.

[0016] Thus, in the first aspect of the disclosure, there is provided an Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof, wherein said polypeptide has binding affinity for an Fc region of an immunoglobulin, and has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:59 for the same VH3 region.

[0017] As an alternative, the Fc binding polypeptides as disclosed herein may be evaluated for binding affinity for a VH3 region of trastuzumab compared to the affinity of SEQ ID NO:58 for the same VH3 region. Thus, as an alternative, there is provided an Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof, wherein said polypeptide has binding affinity for an Fc region of an immunoglobulin, and has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:58 for the same VH3 region. Thus, it will be understood that in some embodiments, there is

provided an Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof as disclosed herein, which has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:58 and of SEQ ID NO:59 for the same VH3 region.

[0018] The skilled person will appreciate that the Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof as disclosed herein may for example, but not necessarily, be derived from any one of domains A (SEQ ID NO:62), B (SEQ ID NO:61), C (SEQ ID NO:60), D (SEQ ID NO:63) and E (SEQ ID NO:64) of SpA or derivatives thereof, such as domain Z (SEQ ID NO:59) or variants thereof, such as SEQ ID NO:58. The Fc binding polypeptide derived from a SpA domain may thus be a derivative, a mutant, a variant or a fragment of an SpA domain as defined above.

[0019] The Fc binding polypeptides of the present disclosure thus are characterized by their retained inherent ability to bind to the Fc (fragment crystallizable region) of antibodies/immunoglobulins, in particular to the Fc region of the antibody subclasses IgG1, IgG2 and IgG4, and their reduced or abolished binding affinity for the VH3 region of the antibody trastuzumab. The skilled person appreciates that similar effect will be observed for antibodies or fragments thereof containing the same or similar VH3 region as trastuzumab.

[0020] As used herein the term VH3 in the context of binding affinity for VH3 refers to the VH3 region of the antibody trastuzumab (trade names Herceptin™ (Roche), Trazimera™ (Pfizer)) which region comprises the amino acid residues in the following positions according to the Kabat numbering system: H15:G; H17:S; H19:R; H57:T; H59:Y; H64:K; H65:G; H66:R; H68:T; H70:S; H81:Q; H82a:N and H82b:S. For avoidance of any doubt, the term “VH3 region of trastuzumab” and the term “VH3” (without any additional specification) are used to refer to the VH3 region of the antibody trastuzumab.

[0021] The present inventors have surprisingly found that by introducing mutations in positions X₃₃, X₄₀ and X₅₁ of Fc binding polypeptides derived from SpA or domains thereof, the binding affinity for VH3 of trastuzumab is significantly reduced or abolished. X₃₃, X₄₀ and X₅₁ refer to positions 33, 40 and 51 in a SpA domain, such as SEQ ID NO:58 and 59. Thus, the presently identified mutations in positions X₃₃, X₄₀ and X₅₁ of the Fc binding polypeptides as defined herein significantly reduce or abolish binding affinity for said VH3. The Fc binding polypeptides comprising amino acids as defined herein in positions X₃₃, X₄₀ and X₅₁ lose at least a significant part of their affinity for VH3 while retaining the affinity for the Fc region.

[0022] According to a second aspect, there is provided an Fc binding polypeptide comprising a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii),

[0023] wherein i), ii) and iii) are defined as follows:

[0024] v) EX₂₅QX₂₇X₂₈X₂₉X₃₀X₃₂X₃₃
LX₃₅X₃₆X₃₇PSX₄₀SX₄₂X₄₃X₄₄LX₄₆EAX₄₉X₅₀X₅₁N₂X₅₃
X₅₄

wherein, independently from each other,

[0025] X₂₅ is selected from E and D;

[0026] X₂₇ is selected from R and H;

[0027] X₂₈ is selected from N, A, S, H and W;

[0028] X₂₉ is selected from A, G and K;

[0029] X₃₀ is selected from F and A;

[0030] X₃₂ is selected from Q and H;

[0031] X₃₅ is selected from K, R and H;

[0032] X₃₆ is selected from D and H;

[0033] X₃₇ is selected from D and E;

[0034] X₄₂ is selected from A, K, L, R and T;

[0035] X₄₃ is selected from N, E, A, K and S;

[0036] X₄₄ is selected from L, I and V;

[0037] X₄₆ is selected from A, G and K;

[0038] X₄₉ is selected from K, Q and R;

[0039] X₅₀ is selected from K and R;

[0040] X₅₃ is selected from D, E and K; and

[0041] X₅₄ is selected from A and S;

[0042] ii) an amino acid sequence which has at least 83% identity to a sequence defined by i)

[0043] iii) an amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, residues 24-54 in SEQ ID NO:60, residues 24-54 in SEQ ID NO:61, residues 24-54 in SEQ ID NO:62, residues 27-57 in SEQ ID NO:63 and residues 22-52 in SEQ ID NO:64,

[0044] wherein additionally, in each of i), ii) and iii) independently from each other,

[0045] X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0046] X₄₀ is selected from E, G, R, D, K, Q, N, H and S, and

[0047] X₅₁ is selected from L, V, S, I, R and G;

with the proviso that when X₅₁ is L then X₃₃X₄₀ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X₅₁ is G then X₃₃X₄₀ is TK.

[0048] In one embodiment of the second aspect, said Fc binding polypeptide comprises a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii) as defined above, and wherein additionally, in each of i), ii) and iii) independently from each other,

[0049] X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0050] X₄₀ is selected from E, G, R, D, K, Q, N, H and S, and

[0051] X₅₁ is selected from L, V, S, I, R and G;

[0052] with the proviso that when X₅₁ is L then X₃₃X₄₀ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X₅₁ is G then X₃₃X₄₀ is TK;

and

[0053] wherein said Fc binding polypeptide has a lower affinity for VH3 region of trastuzumab than SEQ ID NO:59.

[0054] As explained above, the Fc binding polypeptides as disclosed herein may be evaluated for binding affinity for a VH3 region of trastuzumab compared to the affinity of SEQ ID NO:58 for the same VH3 region. Thus, as an alternative, there is provided an Fc binding polypeptide comprising a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii) as defined above, and wherein additionally, in each of i), ii) and iii) independently from each other,

[0055] X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0056] X₄₀ is selected from E, G, R, D, K, Q, N, H and S, and

[0057] X₅₁ is selected from L, V, S, I, R and G;

[0058] with the proviso that when X_{51} is L then $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X_{51} is G then $X_{33}X_{40}$ is TK.

[0059] As an additional alternative, there is provided an Fc binding polypeptide comprising a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii) as defined above, and wherein additionally, in each of i), ii) and iii) independently from each other,

[0060] X_{33} is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0061] X_{40} is selected from E, G, R, D, K, Q, N, H and S, and

[0062] X_{51} is selected from L, V, S, I, R and G; with the proviso that when X_{51} is L then $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X_{51} is G then $X_{33}X_{40}$ is TK; and

[0063] wherein said Fc binding polypeptide has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:58 for the same VH3 region.

[0064] Again, it will be understood that in some embodiments, there is provided an Fc binding polypeptide as disclosed herein, which has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:58 and of SEQ ID NO:59 for the same VH3 region. It will be understood by the skilled person that the comparisons of binding affinity for the VH3 region of trastuzumab exhibited by the Fc binding polypeptides of the present disclosure may be made using SEQ ID NO:59 and/or SEQ ID NO:58 as a reference.

[0065] To clarify, it will be appreciated that the % identity in ii) and iii) does not relate to X_{33} , X_{40} and X_{51} . Thus, the amino residues in positions X_{33} , X_{40} and X_{51} are as defined above in Fc binding polypeptides encompassed by the definition according to ii) and iii).

[0066] The above definition of a class of sequence related, Fc binding polypeptides is based on a statistical analysis of a number of Fc binding polypeptide variants of a parent scaffold, that were selected for the properties of at least maintained affinity for Fc and significantly reduced affinity for VH3 in selection experiments.

[0067] It is also envisioned that said Fc binding polypeptide will exhibit a lower affinity for VH3 region of trastuzumab than otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SQL. Thus a comparison may be made between any otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SQL and the Fc binding polypeptide of the present disclosure. In one embodiment, said Fc binding polypeptide has a lower affinity for the VH3 region of trastuzumab than an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SQL. In yet another embodiment, said Fc binding polypeptide has a lower affinity for the VH3 region of trastuzumab than an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SQL, and than SEQ ID NO:59.

[0068] It is also envisioned that said Fc binding polypeptide will exhibit a lower affinity for VH3 region of trastuzumab than otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL. Thus, in an alternative embodiment, a comparison may be made between any otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL and the Fc binding polypeptide of the present disclosure. In an alternative embodiment, said Fc binding polypeptide has

a lower affinity for the VH3 region of trastuzumab than an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL. In yet another embodiment, said Fc binding polypeptide has a lower affinity for the VH3 region of trastuzumab than an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL, and than SEQ ID NO:58.

[0069] The Fc binding polypeptides may comprise the identified sequence A, which corresponds to the region which interacts and binds to VH3 of the parent scaffold. This region constitutes two alpha helices, namely helix 2 and helix 3, within a three-helical bundle protein domain. Helix 1 and helix 2 of this three-helical bundle protein constitute a binding surface for interaction with the Fc region of antibodies.

[0070] As the skilled person will realize, the function of any polypeptide is dependent on the tertiary structure of the polypeptide. It is therefore possible to make minor changes to the sequence of amino acids in a polypeptide without affecting the function thereof. Thus, the disclosure encompasses modified variants of the Fc binding polypeptide, which are such that the Fc binding characteristics are retained, while the binding characteristics for VH3 are significantly reduced or abolished.

[0071] The modification of VH3 binding affinity identified by the present inventors, is applicable to Fc binding polypeptides based on the different three-helical domains of Protein A from *Staphylococcus aureus*, such as any one of the domains A (SEQ ID NO:62), B (SEQ ID NO:61), C (SEQ ID NO:60), D (SEQ ID NO:63) and E (SEQ ID NO:64), in particular domain B, and derivatives thereof. In particular, said modification may be applicable to Fc binding polypeptides based on the three-helical bundle protein domain Z (SEQ ID NO:59), which domain Z is derived from domain B of staphylococcal Protein A or to the variant SEQ ID NO:58. Such Fc binding polypeptides have been described in: WO2003080655, EP230869A2, Sjö Dahl, Eur J Biochem 1977 September; 78(2):471-90, WO2008039141, WO2015005859, U.S. Ser. No. 10/308,690, WO2016079033, WO2017194596, U.S. Pat. Nos. 8,859, 726, 9,187,555, 9,683,013, U.S. Ser. No. 10/189,891, U.S. Pat. No. 9,663,558, WO2016079034, JP2006304633A, WO2007097361, EP1992692A1, EP2202310A2, WO2012083425, WO2012086660, US20120208234, EP2495254A1, WO2012133342, WO2013109302A2, WO2015034000, WO2015034056, CN105481954A, WO2016152946A1, WO2017009421, WO2017014261, WO2017014260, WO2018009006, WO2018029158, WO2018029157, WO2019030156, CN109721645A, WO2019093439, WO2020040307 and WO2020157281, which are hereby incorporated by reference.

[0072] Thus, the present inventors envision that the mutations in positions X_{33} , X_{40} and X_{51} as disclosed herein may be introduced into any one of the previously known Fc binding polypeptides to achieve the effect of significantly reduced or abolished binding affinity for VH3, while retaining the Fc binding characteristics.

[0073] Also encompassed by the present disclosure is an Fc binding polypeptide comprising a Sequence A with at least 83% identity to a polypeptide as defined in i). In some embodiments, the polypeptide may comprise a sequence which is at least 87%, such as at least 90%, such as at least 93%, such as at least 96% identical to a polypeptide as defined in i). For example, it is possible that an amino acid

residue belonging to a certain functional grouping of amino acid residues (e.g. hydrophobic, hydrophilic, polar etc.) could be exchanged for another amino acid residue from the same functional group. In some embodiments, such changes may be made in any position except for X_{33} , X_{40} and X_{51} of the sequence of the Fc binding polypeptide as disclosed herein. In other embodiments, such changes may be made only in the non-variable positions, also denoted scaffold amino acid residues. In such cases, changes are not allowed in the variable positions, i.e. positions denoted with an “X” in sequence i).

[0074] Also encompassed by the present disclosure is an Fc binding polypeptide comprising a Sequence A which Sequence A has at least 70% identity to any sequence selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, residues 24-54 in SEQ ID NO:60, residues 24-54 in SEQ ID NO:61, residues 24-54 in SEQ ID NO:62, residues 27-57 in SEQ ID NO:63 and residues 22-52 in SEQ ID NO:64. In some embodiments, the polypeptide may comprise a sequence which is at least 74%, such as at least 77%, such as at least 80%, such as at least 83%, such as at least 87%, such as at least 90%, such as at least 93%, such as at least 96%, identical to any sequence selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, residues 24-54 in SEQ ID NO:60, residues 24-54 in SEQ ID NO:61, residues 24-54 in SEQ ID NO:62, residues 27-57 in SEQ ID NO:63 and residues 22-52 in SEQ ID NO:64. In particular embodiments, said sequence is selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, and residues 24-54 in SEQ ID NO:61, such as the group consisting of: residues 24-54 in SEQ ID NO:58 and residues 24-54 in SEQ ID NO:59. In one embodiment, said sequence is residues 24-54 in SEQ ID NO:58.

[0075] The term “% identity”, as used throughout the disclosure, may for example be calculated as follows. The query sequence is aligned to the target sequence using the CLUSTAL W algorithm (Thompson et al, Nucleic Acids Research, 22: 4673-4680 (1994)). A comparison is made over the window corresponding to the shortest of the aligned sequences. The shortest of the aligned sequences may in some instances be the target sequence. In other instances, the query sequence may constitute the shortest of the aligned sequences. The amino acid residues at each position are compared and the percentage of positions in the query sequence that have identical correspondences in the target sequence is reported as % identity.

[0076] In one embodiment, the Fc binding polypeptide comprising a Sequence A, which consists of an amino acid sequence selected from i), ii) and iii) comprises amino acid residues selected independently from T, S, G, Q, A, E, H, R, P, D, K and N for X_{33} ; from E, G, R, D, K, Q, N, H and S for X_{40} ; and L, V, S, I, R and G for X_{51} ; and in the case wherein X_{51} is L then $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X_{51} is G then $X_{33}X_{40}$ is TK.

[0077] In one particular embodiment provided herein, in each of i), ii) and iii) independently from each other,

[0078] X_{33} is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0079] X_{40} is selected from E, G, R, D, K, Q, N, H and S; and

[0080] X_{51} is selected from L, V, S, I and R;

with the proviso that when X_{51} is L, $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and wherein said Fc binding polypeptide has a lower affinity for VH3 of trastuzumab than SEQ ID NO:58.

[0081] As explained above, the comparison may alternatively be made with an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL; or with both an otherwise identical Fc binding polypeptide wherein $X_{33}X_{40}X_{51}$ is SVL and SEQ ID NO:58.

[0082] In one embodiment of the Fc binding polypeptides as disclosed herein, said Sequence A fulfills the criteria i) and ii), or the criteria i) and iii), or the criteria ii) and iii). In one embodiment, said Sequence A fulfills all criteria i), ii) and iii).

[0083] In one embodiment according to the second aspect as disclosed herein, X_{33} is selected from T, S, G, Q, A, E, H, R, P, D and K, such as selected from T, S, G, Q, A, H, R, P, D and K, such as selected from T, S, G, Q, A, H, R, P and D or selected from T, S, G, Q, A, H, R, P and K; such as selected from T, S, G, Q, A, H, R and P; such as selected from T, S, G, Q, A, H and P or selected from T, G, Q, A, H, R and P; such as selected from T, G, Q, P, A and H; such as selected from T, G, Q, H and P; such as selected from T, G, Q and P or selected from G, Q, H and P or selected from T, G, H and P; such as the group selected from G and P; such as G. In one embodiment, X_{33} is selected from T, S, G, Q, A, E and H, such as selected from T, S, G, Q, A and E, such as selected from S, G, Q and A. In one embodiment, X_{33} is selected from S, G, A, E and H, such as selected from S, G, A and E. In one embodiment, X_{33} is selected from A, S and E. In one embodiment, X_{33} is selected from A and S or selected from A and E or selected from S and E. In one embodiment, X_{33} is S.

[0084] In one embodiment, X_{40} is selected from E, G, R, D, K, Q, N and S or selected from E, G, R, D, K, Q, H and S or selected from G, R, D, K, Q, N, H and S, such as selected from G, R, D, K, Q, and S, such as selected from G, R, D, K and Q; such as selected from R, D, K and Q; such as selected from R, K and Q; such as selected from R and K; such as is R or is K. In one embodiment, X_{40} is selected from E, G, R, D, K and Q, such as selected from E, G, R, D, K and Q, such as selected from E, G, R, D and K or selected from E, G, R, D and Q. In one embodiment, X_{40} is selected from G, R, E and D. In one embodiment, X_{40} is selected from G, R and D. In one embodiment, X_{40} is selected from G, R and K. In one embodiment, X_{40} is selected from G and R. In one embodiment, X_{40} is G. In one embodiment, X_{40} is R.

[0085] In one embodiment, X_{51} is selected from L, V, I, R and S. In one embodiment X_{51} is selected from L, I, R and S. In one embodiment, X_{51} is selected from L, V, I and R. In one embodiment, X_{51} is selected from I, V and R; or is selected from L, I and R; or is selected from V, L and I; or is selected from R, I and L. In one embodiment, X_{51} is selected from V and L; or is selected from V and I; or is selected from V and R; or is selected from L and I; or is selected from L and R; or is selected from I and R. In one embodiment, X_{51} is V or L or I or R. In one embodiment, X_{51} is 1. In one embodiment, X_{51} is V.

[0086] In one embodiment, X_{51} is selected from L, V, S, I and R with the proviso that when X_{51} is L then $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and

KK. In another embodiment, X_{51} is selected from L, V, S, I and R with the proviso that when X_{51} is L then $X_{33}X_{40}$ are AD or HK.

[0087] In one embodiment of the Fc binding polypeptide, at least one, such as two or all three, of amino acid residues X_{33} , X_{40} and X_{51} is/are (an) uncharged amino acid residue (s). In one embodiment, the amino acid residue in X_{33} is an uncharged amino acid. In one embodiment, the amino acid residue in X_{40} is an uncharged amino acid.

[0088] In one particular embodiment there is provided an Fc binding polypeptide, wherein at least two of the amino acid residues in positions X_{33} , X_{40} and X_{51} are mutated compared to SEQ ID NO:58 or 59. In particular, positions X_{40} and X_{51} may be mutated compared to SEQ ID NO:58 or 59.

[0089] In a more specific embodiment defining a sub-class of Fc binding polypeptides as disclosed herein, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GGS, GNI, GQI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, PQR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, ROS, RRI, RRL, SGV, SRR, SRS, SRV, TER, TKG, TKI, TQI, TRR and TSR, such as the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GNI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, PQR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, ROS, RRI, RRL, SGV, SRR, SRS, SRV, TER, TKI, TQI, TRR and TSR. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGS, GGI, SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV, GQI and QDI, or the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV and QDI. In another embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, AGR and GDV. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, ARI, SGV, AGR, EGL and HKL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, TER, AGR, EGL and GDV. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, QRI, ARI, SGV, AGR and GDV. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV, GQI and QDI. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, GRL, ARI, SGV, AGR and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI, SGV, AGR and EGL. In another embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of SGV, AGR and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI, AGR and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI, SGV and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI, SGV and AGR.

In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI and SVG. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI and AGR, or the group consisting of ARI and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of SGV

and AGR. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of SGV and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ are selected from the group consisting of AGR and EGL. In one embodiment, $X_{33}X_{40}X_{51}$ is ARI or SGV or AGR or EGL. In one particular embodiment, $X_{33}X_{40}X_{51}$ is SGV.

[0090] Herein, " X_n " and " X_m " are used to indicate amino acid residues in positions n and m corresponding to positions n and m in e.g. SEQ ID NO 59 as defined herein, wherein n and m are integers which indicate the position of an amino acid residue within said sequence as counted from the N-terminal end of said sequence. For example, X_{24} and X_{13} indicate the amino acid residue in positions that corresponds to positions twenty-four and thirteen, respectively, from the N-terminal end of e.g. SEQ ID NO 59 as defined herein, in an alignment. To clarify the X numbering as used herein is based on the full-length scaffold comprising 58 amino acid residues. Hence X_{24} is to be understood as being the 24th amino acid residue in the 58-mer. The skilled person is able to make alignments of sequences to determine the position of an amino acid in accordance with the above.

As described in detail in the experimental section to follow, the selection of Fc binding polypeptide variants has led to the identification of a number of individual amino acid sequences according to Sequence A as defined herein. These sequences constitute individual embodiments of sequence i) according to this aspect. The sequences of individual amino acid sequences according to Sequence A correspond to amino acid positions 24-54 in SEQ ID NO:1-57 and SEQ ID NO:208-264 presented in FIG. 10, as well as SEQ ID NO:269-305. Hence, in one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57, 208-264 and 269-305, such as SEQ ID NO:1-56 and 208-263. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO: SEQ ID NO:2-57, 209-264 and 269-305. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223 or the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO: SEQ ID NO:2-16 and 209-223. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:7 and 214.

[0091] In one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:208-264 and 269-305, such as SEQ ID NO:208-263. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209-264 and 269-305. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:208-223 or the group consisting of SEQ ID NO:209-221 and 223. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209-223. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:213, 214, 218 and 219. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:213, 214 and 218. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in SEQ ID NO:214.

[0092] In one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2-57. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 or the group consisting of SEQ ID NO:2-14 and 16. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2-16. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11 and 12. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7 and 11. In one embodiment, sequence i) corresponds to the sequence from position 24 to position 54 in SEQ ID NO:7.

[0093] In some embodiments of the present disclosure, the polypeptide derived from SpA or Sequence A as defined above “forms part of” a three-helix bundle protein domain. This is understood to mean that the sequence of the SpA derived polypeptide or Sequence A is “inserted” into or

“grafted” onto the sequence of the original three-helix bundle domain, such that the grafted sequence replaces a similar structural motif in the original domain. For example, without wishing to be bound by theory, the Sequence A is thought to constitute two of the three helices of a three-helix bundle and can therefore replace such a two-helix motif within any three-helix bundle. As the skilled person will realize, the replacement of two helices of the three-helix bundle domain by the two Sequence A helices has to be performed so as not to affect the basic structure of the polypeptide. That is, the overall folding of the Ca backbone of the polypeptide according to this embodiment of the invention is substantially the same as that of the three-helix bundle protein domain of which it forms a part, e.g. having the same elements of secondary structure in the same order etc. Thus, a Sequence A according to the disclosure “forms part” of a three-helix bundle domain if the polypeptide according to this embodiment has the same fold as the original domain, implying that the basic structural properties are shared, those properties e.g. resulting in similar CD spectra. The skilled person is aware of other parameters that are relevant.

[0094] Thus, in one embodiment the SpA derived polypeptide or Sequence A forms part of a three-helix bundle protein domain. In some embodiments, said Sequence A forms, or essentially forms, part of two helices with an interconnecting loop, within said three-helix bundle protein domain. In particular, said three-helix bundle protein domain may be selected from domains from bacterial receptor proteins, in other words from bacterial receptor domains. Non-limiting examples of such domains are the five different three-helical domains of Protein A from *Staphylococcus aureus*, such as domain A, B, C, D and E, in particular domain B and derivatives thereof and domain C and derivatives thereof. An example of a derivative of said domain B is domain Z. Another example of a derivative of domain C is domain C with the mutation G29A. In one embodiment, said three-helix bundle protein domain is selected from domains of protein A from *Staphylococcus aureus* or derivatives thereof. In some embodiments, the three-helix bundle protein domain is a variant of domain Z, which is derived from domain B of staphylococcal Protein A. In some embodiments, the three-helical bundle protein domain is a domain C comprising the mutation G29A.

[0095] Thus, in one embodiment, there is provided an Fc binding polypeptide as defined herein, further comprising a Sequence B arranged N-terminally of said Sequence A, which Sequence B consists of an amino acid sequence selected from iv) and v), and wherein iv) and v) are defined as follows:

[0096] iv) $X_8X_9X_{10}X_{11}AFYX_{15}IX_{17}X_{18}X_{19}PX_{21}LX_{23}$, wherein, independently from each other,

[0097] X_8 is selected from E, D and A;

[0098] X_9 is selected from Q, A, L, W, E, V, K, T and H;

[0099] X_{10} is selected from Q and H;

[0100] X_{11} is selected from N, A, S, E, K, H, Q, Y, T, F, L, W, I, M, V and R;

[0101] X_{15} is selected from E, H and Q;

[0102] X_{17} is selected from L and H;

[0103] X_{18} is selected from H, N and K;

[0104] X_{19} is selected from L and M;

[0105] X_{21} is selected from N, Y and S;

[0106] X_{23} is selected from T and N;

[0107] v) an amino acid sequence which has at least 75% identity to a sequence defined by iv).

[0108] In some embodiments where the sequence of Fc binding polypeptide as disclosed herein forms part of a three-helix bundle protein domain, it essentially forms part of at least three helices and the interconnecting loop between helix 1 and helix 2 (referred to as loop 1 (L1)) and the interconnecting loop between helix 2 and helix 3 (referred to as loop 1 (L2)). In this case, Sequence B essentially corresponds to helix 1 and the and L1. As used herein, the term “essentially corresponds to” is to be interpreted as making up the main part of the structure, in other words a few amino acid residues (for example, but not limited to, 1, 2, 3, 4 or 5 residues) may be missing compared to the original three-helix bundle protein domain.

[0109] As discussed above, polypeptides comprising minor changes as compared to the above amino acid sequences without largely affecting the tertiary structure and the function thereof are also within the scope of the present disclosure. Thus, in some embodiments, sequence v) has at least 87% identity, such as at least 90% identity, such as at least 93% identity, such as at least 96% identity with the sequence iv).

[0110] As explained above, “X_n” and “X_m” are used to indicate amino acid residues in positions n and m corresponding to positions n and m in SEQ ID NO 59 as defined above, wherein n and m are integers which indicate the position of an amino acid residue within said sequence as counted from the N-terminal end of said sequence. Again, the X numbering as used herein is based on the full-length scaffold comprising 58 amino acid residues. Hence X₈ is to be understood as being the 8th amino acid residue in the 58-mer. The skilled person is able to make alignments of sequences to determine the position of an amino acid in accordance with the above.

[0111] Thus in yet another embodiment, there is provided an Fc binding polypeptide as defined herein, which comprises a binding module sequence C, which Sequence C comprises or consists of Sequence A as defined herein and Sequence B as defined herein, in the following order from the N-terminus to the C-terminus

[0112] [Sequence B]-[Sequence A]

or comprises or consists of any amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, residues 8-54 in SEQ ID NO:60, residues 8-54 in SEQ ID NO:61, residues 8-54 in SEQ ID NO:62, residues 11-57 in SEQ ID NO:63 and residues 6-52 in SEQ ID NO:64 wherein, independently from each other,

[0113] X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0114] X₄₀ is selected from E, G, R, D, K, Q, N, H and S, and

[0115] X₅₁ is selected from L, V, S, I, R and G;

with the proviso that when X₅₁ is L then X₃₃X₄₀ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X₅₁ is G then X₃₃X₄₀ is TK.

[0116] In one embodiment, said sequence C is selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, and residues 8-54 in SEQ ID NO:61, such as the group consisting of: residues

8-54 in SEQ ID NO:58 and residues 8-54 in SEQ ID NO:59. In one embodiment, said sequence is residues 8-54 in SEQ ID NO:58.

[0117] In one embodiment, said sequence C has at least 70% identity to any sequence selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, residues 8-54 in SEQ ID NO:60, residues 8-54 in SEQ ID NO:61, residues 8-54 in SEQ ID NO:62, residues 11-57 in SEQ ID NO:63 and residues 6-52 in SEQ ID NO:64, has at least 72%, such as at least 74%, such as at least 76%, such as at least 78%, such as at least 80%, such as at least 82%, such as at least 85%, such as at least 87%, such as at least 89%, such as at least 91%, such as at least 93%, such as at least 85%, such as at least 97%, identity to any sequence selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, residues 8-54 in SEQ ID NO:60, residues 8-54 in SEQ ID NO:61, residues 8-54 in SEQ ID NO:62, residues 11-57 in SEQ ID NO:63 and residues 6-52 in SEQ ID NO:64. As previously explained, the % identity does not apply to the amino acid residues in positions X₃₃, X₄₀ and X₅₁. The amino acid residues in said positions are as defined above.

[0118] As discussed above, in some embodiments where the Fc binding polypeptides as disclosed herein form part of a three-helix bundle protein domain, they essentially form part of at least the three helices and the interconnecting loop between helix 1 and helix 2 (referred to as loop 1 (L1)) and the interconnecting loop between helix 2 and helix 3 (referred to as loop 1 (L2)). Herein, binding module Sequence C essentially forms part of at least the three helices and the interconnecting loop between helix 1 and helix 2 (referred to as loop 1 (L1)) and the interconnecting loop between helix 2 and helix 3 (referred to as loop 2 (L2)). It will be appreciated that the length of L1 and L2 may vary, for example by the addition or removal of one or more additional amino acid residues, provided that this variation does not significantly alter the three dimensional structure of polypeptide comprising binding module Sequence C.

[0119] As explained above, Sequence B comprises helix 1 (H1) and L1. The skilled person will appreciate that the amino acid sequences of H1 and L1 may be independently combined while maintaining the three-dimensional structure of the Fc binding polypeptide. Thus, in one embodiment, there is provided an Fc binding polypeptide as defined herein, which comprises a binding module Sequence C, which Sequence C consists of the sequences [H1], [L1] and [Sequence A] in the following order from the N-terminus to the C-terminus; [H1]-[L1]-[Sequence A], wherein [Sequence A] is as defined herein, and wherein, independently from each other, [H1] is selected from the group consisting of:

EQQNAFYEILH (SEQ ID NO:67); DQQAIFYEILH (SEQ ID NO:68); EAQEAIFYEILH (SEQ ID NO:69); DQQSAFYEILH (SEQ ID NO:70); AQQAIFYEILH (SEQ ID NO:71); EQQNAFYEILN (SEQ ID NO:72); EQQAIFYEILH (SEQ ID NO:73); AQQSAFYEILH (SEQ ID NO:74); EQQQAIFYEILH (SEQ ID NO:75); EHQNAFYEILH (SEQ ID NO:76); EAQNAFYEILH (SEQ ID NO:77); ETQNAFYEILH (SEQ ID NO:78); EQQSAFYEILH (SEQ ID NO:79); EAQNAFYKILH (SEQ ID NO:80); EQQEAIFYEILH (SEQ ID NO:81); EAQKAFYEILK (SEQ ID NO:82); EQHNAFYEILH (SEQ ID NO:83); EHHHALYHILH (SEQ ID NO:84); EQQKAFYAILH (SEQ ID NO:85); DQQSAFYEILN (SEQ

ID NO:86); AQQNAFYQVLN (SEQ ID NO:87); EQQKAFYEILH (SEQ ID NO:88); EQQKAFYEILK (SEQ ID NO:89); EQQNAFYEILS (SEQ ID NO:90); EQQNAFYHILH (SEQ ID NO:91); and EQQNAFYHILN (SEQ ID NO:92), and [L1] is selected from the group consisting of:

LPNLT (SEQ ID NO:93); LPNLN (SEQ ID NO:94); MPNLN (SEQ ID NO:95); LPYLT (SEQ ID NO:96); LPSLT (SEQ ID NO:97); MPNLL (SEQ ID NO:98); LPNGN (SEQ ID NO:99); LPNLS (SEQ ID NO:100); GPNAN (SEQ ID NO:101); VPNLN (SEQ ID NO:102); LPNLR (SEQ ID NO:103); and LPNLL (SEQ ID NO:104).

[0120] In one particular embodiment, H1 is selected from the group consisting of EQQNAFYEILH; DQQAIFYEILH; EAQEAIFYEILH; DQQSAIFYEILH; AQQAAIFYEILH; EQQNAFYEILN; EQQAIFYEILH; AQQSAIFYEILH; EQQQAIFYEILH; and EHQNAFYEILH, such as the group consisting of EQQNAFYEILH; DQQAIFYEILH; EAQEAIFYEILH; DQQSAIFYEILH; AQQAAIFYEILH; EQQNAFYEILN; and EQQAIFYEILH, such as the group consisting of EQQNAFYEILH; DQQAIFYEILH; EAQEAIFYEILH; DQQSAIFYEILH; and AQQAAIFYEILH, such as the group consisting of EQQNAFYEILH; DQQAIFYEILH and EAQEAIFYEILH. In one embodiment, H1 comprises or consists of EQQNAFYEILH. In one embodiment, H1 comprises or consists of DQQAIFYEILH. In one embodiment, H1 comprises or consists of EAQEAIFYEILH.

[0121] In one particular embodiment, L1 is selected from the group consisting of LPNLT; LPNLN; MPNLN; LPYLT; LPSLT; MPNLL; LPNGN, and LPNLS, such as the group consisting of LPNLT; LPNLN; MPNLN; LPYLT; PSLT, and MPNLL; such as the group consisting of LPNLT; LPNLN; MPNLN and LPYLT; such as the group consisting of LPNLT; LPNLN and MPNLN; such as the group consisting of LPNLT; and LPNLN. In one embodiment, L1 comprises or consists of LPNLT. In one embodiment, L1 comprises or consists of LPNLN.

[0122] In one embodiment, there is provided an Fc binding polypeptide as disclosed herein, which comprises a binding module Sequence C selected from the group consisting of: EQQNAFYEILHLPNLT-[Sequence A](SEQ ID NO:105); EQQNAFYEILHLPNLN-[Sequence A](SEQ ID NO:106); DQQAIFYEILHLPNLT-[Sequence A](SEQ ID NO:107); EAQEAIFYEILHLPNLT-[Sequence A](SEQ ID NO:108); DQQSAIFYEILHLPNLT-[Sequence A](SEQ ID NO:109); AQQAAIFYEILHLPNLT-[Sequence A](SEQ ID NO:110); EQQAIFYEILHLPNLT-[Sequence A](SEQ ID NO:111); AQQSAIFYEILHLPNLT-[Sequence A](SEQ ID NO:112); EQQNAFYEILNMPNLN-[Sequence A](SEQ ID NO:113); EAQNAFYEILHLPNLT-[Sequence A](SEQ ID NO:114); EQQQAIFYEILHLPNLT-[Sequence A](SEQ ID NO:115); EQQNAFYEILHLPYLT-[Sequence A](SEQ ID NO:116); ETQNAFYEILHLPNLT-[Sequence A](SEQ ID NO:117); AQQNAFYKILHLPNLT-[Sequence A](SEQ ID NO:118); EQQNAFYEILHLPNGN-[Sequence A](SEQ ID NO:119); EHQNAFYEILHLPNLN-[Sequence A](SEQ ID NO:120); EQQNAFYEILNMPNLL-[Sequence A](SEQ ID NO:121); EQQSAIFYEILHLPNLT-[Sequence A](SEQ ID NO:122); EQQAIFYEILHLPNLT-[Sequence A](SEQ ID NO:123); EAQKAFYEILKLPNLT-[Sequence A](SEQ ID NO:124); EQHNAFYEILHLPNLN-[Sequence A](SEQ ID NO:125); EHHHALYHILHLPNLN-[Sequence A](SEQ ID NO:126); AQQSAIFYEILHLPNLT-[Sequence A](SEQ ID

NO:127); EQQKAFYAILHLPYLT-[Sequence A](SEQ ID NO:128); DQQSAFYEILNMPNLN-[Sequence A](SEQ ID NO:129); AQQNAFYQVLNMPNLN-[Sequence A](SEQ ID NO:130); EQQKAFYEILKLPNLT-[Sequence A](SEQ ID NO:131); EQQNAFYEILSLPNLT-[Sequence A](SEQ ID NO:132); EQQNAFYEILHLPNLS-[Sequence A](SEQ ID NO:133); EQQNAFYHILHLPNLN-[Sequence A](SEQ ID NO:134); EHQNAFYEILHLPNLT-[Sequence A](SEQ ID NO:135); EQQAIFYEILHLPNLN-[Sequence A](SEQ ID NO:136); AQQAAIFYEILHLPNLT-[Sequence A](SEQ ID NO:137); wherein [Sequence A] is as defined herein.

[0123] In one particular embodiment, said binding module Sequence C is selected from the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; DQQAIFYEILHLPNLT-[Sequence A]; EAQEAIFYEILHLPNLT-[Sequence A]; DQQSAIFYEILHLPNLT-[Sequence A]; AQQAAIFYEILHLPNLT-[Sequence A]; EQQAIFYEILHLPNLT-[Sequence A]; AQQSAIFYEILHLPNLT-[Sequence A]; EQQNAFYEILNMPNLN-[Sequence A]; EAQNAFYEILHLPNLT-[Sequence A]; EQQQAIFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPYLT-[Sequence A]; ETQNAFYEILHLPNLT-[Sequence A]; EAQNAFYKILHLPNLT-[Sequence A]; EQQNAFYEILHLPNGN-[Sequence A]; EHQNAFYEILHLPNLN-[Sequence A]; and EQQNAFYEILNMPNLL-[Sequence A]; such as the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; DQQAIFYEILHLPNLT-[Sequence A]; EAQEAIFYEILHLPNLT-[Sequence A]; DQQSAIFYEILHLPNLT-[Sequence A]; AQQAAIFYEILHLPNLT-[Sequence A]; EQQAIFYEILHLPNLT-[Sequence A]; AQQSAIFYEILHLPNLT-[Sequence A]; EQQNAFYEILNMPNLN-[Sequence A]; EAQNAFYEILHLPNLT-[Sequence A]; and EQQQAIFYEILHLPNLT-[Sequence A]; such as the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; DQQAIFYEILHLPNLT-[Sequence A]; EAQEAIFYEILHLPNLT-[Sequence A]; DQQSAIFYEILHLPNLT-[Sequence A]; AQQAAIFYEILHLPNLT-[Sequence A]; EQQAIFYEILHLPNLT-[Sequence A]; AQQSAIFYEILHLPNLT-[Sequence A]; and EQQNAFYEILNMPNLN-[Sequence A]; such as the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; DQQAIFYEILHLPNLT-[Sequence A]; EAQEAIFYEILHLPNLT-[Sequence A]; DQQSAIFYEILHLPNLT-[Sequence A]; AQQAAIFYEILHLPNLT-[Sequence A]; and EQQAIFYEILHLPNLT-[Sequence A]; such as the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; DQQAIFYEILHLPNLT-[Sequence A]; EAQEAIFYEILHLPNLT-[Sequence A]; and DQQSAIFYEILHLPNLT-[Sequence A]; such as the group consisting of: EQQNAFYEILHLPNLT-[Sequence A]; EQQNAFYEILHLPNLN-[Sequence A]; and EAQEAIFYEILHLPNLT-[Sequence A]; such as the group

consisting of: EQQNAFYEILHLPNLT-[Sequence A]; and EAQEFYIEILHLPNLT-[Sequence A].

[0124] In one particular embodiment, said binding module Sequence C comprises or consists of EQQNAFYEILHLPNLT-[Sequence A]. In one particular embodiment, said binding module Sequence C comprises or consists of EQQNAFYEILHLPNLT-[Sequence A]. In one particular embodiment, said binding module Sequence C comprises or consists of EAQEFYIEILHLPNLT-[Sequence A].

[0125] In another embodiment, there is provided an Fc binding polypeptide comprising a binding module Sequence C selected from the group consisting of:

vi) (SEQ ID NO: 108)
EAQEFYIEILHLPNLT-[Sequence A]

wherein [Sequence A] is as defined herein;

[0126] vii) an amino acid sequence which has at least 85% identity to the sequence defined in vi);

viii) (SEQ ID NO: 106)
EQQNAFYEILHLPNLT-[Sequence A]

wherein [Sequence A] is as defined herein; and

[0127] ix) an amino acid sequence which has at least 85% identity to the sequence defined in viii).

[0128] Also, in a further embodiment, there is provided an Fc binding polypeptide comprising a binding module Sequence C selected from the group consisting of:

vi) (SEQ ID NO: 108)
EAQEFYIEILHLPNLT-[Sequence A]

wherein [Sequence A] is as defined herein; and

[0129] vii) an amino acid sequence which has at least 85% identity to the sequence defined in vi).

[0130] As discussed above, polypeptides comprising minor changes as compared to the above amino acid sequences without largely affecting the tertiary structure and the function thereof also fall within the scope of the present disclosure. Thus, in some embodiments, the Fc binding polypeptides comprising a binding module Sequence C as defined above may for example have a sequence which is at least 87%, such as at least 89%, such as at least 91%, such as at least 93%, such as at least 95%, such as at least 97% identical to a sequence defined by vi).

[0131] Also, in a further embodiment, there is provided an Fc binding polypeptide comprising a binding module Sequence C selected from the group consisting of:

viii) (SEQ ID NO: 106)
EQQNAFYEILHLPNLT-[Sequence A]

wherein [Sequence A] is as defined herein; and

[0132] ix) an amino acid sequence which has at least 85% identity to the sequence defined in viii).

[0133] In some embodiments, the Fc binding polypeptides comprising a binding module Sequence C as defined above may for example have a sequence which is at least 87%,

such as at least 89%, such as at least 91%, such as at least 93%, such as at least 95%, such as at least 97% identical to a sequence defined by viii).

[0134] As described in detail in the experimental section to follow, the selection of Fc binding polypeptide variants has led to the identification of a number of individual amino acid sequences and said sequences, also referred to herein as binding module Sequences C, constitute individual embodiments of sequence vi) or viii) corresponding to amino acid positions 8-54 in SEQ ID NO:1-57 and 208-264 presented in FIG. 10, as well as SEQ ID NO:269-305. Hence, the sequences of individual amino acid sequences according to Sequence C correspond to amino acid positions 8-54 in SEQ ID NO:1-57 and SEQ ID NO:208-264 presented in FIG. 10, as well as SEQ ID NO: 269-305. Hence, in one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57, 208-264 and 269-305, such as SEQ ID NO:1-56 and 208-263. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2-57, 209-264, and 269-305. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223 or the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO: SEQ ID NO:2-16 and 209-223. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218. In one embodiment, sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:7 and 214.

[0135] In one embodiment, there is provided an Fc binding polypeptide wherein sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of 2-57. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 or the group consisting of SEQ ID NO:2-14 and 16. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2-16. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54

in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11 and 12. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7 and 11. In one embodiment, sequence vi) corresponds to the sequence from position 8 to position 54 in SEQ ID NO:7.

[0136] In one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:208-264 and 269-305, such as SEQ ID NO:208-263. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209-264 and 269-305. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:208-223 or the group consisting of SEQ ID NO:209-221 and 223. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209-223. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:213, 214, 218 and 219. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:213, 214 and 218. In one embodiment, sequence viii) corresponds to the sequence from position 8 to position 54 in SEQ ID NO:214.

[0137] In some embodiments, the binding module Sequence C may form part of a polypeptide comprising an amino acid sequence further comprising Sequence 1 [S1] and/or Sequence 2 [S2]. In one embodiment, there is provided an Fc binding polypeptide as defined herein, comprising a Sequence 1 [S1], [binding module sequence C] and Sequence 2 [S2] in the following order from the N-terminus to the C-terminus

[S1]-[binding module sequence C]-[S2],
wherein [S1] or [S2] may be present or absent and [binding module sequence C] is as defined herein, and
wherein, independently from each other,
[S1] is selected from the group consisting of:

ADNKFNK; (SEQ ID NO: 138)
VDKFDK; (SEQ ID NO: 139)

-continued

VDNKFNK; (SEQ ID NO: 140)
IAAKHDK; (SEQ ID NO: 141)
IAAQHDK; (SEQ ID NO: 142)
ADNNFNK; (SEQ ID NO: 143)
IAAKFDE; (SEQ ID NO: 144)
PAAKHDK; (SEQ ID NO: 145)
ADNAFNT; (SEQ ID NO: 146)
FNK; (SEQ ID NO: 147)
ADNRFNE; (SEQ ID NO: 148)
IDSKFDE; (SEQ ID NO: 149)
ADNRFNR; (SEQ ID NO: 150)
ADNKHNK; (SEQ ID NO: 151)
ADSKFDE; (SEQ ID NO: 152)
IDAKHDE; (SEQ ID NO: 153)
QQNKFNK; (SEQ ID NO: 154)
ADNKFHK; (SEQ ID NO: 155)
KFNK; (SEQ ID NO: 156)
ADNNFNK; (SEQ ID NO: 157)
AAAKHDK; (SEQ ID NO: 158)
IDNKFNK; (SEQ ID NO: 159)
IDAKFDE; (SEQ ID NO: 160)
DNNFNK; (SEQ ID NO: 161)
ADNKFNE; (SEQ ID NO: 162)
AAQHDK; and (SEQ ID NO: 163)
AAAKFDE;

and [S2] is selected from the group consisting of:

(SEQ ID NO: 164)
QAPK;
(SEQ ID NO: 165)
QAPP;
QAP;
(SEQ ID NO: 166)
QAPR;
(SEQ ID NO: 167)
QAPE;
APK;
(SEQ ID NO: 168)
QAPG;
(SEQ ID NO: 169)
QAIK;
and
QA.

[0138] In one embodiment, S1 is selected from the group consisting of ADNKFNK; VDAKFDDK; VDNKFNK; IAAKHDK; IAAQHDK; ADNNFNK; IAAKFDE; PAAKHDK; ADNAFNT; FNK; ADNRFNE; and IDSKFDE, such as the group consisting of ADNKFNK; VDAKFDDK; VDNKFNK; IAAKHDK; IAAQHDK; ADNNFNK; IAAKFDE; and PAAKHDK, such as the group consisting of ADNKFNK; VDAKFDDK; VDNKFNK; IAAKHDK; IAAQHDK; and ADNNFNK, such as the group consisting of ADNKFNK; VDAKFDDK; VDNKFNK; IAAKHDK; and IAAQHDK, such as the group consisting of ADNKFNK; VDAKFDDK; VDNKFNK; and IAAKHDK, such as the group consisting of ADNKFNK; VDAKFDDK; and VDNKFNK. In one embodiment, S1 comprises or consists of VDAKFDDK. In one embodiment, S1 comprises or consists of ADNKFNK.

[0139] In one embodiment, S2 is selected from the group consisting of QAPK; QAPP; QAP; QAPR; and QAPE, such as the group consisting of QAPK; QAPP; QAP; and QAPE, such as the group consisting of QAPK; QAPP; QAP; and QAPR, such as the group consisting of QAPK; QAPP and QAP; such as the group consisting of QAPK; and QAPP. In one embodiment, S2 comprises or consists of QAPK. In one embodiment, S2 comprises or consists of QAPP. In one embodiment, S2 comprises or consists of QAP.

[0140] In some embodiments, the Fc binding polypeptide may be selected from polypeptides of which Sequence A forms a part according to the following. Thus, the Fc binding polypeptide may comprise an amino acid sequence selected from the group consisting of:

(SEQ ID NO: 170)
ADNKFNKQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 171)
ADNKFNKQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 172)
VDAKFDDKEAQAFYEILHLPNLT-[Sequence A]-QAPK;

-continued

(SEQ ID NO: 173)
VDNKFNKQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 174)
VDNKFNKQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 175)
PAAKHDKDQQSAFYEILHLPNLT-[Sequence A]-QAPP;
(SEQ ID NO: 176)
ADNRFNKQQNAFYEILHLPNLT-[Sequence A]-QAPR;
(SEQ ID NO: 177)
VDAKFDDKEAQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 178)
ADNAFNTEQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 179)
VDAKFDDKETQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 180)
ADNKFNKQQNAFYEILHLPNGN-[Sequence A]-QAPK;
(SEQ ID NO: 181)
ADNKFNKQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 182)
FNKEQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 183)
VDAKFDDKEQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 184)
VDAKFDDKEAQKAFYEILKLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 185)
ADNKFNKQHNNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 186)
VDNKFNKQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 187)
VDAKFDDKEQQEAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 188)
VDAKFDDKEQQKAFYEILKLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 189)
ADNKFHKEQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 190)
ADNKFNKQQNAFYHILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 191)
ADNKFNKQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 192)
ADNKHNKHHHLYHILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 193)
KFNKEQQNAFYEILHLPNLT-[Sequence A]-QAPK;
(SEQ ID NO: 194)
ADSKFDEAQQSAFYEILHLPNLT-[Sequence A]-QAPP;
and
(SEQ ID NO: 195)
IDAKFDEAQQAQAFYEILHLPNLT-[Sequence A]-QAPP,
wherein
[Sequence A] is as defined herein.

[0141] In particular embodiments, the Fc binding polypeptide comprises an amino acid sequence selected from the group consisting of:

ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK;
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 PAAKHDKDQQSAFYIELHLPNLN-[Sequence A]-QAPP;
 ADNRFNREQQNAFYIELHLPNLN-[Sequence A]-QAPR;
 VDAKFDKEAQNAFYIELHLPNLN-[Sequence A]-QAPK;
 ADNAPNTEQQNAFYIELHLPNLN-[Sequence A]-QAPK,
 and
 VDAKFDKETQNAFYIELHLPNLN-[Sequence A]-QAPK;
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

such as the group consisting of:

ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK;
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 PAAKHDKDQQSAFYIELHLPNLN-[Sequence A]-QAPP,
 and
 ADNRFNREQQNAFYIELHLPNLN-[Sequence A]-QAPR;

such as the group consisting of:

ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK;
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK,
 and
 VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

such as the group consisting of:

ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;
 and
 VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK.

In one embodiment the Fc binding polypeptide comprises or consists of ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK. In one embodiment the Fc binding polypeptide comprises or consists of ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK. In one embodiment the Fc binding polypeptide comprises or consists of VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK.

[0142] In one embodiment, the Fc binding polypeptide comprises an amino acid sequence selected from:

x)
 (SEQ ID NO: 172)
 VDAKFDKE AQEAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein; and

[0143] xi) an amino acid sequence which has at least 86% identity to the sequence defined in x).

[0144] Again, polypeptides comprising minor changes as compared to the above amino acid sequences x) without largely affecting the tertiary structure and the function thereof are also within the scope of the present disclosure. Thus, in some embodiments, the Fc binding polypeptides as defined above may for example have a sequence which is at least 87%, such as at least 89%, such as at least 91%, such as at least 93%, such as at least 94%, such as at least 96%, such as at least 98% identical to the sequence defined by x).

[0145] Sequence x) in such a polypeptide may be selected from the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:2-57. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-16 or the group consisting of SEQ ID NO:2-14 and 16. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:2-16. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12.

[0146] In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 11 and 12. In one embodiment, sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7 and 11. In another embodiment, sequence x) corresponds to the sequence SEQ ID NO:7.

[0147] In one embodiment, the Fc binding polypeptide comprises an amino acid sequence selected from:

xii)
 (SEQ ID NO: 170)
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein; and

[0148] xiii) an amino acid sequence which has at least 86% identity to the sequence defined in xii).

[0149] In yet another embodiment, the Fc binding polypeptide comprises an amino acid sequence selected from:

xiv)
 (SEQ ID NO: 171)
 ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein; and

[0150] xv) an amino acid sequence which has at least 86% identity to the sequence defined in xiv).

[0151] In some embodiments, the Fc binding polypeptides as defined in xiii) or xv) may for example have a sequence which is at least 87%, such as at least 89%, such as at least

91%, such as at least 93%, such as at least 94%, such as at least 96%, such as at least 98% identical to the sequence defined by xii) or xiv).

[0152] In one embodiment, the Fc binding polypeptide comprises an amino acid sequence selected from:

xvi)
VDNKFNKEQQNAFYIEILHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein; and

[0153] xvii) an amino acid sequence which has at least 86% identity to the sequence defined in xvi).

[0154] In some embodiments, the Fc binding polypeptides as defined in xvii) may for example have a sequence which is at least 87%, such as at least 89%, such as at least 91%, such as at least 93%, such as at least 94%, such as at least 96%, such as at least 98% identical to the sequence defined by xvi).

[0155] Sequence xvi) in such a polypeptide may be selected from the group consisting of SEQ ID NO:208-264 and 269-305, such as SEQ ID NO:208-263. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:209-264 and 269-305. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:208-223 or the group consisting of SEQ ID NO:209-221 and 223. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:209-223. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219.

[0156] In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:213, 214, 218 and 219. In one embodiment, sequence xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:213, 214 and 218. In another embodiment, sequence xvi) corresponds to the sequence SEQ ID NO:214.

[0157] In one embodiment, the Fc binding polypeptide comprises an amino acid sequence selected from

x)
VDAKFDKE AQEAFYIEILHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein;

[0158] xi) an amino acid sequence which has at least 86% identity to the sequence defined in x);

xvi)
VDNKFNKEQQNAFYIEILHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined herein; and

[0159] xvii) an amino acid sequence which has at least 86% identity to the sequence defined in xvi).

[0160] In some embodiments, the Fc binding polypeptides as defined in xi) or xvii) may for example have a sequence which is at least 87%, such as at least 89%, such as at least 91%, such as at least 93%, such as at least 94%, such as at least 96%, such as at least 98% identical to the sequence defined by x) or xvi).

[0161] In one embodiment according to the second aspect as disclosed herein, there is provided an Fc binding polypeptide wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-57 and 208-264 and 269-305, such as SEQ ID NO:1-56 and 208-263. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2-57 and 209-264 and 269-305. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223 or the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2-16 and 209-223. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219. In one embodiment, sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218. In one embodiment, sequence vi) or viii) corresponds to a sequence selected from the group consisting of SEQ ID NO:7 and 214.

[0162] A further polypeptide domain with the same functionality may be attached to the Fc binding polypeptide as defined herein, for example any Fc binding polypeptide as defined in i) to xvii), thus achieving a Fc binding polypeptide multimer. Such multimers may in particular be useful to increase the efficiency of binding the Fc region, for example to ensure that more Fc region containing polypeptides are bound to the Fc binding polypeptide multimer compared to the monomer variants thereof (in other words the Fc binding polypeptides). Thus, in a third aspect of the present disclosure, there is provided an Fc binding polypeptide multimer. Said multimer is understood to comprise at least two Fc binding polypeptides as disclosed herein as monomer units, the amino acid sequences of which may be the same or different. Thus, in one embodiment, there is provided an Fc binding polypeptide multimer wherein each monomer of the multimer comprises a Sequence A which is independently selected from any Sequence A defined herein. In another embodiment, there is provided an Fc binding polypeptide multimer wherein each monomer of the multimer comprises a Fc binding polypeptide which is independently selected from any Fc binding polypeptide defined herein.

[0163] Multimeric forms of the polypeptides may comprise a suitable number of domains, each having an Fc binding activity, and each forming a monomer within the multimer. In one embodiment, said multimer is selected from the group consisting of dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer and decamer, such as selected from the group consisting of tetramer, pentamer, hexamer, heptamer and octamer; such as the group consisting of pentamer, hexamer and heptamer. In one particular embodiment, said multimer is a hexamer.

[0164] These monomers within the multimer may have the same amino acid sequence, but alternatively, they may have different amino acid sequences. Thus the monomers may have any amino acid sequence of the Fc binding polypeptides as defined herein, for example all monomers within the

multimer may have different sequences or only a subset of the monomers in the multimer may have different amino acid sequences while other have the same sequence. In other words, the Fc binding polypeptide of the invention may form homo- or heteromultimers, for example homo- or heterodimers. In one embodiment, there is provided an Fc binding polypeptide multimer, wherein the multimer is a homomer. In another embodiment, there is provided an Fc binding polypeptide multimer, wherein multimer is a heteromer. Said heteromer may comprise at least two or at least three different Sequence A, wherein each Sequence A is as defined herein. In one embodiment, said multimer is a heteromer comprising at least two or at least three different Fc binding polypeptides, each as defined herein.

[0165] In one particular embodiment, said multimer is a homomer comprising at least three, such as at least four, such as at least five, such as at least six, such as at least seven, such as at least eight monomers selected from any one of SEQ ID NO:1-57 and 208-264 and 269-305, such as any one of SEQ ID NO:1-56 and 208-263, such as any one of SEQ ID NO:1-16 and SEQ ID NO:208-223, such as any one of SEQ ID NO:2-14, 16, 209-221 and 223, such as any one of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as any one of SEQ ID NO:2, 4, 6, 7, 12, 209, 211, 213, 214 and 219, such as any one of SEQ ID NO:6, 7, 12, 213, 214 and 219; such as any one of SEQ ID NO:6, 7, 213 and 214; such as any one of SEQ ID NO:7 and SEQ ID NO:214.

[0166] In one particular embodiment, said multimer is a homomer comprising at least three, such as at least four, such as at least five, such as at least six, such as at least seven, such as at least eight monomers selected from any one of SEQ ID NO:208-264 and 269-305, such as any one of SEQ ID NO: 208-263, such as any one of SEQ ID NO:208-223, such as any one of SEQ ID NO: 209-221 and 223, such as any one of SEQ ID NO: 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as any one of SEQ ID NO:209, 211, 213, 214 and 219, such as any one of SEQ ID NO: 213, 214 and 219; such as any one of SEQ ID NO:213 and 214; such as of SEQ ID NO:214.

[0167] In one particular embodiment, said multimer is a homomer comprising at least three, such as at least four, such as at least five, such as at least six, such as at least seven, such as at least eight monomers selected from any one of SEQ ID NO:1-57, such as any one of SEQ ID NO:1-56, such as any one of SEQ ID NO:1-16, such as any one of SEQ ID NO:2-14 and 16, such as any one of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14; such as any one of SEQ ID NO:2, 4, 6, 7 and 12, such as any one of SEQ ID NO:6, 7 and 12; such as any one of SEQ ID NO:6 and 7; such as of SEQ ID NO:7.

[0168] In one particular embodiment, said multimer is a homomer comprising six monomers of SEQ ID NO:7, in other words said multimer is a homoheptamer of SEQ ID NO:7. In another embodiment, the multimer is a homoheptamer of SEQ ID NO:6. In one embodiment, the multimer is a homoheptamer of SEQ ID NO:12. In one embodiment, said multimer is a homomer comprising six monomers of SEQ ID NO:214, in other words said multimer is a homoheptamer of SEQ ID NO:214. In another embodiment, the multimer is a homoheptamer of SEQ ID NO:213. In one embodiment, the multimer is a homoheptamer of SEQ ID 219.

[0169] In particular embodiments, the multimer comprises a sequence selected from the group consisting of SEQ ID

NO:203, 204 and 205. In one embodiment, said multimer comprises the sequence SEQ ID NO: 204.

[0170] It will be appreciated that monomers in the multimers discussed herein, which multimers comprise Sequence A or comprise Fc binding polypeptides as disclosed herein, may be directly coupled to each other or spaced apart by linker sequences. The skilled person appreciates that the presence or absence of linker sequences may be different between different monomers moieties in a multimer, and if linkers are present, the sequence of each individual linker may be the same or different.

[0171] As the skilled person understands, the construction of a multimer, for example as a fusion protein, often involves use of linkers between the monomer moieties to be fused. The skilled person is aware of different kinds of linkers with different properties, such as flexible amino acid linkers, rigid amino acid linkers and cleavable amino acid linkers. Linkers may be used in order to for example increase stability or improve folding of fusion proteins, to increase expression or to improve activity, affinity and/or binding capacity. Thus, in one embodiment, the Fc binding polypeptide multimer as defined herein further comprises at least one linker. In one embodiment, a linker is present between each monomer within the multimer. The linker may for example be selected from the group consisting of flexible amino acid linkers, rigid amino acid linkers and cleavable amino acid linkers. Alternatively, the linker may be a non-peptidic linker. Thus, the Fc binding polypeptides disclosed herein may be linked to each directly by peptide bonds between the C-terminal and N-terminal ends of the polypeptides. Alternatively, two or more monomers, in other words monomer units or moieties, within the multimer can be linked by elements comprising oligomeric or polymeric species, such as elements comprising up to 15 or 30 amino acids, such as 1-5, 1-10 or 5-10 amino acids. The nature of such a link should preferably not destabilize the spatial conformation of the protein units, that is of the Fc binding polypeptide monomers within the multimer. This can e.g. be achieved by avoiding the presence of proline in the linkers. Furthermore, said linkers should preferably also be sufficiently stable in alkaline environments not to impair the properties of the mutated protein units. For this purpose, it is advantageous if the linkers do not contain asparagine. It can additionally be advantageous if the linker do not contain glutamine. The multimer may further at the N-terminal end comprise a plurality of amino acid residues originating from the cloning process or constituting a residue from a cleaved off signaling sequence. The number of additional amino acid residues may e.g. be 15 or less, such as 10 or less or 5 or less.

[0172] The skilled person will understand that various modifications and/or additions can be made to an Fc binding polypeptide or to an Fc binding polypeptide multimer according to any aspect disclosed herein in order to tailor the polypeptide or multimer to a specific application without departing from the scope of the present disclosure.

[0173] For example, in one embodiment there is provided an Fc binding polypeptide or an Fc binding polypeptide multimer as described herein, which polypeptide has been extended by and/or comprises additional amino acids at the C terminus and/or N terminus. Such a polypeptide or multimer should be understood as a polypeptide or multimer having one or more additional amino acid residues at the very first and/or the very last position in the polypeptide chain, i.e. at the N- and/or C-terminus of the polypeptide or

multimer. For example said additional amino acid residues may be at the N- and/or C-terminus any one of Sequence A according to i), ii) or iii), binding module Sequence C according to vi), vii), viii) or ix) or sequence x), xi), xii), xiii), xiv), xv) xvi) or xvii). Thus, the Fc binding polypeptides as defined herein may comprise any suitable number of additional amino acid residues, for example one, two, three, four, five, six, seven, eight, nine, ten or more additional amino acid residues. Said amino acid residues may individually or collectively improve production, purification, stabilization in vitro or coupling of the polypeptide to substrates of interest, for example to a solid support, such as a solid support described in connection to the aspect of a separation matrix.

[0174] Said additional amino residues may be coupled to the Fc binding polypeptide or multimer by means of chemical conjugation (using known organic chemistry methods) or by any other means, such as expression of the Fc binding polypeptide or multimer as a fusion protein or joined in any other fashion, either directly or via a linker, for example an amino acid linker as described above.

[0175] In some embodiments, the Fc binding polypeptides and/or multimer, as disclosed above, further comprises at the C-terminal or N-terminal end one or more coupling elements, selected from the group consisting of a cysteine residue, a plurality of lysine residues and a plurality of histidine residues. The coupling element may e.g. be a single cysteine at the C-terminal end. The coupling element(s) may be directly linked to the C- or N-terminal end, or it/they may be linked via a linker comprising up to 15 amino acids, such as 1-5, 1-10 or 5-10 amino acids. This stretch should preferably also be sufficiently stable in alkaline environments not to impair the properties of the mutated protein. For this purpose, it is advantageous if the stretch does not contain asparagine. It can additionally be advantageous if the stretch does not contain glutamine. An advantage of having a C-terminal cysteine is that endpoint coupling of the protein can be achieved through reaction of the cysteine thiol with an electrophilic group on a support. This provides excellent mobility of the coupled protein which is important for the binding capacity.

[0176] In a fourth aspect of the present disclosure, there is provided a polynucleotide encoding an Fc binding polypeptide or an Fc binding polypeptide multimer as described herein; an expression vector comprising said polynucleotide; and a host cell comprising said expression vector.

[0177] Also encompassed by this disclosure is a method of producing the Fc binding polypeptide or the Fc binding polypeptide multimer as described herein, comprising culturing said host cell under conditions permissive of expression of said polypeptide or multimer from its expression vector, and isolating the polypeptide or multimer.

[0178] The Fc binding polypeptide or multimer as disclosed herein may alternatively be produced by non-biological peptide synthesis using amino acids and/or amino acid derivatives having protected reactive side-chains, the non-biological peptide synthesis comprising

[0179] step-wise coupling of the amino acids and/or the amino acid derivatives to form a polypeptide according to the second aspect or a multimer according to the third aspect having protected reactive side-chains,

[0180] removal of the protecting groups from the reactive side-chains of the polypeptide or multimer, and

[0181] folding of the polypeptide or multimer in aqueous solution.

[0182] The terms “Fc binding” and “binding affinity for Fc” as used in this disclosure refer to a property of a polypeptide which may be tested for example by ELISA or the use of surface plasmon resonance (SPR) technology.

[0183] Fc binding affinity may be tested in an experiment in which Fc, or a fragment thereof, is immobilized on a sensor chip of the surface plasmon resonance (SPR) instrument, and the sample containing the polypeptide to be tested is passed over the chip. Alternatively, the polypeptide to be tested is immobilized on a sensor chip of the instrument, and a sample containing Fc, or a fragment thereof, is passed over the chip. The skilled person may then interpret the results obtained by such experiments to establish at least a qualitative measure of the binding affinity of the polypeptide for Fc. If a quantitative measure is desired, for example to determine a K_D value for the interaction, surface plasmon resonance methods may also be used. Binding values may for example be defined in a Biacore (Cytiva) or ProteOn XPR 36 (Bio-Rad) instrument. Fc is suitably immobilized on a sensor chip of the instrument, and samples of the polypeptide whose affinity is to be determined are prepared by serial dilution and injected in random order. K_D values may then be calculated from the results using for example the 1:1 Langmuir binding model of the BIAevaluation 4.1 software, or other suitable software, provided by the instrument manufacturer.

[0184] As explained above the Fc binding polypeptide or Fc binding polypeptide multimer of the present disclosure are able to bind to the Fc region of either IgG1, IgG2 or IgG4 with said affinity.

[0185] Thus in one embodiment, there is provided an Fc binding polypeptide or Fc binding polypeptide multimer as disclosed herein, which is capable of binding to Fc such as that the K_D value of the interaction is at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M, such as at most 1×10^{-11} M.

[0186] It will be appreciated that the Fc binding polypeptides and/or the Fc binding polypeptide multimers exhibit a similar K_D value of the interaction with Fc as SEQ ID NO:58 and/or SEQ ID NO:59. In particular, the K_D value of the interaction with Fc of the Fc binding polypeptide or Fc binding polypeptide multimer as disclosed herein may differ with one order of magnitude from the K_D value of the interaction of SEQ ID NO:58 and/or SEQ ID NO:59 with Fc. For example, if the K_D value of the interaction of SEQ ID NO:58 and/or SEQ ID NO:59 with a particular Fc is 9×10^{-10} M in an assay, then the K_D value of the interaction with Fc of the Fc binding polypeptide or Fc binding polypeptide multimer may be in the range of from 9×10^{-9} M to 9×10^{-11} M (one order of magnitude), in that assay. The skilled person will appreciate that it may be useful to compare K_D values obtained using the same assay and while some variation may occur between different assays, intra-assay comparisons generally demonstrate the same trends independent of the assay employed. Assays may for example differ depending on if it is the Fc containing polypeptide or the Fc binding polypeptide which is immobilized on a chip. In one embodiment, said K_D value of the interaction with Fc is measured in an assay wherein the Fc binding polypeptide is immobilized on the chip. In a different embodiment, the

K_D value of the interaction with Fc is measured in an assay wherein the Fc containing polypeptide is immobilized on the chip.

[0187] In another embodiment, there is provided an Fc binding polypeptide or Fc binding polypeptide multimer as disclosed herein, which is capable of binding to Fc such as that the K_D value of the interaction is at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 90% of the K_D value of the interaction of SEQ ID NO:58 with Fc. In one embodiment, said Fc binding polypeptide or Fc binding polypeptide multimer is capable of binding to Fc such as that the K_D value of the interaction is at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 90% of the K_D value of the interaction of SEQ ID NO:59 with Fc. In one embodiment, said Fc binding polypeptide or Fc binding polypeptide multimer is capable of binding to Fc such as that the K_D value of the interaction is at least 50%, such as at least 60%, such as at least 70%, such as at least 80%, such as at least 90% of the K_D value of the interaction of SEQ ID NO:58 and SEQ ID NO:59 with Fc.

[0188] Similarly, the terms “VH3 binding” and “binding affinity for VH3” as used in this disclosure refer to a property of a polypeptide which may be tested for example by ELISA or the use of surface plasmon resonance (SPR) technology. VH3 binding affinity may be tested in an experiment in which VH3, or a fragment thereof, is immobilized on a sensor chip of the surface plasmon resonance (SPR) instrument, and the sample containing the polypeptide to be tested is passed over the chip. Alternatively, the polypeptide to be tested (in this case the Fc binding polypeptide) may be immobilized on a sensor chip of the instrument, and a sample containing VH3, or a fragment thereof, is passed over the chip. The skilled person appreciates that essentially an absence of binding is to be detected if the binding to VH3 is abolished or significantly reduced. The skilled person may then interpret the results obtained by such experiments to establish at least a qualitative measure of the binding affinity of the polypeptide for VH3. If a quantitative measure is desired, for example to determine a K_D value for the interaction, surface plasmon resonance methods may also be used in an analogous manner to what is explained above of Fc binding.

[0189] As explained in connection with the second aspect of the present disclosure, the presently identified Fc binding polypeptides and Fc binding polypeptide multimers as defined herein have significantly reduced or abolished binding affinity for VH3 of trastuzumab. The skilled person will appreciate that the present Fc binding polypeptides and Fc binding polypeptide multimers do not significantly interact with VH3, and thus their capacity to bind to VH3 is low.

[0190] In one embodiment, there is provided an Fc binding polypeptide or Fc binding polypeptide multimer as disclosed herein, which is not capable of binding to VH3 with a K_D value of the interaction of less than 1×10^{-4} M, such as less than 1×10^{-3} M.

[0191] In addition, the Fc binding polypeptide displays improved alkali stability. In embodiments where the Fc binding polypeptide is used for separation or isolation of e.g. antibodies/immunoglobulins having an Fc region, high alkali stability will allow for use of highly alkaline conditions during cleaning, essential for long-term repeated use in a bioprocess separation setting.

[0192] In one embodiment, there is provided an Fc binding polypeptide or Fc binding polypeptide multimer as disclosed herein which has an alkali stability of at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 89%, such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99% of the alkali stability of SEQ ID NO:58 after incubation in 0.5 M NaOH. In one embodiment, said Fc binding polypeptide or Fc binding polypeptide multimer has an alkali stability of at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 89%, such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99% of the alkali stability of SEQ ID NO:59 after incubation in 0.5 M NaOH. In one embodiment, said Fc binding polypeptide or Fc binding polypeptide multimer has an alkali stability of at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 89%, such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99% of the alkali stability of SEQ ID NO:58 and SEQ ID NO:59 after incubation in 0.5 M NaOH. In one embodiment, said incubation is at least for 12 hours, such as at least for 24 hours or more. Said incubation may be for example at $22 \pm 2^\circ$ C.

[0193] In another aspect, the present invention discloses a separation matrix, comprising an Fc binding polypeptide according to any embodiment disclosed above coupled to a solid support. Such a matrix is useful for separation of immunoglobulins or other Fc-containing proteins from e.g. a sample. Since the Fc binding polypeptide has binding affinity to Fc but low or no binding affinity to VH3, the separation matrix may allow for efficient separation of Fc containing proteins from any VH3 containing proteins or impurities. As described above, the Fc binding polypeptide displays improved alkali stability which will allow for use of highly alkaline conditions during cleaning of the separation matrix, essential for long-term repeated use in a bioprocess separation setting. The alkali stability of the matrix can be assessed by measuring the immunoglobulin-binding capacity, or Fc binding capacity, typically using polyclonal human IgG, before and after incubation in alkaline solutions at a specified temperature, e.g. $22 \pm 2^\circ$ C. The incubation can e.g. be performed in 0.5 M or 1.0 M NaOH for a number of 15 min cycles, such as 100, 200 or 300 cycles, corresponding to a total incubation time of 25, 50 or 75 h. The IgG capacity of the matrix after 96-100 15 min incubation cycles or a total incubation time of 24 or 25 h in 0.5 M NaOH at $22 \pm 2^\circ$ C. can be at least 80, such as at least 85, at least 90 or at least 95% of the IgG capacity before the incubation. The capacity of the matrix after a total incubation time of 24 h in 1.0 M NaOH at $22 \pm 2^\circ$ C. can be at least 70, such as at least 80 or at least 90% of the IgG capacity before the incubation. In one embodiment, the separation matrix comprises an Fc binding polypeptide multimer, wherein each monomer comprises an Fc binding polypeptide as defined herein. The multimer, which is a homo-multimer or a hetero-multimer, may be a dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer or decamer. For example, the separation matrix may comprise a hexamer of an Fc binding polypeptide coupled to the solid support. Each monomer of the Fc binding polypeptide multimer may comprise a Sequence A which is independently selected from any Sequence A defined herein. In particular, it may comprise a sequence selected from the group consisting of SEQ ID NO:1-57, 208-264 and 269-305, such as SEQ ID

NO:1-56 and 208-263, such as SEQ ID NO:1-16 and SEQ ID NO:208-223, such as SEQ ID NO:2-14, 16, 209-221 and 223, such as SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as SEQ ID NO:2, 4, 6, 7, 12, 209, 211, 213, 214 and 219, such as SEQ ID NO:6, 7, 12, 213, 214, and 219; such as SEQ ID NO:6, 7, 213 and 214; such as SEQ ID NO:7 and SEQ ID NO:214. It may comprise a sequence selected from the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56. In particular, it may comprise a sequence selected from the group consisting of SEQ ID NO:1-16; such as from the group consisting of SEQ ID NO:2-14 and 16, such as from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14; such as from the group consisting of SEQ ID NO:2, 4, 6, 7 and 12, such as from the group consisting of SEQ ID NO:6, 7 and 12; such as from the group consisting of SEQ ID NO:6 and 7, such as SEQ ID NO:7. In one embodiment, said separation matrix comprises an Fc binding polypeptide multimer, optionally a hexamer, wherein each monomer of said multimer comprises a sequence corresponding to SEQ ID NO:7. It may comprise a sequence selected from the group consisting of SEQ ID NO:208-264 and 269-305, such as from the group consisting of SEQ ID NO:208-263, such as from the group consisting of SEQ ID NO:208-223, such as from the group consisting of SEQ ID NO:209-221 and 223, such as from the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as from the group consisting of SEQ ID NO:209, 211, 213, 214 and 219, such as from the group consisting of SEQ ID NO:213, 214 and 219; such as from the group consisting of SEQ ID NO:213 and 214. In one embodiment, said separation matrix comprises an Fc binding polypeptide multimer, optionally a hexamer, wherein each monomer of said multimer comprises a sequence corresponding to SEQ ID NO:214.

[0194] As the skilled person will understand, the expressed Fc binding polypeptide, in monomeric or multimeric form, should be purified to an appropriate extent before being immobilized to a support. Such purification methods are well known in the field, and the immobilization of protein-based ligands to supports is easily carried out using standard methods. Suitable methods and supports will be discussed below in more detail and are disclosed e.g. in WO16079033 which is incorporated herein by reference.

[0195] The solid support of the matrix according to the invention can be of any suitable well-known kind. As a non-limiting example, a conventional affinity separation matrix is often of organic nature and based on polymers that expose a hydrophilic surface to the aqueous media used, i.e. expose hydroxy (—OH), carboxy (—COOH), carboxamido (—CONH₂, possibly in N-substituted forms), amino (—NH₂, possibly in substituted form), oligo- or polyethylenoxy groups on their external and, if present, also on internal surfaces. The solid support can suitably be porous. The porosity can be expressed as a *K_{av}* or *K_d* value (the fraction of the pore volume available to a probe molecule of a particular size) measured by inverse size exclusion chromatography, e.g. according to the methods described in *Gel Filtration Principles and Methods*, Pharmacia LKB Biotechnology 1991, pp 6-13. By definition, both *K_d* and *K_{av}* values always lie within the range 0-1. The *K_{av}* value can advantageously be 0.6-0.95, e.g. 0.7-0.90 or 0.6-0.8, as measured with dextran of Mw 110 kDa as a probe molecule. An advantage of this is that the support has a large fraction

of pores able to accommodate both the polypeptides/multimers of the invention and immunoglobulins binding to the polypeptides/multimers and to provide mass transport of the immunoglobulins to and from the binding sites.

[0196] The polypeptides or multimers may be attached to the support via conventional coupling techniques utilizing e.g. thiol, amino and/or carboxy groups present in the ligand. Bisepoxides, epichlorohydrin, CNBr, N-hydroxysuccinimide (NHS) etc. are well-known coupling reagents. Between the support and the polypeptide/multimer, a molecule known as a spacer can be introduced, which improves the availability of the polypeptide/multimer and facilitates the chemical coupling of the polypeptide/multimer to the support. Depending on the nature of the polypeptide/multimer and the coupling conditions, the coupling may be a multipoint coupling (e.g. via a plurality of lysines) or a single point coupling (e.g. via a single cysteine).

[0197] Alternatively, the polypeptide/multimer may be attached to the support by non-covalent bonding, such as physical adsorption or biospecific adsorption.

[0198] In some embodiments the matrix comprises 5-25, such as 5-20 mg/ml, 5-15 mg/ml, 5-11 mg/ml or 6-11 mg/ml of the polypeptide or multimer coupled to the support. The amount of coupled polypeptide/multimer can be controlled by the concentration of polypeptide/multimer used in the coupling process, by the activation and coupling conditions used and/or by the pore structure of the support used. As a general rule the absolute binding capacity of the matrix increases with the amount of coupled polypeptide/multimer, at least up to a point where the pores become significantly constricted by the coupled polypeptide/multimer. The relative binding capacity per mg coupled polypeptide/multimer will decrease at high coupling levels, resulting in a cost-benefit optimum within the ranges specified above.

[0199] In certain embodiments the polypeptides or multimers are coupled to the support via thioether bonds. Methods for performing such coupling are well-known in this field and easily performed by the skilled person in this field using standard techniques and equipment. Thioether bonds are flexible and stable and generally suited for use in affinity chromatography. In particular when the thioether bond is via a terminal or near-terminal cysteine residue on the polypeptide or multimer, the mobility of the coupled polypeptide/multimer is enhanced which provides improved binding capacity and binding kinetics. In some embodiments the polypeptide/multimer is coupled via a C-terminal cysteine provided on the protein as described above. This allows for efficient coupling of the cysteine thiol to electrophilic groups, e.g. epoxide groups, halohydrin groups etc. on a support, resulting in a thioether bridge coupling.

[0200] In certain embodiments the solid support comprises a polyhydroxy polymer, such as a polysaccharide. Examples of polysaccharides include e.g. dextran, starch, cellulose, pullulan, agar, agarose etc. Polysaccharides are inherently hydrophilic with low degrees of nonspecific interactions, they provide a high content of reactive (activatable) hydroxyl groups and they are generally stable towards alkaline cleaning solutions used in bioprocessing. In some embodiments the support comprises agar or agarose. The supports used in the present invention can easily be prepared according to standard methods, such as inverse suspension gelation (S Hjerten: *Biochim Biophys Acta* 79(2), 393-398 (1964)). Alternatively, the base matrices are commercially available products, such as crosslinked agarose beads sold

under the name of SEPHAROSE™ FF (Cytiva). In an embodiment, which is especially advantageous for large-scale separations, the support has been adapted to increase its rigidity using the methods described in U.S. Pat. No. 6,602,990 or U.S. Pat. No. 7,396,467, which are hereby incorporated by reference in their entirety, and hence renders the matrix more suitable for high flow rates.

[0201] In certain embodiments the support, such as a polysaccharide or agarose support, is crosslinked, such as with hydroxyalkyl ether crosslinks. Crosslinker reagents producing such crosslinks can be e.g. epihalohydrins like epichlorohydrin, diepoxides like butanediol diglycidyl ether, allylating reagents like allyl halides or allyl glycidyl ether. Crosslinking is beneficial for the rigidity of the support and improves the chemical stability. Hydroxyalkyl ether crosslinks are alkali stable and do not cause significant nonspecific adsorption.

[0202] Alternatively, the solid support is based on synthetic polymers, such as polyvinyl alcohol, polyhydroxyalkyl acrylates, polyhydroxyalkyl methacrylates, polyacrylamides, polymethacrylamides etc. In case of hydrophobic polymers, such as matrices based on divinyl and monovinyl-substituted benzenes, the surface of the matrix is often hydrophilised to expose hydrophilic groups as defined above to a surrounding aqueous liquid. Such polymers are easily produced according to standard methods, see e.g. "Styrene based polymer supports developed by suspension polymerization" (R Arshady: *Chimica e L'Industria* 70(9), 70-75 (1988)). As an alternative, a commercially available product, such as SOURCE™ (Cytiva) may be used. Alternatively, the solid support according to the invention comprises a support of inorganic nature, e.g. silica, zirconium oxide etc.

[0203] In one embodiment, the solid support has the form of a porous monolith. In an alternative embodiment, the support is in beaded or particle form that can be porous or non-porous. Solid supports in beaded or particle form can be used as a packed bed or in a suspended form. Suspended forms include those known as expanded beds and pure suspensions, in which the particles or beads are free to move. In case of monoliths, packed bed and expanded beds, the separation procedure commonly follows conventional chromatography with a concentration gradient. In case of pure suspension, batch-wise mode will be used.

[0204] In one embodiment, the separation matrix is a convection-based chromatography matrix. Such convection-based chromatography matrix may comprise a fibrous support or fibrous substrate. Said fibrous support may be based on electrospun polymeric fibers or cellulose fibers, optionally non-woven fibers. The fibrous support may thus be a fibrous non-woven polymer matrix. The fibers comprised in said fibrous support have a cross-sectional diameter of 10-1000 nm, such as 200-800 nm, 200-400 nm or 300-400 nm. Such a fibrous support can be found in a HiTrap Fibro™ unit (Cytiva). Alternative fibrous supports are disclosed in e.g. WO2019/137869 and WO2018/011600.

[0205] In yet another embodiment, the solid support is in another form such as a surface, a chip, capillaries, or a filter (e.g. a membrane or a depth filter matrix).

[0206] In another aspect, the present invention provides a method of isolating an immunoglobulin, wherein a separation matrix as disclosed above is used. In some embodiments, the method comprises a) contacting a liquid sample comprising an immunoglobulin with a separation matrix as disclosed above. The method may furthermore comprise b)

washing said separation matrix with a washing liquid, c) eluting the immunoglobulin from the separation matrix with an elution liquid, and optionally d) cleaning the separation matrix with a cleaning liquid. The cleaning liquid can alternatively be called a cleaning-in-place (CIP) liquid. The contact (incubation) time may be at least 10 min.

[0207] The method may also comprise steps of, before step a), providing an affinity separation matrix according to any of the embodiments described above and providing a solution comprising an immunoglobulin and at least one other substance as a liquid sample and of, after step c), recovering the eluate and optionally subjecting the eluate to further separation steps, e.g. by anion or cation exchange chromatography, multimodal chromatography and/or hydrophobic interaction chromatography. Suitable compositions of the liquid sample, the washing liquid and the elution liquid, as well as the general conditions for performing the separation are well known in the art of affinity chromatography and in particular in the art of Protein A chromatography. The liquid sample comprising an Fc-containing protein and at least one other substance may comprise host cell proteins (HCP), such as CHO cell, *E. Coli* or yeast proteins. Contents of CHO cell and *E. Coli* proteins can conveniently be determined by immunoassays directed towards these proteins, e.g. the CHO HCP or *E. Coli* HCP ELISA kits from Cygnus Technologies. The host cell proteins or CHO cell/*E. Coli* proteins may be desorbed during step b). The elution may be performed by using any suitable solution used for elution from Protein A media. This can e.g. be a solution or buffer with pH 5 or lower, such as pH 2.5-5 or 3-5. It can also in some cases be a solution or buffer with pH 11 or higher, such as pH 11-14 or pH 11-13. In some embodiments the elution buffer or the elution buffer gradient comprises at least one mono- di- or trifunctional carboxylic acid or salt of such a carboxylic acid. In certain embodiments the elution buffer or the elution buffer gradient comprises at least one anion species selected from the group consisting of acetate, citrate, glycine, succinate, phosphate, and formate.

[0208] In some embodiments, the cleaning liquid is alkaline, such as with a pH of 13-14. Such solutions provide efficient cleaning of the matrix, in particular at the upper end of the interval.

[0209] In certain embodiments, the cleaning liquid comprises 0.1-2.0 M NaOH or KOH, such as 0.5-2.0 or 0.5-1.0 M NaOH or KOH. These are efficient cleaning solutions, and in particular so when the NaOH or KOH concentration is above 0.1 M or at least 0.5 M. The high stability of the polypeptides of the invention enables the use of such strongly alkaline solutions.

[0210] The method may also include a step of sanitizing the matrix with a sanitization liquid, which may e.g. comprise a peroxide, such as hydrogen peroxide and/or a peracid, such as peracetic acid or performic acid.

[0211] In some embodiments, steps a)-d) are repeated at least 10 times, such as at least 50 times, 50-200, 50-300 or 50-500 times. This is important for the process economy in that the matrix can be re-used many times. Steps a)-c) can also be repeated at least 10 times, such as at least 50 times, 50-200, 50-300 or 50-500 times, with step d) being performed after a plurality of instances of step c), such that step d) is performed at least 10 times, such as at least 50 times. Step d) can e.g. be performed every second to twentieth instance of step c).

[02112] The terms an “Fc binding polypeptide” and “Fc binding protein” mean a polypeptide or protein respectively, capable of binding to the crystallizable part (Fc) of an antibody and includes e.g. Protein A, or any fragment or fusion protein thereof that has maintained said binding property.

[02113] The term “linker” herein means an element linking two polypeptide units, monomers or domains to each other in a multimer.

[02114] The term “spacer” herein means an element connecting a polypeptide or a polypeptide multimer to a support.

[02115] The term “VH binding” refers to binding to the variable heavy (VH) chain of an antibody, or the so-called Fab fragment. The term “Fab fragment” and “VH chain” may be used interchangeably herein, and subsequently the term “Fab binding” is used interchangeably with the term “VH binding”.

[02116] While the invention has been described with reference to various exemplary aspects and embodiments, it will be understood by those skilled in the art that various changes may be made and equivalents may be substituted for elements thereof without departing from the scope of the invention. In addition, many modifications may be made to adapt a particular situation or molecule to the teachings of the invention without departing from the essential scope thereof. Therefore, it is intended that the invention not be limited to any particular embodiment contemplated, but that the invention will include all embodiments falling within the scope of the appended claims.

BRIEF DESCRIPTION OF THE FIGURES

[02117] FIG. 1a is an overview of Biacore set-up for screening of library of Fc binding polypeptide variants.

[02118] FIG. 1b shows sensorgrams of Fab and Fc-binding of albumin binding domain-linked domain Z (SEQ ID NO:59-ABD) and an inventive Fc binding polypeptide linked to ABD (candidate-ABD).

[02119] FIG. 2 is an example plot of alkali stability for tested Fc binding polypeptide variants in comparison to SEQ ID NO:58.

[0220] FIG. 3 shows Biacore binding curves (reference-subtracted) following injection of 1 g/L trastuzumab Fab.

[0221] FIG. 4 shows Biacore binding curves (reference-subtracted) following injection of 40 mg/L Fc.

[0222] FIG. 5 shows Biacore binding curves (reference-subtracted) following injection of 50 mg/L trastuzumab.

[0223] FIG. 6 shows Biacore binding curves (reference-subtracted) following injection of 0.5 g/L control antibody which does not contain the VH3 of trastuzumab

[0224] FIG. 7 shows Biacore binding capacity as a function of ligand coupling level for four different target proteins.

[0225] FIG. 8 is a plot of trastuzumab response as a function of number of NaOH injections for hexameric Fc binding polypeptide ligands.

[0226] FIG. 9 is a Plot of trastuzumab response as a function of number of NaOH injections for monomeric Fc binding polypeptide ligands.

[0227] FIG. 10 is a listing of the amino acid sequences of examples of Fc binding polypeptides of the present disclosure (SEQ ID NO:1-57 and SEQ ID NO:208-264) and hexamer variants thereof (SEQ ID NO:203-205), as well amino acid sequences of domain A, B, C, D and E of

Staphylococcal protein A and derivatives thereof (SEQ ID NO:58-64). In the Fc binding polypeptides of the present disclosure, the deduced Sequence A extend from residue 24 to residue 54 in each sequence. The amino acid sequences Sequence B extend from residue 8-23 and binding module Sequence C predicted to constitute the complete three-helix bundle within each of these Fc binding polypeptides extend from residue 8 to residue 54.

[0228] FIG. 11 shows Biacore binding curves (reference-subtracted) for full antibodies trastuzumab and adalimumab. 11a) Domain B_{wt} (SEQ ID NO 61) and trastuzumab; 11b) Domain B_{wt} (SEQ ID NO 61) and adalimumab; 11c) mutated domain B (SEQ ID 275) and trastuzumab; 11d) mutated domain B (SEQ ID 275) and adalimumab.

[0229] FIG. 12 shows Biacore binding curves (reference-subtracted) for Fc fragments from adalimumab and for Fab fragments from trastuzumab. 12a) Domain B_{wt} (SEQ ID NO 61) and Fc fragment; 12b) Domain B_{wt} (SEQ ID NO 61) and Fab fragment; 12c) mutated domain B (SEQ ID 275) and Fc fragment; 12d) mutated domain B (SEQ ID 275) and Fab fragment.

EXAMPLES

Summary of Examples 1 to 3

[0230] This project aimed for providing Fc binding polypeptides with reduced or abolished ability to bind VH3 compared to SEQ ID NO:59 or 58. Three positions (33, 40 and 51) in SEQ ID NO:58 were identified as key positions to knock out VH binding and randomized in a Fc binding polypeptide variant library. The library was screened for the following parameters: no VH binding, high Fc binding and high alkaline stability. The set-up for analysis of the members of the library is outlined in FIG. 2.

[0231] From the screening, a subset of ligands were chosen for further studies on column. The further studies included multimerization of the ligands to hexamers and expression of said hexamers in *E. coli*. On these hexameric Fc binding polypeptide ligands, a primary characterization was performed, wherein the ligands were evaluated for dynamic binding capacity (DBC), pH elution, no binding of Fab and alkaline stability.

[0232] Trastuzumab used in the Examples below was produced/purified from in-house production cell line (CHO-K1 background).

Example 1

Biacore Evaluation of Fc Binding Polypeptide VH Knock-Out Library and Sequencing of Candidates

[0233] The impact of mutations at key amino acid positions was investigated to identify positions that enabled reduced binding to the VH3 region. The three positions with the highest impact on VH3 (in a Fab fragment) binding were identified as amino acid positions 33/40/51 of SEQ ID NO:58.

[0234] To identify optimal Fc binding polypeptide candidates—a small library with a diversity of amino acids at the three identified positions (33/40/51) were designed and constructed (Table 1). The total diversity of the library was 648 possible variants.

TABLE 1

Summary of library randomization. Library randomization		
Position 33	Position 40	Position 51
P	H	L
T	Q	I
A	N	V
H	K	R
Q	D	S
N	E	G
K	R	
D	S	
E	G	
R		
S		
G		

Generation of Candidate Lysates and Colony Storage Plates

[0235] For evaluation of the mutant library all the Fc binding polypeptide variants were constructed as fusion proteins with C-terminal fusion to the Albumin Binding Domain (ABD) (SEQ ID NO:201). This allowed for efficient screening of variants from raw *E. coli* lysates via use of capture to immobilized RSA (Rat Serum Albumin; high affinity for ABD). In short, the generation of protein was achieved by transformation of KCM competent *E. coli* top10 with the library in plasmid form. Colonies representing individual variants were generated by plating the transformation product on agar plates, followed by incubation at 37° C. overnight for individual colonies to appear.

[0236] Individual colonies were transferred from the agar plates to 96-well Deep Well Plates (96DWP, 10 in total) for culture and protein production. Pipette tips used to pipette up/down 10-times to ensure bacterial dispersion and then 3-5 µl from each well was transferred to a marked agar plate in marked 96-grid structure for later retrieval of interesting candidates for sequencing, re-culture or re-cloning (colony storage plate).

[0237] The 96DWPs were incubated in an Infors HT shaking incubator for 24 hours at 27° C. and 300 rpm shaking. Finished cultures were frozen at -20° C.

[0238] To generate crude protein containing lysates; frozen 96DWPs were incubated in 80° C. water bath for 2 hours followed by pelleting of cell debris by centrifugation at 3400xg for 40 min. These crude lysate plates with the raw protein were then kept in cooler until further analysis.

[0239] Raw protein lysates were generated by heat treatment (2 hours at 80° C.) in Biacore running buffer. Variants were then screened for expression level (capture to immobilized RSA), binding to Fab prepared from trastuzumab (SEQ ID NO:198 and 199; Fab fragment of trastuzumab, comprising VH3) and binding to Fc (SEQ ID NO:200) from trastuzumab. FIG. 1 shows an overview of a Biacore set-up.

Pre-Testing of Antibody Fragment Material

[0240] An analysis of the antibody fragment material Fab (comprising SEQ ID NO:198 and 199) and Fc (SEQ ID NO:200) was performed using albumin binding domain-linked domain Z (Z-ABD) (SEQ ID NO:202).

Materials and Methods

[0241] Z-ABD (SEQ ID NO:202), Fab fragment of trastuzumab (comprising SEQ ID NO:198 and 199) and Fc (SEQ ID NO:200) were obtained from preparations generated in-house. Biacore CM5 chips; RSA; Biacore EHS coupling kit; Biacore 8K+ instrument (all from Cytiva, Uppsala, Sweden) were used according to manufacturer's instructions. RSA was immobilized on all channels of FC2 on a CM5 chip (50 ug/ml RSA pH5 in acetate buffer). The immobilized amount was ranging between 13500-14000 Ru. FC1 was blocked with ethanolamine.

[0242] To test binding to available Fab and Fc, an analysis run using Z-ABD was performed. Z-ABD was diluted 1:1 with PBS-P+ (phosphate buffered saline with KCl and Tween20), Fab fragment of trastuzumab (also referred to as Fab below) was diluted to 1.3 g/L in pBS-P+ (Prep 1), and Fc was diluted to 18 µg/ml in PBS-P+. Z was captured on FC2, followed by injection of Fab and then Fc. The resulting sensorgram is shown in FIG. 1B (a).

Biacore Analysis of Fc Binding Polypeptide VH Knock-Out Candidates

Materials and Methods

[0243] CM5 sensor chip with RSA was immobilized on all channels of FC2 (50 ug/ml RSA pH5 in acetate buffer). Biacore 8K+ instrument (Cytiva) was used for analysis, *E. coli* lysates of the VH knock-out candidate Fc binding polypeptides were prepared as described above. Fab fragment of trastuzumab as used at the concentration of 1.3 g/L in pBS-P+ and the Fc preparation used was 18 µg/ml in PBS-P+).

Two methods for the Biacore analysis were used as described below:

Biacore Method 1:

[0244] Biacore method 1 setup was as follows:

- [0245] Running buffer: PBS-P+
- [0246] Flow-rate: 5 µl/min during analysis; 30 µl/min during regeneration
- [0247] Injections: 60 s capture of candidate Fc binding polypeptide fused to ABD (SEQ ID NO:201) (1+2 dilution in PBS-P+), 240 s injection of 1.3 g/L Fab fragment of trastuzumab preparation 1 diluted in PBS-P+, 60 s injection of 18 µg/ml Fc diluted in PBS-P+ (all injections over both flow cells).
- [0248] Regeneration: 2x30 s injections with Glycin-HCl pH 1.5.

Biacore Method 2:

[0249] Biacore method 2 setup was identical to above except a reduction of injection time for Fab fragment of trastuzumab to preserve sample and use of a new Fab fragment of trastuzumab batch (preparation 2). New injection time was 90 s. The resulting sensorgram of a candidate Fc binding polypeptide according to present invention is shown in FIG. 1B. The candidate binds to VH3 with a lower affinity than SEQ ID NO:59, while maintaining affinity for Fc.

Selection of Top Candidate Fc Binding Polypeptides

[0250] Based on screening results, candidates were selected for identification by sequencing using the following criteria:

[0251] (1) All candidates giving a Fab response below zero were taken forward.

[0252] (2) Of these, all candidates giving a capture level above 1600 Ru taken forward.

[0253] (3) Remaining candidates sorted based on Fc to Capture level ratio*.

[0254] (4) Top 70 candidates based on Fc to capture level selected for sequencing.

[0255] *It should be noted that both Fab and Fc interaction was impacted by the amount of candidate captured in the first step. Due to this, selection of candidates based on retained Fc interaction was done using the ratio of the Fc signal and the Capture level (referred to as Fc to Capture).

[0256] Selected candidates had a Fc to capture ratio of 1.35-1.68. The data for Z (SEQ ID NO:59) above gives a Fc to capture ratio of around 1.55-1.6 in comparison.

Sequencing of Selected Fc Binding Polypeptide VH Knock-Out Candidates

Materials and Methods

[0257] For sequencing the material used was: colony storage plates, Agar-MTP plate (Eurofins), and B069 sequencing primer (IDT). Selected colonies were transferred from the colony storage plates to Agar-MTP plates using pipette tips. Agar-MTP plates were sent together with the B069 primer to Eurofins for sequencing. Sequencing results analyzed using Geneious prime software.

Results

[0258] 57 unique Fc binding polypeptides were obtained corresponding to SEQ ID NO:1-57 in FIG. 10. Table 2 shows Biacore data and sequencing results for these candidates. The table is sorted based on Fc to Capture level ratio. Sequence alterations outside targeted region occurred for some variants and are indicated in comments. The most frequently occurring amino acid residues in positions 33, 40 and 51 are summarized in Table 3.

TABLE 2

Biacore data and sequencing results for the 57 candidates.				
Capture level_1	Analyte binding early_1: Fab	Analyte binding late_2: FC	FC to Capture ratio	Seq variant positions 33-40-51
1762	-37.6	2873	1.63	ADL
1787	-42.4	2914	1.63	AGR
1821	-17.2	2925	1.61	AKL
1784	-18.9	2907	1.63	ARI
2084	-2.85	3114	1.49	ARL
1790	-18.8	2881	1.61	DK
1741	-35.1	2860	1.64	DNS
1762	-17.9	2832	1.61	DRV
1713	-32.6	2849	1.66	EGL
1772	-18.3	2845	1.60	ERL
1736	-41.1	2832	1.63	GDR
1738	-19.4	2900	1.67	GDV
1704	-40.8	2862	1.68	GER
1782	-20.3	2875	1.61	GGI
2867	-8.61	4363	1.52	GGS
1704	-33.8	2836	1.66	GNI

TABLE 2-continued

Biacore data and sequencing results for the 57 candidates.				
Capture level_1	Analyte binding early_1: Fab	Analyte binding late_2: FC	FC to Capture ratio	Seq variant positions 33-40-51
1774	83.6	2965	1.67	GQI
2073	-4.06	3224	1.56	GRI
1775	-19.0	2927	1.65	GRL
2126	-3.11	3249	1.53	GRR
2099	-4.41	3117	1.48	HKI
1773	-37.3	2913	1.64	HKL
1740	-28.8	2845	1.64	HQS
1801	-17.9	2907	1.61	HRI
1813	-16.5	2934	1.62	HSR
1818	-15.7	2950	1.62	HSV
1818	-18.5	2916	1.60	KE
2180	-8.37	2972	1.36	KKL
1810	-34.0	2969	1.64	KRI
1763	-40.6	2876	1.63	NHS
1793	-17.3	2923	1.63	PDR
1785	-40.9	2909	1.63	PHR
1730	-40.9	2861	1.65	PHS
1757	-35.4	2891	1.65	PKL
1780	-17.3	2879	1.62	PKS
1729	-33.3	2839	1.64	PQR
1728	-33.0	2847	1.65	PSI
1755	-34.6	2891	1.65	QDI
1753	-17.6	2857	1.63	QDV
1774	-20.9	2861	1.61	QNS
1729	-17.1	2811	1.63	QQI
1787	-19.1	2895	1.62	QRI
1806	-19.0	2901	1.61	QRV
1679	-33.8	2797	1.67	RQR
1786	-17.0	2869	1.61	RQS
2247	-8.19	3026	1.35	RRI
1705	-39.3	2818	1.65	RRL
1801	-18.7	2927	1.63	SGV
1817	-15.8	2927	1.61	SRR
1816	-18.8	2929	1.61	SRS
1856	-19.3	3051	1.64	SRV
1739	-22.6	2916	1.68	TER
1746	-15.6	2864	1.64	TKG
1743	-18.5	2843	1.63	TKI
1753	-21.2	2813	1.61	TQI
1714	-26.5	2827	1.65	TRR
1805	-28.7	2951	1.63	TSR

TABLE 3

Summary of frequently occurring amino acid residues in positions 33 (pos 1), 40 (pos 2) and 51 (pos 51) in the Fc binding polypeptides disclosed herein.

Pos1	Pos2	Pos3
All 12 amino acids seen	All 9 amino acids seen	All 6 amino acids seen
13 of 64 = Glycine (G)	18 of 64 Arginine (R)	20 of 64 Isoleucine (I)
9 of 64 = Proline (P)	10 of 64 Lysine (K)	16 of 64 Arginine (R)
8 of 64 = Threonine (T)	32 of 64 positively charged	11 of 64 Leucine (L)
	41 of 64 charged	only one Glycine (G)
	57 of 64 polar	

Example 2

Biacore Evaluation of Fc Binding Polypeptide VH Knock-Out Candidates and Controls

Background

[0259] The selected 16 Fc binding polypeptide variants from the library screen were evaluated together with the control SEQ ID NO:58.

[0260] The following aspects were evaluated:

[0261] (1) Knock-out of VH3 interaction (tested through high concentration injection of Fab fragment generated from trastuzumab).

[0262] (2) Binding to trastuzumab.

[0263] (3) Kinetics/affinity assessment of Fc interaction (injection of Fc samples generated from trastuzumab)

[0264] (4) NaOH-stability (tested by reduction of trastuzumab binding after increasing number of treatments with 0.5M NaOH).

[0265] The different variants analyzed are summarized in Table 4.

TABLE 4

Overview of candidates evaluated in Biacore. Amino acids in bold are changed compared to SEQ ID NO: 58.			
Fc binding polypeptide variant	SEQ ID NO:	Sequence at aa position 33/40/51	Description
	58	SVL	Positive control for alkali stability and Fc interaction. Positive control for VH3 interaction.
ZGE85	1	GGS	Library candidate
ZGE86	2	GGI	Library candidate
ZGE87	3	SRV	Library candidate
ZGE88	4	GRL	Library candidate
ZGE89	5	QRI	Library candidate
ZGE90	6	ARI	Library candidate
ZGE91	7	SGV	Library candidate
ZGE92	8	TER	Library candidate
ZGE93	9	GER	Library candidate
ZGE94	10	ADL	Library candidate
ZGE95	11	AGR	Library candidate
ZGE96	12	EGL	Library candidate
ZGE97	13	HKL	Library candidate
ZGE98	14	GDV	Library candidate
ZGE99	15	QOI	Library candidate
ZGE100	16	QDI	Library candidate

Generation of his-Purified Candidate Proteins

Materials and Methods

[0266] Materials and equipment used were as follows: glycerol stocks for pAM095 (SEQ ID NO:58), and pAM200-215 (ZGE85-100, corresponding to SEQ ID NO:1-16); 2YT culture medium, Carbenicillin, IPTG, 100 ml baffled glass shake flasks, Infors HT Shaking incubator; His-GraviTrap and kit (Gytiva), PD10 GraviTraps (Gytiva).

[0267] Protein expression culture medium (2YT medium supplemented with 200 µg/ml Carbenicillin and 1 mM IPTG) was prepared and added to filled baffled shake flasks (20 ml per flask, 22 flasks in total). Each flask was inoculated with 5 µl from corresponding glycerol stock. The flasks were incubated in Infors HT shaking incubator for 24 hours at 27° C. and 190 rpm shaking, whereafter the cultures were pelleted and re-suspended in 5 ml PBS in Falcon tubes and frozen.

[0268] To generate crude Fc binding polypeptide variant lysates the frozen Falcon tubes were incubated in 80° C. water bath for 2 hours followed by pelleting of cell debris by centrifugation at 12000×g for 30 min.

[0269] Imidazole was added to the lysates to achieve the concentration recommended by the manufacturer and the samples were purified using His GraviTraps and kit according to kit instructions. Buffer exchange into HBS-P+ (0.1 M

HEPES, 1.5 M NaCl and 0.5% v/v Surfactant P20) (running buffer used in Biacore) was performed using PD10 GraviTraps and kit instructions.

[0270] Concentration of buffer exchanged samples was measured using NanoDrop (Thermo Scientific) according to manufacturer's instructions. Following purification, samples were kept in refrigerator until all measurements performed and were then transferred to eppendorf tubes and kept in freezer (−20° C.).

Ph-Scouting for Immobilization Conditions

Materials and Methods

[0271] Materials and equipment used were as follows: Fc binding polypeptide variants ZGE85-100 (corresponding to SEQ ID NO:1-16, respectively); Biacore CM5 chips (Cytiva); Biacore Acetate-buffers (Cytiva) with different pH; Biacore 8K+ instrument (Cytiva).

[0272] The eight first variants (SEQ ID NO:1-8) were used for pH-scouting (pH 4, pH 4.5, pH 5 and pH 5.5). The pH-scout was performed on Biacore 8K+ using the standard method defined in the software. pH 5 was found to give the best surface attraction for all variants. Hence, pH 5 was selected for immobilization of variants.

Initial Biacore Analysis of VH3 Knock-Out Variants

Material and Methods

[0273] Materials and equipment used were as follows: CM5 sensor chips (Cytiva), Biacore NHS coupling kit (Cytiva), Fc binding polypeptide variants from above; Biacore 8K+ (Cytiva); Fab (prepared from trastuzumab in-house and comprising SEQ ID NO:198 and 199), Fc (SEQ ID NO:200) from trastuzumab.

[0274] Immobilization was performed using a standard method in Biacore software with coupling of Fc binding polypeptide variants in FC2 and activation/inactivation in FC1. Fc binding polypeptide variants were diluted in pH5 Acetate buffer at approximately 30 µg/ml. The immobilization levels did vary somewhat between different Fc binding polypeptide variants (840-1190 RU).

Biacore Method:

[0275] Running buffer: HBS-P+

[0276] Flow rate: 5 µl/min

[0277] Sample injection: 120 s/2 min over both Flow Cells (FC1 and FC2)

[0278] Dissociation time: 30 s

[0279] Regeneration: 10 mM Glycin-HCl pH 1.5, 30 µl/min, 2×30 s

[0280] In each run SEQ ID NO:58 was immobilized in channel1 in FC2 as internal control. Channel3-channel8 was immobilized with the candidate Fc binding polypeptides. Multiple chips were used until all candidates were tested.

[0281] Injections for each channel (cycles) were as follows: buffer, 1 g/L Fab, 40 µg/ml Fc except for immobilized ligands ZGE99-ZGE100. For immobilized ligands ZGE99-ZGE100, injections for each channel (cycles) were as follows: buffer, 1 g/L Fab.

Results

[0282] All sensorgrams were generated as reference subtracted and the responses as the difference between baseline before injection and signal just before end of injection.

[0283] The response levels of the Fab (VH3) interaction of the Fc binding polypeptide variants are given in Table 5. The data shows that the Fc binding polypeptide variants (SEQ ID NO:1-16) exhibited a significantly reduced or abolished binding affinity for Fab, which included the VH3 from trastuzumab. A subset of the Fc binding polypeptide variants were analyzed further.

TABLE 5

Immobilization levels and response values for Fc binding polypeptide variants following injection of 1 g/L Fab and 40 µg/ml Fc, respectively. SEQ ID NO: 58 is a positive control for binding to VH3 from trastuzumab.

Immobilized ligand	Immobilized ligand level (RU)	Fab 1 g/l analyte binding Relative (RU)	40 µg/ml Fc analyte binding Relative (RU)
SEQ ID NO: 58	1169.5	2587.5	2650.1
ZGE85	987.3	39.1	2336.4
ZGE86	1031.2	-18.4	2550.2
ZGE87	1126.9	7.0	2562.2
ZGE88	1188.7	8.3	2647.9
ZGE89	1180.1	6.7	2615.5
ZGE90	1159.3	-0.9	2648.2
ZGE91	889.8	-7.0	1887.1
ZGE92	1135.4	9.0	2433.6
ZGE93	939.1	59.2	2346.0
ZGE94	1073.2	23.4	2490.1
ZGE95	999.5	-0.6	2037.3
ZGE96	839.4	-3.3	2055.8
ZGE97	1100.1	3.2	2546.0
ZGE98	890.0	7.4	2217.3
ZGE99	845.6	105.3	
ZGE100	1007.7	12.5	

Alkali Stability (0.5 M NaOH) Evaluation Using Biacore

Materials and Methods

[0284] Materials and equipment used were as follows: CM5 sensor chips (Cytiva), Biacore NHS coupling kit (Cytiva), Fc binding polypeptide variants ZGE85-ZGE100 (corresponding to SEQ ID NO:1-16, respectively); Biacore 8K+ (Cytiva); trastuzumab, 0.5 M NaOH.

[0285] Immobilization was performed using standard method in Biacore software with coupling of Fc binding polypeptide variants in FC2 and activation/inactivation in FC1. Fc binding polypeptide variants were diluted in pH5 Acetate buffer at approximately 30 µg/ml.

[0286] For each run SEQ ID NO:58 was immobilized in channel 1 (FC1) as internal control. Channel3-channel8 immobilized with other candidates. Multiple chips used until all candidates tested.

[0287] Immobilization levels vary somewhat between different Fc binding polypeptide variants (840-1190 Ru).

Biacore Method (Cycle):

[0288] Running buffer: HBS-P+

[0289] Flow rate: 5 µl/min

[0290] Sample injection 1 (trastuzumab 47 µg/ml): 200 s over both Flow Cells

[0291] Dissociation time 1: 60 s

[0292] Sample injection 2 (0.5 M NaOH): 300 s over both flow cells.

[0293] Regeneration: 10 mM Glycin-HCl pH 1.5, 30 µl/min, 2×30 s

[0294] This cycle was repeated more than 70 times to follow stability of trastuzumab response values.

[0295] All sensorgrams were generated as reference subtracted. Responses as difference between baseline before injection and signal just before end of injection.

Results

[0296] Stability was visualized by plotting normalized trastuzumab response values (% of response in cycle 1) for the first cycle as set out above to the last cycle run. SEQ ID NO:58 present as control in each separate stability run. FIG. 2 shows an example plot of alkali stability for tested Fc binding polypeptide variants in comparison to SEQ ID NO:58. In each cycle, 0.5 M NaOH was injected with a contact time of 5 min at 10 µl/min.

[0297] This data showed that the candidate Fc binding polypeptides SEQ ID NO:1-16 exhibited a stability in 0.5 M NaOH similar to that of SEQ ID NO:58 (see summary in Table 8).

Kinetics of Fc Interaction Using Immobilized Fc Binding Polypeptide Variants

Materials and Methods

[0298] Materials and equipment used were as follows: CM5 sensor chips (Cytiva), Biacore NHS coupling kit (Cytiva), Fc binding polypeptide variants ZGE99 (SEQ ID NO:15), ZGE100 (SEQ ID NO:16), ZGE86 (SEQ ID NO:2), ZGE91 (SEQ ID NO:7), ZGE98 (SEQ ID NO:14), respectively; Biacore 8K+ (Cytiva); Fc (prepared in-house) as above.

[0299] Immobilization was performed using standard method in Biacore software with coupling of Fc binding polypeptide variants in FC2 and activation/inactivation of FC1. Fc binding polypeptide variants were diluted in pH5 Acetate buffer at approximately 30 µg/ml.

[0300] Immobilization levels varied somewhat between different Fc binding polypeptide variants (840-1190 Ru).

Biacore Method:

[0301] Running buffer: HBS-P+

[0302] Used pre-defined method for multi-cycle kinetics on Biacore-8K+ using injections of buffer, 2.4 nM Fc, 12 nM Fc, 60 nM Fc, 300 nM Fc, 1500 nM Fc and 7500 nM Fc.

Results

[0303] Results were analyzed using pre-defined Evaluation method Multi-cycle kinetics. Note: These data were collected using chip immobilized for evaluation of last variants in above tests.

[0304] Kinetic parameters for evaluated Fc binding polypeptide variants shown in Table 6.

[0305] The data shows that in this assay set up, the Fc-binding polypeptides ZGE99 (SEQ ID NO:15), ZGE100 (SEQ ID NO:16), ZGE86 (SEQ ID NO:2), ZGE91 (SEQ ID NO:7), ZGE98 (SEQ ID NO:14) exhibited an K_D value of the interaction with Fc in the range of 10^{-10} M, which is in the same range as that of the control SEQ ID NO:58.

TABLE 6

Kinetic parameters for tested Fc binding polypeptide variants immobilized on surface. Analyte solutions: Fc. Kinetics model 1:1 binding.						
Immobilized ligand	Kinetics Chi ² (RU ²)	ka (1/Ms)	kd (1/s)	KD (M)	Rmax (RU)	tc
SEQ ID NO: 58	2.23e+03	5.22e+05	4.53e-04	8.68e-10	971.1	1.25e+12
ZGE99	1.68e+03	5.89e+05	3.87e-04	6.57e-10	858.4	2.63e+13
ZGE100	1.89e+03	5.35e+05	5.23e-04	9.78e-10	852.5	1.24e+12
ZGE86	1.91e+03	5.55e+05	4.27e-04	7.68e-10	894.8	1.43e+13
ZGE91	1.65e03	6.7e+05	5.12e-04	7.64e-10	785.7	8.34e+08
ZGE98	1.75e+03	5.24e+05	4.22e-04	8.06e-10	856.4	2.75e+13

Kinetics of Fc Interaction of the Fc Binding Polypeptides Using Immobilized Fc

Materials and Methods

[0306] Materials and equipment used were as follows: CM5 sensor chips (Cytiva), Biacore NHS coupling kit (Cytiva), Fc binding polypeptides ZGE86 (SEQ ID NO:2), ZGE87 (SEQ ID NO:3), ZGE88 (SEQ ID NO:4), ZGE89 (SEQ ID NO:5), ZGE90 (SEQ ID NO:6) and ZGE91 (SEQ ID NO:7); Biacore 8K+ (Cytiva); Fc as above (prepared in-house).

Results

[0310] Kinetic parameters for evaluated Fc binding polypeptide variants shown in Table 7.

[0311] The data shows that in this assay set up, the Fc-binding polypeptides, ZGE86 (SEQ ID NO:2), ZGE87 (SEQ ID NO:3), ZGE88 (SEQ ID NO:4), ZGE89 (SEQ ID NO:5), ZGE90 (SEQ ID NO:6) and ZGE91 (SEQ ID NO:7) exhibited an K_D value of the interaction with Fc in the range of 10^{-7} M, which is in the same range as that of the control SEQ ID NO:58.

TABLE 7

Kinetic parameters for tested Fc binding polypeptide variants with Fc immobilized on surface. Immobilized ligand: Fc 20 µg/ml pH 5. Kinetics model 1:1 binding.						
Analyte 2 Solution	Kinetics Chi ² (RU ²)	ka (1/Ms)	kd (1/s)	KD (M)	Rmax (RU)	tc
SEQ ID NO: 58	1.25e+01	2.10e+05	3.50e-02	1.67e-07	248.0	2.67e+09
ZGE86	1.43e+01	1.90e+05	2.90e-02	1.52e-07	257.9	2.66e+09
ZGE87	1.55e+01	1.89e+05	3.41e-02	1.81e-07	272.4	2.85e+09
ZGE88	1.78e+01	2.35e+05	3.05e-02	1.30e-07	278.4	1.23e+08
ZGE89	1.54e+01	2.04e+05	3.64e-02	1.78e-07	271.0	1.27e+08
ZGE90	1.48e+01	2.22e+05	3.79e-02	1.71e-07	262.4	1.33e+08
ZGE91	1.46e+01	1.86e+05	3.35e-02	1.80e-07	263.1	3.22e+12

[0307] Immobilization was performed using standard method in Biacore software with coupling of Fc in FC2 and activation/inactivation of FC1. Fc was diluted in pH5 Acetate buffer to 20 ug/ml. Immobilization levels were around 7500 Ru.

Biacore Method:

[0308] Running buffer: HBS-P+

[0309] Used pre-defined method for multi-cycle kinetics on Biacore-8K+ using injections of buffer, 2.4 nM Fc binding polypeptide variant, 12 nM Fc binding polypeptide variant, 60 nM Fc binding polypeptide variant, 300 nM Fc binding polypeptide variant, 1500 nM Fc binding polypeptide variant. 8 Fc binding polypeptide variants evaluated in parallel. Results were analyzed using pre-defined Evaluation method Multi-cycle kinetics.

Summary of Results and Comments

[0312] Table 8 summarizes the results from the detailed Biacore evaluation as described above. It was observed that all evaluated Fc binding polypeptides ZGE85-100, corresponding to SEQ ID NO:1-16, exhibited significantly reduced or abolished ability to bind to VH3 of trastuzumab. Furthermore, the analyzed Fc binding polypeptides showed a binding affinity for Fc of trastuzumab as well as stability as evaluated in 0.5 M NaOH in the same range as the control SEQ ID NO:58.

[0313] Based on the obtained results, three variants (ZGE90 (SEQ ID NO:6); ZGE91 (SEQ ID NO:7); and ZGE96 (SEQ ID NO:12) were selected for hexamer construction and further analysis (see Example 3). In selection, priority was given to complete absence of VH3 interaction and high alkali stability.

TABLE 8

Summary of results from detailed Biacore evaluation of Fc binding polypeptide variants.
Variants in bold were selected for hexamer construction and further analysis.

Variant	Fab	FC			Bivalent aff.	Stability 0.5M NaOH (% of SEQ ID NO: 58)
	binding	FC ka*	FC kd*	FC		
	RU (% of SEQ ID NO: 58)	(% of SEQ ID NO: 58)	(% of SEQ ID NO: 58)	affinity KD* (% of SEQ ID NO: 58)		
ZGE85	39.1 (1.6)	—	—	—	—	62.6
ZGE86	-18.4 (0)	90.5	120.7	109.9	113% (0.77 nM)	89 (100x) 98.6 (30x)
ZGE87	7.0 (0.27)	90	102.6	92.3	—	100.3 (64x)
ZGE88	8.3 (0.32)	111.9	114.8	128.5	—	100.1 (64x)
ZGE89	6.7 (0.26)	97.1	96.2	93.8	—	100.2 (64x)
ZGE90	-0.9 (0)	105.7	92.3	97.7	—	99.9 (64x)
ZGE91	-7.0 (0)	88.6	104.5	92.8	114% (0.76 nM)	98.4 (64x) 100 (30x)
ZGE92	9.0 (0.35)	83.1	75.3	62.3	—	95.1 (64x)
ZGE93	59.2 (2.5)	—	—	—	—	98.6 (30x)
ZGE94	23.4 (1)	—	—	—	—	99.8 (30x)
ZGE95	-0.6 (0)	98.5	102	100	—	99.5 (30x)
ZGE96	-3.3 (0)	76.4	85.4	65	—	99 (30x)
ZGE97	3.2 (0.14)	—	—	—	—	100.7 (30x)
ZGE98	7.4 (0.32)	82	130.9	106.4	107% (0.81 nM)	98.8 (30x)
ZGE99	105.3 (4.6)	—	—	—	132% (0.66 nM)	100 (30x)
ZGE100	12.5 (0.6)	—	—	—	88.8% (0.98 nM)	99.5 (30x)

*Monovalent affinity; FC immobilized

Example 3

Evaluation of Fc Binding Polypeptides VH Knock-Out Hexamer Candidates and Controls

Expression and Coupling of Four Variants in Hexameric Form

Plasmids and Cell-Banks

[0314] A subset of hexamer candidate variants according to general formula SEQ ID NO:207 were prepared. In particular, four sequences (ZGE90 (SEQ ID NO:6), ZGE91 (SEQ ID NO:91), ZGE96 (SEQ ID NO:12) and SEQ ID NO:58) as hexamers, which comprise the sequences according to SEQ ID NO:203-206, respectively, were ordered as DNA synthesis plasmids from ATUM (Newark, CA, USA) in the pJ401 plasmid backbone. The plasmids were transformed into chemically competent *E. coli* K12-017 cells and research cell-banks were made according to standard procedures.

Fermentation

[0315] 1 L fermentations was performed with the four abovementioned hexamers candidates. The results from protein expression are shown in Table 9.

TABLE 9

Samples were prepared by heat treatment in heat-block at 85° C., 5 min.

Name	SEQ ID NO:	Scale	Expression [g/L]
ZGE90-6	203	1 L	8.9
ZGE91-6	204	1 L	14.7
ZGE96-6	205	1 L	11.1
Hexamer of SEQ ID NO: 58	206	1 L	12.0

[0316] The products of the fermentations were purified on IgG Sepharose® as capture followed by Source™ 15Q as polishing step.

[0317] Coupling was made with epoxy coupling and the ligand density of the coupled prototypes was measured with amino acid analysis (AAA) by QC, and dynamic binding capacity (DBC) was measured with 1 mL cubed gel, see Table 10.

TABLE 10

Ligand density, Free His, Lys and DBC measurements at 6 min residence time and pH at elution.							
Name	SEQ ID NO:	Dry Weight Increase, mg/mL gel	Ligand density, mg/mL gel	Free His, %	Free Lys, %	DBC Qb10% 6 min rt, mg/mL gel	pH at peak elution
ZGE90-6	203	21.9	18.6	101	101	63.6	4.0
ZGE91-6	204	17.6	16	98	101	66.0	4.0
ZGE96-6	205	14.5	13.8	95	102	60.3	4.0
Hexamer of SEQ ID NO: 58	206	NA	16.1	95	100	63.8	3.8

Evaluation of Hexameric Variants by Chromatography

[0318] Coupled variants were initially tested for DBC with trastuzumab and negative binding for a Fab fragment cleaved from trastuzumab. None of the candidate hexamers showed binding to the Fab. Moreover, a pH gradient was run to analyze pH at elution and alkaline stability with an accelerated alkaline stability study. The results are shown in Table 11.

Alkaline Stability with Accelerated Alkaline Stability Study
[0319] Accelerated alkaline stability studies was performed on coupled hexamers with trastuzumab as testing substance. The DBC was initially measured for the candidate hexamers and a reference resin. Then, resins were soaked in 0.5 M NaOH for 4 h contact time (which is comparable to 16 cycles of 15 min contact time). After the NaOH soak, the resins were neutralized in running buffer and a new DBC measurement was performed. This cycling was then iterated for up to 24 h total contact time. The results are shown in Table 11.

TABLE 11

Accelerated alkaline studies. Incubation in 0.5M NaOH, 4 h followed by DBC measurements at 6 min residence time. The study is cycled up to 24 h total incubation time, corresponding to 96 cycles of 15 min contact time.								
Candidate	0 h	4 h	8 h	12 h	16 h	20 h	24 h	Residual capacity [%]
ZGE90-6	60.6	59.9	60.4	59.7	59.7	59.3	53.5	88%
ZGE91-6	61.7	58.5	62.7	61.9	61.9	61.4	56.2	91%
ZGE96-6	62.6	60.4	60.5	58.7	58.7	57.3	51.4	82%
Hexamer of SEQ ID NO: 58	65.9	58.4	60.8	60.5	60.5	60.4	59.1	90%

Results

[0320] All tested candidate hexamers exhibited a residual capacity of over 80% after 24 hours.

Binding Characteristics in Biacore

Materials and Methods

[0321] Hexameric and monomeric Fc binding polypeptide variants, Biacore CM5 chips; EDC/NHS coupling kit, Biacore Acetate-buffers with different pH; PBS-P+ running buffer (Biacore); 10 mM Glycin-HCl pH 1.5; Fab fragment

of trastuzumab, Fc, trastuzumab and unknown antibody (which does not comprise VH3) stock solutions; Biacore 8K+ instrument.

[0322] Fc binding polypeptide variants were immobilized on Biacore CM5 chip FC2 using standard EDC/NHS coupling method (FC1 activated and blocked) with Fc binding polypeptide variants diluted in Biacore acetate buffer with pH5 (410 s contact time). Coupling per channel according to Table 12 below.

TABLE 12

Summary of binding characteristics of monomer and hexamers of selected Fc binding polypeptides.				
Channel	Fc-binding polypeptide variant	SEQ ID NO:	Concentration in Acetate- buffer pH5 (ug/ml)	Coupled amount (Ru)
1	Hexamer of SEQ ID NO: 58	207	60	5461
2	ZGE90-6	203	48	6469
3	ZGE91-6	204	61	5870
4	ZGE96-6	206	51	5682
5	SEQ ID NO:58	58	22	1022
6	ZGE86	2	17	900
7	ZGE90	6	17	1154

[0323] Next a start-up cycle using 125 ng/ml trastuzumab was run (120 s contact time at 10 μ l/min) followed by regeneration using a 30 s pulse of 10 mM Glycin-HCl at 30 μ l/min. The different analytes were then injected over the surface using the same method. Injections summarized in Table 13. The results are displayed as reference subtracted sensorgrams (FC2-FC1).

TABLE 13

Summary of injections	
Injection	Sample
1	1000 μ g/ml Fab (Fab fragment of trastuzumab; comprises VH3)
2	40 μ g/ml Fc (Prepared by digestion from trastuzumab)
3	50 μ g/ml trastuzumab (comprises a VH3 region)
4	500 μ g/ml of unknown antibody, which does not comprise VH3

Results

[0324] FIG. 3 shows Biacore binding curves (reference-subtracted) following injection of 1 g/L Fab from trastuzumab.

[0325] Binding to the three different Fc containing targets (Fc only, trastuzumab and a monoclonal antibody which does not contain VH3) was tested. The Biacore curves can be found in FIG. 4 to 6 (FIG. 4: 40 µg/ml Fc; FIG. 5: 50 µg/ml trastuzumab; FIG. 6: 500 µg/ml of said monoclonal antibody which does not contain VH3). As can be seen, binding response is roughly inversely proportional to the amount of ligand immobilized on chip, with one major exception, SEQ ID NO:58 shows significantly reduced binding to trastuzumab. These results prompted a more thorough investigation of the relation between coupling level and binding response (capacity) for different targets using SEQ ID NO:58 (with dual interaction) and ZGE91 (with pure Fc interaction).

[0326] SEQ ID NO:58 and ZGE91 were immobilized (coupled) at different levels and Fc (400 mg/L), Fab (76.8 mg/L), trastuzumab (470 mg/L) and said monoclonal antibody which does not contain VH3 (510 mg/L) were injected at high concentrations over the surfaces.

[0327] A plot showing binding level as a function of coupled ligand level for the two hexameric ligands and the different targets can be found in FIG. 7.

[0328] Interestingly, the VH3 negative variant ZGE91 (SEQ ID NO:7) showed a higher binding capacity than SEQ ID NO:58, which binds the VH3 and Fc regions of trastuzumab.

NaOH Stability in Biacore Assay

Materials and Methods

[0329] Hexameric and monomeric Fc binding polypeptide variants, Biacore CM5 chips; EDC/NHS coupling kit, Biacore Acetate-buffers with different pH; PBS-P+ running buffer (Biacore); 10 mM Glycin-HCl pH 1.5; trastuzumab and stock solution; Biacore 8K+ instrument.

[0330] Fc binding polypeptide variants were immobilized on Biacore CM5 chip FC2 using standard EDC/NHS coupling method (FC1 activated and blocked) with Fc binding polypeptide variants diluted in Biacore acetate buffer with pH5/pH4.5 (205 s contact time). Coupling per channel was according to Table 14 below. The experiment aimed to reach close to 1000 Ru immobilized for each variant.

TABLE 14

Coupling per channel.					
Channel	Fc binding polypeptide variant	SEQ ID NO:	Concentration in Acetate-buffer (µg/ml)	Contact time in coupling (s)	Coupled amount (Ru)
1	Hexamer of SEQ ID NO:58	207	11.25/pH4.5	205	1004
2	ZGE90-6	203	7.5/pH5	205	1072
3	ZGE91-6	204	7.5/pH5	205	750
4	ZGE96-6	205	7.5/pH5	205	872
5	ZGE91	7	15/pH5	205	832
6	ZGE95	11	15/pH5	205	924
7	ZGE96	12	15/pH5	205	902

Biacore Method

[0331] Injection of 96 µg/ml trastuzumab at 10 µl/min and 300 s contact time, 2×10 mM Glycin-HCl pH 1.5 at 30 µl/min and 30 s contact time, 0.5M NaOH at 10 µl/min and 600 s contact time. The cycle repeated 100x. Results showed are based on reference subtracted data (from Cycle 2 onward).

Results

[0332] Results are shown be found in FIGS. 7 and 8. Coupling level of ligands was adjusted to be in the linear response range. Note that differences in the exact coupling level can have some impact on obtained stability plots.

[0333] Based on the Biacore data (FIG. 7), it seems plausible that ZGE91-6 (SEQ ID NO:204) exhibits a high optimal ligand density and hence a potential for increased capacity. The data in FIG. 7 also indicates that optimal coupling level is dependent on the size of the target protein. It may be that that complex bi-specific formats or large Fc fusion proteins would benefit from a product with lower ligand density.

Example 4

Evaluation of Fc Binding Polypeptides VH Knock-Out Candidates in Different Scaffolds

[0334] It is possible to introduce the mutations in positions 33, 40 and 51 in a different scaffold background. For instance SEQ ID NO:59 may be modified with the amino acid residues according to Table 2 in the three positions. The resulting Fc binding polypeptides are listed in FIG. 10 as SEQ ID NO:208-264.

[0335] Additionally, a selection of mutations as disclosed above may be introduced into any scaffolds, such as any one of domains A (SEQ ID NO:62), B (SEQ ID NO:61), C (SEQ ID NO:60), D (SEQ ID NO:63) and E (SEQ ID NO:64) of SpA or derivatives thereof, such as domain Z (SEQ ID NO:59) or variants thereof, such as SEQ ID NO:58.

[0336] Evaluation and analysis was performed essentially as described in Example 2, section entitled "Initial Biacore analysis of VH3 knock-out and retained Fc binding".

[0337] It is expected that the Fc binding polypeptides SEQ ID NO:208-264 and 269-304 exhibit Fc binding affinity in the same range as SEQ ID NO:59 and significantly reduced or abolished affinity for VH3 of trastuzumab compared to SEQ ID NO:59. Such a comparison can also be made with SEQ ID NO:58.

[0338] For the purpose of showing that the mutations in positions 33, 40 and 51 have the same effect in any scaffold background, three sets of mutations for positions 33, 40 and 51 according to the above, A/R/I, S/G/V, and A/G/R, were introduced into domains A (ZGE0206) (SEQ ID NO:62), B (ZGE0207) (SEQ ID NO:61), C (ZGE0208) (SEQ ID NO:60), D (ZGE0205) (SEQ ID NO:63) and E (ZGE0204) (SEQ ID NO:64), Zwt (ZGE0209) (SEQ ID NO 59) as well as modified versions of protein A such as ZGE0201 (SEQ ID NO 265), ZGE0202 (SEQ ID NO 266), and ZGE0203 (SEQ ID NO 267), and modified version of Z such as Z var (ZGE0210) (SEQ ID NO 268). These constructs were tested in a Biacore as disclosed above, with small modifications as indicated below.

[0339] Monomeric protein A or protein A derived IgG binding polypeptide variants according to the above (in

Biacore 10 mM Acetate-buffers (pH4, pH4.5, pH5 or pH5.5) were immobilized on Biacore CM5 chips FC2 and FC1 as reference (activated and deactivated) using standard EDC/NHS coupling method. Experiments used PBS-P+ running buffer (Biacore); regeneration buffer 10 mM Glycin-HCl pH 1.5; Target molecules trastuzumab, adalimumab and adalimumab Fc fragment at concentrations 0, 5, 25, 50, 100, 200, 400, 800, 1600 nM, and trastuzumab Fab fragment at concentrations 0, 50, 100, 200, 400, 800, 1600 nM were used with a Biacore 8K+ instrument.

[0340] Each cycle used a flowrate of 10 μ l/min with 600 seconds association and disassociation of each target molecule and concentration followed by regeneration using two 30 s pulses of 10 mM Glycin-HCl at 30 μ l/min. The results are presented in Table 15 below for max response of 800 nM of each target molecule, wherein the results have been rounded to the nearest 50 ru. Exemplary samples are displayed as reference subtracted sensorgrams (FC2-FC1) in FIGS. 11 and 12.

TABLE 15

Binding in Ru to trastuzumab, adalimumab, Fc fragment from adalimumab, Fab fragment from trastuzumab, for different scaffolds.						
Sample No.	SEQ ID NO	Trastuzumab (Ru)	Adalimumab (Ru)	Fc fragment (Ru)	Fab fragment (Ru)	Immobilized ligand level (Ru)
ZGE0201	265	5200	5050	1700	150	860
ZGE0211	269	5550	5900	2400	50	1110
ZGE0221	279	3550	3800	1400	50	710
ZGE0231	289	3800	4050	1500	50	780
ZGE0202	266	5850	5250	1800	50	1000
ZGE0212	270	5700	6000	2400	0	1170
ZGE0222	280	5250	5500	2000	0	1050
ZGE0232	290	3600	3800	1400	0	840
ZGE0203	267	4700	3250	1250	100	1120
ZGE0213	271	3250	3550	1650	0	1200
ZGE0223	281	1355	1400	650	0	1280
ZGE0233	291	1400	1550	700	0	1230
Domain E						
ZGE0204	64	5050	4200	1450	850	780
ZGE0214	272	3850	4100	1650	0	780
ZGE0224	282	3550	3650	1350	0	740
ZGE0234	292	2750	2950	1100	0	800
Domain D						
ZGE0205	63	5350	4600	1600	750	830
ZGE0215	273	4700	5000	1950	0	930
ZGE0225	283	4100	4200	1500	0	810
ZGE0235	293	4800	5050	1850	0	1000
Domain A						
ZGE0206	62	4850	4050	1350	1150	710
ZGE0216	274	4350	4550	1850	0	1140
ZGE0226	284	1500	1500	600	0	810
ZGE0236	294	2550	2700	1050	0	750
Domain B						
ZGE0207	61	6250	5500	1750	1700	930
ZGE0217	275	4650	4900	1850	0	1170
ZGE0227	285	3300	3300	1300	0	940
ZGE0237	295	3050	3250	1250	0	800
Domain C						
ZGE0208	60	6000	50000	1750	3000	950
ZGE0218	276	5450	5650	2100	50	1000
ZGE0228	286	3300	3200	1300	0	1060
ZGE0238	296	3650	3750	1500	0	990
Z wt						
ZGE0209	59	4400	4000	1600	50	970
ZGE0219	277	5100	5400	2000	0	1180
ZGE0229	287	3350	3550	1400	0	930
ZGE0239	297	2950	3150	1200	0	850
Z var						
ZGE0210	268	4600	4300	1750	50	1090
ZGE0220	278	4950	5250	2000	0	1230
ZGE0230	288	2650	2850	1150	0	930
ZGE0240	297	3150	3350	1300	0	710

[0341] Each scaffold was tested in a set of four and is grouped accordingly in the table above: 1) wt, 2) with mutation A/R/I, 3) with mutation S/G/V, and 4) with mutation A/G/R. Thus, as an example, for domain C the test set is according to the following: 1) ZGE0208 (wt), 2) ZGE0218 (A/R/I), 3) ZGE228 (S/G/V), 4) ZGE238 (A/G/R).

[0342] In general, the response for Fc-binding is lower than the response for the full antibody, as the Fc fragment is smaller than the full antibody. The response in a Biacore system is directly related to the change in mass concentration on the surface, so that molar responses (i.e. responses for a given number of molecules) are proportional to the size of the molecule involved. Thus, a given number of molecules binding to the surface will give a lower response if the molecule is small.

[0343] As is evident from Table 15, and FIGS. 11 and 12, the tested mutations had the effect of substantially maintaining the Fc binding activity, but substantially abolishing the Fab binding activity for all domains and versions thereof tested, in comparison with the wildtype or a mutated version of the SpA domain lacking the above specified mutations for positions 33, 40 and 51.

[0344] Thus, it is shown that the mutations as disclosed herein for positions 33, 40 and 51, as defined above, retains the inherent ability to bind to the Fc region of antibodies/immunoglobulins, in particular to the Fc region of the antibody subclasses IgG1, IgG2 and IgG4, and show a reduced or abolished binding affinity for the VH3 region of the antibody trastuzumab, for all Protein A scaffolds, such as naturally occurring SpA domains, as well as versions of the same.

ITEMIZED LIST OF EMBODIMENTS

1. Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof, wherein said polypeptide has binding affinity to an Fc region of an immunoglobulin, and has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:59 for the same VH3 region.

2. Fc binding polypeptide comprising a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii), wherein i), ii) and iii) are defined as follows:

i)

(SEQ ID NO: 65)

EX₂₅QX₂₇X₂₈X₂₉X₃₀IX₃₂X₃₃LX₃₅X₃₆X₃₇PSX₄₀SX₄₂X₄₃X₄₄LX₄₆EAX₄₉
X₅₀X₅₁NX₅₃X₅₄

wherein, independently from each other,

[0345] X₂₅ is selected from E and D;

[0346] X₂₇ is selected from R and H;

[0347] X₂₈ is selected from N, A, S, H and W;

[0348] X₂₉ is selected from A, G and K;

[0349] X₃₀ is selected from F and A;

[0350] X₃₂ is selected from Q and H;

[0351] X₃₅ is selected from K, R and H;

[0352] X₃₆ is selected from D and H;

[0353] X₃₇ is selected from D and E;

[0354] X₄₂ is selected from A, K, L, R and T;

[0355] X₄₃ is selected from N, E, A, K and S;

[0356] X₄₄ is selected from L, I and V;

[0357] X₄₆ is selected from A, G and K;

[0358] X₄₉ is selected from K, Q and R;

[0359] X₅₀ is selected from K and R;

[0360] X₅₃ is selected from D, E and K; and

[0361] X₅₄ is selected from A and S;

[0362] ii) an amino acid sequence which has at least 83% identity to a sequence defined by i)

[0363] iii) an amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, residues 24-54 in SEQ ID NO:60, residues 24-54 in SEQ ID NO:61, residues 24-54 in SEQ ID NO:62, residues 27-57 in SEQ ID NO:63 and residues 22-52 in SEQ ID NO:64,

wherein additionally, in each of i), ii) and iii) independently from each other,

[0364] X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;

[0365] X₄₀ is selected from E, G, R, D, K, Q, N, H and S; and

[0366] X₅₁ is selected from L, V, S, I, R and G;

with the proviso that when X₅₁ is L then X₃₃X₄₀ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X₅₁ is G then X₃₃X₄₀ is TK.

3. Fc binding polypeptide according to item 2, wherein said Fc binding polypeptide has a lower binding affinity for the VH3 region of trastuzumab than SEQ ID NO:59.

4. Fc binding polypeptide according to item 2 or 3, wherein X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D and K, such as selected from T, S, G, Q, A, H, R, P, D and K, such as selected from T, S, G, Q, A, H, R, P and D or selected from T, S, G, Q, A, H, R, P and K; such as selected from T, S, G, Q, A, H, R and P; such as selected from T, S, G, Q, A, H and P or selected from T, G, Q, A, H, R and P; such as selected from T, G, Q, P, A and H; such as selected from T, G, Q, H and P; such as selected from T, G, Q and P or selected from G, Q, H and P or selected from T, G, H and P; such as the group selected from G and P; such is G.

5. Fc binding polypeptide according to any one of items 2-4, wherein X₃₃ is selected from T, S, G, Q, A, E and H, such as selected from T, S, G, Q, A and E, such as selected from S, G, Q and A.

6. Fc binding polypeptide according to any one of items 2-5, wherein X₃₃ is selected from S, G, A, E and H, such as selected from S, G, A and E.

7. Fc binding polypeptide according to any one of items 2-6, wherein X₃₃ is selected from A, S and E.

8. Fc binding polypeptide according to any one of items 2-7, wherein X₃₃ is selected from A and S or selected from A and E or selected from S and E.

9. Fc binding polypeptide according to any one of items 2-8, wherein X₃₃ is S.

10. Fc binding polypeptide according to any one of items 2-9, wherein X₄₀ is selected from E, G, R, D, K, Q, N and S or selected from E, G, R, D, K, Q, H and S or selected from G, R, D, K, Q, N, H and S, such as selected from G, R, D, K, Q, and S, such as selected from G, R, D, K and Q; such as selected from R, D, K and Q; such as selected from R, K and Q; such as selected from R and K; such as is R or is K.

11. Fc binding polypeptide according to any one of items 2-10, wherein X₄₀ is selected from E, G, R, D, K and Q, such as selected from E, G, R, D and K or selected from E, G, R, D and Q.

12. Fc binding polypeptide according to any one of items 2-11, wherein X₄₀ is selected from G, R, E and D.

13. Fc binding polypeptide according to any one of items 2-12, wherein X_{40} is selected from G, R and D.
14. Fc binding polypeptide according to any one of items 2-11, wherein X_{40} is selected from G, R and K.
15. Fc binding polypeptide according to any one of items 2-14, wherein X_{40} is selected from G and R.
16. Fc binding polypeptide according to any one of items 2-15, wherein X_{40} is G.
17. Fc binding polypeptide according to any one of items 2-14, wherein X_{40} is R.
18. Fc binding polypeptide according to any one of items 2-17, wherein X_{51} is selected from L, V, I, R and S.
19. Fc binding polypeptide according to any one of items 2-18, wherein X_{51} is selected from L, I, R and S.
20. Fc binding polypeptide according to any one of items 2-18, wherein X_{51} is selected from or selected from L, V, I and R.
21. Fc binding polypeptide according to one of items 2-18 and 20, wherein X_{51} is selected from I, V and R; or is selected from L, I and R; or is selected from V, L and I.
22. Fc binding polypeptide according to any one of item 2-18 and 20-21, wherein X_{51} is selected from V and L; or is selected from V and I; or is selected from V and R; or is selected from L and I; or is selected from L and R; or is selected from I and R.
23. Fc binding polypeptide according to any one of items 2-22, wherein X_{51} is L or R.
24. Fc binding polypeptide according to any one of items 2-22, wherein X_{51} is I.
25. Fc binding polypeptide according to any one of items 2-18 and 20-22, wherein X_{51} is V.
26. Fc binding polypeptide according to any one of items 2-18, wherein X_{51} is selected from L, V, S, I and R with the proviso that when X_{51} is L then $X_{33}X_{40}$ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK.
27. Fc binding polypeptide according to item 26, wherein X_{51} is selected from L, V, S, I and R with the proviso that when X_{51} is L then $X_{33}X_{40}$ are AD or HK.
28. Fc binding polypeptide according to any one of items 2-27, wherein at least one, such as two or all three, of amino acid residues X_{33} , X_{40} and X_{51} is/are an uncharged amino acid residue.
29. Fc binding polypeptide according to item 28, wherein the amino acid residue in X_{33} is an uncharged amino acid.
30. Fc binding polypeptide according to any one of items 28-29, wherein the amino acid residue in X_{40} is an uncharged amino acid.
31. Fc binding polypeptide according to any one of items 2-30, wherein at least two of the amino acid residues in positions X_{33} , X_{40} and X_{51} are mutated compared to SEQ ID NO:59.
32. Fc binding polypeptide according to item 2 or 3, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GGS, GNI, GQI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, POR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, ROS, RRI, RRL, SGV, SRR, SRS, SRV, TER, TKG, TKI, TQI, TRR and TSR, such as the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GNI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, POR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, ROS, RRI, RRL, SGV, SRR, SRS, SRV, TER, TKI, TQI, TRR and TSR.
33. Fc binding polypeptide according to item 32, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGS, GGI, SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV, GQI and QDI or the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV and QDI.
34. Fc binding polypeptide according to any one of items 32-33, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, TER, AGR, EGL, HKL and GDV.
35. Fc binding polypeptide according to any one of items 32-34, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, ARI, SGV, AGR, EGL and HKL.
36. Fc binding polypeptide according to any one of items 32-35, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, ARI, SGV, AGR and EGL.
37. Fc binding polypeptide according to any one of items 32-34, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, GRL, QRI, ARI, SGV, TER, AGR, EGL and GDV.
38. Fc binding polypeptide according to any one of items 32-34 and 37, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, SRV, QRI, ARI, SGV, AGR and GDV.
39. Fc binding polypeptide according to any one of items 32-33, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV, GQI and QDI.
40. Fc binding polypeptide according to any one of items 32-34 and 37, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of GGI, GRL, ARI, SGV, AGR and EGL.
41. Fc binding polypeptide according to any one of items 32-34 and 39-40, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI, SGV, AGR and EGL.
42. Fc binding polypeptide according to any one of items 32-34 and 39-41, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of SGV, AGR and EGL; or the group consisting of ARI, AGR and EGL; or the group consisting of ARI, SGV and EGL; or the group consisting of ARI, SGV and AGR.
43. Fc binding polypeptide according to any one of items 32-34 and 39-42, wherein $X_{33}X_{40}X_{51}$ are selected from the group consisting of ARI and SVG; or the group consisting of ARI and AGR; or the group consisting of ARI and EGL; or the group consisting of SGV and AGR; or the group consisting of SGV and EGL; or the group consisting of AGR and EGL.
44. Fc binding polypeptide according to any one of items 32-34 and 39-42, wherein $X_{33}X_{40}X_{51}$ is ARI or SGV or AGR or EGL.
45. Fc binding polypeptide according to any one of items 2-44, wherein $X_{33}X_{40}X_{51}$ is SGV.
46. Fc binding polypeptide according to any one of items 2-45, wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57 and 208-264; such as the group consisting of SEQ ID NO:1-56 and 208-263; or such as the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56; or such as the group consisting of 208-264; such as SEQ ID NO:208-263.
47. Fc binding polypeptide according to item 46, wherein sequence i) corresponds to the sequence from position 24 to

position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223; such as the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223; or such as the group consisting of SEQ ID NO:1-16, such as SEQ ID NO:2-14 and 16; or such as the group consisting of 208-223; such as SEQ ID NO:209-221 and 223.

48. Fc binding polypeptide according to any one of items 46-47, wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14 or the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221.

49. Fc binding polypeptide according to any one of items 46-48, wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219 or wherein sequence i) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 12, 209, 211, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:2, 4, 6, 7 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214 and 219.

50. Fc binding polypeptide according to any one of items 46-49, wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:6, 7, 11 and 12 or the group consisting of SEQ ID NO:213, 214, 218 and 219 or wherein sequence i) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 12, 213, 214 and 219; such as the group consisting of SEQ ID NO:6, 7 and 12 or the group consisting of SEQ ID NO:213, 214 and 219.

51. Fc binding polypeptide according to any one of items 46-50, wherein sequence i) corresponds to the sequence from position 24 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218; such as the group consisting of SEQ ID NO:6, 7 and 11 or the group consisting of SEQ ID NO:213, 214 and 218 or wherein sequence i) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 213 and 214; such as the group consisting of SEQ ID NO:6 and 7 or the group consisting of SEQ ID NO:213 and 214.

52. Fc binding polypeptide according to any one of items 46-51, wherein sequence i) corresponds to the sequence from position 24 to position 54 in SEQ ID NO:7 or SEQ ID NO:214.

53. Fc binding polypeptide according to any one of items 1-52, wherein said polypeptide derived from SpA or said Sequence A forms part of a three-helix bundle protein domain.

54. Fc binding polypeptide according to item 53, wherein said polypeptide derived from SpA or said Sequence A forms part of two helices with an interconnecting loop, within said three-helix bundle protein domain.

55. Fc binding polypeptide according to any one of items 53-54, wherein said three-helix bundle protein domain is selected from bacterial receptor domains.

56. Fc binding polypeptide according to item 55, wherein said three-helix bundle protein domain is selected from domains of protein A from *Staphylococcus aureus* or derivatives thereof.

57. Fc binding polypeptide according to one of items 2-56, further comprising a Sequence B arranged N-terminally of said sequence A, which Sequence B consists of an amino acid sequence selected from iv) and v), and wherein iv) and v) are defined as follows:

iv) (SEQ ID NO: 66)
X₈X₉X₁₀X₁₁AFYX₁₅IX₁₇X₁₈X₁₉PX₂₁LX₂₃

wherein, independently from each other,

[0367] X₈ is selected from E, D and A;

[0368] X₉ is selected from Q, A, L, W, E, V, K, T and H;

[0369] X₁₀ is selected from Q and H;

[0370] X₁₁ is selected from N, A, S, E, K, H, Q, Y, T, F, L, W, I, M, V and R;

[0371] X₁₅ is selected from E, H and Q;

[0372] X₁₇ is selected from L and H;

[0373] X₁₈ is selected from H, N and K;

[0374] X₁₉ is selected from L and M;

[0375] X₂₁ is selected from N, Y and S;

[0376] X₂₃ is selected from T and N;

[0377] v) an amino acid sequence which has at least 75% identity to a sequence defined by iv).

58. Fc binding polypeptide according to any preceding item, comprising a binding module sequence C, which Sequence C consists of Sequence A according to any one of items 2-56 and Sequence B according to item 57, in the following order from the N-terminus to the C-terminus

[Sequence B]-[Sequence A]

or any amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, residues 8-54 in SEQ ID NO:60, residues 8-54 in SEQ ID NO:61, residues 8-54 in SEQ ID NO:62, residues 11-57 in SEQ ID NO:63 and residues 6-52 in SEQ ID NO:64.

59. Fc binding polypeptide in according to any preceding item, comprising a binding module Sequence C, which Sequence C consists of the sequences [H1], [L1] and [Sequence A] in the following order from the N-terminus to the C-terminus

[H1]-[L1]-[Sequence A],

wherein [Sequence A] is as defined in any one of items 2-56, and wherein, independently from each other, [H1] is selected from the group consisting of

EQQNAFYEILH; (SEQ ID NO: 67)

DQQAIFYEILH; (SEQ ID NO: 68)

EAQEAIFYEILH; (SEQ ID NO: 69)

-continued	and [L1] is selected from the group consisting of:		
DQQSAFYEILH;	(SEQ ID NO: 70)		(SEQ ID NO: 93)
AQQAAFYEILH;	(SEQ ID NO: 71)	LPNLT;	(SEQ ID NO: 94)
EQQNAFYEILN;	(SEQ ID NO: 72)	LPNLN;	(SEQ ID NO: 95)
EQQAAFYEILH;	(SEQ ID NO: 73)	MPNLN;	(SEQ ID NO: 96)
AQQSAFYEILH;	(SEQ ID NO: 74)	LPYLT;	(SEQ ID NO: 97)
EQQAFYEILH;	(SEQ ID NO: 75)	LPSLT;	(SEQ ID NO: 98)
EHQNAFYEILH;	(SEQ ID NO: 76)	MPNLL;	(SEQ ID NO: 99)
EAQNAFYEILH;	(SEQ ID NO: 77)	LPNGN;	(SEQ ID NO: 100)
ETQNAFYEILH;	(SEQ ID NO: 78)	LPNLS;	(SEQ ID NO: 101)
EQQSAFYEILH;	(SEQ ID NO: 79)	GPNAN;	(SEQ ID NO: 102)
EAQNAFYKILH;	(SEQ ID NO: 80)	VPNLN;	(SEQ ID NO: 103)
EQQEAFYEILH;	(SEQ ID NO: 81)	LPNLR; and	(SEQ ID NO: 104)
EAQKAFYEILK;	(SEQ ID NO: 82)	LPNLL.	
EQHNAFYEILH;	(SEQ ID NO: 83)	60. Fc binding polypeptide according to any one of items 58-59, comprising a binding module Sequence C selected from the group consisting of	
EHHHALYHILH;	(SEQ ID NO: 84)	EQQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 105)
EQQKAFYAILH;	(SEQ ID NO: 85)	EQQNAFYEILHLPNLN-[Sequence A];	(SEQ ID NO: 106)
DQQSAFYEILN;	(SEQ ID NO: 86)	DQQAAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 107)
AQQNAFYQVLN;	(SEQ ID NO: 87)	EAQEAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 108)
EQQKAFYEILH;	(SEQ ID NO: 88)	DQQSAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 109)
EQQKAFYEILK;	(SEQ ID NO: 89)	AQQAAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 110)
EQQNAFYEILS;	(SEQ ID NO: 90)	EQQAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 111)
EQQNAFYHILH; and	(SEQ ID NO: 91)	AQQSAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 112)
		EQQNAFYEILNMPNLN-[Sequence A];	(SEQ ID NO: 113)
EQQNAFYHILN;	(SEQ ID NO: 92)	EAQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 114)

-continued

EQQQAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 115)	vi)	(SEQ ID NO: 108)
EQQNAFYEILHLPYLT-[Sequence A];	(SEQ ID NO: 116)	EAQEAFYEILHLPNLT-[Sequence A]	
ETQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 117)	wherein [Sequence A] is as defined in any one of items 2-56;	
EAQNAFYKILHLPNLT-[Sequence A];	(SEQ ID NO: 118)	[0378] vii) an amino acid sequence which has at least 85% identity to the sequence defined in vi);	
EQQNAFYEILHLPNGN-[Sequence A];	(SEQ ID NO: 119)	viii)	(SEQ ID NO: 106)
EHQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 120)	EQQNAFYEILHLPNLT-[Sequence A]	
EQQNAFYEILNMPNLT-[Sequence A];	(SEQ ID NO: 121)	wherein [Sequence A] is as defined in any one of items 2-56; and	
EQQSAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 122)	[0379] ix) an amino acid sequence which has at least 85% identity to the sequence defined in viii).	
EQQEAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 123)	62. Fc binding polypeptide according to item 61, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-57 and 208-264; such as the group consisting of SEQ ID NO:1-56 and 208-263; or such as the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56; or such as the group consisting of SEQ ID NO:208-264; such as SEQ ID NO:208-263.	
EAQKAFYEILKLPNLT-[Sequence A];	(SEQ ID NO: 124)	63. Fc binding polypeptide according to any one of items 61-62, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223; such as the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223; or such as the group consisting of SEQ ID NO:1-16, such as SEQ ID NO:2-14 and 16; or such as the group consisting of 208-223; such as SEQ ID NO:209-221 and 223.	
EQHNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 125)	64. Fc binding polypeptide according to any one of items 61-63, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14 or the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221.	
EHHHALYHILHLPNLT-[Sequence A];	(SEQ ID NO: 126)	65. Fc binding polypeptide according to any one of items 61-64, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219 or wherein sequence vi) or viii) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 12, 209, 211, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:2, 4, 6, 7 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214 and 219.	
AQQSAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 127)	66. Fc binding polypeptide according to any one of items 61-65, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:6, 7, 11 and 12 or the group consisting of SEQ ID NO:213, 214, 218 and 219 or wherein sequence vi) or viii) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 12, 213, 214 and 219; such as the group consisting of SEQ ID NO:6, 7 and 12 or the group consisting of SEQ ID NO:213, 214 and 219.	
EQQKAFYAILHLPYLT-[Sequence A];	(SEQ ID NO: 128)		
DQQSAFYEILNMPNLT-[Sequence A];	(SEQ ID NO: 129)		
AQQNAFYQVLNMPNLT-[Sequence A];	(SEQ ID NO: 130)		
EQQKAFYEILKLPNLT-[Sequence A];	(SEQ ID NO: 131)		
EQQNAFYEILSLPNLT-[Sequence A];	(SEQ ID NO: 132)		
EQQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 133)		
EQQNAFYHILHLPNLT-[Sequence A];	(SEQ ID NO: 134)		
EHQNAFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 135)		
EQQAIFYEILHLPNLT-[Sequence A];	(SEQ ID NO: 136)		
AQQAIFYEILHLPNLT-[Sequence A]; wherein [Sequence A] is as defined in any one of items 2-56.	(SEQ ID NO: 137)		

61. Fc binding polypeptide according to any one of items 58-60, comprising a binding module Sequence C selected from the group consisting of:

67. Fc binding polypeptide according to any one of items 61-66, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218; such as the group consisting of SEQ ID NO:6, 7 and 11 or the group consisting of SEQ ID NO:213, 214 and 218 or wherein sequence vi) or viii) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 213 and 214; such as the group consisting of SEQ ID NO:6 and 7 or the group consisting of SEQ ID NO:213 and 214.

68. Fc binding polypeptide according to any one of items 61-67, wherein sequence vi) or viii) corresponds to the sequence from position 8 to position 54 in SEQ ID NO:7 or SEQ ID NO:214.

69. Fc binding polypeptide according to any preceding item, comprising a sequence [S1], [binding module Sequence C] and [S2] in the following order from the N-terminus to the C-terminus

[S1]-[binding module Sequence C]-[S2],

wherein [S1] or [S2] may be present or absent and [binding module sequence C] is as defined in any one of items 58-68, and

wherein, independently from each other,

[S1] is selected from the group consisting of

ADNKFNK;	(SEQ ID NO: 138)
VDKFDK;	(SEQ ID NO: 139)
VDNKFNK;	(SEQ ID NO: 140)
IAAKHDK;	(SEQ ID NO: 141)
IAAQHDK;	(SEQ ID NO: 142)
ADNNFNK;	(SEQ ID NO: 143)
IAAKFDE;	(SEQ ID NO: 144)
PAAKHDK;	(SEQ ID NO: 145)
ADNAFNT;	(SEQ ID NO: 146)
FNK;	(SEQ ID NO: 147)
ADNRFNE;	(SEQ ID NO: 148)
IDSKFDE;	(SEQ ID NO: 149)
ADNRFNR;	(SEQ ID NO: 150)
ADNKHNK;	(SEQ ID NO: 151)
ADSKFDE;	

-continued

IDAKHDE;	(SEQ ID NO: 152)
QQNKFNK;	(SEQ ID NO: 153)
ADNKFHK;	(SEQ ID NO: 154)
KFNK;	(SEQ ID NO: 155)
ADNNFNR;	(SEQ ID NO: 156)
AAAKHDK;	(SEQ ID NO: 157)
IDNKFNK;	(SEQ ID NO: 158)
IDAKFDE;	(SEQ ID NO: 159)
DNNFNK;	(SEQ ID NO: 160)
ADNKFNE;	(SEQ ID NO: 161)
AAQHDHDK;	(SEQ ID NO: 162)
and	
AAAKFDE;	(SEQ ID NO: 163)

and [S2] is selected from the group consisting of

QAPK;	(SEQ ID NO: 164)
QAPP;	(SEQ ID NO: 165)
QAP;	(SEQ ID NO: 166)
QAPR;	(SEQ ID NO: 167)
QAPE;	(SEQ ID NO: 168)
APK;	(SEQ ID NO: 169)
QAPG;	(SEQ ID NO: 170)
QAIK;	
and	
QA.	

70. Fc binding polypeptide according to any preceding item, comprising a sequence selected from the group consisting of:

ADNKFNKQQNAFYIILHLPNLT-[Sequence A]-QAPK;	(SEQ ID NO: 170)
---	------------------

-continued

(SEQ ID NO: 171)
ADNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 172)
VDAKFDKEAQEAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 173)
VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 174)
VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 175)
PAAKHDKDQQSAFYIELHLPNLN-[Sequence A]-QAPP;

(SEQ ID NO: 176)
ADNRFNREQQNAFYIELHLPNLN-[Sequence A]-QAPR;

(SEQ ID NO: 177)
VDAKFDKEAQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 178)
ADNAFNTEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 179)
VDAKFDKETQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 180)
ADNKFNKEQQNAFYIELHLPNGN-[Sequence A]-QAPKA;

(SEQ ID NO: 181)
ADNKFNKEHQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 182)
FNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 183)
VDAKFDKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 184)
VDAKFDKEQKAFYIELKLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 185)
ADNKFNKEQHNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 186)
VDNKFNKEAQNAFYIKLHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 187)
VDAKFDKEQQEAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 188)
VDAKFDKEQQKAFYIELKLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 189)
ADNKFHKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 190)
ADNKFNKEQQNAFYHILHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 191)
ADNKFNKEHQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 192)
ADNKHNKEHHHALYHILHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 193)
KFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK;

(SEQ ID NO: 194)
ADSKFDEAQQSAFYIELHLPNLT-[Sequence A]-QAPP;

and

(SEQ ID NO: 195)
IDAKFDEAQQAIFYIELHLPNLN-[Sequence A]-QAPP,

wherein [Sequence A] is as defined in any one of items 2-56.

71. Fc binding polypeptide according to any preceding item, comprising a sequence selected from the group consisting of:

x)
(SEQ ID NO: 172)
VDAKFDKE AQEAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined in any one of items 2-56;
[0380] xi) an amino acid sequence which has at least 86% identity to the sequence defined in x),

xvi)
(SEQ ID NO: 173)
VDNKFNKEQQNAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined in any one of items 2-56;
and

[0381] xvii) an amino acid sequence which has at least 86% identity to the sequence defined in xvi).

72. Fc binding polypeptide according to any preceding item, comprising a sequence selected from the group consisting of:

x)
(SEQ ID NO: 172)
VDAKFDKE AQEAFYIELHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined in any one of items 2-56;
and

[0382] xi) an amino acid sequence which has at least 86% identity to the sequence defined in x).

73. Fc binding polypeptide according to item 71 or 72, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-57 and 208-264; such as the group consisting of SEQ ID NO:1-56 and 208-263; or such as the group consisting of SEQ ID NO:1-57, such as SEQ ID NO:1-56; or such as the group consisting of 208-264; such as SEQ ID NO:208-263.

74. Fc binding polypeptide according to any one of items 71-73, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-16 and 208-223; such as the group consisting of SEQ ID NO:2-14, 16, 209-221 and 223; or such as the group consisting of SEQ ID NO:1-16, such as SEQ ID NO:2-14 and 16; or such as the group consisting of 208-223; such as SEQ ID NO:209-221 and 223.

75. Fc binding polypeptide according to any one of items 71-74, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12, 14, 209, 210, 211, 212, 213, 214, 215, 218, 219 and 221; such as the group consisting of SEQ ID NO:2, 3, 4, 5, 6, 7, 8, 11, 12 and 14 or the group consisting of SEQ ID NO:209, 210, 211, 212, 213, 214, 215, 218, 219 and 221.

76. Fc binding polypeptide according to any one of items 71-75, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 11, 12, 209, 211, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:2, 4, 6, 7, 11 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214, 218 and 219 or wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:2, 4, 6, 7, 12, 209, 211, 213, 214, 218 and 219; such as

the group consisting of SEQ ID NO:2, 4, 6, 7 and 12 or the group consisting of SEQ ID NO:209, 211, 213, 214 and 219.

77. Fc binding polypeptide according to any one of items 71-76, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:6, 7, 11 and 12 or the group consisting of SEQ ID NO:213, 214, 218 and 219 or wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 12, 213, 214 and 219; such as the group consisting of SEQ ID NO:6, 7 and 12 or the group consisting of SEQ ID NO:213, 214 and 219.

78. Fc binding polypeptide according to any one of items 71-77, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218; such as the group consisting of SEQ ID NO:6, 7 and 11 or the group consisting of SEQ ID NO:213, 214 and 218 or wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:6, 7, 213 and 214; such as the group consisting of SEQ ID NO:6 and 7 or the group consisting of SEQ ID NO:213 and 214.

79. Fc binding polypeptide according to any one of items 71-78, wherein sequence x) or xvi) corresponds to the sequence SEQ ID NO:7 or SEQ ID NO:214.

80. Fc binding polypeptide according to any one of items 1-70, comprising a sequence selected from the group consisting of:

xii) (SEQ ID NO: 170)
ADNKFNKEQQNAFYEILHLPNLT-[Sequence A]-QAPK

wherein [Sequence A] is as defined in any one of items 2-56; and

[0383] xiii) an amino acid sequence which has at least 86% identity to the sequence defined in xii).

81. Fc binding polypeptide according to any one of items 1-70, comprising a sequence selected from the group consisting of:

xiv) (SEQ ID NO: 171)
ADNKFNKEQQNAFYEILHLPNLN-[Sequence A]-QAPK

wherein [Sequence A] is as defined in any one of items 2-56; and

[0384] xv) an amino acid sequence which has at least 86% identity to the sequence defined in xiv).

82. Fc binding polypeptide multimer, wherein each monomer of the multimer comprises a Fc binding polypeptide which is independently selected from any Fc binding polypeptide defined in any one of items 1-81.

83. Fc binding polypeptide multimer according to item 82, wherein each monomer of the multimer comprises a Sequence A which is independently selected from any Sequence A defined in any one of items 2-56.

84. Fc binding polypeptide multimer according to item 82 or 83, wherein said multimer is selected from the group consisting of dimer, trimer, tetramer, pentamer, hexamer, heptamer, octamer, nonamer and decamer.

85. Fc binding polypeptide multimer according to any one of items 82-84, wherein said multimer is selected from the

group consisting of tetramer, pentamer, hexamer, heptamer and octamer; such as the group consisting of pentamer, hexamer and heptamer.

86. Fc binding polypeptide multimer according to any one of items 82-85, wherein said multimer is a hexamer.

87. Fc binding polypeptide multimer according to any one of items 82-86, wherein said multimer is a homomer.

88. Fc binding polypeptide multimer according to any one of items 82-86, wherein said multimer is a heteromer comprising at least two or at least three different Fc binding polypeptides each defined as in any one of items 1-81.

89. Fc binding polypeptide multimer according to any one of items 82-86 and 88, wherein said multimer is a heteromer comprising at least two or at least three different sequence A, each defined as in any one of items 2-56

90. Fc binding polypeptide multimer according to any one of items 82-89, further comprising at least one linker.

91. Fc binding polypeptide multimer according to item 90, wherein said linker comprises up to 15 amino acid residues.

92. Fc binding polypeptide according to any one of item 1-81 or a Fc binding polypeptide multimer according to any one of items 82-91 which comprises additional amino acids at the C-terminal and/or N-terminal end.

93. Fc binding polypeptide or Fc binding polypeptide multimer according to item 92, wherein said additional amino acid(s) improve(s) production, purification, stabilization in vitro or coupling of the polypeptide.

94. Fc binding polypeptide or Fc binding polypeptide multimer according to item 93, wherein said additional amino acid(s) improve(s) coupling of the polypeptide and are selected from the group consisting of one or more cysteine residues, a plurality of lysine residues and a plurality of histidine residues.

95. Fc binding polypeptide multimer according to any one of items 82-94, wherein said Fc binding polypeptide monomers are expressed as a fusion protein.

96. Fc binding polypeptide multimer according to any one of items 82-94, wherein said Fc binding polypeptide monomer units are covalently coupled together.

97. A polynucleotide encoding an Fc binding polypeptide according to any one of items 1-81 and 92-94 or a Fc binding polypeptide multimer according to any one of items 82-95.

98. Expression vector comprising a polynucleotide according to item 97.

99. Host cell comprising an expression vector according to item 98.

100. Method of producing an Fc binding polypeptide according to any one of items 1-81 and 92-94 or an Fc binding polypeptide multimer according to any one of items 82-95, comprising

[0385] culturing a host cell according to item 99 under conditions permissive of expression of said polypeptide or multimer from said expression vector, and

[0386] isolating said polypeptide or multimer.

101. Fc binding polypeptide according to any one of items 1-81 and 92-94 or Fc binding polypeptide multimer according to any one of items 82-96, which is capable of binding to Fc such that the K_D value of the interaction is at most 1×10^{-7} M, such as at most 1×10^{-8} M, such as at most 1×10^{-9} M, such as at most 1×10^{-10} M, such as at most 1×10^{-11} M.

102. Fc binding polypeptide according to any one of items 1-82 and 92-94 and 101 or Fc binding polypeptide multimer according to any one of items 82-96 and 101, which is capable of binding to Fc such as that the K_D value of the

104. Fc binding polypeptide according to any one of items 1-82 and 92-94 and 101-103 or Fc binding polypeptide multimer according to any one of items 82-96 and 102-103, which exhibits has an alkali stability of at least 60%, such as at least 70%, such as at least 80%, such as at least 85%, such as at least 89%, such as at least 90%, such as at least 95%, such as at least 98%, such as at least 99% of the alkali stability of SEQ ID NO:58 and/or SEQ ID NO:59 after incubation in 0.5 M NaOH.

106. Separation matrix according to item 105 wherein said Fc binding polypeptide multimer is a multimer according to any one of items 82-96 and 101-104.

108. Separation matrix according to any one of items 105-107 wherein said Fc binding polypeptide comprises a sequence selected from the group consisting of SEQ ID NO:1-57 and 208-264, such as the group consisting of SEQ ID NO:1-56 and 208-263; such as the group consisting of SEQ ID NO:1-16 and 208-223; such as the group consisting of SEQ ID NO:6, 7, 11, 12, 213, 214, 218 and 219; such as the group consisting of SEQ ID NO:6, 7, 11, 213, 214 and 218, such as the group consisting of SEQ ID NO:7 and 214;

[0388] SEQ ID NO:1-57, such as selected from the group consisting of SEQ ID NO:1-56, such as the group consisting of SEQ ID NO:1-16, such as selected from the group consisting of SEQ ID NO:6, 7, 11 and 12; such as selected from the group consisting of SEQ ID NO:6, 7 and 11; such as SEQ ID NO:7.

[0390] SEQ ID NO:1-57; such as selected from the group consisting of SEQ ID NO:1-56; such as the group consisting of SEQ ID NO:1-16; such as selected from the group consisting of SEQ ID NO:6, 7 and 12; such as selected from the group consisting of SEQ ID NO:6 and 7; such as SEQ ID NO:7.

111. Separation matrix according to item 110 wherein said polymer is a polysaccharide, preferably a polysaccharide selected from dextran, starch, cellulose, pullulan, agar, and agarose.

112. Separation matrix according to any one of items 105-111 wherein said solid support is in fibrous, beaded or particle form.

113. Separation matrix according to any one of items 105-112, wherein said Fc binding polypeptide is coupled to the solid support by covalent or non-covalent binding.

114. A method of isolating an immunoglobulin comprising
[0391] a) contacting a liquid sample comprising said
immunoglobulin with a separation matrix according to
any one of items 105-113.

115. Method according to item 114 further comprising

[0392] b) washing said separation matrix with a washing liquid, and

[0393] c) eluting the immunoglobulin from the separation matrix with an elution liquid, and optionally

[0394] d) cleaning the separation matrix with a cleaning liquid.

SEQUENCE LISTING

Sequence total quantity: 305

```

SEQ ID NO: 1          moltype = AA  length = 58
FEATURE               Location/Qualifiers
REGION                1..58
                        note = Engineered Fc binding polypeptide
source                1..58
                        mol_type = protein
                        organism = synthetic construct

```

SEQUENCE: 1
VDAKFDKEAO EAFYEILHLP NLTEEORNAF IOGLKDDPSG SKAILAEAKK SNDAOAPK 58

```

SEQ ID NO: 2      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION            1..58
                  note = Engineered Fc binding polypeptide
source            1..58
                  mol_type = protein
                  organism = synthetic construct

```

-continued

SEQUENCE: 2
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQGLKDDPSG SKAILAEAKK INDAQAPK 58

SEQ ID NO: 3 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 3
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQSLKDDPSR SKAILAEAKK VNDAQAPK 58

SEQ ID NO: 4 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 4
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQGLKDDPSR SKAILAEAKK LNDAQAPK 58

SEQ ID NO: 5 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 5
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQQLKDDPSR SKAILAEAKK INDAQAPK 58

SEQ ID NO: 6 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 6
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQALKDDPSR SKAILAEAKK INDAQAPK 58

SEQ ID NO: 7 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 7
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQSLKDDPSG SKAILAEAKK VNDAQAPK 58

SEQ ID NO: 8 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 8
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQTLKDDPSE SKAILAEAKK RNDAQAPK 58

SEQ ID NO: 9 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 9
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQGLKDDPSE SKAILAEAKK RNDAQAPK 58

SEQ ID NO: 10 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58

-continued

source	note = Engineered Fc binding polypeptide 1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 10		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQALKDDPSD SKAILAEAKK LNDAQAPK	58
SEQ ID NO: 11	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 11		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQALKDDPSG SKAILAEAKK RNDAQAPK	58
SEQ ID NO: 12	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 12		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQELKDDPSG SKAILAEAKK LNDAQAPK	58
SEQ ID NO: 13	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 13		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQHLKDDPSK SKAILAEAKK LNDAQAPK	58
SEQ ID NO: 14	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 14		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQGLKDDPSD SKAILAEAKK VNDAQAPK	58
SEQ ID NO: 15	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 15		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQGLKDDPSQ SKAILAEAKK IND AQAPK	58
SEQ ID NO: 16	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 16		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQQLKDDPSD SKAILAEAKK IND AQAPK	58
SEQ ID NO: 17	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 17		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQALKDDPSK SKAILAEAKK LNDAQAPK	58

-continued

```

SEQ ID NO: 18      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 18
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQALKDDPSR SKAILAEAKK LNDQAQPK 58

SEQ ID NO: 19      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 19
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQDLKDDPSK SKAILAEAKK INDAQAPK 58

SEQ ID NO: 20      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 20
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQDLKDDPSN SKAILAEAKK SNDAQAPK 58

SEQ ID NO: 21      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 21
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQDLKDDPSR SKAILAEAKK VNDAQAPK 58

SEQ ID NO: 22      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 22
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQELKDDPSR SKAILAEAKK LNDQAQPK 58

SEQ ID NO: 23      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 23
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQGLKDDPSD SKAILAEAKK RNDAQAPK 58

SEQ ID NO: 24      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 24
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQGLKDDPSN SKAILAEAKK INDAQAPK 58

SEQ ID NO: 25      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION           1..58
                  note = Engineered Fc binding polypeptide
source           1..58

```

-continued

	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 25		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQGLKDDPSR SKAILAEAKK INDAQAPK	58
SEQ ID NO: 26	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 26		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQGLKDDPSR SKAILAEAKK RNDAQAPK	58
SEQ ID NO: 27	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 27		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQHLKDDPSK SKAILAEAKK INDAQAPK	58
SEQ ID NO: 28	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 28		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQHLKDDPSQ SKAILAEAKK SNDAQAPK	58
SEQ ID NO: 29	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 29		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQHLKDDPSR SKAILAEAKK INDAQAPK	58
SEQ ID NO: 30	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 30		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQHLKDDPSS SKAILAEAKK RNDAQAPK	58
SEQ ID NO: 31	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 31		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQHLKDDPSS SKAILAEAKK VNDAQAPK	58
SEQ ID NO: 32	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 32		
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQKLKDDPSE SKAILAEAKK INDAQAPK	58
SEQ ID NO: 33	moltype = AA length = 58	

-continued

FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 33		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQKLKDDPSK SKAILAEAKK LNDQAQPK		58
SEQ ID NO: 34	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 34		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQKLKDDPSR SKAILAEAKK INDAQAPK		58
SEQ ID NO: 35	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 35		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQNLKDDPSH SKAILAEAKK SNDAQAPK		58
SEQ ID NO: 36	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 36		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSD SKAILAEAKK RNDAQAPK		58
SEQ ID NO: 37	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 37		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSH SKAILAEAKK RNDAQAPK		58
SEQ ID NO: 38	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 38		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSH SKAILAEAKK SNDAQAPK		58
SEQ ID NO: 39	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 39		
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSK SKAILAEAKK LNDQAQPK		58
SEQ ID NO: 40	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 40
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSK SKAILAEAKK SNDAQAPK 58

SEQ ID NO: 41 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 41
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSQ SKAILAEAKK RNDAQAPK 58

SEQ ID NO: 42 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 42
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQPLKDDPSS SKAILAEAKK INDAQAPK 58

SEQ ID NO: 43 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 43
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQQLKDDPSD SKAILAEAKK VNDAQAPK 58

SEQ ID NO: 44 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 44
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQQLKDDPSN SKAILAEAKK SNDAQAPK 58

SEQ ID NO: 45 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 45
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQQLKDDPSQ SKAILAEAKK INDAQAPK 58

SEQ ID NO: 46 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 46
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQQLKDDPSR SKAILAEAKK VNDAQAPK 58

SEQ ID NO: 47 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 47
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQRLKDDPSQ SKAILAEAKK RNDAQAPK 58

SEQ ID NO: 48 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58

-continued

source	note = Engineered Fc binding polypeptide 1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 48		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQRLKDDPSQ SKAILAEAKK SNDAQAPK	58
SEQ ID NO: 49	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 49		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQRLKDDPSR SKAILAEAKK INDAQAPK	58
SEQ ID NO: 50	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 50		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQRLKDDPSR SKAILAEAKK LNDQAAPK	58
SEQ ID NO: 51	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 51		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQSLKDDPSR SKAILAEAKK RNDAQAPK	58
SEQ ID NO: 52	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 52		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQSLKDDPSR SKAILAEAKK SNDAQAPK	58
SEQ ID NO: 53	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 53		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQTLKDDPSK SKAILAEAKK INDAQAPK	58
SEQ ID NO: 54	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 54		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQTLKDDPSQ SKAILAEAKK INDAQAPK	58
SEQ ID NO: 55	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
REGION	1..58	
	note = Engineered Fc binding polypeptide	
source	1..58 mol_type = protein organism = synthetic construct	
SEQUENCE: 55		
VDAKFDKEAQ EAFYEILHLP	NLTTEEQRNAF IQTLKDDPSR SKAILAEAKK RNDAQAPK	58

-continued

```

SEQ ID NO: 56      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION            1..58
                  note = Engineered Fc binding polypeptide
source            1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 56
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQTLKDDPSS SKAILAEAKK RNDAQAPK    58

SEQ ID NO: 57      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION            1..58
                  note = Engineered Fc binding polypeptide
source            1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 57
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQTLKDDPSK SKAILAEAKK GNDAQAPK    58

SEQ ID NO: 58      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION            1..58
                  note = Engineered Fc binding polypeptide
source            1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 58
VDAKFDKEAQ EAFYEILHLP NLTEEQRNAF IQSLKDDPSV SKAILAEAKK LNDAQAPK    58

SEQ ID NO: 59      moltype = AA  length = 58
FEATURE           Location/Qualifiers
REGION            1..58
                  note = Engineered Fc binding polypeptide
source            1..58
                  mol_type = protein
                  organism = synthetic construct

SEQUENCE: 59
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQSLKDDPSQ SANLLAEAKK LNDAQAPK    58

SEQ ID NO: 60      moltype = AA  length = 58
FEATURE           Location/Qualifiers
source            1..58
                  mol_type = protein
                  organism = Staphylococcus aureus

SEQUENCE: 60
ADNKFNKEQQ NAFYEILHLP NLTEEQRNGF IQSLKDDPSV SKEILAEAKK LNDAQAPK    58

SEQ ID NO: 61      moltype = AA  length = 58
FEATURE           Location/Qualifiers
source            1..58
                  mol_type = protein
                  organism = Staphylococcus aureus

SEQUENCE: 61
ADNKFNKEQQ NAFYEILHLP NLNEEQRNGF IQSLKDDPSQ SANLLAEAKK LNDAQAPK    58

SEQ ID NO: 62      moltype = AA  length = 58
FEATURE           Location/Qualifiers
source            1..58
                  mol_type = protein
                  organism = Staphylococcus aureus

SEQUENCE: 62
ADNNFNKEQQ NAFYEILHMP NLNEEQRNGF IQSLKDDPSQ SANLLAEAKK LNESQAPK    58

SEQ ID NO: 63      moltype = AA  length = 61
FEATURE           Location/Qualifiers
source            1..61
                  mol_type = protein
                  organism = Staphylococcus aureus

SEQUENCE: 63
ADAQQNKFNK DQSAFYEIL NMPNLNEEQR NGFIQSLKDD PSQSTNVLGE AKKLNESQAP    60
K                                                    61

SEQ ID NO: 64      moltype = AA  length = 56
FEATURE           Location/Qualifiers

```

-continued

```

source                1..56
                      mol_type = protein
                      organism = Staphylococcus aureus

SEQUENCE: 64
AQHDEAQQNA FYQVLNMPNL NADQRNGFIQ SLKDDPSQSA NVLGEAQKLN DSQAPK      56

SEQ ID NO: 65         moltype = AA length = 31
FEATURE              Location/Qualifiers
REGION              1..31
                      note = Engineered Fc binding polypeptide
SITE                2
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              4..7
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              9..10
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              12..14
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                17
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              19..21
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                23
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              26..28
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              30..31
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid

source                1..31
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 65
EXQXXXIXX LXXXPSXSXX XLKEAXXXNX X      31

SEQ ID NO: 66         moltype = AA length = 16
FEATURE              Location/Qualifiers
REGION              1..16
                      note = Engineered Fc binding polypeptide
REGION              1..4
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                8
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              10..12
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                14
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                16
                      note = misc_feature - Xaa can be any naturally occurring
                      amino acid

source                1..16
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 66
XXXXAFYXIX XXPXLX      16

SEQ ID NO: 67         moltype = AA length = 11
FEATURE              Location/Qualifiers
REGION              1..11
                      note = Engineered Fc binding polypeptide

source                1..11
                      mol_type = protein
                      organism = synthetic construct

SEQUENCE: 67

```

-continued

EQQNAFYEIL H	11
SEQ ID NO: 68	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 68	
DQQAAFYEIL H	11
SEQ ID NO: 69	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 69	
EAQEAFYEIL H	11
SEQ ID NO: 70	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 70	
DQQSIFYEIL H	11
SEQ ID NO: 71	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 71	
AQQAAFYEIL H	11
SEQ ID NO: 72	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 72	
EQQNAFYEIL N	11
SEQ ID NO: 73	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 73	
EQQAAFYEIL H	11
SEQ ID NO: 74	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
source	note = Engineered Fc binding polypeptide
	1..11
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 74	
AQQSIFYEIL H	11
SEQ ID NO: 75	moltype = AA length = 11
FEATURE	Location/Qualifiers
REGION	1..11
	note = Engineered Fc binding polypeptide

-continued

source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 75 EQQQAFYEIL H		11
SEQ ID NO: 76 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 76 EHQNAFYEIL H		11
SEQ ID NO: 77 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 77 EAQNAFYEIL H		11
SEQ ID NO: 78 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 78 ETQNAFYEIL H		11
SEQ ID NO: 79 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 79 EQQSAFYEIL H		11
SEQ ID NO: 80 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 80 EAQNAFYKIL H		11
SEQ ID NO: 81 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 81 EQQEAFYEIL H		11
SEQ ID NO: 82 FEATURE REGION	moltype = AA length = 11 Location/Qualifiers 1..11 note = Engineered Fc binding polypeptide	
source	1..11 mol_type = protein organism = synthetic construct	
SEQUENCE: 82 EAQKAFYEIL K		11

-continued

SEQ ID NO: 83	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 83		
EQHNAFYEIL H		11
SEQ ID NO: 84	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 84		
EHHHALYHIL H		11
SEQ ID NO: 85	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 85		
EQQKAFYAIL H		11
SEQ ID NO: 86	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 86		
DQQSAFYEIL N		11
SEQ ID NO: 87	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 87		
AQQNAFYQVL N		11
SEQ ID NO: 88	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 88		
EQQKAFYEIL H		11
SEQ ID NO: 89	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 89		
EQQKAFYEIL K		11
SEQ ID NO: 90	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	

-continued

SEQUENCE: 90	organism = synthetic construct	
EQQNAFYEIL S		11
SEQ ID NO: 91	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 91		
EQQNAFYHIL H		11
SEQ ID NO: 92	moltype = AA length = 11	
FEATURE	Location/Qualifiers	
REGION	1..11	
	note = Engineered Fc binding polypeptide	
source	1..11	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 92		
EQQNAFYHIL N		11
SEQ ID NO: 93	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 93		
LPNLT		5
SEQ ID NO: 94	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 94		
LPNLN		5
SEQ ID NO: 95	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 95		
MPNLN		5
SEQ ID NO: 96	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 96		
LPYLT		5
SEQ ID NO: 97	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 97		
LPSLT		5
SEQ ID NO: 98	moltype = AA length = 5	
FEATURE	Location/Qualifiers	

-continued

REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 98		
MPNLL		5
SEQ ID NO: 99	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 99		
LPNGN		5
SEQ ID NO: 100	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 100		
LPNLS		5
SEQ ID NO: 101	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 101		
GPNAN		5
SEQ ID NO: 102	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 102		
VPNLN		5
SEQ ID NO: 103	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 103		
LPNLR		5
SEQ ID NO: 104	moltype = AA length = 5	
FEATURE	Location/Qualifiers	
REGION	1..5	
	note = Engineered Fc binding polypeptide	
source	1..5	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 104		
LPNLL		5
SEQ ID NO: 105	moltype = AA length = 47	
FEATURE	Location/Qualifiers	
REGION	1..47	
	note = Engineered Fc binding polypeptide	
SITE	18	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23	

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 105	
EQQNAFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX	47
SEQ ID NO: 106	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 106	
EQQNAFYEIL HLPNLNEXQX XXXIXLXXX PSXSXXLXE AXXNXX	47
SEQ ID NO: 107	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid

-continued

REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 107
 DQQAIFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX 47
 SEQ ID NO: 108 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 108
 EAQEAIFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX 47
 SEQ ID NO: 109 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring

-continued

```

REGION          amino acid
                28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source          1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 109
DQQSIFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX          47

SEQ ID NO: 110    moltype = AA  length = 47
FEATURE          Location/Qualifiers
REGION          1..47
                note = Engineered Fc binding polypeptide
SITE            18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source          1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 110
AQQAFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX          47

SEQ ID NO: 111    moltype = AA  length = 47
FEATURE          Location/Qualifiers
REGION          1..47
                note = Engineered Fc binding polypeptide
SITE            18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          28..30

```

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 111		
EQQAIFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX		47
SEQ ID NO: 112	moltype = AA length = 47	
FEATURE	Location/Qualifiers	
REGION	1..47	
	note = Engineered Fc binding polypeptide	
SITE	18	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 112		
AQQAIFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX		47
SEQ ID NO: 113	moltype = AA length = 47	
FEATURE	Location/Qualifiers	
REGION	1..47	
	note = Engineered Fc binding polypeptide	
SITE	18	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30	
	note = misc_feature - Xaa can be any naturally occurring amino acid	

-continued

SITE	33	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47	mol_type = protein organism = synthetic construct
SEQUENCE: 113		
EQQNAFYEIL NMPNLNEXQX	XXXIXXLXXX PSXSXXXLXE AXXXNXX	47
SEQ ID NO: 114		
FEATURE	Location/Qualifiers	
REGION	1..47	note = Engineered Fc binding polypeptide
SITE	18	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47	mol_type = protein organism = synthetic construct
SEQUENCE: 114		
EAQNAFYEIL HLPNLTEXQX	XXXIXXLXXX PSXSXXXLXE AXXXNXX	47
SEQ ID NO: 115		
FEATURE	Location/Qualifiers	
REGION	1..47	note = Engineered Fc binding polypeptide
SITE	18	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33	note = misc_feature - Xaa can be any naturally occurring

-continued

REGION	amino acid 35..37 note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47 note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47 mol_type = protein organism = synthetic construct	
SEQUENCE: 115 EQQQAFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX		47
SEQ ID NO: 116 moltype = AA length = 47		
FEATURE Location/Qualifiers		
REGION	1..47 note = Engineered Fc binding polypeptide	
SITE	18 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30 note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37 note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47 note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47 mol_type = protein organism = synthetic construct	
SEQUENCE: 116 EQQNAFYEIL HLPYLTEXQX XXXIXLXXX PSXSXXLXE AXXNXX		47
SEQ ID NO: 117 moltype = AA length = 47		
FEATURE Location/Qualifiers		
REGION	1..47 note = Engineered Fc binding polypeptide	
SITE	18 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30 note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33 note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37	

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 117	
ETQNAPYEIL HLPNLTEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX	47
SEQ ID NO: 118	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 118	
EAQNAPYKIL HLPNLTEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX	47
SEQ ID NO: 119	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid

-continued

SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 119
 EQQNAFYEIL HLPNGNEXQX XXXIXLXXX PSXSXXLXE AXXNXX 47

SEQ ID NO: 120 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 120
 EHQNIFYEIL HLPNLNEXQX XXXIXLXXX PSXSXXLXE AXXNXX 47

SEQ ID NO: 121 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring

-continued

```

REGION          amino acid
                42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source          1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 121
EQQNAFYEIL NMPNLLEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX          47

SEQ ID NO: 122      moltype = AA length = 47
FEATURE            Location/Qualifiers
REGION            1..47
                note = Engineered Fc binding polypeptide
SITE              18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE              33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE              39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source            1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 122
EQQSIFYEIL HLPNLTEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX          47

SEQ ID NO: 123      moltype = AA length = 47
FEATURE            Location/Qualifiers
REGION            1..47
                note = Engineered Fc binding polypeptide
SITE              18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE              33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE              39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION            42..44

```

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 123	
EQQEAFYEIL HLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXXNXX	47
SEQ ID NO: 124	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	46..47
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..47
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 124	
EAQKAFYEIL KLPNLTEXQX XXXIXLXXX PSXSXXLXE AXXXNXX	47
SEQ ID NO: 125	moltype = AA length = 47
FEATURE	Location/Qualifiers
REGION	1..47
	note = Engineered Fc binding polypeptide
SITE	18
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	20..23
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	25..26
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	28..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	39
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid

-continued

REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 125
 EQHNAFYEIL HLPNLNEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 126 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..47
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 126
 EHHHALYHIL HLPNLNEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 127 moltype = AA length = 47
 FEATURE Location/Qualifiers
 REGION 1..47
 note = Engineered Fc binding polypeptide
 SITE 18
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 39
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring

-continued

```

source          amino acid
                1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 127
AQQSAFYEIL HLPSTLTXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX          47

SEQ ID NO: 128    moltype = AA  length = 47
FEATURE          Location/Qualifiers
REGION           1..47
                note = Engineered Fc binding polypeptide
SITE            18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source          1..47
                mol_type = protein
                organism = synthetic construct

SEQUENCE: 128
EQQKAFYAIL HLPYLTXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX          47

SEQ ID NO: 129    moltype = AA  length = 47
FEATURE          Location/Qualifiers
REGION           1..47
                note = Engineered Fc binding polypeptide
SITE            18
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          20..23
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          25..26
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          28..30
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            33
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          35..37
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
SITE            39
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          42..44
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
REGION          46..47
                note = misc_feature - Xaa can be any naturally occurring
                amino acid
source          1..47

```


-continued

```

mol_type = protein
organism = synthetic construct
SEQUENCE: 129
DQGSAPFYEL NMPNLNEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 130      moltype = AA length = 47
FEATURE            Location/Qualifiers
REGION             1..47
                    note = Engineered Fc binding polypeptide
SITE               18
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             20..23
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             25..26
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             28..30
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
SITE               33
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             35..37
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
SITE               39
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             42..44
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             46..47
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
source             1..47
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 130
AQQNAFYQVL NMPNLNEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 131      moltype = AA length = 47
FEATURE            Location/Qualifiers
REGION             1..47
                    note = Engineered Fc binding polypeptide
SITE               18
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             20..23
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             25..26
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             28..30
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
SITE               33
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             35..37
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
SITE               39
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             42..44
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
REGION             46..47
                    note = misc_feature - Xaa can be any naturally occurring
                    amino acid
source             1..47
                    mol_type = protein
                    organism = synthetic construct

```

-continued

SEQUENCE: 131
EQQKAFYEIL KLPNLTEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 132 moltype = AA length = 47
FEATURE Location/Qualifiers
REGION 1..47
 note = Engineered Fc binding polypeptide
SITE 18
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 39
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
source 1..47
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 132
EQQNAFYEIL SLPNLTEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

SEQ ID NO: 133 moltype = AA length = 47
FEATURE Location/Qualifiers
REGION 1..47
 note = Engineered Fc binding polypeptide
SITE 18
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 20..23
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 25..26
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 28..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 39
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 46..47
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
source 1..47
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 133
EQQNAFYEIL HLPNLSEXQX XXXIXXLXXX PSXSXXXLXE AXXXNXX 47

-continued

```

SEQ ID NO: 134      multype = AA length = 47
FEATURE             Location/Qualifiers
REGION              1..47
                    note = Engineered Fc binding polypeptide
SITE                18
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              20..23
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              25..26
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              28..30
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
SITE                33
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              35..37
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
SITE                39
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              42..44
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
REGION              46..47
                    note = misc_feature - Xaa can be any naturally occurring
                        amino acid
source              1..47
                    mol_type = protein
                    organism = synthetic construct

```

organism = *Synscolecus consociatus*

SEQUENCE: 134
EQONAFYHIL HLPNLNEXQX XXXIXXLXXX PSXSLXXLXE AXXXXNXX 47

```

SEQ ID NO: 135      multype = AA   length = 47
FEATURE             Location/Qualifiers
REGION              1..47
                    note = Engineered Fc binding polypeptide
SITE                18
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              20..23
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              25..26
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              28..30
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                33
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              35..37
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
SITE                39
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              42..44
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
REGION              46..47
                    note = misc_feature - Xaa can be any naturally occurring
                      amino acid
source              1..47
                    mol_type = protein
                    organism = synthetic construct

```

SEQUENCE: 135
EHONAFYEIL HLPNLTEXOX XXXIXXLXXX PSXSLXXLXE AXXXXNXX 47

SEQ ID NO: 136 moltype = AA length = 47

-continued

FEATURE	Location/Qualifiers	
REGION	1..47	
	note = Engineered Fc binding polypeptide	
SITE	18	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 136		
EQQAIFYEIL HLPNLNEXQX XXXIXLXXX PSXXXXLXE AXXXNXX		47
SEQ ID NO: 137	moltype = AA length = 47	
FEATURE	Location/Qualifiers	
REGION	1..47	
	note = Engineered Fc binding polypeptide	
SITE	18	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	20..23	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	25..26	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	28..30	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	33	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	35..37	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
SITE	39	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	42..44	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
REGION	46..47	
	note = misc_feature - Xaa can be any naturally occurring amino acid	
source	1..47	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 137		
AQQAIFYEIL HLPNLTXXQX XXXIXLXXX PSXXXXLXE AXXXNXX		47
SEQ ID NO: 138	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	

-continued

source	note = Engineered Fc binding polypeptide 1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 138 ADNKFNK		7
SEQ ID NO: 139 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 139 VDAKFDK		7
SEQ ID NO: 140 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 140 VDNKFNK		7
SEQ ID NO: 141 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 141 IAAKHDK		7
SEQ ID NO: 142 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 142 IAAQHDK		7
SEQ ID NO: 143 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 143 ADNNFNK		7
SEQ ID NO: 144 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 144 IAAKFDE		7
SEQ ID NO: 145 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Engineered Fc binding polypeptide	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 145 PAAKHDK		7

-continued

SEQ ID NO: 146	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 146		
ADNAFNT		7
SEQ ID NO: 147	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 147		
ADNRFNE		7
SEQ ID NO: 148	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 148		
IDSKFDE		7
SEQ ID NO: 149	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 149		
ADNRFNR		7
SEQ ID NO: 150	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 150		
ADNKHNK		7
SEQ ID NO: 151	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 151		
ADSKFDE		7
SEQ ID NO: 152	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 152		
IDAKHDE		7
SEQ ID NO: 153	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	

-continued

SEQUENCE: 153	mol_type = protein	
QQNKFNK	organism = synthetic construct	7
SEQ ID NO: 154	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Engineered Fc binding polypeptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 154		7
ADNKFHK		
SEQ ID NO: 155	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
source	note = Engineered Fc binding polypeptide	
	1..4	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 155		4
KFNK		
SEQ ID NO: 156	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Engineered Fc binding polypeptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 156		7
ADNNFNR		
SEQ ID NO: 157	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Engineered Fc binding polypeptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 157		7
AAAKHDK		
SEQ ID NO: 158	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Engineered Fc binding polypeptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 158		7
IDNKFNK		
SEQ ID NO: 159	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
source	note = Engineered Fc binding polypeptide	
	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 159		7
IDAKFDE		
SEQ ID NO: 160	moltype = AA length = 6	
FEATURE	Location/Qualifiers	
REGION	1..6	
source	note = Engineered Fc binding polypeptide	
	1..6	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 160		6
DNNFNK		
SEQ ID NO: 161	moltype = AA length = 7	

-continued

FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 161		
ADNKFNE		7
SEQ ID NO: 162	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 162		
AAAQHDK		7
SEQ ID NO: 163	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = Engineered Fc binding polypeptide	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 163		
AAAKFDE		7
SEQ ID NO: 164	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
	note = Engineered Fc binding polypeptide	
source	1..4	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 164		
QAPK		4
SEQ ID NO: 165	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
	note = Engineered Fc binding polypeptide	
source	1..4	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 165		
QAPP		4
SEQ ID NO: 166	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
	note = Engineered Fc binding polypeptide	
source	1..4	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 166		
QAPR		4
SEQ ID NO: 167	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
	note = Engineered Fc binding polypeptide	
source	1..4	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 167		
QAPE		4
SEQ ID NO: 168	moltype = AA length = 4	
FEATURE	Location/Qualifiers	
REGION	1..4	
	note = Engineered Fc binding polypeptide	
source	1..4	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 168
QAPG 4

SEQ ID NO: 169 moltype = AA length = 4
FEATURE Location/Qualifiers
REGION 1..4
note = Engineered Fc binding polypeptide
source 1..4
mol_type = protein
organism = synthetic construct

SEQUENCE: 169
QAIK 4

SEQ ID NO: 170 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid
source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 170
ADNKFNKEQQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 171 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51

-continued

note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid
source 1..58
mol_type = protein
organism = synthetic construct
SEQUENCE: 171
ADNKFNKEQQ NAFYEILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58
SEQ ID NO: 172 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid
source 1..58
mol_type = protein
organism = synthetic construct
SEQUENCE: 172
VDAKFDKEAQ EAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58
SEQ ID NO: 173 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid

-continued

REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..58
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 173
 VDNKFNKEQQ NAFYEILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 174 moltype = AA length = 58
 FEATURE Location/Qualifiers
 REGION 1..58
 note = Engineered Fc binding polypeptide
 SITE 25
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 46
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring amino acid
 source 1..58
 mol_type = protein
 organism = synthetic construct
 SEQUENCE: 174
 VDNKFNKEQQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 175 moltype = AA length = 58
 FEATURE Location/Qualifiers
 REGION 1..58
 note = Engineered Fc binding polypeptide
 SITE 25
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 46
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring amino acid
 REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring

-continued

source	amino acid 1..58 mol_type = protein organism = synthetic construct
SEQUENCE: 175	
PAAKHKDKDQQ SAFYEILHLP NLTEXQXXXX IXXLXXXPSX SXXXLXEAXX XNXXQAPP	58
SEQ ID NO: 176	moltype = AA length = 58
FEATURE	Location/Qualifiers
REGION	1..58 note = Engineered Fc binding polypeptide
SITE	25 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37 note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44 note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54 note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58 mol_type = protein organism = synthetic construct
SEQUENCE: 176	
ADNRFNREQQ NAFYEILHLP NLTEXQXXXX IXXLXXXPSX SXXXLXEAXX XNXXQAPR	58
SEQ ID NO: 177	moltype = AA length = 58
FEATURE	Location/Qualifiers
REGION	1..58 note = Engineered Fc binding polypeptide
SITE	25 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37 note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44 note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51 note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54 note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58

-continued

mol_type = protein
organism = synthetic construct

SEQUENCE: 177
VDAKFDKEAQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 178 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid
source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 178
ADNAPNTEQQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 179 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid
source 1..58
mol_type = protein
organism = synthetic construct

-continued

SEQUENCE: 179
VDAKFDKETQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 180 moltype = AA length = 59
FEATURE Location/Qualifiers
REGION 1..59
 note = Engineered Fc binding polypeptide
SITE 25
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 46
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
source 1..59
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 180
ADNKFNKEQQ NAFYEILHLP NGNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPKA 59

SEQ ID NO: 181 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
 note = Engineered Fc binding polypeptide
SITE 25
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
SITE 46
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 181
ADNKFNKEHQ NAFYEILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

-continued

SEQ ID NO: 182 moltype = AA length = 54
 FEATURE Location/Qualifiers
 REGION 1..54
 note = Engineered Fc binding polypeptide
 SITE 21
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 23..26
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 28..29
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 31..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 36
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 38..40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 42
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 45..47
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 49..50
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 source 1..54
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 182
 FNKEQQNAFY EILHLPNLNE QQXXXXIXXL XXXPSXSSXX LXEAXXNXX QAPK 54

SEQ ID NO: 183 moltype = AA length = 58
 FEATURE Location/Qualifiers
 REGION 1..58
 note = Engineered Fc binding polypeptide
 SITE 25
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 46
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 183
 VDAKFDKEQQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 184 moltype = AA length = 58

-continued

FEATURE	Location/Qualifiers
REGION	1..58
	note = Engineered Fc binding polypeptide
SITE	25
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 184	
VDAKFDKEAQ KAFYEILKLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58	
SEQ ID NO: 185 moltype = AA length = 58	
FEATURE	Location/Qualifiers
REGION	1..58
	note = Engineered Fc binding polypeptide
SITE	25
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 185	
ADNKFNKEQH NAFYEILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58	
SEQ ID NO: 186 moltype = AA length = 58	
FEATURE	Location/Qualifiers
REGION	1..58

-continued

SITE note = Engineered Fc binding polypeptide
 25
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 46
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 186

VDNKFNKEAQ NAFYKILHLP NLTEXQXXXX IXXLXXXP SX XLXEAXX XNXXQAPK 58

SEQ ID NO: 187 moltype = AA length = 58
 FEATURE Location/Qualifiers
 REGION 1..58
 note = Engineered Fc binding polypeptide
 SITE 25
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 27..30
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 32..33
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 35..37
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 42..44
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 SITE 46
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 49..51
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 REGION 53..54
 note = misc_feature - Xaa can be any naturally occurring
 amino acid
 source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 187

VDAKFDKEQQ EAFYEILHLP NLTEXQXXXX IXXLXXXP SX XLXEAXX XNXXQAPK 58

SEQ ID NO: 188 moltype = AA length = 58
 FEATURE Location/Qualifiers
 REGION 1..58
 note = Engineered Fc binding polypeptide
 SITE 25

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 188	
VDAKFDKEQQ KAFYEILKLP NLTEXQXXXX IXLXXXXPSX SXXXLXEAXX XNXXQAPK	58
SEQ ID NO: 189	moltype = AA length = 58
FEATURE	Location/Qualifiers
REGION	1..58
	note = Engineered Fc binding polypeptide
SITE	25
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	27..30
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	32..33
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	35..37
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	40
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	42..44
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	46
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	49..51
	note = misc_feature - Xaa can be any naturally occurring amino acid
REGION	53..54
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 189	
ADNKFHKEQQ NAFYEILHLP NLNEXQXXXX IXLXXXXPSX SXXXLXEAXX XNXXQAPK	58
SEQ ID NO: 190	moltype = AA length = 58
FEATURE	Location/Qualifiers
REGION	1..58
	note = Engineered Fc binding polypeptide
SITE	25
	note = misc_feature - Xaa can be any naturally occurring amino acid

-continued

REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid

SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid

SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid

source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 190

ADNKFNKEQQ NAFYHILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 191 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide

SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 27..30
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 32..33
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 35..37
note = misc_feature - Xaa can be any naturally occurring amino acid

SITE 40
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 42..44
note = misc_feature - Xaa can be any naturally occurring amino acid

SITE 46
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 49..51
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 53..54
note = misc_feature - Xaa can be any naturally occurring amino acid

source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 191

ADNKFNKEHQ NAFYEILHLP NLTEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK 58

SEQ ID NO: 192 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide

SITE 25
note = misc_feature - Xaa can be any naturally occurring amino acid

REGION 27..30
note = misc_feature - Xaa can be any naturally occurring

-continued

```

REGION          amino acid
                 32..33
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION          35..37
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
SITE            40
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION          42..44
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
SITE            46
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION          49..51
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION          53..54
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
source          1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 192
ADNKHNKEHH HALYHILHLP NLNEXQXXXX IXXLXXPSX SXXXLXEAXX XNXXQAPK      58

SEQ ID NO: 193      moltype = AA length = 55
FEATURE            Location/Qualifiers
REGION            1..55
note = Engineered Fc binding polypeptide
SITE              22
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            24..27
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            29..30
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            32..34
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
SITE              37
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            39..41
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
SITE              43
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            46..48
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            50..51
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
source            1..55
mol_type = protein
organism = synthetic construct

SEQUENCE: 193
KFNKEQQNAF YEILHLPNLN EXQXXXXIXX LXXPSXSXX XLXEAXXNX XQAPK      55

SEQ ID NO: 194      moltype = AA length = 58
FEATURE            Location/Qualifiers
REGION            1..58
note = Engineered Fc binding polypeptide
SITE              25
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            27..30
note = misc_feature - Xaa can be any naturally occurring
                 amino acid
REGION            32..33

```

-continued

```

note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring
amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring
amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring
amino acid
source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 194
ADSKFDEAQQ SAFYEILHLP SLTEXQXXXX IXXLXXXPSX SXXXLXEAXX XNXXQAPP 58

SEQ ID NO: 195 moltype = AA length = 58
FEATURE Location/Qualifiers
REGION 1..58
note = Engineered Fc binding polypeptide
SITE 25
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 27..30
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 32..33
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 35..37
note = misc_feature - Xaa can be any naturally occurring
amino acid
SITE 40
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 42..44
note = misc_feature - Xaa can be any naturally occurring
amino acid
SITE 46
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 49..51
note = misc_feature - Xaa can be any naturally occurring
amino acid
REGION 53..54
note = misc_feature - Xaa can be any naturally occurring
amino acid
source 1..58
mol_type = protein
organism = synthetic construct

SEQUENCE: 195
IDAKFDEAQQ AAFYEILHLP NLTEXQXXXX IXXLXXXPSX SXXXLXEAXX XNXXQAPP 58

SEQ ID NO: 196 moltype = AA length = 214
FEATURE Location/Qualifiers
REGION 1..214
note = Humanized mus musculus antibody fragment
source 1..214
mol_type = protein
organism = synthetic construct

SEQUENCE: 196
DIQMTQSPSS LSASVGDVRT ITCRASQDVN TAVAWYQQKP GKAPKLLIYS ASFLYSGVPS 60
RFSGSRSGTD FTLTISSLQP EDFATYICQQ HYTPPTFGQ GTKVEIKRTV AAPSVFIFPP 120
SDEQLKSGTA SVVCLLNIFY PREAKVQWKV DNALQSGNSQ ESVTEQDSKD STYLSSTLT 180
LSKADYEKHK VYACEVTHQG LSSPVTKSFN RGEC 214

```

-continued

SEQ ID NO: 197 moltype = AA length = 450
 FEATURE Location/Qualifiers
 REGION 1..450
 note = Humanized mus musculus antibody fragment
 source 1..450
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 197
 EVQLVESGGG LVQPGGSLRL SCAASGFNIK DTYIHWRQA PGKGLEWVAR IYPTNGYTRY 60
 ADSVKGRFTI SADTSKNTAY LQMNSLRAED TAVYYCSRWG GDGPFYAMDYV GQGTLVTVSS 120
 ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 180
 GLYSLSVVTV VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKTHTCP PCPAPELLGG 240
 PSVFLFPPPK KDTLMISRTPEVTCVVVDVS HEDPEVKFNW YVDGVEVHNA KTKPREEQYN 300
 STYRVSVLT VLHQDWLNGK EYCKKVSNAK LPAPIEKTIS KAKGQPREPQ VYTLPPSREE 360
 MTKNQVSLTC LVKGFYPSDI AVEWESNGQP ENNYKTTTPV LDSGGSFFLY SKLTVDKSRW 420
 QQGNVFSCSV MHEALHNHYT QKSLSLSPGK 450

SEQ ID NO: 198 moltype = AA length = 214
 FEATURE Location/Qualifiers
 REGION 1..214
 note = Humanized mus musculus antibody fragment
 source 1..214
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 198
 DIQMTQSPSS LSASVGDRTV ITCRASQDVN TAVAWYQQKP GKAPKLLIYS ASFLYSGVPS 60
 RFSGSRSGTD FTLTISLQPEDPATYYCQQ HYTTPPTFGQ GTKVEIKRTV AAPSVFIFPP 120
 SDEQLKSGTA SVVCLLNIFY PREAKVQWKV DNALQSGNSQ ESVTEQDSKD STYLSSTLT 180
 LSKADYEKKH VYACEVTHQG LSSPVTKSFN RGEC 214

SEQ ID NO: 199 moltype = AA length = 237
 FEATURE Location/Qualifiers
 REGION 1..237
 note = Humanized mus musculus antibody fragment
 source 1..237
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 199
 EVQLVESGGG LVQPGGSLRL SCAASGFNIK DTYIHWRQA PGKGLEWVAR IYPTNGYTRY 60
 ADSVKGRFTI SADTSKNTAY LQMNSLRAED TAVYYCSRWG GDGPFYAMDYV GQGTLVTVSS 120
 ASTKGPSVFP LAPSSKSTSG GTAALGCLVK DYFPEPVTVS WNSGALTSGV HTFPAVLQSS 180
 GLYSLSVVTV VPSSSLGTQT YICNVNHKPS NTKVDKKVEP KSCDKTHTCP PCPAPEL 237

SEQ ID NO: 200 moltype = AA length = 213
 FEATURE Location/Qualifiers
 REGION 1..213
 note = Humanized mus musculus antibody fragment
 source 1..213
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 200
 LGGSPVFLFP PKPKDTLMIS RTPEVTCVVV DVSHEDPEVK FNWYVDGVEV HNAKTKPREE 60
 QYNSTYRVVS VLTVLHQDWL NGKEYKCKVS NKALPAPIEK TISKAKGQPR EPQVYTLPPS 120
 REEMTKNQVS LTCLVKGFYP SDIAVEWESN GPENNYKTT PPVLDSDGSF FLYSKLTVDK 180
 SRWQQGNVFS CSVMEALHN HYTKSLSLSGK 213

SEQ ID NO: 201 moltype = AA length = 46
 FEATURE Location/Qualifiers
 source 1..46
 mol_type = protein
 organism = Streptococcus sp.

SEQUENCE: 201
 LAEAKVLANR ELDKYGVS DY YKNLINNAKT VEGVKALIDE ILAALP 46

SEQ ID NO: 202 moltype = AA length = 110
 FEATURE Location/Qualifiers
 REGION 1..110
 note = Engineered Fc binding polypeptide
 source 1..110
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 202
 VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQSLKDDPSQ SANLLAEAKK LNDAQAPKAS 60
 GGGSLAEAKV LANRELDKYG VSDYYKNLIN NAKTVEGVKA LIDEILAALP 110

SEQ ID NO: 203 moltype = AA length = 349

-continued

FEATURE Location/Qualifiers
 REGION 1..349
 note = Engineered Fc binding polypeptide
 source 1..349
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 203

VDAKFDKEAQ	EAFYEILHLP	NLTEEQRN	AF	IQALKDDPSR	SKAILAEAKK	INDAQAPKVD	60
AKFDKEAQEA	FYEILHLPLN	L		TEEQQRN	AF	IQ	120
FDKEAQEA	F	Y	E	I	L	H	180
KEAQEA	F	Y	E	I	L	H	240
AQEA	F	Y	E	I	L	H	300
EAFYEILHLP	NLTEEQRN	AF	IQ	ALKDDPSR	SKAILAEAKK	INDAQAPK	349

SEQ ID NO: 204 moltype = AA length = 349
 FEATURE Location/Qualifiers
 REGION 1..349
 note = Engineered Fc binding polypeptide
 source 1..349
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 204

VDAKFDKEAQ	EAFYEILHLP	NLTEEQRN	AF	IQSLKDDPSG	SKAILAEAKK	VND AQAPKVD	60
AKFDKEAQEA	FYEILHLPLN	L		TEEQQRN	AF	IQ	120
FDKEAQEA	F	Y	E	I	L	H	180
KEAQEA	F	Y	E	I	L	H	240
AQEA	F	Y	E	I	L	H	300
EAFYEILHLP	NLTEEQRN	AF	IQ	SLKDDPSG	SKAILAEAKK	VND AQAPK	349

SEQ ID NO: 205 moltype = AA length = 349
 FEATURE Location/Qualifiers
 REGION 1..349
 note = Engineered Fc binding polypeptide
 source 1..349
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 205

VDAKFDKEAQ	EAFYEILHLP	NLTEEQRN	AF	IQELKDDPSG	SKAILAEAKK	LND AQAPKVD	60
AKFDKEAQEA	FYEILHLPLN	L		TEEQQRN	AF	IQ	120
FDKEAQEA	F	Y	E	I	L	H	180
KEAQEA	F	Y	E	I	L	H	240
AQEA	F	Y	E	I	L	H	300
EAFYEILHLP	NLTEEQRN	AF	IQ	ELKDDPSG	SKAILAEAKK	LND AQAPK	349

SEQ ID NO: 206 moltype = AA length = 349
 FEATURE Location/Qualifiers
 REGION 1..349
 note = Engineered Fc binding polypeptide
 source 1..349
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 206

VDAKFDKEAQ	EAFYEILHLP	NLTEEQRN	AF	IQSLKDDPSV	SKAILAEAKK	LND AQAPKVD	60
AKFDKEAQEA	FYEILHLPLN	L		TEEQQRN	AF	IQ	120
FDKEAQEA	F	Y	E	I	L	H	180
KEAQEA	F	Y	E	I	L	H	240
AQEA	F	Y	E	I	L	H	300
EAFYEILHLP	NLTEEQRN	AF	IQ	SLKDDPSV	SKAILAEAKK	LND AQAPK	349

SEQ ID NO: 207 moltype = AA length = 349
 FEATURE Location/Qualifiers
 REGION 1..349
 note = Engineered Fc binding polypeptide
 SITE 33
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 40
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 51
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 91
 note = misc_feature - Xaa can be any naturally occurring amino acid
 SITE 98

-continued

	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	109
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	149
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	156
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	167
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	207
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	214
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	225
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	265
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	272
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	283
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	323
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	330
	note = misc_feature - Xaa can be any naturally occurring amino acid
SITE	341
	note = misc_feature - Xaa can be any naturally occurring amino acid
source	1..349
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 207	
VDAKFDKEAQ EAFYEILHLP	NLTEEQRNAF IQXLKDDPSX SKAILAEAKK XNDAQAPKVD 60
AKFDKEAQEA FYEILHLPNL	TEEQRNAFIQ XLKDDPSXSK AILAEAKKXN DAQAPKVDK 120
FDKEAQEA FYEILHLPNLTE	EQRNAFIQXL KDDPSXSKAI LAEAKKXNDA QAPKVDKFD 180
KEAQEA FYEILHLPNLTEEQ	RNAFIQXLKD DPSXSKAILA EAKKXNDAQA PKVDKFDKE 240
AQEA FYEILHLPNLTEEQRN	AFIQXLKDDP SXSKAILAEA KXNDAQAPK VDAKFDKEAQ 300
EAFYEILHLP NLTEEQRNAF	IQXLKDDPSX SKAILAEAKK XNDAQAPKC 349
SEQ ID NO: 208	
FEATURE	moltype = AA length = 58
source	Location/Qualifiers
	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 208	
VDNKFNKEQQ NAFYEILHLP	NLNTEEQRNFIQ IQGLKDDPSG SANLLAEAKK SNDAQAPK 58
SEQ ID NO: 209	
FEATURE	moltype = AA length = 58
source	Location/Qualifiers
	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 209	
VDNKFNKEQQ NAFYEILHLP	NLNTEEQRNFIQ IQGLKDDPSG SANLLAEAKK INDAQAPK 58
SEQ ID NO: 210	
FEATURE	moltype = AA length = 58
source	Location/Qualifiers
	1..58
	mol_type = protein
	organism = synthetic construct
SEQUENCE: 210	
VDNKFNKEQQ NAFYEILHLP	NLNTEEQRNFIQ IQSLKDDPSR SANLLAEAKK VNDAQAPK 58

-continued

SEQ ID NO: 211	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 211		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSR SANLLAEAKK LNDQAQPK		58
SEQ ID NO: 212	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 212		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 213	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 213		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 214	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 214		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQSLKDDPSG SANLLAEAKK VNDAQAPK		58
SEQ ID NO: 215	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 215		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSE SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 216	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 216		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSE SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 217	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 217		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSD SANLLAEAKK LNDQAQPK		58
SEQ ID NO: 218	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 218		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSG SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 219	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 219		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQELKDDPSG SANLLAEAKK LNDQAQPK		58
SEQ ID NO: 220	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	

-continued

SEQUENCE: 220	organism = synthetic construct	
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSK SANLLAEAKK LNDAPK		58
SEQ ID NO: 221	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 221		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSD SANLLAEAKK VNDAPK		58
SEQ ID NO: 222	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 222		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSQ SANLLAEAKK INDAPK		58
SEQ ID NO: 223	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 223		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSD SANLLAEAKK INDAPK		58
SEQ ID NO: 224	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 224		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSK SANLLAEAKK LNDAPK		58
SEQ ID NO: 225	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 225		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSR SANLLAEAKK LNDAPK		58
SEQ ID NO: 226	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 226		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQDLKDDPSK SANLLAEAKK INDAPK		58
SEQ ID NO: 227	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 227		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQDLKDDPSN SANLLAEAKK SNDAPK		58
SEQ ID NO: 228	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 228		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQDLKDDPSR SANLLAEAKK VNDAPK		58
SEQ ID NO: 229	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 229		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQELKDDPSR SANLLAEAKK LNDAPK		58

-continued

SEQ ID NO: 230	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 230		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSD SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 231	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 231		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSN SANLLAEAKK INDAQAPK		58
SEQ ID NO: 232	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 232		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 233	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 233		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQGLKDDPSR SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 234	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 234		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSK SANLLAEAKK INDAQAPK		58
SEQ ID NO: 235	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 235		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSQ SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 236	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 236		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 237	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 237		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSS SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 238	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 238		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQHLKDDPSS SANLLAEAKK VNDAQAPK		58
SEQ ID NO: 239	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	

-continued

SEQUENCE: 239	organism = synthetic construct	
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQKLKDDPSE SANLLAEAKK INDAQAPK		58
SEQ ID NO: 240	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 240		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQKLKDDPSK SANLLAEAKK LNDAQAPK		58
SEQ ID NO: 241	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 241		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQKLKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 242	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 242		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQNLKDDPSH SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 243	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 243		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSD SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 244	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 244		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSH SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 245	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 245		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSH SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 246	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 246		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSK SANLLAEAKK LNDAQAPK		58
SEQ ID NO: 247	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 247		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSK SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 248	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 248		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSQ SANLLAEAKK RNDAQAPK		58

-continued

SEQ ID NO: 249	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 249		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQPLKDDPSS SANLLAEAKK INDAQAPK		58
SEQ ID NO: 250	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 250		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSD SANLLAEAKK VNDAQAPK		58
SEQ ID NO: 251	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 251		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSN SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 252	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 252		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSQ SANLLAEAKK INDAQAPK		58
SEQ ID NO: 253	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 253		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQQLKDDPSR SANLLAEAKK VNDAQAPK		58
SEQ ID NO: 254	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 254		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQRLKDDPSQ SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 255	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 255		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQRLKDDPSQ SANLLAEAKK SNDAQAPK		58
SEQ ID NO: 256	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 256		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQRLKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 257	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 257		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQRLKDDPSR SANLLAEAKK LNDAQAPK		58
SEQ ID NO: 258	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	

-continued

```

                                organism = synthetic construct
SEQUENCE: 258
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQSLKDDPSR SANLLAEAKK RNDAPK 58

SEQ ID NO: 259      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 259
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQSLKDDPSR SANLLAEAKK SNDAPK 58

SEQ ID NO: 260      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 260
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSK SANLLAEAKK INDAPK 58

SEQ ID NO: 261      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 261
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSQ SANLLAEAKK INDAPK 58

SEQ ID NO: 262      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 262
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSR SANLLAEAKK RNDAPK 58

SEQ ID NO: 263      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 263
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSK SANLLAEAKK RNDAPK 58

SEQ ID NO: 264      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 264
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQTLKDDPSK SANLLAEAKK GNDAPK 58

SEQ ID NO: 265      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 265
IAAQHDKDQQ AAFYEILHLP NLTEEQRNAF IQSLRDDPSQ SLEILAEAKK LNDAPK 58

SEQ ID NO: 266      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 266
IAAQHDKHEQ AAFYEILHLP NLTEDQRNAF IQSLRDDPSQ SLEILGEAKK LNDAPK 58

SEQ ID NO: 267      moltype = AA length = 58
FEATURE            Location/Qualifiers
source             1..58
                   mol_type = protein
                   organism = synthetic construct

SEQUENCE: 267
VDNKFNKEQQ NAFTEILHLP NLTEEQRNAF IQSLKDDPSQ TKEILAEAKK LNDAPK 58

```

-continued

SEQ ID NO: 268	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 268		
VDAKFDKEQQ NAFYEILHLP NLTEEQRNAF IQSLKDDPSQ SANLLAEAKK LNDQAQPK		58
SEQ ID NO: 269	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 269		
IAAQHDKDQQ AAFYEILHLP NLTEEQRNAF IQALRDDPSR SLEILAEAKK INDAQAPK		58
SEQ ID NO: 270	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 270		
IAAQHDKHEQ AAFYEILHLP NLTEDQRNAF IQALRDDPSR SLEILGEAKK INDAQAPK		58
SEQ ID NO: 271	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 271		
VDNKFNKEQQ NAFTEILHLP NLTEEQRNAF IQALKDDPSR TKEILAEAKK INDAQAPK		58
SEQ ID NO: 272	moltype = AA length = 56	
FEATURE	Location/Qualifiers	
source	1..56	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 272		
AQHDEAQQNA FYQVLNMPNL NADQRNGFIQ ALKDDPSRSA NVLGAEQKIN DSQAPK		56
SEQ ID NO: 273	moltype = AA length = 61	
FEATURE	Location/Qualifiers	
source	1..61	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 273		
ADAQQNKFNK DQQSIFYEIL NMPNLNEEQR NGFIQALKDD PSRSTNVLGE AKKINESQAP		60
K		61
SEQ ID NO: 274	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 274		
ADNNFNKEQQ NAFYEILNMP NLNEEQRNGF IQALKDDPSR SANLLAEAKK INESQAPK		58
SEQ ID NO: 275	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 275		
ADNKFNKEQQ NAFYEILHLP NLNEEQRNGF IQALKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 276	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 276		
ADNKFNKEQQ NAFYEILHLP NLTEEQRNGF IQALKDDPSR SKEILAEAKK INDAQAPK		58
SEQ ID NO: 277	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	

-continued

	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 277		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 278	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 278		
VDAKFDKEQQ NAFYEILHLP NLTEEQRNAF IQALKDDPSR SANLLAEAKK INDAQAPK		58
SEQ ID NO: 279	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = unidentified	
SEQUENCE: 279		
IAAQHDKDQQ AAFYEILHLP NLTEEQRNAF IQSLRDDPSG SLEILAEAKK VNDAQAPK		58
SEQ ID NO: 280	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 280		
IAAQHDKHEQ AAFYEILHLP NLTEDQRNAF IQSLRDDPSG SLEILGEAKK VNDAQAPK		58
SEQ ID NO: 281	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 281		
VDNKFNKEQQ NAFYEILHLP NLTEEQRNAF IQSLKDDPSG TKEILAEAKK VNDAQAPK		58
SEQ ID NO: 282	moltype = AA length = 56	
FEATURE	Location/Qualifiers	
source	1..56	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 282		
AQHDEAQQNA FYQVLNMPNL NADQRNGFIQ SLKDDPSGSA NVLGAEQKVN DSQAPK		56
SEQ ID NO: 283	moltype = AA length = 61	
FEATURE	Location/Qualifiers	
source	1..61	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 283		
ADAQQNKFNK DQSAFYEIL NMPNLNEEQR NGFIQSLKDD PSGSTNVLGE AKKVNESQAP		60
K		61
SEQ ID NO: 284	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 284		
ADNNFNKEQQ NAFYEILHLP NLNEEQRNGF IQSLKDDPSG SANLLAEAKK VNESQAPK		58
SEQ ID NO: 285	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 285		
ADNKFNKEQQ NAFYEILHLP NLNEEQRNGF IQSLKDDPSG SANLLAEAKK VNDAQAPK		58
SEQ ID NO: 286	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 286		

-continued

ADNKFNKEQQ	NAFYEILHLP	NLTEEQRNGF	IQSLKDDPSG	SKEILAEAKK	VNDAQAPK	58
SEQ ID NO: 287	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 287						
VDNKFNKEQQ	NAFYEILHLP	NLNEEQRNAF	IQSLKDDPSG	SANLLAEAKK	VNDAQAPK	58
SEQ ID NO: 288	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 288						
VDAKFDKEQQ	NAFYEILHLP	NLTEEQRNAF	IQSLKDDPSG	SANLLAEAKK	VNDAQAPK	58
SEQ ID NO: 289	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 289						
IAAQHDKDQQ	AAFYEILHLP	NLTEEQRNAF	IQALRDDPSG	SLEILAEAKK	RNDAQAPK	58
SEQ ID NO: 290	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 290						
IAAQHDKHQ	AAFYEILHLP	NLTEDQRNAF	IQALRDDPSG	SLEILGEAKK	RNDAQAPK	58
SEQ ID NO: 291	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 291						
VDNKFNKEQQ	NAFYEILHLP	NLTEEQRNAF	IQALKDDPSG	TKEILAEAKK	RNDAQAPK	58
SEQ ID NO: 292	moltype = AA length = 56					
FEATURE	Location/Qualifiers					
source	1..56					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 292						
AQHDEAQQNA	FYQVLNMPNL	NADQRNGFIQ	ALKDDPSGSA	NVLGEAQKRN	DSQAPK	56
SEQ ID NO: 293	moltype = AA length = 61					
FEATURE	Location/Qualifiers					
source	1..61					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 293						
ADAQQNKPNK	DQQSAFYEIL	NMPNLNNEEQR	NGFIQALKDD	PSGSTNVLGE	AKKRNESQAP	60
K						61
SEQ ID NO: 294	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 294						
ADNNFNKEQQ	NAFYEILHLP	NLNEEQRNGF	IQALKDDPSG	SANLLAEAKK	RNESQAPK	58
SEQ ID NO: 295	moltype = AA length = 58					
FEATURE	Location/Qualifiers					
source	1..58					
	mol_type = protein					
	organism = synthetic construct					
SEQUENCE: 295						
ADNKFNKEQQ	NAFYEILHLP	NLNEEQRNGF	IQALKDDPSG	SANLLAEAKK	RNDAQAPK	58
SEQ ID NO: 296	moltype = AA length = 58					

-continued

FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 296		
ADNKFNKEQQ NAFYEILHLP NLTEEQRNGF IQALKDDPSG SKEILAEAKK RNDAQAPK		58
SEQ ID NO: 297	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 297		
VDNKFNKEQQ NAFYEILHLP NLNEEQRNAF IQALKDDPSG SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 298	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 298		
VDAKFDKEQQ NAFYEILHLP NLTEEQRNAF IQALKDDPSG SANLLAEAKK RNDAQAPK		58
SEQ ID NO: 299	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 299		
ADNKFNKDHH AAFYEIHNMP YLNEDQHNAI IHSLRHEPSG SLEVLGEAQR VNKSQAPK		58
SEQ ID NO: 300	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 300		
ADNKFNKAQH NAFYEIHNMP YLNEDQHNAI IHSLRHEPSG SANVLGEAQR VNKSQAPK		58
SEQ ID NO: 301	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 301		
ADNKFNKAKH NAFYEIHNMP YLNEDQHNGA IHSLRHEPSG SAAVLAEAQR VNKSQAPK		58
SEQ ID NO: 302	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 302		
ADNKFNKAQH NAFYEIHNMP YLNEDQHNAI IHSLRHEPSG SAEVLAEAQR VNKSQAPK		58
SEQ ID NO: 303	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 303		
ADNKFNKAQH WAFYEIHNMP YLNEDQHNAI IHSLRHEPSG SAEVLAEAQR VNKSQAPK		58
SEQ ID NO: 304	moltype = AA length = 58	
FEATURE	Location/Qualifiers	
source	1..58	
	mol_type = protein	
	organism = synthetic construct	

-continued

SEQUENCE: 304
ADNKFNKAQH TAFYEIHNP YLNEDQHNA IHS LRHEPSG SAEVLAEQR VNKSQAPK 58

SEQ ID NO: 305 moltype = AA length = 58
FEATURE Location/Qualifiers
source 1..58
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 305
ADNKFNKAQH QAFYEIHNP YLNEDQHNA IHS LRHEPSG SAEVLAEQR VNKSQAPK 58

1. Fc binding polypeptide derived from a *Staphylococcus* Protein A (SpA) or any domain thereof, wherein said polypeptide has binding affinity to an Fc region of an immunoglobulin, and has lower binding affinity for a VH3 region of trastuzumab compared to the binding affinity of SEQ ID NO:59 for the same VH3 region.

2. Fc binding polypeptide comprising a sequence A, which Sequence A consists of an amino acid sequence selected from i), ii) and iii), wherein i), ii) and iii) are defined as follows:

i)

(SEQ ID NO: 65)
EX₂₅QX₂₇X₂₈X₂₉X₃₀IX₃₂X₃₃LX₃₅X₃₆X₃₇PSX₄₀SX₄₂X₄₃LX₄₄EAX₄₉
X₅₀X₅₁NX₅₃X₅₄

wherein, independently from each other,

- X₂₅ is selected from E and D;
- X₂₇ is selected from R and H;
- X₂₈ is selected from N, A, S, H and W;
- X₂₉ is selected from A, G and K;
- X₃₀ is selected from F and A;
- X₃₂ is selected from Q and H;
- X₃₅ is selected from K, R and H;
- X₃₆ is selected from D and H;
- X₃₇ is selected from D and E;
- X₄₂ is selected from A, K, L, R and T;
- X₄₃ is selected from N, E, A, K and S;
- X₄₄ is selected from L, I and V;
- X₄₆ is selected from A, G and K;
- X₄₉ is selected from K, Q and R;
- X₅₀ is selected from K and R;
- X₅₃ is selected from D, E and K; and
- X₅₄ is selected from A and S;

ii) an amino acid sequence which has at least 83% identity to a sequence defined by i);

iii) an amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 24-54 in SEQ ID NO:58, residues 24-54 in SEQ ID NO:59, residues 24-54 in SEQ ID NO:60, residues 24-54 in SEQ ID NO:61, residues 24-54 in SEQ ID NO:62, residues 27-57 in SEQ ID NO:63 and residues 22-52 in SEQ ID NO:64,

wherein additionally, in each of i), ii) and iii) independently from each other,

- X₃₃ is selected from T, S, G, Q, A, E, H, R, P, D, K and N;
 - X₄₀ is selected from E, G, R, D, K, Q, N, H and S; and
 - X₅₁ is selected from L, V, S, I, R and G;
- with the proviso that when X₅₁ is L then X₃₃X₄₀ is selected from AD, HK, EG, ER, GR, AK, AR, PK, RR and KK and when X₅₁ is G then X₃₃X₄₀ is TK.

3. Fc binding polypeptide according to claim 2, wherein said Fc binding polypeptide has a lower binding affinity for the VH3 region of trastuzumab, than SEQ ID NO:59.

4. Fc binding polypeptide according to claim 2, wherein X₃₃ is selected from T, S, G, Q, A, E and H; and/or X₄₀ is selected from E, G, R, D, K and Q; and/or X₅₁ is selected from L, V, S, I and R.

5. Fc binding polypeptide according to claim 2, wherein X₃₃X₄₀X₅₁ are selected from the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GGS, GNI, GQI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, PQR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, RQS, RRI, RRL, RLL, SGV, SRR, SRS, SRV, TER, TKG, TKI, TQI, TRR and TSR, such as the group consisting of ADL, AGR, AKL, ARI, ARL, DKI, DNS, DRV, EGL, ERL, GDR, GDV, GER, GGI, GNI, GRI, GRL, GRR, HKI, HKL, HQS, HRI, HSR, HSV, KEI, KKL, KRI, NHS, PDR, PHR, PHS, PKL, PKS, PQR, PSI, QDI, QDV, QNS, QQI, QRI, QRV, RQR, RQS, RRI, RRL, RLL, SGV, SRR, SRS, SRV, TER, TKI, TQI, TRR and TSR.

6. Fc binding polypeptide according to claim 2, wherein X₃₃X₄₀X₅₁ are selected from the group consisting of GGS, GGI, SRV, GRL, QRI, ARI, SGV, TER, GER, ADL, AGR, EGL, HKL, GDV, GQI and QDI, such as the group consisting of ARI, SGV, AGR and EGL.

7. Fc binding polypeptide according to claim 2, wherein X₃₃X₄₀X₅₁ is SGV.

8. Fc binding polypeptide according claim 2, further comprising a Sequence B arranged N-terminally of said sequence A, which Sequence B consists of an amino acid sequence selected from iv) and v), and wherein iv) and v) are defined as follows:

iv)

(SEQ ID NO: 66)
X₈X₉X₁₀X₁₁AFYX₁₅IX₁₇X₁₈X₁₉PX₂₁LX₂₃

wherein, independently from each other,

- X₈ is selected from E, D and A;
- X₉ is selected from Q, A, L, W, E, V, K, T and H;
- X₁₀ is selected from Q and H;
- X₁₁ is selected from N, A, S, E, K, H, Q, Y, T, F, L, W, I, M, V and R;
- X₁₅ is selected from E, H and Q;
- X₁₇ is selected from L and H;
- X₁₈ is selected from H, N and K;
- X₁₉ is selected from L and M;
- X₂₁ is selected from N, Y and S;

X₂₃ is selected from T and N;

v) an amino acid sequence which has at least 75% identity to a sequence defined by iv).

9. Fc binding polypeptide according to claim 1, comprising a binding module Sequence C, which Sequence C consists of Sequence A and Sequence B, in the following order from the N-terminus to the C-terminus

[Sequence B]-[Sequence A]

or any amino acid sequence which has at least 70% identity to any sequence selected from the group consisting of: residues 8-54 in SEQ ID NO:58, residues 8-54 in SEQ ID NO:59, residues 8-54 in SEQ ID NO:60, residues 8-54 in SEQ ID NO:61, residues 8-54 in SEQ ID NO:62, residues 11-57 in SEQ ID NO:63 and residues 6-52 in SEQ ID NO:64.

10. Fc binding polypeptide according to claim 2, comprising a sequence selected from the group consisting of:

x)

(SEQ ID NO: 172)

VDAKFDKE AQEAFYEILHLPNLT-[Sequence A]-QAPK

wherein [Sequence A] is as defined in claim 2;

xi) an amino acid sequence which has at least 86% identity to the sequence defined in x);

xvi)

(SEQ ID NO: 173)

VDNKFNKEQQNAFYEILHLPNLT-[Sequence A]-QAPK

wherein [Sequence A] is as defined in claim 2; and

xvii) an amino acid sequence which has at least 86% identity to the sequence defined in xvi).

11. Fc binding polypeptide according to claim 10, wherein sequence x) or xvi) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-57, 208-264, 269-305 such as the group consisting of SEQ ID NO:1-16 and 208-223, such as corresponds to SEQ ID NO:7 or 214.

12. Fc binding polypeptide according to claim 10, wherein sequence x) corresponds to a sequence selected from the group consisting of SEQ ID NO:1-57, such as the group consisting of SEQ ID NO:1-16, such as corresponds to SEQ ID NO:7.

13. Fc binding polypeptide multimer, wherein each monomer of the multimer comprises a Fc binding polypeptide which is independently selected from any Fc binding polypeptide as defined in claim 1 and wherein the multimer preferably is a hexamer.

14. Fc binding polypeptide according to claim 1 or Fc binding polypeptide multimer thereof, which is capable of binding to Fc such that the K_D value of the interaction is at most 1×10⁻⁷ M, such as at most 1×10⁻⁸ M, such as at most 1×10⁻⁹ M, such as at most 1×10⁻¹⁰ M, such as at most 1×10⁻¹¹ M.

15. Fc binding polypeptide according to claim 1, or Fc binding polypeptide multimer thereof, which is not capable of binding to said VH3 with a K_D value of the interaction of less than 1×10⁻⁴ M, such as less than 1×10⁻³ M.

16. A separation matrix comprising an Fc binding polypeptide according to claim 1, or an Fc binding polypeptide multimer thereof being coupled to a solid support, which solid support preferably is in fibrous, beaded or particle form.

17. A method of isolating an immunoglobulin comprising a) contacting a liquid sample comprising said immunoglobulin with a separation matrix according claim 16.

* * * * *