



US012416019B2

(12) **United States Patent**
Xiao et al.

(10) **Patent No.:** **US 12,416,019 B2**

(45) **Date of Patent:** ***Sep. 16, 2025**

(54) **EXPRESSION CASSETTE CONTAINING
OVERLAPPING OPEN READING FRAMES
AND APPLICATION**

CN	106544325	3/2017
CN	112553257	3/2021
CN	113897396	1/2022

(71) Applicant: **GENEVOYAGER (WUHAN) CO.,
LTD.**, Hubei (CN)

OTHER PUBLICATIONS

(72) Inventors: **He Xiao**, Hubei (CN); **Xiaobin He**,
Hubei (CN); **Gang Huang**, Hubei
(CN); **Xing Pan**, Hubei (CN); **Yicheng
Zhou**, Hubei (CN); **Liang Du**, Hubei
(CN); **Mengdie Wang**, Hubei (CN);
Huanhuan Zuo, Hubei (CN); **Hao Sun**,
Hubei (CN)

(73) Assignee: **GENEVOYAGER (WUHAN) CO.,
LTD.**, Hubei (CN)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **18/434,746**

(22) Filed: **Feb. 6, 2024**

(65) **Prior Publication Data**

US 2025/0043307 A1 Feb. 6, 2025

Related U.S. Application Data

(63) Continuation of application No.
PCT/CN2023/141032, filed on Dec. 22, 2023.

(30) **Foreign Application Priority Data**

Aug. 3, 2023	(CN)	202310975187.6
Nov. 17, 2023	(CN)	202311544887.6

(51) **Int. Cl.**
C12N 15/86 (2006.01)
C12N 5/07 (2010.01)

(52) **U.S. Cl.**
CPC **C12N 15/86** (2013.01); **C12N 5/0601**
(2013.01); **C12N 2710/14022** (2013.01); **C12N**
2710/14043 (2013.01); **C12N 2710/14052**
(2013.01); **C12N 2750/14144** (2013.01)

(58) **Field of Classification Search**
CPC C12N 15/86
See application file for complete search history.

(56) **References Cited**

FOREIGN PATENT DOCUMENTS

CN	101522903	9/2009
CN	103849629	6/2014

Chen H. (2008, Molecular Cell Therapy, vol. 16(5), pp. 924-930).
(Year: 2008).*

Pyle et al. (1997, Analytical Biochemistry, vol. 253, pp. 253-258).
(Year: 1997).*

Urabe et al. (Human Gene Therapy, vol. 13, pp. 1935-1943). (Year:
2002).*

Seppo Ylä-Herttuala, "Endgame: glybera finally recommended for
approval as the first gene therapy drug in the European union",
Molecular Therapy, Oct. 2012, pp. 1831-1832.

Masashi Urabe et al., "Insect Cells as a Factory to Produce Adeno-
Associated Virus Type 2 Vectors", Human Gene Therapy, Nov.
2002, pp. 1935-1943.

Masashi Urabe et al., "Scalable Generation of High-Titer Recombi-
nant Adeno-Associated Virus Type 5 in Insect Cells", Journal of
Virology, Feb. 2006, pp. 1874-1885.

Erik Kohlbrenner et al., "Successful Production of Pseudotyped
rAAV Vectors Using a Modified Baculovirus Expression System",
Molecular Therapy, Dec. 2005, pp. 1217-1225.

Xiao Xiao et al., "Production of High-Titer Recombinant Adeno-
Associated Virus Vectors in the Absence of Helper Adenovirus",
Journal of Virology, Mar. 1998, pp. 2224-2232.

Mark R. Bruder et al., "Utility of Alternative Promoters for Foreign
Gene Expression Using the Baculovirus Expression Vector Sys-
tem", Viruses, Nov. 2022, pp. 1-18.

* cited by examiner

Primary Examiner — Anoop K Singh

Assistant Examiner — David A Montanari

(74) *Attorney, Agent, or Firm* — JCIPRNET

(57) **ABSTRACT**

An expression cassette containing overlapping open reading
frames and an application thereof are provided. The over-
lapping open reading frames are overlapping open reading
frames of a first ORF and a second ORF and include in
sequence from a 5' end to a 3' end: a first promoter at least
used to drive gene transcription of the first ORF; a 5' part of
a gene of the first ORF; an intron; and a 3' part of a gene of
the second ORF, the intron including a second promoter
used only to drive gene transcription of the second ORF. By
arranging two promoters in a single expression cassette in
the disclosure, the two promoters are used to drive the
expression of proteins of the overlapping reading frames and
regulate the relative expression time and expression inten-
sity of different proteins.

21 Claims, 10 Drawing Sheets

Specification includes a Sequence Listing.




FIG. 1(Prior Art)




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11




FIG. 12




FIG. 13

1

EXPRESSION CASSETTE CONTAINING OVERLAPPING OPEN READING FRAMES AND APPLICATION

CROSS-REFERENCE TO RELATED APPLICATION

This application is a continuation of international application of PCT application serial no. PCT/CN2023/141032, filed on Dec. 22, 2023, which claims the priority benefit of China application no. 202310975187.6, filed on Aug. 3, 2023 and China application no. 202311544887.6, filed on Nov. 17, 2023. The entirety of each of the above-mentioned patent applications is hereby incorporated by reference herein and made a part of this specification.

REFERENCE TO A SEQUENCE LISTING

The instant application contains a Sequencing Listing which has been submitted electronically in XML file and is hereby incorporated by reference in its entirety. Said XML copy, created on Jan. 29, 2024, is named 142172_SEQUENCINGLISTING and is 61,181 bytes in size.

BACKGROUND OF THE INVENTION

Technical Field

The disclosure belongs to the field of genetic engineering, and more specifically, relates to an expression cassette containing a gene with overlapping open reading frames and an application.

Description of Related Art

Adeno-associated virus (AAV), also known as adeno-accompanied virus, belongs to the genus Dependovirus in the family Parvoviridae, is the single-stranded DNA-deficient virus with the simplest structure discovered so far, and requires helper viruses (usually adenoviruses) to participate in replication. The AAV genome is a single-stranded linear DNA, approximately 4700 bp, and includes two upstream and downstream open reading frames (ORFs): Rep and Cap, located between 2 T-shaped inverted terminal repeats (ITR) composed of 145 nucleotides each, as shown in FIG. 1. The function of ITR is to serve as the origin of viral replication and packaging signal. The Rep gene is involved in viral replication and integration and encodes viral replication proteins. The Cap gene is responsible for encoding the three capsid proteins of the virus. Recombinant adeno-associated virus (rAAV) is one of the most promising vectors in the field of gene therapy due to its characteristics of wide host range, low immunogenicity, high safety, and capability of mediating long-term stable expression of exogenous genes in animals. With the approval of the first recombinant adeno-associated virus (rAAV)-mediated gene therapy, there is increasing demand for large-scale AAV vector manufacturing technology (Yla-Herttuala S., 2012, Mol Ther, 20:1831-1832).

At present, there are two main types of rAAV production systems: one is a conventional production system using mammalian cells (e.g., 293 cells, COS cells, HeLa cells, KB cells, etc.), and the other is a production system using insect cells. Herein, the production system using insect cells uses the baculovirus expression vectors (BEVs) carrying the rAAV vector genome, Rep protein, and Cap protein to infect Sf9 insect cells to produce rAAV virus. Taking advantage of

2

the fact that Sf9 cells can express large amounts of recombinant proteins, this system uses insect baculovirus containing rAAV genome, AAV-Rep, and AAV-Cap genes to transduce Sf9 cells, allowing rAAV to be assembled in Sf9 cells, and the purpose of large-scale production of rAAV is ultimately achieved.

In this production system, promoters play an important role. The selected exogenous gene is placed under the control of a native promoter and then expressed during the infection cycle, releasing the protein during cell lysis. The promoter is recognized by RNA polymerase, and transcription begins. In ribonucleic acid (RNA) synthesis, the promoter can interact with the transcription factor that determines the start of transcription, controls the initiation time and extent of gene expression (transcription). It contains the core promoter region and the regulatory region, which is like a "switch" that determines the activity of the gene and thus controls which protein the cell starts to produce. The promoter itself has no compilation function, but it has the function of directing the translation of amino acids in the gene, just like a flag. The core part thereof is the RNA polymerase binding site upstream of the non-coding region. Currently, the most commonly used promoters in BEVs are the polyhedral pol H (polyhedrin) promoter and the p10 promoter, in addition to basic promoters and a few early promoters. The expression levels of the same exogenous gene under the control of different promoters are greatly different.

In order to achieve maximum expression levels, very late promoters are used most of the time to drive the expression of exogenous genes. Although this approach is usually successful when large amounts of recombinant protein are required, but in some cases, the use of alternative promoters with earlier or weaker expression may be beneficial to the quality and yield of the final protein. Proteins that aggregate during rapid expression and proteins that require advanced post-translational modifications have also been shown to yield ideal products through sustained and stable expression using early promoters.

Immediate early and early promoters are activated prior to viral DNA replication. The immediate early promoter (IE) is transcribed by host cell RNA polymerase II and also utilizes host transcription factors. The structural organization of the immediate early and early promoters is quite different from that of the late and very late baculovirus promoters, and is more similar to that of other eukaryotic promoters. The most striking feature of the baculovirus immediate early promoter is the conserved CAGT motif, which serves as the transcription start site and plays an important role in regulating downstream gene expression. In some cases, immediate early promoter contains a TATA-like motif and/or an initiation sequence. This region is recognized by host RNA polymerase II and allows transcription of immediate early viral genes immediately following infection. The IE1 promoter is an immediate early promoter and is the most commonly used constitutive promoter for baculovirus expression because it is active at all time stages as well as in uninfected insect cells. The 39 k promoter is an early promoter that is transactivated by the IE1 protein. When being co-expressed with the IE1 protein, the 39 k promoter can also be used transiently.

The late promoter is transcribed by a viral RNA polymerase that recognizes the conserved (A/G/T) TAAG late/very late promoter motif. The transcription initiates at the second nucleotide within this motif, and its presence is required for promoter activity. Studies investigating the function of the TAAG motif in late promoters have shown

that mutations in this sequence disrupt promoter transcriptional activity. The very late promoter transcriptional activity peaks at 18 to 24 hours after infection and exceeds late promoter activity. At 24 hours after infection, the transcription level of the very late promoter exceeds that of all other promoters in cells.

The P10 protein is expressed at high levels very late in the infection cycle and is thought to play a role in inclusion maturation and cell lysis required for virus release. Weyer and Possee first define the 101 nucleotides upstream of the ATG of the p10 gene as the p10 promoter sequence and suggest that it be used for the expression of exogenous proteins. Since then, the p10 promoter has been widely used to produce proteins in BEVS, such as the production of cauliflower mosaic virus gene 1 and human interleukin 2. p10 can also be used to connect other promoters for co-expression.

In the AAV insect cell production system, for the Cap protein, Urabe et al. replace the start codon AUG of VP1 with the suboptimal start codon ACG to construct one single polycistronic mRNA. This polycistronic mRNA expresses all three AAV2 VP proteins without the need for splicing (Urabe et al., 2002, Hum Gene Ther, 13:1935-1943). However, there are many and increasing serotypes of AAV, and the modification method of Urabe et al. is not suitable for all serotypes. For instance, AAV5 particles produced in the baculovirus system of Urabe et al. using ACG as the start codon of the VP1 capsid protein have poor infectivity. In the method provided by Chinese patent CN106544325A, different suboptimal start codons CTG are used to enhance the expression of VP1. Although this design improves the infectivity of AAV5 particles, this method lacks flexibility in adjusting the relative content of VP1/VP2/VP3. Further, this method also appears to lack the flexibility necessary for serotype-specific sequence adjustment. In the method provided by Chinese patent CN101522903A, an artificial intron containing the polH promoter sequence is inserted into the open reading frame of VP1, and two promoters are used to express VP1 and VP2/VP3. Although this design can also achieve the expression of VP1/VP2/VP3 in the same reading frame, this method cannot effectively adjust the relative amounts of VP1/VP2/VP3. Moreover, the relative expression amounts of VP1/VP2/VP3 vary greatly among different serotypes, resulting in low production efficiency.

For the expression of the Rep protein, Urabe et al. construct two independent expression cassettes containing Rep78 and Rep52 and inserted them into the same baculovirus vector. Herein, Rep52 uses the polH promoter, and Rep78 uses the deltaIE1 promoter, which has weaker promoter activity than the polH promoter (Urabe et al., 2002, Hum Gene Ther, 13:1935-1943). However, the immediate early promoter deltaIE1 is active early in baculovirus infection, and early and abundant expression of the Rep78 protein in insect cells may have a negative impact on the yield of recombinant baculovirus (Urabe et al., 2006, Journal of Virology 80.4:1874-1885). Further, some studies have found that the AAV Rep protein expression method developed by Urabe et al. has the problem of unstable passaging of baculovirus vectors (Kohlbrenner et al., 2005, Mol. Ther. 12: 1217-25). In order to solve the problem of inherent passaging instability of two independent Rep expression cassettes, Chinese patent CN 103849629 A discloses a method for producing AAV in insect cells. The translation initiation codons of the AAV Rep78 protein is replaced with ACG, and the codon is partially skipped by scanning ribosomes so that the translation initiation can also occur at the start codon of the Rep52 protein further downstream. In this method, the

expression of Rep78 and Rep52 proteins in the same reading frame can be achieved, and the stability of the rAAV vector produced in insect cells is significantly improved. However, this method lacks flexibility in adjusting the relative contents of Rep52/Rep78 proteins. The high stoichiometric ratio of Rep52/Rep78 proteins may be a key factor in the high yield of rAAV (Xiao, Xiao et al., 1998, Journal of Virology 72.3 (1998): 2224-2232). In a method for producing AAV in an insect cell disclosed in Chinese patent CN 113897396 A, the inventors achieve the expression of Rep78 and Rep52 proteins in the same reading frame by constructing a Rep expression cassette containing an intron regulatory sequence. In this way, the stability of the rAAV vector produced in the insect cell is improved, and further, the relative content of Rep52/Rep78 proteins can be flexibly adjusted by adjusting the intron splicing efficiency. However, there is a lack of flexibility in controlling the relative expression time and relative expression intensity of the two.

Therefore, the large-scale production of rAAV vectors using baculovirus in insect cells still needs to be further improved. Many studies have shown that earlier promoters may be more efficient in producing some exogenous proteins, particularly those that require extensive post-translational processing and/or secretion. This is because when the latest promoter is most active, the host cell's processing of the exogenous protein may be impaired. For complex viral particles, different proteins may be required in different amounts. By using different baculovirus promoters that are active at different stages of infection, control of the expression time and amount of the target protein can be achieved (Bruder et al., Viruses 2022, 14, 2670). However, the mechanism of synergy between different promoters is still unclear.

SUMMARY OF THE INVENTION

In view of the above defects or improvement needs of the related art, the disclosure provides an expression cassette for a gene containing overlapping open reading frames and an application, aiming to better control the relative expression time and relative intensity of different proteins represented by Rep78 and Rep52 proteins by constructing an expression cassette containing different proteins located in the same overlapping open reading frame and by using different promoters to control the expression of different proteins, and the use of overlapping open reading frames to express proteins not only ensures the stability of a rAAV vector produced in an insect cell, but also improves the yield and quality of the produced rAAV.

To achieve the above, in an aspect of the disclosure, the disclosure provides an expression cassette of a gene containing overlapping open reading frames, including:

- operably linked from a 5' end to a 3' end:
 - an artificially constructed sequence; and
 - an overlapping open reading frame lacking a first translation initiation codon,
- where the artificially constructed sequence includes a first promoter and a second promoter independent of each other, and a complete sequence of the first translation initiation codon is not included between the first promoter and the second promoter.

Preferably, the gene is a parvovirus Rep gene, the first promoter is used to drive expression of a Rep78/68 protein and a Rep52/40 protein, and the second promoter is used to additionally drive the expression of the Rep52/40 protein.

Preferably, the gene is a parvovirus Cap gene, the first promoter is used to drive expression of a VP1 protein and a

VP2/3 protein, and the second promoter is used to additionally drive the expression of the VP2/3 protein.

Preferably, the artificially constructed sequence further includes an intron, and the intron is used to form the first translation initiation codon in the artificially constructed sequence through alternative splicing during post-transcriptional processing.

Further preferably, the intron is located between any two nucleotides of ATG.

Further preferably, the first promoter precedes the 5' end of the intron.

Further preferably, the second promoter is within the intron.

Preferably, the second promoter precedes the 5' end of the first promoter.

Preferably, the first promoter is active before the second promoter.

Further preferably, the first promoter is deltaIE1, IE1, 39K, or p6.9, and the second promoter is polH or p10.

Preferably, the first promoter is p10, and the second promoter is polH.

According to an aspect of the disclosure, the disclosure provides an optimized expression cassette for a gene containing overlapping open reading frames. The overlapping open reading frames are overlapping open reading frames of a first ORF and a second ORF and further include in sequence from a 5' end to a 3' end:

a first promoter used to drive gene transcription of the first ORF;

a 5' part of a gene of the first ORF;

an intron; and

a 3' part of a gene of the second ORF,

where the intron includes a second promoter used to drive gene transcription of the second ORF, the first promoter is an immediate early promoter, an early promoter, or a late promoter, and the second promoter is a very late promoter.

Preferably, the intron is located between any two nucleotides of the translation initiation codons ATG of the first ORF. The intron is used to form the first translation initiation codon in the artificially constructed sequence through alternative splicing during post-transcriptional processing.

Preferably, the intron is located between the translation initiation codons ATG of the first ORF and an exon of the first ORF.

Preferably, the first promoter is deltaIE1, IE1, 39K, or p6.9, and the second promoter is polH or p10.

Preferably, the expression cassette is used to encode a parvovirus Rep gene, the first ORF is used to encode a Rep68/Rep78 protein, and the second ORF is used to encode a Rep40/Rep52 protein.

Preferably, the expression cassette is used to encode a parvovirus Cap gene, the first ORF is used to encode a VP1 protein, and the second ORF is used to encode a P2/VP3 protein.

According to another aspect of the disclosure, the disclosure further provides a nucleic acid molecule including a first expression cassette, and the first expression cassette is any expression cassette of the above.

Preferably, a second expression cassette is further included, and the first expression cassette and the second expression cassette are respectively used to encode the parvovirus Cap gene and the parvovirus Rep gene.

Preferably, the nucleic acid molecule further includes an exogenous gene.

Further preferably, AAV inverted terminal repeats are provided at both ends of the exogenous gene.

Further preferably, the exogenous gene is a reporter gene, and the reporter gene is at least one of a chloramphenicol acetyltransferase encoding gene, a β -galactosidase encoding gene, a β -glucuronidase encoding gene, a *Renilla* luciferase encoding gene, an alkaline phosphatase encoding gene, a firefly luciferase encoding gene, a green fluorescent protein encoding gene, and a red fluorescent protein encoding gene.

Further preferably, the exogenous gene is a gene encoding a drug polypeptide, and the drug polypeptide is at least one of lipoprotein esterase, apolipoprotein, cytokine, interleukin, and interferon.

According to another aspect of the disclosure, the disclosure further provides a recombinant baculovirus vector including the expression cassette according to the above.

Preferably, the recombinant baculovirus vector is an insect cell compatible vector.

According to another aspect of the disclosure, the disclosure further provides an insect cell including the recombinant baculovirus vector according to the above.

Preferably, the insect cell is a *Spodoptera frugiperda* cell, a *Trichopodia* cell, a *Drosophila* cell, or a mosquito cell.

To sum up, in the technical solutions provided by the disclosure when being compared to the related art, two promoters are arranged in a single expression cassette to drive the expression of proteins of the overlapping reading frames and regulate the relative expression time and expression intensity of different proteins. The relative expression time and a wider range of relative expression intensity of the Rep78/68 protein and the Rep52/40 protein in the Rep gene, as well as the relative expression time and a wider range of relative expression intensity of the VP1 protein and the VP2/3 protein in the Cap gene can be better controlled in the disclosure. In this way, not only the efficiency of rAAV vector production in the insect cell is improved, but also the quality of the produced AAV is improved.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic diagram of expression regulation of a Cap gene and a Rep gene in wild-type AAV.

FIG. 2 is a schematic diagram of expression regulation of a RepA-D gene expression cassette I in Examples 1 to 4.

FIG. 3 is a schematic diagram of expression regulation of a Cap gene expression cassette in Example 1.

FIG. 4 is a schematic diagram of expression regulation of a RepI-IV gene expression cassette II in Examples 5 to 8.

FIG. 5 is a schematic diagram of expression regulation of a Rep1-4 gene expression cassette III in Example 5.

FIG. 6 is a Western Blot detection diagram of a recombinant baculovirus vector AcRepA-D expressing a Rep protein in Examples 1 to 4.

FIG. 7 is a Western Blot detection diagram of a recombinant baculovirus vector AcRepI-IV expressing a Rep protein in Examples 5 to 8.

FIG. 8 is a Western Blot detection diagram of a recombinant baculovirus vector AcRep1-4 expressing a Rep protein in Comparative Examples 1 to 4.

FIG. 9 is a growth kinetic curve of sf9 cells infected with constructs CRI-0, CRI-1, and CRI-2 in Verification Example 1.

FIG. 10 is a schematic diagram of expression regulation of a RepE gene expression cassette in Example 9.

FIG. 11 is a schematic structural diagram of a RepF-H gene expression cassette in Examples 10 to 12.

FIG. 12 is a Western Blot detection diagram of a recombinant baculovirus vector AcRepE-H expressing a Rep protein in Examples 9 to 12.

FIG. 13 is a Western Blot detection diagram of a recombinant baculovirus vector AcRep5 expressing a Rep protein in Comparative Example 5.

DETAILED DESCRIPTION OF THE EMBODIMENTS

In order to make the objectives, technical solutions, and advantages of the disclosure clearer and more comprehensible, the disclosure is further described in detail with reference to the drawings and embodiments. It should be understood that the specific embodiments described herein serve to explain the invention merely and are not used to limit the invention. In addition, the technical features involved in the various embodiments of the invention described below can be combined with each other as long as the technical features do not conflict with each other.

For ease of understanding, the terms appearing in this specific embodiment are defined as follows:

The term “polynucleotide”: refers to a biological macromolecule composed of single nucleotides. The polynucleotide molecule mentioned in the disclosure may be a deoxyribonucleic acid (DNA) molecule or a ribonucleic acid (RNA) molecule. A person having ordinary skill in the art knows that the RNA sequence is basically similar to the DNA sequence or has a certain degree of sequence identity. Thymine (T) in the DNA sequence can be considered equivalent to uracil (U) in the RNA sequence.

The term “operably linked”: refers to the connection of polynucleotide (or polypeptide) sequences in a functional relationship. Two nucleotide sequences are “operably linked” when the two nucleotide sequences are placed in a functional relationship. For instance, a transcriptional control sequence (e.g., a promoter) is operably linked to a gene’s coding sequence if it affects the transcription of the gene’s coding sequence.

The term “expression cassette”: refers to a nucleic acid construct including coding and regulatory sequences operably linked when introduced into a host cell, resulting in the transcription and/or translation of RNA or polypeptide, respectively. An expression cassette is understood to include a promoter allowing the initiation of transcription, the open reading frame of the gene of interest, and a transcription terminator. Generally, the promoter sequence is placed upstream of the gene of interest at a distance compatible with expression control. The “expression cassette” may be a multi-piece nucleic acid construct or a one-piece nucleic acid construct.

The term “open reading frame” (ORF): refers to the normal nucleotide sequence of a structural gene with the potential to encode a protein or polypeptide and begins with a start codon and ends with a stop codon with no stop codon in between to interrupt translation. On an mRNA chain, a ribosome starts translation from the start codon, synthesizes a polypeptide chain along the mRNA sequence, and continues to extend. When a stop codon is encountered, the extension reaction of the polypeptide chain is terminated.

The term “vector”: refers to a nucleic acid molecule designed to transport, transfer, and/or store genetic material, as well as to express genetic material and/or integrate genetic material into the chromosomal DNA of a host cell, such as a plasmid vector, a cosmid vector, an artificial chromosome, a phage vector, and other viral vectors. A vector usually consists of at least three basic units, namely a replication source, a selectable marker, and a multiple cloning site.

The term “intron”: also known as spacer sequence, refers to a non-coding segment in a gene or mRNA molecule and is an intervening sequence in the DNA of eukaryotic cells. The intronic sequence is transcribed in the mRNA precursor, is removed by splicing, and is ultimately no longer present in the mature mRNA molecule. Depending on whether the splicing process is spontaneous or processed by the spliceosome, introns are divided into self-splicing introns and spliceosomal introns. A self-splicing intron is a special type of intron, a ribozyme that can be cut out by its own action to leave the mRNA. The intron involved in the disclosure is a spliceosomal intron, and the splicing of such an intron requires the help of the spliceosome. There are a splicing donor sequence and a splicing acceptor sequence at both ends of the intron sequence, which are the sequences on both sides of the cutoff and rejoining sites. The spliceosome is a ribonucleoprotein complex dynamically composed of small nuclear RNAs (snRNAs) and protein factors. The spliceosome recognizes the splicing site of the mRNA precursor, catalyzes the splicing reaction, completely cuts out the intron, and then rejoins the upstream and downstream RNA sequences.

The intron mentioned in the disclosure can be a natural intron, or an intron that has been artificially modified and has splicing activity in insect cells, and the source thereof is not limited to insect cells.

The term “AAV serotype”: since the discovery of adeno-associated viruses (AAVs), more than 100 AAV serotypes or mutants have been isolated from adenovirus, humans, or primates, and some mammals, among which AAV1-9 is mainly used. The main difference between rAAV vectors of different serotypes is the capsid protein. Different AAV serotypes have certain differences in infection efficiency and tissue specificity.

The genome structures of all known adeno-associated virus serotypes are very similar. AAV is a single-stranded DNA virus with a simple genome structure and a full length of approximately 4.7 kb, as shown in FIG. 1. Its genome contains a rep gene expression cassette, a cap gene expression cassette, and AAV inverted terminal repeats (ITRs) located at both ends of the genome. ITR is a 125-nucleotide palindrome structure at both ends of the genome, can form a self-complementary inverted T-shaped hairpin structure, and is a cis-acting element required for the initiation of DNA replication and packaging of recombinant AAV genome into infectious viral particles. Between the ITR sequences is the virus coding region, where the rep gene expression cassette and the cap gene expression cassette are located.

As a defective virus, AAV cannot replicate independently in the absence of helper viruses. Therefore, AAV can only be integrated into the host cell chromosome at a specific site and remain in a latent state. In the presence of helper virus, the increased expression amount of the rep gene can rescue the AAV genome integrated in the host cell chromosome and replicate AAV DNA in large quantities. The single-stranded rAAV genome is packaged into infectious virus particles under the action of VP capsid protein.

The term “Cap gene”: consists of overlapping amino acid sequences encoding the structural VP capsid protein and contains three overlapping open reading frames, encoding three types of subunits: VP1, VP2, and VP3. VP1, VP2, and VP3 contain different start codons and share a stop codon, and VP1 and VP2 share the VP3 sequence. The N-terminus of VP1 has a conserved phospholipase A2 sequence, and this sequence is involved in the escape of the virus from the body and is critical to its infectivity. The VP2 protein is not essential for assembly or infection. The core of the VP3

protein consists of a conserved β -barrel motif, which determines the differences in receptors between different serotypes of AAV and host cells. The correct ratio of the three proteins in wild-type AAV is 6:6:54, which is approximately 1:1:10.

The term “Rep gene”: is used to encode four overlapping multifunctional proteins, Rep78, Rep68, Rep52, and Rep40. Rep78 and Rep68 proteins are involved in the replication and integration of AAV and can bind to specific sequences in the ITR. Rep52 and Rep40 proteins have helicase and ATPase activities and participate in single-stranded genome replication and virus assembly. Whether in mammalian cells or insect cells, unspliced mRNA encoding Rep78 and Rep52 proteins is sufficient to prepare rAAV.

The term “relative expression content”: is the relative proportion of the expression amounts of different proteins. For instance, an increase in the relative expression content of Rep78/Rep52 protein may mean an increase in Rep78 protein expression amount, or it may mean a decrease in Rep52 protein expression amount.

The term “promoter”: is the DNA sequence located upstream of the 5' end of a structural gene and is able to activate RNA polymerase, so that it can accurately bind to the template DNA and have the specificity for transcription initiation.

The term “solid rate”: taking the AAV vector as an example, the solid AAV is the AAV vector carrying the target gene. The empty shell AAV vector is an AAV vector that does not carry the target gene, consists of a serotype-specific AAV protein coat, and is usually produced by incomplete viral packaging during the production process. It must be minimized during purification of gene therapy products because these viral capsids do not contain DNA. When they are infused into patients, they compete for the limited number of receptors on the cell surface on the one hand, and they are not beneficial to gene therapy and may cause side effects on the other hand. Further, the increase in empty shell AAV content will lead to an increase in the dosage. The higher the dosage, the higher the risk of serious side effects. The same is true for AAVs that contain incomplete copies of therapeutic genes.

The solid rate refers to the proportion of solid AAV in the prepared AAV vector. The higher the solid rate, the greater the proportion of solid AAV, which proves that the quality of the prepared AAV vector is higher.

The disclosure provides an expression cassette used to express a gene including overlapping open reading frames in a cell, especially in an insect cell, and the expression cassette includes:

- operably linked from a 5' end to a 3' end:
 - an artificially constructed sequence; and
 - an overlapping open reading frame lacking a first translation initiation codon,
- where the artificially constructed sequence includes a first promoter and a second promoter independent of each other, and a complete sequence of the first translation initiation codon is not included between the first promoter and the second promoter.

In some embodiments, the artificially constructed sequence also includes an intron, and the intron is located between translation initiation codons ATG, for example, between AT and G or between A and TG. The intron is used to form the first translation initiation codon in the artificially constructed sequence through alternative splicing during post-transcriptional processing, so that the overlapping open reading frame can be expressed smoothly.

In some embodiments, the second promoter precedes the first promoter (i.e., precedes the 5' end of the first promoter), and the first promoter precedes the intron. In some other embodiments, the first promoter precedes the intron, and the second promoter is contained within the intron.

In some embodiments, the first promoter is used to express all proteins encoded by the overlapping open reading frame, while the second promoter is used to adjust the relative expression time and relative expression content/expression intensity of these proteins. In some other embodiments, the gene encoded by the expression cassette is a parvovirus Rep gene. The two promoters of the disclosure may be used to adjust the relative expression time and relative expression content/expression intensity of different Rep proteins. Further, a Rep40 protein has a similar function to a Rep52 protein, and a Rep68 protein also has a similar function to a Rep78 protein. Therefore, the first promoter may be used to drive the expression of the Rep68 and/or Rep78 protein(s) and may also be used to drive the expression of the Rep40 protein and/or the Rep52 protein, and the second promoter is used to additionally drive the expression of the Rep40 protein and/or the Rep52 protein.

In some embodiments, the gene encoded by the expression cassette is a parvovirus Cap gene. The two promoters of the disclosure may be used to adjust the relative expression time and relative expression content/expression intensity of different Cap proteins. Further, the first promoter is used to drive the expression of a VP1 protein and may also be used to drive the expression of a VP2/3 protein, and the second promoter is used to additionally drive the expression of the VP2/3 protein.

In some embodiments, the first promoter is active before the second promoter. For instance, the first promoter may be an early and late promoter such as deltaIE1, IE1, 39K, or p6.9, and the second promoter may be a very late promoter such as p10 or polH.

In some other embodiments, the first promoter and the second promoter may produce activity almost simultaneously, for example, the first promoter is p10 and the second promoter is polH.

Since variations in some non-critical sites of the promoters have no impact on their functions, the sequences of the first promoter and the second promoter may also be partially different from the original sequences of the above promoters.

The disclosure further provides an optimized expression cassette for a gene including overlapping open reading frames for use in a cell, especially in an insect cell. The overlapping open reading frames are overlapping open reading frames of a first ORF and a second ORF and include in sequence from a 5' end to a 3' end:

- a first promoter used to drive gene transcription of the first ORF;
 - a 5' part of a gene of the first ORF;
 - an intron; and
 - a 3' part of a gene of the second ORF,
- where the intron includes a second promoter used to drive gene transcription of the second ORF, the first promoter is an immediate early promoter, an early promoter, or a late promoter, the second promoter is a very late promoter, and for example, in some embodiments, the first promoter is deltaIE1, IE1, 39K, or p6.9, and the second promoter is polH or p10.

In some embodiments, the intron is located between any two nucleotides of translation initiation codons ATG of the first ORF. The intron is used to form the first translation

11

initiation codon in the artificially constructed sequence through alternative splicing during post-transcriptional processing.

In some other embodiments, the intron is located between the translation initiation codons ATG of the first ORF and an exon of the first ORF.

In some embodiments, the gene encoded by the expression cassette is a parvovirus Rep gene, the first ORF is used to encode the Rep78 protein, and the second ORF is used to encode the Rep52 protein. The two promoters of the disclosure may be used to adjust the relative expression time and relative expression content/expression intensity of different Rep proteins. Further, the Rep40 protein has a similar function to the Rep52 protein, and the Rep68 protein also has a similar function to the Rep78 protein. Therefore, the first promoter may be used to drive the expression of the Rep68 and/or Rep78 protein(s), and the second promoter is used to drive the expression of the Rep40 protein and/or the Rep52 protein.

In some embodiments, the second promoter contains translation initiation codon and stop codon sequences that are recognized by a ribosome, so that when the intron is not spliced during the post-transcriptional processing of a Rep gene expression cassette driven by the first promoter, the Rep40 protein and/or the Rep52 protein will not be translated and produced. In some embodiments, the gene encoded by the expression cassette is a parvovirus Cap gene. The two promoters of the disclosure may be used to adjust the relative expression time and relative expression content/expression intensity of different Cap proteins. Further, the first promoter is used to drive the expression of the VP1 protein, and the second promoter is used to additionally drive the expression of the VP2 and/or VP3 protein(s).

Since variations in some non-critical sites of the promoters have no impact on their functions, the sequences of the first promoter and the second promoter may also be partially different from the original sequences of the above promoters.

The expression cassette of the Rep gene may be integrated into different baculovirus vectors with the expression cassette of the Cap gene and a sequence of an exogenous gene containing AAV inverted terminal repeats at both ends, or they can be integrated into the same recombinant baculovirus vector. In some embodiments, both the expression cassette of the Rep gene and the expression cassette of the Cap gene can use the abovementioned expression cassette, or one of them can use the abovementioned expression cassette.

In some embodiments, the exogenous gene is a reporter gene and is, for example, a chloramphenicol acetyltransferase encoding gene, a β -galactosidase encoding gene, a β -glucuronidase encoding gene, a *Renilla* luciferase encoding gene, an alkaline phosphatase encoding gene, a firefly luciferase encoding gene, a green fluorescent protein encoding gene, or a red fluorescent protein encoding gene.

In some other embodiments, the exogenous gene is a gene encoding a drug polypeptide, and the drug polypeptide is lipoprotein esterase, apolipoprotein, cytokine, interleukin, or interferon and the like.

In some embodiments, the insect cell is a *Spodoptera frugiperda* cell, a *Trichopodia* cell, a *Drosophila* cell, or a mosquito cell and the like.

Example 1

Step One: Construction of Rep Gene Expression Cassette I and Recombinant Baculovirus Vector Containing Rep Gene Expression Cassette I

With reference to FIG. 2, herein, content A is a schematic diagram of a DNA chain of the Rep gene expression cassette

12

I, content B is a schematic diagram of the translation and expression of the Rep52 protein when the intron is not spliced during post-transcriptional processing driven by the first promoter, content C is a schematic diagram of the translation and expression of the Rep78 protein after the intron is spliced during post-transcriptional processing driven by the first promoter, and content D is a schematic diagram of the translation and expression of the Rep78 protein after transcription driven by the second promoter. The gene expression cassette, from the 5' end to the 3' end, includes in sequence a first promoter sequence, adenine nucleotide (A), a 5' end donor sequence of the intron, a second promoter sequence, a 3' end acceptor sequence of the intron, thymine nucleotides, guanine nucleotides (TG), and a nucleotide sequence encoding the AAV serotype 2 Rep protein lacking only the translation initiation codons ATG of the Rep78 protein. The overlapping open reading frame nucleotide sequence used to encode the Rep protein is shown in SEQ ID No. 1, its entirety is used to encode the Rep78 protein, and the part used to encode the Rep52 protein is shown in SEQ ID No. 2. There are multiple nucleotide mutations between a Rep78 translation initiation codon and a Rep52 translation initiation codon to eliminate possible translation initiation sites in this region. Herein, deltaIE1 is used as the first promoter sequence and polH is used as the second promoter sequence in this example. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain a construct RepA, whose nucleotide sequence is shown in SEQ ID No. 3.

The construct is cloned into a pFastBac vector to prepare a transfer plasmid. The transfer plasmid is transformed into a DH10Bac strain, and a recombinant baculovirus vector AcRepA containing RepA is obtained through the gene transposition reaction mediated by Tn7 transposase.

Step Two: Construction of Cap Gene Expression Cassette

Construction of a Cap8 expression cassette containing the AAV serotype 8 Cap gene: The Cap gene expression cassette is constructed with reference to the method described in Example 3 of Chinese patent (CN202111105263.5). With reference to FIG. 3, the Cap8 expression cassette, from 5' to 3', includes in sequence, a p10 promoter, an artificially constructed sequence, and a nucleotide sequence encoding the AAV serotype 8 VP protein lacking only the translation initiation codons ATG of the VP1 protein. Herein, the artificially constructed sequence contains the translation initiation codons ATG and contains two intron alternative splicing sites. The amino terminus of the AAV serotype 8 VP protein is encoded with at least one nucleotide mutation to eliminate possible splicing sites within the coding sequence. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain a construct Cap8 expression cassette, whose nucleotide sequence is shown in SEQ ID No.4.

Step Three: Construction of Recombinant AAV Bacmid

The method for constructing recombinant AAV bacmid for producing AAV virus in an insect cell is provided with reference to Example 1 in the previously applied patent CN112553257A in this example, and the following steps are included.

13

(1) A homologous recombination vector containing the essential functional elements Cap and Rep gene expression cassettes of AAV is constructed. The essential functional elements in the homologous recombination vector are then inserted into the C-terminus of the essential gene Ac135 (116492 . . . 117391) in the baculovirus genome in *E. coli* through Red homologous recombination. A recombinant baculovirus vector containing Cap and Rep gene expression cassettes is obtained, numbered Bac-Cap-Rep.

(2) A homologous recombination vector containing an ITR core element (ITR-GOI) is constructed. The ITR core element is numbered I-G-1, and its nucleotide sequence is shown in SEQ ID No. 5. The GOI in the ITR core element uses a red fluorescent protein mcherry gene expression cassette, that is, the miniEfla promoter controls mcherry expression to facilitate detection of rAAV activity in this example.

(3) The shuttle vector constructed in the above step (2) is used to transform competent cells containing the Bac-Cap-Rep recombinant bacmid. The essential functional elements in the homologous recombination vector are then inserted into the C-terminus of the essential gene Ac98 (88502 . . . 88464) in the baculovirus genome in *E. coli* through Red homologous recombination. A recombinant bacmid containing the functional protein components necessary for rAAV production and the ITR core element of the recombinant baculovirus genome is finally obtained, numbered Bac-Cap-Rep-ITR-GOI.

Step Four: Preparation of AAV Recombinant Baculovirus

The recombinant AAV bacmid prepared in step three is transfected into a host cell line and cultured to obtain the AAV recombinant baculovirus, and the specific steps are provided as follows.

The above recombinant bacmid DNA was extracted and transfected into Sf9 insect cells to prepare recombinant baculovirus BEV and rAAV. The transfected Sf9 insect cells successfully produced BEV, and further infection with a large number of replicating and proliferating BEV caused obvious cytopathic effect (CPE) in Sf9 cells. The culture supernatant of Sf9 cells that had undergone CPE was collected, which contained a large amount of BEV, which was the 0th generation BEV (P0). At the same time, the Sf9 cells containing a large amount of rAAV were collected. The prepared BEV-P0 was infected into suspension-cultured Sf9 cells at a multiplicity of infection (MOI=1). After 72 hours of infection, the cell activity dropped to less than 50%. The cell culture medium was centrifuged at 1000 g for 5 minutes, and the culture supernatant and cell pellet were collected. The supernatant was labeled as the first generation BEV-P1, and the cells were labeled as rAAV packaged with BEV-P0.

Examples 2 to 4

In Examples 2 to 4, deltaIE1 in Example 1 is replaced with 39K, p6.9, and p10 as the first promoter sequence, and other steps are the same as in Example 1. Constructs RepB, RepC, and RepD, recombinant baculovirus vectors AcRepB, AcRepC, and AcRepD, as well as subsequent recombinant AAV bacmid and AAV recombinant baculovirus are obtained. The nucleotide sequences of RepB-RepD are shown in SEQ ID No. 6-8.

14

Example 5

Step One: Construction of Rep Gene Expression Cassette II and Recombinant Baculovirus Vector Containing Rep Gene Expression Cassette II

With reference to what is shown in FIG. 4, herein, content A is a schematic diagram of a DNA chain of the Rep gene expression cassette II, content B is a schematic diagram of the translation and expression of the Rep52 protein when the intron is not spliced during post-transcriptional processing driven by the first promoter, content C is a schematic diagram of the translation and expression of the Rep78 protein after the intron is spliced during post-transcriptional processing driven by the first promoter, and content D is a schematic diagram of the translation and expression of the Rep52 protein after splicing of the first intron splicing donor and splicing acceptor during post-transcriptional processing driven by the second promoter. A TAA translation termination sequence is inserted between the first promoter and the second promoter. During post-transcriptional processing driven by the second promoter, if the intron is not spliced or the second intron splicing donor and splicing acceptor are spliced, protein translation will be terminated prematurely and the Rep78 protein will not be produced. The gene expression cassette, from the 5' end to the 3' end, includes in sequence a second promoter sequence, a 5' end donor sequence of the first intron, a first promoter sequence, adenine nucleotide (A), a 5' end donor sequence of the second intron, a 3' end acceptor sequence of the intron, thymine nucleotides, guanine nucleotides (TG), and a nucleotide sequence encoding the AAV serotype 2 Rep protein lacking only the translation initiation codons ATG of the Rep78 protein. Herein, deltaIE1 is used as the first promoter sequence and polH is used as the second promoter sequence in this example. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain a construct RepI, whose nucleotide sequence is shown in SEQ ID No. 9.

The construct was cloned into a pFastBac vector to prepare a transfer plasmid. The transfer plasmid was transformed into a DH10Bac strain, and a recombinant baculovirus vector AcRepI containing RepI was obtained through the gene transposition reaction mediated by Tn7 transposase.

Step two to step four are the same as in Example 1.

Examples 6 to 8

In Examples 6 to 8, deltaIE1 in Example 5 is replaced with 39K, p6.9, and p10 as the first promoter sequence, and other steps are the same as in Example 5. Constructs RepII, RepIII, and RepIV (the nucleotide sequences of which are shown in SEQ ID No. 10-12), recombinant baculovirus vectors AcRepII, AcRepIII, and AcRepIV as well as subsequent recombinant AAV bacmid and AAV recombinant baculovirus are obtained.

Comparative Examples 1 to 4

Step one: A Rep gene expression cassette III and a recombinant baculovirus vector containing the Rep gene expression cassette III are constructed, and the Rep gene expression cassette III is constructed with reference to the method described in Example 6 of Chinese patent (CN202111105263.5). With reference to FIG. 5, herein, content A is a schematic diagram of a DNA chain of the Rep

15

gene expression cassette III, content B is a schematic diagram of the translation and expression of the Rep52 protein when the intron is not spliced during post-transcriptional processing, and content C is a schematic diagram of the translation and expression of the Rep78 protein after the intron is spliced during post-transcriptional processing. The gene expression cassette, from the 5' end to the 3' end, includes in sequence a promoter sequence, adenine nucleotide (A), a 5' end donor sequence of the intron, a 3' end acceptor sequence of the intron, thymine nucleotides, guanine nucleotides (TG), and a nucleotide sequence encoding the AAV serotype 2 Rep protein lacking the translation initiation codons ATG of the Rep78 protein. Herein, Comparative Examples 1 to 4 use deltaIE1, 39K, p6.9, and polH as promoter sequences. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain the constructs Rep1-4, whose nucleotide sequences are shown in SEQ ID No. 13-16.

The constructs were cloned into a pFastBac vector to prepare a transfer plasmid. The transfer plasmid was transformed into a DH10Bac strain, and recombinant baculovirus vectors AcRep1-4 respectively containing Rep1-4 were obtained through the gene transposition reaction mediated by Tn7 transposase.

Step two to step four are the same as in Example 1.

Experimental Result Verification

For the ease of explanation of the experimental results, the recombinant AAV bacmids (Bac-Cap-Rep-ITR-GOI) constructed according to Examples 1 to 6 and Comparative Example 4 of the disclosure are numbered according to Table 1 according to their corresponding Cap gene expression cassette, Rep gene expression cassette, and ITR core element. The 7 different recombinant AAV bacmids constructed in this example all contain the Cap gene expression cassette, the Rep gene expression cassette, and the ITR core element. Herein, the recombinant AAV bacmid CRI-0 is a control bacmid containing the Rep4 gene expression cassette, and the Rep78/52 protein in the Rep4 gene expression cassette is driven by a single very late strong promoter polH. The details are shown in Table 1.

TABLE 1

List of components of 7 different recombinant AAV bacmids constructed in the Examples and Comparative Examples			
Recombinant AAV bacmid (Bac-Cap-Rep-ITR-GOI) number	Cap gene expression cassette	Functional protein components and ITR core elements necessary for rAAV production	
		Rep gene expression cassette	ITR core component (ITR-GOI)
CRI-0	Cap8	Rep4	I-G-1
CRI-1	Cap8	RepA	I-G-1
CRI-2	Cap8	RepB	I-G-1
CRI-3	Cap8	RepC	I-G-1
CRI-4	Cap8	RepD	I-G-1
CRI-5	Cap8	RepI	I-G-1
CRI-6	Cap8	RepII	I-G-1

Verification Example 1 detects the expression of the Rep proteins (Rep78 and Rep52), and the specific steps are as follows.

The above recombinant baculovirus vector DNA was extracted and transfected into Sf9 insect cells to prepare the recombinant baculovirus BEV. The transfected Sf9 insect

16

cells successfully produced BEV, and further infection with a large number of replicating and proliferating BEV caused obvious cytopathic effect (CPE) in Sf9 cells. The culture supernatant of Sf9 cells that had undergone CPE was collected, which contained a large amount of BEV, which was the 0th generation BEV (P0). At the same time, the Sf9 cells containing a large amount of rAAV were collected. The prepared BEV-P0 was infected into suspension-cultured Sf9 cells at a multiplicity of infection (MOI=1). After 72 hours of infection, the cell activity dropped to less than 50%. The cell culture medium was centrifuged at 1000 g for 5 minutes, and the culture supernatant and cell pellet were collected. The supernatant was labeled as the first generation BEV-P1. Continuing to expand the culture, the prepared BEV-P1 was infected into suspension-cultured Sf9 cells at a multiplicity of infection (MOI=1). After 72 hours of infection, the cell activity dropped to less than 50%. The cell culture medium was centrifuged at 1000 g for 5 minutes, and the cell pellet was collected for Western Blot to detect the expression of the Rep proteins (Rep78 and Rep52).

FIGS. 6 to 8 are Western Blot detection diagrams of the Rep proteins (Rep78 and Rep52) of the recombinant baculovirus vector containing the Rep gene expression cassette in Examples 1 to 8 and Comparative Examples 1 to 4.

FIG. 6 is a Western Blot detection diagram of the Rep proteins (Rep78 and Rep52) of the recombinant baculovirus vectors AcRepA-D containing the Rep gene expression cassette I, in which lanes 1~4 represent AcRepA-D respectively. It can be seen from FIG. 6 that the above recombinant baculovirus vectors can produce the Rep78 and Rep52 proteins, and the expression amount of the Rep52 protein is significantly higher than that of the Rep78 protein. The high stoichiometric ratio of Rep52 to Rep78 may be a key factor for high rAAV yield. In constructs AcRepA, AcRepB, AcRepC, and AcRepD, the inventors insert the second promoter into the intron sequence to relatively enhance the expression of the Rep52 protein.

FIG. 7 is a Western Blot detection diagram of the Rep proteins (Rep78 and Rep52) of the recombinant baculovirus vectors AcRepI, AcRepII, AcRepIII, and AcRepIV containing the Rep gene expression cassette II, in which lanes 1-4 represent AcRepI-IV respectively. In constructs AcRepI, AcRepII, AcRepIII, and AcRepIV, the inserted second promoter drives the splicing interaction between the first intron splicing donor and the splicing acceptor during post-transcriptional processing to enhance the expression of the Rep52 protein.

FIG. 8 shows the recombinant baculovirus vectors AcRep1-4 containing the Rep gene expression cassette III in Comparative Examples 1-4, in which lanes 1-4 respectively represent the AcRep1-4 in Comparative Examples. It can be seen from the figure that the above recombinant baculovirus vectors can all produce the Rep78 and Rep52 proteins. In constructs AcRep1, AcRep2, AcRep3, and AcRep4, the Rep78/52 protein is expressed through a single promoter. Herein, AcRep1, AcRep2, and AcRep3 use deltaIE1, 39K, and p6.9 promoters to drive the expression of Rep78/52 protein in the immediate early, delayed early, and late stages of baculovirus respectively. Early expression of the Rep78 protein may be beneficial for AAV genome replication and packaging. However, compared with the very late promoter (p10 or polH), the lower promoter strength of deltaIE1, 39K, and p6.9 results in insufficient Rep52 protein, which is not conducive to high rAAV productivity, so it is necessary to enhance the expression of the Rep52 protein.

Verification Example 2 Growth Kinetics Analysis

The immediate early promoter deltaIE1 is active early in baculovirus infection, and early and abundant expression of

the Rep78 protein in insect cells may have a negative impact on the yield of recombinant baculovirus (Urabe et al., 2006, Journal of Virology 80.4:1874-1885). In order to compare the growth kinetics of the recombinant baculoviruses containing different Rep gene expression cassettes in the above Examples and Comparative Examples, the above recombinant baculovirus is used to infect Sf9 cells at a multiplicity of infection (MOI=0.5). The supernatant of the infected cell culture is collected at 24, 48, 72, and 96 hours after infection, and the titer of the second generation BEV-P2 is determined using fluorescence quantitative PCR (qPCR), where the titer unit is expressed in VG/ml (VG, virus genomes). A pair of primers (Q-GP64-F: AACCTGGACAT-TACCCCGCC (SEQ ID NO: 24) and Q-GP64-R: CCGTTGTACGCATACGCCTG (SEQ ID NO: 25)) corresponding to the gp64 gene is used to determine the BEV titer, and titer unit is expressed in VG/ml (VG, virus genomes). The experimental results are shown in FIG. 9. In constructs CRI-0, CRI-1, and CRI-2, the Rep78 protein is expressed in the very late stage, immediate early stage, and delayed early stage respectively. As can be seen from FIG. 9, there are no significant differences in viral titers among CRI-0, CRI-1, and CRI-2, indicating that the early expression of the Rep78 protein in CRI-1 and CRI-2 constructs has less impact on baculovirus replication. In the Rep expression cassette designed by Urabe et al., deltaIE1 alone drives the expression of the Rep78 protein. In CRI-1, the rep gene driven by deltaIE1 promoter is partially transcribed and translated into the Rep78 protein through intron splicing, and partially transcribed and translated into the Rep52 protein, resulting in a relatively lower expression amount of the Rep78 protein, which may be the reason for its reduced impact on baculovirus replication.

Verification Example 3: Purification of Recombinant AAV Virus Particles and Detection of rAAV Virus Titer and Virus Particle Solid Rate

This verification example uses Q-PCR to detect the titer of the harvested rAAV virus, and the titer unit is expressed in VG/ml (VG, virus genomes). rAAV titers are measured using a pair of primers (Q-ITR-F: GGAACCCCTAGT-GATGGAGTT (SEQ ID NO: 26) and Q-ITR-R: CGGCCTCAGTGAGCGA (SEQ ID NO: 27)) targeting the ITR sequence. The specific operation is as follows. The culture was continuously expanded according to step four of Example 1 until the Sf9 cells cultured in the suspension were infected with BEV-P2 seed virus according to the multiplicity of infection (MOI=1) for rAAV packaging. The packaging volume is 300 mL to 400 mL. After 3 days of infection, cell activity was monitored. After the activity was lower than 50%, the cell culture was harvested, 500 μ l of the harvested mixed solution (mixed solution of cells and supernatant) was put into a 1.5 ml EP tube, frozen and thawed four times in liquid nitrogen, and treated in a 37° C. water bath. 0.1 μ l of Benzonase was added to the 500 μ l mixed solution, and the mixed solution was mixed in a 37° C. water bath for 1 hour and then in a 95° C. water bath for 10 minutes. Centrifugation was performed at 2500 g for 10 minutes, and the supernatant was collected in a new 1.5 ml EP tube. 200 μ l of the supernatant was added to 10 μ l of 10% SDS solution to a final concentration of 0.5% SDS. 1.2 μ l Proteinase K (working concentration 112 μ g/ml) was then added, the final volume was approximately 210 μ l, and a 55° C. water bath was performed for 1 hour. After a short centrifugation, a 95° C. water bath was performed for 10 minutes. After mixing, 5 μ l of sample was taken out and 145 μ l of ddH₂O was added

to make the final volume 150 μ l. After mixing, 10 μ l of the sample was taken out and 90 μ l of ddH₂O was added to make the final volume 100 μ l. 2 μ l was used as a Q-PCR template to measure the titer of the virus sample. The experimental results are shown in Table 2.

This example uses analytical ultracentrifugation technology (abbreviated as AUC) to detect the solid rate (complete capsid: total capsid) of the recombinant adeno-associated virus (rAAV).

With the use of the sedimentation rate analysis method in the analytical ultracentrifugation technology, the sample is rotated at high speed in the centrifuge, and each component in the sample moves to the bottom of the sample pool. Different components have different movement rates (i.e., sedimentation coefficients) and have different times to fall to the bottom during centrifugation. During this process, the sample distribution state is scanned, and the sedimentation properties of different components are obtained by analyzing its change process over time. The sedimentation coefficient of each component of the sample depends on the molecular weight, molecular shape, and conformation. AUC is generally used to characterize the solid rate of rAAV.

The collected cell culture was purified using an affinity chromatography column (AVIPure AAV8 Resin). 400 μ l of the purified rAAV sample was diluted to a dilution factor of OD₂₃₀=0.90, and loaded onto the sample. The sedimentation coefficient and peak area percentage of each peak were recorded. Herein, the sedimentation coefficient of empty shell virus was 55S to 65S, the sedimentation coefficient of solid virus was 80S to 110S, 65S to 80S was the partial part, and greater than 110S was the aggregate. The peaks with sedimentation coefficients between 20S and 150S were normalized to obtain the peak area percentage of each peak, and the empty and solid rate of the virus sample was reported based on the results. The experimental results are shown in Table 2.

TABLE 2

Titer and solid rate detection results of rAAV virions produced using 7 different recombinant AAV bacmids		
Recombinant AAV bacmid number	rAAV virus titer (VG/ml)	rAAV solid rate intact capsids: total capsids)
CRI-0	1.12E+12	24%
CRI-1	1.23E+12	37%
CRI-2	1.37E+12	45%
CRI-3	7.37E+11	28%
CRI-4	6.41E+11	27%
CRI-5	1.20E+12	32%
CRI-6	1.27E+12	43%

As can be seen from Table 2, the titers of rAAV virus particles produced using recombinant AAV bacmids CRI-1, CRI-2, CRI-5, and CRI-6 are higher than those of the control bacmid CRI-0, and the solid rate is significantly higher than that of the control bacmid CRI-0. Compared to the control bacmid CRI-0, the early expression of the Rep78 protein and the higher Rep52/Rep78 ratio in bacmids CRI-1, CRI-2, CRI-5, and CRI-6 may be the key factors for their high titers and the high solid rate. The titers of rAAV virus particles produced using the recombinant AAV bacmids CRI-3 and CRI-4 are lower than that of the control bacmid CRI-0, but their solid rates are higher than that of the control bacmid CRI-0. Bacmids CRI-0, CRI-3, and CRI-4 use polH, p6.9, and p10 late promoters respectively to drive the expression of the Rep78 protein, where the polH promoter is the most

active. Compared to the control bacmid CRI-0, the lower Rep78 protein expression and higher Rep52/Rep78 ratio in bacmids CRI-3 and CRI-4 may be responsible for their relatively low titers and high solid rates. Increasing the solid rates of viral particles is beneficial to the rAAV product, so fewer particles can be used to obtain a similar amount of genome copies per kilogram, and it is also beneficial to establish a stable downstream purification process.

Example 9

Step One: Construction of Rep Gene Expression Cassette IV and Recombinant Baculovirus Vector Containing Rep Gene Expression Cassette IV

With reference to FIG. 10, herein, content A is a schematic diagram of a DNA chain of the Rep gene expression cassette IV, content B is a schematic diagram of the translation and expression of the Rep78 protein after the intron is spliced during post-transcriptional processing driven by the first promoter, and content C is a schematic diagram of the translation and expression of the Rep78 protein after transcription driven by the second promoter. The gene expression cassette, from the 5' end to the 3' end, includes in sequence a first promoter sequence, adenine nucleotide (A), a 5' end donor sequence of the intron, a second promoter sequence, a 3' end acceptor sequence of the intron, thymine nucleotides, guanine nucleotides (TG), and a nucleotide sequence encoding the AAV serotype 2 Rep protein lacking only the translation initiation codons ATG of the Rep78 protein. The overlapping open reading frame nucleotide sequence used to encode the Rep protein is shown in SEQ ID No. 1, its entirety is used to encode the Rep78 protein, and the part used to encode the Rep52 protein is shown in SEQ ID No. 2. There are multiple nucleotide mutations between a Rep78 translation initiation codon and a Rep52 translation initiation codon to eliminate possible translation initiation sites in this region. In this example, deltaIE1 is used as the first promoter sequence (the nucleotide sequence is shown in SEQ ID No. 17), and polH is used as the second promoter sequence (the nucleotide sequence is shown in SEQ ID No. 18). Herein, the second promoter contains the translation initiation and stop codon sequences that are recognized by a ribosome, so that when the intron is not spliced during the post-transcriptional processing of a Rep gene expression cassette driven by the first promoter, the Rep52 protein will not be translated and produced. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain a construct RepE, whose nucleotide sequence is shown in SEQ ID No. 19.

The construct was cloned into a pFastBac vector to prepare a transfer plasmid. The transfer plasmid was transformed into a DH10Bac strain, and a recombinant baculovirus vector AcRepE containing RepE was obtained through the gene transposition reaction mediated by Tn7 transposase.

Step two: Construction of Cap gene expression cassette

This step is the same as step two of Example 1.

Step three: Construction of recombinant AAV bacmid

This step is the same as step three of Example 1.

Step four: Preparation of AAV recombinant baculovirus

This step is the same as step four of Example 1.

Examples 10 to 12

In Examples 10 to 12, deltaIE1 in Example 9 is replaced with 39K (SEQ ID No. 20), p6.9 (SEQ ID No. 21), and p10 (SEQ ID No. 22) as the first promoter sequence, and other steps are the same as in Example 9. Constructs RepF, RepG, and RepH, as shown in FIG. 11, corresponding recombinant baculovirus vectors AcRepF, AcRepG, and AcRepH, as well as subsequent recombinant AAV bacmid and AAV recombinant baculovirus are obtained.

Comparative Example 5

Step one: A Rep gene expression cassette V and a recombinant baculovirus vector containing the Rep gene expression cassette V are constructed, and the Rep gene expression cassette V is constructed with reference to the method described in Example 6 of Chinese patent (CN202111105263.5). With reference to FIG. 5, herein, content A is a schematic diagram of a DNA chain of the Rep gene expression cassette V, content B is a schematic diagram of the translation and expression of the Rep52 protein when the intron is not spliced during post-transcriptional processing, and content C is a schematic diagram of the translation and expression of the Rep78 protein after the intron is spliced during post-transcriptional processing. The gene expression cassette, from the 5' end to the 3' end, includes in sequence a promoter sequence, adenine nucleotide (A), a 5' end donor sequence of the intron, a 3' end acceptor sequence of the intron, thymine nucleotides, guanine nucleotides (TG), and a nucleotide sequence encoding the AAV serotype 2 Rep protein lacking the translation initiation codons ATG of the Rep78 protein. Herein, Comparative Example 5 uses polH as the promoter sequence. The above sequences are connected through artificial direct synthesis or overlap extension PCR amplification to obtain a construct Rep5, whose nucleotide sequence is shown in SEQ ID No. 23.

The construct was cloned into a pFastBac vector to prepare a transfer plasmid. The transfer plasmid was transformed into a DH10Bac strain, and a recombinant baculovirus vector AcRep5 containing Rep5 was obtained through the gene transposition reaction mediated by Tn7 transposase.

Step two to step four are the same as in Example 9.

Experimental Result Verification

For the ease of explanation of the experimental results, the recombinant AAV bacmids (Bac-Cap-Rep-ITR-GOI) constructed according to Examples 9 to 12 and Comparative Example 5 of the disclosure are numbered according to Table 3 according to their corresponding Cap gene expression cassette, Rep gene expression cassette, and ITR core element. The 5 different recombinant AAV bacmids constructed in this example all contain the Cap gene expression cassette, the Rep gene expression cassette, and the ITR core element. Herein, the recombinant AAV bacmid CRI-7 is a control bacmid containing the Rep5 gene expression cassette, and the Rep78/52 protein in the Rep5 gene expression cassette is driven by a single very late strong promoter polH. The details are shown in Table 3.

TABLE 3

List of components of 7 different recombinant AAV bacmids constructed in the Examples and Comparative Example 5				
Corresponding	Recombinant AAV bacmid	Functional protein components and ITR core elements necessary for rAAV production		
		Cap gene expression cassette	Rep gene expression cassette	ITR core component (ITR-GOI)
Comparative Example 5	CRI-7	Cap8	Rep5	I-G-1
Example 9	CRI-8	Cap8	RepE	I-G-1
Example 10	CRI-9	Cap8	RepF	I-G-1
Example 11	CRI-10	Cap8	RepG	I-G-1
Example 12	CRI-11	Cap8	RepH	I-G-1

Verification Example 4

The specific steps of the detection of expression of the Rep proteins (Rep78 and Rep52) are as follows.

The above recombinant baculovirus vector DNA was extracted and transfected into Sf9 insect cells to prepare the recombinant baculovirus BEV. The transfected Sf9 insect cells successfully produced BEV, and further infection with a large number of replicating and proliferating BEV caused obvious cytopathic effect (CPE) in Sf9 cells. The culture supernatant of Sf9 cells that had undergone CPE was collected, which contained a large amount of BEV, which was the 0th generation BEV (P0). At the same time, the Sf9 cells containing a large amount of rAAV were collected. The prepared BEV-P0 was infected into suspension-cultured Sf9 cells at a multiplicity of infection (MOI=1). After 72 hours of infection, the cell activity dropped to less than 50%. The cell culture medium was centrifuged at 1000 g for 5 minutes, and the culture supernatant and cell pellet were collected. The supernatant was labeled as the first generation BEV-P1. Continuing to expand the culture, the prepared BEV-P1 was infected into suspension-cultured Sf9 cells at a multiplicity of infection (MOI=1). After 72 hours of infection, the cell activity dropped to less than 50%. The cell culture medium was centrifuged at 1000 g for 5 minutes, and the cell pellet was collected for Western Blot to detect the expression of the Rep proteins (Rep78 and Rep52).

FIGS. 12 to 13 are Western Blot detection diagrams of the Rep proteins (Rep78 and Rep52) of the recombinant baculovirus vector containing the Rep gene expression cassette in Examples 9 to 12 and Comparative Example 5.

FIG. 12 is a Western Blot detection diagram of the Rep proteins (Rep78 and Rep52) of the recombinant baculovirus vectors AcRepE-H containing the Rep gene expression cassette IV, in which lanes 1-4 represent AcRepE-H respectively. It can be seen from FIG. 12 that the above recombinant baculovirus vectors can produce the Rep78 and Rep52 proteins, and the expression amount of the Rep52 protein of AcRepE-G is significantly higher than that of the Rep78 protein. The high stoichiometric ratio of Rep52 and Rep78 proteins may be a key factor in the high yield of rAAV.

FIG. 13 shows the recombinant baculovirus vector AcRep5 containing the Rep gene expression cassette V in Comparative Example 5, in which lane 1 represents the AcRep5 in Comparative Example 5. It can be seen from the figure that the above recombinant baculovirus vectors can produce the Rep78 and Rep52 proteins.

Verification Example 5

Purification of Recombinant AAV Virus Particles and Detection of Raav Virus Titer and Virus Particle Solid Rate

This verification example uses Q-PCR to detect the titer of the harvested rAAV virus, and the titer unit is expressed in VG/ml (VG, virus genomes). rAAV titers are measured using a pair of primers (Q-ITR-F: GGAACCCCTAGT-GATGGAGTT (SEQ ID NO: 26) and Q-ITR-R: CGGCCTCAGTGAGCGA (SEQ ID NO: 27)) targeting the ITR sequence. The specific operation is as follows. The culture was continuously expanded according to step four of Example 9 until the Sf9 cells cultured in the suspension were infected with BEV-P2 seed virus according to the multiplicity of infection (MOI=1) for rAAV packaging. The packaging volume is 300 mL to 400 mL. After 3 days of infection, cell activity was monitored. After the activity was lower than 50%, the cell culture was harvested, 500 μ l of the harvested mixed solution (mixed solution of cells and supernatant) was put into a 1.5 ml EP tube, frozen and thawed four times in liquid nitrogen, and treated in a 37° C. water bath. 0.1 μ l of Benzonase was added to the 500 μ l mixed solution, and the mixed solution was mixed in a 37° C. water bath for 1 hour and then in a 95° C. water bath for 10 minutes. Centrifugation was performed at 2500 g for 10 minutes, and the supernatant was collected in a new 1.5 ml EP tube. 200 μ l of the supernatant was added to 10 μ l of 10% SDS solution to a final concentration of 0.5% SDS. 1.2 μ l Proteinase K (working concentration 112 μ g/ml) was then added, the final volume was approximately 210 μ l, and a 55° C. water bath was performed for 1 hour. After a short centrifugation, a 95° C. water bath was performed for 10 minutes. After mixing, 5 μ l of sample was taken out and 145 μ l of ddH₂O was added to make the final volume 150 μ l. After mixing, 10 μ l of the sample was taken out and 90 μ l of ddH₂O was added to make the final volume 100 μ l. 2 μ l was used as a Q-PCR template to measure the titer of the virus sample. The experimental results are shown in Table 4.

This example uses a technique of analytical ultracentrifugation (abbreviated as AUC) to detect the solid rate (complete capsid: total capsid) of the recombinant adeno-associated virus (rAAV).

Using the sedimentation rate analysis method in the analytical ultracentrifugation technology, the sample is rotated at high speed in the centrifuge, and each component in the sample moves to the bottom of the sample pool. Different components have different movement rates (i.e., sedimentation coefficients) and have different times to fall to the bottom during centrifugation. During this process, the sample distribution state is scanned, and the sedimentation properties of different components are obtained by analyzing its change process over time. The sedimentation coefficient of each component of the sample depends on the molecular weight, molecular shape, and conformation. AUC is generally used to characterize the solid rate of rAAV.

The collected cell culture was purified using an affinity chromatography column (AVIPure AAV8 Resin). 400 μ l of the purified rAAV sample was diluted to a dilution factor of OD₂₃₀=0.90, and loaded onto the sample. The sedimentation coefficient and peak area percentage of each peak were recorded. Herein, the sedimentation coefficient of empty shell virus was 55S to 65S, the sedimentation coefficient of solid virus was 80S to 110S, 65S to 80S was the partial part, and greater than 110S was the aggregate. The peaks with sedimentation coefficients between 20S and 150S were

23

normalized to obtain the peak area percentage of each peak, and the empty and solid rate of the virus sample was reported based on the results. The experimental results are shown in Table 4.

TABLE 4

Titer and solid rate detection results of rAAV virions produced using 5 different recombinant AAV bacmids			
Corresponding Example or Comparative Example	Recombinant bacmid AAV number	rAAV virus titer (VG/ml)	rAAV solid rate (intact capsids: total capsids)
Comparative Example 5	CRI-7	1.12E+12	23%
Example 9	CRI-8	1.23E+12	35%
Example 10	CRI-9	1.37E+12	39%
Example 11	CRI-10	7.37E+11	28%
Example 12	CRI-11	1.08E+12	20%

As can be seen from Table 4, the titers of rAAV virus particles produced using recombinant AAV bacmids CRI-8 and CRI-9 are higher than those of the control bacmid CRI-7, and the solid rate is significantly higher than that of the control bacmid CRI-7. Compared to the control bacmid CRI-7, the early expression of the Rep78 protein and the higher Rep52/Rep78 ratio in bacmids CRI-8 and CRI-9 may

24

be the key factors for their high titers and the high solid rate. The titer of rAAV virus particles produced using the recombinant AAV bacmid CRI-10 is lower than that of the control bacmid CRI-0, but its solid rate is higher than that of the control bacmid CRI-0. Bacmids CRI-7 and CRI-10 use polH and p6.9 late promoters respectively to drive the expression of the Rep78 protein, where the polH promoter is the most active. Compared to the control bacmid CRI-7, the lower Rep78 protein expression and higher Rep52/Rep78 ratio in the bacmid CRI-10 may be responsible for its relatively low titer and high solid rate. The titer of rAAV virus particles produced using the recombinant AAV bacmid CRI-11 is comparable to that of the control bacmid CRI-0, but its solid rate is slightly lower than that of the control bacmid CRI-7, possibly due to its use of a relatively weak very late promoter to drive Rep78 expression. Increasing the solid rates of viral particles is beneficial to the rAAV product, so fewer particles can be used to obtain a similar amount of genome copies per kilogram, and it is also beneficial to establish a stable downstream purification process.

A person having ordinary skill in the art should be able to easily understand that the above description is only preferred embodiments of the disclosure and is not intended to limit the disclosure. Any modifications, equivalent replacements, and modifications made without departing from the spirit and principles of the disclosure should fall within the protection scope of the disclosure.

SEQUENCE LISTING

```

Sequence total quantity: 23
SEQ ID NO: 1          moltype = DNA length = 1866
FEATURE               Location/Qualifiers
source                1..1866
                     mol_type = other DNA
                     note = artificial sequence
                     organism = unidentified

SEQUENCE: 1
atgccgggggt tttacgagat tgtgattaag gtccccagcg accttgacgg gcactctgccc 60
ggcattttctg acagcttttgt gaactgggtg gccgagaagg agtgggagtt gccgccagat 120
tctgacttgg atctgaatct gattgagcag gcacccctga cgtgggcccga gaagctgcag 180
cgcgactttc tgacggagtg gcgccgtgtg agtaaggccc cggaggccctt tttctttgtg 240
caatttgaga agggagagag ctacttccac ttacacgtgc tcgtggaaac caccgggggtg 300
aaatccttag ttttgggacg tttcctgagt cagattcgcg aaaaactgat tcagagaatt 360
taccgcggga tcgagccgac tttgccaaac tggttccgcg tcacaaagac cagaaacggc 420
gccggaggcg ggaacaagggt ggtggacgag tgctacatcc ccaattactt gctcccaaa 480
accagcctg agctccagtg ggctgggact aatttagaac agtatttaag cgctgtttg 540
aatctcacgg agcgtaaacg gttggtggcg cagcatctga cgacgtgtc gcagacgcag 600
gagcagaaca aagagaatca gaatcccaat tctgacgcgc cgtggtatcg atcaaaaact 660
tcagccaggt acatggagct ggtcgggttg ctctgggaca aggggattac ctcgagagaag 720
cagtggatcc aggaggacca ggcctcatac atctccttca atgcggcctc caactcgcgg 780
tcccaaatca aggctgcctt ggacaatgcg ggaagatta tgagcctgac taaaaccgcc 840
cccgactacc tgggtgggcca gcagcccgctg gaggacattt ccagcaatcg gatttataaa 900
attttggaac taaacgggta cgatcccaaa tatgcggcct cgtctttctt gggatggggc 960
acgaaaaagt tcggcaagag gaacaccatc tggctgtttg ggctgcaac taccgggaag 1020
accaacatcg cggaggccat agcccacact gtgcccttct acgggtgcgt aaactggacc 1080
aatgagaact ttcccttcaa cgactgtgtc gacaagatgg tgatctgggt ggaggagggg 1140
aagatgaccg ccaaggtcgt ggagtgcgcc aaagccattc tcggaggaag caaggtgcgc 1200
gtggaccaga aatgcaagtc ctccgcccag atagaccgga ctcccgatg cgtcacctcc 1260
aacaccaaca tgtgcgcgtg gattgacggg aactcaacga ctttcgaaca ccagcagccg 1320
ttgcaagacc ggatgttcaa atttgaactc acccgccgtc tggatcatga ctttgggaag 1380
gtcaccgaag aggaagtcaa agacttttct cgggtgggcaa aggatcacgt ggttgaggtg 1440
gagcatgaat tctacgtcaa aaagggtgga gccaaagaaa gaccgcgccc cagtgcgcga 1500
gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc agccatcgac gtcagacgcg 1560
gaagcttcga tcaactacgc agacaggtag caaaacaaat gttctcgtca cgtgggcatg 1620
aatctgatgc tgtttccctg cagacaatgc gagagaatga atcagaatc aaatatctgc 1680
ttactcacg gacagaaga ctgtttagag tgctttcccg tgcagaatc tcaaccgctt 1740
tctgtcgtca aaaaggcgtg tcgaaactg tgctacatc atcatatcat gggaaagggtg 1800
ccagacgctt gcactgcctg cgatctggtc aatgtggatt tggatgactg catctttgaa 1860
caataa

```

```

SEQ ID NO: 2          moltype = DNA length = 1194
FEATURE               Location/Qualifiers

```

-continued

```

source          1..1194
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

SEQUENCE: 2
atggagctgg tccgggtggct cgtgggacaag gggattacct cggagaagca gtggatccag 60
gaggaccagg cctcatacat ctcttcaat gcggcctcca actcgcggtc ccaaatcaag 120
gctgccttgg acaatgctgg aaagattatg agcctgacta aaaccgcccc cgactacctg 180
gtgggccagc agcccggtga ggacatttcc agcaatcgga ttataaaat ttgggaacta 240
aacgggtacg atccccataa tgcggcttcc gtctttctgg gatgggccac gaaaaagtgc 300
ggcaagagga acacccatctg gctgtttggg cctgcaacta ccgggaagac caacatcgcg 360
gaggccatag cccacactgt gcccttctac ggggtcgtaa actggaccaa tgagaacttt 420
cccttcaacg actgtgtcga caagatgggt atctgggtggg aggaggggaa gatgaccgcc 480
aaggctcggt agtcggccaa agccatttct ggaggaagca aggtgctggt ggaccagaaa 540
tgcaagtcct cggcccagat agacccgact cccgtgatcg tcacctccaa caccaacatg 600
tgcgccgtga ttgacgggaa ctcaacgacc tcgaacacc agcagccgtt gcaagaccgg 660
atgttcaaat ttgaactcac ccgccgtctg gatcatgact ttgggaaggt caccaagcac 720
gaagtcaaaag actttttccg gtgggcaaaag gatcacgtgg ttgaggtgga gcatgaattc 780
tacgtcaaaa aggggtggagc caagaaaaga cccgccccca gtgacgcaga tataagtga 840
cccaaacggg tgcgcgagc agttgcgcag ccatcgagct cagacgcgga agcttcgac 900
aactacgcag acaggtacca aaacaaatgt tctcgtcacg tgggcatgaa tctgatgctg 960
ttccctgca gacaatgcga gagaatgaat cagaattcaa atatctgctt cactcacgga 1020
cagaaagact gtttagagtg ctttcccggt tcagaatctc aaccgcttgc tctcgtcaaa 1080
aaggcgatc agaaactgtg ctacattcat catatcatgg gaaaggtgcc agacgcttgc 1140
actgcctgcg atctggtcaa tgtggatttg gatgactgca tcttgaaca ataa 1194

```

```

SEQ ID NO: 3      moltype = DNA length = 2222
FEATURE          Location/Qualifiers
source           1..2222
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

```

SEQUENCE: 3
aataaacgat aacgcgcttg gtggcgtag gcatgtaaaa ggttacatca ttatcttgtt 60
cgccatccgg ttgtatataa tagacgttca tgttggtttt tgtttcagtt gcaagttggc 120
tgcggcgctg gcagcacctt tgcgggatcc tcaggtagta agtattcatt gtaaatctga 180
tattatttgt attattatag ctacctaat tgcagtgcaag atcattgaga taattaaaat 240
tataaccatc tcgcaataaa ataagtattt tactgttttc gtaacagttt tgtaataaaa 300
aaacctataa atcccatcac gccagataaa attcgttcat tttggatatt gtttcagtgc 360
cggggtttta cgaagtgttg attaagggtc ccagcgacct tgacgggcat ctgcccgga 420
ttcttgacag ctttgtgaac tgggtggcgg agaaggagtg ggagttgccg ccagattctg 480
acttggtatc gaatctgatt gacgaggcac cctgaccgtg ggccgagaag ctgcagcgcg 540
actttctgac ggagtggcgc cgtgtgagta agggcccgga ggcccttttc tttgtgcaat 600
ttgagaaggg agagagctac ttccacttac acgtgctcgt ggaaaccacc ggggtgaaat 660
ccttagtttt gggacgtttc ctgagtcaga ttccgcaaaa actgattcag agaatttacc 720
gcgggatcga cccgactttg ccaaaactgg tgcggtcac aaagaccaga aacggcgccg 780
gaggcgggaa caaggtggtg gacgagtgtc acatccccaa ttacttgctc cccaaaaacc 840
agcctgagct ccagtggggtg tggactaatt tagaacagta ttaagcgcc tgtttgaatc 900
tcacggagcg taacgcgttg gtggcgagc atctgacgca cgtgtcgag acgcaggagc 960
agaacaaaag gaatcagaat cccaattctg acgcgccggt gatcagatca aaaacttcag 1020
ccaggtacat ggagctggtc ggttggtctg tggacaaggg gattacctcg gagaagcagt 1080
ggatccagga ggaccaggcc tcatacatct ccttcaatgc ggcctccaac tcgcggtccc 1140
aaatcaaggc tgccttggac aatgcgggaa agattatgag cctgactaaa accgcccccg 1200
actacctggt gggccagcag cccgtggagg acatttccag caatcggatt tataaaattt 1260
tggaactaaa cgggtacgat cccaatatg cggcttccgt ctttctggga tgggcccaga 1320
aaaagtctcg caagaggaac accatctggc tgtttgggct tgcaactacc ggggaagacca 1380
acatcgcgga ggccatagcc cacactgtgc ccttctacgg gtgcgtaaac tggaccaatg 1440
agaactttcc cttcaacgac tgtgtcgaca agatgggtgat ctggtgggag gaggggaaga 1500
tgaccgccaa ggtcgtggag tcggccaaaag ccattctcgg aggaagcaag gtgcgctgg 1560
accagaaatg caagtctctg gccagatag acccgactcc cgtgatcgtc acctccaaca 1620
ccaacatgtg cgcctgtgatt gacgggaact caacgacctt cgaacaccag cagccgttgc 1680
aagaccggat gttcaaatgt gaactcacc gccgtctgga tcatgacttt gggaaaggtc 1740
ccaagcagga agtcaaaagc tttttccggt gggcaaaagg tcacgtggtt gaggtggagc 1800
atgaattcta cgtcaaaaag ggtggagcca agaaaagacc cgcgccaggt gacgcagata 1860
taagtgagcc caaacgggtg cgcgagtcag ttgcgcagcc atcgacgtca gacgcggaag 1920
cttcgatcaa ctacgcagac aggtacccaa acaaatgttc tcgtcactgt ggcatgaatc 1980
tgatgctgtt ccctgcgaga caatgcgaga gaatgaatca gaattcaaat atctgcttca 2040
ctcagcgaca gaaagactgt tttagagtgt ttcccggtgc agaattcaca cccgtttctg 2100
tcgtcaaaaa ggcgtatcag aaactgtgct acattcatca tatcatggga aaggtgccag 2160
acgcttgcac tgctgcgat ctggtcaatg tggatttggg tgactgcatc tttgaacaat 2220
aa 2222

```

```

SEQ ID NO: 4      moltype = DNA length = 2583
FEATURE          Location/Qualifiers
source           1..2583
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

SEQUENCE: 4

-continued

```

gacctttaat tcaacccaac acaatatatt atagttaaat aagaattatt atcaaatcat 60
ttgtatatta attaaaaatac tatactgtaa attacatttt atttacaatc actcgacgaa 120
gacttgatca cccggggggac aaggtaagta ttcattgttaa atctgatatt atttgtatta 180
ttatacttac ctaatttgca gtgcaggccg ccatggtaag tatcgataac tttgttttct 240
ttcacattta caactccgac atacaaattg taatttttatt actgtttggg ccataaacac 300
ttgtttacca tttctttttt tacaagtttt aatattttct gcatatataa aacattattt 360
atthtgacgg ctgcccagcg ttatctaccc gattggctcg aggacaacct ctctgagggc 420
attcgcgagt ggtggggcgt gaaacctgga gcccgaagc ccaagccaa ccagcaaaag 480
caggacgacg gccggggctc ggtgcttcct ggtacaaagt acctcgagc cttcaacgga 540
ctcgacaagg gggagcccggt caacgcggcg gacgcagcgg ccctcgagca cgacaaggcc 600
tacgaccagc agctgcaggc ggggtgacaat ccgtacctgc ggtataacca cgccgacgcc 660
gagtttcagg agcgtctgca agaagatacg tcttttgagg gcaacctcgg gcgagcagtc 720
ttccaggcca agaagcgggt tctcgaacct ctcggtctgg ttgaggaagg cgctaagacg 780
gtcctcgtaa agaagagcca ggttagagcca tcaccccagc gttctccaga ctctctacg 840
ggcatcgcca agaagcgcca acagcccgcc agaaaaagac tcaattttgg tcagactggc 900
gactcagagt cagttccaga cctcaacct ctcgagaaac ctccagcagc gccctctggt 960
gtgggacctt atacaattgc tgcaggcggg tgccgaccaa tggcagacaa taacgaaggc 1020
gccgacggag tgggtagttc ctcggaatat tggcattgag attccacatg gctggggcag 1080
agagtcacac ccaccagcac ccgaacctgg gccctgcccc cctacaacaa ccacctctac 1140
aagcaaatct ccaacgggac atcgggagga gccaccaaag acaaacacta cttcggtctac 1200
agcaccctct gggggatttt tgactttaac agattccact gccacttttc accacgtgac 1260
tggcagcgac tcataaccaa caactgggga ttccggccca agagactcag cttcaagctc 1320
ttcaacatcc aggtcaagga ggtcacgagc aatgaaggca ccaagaccat cgccaataac 1380
ctcaccagca ccattccagg gtttacggac tcggagtagc agctgcccga cgttctcggc 1440
tctgcccacc agggctgctt gctcctgctt ccggcgagcg tgttcatgat tcccagtag 1500
ggctacctaa cactcaacaa cggtagtcag gccgtgggac gctcctctct ctactgctg 1560
gaatactttc cttcgagat gctgagaacc ggcaacaact tccagtttac ttacaccttc 1620
gaggacgtgc ctttccacag cagctacgcc cacagccaga gcttggaccg gctgatgaat 1680
cctctgattg accagtaact gtactacttg tctcgagtc aaacaacagg aggcacggca 1740
aatacgacga ctctgggctt cagccaaggt gggcctaata caatggccaa tcaggcaaaag 1800
aactggctgc caggacctg ttaccgccaa caacgcgtct caacgacaac cgggcaaaac 1860
aacaatagca cagtttgcct gactgctggg accaaatacc atctgaatgc aagaaattca 1920
ttggctaact ttggcatctc tatggcaaca cacaagacag acgaggagcg ttttttccc 1980
agtaacggga cctgtatttt tggcaaacaa aatgctgcca gagacaatgc ggattacagc 2040
gatgtcatgc tcaccagcga ggaagaaatc aaaaccacta acctgtggc tacagaggaa 2100
tacggtatcg tggcagataa cttgcagcag caaaacacgg ctctcctaat tggaaactgtc 2160
aaacgcccag gggccttacc cgttatggc tggcagaacc gggacgtgta cctgcagggt 2220
cccatctggg ccaagatttc tcacacggac ggcaacttcc acccgtctcc gctgatgggc 2280
ggctttggcc tgaacatctc tccgcctcag atcctgatca agaacacgcc tgtacctgcy 2340
gatctccga ccactctcaa ccagtcaaa ctgaactctt tcatcacgca atacagcacc 2400
ggacagggtc cgtgtgaaat tgaatgggag ctgcagaagg aaaacagcaa gcgctggaac 2460
cccagatcc agtacacctc caactactac aaatctacaa gtgtggactt tgcgtgtaat 2520
acagaaggcg tgtactctga accccgcccc attggcaccg gttacctcac ccgtaatctg 2580
taa 2583

```

```

SEQ ID NO: 5      moltype = DNA length = 2407
FEATURE          Location/Qualifiers
source           1..2407
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

```

SEQUENCE: 5
cctgcaggca gctgcgcgt cgtcgcgtca ctgaggccgc ccgggcaaaag cccgggctgc 60
ggggacattt tggtcgccc gctcagtgta gcgagcgagc gcgcagagag ggagtgccca 120
actccatcac taggggttcc tgcggccgca cgcgcgcccc gtcagtgggc agagcgccaa 180
tcgcccacag tccccagaaa gttgggggga ggggtcgcca attgaacggg tgcctagaga 240
aggtggcgcg gggtaaaact ggaagtgtat gtcgtgactt ggtccgacct ttttcccgag 300
ggtgggggag aaccgtatat aagtgcagta gtcccggtga acgttctttt tcgcaacggg 360
tttgccgcca gaacacgcgt aagggatccg ccacctggtt gagcaagggc gaggaggata 420
acatggccat catcaaggag ttcatgctc tcaagggtga catggagggc tccgtgaacg 480
gccacgagtt cgagatcgag ggcgaggcgc agggccgccc ctacgagggc accgagaccg 540
ccaaagctgaa ggtgaccaag ggtggccccc tgcccttcgc ctgggacatc ctgtccctc 600
agttcatgta cggctccaag gcttacgtga agcaccgccg cgacatcccc gactacttga 660
agctgtcctt ccccgagggc ttcaagtggg agcgcgtgat gaacttcgag gacggcgggc 720
tgggtgacct gacccaggac tctcctctgc aggcggcgca gttcatctac aagggtgaagc 780
tgccgggac caacttcccc tccgacggcc ccgtaatgca gaagaagacc atgggctggg 840
aggctcctc cgagcggagt tcccccgagg acggcgccct gaagggcgag atcaagcaga 900
ggctgaagct gaaggacggc ggcactacg acgtgagggt caagaccacc tacaaggcca 960
agaagcccggt gcagctgccc ggcgcctaca acgtcaacat caagttggac atcacctccc 1020
acaacgagga ctacaccatc gtggaacagt acgaacgcgc cgagggcgcc cactccaccg 1080
gcggcatgga cagactgtac aagtaagaat tcgatatcaa gcttatcgat aatcaacctc 1140
tggattacaa aatttgtgaa agattgactg gtattcttaa ctatgttctt ccttttacgc 1200
tatgtggata cgtgctttta atgcctttgt atcatgctat tgcctcccgat atggctttca 1260
ttttctctc cttgtataaa tctgtgtgac tgtctcttta tgaggagttg tggcccggtg 1320
tcaggcaaac tggcggtggg tgcactgtgt ttgctgacgc aacccccact ggttggggca 1380
ttgccaccac ctgtcagctc ctttccggga ctttccgctt cccctccctc attgccacgg 1440
cggaactcat ccgcgcctgc cttgcccgtc gctggacagg ggctcggtcg ttgggcaactg 1500
acaattccgt ggtgtgtgtc gggaaatcat cgtcctttcc ttggtgctc gctgtgtgtg 1560
ccactcggtat tctgcgctgg acgtccttct gctacgtccc ttcggccctc aatccagcgg 1620

```

-continued

```

accttctcttc cgcgggctcg ctgcgggctc tgcgggctct tccggctctt cgcctctgcc 1680
ctcagacgag tcggatctcc ctttgggccc cctccccgca tcgataccga gcgctgctcg 1740
agagatctac ggggtggcatc cctgtgaccc ctccccagtg cctctctctg ccttggaagt 1800
tgccactcca gtgcccacca gccctgtcct aataaaatta agttgcacat tttgtctga 1860
ctaggtgtcc ttctataata ttatgggggtg gagggggggtg gtatggagca aggggcaagt 1920
tggaagaca accgttaggg cctgcggggt ctattgggaa ccaagctgga gtgcagtggc 1980
acaactcttg ctcaactgcaa tctccgctc ctgggttcaa gcgattctcc tgcctcagcc 2040
tcccagattg ttgggattcc aggcattgcat gaccaggctc agctaatttt tgtttttttg 2100
gtagagacgg ggtttcacca tattggccag gctgtctctc aactccta atctcaggtgat 2160
ctacccacct tggcctccca aattgctggg attacaggcg tgaaccactg ctccctccc 2220
tgtcctcttg atttttagg taaccacgtg cggacgcagc ggccgcagga acccctagt 2280
atggagtttg ccactccctc tctgcgcgct cgctcgctca ctgagggcgg gcgaccaaag 2340
gtcggccgac gcccgggctt tgcccgggcg gcctcagtga gcgagcgagc gcgcagctgc 2400
ctgcagg 2407

```

```

SEQ ID NO: 6      moltype = DNA length = 2406
FEATURE          Location/Qualifiers
source           1..2406
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

```

SEQUENCE: 6
cacatgttgg acatcgtgtc gtttgagcgt ataaaagaat atataagagc taatttaggc 60
catttcacag taatcaccca caaatgttcg aagcgtaagg tgtgtcttca tcacaaacga 120
attgccaggt tgttgggcat taaaaaata tatcatcaag aatacaaacg ggttgtttca 180
aaggtttaca agaagcaaac attgtaaacg tgcgggagca acaatctccg gagactgcgg 240
cgtgtgtcaa aaattaaaag ctgttgtaata aactggaatc gagctcttac aacaaatcca 300
acattgacca gctggcgtt attgtaaaag atcctcaggt agtaagtatt cattgtaaat 360
ctgatattat ttgtattatt atacctacct aatttgcagt gcagatcatt gagataatta 420
aaattataac catctcgcaa ataaataagt attttactgt tttcgtaaca gttttgtaat 480
aaaaaaacat ataaatccca ttacggcagt ataaattcgt tcattttgga tattgtttca 540
gtgcgggggt tttacgagat tgtgattaag gtccccagcg accttgacgg gcactctgcc 600
ggcattttctg acagctttgt gaactgggtg gccgagaagg agtgggagtt gccgccagat 660
tctgacttgg atctgaatct gattgagcag gcaccctga cgtggccga gaagctgcag 720
cgcgacttcc tgacggagtg gcgcggtgtg agtaaggccc cggaggccct tttctttgtg 780
caatttgaga agggagagag ctacttccac ttacacgtgc tcgtggaaac caccggggtg 840
aaatccttag ttttgggacg ttctctgagt cagattccgc aaaaactgat tcagagaatt 900
taccgcgga tcgagccgac tttgccaaac tgggtccggy tcacaaagac cagaaacggc 960
gccggaggcg ggaacaaggt ggtggacgag tgcacatcc ccaattactt gctcccaaaa 1020
accagcctg agctccagtg ggcgtggact aatttagaac agtatttaag gcctgtttg 1080
aatctcaccg agcgtaaacg gttggtggcg cagcatctga cgcacgtgtc gcagacgcag 1140
gagcagaaca aagagaatca gaatcccaat tctgacgcgc cgtgtgatcag atcaaaaact 1200
tcagccaggt acatggagct ggtcgggtgg ctctgggaca aggggattac ctcgagaag 1260
cagtggatcc agggaggcca ggctcatac atctccttca atgcggcctc caactcggcg 1320
tcccaaatca aggtgcctt ggacaatgcg ggaagatta tgagcctgac taaaaccgcc 1380
ccgactacc tgggtggcca gcagcccggtg gaggaacttt ccagcaatcg gatttataaa 1440
attttggaac taaacgggta cgaatcccaa tatgcggctt ccgtctttct gggatgggccc 1500
acgaaaaagt tcggcaagag gaacaccatc tggctgtttg ggctcgcaac taccgggaag 1560
acacaacatg cggaggccat agcccacact gtgcccttct acgggtgcgt aaactggacc 1620
aatgagaact ttcccttcaa cgactgtgtc gacaagatgg tgatctgggt ggaggagggg 1680
aagatgaccg ccaaggtcgt ggaagtcggc aaagccatc tcggaggaag caaggtgcgc 1740
gtggaccaga aatgcaagtc ctcgcccagc atagaccga ctcccgatg cgtcacctcc 1800
aacaccaaca tgtgcgcctg gattgacggg aactcaacga ccttcgaaca ccagcagccg 1860
tgtcaagacc ggaatgtcaa atttgaactc acccgccgtc tggatcatga ctttgggaag 1920
gtcaccagcg aggaagtcaa agacttttct cgggtgggcaa aggatcacgt ggttgaggtg 1980
gagcatgaat tctacgtcaa aaaggggtga gccaaagaaa gaccgcccc cagtgcagca 2040
gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc agccatcgac gtcagacgcg 2100
gaagcttcga tcaactacgc agacaggtac caaaacaaat gttctcgtca cgtgggcagt 2160
aatctgatgc tgtttccctg cagacaatgc gagagaatga atcagaatc aaatatctgc 2220
ttcactcacg gacagaaaga ctgttttagg tgccttcccg tgcagaatc tcaaccctgt 2280
tctgtcgtca aaaaggcgta tcagaaactg tgcacattc atcatatcat gggaaaggtg 2340
ccagacgctt gcaactgcctg cgatctggtc aatgtggatt tggatgactg catctttgaa 2400
caataa 2406

```

```

SEQ ID NO: 7      moltype = DNA length = 2137
FEATURE          Location/Qualifiers
source           1..2137
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

```

SEQUENCE: 7
gggagttcag tcgtcgaatg caaacgtaa aaaatattaa taaggtaaaa actacagctg 60
gatcctcagg tagtaagtat tcattgtaaa tctgatatta tttgtattat tataacctacc 120
taatttcag tttcgtatc ttagataatt aaaattataa ccatctcgca aataaataag 180
tattttactg tttcgttaac agttttgtta taaaaaaacc tataaatccc attacggcag 240
tataaattcg ttcatttttg atattgttcc agtgcggggg ttttacgaga ttgtgattaa 300
ggtccccagc gaccttgacg ggcactctgc cggcatttct gacagctttg tgaactgggt 360
ggccgagaag gagtgggagt tgcccgcaga ttctgacttg gatctgaatc tgattgagca 420
ggcaccctcg accgtggcgg agaagctgca gcgcgacttt ctgacggagt ggccgcgtgt 480

```

-continued

```

gagtaaggcc cccgaggccc tttctttgt gcaatttgag aaggagagaga gctacttcca 540
cttacacgtg ctcgtgga aaacccgggt gaaatoccta gttttgggac gtttccctgag 600
tcagattcgc gaaaaactga ttccagagaat ttaccgcggg atccgagccga ctttgccaaa 660
ctggttcgcg gtcacaaaga ccagaaacgg ccgcccggagg gggaaacaagg tgggtggacga 720
gtgctacatc cccaattact tgctccccaa aaccacagcct gagctccagt gggcgtggac 780
taatttagaa cagtatttaa gcgcctgttt gaatctcacg gagcgtaaac ggttggtggc 840
gcagcatctg acgcacgtgt cgcagacgca ggagcagaac aaagagaatc agaatcccaa 900
ttctgacgcg ccggtgatca gatcaaaaac ttcagccagg tacatggagc tggtcgggtg 960
gctcgtggac aaggggatta cctcggagaa gcagtggatc caggaggacc aggcctcata 1020
catctccttc aatcgggcct ccaactcgcg gtcccaaadc aaggctgcct tggacaatgc 1080
gggaaagatt atgagcctga ctaaaaacgc ccccgactac ctgggtgggc agcagcccgt 1140
ggaggacatt tccagcaatc ggatttataa aattttggaa ctaaacgggt acgatcccca 1200
atatcgccgt cctcgtcttc tgggatgggc cagcaaaaaa ttcggcaaga ggaacacct 1260
ctggctgttt ggcctctgca ctaccgggaa gaccaacatc gcggaggcca tagccacac 1320
tgtgcccttc tacgggtgcg taaactggac caatgagaac tttcccttca acgactgtgt 1380
cgacaagatg gtgatctggt gggaggaggg gaagatgacc gccaaaggtcg tggagtcggc 1440
caaaagccatt ctcggaggaa gcaaggtgcy cgtggaccag aaatgcaagt cctcggccca 1500
gatagaccgc actcccgatg tctcaccctc caacaccaac atgtgcgcgc tgattgacgc 1560
gaactcaacg accttcgaac accagcagcc gttgcaagac cggatgttca aatttgaact 1620
caccgcgcgt ctggatcatg actttgggaa ggtcaccag caggaagtca aagactttt 1680
ccggtgggca aaggatcacg ttggttgaggt ggagcatgaa ttctacgtca aaaaggggtg 1740
agccaagaaa agaccgcgcc ccagtgcgc agatataagt gagcccaaac ggggtgcgcga 1800
gtcagttgcy cagccatcga cgtcagacgc ggaagcttcg atcaactacg cagacaggta 1860
ccaaaacaaa tgtctcgtc acgtgggcat gaactctgat ctgtttccct gcagacaatg 1920
cgagagaatg aatcagaatc caaatatctg cttactcacg ggacagaaag actgtttaga 1980
gtgctttccc gtgtcagaat ctcaaccctg tctgtcgtc aaaaaggcgt atcagaaact 2040
gtgctacatt catcatatca tgggaaaggt gccagacgct tgcactgcct gcgatctggt 2100
caatgtgatg ttgatgact gcatctttga acaataa 2137

```

```

SEQ ID NO: 8      moltype = DNA length = 2188
FEATURE          Location/Qualifiers
source           1..2188
                 mol_type = other DNA
                 note = artificial sequence
                 organism = unidentified

```

```

SEQUENCE: 8
gacctttaat tcaacccaac acaatatatt atagttaaat aagaattatt atcaaatcat 60
ttgtatatta attaaaatac tatactgtaa attacatatt atttacaatc ggatcctcag 120
gtagtaagta ttcatgttaa atctgatatt atttgtatta ttatacctac ctaatttgca 180
gtgcagatca tttagatata taaaattata accatctcgc aaataaataa gtattttact 240
gttttcgtaa cagttttgta ataaaaaac ctataaatcc cattacggca gtataaatc 300
gttcattttg gatattgttt cagtgcgggg gttttacgag attgtgatta aggtcccccag 360
cgacttgac gggcatctgc ccggcatctt tgacagcttt gtgaactggg tggccgagaa 420
ggagtgggag ttgcgcgcag attctgactt ggatctgaat ctgattgagc aggcaccccct 480
gacgctggcc gagaagctgc agcgcgactt tctgacggag tggcgccgctg tgagtgaagg 540
cccgagggcc cttttctttg tgcaatttga gaaggagag agctacttcc acttacacgt 600
gctcgtggaa accaccgggg tgaatcctt agttttggga cgtttcctga gtcagattcg 660
cgaaaaactg attcagagaa ttaccgcgg gatcagaccg actttgcca actggttcgc 720
gggtcaaaaag accagaaaag gcgcggagg cggaacaacg gtgggtggag agtgctacat 780
ccccaattac ttgtcccca aaaccagcc tgagctccag tggcgctgga ctaatttaga 840
acagtattta agcgcctggt tgaatctcac ggagcgtaaa cggttgtggt gcgagcatct 900
gacgcacgtg tccgacagcg aggagcagaa caaagagaat cagaatccca attctgacgc 960
gccggtgatc agatcaaaaa cttcagccag gtacatggag ctgggtcgggt ggctcgtgga 1020
caaggggatt acctcggaga agcagtggat ccaggaggac caggcctcat acatctcctt 1080
caatgcggcc tccaactcgc ggtcccaaat caaggctgcc ttggacaatg cgggaaagat 1140
tatgagcctg actaaaaccg cccccgacta cctgggtggg cagcagcccg tggaggacat 1200
ttccagcaat cggatttata aaatttttga actaaacggg tacgatcccc aatatgcggc 1260
ttccgtcttt ctgggatggg ccacgaaaaa gttcggcaag aggaacacca tctggctggt 1320
tgggcctgca actaccggga agaccaacat ccgaggaggc atagcccaca ctgtgccctt 1380
ctacgggtgc gtaaaactgga ccaatgagaa ctttcccttc aacgactgtg tcgacaagat 1440
gggtgatctg tgggaggagg ggaagatgac cgccaaggtc gtggagtgcg ccaaagccat 1500
tctcggaggga agcaagggtc gcgtggacca gaaatgcaag tctcggccc agatagacc 1560
gactcccggt atcgtcacct ccaacaccaa catgtgcgcc gtgattgacg ggaactcaac 1620
gaccttcgaa caccagcagc cgttgcaaga ccggatgttc aaatttgaa tcaccgcgcg 1680
tctggatcat gactttggga aggtcaccaa gcaggaagtc aaagactttt tccggtgggc 1740
aaaggatcac gtggttgagg tggagcatga attctacgtc aaaaagggtg gagccaagaa 1800
aagaccgcc cccagtgcag cagatataag tgagcccaa cgggtgcgcg agtcagtgc 1860
gcagccatc agtcaagcgc cggaaagcttc gatcaactac gcagacaggt accaaaacaa 1920
atgttctcgt cacgtgggca tgaatctgat gctgtttccc tgcagacaat gcgagagaat 1980
gaatcagaat tcaaatatct gcttactca cggacagaaa gactgtttag agtgctttcc 2040
cgtgtcagaa tctcaaccct tttctgctg caaaaaggcg tatcagaaac tgtgctacat 2100
tcatcatatc atgggaaagg tgccagacgc ttgactgccc tgcgatctgg tcaatgtgga 2160
tttgatgac tgcacttttg aacaataa 2188

```

```

SEQ ID NO: 9      moltype = DNA length = 2379
FEATURE          Location/Qualifiers
source           1..2379
                 mol_type = other DNA
                 note = artificial sequence

```

-continued

```

organism = unidentified
SEQUENCE: 9
atcattgaga taattaaat tataaccatc tcgcaataa ataagtattt tactgttttc 60
gtaacagttt tgtaataaaa aaacctataa atattccgga ttattcatac cgtcccacca 120
tcgggcgcgg atcccagcag gtaagtatcg ataactttgt tttctttcac atttacaact 180
ccgacataca aatatgaata aacgataacg ccgttggttg cgtgaggcat gtaaaagggt 240
acatcattat cttgttcgcc atccggttg tataaataga cgttcattgt ggtttttgt 300
tcagttgcaa gttggctgcg gcgcgcgcag cacctttgcg taagccgcca gtaagtattc 360
attgtaaatc tgatattatt tgtattatta tacctaccta atttgacgtg cagggttaatt 420
ttattactgt ttggtccata aacacttgtt taccatttcc ttttttaca gttttaatat 480
tttctgcata tataaaaacat tattttttt gcagtgcggg gggttttacg gattgtgatt 540
aaggteccca gcgaccttga cgggcacatc ccgcgcattt ctgacagctt tgtgaactgg 600
gtggccgaga aggagtgagg gttgcgcgca gattctgact tggatctgaa tctgattgag 660
caggcaccac tgaccgtggc cgagaagctg cagcgcgact tctgacgga gtggcgccgt 720
gtgagtaagg ccccgaggc ccttttcttt gtgcaatttg agaagggaga gactacttcc 780
cacttacacg tgctcgtgga aaccaccggg gtgaaatcct tagttttggg acgtttctcg 840
agtcagattc gcgaaaaaac gattcagaga atttaccgag ggatcgagcc gactttgcca 900
aactggttcg cggtcacaaa gaccagaaac ggccgcggag gcgggaacaa ggtggtggac 960
gagtgctaca tccccaatc cttgctcccc aaaaccagc ctgagctcca gtggcggtg 1020
actaatttag aacagatttt aagcgctgtt ttgaatctca cggagcgtaa acggttggtg 1080
gcgcagcatc tgacgcacgt gtgcgcagac caggagcaga acaagagaa tccagaatccc 1140
aattctgacg cgcggtgat cagatcaaaa acttcagcca ggtacatgga gctggtcggg 1200
tggtctcgtg acaagggtat tacctcggag aagcagtgga tccaggagga ccaggcctca 1260
tacatctcct tcaatcgggc ctccaaactc cgggtcccaa tcaaggctgc cttggacaat 1320
gcgggaaaga ttatgagctc gactaaaaac gccccgact acctggtggg ccagcagccc 1380
gtggaggaca tttccagcaa tccggtttat aaaaattttg aactaaacgg gtacgactcc 1440
caatatcgcg cttccgtctt tctgggatgg gccacgaaaa agttcggcaa gaggaacacc 1500
atctggctgt ttgggcctgc aactaccggg aagaccaaca tcgcggaggc catagccacc 1560
actgtgcctt tctacgggtg cgtaaactgg accaatgaga actttccctt caacgactgt 1620
gtcgacaaga tgggtgatct gtgggaggag gggaagatga ccgccaaggt cgtggagctg 1680
gccaaagcca tctcggagg aagcaagggt cgcgtggacc agaaatgcaa gtctcggcc 1740
cagatagacc cgaactccgt gatcgtcacc tccaacacca acatgtgcgc cgtgattgac 1800
gggaaactca cgactctcga accaccagc cgtgtgcaag accggtggtt caaatttgaa 1860
ctcaccgcc cttgggatca tgactttggg aaggtcacca agcaggaaat caaagacttt 1920
ttccggtggg caaaggatca cgtggttgag gtggagcatg aattctactt caaaaagggt 1980
ggagccaaga aaagaccgcg ccccgatgac gcagatataa gtgagcccaa acggtgctgc 2040
gagtcagttg acgagccatc gacgtcagac gcggaagctt cgatcaacta cgcagacagg 2100
taccaaaaca aatgtttctg tcacgtgggc atgaatctga tgcgtgttcc ctgcagacaa 2160
tgccagagaa tgaatcagaa ttcaaatatc tgcctcactc acggacagaa agactgttta 2220
gagtgctttc ccgtgtcaga atctcaacc ctttctgtcg tcaaaaaggc gtatcagaaa 2280
ctgtgctaca ttcacatcat catgggaaa gtgccagacg cttgcaactgc ctgcgactgc 2340
gtcaatgtgg atttgatga ctgcactctt gaacaataa 2379

```

```

SEQ ID NO: 10      moltype = DNA length = 2560
FEATURE            Location/Qualifiers
source              1..2560
                    mol_type = other DNA
                    note = artificial sequence
                    organism = unidentified

```

```

SEQUENCE: 10
atcattgaga taattaaat tataaccatc tcgcaataa ataagtattt tactgttttc 60
gtaacagttt tgtaataaaa aaacctataa atattccgga ttattcatac cgtcccacca 120
tcgggcgcgg atcccagcag gtaagtatcg ataactttgt tttctttcac atttacaact 180
ccgacataca aatcacatgt tggacatcgt gtctgttgag cgtataaaag aatatataag 240
agctaattta ggcattttca cagtaatcac cgacaaatgt tcgaagcgta aggtgtgtct 300
tcacacaaaa cgaattgccg ggtgtttggg cattaaaaaa atatatcact aagaatacaa 360
acgggttgtt tcaaaagggtt acaagaagca aacattgtaa acgtgcccga gcaacaatct 420
ccggagactg cggccgtgtg caaaaattaa aagctgttga ataaactgga atcgagctct 480
tacaacaaat ccaacattga ccagctggcc gttattgtta atagccgcc agtaagtatt 540
cattgtaaat ctgatattat ttgtattatt atacctacct aatttgacgt gcagggtaat 600
tttattactg tttggtccat aaacacttgt ttaccatttc cttttttaca agttttaata 660
ttttctgcat atataaaaca ttattttatt tgcagtgcgc gggttttacg agatttgtat 720
taaggtcccc agcgaccttg acgggcatct gcccggcatt tctgacagct ttgtgaactg 780
gggtggccag aaggagtggt agttgcgcgc agattctgac ttggtatctg atctgattga 840
gcaggcacc ctagccgttg ccgagaagct gcagcgcgac tttctgacgg agtggcgccg 900
tgtagtagaa gccccggagg cctttttctt tgtgcaattt gagaagggag agagctactt 960
ccacttacac gtgctcgttg aaaccaccgg ggtgaaatcc ttagtttttg gacgtttcct 1020
gagtcagatt gcgaaaaaac tgattcagag aatttaccgc gggatcgagc cgactttgcc 1080
aaactggttc gcggtcacia agaccagaaa cggcgccgga gcggggaaca aggtggttga 1140
cgagtgttac atccccaatt acttgtctcc caaaaaccag cctgagctcc agtgggctgt 1200
gactaattta gaacagatt taagccctgt tttgaatctc acggagcgta aacggttggt 1260
ggcgcagcat ctgacgcagc tgtcgcagac gcaggagcag aacaaagaga atcagaatcc 1320
caattctgac gcgcgggtga tcagatcaaaa aacttcagcc aggtacatgg agctggtcgg 1380
gtggctcgtg gacaagggga ttacctcgga gaagcagtg atccaggagg accaggctc 1440
atacatctcc ttcaatcggt cctccaaact gcgggtccaa atcaaggctg ccttggaaca 1500
tgccggaaag attatgagc tgactaaaac cggcccgac tacctggtg gccagcagcc 1560
cgtggaggac atttccagca atcggattta taaaattttg gaactaaacg ggtacgatcc 1620
ccaatatcgc gcttcgctct ttctgggatg ggccacgaaa aagttcggca agaggaaac 1680
catctggctg tttgggcctg caactaccgg gaagaccaac atcgcggagg ccatagccca 1740

```

-continued

cactgtgccc	ttctacgggt	gcgtaaactg	gaccaatgag	aactttccct	tcaacgactg	1800
tgtcgacaag	atggtgatct	ggtgggagga	ggggaagatg	accgccaagg	tcgtggagtc	1860
ggccaaagcc	attctcggag	gaagcaaggt	gcgcgtggac	cagaaatgca	agtccctcgc	1920
ccagatagac	ccgactcccc	tgatcgtcac	ctccaacacc	aacatgtgcg	ccgtgattga	1980
cgggaactca	acgaacctcg	aacaccagca	gccgttgcaa	gaccggatgt	tcaaatTTga	2040
actcaccgcc	cgtctggatc	atgactttgg	gaaggtcacc	aagcaggaag	tcaaaagact	2100
tttccggtgg	gcaaaggatc	acgtggttga	ggtggagcat	gaattctacg	tcaaaaaggg	2160
tggagccaag	aaaagacccc	ccccagtgga	cgcagatata	agtgagccca	aacgggtgcy	2220
cgagtcagtt	gcgcagccat	cgacgtcaga	cgcggaagct	tcgatcaact	acgcagacag	2280
gtaccaaacc	aaatgttctc	gtcacgtggg	catgaatctg	atgctgtttc	cctgcagaca	2340
atgcgagaga	atgaatcaga	attcaaatat	ctgcttctac	cacggacaga	aagactgttt	2400
agagtgtctt	cccgtgtcag	aatctcaacc	cgtttctgtc	gtcaaaaagg	cgtatcagaa	2460
actgtgtctac	attcatcata	tcatgggaaa	ggtgccagac	gcttgcaactg	cctgcgatct	2520
ggccaatgtg	gatttgatg	actgcatctt	tgaacaataa			2560

SEQ ID NO: 11 moltype = DNA length = 2294
 FEATURE Location/Qualifiers
 source 1..2294
 mol_type = other DNA
 note = artificial sequence
 organism = unidentified

SEQUENCE: 11

atcattgaga	taattaaaat	tataaccatc	tcgcaaatata	ataagtattt	tactgttttc	60
gtaacagttt	tgtaataaaa	aaacctataa	atattccgga	ttattcatac	cgtcccacca	120
tcgggcccgc	atcccagcag	gtaagtatcg	ataactttgt	tttctttcac	atttacaact	180
ccgacataca	aatatgggga	gttcagtcgt	cgaatgcaaa	acgtaaaaaa	tattataaag	240
gtaaaaacta	cagcttaagc	cgcagctaag	tattcattgt	aaatctgata	ttatttTgat	300
tattatacct	acctaaattt	cagtgcaggg	taatttttatt	actgtttggt	ccataaacac	360
ttgtttacca	tttctttttt	tacaagtttt	aatattttct	gcataataaa	aacattattt	420
atTTTgcagt	gcccgggttt	tacgagattg	tgattaaggt	ccccagcgac	cttgacgggc	480
atctgcccgc	catttctgac	agctttgtga	actgggtggc	cgagaaggag	tgggagttgc	540
cgccagattc	tgactttggat	ctgaatctga	ttgagcaggc	acccctgacc	gtggccgaga	600
agctgcagcg	cgactttctg	acggagtggt	gccgtgtgag	taaggccccc	gaggcccttt	660
tctttgtgca	atttgacaag	ggagagagct	acttccactt	acacgtgctc	gtggaaacca	720
cgggggtgaa	atccttagtt	ttgggacggt	tctctagtc	gattcgcgaa	aaactgattc	780
agagaattta	ccgcgggatc	gagccgactt	tgccaaaactg	gttcgcggtc	acaaagacca	840
gaaacggcgc	cggaggcggg	aacaaggtgg	tggacgagtg	ctacatcccc	aattactttg	900
tcccacaaa	ccagocctgag	ctccagtggt	cgtggactaa	tttgaacag	tatttaagcg	960
cctgtttgaa	tctcacggag	cgtaaaacgt	tgttgggcga	gcattctgacg	cacgtgtcgc	1020
agacgcagga	gcagaacaaa	gagaatcaga	atcccaattc	tgacgcgccg	gtgatcagat	1080
caaaaacttc	agccaggtac	atggagctgg	tcgggtggct	cgtggacaag	gggattacct	1140
cgggagaagca	gtggatccag	gaggaccagg	cctcatacat	ctccttcaat	gcggccctcca	1200
actcgcggtc	ccaaatcaag	gctgccttgg	acaatgcggg	aaagattatg	agcctgacta	1260
aaaccgcccc	cgactacctg	gtggggccagc	agcccggtgga	ggacatttcc	agcaatcgga	1320
tttataaaat	tttggaacta	aacgggtacg	atcccacata	tgccggttcc	gtctttctgg	1380
gatggggcac	gaaaaaggtc	ggcaagagga	acaccatctg	gctgtttggg	cctgcaacta	1440
ccgggaagac	caacatcgcg	gagggccatag	cccacactgt	gcccttctac	gggtgcgtaa	1500
actcgaccaa	tgagaacttt	cccttcaacg	actgtgtcga	caagatgggt	atctggtggg	1560
aggaggggaa	gatgacgcgc	aaggtcgtgg	agtcggccaa	agccattctc	ggagggaagca	1620
agggtgcgct	ggaccagaaa	tgcaggtcct	cgggccagat	agaccgcact	cccgatgatc	1680
tcacctccaa	caccaacatg	tgcgcgtgga	ttgacgggaa	ctcaacgacc	ttcgaacacc	1740
agcagccgtt	gcaagaccgg	atgttcaaat	ttgaactcac	cgcgcgtctg	gatcatgact	1800
ttgggaaggt	caccaagcag	gaagtcaaa	actttttccg	gtgggcaaa	gatcacgtgg	1860
ttgaggtgga	gcattgaattc	tacgtcaaaa	agggtggagc	caagaaaaga	cccgcctcca	1920
gtgacgcaga	tataagttag	cccaaacggg	tgcgcgagtc	agttgcgcag	ccatcgacgt	1980
cagacgcgga	agcttcgatc	aactacgcga	acaggtacca	aaacaaatgt	tctcgtcacg	2040
tgggcatgaa	tctgatgctg	tttccctgca	gacaatgcga	gagaatgaat	cagaatttcaa	2100
atatctgctt	cactcacgga	cagaaagact	gttttagagt	ctttcccggt	tcagaatctc	2160
aaaccggttc	tgtcgtcaaa	aaggcgtatc	agaaactgtg	ctacattcat	catatcatgg	2220
gaaagggtgc	agacgcttgc	actgcctgcg	atctgggtcaa	tgtggatttg	gatgactgca	2280
tctttgaaca	ataa					2294

SEQ ID NO: 12 moltype = DNA length = 2345
 FEATURE Location/Qualifiers
 source 1..2345
 mol_type = other DNA
 note = artificial sequence
 organism = unidentified

SEQUENCE: 12

atcattgaga	taattaaaat	tataaccatc	tcgcaaatata	ataagtattt	tactgttttc	60
gtaacagttt	tgtaataaaa	aaacctataa	atattccgga	ttattcatac	cgtcccacca	120
tcgggcccgc	atcccagcag	gtaagtatcg	ataactttgt	tttctttcac	atttacaact	180
ccgacataca	aatatggacc	tttaattcaa	cccaacacaa	tatattatag	ttaaataaga	240
attattatata	cactcattgt	atattaatta	aaatactata	ctgtaaaatta	cattttattt	300
acaatctaa	ccgccagtaa	gtattcattg	taaactctgat	attatttTga	ttattatacc	360
tacctaaatt	gcagtgacgg	gtaattttat	tactgttttg	tccataaaca	cttggtttacc	420
atTTTctttt	ttacaagttt	taatatTTTt	tgcatatata	aaacattatt	tattttgcag	480
tgccggggtt	ttacgagatt	gtgattaagg	tcccagcgca	ccttgacggg	catctgcgcc	540
gcattttcga	cagctttgtg	aactgggtgg	ccgagaagga	gtgggagttg	ccgccagatt	600

-continued

```

ctgacttgga tctgaatctg attgagcagg cacccttgac cgtggccgag aagetgcage 660
gcgactttct gacggagtgg cgcctgtgta gtaaggcccc ggaggccctt ttctttgtgc 720
aattttgagaa gggagagagc tacttccact tacactgtct cgtggaacc accggggtga 780
aatccttagt tttgggacgt ttcttgagtc agattcgcga aaaactgatt cagagaattt 840
accgcgggat cgagccgact ttgccaaact ggttcgcggt cacaaagacc agaaacggcg 900
cgggagggcg gaacaagggt gtggacgagt gctacatccc caattacttg cccccaaaa 960
cccagcctga gctccagtgg gcgtggacta atttagaaca gtatttaagc gcctgtttga 1020
atctcacgga gcgtaaacgg ttggtggcgc agcatctgac gcacgtgtcg cagacgcagg 1080
agcagaacaa agagaatcag aatcccaatt ctgacgcgcc ggtgatcaga tcaaaaactt 1140
cagccaggta catggagctg gtcgggtggc tcgtggacaa ggggattacc tcggagaagc 1200
agtggatcca ggaggaccag gcctcataca tctccttcaa tgcggcctcc aactcgcggt 1260
cccaaatcaa ggctgccttg gacaatgcgg gaaagattat gagcctgact aaaaccgccc 1320
ccgactacct ggtgggcccag cagcccgtgg aggacatttc cagcaatcgg atttataaaa 1380
ttttggaaact aaacgggtac gatccccaat atgcggcttc cgtctttctg ggatggggca 1440
cgaaaaagt t cggaagagg aacacccatct ggctgtttgg gcctgcaact accgggaaga 1500
ccaacatcgc ggaggccata gccccactg tgcccttcta cgggtgcgta aactggacca 1560
atgagaacta gtgcgcctg gactgtgtcg acaagatggg gatctggtgg gagggggga 1620
agatgaccgc caaggtcgtg gactcggcca aagccattct cggaggaagc aaggtgcgcg 1680
tggaccagaa atgcaagtc tgcggccaga tagaccgcac tcccgtgatc gtcacctcca 1740
acaccaacat gtgcgcctg attgacggga actcaacgac cttcgaacac cagcagccgt 1800
tgcaagaccg gatgttcaaa ttgaaactca cccgcgctct ggatcatgac ttgggaagg 1860
tcaccaagca ggaagtcaaa gactttttcc ggtgggcaaa ggatcacgtg gttgaggtgg 1920
agcatgaatt ctacgtcaaa aaggggtggag ccaagaaaaa acccgccccc agtgacgcag 1980
atataagtga gcccaaacgg gtgcgcgagt cagtgtgcga gccatcgacg tcagacgcgg 2040
aagcttcgat caactacgca gacaggtacc aaaacaaatg ttctcgtcac gtgggcatga 2100
atctgatgct ttctccctgc agacaatgcg agagaatgaa tcagaattca aatatctgct 2160
tcactcacgg acagaagac tgttttagagt gctttcccgt gtcagaatct caaccgcttt 2220
ctgtcgtcaa aaaggcgtat gacaaaactgt gctacattca tcatatcatg ggaaaggtgc 2280
cagacgcttg catcgctgc gatctgggtca atgtgggattt ggatgactgc atctttgaac 2340
aataa 2345

```

```

SEQ ID NO: 13      moltype = DNA length = 2079
FEATURE            Location/Qualifiers
source              1..2079
                    mol_type = other DNA
                    note = artificial sequence
                    organism = unidentified

```

```

SEQUENCE: 13
atcattgaga taattaaaa tataaccatc tcgcaataa ataagtat ttttcttct 60
gtaacagttt tgtataaaaa aaacctataa attgattcct caggtagtaa gtattcattg 120
taaatctgat attatttcta ttattatacc tacctaattt gcagtgcagc ccattacggc 180
agtataaatt cgttcatttt ggatattgtt tcagtgcggg ggttttacga gatttgtgatt 240
aaggtcccca gcgacatttg cgggcatctg cccggcattt ctgacagctt tgtgaattgg 300
gtggccgaga aggagtggga ttgcccgcga gattctgact tggatctgaa tctgattgag 360
caggcaccoc tgaccgtggc cgagaagctg cagcgcgact ttctgacgga gtggcgccgt 420
gtgagtgaag ccccgaggcg ccttttcttt gtgcaatttg agaagggaga gagctacttc 480
cacttacacg tgctcgtgga aaccaccggg gtgaaatcct tagttttggg acgtttcctg 540
agtcagattc gcgaaaaact gattcagaga atttaccgcg ggatcgagcc gactttgcc 600
aaactggctc gcgtcacaaa caccagaaac ggcgcgggag gcgggaacaa ggtggtggac 660
gagtgcatac tccccaatca ttgctcccc aaaacccagc ctgagctcca gtgggctggg 720
actaattaga aacagatttt aagcgcctgt ttgaatctca cggagcgtaa acgggtgggtg 780
gcgcagcatc tgacgcagct gtgcgagacg caggagcaga acaagagaa tcagaatccc 840
aattctgacg gcgcgggtgat cagatcaaaa acttcagcca ggtacatgga gctggctcggg 900
tggtcgtggt acaaggggat tacctcggag aagcagtgga tccaggagga ccaggccctca 960
tacatctcct tcaatgcggc ccccaactcg cgttcccaaa tcaaggctgc cttggacaat 1020
gcccggaaaga ttatgagcct gactaaaacc gcccccgcgt acctggtggg ccagcagccc 1080
gtggaggaca tttccagcaa tcggattttat aaaattttgg aactaaacgg gtacgatccc 1140
caatatcgcg cttccgtctt ttgggatgg gccacgaaaa agttcggcaa gaggaacacc 1200
atctggctgt ttgggctctg aactaccggg aagaccaaca tcgcggaggc catagcccac 1260
actgtgccct tctacgggtg cgtaaactgg accaatgaga actttccctt caacgactgt 1320
gtcgacaaga tggatgatct gtgggaggag gggaagatga ccgccaaggt cgtggagtgc 1380
gccaagacca ttctcggagg aagcaagggt cgcgtggacc agaaatgcaa gtctcggccc 1440
cagatagacc cgactcccgat gatcgtaacc tccaacacca acatgtgcgc cgtgattgac 1500
gggaactcaa cgactctcga acaccagcag ccggttgcaag accggatgtt caaattttaa 1560
ctcaccgccg gtctggatca tgactttggg aaggtcacca agcaggaagt caaagacttt 1620
ttccgggtggg caaaggatca cgtggttgag gtggagcatg aattctactg caaaaagggt 1680
ggagccaaga aaagaccgcg ccccgatgac gcagatataa gtgagcccaa acgggtgcgc 1740
gagtcagttg cgcagccatc gacgtcagac gcggaagctt cgatcaacta cgcagacagg 1800
taccaaaaca aatgttctcg tcacgtgggc atgaatctga tgctgtttcc ctgcagacaa 1860
tgcgagagaa tgaatcagaa ttcaaatatc tgcttcactc acggacagaa agactgttta 1920
gagtgcttcc ccgtgtcaga atctcaaccc gtttctgtcg tcaaaaaggc gtatcagaaa 1980
ctgtgctaca ttcatcatat catgggaaag gtgccagacg cttgcaactg ctgcgatctg 2040
gtcaatgtgg atttgatga ctgcattctt gaacaataa 2079

```

```

SEQ ID NO: 14      moltype = DNA length = 2130
FEATURE            Location/Qualifiers
source              1..2130
                    mol_type = other DNA
                    note = artificial sequence

```

-continued

```

organism = unidentified
SEQUENCE: 14
aataaacgat aacgccgttg gtggcggtgag gcatgtaaaa ggttacatca ttatcttgtt 60
cgccatccgg ttggtataaa tagacgttca tgttggtttt tgtttcagtt gcaagttggc 120
tgccggcgccg gcagcacctt tgccgggatcc tcaggtagta agtattcatt gtaaatctga 180
tattatttgt attattatac ctacctaat tgcagtgtag cccattacgg cagtataaat 240
tcgttcattt tggatattgt ttcagtgcgg ggggttttac agattgtgat taaggteccc 300
agcgaccttg acgggcatct gcccgccatt tctgacagct ttgtgaactg ggtggccgag 360
aaggagtggtg agttgccgcc agattctgac ttggatctga atctgattga gcaggcacc 420
ctgaccgttg ccgagaagct gcagcgcgac ttctcgacgg agtggcgccg tgtgagtaag 480
gccccggagg cccctttctt tgtgcaattt gagaagggag agagctactt ccacttacac 540
gtgctcgttg aaaccacggg ggtgaaatcc ttagttttgg gacgtttcct gagtacagatt 600
cgcgaaaaac tgattcagag aatttaccgc gggatcgagc cgaactttgc aaactgggtc 660
gcggtcacaa agaccagaaa cggcgccgga ggccgggaaca aggtgggtga cgaagtgtac 720
atcccccaat acttgctccc caaaacccag cctgagctcc agtggcggtg gactaattta 780
gaacagttat taagccctg tttgaatctc accggagcgt aacgggttgt ggccgagcat 840
ctgacgcacg tgtcgagcac gcaggagcag aacaagaga atcagaatcc caattctgac 900
gcgcgggtga tcagatcaaa aacttcagcc aggtacatgg agctggctgg gtggtcgtg 960
gacaagggga ttacctcgga gaagcagtg atccaggagg accaggcctc atacctctc 1020
ttcaatgcgg cctccaaact gcgggtcccaa ccttggaaca tgcgggaaa 1080
attatgagcc tgactaaaac cgcgcccgac tacctgggtg gccagcagcc cgtggaggac 1140
atttccagca atcggattta taaaattttg gaactaaacg ggtacgatcc ccaatagcg 1200
gcttcgctct tctgggatg gggcacgaaa aagttcgcca agaggaaac catctggctg 1260
tttgggctg caactaccgg gaagaccaac atcgccggag ccatagccca cactgtgccc 1320
ttctacgggt gcgtaaactg gaccaatgag aactttccct tcaacgactg tgcgacaag 1380
atgggtgatc ggtgggagga ggggaagatg accgccaaag tcgtggagtc ggccaaagcc 1440
attctcggag gaagcaaggt gcgcgtggac cagaaatgca agtccctggc ccagatagac 1500
ccgactcccg tcatcgtcac ctccaacacc aacatgtgcg ccgtgattga cgggaactca 1560
acgaccttcg aacaccgaca gccgttgcaa gaccggatgt tcaaaattga actcaccgcc 1620
cgtctggatc atgactttgg gaaggtcacc aagcagggaag tcaaaagact tttccggtgg 1680
gcaaaaggatc accgtggtga ggtggagcat gaattctacg tcaaaaaggg tggagccaag 1740
aaaaagaccg ccccgagtg cgcagatata agtgagccca aacgggtgcg cgaagtcatg 1800
gcgcagccat cgacgtcaga cgcggaagct tcgatcaact acgcagacag gtacccaaac 1860
aaatgttctc gtcacgtggg catgaatctg atgctgttcc cctgcagaca atgcgagaga 1920
atgaatcaga attcaaatat ctgcttcact cacggacaga aagactgttt agagtcttt 1980
cccggtgcag aatctcaacc cgtttctgtc gtcaaaaagg cgtatcagaa actgtgtctc 2040
attcatcata tcatgggaaa ggtgcagag gcttgccactg cctgcgactc ggtcaatgtg 2100
gatttgatg actgcatctt tgaacaataa 2130

```

```

SEQ ID NO: 15      moltype = DNA length = 2314
FEATURE            Location/Qualifiers
source              1..2314
                    mol_type = other DNA
                    note = artificial sequence
                    organism = unidentified

```

```

SEQUENCE: 15
cacatgttgg acatcgtgtc gtttgagcgt ataaaagaat atataagagc taatttaggc 60
catttcacag taatcaccga caaatgttcg aagcgtaagg tgtgtcttca tcacaaacga 120
attgccagg tgttgggcat taaaaaaata tatcatcaag aatacaaacg ggttgtttca 180
aagggtttca agaagcaaac attgtaaacg tgccggagca acaatctccg gagactgcgg 240
ccgtgtgcaa aaattaaaag ctgttgaata aactggaatc gagctcttac aacaaatcca 300
acattgacca tctggccgtt attgtaaaag atcctcaggt agtaagtatt cattgtaaat 360
ctgatattat ttgtattatt atacctacct aatttgcagt gcagccatt acggcagtat 420
aaatcgttc attttggata ttgtttcagt gccggggttt tacgagattg tgattaaggt 480
ccccagcgac ctgacggggc atctgcccgg catttctgac agcttttgtga actgggtggc 540
cgagaaggag tgggagttgc cgcagattc tgacttggat ctgaatctga ttgagcaggc 600
acccctgacc gtggccgaga agctgcagcg cgactttctg acggagtggc gccgtgtgag 660
taaggccccg gaggcccttt tcttttgtca atttgagaag ggagagagct acttccactt 720
acacgtgtc gtggaaacca cgggggtgaa atccttagtt ttgggagctt tctgagtca 780
gattcgcgaa aaactgatc agagaattta ccgcgggatc gagccgactt tgccaaactg 840
gttcgcggtc acaagacca gaaacggcgc cggaggcggg aacaaggtgg tggacgagt 900
ctacatcccc aattacttgc tccccaaaac ccagcctgag ctccagtggg cgtggactaa 960
tttagaacag tatattaagg ctggtttgaa tctcacggag cgtaaacggt ttgtggcgca 1020
gcatctgacg cacgtgtcgc acagcgagga gcagaacaaa gagaatcaga atcccaattc 1080
tgacgcgcgg gtgatcagat caaaaacttc agccaggtac atggagctgg tcgggtggct 1140
cgtggacaag gggattacct cggagaagca gtggatccag gaggaccagg cctcatcat 1200
ctccttcaat gcggctccca actcgcggtc ccaaatcaag gctgccttgg acaatgcggg 1260
aaagattatg agcctgacta aaaccgcccc cgactacctg gtgggcccag agcccggtga 1320
ggacatttcc agcaatcgga ttataaaaat ttggaaacta aacgggtacg atccccaa 1380
tgccggttcc gtctttcttg gatgggccac gaaaaagttc ggcaagagga acaccatctg 1440
gctgttttgg cgtgcaacta ccgggaagac caacatcgcg gaggccatag cccacactgt 1500
gcccctctac ggtgctgtaa actggaccaa tgagaacttt ccttcaacg actgtgtcga 1560
caagatggtg atctggtggg aggaggggaa gatgaccgcc aaggtcgtgg agtcggccaa 1620
agccattctc ggaggaagca agtgccgct ggaccagaaa tgcaagtcc cgccagat 1680
agaccggact ccggtgatcg tccctccaa caccaacatg tgcgccgtga ttgacgggaa 1740
ctcaacgacc ttgaacacc agcagccgtt gcaagaccgg atgttcaaat ttgaactcac 1800
cgccgctctg gatcatgact ttgggaaggt caccaagcag gaagtcaaa actttttccg 1860
tggggcaaa gatcacgtgg ttgagtgga gcatgaattc tacgtcaaaa aggggtggagc 1920
caagaaaaga cccgccccca gtgacgcaga tataagttag cccaacaggg tgcgcgagtc 1980

```

-continued

```

agttgctgcag ccatcgacgt cagacgcgga agcttcgatc aactacgcag acaggtacca 2040
aaacaaatgt tctcgtcacg tgggcagtaa tctgatgctg ttccctgca gacaatgcga 2100
gagaatgaat cagaatctaa atatctgctt cactcacgga cagaagact gtttagagtg 2160
ctttcccggtg tcagaatctc aaccgcgttc tgcctgcaaa aaggcgtatc agaaactgtg 2220
ctacattcat catatcatgg gaaaggtgcc agacgcttgc actgcctgcg atctggtcaa 2280
tgtggatttg gatgactgca tctttgaaca ataa 2314

```

```

SEQ ID NO: 16      moltype = DNA length = 2045
FEATURE           Location/Qualifiers
source            1..2045
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 16
gggagttcag tcgtcgaatg caaaacgtaa aaaatattaa taaggtaaaa actacagctg 60
gatctcagg tagtaagtat tcattgtaaa tctgatatta ttgtattat tatacctacc 120
taatttgca tgcagcccat tacggcagta taaattcgtt cattttggat attgtttcag 180
tgccgggggtt ttacgagatt gtgattaagg tccccagca ccttgacggg catctgcccg 240
gcatttctga cagctttgtg aactgggtgg ccgagaagga gtgggagttg ccgccagatt 300
ctgactttga tctgaatctg attgagcagg caccctgac cgtggccgag aagctgcagc 360
gcgactttct gacggagtggt gcgcgtgtga gtaaggcccc ggaggccctt ttctttgtgc 420
aatttgagaa gggagagagc tacttccact tacacgtgct cgtggaaacc accggggtga 480
aatccttagt tttgggacgt ttcttgagtc agattcgcga aaaactgatt cagagaattt 540
accgcgggat cgagccgact ttgccaaact ggttcgcggt cacaagacc agaaacggcg 600
ccggaggcgg gaacaagggt gtggacgagt gctacatccc caattacttg ctccccaaaa 660
cccagcctga gctccagtgg gcttggaact atttagaaca gtatttaagc gctgttttga 720
atctcacgga gcgtaaacgg ttggtggcgc agcatctgac gcacgtgtcg cagacgcagg 780
agcagaacaa agagaatcag aatcccaatt ctgacgcgcc ggtgatcaga tcaaaaactt 840
cagccaggtg catggagctg gtcgggtggc tcgtgggcaa ggggattacc tcggagaagc 900
agtggatcca ggaggaccag gcctcataca tctccttcaa tgccgcctcc aactgcgctg 960
cccaaatcaa ggctgccttg gacaatgcgg gaaagattat gagcctgact aaaaccgccc 1020
ccgactacct ggtggggccag cagcccgtgg aggcacattc cagcaatcgg atttataaaa 1080
ttttggaact aaacgggtac gatcccaat atgcgggcttc cgtctttctg ggatgggcca 1140
cgaaaaagtt gggcaaggag aacaccatct ggctgttttg gcctgcaact accgggaaga 1200
ccaacatcgc ggaggccata gccccactg tgccttcta cgggtgcgta aactggacca 1260
atgagaactt tcccttcaac gactgtgtcg acaagatggt gatctgttg ggaggaggga 1320
agatgaccgc caaggtcgtg gactcggcca aagccattct cggagggaagc aaggtgcgcg 1380
tggaccgcaa atgcaagtcc tcgggccaga tagaccgcac tcccgatgc gtcacctcca 1440
acaccaacat gtgcgccgtg attgacggga actcaacgac cttcgaacac cagcagccgt 1500
tgcaagaccg gatgttcaaa ttgaaactca cccgcgtct ggaatcagac ttggaagga 1560
tcaccaagca ggaagtcaaa gactttttcc ggtgggcaaa ggatcacgtg gttgaggtgg 1620
agcatgaatt ctacgtcaaa aaggggtggag ccaagaaaaa acccgcccc agtgacgcag 1680
atataagtga ccccaaacgg gtgcgcgagt cagttgcgca gccatcgacg tcagacgcgg 1740
aagcttcgat caactacgca gacaggtacc aaaacaaatg ttctcgtcac gtgggcattg 1800
atctgatgct gtttccctgc agacaatgcg agagaatgaa tcagaattca aatatctgct 1860
tcactcagg acagaaagac tgttttagagt gctttcccg gtcagaatct caaccggtt 1920
ctgtcgtcaa aaaggcgtat cagaaactgt gctacattca tcatatcatg ggaagggtgc 1980
cagacgcttg cactgcctgc gatctggtca atgtggattt ggatgactgc atctttgaac 2040
ataaa 2045

```

```

SEQ ID NO: 17      moltype = DNA length = 144
FEATURE           Location/Qualifiers
source            1..144
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 17
aataaacgat aacgccgttg gtggcgtgag gcatgtaaaa ggttacatca ttatcttgtt 60
cgccatccgg ttggtataaa tagacgttca tgttggtttt tgtttcagtt gcaagttggc 120
tcggcgccgc gcagcacctt tgcg 144

```

```

SEQ ID NO: 18      moltype = DNA length = 92
FEATURE           Location/Qualifiers
source            1..92
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 18
atcatggaga taattaaaat gataaccatc tcgcaaataa ataagtattt tactgttttc 60
gtaacagttt tgtaataaaa aaacctataa at 92

```

```

SEQ ID NO: 19      moltype = DNA length = 2222
FEATURE           Location/Qualifiers
source            1..2222
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 19
aataaacgat aacgccgttg gtggcgtgag gcatgtaaaa ggttacatca ttatcttgtt 60

```


-continued

```

cgccatccgg ttggtataaa tagacgttca tgttggtttt tgtttcagtt gcaagttggc 120
tgcggcgcgc gcagcacctt tgcgggatcc tcaggtagta agtattcatt gtaaatctga 180
tattatttgt attattatac ctacctaatt tgcagtgacg atcatggaga taattaaaat 240
gataaccatc tcgcaaataa ataagtattt tactgttttc gtaacagttt tgtaataaaa 300
aaacctataa atcccattac ggcagtataa attcgttcat tttggatatt gtttcagtg 360
cggggtttta cgagattgtg attaagggtcc ccagcgacct tgacgggcat ctgcccggca 420
tttctgacag ctttgtgaac tgggtggccg agaaggagtg ggagttgccg ccagattctg 480
acttggatct gaatctgatt gagcaggcac cctgaccgt ggcgagaag ctgcagcgcg 540
actttctgac ggagtgccgc cgtgtgagta agggcccgga ggcccttttc tttgtgcaat 600
ttgagaaggg agagagctac ttccacttac acgtgctcgt ggaaaccacc ggggtgaaat 660
ccttagtttt gggacgtttc ctgagtcaga ttgcgaaaa actgattcag agaatttacc 720
gcgggatcga gccgactttg ccaaaactgt tgcgggtcac aaagaccaga aacggcgccg 780
gaggcgggaa caaggtgggtg gacgagtgct acatcccaa ttacttgctc cccaaaacc 840
agcctgagct ccagtggcg tggaactaatt tagaacagta ttaagcgcc tgtttgaatc 900
tcacggagcg taaacggttg gtggcgacgc atctgacgca cgtgtcgag acgcaggagc 960
agaacaaaga gaatcagaat cccaattctg acgcgccgt gatcagatca aaaacttcag 1020
ccaggtacat ggagctggtc ggggtggctg tggacaagg gattacctcg gagaagcagt 1080
ggatccagga ggaccaggcc tcatacatct ccttcaatgc ggctccaac tcgcggtccc 1140
aaatcaaggc tgccttgagc aatgcgggaa agattatgag cctgactaaa accgccccg 1200
actacctggt gggccagcag cccgtggagg acatttcag caatcgatt tataaaattt 1260
tggaactaaa cgggtacgat ccccaatag cggcttccgt ctttctggga tgggccacga 1320
aaaagtctcg caagaggaac accatctggc tgtttgggcc tgcaactacc ggggaagcca 1380
acatcgcgga ggcacatagc cacactgtgc ccttctacgg gtgcgtaaac tggaccaatg 1440
agaactttcc cttcaacgac tgtgtcgaca agatgggat cgtgtgggag gagggaaga 1500
tgaccgcaa ggtcgtggag tcggccaaag ccattctcgg aggaagcaag gtgcgctgg 1560
accagaaatg caagtctcgc gccagatag acccgactcc cgtgatcgtc acctccaaca 1620
ccaacatgtg cgcgtgatt gacgggaact caacgacct cgaacaccag cagccgttgc 1680
aagaccgat gttcaaattt gaactcacc gccgtctgga tcatgacttt gggaaggtca 1740
ccaagcagga agtcaaaagc tttttccggt gggcaaaagga tcacgtgggt gaggtggagc 1800
atgaattcta cgtcaaaaag ggtggagcca agaaaagacc cgccccagc gacgcagata 1860
taagtgaacc caaacgggtg cgcgagtcag ttgcgcagcc atcgacgtca gacgcggaag 1920
cttcgatcaa ctacgcagac aggtacaaaa acaaatgttc tcgtcacgtg ggcatgaatc 1980
tgatgctgtt tccctgcaga caatgcgaga gaattgaatc gaattcaaat atctgcttca 2040
ctcacggaca gaaagactgt tttagagtgct ttcccgctgc agaactctca cccgtttctg 2100
tcgtcaaaaa ggcgtatcat aaactgtgct acattcatca tatcatggga aaggtgccag 2160
acgcttgcac tgctcgcat ctggtcaatg tggatttgga tgactgcatc tttgaacaat 2220
aa 2222

```

```

SEQ ID NO: 20      moltype = DNA length = 328
FEATURE           Location/Qualifiers
source            1..328
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 20
cacatgttgg acatcgtgtc gtttgagcgt ataaaagaat atataagagc taatttaggc 60
catttcacag taatcacgga caaatgttcg aagcgtaagg tgtgtcttca tcacaaacga 120
attgccaggt tgttgggcat taaaaaata tatcatcaag aatacaaacg ggttgtttca 180
aaggtttaca agaagcaaac attgtaaacg tgccggagca acaatctccg gagactgcgg 240
cgtgtgcaaa aaattaaaag ctgttgaata aactggaatc gagctcttac aacaaatcca 300
acattgacca gctggcgctt attgtaaa 328

```

```

SEQ ID NO: 21      moltype = DNA length = 59
FEATURE           Location/Qualifiers
source            1..59
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 21
gggagttcag tcgtcgaaatg caaacgtaa aaaatattaa taaggtaaaa actacagct 59

```

```

SEQ ID NO: 22      moltype = DNA length = 110
FEATURE           Location/Qualifiers
source            1..110
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 22
gacctttaat tcaacccaac acaatatatt atagttaa atagaattatt atcaaatcat 60
ttgtatatta attaaaatac tatactgtaa attacatttt atttacaatc 110

```

```

SEQ ID NO: 23      moltype = DNA length = 2114
FEATURE           Location/Qualifiers
source            1..2114
                  mol_type = other DNA
                  note = artificial sequence
                  organism = unidentified

```

```

SEQUENCE: 23
atcatggaga taattaaaat gataaccatc tcgcaaataa ataagtattt tactgttttc 60

```

-continued

```

gtaacagttt tgaataaaaa aaacctataa atattccgga ttattcatat cgtcccacca 120
tcgggcgcgg atcctcaggt agtaagtatt cattgtaaat ctgattattat ttgtattatt 180
atacctacct aatttgagct gcagccatt acggcagtat aaattcggtt attttggata 240
ttgtttcagt gccggggttt tacgagattg tgattaaggt ccccgagcag cttgacgggc 300
atctgcccgg catttctgac agctttgtga actgggtggc cgagaaggag tgggagttgc 360
cgccagatct tgacttgat ctgaatctga ttgagcaggc acccctgacc gtggccgaga 420
agctgcagcg cgactttctg acggagtggc gccgtgtgag taaggcccg gaggccttt 480
tctttgtgca atttgagaag ggagagagct acttccactt acacgtgtct gtggaacca 540
cgggggtgaa atccttagtt ttgggacgtt tctctgagtc gattcgcgaa aaactgattc 600
agagaattta ccgcgggatc gagccgactt tgccaaactg gttcgcggtc acaaagacca 660
gaaacggcgc cggaggcggg aacaagggtg tggacgagtg ctacatcccc aattacttgc 720
tccccaaac ccagcctgag ctccagtggt cgtggactaa tttagaacag tatttaagcg 780
cctggttgaa tctcacggag cgtaaacggt tgggtggcga gcatctgacg cacgtgtcgc 840
agacgcagga gcagaacaaa gagaatcaga atcccaattc tgacgcgcgc gtgatcagat 900
caaaaacttc agccaggtac atggagctgg tcgggtggct cgtggacaag gggattacct 960
cggagaagca gtggatccag gaggaccagg cctcatatcat ctcttcaat gcggcctcca 1020
actcgcgttc ccaaatcaag gctgccttgg acaatgcggg aaagattatg agcctgacta 1080
aaaccgcccc cgactacctg gtgggccagc agcccggtga ggacatttcc agcaatcgga 1140
ttataaaat tttggaacta aacgggtacg atccccataa tgcggcttcc gtctttctgg 1200
gatggggcac gaaaaagttc ggcaagagga acaccatctg gctgtttggg cctgcaacta 1260
ccgggaagac caacatcgcg gaggccatag cccacactgt gcccttctac ggggtgcgtaa 1320
actggaccaa tgagaacttt ccctcaacg actgtgtcga caagatggtg atctggtggg 1380
aggaggggaa gatgaccgcc aaggtcgtgg agtcggccaa agccattctg ggaggaagca 1440
aggtgcgcgt ggaccagaaa tgcaagtcct cggcccagat agaccgcact cccgtgatcg 1500
tcactccaa caccacatg tcgcgcgtga ttgacgggaa ctcaacgacc ttcgaacacc 1560
agcagccgtt ccaagaccgg atgttcaaat ttgaactcac ccgcgctctg gatcatgact 1620
ttgggaaggt caccaagcag gaagtcaaa actttttccg gtgggcaaa gatcacgtgg 1680
ttgaggtgga tcatgaattc tacgtcaaaa aggggtggagc caagaaaaga cccgccccca 1740
gtgacgcaga tataagttag cccaaacggg tcgcgcgagtc agttgcgcag ccatcgacgt 1800
cagacgcgga agcttcgatc aactacgcag acaggtacca aaacaaatgt tctcgtcacg 1860
tgggcatgaa tctgatctg ttccctgca gacaatgcga gagaatgaat cagaattcaa 1920
atatctgctt cactcacgga cagaagagct gtttagagtg ctttccgtg tcagaatctc 1980
aaccggttct tgtcgtcaaa aaggcgtatc agaaactgtg ctacattcat catatcatgg 2040
gaaaggtgcc agacgcttgc actgcctgcg atctggtcaa tgtggatttg gatgactgca 2100
tctttgaaca ataa 2114

```

The invention claimed is:

1. An expression cassette containing overlapping open reading frames, the overlapping open reading frames being overlapping open reading frames of a first ORF and a second ORF, comprising a second promoter used only to drive gene transcription of the second ORF and further comprising in sequence from a 5' end to a 3' end:

a first promoter at least used to drive gene transcription of the first ORF;
a 5' part of a gene of the first ORF;
an intron; and
a 3' part of a gene of the second ORF,
wherein the second promoter precedes the 5' end of the first promoter or within the intron,
wherein during post-transcriptional processing, the translation initiation codons ATG of the first ORF is formed through alternative splicing of the intron.

2. The expression cassette according to claim 1, wherein the intron is located between any two nucleotides of the translation initiation codons ATG of the first ORF.

3. The expression cassette according to claim 1, wherein the second promoter is within the intron.

4. The expression cassette according to claim 1, wherein the second promoter precedes the 5' end of the first promoter.

5. The expression cassette according to claim 1, wherein the first promoter is only used to drive gene transcription of the first ORF.

6. The expression cassette according to claim 1, wherein the first promoter is used to drive gene transcription of the first ORF and the second ORF.

7. The expression cassette according to claim 1, wherein the gene of the first ORF is used to express a Rep78 and/or Rep68 protein, and the gene of the second ORF is used to express a Rep52 and/or Rep 40 protein.

8. The expression cassette according to claim 1, wherein the gene of the first ORF is used to express a VP1 protein, and the gene of the second ORF is used to express a VP2 and/or VP3 protein(s).

9. The expression cassette according to claim 1, wherein the first promoter is active before the second promoter.

10. The expression cassette according to claim 9, wherein the first promoter is an immediate early promoter, an early promoter, or a late promoter, and the second promoter is a very late promoter.

11. The expression cassette according to claim 1, wherein the first promoter is deltaIE1, IE1, 39K, or p6.9, and the second promoter is polH or p10.

12. The expression cassette according to claim 1, wherein the first promoter is p10, and the second promoter is polH.

13. A nucleic acid molecule, comprising a Cap gene expression cassette and a Rep gene expression cassette, and the Cap gene expression cassette and/or the Rep gene expression cassette is the expression cassette according to claim 1.

14. The nucleic acid molecule according to claim 13, further comprising an exogenous gene.

15. The nucleic acid molecule according to claim 14, wherein AAV inverted terminal repeats are provided at both ends of the exogenous gene.

16. The nucleic acid molecule according to claim 14, wherein the exogenous gene is a reporter gene, and the reporter gene is at least one of a chloramphenicol acetyltransferase encoding gene, a β -galactosidase encoding gene, a β -glucuronidase encoding gene, a *Renilla luciferase* encoding gene, an alkaline phosphatase encoding gene, a firefly luciferase encoding gene, a green fluorescent protein encoding gene, and a red fluorescent protein encoding gene.

47

48

17. The nucleic acid molecule according to claim 14, wherein the exogenous gene is a gene encoding a drug polypeptide, and the drug polypeptide is at least one of lipoprotein esterase, apolipoprotein, cytokine, interleukin, and interferon.

5

18. A recombinant baculovirus vector, comprising the expression cassette according to claim 1.

19. The recombinant baculovirus vector according to claim 18, wherein the recombinant baculovirus vector is an insect cell compatible vector.

10

20. An isolated insect cell, comprising the recombinant baculovirus vector according to claim 18.

21. The insect cell according to claim 20, wherein the insect cell is a *Spodoptera frugiperda* cell, a *Trichopodia* cell, a *Drosophila* cell, or a mosquito cell.

15

* * * * *