

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12397016
Kind Code	B2
Date of Patent	August 26, 2025
Inventor(s)	Arnold; Katelyn et al.

Sulfated oligosaccharides having anti-inflammatory activity

Abstract

Provided herein are small molecule compounds, including non-anticoagulant heparan sulfate oligosaccharide molecules, having anti-inflammatory properties and capable of interacting with high mobility group box 1 (HMGB1) proteins in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation end products (RAGE). Also provided herein are methods of treating Paracetamol (APAP) overdose in subjects.

Inventors: Arnold; Katelyn (Durham, NC), Xu; Ding (East Amherst, NY), Xu; Yongmei (Durham, NC), Pawlinski; Rafal (Chapel Hill, NC), Liu; Jian (Chapel Hill, NC)

Applicant: The University of North Carolina at Chapel Hill (Chapel Hill, NC); THE RESEARCH FOUNDATION FOR THE STATE UNIVERSITY OF NEW YORK (Amherst, NY)

Family ID: 1000008778531

Assignee: The University of North Carolina at Chapel Hill (Chapel Hill, NC); THE RESEARCH FOUNDATION FOR THE STATE UNIVERSITY OF NEW YORK (Amherst, NY)

Appl. No.: 18/406942

Filed: January 08, 2024

Prior Publication Data

Document Identifier	Publication Date
US 20240216419 A1	Jul. 04, 2024

Related U.S. Application Data

division parent-doc US 16761159 US 11865137 WO PCT/US2018/059152 20181105 child-doc US 18406942

Publication Classification**Int. Cl.:** A61K31/727 (20060101)**U.S. Cl.:**

CPC A61K31/727 (20130101);

Field of Classification Search**USPC:** None

References Cited**U.S. PATENT DOCUMENTS**

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4554101	12/1984	Hopp	N/A	N/A
4865870	12/1988	Hu et al.	N/A	N/A
5527785	12/1995	Bevilacqua	514/56	A61K 31/715
5543403	12/1995	Petitou et al.	N/A	N/A
5817487	12/1997	Kobayashi et al.	N/A	N/A
5834282	12/1997	Habuchi et al.	N/A	N/A
5935824	12/1998	Sgariato	N/A	N/A
6255088	12/2000	Wong et al.	N/A	N/A
6608044	12/2002	Aderka et al.	N/A	N/A
6861254	12/2004	Rosenberg et al.	N/A	N/A
6977248	12/2004	Shukla et al.	N/A	N/A
7101859	12/2005	Yedgar et al.	N/A	N/A
7531338	12/2008	Liu	N/A	N/A
8771995	12/2013	Liu et al.	N/A	N/A
9951149	12/2017	Liu et al.	N/A	N/A
10286047	12/2018	Spirig et al.	N/A	N/A
11203772	12/2020	Xu et al.	N/A	N/A
11633424	12/2022	Liu et al.	N/A	N/A
11865137	12/2023	Arnold et al.	N/A	N/A
11903963	12/2023	Liu et al.	N/A	N/A
11993627	12/2023	Liu et al.	N/A	N/A
2003/0083294	12/2002	Sullenger	N/A	N/A
2003/0099967	12/2002	Deangelis	N/A	N/A
2004/0087492	12/2003	Yedgar et al.	N/A	N/A
2004/0191870	12/2003	Rosenberg et al.	N/A	N/A
2004/0259142	12/2003	Chai et al.	N/A	N/A
2005/0090601	12/2004	Dadalas et al.	N/A	N/A
2005/0090661	12/2004	Asari et al.	N/A	N/A
2005/0101532	12/2004	Yang et al.	N/A	N/A
2005/0191288	12/2004	Bennett et al.	N/A	N/A

2005/0225562	12/2004	Higgins et al.	N/A	N/A
2005/0255562	12/2004	Rosenberg et al.	N/A	N/A
2005/0282775	12/2004	Kennedy	514/56	A61K 31/727
2006/0165673	12/2005	Liu	N/A	N/A
2006/0172931	12/2005	San Antonio et al.	N/A	N/A
2006/0229276	12/2005	Hook et al.	N/A	N/A
2008/0109236	12/2007	DeAngelis	N/A	N/A
2009/0035787	12/2008	Liu	N/A	N/A
2009/0155851	12/2008	Sugiura et al.	N/A	N/A
2009/0197308	12/2008	Liu et al.	N/A	N/A
2010/0125052	12/2009	Lu et al.	N/A	N/A
2010/0298260	12/2009	Sundaram	536/123.13	C12P 19/04
2010/0305022	12/2009	Shriver	N/A	N/A
2011/0054236	12/2010	Yang et al.	N/A	N/A
2011/0281819	12/2010	Kakehi et al.	N/A	N/A
2012/0064044	12/2011	Egan	N/A	N/A
2012/0308546	12/2011	Kizhakkedathu et al.	N/A	N/A
2012/0322114	12/2011	Liu et al.	N/A	N/A
2012/0322760	12/2011	Fier et al.	N/A	N/A
2013/0022647	12/2012	Kizhakkedathu et al.	N/A	N/A
2013/0296540	12/2012	Xu et al.	N/A	N/A
2013/0338097	12/2012	Stephens et al.	N/A	N/A
2016/0122446	12/2015	Liu et al.	N/A	N/A
2021/0137967	12/2020	Liu et al.	N/A	N/A
2021/0169923	12/2020	Arnold et al.	N/A	N/A
2021/0260098	12/2020	Liu et al.	N/A	N/A
2021/0332076	12/2020	Liu et al.	N/A	N/A
2022/0265699	12/2021	Arnold et al.	N/A	N/A
2022/0416486	12/2021	Yamaguchi	N/A	N/A
2024/0066048	12/2023	Liu et al.	N/A	N/A
2024/0309035	12/2023	Liu et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
2003247808	12/2003	AU	N/A
103402526	12/2012	CN	N/A
111601603	12/2019	CN	N/A
112437667	12/2020	CN	N/A
105452479	12/2020	CN	N/A
114980904	12/2021	CN	N/A
0 394 971	12/1989	EP	N/A
0 565 863	12/1992	EP	N/A
1885865	12/2007	EP	N/A
3011043	12/2015	EP	N/A
2910476	12/2021	ES	N/A

2924830	12/2021	ES	N/A
2002-534375	12/2001	JP	N/A
2005-508827	12/2004	JP	N/A
2011-168591	12/2010	JP	N/A
2014-501730	12/2013	JP	N/A
2016-523535	12/2015	JP	N/A
6670235	12/2019	JP	N/A
2021502470	12/2020	JP	N/A
2021528421	12/2020	JP	N/A
2023501568	12/2022	JP	N/A
2023-166387	12/2022	JP	N/A
2024056723	12/2023	JP	N/A
7495061	12/2023	JP	N/A
7538724	12/2023	JP	N/A
WO 89/04328	12/1988	WO	N/A
WO 93/05167	12/1992	WO	N/A
WO 96/14425	12/1995	WO	N/A
WO0151003	12/2000	WO	N/A
WO2003018598	12/2002	WO	N/A
WO 2004/005475	12/2003	WO	N/A
WO 2004/009642	12/2003	WO	N/A
WO 2004/017910	12/2003	WO	N/A
WO 2004/050673	12/2003	WO	N/A
WO2005/118609	12/2004	WO	N/A
WO 2006/124801	12/2005	WO	N/A
WO2009/014715	12/2008	WO	N/A
WO 2009/079693	12/2008	WO	N/A
WO 2012/088416	12/2011	WO	N/A
WO 2012/116048	12/2011	WO	N/A
WO 2014/204929	12/2013	WO	N/A
WO 2018/165656	12/2017	WO	N/A
WO 2019/010216	12/2018	WO	N/A
WO 2019/090203	12/2018	WO	N/A
WO 2019/246264	12/2018	WO	N/A
WO 2021/097345	12/2020	WO	N/A

OTHER PUBLICATIONS

Sommers, Journal of Pharmaceutical and Biomedical Analysis 140 (2017) 113-121. (Year: 2017).
cited by examiner

Nelson, Blood, vol. 82, No. 11 Dec. 1, 1993: pp. 3253-3258. (Year: 1993). cited by examiner

Advisory Action corresponding to U.S. Appl. No. 13/996,930 dated Dec. 9, 2016, 6 Pages. cited by
applicant

Aikawa, J.I., et al., "Molecular Cloning and Expression of a Third Member of the Heparan
Sulfate/Heparin GlcNAc N-Deacetylase/N-Sulfotransferase Family," The Journal of Biological
Chemistry, vol. 274, No. 5, 1999, pp. 2690-2695. cited by applicant

Aikawa, J.I., et al., "Multiple Isozymes of Heparan Sulfate/Heparin GlcNAc N-Deacetylase/GlcN
N-Sulfotransferase," The Journal of Biological Chemistry, vol. 276, No. 8, pp. 5876-5882 (Feb. 23,
2001). cited by applicant

Antoine, D.J., et al., "Mechanistic Biomarkers Provide Early And Sensitive Detection Of
Acetaminophen-Induced Acute Liver Injury At First Presentation To Hospital," Hepatology vol. 58,

pp. 777-787 (2013). cited by applicant

Applicant-Initiated Interview Summary corresponding to U.S. Appl. No. 13/996,930 dated Jan. 23, 2017. cited by applicant

Arepally, G, M., et al., "Clinical Practice. Heparin-Induced Thrombocytopenia," *N. Eng. J. Med.*, vol. 355, pp. No. 8, pp. 809-817, Aug. 24, 2006. cited by applicant

Arnold, K., et al., "Design Of Anti-Inflammatory Heparan Sulfate To Protect Against Acetaminophen-Induced Acute Liver Failure," *Sci. Transl. Med.*, vol. 12, No. 535, pp. 1-26, Article ID eaav8075 (Mar. 18, 2020). cited by applicant

Arnold, K., et al., "Potential Use of Anti-Inflammatory Synthetic Heparan Sulfate to Attenuate Liver Damage," *Biomedicines*, vol. 8, 0503, 1-15 pages, 2020. cited by applicant

Arnold, K., et al., "Synthetic anticoagulant heparan sulfate attenuates liver ischemia reperfusion injury," *Sci. Reports*, vol. 10, Article No. 17187 (10 pages) (2020). cited by applicant

Arungundram, S., et al., "Modular Synthesis of Heparan Sulfate Oligosaccharides for Structure-Activity Relationship Studies," *J. Am. Chem. Soc.*, vol. 131, pp. 17394-17405, Dec. 2, 2009. cited by applicant

Ashikari-Hada, S., et al., "Characterization of growth factor-binding structures in heparin/heparan sulfate using an octasaccharide library," *J. Biol. Chem.*, vol. 279, No. 13, pp. 12346-12354, Mar. 26, 2004. cited by applicant

Axelsson, J., et al., "Inactivation Of Heparan Sulfate 2-O-Sulfotransferase Accentuates Neutrophil Infiltration During Acute Inflammation In Mice," *Blood*, vol. 120, pp. 1742-1751 (2012). cited by applicant

Bailey, G.P., et al., "Delays During The Administration Of Acetylcysteine For The Treatment Of Paracetamol Overdose," *Br. J. Clin. Pharmacol.* vol. 62, pp. cited by applicant

Baleux, F., et al., "A Synthetic CD4-Heparan Sulfate Glycoconjugate Inhibits CCR5 And CXCR4 HIV-1 Attachment and Entry," *Nat. Chem. Biol.*, vol. 5, No. 10, pp. 743-748, Oct. 2009. cited by applicant

Bame, K. J., et al., "Undersulfated Heparan Sulfate In A Chinese Hamster Ovary Cell Mutant Defective In Heparan Sulfate N-Sulfotransferase," *J Biol. Chem*, vol. 264, No. 14, pp. 8059-8065, May 15, 1989. cited by applicant

Beeson, J.G., et al., "Inhibition of Binding of Malaria-Infected Erythrocytes by a Tetradecasaccharide Fraction from Chondroitin Sulfate A," *Infection and Immunity*, vol. 66 No. 7 pp. 3397-3402 (Year: 1998). cited by applicant

Belot, F., et al., "Syntheses Of Chondroitin 4- And 6-Sulfate Pentasaccharide Derivatives Having A Methyl Beta-D-Glucopyranosiduronic Acid At The Reducing End," *Carbohydr. Res.*, vol. 326, pp. 88-97. (Year: 2000). cited by applicant

Bianchi, M.E., et al., "High-Mobility Group Box 1 Protein Orchestrates Responses To Tissue Damage Via Inflammation, Innate And Adaptive Immunity, And Tissue Repair," *Immunol. Rev.* vol. 280, pp. 74-82 (2017). cited by applicant

Bitter T., et al., "A Modified Uronic Acid Carbazole Reaction," *Anal. Biochem.*, vol. 4, pp. 330-334, 1962. cited by applicant

Blieden, M., et al., "A Perspective On The Epidemiology Of Acetaminophen Exposure And Toxicity In The United States," *Expert Rev. Clin. Pharmacol.* vol. 7, pp. 341-348 (2014). cited by applicant

Bourgeaux, V., et al., "Two-Step Enzymatic Synthesis Of UDP-N-Acetylgalactosamine," *Bioorg. Med. Chem. Lett.*, vol. 15, pp. 5459-5462 (2005). cited by applicant

Bowman, K.G., et al., "Carbohydrate Sulfotransferases: Mediators Of Extracellular Communication," *Chemistry & Biology*, vol. 6, pp. R9-R22 (Jan. 1999). cited by applicant

Bradbury, E.J., et al., "Chondroitinase ABC Promotes Functional Recovery After Spinal Cord Injury," *Nature*, vol. 416, pp. 636-640 (2002). cited by applicant

Brinkmann, V., et al., "Neutrophil Extracellular Traps Kill Bacteria," *Science*, vol. 303, pp. 1532-

1535, 2004. cited by applicant

Broun, P., et al., "Catalytic plasticity of fatty acid modification enzymes underlying chemical diversity of plant lipids," *Science*, vol. 282, pp. 1315-1317 (1998). cited by applicant

Brown, J. M., et al., "A Sulfated Carbohydrate Epitope Inhibits Axon Regeneration After Injury," *Proc. Natl. Acad. Sci. USA*, vol. 109, pp. 4768-4773 (2012). cited by applicant

Brown, L., et al., "Cardenolide Analogues. 11. Improved Method For The Use Of Fetizon's Reagent In The Synthesis Of Cardiac Glycosides," *Drug Research*, vol. 31, No. 7, pp. 1059-1064 (1981). cited by applicant

Burkart, M, D., et al., "Regeneration of PAPS for the Enzymatic Synthesis of Sulfated Oligosaccharides," *J. Org. Chem.*, vol. 65, pp. 5565-5574 (2000). cited by applicant

Cai, C., et al., "Towards The Chemoenzymatic Synthesis Of Heparan Sulfate Oligosaccharides: Oxidative Cleavage Of P-Nitrophenyl Group With Ceric Ammonium Salts," *Tetra. Lett.*, vol. 54, No. 33, pp. 4471-4474 (2013). cited by applicant

Capila, I., et al., "Heparin—Protein Interactions," *Angew. Chem. Int. Ed.*, vol. 41, pp. 390-412 (2002). cited by applicant

Carfi, A., et al., "Herpes Simplex Virus Glycoprotein D Bound to the Human Receptor HveA," *Molecular Cell*, vol. 8, pp. 169-179 (Jul. 2001). cited by applicant

Cassinelli, G., et al., "Old And New Applications Of Non-Anticoagulant Heparin," *International Journal of Cardiology*, 212S1 pp. S14-S21 (2016). cited by applicant

Casu, B., et al., "Heparin-Like Compounds Prepared By Chemical Modification Of Capsular Polysaccharide From *E. coli* K5," *Carbohydrate Research* vol. 263, pp. 271-284 (1994). cited by applicant

Chan, S., et al., "Regulation Of Pfemp1-VAR2CSA Translation By A Plasmodium Translation-Enhancing Factor," *Nature Microbiology*, vol. 2, Article No. 17068 (May 8, 2017). cited by applicant

Chen, G, Y., et al., "Sterile Inflammation: Sensing And Reacting To Damage," *Nat. Immunol.* vol. 10, pp. 826-837 (2010). cited by applicant

Chen, J., et al., "Using an Enzymatic Combinatorial Approach to Identify Anticoagulant Heparan Sulfate Structures," *Chemistry and Biology*, *Current Biology*, London, GB, vol. 14., No. 9, pp. 986-993 (Sep. 19, 2007). cited by applicant

Chen, M., et al., "Determination of the Substrate Specificities of N-Acetyl-D-glucosaminyltransferase," *Biochemistry*, vol. 45, pp. 12358-12365, 2006. cited by applicant

Chen, R., et al., "Release And Activity Of Histone In Disease," *Cell Death and Disease*, vol. 5, No. 8, e1370, Aug. 14, 2014. cited by applicant

Chen., et al., *Towards De Novo Synthesis of Structure-Defined Oligosaccharides with Heparan Sulfate Biosynthetic Enzymes*, PhD dissertation. 1-167, (Date Created: Aug. 2008; Date Deposited: Oct. 11, 2010.). cited by applicant

Chica R.A., et al., "Semi-rational Approaches to Engineering Enzyme Activity: Combining the Benefits of Directed Evolution and Rational Design," *Current Opinion in Biotechnology*, vol. 16, No. 4, Aug. 2005, pp. 378-384. cited by applicant

Clark, S. R., et al., "Platelet TLR4 activates neutrophil extracellular traps to ensnare bacteria in septic blood," *Nat Med.*, vol. 13, No. 4, pp. 463-469, 2007. cited by applicant

Cole, C.L., et al., "Synthetic Heparan Sulfate Oligosaccharides Inhibit Endothelial Cell Functions Essential for Angiogenesis" *Plos One*, vol. 5, issue 7, pp. 1-15. (Year: 2010). cited by applicant

Communication of European publication number and information on the application of Article 67(3) EPC corresponding to European Application No. 14812890.3 dated Mar. 31, 2016. cited by applicant

Communication of European publication number corresponding to European Patent application No. 20887629.2 dated Jul. 20, 2022. cited by applicant

Communication of the extended European search report corresponding to European Application

No. 14812890.3 dated Dec. 21, 2016. cited by applicant

Communication of the extended European Search report corresponding to European Patent Application No. 20887629.2 dated Oct. 27, 2023. cited by applicant

Communication pursuant to Article 94(3) EPC Corresponding to European Patent Application No. 19822610.2-1109 dated Sep. 30, 2024, pp. 5. cited by applicant

Communication under Rule 71(3) EPC (Intention to Grant) corresponding to European Patent Application No. 18873131.9-1109 dated Oct. 16, 2024, 7 Pages. cited by applicant

Conrad, H, E., "Heparin-Binding Proteins," J. of Medicinal Chemistry, vol. 42, No. 4, pp. 777-778 (1998). cited by applicant

Copeland, R., et al., "Using a 3-O-Sulfated Heparin Octasaccharide to Inhibit the Entry of Herpes Simplex Virus Type 1," Biochemistry, vol. 47, pp. 5774-5783 (2008). cited by applicant

Corrected Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated Sep. 20, 2022. cited by applicant

Coutant, C., et al., "2-Deoxy-2-Trichloroacetamido-D-Glucopyranose Derivatives In Oligosaccharide Synthesis: From Hyaluronic Acid To Chondroitin 4-Sulfate Trisaccharides" J Chem Soc Perkin Trans 1, pp. 1573-1581 (Year: 1995). cited by applicant

Crowther, M, A., et al., "Mechanisms Responsible For The Failure Of Protamine To Inactivate Low-Molecular-Weight Heparin," British Journal of Hematology, vol. 116, pp. 178-186 (2002). cited by applicant

Darden, T., et al., "Particle Mesh Ewald: An N·Log(N) Method For Ewald Sums In Large Systems," J. Chem. Phys. 1993, vol. 98, No. 12, pp. 10089-10092. cited by applicant

Das, S.K., et al., "Synthesis of Conformationally Locked I-Iduronic Acid Derivatives: Direct Evidence for a Critical Role of the Skew-Boat 280 Conformer in the Activation of Antithrombin by Heparin," Chem. Eur. J., vol. 7, No. 22, pp. 4821-4834 (2001). cited by applicant

Davenport, A., "Review Article: Low-Molecular-Weight Heparin As An Alternative Anticoagulant To Unfractionated Heparin For Routine Outpatient Haemodialysis Treatments," Nephrology, vol. 14, pp. 455-461 (2009). cited by applicant

De Paz, J.L., et al., "Microarrays of Synthetic Heparin Oligosaccharides," Journal of the American Chemical Society, vol. 128, No. 9, Mar. 8, 2006, pp. 2766-2767. cited by applicant

Deagostini, A.I., et al., "Human Follicular Fluid Heparan Sulfate Contains Abundant 3-O-Sulfated Chains with Anticoagulant Activity," J. Biol. Chem., vol. 283, pp. 28115-28124, Oct. 17, 2008. cited by applicant

Decision to Grant corresponding to European Patent Application No. 14812890.3 dated Apr. 14, 2022. cited by applicant

Decision to Grant corresponding to Japanese Patent Application No. 2016521505 dated Feb. 3, 2020. cited by applicant

Decision to Grant corresponding to Japanese Patent Application No. 2019549419 dated Jul. 11, 2023. cited by applicant

Devos D., et al., "Practical Limits of Function Prediction," Proteins: Structure, Function, and Genetics, vol. 41, No. 1, Oct. 1, 2000, pp. 98-107. cited by applicant

Dooley, T.P., "Cloning Of The Human Phenol Sulfotransferase Gene Family: Three Genes Implicated In The Metabolism Of Catecholamines, Thyroid Hormones And Drugs," Chemico-Biological Interactions, vol. 109, pp. 29-41 (1998). cited by applicant

Dou, W., et al., "Role of Deacetylase Activity of N-Deacetylase/N-Sulfotransferase 1 in Forming N-Sulfated Domain in Heparan Sulfate", The Journal of Biological Chemistry, vol. 290, No. 33, pp. 20427-20437 (Aug. 14. 2015). cited by applicant

Edens, R.E., et al., "Gradient Polyacrylamide Gel Electrophoresis for Determination of Molecular Weights of Heparin Preparations and Low-Molecular-Weight Heparin Derivatives," J. Pharm. Sci., vol. 81, No. 8, pp. 823-827 (Aug. 1992). cited by applicant

Eller, S., et al., "Automated Solid-Phase Synthesis of Chondroitin Sulfate Glycosaminoglycans,"

Angew. Chem. Int. Ed., vol. 52, pp. 5858-5861 (2013). cited by applicant

Esko, J.D., et al., "Molecular diversity of heparan sulfate," J. Clin. Invest., vol. 108, pp. 169-173 (2001). cited by applicant

European Search Report corresponding to European Patent Application No. 18764628.6 dated Dec. 2, 2020. cited by applicant

European Search Report corresponding to European Patent Application No. 18873131.9 dated Jul. 12, 2021. cited by applicant

Extended European Search Report Corresponding to European Patent Application No. 19822610.2 dated Mar. 29, 2022. cited by applicant

Extended European Search Report Corresponding to European Patent Application No. 06770377.7 dated Nov. 22, 2010. cited by applicant

Falany, C.N., "Introduction: Changing view of sulfation and the cytosolic Sulfotransferases," vol. 11, The FASEB Journal, pp. 1-2 (Jan. 1997). cited by applicant

Feltracco, P., et al., "Perioperative thrombotic complications in liver transplantation." World J. Gastroenterol., vol. 21, pp. 8004-8013 (2015). cited by applicant

Feng, S., et al., "Characteristics Associated with Liver Graft Failure: The Concept of a Donor Risk Index." Am. J. Transplant., vol. 6, pp. 783-790 (2006). cited by applicant

Feyerabend, T.B., et al., "Heparan sulfate C5-epimerase is essential for heparin biosynthesis in mast cells," Nat. Chem. Biol., vol. 2, No. 4, pp. 195-196 (Apr. 2006). cited by applicant

Fiser, A., et al., "Modeller: Generation and Refinement of Homology-Based Protein Structure Models," Methods Enzymol, vol. 374, pp. 461-491, 2003. cited by applicant

Frank, R.D., et al., "A non-anticoagulant synthetic pentasaccharide reduces inflammation in a murine model of kidney ischemia-reperfusion injury," Thromb Haemost, vol. 96, pp. 802-806. (Dec. 2006). cited by applicant

Freeman, C.G., et al., "The accumulation of circulating histones on heparan sulphate in the capillary glycocalyx of the lungs." Biomater., vol. 34, pp. 5670-5676 (2013). cited by applicant

Fried, M., et al., "Designing a VAR2CSA-based vaccine to prevent placental malaria." Vaccine, vol. 33, pp. 7483-7488 (2015). cited by applicant

Fukuta, M., et al., "Molecular cloning and expression of human chondroitin 6-sulfotransferase," Biochimica et Biophysica Acta, vol. 1399, pp. 57-61 (1998). cited by applicant

Fuster, J. J., et al., "The sweet and sour of cancer: glycans as novel therapeutic targets," Nat. Rev. Cancer, vol. 5, No. 7, pp. 1-27 (Jul. 2005). cited by applicant

Gama, C.I., et al., "Sulfation patterns of glycosaminoglycans encode molecular recognition and activity," Nat Chem. Biol., vol. 2, No. 9, pp. 467-473 (Sep. 2006). cited by applicant

Ganey, P.E., et al. "Role of the Coagulation System in Acetaminophen-Induced Hepatotoxicity in Mice." Hepatology, vol. 46(4), pp. 1177-1186 (2007). cited by applicant

Goddard-Borger, E, D., et al., "An Efficient, Inexpensive and Shelf-Stable Diazotransfer Reagent: Imidazole-1-sulfonyl Azide Hydrochloride." Org. Lett., vol. 9, pp. 3797-3800 (2007). cited by applicant

Guerrini, M., et al., "An unusual antithrombin-binding heparin octasaccharide with an additional 3-O-sulfated glucosamine in the active pentasaccharide sequence," Biochem. J., vol. 449, pp. 343-351, 2013. cited by applicant

Guerrini, M., et al., "Antithrombin-binding oligosaccharides: structural diversities in a unique function?," Glycoconj. J., vol. 31, 409, pp. 9, Aug. 2014. cited by applicant

Guerrini, M., et al., "Oversulfated chondroitin sulfate is a contaminant in heparin associated with adverse clinical events," Nat. Biotechnol., vol. 26, No. 6, pp. 669-675 (Jun. 2008). cited by applicant

Habuchi, H., et al., "Molecular Characterization and Expression of Heparan-sulfate 6-Sulfotransferase—Complete cDNA Cloning in Human and Partial Cloning in Chinese Hamster Ovary Cells," The Journal of Biological Chemistry. vol. 273, No. 15, pp. 9208-9213 (Apr. 10,

1998). cited by applicant

Habuchi, O., et al., "Purification of Chondroitin 6-Sulfotransferase Secreted from Cultured Chick Embryo Chondrocytes," *The Journal of Biological Chemistry*, vol. 268(29), pp. 21968-21974 (1993). cited by applicant

Hagiwara, S., et al., "Danaparoid sodium inhibits systemic inflammation and prevents endotoxin-induced acute lung Injury in rats," *Critical Care*, vol. 12, No. 2, report No. R43, Apr. 2, 2008. cited by applicant

Orgaran, Product Monograph—HIT for danaparoid sodium, pp. 41, Feb. 9, 2018. (Year: 2018). cited by applicant

Hajmohammadi, S., "Normal levels of anticoagulant heparan sulfate are not essential for normal hemostasis," *The Journal of Clinical Investigation*, vol. 111, pp. 989-999, No. 7, Apr. 2003. cited by applicant

Hansen, S. U., et al., "First gram-scale synthesis of a heparin-related dodecasaccharide," *Org. Lett.*, vol. 15, No. 1, pp. 88-91, Jan. 4, 2013. cited by applicant

Hansen, S.U., "Tetrasaccharide iteration synthesis of a heparin-like dodecasaccharide and radiolabelling for in vivo tissue distribution studies," *Nature Communications*, vol. 4, Article No. 2016, pp. 1-9, May 16, 2013. cited by applicant

Harada, N., et al., "Danaparoid sodium reduces ischemia/reperfusion-induced liver injury in rats by attenuating inflammatory responses" *Thromb Haemost*, 2007; 97; pp. 81-87. cited by applicant

Harada, N., et al., "Dalteparin, a low molecular weight heparin, attenuates inflammatory responses and reduces ischemia-reperfusion-induced liver injury in rats." *Crit. Care Med.*, vol. 34, Article No. 8, (2006). cited by applicant

Harder, S., "Renal profiles of anticoagulants," *J. Clin. Pharmacol.*, vol. 52, No. 7, pp. 964-975, Jul. 2012. cited by applicant

Harris, E. N., "The human hyaluronan receptor for endocytosis (HARE/Stabilin-2) is a systemic clearance receptor for heparin," *J. Biol. Chem.*, vol. 283, No. 25, pp. 17341-17350, Jun. 20, 2008. cited by applicant

Harris, E.N., et al., "Endocytic Function, Glycosaminoglycan Specificity, and Antibody Sensitivity of the Recombinant Human 190-kDa Hyaluronan Receptor for Endocytosis (HARE)," *J. Biol. Chem.*, vol. 279, No. 35, pp. 36201-36209 (Aug. 27, 2004). cited by applicant

Heard, K.J., "Acetylcystein for acetaminophen poisoning," *N. Eng. J. Med.* vol. 359, pp. 285-292 (2008). cited by applicant

Hirsch, J., et al., "Beyond Unfractionated Heparin and Warfarin Current and Future Advances," *Circulation*, vol. 116, pp. 552-560 (2007). cited by applicant

Hirsch, J., et al., "Heparin and Low-Molecular-Weight Heparin The Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy," *Chest*, vol. 126, pp. 188S-203S (2004). cited by applicant

Hsieh, P-H., "Uncovering the Relationship between Sulphation Patterns and Conformation of Iduronic Acid in Heparan Sulphate," *J. Sci Rep*, vol. 6, Article No. 29602, pp. 1-8, 2016. cited by applicant

Hsieh, P-H., et al., "Chemoenzymatic synthesis and structural characterization of 2-O-sulfated glucuronic acid containing heparan sulfate hexasaccharides." *Glycobiology* vol. 24, pp. 681-692 (2014). cited by applicant

Hu, Y-P., et al., "Synthesis of 3-O-sulfonated heparan sulfate octasaccharides that inhibit the herpes simplex virus type 1 host-cell interaction," *Nat Chem*, vol. 3, pp. 557-563, Jul. 2011. cited by applicant

Huang, C. C., et al., "Enhancing UCSF Chimera through web services," *Nucleic Acids Res.*, vol. 42, pp. W478- W484, May 26, 2014. cited by applicant

Huebener, P., el al., "The HMGB1/RAGE axis triggers neutrophil-mediated injury amplification following necrosis." *J. Clin. Invest.*, vol. 125, pp. 539-550 (2015). cited by applicant

Humphrey, W., "VMD: Visual Molecular Dynamics," J. Mol. Graph., vol. 14, pp. 33-38, 1996. cited by applicant

Iba et al., "Danaparoid sodium attenuates the increase in inflammatory cytokines and preserves organ function in endotoxemic rats," Critical Care, vol. 12, Article No. R86 (7 pages) (2008). cited by applicant

Iba, T., et al., "Advance in the management of sepsis-induced coagulopathy and disseminated intravascular coagulation." J. Clin. Med., vol. 8, Article No. 728 (16 pages) (2019). cited by applicant

Intention to Grant corresponding to European Patent Application No. 11849994.6 dated Apr. 7, 2021. cited by applicant

Intention to Grant corresponding to European Patent Application No. 11849994.6 dated Sep. 1, 2021. cited by applicant

Intention to Grant corresponding to European Patent Application No. 14812890.3 dated Oct. 27, 2021. cited by applicant

Intention to Grant corresponding to European Patent Application No. 14812890.3 dated Mar. 16, 2022. cited by applicant

International Preliminary Examination Report for corresponding PCT Application No. PCT/US2008/008945 dated Jan. 26, 2010. cited by applicant

International Preliminary Report on Patentability corresponding to International Application No. PCT/US2018/040774 dated Jan. 7, 2020. cited by applicant

International Preliminary Report on Patentability corresponding to International Application No. PCT/US2019/037993 dated Dec. 22, 2020. cited by applicant

International Preliminary Report on Patentability Corresponding to International Patent Application No. PCT/US2020/060581 dated May 17, 2022. cited by applicant

International Preliminary Report on Patentability corresponding to International Application No. PCT/US2014/042683 dated Dec. 22, 2015. cited by applicant

International Preliminary Report on Patentability corresponding to International Application No. PCT/US2011/066843 dated Jun. 25, 2013. cited by applicant

International Preliminary Report on Patentability Corresponding to International application No. PCT/US2018/021986 dated Sep. 10, 2019. cited by applicant

International Preliminary Report on Patentability corresponding to international application No. PCT/US2018/059152 dated May 5, 2020. cited by applicant

International Search Report and the Written Opinion of the International Searching Authority Corresponding to International Application No. PCT/US 2018/040774 dated Sep. 18, 2018. cited by applicant

International Search Report and the Written Opinion of the International Searching Authority corresponding to international application No. PCT/US 2018/059152 dated Mar. 6, 2019. cited by applicant

International Search Report and Written Opinion of the International Searching Authority Corresponding to International Patent Application No. PCT/US2020/060581 dated Feb. 11, 2021. cited by applicant

International Search Report corresponding to International Application No. PCT/US2019/037993 dated Oct. 18, 2019. cited by applicant

International Search Report corresponding to International Application No. PCT/US2014/042683 dated Oct. 9, 2014. cited by applicant

International Search Report Corresponding to International application No. PCT/US 2018/021986 dated Aug. 1, 2018. cited by applicant

Jackson, S.P., et al., "Thromboinflammation: challenges of therapeutically targeting coagulation and other host defense mechanisms." Blood, vol. 133, pp. 906-918 (2019). cited by applicant

Jaeschke, H., et al., "Complement activates Kupffer cells and neutrophils during reperfusion after

hepatic ischemia.” Am. J. Physiol-Gastroint. Liver Physiol., vol. 264, pp. G801-G809 (1993). cited by applicant

Jaimes, F., et al., “Unfractionated heparin for treatment of sepsis: A randomized clinical trial (The HETRASE Study).” Crit. Care Med., vol. 37, pp. 1185-1196 (2009). cited by applicant

Jemth, P., et al., “Oligosaccharide library-based assessment of heparan sulfate 6-O-sulfotransferase substrate specificity,” Journal of Biological Chemistry, vol. 278, No. 27, pp. 24371-24376 (Jul. 4, 2003). cited by applicant

Jin, L., et al., “The anticoagulant activation of antithrombin by heparin,” Proc. Natl. Acad. Sci., vol. 94, pp. 14683-14688, Dec. 1997. cited by applicant

Kahn, S. R. et al., “Prevention of VTE in nonsurgical patients: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines,” Chest 141 (2 suppl), e195S-e226S Feb. 2012. cited by applicant

Kakkar, A. K., et al., “Low Molecular Weight Heparin, Therapy With Dalteparin, and Survival in Advanced Cancer: The Fragmin Advanced Malignancy Outcome Study (Famous),” J. Clin. Oncol., vol. 22, No. 10, pp. 1944-1948 (May 15, 2004). cited by applicant

Kakuta, Y., et al., “Heparan sulphate N-sulphotransferase activity: reaction mechanism and substrate recognition,” Biochem. Soc. Trans., vol. 31 (pt2), pp. 331-334 (2003). cited by applicant

Kamimura, K., et al., “Regulation of Notch signaling by *Drosophila* heparan sulfate 3-O sulfotransferase,” J. Cell Biol, vol. 166, No. 7, pp. 1069-1079, Sep. 27, 2004. cited by applicant

Kaneko, J., et al., “Coagulation and fibrinolytic profiles and appropriate use of heparin after living-donor liver transplantation.” Clin. Transplant., vol. 19, pp. 804-809 (2005). cited by applicant

Kirschner, K. N., et al., “GLYCAM06: A Generalized Biomolecular Force Field. Carbohydrates,” J. Comput. Chem., vol. 29, No. 4, pp. 622-655, Mar. 2008. cited by applicant

Kishimoto, T. K. et al., “Contaminated heparin associated with adverse clinical events and activation of the contact system,” N. Engl. J. Med., vol. 358, No. 23, pp. 2457-2467, Jun. 5, 2008. cited by applicant

Kisselev, L., “Polypeptide release factors in prokaryotes and eukaryotes: same function, different structure,” Structure, vol. 10, pp. 8-9 (2002). cited by applicant

Kobayashi, M., et al., “Purification and characterization of heparan sulfate 2-sulfotransferase from cultured Chinese hamster ovary cells,” J Biol. Chem. vol. 271, No. 13, pp. 7645-7653, Mar. 29, 1996. cited by applicant

Kollman, P., A., et al., “Calculating Structures and Free Energies of Complex Molecules: Combining Molecular Mechanics and Continuum Models,” Acc. Chem. Res., vol. 33, pp. 889-897, 2000. cited by applicant

Konishi, T., et al., “Hepatic ischemia/reperfusion: mechanisms of tissue injury, repair, and regeneration.” Gene Expr., vol. 17, pp. 277-287 (2017). cited by applicant

Kopec, A. K., et al., “Fibrin(ogen) drives repair after acetaminophen-induced liver injury via leukocyte αMb2 integrin-dependent upregulation of Mmp12.” J. Hepatol., vol. 66, pp. 787-797 (2017). cited by applicant

Kreimann, M., et al., “Binding of anti-platelet factor 4/heparin antibodies depends on the thermodynamics of conformational changes in platelet factor 4,” Blood, vol. 124, No. 15, pp. 2442-2449, Oct. 9, 2014. cited by applicant

Kreuger, J., et al., “Interactions between heparan sulfate and proteins: the concept of specificity,” J. Cell Biol., vol. 174, No. 3, pp. 323-327 (Jul. 31, 2006). cited by applicant

Kuberan, B., et al., “Enzymatic synthesis of antithrombin III-binding heparan sulfate pentasaccharide,” Nature Biotechnology, vol. 21, No. 11, 1343-1346 (Nov. 2003). cited by applicant

Kuberan, B., et al., The Journal of Biological Chemistry, “Chemoenzymatic Synthesis of Classic and Non-classical Anticoagulant Heparan Sulfate Polysaccharides”, 2003, vol. 278, No. 52, pp. 52613-52621 (Year: 2003). cited by applicant

Kubes, et al., "Sterile inflammation in the liver." *Gastroenterology*, vol. 143, pp. 1158-1172 (2012). cited by applicant

Langdown, J.; Belzar, K. J.; Savory, W. J., Baglin, T. P.; Huntington, J. A. J. *Mol. Biol.* 2009, 386, 1278. cited by applicant

Laremore, T. et al. "Ionic liquid matrix for direct UV-MALDI-TOF-MS Analysis of Dermatan Sulfate and Chondroitin Sulfate Oligosaccharides." *Anal. Chem.*, vol. 79, pp. 1604-1610. (Year: 2007). cited by applicant

Laurent et al., "The Molecular-Weight-Dependence of the Anti-Coagulant Activity of Heparin," *Biochem. J.*, vol. 175, pp. 691-701 (1978). cited by applicant

Ledin et al., "Heparan Sulfate Structure in Mice with Genetically Modified Heparan Sulfate Production," *J. Biol. Chem.*, vol. 279, No. 41, pp. 42732-42741 (2004). cited by applicant

Lee, "Acetaminophen toxicity: changing perceptions on a social/medical issue." *Hepatology*, vol. 46, pp. 966-970 (2007). cited by applicant

Lee, M.K., and Lander, A.D., (1991) *Proc. Natl. Acad. Sci. USA* 88, 2768-2772. cited by applicant

Li et al., "Biosynthesis of Heparin/Heparan Sulfate cDNA Cloning and Expression of D-Glucuronyl C5-Epimerase From Bovine Lung," *J. Biol. Chem.*, vol. 272, No. 4, pp. 28158-28163 (Oct. 31, 1997). cited by applicant

Li et al., "Enzymatic Synthesis of Homogeneous Chondroitin Sulfate Oligosaccharides." *Angew. Chemie.*, vol. 129(39), pp. 11946-11949 (2017). cited by applicant

Li et al., "Enzymatic synthesis of homogenous chondroitin sulfate e oligosaccharides," *Abstract of Glycobiol.*, vol. 28(12) (2018) [Abstract]. cited by applicant

Li J, Su W, and Liu J. "Enzymatic synthesis of homogeneous chondroitin sulfate oligosaccharides." *Angew Chem Int Ed.* 2017; 56:11784-7. cited by applicant

Li, J., et al., "Enzymatic Synthesis of Chondroitin Sulfate E to Attenuate Bacteria Lipopolysaccharide-induced Organ Damage," *ACS Central Science*, vol. 6, No. 7, pp. 1199-1207, Jul. 1, 2020. cited by applicant

Li, L., et al., "Top-down approach for the direct characterization of low molecular weight heparins using LC-FR-MS," *Anal. Chem.*, vol. 84, No. 20, pp. 8822-8829, Oct. 16, 2012. cited by applicant

Liliensiek et al., "Receptor for advanced glycation end products (RAGE) regulates sepsis but not the adaptive immune response." *J. Clin. Invest.* vol. 113, pp. 1641-1650 (2004). cited by applicant

Lindahl et al., "Regulated Diversity of Heparan Sulfate," *The Journal of Biological Chemistry*, vol. 273, No. 39, pp. 24979-24982 (Sep. 25, 1998). cited by applicant

Lindahl, U.; Backstrom, G.; Thunberg, L.; Leder, I. G. *Proc. Natl. Acad. Sci.* 1980, 77, pp. 6551-6555. cited by applicant

Linhardt et al., "Production and Chemical Processing of Low Molecular Weight Heparins," *Seminars in Thrombosis and Hemostasis*, vol. 25, Suppl.3, pp. 5-16 (1999). cited by applicant

Linhardt, R. J., et al., "Synthetic heparin," *Curr. Opin. Pharmacol*, vol. 12, No. 2, pp. 217-219, Apr. 2012. cited by applicant

Liu et al., "Anticoagulant heparan sulfate: structural specificity and biosynthesis," *Appl Microbiol Biotechnol.*, vol. 74, pp. 263-272 (2007). cited by applicant

Liu et al., "Cell Surface Heparan Sulfate and Its Roles in Assisting Viral Infections," *Medicinal Research Reviews*, vol. 22, No. 1, pp. 1-25 (2002). cited by applicant

Liu et al., "Characterization of a Heparan Sulfate Octasaccharide that Binds to Herpes Simplex Virus Type 1 Glycoprotein D," *The Journal of Biological Chemistry*, vol. 277, No. 36, pp. 33456-33467 (Sep. 6, 2002). cited by applicant

Liu et al., "Expression of Heparan Sulfate D-Glucosaminyl 3-O-Sulfotransferase Isoforms Reveals Novel Substrate Specificities," *The Journal of Biological Chemistry*, vol. 274, No. 8, pp. 5185-5192 (Feb. 19, 1999). cited by applicant

Liu et al., "Heparan Sulfate D-Glucosaminyl 3-O-Sulfotransferase-3A Sulfates N-Unsubstituted Glucosamine Residues," *The Journal of Biological Chemistry* vol. 274, No. 53, pp. 38155-38162

(Dec. 31, 1999). cited by applicant

Liu et al., "Purification of Heparan Sulfate D-Glucosaminyl 3-O-Sulfotransferase," *The Journal of Biological Chemistry*, vol. 271, No. 43, pp. 27072-27082 (Oct. 25, 1996). cited by applicant

Liu et al., *Chemoenzymatic Design of Heparan Sulfate Oligosaccharides*, *J Biol Chem*, vol. 285, No. 44, pp. 34240-34249 (Oct. 29, 2010). cited by applicant

Liu et al., "Enzymatic Placement of 6-O-Sulfo Groups in Heparan Sulfate," *Biochemistry* 2011, 50, 4382-4391. cited by applicant

Liu et al., "Lessons learned from the contamination of heparin," *Nat. Prod. Rep.*, vol. 26, pp. 313-321 (2009). cited by applicant

Liu, J. et al., "Chemoenzymatic synthesis of heparan sulfate and heparin", *Royal Society of Chemistry*, vol. 31, pp. 1676-1685 (Year: 2014). cited by applicant

Loganathan et al., "Structural Variation in the Antithrombin III Binding Site Region and Its Occurrence in Heparin from Different Sources," *Biochemistry*, vol. 29, pp. 4362-4368 (1990). cited by applicant

Lopin et al., "From Polymer to Size-Defined Oligomers: An Expeditious Route for the Preparation of Chondroitin Oligosaccharides." *Angew. Chem. Int. Ed.*, vol. 45, pp. 2574-2578 (2006). cited by applicant

Lopin-Bon et al., "Stereocontrolled preparation of biotinylated chondroitin sulfate E di-, tetra-, and hexasaccharide conjugates." *Carbohydr. Res.*, vol. 402, pp. 35-43 (2015). cited by applicant

Lu et al., "Innate Immune Regulations and Liver Ischemia-Reperfusion Injury." *Trasplantation*, vol. 100, pp. 2601-2610 (2016). cited by applicant

Lundbäck et al., "A novel high mobility group box 1 neutralizing chimeric antibody attenuates drug-induced liver injury and postinjury inflammation in mice." *Hepatology* vol. 64, pp. 1699-1710 (2016). cited by applicant

Ly et al., "The proteoglycan bikunin has a defined sequence." *Nat. Chem. Biol.*, vol. 7, pp. 827-833 (2011). cited by applicant

Macchione et al., "Synthesis of chondroitin sulfate oligosaccharides using N-tetrachlorophthaloyl and N-trifluoroacetyl galactosamine building blocks," *European Journal of Organic Chemistry*, pp. 3868-3884 (2014). cited by applicant

Mackman, "Triggers, targets and treatments for thrombosis," *Nature*, vol. 451, No. 21, pp. 914-918 (Feb. 21, 2008). cited by applicant

Mahe, I.; Chidac, J.; Helfer, H.; Nobel, S. J. *Thromb. Haemost.* 2016, 14, 2017 cited by applicant

Man et al., "Tolerance of the liver to intermittent pringle maneuver in hepatectomy for liver tumors." *JAMA Surgery*, vol. 134, pp. 533-539 (1999). cited by applicant

Marcum, J. A. et al., "The origin of the dispute over the discovery of heparin," *J. Hist. Med. Allied Sci.*, vol. 55, pp. No. 1, 37-66, Jan. 1, 2000. cited by applicant

Martinez-Gonzalez et al., "New Challenges for a Second-Generation Low-Molecular-Weight Heparin: Focus on Bemiparin," *Expert Rev. Cardiovasc. Ther.*, vol. 8, No. 5, pp. 625-634 (2010). cited by applicant

Maza, S., et al., "Synthesis of chondroitin/dermatan sulfate-like oligosaccharides and evaluation of their protein affinity by fluorescence polarization." *Org. Biomol. Chem.*, vol. 11, pp. 3510-3525 (2013). cited by applicant

Mazany et al., "Human chondroitin 6-sulfotransferase: cloning, gene structure, and chromosomal localization," *Biochimica et Biophysica Acta*, vol. 1407, pp. 92-97 (1998). cited by applicant

McGowan, K. E.; Makari, J.; Diamantouros, A.; Bucci, C.; Rempel, P.; Selby, R.; Geerts, W. *Blood* 2016, 127, 1954. cited by applicant

Miyachi et al., "Syntheses of chondroitin sulfate tetrasaccharide structures containing 4,6-disulfate patterns and analysis of their interaction with glycosaminoglycan-binding protein." *Bioorg. Med. Chem. Lett.*, vol. 25, pp. 1552-1555 (2015). cited by applicant

Miyata et al., "Persistent cortical plasticity by upregulation of chondroitin 6-sulfation." *Nat. Neurosci.*, vol. 15, pp. 414-422 (2012). cited by applicant

Mizumoto et al., "Molecular interactions between chondroitin-dermatan sulfate and growth factors/receptors/matrix proteins." *Curr. Opin. Struct. Biol.*, vol. 34, pp. 35-42 (2015). cited by applicant

Monneau et al., "The sweet spot: how GAGs help chemokines guide migrating cells." *J. Leukoc. Biol.* vol. 99, pp. 935-953 (2016). cited by applicant

Moon et al., "Dissecting the substrate recognition of 3-O-sulfotransferase for the biosynthesis of anticoagulant heparin," *Proceedings of the National Academy of Sciences*, vol. 109, No. 14, pp. 5265-5270 (2012). cited by applicant

Mossanen et al., "Acetaminophen-induced acute liver injury in mice." *Lab. Anim.* vol. 49, pp. 30-36 (2015). cited by applicant

Mousa, "Drug Discovery and Evaluation: Pharmacological Assays" (ed. Vogel, H.), 393-456 (Springer-Verlag Berlin, Heidelberg, New York (2008). cited by applicant

Mousa, "Heparin and Low-Molecular Weight Heparins in Thrombosis and Beyond," *Meth. Mol. Biol.*, vol. 663, pp. 109-132 (2010). cited by applicant

Mousa, "In Vitro Methods of Evaluating Antithrombotics and Thrombolytics," *Meth. Mol. Biol.*, vol. 663, pp. 1-28 (2010). cited by applicant

Munoz E., et al., "Affinity, kinetic, and structural study of the interaction of 3-O-sulfotransferase isoform 1 with Heparan Sulfate", *Biochemistry*, vol. 45, No. 16, 2006, pp. 5122-5128. cited by applicant

Munoz et al., "Enzymatic synthesis of heparin related polysaccharides on sensor chips: Rapid screening of heparin-protein interactions." *Biochemical and Biophysical Research Communications*, Academic Press Inc., Orlando, FL, US, vol. 339, No. 2, pp. 597-602 (Jan. 13, 2006). cited by applicant

Nadanaka et al., "Characteristic Hexasaccharide Sequences in Octasaccharides Derived from Shark Cartilage Chondroitin Sulfate D with a Neurite Outgrowth Promoting Activity," *The Journal of Biological Chemistry*, vol. 273(6), pp. 3296-3307 (1998). cited by applicant

Nagano et al., "Chondroitin sulfate protects vascular endothelial cells from toxicities of extracellular histones." *Eur. J. Pharmacol.*, vol. 826, pp. 48-55 (2018). cited by applicant

Nam et al., "Syndecan-1 Limits the Progression of Liver Injury and Promotes Liver Repair in Acetaminophen-Induced Liver Injury in Mice." *Hepatology*, vol. 66(5), pp. 1601-1615, doi: 10.1002/hep.29265 (2017). cited by applicant

Nastuk et al., "Expression Cloning and Characterization of NSIST, a Novel Sulfotransferase Expressed by a Subset of Neurons and Postsynaptic Targets," *The Journal of Neuroscience*, vol. 18, No. 18, pp. 7167-7177 (Sep. 15. 1998). cited by applicant

Noti et al., "Chemical Approaches to Define the Review Structure-Activity Relationship of Heparin-like Glycosaminoglycans," *Chemistry & Biology*, vol. 12, pp. 731-756 (Jul. 2005). cited by applicant

Notice of Allowance and Interview Summary corresponding to U.S. Appl. No. 14/898,865 dated Dec. 15, 2017. cited by applicant

Notice of Allowance and Interview Summary corresponding to U.S. Appl. No. 17/254,145 dated Jan. 30, 2023. cited by applicant

Notice of Allowance corresponding to U.S. Appl. No. 17/254,145 dated Dec. 8, 2022. cited by applicant

Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated Sep. 12, 2022. cited by applicant

Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated Feb. 10, 2023. cited by applicant

Notice of Allowance corresponding to U.S. Appl. No. 13/996,930 dated Aug. 11, 2021. cited by

applicant
Notice of Allowance corresponding to U.S. Appl. No. 16/761,159 dated Aug. 24, 2023. cited by applicant
Notice of Allowance corresponding to U.S. Appl. No. 16/761,159 dated Dec. 4, 2023. cited by applicant
Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated May 24, 2023. cited by applicant
Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated Sep. 7, 2023. cited by applicant
Notice of Allowance corresponding to U.S. Appl. No. 16/492,858 dated Dec. 21, 2023. cited by applicant
Notice of Publication Corresponding to European Patent Application. No. 19822610.2 dated Mar. 31, 2021. cited by applicant
Notice of Publication Corresponding to European Patent application No. 18764628.6 dated Nov. 20, 2019. cited by applicant
Notice of Publication Corresponding to European Patent Application No. 18873131.9 dated Jul. 15, 2020. cited by applicant
Oduah et al., "Heparin: Past, present, and future." Pharmaceuticals (Basel), vol. 9, Article No. 38 (2016). cited by applicant
Office Action (Annex to the communication) corresponding to European Patent Application No. 18873131.9 dated Apr. 4, 2024, 7 Pages. cited by applicant
Office Action (Decision to grant) corresponding to European Patent Application No. 18873131.9 dated Feb. 13, 2025, 3 Pages. cited by applicant
Office Action (Decision of Rejection) corresponding to Chinese Patent Application No. 20180020095.X dated Dec. 1, 2022, pp. 10 (Translation). cited by applicant
Office Action (Decision to grant) corresponding to European Patent Application No. 11849994.6 dated Jan. 13, 2022. cited by applicant
Office Action (European Search Report) corresponding to European Patent Application No. 11849994.6 dated Apr. 1, 2016. cited by applicant
Office Action (Final Rejection) corresponding to U.S. Appl. No. 13/996,930 dated Jun. 22, 2020. cited by applicant
Office Action (Final) corresponding to U.S. Appl. No. 13/417,641 dated Sep. 5, 2013. cited by applicant
Office Action (Final) corresponding to U.S. Appl. No. 16/625,342 dated Nov. 4, 2022. cited by applicant
Office Action (Non- Final Rejection) corresponding to U.S. Appl. No. 18/138,596 dated Jun. 5, 2024. cited by applicant
Office Action (non-final) corresponding to U.S. Appl. No. 13/417,641 dated Feb. 11, 2013. cited by applicant
Office Action (Notice of Allowance) corresponding to U.S. Appl. No. 13/417,641 dated Feb. 27, 2014. cited by applicant
Office Action (Notice of Allowance) corresponding to U.S. Appl. No. 13/996,930 dated Nov. 24, 2021. cited by applicant
Office Action (Notice of Allowance) corresponding to U.S. Appl. No. 16/625,342 dated Jan. 10, 2024. cited by applicant
Office Action (Notice of Allowance) corresponding to U.S. Appl. No. 16/625,342 dated Jan. 19, 2024. cited by applicant
Office Action (Notice of Allowance) corresponding to U.S. Appl. No. 16/625,342 dated Apr. 26, 2024. cited by applicant
Office Action (Notice of Reasons for Rejection) corresponding to Japanese Patent Application No.

2020-570916 dated Jun. 20, 2023. cited by applicant
Office Action (Notice of Reasons for Rejection) corresponding to Japanese Patent Application No. 2020-570916 dated Jan. 16, 2024. cited by applicant
Office Action (Notice of Reasons for Rejection) corresponding to Japanese Patent Application No. 2023-129964 dated Oct. 8, 2024, 14 pages. cited by applicant
Office Action (Notice of Reasons for Rejection) corresponding to Japanese Patent Application No. 2022-527682 dated Oct. 29, 2024, pp. 8. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 16/625,342 dated Dec. 16, 2021. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 17/254,145 dated Nov. 26, 2021. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 16/761,159 dated Jun. 10, 2022. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 16/492,858 dated Jun. 30, 2021. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 11/920,319 dated Jan. 27, 2010. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 13/417,641 dated Dec. 27, 2012. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 13/996,930 dated Mar. 17, 2015. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 17/744,407 dated Nov. 15, 2024. cited by applicant
Non-Final Office Action corresponding to U.S. Appl. No. 17/744,407 dated Feb. 26, 2025. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 18/222,910 dated Dec. 6, 2024, 9 Pages. cited by applicant
Office Action (Restriction Requirement) corresponding to U.S. Appl. No. 18/222,910 dated Feb. 26, 2025, 11 Pages. cited by applicant
Office Action corresponding to Chinese Application No. 202080092829.2 dated Jun. 21, 2023. cited by applicant
Office Action (Decision of Rejection) corresponding to Chinese Application No. 202080092829.2 dated Mar. 3, 2024, pp. 19. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201880020095.X dated Jun. 8, 2022. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201480044429.9 dated Aug. 30, 2018, pp. 43. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201480044429.9 dated Apr. 9, 2019. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201480044429.9 dated Mar. 3, 2020. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201880020095.X dated Sep. 22, 2021. cited by applicant
Office Action corresponding to Chinese Patent Application No. 2018800850125 dated Jan. 20, 2023. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201880020095.X dated Apr. 18, 2023. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201980044697.3 dated Jul. 7, 2023. cited by applicant
Office Action corresponding to Chinese Patent Application No. 201880085012.5 dated Nov. 30,

2023. cited by applicant
Office Action corresponding to Chinese patent Application No. 202310342719.2 dated May 8, 2023. cited by applicant
Office Action corresponding to Chinese Patent Application No. 202080092 dated Jun. 21, 2023. cited by applicant
Office Action corresponding to Chinese patent Application No. 202310342719.2 dated Feb. 28, 2024. cited by applicant
Office Action corresponding to Chinese patent Application No. 202310342719.2 dated Jul. 24, 2024, 8 Pages. cited by applicant
Office Action corresponding to Chinese patent Application No. 202310342719.2 dated Oct. 16, 2024, pp. 11. cited by applicant
Office Action corresponding to European Patent Application No. 11849994.6 dated May 24, 2018. cited by applicant
Office Action corresponding to European Patent Application No. 11849994.6 dated Jan. 22, 2020. cited by applicant
Office Action corresponding to European Patent Application No. 14812890.3 dated Jun. 23, 2020. cited by applicant
Office Action corresponding to European Patent Application No. 18873131.9-1112 dated Aug. 14, 2023. cited by applicant
Office Action corresponding to European Patent Application No. 18764628.6 dated Nov. 16, 2023. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2019-549419 dated Dec. 23, 2022, pp. 3. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2016-521505 dated Jul. 19, 2018. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2016-521505 dated Jun. 21, 2019. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2019-549419 dated Mar. 22, 2022. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2020-544568 dated Jan. 10, 2023. cited by applicant
Office Action corresponding to Japanese Patent Application No. 2024-008526 dated Dec. 24, 2024, pp. 4—English summary provided. cited by applicant
Office Action Corresponding to Japanese Patent Application Serial No. 2020-544568 dated Sep. 26, 2023. cited by applicant
Office Action corresponding to U.S. Appl. No. 12/178,434 dated Oct. 28, 2011. cited by applicant
Office Action corresponding to U.S. Appl. No. 12/178,434 dated Apr. 19, 2011. cited by applicant
Office Action corresponding to U.S. Appl. No. 12/178,434 dated Jan. 26, 2011. cited by applicant
Office Action corresponding to U.S. Appl. No. 16/492,858 dated Jan. 13, 2022. cited by applicant
Office Action corresponding to U.S. Appl. No. 16/625,342 dated Mar. 21, 2022. cited by applicant
Office Action corresponding to U.S. Appl. No. 16/761,159 dated Jan. 11, 2023. cited by applicant
Office Action corresponding to U.S. Appl. No. 17/254,145 dated Feb. 16, 2022. cited by applicant
Office Action corresponding to U.S. Appl. No. 13/996,930 dated Oct. 8, 2015. cited by applicant
Office Action corresponding to U.S. Appl. No. 13/996,930 dated May 26, 2016. cited by applicant
Office Action corresponding to U.S. Appl. No. 13/996,930 dated Dec. 21, 2017. cited by applicant
Office Action corresponding to U.S. Appl. No. 13/996,930 dated Jul. 30, 2018. cited by applicant
Office Action corresponding to U.S. Appl. No. 13/996,930 dated Nov. 22, 2019. cited by applicant
Office Action corresponding to U.S. Appl. No. 16/625,342 dated Jun. 15, 2023. cited by applicant
Official Action (Final) corresponding to U.S. Appl. No. 11/920,319 dated Dec. 16, 2010. cited by applicant

Official Action corresponding to U.S. Appl. No. 14/898,865 dated Mar. 23, 2017. cited by applicant

Oliveira et al., "Neutrophils: a cornerstone of liver ischemia and reperfusion injury." Lab. Invest., vol. 98, pp. 51-62 (2018). cited by applicant

Ong et al., "Expression Cloning of a Human Sulfotransferase that Directs the Synthesis of the HNK-1 Glycan on the Neural Cell Adhesion Molecule and Glycolipids," The Journal of Biological Chemistry, vol. 273, No. 9, pp. 5190-5195 (Feb. 27, 1996). cited by applicant

Onufriev, A; Bashford, D.; Case, D. A Proteins 2004, 55, 383. cited by applicant

Orgueira H.A., et al., "Modular Synthesis of Heparin Oligosaccharides," Chemistry, vol. 09, No. 01, Jan. 3, 2003, pp. 140-169. cited by applicant

Ouyang et al., "Molecular Cloning and Expression of Human and Mouse Tyrosylprotein Sulfotransferase-2 and a Tyrosylprotein Sulfotransferase Homologue in *Caenorhabditis elegans*," The Journal of Biological Chemistry, vol. 273, No. 38, pp. 24770-24774 (Sep. 18, 1998). cited by applicant

Park et al., "Cell surface heparan sulfate proteoglycans: selective regulators of ligand-receptor encounters." J. Biol. Chem. vol. 275, pp. 29923-29926 (2000). cited by applicant

Patel, V. N.; Lombaert, I. M.A.; Cowherd, S. N.; Shworak, N.; Xu, Y.; Liu, J.; Hoffman, M. P. Developmental Cell 2014, 29, 662. cited by applicant

Pempe, et al., "Probing Structural Selectivity of Synthetic Heparin Binding to Stabilin Protein Receptors," Journal of Biol. Chem., vol. 287, No. 25, pp. 20774-20783 (Jun. 15, 2012). cited by applicant

Pettersen, E. F.; Goddard, T. D.; Huang, C. C.; Couch, G. S.; Greenblatt, D. M.; Meng, E. C.; Ferrin, T. E. J. Comp. Chem. 2004, 25, 1605. cited by applicant

Pierce et al., "Inflammatory response to trauma: implications for coagulation and resuscitation." Curr. Opin. Anesthesio., vol. 27, pp. 246-252 (2014). cited by applicant

Polat, T., et al., "Anomeric reactivity-based one-pot synthesis of heparinlike oligosaccharides.," J. Am. Chem. Soc., vol. 129, No. 42, pp. 12795-12800, Oct. 24, 2007. cited by applicant

Proudfoot et al., "Glycosaminoglycan binding and oligomerization are essential for the in vivo activity of certain chemokines." Proc. Natl. Acad. Sci. USA vol. 100, pp. 1885-1890 (2003). cited by applicant

Pulsipher et al., "Directing Neuronal Signaling through Cell-Surface Glycan Engineering." J. Am. Chem. Soc., vol. 136, pp. 6794-6797 (2014). cited by applicant

Raman, R.; Venkataraman, G.; Ernst, S.; Sasisekharan, R. Proc. Natl. Acad. Sci. 2003, 100, 2357. cited by applicant

Razi et al., "Structural and functional properties of heparin analogues obtained by chemical sulphation of *Escherichia coli* K5 capsular polysaccharide," Biochem. J., vol. 389, pp. 465-472 (1995). cited by applicant

Rejection decision corresponding to Chinese Patent Application No. 201880085012.5 dated Feb. 23, 2024. cited by applicant

Rohrmann et al., "Two N-acetylgalactosaminyltransferase are involved in the biosynthesis of chondroitin sulfate," European Journal of Biochemistry, vol. 148, pp. 463-469 (1985). cited by applicant

Roman-Blas et al., "The combined therapy with chondroitin sulfate plus glucosamine sulfate or chondroitin sulfate plus glucosamine hydrochloride does not improve joint damage in an experimental model of knee osteoarthritis in rabbits." Eur. J. Pharmacol., vol. 794, pp. 8-14 (2017). cited by applicant

Rosenberg et al., "Heparan Sulfate Proteoglycans of the Cardiovascular System Specific Structures Emerge But How Is Synthesis Regulated?" J. Clin. Invest., vol. 99, No. 9, pp. 2062-2070 (May 1997). cited by applicant

Saeki et al., "Molecular Cloning, Expression, and Characterization of a Novel Mouse Liver SULTIBI Sulfotransferase," J. Blochem., vol. 124, pp. 55-64 (1998). cited by applicant

Sala et al., "UDP-N-trifluoroacetylglucosamine as an alternative substrate in N-acetylglucosaminyltransferase reactions", *Carbohydrate Research*, vol. 306, pp. 127-136 (1998). cited by applicant

Sarris et al., "Inflammatory chemokines direct and restrict leukocyte migration within live tissues as glycan-bound gradients." *Curr. Biol.* vol. 22, pp. 2375-2382 (2012). cited by applicant

Sattelle, B. M., et al., "Free energy landscapes of iduronic acid and related monosaccharides," *J Am Chem Soc*, vol. 132, pp. 13132-13134, Sep. 29, 2010. cited by applicant

Sattelle, B. M.; Almond, A. *Glycobiology* 2011, 21, 1651. cited by applicant

Schroeder et al., "Protamine neutralization of low molecular weight heparins and their oligosaccharide components," *Anal Bioanal Chem*, vol. 399, pp. 763-771 (2011). cited by applicant

Schworer, R.; Zubkova, O. V.; Turnbull, J. E.; Tyler, P. C. *Chem. Eur. J.* 2013, 19, 6817. cited by applicant

Seffernick, J. L., et al., "Melamine deaminase and atrazine chlorohydrolase: 98 percent identical but functionally different," *Journal of Bacteriology*, vol. 183, No. 8, pp. 2405-2410, Apr. 2001. cited by applicant

Sen S., et al., "Developments in Directed Evolution for Improving Enzyme Functions," *Applied Biochemistry and Biotechnology*, vol. 143, No. 3, Dec. 2007, pp. 212-223. cited by applicant

Sheng et al., "The Dominating Role of N-Deacetylase/N-Sulfotransferase 1 in Forming Domain Structures in Heparan Sulfate," *The Journal of Biological Chemistry*, vol. 286, No. 22, pp. 19768-19776 (Jun. 3, 2011). cited by applicant

Sheng, J., et al., "Uncovering biphasic catalytic mode of C5-epimerase in heparan sulfate biosynthesis," *J. Biol. Chem.*, vol. 287, No. 25, pp. 20996-21002 (Jun. 15, 2012). cited by applicant

Shiori et al., "Sequence determination of synthesized chondroitin sulfate dodecasaccharides." *Glycobiology*, vol. 26, pp. 592-606 (2016). cited by applicant

Shively et al., "Formation of Anhydrosugars in the Chemical Depolymerization of Heparin," *Biochemistry*, vol. 15, No. 18, pp. 3932-3942 (1976). cited by applicant

Shriver et al., "Glycomics: A Pathway to a Class of New and Improved Therapeutics," *Nat. Rev. Drug Discov.*, vol. 3, pp. 863-873 (Oct. 2004). cited by applicant

Shriver, Z., et al., "Heparin and Heparan Sulfate: Analyzing Structure and Microheterogeneity." *Handb Exp. Pharmacol*, 207, pp. 159-176. (2012). cited by applicant

Shukla et al., "A Novel Role for 3-O-Sulfated Heparan Sulfate in Herpes Simplex Virus 1 Entry," *Cell*, vol. 99, pp. 13-22 (Oct. 1, 1999). cited by applicant

Shukla et al., "Herpesviruses and heparan sulfate: an intimate relationship in aid of viral entry," *The Journal of Clinical Investigation*, vol. 108, No. 4, pp. 503-510 (Aug. 2001). cited by applicant

Shworak et al., "Molecular Cloning and Expression of Mouse and Human cDNAs Encoding Heparan Sulfate D-Glucosaminyl 3-O-Sulfotransferase," *The Journal of Biological Chemistry*, vol. 272, No. 44), pp. 28008-28019 (1997). cited by applicant

Silk, E., et al., "The role of extracellular histone in organ injury", *Cell Death & Disease*, vol. 8, No. 5, 1, e2812, pp. 1-11, May 1, 2017. cited by applicant

Singh, A; Tessier, M. B.; Pederson, K.; Wang, X.; Venot, A P.; Boons, G.-J.; Prestegard, J. H.; Woods, R. J. *Can. J. Chem.* 2016, 10.1139/cjc. cited by applicant

Sismey-Ragatz, et al., "Chemoenzymatic Synthesis with Distinct Pasteurella Heparosan Synthases," *J. Biol. Chem.*, vol. 282, No. 39, pp. 28321-28327 (Jul. 11, 2007). cited by applicant

Solera et al., "Chondroitin sulfate tetrasaccharides: synthesis, three-dimensional structure and interaction with midkine." *Chemistry*, vol. 22, pp. 2356-2369 (2016). cited by applicant

Sommers, C.D., et al., "Heparin and homogeneous model heparin oligosaccharides form distinct complexes with protamine: Light scattering and zeta potential analysis," *Journal of Pharmaceutical and Biomedical Analysis*, vol. 140, pp. 113-121. (Year: 2017). cited by applicant

Stabler et al., "Chondroitin sulphate inhibits NF- κ B activity induced by interaction of pathogenic and damage associated molecules." *Osteoarthritis and Cartilage*, vol. 25, pp. 166-174 (2017). cited

by applicant

STN record for Chen et al., dissertation, "Towards de novo synthesis of structure-defined oligosaccharides with heparan sulfate u biosynthetic enzymes", entered into STN: Apr. 20, 2009 1 page. cited by applicant

Sugigura et al., "Molecular dissection of placental malaria protein VAR2CSA interaction with a chemo-enzymatically synthesized chondroitin sulfate library." Glycoconj. J., vol. 33, pp. 985-994 (2016). cited by applicant

Sugiura et al., "Baculovirus Envelope Protein ODV-E66 Is a Novel Chondroitinase with Distinct Substrate Specificity." J. Biol. Chem., vol. 286, pp. 29026-29034 (2011). cited by applicant

Sugiura et al., "Construction of a Chondroitin Sulfate Library with Defined Structures and Analysis of Molecular Interactions." J. Biol. Chem., vol. 287, pp. 43390-43400 (2012). cited by applicant

Sugiura et al., "Sequential synthesis of chondroitin oligosaccharides by immobilized chondroitin polymerase mutants." Glycoconj. J., vol. 25, pp. 521-530 (2008). cited by applicant

Sugumaran, G., et al., "Simultaneous Sulfation of endogenous Chondroitin Sulfate and Chondroitin-derived Oligosaccharides" The Journal of Biological Chemistry vol. 261 No. 27 pp. 12659-12664, Oct. 1986. cited by applicant

Sundaram, M. et al., "Rational design of low-molecular weight heparins with improved in vivo activity," Proc. Natl. Acad. Sci., vol. 100, No. 2, pp. 651-656 (Jan. 21, 2003). cited by applicant
Supplemental Notice of Allowability and Interview Summary corresponding to U.S. Appl. No. 14/898,865 dated Jan. 12, 2018. cited by applicant

Szajek et al., "The US regulatory and pharmacopeia responses to the global heparin contamination crisis." Nat. Biotechnol. vol. 34, pp. 625-630 (2016). cited by applicant

Szatmary, P., et al., "Biology, role and therapeutic potential of circulating histones in acute inflammatory disorders", Journal of Cellular and Molecular Medicine, vol. 22, No. 10, pp. 4617-4629, Aug. 7, 2018. cited by applicant

Takagaki et al., "Enzymatic Reconstruction of a Hybrid Glycosaminoglycan Containing 6-Sulfated, 4-Sulfated, and Unsulfated N-Acetylgalactosamine" Biochemical and Biophysical Research Communications vol. 258 pp. 741-744 (Year: 1999). cited by applicant

Takagaki, K., et al., "Chimeric Glycosaminoglycan Oligosaccharides Synthesized by Enzymatic Reconstruction and Their Use in Substrate Specificity Determination of *Streptococcus hyaluronidase*," The Japanese Biochemical Society, vol. 127, No. 4, pp. 695-702, Apr. 2000. cited by applicant

Tamura et al., "Synthesis of chondroitin sulfate E octasaccharide in a repeating region involving an acetamide auxiliary." Carbohydr. Res., vol. 343, pp. 39-47 (2008). cited by applicant

Tecle, E.; Diaz-Balzac, C. A.; Bulow, H. E. G3 (Bethesda) 2013, 3, 541. cited by applicant

Teng et al., "Molecular functions of syndecan-1 in disease." Matrix Biol., vol. 31, pp. 3-16 (2012). cited by applicant

Thacker, B. E.; Seamen, E.; Lawrence, R.; Parker, M. W.; Xu, Y.; Liu, J.; Vander, K. C. W.; Eska, J. D. ACS Chem. Biol. 2016, 11, 971. cited by applicant

Tohu et al., Anti-Xa and Anti-IIa Drugs Alter International Normalized Ratio Measurements: Potential Problems in the Monitoring of Oral Anticoagulants Clin. Appl. Thrombos Hemostas, vol. 10, pp. 301-309 (2004). cited by applicant

Tsau, C.; Ito, M.; Gromova, A.; Hoffman, M. P.; Meech, R.; Makarenkova, H. P. Development 2011, 138, 3307. cited by applicant

Tsung et al., "HMGB1 release induced by liver ischemia involves Toll-like receptor 4-dependent reactive oxygen species production and calcium-mediated signaling." J. Exp. Med., vol. 204, pp. 2913-2923 (2007). cited by applicant

Tsung et al., "The nuclear factor HMGB1 mediates hepatic injury after murine liver ischemia-reperfusion." J. Exp. Med., vol. 201, pp. 1135-1143 (2005). cited by applicant

Turnbull, J. E., Chemistry "Getting the farm out of pharma for heparin production," Science, vol.

334, No. 6055, pp. 462-463, Oct. 28, 2011. cited by applicant

Uchimura et al., "Molecular Cloning and Characterization of an N-Acetylglucosamine-6-O-sulfotransferase," The Journal of Biological Chemistry, vol. 273, No. 35, pp. 22577-22583 (Aug. 28, 1998). cited by applicant

Vann et al., "The Structure of the Capsular Polysaccharide (K5 Antigen) of Urinary-Tract-Infective *Escherichia coli* 010 : K5 : H4 A Polymer Similar to Desulfo-Heparin," Eur. J. Biochem, vol. 116, pp. 359-364 (1981). cited by applicant

Venereau et al., "HMGB1 as biomarker and drug target." Pharmacol. Res. vol. 111, pp. 534-544 (2016). cited by applicant

Vives, R, R., et al., "Sequence analysis of heparan sulphate and heparin oligosaccharides", Biochem. J, 1999, vol. 339, pp. 767-773. cited by applicant

Wang et al., "*E. coli* K5 fermentation and the Preparation of Heparosan, a Bioengineered Heparin Precursor," Biotechnol. Bioeng, vol. 107, No. 7, pp. 968-977 (Dec. 15, 2010). cited by applicant

Wang et al., "Endothelial heparan sulfate deficiency impairs L-selectin- and chemokine-mediated neutrophil trafficking during inflammatory responses." Nat. Immunol. vol. 6, pp. 902-910 (2005). cited by applicant

Wang, Z. et al., "Preactivation-based one-pot combinatorial synthesis of heparinlike hexasaccharides for the analysis of heparin-protein interactions," Chem. Eur.J., vol. 16, No. 28, pp. 8365-8375, Jul. 26, 2010. cited by applicant

Wang, Z., et al., "Synthesis of 3-O-sulfated oligosaccharides to understand the relationship between structures and functions of heparan sulfate," Journal of the American Chemical Society, Mar. 24, 2017, 139(14), pp. 5249-5256. cited by applicant

Weber et al., "Renal dysfunction in liver transplant recipients: Evaluation of the critical issues." Liver Transplant., vol. 18, pp. 1290-1301 (2012). cited by applicant

Weitz et al., "Beyond heparin and warfarin: the new generation of anticoagulants," Expert Opin. Investig. Drugs, vol. 16, No. 3, pp. 271-282 (2007). cited by applicant

Weitz, "Potential of new anticoagulants in patients with cancer," Thromb. Res., vol. 125 (Suppl 2), pp. S30-S35 (2010). cited by applicant

Whisstock, J, C., et al., "Prediction of protein function from protein sequence and structure," Quarterly Reviews of Biophysics, vol. 36, No. 3, pp. 307-340, Aug. 1, 2003. cited by applicant

Wildhagen et al., "Nonanticoagulant heparin prevents histone-mediated cytotoxicity in vitro and improves survival in sepsis." Blood vol. 123, pp. 1098-1101 (2014). cited by applicant

Written Opinion corresponding to International Application No. PCT/US2019/037993 dated Oct. 18, 2019. cited by applicant

Written Opinion of the International Searching Authority Corresponding to International Application No. PCT/US2018/021986 dated Aug. 1, 2018. cited by applicant

Xu et al., Chemoenzymatic Synthesis of Homogeneous Ultralow Molecular Weight Heparins, Science, vol. 334, No. 6055, pp. 498-501 (Oct. 2011). cited by applicant

Xu et al., "Heparan sulfate is essential for high mobility group protein 1 (HMGB1) signaling by the receptor for advanced glycation end products (RAGE)." J. Biol. Chem. vol. 286, pp. 41736-41744 (2011). cited by applicant

Xu et al., "Homogeneous and reversible low-molecular-weight heparins with reversible anticoagulant activity." Nat. Chem. Biol. vol. 10, pp. 248-250 (2014). cited by applicant

Xu et al., Synthetic oligosaccharides can replace animal-sourced low-molecular weight heparins Sci. Transl. Med. vol. 9, ean5954 (2017). cited by applicant

Xu J, Zhang X, Monestier M, Esmon NL, and Esmon CT. Extracellular histones are mediators of death through TLR2 and TLR4 in mouse fatal liver injury. J Immunol. 2011; 187:2626-31. cited by applicant

Xu J, Zhang X, Pelayo R, Monestier M, Ammollo CT, Semeraro F, et al. Extracellular histones are major mediators of death in sepsis. Nat Med. 2009;15:1318-21. cited by applicant

Xu, D. et al., "Engineering sulfotransferases to modify heparan sulfate," *Nat Chem Biol*, vol. 4, No. 3, pp. 200-202 (Mar. 2008). cited by applicant

Xu, D.; Esko, "Demystifying Heparan Sulfate-Protein Interactions," *J. Annu Rev Biochem*. 2014, 83, 129. cited by applicant

Xu, D.; Olson, J.; Cole, J. N.; van Wijk, X. M.; Brinkmann, V.; Zychlinsky, A.; Nizet, V.; ESKA, J. D.; Chang, Y. C. *Infect. Immun*. 2015, 83, 3648. cited by applicant

Xu, et. al., "Homogeneous low-molecular-weight heparins with reversible anticoagulant activity," *Nat Chem Biol*., vol. 10, pp. 248-252 (2014). cited by applicant

Xu, Y., et al., "Structure Based Substrate Specificity Analysis of Heparan Sulfate 6-O-Sulfotransferases," *ACS Chemical Biology*, Nov. 7, 2016, 12(1), pp. 73-82. cited by applicant

Xue et al., "Impact of donor binding on polymerization catalyzed by KfoC by regulating the affinity of enzyme for acceptor." *Biochim. Biophys. Acta*, vol. 1860, pp. 844-855 (2016). cited by applicant

Yang et al., "An Approach to Synthesize Chondroitin Sulfate-E (CS-E) Oligosaccharide Precursors." *J. Organic Chem.*, vol. 83, pp. 5897-5908 (2018). cited by applicant

Yang et al., "Middle region of the *Borrelia burgdorferi* surface-located protein 1 (Lmp1) interacts with host chondroitin-6-sulfate and independently facilitates infection." *Cell Microbiology*, vol. 18, 97-110 (2016). cited by applicant

Yang, "Inflammation plays a dual role in acetaminophen hepatotoxicity," *Translational Medicine Journal*, vol. 5, No. 3, pp. 129-133 (Jun. 2016). cited by applicant

Yang, J.; Hsieh, P.; Liu, X.; Zhou, W.; Zhang, X.; Zhao, J.; Xu, Y.; Zhang, F.; Linhardt, R. J.; Liu, J. *Chem Comm* 2017, 53, 1743. cited by applicant

Yang, Z.; et al., "UCSF Chimera, MODELLER, and IMP: an Integrated Modeling System," *J. Struct. Biol*. 2012, 179, 269. cited by applicant

Yoshinari et al., "Molecular Cloning, Expression, and Enzymatic Characterization of Rabbit Hydroxysteroid Sulfotransferase AST-RB2 (ST2A8)," *J. Biochem.*, vol. 123, pp. 740-746 (1998). cited by applicant

Yu et al., "Highly Efficient Chemoenzymatic Synthesis of Beta1-3-Linked Galactosides," *Chemical Communications*, vol. 46(40), pp. 7507-7509 (2010). cited by applicant

Yusa et al., "N-Linked Oligosaccharides on Chondroitin 6-Sulfotransferase-1 Are Required for Production of the Active Enzyme, Golgi Localization, and Sulfotransferase Activity toward Keratan Sulfate." *J. Biol. Chem.*, vol. 281, pp. 20393-20403 (2006). cited by applicant

Zhang et al., "6-O-Sulfotransferase-1 Represents a Critical Enzyme in the Anticoagulant Heparan Sulfate Biosynthetic Pathway*," *J. Biol. Chem.*, vol. 276, pp. 42311-42321 (2001). cited by applicant

Zhang et al., "Solution Structures of Chemoenzymatically Synthesized Heparin and Its Precursors," *J. Am. Chem. Soc.*, vol. 130, pp. 12998-13007 (2008). cited by applicant

Zhang et al., "The Effect of Precursor Structures on the Action of Glucosaminyl 3-O-Sulfotransferase-1 and the Biosynthesis of Anticoagulant Heparan Sulfate," *J. Biol. Chem.*, vol. 276, No. 31, pp. 28806-28813 (2001). cited by applicant

Zhang, Z. et al., "Oversulfated chondroitin sulfate: impact of a heparin impurity, associated with adverse clinical events, on low-molecular-weight heparin preparation," *J. Med. Chem.*, vol. 51, No. 18, pp. 5498-5501, Sep. 25. 2008). cited by applicant

Zhao et al. "Enzymatic route to preparative-scale synthesis of UDP-GlcNAc/GalNAc, their analogues and GDP-fucose," *Nat. Protoc.*, vol. 5, No. 4, pp. 636-646 (2010). cited by applicant

Zhou et al. "Expression of heparin sulfate sulfotransferases in *Kluyveromyces lactis* and preparation of 3'-phosphoadenosine-5'-phosphosulfate," *Glycobiology*, vol. 21, No. 6, pp. 771-780 (2011). cited by applicant

Zitvogel et al., "Decoding cell death signals in inflammation and immunity." *Cell*, vol. 140, pp. 798-804 (2010). cited by applicant

Zong, C.; Huang, R.; Condac, E.; Chiu, Y.; Xiao, W.; Li, Z. Q.; Lu, W.; Ishihara, M.; Wang, S.; Ramiah, A.; Stickney, M.; Azadi, P.; Amster, I. J.; Moremen, K. W.; Wang, L.; Sharp, J. S.; Boons, G.-J. J. Am. Chem. Soc. 2016, 138, 13059. cited by applicant
Ham, H., et al., "Design of an Ultralow Molecular Weight Heparin that Resists Heparanase Biodegradation," Journal of Medicinal Chemistry, vol. 66, 2023, pp. 2194-2203. cited by applicant
Patent Certificate for European Patent No. 3691653 dated Mar. 12, 2025. cited by applicant
Notice of Allowance for Japanese Patent Application No. 2024-008526 dated Jul. 4, 2025. cited by applicant

Primary Examiner: Berry; Layla D

Attorney, Agent or Firm: Jenkins, Taylor & Hunt, P.A.

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a divisional application of U.S. patent application Ser. No. 16/761,159, filed May 1, 2020, which is a national phase patent application under 35 U.S.C. Section 371 of PCT International Patent Application No. PCT/US2018/059152, filed Nov. 5, 2018, which claims priority to and benefit of U.S. Provisional Patent Application Ser. No. 62/581,443, filed Nov. 3, 2017. The disclosure of each of these applications is incorporated herein by reference in its entirety.

TECHNICAL FIELD

(1) Disclosed herein in some embodiments are new sulfated heparan sulfate oligosaccharide compounds having anti-inflammatory activity. For example, in some embodiments it is observed that the compounds display a protective effect on drug-induced liver damage. While it is not desired to be bound by any particular theory of operation, the compounds likely inhibit the interaction of HMGB1/RAGE, a key signaling interaction involved in sterile inflammatory injury. This discovery can be applied for a wide range of diseases that are caused by over reaction of inflammation responses.

BACKGROUND

(2) Sterile inflammation is a natural process that initiates tissue repair in response to cellular damage. However, exaggerated inflammation after an initial insult often damages surrounding healthy tissues and is a key contributor to many disease processes. High mobility group box 1 (HMGB1) is a DNA binding protein that regulates transcription and is released from the nucleus during necrotic cell death (Zitvogel, 2010; Chen, 2010). Extracellularly, HMGB1 is a damage associated molecular pattern (DAMP) protein that acts as a pro-inflammatory molecule, orchestrating migration and activation of inflammatory cells to the injury site (Bianchi, 2017). HMGB1-dependent inflammation has been associated with ischemia-reperfusion and drug-induced liver injury (Huebener, 2015).

(3) Acetaminophen (APAP), also known as paracetamol, is a widely used analgesic and is the active pharmaceutical ingredient of Tylenol®. Ingestion of a supratherapeutic dose causes acute liver failure (ALF) (Heard, 2008). The misuse of Vicodin® or Percocet®, co-formulations of opioids and APAP, can also cause ALF. In the US, nearly 50% of drug-induced liver injury has been attributed to APAP toxicity (Lee, 2007), which accounts for ~80,000 emergency room visits annually (Blieden, 2014). The mechanism for APAP toxicity begins with its metabolic conversion to the reactive chemical species, N-acetyl-p-benzoquinone imine (NAPQI), which causes hepatocyte necrosis. (Tacke, 2015). Necrotic hepatocytes release HMGB1 which drives chemotaxis of neutrophils through the receptor for advanced glycation end-products (RAGE), activating sterile

inflammation and amplifying liver injury (Huebener 2015).

(4) However, additional compositions and methods for treating inflammation in subject in need thereof remain a need in the art.

SUMMARY

(5) This summary lists several embodiments of the presently disclosed subject matter, and in many cases lists variations and permutations of these embodiments. This summary is merely exemplary of the numerous and varied embodiments. Mention of one or more representative features of a given embodiment is likewise exemplary. Such an embodiment can typically exist with or without the feature(s) mentioned; likewise, those features can be applied to other embodiments of the presently disclosed subject matter, whether listed in this summary or not. To avoid excessive repetition, this summary does not list or suggest all possible combinations of such features.

(6) In some embodiments, provided herein are small molecule compounds having an anti-inflammatory property, the small molecule compound comprising a non-anticoagulant heparan sulfate oligosaccharide molecule, optionally wherein the non-anticoagulant heparan sulfate oligosaccharide molecule is configured to interact with a high mobility group box 1 (HMGB1) protein in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation end products (RAGE). The oligosaccharide molecule can comprise about 10 to about 20 saccharide units, or about 12 to about 18 saccharide units. The oligosaccharide molecule can comprise about 18 saccharide units.

(7) In some embodiments, provided herein is a small molecule compound comprising a disaccharide structure unit as shown:

(8) ##STR00001##

wherein R=—H or —SO₃H. The disaccharide structure unit can be selected from the group consisting of non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin, and O-desulfated heparin (ODSH). In some embodiments, the oligosaccharide molecule protects against liver injury in vivo. In some embodiments, the oligosaccharide molecule decreases neutrophil infiltration in vivo. In some embodiments, the oligosaccharide molecule decreases inflammation in vivo.

(9) Provided herein are methods of treating a subject, comprising providing a subject to be treated, wherein the subject is suffering from inflammation, and administering to the subject a non-anticoagulant heparan sulfate oligosaccharide molecule having an anti-inflammatory property, optionally wherein the non-anticoagulant heparan sulfate oligosaccharide molecule is configured to interact with a high mobility group box 1 (HMGB1) protein in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation endproducts (RAGE). In some embodiments, the subject suffers from any injury resulting in increased inflammation. In some embodiments, the subject suffers from liver injury. In some embodiments, the subject in need of treatment is a subject suffering from an overdose of Paracetamol (APAP). In some embodiments, the subject in need of treatment is a human subject. In some embodiments, the small molecule compound comprises a disaccharide structure unit selected from the group consisting of non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin, and O-desulfated heparin (ODSH). In some embodiments, the oligosaccharide molecule comprises about 10 to about 20 saccharide units, or about 12 to about 18 saccharide units. In some embodiments, the oligosaccharide molecule comprises about 18 saccharide units. Administration of the oligosaccharide molecule can decrease neutrophil infiltration in the subject. Administration of the oligosaccharide molecule can decrease inflammation in the subject. Administration of the oligosaccharide molecule can protect against liver damage and multi-organ system failure in the subject. In some aspects, the subject to be treated suffers from drug-induced inflammation.

(10) In some aspects, provided herein are methods of treating Paracetamol (APAP) overdose in a subject, the method comprising providing a subject in need of treatment for APAP overdose, administering to the subject a non-anticoagulant heparan sulfate oligosaccharide molecule having

an anti-inflammatory property, optionally wherein the non-anticoagulant heparan sulfate oligosaccharide molecule is configured to interact with a high mobility group box 1 (HMGB1) protein in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation endproducts (RAGE), wherein damage from the APAP overdose in the subject is mitigated. The treatment for the APAP overdose in the subject can in some embodiments be effective between 0 hours and 24 hours after the overdose, or at least within 12 hours after the overdose. The treatment for the APAP overdose in the subject can comprise protection against liver injury and/or multi-organ system failure. The treatment for the APAP overdose in the subject can comprise a decrease in neutrophil infiltration in the subject. The treatment for the APAP overdose in the subject can comprise blocking the interaction between the HMGB1 protein and RAGE. In some aspects, the small molecule compound comprises a disaccharide structure unit as shown:

(11) ##STR00002##

wherein R=—H or —SO₃H. The small molecule compound can comprise a disaccharide structure unit is selected from the group consisting of non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin, and O-desulfated heparin (ODSH). The oligosaccharide molecule comprises about 10 to about 20 saccharide units, or about 12 to about 18 saccharide units. In some aspects, the oligosaccharide molecule comprises about 18 saccharide units.

(12) Accordingly, it is an object of the presently disclosed subject matter to provide sulfated heparan sulfate oligosaccharide compounds having anti-inflammatory activity and methods for making and using the same.

(13) This and other objects are achieved in whole or in part by the presently disclosed subject matter. Further, an object of the presently disclosed subject matter having been stated above, other objects and advantages of the presently disclosed subject matter will become apparent to those skilled in the art after a study of the following description, Drawings and Examples.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) The presently disclosed subject matter can be better understood by referring to the following figures. The components in the figures are not necessarily to scale, emphasis instead being placed upon illustrating the principles of the presently disclosed subject matter (often schematically). In the figures, like reference numerals designate corresponding parts throughout the different views. A further understanding of the presently disclosed subject matter can be obtained by reference to an embodiment set forth in the illustrations of the accompanying drawings. Although the illustrated embodiment is merely exemplary of systems for carrying out the presently disclosed subject matter, both the organization and method of operation of the presently disclosed subject matter, in general, together with further objectives and advantages thereof, may be more easily understood by reference to the drawings and the following description. The drawings are not intended to limit the scope of this presently disclosed subject matter, which is set forth with particularity in the claims as appended or as subsequently amended, but merely to clarify and exemplify the presently disclosed subject matter.

(2) For a more complete understanding of the presently disclosed subject matter, reference is now made to the following drawings in which:

(3) FIGS. 1A-1G are schematic illustrations of a chemoenzymatic synthetic pathway for the synthesis of 18-mer oligosaccharides;

(4) FIG. 2 is a schematic illustration of a chemical structure of an 18-mer oligosaccharide as disclosed herein;

(5) FIGS. 3A through 3D are graphical depictions of data showing the effects of an 18-mer oligosaccharide on liver injury after APAP overdose;

- (6) FIGS. 4A through 4D are graphical depictions of data showing 18-mer oligosaccharide targeting HMGB1 to decrease inflammation;
- (7) FIG. 4E show symbolic structures of HS oligosaccharides;
- (8) FIG. 4F is a legend for FIG. 4E;
- (9) FIGS. 5A through 5D are graphical depictions of data showing the effects of delayed 18-mer oligosaccharide treatment on APAP overdose;
- (10) FIGS. 6A-6D are graphical depictions of data showing biological parameters of APAP overdosed mice;
- (11) FIG. 7 is a schematic of the chemical structures of biotinylated and non-biotinylated oligosaccharides: 6-mer, 12-mer, 18-mer and 18-mer AXa;
- (12) FIGS. 8A-8D are graphical depictions of data showing the protective effects of 6-mer, 12-mer, 18-mer and 18-mer AXa oligosaccharide;
- (13) FIGS. 9A and 9B are graphical depictions of data showing APAP overdose results in increased levels of ALT, migrating neutrophils and shed syndecan-1; and
- (14) FIGS. 10A through 10C are graphical depictions of data based on the analysis of syndecan-1 from ALF patients and the plasma concentration of ALT and HMGB1 in ALF patients.

DETAILED DESCRIPTION

(15) The presently disclosed subject matter now will be described more fully hereinafter, in which some, but not all embodiments of the presently disclosed subject matter are described. Indeed, the presently disclosed subject matter can be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are provided so that this disclosure will satisfy applicable legal requirements.

(16) Heparan sulfate (HS) is a sulfated polysaccharide abundantly present on cell surface and in extracellular matrix. Many DAMPs, including HMGB1, are HS-binding proteins (Xu, 2011). HS participates in various aspects of inflammation including chemokine presentation and neutrophil transendothelial migration (Proudfoot, 2003; Wang, 2005; Axelsson, 2013; and Sarris, 2012). HS is comprised of disaccharide repeating units of glucuronic acid (GlcA) or iduronic acid (IdoA) linked to glucosamine residues that carry sulfo groups. The chain length and sulfation pattern of HS determine its biological functions (Gama, 2006). In accordance with aspects of the presently disclosed subject matter, synthesis of a specially designed HS octadecasaccharide (18-mer) to exploit the anti-inflammatory effect is provided. The 18-mer protects APAP-induced acute liver failure through neutralizing the pro-inflammatory activity of HMGB1 in a murine model. The results presented offer a new chemical space to curb HMGB1-mediated inflammatory diseases.

(17) In some embodiments, a small molecule compound with anti-inflammatory properties is disclosed. In some embodiments, the small molecule compound comprises a non-anticoagulant heparan sulfate oligosaccharide molecule with anti-inflammatory properties. In some embodiments, the non-anticoagulant heparan sulfate oligosaccharide molecule is configured to interact with a high mobility group box 1 (HMGB1) protein in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation endproducts (RAGE).

(18) In some embodiments, a small molecule compound in accordance with the presently disclosed subject matter comprises, or can be part of a composition comprising, a disaccharide structure unit as shown:

(19) ##STR00003##

wherein R=—H or —SO₃H.

By way of example and not limitation, this disaccharide composition can be found in non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin or O-desulfated heparin (ODSH). A composition in accordance with the presently disclosed subject matter can comprise a non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin or O-desulfated heparin (ODSH).

(20) In some embodiments, the oligosaccharide molecule comprises about 10 to about 20

saccharide units, or about 12 to about 18 saccharide units. In some embodiments, the oligosaccharide molecule comprises about 18 saccharide units. In some embodiments, generic structures of heparan sulfate oligosaccharides displaying anti-inflammatory effects comprise:

(21) ##STR00004##

wherein R_{sup}.3 is H or a detectable tag. In some embodiments, the detectable tag comprises para-nitrophenyl.

(22) In some embodiments, the oligosaccharide molecule comprises one of the following structures:

(23) ##STR00005## ##STR00006## ##STR00007##

(24) The compositions of the presently disclosed subject matter comprise in some embodiments a composition that includes a pharmaceutically acceptable carrier. Any suitable pharmaceutical formulation can be used to prepare the compositions for administration to a subject. In some embodiments, the composition and/or carriers can be pharmaceutically acceptable in humans.

(25) For example, suitable formulations can include aqueous and non-aqueous sterile injection solutions that can contain anti-oxidants, buffers, bacteriostatics, bactericidal antibiotics, and solutes that render the formulation isotonic with the bodily fluids of the subject; and aqueous and non-aqueous sterile suspensions that can include suspending agents and thickening agents. The formulations can be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials, and can be stored in a frozen or freeze-dried (lyophilized) condition requiring only the addition of sterile liquid carrier, for example water for injections, immediately prior to use. Some exemplary ingredients are sodium dodecyl sulfate (SDS), in one example in the range of 0.1 to 10 mg/ml, in another example about 2.0 mg/ml; and/or mannitol or another sugar, for example in the range of 10 to 100 mg/ml, in another example about 30 mg/ml; and/or phosphate-buffered saline (PBS).

(26) It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this presently disclosed subject matter can include other agents conventional in the art having regard to the type of formulation in question. For example, sterile pyrogen-free aqueous and non-aqueous solutions can be used.

(27) Methods of treating a subject are also provided in accordance with the presently disclosed subject matter, including but not limited to methods that treat conditions comprising inflammation. The therapeutic methods of the presently disclosed subject matter can comprise administering to the subject a non-anticoagulant heparan sulfate oligosaccharide molecule. In some embodiments, the non-anticoagulant heparan sulfate oligosaccharide molecule is configured to interact with a high mobility group box 1 (HMGB1) protein in a manner sufficient to affect an interaction between the HMGB1 protein and a receptor for advanced glycation endproducts (RAGE).

(28) In some embodiments, the oligosaccharide molecule protects against liver injury in vivo. In some embodiments, the oligosaccharide molecule decreases neutrophil infiltration in vivo. In some embodiments, the oligosaccharide molecule decreases inflammation in vivo.

(29) In some embodiments, the subject is suffering from inflammation. In some embodiments, the subject suffers from any injury resulting in increased inflammation. In some embodiments, the subject suffers from liver injury. In some embodiments, the subject in need of treatment is a subject suffering from an overdose of Paracetamol (acetaminophen or APAP).

(30) In some embodiments, the subject treated in the presently disclosed subject matter is desirably a human subject, although it is to be understood the methods described herein are effective with respect to all vertebrate species (e.g., mammals, birds, etc.), which are intended to be included in the term "subject."

(31) More particularly, provided herein is the treatment of mammals, such as humans, as well as those mammals of importance due to being endangered (such as Siberian tigers), of economical importance (animals raised on farms for consumption by humans) and/or social importance (animals kept as pets or in zoos) to humans, for instance, carnivores other than humans (such as

cats and dogs), swine (pigs, hogs, and wild boars), ruminants (such as cattle, oxen, sheep, giraffes, deer, goats, bison, and camels), and horses. Thus, the methods described herein include the treatment of livestock, including, but not limited to, domesticated swine (pigs and hogs), ruminants, horses, and the like.

(32) In some embodiments, the oligosaccharide molecule decreases neutrophil infiltration in the subject. In some embodiments, the administration of the oligosaccharide molecule decreases inflammation in the subject. In some embodiments, the administration of the oligosaccharide molecule protects against liver damage and multi-organ system failure in the subject. In some embodiments, the subject to be treated suffers from drug-induced inflammation. In some embodiments, damage from the APAP overdose in the subject is mitigated.

(33) In some embodiments, the treatment for the APAP overdose in the subject is effective between 0 hours and 24 hours after the overdose, or in some embodiments at least within 12 hours after the overdose, and in some embodiments within 6 hours after overdose. In some embodiments, the treatment for the APAP overdose in the subject comprises protection against liver injury and/or multi-organ system failure. In some embodiments, the treatment for the APAP overdose in the subject comprises a decrease in neutrophil infiltration in the subject. In some embodiments, the treatment for the APAP overdose in the subject comprises blocking the interaction between the HMGB1 protein and RAGE.

(34) In some embodiments, a small molecule compound in accordance with the presently disclosed subject matter comprises a disaccharide structure unit as shown:

(35) ##STR00008##

wherein R=—H or —SO₂—H.

By way of example and not limitation, this disaccharide structure unit can be found in non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin or O-desulfated heparin (ODSH). A composition employed in accordance with the presently disclosed methods can comprise a non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin or O-desulfated heparin (ODSH). In some embodiments, the composition protects against liver injury in vivo, e.g., provides hepatoprotection in vivo. In some embodiments, the oligosaccharide molecule decreases neutrophil infiltration in vivo. In some embodiments, the subject suffers from liver injury. In some embodiments, the subject in need of treatment is a subject suffering from an overdose of Paracetamol (acetaminophen or APAP).

(36) In some embodiments, the oligosaccharide molecule comprises about 10 to about 20 saccharide units, or about 12 to about 18 saccharide units. In some embodiments, the oligosaccharide molecule comprises about 18 saccharide units. In some embodiments, generic structures of heparan sulfate oligosaccharides displaying anti-inflammatory effects comprise:

(37) ##STR00009##

wherein R_{sup}.3 is H or a detectable tag. In some embodiments, the detectable tag comprises para-nitrophenyl.

(38) In some embodiments, the oligosaccharide molecule comprises one of the following structures:

(39) ##STR00010## ##STR00011## ##STR00012##

Definitions

(40) The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the presently disclosed subject matter.

(41) While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

(42) All technical and scientific terms used herein, unless otherwise defined below, are intended to have the same meaning as commonly understood by one of ordinary skill in the art. References to techniques employed herein are intended to refer to the techniques as commonly understood in the

art, including variations on those techniques or substitutions of equivalent techniques that would be apparent to one skilled in the art. While the following terms are believed to be well understood by one of ordinary skill in the art, the following definitions are set forth to facilitate explanation of the presently disclosed subject matter.

(43) The term “non-anticoagulant” and phrase “without anticoagulant” used in this application describes presently disclosed subject matter having less than or equal to about 50 units/mg anticoagulant activity where the anticoagulant activity is attributed to inhibition of Factor Xa and/or Factor IIa.

(44) In describing the presently disclosed subject matter, it will be understood that a number of techniques and steps are disclosed. Each of these has individual benefit and each can also be used in conjunction with one or more, or in some cases all, of the other disclosed techniques.

(45) Accordingly, for the sake of clarity, this description will refrain from repeating every possible combination of the individual steps in an unnecessary fashion. Nevertheless, the specification and claims should be read with the understanding that such combinations are entirely within the scope of the invention and the claims.

(46) Following long-standing patent law convention, the terms “a”, “an”, and “the” refer to “one or more” when used in this application, including the claims. Thus, for example, reference to “a unit cell” includes a plurality of such unit cells, and so forth.

(47) Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about”. Accordingly, unless indicated to the contrary, the numerical parameters set forth in this specification and attached claims are approximations that can vary depending upon the desired properties sought to be obtained by the presently disclosed subject matter.

(48) As used herein, the term “about,” when referring to a value or to an amount of a composition, mass, weight, temperature, time, volume, concentration, percentage, etc., is meant to encompass variations of in some embodiments $\pm 20\%$, in some embodiments $\pm 10\%$, in some embodiments $\pm 5\%$, in some embodiments $\pm 1\%$, in some embodiments $\pm 0.5\%$, and in some embodiments $\pm 0.1\%$ from the specified amount, as such variations are appropriate to perform the disclosed methods or employ the disclosed compositions.

(49) The term “comprising”, which is synonymous with “including” “containing” or “characterized by” is inclusive or open-ended and does not exclude additional, unrecited elements or method steps. “Comprising” is a term of art used in claim language which means that the named elements are essential, but other elements can be added and still form a construct within the scope of the claim.

(50) As used herein, the phrase “consisting of” excludes any element, step, or ingredient not specified in the claim. When the phrase “consists of” appears in a clause of the body of a claim, rather than immediately following the preamble, it limits only the element set forth in that clause; other elements are not excluded from the claim as a whole.

(51) As used herein, the phrase “consisting essentially of” limits the scope of a claim to the specified materials or steps, plus those that do not materially affect the basic and novel characteristic(s) of the claimed subject matter.

(52) With respect to the terms “comprising”, “consisting of”, and “consisting essentially of”, where one of these three terms is used herein, the presently disclosed and claimed subject matter can include the use of either of the other two terms.

(53) As used herein, the term “and/or” when used in the context of a listing of entities, refers to the entities being present singly or in combination. Thus, for example, the phrase “A, B, C, and/or D” includes A, B, C, and D individually, but also includes any and all combinations and subcombinations of A, B, C, and D.

EXAMPLES

(54) The following examples are included to further illustrate various embodiments of the presently disclosed subject matter. However, those of ordinary skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the presently disclosed subject matter.

Example 1

(55) Acetaminophen/paracetamol (APAP) overdose is the leading cause of drug-induced acute liver failure (ALF) in the US and Europe. The progression of the disease is attributed to sterile inflammation induced by the release of high mobility group box 1 (HMGB1). A specific, effective and safe approach to neutralize the pro-inflammatory activity of HMGB1 is highly desirable. Disclosed herein in some embodiments is a HS octadecasaccharide (18-mer) that displays potent hepatoprotection by targeting to HMGB1. Investigation of the role of endogenous syndecan-1 in response to APAP overdose substantiated the mechanism for 18-mer's protection. The presently disclosed data suggest that 18-mer potentiates the host anti-inflammation effect mediated by syndecan-1. Finally, it is demonstrated that 18-mer administered six hours after APAP overdose is still protective, and therefore offers a therapeutic advantage over N-acetyl cysteine for late-presenting patients. Synthetic HS opens a new approach for the treatment of ALF and other HMGB1-involved inflammatory diseases

(56) A series of novel sulfated heparan sulfate oligosaccharide compounds were synthesized. In particular, chemoenzymatic synthesis was employed to obtain a pure HS/heparin-like oligosaccharide without anticoagulant activity. One exemplary compound, termed 18-mer NS2S, was used for in depth in vivo studies in mice. 18-mer NS2S significantly decreased injury in the context of acetaminophen (APAP)-induced acute liver failure. Although it is not desired to be bound by any particular theory of operation, the mechanism of action of 18-mer NS2S likely involves the HMGB1/RAGE. 18mer NS2S treatment is still effective 6 hours after APAP overdose in mice whereas the clinically used antidote, NAC, loses effectiveness at this time. Additional description of these observations can be found in the Examples that follow.

(57) Chemoenzymatic synthesis was used to generate novel, pure oligosaccharide compounds without anticoagulant activity. Representative synthetic routes are disclosed in the Examples that follow. A series of 18-mer compounds and one 12-mer, and one 6-mer compound were used in vivo. One exemplary 18-mer NS2S was shown to be protective against liver injury in vivo. In this example, 18-mer NS2S significantly increases the survival rate in vivo. 18-mer NS2S decreases neutrophil infiltration. RAGE knockout mice were observed to be insensitive to 18-mer NS2S's protective effects. This observation supports that delayed treatment after acetaminophen (APAP) overdose is possible.

Example 2

Materials and Methods

(58) Study Design

(59) This study was designed to synthesize a HS octadecasaccharide (18-mer) and evaluate its anti-inflammatory effect in an APAP-induced liver failure murine model. The chemoenzymatic synthesis of two HS 18-mers, including 18-mer NS2S (nonanticoagulant compound; referred to as 18-mer) and 18-mer AXa (anticoagulant compound) is demonstrated herein. The structures of two 18-mers were confirmed using both nuclear magnetic resonance (NMR) and high resolution mass spectroscopy (MS). Demonstration of the binding between HMGB1 and biotinylated 18-mer and 18-mer AXa was achieved using avidin-agarose column followed by western analysis. HMGB1 binds to 18-mer and 18-mer AXa in nanomolar ranges as determined by surface plasmon resonance (SPR). The hepatoprotection effect of the synthesized 18-mers was evaluated in a well-established APAP-induced liver failure murine model. The liver damage was assessed by two methods, including the plasma ALT level and examination of hematoxylin and eosin (H&E) stained liver sections. The inflammation responses after APAP overdose were assessed by determining the

neutrophil migration to the injury sites. Factor Xa (FXa) activity was used as a surrogate to assess the anticoagulant activity of 18-mers, 12-mer and 6-mer in both in vitro and in vivo experiments. The number of animals in the tested groups and controls groups as well as statistical analysis are presented in the figure legends. Anonymous patient ALF plasma samples were obtained from the Acute Liver Failure Study Group (ALFSG) biorepository (of the National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, Maryland, United States of America) to determine the plasma level of shed syndecan-1. The analyses were stopped after analyzing 31 patient samples as a clear statistical difference between healthy control group (n=11) and AFL patients was observed.

(60) Expression of HS Biosynthetic Enzymes

(61) A total of nine enzymes was used for the synthesis, including NST, C.sub.5-epi, 2-OST, 6-OST-1, 6-OST-3, 3-OST-1, 3-OST-5, and pmHS2. All enzymes, with the exception of C.sub.5-epi and 2-OST, were expressed in *E. coli* and purified by appropriate affinity chromatography as described previously (Renpeng et al., 2010; Xu et al., 2008). Recombinant C.sub.5-epi and 2-OST were expressed in insect cells using the Bac-to-Bac baculovirus expression approach (from Invitrogen) to obtain high expression levels (19). Three enzyme co-factors, including 3'-phosphoadenosine 5'-phosphosulfate (PAPS), uridine 5'-diphosphoglucuronic acid (UDP-GICA), uridine 5'-diphospho N-trifluoroacetyl glucosamine (UDP-GlcNTFA), were all synthesized in-house using enzymatic approaches as described previously (Nam, 2017).

(62) Synthesis of Non-Biotinylated 18-Mer AXa

(63) The synthesis of 18-mer AXa was initiated from 18-mer. Two enzymatic modifications were added, including 6-O-sulfotransferases (6-OST) modification and 3-O-sulfotransferase isoform 1 (3-OST-1) modification. In the 6-O-sulfation step, the 18-mer (0.5 mM) was incubated in a buffer containing MES (50 mM, pH 7.0), 6-OST-1 (50 µg ml.sup.-1), 6-OST-3 (50 µg ml.sup.-1) and PAPS (1.5 equiv. of 6-hydroxyl group amount) at 37° C. overnight in 60 ml. The product was then purified by Q-Sepharose column for the subsequent 3-O-sulfation step.

(64) The 3-O-sulfation step was completed by 3-OST-1 enzyme. The 6-O-sulfated 18-mer (0.5 mM) was incubated in a solution containing MES (50 mM) buffer (pH 7.0), 3-OST-1 (20 µg ml.sup.-1) and PAPS (0.675 mM) in 20 ml at 37° C. for 2 hours. The product was purified by Q-Sepharose column. During the synthesis, the products were monitored by HPLC using a DEAE-NPR column (4.6 mm×75 mm, from Tosoh).

(65) The synthesis of non-biotinylated 6-mer and non-biotinylated 12-mer was completed using the enzymatic method, and was reported previously (Xu, 2017).

(66) Conversion of Nonbiotinylated Oligosaccharides to Biotinylated Oligosaccharides

(67) Four nonbiotinylated oligosaccharides, 6-mer, 12-mer, 18-mer and 18-mer AXa, were converted to biotinylated counterparts. 6-mer, 12-mer, 18-mer and 18-mer AXa with a pNP tag (5-10 mg) and 0.5 mg Pd/C were dissolved in 20 mM NaOAc, pH 5.0 in a total volume of 4 ml. Reaction mixture was vacuumed and refilled with H.sub.2 three times. The reaction was then incubated at room temperature for 4 h. After that, it was filtered to remove charcoal. The filtered solution was adjusted to pH 8.5 using 500 mM Na.sub.2HPO.sub.4. Succinimidyl 6-azidohexanoate (20 molar equivalent of starting oligosaccharides) was added and incubated at 37° C. overnight. Reaction was purified by DEAE-HPLC column to generate azido tagged oligosaccharides. PBS (pH7.4) buffer was bubbled with N.sub.2 for 5 min to prepare the sample solution of 0.1M CuSO.sub.4, 0.1M Tris(3-hydroxypropyl-triazolylmethyl)amine (THPTA) (Sigma, Burlington, Massachusetts, United States of America), 0.15M sodium ascorbate, 0.01M azido tagged oligosaccharides and 0.02 M biotin-PEG.sub.4-alkyne (Sigma). The mixture of 400 µl THPTA and 80 µl CuSO.sub.4 was vortexed, then 160 µl sodium ascorbate, 200 µl azido tagged oligomers and 200 µl biotin-PEG.sub.4-alkyne was added and bubbled with N.sub.2 for 2 min, then incubated at 37° C. overnight. The reaction was purified by DEAE-HPLC column to generate biotinylated products. The reactions were monitored using HPLC and MS.

(68) HPLC Analysis

(69) Both DEAE-NPR HPLC and polyamine-based anion exchange (PAMN)-HPLC were used to analyze the purity of the products. The elution conditions for the HPLC analysis were described elsewhere (Renpeng et al., 2010). Briefly, for DEAE-HPLC method, the column TSK gel DNA-NPR (4.6×75 mm, from Tosoh Bioscience) was eluted with a linear gradient of NaCl in 20 mM sodium acetate buffer (pH 5.0) from 0 to 1M for 60 min at a flow rate of 0.4 ml min⁻¹. As for PAMN-HPLC, the column (Polyamine II-HPLC, 4.6×250 mm, from YMC) was eluted with a linear gradient of KH.sub.2PO.sub.4 from 0 to 1M for 40 min then remained at 1M for 30 min at a flow rate of 0.5 ml min⁻¹.

(70) MS Analysis of Oligosaccharides

(71) The low-resolution analyses were performed at a Thermo LCQ-Deca. Oligosaccharides were directly diluted in 200:1 of mixture of MeOH/H.sub.2O (9:1, vol/vol). A syringe pump (Harvard Apparatus) was used to introduce the sample by direct infusion (50 µl min⁻¹). Experiments were carried out in negative ionization mode. Synthetic nonsulfated oligosaccharides were diluted in 200 µl of H.sub.2O with the electrospray source set to 5 kV and 275° C. Sulfated oligosaccharides were diluted in 200 µl of 10 mM ammonium bicarbonate with the electrospray source set to 3 KV and 150° C. The automatic gain control was set to 1×10^{sup.7} for full scan MS. The MS data were acquired and processed using Xcalibur 1.3.

(72) High resolution ESI-MS analysis was conducted on Thermo LTQ XL Orbitrap (Bremen, Germany) under the following conditions. A Luna hydrophilic liquid interaction chromatography (HILIC) column (2.0×50 mm.sup.2, 200 Å, Phenomenex, Torrance, California, United States of America) was used to separate the oligosaccharide mixture. Mobile phase A was 5 mM ammonium acetate prepared with high performance liquid chromatography (HPLC) grade water. Mobile phase B was 5 mM ammonium acetate prepared in 98% HPLC grade acetonitrile with 2% of HPLC grade water. After injection of 5.0 µl 12-mer mixture (1.0 µg µl.sup.-1) through an Agilent 1200 autosampler, HPLC binary pump was used to deliver the gradient from 3% A to 80% A over 10 min at a flow rate of 250 µl min⁻¹. The LC column was directly connected online to the standard electrospray ionization source of LTQ-Orbitrap XL Fourier transform (FT) mass spectrometer (MS) (Thermo Fisher Scientific, San-Jose, California, United States of America). The optimized parameters, used to prevent in-source fragmentation, included a spray voltage of 4.2 kV, a capillary voltage of -40 V, a tube lens voltage of -50 V, a capillary temperature of 275° C., a sheath flow rate of 40, and an auxiliary gas flow rate of 20. External calibration of mass spectra routinely produced a mass accuracy of less than 3 ppm. All FT mass spectra were acquired at a resolution of 60,000 with 200-2000 Da mass range.

(73) LC/MS Disaccharide Analysis of Human Plasma Samples

(74) Pooled individual plasma samples of healthy control subjects and APAP-ALF patients were digested in 300 µl digestion buffer (50 mM ammonium acetate containing 2 mM calcium chloride adjusted to pH 7.0). Recombinant heparin lyase I, II, III (pH optima 7.0-7.5) and recombinant chondroitin lyase ABC (10 mU each, pH optimum 7.4) were added to each sample and mixed well. The samples were all placed in a water bath at 37° C. for 12 h, after which enzymatic digestion was terminated by removing the enzymes by centrifugation. The filter unit was washed twice with 250 µl distilled water and the filtrates, containing the disaccharide products, were dried by vacuum centrifuge. The dried samples were AMAC-labeled by adding 10 µl of 0.1 M 2-aminoacridone (AMAC) in DMSO/acetic acid (17/3, vol/vol) incubating at room temperature for 10 min, followed by adding 10 µl of 1 M aqueous sodium cyanoborohydride and incubating for 1 h at 45° C. A mixture containing all 17-disaccharide standards purchased from Iduron (UK) prepared at 0.5 ng µl.sup.-1 was similarly AMAC-labeled and used for each run as an external standard. After the AMAC-labeling reaction, the samples were centrifuged, and each supernatant was recovered. LC was performed on an Agilent 1200 LC system at 45° C. using an Agilent Poroshell 120 ECC18 (2.7 µm, 3.0×50 mm) column. Mobile phase A was 50 mM ammonium acetate aqueous solution, and the mobile phase B was methanol. The mobile phase passed through the column at a flow rate of

300 μ l min⁻¹. The gradient was 0-10 min, 5-45% B; 10-10.2 min, 45-100% B; 10.2-14 min, 100% B; 14-22 min, 100-5% B. Injection volume is 5 μ l. A triple quadrupole mass spectrometry system equipped with an ESI source (Thermo Fisher Scientific) was used as a detector. The online MS analysis was at the Multiple Reaction Monitoring (MRM) mode. MS parameters: negative ionization mode with a spray voltage of 3000 V, a vaporizer temperature of 300° C., and a capillary temperature of 270° C.

(75) NMR Analysis

(76) The NMR spectra of 18-mer and 18-mer AXa were obtained on Bruker 800 MHz standard-bore NMR spectrometer with TopSpin 2.1.6 software (Bruker, Billerica, Massachusetts, United States of America). Samples (3.0 to 6.0 mg) were each dissolved in 0.4 ml of 99.9% D₂O centrifuged at 5000 $\times$ g for 1 min and lyophilized. The process was repeated twice and the final sample was dissolved in 0.45 mL of 99.99% D₂O. ¹H spectroscopy, ¹H-¹H correlated spectroscopy (COSY), ¹H-¹³C heteronuclear single quantum coherence spectroscopy (HSQC), ¹H-¹H total correlation spectroscopy (TOCSY) and ¹H-¹H nuclear Overhauser effect spectroscopy (NOESY) experiments were all carried out at 298 K.

(77) Expression of HMGB1

(78) The complete open reading frame of human HMGB1 was cloned into pcDNA3.1(+)-C-6His (GenScript, Piscataway, New Jersey, United States of America). Transfection was performed using FectoPRO transfection reagent (Polyplus transfection). Recombinant HMGB1-his was produced in 293-freestyle cells (Thermo Fisher Scientific) at 31° C. Purification of HMGB1-his from cell lysate was carried out using Ni Sepharose™ 6 Fast Flow gel (GE Healthcare, Little Chalfont, England) followed by Superdex200 gel filtration chromatography. After purification, HMGB1-his was >99% pure as determined by SDS-PAGE followed by silver staining. Endotoxin removal was performed using Detoxi-Gel (Thermo Fisher Scientific) and the final endotoxin level was <0.1 EU/ μ g protein by a Chromogenic LAL Endotoxin Assay (GenScript, Piscataway, New Jersey, United States of America). The recombinant HMGB1-his had both cytokine and chemokine activities, which were assessed by a cell signaling assay and an air pouch assay (described in detail below), respectively.

(79) Removal of Endotoxin from HS Oligosaccharides

(80) Endotoxin was removed from HS oligosaccharides by using a 50 ml centrifugal filter unit (Amicon Ultra-15, Ultracel-100k; Merck Millipore, Darmstadt, Germany) at 4,000 rpm for 30 min. The process was repeated three times by refilling the filter insert with 1 ml of endotoxin-free water each time. The filtered solution was collected. The level of endotoxin was measured using the Limulus Amebocyte Lysate (LAL) kit (Associated of Cape Cod Inc.). The LAL test was performed by adding 100 μ l of reconstituted Pyrotell (sensitivity 0.03 endotoxin units ml⁻¹) to 100 μ l of 0.1 mg ml⁻¹ HS oligosaccharides in sterile saline which is the concentration used for animal injections. The Pyrotell and HS oligosaccharides were added to a 10 mm $\times$ 75 mm depyrogenated glass reaction tube (Associates of Cape Cod Inc.) and incubated at 37° C. for 1 h. At the end of the incubation time, the tubes were inverted. If a gel clot forms, the sample is positive for endotoxin. If the sample remains in solution, the negative result implies that any endotoxin in the sample is below the sensitivity of Pyrotell. All HS oligosaccharides solutions tested negative.

(81) Histology/Immunohistochemistry

(82) Liver tissues were fixed in 10% neutral buffered formalin for 24 h at room temperature, paraffin-embedded, and sectioned. Liver sections (4 μ m) were stained with hematoxylin-eosin (H&E) or immunostained with monoclonal antibodies anti-neutrophil (Abcam, Ab 2557, NIMP-R14) (Abcam, Cambridge, United Kingdom) or anti-syndecan-1 (StemCell Technologies, 60035, clone 281.2) (StemCell Technologies, Vancouver, Canada) and goat anti-rat biotinylated secondary antibodies (Abcam). For fibrin(ogen) staining, polyclonal anti-fibrin(ogen) (Dako of Agilent, Santa Clara, California, United States of America) was used followed by goat anti-rabbit biotinylated secondary antibody (Sigma). Embedding, sectioning and H&E staining were performed at the

Animal Histopathology and Laboratory Medicine Core Facility at UNC Chapel Hill. H&E analyses were performed by the Translational Pathology Laboratory Core Facility at UNC Chapel Hill using Aperio ImageScope Software (Leica Biosystems, Concord, Canada). IHC images were captured using an HD camera attached to a bright field microscope (Leica DM 1000 LED, Leica Microsystems Inc., IL, USA) and were processed using ImageJ. For neutrophil quantitation, five 100× images were randomly selected for each sample and the average neutrophils/field were reported.

(83) Mouse Models of Inflammation: Peritonitis and Air Pouch

(84) Two models were used to study in vivo neutrophil migration: peritonitis and an air pouch inflammation model. For the peritonitis model, 30 mg of liver lysate was injected into the peritoneal cavity of the mouse in the absence or presence of 20 µg of 18-mer. After 20 h, the mice were euthanized via inhalation of isofluorane and the peritoneal cavity was washed with 10 ml of ice cold PBS. The peritoneal lavage was used to determine neutrophil migration using flow cytometry.

(85) In the air pouch technique, 3 ml of sterile air is injected under the skin on the back of the mouse. Three days later, the pouch is refilled with sterile air. On the sixth day, 5 µg of recombinant HMGB1 in the absence or presence of 22.3 µg of 18-mer were injected into the air pouch. Control mice were injected with 2 mg ml⁻¹ BSA in PBS. After 4 hours, mice were euthanized by isofluorane inhalation and the air pouch was washed with PBS only. The lavage was used to determine neutrophil migration using flow cytometry.

(86) Flow Cytometry

(87) Peritoneal and air pouch lavages were stained with fluorescently labeled antibodies against Ly-6G/Ly-6C (monoclonal anti-mouse RB6-8C5, PE-Cy7, eBioscience) and Cd11b (monoclonal anti-mouse M1/70, FITC, eBioscience). Samples were run on a Stratified S1000Exi flow cytometer.

(88) Immunoblot

(89) Liver lysate was prepared by snap freezing tissue in liquid nitrogen at the time of sacrifice. The tissue was mechanically homogenized in buffer containing 200 mM MES, 500 mM phosphate, and 1 mM EDTA at pH 6 followed by three rounds of freeze thawing. The lysed sample were centrifuged at 10000×g for 15 min at 4° C. Biotinylated HS oligosaccharides (final concentration 15 µM) were mixed with 100 µl of fresh liver lysate (~12.5 mg) and incubated overnight at 4° C. Pierce High Capacity Streptavidin Agarose (Thermo Fisher Scientific) was used to isolate biotinylated HS oligosaccharide bound complexes. After washing with 50 mM MES, 50 mM NaCl pH 6, samples were eluted with LDS buffer. Eluted samples were separated using NuPAGE 4-12% Bis-tris protein gels and assayed using anti-HMGB1 antibody (Abcam, rabbit monoclonal EPR3507) and goat anti-rabbit HRP (Abcam).

(90) Preparation of ³⁵S-Labeled HS and Binding Studies to Recombinant HMGB1

(91) [³⁵S]HS was prepared using [³⁵S]PAPS (4×10⁷ cpm) and N-sulfotransferase (NST) (100 µg ml⁻¹), and HS from bovine kidney in 50 mM MES in a total volume of 2 ml. The [³⁵S]HS (1.35×10⁵ cpm) was incubated with 1 µg of recombinant HMGB1 in 50 mM MES, 70 mM NaCl, 10 mM imidazole at pH 6 and incubated at room temperature for 30 min. 5 µg chromatin immunoprecipitation grade anti-HMGB1 (Abcam) was added and incubated for 1 h at 4° C. Reaction mixtures were purified using Dynabeads Protein A (Thermo) and eluted in a buffer containing 25 mM Tris, 150 mM NaCl at pH 7.5. The eluted samples were measured for [³⁵S] counts using a liquid scintillation analyzer (Packard; GMI, Ramsey, Minnesota, United States of America). [³⁵S]HS in the absence of recombinant HMGB1 serves as a negative control.

(92) Surface Plasmon Resonance

(93) The biotinylated HS 18-mers (18-mer and 18-mer AXa) were immobilized to streptavidin (SA) sensor chips (BIAcore, GE Healthcare, Uppsala, Sweden) based on the manufacturer's protocol. In brief, 20 µl of biotinylated HS oligosaccharides were injected over flow cells 2,3, and 4 (FC2, FC3, and FC4) of the SA chip at a flow rate of 10 µl min⁻¹. The successful immobilization of

biotinylated oligosaccharides were confirmed by the observation of a 563, 823, 505, 553 and 176 resonance unit (RU) increase, respectively on the SA chip. The control flow cell (FC1) was prepared by 1 min injection with saturated biotin.

(94) Recombinant HMGB1 was diluted in 0.01 M HEPES, 0.15 M NaCl, 3 mM EDTA, and 0.005% surfactant P20 at pH 7.4. Recombinant HMGB1 at concentrations of 1000, 500, 250, 125 and 63 nM were injected at a flow rate of 30 μ l min.^{sup.}−1. At the end of the recombinant HMGB1 injection, the same buffer was flowed over the SA surface to facilitate dissociation. After a 3 min dissociation time, the SA surface was regenerated by injecting 30 μ l of 2 M NaCl. The response was monitored as a function of time (sensogram) at 25° C. The sensograms of various recombinant HMGB1 concentrations were globally fitted with 1:1 Langmuir model. SPR measurements were performed on a BIAcore 3000 operated using BIAcore 3000 control and BIAevaluation software (version 4.0.1)

(95) APAP-ALF Patient Plasma

(96) APAP-ALF patient plasmas, containing 1.8 mg ml.^{sup.}−1 EDTA, were obtained from the Acute Liver Failure Study Group (ALFSG) biorepository (of the National Institute of Diabetes and Digestive and Kidney Diseases, Bethesda, Maryland, United States of America). Details on the study design and collection methods are described previously (35). Briefly, starting in 1998, adult patients who met the inclusion and exclusion criteria were enrolled in the ALFSG Registry. Plasma samples were obtained on admission to the Registry. In this study, only the plasma from APAP-overdosed patients was analyzed. Clinical data including the estimated quantity of APAP ingested, estimated time from ingestion to hospitalization, intentionality of overdose and patient demographics (age, gender, race, comorbidities, etc.), were not revealed for this study. Plasma syndecan-1 levels were measured using an ELISA kit (Human syndecan-1 ELISA, CellSciences, Newburyport Massachusetts, United States of America) according to the manufacturer's protocol. Plasma HMGB1 levels were measured using an HMGB1 ELISA according the manufacturer's protocol. Plasma ALT levels were measured using ALT Infinity Reagent.

(97) Determination of the In Vitro and Ex Vivo Anti-FXa Activity

(98) Assays were based on a previously published method (19). Briefly, human FXa (Enzyme Research Laboratories, South Bend, Indiana, United States of America) was diluted to 50 U ml.^{sup.}−1 with PBS. The chromogenic substrate S-2765 (Diapharma, West Chester, Ohio, United States of America) was diluted to 1 mg ml.^{sup.}−1 in water. For in vitro studies, Fondaparinux and HS oligosaccharides (18-, 12-, 6-, and 18-mer AXa) were dissolved in PBS at various concentrations (11-131 nM). 16 μ l of sample was incubated with 60 μ l of 35 μ l ml.^{sup.}−1 antithrombin (Cutter Biologics) for 2 min at room temperature. Next, 100 μ l of FXa was added and incubated for 4 min at room temperature. 30 μ l of S-2765 substrate was added and the absorbance of the reaction mixture was measured at 405 nm continuously for 5 min. PBS serves as a control sample. The maximum slope for each sample was convert to percent FXa activity by dividing by the maximum slope for the control sample.

(99) For ex vivo studies, mouse plasma collected 24 h post-APAP overdose in mice treated with APAP only, 18-mer, and 18-mer AXa were used and the same protocol was following.

(100) Statistical Analysis

(101) All data are expressed as mean $\pm$ SEM. Statistical significance between experimental and control groups were analyzed by two-tailed unpaired Student t test, between multiple groups by one-way ANOVA followed by Dunnett's or Tukey's multiple comparison's test, and Kaplan-Meier survival curves by log-rank test using GraphPad Prism software (version 7.03; GraphPad Software, Inc., LaJolla, California, United States of America).

Example 3

Synthesis of HS 18-mer

(102) The synthesis of 18-mer was completed according to the chemoenzymatic method published previously (17, 31). Briefly, heparosan synthase-2 (PmHS2) from *Pasteurella multocida* was used

to elongate the monosaccharide, GlcA-pNP, to appropriate sized backbones. The backbone was then subjected to the modification of N-sulfotransferase (NST), C.sub.5-epimerase (C.sub.5-epi), and 2-O-sulfotransferase (2-OST). There were five major steps involved in the overall synthesis, including Step a (elongation step to add GlcNTFA), Step b (elongation step to add GlcA), Step c (detrifluoroacetylation using LiOH), step d (N-sulfation step), step e (2-O-sulfation/epimerization) (FIGS. 1A-1G).

(103) Step a was to elongate the oligosaccharide backbone to the desired size, involving the addition of GlcNTFA residue. Briefly, GlcA-pNP (3.2 mM) was dissolved in buffer containing Tris (25 mM, pH 7.2), MnCl.sub.2 (5 mM), pmHS2 (60 µg ml.sup.-1) and UDP-GlcNTFA (4.5 mM), then incubated at 30° C. overnight. The total reaction volume was 4 L. A C.sub.18-column (3×15 cm, or 120 g, Biotage) was used for purification with gradient elution method (0-100% methanol in H.sub.2O, 0.1% trifluoroacetic acid, 5 ml min.sup.-1).

(104) Step b was to elongate with a GlcA residue. A disaccharide (3.2 mM) was dissolved in a buffer containing Tris (25 mM, pH 7.2), MnCl.sub.2 (5 mM), pmHS2 (30 µg ml.sup.-1) and UDP-GlcA (1.5 mM), then incubated at 30° C. overnight. The total reaction volume was 3 L. A C.sub.18-column (3×15 cm, or 120 g, Biotage) was used for purification as described above.

(105) Step c was to convert a GlcNTFA residue to a GlcNH.sub.2 residue under alkaline conditions. The detrifluoroacetylation of oligosaccharide (13 mM) was conducted in 0.1 M LiOH under ice bath for 0.5 h. The products were monitored by electrospray ionization mass spectrometry (ESI-MS). After the reaction was completed, the pH was immediately adjusted to 7.0 using hydrochloric acid (1 M).

(106) Step d was to convert a GlcNH.sub.2 residue to a GlcNS residue using NST enzyme. In one example, the detrifluoroacetylated tetrasaccharide (1.3 mM, GlcNH.sub.2-GlcA-GlcNH.sub.2-GlcA-pNP) incubated with N-sulfotransferase (32 µg ml.sup.-1) and PAPS (1.5 equiv. of free amino group amount) in a solution containing 2-(N-morpholino) ethanesulfonic acid (MES, 50 mM) pH 7.0 at 37° C. overnight in a reaction volume of 5.6 L.

(107) Step e was to convert an internal GlcA residue to an IdoA2S residue and involves both C.sub.5-epimerase and 2-O-sulfotransferase (2-OST). For example, the oligosaccharide GlcNTFA-GICA-GlcNS-GlcA-GlcNS-GlcA-pNP (1.2 mM) was incubated in a solution containing Tris (25 mM) buffer (pH 7.5) and semi-purified C.sub.5-epimerase (3 µg ml.sup.-1), 2-OST (6.5 µg ml.sup.-1) and PAPS (1.8 mM) at 37° C. overnight in a reaction volume of 4.9 L.

(108) The sulfated products were purified using Q-Sepharose fast flow column (from GE Healthcare Life Science) and eluted with a linear gradient 0-100% 2 M NaCl in 20 mM NaOAc-HOAc, pH 5.0 in 3 h. Different sizes of Q-Sepharose columns and salt gradients were chosen based on the binding affinity of the product and reaction scale.

(109) At every elongation synthesis step, the products were monitored by Shimadzu HPLC equipped with a polyamine II column (4.6 mm×250 mm, from YMC). The structures of the intermediates from each step were characterized by electrospray ionization mass spectrometry (ESI-MS).

Example 4

Mouse Model of APAP Liver Injury

(110) All animal experiments were approved by the Institutional Animal Care and Use Committee of University of North Carolina at Chapel Hill (Chapel Hill, North Carolina, United States of America) and the IACUC of the University at Buffalo (Buffalo, New York, United States of America). Ager.sup./- mice were originally gifted by Dr. Angelika Bierhaus (University of Heidelberg, Heidelberg, Germany) (32). C57BL/6J or Ager.sup./- mice were fasted overnight (12-15 h) to deplete glutathione stores before APAP (Sigma) administration. Fresh APAP was dissolved in warm (~50° C.) sterile 0.9% sodium chloride solution (sterile saline), cooled to 37° C., and injected intraperitoneal at 400 or 600 mg kg.sup.-1. In some experiments, mice were injected subcutaneously at 30 min post-APAP with 9.5 µM HS oligosaccharide in ~200 µl sterile saline and

again at 12 hours post-APAP with 4.75 μ M HS oligosaccharide in ~100 μ l or equivalent volumes of sterile saline.

(111) To compare the 18-mer's effectiveness to N-acetyl cysteine (NAC), 300 mg kg.sup.-1 NAC at pH 7.5 in sterile saline was injected intraperitoneal 30 min post-APAP. In a separate experiment, 300 mg kg.sup.-1 NAC or 0.34 mg/kg 18-mer were administered 6 h post-APAP, then at 18 h post-APAP, 0.17 mg kg.sup.-1 18-mer was administered to the 18-mer treated group. At 24 h post-APAP, mice were euthanized and blood and liver tissue were collected.

(112) In the survival study (600 mg kg.sup.-1 APAP), mice were injected 30 min post-APAP with 0.4 mg kg.sup.-1 18-mer followed by repeat injections every 12 h or equivalent volumes of sterile saline for 96 h.

Example 5

Evaluation of APAP-Induced Liver Injury

(113) Plasma ALT was measured using the ALT Infinity reagent (Thermo Fisher Scientific) following the manufacturer's instructions. Plasma TNF- α was measured using Mouse TNF- α DuoSet Kit (R&D Systems, Minneapolis, Minnesota, United States of America) according to the manufacturer's instructions. Plasma HMGB1 levels were determined using HMGB1 ELISA Kit (Tecan US of Tecan Trading AG, Mannedorf, Switzerland) according to the manufacturer's instructions. Plasma syndecan-1 levels were determined using Mouse Syndecan-1 ELISA (CellSciences) according to manufacturer's instructions. Hepatic GSH levels were determined using Glutathione Assay Kit (Cayman Chemicals, Ann Arbor, Michigan, United States of America) according to the manufacturer's instructions.

Example 6

18-Mers Significantly Decrease Liver Injury 36 Hours after APAP Overdose

(114) Similar to the above examples, C57BI/6J male mice, at about 10 weeks of age and about 25 grams of body weight, were overdosed with acetaminophen (APAP; 400 mg/kg) via intraperitoneal injection at time 0 hr.

(115) Mice received repeat subcutaneous injections of 18-mers 30 minutes, 12 hours and 24 hours after APAP. The mice were sacrificed via cervical dislocation, blood was drawn from the vena cava, and liver tissue was harvested for histology. Alanine aminotransferase (ALT) was used as a biomarker for liver injury. Tumor necrosis factor- α (TNF- α) was used as a marker for inflammation. Hematoxylin and Eosin staining (H&E) was used as the histological staining technique to quantitate regions of tissue necrosis.

(116) The resulting data showed that all 18-mer compounds (Table 1) decreased plasma ALT levels. Although all 18-mers tested were effective, the NS2S sulfation pattern provided the most consistent decrease in ALT (largest difference of means compared to APAP control), results of which are presented in Table 1.

(117) TABLE-US-00001 TABLE 1 Difference of Comparison N APAP; N = 19 Means P 18mer NS2S 17 2928 <0.001 18mer NS6S 17 2261 <0.001 18mer NS6S2S 14 1779 <0.001 18mer NS 12 1682 <0.001 18mer NAc 7 1356 0.014

Example 7

Results and Discussion

(118) HS isolated from natural sources is highly complex mixtures with different polysaccharide chain lengths and sulfation patterns. Lack of structurally homogeneous HS oligosaccharides, especially long HS oligosaccharides that display similar functions of full length HS, hampers the efforts to exploit the interests of HS as a therapeutic agent (Liu, 2014). We have recently developed a chemoenzymatic method to synthesize HS oligosaccharides with excellent efficiency (Xu, 2011, Xu 2014, Xu, 2017). Here, HS oligosaccharides, including 323 mg of an HS octadecasaccharide (18-mer), were synthesized (FIGS. 1A-1G and FIG. 2). This represents one of the longest HS oligosaccharides synthesized to date. The structure of 18-mer was confirmed by high resolution mass spectrometry and NMR. The purity was determined to be >98% by high resolution anion

exchange high-performance liquid chromatography.

(119) The hepatoprotective effects of 18-mer were examined in an APAP-induced ALF murine model. Mice treated with 18-mer after APAP overdose had significantly healthier livers than APAP control mice as indicated by a higher population of normal hepatocytes (FIG. 3A) and a lower plasma level of alanine aminotransferase (ALT), a biomarker of liver damage (FIG. 3C). 18-mer treatment showed reduced neutrophil infiltration into the liver and decreased plasma levels of tissue necrosis factor α (TNF- α) (FIGS. 3B and 6A), suggesting attenuation of local and systemic inflammation. 18-mer also decreased mortality following a lethal APAP overdose (600 mg kg.sup.-1), resulting in 90% survival at 96 hours compared to 42% survival in the control group (FIG. 3D). Hepatic glutathione (GSH) levels in APAP control and 18-mer treated mice were essentially the same during the course of the studies (FIG. 6B). GSH levels decrease as NAPQI forms (Tacke, 2015; Nam, 2017), therefore the data suggest that 18-mer does not affect the metabolic conversion of APAP to the cytotoxic intermediate, NAPQI. These results demonstrate that 18-mer ameliorates liver injury by reducing the inflammatory responses rather than by affecting APAP metabolism.

(120) Two lines of evidence suggest that 18-mer targets HMGB1 to reduce inflammation after APAP overdose. First, we discovered that 18-mer diminishes the HMGB1-mediated neutrophil infiltration in two in vivo models. In an air pouch model, injection of recombinant HMGB1 induced extensive neutrophil infiltration (FIG. 4A), an effect that was significantly reduced by co-administration of 18-mer. 18-mer also reduced neutrophil infiltration in a peritonitis model induced by liver lysate (FIG. 6C), a process known to be mediated by HMGB1(4). Second, RAGE knockout mice, or Ager.sup./- mice, were used to demonstrate that 18-mer targets the HMGB1/RAGE axis. Because the interaction of HMGB1 and RAGE is essential for the pro-inflammatory response in APAP overdose (Huebener, 2015), the hepatoprotective effect of 18-mer is expected to be dependent on the presence of RAGE. Hepatic neutrophil infiltration was increased, and the increase was significantly greater in wild type (WT) than Ager.sup./- mice after APAP overdose (FIG. 4B). 18-mer treated Ager.sup./- mice failed to show a decrease in neutrophil infiltration, while WT mice responded to 18-mer treatment (FIG. 4B). Furthermore, 18-mer treatment in Ager.sup./- mice was incapable of lowering the ALT level (FIG. 6D). This data suggest that 18-mer lost its protective effect in the absence of RAGE.

(121) Structure-activity relationship studies were conducted to examine binding to HMGB1 and protection in the APAP model using different forms of HS. Four biotinylated oligosaccharides were synthesized, including 6-, 12-, and two 18-mers (FIG. 4E) (for structures and characterization see FIG. 7). Among these, 6-, 12- and 18-mer varied only in chain length but not in sulfation pattern. These oligosaccharides are non-anticoagulant because they lack anti-factor Xa activity. 18-mer AXa is a highly sulfated anticoagulant octadecasaccharide that possesses potent anti-factor Xa activity (FIGS. 8A and 8B). The 18-mer and 18-mer AXa, but not the 6-mer or 12-mer, were observed to bind HMGB1 from mouse liver lysate, suggesting that there is a minimum chain length requirement for binding (FIG. 4C). The HMGB1-binding constants (KD) were determined to be 186 nM (for 18-mer) and 65 nM (18-mer AXa) by surface plasmon resonance, yet only the 18-mer displayed the hepatoprotective effects as measured by ALT (FIG. 4D) and by neutrophil infiltration (FIG. 8C). It is apparent that the lack of hepatoprotection from the 6- and 12-mer correlates with the inability to bind HMGB1. However, the 18-mer AXa, while capable of binding HMGB1, also lacked hepatoprotection. This may be due to its anticoagulant activity since anticoagulant unfractionated heparin also lacked hepatoprotection after APAP-overdose (FIG. 8D). Kopeck and colleagues have reported that fibrin is required to activate liver repair after APAP overdose (20). Administering 18-mer AXa reduces fibrin formation, and thus, results in loss of hepatoprotection.

(122) To further understand the protective role of 18-mer, the role of syndecan-1, a major HS proteoglycan present on hepatocytes, was investigated in response to APAP overdose. Syndecan-1 comprises a core protein attached with HS chains and is shed from the cell surface by matrix metalloproteases in pathological conditions (Park, 2000). Higher but fluctuating levels of plasma

syndecan-1 were observed over a period of 24 hours after APAP overdose in mice (FIG. 5A). Concurrently, plasma levels of HMGB1 (FIG. 5A), ALT and hepatic neutrophil infiltration increased over time (FIGS. 9A and 9B). Loss of cell surface syndecan-1 was confirmed by immunostaining liver section. Using 35S-labeled HS from bovine kidney as a surrogate molecule for the HS chains on shed syndecan-1, we demonstrated that HMGB1 binds to HS (FIG. 5B), which is consistent with a previous report using HS from different sources (9). These results suggest that shed syndecan-1 binds to HMGB1 through its HS chains. A significant increase in plasma syndecan-1 levels was also observed from patients with APAP-induced ALF (FIG. 5C). Notably, shed syndecan-1 levels in APAP-ALF patients were about 200-fold higher than in APAP overdosed mice as measured by core protein analysis. HS chain analysis confirmed the heightened levels of shed syndecan-1 in ALF patients (FIG. 10A). HMGB1 and ALT in APAP-ALF patients were also higher than the healthy control group (FIGS. 10B and 10C), consistent with a previous report (22).

(123) The relationship between syndecan-1 shedding and HMGB1 release in humans and mice after APAP overdose underscores an endogenous protective pathway. Syndecan-1 sheds after the initial insult and neutralizes the pro-inflammatory activities of HMGB1, thereby limiting sterile inflammation. Indeed, syndecan-1^{sup.-/-} mice are more susceptible to APAP-induced ALF than WT animals (Nam, 2017). During extensive APAP damage, it is likely that shed syndecan-1 is inadequate to neutralize all HMGB1, and the addition of 18-mer, a mimetic of HS on syndecan-1, provides further protection.

(124) Since sterile inflammation occurs after NAPQI-induced cell damage, we examined the possibility of further delaying 18-mer treatment after APAP overdose. N-acetyl-cysteine (NAC) is a NAPQI-neutralizing antioxidant that is the standard of care for the treatment of APAP overdose. However, NAC treatment is only effective if given within 8 hours of APAP ingestion (Bailey, 2016). While NAC loses its protective effect when administered at six hours post-APAP in mice, the 18-mer is still able to decrease ALT (FIG. 5D). These data suggest that 18-mer treatment has a potential advantage for late-presenting APAP overdose patients by providing a wider therapeutic window.

Example 8

Conclusions

(125) The use of synthetic HS to treat APAP-induced ALF by targeting HMGB1-mediated sterile inflammation is disclosed herein. The availability of homogeneous oligosaccharides has made it possible to identify candidate targets and compounds underlying this effect. In addition to neutralizing HMGB1, HS of syndecan-1 can also activate liver repair and modulate the activity of chemokines. 18-mer could also contribute to these functions. HS is well-tolerated by patients (Shriver, 2004) and unfractionated heparin, a highly sulfated form of HS, has been used as an anticoagulant for nearly a century (Szajek, 2016). Now, the synthesis of HS oligosaccharides can be achieved on a large scale and in a cost-effective manner. HMGB1-neutralizing synthetic oligosaccharides should offer a promising therapeutic approach for ALF. Because HMGB1 has been implicated in diverse disease states including cancers, stroke and arthritis (Venereau, 2016), novel HS-based therapeutics provide promising opportunities as inhibitors of HMGB1.

REFERENCES

(126) All references listed herein including but not limited to all patents, patent applications and publications thereof, scientific journal articles, and database entries (e.g., GENBANK® database entries and all annotations available therein) are incorporated herein by reference in their entirety to the extent that they supplement, explain, provide a background for, or teach methodology, techniques, and/or compositions employed herein. 1. L. Zitvogel, O. Kepp, G. Kroemer, Decoding cell death signals in inflammation and immunity. *Cell* 140, 798-804 (2010). 2. G. Y. Chen, G. Nuñez, Sterile inflammation: sensing and reacting to damage. *Nat. Immunol.* 10, 826-837 (2010). 3. M. E. Bianchi et al., High-mobility group box 1 protein orchestrates responses to tissue damage

via inflammation, innate and adaptive immunity, and tissue repair. *Immunol. Rev.* 280, 74-82 (2017). 4. P. Huebener et al., The HMGB1/RAGE axis triggers neutrophil-mediated injury amplification following necrosis. *J. Clin. Invest.* 125, 539-550 (2015). 5. K. J. Heard, Acetylcysteine for acetaminophen poisoning. *N. Eng. J. Med.* 359, 285-292 (2008). 6. W. M. Lee, Acetaminophen toxicity: changing perceptions on a social/medical issue. *Hepatology* 46, 966-970 (2007). 7. M. Blieden, L. C. paramore, D. Shah, R. Ben-Joseph, A perspective on the epidemiology of acetaminophen exposre and toxicity in the United States. *Expert Rev. Clin. Pharmacol.* 7, 341-348 (2014). 8. F. Tacke, J. C. Mossanen, Acetaminophen-induced acute liver injury in mice. *Lab. Anim.* 49, 30-36 (2015). 9. D. Xu, J. Young, D. Song, J. D. Esko, Heparan sulfate is essential for high mobility group protein 1 (HMGB1) signaling by the receptor for advanced glycation end products (RAGE). *J. Biol. Chem.* 286, 41736-41744 (2011). 10. A. E. Proudfoot et al., Glycosaminoglycan binding and oligomerization are essential for the in vivo activity of certain chemokines. *Proc. Natl. Acad. Sci. USA* 100, 1885-1890 (2003). 11. L. Wang, M. Fuster, P. Sriramaraao, J. D. Esko, Endothelial heparan sulfate deficiency impairs L-selectin- and chemokine-mediated neutrophil trafficking during inflammatory responses. *Nat. Immunol.* 6, 902-910 (2005). 12. J. Axelsson et al., Inactivation of heparan sulfate 2-O-sulfotransferase accentuates neutrophil infiltration during acute inflammation in mice. *Blood* 120, 1742-1751 (2013). 13. M. Sarris et al., Inflammatory chemokines direct and restrict leukocyte migration within live tissues as glycan-bound gradients. *Curr. Biol.* 22, 2375-2382 (2012). 14. C. Gama et al., Sulfation patterns of glycosaminoglycans encode molecular recognition and activity. *Nat. Chem. Biol.* 2, 467-473 (2006). 15. J. Liu, R. J. Linhardt Chemoenzymatic synthesis of heparan sulfate and heparin. *Nat. Prod. Rep.* 31, 1676-1685 (2014). 16. Y. Xu et al., Chemoenzymatic synthesis of homogeneous ultra-low molecular weight heparin. *Science* 334, 498-501 (2011). 17. Y. Xu et al., Homogeneous and reversible low-molecular weight heparins with reversible anticoagulant activity. *Nat. Chem. Biol.* 10, 248-250 (2014). 18. Y. Xu et al., Synthetic oligosaccharides can replace animal-sourced low-molecular weight heparins. *Sci. Transl. Med.* 9, eaan5954 (2017). 19. E. J. Nam et al., Syndecan-1 Limits the Progression of Liver Injury and Promotes Liver Repair in Acetaminophen-Induced Liver Injury. *Hepatology*, doi: 10.1002/hep.29265 (2017). 20. A. Kopec et al., Fibrin(ogen) drives repair after acetaminophen-induced liver injury via leukocyte α 5 β 1 integrin-dependent upregulation of Mmp12. *J. Hepatol.* 66, 787-797 (2017). 21. P. W. Park, O. Reizes, M. Bernfield, Cell surface heparan sulfate proteoglycans: selective regulators of ligand-receptor encounters. *J. Biol. Chem.* 275, 29923-29926 (2000). 22. D. J. Antoine et al., Mechanistic biomarkers provide early and sensitive detection of acetaminophen-induced acute liver injury at first presentation to hospital. *Hepatology* 58, 777-787 (2013). 23. G. P. Bailey et al., Delays during the administration of acetylcysteine for the treatment of paracetamol overdose. *Br. J. Clin. Pharmacol.* 62, 1358-1363 (2016). 24. K. C. Wildhagen et al., Nonanticoagulant heparin prevents histone-mediated cytotoxicity in vitro and improves survival in sepsis. *Blood* 123, 1098-1101 (2014). 25. Y. Monneau, F. Arenzana-Seisdedos, H. Lortat-Jacob, The sweet spot: how GAGs help chemokines guide migrating cells. *J. Leukoc. Biol.* 99, 935-953 (2016). 26. P. Lundback et al., A novel high mobility group box 1 neutralizing chimeric antibody attenuates drug-induced liver injury and postinjury inflammation in mice. *Hepatology* 64, 1699-1710 (2016). 27. T. Yamamoto, Y. Tajima, HMGB1 is a promising therapeutic target for acute liver failure. *Expert Rev. Gastroenterol. Hepatol.* 11, 673-682 (2017). 28. Z. Shriver, S. Raguram, R. Sasisekharan, Glycomics: A pathway to a class of new and improved therapeutics. *Nat Rev Drug Discov* 3, 863-873 (2004). 29. A. Szajek et al., The US regulatory and pharmacopeia responses to the global heparin contamination crisis. *Nat. Biotechnol.* 34, 625-630 (2016). 30. E. Venereau et al., HMGB1 as biomarker and drug target. *Pharmacol. Res.* 111, 534-544 (2016). 31. P. Hsieh, Y. Xu, D. A. Keire, J. Liu, Chemoenzymatic synthesis and structural characterization of 2-O-sulfated glucuronic acid containing heparan sulfate hexasaccharides. *Glycobiology* 24, 681-692 (2014). 32. B. Liliensiek et al., Receptor for advanced glycation end products (RAGE) regulates sepsis but not the adaptive immune response. *J. Clin. Invest.* 113, 1641-1650 (2004). 33. PCT Publication No.

WO2012/088416, published Jun. 28, 2012 34. Published U.S. Patent Application No. US2013/0296540, published Nov. 7, 2013 35. PCT Publication No. WO2014204929, published Dec. 24, 2014 36. Published U.S. Patent Application No. US2016/0122446, published May 5, 2016 37. Xu, Y., Chandarajoti, K., Zhang, X., Pagadala, V., Dou, W., Hoppensteadt, D. M., Sparkenbaugh, E., Colley, B., Daily, S., Key, N., Severynse-Stevens, D., Fareed, J., Linhardt, R. J., Pawlinski, R., and Liu, J. (2017) Synthetic oligosaccharides can replace animal-sourced low-molecular weight heparins *Sci. Transl. Med.*, 9, ean5954. 38. Xu, Y, Cai, C., Chandarajoti, K, Li, L., Hsieh, P., Pham, T., Sparkenbaugh, E. M., Sheng, J., Key, N., Pawlinski, R., Harris, E., Linhardt, R. J., and Liu, J. (2014) Homogeneous and reversible low-molecular weight heparins with reversible anticoagulant activity. *Nat. Chem Biol.* 10: 248-250. 39. Renpeng Liu, Yongmei Xu, Miao Chen, Michel Weïwer, Xianxuan Zhou, Arlene S. Bridges, Paul L. DeAngelis, Qisheng Zhang, Robert J. Linhardt and Jian Liu (2010) Chemoenzymatic design of heparan sulfate oligosaccharides *J. Biol. Chem.* 285:34240-34249 40. Xu, D., Moon, A. F., Song, D., Pedersen, L. C. and Liu, J. (2008) Engineering the sulfotransferases to modify heparan sulfate *Nat. Chem. Biol.* 4: 200-202.

(127) It will be understood that various details of the presently disclosed subject matter may be changed without departing from the scope of the presently disclosed subject matter. Furthermore, the foregoing description is for the purpose of illustration only, and not for the purpose of limitation.

Claims

1. A non-anticoagulant heparan sulfate oligosaccharide molecule having an anti-inflammatory property, wherein the oligosaccharide molecule comprises one of the following structures:
##STR00013## ##STR00014## ##STR00015## wherein R^{sup.3} is H or a detectable tag.
 2. The oligosaccharide molecule of claim 1, wherein the oligosaccharide molecule is provided in a composition further comprising a compound comprising a disaccharide structure unit as shown:
##STR00016## wherein R=−H or −SO₃H.
 3. The oligosaccharide molecule of claim 2, wherein the compound comprising a disaccharide structure unit is selected from the group consisting of non-anticoagulant heparin, non-anticoagulant low-molecular weight heparin, and O-desulfated heparin (ODSH).
 4. The oligosaccharide molecule of claim 1, wherein the oligosaccharide molecule protects against liver injury in vivo.
 5. The oligosaccharide molecule of claim 1, wherein the oligosaccharide molecule decreases neutrophil infiltration in vivo.
 6. The oligosaccharide molecule claim 1, wherein the oligosaccharide molecule decreases inflammation in vivo.
 7. The oligosaccharide molecule of claim 1, wherein R^{sup.3} in each structure is a detectable tag.
 8. The oligosaccharide molecule of claim 1, wherein the detectable tag in each structure is p-nitrophenyl.
-