



US012397051B2

(12) **United States Patent**
Nag et al.

(10) **Patent No.: US 12,397,051 B2**

(45) **Date of Patent: Aug. 26, 2025**

(54) **VACCINE FOR USE AGAINST
 CORONAVIRUS AND VARIANTS THEREOF**

FOREIGN PATENT DOCUMENTS

WO WO-2021226436 A1 * 11/2021 A61K 39/215

(71) Applicant: **Globe Biotech Limited**, Dacca (BD)

OTHER PUBLICATIONS

(72) Inventors: **Kakon Nag**, Dacca (BD); **Juwel Chandra Baray**, Dacca (BD); **Maksudur Rahman Khan**, Dacca (BD); **Asif Mahmud**, Dacca (BD); **Enamul Haq Sarker**, Dacca (BD); **Samir Kumar**, Dacca (BD); **Jikrul Islam**, Dacca (BD); **Rony Roy**, Dacca (BD); **Mohammad Mohiuddin**, Dacca (BD); **Naznin Sultana**, Dacca (BD)

Nag K, Chandra Baray J, Rahman Khan M, Mahmud A, Islam J, Myti S, Ali R, Haq Sarker E, Kumar S, Hossain Chowdhury M, Roy R, Islam F, et al. An mRNA-based vaccine candidate against SARS-CoV-2 elicits stable immuno-response with single dose. Vaccine. Jun. 23, 2021;39(28):3745-3755. Epub May 18, 2021. (Year: 2021).*

De Breyne S, Vindry C, Guillin O, Condé L, Mure F, Gruffat H, Chavatte L, Ohlmann T. Translational control of coronaviruses. Nucleic Acids Res. Dec. 16, 2020;48(22):12502-12522. (Year: 2020).*

Park JW, Lagniton PNP, Liu Y, Xu RH. mRNA vaccines for COVID-19: what, why and how. Int J Biol Sci. Apr. 10, 2021;17(6):1446-1460. doi: 10.7150/ijbs.59233. PMID: 33907508; PMCID: PMC8071766. (Year: 2021).*

Granados-Riveron JT, Aquino-Jarquín G. Engineering of the current nucleoside-modified mRNA-LNP vaccines against SARS-CoV-2. Biomed Pharmacother. Oct. 2021;142:111953. doi: 10.1016/j.biopha.2021.111953. Epub Jul. 23, 2021. PMID: 34343897; PMCID: PMC8299225. (Year: 2021).*

Zhang Y, Zhang T, Fang Y, Liu J, Ye Q, Ding L. SARS-CoV-2 spike L452R mutation increases Omicron variant fusogenicity and infectivity as well as host glycolysis. Signal Transduct Target Ther. Mar. 9, 2022;7(1):76. (Year: 2022).*

Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG, Zhang X, Muruato AE, Zou J, Fontes-Garfias CR, Mirchandani D, Scharton D, Bilello JP, et al. Spike mutation D614G alters SARS-CoV-2 fitness. Nature. Apr. 2021;592(7852):116-121. Epub Oct. 26, 2021. Erratum in: Nature. Jul. 2021;595(7865): E1. (Year: 2020).*

Kumar R, Srivastava Y, Muthuramalingam P, Singh SK, Verma G, Tiwari S, Tandel N, Beura SK, Panigrahi AR, Maji S, Sharma P, Rai PK, Prajapati DK, Shin H, Tyagi RK. Understanding Mutations in Human SARS-CoV-2 Spike Glycoprotein: A Systematic Review & Meta-Analysis. Viruses. Mar. 27, 2023;15(4):856. (Year: 2023).*

VanInsberghe D, Neish AS, Lowen AC, Koelle K. Recombinant SARS-COV-2 genomes are currently circulating at low levels. bioRxiv [Preprint]. Mar. 15, 2021:2020.08.05.238386. Update in: Virus Evol. Jul. 15, 2021;7(2):veab059. (Year: 2021).*

Amendola A, et al. Surface glycoprotein, partial [Severe acute respiratory syndrome coronavirus 2]. GenBank: QVK80293.1., Aug. 9, 2021. (Year: 2021).*

* cited by examiner

Primary Examiner — Rachel B Gill

(74) Attorney, Agent, or Firm — Husch Blackwell LLP

(73) Assignee: **Globe Biotech Limited**, Dacca (BD)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 165 days.

(21) Appl. No.: **17/475,001**

(22) Filed: **Sep. 14, 2021**

(65) **Prior Publication Data**

US 2023/0084012 A1 Mar. 16, 2023

(51) **Int. Cl.**

A61K 39/215 (2006.01)

A61K 9/1272 (2025.01)

A61K 9/51 (2006.01)

A61K 39/00 (2006.01)

A61K 48/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/215** (2013.01); **A61K 9/1272** (2013.01); **A61K 9/5123** (2013.01); **A61K 39/00** (2013.01); **A61K 48/0066** (2013.01); **A61K 48/0075** (2013.01); **A61K 2039/55555** (2013.01)

(58) **Field of Classification Search**

CPC .. **A61K 39/215**; **A61K 9/1272**; **A61K 9/5123**; **A61K 39/00**; **A61K 48/0066**; **A61K 48/0075**; **A61K 2039/55555**; **A61K 2039/51**; **A61K 2039/57**; **A61K 2039/575**; **A61K 9/127**; **A61K 39/12**; **A61P 31/14**; **C07K 14/005**; **C12N 15/88**; **C12N 2770/20022**; **C12N 2770/20034**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2002/0123099 A1 * 9/2002 Weiner A61P 9/00 702/19
 2022/0202930 A1 * 6/2022 Roth A61K 9/5123
 2022/0370599 A1 * 11/2022 Abshire A61P 31/16
 2022/0381780 A1 * 12/2022 Wohlstadter G16B 25/10
 2022/0401550 A1 * 12/2022 Simon-Lorieri C07K 14/005
 2023/0285539 A1 * 9/2023 Joyce A61K 39/12
 2023/0346921 A1 * 11/2023 Wei A61P 31/14
 2024/0100151 A1 * 3/2024 Carfi A61K 39/215
 2024/0139309 A1 * 5/2024 Carfi C07K 14/005
 2024/0156946 A1 * 5/2024 Roth A61K 9/5123
 2024/0342269 A1 * 10/2024 Dias A61K 9/5123

(57) **ABSTRACT**

A novel transgene for use to produce a coronavirus vaccine is provided. The transgene encodes: i) an RNA polymerase promoter; ii) a 5' UTR; iii) a secretory sequence; iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant sequence at amino acid position 614 of a native spike protein; and v) a 3' UTR and poly A sequence. A vaccine is also provided comprising the transgene or an mRNA transcript thereof.

30 Claims, 12 Drawing Sheets

Specification includes a Sequence Listing.




FIG. 2A




FIG. 2B




FIG. 2C (lgG2a + lgG2b) : (lgG1 + lgG3)




FIG. 2D





FIG. 2E






FIG. 4A




FIG. 4B





FIG. 4C Pseudovirus neutralization



Neutralization assay

FIG. 4D




FIG. 4E Neutralization assay




FIG. 4F




FIG. 4G




FIG. 4H




FIG. 4I




FIG. 4J




FIG. 4K




FIG. 5A)

148	cagtgtgttaactcttacaaccagaactcaattacccccgcatacactaaltcttccacacgtggtgttt	217
218	attacctgacaaaagtcttcagatcctcagttttacattcaactcaggactgttcttacctttctttc	287
288	caatgttacttggttccatgctatacatgtctctgggaccaatggtactaagagggttgataacctgtc	357
358	ctaccatttaattgatgggtgtttattttgcttccactgagaagtcataacataataagaggctggattttg	427
428	gtactactttagattcgaagaccagtcctctacttattgttaataacgctactaatgttgattaaagt	497
498	ctgtgaatttcaatttgaatgatccattttgggtgtttattaccacaaaaacaacaaagtggatg	567
568	gaaagtgagttcagagtttattctagtgcgaataattgcacttttgaatatgtctctcagccctttctta	637
638	tggacctgaaggaaaacagggttaattcaaaaatcttagggaattgtgtttaa gaatattgatggta	707
708	ttttaaataatatttaagcacacgcciatatttaagtgcgtgatctccctcagggttttccggcttla	777
778	gaaccattggtagatttccaatagggtatttaacatcactagggttcaaaactttactgtttacatagaa	847
848	gttatttgactcctgggtattctcttcagggttgacagctgggtgctgcagcttattatgtgggttatct	917
918	tcaacctaggacttttctataaaatataatgaaaatggaaccattacagatgctgtagactgtgcactt	987
988	gacctctctcagaaacaaagtgtacgttgaaatccttcactgtagaaaaaggaatctatcaaaacttcta	1057
1058	actttagagtcacacacagaatctattgttagatttctaataattacaaactgtgccccttttggtga	1127
1128	agtttttaacgccaccagatttgcactctgtttatgcttggacaggagaatcagcaactgtgttgc	1197
1198	gattattctgtcctataaattccgcacattttccacttttaagtgttatggagtgtctcctactaaat	1267
1268	taaatgatctctgtcttactaatgtctatgcagattcatttgaattagagggtgatgaagtcagacaaat	1337
1338	cgctccaggggcaaacgtgaaagattgctgattataattataaattaccagatgattttacaggctgcgtt	1407
1408	atagcttggaaattcaacaacttgaattctaaaggttgggtgtaattataaattacgtatagatttga	1477
1478	ggaaagtctaattcaaaccttttgagagagatattcaactgaaatctatcaggccggtagcacaccttg	1547
1548	taatgggtgtgaagggttttaattgttactttccttacaatcatatggtttccaaccactaatggtgtt	1617
1618	ggttaccaaccatacagagtagtagtactttcttgaacttctacatgcaccagcaactgtttgtggac	1687
1688	ctaaaaagtctactaatttgggttaaaaacaaatgtgtcaatttcaacttcaatggttlaacaggcacagg	1757
1758	tgttcttactgaagtcaaaaaagtttctgcctttccaacaatttggcagagacattgctgacactact	1827
1828	gatgctgtccgtgatccacagacacttgagattcttgacattacacatgttctttgggtggtgacag	1897
1898	ttataacaccagggaacaaataacttctaaccagggtgctgttctttatcagggtgttaactgcacagaagt	1967
1968	ccctgttctatitcatgcagatcaacttactcctacttggcgtgtttattctacagggttcaatgtttt	2037
2038	caaacacgtgcaggctgtttaaaggggctgaacatgtcaacaactcatatgagtgtgacataccattg	2107
2108	gtgcagggtatattgcgctagttatcagactcagactaattctcctcggcgggcacgtagtgtagctagta	2177
2178	atccatcattgcttacaactatgtcacttgggtcagaaaattcagttgcttacttaataactctattgcc	2247
2248	atacccacaaattttactattagtgttaccacagaaattctaccagtgtctatgaccaagacatcagtag	2317
2318	attgtacaatgtacatttgggtgattcaactgaatgcagcaactctttgttgcaatatggcagttttg	2387
2388	tacacaattaaaccgtgctttaaactggaatagctgttgaacaagacaaaaacaccaagaagttttgca	2457
2458	caagtcaaaacaaatttacaacacaccaccaattaaagattttgggtggttttaatttttcaaaatattac	2527
2528	cagatccatcaaaaaccaagcaagaggtcatttattgaagatctacttttcaacaaagtgcacttgcaga	2597
2598	tgtgtgcttcatcaacaataatgggtgattgccttgggtgatattgctgctagagacctcatitgtgcacaa	2667
2668	aagttttaacggccttactgtttgccaccttggctcagatgaaatgattgctcaatacaacttctgcac	2737
2738	tgttagcgggtacaateacttctgggtggaccttgggtgcagggtgctgcattacaataaccatttctat	2807
2808	gcaaatggccttataggttaattggattggagtacacagaatgttctctatgagaacaaaaattgatt	2877
2878	gccaaaccaatttaatagtctattggcaaaattcaagactcactttcttcacagcaagtgcacttggaa	2947
2948	aacttcaagatgtggtaacccaaaatgcacaagctttaaacaacgcttgttaacaacttagctccaattt	3017
3018	tgggtcaatttcaagtgttttaattgatattctttcacgcttggacctctgaggctgaagtgcacaa	3087
3088	gatagggtgatcacaggcagacttcaaaagtgtgcagacatatgtactcaacaattaattagagctgcag	3157
3158	aaatcagagcttctgctaacttctgctactataaatgtcagagtgtgtgacttggacaatcaaaaagagt	3227
3228	tgatttttggtaaagggtatcatcttatgtccttccctcagtcagcacctcatgggtgtagtcttctg	3297
3298	catgtgacttatgtccctgcacaagaaaagaacttcacaactgtccttgcatttgcattgatggaaaag	3367

FIGURE 5 cont'd

3368	cacactttcctcgtgaagggtgtctttgtttcaaattggcacacactggttgaacacaaaggaatttta	3437
3438	tgaaccacaaatcattactacagacaacacattgtgtctggtaactgtgatgtgtaataggaattgtc	3507
3508	aacaacacagtttatgatcctttgcaacctgaaitagactcattcaaggaggagttagataaatttta	3577
3578	agaatcatacatcaccagatgttgatttaggtgacatctctggcattaatgcttcagttgtaaacattca	3647
3648	aaaagaaattgaccgectcaatgagggtgccaagaatttaaatgaatctctcatcgafctccaagaactt	3717
3718	ggaaagtatgagcagtatataaaatggccatgggtacatttggctaggtttatagctggcttgattgcca	3787
3788	tagtaatggtgacaattatgctttgctgtatgaccagtgtctgtagttgtctcaagggtgtgtgtctg	3857
3858	tggatcctgctgcaaatgtgatgaagacgactctgagccagtgtctcaaaggagtcaaattacattacaca	3927
3928	taa (SEQ ID NO: 8)	3930

FIG. 5B)

148	cagtgcgtgaacctgaccaccaagaacacagctgcctccagcctacaccaacagcttcaccagaggcgtgt	217
218	actaccccgacaagggtgttcagatccagcgtgtgcactctaccaggacctgttctgtcctttcttcag	287
288	caacgtgacctgggtccacgccatccacgtgtccggcacaatggcaccaagagattcgacaaccccggtg	357
358	ctgcccttcaacgacggggtgtactttgccagcaccgagaagtccaacatcatcagaggctggatctcg	427
428	gcaccacactggacagcaagaccagagcctgtgatcgtgaacaacgccaccaacgtggatcaaaagt	497
498	gtgcgagttccagttctgcaacgaccccttctgggcgtctactccaagaacaacaagagctggatg	567
568	gaaagcgagttccgggtgtacagcagcgccaacactgcaccttcgagtacgtgtccagcctttcctga	637
638	tggacctggaaggcaagcaggggcaactcaagaacctgcgcgagttcgtgtcaagaacatcgacggcta	707
708	cttcaagatctacagcaagcacacccctatcaacctcgtgcgggatctgcctcagggttctctgtcttg	777
778	gaacccctgtgtgatctgccatcggtacatcaacatccccgggttcagacactgctggcctgcacagaa	847
848	gctacctgacacctggcgatagcagctctggatggacagctggagccgctgcctactatgtgggatacct	917
918	gcagcctcggaccttctgtctgaagtacaacgagaacggcaccatcaccgacgccgtggattgtgtctg	987
988	gatcctctgagcgagacaaaagtgcacctgaagtccttcaccgtggaaaaaggcatctaccagaccagca	1057
1058	acttcgggtgcagcccacccaatccatcgtgcgggtccccaatataccaatctgtgcccttcgggga	1127
1128	gggtgtcaatgccaccagattgcctctgtgtacgcctggaaacgggaagcggatcagcaattgcgtggcc	1197
1198	gactactcctgtgtgtacaactccgccagcttcagcacttcaagtgtacggcgtgtccctaccaagc	1267
1268	tgaacgacctgtgtcttcaaaaacgtgtacgccgacagcttcgtgatccggggagatgaagtgcggcagat	1337
1338	tgccccctggacagacaggcaagatcgccgactacaactcaagctgcccgacgactcacccgctgtgtg	1407
1408	attgcctggacagcaacaacctggactccaaagtggcggaactacaattacctgtaccggctgttcc	1477
1478	ggaaagtcaatctgaagcccttcgagcgggacatctccaccagatctatcaggccggcagcacccttg	1547
1548	taacggcgtggaaggcttcaactgtaacttccactgcagtcctacggcttcagccaacaaacggcgtg	1617
1618	ggctaccagccttacagagtgggtgtgtgagcttcgagctgtgcatgctcctgccacagtgtgcggcc	1687
1688	ctaagaaaagcacaatctcgtgaagaacaaatgcgtgaacttcaactcaacggcctgacggcgaaccgg	1757
1758	cgtgctgacagagagcaacaagaagtctcctgccattccagcagttcggccgggatatcgccgataccaca	1827
1828	gatgcctgacagatccccagacacttgaaatcctggacatcaccccttcagacttcggcggaagtgtctg	1897
1898	tgatcacccctggcaccacaccagcaatcaggtggcagtgctgtaccaggaatgtgaactgtacagaggt	1967
1968	gccagtggcatttcacgccgatcagctgacccctacttggcgggtgtactccacaggcagcaatgtgttt	2037
2038	cagaccagagcggcgtgtctgatcggagccgagcacgtgaacaatagctacgagtgcgacatccccatcg	2107
2108	gcgctggcatctgcgcctcttaccagacacagacaaacagccacagcggccagatctgtggccagcca	2177
2178	gagcatcattgectacacaatgtctctgggagccgagaacagcgtggcctactccaacaactctatcgt	2247
2248	atccccaccaacttcaecatcagcgtgaccacagagatcctgctgtgtecattgaccaagaccagcgtgg	2317
2318	actgcacatgtacatctgcggcgattccaccagagtgtccaaactgctgtcagtagcggcagcttctg	2387
2388	caccagctgaatagacccctgacaggggatcgccgtggacaggacaagaacaccaagaggtgttcgcc	2457
2458	caagtgaagcagatctacaagacccctctatcaaggacttcggcggcttcaatttcagccagattctgc	2527

FIGURE 5 cont'd

2528	cogatectagcaagcccagcaagcggagctcatcgaggacctgctgticaaaaagtacactggccga	2597
2598	cgcgggttcataagcagtatggcgattgtctggcgacattgccgccagggatctgattggccccag	2667
2668	aagtftaacggactgacagtgtgctcctctgtctgaccgatgagatgategcccagtacacatctgcc	2737
2738	tgctggccggcacaatcacaaaggctggacattggagctggcgctgccctgcagatcccccttgctat	2807
2808	gcagatggcctaccgggtcaacggcatcggagtaccagaatgtgtgtacgagaaccagaagctgac	2877
2878	gccaaccagttcaacagcggcatcggcaagatccaggacagcctgagcagcacagcaagcggcctgggaa	2947
2948	agctgcaggacgtgggtcaaccagaatgccaggcactgaacacctgggtcaagcagctgtctagcaactt	3017
3018	cggagccatcagctctgtgtgaacgatctctgagcagactggacctcctgaggccgaggtgcagatc	3087
3088	gacagactgatcacaggcagactgcagagcctccagacatacgtgacccagcagctgatcagagccg	3157
3158	agattagagcctctgccaatctggccgccaccaagatgtctgagtgtgtgtgtggccagagcaagagat	3227
3228	ggacttttgcggcaagggtaccacctgatgagcttccctcagctgtgacccacacggcgtgtgtgtctg	3297
3298	cacgtgacctacgtgcccgtcaagagaagaattcaccacggctccagccatctgccagacggcaag	3367
3368	ccaacttctagagaaggcgtgtgtgtgtccacggcaccatgtgtgtgtgtgtgtgtgtgtgtgtgtgt	3437
3438	cgaagcccgatcatcaccaccgacaacaccttctgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt	3507
3508	aacaataccgtgtacgacctctgcagcccagctggacagcttcaaaggagaaatggacaagtacttta	3577
3578	agaaccacacaagccccgacgtggacctggggatcatcagggaatcaatgccagcgtgtgtgtgtgtgt	3647
3648	gaaagagatcgaccggctgaacgaggtggccaagaatctgaacgagagcctgatcagctgcaagaactg	3717
3718	gggaagtacgagcagtaacaaagtggccttggatcatctgtgtgtgtgtgtgtgtgtgtgtgtgtgt	3787
3788	tcgtgatgtgtcaaatcatctgt	3857
3858	tggctcctgtgtcaagttcgacgaggacgattctgagccgtgtgtgtgtgtgtgtgtgtgtgtgtgt	3927
3928	tga (SEQ ID NO: 9)	3930

FIG. 5C)

148	cagtgcgtgaacctgaccaccagaacacagctgcctccagcctacaccaacagcttcaccagaggcgtgt	217
218	actaccccgacaagggtgtcagatccagcgt	287
288	caacgtgacctgtgtccacgccatccacgt	357
358	ctgcccttcaacgacggggtgtactttgccagcaccgagaagtccaacatcatcagaggctgtgtgtgt	427
428	gaccacacttgacagcaagaccagagcctgt	497
498	gtgcgagttcagttctgcaacgaccttctgt	567
568	gaaagcgagttccgggtgtacagcagcggcaacactgcaccttcagtagctgtgtgtgtgtgtgtgtgt	637
638	tggacctggaaggcaagcagggcaacttcaagaacctgcgcgagttcgtgtgtgtgtgtgtgtgtgtgt	707
708	cttcaagatctacagcaagcacaacctatcaacctgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgtgt	777
778	gaacctcctgt	847
848	gtacctgacacctggcgatagcagctctgt	917
918	gcagcctcggaccttctgt	987
988	gatcctctgagcagagacaaagtgcacctgaagtccttaccgtgtgtgtgtgtgtgtgtgtgtgtgtgt	1057
1058	acttccgggtgcagccccaccgaatccatcgt	1127
1128	gggtgtcaatgccaccagattcgcctctgt	1197
1198	gactactccgt	1267
1268	tgaacgacctgt	1337
1338	tgcctcgtgacagacaggaagatgcgcgactacaactacaagctgtgtgtgtgtgtgtgtgtgtgtgtgt	1407
1408	attgcctggaacagcaacaacctggactccaaagtcggcggaactacaattacctgtaccggctgtgtgt	1477
1478	ggaaagtcacatctgaagcccttcgagcgggacatctccaccgagatctatcaggccggcagcacccctgt	1547
1548	taacggcgtggaaggcttcaactgtacttccactgcagctctacggcttccagccaacaacggcgtgtgt	1617
1618	ggctaccagccttacagagt	1687
1688	ctaagaaaagcacaatctcgtgaagaacaatgcgtgaacttcaacttcaacggcctgacggcaccgg	1757

FIGURE 5 cont'd

1758	cgtgctgacagagagcaacaagaagttcttgcattccagcagttcggccgggatatcgccgataccaca	1827
1828	gatgccgtcagagatccccagacactggaaatcctggacatcaccccttgcaacttcggcggagtgctg	1897
1898	tgatcacccctggcaccacaccagcaatcagggtggcagtgctgtaccagggcgtgaactgtacagaggt	1967
1968	gccagtggccattcacgcgatcagctgacccctacttggcgggtgtactccacaggcagcaatgtgtt	2037
2038	cagaaccagagccgggtgtctgatcggagccgagcacgtgaacaatagctacgagtgcgacatccccatcg	2107
2108	gcgctggcatctgcgctcttaccagacacagacaaacagcccagacgggccagatctgtggccagcca	2177
2178	gagcatcattgcctacacaatgtctctgggagccgagaacagcgtggcctactccaacaactctatcgt	2247
2248	atccccaccaacttcaaccatcagcgtgaccacagagatcctgctgtgtccatgaccaagaccagcgtgg	2317
2318	actgcaccatgtacatctcgggcgattccaccgagtgctccaacctgctgtcgtcagctacggcagcttctg	2387
2388	caccagctgaatagagccctgacagggatcgccgtggaaacaggacaagaacaccaagaggtgttcgcc	2457
2458	caagtgaagcagatctacaagacccctctatcaaggacttcggcggcttcaattcagccagattctgc	2527
2528	ccgactctagcaagcccagcaagcggagcttcatcgaggacctgctgttcaaaaagtgaacttggccga	2597
2598	cgcgggttcatcaagcagataggcgattgtctgggcgacattgcgcgcaaggatctgatttgcgccag	2667
2668	aagttaaaggactgacagtgtctgctcctctgtgtgaccgatgagatgatcgccagctacacatctgcc	2737
2738	tgctggccggcacatcacaagcggctggacatttggagctggcgctgcccctgcagatcccccttctat	2807
2808	gcagatggcctaccgggtcaacggcatcggagtgaccagaatgtgtgtgacgagaaccagaagctgatc	2877
2878	gccaaccagttcaacagcggcatcggcaagatccaggacagcctgagcagcagcaagcggcctgggaa	2947
2948	agctgcaggacgtgtgtcaaccagaatgccaggcactgaacacctgggtcaagcagctgtctagcaact	3017
3018	cggagccatcagctctgtgtgaacgatatcctgagcagactggacctcctgaggccgaggtgcagatc	3087
3088	gacagactgatcagggcagactgcagagcctccagacatactgacccagcagctgatcagagccgccg	3157
3158	agattagagcctctgccaatctggccgccaccaagatgtctgagtggtgtgtgctggccaagagaggt	3227
3228	ggacttttgcggcaagggttaaccactgatgagcttccctcagctgcaccacacggcgtgtgtgtctg	3297
3298	cacgtgacctacgtgcccgctcaagagaagaatttcaccacccgctccagccatctgccacgacggcaag	3367
3368	cccactttctagagaaggcgtgttcgtgtccaacggcacccattgtgtgtgacacagcggaaacttcta	3437
3438	cgaagcccgatcatcaccaccgacaaacacttctgtgtctggcaactgcgacgtcgtgatcggcattgtg	3507
3508	aacaataccgtgtacgacctctgcagcccagctggacagcttcaaagaggaaactggacaagtacttta	3577
3578	agaaccacacaagccccgacgtggacctggggcagatcagcggaaatcaatgccagcgtcgtgaacatcca	3647
3648	gaaagagatcgaccggctgaacgaggtggccaagaatctgaacgagagcctgatcgaactgcaagaactg	3717
3718	gggaagtacgagcagctacatcaagtggccttggatcatctggctgggctttatcgccggactgattgcca	3787
3788	tcgtgatggtcacaatcatgctgtgttcagaccagctgctgtagctgcctgaagggtgtgtgtgctg	3857
3858	tggtcctgctgcaagttcgacagggacgattctgagccgtgctgaaaggcgtgaagctgcactacacc	3927
3928	tga (SEQ ID NO: 10)	3930

FIG. 5D)

148	cagtgcgtgaacctgaccaccagaacacagctgcctccagcctacaccaacagcttaccagaggcgtgt	217
218	actaccccgacaagggtgtcagatccagcgtgtgcactctaccaggacctgttctgctttcttcag	287
288	caacgtgacctggttccacgcatccacgtgtccggcaccacaaatggcaccagaagatcgacaaccccg	357
358	ctgcccttcaacgacggggtgtactttgccagcaccgagaagtccaacatcatcagaggctggatcttcg	427
428	gcaccactggacagcaagaccagagcctgtgatcgtgaacaacgccaccaacgtggatcatcaaagt	497
498	gtgcgagttcagttctgaacgaacccctctggcgcttactaccacaagaacaacaagagctggatg	567
568	gaaagcgagttccgggtgtacagcagcggcaacaactgcaccttcagctgtgtccagaccttctga	637
638	tggaacctggaaggcaagcagggcaacttcaagaacctgcgcgagttcgtgttcaagaacatcgacggcla	707
708	cttcaagatctacagaagcacaacccctatcaacctcgtgcgggatctgcctcagggttctctgtctg	777
778	gaacccctggttgatctgcccctcggaatcaacatcacccgggttcagacactgctggccctgcacagaa	847
848	gctacctgacacctggcgatagcagctctggatggacagctggagccgctgcctactatgtgggatacct	917
918	gcagcctcggaaccttctgtgaagtacaacgagaacggcaccatcacccagccgtggattgtgtctg	987

FIGURE 5 cont'd

988	gatcctctgagcgagacaaagtgcacccctgaagtccttcaccgtggaaaaggcatctaccagaccagca	1057
1058	acttcegggtgcagcccacgaatccatcgtgcgggtcccaatafcaccaatctgtcccccttcggcga	1127
1128	ggtgttcaatgccaccagattcgcctctgtgtacgectggaaccggaagcgatcagcaattgcgtggcc	1197
1198	gactactccgtgctgtacaactccgccagcttcagcaactcaagtgtaaggcggtgtccctaccaagc	1267
1268	tgaacgacctgtgcttcacaaacgtgtacgccgacagcttcgtgatccggggagatgaagtgcggcagat	1337
1338	tgcccttgacagacaggcaagatgccgactacaactacaagctgcccgaagactaccggctgtgtg	1407
1408	attgcctggaaacagcaacaacctggactccaaagtgcggggcaactacaattacagataccggctgttcc	1477
1478	ggaaagtccaatctgaagcccttcgagcggggacatccacggagatctatcaggccggcagcaccccttg	1547
1548	taacggcgtggaaggcttcaactgtacttccactgcagtcctacggcttcagccaacatacggcggtg	1617
1618	ggctaccagccttacagagtgggtggtgctgagcttcgagctgctgcatgctcctgccacagtgtgcggcc	1687
1688	ctaagaaaagcaccaatctcgtgaagaacaaatcgtgaacttcaactcaacggcctgacgggcacogg	1757
1758	cgtgctgacagagagcaacaagaaagtccctgccattccagcagttcggccgggatacggcgataccaca	1827
1828	gatgccgtcagagatcccagacactggaaatccctggacatcaccccttgcaacttcggcggagtgctg	1897
1898	tgatcccccctggcaccaacaccagcaatcaggtggcagtgctgtaccagggcggtgaactgtacagaggt	1967
1968	gccagtggccattcacgccgatcagctgacccctacttggcgggtgtactccacaggcagcaatgtgtt	2037
2038	cagaccagagccggctgtctgatcggagccgagcacgtgaacaatagtaagagtgcgacatccccatcg	2107
2108	gcgctggcatctgcgcctcttaccagacacagacaaaacagccccagacggccagatctgtggccagcca	2177
2178	gagcatcattgectacacaatgtctctgggagccgagaacagcgtggcctactccaacaactctatcgt	2247
2248	atccccaccaacttcaaccatcagcgtgaccacagagatcctgctgtgtccatgaccaagaccagcgtgg	2317
2318	actgcaaccatgtacatctgcggcgattccaccgagtgtccaaactgctgctgcagtacggcagcttctg	2387
2388	caccagctgaatagagccctgacaggatcggcgtggaacaggacaagaacaccaagaggtgttcgcc	2457
2458	caagtgaagcagatctacaagacccctctatcaaggacttcggcggcttcaatttcagccagattctgc	2527
2528	ccgatcctagcaagcccagcaagcggagcttcacagaggacctgctgttcaacaaagtgacactggccga	2597
2598	cggcggcttcatcaagcagtatggcgattgtctggcgacattgccgcaaggatctgatttgcgccag	2667
2668	aagtttaacggactgacagtgtcctcctctgtgaccgatgagatgacgccagtaacatctgcc	2737
2738	tgctggcgggcacaatcacaagcggcgtggacatttggagctggcgctgcccigcagatccccittgtat	2807
2808	gcagatggcctaccgggtcaacggcatcggagtgaaccagaatgtgtgtgtacgagaaccaagaagctgatc	2877
2878	gccaaccagttcaacagcggccatcggcaagatccaggacagcctgagcagcacagcaagcgcctgggaa	2947
2948	agctgcaggacgtgtgtaaccagaatgccaggcactgaacacccctggtcaagcagctgtctagcaact	3017
3018	cggagccatcagctctgtgtgtgaacgatctctgagcagactggacctcctgagggcgaggtgcagatc	3087
3088	gacagactgatcagggcagactgcagagcctccagacatacgtgacccaagcagctgatcagaccgccg	3157
3158	agattagagcctctgccaatctggccgccaccaagatgtctgagtgtgtgtgtggccagagcaagagagt	3227
3228	ggacttttgcggcaagggtaccacctgagtgttccctcagctgtgaccacacggcggtgtgttctg	3297
3298	caegtgaacctacgtgcccgtcgaagagaagaatttcaccacccgtccagccatctgccacgacggcaag	3367
3368	cccactttcctagagaaggcgtgtgtgttccaaacggcaccctattgggtcgtgacacagcggaaacttcta	3437
3438	cgagccccagatcatcaccaccgacaacaccttcgtgtctggcaactgcgacgtcgtgatcggcattgtg	3507
3508	aacaataccgtgtacgacctctgcagcccagctggacagcttcaaagggaactggacaagtacttta	3577
3578	agaaccacacaagccccgacgtggacctggggcgatatcagcggaaatcaatgccagcgtcgtgaacatcca	3647
3648	gaaagagatcgaccggctgaacgaggtggccaagaatctgaaacgagagcctgatcgaacctgcaagaactg	3717
3718	gggaagtacgagcagtaacatcaagtggccttggtaacatctggtgtgggtttatcgccggactgattgcca	3787
3788	tcgtgatgttcacaatcatgtgtgtgtgcatgaccagctgctgtagctgcctgaagggtgtgtgtgctg	3857
3858	tggctcctgctgcaagttcgacgaggacgattctgagcccgtgctgaaaggcgtgaagctgcactacac	3927
3928	tga (SEQ ID NO: 11)	3930

FIG. 6A)

14	QCVNLTTRTQLPPAYTNSFTRGVYYPDKVFRSSVLHSTQDLFLPFFSNVT	64
64	WFHAIHVSGTNGTKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDSK	113
114	TQSLILIVNNATNVVIVKVEFQFCNDPFLGVYHKNKNSWMESEFRVYSSA	163
164	NNCTFEYVSQPFLLMDLEGKQGNFKNLREFVFKNIDGYFKIYSKHTPINLV	213
214	RDLPOGFSALEPLVDLPIGINITRFQTLLALHRSYLTPGDSSSGWTAGAA	263
264	AYYVGYLQPRTFLLKYNENGTITDAVDCALDPLSETKCTLKSFTVEKGIY	313
314	QTSNFRVQPTESIVRFPNITNLCPFGEVFNATRFASVYAWNRRKRISNCVA	363
364	DYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVQRQIAPG	413
414	QTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGGNYNLYRLFRKSNLKP	463
464	FERDISTEIIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYQPYRVVVL	513
514	SFELLHAPATVCGPKKSTNLVKNKCVNFNFNGLTGTGVLTESNKKFLPFQ	563
564	QFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITPGTNTSNQVAVLYQ	613
614	GVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAHEVNNSYECD	663
664	IPIGAGICASYQTQTNSPRRARVASQSIIAYTMSLGAENSVAYSNNNSIA	713
714	IPTNFTISVTTEILPVSMTKTSVDCTMYICGDSTECSNLLLQYGSFCTQL	763
764	NRALTGIAVEQDKNTQEVFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPS	813
814	KRSFIEDLLFNKVTLADAGFIKQYGDCLGDIAARDLICAQKFNGLTVLPP	863
864	LLTDEMIAQYTSALLAGTITSGWTFGAGAAIQIPFAMQMAYRFNGIGVTQ	913
914	NVLYENQKLIANQFNSAIGKIQDSLSSTASALGKLQDVVNQNAQALNTLV	963
964	KQLSSNFGAISSVLNDILSRDPPEAEVQIDRLITGRLQSLQTYVTQQLI	1013
1014	RAAEIRASANLAATKMSECVLGQSKRVDFCGKGYHLMSFPQSAPHGVVFL	1063
1064	HVTYVPAQEKNFTTAPAICHGKAHFPREGVFVSNNGTHWFVTQRNFYEPQ	1113
1114	IITTDNTFVSGNCDVVIGIVNNTVYDPLQPELDSFKEELDKYFKNHTSPD	1163
1164	VDLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDLQELGKYEQYIKWP	1213
1214	WYIWLGFIAGLIAIVMVTIMLCCMTSCCSCCLKGCCSCGSCCKFDEDDSEP	1263
1264	VLKGVKLHYT (SEQ ID NO: 12)	

FIG. 6B)

14	QCVNLTTRTQLPPAYTNSFTRGVYYPDKVFRSSVLHSTQDLFLPFFSNVT	63
64	WFHAIHVSGTNGTKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDSK	113
114	TQSLILIVNNATNVVIVKVEFQFCNDPFLGVYHKNKNSWMESEFRVYSSA	163
164	NNCTFEYVSQPFLLMDLEGKQGNFKNLREFVFKNIDGYFKIYSKHTPINLV	213
214	RDLPOGFSALEPLVDLPIGINITRFQTLLALHRSYLTPGDSSSGWTAGAA	263
264	AYYVGYLQPRTFLLKYNENGTITDAVDCALDPLSETKCTLKSFTVEKGIY	313
314	QTSNFRVQPTESIVRFPNITNLCPFGEVFNATRFASVYAWNRRKRISNCVA	363
364	DYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVQRQIAPG	413
414	QTGKIADYNYKLPDDFTGCVIAWNSNNLDSKVGGNYNLYRLFRKSNLKP	463
464	FERDISTEIIYQAGSTPCNGVEGFNCYFPLQSYGFQPTNGVGYQPYRVVVL	513
514	SFELLHAPATVCGPKKSTNLVKNKCVNFNFNGLTGTGVLTESNKKFLPFQ	563
564	QFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITPGTNTSNQVAVLYQ	613
614	GVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAHEVNNSYECD	663
664	IPIGAGICASYQTQTNSPRRARVASQSIIAYTMSLGAENSVAYSNNNSIA	713
714	IPTNFTISVTTEILPVSMTKTSVDCTMYICGDSTECSNLLLQYGSFCTQL	763
764	NRALTGIAVEQDKNTQEVFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPS	813
814	KRSFIEDLLFNKVTLADAGFIKQYGDCLGDIAARDLICAQKFNGLTVLPP	863

FIGURE 6 cont'd

864	LLTDEMIAQYTSALLAGTITSGWTFGAGAALQIPFAMQMAYRFNGIGVTQ	913
914	NVLYENQKLIANQFNSAIGKIQDSLSSSTASALGKLQDVVNQNAQALNTLV	963
964	KQLSSNFGAISSVLNDILSRDPPEAEVQIDRLITGRLQSLQTYVTQQLI	1013
1014	RAAEIRASANLAATKMSECVLGQSKRVDFCGKGYHLMSFPQSAPHGVVFL	1063
1064	HVTYVPAQEKNFTTAPAICHGKAHFPREGVFVSNGTHWTFVTQRNFYEPQ	1113
1114	IITTDNTFVSGNCDVVIGIVNNTVYDPLQPELDSFKEELDKYFKNHTSPD	1163
1164	VDLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDLQELGKYEQYIKWP	1213
1214	WYIWLGFIAGLIAIVMTIMLCCMTSCCSCLKGCCSCGSCCKFDEDDSEP	1263
1264	VLKGVKLHYT (SEQ ID NO: 13)	1273

FIG. 6C)

14	QCVNLTRTQLPPAYTNSFTRGVYYPDKVFRSSVLHSTQDLFLPFFSNVT	63
64	WFHAIHVSGTNGTKRFDNPVLPFNDGVYFASTEKSNIIRGWIFGTTLDSK	113
114	TQSLLIIVNNATNVVIKVFCEQFCNDPFLGVYHKNKNSWMESEFRVYSSA	163
164	NNCTFEYVSQPFLMDLEGKQGNFKNLREFVFKNIDGYFKIYSKHTPINLV	213
214	RDLPOGFSALEPLVDLPIGINITRFQTLALHRSYLTPGDSSSGWTAGAA	263
264	AYYVGYLQPRTFLLKYNENGTITDAVDCALDPLSETKCTLKSFTVEKGIY	313
314	QTSNFRVQPTESIVRFPNITNLCPFGEVFNATRFASVYAWNRRKRISNCVA	363
364	DYSVLYNSASFSTFKCYGVSPTKLNDLCFTNVYADSFVIRGDEVQRQIAPG	413
414	QTGKIADYNYKLDDFTGCVIAWNSNNLDSKVGGNYNRYRLFRKSNLKP	463
464	FERDISTEIYQAGSTPCNGVEGFNCYFPLQSYGFQPTYGVGYQPYRVVVL	513
514	SFELLHAPATVCGPKKSTNLVKNKCVNFNFNGLTGTGVLTESNKKFLPFQ	563
564	QFGRDIADTTDAVRDPQTLEILDITPCSFGGVSVITPGTNTSNQVAVLYQ	613
614	GVNCTEVPVAIHADQLTPTWRVYSTGSNVFQTRAGCLIGAEHVNNSYECD	663
664	IPIGAGICASYQTQTNSPRRARSVASQSI IAYTMSLGAENSVAYSNNNSIA	713
714	IPTNFTISVTTEILPVSMTKTSVDCTMYICGDSTECSNLLLQYGSFCTQL	763
764	NRALTGIAVEQDKNTQEVFAQVKQIYKTPPIKDFGGFNFSQILPDPSKPS	813
814	KRSFIEDLLFNKVTADAGFIKQYGDCLGDIAARDLICAQKFNGLTVLPP	863
864	LLTDEMIAQYTSALLAGTITSGWTFGAGAALQIPFAMQMAYRFNGIGVTQ	913
914	NVLYENQKLIANQFNSAIGKIQDSLSSSTASALGKLQDVVNQNAQALNTLV	963
964	KQLSSNFGAISSVLNDILSRDPPEAEVQIDRLITGRLQSLQTYVTQQLI	1013
1014	RAAEIRASANLAATKMSECVLGQSKRVDFCGKGYHLMSFPQSAPHGVVFL	1063
1064	HVTYVPAQEKNFTTAPAICHGKAHFPREGVFVSNGTHWTFVTQRNFYEPQ	1113
1114	IITTDNTFVSGNCDVVIGIVNNTVYDPLQPELDSFKEELDKYFKNHTSPD	1163
1164	VDLGDISGINASVVNIQKEIDRLNEVAKNLNESLIDLQELGKYEQYIKWP	1213
1214	WYIWLGFIAGLIAIVMTIMLCCMTSCCSCLKGCCSCGSCCKFDEDDSEP	1263
1264	VLKGVKLHYT (SEQ ID NO: 14)	

FIGURE 7

1	CACCACCACCACCACCACTGACTCGAGGCTGGAGCCTCGGTGGCCATGCT	50
	4(e) 4(a)	
51	TCTTGCCCCTTGGGCCTCCCCCAGCCCCTCCTCCCCTTCCTGCACCCGT	100
	4(a)	
101	ACCCCGGGTCTTTGAGATCTGGTTACCACTAAACCAGCCTCAAGAACAC	150
	4(a) 4(b)	
151	CCGAATGGAGTCTCTAAGCTACATAATACCAACTTACACTTTACAAAATG	200
	4(b)	
201	TTGTCCCCCAAATGTAGCCATTCGTATCTGCTCCGTTGCCAAAGAAAGT	250
	4(b)	
251	TTCTTCACATTCTAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA	300
	4(b) 4(c)	
301	AA	350
	4(c)	
351	AAAGGC	396
	4(c) 4(d)	

(SEQ ID NO: 15)

1

VACCINE FOR USE AGAINST CORONAVIRUS AND VARIANTS THEREOF

FIELD OF THE INVENTION

The present invention generally relates to vaccines, and in particular, to vaccines that are useful against Coronavirus.

SUBMISSION OF SEQUENCE LISTING

The contents of the electronic submission of the text file Sequence Listing, which is named GOW-41411.Sequenc. List.5.19.2022.txt, which was created May 19, 2022 and is 56 KB in size, is incorporated herein by reference in its entirety.

BACKGROUND

A new infectious coronavirus (SARS-CoV-2) was first reported in Wuhan, China in December, 2019 that causes COVID-19. The World Health Organization (WHO) declared COVID-19 a global public health emergency situation on Feb. 5, 2020 after obtaining growing evidence of continuous person-to-person transmission. The virus spread worldwide quickly, and consequently WHO declared it pandemic in March of 2020. As of Dec. 21, 2020, the pandemic has resulted in 1,701,187 deaths among over 42 77,243,764 patients in 220 countries, with a case-fatality rate of 2.20%.

There will be a risk of pandemic as long as there is a COVID-19 epidemic situation in any part of the world unless people are properly vaccinated. Therefore, effective vaccines against SARS-CoV-2 are required to control morbidity and mortality related to COVID-19. Generally, non-replicating viral vectors, inactivated virus, DNA-based and protein-based vaccines have been the major approaches for the development of stable and effective vaccines; although, they have their inherent limitations. Recently, mRNA-based vaccines have become a promising approach because of their opportunity for rapid development, comparative low dose, and better safety profile in terms of no potential risk of infection or insertional mutagenesis, and low capital expenditure (CAPEX).

Several leading vaccines have been developed as well as others which are currently under development against SARS-CoV-2, including vaccines which are mRNA-based. Lipid nanoparticle technology has been developed for effective delivery of single-stranded therapeutics like siRNA, antisense oligo, mRNA etc. The first RNA-LNP therapeutic was approved in 2018 and has set the example for clinical safety of LNP-formulated RNA. Two approved vaccines against SARS-CoV-2 are mRNA vaccines (developed by Pfizer-BioNTech and Moderna Inc.) and they have been successfully used to vaccinate millions of people.

Coronaviruses have genetic proofreading mechanisms, and SARS-CoV-2 sequence diversity is comparatively low; although, natural selection can adopt rare but favorable mutations. Since the outbreak in China, SARS-CoV-2 has gone through numerous mutations. The D614G amino acid change, among these, in the spike protein of Wuhan reference strain is caused by an A-to-G nucleotide substitution at position 23,403 of the relevant nucleotide sequence. Currently, D614G is the most prevalent circulating isotype of SARS-CoV-2 worldwide (more than 95%). The G614 genotype is associated with increased case fatality rate over D614. Scientific findings have demonstrated that the G614 variant is ~10 times more infectious over the D614 geno-

2

type. It has also been revealed from in vitro and clinical data that the G614 variant has a distinct phenotype, and there is likely be a huge impact of this mutation on infection, transmission, disease onset, disease prognosis, as well as on vaccine and therapeutic development.

To date, there is no published report regarding D614G-relevant vaccine development. A few studies have shown that antibody generated using D614G variant-target did not show a significant difference between D614 and G614 variants in terms of cellular entry. These studies did not use G614-specific antibody, and applied artificial systems for characterizing relevant functional experiments. Furthermore, it is not known how the G614 variant vaccine will behave on immunization in humans and the impact of the relevant antibody on SARS-CoV-2.

Besides D614G, which is a constant mutation in all published variants, other prevalent mutations in the spike protein include: K417N, K417T, L452R, T478K, E484Q, E484K, N501Y, A570D, H655Y, P681R, P681H etc. Among them L452R, E484K, N501Y, A570D, D614G and P681H are referred to as the Alpha variant (B.1.1.7); K417N, E484K, N501Y and D614G are referred to as the Beta variant (B.1.351, B.1.351.2, B.1.351.3); K417T, E484K, N501Y, D614G and H655Y are referred to as the Gamma variant (P.1, P.1.1, P.1.2); and L452R, T478K, D614G and P681R are referred to as the Delta variant (B.1.617.2, AY.1, AY.2). The Delta variant is the most prevalent variant around the globe. However, there is no reported vaccine candidate that has taken these mutations into consideration.

Therefore, development of a G614 variant-specific vaccine is of prime importance.

SUMMARY

A transgene for use in coronavirus vaccine has now been developed incorporating a spike protein variant-targeted nucleic acid sequence, as well as other immunogen-enhancing aspects.

Thus, in one aspect of the invention, a transgene is provided that encodes:

- i) an RNA polymerase promoter;
- ii) a 5' UTR;
- iii) a secretory sequence;
- iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant sequence at amino acid position 614 of a native spike protein, and optionally variant sequences at one or both of amino acid positions 452 and 501 of the native spike protein; and
- v) a 3' UTR and poly A sequence.

In another aspect of the invention, a vaccine incorporating a transgene as described above encoding a spike protein variant sequence at amino acid positions 452, 501 and 614 of a native spike protein is provided.

In another aspect of the invention, a method of vaccinating a mammal is provided comprising administering to the mammal a vaccine incorporating a transgene that encodes: i) an RNA polymerase promoter; ii) a 5' UTR; iii) an IgE-based secretory sequence; iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant sequence at amino acid position 614 of a native spike protein or a position that corresponds with the amino acid position at position 614 of the native spike protein, and optionally variant sequences at one or both of amino acid positions 452 and 501 of the native spike protein, and v) a 3' UTR and poly A sequence.

These and other aspects of the invention are described in the detailed description that follows by reference to the following figures.

BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A-1H respectively provide: FIG. 1A is a schematic of a transgene depicting L452, N501Y, D614G, K986P, and V987P (as shown in SEQ ID NO: 14) in accordance with an embodiment of the invention, FIGS. 1B-1D are DNA sequencing electropherogram data of L452, N501Y and D614G sequence (as shown in SEQ ID NO: 14) in the target respectively, FIG. 1E is a transient extracellular protein expression in Flp-In CHO and Lenti-X 293T cells where lane 1 is the positive control of Spike S2 ECD-His recombinant protein, lane 2-3 are natural spike extracellular protein containing RYG variant in Flp-In CHO and Lenti-X 293T cells respectively, lane 4-5 are GC-rich codon optimized spike extracellular protein containing RYG variant in Flp-In CHO and Lenti-X 293T cells respectively, and lane 6 is the reference protein ladder, FIG. 1F is a IVT optimization where Lane 4 is the optimized condition, FIG. 1G represents identification of purified capped mRNA by SEC-HPLC, and FIG. 1H represents particle size distribution of mRNA-LNP dose formulation;

FIGS. 2A-2E graphically illustrate, respectively: FIG. 2A represents an antibody titer analysis from serum of different groups after 14 and 35 days of immunization (n=6), data were 630 compared by Mann-Whitney test, ****=p-value <0.0001, ***=p-value <0.001, **=p-631 value <0.01; FIG. 2B depicts the ratio of IgG2a and IgG1 in treatment groups, FIG. 2C depicts the ratio of IgG2a+IgG2b and IgG1+IgG3 in treatment groups, FIG. 2D represents serum antibody affinity analysis, and FIG. 2E represents resin pull-down serum antibody affinity analysis.

FIGS. 3A-3M graphically illustrate, respectively, cellular immune response analysis (cellular and secretory cytokine) in control and treatment groups (n=3 for each mentioned otherwise) including: FIG. 3A depicts IFN-gamma expressing cell population percentage at Day 14, treatment n=2; FIG. 3B depicts IL-2 expressing cell population percentage at Day 14, FIG. 3C depicts TNF- α expressing cell population percentage at Day 14, FIG. 3D depicts IL-6 expressing cell population percentage at Day 14, FIG. 3E depicts secretory IFN-gamma concentration at 6 and 18 hours, FIG. 3F depicts secretory IL-2 concentration at 6 and 18 hours, FIG. 3G depicts secretory IL-4 concentration at 6 and 18 hours, FIG. 3H depicts secretory IL-6 concentration at 6 and 18 hours, FIG. 3I depicts CD4+ cell population at Day 91, stimulated with S1 peptide pool, FIG. 3J depicts CD4+ cell population at Day 91, stimulated with S2 peptide pool, FIG. 3K depicts CD8+ cell population at Day 91, stimulated with S1 peptide pool, FIG. 3L depicts CD8+ cell population at Day 91, stimulated with S2 peptide pool, and FIG. 3M depicts CD27+ memory B cell population at Day 91, stimulated with S1 peptide pool (unpaired T-test were performed between control and treatment groups; ***=p-value <0.001, **=p-643 value <0.01, *=p-value <0.05).

FIGS. 4A-4K graphically illustrate, respectively, the results of in vitro neutralization assays including: FIG. 4A represents an image of Green fluorescence protein (GFP) expression after adeno-based SARS-CoV-2 pseudovirus neutralization assay from 2-4 sample dilution, FIG. 4B depicts the correlation between SARS-CoV-2 antibody from mice sera and intensity of GFP in different experimental groups, FIG. 4C depicts adeno-based SARS-CoV-2 pseudovirus neutralization percentage at different sample

dilutions, analyzed by real-time PCR, FIG. 4D depicts HIV-1 based SARS-CoV-2 pseudovirus copy number analysis by real-time PCR, FIG. 4E depicts the inhibition and neutralization assay of the SARS-CoV-2-S pseudo-type virus; all the samples were compared by one-way ANNOVA method, ***=p-664 value <0.001, **=p-value <0.01, *=p-value <0.05, FIG. 4F depicts fluorescence image of lung section of control group mouse, FIG. 4G depicts trans image of lung section of control group mouse, FIG. 4H depicts overlay image of lung section of control group mouse, FIG. 4I depicts fluorescence image of lung section of treatment group mouse, FIG. 4J depicts trans image of lung section of treatment group mouse, and FIG. 4K depicts overlay image of lung section of treatment group mouse, intentional green color enhancement was done to observe any GFP intensity.

FIGS. 5A-5D illustrate nucleotide sequences of exemplary SARS-CoV-2 spike protein components comprising: FIG. 5A depicts D614G, K986P and V987P mutations (SEQ ID No. 8); FIG. 5B depicts GC-rich and codon optimized synthetic sequence (amino acid position 14-1273) with D614, and K986P and V987P mutations (SEQ ID No. 9); FIG. 5C depicts GC-rich and codon optimized synthetic sequence (amino acid position 14-1273) with D614G, K986P and V987P mutations (SEQ ID No. 10); and FIG. 5D depicts GC-rich and codon optimized synthetic sequence (amino acid position 14-1273) with L452R, N501Y, D614G, K986P and V987P mutations (SEQ ID No. 11);

FIGS. 6A-6C illustrate amino acid sequences of exemplary SARS-CoV-2 spike protein components comprising: FIG. 6A depicts D614G, K986P and V987P mutations (SEQ ID No. 12); FIG. 6B depicts D614, and K986P and V987P mutations (SEQ ID No. 13); and FIG. 6C depicts L452R, N501Y, D614G, K986P and V987P mutations (SEQ ID No. 14); and

FIG. 7 illustrates the nucleotide sequence of a 3'-UTR sequence incorporated within a transgene in accordance with an embodiment of the invention, including a spacer sequence at nucleotide positions 1 to 27, modified 3'-UTR of alpha-1-globin at nucleotide positions 28 to 115, modified 3'-UTR of β -globin at positions 116 to 263, a poly A tail followed by a spacer (GGC) (SEQ ID No. 15).

DETAILED DESCRIPTION

A transgene for use in a coronavirus vaccine is provided. The transgene encodes: i) an RNA polymerase promoter; ii) a 5' UTR; iii) a secretory sequence; iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant sequence at amino acid position 614 of a native spike protein or a position that corresponds with the amino acid position 614 of the native spike protein, and optionally variant sequences at one or both of amino acid positions 452 and 501 of the native spike protein; and v) a 3' UTR and poly A sequence.

The coronavirus is an enveloped RNA virus with a nucleocapsid of helical symmetry and club-shaped spikes that project from surface. The coronavirus is one of the largest RNA viruses, having a genome that ranges in size from approximately 26 to 32 kilobases. Coronaviruses comprise four genera: Alphacoronavirus, Betacoronavirus, Gammacoronavirus and Deltacoronavirus, and of these, Alphacoronaviruses and Betacoronaviruses infect mammals. Coronaviruses (CoV) include common cold viruses, such as HCoV-229E, HCoV-HKU1, HCoV-NL63, and HCoV-OC43, as well as SARS-CoV. The term "SARS-CoV" refers

to a coronavirus that causes severe acute respiratory syndrome. Examples include SARS-CoV1, SARS-CoV2 and MERS-CoV.

SARS-CoV-2 is an enveloped RNA virus of ~30 kb comprising a 5'-UTR, two ORFs which encode 16 non-structural proteins such as RNA-dependent RNA polymerase, and a region that encodes structural proteins including a spike protein (S), envelope protein (E), membrane protein (M), and nucleocapsid protein (N). In one embodiment, the SARS-CoV-2 has a genome sequence that essentially corresponds with the reference SARS-CoV-2 sequence deposited at Genbank under accession no. MN997409, a 29882 nucleotide sequence in which the region spanning ribonucleotide positions 21563-25384 encodes the spike (S) protein. The coronavirus spike protein includes a signal peptide located at the N-terminus, the S1 subunit, and the S2 subunit; the last two regions are responsible for receptor binding and membrane fusion, respectively. In the S1 subunit, there is an N-terminal domain and a receptor-binding domain (RBD); while the fusion peptide (FP), heptapeptide repeat sequence 1 (HR1), heptapeptide repeat sequence 2 (HR2), transmembrane (TM) domain, and cytoplasm domain comprise the S2 subunit.

SARS-CoV-2 variants have been identified which incorporate an amino acid change at amino acid position 452 of the native spike protein sequence, referred to herein as a "SARS-CoV-2 452 variant". In one embodiment, the amino acid change is represented by L452R, i.e. substitution of leucine (L) in the native strain with arginine (R) in a variant strain, which results from an T-to-G nucleotide substitution at position 22,917 of the native nucleotide sequence. As used herein, these variant coronaviruses are referred to as the "L452R" or "R452" variant.

SARS-CoV-2 variants have been identified which incorporate an amino acid change at amino acid position 501 of the native spike protein sequence, referred to herein as a "SARS-CoV-2 501 variant". In one embodiment, the amino acid change is represented by N501Y, i.e. substitution of asparagine (N) in the native strain with tyrosine (Y) in a variant strain, which results from an A-to-T nucleotide substitution at position 23,063 of the native nucleotide sequence. As used herein, these variant coronaviruses are referred to as the "N501Y" or "Y501" variant.

SARS-CoV-2 variants have been identified which incorporate an amino acid change at amino acid position 614 of the native spike protein sequence, referred to herein as a "SARS-CoV-2 614 variant". In one embodiment, the amino acid change is represented by D614G, i.e. substitution of aspartic acid (D) in the native strain with glycine (G) in a variant strain, which results from an A-to-G nucleotide substitution at position 23,403 of the native nucleotide sequence. As used herein, these variant coronaviruses are referred to as the "D614G" or "G614" variant.

The present transgene is constructed using well-established methods, and may be constructed for use in a nucleic acid-based vaccine, e.g. an mRNA vaccine or DNA vaccine, a subunit vaccine, a non-replicating viral vaccine, an attenuated viral vaccine and an inactivated viral vaccine.

The transgene encodes: i) an RNA polymerase promoter; ii) a 5' UTR; iii) an IgE-based secretory sequence; iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant at amino acid position 614 of a native spike protein or a position that corresponds with the amino acid position 614 of the native spike protein, and optionally variant sequences at one or both of amino acid positions 452 and 501 of the native spike

protein or a position that corresponds with the amino acid at amino acid positions 452 and/or 501; and v) a 3' UTR and poly A sequence.

The transgene incorporates a sequence encoding a promoter. Generally, the transgene will encode a strong promoter that results in high levels of transcription on binding with a selected RNA polymerase. Examples of suitable promoters include promoters recognized by polymerases of the single-subunit DNA-dependent RNA polymerase (ssRNAP) family, such as the T7 promoter, the T3 promoter, the Sp6 promoter, the mitochondrial RNA polymerase (POLRMT) promoter, and the chloroplastic ssRNAP promoter. The nucleotide sequence encoding the T7 promoter (with an GG cap at position 35-36) is: 5'-TAATACGACTCACTATAGG-3' (SEQ ID No: 1). The nucleotide sequence encoding the T3 promoter (with an GG cap at position 35-36) is: 5'-AATTAACCCTCACTAAAGG-3' (SEQ ID No: 2). The nucleotide sequence encoding the Sp6 promoter (with an AG cap at position 35-36) is: 5'-ATTAGGTGACACTATAG TAG 3' (SEQ ID No: 3).

To facilitate efficient transcription by the promoter, the transgene also incorporates an upstream spacer sequence adjacent to the promoter sequence. The spacer is generally 10-20 nucleotides in length. In one embodiment, the spacer is 17-18 nucleotides in length. Upstream promoter spacer sequences are generally known in the art, and are not particularly restricted with respect to sequence. In one embodiment, the transgene incorporates a promoter spacer sequence that is a random nucleotide sequence. In another embodiment, the promoter spacer sequence is an AT-rich nucleotide sequence, e.g. comprising 40-50% or more adenine (A) and thymine (T) residues. In another embodiment, the promoter spacer has the following nucleotide sequence: GCCTGGCTTATCGAAAT (SEQ ID No: 4).

The transgene incorporates a 5'-UTR sequence that functions to enhance the transcription and translation of the coding region, e.g. for the spike protein component. The 5'-UTR incorporated into the transgene may be the native 5'-UTR for the coding region of the target coronavirus. Native 5'-UTR sequences are generally about 150-200 nucleotides in length. Alternatively, the 5'-UTR may be a synthetic sequence. In one embodiment, the 5'-UTR sequence is:

(SEQ ID No: 5)
5'-GAAATAAGAGAGAAAAGAAGAGTAAGAA
GAAATATAAGAGCTAGCGGTACC-3'.

The transgene incorporates a nucleotide sequence that encodes a coronavirus spike protein component that targets SARS-CoV-2 452, 501 and 614 variants, such as the L452R, N501Y and D614G coronavirus variants. Thus, the nucleotide sequence of the spike protein component comprises a nucleotide sequence that essentially corresponds with a native spike protein-encoding sequence except that the nucleotides encoding the amino acid at position 614 of the spike protein, and optionally the nucleotides encoding the amino acid at positions 452 and 501 of the native spike protein, encode a variant amino acid, such as glycine (614), tyrosine (452) and arginine (501). Nucleotide sequences encoding exemplary spike protein components, and the amino acid sequences encoded, are illustrated in FIGS. 5A-5D and FIGS. 6A-6C, respectively. As the spike protein is common across Coronaviridae, but the corresponding amino acid in the spike protein of coronaviruses other than the SARS-CoV-2 virus may occur at a position offset from

amino acid positions 452, 501 and 614, it is understood herein that such variants are also SARS-CoV variants. The transgene may encode a full-length variant spike protein, or may encode an antigenic fragment thereof that incorporate nucleotides encoding the variant 614 amino acid, an optionally the variant 452 and 501 amino acids. For example, (i) the transgene may encode an N-terminal fragment up to at least amino acid position 614, (ii) the transgene may encode an N-terminal fragment up to at least amino acid position 541, (iii) a C-terminal fragment including the amino acid at position 452, (iv) a C-terminal fragment including the amino acid at position 319, (v) an internal fragment incorporating at least the region R452-G614, and (vi) an internal fragment incorporating at least the region 319-541 of the spike protein. The transgene comprising L452R, N501Y and D614G variations in the spike protein component may be referred to as the 'RYG' spike protein variant.

The transgene may incorporate one or more additional sequence variations to more specifically target known SARS-CoV-2 variants, for example, the transgene may include a sequence that also encodes the amino acid variant, T478K; or the amino acid variant, E484K; or the amino acid variant, E484Q; or the amino acid variant, K986P; or the amino acid variant, V987P or the amino acid variants; or the amino acid variants, K986P and V987P, or any other combination of these variants, or other amino acid variants.

In addition to the spike protein coding sequence, the spike protein component also incorporates a nucleotide sequence that encodes a secretory sequence which is upstream of the coding sequence. The secretory sequence may be the native secretory sequence for the spike protein, or may be a non-native secretory sequence. The secretory sequence is selected based on its efficient cleavage from the spike protein component when translated so as not to compromise the function of the transgene product. Secretory sequences suitable for use include, but are not limited to, the IgE receptor secretory sequence, erythropoietin secretory sequence, growth hormone secretory sequence, granulocyte colony stimulating factor secretory sequence, fibroblast growth factor secretory sequence, vascular endothelial growth factor secretory sequence, and G-protein coupled receptor family secretory sequences. As one of skill in the art will appreciate, the selected secretory sequence may comprise one or more modifications from its native sequence, i.e. one or more modified nucleotides, provided that such modifications do not adversely affect its cleavage from the spike protein component following translation. In one embodiment, the native secretory sequence of the spike protein is replaced with the secretory sequence of the IgE receptor sequence, 5'-MDWTWILFLVAAATRVHS-3' (SEQ ID No: 6), encoded by the nucleotide sequence, atggactggacgtg-gatcctcttcttggtggcagcagccagcgcagtcactcc (SEQ ID No: 7).

The transgene also incorporates a downstream 3' UTR region to facilitate translation and provide stability to the transcript, and may additionally comprise an upstream ribosome-binding site. As one of skill in the art will appreciate, the ribosome binding site is a sequence of nucleotides upstream of the start codon that function to recruit a ribosome to initiate translation. An exemplary ribosome binding site has the sequence, GCCACC. The 3'-UTR region may correspond with the native coronavirus 3'-UTR, or may incorporate one or more modifications from the native sequence which do not adversely impact its function to facilitate translation and to provide transcript stability. For example, the native 3'-UTR may be replaced with the 3'-UTR of another protein. Examples of 3'-UTRs that are suitable for use in the present transgene include, but are not

limited to, one or more of the 3'-UTR of alpha-1-globin, beta-globin, c-fos, metallothionein IG, lipoprotein lipase, glutathione peroxidase 4, glutathione peroxidase 3, glutathione peroxidase 2, c-myc, 15-lipoxygenase, and transferrin receptor 2a. In one embodiment, the 3'-UTR of either or both of alpha-1-globin and beta-globin are included in the present transgene. These 3'-UTR sequences may be used in their native form or modified forms, e.g. including one or more nucleotide modifications that do not adversely affect their function. For example, 3'-UTR of α -1-globin may be modified to delete a "T" residue at position 18, and/or the 3'-UTR of β -globin may be modified to replace the 'TAATAA' sequence with 'GTTGCC' to avoid overlap with the poly-A site. The 3'-UTR region also comprises a poly-A sequence having 50-150 A residues, for example, 100 to 140 A residues, or preferably 120-130 A residues. The 3'-UTR may also comprise an upstream and downstream spacer sequence. In one embodiment, the upstream spacer of the 3'-UTR is a 27 nucleotide sequence and the downstream 3'-UTR sequence is 3 nucleotides in length. In another embodiment, the 3'-UTR incorporates the alpha-1-globin and beta-globin 3'-UTR regions, a 130 residue poly A tail, and spacers at its upstream and downstream ends. An exemplary 3'-UTR is shown in FIG. 7.

For use in an mRNA vaccine, the DNA transgene construct is prepared as described, and mRNA is synthesized therefrom by in vitro transcription of the cDNA template, typically plasmid DNA (pDNA), prepared using methods known in the art. Transcription of the cDNA template is conducted using RNA polymerase that corresponds with the promoter incorporated within the transgene, i.e. T7 RNA polymerase when the T7 promoter is incorporated within the transgene, or Sp6 RNA polymerase when the Sp6 promoter is incorporated within the transgene, etc. For stability and efficient translation, the resultant mRNA strand will include a 3' poly(A) tail, as well as 5' and 3' untranslated regions (UTRs) flanking the coding region, as previously described.

The transcript may also be subjected to post-transcriptional processing to optimize the transcript for therapeutic use. For example, the transcript may be modified to include a 5' mRNA cap to provide protection against enzymatic degradation.

The mRNA vaccine is then formulated for administration. In this regard, mRNA may be complexed with agents or adjuvants which prevent degradation, enhance uptake and promote translation. Examples of such adjuvants include, but are not limited to, cationic polypeptides (e.g. protamine), nanoemulsions, carrier peptides, lipid nanoparticles, liposomes, and immune activator proteins (e.g. CD70, CD40L, TLRs). As will be appreciated by one of skill in the art, the vaccine, and any adjuvants, are administered in a suitable carrier, such as saline or other suitable buffer.

In one embodiment, the mRNA vaccine is formulated for administration in lipid nanoparticles. The lipid nanoparticles will generally comprise: an ionizable cationic lipid (whose positive charge binds to the negatively charged mRNA), a PEGylated lipid (for stability), a phospholipid (for structure), and cholesterol (for structure). The ionizable cationic lipid may be any suitable ionizable cationic lipid such as, but are not limited to, C12-200 (1,1'-((2-(4-(2-((bis (2) amino) ethyl) (2-hydroxydodecyl) amino) ethyl)-piperazin-1-yl) ethyl) azanediy) bis (dodecan-2-ol)); cKK-E12 (3,6-bis (4-(bis (2-hydroxydodecyl)-amino)v butyl) piperazine-2,5-dione); MC3 ([[(6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl]4-(dimethylamino)butanoate]; DOTMA (1,2-di-O-octadecenyl-3-trimethylammonium propane); DDAB (N-decyl-N,N-dimethyldecyl ammonium bromide)N-

decyl-N,N-dimethyldecan-1-ammonium bromide); DOTAP (1,2-dioleoyloxy (trimethylammonium) propane); DC-cholesterol ([10,13-dimethyl-17-(6-methylheptan-2-yl)-2,3,4,7,8,9,11,12,14,15,16,17-dodecahydro-1H-cyclopenta[a]phenanthren-3-yl] N-[2 (dimethyl amino) ethyl] carbamate; hydrochloride); GL67 ([[(3S)-10,13-dimethyl-17-(6-methylheptan-2-yl)-2,3,4,7,8,9,11,12,14,15,16,17-dodecahydro-1H-cyclopenta[a]phenanthren-3-yl] N-(3-aminopropyl)-N-[4-(3-aminopropylamino)butyl] carbamate); and DODMA (N,N-dimethyl-2,3-bis[(Z)-octadecenoxy]propan-1-amine). Phospholipids suitable to provide structure to the lipid nanoparticle include phosphatidylcholines such as distearoylphosphatidylcholine (DSPC) ([[(2R)-2,3-di(octadecanoyloxy)propyl] 2-(trimethylazaniumyl)ethyl phosphate), POPC ([[(2R)-3-hexadecanoyloxy-2-[(Z)-octadec-9-enoyl]oxypropyl] 2-(trimethylazaniumyl)ethyl phosphate), DMPC ([[(2R)-2,3-di(tetradecanoyloxy)propyl] 2-(trimethylazaniumyl)ethyl phosphate), DPPC ([[(2R)-2,3-di(hexadecanoyloxy)propyl] 2-(trimethylazaniumyl)ethyl phosphate) and mixtures thereof. Suitable PEGylated lipids for inclusion in the lipid nanoparticle include, but are not limited to, 1,2-dimyristoyl-rac-glycero-3-methoxypolyethylene glycol-2000 (1,2-DMG PEG2000), 1,3-DMG PEG2000, and mixtures thereof. The lipids are combined in suitable molar ratios to provide the required structure and stability. In one embodiment, the nanoparticle includes the lipids, MC3; DSPC; cholesterol; and DMG-PEG2000 in a molar ratio of MC3:DSPC:cholesterol:PEG of 45:55:8:1:35:40:1:2, such as a ratio of 50:10:38.5:1.5.

To prepare the present nanoparticles, the mRNA in a suitable buffer is combined with the lipids at a ratio of about 3:1 to 7:1 to encapsulate the mRNA. The nanoparticles were treated to obtain a stabilized formulation comprising nanoparticles in the range of 40-180 nm, preferably 70-100 nm, 70-90 nm, or 80-90 nm. For example, the nanoparticles were dialyzed, or buffer exchanged using tangential flow filtration (TFF), to achieve stabilized nanoparticles. Stabilization of the nanoparticles is achieved by a selected treatment for at least about 5 minutes to 24 hours using suitable reagents, including HEPES (2-[4-(2-hydroxyethyl)piperazin-1-yl] ethanesulfonic acid), acetate buffer, phosphate buffered saline (PBS) and Tris (trisaminomethane)-HCl. For example, such as 1 to 100 mM HEPES (pH range: 6.0 to 8.0), 1 to 100 mM Acetate (pH range: 4.0 to 7.5), 1x to 10xPBS (pH range: 6.0 to 8.0) and 1 to 50 mM Tris-HCl (pH range: 6.0 to 8.0), and combinations thereof may be used. The preferable combinations are: 1 to 100 mM HEPES (pH range: 6.0 to 8.0) and 1 to 100 mM Acetate (pH range: 4.0 to 7.5), 1x to 10xPBS (pH range: 6.0 to 8.0) and 1 to 100 mM Acetate (pH range: 4.0 to 7.5), 1 to 50 mM Tris-HCl (pH range: 6.0 to 8.0) and 1 to 100 mM Acetate (pH range: 4.0 to 7.5), 1 to 100 mM HEPES (pH range: 6.0 to 8.0) and 1x to 10xPBS (pH range: 6.0 to 8.0), 1 to 100 mM HEPES (pH range: 6.0 to 8.0) and 1 to 50 mM Tris-HCl (pH range: 6.0 to 8.0), 1 to 100 mM HEPES (pH range: 6.0 to 8.0) and 1 to 100 mM Acetate (pH range: 4.0 to 7.5) with 1x to 10xPBS (pH range: 6.0 to 8.0), 1 to 100 mM HEPES (pH range: 6.0 to 8.0) and 40 to 60 mM Acetate (pH range: 4.0 to 7.5) and 10 to 40 mM Tris-HCl (pH range: 6.0 to 8.0) for at least 5 minutes to 24 hours. More preferably, 40 to 60 mM HEPES and 40 to 60 mM acetate (pH range 6.3 to 7.0), 1x PBS and 40 to 60 mM Acetate (pH range: 6.0 to 7.2), 10 to 40 mM Tris-HCl and 40 to 60 mM Acetate (pH range: 6.0 to 7.5), 40 to 60 mM HEPES and 1xPBS (pH range: 6.0 to 8.0), 40 to 60 mM HEPES and 10 to 40 mM Tris-HCl (pH range: 6.5 to 7.5), 40 to 60 mM HEPES and 40 to 60 mM Acetate and 1xPBS (pH range: 6.5 to 7.0), 40 to 60 mM

HEPES and 40 to 60 mM Acetate and 10 to 40 mM Tris-HCl (pH range: 6.8 to 7.5), for at least 5 minutes to 4 hours.

For clarity, the term "acetate buffer" as used herein refers to sodium acetate, potassium acetate, calcium acetate, magnesium acetate, lithium acetate, barium acetate, cesium acetate, iron acetate, aluminium acetate, zinc acetate, copper acetate, ammonium acetate, triethyl ammonium acetate, diethyl ammonium acetate and other suitable acetate salts of inorganic and organic cations/bases and combinations thereof.

Following stabilization, the final nanoparticle product is combined with a suitable solution for administration, e.g. an aqueous solution such as a saline, buffered saline or Tris or mixtures thereof, adjusted to a pH of 7.2+/-0.2. Preferably, the solution comprises sodium chloride (pH range: 7.2+/-0.2), PBS (pH range: 7.2+/-0.2), and Tris-HCl (pH range: 7.2+/-0.2). The nanoparticle solution may additionally comprise stabilizing agents, such as carbohydrates, glycerol, vitamins, antioxidants, anti-reducing agents, and anti-hydrolyzing agents, to maintain the desired particle size during storage. Exemplary stabilizing agents include sucrose, trehalose, mannitol, PEG, propylene glycol, sorbitol, L-arginine, vitamin-A, vitamin-B, vitamin-C, vitamin-E, iron, aluminium, zinc, calcium, magnesium, sodium, potassium, glycerol and combinations thereof, in amounts of 3-20% by wt.

The DNA transgene construct is then generally adapted for administration. The transgene construct may be formulated for administration as a linear molecule, covalently-closed linear construct or mini-circle. Alternatively, the transgene construct may be incorporated into a vector such as a plasmid or cosmid using techniques well-known in the art and then formulated for administration. The resulting DNA vaccine is formulated for administration. The vaccine may be incorporated within a delivery system adapted to enhance immunogenicity of the vaccine, for example, biodegradable polymeric microparticles (e.g. chitosan, polylactide-co-glycolides, polyethyleneimine, amine-functionalized polymethacrylates, cationic poly(β -amino esters), poloxamers and polyvinylpyrrolidone polymers) or liposomes. Additional delivery systems may also be used such as calcium phosphate, calcium apatite, or calcium hydroxy apatite. The vaccine may also be combined with an adjuvant to enhance immunogenicity, e.g. inorganic compounds such as aluminium-containing compounds and squalene, oils such as paraffin, bacterial products such as toxoids, plant saponins, cytokines such as IL-1, IL-2 or IL-12, a cytosine phosphoguanine (CpG) motif-containing adjuvant, or an adjuvant combination such as Freund's adjuvant.

Viral-vectored vaccines may also be utilized to administer the present transgene construct, including both DNA viral vectors and RNA viral vectors. DNA viral vector vaccines are adapted to expressibly incorporate the present DNA transgene construct, e.g. under the control of a viral promoter. Examples of suitable DNA viruses for use as vaccines include, but are not limited to, poxviruses such as vaccinia virus and modified vaccinia virus, adenoviruses, adeno-associated viruses, herpes simplex virus and cytomegalovirus, and including various serotypes thereof, both replication-competent and replication-deficient. RNA viral vector vaccines may also be adapted to expressibly incorporate an appropriate transcript of the present transgene construct, e.g. positive or negative strand. Examples of suitable RNA viruses for use as a vaccine to deliver the transgene include, but are not limited to, vesicular stomatitis viruses, retroviruses such as MoMLV, lentiviruses, Sendai viruses, measles-derived vaccines, Newcastle disease virus, alphaviruses

11

such as Semliki Forest virus, flaviviruses, or an RNA replicon based on an RNA virus (i.e. derived from alphavirus, flavivirus, etc).

The present vaccine is used in a method of vaccinating a mammal against infection by a coronavirus. The vaccine is administered to the mammal in a therapeutically effective amount, i.e. an amount sufficient to generate in the mammal an immune response. The term "mammal" is used herein to refer to both human and non-human mammals. As one of skill in the art will appreciate, the amount required to generate an immune response will vary with a number of factors, including, for example, the particular transgene/antigens in the vaccine, the vector used to deliver the vaccine, and the mammal to be treated, e.g. species, age, size, etc. The dosage may also vary based on the form of the vaccine, i.e. DNA vaccine vs. mRNA vaccine, as well as the formulated form of the vaccine. In this regard, suitable dosages may be determined using appropriate well-established means. It is expected that a dosage of to 300 µg, for example 10-100 µg, of the mRNA formulated in lipid nanoparticles will generate a suitable immune response in humans.

The present vaccine is administered to a mammal to prevent a coronavirus infection in any one of several administrable routes including, but not limited to, parenteral administration such as intravenously or intramuscularly, intranasally or by inhalation. For nucleic acid-based vaccines, other techniques such as administration by electroporation or using gene gun technology may be utilized.

The vaccine may be administered using a prime/boost protocol. In this regard, the prime and boosting vaccines, which may be the same or different vaccine type (e.g., both the prime and boosting vaccine may be a nucleic acid-based vaccine, or both may be a viral vectored vaccine, or the prime vaccine may be nucleic acid-based and the boosting vaccine may be a viral vector, or vice versa) are administered such that the boost is administered once the immune response to the prime dose has reached a suitable level, e.g., a period of time of about 1-26 weeks following the prime dose. The prime and boosting doses may be administered by the same or different administrable routes.

In an embodiment, the present vaccine is formulated as an mRNA vaccine in a lipid nanoparticle and is administered in two doses, i.e., administration of a prime (first dose) followed by administration of a boost (2nd dose), within about 4-12 weeks, so as to generate a sufficient immune response.

In another embodiment, the present vaccine is formulated as an mRNA vaccine in a lipid nanoparticle and is administered using a single administration, i.e., administration of a boost is not required to generate a sufficient immune response.

The present vaccine advantageously results in a vaccine that generates a strong immune response and neutralization efficacy against SARS-CoV-2 variants, including the Alpha, Beta, Gamma and Delta variants. To date, a vaccine targeting such SARS-CoV-2 variants has not been developed. The present vaccine is based on a transgene designed to yield high levels of mRNA vaccine, with enhanced stability, processability and little or no toxicity. In addition, in an embodiment, the mRNA vaccine may be provided as a lipid nanoparticle vaccine which provides efficient mRNA release on administration and uptake by antigen-processing cells. The present vaccine is confirmed to elicit neutralizing antibodies and a beneficial balanced cellular response (e.g. Th1:Th2 is approximately equal, i.e. about 1:1).

12

Embodiments of the invention are described by reference to the following specific Example which are not to be construed as limiting.

Example

The following work describes the development of an mRNA vaccine in accordance with an embodiment of the present invention.

Methods

Target gene and vector cloning—The target DNA was identified using data mining and analysis using bioinformatics. The target DNA was amplified from a patient sample, sequence confirmed, modified to achieve the desired design architecture such that it harbors a suitable 5' UTR, ORF to express the S protein with R452, Y501, G614 and double proline (2P) mutations (K986P and V987P) with an IgE-90 secretory signal sequence, a special 3' UTR constructed with modified alpha and beta globin in tandem, and finally a 130 residue-long poly-A tail. For overexpression of S protein, a GC-rich codon optimized S gene was prepared using in-house developed using a polymerase chain assembly method. After assembly of the S gene, it was replaced with the above-mentioned construct.

Target selection—As of March 2020, there were 170 surface glycoproteins (partial and complete sequence) out of 1661 SARS-COV-2 proteins posted on NCBI Virus database. A comparative sequence alignment using Clustal Omega (ebi.ac.uk/Tools/msa/clustalo/) showed differences in several regions, notably in position 614 (D>G). A total of 15 glycine containing surface glycoproteins were found. A consensus sequence from multiple sequence alignment was identified using EMBOSS Cons (ebi.ac.uk/Tools/msa/emboss_cons/) and selected as a primary target sequence for vaccine development. Hydrophilicity/hydrophobicity plot analysis was performed using GENETYX Ver8.2.0, protein 3D modeling using Phyre2 and visualized using UCSF Chimera 1.11.2rc. The R452, Y501, D614G and double proline (2P) mutations (K986P and V987P) were incorporated into the target sequence.

Target amplification—Nasopharyngeal and oropharyngeal swab sample were collected from a COVID-19 positive male patient. Virus heat inactivation at 56° C. for 30 minutes and total RNA including virus RNA extraction was performed using TRIzol™ Plus RNA Purification Kit (ThermoFisher, USA). cDNA synthesis was performed using GoScript™ Reverse Transcription System (Promega, USA). S-gene (Surface glycoprotein) was amplified using 3 different sets of primers and Platinum™ SuperFi™ DNA Polymerase (ThermoFisher, USA). Amplified S-gene and polymerase chain reaction (PCR) engineered pET31b(+) (Novagen, Germany) bacterial expression vector were amplified, excised and extracted from agarose gel using GeneJET Gel Extraction and DNA Cleanup Micro Kit (ThermoFisher, USA), and assembled together using NEBuilder® HiFi DNA Assembly Master Mix (NEB, USA). Sub-cloning was performed into DH5α chemical competent cells, miniprep purification was using PureLink™ Quick Plasmid Miniprep Kit (ThermoFisher, USA). S-gene integration check into vector was performed via restriction digestion using XbaI (ThermoFisher, USA) and EcoRI (ThermoFisher, USA), and PCR using primers. DNA sequencing was performed to confirm the complete open reading frame (ORF) compatibility of target S-gene. Finally, sequence confirmed rDNA (rDNA ID: p20004) was further

amplified and purified using PureLink™ HiPure Plasmid Midiprep Kit (ThermoFisher, USA), sequenced, and stored for future purposes. Also, sequence confirmed S-gene was submitted to NCBI (GenBank accession number MT676411.1), where we identified and noted D614G mutation. Supplier's manual with minor modifications were followed for all the methods.

Target modification—An immunoglobulin (Ig) heavy chain (HC) 19 amino acid signal peptide (H1) was assembled, and amplified along with homology arm for incorporating into rDNA p20004, replacing native 13 amino acid leader sequence. Assembled signal peptide was amplified with homology arm and rDNA p20004 was engineered via PCR. New rDNA p20006 was prepared by incorporating signal peptide and engineered p20004 rDNA, using above explained method as p20004 rDNA preparation. S-gene was amplified from rDNA p20006. This gene and pcDNA™5/FRT Mammalian Expression Vector (ThermoFisher, USA) were digested using Acc65I (ThermoFisher, USA) and XhoI (ThermoFisher, USA) and visualized via agarose gel electrophoresis. The desired bands from the gel were excised and purified using GeneJET Gel Extraction and DNA Cleanup Micro Kit and ligated using T4 DNA Ligase (ThermoFisher, USA). After ligation, sub-cloning into DH5α chemical competent cells, plasmid miniprep purification, insert checking, DNA sequencing, plasmid midiprep purification, DNA sequencing and storage (rDNA ID: p20010) were performed. Amino acid mutations at position 452 (L452R), 501 (N501Y), and 2P (double Proline) amino acid mutations at position 986 (K986P) and 987 (V987P) were also performed via site directed mutagenesis. DNA sequencing was performed to confirm desired mutations (rDNA ID: p20015). Finally, a T7 promoter sequence, a synthetic 5'-UTR, an IgE signal peptide replacing native 13 amino acids signal peptide from S-gene, a 3'-UTR (modified alpha globin and modified beta globin), and a 130 bp synthetic poly A-tail (pA-tail) were added. A restriction endonuclease (Sfo I) sequence before T7 promoter sequence and after pA were added for cutting out desired size of DNA for in vitro mRNA synthesis (rDNA ID: p20020.2). A GC-rich codon optimized S gene was assembled via in-house developed polymerase chain assembly method for the purpose of overexpression of S protein. After assembly and amplification of S gene, it was replaced with natural S gene from the rDNA p20020.2 and generated a new rDNA p21004. Final rDNA ID was p20020, p20020.3, p20020.2, p21005, p21006, p21004, p21009 and rDNA construction was performed as mentioned before for p20004 and p20006. Supplier's manual with minor modifications were followed for all the methods.

Sequencing—DNA sequencing was performed as according to supplier's protocol for the final construct p20020, 20020.3, p20020.2, p21005, p21006, p21004, p21009 and other constructs e.g., p20004, p20006, p20010, p20013, p20013.1, p20015, p20026, p20026.1, p20030, p20030.1, p20031, p20031.1, p21010, p21011, p21012, etc. using 3500 Genetic Analyzer (ThermoFisher, USA). DNA sequencing data clearly confirmed the presence of the target sequences and modifications. BigDye® Terminator v1.1 Cycle Sequencing Kits (ThermoFisher, USA) and POP-6 polymer chemistry (ThermoFisher, USA) was used for DNA sequencing reaction.

mRNA production—The in vitro transcription (IVT) of mRNA was performed in the presence with 3'-O-Me-m7G (5)ppp(5')G RNA Cap Structure Analog and S-adenosylmethionine (NEB, USA) using MEGAscript™ T7 Transcription Kit (ThermoFisher, USA), and Ribonucleotide Solution

Set (NEB, USA). IVT mRNA synthesis reaction was optimized into 4 steps (step 1: synthesis time factor, step 2: rNTPs concentration, step 3: RNase inhibitor and pyrophosphatase effect, and step 4: temperature dependency); final concentration of ribonucleotides was as follows: ATP and UTP—13.13 mM, and GTP and CTP—9.38 mM. The reaction was run for 2 hours at 37° C. followed by a DNase treatment at 37° C. for 15 minutes and dephosphorylation using Antarctic Phosphatase (NEB, USA) according to supplier's manual. IVT capped mRNA was purified using phenol:chloroform:isoamyl alcohol, and purified using MEGAclear™ Transcription Clean-Up Kit (ThermoFisher, USA).

mRNA identification—Capped mRNA, purified mRNA, formulated lipid nanoparticles (LNPs) and formulated LNPs, treated with RNase samples, were analyzed by size exclusion chromatography (SEC). SEC was performed in Ultimate 3000 (ThermoFisher, USA) system using 10 mM Disodium hydrogen phosphate (Wako, Japan), 10 mM Sodium dihydrogen phosphate (Wako, Japan), 100 mM Sodium chloride (Merck, Germany), pH 6.6 as mobile phase. Biobasic SEC-300 (300×7.8 mm, particle size; 5 μm, ThermoFisher, USA) column was used with 1.0 mL/minute flow rate, 260 nm wavelength, 10 μL sample injection volume for 20 minutes.

Formulation of mRNA—Purified mRNAs were first diluted with sodium acetate buffer to the desired concentration. The lipid molecules were dissolved in ethanol and mixed well. Lipids (MC3:DSPC:cholesterol:DMG-PEG2000) were combined in a molar ratio of 50:10:38.5:1.5. Then, sodium acetate buffer containing mRNA and the lipid samples were mixed at a ratio of 3:1 to 7:1 and passed through the liposome extruder (Genizer, USA) or were made with another suitable liposome technology, for example, T-mixture, Y-mixture, Herringbone mixture, diffusion film mixture etc. to encapsulate the mRNA. The size distribution was checked after encapsulation of mRNA into nanoparticles. Then, the samples were stabilized through dialysis or other suitable technologies for example, TFF or dilution against 40 to 60 mM HEPES and 40 to 60 mM sodium acetate (pH range 6.5 to 6.9) for 3.0 hours. After completion of stabilization, samples were formulated through dialysis against 1×PBS (pH range: 7.2±0.2) or 20 mM Tris-HCl (pH range: 7.2±0.2). To maintain the desired LNP size during storage, 7 to 10% by wt sucrose was added. The size distribution was checked by Zetasizer Nano ZSP (Malvern Panalytical, UK). LNP samples were analyzed for size distribution in respective buffers. The formulated sample (referred to as GBPD060) was concentrated using Ultra centrifugal filters or TFF membrane and passed through 0.22 micron filter, and stored at 5±3° C., -30±5° C., -80±5° C. The formulation was confirmed by quality control tests for particle size distribution, encapsulation efficiency, endotoxin limit and sterility.

Animal management and vaccination—The study procedures were performed according to local and international regulation. A total of 50 BALB/c Swiss albino mice (male and female) 6-8 weeks old were selected and isolated 5 days before immunization. After careful observation and conditioning, suitable numbers of mice were selected. The temperature in the experimental animal room was 26° C. (±2° C.) and the relative humidity was 60±5%. The room was HVAC controlled ISO class 7 room with 70% fresh air intake and full exhaust. The mice were individually housed in cages with proper water and feed, and kept under 12 hours of day-night cycle. O30 mice were separated into 5 different groups consisting of 6 mice (3 males and 3 females) in each

15

group. GBPD060 formulations and vehicle were administered intramuscularly (IM) in the quadriceps for treatment and placebo, respectively. Each mouse of group 1, 2 and 3 was immunized with sterile 0.1 µg/50 µL, 1 µg/50 µL, 10 µg/50 µL of GBPD060, respectively. Second dose was administered 21 days later. The sample sizes were determined by the resource equation method.

Local tolerance—Local tolerance was confirmed by clinical signs, and macroscopic and histopathology evaluations of injection sites in animals. Euthanasia and evaluation of lesions was performed in one representative mouse from placebo and control group and 3 from the treatment group at 132 hours post treatment. The inner thigh muscle of injected site of each mouse was excised and placed in 10% neutral buffered formalin until adequately fixed. After trimming, processing and paraffin embedding, the sections (6 µm) are HE stained and observed for erythema and edema under microscope.

Immunogenicity—Approximately 200 µL blood was collected from facial vein and centrifuged at 1500×g for 138 serum isolation (10 minutes at 4° C.). All serum was aliquoted, frozen immediately and stored at -80° C. until analysis. The reactivity of the sera from each group of mice immunized with GBPD060 was measured against SARS-CoV-2 S antigen (SinoBiologicals, China). The serum IgG binding endpoint titers (EPTs) were measured in mice immunized with GBPD060. EPTs were observed in the sera of mice at day 7 and day 14 after immunization with a single dose of GBPD060.

Toxicity—Whole blood (~50 µL) from each mouse was collected in 2% EDTA at 3 days pre-immunization and 14 days post-immunization. Complete blood count (CBC) analysis was measured using an auto hematology analyzer BK-6190-Vet (Biobase, China). Samples were used for blood-chemistry analysis viz., alanine transferase (ALT), aspartate transaminase (AST) and blood urea nitrogen (BUN) using semi-automatic chemistry analyzer (Biobase, China).

Pseudovirus preparation and in vitro neutralization—Adenovirus and retrovirus-based pseudovirus were prepared for SARS-CoV-2 in vitro and in vivo neutralization assay. Pseudotyped-SARS-CoV-2 adenovirus was prepared expressing the SARS-CoV-2 surface glycoprotein gene (S gene) on the virus. S gene of SARS-CoV-2 was cloned into pAADV-B02 vector (Genemedi, China) that also contains a GFP gene downstream of the gene of interest. Site directed mutagenesis was performed to incorporate Q498T and P499Y mutations in the S-gene for the purpose of homotypic pseudovirus preparation. After construction, SARS-CoV-2 S gene containing rDNA and adenovirus backbone plasmid pAADV-C01 (Genemedi, China) were co-transfected into HEK293-based adapted viral production cell (VPC) (ThermoFisher, USA). VPCs were seeded in a 6-well TC-treated plate (Nest, China) at a concentration of 6×10^5 cells/well and cultured overnight. Co-transfection were performed using Lipofectamine 3000 (ThermoFisher, USA) reagent according to manufacturer's protocol. Next day 1.25% low melting agarose in DMEM media was spread on the well and incubated until plaques were formed. After formation of plaques, multiple plaques were collected in DMEM media and titers were measured for plaque selection. Then selected plaque was added on the fresh VPC. After few days, cells and supernatant were collected and repeated freeze-thawing was performed for collection of viruses (P1 pseudovirus). These processes were repeated and P4 pseudoviruses were collected; concentration and purification were performed by

16

ultracentrifugation and sucrose gradient. After titer determination, pseudoviruses were stored at -86° C. (ThermoFisher, USA).

Another pseudotyped SARS-CoV-2 retrovirus was prepared as follows. S gene was cloned into pMSCV_Neo vector (TakaRa Bio, USA). After preparation of S gene-expressing plasmid, co-transfection was performed on VPC. pMD2G and pSPAX2 (Genemedi, China) packaging plasmid were used for retro-based pseudovirus preparation. 9×10^6 cells were seeded in a 75 cm² TC-treated flask and cultured overnight. Co-transfection was performed using Lipofectamine 3000 reagent. Media was replaced with complete DMEM after 6 hours. Media was collected after 48 hours and stored at 4° C. 12 mL media was added into the flask, collected the next day and pooled with previously stored media. Then concentration and purification were performed by ultracentrifugation. Pseudoviruses were stored at -86° C. (ThermoFisher, USA) after titer determination.

ACE2-overexpressing HEK293 cells (Innoprot, Spain) were seeded in two 96-well TC-treated plate at a concentration of 2.2×10^4 cells/well and incubated overnight. One plate was used for adeno-based pseudovirus and the other plate for retro-based pseudovirus, respectively. Two separate plates were used for serum preparation. Different rows of the plate were used for different groups, such as A1-A10 for treatment group, B1-B10 for placebo, C1-C10, D1-D10 E1-E10 and F1-F10 for control, CR3022, commercial anti spike and only cell group (negative control). Sera from different mice of the same group were collected and pooled for neutralization assay. 10 µL sera from vaccinated mice was added in 90 µL complete DMEM media, and were serially 2-fold diluted up-to 9 times. 1.2×10^5 pseudovirus in 50 µL was added into different wells that contained serially diluted serum and mixed properly. The SARS-CoV-2 pseudovirus and serum mixture was incubated for 1.5 hour at 37° C. Then, 100 µL of pseudovirus and serum mixture was transferred on pre seeded cells. 5 µg/mL poly L-lysine (Wako, Japan) was added into each well for enhancing the transduction. Then, incubation was performed at 37° C. for 48 hours and GFP-fluorescence were measured using Varioskan LUX (ThermoFisher, USA). Number of virus particle inside the cells were determined by qPCR (QuantStudio 12K Flex, ThermoFisher, USA). For retro-based neutralization assay, qPCR was used to analyze the copy number of S gene in cell. Genomic DNA was extracted by MagMAX Express-96 Standard (ThermoFisher, USA) using Magmax DNA multi-sample ultra-kit. (ThermoFisher, USA) and used for determination of S gene copy number by qPCR.

In vivo neutralization—18 albino male mice 6-8 weeks of age were selected and divided into 6 groups, 1 control and 5 treatment, comprising 3 male mice in each group. The control group mice were immunized intramuscularly with 50 µL of placebo and treatment group mice were immunized with 1 µg/50 µL of GBPD060 vaccine. GFP pseudo-typed homotypic SARS-CoV-2 adenovirus were sprayed in the nasopharynx on 21-day post immunization. Nasopharynx and lung aspirate samples from mice were collected and analyzed for viral copy number using qPCR at indicated time point. Animals were sacrificed and lung sections were performed and microscopic slides were prepared for fluorescence imaging (GFP) to detect viral load.

Antibody analysis by ELISA—Serum from the mice of different groups were analyzed by standard enzyme-linked immunosorbent assay (ELISA) to determine antibody titers. ELISA plate (Corning, USA) was coated with 1 µg/mL SARS-CoV-2 Spike S1+S2 ECD-His recombinant protein (Sino Biological, China) in Dulbecco's phosphate-buffered

saline (DPBS) (ThermoFisher, USA) for 2 hours at room temperature. Plates were washed three times with DPBS+ 0.05% Tween 20 (Scharlau, Spain) and then blocked with PBS+1% BSA (ThermoFisher, USA)+0.050% Tween-20 for 2 hours at 37° C. The plates were washed 3 times then incubated with sera and SARS-CoV-2 Spike antibody (Sino Biological, China) for 2 hours at 37° C. After washing 3 times, the plates were then incubated with HRP conjugated Goat anti-Mouse IgG (H+L) secondary antibody, (ThermoFisher, USA) for 50 minutes at room temperature. Final washing was done 3 times and then the plates were developed for colorimetric detection reaction with Pierce TMB substrate (ThermoFisher, USA) for 10 minutes. The reaction was stopped with 1N hydrochloric acid (HCl) and signals were measured at 450 nm wavelength within 30 minutes. For isotype analysis, Pierce Rapid ELISA Mouse mAb Isotyping kit (ThermoFisher, USA) was used. Serum samples from 4 subjects of treatment 2 and 3 were analyzed. All the steps were followed as per supplier's instructions.

Antibody binding affinity—The BIAcore T200 equipment (GE Healthcare, USA) and Amine coupling kit (GE Healthcare, USA) were used for immobilization of SARS-CoV-2 Spike S1+S2 ECD-His recombinant protein (Sino Biological, China) in Series S Sensor Chips CM5 (GE Healthcare, USA). First, the flow cell surface of Series S Sensor Chips CM5 was activated by injecting a mixture of EDC/NHS (1:1) for 7 minutes. Then 70 μ L of 50 μ g/mL S1+S2 protein was prepared in sodium acetate at pH 5.0 and injected over the activated surface at 10 μ L/min flow rate. Residual NHS-esters were deactivated by a 70 μ L injection of 1 M ethanolamine, pH 8.5. The immobilization procedure was performed by using running buffer HBS-EP, pH 7.4 (GE Healthcare, USA). Sera at 14-day post immunization were used for the experiment; 5 samples, each containing either 1 μ L sera or MabSelect resin- (GE Healthcare, USA) pull-down sera (IgG) were analyzed using surface plasmon resonance (SPR) to analyze the binding affinity of the antibody pool. Samples (1 μ L) were pretreated with 5 μ L of either buffer or SARS-CoV-2 S or SARS-CoV-2 S1 or SARS-CoV-2 S2 protein (500 μ g/mL) at 25° C. for 15 minutes. All samples were diluted in 1xHBS-EP (pH, 7.4) running buffer. Samples were applied through the active flow cell surface of CM5 chip for binding analysis. Glycine-HCl of pH 2.5 was used for regeneration of the chip.

SARS-CoV-2 surface glycoprotein peptide pool generation and mapping—40 μ g of SARS-CoV-2 Spike S1+S2 (S) ECD His recombinant protein (Sino Biological, China), S2 ECD-His Recombinant Protein (Sino Biological, China), and RBD(S1)-His Recombinant Protein (Sino Biological, China) were diluted in 50 mM ammonium bicarbonate (Wako Pure Chemicals Industries Ltd., Japan), pH 8 containing 8 M urea (ThermoFisher Scientific, USA). 500 mM DTT (ThermoFisher Scientific, USA) was added to the solution to a final concentration of 20 mM (1:25 dilution) mixed briefly, and incubated at 60° C. for 1 hour. For alkylation, 1 M IAA (Sigma-Aldrich, USA) was added to the reduced protein sample to a final concentration of 40 mM (1:25 dilution), and the reaction mixture was incubated for 30 minutes protected from light. The reaction was stopped by adding 500 mM DTT solution to a final concentration of 10 mM (1:50 dilution). Enzymes (ThermoFisher Scientific, USA) were added to the sample solution to a final trypsin to protein ratio of 1:23 (w/w). Samples were incubated at 37° C. for 16-24 hours. Reaction was stopped by adding formic acid up-to pH 2.0. C18 spin column (ThermoFisher Scientific, USA) was prepared as per manufacturer's instruction to purify the peptide-pool. Columns were activated by adding

200 μ L 50% acetonitrile (Wako Pure Chemicals Industries Ltd., Japan), and equilibrated using 200 μ L 0.5% formic acid (Wako Pure Chemicals Industries Ltd., Japan) in 5% acetonitrile (Wako Pure Chemicals Industries Ltd., Japan). Samples were applied to the column and eluted using 20 μ L 70% acetonitrile. Samples were dried under low temperature and vacuum and processed for experiments.

SARS-CoV-2 S-ECD His recombinant protein (Sino Biological, China), S2-ECD-His Recombinant Protein (Sino Biological, China), and RBD(S1)-His recombinant proteins (Sino Biological, China) were digested with Serine Protease (MS grade, Pierce, ThermoFisher, USA) and purified according to the supplier's instructions (supplementary method). 1 μ g of digested peptides were loaded into mass spectrometry system (Q Exactive Hybrid Quadrupole-Orbitrap MS, ThermoFisher Scientific, USA). For separation of peptides Hypersil gold C18 (100x2.1 mm; particle size: 1.9 μ m, ThermoFisher, USA) column was used. Column oven temperature was set at 40° C. and eluted in 95—60% mobile phase A (0.1% formic acid in water) and 5-40% mobile phase B (0.1% formic acid in acetonitrile) gradient with 0.300 mL/min flow rate for 65 minutes. Peptide elution were checked by UV absorbance at 214 nm. For peptide identification, data dependent mass spectrometry was performed where full-MS scan range was 350 m/z to 2200 m/z, resolution was 70000, AGC target was 3E6, maximum IT was 100 milliseconds (ms). Data dependent mass spectrometry resolution was 17500, AGC target was 1E5 and maximum IT was 100 ms. Data analysis was performed in BioPharma Finder (ThermoFisher, USA) using variable parameters to get confident data, and then data were combined in one map to visualize complete fragmentation.

Mouse splenocyte isolation, peptide stimulation and flow cytometric analysis of Tcell populations—Harvested spleen was taken in RPMI and mashed within the petri dish. Cells were washed out from slides using micropipette. A 10 ml pipette was used to draw the solution up and down, each time closing the end of the pipette against the bottom of the petri dish to forcefully expel the contents and break up the pieces. Cell suspension was passed through a sterile 40 μ m mesh strainer. Centrifugation was performed for 10 minutes at 250xg, at 4° C. Supernatants were discarded and cells were re-suspended in 10 mL PBS. Samples were subjected for FicolI-Paque PLUS processing. Centrifugation was performed for 30 minutes at 400xg at 20° C. Two thirds of the top layer were aspirated and lymphocyte layer was collected using a micropipette. Cells were washed 3 times with PBS, collected in RPMI, and were subjected to experiments.

Harvested splenocyte were cultured at 37° C. and 5% CO2 in RPMI and were either stimulated with S-peptide pool or buffer. After 6 hours, media were collected, cells were washed twice with PBS and incubated for another 12 hours and were considered as 18-hour sample. Samples were collected for cytokine assay and cells were processed for flow cytometric analysis of T cell populations. For day 14 samples, surface antigen and intracellular cytokine staining of cells were performed with following antibodies according to the supplier's instructions: V500-anti-mouse CD45 (BD Bioscience, USA), FITC-anti-mouse CD4 (ThermoFisher, USA), anti-mouse IFN-gamma (ThermoFisher, USA), Alexa Fluor® 594-secondary antibody (ThermoFisher, USA), in-house developed TNF alpha fusion protein, anti Fc primary antibody (ThermoFisher, USA), Alexa Fluor® 594-secondary antibody (ThermoFisher, USA), anti-mouse IL-2 (ThermoFisher, USA), Alexa Fluor® 594-secondary antibody (ThermoFisher, USA), anti-mouse IL-6 (ThermoFisher, USA), Alexa Fluor® 594-secondary antibody

(ThermoFisher, USA) and no live/dead staining. Cells were washed, fixed, permeabilized, stained and stored at 4° C. After 48 hours, cell-events were acquired using an FACS Lyric (BD Biosciences), followed by FlowJo software (FlowJo LLC, Ashland, OR) analysis. For day-91 samples, surface antigen and intracellular cytokine staining of cells were performed with following antibodies as per supplier's instructions: V500-anti-mouse CD45 (BD Bioscience, USA), Alexa Fluor® 700-anti-mouse CD3 (BD Bioscience, USA), APC-Cy™7-anti-mouse CD4 (BD Bioscience, USA), PE-Cy™7-anti-mouse CD8 (BD Bioscience, USA), FITC-anti-mouse IFN- γ (BD Bioscience, USA), 88700-anti-mouse TNF- α (BD Bioscience, USA), BV421-anti-mouse IL-2 (BD Bioscience, USA), APC-anti-mouse IL-4 (BD Bioscience, USA), PE-anti-mouse IL-17A (BD Bioscience, USA) and no live/dead staining. For memory B cell identification, following antibodies were used: V500 anti-mouse CD45 (BD Bioscience, USA), PE-Cy™7-anti-mouse CD19 (BD Bioscience, USA), BV421-anti-mouse CD27 (BD Bioscience, USA) and no live/dead staining. Cells were washed, fixed/permeabilized, stained and cell-events were acquired using an FACS Lyric (BD Biosciences), followed by FlowJo software (FlowJo LLC, Ashland, OR) analysis.

IL-2 and IL-6 were measured by ELISA using standard protocol. ELISA plate (Corning) was coated with 1 μ g/mL IL-2 polyclonal antibody (ThermoFisher, USA) in Dulbecco's phosphate-buffered saline (DPBS) (ThermoFisher, USA) for 2 hours at room temperature. After coating, plate was washed for 3 times with DPBS+0.05 Tween 20 (Scharlau, Spain) and then blocked with PBS+1% BSA (ThermoFisher, USA)+0.050% Tween 20 for 2 hours at 37° C. After blocking, plates were washed for 3 times and incubated with IL-2 and mouse splenocyte samples for 2 hours at 37° C. Plates were then washed again and incubated with IL-2 monoclonal antibody (ThermoFisher, USA) for 2 hours at 37° C. After washing for 3 times, plates were incubated with Goat anti-Mouse IgG (H+L) HRP-conjugated secondary antibody, (ThermoFisher, USA) for 50 min at room temperature. Final washing was done for 3 times and then developed with Pierce TMB substrate (ThermoFisher, USA) for 10 min. Reaction was stopped with 1N HCl. Signals were measured at 450 nm wavelength within 30 min. For IL-6 analysis, IL-6 Mouse ELISA kit (ThermoFisher, USA) was used, steps were performed as per manufacturer instructions.

Results

Bioinformatics analysis to initiate the designing of 'GBPD060'—The analysis started with alignment of available sequences of SARS-CoV-2 spike (S) protein. A total of 15 D614G sequences out of 170 reference sequences of SARS-CoV-2 were found. Hydropathy profile showed a minor variation in the S protein between D614 and G614 genotypes. Relevant 3D modeling suggested that there might be a higher angular strain on G614 than on the D614, which could affect the stability and atomic distance with the neighboring atoms.

Construction, antigen expression and formulation of 'GBPD060'—The ORF for the SARS-CoV-2 spike protein with G614-translating codon was obtained from a confirmed COVID-19 patient through PCR amplification (Accession No. MT676411.1). Necessary modifications were performed to obtain the desired clone in pET31b vector as described in 'Methods' section. The schematic diagram of the target gene and construction scheme are shown in FIG. 1A. The capped-mRNA with a 130-202 nucleotide residue-long poly A tail was obtained. The mRNA sequence with poly A tail was

confirmed by DNA sequencing after reverse transcription (FIG. 1B-1D); Accession No. MWO45214. The IVT process was tuned and validated to obtain the desired mRNA with high yield and quality (FIGS. 1E and 1F). The mRNA was encapsulated in lipid nanoparticle (LNP) ranging from 60-140 nm with a final pH of 7.2 (FIGS. 1G and H).

A pilot study with limited numbers of mice was conducted to identify the suitable mRNA-LNP size for the formulation. mRNA-LNP either smaller than 60 nm or larger than 120 nm did not generate considerable immunological response even with a dose of 10 μ g/mouse. To obtain the best process control for dose production, the mRNA-LNP size range used was set at 85 $\pm$ 20 nm (FIG. 1e). LNP without SARS-CoV-2 S-mRNA was used as placebo.

Local tolerance and toxicity—Control, treatment and placebo groups comprising 3 male mice each were used for local tolerance testing. No detrimental physical consequences of administration were observed such as, local trauma following injection and/or physicochemical actions of the vaccine from local toxicological or pharmacodynamics effects. No sign of erythema or erythredema was observed in muscle tissue section from the site of injection. Complete blood count (CBC) from different groups indicated good health status of mice; all parameters were in the normal physiological range. There were no signs of anemia, infection, inflammation, and bleeding disorder. Liver function tests (LFTs), such as alanine transaminase (ALT) and aspartate aminotransferase (AST) were performed to confirm clinical suspicion of potential liver injury or disease and to distinguish between hepatocellular damage and cholestasis. Blood urea nitrogen (BUN) was tested to evaluate the health of kidneys, such as kidney disease or damage. Data for ALT, AST and BUN were in the normal range and no significant changes were observed between pre-immunization and after immunization.

'GBPD060' induces high and Th-1 biased antibodies against full-length SARS-CoV2 S-protein—Immunization of mice with mRNA-LNP produced specific titer in a dose dependent manner (FIG. 2a). Low dose (0.1 μ g/mouse) immunization produced a moderate level of antibody response (FIG. 2a, Treatment 1). The best antibody response was obtained with 1 μ g/mouse dose (FIG. 2a, Treatment 2). High dose (10 μ g/mouse) immunization produced a higher level of titer with a wide-spread distribution, and therefore, this dose was considered unacceptable (FIG. 2a, Treatment 3).

The subtyping analysis revealed that the titer contained a balanced ratio of IgG2a and IgG1 in a 7-day post immunization sera, and it remains stable for 14-day post immunization sera (FIG. 2b, Treatment 2). A similar trend was observed for (IgG2a+IgG2b) and (IgG1+IgG3) (FIG. 2c, Treatment 2), which has indicated that the antigenic response was CD4+Th1-biased. High dose (10 μ g/mouse) injected mice sera also produced similar response (FIGS. 2b and c, Treatment 3). To check whether the immunization generated antibody pool spanning for the whole antigen or for any specific domain (either S1 or S2), a surface plasmon resonance (SPR) experiment was conducted. The S protein chip recognized high-affinity antibody from the anti-sera pool (FIG. 2d). The response was attenuated significantly for S-protein(s) (S, S1 and S2) pretreated sera (FIG. 2d). S and S1 pretreatment showed strong inhibitory responses, while S2 pretreatment showed a comparatively moderate inhibitory response. The purified Ig from the pooled anti-sera produced a significantly pronounced response (FIG. 2e). The SPR data clearly showed that the vaccination produced a specific antibody pool against the full-length S protein.

Cellular and cytokine responses to 'GBPD060'—The cellular response and induction of specific cytokines in response to vaccination was also characterized after 14 and 91 days of the first immunization. Splenocytes obtained from vaccinated mice were re-stimulated with a library of SARS-CoV-2-S peptide pool. 14 days after first immunization, the stimulated splenocytes generated a significantly higher population of CD4+Th1 cytokine IFN- γ , IL-2 and TNF- α expressing cells (0.21 ± 0.03 , 0.61 ± 256.03 and 0.95 ± 0.11 , respectively) compared to the placebo treated group (0.06 ± 0.03 , 0.18 ± 257.07 and 0.52 ± 0.06 , respectively) (FIG. 3a-3c.). CD4+Th2 cytokine IL-6-expressing cells were moderately increased in stimulated splenocytes of vaccinated mice compared to the placebo-injected mice (0.42 ± 0.08 and 0.23 ± 0.02 , respectively) (FIG. 3d). The amount of secreted cytokines in treated vs. placebo groups were as follows: IFN- γ (treatment group, 559.87 ± 70.76 pg/ml and 303.47 ± 156.53 pg/ml; placebo, 28.29 ± 2.03 pg/ml and 262.16 ± 2.52 pg/ml), IL2 (treatment group, 499.10 ± 30.80 pg/ml and 345.17 ± 22.85 pg/ml; placebo group, 175.71 ± 21.92 pg/ml and 136.87 ± 15.18 pg/ml), IL4 (treatment group, $77.94\pm264\pm7.7$ pg/ml and 46.36 ± 3.7 pg/ml; placebo group, 22.5 ± 3.25 pg/ml and 9.5 ± 1.08 pg/ml) and IL6 (treatment group, 45.78 ± 15.52 pg/ml and 32.61 ± 15.52 pg/ml; placebo group, 266.16 ± 3.53 pg/ml and 14.87 ± 3.08 pg/ml), respectively for 6- and 18-hr, which indicated a vaccine specific response (FIG. 3e-3h).

At 91 days after first immunization, the stimulated (S1 and S2) splenocytes generated a significantly higher population of CD4+Th1 cytokine IFN- γ , TNF- α and IL-2-expressing cells (0.39 ± 0.01 and 0.19 ± 0.08 , 0.6 ± 0.17 and 1.13 ± 0.21 , and 0.07 ± 0.07 and 0.11 ± 271.01 , respectively) compared to the placebo treated group (0.14 ± 0.03 and 0.18 ± 0.08 , 0.36 ± 272.05 and 0.69 ± 0.04 , and 0.07 ± 0.07 and 0.06 ± 0.05 , respectively) (FIGS. 3i and 3j). Similarly, S1 and S2 stimulated splenocytes also generated a significantly higher population of CD8+Th1 cytokine IFN- γ , TNF- α and IL-2-expressing cells (0.91 ± 0.08 and 0.48 ± 0.27 , 0.58 ± 0.25 and 1.3 ± 0.21 , and 0.3 ± 0.08 and 0.05 ± 0.05 , respectively) compared to the placebo treated group (0.24 ± 0.13 and 0.25 ± 0.1 , 0.26 ± 0.15 and 0.55 ± 0.16 , and 0.03 ± 277.03 and 0.03 ± 0.03 , respectively) (FIGS. 3k and 3l). S1 and S2 stimulated splenocytes generated moderate CD4+ and CD8+Th2 cytokine IL-4 and IL-17A-expressing cells compared to the placebo-injected mice (FIG. 3i-3l). Higher levels of sustained Th1 specific cytokine response over Th2 specific cytokine suggested a stable and balanced Th1-biased immunologic response after administration of GBPD060' vaccine. S1 stimulated splenocytes generated a significantly higher population of CD19+CD27+ expressing cells (1.57 ± 0.17) compared to the placebo treated group (0.72 ± 0.12) (FIG. 3m).

'GBPD060' induces neutralization of SARS-CoV-2-S pseudo-type viruses—Sera of vaccinated mice inhibited infection of GFP-expressing pseudo-type SARS-CoV-2-S adenovirus in hACE2-expressing HEK293 (ACE2-HEK293) cells in a dose dependent manner (FIG. 4a). A neutralization assay demonstrated that there was a correlation between the intensity of GFP and SARS-CoV-2 specific antibody for vaccinated mice. A higher concentration of SARS-CoV-2 antibody efficiently neutralized the entry of the pseudovirus into the ACE2-HEK293 cell. The IC50 value for GFP-inhibition was found to be significantly higher for the anti-sera (~ 3 μ g/mL) compared to the CR3022, which is an anti-SARS-CoV monoclonal antibody

(mAb), and a commercially available polyclonal mouse antibody against S-protein (~ 7 μ g/ml).

In parallel with the GFP analysis, to confirm the virion copy number, real-time PCR was implemented as an orthologous method. Copy number analysis was performed using GoTaq® qPCR master mix (Promega, USA) with primers. A 5-fold serially diluted known concentration of pseudovirus was used for standard curve generation. GoTaq® DNA Polymerase activation was done at 95° C. for 10 minutes. Denaturation was done at 95° C. for 10 seconds, annealing at 51° C. for 30 seconds, extension at 72° C. for 30 seconds for 40 cycles. After completion of PCR cycle, melt curve was done for sample integrity checking. The data showed correlation with the GFP and gene copy analysis (FIG. 4b). HIV1-based SARS-CoV-2-S pseudo type virus infection was also significantly inhibited by 1 μ g/mouse dose anti-sera compared to the placebo anti-sera (FIG. 4c, Serum). Either S1 or S2 protein pretreatment nullified the inhibition capacity of anti-sera (FIG. 4c, Serum+S1 and Serum+S2) confirming that the inhibition property for the HIV1-based SARS-CoV-2-S pseudo type virus is related to the vaccination.

It was then determined whether or not the immunization could protect mice from GFP-expressing homotypic pseudo-type adenovirus. Virus was sprayed into the nasopharyngeal space of mice, either in buffer or pretreated with immunized sera. The anti-sera-treated SARS-CoV-2-S adenovirus produced lower copy of virus compared to the buffer-treated virus (FIG. 4d, Treatment 2 and Treatment 1, respectively). The copy number of virus was found to be reduced further from day 2 to day 3 (FIG. 4d, Treatment 4 and Treatment 3, respectively). These data clearly revealed that though the anti-serum exhibited significant inhibitory capacity against viral infection, systemic immune-protection provides enhanced protection. Lower copy number of virus over time indicated that significance of cellular immunity, along with humoral immunity, for efficient viral clearance. The anti-sera treated with S1+S2 protein failed to inhibit SARS-CoV-2-S adenovirus infection in the placebo-injected mice (FIG. 4d, Treatment 5), and confirming that the inhibition and neutralization of the SARS-CoV-2-S pseudo-type virus is correlated with the immunogenic response generated by the GBPD060' vaccine.

The lungs of control- and GBPD060-immunized mice (single-dose, day 14 post-immunization), which were challenged with homotypic SARS-CoV-2 pseudovirus (day-5 post challenge) were collected. GFP-labelled viruses were clearly found spread throughout the lungs of control group mice. Peritracheal viral infection was predominant and suggested that the viral entry pathway is the airways of the lung (FIG. 4e-4g). The immunized mice lungs showed a few viral-spots on day-5 post challenge tissue, which completely disappeared by day-6 of post challenge. No peritracheal viral signal was observed in these lungs (FIG. 4h-4j). The data provided visual and conclusive evidence that GBPD060 immunization neutralizes SARS-COV2 virus and protects against infection by the virus within the lungs.

A transgene comprising a single variation in the S-protein component, i.e. only the G614 variant was also prepared, and used to make an mRNA vaccine as above described. This vaccine exhibited similar protection against infection by SARS-CoV-2.

Discussion

A vaccine targeting the G614 mutant has been developed. The vaccine incorporates the high-expressing spike protein

23

as antigen in a putative perfusion stabilized condition. Comparative design features of 'GBPD060' are shown in Table 1. 'GBPD060' mRNA has features along with the G614-targeted mutation, which are different from other vaccines. The design comprises R452, Y501 and G614 (named 'RYG') in the final construct which is unique compared to any current vaccine. The RYG-variant showed similar biochemical and biological properties to our developed G614 variant. The ribosome binding site, IgE-signal sequence (by replacing the native 13 amino acids from the N-terminal of the SARS-CoV-2 S protein), and the 3' UTR which comprises the 3'UTRs of the alpha and beta globin gene in tandem are features of the vaccine. The vaccine is formulated for delivery in LNPs which are 65-105 nm in size to provide a suitable antibody response. Further, the pH (7.2) of the formulation buffer for the mRNA-LNP vaccine was lower than that in other vaccines to aid in quick release of the cargo from endosomal compartment and protects the mRNA from acid hydrolysis and lysosomal digestion in the intracellular milieu. The vaccine was found to elicit a stable Th1-IgG2-biased antibody response.

'GBPD060' immunization did not produce any noticeable effect for local or systemic toxicity as was primarily evident by the absence of four cardinal signs of inflammation: redness (Latin rubor), heat (calor), swelling (tumor), and pain (dolor). There was no erythema or erythredema as well in any injection site. Vaccination did not significantly alter CBC and blood chemistry indicating the vaccine is safe for use in mammals.

A balanced response between Th1 and Th2 is desired to achieve safe and effective humoral immunity performance. 'GBPD060' produced a well-balanced IgG1 and IgG2 response at 7 days post-immunization, which was maintained in 14th day post-immunization sera, indicating a stable antibody response. GBPD060' also elicited a higher ratio of IgG2 to IgG1 indicating a higher capacity of the antibody pool to clear antigen from the system. The ratio of IgG2a to IgG1, and cytokine-stained CD4+ and CD8+ T cell population showed a Th1-bias response. Since mouse IgG2 is equivalent to human IgG1, 'GBPD060' will elicit effective cellular and humoral response against SARS-CoV-2 in human.

Importantly, 'GBPD060' has elicited a high level of specific antibody with a single immunization, which is

24

comparable with the level of antibody response observed after administration of a 2nd boosting dose in the mRNA vaccine developed by others. Additionally, a significant level of 51 peptide-pool-specific CD27+ memory B cells were observed 91 days following a first immunization. Therefore, GBPD060 provides suitable protection against SARS-CoV-2 in human with a single dose of injection. A single dose for effective immunization against SARS-CoV-2 is highly desirable to achieve mass global vaccination.

TABLE 1

Comparative design features of 'GBPD060'		
Parameter	GBPD060	Others
Construct	T7 promoter	T7 promoter[6]
	51 bp 5'-UTR	5'-UTR[5-6]
	Ribosome binding sequence	Not specified
	IgE signal peptide in the ORF	ORF[5-8]
	L452R, N501Y and D614G mutations	Not in consideration
	K986P and V987P mutations	K986P and V987P mutations[5-6]
LNP	Modified alpha and beta globin in 3'-UTR	3'-UTR[5-6]
	130 bp poly A tail	Poly A tail[5], [7]
	LNP composition: MC3, DSPC, Cholesterol and DMG-PEG2000 (50:10:38:5:1.5). Stabilization buffer: 1x PBS, pH 7.2	LNP composition: ionizable lipid, DSPC, Cholesterol and PEG2000-DMG[7] Stabilization buffer: HEPES buffer, pH 8.0; [7]
IgG2a/IgG1 ratio	LNP size range: 85 ± 20 nm	LNP Size: ~75 nm[8] and average size 100 nm[32]
	~1.0	~0.8, [5] 1.6[8]

The findings clearly demonstrate that 'GBPD060' is safe for in vivo administration, and elicits balanced and stable cellular and humoral response that neutralize SARS-CoV-2 spike protein-mediated infection.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 15

<210> SEQ ID NO 1

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: nucleotide sequence encoding the T7 promoter

<400> SEQUENCE: 1

taatacagact cactatagg

19

<210> SEQ ID NO 2

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: nucleotide sequence encoding the T3 promoter

<400> SEQUENCE: 2

-continued

aattaaccct cactaaagg 19

<210> SEQ ID NO 3
 <211> LENGTH: 18
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: nucleotide sequence encoding the Sp6 promoter
 <400> SEQUENCE: 3

atttaggtga cactatag 18

<210> SEQ ID NO 4
 <211> LENGTH: 17
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence
 <400> SEQUENCE: 4

gcctggctta tcgaaat 17

<210> SEQ ID NO 5
 <211> LENGTH: 51
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence
 <400> SEQUENCE: 5

gaaataagag agaaaagaag agtaagaaga aatataagag ctacgggtac c 51

<210> SEQ ID NO 6
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence
 <400> SEQUENCE: 6

Met Asp Trp Thr Trp Ile Leu Phe Leu Val Ala Ala Ala Thr Arg Val
 1 5 10 15

His Ser

<210> SEQ ID NO 7
 <211> LENGTH: 54
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence
 <400> SEQUENCE: 7

atggactgga cctggatcct cttcttggtg gcagcagcca cgcgagtcca ctcc 54

<210> SEQ ID NO 8
 <211> LENGTH: 3783
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence
 <400> SEQUENCE: 8

cagtgtgtta atcttacaac cagaactcaa ttacccctg catacactaa ttctttcaca 60

cgtggtgttt attaccctga caaagttttc agatcctcag ttttacattc aactcaggac 120

-continued

ttgttcttac	ctttcttttc	caatgttact	tggttocatg	ctatacatgt	ctctgggacc	180
aatggtacta	agaggtttga	taaccctgtc	ctaccattta	atgatggtgt	ttattttgct	240
tccactgaga	agtcatacat	aataagaggc	tggatttttg	gtactacttt	agattcgaag	300
accaggtccc	tacttattgt	taataacgct	actaatgttg	ttattaaagt	ctgtgaattt	360
caattttgta	atgatccatt	tttgggtgtt	tattaccaca	aaaacaacaa	aagttggatg	420
gaaagtgagt	tcagagttta	ttctagtgcg	aataattgca	cttttgaata	tgtctctcag	480
ccttttctta	tggaccttga	aggaaaacag	ggtaatttca	aaaatcttag	ggaatttgtg	540
tttaagaata	ttgatggtta	ttttaaaata	tattctaagc	acacgcctat	taatttagtg	600
cgtgatctcc	ctcagggttt	ttcggcttta	gaaccattgg	tagatttgcc	aatagggtatt	660
aacatcacta	ggtttcaaac	tttacttgct	ttacatagaa	gttatttgac	tcctgggtgat	720
tcttcttcag	gttggacagc	tgggtctgca	gcttattatg	tgggttatct	tcaacctagg	780
acttttctat	taaaaataaa	tgaaaatgga	accattacag	atgctgtaga	ctgtgcactt	840
gacctctctc	cagaacaacaa	gtgtacgttg	aaatccttca	ctgtagaaaa	aggaatctat	900
caaacttcta	acttttagagt	ccaaccaaca	gaatctattg	ttagatttcc	taatattaca	960
aacttggtgc	cttttggtga	agtttttaac	gccaccagat	ttgcatctgt	ttatgcttgg	1020
aacaggaaga	gaatcagcaa	ctgtgttgct	gattattctg	tcctatataa	ttccgcatca	1080
ttttccactt	ttaagtgtta	tggagtgtct	cctactaaat	taaatgatct	ctgctttact	1140
aatgtctatg	cagattcatt	tgtaattaga	ggtgatgaag	tcagacaaat	cgctccaggg	1200
caaaactggaa	agattgctga	ttataattat	aaattaccag	atgattttac	aggctgcgtt	1260
atagcttgga	attctaacaa	tcttgattct	aagggtgggtg	gtaattataa	ttacctgtat	1320
agattgttta	ggaagtctaa	tctcaaacct	tttgagagag	atatttcaac	tgaatctat	1380
caggccggta	gcacaccttg	taatgggtgt	gaaggtttta	attgttactt	tcctttacaa	1440
tcatatggtt	tccaaccac	taatgggtgt	ggttaccac	catacagagt	agtagtactt	1500
tcttttgaac	ttctacatgc	accagcaact	gtttgtggac	ctaaaaagtc	tactaatttg	1560
gttaaaaaa	aatgtgtcaa	tttcaacttc	aatggtttaa	caggcacagg	tggtcttact	1620
gagtctaaca	aaaagtttct	gcctttccaa	caatttggca	gagacattgc	tgacactact	1680
gatgtgtccc	gtgatccaca	gacacttgag	attcttgaca	ttacaccatg	ttcttttgggt	1740
ggtgtcagtg	ttataacacc	aggaacaaat	acttctaacc	aggttgctgt	tctttatcag	1800
ggtgttaact	gcacagaagt	cctgttgct	attcatgcag	atcaacttac	tcctacttgg	1860
cgtgtttatt	ctacagggtc	taatgttttt	caaacacgtg	caggctgttt	aataggggct	1920
gaacatgtca	acaactcata	tgagtgtgac	ataccatttg	gtgcagggtat	atgcgctagt	1980
tatcagactc	agactaattc	tcctcggcgg	gcacgtagtg	tagctagtca	atccatcatt	2040
gcctacacta	tgtcacttgg	tgcaaaaaat	tcagttgctt	actctaataa	ctctattgcc	2100
ataccacaaa	attttactat	tagtgttacc	acagaaattc	taccagtgtc	tatgaccaag	2160
acatcagtag	attgtacaat	gtacatttgt	ggtgattcaa	ctgaatgcag	caatcttttg	2220
ttgcaatatg	gcagtttttg	tacacaatta	aaccgtgctt	taactggaat	agctgttgaa	2280
caagacaaaa	acaccaaga	agtttttgca	caagtcaaac	aaatttacaa	aacaccacca	2340
attaagatt	ttggtggttt	taatttttca	caaattattac	cagatccatc	aaaaccaagc	2400
aagaggtcat	ttattgaaga	tctacttttc	aacaaagtga	cacttgacga	tgctggcttc	2460

-continued

atcaaaacaat atggtgattg ccttggatgat attgctgcta gagacctcat ttgtgcacaa	2520
aagtttaacg gccttactgt ttggccacct ttgctcacag atgaaatgat tgctcaatac	2580
acttctgcac tgttagcggg tacaatcact tctggttga cctttggtgc aggtgctgca	2640
ttacaaatac catttgctat gcaaatggct tataggttta atggtattgg agttacacag	2700
aatgttctct atgagaacca aaaattgatt gccaaccaat ttaatagtgc tattggcaaa	2760
attcaagact cactttcttc cacagcaagt gcacttgaa aacttcaaga tgtggtcaac	2820
caaatgcac aagctttaaa cacgcttgtt aaacaactta gctccaattt tgggtcaatt	2880
tcaagtgttt taaatgatat cctttcacgt ctgaccctc ctgaggctga agtgcaaat	2940
gatagggtta tcacaggcag acttcaaagt ttgcagacat atgtgactca acaattaatt	3000
agagctgcag aaatcagagc ttctgctaatt ctgctgcta ctaaatgtc agagtgtgta	3060
cttgacaat caaaaagagt tgatttttgt ggaaagggct atcatcttat gtccttcct	3120
cagtcagcac ctcatggtgt agtctcttg catgtgactt atgtccctgc acaagaaaag	3180
aaactcaca ctgctcctgc catttgctcat gatggaaaag cacactttcc tcgtgaaggt	3240
gtctttgttt caaatggcac aactggttt gtaacacaaa ggaattttta tgaaccacaa	3300
atcattacta cagacaacac atttgtgtct ggtaactgtg atgttgtaat aggaattgtc	3360
aacaacacag tttatgatcc ttgcaacct gaattagact cattcaagga ggagttagat	3420
aaatatttta agaatcatc atcaccagat gttgatttag gtgacatctc tggcattaat	3480
gcttcagttg taaacattca aaaagaaatt gaccgctca atgaggttgc caagaattta	3540
aatgaatctc tcatgatct ccaagaactt ggaaagtatg agcagtatat aaaatggcca	3600
tggtagattt ggctaggttt tatagctggc ttgattgcca tagtaatggt gacaattatg	3660
ctttgctgta tgaccagttg ctgtagttgt ctcaagggct gttgttcttg tggatcctgc	3720
tgcaaatgtg atgaagacga ctctgagcca gtgctcaaag gagtcaaatt acattacaca	3780
taa	3783

<210> SEQ ID NO 9

<211> LENGTH: 3783

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 9

cagtgctgta acctgaccac cagaacacag ctgcctccag cctacaccaa cagcttcacc	60
agaggcgtgt actaccccg caaggtgttc agatccagcg tgctgcactc taccaggac	120
ctgttctctc cttttctcag caacgtgacc tggttccacg ccatccacgt gtcggaacc	180
aatggcacca agagattcga caacccctg ctgcccttca acgacggggt gtactttgcc	240
agcaccgaga agtccaacat catcagaggc tggatcttcg gcaccacact ggacagcaag	300
acccagagcc tgctgatcgt gaacaacgcc accaacgtgg tcatcaaagt gtgcgagttc	360
cagttctgca acgacccctt cctgggcgtc tactaccaca agaacaacaa gagctggatg	420
gaaagcgagt tccgggtgta cagcagcgcc aacaactgca ccttogagta cgtgtcccag	480
cctttcctga tggacctgga aggcaagcag ggcaacttca agaacctgcg cgagtctctg	540
ttcaagaaca tcagcgcta cttcaagatc tacagcaagc acaccctat caacctctg	600
cgggatctgc ctacgggctt ctctgctctg gaaccctgg tggatctgcc catcggcac	660
aacatcaccg ggtttcagac actgctggcc ctgcacagaa gctacctgac acctggcgat	720

-continued

agcagctctg gatggacagc tggagccgct gcctactatg tgggatacct gcagcctcgg	780
accttctctg tgaagtacaa cgagaacggc accatcaccg acgccgtgga ttgtgctctg	840
gacctctctg gcgagacaaa gtgcaccctg aagtccttca ccgtggaaaa gggcatctac	900
cagaccagca acttccgggt gcagcccacc gaatccatcg tgcgggtccc caatatacc	960
aatctgtgcc ccttcggcga ggtgttcaat gccaccagat tcgcctctgt gtaacgcctgg	1020
aaccggaagc ggatcagcaa ttgcgtggcc gactactccg tgctgtacaa ctccgccagc	1080
ttcagcacct tcaagtgeta cggcgtgtcc cctaccaagc tgaacgacct gtgcttcaca	1140
aacgtgtacg ccgacagctt cgtgatccgg ggagatgaag tgcggcagat tgcccctgga	1200
cagacaggca agatcgccga ctacaactac aagctgcccg acgacttcac cggctgtgtg	1260
attgcctgga acagcaacaa cctggactcc aaagtccggc gcaactacaa ttacctgtac	1320
cggctgttcc ggaagtccaa tctgaagccc ttcgagcggg acatctccac cgagatctat	1380
caggccggca gccccccttg taacggcgtg gaaggcttca actgctactt cccactgcag	1440
tcctacggct tccagccaac aaacggcgtg ggctaccagc cttacagagt ggtggtgctg	1500
agcttcgagc tgctgcatgc tcctgccaca gtgtgcggcc ctaagaaaag caccaatctc	1560
gtgaagaaca aatcgtgtaa cttcaacttc aacggcctga ccggcacccg cgtgctgaca	1620
gagagcaaca agaagtctct gccattccag cagttcggcc gggatatcgc cgataccaca	1680
gatgccgtca gagatcccca gacactggaa atcctggaca tcacccttg cagcttcggc	1740
ggagtgtctg tgatcacccc tggcaccaac accagcaatc aggtggcagt gctgtaccag	1800
gatgtgaact gtacagaggt gccagtggcc attcacgcgc atcagctgac ccctacttgg	1860
cgggtgtact ccacaggcag caatgtgttt cagaccagag ccggtgtgtc gatcggagcc	1920
gagcacgtga acaatagcta cgagtgcgac atccccatcg gcgctggcat ctgcgcctct	1980
taccagacac agacaaacag cccagacgg gccagatctg tggccagcca gagcatcatt	2040
gcctacacaa tgtctctggg agccgagaa agcgtggcct actccaacaa ctctatcgct	2100
atccccacca acttcacat cagcgtgacc acagagatcc tgctgtgtc catgaccaag	2160
accagcgtgg actgcacat gtacatctgc ggcgattcca ccgagtgtc caacctgctg	2220
ctgcagtacg gcagcttctg caccagctg aatagagccc tgacagggat ccgcgtggaa	2280
caggacaaga acacccaaga ggtgttcgcc caagtgaagc agatctacaa gacccctcct	2340
atcaaggact tcggcggctt caatttcagc cagattctgc ccgatactag caagcccagc	2400
aagcggagct tcategagga cctgtgttc aacaaagtga cactggccga ccgcggcttc	2460
atcaagcagt atggcgattg tctgggcgac attgcccca gggatctgat ttgcgccag	2520
aagttaacg gactgacagt gctgcctcct ctgctgaccg atgagatgat cggccagtac	2580
acatctgccc tgctggccgg cacaatcaca agcggctgga catttgagc tggcgtgcc	2640
ctgcagatcc cctttgtat gcagatggcc taccggttca acggcatcgg agtgaccag	2700
aatgtgctgt acgagaacca gaagctgac gccaaacagt tcaacagcgc catcggcaag	2760
atccaggaca gcctgagcag cacagcaagc gccctgggaa agctgcagga cgtggtcaac	2820
cagaatgccc aggcactgaa caccctgtc aagcagctgt ctageaactt cggagccatc	2880
agctctgtgc tgaacgatat cctgagcaga ctggaccctc ctgaggccga ggtgcagatc	2940
gacagactga tcacaggcag actgcagagc ctccagacat acgtgaccca gcagctgac	3000
agagccggc agattagagc ctctgccaat ctggccgcca ccaagatgtc tgagtgtgtg	3060

-continued

ctggggccaga gcaagagagt ggacttttgc ggcaagggct accacctgat gagcttcct	3120
cagctctgcac cacacggcgt ggtgtttctg cacgtgacct acgtgccgc tcaagagaag	3180
aatttcacca cgcctccagc catctgccac gacggcaaag cccactttcc tagagaaggc	3240
gtgttcgtgt ccaacggcac ccattgggtc gtgacacagc ggaacttcta cgagcccag	3300
atcatcacca ccgacaacac ctctgtgtct ggcaactgcg acgtcgtgat cggcattgtg	3360
aacaataccg tgtacgaccc tctgcagccc gagctggaca gcttcaaaga ggaactggac	3420
aagtacttta agaaccacac aagccccgac gtggacctgg gcgatatcag cggaatcaat	3480
gccagcgtcg tgaacatcca gaaagagatc gaccggctga acgaggtggc caagaatctg	3540
aacgagagcc tgatcgacct gcaagaactg gggaagtacg agcagtacat caagtggcct	3600
tggtacatct ggctgggctt tatcgccgga ctgattgcca tcgtgatggc cacaatcatg	3660
ctgtgttgca tgaccagctg ctgtagctgc ctgaagggct gttgtagctg tggctcctgc	3720
tgcaagttcg acgaggacga ttctgagccc gtgctgaaag gcgtgaagct gcactacacc	3780
tga	3783

<210> SEQ ID NO 10

<211> LENGTH: 3783

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 10

cagtgcgtga acctgaccac cagaacacag ctgcctccag cctacaccaa cagcttcacc	60
agaggcgtgt actaccccgca caagggtgtc agatccagcg tgctgcactc taccaggac	120
ctgttctctg cttttcttcag caacgtgacc tggttccacg ccattccacgt gtccggcacc	180
aatggcacca agagattcga caaccctgtg ctgcccttca acgacggggg gtactttgcc	240
agcaccgaga agtccaacat catcagaggc tggatcttcg gcaccacact ggacagcaag	300
accagagacc tgctgatcgt gaacaacgcc accaacgtgg tcatcaaagt gtgcgagttc	360
cagttctgca acgacccctt cctgggcgtc tactaccaca agaacaacaa gagctggatg	420
gaaagcgagt tccgggtgta cagcagcgcc aacaactgca ccttcagta cgtgtcccag	480
cctttctctg tggacctgga aggcaagcag ggcaacttca agaacctgcg cgagtctctg	540
ttcaagaaca tcgacggcta cttcaagatc tacagcaagc acaccctat caacctcgtg	600
cgggatctgc ctcagggtt cctgtctctg gaaccctctg tggatctgcc catcggcac	660
aacatcaccc ggtttcagac actgctggcc ctgcacagaa gctacctgac acctggcgat	720
agcagctctg gatggacagc tggagccgct gcctactatg tgggatacct gcagcctcgg	780
accttctctg tgaagtacaa cgagaacggc accatcaccc acgccgtgga ttgtgctctg	840
gatcctctga gcgagacaaa gtgcacctg aagtcttca ccgtggaaaa ggcatctac	900
cagaccagca acttcgggt gcagcccacc gaatccatcg tgcgggtccc caatatacc	960
aatctgtgcc ccttcggcga ggtgttcaat gccaccagat tcgcctctgt gtacgcctgg	1020
aaccggaagc ggatcagcaa ttgcgtggcc gactactccg tgctgtacaa ctccgccagc	1080
ttcagcacct tcaagtgcta cggcgtgtcc cctaccaagc tgaacgacct gtgcttcaca	1140
aacgtgtacg ccgacagctt cgtgatccg ggagatgaag tgcggcagat tgcctcggga	1200
cagacaggca agatcgccga ctacaactac aagctgcccg acgacttcac cggctgtgtg	1260
attgcctgga acagcaacaa cctggactcc aaagtcggcg gcaactacaa ttacctgtac	1320

-continued

cggtgtttcc ggaagtccaa tctgaagccc ttcgagcggg acatctccac cgagatctat	1380
caggccggca gcaccccttg taacggcgtg gaaggettca actgtactt cccactgcag	1440
tcctacggct tcacagcaac aaacggcgtg ggctaccagc cttacagagt ggtggtgctg	1500
agcttcgagc tgctgcatgc tectgccaca gtgtgcggcc ctaagaaaag caccaatctc	1560
gtgaagaaca aatgcgtgaa cttcaacttc aacggcctga ccggcaccgg cgtgctgaca	1620
gagagcaaca agaagttcct gccattccag cagttcggcc gggatatcgc cgataccaca	1680
gatgccgtca gagatcccca gacactggaa atcctggaca tcaccccttg cagcttcggc	1740
ggagtgtctg tgatcacccc tggcaccaac accagcaatc aggtggcagt gctgtaccag	1800
ggcgtgaact gtacagaggt gccagtggcc attcacgccg atcagctgac ccctacttgg	1860
cgggtgtact ccacaggcag caatgtgttt cagaccagag ccggctgtct gatcggagcc	1920
gagcacgtga acaatagcta cgagtgcgac atccccatcg gcgctggcat ctgcgcctct	1980
taccagacac agacaaaacag cccagacgg gccagatctg tggccagcca gagcatcatt	2040
gcctacacaa tgtctctggg agccgagaac agcgtggcct actccaacaa ctctatcgct	2100
atccccacca acttcacat cagcgtgacc acagagatcc tgctgtgtc catgaccaag	2160
accagcgtgg actgcacat gtacatctgc ggcgattcca ccgagtgtc caacctgctg	2220
ctgcagtacg gcagctttctg caccagctg aatagagccc tgacagggat cgcgctggaa	2280
caggacaaga acacccaaga ggtgttcgcc caagtgaagc agatctacaa gacccctcct	2340
atcaaggact tcggcggctt caatttcagc cagattctgc ccgatcctag caagcccagc	2400
aagcggagct tcctcgagga cctgctgttc aacaaagtga cactggccga cgcggcttc	2460
atcaagcagt atggcgattg tctggcgac attgccgcca gggatctgat ttgcgccag	2520
aagtttaacg gactgacagt gctgctcct ctgctgaccg atgagatgat cccccagta	2580
acatctgccc tgctggccgg cacaatcaca agcggctgga catttgagc tggcgtgcc	2640
ctgcagatcc cctttgctat gcagatggcc taccggttca acggcatcgg agtgacccag	2700
aatgtgctgt acgagaacca gaagctgac gccaacagct tcaacagcgc catcggcaag	2760
atccaggaca gcctgagcag cacagcaagc gccctgggaa agctgcagga cgtggtaac	2820
cagaatgccc aggcactgaa caccctggtc aagcagctgt ctagcaactt cggagccatc	2880
agctctgtgc tgaacgatat cctgagcaga ctggaccctc ctgagggcca ggtgcagatc	2940
gacagactga tcacaggcag actgcagagc ctccagacat acgtgaccca gcagctgac	3000
agagccgccg agattagagc ctctgccaat ctggccgcca ccaagatgtc tgagtgtgtg	3060
ctgggccaga gcaagagagt ggacttttgc ggcaagggt accacctgat gagcttcct	3120
cagtctgcac cacacggcgt ggtgtttctg cacgtgacct acgtgcccgc tcaagagaag	3180
aatttcacca ccgctccagc catctgccac gacggcaaaag cccactttcc tagagaaggc	3240
gtgttcgtgt ccaacggcac ccattggttc gtgacacagc ggaacttcta cgagccccag	3300
atcatcacca ccgacaacac cttcgtgtct ggcaactgcg acgtcgtgat cggcattgtg	3360
aacaataccg tgtaacgccc tctgcagccc gagctggaca gcttcaaaga ggaactggac	3420
aagtacttta agaaccacac aagccccgac gtggacctgg gcgatatcag cggaatcaat	3480
gccagcgtcg tgaacatcca gaaagagatc gaccggctga acgaggtggc caagaatctg	3540
aacgagagcc tgatcgacct gcaagaactg ggaagtacg agcagtacat caagtggcct	3600
tggtacatct ggctgggctt tatcgccgga ctgattgcca tcgtgatggt cacaatcatg	3660

-continued

ctgtgttgca tgaccagctg ctgtagctgc ctgaagggt gttgtagctg tggtccctgc	3720
tgcaagttcg acgaggacga ttctgagccc gtgctgaaag gcgtgaagct gcactacacc	3780
tga	3783

<210> SEQ ID NO 11
 <211> LENGTH: 3783
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 11

cagtgcgtga acctgaccac cagaacacag ctgcctccag cctacaccaa cagcttcacc	60
agaggcgtgt actaccccgga caaggtgttc agatccagcg tgctgcactc taccaggac	120
ctgttctctgc ctttcttcag caacgtgacc tggttccacg ccatccacgt gtcggcacc	180
aatggcacca agagattcga caaccccggtg ctgcccttca acgacgggggt gtactttgcc	240
agcaccgaga agtccaacat catcagaggc tggatcttcg gcaccacact ggacagcaag	300
acccagagcc tgctgatcgt gaacaacgcc accaacgtgg tcatcaaagt gtgcgagttc	360
cagttctgca acgaccctt cctgggcgtc tactaccaca agaacaacaa gagctggatg	420
gaaagcgagt tccgggtgta cagcagcgcc aacaactgca ccttcgagta cgtgtcccag	480
ccttctctga tggacctgga aggcaagcag ggcaacttca agaacctgcg cgagtctgtg	540
ttcaagaaca tcgacggcta cttcaagatc tacagcaagc acaccttat caacctcgtg	600
cgggatctgc ctacgggctt ctctgctctg gaacctctgg tggatctgcc catcggcac	660
aacatcaccg gggttcagac actgctggcc ctgcacagaa gctacctgac acctggcgat	720
agcagctctg gatggacagc tggagccgct gcctactatg tgggatacct gcagcctcgg	780
accttctctg tgaagtacaa cgagaacggc accatcaccg acgcccgtgga ttgtgctctg	840
gatcctctga gcgagacaaa gtgcacccctg aagtccttca ccgtggaaaa gggcatctac	900
cagaccagca acttcgggtg gcagcccacc gaatccatcg tgcgggtccc caatatcacc	960
aatctgtgcc ccttcggcga ggtgttcaat gccaccagat tcgcctctgt gtacgcctgg	1020
aaccggaagc ggatcagcaa ttgcgtggcc gactactccg tgctgtacaa ctccgccagc	1080
ttcagcacct tcaagtgcta cggcgtgtcc cctaccaagc tgaacgacct gtgcttcaca	1140
aacgtgtacg ccgacagctt cgtgatccgg ggagatgaag tgcggcagat tgcccctgga	1200
cagacaggca agatcgccga ctacaactac aagctgcccg acgacttcac cggctgtgtg	1260
attgcctgga acagcaacaa cctggactcc aaagtccggc gcaactacaa ttacagatac	1320
cggctgttcc ggaagtccaa tctgaagccc ttcgagcggg acatctccac cgagatctat	1380
caggccggca gcaacccttg taacggcgtg gaaggcttca actgctactt cccactgcag	1440
tcctacggct tccagccaac atacggcgtg ggctaccagc cttacagagt ggtggtgctg	1500
agcttcgagc tgctgcatgc tcttgccaca gtgtgcggcc ctaagaaaag caccaatctc	1560
gtgaagaaca aatgcgtgaa cttcaacttc aacggcctga ccggcaccgg cgtgctgaca	1620
gagagcaaca agaagttcct gccattccag cagttcggcc gggatatcgc cgataccaca	1680
gatgccgtca gagatcccca gacactggaa atcctggaca tcacccttg cagcttcggc	1740
ggagtgtctg tgatcacccc tggcaccaac accagcaatc aggtggcagt gctgtaccag	1800
ggcgtgaact gtacagaggt gccagtggcc attcacgccg atcagctgac ccctacttgg	1860
cgggtgtact ccacaggcag caatgtgttt cagaccagag ccggtgtctc gatcggagcc	1920

-continued

```

gagcacgtga acaatagcta cgagtgcgac atccccatcg gcgtggcat ctgcgcctct 1980
taccagacac agacaaacag ccccgacgg gccagatctg tggccagcca gagcatcatt 2040
gcctacacaa tgtctctggg agccgagAAC agcgtggcct actccaacaa ctctatcgct 2100
atccccacca acttcaccat cagcgtgacc acagagatcc tgectgtgtc catgaccaag 2160
accagcgtgg actgcaccat gtacatctgc ggcgattcca ccgagtgtc caacctgtg 2220
ctgcagtacg gcagcttctg cccccagctg aatagagccc tgacagggat cgccgtggaa 2280
caggacaaga acaccaaga ggtgttcgcc caagtgaagc agatctacaa gaccctcct 2340
atcaaggact tcggcggctt caatttcagc cagattctgc ccgatcctag caagcccagc 2400
aagcggagct tcctcgagga cctgtgttc aacaaagtga cactggccga cgccggcttc 2460
atcaagcagt atggcgattg tctggcgac attgcgcga gggatctgat ttgcgccag 2520
aagttaacg gactgacagt gctgcctcct ctgtgaccg atgagatgat cgcccagtac 2580
acatctgccc tgctggccgg cacaatcaca agcggctgga catttgagc tggcgtgccc 2640
ctgcagatcc ccttctgtat gcagatggcc taccgggttc acggcatcgg agtgaccag 2700
aatgtgctgt acgagaacca gaagctgatc gccaacagc tcaacagcgc catcgcaag 2760
atccaggaca gcctgagcag cacagcaagc gccctgggaa agctgcagga cgtggccaac 2820
cagaatgccc aggcactgaa caccctgttc aagcagctgt ctagcaactt cggagccatc 2880
agctctgtgc tgaacgatat cctgagcaga ctggaccctc ctgaggccga ggtgcagatc 2940
gacagactga tcacaggcag actgcagagc ctccagacat acgtgaccca gcagctgatc 3000
agagccgccc agattagagc ctctgccaat ctggccgcca ccaagatgtc tgagtgtgtg 3060
ctgggccaga gcaagagagt ggacttttgc ggcaagggt accacctgat gagcttcct 3120
cagctctgac cacacggcgt ggtgtttctg cacgtgacct acgtgcccgc tcaagagaag 3180
aatctcaca ccgtccagc catctgccac gacggcaag cccactttcc tagagaaggc 3240
gtgttcgtgt ccaacggcac ccattggttc gtgacacagc ggaacttcta cgagcccag 3300
atcatcaca ccgacaacac ctctgtgtct ggcaactcgc acgtcgtgat cggcattgtg 3360
aacaataccg tgtaacgccc tctgcagccc gagctggaca gcttcaaga ggaactggac 3420
aagtacttta agaaccacac aagccccgac gtggacctgg gcgatatcag cggaatcaat 3480
gccagcgtcg tgaacatcca gaaagagatc gaccggctga acgaggtggc caagaatctg 3540
aacgagagcc tgatcgacct gcaagaactg ggggaagtac agcagtacat caagtggcct 3600
tggtacatct ggctgggctt tatcgccgga ctgattgcca tcgtgatggt cacaatcatg 3660
ctgtgttgca tgaccagctg ctgtagctgc ctgaagggtc gttgtagctg tggctcctgc 3720
tgcaagttcg acgaggacga ttctgagccc gtgctgaaag gcgtgaagct gcactacacc 3780
tga 3783

```

<210> SEQ ID NO 12

<211> LENGTH: 1259

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 12

```

Gln Cys Val Asn Leu Thr Thr Arg Thr Gln Leu Pro Pro Ala Tyr Thr
1           5           10           15

```

```

Asn Ser Phe Thr Arg Gly Val Tyr Tyr Pro Asp Lys Val Phe Arg Ser

```

-continued

20					25					30					
Ser	Val	Leu	His	Ser	Thr	Gln	Asp	Leu	Phe	Leu	Pro	Phe	Phe	Ser	Asn
		35					40					45			
Val	Thr	Trp	Phe	His	Ala	Ile	His	Val	Ser	Gly	Thr	Asn	Gly	Thr	Lys
		50				55					60				
Arg	Phe	Asp	Asn	Pro	Val	Leu	Pro	Phe	Asn	Asp	Gly	Val	Tyr	Phe	Ala
65					70					75					80
Ser	Thr	Glu	Lys	Ser	Asn	Ile	Ile	Arg	Gly	Trp	Ile	Phe	Gly	Thr	Thr
				85					90					95	
Leu	Asp	Ser	Lys	Thr	Gln	Ser	Leu	Leu	Ile	Val	Asn	Asn	Ala	Thr	Asn
			100					105						110	
Val	Val	Ile	Lys	Val	Cys	Glu	Phe	Gln	Phe	Cys	Asn	Asp	Pro	Phe	Leu
		115						120				125			
Gly	Val	Tyr	Tyr	His	Lys	Asn	Asn	Lys	Ser	Trp	Met	Glu	Ser	Glu	Phe
	130					135					140				
Arg	Val	Tyr	Ser	Ser	Ala	Asn	Asn	Cys	Thr	Phe	Glu	Tyr	Val	Ser	Gln
145					150					155					160
Pro	Phe	Leu	Met	Asp	Leu	Glu	Gly	Lys	Gln	Gly	Asn	Phe	Lys	Asn	Leu
				165					170					175	
Arg	Glu	Phe	Val	Phe	Lys	Asn	Ile	Asp	Gly	Tyr	Phe	Lys	Ile	Tyr	Ser
			180						185					190	
Lys	His	Thr	Pro	Ile	Asn	Leu	Val	Arg	Asp	Leu	Pro	Gln	Gly	Phe	Ser
		195					200					205			
Ala	Leu	Glu	Pro	Leu	Val	Asp	Leu	Pro	Ile	Gly	Ile	Asn	Ile	Thr	Arg
	210					215					220				
Phe	Gln	Thr	Leu	Leu	Ala	Leu	His	Arg	Ser	Tyr	Leu	Thr	Pro	Gly	Asp
225					230					235					240
Ser	Ser	Ser	Gly	Trp	Thr	Ala	Gly	Ala	Ala	Ala	Tyr	Tyr	Val	Gly	Tyr
				245					250					255	
Leu	Gln	Pro	Arg	Thr	Phe	Leu	Leu	Lys	Tyr	Asn	Glu	Asn	Gly	Thr	Ile
			260					265						270	
Thr	Asp	Ala	Val	Asp	Cys	Ala	Leu	Asp	Pro	Leu	Ser	Glu	Thr	Lys	Cys
		275					280						285		
Thr	Leu	Lys	Ser	Phe	Thr	Val	Glu	Lys	Gly	Ile	Tyr	Gln	Thr	Ser	Asn
	290					295					300				
Phe	Arg	Val	Gln	Pro	Thr	Glu	Ser	Ile	Val	Arg	Phe	Pro	Asn	Ile	Thr
305					310					315					320
Asn	Leu	Cys	Pro	Phe	Gly	Glu	Val	Phe	Asn	Ala	Thr	Arg	Phe	Ala	Ser
				325					330					335	
Val	Tyr	Ala	Trp	Asn	Arg	Lys	Arg	Ile	Ser	Asn	Cys	Val	Ala	Asp	Tyr
			340					345					350		
Ser	Val	Leu	Tyr	Asn	Ser	Ala	Ser	Phe	Ser	Thr	Phe	Lys	Cys	Tyr	Gly
	355						360					365			
Val	Ser	Pro	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Thr	Asn	Val	Tyr	Ala
	370					375					380				
Asp	Ser	Phe	Val	Ile	Arg	Gly	Asp	Glu	Val	Arg	Gln	Ile	Ala	Pro	Gly
385					390					395					400
Gln	Thr	Gly	Lys	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro	Asp	Asp	Phe
			405						410					415	
Thr	Gly	Cys	Val	Ile	Ala	Trp	Asn	Ser	Asn	Asn	Leu	Asp	Ser	Lys	Val
			420					425					430		
Gly	Gly	Asn	Tyr	Asn	Tyr	Leu	Tyr	Arg	Leu	Phe	Arg	Lys	Ser	Asn	Leu
	435						440					445			

-continued

Lys Pro Phe Glu Arg Asp Ile Ser Thr Glu Ile Tyr Gln Ala Gly Ser
 450 455 460
 Thr Pro Cys Asn Gly Val Glu Gly Phe Asn Cys Tyr Phe Pro Leu Gln
 465 470 475 480
 Ser Tyr Gly Phe Gln Pro Thr Asn Gly Val Gly Tyr Gln Pro Tyr Arg
 485 490 495
 Val Val Val Leu Ser Phe Glu Leu Leu His Ala Pro Ala Thr Val Cys
 500 505 510
 Gly Pro Lys Lys Ser Thr Asn Leu Val Lys Asn Lys Cys Val Asn Phe
 515 520 525
 Asn Phe Asn Gly Leu Thr Gly Thr Gly Val Leu Thr Glu Ser Asn Lys
 530 535 540
 Lys Phe Leu Pro Phe Gln Gln Phe Gly Arg Asp Ile Ala Asp Thr Thr
 545 550 555 560
 Asp Ala Val Arg Asp Pro Gln Thr Leu Glu Ile Leu Asp Ile Thr Pro
 565 570 575
 Cys Ser Phe Gly Gly Val Ser Val Ile Thr Pro Gly Thr Asn Thr Ser
 580 585 590
 Asn Gln Val Ala Val Leu Tyr Gln Gly Val Asn Cys Thr Glu Val Pro
 595 600 605
 Val Ala Ile His Ala Asp Gln Leu Thr Pro Thr Trp Arg Val Tyr Ser
 610 615 620
 Thr Gly Ser Asn Val Phe Gln Thr Arg Ala Gly Cys Leu Ile Gly Ala
 625 630 635 640
 Glu His Val Asn Asn Ser Tyr Glu Cys Asp Ile Pro Ile Gly Ala Gly
 645 650 655
 Ile Cys Ala Ser Tyr Gln Thr Gln Thr Asn Ser Pro Arg Arg Ala Arg
 660 665 670
 Ser Val Ala Ser Gln Ser Ile Ile Ala Tyr Thr Met Ser Leu Gly Ala
 675 680 685
 Glu Asn Ser Val Ala Tyr Ser Asn Asn Ser Ile Ala Ile Pro Thr Asn
 690 695 700
 Phe Thr Ile Ser Val Thr Thr Glu Ile Leu Pro Val Ser Met Thr Lys
 705 710 715 720
 Thr Ser Val Asp Cys Thr Met Tyr Ile Cys Gly Asp Ser Thr Glu Cys
 725 730 735
 Ser Asn Leu Leu Leu Gln Tyr Gly Ser Phe Cys Thr Gln Leu Asn Arg
 740 745 750
 Ala Leu Thr Gly Ile Ala Val Glu Gln Asp Lys Asn Thr Gln Glu Val
 755 760 765
 Phe Ala Gln Val Lys Gln Ile Tyr Lys Thr Pro Pro Ile Lys Asp Phe
 770 775 780
 Gly Gly Phe Asn Phe Ser Gln Ile Leu Pro Asp Pro Ser Lys Pro Ser
 785 790 795 800
 Lys Arg Ser Phe Ile Glu Asp Leu Leu Phe Asn Lys Val Thr Leu Ala
 805 810 815
 Asp Ala Gly Phe Ile Lys Gln Tyr Gly Asp Cys Leu Gly Asp Ile Ala
 820 825 830
 Ala Arg Asp Leu Ile Cys Ala Gln Lys Phe Asn Gly Leu Thr Val Leu
 835 840 845
 Pro Pro Leu Leu Thr Asp Glu Met Ile Ala Gln Tyr Thr Ser Ala Leu
 850 855 860

-continued

Leu	Ala	Gly	Thr	Ile	Thr	Ser	Gly	Trp	Thr	Phe	Gly	Ala	Gly	Ala	Ala	865	870	875	880
Leu	Gln	Ile	Pro	Phe	Ala	Met	Gln	Met	Ala	Tyr	Arg	Phe	Asn	Gly	Ile	885	890	895	
Gly	Val	Thr	Gln	Asn	Val	Leu	Tyr	Glu	Asn	Gln	Lys	Leu	Ile	Ala	Asn	900	905	910	
Gln	Phe	Asn	Ser	Ala	Ile	Gly	Lys	Ile	Gln	Asp	Ser	Leu	Ser	Ser	Thr	915	920	925	
Ala	Ser	Ala	Leu	Gly	Lys	Leu	Gln	Asp	Val	Val	Asn	Gln	Asn	Ala	Gln	930	935	940	
Ala	Leu	Asn	Thr	Leu	Val	Lys	Gln	Leu	Ser	Ser	Asn	Phe	Gly	Ala	Ile	945	950	955	960
Ser	Ser	Val	Leu	Asn	Asp	Ile	Leu	Ser	Arg	Leu	Asp	Pro	Pro	Glu	Ala	965	970	975	
Glu	Val	Gln	Ile	Asp	Arg	Leu	Ile	Thr	Gly	Arg	Leu	Gln	Ser	Leu	Gln	980	985	990	
Thr	Tyr	Val	Thr	Gln	Gln	Leu	Ile	Arg	Ala	Ala	Glu	Ile	Arg	Ala	Ser	995	1000	1005	
Ala	Asn	Leu	Ala	Ala	Thr	Lys	Met	Ser	Glu	Cys	Val	Leu	Gly	Gln		1010	1015	1020	
Ser	Lys	Arg	Val	Asp	Phe	Cys	Gly	Lys	Gly	Tyr	His	Leu	Met	Ser		1025	1030	1035	
Phe	Pro	Gln	Ser	Ala	Pro	His	Gly	Val	Val	Phe	Leu	His	Val	Thr		1040	1045	1050	
Tyr	Val	Pro	Ala	Gln	Glu	Lys	Asn	Phe	Thr	Thr	Ala	Pro	Ala	Ile		1055	1060	1065	
Cys	His	Asp	Gly	Lys	Ala	His	Phe	Pro	Arg	Glu	Gly	Val	Phe	Val		1070	1075	1080	
Ser	Asn	Gly	Thr	His	Trp	Phe	Val	Thr	Gln	Arg	Asn	Phe	Tyr	Glu		1085	1090	1095	
Pro	Gln	Ile	Ile	Thr	Thr	Asp	Asn	Thr	Phe	Val	Ser	Gly	Asn	Cys		1100	1105	1110	
Asp	Val	Val	Ile	Gly	Ile	Val	Asn	Asn	Thr	Val	Tyr	Asp	Pro	Leu		1115	1120	1125	
Gln	Pro	Glu	Leu	Asp	Ser	Phe	Lys	Glu	Glu	Leu	Asp	Lys	Tyr	Phe		1130	1135	1140	
Lys	Asn	His	Thr	Ser	Pro	Asp	Val	Asp	Leu	Gly	Asp	Ile	Ser	Gly		1145	1150	1155	
Ile	Asn	Ala	Ser	Val	Val	Asn	Ile	Gln	Lys	Glu	Ile	Asp	Arg	Leu		1160	1165	1170	
Asn	Glu	Val	Ala	Lys	Asn	Leu	Asn	Glu	Ser	Leu	Ile	Asp	Leu	Gln		1175	1180	1185	
Glu	Leu	Gly	Lys	Tyr	Glu	Gln	Tyr	Ile	Lys	Trp	Pro	Trp	Tyr	Ile		1190	1195	1200	
Trp	Leu	Gly	Phe	Ile	Ala	Gly	Leu	Ile	Ala	Ile	Val	Met	Val	Thr		1205	1210	1215	
Ile	Met	Leu	Cys	Cys	Met	Thr	Ser	Cys	Cys	Ser	Cys	Leu	Lys	Gly		1220	1225	1230	
Cys	Cys	Ser	Cys	Gly	Ser	Cys	Cys	Lys	Phe	Asp	Glu	Asp	Asp	Ser		1235	1240	1245	
Glu	Pro	Leu	Lys	Gly	Val	Lys	Leu	His	Tyr	Thr						1250	1255		

-continued

```

<210> SEQ ID NO 13
<211> LENGTH: 1260
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 13

Gln Cys Val Asn Leu Thr Thr Arg Thr Gln Leu Pro Pro Ala Tyr Thr
1          5          10          15

Asn Ser Phe Thr Arg Gly Val Tyr Tyr Pro Asp Lys Val Phe Arg Ser
20        25        30

Ser Val Leu His Ser Thr Gln Asp Leu Phe Leu Pro Phe Phe Ser Asn
35        40        45

Val Thr Trp Phe His Ala Ile His Val Ser Gly Thr Asn Gly Thr Lys
50        55        60

Arg Phe Asp Asn Pro Val Leu Pro Phe Asn Asp Gly Val Tyr Phe Ala
65        70        75        80

Ser Thr Glu Lys Ser Asn Ile Ile Arg Gly Trp Ile Phe Gly Thr Thr
85        90        95

Leu Asp Ser Lys Thr Gln Ser Leu Leu Ile Val Asn Asn Ala Thr Asn
100       105       110

Val Val Ile Lys Val Cys Glu Phe Gln Phe Cys Asn Asp Pro Phe Leu
115       120       125

Gly Val Tyr Tyr His Lys Asn Asn Lys Ser Trp Met Glu Ser Glu Phe
130       135       140

Arg Val Tyr Ser Ser Ala Asn Asn Cys Thr Phe Glu Tyr Val Ser Gln
145       150       155       160

Pro Phe Leu Met Asp Leu Glu Gly Lys Gln Gly Asn Phe Lys Asn Leu
165       170       175

Arg Glu Phe Val Phe Lys Asn Ile Asp Gly Tyr Phe Lys Ile Tyr Ser
180       185       190

Lys His Thr Pro Ile Asn Leu Val Arg Asp Leu Pro Gln Gly Phe Ser
195       200       205

Ala Leu Glu Pro Leu Val Asp Leu Pro Ile Gly Ile Asn Ile Thr Arg
210       215       220

Phe Gln Thr Leu Leu Ala Leu His Arg Ser Tyr Leu Thr Pro Gly Asp
225       230       235       240

Ser Ser Ser Gly Trp Thr Ala Gly Ala Ala Ala Tyr Tyr Val Gly Tyr
245       250       255

Leu Gln Pro Arg Thr Phe Leu Leu Lys Tyr Asn Glu Asn Gly Thr Ile
260       265       270

Thr Asp Ala Val Asp Cys Ala Leu Asp Pro Leu Ser Glu Thr Lys Cys
275       280       285

Thr Leu Lys Ser Phe Thr Val Glu Lys Gly Ile Tyr Gln Thr Ser Asn
290       295       300

Phe Arg Val Gln Pro Thr Glu Ser Ile Val Arg Phe Pro Asn Ile Thr
305       310       315       320

Asn Leu Cys Pro Phe Gly Glu Val Phe Asn Ala Thr Arg Phe Ala Ser
325       330       335

Val Tyr Ala Trp Asn Arg Lys Arg Ile Ser Asn Cys Val Ala Asp Tyr
340       345       350

Ser Val Leu Tyr Asn Ser Ala Ser Phe Ser Thr Phe Lys Cys Tyr Gly
355       360       365

Val Ser Pro Thr Lys Leu Asn Asp Leu Cys Phe Thr Asn Val Tyr Ala

```

-continued

370	375	380
Asp Ser Phe Val Ile Arg Gly Asp Glu Val Arg Gln Ile Ala Pro Gly 385 390 395 400		
Gln Thr Gly Lys Ile Ala Asp Tyr Asn Tyr Lys Leu Pro Asp Asp Phe 405 410 415		
Thr Gly Cys Val Ile Ala Trp Asn Ser Asn Asn Leu Asp Ser Lys Val 420 425 430		
Gly Gly Asn Tyr Asn Tyr Leu Tyr Arg Leu Phe Arg Lys Ser Asn Leu 435 440 445		
Lys Pro Phe Glu Arg Asp Ile Ser Thr Glu Ile Tyr Gln Ala Gly Ser 450 455 460		
Thr Pro Cys Asn Gly Val Glu Gly Phe Asn Cys Tyr Phe Pro Leu Gln 465 470 475 480		
Ser Tyr Gly Phe Gln Pro Thr Asn Gly Val Gly Tyr Gln Pro Tyr Arg 485 490 495		
Val Val Val Leu Ser Phe Glu Leu Leu His Ala Pro Ala Thr Val Cys 500 505 510		
Gly Pro Lys Lys Ser Thr Asn Leu Val Lys Asn Lys Cys Val Asn Phe 515 520 525		
Asn Phe Asn Gly Leu Thr Gly Thr Gly Val Leu Thr Glu Ser Asn Lys 530 535 540		
Lys Phe Leu Pro Phe Gln Gln Phe Gly Arg Asp Ile Ala Asp Thr Thr 545 550 555 560		
Asp Ala Val Arg Asp Pro Gln Thr Leu Glu Ile Leu Asp Ile Thr Pro 565 570 575		
Cys Ser Phe Gly Gly Val Ser Val Ile Thr Pro Gly Thr Asn Thr Ser 580 585 590		
Asn Gln Val Ala Val Leu Tyr Gln Asp Val Asn Cys Thr Glu Val Pro 595 600 605		
Val Ala Ile His Ala Asp Gln Leu Thr Pro Thr Trp Arg Val Tyr Ser 610 615 620		
Thr Gly Ser Asn Val Phe Gln Thr Arg Ala Gly Cys Leu Ile Gly Ala 625 630 635 640		
Glu His Val Asn Asn Ser Tyr Glu Cys Asp Ile Pro Ile Gly Ala Gly 645 650 655		
Ile Cys Ala Ser Tyr Gln Thr Gln Thr Asn Ser Pro Arg Arg Ala Arg 660 665 670		
Ser Val Ala Ser Gln Ser Ile Ile Ala Tyr Thr Met Ser Leu Gly Ala 675 680 685		
Glu Asn Ser Val Ala Tyr Ser Asn Asn Ser Ile Ala Ile Pro Thr Asn 690 695 700		
Phe Thr Ile Ser Val Thr Thr Glu Ile Leu Pro Val Ser Met Thr Lys 705 710 715 720		
Thr Ser Val Asp Cys Thr Met Tyr Ile Cys Gly Asp Ser Thr Glu Cys 725 730 735		
Ser Asn Leu Leu Leu Gln Tyr Gly Ser Phe Cys Thr Gln Leu Asn Arg 740 745 750		
Ala Leu Thr Gly Ile Ala Val Glu Gln Asp Lys Asn Thr Gln Glu Val 755 760 765		
Phe Ala Gln Val Lys Gln Ile Tyr Lys Thr Pro Pro Ile Lys Asp Phe 770 775 780		
Gly Gly Phe Asn Phe Ser Gln Ile Leu Pro Asp Pro Ser Lys Pro Ser 785 790 795 800		

-continued

Lys Arg Ser Phe Ile Glu Asp Leu Leu Phe Asn Lys Val Thr Leu Ala
 805 810 815
 Asp Ala Gly Phe Ile Lys Gln Tyr Gly Asp Cys Leu Gly Asp Ile Ala
 820 825 830
 Ala Arg Asp Leu Ile Cys Ala Gln Lys Phe Asn Gly Leu Thr Val Leu
 835 840 845
 Pro Pro Leu Leu Thr Asp Glu Met Ile Ala Gln Tyr Thr Ser Ala Leu
 850 855 860
 Leu Ala Gly Thr Ile Thr Ser Gly Trp Thr Phe Gly Ala Gly Ala Ala
 865 870 875 880
 Leu Gln Ile Pro Phe Ala Met Gln Met Ala Tyr Arg Phe Asn Gly Ile
 885 890 895
 Gly Val Thr Gln Asn Val Leu Tyr Glu Asn Gln Lys Leu Ile Ala Asn
 900 905 910
 Gln Phe Asn Ser Ala Ile Gly Lys Ile Gln Asp Ser Leu Ser Ser Thr
 915 920 925
 Ala Ser Ala Leu Gly Lys Leu Gln Asp Val Val Asn Gln Asn Ala Gln
 930 935 940
 Ala Leu Asn Thr Leu Val Lys Gln Leu Ser Ser Asn Phe Gly Ala Ile
 945 950 955 960
 Ser Ser Val Leu Asn Asp Ile Leu Ser Arg Leu Asp Pro Pro Glu Ala
 965 970 975
 Glu Val Gln Ile Asp Arg Leu Ile Thr Gly Arg Leu Gln Ser Leu Gln
 980 985 990
 Thr Tyr Val Thr Gln Gln Leu Ile Arg Ala Ala Glu Ile Arg Ala Ser
 995 1000 1005
 Ala Asn Leu Ala Ala Thr Lys Met Ser Glu Cys Val Leu Gly Gln
 1010 1015 1020
 Ser Lys Arg Val Asp Phe Cys Gly Lys Gly Tyr His Leu Met Ser
 1025 1030 1035
 Phe Pro Gln Ser Ala Pro His Gly Val Val Phe Leu His Val Thr
 1040 1045 1050
 Tyr Val Pro Ala Gln Glu Lys Asn Phe Thr Thr Ala Pro Ala Ile
 1055 1060 1065
 Cys His Asp Gly Lys Ala His Phe Pro Arg Glu Gly Val Phe Val
 1070 1075 1080
 Ser Asn Gly Thr His Trp Phe Val Thr Gln Arg Asn Phe Tyr Glu
 1085 1090 1095
 Pro Gln Ile Ile Thr Thr Asp Asn Thr Phe Val Ser Gly Asn Cys
 1100 1105 1110
 Asp Val Val Ile Gly Ile Val Asn Asn Thr Val Tyr Asp Pro Leu
 1115 1120 1125
 Gln Pro Glu Leu Asp Ser Phe Lys Glu Glu Leu Asp Lys Tyr Phe
 1130 1135 1140
 Lys Asn His Thr Ser Pro Asp Val Asp Leu Gly Asp Ile Ser Gly
 1145 1150 1155
 Ile Asn Ala Ser Val Val Asn Ile Gln Lys Glu Ile Asp Arg Leu
 1160 1165 1170
 Asn Glu Val Ala Lys Asn Leu Asn Glu Ser Leu Ile Asp Leu Gln
 1175 1180 1185
 Glu Leu Gly Lys Tyr Glu Gln Tyr Ile Lys Trp Pro Trp Tyr Ile
 1190 1195 1200

-continued

Trp Leu Gly Phe Ile Ala Gly Leu Ile Ala Ile Val Met Val Thr
 1205 1210 1215

Ile Met Leu Cys Cys Met Thr Ser Cys Cys Ser Cys Leu Lys Gly
 1220 1225 1230

Cys Cys Ser Cys Gly Ser Cys Cys Lys Phe Asp Glu Asp Asp Ser
 1235 1240 1245

Glu Pro Val Leu Lys Gly Val Lys Leu His Tyr Thr
 1250 1255 1260

<210> SEQ ID NO 14
 <211> LENGTH: 1260
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 14

Gln Cys Val Asn Leu Thr Thr Arg Thr Gln Leu Pro Pro Ala Tyr Thr
 1 5 10 15

Asn Ser Phe Thr Arg Gly Val Tyr Tyr Pro Asp Lys Val Phe Arg Ser
 20 25 30

Ser Val Leu His Ser Thr Gln Asp Leu Phe Leu Pro Phe Phe Ser Asn
 35 40 45

Val Thr Trp Phe His Ala Ile His Val Ser Gly Thr Asn Gly Thr Lys
 50 55 60

Arg Phe Asp Asn Pro Val Leu Pro Phe Asn Asp Gly Val Tyr Phe Ala
 65 70 75 80

Ser Thr Glu Lys Ser Asn Ile Ile Arg Gly Trp Ile Phe Gly Thr Thr
 85 90 95

Leu Asp Ser Lys Thr Gln Ser Leu Leu Ile Val Asn Asn Ala Thr Asn
 100 105 110

Val Val Ile Lys Val Cys Glu Phe Gln Phe Cys Asn Asp Pro Phe Leu
 115 120 125

Gly Val Tyr Tyr His Lys Asn Asn Lys Ser Trp Met Glu Ser Glu Phe
 130 135 140

Arg Val Tyr Ser Ser Ala Asn Asn Cys Thr Phe Glu Tyr Val Ser Gln
 145 150 155 160

Pro Phe Leu Met Asp Leu Glu Gly Lys Gln Gly Asn Phe Lys Asn Leu
 165 170 175

Arg Glu Phe Val Phe Lys Asn Ile Asp Gly Tyr Phe Lys Ile Tyr Ser
 180 185 190

Lys His Thr Pro Ile Asn Leu Val Arg Asp Leu Pro Gln Gly Phe Ser
 195 200 205

Ala Leu Glu Pro Leu Val Asp Leu Pro Ile Gly Ile Asn Ile Thr Arg
 210 215 220

Phe Gln Thr Leu Leu Ala Leu His Arg Ser Tyr Leu Thr Pro Gly Asp
 225 230 235 240

Ser Ser Ser Gly Trp Thr Ala Gly Ala Ala Ala Tyr Tyr Val Gly Tyr
 245 250 255

Leu Gln Pro Arg Thr Phe Leu Leu Lys Tyr Asn Glu Asn Gly Thr Ile
 260 265 270

Thr Asp Ala Val Asp Cys Ala Leu Asp Pro Leu Ser Glu Thr Lys Cys
 275 280 285

Thr Leu Lys Ser Phe Thr Val Glu Lys Gly Ile Tyr Gln Thr Ser Asn
 290 295 300

-continued

Phe	Arg	Val	Gln	Pro	Thr	Glu	Ser	Ile	Val	Arg	Phe	Pro	Asn	Ile	Thr	305	310	315	320
Asn	Leu	Cys	Pro	Phe	Gly	Glu	Val	Phe	Asn	Ala	Thr	Arg	Phe	Ala	Ser	325	330	335	
Val	Tyr	Ala	Trp	Asn	Arg	Lys	Arg	Ile	Ser	Asn	Cys	Val	Ala	Asp	Tyr	340	345	350	
Ser	Val	Leu	Tyr	Asn	Ser	Ala	Ser	Phe	Ser	Thr	Phe	Lys	Cys	Tyr	Gly	355	360	365	
Val	Ser	Pro	Thr	Lys	Leu	Asn	Asp	Leu	Cys	Phe	Thr	Asn	Val	Tyr	Ala	370	375	380	
Asp	Ser	Phe	Val	Ile	Arg	Gly	Asp	Glu	Val	Arg	Gln	Ile	Ala	Pro	Gly	385	390	395	400
Gln	Thr	Gly	Lys	Ile	Ala	Asp	Tyr	Asn	Tyr	Lys	Leu	Pro	Asp	Asp	Phe	405	410	415	
Thr	Gly	Cys	Val	Ile	Ala	Trp	Asn	Ser	Asn	Asn	Leu	Asp	Ser	Lys	Val	420	425	430	
Gly	Gly	Asn	Tyr	Asn	Tyr	Arg	Tyr	Arg	Leu	Phe	Arg	Lys	Ser	Asn	Leu	435	440	445	
Lys	Pro	Phe	Glu	Arg	Asp	Ile	Ser	Thr	Glu	Ile	Tyr	Gln	Ala	Gly	Ser	450	455	460	
Thr	Pro	Cys	Asn	Gly	Val	Glu	Gly	Phe	Asn	Cys	Tyr	Phe	Pro	Leu	Gln	465	470	475	480
Ser	Tyr	Gly	Phe	Gln	Pro	Thr	Tyr	Gly	Val	Gly	Tyr	Gln	Pro	Tyr	Arg	485	490	495	
Val	Val	Val	Leu	Ser	Phe	Glu	Leu	Leu	His	Ala	Pro	Ala	Thr	Val	Cys	500	505	510	
Gly	Pro	Lys	Lys	Ser	Thr	Asn	Leu	Val	Lys	Asn	Lys	Cys	Val	Asn	Phe	515	520	525	
Asn	Phe	Asn	Gly	Leu	Thr	Gly	Thr	Gly	Val	Leu	Thr	Glu	Ser	Asn	Lys	530	535	540	
Lys	Phe	Leu	Pro	Phe	Gln	Gln	Phe	Gly	Arg	Asp	Ile	Ala	Asp	Thr	Thr	545	550	555	560
Asp	Ala	Val	Arg	Asp	Pro	Gln	Thr	Leu	Glu	Ile	Leu	Asp	Ile	Thr	Pro	565	570	575	
Cys	Ser	Phe	Gly	Gly	Val	Ser	Val	Ile	Thr	Pro	Gly	Thr	Asn	Thr	Ser	580	585	590	
Asn	Gln	Val	Ala	Val	Leu	Tyr	Gln	Gly	Val	Asn	Cys	Thr	Glu	Val	Pro	595	600	605	
Val	Ala	Ile	His	Ala	Asp	Gln	Leu	Thr	Pro	Thr	Trp	Arg	Val	Tyr	Ser	610	615	620	
Thr	Gly	Ser	Asn	Val	Phe	Gln	Thr	Arg	Ala	Gly	Cys	Leu	Ile	Gly	Ala	625	630	635	640
Glu	His	Val	Asn	Asn	Ser	Tyr	Glu	Cys	Asp	Ile	Pro	Ile	Gly	Ala	Gly	645	650	655	
Ile	Cys	Ala	Ser	Tyr	Gln	Thr	Gln	Thr	Asn	Ser	Pro	Arg	Arg	Ala	Arg	660	665	670	
Ser	Val	Ala	Ser	Gln	Ser	Ile	Ile	Ala	Tyr	Thr	Met	Ser	Leu	Gly	Ala	675	680	685	
Glu	Asn	Ser	Val	Ala	Tyr	Ser	Asn	Asn	Ser	Ile	Ala	Ile	Pro	Thr	Asn	690	695	700	
Phe	Thr	Ile	Ser	Val	Thr	Thr	Glu	Ile	Leu	Pro	Val	Ser	Met	Thr	Lys	705	710	715	720
Thr	Ser	Val	Asp	Cys	Thr	Met	Tyr	Ile	Cys	Gly	Asp	Ser	Thr	Glu	Cys				

-continued

725							730					735			
Ser	Asn	Leu	Leu	Gln	Tyr	Gly	Ser	Phe	Cys	Thr	Gln	Leu	Asn	Arg	
		740					745					750			
Ala	Leu	Thr	Gly	Ile	Ala	Val	Glu	Gln	Asp	Lys	Asn	Thr	Gln	Glu	Val
		755					760					765			
Phe	Ala	Gln	Val	Lys	Gln	Ile	Tyr	Lys	Thr	Pro	Pro	Ile	Lys	Asp	Phe
	770					775					780				
Gly	Gly	Phe	Asn	Phe	Ser	Gln	Ile	Leu	Pro	Asp	Pro	Ser	Lys	Pro	Ser
785					790					795					800
Lys	Arg	Ser	Phe	Ile	Glu	Asp	Leu	Leu	Phe	Asn	Lys	Val	Thr	Leu	Ala
				805					810					815	
Asp	Ala	Gly	Phe	Ile	Lys	Gln	Tyr	Gly	Asp	Cys	Leu	Gly	Asp	Ile	Ala
			820						825				830		
Ala	Arg	Asp	Leu	Ile	Cys	Ala	Gln	Lys	Phe	Asn	Gly	Leu	Thr	Val	Leu
		835					840					845			
Pro	Pro	Leu	Leu	Thr	Asp	Glu	Met	Ile	Ala	Gln	Tyr	Thr	Ser	Ala	Leu
		850				855					860				
Leu	Ala	Gly	Thr	Ile	Thr	Ser	Gly	Trp	Thr	Phe	Gly	Ala	Gly	Ala	Ala
865					870					875					880
Leu	Gln	Ile	Pro	Phe	Ala	Met	Gln	Met	Ala	Tyr	Arg	Phe	Asn	Gly	Ile
				885					890					895	
Gly	Val	Thr	Gln	Asn	Val	Leu	Tyr	Glu	Asn	Gln	Lys	Leu	Ile	Ala	Asn
			900						905				910		
Gln	Phe	Asn	Ser	Ala	Ile	Gly	Lys	Ile	Gln	Asp	Ser	Leu	Ser	Ser	Thr
		915						920					925		
Ala	Ser	Ala	Leu	Gly	Lys	Leu	Gln	Asp	Val	Val	Asn	Gln	Asn	Ala	Gln
		930					935					940			
Ala	Leu	Asn	Thr	Leu	Val	Lys	Gln	Leu	Ser	Ser	Asn	Phe	Gly	Ala	Ile
945					950						955				960
Ser	Ser	Val	Leu	Asn	Asp	Ile	Leu	Ser	Arg	Leu	Asp	Pro	Pro	Glu	Ala
				965					970					975	
Glu	Val	Gln	Ile	Asp	Arg	Leu	Ile	Thr	Gly	Arg	Leu	Gln	Ser	Leu	Gln
			980						985				990		
Thr	Tyr	Val	Thr	Gln	Gln	Leu	Ile	Arg	Ala	Ala	Glu	Ile	Arg	Ala	Ser
		995					1000						1005		
Ala	Asn	Leu	Ala	Ala	Thr	Lys	Met	Ser	Glu	Cys	Val	Leu	Gly	Gln	
	1010						1015					1020			
Ser	Lys	Arg	Val	Asp	Phe	Cys	Gly	Lys	Gly	Tyr	His	Leu	Met	Ser	
	1025						1030					1035			
Phe	Pro	Gln	Ser	Ala	Pro	His	Gly	Val	Val	Phe	Leu	His	Val	Thr	
	1040						1045					1050			
Tyr	Val	Pro	Ala	Gln	Glu	Lys	Asn	Phe	Thr	Thr	Ala	Pro	Ala	Ile	
	1055						1060					1065			
Cys	His	Asp	Gly	Lys	Ala	His	Phe	Pro	Arg	Glu	Gly	Val	Phe	Val	
	1070						1075					1080			
Ser	Asn	Gly	Thr	His	Trp	Phe	Val	Thr	Gln	Arg	Asn	Phe	Tyr	Glu	
	1085						1090					1095			
Pro	Gln	Ile	Ile	Thr	Thr	Asp	Asn	Thr	Phe	Val	Ser	Gly	Asn	Cys	
	1100						1105					1110			
Asp	Val	Val	Ile	Gly	Ile	Val	Asn	Asn	Thr	Val	Tyr	Asp	Pro	Leu	
	1115						1120					1125			
Gln	Pro	Glu	Leu	Asp	Ser	Phe	Lys	Glu	Glu	Leu	Asp	Lys	Tyr	Phe	
	1130						1135					1140			

-continued

Lys Asn His Thr Ser Pro Asp Val Asp Leu Gly Asp Ile Ser Gly
1145 1150 1155

Ile Asn Ala Ser Val Val Asn Ile Gln Lys Glu Ile Asp Arg Leu
1160 1165 1170

Asn Glu Val Ala Lys Asn Leu Asn Glu Ser Leu Ile Asp Leu Gln
1175 1180 1185

Glu Leu Gly Lys Tyr Glu Gln Tyr Ile Lys Trp Pro Trp Tyr Ile
1190 1195 1200

Trp Leu Gly Phe Ile Ala Gly Leu Ile Ala Ile Val Met Val Thr
1205 1210 1215

Ile Met Leu Cys Cys Met Thr Ser Cys Cys Ser Cys Leu Lys Gly
1220 1225 1230

Cys Cys Ser Cys Gly Ser Cys Cys Lys Phe Asp Glu Asp Asp Ser
1235 1240 1245

Glu Pro Val Leu Lys Gly Val Lys Leu His Tyr Thr
1250 1255 1260

<210> SEQ ID NO 15
<211> LENGTH: 396
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthesized sequence

<400> SEQUENCE: 15

caccaccacc accaccactg actcgaggct ggagcctcgg tggccatgct tcttgcccct 60
tgggcctccc ccagcccct cctccccttc ctgcacccgt acccccgggt ctttgagatc 120
tggttaccac taaaccagcc tcaagaacac ccgaatggag tctctaagct acataatacc 180
aacttacact ttacaaaatg ttgtccccc aatgtagcc attcgatatc gctccgttgc 240
caaagaaagt ttcttcacat tctaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 300
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 360
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaggc 396

The invention claimed is:

1. A transgene that encodes:

- i) an RNA polymerase promoter recognized by a polymerase of a single-subunit DNA-dependent RNA polymerase (ssRNAP) family;
- ii) a 5' UTR, wherein the 5'-UTR sequence is:

5' -GAATAAGAGAGAAAAGAAGAGTAAGAAGA

AATATAAGAGCTAGCGGTAC-3'.

iii) a secretory sequence;

- iv) a coronavirus spike protein component, wherein the spike protein component incorporates a variant sequence at amino acid position 614 of a native spike protein or a position that corresponds with the amino acid position at position 614 of the native spike protein, and additionally comprises a variant sequence at one or both of amino acid positions 452 and 501 of the native spike protein, wherein the amino acid position of the variant sequence is determined from a methionine start codon of the native spike protein and wherein the variant sequence at position 452 is L452R, the variant sequence at position 501 is N501Y and the variant sequence at position 614 is D614G; and

v) a 3' UTR and poly A sequence.

2. The transgene of claim 1, wherein the promoter is selected from a T7 promoter, T3 promoter, Sp6 promoter, mitochondrial RNA polymerase (POLRMT) promoter and chloroplastic ssRNAP promoter.

3. The transgene of claim 1, wherein the promoter is the T7 promoter.

4. The transgene of claim 1, wherein the spike protein component is a full-length SARS-COV2 spike protein or an antigenic fragment thereof.

5. The transgene of claim 4, wherein the antigenic fragment comprises an N-terminal fragment up to at least amino acid position 614 of the spike protein, a C-terminal fragment including the amino acid at position 614 of the spike protein, or an internal fragment incorporating at least the region V595-G614 or the region G614-A647 of the spike protein.

6. The transgene of claim 1, wherein the protein spike component additionally comprises a variant amino acid sequence selected from the group of: T478K, E484K, E484Q, P681R, P681H, K986P and V987P.

7. The transgene of claim 1, comprising an upstream promoter spacer sequence.

8. The transgene of claim 1, wherein the 5'-UTR sequence is a native coronavirus 5'-UTR.

61

9. The transgene of claim 1, wherein 3'-UTR is a native coronavirus 3'-UTR.

10. The transgene of claim 1, wherein 3'-UTR comprises one or more of the 3'-UTR of alpha-1-globin, beta-globin, c-fos, metallothionein IG, lipoprotein lipase, glutathione peroxidase 4, glutathione peroxidase 3, glutathione peroxidase 2, c-myc, 15-lipoxygenase, and transferrin receptor 2a.

11. The transgene of claim 10, wherein the 3'-UTR comprises the 3'-UTR of at least one of alpha-1-globin and beta-globin.

12. The transgene of claim 1, wherein the secretory sequence is selected from an IgE receptor secretory sequence, erythropoietin secretory sequence, growth hormone secretory sequence, granulocyte colony stimulating factor secretory sequence, fibroblast growth factor secretory sequence, vascular endothelial growth factor secretory sequence, a G-protein coupled receptor secretory sequence, or a modified secretory sequence thereof that retains the ability to cleave from the spike protein component.

13. The transgene of claim 1, comprising a T7 promoter, an IgE receptor secretory sequence, and a 3'-UTR comprising the 3'-UTR of alpha-1-globin and the 3'-UTR of beta-globin.

14. The transgene of claim 1, wherein the spike protein component additionally comprises the variants, K986P and V987P.

15. The transgene of claim 13, wherein the 3'-UTR comprises the nucleotide sequence of SEQ ID NO: 7.

16. A vaccine incorporating a transgene as defined in claim 1 or an mRNA transcript thereof.

17. The vaccine of claim 16, which is an mRNA vaccine, DNA vaccine, subunit vaccine, non-replicating viral vaccine, attenuated viral vaccine or inactivated viral vaccine.

18. The vaccine of claim 17, which is an mRNA vaccine.

62

19. The vaccine of claim 18, wherein the mRNA comprises a 5'-methylated cap.

20. The vaccine of claim 18, formulated for administration in lipid nanoparticles.

21. The vaccine of claim 18, comprising the mRNA in a buffer combined with lipids in a ratio of 3:1 to 7:1.

22. The vaccine of claim 18, wherein the pH of the formulation is 7.2 ± 0.2 .

23. The vaccine of claim 20, wherein the lipid nanoparticles have an average particle size of 85 ± 20 nm.

24. The vaccine of claim 20, wherein the lipid nanoparticles comprise an ionizable cationic lipid, a PEGylated lipid, a phospholipid, and cholesterol.

25. The vaccine of claim 20, wherein the lipid nanoparticles comprise MC3 ([[(6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl] 4-(dimethylamino) butanoate), DSPC ([[(2R)-2,3-di (octadecanoyloxy) propyl] 2-(trimethylazaniumyl) ethyl phosphate), dimyristoyl-rac-glycero-3-methoxypolyethylene glycol-2000 and cholesterol.

26. The vaccine of claim 25, comprising MC3; DSPC; cholesterol; and DMG-PEG2000 in a molar ratio of 50:10:38.5:1.5.

27. A method of vaccinating a mammal against a coronavirus infection comprising administering to the mammal a vaccine incorporating a transgene as defined in claim 1 or an mRNA transcript thereof.

28. The method of claim 27, wherein the vaccine is an mRNA vaccine formulated in lipid nanoparticles.

29. The method of claim 27, wherein the vaccine is administered at a dosage of 1-300 μ g.

30. The method of claim 27, wherein the mammal is a human.

* * * * *