

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12414563
Kind Code	B2
Date of Patent	September 16, 2025
Inventor(s)	Zhang; Hong et al.

Encapsulation of cyclohexanediones process and product

Abstract

Encapsulation of cyclohexanediones process and product. A composition particularly adapted for herbicidal use is described, including biopolymer micro-capsules containing one or more cyclohexanediones encapsulated therein, resulting in enhanced chemical stability and dispersibility of the encapsulated cyclohexanediones for such use. A complex coacervation method of forming such microcapsules is also described.

Inventors:	Zhang; Hong (Cary, NC), Branaghan; Rachel Louise (Fishbourne, GB), Maude; Sarah Jane (Chadbury, GB)
Applicant:	Arysta LifeScience Inc. (Cary, NC)
Family ID:	1000008820743
Appl. No.:	17/291508
Filed (or PCT Filed):	November 12, 2019
PCT No.:	PCT/US2019/060981
PCT Pub. No.:	WO2020/102232
PCT Pub. Date:	May 22, 2020

Prior Publication Data

Document Identifier	Publication Date
US 20220000105 A1	Jan. 06, 2022

Related U.S. Application Data

us-provisional-application US 62760565 20181113

Publication Classification

Int. Cl.: **A01N35/10** (20060101); **A01N25/02** (20060101); **A01N25/10** (20060101); **A01N25/22** (20060101); **A01N25/28** (20060101)

U.S. Cl.:

CPC **A01N35/10** (20130101); **A01N25/02** (20130101); **A01N25/10** (20130101); **A01N25/22** (20130101); **A01N25/28** (20130101);

Field of Classification Search

CPC: A01N (25/28); A01N (35/10)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
4952722	12/1989	Serban et al.	N/A	N/A
5922357	12/1998	Coombes et al.	N/A	N/A
2010/0248963	12/2009	Becher	504/144	A01N 37/26
2011/0301036	12/2010	Tank et al.	N/A	N/A
2014/0080712	12/2013	Lao et al.	N/A	N/A
2018/0139956	12/2017	Ishibashi	N/A	A01N 47/12

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
101427672	12/2008	CN	N/A
102388864	12/2011	CN	N/A
101396018	12/2011	CN	N/A
101990890	12/2012	CN	N/A
2488437	12/2012	RU	N/A
2005102045	12/2004	WO	N/A
2010093970	12/2009	WO	N/A
2012080208	12/2011	WO	N/A
2013158620	12/2012	WO	N/A

OTHER PUBLICATIONS

International Search Report and Written Opinion for International Application PCT/US2019/060981; International Filing Date: Nov. 11, 2019; Date of Mailing: Jan. 31, 2020; 10 pages. cited by applicant

Primary Examiner: Justice; Gina C

Attorney, Agent or Firm: CANTOR COLBURN LLP

Background/Summary

CROSS REFERENCE TO RELATED APPLICATIONS (1) This application is a National Stage application of PCT/US2019/060981, filed Nov. 12, 2019, which claims the benefit of priority to U.S. Provisional Patent Application 62/760,565, filed Nov. 13, 2018, both of which are incorporated by reference in their entirety herein.

(1) The field of art to which this invention generally pertains is products and methods of making products which are useful in fungicidal, insecticidal and herbicidal compositions.

TECHNICAL FIELD

Background

(2) While many chemical products are known to be very useful as materials for herbicidal and insecticidal applications, because of their complexity, there is a constant search for ways to improve the efficiency and effectiveness of such products such as, for example, improving their stability and dispersibility.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) The FIGURE depicts a representative complex coacervation process for making products as described herein.

BRIEF SUMMARY

(2) A composition particularly adapted for herbicidal use is described including one or more biopolymer containing micro-capsules containing one or more cyclohexanediones encapsulated therein, resulting in enhanced chemical stability and dispersibility of the encapsulated cyclohexanediones for such use.

(3) Additional embodiments include, the composition described above where the cyclohexanedione comprises Clethodim; the composition described above where the biopolymer comprises polysaccharide and/or cellulose; the composition described above where the polysaccharide comprises Inulin; the composition described above including one or more surfactants; the composition described above where the cyclohexanediones are present in the micro-capsules in an amount of 1% to 70% by weight; the composition described above made by complex coacervation; the composition described above where the composition optionally contains one or more natural gum, starch, protein, alcohol, water soluble polymer, and/or an oil; the composition described above where the alcohol comprises Sorbitol; the composition described above where the natural gum comprises Gum Arabic; the composition described above where the surfactant comprises one or more polyalkylene oxide block copolymers; a water dispersion containing the composition described above; and an oil dispersion containing the composition described above; the composition described above where the oil comprises vegetable oil; the composition described above where the oil comprises soybean oil, pine oil, and/or sunflower oil; the composition described above where the oil comprises paraffinic oil; the composition described above where the oil comprises an aromatic solvent; the composition described above where the solvent comprises C.sub.9 to C.sub.10 dialkyl and trialkyl benzenes (such as, for example, Aromatic 100, Aromatic 150 or Aromatic 200); the composition described above where the vegetable oil comprises rapeseed oil/or methylated rapeseed oil; the composition described above where the oil comprises a polyoxyethylene sorbitan monolaurate with a carbon chain length varying from C.sub.10 to C.sub.20 (such as, for example, Tween® 80, or Tween® 20, or Tween® 85, or Tween® 60 (Sigma-Aldrich); the composition described above where the polyoxyethylene sorbitan is a monolaurate, monostearate, tristearate, and/or trioleate.

(4) A method of controlling insects and/or weeds comprising applying the composition described above is also described.

(5) Additional embodiments described include the method described above where the applying step is performed post-emergently after the crop breaks ground; and the method described above where the applying step is performed pre-emergently before the crop breaks ground.

(6) These, and additional embodiments, will be apparent from the following descriptions.

DETAILED DESCRIPTION

(7) The particulars shown herein are by way of example and for purposes of illustrative discussion of the various embodiments of the present invention only and are presented in the cause of providing what is believed to be the most useful and readily understood description of the principles and conceptual aspects of the invention. In this regard, no attempt is made to show details of the invention in more detail than is necessary for a fundamental understanding of the invention, the description making apparent to those skilled in the art how the several forms of the invention may be embodied in practice

(8) The present invention will now be described by reference to more detailed embodiments. This invention may, however, be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art.

(9) Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for describing particular embodiments only and is not intended to be limiting of the invention. As used in the description of the invention and the appended claims, the singular forms “a,” “an,” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise. All publications, patent applications, patents, and other references mentioned herein are expressly incorporated by reference in their entirety.

(10) Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the following specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention. At the very least, and not as an attempt to limit the application of the doctrine of equivalents to the scope of the claims, each numerical parameter should be construed in light of the number of significant digits and ordinary rounding approaches.

(11) Notwithstanding that the numerical ranges and parameters setting forth the broad scope of the invention are approximations, the numerical values set forth in the specific examples are reported as precisely as possible. Any numerical value, however, inherently contains certain errors necessarily resulting from the standard deviation found in their respective testing measurements. Every numerical range given throughout this specification will include every narrower numerical range that falls within such broader numerical range, as if such narrower numerical ranges were all expressly written herein.

(12) Additional advantages of the invention will be set forth in part in the description which follows, and in part will be obvious from the description, or may be learned by practice of the invention. It is to be understood that both the foregoing general description and the following detailed description are exemplary and explanatory only and are not restrictive of the invention, as claimed.

(13) What is described herein is the encapsulation of cyclohexanediones to improve both their chemical stability and dispersibility, including improving their compatibility with other premix molecules and different mixtures, sustainable release, extend release profile, and longer crop protection, among others. Through the use of biopolymer encapsulation materials (such as polysaccharides and cellulose, for example—see also the tables), cyclohexanediones (such as

Clethodim, for example) microcapsules are formed through conventional complex coacervation techniques. However, by carefully selecting this encapsulation technology and one or more appropriate surfactants, the new delivery system demonstrates many advantages, including the following to name just a few: significant chemical stability (e.g., for Clethodim, at least 2 weeks at 54° C. as compared to currently available Clethodim products); stability of microcapsules after aging at accelerated temperatures in either a dry form or an oil dispersion (e.g., for Clethodim products); high dispersibility of microcapsules in either aqueous phase or oil phase; controlled release of the encapsulated active ingredient; significantly high loading of active ingredient in either a dry form or an oil dispersion with a target of a practical formulation volume for the particular application method and, the oil phase acts as a built-in adjuvant for improving bio-efficacy for oil dispersions.

Example 1

(14) A representative complex coacervation process with the selected appropriate surfactants is described below. See, also, the FIGURE. The composition materials are first hydrated and thoroughly mixed (1). The solutions are prepared by dispersing the composition materials in a concentration from 2% to 50% by weight based on their molecular weights and the desired viscosity of the final solutions, i.e. balanced viscosity for product handling and minimized phase separation. The solutions are gently stirred and stored, for example, overnight to ensure complete hydration of the biopolymers (2). Heat may be applied to enhance dissolving if necessary. The biopolymer solution or a mixture of biopolymer solutions are mixed in a beaker with appropriate surfactants (3). A Clethodim solution in a concentration from 50% to 80% is then metered into the breaker (4) and homogenized (5) for 10 minutes at 5,000 to 10,000 rpm (revolutions per minute) in a conventional mixer to generate micro-emulsions. The micro-emulsion mixture is further spray-dried (6) with an air flow whose inlet temperature is from 110° C. to 120° C. The microcapsule powder is collected and analyzed.

Example 2

(15) Alternatively, the microcapsule powder can be further made into an Oil Dispersion by suspending the powder into one or more selected oils in a concentration from 20% to 90% under agitation. Optionally, one or more surfactants including dispersants, emulsifiers, rheology aids are added into the Oil Dispersion to stabilize the mixture. The oil dispersion is analyzed below.

Example 3

(16) Representative encapsulation efficiency with various deliver systems is demonstrated below. To determine total oil content, 0.5 g (grams) the encapsulated powder is mixed with 10 ml (milliliters) of water and stirred for 2 minutes. 10 ml of isopropanol and hexane (1:1 to 1:3 ratio) is added and stirred for 5 minutes. The solution is then centrifuged, filtered and placed in a water bath at 70° C. to evaporate the solvent. The final weight of the oil is recorded. To determine surface oil content, 0.5 g of encapsulated powder is mixed with hexane, centrifuged and filtered. The supernatant is evaporated as above and the quantity of oil recorded. Results are given in Table 1 below.

(17) TABLE-US-00001 TABLE 1 Sample A B C D E Encapsulation Efficiency 95.5% 95.1% 96.1% 95.8% 96.2%

Example 4

(18) Chemical retention of encapsulated Clethodim powder stored at 54° C. is compared with some commercial EC (emulsifiable concentrate) products currently on the market,

(19) TABLE-US-00002 TABLE 2 Sample Commercial 1 .sup.(1) Commercial 2 .sup.(2) F G H I J K Day 7 87.3% 90.7% 96.0% 99.7% 98.5% 97.7% >100% >100% Day 14 87.2% 88.2% 92.4% 94.3% 94.6% 93.5% 95.3% 96.1% Notes: .sup.(1) The commercial EC product contains 120 g/L of Clethodim .sup.(2) The commercial EC product contains 360 g/L of Clethodim

(20) Chemical retention of encapsulated Clethodim oil dispersions stored at 54° C. is demonstrated in Table 3.

(21) TABLE-US-00003 TABLE 3 Sample L M N O Stored for 7 Days 97.3% >100% 99.8%

>100% Stored for 14 Days 97.7% >100% >100% 98.6%

(22) Dispersibility at 2.5% OD (oil dispersion) in dilution in ASTM 100 milliliter graduated centrifuge tubes is demonstrated in Table 4.

(23) TABLE-US-00004 TABLE 4 Sedimentation (ml) at time Comments Number of inversions Number of period (hours) after for Re-Dispersion Sample Bloom inversions* 1 2 24 24 hours (ASTM Testing Tube) P Moderate 3 0 0 0 Slight 1 creaming Q Good 1 0 0 0 Slight 1 creaming

*Inversion is to flip over testing tubes upside down to mix the solution to re-suspend or re-mix. The number of inversions is to evaluate the ease of re-mixing the solution. The lower the better.

Example 5

(24) Representative emulsion-in-water compositions (before spray-drying) are shown in Tables 5 and 6. Representative compositions of encapsulated powder suspended in oil to form oil dispersions are shown in Table 7.

(25) TABLE-US-00005 TABLE 5 Formulation Component 1 2 3 4 5 6 7 8 9 10 11 12 13 14 Clethodim 20 20 20 20 20 20 20 20 20 20 20 20 20 20 (Cyclohexanedione) Inulin Polysaccharide 5 10 15 — — — 2.5 6.25 3.13 5 5 5 5 5 Gum Arabic Natural Gum 15 10 5 15 10 5 17.5 18.75 21.87 15 15 15 15 15 Maltodextrin Polysaccharide — — — 5 10 15 — — — 2.5 — — — —

Waxy Maize Starch/ — — — — — — — — — — — — — — Polysaccharide Whey Protein — — — — — — — — — — — — — — 2.5 — — — — Xylitol Sugar Alcohol — — — — — — — — — — — — — —

2.5 — — — Sorbitol Sugar Alcohol — — — — — — — — — — — — — — 2.5 — — — Mannitol Sugar Alcohol — — — — — — — — — — — — — — 2.5 Starch Polysaccharide — — — — — — — — — — — — — —

— Cyclodextrin Cyclic oligosaccharide Polyoxyethylene (20) — sorbitan

monolaurate surfactant Alkylated vinyl- — pyrrolidone polymer surfactant Isopropyl alcohol — EO/PO (ethylene

— oxide/propylene oxide) block copolymer surfactant EO/PO block copolymer — surfactant EO/PO block

copolymer — surfactant EO/PO block copolymer — surfactant EO/PO block copolymer —

— surfactant EO/PO block copolymer — surfactant EO/PO block copolymer —

surfactant EO/PO block copolymer — surfactant Polyalkylene oxide block — copolymer surfactant EO/PO block copolymer — surfactant Nonionic block copolymer

— surfactant Butyl block copolymer — surfactant Polyvinylpyrrolidone —

Water soluble polymer Water 60 60 60 60 60 60 60 60 55 55 57.5 57.5 57.5 57.5 57.5 Total 100 100 100 100 100 100 100 100 100 100 100 100 100 100 100 Formulation Component 15 16 17 18 19 20 21 22 23 24 25 26 27 Clethodim 20 20 20 20 20 20 20 25 30 20 20 20 20 20 (Cyclohexanedione) Inulin Polysaccharide 5 — 5 5 5 5 5 5 5 5 5 5 5 5 Gum Arabic Natural Gum 15 15 10 15 15 10 10 15 15 15 15 15 15 Maltodextrin Polysaccharide — — — — — — — — — — — — — —

Waxy Maize Starch/ — Polysaccharide Whey Protein — Xylitol Sugar Alcohol —

Sorbitol Sugar Alcohol — — 5 5 — — — 5 5 5 5 5 5 Mannitol Sugar Alcohol — Starch Polysaccharide 2.5 —

Cyclodextrin Cyclic — 20 — oligosaccharide Polyoxyethylene (20) — 1 — — — — — — — 5 — sorbitan monolaurate surfactant Alkylated vinyl- —

— 1 — pyrrolidone polymer surfactant Isopropyl alcohol — 5 — EO/PO (ethylene oxide) block copolymer surfactant EO/PO block copolymer — 5 — surfactant EO/PO block copolymer — surfactant EO/PO

surfactant EO/PO block copolymer — surfactant EO/PO

surfactant EO/PO block copolymer — surfactant EO/PO

surfactant EO/PO block copolymer — surfactant EO/PO

[illegible]

[illegible]

(28) As demonstrated by the tables above, the water based dispersions can contain functional materials such as: Cyclohexanediones such as Clethodim, Polysaccharides such as Inulin, Natural gums such as Gum Arabic, Polysaccharides such as maltodextrin, starches and polysaccharides such as Waxy Maize Starch, proteins such as Whey Protein, Sugar Alcohols such as Sorbitol and Mannitol, Polysaccharide starches, Cyclic oligosaccharide such as Cyclodextrin; conventional surfactants such as Polyoxyethylene (20) sorbitan monolaurate, Alkylated vinyl-pyrrolidone polymer, Isopropyl alcohol, EO/PO (ethylene oxide/propylene oxide) block copolymer, Anionic polymeric dispersant Polyalkylene oxide block copolymer, nonionic block copolymer, Butyl block copolymer. and Polyvinylpyrrolidone Water soluble polymer.

(29) As also demonstrated by the tables above, the oil based dispersions can also contain functional materials such as: conventional surfactants such as EO/PO (ethylene oxide/propylene oxide) block copolymer, Polyalkylene oxide block copolymer, EO/PO block copolymer, nonionic block copolymer, Butyl block copolymer, Polyvinylpyrrolidone Water soluble polymer, anionic polymeric dispersant, cetyl ether phosphate, oleyl ether phosphate, polyoxyethylene (5) oleyl mono/di-phosphate, polyoxyethylene (3) oleyl mono/di-phosphate, sodium dioctyl-sulfosuccinate, polyoxyethylene (40) sorbitol hexaoleate, ethoxylated castor oil and rapeseed oil fatty acid methyl ester.

(30) These examples are merely illustrations and are not to be understood as limiting the scope and underlying principles of the invention in any way. Various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art form after the following examples and foregoing description. Such modifications are also intended to fall within the scope of the appended claims.

(31) As described herein, these problems and others in this area are addressed by the invention described herein. Thus, the scope of the invention shall include all modifications and variations that may fall within the scope of the attached claims. Other embodiments of the invention will be apparent to those skilled in the art from consideration of the specification and practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with a true scope and spirit of the invention being indicated by the following claims.

Claims

1. A cyclohexanedione composition for herbicidal use, comprising clethodim encapsulated with a biopolymer microcapsule, wherein the biopolymer microcapsule is made by complex coacervation and the biopolymer comprises inulin.
2. The composition of claim 1, further including one or more surfactants, optionally wherein the one or more surfactants comprise one or more polyalkylene oxide block copolymers.
3. The composition of claim 1, wherein the clethodim is present in the micro-capsules in an amount of 1% to 70% by weight.
4. The composition of claim 1, wherein the composition further contains one or more of a natural gum, starch, a protein, an alcohol, a water soluble polymer, and/or an oil.
5. The composition of claim 4, wherein the alcohol comprises sorbitol.
6. The composition of claim 4 wherein the natural gum comprises gum arabic.

7. A water dispersion containing the composition of claim 1.
 8. An oil dispersion containing the composition of claim 1.
 9. The oil dispersion of claim 8, wherein the oil comprises a vegetable oil, soybean oil, pine oil, sunflower oil, rapseed oil, and/or methylated rapseed oil, paraffinic oil or combinations comprising at least one of the foregoing.
 10. The oil dispersion of claim 8, wherein the oil comprises an aromatic solvent, optionally wherein the solvent comprises C.sub.9 to C.sub.10 dialkyl and trialkyl benzenes.
 11. The oil dispersion of claim 8, wherein the oil comprises a polyoxyethylene sorbitan monolaurate with a carbon chain length varying from C.sub.10 to C.sub.20; optionally wherein the polyoxyethylene sorbitan is a monolaurate, monostearate, tristearate, and/or trioleate.
 12. A method of controlling insects and/or weeds in a crop comprising applying the composition of claim 1 to the crop.
 13. The method of claim 12, where the applying step is performed post-emergently after the crop breaks ground.
 14. The method of claim 12, where the applying step is performed pre-emergently before the crop breaks ground.
-