

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent Application Publication

20250276098

Kind Code

A1

Publication Date

September 04, 2025

Inventor(s)

McClain; Ethan S. et al.

BREATH TEST METHOD AND TEST MEAL

Abstract

A breath test method is provided that includes the steps of (a) obtaining a pre-meal baseline breath sample from a test subject, (b) providing a test meal comprising hypoallergenic formula and .sup.13C-Spirulina, the .sup.13C-Spirulina being bound to components of the hypoallergenic formula, (c) allowing the test subject to consume the test meal, (d) obtaining breath samples from the test subject at a series of time points after consumption of the test meal, and (e) analyzing the breath samples.

Inventors: McClain; Ethan S. (Franklin, TN), Giffin; Nicole L. (Gallatin, TN), Williams; Catherine E. (Fairview, TN), Bush; Kerry C. (Franklin, TN)

Applicant: Advanced Breath Diagnostics, LLC (Brentwood, TN)

Family ID: 96881663

Appl. No.: 19/065424

Filed: February 27, 2025

Related U.S. Application Data

us-provisional-application US 63559645 20240229

Publication Classification

Int. Cl.: A61K49/00 (20060101); G01N33/497 (20060101)

U.S. Cl.:

CPC A61K49/0004 (20130101); G01N33/497 (20130101);

Background/Summary

RELATED APPLICATION [0001] The present application claims priority to U.S. provisional application No. 63/559,645, the entire contents of which is incorporated by reference.

FIELD OF THE INVENTION

[0002] This invention relates generally to a test meal including an edible food component that includes a marker. The invention also relates to gastric emptying breath tests for evaluating gastric emptying rates in test subjects.

BACKGROUND OF THE INVENTION

[0003] Gastroparesis (or delayed gastric emptying) is a condition that affects motility in the stomach and causes food to stay in the stomach for too long. Normally, the stomach muscles contract to propel food through the digestive tract, moving food from the stomach to the small intestine. However, in a subject with gastroparesis, the stomach muscles do not contract normally, thereby causing a delay in the rate of gastric emptying. Gastroparesis can be diagnosed by evaluating a gastric emptying rate in a subject. Currently, there are two types of tests for measuring gastric emptying rate: gastric scintigraphy tests and breath tests.

[0004] In a scintigraphy test, a subject ingests a test meal including a component that has been radiolabeled with a radioisotope. The gamma emission from the radiolabel is measured by a scintillation camera it moves from the stomach to the small intestine. Several disadvantages are associated with the scintigraphy test. First, subjects must be subjected to radioisotopes. This is particularly problematic for women of childbearing age or children. Further, the test must be carried out at specialized nuclear medicine facilities. Scintigraphy tests cannot be administered in a regular clinical setting or at home through virtual telehealth due to radiation risks and requirements for specially licensed personnel and facilities to handle radioactive materials.

[0005] In a breath test, a subject ingests a test meal including a component that has been labeled with a non-radioactive stable isotope. While a number of breath tests have been developed, there is currently only one standardized, FDA-approved gastric emptying breath test (“GEBT method”) for evaluating a gastric emptying rate in a subject. The GEBT method is disclosed in PMA No. P110015 and approved by the FDA on Apr. 6, 2015. This standardized GEBT method uses an egg-based test meal that contains reconstituted lyophilized whole eggs having components bound to ^{13}C -Spirulina (“egg-based test meal”). The whole eggs are part of a formula containing nonfat dry milk. The egg-based test meal also includes Nabisco PREMIUM saltine crackers.

[0006] One drawback with the egg-based test meal is that it includes whole eggs. Some subjects may have an allergy to whole eggs and therefore be ineligible to use this test. Further, the incidence of allergies to whole eggs is higher in infants and children and it is often unknown whether the infant or child has an allergy or intolerance to whole eggs, dairy and/or gluten. If such an unknown allergy or intolerance exists, it may have allergenic risks and/or impact the test results.

[0007] It would therefore be desirable to provide an alternative breath test meal to the egg-based test meal that does not include eggs, dairy or gluten. It would also be desirable to be able to use such an alternative breath test meal in the standardized, FDA-approved GEBT method. It would also be desirable to provide an option to elect between different test meals for use in the GEBT method. Further, it would be desirable to provide a meal capable for use in an infant.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

[0008] FIG. 1 is a graph showing $^{13}\text{CO}_2$ excretion rates at a series of time points for healthy subjects administered a formula-based meal.

[0009] FIG. 2 is a graph showing .sup.13CO.sub.2 excretion rates at a series of time points for healthy subjects administered a standardized egg-based test meal.

[0010] FIG. 3 is a graph showing average .sup.13CO.sub.2 excretion rates at a series of time points for healthy subjects administered a formula-based meal and a standardized egg-based test meal.

SUMMARY

[0011] Certain embodiments provide a gastric emptying breath test meal. In some cases, the breath test meal includes hypoallergenic formula, water and .sup.13C-Spirulina, and the .sup.13C-Spirulina is bound to components of the hypoallergenic formula. In some cases, the hypoallergenic formula has no protein sources other than 100% free amino acids. One exemplary hypoallergenic formula is the EleCareJr amino acid-based nutrition powder.

[0012] The mixture can include hypoallergenic formula in an amount from 20 to 80 grams, such as from 30 to 70 grams, from 40 to 60 grams, from 45 to 55 grams, from 50 grams to 52 grams (e.g., 51 grams), water in an amount from 95 to 375 mL, from 140 mL to 340 mL, from 200 mL to 300 mL, from 225 mL to 275 mL, from 230 mL to 250 mL, from 235 mL to 245 mL, from 239 mL to 241 mL (e.g., 240 mL), and .sup.13C-Spirulina in an amount from 50 to 200 mg, such as from 75 mg to 125 mg, from 95 mg to 105 mg or from 99 mg to 101 mg (e.g., 100 mg). The formula-based meal can also include an additional serving of water in an amount from 120 to 240 mL, such as from 160 mL to 200 mL, from 170 mL to 190 mL, from 175 mL to 185 mL, or from 179 mL to 181 mL (e.g., 180 mL). The additional serving of water is consumed along with the formula-based meal but is not mixed with it.

[0013] In some cases, the breath test meal includes hypoallergenic formula in an amount from 20 grams to 80 grams, water in an amount from 95 mL to 375 mL and .sup.13C-Spirulina in an amount from 50 mg to 200 mg. Each component can be present in any range described herein. In some cases, the hypoallergenic formula is present in an amount from 45 grams to 55 grams, water is present in an amount from 235 mL to 245 mL and .sup.13C-Spirulina is present in an amount from 95 mg to 105 mg. In certain cases, an additional serving of water is present in an amount from 120 to 240 mL, such as from 175 mL to 185 mL.

[0014] Other embodiments provide a breath test method. The method can include the steps of (a) obtaining a pre-meal baseline breath sample from a test subject, (b) providing a breath test meal comprising hypoallergenic formula and .sup.13C-Spirulina, the .sup.13C-Spirulina being bound to components of the hypoallergenic formula, (c) allowing the test subject to consume the breath test meal, (d) obtaining breath samples from the test subject at a series of time points after consumption of the breath test meal, and (e) analyzing the breath samples. The breath test meal can include any breath test meal described herein.

[0015] In some cases, the series of time points includes 45, 90, 120, 150, 180 and 240 minute time points. In certain cases, the series of time points can also include a 15 minute time point, a 30 minute time point, a 45 minute time point, a 60 minute time point and/or a 210 minute time point. In specific cases, the series of time points can include 15, 30, 45, 60, 90, 120, 150, 180, 210 and 240 minute time points.

[0016] Also, in some cases, the step of obtaining a pre-meal baseline breath sample takes place after the test subject fasts for at least 8 hours. Further, in some cases, the step of allowing the test subject to consume the breath test meal includes prompting the test subject to consume the breath test meal within 10 minutes. Even further, in some cases, the step of collecting the breath test samples from the test subject includes depositing breath samples into collection tubes and then capping the collection tubes. Finally, in some cases, the step of analyzing the breath samples can include using Gas Isotope Ratio Mass Spectrometry to determine the ratio of .sup.13CO.sub.2/.sup.12CO.sub.2 in each breath sample.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[0017] The following detailed description is to be read with reference to the drawings, in which like elements in different drawings have like reference numerals. The drawings, which are not

necessarily to scale, depict selected embodiments and are not intended to limit the scope of the invention. Skilled artisans will recognize that the examples provided herein have many useful alternatives that fall within the scope of the invention.

[0018] Applicant has developed an alternative test meal that can be used in the standardized, FDA-approved gastric emptying breath test (“GEBT method”) disclosed in PMA No. P110015 and approved by the FDA on Apr. 6, 2015, the entire contents of which are incorporated herein by reference.

[0019] The GEBT method currently uses an egg-based test meal consisting of 27 grams of re-hydrated, pasteurized scrambled egg mix containing a dose of 43 mg of .sup.13C (provided by approximately 100 mg of .sup.13C-Spirulina), 6 saltine crackers, and 180 mL of water. The pasteurized scrambled egg mix includes whole eggs, water, nonfat dry milk, salt, and smoke flavoring. The .sup.13C-Spirulina is cultured in a >95% .sup.13C enriched medium, resulting in full .sup.13C-labeling of nearly all of the algae's carbon containing nutrients.

[0020] Applicant has discovered a new test meal that can be used in the standardized GEBT method as an alternative to the egg-based test meal. The new test meal comprises hypoallergenic infant formula and .sup.13C-Spirulina, wherein the .sup.13C-Spirulina is bound to components of the infant formula (“formula-based test meal”). In some cases, the hypoallergenic formula has no protein sources other than 100% free amino acids. The .sup.13C-Spirulina in the formula-based test meal can also be cultured in a >95% .sup.13C enriched medium, resulting in full .sup.13C-labeling of nearly all of the algae's carbon containing nutrients.

[0021] When the formula-based test meal is administered according to the standardized GEBT method, it surprisingly provides results comparable to results obtained by administering the egg-based test meal. This was an unexpected discovery because the formula-based test meal includes a completely different matrix than the egg-based test meal. It was previously believed that whole eggs in the egg-based test meal provided a unique protein/fat matrix that allowed the .sup.13C-Spirulina to remain bound to whole egg components while travelling through the subject's digestive tract. However, Applicant surprisingly discovered that .sup.13C-Spirulina also remains bound to the matrix of the formula-based test meal while travelling through a digestive tract.

[0022] The formula-based test meal is beneficial because it is hypoallergenic and does not include eggs, dairy or gluten. Also, the formula-based meal has a liquid-like consistency and therefore can be administered to infants in a bottle. The formula-based test meal is also advantageous because it can be used in the standardized GEBT method. This provides quite an advantage because clinicians do not need to use an entirely different breath test method to use the formula-based test meal to assess gastric emptying in infants.

[0023] Some embodiments provide a gastric emptying breath test meal comprising a mixture including hypoallergenic formula, water and .sup.13C-Spirulina, wherein the .sup.13C-Spirulina is bound to components of the hypoallergenic formula. In some cases, the hypoallergenic formula has no protein sources other than 100% free amino acids. One exemplary hypoallergenic formula is the EleCare Jr amino acid-based nutrition powder. The mixture can include hypoallergenic formula in an amount from 20 to 80 grams, such as from 30 to 70 grams, from 40 to 60 grams, from 45 to 55 grams, from 50 grams to 52 grams (e.g., 51 grams), water in an amount from 95 to 375 mL, from 140 mL to 340 mL, from 200 mL to 300 mL, from 225 mL to 275 mL, from 230 mL to 250 mL, from 235 mL to 245 mL, from 239 mL to 241 mL (e.g., 240 mL), and .sup.13C-Spirulina in an amount from 50 to 200 mg, such as from 75 mg to 125 mg, from 95 mg to 105 mg or from 99 mg to 101 mg (e.g., 100 mg). The formula-based meal can also include an additional serving of water in an amount from 120 to 240 mL, such as from 160 mL to 200 mL, from 170 mL to 190 mL, from 175 mL to 185 mL, or from 179 mL to 181 mL (e.g., 180 mL). The additional serving of water is consumed along with the formula-based meal but is not mixed with it.

[0024] Other embodiments provide a breath test method using a formula-based test meal. The formula-based test meal can include any embodiment described herein. The method can include

steps of (1) collecting a pre-meal fasting breath sample from subject, (2) preparing a formula-based test meal, (3) allowing a subject to consume the formula-based test meal, and (4) collecting breath samples at a series of time points after the subject consumes the formula-based test meal, (5) and analyzing the breath samples at each time point. The method can be performed by a test administrator such as a clinician in a clinical setting. In some cases, the method can be performed virtually via telehealth. In some cases, the series of time points include any of 15, 30, 45, 60, 90, 120, 150, 180, 210 and 240 minute time points. In certain cases, the series of time points include the 45, 90, 120, 150, 180 and 240 minute time points. In specific cases, the series of time points include all 15, 30, 45, 60, 90, 120, 150, 180, 210 and 240 minute time points.

[0025] One exemplary breath test method using the formula-based test meal will now be described.

[0026] 1. Prompting a test subject to fast overnight for at least 8 hours. No solid foods or liquids other than 120 mL of water up to 1 hour before the test is allowed. [0027] 2. Collecting a pre-meal, baseline breath sample. Any method of collecting breath samples can be used but in preferred cases, the breath samples are deposited into collection tubes and then the collection tubes are capped. [0028] 3. Preparing the formula-based test meal. [0029] 4. Prompting the test subject to ingest the formula-based test meal along with an additional serving of water. The test subject is also prompted to consume the test meal and water within 10 minutes. [0030] 5. Collecting a breath sample at 15, 30, 45, 60, 90, 120, 150, 180, 210 and 240 minute time points after the test subject consumes the formula-based test meal. [0031] 6. Using a Gas Isotope Ratio Mass Spectrometry to determine the ratio of $\text{sup.13CO.sub.2}/\text{sup.12CO.sub.2}$ in each breath sample. The ratio is then used to calculate the sup.13CO.sub.2 excretion rate for each breath sample. The rate of sup.13CO.sub.2 excretion at each GEBT measurement time is directly proportional to rate of gastric emptying. Preferably, the excretion rate for each breath sample is reported using a kPCD metric. The acronym kPCD stands for “1000 X Percent Carbon-13 Dose (PCD) excreted per minute.” A kPCD value at each time point (kPCD.sub.t) is calculated at each GEBT measurement time using the following formula:

[00001]
$$\text{kPCD}_t = \left[\frac{\text{DOB} \times \text{CO}_2\text{PR} \times R_s \times 13}{10 \times \text{dose}} \right] \times 1000$$
 [0032] Where: [0033] DOB=The measured difference in the ratio $[\text{sup.13CO.sub.2}/\text{sup.12CO.sub.2}]$ between a breath sample at a time point and the baseline breath sample. [0034] CO.sub.2PR = CO.sub.2 production rate (mmol $\text{CO.sub.2}/\text{min}$) calculated using Schofield equations, which incorporate the subject's age, sex, height, and weight. [0035] $R_{\text{sub.s}}$ =The ratio $[\text{sup.13CO.sub.2}/\text{sup.12CO.sub.2}]$ in a reference standard (Pee Dee belemnite) for these measurements. $R_{\text{sub.s}}=0.0112372$ [0036] 13=the atomic weight of Carbon-13 [0037] 10=A constant factor for converting units [0038] dose=the weight (mg) of Carbon-13 in the dose of sup.13C -Spirulina administered to the test subject. Since sup.13C -Spirulina is approximately 43% Carbon-13, a 100 mg dose of sup.13C -Spirulina contains approximately 43 mg of Carbon-13. [0039] 7. The kPCD values for each time point (pre-meal, 15, 30, 45, 60, 90, 120, 150, 180, 210 and 240 time points) are provided. Preferably, the kPCD values are provided in a graphical display showing the subject's kPCD at each time point. The graphical display can also include a reference range cutoff point at each time point. The cutoff point demarcates normal from delayed gastric emptying. [0040] 8. The test subject's kPCD values are compared to the cutoff points of the reference range to qualitatively classify the subject as having normal or delayed gastric emptying. Generally, subjects classified as having normal gastric emptying typically display kPCD values that exceed time-specific cutoff points, reach a maximum kPCD value from 120-180 minutes, and then decline. In contrast, subjects classified as having delayed gastric emptying typically display kPCD values that are lower and rise continuously throughout the four-hour evaluation period. As a result, their highest kPCD value often occurs at four hours.

Note that this exemplary test method can be performed in any desired setting, such as a clinical setting or a telehealth setting. In some cases, a clinician prepares the test meal whereas in other cases, the test subject prepares the test meal. Also, in some cases, the breath samples are analyzed in an outside lab by a lab technician. This exemplary method therefore is not limited to a particular

setting and various steps can be performed in a single setting or across multiple settings.

EXAMPLE 1

[0041] The GEBT method was performed on seven healthy subjects using the formula-based test meal. In this experiment, the formula-based test meal included a mixture including 51 grams of EleCare, 240 mL of water and 100 mg of .sup.13C-Spirulina. The subjects were also given 180 mL of water along with the mixture. The results are shown in Table 2 and plotted on a curve shown in FIG. 1.

[illegible]

EXAMPLE 2

[0042] The GEBT method also was performed on six of the same healthy subjects in Example 1, using the egg-based test meal. In this experiment, the egg-based test meal included 27 grams of re-hydrated, pasteurized scrambled egg mix containing a dose of 43 mg of .sup.13C (provided by approximately 100 mg of .sup.13C-Spirulina), 6 saltine crackers, and 180 mL of water. The results are shown in Table 2 and plotted on a curve shown in FIG. 2.

[illegible]

[0043] FIG. 3 illustrates a curve showing average kPCD values for the six healthy subjects that underwent the GEBT method for both the formula-based meal and the egg-based meal. The results shown in Examples 2-3 show that the kPCD values surprisingly reflected similar rates and patterns of gastric emptying in subjects across both meals.

[0044] While some preferred embodiments of the invention have been described, it should be understood that various changes, adaptations and modifications may be made therein without departing from the spirit of the invention and the scope of the appended claims.

Claims

1. A breath test method, comprising the steps of: obtaining a pre-meal baseline breath sample from a test subject; providing a test meal comprising a mixture including hypoallergenic formula, water and .sup.13C-Spirulina, wherein the .sup.13C-Spirulina is bound to components of the hypoallergenic formula; allowing the test subject to consume the test meal; obtaining breath samples from the test subject at a series of time points after consumption of the test meal; and analyzing the breath samples.
2. The breath test method of claim 1 wherein the series of time points includes 45, 90, 120, 150, 180 and 240 minute time points.
3. The breath test method of claim 1 wherein the .sup.13C-Spirulina is present in an amount from 50 mg to 200 mg.
4. The breath test method of claim 3 wherein the .sup.13C-Spirulina is present in an amount from 95 mg to 105 mg.
5. The breath test method of claim 1 wherein the hypoallergenic formula is present in an amount from 20 grams to 80 grams.

- 6.** The breath test method of claim 5 wherein the hypoallergenic formula is present in an amount from 45 grams to 55 grams.
 - 7.** The breath test method of claim 1 wherein the water is present in an amount from 95 mL to 275 mL.
 - 8.** The breath test method of claim 7 wherein the water is present in an amount from 235 mL to 245 mL.
 - 9.** The breath test method of claim 1 further comprising an additional serving of water that is not part of the mixture.
 - 10.** The breath test method of claim 9 wherein the additional serving of water is present in an amount from 120 ml to 240 mL.
 - 11.** The breath test method of claim 10 wherein the additional serving of water is present in an amount from 175 ml to 185 mL.
 - 12.** The breath test method of claim 1 wherein the mixture comprises hypoallergenic formula in an amount from 20 grams to 80 grams, water in an amount from 95 mL to 375 mL and .sup.13C-Spirulina in an amount from 50 mg to 200 mg.
 - 13.** The breath test method of claim 12 wherein the mixture comprises hypoallergenic formula in an amount from 45 grams to 55 grams, water in an amount from 235 mL to 245 mL and .sup.13C-Spirulina in an amount from 95 mg to 105 mg.
 - 14.** The breath test method of claim 13 further comprising water.
 - 15.** The breath test method of claim 14 wherein the water is present in an amount from 175 mL to 185 mL.
 - 16.** The breath test method of claim 1 wherein the hypoallergenic formula has no protein sources other than 100% free amino acids
-