



US012385020B1

(12) **United States Patent**
Lee et al.

(10) **Patent No.:** **US 12,385,020 B1**

(45) **Date of Patent:** **Aug. 12, 2025**

(54) **5-DEHYDRO-2-DEOXYGLUCONOKINASE
VARIANT AND METHOD FOR PRODUCING
5'-INOSINIC ACID USING SAME**

(58) **Field of Classification Search**

CPC C12N 9/1205

See application file for complete search history.

(71) Applicant: **DAESANG CORPORATION**, Seoul
(KR)

(56) **References Cited**

(72) Inventors: **Sun Hee Lee**, Seoul (KR); **Tae Yeol
Choi**, Seoul (KR); **Hyun Ho Kim**,
Seoul (KR); **Dong Hyun Kim**, Seoul
(KR); **Hyun Sook Kim**, Seoul (KR);
Jong Hwan Shin, Seoul (KR)

FOREIGN PATENT DOCUMENTS

KR	10-1166027	7/2012
KR	10-2016-0078694	7/2016
KR	10-1904675	10/2018
KR	10-1916611	11/2018
KR	10-1916622	11/2018

(73) Assignee: **DAESANG CORPORATION**, Seoul
(KR)

OTHER PUBLICATIONS

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

International Search Report issued Jan. 19, 2024 in International
(PCT) Application No. PCT/KR2023/012969.
Zuniga-Soto, Evelyn et al., "Insights into the transcriptomic response
of the plant engineering bacterium *Ensifer adhaerens* OV14 during
transformation", Scientific Reports, 2019, vol. 9, No. 10344, pp.
1-17.

(21) Appl. No.: **18/879,995**

Primary Examiner — Suzanne M Noakes

(22) PCT Filed: **Aug. 31, 2023**

(74) *Attorney, Agent, or Firm* — Wenderoth, Lind &
Ponack, L.L.P.

(86) PCT No.: **PCT/KR2023/012969**

§ 371 (c)(1),

(2) Date: **Dec. 30, 2024**

(87) PCT Pub. No.: **WO2024/210281**

PCT Pub. Date: **Oct. 10, 2024**

(57) **ABSTRACT**

The present invention relates to a novel 5-dehydro-2-deoxy-
gluconokinase variant and a method of producing 5'-inosinic
acid using the same. The 5-dehydro-2-deoxygluconokinase
variant is obtained by substituting one or more amino acids
in the amino acid sequence constituting 5-dehydro-2-deoxy-
gluconokinase to change the activity of the protein, and a
recombinant microorganism comprising the 5-dehydro-2-
deoxygluconokinase variant is capable of efficiently produc-
ing 5'-inosinic acid.

(30) **Foreign Application Priority Data**

Apr. 6, 2023 (KR) 10-2023-0045161

(51) **Int. Cl.**

C12N 9/12 (2006.01)


C12P 19/32 (2006.01)

(52) **U.S. Cl.**

CPC **C12N 9/1205** (2013.01); **C12P 19/32**
(2013.01); **C12Y 207/01092** (2013.01)

7 Claims, 1 Drawing Sheet

Specification includes a Sequence Listing.



1

5-DEHYDRO-2-DEOXYGLUCONOKINASE VARIANT AND METHOD FOR PRODUCING 5'-INOSINIC ACID USING SAME

SEQUENCE LISTING

A sequence listing in electronic (XML file) format is filed with this application and incorporated herein by reference. The name of the XML file is "Sequence_Listing-1948A.xml"; the file was created on Dec. 29, 2024; the size of the file is 17.022 bytes.

TECHNICAL FIELD

The present invention relates to a novel 5-dehydro-2-deoxygluconokinase variant and a method of producing 5'-inosinic acid using the same.

BACKGROUND ART

5'-inosinic acid (or inosine monophosphate (IMP)) is an intermediate in the metabolic system of nucleic acid biosynthesis, and not only plays an important physiological role in the bodies of plants and animals, but is also used in various applications, including food, medicine, and various medical applications. In particular, 5'-inosinic acid is a nucleic acid-based seasoning, which has drawn much attention as a savory seasoning, because it has significant synergistic effects on taste when used together with monosodium glutamate (MSG).

Methods for producing 5'-inosinic acid include a method of enzymatically degrading ribonucleic acid extracted from yeast cells, a method of chemically phosphorylating inosine produced by fermentation, etc. Recently, a method of culturing a 5'-inosinic acid-producing microorganism and recovering 5'-inosinic acid accumulated in the medium has been mainly used.

For the production of 5'-inosinic acid using microorganisms, in order to improve the efficiency of production of 5'-inosinic acid, there has been development of a variety of recombinant strains or mutant strains having excellent 5'-inosinic acid productivity by applying genetic recombination technology to microorganisms such as *Escherichia coli* and *Corynebacterium*, which are widely used in the production of useful substances such as nucleic acids or L-amino acids, and methods of producing 5'-inosinic acid using the same. In particular, there have been attempts to increase the production of 5'-inosinic acid by targeting genes such as enzymes, transcription factors and transport proteins, which are involved in the biosynthetic pathway of 5'-inosinic acid, or by inducing mutations in promoters that regulate the expression of these genes. However, there are dozens to hundreds of types of proteins such as enzymes, transcription factors and transport proteins, which are involved directly or indirectly in the production of 5'-inosinic acid, and thus much research is still needed on the increase in 5'-inosinic acid productivity by changes in the activity of these proteins.

PRIOR ART DOCUMENTS

Patent Documents

Korean Patent No. 10-116602

DISCLOSURE

Technical Problem

An object of present invention is to provide a novel 5-dehydro-2-deoxygluconokinase variant.

2

Another object of the present invention is to provide a polynucleotide encoding the variant.

Still another object of the present invention is to provide a transformant comprising the variant or polynucleotide.

Yet another object of the present invention is to provide a method of producing 5'-inosinic acid using the transformant.

Technical Solution

One aspect of the present invention provides a 5-dehydro-2-deoxygluconokinase variant consisting of the amino acid sequence of SEQ ID NO: 2 in which serine at position 146 in the amino acid sequence of SEQ ID NO: 4 is substituted with asparagine.

As used in the present invention, the term "5-dehydro-2-deoxygluconokinase catalyzes a reaction that produces ADP and 6-phospho-5-dehydro-2-deoxy-D-gluconate using ATP and 5-dehydro-2-deoxy-D-gluconate as substrates, and it may be a polypeptide or protein consisting of the amino acid sequence of SEQ ID NO: 4 and having 5-dehydro-2-deoxygluconokinase activity.

Information on the nucleic acid and protein sequences of the 5-dehydro-2-deoxygluconokinase is available from known sequence databases (e.g., GenBank, UniProt).

According to one embodiment of the present invention, the 5-dehydro-2-deoxygluconokinase may be encoded by the nucleotide sequence of SEQ ID NO: 3.

The amino acid sequence of the 5-dehydro-2-deoxygluconokinase according to the present invention or the nucleotide sequence encoding the same may include a nucleotide sequence or amino acid sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99 or 100% homology or identity to the amino acid sequence of SEQ ID NO: 4 or the nucleotide sequence of SEQ ID NO: 3. As used herein, the term "homology" or "identity" means the percentage rate of identity between two sequences, which is determined by aligning the reference nucleotide sequence or amino acid sequence and any other nucleotide sequence or amino acid sequence to correspond to each other as much as possible and analyzing the aligned sequences.

According to one embodiment of the present invention, the 5-dehydro-2-deoxygluconokinase may be derived from wild-type *Corynebacterium stationis*.

As used in the present invention, the term "variant" refers to a protein that has an amino acid sequence different from the amino acid sequence before mutation by the conservative substitution and/or modification of one or more amino acids at the N-terminus, C-terminus of and/or within the amino acid sequence, which result(s) from mutation in the nucleotide sequence of the gene encoding the protein, but retains the functions or properties of the protein before mutation. As used herein, the term "conservative substitution" means substituting one amino acid with another amino acid having similar structural and/or chemical properties.

The conservative substitution may have little or no impact on the activity of the protein or polypeptide. In addition, the term "modification" refers to substitution, insertion, deletion, or the like of one or more amino acids. The amino acid is selected from among alanine (Ala, A), isoleucine (Ile, I), valine (Val, V), leucine (Leu, L), methionine (Met, M), asparagine (Asn, N), cysteine (Cys, C), glutamine (Gln, Q), serine (Ser, S), threonine (Thr, T), phenylalanine (Phe, F), tryptophan (Trp, W), tyrosine (Tyr, Y), aspartic acid (Asp, D), glutamic acid (Glu, E), arginine (Arg, R), histidine (His, H), lysine (Lys, K), glycine (Gly, G), and proline (Pro, P).

In addition, some variants include those in which one or more portions, such as an N-terminal leader sequence or

transmembrane domain, have been removed, or those in which a portion has been removed from the N- and/or C-terminus of a mature protein.

The variant may have increased (enhanced), unchanged, or decreased (weakened) ability compared to that of the protein before mutation. Here, the term “increased or enhanced” includes: a case in which the activity of the protein itself has increased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is higher than that in the wild-type strain or the strain expressing the protein before mutation due to increased expression or translation of the gene encoding the protein; and a combination thereof. In addition, the term “decreased or weakened” includes: a case in which the activity of the protein itself has decreased compared to the activity of the protein before mutation; a case in which the overall activity of the protein in the cell is lower than that in the wild-type strain or the strain expressing the protein before mutation due to reduced expression or translation of the gene encoding the protein; and a combination thereof. In the present invention, the term “variant” may be used interchangeably with terms such as variant type, modification, variant polypeptide, mutated protein, mutation, and the like.

The 5-dehydro-2-deoxygluconokinase variant according to the present invention may comprise an amino acid sequence having at least 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98s, 99% or 100% homology or identity to the amino acid sequence of SEQ ID NO: 2.

Another aspect of the present invention provides a polynucleotide encoding the 5-dehydro-2-deoxygluconokinase variant.

As used in the present invention, the term “polynucleotide” refers to a DNA or RNA strand having a certain length or more, which is a long-chain polymer of nucleotides formed by linking nucleotide monomers via covalent bonds. More specifically, the term “polynucleotide” refers to a polynucleotide fragment encoding the variant.

According to one embodiment, the polynucleotide may comprise a nucleotide sequence encoding the amino acid sequence of SEQ ID NO: 2.

More specifically, the polynucleotide may comprise the nucleotide sequence of SEQ ID NO: 1 in which nucleotide “g” at position 437 in the nucleotide sequence of SEQ ID NO: 3 encoding 5-dehydro-2-deoxygluconokinase is substituted with nucleotide “a”.

Still another aspect of the present invention provides a vector comprising a polynucleotide encoding the 5-dehydro-2-deoxygluconokinase variant.

Yet another aspect of the present invention provides a transformant comprising the 5-dehydro-2-deoxygluconokinase variant or the polynucleotide.

As used in the present invention, the term “vector” refers to any type of nucleic acid sequence transfer structure that is used as a means for transferring and expressing a gene of interest in a host cell. Unless otherwise specified, the term “vector” may mean one allowing the nucleic acid sequence contained therein to be expressed after insertion into the host cell genome and/or one allowing the nucleic acid sequence to be expressed independently. This vector comprises essential regulatory elements operably linked so that an inserted gene can be expressed. As used herein, the term “operably linked” means that a gene of interest and regulatory sequences thereof are functionally linked together in a manner enabling gene expression, and the “regulatory elements” include a promoter for initiating transcription, any operator sequence for regulating transcription, a sequence

encoding suitable mRNA ribosome-binding sites, and a sequence for regulating termination of transcription and translation.

The vector in the present invention is not particularly limited as long as it may replicate in a host cell, and any vector known in the art may be used. Examples of the vector include a natural or recombinant plasmid, cosmid, virus and bacteriophage. Examples of a phage vector or cosmid vector include, but are not limited to, pWE15, M13, λMBL3, λMBL4, λIXII, λASHII, λAPII, λt10, λt11, Charon4A, and Charon21A, and examples of a plasmid vector include, but are not limited to, pBR series, pUC series, pBluescriptII series, pGEM series, pTZ series, pCL series, and pET series.

The vector may typically be constructed as a vector for cloning or as a vector for expression. The vector for expression may be a conventional vector that is used in the art to express a foreign gene or protein in a plant, animal, or microorganism, and may be constructed through various methods known in the art.

As used in the present invention, the term “recombinant vector” may be transformed into a suitable host cell, and then may replicate regardless of the genome of the host cell or may be integrated into the genome itself. In this case, the “suitable host cell” may contain a replication origin, which is a particular nucleotide sequence which enables the vector to replicate in the suitable host cell and from which replication starts. For example, when the vector used is an expression vector and uses a prokaryotic cell as a host, the vector generally comprises a strong promoter capable of promoting transcription (e.g., pLλ promoter, CMV promoter, trp promoter, lac promoter, tac promoter, T7 promoter, etc.), a ribosome binding site for initiation of translation, and a transcription/translation termination sequence. When a eukaryotic cell is used as a host, the vector comprises a replication origin operating in the eukaryotic cell, and examples of the replication origin include, but are not limited to, an f1 replication origin, an SV40 replication origin, a pMB1 replication origin, an adeno replication origin, an AAV replication origin, and a BBV replication origin. In addition, the recombinant vector may comprise a promoter derived from the genome of a mammalian cell (e.g., metallothionein promoter) or a promoter derived from a mammalian virus (e.g., adenovirus late promoter, vaccinia virus 7.5K promoter, SV40 promoter, cytomegalovirus promoter, HSV-tk promoter, etc.), and generally has a polyadenylation sequence as a transcription termination sequence.

The recombinant vector may comprise a selection marker. The selection marker serves to select a transformant (host cell) transformed with the vector, and since only cells expressing the selection marker can survive in the medium treated with the selection marker, it is possible to select transformed cells. Representative examples of the selection marker include, but are not limited to, kanamycin, streptomycin, and chloramphenicol.

The transformant may be produced by inserting the recombinant vector into a host cell, and the transformant may be obtained by introducing the recombinant vector into an appropriate host cell. The host cell is a cell capable of stably and continuously cloning or expressing the expression vector, and any host cell known in the art may be used.

Where the vector is transformed into prokaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, *E. coli* sp. strains such as *E. coli* DH5α, *E. coli* JM109, *E. coli* BL21, *E. coli* RR1, *E. coli* LE392, *E. coli* B, *E. coli* X 1776, *E. coli* W3110, and *E. coli* XL1-Blue, *Bacillus* sp. strains such as *Bacillus subtilis* and *Bacillus thuringiensis*, *Coryne-*

bacterium sp. strains such as *Corynebacterium glutamicum* and *Corynebacterium stationis*, and various Enterobacteriaceae strains such as *Salmonella typhimurium*, *Serratia marcescens*, and *Pseudomonas* species.

Where the vector is transformed into eukaryotic cells to generate recombinant microorganisms, examples of host cells that may be used include, but are not limited to, yeast (e.g., *Saccharomyces cerevisiae*), insect cells, plant cells and animal cells, such as Sp2/0, CHO K1, CHO DG44, PER.C6, W138, BHK, COS7, 293, HepG2, Huh7, 3T3, RIN, and MDCK cell lines.

As used in the present invention, the term “transformation” refers to a phenomenon in which external DNA is introduced into a host cell, thereby artificially causing genetic changes, and the term “transformant” refers to a host cell into which external DNA has been introduced and in which the expression of the gene of interest is stably maintained.

The transformation may be performed using a suitable vector introduction technique selected depending on the host cell, so that the gene of interest or a recombinant vector comprising the same may be expressed in the host cell. For example, introduction of the vector may be performed by electroporation, heat-shock, calcium phosphate (CaPO₄) precipitation, calcium chloride (CaCl₂) precipitation, micro-injection, polyethylene glycol (PEG) method, DEAE-dextran method, cationic liposome method, lithium acetate-DMSO method, or any combination thereof, without being limited thereto. As long as the transformed gene may be expressed in the host cell, it may be inserted into the chromosome of the host cell, or may exist extrachromosomally, without being limited thereto.

The transformant may include a cell transfected, transformed, or infected with the recombinant vector of the present invention in vivo or in vitro, and may be used in the same sense as a recombinant host cell, a recombinant cell, or a recombinant microorganism.

Genes inserted into the recombinant vector of the present invention may be introduced into a host cell such as a *Corynebacterium* sp. strain by homologous recombination crossover.

According to one embodiment of the present invention, the transformant may be a *Corynebacterium* sp. microorganism.

The *Corynebacterium* sp. microorganism may be, but is not limited to, *Corynebacterium glutamicum*, *Corynebacterium crudilactis*, *Corynebacterium deserti*, *Corynebacterium callunae*, *Corynebacterium suranareae*, *Corynebacterium lubricantis*, *Corynebacterium doosanense*, *Corynebacterium efficiens*, *Corynebacterium uterequi*, *Corynebacterium stationis*, *Corynebacterium pacaense*, *Corynebacterium singulare*, *Corynebacterium humireducens*, *Corynebacterium marinum*, *Corynebacterium halotolerans*, *Corynebacterium spheniscorum*, *Corynebacterium freiburgense*, *Corynebacterium striatum*, *Corynebacterium canis*, *Corynebacterium ammoniagenes*, *Corynebacterium renale*, *Corynebacterium pollutisoli*, *Corynebacterium imitans*, *Corynebacterium caspium*, *Corynebacterium testudinatoris*, *Corynebacterium pseudopelargi*, or *Corynebacterium flavescens*.

The transformant in the present invention may be a strain either comprising the above-described 5-dehydro-2-deoxygluconokinase variant or a polynucleotide encoding the same or comprising the vector comprising the same, a strain expressing the 5-dehydro-2-deoxygluconokinase variant or

the polynucleotide, or a strain having activity for the 5-dehydro-2-deoxygluconokinase variant, without being limited thereto.

The transformant of the present invention may comprise other protein variants or genetic mutations, in addition to the 5-dehydro-2-deoxygluconokinase variant.

According to one embodiment of the present invention, the transformant may have the ability to produce inosinic acid.

The 5'-inosinic acid is a nucleic acid-based compound that gives flavor to food, especially umami (savory) taste, and is used with the same meaning as inosine monophosphate (IMP).

The transformant may naturally have the ability to produce 5'-inosinic acid or may be one artificially endowed with the ability to produce 5'-inosinic acid.

According to one embodiment of the present invention, the transformant may have an increased ability to produce 5'-inosinic acid, due to a change in 5-dehydro-2-deoxygluconokinase activity.

As used in the present invention, the term “increased ability to produce” means that 5'-inosinic acid productivity has increased compared to that of the parent strain. As used herein, the term “parent strain” refers to a wild-type strain or mutant strain to be mutated, and includes a strain that is to be mutated directly or to be transformed with a recombinant vector or the like. In the present invention, the parent strain may be a wild-type *Corynebacterium* sp. strain or a *Corynebacterium* sp. strain mutated from the wild-type strain.

The transformant according to the present invention exhibits an increased ability to produce 5'-inosinic acid compared to the parent strain, due to the change in 5-dehydro-2-deoxygluconokinase activity caused by introduction of the 5-dehydro-2-deoxygluconokinase variant thereto. More specifically, the amount of 5'-inosinic acid produced by the transformant may be at least 1%, 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100% higher than that produced by the parent strain, or may be 1.1-fold, 1.5-fold, 2-fold, 2.5-fold, 3-fold, 3.5-fold, 4-fold, 4.5-fold, 5-fold, 5.5-fold, 6-fold, 6.5-fold, 7-fold, 7.5-fold, 8-fold, 8.5-fold, 9-fold, 9.5-fold, or 10-fold higher than that produced by the parent strain, without being limited thereto. For example, the amount of 5'-inosinic acid produced by the transformant comprising the 5-dehydro-2-deoxygluconokinase variant may be at least 5%, specifically 5 to 50% (preferably 10 to 40%) higher than that produced by the parent strain.

Still yet another aspect of the present invention provides a method for producing 5'-inosinic acid, comprising steps of: culturing the transformant in a medium; and recovering 5'-inosinic acid from the transformant or the medium in which the transformant has been cultured.

The culturing may be performed using a suitable medium and culture conditions known in the art, and any person skilled in the art may easily adjust and use the medium and the culture conditions. Specifically, the medium may be a liquid medium, without being limited thereto. Examples of the culturing method include, but are not limited to, batch culture, continuous culture, fed-batch culture, or a combination thereof.

According to one embodiment of the present invention, the medium should meet the requirements of a specific strain in a proper manner, and may be appropriately modified by a person skilled in the art. For culture media for *Escherichia* sp. strains, reference may be made to, but not limited to, a

known document (Manual of Methods for General Bacteriology, American Society for Bacteriology, Washington D.C., USA, 1981).

According to one embodiment of the present invention, the medium may contain various carbon sources, nitrogen sources, and trace element components. Examples of carbon sources that may be used include: sugars and carbohydrates such as glucose, sucrose, lactose, fructose, maltose, starch, and cellulose; oils and fats such as soybean oil, sunflower oil, castor oil, and coconut oil; fatty acids such as palmitic acid, stearic acid, and linoleic acid; alcohols such as glycerol and ethanol; and organic acids such as acetic acid. These substances may be used individually or as a mixture, without being limited thereto. Examples of nitrogen sources that may be used include peptone, yeast extract, meat extract, malt extract, corn steep liquor, soybean meal, urea, or inorganic compounds such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, and ammonium nitrate. The nitrogen sources may also be used individually or as a mixture, without being limited thereto. Examples of phosphorus sources that may be used include, but are not limited to, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts. In addition, the culture medium may contain, but is not limited to, metal salts such as magnesium sulfate or iron sulfate, which are required for growth. In addition, the culture medium may contain essential growth substances such as amino acids and vitamins. Moreover, suitable precursors may be used in the culture medium. The medium or individual components may be added to the culture medium batchwise or in a continuous manner by a suitable method during culturing, without being limited thereto.

According to one embodiment of the present invention, the pH of the culture medium may be adjusted by adding compounds such as ammonium hydroxide, potassium hydroxide, ammonia, phosphoric acid and sulfuric acid to the microorganism culture medium in an appropriate manner during the culturing. In addition, during the culturing, foaming may be suppressed using an anti-foaming agent such as a fatty acid polyglycol ester. Additionally, to keep the culture medium in an aerobic condition, oxygen or an oxygen-containing gas (for example, air) may be injected into the culture medium. The temperature of the culture medium may be generally 20° C. to 45° C., for example, 25° C. to 40° C. The culturing may be continued until a desired amount of a useful substance is produced. For example, the culturing time may be 10 hours to 160 hours.

According to one embodiment of the present invention, in the step of recovering 5'-inosinic acid from the cultured transformant or the medium in which the transformant has been cultured, the produced 5'-inosinic acid may be collected or recovered from the medium using a suitable method known in the art depending on the culture method. Examples of a method that may be used to recover the produced 5'-inosinic acid include, but are not limited to, centrifugation, filtration, extraction, spraying, drying, evaporation, precipitation, crystallization, electrophoresis, fractional dissolution (e.g., ammonium sulfate precipitation), chromatography (e.g., ion exchange, affinity, hydrophobicity and size exclusion), and the like.

According to one embodiment of the present invention, the step of recovering 5'-inosinic acid may be performed by centrifuging the culture medium at a low speed to remove biomass and separating the obtained supernatant through ion-exchange chromatography.

According to one embodiment of the present invention, the step of recovering 5'-inosinic acid may include a process of purifying the 5'-inosinic acid.

Advantageous Effects

The 5-dehydro-2-deoxygluconokinase variant according to the present invention is obtained by substituting one or more amino acids in the amino acid sequence constituting 5-dehydro-2-deoxygluconokinase to change the activity of the protein, and a recombinant microorganism comprising the 5-dehydro-2-deoxygluconokinase variant is capable of efficiently producing 5'-inosinic acid.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 shows the structure of a pK19msb plasmid according to one embodiment of the present invention.

MODE FOR INVENTION

Hereinafter, the present invention will be described in more detail. However, this description is merely presented by way of example to facilitate the understanding of the present invention, and the scope of the present invention is not limited by this exemplary description.

Example 1. Construction of Strain Expressing 5-Dehydro-2-Deoxygluconokinase Variant

To evaluate the effect of a variant (SEQ ID NO: 2) having a substitution of asparagine (N) for serine (S) at position 146 in the amino acid sequence of 5-dehydro-2-deoxygluconokinase (SEQ ID NO: 4) on the production of 5'-inosinic acid, the present inventors constructed a vector for expressing the 5-dehydro-2-deoxygluconokinase variant and a strain into which the vector has been introduced.

1-1. Construction of Vector for Expression of 5-Dehydro-2-Deoxygluconokinase Variant

Using the genomic DNA of wild-type *Corynebacterium stationis* ATCC6872 as a template, PCR reactions were performed using a primer pair of primers 1 and 2 and a primer pair of primers 3 and 4, respectively. Thereafter, using the two PCR products as templates, overlapping PCR was performed using a primer pair of primers 1 and 4 to obtain a single fragment. The PCR fragment and a pK19msb plasmid (SEQ ID NO: 5) were treated with the restriction enzyme *smal* (NEB) and ligated together using T4 ligase. The resulting plasmid was named pK_DD.

The PCR amplification was performed using Pfu PreMix (Bioneer) under the following conditions: denaturation at 95° C. for 5 min, and then 30 cycles, each consisting of 95° C. for 30 sec, 58° C. for 30 sec, and 72° C. for 1 min and 30 sec, followed by reaction at 72° C. for 5 min.

The primer sequences used for plasmid construction are shown in Table 1 below.

TABLE 1

Primer name	SEQ ID NO.	Primer sequence (5'→3')
Primer 1	6	AATGTGCTATTGTCAGGACATGC
Primer 2	7	GGCTCTTCGTTTAAACCGGTCA

TABLE 1-continued

Primer name	SEQ ID NO.	Primer sequence (5'→3')
Primer 3	8	TGACCGGTTTAAACGAAGAGCC
Primer 4	9	ACCTTCTGACTTAGGGCTGCTT

1-2. Construction of Mutant Strain into Which 5-Dehydro-2-Deoxygluconokinase Variant Has Been Introduced

An electrocompetent cell preparation method, a modification of the method of van der Rest et al., was used as a method for transformation of *Corynebacterium stationis* KCCM13339P.

First, *Corynebacterium stationis* KCCM13339P was primarily cultured in 10 mL of 2YT medium (containing 16 g/l of tryptone, 10 g/l of yeast extract, and 5 g/l of sodium chloride) supplemented with 2% glucose, thus preparing a seed culture. Isonicotinic acid hydrazine at a concentration of 1 mg/ml and 2.5% glycine were added to 100 ml of 2YT medium free of glucose. Next, the seed culture was inoculated into the 2YT medium to reach an OD₆₁₀ value of 0.3, and then cultured at 30° C. and 180 rpm for 5 to 8 hours so that the OD₆₁₀ value reached 0.6 to 0.7. The culture was kept on ice for 30 minutes, and then centrifuged at 3,500 rpm at 4° C. for 10 minutes. Thereafter, the supernatant was discarded and the precipitated *Corynebacterium stationis* KCCM13339P was washed 4 times with a 10% glycerol solution and finally re-suspended in 0.5 ml of a 10% glycerol solution, thereby preparing competent cells. Electroporation was performed using a Bio-Rad electroporator. The prepared competent cells and the constructed pK_DD vector were placed in an electroporation cuvette (0.2 mm), and then subjected to electroporation under conditions of 2.5 kV, 200Ω and 12.5 pF. Immediately after completion of the electroporation, 1 ml of a regeneration (RG) medium (containing 18.5 g/l brain heart infusion and 0.5 M sorbitol) was added to the cells which were then heat-treated at 46° C. for 6 minutes. Next, the cells were cooled at room temperature, transferred into a 15-ml cap tube, incubated at 30° C. for 2 hours, and plated on a selection medium (containing 5 g/l tryptone, 5 g/l NaCl, 2.5 g/l yeast extract, 18.5 g/l brain heart infusion powder, 15 g/l agar, 91 g/l sorbitol, and 20 pg/l kanamycin). The cells were cultured at 30° C. for 72 hours, and the generated colonies were cultured in medium until the stationary phase to induce secondary recombination. Then, the cells were diluted to 10⁻⁵ to 10⁻⁷, and plated on an antibiotic-free plate medium (containing 10% sucrose), and a strain having no kanamycin resistance and grown on the medium containing 10% sucrose was selected and named IDD-1.

Experimental Example 1. Evaluation of 5'-Inosinic Acid Productivity of Strain Expressing 5-Dehydro-2-Deoxygluconokinase Variant

5'-Inosinic acid productivity was compared between the parent strain KCCM13339P and the mutant strain IDD-1 into which the 5-dehydro-2-deoxygluconokinase variant has been introduced.

Each strain (parent strain or mutant strain) was inoculated at 1% by volume into a 100-mL flask containing 10 mL of the medium for 5'-inosinic acid production shown in Table 2 below, and cultured with shaking at 200 rpm at 34° C. for 45 hours. After completion of the culturing, the concentration of 5'-inosinic acid in the medium was measured using HPLC (Agilent), and the results are shown in Table 3 below.

TABLE 2

Component	Content
Glucose	70 g/L
(NH ₄) ₂ SO ₄	2 g/L
MgSO ₄	1 g/L
Urea	2 g/L
Yeast extract	20 g/L
KH ₂ PO ₄	2 g/L
FeSO ₄	10 mg/L
MnSO ₄	10 mg/L
Thiamine_HCl	5 mg/L
biotin	20 ug/L
Cystein	20 mg/L
Bata-alanine	20 mg/L
Adenine	30 mg/L

TABLE 3

Strain	5'-inosinic acid production (g/L)
KCCM13339P	19.8
IDD-1	23.0

As shown in Table 3 above, it was confirmed that the amount of 5'-inosinic acid produced by the mutant strain into which the 5-dehydro-2-deoxygluconokinase variant has been introduced was increased by about 16% compared to that produced by the parent strain, due to substitution of serine at position 146 with asparagine. These results suggest that introduction of a point mutation into 5-dehydro-2-deoxygluconokinase provides a significant effect on 5'-inosinic acid productivity.

So far, the present invention has been described with reference to the preferred embodiments. Those of ordinary skill in the art to which the present invention pertains will appreciate that the present invention may be embodied in modified forms without departing from the essential characteristics of the present invention. Therefore, the disclosed embodiments should be considered from an illustrative point of view, not from a restrictive point of view. The scope of the present invention is defined by the claims rather than the foregoing description, and all differences within the scope equivalent thereto should be construed as being included in the present invention.

Accession Number

Depository Authority: Korean Culture Center of Microorganisms (KCCM)

Accession Number: KCCM13339P

Deposit Date: Mar. 29, 2023

SEQUENCE LISTING

Sequence total quantity: 9

SEQ ID NO: 1 moltype = DNA length = 957

-continued

FEATURE
source

Location/Qualifiers
1..957
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 1

atggcgatt	taactagcgc	tcacgaagtt	ctggccatcg	gcaggcttgg	cgtggatatt	60
tatccactac	agtcggcggt	tgggctagaa	gaggtatctt	ctttcgccaa	atacctaggc	120
ggatccgcag	caaatgtctc	tgttcgagcc	gcacggcacg	ggcacaattc	cgcactgcta	180
tcacgagtag	gcaatgaccc	actgggtaag	tacctgctca	atgagctgga	aaggctcggg	240
gtagataatc	agtaacgtgc	tactgatccg	acttataaaa	ccccgctgac	tttgtgcgaa	300
attttccac	cagatgattt	cccgtgtgac	ttctaccgcg	agcccaaaag	tccggatatg	360
tgcacgaggt	cctctgacgt	caacctcgaa	gatgtccgcg	atgcagaaat	cttgtgggtt	420
accctgacgg	gtttaaaccg	agagccaagc	cgcggcgcac	atcaggaaat	cctgaacacc	480
cgcgccccgc	gcaagcacac	catcttggac	ttggactacc	gcgacatgtt	ctggaattcc	540
aagcaggaag	caacagagca	ggcacagtgg	gcaactcgagc	atgccaccgt	tcgggtcggg	600
aacaaggaag	aatgcgaggt	cgccatcggc	gaaaccgaac	cagaacgcgc	cggcaaggcg	660
ttgttggaa	atggcgtaga	gttggccatc	gttaagcaag	gccccaaagg	cgtgctggcc	720
atgactgaag	atgaaaccgt	cgaagtgcc	ccggtgtggg	tcgatgttgt	caacggcctc	780
ggcgctggcg	atgccttcgg	tggcgcgctg	tgccacggtc	tggtgtccga	ctggccactc	840
gaaaaagtc	tgcgctttgc	caacgcggca	ggctcgggtg	ttgcccggag	cttagagtgc	900
agcaccgcc	tgccaaccac	tgaagaagtt	gaagcagccc	taagtcagaa	ggtctag	957

SEQ ID NO: 2

moltype = AA length = 318

FEATURE
source

Location/Qualifiers
1..318
mol_type = protein
organism = synthetic construct

SEQUENCE: 2

MAYLTSABEV	LAIGRLGVDI	YPLQSGVGL	EVSSFGKYL	GSAANVSVA	ARHGHSALL	60
SRVGNLPLG	YLLNELERL	VDNQYVATD	TYKTPLTLC	IFPDDDFLY	FYREPKAPD	120
CISSSDVNL	DVRDAEILWF	TLTGLNEEPS	RGHQEILNT	RARRKHTILD	LDYRDMFWNS	180
KQATEQAQW	ALEHATVAVG	NKEECEVAIG	ETEPERAGKA	LLEHGKVLAI	VKQGPKGVL	240
MTEDTVEVP	PVWVDVNG	GAGDAFGGAL	CHGLLSDWPL	EKVLRFANAA	GSVVAGRLEC	300
STAMPTTEEV	EAALSQKV					318

SEQ ID NO: 3

moltype = DNA length = 957

FEATURE
source

Location/Qualifiers
1..957
mol_type = genomic DNA
organism = Corynebacterium stationis

SEQUENCE: 3

atggcgatt	taactagcgc	tcacgaagtt	ctggccatcg	gcaggcttgg	cgtggatatt	60
tatccactac	agtcggcggt	tgggctagaa	gaggtatctt	ctttcgccaa	atacctaggc	120
ggatccgcag	caaatgtctc	tgttcgagcc	gcacggcacg	ggcacaattc	cgcactgcta	180
tcacgagtag	gcaatgaccc	actgggtaag	tacctgctca	atgagctgga	aaggctcggg	240
gtagataatc	agtaacgtgc	tactgatccg	acttataaaa	ccccgctgac	tttgtgcgaa	300
attttccac	cagatgattt	cccgtgtgac	ttctaccgcg	agcccaaaag	tccggatatg	360
tgcacgaggt	cctctgacgt	caacctcgaa	gatgtccgcg	atgcagaaat	cttgtgggtt	420
accctgacgg	gtttaaaccg	agagccaagc	cgcggcgcac	atcaggaaat	cctgaacacc	480
cgcgccccgc	gcaagcacac	catcttggac	ttggactacc	gcgacatgtt	ctggaattcc	540
aagcaggaag	caacagagca	ggcacagtgg	gcaactcgagc	atgccaccgt	tcgggtcggg	600
aacaaggaag	aatgcgaggt	cgccatcggc	gaaaccgaac	cagaacgcgc	cggcaaggcg	660
ttgttggaa	atggcgtaga	gttggccatc	gttaagcaag	gccccaaagg	cgtgctggcc	720
atgactgaag	atgaaaccgt	cgaagtgcc	ccggtgtggg	tcgatgttgt	caacggcctc	780
ggcgctggcg	atgccttcgg	tggcgcgctg	tgccacggtc	tggtgtccga	ctggccactc	840
gaaaaagtc	tgcgctttgc	caacgcggca	ggctcgggtg	ttgcccggag	cttagagtgc	900
agcaccgcc	tgccaaccac	tgaagaagtt	gaagcagccc	taagtcagaa	ggtctag	957

SEQ ID NO: 4

moltype = AA length = 318

FEATURE
source

Location/Qualifiers
1..318
mol_type = protein
organism = Corynebacterium stationis

SEQUENCE: 4

MAYLTSABEV	LAIGRLGVDI	YPLQSGVGL	EVSSFGKYL	GSAANVSVA	ARHGHSALL	60
SRVGNLPLG	YLLNELERL	VDNQYVATD	TYKTPLTLC	IFPDDDFLY	FYREPKAPD	120
CISSSDVNL	DVRDAEILWF	TLTGLSEEPS	RGHQEILNT	RARRKHTILD	LDYRDMFWNS	180
KQATEQAQW	ALEHATVAVG	NKEECEVAIG	ETEPERAGKA	LLEHGKVLAI	VKQGPKGVL	240
MTEDTVEVP	PVWVDVNG	GAGDAFGGAL	CHGLLSDWPL	EKVLRFANAA	GSVVAGRLEC	300
STAMPTTEEV	EAALSQKV					318

SEQ ID NO: 5

moltype = DNA length = 5719

FEATURE
source

Location/Qualifiers
1..5719
mol_type = other DNA
organism = synthetic construct

SEQUENCE: 5

tgccgcaagc	actcaggggc	caagggtctg	taagggaagc	ggaacacgta	gaaagccagt	60
ccgcagaaac	gggtctgacc	ccggatgaat	gtcagctact	gggctatctg	gacaaggga	120

-continued

aacgcaagcg	caaagagaaa	gcaggtagct	tgcagtgggc	ttacatggcg	atagctagac	180
tgggcggttt	tatggacagc	aagcgcaaccg	gaattgccag	ctggggcgcc	ctctggttaag	240
gttgggaagc	cctgcaaaagt	aaactggatg	gctttcttgc	cgccaaggat	ctgatggcgc	300
aggggatcaa	gatctgatca	agagacagga	tgaggatcgt	ttcgcatgat	tgaacaagat	360
ggattgcacg	caggtttctcc	ggcgccttgg	gtggagaggc	tattcggtca	tgactgggca	420
caacagacaa	tcggctgctc	tgatgccgcc	gtgttccggc	tgtcagcgca	ggggcgcccc	480
gttctttttg	tcaagaccga	cctgtccggg	gccctgaatg	aactccaaga	cgaggcagcg	540
cggtatcgt	ggctggccac	gacgggcgtt	ccttgccgag	ctgtgctcga	cgttgtcact	600
gaagcgggaa	gggactggct	gctattgggc	gaagtgcggg	ggcaggatct	cctgtcatct	660
caacctgtct	ctgccgagaa	agtatccatc	atggctgatg	caatgcggcg	gctgcatacg	720
cttgatccgg	ctacotgcc	attcgaccac	caagcgaaac	atcgcatcga	gcgagcacgt	780
actcggatgg	aagccggtct	tgtcgatcag	gatgatctgg	acgaagagca	tcaggggctc	840
gcgcagccg	aactgttcgc	caggctcaag	gcgcggatgc	ccgaacggcg	ggatctcgtc	900
gtgacccatg	cgatgtccgc	cttgccgaat	atcatgggtg	aaaatggccg	cttttctgga	960
ttcatcgact	gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttggctacc	1020
cgtgatattg	ctgaagagct	tgccggcgaa	tgggctgacc	gcttctcgt	gctttaacgg	1080
atcgccgctc	cgatttcgca	gcgcctcgcc	ttctatcgcc	ttcttgacga	gttctctga	1140
ggcggactct	gggttctgct	agaggatcga	tcctttttaa	cccatcacat	atacctgccg	1200
ttcactatta	tttagtgaaa	tgagatatta	tgatattttc	tgaattgtga	ttaaaaaggc	1260
aactttatgc	ccatgcaaca	gaaactataa	aaaatacaga	gaatgaaaag	aaacagatag	1320
atttttttagt	tccttagggc	cgtagtctgc	aaatcctttt	atgattttct	atcaaacaaa	1380
agaggaaaat	agaccagttg	caatccaaac	gagagtctaa	tagaatgagg	tcgaaaagta	1440
aatcgccggg	gttgtttact	gataaagcag	gcaagacctc	aaatgtgtta	agggcgaagt	1500
gtatactttg	gcgtcaccct	ttacatattt	taggtctttt	tttatgttgc	gtaactaact	1560
tgccatcttc	aaaacaggag	gctggaagaa	gcagaccgct	aacacagtac	ataaaaaagg	1620
agacatgaac	gatgaacatc	aaaaagtttg	caaaaacagc	aacagtatta	acctttacta	1680
ccgcaactgt	ggcaggagcg	gcaactcaag	cgtttgcgaa	agaaacgaac	caaaagccat	1740
ataaggaaaac	atacgggcat	tcctcatatta	cacgccatga	tatgctgcaa	atccctgaac	1800
agcaaaaaaa	tgaaaaatat	caagtttctg	aatttgattc	gtccacaatt	aaaaatatct	1860
cttctgcaaa	aggcctggac	gcttgggaca	gctggccatt	acaaaacgct	gacggcactg	1920
tcgcaaaacta	tcacggctac	ccatcgtctc	ttgcattagc	cggagatcct	aaaaatgcgg	1980
atgacacatc	gattttacatg	ttctatcaaa	aagtcggcga	aacttctatt	gacagctgga	2040
aaaaacgctg	ccgcgtcttt	aaagacagcg	acaaattcga	tgcaaatgat	tcctatcctaa	2100
aaagccaaaac	acaagaatgg	tcagggttcag	ccacattttac	atctgacgga	aaaactcgtt	2160
tattctacac	tgattttctc	ggttaaacatt	acggcaaaac	aacactgaca	actgcacaag	2220
ttaacgtatc	agcatcagac	agctctttga	acatcaacgg	tgtagaggat	tataaatcaa	2280
tcctttgacgg	tgacggaaaa	acgtatcaaa	atgtacagca	gttcatcgat	gaaggcaact	2340
acagctcagg	cgacaacccat	acgctgagag	atcctcacta	cgtagaagat	aaaggccaca	2400
aatacttagt	atttgaagca	aacactggaa	ctgaagatgg	ctaccaaggc	gaagaatctt	2460
tatttaacaa	agcatactat	ggcaaaaagca	catcattctt	ccgtcaagaa	agtcaaaaac	2520
ttctgcaaa	cgataaaaaa	cgcacggctg	agttagcaaa	cgccgctctc	ggtagatttg	2580
agctaaacga	tgatttacaca	ctgaaaaaag	tgatgaaacc	gctgattgca	tcatacacag	2640
taacagatga	aattgaacgc	gcgaacgtct	ttaaaatgaa	cggcaaatgg	tacctgttca	2700
ctgactcccg	cggatcaaaa	atgacgattg	acggcattac	gtctaacgat	atttacatgc	2760
ttggttatgt	ttctaatctc	ttaaactggc	catacaagcc	gctgaacaaa	actggccttg	2820
tgttaaaaat	ggatcttgat	cctaacgatg	taacctttac	ttactcacac	ttcgtgtgac	2880
ctcaagcgaa	aggaacaacat	gtcgtgatta	caagctatat	gacaaacaga	ggattctacg	2940
cagacaacaa	atcaacggtt	gcgcggagct	tcctgctgaa	catcaaaagg	aagaaaaacat	3000
ctggtgtcaa	agacagcatc	cttgaacaa	gacaattaac	agttaaacaa	taaaaacgca	3060
aaagaaaaat	ccgatgggta	ccgagcgaaa	tgaccgacca	agcgaacgcc	aaactgccat	3120
cacgagattt	cgattccacc	gcgcgcttct	atgaaagggt	gggcttcgga	atcgttttcc	3180
gggacgcccc	cgccgagctg	ctcatagtcc	acgacgcccc	tgattttgta	gccctggccg	3240
acggccagca	ggtaggccga	caggctcatg	ccggcccgcc	ccgccttttc	ctcaatcgct	3300
cttcggtcgt	ctggaaggca	gtacaccttg	ataggtgggc	tgcccttcc	gggtggcttg	3360
gtttcatcag	ccatccgctt	gcctcatctc	gttacgcccc	cggtagcccg	ccagcctcgc	3420
agagcaggat	tcctgttgag	acccgccagg	tgccaataag	ggacagtgaa	gaaggaaacac	3480
ccgctccgcg	gtgggcctac	ttcacctatc	ctgcccggtc	gacgcggttg	gatacaccaa	3540
ggaaagtcta	cacgaacctc	ttggcaaaaat	cctgtatatc	gtgcgaaaaa	ggatggatat	3600
accgaaaaaa	tcgtataaat	gaccccgaa	cagggttatg	cagcgaaaaa	gcgctgcttc	3660
cctgctgttt	tgtggaatat	ctaccgactg	gaaacaggca	aatcgaggaa	attactgaac	3720
tgaggggaca	ggcgagagac	agctcctgaa	aatctcgata	actcaaaaaa		3780
tacgccccgt	agtgatctta	tttcattatg	gtgaaagtgt	gaacctctta	cgtgcgatc	3840
aacgtctcat	tttcgcaaaa	agttggccca	gggcttcccg	gtatcaacag	ggacaccagg	3900
atttatttat	ctcgcgaagt	gatcttccgt	cacaggatatt	tattcgccgc	aaagtgcgtc	3960
gggtgatgct	gccaaacttac	tgatttagtg	tatgatgggt	tttttgaggt	gctccagtggt	4020
cttctgtttc	tatcagctcc	tgaaaaatctc	gataaactcaa	aaaatacgcc	cggtagtgtat	4080
cttatttcat	tatggtgaaa	ggtggaacct	cttactgtcc	gatcaacgtc	tcattttcgc	4140
caaaagttyg	cccagggtct	ccgggtatca	acagggacac	caggatttat	ttattctgct	4200
aagtgatctt	ccgtcacagg	tatttatctg	gcgcaaaagt	cgtcgggtga	tgctgccaac	4260
ttactgattt	agtgatgat	gggtgttttg	aggtgctcca	gtggcttctg	tttctatcag	4320
ggctggatga	tcctccagcg	cggggatctc	atgctggagt	tcctcgccca	ccccaaaagg	4380
atctaggatga	agatcctttt	tgataatctc	atgacaaaaa	tcctttaacg	tgagttttcg	4440
ttccactgag	cgtcagaccc	cgtagaaaaag	atcaaaaggat	cttcttgaga	tccttttttt	4500
ctcgcgctaa	gtctgctctt	gcaaaacaaa	aaaccaccgc	taccagcggt	ggtttgtttg	4560
ccggatcaag	agctaccaac	tcctttttccg	aaggtaactg	gcttcagcag	agcgcagata	4620
ccaaataactg	ttcttctagt	gtagccgtag	ttagccacc	acttcaagaa	ctctgtagca	4680
ccgctacat	acctcgctct	gctaactctg	ttaccagtggt	ctgctgccag	tgccgataag	4740
tcgtgtctta	cggggttgga	ctcaagacga	tagttaccgg	ataaggcgca	gcggtcgggc	4800
tgaacggggg	gttcgtgcac	acagcccgac	ttggagcgaa	gcacctacac	cgaactgaga	4860

-continued

```

tacctacagc gtgagctatg agaagcgcc acgcttcccg aagggagaaa ggcggacag 4920
tatccggtaa gcggcagggg cggaacagga gagcgacga gggagcttcc agggggaaac 4980
gcctgggtatc tttatagtc tgcgggttt cgccacctct gacttgagcg tcgatttttg 5040
tgatgctcgt caggggggcg gagcctatgg aaaaacgcca gcaacgcggc ctttttacgg 5100
ttcctggcct tttgctggcc ttttgctcac atgttctttc ctgctgtatc cctgattct 5160
gtggataaac gtattaccgc ctttgagtga gctgataccg ctgcgcgcag ccgaacgacc 5220
gagcgacagc agtcagtgag cgaggaagcg gaagagcgcc caatacgcaa accgcctctc 5280
cccgcgctgt ggccgattca ttaatgcagc tggcacgaca gggttccgca ctggaaagcg 5340
ggcagtgagc gcaacgcaat taatgtgagt tagctcactc attaggcacc ccaggcttta 5400
cactttatgc ttccggctcg tatgttgtgt ggaattgtga gcggataaca atttcacaca 5460
ggaaacagct atgacatgat tacgccaaagc ttgcatgcct gcaggtcgac tctagaggat 5520
ccccgggtac cgagctcgaa ttcaactggc gtctgttttac aacgtcgtga ctgggaaaac 5580
cctggcggtta cccaacttaa tcgccttgca gcacatcccc ctttcgccag ctggcgtaat 5640
agcgaagagg ccgcaccca tcgcccttcc caacagttgc gcagcctgaa tggcgaatga 5700
cgataagcta gcttcacgc                                     5719

```

```

SEQ ID NO: 6      moltype = DNA length = 23
FEATURE          Location/Qualifiers
source          1..23
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 6
aatgtgctat tgtcaggaca tgc                                     23

```

```

SEQ ID NO: 7      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source          1..22
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 7
ggctcttcgt ttaaaccggt ca                                     22

```

```

SEQ ID NO: 8      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source          1..22
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 8
tgaccggttt aaacgaagag cc                                     22

```

```

SEQ ID NO: 9      moltype = DNA length = 22
FEATURE          Location/Qualifiers
source          1..22
                mol_type = other DNA
                organism = synthetic construct

```

```

SEQUENCE: 9
accttctgac ttagggctgc tt                                     22

```

The invention claimed is:

1. A 5-dehydro-2-deoxygluconokinase variant consisting of the amino acid sequence of SEQ ID NO: 2 in which serine at position 146 in the amino acid sequence of SEQ ID NO: 4 is substituted with asparagine.
2. A polynucleotide encoding the variant of claim 1.
3. A transformant comprising the variant of claim 1.
4. The transformant of claim 3, which is a *Corynebacterium* sp. microorganism.

5. The transformant of claim 3, which has ability to produce 5'-inosinic acid.

6. A method for producing 5'-inosinic acid, comprising steps of:

culturing the transformant of claim 3 in a medium; and recovering 5'-inosinic acid from the transformant or the medium in which the transformant has been cultured.

7. A transformant comprising the polynucleotide of claim 2.

* * * * *