

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12403155
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Pauza; Charles David et al.

Methods and compositions for the activation of gamma-delta T-cells

Abstract

The present invention relates generally to methods and compositions for gene therapy and immunotherapy that activate gamma delta T-cells, and in particular, can be used in the treatment of various cancers and infectious diseases.

Inventors: Pauza; Charles David (Rockville, MD), Li; Haishan (Rockville, MD), Lahusen; Tyler (Rockville, MD), Liou; Mei-Ling (Rockville, MD)

Applicant: American Gene Technologies International Inc. (Rockville, MD)

Family ID: 63446778

Assignee: American Gene Technologies International Inc. (Rockville, MD)

Appl. No.: 18/055525

Filed: November 15, 2022

Prior Publication Data

Document Identifier	Publication Date
US 20230277579 A1	Sep. 07, 2023

Related U.S. Application Data

continuation parent-doc US 16988427 20200807 US 11534450 child-doc US 18055525
continuation parent-doc US 16530908 20190802 US 10772905 20200915 child-doc US 16988427
continuation parent-doc US 16132247 20180914 US 10420789 20190914 child-doc US 16530908
continuation parent-doc US 15904131 20180223 US 10137144 20181127 child-doc US 16132247
continuation parent-doc WO PCT/US2017/013399 20170113 PENDING child-doc US 15652080
continuation-in-part parent-doc US 15652080 20170717 US 9914938 20180313 child-doc US 15904131
us-provisional-application US 62279474 20160115

Publication Classification

Int. Cl.: C12N15/113 (20100101); A61K31/675 (20060101); A61K31/7105 (20060101); A61K47/68 (20170101); A61K47/69 (20170101); A61P35/02 (20060101); C12N15/86 (20060101); C12Q1/68 (20180101)

U.S. Cl.:

CPC A61K31/7105 (20130101); A61K31/675 (20130101); A61K47/6807 (20170801); A61K47/6901 (20170801); A61P35/02 (20180101); C12N15/1137 (20130101); C12N15/86 (20130101); C12Y205/0101 (20130101); A61K2300/00 (20130101); C12N2310/14 (20130101); C12N2310/531 (20130101); C12N2330/51 (20130101); C12N2740/16032 (20130101); C12N2740/16043 (20130101); C12N2740/16045 (20130101); C12N2750/14143 (20130101); C12N2810/6072 (20130101)

Field of Classification Search

USPC: None

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5668255	12/1996	Murphy	N/A	N/A
8124752	12/2011	Bumcrot et al.	N/A	N/A
8287857	12/2011	Dudley et al.	N/A	N/A
8993532	12/2014	Hannon et al.	N/A	N/A
9522176	12/2015	DeRosa et al.	N/A	N/A
9834790	12/2016	Pauza et al.	N/A	N/A
9914938	12/2017	Pauza et al.	N/A	N/A
10023880	12/2017	Pauza et al.	N/A	N/A
10036040	12/2017	Pauza et al.	N/A	N/A
10137144	12/2017	Pauza et al.	N/A	N/A
10208295	12/2018	DeRosa et al.	N/A	N/A
10420789	12/2018	Pauza et al.	N/A	N/A
10428350	12/2018	Pauza et al.	N/A	N/A
10472649	12/2018	Pauza et al.	N/A	N/A
10767183	12/2019	Lahusen et al.	N/A	N/A
10772905	12/2019	Pauza et al.	N/A	N/A
2004/0142416	12/2003	Laipis et al.	N/A	N/A
2004/0180847	12/2003	Dobie et al.	N/A	N/A
2004/0192629	12/2003	Xu et al.	N/A	N/A
2004/0248296	12/2003	Beresford et al.	N/A	N/A
2004/0265306	12/2003	Arthos et al.	N/A	N/A
2005/0019927	12/2004	Markus et al.	N/A	N/A
2005/0138677	12/2004	Pfister et al.	N/A	N/A
2006/0057553	12/2005	Aguiar-Cordova	N/A	N/A
2006/0073576	12/2005	Barnett et al.	N/A	N/A
2006/0246520	12/2005	Champagne et al.	N/A	N/A

2007/0141679	12/2006	Sodroski	N/A	N/A
2008/0003225	12/2007	Vie et al.	N/A	N/A
2008/0003682	12/2007	Lois-Caballe et al.	N/A	N/A
2008/0153737	12/2007	Lieberman et al.	N/A	N/A
2008/0199961	12/2007	Rasko et al.	N/A	N/A
2008/0227736	12/2007	Chen et al.	N/A	N/A
2008/0293142	12/2007	Liu et al.	N/A	N/A
2009/0148936	12/2008	Stout et al.	N/A	N/A
2009/0304688	12/2008	Fournie et al.	N/A	N/A
2010/0286166	12/2009	Rodriguez et al.	N/A	N/A
2010/0316676	12/2009	Sanders et al.	N/A	N/A
2011/0008803	12/2010	Stockwell	N/A	N/A
2012/0027725	12/2011	Galvin et al.	N/A	N/A
2013/0090371	12/2012	Lu	N/A	N/A
2013/0122380	12/2012	Visco et al.	N/A	N/A
2013/0211380	12/2012	Cabrera Aquino et al.	N/A	N/A
2014/0155468	12/2013	Gregory et al.	N/A	N/A
2014/0178340	12/2013	Robbins et al.	N/A	N/A
2014/0348794	12/2013	Chiorini et al.	N/A	N/A
2015/0010578	12/2014	Balazs et al.	N/A	N/A
2015/0018539	12/2014	Fellmann	N/A	N/A
2015/0126580	12/2014	DePinho et al.	N/A	N/A
2016/0060707	12/2015	Goldenberg et al.	N/A	N/A
2016/0243169	12/2015	Chen et al.	N/A	N/A
2016/0287635	12/2015	Hariri et al.	N/A	N/A
2017/0015976	12/2016	Nelson	N/A	N/A
2017/0028036	12/2016	Mingozzi et al.	N/A	N/A
2017/0335344	12/2016	Pauza et al.	N/A	N/A
2018/0142257	12/2017	Pauza et al.	N/A	N/A
2018/0142258	12/2017	Pauza et al.	N/A	N/A
2018/0195050	12/2017	Szalay et al.	N/A	N/A
2018/0256624	12/2017	Pauza et al.	N/A	N/A
2018/0305716	12/2017	Pauza et al.	N/A	N/A
2018/0355032	12/2017	Roberts et al.	N/A	N/A
2019/0062786	12/2018	Pauza et al.	N/A	N/A
2019/0078096	12/2018	Lahusen et al.	N/A	N/A
2019/0083523	12/2018	Pauza et al.	N/A	N/A
2019/0218573	12/2018	Pauza et al.	N/A	N/A
2019/0388456	12/2018	Pauza et al.	N/A	N/A
2020/0017570	12/2019	Walcheck et al.	N/A	N/A
2020/0063161	12/2019	Pauza et al.	N/A	N/A
2020/0087682	12/2019	Lahusen et al.	N/A	N/A
2020/0181645	12/2019	Pauza et al.	N/A	N/A
2020/0318081	12/2019	Lahusen et al.	N/A	N/A
2020/0354679	12/2019	Niazi	N/A	N/A
2021/0047644	12/2020	Lahusen et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
101516365	12/2008	CN	N/A

101679466	12/2009	CN	N/A
101805750	12/2009	CN	N/A
105112370	12/2014	CN	N/A
108883100	12/2017	CN	N/A
3402483	12/2017	EP	N/A
3426777	12/2018	EP	N/A
2007527240	12/2006	JP	N/A
2008538174	12/2007	JP	N/A
2013530152	12/2012	JP	N/A
2015518838	12/2014	JP	N/A
2016502404	12/2015	JP	N/A
WO 2002020554	12/2001	WO	N/A
WO 2003093436	12/2002	WO	N/A
WO 2005033282	12/2004	WO	N/A
WO-2006039721	12/2005	WO	N/A
WO-2006089001	12/2005	WO	N/A
WO 2007000668	12/2006	WO	N/A
WO-2007015122	12/2006	WO	N/A
WO-2007132292	12/2006	WO	N/A
WO-2008025025	12/2007	WO	N/A
WO 2009001224	12/2007	WO	N/A
WO 2009100928	12/2008	WO	N/A
WO 2009147445	12/2008	WO	N/A
WO 2010051521	12/2009	WO	N/A
WO-2010111522	12/2009	WO	N/A
WO 2010117974	12/2009	WO	N/A
WO-2010119039	12/2009	WO	N/A
WO 2010127166	12/2009	WO	N/A
WO 2011008348	12/2010	WO	N/A
WO-2011148194	12/2010	WO	N/A
WO-2012048303	12/2011	WO	N/A
WO 2012061075	12/2011	WO	N/A
WO-2012071559	12/2011	WO	N/A
WO-2012145624	12/2011	WO	N/A
WO-2013056148	12/2012	WO	N/A
WO-2013096455	12/2012	WO	N/A
WO-2014117050	12/2013	WO	N/A
WO 2014187881	12/2013	WO	N/A
WO-2014195159	12/2013	WO	N/A
WO-2015017755	12/2014	WO	N/A
WO 2015061491	12/2014	WO	N/A
WO 2015078999	12/2014	WO	N/A
WO-2015164759	12/2014	WO	N/A
WO 2016046234	12/2015	WO	N/A
WO 2016122058	12/2015	WO	N/A
WO-2017068077	12/2016	WO	N/A
WO 2017100551	12/2016	WO	N/A
WO-2017123918	12/2016	WO	N/A
WO-2017156311	12/2016	WO	N/A
WO-2017165641	12/2016	WO	N/A

WO-2017173453	12/2016	WO	N/A
WO 2018126112	12/2017	WO	N/A
WO-2018148443	12/2017	WO	N/A
WO 2018187231	12/2017	WO	N/A
WO-2018232359	12/2017	WO	N/A
WO 2019070674	12/2018	WO	N/A
WO-2020011247	12/2019	WO	N/A
WO-2020097049	12/2019	WO	N/A
WO 2020243717	12/2019	WO	N/A

OTHER PUBLICATIONS

Nielson et al., Molecular Therapy vol. 15, Suppl. 1:S270, May 2007. cited by examiner
JP Office Action in Japanese Application No. 2022-006999, dated Jan. 5, 2023, 20 pages (with English translation). cited by applicant
IL Office Action issued in Application No. 271274 on Aug. 6, 2023, 11 pages. cited by applicant
JP; Office Action issued in Application No. 2022-006999 on Sep. 20, 2023. cited by applicant
USPTO; Examiner's Answer in U.S. Appl. No. 16/614,682, dated Sep. 27, 2023. cited by applicant
Brake et al., "Lentiviral Vector Design for Multiple shRNA Expression and Durable HIV-1 Inhibition," Molecular Therapy, 16(3), 557-564, 2008. cited by applicant
KR; Office Action issued Oct. 20, 2023 in Application No. 10-2020-7000631. cited by applicant
UAE; Office Action issued Oct. 20, 2023 in Application No. P6001801/2019. cited by applicant
JP; Office Action issued Oct. 19, 2023 in Application No. 2021-523916. cited by applicant
US; Restriction Requirement issued in U.S. Appl. No. 17/570,313 on Nov. 15, 2023. cited by applicant
CN; Office Action issued in Application No. 201880039828.4 on Nov. 30, 2023. cited by applicant
IL; Notice of Allowance issued in Application No. 297238 on Dec. 11, 2023. cited by applicant
Altschul S.F., et al., "Basic Local Alignment Search Tool," Journal Molecular Biology, 1990, vol. 215, pp. 403-410. cited by applicant
Altschul S.F., et al., "Gapped BLAST and PSI-BLAST: A New Generation of Protein Database Search Programs," Nucleic Acids Research, Jul. 1997, vol. 25, No. 17, pp. 3389-3402. cited by applicant
Ausubel F.M., et al., "Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology," Wiley, John & Sons, Inc., 1995, 1 Page. cited by applicant
Benyamine A., et al., "BTN3A Molecules Considerably Improve Vy9Vδ2T Cells-based Immunotherapy in Acute Myeloid Leukemia," Oncoimmunology, Oct. 2, 2016, vol. 5, No. 10, 10 Pages, (E1146843) *the whole document*. cited by applicant
Berge et al., "Pharmaceutical Salts" Jan. 1977, Journal of Pharmaceutical Sciences, 66(1):1-19. cited by applicant
Capietto A-H., et al., "Stimulated Yδ T Cells Increase the in Vivo Efficacy of Trastuzumab in HER-2+ Breast Cancer," Journal of Immunology, 2011, vol. 187(2), pp. 1031-1038. cited by applicant
Chen H.C., et al., "An Unconventional TRAIL to Cancer Therapy," European Journal of Immunology, 2013, vol. 43, No. 12, pp. 3159-3162, DOI:10.1002/eji.201344105, ISSN 0004789814, XP071226184. cited by applicant
Chen Z., et al., "CD16+ Gammadelta T Cells Mediate Antibody Dependent Cellular Cytotoxicity: Potential Mechanism In The Pathogenesis Of Multiple Sclerosis," Clinical Immunology, 2008, vol. 128(2), pp. 219-227. cited by applicant
Cheng M., et al., "Establishment, Characterization, and Successful Adaptive Therapy Against Human Tumors of NKG Cell, a New Human NK Cell Line", Cell Transplantation, Jun. 2011, vol. 20, pp. 1731-1746. cited by applicant

Coligan J.E., et al., "Current Protocols in Protein Science," Short Protocols in Protein Science, 1996, vol. 24, No. 409, 1 Page. cited by applicant

Corrected Notice of Allowance for U.S. Appl. No. 16/563,738, mailed Aug. 31, 2022, 5 Pages. cited by applicant

Couzi L., et al., "Antibody-Dependent Anti-Cytomegalovirus Activity of Human Gammadelta T Cells Expressing CD16 (FcgammaRIIIa)," Blood, 2012, vol. 119(6), pp. 1418-1427. cited by applicant

Davis-Gardner M.E., et al., "eCD4-Ig promotes ADCC activity of sera from HIV-1-infected patients", Department of Immunology and Microbiology, The Scripps Research Institute, PLOS Pathogen, Dec. 18, 2017, 19 Pages, Retrieved from URL: <https://doi.org/10.1371/journal.ppat.1006786>. cited by applicant

Deveraux J., et al., "A Comprehensive Set of Sequence Analysis Programs for the VAX," Nucleic Acids Research, 1984, vol. 12, No. 1, pp. 387-395. cited by applicant

Dieli F., et al., "Targeting Human $\gamma\delta$ T Cells with Zoledronate and Interleukin-2 for Immunotherapy of Hormone-Refractory Prostate Cancer," Europe PMC Funders Group, Cancer Research, Aug. 1, 2007, vol. 67, No. 15, pp. 7450-7457. cited by applicant

Extended European Search Report for European Application No. 19883230.5, mailed Jul. 21, 2022, 9 Pages. cited by applicant

Extended European Search Report for European Application No. 17764128.9, mailed Aug. 12, 2019, 8 Pages. cited by applicant

Extended European Search Report for European Application No. 18817253.0, mailed Feb. 10, 2021, 8 Pages. cited by applicant

Extended European Search Report for European Application No. 22154806.8, mailed Jul. 4, 2022, 8 Pages. cited by applicant

Extended European Search Report for European Application No. 17739028.3, mailed Jun. 6, 2019, 8 Pages. cited by applicant

Final Office Action for U.S. Appl. No. 16/132,247, mailed Jul. 1, 2019, 7 Pages. cited by applicant

Final Office Action for U.S. Appl. No. 16/182,443, mailed May 2, 2019, 07 Pages. cited by applicant

Final Office Action for U.S. Appl. No. 16/614,682, mailed Aug. 2, 2022, 34 Pages. cited by applicant

First Office Action in the CN Application No. 201780017712.6, mailed May 8, 2020, 10 Pages. cited by applicant

Fisher J.P.H., et al., "Effective Combination Treatment of GD2-Expressing Neuroblastoma and Ewing's Sarcoma Using Anti-GD2 ch14.18/CHO Antibody with Vy9V52+ γ T Cells," Oncolmmunology, 2016, vol. 5, Issue No. 1, e1025194, 32 Pages. cited by applicant

Fujiwara Y., et al., "A Nucleolar Stress-Specific p53-miR-101 Molecular Circuit Functions as an Intrinsic Tumor-Suppressor Network," EBiomedicine, NL, Jul. 7, 2018, vol. 33, pp. 33-48, DOI: 10.1016/j.ebiom.2018.06.031, ISSN: 2352-3964, XP055939874, Retrieved from URL: <https://www.sciencedirect.com/science/article/pii/S235239641830238X?via%3Dihub>. cited by applicant

Gagniuc P., et al., "Eukaryotic Genomes May Exhibit up to 10 Generic Classes of Gene Promoters," BMC Genomics, 2012, vol. 13, 17 Pages, DOI:10.1186/1471-2164-13-512, XP021134695. cited by applicant

Gennaro A.R., "Remington's Pharmaceutical Sciences," 17th edition, Mack Publishing Company, Easton, Pa., Oct. 1985, vol. 74, No. 10, pp. 1143-1144. cited by applicant

Gertner-Dardenne J., et al., "Bromohydrin Pyrophosphate Enhances Antibody-Dependent Cell-Mediated Cytotoxicity Induced by Therapeutic Antibodies," Blood, 2009, vol. 113(20), pp. 4875-4884. cited by applicant

Harlow E., et al., "Antibodies: A Laboratory Manual," Cold Spring Harbor Laboratory Press, 1988,

1 Page. cited by applicant

Harly C., et al., "Key Implication of CD277/butyrophilin-3 (BTN3A) in Cellular Stress Sensing by a Major Human T-cell Subset," Blood, Sep. 13, 2012, vol. 120, No. 11, pp. 2269-2279, DOI:10.1182/blood-2012-05-430470, ISSN 00064971, XP055081172. cited by applicant

Hassan G., et al., "Isolation of Umbilical Cord Mesenchymal Stem Cells Using Human Blood Derivative Accompanied With Explant Method," Stem Cell Investigation, 2019, pp. 1-8. cited by applicant

Herrera L., et al., "Adult Peripheral Blood and Umbilical Cord Blood NK Cells Are Good Sources for Effective CAR Therapy Against CD19 Positive Leukemic Cells," Scientific Reports, Dec. 2019, vol. 9, Article. 18729, 2 Pages. cited by applicant

Huang Q., et al., "An Efficient Protocol to Generate Placental Chorionic Plate-derived Mesenchymal Stem Cells with Superior Proliferative and Immunomodulatory Properties," Stem Cell Research & Therapy, 2019, vol. 10(301), pp. 1-15. cited by applicant

International Search Report and Written Opinion for International Application No. PCT/US2021/020721, mailed Jul. 21, 2021, 10 Pages. cited by applicant

International Search Report and Written Opinion for International Application No. PCT/US2019/059828, mailed Feb. 14, 2020, 12 Pages. cited by applicant

International Search Report and Written Opinion for International Application No. PCT/US2022/013422, mailed May 13, 2022, 19 Pages. cited by applicant

International Search Report for International Application No. PCT/US2017/013399, mailed May 26, 2017, 4 Pages. cited by applicant

International Search Report for International Application No. PCT/US2017/021639, mailed Aug. 25, 2017, 5 Pages. cited by applicant

International Search Report for International Application No. PCT/US2018/037924, mailed Nov. 9, 2018, 7 Pages. cited by applicant

Invitation to Pay Additional Fees and, where Applicable, Protest Fee for International Application No. PCT/US2018/037924, mailed Sep. 11, 2018, 3 Pages. cited by applicant

Jiang X., et al., "A Novel EST-derived RNAi Screen Reveals a Critical Role For Farnesyl Diphosphate Synthase in Beta 2-adrenergic Receptor Internalization and Down-regulation," The FASEB Journal, Published Online on Jan. 27, 2012, vol. 26(5), pp. 1995-2007. cited by applicant

Li J., et al., "Reduced Expression of the Mevalonate Pathway Enzyme Farnesyl Pyrophosphate Synthase Unveils Recognition of Tumor Cells by V[gamma]9V[delta]2 T Cells," The Journal of Immunology, US, (Jun. 3, 2009), Jun. 2009, vol. 182, No. 12, pp. 8118-8124, doi: 10.4049/jimmunol.0900101, ISSN 0022-1767, XP055605150. cited by applicant

Li Z., et al., "Inhibition of Farnesyl Pyrophosphate Synthase Prevents Angiotensin II-induced Cardiac Fibrosis In Vitro," Clinical & Experimental Immunology, 2014, vol. 176, pp. 429-437. cited by applicant

Mensali N., et al., "NK Cells Specifically TCR-dressed to Kill Cancer Cells", EBioMedicine, Jan. 2019, vol. 40, pp. 106-117. cited by applicant

Miettinen T.P., et al., "Mevalonate Pathway Regulates Cell Size Homeostasis and Proteostasis Through Autophagy," Cell Reports, Dec. 22, 2015, vol. 13(11), pp. 2610-2620. cited by applicant

Myers E.W., et al., "Optimal Alignments in Linear Space," Cabios, 1988, vol. 4, No. 1, pp. 11-17. cited by applicant

Nada M.H., et al., "Enhancing Adoptive Cancer Immunotherapy with Vγ2Vδ2 T Cells Through Pulse Zoledronate Stimulation," Journal for Immunotherapy of Cancer, Feb. 21, 2017, vol. 5, No. 1, pp. 1-23, DOI: 10.1186/S40425-017-0209-6, XP021242440. cited by applicant

Needleman et al. "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins" Mar. 28, 1970, J. Molecular Biology 48(3):443-453. cited by applicant

Non-Final Office Action for U.S. Appl. No. 15/849,062, mailed Feb. 22, 2018, 05 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 15/850,937, mailed Feb. 22, 2018, 05 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 15/904,131, mailed Jun. 15, 2018, 07 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/008,991, mailed May 7, 2019, 07 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/083,384, mailed Mar. 16, 2020, 9 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/132,247, mailed May 16, 2019, 06 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/182,443, mailed Dec. 31, 2018, 10 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/530,908, mailed Jun. 1, 2020, 6 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/563,738, mailed Mar. 12, 2021, 11 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/614,682, mailed Feb. 28, 2022, 75 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 16/943,800, mailed Nov. 25, 2020, 08 Pages. cited by applicant

Non-Final Office Action for U.S. Appl. No. 17/198,017, mailed Jul. 20, 2021, 7 Pages. cited by applicant

Notice of Allowance for Chinese Application No. 201780017712.6, dated Aug. 25, 2022, 4 Pages. (with English translation). cited by applicant

Notice of Allowance for Japanese Patent Application No. 2018-547354, dated Dec. 17, 2021, 6 Pages. (with English translation). cited by applicant

Notice of Allowance for U.S. Appl. No. 14/706,481, mailed Oct. 13, 2019, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 15/652,080, mailed Nov. 2, 2017, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 15/849,062, mailed Apr. 26, 2018, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 15/850,937, mailed Apr. 23, 2018, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/008,991, mailed Aug. 14, 2019, 5 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/083,384, mailed May 18, 2020, 8 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/132,247, mailed Jul. 19, 2019, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/182,443, mailed Jul. 3, 2019, 3 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/182,443, mailed Jun. 18, 2019, 5 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/530,908, mailed Jul. 10, 2020, 05 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/943,800, mailed Feb. 10, 2021, 7 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 16/988,427, mailed Aug. 26, 2022, 5 pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 17/198,017, mailed Nov. 3, 2021, 6 Pages. cited by applicant

Notice of Allowance for U.S. Appl. No. 17/289,653, mailed Jan. 5, 2022, 9 Pages. cited by applicant

Notice of Final Rejection for Japanese Patent Application No. 2018-536892, dated Nov. 16, 2020, 8 Pages. cited by applicant

Notice of Publication in the PCT Application No. PCT/US2021/020721, dated Sep. 10, 2021. cited by applicant

Notice of Reasons for Refusal for Japanese Patent Application No. 2018-536892, dated Jul. 11, 2022, 21 Pages. cited by applicant

Notice of Reasons for Refusal for Japanese Patent Application No. 2021-045605, dated Apr. 1, 2022, 5 Pages. (with English translation). cited by applicant

Notice of Reasons for Rejection for Japanese Patent Application No. 2018-547354, mailed Feb. 16, 2021, 22 Pages. cited by applicant

Notice of Reasons for Rejection for Japanese Patent Application No. 2021-523916, mailed Jul. 12, 2022, 7 Pages. cited by applicant

Nucleotide: “{Long Control Region} [Human Papillomavirus, type 16, Genomic, 660 nt],” Accession S60559, Publication (online), May 7, 1993 [Retrieved May 9, 2017], 1 Page, XP055455689, Retrieved from URL: [https://www.ncbi.nlm.nih.gov/nucleotide/237343?report=genbank&log\\$=nucldp&blast_rank=1&RID=H3FCKA00014](https://www.ncbi.nlm.nih.gov/nucleotide/237343?report=genbank&log$=nucldp&blast_rank=1&RID=H3FCKA00014). cited by applicant

Office Action for Chinese Application No. 202010396594.8, mailed Jan. 15, 2021, 12 Pages. cited by applicant

Office Action for Chinese Patent Application No. 201780017712.6, dated Nov. 3, 2021, 16 Pages. (with English translation). cited by applicant

Office Action for European Patent Application No. 17739028.3, mailed May 22, 2023, 125 pages. cited by applicant

Office Action for Japanese Application No. 2018-536892, mailed Jun. 26, 2020, 07 Pages. cited by applicant

Office Action for Japanese Patent Application No. 2021045605, mailed Nov. 2, 2022, 8 Pages. cited by applicant

Ostertag D., et al., “Brain Tumor Eradication and Prolonged Survival from Intratumoral Conversion of 5-Fluorocytosine to 5-fluorouracil Using a Nonlytic Retroviral Replicating Vector,” *Neuro-Oncology*, Feb. 2012, vol. 14(2), pp. 145-159. cited by applicant

Pauza C.D., et al., “ $\gamma\delta$ T cells in HIV Disease: Past, Present, and Future,” *Frontiers in Immunology*, Jan. 30, 2015, vol. 5, No. 687, 12 Pages. cited by applicant

Pauza C.D., et al., “Evolution and Function of the TCR Vgamma9 Chain Repertoire: It's Good to be Public,” *Cell Immunology*, Jul. 2015, vol. 296, No. 1, pp. 22-30. cited by applicant

Pearson et al. “Improved Tools for Biological Sequence Comparison” Apr. 1988, *Proceedings of the National Academy of Sciences of the United States of America* 85(8):2444-2448. cited by applicant

Poonia B., et al., “Gamma Delta T Cells From HIV+ Donors Can Be Expanded In Vitro By Zoledronate/Interleukin-2 to Become Cytotoxic Effectors for Antibody-dependent Cellular Cytotoxicity,” *Cytotherapy*, 2012, vol. 14, No. 2, pp. 173-181. cited by applicant

Restriction Requirement for U.S. Appl. No. 16/083,384, mailed Nov. 7, 2019, 8 Pages. cited by applicant

Restriction Requirement for U.S. Appl. No. 16/563,738, mailed Dec. 8, 2020, 6 Pages. cited by applicant

Riano F., et al., “V γ 9V δ 2 TCR-activation by Phosphorylated Antigens Requires Butyrophilin 3 A1 (BTN3A1) and Additional Genes on Human Chromosome 6,” *European Journal of Immunology*, 2014, vol. 44, pp. 2571-2576. cited by applicant

Roden C., et al., "Novel Determinants of Mammalian Primary microRNA Processing Revealed by Systematic Evaluation of Hairpin-Containing Transcripts and Human Genetic Variation," Cold Spring Harbor Laboratory Press, 2017, vol. 27, pp. 374-384, ISSN 1088-9051/17, Retrieved from URL: www.genome.org. cited by applicant

Sambrook J., et al., "Molecular Cloning: A Laboratory Manual," 3rd edition, Cold Spring Harbor Laboratory Press, 2000, 2272 Pages. cited by applicant

Sandstorm A., et al., "The Intracellular B30.2 Domain of Butyrophilin 3A1 Binds Phosphoantigens to Mediate Activation of Hu," *Immunity*, 2014, vol. 40, No. 4, pp. 490-500, DOI: 10.1016/j.immuni.2014.03.003, ISSN 0004789818, XP055481379. cited by applicant

Schiller C.B., et al., "CD19-Specific Triplebody SPM-1 Engages NK and y T Cells for Rapid and Efficient Lysis of Malignant B-Lymphoid Cells," *Oncotarget*, 2016, vol. 7(50), pp. 83392-83408. cited by applicant

Second Office Action for Chinese Application No. 201780017712.6, dated Feb. 3, 2021, 10 Pages. (with English translation). cited by applicant

Selbach M., et al., "Widespread Changes in Protein Synthesis Induced by MicroRNAs," *Nature*, Sep. 4, 2008, vol. 455, pp. 58-63, DOI: 10.1038/nature07228. cited by applicant

Shalova I.N., et al., "CD16 Regulates TRIF-Dependent TLR4 Response in Human Monocytes and Their Subsets", *The Journal of Immunology*, 2012, vol. 188, pp. 3584-3593. cited by applicant

Smith T.F., et al., "Comparison of Biosequences," *Advances in Applied Mathematics*, 1981, vol. 2, pp. 482-489. cited by applicant

Third Office Action for Chinese Application No. 201780017712.6, dated May 14, 2021, 8 Pages. (with English translation). cited by applicant

Tian Y., et al., "MicroRNA-30a Promotes Chondrogenic Differentiation of Mesenchymal Stem Cells Through Inhibiting Delta-like 4 Expression," *Life Science*, Mar. 1, 2016, vol. 148, 36 Pages, DOI:10.1016/J.LFS.2016.02.031, ISSN 0024-3205, XP029460066. cited by applicant

Tokuyama H., et al., "Vy9Vδ2 T Cell Cytotoxicity Against Tumor Cells is Enhanced by Monoclonal Antibody Drugs—Rituximab and Trastuzumab," *International Journal of Cancer*, 2008, vol. 122 (11), pp. 2526-2534. cited by applicant

Tolmachov O.E., et al., "Designing Lentiviral Gene Vectors," *Viral Gene Therapy*, Jul. 20, 2011, Chapter. 13, 23 Pages, ISBN: 978-953-307-539-6, Retrieved from URL: <http://www.intechopen.com/books/viral-gene-therapy/designing-lentiviral-gene-vectors>. cited by applicant

Tracey A., "Human DNA Sequence from Clone RP1-288M22 on Chromosome 6q 12-13, Complete Sequence," National Center for Biotechnology, GenBank Entry, Jan. 24, 2013, pp. 1-34, Retrieved from URL:[https://www.ncbi.nlm.nih.gov/nucleotide/AL035467.23?report=genbank&log\\$=nucldtop&blast_rank=1&RID=UUD4GX2DO14](https://www.ncbi.nlm.nih.gov/nucleotide/AL035467.23?report=genbank&log$=nucldtop&blast_rank=1&RID=UUD4GX2DO14). cited by applicant

Twitty C.G., et al., "Retroviral Replicating Vectors Deliver Cytosine Deaminase Leading to Targeted 5- Fluorouracil-Mediated Cytotoxicity in Multiple Human Cancer Types," *Human Gene Therapy Methods*, Feb. 1, 2016, vol. 27, No. 1, pp. 17-31. cited by applicant

Ueda M., et al., "CD47-Dependent Molecular Mechanisms of Blood Outgrowth Endothelial Cell Attachment on Cholesterol-Modified Polyurethane," *Biomaterials*, Elsevier, Amsterdam, NL, Sep. 1, 2010, vol. 31, No. 25, pp. 6394-6399, ISSN: 0142-9612, XP027102945. cited by applicant

Wang B., et al., "Kinesin Family Member 11 is a Potential Therapeutic Target and is Suppressed by MicroRNA-30a in Breast Cancer," *Molecular Carcinogenesis*, Aug. 2020, vol. 59, No. 8, pp. 908-922. cited by applicant

Wang Y., et al., "Intravenous Delivery of SIRNA Targeting CD47 Effectively Inhibits Melanoma TumorGrmYth and Lung Metastasis," *Molecular Therapy*, Oct. 2013, vol. 21, No. 10, pp. 1919-1929. cited by applicant

Wilkin D.J., et al., "Isolation and Sequence of the Human Farnesyl Pyrophosphate Synthetase eDNA," *The Journal of Biological Chemistry*, Mar. 15, 1990, vol. 265. No. 8, pp. 4607-4614. cited

by applicant

Written Opinion for International Application No. PCT/US2017/013399, mailed May 26, 2017, 8 Pages. cited by applicant

Written Opinion for International Application No. PCT/US2017/021639, mailed Aug. 25, 2017, 7 Pages. cited by applicant

Written Opinion for International Application No. PCT/US2018/037924, mailed Nov. 9, 2018, 11 Pages. cited by applicant

Yang J., et al., "Lentiviral-Mediated Silencing of Famelsy1 Pyrophosphate Synthase through RNA Interference in Mice," Biomed Research International, 2015, vol. 2015, Article ID. 914026, 07 pages. cited by applicant

Ye Y., et al., "Knockdown of Farnesylpyrophosphate Synthase Prevents Angiotensin II-Medicated Cardiac Hypertrophy," The International Journal of Biochemistry & Cell Biology, 2010, vol. 42, pp. 2056-2064. cited by applicant

AU; Examination Report issued in Application No. 2021203836 on Jan. 30, 2024. cited by applicant

EP; Search Report issued in Application No. 23199847.7 on Mar. 5, 2024. cited by applicant
{Long control region} [human papillomavirus, type 16, Genomic, 860 nt]; Accession S60559. Publication [online]. May 7, 1993, [https://www.ncbi.nlm.nih.gov/nucleotide/237343?](https://www.ncbi.nlm.nih.gov/nucleotide/237343?report=genbank&log$=nuclopt&blast_rank=1&RID=H3FCKA00014)

report=genbank&log\$=nuclopt&blast_rank=1&RID=H3FCKA00014; pp. 1. cited by applicant

Bartholome, "Genetics and Biochemistry of the Phenylketonuria-Present State," Human Genetics, vol. 51(3), pp. 241-245, (1979). cited by applicant

Blau et al., "Phenylketonuria," The Lancet, vol. 376(9750), pp. 1417-1427, (2010). cited by applicant

Chandler et al., "Vector Design Influences Hepatic Genotoxicity After Adeno-Associated Virus Gene Therapy," Journal of Clinical Investigation, vol. 125(2), pp. 870-880, (2015). cited by applicant

Charron et al., "Dominant-Negative Interference in the Pahenu2 Mouse Model of PKU: Effectiveness of Vectors Expressing Either Modified Forms of Phenylalanine Hydroxylase (PAH) or Ribozymes Plus a Hardened PAH mRNA," Molecular Therapy, vol. 11, pp. S163-S164, (2005). cited by applicant

Charron, "Gene Therapy for Phenylketonuria: Dominant-Negative Interference in a Recessive Disease," Dissertation, University of Florida 2005, http://etd.fcla.edu/UF/UFE0011392/charron_c.pdf, (retrieved Jul. 26, 2018) (2005). cited by applicant

Christophersen et al., "A Technique of Transumbilical Portal Vein Catheterization in Adults," The Archives of Surgery, vol. 95(6), pp. 960-963, (1967). (Abstract Only). cited by applicant

Condiotti et al., "Prolonged Liver-Specific Transgene Expression by a Non-Primate Lentiviral Vector," Biochemical and Biophysical Research Communications, vol. 320(3), pp. 998-1006, (Jul. 30, 2004). cited by applicant

Daryl S. Schiller, "Parameters Influencing Measurement of the Gag Antigen-Specific T-Proliferative Response to HIV Type 1 Infection," AIDS Research and Human Retroviruses, vol. 16, No. 3, pp. 259-271, (2000). cited by applicant

Ding et al., "Administration-Route and Gender-Independent Longterm Therapeutic Correction of Phenylketonuria (PKU) in a Mouse Model by Recombinant Adeno-Associated Virus 8 Pseudotyped Vector-Mediated Gene Transfer," Gene Therapy, vol. 13, pp. 583-587, (Dec. 1, 2005). cited by applicant

Donsante et al., "AAV Vector Integration Sites in Mouse Hepatocellular Carcinoma," Science, vol. 317(5837), p. 477, (2007). cited by applicant

Eisensmith et al., "Multiple Origins for Phenylketonuria in Europe," American Journal of Human Genetics, vol. 51(6), pp. 1355-1365, (1992). cited by applicant

EP; Supplementary Search Report in the EP Application No. 18781288.8 dated Dec. 8, 2020. cited by applicant

Fisher et al., "The Inhibition of Phenylalanine and Tyrosine Hydroxylases by High Oxygen Levels," *Journal of Neurochemistry*, vol. 19(5), pp. 1359-1365, (1972). (Abstract Only). cited by applicant

Fusetti, et al., "Structure of Tetrameric Human Phenylalanine Hydroxylase and Its Implications for Phenylketonuria," *J. Bio. Chem.*, vol. 273, No. 27, p. 16962-16967 (1998). cited by applicant
GenBank Accession No. JG619773, MNESC1NG-T3-001_L15_6FEB2009_054 MNESC1NG cell culture from Mahonia nervosa Berberis nervosa cDNA, mRNA sequence, Feb. 13, 2014 (online). [Retrieved on Dec. 5, 2017]. Retrieved from the internet:<URL:

<https://www.ncbi.nlm.nih.gov/nucest/JG619773> > entire document. cited by applicant

GenBank Sequence M65141.1 Retrieved from the Internet <URL:

<https://www.ncbi.nlm.nih.gov/nuccore/M65141.1>. Especially Sequence, nt 301-420, (Retrieved Mar. 31, 2019). cited by applicant

Gober et al., "Human T Cell Receptor $\gamma\delta$ Cells Recognize Endogenous Mevalonate Metabolites in Tumor Cells," *J. of Experimental Med.*, Jan. 20, 2003, vol. 197, pp. 163-168. cited by applicant

Grisch-Chan et al., "Low-Dose Gene Therapy for Murine PKU Using Episomal Naked DNA Vectors Expressing PAH from Its Endogenous Liver Promoter," *Molecular Therapy Nucleic Acids*, vol. 7, pp. 339-349, (2017). cited by applicant

Guldborg et al., "Aberrant Phenylalanine Metabolism in Phenylketonuria Heterozygotes," *Journal of Inherited Metabolic Disease*, vol. 21(4), pp. 365-372, (1998). cited by applicant

Hafid et al., "Phenylketonuria: A Review of Current and Future Treatments," *Translational Pediatrics*, vol. 4(4), pp. 304-317, (2015). cited by applicant

Harding et al., "Complete correction of hyperphenylalaninemia following liver-directed, recombinant AAV2/8 vector-mediated gene therapy in murine phenylketonuria", *Gene Ther.*, Mar. 2006, 13(5):457-462. cited by applicant

Hee Yeon Kim., "Farnesyl diphosphate synthase is important for the maintenance of glioblastoma stemness," *Experimental & Molecular Medicine*, (2018). cited by applicant

Ho et al., *Translational Pediatrics*, 2014, 3(2): 49-62. (Year 2014. cited by applicant

Hong Wang., "Indirect Stimulation of Human V2V2 Cells Through Alterations in Isoprenoid Metabolism," *The Journal of Immunology*, (2011). cited by applicant

Human papillomavirus type 16 (HPV16), complete genome; GenBank: K02718.1; Publication [online], Mar. 18, 1994, [https://www.ncbi.nlm.nih.gov/nucleotide/333031?](https://www.ncbi.nlm.nih.gov/nucleotide/333031?report=genbank&log$=nuclopt&blast_rank=22&RID=H3E1THFU014)

[report=genbank&log\\$=nuclopt&blast_rank=22&RID=H3E1THFU014](https://www.ncbi.nlm.nih.gov/nucleotide/333031?report=genbank&log$=nuclopt&blast_rank=22&RID=H3E1THFU014); pp. 1-4. cited by applicant
JP Office Action in Japanese Application No. 2019-554397, dated Nov. 21, 2022, 8 pages (with English translation). cited by applicant

JP Office Action in Japanese Application No. 2020-518812, dated Aug. 25, 2022, 13 pages (with English translation). cited by applicant

Kaufman et al., "A Model of Human Phenylalanine Metabolism in Normal Subjects and in Phenylketonuric Patients," *Proceedings of the National Academy of Sciences USA*, vol. 96(6), pp. 3160-3164, (1999). cited by applicant

Kaufman et al., "Phenylalanine Hydroxylase Activity in Liver Biopsies from Hyperphenylalaninemia Heterozygotes: Deviation from Proportionality with Gene Dosage," *Pediatric Research*, vol. 9(8), pp. 632-634, (1975). cited by applicant

Ledley et al., "Retroviral-mediated gene transfer of human phenylalanine hydroxylase into NIH 3T3 and hepatoma cells", *Proceedings of the National Academy of Sciences*, vol. 83, No. 2. (Jan. 1, 1986), pp. 409-413, XP002583115. cited by applicant

Longo et al., "Single-Dose, Subcutaneous Recombinant Phenylalanine Ammonia Lyase Conjugated with Polyethylene Glycol in Adult Patients with Phenylketonuria: An Open-Label, Multicentre, Phase 1 Dose-Escalation Trial," *The Lancet*, vol. 384(9937), pp. 37-44, (2014). cited by applicant

Lu et al., “Anti-sense-Mediated Inhibition of Human Immunodeficiency Virus (HIV) Replication by Use of an HIV Type 1-Based Vector Results in Severely Attenuated Mutants Incapable of Developing Resistance,” *Journal of Virology*, vol. 79, No. 13, pp. 7079-7088 (Jul. 2004). cited by applicant

Makinen et al., “stable RNA Interference: Comparison of U6 and H1 Promoters in Endothelial Cells and in Mouse Brain,” *Journal of Gene Medicine*, vol. 8, pp. 433-441, 2006. cited by applicant

Mochizuki et al., “Long-Term Correction of Hyperphenylalaninemia by AAV-Mediated Gene Transfer Leads to Behavioral Recovery in Phenylketonuria Mice,” *Gene Therapy*, vol. 11(13), pp. 1081-1086, (2004). cited by applicant

Moser et al., “ $\gamma\delta$ T cells: novel initiators of adaptive immunity,” *Immunological Reviews*, vol. 215, pp. 89-102 (Feb. 2, 2007). cited by applicant

Nault et al., “Adeno-Associated Virus Type 2 as an Oncogenic Virus in Human Hepatocellular Carcinoma,” *Molecular & Cellular Oncology*, vol. 3(2), p. e1095271, (2016). cited by applicant

Nowacki et al., “The PAH Mutation Analysis Consortium Database: Update 1996,” *Nucleic Acid Research*, vol. 25(1), pp. 139-142, (Jan. 1, 1997). cited by applicant

Oh et al. “Lentiviral Vector Design Using Alternative RNA Export Elements,” *Retrovirology*, vol. 4:38, pp. 1-10, (2007). cited by applicant

Oh et al., “Long-Term Enzymatic and Phenotypic Correction in the Phenylketonuria Mouse Model by Adeno-Associated Virus Vector-Mediated Gene Transfer,” *Pediatric Research*, vol. 56(2), pp. 278-284, (2004). cited by applicant

Oh et al., “Reversal of Gene Expression Profile in the Phenylketonuria Mouse Model After Adeno-Associated Virus Vector-Mediated Gene Therapy,” *Molecular Genetics and Metabolism*, vol. 86(Supp. 1), pp. S124-132, (2005). cited by applicant

Pan et al., “Biodistribution and Toxicity Studies of VSVG-Pseudotyped Lentiviral Vector After Intravenous Administration in Mice with the Observation of in Vivo Transduction of Bone Marrow,” *Molecular Therapy*, vol. 6(1), pp. 19-29, (2002). cited by applicant

PCT; International Preliminary Report on Patentability dated Oct. 8, 2019 in the Application No. PCT/ US2018/025733. cited by applicant

PCT; International Search Report dated Apr. 12, 2019 in Application No. PCT/ US2018/053919. cited by applicant

PCT; International Search Report dated Sep. 24, 2018 in Application No. PCT/US2018/025733. cited by applicant

PCT; Invitation to Pay Additional Fees in Application No. PCT/US2018/053919 dated Feb. 22, 2019. cited by applicant

PCT; Written Opinion dated Apr. 12, 2019 in Application No. PCT/US2018/053919. cited by applicant

PCT; Written Opinion dated Sep. 24, 2018 in Application No. PCT/US2018/025733. cited by applicant

Shedlovsky et al., “Mouse Models of Human Phenylketonuria,” *Genetics*, vol. 134(4), pp. 1205-1210, (1993). cited by applicant

Stunkel et al., “The Chromatin Structure of the Long Control Region of Human Papillomavirus Type 16 Repress Viral Oncoprotein Expression,” *Journal of Virology*, vol. 73, No. 3, pp. 1918-1930 (Mar. 1999). cited by applicant

Thompson et al., “Alkylamines cause V γ 9V δ 2 T-cell activation and proliferation by inhibiting the mevalonate pathway,” *Blood*, Jan. 15, 2006, vol. 107, pp. 651-654. cited by applicant

US Non-Final Office Action in U.S. Appl. No. 16/494,196, dated Dec. 6, 2022, 51 pages. cited by applicant

US Restriction Requirement in U.S. Appl. No. 16/652,867, dated Sep. 9, 2022, 9 pages. cited by applicant

USPTO; Invitation to Pay Additional Fees And, Where Applicable, Protest Fee dated Jul. 17, 2018

in Application No. PCT/US2018/25733. cited by applicant
Wang et al., “Butyrophilin 3A1 Plays an Essential Role in Prenyl Pyrophosphate Stimulation of Human Vg2Vd2 T Cells,” Journal of Immunology, vol. 191(3), pp. 1029-1042, (Jul. 5, 2013). cited by applicant
Yagi et al., “Complete Restoration of Phenylalanine Oxidation in Phenylketonuria Mouse by a Self-Complementary Adeno-Associated Virus Vector,” Journal of Gene Medicine, vol. 13(2), pp. 114-122, (2011). cited by applicant
Yano et al., “Evaluation of Tetrahydrobiopterin Therapy with Large Neutral Amino Acid Supplementation in Phenylketonuria: Effects on Potential Peripheral Biomarkers, Melatonin and Dopamine, for Brain Monoamine Neurotransmitters,” PLoS One, vol. 11(8), p. e0160892, (2016). cited by applicant
Zhaobing Ding et al., “Liver -Directed, AAV-and Lentivirus-Mediated Gene Therapy in the Phenylketonuria Mouse Model Pah-enu2”, Molecular Therapy, vol. 11, Supp. 1. (May 2005) XP055751452. cited by applicant
Thompson et al., “Alkylamines cause Vγ9Vδ2 T-cell activation and proliferation by inhibiting the mevalonate pathway,” Blood, Jan. 15, 2006, vol. 107, pp. 651-654. cited by applicant
CN Office Action in Chinese Application No. 201880039828.4, dated Mar. 1, 2023, 19 pages (with English translation). cited by applicant
JP Notice of Allowance in Japanese Application No. 2018-536892, dated Mar. 29, 2023, 4 pages (with English translation). cited by applicant
JP Office Action in Japanese Application No. 2019-569226, dated Mar. 20, 2023, 5 pages (with English translation). cited by applicant
JP Office Action in Japanese Application No. 2021-523916, dated Apr. 18, 2023. cited by applicant
JP Office Action in Japanese Application No. 2021-045605, dated Apr. 19, 2023. cited by applicant
CA Office Action in Canadian Application No. 3,011,529, dated Feb. 21, 2023, 7 pages. cited by applicant
JP Office Action in Japanese Application No. 2018-536892, dated Jan. 30, 2023, 4 pages (with English translation). cited by applicant

Primary Examiner: McGarry; Sean

Attorney, Agent or Firm: COOLEY LLP

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a continuation of U.S. patent application Ser. No. 16/988,427 filed on Aug. 7, 2020 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS”, which is a continuation of U.S. patent application Ser. No. 16/530,908 filed on Aug. 2, 2019 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS”, which is a continuation of U.S. patent application Ser. No. 16/132,247 filed on Sep. 14, 2018 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS,” which is a continuation of U.S. patent application Ser. No. 15/904,131 filed on Feb. 23, 2018 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS”, which is a continuation in part of U.S. patent application Ser. No. 15/652,080 filed on Jul. 17, 2017 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS”, which is a continuation of International Application No. PCT/US17/13399 filed on Jan. 13, 2017 entitled “METHODS AND COMPOSITIONS FOR THE

ACTIVATION OF GAMMA-DELTA T-CELLS”, which claims priority to U.S. Provisional Patent Application No. 62/279,474 filed on Jan. 15, 2016 entitled “METHODS AND COMPOSITIONS FOR THE ACTIVATION OF GAMMA-DELTA T-CELLS”, the disclosures of which are incorporated herein by reference.

SEQUENCE LISTING

(1) A Sequence Listing is enclosed with this application and is incorporated herein by reference. The text file of the Sequence Listing is named 436313003138_SL.xml and the file size is 119 kilobytes.

FIELD OF THE INVENTION

(2) The present disclosure relates generally to the fields of gene therapy and immunotherapy, specifically in relation to increased activation and effector cell function of gamma delta (“GD”) T cells.

BACKGROUND

(3) Human T cells are distinguished on the basis of T cell receptor structure. The major populations, including CD4+ and CD8+ subsets, express a receptor composed of alpha and beta chains. A smaller subset expresses T cell receptor made from gamma and delta chains. Gamma delta (“GD”) T cells make up 3-10% of circulating lymphocytes, and a Vδ2+ subset makes up 75% of GD T cells in blood. Vδ2+ cells recognize non-peptide epitopes and do not require antigen presentation by major histocompatibility complexes (“MHC”) or human leukocyte antigen (“HLA”). The majority of Vδ2+ T cells also express a Vγ9 chain and are stimulated by exposure to 5-carbon pyrophosphate compounds that are intermediates in mevalonate and non-mevalonate sterol/isoprenoid synthesis pathways. The response to isopentenyl pyrophosphate (5-carbon) is universal among healthy human beings.

(4) Another subset of GD T cells, Vδ1+, make up a much smaller percentage of the T cells circulating in the blood, but Vδ1+ cells are commonly found in the epithelial mucosa and the skin.

(5) In general, GD T cells have several functions, including killing tumor cells and pathogen-infected cells. Stimulation through their unique T cell receptor (“TCRs”) composed of two glycoprotein chains, γ and δ, improves the capacity for cellular cytotoxicity, cytokine secretion and other effector functions. The TCRs of GD T cells have unique specificities and the cells themselves occur in high clonal frequencies, thus allowing rapid innate-like responses to tumors and pathogens.

(6) Bisphosphonate drugs and other inhibitors of farnesyl diphosphate synthase (“FDPS”), which are downstream from isopentenyl pyrophosphate (“IPP”) in the mevalonate pathway (see, for e.g., FIG. 1), have been used to treat various diseases, including cancers, specifically those involving bone metastasis. Bisphosphonate drugs include, for example, Zometa® (Novartis) and Fosamax® (Merck).

(7) Certain bisphosphonates have also been investigated for stimulation of GD T cells. This may be because when FDPS is inhibited in myeloid cells, IPP begins to accumulate and geranylgeranyl pyrophosphate (“GGPP”), a downstream product of FDPS that suppresses activation of the inflammasome pathway, is reduced. The reduction in GGPP removes an inhibitor of the caspase-dependent inflammasome pathway and allows secretion of mature cytokines including interleukin-beta and interleukin-18, the latter being especially important for gamma delta T cell activation.

(8) Thus, when FDPS is blocked, the increased IPP and decreased GGPP combine to activate Vδ2+ T cells. Vδ2+ cells activated by IPP or bisphosphonates will proliferate rapidly, express a number of cytokines and chemokines, and can function to cytotoxically destroy tumor cells or cells infected with pathogenic microorganisms.

(9) However, bisphosphonates are associated with inflammation and osteonecrosis, as well as having poor bioavailability due to their chemistry. Likewise, IPP has a very short half-life and is difficult to synthesize. Both types of compounds require systemic administration in an individual.

Accordingly, both bisphosphonates in general, and IPP specifically, leave a great deal to be desired for therapeutic purposes involving activation of GD T cells.

SUMMARY OF THE INVENTION

(10) In one aspect, a method of activating a GD T cell is provided. The method includes infecting, in the presence of the GD T cell, a target cell with a viral delivery system that encodes at least one genetic element. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In embodiments, the enzyme is FDPS. In embodiments, when the enzyme is inhibited in the target cell, the target cell subsequently activates the GD T cell. In embodiments, the target cell is a cancer cell or a cell that has been infected with an infectious agent. In a preferred embodiment, the activation of the GD T cell results in the GD T cell killing the cancer cell or the cell infected with an infectious agent. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA. In further embodiments, the target cell is also contacted with a bisphosphonate drug. In embodiments, the bisphosphonate drug is zoledronic acid.

(11) In another aspect, a method of treating cancer in a subject is provided. The method includes administering to the subject a therapeutically-effective amount of a viral delivery system that encodes at least one genetic element. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In further embodiments, when the enzyme is inhibited in a cancer cell in the presence of a GD T cell, the cancer cell activates the GD T cell, to thereby treat the cancer. In embodiments, the enzyme is FDPS. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA. In further embodiments, the target cell is also contacted with a bisphosphonate drug. In embodiments, the bisphosphonate drug is zoledronic acid.

(12) In another aspect, a method of treating an infectious disease in a subject is provided. The method includes administering to the subject a therapeutically-effective amount of a viral delivery system that encodes at least one genetic element. In embodiments, the at least one genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In further embodiments, when the enzyme is inhibited in a cell that is infected with an infectious agent in the presence of a GD T cell, the infected cell activates the GD T cell, to thereby treat the infected cell, and the infectious disease. In embodiments, the enzyme is FDPS. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA. In further embodiments, the target cell is also contacted with a bisphosphonate drug. In embodiments, the bisphosphonate drug is zoledronic acid.

(13) In another aspect, the at least one encoded genetic element includes a shRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with GTCCTGGAGTACAATGCCATTCTCGAGAATGGCATTGTACTCCAGGACTTTTT (SEQ ID NO: 1); GCAGGATTTTCGTTTCAGCACTTCTCGAGAAGTGCTGAACGAAATCCTGCTTTTT (SEQ ID NO: 2); GCCATGTACATGGCAGGAATTCTCGAGAA TTCCTGCCATGTACATGGCTTTTT (SEQ ID NO: 3); or GCAGAAGGAGGCTGA GAAAGTCTCGAGACTTTCTCAGCCTCCTTCTGCTTTTT (SEQ ID NO: 4). In a preferred embodiment, the shRNA includes GTCCTGGAGTACAATGCCATTCTCGAG AATGGCATTGTACTCCAGGACTTTTT (SEQ ID NO: 1); GCAGGATTTTCGTTTC ACACTTCTCGAGAAGTGCTGAACGAAATCCTGCTTTTT (SEQ ID NO: 2); GCCA TGTACATGGCAGGAATTCTCGAGAATTCCTGCCATGTACATGGCTTTTT (SEQ ID NO: 3); or GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCCTCCTT CTGCTTTTT (SEQ ID NO: 4).

(14) In another aspect, the at least one encoded genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGCGTGAAGC CACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGCCTCGGACTTCAAGGG

GCT (SEQ ID NO: 5); AAGGTATATTGCTGTTGACAGTGAGCGACACT
TTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGGCTGAGAAAGTGCTGC
CTACTGCCTCGGACTTCAAGGGGCT (SEQ ID NO: 6); TGCTGTTGACAGTG
AGCGACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGAGGCTGAG
AAAGTTGCCTACTGCCTCGGA (SEQ ID NO: 7); CCTGGAGGCTTGCTGAAG
GCTGTATGCTGACTTTCTCAGCCTCCTTCTGCTTTTGGCCACTGACTGAGCAGAAGGG
CTGAGAAAGTCAGGACACAAGGCCTGTTACTAGCACTCA (SEQ ID NO: 8);
CATCTCCATGGCTGTACCACCTTGTCTGGGACTTTCTCAGCCTCCTTCTGCCTGTTGAA
TCTCATGGCAGAAGGAGGCGAGAAAGTCTGACATTTTGGTATCTTTCATCTGACCA
(SEQ ID NO: 9); or GGGCCTGGCTCGAGCAGGGGGGCGAGGGATACTTTCT
CAGCCTCCTTCTGCTGGTCCCCCTCCCCGCAGAAGGAGGCTGAGAAAGTCCTTCCCTC
CCAATGACCGCGTCTTCGTCG (SEQ ID NO: 10). In a preferred embodiment, the microRNA
includes AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCT
CCTTCTGCGTGAAGCCACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGC
CTCGGACTTCAAGGGGCT (SEQ ID NO: 5); AAGGTATATTGCTGTTGACAGT
GAGCGACACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGGGCTGA
GAAAGTGCTGCCTACTGCCTCGGACTTCAAGGGGCT (SEQ ID NO: 6); TGCTG
TTGACAGTGAGCGACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGG
AGGCTGAGAAAGTTGCCTACTGCCTCGGA (SEQ ID NO: 7); CCTGGAGGCT
TGCTGAAGGCTGTATGCTGACTTTCTCAGCCTCCTTCTGCTTTTGGCCACTGACTGAG
CAGAAGGGCTGAGAAAGTCAGGACACAAGGCCTGTTACTAGCACTCA (SEQ ID NO: 8);
CATCTCCATGGCTGTACCACCTTGTCTGGGACTTTCTCAGCCTCCTT
CTGCCTGTTGAATCTCATGGCAGAAGGAGGCGAGAAAGTCTGACATTTTGGTATCTT
TCATCTGACCA (SEQ ID NO: 9); or GGGCCTGGCTCGAGCAGGGGGGCGAGGG
ATACTTTCTCAGCCTCCTTCTGCTGGTCCCCCTCCCCGCAGAAGGAGGCTGAGAAAGT
CCTTCCCTCCCAATGACCGCGTCTTCGTCG (SEQ ID NO: 10).

(15) In another aspect, a viral vector comprising at least one encoded genetic element is provided. The at least one encoded genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In embodiments, the enzyme involved in the mevalonate pathway is farnesyl diphosphate synthase (FDPS). In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA.

(16) In another aspect, the at least one encoded genetic element includes a shRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with v In a preferred embodiment, the shRNA includes SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4.

(17) In another aspect, the at least one encoded genetic element includes a microRNA having at least 80%, or at least 85%, or at least 90%, or at least 95% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In a preferred embodiment, the microRNA includes SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10.

(18) In embodiments, the viral vector is comprised of any vector that can effectively transduce the small RNA into a target cell. In embodiments, the viral vector is a lentiviral vector. In other embodiments, the viral vector is an adeno-associated virus vector.

(19) In another aspect, the viral vector includes a second encoded genetic element. In embodiments, the second genetic element includes at least one cytokine or chemokine. In embodiments, the at least one cytokine is selected from the group consisting of: IL-18, TNF- α , interferon- γ , IL-1, IL-2, IL-15, IL-17, and IL-12. In embodiments, the at least one chemokine is a CC chemokine or a CXC chemokine. In further embodiments, the at least one chemokine is RANTES.

(20) In another aspect, a lentiviral vector system for expressing a lentiviral particle is provided. The

system includes a lentiviral vector, at least one envelope plasmid for expressing an envelope protein optimized for infecting a cell; and at least one helper plasmid for expressing gag, pol, and rev genes. When the lentiviral vector, the at least one envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell, a lentiviral particle is produced by the packaging cell. In embodiments, the lentiviral particle is capable of infecting a targeting cell, and inhibiting an enzyme involved in the mevalonate pathway within the target cell. In embodiments, the enzyme involved in the mevalonate pathway is FDPS. In embodiments, the lentiviral vector system includes a first helper plasmid for expressing the gag and pol genes, and a second helper plasmid for expressing the rev gene. In embodiments, the envelope protein is preferably optimized for infecting a target cell. In embodiments, the target cell is a cancer cell. In other embodiments, the target cell is a cell that is infected with an infectious agent.

(21) In another aspect a pharmaceutical combination is disclosed which includes a bisphosphonate compound; and a lentiviral particle produced by a packaging cell and capable of infecting a target cell. The lentiviral particle comprises an envelope protein capable of infecting the target cell, and: at least one encoded shRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded shRNA comprises a sequence having at least 80% percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4; or at least one encoded microRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded microRNA comprises a sequence having at least 80% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10, wherein the pharmaceutical combination is at least one of fixed and non-fixed. In embodiments, the at least one encoded shRNA comprises a sequence having at least 85% or at least 90% or at least 95% percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4; or the at least one encoded microRNA comprises a sequence having at least 85% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the at least one encoded shRNA comprises SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4 or the at least one encoded microRNA comprises SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the pharmaceutical composition comprises a fixed combination. In embodiments, the pharmaceutical composition comprises a non-fixed combination. In embodiments, the bisphosphonate drug comprises zoledronic acid. In embodiments, the bisphosphonate drug and the lentiviral particle are present in synergistically effective amounts. In embodiments, the target cell is one or more cancer cells that are present in a cancer selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof. In embodiments, the target cell is one or more cancer cells that are present in a hepatocellular carcinoma. In embodiments, the target cell is capable of activating a gamma delta T cell following infection with the lentiviral particle. In embodiments, the enzyme is FDPS.

(22) In another aspect, a method of treating a cancer in a subject using an immunotherapy-based composition is disclosed. The method includes administering a therapeutically-effective amount of a bisphosphonate drug to the subject; and administering a therapeutically-effective amount of the immunotherapy-based composition to the subject, wherein the immunotherapy-based composition comprises a lentiviral particle. The lentiviral particle comprises an envelope protein capable of infecting one or more cancer cells, and at least one encoded shRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded shRNA comprises a sequence having at least 80% percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4 or at least one encoded microRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded microRNA comprises a sequence having at least 80% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the at least one encoded shRNA comprises a sequence having at least 85% or at least 90% or at least 95%

percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4, or at least one encoded microRNA comprises a sequence having at least 85% or at least 90% or at least 95% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the at least one encoded shRNA comprises SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4. In embodiments, the at least one encoded microRNA comprises SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the one or more cancer cells are present in a cancer selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof. In embodiments, the bisphosphonate drug comprises zoledronic acid. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in a fixed combination. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in a non-fixed combination. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered simultaneously. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered sequentially. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in synergistically effective amounts. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered at a synergistically effective time interval. In embodiments, the one or more cancer cells are capable of activating a gamma delta T cell resident in the subject following infection of the one or more cancer cells with the immunotherapy-based composition. In embodiments, activating the gamma delta T cell comprises increasing tumor necrosis factor (TNF)- α expression by the gamma delta T cell. In embodiments, activating the gamma delta T cell comprises increasing expression and/or secretion of cytokines, chemokines, and/or cell death ligands including but not limited to FasL and TRAIL. In embodiments, the enzyme of the mevalonate pathway is farnesyl diphosphate synthase (FDPS).

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) FIG. 1 depicts an overview of the major steps in the mevalonate pathway for biosynthesis of steroids and isoprenoids.
- (2) FIG. 2 depicts an exemplary 3-vector lentiviral vector system in a circularized form.
- (3) FIG. 3 depicts an exemplary 4-vector lentiviral vector system in a circularized form.
- (4) FIG. 4 depicts: (A) a linear map of a lentiviral vector expressing a FDPS shRNA targeting sequence; and (B) a linear map of a lentiviral vector expressing a synthetic microRNA with a FDPS targeting sequence.
- (5) FIG. 5 depicts data demonstrating activation of V δ 2⁺ T cells THP-1 leukemia cells with a lentivirus expressing FDPS shRNA #4 (SEQ ID NO: 4), as described herein.
- (6) FIG. 6 depicts data demonstrating activation of V δ 2⁺ T cells by THP-1 leukemia cells with a lentivirus expressing FDPS shRNA #4 (SEQ ID NO: 4), as described herein.
- (7) FIG. 7 depicts data demonstrating activation of V δ 2⁺ T cells by PC3 prostate carcinoma cells with a lentivirus expressing FDPS shRNA #1 (SEQ ID NO: 1), as described herein.
- (8) FIG. 8 depicts data demonstrating activation of V δ 2⁺ T cells by PC3 prostate carcinoma cells with a lentivirus expressing FDPS shRNA #4 (SEQ ID NO: 4), as described herein.
- (9) FIG. 9 depicts data demonstrating activation of V δ 2⁺ T cells by HepG2 carcinoma cells with a lentivirus expressing FDPS shRNA #1 (SEQ ID NO: 1) or FDPS shRNA #4 (SEQ ID NO: 4), as described herein.
- (10) FIG. 10 depicts data demonstrating activation of V δ 2⁺ T cells by THP-1 leukemia cells with a lentivirus expressing miR30 FDPS #1 (SEQ ID NO: 5), as described herein.

(11) FIG. 11 depicts data demonstrating the percent of specific lysis versus an E:T ratio for a variety of experimental conditions, as described herein.

(12) FIG. 12 depicts data demonstrating lentiviral-delivered shRNA-based RNA interference targeting the human FDPS gene.

(13) FIG. 13 depicts data demonstrating lentiviral-delivered miR-based RNA interference targeting the human FDPS gene.

(14) FIG. 14 depicts data demonstrating activation of Vδ2+ T cells by HepG2 carcinoma cells with an adeno-associated virus expressing FDPS shRNA #4 (SEQ ID NO: 4), as described herein.

(15) FIG. 15 depicts immunoblot data demonstrating lack of RAP1 prenylation in the cells transduced with LV-shFDPS and treated with zoledronic acid.

DETAILED DESCRIPTION

(16) Overview of Disclosure

(17) The present disclosure relates to gene therapy constructs and delivery of the same to cells, resulting in suppression of Farnesyl diphosphate synthase (“FDPS”), which is necessary to convert isopentenyl phosphate (IPP) to farnesyl diphosphate (FDP), as shown, for example, in FIG. 1. In embodiments, one or more viral vectors are provided with microRNAs or short hairpin RNAs (shRNA) that target FDPS, thereby reducing expression levels of this enzyme. The viral vectors include lentiviral vectors and AAV vectors. A consequence of modulating expression of FDPS is to increase the accumulation of IPP, which is a stimulator of GD T cell proliferation and differentiation. Accordingly, the constructs provided herein are used to activate GD T cells, and are used to treat cancers and infectious diseases.

(18) Definitions and Interpretation

(19) Unless otherwise defined herein, scientific and technical terms used in connection with the present disclosure shall have the meanings that are commonly understood by those of ordinary skill in the art. Further, unless otherwise required by context, singular terms shall include pluralities and plural terms shall include the singular. Generally, nomenclature used in connection with, and techniques of, cell and tissue culture, molecular biology, immunology, microbiology, genetics and protein and nucleic acid chemistry and hybridization described herein are those well-known and commonly used in the art. The methods and techniques of the present disclosure are generally performed according to conventional methods well-known in the art and as described in various general and more specific references that are cited and discussed throughout the present specification unless otherwise indicated. See, e.g.: Sambrook J. & Russell D. *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (2000); Ausubel et al., *Short Protocols in Molecular Biology: A Compendium of Methods from Current Protocols in Molecular Biology*, Wiley, John & Sons, Inc. (2002); Harlow and Lane *Using Antibodies: A Laboratory Manual*; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1998); and Coligan et al., *Short Protocols in Protein Science*, Wiley, John & Sons, Inc. (2003). Any enzymatic reactions or purification techniques are performed according to manufacturer's specifications, as commonly accomplished in the art or as described herein. The nomenclature used in connection with, and the laboratory procedures and techniques of, analytical chemistry, synthetic organic chemistry, and medicinal and pharmaceutical chemistry described herein are those well-known and commonly used in the art.

(20) As used in the description and the appended claims, the singular forms “a”, “an” and “the” are used interchangeably and intended to include the plural forms as well and fall within each meaning, unless the context clearly indicates otherwise. Also, as used herein, “and/or” refers to and encompasses any and all possible combinations of one or more of the listed items, as well as the lack of combinations when interpreted in the alternative (“or”).

(21) All numerical designations, e.g., pH, temperature, time, concentration, and molecular weight, including ranges, are approximations which are varied (+) or (−) by increments of 0.1. It is to be understood, although not always explicitly stated that all numerical designations are preceded by

the term “about”. The term “about” also includes the exact value “X” in addition to minor increments of “X” such as “X+0.1” or “X-0.1.” It also is to be understood, although not always explicitly stated, that the reagents described herein are merely exemplary and that equivalents of such are known in the art.

(22) As used herein, the term “about” will be understood by persons of ordinary skill in the art and will vary to some extent depending upon the context in which it is used. If there are uses of the term which are not clear to persons of ordinary skill in the art given the context in which it is used, “about” will mean up to plus or minus 10% of the particular term.

(23) As used herein, the terms “administration of” or “administering” refer to providing an active agent to a subject in need of treatment in a form that can be introduced into that individual's body in a therapeutically useful form and therapeutically effective amount.

(24) As used herein, the terms “bisphosphonates” and “bisphosphonate drugs” refer to therapeutic agents of various embodiments, and encompass any of aminobisphosphonates, diphosphonates, biposphonic acids, and diphosphonic acids, as well as pharmaceutically acceptable salts and derivatives thereof. The use of a specific nomenclature in referring to bisphosphonates is not meant to limit the scope of the present invention, unless specifically indicated.

(25) As used herein, the terms “co-administration” or “combined administration” or “combined use” or “combination therapy” or the like as utilized herein refer to administration of a therapeutic vector or a lentiviral particle and a bisphosphonate drug or a therapeutic vector or a lentiviral particle and an antibody or a therapeutic vector or a lentiviral particle and a bisphosphonate drug and an antibody to a single subject in need thereof (e.g., a patient), and are intended to include treatment regimens in which the agents are not necessarily administered by the same route of administration and/or at the same time.

(26) As used herein, the term “fixed combination,” refers to two or more active ingredients or components, including any of their respective compositions, formulations or drug forms, e.g., a therapeutic vector or a lentiviral particle and a bisphosphonate drug or any combination of these, that are administered essentially in combination to a patient, for example essentially simultaneously, in the form of a single entity or dosage or combined entities or dosages, e.g., in one tablet or in one capsule or in combined tablets or capsules or combined liquid forms.

(27) As used herein, the term “non-fixed combination,” refers to two or more active ingredients or components, including any of their respective compositions, formulations or drug forms, e.g., a therapeutic vector or a lentiviral particle and a bisphosphonate drug or any combination of these, that are administered in combination to a patient as separate entities either simultaneously, concurrently or sequentially with no specific time limits, wherein such administration provides therapeutically effective levels of the active components in the patient. The non-fixed combination can be dosed independently of each other or by use of different fixed combinations e.g., simultaneously or at different time points. The active components may be administered as separate pharmaceutical dosage forms or pharmaceutical formulations that may be, for example, sold independently of each other, with or without label instructions concerning the possibility of a combined use. Such instructions may be provided in the package equipment, e.g., leaflet or the like, or in other information, e.g., provided to physicians and medical staff. A non-fixed combination, its respective active ingredients or components, including any of their respective compositions, formulations or drug forms, or the parts thereof, can be administered simultaneously or chronologically staggered, e.g., at different time points and with equal or different time intervals for any part of the administration. Such time intervals may be chosen such that the effect on the treated disease, when treated in combination, is more effective than would be obtained by use of only any one of the active components.

(28) As used herein, the terms “combination,” “in combination” and “combination therapies,” may refer generally to any or both of the “fixed combination” and “non-fixed combination” definitions and embodiments described above.

(29) As used herein, the transitional term “comprising,” when used to define compositions and methods, means that the compositions and methods include the recited elements, but does not exclude others. As used herein, “consisting essentially of,” when used to define compositions and methods, means that the composition and methods include additional elements, but only if those additional elements do not materially affect the basic and novel characteristics of the composition or methods. As used herein, “consisting of,” when used to define compositions and methods, means that the compositions and methods exclude more than trace elements of other ingredients for compositions and substantial method steps. Embodiments defined by each of these transitional terms are within the scope of this disclosure. For example, it is intended that the methods and compositions can include additional steps and components (comprising) or alternatively including steps and compositions of no significance (consisting essentially of) or alternatively, intending only the stated method steps or compositions (consisting of).

(30) As used herein, the terms “expression,” “expressed,” or “encodes” refer to a process by which polynucleotides are transcribed into mRNA and/or the process by which the transcribed mRNA is subsequently being translated into peptides, polypeptides, or proteins. Expression may include splicing of the mRNA in a eukaryotic cell or other forms of post-transcriptional modification or post-translational modification.

(31) As used herein, the term “farnesyl diphosphate synthase” may also be referred to herein as FDPS, and may also be referred to herein as farnesyl pyrophosphate synthase or FPPS.

(32) As used herein, the term “gamma delta T cell” may also be referred to herein as a $\gamma\delta$ T cell, or further as a GD T cell. The term “gamma delta T cell activation” refers to any measurable biological phenomenon associated with a gamma delta T cell that is representative of such T cell being activated. Non-limiting examples of such a biological phenomenon include an increase of cytokine production, changes in the qualitative or quantitative composition of cell surface proteins, an increase in T cell proliferation, and/or an increase in T cell effector function, such killing or a target cell or assisting another effector cell to kill a target cell.

(33) As used herein, the terms “individual,” “subject,” and “patient” are used interchangeably herein, and refer to any individual mammal subject, e.g., bovine, canine, feline, equine, and/or human.

(34) As used herein, the term “miRNA” refers to a microRNA, and also may be referred to herein as “miR”.

(35) The term “packaging cell line” refers to any cell line that can be used to express a lentiviral particle.

(36) As used herein, the term “homology” refers to the percentage number of amino acids, nucleic acids, or analogs thereof, that are identical or constitute conservative substitutions. Homology may be determined using sequence comparison programs such as GAP (Deveraux et al., 1984, Nucleic Acids Research 12, 387-395). In this way sequences of a similar or substantially different length to those cited herein could be compared by insertion of gaps into the alignment, such gaps being determined, for example, by the comparison algorithm used by GAP.

(37) As used herein, the term “sequence identity,” which also may appear in the non-limiting context of “a sequence 50% identical to,” and “having at least 80%, or at least 85%, or at least 90%, or at least 95% identity with” a given sequence, as similar phrasings, as used herein, refers to the extent that sequences are identical on a nucleotide-by-nucleotide basis or an amino acid-by-amino acid basis over a window of comparison. Thus, a “percentage of sequence identity” may be calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, I) or the identical amino acid residue (e.g., Ala, Pro, Ser, Thr, Gly, Val, Leu, Ile, Phe, Tyr, Trp, Lys, Arg, His, Asp, Glu, Asn, Gln, Cys and Met) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of

sequence identity. Optimal alignment of sequences for aligning a comparison window may be conducted by computerized implementations of algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Drive Madison, Wis., USA) or by inspection and the best alignment (i.e., resulting in the highest percentage homology over the comparison window) generated by any of the various methods selected. Reference also may be made to the BLAST family of programs as for example disclosed by Altschul et al., Nucl. Acids Res. 25:3389, 1997.

(38) As used here, the term “percent identity,” which may be used interchangeably with the term “sequence identity”, in the context of two or more nucleic acid or polypeptide sequences, refer to two or more sequences or subsequences that have a specified percentage of nucleotides or amino acid residues that are the same, when compared and aligned for maximum correspondence, as measured using one of the sequence comparison algorithms described below (e.g., BLASTP and BLASTN or other algorithms available to persons of skill) or by visual inspection. Depending on the application, the “percent identity” can exist over a region of the sequence being compared, e.g., over a functional domain, or, alternatively, exist over the full length of the two sequences to be compared. For sequence comparison, typically one sequence acts as a reference sequence to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are input into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. The sequence comparison algorithm then calculates the percent sequence identity for the test sequence(s) relative to the reference sequence, based on the designated program parameters.

(39) Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, Adv. Appl. Math. 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, J. Mol. Biol. 48:443 (1970), by the search for similarity method of Pearson & Lipman, Proc. Nat'l. Acad. Sci. USA 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by visual inspection (see generally Ausubel et al., *infra*).

(40) Suitable algorithms for determining percent sequence identity include the BLAST algorithm, which is described in Altschul et al., J. Mol. Biol. 215:403-410 (1990). Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information website.

(41) The percent identity between two nucleotide sequences can be determined using the GAP program in the GCG software package (available at <http://www.gcg.com>), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. The percent identity between two nucleotide or amino acid sequences can also be determined using the algorithm of E. Meyers and W. Miller (CABIOS, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. In addition, the percent identity between two amino acid sequences can be determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available at <http://www.gcg.com>), using either a Blossum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

(42) The nucleic acid and protein sequences of the present disclosure can further be used as a “query sequence” to perform a search against public databases to, for example, identify related sequences. Such searches can be performed using the NBLAST and XBLAST programs (version 2.0) of Altschul, et al. (1990) *J. Mol. Biol.* 215:403-10. BLAST nucleotide searches can be performed with the NBLAST program, score=100, word length=12 to obtain nucleotide sequences homologous to the nucleic acid molecules provided in the disclosure. BLAST protein searches can be performed with the XBLAST program, score=50, wordlength=3 to obtain amino acid sequences

homologous to the protein molecules of the disclosure. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul et al., (1997) *Nucleic Acids Res.* 25(17):3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>.

(43) As used herein, the term “pharmaceutically acceptable” refers to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues, organs, and/or bodily fluids of human beings and animals without excessive toxicity, irritation, allergic response, or other problems or complications commensurate with a reasonable benefit/risk ratio.

(44) As used herein, the term “pharmaceutically acceptable carrier” refers to, and includes, any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. The compositions can include a pharmaceutically acceptable salt, e.g., an acid addition salt or a base addition salt (see, e.g., Berge et al. (1977) *J Pharm Sci* 66:1-19).

(45) As used herein, the term “pharmaceutically acceptable salt” refers to derivatives of compounds or other active ingredients, wherein the parent compound or active ingredient is modified by converting an existing acid or base moiety to its salt form. Non-limiting examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; alkali metal, alkaline metal, ammonium, and mono-, di, tri-, or tetra-C1-C30-alkyl-substituted ammonium; and the like. The pharmaceutically acceptable salts of various embodiments include the conventional non-toxic salts of the compound or active ingredient formed, for example, from nontoxic inorganic or organic acids. Suitable organic acids are, e.g., carboxylic acids or sulfonic acids, such as acetic acid, succinic acid, fumaric acid or methanesulfonic acid. The pharmaceutically acceptable salts herein can be synthesized from the parent compound or active ingredient which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, Pa., 1985, p. 1418 and Journal of Pharmaceutical Science, 66, 2 (1977), each of which is incorporated herein by reference in its entirety.

(46) As used herein, the term “SEQ ID NO” is synonymous with the term “Sequence ID No.”

(47) As used herein, the term “small RNA” refers to non-coding RNA that are generally less than about 200 nucleotides or less in length and possess a silencing or interference function. In embodiments, the small RNA is about 175 nucleotides or less, about 150 nucleotides or less, about 125 nucleotides or less, about 100 nucleotides or less, or about 75 nucleotides or less in length. Such RNAs include microRNA (miRNA), small interfering RNA (siRNA), double stranded RNA (dsRNA), and short hairpin RNA (shRNA). In embodiments, “small RNA” are capable of inhibiting or knocking-down gene expression of a target gene, generally through pathways that result in the inhibitions or destruction of the target gene mRNA.

(48) As used herein, the term “therapeutically effective amount” refers to a sufficient quantity of the active agents of the present disclosure, in a suitable composition, and in a suitable dosage form to treat or prevent the symptoms, progression, or onset of the complications seen in patients suffering from a given ailment, injury, disease, or condition. The therapeutically effective amount will vary depending on the state of the patient's condition or its severity, and the age, weight, etc., of the subject to be treated. A therapeutically effective amount can vary, depending on any of a number of factors, including, e.g., the route of administration, the condition of the subject, as well as other factors understood by those in the art.

(49) As used herein, the term “therapeutic vector” includes, without limitation, reference to a lentiviral vector or an AAV vector.

(50) As used herein, the terms “treatment” and “treating” refer to the intended targeting of a disease state and combatting of it, i.e., ameliorating or preventing the disease state. A particular treatment thus will depend on the disease state to be targeted and the current or future state of medicinal therapies and therapeutic approaches. A treatment may have associated toxicities.

(51) Desirable effects include, but are not limited to, preventing occurrence or recurrence of disease, alleviating symptoms, suppressing, diminishing or inhibiting any direct or indirect pathological consequences of the disease, ameliorating or palliating the disease state, and causing remission or improved prognosis.

(52) Description of Aspects of the Disclosure

(53) In one aspect, a method of activating a GDT cell is provided. The method includes infecting, in the presence of the GD T cell, a target cell with a viral delivery system encoding at least one genetic element. In embodiments, the at least one encoded genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In embodiments, the enzyme is FDPS. In embodiments, when the enzyme is inhibited in the target cell, the target cell activates the GD T cell. In embodiments, the target cell is a cancer cell or a cell that has been infected with an infectious agent. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA.

(54) In embodiments, the at least one encoded genetic element includes a shRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with

GTCCTGGAGTACAATGCCATTCTCGAGAATGGCATTGTACTCCAGGACTTTTT (SEQ ID NO: 1); GCAGGATTTCGTTTCAGCACTTCTCGAGAAGTGCTGAACGAAATCCTGCTTTTT (SEQ ID NO: 2); GCCATGTACATGGCAGGAATTCTCGAGAA TTCCTGCCATGTACATGGCTTTTT (SEQ ID NO: 3); or GCAGAAGGAGGCTGA GAAAGTCTCGAGACTTTCTCAGCCTCCTTCTGCTTTTT (SEQ ID NO: 4). In a preferred embodiment, the shRNA includes GTCCTGGAGTACAATGCCATTCTCGAG AATGGCATTGTACTCCAGGACTTTTT (SEQ ID NO: 1); GCAGGATTTCGTTCA GCACTTCTCGAGAAGTGCTGAACGAAATCCTGCTTTTT (SEQ ID NO: 2); GCCA TGTACATGGCAGGAATTCTCGAGAATTCCTGCCATGTACATGGCTTTTT (SEQ ID NO: 3); or GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCCTCCTT CTGCTTTTT (SEQ ID NO: 4).

(55) In another aspect, the at least one encoded genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with

AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGCGTGAAGC CACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGCCTCGGACTTCAAGGG GCT (SEQ ID NO: 5); AAGGTATATTGCTGTTGACAGTGAGCGACACT TTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGGCTGAGAAAGTGCTGC CTACTGCCTCGGACTTCAAGGGGCT (SEQ ID NO: 6); TGCTGTTGACAGTG AGCGACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGAGGCTGAG AAAGTTGCCTACTGCCTCGGA (SEQ ID NO: 7); CCTGGAGGCTTGCTGAAG GCTGTATGCTGACTTTCTCAGCCTCCTTCTGCTTTTGGCCACTGACTGAGCAGAAGGG CTGAGAAAGTCAGGACACAAGGCCTGTTACTAGCACTCA (SEQ ID NO: 8); CATCTCCATGGCTGTACCACCTTGTCTGGGACTTTCTCAGCCTCCTTCTGCCTGTTGAA TCTCATGGCAGAAGGAGGCGAGAAAGTCTGACATTTTGGTATCTTTTCATCTGACCA (SEQ ID NO: 9); or GGGCCTGGCTCGAGCAGGGGGCGAGGGATACTTTCT

CAGCTTCCTTCTGCTGGTCCCTCCCTCCCGCAGAAGGAGGCTGAGAAAGTCCTTCCCTC
CCAATGACCGCGTCTTTCGTCG (SEQ ID NO: 10). In a preferred embodiment, the microRNA
includes AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCT
CCTTCTGCGTGAAGCCACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGC
CTCGGACTTCAAGGGGCT (SEQ ID NO: 5); AAGGTATATTGCTGTTGACAGT
GAGCGACACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGGGCTGA
GAAAGTGCTGCCTACTGCCTCGGACTTCAAGGGGCT (SEQ ID NO: 6); TGCTG
TTGACAGTGAGCGACTTTCTCAGCCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGG
AGGCTGAGAAAGTTGCCTACTGCCTCGGA (SEQ ID NO: 7); CCTGGAGGCT
TGCTGAAGGCTGTATGCTGACTTTCTCAGCCTCCTTCTGCTTTTGGCCACTGACTGAG
CAGAAGGGGCTGAGAAAGTCAGGACACAAGGCCTGTTACTAGCACTCA (SEQ ID NO: 8);
CATCTCCATGGCTGTACCACCTTGTCGGGACTTTCTCAGCCTCCTT
CTGCCTGTTGAATCTCATGGCAGAAGGAGGCGAGAAAGTCTGACATTTTGGTATCTT
TCATCTGACCA (SEQ ID NO: 9); or GGGCCTGGCTCGAGCAGGGGGCGAGGG
ATACTTTCTCAGCCTCCTTCTGCTGGTCCCCTCCCCGCAGAAGGAGGCTGAGAAAGT
CCTTCCCTCCCAATGACCGCGTCTTCGTCG (SEQ ID NO: 10).

(56) In another aspect, the target cell is also contacted with a bisphosphonate drug. In a preferred embodiment, the bisphosphonate drug is zoledronic acid. The bisphosphonate drug may be a pharmaceutically acceptable salt, hydrate or a solvate thereof.

(57) In another aspect, a method of treating cancer in a subject is provided. The method includes administering to the subject a therapeutically-effective amount of a viral delivery system encoding at least one genetic element. In embodiments, the at least one encoded genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In further embodiments, when the enzyme is inhibited in a cancer cell in the presence of a GD T cell, the cancer cell activates the GD T cell, to thereby treat the cancer. In embodiments, the enzyme is FDPS. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA.

(58) In another aspect, a method of treating an infectious disease in a subject is provided. The method includes administering to the subject a therapeutically-effective amount of a viral delivery system encoding at least one genetic element. In embodiments, the at least one encoded genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In further embodiments, when the enzyme is inhibited in a cell that is infected with an infectious agent and is in the presence of a GD T cell, the infected cell activates the GD T cell, to thereby treat the infected cell, and the infectious disease. In embodiments, the enzyme is FDPS. In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA.

(59) In embodiments, the at least one encoded genetic element includes a shRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4. In a preferred embodiment, the shRNA includes SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4.

(60) In other embodiments, the at least one encoded genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In a preferred embodiment, the microRNA includes SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10.

(61) In another aspect, a viral vector comprising at least one encoded genetic element is provided.

The at least one encoded genetic element includes a small RNA capable of inhibiting production of an enzyme involved in the mevalonate pathway. In embodiments, the enzyme involved in the mevalonate pathway is farnesyl diphosphate synthase (FDPS). In embodiments, the at least one encoded genetic element includes a microRNA or a shRNA.

(62) In another aspect, the at least one encoded genetic element includes a shRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4. In a preferred embodiment, the shRNA includes SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4.

(63) In another aspect, the at least one encoded genetic element includes a microRNA having at least 80%, at least 81%, at least 82%, at least 83%, at least 84%, at least 85%, at least 86%, at least 87%, at least 88%, at least 89%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95% or more percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In a preferred embodiment, the microRNA includes SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10.

(64) In embodiments, the viral vector includes any vector that can effectively transduce the small RNA. In embodiments, the viral vector is a lentiviral vector. In other embodiments, the viral vector is an adeno-associated virus (AAV) vector.

(65) In another aspect, the viral vector includes a second encoded genetic element. In embodiments, the second genetic element includes at least one cytokine or chemokine. In embodiments, the at least one cytokine is selected from the group consisting of: IL-18, TNF- α , interferon- γ , IL-1, IL-2, IL-15, IL-17, and IL-12. In embodiments, the at least one chemokine is a CC chemokine, CXC chemokine, a CX3 chemokine or a XC chemokine. In a further embodiment, the at least one chemokine is the CC chemokine, RANTES.

(66) In another aspect, a lentiviral vector system for expressing a lentiviral particle is provided. The system includes a lentiviral vector, at least one envelope plasmid for expressing an envelope protein optimized for infecting a cell; and at least one helper plasmid for expressing gag, pol, and rev genes. When the lentiviral vector, the at least one envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell, a lentiviral particle is produced by the packaging cell. In embodiments, the lentiviral particle is capable of infecting a targeting cell, and inhibiting an enzyme involved in the mevalonate pathway within the target cell. In embodiments, the enzyme involved in the mevalonate pathway is FDPS. In embodiments, the lentiviral vector system includes a first helper plasmid for expressing the gag and pol genes, and a second helper plasmid for expressing the rev gene. In embodiments, the envelope protein is preferably optimized for infecting a target cell. In embodiments, the target cell is a cancer cell. In other embodiments, the target cell is a cell that is infected with an infectious disease.

(67) In another aspect a pharmaceutical combination is disclosed which includes a bisphosphonate compound; and a lentiviral particle produced by a packaging cell and capable of infecting a target cell. The lentiviral particle comprises an envelope protein capable of infecting the target cell, and: at least one encoded shRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded shRNA comprises a sequence having at least 80%, or at least 81%, or at least 82%, or at least 83%, or at least 84%, or at least 85%, or at least 86%, or at least 87%, or at least 88%, or at least 89%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98% or at least 99% percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4; or at least one encoded microRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded microRNA comprises a sequence having at least 80%, or at least 81%, or at least 82%, or at least 83%, or at least 84%, or at least 85%, or at

least 86%, or at least 87%, or at least 88%, or at least 89%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98% or at least 99% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10, wherein the pharmaceutical combination is at least one of fixed and non-fixed. In embodiments, the at least one encoded shRNA comprises SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4 or the at least one encoded microRNA comprises SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the pharmaceutical composition comprises a fixed combination. In embodiments, the pharmaceutical composition comprises a non-fixed combination. In embodiments, the bisphosphonate drug comprises zoledronic acid. In embodiments, the bisphosphonate drug and the lentiviral particle are present in synergistically effective amounts. In embodiments, the target cell is one or more cancer cells that are present in a cancer selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof. In embodiments, the target cell is one or more cancer cells that are present in a hepatocellular carcinoma. In embodiments, the target cell is capable of activating a gamma delta T cell following infection with the lentiviral particle. In embodiments, the enzyme is FDPS.

(68) In another aspect, a method of treating a cancer in a subject using an immunotherapy-based composition is disclosed. The method includes administering a therapeutically-effective amount of a bisphosphonate drug to the subject; and administering a therapeutically-effective amount of the immunotherapy-based composition to the subject, wherein the immunotherapy-based composition comprises a lentiviral particle. The lentiviral particle comprises an envelope protein capable of infecting one or more cancer cells, and at least one encoded shRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded shRNA comprises a sequence having at least 80%, or at least 81%, or at least 82%, or at least 83%, or at least 84%, or at least 85%, or at least 86%, or at least 87%, or at least 88%, or at least 89%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98% or at least 99% percent identity with SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4 or at least one encoded microRNA capable of inhibiting production of an enzyme of the mevalonate pathway, wherein the at least one encoded microRNA comprises a sequence having at least 80%, or at least 81%, or at least 82%, or at least 83%, or at least 84%, or at least 85%, or at least 86%, or at least 87%, or at least 88%, or at least 89%, or at least 90%, or at least 91%, or at least 92%, or at least 93%, or at least 94%, or at least 95%, or at least 96%, or at least 97%, or at least 98% or at least 99% percent identity with SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the at least one encoded shRNA comprises SEQ ID NO: 1; SEQ ID NO: 2; SEQ ID NO: 3; or SEQ ID NO: 4. In embodiments, the at least one encoded microRNA comprises SEQ ID NO: 5; SEQ ID NO: 6; SEQ ID NO: 7; SEQ ID NO: 8; SEQ ID NO: 9; or SEQ ID NO: 10. In embodiments, the one or more cancer cells are present in a cancer selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof. In embodiments, the bisphosphonate drug comprises zoledronic acid. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in a fixed combination. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in a non-fixed combination. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered simultaneously. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered sequentially. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered in synergistically effective amounts. In embodiments, the bisphosphonate drug and the immunotherapy-based composition are administered at a synergistically effective time interval. In embodiments, the one or more cancer

cells are capable of activating a gamma delta T cell resident in the subject following infection of the one or more cancer cells with the immunotherapy-based composition. In embodiments, activating the gamma delta T cell comprises increasing tumor necrosis factor (TNF)- α expression by the gamma delta T cell. In embodiments, the enzyme of the mevalonate pathway is farnesyl diphosphate synthase (FDPS).

(69) Cancer

(70) The compositions and methods provided herein are used to treat cancer. A cell, tissue, or target may be a cancer cell, a cancerous tissue, harbor cancerous tissue, or be a subject or patient diagnosed or at risk of developing a disease or condition. In certain aspects, a cell may be an epithelial, an endothelial, a mesothelial, a glial, a stromal, or a mucosal cell. The cancer cell population can include, but is not limited to a brain, a neuronal, a blood, an endometrial, a meninges, an esophageal, a lung, a cardiovascular, a liver, a lymphoid, a breast, a bone, a connective tissue, a fat, a retinal, a thyroid, a glandular, an adrenal, a pancreatic, a stomach, an intestinal, a kidney, a bladder, a colon, a prostate, a uterine, an ovarian, a cervical, a testicular, a splenic, a skin, a smooth muscle, a cardiac muscle, or a striated muscle cell, and can also include a cancer cell population from any of the foregoing, and can be associated with one or more of carcinomas, sarcomas, myelomas, leukemias, lymphomas, mixed types or mixtures of the foregoing. In still a further aspect cancer includes, but is not limited to astrocytoma, acute myeloid leukemia, anaplastic large cell lymphoma, acute lymphoblastic leukemia, angiosarcoma, B-cell lymphoma, Burkitt's lymphoma, breast carcinoma, bladder carcinoma, carcinoma of the head and neck, cervical carcinoma, chronic lymphoblastic leukemia, chronic myeloid leukemia, colorectal carcinoma, endometrial carcinoma, esophageal squamous cell carcinoma, Ewing's sarcoma, fibrosarcoma, glioma, glioblastoma, gastrinoma, gastric carcinoma, hepatoblastoma, hepatocellular carcinoma, Kaposi's sarcoma, Hodgkin lymphoma, laryngeal squamous cell carcinoma, larynx carcinoma, leukemia, leiomyosarcoma, lipoma, liposarcoma, melanoma, mantle cell lymphoma, medulloblastoma, mesothelioma, myxofibrosarcoma, myeloid leukemia, mucosa-associated lymphoid tissue B cell lymphoma, multiple myeloma, high-risk myelodysplastic syndrome, nasopharyngeal carcinoma, neuroblastoma, neurofibroma, high-grade non-Hodgkin lymphoma, non-Hodgkin lymphoma, lung carcinoma, non-small cell lung carcinoma, ovarian carcinoma, esophageal carcinoma, osteosarcoma, pancreatic carcinoma, pheochromocytoma, prostate carcinoma, renal cell carcinoma, retinoblastoma, rhabdomyosarcoma, salivary gland tumor, schwannoma, small cell lung cancer, squamous cell carcinoma of the head and neck, testicular tumor, thyroid carcinoma, urothelial carcinoma, and Wilms tumor.

(71) The compositions and methods provided herein are also used to treat NSCLC (non-small cell lung cancer), pediatric malignancies, cervical and other tumors caused or promoted by human papilloma virus (HPV), melanoma, Barrett's esophagus (pre-malignant syndrome), adrenal and skin cancers and auto immune, neoplastic cutaneous diseases.

(72) Infectious Diseases

(73) The compositions and methods disclosed herein can be used to treat infectious diseases. The term "infectious disease" includes any disease that is caused by an infectious agent. An "infectious agent" includes any exogenous pathogen including, without limitation, bacteria, fungi, viruses, mycoplasma, and parasites. Infectious agents that may be treated with compositions provided for in this disclosure include any art-recognized infectious organisms that cause pathogenesis in an animal, including such organisms as bacteria that are gram-negative or gram-positive cocci or bacilli, DNA and RNA viruses, including, but not limited to, DNA viruses such as papilloma viruses, parvoviruses, adenoviruses, herpesviruses and vaccinia viruses, and RNA viruses, such as arenaviruses, coronaviruses, rhinoviruses, respiratory syncytial viruses, influenza viruses, picomaviruses, paramyxoviruses, reoviruses, retroviruses, and rhabdoviruses. Examples of fungi that may be treated with the compositions and methods of the disclosure include fungi that grow as molds or are yeast-like, including, for example, fungi that cause diseases such as ringworm,

histoplasmosis, blastomycosis, aspergillosis, cryptococcosis, sporotrichosis, coccidioidomycosis, paracoccidio-idomycosis, and candidiasis. Compositions and methods provided for herein may be utilized to treat parasitic infections including, but not limited to, infections caused by somatic tapeworms, blood flukes, tissue roundworms, ameba, and *Plasmodium*, *Trypanosoma*, *Leishmania*, and *Toxoplasma* species.

(74) Methods of GD T Cell Activation

(75) Provided herein are compositions and methods for activating GD T cells in an individual, as well as methods for treating tumors and infectious diseases. For instance, in embodiments, the compositions and methods provided herein can be used in methods to treat all known cancers because activated GD T cells comprise a natural mechanism for immune surveillance of tumors (See for e.g.: Pauza et al. 2014 *Frontiers in Immunol.* 5:687). Likewise, in embodiments, the compositions and methods provided herein can be used to treat infectious diseases, including but not limited to flavivirus, influenza virus, human retrovirus, mycobacteria, plasmodia and a variety of other viral, fungal and bacterial infections. (See for e.g.: Pauza and Cairo, 2015 *Cell Immunol.* 296(1).

(76) In general, a vector system is administered to an individual to transfect or transduce a target cell population with the disclosed constructs for decreasing expression of FDPS and, in other embodiments, increasing expression of chemokines or cytokines. Administration and transfection/transduction can occur in vivo or ex vivo, with the transfected cells later administered back into the subject in the latter scenario.

(77) Administration of the disclosed vectors and transfection or transduction of the disclosed constructs into a subject's cells result in decreased expression of FDPS, increased expression of cytokines or chemokines, accumulation of IPP and in many cases, reduced growth rates for genetically modified tumor cells. All of these features work together to activate and co-localize GD T cells to the site of a tumor or infection.

(78) The disclosed methods can also increase the capacity of NK cells to recognize and destroy tumor cells and/or infected cells. Crosstalk between GD T cells and NK cells is an important aspect of regulating the immune and inflammatory responses. Further, GD T cells are known to trigger dendritic cell maturation, recruit B cells and macrophages, and participate in a variety of cytolytic activities, such as secretion of interferon- γ and TNF- α .

(79) In embodiments, the disclosed compositions and methods provided herein comprise a form of gene therapy for activating GD T cells at the site of tumor or infectious disease pathology. In an aspect, the compositions and methods provided herein activate GD T cells and support their proliferation, differentiation, and functional capacities by promoting the production of specific cytokines needed for cytolytic activity capable of killing cancer cells or treating infectious diseases.

(80) In embodiments the gene therapy sequences (e.g., FDPS shRNAs) are carried by therapeutic vectors, including but not limited to viral vectors such as lentiviruses or adeno-associated viruses, although other viral vectors can also be suitable. Gene therapy constructs may also be delivered in the form of DNA or RNA, including but not limited to plasmid forms. In embodiments, the disclosed gene therapy constructs may also be delivered in the form of protein-nucleic acid complexes or lipid nucleic acid complexes and mixtures of these formulations. For instance, a protein-nucleic acid complex can comprise nucleic acids of interest in a complex with cationic peptides such as lysine and arginine. Lipid-nucleic acids complexes can comprise lipid emulsions, micelles, liposomes, and/or mixtures of neutral and cationic lipids such as DOTMA, DOSPA, DOTAP, and DMRIE.

(81) In embodiments, therapeutic vectors may comprise a single construct or at least two, at least three, at least four, or at least five different constructs. When more than one construct is present in a vector the constructs may be identical, or they may be different. For instance, the constructs may vary in terms of their promoters, the presence or absence of integrating elements, and/or their sequences. In some embodiments, a therapeutic vector will comprise at least one construct that

encodes a small RNA capable of knocking down the expression of FDPS. In embodiments, the therapeutic vector will also encode a specific cytokine(s) and/or chemokine(s), including but not limited to TNF- α , interferon- γ , IL-1, IL-2, IL-15, IL-17, IL-18 or IL-12. In some embodiments, a single construct may encode both small RNAs capable of knocking down the expression of FDPS and specific cytokines or chemokines, including but not limited to TNF- α , interferon- γ , IL-1, IL-2, IL-15, IL-17, IL-18 or IL-12.

(82) In embodiments, viral vectors may introduce nucleic acid constructs that become integrated into the host chromosome. Alternately, transient delivery vectors may be used to prevent chromosomal integration and limit the lifespan of gene therapy constructs.

(83) In embodiments, the disclosed constructs and vectors comprise short hairpin RNA ("shRNA"), micro RNA ("miRNA"), or siRNA capable of reducing or knocking down expression of FDPS and/or geranyl pyrophosphate synthase ("GPPS") and/or farnesyl transferase ("FT") genes. By down regulating these genes, which control steroid and isoprenoid synthesis, isopentenyl pyrophosphate ("IPP") levels are elevated. Elevation and accumulation of IPP is a known mechanism for increasing GD T cells activation. Further, down regulation of these pyrophosphate synthase genes removes an important negative regulator of inflammasome function that in turn results in increased expression of cytokines that are important for GD T cell activation and effector cell function.

(84) In embodiments, the disclosed constructs are regulated by specific promoters that are capable of producing interleukin-2 and/or interleukin-15 to sustain GD T cell proliferation. In addition, the disclosed constructs may be regulated by specific promoters that are capable of producing interleukin-1 beta and/or interleukin-18 and/or interferon-gamma required for GD T cell differentiation and acquisition of all effector cell function. Desirable effector cell functions include the capacity for direct cytotoxic cell killing of tumors and/or infected cells, secretion of beneficial cytokines and/or chemokines, increased expression of NK receptors required to recognize cancerous or infected cells, and increased expression of Fc receptors needed to bind targeting antibodies in order to co-localize GD T cells with cancerous or infected cell targets.

(85) In embodiments, the disclosed methods activate GD T cells, resulting in the indirect effect of increasing the capacity for NK cells to attack and destroy cancerous cells, tumors, or infected cells. The activation of NK cells requires GD T cells that are stimulated to proliferate and differentiate, and to express 4-1BBL costimulatory ligand needed to engage the 4-1BB costimulatory receptor on NK cells. This form of crosstalk is known as an important mechanism for activating NK cells and is achieved here through the action of the disclosed methods and compositions.

(86) In another aspect, crosstalk between GD T cells and NK cells is an important mechanism for eliminating inflammatory dendritic cells that accumulate in diseased tissues. Alone, neither GD T cells nor NK cells are capable of destroying dendritic cells, but once the aforementioned crosstalk interactions have occurred, NK cells are altered to become cytotoxic against inflammatory dendritic cells. This immuno-regulatory mechanism depends on strong activation and proliferation of GD T cells.

(87) In embodiments, the disclosed methods for activation of GD T cells further comprise a step of suppressing pathologic inflammatory responses that may include cellular proliferation leading to atherosclerosis, chronic immune activation that stimulates tumor growth, autoimmune diseases including psoriasis and other presentations in the epidermis, inflammatory diseases of the central nervous system, and arthritis and other diseases of unregulated immune responses.

(88) In embodiments, therapeutic vectors are administered in combination with bisphosphonate drugs. In various embodiments, such combinations achieve synergistic, positive or heightened activation of gamma delta T cells. Such positive activation may allow alternate, modified or reduced doses of bisphosphonates and may decrease adverse reactions to bisphosphonates including acute inflammatory responses and chronic diseases. Combinations of therapeutic vectors with bisphosphonates may be together or separate, with or without instructions for combined use or

to combination products. The therapeutic vectors and/or bisphosphonates may be administered entirely separately and may be formulated in entirely distinct pharmaceutical dosage forms. The therapeutic vectors and/or bisphosphonates may be sold independently of each other, with or without label instructions concerning the possibility of a combined use. Such instructions also may be provided in the package equipment, e.g., leaflet or the like, or in other information, e.g., provided to physicians and medical staff (e.g., oral communications, communications in writing or the like). Such labels or other instructions can refer to either a fixed combination in one dosage unit form, or a non-fixed combination as a kit of parts for the combined administration where the therapeutic vector may be administered independently of the bisphosphonate drug, at the same time, or separately within time intervals. In various embodiments, the combination exhibits a cooperative or joint effect, or a decrease in toxicity or complications of treatment. In one embodiment the effect of the combination is synergistic. A synergistic effect is achieved when the active ingredients used together is greater than the sum of the effects that results from using the compounds separately. A synergistic effect may be attained when the active ingredients are: (1) co-formulated and administered or delivered simultaneously in a combined, unit dosage formulation; (2) delivered by alternation or in parallel as separate formulations; or (3) by some other regimen. When delivered in alternation therapy, a synergistic effect may be attained when the compounds are administered or delivered sequentially, e.g., by different injections in separate syringes. In general, during alternation therapy, an effective dosage of each active ingredient is administered sequentially, i.e., serially, whereas in combination therapy, effective dosages of two or more active ingredients are administered together, albeit subject to potential variances in timing as detailed herein.

(89) The combinations herein may be manufactured and/or formulated by the same or different manufacturers. The active ingredients may be brought together into a combination therapy: (i) prior to release of the combination product to physicians (e.g., in the case of a kit comprising the compound of the disclosure and the other therapeutic agent); (ii) by the treating physician (or under the guidance of a physician) shortly before administration; (iii) in the actual patient, e.g., during sequential administration of the active ingredients disclosed herein.

(90) In embodiments, a therapeutically effective amount of each of the combinations may be administered simultaneously or sequentially and in any order, and the components may be administered together or separate. For example, the method of treating a proliferative disease according to the disclosure may comprise (i) administration of a first agent such as a therapeutic vector that forms part of a lentiviral particle, and (ii) administration of a second agent such as a bisphosphonate drug in free or pharmaceutically acceptable salt form, simultaneously or sequentially in any order, in jointly therapeutically effective amounts, preferably in cooperative, jointly effective, and/or synergistically effective, amounts, e.g., in daily or intermittent dosages corresponding to the amounts described herein. The combinations may be administered separately at different times during the course of therapy or concurrently in divided or single drug forms. Furthermore, the term “administering” also encompasses the use of a pro-drug of a combination partner that converts in vivo to the combination partner as such. The instant disclosure is therefore to be understood as embracing all such regimens of simultaneous or alternating treatment and the term “administering” is to be interpreted accordingly.

(91) In embodiments, agents (i) and (ii) can be administered using any pharmaceutically acceptable method, such as intranasal, buccal, sublingual, oral, rectal, ocular, parenteral (intravenously, intradermally, intramuscularly, subcutaneously, intraperitoneally), pulmonary, intravaginal, locally administered, topically administered, topically administered after scarification, mucosally administered, via an aerosol, in semi-solid media such as agarose or gelatin, or via a buccal or nasal spray formulation, and/or in solid media such as granules or powders including inert excipients. For example, a therapeutic vector and/or bisphosphonate drug may be administered intravenously. Further, agents (i) and (ii) can be formulated into any pharmaceutically acceptable dosage form,

such as a solid dosage form, tablet, pill, lozenge, capsule, liquid dispersion, gel, aerosol, pulmonary aerosol, nasal aerosol, ointment, cream, semi-solid dosage form, a solution, an emulsion, and a suspension. For example, a bisphosphonate drug may be formulated into a tablet and administered orally.

(92) A combination therapy according to the disclosure can besides or in addition be administered especially for cancer therapy in combination with chemotherapy, radiotherapy, immunotherapy, surgical intervention, or a combination of these. Long-term therapy is equally possible as is adjuvant therapy in the context of other treatment strategies, as described above. In embodiments, a combination therapy can also include immune adjuvants (e.g., Toll-like receptor ligands), immune stimulating toxins, or stimulatory protozoans or stimulatory bacilli (e.g., bacille Calmette-Guerin), cancer therapeutic drugs, cell-based therapies (gamma delta T cell or other cell types known to be in use or under evaluation for tumor therapy and may also include natural or genetically-engineered cells and cells cultured under) ionizing radiation or surgery. Other possible treatments are therapy to maintain the patient's status after tumor regression, or even chemo-preventive therapy, for example in patients at risk.

(93) Constructs for GD T Cell Activation

(94) Inhibition of FDPS results in IPP accumulation, resulting in activation of V δ 2+GD T cells and expression of IL-18, which is also important in activating GD T cells. Inhibition of farnesyl transferase results in decreased prenylation of proteins. The disclosed constructs can be transfected or transduced into specific target cells, like tumor cells or infected cells, where they can express RNA sequences (i.e., siRNA, shRNA or microRNA) that will inhibit translation of FDPS as well as encode and express cytotoxic cytokines or chemokines.

(95) Disclosed herein are constructs for decreasing expression of FDPS and/or FT, increasing expression of cytokines, and increasing expression of chemokines including RANTES. For instance, in some embodiments the constructs may encode for interferon-gamma, IL-1, IL-2, IL-15, IL-17, IL-18 or IL-12.

(96) Expression of cytokines and chemokines, like those listed above, will result in localized cytotoxic destruction of tumor cells or cells infected with pathogenic organisms. Accordingly, expression of such constructs by a tumor cell or an infected cell will result in the unwanted cells assisting in its own destruction.

(97) Likewise, if the disclosed constructs are expressed in a tumor cell or infected cell, decreasing the expression of FDPS and FT will result in activation and recruitment of GD T cells to the tumor site of site of cell infection. Increasing expression of RANTES will further attract GD T cells to intended tissue location. Because GD T cells can kill a broad range of tumors of epithelial origin as well as many leukemias and lymphomas, and are further able to produce high levels of the anti-tumor cytokine, IFN γ , recruitment of GD T cells to the site of a tumor can be a particularly effective means of inducing anti-tumor immunity.

(98) Decreased expression of FDPS can be achieved via shRNA, microRNA, siRNA, or other means known in the art. For instance, shRNAs according to SEQ ID NOS: 1, 2, 3, or 4, or variants thereof can be used in the disclosed constructs and methods, although this example is not limiting. The coding regions for RNAs to decrease expression of FDPS and FT and the coding regions of cytokine and chemokines may be in the same construct or on different constructs.

(99) The classical approach for the production of recombinant polypeptides or gene regulatory molecules including small RNA is the use of stable expression constructs. These constructs are based upon chromosomal integration of a transduced expression plasmid (or at least a portion thereof) into the genome of the host cell, short-duration plasmid transfection, or non-integrating viral vectors also with limited half-life. The sites of gene integration are generally random, and the number and ratio of genes integrating at any particular site are often unpredictable; likewise, non-integrating plasmids or viral vectors also generate nuclear DNA but these species usually lack sequences required for DNA replication and continuous maintenance. Thus, constructs that rely on

chromosomal integration result in permanent maintenance of the recombinant gene that may exceed the therapeutic interval.

(100) An alternative to stable expression constructs for gene expression are transient expression constructs. The expression of the latter gene expression construct is based on non-integrated plasmids, and hence the expression is typically lost as the cell undergoes division or the plasmid vectors are destroyed by endogenous nucleases.

(101) The disclosed constructs are preferably episomal constructs that are transiently expressed. Episomal constructs are degraded or diluted over time such that they do not make permanent changes to a subject's genome, nor are they incorporated into the chromosome of a target cell. The process of episomal replication typically incorporates both host cell replication machinery and viral trans-acting factors.

(102) Avoiding chromosomal integration reduces certain barriers to in vivo gene delivery. However, even integration-defective constructs can have a background frequency of integration, and any DNA molecule can find rare homologies to recombine with host sequences; but these rates of integration are exceptionally rare and generally not clinically significant.

(103) Thus, in some embodiments, the disclosed vectors support active gene and/or small RNA delivery over a period of about 1, about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, about 10, about 11, or about 12 weeks. In some embodiments, the disclosed vectors support active gene and/or small RNA delivery over a period of about 1 month, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, or longer. Any combination of these time periods can also be used in the methods of the invention, e.g., 1 month and 1 week, or 3 months and 2 weeks.

(104) However, in some embodiments, the constructs comprise integrating elements that depend on a retroviral integrase gene, such that the construct becomes integrated into the subject's chromosome. Retrotransposition and transposition are additional examples of mechanisms whereby mobile genetic elements become integrated or inserted into the chromosome. Plasmids may become integrated into the chromosome by recombination, and gene editing technologies including CRISPR and TALEN utilize guide RNA sequences and alter chromosomal loci by gene conversion mechanisms.

(105) Constructs may comprise specific promoters for expressing cytokines involved in the maintenance of GD T cells (i.e., IL-2, IL-7, IL-17, and IL-15). For example, promoters that may be incorporated into the disclosed constructs include but are not limited to TATA-box promoters, CpG-box promoters, CCAAT-box promoters, TTGACA-box promoters, BRE-box promoters, INR-box promoters, AT-based promoters, CG-based promoters, ATCG-compact promoters, ATCG-balanced promoters, ATCG-middle promoters, ATCG-less promoters, AT-less promoters, CG-less promoters, AT-spike promoters, and CG-spike promoters. See Gagniuc and Ionescu-Tirgoviste, Eukaryotic genomes may exhibit up to 10 generic classes of gene promoters, BMC GENOMICS 13:512 (2012).

(106) Therapeutic Vectors

(107) The construct can be delivered via known transfection and/or transduction vectors, including but not limited to lentiviral vectors, gamma-retroviruses, adeno-associated virus, poxvirus, herpesvirus vectors, protein and/or lipid complexes, liposomes, micelles, and the like.

(108) Viral vectors can be preferentially targeted to cell types that are useful for the disclosed methods (i.e., tumor cells or myeloid cells). Viral vectors can be used to transduce genes into target cells owing to specific virus envelope-host cell receptor interactions and viral mechanisms for gene expression. As a result, viral vectors have been used as vehicles for the transfer of genes into many different cell types including whole embryos, fertilized eggs, isolated tissue samples, tissue targets in situ, and cultured cell lines. The ability to introduce and express foreign genes in a cell is useful for the study of gene expression, and the elucidation of cell lineages as well as providing the potential for therapeutic interventions such as gene therapy, somatic cell reprogramming of induced

pluripotent stem cells, and various types of immunotherapy. Viral components from viruses like Papovaviridae (e.g. bovine papillomavirus or BPV) or Herpesviridae (e.g. Epstein Barr Virus or EBV) or Hepadnaviridae (e.g. Hepatitis B Virus or HBV) or pox vectors including vaccinia may be used in the disclosed vectors.

(109) Lentiviral vectors are a preferred type of vector for the disclosed compositions and methods, although the disclosure is not specifically limited to lentiviral vectors. Lentivirus is a genus of viruses that can deliver a significant amount of viral nucleic acid into a host cell. Lentiviruses are characterized as having a unique ability to infect/transduce non-dividing cells, and following transduction, lentiviruses integrate their nucleic acid into the host cell's chromosomes.

(110) Infectious lentiviruses have three main genes coding for the virulence proteins gag, pol, and env, and two regulatory genes including tat and rev. Depending on the specific serotype and virus, there may be additional accessory genes that code for proteins involved in regulation, synthesis, and/or processing viral nucleic acids and other replicative functions.

(111) Moreover, lentiviruses contain long terminal repeat (LTR) regions, which may be approximately 600 nt long. LTRs may be segmented into U3, R, and U5 regions. LTRs can mediate integration of retroviral DNA into the host chromosome via the action of integrase. Alternatively, without functioning integrase, the LTRs may be used to circularize the viral nucleic acid.

(112) Viral proteins involved in early stages of lentivirus replication include reverse transcriptase and integrase. Reverse transcriptase is the virally encoded, RNA-dependent DNA polymerase. The enzyme uses a viral RNA genome as a template for the synthesis of a complementary DNA copy. Reverse transcriptase also has RNaseH activity for destruction of the RNA-template. Integrase binds both the viral cDNA generated by reverse transcriptase and the host DNA. Integrase processes the LTR before inserting the viral genome into the host DNA. Tat acts as a trans-activator during transcription to enhance initiation and elongation. The rev responsive element acts post-transcriptionally, regulating mRNA splicing and transport to the cytoplasm.

(113) Viral vectors, in general, comprise glycoproteins and the various glycoproteins may provide specific affinities. For instance, VSV-G peptides can increase transfection into myeloid cells. Alternatively, viral vectors can also have targeting moieties, such as antibodies, attached to their shell peptides. Targeting antibodies can be specific for antigens that are overexpressed on a tumor, for instance, like HER-2, PSA, CEA, M2-PK, and CA19-9.

(114) Other viral vector specificities are also known in the art and can be used to target particular populations of cells. For example, poxvirus vectors target to macrophages and dendritic cells.

(115) Lentiviral Vector System

(116) A lentiviral virion (particle) is expressed by a vector system encoding the necessary viral proteins to produce a virion (viral particle). There is at least one vector containing a nucleic acid sequence encoding the lentiviral pol proteins necessary for reverse transcription and integration, operably linked to a promoter. In another embodiment, the pol proteins are expressed by multiple vectors. There is also a vector containing a nucleic acid sequence encoding the lentiviral gag proteins necessary for forming a viral capsid operably linked to a promoter. In an embodiment, this gag nucleic acid sequence is on a separate vector than at least some of the pol nucleic acid sequence. In another embodiment, the gag nucleic acid is on a separate vector from all the pol nucleic acid sequences that encode pol proteins.

(117) Numerous modifications can be made to the vectors, which are used to create the particles to further minimize the chance of obtaining wild type revertants. These include, but are not limited to deletions of the U3 region of the LTR, tat deletions and matrix (MA) deletions.

(118) The gag, pol and env vector(s) do not contain nucleotides from the lentiviral genome that package lentiviral RNA, referred to as the lentiviral packaging sequence.

(119) The vector(s) forming the particle preferably do not contain a nucleic acid sequence from the lentiviral genome that expresses an envelope protein. Preferably, a separate vector that contains a nucleic acid sequence encoding an envelope protein operably linked to a promoter is used. This env

vector also does not contain a lentiviral packaging sequence. In one embodiment the env nucleic acid sequence encodes a lentiviral envelope protein.

(120) In another embodiment the envelope protein is not from the lentivirus, but from a different virus. The resultant particle is referred to as a pseudotyped particle. By appropriate selection of envelopes one can “infect” virtually any cell. For example, one can use an env gene that encodes an envelope protein that targets an endocytic compartment such as that of the influenza virus, VSV-G, alpha viruses (Semliki forest virus, Sindbis virus), arenaviruses (lymphocytic choriomeningitis virus), flaviviruses (tick-borne encephalitis virus, Dengue virus, hepatitis C virus, GB virus), rhabdoviruses (vesicular stomatitis virus, rabies virus), paramyxoviruses (mumps or measles) and orthomyxoviruses (influenza virus). Other envelopes that can preferably be used include those from Moloney Leukemia Virus such as MLV-E, MLV-A and GALV. These latter envelopes are particularly preferred where the host cell is a primary cell. Other envelope proteins can be selected depending upon the desired host cell. For example, targeting specific receptors such as a dopamine receptor can be used for brain delivery. Another target can be vascular endothelium. These cells can be targeted using a filovirus envelope. For example, the GP of Ebola, which by post-transcriptional modification become the GP, and GP.sub.2 glycoproteins. In another embodiment, one can use different lentiviral capsids with a pseudotyped envelope (for example, FIV or SHIV [U.S. Pat. No. 5,654,195]). A SHIV pseudotyped vector can readily be used in animal models such as monkeys.

(121) As detailed herein, a lentiviral vector system typically includes at least one helper plasmid comprising at least one of a gag, pol, or rev gene. Each of the gag, pol and rev genes may be provided on individual plasmids, or one or more genes may be provided together on the same plasmid. In one embodiment, the gag, pol, and rev genes are provided on the same plasmid (e.g., FIG. 2). In another embodiment, the gag and pol genes are provided on a first plasmid and the rev gene is provided on a second plasmid (e.g., FIG. 3). Accordingly, both 3-vector and 4-vector systems can be used to produce a lentivirus as described in the Examples section and elsewhere herein. The therapeutic vector, the envelope plasmid and at least one helper plasmid are transfected into a packaging cell line. A non-limiting example of a packaging cell line is the 293T/17 HEK cell line. When the therapeutic vector, the envelope plasmid, and at least one helper plasmid are transfected into the packaging cell line, a lentiviral particle is ultimately produced.

(122) In another aspect, a lentiviral vector system for expressing a lentiviral particle is disclosed. The system includes a lentiviral vector as described herein; an envelope plasmid for expressing an envelope protein optimized for infecting a cell; and at least one helper plasmid for expressing gag, pol, and rev genes, wherein when the lentiviral vector, the envelope plasmid, and the at least one helper plasmid are transfected into a packaging cell line, a lentiviral particle is produced by the packaging cell line, wherein the lentiviral particle is capable of inhibiting production of chemokine receptor CCR5 or targeting an HIV RNA sequence.

(123) In another aspect, and as detailed in FIG. 2, the lentiviral vector, which is also referred to herein as a therapeutic vector, can include the following elements: hybrid 5' long terminal repeat (RSV/5' LTR) (SEQ ID NOS: 11-12), Psi sequence (RNA packaging site) (SEQ ID NO: 13), RRE (Rev-response element) (SEQ ID NO: 14), cPPT (polypurine tract) (SEQ ID NO: 15), H1 promoter (SEQ ID NO: 16), FDPS shRNA (SEQ ID NOS: 1, 2, 3, 4), Woodchuck Post-Transcriptional Regulatory Element (WPRE) (SEQ ID NO: 17), and 3' Delta LTR (SEQ ID NO: 18). In another aspect, sequence variation, by way of substitution, deletion, addition, or mutation can be used to modify the sequences references herein.

(124) In another aspect, and as detailed herein, a helper plasmid has been designed to include the following elements: CAG promoter (SEQ ID NO: 19); HIV component gag (SEQ ID NO: 20); HIV component pol (SEQ ID NO: 21); HIV Int (SEQ ID NO: 22); HIV RRE (SEQ ID NO: 23); and HIV Rev (SEQ ID NO: 24). In another aspect, the helper plasmid may be modified to include a first helper plasmid for expressing the gag and pol genes, and a second and separate plasmid for expressing the rev gene. In another aspect, sequence variation, by way of substitution, deletion,

addition, or mutation can be used to modify the sequences references herein.

(125) In another aspect, and as detailed herein, an envelope plasmid has been designed to include the following elements being from left to right: RNA polymerase II promoter (CMV) (SEQ ID NO: 25) and vesicular stomatitis virus G glycoprotein (VSV-G) (SEQ ID NO: 26). In another aspect, sequence variation, by way of substitution, deletion, addition, or mutation can be used to modify the sequences references herein.

(126) In another aspect, the plasmids used for lentiviral packaging can be modified with similar elements and the intron sequences could potentially be removed without loss of vector function. For example, the following elements can replace similar elements in the plasmids that comprise the packaging system: Elongation Factor-1 (EF-1), phosphoglycerate kinase (PGK), and ubiquitin C (UbC) promoters can replace the CMV or CAG promoter. SV40 poly A and bGH poly A can replace the rabbit beta globin poly A. The HIV sequences in the helper plasmid can be constructed from different HIV strains or clades. The VSV-G glycoprotein can be substituted with membrane glycoproteins from feline endogenous virus (RD114), gibbon ape leukemia virus (GALV), Rabies (FUG), lymphocytic choriomeningitis virus (LCMV), influenza A fowl plague virus (FPV), Ross River alphavirus (RRV), murine leukemia virus 10A1 (MLV), or Ebola virus (EboV).

(127) Of note, lentiviral packaging systems can be acquired commercially (e.g., Lenti-vpak packaging kit from OriGene Technologies, Inc., Rockville, Md.), and can also be designed as described herein. Moreover, it is within the skill of a person skilled in the art to substitute or modify aspects of a lentiviral packaging system to improve any number of relevant factors, including the production efficiency of a lentiviral particle.

(128) Doses and Dosage Forms

(129) The disclosed vectors allow for short, medium, or long-term expression of genes or sequences of interest and episomal maintenance of the disclosed vectors. Accordingly, dosing regimens may vary based upon the condition being treated and the method of administration.

(130) In one embodiment, transduction vectors may be administered to a subject in need in varying doses. Specifically, a subject may be administered about $\geq 10^6$ infectious doses (where 1 dose is needed on average to transduce 1 target cell). More specifically, a subject may be administered about $\geq 10^7$, about $\geq 10^8$, about $\geq 10^9$, or about $\geq 10^{10}$ infectious doses, or any number of doses in-between these values. Upper limits of transduction vector dosing will be determined for each disease indication and will depend on toxicity/safety profiles for each individual product or product lot.

(131) Additionally, a vector of the present disclosure may be administered periodically, such as once or twice a day, or any other suitable time period. For example, vectors may be administered to a subject in need once a week, once every other week, once every three weeks, once a month, every other month, every three months, every six months, every nine months, once a year, every eighteen months, every two years, every thirty months, or every three years.

(132) In one embodiment, the disclosed vectors are administered as a pharmaceutical composition. In some embodiments, the pharmaceutical composition comprising the disclosed vectors can be formulated in a wide variety of dosage forms, including but not limited to nasal, pulmonary, oral, topical, or parenteral dosage forms for clinical application. Each of the dosage forms can comprise various solubilizing agents, disintegrating agents, surfactants, fillers, thickeners, binders, diluents such as wetting agents or other pharmaceutically acceptable excipients. The pharmaceutical composition comprising a vector can also be formulated for injection, insufflation, infusion, or intradermal exposure. For instance, an injectable formulation may comprise the disclosed vectors in an aqueous or non-aqueous solution at a suitable pH and tonicity.

(133) The disclosed vectors may be administered to a subject via direct injection into a tumor site or at a site of infection. In some embodiments, the vectors can be administered systemically. In some embodiments, the vectors can be administered via guided cannulation to tissues immediately surrounding the sites of tumor or infection.

(134) The disclosed vector compositions can be administered using any pharmaceutically acceptable method, such as intranasal, buccal, sublingual, oral, rectal, ocular, parenteral (intravenously, intradermally, intramuscularly, subcutaneously, intraperitoneally), pulmonary, intravaginal, locally administered, topically administered, topically administered after scarification, mucosally administered, via an aerosol, in semi-solid media such as agarose or gelatin, or via a buccal or nasal spray formulation.

(135) Further, the disclosed vector compositions can be formulated into any pharmaceutically acceptable dosage form, such as a solid dosage form, tablet, pill, lozenge, capsule, liquid dispersion, gel, aerosol, pulmonary aerosol, nasal aerosol, ointment, cream, semi-solid dosage form, a solution, an emulsion, and a suspension. Further, the composition may be a controlled release formulation, sustained release formulation, immediate release formulation, or any combination thereof. Further, the composition may be a transdermal delivery system.

(136) In some embodiments, the pharmaceutical composition comprising a vector can be formulated in a solid dosage form for oral administration, and the solid dosage form can be powders, granules, capsules, tablets or pills. In some embodiments, the solid dosage form can include one or more excipients such as calcium carbonate, starch, sucrose, lactose, microcrystalline cellulose or gelatin. In addition, the solid dosage form can include, in addition to the excipients, a lubricant such as talc or magnesium stearate. In some embodiments, the oral dosage form can be immediate release, or a modified release form. Modified release dosage forms include controlled or extended release, enteric release, and the like. The excipients used in the modified release dosage forms are commonly known to a person of ordinary skill in the art.

(137) In a further embodiment, the pharmaceutical composition comprising a vector can be formulated as a sublingual or buccal dosage form. Such dosage forms comprise sublingual tablets or solution compositions that are administered under the tongue and buccal tablets that are placed between the cheek and gum.

(138) In some embodiments, the pharmaceutical composition comprising a vector can be formulated as a nasal dosage form. Such dosage forms of the present invention comprise solution, suspension, and gel compositions for nasal delivery.

(139) In some embodiments, the pharmaceutical composition comprising a vector can be formulated in a liquid dosage form for oral administration, such as suspensions, emulsions or syrups. In some embodiments, the liquid dosage form can include, in addition to commonly used simple diluents such as water and liquid paraffin, various excipients such as humectants, sweeteners, aromatics or preservatives. In particular embodiments, the composition comprising vectors can be formulated to be suitable for administration to a pediatric patient.

(140) In some embodiment, the pharmaceutical composition can be formulated in a dosage form for parenteral administration, such as sterile aqueous solutions, suspensions, emulsions, non-aqueous solutions or suppositories. In some embodiments, the solutions or suspensions can include propylene glycol, polyethylene glycol, vegetable oils such as olive oil or injectable esters such as ethyl oleate.

(141) The dosage of the pharmaceutical composition can vary depending on the patient's weight, age, gender, administration time and mode, excretion rate, and the severity of disease.

(142) In some embodiments, the treatment of cancer is accomplished by guided direct injection of the disclosed vector constructs into tumors, using needle, or intravascular cannulation. In some embodiments, the disclosed vectors are administered into the cerebrospinal fluid, blood or lymphatic circulation by venous or arterial cannulation or injection, intradermal delivery, intramuscular delivery or injection into a draining organ near the site of disease.

(143) The following examples are given to illustrate the present invention. It should be understood, however, that the invention is not to be limited to the specific conditions or details described in these examples. All printed publications referenced herein are specifically incorporated by reference.

EXAMPLES

Example 1: Development of a Lentiviral Vector System

(144) A lentiviral vector system was developed as summarized in FIG. 4 (circularized form). Lentiviral particles were produced in 293T/17 HEK cells (purchased from American Type Culture Collection, Manassas, Va.) following transfection with the therapeutic vector, the envelope plasmid, and the helper plasmid. The transfection of 293T/17 HEK cells, which produced functional viral particles, employed the reagent Poly(ethylenimine) (PEI) to increase the efficiency of plasmid DNA uptake. The plasmids and DNA were initially added separately in culture medium without serum in a ratio of 3:1 (mass ratio of PEI to DNA). After 2-3 days, cell medium was collected and lentiviral particles were purified by high-speed centrifugation and/or filtration followed by anion-exchange chromatography. The concentration of lentiviral particles can be expressed in terms of transducing units/ml (TU/ml). The determination of TU was accomplished by measuring HIV p24 levels in culture fluids (p24 protein is incorporated into lentiviral particles), measuring the number of viral DNA copies per cell by quantitative PCR, or by infecting cells and using light (if the vectors encode luciferase or fluorescent protein markers).

(145) As mentioned above, a 3-vector system (i.e., a 2-vector lentiviral packaging system) was designed for the production of lentiviral particles. A schematic of the 3-vector system is shown in FIG. 2. Briefly, and with reference to FIG. 2, the top-most vector is a helper plasmid, which, in this case, includes Rev. The vector appearing in the middle of FIG. 2 is the envelope plasmid. The bottom-most vector is the therapeutic vector, as described herein.

(146) Referring more specifically to FIG. 2, the Helper plus Rev plasmid includes a CAG enhancer (SEQ ID NO: 27); a CAG promoter (SEQ ID NO: 19); a chicken beta actin intron (SEQ ID NO: 28); a HIV gag (SEQ ID NO: 20); a HIV Pol (SEQ ID NO: 21); a HIV Int (SEQ ID NO: 22); a HIV RRE (SEQ ID NO: 23); a HIV Rev (SEQ ID NO: 24); and a rabbit beta globin poly A (SEQ ID NO: 29).

(147) The Envelope plasmid includes a CMV promoter (SEQ ID NO: 25); a beta globin intron (SEQ ID NO: 30); a VSV-G (SEQ ID NO: 28); and a rabbit beta globin poly A (SEQ ID NO: 31).

(148) Synthesis of a 2-Vector Lentiviral Packaging System Including Helper (Plus Rev) and Envelope Plasmids.

(149) Materials and Methods:

(150) Construction of the helper plasmid: The helper plasmid was constructed by initial PCR amplification of a DNA fragment from the pNL4-3 HIV plasmid (NIH Aids Reagent Program) containing Gag, Pol, and Integrase genes. Primers were designed to amplify the fragment with EcoRI and NotI restriction sites which could be used to insert at the same sites in the pCDNA3 plasmid (Invitrogen). The forward primer was (5'-TAAGCAGAATTC

ATGAATTTGCCAGGAAGAT-3') (SEQ ID NO: 32) and reverse primer was (5'-CCATACAATGAATGGACACTAGGCGGCCGCACGAAT-3') (SEQ ID NO: 33).

(151) The sequence for the Gag, Pol, Integrase fragment was as follows:

(152) TABLE-US-00001 (SEQ ID NO: 34)

GAATTCATGAATTTGCCAGGAAGATGGAAACCAAAAATGATAGGGGGAAT
TGGAGGTTTTATCAAAGTAAGACAGTATGATCAGATACTCATAGAAATCT
GCGGACATAAAGCTATAGGTACAGTATTAGTAGGACCTACACCTGTCAAC
ATAATTGGAAGAAATCTGTTGACTCAGATTGGCTGCACTTTAAATTTTCC
CATTAGTCCTATTGAGACTGTACCAGTAAAATTAAAGCCAGGAATGGATG
GCCCCAAAAGTTAAACAATGGCCATTGACAGAAGAAAAAATAAAAGCATT
GTAGAAATTTGTACAGAAATGGAAAAGGAAGGAAAAATTTCAAAAATTGG
GCCTGAAAATCCATACAATACTCCAGTATTTGCCATAAAGAAAAAAGACA
GTACTAAATGGAGAAAATTAGTAGATTTTCAGAGAACTTAATAAGAGAACT
CAAGATTTCTGGGAAGTTCAATTAGGAATACCACATCCTGCAGGGTTAA
ACAGAAAAAATCAGTAACAGTACTGGATGTGGGCGATGCATATTTTTCAG

TTCCCTTAGATAAAGACTTTCAGTACTGCATTTACCATACCTAGT
 ATAAACAATGAGACACCAGGGATTAGATATCAGTACAATGTGCTTCCACA
 GGGATGGAAAGGATCACCAGCAATATTCCAGTGTAGCATGACAAAAATCT
 TAGAGCCTTTTAGAAAAACAAAATCCAGACATAGTCATCTATCAATACATG
 GATGATTTGTATGTAGGATCTGACTTAGAAATAGGGCAGCATAGAACAAA
 AATAGAGGAACCTGAGACAACATCTGTTGAGGTGGGGATTACCACACCAG
 AAAAAAACATCAGAAAGAACCTCCATTTCCTTTGGATGGGTATGAACTC
 CATCCTGATAAATGGACAGTACAGCCTATAGTGCTGCCAGAAAAGGACAG
 CTGGACTGTCAATGACATACAGAAATTAGTGGGAAAATTGAATTGGGCAA
 GTCAGATTTATGCAGGGATTAAAGTAAGGCAATTATGTAAACTTCTTAGG
 GGAACCAAAGCACTAACAGAAGTAGTACCCTAACAGAAGAAGCAGAGCT
 AGAACTGGCAGAAAACAGGGAGATTCTAAAAGAACCGGTACATGGAGTGT
 ATTATGACCCATCAAAGACTTAATAGCAGAAATACAGAAGCAGGGGCAA
 GGCCAATGGACATATCAAATTTATCAAGAGCCATTAAAAATCTGAAAAC
 AGGAAAGTATGCAAGAATGAAGGGTGCCCACTAATGATGTGAAACAAT
 TAACAGAGGCAGTACAAAAAATAGCCACAGAAAGCATAGTAATATGGGGA
 AAGACTCCTAAATTTAAATTACCCATACAAAAGGAAACATGGGAAGCATG
 GTGGACAGAGTATTGGCAAGCCACCTGGATTCTGAGTGGGAGTTTGTCA
 ATACCCCTCCCTTAGTGAAGTTATGGTACCAGTTAGAGAAAGAACCCATA
 ATAGGAGCAGAACTTTCTATGTAGATGGGGCAGCCAATAGGGAACTAA
 ATTAGGAAAAGCAGGATATGTAAGTACAGAGGAAGACAAAAAGTTGTCC
 CCTAACGGACACAACAAATCAGAAGACTGAGTTACAAGCAATTCATCTA
 GCTTTGCAGGATTTCGGGATTAGAAGTAAACATAGTGACAGACTCACAATA
 TGCATTGGGAATCATTCAAGCACAACCAGATAAGAGTGAATCAGAGTTAG
 TCAGTCAAATAATAGAGCAGTTAATAAAAAAGGAAAAAGTCTACCTGGCA
 TGGGTACCAGCACACAAAGGAATTGGAGGAAATGAACAAGTAGATAAATT
 GGTCAGTGCTGGAATCAGGAAAGTACTATTTTATAGATGGAATAGATAAGG
 CCCAAGAAGAACATGAGAAATATCACAGTAATTGGAGAGCAATGGCTAGT
 GATTTTAACCTACCACCTGTAGTAGCAAAAGAAATAGTAGCCAGCTGTGA
 TAAATGTCAGCTAAAAGGGGAAGCCATGCATGGACAAGTAGACTGTAGCC
 CAGGAATATGGCAGCTAGATTGTACACATTTAGAAGGAAAAGTTATCTTG
 GTAGCAGTTCATGTAGCCAGTGGATATATAGAAGCAGAAGTAATTCCAGC
 AGAGACAGGGCAAGAAACAGCATACTTCCTCTTAAAATTAGCAGGAAGAT
 GGCCAGTAAAAACAGTACATACAGACAATGGCAGCAATTTACCAGTACT
 ACAGTTAAGGCCGCTGTTGGTGGGCGGGGATCAAGCAGGAATTTGGCAT
 TCCCTACAATCCCCAAAGTCAAGGAGTAATAGAATCTATGAATAAAGAAT
 TAAAGAAAATTATAGGACAGGTAAGAGATCAGGCTGAACATCTTAAGACA
 GCAGTACAAATGGCAGTATTCATCCACAATTTTAAAAGAAAAGGGGGGAT
 TGGGGGGTACAGTGCAGGGGAAAGAATAGTAGACATAATAGCAACAGACA
 TACAACTAAAGAATTACAAAAACAAATTACAAAAATTCAAATTTTCGG
 GTTTATTACAGGGACAGCAGAGATCCAGTTTGGAAGGACCAGCAAAGCT
 CCTCTGGAAAGGTGAAGGGGCAGTAGTAATACAAGATAATAGTGACATAA
 AAGTAGTGCCAAGAAGAAAAGCAAAGATCATCAGGGATTATGGAAAACAG
 ATGGCAGGTGATGATTGTGTGGCAAGTAGACAGGATGAGGATTAA

(153) Next, a DNA fragment containing the Rev, RRE, and rabbit beta globin poly A sequence with XbaI and XmaI flanking restriction sites was synthesized by MWG Operon. The DNA fragment was then inserted into the plasmid at the XbaI and XmaI restriction sites. The DNA sequence was as follows:

(154) TABLE-US-00002 (SEQ ID NO: 35)

TCTAGAATGGCAGGAAGAAGCGGAGACAGCGACGAAGAGCTCATCAGAAC

AGTCACTCATCTACCTATCTCTCAAAACCCACCTCCCCAATCCCCG
 AGGGGACCCGACAGGCCCGAAGGAATAGAAGAAGAAGGTGGAGAGAGAGA
 CAGAGACAGATCCATTCTGATTAGTGAACGGATCCTTGGCACTTATCTGGG
 ACGATCTGCGGAGCCTGTGCCTCTTCAGCTACCACCGCTTGAGAGACTTA
 CTCTTGATTGTAACGAGGATTGTGGAACCTTCTGGGACGCAGGGGGGTGGGA
 AGCCCTCAAATATTGGTGGAAATCTCCTACAATATTGGAGTCAGGAGCTAA
 AGAATAGAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACT
 ATGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTC
 TGGTATAGTGCAGCAGCAGAACAATTTGCTGAGGGCTATTGAGGCGCAAC
 AGCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGA
 ATCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTAGATCTTTT
 TCCCTCTGCCAAAAATTATGGGGACATCATGAAGCCCCTTGAGCATCTGA
 CTTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTTGGAAT
 TTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTAAA
 ACATCAGAATGAGTATTTGGTTTAGAGTTTGGCAACATATGCCATATGCT
 GGCTGCCATGAACAAAGGTGGCTATAAAGAGGTCATCAGTATATGAAACA
 GCCCCCTGCTGTCCATTCTTATTCCATAGAAAAGCCTTGACTTGAGGTT
 AGATTTTTTTTTATATTTTGTGTTTGTGTTATTTTTTTCTTTAACATCCCTA
 AAATTTTCCTTACATGTTTTACTAGCCAGATTTTTCTCCTCTCCTGACT
 ACTCCCAGTCATAGCTGTCCCTCTTCTCTTATGAAGATCCCTCGACCTGC
 AGCCCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTG
 TTATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
 AAGCCTGGGGTGCCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGC
 TCACTGCCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAGCGGATCCGCAT
 CTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCCCGCC
 CCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCCATGGCTGACTAATTT
 TTTTTATTTATGCAGAGGCCGAGGCCGCCTCGGCCTCTGAGCTATTCCAG
 AAGTAGTGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTAAC
 TTGTTTATTGCACTTATAATGGTTACAAATAAAGCAATAGCATCACAAA
 TTTCACAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGTTTGTCCA
 AACTCATCAATGTATCTTATCAGCGGCCGCCCCGGG

(155) Finally, the CMV promoter of pCDNA3.1 was replaced with the CAG enhancer/promoter plus a chicken beta actin intron sequence. A DNA fragment containing the CAG enhancer/promoter/intron sequence with MluI and EcoRI flanking restriction sites was synthesized by MWG Operon. The DNA fragment was then inserted into the plasmid at the MluI and EcoRI restriction sites. The DNA sequence was as follows:

(156) TABLE-US-00003 (SEQ ID NO: 36)
 ACGCGTTAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCATAGCC
 CATATATGGAGTTCCGCGTTACATAACTTACGGTAAATGGCCCGCCTGGC
 TGACCGCCCAACGACCCCCGCCCATTGACGTCAATAATGACGTATGTTCC
 CATAGTAACGCCAATAGGGACTTTCCATTGACGTCAATGGGTGGACTATT
 TACGGTAAACTGCCCACTTGGCAGTACATCAAGTGTATCATATGCCAAGT
 ACGCCCCCTATTGACGTCAATGACGGTAAATGGCCCGCCTGGCATTATGC
 CCAGTACATGACCTTATGGGACTTTCTACTTGGCAGTACATCTACGTAT
 TAGTCATCGCTATTACCATGGGTTCGAGGTGAGCCCCACGTTCTGCTTCAC
 TCTCCCCATCTCCCCCCCCCTCCCCACCCCCAATTTTGTATTATTTATTT
 TTTAATTATTTTGTGCAGCGATGGGGGCGGGGGGGGGGGGGGGCGCGGCC
 AGGCGGGGCGGGGCGGGGCGAGGGGCGGGGCGGGGCGAGGCGGAGAGGTG
 CGGCGGCAGCCAATCAGAGCGGCGCGCTCCGAAAGTTTCTTTTATGGCG
 AGGCGGCGGCGGGCGGGCGGCCCTATAAAAAGCGAAGCGCGCGGGCGGG

AGTCGCTGCTTCCTTCGCCCCGTGCCCGCTCCCGCCGCTCGCGCC
GCCCCGCCCCGGCTCTGACTGACCGCGTTACTCCCACAGGTGAGCGGGCGG
GACGGCCCTTCTCCTCCGGGCTGTAATTAGCGCTTGGTTTAATGACGGCT
CGTTTCTTTTCTGTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCC
CTTTGTGCGGGGGGGAGCGGCTCGGGGGGGTGCCTGCGTGTGTGTGTGCGT
GGGGAGCGCCGCGTGCGGCCCGCGCTGCCCGGCGGCTGTGAGCGCTGCGG
GCGCGGCGCGGGGCTTTGTGCGCTCCGCGTGTGCGCGAGGGGAGCGCGGC
CGGGGGCGGTGCCCCGCGGTGCGGGGGGGCTGCGAGGGGAACAAAGGCTG
CGTGCGGGGTGTGTGCGTGGGGGGGTGAGCAGGGGGTGTGGGCGCGGCGG
TCGGGCTGTAACCCCCCTGCACCCCCCTCCCCGAGTTGCTGAGCACGG
CCCGGCTTCGGGTGCGGGGCTCCGTGCGGGGCGTGCGCGGGGGCTCGCCG
TGCCGGGCGGGGGGTGGCGGCAGGTGGGGGTGCCGGGCGGGGCGGGGCCG
CCTCGGGCCGGGGAGGGCTCGGGGGAGGGGCGCGGCGGCCCGGAGCGCC
GGCGGCTGTGAGGCGCGGCGAGCCGCAGCCATTGCCTTTTATGGTAATC
GTGCGAGAGGGCGCAGGGACTTCCTTTGTCCCAAATCTGGCGGAGCCGAA
ATCTGGGAGGCGCCGCCGCACCCCCCTCTAGCGGGCGCGGGCGAAGCGGTG
CGGCGCCGGCAGGAAGGAAATGGGCGGGGAGGGCCTTCGTGCGTCGCCGC
GCCGCCGTCCCCTTCTCCATCTCCAGCCTCGGGGCTGCCGCAGGGGGACG
GCTGCCTTCGGGGGGGACGGGGCAGGGCGGGGTTCGGCTTCTGGCGTGTG
ACCGGCGGGAATTC

Construction of the VSV-G Envelope Plasmid:

(157) The vesicular stomatitis Indiana virus glycoprotein (VSV-G) sequence was synthesized by MWG Operon with flanking EcoRI restriction sites. The DNA fragment was then inserted into the pCDNA3.1 plasmid (Invitrogen) at the EcoRI restriction site and the correct orientation was determined by sequencing using a CMV specific primer. The DNA sequence was as follows:

(158) TABLE-US-00004 (SEQ ID NO: 37)

GAATTCATGAAGTGCCTTTTGTACTTAGCCTTTTATTTCATTGGGGTGAA
TTGCAAGTTCACCATAGTTTTTCCACACAACCAAAAAGGAAACTGGAAAA
ATGTTCTTCTAATTACCATATTGCCCCGTCAAGCTCAGATTTAAATTGG
CATAATGACTTAATAGGCACAGCCTTACAAGTCAAAATGCCCAAGAGTCA
CAAGGCTATTCAAGCAGACGGTTGGATGTGTGTCATGCTTCCAAATGGGTCA
CTACTTGTGATTTCCGCTGGTATGGACCGAAGTATATAACACATTCCATC
CGATCCTTCACTCCATCTGTAGAACAATGCAAGGAAAGCATTGAACAAAC
GAAACAAGGAACTTGGCTGAATCCAGGCTTCCCTCCTCAAAGTTGTGGAT
ATGCAACTGTGACGGATGCCGAAGCAGTGATTGTCCAGGTGACTCCTCAC
CATGTGCTGGTTGATGAATACACAGGAGAATGGGTTGATTCACAGTTCAT
CAACGGAAAATGCAGCAATTACATATGCCCCACTGTCCATAACTCTACAA
CCTGGCATTCTGACTATAAGGTCAAAGGGCTATGTGATTCTAACCTCATT
TCCATGGACATCACCTTCTTCTCAGAGGACGGAGAGCTATCATCCCTGGG
AAAGGAGGGCACAGGGTTCAGAAGTAACTACTTTGCTTATGAAACTGGAG
GCAAGGCCTGCAAAATGCAATACTGCAAGCATTGGGGAGTCAGACTCCCA
TCAGGTGTCTGGTTCGAGATGGCTGATAAGGATCTCTTTGCTGCAGCCAG
ATTCCCTGAATGCCCAGAAGGGTCAAGTATCTCTGCTCCATCTCAGACCT
CAGTGGATGTAAGTCTAATTCAGGACGTTGAGAGGATCTTGGATTATTCC
CTCTGCCAAGAAACCTGGAGCAAAATCAGAGCGGGTCTTCCAATCTCTCC
AGTGGATCTCAGCTATCTTGCTCCTAAAAACCCAGGAACCGGTCCTGCTT
TCACCATAATCAATGGTACCCTAAAATACTTTGAGACCAGATACATCAGA
GTCGATATTGCTGCTCCAATCCTCTCAAGAATGGTCGGAATGATCAGTGG
AACTACCACAGAAAGGGAACCTGTGGGATGACTGGGCACCATATGAAGACG
TGGA AATTGGACCCAATGGAGTTCTGAGGACCAGTTCAGGATATAAGTTT

CCTTTATACATGATTGGACATGGTATGTTGGACTCCGATCTTCATCTTAG
CTCAAAGGCTCAGGTGTTTCGAACATCCTCACATTCAAGACGCTGCTTCGC
AACTTCCTGATGATGAGAGTTTATTTTTTTGGTGATACTGGGCTATCCAAA
AATCCAATCGAGCTTGTAGAAGGTTGGTTCAGTAGTTGGAAAAGCTCTAT
TGCCTCTTTTTTCTTTATCATAGGGTTAATCATTGGACTATTCTTGGTTC
TCCGAGTTGGTATCCATCTTTGCATTAAATTAAGCACACCAAGAAAAGA
CAGATTTATACAGACATAGAGATGAGAATTC

(159) A 4-vector system (i.e., a 3-vector lentiviral packaging system) has also been designed and produced using the methods and materials described herein. A schematic of the 4-vector system is shown in FIG. 3. Briefly, and with reference to FIG. 3, the top-most vector is a helper plasmid, which, in this case, does not include Rev. The vector second from the top is a separate Rev plasmid. The vector second from the bottom is the envelope plasmid. The bottom-most vector is the previously described therapeutic vector.

(160) Referring, in part, to FIG. 2, the Helper plasmid includes a CAG enhancer (SEQ ID NO: 27); a CAG promoter (SEQ ID NO: 19); a chicken beta actin intron (SEQ ID NO: 28); a HIV gag (SEQ ID NO: 20); a HIV Pol (SEQ ID NO: 21); a HIV Int (SEQ ID NO: 22); a HIV RRE (SEQ ID NO: 23); and a rabbit beta globin poly A (SEQ ID NO: 29).

(161) The Rev plasmid includes a RSV promoter and a HIV Rev (SEQ ID NO: 38); and a rabbit beta globin poly A (SEQ ID NO: 29).

(162) The Envelope plasmid includes a CMV promoter (SEQ ID NO: 25); a beta globin intron (SEQ ID NO: 30); a VSV-G (SEQ ID NO: 28); and a rabbit beta globin poly A (SEQ ID NO: 29).

(163) Synthesis of a 3-Vector Lentiviral Packaging System Including Helper, Rev, and Envelope Plasmids.

(164) Materials and Methods:

(165) Construction of the Helper Plasmid without Rev:

(166) The Helper plasmid without Rev was constructed by inserting a DNA fragment containing the RRE and rabbit beta globin poly A sequence. This sequence was synthesized by MWG Operon with flanking XbaI and XmaI restriction sites. The RRE/rabbit poly A beta globin sequence was then inserted into the Helper plasmid at the XbaI and XmaI restriction sites. The DNA sequence is as follows:

(167) TABLE-US-00005 (SEQ ID NO: 65)

TCTAGAAGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAGCACTA
TGGGCGCAGCGTCAATGACGCTGACGGTACAGGCCAGACAATTATTGTCT
GGTATAGTGCAGCAGCAGAACAAATTTGCTGAGGGCTATTGAGGCGCAACA
GCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGAA
TCCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCTAGATCTTTTT
CCCTCTGCCAAAAATTATGGGGACATCATGAAGCCCCTTGAGCATCTGAC
TTCTGGCTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTTGGAATT
TTTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTAAAA
CATCAGAATGAGTATTTGGTTTAGAGTTTGGCAACATATGCCATATGCTG
GCTGCCATGAACAAAGGTGGCTATAAAGAGGTCATCAGTATATGAAACAG
CCCCCTGCTGTCCATTCTTATTCCATAGAAAAGCCTTGACTTGAGGTTA
GATTTTTTTTATATTTTGTGTTATTTTTTTCTTTAACATCCCTAA
AATTTTCCTTACATGTTTTACTAGCCAGATTTTTCCTCCTCTCCTGACTA
CTCCCAGTCATAGCTGTCCCTCTTCTCTTATGAAGATCCCTCGACCTGCA
GCCCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTGT
TATCCGCTCACAATTCCACACAACATACGAGCCGGAAGCATAAAGTGTA
AGCCTGGGGTGCTAATGAGTGAGCTAACTCACATTAATTGCGTTGCGCT
CACTGCCCCGCTTTCCAGTCGGGAAACCTGTCGTGCCAGCGGATCCGCATC
TCAATTAGTCAGCAACCATAGTCCCGCCCCTAACTCCGCCCATCCCGCCC

CTAACCCTCCCGAGTTCCCGCCCATTCGCGCCCATGGCTGACTAATTTT
TTTTATTTATGCAGAGGCCGAGGCCGCTCGGCCTCTGAGCTATTCCAGA
AGTAGTGAGGAGGCTTTTTTGGAGGCCCTAGGCTTTTGCAAAAAGCTAACT
TGTTTATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCATCACAAAT
TTCACAAATAAAGCATTTTTTTCCTGACTGCAATTCTAGTTGTGGTTTGTCCAA
ACTCATCAATGTATCTTATCACCCGGG

Construction of the Rev Plasmid:

(168) The RSV promoter and HIV Rev sequence was synthesized as a single DNA fragment by MWG Operon with flanking MfeI and XbaI restriction sites. The DNA fragment was then inserted into the pCDNA3.1 plasmid (Invitrogen) at the MfeI and XbaI restriction sites in which the CMV promoter is replaced with the RSV promoter. The DNA sequence was as follows:

(169) TABLE-US-00006 (SEQ ID NO: 38)

CAATTGCGATGTACGGGGCCAGATATACGCGTATCTGAGGGGACTAGGGTG
TGTTTAGGGCGAAAAGCGGGGCTTCGGTTGTACGCGGTTAGGAGTCCCCTC
AGGATATAGTAGTTTCGCTTTTGCATAGGGAGGGGGAAATGTAGTCTTAT
GCAATACACTTGTAGTCTTGAACATGGTAACGATGAGTTAGCAACATGC
CTTACAAGGAGAGAAAAAGCACCGTGCATGCCGATTGGTGGAAAGTAAGGT
GGTACGATCGTGCCTTATTAGGAAGGCAACAGACAGGTCTGACATGGATT
GGACGAACCACTGAATTCGCGATTGCAGAGATAATTGTATTTAAGTGCCT
AGCTCGATACAATAAACGCCATTTGACCATTACCCACATTGGTGTGCACC
TCCAAGCTCGAGCTCGTTTAGTGAACCGTCAGATCGCCTGGAGACGCCAT
CCACGCTGTTTTGACCTCCATAGAAGACACCGGGACCGATCCAGCCTCCC
CTCGAAGCTAGCGATTAGGCATCTCCTATGGCAGGAAGAAGCGGAGACAG
CGACGAAGAACTCCTCAAGGCAGTCAGACTCATCAAGTTTCTCTATCAA
GCAACCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAAGGAATAGA
AGAAGAAGGTGGAGAGAGAGACAGAGACAGATCCATTGATTAGTGAACG
GATCCTTAGCACTTATCTGGGACGATCTGCGGAGCCTGTGCCTCTTCAGC
TACCACCGCTTGAGAGACTTACTCTTGATTGTAACGAGGATTGTGGAAGT
TCTGGGACGCAGGGGGTGGGAAGCCCTCAAATATTGGTGGAAATCTCCTAC
AATATTGGAGTCAGGAGCTAAAGAATAGTCTAGA

(170) The plasmids for the 2-vector and 3-vector packaging systems could be modified with similar elements and the intron sequences could potentially be removed without loss of vector function. For example, the following elements could replace similar elements in the 2-vector and 3-vector packaging system:

(171) Promoters: Elongation Factor-1 (EF-1) (SEQ ID NO: 39), phosphoglycerate kinase (PGK) (SEQ ID NO: 40), and ubiquitin C (UbC) (SEQ ID NO: 41) can replace the CMV (SEQ ID NO: 25) or CAG promoter (SEQ ID NO: 19). These sequences can also be further varied by addition, substitution, deletion or mutation.

(172) Poly A sequences: SV40 poly A (SEQ ID NO: 42) and bGH poly A (SEQ ID NO: 43) can replace the rabbit beta globin poly A (SEQ ID NO: 29). These sequences can also be further varied by addition, substitution, deletion or mutation.

(173) HIV Gag, Pol, and Integrase sequences: The HIV sequences in the Helper plasmid can be constructed from different HIV strains or clades. For example, HIV Gag (SEQ ID NO: 20); HIV Pol (SEQ ID NO: 21); and HIV Int (SEQ ID NO: 22) from the Bal strain can be interchanged with the gag, pol, and int sequences contained in the helper/helper plus Rev plasmids as outlined herein. These sequences can also be further varied by addition, substitution, deletion or mutation.

(174) Envelope: The VSV-G glycoprotein can be substituted with membrane glycoproteins from feline endogenous virus (RD114) (SEQ ID NO: 44), gibbon ape leukemia virus (GALV) (SEQ ID NO: 45), Rabies (FUG) (SEQ ID NO: 46), lymphocytic choriomeningitis virus (LCMV) (SEQ ID NO: 47), influenza A fowl plague virus (FPV) (SEQ ID NO: 48), Ross River alphavirus (RRV)

(SEQ ID NO: 49), murine leukemia virus 10A1 (MLV) (SEQ ID NO: 50), or Ebola virus (EboV) (SEQ ID NO: 51). Sequences for these envelopes are identified in the sequence portion herein. Further, these sequences can also be further varied by addition, substitution, deletion or mutation. (175) In summary, the 3-vector versus 4-vector systems can be compared and contrasted, in part, as follows. The 3-vector lentiviral vector system contains: 1. Helper plasmid: HIV Gag, Pol, Integrase, and Rev/Tat; 2. Envelope plasmid: VSV-G/FUG envelope; and 3. Therapeutic vector: RSV 5'LTR, Psi Packaging Signal, Gag fragment, RRE, Env fragment, cPPT, WPRE, and 3' delta LTR. The 4-vector lentiviral vector system contains: 1. Helper plasmid: HIV Gag, Pol, and Integrase; 2. Rev plasmid: Rev; 3. Envelope plasmid: VSV-G/FUG envelope; and 4. Therapeutic vector: RSV 5'LTR, Psi Packaging Signal, Gag fragment, RRE, Env fragment, cPPT, WPRE, and 3' delta LTR. Sequences corresponding with the above elements are identified in the sequence listings portion herein.

Example 2: Development of a Lentiviral Vector that Expresses FDPS

(176) The purpose of this Example was to develop an FDPS lentivirus vector.

(177) Inhibitory RNA Design: The sequence of *Homo sapiens* Farnesyl diphosphate synthase (FDPS) (NM_002004.3) mRNA was used to search for potential siRNA or shRNA candidates to knockdown FDPS levels in human cells. Potential RNA interference sequences were chosen from candidates selected by siRNA or shRNA design programs such as from GPP Web Portal hosted by the Broad Institute (<http://portals.broadinstitute.org/gpp/public/>) or the BLOCK-iT RNAi Designer from Thermo Scientific (<https://rnaidesigner.thermofisher.com/rnaiexpress/>). Individual selected shRNA sequences were inserted into a lentiviral vector immediately 3 prime to a RNA polymerase III promoter such as H1 (SEQ ID NO: 16), U6 (SEQ ID NO: 52), or 7SK (SEQ ID NO: 53) to regulate shRNA expression. These lentivirus shRNA constructs were used to transduce cells and measure the change in specific mRNA levels. The shRNA most potent for reducing mRNA levels were embedded individually within a microRNA backbone to allow for expression by either the EF-1alpha or CMV RNA polymerase II promoters. The microRNA backbone was selected from mirbase.org. RNA sequences were also synthesized as synthetic siRNA oligonucleotides and introduced directly into cells without using a lentiviral vector.

(178) Vector Construction: For FDPS shRNA, oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by Eurofins MWG Operon. Overlapping sense and antisense oligonucleotide sequences were mixed and annealed during cooling from 70 degrees Celsius to room temperature. The lentiviral vector was digested with the restriction enzymes BamHI and EcoRI for one hour at 37 degrees Celsius. The digested lentiviral vector was purified by agarose gel electrophoresis and extracted from the gel using a DNA gel extraction kit from Thermo Scientific. The DNA concentrations were determined and vector to oligo (3:1 ratio) were mixed, allowed to anneal, and ligated. The ligation reaction was performed with T4 DNA ligase for 30 minutes at room temperature. 2.5 microliters of the ligation mix were added to 25 microliters of STBL3 competent bacterial cells. Transformation was achieved after heat-shock at 42 degrees Celsius. Bacterial cells were spread on agar plates containing ampicillin and drug-resistant colonies (indicating the presence of ampicillin-resistance plasmids) were recovered and expanded in LB broth. To check for insertion of the oligo sequences, plasmid DNA was extracted from harvested bacteria cultures with the Thermo Scientific DNA mini prep kit. Insertion of shRNA sequences in the lentiviral vector was verified by DNA sequencing using a specific primer for the promoter used to regulate shRNA expression. Using the following target sequences, exemplary shRNA sequences were determined to knock-down FDPS:

(179) TABLE-US-00007 (FDPS target sequence #1; SEQ ID NO: 54)
GTCCTGGAGTACAATGCCATT; (FDPS shRNA sequence #1; SEQ ID NO: 1)
GTCCTGGAGTACAATGCCATTCTCGAGAATGGCATTGTACTCCAGGACTT TTT;
(FDPS target sequence #2; SEQ ID NO: 55) GCAGGATTTCGTTCAGCACTT;
(FDPS shRNA sequence #2; SEQ ID NO: 2)

GCAGGATTCGTTTCAGCACTTCTCGAGAAGTGCTGAACGAAATCCTGCTT TTT;
 (FDPS target sequence #3; SEQ ID NO: 56) GCCATGTACATGGCAGGAATT;
 (FDPS shRNA sequence #3; SEQ ID NO: 3)
 GCCATGTACATGGCAGGAATTCTCGAGAATTCCTGCCATGTACATGGCTT TTT;
 (FDPS target sequence #4; SEQ ID NO: 57)
 GCAGAAGGAGGCTGAGAAAGT; and (FDPS shRNA sequence #4; SEQ ID
 NO: 4) GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCCTCCTTCTGCTT
 TTT.

(180) shRNA sequences were then assembled into a synthetic microRNA (miR) under control of the EF-1 alpha promoter. Briefly, a miR hairpin sequences, such as miR30, miR21, or miR185 as detailed below, was obtained from mirbase.org. The 19-22mer shRNA target sequence was used to construct the synthetic miR sequence. The miR sequence was arranged as an anti-sense-target-sequence-hairpin loop sequence (specific for each microRNA)-sense target sequence.

(181) The following miR sequences were developed:

(182) TABLE-US-00008 (miR30 FDPS sequence #1; SEQ ID NO: 5)
 AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGC
 GTGAAGCCACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGCCT
 CGGACTTCAAGGGGCT (miR30 FDPS sequence #2; SEQ ID NO: 6)
 AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGC
 GTGAAGCCACAGATGGCAGAAGGGCTGAGAAAGTGCTGCCTACTGCCTCG
 GACTTCAAGGGGCT (miR30 FDPS sequence #3; SEQ ID NO: 7)
 TGCTGTTGACAGTGAGCGACTTTCTCAGCCTCCTTCTGCGTGAAGCCACA
 GATGGCAGAAGGAGGCTGAGAAAGTTGCCTACTGCCTCGGA (miR155 FDPS
 sequence #1; SEQ ID NO: 8)
 CCTGGAGGCTTGCTGAAGGCTGTATGCTGACTTTCTCAGCCTCCTTCTGC
 TTTTGGCCACTGACTGAGCAGAAGGGCTGAGAAAGTCAGGACACAAGGCC
 TGTTACTAGCACTCA (miR21 FDPS sequence #1; SEQ ID NO: 9)
 CATCTCCATGGCTGTACCACCTTGTCTGGGACTTTCTCAGCCTCCTTCTGC
 CTGTTGAATCTCATGGCAGAAGGAGGCGAGAAAGTCTGACATTTTGGTAT
 CTTTCATCTGACCA (miR185 FDPS sequence #1; SEQ ID NO: 10)
 GGGCCTGGCTCGAGCAGGGGGCGAGGGATACTTTCTCAGCCTCCTTCTGC
 TGGTCCCCTCCCCGCAGAAGGAGGCTGAGAAAGTCCTTCCCTCCCAATGA
 CCGCGTCTTCGTCG

Example 3—Knock-Down of FDPS for 3 Days in THP1 Monocytic Leukemia by shRNA #4

(183) This Example illustrates that knock-down of FDPS in THP1 monocytic leukemia cells by lentiviral (LV)-expressing FDPS shRNA #4 stimulates TNF- α expression in gamma delta T cells, as shown in FIG. 5.

(184) THP1 cells (1×10^5 cells) were transduced with LV-control or LV-FDPS shRNA #4 for 3 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced THP-1 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 3.1% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 stimulated 5%. With zoledronic acid treatment, LV-control stimulated 7.2% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 stimulated 56.2%.

Example 4—Knock-Down of FDPS for 14 Days in THP1 Leukemia Cells by shRNA #4

(185) This Example illustrates that Knock-down of FDPS for 14 days in THP1 leukemia cells by

lentiviral (LV)-expressing FDPS shRNA #4 stimulates TNF- α expression in GD T cells, as shown in FIG. 6.

(186) THP1 cells (1×10^5 cells) were transduced with LV-control or LV-FDPS shRNA #4 for 14 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced THP-1 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 0.9% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 (SEQ ID NO: 4) stimulated 15.9%. With zoledronic acid treatment, LV-control stimulated 4.7% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 (SEQ ID NO: 4) stimulated 76.2%.

Example 5—Knock-Down of FDPS for 3 Days in PC3 Prostate Carcinoma Cells by shRNA #1

(187) This Example illustrates that knock-down of FDPS for 3 days in PC3 prostate carcinoma cells by lentiviral (LV)-expressing FDPS shRNA #1 stimulates TNF- α expression in GD T cells, as shown in FIG. 7.

(188) PC3 cells were transduced with LV-control or LV-FDPS shRNA #1 (SEQ ID NO: 1) for 3 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced PC3 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 0.2% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #1 stimulated 0.5%. With zoledronic acid treatment, LV-control stimulated 1.7% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #1 (SEQ ID NO: 1) stimulated 32.2%.

Example 6—Knock-Down of FDPS for 3 Days in PC3 Prostate Carcinoma Cells by shRNA #4

(189) This Example illustrates that Knock-down of FDPS for 3 days in PC3 prostate carcinoma cells by lentiviral (LV)-expressing FDPS shRNA #4 stimulates TNF- α expression in GD T cells, as shown in FIG. 8.

(190) PC3 cells were transduced with LV-control or LV-FDPS shRNA #4 (SEQ ID NO: 4) for 3 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced PC3 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 0.5% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 (SEQ ID NO: 4) stimulated 1.9%. With zoledronic acid treatment, LV-control stimulated 2.1% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #4 stimulated 28.7%.

Example 7—Knock-Down of FDPS for 3 Days in HepG2 Liver Carcinoma Cells by shRNA #1 and #4

(191) This Example illustrates that Knock-down of FDPS for 3 days in HepG2 liver carcinoma cells by lentiviral (LV)-expressing FDPS shRNA #1 (SEQ ID NO: 1) and shRNA #4 (SEQ ID NO: 4) stimulates TNF- α expression in GD T cells, as shown in FIG. 9.

(192) HepG2 cells were transduced with LV-control, LV-FDPS shRNA #1 (SEQ ID NO: 1), or LV-

FDPS shRNA #4 (SEQ ID NO: 4) for 3 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced HepG2 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 0.4% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #1 (SEQ ID NO: 1) and #4 (SEQ ID NO: 4) stimulated 0.7% and 0.9%, respectively. With zoledronic acid treatment, LV-control stimulated 6.9% of TNF- α expressing V γ 9V δ 2 T cells and LV-FDPS shRNA #1 and #4 stimulated 7.6% and 21.1%, respectively.

Example 8—Knock-Down of FDPS for 3 Days in THP1 Leukemia by microRNA-30

(193) This Example illustrates that Knock-down of FDPS for 3 days in THP1 leukemia cells by lentiviral (LV)-expressing FDPS-targeted synthetic microRNA-30 stimulates TNF- α expression in gamma delta T cells, as shown in FIG. 10.

(194) THP1 cells (1×10^5 cells) were transduced with LV-control or LV-miR30 FDPS #1 (SEQ ID NO: 5) for 3 days. Two days after transduction, cells were treated with or without 104 zoledronic acid. After 24 hours, the transduced THP-1 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2+ and TNF- α + cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms. Without zoledronic acid, LV-control stimulated 0.2% of TNF- α expressing V γ 9V δ 2 T cells and LV-miR30 FDPS stimulated 8.1%. With zoledronic acid treatment, LV-control stimulated 5.3% of TNF- α expressing V γ 9V δ 2 T cells and LV-miR30 FDPS #1 (SEQ ID NO: 5) stimulated 67.3%.

Example 9: E:T Ratios Resulting from Mixture of THP-1 Cells, Cultured Human GD T Cells, and/or Zometa (Zol)

(195) This Example demonstrates results from mixing treated THP-1 monocytoid tumor cells with cultured human GD T cells, as shown in FIG. 11.

(196) The monocytoid cell line THP-1 was treated with control lentivirus vector (LV), LV suppressing farnesyl diphosphate synthase gene expression (LV-FDPS), zoledronic acid (Zol) or combinations. The legend, as shown in FIG. 11, was: lentiviral control vectors (LV-Control), lentiviral vectors expressing microRNA to down regulate FDPS (LV-FPPS), Zometa (Zol), Zometa plus lentiviral control (Zol+LV-Control), or Zometa plus lentiviral vectors expressing microRNA to down regulate FPPS (Zol+LV-FPPS).

(197) Human GD T cells were cultured from an anonymous donor and added to treated THP-1 cells in 4:1, 2:1 or 1:1 ratios (GD T:THP-1) for 4 hours. Cell killing was measured by a fluorescence assay. When THP-1 cells were treated with a combination of LV-FDPS and Zol, cytotoxic T cell killing by GD T cells was increased greatly compared to either treatment alone. When LV-FDPS treatment alone was compared to Zol treatment alone, the LV-FDPS lead to greater killing but was >3-fold below tumor cell killing after combination treatment. The combined LV-FDPS plus Zol treatment caused nearly 70% tumor cell killing with 4:1 ratio; this was more than 3-fold higher than the second best treatment (LV-FDPS alone).

Example 10—Lentiviral-Delivered shRNA-Based RNA Interference Targeting the Human Farnesyl Diphosphate Synthase (FDPS) Gene

(198) HepG2 human hepatocellular carcinoma cells were infected with lentiviral vectors containing the H1 promoter and either a non-targeting or four different FDPS shRNA sequences, as shown in

FIG. 12. After 48 hours, RNA was extracted from the cells and converted to cDNA. Expression of FDPS cDNA was determined by quantitative PCR using SYBR Green and FDPS primers. FDPS expression was normalized to actin levels for each sample.

(199) FDPS-targeting lentiviral vectors containing the H1 promoter and either a non-targeting sequence

(200) TABLE-US-00009 (SEQ ID NO: 58) (5'-GCCGCTTTGTAGGATAGAGCTCGAGCTCTATCCTACAAAGCGGCTTTTT- 3')

or one of four different FDPS shRNA sequences

(201) TABLE-US-00010 (FDPS shRNA sequence #1; SEQ ID NO: 1) GTCCTGGAGTACAATGCCATTCTCGAGAATGGCATTGTACTCCAGGACTT TTT;

(FDPS shRNA sequence #2; SEQ ID NO: 2) GCAGGATTTTCGTTTCAGCACTTCTCGAGAAGTGCTGAACGAAATCCTGCTT TTT;

(FDPS shRNA sequence #3; SEQ ID NO: 3) GCCATGTACATGGCAGGAATTCTCGAGAATTCCTGCCATGTACATGGCTT TTT; and

(FDPS shRNA sequence #4; SEQ ID NO: 4) GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCCTCCTTCTGCTT TTT

were produced in 293 T cells.

(202) HepG2 human hepatocellular carcinoma cells were then infected with lentiviral vectors to determine the efficacy of FDPS knock-down. After 48 hours, RNA was extracted from the cells using the RNeasy RNA isolation kit (Qiagen) and converted to cDNA with the SuperScript VILO cDNA synthesis kit (Thermo Scientific). Expression of FDPS cDNA was determined by quantitative PCR on an Applied Biosystems StepOne qPCR machine using a SYBR Green PCR mix (Thermo Scientific) and FDPS primers (Forward primer: 5'-

AGGAATTGATGGCGAGAAGG-3' (SEQ ID NO: 59) and Reverse primer: 5'-CCCAAAGAGGTCAAGGTAATCA-3' (SEQ ID NO: 60)). FDPS expression was normalized to actin levels for each sample using the actin primers (Forward primer: 5'-

AGCGCGGCTACAGCTTCA-3' (SEQ ID NO: 61) and Reverse primer: 5'-GGCGACGTAGCACAGCTTCT-3') (SEQ ID NO: 62). The relative FDPS RNA expression of the shCon sample is set at 100%. There was an 85% (FDPS sequence #1), 89% (FDPS sequence #2), 46% (FDPS sequence #3), and 98% (FDPS sequence #4) decrease in FDPS expression.

Example 11—Lentiviral-Delivered miR-Based RNA Interference Targeting the Human Farnesyl Diphosphate Synthase (FDPS) Gene

(203) As shown in FIG. 13, HepG2 human hepatocellular carcinoma cells were infected with lentiviral vectors containing either the H1 promoter (SEQ ID NO: 16) the FDPS shRNA #4 (SEQ ID NO: 4) sequence or the EF-1a promoter (SEQ ID NO: 40) and miR30-based FDPS sequences. After 48 hours, cells were lysed and an immunoblot was performed using an anti-FDPS (Thermo Scientific) and an anti-actin (Sigma) antibody as a protein loading control.

(204) More specifically, HepG2 human hepatocellular carcinoma cells were infected with lentiviral vectors containing either the H1 promoter (SEQ ID NO: 16) and the FDPS shRNA sequence

(205) TABLE-US-00011 (FDPS shRNA sequence #4; SEQ ID NO: 4) GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCCTCCTTCTGCTT TTT or the EF-1alpha promoter (SEQ ID NO: 39) and miR30-based FDPS sequences (miR30 FDPS sequence #1; SEQ ID NO: 5)

AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGC GTGAAGCCACAGATGGCAGAAGGAGGCTGAGAAAGTGCTGCCTACTGCCT CGGACTTCAAGGGGCT and (miR30 FDPS sequence #2; SEQ ID NO: 6) AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAGCCTCCTTCTGCG TGAAGCCACAGATGGCAGAAGGGCTGAGAAAGTGCTGCCTACTGCCTCGGA CTTCAAGGGGCT.

(206) After 48 hours, cells were lysed with NP-40 lysis buffer and protein was quantified with the

Bio-Rad protein assay reagent. Protein samples at 50 micrograms were electrophoresed on 4-12% Bis-Tris gels (Thermo Scientific) and transferred to PVDF membranes (EMD Millipore). An immunoblot was performed using an anti-FDPS (Thermo Scientific) and an anti-actin (Sigma) antibody as a protein loading control. Antibodies were bound with HRP-conjugated secondary antibodies and detected with a Licor c-DiGit Blot scanner using the Immobilon Western ECL reagent (EMD Millipore). The densitometry of the immunoblot bands were quantified with the NIH image software. The LV control with the EF-1 promoter was set at 100%. There was a 68% (LV-shFDPS #4), 43% (LV-miR FDPS #1), and 38% (LV-miR FDPS #3) reduction of FDPS protein expression.

Example 12—Knock-Down of FDPS for 3 Days in HepG2 Liver Carcinoma Cells by Adeno-Associated Virus (AAV)-Expressing FDPS shRNA #4

(207) This Example illustrates that knock-down of FDPS for 3 days in HepG2 liver carcinoma cells by adeno-associated virus (AAV)-expressing FDPS shRNA #4 (SEQ ID NO: 4) stimulates TNF- α expression in GD T cells (FIG. 14, Panel B).

(208) HepG2 cells were transduced with control or AAV-FDPS shRNA #4 (SEQ ID NO: 8) for 3 days. Two days after transduction, cells were treated with or without 1 μ M zoledronic acid. After 24 hours, the transduced HepG2 cells were co-cultured with 5×10^5 PBMC cells and IL-2 in a round bottom 96 well plate for 4 hours. The PBMC cells were pre-stimulated with zoledronic acid and IL-2 for 11 days to expand V γ 9V δ 2 T cells. After staining for V γ 9V δ 2 and TNF- α using fluorophore-conjugated anti TCR-V δ 2 and anti-TNF- α antibody, cells were analyzed via flow cytometry. Live cells were gated, and V δ 2⁺ and TNF- α ⁺ cells were selected on a dot blot. The activated cytotoxic V γ 9V δ 2 T cells appeared in the upper right quadrant of flow cytograms (FIG. 14, Panel B).

(209) AAV Vector Construction. FDPS shRNA sequence #4 (SEQ ID NO: 4) was inserted into the pAAV plasmid (Cell Biolabs). FDPS oligonucleotide sequences containing BamHI and EcoRI restriction sites were synthesized by Eurofins MWG Operon. Overlapping sense and antisense oligonucleotide sequences were mixed and annealed during cooling from 70 degrees Celsius to room temperature. The pAAV was digested with the restriction enzymes BamHI and EcoRI for one hour at 37 degrees Celsius. The digested pAAV plasmid was purified by agarose gel electrophoresis and extracted from the gel using a DNA gel extraction kit from Thermo Scientific. The DNA concentrations were determined and vector to oligo (3:1 ratio) were mixed, allowed to anneal, and ligated. The ligation reaction was performed with T4 DNA ligase for 30 minutes at room temperature. 2.5 microliters of the ligation mix were added to 25 microliters of STBL3 competent bacterial cells. Transformation was achieved after heat-shock at 42 degrees Celsius. Bacterial cells were spread on agar plates containing ampicillin and drug-resistant colonies (indicating the presence of ampicillin-resistance plasmids) were recovered and expanded in LB broth. To check for insertion of the oligo sequences, plasmid DNA was extracted from harvested bacteria cultures with the Thermo Scientific DNA mini prep kit. Insertion of shRNA sequences in the pAAV plasmid was verified by DNA sequencing using a specific primer for the promoter used to regulate shRNA expression. An exemplary AAV vector with a H1 promoter (SEQ ID NO: 16), shFDPS sequence (e.g., SEQ ID NO: 4), Left Inverted Terminal Repeat (Left ITR; SEQ ID NO: 63), and Right Inverted Terminal Repeat (Right ITR; SEQ ID NO: 64) can be found in FIG. 14, Panel A).

(210) Production of AAV particles. The AAV-FDPS shRNA plasmid was combined with the plasmids pAAV-RC2 (Cell Biolabs) and pHelper (Cell Biolabs). The pAAV-RC2 plasmid contains the Rep and AAV2 capsid genes and pHelper contains the adenovirus E2A, E4, and VA genes. To produce AAV particles, these plasmids were transfected in the ratio 1:1:1 (pAAV-shFDPS: pAAV-RC2: pHelper) into 293T cells. For transfection of cells in 150 mm dishes (BD Falcon), 10 micrograms of each plasmid were added together in 1 ml of DMEM. In another tube, 60 microliters of the transfection reagent PEI (1 microgram/ml) (Polysciences) was added to 1 ml of DMEM. The two tubes were mixed together and allowed to incubate for 15 minutes. Then the transfection

mixture was added to cells and the cells were collected after 3 days. The cells were lysed by freeze/thaw lysis in dry ice/isopropanol. Benzonase nuclease (Sigma) was added to the cell lysate for 30 minutes at 37 degrees Celsius. Cell debris were then pelleted by centrifugation at 4 degrees Celsius for 15 minutes at 12,000 rpm. The supernatant was collected and then added to target cells.

Example 13—Decreased RAP1 Prenylation in the Cells Transduced with LV-shFDPS and Treated with Zoledronic Acid

(211) This Example illustrates that lentiviral-delivered shRNA targeting the human farnesyl diphosphate synthase (FDPS) gene and zoledronic acid synergize to inhibit farnesyl diphosphate production.

(212) FDPS is an enzyme in the isoprenoid synthesis pathway that catalyzes the production of farnesyl diphosphate. Inhibiting the enzyme activity of FDPS by zoledronic acid or reduced protein expression by shRNA-mediated knock-down will result in reduced farnesyl diphosphate levels. Farnesylation of cellular proteins requires farnesyl diphosphate. RAP1A is a protein that is modified by farnesylation, which can be used as a biomarker for levels of cellular farnesyl diphosphate. An antibody that specifically recognizes reduced RAP1A farnesylation was used to measure FDPS activity after transduction with LV-shFDPS alone or in combination with zoledronic acid. HepG2 human hepatocellular carcinoma cells were infected with lentiviral vectors containing FDPS shRNA sequence #4. For the zoledronic acid treated cells, zoledronic acid (Sigma) was added for the last 24 hours. After 48 hours, cells were lysed with NP-40 lysis buffer and protein was quantified with the Bio-Rad protein assay reagent. Protein samples at 50 micrograms were electrophoresed on 4-12% Bis-Tris gels (Thermo Scientific) and transferred to PVDF membranes (EMD Millipore). As shown in FIG. 15, an immunoblot was performed using an anti-FDPS (Thermo Scientific), anti-RAP1A (Santa Cruz), and an anti-actin (Sigma) antibody as a protein loading control. Antibodies were bound with HRP-conjugated secondary antibodies and detected with a Licor c-DiGit Blot scanner using the Immobilon Western ECL reagent (EMD Millipore). An increase in the RAP1A band intensity correlates with reduced farnesylation. RAP1A defarnesylation occurred only in the cells transduced with LV-shFDPS and treated with zoledronic acid.

Example 14—Treatment of a Subject with Cancer

(213) LV-FDPS is a Genetic Medicine Delivered by a Lentivirus Vector Via Local Administration to the Site of Late Stage, Non-Resectable Hepatocellular Carcinoma

(214) A Phase I clinical trial will test safety and feasibility of delivering LV-FDPS to the site of hepatocellular carcinoma (HCC) using ultrasound guided cannulation of the liver in patients without concomitant radiotherapy or chemotherapy. It is rationally predicted that this study will result in the successful treatment of HCC. The study is an open label, 4×3 dose escalation (4 dose ranges, up to 3 subjects per dose) to identify the maximum tolerable dose of LV-FDPS in patients 18 years or older with Stage III/IV non-resectable HCC.

(215) LV-FDPS is a genetic therapy designed to reduce expression in tumor cells of the enzyme farnesyl diphosphate synthase. Experimental studies show that tumor cells modified by LV-FDPS induce the anti-tumor activity of human gamma delta T cells, including the capacity for tumor killing by cellular cytotoxicity.

(216) Subjects with target lesions ≥ 1 cm in longest diameter (measured by helical CT) and ≤ 4.9 cm maximum diameter and meeting inclusion and exclusion criteria detailed below, are enrolled into the next available dosing category. A maximum of 3 subjects are recruited for each dosage group. The dose is number of transducing units of LV-FDPS as described in the product release criteria, delivered via intrahepatic cannulation in a single bolus with volume not to exceed 25 mL. The minimum dose is 1×10^9 transducing units and escalation is 10-fold to a next dose of 1×10^{10} transducing units, the next dose is 1×10^{11} transducing units, and a maximum dose of 1×10^{12} transducing units based on reported experience with recombinant adenovirus therapy for HCC (Sangro et al., A phase I clinic trial of thymidine kinase-based gene therapy in

advanced hepatocellular carcinoma, 2010, *Cancer Gene Ther.* 17:837-43). Subjects are enrolled, treated and evaluated for 3 months. All safety evaluations are completed for each group prior to enrolling and treating subjects at the next higher dose level. Enrollment and dose escalation continue until a maximum tolerable dose is achieved or the study is terminated.

(217) Cannulation is via the left subclavian artery until tip of catheter is at the proper hepatic artery junction. Cannulation is guided by ultrasonography as described (Lin et al., Clinical effects of intra-arterial infusion chemotherapy with cisplatin, mitomycin C, leucovorin and 5-Fluorouracil for unresectable advanced hepatocellular carcinoma, 2004, *J. Chin. Med. Assoc.* 67:602-10).

(218) Primary Outcome Measures

(219) Safety: Systemic and locoregional adverse events are graded according to CTCAS and coded according to MedRA. The adverse events data for all subjects at a single dose range will be evaluated prior to dose escalation. The final safety assessment incorporates data from all dose ranges.

(220) Secondary Outcome Measures

(221) Lesion distribution and retention of LV-FDPS following locoregional administration and subsequent biopsy or necropsy to obtain tissues. Objective response rate (ORR) in target and measurable non-local lesions (if present) by physical analysis, medical imaging or biopsy during 3 months after treatment. Levels of LV-FDPS in blood stream during 10 minutes, 30 minutes, 1 hour and 1 day after local injection. Changes in markers of hepatic function including ALP, ALT, ASAT, total bilirubin and GGT during 3 months after treatment. Disease free survival beyond historical control (no LV-FDPS) patients in ad hoc analysis.

Inclusion Criteria Greater than 18 years and including both males and females. Diagnosis confirmed by histology or cytology or based on currently accepted clinical standards of hepatocellular carcinoma of parenchyma cell origin that is not amenable, at the time of screening, to resection, transplant or other potentially curative therapies. Treating physician determines that the lesion is amenable to locoregional targeted delivery. Target lesion must represent measurable disease with a unidimensional longest diameter of ≥ 1.0 cm by computed tomography; the maximum longest diameter is ≤ 5.0 cm. Karnofsky performance score 60-80% of ECOG values. Life expectancy ≥ 12 weeks. Hematopoietic function: WBC $\geq 2,500/\text{mm}^3$; ANC $\geq 1000/\text{mm}^3$; Hemoglobin ≥ 8 g/dL; Platelet count $\geq 50,000/\text{mm}^3$; Coagulation INR ≤ 1.3 . AST and ALT < 5 times ULN; ALP < 5 times ULN. Bilirubin ≤ 1.5 times ULN; Creatinine ≤ 1.5 times ULN and eGFR ≥ 50 . Thyroid function: Total T3 or free T3, total T4 or free T4 and TSH $\leq$ CTCAE Grade 2 abnormality. Renal, cardiovascular and respiratory function adequate in the opinion of the attending physician. Immunological function: Circulating Vgamma9Vdelta2+ T cells $\geq 30/\text{mm}^3$; no immunodeficiency disease. Negative for HIV by serology and viral RNA test. Written informed consent.

Exclusion Criteria Target lesion contiguous with, encompasses or infiltrates blood vessel. Primary HCC amenable to resection, transplantation or other potentially curative therapies. Hepatic surgery or chemoembolization within the past 4 months. Hepatic radiation or whole body radiation therapy within past 4 months. Chemotherapy with 4 weeks or any use of nitrosourea, mitomycin C or cisplatin. Current or within past 4 weeks receipt of bisphosphonate therapy Investigational agents within 4 weeks or < 5 drug half-lives. Impaired wound healing due to diabetes. Significant psychiatric illness, alcohol dependence or illicit drug use. Unwilling to comply with study protocols and reporting requirements. Bisphosphonate treatment within past 4 months. Presence of clinically significant cardiovascular, cerebrovascular (stroke), immunological (except hepatitis B or C virus infection, viral hepatitis or cirrhosis), endocrine or central nervous system disorders; current encephalopathy; variceal bleeding requiring hospitalization or transfusion within past 4 months. History of HIV or acquired immune deficiency syndrome. Current or prior treatment with antiretroviral medications. Pregnant, lactating or refusal to adopt barrier or chemical contraceptive use throughout trial and follow-up interval.

LV-FDPS is a Genetic Medicine Delivered by a Lentivirus Vector Via Local Administration to the Site of Late Stage, Non-Resectable Hepatocellular Carcinoma—Adjunct Administration of Bisphosphonate

(222) A Phase I clinical trial will test safety and feasibility of delivering LV-FDPS to the site of hepatocellular carcinoma (HCC) using ultrasound guided cannulation of the liver in patients with concomitant bisphosphonate chemotherapy. It is rationally predicted that this study will result in the successful treatment of HCC. The study is an open label, 4×3 dose escalation (4 dose ranges, up to 3 subjects per dose) to identify the maximum tolerable dose of LV-FDPS in patients 18 years or older with Stage III/IV non-resectable HCC.

(223) LV-FDPS is a genetic therapy designed to reduce expression in tumor cells of the enzyme farnesyl diphosphate synthase. Experimental studies show that tumor cells modified by LV-FDPS induce the anti-tumor activity of human gamma delta T cells, including the capacity for tumor killing by cellular cytotoxicity. Prior experimental studies also showed the potential for positive interactions of LV-FDPS and specific bisphosphonate drugs that may be prescribed in primary or metastatic diseases. For this study, subjects will receive dose escalating amounts of LV-FDPS with continuous standard of care dosing with Aredia® (pamidronate), Zometa® (zoledronic acid) or Actonel® (risedronate) according to physician advice and subject preference.

(224) Subjects with target lesions ≥ 1 cm in longest diameter (measured by helical CT) and ≤ 4.9 cm maximum diameter and meeting inclusion and exclusion criteria detailed below, are enrolled and started on bisphosphonate therapy. 30 days later size of the target lesion is re-evaluated to ensure subjects still meet starting criteria for LV-FDPS. Subjects without objective clinical response on bisphosphonate are enrolled into the next available LV-FDPS dosing category. A maximum of 3 subjects are recruited for each dosage group and all continue on bisphosphonate for the study duration unless otherwise advised by the attending physician. The LV-FDPS dose is a number of transducing units of LV-FDPS as described in the product release criteria, delivered via intrahepatic cannulation in a single bolus with volume not to exceed 25 mL. The minimum dose is 1×10^9 transducing units and escalation is 10-fold to a next dose of 1×10^{10} transducing units, the next dose is 1×10^{11} transducing units, and a maximum dose of 1×10^{12} transducing units based on reported experience with recombinant adenovirus therapy for HCC (Sangro, et al., A phase I clinic trial of thymidine kinase-based gene therapy in advanced hepatocellular carcinoma, 2010, *Cancer Gene Ther.* 17:837-43). Subjects are enrolled, treated and evaluated for 3 months. All safety evaluations are completed for each group prior to enrolling and treating subjects at the next higher dose level. Enrollment and dose escalation continue until a maximum tolerable dose is achieved or the study is terminated.

(225) Cannulation is via the left subclavian artery until tip of catheter is at the proper hepatic artery junction. Cannulation is guided by ultrasonography as described (Lin et al., Clinical effects of intra-arterial infusion chemotherapy with cisplatin, mitomycin C, leucovor and 5-Fluorouracil for unresectable advanced hepatocellular carcinoma, 2004, *J. Chin. Med. Assoc.* 67:602-10).

(226) Primary Outcome Measures

(227) Safety: Systemic and locoregional adverse events are graded according to CTCAS and coded according to MedRA. The adverse events data for all subjects at a single dose range will be evaluated prior to dose escalation. The final safety assessment incorporates data from all dose ranges.

(228) Secondary Outcome Measures

(229) Lesion distribution and retention of LV-FDPS following locoregional administration and subsequent biopsy or necropsy to obtain tissues. Objective response rate (ORR) in target and measurable non-local lesions (if present) by physical analysis, medical imaging or biopsy during 3 months after treatment. Levels of LV-FDPS in blood stream during 10 minutes, 30 minutes, 1 hour and 1 day after local injection. Changes in markers of hepatic function including ALP, ALT, ASAT, total bilirubin and GGT during 3 months after treatment. Disease free survival beyond historical

control (no LV-FDPS) patients in ad hoc analysis.

Inclusion Criteria Greater than 18 years and including both males and females. Diagnosis confirmed by histology or cytology or based on currently accepted clinical standards of hepatocellular carcinoma of parenchyma cell origin that is not amenable, at the time of screening, to resection, transplant or other potentially curative therapies. Treating physician determines that the lesion is amenable to locoregional targeted delivery. Target lesion must represent measurable disease with a unidimensional longest diameter of ≥ 1.0 cm by computed tomography; the maximum longest diameter is ≤ 5.0 cm. Karnofsky performance score 60-80% of ECOG values. Life expectancy ≥ 12 weeks. Hematopoietic function: WBC $\geq 2,500/\text{mm}^3$; ANC $\geq 1000/\text{mm}^3$; Hemoglobin ≥ 8 g/dL; Platelet count $\geq 50,000/\text{mm}^3$; Coagulation INR ≤ 1.3 . AST and ALT < 5 times ULN; ALPS < 5 time ULN. Bilirubin ≤ 1.5 times ULV; Creatine ≤ 1.5 times ULN and eGFR ≥ 50 . Thyroid function: Total T3 or free T3, total T4 or free T4 and THCAE Grade 2 abnormality. Renal, cardiovascular and respiratory function adequate in the opinion of the attending physician. Immunological function: Circulating Vgamma9Vdelta2+ T cells $\geq 30/\text{mm}^3$; no immunodeficiency disease. Negative for HIV by serology and viral RNA test. Written informed consent.

Exclusion Criteria Intolerant to or unwilling to continue bisphosphonate adjunct therapy. Objective clinical response after bisphosphonate therapy. Target lesion contiguous with, encompasses or infiltrates blood vessel. Primary HCC amenable to resection, transplantation or other potentially curative therapies. Hepatic surgery or chemoembolization within the past 4 months. Hepatic radiation or whole body radiation therapy within past 4 months. Chemotherapy excluding bisphosphonate, within 4 weeks or any use of nitrosourea, mitomycin C or cisplatin. Investigational agents within 4 weeks or < 5 drug half-lives. Impaired wound healing due to diabetes. Significant psychiatric illness, alcohol dependence or illicit drug use. Unwilling to comply with study protocols and reporting requirements. Presence of clinically significant cardiovascular, cerebrovascular (stroke), immunological (except hepatitis B or C virus infection, viral hepatitis or cirrhosis), endocrine or central nervous system disorders; current encephalopathy; variceal bleeding requiring hospitalization or transfusion within past 4 months. History of HIV or acquired immune deficiency syndrome. Current or prior treatment with antiretroviral medications. Pregnant, lactating or refusal to adopt barrier or chemical contraceptive use throughout trial and follow-up interval.

Example 15—Treatment of a Subject with Chronic Viral Disease(s) of the Liver

(230) LV-FDPS is a Genetic Medicine Delivered by a Lentivirus Vector Via Local Administration to Liver for the Treatment of Hepatitis B Virus, Hepatitis C Virus, HIV or Other Viral Infection of the Liver

(231) A Phase I clinical trial will test safety and feasibility of delivering LV-FDPS to virally infected liver using ultrasound guided cannulation. It is rationally predicted that this study will result in the successful treatment of infections of the liver. The study is an open label, 4×3 dose escalation (4 dose ranges, up to 3 subjects per dose) to identify the maximum tolerable dose of LV-FDPS in patients 18 years or older with chronic viral disease of the liver that is resistant to chemotherapy.

(232) LV-FDPS is a genetic therapy designed to reduce expression in tumor cells of the enzyme farnesyl diphosphate synthase. Experimental studies show that tumor cells modified by LV-FDPS induce human gamma delta T cells, including a capacity for cellular cytotoxicity against virally-infected cells.

(233) Subjects with confirmed viral infection of the liver including hepatitis B virus, hepatitis C virus, HIV or other viruses are enrolled into the next available LV-FDPS dosing category. A maximum of 3 subjects are recruited for each dosage group. The LV-FDPS dose is a number of transducing units of LV-FDPS as described in the product release criteria, delivered via intrahepatic cannulation in a single bolus with volume not to exceed 25 mL. The minimum dose is 1×10^9 transducing units and escalation is 10-fold to a next dose of 1×10^{10} transducing units, the next

dose is 1×10^8 sup.11 transducing units, and a maximum dose of 1×10^8 sup.12 transducing units based on reported experience with recombinant adenovirus therapy for HCC (Sangro, et al., A phase I clinic trial of thymidine kinase-based gene therapy in advanced hepatocellular carcinoma, 2010, *Cancer Gene Ther.* 17:837-43). Subjects are enrolled, treated and evaluated for 3 months. All safety evaluations are completed for each group prior to enrolling and treating subjects at the next higher dose level. Enrollment and dose escalation continue until a maximum tolerable dose is achieved or the study is terminated.

(234) Cannulation is via the left subclavian artery until tip of catheter is at the proper hepatic artery junction. Cannulation is guided by ultrasonography as described (Lin et al., Clinical effects of intra-arterial infusion chemotherapy with cisplatin, mitomycin C, leucovorin and 5-Fluorouracil for unresectable advanced hepatocellular carcinoma, 2004, *J. Chin. Med. Assoc.* 67:602-10).

(235) Primary Outcome Measures

(236) Safety: Systemic and locoregional adverse events are graded according to CTCAS and coded according to MedRA. The adverse events data for all subjects at a single dose range will be evaluated prior to dose escalation. The final safety assessment incorporates data from all dose ranges.

(237) Secondary Outcome Measures

(238) Lesion distribution and retention of LV-FDPS following locoregional administration and subsequent biopsy or necropsy to obtain tissues. Objective response rate (ORR) measured as a Sustained Viral Response (SVR) within the organ or systemically during 3 months after treatment. Levels of LV-FDPS in blood stream during 10 minutes, 30 minutes, 1 hour and 1 day after local injection. Changes in markers of hepatic function including ALP, ALT, ASAT, total bilirubin and GGT during 3 months after treatment. Disease free survival beyond historical control (no LV-FDPS) patients in ad hoc analysis.

Inclusion Criteria Greater than 18 years and including both males and females. Diagnosis confirmed by histology or cytology or based on currently accepted clinical standards of chronic viral infection of the liver that is not amenable, at the time of screening, to resection, transplant or other potentially curative therapies. Treating physician determines that the lesion is amenable to locoregional targeted delivery. Karnofsky performance score 60-80% of ECOG values. Life expectancy ≥ 12 weeks. Hematopoietic function: WBC $\geq 2,500/\text{mm}^3$; ANC $\geq 1000/\text{mm}^3$; Hemoglobin ≥ 8 g/dL; Platelet count $\geq 50,000/\text{mm}^3$; Coagulation INR ≤ 1.3 . AST and ALT < 5 times ULN; ALPS < 5 times ULN. Bilirubin ≤ 1.5 times ULN; Creatine ≤ 1.5 times ULN and eGFR ≥ 50 . Thyroid function: Total T3 or free T3, total T4 or free T4 and TSH $\leq$ CTCAE Grade 2 abnormality. Renal, cardiovascular and respiratory function adequate in the opinion of the attending physician. Immunological function: Circulating Vgamma9Vdelta2+ T cells $\geq 30/\text{mm}^3$; no immunodeficiency disease. Negative for HIV by serology and viral RNA test. Written informed consent.

Exclusion Criteria Chronic viral disease amenable to resection, transplantation or other potentially curative therapies. Hepatic surgery or chemoembolization within the past 4 months. Hepatic radiation or whole body radiation therapy within past 4 months. Investigational agents within 4 weeks or < 5 drug half-lives. Current (within past 4 weeks) or ongoing receipt of bisphosphonate therapy. Impaired wound healing due to diabetes. Significant psychiatric illness, alcohol dependence or illicit drug use. Unwilling to comply with study protocols and reporting requirements. Presence of clinically significant cardiovascular, cerebrovascular (stroke), immunological (except virus infection, viral hepatitis or cirrhosis), endocrine or central nervous system disorders; current encephalopathy; variceal bleeding requiring hospitalization or transfusion within past 4 months. Pregnant, lactating or refusal to adopt barrier or chemical contraceptive use throughout trial and follow-up interval.

LV-FDPS is a Genetic Medicine Delivered by a Lentivirus Vector Via Local Administration to Liver for the Treatment of Hepatitis B Virus, Hepatitis C Virus, HIV or Other Viral Infection of the

Liver—Concomitant Adjunct Bisphosphonate Therapy

(239) A Phase I clinical trial will test safety and feasibility of delivering LV-FDPS to virally infected liver using ultrasound guided cannulation. It is rationally predicted that this study will result in the successful treatment of infections of the liver. The study is an open label, 4×3 dose escalation (4 dose ranges, up to 3 subjects per dose) to identify the maximum tolerable dose of LV-FDPS in patients 18 years or older with chronic viral disease of the liver that is resistant to chemotherapy.

(240) LV-FDPS is a genetic therapy designed to reduce expression in tumor cells of the enzyme farnesyl diphosphate synthase. Experimental studies show that tumor cells modified by LV-FDPS induce human gamma delta T cells, including a capacity for cellular cytotoxicity against virally-infected cells. Prior experimental studies also showed the potential for positive interactions of LV-FDPS and specific bisphosphonate drugs that may be prescribed during infectious disease. For this study, subjects will receive dose escalating amounts of LV-FDPS with continuous standard of care dosing with Aredia® (pamidronate), Zometa® (zoledronic acid) or Actonel® (risedronate) according to physician advice and subject preference.

(241) Subjects with confirmed viral infection of the liver including hepatitis B virus, hepatitis C virus, HIV or other viruses will initiate bisphosphonate therapy for 45 days before re-screening to meet enrollment criteria for LV-FDPS treatment of infectious disease. Eligible subjects are enrolled into the next available LV-FDPS dosing category. A maximum of 3 subjects are recruited for each dosage group. The LV-FDPS dose is a number of transducing units of LV-FDPS as described in the product release criteria, delivered via intrahepatic cannulation in a single bolus with volume not to exceed 25 mL. The minimum dose is 1×10^9 transducing units and escalation is 10-fold to a next dose of 1×10^{10} transducing units, the next dose is 1×10^{11} transducing units, and a maximum dose of 1×10^{12} transducing units based on reported experience with recombinant adenovirus therapy for HCC (Sangro, et al., A phase I clinic trial of thymidine kinase-based gene therapy in advanced hepatocellular carcinoma, 2010, *Cancer Gene Ther.* 17:837-43). Subjects are enrolled, treated and evaluated for 3 months. All safety evaluations are completed for each group prior to enrolling and treating subjects at the next higher dose level. Enrollment and dose escalation continue until a maximum tolerable dose is achieved or the study is terminated.

(242) Cannulation is via the left subclavian artery until tip of catheter is at the proper hepatic artery junction. Cannulation is guided by ultrasonography as described (Lin et al., Clinical effects of intra-arterial infusion chemotherapy with cisplatin, mitomycin C, leucovor and 5-Fluorouracil for unresectable advanced hepatocellular carcinoma, 2004, *J. Chin. Med. Assoc.* 67:602-10).

(243) Primary Outcome Measures

(244) Safety: Systemic and locoregional adverse events are graded according to CTCAS and coded according to MedRA. The adverse events data for all subjects at a single dose range will be evaluated prior to dose escalation. The final safety assessment incorporates data from all dose ranges.

(245) Secondary Outcome Measures

(246) Lesion distribution and retention of LV-FDPS following locoregional administration and subsequent biopsy or necropsy to obtain tissues. Objective response rate (ORR) measured as a Sustained Viral Response (SVR) within the organ or systemically during 3 months after treatment. Levels of LV-FDPS in blood stream during 10 minutes, 30 minutes, 1 hour and 1 day after local injection. Changes in markers of hepatic function including ALP, ALT, ASAT, total bilirubin and GGT during 3 months after treatment. Disease free survival beyond historical control (no LV-FDPS) patients in ad hoc analysis.

Inclusion Criteria Greater than 18 years and including both males and females. Diagnosis confirmed by histology or cytology or based on currently accepted clinical standards of chronic viral infection of the liver that is not amenable, at the time of screening, to resection, transplant or other potentially curative therapies. Treating physician determines that the lesion is amenable to

locoregional targeted delivery. Karnofsky performance score 60-80% of ECOG values. Life expectancy ≥ 12 weeks. Hematopoietic function: WBC $\geq 2,500/\text{mm}^3$; ANC $\geq 1000/\text{mm}^3$; Hemoglobin ≥ 8 g/dL; Platelet count $\geq 50,000/\text{mm}^3$; Coagulation INR ≤ 1.3 . AST and ALT < 5 times ULN; ALPS < 5 time ULN. Bilirubin ≤ 1.5 times ULV; Creatine ≤ 1.5 times ULN and eGFR ≥ 50 . Thyroid function: Total T3 or free T3, total T4 or free T4 and $\text{THC} \leq \text{CTCAE}$ Grade 2 abnormality. Renal, cardiovascular and respiratory function adequate in the opinion of the attending physician. Immunological function: Circulating $\text{V}\gamma 9\text{V}\delta 2^+$ T cells $\geq 30/\text{mm}^3$; no immunodeficiency disease. Negative for HIV by serology and viral RNA test. Written informed consent.

Exclusion Criteria Chronic viral disease amenable to resection, transplantation or other potentially curative therapies. Hepatic surgery or chemoembolization within the past 4 months. Hepatic radiation or whole body radiation therapy within past 4 months. Investigational agents within 4 weeks or < 5 drug half-lives. Impaired wound healing due to diabetes. Significant psychiatric illness, alcohol dependence or illicit drug use. Unwilling to comply with study protocols and reporting requirements. Presence of clinically significant cardiovascular, cerebrovascular (stroke), immunological (except virus infection, viral hepatitis or cirrhosis), endocrine or central nervous system disorders; current encephalopathy; variceal bleeding requiring hospitalization or transfusion within past 4 months. Pregnant, lactating or refusal to adopt barrier or chemical contraceptive use throughout trial and follow-up interval

Sequences

(247) The following sequences are referred to herein:

(248) TABLE-US-00012 SEQ ID NO: Description Sequence 1 FDPS shRNA
GTCCTGGAGTACAATGCCATTCTCGAGAATGGCATTGTA sequence #1
CTCCAGGACTTTTT 2 FDPS shRNA
GCAGGATTCGTTTCAGCACTTCTCGAGAAGTGCTGAACG sequence #2
AAATCCTGCTTTTT 3 FDPS shRNA
GCCATGTACATGGCAGGAATTCTCGAGAATTCCTGCCAT sequence #3
GTACATGGCTTTTT 4 FDPS shRNA
GCAGAAGGAGGCTGAGAAAGTCTCGAGACTTTCTCAGCC sequence #4
TCCTTCTGCTTTTT 5 miR30 FDPS
AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAG sequence #1
CCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGAGGCT
GAGAAAGTGCTGCCTACTGCCTCGGACTTCAAGGGGCT 6 miR30 FDPS
AAGGTATATTGCTGTTGACAGTGAGCGACACTTTCTCAG sequence #2
CCTCCTTCTGCGTGAAGCCACAGATGGCAGAAGGGGCTGA
GAAAGTGCTGCCTACTGCCTCGGACTTCAAGGGGCT 7 miR30 FDPS
TGCTGTTGACAGTGAGCGACTTTCTCAGCCTCCTTCTGCG sequence #3
TGAAGCCACAGATGGCAGAAGGAGGCTGAGAAAGTTGC CTACTGCCTCGGA 8
miR155 FDPS CCTGGAGGCTTGCTGAAGGCTGTATGCTGACTTTCTCAG sequence #1
CCTCCTTCTGCTTTTTGGCCACTGACTGAGCAGAAGGGGCT
GAGAAAGTCAGGACACAAGGCCTGTTACTAGCACTCA 9 miR21 FDPS
CATCTCCATGGCTGTACCACCTTGTCGGGACTTTCTCAGC sequence #1
CTCCTTCTGCCTGTTGAATCTCATGGCAGAAGGAGGCGA
GAAAGTCTGACATTTTGGTATCTTTCATCTGACCA 10 miR185 FDPS
GGGCCTGGCTCGAGCAGGGGGCGAGGGATACTTTCTCAG sequence #1
CCTCCTTCTGCTGGTCCCCTCCCCGCAGAAGGAGGCTGA
GAAAGTCCTTCCCTCCCAATGACCGCGTCTTCGTCG 11 Rous Sarcoma
GTAGTCTTATGCAATACTCTTGTAAGTCTTGCAACATGGTA virus (RSV)
ACGATGAGTTAGCAACATGCCTTACAAGGAGAGAAAAA promoter
GCACCGTGCATGCCGATTGGTGGAAGTAAGGTGGTACGA

TCGTGCTTACTTACCAACAGACAGCGGTCTGACATG
GATTGGACGAACCACTGAATTGCCGCATTGCAGAGATAT
TGTATTTAAGTGCCTAGCTCGATACAATAAACG 12 5' Long terminal
GGTCTCTCTGGTTAGACCAGATCTGAGCCTGGGAGCTCT repeat (LTR)
CTGGCTAACTAGGGAACCCACTGCTTAAGCCTCAATAAA
GCTTGCCTTGAGTGCTTCAAGTAGTGTGTGCCCCGTCTGTT
GTGTGACTCTGGTAACTAGAGATCCCTCAGACCCTTTTA
GTCAGTGTGGAAAATCTCTAGCA 13 Psi Packaging
TACGCCAAAAATTTGACTAGCGGAGGCTAGAAGGAGA signal GAG 14 Rev response
AGGAGCTTTGTTCCCTTGGGTCTTGGGAGCAGCAGGAAG element (RRE)
CACTATGGGCGCAGCCTCAATGACGCTGACGGTACAGGC
CAGACAATTATTGTCTGGTATAGTGCAGCAGCAGAACAA
TTTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCA
ACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGAAT
CCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCC 15 Central polypurine
TTTTAAAAGAAAAGGGGGGATTGGGGGGGTACAGTGCAG tract (cPPT)
GGGAAAGAATAGTAGACATAATAGCAACAGACATACAA
ACTAAAGAATTACAAAAACAAATTACAAAATTCAAAATT TTA 16 Polymerase III
GAACGCTGACGTCATCAACCCGCTCCAAGGAATCGCGGG shRNA promoters;
CCCAGTGTCACTAGGCGGGAACACCCAGCGCGCGTGC GC H1 promoter
CCTGGCAGGAAGATGGCTGTGAGGGACAGGGGAGTGGC
GCCCTGCAATATTTGCATGTGCTATGTGTTCTGGGAAAT
CACCATAAACGTGAAATGTCTTTGGATTTGGGAATCTTA
TAAGTTCTGTATGAGACCACTT 17 Long WPRE
AATCAACCTCTGATTACAAAATTTGTGAAAGATTGACTG sequence
GTATTCTTAACTATGTTGCTCCTTTTACGCTATGTGGATA
CGCTGCTTTAATGCCTTTGTATCATGCTATTGCTTCCCGT
ATGGCTTTTCATTTTCTCCTCCTTGTATAAATCCTGGTTGC
TGTCTCTTTATGAGGAGTTGTGGCCCGTTGTCAGGCAAC
GTGGCGTGGTGTGCACTGTGTTTGCTGACGCAACCCCCA
CTGGTTGGGGCATTGCCACCACCTGTCAGCTCCTTTCCGG
GACTTTCGCTTTCCCCCTCCCTATTGCCACGGCGGAATC
ATCGCCGCCTGCCTTGCCCGCTGCTGGACAGGGGCTCGG
CTGTTGGGCACTGACAATTCCGTGGTGTGTCGGGGAAA
TCATCGTCCTTTCCCTTGGCTGCTCGCCTGTGTTGCCACCT
GGATTCTGCGCGGGACGTCCTTCTGCTACGTCCCTTCGGC
CCTCAATCCAGCGGACCTTCCTTCCCGCGGCCTGCTGCC
GGCTCTGCGGCCTCTTCCGCGTCTTCGCCTTCGCCCTCAG
ACGAGTCGGATCTCCCTTTGGGCCGCCTCCCCGCCT 18 3' delta LTR
TGGAAGGGCTAATCACTCCCAACGAAGATAAGATCTGC
TTTTTGCTTGTACTGGGTCTCTCTGGTTAGACCAGATCTG
AGCCTGGGAGCTCTCTGGCTAACTAGGGAACCCACTGCT
TAAGCCTCAATAAAGCTTGCCTTGAGTGCTTCAAGTAGT
GTGTGCCCCGTCTGTTGTGTGACTCTGGTAACTAGAGATC
CCTCAGACCCTTTTAGTCACTGTGGAAAATCTCTAGCAG TAGTAGTTCATGTCA 19
Helper/Rev; GCTATTACCATGGGTGCGAGGTGAGCCCCACGTTCTGCTT Chicken beta
actin CACTCTCCCCATCTCCCCCCCCCTCCCCACCCCCAATTTTG (CAG) promoter;
TATTTATTTATTTTAAATTATTTTGTGTCAGCGATGGGGG Transcription
CGGGGGGGGGGGGGGGGGGGCGCGCGCCAGGCGGGGGCGGGGC
GGGGCGAGGGGCGGGGCGGGGCGAGGCGGAGAGGTGC

GGCGGCGACCAATACAGGACGCGCGCGCTCCGAAAGTTCC
TTTTATGGCGAGGCGGCGGCGGCGGCCCTATAAAAA
GCGAAGCGCGCGGCGGGCG 20 Helper/Rev; HIV
ATGGGTGCGAGAGCGTCAGTATTAAGCGGGGGAGAATT Gag; Viral capsid
AGATCGATGGGAAAAAATTCGGTTAAGGCCAGGGGGAA
AGAAAAAATATAAATTAACATATAGTATGGGCAAGC
AGGGAGCTAGAACGATTTCGCAGTTAATCCTGGCCTGTTA
GAAACATCAGAAGGCTGTAGACAAATACTGGGACAGCT
ACAACCATCCCTTCAGACAGGATCAGAAGAAGCTTAGATC
ATTATATAATACAGTAGCAACCCTCTATTGTGTGCATCA
AAGGATAGAGATAAAAGACACCAAGGAAGCTTTAGACA
AGATAGAGGAAGAGCAAAACAAAAGTAAGAAAAAAGC
ACAGCAAGCAGCAGCTGACACAGGACACAGCAATCAGG
TCAGCCAAAATTACCCTATAGTGCAGAACATCCAGGGGC
AAATGGTACATCAGGCCATATCACCTAGAACTTTAAATG
CATGGGTAAAAGTAGTAGAAGAGAAGGCTTTCAGCCCA
GAAGTGATACCCATGTTTTTCAGCATTATCAGAAGGAGCC
ACCCACAAGATTTAAACACCATGCTAAACACAGTGGGG
GGACATCAAGCAGCCATGCAAATGTTAAAAGAGACCAT
CAATGAGGAAGCTGCAGAATGGGATAGAGTGCATCCAG
TGCATGCAGGGCCTATTGCACCAGGCCAGATGAGAGAA
CCAAGGGGAAGTGACATAGCAGGAAGCTACTAGTACCCTT
CAGGAACAAATAGGATGGATGACACATAATCCACCTATC
CCAGTAGGAGAAATCTATAAAAGATGGATAATCCTGGG
ATTAAATAAAATAGTAAGAATGTATAGCCCTACCAGCAT
TCTGGACATAAGACAAGGACCAAAGGAACCCTTTAGAG
ACTATGTAGACCGATTCTATAAACTCTAAGAGCCGAGC
AAGCTTCACAAGAGGTAAAAAATTGGATGACAGAAACC
TTGTTGGTCCAAAATGCGAACCCAGATTGTAAGACTATT
TTAAAAGCATTGGGACCAGGAGCGACACTAGAAGAAAT
GATGACAGCATGTCAGGGAGTGGGGGGACCCGGCCATA
AAGCAAGAGTTTTTGGCTGAAGCAATGAGCCAAGTAACA
AATCCAGCTACCATAATGATACAGAAAGGCAATTTTAGG
AACCAGAGAAAGACTGTTAAGTGTTTCAATTGTGGCAAA
GAAGGGCACATAGCCAAAAATTGCAGGGCCCCTAGGAA
AAAGGGCTGTTGGAAATGTGGAAAGGAAGGACACCAA
TGAAAGATTGTACTGAGAGACAGGCTAATTTTTTTAGGGA
AGATCTGGCCTTCCCACAAGGGAAGGCCAGGGAATTTTC
TTCAGAGCAGACCAGAGCCAACAGCCCCACCAGAAGAG
AGCTTCAGGTTTGGGGAAGAGACAACAACCTCCCTCTCAG
AAGCAGGAGCCGATAGACAAGGAACTGTATCCTTTAGCT
TCCCTCAGATCACTCTTTGGCAGCGACCCCTCGTCACAAT AA 21 Helper/Rev; HIV
ATGAATTTGCCAGGAAGATGGAAACCAAAAATGATAGG Pol; Protease and
GGGAATTGGAGGTTTTATCAAAGTAGGACAGTATGATCA reverse
GATACTCATAGAAATCTGCGGACATAAAGCTATAGGTAC transcriptase
AGTATTAGTAGGACCTACACCTGTCAACATAATTGGAAG
AAATCTGTTGACTCAGATTGGCTGCACTTTAAATTTTCCC
ATTAGTCCTATTGAGACTGTACCAGTAAAATTAAAGCCA
GGAATGGATGGCCCAAAAGTTAAACAATGGCCATTGAC
AGAAGAAAAAATAAAAGCATTAGTAGAAATTTGTACAG

AAATGGGAAAGAAAATTTCAAAAATTGGGCCT
GAAAATCCATACAATACTCCAGTATTTGCCATAAAGAAA
AAAGACAGTACTAAATGGAGAAAATTAGTAGATTTTCAG
AGAACTTAATAAGAGAACTCAAGATTTCTGGGAAGTTCA
ATTAGGAATACCACATCCTGCAGGGTTAAAACAGAAAA
AATCAGTAACAGTACTGGATGTGGGCGATGCATATTTTT
CAGTTCCCTTAGATAAAGACTTCAGGAAGTATACTGCAT
TTACCATAACCTAGTATAAACAATGAGACACCAGGGATTA
GATATCAGTACAATGTGCTTCCACAGGGATGGAAAGGAT
CACCAGCAATATTCCAGTGTAGCATGACAAAAATCTTAG
AGCCTTTTAGAAAACAAAATCCAGACATAGTCATCTATC
AATACATGGATGATTTGTATGTAGGATCTGACTTAGAAA
TAGGGCAGCATAGAACAAAAATAGAGGAACTGAGACAA
CATCTGTTGAGGTGGGGATTTACCACACCAGACAAAAAA
CATCAGAAAGAACCTCCATTCTTTGGATGGGTATGAA
CTCCATCCTGATAAATGGACAGTACAGCCTATAGTGCTG
CCAGAAAAGGACAGCTGGACTGTCAATGACATACAGAA
ATTAGTGGGAAAATTGAATTGGGCAAGTCAGATTTATGC
AGGGATTAAAGTAAGGCAATTATGTAAACTTCTTAGGGG
AACCAAAGCACTAACAGAAGTAGTACCACTAACAGAAG
AAGCAGAGCTAGAACTGGCAGAAAACAGGGAGATTCTA
AAAGAACCGGTACATGGAGTGTATTATGACCCATCAAAA
GACTTAATAGCAGAAATACAGAAGCAGGGGGCAAGGCCA
ATGGACATATCAAATTTATCAAGAGCCATTTAAAAATCT
GAAAACAGGAAAATATGCAAGAATGAAGGGTGCCCACA
CTAATGATGTGAAACAATTAACAGAGGCAGTACAAAAA
ATAGCCACAGAAAGCATAGTAATATGGGGAAAGACTCC
TAAATTTAAATTACCCATACAAAAGGAAACATGGGAAG
CATGGTGGACAGAGTATTGGCAAGCCACCTGGATTCTTG
AGTGGGAGTTTGTCAATACCCCTCCCTTAGTGAAGTTAT
GGTACCAGTTAGAGAAAGAACCCATAATAGGAGCAGAA
ACTTTCTATGTAGATGGGGCAGCCAATAGGGAAACTAAA
TTAGGAAAAGCAGGATATGTAAGTACAGAGGAAGACA
AAAAGTTGTCCCCCTAACGGACACAACAAATCAGAAGA
CTGAGTTACAAGCAATTCATCTAGCTTTGCAGGATTCGG
GATTAGAAGTAAACATAGTGACAGACTCACAATATGCAT
TGGGAATCATTCAAGCACAACCAGATAAGAGTGAATCA
GAGTTAGTCAGTCAAATAATAGAGCAGTTAATAAAAAA
GGAAAAAGTCTACCTGGCATGGGTACCAGCACACAAAG
GAATTGGAGGAAATGAACAAGTAGATGGGTTGGTCAGT
GCTGGAATCAGGAAAGTACTA 22 Helper Rev; HIV
TTTTTAGATGGAATAGATAAGGCCCAAGAAGAACATGA Integrase;
GAAATATCACAGTAATTGGAGAGCAATGGCTAGTGATTT Integration of viral
TAACCTACCACCTGTAGTAGCAAAAGAAATAGTAGCCAG RNA
CTGTGATAAATGTCAGCTAAAAGGGGAAGCCATGCATG
GACAAGTAGACTGTAGCCCAGGAATATGGCAGCTAGATT
GTACACATTTAGAAGGAAAAGTTATCTTGGTAGCAGTTC
ATGTAGCCAGTGGATATATAGAAGCAGAAGTAATTCCAG
CAGAGACAGGGCAAGAAACAGCATACTTCCTCTTAAAT
TAGCAGGAAGATGGCCAGTAAAAACAGTACATACAGAC

AATGGCAGCAATTTACCACTACTACAGTTAAGGCCGCC
TGTTGGTGGGCGGGGATCAAGCAGGAATTTGGCATTCCC
TACAATCCCCAAAGTCAAGGAGTAATAGAATCTATGAAT
AAAGAATTAAAGAAAATTATAGGACAGGTAAGAGATCA
GGCTGAACATCTTAAGACAGCAGTACAAATGGCAGTATT
CATCCACAATTTTAAAAGAAAAGGGGGGATTGGGGGGT
ACAGTGCAGGGGAAAGAATAGTAGACATAATAGCAACA
GACATACAACTAAAGAATTACAAAAACAAATTACAAA
AATTCAAAATTTTCGGGTTTATTACAGGGACAGCAGAGA
TCCAGTTTGGAAGGACCAGCAAAGCTCCTCTGGAAAGG
TGAAGGGGCAGTAGTAATACAAGATAATAGTGACATAA
AAGTAGTGCCAAGAAGAAAAGCAAAGATCATCAGGGAT
TATGGAAAACAGATGGCAGGTGATGATTGTGTGGCAAGT AGACAGGATGAGGATTAA
23 Helper/Rev; HIV AGGAGCTTTGTTCTTGGGTTCTTGGGAGCAGCAGGAAG RRE;
Binds Rev CACTATGGGCGCAGCGTCAATGACGCTGACGGTACAGGC element
CAGACAATTATTGTCTGGTATAGTGACAGCAGCAGAACAA
TTTGCTGAGGGCTATTGAGGCGCAACAGCATCTGTTGCA
ACTCACAGTCTGGGGCATCAAGCAGCTCCAGGCAAGAAT
CCTGGCTGTGGAAAGATACCTAAAGGATCAACAGCTCCT 24 Helper/Rev; HIV
ATGGCAGGAAGAAGCGGAGACAGCGACGAAGAAGCTCCT Rev; Nuclear
CAAGGCAGTCAGACTCATCAAGTTTCTCTATCAAAGCAA export and
CCCACCTCCCAATCCCGAGGGGACCCGACAGGCCCGAA stabilize viral
GGAATAGAAGAAGAAGGTGGAGAGAGAGACAGAGACA mRNA
GATCCATTCGATTAGTGAACGGATCCTTAGCACTTATCT
GGGACGATCTGCGGAGCCTGTGCCTCTTCAGCTACCACC
GCTTGAGAGACTTACTCTTGATTGTAACGAGGATTGTGG
AACTTCTGGGACGCAGGGGGTGGGAAGCCCTCAAATATT
GGTGGAATCTCCTACAATATTGGAGTCAGGAGCTAAAGA ATAG 25 Envelope; CMV
ACATTGATTATTGACTAGTTATTAATAGTAATCAATTACG promoter;
GGGTCATTAGTTCATAGCCCATATATGGAGTTCCGCGTT Transcription
ACATAACTTACGGTAAATGGCCCGCCTGGCTGACCGCCC
AACGACCCCCGCCATTGACGTCAATAATGACGTATGTT
CCCATAGTAACGCCAATAGGGACTTTCCATTGACGTCAA
TGGGTGGAGTATTTACGGTAAACTGCCCACTTGGCAGTA
CATCAAGTGTATCATATGCCAAGTACGCCCCCTATTGAC
GTCAATGACGGTAAATGGCCCGCCTGGCATTATGCCCAG
TACATGACCTTATGGGACTTTCTACTTGGCAGTACATCT
ACGTATTAGTCATCGCTATTACCATGGTGATGCGGTTTTG
GCAGTACATCAATGGGCGTGGATAGCGGTTTGACTCACG
GGGATTTCCAAGTCTCCACCCCATGACGTCAATGGGAG
TTTGTTTTTGGCACCAAAATCAACGGGACTTTCCAAAATG
TCGTAACAACTCCGCCCCATTGACGCAAATGGGCGGTAG
GCGTGTACGGTGGGAGGTCTATATAAGC 26 Envelope; VSV-
ATGAAGTGCCTTTTTGTACTTAGCCTTTTTTATTCATTGGGG G; Glycoprotein
TGAATTGCAAGTTCACCATAGTTTTTCCACACAACCAAA envelope-cell
AAGGAAACTGGAAAAATGTTCTTCTAATTACCATTATT entry
GCCCCGTCAAGCTCAGATTTAAATTGGCATAATGACTTAA
TAGGCACAGCCTTACAAGTCAAAATGCCCAAGAGTCACA
AGGCTATTCAAGCAGACGGTTGGATGTGTTCATGCTTCCA
AATGGGTCACTACTTGTGATTTCCGCTGGTATGGACCGA

AGTATATAACACATCCGATCCCTTCACTCCATCTGT
AGAACAATGCAAGGAAAGCATTGAACAAACGAAACAAG
GAACTTGGCTGAATCCAGGCTTCCCTCCTCAAAGTTGTG
GATATGCAACTGTGACGGATGCCGAAGCAGTGATTGTCC
AGGTGACTCCTCACCATGTGCTGGTTGATGAATACACAG
GAGAATGGGTTGATTCACAGTTCATCAACGGAAAATGCA
GCAATTACATATGCCCCACTGTCCATAACTCTACAACCT
GGCATTCTGACTATAAGGTCAAAGGGCTATGTGATTCTA
ACCTCATTTCCATGGACATCACCTTCTTCTCAGAGGACG
GAGAGCTATCATCCCTGGGAAAGGAGGGGCACAGGGTTC
AGAAGTAACTACTTTGCTTATGAAACTGGAGGCAAGGCC
TGCAAAATGCAATACTGCAAGCATTGGGGAGTCAGACTC
CCATCAGGTGTCTGGTTTCGAGATGGCTGATAAGGATCTC
TTTGCTGCAGCCAGATTCCCTGAATGCCCAGAAGGGTCA
AGTATCTCTGCTCCATCTCAGACCTCAGTGGATGTAAGT
CTAATTCAGGACGTTGAGAGGATCTTGGATTATTCCCTCT
GCCAAGAAACCTGGAGCAAAATCAGAGCGGGTCTTCCA
ATCTCTCCAGTGGATCTCAGCTATCTTGCTCCTAAAAACC
CAGGAACCGGTCTGCTTTCACCATAATCAATGGTACCC
TAAAATACTTTGAGACCAGATACATCAGAGTCGATATTG
CTGCTCCAATCCTCTCAAGAATGGTCGGAATGATCAGTG
GAACTACCACAGAAAGGGAAGTGTGGGATGACTGGGCA
CCATATGAAGACGTGGAAATTGGACCCAATGGAGTTCTG
AGGACCAGTTCAGGATATAAGTTTCCTTTATACATGATT
GGACATGGTATGTTGGACTCCGATCTTCATCTTAGCTCA
AAGGCTCAGGTGTTGGAACATCCTCACATTCAAGACGCT
GCTTCGCAACTTCCTGATGATGAGAGTTTATTTTTTTGGTG
ATACTGGGCTATCCAAAAATCCAATCGAGCTTGTAGAAG
GTTGGTTCAGTAGTTGGAAAAGCTCTATTGCCTCTTTTTT
CTTTATCATAGGGTTAATCATTGGACTATTCTTGGTTCTC
CGAGTTGGTATCCATCTTTGCATTAAATTAAAGCACACC
AAGAAAAGACAGATTTATACAGACATAGAGATGA 27 Helper/Rev; CMV
TAGTTATTAATAGTAATCAATTACGGGGTCATTAGTTCAT early (CAG)
AGCCCATATATGGAGTTCCGCGTTACATAACTTACGGTA enhancer; Enhance
AATGGCCCGCCTGGCTGACCGCCCAACGACCCCCGCCCA Transcription
TTGACGTCAATAATGACGTATGTTCCCATAGTAACGCCA
ATAGGGACTTTCCATTGACGTCAATGGGTGGACTATTTA
CGGTAAACTGCCCACCTTGGCAGTACATCAAGTGTATCAT
ATGCCAAGTACGCCCCCTATTGACGTCAATGACGGTAAA
TGGCCCGCCTGGCATTATGCCCAGTACATGACCTTATGG
GACTTTCCTACTTGGCAGTACATCTACGTATTAGTCATC 28 Helper/Rev;
GGAGTCGCTGCGTTGCCTTCGCCCCGTGCCCCGCTCCGC Chicken beta actin
GCCGCTCGCGCCGCCCCGCCCGGGCTCTGACTGACCGCG intron; Enhance
TACTCCCACAGGTGAGCGGGCGGGACGGCCCTTCTCCT gene expression
CCGGGCTGTAATTAGCGCTTGGTTTAATGACGGCTCGTTT
CTTTTCTGTGGCTGCGTGAAAGCCTTAAAGGGCTCCGGG
AGGGCCCTTTGTGCGGGGGGGAGCGGCTCGGGGGGTGC
GTGCGTGTGTGTGTGCGTGGGGAGCGCCGCGTGCGGCC
GCGCTGCCCGGCGGCTGTGAGCGCTGCGGGCGCGGCGC
GGGGCTTTGTGCGCTCCGCGTGTGCGCGAGGGGAGCGCG

GCCGGCGGCGGTGCCCGGCTGCGGGGGGGGCTGCGAG
GGGAACAAAGGCTGCGTGCGGGGTGTGTGCGTGGGGGG
GTGAGCAGGGGGTGTGGGCGCGGCGGTCGGGCTGTAAC
CCCCCCCCTGCACCCCCCTCCCCGAGTTGCTGAGCACGGC
CCGGCTTCGGGTGCGGGGGCTCCGTGCGGGGGCGTGGCGCG
GGGCTCGCCGTGCCGGGGCGGGGGGTGGCGGCAGGTGGG
GGTGCCGGGGCGGGGGCGGGGGCCGCTCGGGCCGGGGAGG
GCTCGGGGGAGGGGGCGCGGCGGCCCCCGAGCGCCGGCG
GCTGTGAGGCGCGGCGAGCCGCAGCCATTGCCTTTTAT
GGTAATCGTGCGAGAGGGGCGCAGGGACTTCCTTTGTCCC
AAATCTGGCGGAGCCGAAATCTGGGAGGCGCCGCCGCA
CCCCCTCTAGCGGGCGCGGGCGAAGCGGTGCGGCGCCG
GCAGGAAGGAAATGGGCGGGGAGGGCCTTCGTGCGTCG
CCGCGCCGCGCGTCCCCCTTCTCCATCTCCAGCCTCGGGGCT
GCCGCAGGGGGACGGCTGCCTTCGGGGGGGGACGGGGCA
GGGCGGGGTTCGGCTTCTGGCGTGTGACCGGCGG 29 Helper/Rev;
AGATCTTTTTCCCTCTGCCAAAAATTATGGGGACATCAT Rabbit beta globin
GAAGCCCCTTGAGCATCTGACTTCTGGCTAATAAAGGAA poly A; RNA
ATTTATTTTCATTGCAATAGTGTGTTGGAATTTTTTGTGT stability
CTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTA
AACATCAGAATGAGTATTTGGTTTAGAGTTTGGCAACA
TATGCCATATGCTGGCTGCCATGAACAAAGGTGGCTATA
AAGAGGTCATCAGTATATGAAACAGCCCCCTGCTGTCCA
TTCCTTATTCCATAGAAAAGCCTTGACTTGAGGTTAGATT
TTTTTTATATTTTGTTTTGTGTTATTTTTTTCTTTAACATCC
CTAAAATTTTCCTTACATGTTTTACTAGCCAGATTTTTCC
TCCTCTCCTGACTACTCCCAGTCATAGCTGTCCCTCTTCT CTTATGAAGATC 30
Envelope; Beta GTGAGTTTGGGGACCCTTGATTGTTCTTTCTTTTTTCGCTA globin intron;
TTGTAAAATTCATGTTATATGGAGGGGGCAAAGTTTTCA Enhance gene
GGGTGTTGTTTAGAATGGGAAGATGTCCCTTGATCACC expression
ATGGACCCTCATGATAATTTTGTTTCTTTCACTTTCTACT
CTGTTGACAACCATTGTCTCCTCTTATTTTCTTTTCATTTT
CTGTAACTTTTTCGTTAAACTTTAGCTTGCATTTGTAACG
AATTTTTTAAATTCACTTTTGTTTATTTGTCAGATTGTAAG
TACTTTCTCTAATCACTTTTTTTTTCAAGGCAATCAGGGTA
TATTATATTGTACTTCAGCACAGTTTTAGAGAACAATTGT
TATAATTAAATGATAAGGTAGAATATTTCTGCATATAAA
TTCTGGCTGGCGTGGAAATATTCTTATTGGTAGAAACAA
CTACACCCTGGTCATCATCCTGCCTTTCTCTTTATGGTTA
CAATGATATACACTGTTTGAGATGAGGATAAAATACTCT
GAGTCCAAACCGGGCCCCCTCTGCTAACCATGTTTCATGCC TTCTTCTCTTTCCTACAG 31
Envelope; Rabbit AGATCTTTTTCCCTCTGCCAAAAATTATGGGGACATCAT beta
globin poly GAAGCCCCTTGAGCATCTGACTTCTGGCTAATAAAGGAA A; RNA
stability ATTTATTTTCATTGCAATAGTGTGTTGGAATTTTTTGTGT
CTCTCACTCGGAAGGACATATGGGAGGGCAAATCATTTA
AACATCAGAATGAGTATTTGGTTTAGAGTTTGGCAACA
TATGCCCATATGCTGGCTGCCATGAACAAAGGTGGCTA
TAAAGAGGTCATCAGTATATGAAACAGCCCCCTGCTGTC
CATTCCTTATTCCATAGAAAAGCCTTGACTTGAGGTTAG
ATTTTTTTTTATATTTTGTTTTGTGTTATTTTTTTCTTTAACA

TCCCTAAATTCCTTACTGTTTTACTAGCCAGATTTT
TCCTCCTCTCCTGACTACTCCCAGTCATAGCTGTCCCTCT TCTCTTATGGAGATC 32
Primer TAAGCAGAATTCATGAATTTGCCAGGAAGAT 33 Primer
CCATACAATGAATGGACACTAGGCGGCCGCACGAAT 34 Gag, Pol,
GAATTCATGAATTTGCCAGGAAGATGGAAACCAAAAAT Integrase fragment
GATAGGGGGAATTGGAGGTTTTATCAAAGTAAGACAGT
ATGATCAGATACTCATAGAAATCTGCGGACATAAAGCTA
TAGGTACAGTATTAGTAGGACCTACACCTGTCAACATAA
TTGGAAGAAATCTGTTGACTCAGATTGGCTGCACTTTAA
ATTTTCCCATTAGTCCTATTGAGACTGTACCAGTAAAATT
AAAGCCAGGAATGGATGGCCCAAAGTTAAACAATGGC
CATTGACAGAAGAAAAAATAAAAGCATTAGTAGAAATT
TGTACAGAAATGGAAAAGGAAGGAAAAATTTCAAAAAT
TGGGCCTGAAAATCCATACAATACTCCAGTATTTGCCAT
AAAGAAAAAAGACAGTACTAAATGGAGAAAATTAGTAG
ATTTCAGAGAACTTAATAAGAGAACTCAAGATTTCTGGG
AAGTTCAATTAGGAATACCACATCCTGCAGGGTTAAAC
AGAAAAAATCAGTAACAGTACTGGATGTGGGCGATGCA
TATTTTTTCAGTTCCCTTAGATAAAGACTTCAGGAAGTATA
CTGCATTTACCATACCTAGTATAAACAATGAGACACCAG
GGATTAGATATCAGTACAATGTGCTTCCACAGGGATGGA
AAGGATCACCAGCAATATTCCAGTGTAGCATGACAAAA
ATCTTAGAGCCTTTTAGAAAACAAAATCCAGACATAGTC
ATCTATCAATACATGGATGATTTGTATGTAGGATCTGAC
TTAGAAATAGGGCAGCATAGAACAAAAATAGAGGAACT
GAGACAACATCTGTTGAGGTGGGGATTTACCACACCAGA
CAAAAAACATCAGAAAGAACCTCCATTCCTTTGGATGGG
TTATGAACTCCATCCTGATAAATGGACAGTACAGCCTAT
AGTGCTGCCAGAAAAGGACAGCTGGACTGTCAATGACA
TACAGAAATTAGTGGGAAAATTGAATTGGGCAAGTCAG
ATTTATGCAGGGATTAAAGTAAGGCAATTATGTAAACTT
CTTAGGGGAACCAAAGCACTAACAGAAGTAGTACCACT
AACAGAAGAAGCAGAGCTAGAACTGGCAGAAAACAGGG
AGATTCTAAAAGAACCGGTACATGGAGTGTATTATGACC
CATCAAAAGACTTAATAGCAGAAATACAGAAGCAGGGG
CAAGGCCAATGGACATATCAAATTTATCAAGAGCCATTT
AAAAATCTGAAAACAGGAAAGTATGCAAGAATGAAGGG
TGCCCACTAATGATGTGAAACAATTAACAGAGGCAGT
ACAAAAAATAGCCACAGAAAGCATAGTAATATGGGGAA
AGACTCCTAAATTTAAATTACCCATACAAAAGGAAACAT
GGGAAGCATGGTGGACAGAGTATTGGCAAGCCACCTGG
ATTCCTGAGTGGGAGTTTGTCAATACCCCTCCCTTAGTGA
AGTTATGGTACCAGTTAGAGAAAGAACCCATAATAGGA
GCAGAAACTTTCTATGTAGATGGGGCAGCCAATAGGGA
AACTAAATTAGGAAAAGCAGGATATGTAAGTACAGAG
GAAGACAAAAAGTTGTCCCCCTAACGGACACAACAAT
CAGAAGACTGAGTTACAAGCAATTCATCTAGCTTTGCAG
GATTCGGGATTAGAAGTAAACATAGTGACAGACTCACA
ATATGCATTGGGAATCATTCAAGCACAACCAGATAAGAG
TGAATCAGAGTTAGTCAGTCAAATAATAGAGCAGTTAAT

AAAAAAGGAATTGGAGGAAATGAACAAGTAGATAAATTG
GTCAGTGCTGGAATCAGGAAAGTACTATTTTTTAGATGGA
ATAGATAAGGCCCAAGAAGACATGAGAAATATCACAG
TAATTGGAGAGCAATGGCTAGTGATTTTAACCTACCACC
TGTAGTAGCAAAAGAAATAGTAGCCAGCTGTGATAAAT
GTCAGCTAAAAGGGGAAGCCATGCATGGACAAGTAGAC
TGTAGCCCAGGAATATGGCAGCTAGATTGTACACATTTA
GAAGGAAAAGTTATCTTGGTAGCAGTTCATGTAGCCAGT
GGATATATAGAAGCAGAAGTAATTCCAGCAGAGACAGG
GCAAGAAACAGCATACTTCCTCTTAAAATTAGCAGGAAG
ATGGCCAGTAAAAACAGTACATACAGACAATGGCAGCA
ATTTACCAGTACTACAGTTAAGGCCGCCTGTTGGTGGG
CGGGGATCAAGCAGGAATTTGGCATTCCCTACAATCCCC
AAAGTCAAGGAGTAATAGAATCTATGAATAAAGAATTA
AAGAAAATTATAGGACAGGTAAGAGATCAGGCTGAACA
TCTTAAGACAGCAGTACAAATGGCAGTATTCATCCACAA
TTTTAAAAGAAAAGGGGGGATTGGGGGGGTACAGTGCAG
GGGAAAGAATAGTAGACATAATAGCAACAGACATACAA
ACTAAAGAATTACAAAAACAAATTACAAAAATTCAAAA
TTTTCGGGTTTATTACAGGGACAGCAGAGATCCAGTTTG
GAAAGGACCAGCAAAGCTCCTCTGGAAAGGTGAAGGGG
CAGTAGTAATACAAGATAATAGTGACATAAAAGTAGTG
CCAAGAAGAAAAGCAAAGATCATCAGGGATTATGGAAA
ACAGATGGCAGGTGATGATTGTGTGGCAAGTAGACAGG ATGAGGATTAA 35 DNA
Fragment TCTAGAATGGCAGGAAGAAGCGGAGACAGCGACGAAGA containing Rev,
GCTCATCAGAACAGTCAGACTCATCAAGCTTCTCTATCA RRE and rabbit
AAGCAACCCACCTCCCAATCCCGAGGGGACCCGACAGG beta globin poly A
CCCGAAGGAATAGAAGAAGAAGGTGGAGAGAGAGACA
GAGACAGATCCATTTCGATTAGTGAACGGATCCTTGGCAC
TTATCTGGGACGATCTGCGGAGCCTGTGCCTCTTCAGCT
ACCACCGCTTGAGAGACTTACTCTTGATTGTAACGAGGA
TTGTGGAACCTTCTGGGACGCAGGGGGTGGGAAGCCCTCA
AATATTGGTGGAAATCTCCTACAATATTGGAGTCAGGAGC
TAAAGAATAGAGGAGCTTTGTTCCTTGGGTTCTTGGGAG
CAGCAGGAAGCACTATGGGCGCAGCGTCAATGACGCTG
ACGGTACAGGCCAGACAATTATTGTCTGGTATAGTGCAG
CAGCAGAACAATTTGCTGAGGGCTATTGAGGCGCAACA
GCATCTGTTGCAACTCACAGTCTGGGGCATCAAGCAGCT
CCAGGCAAGAATCCTGGCTGTGGAAAGATACCTAAAGG
ATCAACAGCTCCTAGATCTTTTTCCCTCTGCCAAAATTA
TGGGGACATCATGAAGCCCCTTGAGCATCTGACTTCTGG
CTAATAAAGGAAATTTATTTTCATTGCAATAGTGTGTTG
GAATTTTTTGTGTCTCTCACTCGGAAGGACATATGGGAG
GGCAAATCATTTAAAACATCAGAATGAGTATTTGGTTTA
GAGTTTGGCAACATATGCCATATGCTGGCTGCCATGAAC
AAAGGTGGCTATAAAGAGGTCATCAGTATATGAAACAG
CCCCCTGCTGTCCATTTCCTTATTCCATAGAAAAGCCTTGA
CTTGAGGTTAGATTTTTTTTTTATATTTTGTGTTTGTGTTATTT
TTTTCTTTAACATCCCTAAAATTTTCCTTACATGTTTTACT

AGCCAGATTTTCTTCTCTTCTTATGAAGATCCCTCGACCTGCAGC
GCTGTCCCTCTTCTCTTATGAAGATCCCTCGACCTGCAGC
CCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCTGTGT
GAAATTGTTATCCGCTCACAATTCCACACAACATACGAG
CCGGAAGCATAAAGTGTAAGCCTGGGGTGCCTAATGA
GTGAGCTAACTCACATTAATTGCGTTGCGCTCACTGCCC
GCTTTCCAGTCGGGAAACCTGTCGTGCCAGCGGATCCGC
ATCTCAATTAGTCAGCAACCATAGTCCCGCCCCCTAACTC
CGCCCATCCCGCCCCCTAACTCCGCCAGTTCCGCCCATTC
TCCGCCCCATGGCTGACTAATTTTTTTTTTATTTATGCAGAG
GCCGAGGCCGCCCTCGGCCTCTGAGCTATTCCAGAAGTAG
TGAGGAGGCTTTTTTGGAGGCCTAGGCTTTTGCAAAAAG
CTAACTTGTTTATTGCAGCTTATAATGGTTACAAATAAA
GCAATAGCATCACAAATTCACAAATAAAGCATTTTTTT
CACTGCATTCTAGTTGTGGTTTGTCCAAACTCATCAATGT
ATCTTATCAGCGGCCGCCCGGG 36 DNA fragment

ACGCGTTAGTTATTAATAGTAATCAATTACGGGGTCATT containing the
AGTTCATAGCCCATATATGGAGTTCCGCGTTACATAACT CAG
TACGGTAAATGGCCCGCCTGGCTGACCGCCCAACGACCC enhancer/promoter/
CCGCCCATTGACGTCAATAATGACGTATGTTCCCATAGT intron sequence
AACGCCAATAGGGACTTTCATTGACGTCAATGGGTGGA
CTATTTACGGTAAACTGCCCCACTTGGCAGTACATCAAGT
GTATCATATGCCAAGTACGCCCCCTATTGACGTCAATGA
CGGTAAATGGCCCGCCTGGCATTATGCCCAGTACATGAC
CTTATGGGACTTTCCTACTTGGCAGTACATCTACGTATTA
GTCATCGCTATTACCATGGGTGAGGTGAGCCCCACGTT
CTGCTTCACTCTCCCCATCTCCCCCCCCCTCCCCACCCCCA
ATTTTGTATTTATTTATTTTTTAATTATTTTGTGCAGCGAT
GGGGGCGGGGGGGGGGGGGGGGGGGCGCGCGCCAGGCGGGGC
GGGGCGGGGCGAGGGGCGGGGCGGGGCGAGGCGGAGA
GGTGCGGCGGCAGCCAATCAGAGCGGCGCGCTCCGAAA
GTTTCCTTTTATGGCGAGGCGGCGGCGGCGGCGGCCCTA
TAAAAAGCGAAGCGCGCGGGCGGGCGGGAGTCGCTGCGT
TGCCTTCGCCCCGTGCCCCGCTCCGCGCCGCTCGCGCC
GCCCCGCCCCGGCTCTGACTGACCGCGTTACTCCACAGG
TGAGCGGGCGGGACGGCCCTTCTCCTCCGGGCTGTAATT
AGCGCTTGGTTTAATGACGGCTCGTTTCTTTTCTGTGGCT
GCGTGAAAGCCTTAAAGGGCTCCGGGAGGGCCCTTTGTG
CGGGGGGGAGCGGCTCGGGGGGTGCGTGCGTGTGTGTG
TGCGTGGGGAGCGCCGCGTGCGGCCCGCGCTGCCCGGCG
GCTGTGAGCGCTGCGGGCGCGGCGCGGGGCTTTGTGCGC
TCCGCGTGTCGCGAGGGGAGCGCGGCCGGGGGCGGTG
CCCCGCGGTGCGGGGGGGCTGCGAGGGGAACAAAGGCT
GCGTGCGGGGTGTGTGCGTGGGGGGGTGAGCAGGGGGT
GTGGGCGCGGCGGTGCGGGCTGTAACCCCCCTGCACCC
CCCTCCCCGAGTTGCTGAGCACGGCCCGGCTTCGGGTGC
GGGGCTCCGTGCGGGGCGTGGCGCGGGGCTCGCCGTGCC
GGGCGGGGGGTGGCGGCAGGTGGGGGTGCCGGGCGGGG
CGGGGCCGCCCTCGGGCCGGGGAGGGCTCGGGGGAGGGG
CGCGGCGGCCCGGAGCGCCGGCGGCTGTGAGGCGCG

GCGAGCGCGAGCTTTTATGGTAAATCGTGCGAG
AGGGCGCAGGGACTTCCTTTGTCCCAAATCTGGCGGAGC
CGAAATCTGGGAGGGCGCCGCCGCACCCCTCTAGCGGGC
GCGGGCGAAGCGGTGCGGGCGCCGGCAGGAAGGAAATGG
GCGGGGAGGGCCTTCGTGCGTCGCCGCGCCGCGTCCCC
TTCTCCATCTCCAGCCTCGGGGGCTGCCGCAGGGGGACGG
CTGCCTTCGGGGGGGACGGGGCAGGGCGGGGTTCTGGCTT
CTGGCGTGTGACCGGCGGGGAATTC 37 DNA fragment
GAATTCATGAAGTGCCTTTTGTACTTAGCCTTTTTATTCA containing VSV-G
TTGGGGTGAATTGCAAGTTCACCATAGTTTTTCCACACA
ACCAAAAAGGAAACTGGAAAAATGTTCTTCTAATTACC
ATTATTGCCCCGTCAAGCTCAGATTTAAATTGGCATAATG
ACTTAATAGGCACAGCCTTACAAGTCAAAATGCCCAAGA
GTCACAAGGCTATTCAAGCAGACGGTTGGATGTGTCATG
CTTCCAAATGGGTCACTACTTGTGATTTCCGCTGGTATGG
ACCGAAGTATATAACACATTCCATCCGATCCTTCACTCC
ATCTGTAGAACAATGCAAGGAAAGCATTGAACAAACGA
AACAAGGAACTTGGCTGAATCCAGGCTTCCCTCCTCAAA
GTTGTGGATATGCAACTGTGACGGATGCCGAAGCAGTGA
TTGTCCAGGTGACTCCTCACCATGTGCTGGTTGATGAAT
ACACAGGAGAATGGGTTGATTCACAGTTCATCAACGGAA
AATGCAGCAATTACATATGCCCCACTGTCCATAACTCTA
CAACCTGGCATTCTGACTATAAGGTCAAAGGGCTATGTG
ATTCTAACCTCATTTCATGGACATCACCTTCTTCTCAGA
GGACGGAGAGCTATCATCCCTGGGAAAGGAGGGCACAG
GGTTCAGAAGTAACTACTTTGCTTATGAAACTGGAGGCA
AGGCCTGCAAAATGCAATACTGCAAGCATTGGGGAGTC
AGACTCCCATCAGGTGTCTGGTTCGAGATGGCTGATAAG
GATCTCTTTGCTGCAGCCAGATTCCCTGAATGCCCAGAA
GGGTCAAGTATCTCTGCTCCATCTCAGACCTCAGTGGAT
GTAAGTCTAATTCAGGACGTTGAGAGGATCTTGGATTAT
TCCCTCTGCCAAGAAACCTGGAGCAAAATCAGAGCGGGT
CTTCCAATCTCTCCAGTGGATCTCAGCTATCTTGCTCCTA
AAAACCCAGGAACCGGTCCTGCTTTCACCATAATCAATG
GTACCCTAAAATACTTTGAGACCAGATACATCAGAGTCG
ATATTGCTGCTCCAATCCTCTCAAGAATGGTCGGAATGA
TCAGTGGAACCTACCACAGAAAGGGAACTGTGGGATGAC
TGGGCACCATATGAAGACGTGGAAATTGGACCCAATGG
AGTTCTGAGGACCAGTTCAGGATATAAGTTTCCTTTATA
CATGATTGGACATGGTATGTTGGACTCCGATCTTCATCTT
AGCTCAAAGGCTCAGGTGTTTGAACATCCTCACATTCAA
GACGCTGCTTCGCAACTTCCTGATGATGAGAGTTTATTTT
TTGGTGATACTGGGCTATCCAAAAATCCAATCGAGCTTG
TAGAAGGTTGGTTCAGTAGTTGGAAAAGCTCTATTGCCT
CTTTTTTCTTTATCATAGGGTTAATCATTGGACTATTCTT
GGTTCTCCGAGTTGGTATCCATCTTTGCATTAAATTAAAG
CACACCAAGAAAAGACAGATTTATACAGACATAGAGAT GAGAATTC 38 RSV
promoter and CAATTGCGATGTACGGGCCAGATATACGCGTATCTGAGG HIV Rev
GGACTAGGGTGTGTTTAGGCGAAAAGCGGGGCTTCGGTT
GTACGCGGTTAGGAGTCCCCTCAGGATATAGTAGTTTCG

CTTTTGTAGGTAGGAGGAAATGTAGTCTTATGCAATA
CACTTGTAGTCTTGCAACATGGTAACGATGAGTTAGCAA
CATGCCTTACAAGGAGAGAAAAAGCACCGTGCATGCCG
ATTGGTGGAAGTAAGGTGGTACGATCGTGCCTTATTAGG
AAGGCAACAGACAGGTCTGACATGGATTGGACGAACCA
CTGAATTCGCGATTGCAGAGATAATTGTATTTAAGTGCC
TAGCTCGATAACAATAAACGCCATTTGACCATTACACCACA
TTGGTGTGCACCTCCAAGCTCGAGCTCGTTTAGTGAACC
GTCAGATCGCCTGGAGACGCCATCCACGCTGTTTTGACC
TCCATAGAAGACACCGGGACCGATCCAGCCTCCCCTCGA
AGCTAGCGATTAGGCATCTCCTATGGCAGGAAGAAGCG
GAGACAGCGACGAAGAAGTCTCAAGGCAGTCAGACTC
ATCAAGTTTCTCTATCAAAGCAACCCACCTCCCAATCCC
GAGGGGACCCGACAGGCCCGAAGGAATAGAAGAAGAA
GGTGGAGAGAGAGACAGAGACAGATCCATTTCGATTAGT
GAACGGATCCTTAGCACTTATCTGGGACGATCTGCGGAG
CCTGTGCCTCTTCAGCTACCACCGCTTGAGAGACTTACTC
TTGATTGTAACGAGGATTGTGGAAGTTCTGGGACGCAGG
GGGTGGAAGCCCTCAAATATTGGTGGAAATCTCCTACAA
TATTGGAGTCAGGAGCTAAAGAATAGTCTAGA 39 Elongation Factor-
CCGGTGCCTAGAGAAGGTGGCGCGGGGTAAACTGGGAA 1 alpha (EF1-
AGTGATGTCGTGTACTGGCTCCGCCTTTTTTCCCGAGGGTG alpha) promoter
GGGGAGAACCGTATATAAGTGCAGTAGTCGCCGTGAAC
GTTCTTTTTTCGCAACGGGTTTGCCGCCAGAACACAGGTA
AGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACG
GGTTATGGCCCTTGCGTGCCTTGAATTACTTCCACGCCCC
TGGCTGCAGTACGTGATTCTTGATCCCGAGCTTCGGGTT
GGAAGTGGGTGGGAGAGTTTCGAGGCCTTGCGCTTAAGG
AGCCCCCTTCGCCTCGTGCTTGAGTTGAGGCCTGGCCTGG
GCGCTGGGGCCGCGCGGTGCGAATCTGGTGGCACCTTCG
CGCCTGTCTCGCTGCTTTCGATAAGTCTCTAGCCATTTAA
AATTTTTGATGACCTGCTGCGACGCTTTTTTTCTGGCAAG
ATAGTCTTGTAATGCGGGCCAAGATCTGCACACTGGTA
TTTCGGTTTTTTGGGGCCGCGGGCGGCGACGGGGCCCGTG
CGTCCCAGCGCACATGTTTCGGCGAGGCGGGGCCTGCGAG
CGCGGCCACCGAGAATCGGACGGGGGTAGTCTCAAGCT
GGCCGGCCTGCTCTGGTGCCTGGCCTCGCGCCGCGGTGT
ATCGCCCCGCCCTGGGCGGCAAGGCTGGCCCCGGTCGGCA
CCAGTTGCGTGAGCGGAAAGATGGCCGCTTCCCGGCCCT
GCTGCAGGGAGCTCAAATGGAGGACGCGGCGCTCGGG
AGAGCGGGCGGGTGAGTCACCCACACAAAGGAAAAGGG
CCTTTCCGTCTCAGCCGTCGCTTCATGTGACTCCACGGA
GTACCGGGCGCCGTCCAGGCACCTCGATTAGTTCTCGAG
CTTTTGGAGTACGTGCTCTTTAGGTTGGGGGGAGGGGTT
TTATGCGATGGAGTTTCCCCACACTGAGTGGGTGGAGAC
TGAAGTTAGGCCAGCTTGGCACTTGATGTAATTCTCCTTG
GAATTTGCCCTTTTTGAGTTTGGATCTTGGTTCATTCTCA
AGCCTCAGACAGTGGTTCAAAGTTTTTTCTTCCATTTCA GGTGTCGTGA 40
Promoter; PGK GGGGTTGGGGTTGCGCCTTTTTCCAAGGCAGCCCTGGGTT
TGCGCAGGGACGCGGCTGCTCTGGGCGTGGTTCCGGGAA

[illegible]

ATGAAACTCCCAAGCAATGGTCAATTTTATGTAGCCTA
ATAATAGTTCGGGCAGGGTTTGACGACCCCCGCAAGGCT
ATCGCATTAGTACAAAAACAACATGGTAAACCATGCGA
ATGCAGCGGAGGGCAGGTATCCGAGGCCCCACCGAACT
CCATCCAACAGGTAACTTGCCCAGGCAAGACGGCCTACT
TAATGACCAACCAAAAATGGAAATGCAGAGTCACTCCA
AAAAATCTCACCCCTAGCGGGGGGAGAACTCCAGAACTG
CCCCTGTAACTTTCCAGGACTCGATGCACAGTTCTTGT
TATACTGAATACCGGCAATGCAGGGCGAATAATAAGAC
ATACTACACGGCCACCTTGCTTAAAATACGGTCTGGGAG
CCTCAACGAGGTACAGATATTACAAAACCCCAATCAGCT
CCTACAGTCCCCTTGTAGGGGCTCTATAAATCAGCCCGT
TTGCTGGAGTGCCACAGCCCCCATCCATATCTCCGATGG
TGGAGGACCCCTCGATACTAAGAGAGTGTGGACAGTCCA
AAAAAGGCTAGAACAAATTCATAAGGCTATGCATCCTGA
ACTTCAATACCACCCCTTAGCCCTGCCCCAAAGTCAGAGA
TGACCTTAGCCTTGATGCACGGACTTTTGATATCCTGAAT
ACCACTTTTAGGTTACTCCAGATGTCCAATTTTAGCCTTG
CCAAGATTGTTGGCTCTGTTTAAAACTAGGTACCCCTA
CCCCTCTTGCGATACCCACTCCCTCTTTAACCTACTCCCT
AGCAGACTCCCTAGCGAATGCCTCCTGTCAGATTATACC
TCCCCTCTTGGTTCAACCGATGCAGTTCTCCAACTCGTCC
TGTTTATCTTCCCCTTTTCATTAACGATACGGAACAAATAG
ACTTAGGTGCAGTCACCTTTACTAACTGCACCTCTGTAGC
CAATGTCAGTAGTCCTTTATGTGCCCTAAACGGGTCAGT
CTTCCTCTGTGGAATAACATGGCATAACCTATTTACCC
CAAACTGGACAGGACTTTGCGTCCAAGCCTCCCTCCTC
CCCGACATTGACATCATCCCGGGGGATGAGCCAGTCCCC
ATTCCTGCCATTGATCATTATATACATAGACCTAAACGA
GCTGTACAGTTCATCCCTTTACTAGCTGGACTGGGAATC
ACCGCAGCATTACCACCGGAGCTACAGGCCTAGGTGTC
TCCGTCACCCAGTATACAAAATTATCCCATCAGTTAATA
TCTGATGTCCAAGTCTTATCCGGTACCATACAAGATTTAC
AAGACCAGGTAGACTCGTTAGCTGAAGTAGTTCTCCAAA
ATAGGAGGGGACTGGACCTACTAACGGCAGAACAAAGGA
GGAATTTGTTTAGCCTTACAAGAAAAATGCTGTTTTTATG
CTACAAGTCAGGAATTGTGAGAAACAAAATAAGAACC
CTACAAGAAGAATTACAAAACGCAGGGAAAGCCTGGC
ATCCAACCTCTCTGGACCGGGCTGCAGGGCTTTCTTCC
GTACCTCCTACCTCTCCTGGGACCCCTACTCACCTCCTA
CTCATACTAACCATTGGGCCATGCGTTTTCAATCGATTGG
TCCAATTTGTAAAGACAGGATCTCAGTGGTCCAGGCTC
TGGTTTTGACTCAGCAATATCACCAGCTAAAACCCATAG AGTACGAGCCATGA 45
Envelope; GALV ATGCTTCTCACCTCAAGCCCGCACACCTTCGGCACCCAG
ATGAGTCCTGGGAGCTGGAAAAGACTGATCATCCTCTTA
AGCTGCGTATTCGGAGACGGCAAAACGAGTCTGCAGAA
TAAGAACCCCCACCAGCCTGTGACCCTCACCTGGCAGGT
ACTGTCCCAAACCTGGGGACGTTGTCTGGGACAAAAAGGC
AGTCCAGCCCCTTTGGACTTGGTGGCCCTCTCTTACACCT
GATGTATGTGCCCTGGCGGCCGGTCTTGAGTCCTGGGAT

ATCCCGGATGTGCTCCTCTAAAAGAGTTAGA
CCTCCTGATTCAGACTATACTGCCGCTTATAAGCAAATC
ACCTGGGGAGCCATAGGGTGCAGCTACCCTCGGGCTAGG
ACCAGGATGGCAAATCCCCCTTCTACGTGTGTCCCCGA
GCTGGCCGAACCCATTCAGAAGCTAGGAGGTGTGGGGG
GCTAGAATCCCTATACTGTAAAGAATGGAGTTGTGAGAC
CACGGGTACCGTTTATTGGCAACCCAAGTCCTCATGGGA
CCTCATAACTGTAAAATGGGACCAAAATGTGAAATGGG
AGCAAAAATTTCAAAAAGTGTGAACAAACCGGCTGGTGT
AACCCCCTCAAGATAGACTTCACAGAAAAAGGAAAACCT
CTCCAGAGATTGGATAACGGAAAAAACCTGGGAATTAA
GGTTCTATGTATATGGACACCCAGGCATACAGTTGACTA
TCCGCTTAGAGGTCACATAACATGCCGGTTGTGGCAGTGG
GCCCAGACCCTGTCCTTGCGGAACAGGGACCTCCTAGCA
AGCCCCCTCACTCTCCCTCTCTCCCCACGGAAAGCGCCGC
CCACCCCTCTACCCCCGGCGGCTAGTGAGCAAACCCCTG
CGGTGCATGGAGAAACTGTTACCCTAAACTCTCCGCCTC
CCACCAGTGGCGACCGACTCTTTGGCCTTGTGCAGGGGG
CCTTCCTAACCTTGAATGCTACCAACCCAGGGGGCCACTA
AGTCTTGCTGGCTCTGTTTGGGCATGAGCCCCCCTTATTA
TGAAGGGATAGCCTCTTCAGGAGAGGTGCTTATACCTC
CAACCATACCCGATGCCACTGGGGGGCCCAAGGAAAGC
TTACCCTCACTGAGGTCTCCGGACTCGGGTCATGCATAG
GGAAGGTGCCTCTTACCCATCAACATCTTTGCAACCAGA
CCTTACCCATCAATTCTCTAAAAACCATCAGTATCTGCT
CCCCTCAAACCATAGCTGGTGGGCCTGCAGCACTGGCCT
CACCCCCTGCCTCTCCACCTCAGTTTTTAATCAGTCTAAA
GACTTCTGTGTCCAGGTCCAGCTGATCCCCCGCATCTATT
ACCATTCTGAAGAAACCTTGTTACAAGCCTATGACAAAT
CACCCCCCAGGTTTAAAAGAGAGCCTGCCTCACTTACCC
TAGCTGTCTTCCTGGGGTTAGGGATTGCGGCAGGTATAG
GTACTGGCTCAACCGCCCTAATTAAAGGGCCCATAGACC
TCCAGCAAGGCCTAACAGCCTCCAAATCGCCATTGACG
CTGACCTCCGGGGCCCTTCAGGACTCAATCAGCAAGCTAG
AGGACTCACTGACTTCCCTATCTGAGGTAGTACTCCAAA
ATAGGAGAGGCCTTGACTTACTATTCCTTAAAGAAGGAG
GCCTCTGCGCGGCCCTAAAAGAAGAGTGCTGTTTTTATG
TAGACCACTCAGGTGCAGTACGAGACTCCATGAAAAAA
CTTAAAGAAAGACTAGATAAAAGACAGTTAGAGCGCCA
GAAAAACCAAAACTGGTATGAAGGGTGGTTCAATAACT
CCCCTTGGTTTACTACCCTACTATCAACCATCGCTGGGCC
CCTATTGCTCCTCCTTTTTGTTACTCACTCTTGGGCCCTGC
ATCATCAATAAATTAATCCAATTCATCAATGATAGGATA
AGTGCAGTCAAAATTTTAGTCCTTAGACAGAAATATCAG
ACCCTAGATAACGAGGAAAACCTTTAA 46 Envelope; FUG
ATGGTTCCGCAGGTTCTTTTGTGTGTACTCCTTCTGGGTT
TTTCGTTGTGTTTCGGGAAGTTCCCCATTTACACGATACC
AGACGAACCTTGGTCCCTGGAGCCCTATTGACATACACCA
TCTCAGCTGTCCAAATAACCTGGTGTGTGGAGGATGAAGG
ATGTACCAACCTGTCCGAGTTCTCCTACATGGAACCTCAA

AGTGGGATCATCAAAAGTGAAACGGGGTTCAC
TTGCACAGGTGTTGTGACAGAGGCAGAGACCTACACCAA
CTTTGTTGGTTATGTCACAACCACATTCAAGAGAAAGCA
TTTCCGCCCCACCCCAGACGCATGTAGAGCCGCGTATAA
CTGGAAGATGGCCGGTGACCCCAGATATGAAGAGTCCCT
ACACAATCCATACCCCGACTACCACTGGCTTCGAACTGT
AAGAACCACCAAAGAGTCCCTCATTATCATATCCCCAAG
TGTGACAGATTTGGACCCATATGACAAATCCCTTCACTC
AAGGGTCTTCCCTGGCGGAAAGTGCTCAGGAATAACGGT
GTCCTCTACCTACTGCTCAACTAACCATGATTACACCAT
TGGATGCCCCGAGAATCCGAGACCAAGGACACCTTGTGAC
ATTTTTACCAATAGCAGAGGGAAGAGAGCATCCAACGG
GAACAAGACTTGCGGCTTTGTGGATGAAAGAGGCCTGTA
TAAGTCTCTAAAAGGAGCATGCAGGCTCAAGTTATGTGG
AGTTCTTGGACTTAGACTTATGGATGGAACATGGGTTCG
GATGCAAACATCAGATGAGACCAAATGGTGCCCTCCAG
ATCAGTTGGTGAATTTGCACGACTTTTCGCTCAGACGAGA
TCGAGCATCTCGTTGTGGAGGAGTTAGTTAAGAAAAGAG
AGGAATGTCTGGATGCATTAGAGTCCATCATGACCACCA
AGTCAGTAAGTTTCAGACGTCTCAGTCACCTGAGAAAAC
TTGTCCCAGGGTTTGGAAAAGCATATAACCATATTCAACA
AAACCTTGATGGAGGCTGATGCTCACTACAAGTCAGTCC
GGACCTGGAATGAGATCATCCCCTCAAAGGGTGTTTGA
AAGTTGGAGGAAGGTGCCATCCTCATGTGAACGGGGTGT
TTTTCAATGGTATAATATTAGGGCCTGACGACCATGTCCT
AATCCCAGAGATGCAATCATCCCTCCTCCAGCAACATAT
GGAGTTGTTGGAATCTTCAGTTATCCCCCTGATGCACCCC
CTGGCAGACCCTTCTACAGTTTTCAAAGAAGGTGATGAG
GCTGAGGATTTTGTGTAAGTTCACCTCCCCGATGTGTAC
AAACAGATCTCAGGGGTGACCTGGGTCTCCCGAACTGG
GGAAAGTATGTATTGATGACTGCAGGGGGCCATGATTGGC
CTGGTGTTGATATTTTCCCTAATGACATGGTGCAGAGTTG
GTATCCATCTTTGCATTAAATTAAGCACACCAAGAAAA
GACAGATTTATACAGACATAGAGATGAACCGACTTGGA AAGTAA 47 Envelope; LCMV
ATGGGTCAGATTGTGACAATGTTTGAGGCTCTGCCTCAC
ATCATCGATGAGGTGATCAACATTGTCATTATTGTGCTTA
TCGTGATCACGGGTATCAAGGCTGTCTACAATTTTGCCA
CCTGTGGGATATTCGCATTGATCAGTTTCCTACTTCTGGC
TGGCAGGTCCTGTGGCATGTACGGTCTTAAGGGACCCGA
CATTTACAAAGGAGTTTACCAATTTAAGTCAGTGGAGTT
TGATATGTCACATCTGAACCTGACCATGCCCAACGCATG
TTCAGCCAACAACCTCCCACCATTACATCAGTATGGGGAC
TTCTGGACTAGAATTGACCTTCACCAATGATTCCATCATC
AGTCACAACTTTTGCAATCTGACCTCTGCCTTCAACAAA
AAGACCTTTGACCACACACTCATGAGTATAGTTTTCGAGC
CTACACCTCAGTATCAGAGGGAACCTCCAACCTATAAGGCA
GTATCCTGCGACTTCAACAATGGCATAACCATCCAATAC
AACTTGACATTCTCAGATCGACAAAGTGCTCAGAGCCAG
TGTAGAACCTTCAGAGGTAGAGTCCTAGATATGTTTAGA
ACTGCCTTCGGGGGGGAAATACATGAGGAGTGGCTGGGG

CTGGAGCAGGACACCTGGTGTAGCCA
GACGAGTTACCAATACCTGATTATACAAAATAGAACCTG
GGAAAACCACTGCACATATGCAGGTCCTTTTGGGATGTC
CAGGATTCTCCTTTCCCAAGAGAAGACTAAGTTCTTCAC
TAGGAGACTAGCGGGCACATTCACCTGGACTTTGTCAGA
CTCTTCAGGGGTGGAGAATCCAGGTGGTTATTGCCTGAC
CAAATGGATGATTCTTGCTGCAGAGCTTAAGTGTTTCGG
GAACACAGCAGTTGCGAAATGCAATGTAAATCATGATGC
CGAATTCTGTGACATGCTGCGACTAATTGACTACAACAA
GGCTGCTTTGAGTAAGTTCAAAGAGGACGTAGAATCTGC
CTTGCACTTATTCAAAACAACAGTGAATTCTTTGATTTC
GATCAACTACTGATGAGGAACCACTTGAGAGATCTGATG
GGGGTGCCATATTGCAATTACTCAAAGTTTTGGTACCTA
GAACATGCAAAGACCGGCGAAACTAGTGTCCCCAAGTG
CTGGCTTGTCACCAATGGTTCTTACTTAAATGAGACCCA
CTTCAGTGATCAAATCGAACAGGAAGCCGATAACATGAT
TACAGAGATGTTGAGGAAGGATTACATAAAGAGGCAGG
GGAGTACCCCCCTAGCATTGATGGACCTTCTGATGTTTT
CACATCTGCATATCTAGTCAGCATCTTCCTGCACCTTGTC
AAAATACCAACACACAGGCACATAAAAGGTGGCTCATG
TCCAAAGCCACACCGATTAAACCAACAAAGGAATTTGTAG
TTGTGGTGCATTTAAGGTGCCTGGTGTA AAAACCGTCTG GAAAAGACGCTGA 48
Envelope; FPV ATGAACACTCAAATCCTGGTTTTTCGCCCTTGTGGCAGTC
ATCCCCACAAATGCAGACAAAATTTGTCTTGGACATCAT
GCTGTATCAAATGGCACCAAAGTAAACACACTCACTGAG
AGAGGAGTAGAAGTTGTCAATGCAACGGAAACAGTGGA
GCGGACAAACATCCCCAAAATTTGCTCAAAAGGGAAAA
GAACCACTGATCTTGGCCAATGCGGACTGTTAGGGACCA
TTACCGGACCACCTCAATGCGACCAATTTCTAGAATTTTC
AGCTGATCTAATAATCGAGAGACGAGAAGGAAATGATG
TTTGTTACCCGGGGAAGTTTGTTAATGAAGAGGCATTGC
GACAAATCCTCAGAGGATCAGGTGGGATTGACAAAGAA
ACAATGGGATTCACATATAGTGGAATAAGGACCAACGG
AACAAC TAGTG CATGTAGAAGATCAGGGTCTTCATTCTA
TGCAGAAATGGAGTGGCTCCTGTCAAATACAGACAATGC
TGCTTTCCCACAAATGACAAAATCATAAAAAACACAAG
GAGAGAATCAGCTCTGATAGTCTGGGGAATCCACCATTC
AGGATCAACCACCGAACAGACCAA ACTATATGGGAGTG
GAAATAAACTGATAACAGTCGGGAGTTCCAAATATCATC
AATCTTTTGTGCCGAGTCCAGGAACACGACCGCAGATAA
ATGGCCAGTCCGGACGGATTGATTTTCATTGGTTGATCTT
GGATCCCAATGATACAGTTACTTTTAGTTTCAATGGGGC
TTTCATAGCTCCAAATCGTGCCAGCTTCTTGAGGGGAAA
GTCCATGGGGATCCAGAGCGATGTGCAGGTGATGCCAA
TTGCGAAGGGGAATGCTACCACAGTGGAGGGACTATAA
CAAGCAGATTGCCTTTTCAAAACATCAATAGCAGAGCAG
TTGGCAAATGCCAAGATATGTAAAACAGGAAAGTTTAT
TATTGGCAACTGGGATGAAGAACGTTCCCGAACCTTCCA
AAAAAAGGAAAAAAAGAGGCCTGTTTGGCGCTATAGCA
GGGTTTATTGAAAATGGTTGGGAAGGTCTGGTCGACGGG

TGGTAGTTTGCAGTAATGCACAAGGAGAAAGG
AACTGCAGCAGACTACAAAAGCACCCAATCGGCAATTG
ATCAGATAACCGGAAAGTTAAATAGACTCATTGAGAAA
ACCAACCAGCAATTTGAGCTAATAGATAATGAATTCACT
GAGGTGGAAAAGCAGATTGGCAATTTAATTAAGTGGACC
AAAGACTCCATCACAGAAGTATGGTCTTACAATGCTGAA
CTTCTTGTTGGCAATGGAAAACCAGCACACTATTGATTTG
GCTGATTCAGAGATGAACAAGCTGTATGAGCGAGTGAG
GAAACAATTAAGGGAAAATGCTGAAGAGGATGGCACTG
GTTGCTTTGAAATTTTTCATAAATGTGACGATGATTGTAT
GGCTAGTATAAGGAACAATACTTATGATCACAGCAAATA
CAGAGAAGAAGCGATGCAAAATAGAATACAAATTGACC
CAGTCAAATTGAGTAGTGGCTACAAAGATGTGATACTTT
GGTTTAGCTTCGGGGCATCATGCTTTTTTGCTTCTTGCCAT
TGCAATGGGCCTTGTTTTTCATATGTGTGAAGAACGGAAA
CATGCGGTGCACTATTTGTATATAA 49 Envelope; RRV
AGTGTAACAGAGCACTTTAATGTGTATAAGGCTACTAGA
CCATACCTAGCACATTGCGCCGATTGCGGGGACGGGTAC
TTCTGCTATAGCCCAGTTGCTATCGAGGAGATCCGAGAT
GAGGCGTCTGATGGCATGCTTAAGATCCAAGTCTCCGCC
CAAATAGGTCTGGACAAGGCAGGCACCCACGCCCACAC
GAAGCTCCGATATATGGCTGGTCATGATGTTTCAGGAATC
TAAGAGAGATTCTTTGAGGGTGTACACGTCCGCAGCGTG
CTCCATACATGGGACGATGGGACACTTCATCGTCGCACA
CTGTCCACCAGGCGACTACCTCAAGGTTTCGTTTCGAGGA
CGCAGATTTCGCACGTGAAGGCATGTAAGGTCCAATACAA
GCACAATCCATTGCCGGTGGGTAGAGAGAAGTTTCGTGGT
TAGACCACACTTTGGCGTAGAGCTGCCATGCACCTCATA
CCAGCTGACAACGGCTCCCACCGACGAGGAGATTGACAT
GCATACACCGCCAGATATACCGGATCGCACCCCTGCTATC
ACAGACGGCGGGCAACGTCAAAATAACAGCAGGCGGCA
GGACTATCAGGTACAACCTGTACCTGCGGCCGTGACAACG
TAGGCACTACCAGTACTGACAAGACCATCAACACATGCA
AGATTGACCAATGCCATGCTGCCGTCACCAGCCATGACA
AATGGCAATTTACCTCTCCATTTGTTCCCAGGGCTGATCA
GACAGCTAGGAAAGGCAAGGTACACGTTCCGTTCCCTCT
GACTAACGTCACCTGCCGAGTGCCGTTGGCTCGAGCGCC
GGATGCCACCTATGGTAAGAAGGAGGTGACCCTGAGATT
ACACCCAGATCATCCGACGCTCTTCTCCTATAGGAGTTT
AGGAGCCGAACCGCACCCGTACGAGGAATGGGTGACA
AGTTCTCTGAGCGCATCATCCAGTGACGGAAGAAGGGA
TTGAGTACCAGTGGGGCAACAACCCGCCGGTCTGCCTGT
GGGCGCAACTGACGACCGAGGGCAAACCCCATGGCTGG
CCACATGAAATCATTAGTACTATTATGGACTATACCCC
GCCGCCACTATTGCCGCAGTATCCGGGGCGAGTCTGATG
GCCCTCCTAACTCTGGCGGCCACATGCTGCATGCTGGCC
ACCGCGAGGAGAAAGTGCCTAACACCGTACGCCCTGAC
GCCAGGAGCGGTGGTACCGTTGACACTGGGGCTGCTTTG
CTGCGCACCGAGGGCGAATGCA 50 Envelope; MLV
ATGGAAGGTCCAGCGTTCTCAAAACCCCTTAAAGATAAG 10A1

ATTAACCGTCTGCTTAATGGTCATGGGGGTCTAT
TTAAGAGTAGGGATGGCAGAGAGCCCCATCAGGTCTTT
AATGTAACCTGGAGAGTCACCAACCTGATGACTGGGCGT
ACCGCCAATGCCACCTCCCTTTTAGGAACTGTACAAGAT
GCCTTCCCAAGATTATATTTTGATCTATGTGATCTGGTCG
GAGAAGAGTGGGACCCTTCAGACCAGGAACCATATGTC
GGGTATGGCTGCAAATACCCCGGAGGGAGAAAGCGGAC
CCGGACTTTTGACTTTTACGTGTGCCCTGGGCATACCGTA
AAATCGGGGTGTGGGGGGCCAAGAGAGGGGCTACTGTGG
TGAATGGGGTTGTGAAACCACCGGACAGGCTTACTGGAA
GCCCACATCATCATGGGACCTAATCTCCCTTAAGCGCGG
TAACACCCCCTGGGACACGGGATGCTCCAAAATGGCTTG
TGGCCCCTGCTACGACCTCTCCAAAGTATCCAATTCCTTC
CAAGGGGCTACTCGAGGGGGCAGATGCAACCCTCTAGTC
CTAGAATTCACTGATGCAGGAAAAAAGGCTAATTGGGA
CGGGCCCAAATCGTGGGGACTGAGACTGTACCGGACAG
GAACAGATCCTATTACCATGTTCTCCCTGACCCGCCAGG
TCCTCAATATAGGGCCCCGCATCCCCATTGGGCCTAATC
CCGTGATCACTGGTCAACTACCCCCCTCCCGACCCGTGC
AGATCAGGCTCCCCAGGCCTCCTCAGCCTCCTCCTACAG
GCGCAGCCTCTATAGTCCCTGAGACTGCCCCACCTTCTC
AACAACCTGGGACGGGAGACAGGCTGCTAAACCTGGTA
GAAGGAGCCTATCAGGCGCTTAACCTCACCAATCCCGAC
AAGACCCAAGAATGTTGGCTGTGCTTAGTGTCGGGACCT
CCTTATTACGAAGGAGTAGCGGTCGTGGGCACTTATACC
AATCATTCTACCGCCCCGGCCAGCTGTACGGCCACTTCC
CAACATAAGCTTACCCTATCTGAAGTGACAGGACAGGGC
CTATGCATGGGAGCACTACCTAAAACTCACCAGGCCTTA
TGTAACACCACCCAAAGTGCCGGCTCAGGATCCTACTAC
CTTGACAGACCCGCTGGAACAATGTGGGCTTGTAGCACT
GGATTGACTCCCTGCTTGTCCACCACGATGCTCAATCTA
ACCACAGACTATTGTGTATTAGTTGAGCTCTGGCCCAGA
ATAATTTACCACTCCCCCGATTATATGTATGGTCAGCTTG
AACAGCGTACCAAATATAAGAGGGAGCCAGTATCGTTG
ACCCTGGCCCTTCTGCTAGGAGGATTAACCATGGGAGGG
ATTGCAGCTGGAATAGGGACGGGGACCACTGCCCTAATC
AAAACCCAGCAGTTTGAGCAGCTTCACGCCGCTATCCAG
ACAGACCTCAACGAAGTCGAAAAATCAATTACCAACCTA
GAAAAGTCACTGACCTCGTTGTCTGAAGTAGTCCTACAG
AACCGAAGAGGCCTAGATTTGCTCTTCCTAAAAGAGGGA
GGTCTCTGCGCAGCCCTAAAAGAAGAATGTTGTTTTAT
GCAGACCACACGGGACTAGTGAGAGACAGCATGGCCAA
ACTAAGGGAAAGGCTTAATCAGAGACAAAAACTATTTG
AGTCAGGCCAAGGTTGGTTTCGAAGGGCAGTTTAATAGAT
CCCCCTGGTTTACCACCTTAATCTCCACCATCATGGGACC
TCTAATAGTACTCTTACTGATCTTACTCTTTGGACCCTGC
ATTCTCAATCGATTGGTCCAATTTGTAAAGACAGGATC
TCAGTGGTCCAGGCTCTGGTTTTGACTCAACAATATCAC
CAGCTAAAACCTATAGAGTACGAGCCATGA 51 Envelope; Ebola
ATGGGTGTTACAGGAATATTGCAGTTACCTCGTGATCGA

TTCAAAGAGGACATTTTCTTTGGGTAATTATCCTTT
TCCAAAGAACATTTTCCATCCCACCTTGGAGTCATCCACA
ATAGCACATTACAGGTTAGTGATGTCGACAAACTGGTTT
GCCGTGACAAACTGTCATCCACAAATCAATTGAGATCAG
TTGGACTGAATCTCGAAGGGAATGGAGTGGCAACTGAC
GTGCCATCTGCAACTAAAAGATGGGGCTTCAGGTCCGGT
GTCCCACCAAAGGTGGTCAATTATGAAGCTGGTGAATGG
GCTGAAAACCTGCTACAATCTTGAAATCAAAAAACCTGAC
GGGAGTGAGTGTCTACCAGCAGCGCCAGACGGGATTCTG
GGGCTTCCCCCGGTGCCGGTATGTGCACAAAGTATCAGG
AACGGGACCGTGTGCCGGAGACTTTGCCTTCCACAAAGA
GGGTGCTTTCTTCCTGTATGACCGACTTGCTTCCACAGTT
ATCTACCGAGGAACGACTTTCGCTGAAGGTGTCGTTGCA
TTTCTGATACTGCCCCAAGCTAAGAAGGACTTCTTCAGC
TCACACCCCTTGAGAGAGCCGGTCAATGCAACGGAGGA
CCCGTCTAGTGGCTACTATTCTACCACAATTAGATATCA
AGCTACCGGTTTTTGGAACCAATGAGACAGAGTATTTGTT
CGAGGTTGACAATTTGACCTACGTCCAACCTTGAATCAAG
ATTCACACCACAGTTTCTGCTCCAGCTGAATGAGACAAT
ATATACAAGTGGGAAAAGGAGCAATACCACGGGAAAAC
TAATTTGGAAGGTCAACCCCGAAATTGATACAACAATCG
GGGAGTGGGCCTTCTGGGAAACTAAAAAACCTCACTA
GAAAAATTCGCAGTGAAGAGTTGTCTTTCACAGCTGTAT
CAAACAGAGCCAAAAACATCAGTGGTCAGAGTCCGGCG
CGAACTTCTTCCGACCCAGGGACCAACACAACAACTGAA
GACCACAAAATCATGGCTTCAGAAAATTCCTCTGCAATG
GTTCAAGTGCACAGTCAAGGAAGGGAAGCTGCAGTGTC
GCATCTGACAACCTTGCCACAATCTCCACGAGTCCTCA
ACCCCCACAACCAAACAGGTCCGGACAACAGCACCC
ACAATACACCCGTGTATAAACTTGACATCTCTGAGGCAA
CTCAAGTTGAACAACATCACCGCAGAACAGACAACGAC
AGCACAGCCTCCGACACTCCCCCGCCACGACCGCAGCC
GGACCCCTAAAAGCAGAGAACACCAACACGAGCAAGGG
TACCGACCTCCTGGACCCCGCCACCACAACAAGTCCCCA
AAACCACAGCGAGACCGCTGGCAACAACAACACTCATC
ACCAAGATACCGGAGAAGAGAGTGCCAGCAGCGGGAAG
CTAGGCTTAATTACCAATACTATTGCTGGAGTCGCAGGA
CTGATCACAGGCGGGAGGAGAGCTCGAAGAGAAGCAAT
TGTCAATGCTCAACCCAAATGCAACCCTAATTTACATTA
CTGGACTACTCAGGATGAAGGTGCTGCAATCGGACTGGC
CTGGATACCATATTTTCGGGCCAGCAGCCGAGGGAATTTA
CATAGAGGGGCTGATGCACAATCAAGATGGTTTAATCTG
TGGGTTGAGACAGCTGGCCAACGAGACGACTCAAGCTCT
TCAACTGTTCTGAGAGCCACAACCGAGCTACGCACCTT
TTCAATCCTCAACCGTAAGGCAATTGATTTCTTGCTGCAG
CGATGGGGCGGCACATGCCACATTTTGGGACCGGACTGC
TGTATCGAACCACATGATTGGACCAAGAACATAACAGAC
AAAATTGATCAGATTATTCATGATTTTGTTGATAAAACC
CTTCCGGACCAGGGGGACAATGACAATTGGTGGACAGG
ATGGAGACAATGGATACCGGCAGGTATTGGAGTTACAG

GCCTTATAATTGCTTATGCTTATTCGTATGCTAA ATTTGTCTTTTAG 52
Polymerase III TTTCCCATGATTCCTTCATATTTGCATATACGATACAAGG shRNA
promoters; CTGTTAGAGAGATAATTGGAATTAATTTGACTGTAAACA U6 promoter
CAAAGATATTAGTACAAAATACGTGACGTAGAAAGTAA
TAATTTCTTGGGTAGTTTGCAGTTTTTAAAATTATGTTTTA
AAATGGACTATCATATGCTTACCGTAACTTGAAAGTATT
TCGATTTCTTGGCTTTATATATCTTGTGGAAGGACGAA AC 53 Polymerase III
CTGCAGTATTTAGCATGCCCCACCCATCTGCAAGGCATT shRNA promoters;
CTGGATAGTGTCAAAACAGCCGGAATCAAGTCCGTTTA 7SK promoter
TCTCAAACCTTTAGCATTTTTGGGAATAAATGATATTTGCTA
TGCTGGTTAAATTAGATTTTAGTTAAATTTCTGCTGAAG
CTCTAGTACGATAAGCAACTTGACCTAAGTGTAAGTTG
AGATTCCTTCAGGTTTATATAGCTTGTGCGCCGCTGGC TACCTC 54 FDPS target
GTCCTGGAGTACAATGCCATT sequence #1 55 FDPS target
GCAGGATTTTCGTTTCAGCACTT sequence #2 56 FDPS target
GCCATGTACATGGCAGGAATT sequence #3 57 FDPS target
GCAGAAGGAGGCTGAGAAAGT sequence #4 58 Non-targeting
GCCGCTTTGTAGGATAGAGCTCGAGCTCTATCCTACAAA sequence GCGGCTTTTT 59
Forward primer AGGAATTGATGGCGAGAAGG 60 Reverse primer
CCCAAAGAGGTCAAGGTAATCA 61 Forward primer AGCGCGGCTACAGCTTCA 62
Reverse primer GGCGACGTAGCACAGCTTCT 63 Left Inverted
CCTGCAGGCAGCTGCGCGCTCGCTCGCTCACTGAGGCCG Terminal Repeat
CCCGGGCGTCGGGCGACCTTTGGTTCGCCCCGGCCTCAGTG (Left ITR)
AGCGAGCGAGCGCGCAGAGAGGGAGTGGCCAACTCCAT CACTAGGGGTTTCCT 64
Right Inverted GAGCGGCCGCAGGAACCCCTAGTGATGGAGTTGGCCACT Terminal
Repeat CCCTCTCTGCGCGCTCGCTCGCTCACTGAGGCCGGGCGA (Right ITR)
CCAAAGGTCGCCCCGACGCCCCGGGCTTTGCCCCGGGCGGCC
TCAGTGAGCGAGCGAGCGCGCAGCTGCCTGCAGG 65 RRE/rabbit poly A
TCTAGAAGGAGCTTTGTTTCTTGGGTTCTTGGGAGCAGC beta globin
AGGAAGCACTATGGGCGCAGCGTCAATGACGCTGACGG
TACAGGCCAGACAATTATTGTCTGGTATAGTGCAGCAGC
AGAACAATTTGCTGAGGGCTATTGAGGCGCAACAGCATC
TGTTGCAACTCACAGTCTGGGGCATCAAGCAGCTCCAGG
CAAGAATCCTGGCTGTGGAAGATACCTAAAGGATCAA
CAGCTCCTAGATCTTTTTCCCTCTGCCAAAATTATGGGG
ACATCATGAAGCCCCTTGAGCATCTGACTTCTGGCTAAT
AAAGGAAATTTATTTTCATTGCAATAGTGTGTTGGAATTT
TTTGTGTCTCTCACTCGGAAGGACATATGGGAGGGCAA
TCATTTAAAACATCAGAATGAGTATTTGGTTTAGAGTTT
GGCAACATATGCCATATGCTGGCTGCCATGAACAAAGGT
GGCTATAAAGAGGTCATCAGTATATGAAACAGCCCCCTG
CTGTCCATTCCTTATTCCATAGAAAAGCCTTGACTTGAGG
TTAGATTTTTTTTATATTTTGTGTTTATTTTTTTCTTT
AACATCCCTAAAATTTTCTTACATGTTTTACTAGCCAGA
TTTTTCCTCCTCTCCTGACTACTCCCAGTCATAGCTGTCC
CTCTTCTCTTATGAAGATCCCTCGACCTGCAGCCCAAGCT
TGGCGTAATCATGGTCATAGCTGTTTCCTGTGTGAAATTG
TTATCCGCTCACAATTCCACACAACATACGAGCCGGAAG
CATAAAGTGTAAGCCTGGGGTGCCTAATGAGTGAGCTA
ACTCACATTAATTGCGTTGCGCTCACTGCCCCGCTTTCCAG

TCGGGAACCTGTCGTGCCAGCGGATCCGCATCTCAATT
AGTCAGCAACCATAGTCCCGCCCCCTAACTCCGCCCATCC
CGCCCCTAACTCCGCCCAGTTCCGCCCATTCTCCGCCCCA
TGGCTGACTAATTTTTTTTTTATTTATGCAGAGGCCGAGGCC
GCCTCGGCCTCTGAGCTATTCCAGAAGTAGTGAGGAGGC
TTTTTTGGAGGCCTAGGCTTTTGCAAAAAGCTAACTTGTT
TATTGCAGCTTATAATGGTTACAAATAAAGCAATAGCAT
CACAAATTTACAAATAAAGCATTTTTTTTTCACTGCATTCT
AGTTGTGGTTTGTCCAACTCATCAATGTATCTTATCACC CGGG

(249) While certain of the preferred embodiments of the present invention have been described and specifically exemplified above, it is not intended that the invention be limited to such embodiments. Various modifications may be made thereto without departing from the scope and spirit of the present invention.

Claims

1. A pharmaceutical combination, comprising: (a) a bisphosphonate drug; and (b) a lentiviral particle comprising an envelope protein and a therapeutic vector comprising: an EF-1 alpha promoter and at least one encoded shRNA or microRNA that, when expressed, inhibits production of farnesyl diphosphate synthase (FDPS) wherein the bisphosphonate drug and lentiviral particle are administered together or separately, wherein the at least one encoded shRNA or microRNA comprises a sequence having at least 80 percent identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, or SEQ ID NO: 10.
2. The pharmaceutical combination of claim 1, wherein the at least one encoded shRNA or microRNA activates a GD T cell.
3. The pharmaceutical combination of claim 1, wherein the at least one encoded shRNA or microRNA activates a GD T cell.
4. The pharmaceutical combination of claim 1, wherein the envelope protein is capable of infecting a cancer cell.
5. The pharmaceutical combination of claim 1, wherein the bisphosphonate drug comprises zoledronic acid.
6. A pharmaceutical combination, comprising: (a) a bisphosphonate drug; and (b) a lentiviral particle comprising an envelope protein and a therapeutic vector comprising: at least one encoded shRNA or microRNA that, when expressed, inhibits production of farnesyl diphosphate synthase (FDPS) and activates a GD T cell wherein the bisphosphonate drug and lentiviral particle are administered together or separately, wherein the at least one encoded shRNA or microRNA comprises a sequence having at least 80 percent identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, or SEQ ID NO: 10.
7. The pharmaceutical combination of claim 6, wherein the envelope protein is capable of infecting a cancer cell.
8. The pharmaceutical combination of claim 6, wherein the bisphosphonate drug comprises zoledronic acid.
9. A method of treating a cancer in a subject in need thereof, using an immunotherapy-based composition, the method comprising: (a) administering a therapeutically-effective amount of an bisphosphonate drug to the subject; and (b) administering a therapeutically-effective amount of the immunotherapy-based composition to the subject, wherein the immunotherapy-based composition comprises a lentiviral particle, the lentiviral particle comprising an envelope protein and a therapeutic vector comprising: an EF-1 alpha promoter; and, at least one encoded shRNA or

microRNA that, when expressed, inhibits production of farnesyl diphosphate synthase (FDPS), wherein the at least one encoded shRNA or microRNA comprises a sequence having at least 80 percent identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, or SEQ ID NO: 10.

10. The method of claim 9, wherein the at least one encoded shRNA or microRNA activates a GD T cell.

11. The method of claim 9, wherein the at least one encoded shRNA or microRNA activates a GD T cell.

12. The method of claim 9, wherein the cancer is selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof.

13. The method of claim 9, wherein the bisphosphonate drug comprises zoledronic acid.

14. A method of treating a cancer in a subject in need thereof, using an immunotherapy-based composition, the method comprising: (a) administering a therapeutically-effective amount of an bisphosphonate drug to the subject; and (b) administering a therapeutically-effective amount of the immunotherapy-based composition to the subject, wherein the immunotherapy-based composition comprises a lentiviral particle, the lentiviral particle comprising an envelope protein and a therapeutic vector comprising: at least one encoded shRNA or microRNA that, when expressed, inhibits production of farnesyl diphosphate synthase (FDPS) and activates a GD T cell, wherein the at least one encoded shRNA or microRNA comprises a sequence having at least 80 percent identity with SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3, SEQ ID NO: 4, SEQ ID NO: 5, SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8, SEQ ID NO: 9, or SEQ ID NO: 10.

15. The method of claim 14, wherein the cancer is selected from one or more of a carcinoma, a leukemia, a lymphoma, a sarcoma, a myeloma, a mesothelioma, a mixed type, or mixtures thereof.

16. The method of claim 14, wherein the bisphosphonate drug comprises zoledronic acid.
