

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12404339
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Ober; Elizabeth Sally Ward et al.

Method of depleting target antigen-specific antibody from a patient by administering a fusion protein (Seldeg) for selectively depleting antigen-specific antibodies

Abstract

The present disclosure includes a fusion protein, called a “Seldeg”, including a targeting component that specifically binds to a cell surface receptor or other cell surface molecule at near-neutral pH, and an antigen component fused directly or indirectly to the targeting component. The antigen component is configured to specifically bind a target antigen-specific antibody. The present disclosure also includes a method of depleting a target antigen-specific antibody from a patient by administering to the patient a Seldeg having an antigen component configured to specifically bind the target antigen-specific antibody.

Inventors: Ober; Elizabeth Sally Ward (Southampton, GB), Devanaboyina; Venkata Siva Charan (College Station, TX), Ober; Raimund Johannes (Southampton, GB)

Applicant: THE TEXAS A&M UNIVERSITY SYSTEM (College Station, TX)

Family ID: 62242041

Assignee: The Texas A&M University System (College Station, TX)

Appl. No.: 17/930545

Filed: September 08, 2022

Prior Publication Data

Document Identifier	Publication Date
US 20230087965 A1	Mar. 23, 2023

Related U.S. Application Data

continuation parent-doc US 16465975 US 11459396 WO PCT/US2017/064186 20171201 child-doc US 17930545

Publication Classification

Int. Cl.: **C07K16/28** (20060101); **A61K38/00** (20060101); **A61K39/00** (20060101); **C07K14/71** (20060101); A61K51/10 (20060101)

U.S. Cl.:

CPC **C07K16/2881** (20130101); **A61K39/00** (20130101); **A61K39/0011** (20130101); **C07K14/71** (20130101); A61K38/00 (20130101); A61K51/1027 (20130101); C07K2317/52 (20130101); C07K2317/569 (20130101); C07K2317/71 (20130101); C07K2317/72 (20130101); C07K2317/76 (20130101); C07K2317/77 (20130101); C07K2317/92 (20130101); C07K2319/30 (20130101)

Field of Classification Search

CPC: C07K (16/2881); C07K (2317/52); C07K (2317/569); C07K (2317/71); C07K (2317/72); C07K (2317/76); C07K (2317/77); C07K (2317/92); C07K (2319/30); A61K (38/00)

References Cited**U.S. PATENT DOCUMENTS**

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
2005/0249723	12/2004	Lazar	N/A	N/A
2006/0235208	12/2005	Lazar et al.	N/A	N/A
2016/0060358	12/2015	Hay	N/A	N/A
2017/0334962	12/2016	Ober et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
0216414	12/2001	WO	N/A
2002044329	12/2001	WO	N/A
2006130834	12/2005	WO	N/A
2012142515	12/2011	WO	N/A
20120150319	12/2011	WO	N/A
20150100299	12/2014	WO	N/A

OTHER PUBLICATIONS

Devanaboyina et al., Nature Communications, May 31, 2017, 6 pages plus 8 supplemental pages; DOI: 10.1038/ncomms15314). cited by applicant

Sun et al., Molecular Therapy vol. 29(3), Mar. 1, 2021, 2 pages plus 6 supplemental pages; (<https://doi.org/10.1016/j.ymthe.2020.11.017>). cited by applicant

Khare et al., (2021) 13(1), 6 pages (<https://doi.org/10.1080/19420862.2021.1976705>). cited by applicant

International Search Report and Written Opinion, PCT/US2017/064186, Mar. 15, 2018. cited by applicant

Battaini, et al., "Antibody response after vaccination with antigen-pulsed dendritic cells", International Journal of VBiological Markers, vol. 19, No. 3, pp. 213-220, (Jul.-Sep. 2004). cited by applicant

Challa et al., "Antigen dynamics govern the induction of CD4+ T cell tolerance during autoimmunity", Journal of Autoimmunity, vol. 72, pp. 84-94, published online: May 25, 2016. cited by applicant

Ha et al., "Immunoglobulin Fe Heterodimer Platform Technology: From Design to Applications in Therapeutic Antibodies and Proteins", Frontiers in Immunology, vol. 7, Art 394, pp. 1-6, publishes; Oct. 6, 2016. cited by applicant

Mann et al., "Transferrin conjugation confers mucosal molecular targeting to a model HIV-1 trimeric gp140 vaccine antigen", (2012), Journal of Controlled Release, vol. 158, pp. 240-249. cited by applicant

Neves et al., "Imaging cell death", (2014) Journal of Nuclear Medicine, vol. 55, No. 1, pp. 1-4. cited by applicant

Saito et al., "Vector-mediated delivery of 125I-labeled b-amyloid peptide Abl-40 through the blood-brain Barrier and binding to Alzheimer disease amyloid of the Abl-40/vector complex", Proc. Natl. Acad. Sci. USA, vol. 92, pp. 10227-10231, (Oct. 1995). cited by applicant

Wang et al., (2016), "Nanobody-derived nano biotechnology tool kits for diverse biomedical and biotechnology applications" International Journal of Nanomedicine, vol. 11, pp. 3287-3303, published Jul. 21, 2016. cited by applicant

Ward et al., (2015), "Targeting FcRn for the modulation of antibody dynamics", Molecular Immunology, vol. 67, pp. 131-141. cited by applicant

Challa et al., "Autoantibody depletion ameliorates disease in murine experimental autoimmune encephalomyelitis" mAbs, Sep./Oct. 2013, pp. 655-659, vol. 5, Issue 5. cited by applicant

Primary Examiner: Landsman; Robert S

Attorney, Agent or Firm: Edell, Shapiro & Finnan, LLC

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application is a continuation of U.S. patent application Ser. No. 16/465,975, filed on May 31, 2019, now U.S. Pat. No. 11,459,396, issued on Oct. 4, 2022, which in turn is a National Stage Entry of International Application PCT/US2017/064186, with an international filing date of Dec. 1, 2017, which claims benefit to U.S. Provisional Patent Application No. 62/429,367, filed on Dec. 2, 2016, the contents of each of which are incorporated by reference herein in their entirety.

SEQUENCE LISTING

(1) The instant application contains a Sequence Listing, which has been submitted electronically in ST.26 (XML) format and is hereby incorporated by reference in its entirety. Said Sequence List, created on Nov. 1, 2022, is named 16810002CON_SEQ_LIST_XML.1 and is 69 kilobytes in size.

TECHNICAL FIELD

(2) This disclosure relates to engineered proteins, and more specifically, to fusion proteins that selectively deplete target antigen-specific antibodies from the body ("Seldegs").

BACKGROUND

(3) Antibodies are Y-shaped proteins present in blood and other body fluids of the human body and the bodies of mammals. Antibodies are a critical component of the body's immune system. They function by recognizing a unique part of a foreign target, called the antigen. An antibody is able to selectively recognize and trigger an immune response to an antigen through its two antigen-binding sites. Each antigen-binding site is at the end of each upper tip of the antibody's Y-shape. The target antigen may bind one or both antigen-binding sites. The base of an antibody's Y-shape is called an

Fc fragment. When an antibody binds to its target, the Fc region can bring about target clearance through antibody effector functions. Such responses can include cellular processes to destroy the antigen. In certain autoimmune diseases and other illnesses, pathogenic antibodies may be created that target self-antigens in the body, contributing to pathogenesis. An antibody may be in either of two physical forms, a soluble form that is secreted from the cell and is free in the blood plasma, or a membrane-bound form that is attached to the outer-membrane of a B cell. The secreted antibodies cause pathology in diseases involving autoreactive antibodies. They can also contribute to transplant rejection or the elimination of protein-based therapeutics.

(4) Due to their ability to bind specifically to target molecules, antibodies can be used to treat diseases such as cancer and autoimmunity. They also have applications in the detection of tumors during whole body imaging using, for example, radiolabeled antibodies in positron emission tomography (PET). However, their relatively long in vivo persistence can lead to high background in non-tumor tissue, resulting in poor contrast for tumor imaging and undesirable off-target effects.

SUMMARY

(5) The present disclosure includes fusion proteins, herein referred to as “Seldegs”, that are configured to allow selective clearance of antigen-specific antibodies. A Seldeg includes a targeting component that is configured to specifically bind to a cell surface receptor or other cell surface molecule, and an antigen component that is configured to specifically bind to an antigen-specific antibody or a variant thereof.

(6) The targeting component of the Seldeg includes a protein or a protein fragment that is configured to specifically bind to a cell surface receptor or other cell surface molecule. The antigen component of the Seldeg includes one molecule of an antigen or antigen fragment or antigen mimetic configured to specifically bind a target antigen-specific antibody. The antigen component is fused directly or indirectly to the targeting component.

(7) The present disclosure also includes a method of depleting a target antigen-specific antibody from a patient by administering to the patient a Seldeg in an amount sufficient to remove at least 50% of the target antigen-specific antibody from the circulation or a target tissue in the patient.

(8) The above Seldegs and methods may further include the following details, which may be combined with one another unless clearly mutually exclusive: i) the targeting component can bind to the cell surface receptor or cell surface molecule with a dissociation constant of less than 10 μM at near-neutral pH; ii) near-neutral pH may be greater than 6.8 and less than 7.5; iii) the Seldeg can comprise at least a first targeting component and a second targeting component, wherein the protein or protein fragment of the first targeting component is configured to bind to a different cell surface receptor or a different cell surface molecule than the protein or protein fragment of the second targeting component iv) the targeting component may include a heterodimer of two immunoglobulin Fc fragments in which one immunoglobulin Fc fragment of the heterodimer is fused to the antigen component and the other immunoglobulin Fc fragment may not be; v) the immunoglobulin Fc fragment may have substantially reduced binding or no detectable binding to Fc gamma receptors; vi) the immunoglobulin Fc fragments can be derived from an immunoglobulin class or isotype that does not bind to Fc gamma receptors or complement; vii) the immunoglobulin Fc fragments can be configured to bind to Fc gamma receptors and complement; viii) at least one of the immunoglobulin Fc fragments can be modified to have a higher binding affinity for FcRn at near-neutral pH than an unmodified immunoglobulin Fc fragment; ix) the antigen component may be fused to one immunoglobulin Fc fragment at an N-terminus or a C-terminus of a hinge-CH.sub.2—CH.sub.3 domain of the immunoglobulin Fc fragment; x) the immunoglobulin Fc fragments may be modified to have no binding affinity for Fc gamma receptors and/or complement (C1q), or lower binding affinity for Fc gamma receptors and/or complement (C1q) than unmodified immunoglobulin Fc fragments; xi) the targeting component may include one or more antibody variable regions or fragments thereof that are configured to specifically bind to the cell surface receptor or the cell surface molecule; xii) the antibody variable region or

fragment thereof may include at least one nanobody; xiii) the nanobody may be a nanobody multimer in which one nanobody is fused to the antigen component and all other nanobodies in the nanobody multimer may not be; xiv) the targeting component may be configured to dissociate from the cell surface receptor or cell surface molecule following entry into an endosome of a complex comprising the Seldeg and the cell surface receptor or cell surface molecule; xv) the antigen component may be fused to an N-terminal location or a C-terminal location on the targeting component; xvi) the antigen component may be fused to a non-terminal location on the targeting component; xvii) the antigen component may be fused to the targeting component via a chemical reaction, through a linker, or during formation of a single combined antigen component-targeting component fusion protein; xviii) the targeting component can be one or more albumin molecules, albumin fragments or mutated albumin variants that are configured to specifically bind to a FcRn; xix) the targeting component can include one or more antibody variable domains or nanobodies that are configured to bind to a transferrin receptor; xx) the targeting component can include one or more protein molecules or protein domains configured to bind to a transferrin receptor; xxi) the targeting component can include one or more protein molecules or protein domains configured to bind to a phosphatidylserine; xxii) the targeting protein component can include one or more antibody variable domains or nanobodies configured to bind to a phosphatidylserine; xxiii) the one or more protein molecules or protein domains can be configured to bind the phosphatidylserine via a calcium-dependent mechanism; xxiv) the targeting component can include a C2A domain of synaptotagmin 1; xxv) the Seldeg can include at least a first antigen component and a second antigen component, wherein the one molecule of the antigen, antigen fragment or antigen mimetic of the first antigen component is different to the one molecule of the antigen molecule, antigen fragment or antigen mimetic of the second antigen component; xxvi) the Seldeg can include at least a first antigen component and a second antigen component, wherein the one molecule of the antigen, antigen fragment or antigen mimetic of the first antigen component is the same as the one molecule of the antigen molecule, antigen fragment or antigen mimetic of the second antigen component; xxvii) the method may include administering an amount sufficient of Seldeg to remove at least 50% of the target antigen-specific antibody from the circulation or the target tissue in the patient within five hours of administration; xxviii) the method may include administering a Seldeg having a targeting component that includes a protein or protein fragment configured to bind to the cell surface receptor or other cell surface molecule with a dissociation constant of less than 10 μ M at near neutral pH; xxix) the amount sufficient of Seldeg administered may be an amount at least equimolar to the amount of target antigen-specific antibody to be depleted; xxx) the method may include administering the Seldeg in an amount sufficient to remove at least 90% of the target antigen-specific antibody from the circulation or target tissue in the patient within two hours of administration; xxxi) the Seldeg may be administered in an amount sufficient to remove at least 50% of the target antigen-specific antibody from the circulation or target tissue in the patient within one hour of administration; xxxii) the Seldeg may be re-administered whenever 50% of patients are expected to have regenerated a threshold amount of target antigen-specific antibody in the circulation or target tissue; xxxiii) the Seldeg may remove less than 10% of non-target antibodies in the circulation or in a tissue targeted by the target antigen-specific antibody; xxxiv) the Seldeg may remove an amount of non-targeted antibodies in the circulation or in the target tissue of the patient that does not cause a clinically adverse effect in the patient; xxxv) the Seldeg may remove less than 1% of non-target antibodies in the circulation or in a tissue targeted by the target antigen-specific antibody; xxxvi) the Seldeg may cause degradation of the target antigen-specific antibody by a cell expressing the cell surface receptor or cell surface molecule; xxxvii) the Seldeg may be administered to a patient with an autoimmune disease and the target antigen-specific antibody may specifically bind to an autoantigen; xxxviii) the Seldeg may be administered to a patient receiving a transplanted organ and the target antigen-specific antibody may specifically bind to an antigen on the transplanted organ; xxxvix) the Seldeg may be administered to increase contrast during tumor

imaging and the target antigen-specific antibody may specifically bind to a tumor antigen; xl) the Seldeg may be administered to a patient who has received a biologic and the target antigen-specific antibody may be the biologic; xli) the Seldeg may be administered to a patient prior to the delivery of a therapeutic agent, if the patient has antibodies specific for the therapeutic agent, if the patient has antibodies specific for the therapeutic agent, and the Seldeg is configured to target the antibodies specific for the therapeutic agent; xlii) the Seldeg may be administered to provide a PET image contrast agent; xliii) the target antigen-specific antibody may be an anti-MOG antibody; xliv) the target antigen-specific antibody may be an anti-HER2 antibody; xlv) the Seldeg may include proteins having amino acid sequences of at least one of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, SEQ ID NO:10, SEQ ID NO:12, SEQ ID NO:14, SEQ ID NO:16, SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO:22, SEQ ID NO:24, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:30, SEQ ID NO:32, or SEQ ID NO:34, or a homolog thereof, xlvi) the Seldeg may include a heterodimer of proteins having amino acid sequences of i) SEQ ID NO: 2 plus SEQ ID NO: 6, ii) SEQ ID NO: 4 plus SEQ ID NO:6, iii) SEQ ID NO:8 plus SEQ ID NO:10, iv) SEQ ID NO:12 plus SEQ ID NO:14, v) SEQ ID NO:14 plus SEQ ID NO:16 plus SEQ ID NO:20, vi) SEQ ID NO: 22 plus SEQ ID NO: 24, vii) SEQ ID NO: 26 plus SEQ ID NO: 28, viii) SEQ ID NO: 30 plus SEQ ID NO: 6, ix) SEQ ID NO: 32 plus SEQ ID NO: 6, or x) SEQ ID NO: 34 plus SEQ ID NO: 6, or homologs thereof.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

- (1) For a more complete understanding of the present invention and its features and advantages, reference is now made to the following description, taken in conjunction with the accompanying drawings, which are not to scale, in which like numerals refer to like features, and in which:
- (2) FIG. 1 is a schematic diagram of selected cellular events that lead to the degradation of antigen-specific antibodies in the presence of Seldegs;
- (3) FIG. 2A is a schematic diagram of a Seldeg including an antigen fused to a N-terminal location of an Fc fragment;
- (4) FIG. 2B is a schematic diagram of a Seldeg including an antigen fused to a C-terminal location of an Fc fragment;
- (5) FIG. 2C is a schematic diagram of a Seldeg including an antigen fused to a non-terminal location of an Fc fragment;
- (6) FIG. 2D is a schematic diagram of a Seldeg including an antigen fused to a terminal location of a protein or protein fragment that binds to a cell surface receptor or cell surface molecule;
- (7) FIG. 2E is a schematic diagram of a Seldeg including an antigen fused to a non-terminal location of a protein or protein fragment that binds to a cell surface receptor or cell surface molecule;
- (8) FIG. 2F is a schematic diagram of a Seldeg including an antigen fused to a N-terminal location of an Fc fragment and protein or protein fragments that bind to a cell surface protein or cell surface receptor to the fused C-termini of the Fc fragment;
- (9) FIG. 2G is a schematic diagram of a Seldeg including an antigen fused to a C-terminal location of an Fc fragment and protein or protein fragments that bind to a cell surface protein or cell surface receptor fused to the N-termini of the Fc fragment;
- (10) FIG. 2H is a schematic diagram of a Seldeg including an antigen fused to a C-terminal location of an antibody that binds to a cell surface protein or cell surface receptor;
- (11) FIG. 2I is a schematic diagram of a Seldeg including an antigen fused to a C-terminal location of an Fc fragment and scFv fragments that bind to a cell surface protein or cell surface receptor fused to the N-termini of the Fc fragment;

(12) FIG. 2J is a schematic diagram of a Seldeg including an antigen fused to a N-terminal location of an Fc fragment and scFv fragments that bind to a cell surface protein or cell surface receptor fused to the C-termini of the Fc fragment;

(13) FIG. 2K is a schematic diagram of a Seldeg including two different antigens fused to the N-terminal locations of an Fc fragment;

(14) FIG. 2L is a schematic diagram of a Seldeg comprising two different antigens fused to the N-terminal locations of an Fc fragment and protein or protein fragments that bind to a cell surface protein or cell surface receptor fused to the C-termini of the Fc fragment;

(15) FIG. 2M is a schematic diagram of a Seldeg comprising two antigen molecules fused to the N-terminal locations of an Fc fragment;

(16) FIG. 2N is a schematic diagram of a Seldeg comprising two antigen molecules fused to the N-terminal locations of an Fc fragment and protein or protein fragments that bind to a cell surface protein or cell surface receptor fused to the C-termini of the Fc fragment;

(17) FIG. 3A is a schematic diagram of two exemplary FcRn-targeting Seldegs, a human epidermal growth factor receptor 2 Seldeg (“HER2-Seldeg”) and a myelin oligodendrocyte glycoprotein Seldeg (“MOG-Seldeg”);

(18) FIG. 3B shows the increased binding of an exemplary FcRn-targeting Seldeg to FcRn at pH 6.0 and 7.4.

(19) FIG. 3C shows HPLC analyses of two exemplary FcRn-targeting Seldegs following incubation at 4° C. (30 days) and 37° C. (5 days) to evaluate their storage stability.

(20) FIG. 3D shows a graph reporting exemplary normalized body counts versus time, showing clearance of an antigen-specific antibody by an exemplary FcRn-targeting Seldeg;

(21) FIG. 3E shows additional graphs reporting exemplary normalized blood and body count versus time, showing clearance of an antigen-specific antibody by an exemplary FcRn-targeting Seldeg;

(22) FIG. 3F shows additional graphs reporting exemplary normalized blood and body count versus time, showing clearance of an antigen-specific antibody by an exemplary FcRn-targeting Seldeg;

(23) FIG. 4A shows in the upper left panel a schematic diagram of an exemplary Seldeg referred to as MOG-Seldeg-PS comprising an antigen fused to targeting protein (C2A domain of synaptotagmin 1, Syt1) that binds to phosphatidylserine (PS); in the upper right panel FIG. 4A shows exemplary SDS-PAGE gels (reducing and non-reducing conditions) of recombinant proteins of MOG-Seldeg-PS, MOG-Seldeg-PS(DN) with mutations that substantially reduce binding to PS, and Fc-Syt1 that has no antigen (MOG) attached; in the lower panel, FIG. 4A shows exemplary HPLC profiles of the recombinant proteins that are shown in the upper right FIG. 4A, MOG-Seldeg-PS, MOG-Seldeg-PS(DN) and Fc-Syt1;

(24) FIG. 4B shows additional graphs reporting exemplary normalized blood and body count versus time, showing clearance of an antigen-specific antibody by an exemplary PS-targeting Seldeg;

(25) FIG. 5A shows graphs reporting exemplary data showing the accumulation of antigen-specific antibodies in cells in the presence of exemplary FcRn-targeting Seldegs and control proteins;

(26) FIG. 5B is an exemplary series of microscopic images of an exemplary FcRn-targeting Seldeg and control protein in the presence of target antigen-specific antibodies, with microscopic images of representative endosomes cropped, expanded, and presented in the top right-corner insets;

(27) FIG. 5C is another series of exemplary microscopic images of an exemplary FcRn-targeting Seldeg and control protein in the presence of target antigen-specific antibodies, with microscopic images of representative endosomes cropped, expanded, and presented in the top right-corner insets;

(28) FIG. 6A is another series of exemplary microscopic images of an exemplary FcRn-targeting Seldeg and control protein in the presence of target antigen-specific antibodies, with microscopic images of representative lysosomes cropped, expanded, and presented in the top right-corner insets;

(29) FIG. 6B is another series of exemplary microscopic images of exemplary FcRn-targeting Seldegs in the presence of antigen-specific antibodies that do not recognize the antigen that is being targeted by the Seldeg, with microscopic images of representative lysosomes cropped, expanded, and presented in the top right-corner insets;

(30) FIG. 7 shows graphs reporting exemplary data showing the accumulation of antigen-specific antibodies in cells in the presence of exemplary PS-targeting Seldegs and control proteins;

(31) FIG. 8 is a schematic diagram of an exemplary Seldeg including an antigen fused to targeting protein (antibody) that binds to the transferrin receptor (TfR); HPLC profiles of the recombinant proteins are shown, including an analysis of the targeting protein (antibody) without antigen (MOG) attached;

(32) FIG. 9 is a graph reporting exemplary data showing the accumulation of antigen-specific antibodies in cells in the presence of an exemplary TfR-targeting Seldeg and control protein;

(33) FIG. 10A is an exemplary series of positron emission tomography (PET) analyses of tumors in mice following delivery of radiolabeled HER2-specific antibody and treatment with an exemplary FcRn-targeting Seldeg, control protein or vehicle control.

(34) FIG. 10B shows a graph reporting contrast measures for tumor:thoracic regions of tumor-bearing mice following delivery of radiolabeled HER2-specific antibody and treatment with an exemplary FcRn-targeting Seldeg, control protein or vehicle control.

DETAILED DESCRIPTION

(35) This disclosure relates to engineered proteins, and more specifically, to Seldegs, which are fusion proteins that are configured to selectively target antigen-specific antibodies for depletion from the body. Seldegs cause the selective degradation of the targeted antigen-specific antibodies by binding to the antigen-specific antibodies and directing them to late endosomes or lysosomes, which contain degradative enzymes. A Seldeg is a fusion protein or molecule that includes at least a targeting component and an antigen component. The targeting component includes a protein or protein fragment or other molecule that is configured to bind to a cell surface receptor or other cell surface molecule. The antigen component includes one molecule of an antigen, antigen fragment or antigen mimetic that is recognized by the targeted antigen-specific antibody.

(36) Upon binding of the antigen-specific antibody to the antigen component, a complex is formed comprising the Seldeg and the antigen-specific antibody. The complex is also configured to bind to the cell surface receptor or other cell surface molecule, allowing cellular internalization of a complex that includes the Seldeg, the antigen-specific antibody, and the targeted cell surface receptor or other cell surface molecule (see FIG. 1). The targeted cell surface receptor or cell surface molecule may dissociate from the complex upon entry into the endosomes, due to acidic pH, low calcium concentrations and/or other conditions that distinguish the endosomal environment from the extracellular environment. Internalization into endosomes and lysosomal entry results in the selective degradation of the complex.

(37) The term “antigen-specific antibody” as used herein refers to an antibody or antibody that binds to a particular antigen, antigen fragment or antigen mimetic.

(38) The term “antigen fragment” as used herein refers to a part of the antigen that can be recognized by the antigen-specific antibody.

(39) The term “antigen mimetic” as used herein refers to a protein, protein fragment, peptide or other molecule that has the same overall shape and properties as the part of the antigen that is recognized by an antigen-specific antibody.

(40) The term “cell surface molecule” as used herein refers to a protein or other biological molecule (e.g. phospholipid, carbohydrate) that is exposed on the plasma membrane of a cell.

(41) Seldegs may include an antigen fused to an Fc fragment of an IgG antibody (herein also referred to as “immunoglobulin Fc fragment”), an FcRn-specific nanobody-antigen fusion molecule, an FcRn-specific antibody that binds to FcRn through its variable region fused to an antigen, an albumin-antigen fusion protein, a PS-binding protein, a TfR-specific antibody or other

protein, protein fragment or other molecule that is configured to bind to a cell surface receptor or other cell surface molecule identifiable by skilled persons upon reading of the present disclosure. (42) Examples of Seldegs described herein include targeting components that are configured to bind to cell surface molecules such as human FcRn, exposed phosphatidylserine (PS) or the transferrin receptor (TfR) with affinities (dissociation constants) of less than 10 μ M at near neutral pH.

(43) FcRn and TfR are proteins, and PS is a phospholipid, that may be found on the surface and within multiple different cell types within the body. This invention is not limited to targeting these receptors or cell surface molecules, and many other targets could be envisaged such as the low density lipoprotein receptor, high density lipoprotein receptor, asialoglycoprotein receptor, inhibitory Fc gamma receptors, T cell receptor, B cell receptor, G-protein coupled receptors, insulin receptor, glucagon receptors, galactose receptors, mannose receptors, VEGF receptors, mannose receptors among others identifiable by those skilled in the art. Other targets can be identified in, for example, the following publications or databases: Cell surface receptor protein atlas (Bausch-Fluck, D., Hofmann, A., Bock, T., Frei, A. P., Cerciello, F., Jacobs, A., Moest, H., Omasits, U., Gundry, R. L., Yoon, C., Schiess, R., Schmidt, A., Mirkowska, P., Härtlova, A., Van Eyk, J. E., Bourquin, J-P., Aebersold, R., Boheler, K. R., Zandstra, P., Wollscheid, B. (2015) A mass spectrometric-derived cell surface protein atlas. PLoS One 10: e0121314), and the Human protein atlas (<https://www.proteinatlas.org/humanproteome/secretome>).

(44) The targeting component can bind the cell surface receptor or other cell surface molecule with an affinity (dissociation constant) of less than 10 μ M at near-neutral pH.

(45) Accordingly, the targeting component of a Seldeg can include any type of molecule that is configured to specifically bind to a cell surface receptor or other cell surface molecule. Such molecules can include proteins, protein fragments, polynucleotides such as ribonucleic acids or deoxyribonucleic acids, polypeptides, polysaccharides, lipids, amino acids, peptides, sugars and/or other small or large molecules and/or polymers identifiable by skilled persons upon reading of the present disclosure. For example, the targeting component of a Seldeg can include a polynucleotide that is a ligand for a cellular receptor.

(46) The Seldeg can comprise at least a first targeting component and a second targeting component, wherein the protein or protein fragment of the first targeting component is configured to bind to a different cell surface receptor or a different cell surface molecule than the protein or protein fragment of the second targeting component.

(47) As shown in FIG. 1, a Seldeg **20** may reversibly bind to cell surface receptor or other molecule **30** on the surface of a cell **10**. A target antigen-specific antibody **40**, present in extracellular space **50**, may reversibly bind to Seldeg **20**. This binding typically occurs at near-neutral pH, such as at a pH greater than 6.8 and less than 7.5, because that is the typical pH of extracellular space **50**. Non-target antibodies **60** do not bind to Seldeg **20**, or bind with such low affinities that any binding is non-specific. Cell surface receptor or molecule **30** with attached Seldeg **20** and target antigen-specific antibody **40** are internalized into the cell **10** through receptor-mediated uptake into endosome **70** via pathway A. The receptor or other molecule can be recycled back to the surface of the cell. Accordingly, on pathway B, receptor or molecule **30** with attached Seldeg **20** and target antigen-specific antibody **40** are cycled back to the surface of cell **10**. Target antigen-specific antibody **40** may then be released and may reattach from Seldeg **20** to the same or another Seldeg, or it may remain attached. Similarly Seldeg **20** may then be released from receptor or molecule **30** and may reattach to the same or another cell surface receptor or molecule. For some Seldegs, the complex of Seldeg **20** and antibody **40** may dissociate from the receptor or other molecule **30** in the early or late endosome due to the acidic pH or low Ca^{2+} concentrations in this compartment (pathway C). Accordingly, in pathway D, receptor or cell surface molecule may recycle back to the cell surface whereas the Seldeg **20** bound to antibody **40** enters the lysosomes and is degraded into fragments **80** in pathway E. For some Seldegs, at some point following internalization into cell **10**,

receptor or molecule **30** with attached Seldeg **20** and target antigen-specific antibody **40** enters the late endosomal/lysosomal in pathway F, in which at least target antigen-specific antibody **40** is degraded into fragments **80**. This entry into lysosomes is expected to be increased if Seldeg **20** cross-links receptor or molecule **30** into dimers or higher order aggregates.

(48) Through this mechanism of selective depletion, Seldegs target and selectively deplete antigen-specific antibodies from the body without adversely affecting the levels of antibodies of non-targeted specificities.

(49) In particular, Seldegs as described herein can target and selectively deplete antigen-specific antibodies from the body without having an adverse clinical effect in the patient due to the depletion of antibodies of non-targeted specificities. Such adverse clinical effects include, for example, immunosuppression, and symptoms thereof, such as pinkeye, bronchitis, ear infections, sinus infections, cold, diarrhea, pneumonia, yeast infection, meningitis, skin infections, and other opportunistic infections, particularly opportunistic infection normally controlled through anti-body mediated immune responses; and blood disorders, such as low platelet counts or anemia, and hypogammaglobulinemia and symptoms thereof, such as abdominal pain, bloating, nausea, vomiting, diarrhea, or weight loss.

(50) In general, Seldegs according to this disclosure are configured to specifically bind a cell surface receptor/molecule at near-neutral pH via a targeting component and also specifically bind to an antigen-specific antibody at near-neutral pH via an antigen component that is fused directly or indirectly to the targeting component. The term “specifically bind” as used herein refers to a detectable selective inter-molecular interaction between the targeting component and the cell surface receptor/molecule, or between the antigen component and the antigen-specific antibody. For example, to specifically bind, the antigen needs to show a detectable interaction with the antibodies that are being targeted, whilst not showing a detectable interaction with other antibodies. Techniques for detecting specific binding are known within the art, such as ELISA, surface plasmon resonance analysis and other methods identifiable by skilled persons

(51) Accordingly, Seldegs allow at least a portion of the antigen-specific antibody in the circulation of a patient to be internalized into cells that express the targeted cell surface receptor or targeted other cell surface molecule and thereafter intracellularly degraded.

(52) Seldegs according to this disclosure may avoid immune responses by containing one copy, otherwise referred herein as “one molecule”, of each type of antigen, antigen fragment or antigen mimetic, per Seldeg, which in combination with the insertion of mutations to reduce or eliminate Fc gamma receptor binding and/or complement binding, is expected to decrease antibody cross-linking and the formation of potentially inflammatory immune complexes. In particular, at least 99% of Seldegs, at least 99.5% of Seldegs, or at least 99.9% of Seldegs may contain only one copy of antigen, antigen fragment, or antigen mimetic per Seldeg at near-neutral pH. Other Seldegs according to the present disclosure can contain more than one molecule of an antigen, antigen fragment, or antigen mimetic. The bivalent nature of the antibodies that are bound by Seldegs may result in complexes of two Seldeg molecules per antibody, which through target receptor dimerization is expected to increase the efficiency of lysosomal delivery of Seldeg-antibody complexes.

(53) Seldegs can include at least a first antigen component and a second antigen component, wherein the one molecule of the antigen, antigen fragment or antigen mimetic of the first antigen component is different to the one molecule of the antigen molecule, antigen fragment or antigen mimetic of the second antigen component. Accordingly, Seldegs comprising at least a first antigen component and a second antigen component allow clearance of antigen-specific antibodies of more than one specificity.

(54) Seldegs can include at least a first antigen component and a second antigen component, wherein the one molecule of the antigen, antigen fragment or antigen mimetic of the first antigen component is the same as the one molecule of the antigen molecule, antigen fragment or antigen

mimetic of the second antigen component.

(55) Accordingly, Seldegs can include, for example, one or more antigen components fused to a C-terminus and/or an N-terminus of a targeting component, wherein the one molecule of the antigen mimetic antigen components of the respective antigen components of the respective antigen components can be the same or different.

(56) In addition, Seldegs may contain human or humanized proteins or protein fragments to avoid or decrease the possibility of an immune reaction to the Seldeg when administered to a human. The antigen, antigen fragment, or antigen mimetic of the antigen component is preferably a human protein or protein fragment for administration of the Seldeg to a human. The targeting component is also preferably a human protein or protein fragment, such as a human antibody fragment or human albumin or albumin fragment, or a humanized antibody or humanized antibody fragment for administration of the Seldeg to a human. If a Seldeg is developed for use in a non-human animal, then proteins or protein fragments derived from or engineered to be immunologically compatible with that animal may be used instead.

(57) FIG. 2A is a schematic representation of the activity of a Seldeg **20a** including antigen **100** fused to a targeting component having the Fc fragment of IgG **110**. As understood by persons skilled in the art, the Fc fragment of an IgG is all of the lower base of the antibody's Y-shape, which is the sulfhydryl-bridged hinge region and the CH2 and CH3 domains. Seldegs can have an Fc fragment that does not have the hinge region, or the hinge region does not have sulfhydryl bridges. Fc fragment **110** allows Seldeg **20a** to bind an FcRn molecule on an FcRn-expressing cell. In the example shown in FIG. 2A, antigen **100** may be fused to Fc fragment **110a** at the N-terminus of the hinge-CH.sub.2—CH.sub.3 **120**. When antigen **100** is fused to Fc fragment **110a** and the resulting antigen-Fc fragment dimerizes with another Fc fragment **110b** lacking an antigen, using the knobs-into-holes strategy (for example, as described in Moore, G. L., Bautista, C., Pong, E., Nguyen, D. H., Jacinto, J., Eivazi, A., Muchhal, U. S., Karki, S., Chu, S. Y., Lazar, G. A., 2011). A novel bispecific antibody format enables simultaneous bivalent and monovalent co-engagement of distinct target antigens. MAb 3, 546-557), a heterodimeric Seldeg molecule **20a** as shown is produced. Seldeg **20a** has an Fc fragment with a monomeric display of antigen **100**, which avoids the formation of multimeric immune complexes that can cause inflammation and other adverse effects. Although Seldegs containing only antigen **100** fused to Fc fragment **110a** may be produced and used in some circumstances, due to the tendency of Fc fragment to dimerize, a dimer will typically be produced. In order to avoid Fc fragment dimers in which both Fc fragments **110a** have a fused antigen **100**, which can lead to the formation of multimeric immune complexes, Seldegs are designed with knobs-into-holes mutations and/or electrostatic steering mutations (for example, as described in Gunasekaran, K., Pentony, M., Shen, M., Garrett, L., Forte, C., Woodward, A., Ng, S. B., Born, T., Retter, M., Manchulenko, K., Sweet, H., Foltz, I. N., Wittekind, M., Yan, W. (2010) Enhancing antibody Fc heterodimer formation through electrostatic steering effects: applications to bispecific molecules and monovalent IgG. J Biol Chem 285, 19637-19646) to promote heterodimer formation, so there is only one Fc with one antigen fused. Other approaches can also be used to generate heterodimers, such as the insertion of a (G.sub.4S).sub.13 linker peptide between the C-terminus of the antigen-Fc fusion and N-terminus of a second Fc fragment (for example, as described in Zhou, L., Wang, H-Y., Tong, S., Okamoto, C. T., Shen, W-C., Zaro, J. L. (2016) Single chain Fc-dimer-human growth hormone fusion protein for improved drug delivery. Biomaterials, 117, 24-31]. DNA and protein sequences of several examples of Seldegs comprising knobs-into-holes mutations, electrostatic steering mutations, and/or arginine mutations or other mutations that reduce Fc gamma receptor and complement binding are described in Example 10.

(58) Additional examples of knobs-into-holes mutations include Y349T/T394F: S364H/F405A and Y349T/F405F: S364H/T394F (for example as described in Moore, G. L., Bautista, C., Pong, E., Nguyen, D. H., Jacinto, J., Eivazi, A., Muchhal, U. S., Karki, S., Chu, S. Y., Lazar, G. A. (2011) A novel bispecific antibody format enables simultaneous bivalent and monovalent co-engagement of

distinct target antigens. MAbs 3, 546-557) and T366S:L368A/Y407V (for example as described in Atwell, S., Ridgway, J. B. B., Wells, J. A., Carter, P (1997) Stable heterodimers from remodeling the domain interface of a homodimer using a phage display library. J. Mol. Biol., 270, 26-35) among others identifiable by persons skilled in the art. The residue numbering of these exemplary knobs into holes mutations refers to the EU antibody numbering system, as would be understood by persons skilled in the art.

(59) Additional examples of electrostatic steering mutations include E356K/D399K:K392D/K409D and K409D/K370D:D357K/D399K (for example as described in Gunasekaran, K., Pentony, M., Shen, M., Garrett, L., Forte, C., Woodward, A., Ng, S. B., Born, T., Retter, M., Manchulenko, K., Sweet, H., Foltz, I. N., Wittekind, M., Yan, W. (2010). Enhancing antibody Fc heterodimer formation through electrostatic steering effects: applications to bispecific molecules and monovalent IgG. J Biol Chem 285, 19637-19646.) among others identifiable by persons skilled in the art. The residue numbering of these exemplary electrostatic steering mutations refers to the EU antibody numbering system, as would be understood by persons skilled in the art.

(60) Additional examples of arginine mutations or other mutations to reduce binding to Fc gamma receptors and complement (C1q) include G236R/L328R (for example as described in Horton, H. M., Bernett, M. J., Pong, E., Peipp, M., Karki, S., Chu, S. Y., Richards, J. O., Vostiar, I., Joyce, P. F., Repp, R., Desjarlais, J. R., Zhukosky, E. (2010) Potent in vitro and in vivo activity of an Fc-engineered anti-CD19 monoclonal antibody against lymphoma and leukemia. Cancer Res., 68, 8049-8057; Moore, G. L., Bautista, C., Pong, E., Nguyen, D. H., Jacinto, J., Eivazi, A., Muchhal, U. S., Karki, S., Chu, S. Y., Lazar, G. A. (2011). A novel bispecific antibody format enables simultaneous bivalent and monovalent co-engagement of distinct target antigens. MAbs 3, 546-557), N297A or N297Q (for example as described in Tao, M-H., Morrison, S. L. (1989) Studies of aglycosylated chimeric mouse-human IgG: role of carbohydrate in the structure and effector functions mediated by the human IgG constant region. J. Immunol., 143, 2595-2601; Lux, A., Yu, X., Scanlan, C. N., Nimmerjahn, F. (2013) Impact of immune complex size and glycosylation on IgG binding to human FcγRs. J. Immunol., 190, 4315-4323), D265A (for example as described in Lux, A., Yu, X., Scanlan, C. N., Nimmerjahn, F. (2013) Impact of immune complex size and glycosylation on IgG binding to human FcγRs. J. Immunol., 190, 4315-4323; Clynes, R. A., Towers, T. L., Presta, L. G., Ravetch, J. V. (2000) Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. Nat. Med. 6, 443-446), L234A/L235A (for example as described in Wines, B. D., Powell, M. S., Parren, P. W. H. I., Barnes, N., Hogarth, P. M. (2000) The IgG Fc contains distinct Fc receptor (FcR) binding sites: the leukocyte receptors FcγRI and FcγRIIa bind to a region in the Fc distinct from that recognized by neonatal FcR and protein A. J. Immunol., 164, 5313-5318), and L234A/L235A/P329G (for example as described in Schlothauer, T., Herter, S., Koller, C. F., Grau-Richards, S., Steinhart, V., Spick, C., Kubbies, M., Klein, C., Umana, P., Mossner, E. (2016) Novel human IgG1 and IgG4 Fc-engineered antibodies with completely abolished effector functions. PEDS, 29, 457-466), among others identifiable by persons skilled in the art. The residue numbering of these exemplary arginine mutations or other mutations to reduce binding to Fc gamma receptors and complement (C1q) refers to the EU antibody numbering system, as would be understood by persons skilled in the art.

(61) Other mutations to ablate FcγR and/or complement binding that target residues at, or in proximity to, the location of the FcγR and complement binding sites can be used. These sites on the Fc region of IgG have been localized (for example, as described in Jefferis, R., Lund, J. (2002) Interaction sites on human IgG-Fc for FcγR: current models. Immunol. Letts., 82, 57-65; Duncan, A. R., Winter, G. (1988) The binding site for C1q on IgG. Nature, 332, 738-740; Idusogie, E. E., Presta, L. G., Gazzano-Santoro, H., Totpal, K., Wong, P. Y., Ultsch, M., Meng, G., Mulkerrin, M. G. (2000) Mapping of the C1q binding site on rituxan, a chimeric human antibody with a human IgG1 Fc. J. Immunol., 164, 4178-4184; Hogarth, P. M., Anania, J., Wines, B. D. (2014) The FcγR of humans and non-human primates and their interaction with IgG: implications for induction of

inflammation, resistance to infection and the use of therapeutic monoclonal antibodies. Curr. Top. Microbiol. Immunol., 382, 321-352).

(62) Seldegs may include Fc fragments derived from immunoglobulin classes or isotypes that do not bind, or have very weak binding, to Fc gamma receptors or complement such as human IgG2 or human IgG4.

(63) For some applications such as diagnostic imaging, Seldegs may include Fc fragments with binding sites for Fc gamma receptors and/or complement to increase inflammatory responses against the antigen that is present in the Seldeg.

(64) Fc fragment **110** may be modified to substantially increase its binding affinity for FcRn at near-neutral pH as compared to unmodified Fc fragments. For example, the dissociation constant between Fc fragment **110** and FcRn at a pH greater than 6.8 and less than 7.5 may be less than 10 μ M as determined by surface plasmon resonance or other biophysical method. However, Fc fragment **110** may have a similar or increased affinity for FcRn as compared to an unmodified Fc fragment at acidic endosomal pH (about 6.0), or it may be modified to have a much lower or negligible binding affinity for FcRn at endosomal pH as compared to an unmodified Fc fragment. This increase in binding affinity at near neutral pH allows each Seldeg to cause its bound target antigen-specific antibody to be efficiently internalized and trafficked into late endosomes or lysosomes in FcRn-expressing cells. Enhanced binding affinity of the Fc fragment for FcRn may be achieved by insertion of mutations. Naturally-occurring IgGs have a substantially higher binding affinity for FcRn at acidic pH levels as opposed to near-neutral pH. This property is essential for the recycling and transport of IgG within FcRn-expressing cells. In contrast, an increase in binding affinity for FcRn at pH 7.4, for example, results in receptor-mediated internalization into cells and lysosomal delivery.

(65) Fc fragment **110** may also be modified to eliminate or substantially reduce the binding affinity for Fc gamma receptors and complement (C1q). This modification prevents inflammatory responses caused by the formation of multimeric immune complexes. For example, as described in Example 10, the Fc regions can be mutated (G236R/L328R; EU numbering) (for example as described in Moore, G. L., Bautista, C., Pong, E., Nguyen, D. H., Jacinto, J., Eivazi, A., Muchhal, U. S., Karki, S., Chu, S. Y., Lazar, G. A. (2011). A novel bispecific antibody format enables simultaneous bivalent and monovalent co-engagement of distinct target antigens. MAbs 3, 546-557) (EU numbering), herein also referred to as "arginine mutations", so that they do not bind Fc gamma receptors. In Example 10, these mutations correspond to residues 22 and 114 of Fc-Syt1 (see SEQ ID NO:10), and residues 114 and 236 of MOG-Seldeg-PS (see SEQ ID NO: 8). Other examples of mutations that substantially reduce or ablate binding to Fc gamma receptors and complement include N297A or N297Q) (EU numbering; for example as described in Tao, M-H., Morrison, S. L. (1989) Studies of aglycosylated chimeric mouse-human IgG: role of carbohydrate in the structure and effector functions mediated by the human IgG constant region. J. Immunol., 143, 2595-2601; Lux, A., Yu, X., Scanlan, C. N., Nimmerjahn, F. (2013) Impact of immune complex size and glycosylation on IgG binding to human Fc γ Rs. J. Immunol., 190, 4315-4323), D265A (for example as described in Lux, A., Yu, X., Scanlan, C. N., Nimmerjahn, F. (2013) Impact of immune complex size and glycosylation on IgG binding to human Fc γ Rs. J. Immunol., 190, 4315-4323; Clynes, R. A., Towers, T. L., Presta, L. G., Ravetch, J. V. (2000) Inhibitory Fc receptors modulate in vivo cytotoxicity against tumor targets. Nat. Med. 6, 443-446), L234A/L235A (for example as described in Wines, B. D., Powell, M. S., Parren, P. W. H. I., Barnes, N., Hogarth, P. M. (2000) The IgG Fc contains distinct Fc receptor (FcR) binding sites: the leukocyte receptors Fc γ RI and Fc γ RIIa bind to a region in the Fc distinct from that recognized by neonatal FcR and protein A. J. Immunol., 164, 5313-5318), L234A/L235A/P329G (for example as described in Schlothauer, T., Herter, S., Koller, C. F., Grau-Richards, S., Steinhart, V., Spick, C., Kubbies, M., Klein, C., Umana, P., Mossner, E. (2016) Novel human IgG1 and IgG4 Fc-engineered antibodies with completely abolished effector functions. PEDS, 29, 457-466), (EU numbering) among others identifiable by

persons skilled in the art. A reduction in binding affinity for Fc gamma receptors of at least 10-fold is preferred.

(66) As shown in FIG. 2B, in Seldeg **20b** antigen **100** may be attached to Fc fragment **110a** at another terminal location, or as shown in FIG. 2C, in Seldeg **20c**, antigen **100** may be attached at a non-terminal location. Any location that does not prevent specific FcRn binding is suitable. Such locations include amino acid residues that are sufficiently distant from the FcRn interaction site (encompassing residues 252-256, 309-311, 433-436 at the CH2-CH3 domain interface; EU numbering so as not to either directly or sterically block FcRn binding, as would be identifiable by skilled persons.

(67) Antigen **100** may be fused to Fc fragment **110** in any suitable manner, including attachment via a chemical reaction, attachment through a linker, or during formation of a single combined antigen-Fc fragment protein. Examples of chemical coupling that could be used are: amine-to-amine (NHS esters), sulfhydryl-to-sulfhydryl (maleimide), amine-to-sulfhydryl (NHS ester/maleimide), sulfhydryl-to-carbohydrate (maleimide/hydrazide), or attachment to via an unnatural amino acid with the desired chemical reactivity. This unnatural amino acid can be inserted during recombinant production of the Fc fragment and/or antigen. Polyethyleneglycol (PEG) spacers can also be inserted between the chemically conjugated proteins, protein fragments or other molecules. Possible linkers include repeats of glycine-serine linker peptides, or other more rigid linker peptides, that are encoded in the recombinant expression plasmid for the antigen-Fc fusion. Linkage chemistry, sites of linkage and choice of peptide can be guided by molecular modeling, and can be designed to minimize loss of binding activity of the antigen or the protein/protein fragment targeting the cell surface molecule, as would be understood by skilled persons.

(68) FIG. 2D is a schematic representation of Seldeg **20d** in which antigen **100** is attached to antibody variable region **130**. Antibody variable region **130** specifically binds to a cell surface receptor or cell surface molecule. Antibody variable region **130** may be an entire variable region or a fragment thereof, so long as it can specifically bind to a cell surface receptor or cell surface molecule. Antibody variable region **130** may include portions of a non-variable region of an antibody that is configured to bind to a cell surface receptor or cell surface molecule. For example, antibody variable region **130** may be a single-domain antibody (sdAb) or camelid-derived VHH domain (also commonly referred to as a nanobody). Such variable regions have the overall fold of an immunoglobulin domain, comprising two anti-parallel β -sheets, and can also include domains from other members of the immunoglobulin superfamily such as T cell receptor variable domains, constant region domains of antibodies or domains of the coreceptor, CD4, among others identifiable by persons skilled in the art. Antibody variable region **130** may be present as a monomer as shown in FIG. 2D, or as a multimer. For example, if antibody variable region **130** is present as a nanobody, it may be engineered with a linker peptide such as GSSGSGGGGS (SEQ ID NO:35) between the C-terminus of the first nanobody and the N-terminus of the second nanobody to form a dimer, resulting in increased binding avidity for target receptor/molecule. If antibody variable region **130** is a nanobody or another protein that is engineered to form multimers, variants without antigen **100** may be included during Seldeg formation so that multimers contain only one copy of antigen **100**, just as described above for Seldegs containing Fc fragments. Antibody variable regions can also include heterodimers of heavy chain variable (VH) domains linked by peptide linkers to light chain variable (VL) domains to form scFv fragments. The linker sequences that are used to link VH and VL domains are well known to those with skill in the art and include the GGGSGGGSGGGGS [(G.sub.4S).sub.3](SEQ ID NO:36) sequence that connects the C-terminus of the VH domain to the N-terminus of the VL domain. In some embodiments, the C-terminus of the VL domain can be connected to the N-terminus of the VH domain with similar linker sequences. ScFvs that bind to a cell surface receptor or other cell surface molecule can be isolated from libraries of scFvs using phage display, yeast display or other antibody display approaches. The targeting protein component of a Seldeg could also include Fab

fragments of an antibody that can be isolated from libraries of Fab fragments using phage display, yeast display etc. For nanobodies, scFvs and Fab fragments, affinities for binding to a cell surface receptor or cell surface molecule can be increased by randomly mutating residues in the complementarity determining regions (CDRs), or by using error-prone polymerase chain reaction, to generate libraries of mutated nanobodies or variable domains. Exemplary CDR residues that would be targeted are those in CDR3 of the light chain variable domain (residues 89-97; Kabat numbering) and CDR3 of the heavy chain variable domain (residues 95-102; Kabat numbering). These libraries can be displayed on phage or yeast and higher affinity variants selected using approaches known to those with skill in the art.

(69) Although FIG. 2D illustrates antigen **100** at a terminal location of antibody variable region **130**, it may instead be located at a non-terminal location. Antigen **100** may be fused to antibody variable region **110** in any suitable manner, including attachment via a chemical reaction, attachment through a linker, or during formation of a single combined antigen-antibody variable region fusion protein.

(70) A Seldeg may also contain an antigen component fused to targeting component that includes a protein other than an antibody or antibody fragment, providing that this protein is configured to bind to a cell surface receptor or other cell surface molecule. For example, as shown in FIG. 2E, Seldeg **20e** includes antigen **100** fused to albumin or an albumin fragment **140** able to bind FcRn. The albumin or albumin fragment may be mutated or modified so that it binds with increased affinity to FcRn. For example, mutations can be inserted into the FcRn binding domain (DIII) of (human serum) albumin using error prone PCR followed by display of libraries of mutated albumin variants on yeast or phage, and selection of higher affinity variants. Alternatively, higher affinity variants can be generated by mutating residues at or near the albumin:FcRn interface and either selecting or screening for albumin variants with increased binding affinity. Although FIG. 2E illustrates antigen **100** at a non-terminal location of albumin or albumin fragment **140**, it may instead be located at a terminal location. Antigen **100** may be fused to albumin or albumin fragment **140** in any suitable manner, including attachment via a chemical reaction, attachment through a linker, or during formation of a single combined antigen-FcRn-binding protein.

(71) FIG. 2F is a schematic representation of exemplary Seldeg **20f** including antigen **100** attached to the N-terminus of an Fc fragment **150**. In the example shown in FIG. 2F, protein or protein fragments **160** that bind to a cell surface receptor or cell surface molecule are attached to the C-terminus of Fc fragment **150a**. For example, the protein or protein fragment may be the C2A domain of synaptotagmin that binds to phosphatidylserine (PS). Fc fragment **150** can be engineered to bind to FcRn with increased affinity and may be mutated so that it binds to Fc gamma receptors and complement with very low or no detectable binding affinity. In order to avoid Fc fragment homodimers having two Fc fragments **150a** with fused antigen **100**, which can lead to the formation of multimeric immune complexes, Seldegs as shown in FIG. 2F are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is only one Fc with one Fc-antigen. In FIG. 2F, both Fc fragments **150a** and **150b** have proteins or protein fragments that bind to the cell surface protein or other cell surface molecule fused to them; alternatively, only one such protein or protein fragment may be present. In the exemplary Seldeg shown in FIG. 2G, both the antigen **100** and protein or protein fragments **160** that bind to a cell surface receptor or cell surface molecule are fused to the C- and N-termini of the Fc fragments **150a** and **150b**, respectively, to generate Seldeg **20g**.

(72) FIG. 2H is a schematic representation of exemplary Seldeg **20h** including antigen **100** attached to the C-terminus of an antibody **170** that binds to a cell surface protein or cell surface molecule. The Fc fragment (Fc) in the antibody can be engineered to bind to FcRn with increased affinity and may be mutated so that it binds to Fc gamma receptors and complement with very low or no detectable binding affinity. In order to avoid antibody homodimers in which both Fc fragments have a fused antigen **100**, which can lead to the formation of multimeric immune complexes,

Seldegs as shown in FIG. 2H are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is only one antibody heavy chain per antibody molecule with attached antigen **100**. Both Fab fragments of the antibody may bind to the same cell surface protein or other cell surface molecule; alternatively, they could bind to two or more different cell surface proteins or molecules.

(73) FIG. 2I is a schematic representation of exemplary Seldeg **20i** comprising antigen **100** attached to the C-terminus of a scFv (**180**)-Fc fusion that binds to a cell surface protein or cell surface molecule. The Fc fragment (Fc) in the antibody can be engineered to bind to FcRn with increased affinity and may be mutated so that it binds to Fc gamma receptors and complement with very low or no detectable binding affinity. In order to avoid antibody homodimers in which both Fc fragments have a fused antigen **100**, which can lead to the formation of multimeric immune complexes, Seldegs as shown in FIG. 2I are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is only one antibody heavy chain-scFv fusion per molecule with attached antigen **100**. Both scFv fragments of the antibody may bind to the same cell surface protein or other cell surface molecule; alternatively, they could bind to two or more different cell surface proteins or molecules.

(74) FIG. 2J is a schematic representation of exemplary Seldeg **20j** comprising antigen **100** attached to the N-terminus of an Fc-scFv (**180**) fusion that binds to a cell surface protein or cell surface molecule. The Fc fragment (Fc) in the antibody can be engineered to bind to FcRn with increased affinity and may be mutated so that it binds to Fc gamma receptors and complement with very low or no detectable binding affinity. In order to avoid antibody homodimers in which both Fc fragments have a fused antigen **100**, which can lead to the formation of multimeric immune complexes, Seldegs as shown in FIG. 2I are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is only one antibody heavy chain-scFv fusion per molecule with attached antigen **100**. Both scFv fragments of the antibody may bind to the same cell surface protein or other cell surface molecule; alternatively, they could bind to two or more different cell surface proteins or molecules.

(75) As shown in FIG. 2K, in exemplary Seldeg **20k** two antigen components (**100**, **190**) may be attached to a targeting component, for example Fc fragment **110a** and Fc fragment **110b** at an N-terminal or other location to generate a Seldeg that can clear antigen-specific antibodies of different specificities. Seldegs as shown in FIG. 2K are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is an antigen molecule of each type (**100**, **190**) in each Seldeg molecule.

(76) FIG. 2L is a schematic representation of exemplary Seldeg **20l** comprising antigen **100** and antigen **190** attached to the N-termini of an Fc fragment **150**. In the exemplary embodiment shown in FIG. 2L, protein or protein fragments **160** that bind to a cell surface receptor or cell surface molecule are attached to the C-termini of Fc fragment **150a** and **150b**. Seldegs as shown in FIG. 2L are designed with knobs-into-holes mutations and/or electrostatic steering mutations to promote heterodimer formation, so there is an antigen molecule of each type (**100**, **190**) in each Seldeg molecule. In FIG. 2L, both Fc fragments **150a** and **150b** have proteins or protein fragments that bind to the cell surface protein or other cell surface molecule fused to them, but in other embodiments, only one such protein or protein fragment may be present.

(77) As shown in FIG. 2M, in exemplary Seldeg **20m** two molecules of the same antigen (**100**) may be attached to Fc fragment **200** at N-terminal or other locations to generate exemplary Seldeg **20m**. This exemplary Seldeg is a homodimer that contains mutations to enhance binding to FcRn, and does not contain knobs-into-holes and/or electrostatic steering mutations.

(78) FIG. 2N is a schematic representation of exemplary Seldeg **20n** comprising two molecules of the same antigen (**100**) attached to the N-termini of an Fc fragment **210**. In the exemplary embodiment shown in FIG. 2N, protein or protein fragments **160** that bind to a cell surface receptor or cell surface molecule are attached to the C-terminus of Fc fragment **210**. This exemplary Seldeg

is a homodimer and does not contain knobs-into-holes and/or electrostatic steering mutations. In FIG. 2N, the homodimeric Fc fragment **200** has proteins or protein fragments that bind to the cell surface protein or other cell surface molecule fused to both polypeptide chains, but in other embodiments, only one such protein or protein fragment may be present.

(79) For Seldegs that target FcRn, similar principles may be applied to other proteins able to bind FcRn. In addition, the FcRn-targeting Seldeg or methods of forming it may be affected by properties of the FcRn-binding protein. Although albumin tends to not form dimers or other multimers, other FcRn-binding proteins may, in which case the final Seldeg may be formed in a manner to those containing antibody fragments so that each Seldeg contains only one copy of the antigen. In the examples shown in FIGS. 2A, 2B, 2C, 2F, 2G, 2H, 2I, 2J, 2K, 2L, the Seldeg has two antibody Fc fragments that are engineered with knobs-into-holes mutations and/or electrostatic steering mutations to drive the formation of heterodimers comprising one antigen linked to one Fc fragment and one Fc fragment with no antigen attached. The Fc fragment can be further engineered to bind to FcRn with increased affinity at near neutral pH (FIGS. 2A, 2B, 2C, 2K, 2M) or connected to one or more proteins, scFv fragments, Fab fragments or other molecules that target one or more cell surface receptors or molecules (FIGS. 2F, 2G, 2H, 2I, 2J, 2L, 2N). The Fc fragments in the examples shown in FIGS. 2F, 2G, 2H, 2I, 2J, 2L, 2N can also be engineered to bind with higher affinity to FcRn so that they target both FcRn and one or more cell surface receptors or molecules.

(80) Albumin binds more strongly to FcRn at acidic pH than at neutral pH. However, albumin may also be modified to alter its binding affinities at near-neutral or endosomal pH to encourage degradation of the target antigen-specific antibody and recycling of the Seldeg. Similarly, antibody variable region FcRn-binding proteins may be affected by pH in a manner specific to that protein, but they may still be modified to alter its binding affinities at near-neutral or endosomal pH to encourage degradation of the target antigen-specific antibody. These FcRn-binding proteins can be isolated from libraries of immunoglobulin variable domains, scFv (VH:VL heterodimers in which VH and VL domains are connected to each other by linker peptides such as GGGGSGGGGSGGGGS)(SEQ ID NO:36) or Fab fragments using phage display, yeast display or other technologies known to those with skill-in-the-art. These libraries can either be derived from naturally occurring antibody variable genes, or can be generated using approaches that result in 'semi-synthetic' libraries wherein complementarity determining regions (CDRs) are produced using randomized oligonucleotide sequences. Further increases to their affinities can be achieved by, for example, inserting random mutations in the CDRs using error-prone PCR followed by selection using phage display or yeast display. Exemplary CDR residues that would be targeted are those in CDR3 of the light chain variable domain (residues 89-97; Kabat numbering) and CDR3 of the heavy chain variable domain (residues 95-102; Kabat numbering). Similar methods can be used to isolate antibody-based proteins or scaffold-based proteins that bind to other cell surface receptors/molecules.

(81) Seldegs may include any targeting component that is configured to specifically bind to a receptor or other molecule on the cell surface (FIG. 2D, 2E, 2F, 2G, 2H, 2I, 2J, 2L, or 2N). The targeting component is fused directly or indirectly (e.g., via a linker) to an antigen component having one molecule of each type of antigen, antigen fragment or antigen mimetic to reduce antibody-mediated crosslinking. The term "type of antigen" as used herein refers to an antigen that binds to a specific antibody. Accordingly, a Seldeg can include more than one antigen type, wherein each Seldeg has only one molecule of each antigen type. If the targeting protein contains an immunoglobulin-derived Fc fragment, the Fc region can be mutated so that it does not bind, or binds at substantially reduced levels, to Fc gamma receptors and complement. Several different possible configurations of Seldegs are shown in FIG. 2A-N; these are shown as examples and are not limiting, since multiple other configurations can also be envisaged by those with skill-in-the-art.

(82) Seldegs may include Fc fragments that bind to Fc gamma receptors and complement. The presence of the Fc gamma and complement binding sites may be desired in the context of particular application areas, when an immune response against the antigen in the Seldeg is desirable (for example, in tumor imaging). In such applications, Seldegs that are configured to bind to Fc gamma receptors and complement may be preferred. For example, Seldegs that include Fc fragments lacking engineered mutations to have decreased binding affinity or no binding affinity for Fc gamma receptors and/or complement (C1q) described herein, such as arginine mutations, may be configured to elicit such an immune response. Seldegs that include Fc fragments with mutations known to those with skill in the art to increase binding **20** to Fc gamma receptors and or complement (C1q) may also be configured to elicit such an immune response. Such Seldegs may also be configured to comprise more than one antigen molecule of the same type per Seldeg (FIG. 2K, 2L, 2M, or 2N) to enhance immune complex formation.

(83) For example, Seldegs can have variations in numbers of targeting domains or antibody fragments (e.g. Fab fragments or scFv fragments) that range from 1-3 targeting domains or antibody fragments (FIG. 2). These targeting domains or antibody fragments can be linked to immunoglobulin Fc fragments, whereas in others, the targeting domains or antibody fragments may be linked to each other; the antigen and antibody fragments can be fused to Fc fragments or each other in different orientations (FIG. 2); Seldegs can include linker sequences that vary in length and composition between the fusion proteins, domains or fragments can be used e.g. IEGRMD (SEQ ID NO:37), GGGGS (SEQ ID NO:38) or 2-3 repeats of this linker; antigen mimetics such as small molecules or peptides can be used; the Fc fragment in Seldegs may be mutated so that it has substantially reduced binding affinity for Fc gamma receptors, complement, and increased affinity for binding to FcRn; The Fc fragments of a Seldeg may have mutations such as knobs-into-holes and/or electrostatic steering mutations so that heterodimers of antigen-linked and non-antigen-linked Fc fragments or containing two different antigens are formed.

(84) Seldegs can include the following antigens that include proteins, glycoproteins and nucleic acids associated with autoimmune disease, including autoimmune encephalitides: myelin oligodendrocyte glycoprotein (MOG), myelin basic protein, proteolipid protein, myelin-associated glycoprotein, myelin-associated oligodendrocyte basic protein, transaldolase, acetylcholine receptor, muscle specific kinase, low density lipoprotein receptor related protein 4, insulin, islet antigen 2, glutamic acid decarboxylase 65, zinc transporter 8, citrullinated antigens, carbamylated antigens, collagen, cartilage gp39, gp130-RAPS, 65 kDa heat shock protein, fibrillarin, small nuclear protein (snoRNP), aquaporin 4, thyroid stimulating factor receptor, nuclear antigens, DNA, histones, glycoprotein gp70, ribosomes, pyruvate dehydrogenase dehydroliamide acetyltransferase, hair follicle antigens, human tropomyosin isoform 5, N-Methyl-D-aspartate (NMDA) receptor, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA) receptor, GABA.sub.A and GABA.sub.B receptors, glycine receptor, dipeptidyl-peptidase-like protein 6 (DPPX), glutamate receptor (GluR5), voltage gated potassium channel, Hu, Thyroid peroxidase, thyroglobulin, thyroid stimulating hormone (TSH) receptor, thyroid hormones T3 and T4, desmoglein 1 and 3, among others identifiable by skilled persons. The following antigens are examples of antigens that are associated with tumors and can be incorporated into Seldegs to clear tumor-specific antibodies during diagnostic imaging: HER2, prostate specific membrane antigen (PSMA), prostate stem cell associated antigen (PSCA), c-Met, EpCAM, carcinoembryonic antigen (CEA). Other antigens include therapeutics for which patients have specific antibodies, or transplantation antigens that are recognized by antibodies in transplant recipients. In addition, it is possible to generate molecular mimics (synthetic, protein fragments etc.) of antigens, and these can also be used to generate Seldegs. The above antigens are examples and are not limiting to additional types of possible antigens identifiable by persons skilled in the art upon reading of the present disclosure.

(85) In several examples described herein, the Seldeg can be a heterodimer of fusion proteins

comprising the amino acid sequences of SEQ ID NO: 2 plus SEQ ID NO: 6, SEQ ID NO: 4 plus SEQ ID NO: 6, SEQ ID NO: 8 plus SEQ ID NO: 10, SEQ ID NO: 12 plus SEQ ID NO: 14, SEQ ID NO: 16 plus SEQ ID NO: 18 plus the antibody light chain SEQ ID NO: 20, SEQ ID NO: 22 plus SEQ ID NO: 24 plus the antibody light chain SEQ ID NO: 20, SEQ ID NO: 26 plus SEQ ID NO: 28, SEQ ID NO: 30 plus SEQ ID NO: 6, SEQ ID NO: 32 plus SEQ ID NO: 6, or SEQ ID NO: 34 plus SEQ ID NO: 6, or homologs thereof.

(86) The Seldeg can be a fusion protein comprising an amino acid sequence having at least 50% identity with SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO: 6, SEQ ID NO: 8, SEQ ID NO: 10, SEQ ID NO: 12, SEQ ID NO: 14, SEQ ID NO: 16, SEQ ID NO: 18, SEQ ID NO: 20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, or SEQ ID NO: 34.

(87) As used herein, “sequence identity” or “identity” in the context of two nucleic acid or polypeptide sequences makes reference to the nucleotide bases or residues in the two sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity or similarity is used in reference to proteins, it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted with a functionally equivalent residue of the amino acid residues with similar physiochemical properties and therefore do not change the functional properties of the molecule.

(88) A functionally equivalent residue of an amino acid used herein typically refers to other amino acid residues having physiochemical and stereochemical characteristics substantially similar to the original amino acid. The physiochemical properties include water solubility (hydrophobicity or hydrophilicity), dielectric and electrochemical properties, physiological pH, partial charge of side chains (positive, negative or neutral) and other properties identifiable to a person skilled in the art. The stereochemical characteristics include spatial and conformational arrangement of the amino acids and their chirality. For example, glutamic acid is considered to be a functionally equivalent residue to aspartic acid in the sense of the current disclosure. Tyrosine and tryptophan are considered as functionally equivalent residues to phenylalanine. Arginine and lysine are considered as functionally equivalent residues to histidine.

(89) A person skilled in the art would understand that similarity between sequences is typically measured by a process that includes the steps of aligning the two polypeptide or polynucleotide sequences to form aligned sequences, then detecting the number of matched characters, i.e. characters similar or identical between the two aligned sequences, and calculating the total number of matched characters divided by the total number of aligned characters in each polypeptide or polynucleotide sequence, including gaps. The similarity result is expressed as a percentage of identity.

(90) As used herein, “percentage of sequence identity” means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may include additions or deletions (gaps) as compared to the reference sequence (which does not include additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

(91) As used herein, “reference sequence” is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length protein or protein fragment. A reference sequence can be, for example, a sequence identifiable in a database such as GenBank and UniProt and others identifiable to those skilled in the art.

(92) As understood by those skilled in the art, determination of percent identity between any two sequences can be accomplished using a mathematical algorithm. Computer implementations of suitable mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include, but are not limited to: CLUSTAL, ALIGN, GAP, BESTFIT, BLAST, FASTA, among others identifiable by skilled persons.

(93) For example, Seldegs according to the present disclosure can have an amino acid sequence having at least 50% sequence identity, preferably at least 80%, more preferably at least 90%, most preferably at least 95% sequence identity compared to SEQ ID NO: 2, SEQ ID NO: 4, SEQ ID NO:6, or SEQ ID NO:8, or SEQ ID NO:10, or SEQ ID NO:12, or SEQ ID NO:14, or SEQ ID NO:16, or SEQ ID NO:18, SEQ ID NO:20, SEQ ID NO: 22, SEQ ID NO: 24, SEQ ID NO: 26, SEQ ID NO: 28, SEQ ID NO: 30, SEQ ID NO: 32, or SEQ ID NO: 34.

(94) The antigen-specific antibodies that are targeted by Seldegs can be autoantibodies that are present in a patient, antibodies that bind to therapeutics, antibodies that recognize transplant and modified (e.g. radiolabeled) antibodies or fragments/engineered forms that are used in diagnostic imaging, among others identifiable by persons skilled in the art.

(95) As shown in the examples below, Seldegs are able to selectively deplete target antigen-specific antibodies with specificity for their fused antigen, for example, the target antigen-specific antibodies HER2-specific trastuzumab or pertuzumab (“TZB” or “PZB”) and MOG-specific antibody (“8-18C5”). As shown in the examples below, Seldegs are able to selectively deplete the target antigen-specific antibodies without negatively affecting the total IgG levels or eliciting an adverse immune reaction. These findings stand in contrast to other approaches, in which treatment results in depletion of total IgGs, through the use of FcRn inhibitors or antibodies that destroy B-cells. Such approaches adversely affect antibodies of non-targeted specificities or B-cell function because they lack the selectivity provided by Seldegs.

(96) Based on this unique selectivity, the Seldeg platform has many applications because Seldegs may be created with many other targeting proteins and antigens. Examples of such applications include treatment of autoimmune disorders, treatment of antibody-mediated rejection prior to or during transplantation, increasing contrast during whole body imaging of tumors (e.g., the tumor antigens PSMA, EpCAM, and CEA may be used as antigens for developing additional Seldegs), depleting the bodily concentration of a particular biologic after administration, for instance, if an adverse reaction is observed, removal of antibodies to clear antibodies that recognize a therapeutic prior to delivery of the therapeutic.

(97) Seldegs may be administered in any way able to deliver them to cells expressing the receptor or other molecule on the cell surface that is being targeted, such as via injection, particularly intravenous, subcutaneous or intramuscular injection, or injection into a tissue targeted by the antigen-specific antibody that is to be depleted. Seldegs can also be expressed in cells that have been genetically engineered to contain expression constructs encoding Seldegs. In particular, cells can be genetically engineered by introducing expression constructs that encode Seldeg proteins that include proteins or peptides that allow secretion of Seldegs from the engineered cells in situ. For example, patient derived cells could be transfected with expression constructs encoding Seldegs and the transfected cells delivered back into the patient, using similar approaches to those described for chimeric antigen receptor (CAR) T cell therapy. The expression constructs would comprise an expression vector known to those skilled in the art and may, for example, contain genes encoding MOG Seldeg (SEQ ID NO: 1 and 5), with secretion leader peptides such as those derived from immunoglobulin genes linked to the 5' end of the coding sequence for the mature MOG Seldeg (SEQ ID NO: 1) and Fc (SEQ ID NO: 5).

(98) Seldegs may be administered in an amount that does not block every targeted receptor/cell surface molecule, allowing normal function of the cell surface receptor/molecule. The dose of Seldeg used may be similar to the amount of antibody being targeted for clearance, which will depend, for example, on the specifics of the antibody-mediated disease, or whether Seldegs are

being used to improve contrast in diagnostic imaging. The amount of Seldeg used is expected to be less than the total number of receptor types being targeted, so that the normal function of the targeted receptor is not adversely affected. In addition, Seldegs can be designed so that they do not compete with the natural ligand of the cell surface receptor or cell surface molecule for binding, for example, by using nanobodies (VHH) that bind to FcRn at a site that does not overlap with the IgG binding site (for example as described in Andersen, J. T., Gonzalez-Pajuelo, M., Foss, S., Landsverk, O. J. B., Pinto, D., Szyroki, A., de Haard, H. J., Saunders, M., Vanlandshoot, P., Sandlie, I. (2012) Selection of nanobodies that target human neonatal receptor. *Sci. Rep.*, 3, 1118). In addition, Seldegs may remove less than 10%, less than 5%, less than 1%, or less than 0.1% of non-targeted antibodies in the circulation or in a tissue targeted by the antigen-specific antibody that is to be depleted. Retention of non-targeted antibodies during and after Seldeg treatment may be important in normal immune function and the avoidance of infections, among other effects as described herein.

(99) Seldegs may be administered daily, weekly, or whenever 50% of patients are expected to have regenerated a threshold amount of targeted antigen-specific antibody in the circulation or in a tissue recognized by the targeted antigen-specific antibody. The levels of targeted antigen-specific antibody can be determined by using enzyme-linked immunosorbent assays (ELISAs) to analyze serum samples. Alternatively, other methods that are well known to those with skill in the art can be used.

(100) In transplant patients at risk for antibody-mediated rejection, the Seldegs may be administered before or after transplantation. An emergency dose of Seldegs may be administered if the target antigen-specific antibody reaches a threshold amount of target antigen-specific antibody in the circulation or in a tissue recognized by the target antigen-specific antibody. The levels of targeted antigen-specific antibody can be determined by using enzyme-linked immunosorbent assays (ELISAs) to analyze serum samples. Alternatively, other methods that are well known to those with skill in the art can be used.

(101) In patients that have antibodies specific for a therapeutic, such as a protein-based therapeutic, the Seldegs may be administered before the delivery of the therapeutic to deplete such antibodies. This is expected to overcome problems associated with rapid antibody-mediated clearance of therapeutics if they have elicited an immune response during prior delivery, or if preexisting antibodies specific for the therapeutic are present in the patient. Preexisting or induced antibodies specific for the protein-based therapeutic can be detected using a number of different methods (for example such as those described in Xue, L., Clements-Egan, A., Amaravadi, L., Birchler, M., Gorovits, B., Liang, M., Myler, H., Purushothama, S., Manning, M. S., Sung, C. (2017) Recommendations for the assessment and management of preexisting drug-reactive antibodies during biotherapeutic development. *AAPS*, 19, 1576-1586).

(102) In diagnostic/theranostic imaging, it is expected that delivery of the radiolabeled imaging antibody will be followed by a period to allow tumor localization. Subsequently, Seldegs can be used to clear radiolabeled antibody from off-target sites (e.g. circulation) to result in improved contrast. For example, this approach can include the following steps: first, a patient is injected with radiolabeled (or other label) antibody that binds to a tumor antigen. Second, following a period (e.g. 16-24 hours) to allow the radiolabeled antibody to bind to the tumor, the Seldeg is injected in an amount that is equivalent in molar amounts to the injected dose of imaging agent. Following a clearance period (e.g. 8-24 hours), the patient is imaged using positron emission tomography or other whole body imaging modality.

(103) The Seldeg may be administered in an amount sufficient to deplete at least 50%, at least 80% or at least 90% of the concentration of the target antigen-specific antibody in the circulation or in a tissue recognized by the target antigen-specific antibody within one hour, two hours, five hours, 24 hours or 48 hours or longer of administration. The persistence of the Seldeg in the body will be a determinant of how long it has activity in depleting antigen-specific antibody. Seldegs can be

designed to have different in vivo half-lives by the behavior of the cell surface receptor or cell surface molecule that they target. The affinity of the Seldeg for this cell surface receptor or cell surface molecule can also be modified using mutagenesis and approaches known to those with skill in the art, to result in Seldegs that have different persistence in the circulation and/or tissues. In particular, the Seldeg may be administered in an amount at least approximately equimolar with the amount of antigen-specific antibody to be depleted. The Seldeg can be administered in an amount that is, for example, in an approximately 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1 or higher molar ratio to the target antigen specific antibody. In particular, for example, wherein a Seldeg targets an antibody that is administered to a patient (e.g. anti MOG, or anti HER2 antibodies, and the like) a Seldeg can be administered in a dose that is for example, in an approximately 1:1, 2:1, 3:1, 4:1, 5:1, 6:1, 7:1, 8:1, 9:1, 10:1 or higher molar ration to the target antigen specific antibody.

EXAMPLES

(104) The following examples are provided to further illustrate specific embodiments of the disclosure. They are not intended to disclose or describe each and every aspect of the disclosure in complete detail and should be not be so interpreted. Unless otherwise specified, designations of cells lines and compositions are used consistently throughout these examples.

Example 1—Expression and Purification of Seldegs that Bind to FcRn

(105) Seldegs that target FcRn on cells contain a recombinant antigen as a monomer linked to a dimeric, human IgG1-derived Fc fragment (FIG. 3A) with mutations to eliminate interactions with human FcγRs and to enhance the binding affinity of the Seldeg Fc fragment to FcRn at near-neutral pH. Heterodimer formation of these two Seldegs is achieved by inserting ‘knobs-into-holes’ mutations in the CH3 domains.

(106) Expression constructs to generate the exemplary HER2-Seldeg (SEQ ID NOS: 3, 4, 5 and 6) were made as follows: to express the polypeptide chain with HER2 fused to an engineered Fc fragment (SEQ ID NO: 3), the gene encoding the HER2 leader peptide and extracellular domain (ECD consisting of 630 residues) was isolated from a HER2-overexpressing breast cancer cell line (BT-474) employing standard molecular biology techniques. This gene was fused via a IEGRMD linker peptide to the N-terminus of the hinge region of a gene encoding the human IgG1-derived Fc fragment using splicing by overlap extension. Mutations to ablate binding to FcγRs (G.sub.236R/L328R; EU numbering), enhance binding to FcRn (MST-HN; M252Y/S254T/T256E/H433K/N434F; EU numbering) and generate ‘knobs-into-holes’ (Y349T/T394F; EU numbering) were inserted into the Fc fragment gene using standard methods. Cysteine (C220; EU numbering) in the hinge region that bridges with cysteine in the light chain constant region was also mutated to serine. Fc fragment genes with FcRn-enhancing mutations, mutations to ablate FcγR binding and without fused antigen were generated with complementary knobs-into-holes mutations (S364H/F405A; EU numbering) (SEQ ID NO: 5). Recombinant proteins were expressed in HEK-293F (Life Technologies) cells following transient transfection with the Gibco Expi293™ expression system kit (Life Technologies). The HER2-Seldeg was purified from culture supernatants using an anion exchange column (SOURCE-15Q, GE Healthcare) at pH 8.0 and a linear salt gradient (0-0.5 M NaCl). Alternatively, HER2-Seldeg can be purified using protein A-Sepharose and standard methods. Following elution from the column (ion exchange or protein A-Sepharose) HER2-Seldeg was dialyzed against phosphate buffered saline (PBS). HER2-Seldeg was further purified using size exclusion chromatography (SEC) (GE Healthcare) in PBS (Lonza) prior to use in experiments. Expression plasmids for other Seldegs were made using analogous methods, and recombinant proteins expressed in transfected HEK-293F cells. Seldegs without FcRn-enhancing mutations (e.g. MOG-Seldeg-PS; MOG-Seldeg-TfRSEQ ID NOS: 7, 8, 9, 10, 11, 12, 13 and 14, 15, 16, 17, 18, 19, and 20) were purified using protein G-Sepharose.

(107) Seldegs targeting antibodies specific for two antigens were generated: (1) the HER2-Seldeg; and (2) the MOG-Seldeg. The antigen HER2 is a well-defined target for therapy and also for

diagnostic imaging of HER2-overexpressing tumors with HER2-specific antibodies, such as trastuzumab (TZB) or pertuzumab (PZB). The antigen MOG is recognized by autoreactive antibodies in both animal models of multiple sclerosis (MS) and MS in humans.

(108) The HER2-Seldeg and MOG-Seldeg maintain a significantly higher binding affinity for FcRn at neutral pH and at an acidic pH, in contrast to recombinant fusion proteins comprising HER2 or MOG fused to Fc fragments to generate analogous constructs (“HER2-WT” and “MOG-WT”, respectively) to the HER2-Seldeg and MOG-Seldeg, except they lack the FcRn-enhancing MST-HN mutations (M252Y, S254T, T256E, H433K, N434F; EU numbering) that increase binding affinity at near-neutral pH (FIG. 3B). Surface plasmon resonance experiments to analyze the interactions of the recombinant proteins with FcRn were carried out using a BIAcore T200 (GE Healthcare). Binding of Seldeg/WT to recombinant mouse FcRn was analyzed by injecting 100 nM MOG/HER2-Seldeg or MOG/HER2-WT over immobilized FcRn (coupled at ~600 RU on a CM5 sensor chip) in PBS (pH 6.0 or 7.4) plus 0.01% v/v Tween-20 at a flow rate of 10 µl/minute. Flow cells were regenerated following each injection and dissociation phase using 0.15 M NaCl, 0.1 M sodium bicarbonate, pH 8.5. Data were zero-adjusted and background-subtracted (background obtained by injection over a flow cell coupled with buffer only during coupling reaction).

(109) Mutations to increase FcRn such as MST-HN were identified using the following approach: residues in proximity to amino acids (e.g. 253, 435) that are known to be essential for FcRn binding were randomly mutated in an Fc fragment gene and the libraries of mutated Fc fragments displayed on phage. Fc fragments with increased binding affinity for FcRn were selected using phage display technology (Ghetie, V., Popov, S., Borvak, J., Radu, C., Matesoi, D., Medesan, C., Ober, R. J., Ward, E. S. (1997) Increasing the serum persistence of an IgG fragment by random mutagenesis, *Nature Biotech.*, 15, 637-640; Dall'Acqua, W. F., Woods, R. M., Ward, E. S., Palaszynski, S. R., Patel, N. K., Brrewah, Y. A., Wu, H., Kiener, P. A., Langermann, S. (2001) Increasing the affinity of a human IgG1 for the neonatal receptor: biological consequences, *J. Immunol.*, 169, 5171-5180). Alternatively, these residues can be mutated to every other possible amino acid and Fc fragments with higher affinity for FcRn identified using methods such as ELISA or surface plasmon resonance binding analyses.

(110) Size exclusion analyses indicate that the recombinant proteins comprising the Seldegs do not form aggregates following incubation in phosphate buffered saline when incubated for up to 30 days at 4° C. or 5 days at 37° C. (FIG. 3C).

Example 2—Ability of Seldegs that Target FcRn to Deplete Target Antigen-Specific Antibodies in Mice that Transgenically Express Human FcγRs (huFcγR Mice)

(111) To determine the ability of Seldegs to deplete target antigen-specific antibodies, mice that transgenically express human FcγRs (huFcγR mice; Smith, P., DiLillo, D. J., Bournazos, S., Li, F., Ravetch, J. V. (2012). Mouse model recapitulating human Fc receptor structural and functional diversity. *Proc. Natl. Acad. Sci. USA* 109, 6181-6186) were injected with 15 µg of radiolabeled (.sup.125I-labeled) MOG-specific antibody 8-18C5. 24-hours after the huFcγR mice were injected with 8-18C5, 125 µg or 31 µg of MOG-Seldeg or a control was administered via injection. In FIG. 3D, “MOG-Seldeg-High” corresponds to the 125 µg dose and “MOG-Seldeg-Low” corresponds to the 31 µg dose. The controls are phosphate-buffered saline (“PBS”) and unmodified MOG-Fc fusion protein (“MOG-WT”). MOG-WT is an analogous construct to the MOG-Seldeg, except it lacks the FcRn-enhancing MST-HN mutations that increase binding affinity at near-neutral pH.

(112) To generate the data of FIG. 3D whole body counts of radiolabeled 8-18C5 were taken at the indicated times. Counts per minute (“CPM”) obtained 24 hours after 8-18C5 administration, immediately prior to Seldeg or control delivery, are taken as 100% for each mouse and CPM of whole body obtained thereafter are normalized to this time point. The error bars indicate standard deviations and statistically significant differences were determined using two-way ANOVA with Tukey's multiple comparisons, with $p < 0.05$ and $n = 6$ mice/group.

(113) Doses of 125 µg and 31 µg of MOG-Seldeg are approximately 2 to 8-fold lower (on a molar

basis) than a dose of 500 µg, a MST-HN Abdeg, which has been shown to globally eliminate IgG levels in mice. Such lower doses of MOG-Seldeg were used to minimize effects on IgGs that were not specific for the fused antigen through competition with the Seldeg for FcRn binding. The doses of 125 µg or 31 µg represent an approximate 16 and 4-fold molar excess, respectively, over the 8-18C5 target antigen-specific antibody.

(114) FIG. 3D provides a graph of normalized body counts of radiolabeled 8-18C5 (the target antigen-specific antibody) versus time. Administration of MOG-Seldeg resulted in a rapid, dose-dependent decrease in 8-18C5 in normalized whole body counts (FIG. 3D), which demonstrates the ability of MOG-Seldeg to selectively deplete 8-18C5 from the body. In contrast, the injection of control MOG-WT, lacking the FcRn-enhancing MST-HN mutations, had no effect on 8-18C5 antibody depletion. This demonstrates the importance of the MST-HN mutations to confer a high binding affinity for FcRn at pH 6.0-7.4.

Example 3—Specificity of Seldegs that Target FcRn and their Ability to Induce a Significant Decrease in Target Antigen-Specific Antibodies Levels, Both in Blood and Throughout the Body

(115) To analyze the specificity of Seldegs and their effect on antibodies with different antigen recognition properties, the behavior of the HER2-specific antibody trastuzumab (TZB) was investigated in the presence of both HER2-Seldeg and MOG-Seldeg (FIG. 3E). To generate the data of FIG. 3E, huFcγR mice were intravenously injected with 15 µg of radiolabeled (.sup.125I-labeled) TZB and subsequently with a 4-fold molar excess of HER2-Seldeg, or, as a control for antigen specificity, MOG-Seldeg. Additional controls are phosphate-buffered saline (“PBS”), Abdeg and a recombinant fusion protein comprising HER2 fused to an Fc fragment to generate an analogous construct (“HER2-WT”) to the HER2-Seldeg, except it lacks the FcRn-enhancing MST-HN mutations that increase binding affinity at near-neutral pH. In contrast to a Seldeg, an “Abdeg” is a human IgG1-derived antibody with the MST-HN mutations that non-selectively depletes antibodies. Such proteins are called Abdegs because they are antibodies that generally cause IgG degradation.

(116) To generate the data of FIG. 3E, the mice were bled and blood and whole body counts were taken at the indicated times. Counts per minute (“CPM”) obtained 24-hours after TZB administration, immediately prior to Seldeg or control delivery, are taken as 100% for each mouse and CPM of blood and whole body obtained thereafter are normalized to this time point. The error bars indicate standard deviations and statistically significant differences were determined using two-way ANOVA with Tukey's multiple comparisons, with $p < 0.05$ and $n = 6$ mice/group.

(117) FIG. 3E provides graphs of normalized blood and body counts of radiolabeled TZB (the target antigen-specific antibody) versus time. Administration of HER2-Seldeg caused a significant decrease in normalized body counts of TZB, both in blood and whole body counts, which demonstrates the ability of HER2-Seldeg to selectively deplete TZB from the body. In contrast, the control proteins demonstrated similar behavior to that observed for the PBS control (FIG. 3E).

Example 4—Ability of Seldegs that Target FcRn to Rapidly Deplete Target Antigen-Specific Antibodies

(118) Mice were intravenously administered with radiolabeled (.sup.125I) TZB (15 µg) and 24 hours later HER2-Seldeg (51 µg; four-fold molar excess over TZB) was delivered intravenously. Additional controls are phosphate-buffered saline (“PBS”) and Abdeg (60 µg).

(119) To generate the data of FIG. 3F, the mice were bled and blood and whole body counts were taken at the indicated times. Counts per minute (“CPM”) obtained 24-hours after TZB administration, immediately prior to Seldeg or control delivery, are taken as 100% for each mouse and CPM of blood and whole body obtained thereafter are normalized to this time point. The error bars indicate standard deviations and statistically significant differences were determined using two-way ANOVA with Tukey's multiple comparisons, with $p < 0.05$ and $n = 5-6$ mice/group.

(120) FIG. 3F provides graphs of normalized blood and body counts of radiolabeled TZB (the target antigen-specific antibody) versus time. Administration of HER2-Seldeg caused a rapid

decrease in normalized body and blood counts of TZB, which demonstrates the ability of the HER2-Seldeg to rapidly deplete TZB from the body. Notably, the blood counts of TZB were reduced close to background levels within two hours of Seldeg administration (FIG. 3F), further supporting the ability of the HER2-Seldeg to deplete target antigen-specific antibodies rapidly from the blood. In contrast, at a dose of 60 µg/mouse, delivery of Abdeg resulted in similar behavior to that observed for the PBS control (FIG. 3F).

Example 5—Activity of Seldegs that Target Exposed Phosphatidylserine (PS) on the Cell Surface (121) We also investigated the ability of a Seldeg in which the targeting protein is the C2 domain of synaptotagmin 1 (Syt1) to selectively deplete antibodies specific for MOG in huFcγR mice. Syt1 binds to exposed PS on cells, and the Seldeg is therefore designed MOG-Seldeg-PS and is shown schematically (FIG. 4A). MOG-Seldeg-PS(DN) that does not bind to PS due to the presence of the ‘DN’ mutations (D173N, D179N, D231N, D233N and D239N) was also generated. MOG Seldeg PS and MOG Seldeg PS(DN) were purified from culture supernatants of transfected HEK 293F cells using protein G-Sepharose and standard methods. Heterodimer formation was achieved by inserting knobs-into-holes and electrostatic steering mutations into the Fc regions, and size exclusion analyses indicate that the recombinant proteins are well behaved (FIG. 4A).

(122) To generate the data shown in FIG. 4B, mice were intravenously injected with radiolabeled (.sup.125I) chimeric 8-18C5 (human constant/mouse variable domains; MOG-specific; 15-20 µg) and 24 hours later phosphate-buffered saline (PBS), 40 µg MOG-Seldeg-PS or as controls, 34 µg Fc-Syt1 (no MOG attached) or 40 µg MOG-Seldeg-PS(DN) were delivered intravenously. Radioactivity levels were determined at the indicated times. Whole body CPM or blood CPM levels obtained immediately prior to MOG-Seldeg-PS or control delivery were taken as 100% and all subsequent CPM that were obtained were normalized against these CPM levels. Error bars indicate the standard error of the mean (SEM) and statistically significant differences between MOG-Seldeg-PS treated group and control groups (Fc-Syt1, MOG-Seldeg-PS(DN) and PBS) are indicated by * (p<0.05, two-way ANOVA with Tukey's multiple comparisons; n=5-6 mice/group).

(123) Administration of MOG-Seldeg-PS caused a significant decrease in normalized body counts of 8-18C5, both in blood and whole body counts, which demonstrates the ability of MOG-Seldeg-PS to selectively deplete 8-18C5 from the body. In contrast, the control protein Fc-Syt1 demonstrated similar behavior to that observed for the PBS control, and although MOG-Seldeg-PS(DN) induced a decrease in 8-18C5 (due to residual binding to PS), the effect was much lower than that of MOG-Seldeg-PS (FIG. 4B).

Example 6—Ability of Seldegs that Target FcRn to Efficiently Internalize Target Antigen-Specific Antibodies into Endosomes and Cause Degradation Via Delivery to Lysosomes

(124) To determine the mechanism of activity of Seldegs at the cellular level, flow cytometry and fluorescence microscopy were performed to analyze the effects of Seldegs on the internalization and accumulation of target antigen-specific antibodies TZB and 8-18C5. First, human endothelial cells (HMEC-1) were transfected with a human FcRn-GFP expression construct, which was mutated so that its binding properties are analogous to mouse FcRn.

(125) The Seldegs efficiently internalized bound target antigen-specific antibodies into endosomes within cells, and following 8 hours of incubation, the target antigen-specific antibodies were delivered to lysosomes.

(126) FIG. 5A is a graph of mean fluorescence intensity (“MFI”) and FIG. 5B and FIG. 5C are microscopic images of Seldeg activity in the presence of target antigen-specific antibodies, with microscopic images of representative endosomes cropped, expanded, and presented in the top right-corner insets. FIG. 5A shows a greater MFI in the presence of, respectively, Alexa 647-labeled TZB with HER2-Seldeg and Alexa 647-labeled 8-18C5 with MOG-Seldeg. This greater MFI indicates that co-incubation of HER2- and MOG-Seldeg, respectively, with TZB and 8-18C5 (400 nM Seldeg or WT control plus 100 nM antigen-specific antibody; 30 minute pulse, no chase or 60 minute chase, labeled 30' P or 30' P 60' C respectively, in FIGS. 5A, 5B, and 5C) results in high

levels of TZB and 8-18C5 entering cells. Similar results were obtained when the HER2-specific antibody pertuzumab (PZB) was used instead of TZB in the presence of HER2-Seldeg (FIG. 5A). Further, the majority of each target antigen-specific antibody is retained by the cells during a 60 minute chase. In contrast, the accumulation of target antigen-specific antibodies within cells is substantially lower in the presence of negative controls HER2-WT (selective for TZB or PZB) and MOG-WT (selective for 8-18C5).

(127) FIGS. 5B and C show that antigen-specific antibodies accumulate in cells and are associated with FcRn in endosomes in the presence of Seldegs. To generate the microscopic images of FIG. 5B, HMEC-1 cells were pulsed with 100 nM Alexa 647-labeled TZB (HER2-specific) in complex with 400 nM Alexa 555-labeled HER2-Seldeg or HER2-WT (as indicated) for 30 minutes, washed and either immediately fixed, or washed, chased in medium for 60 minutes, followed by fixation. Fixed cells were imaged using fluorescence microscopy. The data in FIG. 5C was generated using the same approach, except that 100 nM Alexa 647-labeled 8-18C5 (MOG-specific) in complex with 400 nM Alexa 555-labeled MOG-Seldeg or MOG-WT was used. Bars in each of the microscopic images are 5 μ m, and bars in each of the microscopic image insets are 0.25 μ m. The data shown in FIGS. 5B and 5C demonstrates that antigen-specific antibodies accumulate to substantially higher levels in FcRn-expressing cells in the presence of Seldegs that bind to FcRn compared with in the presence of control (WT) proteins.

(128) FIG. 6A shows microscopic images of Seldeg activity in the presence of target antigen-specific antibodies, with microscopic images of representative lysosomes cropped, expanded, and presented in the top right-corner insets. For FIG. 6A, HMEC-1 cells were pre-pulsed with Alexa 555-labeled Dextran for 2 hours and washed, and then pulsed with 100 nM Alexa 647-labeled 8-18C5 (MOG-specific) in complex with 400 nM MOG-Seldeg or MOG-WT (as indicated) for 30 minutes, followed by an 8 hour chase and then washed, fixed and imaged. To generate the microscopic images of FIG. 6B, HMEC-1 cells were pre-pulsed with Alexa 555-labeled Dextran for 2 hours and washed, and then pulsed with either 100 nM Alexa 647-labeled TZB or 8-18C5 antibodies, and 400 nM MOG-Seldeg or HER2-Seldeg (as indicated) for 30 minutes followed by an 8 hour chase, and then washed, fixed and imaged. For each overlay image (shown in the right column), GFP, Alexa 555 and Alexa 647 are pseudo-colored green, red and blue, respectively. Bars in each of the microscopic images are 5 μ m, and bars in each of the microscopic image insets are 0.25 μ m. Notably, FIG. 6B shows that the cells do not accumulate TZB and 8-18C5 in lysosomes in the presence of MOG-Seldeg or HER2-Seldeg, respectively, which indicates the antigen specificity of their effects.

Example 7—Ability of Seldegs that Target Phosphatidylserine to Efficiently Internalize Target Antigen-Specific Antibodies into Endosomes and Cause Degradation Via Delivery to Lysosomes

(129) FIG. 7 shows that MOG-Seldeg-PS efficiently internalizes MOG-specific antibodies (chimeric 8-18C5) into endothelial cells and macrophages and uptake is dependent on PS-binding. Endothelial cells (2H11) or macrophages (RAW264.7) that expose PS were incubated with 10 nM Alexa 647-labeled chimeric 8-18C5 mixed with 20 nM MOG-Seldeg-PS, MOG-Seldeg-PS(DN) or Fc-Syt1 (no MOG) for 2 hours. Mean fluorescence intensities (MFI) of Alexa 647-labeled chimeric 8-18C5 (made by fusing the variable domains of the mouse antibody, 8-18C5, to the constant regions of human IgG1, C κ , using methods well known to those skilled in the art) for triplicate samples were determined by flow cytometry.

(130) This exemplary PS-binding Seldeg is calcium-dependent and therefore dissociates in endosomes where the calcium concentration is much lower. Not all PS-targeting Seldegs are expected to be calcium-dependent, but some, also such as those comprising annexin V will have this property, as would be understood by skilled persons. In addition, several antibodies can bind to beta-2 glycoprotein 1 that can in turn bind to PS.

Example 8—Ability of Seldegs that Target the Transferrin Receptor to Efficiently Internalize Target Antigen-Specific Antibodies into Cells Expressing the Human Transferrin Receptor

(131) A Seldeg comprising antigen (MOG) fused to an antibody that targets the transferrin receptor (MOG-Seldeg-TfR) has been generated and is shown schematically in FIG. 8. MOG-Seldeg-TfR and control antibody, Ab-TfR (no MOG present), were purified from culture supernatants of transfected HEK-293F cells using protein G-Sepharose and standard methods. Seldeg-TfR efficiently internalizes MOG-specific antibodies (chimeric 8-18C5) into endothelial (HMEC-1) cells and uptake is dependent on the presence of MOG in the Seldeg (FIG. 9). HMEC-1 cells that express human TfR were incubated with 50 nM Alexa 647-labeled chimeric 8-18C5 (specific for MOG) mixed with 200 nM MOG-Seldeg-TfR, or TfR-specific antibody (Ab-TfR; no MOG) for 30 minutes followed by washing and no chase (30'P), or for 30 minutes followed by washing and a 60 minute chase period (30'P, 60'C). Mean fluorescence intensities (MFI) of Alexa 647-labeled 8-18C5 for triplicate samples were determined by flow cytometry (FIG. 9).

Example 9—Ability to Seldegs that Target FcRn to Clear Background During Positron Emission Tomographic Analyses of Tumors Using Radiolabeled Antibodies

(132) Female BALB/c SCID mice were implanted in the mammary fat pad (0.5×10^6 cells/mouse; cells were suspended in RPMI-1640/Matrigel prior to injection) with the HER2-positive breast cancer cell line, HCC1954. 6-7 days later, mice were injected with ^{124}I labeled TZB (60 $\mu\text{g}/\text{mouse}$) and analyzed using positron emission tomography (PET) 22 hours later. Mice were injected with a molar equivalent of HER2-Seldeg (51 $\mu\text{g}/\text{mouse}$) or, as controls, MOG-Seldeg (31 $\mu\text{g}/\text{mouse}$) or PBS vehicle ($n=3$ mice/group). Mice were analyzed using PET 4 hours following injection of the Seldegs or PBS. Imaging of mice was carried out using a Siemens Inveon PET-computed tomography (CT) Multimodality System.

(133) To generate the data shown in FIG. 10A, mice were imaged at 22 hours following injection of ^{124}I labeled TZB (22 h) and at 4.5-5 hours post-injection ('4.5 h clearing') of HER2-Seldeg, MOG-Seldeg or PBS. PET and CT images were co-registered in the AMIDE software package. FIG. 10B provides graphs of contrast measures for radiolabel intensity for 22 hours post-injection of ^{124}I labeled TZB and 4 hours post-injection of HER2-Seldeg, MOG-Seldeg or PBS. To determine contrast measures, two regions of interest were identified in the thoracic region (background) and tumor. The mean values ($n=3$ mice/group) for the ratios of mean intensity in the tumor to mean intensity in the thoracic regions are plotted. Error bars indicate standard errors. Statistically significant differences ($p<0.05$) between mice treated with HER2-Seldeg vs. MOG-Seldeg or PBS are shown. The data show that delivery of HER2-Seldeg reduces the background in the thoracic region of the mice.

Example 10—DNA and Proteins Sequences of Exemplary Seldegs and Controls and Variants Thereof Having Mutations Such as Knobs-into-Holes Mutations, Arginine Mutations and Electrostatic Steering Mutations

(134) Table 1 shows DNA sequences of polynucleotides encoding exemplary proteins described herein, and Table 2 shows amino acid sequences of exemplary proteins encoded by the polynucleotides shown in Table 1, wherein the DNA sequence of SEQ ID NO: 1 encodes the protein of SEQ ID NO:2, the DNA sequence of SEQ ID NO: 3 encodes the protein of SEQ ID NO:4, the DNA sequence of SEQ ID NO: 5 encodes the protein of SEQ ID NO: 6, the DNA sequence of SEQ ID NO: 7 encodes the protein of SEQ ID NO: 8, the DNA sequence of SEQ ID NO: 9 encodes the protein of SEQ ID NO: 10, the DNA sequence of SEQ ID NO: 11 encodes the protein of SEQ ID NO: 12, the DNA sequence of SEQ ID NO: 13 encodes the protein of SEQ ID NO: 14, the DNA sequence of SEQ ID NO: 15 encodes the protein of SEQ ID NO: 16, the DNA sequence of SEQ ID NO: 17 encodes the protein of SEQ ID NO: 18, the DNA sequence of SEQ ID NO: 19 encodes the protein of SEQ ID NO: 20, the DNA sequence of SEQ ID NO: 21 encodes the protein of SEQ ID NO: 22, the DNA sequence of SEQ ID NO: 23 encodes the protein of SEQ ID NO: 24, the DNA sequence of SEQ ID NO: 25 encodes the protein of SEQ ID NO: 26, the DNA sequence of SEQ ID NO: 27 encodes the protein of SEQ ID NO: 28, the DNA sequence of SEQ ID NO: 29 encodes the protein of SEQ ID NO: 30, the DNA sequence of SEQ ID NO: 31 encodes the

protein of SEQ ID NO: 32, the DNA sequence of SEQ ID NO: 33 encodes the protein of SEQ ID NO: 34.

(135) TABLE-US-00001 TABLE 1 DNA sequences of polynucleotides encoding exemplary proteins. Polynucleotide encoding protein DNA sequence SEQ ID NO: MOG-Seldeg GGACAATTCAGAGTGATAGGACCAGGGTATCCCATCCGG 1 with MST-HN, GCTTTAGTTGGGGATGAAGCAGAGCTGCCGTGCCGCATC knobs-into-holes TCTCCTGGGAAAAATGCCACGGGCATGGAGGTGGGTTGG and arginine TACCGTTCTCCCTTCTCAAGAGTGGTTCACCTCTACCGAA mutations ATGGCAAGGACCAAGATGCAGAGCAAGCACCTGAATACC GGGGACGCACAGAGCTTCTGAAAGAGACTATCAGTGAGG GAAAGGTTACCCTTAGGATTCAGAACGTGAGATTCTCAG ATGAAGGAGGCTACACCTGCTTCTTCAGAGACCACTCTTA CCAAGAAGAGGCAGCAATGGAGTTGAAAGTGGAAGATG GAGGCGGTGGATCAGTTGAGCCCAAATCTTCTGACAAAA CTCACACATGCCCACCGTGCCCAGCACCTGAACTCCTGA GGGGACCGTCAGTCTTCCTCTTCCCCCCTAAAACCCAAGGA CACCCTCTACATCACTCGGGAACCTGAGGTCACATGCGTG GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGA ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC GCCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAG GGCAGCCCCGAGAACCACAGGTGACCACCCTGCCCCCAT CCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCT GCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGG AGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAG ACCTTCCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCT CTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCA GGGGAACGTCTTCTCATGCTCTGTGATGCATGAGGCTCTG AAATTCCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG GTAAA HER2-Seldeg ACCCAAGTGTGCACCGGCACAGACATGAAGCTGCGGCTC 3 with MST-HN, CCTGCCAGTCCCGAGACCCACCTGGACATGCTCCGCCAC knobs-into-holes CTCTACCAGGGCTGCCAGGTGGTGCAGGGAAACCTGGAA and arginine CTCACCTACCTGCCCACCAATGCCAGCCTGTCTTCCTGC mutations AGGATATCCAGGAGGTGCAGGGCTACGTGCTCATCGCTC ACAACCAAGTGAGGCAGGTCCCCTGCAGAGGCTGCGGA TTGTGCGAGGCACCCAGCTCTTTGAGGACAACCTATGCCCT GGCCGTGCTAGACAATGGAGACCCGCTGAACAATACCAC CCCTGTCACAGGGGCCTCCCCAGGAGGCCTGCGGGAGCT GCAGCTTCGAAGCCTCACAGAGATCTTGAAAGGAGGGGT CTTGATCCAGCGGAACCCCCAGCTCTGCTACCAGGACAC GATTTTGTGGAAGGACATCTTCCACAAGAACAACAGCT GGCTCTCACACTGATAGACACCAACCGCTCTCGGGCCTGC CACCCCTGTTCTCCGATGTGTAAGGGCTCCCGCTGCTGGG GAGAGAGTTCTGAGGATTGTCAGAGCCTGACGCGCACTG TCTGTGCCGGTGGCTGTGCCCCGCTGCAAGGGGCCACTGCC CACTGACTGCTGCCATGAGCAGTGTGCTGCCGGCTGCAC GGGCCCCAAGCACTCTGACTGCCTGGCCTGCCTCCACTTC AACCACAGTGGCATCTGTGAGCTGCACTGCCCAGCCCTG

GTCACTACACAGAGCTTTGAGTCCATGCCCAATC
CCGAGGGCCGGTATACATTCGGCGCCAGCTGTGTGACTG
CCTGTCCCTACAACCTTTCTACGGACGTGGGATCCTG
CACCTCTGTCTGCCCCCTGCACAACCAAGAGGTGACAGC
AGAGGATGGAACACAGCGGTGTGAGAAGTGCAGCAAGC
CCTGTGCCCCGAGTGTGCTATGGTCTGGGCATGGAGCACTT
GCGAGAGGTGAGGGCAGTTACCAGTGCCAATATCCAGGA
GTTTGCTGGCTGCAAGAAGATCTTTGGGAGCCTGGCATT
CTGCCGGAGAGCTTTGATGGGGACCCAGCCTCCAACACT
GCCCCGCTCCAGCCAGAGCAGCTCCAAGTGTTTGAGACT
CTGGAAGAGATCACAGGTTACCTATACATCTCAGCATGG
CCGGACAGCCTGCCTGACCTCAGCGTCTTCCAGAACCTGC
AAGTAATCCGGGGACGAATTCTGCACAATGGCGCCTACT
CGCTGACCCTGCAAGGGCTGGGCATCAGCTGGCTGGGGC
TGCGCTCACTGAGGGAACCTGGGCAGTGGACTGGCCCTCA
TCCACCATAACACCCACCTCTGCTTCGTGCACACGGTGCC
CTGGGACCAGCTCTTTCGGAACCCGCACCAAGCTCTGCTC
CACACTGCCAACC GGCCAGAGGACGAGTGTGTGGGCGAG
GGCCTGGCCTGCCACCAGCTGTGCGCCCGAGGGCACTGC
TGGGGTCCAGGGGCCACCCAGTGTGTCAACTGCAGCCAG
TTCCTTCGGGGCCAGGAGTGCGTGGAGGAATGCCGAGTA
CTGCAGGGGCTCCCCAGGGAGTATGTGAATGCCAGGCAC
TGTTTGCCGTGCCACCCTGAGTGTGAGCCCCAGAATGGCT
CAGTGACCTGTTTTGGACCGGAGGCTGACCAGTGTGTGG
CCTGTGCCCACTATAAGGACCCTCCCTTCTGCGTGGCCCG
CTGCCCCAGCGGTGTGAAACCTGACCTCTCCTACATGCCC
ATCTGGAAGTTTCCAGATGAGGAGGGCGCATGCCAGCCT
TGCCCCATCAACTGCACCCACTCCTGTGTGGACCTGGATG
ACAAGGGCTGCCCCGCGGAGCAGAGAGCCAGCCCTCTGA
CGATTGAAGGCCGCATGGATCCCAAATCTTCTGACAAA
CTCACACATGCCACCGTGCCCAGCACCTGAACTCCTGA
GGGGACCGTCAGTCTTCCTCTTCCCCCCAAAACCCAAGGA
CACCTCTACATCACTCGGGAACCTGAGGTCACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC
AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG
ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT
GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGA
ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
GCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAG
GGCAGCCCCGAGAACCACAGGTGACCACCCTGCCCCCAT
CCCGGGATGAGCTGACCAAGAACCAGGTGAGCCTGACCT
GCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGG
AGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAG
ACCTTCCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCT
CTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCA
GGGGAACGTCTTCTCATGCTCTGTGATGCATGAGGCTCTG
AAATTCCACTACACGCAGAAGAGCCTCTCCCTGTCTCCTG GTAAA Fc with MST-
GTTGAGCCCAAATCTTCTGACAAAACCTCACACATGCCAC 5 HN, knobs-into-
CGTGCCAGCACCTGAACTCCTGAGGGGACCGTCAGTCTT holes and
CCTCTTCCCCCCAAAACCCAAGGACACCCTCTACATCACT arginine

CGGGAACCTACATGCTGGTGGTGGACGTGAGC mutations
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGAC
GGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA
GGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCT
CACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAAACAAAGCCCCGCCAGCCCCCAT
CGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGA
ACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCT
GACCAAGAACCAGGTCCACCTGACCTGCCTGGTCAAAGG
CTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAA
TGGGCAGCCGGAGAACAATAAGACCACGCCTCCCGT
GCTGGACTCCGACGGCTCCTTCGCCCTCTACAGCAAGCTC
ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC
TCATGCTCCGTGATGCATGAGGCTCTGAAATTCCACTACA
CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA MOG-Seldeg-PS
GGACAATTCAGAGTGATAGGACCAGGGTATCCCATCCGG 7 with knobs-into-
GCTTTAGTTGGGGATGAAGCAGAGCTGCCGTGCCGCATC holes and
TCTCCTGGGAAAAATGCCACGGGCATGGAGGTGGGTTGG arginine
TACCGTTCTCCCTTCTCAAGAGTGGTTCACCTCTACCGAA mutations
ATGGCAAGGACCAAGATGCAGAGCAAGCACCTGAATACC
GGGGACGCACAGAGCTTCTGAAAGAGACTATCAGTGAGG
GAAAGGTTACCCTTAGGATTCAGAACGTGAGATTCTCAG
ATGAAGGAGGCTACACCTGCTTCTTCAGAGACCACTCTTA
CCAAGAAGAGGCAGCAATGGAGTTGAAAGTGGAAGATG
GAGGCGGTGGATCAGTTGAGCCCAAATCTTCTGACAAAA
CTCACACATGCCACCGTGCCCAGCACCTGAACTCCTGA
GGGGACCGTCAGTCTTCCTCTTCCCCCCAAAACCCAAGGA
CACCTCATGATCTCCCGGACCCCTGAGGTACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC
AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG
ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT
GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGA
ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
GCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAG
GGCAGCCCCGAGAACCACAGGTGACCACCCTGCCCCCAT
CCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCT
GCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGG
AGTGGGAGAGCAATGGGCAGCCGGAGAACAATACTACAAG
ACCTTCCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCT
CTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCA
GGGGAACGTCTTCTCATGCTCTGTGATGCATGAGGCTCTG
CATAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
GTAAAGGAGGCGGTGGATCAGAGAAACTGGGAAAACCTT
AGTATTCAGTGGATTATGATTTCCAAAATAACCAGCTGCT
GGTAGGGATCATTACAGGCTGCCGAACCTGCCCGCCTTGGA
CATGGGGGGGCACATCTGATCCTTACGTGAAAGTGTTTCTG
CTACCTGATAAGAAGAAGAAATTTGAGACAAAAGTCCAC
CGAAAAACCTTAATCCTGTCTTCAATGAGCAATTTACTT
TCAAGGTACCATACTCGGAATTGGGTGGCAAAACCTTAG
TGATGGCTGTATATGATTTTGATCGTTTCTCTAAGCATGA

CATCATTTGGAAATGGATCCCTATGGAACACAGTGGA
TTTTGGCCATGTAACCTGAGGAATGGCGTGACCTGCAAAG TGCT Fc-Syt1 with
GTTGAGCCCAAATCTTCTGACAAAACCTCACACATGCCAC 9 knobs-into-holes
CGTGCCACGACCTGAACTCCTGAGGGGACCGTCAGTCTT and arginine
CCTCTTCCCCC AAAACCCAAGGACACCCTCATGATCTCC mutations
CGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGC
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGAC
GGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGA
GGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCT
CACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAACAAAGCCCCGCCAGCCCCCAT
CGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGA
ACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCT
GACCAAGAACCAGGTCCACCTGACCTGCCTGGTCAAAGG
CTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAA
TGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGT
GCTGGACTCCGACGGCTCCTTCGCCCTCTACAGCAAGCTC
ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC
TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA
CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAGGAGGCG
GTGGATCAGAGAACTGGGAAAACCTTCAGTATTCACTGG
ATTATGATTTCCAAAATAACCAGCTGCTGGTAGGGATCAT
TCAGGCTGCCGAACCTGCCCGCCTTGGACATGGGGGGCAC
ATCTGATCCTTACGTGAAAGTGTTTCTGCTACCTGATAAG
AAGAAGAAATTTGAGACAAAAGTCCACCGAAAAACCCTT
AATCCTGTCTTCAATGAGCAATTTACTTTCAAGGTACCAT
ACTCGGAATTGGGTGGCAAAACCCTAGTGATGGCTGTAT
ATGATTTTGATCGTTTCTCTAAGCATGACATCATTGGAGA
ATTTAAAGTCCCTATGAACACAGTGGATTTTGGCCATGTA
ACTGAGGAATGGCGTGACCTGCAAAGTGCT MOG-Seldeg-PS
GGACAATTCAGAGTGATAGGACCAGGGTATCCCATCCGG 11 with knobs-into-
GCTTTAGTTGGGGATGAAGCAGAGCTGCCGTGCCGCATC holes,
TCTCCTGGGAAAAATGCCACGGGCATGGAGGTGGGTGG electrostatic
TACCGTTCTCCCTTCTCAAGAGTGGTTCACCTCTACCGAA steering and
ATGGCAAGGACCAAGATGCAGAGCAAGCACCTGAATACC arginine
GGGGACGCACAGAGCTTCTGAAAGAGACTATCAGTGAGG mutations
GAAAGGTTACCCTTAGGATTCAGAACGTGAGATTCTCAG
ATGAAGGAGGCTACACCTGCTTCTTCAGAGACCACTCTTA
CCAAGAAGAGGCAGCAATGGAGTTGAAAGTGGAAGATG
GAGGCGGTGGATCAGTTGAGCCCAAATCTTCTGACAAAA
CTCACACATGCCCACCGTGCCCAGCACCTGAACTCCTGA
GGGGACCGTCAGTCTTCCTCTTCCCCC AAAACCCAAGGA
CACCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC
AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG
ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT
GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGA
ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
GCCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAG
GGCAGCCCCGAGAACCACAGGTGACCACCCTGCCCCCAT

CCCGGGATGAGAACCAAGGTCAGCCTGACCT
GCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGG
AGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACGAC
ACCTTCCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCT
CTACAGCGACCTCACCGTGGACAAGAGCAGGTGGCAGCA
GGGGAACGTCTTCTCATGCTCTGTGATGCATGAGGCTCTG
CATAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGG
GTAAAGGAGGCGGTGGATCAGAGAAACTGGGAAAACCTT
AGTATTCAGTGGATTATGATTTCCAAAATAACCAGCTGCT
GGTAGGGATCATTAGGCTGCCGAAGTGGCCGCCTTGGA
CATGGGGGGGCACATCTGATCCTTACGTGAAAGTGTTTCTG
CTACCTGATAAGAAGAAGAAATTTGAGACAAAAGTCCAC
CGAAAAACCCCTTAATCCTGTCTTCAATGAGCAATTTACTT
TCAAGGTACCACTCTCGGAATTGGGTGGCAAAACCCCTAG
TGATGGCTGTATATGATTTTGATCGTTTCTCTAAGCATGA
CATCATTGGAGAATTTAAAGTCCCTATGAACACAGTGGA
TTTTGGCCATGTAAGTGGGAATGGCGTGACCTGCAAAG TGCT Fc-Syt1 with
GTTGAGCCCAAATCTTCTGACAAAACCTCACACATGCCAC 13 knobs-into-
CGTGCCAGCACCTGAACTCCTGAGGGGACCGTCAGTCTT holes,
CCTCTTCCCCCAAAACCCAAGGACACCCTCATGATCTCC electrostatic
CGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGC steering and
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGAC arginine
GGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA mutations
GGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCT
CACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAACAAAGCCCCGCCAGCCCCCAT
CGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGA
ACCACAGGTGTACACCCTGCCCCCATCCCGGGATAAGCT
GACCAAGAACCAGGTCCACCTGACCTGCCTGGTCAAAGG
CTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAA
TGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGT
GCTGAAGTCCGACGGCTCCTTCGCCCTCTACAGCAAGCTC
ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC
TCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACA
CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAGGAGGCG
GTGGATCAGAGAAACTGGGAAAACCTTCAGTATTCAGTGG
ATTATGATTTCCAAAATAACCAGCTGCTGGTAGGGATCAT
TCAGGCTGCCGAAGTGGCCGCCTTGACATGGGGGGCAC
ATCTGATCCTTACGTGAAAGTGTTTCTGCTACCTGATAAG
AAGAAGAAATTTGAGACAAAAGTCCACCGAAAAACCCCTT
AATCCTGTCTTCAATGAGCAATTTACTTTCAAGGTACCAT
ACTCGGAATTGGGTGGCAAAACCCCTAGTGATGGCTGTAT
ATGATTTTGATCGTTTCTCTAAGCATGACATCATTGGAGA
ATTTAAAGTCCCTATGAACACAGTGGATTTTGGCCATGTA
ACTGAGGAATGGCGTGACCTGCAAAGTGCT MOG-Seldeg-
GAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAG 15 TfR with knobs-
CCCGGCGCCTCCGTGAAGGTGTCCTGCAAGGCCTCCGGCT into-holes
ACACCTTCACCTCCTACTGGATGCACTGGGTGCGGCAGGC mutations
CCCCGGCCAGCGGCTGGAGTGGATCGGCGAGATCAACCC
CACCAACGGCCGGACCAACTACATCGAGAAGTTCAAGTC

CCGGCCAGGTCACCAAGTCCGGCCTCCACCGC
CTACATGGAGCTGTCCTCCCTGCGGTCCGAGGACACCGCC
GTGTACTACTGCGCCCCGGGGCACCCGGGCCTACCACTACT
GGGGCCAGGGCACCATGGTGACCGTGTCTCTCCGCCTCCA
CCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAA
GAGCACCTCTGGGGGGCACAGCGGCCCTGGGCTGCCTGGT
CAAGGACTACTTCCCCGAACCGGTGACGGTGTTCGTGGAA
CTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGC
TGTCTACAGTCTCAGGACTCTACTCCCTCAGCAGCGTG
GTGACTGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTG
GACAAGAAAGTTGAGCCCCAAATCTTGTGACAAAACCTCAC
ACATGCCCACCGTGCCCAGCACCTGAACTCCTGGGGGGA
CCGTCAGTCTTCTCTTCCCCCCAAAACCCAAGGACACCC
TCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGT
GGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTG
GTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAA
GCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGT
CAGCGTCCTTACCGTCCCTGCACCAGGACTGGCTGAATGGC
AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCA
GCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAG
CCCCGAGAACCACAGGTGACCACCCTGCCCCCATCCCGG
GATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTG
GTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGG
GAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCTTC
CCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCTCTACA
GCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGG
AACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACA
ACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTA
AAGGAGGCGGTGGATCAGGACAATTCAGAGTGATAGGAC
CAGGGTATCCCATCCGGGCTTTAGTTGGGGATGAAGCAG
AGCTGCCGTGCCGCATCTCTCCTGGGAAAAATGCCACGG
GCATGGAGGTGGGTTGGTACCGTTCTCCCTTCTCAAGAGT
GGTTCACCTCTACCGAAATGGCAAGGACCAAGATGCAGA
GCAAGCACCTGAATACCGGGGACGCACAGAGCTTCTGAA
AGAGACTATCAGTGAGGGGAAAGGTTACCCTTAGGATTCA
GAACGTGAGATTCTCAGATGAAGGAGGCTACACCTGCTT
CTTCAGAGACCACTCTTACCAAGAAGAGGCAGCAATGGA GTTGAAAGTGGAAGAT
TfR Ab with GAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAG 17 knobs-
into-holes CCCGGCGCCTCCGTGAAGGTGTCCTGCAAGGCCTCCGGCT mutations
ACACCTTCACCTCCTACTGGATGCACTGGGTGCGGCAGGC
CCCCGGCCAGCGGCTGGAGTGGATCGGCGAGATCAACCC
CACCAACGGCCGGACCAACTACATCGAGAAGTTCAAGTC
CCGGGCCACCCTGACCGTGGACAAGTCCGCCTCCACCGC
CTACATGGAGCTGTCCTCCCTGCGGTCCGAGGACACCGCC
GTGTACTACTGCGCCCCGGGGCACCCGGGCCTACCACTACT
GGGGCCAGGGCACCATGGTGACCGTGTCTCTCCGCCTCCA
CCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAA
GAGCACCTCTGGGGGGCACAGCGGCCCTGGGCTGCCTGGT
CAAGGACTACTTCCCCGAACCGGTGACGGTGTTCGTGGAA

CTCAGCGCGGCGGCGCTGCACACCTTCCC
TGTCCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG
GTGACTGTGCCCTCCAGCAGCTTGGGCACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTG
GACAAGAAAGTTGAGCCCAAATCTTGTGACAAAACCTCAC
ACATGCCCACCGTGCCCAGCACCTGAACTCCTGGGGGGA
CCGTCAGTCTTCCTCTTCCCCCAAAACCCAAGGACACCC
TCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGT
GGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTG
GTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAA
GCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGT
CAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGG
CAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCC
AGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCA
GCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCG
GGATGAGCTGACCAAGAACCAGGTCCACCTGACCTGCCT
GGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTG
GGAGAGCAATGGGCAGCCGGAGAACAACCTACAAGACCA
CGCCTCCCGTGCTGGACTCCGACGGCTCCTTCGCCCTCTA
CAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGG
GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA
CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT AAA TfR Ab LC
GACATCCAGATGACCCAGTCCCCCTCCTCCCTGTCCGCCT 19
CCGTGGGCGACCGGGTGACCATCACCTGCCGGGCCTCCG
ACAACCTGTACTCCAACCTGGCCTGGTACCAGCAGAAGC
CCGGCAAGTCCCCCAAGCTGCTGGTGTACGACGCCACCA
ACCTGGCCGACGGCGTGCCCTCCCGGTTCTCCGGCTCCGG
CTCCGGCACC GACTACACCCTGACCATCTCCTCCCTGCAG
CCCGAGGACTTCGCCACCTACTACTGCCAGCACTTCTGGG
GCACCCCCCTGACCTTCGGCCAGGGCACCAAGGTGGAGA
TCAAGACTGTGGCTGCACCATCTGTCTTCATCTTCCCGCC
ATCTGATGAGCAGTTGAAATCTGGAACCTGCCTCTGTTGTG
TGCCTGCTGAATAACTTCTATCCCAGAGAGGCCAAAGTA
CAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCC
CAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGCAC
CTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGA
CTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCA
TCAGGGCCTGAGTTCGCCCGTCACAAAGAGCTTCAACAG GGGAGAGTGT MOG-Seldeg-
GAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAG 21 TfR with knobs-
CCCGGCGCCTCCGTGAAGGTGTCCTGCAAGGCCTCCGGCT into-holes and
ACACCTTCACCTCCTACTGGATGCACTGGGTGCGGCAGGC arginine
CCCCGGCCAGCGGCTGGAGTGGATCGGCGAGATCAACCC mutations
CACCAACGGCCGGACCAACTACATCGAGAAGTTCAAGTC
CCGGGCCACCCTGACCGTGGACAAGTCCGCCTCCACCGC
CTACATGGAGCTGTCCTCCCTGCGGTCCGAGGACACCGCC
GTGTACTACTGCGCCCGGGGCACCCGGGCCTACCACTACT
GGGGCCAGGGCACCATGGTGACCGTGTCTCCGCCTCCA
CCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCA
GAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGT
CAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGAA

CTCAGCGCGCCGCGCTGCACACCTTCCC GGC
TGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG
GTGACTGTGCCCTCCAGCAGCTTGGG CACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTG
GACAAGAAAGTTGAGCCCAAATCTTGTGACAAA ACTCAC
ACATGCCCACCGTGCCCAGCACCTGAACTCCTGAGGGGA
CCGTCAGTCTTCCTCTTCCCCC CAAAACCCAAGGACACCC
TCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGT
GGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTG
GTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAA
GCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGT
CAGCGTCCTTACCGTCCTGCACCAGGACTGGCTGAATGGC
AAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCCGCCCA
GCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAG
CCCCGAGAACCACAGGTGACCACCCTGCCCCCATCCCGG
GATGAGCTGACCAAGAACCAGGTCAGCCTGACCTGCCTG
GTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTGG
GAGAGCAATGGGCAGCCGGAGAACA ACTACAAGACCTTC
CCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCTCTACA
GCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGG
AACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACA
ACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGTA
AAGGAGGCGGTGGATCAGGACAATTCAGAGTGATAGGAC
CAGGGTATCCCATCCGGGCTTTAGTTGGGGATGAAGCAG
AGCTGCCGTGCCGCATCTCTCCTGGGAAAAATGCCACGG
GCATGGAGGTGGGTTGGTACCGTTCTCCCTTCTCAAGAGT
GGTTCACCTCTACCGAAATGGCAAGGACCAAGATGCAGA
GCAAGCACCTGAATACCGGGGACGCACAGAGCTTCTGAA
AGAGACTATCAGTGAGGGAAAGGTTACCCTTAGGATTCA
GAACGTGAGATTCTCAGATGAAGGAGGCTACACCTGCTT
CTTCAGAGACCACTCTTACCAAGAAGAGGCAGCAATGGA GTTGAAAGTGGAAGAT
TfR Ab with GAGGTGCAGCTGGTGCAGTCCGGCGCCGAGGTGAAGAAG 23 knobs-
into-holes CCCGGCGCCTCCGTGAAGGTGTCCTGCAAGGCCTCCGGCT and arginine
ACACCTTCACCTCCTACTGGATGCACTGGGTGCGGCAGGC mutations
CCCCGGCCAGCGGCTGGAGTGGATCGGCGAGATCAACCC
CACCAACGGCCGGACCAACTACATCGAGAAGTTCAAGTC
CCGGGCCACCCTGACCGTGGACAAGTCCGCCTCCACCGC
CTACATGGAGCTGTCCTCCCTGCGGTCCGAGGACACCGCC
GTGTACTACTGCGCCCGGGGCACCCGGGCCTACCACTACT
GGGGCCAGGGCACCATGGTGACCGTGTCTTCCGCCTCCA
CCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAA
GAGCACCTCTGGGGGCACAGCGGCCCTGGGCTGCCTGGT
CAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGGAA
CTCAGGCGCCCTGACCAGCGGCGTGCACACCTTCCCGGC
TGTCTACAGTCCTCAGGACTCTACTCCCTCAGCAGCGTG
GTGACTGTGCCCTCCAGCAGCTTGGG CACCCAGACCTAC
ATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTG
GACAAGAAAGTTGAGCCCAAATCTTGTGACAAA ACTCAC
ACATGCCCACCGTGCCCAGCACCTGAACTCCTGAGGGGA
CCGTCAGTCTTCCTCTTCCCCC CAAAACCCAAGGACACCC

TCATGATCTCCGTCGACCTGAGGTTCACATGCCGTGGTGGT
GGACGTGAGCCACGAAGACCCTGAGGTCAAGTTCAACTG
GTACGTGGACGGCGTGGAGGTGCATAATGCCAAGACAAA
GCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGT
CAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGAATGG
CAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCGCCC
AGCCCCCATCGAGAAAACCATCTCCAAGCCAAAGGGCA
GCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCG
GGATGAGCTGACCAAGAACCAGGTCCACCTGACCTGCCT
GGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGGAGTG
GGAGAGCAATGGGCAGCCGGAGAACAACACTACAAGACCA
CGCCTCCCGTGCTGGACTCCGACGGCTCCTTCGCCCTCTA
CAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGG
GGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCA
CAACCACTACACGCAGAAGAGCCTCTCCCTGTCTCCGGGT AAA HER2-ECD-Fc
ACCCAAGTGTGCACCGGCACAGACATGAAGCTGCGGCTC 25 with MST-HN
CCTGCCAGTCCCGAGACCCACCTGGACATGCTCCGCCAC and knobs-into-
CTCTACCAGGGCTGCCAGGTGGTGCAGGGAAACCTGGAA holes
CTCACCTACCTGCCCACCAATGCCAGCCTGTCCTTCCTGC
AGGATATCCAGGAGGTGCAGGGCTACGTGCTCATCGCTC
ACAACCAAGTGAGGCAGGTCCCCTGCAGAGGCTGCGGA
TTGTGCGAGGCACCCAGCTCTTTGAGGACAACCTATGCCCT
GGCCGTGCTAGACAATGGAGACCCGCTGAACAATACCAC
CCCTGTCACAGGGGCCTCCCCAGGAGGCCTGCGGGAGCT
GCAGCTTCGAAGCCTCACAGAGATCTTGAAAGGAGGGGT
CTTGATCCAGCGGAACCCCCAGCTCTGCTACCAGGACAC
GATTTTGTGGAAGGACATCTTCCACAAGAACAACCAGCT
GGCTCTCACACTGATAGACACCAACCGCTCTCGGGCCTGC
CACCCCTGTTCTCCGATGTGTAAGGGCTCCCGCTGCTGGG
GAGAGAGTTCTGAGGATTGTCAGAGCCTGACGCGCACTG
TCTGTGCCGGTGGCTGTGCCCCGCTGCAAGGGGGCCACTGCC
CACTGACTGCTGCCATGAGCAGTGTGCTGCCGGCTGCAC
GGGCCCCAAGCACTCTGACTGCCTGGCCTGCCTCCACTTC
AACCACAGTGGCATCTGTGAGCTGCACTGCCCAGCCCTG
GTCACCTACAACACAGACACGTTTGAGTCCATGCCCAATC
CCGAGGGGCCGGTATACATTCGGCGCCAGCTGTGTGACTG
CCTGTCCCTACAACCTACCTTTCTACGGACGTGGGATCCTG
CACCTCGTCTGCCCCCTGCACAACCAAGAGGTGACAGC
AGAGGATGGAACACAGCGGTGTGAGAAGTGCAGCAAGC
CCTGTGCCCCGAGTGTGCTATGGTCTGGGCATGGAGCACTT
GCGAGAGGTGAGGGCAGTTACCAGTGCCAATATCCAGGA
GTTTGCTGGCTGCAAGAAGATCTTTGGGAGCCTGGCATT
CTGCCGGAGAGCTTTGATGGGGACCCAGCCTCCAACACT
GCCCCGCTCCAGCCAGAGCAGCTCCAAGTGTTTGAGACT
CTGGAAGAGATCACAGGTTACCTATACATCTCAGCATGG
CCGGACAGCCTGCCTGACCTCAGCGTCTTCCAGAACCTGC
AAGTAATCCGGGGACGAATTCTGCACAATGGCGCCTACT
CGCTGACCCTGCAAGGGCTGGGCATCAGCTGGCTGGGGC
TGCGCTCACTGAGGGAACTGGGCAGTGGACTGGCCCTCA
TCCACCATAACACCCACCTCTGCTTCGTGCACACGGTGCC

CTGGGACAGCTTCTGCTC
CACACTGCCAACCGGCCAGAGGACGAGTGTGTGGGCGAG
GGCCTGGCCTGCCACCAGCTGTGCGCCCGAGGGCACTGC
TGGGGTCCAGGGCCACCCAGTGTGTCAACTGCAGCCAG
TTCCTTCGGGGCCAGGAGTGCGTGGAGGAATGCCGAGTA
CTGCAGGGGCTCCCCAGGGAGTATGTGAATGCCAGGCAC
TGTTTGCCGTGCCACCCTGAGTGTGAGCCCCAGAATGGCT
CAGTGACCTGTTTTGGACCGGAGGCTGACCAGTGTGTGG
CCTGTGCCCACTATAAGGACCCTCCCTTCTGCGTGGCCCG
CTGCCCCAGCGGTGTGAAACCTGACCTCTCCTACATGCCC
ATCTGGAAGTTTCCAGATGAGGAGGGCGCATGCCAGCCT
TGCCCCATCAACTGCACCCACTCCTGTGTGGACCTGGATG
ACAAGGGCTGCCCCGCCGAGCAGAGAGCCAGCCCTCTGA
CGATTGAAGGCCGCATGGATCCCAAATCTTCTGACAAA
CTCACACATGCCACCGTGCCCAGCACCTGAACTCCTGG
GGGGACCGTCAGTCTTCCTCTTCCCCCCAAAACCCAAGGA
CACCTCTACATCACTCGGGAACCTGAGGTCACATGCGTG
GTGGTGGACGTGAGCCACGAAGACCCTGAGGTCAAGTTC
AACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG
ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGT
GTGGTCAGCGTCCTCACCGTCCTGCACCAGGACTGGCTGA
ATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC
TCCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAAG
GGCAGCCCCGAGAACCACAGGTGACCACCCTGCCCCCAT
CCCGGGATGAGCTGACCAAGAACCAGGTCAGCCTGACCT
GCCTGGTCAAAGGCTTCTATCCCAGCGACATCGCCGTGG
AGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTACAAG
ACCTTCCCTCCCGTGCTGGACTCCGACGGCTCCTTCTTCCT
CTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCA
GGGGAACGTCTTCTCATGCTCTGTGATGCATGAGGCTCTG
AAATTCCACTACACGCAGAAGAGCCTCTCCCTGTCTCCTG GTAAA Fc with MST-
GTTGAGCCCAAATCTTCTGACAAAACCTCACACATGCCAC 27 HN and knobs-
CGTGCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTT into-holes
CCTCTTCCCCCCAAAACCCAAGGACACCCTCTACATCACT mutations
CGGGAACCTGAGGTCACATGCGTGGTGGTGGACGTGAGC
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGAC
GGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA
GGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCT
CACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCCAT
CGAGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGA
ACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCT
GACCAAGAACCAGGTCCACCTGACCTGCCTGGTCAAAGG
CTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAA
TGGGCAGCCGGAGAACAACACTACAAGACCACGCCTCCCGT
GCTGGACTCCGACGGCTCCTTCGCCCTCTACAGCAAGCTC
ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC
TCATGCTCCGTGATGCATGAGGCTCTGAAATTCCACTACA
CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAA PSMA-Seldeg
GTTGAGCCCAAATCTTCTGACAAAACCTCACACATGCCAC 29 with MST-HN,

CGTGCCCGCCAGGAGGACCCGTCAGTCTT knobs-into-holes
CCTCTTCCCCC AAAACCCAAGGACACCCTCTACATCACT and arginine
CGGGAACCTGAGGTCACATGCGTGGTGGTGGACGTGAGC mutations
CACGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGAC
GGCGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGA
GGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCT
CACCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTA
CAAGTGCAAGGTCTCCAACAAAGCCCCGCCAGCCCCCAT
CGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGA
ACCACAGGTGACCACCCTGCCCCCATCCCGGGATGAGCT
GACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGG
CTTCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAA
TGGGCAGCCGGAGAACAACTACAAGACCTTCCCTCCCGT
GCTGGACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTC
ACCGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTC
TCATGCTCTGTGATGCATGAGGCTCTGAAATTCCACTACA
CGCAGAAGAGCCTCTCCCTGTCTCCGGGTAAAGGAGGCG
GTGGATCAAAATCCTCCAATGAAGCTACTAACATTACTCC
AAAGCATAATATGAAAGCATTTTTGGATGAATTGAAAGC
TGAGAACATCAAGAAGTTCTTATATAATTTTACACAGATA
CCACATTTAGCAGGAACAGAACAAAACCTTTCAGCTTGCA
AAGCAAATTCAATCCCAGTGGAAAGAATTTGGCCTGGAT
TCTGTTGAGCTAGCACATTATGATGTCCTGTTGTCCTACC
CAAATAAGACTCATCCCAACTACATCTCAATAATTAATGA
AGATGGAAATGAGATTTTCAACACATCATTATTTGAACCA
CCTCCTCCAGGATATGAAAATGTTTCGGATATTGTACCAC
CTTTCAGTGCTTTCTCTCCTCAAGGAATGCCAGAGGGCGA
TCTAGTGTATGTAACTATGCACGAACTGAAGACTTCTTT
AAATTGGAACGGGACATGAAAATCAATTGCTCTGGGAAA
ATTGTAATTGCCAGATATGGGAAAGTTTTTCAGAGGAAAT
AAGGTAAAAATGCCCAGCTGGCAGGGGCCAAAGGAGTC
ATTCTCTACTCCGACCCTGCTGACTACTTTGCTCCTGGGG
TGAAGTCCTATCCAGATGGTTGGAATCTTCCTGGAGGTGG
TGTCCAGCGTGGAAATATCCTAAATCTGAATGGTGCAGG
AGACCCTCTCACACCAGGTTACCCAGCAAATGAATATGC
TTATAGGCGTGGAAATTGCAGAGGCTGTTGGTCTTCCAAGT
ATTCCTGTTCAATCAATTGGATACTATGATGCACAGAAGC
TCCTAGAAAAAATGGGTGGCTCAGCACACCAGATAGCA
GCTGGAGAGGAAGTCTCAAAGTGCCCTACAATGTTGGAC
CTGGCTTTACTGGAACTTTTCTACACAAAAAGTCAAGAT
GCACATCCACTCTACCAATGAAGTGACAAGAATTTACAA
TGTGATAGGTACTCTCAGAGGAGCAGTGGAACCAGACAG
ATATGTCATTCTGGGAGGTCACCGGGACTCATGGGTGTTT
GGTGGTATTGACCCTCAGAGTGGAGCAGCTGTTGTTCATG
AAATTGTGAGGAGCTTTGGAACACTGAAAAAGGAAGGGT
GGAGACCTAGAAGAACAATTTTGTGTTGCAAGCTGGGATG
CAGAAGAATTTGGTCTTCTTGGTTCTACTGAGTGGGCAGA
GGAGAATTCAAGACTCCTTCAAGAGCGTGGCGTGGCTTA
TATTAATGCTGACTCATCTATAGAAGGAACTACACTCTG
AGAGTTGATTGTACACCGCTGATGTACAGCTTGGTACACA

ACCTAACAAAGCCCTGATGAAGGCTTTG
AAGGCAAATCTCTTTATGAAAGTTGGACTAAAAAAGTC
CTTCCCCAGAGTTCAGTGGCATGCCCAGGATAAGCAAAT
TGGGATCTGGAAATGATTTTGAGGTGTTCTTCCAACGACT
TGGAATTGCTTCAGGCAGAGCACGGTATACTAAAAATTG
GGAAACAAACAAATTCAGCGGCTATCCACTGTATCACAG
TGTCTATGAAACATATGAGTTGGTGGAAAAGTTTTATGAT
CCAATGTTTAAATATCACCTCACTGTGGCCCAGGTTCGAG
GAGGGATGGTGTGTTGAGCTAGCCAATTCCATAGTGCTCCC
TTTTGATTGTCGAGATTATGCTGTAGTTTTTAAGAAAGTAT
GCTGACAAAATCTACAGTATTTCTATGAAACATCCACAG
GAAATGAAGACATACAGTGTATCATTTGATTCACTTTTTT
CTGCAGTAAAGAATTTTACAGAAATTGCTTCCAAGTTCAG
TGAGAGACTCCAGGACTTTGACAAAAGCAACCCAATAGT
ATTAAGAATGATGAATGATCAACTCATGTTTCTGGAAAG
AGCATTTATTGATCCATTAGGGTTACCAGACAGGCCTTTT
TATAGGCATGTCATCTATGCTCCAAGCAGCCACAACAAG
TATGCAGGGGAGTCATTCCCAGGAATTTATGATGCTCTGT
TTGATATTGAAAGCAAAGTGGACCCTTCCAAGGCCTGGG
GAGAAGTGAAGAGACAGATTTATGTTGCAGCCTTCACAG
TGCAGGCAGCTGCAGAGACTTTGAGTGAAGTAGCC GAD65-Seldeg
ATGGCATCTCCGGGCTCTGGCTTTTGGTCTTTCGGGTCGG 31 with MST-HN,
AAGATGGCTCTGGGGATTCCGAGAATCCCGGCACAGCGC knobs-into-holes
GAGCCTGGTGCCAAGTGGCTCAGAAGTTCACGGGCGGCA and arginine
TCGGAACAAACTGTGCGCCCTGCTCTACGGAGACGCCG mutations
AGAAGCCGGCGGAGAGCGGCGGGAGCCAACCCCCGCGG
GCCGCCGCCCGGAAGGCCGCTGCGCCTGCGACCAGAAG
CCCTGCAGCTGCTCCAAAGTGGATGTCAACTACGCGTTTC
TCCATGCAACAGACCTGCTGCCGGCGTGTGATGGAGAAA
GGCCCACTTTGGCGTTTCTGCAAGATGTTATGAACATTTT
ACTTCAGTATGTGGTGAAAAGTTTCGATAGATCAACCAA
AGTGATTGATTTCCATTATCCTAATGAGCTTCTCCAAGAA
TATAATTGGGAATTGGCAGACCAACCACAAAATTTGGAG
GAAATTTTGATGCATTGCCAAACAACCTCTAAAATATGCA
ATTAACACAGGGCATCCTAGATACTTCAATCAACTTTCTA
CTGGTTTGGATATGGTTGGATTAGCAGCAGACTGGCTGAC
ATCAACAGCAAATACTAACATGTTACCTATGAAATTGCT
CCAGTATTTGTGCTTTTGGAAATATGTCACACTAAAGAAAA
TGAGAGAAATCATTGGCTGGCCAGGGGGCTCTGGCGATG
GGATATTTTCTCCCGGTGGCGCCATATCTAACATGTATGC
CATGATGATCGCACGCTTTAAGATGTTCCCAGAAGTCAA
GGAGAAAGGAATGGCTGCTCTTCCCAGGCTCATTGCCTTC
ACGTCTGAACATAGTCATTTTTCTCTCAAGAAGGGAGCTG
CAGCCTTAGGGATTGGAACAGACAGCGTGATTCTGATTA
AATGTGATGAGAGAGGGAAAATGATTCCATCTGATCTTG
AAAGAAGGATTCTTGAAGCCAAACAGAAAGGGTTTGTTT
CTTTCCTCGTGAGTGCCACAGCTGGAACCAACCGTGTACGG
AGCATTTGACCCCCTCTTAGCTGTCGCTGACATTTGCAAA
AAGTATAAGATCTGGATGCATGTGGATGCAGCTTGGGGT
GGGGGATTACTGATGTCCCGAAAACACAAGTGGAAACTG

AGTGGCGTGGAGGCAACTCTGTGACGTGGAATCCA
CACAAGATGATGGGAGTCCCTTTGCAGTGCTCTGCTCTCC
TGGTTAGAGAAGAGGGATTGATGCAGAATTGCAACCAAA
TGCATGCCTCCTACCTCTTTTCAGCAAGATAAACATTATGA
CCTGTCCTATGACACTGGAGACAAGGCCTTACAGTGCGG
ACGCCACGTTGATGTTTTTAAACTATGGCTGATGTGGAGG
GCAAAGGGGACTACCGGGTTTGAAGCGCATGTTGATAAA
TGTTTGGAGTTGGCAGAGTATTTATAACAATCATAAAAA
ACCGAGAAGGATATGAGATGGTGTGTTTATGATGGGAAGCCTC
AGCACACAAATGTCTGCTTCTGGTACATTCTCCAAGCTT
GCGTACTCTGGAAGACAATGAAGAGAGAATGAGTCGCCT
CTCGAAGGTGGCTCCAGTGATTAAAGCCAGAATGATGGA
GTATGGAACCACAATGGTCAGCTACCAACCCTTGGGAGA
CAAGGTCAATTTCTTCCGCATGGTCATCTCAAACCCAGCG
GCAACTCACCAAGACATTGACTTCCTGATTGAAGAAATA
GAACGCCTTGGACAAGATTTAGGAGGCGGTGGATCAGTT
GAGCCCAAATCTTCTGACAAAACCTCACACATGCCCACCG
TGCCCAGCACCTGAACTCCTGAGGGGACCGTCAGTCTTCC
TCTTCCCCCCCCAAAACCCAAGGACACCCTCTACATCACTCG
GGAACCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCA
CGAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGG
CGTGGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGG
AGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCA
CCGTCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACA
AGTGCAAGGTCTCCAACAAAGCCCGCCCAGCCCCATCG
AGAAAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAAC
CACAGGTGACCACCCTGCCCCCATCCCGGGATGAGCTGA
CCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCT
TCTATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATG
GGCAGCCGGAGAACAACACTACAAGACCTTCCCTCCCGTGC
TGGACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCAC
CGTGGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTC
ATGCTCTGTGATGCATGAGGCTCTGAAATTCCACTACACG
CAGAAGAGCCTCTCCCTGTCTCCGGGTAAA AQP4-Seldeg
ATGAGTGACAGACCCACAGCAAGGCGGTGGGGTAAGTGT 33 with MST-HN,
GGACCTTTGTGTACCAGAGAGAACATCATGGTGGCTTTCA knobs-into-holes
AAGGGGTCTGGACTCAAGCTTTCTGGAAAGCAGTCACAG and arginine
CGGAATTTCTGGCCATGCTTATTTTTGTTCTCCTCAGCCTG mutations
GGATCCACCATCAACTGGGGTGGAAACAGAAAAGCCTTTA
CCGGTCGACATGGTTCTCATCTCCCTTTGCTTTGGACTCA
GCATTGCAACCATGGTGCAGTGCTTTGGCCATATCAGCGG
TGGCCACATCAACCCTGCAGTGACTGTGGCCATGGTGTGC
ACCAGGAAGATCAGCATCGCCAAGTCTGTCTTCTACATCG
CAGCCCAGTGCCTGGGGGCCATCATTGGAGCAGGAATCC
TCTATCTGGTCACACCTCCCAGTGTGGTGGGAGGCCTGGG
AGTCACCATGGTTCATGGAAATCTTACCGCTGGTCATGGT
CTCCTGGTTGAGTTGATAATCACATTTCAATTGGTGTTTA
CTATCTTTGCCAGCTGTGATTCCAAACGGACTGATGTCAC
TGGCTCAATAGCTTTAGCAATTGGATTTTCTGTTGCAATT
GGACATTTATTTGCAATCAATTATACTGGTGCCAGCATGA

ATCCCGCCGCTCCGAGTTATCATGGGAAA
TTGGGAAAACCATTTGGATATATTGGGTTGGGCCCATCATA
GGAGCTGTCCTCGCTGGTGGCCTTTATGAGTATGTCTTCT
GTCCAGATGTTGAATTCAAACGTCGTTTTAAAGAAGCCTT
CAGCAAAGCTGCCCAGCAAACAAAAGGAAGCTACATGG
AGGTGGAGGACAACAGGAGTCAGGTAGAGACGGATGAC
CTGATTCTAAAACCTGGAGTGGTGCATGTGATTGACGTTG
ACCGGGGAGAGGAGAAGAAGGGGAAAGACCAATCTGGA
GAGGTATTGTCTTCAGTAGGAGGCGGTGGATCAGTTGAG
CCCAAATCTTCTGACAAAACCTCACACATGCCCACCGTGCC
CAGCACCTGAACTCCTGAGGGGACCGTCAGTCTTCCTCTT
CCCCC AAAACCCAAGGACACCCTCTACATCACTCGGGA
ACCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGA
AGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGT
GGAGGTGCATAATGCCAAGACAAAGCCGCGGGAGGAGC
AGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCG
TCCTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGT
GCAAGGTCTCCAACAAAGCCCGCCCAGCCCCCATCGAGA
AAACCATCTCCAAGCCAAAGGGCAGCCCCGAGAACCAC
AGGTGACCACCCTGCCCCCATCCCGGGATGAGCTGACCA
AGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGCTTCT
ATCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGC
AGCCGGAGAACAACACTACAAGACCTTCCCTCCCGTGCTGG
ACTCCGACGGCTCCTTCTTCCTCTACAGCAAGCTCACCGT
GGACAAGAGCAGGTGGCAGCAGGGGAACGTCTTCTCATG
CTCTGTGATGCATGAGGCTCTGAAATTCCACTACACGCAG
AAGAGCCTCTCCCTGTCTCCGGGTAAA

(136) TABLE-US-00002 TABLE 2 Amino acid sequences of exemplary proteins.

Protein Amino acid sequence SEQ ID NO: MOG-Seldeg
GQFRVIGPGYPIRALVGDEAELPCRISPGKNATGMEVGWYRS 2 with MST-HN,
PFSRVVHLYRNGKDQDAEQAPEYRGRTELLKETISEGKVTLRI knobs-into-
QNVRFSDGGYTCFFRDHSYQEEAAMELKVEDGGGGSVEPK holes and
SSDKTHTCPPCPAPELLRGPVFLFPPKPKDTLYITREPEVTCV arginine
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV mutations
VSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPR
EPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP
ENNYKTFPPVLDSDGSFFLYSKLTVDKSRWQQGNVFCSSVMH EALKFHYTQKSLSLSPGK
HER2-Seldeg TQVCTGTDMLRLPASPEHLDMRLHLYQGCQVQGNLELT 4 with
MST-HN, YLPTNASLSFLQDIQEVQGYVLIAHNQVRQVPLQRLRIVRGTQ knobs-into-
LFEDNYALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEIL holes and
KGGVLIQRNPQLCYQDTILWKDIFHKNNQLALTLIDTNRSRAC arginine
HPCSPMCKGSRCWGESSEDCQSLTRTVCAAGGCARCKGPLPTD mutations
CCHEQCAAGCTGPKHSDCLACLHFNHSGICELHCPALVTYNT
DTFESMPNPEGRYTFGASCVTACPYNYLSTDVGSC TLVCPLH
NQEVTAEDGTQRCEKCSKPCARVCYGLGMEHLREVRAV TSA
NIQEFAGCKKIFGSLAFLPESFDGDPASNTAPLQPEQLQVFETL
EETGYLYISAWPDSL PDL SVFQNLQVIRGRILHNGAYSLTLQG
LGISWLGLRSLRELGSGLALIHNTHLCFVHTVPWDQLFRNP
HQALLHTANRPEDECVGEGLACHQLCARGHCWGPPTQCVN
CSQFLRGQECVEECRVLQGLPREYVNARHCLPCHPECQPQNG

SVTCFGEADQCVCAPFCVARCPGKPDLSYMPIW
KFPDEEGACQPCPINCTHSCVDLDDKGCPAEQRASPLTIEGRM
DPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLYITREPEV
TCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST
YRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAK
GQPREPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAVEWES
NGQPENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS
SVMHEALKFHYTQKSLSLSPGK Fc with MST-
VEPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLYITREPE 6 HN, knobs-
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS into-holes and
TYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAK arginine
GQPREPQVYTLPPSRDELTKNQVHSLTCLVKGFYPSDIAVEWE mutations
SNGQPENNYKTTPPVLDSDGSFALYSKLTVDKSRWQQGNVFS
CSVMHEALKFHYTQKSLSLSPGK MOG-Seldeg-
GQFRVIGPGYPIRALVGDEAELPCRISPGKNATGMEVGWYRS 8 PS with knobs-
PFSRVVHLYRNGKDQDAEQAPEYRGRTELLKETISEGKVTLRI into-holes and
QNVRFSDGGGYTCFFRDHSYQEEAAMELKVEDGGGGGSVEPK arginine
SSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCV mutations
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV
VSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPR
EPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP
ENNYKTTPPVLDSDGSFFLYSKLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPGKGGGGSEKLGKLQYSLDYDFQNNQ
LLVGIIQAAELPALDMGGTSDPYVKVFLLPDKKKKFETKVHR
KTLNPVFNEQFTFKVPYSELGGKTLVMAVYDFDRFSKHDIIGE
FKVPMNTVDFGHVTEEWRDLQSA Fc-Sytl with
VEPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPE 10 knobs-into-
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS holes and
TYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAK arginine
GQPREPQVYTLPPSRDELTKNQVHSLTCLVKGFYPSDIAVEWE mutations
SNGQPENNYKTTPPVLDSDGSFALYSKLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPGKGGGGSEKLGKLQYSLDYD
FQNNQLLVGIIQAAELPALDMGGTSDPYVKVFLLPDKKKKFE
TKVHRKTLNPVFNEQFTFKVPYSELGGKTLVMAVYDFDRFSK
HDIIGEFKVPMNTVDFGHVTEEWRDLQSA MOG-Seldeg-
GQFRVIGPGYPIRALVGDEAELPCRISPGKNATGMEVGWYRS 12 PS with knobs-
PFSRVVHLYRNGKDQDAEQAPEYRGRTELLKETISEGKVTLRI into-holes,
QNVRFSDGGGYTCFFRDHSYQEEAAMELKVEDGGGGGSVEPK electrostatic
SSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPEVTCV steering and
VVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRV arginine
VSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAKGQPR mutations
EPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQP
ENNYDTFPPVLDSDGSFFLYSDLTVDKSRWQQGNVFS
CSVMHEALHNHYTQKSLSLSPGKGGGGSEKLGKLQYSLDYDFQNNQ
LLVGIIQAAELPALDMGGTSDPYVKVFLLPDKKKKFE
KTLNPVFNEQFTFKVPYSELGGKTLVMAVYDFDRFSKHDIIGE
FKVPMNTVDFGHVTEEWRDLQSA Fc-Sytl with
VEPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMISRTPE 14 knobs-into-
VTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS holes,
TYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAIEKTISKAK electrostatic

GQPREPQVYTLPPSRDELTKNQLVKGFLYPSDIAVEWE steering and
SNGQPENNYKTTPPVLKSDGSFALYSKLTVDKSRWQQGNVFS arginine
CSVMHEALHNHYTQKSLSLSPGKGGGGSEKLGKLQYSLDYD mutations
FQNNQLLVGIIQAAELPALDMGGTSDPYVKVFLLPDKKKKFE
TKVHRKTLNPFVNEQFTFKVPYSELGGKTLVMAVYDFDRFSK
HDIIGEFKVPMTVDVDFGHVTEEWRLQSA MOG-Seldeg-
EVQLVQSGAEVKKPGASVKVSCKASGYTFTSYWMHWVRQA 16 TfR with
PGQRLEWIGEINPTNGRTNYIEKFKSRATLTVDKSASTAYMEL knobs-into-
SSLRSED TAVYYCARGTRAYHYWGQGTMTVTVSSASTKGPSV holes
FPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV mutations
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTK
VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI
SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAV
EWESNGQPENNYKTFFPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMEALHNHYTQKSLSLSPGKGGGGSGQFRVIGPGY
PIRALVGDEAEAPCRISPGKNATGMEVGWYRSPFSRVVHLYR
NGKDQDAEQAPEYRGRTSELLKETISEGKVTLRIQNVRFSDG
GYTCFFRDHSYQEEAAMELKVED TfR Ab with
EVQLVQSGAEVKKPGASVKVSCKASGYTFTSYWMHWVRQA 18 knobs-into-
PGQRLEWIGEINPTNGRTNYIEKFKSRATLTVDKSASTAYMEL holes
SSLRSED TAVYYCARGTRAYHYWGQGTMTVTVSSASTKGPSV mutations
FPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTK
VDKKVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI
SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTI
SKAKGQPREPQVYTLPPSRDELTKNQVHLTCLVKGFYPSDIA
VEWESNGQPENNYKTTPPVLDSDGSFALYSKLTVDKSRWQQ
GNVFSCSVMEALHNHYTQKSLSLSPGK TfR Ab LC
DIQMTQSPSSLSASVGDRVTITCRASDNLYSNLAWYQQKPGK 20
SPKLLVYDADTGLADGVPSRFSGSGSGTDYTLTISSLQPEDFAT
YYCQHFHWGTPLTFGQGTKVEIKTVAAPSVFIFPPSDEQLKSGT
ASVVCCLNNFYPPREAKVQWKVDNALQSGNSQESVTEQDSKD
STYLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC MOG-Seldeg-
EVQLVQSGAEVKKPGASVKVSCKASGYTFTSYWMHWVRQA 22 TfR with
PGQRLEWIGEINPTNGRTNYIEKFKSRATLTVDKSASTAYMEL knobs-into-
SSLRSED TAVYYCARGTRAYHYWGQGTMTVTVSSASTKGPSV holes and
FPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV arginine
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTK mutations
VDKKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMI
SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAPIEKTI
SKAKGQPREPQVTTLPSPRDELTKNQVSLTCLVKGFYPSDIAV
EWESNGQPENNYKTFFPVLDSDGSFFLYSKLTVDKSRWQQG
NVFSCSVMEALHNHYTQKSLSLSPGKGGGGSGQFRVIGPGY
PIRALVGDEAEAPCRISPGKNATGMEVGWYRSPFSRVVHLYR
NGKDQDAEQAPEYRGRTSELLKETISEGKVTLRIQNVRFSDG
GYTCFFRDHSYQEEAAMELKVED TfR Ab with

EVQLVQSGAEVQKASGYSCKASGYTFTSYWMHWVRQA 24 knobs-into-
PGQRLEWIGEINPTNGRTNYIEKFKSRA TLTVDKSASTAYMEL holes and
SSLRSED TAVYYCARGTRAYHYWGQGTMTVTVSSASTKGPSV arginine
FPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWNSGALTSGV mutations
HTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTK
VDKKVEPKSCDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLMI
SRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREE
QYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAPIEKTI
SKAKGQPREPQVYTLPPSRDELTKNQVH L TCLVKGFYPSDIA
VEWESNGQPENNYKTTPPVLDSDGSFALYSKLTVDKSRWQQ
GNVFSCSVMHEALHNHYTQKSLSLSPGK HER2-ECD-Fc
TQVCTGTDMKLR L PASPETHLDMLRHLYQGCQVVQGNLELT 26 with MST-HN
YLPTNASLSFLQDIQEVQGYVLI AHNQVRQVPLQRLRIVRG TQ and knobs-
LFEDNYALAVLDNGDPLNNTTPVTGASPGGLRELQLRSLTEIL into-holes
KGGVLIQRNPQLCYQDTILWKDIFHKNNQLALT LIDTNRSRAC
HPCSPMCKGSR CWGESSEDCQSLTRTV CAGGCARCKGPLPTD
CCHEQCAAGCTGPKHSDCLAC LHFNHSGICELHCPALVTYNT
DTFESMPNPEG RYTFGASCVTACPYNYLSTDVGSCTLV CPLH
NQEVT AEDGTQRCEKCSKPCARVCYGLGMEHLREVRAV TSA
NIQEFAGCKKIFGSLAFLPESFDGD PASNTAPLQPEQLQVFETL
EETGYLYISAWPDSL PDL SVFQNLQVIRGRILHNGAYS LTLQG
LGISWLGLRSLRELGSG LALIHNTHLCFVHTVPWDQLFRNP
HQALLHTANRPEDECVGEGLACHQLCARGHCWGPGPTQCVN
CSQFLRGQECVEECRVLQGLPREYVNARHCLPCHPECQPQNG
SVTCFGPEADQC VACA HYKDP PFCVARCPSGVKPDLSYMPIW
KFPDEEGACQPCPINCTHSCVDLDDKGC PAEQRASPLTIEGRM
DPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLYITREPEV
TCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNST
YRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKG
QPREPQVTTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESN
GQPENNYKTFFPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCS
VMHEALKFHYTQKSLSLSPGK Fc with MST-
VEPKSSDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLYITREPE 28 HN and knobs-
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS into-holes
TYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK mutations
GQPREPQVYTLPPSRDELTKNQVH L TCLVKGFYPSDIAVEWE
SNGQPENNYKTTPPVLDSDGSFALYSKLTVDKSRWQQGNVFS
CSVMHEALKFHYTQKSLSLSPGK PSMA-Seldeg
VEPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPKDTLYITREPE 30 with MST-HN,
VTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNS knobs-into-
TYRVVSVLTVLHQDWLNGKEYKCKVSNKARPAPIEKTISKAK holes and
GQPREPQVTTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWES arginine
NGQPENNYKTFFPPVLDSDGSFFLYSKLTVDKSRWQQGNVFSC mutations
SVMHEALKFHYTQKSLSLSPGKGGGGSKSSNEATNITPKHNM
KAFLDELKAENIKKFLYNFTQIPHLAGTEQNFQLAKQIQSQW
KEFGLDSVELAHYDVLLSYPNKTHPNYISIINEDGNEIFNTSLF
EPPPPGYENVSDIVPPFSAFSPQGMPEGDLVYVNYARTEDFFK
LERDMKINCSGKIV IARYGKVFRGNKVKNAQLAGAKGVILYS
DPADYFAPGVKSYPDGWNLPGGGVQRG N ILNLNGAGDPLTP
GYPAN EYAYRRGIAEAVGLPSIPVHPIGYYDAQKLLEKMGG S

APPDSSWSLKVPPYVNFSTQKVKMHIHSTNEVTR
 IYNVIGTLRGAVEPDRYVILGGHRDSWVFGGIDPQSGAAVVH
 EIVRSFGTLKKEGWRPRRTILFASWDAEEFGLLGSTEWAEENS
 RLLQERGVAYINADSSIEGNYTLRVDCTPLMYSLVHNLTKEL
 KSPDEGFEGKSLYESWTKKSPSPEFSGMPRISKLGSGNDFEVF
 FQRLGIASGRARYTKNWETNKFSGYPLYHSVYETYELVEKFY
 DPMFKYHLTVAQVRGGMVFELANSIVLPFDCRDYAVVLRKY
 ADKIYSISMKHPQEMKTYSVSFDSLFSVKNFTEIASKFSERLQ
 DFDKSNPIVLRMMNDQLMFLERAFIDPLGLPDRPFYRHVIYAP
 SSHNKYAGESFPGIYDALFDIESKVDPSKAWGEVKRQIYVAAF TVQAAAETLSEVA
 GAD65- MASPSSGFWSEFGSEDGSGDSENPGTARAWCQVAQKFTGGIG 32 Seldeg with
 NKLCALLYGDAEKPAESGGSQPPRAAARKAACACDQKPCSC MST-HN,
 SKVDVNYAFLHATDLLPACDGERPTLAFLQDVMNILLQYVV knobs-into-
 KSFDRSTKVIDFHYPNELLQEYNWELADQPQNLEEILMHCQT holes and
 TLKYAIKTGHPRYFNQLSTGLDMVGLAADWLTSTANTNMFT arginine
 YEIAPVFVLLLEYVTLKKMREIIGWPGSGDGIFSPGGAISNMY mutations
 AMMIARFKMFPEVKEKGMAALPRLIAFTSEHSHFSLKKGAAA
 LGIGTDSVILIKCDERGKMIPSDLERRILEAKQKGFVPFLVSAT
 AGTTVYGAFDPLLAVADICKKYKIWMHVDAAWGGGLMSR
 KHKWKLSGVERANSVTWNPHKMMGVPLQCSALLVREEGLM
 QNCNQMHASYLFQQDKHYDLSYDTGDKALQCGRHVDVFKL
 WLMWRAKGTGFEAHVDKCLELAEYLYNIIKNREGYEMVFD
 GKPQHTNVCFWYIPPSLRTLEDNEERMSRLSKVAPVIKARMM
 EYGTTMVSYQPLGDKVNFFRMVISNPAATHQDIDFLIEEIERL
 GQDLGGGGSVEPKSSDKTHTCPPCPAPELLRGPSVFLFPPKPK
 DTLYITREPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKT
 KPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKARPA
 PIEKTISKAKGQPREPQVTTTLPPSRDELTKNQVSLTCLVKGFYP
 SDIAVEWESNGQPENNYKTFFPPVLDSDGSFFLYSKLTVDKSR
 WQQGNVFSCSVMHEALKFHYTQKSLSLSPGK AQP4-Seldeg
 MSDRPTARRWGKCGPLCTRENIMVAFKGVWTQAFWKAVTA 34 with MST-HN,
 EFLAMLIFVLLSLGSTINWGGTEKPLPVDMLISLCFGLSIATM knobs-into-
 VQCFGHISGGHINPAVTVMVCTRKISIAKSVFYIAAQCLGAI holes and
 GAGILYLVTPPSVVGGLGVTMVHGNLTAGHGALLVELIITFQLV arginine
 FTIFASCDKRTDVTGSIALAIGFSVAIGHLFAINYTGASMNPA mutations
 RSFGPAVIMGNWENHWIYWVGPIIGAVLAGGLYEYVFCPDVE
 FKRRFKEAFSKAAQQTGKSYMEVEDNRSQVETDDLILKPGVV
 HVIDVDRGEEKKGKDQSGEVLSSVGGGGSVEPKSSDKTHTCP
 PCPAPELLRGPSVFLFPPKPKDTLYITREPEVTCVVVDVSHEDP
 EVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ
 DWLNGKEYKCKVSNKARPAPIEKTISKAKGQPREPQVTTTLPPS
 RDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTFFP
 VLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALKFHYT QKSLSLSPGK

(137) The DNA sequence of SEQ ID NO:1 is of a polynucleotide encoding an exemplary MOG-Seldeg fusion protein (SEQ ID NO:2) having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations. The exemplary DNA and amino acid sequences of the MOG respectively of SEQ ID NO: 1 and 2 are of mouse origin.

(138) In particular, the amino acid sequence of the exemplary MOG-Seldeg of SEQ TD NO: 2 forms a fusion protein having, in order from N- to C-terminus, residues 1-117 of mouse (in) MOG, a first linker at residues 118-122, an immunoglobulin hinge (human IgG1-derived) at residues 123-

138, an immunoglobulin CH2 domain (human IgG1-derived) at residues 139-248, an immunoglobulin CH3 domain (human IgG1-derived) at residues 249-355. The exemplary MOG-Seldeg of SEQ ID NO:2 has mutations that increase FcRn binding at residues 160, 162, 164, 341 and 342, arginine mutations at residues 144 and 236, and 'knobs-into-holes' mutations at residues 257 and 302. The cysteine residue (128) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:2 are those of the protein sequence, and do not refer to the EU numbering convention.

(139) The DNA sequence of SEQ ID NO:3 is of a polynucleotide encoding an exemplary HER2-Seldeg fusion protein (SEQ ID NO:4) having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations.

(140) In particular, the amino acid sequence of the exemplary HER2-Seldeg of SEQ ID NO: 4 forms a fusion protein having, in order from N- to C-terminus, residues 1-630 of HER2, a first linker at residues 631-636, an immunoglobulin hinge (human IgG1-derived) at residues 637-650, an immunoglobulin CH2 domain (human IgG1-derived) at residues 651-760, an immunoglobulin CH3 domain (human IgG1-derived) at residues 761-867. The exemplary HER2-Seldeg of SEQ ID NO:4 has mutations that increase FcRn binding at residues 672, 674, 676, 853 and 854, arginine mutations at residues 656 and 748, and 'knobs-into-holes' mutations at residues 769 and 814. The cysteine residue (640) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:4 are those of the protein sequence, and do not refer to the EU numbering convention.

(141) The DNA sequence of SEQ ID NO:5 is of a polynucleotide encoding an exemplary Fc fragment (SEQ ID NO:6), having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations. The fusion protein of SEQ ID NO:6 is configured, for example, for heterodimer formation with the MOG-Seldeg (SEQ ID NO:2) or HER2-Seldeg (SEQ ID NO:4) fusion.

(142) In particular, the amino acid sequence of the exemplary Fc fragment of SEQ ID NO: 6 has, in order from N- to C-terminus, an immunoglobulin hinge (human IgG1-derived) at residues 1-16, an immunoglobulin CH2 domain (human IgG1-derived) at residues 17-126, an immunoglobulin CH3 domain (human IgG1-derived) at residues 127-233. The exemplary Fc fragment of SEQ ID NO:6 has mutations that increase FcRn binding at residues 38, 40, 42, 219 and 220, arginine mutations at residues 22 and 114, and 'knobs-into-holes' mutations at residues 150 and 191. The cysteine residue (6) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:6 are those of the protein sequence, and do not refer to the EU numbering convention.

(143) The DNA sequence of SEQ ID NO:7 is of a polynucleotide encoding an exemplary MOG-Seldeg-PS fusion having knobs-into-holes mutations and arginine mutations, and the corresponding amino acid sequence of the encoded fusion protein is SEQ ID NO:8.

(144) In particular, the amino acid sequence of the exemplary MOG-Seldeg-PS fusion of SEQ ID NO: 8 has, in order from N- to C-terminus, residues 1-117 of mMOG, a first linker at residues 118-122, an immunoglobulin hinge (human IgG1-derived) at residues 123-138, an immunoglobulin CH2 domain (human IgG1-derived) at residues 139-248, an immunoglobulin CH3 domain (human IgG1-derived) at residues 249-355. The exemplary MOG-Seldeg-PS of SEQ ID NO: 8 has arginine mutations at residues 144 and 236, and 'knobs-into-holes' mutations at residues 257 and 302. The cysteine residue (128) that pairs with an immunoglobulin light chain is mutated to serine. Residues 141-266 of the C2A PS-binding domain of synaptotagmin (Syt1) (shown as residues 361-486) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 356-360). The amino acid residue numbers referred to in SEQ ID NO:8 are those of the protein sequence, and do not refer to the EU numbering convention.

(145) The DNA sequence of SEQ ID NO:9 is of a polynucleotide encoding an exemplary Fc-Syt1 fusion (SEQ ID NO:10) having knobs-into-holes mutations and arginine mutations, configured for

heterodimer formation, for example, with the MOG-Seldeg-PS (SEQ ID NO: 8).

(146) In particular, the amino acid sequence of the exemplary Fc-Syt1 fusion of SEQ ID NO: 10 has, in order from N- to C-terminus, an immunoglobulin hinge (human IgG1-derived) at residues 1-16, an immunoglobulin CH2 domain (human IgG1-derived) at residues 17-126, an immunoglobulin CH3 domain (human IgG1-derived) at residues 126-233. The exemplary Fc-Syt1 fusion protein of SEQ ID NO:10 has arginine mutations at residues 22 and 114 and 'knobs-into-holes' mutations at residues 150 and 191. The cysteine residue (6) that usually pairs with an immunoglobulin light chain is mutated to serine. Residues 141-266 of the C2A PS-binding domain of synaptotagmin (Syt1) (shown as residues 239-364) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 234-238). The amino acid residue numbers referred to in SEQ ID NO:10 are those of the protein sequence, and do not refer to the EU numbering convention.

(147) The DNA sequence of SEQ ID NO:11 is of a polynucleotide encoding an exemplary MOG-Seldeg-PS fusion (SEQ ID NO:12) having knobs-into-holes mutations, electrostatic steering mutations and arginine mutations.

(148) In particular, the amino acid sequence of the exemplary MOG-Seldeg-PS fusion of SEQ ID NO: 12 has, in order from N- to C-terminus, residues 1-117 of mMOG, a first linker at residues 118-122, an immunoglobulin hinge (human IgG1-derived) at residues 123-138, an immunoglobulin CH2 domain (human IgG1-derived) at residues 139-248, an immunoglobulin CH3 domain (human IgG1-derived) at residues 249-355. The exemplary MOG-Seldeg-PS of SEQ ID NO: 12 has arginine mutations at residues 144 and 236, electrostatic steering mutations at residues 300 and 317 and 'knobs-into-holes' mutations at residues 257 and 302. The cysteine residue (128) that pairs with an immunoglobulin light chain is mutated to serine. Residues 141-266 of the C2A PS-binding domain of synaptotagmin (Syt1) (shown as residues 361-486) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 356-360). The amino acid residue numbers referred to in SEQ ID NO:12 are those of the protein sequence, and do not refer to the EU numbering convention.

(149) The DNA sequence of SEQ ID NO:13 is of a polynucleotide encoding an exemplary Fc-Syt1 fusion (SEQ ID NO:14) having knobs-into-holes mutations, electrostatic steering mutations and arginine mutations, configured, for example, for heterodimer formation with the MOG-Seldeg-PS (SEQ ID NO:12).

(150) In particular, the amino acid sequence of the exemplary Fc-Syt1 fusion protein of SEQ ID NO: 14 has, in order from N- to C-terminus, an immunoglobulin hinge (human IgG1-derived) at residues 1-16, an immunoglobulin CH2 domain (human IgG1-derived) at residues 17-126, an immunoglobulin CH3 domain (human IgG1-derived) at residues 127-233. The exemplary MOG-Seldeg-PS of SQ ID NO:14 has arginine mutations at residues 22 and 114, electrostatic steering mutations at residues 143 and 185 and 'knobs-into-holes' mutations at residues 150 and 191. The cysteine residue (6) that pairs with an immunoglobulin light chain is mutated to serine. Residues 141-266 of the C2A PS-binding domain of synaptotagmin (Syt1) (shown in residues 239-364) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 234-238). The amino acid residue numbers referred to in SEQ ID NO:14 are those of the protein sequence, and do not refer to the EU numbering convention.

(151) The DNA sequence of SEQ ID NO:15 is of a polynucleotide encoding an exemplary MOG-Seldeg-TfR fusion protein (SEQ ID NO:16) comprising a TfR-specific antibody heavy chain with knobs into holes mutations.

(152) In particular, the amino acid sequence of the exemplary TfR-specific antibody heavy chain-MOG fusion (MOG-Seldeg-TfR) of SEQ ID NO: 16 has, in order from N- to C-terminus, a TfR-specific VH domain at residues 1-116, an immunoglobulin CH1 domain (human IgG1-derived) at residues 117-213, an immunoglobulin hinge (human IgG1-derived) at residues 214-229, an immunoglobulin CH2 domain (human IgG1-derived) at residues 230-337, an immunoglobulin CH3

domain (human IgG1-derived) at residues 237-446. The exemplary Tfr-specific antibody heavy chain of SEQ ID NO:16 has 'knobs-into-holes' mutations at residues 348 and 393. Residues 1-117 of mMOG (shown as residues 452-568) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 447-451). The amino acid residue numbers referred to in SEQ ID NO:16 are those of the protein sequence, and do not refer to the EU numbering convention.

(153) The DNA sequence of SEQ ID NO:17 is of a polynucleotide encoding an exemplary Tfr-specific antibody heavy chain. The encoded fusion protein (SEQ ID NO:18) having knobs into holes mutations is configured, for example, for heterodimer formation with the MOG-Seldeg-Tfr fusion (SEQ ID NO:16).

(154) In particular, the amino acid sequence of the exemplary Tfr-specific antibody heavy chain of SEQ ID NO: 18 for heterodimer formation with the Tfr-specific heavy chain-MOG fusion (SEQ ID NO: 16) has, in order from N- to C-terminus, of a Tfr-specific VH domain at residues 1-116, an immunoglobulin CH1 domain (human IgG1-derived) at residues 117-213, an immunoglobulin hinge (human IgG1-derived) at residues 214-229 an immunoglobulin CH2 domain (human IgG1-derived) at residues 230-339, an immunoglobulin CH3 domain (human IgG1-derived) at residues 340-446. The exemplary Tfr-specific antibody heavy chain of SEQ ID NO: 18 has 'knobs-into-holes' mutations at residues 363 and 404. The amino acid residue numbers referred to in SEQ ID NO:18 are those of the protein sequence, and do not refer to the EU numbering convention.

(155) The DNA sequence of SEQ ID NO:19 is of a polynucleotide encoding an exemplary light chain of a Tfr-specific antibody (SEQ ID NO:20) and is configured, for example, for heterodimer formation with the MOG-Seldeg-Tfr fusion (SEQ ID NO:16) and Tfr-specific antibody heavy chain (SEQ ID NO:18).

(156) In particular, the amino acid sequence of the exemplary Tfr-specific antibody light chain of SEQ ID NO: 20 has, in order from N- to C-terminus a Tfr-specific VL domain at residues 1-107, and an immunoglobulin CL domain (human Cκ) at residues 108-213. The amino acid residue numbers referred to in SEQ ID NO:20 are those of the protein sequence, and do not refer to the EU numbering convention.

(157) The DNA sequence of SEQ ID NO: 21 is of a polynucleotide encoding an exemplary MOG-Seldeg-Tfr fusion protein (SEQ ID NO: 22) having a Tfr-specific antibody heavy chain with arginine mutations, and knobs-into-holes mutations.

(158) In particular, the amino acid sequence of the exemplary Tfr-specific antibody heavy chain-MOG fusion (MOG-Seldeg-Tfr) of SEQ ID NO: 22 has, in order from N- to C-terminus, a Tfr-specific VH domain at residues 1-116, an immunoglobulin CH1 domain (human IgG1-derived) at residues 117-213, an immunoglobulin hinge (human IgG1-derived) at residues 214-229, an immunoglobulin CH2 domain (human IgG1-derived) at residues 230-339, an immunoglobulin CH3 domain (human IgG1-derived) at residues 340-446. The exemplary Tfr-specific antibody heavy chain of SEQ ID NO: 22 has arginine mutations at residues 235 and 327, and 'knobs-into-holes' mutations at residues 348 and 393. Residues 1-117 of mMOG (shown as residues 452-568) are fused to the C-terminus of the CH3 domain via a GGGGS linker peptide (residues 447-451). The amino acid residue numbers referred to in SEQ ID NO:22 are those of the protein sequence, and do not refer to the EU numbering convention.

(159) The DNA sequence of SEQ ID NO: 23 is of a polynucleotide encoding an exemplary Tfr-specific antibody heavy chain (SEQ ID NO: 24) having arginine mutations, and knobs-into-holes mutations, and is configured, for example, for heterodimer formation with the MOG-Seldeg-Tfr fusion (SEQ ID NO: 22).

(160) In particular, the amino acid sequence of the exemplary Tfr-specific antibody heavy chain of SEQ ID NO: 24 has, in order from N- to C-terminus, a Tfr-specific VH domain at residues 1-116, an immunoglobulin CH1 domain (human IgG1-derived) at residues 117-213, an immunoglobulin hinge (human IgG1-derived) at residues 214-229 an immunoglobulin CH2 domain (human IgG1-derived) at residues 230-339, an immunoglobulin CH3 domain (human IgG1-derived) at residues

340-446. The TfR-specific antibody heavy chain of SEQ ID NO:24 has arginine mutations at residues 235 and 327, and 'knobs-into-holes' mutations at residues 363 and 404.). The amino acid residue numbers referred to in SEQ ID NO:24 are those of the protein sequence, and do not refer to the EU numbering convention.

(161) The DNA sequence of SEQ ID NO: 25 is of a polynucleotide encoding an exemplary HER2-Seldeg fusion protein (SEQ ID NO: 26) having mutations to increase FcRn binding, and knobs-into-holes mutations.

(162) In particular, the amino acid sequence of the exemplary variant HER2-Seldeg of SEQ ID NO: 26 forms a fusion protein having, in order from N- to C-terminus, residues 1-630 of HER2, a first linker at residues 631-636, an immunoglobulin hinge (human IgG1-derived) at residues 637-650, an immunoglobulin CH2 domain (human IgG1-derived) at residues 651-760, an immunoglobulin CH3 domain (human IgG1-derived) at residues 761-867. The HER2-Seldeg of SEQ ID NO: 26 has mutations that increase FcRn binding at residues 672, 674, 676, 853 and 854, and 'knobs-into-holes' mutations at residues 769 and 814. The cysteine residue (640) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:26 are those of the protein sequence, and do not refer to the EU numbering convention.

(163) The DNA sequence of SEQ ID NO: 27 is of a polynucleotide encoding an exemplary Fc fragment (SEQ ID NO:28) having mutations to increase FcRn binding, and knobs-into-holes mutations, and is configured, for example, for heterodimer formation with the HER2-Seldeg (SEQ ID NO: 26).

(164) In particular, the amino acid sequence of the exemplary variant Fc fragment of SEQ ID NO: 28 has, in order from N- to C-terminus, an immunoglobulin hinge (human IgG1-derived) at residues 1-16, an immunoglobulin CH2 domain (human IgG1-derived) at residues 17-126, an immunoglobulin CH3 domain (human IgG1-derived) at residues 127-233. The variant Fc fragment of SEQ ID NO: 28 has mutations that increase FcRn binding at residues 38, 40, 42, 219 and 220, and 'knobs-into-holes' mutations at residues 150 and 191. The cysteine residue (6) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:28 are those of the protein sequence, and do not refer to the EU numbering convention.

(165) The DNA sequence of SEQ ID NO: 29 is of a polynucleotide encoding an exemplary prostate-specific membrane antigen (PSMA)-Seldeg fusion protein (SEQ ID NO: 30) having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations. This fusion protein will form heterodimers, for example, with an exemplary Fc fragment (SEQ ID NO: 6).

(166) In particular, the amino acid sequence of the exemplary variant PSMA-Seldeg of SEQ ID NO: 30 forms a fusion protein having, in order from N- to C-terminus, an immunoglobulin hinge (human IgG1-derived) at residues 1-16, an immunoglobulin CH2 domain (human IgG1-derived) at residues 17-126, an immunoglobulin CH3 domain (human IgG1-derived) at residues 127-233 fused at the C-terminus via a linker at residues 234-238 to the extracellular domain (residues 239-945) of PSMA. The variant PSMA-Seldeg of SEQ ID NO: 30 has mutations that increase FcRn binding at residues 38, 40, 42, 219 and 220, arginine mutations at residues 22 and 114, and 'knobs-into-holes' mutations at residues 135 and 180. The cysteine residue (6) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:30 are those of the protein sequence, and do not refer to the EU numbering convention.

(167) The DNA sequence of SEQ ID NO: 31 is of a polynucleotide encoding an exemplary GAD65-Seldeg fusion protein (SEQ ID NO: 32) having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations. This fusion protein will form heterodimers, for example, with an exemplary Fc fragment (SEQ ID NO: 6).

(168) In particular, the amino acid sequence of the exemplary variant GAD65-Seldeg of SEQ ID NO: 32 forms a fusion protein having, in order from N- to C-terminus, residues 1-585 of human glutamic acid carboxylase 65 (GAD65), a first linker at residues 586-590, an immunoglobulin hinge (human IgG1-derived) at residues 591-606, an immunoglobulin CH2 domain (human IgG1-derived) at residues 607-716, an immunoglobulin CH3 domain (human IgG1-derived) at residues 717-823. The variant GAD65-Seldeg of SEQ ID NO:32 has mutations that increase FcRn binding at residues 628, 630, 632, 809 and 810, arginine mutations at residues 612 and 704, and ‘knobs-into-holes’ mutations at residues 725 and 770. The cysteine residue (596) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:32 are those of the protein sequence, and do not refer to the EU numbering convention.

(169) The DNA sequence of SEQ ID NO: 33 is of a polynucleotide encoding an exemplary aquaporin 4-Seldeg fusion protein (SEQ ID NO: 34) having mutations to increase FcRn binding, knobs-into-holes mutations and arginine mutations. This fusion protein will form heterodimers, for example, with an exemplary Fc fragment (SEQ ID NO: 6).

(170) In particular, the amino acid sequence of the exemplary variant aquaporin 4 (AQP4)-Seldeg of SEQ ID NO: 34 forms a fusion protein having, in order from N- to C-terminus, residues 1-323 of human aquaporin 4, a first linker at residues 324-328, an immunoglobulin hinge (human IgG1-derived) at residues 329-344, an immunoglobulin CH2 domain (human IgG1-derived) at residues 345-454, an immunoglobulin CH3 domain (human IgG1-derived) at residues 455-561. The variant AQP4-Seldeg of SEQ ID NO:34 has mutations that increase FcRn binding at residues 366, 368, 370, 547,548, arginine mutations at residues 350 and 442, and ‘knobs-into-holes’ mutations at residues 463 and 508. The cysteine residue (334) that usually pairs with an immunoglobulin light chain is mutated to serine. The amino acid residue numbers referred to in SEQ ID NO:34 are those of the protein sequence, and do not refer to the EU numbering convention.

(171) The above disclosed subject matter is to be considered illustrative, and not restrictive, and the appended claims are intended to cover all such modifications, enhancements, and other embodiments which fall within the true spirit and scope of the present disclosure. Thus, to the maximum extent allowed by law, the scope of the present disclosure is to be determined by the broadest permissible interpretation of the following claims and their equivalents, and shall not be restricted or limited by the foregoing detailed description.

(172) As used in this specification and the appended claims, the singular forma “a”, “an”, and “the” include plural references unless the content clearly dictates otherwise. The term “plurality” includes two or more referents unless the content clearly dictates otherwise.

(173) Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the disclosure pertains.

Claims

1. A method of depleting target antigen-specific antibody from a patient, the method comprising: administering to the patient a Seldeg in an amount sufficient to remove at least 50% of the target antigen-specific antibody from a circulation or a target tissue in the patient, wherein the Seldeg comprises a targeting component having a protein or a protein fragment configured to specifically bind to an internalizing cell surface receptor or internalizing surface molecule; and an antigen component, having one molecule of an antigen, an antigen fragment or an antigen mimetic configured to specifically bind a target antigen-specific antibody or a variant thereof; wherein the targeting protein component is fused directly or indirectly to the antigen component.
2. The method of claim 1, comprising administering the Seldeg in an amount sufficient to remove at least 50% of the target antigen-specific antibody from the circulation or the target tissue in the

patient within five hours of administration.

3. The method of claim 1, wherein the protein or the protein fragment is configured to bind to the internalizing cell surface receptor or other internalizing cell surface molecule with a dissociation constant of less than 10 μ M at near neutral pH.

4. The method of claim 1, wherein the amount sufficient of Seldeg is an amount at least equimolar to the amount of target antigen-specific antibody to be depleted.

5. The method of claim 1, comprising administering the Seldeg in an amount sufficient to remove at least 90% of the target antigen-specific antibody from the circulation or the target tissue in the patient within two hours of administration.

6. The method of claim 1, comprising administering the Seldeg in an amount sufficient to remove at least 50% of the target antigen-specific antibody from the circulation or the target tissue in the patient within one hour of administration.

7. The method of claim 1, further comprising re-administering the Seldeg whenever 50% of patients are expected to have regenerated a threshold amount of target antigen-specific antibody in the circulation of target tissue.

8. The method of claim 1, wherein the Seldeg removes less than 10% of non-target antibodies in the circulation or in the tissue targeted by the target antigen-specific antibody.

9. The method of claim 1, wherein the Seldeg removes an amount of non-targeted antibodies in the circulation or in the target tissue of the patient that does not cause a clinically adverse effect in the patient.

10. The method of claim 1, wherein the Seldeg removes less than 1% of non-target antibodies in the circulation or in a tissue targeted by the target antigen-specific antibody.

11. The method of claim 1, wherein the target antigen-specific antibody is delivered to lysosomes within the cell following Seldeg-mediated internalization of the target antibody by an internalizing cell surface receptor.

12. The method of claim 1, wherein the Seldeg is administered to a patient with an autoimmune disease and the target antigen-specific antibody specifically binds to an autoantigen.

13. The method of claim 1, wherein the Seldeg is administered to a patient receiving a transplanted organ and the target antigen-specific antibody specifically binds to an antigen on the transplanted organ.

14. The method of claim 1, wherein the Seldeg is administered to increase contrast during tumor imaging and the target antigen-specific antibody specifically binds to a tumor antigen.

15. The method of claim 1, wherein the Seldeg is administered to a patient who has received a biologic and the target antigen-specific antibody is the biologic.

16. The method of claim 1, wherein the Seldeg is administered prior to the delivery of a therapeutic agent, if the patient has antibodies specific for the therapeutic agent, and the Seldeg is configured to target the antibodies specific for the therapeutic agent.

17. The method of claim 1, wherein the Seldeg is administered to provide a PET image contrast agent.

18. The method of claim 1, wherein said targeting component is selected from the group consisting of: (1) an antibody or antibody fragment, having (i) a dissociation constant for FcRn at a pH greater than 6.8 and less than 7.5 of less than 10 μ M; and/or (ii) an altered binding affinity for Fc γ Rs and/or complement (C1q), compared with a wild-type Fc fragment or is an Fc fragment derived from IgG2 or IgG4; (2) an antibody or antibody fragment specific for an internalizing cell surface receptor, wherein the Fc fragment of said antibody or antibody fragment is: (i) engineered to have an altered binding affinity for Fc γ Rs and/or complement (C1q), compared with a wild-type Fc fragment; or (ii) an Fc fragment derived from IgG2 or IgG4; (3) a ligand for an internalizing cell surface receptor, wherein said ligand is fused to an antibody Fc fragment which is (i) engineered to have an altered binding affinity for Fc γ Rs and/or complement (C1q), compared with a wild-type Fc fragment; or (ii) derived from IgG2 or IgG4; (4) an antibody or antibody fragment specific for

transferrin receptor, phosphatidylserine, asialoglycoprotein receptor, inhibitory FcγR or mannose-6-phosphate receptor; and (5) a ligand for FcRn, transferrin receptor, phosphatidylserine, asialoglycoprotein receptor, inhibitory FcγR or mannose-6-phosphate receptor.
