

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent

12404488

Kind Code

B2

Date of Patent

September 02, 2025

Inventor(s)

Chien; Kenneth R. et al.

Use of neuropilin-1 (NRP1) as a cell surface marker for isolating human cardiac ventricular progenitor cells

Abstract

The present invention provides NRP1 as a cell surface marker for isolating human cardiomyogenic ventricular progenitor cells (HVPs), in particular progenitor cells that preferentially differentiate into cardiac ventricular muscle cells. Additional HVP cell surface markers identified by single cell sequencing are also provided. The invention provides in vitro methods of the separation of NRP1+ ventricular progenitor cells, and the large scale expansion and propagation thereof. Large clonal populations of isolated NRP1+ ventricular progenitor cells are also provided. Methods of in vivo use of NRP1+ ventricular progenitor cells for cardiac repair or to improve cardiac function are also provided. Methods of using the NRP1+ ventricular progenitor cells for cardiac toxicity screening of test compounds are also provided.

Inventors: Chien; Kenneth R. (Cambridge, MA), Clarke; Jonathan (Stockholm, SE), Leung; Chuen Yan (Stockholm, SE)

Applicant: Procella Therapeutics AB (Tullinge, SE)

Family ID: 63794530

Assignee: Procella Therapeutics AB (Tullinge, SE)

Appl. No.: 17/507454

Filed: October 21, 2021

Prior Publication Data

Document Identifier

US 20220112458 A1

Publication Date

Apr. 14, 2022

Related U.S. Application Data

Publication Classification

Int. Cl.: C12N5/073 (20100101); A01K67/0275 (20240101); C07K16/28 (20060101);
C12N5/077 (20100101); C12N15/00 (20060101); G01N33/50 (20060101)

U.S. Cl.:

CPC C12N5/0605 (20130101); A01K67/0275 (20130101); C07K16/2863 (20130101);
C12N5/0657 (20130101); C12N15/00 (20130101); G01N33/5014 (20130101);
G01N33/5044 (20130101); A01K2227/105 (20130101); A01K2267/0375 (20130101);
A01K2267/0393 (20130101); C12N2500/98 (20130101); C12N2501/415 (20130101);
C12N2501/727 (20130101); C12N2506/02 (20130101); C12N2506/45 (20130101);
C12N2533/90 (20130101)

Field of Classification Search

CPC: C12N (5/0605); C12N (5/0657); C12N (15/00); C12N (2500/98); C12N (2501/415);
C12N (2501/727); C12N (2506/02); C12N (2506/45); C12N (2533/90); A01K (67/0275);
A01K (2227/105); A01K (2267/0375); A01K (2267/0393); C07K (16/2863); G01N
(33/5014); G01N (33/5044)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5843780	12/1997	Thomson	N/A	N/A
5945577	12/1998	Stice et al.	N/A	N/A
5994619	12/1998	Stice et al.	N/A	N/A
6200806	12/2000	Thomson	N/A	N/A
6235970	12/2000	Stice et al.	N/A	N/A
8765117	12/2013	Chien et al.	N/A	N/A
8951798	12/2014	Palecek et al.	N/A	N/A
9453201	12/2015	Palecek et al.	N/A	N/A
9663764	12/2016	Palecek et al.	N/A	N/A
9717762	12/2016	Wiencierz et al.	N/A	N/A
9765299	12/2016	Palecek et al.	N/A	N/A
10508263	12/2018	Chien et al.	N/A	N/A
10596200	12/2019	Chien et al.	N/A	N/A
10597637	12/2019	Chien et al.	N/A	N/A
10612094	12/2019	Leung et al.	N/A	N/A
11186820	12/2020	Chien et al.	N/A	N/A
11401508	12/2021	Chien et al.	N/A	N/A
11725244	12/2022	Leung et al.	N/A	N/A
2004/0180043	12/2003	Sabbah et al.	N/A	N/A
2005/0214260	12/2004	Franco	N/A	N/A

2006/0246446	12/2005	Evans et al.	N/A	N/A
2008/0038229	12/2007	Minguell et al.	N/A	N/A
2008/0182328	12/2007	Snyder et al.	N/A	N/A
2009/0162326	12/2008	Siemonsmeier et al.	N/A	N/A
2009/0162329	12/2008	Anversa et al.	N/A	N/A
2010/0093089	12/2009	Marban	N/A	N/A
2010/0166714	12/2009	Chien et al.	N/A	N/A
2010/0297124	12/2009	Tosato et al.	N/A	N/A
2011/0033430	12/2010	Chien et al.	N/A	N/A
2012/0009158	12/2011	Chien et al.	N/A	N/A
2012/0027807	12/2011	Chien et al.	N/A	N/A
2012/0301445	12/2011	Blanpain et al.	N/A	N/A
2013/0115626	12/2012	Schmidt et al.	N/A	N/A
2013/0189785	12/2012	Palecek et al.	N/A	N/A
2013/0309769	12/2012	Benvenisty et al.	N/A	N/A
2014/0134733	12/2013	Wu et al.	N/A	N/A
2015/0152389	12/2014	Palecek et al.	N/A	N/A
2015/0252117	12/2014	Chinn et al.	N/A	N/A
2015/0297794	12/2014	Yamashita et al.	N/A	N/A
2016/0053229	12/2015	Chien et al.	N/A	N/A
2016/0068814	12/2015	Palecek et al.	N/A	N/A
2016/0108363	12/2015	Chien et al.	N/A	N/A
2016/0362661	12/2015	O'Sullivan et al.	N/A	N/A
2017/0002325	12/2016	Palecek et al.	N/A	N/A
2017/0067023	12/2016	Yamashita et al.	N/A	N/A
2017/0153236	12/2016	Martin et al.	N/A	N/A
2017/0239298	12/2016	Keith et al.	N/A	N/A
2017/0240964	12/2016	Leung et al.	N/A	N/A
2018/0148691	12/2017	Chien et al.	N/A	N/A
2019/0062696	12/2018	Chien et al.	N/A	N/A
2020/0140819	12/2019	Chien et al.	N/A	N/A
2020/0268803	12/2019	Chien et al.	N/A	N/A
2020/0270685	12/2019	Leung et al.	N/A	N/A
2023/0383345	12/2022	Leung et al.	N/A	N/A

FOREIGN PATENT DOCUMENTS

Patent No.	Application Date	Country	CPC
20120124596	12/2011	KR	N/A
9717762	12/1996	WO	N/A
2009/017460	12/2008	WO	N/A
2011/091944	12/2010	WO	N/A
2011/153236	12/2010	WO	N/A
201216274	12/2011	WO	N/A
2012074116	12/2011	WO	N/A
2012/162740	12/2011	WO	N/A
2012/162741	12/2011	WO	N/A
2013/056072	12/2012	WO	N/A
2014/150602	12/2013	WO	N/A
2014148562	12/2013	WO	N/A
2015/058117	12/2014	WO	N/A

2016/029122	12/2015	WO	N/A
2016/131137	12/2015	WO	N/A
2017/172086	12/2016	WO	N/A
2018/100433	12/2017	WO	N/A

OTHER PUBLICATIONS

Li et al. "Higher mortality in heterozygous neuropilin-1 mice after cardiac pressure overload." *Biochem Biophys Res Commun*. May 30, 2008;370(2):317-21. (Year: 2008). cited by examiner

Smits et al. "Human cardiomyocyte progenitor cell transplantation preserves long-term function of the infarcted mouse myocardium." *Cardiovasc Res* . Aug. 1, 2009;83(3):527-35. (Year: 2009). cited by examiner

Posch et al. "Cardiac alpha-myosin (MYH6) is the predominant sarcomeric disease gene for familial atrial septal defects." *PLoS One*. 2011;6(12):e28872. (Year: 2011). cited by examiner

Andersson, E.R., et al., "Therapeutic modulation of Notch signalling—are we there yet?" *Nat. Rev. Drug Discov*. vol. 13:357-378 (2014). cited by applicant

Ardehali, R. et al., "Prospective isolation of human embryonic stem cell-derived cardiovascular progenitors that integrate into human fetal heart tissue," *PNAS*, vol. 110(9): 3405-3410 (2013). cited by applicant

Ardehali, R. et al., "Prospective isolation of human embryonic stem cell-derived cardiovascular progenitors that integrate into human fetal heart tissue," *Supporting Information, PNAS*, 12 pages (2013). cited by applicant

Arlas.Romero, L. et al., "Targeting Cdc42 in Cancer," *Expert Opin Therp—Targets*, vol. 17(11):1263-1273 (2013). cited by applicant

Badcock, G. et al., "The human embryonal carcinoma marker antigen TRA-1-60 is a sialylated keratan sulfate proteoglycan," *Cancer Res.*, vol. 59:4715-4719 (1999). cited by applicant

Barbash, I.M. et al. "Systemic delivery of bone marrow-derived mesenchymal stem cells to the infarcted myocardium: feasibility, cell migration, and body distribution," *Circulation*, vol. 108, pp. 863-868 (2003). cited by applicant

Basavarajaiah, S., et al., "Physiological upper limits of left ventricular dimensions in highly trained junior tennis players," *Br. J. Sports Med.*, vol. 41(11): 784-788. (2007). cited by applicant

Bash, J. et al., "Rel/NF-kappaB can trigger the Notch signaling pathway by inducing the expression of Jagged1, a ligand for Notch receptors," *The EMBO Journal*, vol. 18(10) pp. 2803-2811 (1999). cited by applicant

Bearzi, C. et al., "Human cardiac stem cells," *PNAS*, vol. 104 (35):14068-14073 (2007). cited by applicant

Bergmann, O., et al., "Dynamics of Cell Generation and Turnover in the Human Heart," *Cell*, vol. 161(17):1566-1575 (2015). cited by applicant

Bergmann, O., et al., "Evidence for cardiomyocyte renewal in humans," *Science*, vol. 324(5923), pp. 98-102 (2009). cited by applicant

Bhattacharya, et al. "High Efficiency Differentiation of Human Pluripotent Stem Cells to Cardiomyocytes and Characterization by Flow Cytometry." *J Vis Exp.*, (91): 52010. (2014). cited by applicant

Birket, M. et al., "Expansion and patterning of cardiovascular progenitors derived from human pluripotent stem cells," *Nature Biotechnology*, vol. 33(9), pp. 970-979(2015). cited by applicant

Bittira, B. et al., "Mobilization and homing of bone marrow stromal cells in myocardial infarction," *Eur. J. Cardiothorac. Surg.*, vol. 24(3) pp. 393-398 (2003). cited by applicant

Blin, G. et al., "A purified population of multipotent cardiovascular progenitors derived from primate pluripotent stem cells engrafts in postmyocardial infarcted nonhuman primates," *The Journal of Clinical Investigation*, vol. 120(4) pp. 1125-1139 (2010). cited by applicant

Brette, F., et al., "T-tubule function in mammalian cardiac myocytes," *Circulation Research*, vol.

92(11) pp. 1182-1192 (2003). cited by applicant

Bu, L., et al., "Human ISL1 heart progenitors generate diverse multipotent cardiovascular cell lineages," *Nature*, vol. 460, pp. 113-117 (2009). cited by applicant

Bui, A.L. et al., "Epidemiology and risk profile of heart failure," *Nat. Rev. Cardiol*, vol. 8(1) pp. 30-41 (2011). cited by applicant

Buikema, J. et al., "Wnt/beta-catenin signaling directs the regional expansion of first and second heart field-derived ventricular cardiomyocytes," *Development*, vol. 140, pp. 4165-4176 (2013). cited by applicant

Burridge, P. et al., "Chemically defined generation of human cardiomyocytes,"; *Nature Methods*, vol. 11 (8), pp. 855-860 (2014). cited by applicant

Cai, C.L. et al., "Isl1 identifies a cardiac progenitor population that proliferates prior to differentiation and contributes a majority of cells to the heart," *Developmental Cell*, vol. 5, pp. 877-889 (2003). cited by applicant

Campagnolo, P. et al., "c-Kit⁺ progenitors generate vascular cells for tissue-engineered grafts through modulation of the Wnt/Klf4 pathway," *Biomaterials*, vol. 60, pp. 53-61 (2015). cited by applicant

Chen, G. et al., "Chemically defined conditions for human iPSC derivation and culture," *Nature Methods*, vol. 8 (5):424-429 (2011). cited by applicant

Chong, J. et al., "Human Embryonic Stem Cell-Derived Cardiomyocytes Regenerate Non-Human Primate Hearts," *Nature*, vol. 510(7504): 273-277 (2014). cited by applicant

Chong, J. et al., "Human embryonic-stem-cell-derived cardiomyocytes regenerate non-human primate hearts," *Nature*, vol. 510, pp. 273-277, (2014). cited by applicant

Christoforou, N. et al., "Implantation of Mouse Embryonic Stem Cell-Derived Cardiac Progenitor Cells Preserves Function of Infarcted Murine Hearts," *PLOS One*, vol. 5(7):e11536, 14 pages (2010). cited by applicant

Collesi, C. et al. "Notch1 signaling stimulates proliferation of immature cardiomyocytes," *The Journal of Cell Biology*, Sep. 29, 2008, 12 pages. cited by applicant

Conradi, L., et al. "Immunobiology of fibrin-based engineered heart tissue," *Stem Cells Transl. Med.*, vol. 4(6), pp. 625-631 (2015). cited by applicant

Den Hartogh, S. et al., "A comprehensive gene expression analysis at sequential stages of in vitro cardiac differentiation from isolated MESP1-expressing-mesoderm progenitors," *Scientific Reports*, vol. 6 (19386) 16 pages (2016) doi: 10.1038/srep19386. cited by applicant

Di Giorgio, F.P. et al., "Human embryonic stem cell-derived motor neurons are sensitive to the toxic effect of glial cells carrying an ALS-causing mutation," *Cell Stem Cell*, vol. 3, pp. 637-648 (2008). cited by applicant

Domian, I.J., et al., "Generation of functional ventricular heart muscle from mouse ventricular progenitor cells," *Science* vol. 326(5951), pp. 426-429 (2009). cited by applicant

Elliot, D. et al., "NKX2-5eGFP/w hESCs for isolation of human cardiac progenitors and cardiomyocytes," *Nature Methods*, vol. 8(12), pp. 1037-1043 (2011). cited by applicant

Extended European Search Report, European Application No. 18211720.0 , dated Jul. 12, 2019, 9 pages. cited by applicant

Extended European Search Report, European Application No. 19210435.4, dated Mar. 6, 2020, 8 pages. cited by applicant

Fernandes, S. et al., "Comparison of Human Embryonic Stem Cell-Derived Cardiomyocytes, Cardiovascular Progenitors, and Bone Marrow Mononuclear Cells for Cardiac Repair," *Stem Cell Reports*, vol. 5, pp. 753-762 (2015). cited by applicant

Fong, C. Y. et al., "Separation of SSEA-4 and TRA-1-60 labelled undifferentiated human embryonic stem cells from a heterogeneous cell population using magnetic-activated cell sorting (MACS) and fluorescence-activated cell sorting (FACS)," *Stem Cell Rev.*, vol. 5:72-80 (2009). cited by applicant

Foo, K. et al., "Human ISL1+ Ventricular Progenitors Self-Assemble into an In Vivo Functional Heart Patch and Preserve Cardiac Function Post Infarction," *Molecular Therapy*, vol. 26(7): 1644-1659 (2018). cited by applicant

Forbes, S.J. et al., "Preparing the ground for tissue regeneration: from mechanism to therapy," *Nat. Med.*, vol. 20(8), pp. 857-869 (2014). cited by applicant

Funakoshi, S. et al., "Enhanced engraftment, proliferation, and therapeutic potential in heart using optimized human PSC-derived cardiomyocytes," *Scientific Reports*, vol. 6:19111, p. 1-14 (2016). cited by applicant

Gaetani, R., et al., "Epicardial application of cardiac progenitor cells in a 3D-printed gelatin/hyaluronic acid patch preserves cardiac function after myocardial infarction," *Biomaterials*, vol. 61, 339-348. (2015). cited by applicant

Gama-Carvalho, M., et al., "Regulation of Cardiac Cell Fate by microRNAs: Implications for Heart Regeneration," *Cells*, vol. 3(4), pp. 996-1026 (2014). cited by applicant

Gao, L. et al., "Myocardial Tissue Engineering With Cells Derived from Human Induced-Pluripotent Stem Cells and a Native-Like, High-Resolution, 3-Dimensionally Printed Scaffold," *Circ Res.*, vol. 120(8): 1318-1325 (2017). cited by applicant

Gao, L. et al., "Supplemental Material Myocardial tissue engineering with cardiac cells derived from human induced-pluripotent stem cells and a native-like, high-resolution, 3-dimensionally printed scaffold," *Circ Res.*, vol. 120 (8):1318-1325 (2017). cited by applicant

Gearing, D. et al., "Leukemia inhibitory factor receptor is structurally related to the IL-6 signal transducer, gp130," *EMBO*, vol. 10(10):2839-2848 (1991). cited by applicant

Giordano, F.J., et al., "A cardiac myocyte vascular endothelial growth factor paracrine pathway is required to maintain cardiac function," *PNAS, U. S. A.*, vol. 98(10), pp. 5780-5785 (2001). cited by applicant

Guo, Y. et al., "Targeted deletion of the A3 adenosine receptor confers resistance to myocardial ischemic injury and does not prevent early preconditioning," *J Mol Cell Cardiol.*, vol. 33(4), pp. 825-830 (2001). cited by applicant

Harrison, R.H., et al., "Tissue engineering and regenerative medicine: a year in review," *Tissue Eng. Part B.*, vol. 20, 1-16. (2014). cited by applicant

Headrick, J.P, et al., "Acute adenosinergic cardioprotection in ischemic-reperfused hearts," *American Journal of Physiology Heart and Circulatory Physiology*, vol. 285; pp. H1797-H1818, (2003). cited by applicant

High, F. et al., "Endothelial expression of the Notch ligand Jagged1 is required for vascular smooth muscle development," *Proceedings of the National Academy of Sciences of the United States of America*, vol. 105(6), pp. 1955-1959 (Feb. 2008). cited by applicant

Horvat, R. et al., "Endothelial Cell Membranes Contain Podocalyxin the Major Sialoprotein of Visceral Glomerular Epithelial Cells," *The Journal of Cell Biology*, vol. 102: 484-491 (1986). cited by applicant

Beare, A. et al "The CD System of Leukocyte Surface Molecules: Monoclonal Antibodies to Human Cell Surface Antigens," *Current Protocols in Immunology*, Appendix 4(1):4A, 73 pages (2008). cited by applicant

Chong, J. et al., "Progenitor Cells Identified by PDGFR-Alpha Expression in the Developing and Diseased Human Heart," *Stem Cells and Development* vol. 22(13):1932-1943 (2013). cited by applicant

Gupta, S. et al "FZD4 as a Mediator of ERG Oncogene-Induced WNT Signaling and Epithelial-to-Mesenchymal Transition in Human Prostate Cancer Cells," *Cancer Res.*, vol. 70(17): 6735-6745 (2010). cited by applicant

Pearl, J. et al "Short-Term Immunosuppression Promotes Engraftment of Embryonic and Induced Pluripotent Stem Cells," *Cell Stem Cell*, vol. 8(3): 309-317 (2011). cited by applicant

Schenke-Layland, K. et al., "Recapitulation of the embryonic cardiovascular progenitor cell niche,"

Biomaterials, vol. 32(11):2748-2756 (2011). cited by applicant

Huang, J. et al., "Tissue Engineering and Regenerative Medicine In Basic Research: A Year in Review 2014," Tissue Engineering: Part B, vol. 21(2) pp. 1-11 (2015). cited by applicant

International Preliminary Report on Patentability, PCT/2015/046309, dated Feb. 28, 2017, 11 pages. cited by applicant

International Preliminary Report on Patentability, PCT/IB2017/001638, dated Jun. 4, 2019, 12 pages. cited by applicant

International Preliminary Report on Patentability, PCT/IB2018/001026, dated Feb. 25, 2020, 6 pages. cited by applicant

International Preliminary Report on Patentability, PCT/US2017/017952, dated Aug. 21, 2018, 7 pages. cited by applicant

International Search Report and Written Opinion, PCT/2015/046309, dated Jan. 19, 2016, 19 pages. cited by applicant

International Search Report and Written Opinion, PCT/IB2017/001638, dated Mar. 22, 2018, 17 pages. cited by applicant

International Search Report and Written Opinion, PCT/IB2017/001638, dated Mar. 23, 2018, 17 pages. cited by applicant

International Search Report and Written Opinion, PCT/IB2018/001026, dated Jan. 16, 2019, 13 pages. cited by applicant

International Search Report and Written Opinion, PCT/US2017/017952, dated May 3, 2017, 14 pages. cited by applicant

Invitation to Pay Additional Fees and, where applicable, Protest Fee, PCT/US2015/046309, dated Nov. 6, 2015, pp. 1-7. cited by applicant

Jackson, K.A. et al., "Regeneration of ischemic cardiac muscle and vascular endothelium by adult stem cells," J. Clin. Invest., vol. 107(11), pp. 1395-1402 (2001). cited by applicant

Jones, E.A. et al. "Jagged1 expression in human embryos: correlation with the Alagille syndrome phenotype," J. Med. Genet. vol. 37, pp. 658-662 (2000). cited by applicant

Kaiser, N.J., et al., "Physiologically inspired cardiac scaffolds for tailored in vivo function and heart regeneration," Biomed. Mater. vol. 10(3) 034003. (2015). cited by applicant

Karakikes, I. et al., "Small molecule-mediated directed differentiation of human embryonic stem cells toward ventricular cardiomyocytes," Stem Cells Transl Med. <[https://www.ncbi.nlm.nih.gov/pubmed/?term=Karakikes+\(2014\)+Stem+Cell+Transl+Med](https://www.ncbi.nlm.nih.gov/pubmed/?term=Karakikes+(2014)+Stem+Cell+Transl+Med)>, vol. 3(1):18-31 (2014). cited by applicant

Keegan, K. et al., "Isolation of an additional member of the fibroblast growth factor receptor family, FGFR-3," Proc. Natl. Acad. Sci. USA, vol. 88(4):1095-1099 (1991). cited by applicant

Khan, M., et al., "Evaluation of Changes in Morphology and Function of Human Induced Pluripotent Stem Cell Derived Cardiomyocytes (HiPSC-CMs) Cultured on an Aligned-Nanofiber Cardiac Patch," PLoS One 10, e0126338, 19 pages, (2015). cited by applicant

Kim, M.S., et al., "Activin-A and Bmp4 levels modulate cell type specification during CHIR-induced cardiomyogenesis," PLOS One, vol. 10(2):e01186701, 16 pages (2015). cited by applicant

Kirikoshi, H., et al. "Molecular cloning and characterization of human Frizzled-4 on chromosome 11q14-q21," Biochem. Biophys. Res. Commun., vol. 264(3), pp. 955-961(1999). cited by applicant

Koo, B.-K., et al., "Controlled gene expression in primary Lgr5 organoid cultures," Nat. Methods vol. 9(1), pp. 81-83 (2012). cited by applicant

Koyanagi, M. et al., "Differentiation of circulating endothelial progenitor cells to a cardiomyogenic phenotype depends on E-cadherin," FEBS Letters, vol. 579: 6060-6066 (2005). cited by applicant

Kraehenbuehl, T.P, et al., "Three-dimensional biomaterials for the study of human pluripotent stem cells," Nat. Methods vol. 8(9), pp. 731-736. (2011). cited by applicant

Krishnan, A, et al., "A detailed comparison of mouse and human cardiac development," Pediatr. Research, vol. 76 (6) pp. 500-507 (2014). cited by applicant

Kwon, C. et al., "Canonical Wnt signaling is a positive regulator of mammalian cardiac progenitors," PNAS, USA, vol. 104(26), pp. 10894-10899 (2007). cited by applicant

Laflamme, M.A., et al., "Cardiomyocytes derived from human embryonic stem cells in pro-survival factors enhance function of infarcted rat hearts," Nat. Biotechnol., vol. 25(9), pp. 1015-1024 (2007). cited by applicant

Laflamme, M.A., et al., "Formation of human myocardium in the rat heart from human embryonic stem cells." Am. J. Pathol. vol. 167(3)pp. 663-671 (2005). cited by applicant

Laflamme, M.A., et al., "Heart regeneration," Nature, vol. 473(7347) pp. 326-335 (2011). cited by applicant

Lalit, P. et al., "Induced Pluripotent Stem Cells for Post-Myocardial Infarction Repair, Remarkable Opportunities and Challenges," Circulation Research, vol. 114, pp. 1328-1345 (2014). cited by applicant

Lancaster, M.A., et al., "Cerebral organoids model human brain development and microcephaly," Nature, vol. 501 (7467), pp. 373-379, (2013). cited by applicant

Laugwitz, K.-L., et al., "Postnatal isl1+ cardioblasts enter fully differentiated cardiomyocyte lineages," Nature 433, 647-653.(2005). cited by applicant

Lauss, M. et al. "Single inner cell masses yield embryonic stem cell lines differing in l1 expression and their developmental potential," Biochemical and Biophysical Research Communications, vol. 331:1577-1586 (2005). cited by applicant

Li, L. et al., The human homolog of rat Jagged1 expressed by marrow stroma inhibits differentiation of 32D cells through interaction with Notch1, Immunity, vol. 8(1) pp. 43-55 (1998). cited by applicant

Li, L., et al., "Alagille syndrome is caused by mutations in human Jagged1, which encodes a ligand for Notch1," Nat. Genet., vol. 16, 243-251.(1997). cited by applicant

Lian, Q. et al., "Establishing Clonal Cell Lines with Endothelial-Like Potential from CD9hi, SSEA-12 Cells in Embryonic Stem Cell-Derived Embryoid Bodies," PLOS One, Issue 1(1):e6, 10 pages (2006) doi:10.1371/journal.pone.0000006. cited by applicant

Lian, X. et al., "Directed cardiomyocyte differentiation from human pluripotent stem cells by modulating Wnt/?-catenin signaling under fully defined conditions," Nature Protocols, vol. 8(1):162-175 (2012). cited by applicant

Lian, X., et al., "Insulin inhibits cardiac mesoderm, not mesendoderm, formation during cardiac differentiation of human pluripotent stem cells and modulation of canonical wnt signaling can rescue this inhibition," Stem Cells, vol. 31 (3), pp. 447-457. (2013). cited by applicant

Lian, X.J., et al., "Robust cardiomyocyte differentiation from human pluripotent stem cells via temporal modulation of canonical Wnt signaling," PNAS, U. S. A., vol. 109(27), pp. E1848-E1857. (2012). cited by applicant

Lin, B. et al., High-purity enrichment of functional cardiovascular cells from human iPS cells, Cardiovascular Research, vol. 95(3), pp. 327-335 (Aug. 8, 2012). cited by applicant

Lin, Z. et al., "Strategies for cardiac regeneration and repair," Sci. Transl. Med., vol. 6, (239) 23 pages (2014). cited by applicant

Lui, K.O., et al., "Driving vascular endothelial cell fate of human multipotent Isl1+ heart progenitors with VEGF modified mRNA," Cell Res., vol. 23, pp. 1172-1186. (2013). cited by applicant

Marks, A.R., "Calcium cycling proteins and heart failure: mechanisms and therapeutics," J. Clin. Invest., vol. 123, pp. 46-52. (2013). cited by applicant

Marrink, J. et al., "TRA-1-60: a new serum marker in patients with germ-cell tumors," Int. J. Cancer, vol. 49:368-372 (1991). cited by applicant

Martin, U., "New muscle for old hearts: engineering tissue from pluripotent stem cells," Hum. Gene Ther., vol. 26, pp. 305-311. (2015). cited by applicant

Masino, A. et al., "Transcriptional Regulation of Cardiac Progenitor Cell Populations," Circ Res.,

vol. 95:389-397 (2004). cited by applicant

Masters, M., et al., "The epicardium signals the way towards heart regeneration," *Stem Cell Res.*, vol. 13, 683-692 (2014). cited by applicant

Masuda S., et al., "Eliminating residual iPS cells for safety in clinical Application," *Protein & Cell*, vol. 6 (7):469-471 (2015). cited by applicant

Masuda, S. et al., "Three-dimensional cardiac tissue fabrication based on cell sheet technology," *Adv. Drug Deliv. Rev.*, vol. 96, pp. 103-109 (2016). cited by applicant

Masumoto, H., et al., "Human iPS cell-engineered cardiac tissue sheets with cardiomyocytes and vascular cells for cardiac regeneration," *Sci. Rep.*, vol. 4 (6716) pp. 1-7 (2014). cited by applicant

Menasche, P. et al., "Human embryonic stem cell-derived cardiac progenitors for severe heart failure treatment: first clinical case report," *European Heart Journal*, vol. 36:2011-2017 (2015). cited by applicant

Menasche', P. et al., "Human embryonic stem cell-derived cardiac progenitors for severe heart failure treatment: first clinical case report," *European Heart Journal*, vol. 36, pp. 2011-2017 (2015). cited by applicant

Menon, V. et al., "Flow Cytometry Protocols for Surface and Intracellular Antigen Analyses of Neural Cell Types," *J. Vis. Exp.*, vol. 94 (e52241) 11 pages (2014) doi:10.3791/52241. cited by applicant

Min, J.Y. et al. "Significant improvement of heart function by cotransplantation of human mesenchymal stem cells and jetal cardiomyocytes in postinfarcted pigs," *Ann. Thorac. Surg.*, vol. 74 (5), pp. 1568-1575 (2002). cited by applicant

Molkentin and Houser, "Are Resident c-Kit+ Cardiac Stem Cells Really All That Are Needed to Mend a Broken Heart?," *Circ Res.*, vol. 113:1037-1039(2013). cited by applicant

Moretti, A., et al., "Multipotent embryonic isl1+ progenitor cells lead to cardiac, smooth muscle, and endothelial cell diversification," *Cell*, vol. 127(6), pp. 1151-1165 (2006). cited by applicant

Musunuru, K. et al., "Stem Cell Models of Cardiac Development and Disease," *Annu Rev Cell Dev Biol.*, vol. 26, p. 667-687 (2010). cited by applicant

Nazarov, I. et al., "Multipotent Stromal Stem Cells from Human Placenta Demonstrate High Therapeutic Potential," *Stem Cells Translational Medicine*, vol. 1: 359-372 (2012). cited by applicant

Niida, A. et al., "DKK1, a negative regulator of Wnt signaling, is a target of the b-catenin/TCF pathway," *Oncogene*, vol. 23: 8520-8526 (2004). cited by applicant

Niu, Y., et al., "Generation of gene-modified cynomolgus monkey via Cas9/RNA-mediated gene targeting in one-cell embryos," *Cell*, vol. 156, pp. 836-843. (2014). cited by applicant

Nsair, A. et al., "Characterization and Therapeutic Potential of Induced Pluripotent Stem Cell-Derived Cardiovascular Progenitor Cells," *PLOS One*, vol. 7(10): e45603, 12 pages (2012). cited by applicant

O'Brien, T.X., et al., "Positional specification of ventricular myosin light chain 2 expression in the primitive murine heart tube," *PNAS, U. S. A.*, vol. 90, pp. 5157-5161 (1993). cited by applicant

Oda, T. et al. "Mutations in the human Jagged1 gene are responsible for Alagille syndrome," *Nat. Genet.*, vol. 16, p. 235-242 (1997). cited by applicant

Oda, T. et al. Identification and cloning of the human homolog (JAG1) of the rat Jagged1 gene from the Alagille syndrome critical region at 20p12, *Genomics*, vol. 43(3) pp. 376-379 (1997). cited by applicant

Orchard, C., "T-tubule function in mammalian cardiac myocytes," *Circ. Res.*, vol. 92(11):1182-1192 (2003). cited by applicant

Orlic, D. et al., "Bone marrow cells regenerate infarcted myocardium" *Nature*, vol. 410, pp. 701-705 (2001). cited by applicant

Ott, H.C., et al., "Perfusion-decellularized matrix: using nature's platform to engineer a bioartificial heart," *Nat. Med.*, vol. 14, pp. 213-221, (2008). cited by applicant

Pagliuca, F.W., et al., "Generation of Functional Human Pancreatic beta Cells In Vitro," *Cell*, vol. 159(2), pp. 428-439 (2014). cited by applicant

Parikh, A. et al., "Signaling Pathways and Gene Regulatory Networks in Cardiomyocyte Differentiation," *Tissue Engineering: Part B*, vol. 21(4):377-392 (2015). cited by applicant

Patel, A.K., et al., "A defined synthetic substrate for serum free culture of human stem cell derived cardiomyocytes with improved functional maturity identified using combinatorial materials microarrays," *Biomaterials*, vol. 61, pp. 257-265 (2015). cited by applicant

Perin, E.C. et al., "Transendocardial, autologous bone marrow cell transplantation for severe, chronic ischemic heart failure," *Circulation*, vol. 107, pp. 2294-2302 (2003). cited by applicant

Pittenger, M.F., et al., "Mesenchymal Stem Cells and Their Potetial as Cardiac Therapeutics," *Circ. Res.* vol. 95, pp. 9-20 (2003). cited by applicant

Plein, A. et al., "Neural crest-derived SEMA3C activates endothelial NRP1 for cardiac outflow tract septation," *Journal of Clinical Investigation*, vol. 125 (7): 2661-2676 (2015). cited by applicant

Qyang, Y., et al. "The renewal and differentiation of Isl1+ cardiovascular progenitors are controlled by a Wnt/beta-catenin pathway," *Cell Stem Cell*, vol. 1, pp. 165-179. (2007). cited by applicant

Robitaille, J. et al. "Mutant frizzled-4 disrupts retinal angiogenesis in familial exudative vitreoretinopathy," *Nat. Genet.*, vol. 32, pp. 326-330 (2002). cited by applicant

Rodrigues, G. et al., "Purification of Human Induced Pluripotent Stem Cell-Derived Neural Precursors Using Magnetic Activated Cell Sorting," *Methods in Molecular Biology*, vol. 283:137-145 (2015). cited by applicant

Ruan, J.-L., et al., "Mechanical Stress Promotes Maturation of Human Myocardium from Pluripotent Stem Cell-Derived Progenitors," *Stem Cells*, vol. 33(7), pp. 2148-2157, (2015). cited by applicant

Rubin, L.L., "Stem cells and drug discovery: the beginning of a new era?" *Cell*, vol. 132(4), pp. 549-552. (2008). cited by applicant

Ruvinov, E., et al., "Alginate biomaterial for the treatment of myocardial infarction: Progress, translational strategies, and clinical outlook: From ocean algae to patient bedside," *Adv. Drug Deliv. Rev.*, vol. 96, pp. 54-76 (2016). cited by applicant

Sahara, M., et al., "Programming and reprogramming a human heart cell," *EMBO J.*, vol. 34(6), pp. 710-738 (2015). cited by applicant

Sandstedt, J. et al., "Human C-kit+CD45- cardiac stem cells are heterogeneous and display both cardiac and endothelial commitment by single-cell qPCR analysis," *Biochemical and Biophysical Research Communications*, vol. 443: 234-238 (2014). cited by applicant

Schaaf, S., et al., "Human engineered heart tissue as a versatile tool in basic research and preclinical toxicology," *PLoS One*, vol. 6(10), e26397, pp. 1-11 (2011). cited by applicant

Schiemann, W.P. et al., "Phosphorylation of the human leukemia inhibitory factor (LIF) receptor by mitogen-activated protein kinase and the regulation of LIF receptor function by heterologous receptor activation," *Proc. Natl.. Acad. Sci. USA*, vol. 92(12):5361-5365 (1995). cited by applicant

Schopperle, W.M. et al., "The TRA-1-60 and TRA-1-81 human pluripotent stem cell markers are expressed on podocalyxin in embryonal carcinoma," *Stem Cells*, vol. 25:723-730 (2007). cited by applicant

Schwan, J., et al., "Prospects for In Vitro Myofilament Maturation in Stem Cell-Derived Cardiac Myocytes, Biomarker," *Insights*, vol. 10, pp. 91-103 (2015). cited by applicant

Segers, V.F.M., et al., "Stem-cell therapy for cardiac disease," *Nature*, vol. 451, pp. 937-942 (2008). cited by applicant

Senyo, S.E., et al., "Mammalian heart renewal by pre-existing cardiomyocytes," *Nature*, vol. 493(7432), pp. 433-436. (2013). cited by applicant

Shiang, R. et al., "Mutations in the transmembrane domain of FGFR3 cause the most common genetic form of dwarfism, achondroplasia," *Cell*, vol. 78:335-343 (1994). cited by applicant

Shiba, Y. et al., "Allogeneic transplantation of iPS cell-derived cardiomyocytes regenerates primate hearts," *Nature*, vol. 538: 388-404 (2016). cited by applicant

Shultz, L.D., et al., "Human lymphoid and myeloid cell development in NOD/LtSz-scid IL2R gamma null mice engrafted with mobilized human hemopoietic stem cells," *J. Immunol.*, vol. 174, pp. 6477-6489 (2005). cited by applicant

Sinzou [Heart], Feb. 2014, vol. 46, No. 2, p. 170-176 (translator's note: no English language counterpart could be located). cited by applicant

Soh, B-S, et al. "Endothelin-1 supports clonal derivation and expansion of cardiovascular progenitors derived from human embryonic stem cells," *Nature Communications*, vol. 7 (10774) 10 pages (2016) doi: 10.1038/ncomms10774. cited by applicant

Später, D., et al., "How to make a cardiomyocyte," *Development*, vol. 141(23), pp. 4418-4431 (2014). cited by applicant

Später, D., et al., "A HCN4+ cardiomyogenic progenitor derived from the first heart field and human pluripotent stem cells," *Nat. Cell Biol.*, vol. 15, pp. 1098-1106 (2013). cited by applicant

Stamm, C. et al. "Autologous bone-marrow stem-cell transplantation for myocardial regeneration," *Lancet*, vol. 361, pp. 45-46 (2003). cited by applicant

Stepniewski, J. et al., "Induced pluripotent stem cells as a model for diabetes investigation," *Scientific Reports*, vol. 5 (8597): 14 pages (2015) DOI: 10.1038/srep08597. cited by applicant

Stevens, K.R. et al., "Physiological function and transplantation of scaffold-free and vascularized human cardiac muscle tissue," *PNAS*, vol. 106(39): 16568-16573 (2009). cited by applicant

Stevens, K.R. et al., "Physiological function and transplantation of scaffold-free and vascularized human cardiac muscle tissue," *Supporting Information, PNAS*, vol. 106(39): 12 pages (2009). cited by applicant

Sultana, N. et al., "Resident c-kit+ cells in the heart are not cardiac stem cells," *Nature Communications*, vol. 6 (8701) 10 pages (2015). cited by applicant

Tanaka, S. et al., "A novel frizzled gene identified in human esophageal carcinoma mediates APC/beta-catenin signals," *PNAS USA*, vol. 95, pp. 10164-10169 (1998). cited by applicant

Thomson, J.A. et al., "Embryonic stem cell lines derived from human blastocysts," *Science*, vol. 282, pp. 1145-1147 (1998). cited by applicant

Tulloch, N.L., et al., "Growth of engineered human myocardium with mechanical loading and vascular co-culture," *Circ. Res.*, vol. 109(1), pp. 47-59.(2011). cited by applicant

Van Berlo, J. et al., "An emerging consensus on cardiac regeneration," *Nat. Med.*, vol. 20(12): 1386-1393 (2014). cited by applicant

Van Berlo, J. et al., "c-kit+ cells minimally contribute cardiomyocytes to the heart," *Nature*, vol. 509: 337-351 (2014). cited by applicant

Van Laake, L.W., et al., "Human embryonic stem cell-derived cardiomyocytes survive and mature in the mouse heart and transiently improve function after myocardial infarction," *Stem Cell Res.*, vol. 1, pp. 9-24. (2007). cited by applicant

Vanhoof, D. et al., "Identification of Cell Surface Proteins for Antibody-Based Selection of Human Embryonic Stem Cell-Derived Cardiomyocytes," *Journal of Proteome Research*, vol. 9(3), pp. 1610-1618 (Mar. 2010). cited by applicant

Vunjak-Novakovic, G., et al., "Bioengineering heart muscle: a paradigm for regenerative medicine," *Annu. Rev. Biomed. Eng.*, vol. 13, pp. 245-267. (2011). cited by applicant

Wang, G., et al., "Modeling the mitochondrial cardiomyopathy of Barth syndrome with iPSC and heart-on-chip technologies," *Nat. Med.*, vol. 20(6), pp. 616-623 (2014). cited by applicant

Wang, J., et al., "Cardiomyopathy associated with microcirculation dysfunction in laminin alpha4 chain-deficient mice," *J. Biol. Chem.*, vol. 281, pp. 213-220 (2006). cited by applicant

Webber, M.J., et al., "A Perspective on the Clinical Translation of Scaffolds for Tissue Engineering," *Ann. Biomed. Eng.*, vol. 43, pp. 641-656. (2014). cited by applicant

Wiencierz et al. "Differential Expression Levels of Integrin $\alpha 6$ Enable the Selective Identification

and Isolation of Atrial and Ventricular Cardiomyocytes,” PLoS One, vol. 10(11):e0143538. (2015). cited by applicant

Wojakowski, W., et al. “The role of CXCR4/SDF-1, CD117/SCF, and c-met/HGF chemokine signalling in the mobilization of progenitor cells and the parameters of the left ventricular function, remodelling, and myocardial perfusion following acute myocardial infarction,” European Heart Journal Supplements, vol. 10 (Supplement K): K16-K23 (2008). cited by applicant

Yang, L., et al., “Human cardiovascular progenitor cells develop from a KDR+ embryonic-stem-cell-derived population,” Nature 453, 524-528. (2008). cited by applicant

Yang, Z., “Cardiac overexpression of A1-adenosine receptor protects intact mice against myocardial infarction,” Am J Physiol Heart Circ Physiol , vol. 282, pp. H949-H955 (2002). cited by applicant

Ye, L., et al., “Fabrication of a myocardial patch with cells differentiated from human-induced pluripotent stem cells,” Methods Mol. Biol., vol. 1299, pp. 103-114.(2015). cited by applicant

Yi, A. et al., “Pregenerative medicine: developmental paradigms in the biology of cardiovascular regeneration,” The Journal of Clinical Investigation, vol. 120(1) pp. 20-28 (2010). cited by applicant

Yin, L. et al., “Induction of Vascular Progenitor Cells from Endothelial Cells Stimulates Coronary Collateral Growth,” Circ Res., vol. 110(2): 241-252 (2012). cited by applicant

Yu, J. et al., “Human induced pluripotent stem cells free of vector and transgene sequences,” Science, vol. 324 (5928), pp. 797-801 (2009). cited by applicant

Zangi, L., et al., “Modified mRNA directs the fate of heart progenitor cells and induces vascular regeneration after myocardial infarction,” Nat. Biotechnol., vol. 31(10), pp. 898-907 (2013). cited by applicant

Zentilin et al. “Cardiomyocyte VEGFR-1 activation by VEGF-B induces compensatory hypertrophy and preserves cardiac function after myocardial infarction,” FASEB, vol. 24(5):1467-1478 (2010). cited by applicant

Zhang, J. et al., “Extracellular matrix promotes highly efficient cardiac differentiation of human pluripotent stem cells: The matrix sandwich method,” Circ. Res., vol. 111, pp. 1125-1136 (2012). cited by applicant

Zhou, B., et al., “Epicardial progenitors contribute to the cardiomyocyte lineage in the developing heart,” Nature, vol. 454, pp. 109-113 (2008). cited by applicant

Primary Examiner: Moloye; Titilayo

Attorney, Agent or Firm: Nelson Mullins Riley & Scarborough LLP

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATION (1) This application is a divisional of U.S. patent application Ser. No. 16/109,218, filed Aug. 22, 2018, which claims priority to, and the benefit of, U.S. Provisional Application No. 62/549,345, filed on Aug. 23, 2017, the entire contents of which is incorporated herein by reference.

BACKGROUND OF THE INVENTION

(1) Heart failure, predominantly caused by myocardial infarction, is the leading cause of death in both adults and children worldwide and is increasing exponentially worldwide (Bui, A. L. et al. (2011) *Nat. Rev. Cardiol.* 8:30-41). The disease is primarily driven by the loss of ventricular muscle that occurs during myocardial injury (Lin, Z. and Pu, W. T. (2014) *Sci. Transl. Med.* 6:239rv1) and is compounded by the negligible ability of the adult heart to mount a regenerative response

(Bergmann, O. et al. (2009) *Science* 324:98-102; Senyo, S. E. et al. (2013) *Nature* 493:433-436). Although heart transplantation can be curative, the markedly limited availability of human heart organ donors has led to a widespread unmet clinical need for a renewable source of pure, mature and functional human ventricular muscle tissue (Segers, V. F. M. and Lee, R. J. (2008) *Nature* 451:937-942; Später, D. et al. (2014) *Development* 141:4418-4431).

(2) Human pluripotent stem cells (hPSCs) offer the potential to generate large numbers of functional cardiomyocytes for potential clinical restoration of function in damaged or diseased hearts. Transplantation of stem cells into the heart to improve cardiac function and/or to enrich and regenerate damaged myocardium has been proposed (see e.g., U.S. Patent Publication 20040180043). Combination therapy, in which adult stem cells are administered in combination with treatment with growth factor proteins has also been proposed (see e.g., U.S. Patent Publication 20050214260).

(3) While cell transplantation into the heart offers a promising approach for improving cardiac function and regenerating heart tissue, the question of what type(s) of cells to transplant has been the subject of much investigation. Types of cells investigated for use in regenerating cardiac tissue include bone marrow cells (see e.g., Orlic, D. et al. (2001) *Nature* 410:701-705; Stamm, C. et al. (2003) *Lancet* 361:45-46; Perin, E. C. et al. (2003) *Circulation* 107:2294-2302), adult stem cells (see e.g., Jackson, K. A. et al. (2001) *J. Clin. Invest.* 107:1395-1402), bone marrow-derived mesenchymal stem cells (see e.g., Barbash, I. M. et al. (2003) *Circulation* 108:863; Pettinger, M. F. and Martin, B. J. (2003) *Circ. Res.* 95:9-20), bone marrow stromal cells (Bittira, B. et al. (2003) *Eur. J. Cardiothorac. Surg.* 24:393-398), a combination of mesenchymal stem cells and fetal cardiomyocytes (see e.g., Min, J. Y. et al. (2002) *Ann. Thorac. Surg.* 74:1568-1575) and a combination of bone marrow-derived mononuclear cells and bone marrow-derived mesenchymal stem cells (see e.g., U.S. Patent Publication 20080038229). Dedifferentiation of adult mammalian cardiomyocytes in vitro to generate cardiac stem cells for transplantation has also been proposed (see e.g., U.S. Patent Publication 20100093089).

(4) A significant advancement in the approach of cell transplantation to improve cardiac function and regenerate heart tissue was the identification and isolation of a family of multipotent cardiac progenitor cells that are capable of giving rise to cardiac myocytes, cardiac smooth muscle and cardiac endothelial cells (Cal, C. L. et al. (2003) *Dev. Cell.* 5:877-889; Moretti, A. et al. (2006) *Cell* 127:1151-1165; Bu, L. et al. (2009) *Nature* 460:113-117; U.S. Patent Publication 20060246446). These cardiac progenitor cells are characterized by the expression of the LIM homeodomain transcription factor Islet 1 (Isl1) and thus are referred to as Isl1⁺ cardiac progenitor cells. (Ibid). In contrast, Isl1 is not expressed in differentiated cardiac cells. Additional markers of the Isl1⁺ cardiac progenitor cells that arise later in differentiation than Isl1 have been described and include Nkx2.5 and flk1 (see e.g., U.S. Patent Publication 20100166714).

(5) The renewal and differentiation of Isl1⁺ cardiac progenitor cells has been shown to be regulated by a Wnt/beta-catenin signaling pathway (see e.g., Qyang, Y. et al. (2007) *Cell Stem Cell.* 1:165-179; Kwon, C. et al. (2007) *Proc. Natl. Acad. Sci. USA* 104:10894-10899). This has led to the development of methods to induce a pluripotent stem cell to enter the Isl1⁺ lineage and for expansion of the Isl1⁺ population through modulation of Wnt signaling (see e.g., Lian, X. et al. (2012) *Proc. Natl. Acad. Sci. USA* 109:E1848-57; Lian, X. et al. (2013) *Nat. Protoc.* 8:162-175; U.S. Patent Publication 20110033430; U.S. Patent Publication 20130189785).

(6) While human pluripotent stem cells hold great promise, a significant challenge has been the ability to move from simply differentiation of diverse cardiac cells to forming a larger scale pure 3D ventricular muscle tissue in vivo, which ultimately requires vascularization, assembly and alignment of an extracellular matrix, and maturation. Toward that end, a diverse population of cardiac cells (atrial, ventricular, pacemaker) has been coupled with artificial and decellurized matrices (Masumoto, H. et al. (2014) *Sci. Rep.* 4:5716; Ott, H. C. et al. (2008) *Nat. Med.* 14:213-221; Schaaf, S. et al. (2011) *PLoS One* 6:e26397), vascular cells and conduits (Tulloch, N. L. et al.

(2011) *Circ. Res.* 109:47-59) and cocktails of microRNAs (Gama-Garvalho, M. et al. (2014) *Cells* 3:996-1026) have been studied, but the goal remains elusive.

(7) While the identification of Isl1 as a marker of cardiac progenitor cells was a significant advance, since Isl1 is an intracellular protein it is not a suitable marker for use in isolating large quantities of viable cells. Rather, a cell surface marker(s) is still needed. Furthermore, Isl1 as a marker identifies a population that can differentiate into multiple cell types within the cardiac lineage, and thus there is still a need for markers that identify cardiac progenitor cells that are biased toward a particular cell type within the cardiac lineage, in particular for progenitor cells that differentiate into ventricular cells. Accordingly, there is still a great need in the art for additional markers of cardiac progenitor cells, in particular cell-surface markers of cardiac progenitor cells, that predominantly give rise to cardiomyocytes and that would allow for rapid isolation and large scale expansion of cardiomyogenic progenitor cells. Furthermore, there is still a great need in the art for methods and compositions for isolating cardiac ventricular progenitors, which differentiate into ventricular muscle cells in vivo, thereby allowing for transplantation of ventricular progenitors or ventricular muscle cells in vivo to enhance cardiac function.

SUMMARY OF THE INVENTION

(8) This invention describes the use of Neuropilin-1 (NRP1) as a cell surface marker for isolating human cardiac ventricular progenitor cells. Furthermore, additional cell surface markers suitable for use in isolating human cardiac ventricular progenitor cells are provided, as shown in Tables 1, 5 and 10. These human cardiac progenitor cells are biased toward the ventricular lineage such that they differentiate predominantly into ventricular muscle cells both in vitro and in vivo. That is, these cardiac progenitor cells can be cultured under conditions in vitro such that they are biased toward the ventricular lineage and thus are human ventricular progenitor (HVP) cells. Moreover, when introduced into the ventricular region of the heart in a subject, these progenitor cells differentiate almost exclusively into ventricular muscle cells that function according to their ventricular programming. In particular, the human ventricular progenitor cells provided herein utilize a cell autonomous pathway by which these cells can build a pure 3-D vascularized, functional and mature ventricular cell wall in vivo on the surface of normal murine kidney or heart, thereby allowing for organ-on-organ in vivo tissue engineering.

(9) Using single cell sequencing of cardiac progenitors at different stages of differentiation, a panel of genes differentially expressed in HVPs was identified, as described in Example 14 and shown in Table 10. Within this panel, day 5 progenitor cells were shown to express both Islet 1 (Isl1) and NRP1. This identification of a key cell surface marker of cardiac ventricular progenitor cells allows for easy and rapid isolation of the cells. Furthermore, determination of culture conditions for expansion and ventricular lineage bias of the cells allows for the preparation of large cultures (a billion or more cells) of a clonal population of cardiac ventricular progenitor cells. These cells can be used, for example, to improve function in a damaged heart in a subject, particularly damage in the ventricular region. The progenitor cells can be transplanted in vivo for differentiation into ventricular cells in situ or, alternatively, a heart muscle patch, comprising ventricular muscle cells, can be prepared in vitro from the progenitors for subsequent transplantation in vivo. The cells also can be used, for example, in in vitro toxicity screening assays to evaluate the cardiac toxicity of test compounds, as well as for biochemical studies to identify relevant pathways used in cardiac maturation and differentiation.

(10) Thus, the invention provides human cardiac ventricular progenitor cells in purified form. The human cardiac ventricular progenitors are capable of differentiation into ventricular muscle cells in vitro and in vivo. These progenitor cells can be expanded to large numbers of cells in vitro and when transplanted into the ventricular region of the heart in vivo they differentiate essentially exclusively into ventricular muscle cells. Still further, the cells have the capacity to migrate in vivo to different sites and, when transplanted in vivo the cells do what they are programmed to do as a ventricular cell (as opposed to a cardiac myocyte which simply contracts). Thus, the ventricular

progenitor cells can be grafted to native tissue to enhance ventricular function and have the ability to call in vasculature into the new ventricular tissue.

(11) Accordingly, in one aspect, the invention pertains to a method for isolating human cardiac ventricular progenitor cells, the method comprising: contacting a culture of cells containing human cardiac progenitor cells with one or more agents reactive with NRP1; and separating NRP1 reactive positive cells from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(12) In one embodiment, the human cardiac progenitor cells are contacted both with an agent reactive with NRP1 and with at least one second agent that binds to an HVP marker(s) (as described herein) to thereby separate NRP1/second agent reactive positive cells from non-reactive cells.

(13) Preferably, the human cardiac progenitor cells are Islet 1+ human cardiac progenitor cells. Preferably, the human cardiac progenitor cells are JAG1+, FZD4+, LIFR+, FGFR3+ and/or TNFSF9+. In another embodiment, the culture of cells is also contacted with at least one second agent; and NRP1 reactive/second agent reactive positive cells are separated from non-reactive cells to thereby isolate cardiac ventricular progenitor cells. In one embodiment, the at least one second agent binds a marker selected from JAG1+, FZD4+, LIFR+, FGFR3+ and/or TNFSF9+. The culture of cells can be simultaneously contacted with the agent reactive with NRP1 and the at least one second agent. Alternatively, the culture of cells can be contacted with the at least one second agent before contacting with the agent reactive with NRP1. Alternatively, the culture of cells can be contacted with the agent reactive with NRP1 before contacting with the at least one second agent. In another embodiment, the culture of cells is also negatively selected for lack of expression of at least one marker of pluripotent stem cells, such as TRA-1-60. In another embodiment, the human cardiac ventricular progenitor cells are further cultured and differentiated such that they express the ventricular marker MLC2v. In certain embodiments, the starting culture of cells containing human cardiac progenitor cells is derived from human embryonic stem cells or human induced pluripotent cells.

(14) In another aspect, the invention pertains to a method for isolating human cardiac ventricular progenitor cells, the method comprising: culturing human pluripotent stem cells under conditions that generate cardiac progenitor cells to obtain a cultured cell population; contacting the cultured cell population with one or more agents reactive with NRP1; and separating NRP1 reactive positive cells from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(15) In another embodiment, the culture of cells is also contacted with at least one second agent; and NRP1 reactive/second agent reactive positive cells are separated from non-reactive cells to thereby isolate cardiac ventricular progenitor cells. In one embodiment, the at least one second agent binds a marker selected from JAG1+, FZD4+, LIFR+, FGFR3+ and/or TNFSF9+. The culture of cells can be simultaneously contacted with the agent reactive with NRP1 and the at least one second agent. Alternatively, the culture of cells can be contacted with the at least one second agent before contacting with the agent reactive with NRP1. Alternatively, the culture of cells can be contacted with the agent reactive with NRP1 before contacting with the at least one second agent. In another embodiment, the culture of cells is also negatively selected for lack of expression of at least one marker of pluripotent stem cells, such as TRA-1-60. In another embodiment, the human cardiac ventricular progenitor cells are further cultured and differentiated such that they express the ventricular marker MLC2v. In certain embodiments, the starting culture of cells containing human cardiac progenitor cells is derived from human embryonic stem cells or human induced pluripotent cells.

(16) In the methods for isolating human cardiac ventricular progenitor cells, various types of agents that bind to NRP1 can be used as the agent(s) reactive with NRP1. For example, in one embodiment, the agent reactive with NRP1 is an anti-NRP1 antibody, such as a monoclonal antibody. In another embodiment, the agent reactive with NRP1 is a soluble NRP1 ligand, such as a NRP1 ligand fusion protein. For example, the agent reactive with NRP1 can comprise the NRP1

ligand VEGF-A or Sema3A, such as a soluble fusion protein (e.g., an Ig fusion protein).

(17) In the methods for isolating human cardiac ventricular progenitor cells, various types of separation methods can be used to separate NRP1 reactive positive cells from non-reactive cells. For example, in one embodiment, the reactive positive cells are separated from the non-reactive cells by fluorescence activated cell sorting (FACS). In another embodiment, the reactive positive cells are separated from the non-reactive cells by magnetic activated cells sorting (MACS).

(18) In yet another aspect, the invention pertains to a method of obtaining a clonal population of human cardiac ventricular progenitor cells, the method comprising: isolating a single NRP1+ human cardiac ventricular progenitor cell; and culturing the single NRP1+ human cardiac ventricular progenitor cell under conditions such that the cell is expanded to at least 1×10^9 cells to thereby obtain a clonal population of human cardiac ventricular progenitor cells.

(19) In one embodiment, the single NRP1+ human cardiac ventricular progenitor cell is Islet 1 positive, Nkx2.5 negative and flk1 negative at the time of initial culture. The single NRP1+ human cardiac ventricular progenitor cell can be isolated by methods such as those described above (e.g., FACS or MACS). The single NRP1+ human cardiac ventricular progenitor cell can be isolated using a reagent(s) reactive with NRP1, such as those described above (e.g., anti-NRP1 antibodies, soluble NRP1 ligands, such as ligand fusion proteins). Upon further culture and differentiation, the clonal population of human cardiac ventricular progenitor cells can express the ventricular marker MLCV2.

(20) In a preferred embodiment, the single NRP1+ human cardiac ventricular progenitor cell is cultured in vitro under conditions such that the cell is biased toward ventricular differentiation. For example, the single NRP1+ human cardiac ventricular progenitor cell can be cultured in Cardiac Progenitor Culture (CPC) medium (80% advanced DMEM/F12 supplemented with 20% KnockOut Serum Replacement, 2.5 mM GlutaMax and 100 μ g/ml Vitamin C), which allows for differentiation of the cells into ventricular cells expressing the MLC2v ventricular marker. In various embodiments, the single NRP1+ human cardiac ventricular progenitor cell is expanded to a clonal population of, for example, at least 1×10^9 cells, at least 2×10^9 cells, at least 5×10^9 cells or at least 10×10^9 cells.

(21) Accordingly in another aspect, the invention pertains to a clonal population of isolated NRP1+ human cardiac ventricular progenitor cells. In various embodiments, this clonal population comprises, for example, at least 1×10^9 cells, at least 2×10^9 cells, at least 5×10^9 cells or at least 10×10^9 cells. In a preferred embodiment, this clonal population comprises at least 1×10^9 NRP1+ human cardiac ventricular progenitor cells.

(22) In yet another aspect, the invention pertains to a method of enhancing cardiac function in a subject using the NRP1+ human cardiac ventricular progenitor cells described herein. For example, in one embodiment, the invention provides a method of enhancing cardiac function in a subject, the method comprising administering a pharmaceutical composition comprising a clonal population NRP1+ human cardiac ventricular progenitor cells, such as a clonal population of at least at least 1×10^9 cells, at least 2×10^9 cells, at least 5×10^9 cells or at least 10×10^9 cells. In one embodiment, the clonal population is administered directly into the heart of the subject. For example, the clonal population can be administered directly into a ventricular region of the heart of the subject. In one embodiment, the pharmaceutical composition administered to the subject comprises the clonal population formulated onto a three dimensional matrix, such as a heart muscle patch comprising ventricular muscle cells. The subject is one in need of enhancement of cardiac function, for example someone who has suffered a myocardial infarction or someone who has a congenital heart disorder.

(23) In yet another aspect, the invention pertains to a method for generating human ventricular tissue comprising: transplanting NRP1+ human cardiac ventricular progenitor cells into an organ of a non-human animal; and allowing the progenitor cells to grow in vivo such that human ventricular tissue is generated.

(24) The non-human animal can be, for example, an immunodeficient mouse. The organ can be, for example, the kidney (e.g., the cells are transplanted under the kidney capsule) or the heart. In one embodiment, the cells are transplanted at a time when one, two, three, four or five of the following cell marker patterns are present: (i) after peak of cardiac mesoderm formation; (ii) at time of peak Islet-1 expression; (iii) before peak of NKX2.5 expression; (iv) before peak expression of downstream genes MEF-2 and TBX-1; and (v) before expression of differentiated contractile protein genes. In one embodiment, the cells are transplanted between day 5 and day 7 (inclusive) of in vitro culture of human pluripotent stem cells under conditions to generate human ventricular progenitor cells. In another embodiment, the cells are transplanted on day 6 of in vitro culture of human pluripotent stem cells under conditions to generate human ventricular progenitor cells. The method can further include harvesting the human ventricular tissue generated in the non-human animal.

(25) In still another aspect of the invention, the human cardiac ventricular progenitor cells described herein can be used in screening assays to evaluate the cardiac toxicity of a test compound. Accordingly, the invention provides a method of screening for cardiac toxicity of test compound, the method comprising: providing NRP1+ cardiac ventricular progenitor cells; contacting the cells with the test compound; and measuring toxicity of the test compound for the cells, wherein toxicity of the test compound for the cells indicates cardiac toxicity of the test compound. The toxicity of the test compound for the cells can be measured, for example, by assessing cell viability or other physiological parameters of the cell.

(26) Culturing methods for generating human ventricular progenitor cells are also provided. For example, in one embodiment, the invention pertains to a method of generating human ventricular progenitors (HVPs) comprising: culturing human pluripotent stem cells (hPSCs) in a medium comprising CHIR98014 such that cells expressing cardiac mesodermal markers are generated; and culturing the cells expressing cardiac mesodermal markers in a medium comprising Wnt-059 such that HVPs are generated.

(27) Other features and advantages of the invention will be apparent from the following detailed description and claims.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) FIG. 1 is a schematic diagram of an exemplary culturing protocol for generating human Isl1+ cardiomyogenic progenitor cells from human pluripotent stem cells (hPSCs).

(2) FIG. 2 shows the results of Western blot analysis of protein expression during cardiac differentiation of hPSCs, showing expression of Isl1, Nkx2.5 and cTn1. GAPDH was used as a control.

(3) FIG. 3 shows the results of flow cytometry analysis of cardiomyogenic progenitor cells, showing expression of Isl1 on cells at day 6 of differentiation.

(4) FIG. 4 shows the results of double staining flow cytometry analysis of cardiomyogenic progenitor cells, showing coexpression of Isl1 and Jag1 on cells at day 6 of differentiation.

(5) FIG. 5 shows the results of Western blot analysis of protein expression during cardiac differentiation of hPSCs, showing expression of FZD4. GAPDH was used as a control.

(6) FIG. 6 shows the results of double staining flow cytometry analysis of cardiomyogenic progenitor cells, showing coexpression of Isl1 and FZD4 on cells at day 5 of differentiation.

(7) FIG. 7 is a schematic diagram of the generation of human ventricular progenitor (HVP) cells, their ultimate differentiation into ventricular myocytes, their antibody purification and their use in transplantation.

(8) FIGS. 8A and 8B are schematic diagrams of the transplantation of HPVs into the renal capsule

- (FIG. 8A) or intra-myocardially (FIG. 8B) for organ-on-organ tissue engineering.
- (9) FIG. 9 shows the results of double staining flow cytometry analysis of human ventricular progenitor (HVP) cells, showing coexpression of Isl1 and LIFR on the cells.
- (10) FIGS. 10A and 10B show the results of flow cytometry analysis of the expression of LIFR and FGFR3 on human ventricular progenitor cells (FIG. 10B) as compared to undifferentiated embryonic stem (ES) cells (FIG. 10A).
- (11) FIG. 11 is a tSNE plot of day 5 cells from single cell sequencing, showing two clusters of cells based on differential gene expression, labeled 0 and 1.
- (12) FIGS. 12A and 12B are tSNE plots of Isl1 expression (FIG. 12A) and NRP1 expression (FIG. 12B) on day 5 cells from single cell sequencing. Dark grey denotes high expression, middle grey denotes low expression and light grey denotes no expression.
- (13) FIG. 13 is a violin plot of expression levels of Isl1 and NRP1 in clusters of day 5 cells from single cell sequencing, showing that NRP1+ and ISL1+ cells are in the same cluster, since they have a similar gene expression profile.
- (14) FIGS. 14A, 14B and 14C show the results of flow cytometric analysis of the expression of NRP1 (FIG. 14A), TRA-1-60 (FIG. 14B) and both NRP1 and TRA-1-60 (FIG. 14C) on day 6 HVPs generated from H9 stem cells.
- (15) FIGS. 15A, 15B and 15C show the results of flow cytometric analysis of the expression of NRP1 (FIG. 15A), ISL1 (FIG. 15B) and both NRP1 and ISL1 (FIG. 15C) on day 6 HVPs generated from H9 stem cells.

DETAILED DESCRIPTION OF THE INVENTION

- (16) The invention provides methods of isolating human cardiomyogenic progenitor cells, in particular cells that are biased to the ventricular lineage, as well as isolated clonal populations of such progenitor cells, based on the discovery that NRP1 is a cell surface marker for cardiac ventricular progenitor cells (HVPs). Additional suitable cells surface markers for HVPs are shown in, for example, Tables 1, 5 and 10. In vitro and in vivo uses for these cardiac ventricular progenitor cells are also provided.
- (17) HVPs have previously been shown to express cell surface markers such as JAG1, FZD4, LIFR, FGFR3 and/or TNFSF9, as well as the intracellular marker Islet 1 (see U.S. Ser. No. 14/832,324, filed Aug. 21, 2015, and U.S. Ser. No. 14/984,783, filed Dec. 30, 2015, the entire contents of each of which are expressly incorporated herein by reference). Furthermore, various positive and negative engraftment markers of HVPs have been identified (see U.S. Ser. No. 15/433,713, filed Feb. 15, 2017, the entire contents of which is expressly incorporated herein by reference).
- (18) In order that the present invention may be more readily understood, certain terms are first defined. Additional definitions are set forth throughout the detailed description.
- (19) As used herein, the terms “Neuropilin-1”, “NRP1” are used interchangeably to refer to a protein known in the art that has been described in, for example, Soker, S. et al. (1998) *Cell* 92:735-745 and Rossignol, M. et al. (2000) *Genomics* 70:211-222. NRP1 is also referred to in the art as BDCA4, VEGF165R and CD304. A non-limiting example of an NRP1 protein is the human protein having the amino acid sequence set forth in Genbank Accession Number NP_003864.4.
- (20) As used herein, the term “stem cells” is used in a broad sense and includes traditional stem cells, progenitor cells, pre-progenitor cells, reserve cells, and the like. The term “stem cell” or “progenitor” are used interchangeably herein, and refer to an undifferentiated cell which is capable of proliferation and giving rise to more progenitor cells having the ability to generate a large number of mother cells that can in turn give rise to differentiated, or differentiable daughter cells. The daughter cells themselves can be induced to proliferate and produce progeny that subsequently differentiate into one or more mature cell types, while also retaining one or more cells with parental developmental potential. The term “stem cell” refers then, to a cell with the capacity or potential, under particular circumstances, to differentiate to a more specialized or differentiated phenotype,

and which retains the capacity, under certain circumstances, to proliferate without substantially differentiating. In one embodiment, the term progenitor or stem cell refers to a generalized mother cell whose descendants (progeny) specialize, often in different directions, by differentiation, e.g., by acquiring completely individual characters, as occurs in progressive diversification of embryonic cells and tissues. Cellular differentiation is a complex process typically occurring through many cell divisions. A differentiated cell may derive from a multipotent cell which itself is derived from a multipotent cell, and so on. While each of these multipotent cells may be considered stem cells, the range of cell types each can give rise to may vary considerably. Some differentiated cells also have the capacity to give rise to cells of greater developmental potential. Such capacity may be natural or may be induced artificially upon treatment with various factors. In many biological instances, stem cells are also “multipotent” because they can produce progeny of more than one distinct cell type, but this is not required for “stem-ness.” Self-renewal is the other classical part of the stem cell definition, and it is essential as used in this document. In theory, self-renewal can occur by either of two major mechanisms. Stem cells may divide asymmetrically, with one daughter retaining the stem state and the other daughter expressing some distinct other specific function and phenotype. Alternatively, some of the stem cells in a population can divide symmetrically into two stems, thus maintaining some stem cells in the population as a whole, while other cells in the population give rise to differentiated progeny only. Formally, it is possible that cells that begin as stem cells might proceed toward a differentiated phenotype, but then “reverse” and re-express the stem cell phenotype, a term often referred to as “dedifferentiation”.

(21) The term “progenitor cell” is used herein to refer to cells that have a cellular phenotype that is more primitive (e.g., is at an earlier step along a developmental pathway or progression than is a fully differentiated cell) relative to a cell which it can give rise to by differentiation. Often, progenitor cells also have significant or very high proliferative potential. Progenitor cells can give rise to multiple distinct differentiated cell types or to a single differentiated cell type, depending on the developmental pathway and on the environment in which the cells develop and differentiate.

(22) The term “pluripotent” as used herein refers to a cell with the capacity, under different conditions, to differentiate to cell types characteristic of all three germ cell layers (endoderm, mesoderm and ectoderm). Pluripotent cells are characterized primarily by their ability to differentiate to all three germ layers, using, for example, a nude mouse and teratomas formation assay. Pluripotency is also evidenced by the expression of embryonic stem (ES) cell markers, although the preferred test for pluripotency is the demonstration of the capacity to differentiate into cells of each of the three germ layers. In some embodiments, a pluripotent cell is an undifferentiated cell.

(23) The term “pluripotency” or a “pluripotent state” as used herein refers to a cell with the ability to differentiate into all three embryonic germ layers: endoderm (gut tissue), mesoderm (including blood, muscle, and vessels), and ectoderm (such as skin and nerve), and typically has the potential to divide in vitro for a long period of time, e.g., greater than one year or more than 30 passages.

(24) The term “multipotent” when used in reference to a “multipotent cell” refers to a cell that is able to differentiate into some but not all of the cells derived from all three germ layers. Thus, a multipotent cell is a partially differentiated cell. Multipotent cells are well known in the art, and examples of multipotent cells include adult stem cells, such as for example, hematopoietic stem cells and neural stem cells. Multipotent means a stem cell may form many types of cells in a given lineage, but not cells of other lineages. For example, a multipotent blood stem cell can form the many different types of blood cells (red, white, platelets, etc.), but it cannot form neurons.

(25) The term “embryonic stem cell” or “ES cell” or “ESC” are used interchangeably herein and refer to the pluripotent stem cells of the inner cell mass of the embryonic blastocyst (see U.S. Pat. Nos. 5,843,780, 6,200,806, which are incorporated herein by reference). Such cells can similarly be obtained from the inner cell mass of blastocysts derived from somatic cell nuclear transfer (see, for example, U.S. Pat. Nos. 5,945,577, 5,994,619, 6,235,970, which are incorporated herein by

reference). The distinguishing characteristics of an embryonic stem cell define an embryonic stem cell phenotype. Accordingly, a cell has the phenotype of an embryonic stem cell if it possesses one or more of the unique characteristics of an embryonic stem cell such that that cell can be distinguished from other cells. Exemplary distinguishing embryonic stem cell characteristics include, without limitation, gene expression profile, proliferative capacity, differentiation capacity, karyotype, responsiveness to particular culture conditions, and the like. In some embodiments, an ES cell can be obtained without destroying the embryo, for example, without destroying a human embryo.

(26) The term “adult stem cell” or “ASC” is used to refer to any multipotent stem cell derived from non-embryonic tissue, including fetal, juvenile, and adult tissue. Stem cells have been isolated from a wide variety of adult tissues including blood, bone marrow, brain, olfactory epithelium, skin, pancreas, skeletal muscle, and cardiac muscle. Each of these stem cells can be characterized based on gene expression, factor responsiveness, and morphology in culture. Exemplary adult stem cells include neural stem cells, neural crest stem cells, mesenchymal stem cells, hematopoietic stem cells, and pancreatic stem cells. As indicated above, stem cells have been found resident in virtually every tissue. Accordingly, the present invention appreciates that stem cell populations can be isolated from virtually any animal tissue.

(27) The term “human pluripotent stem cell” (abbreviated as hPSC), as used herein, refers to a human cell that has the capacity to differentiate into a variety of different cell types as discussed above regarding stem cells and pluripotency. Human pluripotent human stem cells include, for example, induced pluripotent stem cells (iPSC) and human embryonic stem cells, such as ES cell lines.

(28) The term “human cardiac progenitor cell”, as used herein, refers to a human progenitor cell that is committed to the cardiac lineage and that has the capacity to differentiate into all three cardiac lineage cells (cardiac muscle cells, endothelial cells and smooth muscle cells). A culture of human cardiac progenitor cells can be obtained by, for example, culturing stem cells under conditions that bias the stem cells toward differentiation to the cardiac lineage. In certain embodiments, the stem cells that are cultured to generate human cardiac progenitor cells are human embryonic stem cells or human induced pluripotent cells. In certain embodiments, c-kit⁺ adult progenitor cells are explicitly excluded for use in generating human cardiac progenitor cells.

(29) The term “human cardiomyogenic progenitor cell”, as used herein, refers to a human progenitor cell that is committed to the cardiac lineage and that predominantly differentiates into cardiac muscle cells (i.e., more than 50% of the differentiated cells, preferably more than 60%, 70%, 80% or 90% of the differentiated cells, derived from the progenitor cells are cardiac muscle cells).

(30) The term “cardiac ventricular progenitor cell”, as used herein, refers to a progenitor cell that is committed to the cardiac lineage and that predominantly differentiates into cardiac ventricular muscle cells (i.e., more than 50% of the differentiated cells, preferably more than 60%, 70%, 80% or 90% of the differentiated cells, derived from the progenitor cells are cardiac ventricular muscle cells). This type of cell is also referred to herein as a human ventricular progenitor, or HVP, cell.

(31) The term “cardiomyocyte” refers to a muscle cell of the heart (e.g. a cardiac muscle cell). A cardiomyocyte will generally express on its cell surface and/or in the cytoplasm one or more cardiac-specific marker. Suitable cardiomyocyte-specific markers include, but are not limited to, cardiac troponin I, cardiac troponin-C, tropomyosin, caveolin-3, GATA-4, myosin heavy chain, myosin light chain-2a, myosin light chain-2v, ryanodine receptor, and atrial natriuretic factor.

(32) The term “derived from” used in the context of a cell derived from another cell means that a cell has stemmed (e.g. changed from or produced by) a cell that is a different cell type. The term “derived from” also refers to cells that have been differentiated from a progenitor cell.

(33) The term “Isl1⁺ cardiac progenitor cell”, as used herein, refers to a human progenitor cell that is committed to the cardiac lineage and that expresses Islet 1.

- (34) The term “Isl1+ NRP1+ cardiac progenitor cell”, as used herein, refers to a human progenitor cell that is committed to the cardiac lineage and that expresses both Islet 1 and NRP1.
- (35) With respect to cells in cell cultures or in cell populations, the term “substantially free of” means that the specified cell type of which the cell culture or cell population is free, is present in an amount of less than about 10%, less than about 9%, less than about 8%, less than about 7%, less than about 6%, less than about 5%, less than about 4%, less than about 3%, less than about 2% or less than about 1% of the total number of cells present in the cell culture or cell population.
- (36) In the context of cell ontogeny, the adjective “differentiated”, or “differentiating” is a relative term. A “differentiated cell” is a cell that has progressed further down the developmental pathway than the cell it is being compared with. Thus, stem cells can differentiate to lineage-restricted precursor cells (such as a mesodermal stem cell), which in turn can differentiate into other types of precursor cells further down the pathway (such as an cardiomyocyte precursor), and then to an end-stage differentiated cell, which plays a characteristic role in a certain tissue type, and may or may not retain the capacity to proliferate further.
- (37) The term “differentiation” in the present context means the formation of cells expressing markers known to be associated with cells that are more specialized and closer to becoming terminally differentiated cells incapable of further differentiation. The pathway along which cells progress from a less committed cell, to a cell that is increasingly committed to a particular cell type, and eventually to a terminally differentiated cell is referred to as progressive differentiation or progressive commitment. Cell which are more specialized (e.g., have begun to progress along a path of progressive differentiation) but not yet terminally differentiated are referred to as partially differentiated. Differentiation is a developmental process whereby cells assume a specialized phenotype, e.g., acquire one or more characteristics or functions distinct from other cell types. In some cases, the differentiated phenotype refers to a cell phenotype that is at the mature endpoint in some developmental pathway (a so called terminally differentiated cell). In many, but not all tissues, the process of differentiation is coupled with exit from the cell cycle. In these cases, the terminally differentiated cells lose or greatly restrict their capacity to proliferate. However, we note that in the context of this specification, the terms “differentiation” or “differentiated” refer to cells that are more specialized in their fate or function than at a previous point in their development, and includes both cells that are terminally differentiated and cells that, although not terminally differentiated, are more specialized than at a previous point in their development. The development of a cell from an uncommitted cell (for example, a stem cell), to a cell with an increasing degree of commitment to a particular differentiated cell type, and finally to a terminally differentiated cell is known as progressive differentiation or progressive commitment. A cell that is “differentiated” relative to a progenitor cell has one or more phenotypic differences relative to that progenitor cell. Phenotypic differences include, but are not limited to morphologic differences and differences in gene expression and biological activity, including not only the presence or absence of an expressed marker, but also differences in the amount of a marker and differences in the co-expression patterns of a set of markers.
- (38) The term “differentiation” as used herein refers to the cellular development of a cell from a primitive stage towards a more mature (i.e. less primitive) cell.
- (39) As used herein, “proliferating” and “proliferation” refers to an increase in the number of cells in a population (growth) by means of cell division. Cell proliferation is generally understood to result from the coordinated activation of multiple signal transduction pathways in response to the environment, including growth factors and other mitogens. Cell proliferation may also be promoted by release from the actions of intra- or extracellular signals and mechanisms that block or negatively affect cell proliferation.
- (40) The terms “renewal” or “self-renewal” or “proliferation” are used interchangeably herein, and refers to a process of a cell making more copies of itself (e.g. duplication) of the cell. In some embodiments, cells are capable of renewal of themselves by dividing into the same undifferentiated

cells (e.g. progenitor cell type) over long periods, and/or many months to years. In some instances, proliferation refers to the expansion of cells by the repeated division of single cells into two identical daughter cells.

(41) The term “lineages” as used herein refers to a term to describe cells with a common ancestry or cells with a common developmental fate, for example cells that have a developmental fate to develop into ventricular cardiomyocytes.

(42) The term “clonal population”, as used herein, refers to a population of cells that is derived from the outgrowth of a single cell. That is, the cells within the clonal population are all progeny of a single cell that was used to seed the clonal population.

(43) The term “media” as referred to herein is a medium for maintaining a tissue or cell population, or culturing a cell population (e.g. “culture media”) containing nutrients that maintain cell viability and support proliferation. The cell culture medium may contain any of the following in an appropriate combination: salt(s), buffer(s), amino acids, glucose or other sugar(s), antibiotics, serum or serum replacement, and other components such as peptide growth factors, etc. Cell culture media ordinarily used for particular cell types are known to those skilled in the art.

(44) The term “phenotype” refers to one or a number of total biological characteristics that define the cell or organism under a particular set of environmental conditions and factors, regardless of the actual genotype.

(45) A “marker” as used herein describes the characteristics and/or phenotype of a cell. Markers can be used for selection of cells comprising characteristics of interest. Markers will vary with specific cells. Markers are characteristics, whether morphological, functional or biochemical (enzymatic) characteristics particular to a cell type, or molecules expressed by the cell type. Preferably, such markers are proteins, and more preferably, possess an epitope for antibodies or other binding molecules available in the art. However, a marker may consist of any molecule found in a cell including, but not limited to, proteins (peptides and polypeptides), lipids, polysaccharides, nucleic acids and steroids. Examples of morphological characteristics or traits include, but are not limited to, shape, size, and nuclear to cytoplasmic ratio. Examples of functional characteristics or traits include, but are not limited to, the ability to adhere to particular substrates, ability to incorporate or exclude particular dyes, ability to migrate under particular conditions, and the ability to differentiate along particular lineages. Markers may be detected by any method available to one of skill in the art.

(46) The term “isolated cell” as used herein refers to a cell that has been removed from an organism in which it was originally found or a descendant of such a cell. Optionally the cell has been cultured in vitro, e.g., in the presence of other cells. Optionally the cell is later introduced into a second organism or re-introduced into the organism from which it (or the cell from which it is descended) was isolated.

(47) The term “isolated population” with respect to an isolated population of cells as used herein refers to a population of cells that has been removed and separated from a mixed or heterogeneous population of cells. In some embodiments, an isolated population is a substantially pure population of cells as compared to the heterogeneous population from which the cells were isolated or enriched from.

(48) The term “substantially pure”, with respect to a particular cell population, refers to a population of cells that is at least about 75%, preferably at least about 85%, more preferably at least about 90%, and most preferably at least about 95% pure, with respect to the cells making up a total cell population.

(49) The terms “subject” and “individual” are used interchangeably herein, and refer to an animal, for example a human, to whom cardiac ventricular progenitor cells as disclosed herein can be implanted into, for e.g. treatment, which in some embodiments encompasses prophylactic treatment or for a disease model, with methods and compositions described herein, is or are provided. For treatment of disease states that are specific for a specific animal such as a human subject, the term

“subject” refers to that specific animal. The terms “non-human animals” and “non-human mammals” are used interchangeably herein, and include mammals such as rats, mice, rabbits, sheep, cats, dogs, cows, pigs, and non-human primates. The term “subject” also encompasses any vertebrate including but not limited to mammals, reptiles, amphibians and fish. However, advantageously, the subject is a mammal such as a human, or other mammals such as a domesticated mammal, e.g. dog, cat, horse, and the like, or production mammal, e.g. cow, sheep, pig, and the like are also encompassed in the term subject.

(50) As used herein, the term “recipient” refers to a subject that will receive a transplanted organ, tissue or cell.

(51) The term “three-dimensional matrix” or “scaffold” or “matrices” as used herein refers in the broad sense to a composition comprising a biocompatible matrix, scaffold, or the like. The three-dimensional matrix may be liquid, gel, semi-solid, or solid at 25° C. The three-dimensional matrix may be biodegradable or non-biodegradable. In some embodiments, the three-dimensional matrix is biocompatible, or bioresorbable or bioreplacable. Exemplary three-dimensional matrices include polymers and hydrogels comprising collagen, fibrin, chitosan, MATRIGEL™, polyethylene glycol, dextrans including chemically crosslinkable or photocrosslinkable dextrans, processed tissue matrix such as submucosal tissue and the like. In certain embodiments, the three-dimensional matrix comprises allogeneic components, autologous components, or both allogeneic components and autologous components. In certain embodiments, the three-dimensional matrix comprises synthetic or semi-synthetic materials. In certain embodiments, the three-dimensional matrix comprises a framework or support, such as a fibrin-derived scaffold.

(52) As used herein, the terms “administering,” “introducing” and “transplanting” are used interchangeably and refer to the placement of cardiomyogenic progenitor cells and/or cardiomyocytes differentiated as described herein into a subject by a method or route which results in at least partial localization of the cells at a desired site. The cells can be administered by any appropriate route that results in delivery to a desired location in the subject where at least a portion of the cells remain viable. The period of viability of the cells after administration to a subject can be as short as a few hours, e.g. twenty-four hours, to a few days, to as long as several years.

(53) The term “statistically significant” or “significantly” refers to statistical significance and generally means a two standard deviation (2SD) below normal, or lower, concentration of the marker. The term refers to statistical evidence that there is a difference. It is defined as the probability of making a decision to reject the null hypothesis when the null hypothesis is actually true. The decision is often made using the p-value. The term “substantially” or “predominantly” as used herein means a proportion of at least about 60%, or preferably at least about 70% or at least about 80%, or at least about 90%, at least about 95%, at least about 97% or at least about 99% or more, or any integer between 70% and 100%.

(54) The term “disease” or “disorder” is used interchangeably herein, and refers to any alternation in state of the body or of some of the organs, interrupting or disturbing the performance of the functions and/or causing symptoms such as discomfort, dysfunction, distress, or even death to the person afflicted or those in contact with a person. A disease or disorder can also related to a distemper, ailing, ailment, malady, disorder, sickness, illness, complaint, indisposition or affection.

(55) As used herein, the phrase “cardiovascular condition, disease or disorder” is intended to include all disorders characterized by insufficient, undesired or abnormal cardiac function, e.g. ischemic heart disease, hypertensive heart disease and pulmonary hypertensive heart disease, valvular disease, congenital heart disease and any condition which leads to congestive heart failure in a subject, particularly a human subject. Insufficient or abnormal cardiac function can be the result of disease, injury and/or aging. By way of background, a response to myocardial injury follows a well-defined path in which some cells die while others enter a state of hibernation where they are not yet dead but are dysfunctional. This is followed by infiltration of inflammatory cells, deposition of collagen as part of scarring, all of which happen in parallel with in-growth of new

blood vessels and a degree of continued cell death. As used herein, the term “ischemia” refers to any localized tissue ischemia due to reduction of the inflow of blood. The term “myocardial ischemia” refers to circulatory disturbances caused by coronary atherosclerosis and/or inadequate oxygen supply to the myocardium. For example, an acute myocardial infarction represents an irreversible ischemic insult to myocardial tissue. This insult results in an occlusive (e.g., thrombotic or embolic) event in the coronary circulation and produces an environment in which the myocardial metabolic demands exceed the supply of oxygen to the myocardial tissue.

(56) As used herein, the term “treating” or “treatment” are used interchangeably herein and refers to reducing or decreasing or alleviating or halting at least one adverse effect or symptom of a cardiovascular condition, disease or disorder, i.e., any disorder characterized by insufficient or undesired cardiac function. Adverse effects or symptoms of cardiac disorders are well-known in the art and include, but are not limited to, dyspnea, chest pain, palpitations, dizziness, syncope, edema, cyanosis, pallor, fatigue and death. In some embodiments, the term “treatment” as used herein refers to prophylactic treatment or preventative treatment to prevent the development of a symptom of a cardiovascular condition in a subject.

(57) Treatment is generally “effective” if one or more symptoms or clinical markers are reduced as that term is defined herein. Alternatively, a treatment is “effective” if the progression of a disease is reduced or halted. That is, “treatment” includes not just the improvement of symptoms or decrease of markers of the disease, but also a cessation or slowing of progress or worsening of a symptom that would be expected in absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, and remission (whether partial or total), whether detectable or undetectable. “Treatment” can also mean prolonging survival as compared to expected survival if not receiving treatment. Those in need of treatment include those already diagnosed with a cardiac condition, as well as those likely to develop a cardiac condition due to genetic susceptibility or other factors such as weight, diet and health. In some embodiments, the term to treat also encompasses preventative measures and/or prophylactic treatment, which includes administering a pharmaceutical composition as disclosed herein to prevent the onset of a disease or disorder.

(58) A therapeutically significant reduction in a symptom is, e.g. at least about 10%, at least about 20%, at least about 30%, at least about 40%, at least about 50%, at least about 60%, at least about 70%, at least about 80%, at least about 90%, at least about 100%, at least about 125%, at least about 150% or more in a measured parameter as compared to a control or non-treated subject. Measured or measurable parameters include clinically detectable markers of disease, for example, elevated or depressed levels of a biological marker, as well as parameters related to a clinically accepted scale of symptoms or markers for a disease or disorder. It will be understood, that the total daily usage of the compositions and formulations as disclosed herein will be decided by the attending physician within the scope of sound medical judgment. The exact amount required will vary depending on factors such as the type of disease being treated.

(59) With reference to the treatment of a cardiovascular condition or disease in a subject, the term “therapeutically effective amount” refers to the amount that is safe and sufficient to prevent or delay the development of a cardiovascular disease or disorder. The amount can thus cure or cause the cardiovascular disease or disorder to go into remission, slow the course of cardiovascular disease progression, slow or inhibit a symptom of a cardiovascular disease or disorder, slow or inhibit the establishment of secondary symptoms of a cardiovascular disease or disorder or inhibit the development of a secondary symptom of a cardiovascular disease or disorder. The effective amount for the treatment of the cardiovascular disease or disorder depends on the type of cardiovascular disease to be treated, the severity of the symptoms, the subject being treated, the age and general condition of the subject, the mode of administration and so forth. Thus, it is not

possible to specify the exact “effective amount”. However, for any given case, an appropriate “effective amount” can be determined by one of ordinary skill in the art using only routine experimentation. The efficacy of treatment can be judged by an ordinarily skilled practitioner, for example, efficacy can be assessed in animal models of a cardiovascular disease or disorder as discussed herein, for example treatment of a rodent with acute myocardial infarction or ischemia-reperfusion injury, and any treatment or administration of the compositions or formulations that leads to a decrease of at least one symptom of the cardiovascular disease or disorder as disclosed herein, for example, increased heart ejection fraction, decreased rate of heart failure, decreased infarct size, decreased associated morbidity (pulmonary edema, renal failure, arrhythmias) improved exercise tolerance or other quality of life measures, and decreased mortality indicates effective treatment. In embodiments where the compositions are used for the treatment of a cardiovascular disease or disorder, the efficacy of the composition can be judged using an experimental animal model of cardiovascular disease, e.g., animal models of ischemia-reperfusion injury (Headrick J P, Am J Physiol Heart circ Physiol 285; H1797; 2003) and animal models acute myocardial infarction. (Yang Z, Am J Physiol Heart Circ. Physiol 282:H949:2002; Guo Y, J Mol Cell Cardiol 33; 825-830, 2001). When using an experimental animal model, efficacy of treatment is evidenced when a reduction in a symptom of the cardiovascular disease or disorder, for example, a reduction in one or more symptom of dyspnea, chest pain, palpitations, dizziness, syncope, edema, cyanosis, pallor, fatigue and high blood pressure which occurs earlier in treated, versus untreated animals. By “earlier” is meant that a decrease, for example in the size of the tumor occurs at least 5% earlier, but preferably more, e.g., one day earlier, two days earlier, 3 days earlier, or more.

(60) As used herein, the term “treating” when used in reference to a treatment of a cardiovascular disease or disorder is used to refer to the reduction of a symptom and/or a biochemical marker of a cardiovascular disease or disorder, for example a reduction in at least one biochemical marker of a cardiovascular disease by at least about 10% would be considered an effective treatment. Examples of such biochemical markers of cardiovascular disease include a reduction of, for example, creatine phosphokinase (CPK), aspartate aminotransferase (AST), lactate dehydrogenase (LDH) in the blood, and/or a decrease in a symptom of cardiovascular disease and/or an improvement in blood flow and cardiac function as determined by someone of ordinary skill in the art as measured by electrocardiogram (ECG or EKG), or echocardiogram (heart ultrasound), Doppler ultrasound and nuclear medicine imaging. A reduction in a symptom of a cardiovascular disease by at least about 10% would also be considered effective treatment by the methods as disclosed herein. As alternative examples, a reduction in a symptom of cardiovascular disease, for example a reduction of at least one of the following; dyspnea, chest pain, palpitations, dizziness, syncope, edema, cyanosis etc. by at least about 10% or a cessation of such systems, or a reduction in the size one such symptom of a cardiovascular disease by at least about 10% would also be considered as affective treatments by the methods as disclosed herein. In some embodiments, it is preferred, but not required that the therapeutic agent actually eliminate the cardiovascular disease or disorder, rather just reduce a symptom to a manageable extent.

(61) Subjects amenable to treatment by the methods as disclosed herein can be identified by any method to diagnose myocardial infarction (commonly referred to as a heart attack) commonly known by persons of ordinary skill in the art are amenable to treatment using the methods as disclosed herein, and such diagnostic methods include, for example but are not limited to; (i) blood tests to detect levels of creatine phosphokinase (CPK), aspartate aminotransferase (AST), lactate dehydrogenase (LDH) and other enzymes released during myocardial infarction; (ii) electrocardiogram (ECG or EKG) which is a graphic recordation of cardiac activity, either on paper or a computer monitor. An ECG can be beneficial in detecting disease and/or damage; (iii) echocardiogram (heart ultrasound) used to investigate congenital heart disease and assessing abnormalities of the heart wall, including functional abnormalities of the heart wall, valves and

blood vessels; (iv) Doppler ultrasound can be used to measure blood flow across a heart valve; (v) nuclear medicine imaging (also referred to as radionuclide scanning in the art) allows visualization of the anatomy and function of an organ, and can be used to detect coronary artery disease, myocardial infarction, valve disease, heart transplant rejection, check the effectiveness of bypass surgery, or to select patients for angioplasty or coronary bypass graft.

(62) The terms “coronary artery disease” and “acute coronary syndrome” as used interchangeably herein, and refer to myocardial infarction refer to a cardiovascular condition, disease or disorder, include all disorders characterized by insufficient, undesired or abnormal cardiac function, e.g. ischemic heart disease, hypertensive heart disease and pulmonary hypertensive heart disease, valvular disease, congenital heart disease and any condition which leads to congestive heart failure in a subject, particularly a human subject. Insufficient or abnormal cardiac function can be the result of disease, injury and/or aging. By way of background, a response to myocardial injury follows a well-defined path in which some cells die while others enter a state of hibernation where they are not yet dead but are dysfunctional. This is followed by infiltration of inflammatory cells, deposition of collagen as part of scarring, all of which happen in parallel with in-growth of new blood vessels and a degree of continued cell death.

(63) As used herein, the term “ischemia” refers to any localized tissue ischemia due to reduction of the inflow of blood. The term “myocardial ischemia” refers to circulatory disturbances caused by coronary atherosclerosis and/or inadequate oxygen supply to the myocardium. For example, an acute myocardial infarction represents an irreversible ischemic insult to myocardial tissue. This insult results in an occlusive (e.g., thrombotic or embolic) event in the coronary circulation and produces an environment in which the myocardial metabolic demands exceed the supply of oxygen to the myocardial tissue.

(64) The terms “composition” or “pharmaceutical composition” used interchangeably herein refer to compositions or formulations that usually comprise an excipient, such as a pharmaceutically acceptable carrier that is conventional in the art and that is suitable for administration to mammals, and preferably humans or human cells. In some embodiments, pharmaceutical compositions can be specifically formulated for direct delivery to a target tissue or organ, for example, by direct injection or via catheter injection to a target tissue. In other embodiments, compositions can be specifically formulated for administration via one or more of a number of routes, including but not limited to, oral, ocular parenteral, intravenous, intraarterial, subcutaneous, intranasal, sublingual, intraspinal, intracerebroventricular, and the like. In addition, compositions for topical (e.g., oral mucosa, respiratory mucosa) and/or oral administration can form solutions, suspensions, tablets, pills, capsules, sustained-release formulations, oral rinses, or powders, as known in the art are described herein. The compositions also can include stabilizers and preservatives. For examples of carriers, stabilizers and adjuvants, University of the Sciences in Philadelphia (2005) Remington: The Science and Practice of Pharmacy with Facts and Comparisons, 21st Ed.

(65) As used herein, the terms “administering,” “introducing” and “transplanting” are used interchangeably and refer to the placement of a pharmaceutical composition comprising cardiomyogenic progenitor cells, or a composition comprising a population of differentiated cardiomyocytes (e.g., ventricular cardiomyocytes) as described herein, into a subject by a method or route which results in at least partial localization of the pharmaceutical composition, at a desired site or tissue location. In some embodiments, the pharmaceutical composition can be administered by any appropriate route which results in effective treatment in the subject, i.e. administration results in delivery to a desired location or tissue in the subject where at least a portion of the cells are located at a desired target tissue or target cell location.

(66) The phrases “parenteral administration” and “administered parenterally” as used herein mean modes of administration other than enteral and topical administration, usually by injection, and includes, without limitation, intravenous, intramuscular, intraarterial, intrathecal, intraventricular, intracapsular, intraorbital, intracardiac, intradermal, intraperitoneal, transtracheal, subcutaneous,

subcuticular, intraarticular, sub capsular, subarachnoid, intraspinal, intracerebro spinal, and intrasternal injection and infusion. The phrases “systemic administration,” “administered systemically,” “peripheral administration” and “administered peripherally” as used herein mean the administration of cardiovascular stem cells and/or their progeny and/or compound and/or other material other than directly into the cardiac tissue, such that it enters the animal's system and, thus, is subject to metabolism and other like processes, for example, subcutaneous or intravenous administration.

(67) The phrase “pharmaceutically acceptable” is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

(68) The phrase “pharmaceutically acceptable carrier” as used herein means a pharmaceutically acceptable material, composition or vehicle, such as a liquid or solid filler, diluent, excipient, solvent or encapsulating material, involved in carrying or transporting the subject agents from one organ, or portion of the body, to another organ, or portion of the body. Each carrier must be “acceptable” in the sense of being compatible with the other ingredients of the formulation.

(69) The term “drug” or “compound” or “test compound” as used herein refers to a chemical entity or biological product, or combination of chemical entities or biological products, administered to a subject to treat or prevent or control a disease or condition. The chemical entity or biological product is preferably, but not necessarily a low molecular weight compound, but may also be a larger compound, for example, an oligomer of nucleic acids, amino acids, or carbohydrates including without limitation proteins, oligonucleotides, ribozymes, DNazymes, glycoproteins, siRNAs, lipoproteins, aptamers, and modifications and combinations thereof.

(70) The term “transplantation” as used herein refers to introduction of new cells (e.g. reprogrammed cells), tissues (such as differentiated cells produced from reprogrammed cells), or organs into a host (i.e. transplant recipient or transplant subject).

(71) The term “agent reactive with NRP1”, as used herein, refers to an agent that binds to or otherwise interacts with NRP1. Preferably, the agent “specifically” binds or otherwise interacts with NRP1 such that it does not bind or interact with other non-NRP1 proteins.

(72) The term “antibody”, as used herein, includes whole antibodies and any antigen binding fragment (i.e., “antigen-binding portion”) or single chain thereof. An “antibody” refers, in one preferred embodiment, to a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds, or an antigen binding portion thereof. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as V.sub.H) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as V.sub.L) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The term “antigen-binding portion” of an antibody (or simply “antibody portion”), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen.

(73) The term “monoclonal antibody,” as used herein, refers to an antibody that displays a single binding specificity and affinity for a particular epitope.

(74) The term “human monoclonal antibody”, as used herein, refers to an antibody which displays a single binding specificity and which has variable and optional constant regions derived from human germline immunoglobulin sequences.

(75) The term “humanized monoclonal antibody”, as used herein, refers to an antibody which displays a single binding specificity and which has heavy and light chain CDR1, 2 and 3 from a non-human antibody (e.g., a mouse monoclonal antibody) grafted into human framework and constant regions.

(76) The term “chimeric monoclonal antibody”, as used herein, refers to an antibody which displays a single binding specificity and which has heavy and light chain variable regions from one species linked to constant regions from another species.

(77) The term “fusion protein”, as used herein, refers to a composite protein, typically made using recombinant DNA technology, in which two different proteins, or portions thereof, are operatively linked together. A non-limiting example is an Fc fusion protein in which a non-immunoglobulin protein is operatively linked to an immunoglobulin Fc region.

(78) Various aspects of the invention are described in further detail in the following subsections.

(79) Methods of Isolating Human Cardiac Ventricular Progenitor Cells

(80) In one aspect, the invention pertains to methods of isolating human cardiac ventricular progenitor cells. As described in the Examples, NRP1 has now been identified as a cell surface marker of human cardiac ventricular progenitor cells and thus this marker can be used to facilitate isolation of these progenitor cells. Alternative to NRP1, additional markers for human cardiac ventricular progenitor cells are provided herein, including the markers shown in Tables 1, 5 and 10. Any of the proteins in these tables that is a cell surface protein can be used in the isolation of HVPs as described herein.

(81) Accordingly, in one embodiment, the invention provides a method for isolating human cardiac ventricular progenitor cells, the method comprising: contacting a culture of human cells containing cardiac progenitor cells with one or more agents reactive with NRP1; and separating NRP1 reactive positive cells from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(82) Alternatively, after the contacting step, the method can comprise isolating NRP1 reactive positive cells from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(83) Also as described in the Examples, Islet 1 is a marker that is co-expressed with NRP1 by the cardiac ventricular progenitor cells and thus both markers can be used to facilitate isolation of these progenitor cells. Accordingly, in another embodiment of the above method, the culture of human cells is also contacted with an agent reactive with Islet 1; and NRP1 reactive/Islet 1 reactive positive cells are separated from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells. Alternatively or additionally, agents reactive with other HVP markers, including but not limited to JAG1, FZD4, LIFR, FGFR3 and/or TNFSF9, can be used in combination with NRP1 in the isolation of HVPs. Accordingly, the culture of human cells can be simultaneously contacted with the agent(s) reactive with NRP1 and/or at least one second agent reactive with an HVP marker(s), including but not limited to JAG1, FZD4, LIFR, FGFR3 and/or TNFSF9. In one embodiment, the culture of human cells is contacted with the agent reactive with NRP1 before contacting with the second agent(s) reactive with an HVP marker(s). In another embodiment, the culture of human cells is contacted with the second agent(s) reactive with an HVP marker(s) before contacting with the agent reactive with NRP1.

(84) In another embodiment, the method of isolating HVPs further comprises negatively selecting for at least one marker expressed on the surface of human pluripotent stem cells, such as TRA-1-60. The use of negative selection (in addition to positive selection for NRP1 expression, and/or expression of other positive HVP markers) ensures a rigorous definition of the HVP population as well as eliminating batch variation and potential teratoma-causing cells. Accordingly, in one embodiment, cardiac progenitor cells are selected for lack of expression of at least one marker expressed on the surface of human pluripotent stem cells (negative selection), such as TRA-1-60, to thereby isolate a highly purified population of HVPs. Non-limiting examples of markers expressed on the surface of human pluripotent stem cells that can be used for negative selection include: TRA-1-60, TRA-1-81, TRA-2-54, SSEA1, SSEA3, SSEA4, CD9, CD24, E-cadherin and Podocalyxin, and combinations thereof.

(85) In another embodiment, the invention provides a method for isolating human cardiac ventricular progenitor cells, the method comprising: culturing human pluripotent stem cells under conditions that generate cardiac progenitor cells to obtain a culture of cells; contacting the culture

of cells with one or more agents reactive with NRP1; and separating NRP1 reactive positive cells from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(86) Alternatively, after the culturing and contacting steps, the method can comprise isolating NRP1 from non-reactive cells to thereby isolate human cardiac ventricular progenitor cells.

(87) Alternatively or additionally, second agents reactive with other HVP markers, including but not limited to JAG1, FZD4, LIFR, FGFR3 and/or TNFSF9, can be used in combination with NRP1 in the isolation of HVPs. Accordingly, the culture of human cells can be simultaneously contacted with the agent(s) reactive with NRP1 and/or at least one second agent reactive with an HVP marker(s), including but not limited to JAG1, FZD4, LIFR, FGFR3 and/or TNFSF9. In one embodiment, the culture of human cells is contacted with the agent reactive with NRP1 before contacting with the second agent(s) reactive with an HVP marker(s). In another embodiment, the culture of human cells is contacted with the second agent(s) reactive with an HVP marker(s) before contacting with the agent reactive with NRP1.

(88) In another embodiment, the method of isolating HVPs further comprises negatively selecting for at least one marker expressed on the surface of human pluripotent stem cells, such as TRA-1-60. The use of negative selection (in addition to positive selection for NRP1 expression, and/or expression of other positive HVP markers) ensures a rigorous definition of the HVP population as well as eliminating batch variation and potential teratoma-causing cells. Accordingly, in one embodiment, cardiac progenitor cells are selected for lack of expression of at least one marker expressed on the surface of human pluripotent stem cells (negative selection), such as TRA-1-60, to thereby isolate a highly purified population of HVPs. Non-limiting examples of markers expressed on the surface of human pluripotent stem cells that can be used for negative selection include: TRA-1-60, TRA-1-81, TRA-2-54, SSEA1, SSEA3, SSEA4, CD9, CD24, E-cadherin and Podocalyxin, and combinations thereof.

(89) In a preferred embodiment, the agent reactive with NRP1 is an anti-NRP1 antibody, such as a monoclonal antibody. Non-limiting examples include murine, rabbit, human, humanized or chimeric monoclonal antibodies with binding specificity for NRP1. Anti-NRP1 monoclonal antibodies are commercially available in the art. Moreover, anti-NRP1 antibodies can be prepared using standard techniques well established in the art using NRP1 as the antigen.

(90) In another embodiment, the agent reactive with NRP1 is a NRP1 ligand, such as a soluble NRP1 ligand or a soluble NRP1 ligand fusion protein. Non-limiting examples of NRP1 ligands include VEGF-A and Sema3A. Soluble NRP1 ligands can be prepared using standard recombinant DNA techniques, for example by deletion of the transmembrane and cytoplasmic domains. A soluble ligand can be transformed into a soluble ligand fusion protein also using standard recombinant DNA techniques. A fusion protein can be prepared in which fusion partner can comprise a binding moiety that facilitates separation of the fusion protein.

(91) In order to separate the NRP1 reactive positive cells from non-reactive cells, one of a variety of different cell separation techniques known in the art can be used. Preferably, the NRP1 reactive positive cells are separated from non-reactive cells by fluorescence activated cell sorting (FACS). The FACS technology, and apparatuses for carrying it out to separate cells, is well established in the art. When FACS is used for cell separation, preferably the agent(s) reactive with NRP1 that is used is a fluorescently-labeled anti-NRP1 monoclonal antibody. Alternatively, cell separation can be achieved by, for example, magnetic activated cell sorting (MACS). When MACS is used for cell separation, preferably the agent reactive with NRP1 that is used is magnetic nanoparticles coated with anti-NRP1 monoclonal antibody. Alternatively, other single cell sorting methodologies known in the art can be applied to the methods of isolating cardiac ventricular progenitor cells of the invention, including but not limited to IsoRaft array and DEPArray technologies.

(92) Prior to contact with the agent(s) reactive with NRP1, and separation of NRP1 reactive cells, human pluripotent stem cells can be cultured under conditions that lead to the generation of cardiac progenitor cells. Culture conditions for generating cardiac progenitor cells have been described in

the art (see e.g., Lian, X. et al. (2012) *Proc. Natl. Acad. Sci. USA* 109:E1848-1857; U.S. Patent Publication No. 20130189785) and also are described in detail in Example 1 and FIG. 1, as well as in Example 10. Typically, Wnt/ β -catenin signaling is first activated in the hPSCs, followed by an incubation period, followed by inhibition of Wnt/ β -catenin signaling. Activation of Wnt/ β -catenin signaling is achieved by incubation with a Gsk3 inhibitor, preferably CHIR98014 (CAS 556813-39-9). Inhibition of Wnt/ β -catenin signaling is achieved by incubation with a Porcn inhibitor, preferably Wnt-059 (CAS 1243243-89-1). Suitable hPSCs for use in the methods of the invention include induced pluripotent stem cells (iPSC), such as 19-11-1, 19-9-7 or 6-9-9 cells (Yu, J. et al. (2009) *Science* 324:797-801), and human embryonic stem cell lines, such as ES03 cells (WiCell Research Institute) or H9 cells (Thomson, J. A. et al. (1998) *Science* 282:1145-1147). Suitable culture media for generating cardiomyogenic progenitors include E8 medium, mTeSR1 medium and RPMI/B27 minus insulin, each described further in Example 1 and/or Example 10.

(93) Preferably, the human cardiomyogenic progenitor cells are ventricular progenitor cells. Culture conditions have now been determined that bias the cardiomyogenic progenitor cells to the ventricular lineage. These ventricular cardiomyogenic progenitor cells can be cultured in RPMI/B27 medium and they can further differentiate into ventricular muscle cells. A preferred medium for culturing the cardiac ventricular progenitor cells in vitro such that they differentiate into ventricular cells in vitro (e.g., expressing the MLC2v marker described below) is the Cardiac Progenitor Culture (CPC) medium (advanced DMEM/F12 supplemented with 20% KnockOut Serum Replacement, 2.5 mM GlutaMAX and 100 μ g/ml Vitamin C).

(94) Known markers of differentiated cardiac cells can be used to identify the type(s) of cells that are generated by differentiation of the cardiac progenitor cells. For example, cardiac troponin I (cTnI) can be used as a marker of cardiomyocyte differentiation. CD144 (VE-cadherin) can be used as a marker of endothelial cells. Smooth muscle actin (SMA) can be used as a marker of smooth muscle cells. MLC2v can be used as a marker of ventricular muscle cells. MLC2a, which is expressed on both immature ventricular muscle cells and atrial muscle cells, can be used as a marker for those cell types. Additionally, sarcolipin, which is specifically expressed in atrial muscle cells, can be used as a marker for atrial muscle cells. Phospholamban, which is expressed predominantly in the ventricles and, to a lesser extent, in the atria, can also be used as a marker. Hairy-related transcription factor 1 (HRT1), also called Hey1, which is expressed in atrial cardiomyocytes, can be used as a marker for atrial cardiomyocytes. HRT2 (Hey2), which is expressed in ventricular cardiomyocytes, can be used as a marker for ventricular cardiomyocytes. In addition, IRX4 has a ventricular-restricted expression pattern during all stages of development, and thus can be used as a ventricular lineage marker. In summary, the genes expressed in the ventricles, and thus which are appropriate ventricular markers, are: MLC2v, IRX4 and HRT2, while genes expressed in the atria, and thus which are appropriate atrial markers are: MLC2a, HRT1, Sarcolipin and ANF (atrial natriuretic factor). The preferred marker of ventricular differentiation is MLC2v.

(95) Clonal Populations of Human Cardiac Ventricular Progenitor Cells

(96) In another aspect, the invention provides methods for obtaining a clonal population of human cardiac ventricular progenitor cells, as well as isolated clonal populations of such progenitors. The invention allows for the expansion and propagation of the cardiac ventricular progenitor cells such that a clonal population of a billion or more cells can be achieved. The ability to clonally expand the NRP1+ cardiac ventricular progenitor cells to such large numbers is a necessary feature for successful use of these cells in vivo to enhance cardiac function, since such a use requires on the order of a billion or more cells.

(97) Accordingly, in another aspect, the invention provides a method for obtaining a clonal population of human cardiac ventricular progenitor cells, the method comprising: isolating a single NRP1+ human cardiac ventricular progenitor cell; and culturing the single NRP1+ human cardiac ventricular progenitor cell under conditions such that the cell is expanded to at least 1×10^9 .

cells to thereby obtain a clonal population of human cardiac ventricular progenitor cells.

(98) In a preferred embodiment, the single NRP1⁺ human cardiac ventricular progenitor cell is Isl1 positive, Nkx2.5 negative and flk1 negative at the time of initial culture. As described further in the Examples, such a single cell can be obtained at approximately day 6 of the culture under conditions that promote the generation of cardiomyogenic progenitors. The clonal population of human cardiac ventricular progenitors can be further cultured and differentiated in vitro such that the cells express the ventricular marker MLC2v.

(99) Preferably, the single NRP1⁺ human cardiac ventricular progenitor cell is isolated by fluorescence activated cell sorting. Alternatively, the cell can be isolated by MACS or by other cell sorting methods known in the art and/or described herein.

(100) Preferably, the single NRP1⁺ human cardiac ventricular progenitor cell is isolated using one or more agents reactive with NRP1, such as an anti-NRP1 antibody or other agent reactive with NRP1 as described hereinbefore.

(101) In other embodiments, the clonal population of human cardiac ventricular progenitor cells is NRP1⁺ and positive for at least one second marker of HVPs (e.g., Isl1, FZD4, LIFR, FGFR3, TNFSF9). Such double-positive cells can be isolated and clonally expanded as described herein before using both an agent reactive with NRP1 and an agent reactive with the second marker.

(102) In a preferred embodiment, the single NRP1⁺ human cardiac ventricular progenitor cell is cultured in Cardiac Progenitor Culture (CPC) medium, as described hereinbefore.

(103) In a preferred embodiment, the single NRP1⁺ human cardiac ventricular progenitor cell is cultured under conditions such that the cell is biased toward ventricular differentiation. Preferred culture conditions include culture in CPC medium.

(104) In various embodiments, the single NRP1⁺ human cardiac ventricular progenitor cell can be expanded to at least 1×10^9 cells, at least 2×10^9 cells, at least 3×10^9 cells, at least 4×10^9 cells, at least 5×10^9 cells, at least 6×10^9 cells, at least 7×10^9 cells, at least 8×10^9 cells, at least 9×10^9 cells or at least 10×10^9 cells.

(105) Accordingly, the invention also provides a clonal population of at least 1×10^9 NRP1⁺ human cardiac ventricular progenitor cells, which are obtainable or obtained by the methods of the invention for obtaining a clonal population of human cardiac ventricular progenitor cells. In various embodiments, the clonal population of NRP1⁺ human cardiac ventricular progenitor cells comprises at least 1×10^9 cells, at least 2×10^9 cells, at least 3×10^9 cells, at least 4×10^9 cells, at least 5×10^9 cells, at least 6×10^9 cells, at least 7×10^9 cells, at least 8×10^9 cells, at least 9×10^9 cells or at least 10×10^9 cells. Differentiation of the progenitor cells to the ventricular lineage in vitro can be achieved by culture under conditions described herein for biasing toward the ventricular lineage. Furthermore, transplantation of the cardiac ventricular progenitor cells in vivo leads to ventricular differentiation in vivo.

(106) The invention also provides pharmaceutical compositions comprising the clonal population of cardiac ventricular progenitor cells. The pharmaceutical compositions typically are sterile and can comprise buffers, media, excipients and the like suitable for pharmaceutical administration. In one embodiment, the pharmaceutical composition comprising the clonal population is formulated onto a three dimensional (3D) matrix. Compositions formulated onto a 3D matrix are particularly preferred for formation of a heart muscle cell patch that can be transplanted in vivo for heart muscle repair. Furthermore, the compositions can be formulated into two dimensional (2D) sheets of cells, such as a muscular thin film (MTF) as described in Domian, I. J. et al. (2009) *Science* 326:426-429. Such 2D sheets of cell tissue also can be used in the formation of a heart muscle cell patch that can be transplanted in vivo for heart muscle repair.

(107) Generation of Human Ventricular Progenitors (HVPs)

(108) Prior to isolation by the aforementioned methods, and optionally obtaining a clonal population by the aforementioned methods, a non-clonal population of human ventricular progenitors (HVPs) can be obtained by culture of human pluripotent stem cells (hPSCs) under

appropriate culture conditions to generate the HVPs. An exemplary set of culture conditions, and suitable starting cells, is described in detail in Example 1 and Example 10, also referred to herein as the Human Ventricular Progenitor Generation (HVPG) protocol. Suitable hPSC starting cells include induced pluripotent stem cells (iPSC) and human embryonic stem cells, such as ES cell lines. For the protocol, Wnt/ β -catenin signaling first is activated in the hPSCs, followed by an incubation period, followed by inhibition of Wnt/ β -catenin signaling. Wnt/ β -catenin signaling activation is achieved by incubation with a Gsk3 inhibitor, preferably CHIR98014 (CAS 556813-39-9; commercially available from, e.g., Selleckchem). Wnt/ β -catenin signaling inhibition is achieved by incubation with a Porcn inhibitor, preferably Wnt-059 (CAS 1243243-89-1; commercially available from, e.g., Selleckchem or Tocris). The Gsk3 inhibitor is used to promote cardiac mesodermal differentiation, whereas the Porcn inhibitor is used to enhance ventricular progenitor differentiation from mesoderm cells.

(109) Accordingly, in another aspect, the invention provides a method of generating human ventricular progenitors (HVPs) comprising culturing human pluripotent stems cells (hPSCs) in a medium comprising a Gsk3 inhibitor, preferably CHIR98014, for at least 24 hours, more preferably for 2 days or 3 days, followed by culturing the hPSCs in a medium comprising a Porcn inhibitor, preferably Wnt-059 (and lacking the Gsk3 inhibitor), for at least 48 hours such that HVPs are generated. Experiments showed that after 24-hour treatment with CHIR-98014, more than 99% of hPSCs expressed the mesoderm marker Brachyury, and three days later after treatment with CHIR-98014, more than 95% of differentiated cells expressed *Mesp1*, which marks the cardiac mesoderm. Furthermore, 48-hour treatment with Wnt-059 enhanced ventricular progenitor differentiation from mesoderm cells.

(110) Accordingly, with regard to timing of the use of the Gsk3 and Porcn inhibitors, typically, at day 0 of culture, the hPSCs are cultured with the Gsk3 inhibitor, at day 3 of culture the medium is changed to remove the Gsk3 inhibitor and the cells are then cultured with media containing the Porcn inhibitor through day 5 of culture. HVP generation is optimal between days 5 and 7 (inclusive) in culture and peaks at day 6 of culture. Other non-limiting, exemplary details on culture conditions and timing of the use of the Gsk3 and Porcn inhibitors are described in detail in Examples 1 and 10.

(111) In Vivo Tissue Engineering

(112) In vivo transplantation studies described in Example 6 and 7 in which the human ventricular progenitors (HVPs) were transplanted under the kidney capsule in nude mice document the ability of the HVPs to spontaneously assemble into a large wall of mature, functional, human ventricular muscle on the surface of the kidney capsule. Vascularization occurs via a paracrine pathway by calling the murine vasculature to the ventricular muscle wall, while a matrix is generated via a cell autonomous pathway from the progenitors themselves. In vivo intra-myocardial transplantation studies described in Example 8 in which the HVPs were transplanted into the normal murine heart document that the HVPs spontaneously migrate to the epicardial surface, where they expand, subsequently differentiate, and mature into a wall of human ventricular muscle on the surface of the epicardium. Taken together, these studies show that human ventriculogenesis can occur via a completely cell autonomous pathway in vivo via purified HVPs, thereby allowing their use in organ-on-organ in vivo tissue engineering.

(113) The human ventricular myocardium has a limited capacity for regeneration, most of which is lost after 10 years of age (Bergmann, O. et al. (2015) *Cell* 161:1566-1575). As such, new strategies to generate heart muscle repair, regeneration, and tissue engineering approaches during cardiac injury have been a subject of intense investigation in regenerative biology and medicine (Sahara, M. et al. (2015) *EMBO J.* 34:710-738; Segers, V. F. M. and Lee, R. T. (2008) *Nature* 451:937-942). Given the need to achieve coordinated vascularization and matrix formation during tissue engineering of any solid organ, the assumption has been that the formation of an intact 3-D solid organ in vivo will ultimately require the addition of vascular cells and/or conduits, as well as

biomaterials and/or decellularized matrix that will allow alignment and the generation of contractile force (Forbes, S. J. and Rosenthal, N. (2014) *Nature Med.* 20:857-869; Harrison, R. H. et al. (2014) *Tissue Eng. Part B Rev.* 20:1-16). The complexity of adding these various components to achieve the formation of a functional solid organ has confounded attempts to reduce this to clinical practice (Webber, M. J. et al. (2014) *Ann. Biomed. Eng.* 43:641-656). Although hPSCs hold great promise, to date, it has not been possible to build a pure, vascularized, fully functional, and mature 3-D human ventricular muscle organ in vivo on the surface of a heart in any mammalian system (Vunjak-Novakovic, G. et al. (2011) *Annu. Rev. Biomed. Eng.* 13:245-267).

(114) The ability of generate billions of purified HVPs from a renewable source of either human ES or iPS cell lines represent a new approach to the generation of functional ventricular muscle in the setting of advanced heart failure. The progenitors can be delivered by intramyocardial injection and then self-migrate to the epicardial surface where they expand and differentiate, losing progenitor markers. Over the course of several week, the cells exit the cell cycle, and proceed to form adult rod-shaped cells that display several independent markers of mature ventricular myocardium including the formation of T tubules, catecholamine responsiveness, loss of automaticity, adult rod shaped conformation with aligned sarcomenric structures, and the ability to generate force that is comparable to other heart muscle patches derived from hPSCs differentiated cardiomyocytes (Tulloch, N. L. et al. (2011) *Circ. Res.* 109:47-59). The scalability of this cell autonomous pathway has allowed the ectopic generation of human ventricular muscle that has a combined thickness in excess of 1.5 cm in thickness, approaching levels that correspond to the human ventricular free wall (Basavarajaiah, S. et al. (2007) *Br. J. Sports Med.* 41:784-788).

(115) The ability to migrate to the epicardial niche, the site of most of the adult heart progenitors at later stages, is a unique feature of HVPs, and mimics the normal niche of these cells during expansion of the ventricular compact zone during ventriculogenesis. Previous studies have shown that the generation of acute ischemic injury and a breakdown in vascular permeability are a pre-requisite for the grafting of relatively small numbers of ES cell derived cardiomyocytes into injured myocardium (van Laake, L. W. et al. (2007) *Stem Cell Res.* 1:9-24; Laflamme, M. A. et al. (2007) *Nat. Biotechnol.* 25:1015-1024), and even then the survival rate is low (<5%) (Laflamme, M. A. and Murry, C. E. (2011) *Nature* 473:326-335; Laflamme, M. A. et al. (2005) *Am. J. Pathol.* 167:663-671). The ability of intra-myocardial HVPs to form an extensive ventricular patch on the epicardial surface in the absence of acute ischemic injury provides a new therapeutic strategy for dilated cardiomyopathy without the need for additional biomaterials, cells, or transfer of exogenous genes and/or RNAs.

(116) The ability to form a 3-D ventricular muscle wall on the epicardial surface of the in vivo normal heart is a unique feature of the ventricular progenitors as later stage progenitors do not display the ability for the formation of three-dimensional ventricular tissue in either the cardiac or non-cardiac context, emphasizing the importance of generating a committed ventricular lineage as well as purifying the specific ventricular progenitor at a specific stage of ventriculogenesis.

(117) Accordingly, the invention provides methods for generating human ventricular tissue in vivo using the HVPs described herein. In one embodiment, the method comprises transplanting the NRP1+ progenitors into an organ of a non-human animal and allowing the progenitors to grow in vivo such that human ventricular tissue is generated. Preferably, the non-human animal is immunodeficient such that it cannot mount an immune response against the human progenitor cells. In one embodiment, the non-human animal is a mouse, such as an immunodeficient NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ mouse or an immunodeficient SCID-beige mouse (commercially available from Charles River France). In one embodiment, the organ is a kidney (e.g., the cells are transplanted under the kidney capsule). In another embodiment, the organ is a heart. In various embodiments, at least $1 \times 10^{sup.6}$ cells, at least $2 \times 10^{sup.6}$ cells, at least $3 \times 10^{sup.6}$ cells, at least $4 \times 10^{sup.6}$ cells, at least $5 \times 10^{sup.6}$ cells, at least $1 \times 10^{sup.7}$ cells, at least $5 \times 10^{sup.7}$ cells, at least $1 \times 10^{sup.8}$ cells, at least $1 \times 10^{sup.9}$ cells are transplanted.

(118) To obtain HVPs for transplantation, human pluripotent stem cells (hPSCs) can be cultured in vitro under conditions leading to the generation of HVPs, as described herein (referred to herein as the HVPG protocol). Regarding the timing of transplanting HVPs post in-vitro culture, for optimal ventricular tissue generation the cells should be transplanted at a stage that can be defined based on the cellular markers expressed by the HVPs at the time of transplantation, determined at days post the start of culture, which is defined as day 0 of the HVPG protocol. In one embodiment, the cells are transplanted after the peak of cardiac mesoderm formation, which can be defined as peak expression of the mesodermal marker MESP1. Typically, MESP1 expression is between day 2 and day 4 of culture (inclusive) and peaks at day 3. In one embodiment, the cells are transplanted at the time corresponding to peak Islet-1 expression. Typically, Islet 1 is expressed between day 4 to day 8 of culture (inclusive) and peaks at day 6 of culture. In one embodiment, the cells are transplanted before the peak of NKX2.5 expression. Typically, NKX2.5 expression starts at day 6 of culture, peaks at day 10 of culture and is then maintained afterwards. In one embodiment, the cells are transplanted prior to the peak expression of the downstream genes MEF-2 and TBX-1. Typically, these downstream genes are expressed between day 5 and day 15 of culture (inclusive) and peaks at day 8 of culture. In one embodiment, the cells are transplanted prior to the expression of differentiated contractile protein genes. Typically, the expression of contractile protein genes (including TNNT2 and MYH6) starts from day 10 of culture onward. In certain embodiments, the cells are transplanted at a time when two, three or four of the aforementioned marker patterns are present. In another embodiment, the cells are transplanted at a time when all five of the aforementioned marker patterns are present. In one embodiment, the cells are transplanted between day 4 to day 8 (inclusive) of culture. In a more preferred embodiment, the cells are transplanted between day 5 to day 7 (inclusive) of culture. In the most preferred embodiment, the cells are transplanted on day 6 of culture.

(119) The transplanted cells can be allowed to grow in the non-human animal for a suitable period time to allow for the generation of the desired size, amount or thickness of ventricular tissue. In various embodiments, the cells are allowed to grow for one week, two weeks, one month, two months, three months, four months, five months or six months. The method can further comprise harvesting ventricular tissue from the non-human animal after growth of the cells and differentiation into ventricular tissue.

(120) Methods of Enhancing Cardiac Function

(121) The cardiac ventricular progenitor cells of the invention can be used in vivo to enhance cardiac function by transplanting the cells directly into the heart. It has now been shown that the NRP1+ progenitors have the capacity to differentiate into all three types of cardiac lineage cells (cardiac myocytes, endothelial cells and smooth muscle cells) (see Example 3). Furthermore, when cultured under conditions that bias toward the ventricular lineage, the NRP1+ progenitors have now been shown to adopt a predominantly ventricular muscle phenotype when transplanted into the natural ventricle environment in vivo, demonstrating that these progenitor cells “recognize” the ventricular environment and respond and differentiate appropriately in vivo. Since damage to the ventricular environment is largely responsible for the impaired cardiac function in cardiac diseases and disorders, the ability to restore ventricular muscle cells using the ventricular progenitor cells of the invention represents a significant advance in the art.

(122) Accordingly, in another aspect, the invention provides a method of enhancing cardiac function in a subject, the method comprising administering a pharmaceutical composition comprising the clonal population of NRP1+ cardiac ventricular progenitor cells of the invention to the subject. Preferably, the clonal population is administered directly into the heart of the subject. More preferably, the clonal population is administered directly into a ventricular region of the heart of the subject. In one embodiment, the pharmaceutical composition administered to the subject comprises the clonal population formulated onto a three dimensional matrix.

(123) The methods of the invention for enhancing cardiac function in a subject can be used in a

variety of clinical situations involving damage to the heart or reduced or impaired cardiac function. Non-limiting examples of such clinical situations include a subject who has suffered a myocardial infarction and a subject who has a congenital heart disorder. Thus, in another aspect, the invention provides a method of treating a cardiovascular condition, disease or disorder in a subject, the method comprising administering a pharmaceutical composition comprising the clonal population of NRP1+ cardiac ventricular progenitor cells of the invention to the subject. A therapeutically effective amount of cardiac ventricular progenitor cells can be administered for the treatment of a cardiovascular condition, disease or disorder. Examples of preferred cardiovascular conditions, diseases or disorders include coronary artery disease and acute coronary syndrome.

(124) Methods of Use of Cardiac Ventricular Progenitor Cells In Vitro

(125) The cardiac ventricular progenitor cells of the invention can be used in vitro in the study of various aspects of cardiac maturation and differentiation, in particular in identifying the cells signaling pathways and biological mediators involved in the process of cardiac maturation and differentiation.

(126) Furthermore, since the NRP1+ cardiac ventricular progenitor cells of the invention are committed to the cardiac lineage and, moreover, are biased toward ventricular differentiation, these progenitor cells also are useful for evaluating the cardiac toxicity of test compounds. All potential new drugs and therapeutics must be evaluated for their toxicity to cardiac cells, before they can be deemed safe for use in humans. Thus, the ability to assess cardiac toxicity in an in vitro culture system is very advantageous. Accordingly, in another aspect, the invention provides a method of screening for cardiac toxicity of test compound, the method comprising: providing NRP1+ human cardiac ventricular progenitor cells; contacting the cells with the test compound; and measuring toxicity of the test compound for the cells, wherein toxicity of the test compound for the cells indicates cardiac toxicity of the test compound.

(127) In a preferred embodiment, the NRP1+ human cardiac ventricular progenitor cells are provided by isolating the cells according to the methods described herein. In a particularly preferred embodiment, the cells are isolated by separating NRP1+ cells from a cell culture comprising cardiac progenitor cells using an anti-NRP1 antibody. Preferably, the cells are isolated using FACS or MACS as described herein. In yet another embodiment, the NRP1+ human cardiac ventricular progenitor cells are further cultured and differentiation into MLC2v+ ventricular cells prior to contacting with the test compound.

(128) The toxicity of the test compound for the cells can be measured by one or more of a variety of different methods for assessing cell viability or other physiological functions. Preferably, the effect of the test compound on cell viability is measured using a standard cell viability assay, wherein reduced cell viability in the presence of the test compound is indicative of cardiac toxicity of the test compound. Additionally or alternatively, cell growth can be measured. Additionally or alternatively, other indicators of physiological functions can be measured, such as cell adhesion, cell signaling, surface marker expression, gene expression and the like. Similarly, a negative effect of the test compound on any of these indicators of physiological function is indicative of cardiac toxicity of the test compound.

(129) The invention further provides a method of identifying a compound that modulates human cardiac ventricular progenitor cell differentiation, the method comprising: providing NRP1+ human cardiac ventricular progenitor cells; culturing the cells in the presence or absence of a test compound; measuring differentiation of the cells in the presence or absence of the test compound; and selecting a test compound that modulates human cardiac ventricular progenitor cell differentiation, as compared to differentiation in the absence of the test compound, to thereby identify a compound that modulates human cardiac ventricular progenitor cell differentiation.

(130) In one embodiment, the test compound stimulates human cardiac ventricular progenitor cell differentiation. In another embodiment, the test compound inhibits human cardiac ventricular progenitor cell differentiation. Differentiation of the cells can be measured by, for example,

measurement of the expression of differentiation markers appearing on the cultured cells over time, as described herein. In a preferred embodiment, the NRP1+ human cardiac ventricular progenitor cells are provided by isolating the cells according to the methods described herein. In a particularly preferred embodiment, the cells are isolated by separating NRP1+ cells from a cell culture comprising cardiac progenitor cells using an anti-NRP1 antibody. Preferably, the cells are isolated using FACS or MACS as described herein.

(131) The invention further provides a method of identifying a compound that modulates human ventricular cardiomyocyte function, the method comprising: providing NRP1+ human cardiac ventricular progenitor cells; culturing the cells in the presence or absence of a test compound under conditions that generate human ventricular cardiomyocytes; measuring function of the human ventricular cardiomyocytes in the presence or absence of the test compound; and selecting a test compound that modulates human ventricular cardiomyocyte function, as compared to function in the absence of the test compound, to thereby identify a compound that modulates human ventricular cardiomyocyte function.

(132) In one embodiment, the test compound stimulates human ventricular cardiomyocyte function. In another embodiment, the test compound inhibits human ventricular cardiomyocyte function. Function of the cells can be measured by measurement of any suitable indicator of ventricular cell function, including but not limited to, for example, formation of T tubules, acquisition of adult-rod shaped ventricular cardiomyocytes, and ability to generate force in response to electrical stimulation. Suitable assays for measuring such indicators of ventricular cell function are known in the art. In a preferred embodiment, the NRP1+ human cardiac ventricular progenitor cells are provided by isolating the cells according to the methods described herein. In a particularly preferred embodiment, the cells are isolated by separating NRP1+ cells from a cell culture comprising cardiac progenitor cells using an anti-NRP1 antibody. Preferably, the cells are isolated using FACS or MACS as described herein.

(133) In Vivo Animal Models Using Human Ventricular Progenitor Cells

(134) The development of human iPS and ES cell based models of cardiac disease has opened new horizons in cardiovascular drug development and discovery. However, to date, these systems have had the limitations of being based on 2D structures in cultured cell systems. In addition, the fetal and immature properties of the cells limit their utility and fidelity to the adult heart. Human cardiac disease, in particular heart failure, is a complex, multifactorial, multi-organ disease, that is influenced by environmental, hormonal, and other key organs that are known sites for therapeutic targets, such as the kidney. The ability to build a mature functional human ventricular organ either ectopically or on the surface of the intact normal murine heart opens up a new in vivo model system to allow studies that normally could only be assayed on a mature human ventricular muscle chamber, such as ventricular arrhythmias, generation of contractile force, fibrosis, and the potential for regeneration. Accordingly, the option to study human cardiac disease outside of the in vitro tissue culture systems, and directly in the context of heart failure in vivo, is now clearly possible.

(135) Thus, the human ventricular progenitor cells also can be used to create animal models that allow for in vivo assessment of human cardiac tissue function and for in vivo screening of compounds, such as to determine the cardiac toxicity of a test compound in vivo or to identify compounds that modulate human cardiac tissue differentiation or function in vivo. Accordingly, the invention provides methods for testing the effects of test compounds on human ventricular tissue in vivo using the HVPs described herein. In one embodiment, the method comprises: transplanting NRP1+ human ventricular progenitors into an organ of a non-human animal; allowing the progenitors to grow in vivo such that human ventricular tissue is generated; administering a test compound to the non-human animal; and evaluating the effect of the test compound on the human ventricular tissue in the non-human animal.

(136) In another embodiment, the method comprises: administering a test compound to a non-human animal, wherein the non-human animal comprises NRP1+ human ventricular progenitors

transplanted into an organ of the non-human animal; and evaluating the effect of the test compound on the NRP1+ human ventricular progenitors in the non-human animal.

(137) In one embodiment, the cardiac toxicity of the test compound is evaluated, for example by measuring the effect of the test compound on the viability of the human ventricular tissue or the NRP1+ human ventricular progenitors in the non-human animal (as compared to the viability of the tissue or progenitors in the absence of the test compound). Cell viability can be assessed by standard methods known in the art.

(138) In another embodiment, the ability of a test compound to modulate cardiac differentiation can be evaluated, for example by measuring the effect of the test compound on the differentiation of the human ventricular tissue or NRP1+ progenitors in the non-human animal (as compared to the differentiation of the tissue or progenitors in the absence of the test compound). Differentiation of the cells can be measured by, for example, measurement of the expression of differentiation markers appearing on the cells over time.

(139) In another embodiment, the ability of a test compound to modulate cardiac function can be evaluated, for example by measuring the effect of the test compound on the function of the human ventricular tissue or NRP1+ human progenitors in the non-human animal (as compared to the function of the tissue or progenitors in the absence of the test compound). Function of the tissue or progenitors can be measured by measurement of any suitable indicator of ventricular cell function, including but not limited to, for example, formation of T tubules, acquisition of adult-rod shaped ventricular cardiomyocytes, and ability to generate force in response to electrical stimulation. Suitable assays for measuring such indicators of ventricular cell function are known in the art.

(140) Preferably, the non-human animal is immunodeficient such that it cannot mount an immune response against the human progenitor cells. In one embodiment, the non-human animal is a mouse, such as an immunodeficient NOD.Cg-Prkdcscid Il2rgtm1Wjl/SzJ mouse or an immunodeficient SCID-beige mouse (commercially available from Charles River France). In one embodiment, the organ is a kidney (e.g., the cells are transplanted under the kidney capsule). In another embodiment, the organ is a heart. In various embodiments, at least 1×10^6 cells, at least 2×10^6 cells, at least 3×10^6 cells, at least 4×10^6 cells, at least 5×10^6 cells, at least 1×10^7 cells, at least 5×10^7 cells, at least 1×10^8 cells, at least 1×10^9 cells are transplanted.

(141) To create the animal models, HVPs for transplantation can be obtained as described above by culturing of hPSCs in vitro under conditions leading to the generation of HVPs. Regarding the timing of transplanting HVPs post in-vitro culture, for optimal ventricular tissue generation the cells should be transplanted at a stage that can be defined based on the cellular markers expressed by the HVPs at the time of transplantation, determined at days post the start of culture, which is defined as day 0 of the HVPG protocol. In one embodiment, the cells are transplanted after the peak of cardiac mesoderm formation, which can be defined as peak expression of the mesodermal marker MESP1. Typically, MESP1 expression is between day 2 and day 4 of culture (inclusive) and peaks at day 3. In one embodiment, the cells are transplanted at the time corresponding to peak Islet-1 expression. Typically, Islet 1 is expressed between day 4 to day 8 of culture (inclusive) and peaks at day 6 of culture. In one embodiment, the cells are transplanted before the peak of NKX2.5 expression. Typically, NKX2.5 expression starts at day 6 of culture, peaks at day 10 of culture and is then maintained afterwards. In one embodiment, the cells are transplanted prior to the peak expression of the downstream genes MEF-2 and TBX-1. Typically, these downstream genes are expressed between day 5 and day 15 of culture (inclusive) and peaks at day 8 of culture. In one embodiment, the cells are transplanted prior to the expression of differentiated contractile protein genes. Typically, the expression of contractile protein genes (including TNNT2 and MYH6) starts from day 10 of culture onward. In certain embodiments, the cells are transplanted at a time when two, three or four of the aforementioned marker patterns are present. In another embodiment, the cells are transplanted at a time when all five of the aforementioned marker patterns are present. In

one embodiment, the cells are transplanted between day 4 to day 8 (inclusive) of culture. In a more preferred embodiment, the cells are transplanted between day 5 to day 7 (inclusive) of culture. In the most preferred embodiment, the cells are transplanted on day 6 of culture.

(142) The transplanted cells can be allowed to grow in the non-human animal for a suitable period time to allow for the generation of the desired size, amount or thickness of ventricular tissue, prior to administration of the test compound(s). In various embodiments, the cells are allowed to grow for one week, two weeks, one month, two months, three months, four months, five months or six months.

(143) The present invention is further illustrated by the following examples, which should not be construed as further limiting. The contents of figures and all references, patents and published patent applications cited throughout this application are expressly incorporated herein by reference.

EXAMPLES

Example 1: Generation of Human Isl1+ Cardiomyogenic Progenitor Cells by Modulation of Wnt Signaling in Human Pluripotent Stem Cells

(144) Temporal modulation of canonical Wnt signaling has been shown to be sufficient to generate functional cardiomyocytes at high yield and purity from numerous hPSC lines (Lian, X. et al.

(2012) *Proc. Natl. Acad. Sci. USA* 109:E1848-1857; Lian, X. et al. (2013) *Nat. Protoc.* 8:162-175).

In this approach, Wnt/ β -catenin signaling first is activated in the hPSCs, followed by an incubation period, followed by inhibition of Wnt/ β -catenin signaling. In the originally published protocol, Wnt/ β -catenin signaling activation was achieved by incubation with the Gsk3 inhibitor CHIR99021 (GSK-3 α , IC₅₀=10 nM; GSK-3, β IC₅₀=6.7 nM) and Wnt/ β -catenin signaling inhibition was achieved by incubation with the Porcn inhibitor IWP2 (IC₅₀=27 nM). Because we used Gsk3 inhibitor and Wnt production inhibitor for cardiac differentiation, this protocol was termed GiWi protocol. To improve the efficiency of the original protocol and reduce the potential side effects of the small molecules used in the original protocol, a second generation protocol was developed that uses another set of small molecules with higher inhibition potency. In this second generation GiWi protocol, Wnt/ β -catenin signaling activation was achieved by incubation with the Gsk3 inhibitor CHIR98014 (CAS 556813-39-9; commercially available from, e.g., Selleckchem) (GSK-3 α , IC₅₀=0.65 nM; GSK-3, β IC₅₀=0.58 nM) and Wnt/ β -catenin signaling inhibition was achieved by incubation with the Porcn inhibitor Wnt-059 (CAS 1243243-89-1; commercially available from, e.g., Selleckchem or Tocris) (IC₅₀=74 pM). The Gsk3 inhibitor CHIR98014 was used to promote cardiac mesodermal differentiation, whereas the Porcn inhibitor Wnt-059 was used to enhance ventricular progenitor differentiation from mesoderm cells.

(145) For cardiomyocyte differentiation via the use of these small molecules, hPSCs were maintained on Matrigel (BD Biosciences) coated plates (Corning) in E8 medium (described in Chen, G. et al. (2011) *Nature Methods*, 8:424-429; commercially available; STEMCELL Technologies) or mTeSR1 medium (commercially available; STEMCELL Technologies). Suitable hPSCs include induced pluripotent stem cells (iPSCs) such as 19-11-1, 19-9-7 or 6-9-9 cells (Yu, J. et al. (2009) *Science*, 324:797-801) and human embryonic stem cells (hESCs), such as ES03 (WiCell Research Institute) and H9 cells (Thomson, J. A. et al. (1998) *Science*, 282:1145-1147).

(146) hPSCs maintained on a Matrigel-coated surface in mTeSR1 medium were dissociated into single cells with Accutase (Life Technologies) at 37° C. for 5 minutes and then seeded onto a Matrigel-coated cell culture dish at 100,000-200,000 cells/cm² in mTeSR1 medium supplemented with 5 μ M ROCK inhibitor Y-27632 (Selleckchem)(day -2) for 24 hours. Cells were then cultured in mTeSR1, changed daily. At day 0, cells were then treated with 1 μ M Gsk3 inhibitor CHIR98014 (Selleckchem) for 24 hours (day 0 to day 1) in RPMI/B27-ins (500 ml RPMI with 10 ml B27 supplement without insulin). The medium was then changed to the corresponding medium containing 2 μ M the Porcn inhibitor Wnt-059 (Selleckchem) at day 3, which was then removed during the medium change on day 5. Cells were maintained in RPMI/B27 (stock solution: 500 ml RPMI medium+10 ml B27 supplement) starting from day 7, with the medium changed every three

days. This exemplary culturing protocol for generating cardiomyogenic progenitor cells is illustrated schematically in FIG. 1.

(147) Flow cytometry and immunostaining were preformed to examine the expression of particular lineage markers. After 24 hour treatment with CHIR-98014, more than 99% of the hPSCs expressed the mesoderm marker Brachyury. Three days after treatment with CHIR-98014, more than 95% of differentiated cells expressed Mesp1, which marks the cardiac mesoderm. The culture protocol not only allowed the cells to synchronously differentiate into the cardiac mesodermal lineage, but also reproducibly generated more than 90% of ventricular myocytes after 14 days of differentiation, as determined by cTnT flow cytometry and electrophysiology analysis.

(148) To further assess cardiac differentiation of the hPSCs over time, Western blot analysis was performed on days 0-7 and d11 to examine the expression of Isl1 and Nkx2.5 (cardiomyogenic progenitor markers) and cTnI (a cardiac myocyte marker). Cells were lysed in M-PER Mammalian Protein Extraction Reagent (Pierce) in the presence of Halt Protease and Phosphatase Inhibitor Cocktail (Pierce). Proteins were separated by 10% Tris-Glycine SDS/PAGE (Invitrogen) under denaturing conditions and transferred to a nitrocellulose membrane. After blocking with 5% dried milk in TBST, the membrane was incubated with primary antibody overnight at 4° C. The membrane was then washed, incubated with an anti-mouse/rabbit peroxidase-conjugated secondary antibody at room temperature for 1 hour, and developed by SuperSignal chemiluminescence (Pierce). The results are shown in FIG. 2. During cardiac differentiation of hPSCs, Isl1 expression started on day 4 and increased to its maximum expression on day 6, whereas NKx2.5 only started to express on day 6 and reached its maximum expression after day 10. Cardiomyoctes (cTnI+ cells) were not induced until day 11 of differentiation.

(149) In addition, immunostaining of the day 6 cells was performed for Isl1 expression. Cells were fixed with 4% formaldehyde for 15 minutes at room temperature and then stained with primary (anti-Isl1) and secondary antibodies in PBS plus 0.4% Triton X-100 and 5% non-fat dry milk (Bio-Rad). Nuclei were stained with Gold Anti-fade Reagent with DAPI (Invitrogen). An epifluorescence microscope (Leica DM IRB) with a QImaging® Retiga 4000R camera was used for imaging analysis. The results showed substantial numbers of Isl1+ cells.

(150) Flow cytometry analysis of day 6 cells for Isl1 expression also was performed. Cells were dissociated into single cells with Accutase for 10 minutes and then fixed with 1% paraformaldehyde for 20 minutes at room temperature and stained with primary and secondary antibodies in PBS 0.1% Triton X-100 and 0.5% BSA. Data were collected on a FACSCaliber flow cytometer (Beckton Dickinson) and analyzed using FloJo. The results, shown in FIG. 3, showed that more than 95% of cells expressed Isl1 at this stage.

(151) In summary, this example provides a protocol for human ventricular progenitor generation (HVPG protocol) that allows for the large-scale production of billions of Isl1+ human HPVs efficiently within 6 days.

Example 2: Identification of Jagged 1 as a Cell Surface Marker of Cardiac Progenitor Cells

(152) To profile the transcriptional changes that occur during the cardiac differentiation process at a genome-scale level, RNA sequencing (RNA-seq) was performed at different time points following differentiation to build cardiac development transcriptional landscapes. We performed RNA-seq experiments on day 0 to day 7 samples, as well as day 19 and day 35 samples (two independent biological replicates per time point). Two batches of RNA-seq (100 bp and 50 bp read length) were performed using the illumine Hiseq 2000 platform. In total, 20 samples were examined. Bowtie and Tophat were used to map our reads into a reference human genome (hg19) and we calculate each gene expression (annotation of the genes according to Refseq) using RPKM method (Reads per kilobase transcript per million reads). Differentiation of hPSCs to cardiomyocytes involves five major cell types: pluripotent stem cells (day 0), mesoderm progenitors (day 1 to day 2), cardiac mesoderm cells (day 3 to day 4), heart field progenitors (day 5, day 6 and day 7), and cardiomyocytes (day 10 after).

(153) Molecular mRNA analysis of cardiac differentiation from hPSCs using the HVPG protocol revealed dynamic changes in gene expression, with down-regulation of the pluripotency markers OCT4, NANOG and SOX2 during differentiation. Induction of the primitive streak-like genes T and MIXL1 occurred within the first 24 hours following CHIR-98014 addition, and was followed by upregulation of the cardiac mesodermal marker MESP1 on day 2 and day 3. Expression of the cardiac muscle markers TNNT2, TNNC1, MYL2, MYL7, MYH6, MYH7 and IRX4 was detected at later stage of differentiation (after day 10).

(154) By this analysis, genes enriched at each differentiation stage, including mesoderm cells, cardiac progenitors and cardiomyocytes, were identified. Mesoderm cells, which are related to day 1 differentiated cells, express brachyury. We identified potential surface markers for mesoderm cells, including: FZD10, CD48, CD1D, CD8B, IL15RA, TNFRSF1B, TNFSF13, ICOSLG, SEMA7A, SLC3A2, SDC1, HLA-A. Through similar analysis, we also identified surface markers for cardiac mesoderm mesp1 positive cells, including: CXCR4, ANPEP, ITGA5, TNFRSF9, FZD2, CD1D, CD177, ACVRL1, ICAM1, L1CAM, NGFR, ABCG2, FZD7, TNFRSF13C, TNFRSF1B.

(155) Consistent with western blot analysis, ISL1 mRNA was expressed as early as day 4 and peaked on day 5, one day before its protein expression reached its peak. On day 5 of differentiation (the cardiac progenitor stage, isl1 mRNA expression maximum on day 5, isl1 protein expression maximum on day 6), the day 5 enriched genes were compared with an anti-CD antibody array (a panel of 350 known CD antibodies) and a number of potential cell-surface protein markers were identified. We identified many cell-surface proteins expressed at this stage, including: FZD4, JAG1, PDGFRA, LIFR (CD118), TNFSF9, FGFR3.

(156) The cell surface protein Jagged 1 (JAG1) and Frizzled 4 (FZD4) were selected for further analysis. Jagged 1 expression was further studied as described below and in Examples 3 and 4. Frizzled 4 expression was further studied as described in Example 5.

(157) Firstly, the expression of Isl1 and Jag1 was profiled using the double staining flow cytometry technique. Flow cytometric analysis was carried out essentially as described in Example 1, using anti-Isl1 and anti-Jag1 antibodies for double staining. The results are shown in FIG. 4. Jagged 1 expression was found to trace the expression of Islet 1 and on day 6 of differentiation, all of the Islet 1 positive cells also expressed Jagged 1, and vice versa. Because of the co-expression pattern of these two markers, a Jagged 1 antibody was used to enrich the 94.1% Islet 1+ cells differentiated population to 99.8% purity of Islet1+Jagged1+ cells.

(158) It also was confirmed that Islet 1 is an earlier developmental gene than the Nkx2.5 gene using double immunostaining of ISL1 and NKX2.5 expression in HVPs. The purified HVPs uniformly express the ISL1 gene, but at this stage, only a few of the cells started to express Nkx2.5.

(159) Furthermore, immunostaining with both anti-Isl1 and anti-Jag 1 was performed, essentially as described in Example 1, on week 4 human fetal heart tissue, neonatal heart tissue and 8-year old heart tissue. The results revealed that in the in vivo fetal heart, all of the Islet 1 positive cells also expressed Jagged 1. However, the neonatal heart and 8-year old heart did not express Islet 1 or Jagged 1. In the ventricle of week 4 human fetal heart, cardiac Troponin T (cTnT) staining revealed visible sarcomere structures. In addition, over 50% of ventricular cells in the week 4 fetal heart expressed both Islet1 and Jagged1, which was markedly decreased during subsequent maturation, with the loss of expression of both Islet1 and Jagged1 in the ventricular muscle cells of the human neonatal hearts.

(160) The above-described experiments demonstrate that Jagged 1 is a cell surface marker for Islet 1 positive cardiomyogenic progenitor cells.

Example 3: Clonal Differentiation of Isl1+Jag1+ Cardiac Progenitor Cells

(161) To characterize the clonal differentiation potential of Isl1+Jag1+ cells, cardiomyogenic progenitor cells were generated by the culturing protocol described in Example 1, and one single Isl1+Jag1+ cell was seeded into one well of a Matrigel-coated 48-well plate. Cells were purified with antibody of Jag1 and then one single cell was seeded into one well. The single cells were then

cultured for 3 weeks in Cardiac Progenitor Culture (CPC) medium (advanced DMEM/F12 supplemented with 2.5 mM GlutaMAX, 100 µg/ml Vitamin C, 20% Knockout Serum Replacement).

(162) Immunostaining of the 3-week differentiation cell population was then performed with three antibodies: cardiac troponin I (cTnI) for cardiomyocytes, CD144 (VE-cadherin) for endothelial cells and smooth muscle actin (SMA) for smooth muscle cells. The results showed that the single cell-cultured, Isl1+Jag1+ cells gave rise to cTnI positive and SMA positive cells, but not VE-cadherin positive endothelial cells, indicating these generated Islet1+ cells are heart muscle progenitors that have limited differentiation potential to endothelial lineages. Purified Islet1+Jagged1+ cells differentiated with the HVPG protocol from human induced pluripotent stem cells (iPSC 19-9-11 line) also showed similar in vitro differentiation potential and predominantly differentiate to cTnI+SMA+ cells, but not VE-cadherin+ cells. Over the course of several weeks, the cells expressed the ventricular specific marker MLC2v, indicating that the initial ISL1+ subset was already committed to the ventricular cell fate. Because of the limited vascular differentiation potential of Islet1+ cells generated using the HVPG protocol, these generated Islet1+ cells might represent a distinct progenitor population from the previously reported KDR+ population (Yang, L. et al. (2008) *Nature* 453:524-528) or multipotent ISL1+ cells (Bu, L. et al. (2009) *Nature* 460:113-117; Moretti, A. et al. (2006) *Cell* 127:1151-1165), which can give rise to all three lineages of cardiovascular cells.

(163) These results demonstrated that the Isl1+Jag1+ cardiomyogenic progenitor cells can be successfully cultured in vitro from a single cell to a significantly expanded cell population (1×10^9 cells or greater) that contains all three types of cardiac lineage cells, with a predominance of cardiomyocytes. Furthermore, these cells can be cultured in vitro for extended periods of time, for at least 2-3 weeks, and even for months (e.g., six months or more). Since the cardiomyogenic progenitor cells gradually differentiate into cardiomyocytes, which do not proliferate, a culture period of approximately 2-3 weeks is preferred.

Example 4: In Vivo Developmental Potential of Isl1+Jag1+ Cardiac Progenitor Cells

(164) The ES03 human embryonic stem cell (hESC) line (obtained from WiCell Research Institute) expresses green fluorescent protein (GFP) driven by the cardiac-specific cTnT promoter. ES03 cells were used to generate Isl1+Jag1+ cardiomyogenic progenitor cells using the culturing protocol described in Example 1. The Isl1+Jag1+ cardiomyogenic progenitor cells were transplanted into the hearts of severe combined immunodeficient (SCID) beige mice to document their developmental potential in vivo.

(165) Briefly, Isl1+Jag1+ cells were injected (1,000,000 cells per recipient) directly into the left ventricular wall of NOD/SCID-gamma mice in an open-chest procedure. Hearts were harvested 2-3 weeks post-surgery, fixed in 1% PFA and sectioned at 10 µm (n=12). Histological analyses of the hearts of the transplanted mice revealed the presence of GFP+ donor cells, detected by epifluorescence and by staining with an anti-GFP antibody, demonstrating that the Isl1+Jag1+ cardiomyogenic progenitor cells were capable of differentiating into cardiomyocytes when transplanted in vivo.

(166) The Isl1+Jag1+ cardiomyogenic progenitor cells were also transplanted directly into infarcted hearts of SCID beige mice ("injured mice"), as compared to similarly transplanted normal mice. When analyzed two weeks later, injured mice transplanted with the Isl1+Jag1+ cardiomyogenic progenitor cells had a larger graft size than the normal mice similarly transplanted, demonstrating the cardiomyocyte regeneration capacity of the Isl1+Jag1+ cardiomyogenic progenitor cells in vivo.

Example 5: Identification of Frizzled 4 as a Cell Surface Marker of Cardiac Progenitor Cells

(167) As described in Example 2, Frizzled 4 (FZD4) was identified by RNA-seq analysis as being expressed in cardiac progenitor cells. Thus, to confirm FZD4 as a cell surface marker of cardiac progenitor cells, FZD4 expression was assessed during cardiac differentiation via Western blot

analysis. The results, as shown in FIG. 5, demonstrated that FZD4 was not expressed in pluripotent stem cells and the first 3 days of differentiated cells. However, FZD4 started to express on day 4 and maximize its expression on day 5 of expression.

(168) In order to quantify the co-expression pattern of FZD4 and Isl1 at the single cell level, FACS analysis was performed. As shown in FIG. 6, on day 5 of differentiation, more than 83% of cells express both Isl1 and FZD4, demonstrating that FZD4 is a cell surface marker for Isl1 positive cells during cardiac progenitor differentiation using the GiWi protocol.

(169) In order to confirm that both JAG1 and FZD4 were indeed co-expressed with ISL1 on the human ventricular progenitor cells, triple immunofluorescence analysis of day 6 differentiated cells from hPSCs was performed with antibodies to Islet 1, Jagged 1 and Frizzled 4. The triple staining experiment demonstrated that Isl1+ cells expressed both Jagged 1 and Frizzled 4.

Example 6: Human Ventricular Progenitors (HVPs) Generate a 3-D Ventricular Heart Muscle Organ In Vivo

(170) The building of the ventricular heart muscle chamber is one of the most critical and earliest steps during human organogenesis, and requires a series of coordinated steps, including migration, proliferation, vascularization, assembly, and matrix alignment. To test the capacity of HVPs to drive ventriculogenesis in vivo, we transplanted purified HVPs or unpurified HVPs (92.0±1.9% ISL1+) under the kidney capsule of immunocompromised mice. After 2 months post-transplantation, animals transplanted with unpurified HVPs formed tumors, resulting in a tumor formation efficiency of 100% (100%, 4/4), whereas animals transplanted with purified HVPs did not form any tumors (0%, 0/10).

(171) The engrafted kidneys with purified HVPs were further assayed for histological analysis. Hematoxylin and Eosin (H&E) staining revealed an organ that exceeded 0.5 cm in length with more than 1 mm thickness on the surface of the mouse kidney, and that uniformly expressed the ventricular specific marker MLC2v (O'Brien, T. X. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:5157-5161). The resulting human muscle organ was fully vascularized and red blood cells could be detected in the blood vessels. Analysis of cTnT, MLC2v, and MLC2a immunostaining further revealed that the transplanted HVPs not only differentiated into cardiac muscle cells (cTnT+ cells), but also further mature to become MLC2v+ ventricular myocytes that are negative for MLC2a expression. The resulting ventricular muscle organ is fully vascularized by murine derived vascular cells, consistent with the notion that its vascularization occurred via paracrine cues derived from the HVPs.

(172) The blood vessel structure was revealed by immunostaining analysis of antibodies directed against VE-cadherin and smooth muscle actin expression. In addition, using a human specific monoclonal laminin antibody targeting laminin γ -1 chain, the HVPs secreted their own human laminin as their extracellular matrix (the mouse kidney region is negative for human laminin immunostaining). In addition, we found human fibronectin expression is restricted to areas near the blood vessels using a monoclonal human fibronectin antibody.

(173) To assess the capacity of late stage cardiac cells to drive ventriculogenesis, NKX2.5+ cells (day 10 after differentiation) were transplanted under the kidney capsule of immunocompromised NSG mice. At three weeks post-transplantation, animals transplanted with NKX2.5+ cells did not form any visible human muscle graft, indicating that HVPs lose their ability for in vivo ventriculogenesis following peak Islet-1 expression.

(174) Taken together, these studies indicate that the HVPs can synthesize and release their own cardiac laminin-derived matrix, as well as fibronectin which serves to stabilize the vasculature to the nascent ventricular organ.

Example 7: HVPs Create a Mature, Functioning Ventricular Muscle Organ In Vivo Via a Cell Autonomous Pathway

(175) One of the critical limitations for the utility of hPSCs for studies of human cardiac biology and disease is their lack of maturity and persistence of expression of fetal isoforms. To determine if

the HVP derived organs could become functional mature ventricular muscle, long term transplantation studies were performed followed by detailed analyses of a panel of well accepted features of adult ventricular myocardium including formation of T tubules (Brette, F. and Orchard, C. (2003) *Circ. Res.* 92:1182-1192; Marks, A. R. (2013) *J. Clin. Invest.* 123:46-52), ability to generate force comparable to other studies of engineered ventricular tissue, loss of automaticity, and acquisition of adult-rod shaped ventricular cardiomyocytes.

(176) After 5 months post-transplantation of purified HVPs, no tumors formed in all of our animals. Animals were sacrificed and the engrafted kidneys were removed for further analysis. The 5-month human graft was a hemisphere structure with the radius of 0.4 cm (diameter of 0.8 cm). The volume for the 5-month human graft was around 0.13 cm³ for one kidney, a volume that suggests feasibility for generating human ventricular muscle that achieves a thickness comparable to the in vivo human adult heart. Rod-shaped mature human ventricular myocytes were observed in the human muscle organ. In addition, muscle strips taken from our mature human muscle organ generated forces (0.36 ± 0.04 mN) in response to electric stimulation and increased their force generation after treatment with a β -adrenergic agonist isoprenaline (0.51 ± 0.02 mN, $p < 0.05$ compared to control). Taken together, these studies indicate that the HVPs are capable of generating a fully functional, mature human ventricular muscle organ in vivo via a cell autonomous pathway, i.e., without the addition of other cells, genes, matrix proteins, or biomaterials.

Example 8: HVPs Migrate Towards an Epicardial Niche and Spontaneously Form a Human Ventricular Muscle Patch on the Surface of a Normal Murine Heart In Vivo

(177) The epicardium is a known niche for heart progenitors, driving the growth of the ventricular chamber during compact zone expansion, as well as serving as a home to adult epicardial progenitors that can expand after myocardial injury and that can drive vasculogenesis in response to known vascular cell fate switches, such as VEGF (Giordano, F. J. et al. (2001) *Proc. Natl. Acad. Sci. USA* 98:5780-5785; Masters, M. and Riley, P. R. (2014) *Stem Cell Res.* 13:683-692; Zangi, L. et al. (2013) *Nat. Biotechnol.* 31:898-907). To determine if the HVPs might migrate spontaneously to the epicardial surface of the normal heart, purified green fluorescent protein (GFP)-labeled HVPs were injected intra-myocardially into the hearts of immunocompromised mice. After one week or one month post-transplantation, animals were sacrificed and the engrafted hearts were removed for histology. After one week post-transplantation, the majority of GFP⁺ cells were retained in the myocardium. However, almost all the GFP⁺ cells migrated to the epicardium after one month post-transplantation. In addition, GFP⁺ cells were ISL1⁺ and Ki67⁺ after one week post-transplantation.

(178) In order to trace the differentiation potential of Isl1⁺ cells, the purified ISL1⁺JAG1⁺ cells generated from a cTnT promoter driven green fluorescent protein (GFP)-expressing hESC line (H9-cTnT-GFP) were transplanted into the hearts of severe combined immunodeficient (SCID) beige mice to document their developmental potential in vivo. One month after transplantation of Isl1⁺Jag1⁺ cells directly into the ventricle of the hearts of SCID beige mice, Hematoxylin and eosin staining revealed a human muscle strip graft present in the epicardium of the murine heart. In addition, immunohistological analyses revealed the presence of GFP⁺ donor cells detected by epifluorescence and by staining with an anti-GFP antibody. More importantly, when analysed with antibodies of MLC2v and MLC2a, the grafted human muscle strip is positive for MLC2v (100% of cells+), and negative for the atrial marker MLC2a, indicating the transplanted ISL1⁺ cells not only further differentiated to cardiac muscle cells, but also became ventricular muscle cells.

(179) Taken together, these studies indicate that the HVPs can migrate to an epicardial niche, where they expand, and subsequently differentiate in to a homogenous ventricular muscle patch, again without the addition of exogenous cells, genes, matrices, or biomaterials.

Example 9: Additional Experimental Materials and Methods

(180) In this example, additional details on the experimental materials and methods used in Examples 1-8 are provided.

(181) Maintenance of hPSCs

(182) hESCs (ES03, H9) and human iPSCs (19-9-11) were maintained on Matrigel (BD Biosciences) coated plates in mTeSR1 medium (STEMCELL Technologies) according to previous published methods (Lian, X. et al. (2013) *Nat. Proc.* 8:162-175; Lian, X. et al. (2013) *Stem Cells* 31:447-457).

(183) Human Ventricular Progenitor Generation (HVPG) Protocol

(184) hPSCs maintained on a Matrigel-coated surface in mTeSR1 were dissociated into single cells with Accutase at 37° C. for 10 min and then seeded onto a Matrigel-coated cell culture dish at 100,000-200,000 cell/cm² in mTeSR1 supplemented with 5 μM ROCK inhibitor Y-27632 (day -2) for 24 hours. At day -1, cells were cultured in mTeSR1. At day 0, cells were treated with 1 μM CHIR-98014 (Selleckchem) in RPMI supplemented with B27 minus insulin (RPMI/B27-ins) for 24 hours (day 0 to day 1), which was then removed during the medium change on day 1. At day 3, half of the medium was changed to the RPMI/B27-ins medium containing 2 μM Wnt-C59 (Selleckchem), which was then removed during the medium change on day 5. At day 6, cells were dissociated into single cells and purified with anti-JAG1 or anti-FZD4 antibody.

(185) RNA-Seq Library Construction

(186) RNA was isolated (RNeasy Mini kit, Qiagen), quantified (Qubit RNA Assay Kit, Life Technologies) and quality controlled (BioAnalyzer 2100, Agilent). RNA (800 ng) from each sample was used as input for the Illumina TruSeq mRNA Sample Prep Kit v2 (Illumina) and sequencing libraries were created according to the manufacturer's protocol. Briefly, poly-A containing mRNA molecules were purified using poly-T oligo-attached magnetic beads. Following purification, the mRNA was fragmented and copied into first strand complementary DNA using random primers and reverse transcriptase. Second strand cDNA synthesis was then done using DNA polymerase I and RNase H. The cDNA was ligated to adapters and enriched with PCR to create the final cDNA library. The library was pooled and sequenced on a HiSeq 2000 (Illumina) instrument per the manufacturer's instructions.

(187) RNA-Seq Data Processing

(188) The RNA-seq reads were trimmed and mapped to the hg19 reference using Tophat 2. On average, approximately 23 million reads were generated per sample, and 76% of these reads were uniquely mapped. Expression levels for each gene were quantified using the python script *rpkmforgenes* and annotated using RefSeq. Genes without at least one sample with at least ten reads were removed from the analysis. Principle Component Analysis and heatmaps were constructed using the R and Gene-E respectively.

(189) Transplantation

(190) Aliquots of 2 million purified HVPs were collected into an eppendorf tube. Cells were spun down, and the supernatant was discarded. Each tube of cells was transplanted under the kidney capsule, or intra-myocardially injected into the heart of the immunodeficient mice, NOD.Cg-Prkdcscid Il2rgtmlWjl/SzJ or SCID-Beige respectively (Charles River France), following a previously described protocol (Shultz, L. D. et al. (2005) *J. Immunol.* 174:6477-6489). Engrafted Kidneys or hearts are harvested at various time intervals for histological and physiological analysis.

(191) Flow Cytometry

(192) Cells were dissociated into single cells with Accutase for 10 min and then fixed with 1% paraformaldehyde for 20 min at room temperature and stained with primary and secondary antibodies in PBS plus 0.1% Triton X-100 and 0.5% BSA. Data were collected on a FACSCaliber flow cytometer (Beckton Dickinson) and analyzed using FlowJo.

(193) Immunostaining

(194) Cells were fixed with 4% paraformaldehyde for 15 min at room temperature and then stained with primary and secondary antibodies in PBS plus 0.4% Triton X-100 and 5% non-fat dry milk (Bio-Rad). Nuclei were stained with Gold Anti-fade Reagent with DAPI (Invitrogen). An epifluorescence microscope and a confocal microscope (ZEISS, LSM 700) were used for imaging

analysis.

(195) Western Blot Analysis

(196) Cells were lysed in M-PER Mammalian Protein Extraction Reagent (Pierce) in the presence of Halt Protease and Phosphatase Inhibitor Cocktail (Pierce). Proteins were separated by 10% Tris-Glycine SDS/PAGE (Invitrogen) under denaturing conditions and transferred to a nitrocellulose membrane. After blocking with 5% dried milk in TBST, the membrane was incubated with primary antibody overnight at 4° C. The membrane was then washed, incubated with an anti-mouse/rabbit peroxidase-conjugated secondary antibody at room temperature for 1 hour, and developed by SuperSignal chemiluminescence (Pierce).

(197) Electrophysiology (Patch Clamping)

(198) Beating ventricular myocyte clusters were microdissected and replated onto glass coverslips before recording. Action potential activity was assessed using borosilicate glass pipettes (4-5 M Ohm resistance) filled with intracellular solution consisting of 120 mM K D-gluconate, 25 mM KCl, 4 mM MgATP, 2 mM NaGTP, 4 mM Na₂-phospho-creatin, 10 mM EGTA, 1 mM CaCl₂, and 10 mM HEPES (pH 7.4 adjusted with HCl at 25° C.). Cultured cardiomyocytes seeded on coverslip dishes were submerged in extracellular solution (Tyrode's solution) containing 140 mM NaCl, 5.4 mM KCl, 1 mM MgCl₂, 10 mM glucose, 1.8 mM CaCl₂, and 10 mM HEPES (pH 7.4 adjusted with NaOH at 25° C.). Spontaneous action potentials were recorded at 37° C. using patch clamp technique (whole-cell, current clamp configuration) performed using a Multiclamp 700B amplifier (Molecular Devices, CA, USA) software low-pass filtered at 1 kHz, digitized and stored using a Digidata 1322A and Clampex 9.6 software (Molecular Devices, CA, USA).

(199) Statistics

(200) Data are presented as mean±standard error of the mean (SEM). Statistical significance was determined by Student's t-test (two-tail) between two groups. P<0.05 was considered statistically significant.

Example 10: Xeno-Free Human Ventricular Progenitor Differentiation Protocol

(201) In this example, an alternative differentiation protocol for differentiation of human ventricular progenitors is provided, which utilizes a defined, xeno-free culture medium, Essential 8. The Essential 8 medium was developed for growth and expansion of human pluripotent stem cells (hPSCs) and is described further in Chen, G. et al. (2011) Nat. Methods 8:424-429 (referred to therein as “E8” medium).

(202) hPSCs maintained on a Vitronectin (or Laminin 521)-coated surface in Essential 8 medium were dissociated into single cells with Versene solution at 37° C. for 10 min and then seeded onto a Vitronectin (or Laminin 521)-coated cell culture dish at 100,000-200,000 cell/cm.² in Essential 8 medium supplemented with 5 μM ROCK inhibitor Y-27632 (day -2) for 24 hours. At day -1, cells were cultured in Essential 8 medium. At day 0, cells were treated with 0.5 μM CHIR-98014 in RPMI for 24 hours (day 0 to day 1), which was then removed during the medium change on day 1. At day 3, half of the medium was changed to the RPMI medium containing 0.5 μM Wnt-059, which was then removed during the medium change on day 5. At day 6, cells (human ventricular progenitors) were dissociated into single cells and purified with anti-JAG1 or anti-FZD4 antibody. Alternatively cells are purified with anti-LIFR or anti-FGFR3 antibody.

Example 11: Angiogenic Markers for Engraftable Human Ventricular Progenitor Cells

(203) In this example, genes in the angiogenic family that are expressed in human ventricular progenitor cells (HVPs) were identified. HVPs were generated as described in Examples 1 or 10 and RNA sequencing (RNA-seq) was performed at different time points following differentiation as described in Example 1. Cluster analysis of gene expression profiles at different time points during HVP differentiation identified stage-specific signature genes. These genes were clustered hierarchically on the basis of the similarity of their expression profiles. First, genes showing expression in four different categories were identified: (i) cell surface expression; (ii) co-expression with Islet 1; (iii) high expression on day 5 of differentiation; and (iv) high d5/d0 ratio. This analysis

confirmed the cell surface markers for HVPs of: JAG1, FZD4, FGFR3, LIFR (CD118) and TNFSF9. Next, from this same population of HVPs that identified the cell surface markers, gene ontology searches were performed to identify angiogenic family genes that were expressed in this population of HVPs, to thereby identify a gene fingerprint profile that identifies genes critical for cell engraftment.

(204) Statistically, Pearson's correlation with Isl1 expression was used to identify those angiogenic genes whose expression in the HVPs best correlated with Isl1 expression. Table 1 below lists the angiogenic genes that correlate with Isl1 expression with a Pearson's correlation of 0.50 or higher.

(205) TABLE-US-00001 TABLE 1 Angiogenic genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of 0.50 or greater Pearson's Correlation with Isl1 Gene Angiogenic genes (GO: 0001525) Expression FGF10 fibroblast growth factor 10 0.98 PRKD1 protein kinase D1 0.95 CCBE1 collagen and calcium binding EGF domains 1 0.94 PDGFRA platelet-derived growth factor receptor, alpha 0.94 polypeptide EPHB2 EPH receptor B2 0.92 GATA2 GATA binding protein 2 0.92 NTRK1 neurotrophic tyrosine kinase, receptor, type 1 0.92 PTGIS prostaglandin I2 (prostacyclin) synthase 0.87 BMPER BMP binding endothelial regulator 0.85 BMP4 bone morphogenetic protein 4 0.84 C1GALT1 core 1 synthase, glycoprotein-N- 0.84 acetylgalactosamine 3-beta- galactosyltransferase 1 MEIS1 Meis homeobox 1 0.83 TBX1 T-box 1 0.83 PKNOX1 PBX/knotted 1 homeobox 1 0.83 ID1 inhibitor of DNA binding 1, dominant 0.82 negative helix-loop-helix protein TCF21 transcription factor 21 0.82 HEY1 hes-related family bHLH transcription factor 0.80 with YRPW motif 1 HOXB3 homeobox B3 0.78 JAG1 jagged 1 0.75 HGF hepatocyte growth factor (hepapoietin A; 0.74 scatter factor) IL6 interleukin 6 0.74 GHRL ghrelin/obestatin prepropeptide 0.73 IHH indian hedgehog 0.70 SRPK2 SRSF protein kinase 2 0.70 GATA6 GATA binding protein 6 0.69 HAND1 heart and neural crest derivatives expressed 1 0.69 AMOT angiomin 0.69 NRP2 neuropilin 2 0.65 PTEN phosphatase and tensin homolog 0.65 SEMA3E sema domain, immunoglobulin domain (Ig), 0.64 short basic domain, secreted, (semaphorin) 3E APOLD1 apolipoprotein L domain containing 1 0.62 SETD2 SET domain containing 2 0.62 DAB2IP DAB2 interacting protein 0.61 KDR kinase insert domain receptor 0.60 PGF placental growth factor 0.60 EMP2 epithelial membrane protein 2 0.59 TAL1 T-cell acute lymphocytic leukemia 1 0.58 ACVR1 activin A receptor, type I 0.58 HIPK2 homeodomain interacting protein kinase 2 0.56 CSPG4 chondroitin sulfate proteoglycan 4 0.55 TNFAIP3 tumor necrosis factor, alpha-induced protein 3 0.55 NRP1 neuropilin 1 0.55 NFATC4 nuclear factor of activated T-cells, 0.54 cytoplasmic, calcineurin-dependent 4 CDC42 cell division cycle 42 0.54 ANGPTL4 angiopoietin-like 4 0.53 BCAS3 breast carcinoma amplified sequence 3 0.53 HIPK1 homeodomain interacting protein kinase 1 0.53 NRXN3 neurexin 3 0.52 FZD5 frizzled class receptor 5 0.52 HHEX hematopoietically expressed homeobox 0.50

Table 2 below lists the angiogenic genes that correlate with Isl1 expression with a Pearson's correlation of 0.49-0.00.

(206) TABLE-US-00002 TABLE 2 Angiogenic genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of 0.49 to 0.00 Pearson's Correlation with Isl1 Gene Angiogenic genes (GO: 0001525) Expression ACVRL1 activin A receptor type II-like 1 0.49 ENPEP glutamyl aminopeptidase (aminopeptidase A) 0.49 EFNA1 ephrin-A1 0.49 CHRNA7 cholinergic receptor, nicotinic, alpha 7 0.49 (neuronal) TMEM100 transmembrane protein 100 0.48 NOS3 nitric oxide synthase 3 (endothelial cell) 0.47 LEF1 lymphoid enhancer-binding factor 1 0.47 NRXN1 neurexin 1 0.46 EPHB3 EPH receptor B3 0.44 ROCK1 Rho-associated, coiled-coil containing 0.42 protein kinase 1 NF1 neurofibromin 1 0.42 CYSLTR2 cysteinyl leukotriene receptor 2 0.42 FGFR2 fibroblast growth factor receptor 2 0.41 GATA4 GATA binding protein 4 0.40 FMNL3 formin-like 3 0.40 C3 complement component 3 0.40 WASF2 WAS protein family, member 2 0.40 CALCRL calcitonin receptor-like 0.39 HIF1A hypoxia inducible factor 1, alpha subunit 0.39 (basic helix-loop-helix transcription factor) VEGFA vascular endothelial growth factor A 0.39 KRIT1 KRIT1, ankyrin repeat containing 0.39 CDH13 cadherin 13 0.39 COL18A1 collagen, type XVIII, alpha 1

0.39 STK4 serine/threonine kinase 4 0.38 C5 complement component 5 0.38 HDAC7 histone deacetylase 7 0.38 ANGPT2 angiopoietin 2 0.38 PLCG1 phospholipase C, gamma 1 0.37 EDNRA endothelin receptor type A 0.35 TGFB2 transforming growth factor, beta 2 0.35 HAND2 heart and neural crest derivatives expressed 2 0.35 CD34 CD34 molecule 0.35 BTG1 B-cell translocation gene 1, anti-proliferative 0.34 TGFBR1 transforming growth factor, beta receptor 1 0.33 FGFR1 fibroblast growth factor receptor 1 0.33 FN1 fibronectin 1 0.31 TWIST1 twist family bHLH transcription factor 1 0.31 ELK3 ELK3, ETS-domain protein (SRF accessory 0.30 protein 2) THSD7A thrombospondin, type I, domain containing 0.30 7A RGCC regulator of cell cycle 0.30 PLCD1 phospholipase C, delta 1 0.29 SPARC secreted protein, acidic, cysteine-rich 0.29 (osteonectin) TBX20 T-box 20 0.28 PIK3CA phosphatidylinositol-4,5-bisphosphate 0.27 3-kinase, catalytic subunit alpha MMRN2 multimerin 2 0.27 FOXO4 forkhead box O4 0.26 RAMP2 receptor (G protein-coupled) activity 0.25 modifying protein 2 FLT1 fms-related tyrosine kinase 1 0.25 ADRB2 adrenoceptor beta 2, surface 0.25 SLC12A6 solute carrier family 12 (potassium/chloride 0.25 transporter), member 6 ADM adrenomedullin 0.25 NPPB natriuretic peptide B 0.24 SPINK5 serine peptidase inhibitor, Kazal type 5 0.24 MAPK14 mitogen-activated protein kinase 14 0.24 MMP2 matrix metalloproteinase 2 0.24 PTPRM protein tyrosine phosphatase, receptor 0.23 type, M OVOL2 ovo-like zinc finger 2 0.23 CTNNB1 catenin (cadherin-associated protein), beta 1, 0.22 88 kDa OTULIN OTU deubiquitinase with linear linkage 0.21 specificity B4GALT1 UDP-Gal: betaGlcNAc beta 1,4- 0.21 galactosyltransferase, polypeptide 1 PDGFRB platelet-derived growth factor receptor, beta 0.20 polypeptide F3 coagulation factor III (thromboplastin, tissue 0.20 factor) PRKCA protein kinase C, alpha 0.20 LRP5 low density lipoprotein receptor-related 0.20 protein 5 MAP3K7 mitogen-activated protein kinase kinase 0.20 kinase 7 NRCAM neuronal cell adhesion molecule 0.19 MAP2K5 mitogen-activated protein kinase kinase 5 0.18 S1PR1 sphingosine-1-phosphate receptor 1 0.18 NFATC3 nuclear factor of activated T-cells, 0.18 cytoplasmic, calcineurin-dependent 3 TSPAN12 tetraspanin 12 0.18 LAMAS laminin, alpha 5 0.17 LOXL2 lysyl oxidase-like 2 0.17 ANGPT1 angiopoietin 1 0.17 GTF2I general transcription factor Iii 0.16 E2F8 E2F transcription factor 8 0.16 PDE3B phosphodiesterase 3B, cGMP-inhibited 0.15 SHB Src homology 2 domain containing adaptor 0.14 protein B MYH9 myosin, heavy chain 9, non-muscle 0.14 FZD8 frizzled class receptor 8 0.14 NOV nephroblastoma overexpressed 0.14 SH2D2A SH2 domain containing 2A 0.14 FGF8 fibroblast growth factor 8 (androgen-induced) 0.13 TIE1 tyrosine kinase with immunoglobulin-like 0.13 and EGF-like domains 1 EGLN1 egl-9 family hypoxia-inducible factor 1 0.12 RORA RAR-related orphan receptor A 0.11 MFGE8 milk fat globule-EGF factor 8 protein 0.11 ARHGAP24 Rho GTPase activating protein 24 0.10 ITGA5 integrin, alpha 5 (fibronectin receptor, alpha 0.10 polypeptide) PARVA parvin, alpha 0.10 ADIPOR2 adiponectin receptor 2 0.09 NPR1 natriuretic peptide receptor 1 0.09 ITGB1 integrin, beta 1 (fibronectin receptor, beta 0.09 polypeptide, antigen CD29 includes MDF2, MSK12) HIF3A hypoxia inducible factor 3, alpha subunit 0.08 EPAS1 endothelial PAS domain protein 1 0.08 FOXC2 forkhead box C2 0.07 ANXA2 annexin A2 0.06 RBM15 RNA binding motif protein 15 0.06 PITX2 paired-like homeodomain 2 0.06 FOXC1 forkhead box C1 0.06 SRF serum response factor 0.06 ECSCR endothelial cell surface expressed chemotaxis 0.05 and apoptosis regulator SOX17 SRY (sex determining region Y)-box 17 0.04 HDAC5 histone deacetylase 5 0.04 LRG1 leucine-rich alpha-2-glycoprotein 1 0.04 ADAM8 ADAM metalloproteinase domain 8 0.03 UBP1 upstream binding protein 1 (LBP-1a) 0.02 VASH1 vasohibin 1 0.02 ANXA3 annexin A3 0.01 RRAS related RAS viral (r-ras) oncogene homolog 0.01 TYMP thymidine phosphorylase 0.01 PRCP prolylcarboxypeptidase (angiotensinase C) 0.01 SEMA5A sema domain, seven thrombospondin repeats 0.00 (type 1 and type 1-like), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 5A GREM1 gremlin 1, DAN family BMP antagonist 0.00

(207) Angiogenic genes whose expression negatively correlated with Isl1 expression in the HVPs were also identified. Table 3 below lists the angiogenic genes that negatively correlate with Isl1

expression with a Pearson's correlation of -0.50 or less.

(208) TABLE-US-00003 TABLE 3 Angiogenic genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of -0.50 or less Pearson's Correlation with Gene Angiogenic genes (GO: 0001525) Isl1 Expression ETS1 v-ets avian erythroblastosis virus E26 -0.50 oncogene homolog 1 BAX BCL2-associated X protein -0.50 XBP1 X-box binding protein 1 -0.52 TDGF1 teratocarcinoma-derived growth factor 1 -0.53 C5AR1 complement component 5a receptor 1 -0.53 EPHA1 EPH receptor A1 -0.53 HS6ST1 heparan sulfate 6-O-sulfotransferase 1 -0.56 SHC1 SHC (Src homology 2 domain containing) -0.56 transforming protein 1 SP100 SP100 nuclear antigen -0.58 JAM3 junctional adhesion molecule 3 -0.58 CASP8 caspase 8, apoptosis-related cysteine -0.60 peptidase FLT4 fms-related tyrosine kinase 4 -0.60 SFRP2 secreted frizzled-related protein 2 -0.61 HPSE heparanase -0.61 BAK1 BCL2-antagonist/killer 1 -0.65 GPX1 glutathione peroxidase 1 -0.65 VAV3 vav 3 guanine nucleotide exchange factor -0.70 VAV2 vav 2 guanine nucleotide exchange factor -0.72 EGF epidermal growth factor -0.72 ADAM15 ADAM metallopeptidase domain 15 -0.73 AGGF1 angiogenic factor with G patch and FHA -0.76 domains 1

Table 4 below lists the angiogenic genes that negatively correlate with Isl1 expression with a Pearson's correlation of -0.01 to -0.49 .

(209) TABLE-US-00004 TABLE 4 Angiogenic genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of -0.01 to -0.49 Pearson's Correlation with Gene Angiogenic genes (GO: 0001525) Isl1 Expression EIF2AK3 eukaryotic translation initiation factor -0.01 2-alpha kinase 3 ROCK2 Rho-associated, coiled-coil containing -0.01 protein kinase 2 WNT5A wingless-type MMTV integration site -0.02 family, member 5A NR4A1 nuclear receptor subfamily 4, group A, -0.02 member 1 CYP1B1 cytochrome P450, family 1, subfamily B, -0.02 polypeptide 1 PTK2 protein tyrosine kinase 2 -0.03 SFRP1 secreted frizzled-related protein 1 -0.04 STAT1 signal transducer and activator of -0.04 transcription 1, 91 kDa ITGAV integrin, alpha V -0.04 EPHB4 EPH receptor B4 -0.05 CYR61 cysteine-rich, angiogenic inducer, 61 -0.05 TEK TEK tyrosine kinase, endothelial -0.06 COL15A1 collagen, type XV, alpha 1 -0.06 COL4A1 collagen, type IV, alpha 1 -0.07 ANG angiogenin, ribonuclease, RNase A -0.07 family, 5 HSPB1 heat shock 27 kDa protein 1 -0.07 PLXND1 plexin D1 -0.08 HSPG2 heparan sulfate proteoglycan 2 -0.09 VEGFC vascular endothelial growth factor C -0.09 SYNJ2BP synaptojanin 2 binding protein -0.09 THBS1 thrombospondin 1 -0.09 CTGF connective tissue growth factor -0.10 ITGB3 integrin, beta 3 (platelet glycoprotein Ma, -0.12 antigen CD61) AAMP angio-associated, migratory cell protein -0.12 GJA5 gap junction protein, alpha 5, 40 kDa -0.12 PRKCB protein kinase C, beta -0.13 EGR3 early growth response 3 -0.13 JMJD6 jumonji domain containing 6 -0.13 TGFBI transforming growth factor, beta-induced, -0.14 68 kDa SIRT1 sirtuin 1 -0.14 ANGPTL3 angiopoietin-like 3 -0.14 ACKR3 atypical chemokine receptor 3 -0.14 SAT1 spermidine/spermine N1-acetyltransferase -0.15 1 VEGFB vascular endothelial growth factor B -0.16 UTS2 urotensin 2 -0.16 JUN jun proto-oncogene -0.16 TNFSF12 tumor necrosis factor (ligand) -0.16 superfamily, member 12 EGFL7 EGF-like-domain, multiple 7 -0.17 MED1 mediator complex subunit 1 -0.17 SLIT2 slit guidance ligand 2 -0.17 SERPINF1 serpin peptidase inhibitor, clade F -0.18 (alpha-2 antiplasmin, pigment epithelium derived factor), member 1 NOTCH3 notch 3 -0.18 FGF9 fibroblast growth factor 9 -0.19 DLL4 delta-like 4 (Drosophila) -0.19 CCL2 chemokine (C-C motif) ligand 2 -0.19 MMP14 matrix metallopeptidase 14 (membrane- -0.19 inserted) TMPRSS6 transmembrane protease, serine 6 -0.19 EPGN epithelial mitogen -0.20 RBPJ recombination signal binding protein for -0.20 immunoglobulin kappa J region COL4A2 collagen, type IV, alpha 2 -0.20 PRKD2 protein kinase D2 -0.20 ALOX12 arachidonate 12-lipoxygenase -0.21 RNH1 ribonuclease/angiogenin inhibitor 1 -0.21 APOH apolipoprotein H (beta-2-glycoprotein I) -0.21 CHI3L1 chitinase 3-like 1 (cartilage glycoprotein- -0.21 39) ESM1 endothelial cell-specific molecule 1 -0.22 PTGS2 prostaglandin-endoperoxide synthase 2 -0.22 (prostaglandin G/H synthase and cyclooxygenase) ANPEP alanyl (membrane)

aminopeptidase -0.22 LEMD3 LEM domain containing 3 -0.22 UTS2R uterine 2 receptor
-0.22 CIB1 calcium and integrin binding 1 (calmyrin) -0.22 ITGB1BP1 integrin beta 1 binding
protein 1 -0.22 AQP1 aquaporin 1 (Colton blood group) -0.22 IL18 interleukin 18 -0.22 EPHA2
EPH receptor A2 -0.22 EPHB1 EPH receptor B1 -0.22 AGT angiotensinogen (serpin peptidase
-0.22 inhibitor, clade A, member 8) PLAU plasminogen activator, urokinase -0.22 VEZF1
vascular endothelial zinc finger 1 -0.23 SPHK1 sphingosine kinase 1 -0.23 SRPX2 sushi-repeat
containing protein, -0.23 X-linked 2 PDCL3 phosducin-like 3 -0.23 COL8A1 collagen, type VIII,
alpha 1 -0.24 HDAC9 histone deacetylase 9 -0.24 CTSH cathepsin H -0.24 EDN1 endothelin 1
-0.24 CXCL8 chemokine (C-X-C motif) ligand 8 -0.24 ECM1 extracellular matrix protein 1 -0.24
BRCA1 breast cancer 1, early onset -0.24 EFNB2 ephrin-B2 -0.25 SERPINE1 serpin peptidase
inhibitor, clade E (nexin, -0.25 plasminogen activator inhibitor type 1), member 1 SASH1 SAM
and SH3 domain containing 1 -0.25 WNT7B wingless-type MMTV integration site -0.25 family,
member 7B RAMP1 receptor (G protein-coupled) activity -0.26 modifying protein 1 SCG2
secretogranin II -0.26 COL8A2 collagen, type VIII, alpha 2 -0.26 SULF1 sulfatase 1 -0.26 CLIC4
chloride intracellular channel 4 -0.26 FGF1 fibroblast growth factor 1 (acidic) -0.27 NODAL
nodal growth differentiation factor -0.27 RASIP1 Ras interacting protein 1 -0.28 RLN2 relaxin 2
-0.28 POFUT1 protein O-fucosyltransferase 1 -0.28 FGF18 fibroblast growth factor 18 -0.28
AIMP1 aminoacyl tRNA synthetase complex- -0.28 interacting multifunctional protein 1 TGFBR2
transforming growth factor, beta receptor -0.28 II (70/80 kDa) RHOB ras homolog family member
B -0.28 GBX2 gastrulation brain homeobox 2 -0.28 ENPP2 ectonucleotide -0.29
pyrophosphatase/phosphodiesterase 2 MAPK7 mitogen-activated protein kinase 7 -0.30 PROK2
prokineticin 2 -0.30 E2F7 E2F transcription factor 7 -0.30 ERAP1 endoplasmic reticulum
aminopeptidase 1 -0.31 MTDH metadherin -0.31 KLF5 Kruppel-like factor 5 (intestinal) -0.31
DICER1 dicer 1, ribonuclease type III -0.32 LECT1 leukocyte cell derived chemotaxin 1 -0.32
CX3CL1 chemokine (C-X3-C motif) ligand 1 -0.32 PTK2B protein tyrosine kinase 2 beta -0.33
SEMA4A sema domain, immunoglobulin domain -0.34 (Ig), transmembrane domain (TM) and
short cytoplasmic domain, (semaphorin) 4A ARHGAP22 Rho GTPase activating protein 22 -0.34
RSPO3 R-spondin 3 -0.34 KLF4 Kruppel-like factor 4 (gut) -0.34 ROBO1 roundabout guidance
receptor 1 -0.34 GPLD1 glycosylphosphatidylinositol specific -0.35 phospholipase D1 NUS1
NUS1 dehydrolipidyl diphosphate -0.35 synthase subunit NRARP NOTCH-regulated ankyrin
repeat protein -0.35 PDCD10 programmed cell death 10 -0.36 PF4 platelet factor 4 -0.36 PRKX
protein kinase, X-linked -0.36 PML promyelocytic leukemia -0.36 ATP5B ATP synthase, H+
transporting, -0.36 mitochondrial F1 complex, beta polypeptide TNFRSF12A tumor necrosis
factor receptor -0.36 superfamily, member 12A ENG endoglin -0.37 THY1 Thy-1 cell surface
antigen -0.37 FGF2 fibroblast growth factor 2 (basic) -0.37 CXCL12 chemokine (C-X-C motif)
ligand 12 -0.37 CAV1 caveolin 1, caveolae protein, 22 kDa -0.38 PDGFA platelet-derived growth
factor alpha -0.38 polypeptide PNPLA6 patatin-like phospholipase domain -0.38 containing 6
PLCD3 phospholipase C, delta 3 -0.38 DDAH1 dimethylarginine -0.39 dimethylaminohydrolase 1
GNA13 guanine nucleotide binding protein (G -0.39 protein), alpha 13 ADM2 adrenomedullin 2
-0.39 HMOX1 heme oxygenase 1 -0.40 MCAM melanoma cell adhesion molecule -0.41
RAPGEF3 Rap guanine nucleotide exchange factor -0.41 (GEF) 3 TNFAIP2 tumor necrosis factor,
alpha-induced -0.41 protein 2 HTATIP2 HIV-1 Tat interactive protein 2, 30 kDa -0.42 NCL
nucleolin -0.42 ERBB2 erb-b2 receptor tyrosine kinase 2 -0.43 NAA15 N(alpha)-acetyltransferase
15, NatA -0.43 auxiliary subunit ATP1F1 ATPase inhibitory factor 1 -0.43 THBS4
thrombospondin 4 -0.43 SYK spleen tyrosine kinase -0.44 LIF leukemia inhibitory factor -0.44
THBS2 thrombospondin 2 -0.44 PPP1R16B protein phosphatase 1, regulatory subunit -0.44 16B
NOTCH1 notch 1 -0.44 RUNX1 runt-related transcription factor 1 -0.45 PDCD6 programmed cell
death 6 -0.45 VASH2 vasohibin 2 -0.45 GPI glucose-6-phosphate isomerase -0.46 ZC3H12A zinc
finger CCCH-type containing 12A -0.46 WARS tryptophanyl-tRNA synthetase -0.46 HYAL1
hyaluronoglucosaminidase 1 -0.47 PIK3CB phosphatidylinositol-4,5-bisphosphate -0.47 3-kinase,

catalytic subunit beta TNMD tenomodulin -0.49

Example 12: Extracellular Matrix Markers for Engraftable Human Ventricular Progenitor Cells (210) In this example, genes in the extracellular matrix family that are expressed in human ventricular progenitor cells (HVPs) were identified. HVPs were generated as described in Examples 1 or 10 and RNA sequencing (RNA-seq) was performed at different time points following differentiation as described in Example 1. Cluster analysis of gene expression profiles at different time points during HVP differentiation identified stage-specific signature genes. These genes were clustered hierarchically on the basis of the similarity of their expression profiles. First, genes showing expression in four different categories were identified: (i) cell surface expression; (ii) co-expression with Isl1; (iii) high expression on day 5 of differentiation; and (iv) high d5/d0 ratio. This analysis confirmed the cell surface markers for HVPs of: JAG1, FZD4, FGFR3, LIFR (CD118) and TNFSF9. Next, from this same population of HVPs that identified the cell surface markers, gene ontology searches were performed to identify extracellular matrix family genes that were expressed in this population of HVPs, to thereby identify a gene fingerprint profile that identifies genes critical for cell engraftment.

(211) Statistically, Pearson's correlation with Isl1 expression was used to identify those extracellular matrix genes whose expression in the HVPs best correlated with Isl1 expression. Table 5 below lists the extracellular matrix genes that correlate with Isl1 expression with a Pearson's correlation of 0.50 or higher.

(212) TABLE-US-00005 TABLE 5 Extracellular matrix genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of 0.50 or greater Pearson's Correlation with Isl1 Gene Extracellular matrix genes (GO: 0031012) Expression FGF10 fibroblast growth factor 10 0.98 SMOG1 SPARC related modular calcium binding 1 0.97 CCBE1 collagen and calcium binding EGF domains 1 0.94 COL6A6 collagen, type VI, alpha 6 0.89 ADAMTS12 ADAM metalloproteinase with thrombospondin 0.85 type 1 motif, 12 COL19A1 collagen, type XIX, alpha 1 0.85 LAMA1 laminin, alpha 1 0.85 BMP4 bone morphogenetic protein 4 0.84 FBLN7 fibulin 7 0.81 FBLN2 fibulin 2 0.81 NDNF neuron-derived neurotrophic factor 0.80 HTRA1 HtrA serine peptidase 1 0.80 HAPLN1 hyaluronan and proteoglycan link protein 1 0.79 EMILIN1 elastin microfibril interfacer 1 0.79 SPOCK3 sparc/osteonectin, cwcv and kazal-like 0.76 domains proteoglycan (testican) 3 PODNL1 podocan-like 1 0.73 IHH indian hedgehog 0.70 ACAN aggrecan 0.69 NID2 nidogen 2 (osteonidogen) 0.69 COL4A6 collagen, type IV, alpha 6 0.68 LAMC1 laminin, gamma 1 (formerly LAMB2) 0.65 FMOD fibromodulin 0.65 MUC4 mucin 4, cell surface associated 0.64 EMID1 EMI domain containing 1 0.62 HMCN1 hemiscentin 1 0.61 NID1 nidogen 1 0.60 VCAN versican 0.58 CILP2 cartilage intermediate layer protein 2 0.57 SOD3 superoxide dismutase 3, extracellular 0.56 ADAMTS3 ADAM metalloproteinase with thrombospondin 0.54 type 1 motif, 3 ZP3 zona pellucida glycoprotein 3 (sperm receptor) 0.54 ANGPTL4 angiopoietin-like 4 0.53 CRTAC1 cartilage acidic protein 1 0.52 LTBP4 latent transforming growth factor beta binding 0.50 protein 4 FREM1 FRAS1 related extracellular matrix 1 0.50

Table 6 below lists the extracellular matrix genes that correlate with Isl1 expression with a Pearson's correlation of 0.49-0.00.

(213) TABLE-US-00006 TABLE 6 Extracellular matrix genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of 0.49 to 0.00 Pearson's Correlation with Isl1 Gene Extracellular matrix genes (GO: 0031012) Expression SSC5D scavenger receptor cysteine rich family, 5 0.49 domains GPC6 glypican 6 0.49 COL1A1 collagen, type I, alpha 1 0.49 ADAMTSL3 ADAMTS-like 3 0.48 FLRT3 fibronectin leucine rich transmembrane 0.48 protein 3 FBLN1 fibulin 1 0.48 ADAMTS9 ADAM metalloproteinase with thrombospondin 0.48 type 1 motif, 9 COL27A1 collagen, type XXVII, alpha 1 0.47 RELN reelin 0.46 COL9A2 collagen, type IX, alpha 2 0.46 EFEMP2 EGF containing fibulin-like extracellular 0.45 matrix protein 2 AGRN agrin 0.44 PCOLCE procollagen C-endopeptidase enhancer 0.44 NTN4 netrin 4 0.44 CD248 CD248

molecule, endoglin 0.44 TGF β 1 transforming growth factor, β 1 0.43 ADAMTS2 ADAM metalloproteinase with thrombospondin 0.43 type 1 motif, 2 CTHRC1 collagen triple helix repeat containing 1 0.42 FGFR2 fibroblast growth factor receptor 2 0.41 APOE apolipoprotein E 0.41 MMP11 matrix metalloproteinase 11 0.41 MMP15 matrix metalloproteinase 15 0.41 (membrane-inserted) PODN podocan 0.39 VEGFA vascular endothelial growth factor A 0.39 COL18A1 collagen, type XVIII, α 1 0.39 GLG1 golgi glycoprotein 1 0.39 GPC2 glypican 2 0.37 DAG1 dystroglycan 1 (dystrophin-associated 0.35 glycoprotein 1) TGF β 2 transforming growth factor, β 2 0.35 PRELP proline/arginine-rich end leucine-rich repeat 0.35 protein CHAD chondroadherin 0.33 COL2A1 collagen, type II, α 1 0.33 FN1 fibronectin 1 0.31 SMC3 structural maintenance of chromosomes 3 0.31 COL4A5 collagen, type IV, α 5 0.30 FBN3 fibrillin 3 0.30 MMP23B matrix metalloproteinase 23B 0.30 CCDC80 coiled-coil domain containing 80 0.29 SPARC secreted protein, acidic, cysteine-rich 0.29 (osteonectin) TNXB tenascin XB 0.28 COL6A2 collagen, type VI, α 2 0.28 ADAMTS13 ADAM metalloproteinase with thrombospondin 0.28 type 1 motif, 13 LOXL1 lysyl oxidase-like 1 0.28 HAPLN2 hyaluronan and proteoglycan link protein 2 0.28 TNC tenascin C 0.28 ENTPD2 ectonucleoside triphosphate 0.28 diphosphohydrolase 2 TGF β 3 transforming growth factor, β 3 0.28 MFAP4 microfibrillar-associated protein 4 0.27 VWF von Willebrand factor 0.27 WNT2 wingless-type MMTV integration site family 0.27 member 2 MMRN2 multimerin 2 0.27 SPON1 spondin 1, extracellular matrix protein 0.26 ADAMTS1 ADAM metalloproteinase with thrombospondin 0.26 type 1 motif, 1 F2 coagulation factor II (thrombin) 0.26 FLRT2 fibronectin leucine rich transmembrane 0.25 protein 2 MMP2 matrix metalloproteinase 2 0.24 COL26A1 collagen, type XXVI, α 1 0.24 CASK calcium/calmodulin-dependent serine protein 0.24 kinase (MAGUK family) NTN3 netrin 3 0.23 SLC1A3 solute carrier family 1 (glial high affinity 0.22 glutamate transporter), member 3 F3 coagulation factor III (thromboplastin, tissue 0.20 factor) ADAMTS6 ADAM metalloproteinase with thrombospondin 0.20 type 1 motif, 6 COL5A2 collagen, type V, α 2 0.19 ERBB2IP erbb2 interacting protein 0.18 LAMB1 laminin, β 1 0.18 COLQ collagen-like tail subunit (single strand of 0.18 homotrimer) of asymmetric acetylcholinesterase LAMAS laminin, α 5 0.17 LOXL2 lysyl oxidase-like 2 0.17 WNT11 wingless-type MMTV integration site family, 0.17 member 11 LAMB2 laminin, β 2 (laminin S) 0.17 COL5A1 collagen, type V, α 1 0.17 AEBP1 AE binding protein 1 0.17 COL9A3 collagen, type IX, α 3 0.16 CTSD cathepsin D 0.16 COL21A1 collagen, type XXI, α 1 0.16 EGFLAM EGF-like, fibronectin type III and laminin G 0.16 domains FBN2 fibrillin 2 0.15 NAV2 neuron navigator 2 0.15 EMILIN2 elastin microfibril interfacier 2 0.14 WNT9B wingless-type MMTV integration site family, 0.14 member 9B NOV neuroblastoma overexpressed 0.14 CHL1 cell adhesion molecule L1-like 0.13 DLG1 discs, large homolog 1 (Drosophila) 0.11 MFGE8 milk fat globule-EGF factor 8 protein 0.11 TIMP1 TIMP metalloproteinase inhibitor 1 0.11 CST3 cystatin C 0.10 APLP1 amyloid β (A β) precursor-like protein 1 0.10 PRTN3 proteinase 3 0.10 ADAMTS10 ADAM metalloproteinase with thrombospondin 0.09 type 1 motif, 10 ILK integrin-linked kinase 0.09 FRAS1 Fraser extracellular matrix complex subunit 1 0.09 ANXA2P2 annexin A2 pseudogene 2 0.08 SMOC2 SPARC related modular calcium binding 2 0.07 ANXA2 annexin A2 0.06 ODAM odontogenic, ameloblast associated 0.06 FREM2 FRAS1 related extracellular matrix protein 2 0.05 HAPLN3 hyaluronan and proteoglycan link protein 3 0.05 GPC3 glypican 3 0.03 LGALS1 lectin, galactoside-binding, soluble, 1 0.02 ADAMTS8 ADAM metalloproteinase with thrombospondin 0.02 type 1 motif, 8 LUM lumican 0.01 HSP90B1 heat shock protein 90 kDa β (Grp94), 0.00 member 1 HAPLN4 hyaluronan and proteoglycan link protein 4 0.00 MATN2 matrilin 2 0.00

(214) Extracellular matrix genes whose expression negatively correlated with Isl1 expression in the HVPs were also identified. Table 7 below lists the extracellular matrix genes that negatively correlate with Isl1 expression with a Pearson's correlation of -0.50 or less.

(215) TABLE-US-00007 TABLE 7 Extracellular matrix genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of -0.50 or less Pearson's Correlation with Isl1 Gene Extracellular

matrix genes (GO: 0031012) Expression FKBP1A FK506 binding protein 1A, 12 kDa -0.51 CLU clusterin -0.52 TFPI2 tissue factor pathway inhibitor 2 -0.52 PLSCR1 phospholipid scramblase 1 -0.53 FBLN5 fibulin 5 -0.53 VWA1 von Willebrand factor A domain containing 1 -0.54 ADAMTS16 ADAM metalloproteinase with thrombospondin -0.55 type 1 motif, 16 MMP25 matrix metalloproteinase 25 -0.55 SFRP2 secreted frizzled-related protein 2 -0.61 SOD1 superoxide dismutase 1, soluble -0.68

Table 8 below lists the extracellular matrix genes that negatively correlate with Isl1 expression with a Pearson's correlation of -0.01 to -0.49.

(216) TABLE-US-00008 TABLE 8 Extracellular matrix genes expressed in HVPs with a Pearson Correlation with Isl1 Expression of -0.01 to -0.49 Pearson's Correlation with Isl1 Gene

Extracellular matrix genes (GO: 0031012) Expression PAPLN papilin, proteoglycan-like sulfated -0.01 glycoprotein SOST sclerostin -0.01 CDON cell adhesion associated, oncogene -0.02 regulated HMCN2 hemicentin 2 -0.02 WNT5A wingless-type MMTV integration site -0.02 family, member 5A PCSK6 proprotein convertase subtilisin/kexin -0.02 type 6 GSTO1 glutathione S-transferase omega 1 -0.02 LTBP1 latent transforming growth factor beta -0.03 binding protein 1 KAZALD1 Kazal-type serine peptidase inhibitor -0.03 domain 1 LTBP2 latent transforming growth factor beta -0.03 binding protein 2 SFRP1 secreted frizzled-related protein 1 -0.04 ADAM11 ADAM metalloproteinase domain 11 -0.05 COL6A1 collagen, type VI, alpha 1 -0.05 COL22A1 collagen, type XXII, alpha 1 -0.05 CYR61 cysteine-rich, angiogenic inducer, 61 -0.05 ELN elastin -0.06 COL9A1 collagen, type IX, alpha 1 -0.06 VTN vitronectin -0.06 COL15A1 collagen, type XV, alpha 1 -0.06 COL4A1 collagen, type IV, alpha 1 -0.07 ANG angiogenin, ribonuclease, RNase A -0.07 family, 5 HSPG2 heparan sulfate proteoglycan 2 -0.09 CRIP2 cysteine-rich protein 2 -0.09 CD151 CD151 molecule (Raph blood group) -0.09 THBS1 thrombospondin 1 -0.09 ADAMTS4 ADAM metalloproteinase with -0.09 thrombospondin type 1 motif, 4 CTGF connective tissue growth factor -0.10 CRISPLD2 cysteine-rich secretory protein LCCL -0.10 domain containing 2 BMP7 bone morphogenetic protein 7 -0.11 COL6A3 collagen, type VI, alpha 3 -0.11 COL3A1 collagen, type III, alpha 1 -0.11 COL14A1 collagen, type XIV, alpha 1 -0.11 MATN3 matrilin 3 -0.11 CPZ carboxypeptidase Z -0.11 BMP1 bone morphogenetic protein 1 -0.11 WISP1 WNT1 inducible signaling pathway -0.12 protein 1 ADAMTS18 ADAM metalloproteinase with -0.12 thrombospondin type 1 motif, 18 COL7A1 collagen, type VII, alpha 1 -0.12 IGFBP7 insulin-like growth factor binding protein 7 -0.12 COCH cochlin -0.13 ADAMTS5 ADAM metalloproteinase with -0.13 thrombospondin type 1 motif, 5 COL11A2 collagen, type XI, alpha 2 -0.13 TGFB1 transforming growth factor, beta-induced, -0.14 68 kDa COL16A1 collagen, type XVI, alpha 1 -0.14 ACHE acetylcholinesterase (Yt blood group) -0.14 THSD4 thrombospondin, type I, domain -0.15 containing 4 DGCR6 DiGeorge syndrome critical region gene 6 -0.15 TGFB111 transforming growth factor beta 1 induced -0.15 transcript 1 ADAMTSL1 ADAMTS-like 1 -0.15 SERPINA1 serpin peptidase inhibitor, clade A (alpha-1 -0.16 antiproteinase, antitrypsin), member 1 MAMDC2 MAM domain containing 2 -0.16 LAMA4 laminin, alpha 4 -0.17 LTBP3 latent transforming growth factor beta -0.17 binding protein 3 EGFL7 EGF-like-domain, multiple 7 -0.17 NPNT nephronectin -0.17 SERPINF1 serpin peptidase inhibitor, clade F (alpha-2 -0.18 antiplasmin, pigment epithelium derived factor), member 1 ABI3BP ABI family, member 3 (NESH) binding -0.18 protein SERPINE2 serpin peptidase inhibitor, clade E (nexin, -0.18 plasminogen activator inhibitor type 1), member 2 WNT6 wingless-type MMTV integration site -0.19 family, member 6 TIMP3 TIMP metalloproteinase inhibitor 3 -0.19 SNCA synuclein, alpha (non A4 component of -0.19 amyloid precursor) PKM pyruvate kinase, muscle -0.19 FGF9 fibroblast growth factor 9 -0.19 VIT vitrin -0.19 WNT1 wingless-type MMTV integration site -0.19 family, member 1 LAMC3 laminin, gamma 3 -0.19 MMP14 matrix metalloproteinase 14 (membrane- -0.19 inserted) PXDN peroxidase -0.19 HNRNPM heterogeneous nuclear ribonucleoprotein M -0.19 FBN1 fibrillin 1 -0.20 ASPN asporin -0.20 ADAMTSL5 ADAMTS-like 5 -0.20 SPON2 spondin 2, extracellular matrix protein -0.20

COL1A2 collagen, type I, alpha 2 -0.20 BGN biglycan -0.20 COL4A2 collagen, type IV, alpha 2 -0.20 ADAMTSL4 ADAMTS-like 4 -0.21 APOH apolipoprotein H (beta-2-glycoprotein I) -0.21 CHI3L1 chitinase 3-like 1 (cartilage glycoprotein- -0.21 39) ADAMTS7 ADAM metalloproteinase with -0.22 thrombospondin type 1 motif, 7 CALR calreticulin -0.22 MMP9 matrix metalloproteinase 9 -0.22 MMP24 matrix metalloproteinase 24 (membrane- -0.22 inserted) SPOCK2 sparc/osteonectin, cwcvc and kazal-like -0.22 domains proteoglycan (testican) 2 COL11A1 collagen, type XI, alpha 1 -0.23 MMP7 matrix metalloproteinase 7 -0.23 MMP16 matrix metalloproteinase 16 (membrane- -0.23 inserted) MFAP2 microfibrillar-associated protein 2 -0.23 POSTN periostin, osteoblast specific factor -0.24 COL8A1 collagen, type VIII, alpha 1 -0.24 WNT2B wingless-type MMTV integration site -0.24 family, member 2B DCN decorin -0.24 EGFL6 EGF-like-domain, multiple 6 -0.24 MMP10 matrix metalloproteinase 10 -0.24 MGP matrix Gla protein -0.24 ECM1 extracellular matrix protein 1 -0.24 SERPINE1 serpin peptidase inhibitor, clade E (nexin, -0.25 plasminogen activator inhibitor type 1), member 1 MMP1 matrix metalloproteinase 1 -0.25 WNT10A wingless-type MMTV integration site -0.25 family, member 10A B4GALT7 xylosylprotein beta 1,4-g alactosyltransferase, polypeptide 7 COL12A1 collagen, type XII, alpha 1 -0.25 LAMA3 laminin, alpha 3 -0.25 LAMA2 laminin, alpha 2 -0.25 LAMB3 laminin, beta 3 -0.25 WNT7B wingless-type MMTV integration site -0.25 family, member 7B FLRT1 fibronectin leucine rich transmembrane -0.25 protein 1 ADAMTS15 ADAM metalloproteinase with -0.26 thrombospondin type 1 motif, 15 COL8A2 collagen, type VIII, alpha 2 -0.26 MFAP1 microfibrillar-associated protein 1 -0.26 TINAGL1 tubulointerstitial nephritis antigen-like 1 -0.26 FGF1 fibroblast growth factor 1 (acidic) -0.27 OLFML2A olfactomedin-like 2A -0.27 CPA6 carboxypeptidase A6 -0.27 COL17A1 collagen, type XVII, alpha 1 -0.27 SPARCL1 SPARC-like 1 (hevin) -0.27 MFAP5 microfibrillar associated protein 5 -0.27 COL4A4 collagen, type IV, alpha 4 -0.28 WNT8B wingless-type MMTV integration site -0.28 family, member 8B ADAMTS19 ADAM metalloproteinase with -0.29 thrombospondin type 1 motif, 19 CRTAP cartilage associated protein -0.29 WNT5B wingless-type MMTV integration site -0.30 family, member 5B WNT3 wingless-type MMTV integration site -0.30 family, member 3 UCMA upper zone of growth plate and cartilage -0.30 matrix associated GPC1 glypican 1 -0.30 TIMP2 TIMP metalloproteinase inhibitor 2 -0.30 ALPL alkaline phosphatase, liver/bone/kidney -0.30 LECT1 leukocyte cell derived chemotaxin 1 -0.32 GPC4 glypican 4 -0.32 SPOCK1 sparc/osteonectin, cwcvc and kazal-like -0.32 domains proteoglycan (testican) 1 HSD17B12 hydroxysteroid (17-beta) dehydrogenase 12 -0.32 LGALS3 lectin, galactoside-binding, soluble, 3 -0.33 EMILIN3 elastin microfibril interfacer 3 -0.34 GFOD2 glucose-fructose oxidoreductase domain -0.34 containing 2 VWC2 von Willebrand factor C domain -0.34 containing 2 SERAC1 serine active site containing 1 -0.34 WNT8A wingless-type MMTV integration site -0.34 family, member 8A LMCD1 LIM and cysteine-rich domains 1 -0.34 CPXM2 carboxypeptidase X (M14 family), -0.34 member 2 ADAMTS14 ADAM metalloproteinase with -0.34 thrombospondin type 1 motif, 14 GPLD1 glycosylphosphatidylinositol specific -0.35 phospholipase D1 FGFBP3 fibroblast growth factor binding protein 3 -0.35 BCAN brevican -0.35 ITGB4 integrin, beta 4 -0.35 LGALS3BP lectin, galactoside-binding, soluble, 3 -0.36 binding protein LPL lipoprotein lipase -0.38 LAD1 laminin 1 -0.39 WNT3A wingless-type MMTV integration site -0.39 family, member 3A TGFR3 transforming growth factor, beta receptor -0.39 III DST dystonin -0.40 WNT10B wingless-type MMTV integration site -0.40 family, member 10B LEFTY2 left-right determination factor 2 -0.41 TNFRSF11B tumor necrosis factor receptor superfamily, -0.41 member 11b WNT9A wingless-type MMTV integration site -0.41 family, member 9A TIMP4 TIMP metalloproteinase inhibitor 4 -0.42 WNT4 wingless-type MMTV integration site -0.42 family, member 4 NCAN neurocan -0.42 ADAMTS20 ADAM metalloproteinase with -0.43 thrombospondin type 1 motif, 20 ITGA6 integrin, alpha 6 -0.43 LOX lysyl oxidase -0.43 THBS4 thrombospondin 4 -0.43 THBS2 thrombospondin 2 -0.44 ADAMTSL2 ADAMTS-like 2 -0.44 ENTPD1 ectonucleoside triphosphate -0.45 diphosphohydrolase 1 RUNX1 runt-related

transcription factor 1 -0.45 VWA2 von Willebrand factor A domain -0.45 containing 2 REL2
 RELT-like 2 -0.46 PTPRZ1 protein tyrosine phosphatase, receptor-type, -0.46 Z polypeptide 1
 LAMC2 laminin, gamma 2 -0.46 DST dystonin -0.40 WNT10B wingless-type MMTV integration
 site -0.40 family, member 10B LEFTY2 left-right determination factor 2 -0.41 TNFRSF11B
 tumor necrosis factor receptor superfamily, -0.41 member 11b WNT9A wingless-type MMTV
 integration site -0.41 family, member 9A TIMP4 TIMP metalloproteinase inhibitor 4 -0.42

Example 13: Gene Expression Profile for Day 6 Islet 1 Negative Cells

(217) In this example, the gene expression profile was determined for Islet 1 negative cells within the Day 6 HVP population to further characterize a subpopulation of cells within the Day 6 population that do not express the necessary markers to qualify as engraftable HVPs. Day 6 HVP populations were generated as described in Examples 1 or 10 and RNA sequencing (RNA-seq) was performed following differentiation as described in Example 2. Cells that were Islet 1 negative (Isl1-) were further analyzed with respect to their gene expression profile. Genes expressed in the Isl1-cells with an average RNA copy number of 2000 or higher are shown below in Table 9.

(218) TABLE-US-00009 TABLE 9 Gene Expression Profile of Day 6 Islet 1 Negative Cells

Gene	Sample #1	Sample #2	Avg. RNA Copy #
ACTB	12288	28126	20207
MTRNR2L2	24511	9774	17142.5
MALAT1	14163	18092	16127.5
EEF1A1	11663	12456	12059.5
KRT8	8884	14087	11485.5
MTRNR2L8	12836	7688	10262
KRT18	5215	10552	7883.5
FN1	4900	10581	7740.5
MTRNR2L1	8550	5719	7134.5
TTN	3149	9601	6375
GAPDH	4907	6391	5649
YWHAZ	5349	5414	5381.5
MTRNR2L9	5673	3850	4761.5
RPL3	3346	5911	4628.5
AHNAK	6727	2197	4462
KCNQ1OT1	5835	3064	4449.5
TUBB	4870	3936	4403
SLC2A3	3311	4301	3806
FTL	3484	4021	3752.5
HSP90B1	4173	2778	3475.5
KRT19	3502	3202	3352
HSPA8	3455	2903	3179
MYL6	1898	4375	3136.5
RPLP0	2319	3922	3120.5
BSG	2519	3593	3056
COL3A1	5312	695	3003.5
TPM1	2938	3059	2998.5
VCAN	2563	3422	2992.5
ENO1	2449	3535	2992
RPL4	2619	3328	2973.5
ACTG1	2687	3253	2970
MTRNR2L10	3409	2487	2948
HMGN2	2684	3153	2918.5
PRTG	2594	2980	2787
TPI1	2418	3113	2765.5
HMGB1	2577	2880	2728.5
VIM	2621	2704	2662.5
ATP5B	3000	2219	2609.5
HSP90AB1	2735	2419	2577
RPL7	2132	2896	2514
CBX5	2799	2219	2509
MYL7	1614	3382	2498
SERPINH1	2547	2327	2437
HNRNPK	2878	1932	2405
SRRM2	2758	2046	2402
PODXL	3683	1112	2397.5
EEF2	2579	2119	2349
SPARC	3026	1645	2335.5
ACTC1	437	4152	2294.5
HUWE1	2583	1977	2280
COL1A2	3544	941	2242.5
LINC00506	2965	1496	2230.5
HSPA5	2078	2356	2217
MDK	2223	2144	2183.5
HNRNPC	2292	2074	2183
HSP90AA1	2220	2138	2179
RGS5	2180	2150	2165
LAMC1	2757	1565	2161
APLNR	868	3246	2057
UGDH-AS1	2633	1457	2045
RPS3A	1601	2399	2000

(219) Accordingly, the data shown in Table 9 provides a gene expression profile for Islet 1 negative, non-engraftable cells within a Day 6 HVP population that are not suitable for transplantation and thus are to be selected against when choosing cells for transplantation and engraftment.

Example 14: Single Cell Sequencing of HVPs

(220) In this example, single cell sequencing was performed on HVPs from day 4, 5, 6, 8, 9 or 15 of differentiation to thereby identify expressed genes in individual cells at different stages of differentiation. Sequencing cell populations in bulk can mask cellular subtypes within that population. Single cell sequencing offers an unbiased approach to classify cell types within a population.

(221) The single cell sequencing approach used herein comprised the three steps: library preparation, sequencing and analysis. For library preparation, a single cell was collected and lysed. The RNA transcriptome was captured, converted to DNA and prepped for sequencing. The captured transcriptome of a cell is termed a “library”. Sequencing was performed using a next-generation sequencer. For analysis, the sequences were read and mapped to genes. This created a transcriptome-wide expression profile for each cell. The expression profiles were then compared with each other. Similar cells were “clustered” together to represent a cell type.

SLC40A1 1.11E-23 0.439358025 SDC4 8.23E-17 0.437190914 IGFBP3 6.90E-07 0.436357742
SERINC3 1.64E-23 0.425115033 GUCY1A3 7.48E-28 0.424214515 CCDC3 1.41E-35
0.422569528 TGFBI 6.52E-16 0.41635348 LAMPS 8.54E-06 0.415693362 COL6A1 2.88E-25
0.407932515 PLS3 8.22E-12 0.406911474 LAPTM4A 1.16E-25 0.405037615 RUNX1T1
7.93E-78 0.403683684 FGF10 2.18E-13 0.401291848 LRIG3 1.89E-30 0.38790471 DSG2
6.23E-28 0.387712513 ARL2BP 7.58E-14 0.38444917 TCEAL9 1.70E-21 0.38384913 PAPSS1
1.40E-14 0.378653665 ANXA6 2.68E-21 0.378214393 LGALS1 2.64E-06 0.378105889 FSTL1
2.78E-24 0.376247551 MOR198 2.78E-24 0.376247551 TMEM98 8.37E-19 0.376170536
SARAF 4.16E-19 0.375950055 TP53111 6.05E-17 0.373187071 CYB5D1 1.63E-23
0.372197782 IFITM3 7.56E-38 0.371288978 LGR5 1.34E-16 0.369873765 WASF1 9.73E-17
0.368716853 ECE1 9.71E-35 0.364724949 MXRA8 2.10E-12 0.363862558 CSRP2 2.60E-17
0.362236698 SDCBP 6.33E-19 0.361772835 SLC16A1 9.31E-11 0.355359606 COL6A2
1.73E-24 0.353551929 EMP2 3.26E-30 0.352860372 LEPROTL1 2.29E-27 0.352310781 CA2
5.04E-08 0.349789326 CCBE1 9.52E-35 0.349748013 SMOC2 1.94E-17 0.347190203 HOXB1
2.81E-17 0.346717419 GATA5 7.66E-10 0.344901285 RBP1 8.58E-21 0.34419898 PHLDA1
3.77E-28 0.343778762 HAND2 2.82E-24 0.343032467 SLC30A3 2.09E-21 0.335148941
SERPINE2 5.79E-22 0.334593289 SYT10 1.62E-39 0.332905271 RHOC 1.16E-16 0.332380866
MPZL1 1.54E-20 0.330849688 LAPTM4B 1.72E-11 0.330782753 LRRC32 1.75E-22
0.327573069 CGNL1 4.43E-31 0.32754955 CXCL12 1.71E-14 0.326028623 NODAL 7.70E-06
0.324782701 DOK4 5.00E-10 0.322776109 ALDH2 1.15E-24 0.322496612 LOC101927497
5.25E-23 0.321216676 RAMP2 3.25E-12 0.320975405 PLSCR5 5.92E-24 0.320022623
COL5A2 4.08E-28 0.319401842 FAM114A1 7.04E-28 0.319200887 METTL7A 1.05E-53
0.319011726 CYYR1 1.65E-18 0.318957318 TOMM34 1.96E-07 0.318041963 CDH11 2.13E-21
0.317968138 FLRT3 8.21E-08 0.316560332 LYPD6 9.56E-23 0.316391668 AFAP1L2 5.04E-32
0.315613744 SERPINB9 5.60E-12 0.315004476 DYNC1I1 1.11E-20 0.313338566 MGARP
1.62E-09 0.311406096 PHGDH 1.17E-09 0.310776152 FHL3 8.79E-08 0.309585103 NPY
2.27E-05 0.309536196 PPP1R15A 2.34E-09 0.309484964 SPIN4 4.33E-24 0.309135856
FXVD6.FXVD2 5.50E-10 0.308833983 P3H2 9.86E-16 0.307905833 FMOD 1.98E-12
0.307428412 FAM213A 8.18E-22 0.305776171 DHDH 3.35E-08 0.303929476 ARSE 3.34E-11
0.302502476 BRINP1 2.18E-16 0.300841535 MYH10 6.61E-15 0.299656764 HSPA5 2.65E-11
0.29901021 ARF4 1.65E-12 0.297788482 CD74 2.28E-11 0.296915796 WNT5A 2.27E-44
0.29686904 IL6ST 1.62E-40 0.296859117 B2M 6.62E-25 0.296077624 MDK 1.00E-32
0.295677643 TIMP1 4.82E-11 0.294947926 PDIA4 1.34E-08 0.294630611 NPPB 0.012639596
0.29392058 RAB34 8.90E-13 0.293594025 TAX1BP3 2.28E-06 0.293246644 SVEP1 1.02E-62
0.291659157 CADM1 4.45E-19 0.29143391 EMP3 2.86E-12 0.291231978 ISL1 9.26E-08
0.289460012 CSNK2A2 1.33E-21 0.288992825 SERPING1 6.85E-12 0.287793225 TECRL
4.21E-05 0.286902439 SLC7A2 2.65E-37 0.286569819 GPR108 1.88E-10 0.284415977
MIR6791 1.88E-10 0.284415977 KDM6A 4.24E-13 0.283535404 RDH10 6.97E-33
0.283087084 PLOD1 3.49E-09 0.280933842 COPG1 5.98E-11 0.279866228 CASP6 6.03E-09
0.279773211 THBS1 8.65E-06 0.279660845 H2AFY2 1.19E-06 0.278216329 GOLGA7
1.10E-10 0.276228148 SELENOF 7.41E-11 0.274011316 FXVD6 3.31E-10 0.273563514
POSTN 2.13E-28 0.273267932 EFEMP2 3.05E-20 0.273241569 TGFB2 2.05E-13 0.273044307
TPM4 1.38E-14 0.272429895 RAB38 0.000686908 0.271918483 NCCRP1 1.79E-11
0.271623914 SH3GLB1 1.51E-18 0.269224524 PMP22 1.57E-07 0.268878282 PLLP 4.96E-35
0.268507068 SCMH1 2.01E-09 0.267191985 MIR503HG 1.86E-07 0.265375187 M1R424
1.86E-07 0.265375187 KDEL2 9.12E-10 0.264599726 DUSP6 0.000293032 0.263830821
SERPINH1 3.02E-17 0.263654992 PTGES 2.95E-05 0.263223829 TMEM120A 4.23E-07
0.263152389 ADAM19 5.67E-22 0.261893827 CRB2 1.47E-12 0.261335033 ADA 2.80E-10
0.260801645 PGF 3.62E-22 0.259993454 NCAM1 1.73E-28 0.259741889 NKX3.1 1.80E-15
0.259560142 COLEC11 1.56E-26 0.259066004 LOC101927230 2.52E-09 0.257688783 RCN3

2.90E-06 0.256437487 ADAMTS12 1.28E-25 0.255820711 SNORD36B 0.01966799
0.255563423 HOXB.AS3 2.31E-58 0.254873973 STAR 9.39E-29 0.254111765 TMED2
8.28E-19 0.253051723 CAPG 2.11E-08 0.252316312 APOC1 1.79E-08 0.25228836 FKBP7
6.16E-14 0.25196672 PTPN13 2.10E-14 0.250842517 EHD2 4.27E-14 0.250458307 Clorf168
2.14E-15 0.250376891

(225) From the list of genes, neuropilin 1 (NRP1) was selected as a candidate for positive sorting of HVPs. An anti-NRP1 antibody compatible for MACS sorting is commercially available.

(226) The day 5 dataset was further examined, since RNA expressed on day 5 would be translated into protein on day 6, the key window where these cells could engraft. The tSNE plot of the day 5 dataset, shown in FIG. 11, reveals two clusters of cells with similar expression patterns, arbitrarily labeled 0 and 1. Cells within a cluster express similar genes. Cells in different clusters express different genes. The distance between each cell on the tSNE plot represents the degree of difference in their gene expression profile. The cluster labeled 1 has high levels of Isl1 expression.

(227) FIG. 12 shows the expression of Isl1 and NRP1 on day 5 cells, wherein dark grey denotes high expression, middle grey denotes low expression and light grey denotes no expression. Thus, FIG. 12 demonstrates that many of the cells with high levels of Isl1 expression also have high levels of NRP1 expression.

(228) This was confirmed by the violin plot of expression levels of Isl1 and NRP1 shown in FIG. 13. Cells within cluster 1 (shown in FIG. 11) have high levels of expression of Isl1. The violin plot of FIG. 13 demonstrates that Isl1+ cells and NRP1+ cells are in the same cluster, since they have a similar gene expression profile. Accordingly, NRP1 is a suitable cell surface marker for positive selection of HVPs (e.g., day 5-6 Isl1+ HVPs).

Example 15: Further Characterization of NRP1+ HVPs

(229) In this example, the expression of NRP1 on a human embryonic stem cell line was examined further. The H9 stem cell line (also known in the art as WA09; Thomson, J. A. et al. (1998) *Science* 282:1145-1147; WiCell Research Institute) was used for these experiments. H9 cells were cultured under cardiac differentiation conditions (e.g., as described herein in Examples 1 and 10) to generate day 6 human ventricular progenitor cells (HVPs).

(230) In a first set of experiments, day 6 HVPs from H9 cells were stained with anti-NRP1, anti-TRA-1-60 or both anti-NRP1 and anti-TRA-160, followed by flow cytometry analysis. The results are shown in FIG. 14A (single staining for NRP1+ cells), FIG. 14B (single staining for TRA-1-60+ cells) and FIG. 14C (double staining for NRP1+TRA-1-60+ cells). The results determined that 92% of the day 6 HVPs were NRP1+ and 17% of the day 6 HVPs were TRA-1-60+. The double staining analysis determined that within the NRP1+ population, 15.5% of those cells are TRA-1-60+.

(231) In a second set of experiments, day 6 HVPs from H9 cells were stained with anti-NRP1, anti-ISL1 or both anti-NRP1 and anti-ISL1, followed by flow cytometry analysis. The results are shown in FIG. 15A (single staining for NRP1+ cells), FIG. 15B (single staining for ISL1+ cells) and FIG. 15C (double staining for NRP1+ ISL1+ cells). The results determined that 78% of the day 6 HVPs were NRP1+, 80% of the day 6 HVPs were ISL1+ and 73% of the day 6 HVPs were NRP1+ ISL1+.

(232) These results confirm the expression of NRP1 on the surface of a large majority of the day 6 HVPs, as well as coexpression with ISL1 and with TRA-1-60, thereby confirming the suitability of NRP1 as a positive selection, ISL1 as a positive co-selection marker, and TRA-1-60 as a negative selection marker.

EQUIVALENTS

(233) Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents of the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

Claims

1. A method for generating human ventricular tissue comprising: subjecting human pluripotent stem cells to Wnt/ β -catenin signaling activation on day 0 of culture, followed by inhibition of Wnt/ β -catenin signaling activation on day 3 to day 5 of culture to obtain a culture comprising NRP1+ human cardiac ventricular progenitor cells; transplanting NRP1+ human cardiac ventricular progenitor cells on day 5-8 of culture into an organ of a non-human animal; and allowing the progenitor cells to grow in vivo such that human ventricular tissue is generated.
 2. The method of claim 1, wherein the non-human animal is an immunodeficient mouse.
 3. The method of claim 1, wherein the organ is a kidney or a heart.
 4. The method of claim 1, wherein the cells are transplanted at a time when one, two, three, four or five of the following cell marker patterns are present: (i) after peak of cardiac mesoderm formation; (ii) at time of peak Islet-1 expression; (iii) before peak of NKX2.5 expression; (iv) before peak expression of downstream genes MEF-2 and TBX-1; and (v) before expression of differentiated contractile protein genes.
 5. The method of claim 1, wherein the cells are transplanted between day 5 and day 7 (inclusive) of in vitro culture of human pluripotent stem cells under conditions to generate human ventricular progenitor cells.
 6. The method of claim 5, wherein the cells are transplanted on day 6 of in vitro culture of human pluripotent stem cells under conditions to generate human ventricular progenitor cells.
-