

US Patent & Trademark Office

Patent Public Search | Text View

United States Patent	12404553
Kind Code	B2
Date of Patent	September 02, 2025
Inventor(s)	Osborn; Terry W. et al.

Biomarkers for bipolar disorder and schizophrenia

Abstract

The present invention provides combinations of biomarkers that can be used in the diagnosis and differentiation of bipolar disorder and schizophrenia. The present invention therefore provides methods of differentiating, diagnosing and treating bipolar disorder and schizophrenia, by examining relevant proteins and RNA in a patient sample.

Inventors: Osborn; Terry W. (The Villages, FL), Vawter; Marquis (Laguna Niguel, CA)

Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA (Oakland, CA);
LAGUNA DIAGNOSTICS, LLC (The Villages, FL)

Family ID: 61619268

Assignee: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA (Oakland, CA);
LAGUNA DIAGNOSTICS, LLC (The Villages, FL)

Appl. No.: 18/333231

Filed: June 12, 2023

Prior Publication Data

Document Identifier	Publication Date
US 20230383354 A1	Nov. 30, 2023

Related U.S. Application Data

continuation parent-doc US 16333626 US 11713486 WO PCT/US2017/051716 20170915 child-
doc US 18333231
us-provisional-application US 62395159 20160915

Publication Classification

Int. Cl.: C12Q1/6883 (20180101); **G01N33/68** (20060101)

U.S. Cl.:

CPC **C12Q1/6883** (20130101); **G01N33/6896** (20130101); C12Q2600/158 (20130101);
G01N2800/302 (20130101); G01N2800/304 (20130101); G01N2800/60 (20130101)

Field of Classification Search

CPC: C12Q (1/6883); C12Q (2600/158); G01N (33/6896); G01N (2800/302); G01N
(2800/304); G01N (2800/60); G01N (33/50)

References Cited

U.S. PATENT DOCUMENTS

Patent No.	Issued Date	Patentee Name	U.S. Cl.	CPC
5143854	12/1991	Pirrung et al.	N/A	N/A
5384261	12/1994	Winkler et al.	N/A	N/A
5424186	12/1994	Fodor et al.	N/A	N/A
5445934	12/1994	Fodor et al.	N/A	N/A
5677195	12/1996	Winkler et al.	N/A	N/A
5708153	12/1997	Dower et al.	N/A	N/A
5744305	12/1997	Fodor et al.	N/A	N/A
5770358	12/1997	Dower et al.	N/A	N/A
5789162	12/1997	Dower et al.	N/A	N/A
5800992	12/1997	Fodor et al.	N/A	N/A
5856174	12/1998	Lipshutz et al.	N/A	N/A
5922591	12/1998	Anderson et al.	N/A	N/A
6040193	12/1999	Winkler et al.	N/A	N/A
2008/0199866	12/2007	Akil et al.	N/A	N/A
2014/0141986	12/2013	Spetzler et al.	N/A	N/A

OTHER PUBLICATIONS

Mamdani, F., et al. 2015. Variable telomere length across post-mortem human brain regions and specific reduction in the hippocampus of major depressive disorder, *Transl Psychiatry*, 5: e636. cited by applicant

Martin, M. V., et al. 2009. Exon expression in lymphoblastoid cell lines from subjects with schizophrenia before and after glucose deprivation, *BMC Med Genomics*, 2: 62. cited by applicant

Middleton, F. A., et al. 2005. Gene expression analysis of peripheral blood leukocytes from discordant sib-pairs with schizophrenia and bipolar disorder reveals points of convergence between genetic and functional genomic approaches, *Am J Med Genet B Neuropsychiatr Genet*, 136B: 12-25. cited by applicant

Morgan, L. Z., et al. 2016. Quantitative Trait Locus and Brain Expression of HLA-DPA1 Offers Evidence of Shared Immune Alterations in Psychiatric Disorders, *Microarrays (Basel)*, 5. cited by applicant

Munkholm, K., et al. 2014. Reduced mRNA expression of PTGDS in peripheral blood mononuclear cells of rapid-cycling bipolar disorder patients compared with healthy control subjects, *Int J Neuropsychopharmacol*, 18. cited by applicant

Naydenov, A. V., et al. 2007. Differences in lymphocyte electron transport gene expression levels

between subjects with bipolar disorder and normal controls in response to glucose deprivation stress, *Arch Gen Psychiatry*, 64: 555-64. cited by applicant

Nishimura, Y., et al. 2007. Genome-wide expression profiling of lymphoblastoid cell lines distinguishes different forms of autism and reveals shared pathways, *Hum Mol Genet*, 16: 1682-98. cited by applicant

Perl, O., et al. 2006. Low levels of alpha7-nicotinic acetylcholine receptor mRNA on peripheral blood lymphocytes in schizophrenia and its association with illness severity, *Neuropsychobiology*, 53: 88-93. cited by applicant

Philibert, R. A., et al. 2007. Transcriptional profiling of lymphoblast lines from subjects with panic disorder, *Am J Med Genet B Neuropsychiatr Genet*, 144B: 674-82. cited by applicant

Philibert, R. A., et al. 2007. Transcriptional profiling of subjects from the Iowa adoption studies, *Am J Med Genet B Neuropsychiatr Genet*, 144B: 683-90. cited by applicant

Pradervand, S., et al. 2008. Affymetrix Whole-Transcript Human Gene 1.0 ST array is highly concordant with standard 3' expression arrays, *Biotechniques*, 44: 759-62. cited by applicant

Ramos, E. I., et al. 2014. Genetic variation in MKL2 and decreased downstream PCTAIRE1 expression in extreme, fatal primary human microcephaly, *Clin Genet*, 85: 423-32. cited by applicant

Richard, H., et al. 2010. Prediction of alternative isoforms from exon expression levels in RNA-Seq experiments, *Nucleic Acids Res*, 38: e112. cited by applicant

Rollins, B., et al. 2010. Analysis of whole genome biomarker expression in blood and brain, *Am J Med Genet B Neuropsychiatr Genet*, 153B: 919-36. cited by applicant

Ruderfer, D. M., et al. 2014. Polygenic dissection of diagnosis and clinical dimensions of bipolar disorder and schizophrenia, *Mol Psychiatry*, 19: 1017-24. cited by applicant

Sanders, A. R., et al. 2013. Transcriptome study of differential expression in schizophrenia, *Hum Mol Genet*, 22: 5001-14. cited by applicant

Scarr, E., et al. 2015. Biomarkers for Psychiatry: The Journey from Fantasy to Fact, a Report of the 2013 CINP Think Tank, *Int J Neuropsychopharmacol*, 18: pyv042. cited by applicant

Schizophrenia Working Group of the Psychiatric Genomics Consortium. 2014. Biological insights from 108 schizophrenia-associated genetic loci, *Nature*, 511: 421-7. cited by applicant

Schoenbaum, M., et al. 2017. Twelve-Month Health Care Use and Mortality in Commercially Insured Young People With Incident Psychosis in the United States, *Schizophr Bull*, 43: 1262-72. cited by applicant

Segman, R. H., et al. 2005. Peripheral blood mononuclear cell gene expression profiles identify emergent post-traumatic stress disorder among trauma survivors, *Mol Psychiatry*, 10: 500-13, 425. cited by applicant

Sekar, A., et al. 2016. Schizophrenia risk from complex variation of complement component 4, *Nature*, 530: 177-83. cited by applicant

Sequeira, A., et al. 2008. Coding SNPs included in exon arrays for the study of psychiatric disorders, *Mol Psychiatry*, 13: 363-5. cited by applicant

Shen, S., et al. 2010. MADS+: discovery of differential splicing events from Affymetrix exon junction array data, *Bioinformatics*, 26: 268-9. cited by applicant

Tang, Y., et al. 2005. Blood gene expression profiling of neurologic diseases: a pilot microarray study, *Arch Neurol*, 62: 210-5. cited by applicant

Tomasik, J., et al. 2012. Blood test for schizophrenia, *Eur Arch Psychiatry Clin Neurosci*, 262 Suppl 2: S79-83. cited by applicant

Vawter, M. P., et al. 2007. Dysregulation of X-linked gene expression in Klinefelter's syndrome and association with verbal cognition, *Am J Med Genet B Neuropsychiatr Genet*, 144B: 728-34. cited by applicant

Vawter, M. P., et al. 2011. An integrative functional genomics approach for discovering biomarkers in schizophrenia, *Brief Funct Genomics*, 10: 387-99. cited by applicant

Weber, M. D., et al. 2017. Repeated Social Defeat, Neuroinflammation, and Behavior: Monocytes Carry the Signal, *Neuropsychopharmacology*, 42: 46-61. cited by applicant

Wu, J. Q., et al. 2016. Altered neural signaling and immune pathways in peripheral blood mononuclear cells of schizophrenia patients with cognitive impairment: A transcriptome analysis, *Brain Behav Immun*, 53: 194-206. cited by applicant

Xing, Y., et al. 2006. Probe selection and expression index computation of Affymetrix Exon Arrays, *PLoS One*, 1: e88. cited by applicant

Xing, Y., et al. 2007. Assessing the conservation of mammalian gene expression using high-density exon arrays, *Mol Biol Evol*, 24: 1283-5. cited by applicant

Xing, Y., et al. 2008. MADS: a new and improved method for analysis of differential alternative splicing by exon-tiling microarrays, *RNA*, 14: 1470-9. cited by applicant

Xu, W., et al. 2011. Human transcriptome array for high-throughput clinical studies, *Proc Natl Acad Sci U S A*, 108: 3707-12. cited by applicant

Xu, Y., et al. 2016. Altered expression of mRNA profiles in blood of early-onset schizophrenia, *Sci Rep*, 6: 16767. cited by applicant

Yao, Y., et al. 2008. Verification of proposed peripheral biomarkers in mononuclear cells of individuals with schizophrenia, *J Psychiatr Res*, 42: 639-43. cited by applicant

Vawter, M.P et al., Exon Array Biomarkers for the Differential Diagnosis of Schizophrenia and Bipolar Disorder, *Molecular Neuropsychiatry* 2017; 3:197-213, Apr. 10, 2018. cited by applicant

Cohen, O.S. et al., Transcriptomic Analysis of Postmortem Brain identifies Dysregulated Splicing Events in Novel Candidate Genes for Schizophrenia, National Institutes of Health, *Schizophr Res*. Dec. 2012; 142(1-3); 188-199, 30 pages. cited by applicant

Citrome, L., "A systematic review of meta-analyses of the efficacy of oral atypical antipsychotics for the treatment of adult patients with schizophrenia", *Expert Opinion on Pharmacotherapy*, vol. 13, Issue 11, Aug. 2012, 1545-7153. cited by applicant

Ogawa et al., "Mood Stabilizers and Antipsychotics for Acute Mania: A Systematic Review and Meta-Analysis of Combination/Augmentation Therapy Versus Monotherapy", *International Journal of Neuropsychopharmacology*, vol. 25, Issue 10, Oct. 2022, 839-852. cited by applicant

Enard et al. (*Science*. Apr. 12, 2002.; 296(5566):340-43). cited by applicant

Zhang (*European Archives of Psychiatry and Clinical Neuroscience*, 2002, 272:591-602). cited by applicant

Cheung (*Cold Spring Harbor Symposia on Quant Bio.*, 2003, vol. LXVIII, pp. 403-407). cited by applicant

Cohen et al. *Schizophrenia Research*, 2012, vol. 142: 188-199. cited by applicant

Agarwal, A., et al. 2010. Comparison and calibration of transcriptome data from RNA-Seq and tiling arrays, *BMC Genomics*, 11: 383. cited by applicant

Anand, A., et al. 2016. Effects of Lithium Monotherapy for Bipolar Disorder on Gene Expression in Peripheral Lymphocytes, *Mol Neuropsychiatry*, 2: 115-23. cited by applicant

Begemann, M., et al. 2008. Episode-specific differential gene expression of peripheral blood mononuclear cells in rapid cycling supports novel treatment approaches, *Mol Med*, 14: 546-52. cited by applicant

Bemmo, A., et al. 2008. Gene expression and isoform variation analysis using Affymetrix Exon Arrays, *BMC Genomics*, 9: 529. cited by applicant

Benovoy, D., et al. 2008. Effect of polymorphisms within probe-target sequences on oligonucleotide microarray experiments, *Nucleic Acids Res*, 36: 4417-23. cited by applicant

Bomprezzi, R., et al. 2003. Gene expression profile in multiple sclerosis patients and healthy controls: identifying pathways relevant to disease, *Hum Mol Genet*, 12: 2191-9. cited by applicant

Borovecki, F., et al. 2005. Genome-wide expression profiling of human blood reveals biomarkers for Huntington's disease, *Proc Natl Acad Sci U S A*, 102: 11023-8. cited by applicant

Bowden, N. A., et al. 2006. Preliminary investigation of gene expression profiles in peripheral

blood lymphocytes in schizophrenia, *Schizophr Res*, 82: 175-83. cited by applicant

Bradford, J. R., et al. 2010. A comparison of massively parallel nucleotide sequencing with oligonucleotide microarrays for global transcription profiling, *BMC Genomics*, 11: 282. cited by applicant

Brea, D., et al. 2009. Inflammatory and neuroimmunomodulatory changes in acute cerebral ischemia, *Cerebrovasc Dis*, 27 Suppl 1: 48-64. cited by applicant

Chan, M. K., et al. 2014. Applications of blood-based protein biomarker strategies in the study of psychiatric disorders, *Prog Neurobiol*, 122: 45-72. cited by applicant

Chan, M. K., et al. 2015. Development of a blood-based molecular biomarker test for identification of schizophrenia before disease onset, *Transl Psychiatry*, 5: e601. cited by applicant

Cloutier, M., et al. 2016. The Economic Burden of Schizophrenia in the United States in 2013, *J Clin Psychiatry*, 77: 764-71. cited by applicant

Cole, S. W., et al. 2007. Social regulation of gene expression in human leukocytes, *Genome Biol*, 8: R189. cited by applicant

De Jong, S., et al. 2012. A gene co-expression network in whole blood of schizophrenia patients is independent of antipsychotic-use and enriched for brain-expressed genes, *PLoS One*, 7: e39498. cited by applicant

Dilsaver, S. C. 2011. An estimate of the minimum economic burden of bipolar I and II disorders in the United States: 2009, *J Affect Disord*, 129: 79-83. cited by applicant

Downes, C. E., et al. 2010. Neural injury following stroke: are Toll-like receptors the link between the immune system and the CNS?, *Br J Pharmacol*, 160: 1872-88. cited by applicant

Duan, S., et al. 2008. Genetic architecture of transcript-level variation in humans, *Am J Hum Genet*, 82: 1101-13. cited by applicant

Duan, S., et al. 2009. Expression and alternative splicing of folate pathway genes in HapMap lymphoblastoid cell lines, *Pharmacogenomics*, 10: 549-63. cited by applicant

Duan, S., et al. 2008. SNPInProbe_1.0: a database for filtering out probes in the Affymetrix GeneChip human exon 1.0 ST array potentially affected by SNPs, *Bioinformatics*, 2: 469-70. cited by applicant

Evans, S. J., et al. 2012. Fats and factors: lipid profiles associate with personality factors and suicidal history in bipolar subjects, *PLoS One*, 7: e29297. cited by applicant

Fodor, S. P., et al. 1991. Light-directed, spatially addressable parallel chemical synthesis, *Science*, 251: 767-73. cited by applicant

Fraser, H. B., et al. 2009. Common polymorphic transcript variation in human disease, *Genome Res*, 19: 567-75. cited by applicant

Fromer, M., et al. 2016. Gene expression elucidates functional impact of polygenic risk for schizophrenia, *Nat Neurosci*, 19: 1442-53. cited by applicant

Gamazon, E. R., et al. 2010. Comprehensive survey of SNPs in the Affymetrix exon array using the 1000 Genomes dataset, *PLoS One*, 5: e9366. cited by applicant

Gamazon, E. R., et al. 2010. SCAN: SNP and copy number annotation, *Bioinformatics*, 26: 259-62. cited by applicant

Gelderblom, M., et al. 2009. Temporal and spatial dynamics of cerebral immune cell accumulation in stroke, *Stroke*, 40: 1849-57. cited by applicant

Giannone, S., et al. 2004. Gene expression profile analysis in human T lymphocytes from patients with Down Syndrome, *Ann Hum Genet*, 68: 546-54. cited by applicant

Glatt, S. J., et al. 2009. Alternatively Spliced Genes as Biomarkers for Schizophrenia, Bipolar Disorder and Psychosis: A Blood-Based Spliceome-Profiling Exploratory Study, *Curr Pharmacogenomics Person Med*, 7: 164-88. cited by applicant

Glatt, S. J., et al. 2005. Comparative gene expression analysis of blood and brain provides concurrent validation of SELENBP1 up-regulation in schizophrenia, *Proc Natl Acad Sci U S A*, 102: 15533-8. cited by applicant

Glatt, S. J., et al. 2011. Similarities and differences in peripheral blood gene-expression signatures of individuals with schizophrenia and their first-degree biological relatives, *Am J Med Genet B Neuropsychiatr Genet*, 156B: 869-87. cited by applicant

Haenisch, F., et al. 2016. Towards a blood-based diagnostic panel for bipolar disorder, *Brain Behav Immun*, 52: 49-57. cited by applicant

Hawrylycz, M. J., et al. 2012. An anatomically comprehensive atlas of the adult human brain transcriptome, *Nature*, 489: 391-99. cited by applicant

Heinzen, E. L., et al. 2008. Tissue-specific genetic control of splicing: implications for the study of complex traits, *PLoS Biol*, 6: e1. cited by applicant

Hess, J. L., et al. 2016. Transcriptome-wide mega-analyses reveal joint dysregulation of immunologic genes and transcription regulators in brain and blood in schizophrenia, *Schizophr Res*, 176: 114-24. cited by applicant

Horvath, S., et al. 2014. Immune system disturbances in schizophrenia, *Biol Psychiatry*, 75: 316-23. cited by applicant

Hou, L., et al. 2016. Genome-wide association study of 40,000 individuals identifies two novel loci associated with bipolar disorder, *Hum Mol Genet*, 25: 3383-94. cited by applicant

International Schizophrenia Consortium, et al. 2009. Common polygenic variation contributes to risk of schizophrenia and bipolar disorder, *Nature*, 460: 748-52. cited by applicant

Kapur, K., et al. 2008. Cross-hybridization modeling on Affymetrix exon arrays, *Bioinformatics*, 24: 2887-93. cited by applicant

Kapur, K., et al. 2007. Exon arrays provide accurate assessments of gene expression, *Genome Biol*, 8: R82. cited by applicant

Kumarasinghe, N., et al. 2013. Gene expression profiling in treatment-naïve schizophrenia patients identifies abnormalities in biological pathways involving AKT1 that are corrected by antipsychotic medication, *Int J Neuropsychopharmacol*, 16: 1483-503. cited by applicant

Kwan, T., et al. 2008. Genome-wide analysis of transcript isoform variation in humans, *Nat Genet*, 40: 225-31. cited by applicant

Le-Niculescu, H., et al. 2011. Convergent functional genomics of anxiety disorders: translational identification of genes, biomarkers, pathways and mechanisms, *Transl Psychiatry*, 1: e9. cited by applicant

Liew, C. C., et al. 2006. The peripheral blood transcriptome dynamically reflects system wide biology: a potential diagnostic tool, *J Lab Clin Med*, 147: 126-32. cited by applicant

Lin, L., et al. 2009. Using high-density exon arrays to profile gene expression in closely related species, *Nucleic Acids Res*, 37: e90. cited by applicant

Liu, S., et al. 2011. A comparison of RNA-Seq and high-density exon array for detecting differential gene expression between closely related species, *Nucleic Acids Res*, 39: 578-88. cited by applicant

Liu, Y., et al. 2009. Elevated delta-6 desaturase (FADS2) expression in the postmortem prefrontal cortex of schizophrenic patients: relationship with fatty acid composition, *Schizophr Res*, 109: 113-20. cited by applicant

Liu, Y., et al. 2011. Elevated Delta-6 desaturase (FADS2) gene expression in the prefrontal cortex of patients with bipolar disorder, *J Psychiatr Res*, 45: 269-72. cited by applicant

Maes, O. C., et al. 2007. Transcriptional profiling of Alzheimer blood mononuclear cells by microarray, *Neurobiol Aging*, 28: 1795-809. cited by applicant

Mamdani, F., et al. 2013. Coding and noncoding gene expression biomarkers in mood disorders and schizophrenia, *Dis Markers*, 35: 11-21. cited by applicant

Geddes et al. 2013, Comparative efficacy and tolerability of 15 antipsychotic drugs in schizophrenia: a multiple-treatments meta-analysis, *Lancet* 2013; 382: 951-62. cited by applicant

Perlis et al. 2006, Atypical Antipsychotics in the treatment of mania: a meta-analysis of randomized, placebo-controlled trials, *J Clin Psychiatry* 2006; 67:509-516, 509. cited by applicant

Background/Summary

CROSS-REFERENCE TO RELATED APPLICATIONS (1) This application claims the benefit of U.S. application Ser. No. 16/333,626, filed Mar. 15, 2019, which is a U.S. national phase application filed under 35 U.S.C. § 371 claiming priority to International Patent Application No. PCT/US17/51716, filed Sep. 15, 2017, which claims priority to U.S. Provisional Application No. 63/395,159, filed Sep. 15, 2016, the contents of each of which are incorporated by reference herein in their entireties.

BACKGROUND OF THE INVENTION

(1) Schizophrenia and bipolar disorder are chronic, severe and disabling brain disorders that affect about 1 and 2 percent of age 18 and older U.S. population, respectively. Despite moderately effective treatments, such as antipsychotic medications and psychosocial interventions, people with schizophrenia (SZ) and bipolar disorder (BD) often do not receive timely treatment because of misdiagnosis until the disease is already well-established with recurrent episodes of psychosis and mood dysregulation. These episodes result in costly multiple hospitalizations and disabilities that can last for decades. Ideally, successful diagnostic tests could address the significant clinical problem of early identification and enable more timely initiation of treatments.

(2) Over 2,000,000 individuals are clinically diagnosed as suffering with schizophrenia (SZ) in the U.S. Over 100,000 adolescent Americans suffer from an initial episode of psychosis each year. Currently, no 'objective' clinical laboratory test exists to accurately diagnose their disease, and there are no FDA approved biomarkers for psychotic disorders such as SZ or mood disorders associated with psychosis such as bipolar disorder (BD). Physicians cannot use brain biopsies of living patients for diagnosis of neuropsychiatric disorders. Instead, physicians rely upon clinical observation and the patient's history of reported symptoms. Consequently, if physicians misdiagnose similarly presenting diseases like SZ and BD, there can be a lag in treatment and increase in the suicide rate. Following an initial episode of psychosis among individuals aged 16-30, there is a 24-fold increase in the risk of death in the following year (Schoenbaum, Twelve-Month Health Care Use and Mortality in Commercially Insured Young People With Incident Psychosis in the United States. Schizophrenia Bulletin 2017). This study points towards a lack of treatment (61% did not receive any antipsychotic medication) after initial presentation with psychosis and even higher rates in those dying within 12 months of an initial episode of psychosis (Schoenbaum, Twelve-Month Health Care Use and Mortality in Commercially Insured Young People With Incident Psychosis in the United States. Schizophrenia Bulletin 2017). Through clinical observations, these diseases take months or even years to diagnose definitively and to appropriately prescribe disease-matched medications for effective treatment. The mental health field could benefit greatly from commercial blood-based biomarker tests that discriminate between patients without a psychiatric disorder and those with SZ or BD.

(3) A growing body of work has demonstrated the potential utility of RNA diagnostic tools with peripheral samples such as whole blood, peripheral blood mononuclear cells, and lymphoblastic cell lines in multiple studies of SZ and BD (Begemann et al., Mol Med 2008; 14(9-10): 546-552; Bowden et al., Schizophr Res 2006; 82(2-3): 175-183; de Jong S et al., PLoS One 2012; 7(6): e39498; Glatt et al., Proc Natl Acad Sci USA 2005; 102(43): 15533-15538; Middleton et al., Am J Med Genet B Neuropsychiatr Genet 2005; 136B(1): 12-25; Naydenov et al., Arch Gen Psychiatry

2007; 64(5): 555-564; Merl et al., Neuropsychobiology 2006; 53(2): 88-93; Sanders et al., Hum Mol Genet 2013; 22(24): 5001-5014; Yao et al., J Psychiatr Res 2008; 42(8): 639-643). There have also been large studies that have used whole genome RNA expression to compare healthy controls and disorders such as Alzheimer's disease (Maes et al., Neurobiol Aging 2007; 28(12): 1795-1809), autism (Nishimura et al., Hum Mol Genet 2007; 16(14): 1682-1698), Down's Syndrome (Giannone et al., Ann Hum Genet 2004; 68(Pt 6): 546-554), epilepsy (Tang et al., Arch Neurol 2005; 62(2): 210-215), Tourette's Syndrome (Tang et al., Arch Neurol 2005; 62(2): 210-215), Huntington's Disease (Borovecki et al., Proc Natl Acad Sci USA 2005; 102(31): 11023-11028), Klinefelter's Syndrome (KS) (Vawter et al., Am J Med Genet B Neuropsychiatr Genet 2007; 144B(6): 728-734), multiple sclerosis (Bomprezzi et al., Hum Mol Genet 2003; 12(17): 2191-2199), smoking and major depression (Philibert et al., Am J Med Genet B Neuropsychiatr Genet 2007; 144B(5): 683-690), panic disorder (Philibert et al., Am J Med Genet B Neuropsychiatr Genet 2007; 144B(5): 674-682), post-traumatic stress disorder (Segman et al., Mol Psychiatry 2005; 10(5): 500-513, 425), and subjective social isolation (loneliness) (Cole et al., Genome Biol 2007; 8(9): R189).

(4) A tremendous effort has been expended into GWAS of schizophrenia (Consortium. Nature 2014; 511(7510): 421-427) and bipolar disorder (Hou et al., Hum Mol Genet 2016; 25(15): 3383-3394), however, there is a lack of consensus regarding the specific genes that cause schizophrenia or bipolar disorder; with shared genetic factors across these disorders (Iftuderfer et al., Mol Psychiatry 2014; 19(9): 1017-1024). More importantly, which combinations of interacting genes that actually cause each illness as opposed to polygenic susceptibilities for psychiatric endophenotypes are unknown. Estimates of several hundred genes of small effect size were published from the largest international genetic study of SZ (Purcell et al., Nature 2009; 460(7256): 748-752) to the possibility that thousands of genes are involved in the pathogenesis of schizophrenia (Profler et al., Nat Neurosci 2016; 19(11): 1442-1453). Dysregulation of miRNA could potentially help to define sets of genes relevant to pathophysiology, treatment, or secondary to these causes.

(5) Thus, there is an urgent need in the art for compositions and methods for objectively diagnosing SZ and BD, to reduce duration of untreated psychosis by earlier detection to help establish rapid and informative patient decisions. The present invention addresses these needs.

SUMMARY

(6) In one embodiment, the invention relates to a method of diagnosing schizophrenia (SZ) or bipolar disorder (BD) in a subject, the method comprising: a) determining the expression level of at least two biomarker genes selected from the group consisting of SH3Y1,1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5 in a sample of the subject, b) determining the probability of the sample being from a subject with or without SZ or BD, and c) diagnosing the subject as having SZ or BD on the basis of the determined result from the sample as compared to a pre-determined cut-off.

(7) In one embodiment, the method comprises evaluating the expression levels of at least two of TCEA3, SLC44A5, IL5RA, GYLTL1 B and DDX5, determining the probability of the sample being from a subject with schizophrenia, and diagnosing the subject with SZ when the probability of the sample being from a subject with schizophrenia is greater than 0.499.

(8) In one embodiment, the method comprises evaluating the expression levels of at least two of HPR, TREML4, PTGDS, CPA3, TRIM4 and SLC44A5, determining the probability of the sample being from a subject with schizophrenia, and diagnosing the subject with SZ when the probability of the sample being from a subject with schizophrenia is greater than 0.549.

(9) In one embodiment, the method comprises evaluating the expression levels of at least two of SLC44A5, CPA3, TREML4, TRIM4, PTGDS and SH3YL1, determining the probability of the sample being from a subject with schizophrenia, and diagnosing the subject with SZ when the probability of the sample being from a subject with schizophrenia is greater than or equal to 0.411.

(10) In one embodiment, the method comprises evaluating the expression levels of at least two of PTGDS, HLA-DRB5, OXTR and FADS2, determining the probability of the sample being from a healthy subject, and diagnosing the subject with BD when the probability of the sample being from a healthy subject is less than or equal to 0.659.

(11) In one embodiment, the method comprises evaluating the expression levels of at least two of CRIP2, CPA3, OXTR, TRIM4, PTGDS and SH3YL1, determining the probability of the sample being from a subject with BD, and diagnosing the subject with BD when the probability of the sample being from a subject with BD is greater than or equal to 0.452.

(12) In one embodiment, the method comprises evaluating the expression levels of at least two of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8, determining the probability of the sample being from a healthy subject, and diagnosing the subject with SZ or BD when the probability of the sample being from a healthy subject is less than or equal to 0.1518. In one embodiment, the method further comprises evaluating the expression levels of at least two of CRIP2, OXTR and FADS2 in the sample from the subject, wherein the probability of the sample being from a healthy subject was determined as less than or equal to 0.1518, determining the probability of the sample being from a BD subject, diagnosing the subject with SZ when the probability of the sample being from a BD subject is less than or equal to 0.2857, and diagnosing the subject with RD when the probability of the sample being from a BD subject is greater than 0.2857. In one embodiment, the method further comprises evaluating the expression levels of at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5 in the sample from the subject, wherein the probability of the sample being from a healthy subject was determined as greater than 0.1518, determining the probability of the sample being from a subject with schizophrenia, and diagnosing the subject with SZ when the probability of the sample being from a subject with schizophrenia is greater than 0.499. In one embodiment, the method further comprises evaluating the expression levels of at least two of PTGDS, HLA-DRB5, OXTR and FADS2 in the sample from the subject, wherein the probability of the sample being from a healthy subject was determined as greater than 0.1518, determining the probability of the sample being from a healthy subject, and diagnosing the subject with BD when the probability of the sample being from a healthy subject is less than or equal to 0.659.

(13) In one embodiment, the method comprises evaluating the expression levels of at least two of SLC44A5, CPA3, CRIP2, TRIM4, PTGDS and SH3YL1, determining the probability of the sample being from a subject having SZ or RD, and diagnosing the subject with SZ or BD when the probability of the sample being from a subject having SZ or BD is greater than or equal to 0.466. In one embodiment, the method further comprises evaluating the expression levels of at least two of SH3YL1, OXTR, PTGDS, CPA3, TBC1D1, and TCEA3, determining the probability of the sample being from a subject with SZ, diagnosing the subject with SZ when the probability of the sample being from a subject with SZ is greater than or equal to 0.584, and diagnosing the subject with BD when the probability of the sample being from a subject with SZ is less than 0.584.

(14) In one embodiment, the method comprises evaluating the expression levels of at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5, determining the probability of the sample being from a healthy subject, and diagnosing the subject with SZ when the probability of the sample being from a healthy subject is less than or equal to 0.3323. In one embodiment, the expression level of at least two biomarker genes is determined from data generated from the Nanostring platform.

(15) In one embodiment, the method further comprises treating the subject for the diagnosed SZ or BD.

(16) In one embodiment, the expression level of at least two biomarker genes is determined from data generated from a platform selected from Affymetrix exon array and Nanostring.

(17) In one embodiment, the invention relates to a method of identifying a subject as belonging to the normal population with respect to BD or SZ, the method comprising: a) determining the

expression level of at least two biomarker genes selected from the group consisting of SH3YL1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5 in a sample of the subject, b) determining the probability of the sample being from a subject with or without SZ or BD, and c) identifying the subject as belonging to the normal population on the basis of the determined result from the sample as compared to a pre-determined cut-off.

(18) In one embodiment, the method comprises evaluating the expression levels of at least two of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8, determining the probability of the sample being from a healthy subject, and identifying the subject as being from the normal population with respect to BD and SZ when the probability of the sample being from a healthy subject is greater than 0.1518. In one embodiment, the method further comprises evaluating the expression levels of at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5 in the sample from the subject, wherein the probability of the sample being from a healthy subject was determined as greater than 0.1518, determining the probability of the sample being from a subject with schizophrenia, and identifying the subject as being from the normal population with regard to SZ when the probability of the sample being from a subject with schizophrenia is less than or equal to 0.499. In one embodiment, the method further comprises evaluating the expression levels of at least two of PTGDS, HLA-DRB5, OXTR and FADS2 in the sample from the subject, wherein the probability of the sample being from a healthy subject was determined as greater than 0.1518, determining the probability of the sample being from a healthy subject, and identifying the subject as being from the normal population with regard to BD when the probability of the sample being from a healthy subject is greater than 0.659.

(19) In one embodiment, the expression level of at least two biomarker genes is determined from data generated from a platform selected from Affymetrix exon array and Nanostring.

(20) In one embodiment, the method comprises evaluating the expression levels of at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5, determining the probability of the sample being from a healthy subject, and identifying the subject as being from the normal population with respect to SZ when the probability of the sample being from a healthy subject is greater than 0.3323. In one embodiment, the expression level of at least two biomarker genes is determined from data generated from the Nanostring platform.

(21) In one embodiment, the invention relates to a method of differentially diagnosing a subject in need thereof as having a disorder selected from the group consisting of SZ and BD, the method comprising: a) determining the expression level of at least two biomarker genes selected from the group consisting of SH3YL1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5 in a sample of the subject; b) determining the probability of the sample being from a subject having a disorder selected from the group consisting of SZ and BD; and c) differentially diagnosing the subject as having a disorder selected from the group consisting of SZ and BD on the basis of the determined result from the sample as compared to a pre-determined cut-off.

(22) In one embodiment, the method comprises evaluating the expression levels of at least two of CRTP2, OXTR and FADS2 in the sample from the subject, determining the probability of the sample being from a BD subject, diagnosing the subject with SZ when the probability of the sample being from a BD subject is less than or equal to 0.2857, and diagnosing the subject with BD when the probability of the sample being from a BD subject is greater than 0.2857.

(23) In one embodiment, the subject has a prior diagnosis of a disorder selected from the group consisting of SZ and BD.

(24) In one embodiment, the method further comprises treating the subject for the diagnosed SZ or BD.

(25) In one embodiment, the expression level of at least two biomarker genes is determined from data generated from a platform selected from Affymetrix exon array and Nanostring.

Description

BRIEF DESCRIPTION OF THE DRAWINGS

(1) The following detailed description of embodiments of the invention will be better understood when read in conjunction with the appended drawings. For the purpose of illustrating the invention, drawings are included. It should be understood, however, that the invention is not limited to the precise arrangements and instrumentalities of the embodiments shown in the drawings.

(2) FIG. 1 depicts a schematic of a clinical study design to evaluate potential RNA gene biomarkers.

(3) FIG. 2 depicts the results of exemplary experiments demonstrating the fold change ratio (SZ compared to normal control (NC)) for exon microarray results as compared to fold change ratio (SZ compared to NC) of qPCR.

(4) FIG. 3 depicts the results of exemplary experiments demonstrating the REL transcription factor is a central hub for genes in the top 122 differentially expressed list. Left panel shows relationship for BD compared to controls, right panel shows relationship for SZ compared to controls. Many of the genes are oppositely regulated across BD and SZ.

(5) FIG. 4 depicts the results of exemplary experiments demonstrating the MKL2 proto-oncogene is a central hub for genes in the top 122 differentially expressed list. Left panel shows relationship for BD compared to controls, right panel shows relationship for SZ compared to controls. Many of the genes are oppositely regulated across BD and SZ.

(6) FIG. 5 is a graph of experimental data demonstrating the ROC curve for visits 2 and 4.

(7) FIG. 6 is a graph of experimental data demonstrating logistic model estimated probabilities for visits 2 and 4.

(8) FIG. 7 is a graph of experimental data demonstrating the logistic model estimated probabilities for each group, for each visit along with the reference line for the 0.1518 cut-point.

(9) FIG. 8 is a graph of experimental data demonstrating logistic model estimated probabilities for visits 2 and 4.

(10) FIG. 9 is a graph of experimental data demonstrating the ROC curve for visits 2 and 4 for BD vs SZ.

(11) FIG. 10 is a graph of experimental data demonstrating the logistic model estimated probabilities for each group.

(12) FIG. 11 is a graph of experimental data demonstrating the ROC curve for visits 2 and 4 for SZ vs NC.

(13) FIG. 12 is a graph of experimental data demonstrating the logistic model estimated probabilities for each group.

DETAILED DESCRIPTION

(14) The present invention provides biomarkers that can discriminate between normal, BD and SZ subjects. The biomarkers of the invention can be used to screen, assess risk, diagnose and monitor the onset or progression of psychotic disorders and mood disorders. The biomarkers of the invention can be used to identify subjects in need of treatment for BD and SZ.

(15) The present invention therefore provides compositions and methods of diagnosing a subject as having SZ or BD, by examining relevant biomarkers and their expression. In one embodiment, biomarker expression includes transcription into messenger RNA (mRNA) and/or translation into protein, as well as transcription into types of RNA such as transfer RNA (tRNA) and ribosomal RNA (rRNA) that are not translated into protein.

(16) In one embodiment, the invention provides a method for diagnosing a subject with SZ or BD.

In one embodiment, the assay includes detecting expression of relevant biomarkers. In one embodiment, the method includes detecting expression of a combination of biomarker genes. In one embodiment, the combination of biomarker genes is at least two genes selected from the group SH3YL1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5. In one embodiment, the combination of genes is at least two genes selected from the group TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5.

(17) In one embodiment, the invention provides method for diagnosing a subject with SZ. In one embodiment, the method includes evaluating expression of one or more relevant biomarkers. In one embodiment, the method includes detecting expression of a combination of biomarker genes. In one embodiment, the combination of biomarker genes is at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5. In one embodiment, the combination of genes is at least two of HPR, TREML4, PTGDS, CPA3, TRIM4 and SLC44A5. In one embodiment, the combination of genes is at least two of SLC44A5, CPA3, TREML4, TRIM4, PTGDS and SH3YL1. In one embodiment, expression of the combination of genes is used to determine the probability of a patient having SZ. In one embodiment, a patient is diagnosed as having SZ on the basis of the probability of the condition as compared to a pre-determined cut-off from a logistical regression model for the specific set of genes analyzed.

(18) In one embodiment, the invention provides a method for diagnosing a subject with BD. In one embodiment, the method includes evaluating expression of one or more relevant biomarkers as compared to a comparator control. In one embodiment, the method includes detecting expression of a combination of biomarker genes. In one embodiment, the combination of biomarker genes is at least two of PTGDS, HLA-DRB5, OXTR and FADS2. In one embodiment, the combination of biomarker genes is at least two of CRIP2, CPA3, OXTR, TRIM41, PTGDS and SH3YL1. In one embodiment, expression of the combination of biomarker genes is used to determine the probability of a patient having BD. In one embodiment, a patient is diagnosed as having BD on the basis of the probability of the condition as compared to a pre-determined cut-off from a logistical regression model for the specific set of genes analyzed.

(19) In one embodiment, the invention provides a method for differentiating between a subject with SZ or BD and a healthy subject. In one embodiment, the method includes evaluating expression of one or more relevant biomarkers as compared to a comparator control. In one embodiment, the method includes detecting expression of a combination of biomarker genes. In one embodiment, the combination of biomarker genes is at least two of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8. In one embodiment, the combination of biomarker genes is at least two of SLC44A5, CPA3, CRIP2, TRIM4, PTGDS and SH3YL1. In one embodiment, a patient is diagnosed as having SZ or BD on the basis of the probability of having one of the conditions as compared to a pre-determined cut-off from a logistical regression model for the specific set of genes analyzed.

(20) In one embodiment, the method further provides for differentially diagnosing a subject characterized as having “SZ or BD” as having “BD” or “SZ.” In one embodiment, the method comprises evaluating expression of a combination of relevant biomarkers in a subject having been identified as having “SZ or BD”. In one embodiment, the combination of biomarker genes is at least two of CRIP2, OXTR and FADS2. In one embodiment, the combination of biomarker genes is at least two of SH3YL1, OXTR, PTGDS, CPA3, TBC1D1, and TCEA3. In one embodiment, expression of the combination of biomarker genes is used to determine the probability of a patient having “BD” or “SZ”. In one embodiment, a patient is diagnosed as having SZ or BD on the basis of the probability of each condition as compared to a pre-determined cut-off from a logistical regression model for the specific set of genes analyzed.

(21) In one embodiment, the method is useful for differentiating between “SZ” and “BD” in a

subject. In one embodiment, the subject has a prior diagnosis of “SZ” or “BD”. In one embodiment, a subject has no prior diagnosis of either “SZ” or “BD”.

(22) In one embodiment, a prior diagnosis of either “SZ” or “BD” is confirmed using the methods of the invention. In one embodiment, a prior diagnosis of either “SZ” or “BD” is identified as being a misdiagnosis either “SZ” or “BD” based on the methods of the invention. Therefore, in one embodiment, the invention provides a method of correctly diagnosing a subject with a prior diagnosis of “SZ” as having “BD,” In an alternative embodiment, the invention provides a method of correctly diagnosing a subject with a prior diagnosis of “BD” as having “SZ.”

(23) In one embodiment, the invention provides a multi-step method for differentiating or diagnosing a subject as having “SZ” or “BD.” In one embodiment, the invention comprises a first step of distinguishing a subject having “SZ or BD” from the normal population. In one embodiment, the method comprises a further step of differentially diagnosing a subject identified as having as having “SZ or BD” as having either “SZ” or “BD.” In one embodiment, the method comprises a further step of performing a secondary analysis for “SZ” on a subject identified as having as belonging to the normal population. In one embodiment, the method comprises a further step of performing a secondary analysis for “BD” on a subject identified as having as belonging to the normal population. In one embodiment, the method includes detecting expression of different combinations of relevant biomarkers for each determination. In one embodiment, the method further includes using logistic regression models to identify whether expression of a combination of biomarkers is above or below a predetermined cut-off.

(24) In one exemplary embodiment, the method includes detecting expression of a first combination of genes to distinguishing a subject having “SZ or BD” from the normal population. In one embodiment, the first combination of genes is at least two of SLC44A5, HAMA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8. In one embodiment, a result of a logistic regression model, based on the expression at a combination of genes, is determined, wherein the result is the probability of a sample being from a healthy subject. In one embodiment, if the probability of a sample being from a healthy subject is less than or equal to a pre-determined cut-off then the sample is identified as being from a subject having SZ or RD.

(25) In one embodiment, a pre-determined cut-off is 0.1518.

(26) In one embodiment, a subject identified as having “SZ or BD” is further evaluated at a second combination of genes to differentially diagnose the subject as having “SZ” or “BD.” In one embodiment, the second combination of genes is at least two of CRIP2, OXTR and FADS2. In one embodiment, a result of a logistic regression model, based on the expression at a combination of genes, is determined, wherein the result is the probability of a sample being from a subject with BD. In one embodiment, if the probability of a sample being from a subject with BD is less than or equal to a pre-determined cut-off then the sample is identified as being from a subject having SZ. In one embodiment, if the probability of a sample being from a subject with BD is greater than a pre-determined cut-off then the sample is identified as being from a subject having BD. In one embodiment, a pre-determined cut-off is 0.2857.

(27) In one embodiment, a subject identified as likely being from the normal population is further evaluated at one or more additional combination of genes useful for diagnosing the subject as having “SZ” or “BD.” In one embodiment, an additional combination of genes useful for diagnosing “SZ” is at least two of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5. In one embodiment, an additional combination of genes useful for diagnosing “BD” is at least two of PTGDS, HLA-DRB5, OXTR and FADS2. In one embodiment, a result of a logistic regression model, based on the expression at a combination of genes, is determined, wherein the result is the probability of a sample being from a subject with “BD” or “SZ.” In one embodiment, if the probability of a sample being from a healthy subject is less than or equal to a pre-determined cut-off then the sample is identified as being from a subject having “BD” or “SZ.” In one embodiment, if the probability of a sample being from a subject with “BD” or “SZ” is greater than a pre-

determined cut-off then the sample is identified as being from a subject having “BD” or “SZ”.

(28) Accordingly, in some embodiments of the invention, methods for diagnosing SZ or BD in a subject is provided. The methods comprise a) providing a sample from the subject; b) analyzing the sample with an assay that specifically detects a combination of biomarkers of the invention in the sample; c) evaluating gene expression at one or more combination of biomarkers and d) diagnosing SZ or BD in the subject.

(29) In one embodiment, the step of analyzing the sample with an assay that specifically detects a combination of biomarkers of the invention in the sample comprises contacting a sample from a subject with an assay for detecting the expression levels of at least two biomarkers selected from the group SH3YL1, TBC1D1, TCEA3, SLC44A5, HAMA, CPA3, IL5RA, OXTR, CCDC10913, TREM4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDXS, EEF2, ZMYND8 and HLA-DRB5 in the sample. In one embodiment, the assay detected the expression levels of at least two of the biomarkers selected from the group TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group TCEA3, SIX44A5, IL5RA, GYLTL1B and DDX5. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group HPR, TREML4, PTGDS, CPA3, TRIM4 and SLC44A5. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group SLC44A5, CPA3, TREML4, TRIM4, PTGDS and SH3YL1. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group PTGDS, HLA-DRB5, OXTR and FADS2. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group CRIP2, CPA3, OXTR, TRIM4, PTGDS and SH3YL1. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group SLC44A5, CPA3, CRIP2, TRIM4, PTGDS and SH3YL1. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group CRIP2, OXTR and FADS2. In one embodiment, the assay detects the expression levels of at least two of the biomarkers selected from the group SH3YL1, OXTR, PTGDS, CPA3, TBC1D1, and TCEA3.

(30) In one embodiment, the step of evaluating gene expression of one or more combinations of biomarkers comprises comparing the expression levels of the combination of at least two biomarkers selected from the group SH3YL1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EEF2, ZMYND8 and HLA-DRB5 between the sample and a comparator control. In one embodiment, the comparator control is expression levels in a normal subject, or a healthy profile. In one embodiment, the comparator control is a predetermined probability cut-off based on logistical regression analysis.

(31) In one embodiment, expression of the full length protein is detected. In one embodiment, expression of a fragment of the full length protein is detected.

(32) In one embodiment, the biomarker types comprise mRNA biomarkers. In various embodiments, the mRNA is detected by at least one of mass spectroscopy, PCR microarray, thermal sequencing, capillary array sequencing, solid phase sequencing, and the like.

(33) In another embodiment, the biomarker types comprise polypeptide biomarkers. In various embodiments, the polypeptide is detected by at least one of ELISA, Western blot, flow cytometry, immunofluorescence, immunohistochemistry, mass spectroscopy, and the like.

(34) Definitions

(35) Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains.

Although any methods and materials similar or equivalent to those described herein can be used in the practice for testing of the present invention, the preferred materials and methods are described herein. In describing and claiming the present invention, the following terminology will be used.

(36) It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting.

(37) The articles “a” and “an” are used herein to refer to one or to more than one (i.e., to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

(38) “About” as used herein when referring to a measurable value such as an amount, a temporal duration, and the like, is meant to encompass non-limiting variations of $\pm 40\%$ or $\pm 20\%$ or $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, or $\pm 0.1\%$ from the specified value, as such variations are appropriate.

(39) The term “amplification” refers to the operation by which the number of copies of a target nucleotide sequence present in a sample is multiplied.

(40) As used herein, the term “marker” or “biomarker” is meant to include a parameter (e.g., RNA, polypeptide, etc.) which is useful according to this invention for determining the presence and/or severity and/or stage of SZ or BD.

(41) The term “control or reference standard or comparator” describes a material comprising none, or a normal, low, or high level of one or more of the marker (or biomarker) expression products of one or more the markers (or biomarkers) of the invention, such that the control or reference standard or comparator may serve as a comparator against which a sample can be compared.

(42) By the phrase “determining the level of marker (or biomarker) expression” is meant an assessment of the degree of expression of a marker in a sample at the nucleic acid or protein level, using technology available to the skilled artisan to detect a sufficient portion of any marker expression product.

(43) The level of a marker or biomarker “significantly” differs from the level of the marker or biomarker in a reference sample or comparator if the level of the marker in a sample from the patient differs from the level in a reference sample or comparator by an amount greater than the standard error of the assay employed to assess the marker, and preferably at least 10% or more, for example, 20%, 30%, 40%, or 50%, 60%, 70%, 80%, 90% different or more, and/or 1.1 fold, 1.2 fold, 1.4 fold, 1.6 fold, 1.8 fold, 2.0 fold different or more.

(44) “Differentially increased expression” or “up regulation” refers to biomarker product levels which are at least 10% or more, for example, 20%, 30%, 40%, or 50%, 60%, 70%, 80%, 90% higher or more, and/or 1.1 fold, 1.2 fold, 1.4 fold, 1.6 fold, 1.8 fold, 2.0 fold higher or more, and any and all whole or partial increments therebetween than a control.

(45) “Differentially decreased expression” or “down regulation” refers to biomarker product levels which are at least 10% or more, for example, 20%, 30%, 40%, or 50%, 60%, 70%, 80%, 90% lower or less, and/or 2.0 fold, 1.8 fold, 1.6 fold, 1.4 fold, 1.2 fold, 1.1 fold or less lower, and any and all whole or partial increments therebetween than a control.

(46) A “disease” is a state of health of an animal wherein the animal cannot maintain homeostasis, and wherein if the disease is not ameliorated then the animal's health continues to deteriorate.

(47) As used herein, an “immunoassay” refers to a biochemical test that measures the presence or concentration of a substance in a sample, such as a biological sample, using the reaction of an antibody to its cognate antigen, for example the specific binding of an antibody to a protein. Both the presence of the antigen or the amount of the antigen present can be measured.

(48) As used herein, an “instructional material” includes a publication, a recording, a diagram, or any other medium of expression which can be used to communicate the usefulness of a component of the invention in a kit for detecting biomarkers disclosed herein. The instructional material of the kit of the invention can, for example, be affixed to a container which contains the component of the invention or be shipped together with a container which contains the component. Alternatively, the instructional material can be shipped separately from the container with the intention that the

instructional material and the component be used cooperatively by the recipient.

(49) The term “label” when used herein refers to a detectable compound or composition that is conjugated directly or indirectly to a probe to generate a “labeled” probe. The label may be detectable by itself (e.g., radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition that is detectable (e.g., avidin-biotin). In some instances, primers can be labeled to detect a PCR product.

(50) The “level” of one or more biomarkers means the absolute amount or relative amount or concentration of the biomarker in the sample.

(51) The term “marker (or biomarker) expression” as used herein, encompasses the transcription, translation, post-translation modification, and phenotypic manifestation of a gene, including all aspects of the transformation of information encoded in a gene into RNA or protein. By way of non-limiting example, biomarker expression includes transcription into messenger RNA (mRNA) and translation into protein, as well as transcription into types of RNA such as transfer RNA (tRNA) and ribosomal RNA (rRNA) that are not translated into protein.

(52) The terms “microarray” and “array” refers broadly to both “DNA microarrays” and “DNA chip(s),” and encompasses all art-recognized solid supports, and all art-recognized methods for affixing nucleic acid molecules thereto or for synthesis of nucleic acids thereon. Preferred arrays typically comprise a plurality of different nucleic acid probes that are coupled to a surface of a substrate in different, known locations. These arrays, also described as “microarrays” or colloquially “chips” have been generally described in the art, for example, U.S. Pat. Nos. 5,143,854, 5,445,934, 5,744,305, 5,677,195, 5,800,992, 6,040,193, 5,424,186 and Fodor et al., Science, 1991; 251:767-777, each of which is incorporated by reference in its entirety for all purposes. Arrays may generally be produced using a variety of techniques, such as mechanical synthesis methods or light directed synthesis methods that incorporate a combination of photolithographic methods and solid phase synthesis methods. Techniques for the synthesis of these arrays using mechanical synthesis methods are described in, e.g., U.S. Pat. Nos. 5,384,261, and 6,040,193, which are incorporated herein by reference in their entirety for all purposes. Although a planar array surface is preferred, the array may be fabricated on a surface of virtually any shape or even a multiplicity of surfaces. Arrays may be nucleic acids on beads, gels, polymeric surfaces, fibers such as fiber optics, glass or any other appropriate substrate. (See U.S. Pat. Nos. 5,770,358, 5,789,162, 5,708,153, 6,040,193 and 5,800,992, which are hereby incorporated by reference in their entirety for all purposes.) Arrays may be packaged in such a manner as to allow for diagnostic use or can be an all-inclusive device; e.g., U.S. Pat. Nos. 5,856,174 and 5,922,591 incorporated in their entirety by reference for all purposes. Arrays are commercially available from, for example, Affymetrix (Santa Clara, Calif.) and Applied Biosystems (Foster City, Calif.), and are directed to a variety of purposes, including genotyping, diagnostics, mutation analysis, biomarker expression, and gene expression monitoring for a variety of eukaryotic and prokaryotic organisms. The number of probes on a solid support may be varied by changing the size of the individual features. In one embodiment the feature size is 20 by 25 microns rectangle, in other embodiments features may be, for example, 8 by 8, 5 by 5 or 3 by 3 microns rectangle, resulting in about 2,600,000, 6,600,000 or 18,000,000 individual probe features.

(53) “Measuring” or “measurement,” or alternatively “detecting” or “detection,” means assessing the presence, absence, quantity or amount (which can be an effective amount) of either a given substance within a clinical or subject-derived sample, including the derivation of qualitative or quantitative concentration levels of such substances, or otherwise evaluating the values or categorization of a subject's clinical parameters.

(54) A “healthy” or “normal” subject does not have any form of schizophrenia or bipolar disorder.

(55) The terms “patient,” “subject,” “individual,” and the like are used interchangeably herein, and refer to any animal, or cells thereof whether in vitro or in situ, amenable to the methods described herein. In certain non-limiting embodiments, the patient, subject or individual is a human.

(56) A “reference level” of a biomarker means a level of the biomarker that is indicative of a particular disease state, phenotype, or lack thereof, as well as combinations of disease states, phenotypes, or lack thereof. A “positive” reference level of a biomarker means a level that is indicative of a particular disease state or phenotype. A “negative” reference level of a biomarker means a level that is indicative of a lack of a particular disease state or phenotype.

(57) The term “risk stratification,” according to the invention, comprises finding schizophrenic or bipolar patients, particularly those having an early or first psychotic episode, for the purpose of diagnosis and therapy/treatment of the schizophrenic or bipolar condition, with the goal of allowing as advantageous a course of the schizophrenic or bipolar condition as possible.

(58) “Sample” or “biological sample” as used herein means a biological material isolated from a subject. The biological sample may contain any biological material suitable for detecting the desired biomarkers, and may comprise cellular and/or non-cellular material obtained from the subject. One example of a biological sample is a whole blood sample. Another example of a biological sample is a cell-free serum sample.

(59) “Standard control value” as used herein refers to a predetermined amount of a particular protein or nucleic acid that is detectable in a control sample. The standard control value is suitable for the use of a method of the present invention, in order for comparing the amount of a protein or nucleic acid of interest (e.g., marker, biomarker) that is present in a patient sample. An established sample serving as a standard control provides an typical amount of the protein or nucleic acid of interest in a sample type that is typical for an typical, healthy person of reasonably matched background, e.g., gender, age, ethnicity, and medical history. A standard control value may vary depending on the protein or nucleic acid of interest and the nature of the sample (e.g., whole blood, serum, etc.).

(60) Ranges: throughout this disclosure, various aspects of the invention can be presented in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the invention. Accordingly, the description of a range should be considered to have specifically disclosed all the possible subranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed subranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 2.7, 3, 4, 5, 5.3, and 6. This applies regardless of the breadth of the range.

(61) Description

(62) The present invention is based on the identification of biomarkers, the expression levels of which can discriminate between normal, SZ and BD subjects in a biological sample of a subject.

(63) In one embodiment, the invention provides a combination of biomarkers for the diagnosis of SZ. In one embodiment, the invention provides a combination of biomarkers for the diagnosis of BD. In one embodiment, the invention provides a combination of biomarkers for the diagnosis of “SZ or BD.” In one embodiment, the invention provides a combination of biomarkers for differentially diagnosing a subject originally diagnosed as having “SZ or BD” with “SZ” or “BD.”

(64) Identifying a Biomarker

(65) The invention includes methods for the identification of biomarkers differentially expressed between samples of normal, SZ and BD subjects, as well as methods for the detection of the expression products of differentially expressed biomarkers of normal, SZ and BD subjects.

(66) The invention contemplates the identification of differentially expressed biomarkers by whole genome nucleic acid microarray, to identify biomarkers differentially expressed between normal, SZ and BD subjects. The invention further contemplates using methods known to those skilled in the art to detect and to measure the level of differentially expressed biomarker expression products, such as RNA and protein, to measure the level of one or more differentially expressed biomarker expression products. In certain embodiments, the expression level of one or more regions or

fragments of a gene are more informative than the expression level of the entire gene, and therefore in one embodiment, a biomarker expression product is the expression of a fragment or region of a gene.

(67) Methods of detecting or measuring gene expression may utilize methods that focus on cellular components (cellular examination), or methods that focus on examining extracellular components (fluid examination). Because gene expression involves the ordered production of a number of different molecules, a cellular or fluid examination may be used to detect or measure a variety of molecules including RNA, protein, and a number of molecules that may be modified as a result of the protein's function. Typical diagnostic methods focusing on nucleic acids include amplification techniques such as PCR and RT-PCR (including quantitative variants), and hybridization techniques such as in situ hybridization, microarrays, blots, and others. Typical diagnostic methods focusing on proteins include binding techniques such as ELISA, immunohistochemistry, microarray and functional techniques such as enzymatic assays.

(68) The genes identified as being differentially expressed may be assessed in a variety of nucleic acid detection assays to detect or quantify the expression level of a gene or multiple genes in a given sample. For example, traditional Northern blotting, nuclease protection, RT-PCR, microarray, and differential display methods may be used for detecting gene expression levels. Methods for assaying for mRNA include Northern blots, slot blots, dot blots, and hybridization to an ordered array of oligonucleotides. Any method for specifically and quantitatively measuring a specific protein or mRNA or DNA product can be used. However, methods and assays are most efficiently designed with array or chip hybridization-based methods for detecting the expression of a large number of genes. Any hybridization assay format may be used, including solution-based and solid support-based assay formats.

(69) The protein products of the genes identified herein can also be assayed to determine the amount of expression. Methods for assaying for a protein include Western blot, immunoprecipitation, and radioimmunoassay. The proteins analyzed may be localized intracellularly (most commonly an application of immunohistochemistry) or extracellularly (most commonly an application of immunoassays such as ELISA).

(70) Biological samples may be of any biological tissue or fluid. Frequently the sample will be a "clinical sample" which is a sample derived from a patient. The biological sample may contain any biological material suitable for detecting the desired biomarkers, and may comprise cellular and/or non-cellular material obtained from the subject. One example of a biological sample is a whole blood sample. Another example of a biological sample is a serum sample. Another example of a biological sample is a saliva sample. Another example of a biological sample is a urine sample.

(71) Controls group samples may either be from a normal subject or from a subject with a known stage of SZ or BD. As described below, comparison of the expression patterns of the sample to be tested with those of the controls can be used to diagnose, or distinguish between, normal, SZ and BD subjects. In some instances, the control groups are only for the purposes of establishing initial cutoffs or thresholds for the assays of the invention. Therefore, in some instances, the systems and methods of the invention can diagnose between normal, SZ and BD subjects without the need to compare with a control group.

(72) Methods of Differentiation and Diagnosis

(73) The present invention relates to the identification of biomarkers associated with SZ and BD. Accordingly, the present invention features methods for identifying subjects who have developed SZ and BD, and for differentiating between SZ and BD in a subject. Subjects include those subjects who are suffering a first psychotic episode or have not yet been fully clinically characterized as having SZ or BD and subjects who have already been diagnosed as having SZ or BD. In one embodiment, the methods may be useful for identifying a subject as having a risk of developing SZ or BD. In one embodiment, the risk of SZ or BD can be assessed by measuring a combination of the biomarkers described herein, and comparing the measured values to comparator values,

reference values, or index values. Such a comparison can be undertaken with mathematical algorithms or formula in order to combine information from results of multiple individual biomarkers and other parameters into a single measurement or index. Subjects identified as having an increased risk of developing SZ or BD can optionally be selected to receive treatment regimens, such as administration of prophylactic or therapeutic compounds or implementation of exercise regimens or dietary supplements to prevent, treat or delay the onset of SZ or BD.

(74) Identifying a subject before they develop SZ or BD, or shortly after a first psychotic episode, enables the selection and initiation of various therapeutic interventions or treatment regimens in order to delay, reduce or prevent the manifestation and progression of the disorder in the subject.

(75) The biomarkers of the present invention can thus be used to generate a biomarker profile or signature of the subjects: (i) who do not have and are not expected to develop SZ or BD and/or (ii) who have SZ or BD. The biomarker profile of a subject can be compared to a predetermined or comparator biomarker profile or reference biomarker profile to diagnose or identify subjects at risk of developing SZ or BD, to monitor the progression of the disorder, and to monitor the effectiveness of treatments. Data concerning the biomarkers of the present invention can also be combined or correlated with other data or test results, such as, without limitation, measurements of clinical parameters or other algorithms for SZ or BD. Other data includes, but is not limited to, gender, age and ethnicity. The machine-readable media can also comprise subject information such as medical history and any relevant family history.

(76) In one embodiment, the invention is a method of diagnosing SZ. In one embodiment, the invention is a method of diagnosing BD. In one embodiment, the method includes differentiating or distinguishing between normal, SZ and BD subjects.

(77) In various embodiments, methods are disclosed herein that may be of use to determine whether a subject has a SZ or BD. In some embodiments, these methods may utilize a biological sample (such as urine, saliva, blood, serum, amniotic fluid, or tears), for the detection of a combination of biomarkers of the invention in the sample.

(78) In one embodiment, the method includes evaluating the expression level of a combination of biomarkers useful for distinguishing between normal, SZ and BD. In various embodiments, the combination of biomarkers is two or more biomarkers selected from the group consisting of SH3YL1, TBC1D1, TCEA3, SLC44A5, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EE2, ZMYND8 and HLA-DRB5.

(79) In one embodiment, the method includes evaluating the expression level of a combination of biomarkers for diagnosis of SZ. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of TCEA3, SLC44A5, IL5RA, GYLTL1B and DDX5. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of HPR, PTGDS, CPA3, TRIM4 and SLC44A5. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of SLC44A5, CPA3, TREML4, TRIM4, PTGDS and SH3YL1.

(80) In one embodiment, the method includes evaluating the expression level of a combination of biomarkers for the diagnosis of BD. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of PTGDS, HLA-DRB5, OXTR and FADS2. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of CRIP2, CPA3, OXTR, TRIM4, PTGDS and SH3YL1.

(81) In one embodiment, the method includes evaluating the expression level of a combination of biomarkers for distinguishing a subject having SZ or BD from a healthy subject. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of SLC44A5, CPA3, CRIP2, TRIM4, PTGDS and SH3YL1.

(82) In one embodiment, the method includes evaluating the expression level of a combination of

biomarkers for differentially diagnosing a subject identified as having “SZ or BD” as having “SZ” or “BD.” In one embodiment, the combination of biomarkers is at least two selected from the group consisting of CRIP2, OXTR and FADS2. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of SH3YL1, OXTR, PTGDS, CPA3, TBC1D1, and TCEA3.

(83) In one embodiment, the method includes evaluating the expression level of a combination of biomarkers for identifying a healthy subject with respect to BD or SZ. In one embodiment, the combination of biomarkers is at least two selected from the group consisting of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, HPR and ZMYND8.

(84) In one embodiment, the method comprises detecting a combination of biomarkers in a biological sample of the subject. In one embodiment, the biological sample is blood. In various embodiments, the level of one or more of biomarkers of the invention in the biological sample of the subject is compared with the level of the biomarker in a comparator. Non-limiting examples of comparators include, but are not limited to, a negative control, a positive control, standard control, standard value, an expected normal background value of the subject, a historical normal background value of the subject, a reference standard, a reference level, an expected normal background value of a population that the subject is a member of, or a historical normal background value of a population that the subject is a member of.

(85) In various embodiments, the subject is a human subject, and may be of any race, sex and age. In one embodiment, the subject is a human male less than 30 years old.

(86) Information obtained from the methods of the invention described herein can be used alone, or in combination with other information (e.g., disorder status, disorder history, vital signs, blood chemistry, etc.) from the subject or from the biological sample obtained from the subject.

(87) In various embodiments of the methods of the invention, the level of one or more biomarkers of the invention is determined to be increased when the level of one or more of the biomarkers of the invention is increased by at least 10%, by at least 20%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, or by at least 100%, when compared to with a comparator.

(88) In other various embodiments of the methods of the invention, the level of one or more biomarkers of the invention is determined to be decreased when the level of one or more of the biomarkers of the invention is decreased by at least 10%, by at least 20%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 80%, by at least 90%, or by at least 100%, when compared to with a comparator.

(89) In the methods of the invention, a biological sample from a subject is assessed for the level of one or more of the biomarkers of the invention in the biological sample obtained from the patient. The level of one or more of the biomarkers of the invention in the biological sample can be determined by assessing the amount of polypeptide of one or more of the biomarkers of the invention in the biological sample, the amount of mRNA of one or more of the biomarkers of the invention in the biological sample, the amount of enzymatic activity of one or more of the biomarkers of the invention in the biological sample, or a combination thereof.

(90) Detecting a Biomarker

(91) In one embodiment, the invention includes detecting an mRNA in a bodily fluid, wherein the bodily fluid is blood and the mRNA is detected in blood. In some embodiments, detection of mRNAs is performed in a portion of a blood sample (e.g., serum).

(92) In one embodiment, detecting mRNAs, is performed in a bodily fluid, e.g. saliva or urine, which meets the demands of an inexpensive, non-invasive and accessible bodily fluid to act as an ideal medium for investigative analysis.

(93) Biomarkers generally can be measured and detected through a variety of assays, methods and detection systems known to one of skill in the art. Various methods include but are not limited to

refractive index spectroscopy (RI), ultra-violet spectroscopy (UV), fluorescence analysis, electrochemical analysis, radiochemical analysis, near-infrared spectroscopy (near-IR), infrared (IR) spectroscopy, nuclear magnetic resonance spectroscopy (NMR), light scattering analysis (LS), mass spectrometry, pyrolysis mass spectrometry, nephelometry, dispersive Raman spectroscopy, gas chromatography, liquid chromatography, gas chromatography combined with mass spectrometry, liquid chromatography combined with mass spectrometry, matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) combined with mass spectrometry, ion spray spectroscopy combined with mass spectrometry, capillary electrophoresis, colorimetry and surface plasmon resonance (such as according to systems provided by Biacore Life Sciences). See also PCT Publications WO/02004/056456 and WO/2004/088309. In this regard, biomarkers can be measured using the above-mentioned detection methods, or other methods known to the skilled artisan. Other biomarkers can be similarly detected using reagents that are specifically designed or tailored to detect them.

(94) Different types of biomarkers and their measurements can be combined in the compositions and methods of the present invention. In various embodiments, the protein form of the biomarkers is measured. In various embodiments, the nucleic acid form of the biomarkers is measured. In exemplary embodiments, the nucleic acid form is mRNA. In various embodiments, measurements of protein biomarkers are used in conjunction with measurements of nucleic acid biomarkers.

(95) Methods for detecting mRNA, such as RT-PCR, real time PCR, branch DNA, NASBA, RNA-Seq, digital droplet PCR, and others, are well known in the art. Using sequence information provided by the database entries for the biomarker sequences, expression of the biomarker sequences can be detected (if present) and, measured using techniques well known to one of ordinary skill in the art. For example, sequences in sequence database entries or sequences disclosed herein can be used to construct probes for detecting biomarker RNA sequences in, e.g., Northern blot hybridization analyses or methods which specifically, and, preferably, quantitatively amplify specific nucleic acid sequences. As another example, the sequences can be used to construct primers for specifically amplifying the biomarker sequences in, e.g., amplification-based detection methods such as reverse-transcription based polymerase chain reaction (RT-PCR). When alterations in gene expression are associated with gene amplification, deletion, polymorphisms and mutations, sequence comparisons in test and reference populations can be made by comparing relative amounts of the examined DNA sequences in the test and reference cell populations. In addition to Northern blot and RT-PCR, RNA can also be measured using, for example, other target amplification methods (e.g., TMA, SDA, NASBA), signal amplification methods (e.g., bDNA), nuclease protection assays, in situ hybridization and the like.

(96) The concentration of the biomarker in a sample may be determined by any suitable assay. A suitable assay may include one or more of the following methods, an enzyme assay, an immunoassay, mass spectrometry, chromatography, electrophoresis or an antibody microarray, or any combination thereof. Thus, as would be understood by one skilled in the art, the system and methods of the invention may include any method known in the art to detect a biomarker in a sample.

(97) The invention described herein also relates to methods for a multiplex analysis platform. In one embodiment, the method comprises an analytical method for multiplexing analytical measurements of biomarkers. In another embodiment, the method comprises a set of compatible analytical strategies for multiplex measurements of biomarkers and/or metabolites in a sample.

(98) Evaluating Expression Level of a Biomarker

(99) In one embodiment, the expression of a combination of biomarkers of the invention is representative of a health state and is diagnostically useful for determining the health state of a subject. One or more statistical methods, as disclosed herein, can be used to evaluate the expression level of a combination of biomarkers of the invention. Exemplary statistical methods include, for example, discriminant analysis, classification analysis, cluster analysis, analysis of variance

(ANOVA), regression analysis, regression trees, decision trees, nearest neighbor algorithms, principal components, factor analysis, multidimensional scaling and other methods of dimensionality reduction, likelihood models, hypothesis testing, kernel density estimation and other smoothing techniques, cross-validation and other methods to guard against overfitting of the data, the bootstrap and other statistical resampling techniques, artificial intelligence, including artificial neural networks, machine learning, data mining, and boosting algorithms, and Bayesian analysis using prior probability distributions.

(100) In one embodiment, a logistic regression model is used to evaluate the expression level of a combination of biomarkers of the invention. In one embodiment, a diagnostic cut-off for SZ, BD, control or a combination thereof is determined from a logistic regression model and used to diagnose a subject as having SZ, BD, "SZ or BD," or control. Due to alterations in gene expression by white blood cell counts, age and gender, and body-mass index, normative ranges may be adjusted for these covariates in logistic regression analysis.

(101) Kits

(102) The present invention also pertains to kits useful in the methods of the invention. Such kits comprise various combinations of components useful in any of the methods described elsewhere herein, including for example, materials for quantitatively analyzing a biomarker of the invention (e.g., polypeptide and/or nucleic acid), materials for assessing the activity of a biomarker of the invention (e.g., polypeptide and/or nucleic acid), and instructional material. For example, in one embodiment, the kit comprises components useful for the quantification of a desired nucleic acid in a biological sample. In another embodiment, the kit comprises components useful for the quantification of a desired polypeptide in a biological sample. In a further embodiment, the kit comprises components useful for the assessment of the activity (e.g., enzymatic activity, substrate binding activity, etc.) of a desired polypeptide in a biological sample.

(103) In a further embodiment, the kit comprises components for determining the level of a combination of biomarkers of the invention in a biological sample obtained from the subject. In one embodiment, a kit is a diagnostic kit for SZ or BD and comprises at least two probes for detecting expression of at least two biomarkers selected from the group consisting of SH3 YL1, TBC1D1, TCEA3, SLC44AS, HADHA, CPA3, IL5RA, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, GYLTL1B, FADS2, CRIP2, HPR, DDX5, EFF2, ZMYND8 and HLA-DRB5.

(104) In various embodiments, the kit comprises instructional material to determine whether the level of a biomarker of the invention is modulated in a biological sample obtained from the subject, as compared with the level of at least one comparator control contained in the kit, such as a positive control, a negative control, a historical control, a historical norm, or the level of another reference molecule in the biological sample.

(105) Treatments

(106) In certain embodiments, treatment comprises administering a disorder-modulating treatment to a subject. In some embodiments, the disorder-modulating treatment is a drug. The drug can be a therapeutic or prophylactic used in subjects diagnosed or identified with SZ or BD, or at risk of having SZ or BD. In certain embodiments, treatment comprises modifying a therapy used in subjects diagnosed or identified with SZ or BD, or at risk of having SZ or BD. In one embodiment, modifying therapy refers to altering the drug, dosage, duration, frequency or intensity of therapy, or for example, altering the type of therapy provided to the subject.

(107) In various embodiments, effecting a therapy comprises causing a subject to or communicating to a subject the need to make a change in lifestyle, for example, increasing exercise, changing diet, reducing or eliminating smoking, taking a drug, and so on.

(108) In various exemplary embodiments, effecting a therapy comprises administering a disorder-modulating drug to the subject. Any drug or combination of drugs useful for treating or mediating SZ or BD may be administered to a subject on the basis of the diagnosis of SZ or BD. The drugs can be formulated in any number of ways, often according to various known formulations in the art

or as disclosed or referenced herein.

(109) In various embodiments, a drug or combination of drugs is not administered to a subject to treat a disorder. In these embodiments, the practitioner may refrain from administering the drug or combination of drugs, may recommend that the subject not be administered the drug or combination of drugs or may prevent the subject from being administered the drug or combination of drugs.

(110) In various embodiments, one or more additional drugs may be optionally administered in addition to those that are recommended or have been administered. An additional drug will typically not be any drug that is not recommended or that should be avoided. In exemplary embodiments, one or more additional drugs comprise one or more drugs approved for the treatment or mediation of SZ or BD.

(111) Thus, in various methods of diagnosis of the invention, the method also includes the step of administering a treatment to the subject.

EXPERIMENTAL EXAMPLES

(112) The invention is further described in detail by reference to the following experimental examples. These examples are provided for purposes of illustration only, and are not intended to be limiting unless otherwise specified. Thus, the invention should in no way be construed as being limited to the following examples, but rather, should be construed to encompass any and all variations which become evident as a result of the teaching provided herein.

Example 1: Exon Array Biomarkers to Diagnose and Differentiate Schizophrenia and Bipolar Disorder

(113) Without being bound by any particular theory, it was hypothesized that for SZ and BL) there are unique and global sets of chronic differentially expressed genes in blood for each disorder. This hypothesis was based, in part, on the concept that circulating blood reflects the changing health of the body, i.e the “Sentinel Principle” (Liew et al., *J Lab Clin Med* 2006; 147(3): 126-132). As blood circulates through the brain, communication occurs between cells in blood and the brain (Weber et al., *Neuropsychopharmacology* 2017; 42(1): 46-61). Following a brain injury, neutrophils, macrophages, lymphocytes, and dendritic cells can extravasate into the brain from the blood (Gelderblom et al., *Stroke* 2009; 40(5): 1849-1857). These extravasated cells can induce changes in gene expression and protein as reported in neuroimmune studies (Downes and Crack, *Br J Pharmacol* 2010; 160(8): 1872-1888; Brea et al., *Cerebrovasc Dis* 2009; 27 Suppl 1: 48-64). In prior work using blood-brain samples from the same subjects, about 20% of the transcriptome was expressed at comparable levels and significantly correlated in both tissues from the same subjects (Rollins et al., *Am J Med Genet B Neuropsychiatr Genet* 2010; 153B(4): 919-936). This correlation supports the idea that important biomarkers of either SZ or BD could be expressed peripherally, and these would also have a connection to the central modulation of neuroimmune responses.

Advantages of a peripheral transcriptomics study are the ease of access to whole blood and the fact that immune genes are highly expressed. For example, multiple HLA region genes with genome-wide significance are expressed in peripheral blood samples, such as C4 (Sekar et al., *Nature* 2016; 530(7589): 177-183) and HLA-DPA1 (Morgan et al., *Microarrays (Basel)* 2016; 5(1)).

(114) To date, there are no validated biomarker studies of schizophrenia or bipolar disorder using peripheral blood gene expression, although a number of serum proteins have been advanced as validated biomarkers (Chan et al., *Prog Neurobiol* 2014; 122: 45-72; Chan et al., *Transl Psychiatry* 2015; 5: e601; Haenisch et al., *Brain Behav Immun* 2016; 52: 49-57; Scarr et al., *Int J Neuropsychopharmacol* 2015; 18(10): pyv042; Tomasik et al., *Eur Arch Psychiatry Clin Neurosci* 2012; 262 Suppl 2: 579-83). A recent mega-analysis of blood samples comparing SZ and controls was conducted with a total of 578 subjects in nine studies. The mega-analysis of blood transcriptome, showed that 220 genes reached a Bonferroni-corrected level of significance (Hess et al., *Schizophr Res* 2016; 176(2-3): 114-124), indicating the utility of analysis of blood transcriptome for finding differentially expressed genes. There is little agreement among

researchers of which mRNA and protein are jointly dysregulated in the blood of neuropsychiatric patients leaving open the question of which are the most important genes and proteins still to be explored (Sanders et al., *Hum Mol Genet* 2013; 22(24): 5001-5014; Hess et al., *Schizophr Res* 2016; 176(2-3): 114-124; Glatt et al., *Curr Pharmacogenomics Person Med* 2009; 7(3): 164-188; Glatt et al., *Am J Med Genet B Neuropsychiatr Genet* 2011; 156B(8): 869-887; Horvath and Mimics, *Biol Psychiatry* 2014; 75(4): 316-323; Kumarasinghe et al., *Int J Neuropsychopharmacol* 2013; 16(7): 1483-1503; Wu et al., *Brain Behav Immun* 2016; 53: 194-206; Xu et al., *Sci Rep* 2016; 6: 16767; Martin et al., *BMC Med Genomics* 2009; 2: 62; Vawter et al., *Brief Funct Genomics* 2011; 10(6): 387-399; Manidani et al., *Dis Markers* 2013; 35(1): 11-21). The stability of this potential dysregulation has not been tested across illness progression.

(115) The objective of the present study is to test subjects at multiple time points by enrolling SZ (n=30), BD (n=30), and normal control (NC; n=30) subjects at one clinical site to eliminate potential sources of variation. The expression levels of panels of genes were used to define which disorder the patient was diagnosed with the highest likelihood. Tempus Blood RNA tubes and Affymetrix exon arrays were previously validated in a study that measured the sources of variation for 8 healthy controls at 9 sequential blood draws every 6 hours (Rollins et al., *Am J Med Genet B Neuropsychiatr Genet* 2010; 153B(4): 919-936). The resulting data showed that about 20% of the transcripts measured on the Affymetrix exon array did not significantly change over 9 blood draws (Rollins et al., *Am J Med Genet B Neuropsychiatr Genet* 2010; 153B(4): 919-936). The remaining 80% of transcripts were changed significantly during this experiment (Rollins et al., *Am J Med Genet B Neuropsychiatr Genet* 2010; 153B(4): 919-936). Taken together, it is expected that 20% of transcripts will be stable over longer periods of time and do not significantly fluctuate by time of day of the blood draw. This data reinforces the purpose of the present study to test whether stable expression of genes occurs over months instead of days that can be used as a biomarker for SZ and BD to differentiate from controls. In this study RNA expression was measured with the Affymetrix exon array 1.0 ST for biomarker screening. Exon arrays have been widely used for the study of genetic variation in coding regions (Bemmo et al., *BMC Genomics* 2008; 9: 529; Benovoy et al., *Nucleic Acids Res* 2008; 36(13): 4417-4423; Duan et al., *Bioinformatics* 2008; 24(10): 469-470; Heinzen et al., *PLoS Biol* 2008; 6(12): e12; Kwan et al., *Nat Genet* 2008; 40(2): 225-231; Sequeira et al., *Mol Psychiatry* 2008; 13(4): 363-365; Duan et al., *Pharmacogenomics* 2009; 10(4): 549-563; Fraser and Xie, *Genome Res* 2009; 19(4): 567-575; Martin et al., *BMC Med Genomics* 2009; 2: 62; Gamazon et al., *PLoS One* 2010; 5(2): e9366; Gamazon et al., *Bioinformatics* 2010; 26(2): 259-262; Pradervand et al., *BioTechniques* 2008; 44(6): 759-762). Exon array findings correlate positively with RNA-Seq across most levels of transcript expression (Agarwal et al., *BMC Genomics* 2010; 11: 383), and in some cases have less false-positive detection than RNA-Seq (Bradford et al., *BMC Genomics* 2010; 11: 282; Liu et al., *Nucleic Acids Res* 2011; 39(2): 578-88; Richard et al., *Nucleic Acids Res* 2010; 38(10): e112). The analytical techniques for the exon arrays have been well established (Kapur et al., *Bioinformatics* 2008; 24(24): 2887-2893; Kapur et al., *Genome Biol* 2007; 8(5): R82; Xing et al., *PLoS One* 2006; 1: e88; Xing et al., *Mol Biol Evol* 2007; 24(6): 1283-1285; Xing et al., *RNA* 2008; 14(8): 1470-1479; Lin et al., *Nucleic Acids Res* 2009; 37(12): e90; Shen et al., *Bioinformatics* 2010; 26(2): 268-269; Liu et al., *Nucleic Acids Res* 2011; 39(2): 578-588).

(116) To determine stable temporal biomarkers, this study evaluated whole blood gene expression at two different time points using the same subjects to differentiate schizophrenia, bipolar disorder type I, and controls. The diagnostic algorithm uses logistic regression modeling and a total of 18 unique expressed exons within known mRNA transcripts. The model discriminated schizophrenia and bipolar disorder from each other, as well as both from healthy controls in four steps. The upper limit of accuracy achieved in this study was 88%, using the same patients, gene expression platform, and biomarker panel. It is expected that application of these panels to first-episode or prodromal subjects may improve the prediction for those subjects that ultimately convert to either

illness. This will require an additional validation study of the biomarker signatures with a larger cohort size, which was estimated using a power analysis as 310 subjects in a follow-on project for prediction of first-episode or prodromal patients.

(117) The expression differences of three genes related to polyunsaturated fatty acids (PUFAs) and prostaglandin biosynthesis was used in the final biomarker panels to differentiate schizophrenia, bipolar disorder, and controls (PTGDS, FADS2, HADHA). Previously these genes have been associated in psychiatric disorders such as bipolar disorder, major affective disorder, schizophrenia, and anxiety. PTGDS is involved in synthesis of PGD₂ from PGH₂, the cyclooxygenase-mediated product of arachidonic acid which is a PUFA (Begemann et al., *Mol Med* 2008; 14(9-10): 546-552). PTGDS is a top anxiety gene modulated by changes in PUFA (omega-3 fatty acid docosahexaenoic acid) (Le-Niculescu et al., *Transl Psychiatry* 2011; 1: e9) on the convergent functional genomics scale. Increased expression of FADS2 has been found in schizophrenia and bipolar disorder postmortem brain (Liu and McNamara, *Psychiatr Res* 2011; 45(2): 269-272; Liu et al., *Schizophr Res* 2009; 109(1-3): 113-120). FADS2 activity was increased in bipolar disorder and was associated with suicidal behavior (Evans et al., *PLoS One* 2012; 7(1): e29297). In the present study increased expression of FADS2 was found in bipolar disorder, in agreement with FADS2 findings reported. The increased activity of FADS2 could reduce PUFA levels of both AA and EPA by promoting conversion to longer chain fatty acids. Thus studies of PUFA supplementation in mood disorder with n-3 fatty acids was effective in 4 out of 7 well controlled studies in reducing mood symptoms (Evans et al., *PLoS One* 2012; 7(1): e29297). The expression data for FADS2, while interesting, could be subject to dietary influence, such as amounts and types of daily dietary intake of fatty acids, timing of intake, and also medication effects on these genes. Further, genetics plays a significant role, especially in modulating levels of fatty acids and FADS2 expression.

(118) From GWAS of schizophrenia and bipolar disorder, there is a large number of variants contributing to the polygenic susceptibility for these disorders. It is likely that we have highlighted several genes that contribute towards this susceptibility using transcription analysis, as some but not all polygenic effects such as expression quantitative trait loci might be consistent across blood and brain (Vawter et al., *Brief Funct Genomics* 2011; 10(6): 387-399; Mamdani et al., *Dis Markers* 2013; 35(1): 11-21) Additionally two upstream transcription factor merit further investigation, MKL2 and REL, which appear to oppositely regulate genes in BD and SZ, resulting in the ability to differentially diagnose these subjects. These proto-oncogenes have not been previously associated with psychiatric disorders,

(119) There was a consistent increase in expression of IL5RA (interleukin 5 receptor, alpha) in lithium treated subjects with bipolar disorder in PBMCs that passed strict FDR (Anand et al., *Mol Neuropsychiatry* 2016; 2(3): 115-123). In the final biomarker panel, a trend for an increase in IL5RA expression in BD (p-value for BD was 0.056 and fold change was 1.16) was found, however IL5RA expression in SZ was significantly decreased and passing FDR (p-value for SZ vs. NC, 2.65E-08, fold change -1.59). For PLB1 (phospholipase B1), in PBMC, lithium decreased expression (-1.17 fold change) passing FDR 82 in PBMCs, while in the present study, PLB1 was significantly decreased in BD by -1.19 and was not significantly increased in SZ (p=0.06, fold change 1.16). It is unlikely that some expression in the biomarker panel might be due to lithium treatments, since only 2 BD subject were lithium-treated in this study. Another ramification of using biomarkers is to provide a method to monitor drug efficacy and other appropriate early psychosocial interventions for mental disorders. In a separate analysis, the phenotypic neuropsychiatric ratings data obtained from each patient are correlated with the gene expression data.

(120) Over 100,000 adolescent Americans suffer from symptoms of psychosis each year and, currently, there are no biomarkers tests that are FDA approved to classify SZ or BD. There is a need for an 'objective' clinical laboratory test for an early diagnosis of these mental disorders since these disorders may take months or even years to arrive at a diagnosis and for patients to receive

effective treatments. The lag in treatment is associated with an increase in the suicide rate and recurrent episodes of psychosis and mood dysregulation. There is a large increase in deaths reported in first episode psychosis subjects due to lack of treatment after the first year of illness (Schoenbaum, Twelve-Month Health Care Use and Mortality in Commercially Insured Young People With Incident Psychosis in the United States. *Schizophrenia Bulletin* 2017). Thus, it is imperative to have objective biomarkers to help implement treatment at an early stage. One estimate of the direct and indirect annual costs in the USA for schizophrenia is \$174 billion (Cloutier et al., *J Clin Psychiatry* 2016; 77(6): 764-771) plus additional cost of \$151 billion for bipolar disorder (Dilsaver, *J Affect Disord* 2011; 129(1-3): 79-83). The biomarker signatures could lead to faster and more accurate diagnoses, reducing the duration of untreated psychosis, reduction in suicidality, reduction in cognitive decline and add to the understanding of shared and unique pathophysiologies of each disorder. The blood test results that are described offer molecular diagnostic support for a psychiatrist's clinical evaluation with rapid clinical laboratory test results. (121) The materials and methods employed in these experiments are now described.

(122) Subject Enrollment

(123) Subject enrollment occurred at a single clinical site at the University of Iowa in an institutional review board (IRB) approved study. Chronic SZ and BD type I outpatients ages 18-45, provided consent for the study. All subjects met DSM-IV-R criteria and completed the study: SZ (n=30), BD (n=30), and NC (n=30). Clinical assessments included the Scale for the Assessment of Positive Symptoms and Scale for the Assessment of Negative Symptoms (SAPS, SANS), medications and drugs for SZ and BD subjects; Young Mania Rating Scale (YMRS), and Hamilton Rating Scale for Depression (HAM-D or HRSD) for BD subjects. This neuropsychiatric assessment data will be analyzed and reported separately for state biomarker relationships. A mental status exam for normal controls (NC) consisted of the Mini-Mental Status Exam. The outline of the study is shown in FIG. 1.

(124) Whole blood samples were collected in Tempus Blood RNA tubes (Applied Biosystems) from SZ, BD, and NC subjects at 3 visits spanning 3 months. For this report, Tempus tubes from visits 2 and 4 were extracted and RNA gene expression was measured using Affymetrix exon arrays for both visits on all 90 subjects. High-quality RNA was extracted from the Tempus tubes using the manufacturer's protocol and quality was assessed on an Agilent Bioanalyzer using RNA Integrity Number (RIN).

(125) Human Exon Array for Biomarker Profile

(126) There are advantages to using the Affymetrix exon arrays (Xu et al., *Proc Natl Acad Sci USA* 2011; 108(9): 3707-3712) compared to whole transcriptome shotgun sequencing (RNA-Seq). At the time of sample collection, the cost factor was favorable for future clinical biomarker trials that would require hundreds of arrays compared to the cost of RNA-Seq for the entire validation. The processing time and data storage requirements are more favorable for a study this size using exon arrays. Although, previous studies have found that SNPs can affect probe hybridization and consequently alter expression (Benovoy et al., *Nucleic Acids Res* 2008; 36(13): 4417-4423; Fraser and Xie, *Genome Res* 2009; 19(4): 567-575; Gamazon et al, *PLoS One* 2010; 5(2): e9366; Gamazon et al., *Bioinformatics* 2010; 26(2): 259-262; Duan et al., *Am J Hum Genet* 2008; 82(5): 1101-1113), those probesets with common SNPs were eliminated from the final dataset. The exon arrays were run at the Functional Genomics Laboratory, University of California, using the manufacturer's protocol (Affymetrix, Santa Clara, CA). The Functional Genomics Laboratory has run over 1,000 Affymetrix arrays with high-quality call rates.

(127) Data Analysis

(128) The Affymetrix exon array GEL files were imported into Partek Genomics using batch effect removal. The batch effect was based upon exon array scan dates as usually 12 arrays were scanned in a single day. The mean intensity of probes was summarized at the probeset level. Probesets containing common SNPs were excluded from the import and downstream analysis. The resulting

probesets were then median centered within each exon array sample individually (n=180). A two-factor ANOVA was run for each probeset, using diagnosis, visit, and diagnosis x visit interaction. A false discovery rate of 6×10^{-8} was established for diagnosis effect based upon 835,000 probesets. Three filters were used to select probesets from the ANOVA results that passed FDR for diagnosis and were in the most significant p-values for BD compared to NC, BD compared to SZ, and SZ compared to NC. The resulting list of top probesets was filtered to reveal probesets mapping to known RefSeq genes. These probesets were ranked, and the top 300 probesets were evaluated for biomarker signature.

(129) Biomarker Signature

(130) The modeling proceeded in four steps to select the most predictive panel of probesets out of the top 300 in each step for discriminating between groups: Step 1: NC vs, BD+SZ Step 2: NC vs. SZ Step 3: NC vs. BD Step 4: SZ vs. BD

(131) Multivariate logistic regression modeling with forward stepwise selection (SAS PROC LOGISTIC) was used on the combined visit 2 and 4 data from the groups included in the step to select the probesets that were most discriminating between the groups. A probeset was added into the model if the estimate was the most significant with $p < 0.001$ and the resulting ROC AUC also retained statistical significance. Forward selection stopped when potential probesets were no longer statistically significant or did not appreciably improve the ROC AUC. Processing for each step resulted in a subset of the 300 probesets where each probeset contributed to the model significantly and the panel represented the smallest number of probesets that had very high diagnostic utility based on the ROC AUC.

(132) Modeling for the diagnostic for each step was applied to the Visit 2 data using the identified probesets. The optimal cut-point for discriminating between the groups based on the logistic model prediction was obtained by maximizing the Youden Index J 76, where $J = \text{True Positive Rate (TPR)} - \text{False Positive Rate (FPR)}$. The Visit 2 prediction model was then applied to the Visit 4 data to assess utility for a second set of data which included stability over time.

(133) Further validation for each of the four panels included “leave one out” cross-validation where one subject was sequentially left out of the logistic model fit using the remaining subjects and then the predictability of the model for the excluded subject was assessed.

(134) Quantitative PCR

(135) Transcripts were selected for qPCR validation based upon significant differences using the ANOVA-filter. Three filters were used to select probesets that represented a combination of the most significant ANOVA p-values for BD compared to NC, BD compared to SZ, and SZ compared to NC. Standard SybrGreen qPCR methods previously described by the Functional Genomics Laboratory (UC Irvine) were used to confirm gene expression derived from the exon array dataset (Morgan et al., Microarrays (Basel) 2016; 5(1)).

(136) NanoString Gene Expression Platform

(137) A non-PCR based approach of measuring RNA as technical validation of the findings was used based upon NanoString technology (NanoString, Seattle, WA) (Mamdani et al., Transl Psychiatry 2015; 5: e636). The Nanostring platform requires a small quantity of RNA and provides digital counts of hybridization of mRNA to targets. The NanoString multiplex assay uses 100 ng of total RNA and all of the RNA samples were processed at the UCI Genomics High-Throughput Facility. A total of 50 target genes were selected for technical validation, including 44 custom Nanostring probes designed to match the closest probeset on the Affymetrix exon array that was in the biomarker panel, and 6 Nanostring probes for housekeeping genes. The resulting NanoString data was processed according to manufacturer's suggested protocol as outlined. Each data point was preprocessed by the six positive controls, the eight negative controls and the five housekeeping genes as follows: i) Calculate for each subject/visit, the sum of the positive controls, the mean of the negative controls, and the sum of the housekeeping genes. ii) Calculate across all subjects/visits, the mean of the positive control sums, and the mean of the housekeeping gene

sums. iii) For each subject/visit, multiply the data point by (sum of positive controls/mean of positive control sums), subtract off the mean of the negative controls, divide by (sum of housekeeping genes/mean of housekeeping gene sums). If the normalized data point was negative, it was set to zero. The normalized Nanostring data were then analyzed for diagnostic accuracy using a model fitting approach. A potential gene probe set was selected by including all probe sets that had $p < 0.1$ in a univariate logistic regression fit of the probe set on at least one of the diagnosis comparisons (NC vs (SZ+BD), NC vs SZ, NC vs BD, SZ vs BD) for the visit 2 data.

(138) The Results of the Experiments are Now Described

(139) The top 300 probesets from the Affymetrix exon microarray based upon ANOVA significance were evaluated for biomarker signature (as described in Methods) for differentiating BD, SZ, and NC subjects. The resulting biomarker signature was composed of 23 probesets that condensed into 18 known RefSeq genes (biomarker panel Table 1). The diagnostic logistic model was built in four steps, using Visit 2 transcripts shown in Table 1. The resulting logistic predictive model based on Visit 2 was then applied to the Visit 4 data. The summary of individual steps in the construction of the biomarker gene panels are shown (Table 2).

(140) TABLE-US-00001 TABLE 1 Probesets that were found to reliably discriminate BD, SZ, and NC were assigned to known RefSeq transcripts. Biomarker Panel Affymetrix Exon Microarray (Comparisons) Transcript ID Gene BD-NC 2661992 OXTR BD-NC 3195034 PTGDS BD-NC 3333247 FADS2 BD-NC 4048241 HLA-DRB5 BD-SZ 2661992 OXTR BD-SZ 3333247 FADS2 BD-SZ 3554818 CRIP2 BD-SZ-NC 2418570 SLC44A5 BD-SZ-NC 2545092 HADHA BD-SZ-NC 2647109 CPA3 BD-SZ-NC 2661992 OXTR BD-SZ-NC 2739160 CCDC109B BD-SZ-NC 2906720 TREML4 BD-SZ-NC 3063536 TRIM4 BD-SZ-NC 3195034 PTGDS BD-SZ-NC 3667890 HPR BD-SZ-NC 3846538 EEF2 BD-SZ-NC 3908149 ZMYND8 SZ-NC 2401347 TCEA3 SZ-NC 2418570 SLC44A5 SZ-NC 2660617 IL5RA SZ-NC 3329099 GYLTL1B SZ-NC 3766893 DDX5

(141) The diagnostics algorithm uses a four step decision model: Step 1, BD and SZ vs. NC; Step 2, SZ vs. NC; Step 3, BD vs. NC; and Step 4, SZ vs. BD (Table 2).

(142) TABLE-US-00002 TABLE 2 Prediction model trained on Visit 2 and then applied to Visit 4. The confusion matrices show the numbers of subjects correctly and incorrectly classified for each iteration of the classification. Step 1 SZ & Step 2 Step 3 Step 4 Final Call Actual NC BD NC SZ NC BD SZ BD NC SZ BD Training Visit 2 30 NC 30 0 24 6 26 4 21 5 4 30 SZ 3 27 1 29 29 1 3 26 1 30 BD 1 29 1 29 1 29 1 1 28 Testing Visit 4 30 NC 28 2 23 7 28 2 22 6 2 30 SZ 1 29 0 30 29 1 1 28 1 30 BD 1 29 0 30 0 30 1 0 29 Step 1: 11 gene diagnostic differentiating BD + SZ vs. NC Step 2: 5 gene diagnostic differentiating SZ vs. NC Step 3: 4 gene diagnostic differentiating BD vs. NC Step 4: 3 gene diagnostic differentiating SZ vs. BD Final Call: Summary of Steps 1, 2, 3, and 4 classifications.

(143) TABLE-US-00003 TABLE 3 The overall results show a stable clinical biomarker signature of mRNA expression across a 90-day test-retesting period with an accuracy of 88% on the retesting data. Visit 2 Actual Visit 4 Actual Both Actual Predicted BD SZ NC BD SZ NC BD SZ NC BD 28 1 4 29 1 2 57 2 6 SZ 1 26 5 0 28 6 1 54 11 NC 1 3 21 1 1 22 2 4 43 Accuracy 83% 88% 86% Sensitivity (SZ) 87% 93% 90% Sensitivity (BD) 93% 97% 95% Specificity (NC) 70% 73% 72%

(144) The 18-gene biomarker panels, using logistic regression modeling, correctly differentiated the three groups of subjects: SZ (n=30), BD type I (n=30) and NC (n=30) with high accuracy at Visit 2 and Visit 4. The Visit 2 mRNA biomarker levels were significantly correlated with Visit 4 levels ($p < 0.0001$) showing temporal stability.

(145) The initial model was developed for selecting stable probesets across visits and incorporated all subjects and visits to select the most informative probesets. To test that no single subject was overly influential in determining the model, the initial probesets were evaluated in a "leave one out" method, whereby a new model is fit to the remaining subjects, and the left out subject is identified. "Leave one out" cross validation is a model validation technique for assessing how the

results of a statistical analysis will generalize to an independent data set. It is mainly used in settings where the goal is prediction to estimate how accurately a predictive model will perform in practice. This cross validation was applied to the Visit 2 data from each of the four probeset panels (Table 4).

(146) TABLE-US-00004 TABLE 4 The overall results fitting a leave-one out validation model to remaining subjects show a stable clinical biomarker signature of mRNA expression across a 90-day test-retesting period with an accuracy greater than 87% on the retesting data at each of the model classification steps. BD & SZ vs. NC SZ vs. NC BD vs. NC SZ vs. BD 11-Gene 5-Gene 4-Gene 3-Gene Panel Panel Panel Panel Actual Visit 2 Data Sensitivity (SZ) 93% 97% 97% Sensitivity (BD) 93% 97% 97% Specificity (NC) 100% 80% 87% Accuracy 96% 88% 92% 97% Leave One Out X- Validation Sensitivity (SZ) 90% 93% 90% Sensitivity (BD) 90% 93% 93% Specificity (NC) 83% 80% 80% Accuracy 88% 87% 87% 92%

(147) The results are very consistent between the actual data and the “leave one out” analyses indicating that the models should be predictive as applied to independent data cohorts (Table 4).

(148) The area under the curve for each step was greater than 0.95, which is an indication of high combined sensitivity and specificity of classification into three groups (Table 5). When analyzing the same 1.8-gene biomarker panel and including the white blood cell counts as a covariate, the analysis slightly improved the diagnostic predictability of the SZ vs. NC, and BD and SZ vs. NC.

(149) TABLE-US-00005 TABLE 5 The diagnostic algorithm uses four individual steps, shown in each column. BD & SZ vs. NC SZ vs. NC BD vs. NC SZ vs. BD 11-Gene Panel 5-Gene Panel 4-Gene Panel 3-Gene Panel *AUC = 0.994 AUC = 0.954 AUC = 0.974 AUC = 0.998 (p < 0.0001) (p < 0.0001) (p < 0.0001) (p < 0.0001) *area under the curve (AUC) of the receiver operating characteristic (ROC)

(150) To account for potential medication effects (since many of the SZ and BD subjects were on stable dosages of antipsychotic or mood stabilizer medications at the time of blood draw) the possibility that these medications could drive some of the differential transcription signatures was analyzed. A subset of non-medicated BD (n=3) and SZ (n=1) cases were analyzed using the 18-gene diagnostics, which produced 100% accuracy in these four cases. In addition, to test this possible explanation of the gene signature, an even larger cohort of antipsychotic-free SZ patients was re-analyzed using the Illumina gene expression microarray datasets (de Song et al., PLoS One 2012; 7(6): e39498). The genome-wide RNA expression profiling was obtained with the Illumina HumanRef-8 V3 arrays for batch 1 and HumanRef-12 V3 arrays for batch 2 using Illumina's standard protocol at UCLA's Illumina facility. The raw microarray data were available at gene expression omnibus (GEO) under accession GSE38485. Dataset includes schizophrenia patients on antipsychotic-free (n=15) and healthy controls (n=22). The 18 gene signature was extracted from the Illumina dataset and analyzed by the same logistic regression equations derived from the fit to the original data. Using this independent Illumina dataset of gene expression data from antipsychotic-free SZ subjects, the 18-gene diagnostics produced a ROC accuracy of 0.866 in patients less than 30 years of age and medication free (Table 6). These results were achieved with different gene expression technology, different blood collection tubes, and different mRNA extractions technique. The details on the methods and analysis of the antipsychotic-free patients are in Example 8.

(151) TABLE-US-00006 TABLE 6 The 18 gene signature was extracted from an independent dataset of gene expression and analyzed by the same logistic regression equations derived from the fit to the original data. Using this dataset from antipsychotic-free SZ subjects, the 18-gene diagnostics produced a ROC accuracy of 0.866 in patients less than 30 years of age and medication free. Classification of Antipsychotic free patients with SZ using data from Illumina HT-8 array NC SZ Subjects (N) (N) ROC AUC Sensitivity Specificity Antipsychotic-free, 22 15 0.642 80.00% 59.10% all subjects Antipsychotic-free, 14 8 0.866 100.00% 71.40% subjects ages <30 QPCR Validation of Exon Array

(152) Transcripts were validated for schizophrenia and controls using qPCR. Those selected transcripts are shown in Table 7.

(153) TABLE-US-00007 TABLE 7 QPCR results for candidate gene expression differences in schizophrenia and controls. p-value Exon p-value (SZ Ratio (SZ (SZ Ratio (SZ vs. Array vs. NC) vs. NC) qPCR vs. NC) NC) Gene Symbol EDIL3 1.04E-13 0.446 ED1L3 0.01418 0.425 NRCAM 1.37E-02 0.525 NRCAM 0.06127 0.663 PTGDS 1.41E-14 0.722 PTGDS 0.04731 0.695 DSC2 6.86E-04 1.592 DSC2 0.00005 2.021 NRG1 1.14E-02 2.062 NRG1 0.01021 1.979 ITGA2B 3.71E-02 2.088 ITGA2B 0.00007 2.269 ITGB3 2.65E-02 2.601 ITGB3 0.00005 2.065

(154) The fold change ratio (SZ compared to NC) for exon microarray results were compared to fold change ratio (SZ compared to NC) of qPCR. The fold changes were highly correlated (FIG. 2).

(155) Nanostring Platform Accuracy for Diagnostic Classification

(156) TABLE-US-00008 TABLE 8 The Nanostring data set was analyzed by ANOVA, and the following 23 probe sets representing 11 genes passed the initial ANOVA filter ($p < 0.1$). 11 genes passed the initial ANOVA using Nanostring *2537112_SH3YL1 3554838_CRIP2 (39)

2661997_OXTR (43) 2723770_TBC1D1 (30) (55) 2537128_SH3YL1 3554839_CRIP2 (58) 2906726_TREML4 4048243_HLA-DRB5 (61) (45) (57) 2647127_CPA3 3554833_CRIP2 (68) 2906736_TREML4 3195045_PTGDS (35) (50) (59) 2647122_CPA3 2418615_SLC44A5 2906735_TREML4 2401364_TCEA3 (47) (40) (53) (64) 2647124_CPA3 2418581_SLC44A5 2906733_TREML4 2401362_TCEA3 (65) (46) (62) (70) 2647119_CPA3 2418590_SLC44A5 3063538_TRIM4 (69) (51) (52) *Format is Affymetrix Exon Array Probeset ID_Gene

Symbol_V Variable Number Entered Into Regression

(157) For Nanostring data, the best multivariate logistic regression model for diagnosis at visit 2 was obtained by stepwise backward elimination from the full 23 probe set multivariate model to a reduced model with all included probe sets significant with $p < 0.05$ or $p < 0.1$ to maintain $AUC > 0.9$. The cut point for each reduced diagnostic model was found which optimized both sensitivity and specificity based on the visit 2 data, this reduced the useful probe set number to 12 (Table 9). The diagnostic model and cut point were then applied to the visit 4 data to estimate the visit 4 sensitivity and specificity and the agreement between the visit 2 and visit 4 diagnostic predictions.

(158) TABLE-US-00009 TABLE 9 Summary of Nanostring Cut Point Diagnostic Accuracy NC vs SZ/BD NC vs SZ NC vs BD SZ vs BD Sig Probe sets 2537112_SH3YL1 (30) x X x x 2537128_SH3YL1 (61) X x x 2647124_CPA3 (65) x X x x 2647119_CPA3 (69) x X x x 3554833_CRIP2 (68) x x 2418590_SLC44A5 (51) x x 2661997_OXTR (43) x x 2906733_TREML4 (62) x 3063538_TRIM4 (52) x x x 2723770_TBC1D1 (55) x 3195045_PTGDS (59) x x x x 2401362_TCEA3 (70) x AUC 0.913 0.990 0.953 0.905 Visit 2 Sensitivity 98.3% 96.6% 89.3% 86.2% Visit 2 Specificity 75.0% 96.4% 92.9% 82.1% Visit 4 Sensitivity 87.5% 75.0% 75.0% 57.1% Visit 4 Specificity 60.0% 80.0% 76.0% 85.7% Visit 2 4 Agreement 77.6% 76.0% 75.5% 64.2%

Comparison of Affymetrix Exon Array and Nanostring Results

(159) The prediction accuracies for diagnosis were compared between the Affymetrix exon array and Nanostring platform. In general, for the exon array, the model performed equally well for Visit 2 and Visit 4 data due to the data driving the selection algorithm from the original platform. The Visit 2 and Visit 4 agreement for the Affymetrix exon array was significantly higher than Nanostring reproducibility diagnosis classification (two tailed paired t-test, $p = 0.046$). These results show that the test-retest reproducibility of the algorithm using the same subjects was best when using the original platform.

(160) TABLE-US-00010 TABLE 10 Comparison of Affymetrix and Nanostring results showed that Affymetrix exon array had higher sensitivity and specificity for prediction. Genes Included in Each Predictive Model Nanostring NC vs SZ/BD NC vs SZ NC vs BD SZ vs BD Probe Set Affy Affy Affy Affy IDs ID Nanostring ID Nanostring ID Nanostring ID Nanostring 2537112_SH3YL1 x x x x (30) 2537128_SH3YL1 x x x (61) 2647124_CPA3 (65) x x x x 2647119_CPA3 (69) 2647109 x x

x x 3554833_CRIIP2 (68) x x 3554818 2418590_SLC44A5 2418570 x 2418570 x (51)
 2661997_OXTR (43) 2661992 2661992 x 2661992 x 2906733_TREML4 2906720 x (62)
 3063538_TRIM4 (52) 3063536 x x x 2723770_TBC1D1 x (55) 3195045_PTGDS (59) 3195034 x
 x 3195034 x x 2401362_TCEA3 (70) 2401347 x 3667896, 97_HPR 3667890 3908171_ZMYND8
 3908149 2545100_HADHA 2545092 2739191_CCDC109B 2739160 3846545_EEF2 3846538
 3333251, 56, 58, 62, 3333247 3333247 69, 70, 74_FA DS2 4048243, 4048241 52_HLA-DRB5
 3329128_GYLTL1B 3329099 2660633, 41_IL5RA 2660617 3766938_DDX5 3766893 *AUC
 0.994 0.913 0.954 0.990 0.974 0.953 0.998 0.905 Visit 2 93.3% 98.3% 96.7% 96.6% 96.7% 89.3%
 96.7% 86.2% Sensitivity Visit 2 100% 75.0% 83.3% 96.4% 86.7% 92.9% 100% 82.1%
 Specificity Visit 4 96.7% 87.5% 100% 75.0% 93.3% 75.0% 96.7% 57.1% Sensitivity Visit 4
 93.3% 60.0% 80% 80.0% 100.0% 76.0% 100% 85.7% Specificity Visit 2 4 93.3% 77.6% 86.7%
 76.0% 88.3% 75.5% 100% 64.2% Agreement *AUC was for combined visit 2 and 4 modeling and
 the each visit was evaluated for predictiveness individually

Bioinformatics Analysis of Biomarker Panel

(161) A recent mega-analysis of differentially expressed genes in SZ across 9 studies was conducted in blood based transcriptomics (Chan et al., Transl Psychiatry 2015; 5: e601). There were 1624 genes that survived FDR that were compared to the top 122 genes identified by ANOVA in the present study. The overlap between studies was not enriched, as 8.7 genes were expected and 9 genes were observed. Interestingly, two mitochondria genes were found in these top 9 and agreed in fold change direction across both studies (Table 11), mitochondrial ribosomal protein L42 (MRPL42), and transcription factor B1, mitochondrial (TFB1M).

(162) TABLE-US-00011 TABLE 11 Overlap with mega-analysis of blood dysregulated genes (Chan et al., Transl Psychiatry 2015; 5: e601) and current study. P- FDR value (SZ Ratio (SZ Gene Mean q- Gene vs. vs. Symbol Gene Product Difference* Value Symbol NC) NC) FAM118A family with -0.37 5.90E-04 FAM118A 4.17E-13 0.58 sequence similarity 118, member A MRPL42 mitochondrial 0.27 1.90E-02 MRPL42 0.000535 1.24 ribosomal protein L42 PHF14 PHD finger protein 0.26 2.50E-02 PHF14 3.09E-14 0.49 14 PHIP pleckstrin 0.29 9.10E-03 PHIP 4.66E-29 0.59 homology domain interacting protein PLB1 phospholipase B1 0.26 3.40E-02 PLB1 2.70E-13 1.52 SLC22A4 solute carrier 0.34 3.30E-03 SLC22A4 0.039619 1.14 family 22 (organic cation/zwitterion transporter), member 4 STX2 syntaxin 2 0.28 1.30E-02 STX2 1.49E-05 0.81 TFB1M transcription -0.5 3.60E-02 TFB1M 1.11E-05 0.79 factor B1, mitochondrial TNFRSF21 tumor necrosis -0.31 3.70E-03 TNFRSF21 1.96E-13 0.53 factor receptor superfamily, member 21

(163) An IPA analysis of the top 122 genes for the BD and SZ biomarker panel showed some overlap in a network related to two different proto-oncogenes, REL and MKL2. The REL gene (REL proto-oncogene, NF-kB subunit) encodes a protein that belongs to the Rel homology domain/immunoglobulin-like fold, plexin, transcription factor (RI-ID/IPT) family. This proto-oncogene plays a role in the survival and proliferation of B lymphocytes. Single nucleotide polymorphisms in this gene are associated with susceptibility to ulcerative colitis and rheumatoid arthritis. The direct REL gene targets were differentially expressed in BD and SZ (FIG. 3).

(164) As examples, the opposite fold change genes related to REL were seen in BD and SZ, ATP11A was upregulated in SZ (1.68) and down-regulated in BL) (-1.63), while an opposite relationship was seen for PHF14 downregulated in SZ (-2.01) and upregulated in BD (2.95).

(165) MKL2 is a proto-oncogene, which is widely distributed, highly expressed in brain regions such as dentate gyrus (Allen Institute Brain Science (Hawrylycz et al., Nature 2012; 489(7416): 391-399)) and associated with microcephaly (Ramos et al., (Clin Genet 2014; 85(5): 423-432). The MKL2 gene consists of multiple exons, many of those exons are down-regulated in SZ and not changed significantly in BD. The exon levels in blood for MKL2 were significantly differentially expressed in SZ, although not in the top 300 genes (FIG. 4). The downstream effects on MKL2 regulated genes in the top biomarker classification genes were in the expected directions, e.g. there were decreased fold changes seen in genes regulated by MKL2 in SZ (ACTA2, FADS2, CPA3),

while the same set was up-regulated in BD.

(166) Two additional genes in the biomarker panels PTGDS (prostaglandin D2 synthase) and FADS2 (fatty acid desaturase 2) were also found to be candidate genes in the literature. PTGDS expression was reduced in BD PBMC (Munkholm et al., *Int J Neuropsychopharmacol* 2014; 18(5):pyu101), and also in the present study (p =fold change -1.23) and also down in SZ ($p=9.04E-12$, fold change -1.52). In BD and SZ there appears to be a down regulation of PTGDS, which also was ranked high in the convergent functional genomics paradigm for anxiety (Le-Niculescu et al., *Transl Psychiatry* 2011; 1: e9).

(167) FADS2 expression was decreased in schizophrenia in this study (p -value $9.20E-09$, fold change -1.80) and increased in BD (p -value $2.79E-06$, fold change $=1.6$). The FADS2 and PTGDS genes work in the biosynthesis of fatty acids pathway, and converge on key molecules in BD such as arachidonic acid. HADHA was significantly increased in BD ($p=4.74E-08$, fold change 1.67) and was not changed in SZ. HADHA is closely related in the fatty acid pathway with FADS2 and HADHA along with PTGDS participating in biosynthesis and degradation of unsaturated fatty acids, an important pathway implicated in BD and SZ.

Example 2: BD-SZ-NC Comparison

(168) These results demonstrate the identification of a gene panel to discriminate bipolar and schizophrenic subjects from normal controls from 122 top known genes. Forward stepwise variable selection with logistic regression modeling was used to identify a set of genes which would significantly differentiate the combined bipolar and schizophrenic subjects from the normal controls.

(169) Table 12 includes the set of 11 genes that are diagnostic for this purpose.

(170) TABLE-US-00012
TABLE 12 Marker Transcript ID Gene
77 3667890 HPR 5 2906720
TREML4 52 3908149 ZMYND8 36 3195034 PTGDS 100 2647109 CPA3 82 3063536 TRIM4 41
2418570 SLC44A5 97 2661992 OXTR 43 2545092 HADHA 59 2739160 CCDC109B 117
3846538 EEF2

(171) Table 13 includes the model fit for visits 2 and 4.

(172) TABLE-US-00013
TABLE 13 Visits 2 and 4 AIC Intercept only 231.145 AIC with genes
57.157 Estimate P-value intercept β .sub.0 -66.4124 0.0048 marker 77 coefficient β .sub.1 -5.2531
0.0001 marker 5 coefficient β .sub.2 2.0403 0.0008 marker 52 coefficient β .sub.3 -3.0311 0.0010
marker 36 coefficient β .sub.4 8.9782 0.0003 marker 100 coefficient β .sub.5 4.9250 0.0008 marker
82 coefficient β .sub.6 -2.4530 0.0032 marker 41 coefficient β .sub.7 3.1667 0.0020 marker 97
coefficient β .sub.8 2.3342 0.0030 marker 43 coefficient β .sub.9 2.3378 0.0007 marker 59
coefficient β .sub.10 1.9140 0.0016 marker 117 coefficient β .sub.11 -7.0097 0.0019 ROC AUC
0.995 <0.0001

(173) $\text{Prob of Normal Control from Logistic} = \frac{1}{1 + \exp[-(\beta_0 + \beta_1 M_{77} + \beta_2 M_{52} + \beta_3 M_{52} + \beta_4 M_{36} + \beta_5 M_{100} + \beta_6 M_{82} + \beta_7 M_{41} + \beta_8 M_{97} + \beta_9 M_{43} + \beta_{10} M_{59} + \beta_{11} M_{117})]}$

(174) Optimal cut-point for discriminating two populations is Schiz or Bipolar if $\text{Prob(NC) from Logistic} \leq 0.1518$

(175) At this cut-point, across both visits Sensitivity=93.33 (95% CI 87.3–97.1) Specificity=100.00 (95% CI 94.0–100)

(176) FIG. 6 shows the ROC curve for visits 2 and 4.

(177) Table 14 includes data using the 0.1518 cut-point for a positive test.

(178) TABLE-US-00014
TABLE 14 Visit 2 Visit 4 Gene Model NC BD/SZ NC BD/SZ ≤ 0.1518 0
28/26 0 29/29 >0.1518 30 2/4 30 1/1

(179) FIG. 7 shows the logistic model estimated probabilities for each group, for each visit along with the reference line for the 0.1518 cut-point. The visit 2 probabilities are significantly correlated with the visit 4 probabilities ($p<0.0001$) with $r=0.89$ (95% CI 0.83-0.92).

(180) Table 15 includes data demonstrating that 93% of the subjects (84/90) received the same

diagnostic outcome from visit 2 to visit 4.

(181) TABLE-US-00015 TABLE 15 Visit 4 All Visit 2 ≤ 0.1518 > 0.1518 < 0.1518 53 1 > 0.1518 5
31 NC Visit 2 ≤ 0.1518 > 0.1518 ≤ 0.1518 > 0.1518 30 BD Visit 2 ≤ 0.1518 > 0.1518 ≤ 0.1518 27 1
 > 0.1518 2 SZ Visit 2 ≤ 0.1518 > 0.1518 ≤ 0.1518 26 > 0.1518 3 1

Example 3: BD-NC Comparison

(182) These results demonstrate the identification of a gene panel to discriminate bipolar subjects from normal controls from 122 top known genes. Forward stepwise variable selection with logistic regression modeling was used to identify a set of genes which would significantly differentiate the bipolar subjects from the normal controls.

(183) The set of 4 genes that may be diagnostic for this purpose are listed in Table 16.

(184) TABLE-US-00016 TABLE 16 Marker Transcript ID Gene 97 2661992 OXTR 76 4048241
HLA-DRB5 50 3333247 FADS2 36 3195034 PTGDS

(185) Table 17 includes the model fit for visits 2 and 4.

(186) TABLE-US-00017 TABLE 17 Visits 2 and 4 AIC Intercept only 168.355 AIC with genes
45.986 Estimate P-value intercept $\beta_{\text{sub.0}}$ -39.9111 0.0006 marker 97 coefficient $\beta_{\text{sub.1}}$ 5.4440
 < 0.0001 marker 76 coefficient $\beta_{\text{sub.2}}$ -0.6870 0.0012 marker 50 coefficient $\beta_{\text{sub.3}}$ -3.7201
0.0010 marker 36 coefficient $\beta_{\text{sub.4}}$ 3.5108 0.0026 ROC AUC 0.987 < 0.0001

(187) $\text{ProbofNormalfromLogistic} = \frac{1}{1 + \exp[-(\beta_0 + \beta_1 M_{97} + \beta_2 M_{76} + \beta_3 M_{50} + \beta_4 M_{36})]}$

(188) Optimal cut-point for discriminating two populations is BD if Prob(NC) from Logistic ≤ 0.659

(189) At this cut-point, across both visits Sensitivity=98.33 (95% CI 91.1-100) Specificity=90.00 (95% CI 79.5-96.2)

(190) The ROC Curve for visits 2 and 4 is shown in FIG. 5.

(191) Table 18 shows the data using the 0.659 cut-point for a positive test.

(192) TABLE-US-00018 TABLE 18 Visit 2 Visit 4 Gene Model NC BD NC BD ≤ 0.659 4 29 2 30
 > 0.659 26 1 28 0

(193) Visit specific metrics:

(194) TABLE-US-00019 Sensitivity 0.97 1.00 Specificity 0.87 0.93 Accuracy 0.92 0.97

(195) FIG. 8 shows the logistic model estimated probabilities for each group, for each visit along with the reference line for the 0.659 cut-point. The visit 2 probabilities are significantly correlated with the visit 4 probabilities ($p < 0.0001$) with $r = 0.86$ (95% CI 0.78-0.92).

(196) 92% of the subjects (55/60) received the same diagnostic outcome from visit 2 to visit 4 (Table 19).

(197) TABLE-US-00020 TABLE 19 Visit 4 Visit 2 ≤ 0.659 > 0.659 ≤ 0.659 30 3 > 0.659 2 25

Example 4: BD-SZ Comparison

(198) These results demonstrate the identification of a gene panel to discriminate bipolar subjects from schizophrenic subjects from 122 top known genes. Forward stepwise variable selection with logistic regression modeling was used to identify a set of genes which would significantly differentiate the bipolar subjects from the schizophrenic subjects.

(199) The set of 3 genes that may be diagnostic for this purpose are listed in Table 20. Markers 97 and 50 were also identified in the BD-NC panel.

(200) TABLE-US-00021 TABLE 20 Marker Transcript ID Gene 37 3554818 CRIP2 97 2661992
OXTR 50 3333247 FADS2

(201) Table 21 includes the model fit for visits 2 and 4.

(202) TABLE-US-00022 TABLE 21 Visits 2 and 4 AIC Intercept only 168.355 AIC with genes
24.662 Estimate P-value intercept $\beta_{\text{sub.0}}$ -28.7571 0.0403 marker 37 coefficient $\beta_{\text{sub.1}}$ 7.3345
0.0123 marker 97 coefficient $\beta_{\text{sub.2}}$ -8.5153 0.0054 marker 50 coefficient $\beta_{\text{sub.3}}$ 4.1472 0.0044
ROC AUC 0.996 < 0.0001

$$(203) \text{ ProbofBipolarfromLogistic} = \frac{1}{1 + \exp[-(\frac{1}{2}M_{97} + \frac{1}{3}M_{50})]}$$

(204) Optimal cut-point for discriminating two populations is Schiz if Prob(BD) from Logistic ≤ 0.2857

(205) At this cut-point, across both visits Sensitivity=96.67 (95% CI 88.5-99.6) Specificity=100 (95% CI 94-100)

(206) FIG. 9 includes the ROC curve for visits 2 and 4.

(207) Table 22 includes data for using the 0.2857 cut-point for a positive SZ test.

(208) TABLE-US-00023 TABLE 22 Visit 2 Visit 4 Gene Model BD SZ BD SZ ≤ 0.2857 0 29 0 29
 > 0.2857 30 1 30 1

(209) Visit specific metrics: Visit 2 Visit 4

(210) TABLE-US-00024 Sensitivity (SZ) 0.97 0.97 Specificity (BD) 100 100 Accuracy 0.98 0.98

(211) FIG. 10 shows the logistic model estimated probabilities for each group, for each visit along with the reference line for the 0.2857 cut-point. The NC subjects are added in using the predictive model for the BD vs SZ. The visit 2 probabilities are significantly correlated with the visit 4 probabilities ($p < 0.0001$) with $r = 0.96$ (95% CI 0.94-0.98).

(212) Table 23 includes data demonstrating that 100% of the subjects (60/60) received the same diagnostic outcome from visit 2 to visit 4.

(213) TABLE-US-00025 TABLE 23 Visit 4 Visit 2 ≤ 0.2857 > 0.2857 ≤ 0.2857 29 0 > 0.2857 0 31
 Leave One Out Cross Validation

(214) Each subject was sequentially removed, the logistic model was fit with the remaining 59 subjects and the model fit to predict the subject who was left out was used. (Table 24). The leave-one-out model used for this analysis has higher accuracy because the model was based upon combined Visit 2 and Visit 4 data, then run on each individual visit.

(215) TABLE-US-00026 TABLE 24 V2 Actual V4 Actual Both Actual Predicted BD SZ BD SZ
 BD SZ BD 27 1 30 1 57 2 SZ 3 29 0 29 3 58 Accuracy 93% Accuracy 98% Accuracy 96%
 Sensitivity (SZ) 97% Sensitivity (SZ) Sensitivity (SZ) 97% 97% Specificity (BD) 90% Specificity (BD) Specificity (BD) 100% 95%

Example 5: Analyses of Un-Normalized Gene Expression Data

(216) TABLE-US-00027 TABLE 25 (BD + SZ) vs. NC Comparison (visits 2 and 4) Parameter Estimate Std Err P-value Intercept 44.9621 10.2406 $< .0001$ TREML4 -0.6979 0.2658 0.0087
 PTGDS -4.3427 0.9990 $< .0001$ SLC44A5 -1.9189 0.4810 $< .0001$ HADHA -0.9526 0.3386
 0.0049 ZMYND8 0.8177 0.3431 0.0171 GYLTL1B 3.2553 0.7704 $< .0001$ CCDC109B -1.0542
 0.2868 0.0002 HPR 2.5108 0.5648 $< .0001$ TRIM4 2.1397 0.5594 0.0001 OXTR -2.5848 0.6609
 $< .0001$ CPA3 -1.1792 0.4616 0.0106 ROC AUC 0.973 P(model) = 0.0005

(217) TABLE-US-00028 TABLE 26 BD vs. NC Comparison (visits 2 and 4) Parameter Estimate Std Err P-value Intercept 39.6860 9.1607 $< .0001$ OXTR -2.7173 0.6667 $< .0001$ PTGDS -3.2692
 0.8294 $< .0001$ TREML4 -0.6789 0.2640 0.0101 SLC44A5 -1.9348 0.4797 $< .0001$ GYLTL1B
 1.3459 0.6449 0.0369 HPR 1.4963 0.4466 0.0008 TRIM4 1.6745 0.4938 0.0007 ROC AUC 0.951
 P(model) = 0.0007

(218) TABLE-US-00029 TABLE 27 SZ vs. NC Comparison (visits 2 and 4) Parameter Estimate Std Err P-value Intercept -13.4445 6.7739 0.0472 TREML4 0.5711 0.2415 0.0181 PTGDS 3.1815
 0.9906 0.0013 SLC44A5 1.7258 0.5215 0.0009 GYLTL1B -3.1913 0.9423 0.0007 HPR -1.7817
 0.4715 0.0002 TRIM4 -1.3580 0.4218 0.0013 CPA3 1.8422 0.5090 0.0003 ROC AUC 0.952
 P(model) = 0.0003

(219) TABLE-US-00030 TABLE 28 SZ vs. BD Comparison (visits 2 and 4) Parameter Estimate Std Err P-value PTGDS 2.5861 0.7034 0.0002 SLC44A5 1.9851 0.5742 0.0005 GYLTL1B
 -3.5039 0.9396 0.0002 HPR -2.2558 0.5812 0.0001 TRIM4 -1.9102 0.4765 $< .0001$ CPA3 1.5558
 0.4348 0.0003 CCDC109B 0.8509 0.2720 0.0018 ROC AUC 0.961 P(model) = 0.0006

(220) TABLE-US-00031 TABLE 29 Old and New Model Fit Comparison New Un-normalized Gene Old Normalized Gene Expression Expression SZ BD BD SZ BD BD BDSZ vs vs vs BDSZ vs vs vs vs NC NC NC SZ vs NC NC NC SZ HPR X X X X X TREML4 X X X X ZMYND8 X X PTGDS X X X X X CPA3 X X X X TRIM4 X X X X X SLC44A5 X X X X X OXTR X X X X X HADHA X X CCDC109B X X X GYLTL1B X X X X X EEF2 X TCEA3 X CRIP2 X FADS2 X X IL5RA X HLA_DRB5 X DDX5 X AUC 0.973 0.952 0.951 0.961 0.995 0.967 0.987 0.996 P(Model) 0.0005 0.0003 0.0007 0.0006 <0.0001 <0.0001 <0.0001 <0.0001

Example 6: Analysis of Illumina Data for Neuroleptic-Free Subjects and Other Analyses

(221) Table 30 includes data of Illumina SZ genes, logistic regression fit to data, subgrouped by age <30 and/or neuroleptic-free status.

(222) SZ genes: TREML4, PTGDS, SLC44A5, GYLTL1B, HPR, TRIM4, CPA3

(223) Second HPR probe was used because first has missing data. First TRIM4 probe was used as it fit better than second AUC.

(224) TABLE-US-00032 TABLE 30 ROC Illumina SZ genes, logistic NC SZ regression Subjects N N fit to data, Sensitivity Specificity Cutpoint All Illumina Subjects 118 121 0.686 84.3% 36.4% 0.423 Illumina Subjects 44 31 0.782 58.1% 90.9% 0.551 with Age <30 Neuroleptic-free 22 15 0.642 80.0% 59.1% 0.364 Illumina Subjects Neuroleptic-free, 14 8 0.866 100.0% 71.4% 0.308 Age <30 Subjects

(225) Table 31 includes data from the first “All Illumina Subjects” analysis, the predicted classification of the neuroleptic-free subjects.

(226) TABLE-US-00033 TABLE 31 Actual SZ Actual NC N = 15 N = 22 Predicted NC 8 18 Predicted SZ 7 4 Sensitivity = 46.7% Specificity = 81.8%

(227) Logistic Model with 7 SZ genes, age indicator (<30, ≥30), neuroleptic-free (Y,N) ROC AUC=0.702 Sensitivity=64.5% Specificity=69.5% Significance of age indicator term in model p=0.1163 Significance of neuroleptic-free term in model p=0.0362

(228) 7 SZ genes from “New” panel were compared with all 10* genes from “New” panel vs. 17* genes from “Old” panel (Table 32). 7 SZ genes from New panel: TREML4, PTGDS, SLC44A5, GYLTL1B, HPR, TRIM4, CPA3 Additional 3 genes from New panel: ZMYND8, OXTR, HADHA Additional 7 genes from Old panel: EEF2, TCEA3, CRIP2, FADS2, IL5RA (2 different probes included), HLA-DRB5, DDX5 *CCDC109B which was included on both new and old panels is not available on Illumina dataset

(229) Note: used second HPR probe as first has missing data, used first TRIM4 probe as better fitting than second.

(230) TABLE-US-00034 TABLE 32 NC SZ ROC Subjects Gene Set N N AUC Sensitivity Specificity Cutpoint All 7 SZ from 118 121 0.686 84.3% 36.4% 0.423 Subjects New All All 10 from 118 121 0.695 81.8% 30.5% 0.406 Subjects New All All 17 from 118 121 0.731 59.5% 80.5% 0.574 Subjects Old NL Free 7 SZ from 22 15 46.7% 81.8% Subj New NL Free All 10 from 22 15 46.7% 72.7% Subj New NL Free All 17 from 22 15 20.0% 81.8% Subj Old

Example 7: Comparison of SZ Genes from 11-Gene Panel vs 18-Gene Panel in Original Median-Normalized Affymetrix Exon Array Data and Performance in Gender, Age Subgroups

(231) SZ genes From 18 gene panel: SLC44A5, GYLTL1B, TCEA3, IL5RA, DDX5 From 11-gene panel: HPR, TREML4, PTGDS, CPA3, TRIM4, SLC44A5

(232) TABLE-US-00035 TABLE 33 Comparison of SZ genes from 11-gene panel vs 18-gene panel Group (visits 2 NC SZ ROC Cut- Panel and 4 combined) N N AUC Sensitivity Specificity point 18- All subjects 60 60 0.967 90.0% 88.3% 0.499 gene Male, <30 yrs 16 10 0.913 90.0% 81.3% 0.572 Male, ≥30 yrs 28 34 0.993 97.1% 100.0% 0.411 Female 16 16 0.996 100.0% 93.8% 0.659 11- All subjects 60 60 0.975 93.3% 90.0% 0.549 gene Male, <30 yrs 16 10 0.944 90.0% 81.3% 0.774 Male, ≥30 yrs 28 34 0.989 97.1% 89.3% 0.574 Female (only 16 16 0.926 100.0% 75.0% 0.694 HPR and CPA3) A better diagnostic was not observed in young males as seen in Illumina data (not enough females to split by age)

(233) The two panels were very close, with slightly better diagnostic in all subjects with 11-gene panel.

Example 8: Independent Study of Biomarker Signature Validation

(234) The de Jong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498) results provide an independent validation of the SZ signature, with a high ROC shown in subjects with schizophrenia <30 years of age and in antipsychotic-free subjects with schizophrenia <30 years of age. These results suggest that the signature can be analyzed in young patients that are medication free at the time of testing as well as patients being treated with antipsychotic medications at the time of testing.

(235) The Materials and Methods are Now Described

(236) The initial whole blood exon array signature from Example 1 was tested against an independent dataset referred to in deJong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498) and made available under GEO dataset omnibus accession GSE38485.

(237) deJong et al. applied a systems biology approach to genome-wide expression data from whole blood of 92 medicated and 29 antipsychotic-free schizophrenia patients and 118 healthy controls. They showed that gene expression profiling in whole blood can identify twelve large gene co-expression modules associated with schizophrenia.

(238) The Results are Now Described

(239) The three datasets described in Table 33 reported by deJong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498) were independently reanalyzed for the studies presented herein. Dataset 1 includes schizophrenia patients on antipsychotics (n=92) and healthy controls (n=78). Dataset 2 consists of n=29 antipsychotic-free schizophrenia patients and n=40 healthy controls. The demographic information for both datasets is given in Table 34.

(240) TABLE-US-00036 TABLE 34 Description of datasets from de Jong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498). Schizophrenia dataset Antipsychotic-free dataset Controls
Cases Controls Cases Total 78 92 40 29 Mean age 41 yrs 41 yrs 30 yrs 31 yrs Gender 31M, 47F 66M, 26F 27M, 13F 21M, 8F Batch 1 22 15 Batch 2 78 92 18 14 Country* 22 DK, 92 NL 6 DK, 6 DK, 56 NL 34 NL 23 NL Expression Illumina H-12 Illumina H-8 & H-12 Array (16,707 genes) (12,704 genes)

Two datasets are shown, schizophrenia cases and controls, and antipsychotic-free schizophrenia and control dataset. Age and gender information is given for cases and controls separately for antipsychotic free subjects in Table 34. Gene expression data was generated in two batches (batch 1: Illumina H-8 and batch 2: Illumina H-12) and collected at different sites, information given in the fourth and fifth row). The number of expressed genes is given in the last row. *DK=Denmark and ML=The Netherlands.

Subjects in de Jong et al., 2012

(241) Participants were recruited from three sources: i) the Department of Psychiatry of the University Medical Center Utrecht (90 controls and 113 cases), ii) Parnassia PsychoMedical Center in the The Netherlands (2 cases) and iii) the Center for Neuropsychiatric Schizophrenia Research, Psychiatric Center Glostnip, Denmark (28 controls and 6 cases). Diagnoses were determined by Standardized Psychiatric interviews either The Comprehensive Assessment of Symptoms and History (CASH) or the Composite international diagnostic interview (CIDI) by trained clinicians. Schizophrenia was defined by a DSM-IV-TR diagnosis of #295.0-295.89, and #298.9.

(242) Antipsychotic-free patients were not on antipsychotics during the six-month-period prior to blood sampling. Only cases with a DSM IV #295.0-295.89 and #298.9 diagnoses were included to increase clinical homogeneity. Since ethnic heterogeneity and relatedness may affect the distribution of genetic variation and consequently gene expression, de Jong et al. removed non-Caucasian subjects by principal component analysis of SNP array data.

(243) Analysis of Expression Data

(244) The raw microarray data is MIAME compliant and made available at gene expression

omnibus (GEO) under accession GSE38485. First the raw data was (pantile-normalized to correct for overall signal intensity differences among the Illumina human gene expression arrays. Potential batch effects due to date of generation of batch, and because of the use of two different platforms (Illumina HumanRef-8 V3 arrays for batch 1 and HumanRef-12 V3 arrays) were removed by batch analysis in Partek Genomics Suite. After removal of batch effects, the list of probesets that was determined for a signature to be used for identification of chronic cases of schizophrenia from bipolar disorder and controls was extracted.

(245) Reanalysis of Illumina Dataset

(246) In the independent analysis, an accurate prediction of cases from controls of 70%-80% ROC AUC in the deJong et al dataset was independently arrived at using 20 probesets from the signature analysis. This result is important, as deJong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498) included two platforms different from the original exon array analysis that were used, a refined homogenous ethnic background, and a different whole blood RNA extraction methodology. This study was able to definitively ascertain which cases were free of antipsychotics, this first analysis contains both medicated (n=92) and non-medicated cases (n=29); thus most medicated and non-medicated cases were classified correctly, indicating that the signature is not reliant on a 'treatment' artifact. Two extra probe sets that mapped to the same transcripts as the exon array were included in the analysis. Note that there are 2 probes for genes HPR, TRIM4. Both probes are included in the diagnostic analysis. No probe sets were found for gene CCDC109B.

(247) TABLE-US-00037 TABLE 35 The gene expression for the following 20 probe sets for genes identified in the phase I Affymetrix Whole Blood Exon Array dataset as diagnostic for SZ vs BD vs C. The following transcripts were used from GEO dataset GSE38485. Column # Probeset ID Gene Transcript p (SZ vs C)

21001	ILMN_1766551	CPA3	ILMN_19388	0.0220352	ILMN_19388	9052
ILMN_1694432	CRIP2	ILMN_29728	0.020197	ILMN_29728	27078	ILMN_1805344
DDX5	ILMN_20253	0.041188	ILMN_20253	16539	ILMN_1738383	EEF2
ILMN_137242	0.00422672	ILMN_163595	42174	ILMN_2075065	FADS2	ILMN_18999
0.337777	ILMN_18999	9711	ILMN_1697916	GYLTL1B	ILMN_23778	0.0598434
ILMN_23778	12343	ILMN_1712751	HADHA	ILMN_19990	0.319209	ILMN_172930
9628	ILMN_1697499	HLA-DRB5	ILMN_3178	0.23479	ILMN_3178	27713
ILMN_1809212	HPR	ILMN_169425	0.48257	43600	ILMN_2155452	HPR
ILMN_169425	0.693287	ILMN_169425	19423	ILMN_1756455	IL5RA	ILMN_1894
0.6403	ILMN_17920	46474	ILMN_2327812	IL5RA	ILMN_1894	0.572508
ILMN_1894	27014	ILMN_1804929	OXTR	ILMN_7313	0.299299	ILMN_7313
3026	ILMN_1664464	PTGDS	ILMN_19248	6.55E-06	ILMN_19248	17495
ILMN_1744003	SLC44A5	ILMN_13591	0.971099	ILMN_13591	14712	ILMN_1726928
TCEA3	ILMN_27218	0.0850903	ILMN_27218	44519	ILMN_2205322	TREML4
ILMN_24817	0.542362	ILMN_24817	25033	ILMN_1792265	TRIM4	ILMN_5721
0.0626756	ILMN_8530	46384	ILMN_2323385	TRIM4	ILMN_8530	0.938466
ILMN_8530	47890	ILMN_2386179	ZMYND8	ILMN_26803	0.00222525	ILMN_26803

(248) First, reproducibility of the 5 gene diagnostic (TCEA3, GYLTL1B, SLC44A5, IL5RA, DDX5) using the two probe sets for IL5RA was analyzed. The overall ROC was marginally diagnostic with AUC=0.637. However, there appeared to be good reproducibility of the diagnostic signature in males under 30 yrs of age. In the young male subgroup, the two IL5RA probe sets and the DDXS probe set were statistically significant contributors in the logistic model. In Table 36, are results for all subjects broken down by age and gender from the independent deJong et al., 2012 study (de Jong et al., PLoS One 2012; 7(6): e39498).

(249) TABLE-US-00038 TABLE 36 ROC characteristics of de Jong et al., 2012 study using 5 genes diagnostic (TCEA3, GYLTL1B, SLC44A5, IL5RA, DDX5) to diagnose SZ compared to control (C). ROC Gender Age # SZ # C AUC Sensitivity Specificity

Male <30	21	25	0.802	85.71	
64.00	Female <30	10	19	0.684	70.00
68.42	Male ≥30	66	33	0.684	45.45
90.91	Female ≥30	24	41	0.697	45.83
90.24					

(250) Next, the reproducibility of the 11 genes BD+SZ vs NC diagnostic (HPR, TREML4,

ZMYND8, PTGDS, CPA3, TRIM4, SLC44A5, HADHA, CCDC109B, EEF2) was analyzed using the two probe sets for TRIM4. Only one of the probe sets for HPR, had gene expression data for all subjects so this was the one included. Again, there was no Illumina expression for CCDC109B to include. The overall ROC was marginally diagnostic with AUC=0.693. However, there appears to be good reproducibility of the diagnostic in all subgroups when stratified by age and gender. In the young (<30) subgroup, the TREML4, PTGDS and one of the TRIM4 probe sets were statistically significant contributors in the logistic model. In Table 37 are the results for all subjects broken down by age and gender.

(251) TABLE-US-00039 TABLE 37 ROC characteristics of de Jong et al., 2012 study using gene probesets to test the reproducibility of the 11 genes BD + SZ vs NC diagnostic (HPR, TREML4, ZMYND8, PTGDS, CPA3, TRIM4, SLC44A5, OXTR, HADHA, CCDC109B, EEF2). ROC Gender Age # SZ # C AUC Sensitivity Specificity Male <30 21 25 0.857 76.19 84.00 Female <30 10 19 0.916 100.0 78.95 Male ≥30 66 33 0.748 74.24 78.79 Female ≥30 24 41 0.806 79.17 80.49

(252) Finally the antipsychotic-free patients and controls were considered using lumina SZ genes, logistic regression fit to data, subgrouped by age <30 and/or neuroleptic-free status (Table 38).

(253) TABLE-US-00040 TABLE 38 The ROC to diagnose antipsychotic-free patients and control subjects from de Jong et al., 2012 (de Jong et al., PLoS One 2012; 7(6): e39498) gene expression were TREML4, PTGDS, SLC44A5, GYLTL1B, HPR, TRIM4, CPA3 ROC Subjects NC N SZ N AUC Sensitivity Specificity Neuroleptic-free Illumina 22 15 0.642 80.0% 59.1% Subjects Neuroleptic-free, Age <30 14 8 0.866 100.0% 71.4% Subjects

Example 9: SZ-NC Comparison

(254) These results demonstrate the identification of a gene panel to discriminate bipolar schizophrenics from normal controls from 122 top known genes. Forward stepwise variable selection with logistic regression modeling was used to identify a set of genes which would significantly differentiate the bipolar subjects from the normal controls.

(255) Table 39 provides a listing of the set of 5 genes diagnostic for this purpose:

(256) TABLE-US-00041 TABLE 39 Marker Transcript ID Gene 28 2401347 TCEA3 53 3329099 GYLTL1B 41 2418570 SLC44A5 72 2660617 IL5RA 106 3766893 DDX5

Table 40 provides the model it for visits 2 and 4.

(257) TABLE-US-00042 TABLE 40 Visits 2 and 4 AIC Intercept only 168.355 AIC with genes 67.385 Estimate P-value intercept $\beta_{\text{sub.0}}$ -40.2220 <0.0001 marker 28 coefficient $\beta_{\text{sub.1}}$ 6.4838 <0.0001 marker 53 coefficient $\beta_{\text{sub.2}}$ -4.7465 <0.0001 marker 41 coefficient $\beta_{\text{sub.3}}$ 2.3179 <0.0001 marker 72 coefficient $\beta_{\text{sub.4}}$ 2.8666 <0.0001 marker 106 coefficient $\beta_{\text{sub.5}}$ -2.5694 0.0005 ROC AUC 0.967 <0.0001

(258) Prob of Normal from Logistic =
$$\frac{1}{1 + \exp[-(\beta_0 + \beta_1 M_{28} + \beta_2 M_{53} + \beta_3 M_{41} + \beta_4 M_{72} + \beta_5 M_{106})]}$$

(259) The diagnostic measure for discriminating two populations is that a patient is diagnosed as having Schiz if the Prob(NC) from Logistic 0.3323 (optimal cut-point). At this cut-point, across both visits, Sensitivity=86.67 (95% CI 75.4–94.1) and Specificity=93.33 (95% CI 83.8–98.2)

(260) FIG. 11 depicts the ROC Curve for visits 2 and 4.

(261) Table 41 depicts the diagnostic of patients using the 0.3323 cut-point for a positive test.

(262) TABLE-US-00043 TABLE 41 Visit 2 Visit 4 Gene Model NC SZ NC SZ ≤0.3323 3 24 1 28 >0.3323 27 6 29 2

Visit Specific Metrics:

(263) TABLE-US-00044 Sensitivity 0.800 0.933 Specificity 0.900 0.967 Accuracy 0.850 0.964

(264) FIG. 12 shows the logistic model estimated probabilities for each group, for each visit along with the reference line for the 0.3323 cut-point. The visit 2 probabilities are significantly correlated with the visit 4 probabilities (p<0,0001) with r=0.74 (95% CI 0.59–0.83).

(265) Table 42 demonstrates that 83% of the subjects (50/60) received the same diagnostic outcome

from visit 2 to visit 4.

(266) TABLE-US-00045 TABLE 42 Visit 4 Visit 2 ≤ 0.3323 > 0.3323 ≤ 0.3323 27 6 > 0.3323 4 23

(267) The disclosures of each and every patent, patent application, and publication cited herein are hereby incorporated herein by reference in their entirety. While this invention has been disclosed with reference to specific embodiments, it is apparent that other embodiments and variations of this invention may be devised by others skilled in the art without departing from the true spirit and scope of the invention. The appended claims are intended to be construed to include all such embodiments and equivalent variations.

Claims

1. A method of reducing the manifestation and/or progression of bipolar disorder (BD) and schizophrenia (SZ) in a human subject having BD or SZ comprising administering to the human subject one or more clinically approved antipsychotic drugs, wherein the human subject has been selected for administration of the one or more clinically approved antipsychotic drugs after completion of a 4-part gene expression assay on a sample of blood or cells from the human subject, wherein the human subject has been classified as having BD or SZ prior to the administration of the one or more antipsychotic drugs after (a) measuring the mRNA expression levels of each gene in the group consisting of SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, HPR, EEF2, ZMYND8, TCEA3, IL5RA, GYLTL1B, FADS2, CRIP2, DDX5, and HLA-DRB5 in the sample and (b) correlating those measured levels with either BD or SZ by a logistic regression method comprising the following steps: (i) first, the measured mRNA levels of each of the genes SLC44A5, HADHA, CPA3, OXTR, CCDC109B, TREML4, TRIM4, PTGDS, HPR, EEF2, and ZMYND8 are each multiplied by an associated weight factor and then added to form a weighted sum value; and the weighted sum value compared against a reference threshold, wherein a weighted sum value equal to or below this reference threshold identifies the subject as suspected of developing or having BD or SZ; then, second, (ii) if the human subject is suspected of developing or having BD or SZ after step (i), the weighted sum value of the measured mRNA expression levels of each of the genes OXTR, PTGDS, FADS2, and HLA-DRB5 in the sample is calculated and compared against a reference threshold, wherein a weighted sum value equal to or below this reference threshold identifies the human subject as suspected of developing or having BD; and then, third, (iii) if the human subject is suspected of developing or having BD or SZ after step (i), the weighted sum value of the measured mRNA expression levels of the genes TCEA3, SLC44A5, IL5RA, GYLTL1B, and DDX5 in the sample is calculated and compared against a reference threshold, wherein a weighted sum value equal to or below this reference threshold identifies the human subject as suspected of developing or having SZ; and then, fourth, (iv(a)) if the human subject has been identified as suspected of developing or having BD after steps (i) and (ii), the weighted sum value of the measured mRNA expression levels of the genes OXTR, FADS2, and CRIP2 in the sample is calculated and compared against a reference threshold, wherein a weighted sum value above this reference threshold differentiates and classifies the human subject as having BD and not SZ; and (iv(b)) if the human subject has been identified as suspected of developing or having SZ after steps (i) and (iii), the weighted sum value of the measured mRNA expression levels of the genes OXTR, FADS2, and CRIP2 in the sample is calculated and compared against a reference threshold, wherein a weighted sum value equal to or below this reference threshold differentiates and classifies the human subject as having SZ and not BD; wherein the multiplication of mRNA levels by associated weight factors, the weighted sum values, and the reference thresholds are parts of the logistic regression method.

2. The method of claim 1, wherein the human subject is less than 30 years of age, prior to the gene expression measurements.

3. The method of any one of claims 1 and 2, wherein the human subject is displaying psychotic

feature(s), prior to the gene expression measurements.

4. The method of any one of claims 1 and 2, wherein the subject is displaying BD and/or SZ features selected from those identified based on the Diagnostic and Statistical Manual of Mental Illnesses (DSM) criteria, prior to the gene expression measurements.

5. The method of any one of claims 1 and 2, wherein the subject is at high clinical risk, in a prodromal phase, and not yet diagnosed with schizophrenia, prior to the gene expression measurements.

6. The method of any one of claims 1 and 2, wherein the subject is at high clinical risk, in a prodromal phase, and not yet diagnosed with bipolar disorder, prior to the gene expression measurements.

7. The method of any one of claims 1 and 2, wherein the mRNA expression level of each gene is measured by hybridization of a gene array consisting of probes for said genes, RT-PCR, northern blot, nuclease protection, real time PCR, branched DNA, nucleic acid sequence-based amplification (NASBA), RNA Sequencing, digital droplet PCR, and/or differential display.

8. The method of any one of claims 1 and 2, wherein the cells are isolated from the blood.

9. The method of any one of claims 1 and 2, wherein the sample is a blood sample.

10. The method of claim 8, wherein the mRNA expression level of each gene is measured by hybridization of a gene array consisting of probes for said genes, RT-PCR, northern blot, nuclease protection, real time PCR, branched DNA, nucleic acid sequence-based amplification (NASBA), RNA Sequencing, digital droplet PCR, and/or differential display.

11. The method of claim 9, wherein the mRNA expression level of each gene is measured by hybridization of a gene array consisting of probes for said genes, RT-PCR, northern blot, nuclease protection, real time PCR, branched DNA, nucleic acid sequence-based amplification (NASBA), RNA Sequencing, digital droplet PCR, and/or differential display.

12. The method of claim 5, wherein the mRNA expression level of each gene is measured by hybridization of a gene array consisting of probes for said genes, RT-PCR, northern blot, nuclease protection, real time PCR, branched DNA, nucleic acid sequence-based amplification (NASBA), RNA Sequencing, digital droplet PCR, and/or differential display.

13. The method of claim 6, wherein the mRNA expression level of each gene is measured by hybridization of a gene array consisting of probes for said genes, RT-PCR, northern blot, nuclease protection, real time PCR, branched DNA, nucleic acid sequence-based amplification (NASBA), RNA Sequencing, digital droplet PCR, and/or differential display.

14. The method of claim 1, wherein the subject is antipsychotic drug-free, prior to the gene expression measurements.
