



US 20250282860A1

(19) **United States**

(12) **Patent Application Publication**
Ji et al.

(10) **Pub. No.: US 2025/0282860 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **INTERLEUKIN-2 MUTANT AND APPLICATION THEREOF**

(52) **U.S. Cl.**
CPC **C07K 16/246** (2013.01)

(71) Applicant: **Suzhou Forlong Biotechnology Co., Ltd.**, Jiangsu (CN)

(57) **ABSTRACT**

(72) Inventors: **En Ji**, Jiangsu (CN); **Chunyu Wang**, Jiangsu (CN)

The present invention relates to an interleukin-2 (IL-2) mutant and an application thereof. Key sites were selected by means of computer-aided molecular design, and an IL-2 mutant phage display library was constructed based on synthetic biology techniques and combined with forward and reverse screening strategies to screen for a biased IL-2 mutant. Compared with wild-type IL-2, the IL-2 mutant of the present invention has reduced IL-2R α receptor affinity and increased or equivalent IL-2R β receptor affinity, and can activate effector T cells, activate NK cells, and stimulate the release of interferon γ in a biased manner. Also provided in the present invention are a nucleic acid encoding the novel IL-2 mutant and a vector and a host cell comprising the nucleic acid, and further provided are a method for preparing the novel IL-2 mutant, a pharmaceutical combination comprising the novel IL-2 mutant, and a therapeutic use of the novel IL-2 mutant.

(21) Appl. No.: **18/554,874**

(22) PCT Filed: **Apr. 12, 2022**

(86) PCT No.: **PCT/CN2022/086279**

§ 371 (c)(1),

(2) Date: **May 8, 2024**


(30) **Foreign Application Priority Data**

Apr. 13, 2021 (CN) 202110387222.3

Publication Classification

(51) **Int. Cl.**
C07K 16/24 (2006.01)

Specification includes a Sequence Listing.






FIG 1




FIG 2a




FIG 2b




FIG 3a

FIG 3b




FIG 3c




FIG 3d




FIG 3e




FIG 3f




FIG 3g




FIG 3h




FIG 3i




FIG 3j




FIG 3k




FIG 3l




FIG 3m




FIG 3n




FIG 3o




FIG 4a




FIG 4b




FIG 4c




FIG 4d




FIG 4e




FIG 4f




FIG 4g




FIG 4h




FIG 4i




FIG 4j




FIG 4k




FIG 4l




FIG 4m




FIG 4n




FIG 4o




FIG 5a




FIG 5b




FIG 6




FIG 7




FIG 8

INTERLEUKIN-2 MUTANT AND APPLICATION THEREOF

CROSS REFERENCE TO RELATED APPLICATIONS

[0001] This present application is a section 371 of International Application No. PCT/CN2022/086279 filed Apr. 12, 2022, which was published in the Chinese language on Oct. 20, 2022, under International Publication No. WO 2022/218292 A1, which claims the priority of Chinese Patent Application No. 2021103872223 filed with the China National Intellectual Property Administration on Apr. 13, 2021, entitled “Interleukin-2 Mutant and Application Thereof”, the entire content of which is incorporated into the present application by reference.

REFERENCE TO SEQUENCE LISTING SUBMITTED ELECTRONICALLY

[0002] This application contains a sequence listing, which is submitted electronically via EFS-Web as an ASCII formatted sequence listing with a file name “069777-2US1 Substitute Sequence Listing” and a creation date of Apr. 19, 2024, and having a size of 41.1 kb. The sequence listing submitted via EFS-Web is part of the specification and is herein incorporated by reference in its entirety.

TECHNICAL FIELD

[0003] The present invention belongs to the field of biotechnology, and particularly relates to an interleukin-2 mutant, a preparation method therefor, a nucleic acid, plasmid, and host cell required for constructing the interleukin-2 mutant, a corresponding pharmaceutical composition, and an application thereof.

BACKGROUND ART

[0004] Interleukin-2 (IL-2) was discovered in 1976, and it was also called T cell growth factor (TCGF) at that time. It is mainly secreted by activated type I helper lymphocytes TH1 cells. In a precursor expressed by the interleukin gene, which consists of 153 residues, 20 amino acid residues at the N-terminal form a signal peptide, which is excised during secretion to produce mature secreted IL-2. IL-2 has four antiparallel, amphipathic helices, which form a quaternary structure essential for its function.

[0005] IL-2 functions mainly via an IL-2 receptor on the membrane of an effector cell. The IL-2 receptor (IL-2R) has three subunits, P55, P75 and P64, named IL-2 receptor α (IL-2R α , CD25), IL-2 receptor β (IL-2R β , CD122), and IL-2 receptor γ (IL-2R γ , CD132), respectively. IL-2R α binds to IL-2 with a low affinity (K_D of about 1×10^{-8} mol/L) and has no signaling function. IL-2R γ does not bind to IL-2 per se, but shows a high affinity to IL-2 only after IL-2Rs α , β and γ combine (K_D of about 1×10^{-11} mol/L) or β and γ combine (K_D of about 1×10^{-9} mol/L). The signaling function is related to β and γ , and more related to γ . IL-2R α is mainly expressed on the surface of inhibitory regulatory T cells (Tregs) and some endothelial cells, while IL-2R β and IL-2R γ are highly expressed on the surface of effector T cells (Teff) and NK cells. Quiescent effector T cells and NK cells are relatively insensitive to IL-2 because they have no IL-2R α on the cell surface; however, Treg cells are more sensitive to low levels of IL-2 due to their high expression of IL-2R α . Therefore, under normal circumstances, IL-2

will preferentially stimulate the proliferation of Treg cells. During the process of traditional tumor immunotherapies, in order to activate more Teff cells, it is necessary to use a high dose of IL-2, thus inevitably activating a large number of Tregs, leading to severe vascular leakage syndrome (VLS).

SUMMARY OF THE INVENTION

[0006] In order to solve the above technical problems, the present invention discloses a modification method for novel interleukin-2 mutants (IL-2 mutants) and a range of novel interleukin-2 mutants, and further discloses nucleic acid sequences encoding these interleukin-2 mutants and amino acid sequences thereof, vectors containing these nucleic acids, and host cells containing these vectors. The present invention further discloses a pharmaceutical composition of a novel interleukin-2 mutant. The IL-2 mutant and composition disclosed by the present invention can be used for the diagnosis and treatment of a disease. In other embodiments, these novel interleukin-2 mutants, nucleic acids, plasmids, and vectors are used for the diagnosis or treatment of a related disease such as an autoimmune disease or cancer.

[0007] In order to achieve the above object, the present invention provides the following technical solution:

[0008] An interleukin-2 mutant, comprising an amino acid sequence set forth in SEQ ID NO: 1, wherein

[0009] X1 is selected from any amino acid of L, M, E, G, S, D, and Y;

[0010] X2 is any amino acid selected from D and E;

[0011] X3 is any amino acid selected from F, L, V, Y, and I;

[0012] X4 is any amino acid selected from K, H, G, Q, R, N, and S;

[0013] X5 is any amino acid selected from Y, F, and D;

[0014] X6 is any amino acid selected from F, A, Q, S, G, and L; and

[0015] X7 is any amino acid selected from C and A.

[0016] As a preferred embodiment of the interleukin-2 mutant described above, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is L, X2 is D, X3 is L, X4 is H, X5 is Y, X6 is F, and X7 is C; preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is M, X2 is D, X3 is V, X4 is G, X5 is F, X6 is F, and X7 is C; more preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is any amino acid selected from D and E, X3 is F, X4 is K, X5 is Y, X6 is any amino acid selected from A and G, and X7 is any amino acid selected from C and A.

[0017] As a preferred embodiment of the interleukin-2 mutant described above, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is C; preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is C; more preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is G, and X7 is C; further preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is A; more further preferably, the interleukin-2 mutant

comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is G, and X7 is A.

[0018] As a preferred embodiment of the interleukin-2 mutant described above, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is any amino acid selected from I, F, and Y, X4 is any amino acid selected from Q, N, and S, X5 is any amino acid selected from F, Y, and D, X6 is any amino acid selected from F and L, and X7 is any amino acid selected from C and A; preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is I, X4 is Q, X5 is F, X6 is F, and X7 is C; more preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is F, X4 is N, X5 is Y, X6 is F, and X7 is C; further preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is Y, X4 is S, X5 is D, X6 is L, and X7 is C; more further preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is F, X4 is N, X5 is Y, X6 is F, and X7 is A;

[0019] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is S, X2 is D, X3 is L, X4 is Q, X5 is Y, X6 is F, and X7 is C;

[0020] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is any amino acid selected from D and E, X3 is any amino acid selected from L and F, X4 is any amino acid selected from R and K, X5 is any

amino acid selected from F and Y, X6 is any amino acid selected from F, Q, S, and A, and X7 is any amino acid selected from C and A;

[0021] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is L, X4 is R, X5 is F, X6 is F, and X7 is C;

[0022] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is Q, and X7 is C;

[0023] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is S, and X7 is C;

[0024] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is C;

[0025] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is Q, and X7 is A;

[0026] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is S, and X7 is A;

[0027] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is A; and

[0028] preferably, the interleukin-2 mutant comprises an amino acid sequence set forth in SEQ ID NO: 1, in which X1 is Y, X2 is D, X3 is L, X4 is G, X5 is Y, X6 is F, and X7 is C.

(the amino acid sequence of IL-2 mutant)

SEQ ID NO: 1

APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPX1LTX2MLTX3X4FX5MPKKATELK
HLQCLEELKPLEEVLNX6AQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETA
TIVEFLNRWITFX7QSIISTLT

(the amino acid sequence of mutant 21-C1)

SEQ ID NO: 2

APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPLLDMLTLHFYMPKKATELKHLQCL
EEELKPLEEVLNFAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFL
NRWITFCQSIISTLT

(the amino acid sequence of mutant 21-F9)

SEQ ID NO: 3

APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPMLDMLTVGFFMPKKATELKHLQCL
EEELKPLEEVLNFAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFL
NRWITFCQSIISTLT

(the amino acid sequence of mutant 21-F10)

SEQ ID NO: 4

APTSSSTKKTQLQLEHLLLDLQMLNGINNYKNPELTDMLTFKFYMPKKATELKHLQCL
EEELKPLEEVLNAAQSKNFHLRPRDLISNINVIVLELKGSETTFMCEYADETATIVEFL
NRWITFCQSIISTLT

-continued

(the amino acid sequence of mutant 22-A4)

SEQ ID NO: 5

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPGLTDMLTIQFFMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 22-B2)

SEQ ID NO: 6

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPSLTDMLTLQFYMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 22-D11)

SEQ ID NO: 7

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPDLTDMLTLRFFMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 22-E8)

SEQ ID NO: 8

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPDLTEMLTFKFYMPKKATELKHLQCL

EEELKPLEEVLNQAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 23-A5)

SEQ ID NO: 9

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPYLTDMLTLGFYMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 23-C8)

SEQ ID NO: 10

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPGLTDMLTFNFYMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 23-H9)

SEQ ID NO: 11

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPGLTDMLTYSFDMPPKKATELKHLQCL

EEELKPLEEVLNLAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 24-C2)

SEQ ID NO: 12

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPDLTEMLTFKFYMPKKATELKHLQCL

EEELKPLEEVLNSAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 24-C6)

SEQ ID NO: 13

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPELTEMLTFKFYMPKKATELKHLQCL

EEELKPLEEVLNAAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 24-D2)

SEQ ID NO: 14

APTSSSTKKTQLQLEHLLLDLQMILNGINNYKNPELTEMLTFKFYMPKKATELKHLQCL

EEELKPLEEVLNGAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

-continued

NRWITFCQSIISTLT

(the amino acid sequence of mutant 24-E4)

SEQ ID NO: 15

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPDLTDMLTFFKPYMPKKATELKHLQCL

EEELKPLEEVLNAAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFCQSIISTLT

(the amino acid sequence of mutant 21-F10-C125A)

SEQ ID NO: 16

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPELTDMLTFKPYMPKKATELKHLQCL

EEELKPLEEVLNAAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

(the amino acid sequence of mutant 22-E8-C125A)

SEQ ID NO: 17

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPDLTEMLTFKPYMPKKATELKHLQCL

EEELKPLEEVLNQAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

(the amino acid sequence of mutant 23-C8-C125A)

SEQ ID NO: 18

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPGLTDMLTFFKPYMPKKATELKHLQCL

EEELKPLEEVLNFAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

(the amino acid sequence of mutant 24-C2-C125A)

SEQ ID NO: 19

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPDLTEMLTFKPYMPKKATELKHLQCL

EEELKPLEEVLNQAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

(the amino acid sequence of mutant 24-D2-C125A)

SEQ ID NO: 20

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPELTDMLTFKPYMPKKATELKHLQCL

EEELKPLEEVLNQAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

(the amino acid sequence of mutant 24-E4-C125A)

SEQ ID NO: 21

APTSSTTKKTQLQLEHLLLDLQMILNGINNYKNPDLTDMLTFFKPYMPKKATELKHLQCL

EEELKPLEEVLNAAQSKNFHLRPRDLISININVIVLELKGSETTFMCEYADETATIVEFL

NRWITFAQSIISTLT

[0029] In a second aspect of the present invention, there is provided a nucleic acid molecule encoding the interleukin-2 mutant.

[0030] In a third aspect of the present invention, there is provided a plasmid containing the nucleic acid molecule, which is preferably operably linked to a regulatory sequence.

[0031] In a fourth aspect of the present invention, there are provided a host cell containing the plasmid, and a method for producing and, optionally, recovering the interleukin-2 mutant.

[0032] The host cell described in the present invention can be any prokaryotic cell or eukaryotic cell, including but not limit to bacterial cells (e.g., *Escherichia coli*), insect cells (e.g., using a baculovirus expression system), yeast or mammalian cells (e.g., CHO or BHK cell lines). Other

suitable host cells are known to those skilled in the art. A preferred host cell is *Escherichia coli* cell, which has a short culture period and a relatively low production cost.

[0033] In a fifth aspect of the present invention, there is provided a pharmaceutical composition, wherein the pharmaceutical composition comprises a prophylactically or therapeutically effective dose of the interleukin-2 mutant according to any one of claims 1-4, the nucleic acid molecule according to claim 5, or the plasmid according to claim 6, and a physiologically or pharmaceutically acceptable carrier, and is prepared by means of mixing with an excipient or a stabilizer.

[0034] The pharmaceutical composition includes, but is not limited to, a freeze-dried dosage form, an aqueous solution dosage form, a liposome, a capsule dosage form, etc. The concentration of the interleukin-2 mutant of the

present invention or the nucleic acid molecule or plasmid thereof may vary from about 0.1% to 100% (by weight).

[0035] In a sixth aspect of the present invention, there is provided the use of an effective dose of the interleukin-2 mutant according to any one of claims 1-4, the nucleic acid molecule according to claim 5, the plasmid according to claim 6, or the pharmaceutical composition according to claim 8 in the preparation of a medicament for the treatment of a biological material.

[0036] The present invention further provides a method for the diagnosis, prevention or treatment of a disease, comprising administering the interleukin-2 mutant, nucleic acid molecule, plasmid and pharmaceutical composition of the present invention to a subject. The disease is an autoimmune disease, an inflammatory disease, a neurodegenerative disease, a cancer, or a pathogen infection.

[0037] Preferably, the interleukin-2 mutant of the present invention can be used for treating cancer. The cancers described in the present invention include, but are not limited to, lymphoma, blastocytoma, sarcoma (including liposarcoma), neuroendocrine tumors, mesothelioma, schwannoma, meningioma, adenoma, melanoma, and aleukemic leukemia, or lymphatic malignancy. More specific examples of the above cancers include squamous cell carcinoma (e.g., squamous epithelial cell carcinoma), lung cancer, small cell lung cancer, non-small cell lung cancer, lung adenocarcinoma and lung squamous cell carcinoma, peritoneal cancer, hepatocellular carcinoma, gastric cancer, gastrointestinal cancer, pancreatic cancer, malignant glioma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatocellular carcinoma, breast cancer, colon cancer, rectal cancer, colorectal cancer, endometrial or uterine cancer, salivary gland cancer, renal cancer, prostate cancer, vulvar cancer, thyroid cancer, liver cancer, anal cancer, penile cancer, testicular cancer, esophageal cancer, bile duct tumor, head cancer, neck cancer, bone marrow stromal tumor, osteoclastoma, multiple myeloma, osteolytic bone cancers, central nervous system tumor, brain tumor (glioma, neuroblastoma, astrocytoma, medulloblastoma, ependymoma, and retinal neuroblastoma), nasopharyngeal carcinoma, basal cell carcinoma, cholangiocarcinoma, Kaposi's sarcoma, primary liver cancer or endometrial carcinoma, and vascular system tumor (angiosarcoma and hemangiopericytoma).

[0038] Preferably, the interleukin-2 mutant of the present invention can be used for treating a pathogen infection. The pathogens described in the present invention include, but are not limited to, bacteria, fungi, viruses, and parasites.

[0039] The interleukin-2 mutant of the present invention can be used for treating diseases including but not limited to congestive heart failure (CHF), vasculitis, rosacea, acne, eczema, myocarditis and other myocardial diseases, systemic lupus erythematosus, diabetes mellitus, spondylopathy, synovial fibroblast proliferation, bone loss, Paget's disease, disuse osteopenia, malnutrition, periodontal disease, familial spleen anemia, Langerhans cell histiocytosis, spinal cord injury, acute septic arthritis, osteomalacia, hypercortisolism, monostotic fibrous dysplasia, multiple osteofibrous dysplasia, periodontal reconstruction and fracture, sarcoidosis, bone metastasis/osteodynia treatment and humoral hypercalcemia of malignancy, ankylosing spondylitis and other spinal arthropathy, transplant rejection, virus infection, hematologic neoplasm, Hodgkin's lymphoma, and non-Hodgkin's lymphoma (Burkitt's lymphoma, small

lymphocytic lymphoma/chronic lymphocytic leukemia, mycosis fungoides, mantle cell lymphoma, follicular lymphoma, diffuse giant B cell lymphoma, marginal zone lymphoma, hairy cell leukemia and lymphoplasmacytic leukemia), lymphocyte precursor cell tumor, B cell acute lymphoblastic aleukemic leukemia/lymphoma, T cell acute lymphoblastic aleukemic leukemia/lymphoma, thymoma, mature T and NK cell tumors, peripheral T cell aleukemic leukemia, mature T cell aleukemic leukemia/T cell lymphoma, large granular lymphocytic leukemia, Langerhans cell histiocytosis, myeloma of acute myelogenous granulocytic leukemia, mature acute myelogenous leukemia (AML), differentiated acute myelogenous leukemia, acute promyelocytic leukemia, acute myelomonocytic leukemia, acute monocytic leukemia, myelodysplastic syndrome, chronic myeloproliferative disease, chronic myelocytic leukemia, osteoporosis, hepatitis, HIV, AIDS, spinal arthritis, rheumatoid arthritis, inflammatory bowel disease (IBD), sepsis and septic shock, segmental ileitis, psoriasis, scleroderma, graft-versus-host disease (GVHD), allogeneic islet graft rejection, hematological malignancies such as multiple myeloma (MM), myelodysplastic syndrome (MDS) and acute myelogenous leukemia (AML), and tumor-related inflammation, peripheral nerve injury, or demyelinating diseases.

[0040] In a seventh aspect of the present invention, there is provided a detection kit, comprising the interleukin-2 mutant, nucleic acid molecule or plasmid of the present invention. The detection kit is used for detecting pathogens and tumor cells. The pathogens include viruses, bacteria, fungi, parasitic infections, etc. The tumor cells include various benign tumor cells, malignant tumor cells (i.e., cancer cells), solid tumor cells, and hematogenous carcinoma cells.

[0041] Compared with the prior art, the beneficial effects of the present invention lie in that:

[0042] 1. Based on the research of the biological function of IL-2, the present invention develops an IL-2 mutant which selectively agonizes IL-2R $\beta\gamma$ in a biased manner without binding to IL-2R α , and can obtain the potential of reducing the toxicity of IL-2 treatment, such as avoiding VLS, and improving its clinical therapeutic efficacy, such as specifically activating Teff to kill tumors.

[0043] 2. In the present invention, based on the synthetic biology technology, key sites of action of IL-2 against IL-2R α are creatively selected by means of computer-aided molecular design, a phage display library of IL-2 site-directed random mutation is innovatively constructed, and forward and reverse screening strategies are combined to screen for a biased novel IL-2. Compared with wild-type IL-2, the screened novel IL-2 has a higher expression yield.

[0044] 3. The novel IL-2 described in the present invention has a significantly reduced affinity to IL-2R α , and instead of affecting the affinity of IL-2 to IL-2R β , the affinity of the novel IL-2 to IL-2R β is enhanced due to the change of protein conformation. In addition, the novel IL-2 can also significantly activate Teff and NK cells and the release of interferon γ . It shows the effectiveness of the possible anti-tumor treatment thereof and reduces side effects of the treatment in tumor patients.

[0045] 4. The IL-2 mutant and composition described by the present invention can be used for the diagnosis and treatment of a disease. In other embodiments, these novel interleukin-2 mutants, nucleic acids, plasmids, and vectors are used for the diagnose or treatment of a related disease such as an autoimmune disease or cancer.

BRIEF DESCRIPTION OF THE DRAWINGS

[0046] FIG. 1 shows the diversity of the construction of an IL-2 mutant phage library;

[0047] FIG. 2 shows various screening protocols for novel IL-2 and polyclonal phage ELISA data;

[0048] FIG. 2*a* shows the affinity enrichment of antibodies resulting from polyclonal ELISA detection and screening with IL-2R α mutant;

[0049] FIG. 2*b* shows the affinity enrichment of antibodies resulting from polyclonal ELISA detection and screening with IL-2R β mutant;

[0050] FIGS. 3*a-3o* sequentially show the affinity of the IL-2 mutants to IL-2R α in Table 3, as detected by bio-layer interferometry (BLI);

[0051] FIGS. 4*a-4o* sequentially show the BLI-detected affinity of the IL-2 mutants to IL-2R β in Table 3, as detected by bio-layer interferometry (BLI);

[0052] FIG. 5*a* shows the function of IL-2 mutants to activate Teff (CD4 positive T cells);

[0053] FIG. 5*b* shows the function of IL-2 mutants to activate Teff (CD8 positive T cells);

[0054] FIG. 6 shows the function of the IL-2 mutants to activate NK cells;

[0055] FIG. 7 shows the function of the IL-2 mutants to stimulate the release of interferon γ ; and

[0056] FIG. 8 shows the affinity of the IL-2 mutant C125A site mutation proteins to IL-2R α and IL-2R β , as detected by bio-layer interferometry (BLI).

DETAILED DESCRIPTION OF EMBODIMENTS

[0057] In order to enable a person skilled in the art to better understand the solution of the present application, the technical solutions in embodiments of the present application will be described below clearly and completely in conjunction with examples. Obviously, the described embodiments are only some, rather than all, of the embodiments of the present application. Based on the examples of the present application, all other examples obtained by those of ordinary skill in the art without involving any inventive effort should belong to the scope of protection of the present application.

[0058] In order to understand the present invention more thoroughly, some definitions are listed below. The above definitions are intended to include grammatical equivalents.

[0059] As used herein, the term “IL-2” means that interleukin-2 is a cytokine of the chemokine family. It is a cytokine derived from a variety of cells (mainly produced by activated T cells) and also has multi-directional effects (mainly promoting the growth, proliferation and differentiation of lymphocytes). It plays an important role in immune responses, resistance to viral infections, etc., in the body, and can stimulate the proliferation of T cells that have been initiated by specific antigens or mitogenic factors; can activate T cells and promote the production of cytokines; stimulate NK cell proliferation, enhance the killing activity

of NK cells and produce cytokines to induce LAK cell production; promote the proliferation of B cells and the secretion of antibodies; and activate macrophages.

[0060] As used herein, the term “mutant” means an individual in which mutation has occurred and which features a different phenotype from the wild type.

[0061] As used herein, the term “amino acid” means one of the 20 naturally occurring amino acids or any unnatural analogues, which can be located in a specified position. The term “protein” herein means at least two covalently linked amino acids, including proteins, polypeptides, oligopeptides, and peptides. A protein can be composed of naturally occurring amino acids and peptide bonds, or of synthetic peptide mimetic structures, which are called “analogues”. Therefore, the term “amino acid” or “peptide residue” used herein means naturally occurring and synthetic amino acids. By way of example, for the purpose of the present invention, homophenylalanine, citrulline and norleucine are considered as amino acids for the purpose of the present invention. The term “amino acid” also includes imino acid residues such as proline and hydroxyproline. A side chain may be in the (R) or (S) configuration. In a preferred embodiment, an amino acid exists in the (S) or L-configuration. If non-naturally occurring side chains are used, non-amino acid substitutions can be used, for example, to prevent or delay in vivo degradation.

[0062] As used herein, the term “nucleic acid” means a polymer composed of nucleotide units (ribonucleotides, deoxyribonucleotides, related naturally occurring structural variants and their synthetic non-naturally occurring analogues) via phosphodiester bonds. Therefore, the term includes nucleotide polymers, in which nucleotides and bonds therebetween include non-naturally occurring synthetic analogues, such as but not limited to, thiophosphates, aminophosphates, methylphosphates, chiral methylphosphates, 2'-O-methylribonucleotides, peptide nucleic acids (PNAs), etc. For example, these polynucleotides can be synthesized using an automatic DNA synthesizer. The term “oligonucleotide” generally refers to a short polynucleotide, usually no more than about 50 nucleotides. It should be understood that when the nucleotide sequence is represented by a DNA sequence (i.e., A, T, G, and C), this also includes an RNA sequence in which “T” is replaced by “U” (i.e., A, U, G, and C).

[0063] Herein, conventional symbols are used to describe nucleotide sequences: the left-hand end of a single-stranded nucleotide sequence is the 5' end; and the left-hand direction of a double-stranded nucleotide sequence is called 5' direction. The direction in which 5' to 3' nucleotides are added to the newborn RNA transcript is called the transcription direction. A DNA strand with the same sequence as mRNA is called a coding strand.

[0064] As used herein, the term “plasmid” means a plasmid artificially constructed on the basis of a natural plasmid to adapt to laboratory operations. A nucleic acid molecule can be introduced into a host cell, thereby producing a transformed host cell. A vector may include a nucleic acid sequence that allows it to replicate in a host cell, such as origin of replication, and may also include one or more selectable marker genes and other genetic elements known in the art.

[0065] As used herein, the term “host cell”, also called recipient cell, refers to a host cell that receives a foreign gene during transformation and transduction (infection).

[0066] As used herein, the term “pharmaceutically acceptable carrier” means a conventional pharmaceutically acceptable carrier. Remington’s Pharmaceutical Sciences, E W Martin, Mack Publishing Co., Easton, Pa., 15th edition (1975), describes compositions and preparations suitable for drug delivery of one or more therapeutic compounds or molecules (such as one or more antibodies), and additional agents.

[0067] As used herein, the “diagnosis” of a disease refers to the diagnosis of a condition and the development thereof in a patient after examination. The “prevention” of a disease refers to inhibiting the complete development of the disease. The term “treatment” refers to therapeutic intervention to improve the signs or symptoms of a disease or a pathological condition after it begins to develop.

[0068] The term “administration” herein means selecting an appropriate route to introduce the substance into a subject. For example, if the selected route is intravenous, the composition is administered by introducing the substance into the vein of the subject.

[0069] The term “prophylactically/therapeutically effective dose” herein means an amount of a specific agent sufficient to achieve a desired effect in a subject treated with the agent. An exact dosage will depend on the purpose of treatment and can be determined by those skilled in the art by using well-known techniques. The dosage range can be 0.01-100 mg/kg body weight or more, such as 0.1, 1, 10, or 50 mg/kg body weight, preferably 1-10 mg/kg. As is well known in the art, in terms of the degradation of an antibody or Fc fusion, systemic or local drug delivery and new protease synthesis rate, as well as age, weight, general health, sex, diet, administration time, drug interaction, and the severity of the disease, adjustment may be necessary and can be determined by those skilled in the art by means of conventional experimental methods. Such agents include the monomeric Fc domain molecules described herein. In one non-limiting example, this may be the amount of an HIV-specific monomeric Fc domain (or HIV-specific CH3 domain molecule) for preventing, treating or ameliorating HIV infection. Ideally, a therapeutically effective amount of antibody is an amount sufficient to prevent, treat or ameliorate an infection or disease, such as that ascribable to HIV infection in a subject, without causing significant cytotoxic effects in the subject. A therapeutically effective amount of an agent for preventing, ameliorating and/or treating a subject will depend on the subject being treated, the type and severity of the indisposition, and the mode of administration of a therapeutic composition.

[0070] As used herein, the term “residue” means the position in a protein along with the amino acid identity associated therewith. For example, aspartic acid 297 (also known as Asn297 and N297) is a residue in human antibody IgG1.

[0071] As used herein, the term “encoding” means the inherent property of a specific nucleotide sequence in a polynucleotide, such as a gene, a cDNA or an mRNA, to serve as a template for the synthesis of other polymers and macromolecules by biological processes with either a defined nucleotide sequence or a defined amino acid sequence, and the biological properties resulting therefrom. Therefore, if the transcription and translation of mRNA produced from this gene result in a protein in cells or other biological systems, the gene encodes the protein. A coding strand, the nucleotide sequence of which is identical to

mRNA and is usually provided in a sequence listing, and a non-coding strand, used as a transcription template, for a gene or a cDNA, can be referred to as encoding the protein or other product of that gene or cDNA. Unless otherwise specified, the term “nucleotide sequence encoding an amino acid sequence” includes all nucleotide sequences which are degenerate with each other and encode the same amino acid sequence. A nucleotide sequence that encodes a protein or an RNA may also include introns.

[0072] As used herein, the term “operably linked” means that when a first nucleic acid sequence is in a functional relationship with a second nucleic acid sequence, the first nucleic acid sequence is operably linked to the second nucleic acid sequence. For example, if a promoter affects the transcription or expression of a coding sequence, the promoter is operably linked to the coding sequence. Usually, the operably linked DNA sequence is continuous, and when necessary, two protein coding regions are linked in the same reading frame.

[0073] Unless otherwise specified, all the technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the field to which the present disclosure belongs. Unless otherwise explicitly specified in the context, the singular terms “a”, “an”, and “the” include plural indicators. It is to be understood that all base sizes or amino acid sizes given for nucleic acids or polypeptides, as well as all molecular weights or molecular weight values, are approximate and provided for description. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present disclosure, suitable methods and materials are described below. The term “comprise” means “include”. All publications, patent applications, patents and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present description (including the explanation of terms) shall prevail. In addition, the materials, methods and examples are only illustrative and not restrictive.

[0074] The standard recombinant DNA technologies and molecular cloning technologies used in the examples are well known in the art (Ausubel, F. M. et al., Current Protocols in Molecular Biology, published by Greene Publishing Assoc. and Wiley-Interscience), and the materials and methods suitable for microbial growth are well known in the art. Main chemical and biochemical reagents are purchased from KAPA Biosystems, New England Biolabs, TransGen Biotech, Thermo Fisher Scientific, OMEGA biotek, etc.

[0075] The present invention discloses an IL-2 mutant comprising an amino acid sequence set forth in SEQ ID NO: 1, wherein

[0076] X1 is selected from any amino acid of L, M, E, G, S, D, and Y;

[0077] X2 is any amino acid selected from D and E;

[0078] X3 is any amino acid selected from F, L, V, Y, and I;

[0079] X4 is any amino acid selected from K, H, G, Q, R, N, and S;

[0080] X5 is any amino acid selected from Y, F, and D;

[0081] X6 is any amino acid selected from F, A, Q, S, G, and L; and

[0082] X7 is any amino acid selected from C and A.

[0083] The present invention discloses a nucleic acid molecule that encodes the IL-2 mutant;

[0084] the present invention discloses a plasmid containing the above nucleic acid molecule;

[0085] the present invention discloses a host cell containing the above plasmid;

[0086] the present invention discloses a pharmaceutical composition comprising a prophylactically or therapeutically effective dose of the above IL-2 mutant, the above nucleic acid molecule, or the above plasmid, and a physiologically or pharmaceutically acceptable carrier;

[0087] the present invention discloses the use of the above IL-2 mutant, the above nucleic acid molecule, the above plasmid, or the above pharmaceutical composition in the preparation of a medicament for the treatment of a biological material; and

[0088] the present invention discloses a detection kit, wherein the detection kit comprises the above IL-2 mutant, the above nucleic acid molecule, or the above plasmid; and preferably, the detection kit is used for detecting pathogens and tumor cells.

[0089] The present invention will be illustrated in detail below in conjunction with specific examples.

EXAMPLES

Example 1

Screening of IL-2 Mutants

[0090] Key sites of action (K35, R38, F42, K43, Y45, E62, and L72) of IL-2 against IL-2R α were selected by means of computer-aided molecular design, and these seven sites were randomly mutated, and an IL-2 mutant display library with a library capacity of 1.28×10^9 was constructed by means of synthetic biology technology (as shown in FIG. 1) (see the literature PMID: 22518843 for the specific method).

[0091] Novel IL-2 mutants were screened against biotin-labeled soluble IL-2R α and IL-2R β proteins by means of magnetic bead affinity panning method (for the specific method, referring to the literature PMID: 22518843).

[0092] By means of a new strategy of double antigen alternate screening, the biotin-labeled soluble IL-2R α and IL-2R β were successively allowed to specifically bind to streptavidin-coated magnetic beads (purchased from Invitrogen), and four protocols were used for forward and reverse screening (as shown in Table 1). After four rounds of double antigen alternate screening, polyclonal phage ELISA and monoclonal phage ELISA were further performed for screening (for the specific screening methods, referring to PMID: 28966056), a range of the finally screened novel IL-2 that bound IL-2R β with high affinity and bound IL-2R α with low affinity or did not bind IL-2R α were subjected to gene sequencing, and the enriched eligible novel IL-2 mutants were selected (as shown in FIGS. 2a and 2b). After sequencing, the key amino acid site mutations in the screened novel IL-2 mutants were analyzed (as shown in Table 2), and it was found that the three sites K35, R38 and L72 were the most important for binding to IL-2R α , that is, the amino acid changes at these corresponding sites could completely destroy the binding to IL-2R α .

TABLE 1

A	B	C	D	ug
R α	R β	R α	R β	1
R β	R α	R α	R β	1
R α	R β	R β	R α	1
R β	R α	R β	R α	1

TABLE 2

IL-2 mutant sequence (WT-KRFGYEL)							
Name	K35	R38	F42	K43	Y45	E62	L72
1.21-C1	L	D	L	H	Y	E	F
2.21-F9	M	D	V	G	F	E	F
3.21-F10	E	D	F	K	Y	E	A
4.22-A4	G	D	I	Q	F	E	F
5.22-B2	S	D	L	Q	Y	E	F
6.22-D11	D	D	L	R	F	E	F
7.22-E8	D	E	F	K	Y	E	Q
8.23-A5	Y	D	L	G	Y	E	F
9.23-C8	G	D	F	N	Y	E	F
10.23-H9	G	D	Y	S	D	E	L
11.24-C2	D	E	F	K	Y	E	S
12.24-C6	E	E	F	K	Y	E	A
13.24-D2	E	D	F	K	Y	E	G
14.24-E4	D	D	F	K	Y	E	A

Example 2

[0093] The binding capacities of IL-2 mutants to IL-2R α and IL-2R β were detected by bio-layer interferometry (BLI).

[0094] The binding capacities of the IL-2 mutants to IL-2R α and IL-2R β were respectively detected by means of Anti-human IgG Fc Capture (AHC) probes (purchased from GE). The soluble expression and preparation of the IL-2 mutants were basically carried out according to the literature (YING, Tianlei et al., Journal of Biological Chemistry, 2012, 287 (23): 19399-19408).

[0095] The specific detection method involved diluting the IL-2 mutant protein by a factor of 3 (for the detection of the affinity to IL-2R α , dilution was done from an initial concentration of 100 nM to 1.23 nM, and for the detection of the affinity to IL-2R β , dilution was done from an initial concentration of 1000 nM to 12.35 nM).

[0096] As shown in the results of FIGS. 3a-3o, due to the mutations at the key amino acid sites involved in binding to IL-2R α , the IL-2 mutants essentially did not bind to IL-2R α .

[0097] As shown in the results of FIGS. 4a-4o, due to the mutations at the key amino acid sites that did destroy the binding to IL-2R β or due to the conformational change, the IL-2 mutants maintained or even improved the capacity of binding to IL-2R β .

[0098] In an experiment carried out as described in the above assay, the affinity K_D values of fifteen IL-2 mutants expressed by Expi293 (purchased from Thermo Fisher Company) to their receptors were as shown in Table 3:

TABLE 3

Affinity K_D values of IL-2 mutants to their receptors					
Novel IL-2	Affinity to IL-2R α (M)	Affinity to IL-2R β (M)	Novel IL-2	Affinity to IL-2R α (M)	Affinity to IL-2R β (M)
IL-2WT	2.55E-9	7.94E-8	23-C8	N.B	6.78E-7
IL-2V	N.B	2.14E-7	23-H9	N.B	2.80E-5
21-C1	N.B	6.62E-7	24-C2	N.B	9.14E-9
21-F9	N.B	5.00E-8	24-C6	N.B	2.48E-8
21-F10	N.B	8.16E-8	24-D2	N.B	1.72E-7
22-A4	N.B	3.27E-8	24-E4	N.B	8.75E-8
22-B2	N.B	8.66E-5			
22-D11	N.B	1.47E-8			
22-E8	N.B	4.76E-8			

[0099] As could be seen from the shown results: (1) the IL-2 mutants obtained from the phage library by means of double antigen alternate screening were detected to hardly bind to IL-2R α ; and (2) except for 22-B2 and 23-H9, the binding affinities of which to IL-2R β significantly decreased, 24-C2 had significantly increased binding affinity to IL-2R β , 21-C1, 23-C8, and 24-D2 bound to IL-2R β with affinities comparable to IL-2V, and for the remaining IL-2 mutants against IL-2R β , the affinity of IL-2 to IL-2R β stayed essentially unchanged.

Example 3

Experiment of the Function of IL-2 Mutants to Stimulate Effector Cells (Teffs)

[0100] PBMC cells (Allcells Article No.: PB005F, 100 M packaged) were taken out of liquid nitrogen, quickly placed in a water bath at 37° C. for about 1.5 min, and kept spinning continuously, without submerging the whole cryopreservation tube, and the PBMC cells were resuscitated.

[0101] The cells were placed in a biosafety cabinet, and the outside of the cryopreservation tube was wiped with 75% ethanol. The PBMC cells were transferred with a 5 ml pipette to a 50 mL conical tube, in which hot RPMI (purchased from Gibco) with 10% FBS had been added in advance, and DMSO for cell freezing was removed by centrifugation. 20 mL of RPMI medium with 10% FBS was added to re-suspend the cells and was left to stand in a carbon dioxide incubator at 37° C. for culturing overnight. The next day, RPMI medium containing 2% FBS was used to adjust the cell density to 2×10⁶/well, with a volume of 100 ul, and a flat-bottomed 96-well plate (Thermo edge 96 flat) was fully plated. Thereafter, the IL-2 mutant was diluted to four final concentrations, i.e., 100, 10, 1, and 0.1 nM, with medium 1640 containing 2% FBS, with 100 ul/well, and incubated for totally 48 h. Cells were collected by centrifugation and subjected to flow cytometry staining for CD3+CD4+/CD8+T (BioLegend's Cell Surface Flow Cytometry Protocol), and the collected data results were as shown in FIGS. 5a and 5b.

[0102] The IL-2 mutants 21-F10, 22-E8, 23-C8, 24-C2, 24-D2, 24-E4 had good effects on Teff activation, and these mutants were theoretically less immunogenic due to the fact that they had mutations at only three sites relative to wild-type IL-2.

Example 4

[0103] Experiment of the function of IL-2 mutants to stimulate NK cells.

[0104] The method for the resuscitation of PBMC cells was as above.

[0105] To detect the activity of stimulating NK cells, the novel IL-2 mutant was diluted to four final concentrations, i.e., 100, 10, 1, and 0.1 nM, with medium 1640 containing 2% FBS, with 100 ul/well, and co-incubated with the resuscitated PBMC cells for 48 h, cells were collected by centrifugation and subjected to flow cytometry staining for CD3-CD56+NK cells (BioLegend's Cell Surface Flow Cytometry Protocol). The collected data results were as shown in FIG. 6.

[0106] Similarly to the effect of activating Teffs, the IL-2 mutants 21-F10, 22-E8, 23-C8, 24-C2, 24-D2, and 24-E4 also had better NK cell activation effects than the other IL-2 mutants.

Example 5

Experiment of the Function of IL-2 Mutants to Stimulate Interferon γ (IFN γ)

[0107] The method for the resuscitation of PBMC cells was as above.

[0108] The IL-2 mutants were diluted with medium 1640 containing 2% FBS to four final concentrations, i.e., 100, 10, 1, and 0.1 nM, with 100 ul/well, and co-incubated with the resuscitated PBMC cells for 48 h. 50 uL supernatant was taken to detect the release level of IFN γ (R&D DY285B), and the content of IFN γ was determined by ELISA method. As shown in FIG. 7, the IL-2 mutants 21-F10, 22-E8, 23-C8, 24-C2, 24-D2, and 24-E4 had the most significant effect on stimulating the cells to secrete IFN γ .

Example 6

Construction of IL-2 Mutant C125A Site Mutation and the Binding Capacity of the Mutants to IL-2R α and IL-2R β

[0109] The process of the construction of C125A site mutation was as follows: firstly, primers were designed according to the introductions of Beyotime QuickMutation™ Plus Gene Site-directed Mutation Kit (Article No.: D0208S): IL2new C125A-F: ACAGATGGAT-TACCTTTGCTCAGAGCATTATTAGCAC (SEQ ID NO: 43); and IL2new C125A-R: GTGCTAATAATGCTCT-GAGCAAAGGTAATCCATCTGT (SEQ ID NO: 44). For the specific process, reference could be made to the introductions thereof.

[0110] The soluble expression and preparation of IL-2 mutant C125A mutation were basically carried out according to the literature (YING, Tianlei et al., Journal of Biological Chemistry, 2012, 287 (23): 19399-19408). The specific detection method involved using Anti-human IgG Fc Capture (AHC) probes (purchased from GE) to detect the binding capacity of the IL-2 mutant to IL-2R α and IL-2R β , respectively, and diluting the IL-2 mutant C125A site mutation protein by a factor of 3 (for the detection of the affinity to IL-2R α , dilution was done from an initial concentration of 100 nM to 1.23 nM, and for the detection of the affinity to IL-2R β , dilution was done from an initial concentration of 1000 nM to 12.35 nM).

[0111] As shown in the results of FIG. 8, due to the mutations at the key amino acid sites involved in binding to IL-2R α , the IL-2 mutants essentially did not bind to IL-2R α . As shown in the results of FIG. 10, after C125A site mutation, the binding capacity of the IL-2 mutant to IL-2R β

was essentially slightly superior to that of the IL-2 mutant without mutated C125 site, possibly because the mutation at C125 site increased the stability of the protein, so the binding thereof to IL-2R β would be stronger.

[0112] The above description of the disclosed examples enables those skilled in the art to implement or use the present invention. The above description is only preferred embodiments of the present invention, and it should be pointed out that those skilled in the art can also make several improvements and supplements without departing from the

principles of the present invention, and these improvements and supplements should also be regarded as falling within the scope of protection of the present invention. Many modifications to these examples will be obvious to those skilled in the art, and the general principles defined herein can be implemented in other examples without departing from the spirit or scope of the present invention. Therefore, the present invention is not to be limited to the examples shown herein but is to be accorded the widest scope consistent with the principles and novel features disclosed herein.

SEQUENCE LISTING

```
<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (35)..(35)
<223> OTHER INFORMATION: Xaa can be Leu, Met, Glu, Gly, Ser, Asp, or Tyr
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (38)..(38)
<223> OTHER INFORMATION: Xaa can be Asp or Glu
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (42)..(42)
<223> OTHER INFORMATION: Xaa can be Phe, Leu, Val, Tyr, or Ile
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (43)..(43)
<223> OTHER INFORMATION: Xaa can be Lys, His, Gly, Gln, Arg, Asn, or Ser
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (45)..(45)
<223> OTHER INFORMATION: Xaa can be Tyr, Phe, or Asp
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (72)..(72)
<223> OTHER INFORMATION: Xaa can be Phe, Ala, Gln, Ser, Gly, or Leu
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (125)..(125)
<223> OTHER INFORMATION: Xaa can be Cys or Ala

<400> SEQUENCE: 1

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1          5          10          15

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20          25          30

Asn Pro Xaa Leu Thr Xaa Met Leu Thr Xaa Xaa Phe Xaa Met Pro Lys
35          40          45

Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Leu Lys
50          55          60

Pro Leu Glu Glu Val Leu Asn Xaa Ala Gln Ser Lys Asn Phe His Leu
65          70          75          80

Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85          90          95

Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100         105         110

Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Xaa Gln Ser Ile
```

-continued

115	120	125
Ile Ser Thr Leu Thr		
130		
<210> SEQ ID NO 2		
<211> LENGTH: 133		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: interleukin-2 mutant		
<400> SEQUENCE: 2		
Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His		
1 5 10 15		
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys		
20 25 30		
Asn Pro Leu Leu Thr Asp Met Leu Thr Leu His Phe Tyr Met Pro Lys		
35 40 45		
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys		
50 55 60		
Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu		
65 70 75 80		
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu		
85 90 95		
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala		
100 105 110		
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile		
115 120 125		
Ile Ser Thr Leu Thr		
130		
<210> SEQ ID NO 3		
<211> LENGTH: 133		
<212> TYPE: PRT		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: interleukin-2 mutant		
<400> SEQUENCE: 3		
Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His		
1 5 10 15		
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys		
20 25 30		
Asn Pro Met Leu Thr Asp Met Leu Thr Val Gly Phe Phe Met Pro Lys		
35 40 45		
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys		
50 55 60		
Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu		
65 70 75 80		
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu		
85 90 95		
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala		
100 105 110		
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile		
115 120 125		

-continued

Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 4
 <211> LENGTH: 133
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 4

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
 1 5 10 15
 Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
 20 25 30
 Asn Pro Glu Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
 35 40 45
 Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
 50 55 60
 Pro Leu Glu Glu Val Leu Asn Ala Ala Gln Ser Lys Asn Phe His Leu
 65 70 75 80
 Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
 85 90 95
 Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
 100 105 110
 Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
 115 120 125
 Ile Ser Thr Leu Thr
 130

<210> SEQ ID NO 5
 <211> LENGTH: 133
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 5

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
 1 5 10 15
 Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
 20 25 30
 Asn Pro Gly Leu Thr Asp Met Leu Thr Ile Gln Phe Phe Met Pro Lys
 35 40 45
 Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
 50 55 60
 Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu
 65 70 75 80
 Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
 85 90 95
 Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
 100 105 110
 Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
 115 120 125
 Ile Ser Thr Leu Thr
 130

```

<210> SEQ ID NO 7
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 7

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1          5          10          15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20          25          30
Asn Pro Asp Leu Thr Asp Met Leu Thr Leu Arg Phe Phe Met Pro Lys
35          40          45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50          55          60
Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu
65          70          75          80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85          90          95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100         105         110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115         120         125
Ile Ser Thr Leu Thr
130

```


-continued

<210> SEQ ID NO 8
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 8

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30

Asn Pro Asp Leu Thr Glu Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
35 40 45

Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60

Pro Leu Glu Glu Val Leu Asn Gln Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80

Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95

Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110

Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125

Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 9
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 9

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30

Asn Pro Tyr Leu Thr Asp Met Leu Thr Leu Gly Phe Tyr Met Pro Lys
35 40 45

Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60

Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80

Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95

Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110

Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125

Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 10
<211> LENGTH: 133

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 10

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30

Asn Pro Gly Leu Thr Asp Met Leu Thr Phe Asn Phe Tyr Met Pro Lys
35 40 45

Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60

Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80

Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95

Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110

Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125

Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 11
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 11

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30

Asn Pro Gly Leu Thr Asp Met Leu Thr Tyr Ser Phe Asp Met Pro Lys
35 40 45

Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60

Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80

Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95

Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110

Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125

Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 12
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 12

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30
Asn Pro Asp Leu Thr Glu Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
35 40 45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60
Pro Leu Glu Glu Val Leu Asn Ser Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125
Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 13

<211> LENGTH: 133

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 13

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1 5 10 15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20 25 30
Asn Pro Glu Leu Thr Glu Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
35 40 45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50 55 60
Pro Leu Glu Glu Val Leu Asn Ala Ala Gln Ser Lys Asn Phe His Leu
65 70 75 80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85 90 95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100 105 110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
115 120 125
Ile Ser Thr Leu Thr
130

<210> SEQ ID NO 14

<211> LENGTH: 133

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

-continued

<400> SEQUENCE: 14

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
 1 5 10 15
 Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
 20 25 30
 Asn Pro Glu Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
 35 40 45
 Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
 50 55 60
 Pro Leu Glu Glu Val Leu Asn Gly Ala Gln Ser Lys Asn Phe His Leu
 65 70 75 80
 Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
 85 90 95
 Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
 100 105 110
 Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
 115 120 125
 Ile Ser Thr Leu Thr
 130

<210> SEQ ID NO 15

<211> LENGTH: 133

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 15

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
 1 5 10 15
 Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
 20 25 30
 Asn Pro Asp Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
 35 40 45
 Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
 50 55 60
 Pro Leu Glu Glu Val Leu Asn Ala Ala Gln Ser Lys Asn Phe His Leu
 65 70 75 80
 Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
 85 90 95
 Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
 100 105 110
 Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
 115 120 125
 Ile Ser Thr Leu Thr
 130

<210> SEQ ID NO 16

<211> LENGTH: 133

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 16

-continued

```

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1          5          10          15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20          25          30
Asn Pro Glu Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
35          40          45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50          55          60
Pro Leu Glu Glu Val Leu Asn Ala Ala Gln Ser Lys Asn Phe His Leu
65          70          75          80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85          90          95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100         105         110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile
115         120         125
Ile Ser Thr Leu Thr
130

```

```

<210> SEQ ID NO 17
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 17

```

```

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
1          5          10          15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
20          25          30
Asn Pro Asp Leu Thr Glu Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
35          40          45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
50          55          60
Pro Leu Glu Glu Val Leu Asn Gln Ala Gln Ser Lys Asn Phe His Leu
65          70          75          80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
85          90          95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
100         105         110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile
115         120         125
Ile Ser Thr Leu Thr
130

```

```

<210> SEQ ID NO 18
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 18

```

```

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His

```

-continued

1	5	10	15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys	20	25	30
Asn Pro Gly Leu Thr Asp Met Leu Thr Phe Asn Phe Tyr Met Pro Lys	35	40	45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys	50	55	60
Pro Leu Glu Glu Val Leu Asn Phe Ala Gln Ser Lys Asn Phe His Leu	65	70	80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu	85	90	95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala	100	105	110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile	115	120	125
Ile Ser Thr Leu Thr	130		

<210> SEQ ID NO 19
 <211> LENGTH: 133
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 19

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His	1	5	10	15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys	20	25	30	
Asn Pro Asp Leu Thr Glu Met Leu Thr Phe Lys Phe Tyr Met Pro Lys	35	40	45	
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys	50	55	60	
Pro Leu Glu Glu Val Leu Asn Ser Ala Gln Ser Lys Asn Phe His Leu	65	70	75	80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu	85	90	95	
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala	100	105	110	
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile	115	120	125	
Ile Ser Thr Leu Thr	130			

<210> SEQ ID NO 20
 <211> LENGTH: 133
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 20

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His	1	5	10	15
---	---	---	----	----

-continued

```

Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
    20                      25                      30
Asn Pro Glu Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
    35                      40                      45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
    50                      55                      60
Pro Leu Glu Glu Val Leu Asn Gly Ala Gln Ser Lys Asn Phe His Leu
    65                      70                      75                      80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
    85                      90                      95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
    100                      105                      110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile
    115                      120                      125
Ile Ser Thr Leu Thr
    130

```

```

<210> SEQ ID NO 21
<211> LENGTH: 133
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 21

```

```

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
  1          5          10          15
Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
    20                      25                      30
Asn Pro Asp Leu Thr Asp Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
    35                      40                      45
Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
    50                      55                      60
Pro Leu Glu Glu Val Leu Asn Ala Ala Gln Ser Lys Asn Phe His Leu
    65                      70                      75                      80
Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
    85                      90                      95
Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
    100                      105                      110
Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Ala Gln Ser Ile
    115                      120                      125
Ile Ser Thr Leu Thr
    130

```

```

<210> SEQ ID NO 22
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (103)..(104)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (112)..(113)
<223> OTHER INFORMATION: n is a, c, g, or t

```

-continued

```

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (124)..(125)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (127)..(128)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (133)..(134)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (184)..(185)
<223> OTHER INFORMATION: n is a, c, g, or t
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (214)..(215)
<223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 22

gcccccaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaaacc cannbctgac tnnbatgctg      120
accnnbnnbt ttnnbatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180
gaannbctga aaccactgga agaagtgtctg aacnnbgcac agagcaaaaa ctttcatctg      240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa      300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga      360
tggattacct tttgtcagag cattatttagc accctgacc                               399

<210> SEQ ID NO 23
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 23

gcccccaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaaacc cactgctgac tgacatgctg      120
accttgcatt tttacatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180
gaagagctga aaccactgga agaagtgtctg aactttgcac agagcaaaaa ctttcatctg      240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa      300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga      360
tggattacct tttgtcagag cattatttagc accctgacc                               399

<210> SEQ ID NO 24
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 24

gcccccaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaaacc caatgctgac tgatatgctg      120
accgtgggct tttttatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180

```


-continued

```

gaagagctga aacctactgga agaagtgtctg aactttgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct tttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 25
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 25

```

```

gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagagctgac tgacatgctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagagctga aacctactgga agaagtgtctg aacgtgtcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct tttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 26
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 26

```

```

gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagggtgac tgatatgctg    120
accattcagt ttttatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagagctga aacctactgga agaagtgtctg aactttgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct tttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 27
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 27

```

```

gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc caagcctgac tgacatgctg    120
accttcagt tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagagctga aacctactgga agaagtgtctg aactttgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300

```

-continued

accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360

tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 28

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 28

gccccacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60

ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagatctgac tgatatgctg 120

accctccggt tttttatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180

gaagagctga aaccactgga agaagtgctg aactttgcac agagcaaaaa ctttcatctg 240

agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300

accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360

tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 29

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 29

gccccacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60

ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagatctgac tgagatgctg 120

acctttaaat ttttatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180

gaagaactga aaccactgga agaagtgctg aaccaggcac agagcaaaaa ctttcatctg 240

agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300

accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360

tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 30

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 30

gccccacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60

ctgcagatga ttctgaatgg cattaacaac tataaaaacc catacctgac tgacatgctg 120

accctgggtt ttttatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180

gaagagctga aaccactgga agaagtgctg aacttcgcac agagcaaaaa ctttcatctg 240

agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300

accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360

tggattacct tttgtcagag cattattagc accctgacc 399

-continued

<210> SEQ ID NO 31
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 31
gccccaaact gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagggtgac tgacatgctg 120
acctttaact tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180
gaagagctga aacctctgga agaagtgtctg aacttcgcac agagcaaaaa ctttcatctg 240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360
tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 32
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 32
gccccaaact gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagggtgac tgacatgctg 120
acctactcct ttgacatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180
gaagagctga aacctctgga agaagtgtctg aacttggcac agagcaaaaa ctttcatctg 240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360
tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 33
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 33
gccccaaact gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat 60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagatctgac tgagatgctg 120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa 180
gaagaactga aacctctgga agaagtgtctg aactcggcac agagcaaaaa ctttcatctg 240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa 300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga 360
tggattacct tttgtcagag cattattagc accctgacc 399

<210> SEQ ID NO 34
<211> LENGTH: 399

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 34

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagagctgac tgagatgctg      120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180
gaagaactga aacctctgga agaagtgtctg aacgctgcac agagcaaaaa ctttcatctg      240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa      300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga      360
tggattacct tttgtcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 35
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 35

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagagctgac tgacatgctg      120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180
gaagaactga aacctctgga agaagtgtctg aacggtgcac agagcaaaaa ctttcatctg      240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa      300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga      360
tggattacct tttgtcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 36
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 36

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat      60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagatctgac tgatattgctg      120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa      180
gaagaactga aacctctgga agaagtgtctg aacgcggcac agagcaaaaa ctttcatctg      240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa      300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga      360
tggattacct tttgtcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 37
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

-continued

<400> SEQUENCE: 37

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagagctgac tgacatgctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagagctga aaccactgga agaagtgctg aacgctgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgctcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 38

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 38

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagatctgac tgagatgctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagaactga aaccactgga agaagtgctg aaccaggcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgctcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 39

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 39

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaacc cagggtgac tgacatgctg    120
acctttaact tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagagctga aaccactgga agaagtgctg aacttcgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgctcagag cattattagc accctgacc                                399
```

<210> SEQ ID NO 40

<211> LENGTH: 399

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: interleukin-2 mutant

<400> SEQUENCE: 40

```
gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
```

-continued

```

ctgcagatga ttctgaatgg cattaacaac tataaaaaacc cagatctgac tgagatgctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagaactga aaccactgga agaagtgtgtg aactcggcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 41
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 41

```

```

gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaaacc cagagctgac tgacatgctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagaactga aaccactgga agaagtgtgtg aacggtgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 42
<211> LENGTH: 399
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: interleukin-2 mutant

```

```

<400> SEQUENCE: 42

```

```

gccccaaacta gcagcagcac caaaaaaact cagctgcagc tggaacatct gctgctggat    60
ctgcagatga ttctgaatgg cattaacaac tataaaaaacc cagatctgac tgatagctg    120
acctttaaat tttatatgcc aaaaaaagcc actgaactga aacatctgca gtgcctggaa    180
gaagaactga aaccactgga agaagtgtgtg aacggtgcac agagcaaaaa ctttcatctg    240
agaccaagag atctgattag caacattaat gtgattgtgc tggaactgaa aggcagtgaa    300
accaccttta tgtgtgaata tgcagatgaa actgccacca ttgtggaatt tctgaacaga    360
tggattacct ttgtcagag cattattagc accctgacc                                399

```

```

<210> SEQ ID NO 43
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: C125A-F primer

```

```

<400> SEQUENCE: 43

```

```

acagatggat tacctttgct cagagcatta ttagcac                                37

```

```

<210> SEQ ID NO 44
<211> LENGTH: 37

```

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: C125A-R primer

<400> SEQUENCE: 44

gtgctaataa tgctotgagc aaaggtaatc catctgt

37

1. An interleukin-2 mutant, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, wherein

X1 is selected from any amino acid of L, M, E, G, S, D, and Y;

X2 is any amino acid selected from D and E;

X3 is any amino acid selected from F, L, V, Y, and I;

X4 is any amino acid selected from K, H, G, Q, R, N, and S;

X5 is any amino acid selected from Y, F, and D;

X6 is any amino acid selected from F, A, Q, S, G, and L; and

X7 is any amino acid selected from C and A.

2. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is L, X2 is D, X3 is L, X4 is H, X5 is Y, X6 is F, and X7 is C.

3. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is E, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is C.

4. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is G, X2 is D, X3 is any amino acid selected from I, F, and Y, X4 is any amino acid selected from Q, N, and S, X5 is any amino acid selected from F, Y, and D, X6 is any amino acid selected from F and L, and X7 is any amino acid selected from C and A.

5. A nucleic acid molecule encoding the interleukin-2 mutant according to claim 1.

6. A plasmid, characterized by comprising the nucleic acid molecule according to claim 5.

7. A host cell comprising the plasmid according to claim 6.

8. A pharmaceutical composition comprising a prophylactically or therapeutically effective dose of the interleukin-2 mutant according to claim 1, and a physiologically or pharmaceutically acceptable carrier.

9.-10. (canceled)

11. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is M, X2 is D, X3 is V, X4 is G, X5 is F, X6 is F, and X7 is C.

12. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in which X1 is E, X2 is any amino acid selected from D and E, X3 is F, X4 is K, X5 is Y, X6 is any amino acid selected from A and G, and X7 is any amino acid selected from C and A.

13. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is S, X2 is D, X3 is L, X4 is Q, X5 is Y, X6 is F, and X7 is C.

14. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is L, X4 is R, X5 is F, X6 is F, and X7 is C.

15. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is Q, and X7 is C.

16. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is S, and X7 is C.

17. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is C.

18. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is Q, and X7 is A.

19. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is E, X3 is F, X4 is K, X5 is Y, X6 is S, and X7 is A.

20. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is D, X2 is D, X3 is F, X4 is K, X5 is Y, X6 is A, and X7 is A.

21. The interleukin-2 mutant according to claim 1, wherein the interleukin-2 mutant comprises the amino acid sequence set forth in SEQ ID NO: 1, in which X1 is Y, X2 is D, X3 is L, X4 is G, X5 is Y, X6 is F, and X7 is C.

* * * * *