



US 20250282879A1

(19) **United States**

(12) **Patent Application Publication**
TIETZ et al.

(10) **Pub. No.: US 2025/0282879 A1**

(43) **Pub. Date: Sep. 11, 2025**

(54) **MULTISPECIFIC ANTIBODIES HAVING
SPECIFICITY FOR IL-4R AND IL-31**

Publication Classification

(51) **Int. Cl.**

C07K 16/28 (2006.01)

C07K 16/24 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 16/2866** (2013.01); **C07K 16/244**
(2013.01); **C07K 2317/31** (2013.01); **C07K**
2317/33 (2013.01); **C07K 2317/622** (2013.01);
C07K 2317/76 (2013.01); **C07K 2317/92**
(2013.01)

(71) Applicant: **Numab Therapeutics AG**, Horgen
(CH)

(72) Inventors: **Julia TIETZ**, Schlieren (CH); **Tea**
GUNDE, Zurich (CH); **Maria**
JOHANSSON, Denges (CH); **Stefan**
WARMUTH, Au (CH); **Alexandre**
SIMONIN, Rosenau (FR); **Christian**
HESS, Zurich (CH); **Noriko**
SHIRAISHI, Tokyo (JP); **Yoshio**
ARAKAWA, Tokyo (JP); **Yoshihide**
MIYAKE, Tokyo (JP)

(57)

ABSTRACT

The present invention relates to a multispecific antibody comprising one or two binding domains, which specifically bind to IL-4R (IL4R-BDs), and one or two binding domains, which specifically bind to IL-31 (IL31-BDs), wherein the multispecific antibody comprises an immunoglobulin Fc region. The present invention further relates to nucleic acids encoding said multispecific antibody, vector(s) comprising said nucleic acids, host cell(s) comprising said nucleic acids or said vector(s), and a method of producing said multispecific antibody. Additionally, the present invention relates to pharmaceutical compositions comprising said multispecific antibody and methods of use thereof.

(21) Appl. No.: **18/258,858**

(22) PCT Filed: **Dec. 23, 2021**

(86) PCT No.: **PCT/EP2021/087561**

§ 371 (c)(1),


(2) Date: **Jun. 22, 2023**

(30) **Foreign Application Priority Data**

Dec. 23, 2020 (EP) 20216957.9


Specification includes a Sequence Listing.

A



	Dupilumab + BMS-981164	PRO2198-MorrH	PRO2199-MorrL
IC50	0.3082	0.1960	0.2408

Figure 1:





[illegible]

Figure 3:

A inhibition of IL-4 induced SEAP secretion



B inhibition of IL-13 induced SEAP secretion




Figure 4:




Figure 5:





Figure 6:




Figure 8:




Figure 9:




Figure 10:




Figure 11:




Figure 12:





Figure 13:



MULTISPECIFIC ANTIBODIES HAVING SPECIFICITY FOR IL-4R AND IL-31

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a 371 National Phase of International Patent Application No. PCT/EP2021/087561 filed on Dec. 23, 2021, which claims priority to European Patent Application No. 20216957.9 filed on Dec. 23, 2020, the content of each of which applications is incorporated herein by reference.

INCORPORATION OF SEQUENCE LISTING

[0002] This application contains a sequence listing submitted electronically via EFS-web, which serves as both the paper copy and the computer readable form (CRF) and consists of a file entitled “WRN23NP_seqlist.txt”, which was created on Jun. 21, 2023, which is 76,839 bytes in size, and which is herein incorporated by reference in its entirety.

FIELD OF THE INVENTION

[0003] The present invention relates to a multispecific antibody comprising one or two binding domains, which specifically bind to IL-4R (IL4R-BDs), and one or two binding domains, which specifically bind to IL-31 (IL31-BDs), wherein the multispecific antibody comprises an immunoglobulin Fc region. The present invention further relates to nucleic acids encoding said multispecific antibody, vector(s) comprising said nucleic acids, host cell(s) comprising said nucleic acids or said vector(s), and a method of producing said multispecific antibody. Additionally, the present invention relates to pharmaceutical compositions comprising said multispecific antibody and methods of use thereof.

BACKGROUND OF THE INVENTION

[0004] The interleukin-4 receptor IL-4R α , herein also referred to as IL-4R or IL4R, is a type I receptor. It binds interleukin 4 (IL-4) either upon direct interaction with the common cytokine receptor gamma chain (γ_c) or by forming receptor complexes with type II receptors, such as in particular with the type II receptor IL-13R α 1. This IL-4R/IL-13R α 1 receptor complex can bind interleukin 4 (IL-4) as well as interleukin 13 (IL-13) to regulate IgE antibody production in B cells. IL-4 binding to IL-4R is further known to activate macrophages and to promote differentiation of type 2 helper T-cells (Th2-cells), leading to Th2-driven inflammation. IL-4 and/or IL-13 binding to IL-4R α / γ_c and/or IL-4R/IL-13R α 1 leads to the activation of Janus kinase (JAK)1, JAK2, signal transducer and activator of transcription (STAT)1, STAT3, STAT6, and STAT dimerization, which in turn induces the transcription of specific genes.

[0005] IL-4R signaling has been shown to be associated with a number of human diseases, such as inflammatory and autoimmune disorders, including atopic dermatitis, prurigo nodularis, nasal polyposis, chronic urticaria, asthma and chronic obstructive pulmonary disease.

[0006] Several monoclonal antibodies, which reduce or block IL4R-mediated signaling by either binding the cytokines IL-4 and/or IL-13 or to the IL-4R, have been described in the prior art for the treatment of such diseases. For example, dupilumab (Dupixent®) developed by Regeneron

Pharmaceuticals and Sanofi Genzyme, which bind to and antagonizes IL-4R, has been approved for the treatment of moderate-to-severe atopic dermatitis and for the treatment of chronic rhinosinusitis with nasal polyposis (CRSwNP). It is also being evaluated for treatment of persistent asthma in adults and adolescents.

[0007] Atopic dermatitis (AD) is a chronic inflammatory skin disease characterized by intense pruritus (e. g., severe itch) and by scaly and dry eczematous lesions. Severe disease can be extremely disabling due to major psychological problems, significant sleep loss, and impaired quality of life, leading to high socioeconomic costs. AD often begins in childhood before age 5 and may persist into adulthood.

[0008] The pathophysiology of AD is influenced by a complex interplay between immunoglobulin E (IgE)-mediated sensitization, the immune system, and environmental factors. The primary skin defect may be an immunological disturbance that causes IgE-mediated sensitization, with epithelial-barrier dysfunction that is the consequence of both genetic mutations and local inflammation.

[0009] Typical treatments for AD include topical lotions and moisturizers, topical corticosteroid ointments, creams or injections. Most treatment options, however, offer only temporary, incomplete symptom relief. Moreover, many patients with moderate-to-severe AD become resistant to treatment by topical corticosteroids or by calcineurin inhibitors. Therefore, more targeted therapies for the treatment and/or prevention of AD have been developed, such as monoclonal anti-IL4R antibodies that antagonize IL4R function.

[0010] For example, WO 2014/039461 describes anti-IL4R antagonistic antibodies for use in the treatment of atopic dermatitis.

[0011] Furthermore, as already indicated above, the anti-IL-4R monoclonal antibody dupilumab (Dupixent®) received approval from the United States Food and Drug Administration for moderate-to-severe atopic dermatitis in 2017 and for asthma in 2018.

[0012] However, anti-IL-4R therapies have significant limitations. For example, their efficacy in the atopic diseases of interest is often limited and the response rates are often low to moderate. Furthermore, anti-IL-4R therapies do not directly address itching. Itching, however, is a common symptom in skin-associated inflammatory and autoimmune disorders, such as in the case of AD. Itch-associated scratching generally promotes inflammation and worsening of the disease-related symptoms. Thus, there is a critical need for additional treatment options for patients living with moderate to severe atopic dermatitis and other itch-causing inflammatory and autoimmune disorders.

[0013] More specifically, there is a clear need for novel and improved therapeutic antibodies being able to more efficiently reduce or eliminate IL-4- and/or IL-13-mediated signaling activity in said allergic, inflammatory and autoimmune diseases, but also address the itching symptoms (pruritus) associated with these diseases.

[0014] Said therapeutic antibodies should have a high potency as well as a tolerable safety profile to keep dose limiting side effects at a low level. This is particularly important in case of the treatment of chronic diseases, which typically require long term application of said therapeutic antibodies. Furthermore, it is desirable that said therapeutic antibodies have superior biophysical properties, such as a

high stability, in order to facilitate developability and producibility in high yields. In particular, they should have a good storage stability at high concentrations to allow the preparation of highly concentrated formulations for subcutaneous injection applications. Subcutaneous injection is highly desirable and a necessary prerequisite to enable patients to apply the medication themselves.

SUMMARY OF THE INVENTION

[0015] It is an object of the present invention to provide a medicament to improve treatment of allergic, inflammatory and autoimmune diseases, particularly to provide a new, superior treatment modality for pruritus-causing inflammatory and autoimmune diseases, such as moderate to severe atopic dermatitis. More specifically, it is an object of the present invention to provide novel and improved therapeutic antibodies that are able to efficiently reduce or eliminate IL-4- and IL-13-mediated activity in said diseases and that are also able to address the itching symptoms associated with these diseases. It is a further object of the present invention that said novel and improved therapeutic antibodies exhibit high stability at protein concentrations above 100 mg/ml to allow their application by subcutaneous injection.

[0016] The inventors have now surprisingly found that multispecific antibodies comprising one or two IL-4R antagonizing binding domains (IL4R-BDs) and one or two IL-31 neutralizing binding domains (IL31-BDs) can simultaneously block IL-4-/IL-13- as well as IL-31-induced signaling in a highly efficient manner, which allows the simultaneous reduction of IL-4-/IL-13-related symptoms as well as of the itch-related IL-31 activity.

[0017] The inventors have further surprisingly found through testing of many different antibody formats, that said anti-IL-4R $\times$ IL-31 multispecific antibodies, when comprising a functional Fc region, in particular an IgG region comprising a functional Fc region, can block IL-4- and IL-13-induced signaling even better and have very advantageous biophysical properties, in particular an outstanding stability, which allows the preparation of storage stable formulations for antibody concentrations well above 100 mg/ml. Tested formats that did not exhibit the desired combined properties, i. e. the desired high inhibition potency and at the same time the desired high storage stability, were in particular MATCH3, scMATCH3, (scFv)₂Fc(scFv)₂ and Morrison IgG1 (silenced). These multispecific antibody formats were studied intensively. The tested MATCH3- and scMATCH3-based multispecific antibodies exhibited good stability but lacked efficacy in inhibiting IL-4R- and IL-31-mediated signaling. The tested (scFv)₂Fc(scFv)₂-based multispecific antibodies showed improved inhibition potency but was only moderately stable when stored at concentrations above 100 mg/ml for longer time periods. The tested silenced IgG1 Morrison-based multispecific antibodies on the other hand exhibited excellent storage stability at high concentrations but did not completely reach the desired inhibition potency. It was concluded that a functional Fc region, in particular a functional IgG region is necessary for the desired inhibition potency.

[0018] Accordingly, in a first aspect, the present invention relates to a multispecific antibody comprising:

[0019] a) one or two binding domains, which specifically bind to IL-4R (IL4R-BD), and

[0020] b) one or two binding domains, which specifically bind to IL-31 (IL31-BD),

[0021] wherein the multispecific antibody comprises an immunoglobulin Fc region, in particular an IgG region.

[0022] In a second aspect, the present invention relates to a nucleic acid or two nucleic acids encoding the multispecific antibody of the present invention.

[0023] In a third aspect, the present invention relates to a vector or two vectors comprising the nucleic acid or the two nucleic acids of the present invention.

[0024] In a fourth aspect, the present invention relates to a host cell or host cells comprising the vector or the two vectors of the present invention.

[0025] In a fifth aspect, the present invention relates to a method for producing the multispecific antibody of the present invention, comprising (i) providing the nucleic acid or the two nucleic acids of the present invention, or the vector or the two vectors of the present invention, expressing said nucleic acid or said two nucleic acids, or said vector or vectors, and collecting said multispecific antibody from the expression system, or (ii) providing a host cell or host cells of the present invention, culturing said host cell or said host cells; and collecting said multispecific antibody from the cell culture.

[0026] In a sixth aspect, the present invention relates to a pharmaceutical composition comprising the multispecific antibody of the present invention and a pharmaceutically acceptable carrier.

[0027] In a seventh aspect, the present invention relates to a multispecific antibody of the present invention for use as a medicament.

[0028] In an eighth aspect, the present invention relates to a multispecific antibody of the present invention for use in the treatment of a disease, particularly a human disease, more particularly a human disease selected from allergic, inflammatory and autoimmune diseases, particularly from inflammatory and autoimmune diseases.

[0029] In a ninth aspect, the present invention relates to a method for the treatment of a disease, particularly a human disease, more particularly a human disease selected from allergic, inflammatory and autoimmune diseases, particularly from inflammatory and autoimmune diseases, comprising the step of administering the multispecific antibody of the present invention to a patient in need thereof.

[0030] The aspects, advantageous features and preferred embodiments of the present invention summarized in the following items, respectively alone or in combination, further contribute to solving the object of the invention:

[0031] 1. A multispecific antibody comprising:

[0032] a) one or two binding domains, which specifically bind to IL-4R (IL4R-BD), in particular two identical IL4R-BDs, and

[0033] b) one or two binding domains, which specifically bind to IL-31 (IL31-BD), in particular two identical IL31-BDs,

[0034] wherein the multispecific antibody comprises an immunoglobulin Fc region, in particular a functional Fc region.

[0035] 2. The multispecific antibody of item 1, wherein

[0036] each of said IL4R-BDs comprises

[0037] the HCDR1 sequence of SEQ ID NO: 1,

[0038] the HCDR2 sequences of SEQ ID NO: 2,

[0039] the HCDR3 sequence of SEQ ID NO: 3,

[0040] the LCDR1 sequence of SEQ ID NO: 7,

[0041] the LCDR2 sequence of SEQ ID NO: 8, and

- [0042] the LCDR3 sequences of SEQ ID NO: 9; and
 [0043] each of said IL31-BDs comprises
 [0044] the HCDR1 sequence of SEQ ID NO: 12,
 [0045] the HCDR2 sequence of SEQ ID NO: 13,
 [0046] the HCDR3 sequences of SEQ ID NO: 14,
 [0047] the LCDR1 sequence of SEQ ID NO: 17 or 18,
 [0048] the LCDR2 sequence of SEQ ID NO: 19, and
 [0049] the LCDR3 sequences of SEQ ID NOs: 20.
- [0050] 3. The multispecific antibody of item 1 or 2, wherein the format of said multispecific antibody is selected from (scFv)₂-Fc-(scFv)₂ fusion (ADAPTIR); DVD-Ig; a DARTTM and a TRIDENTTM.
- [0051] 4. The multispecific antibody of item 1 or 2, wherein the multispecific antibody comprises an IgG region.
- [0052] 5. The multispecific antibody of item 4, wherein the format of said multispecific antibody is selected from bivalent bispecific IgG formats, trivalent bispecific IgG formats and tetravalent bispecific IgG formats;
 [0053] in particular wherein the format of said multispecific antibody is selected from KiH-based IgGs; DVD-Ig; IgG-scFv fusions, such as CODV-IgG, Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)), bsAb (scFv linked to C-terminus of light chain), Bs1Ab (scFv linked to N-terminus of light chain), Bs2Ab (scFv linked to N-terminus of heavy chain), Bs3Ab (scFv linked to C-terminus of heavy chain), Ts1Ab (scFv linked to N-terminus of both heavy chain and light chain) and Ts2Ab (dsScFv linked to C-terminus of heavy chain);
 [0054] more particularly wherein the format of said multispecific antibody is selected from KiH-based IgGs; DVD-Ig; CODV-IgG and Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)), even more particularly from DVD-Ig and Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)).
- [0055] 6. The multispecific antibody of item 5, wherein the format of said multispecific antibody is selected from Morrison formats.
- [0056] 7. The multispecific antibody of any one of the preceding items, wherein said multispecific antibody comprises two IL4R-BDs.
- [0057] 8. The multispecific antibody of any one of the preceding items, wherein said multispecific antibody comprises two IL31-BDs.
- [0058] 9. The multispecific antibody of item 1, wherein said multispecific antibody comprises:
 [0059] a) two binding domains, which specifically bind to IL-4R (IL4R-BD), and
 [0060] b) two binding domains, which specifically bind to IL-31 (IL31-BD),
 [0061] wherein said multispecific antibody comprises an IgG region, particularly wherein the format of the multispecific antibody is selected from a Morrison format;
 [0062] and wherein
 [0063] each of said IL4R-BDs comprise
 [0064] the HCDR1 sequence of SEQ ID NO: 1,
 [0065] the HCDR2 sequences of SEQ ID NO: 2,
 [0066] the HCDR3 sequence of SEQ ID NO: 3,
 [0067] the LCDR1 sequence of SEQ ID NO: 7,
 [0068] the LCDR2 sequence of SEQ ID NO: 8,
 and
 [0069] the LCDR3 sequences of SEQ ID NO: 9;
 and
 [0070] each of said IL31-BDs comprise
 [0071] the HCDR1 sequence of SEQ ID NO: 12,
 [0072] the HCDR2 sequence of SEQ ID NO: 13,
 [0073] the HCDR3 sequences of SEQ ID NO: 14,
 [0074] the LCDR1 sequence of SEQ ID NO: 17 or 18,
 [0075] the LCDR2 sequence of SEQ ID NO: 19,
 and
 [0076] the LCDR3 sequences of SEQ ID NOs: 20.
- [0077] 10. The multispecific antibody of any one of the preceding items, wherein said multispecific antibody acts as an antagonist of IL4R.
- [0078] 11. The multispecific antibody of item 10, wherein said multispecific antibody blocks the binding of IL4 and IL13 to IL4R.
- [0079] 12. The multispecific antibody of any one of the preceding items, wherein said multispecific antibody blocks the binding of IL-31 to the interleukin 31 receptor alpha (IL-31RA)/oncostatin M receptor (OSMR) complex (IL-31RA/OSMR complex).
- [0080] 13. The multispecific antibody of any one of items 1 to 12, wherein said multispecific antibody has a solubility of at least 100 mg/ml, particularly of at least 120 mg/ml, when formulated in 20 mM acetate buffer at pH 5.5.
- [0081] 14. The multispecific antibody of any one of items 1 to 12, wherein said multispecific antibody, when formulated in 20 mM acetate buffer at pH 5.5, can be concentrated to 100 mg/ml and more, particularly to 100-200 mg/ml, particularly to 100-150 mg/ml, with less than 2%, particularly with less than 1%, loss of monomeric content, as determined by SE-chromatography.
- [0082] 15. The multispecific antibody of any one of items 1 to 12, wherein said multispecific antibody, when formulated in 20 mM acetate buffer at pH 5.5, has a monomeric content of at least 95%, particularly of at least 97%, at concentrations of 100 mg/ml and above, particularly at concentrations in the range of from 100-150 mg/ml.
- [0083] 16. The multispecific antibody of items 1 to 12, wherein said multispecific antibody has a loss in monomer content, after storage at 4° C. for four weeks, of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said multispecific antibody is at a starting concentration of 100 mg/ml, in particular wherein said multispecific antibody is formulated in 20 mM acetate buffer at pH 5.5.
- [0084] 17. The multispecific antibody of any one of the preceding items, wherein the amount of CCL2 secretion of BEAS-2B cells, that are treated with 1 ng/ml IL-4 and 10 ng/ml IL-31, in the presence of 1 µg/ml of said multispecific antibodies is either decreased or only increased by a factor of 1.00 to 1.50, particularly by a factor of 1.00 to 1.4, particularly by a factor of 1.00 to 1.30, particularly by a factor of 1.00 to 1.20, when compared to the amount of CCL2 secretion of untreated BEAS-2B cells.

- [0085] 18. The multispecific antibody of any one of the preceding items, wherein the immunoglobulin Fc region is selected from an IgG subclass, particularly from IgG subclasses IgG1 and IgG4, particularly from IgG4.
- [0086] 19. The multispecific antibody of item 6, wherein the IL4R-BDs are located in the Fab arms, and the IL31-BDs are located in the scFv part of the Morrison format.
- [0087] 20. The multispecific antibody of any one of items 6 and 19, wherein the format of the multispecific antibody is selected from a Morrison-L and a Morrison-H format, in particular wherein the format is a Morrison-H format.
- [0088] 21. The multispecific antibody of any of the preceding items, wherein said IL4R-BDs:
- [0089] a. bind to human IL-4R with a monovalent dissociation constant (K_D) of 0.1 to 200 pM, particularly with a K_D of 0.1 to 100 pM, particularly of 0.1 to 50 pM, as measured by surface plasmon resonance (SPR), particularly wherein said IL4R-BDs are in an scFv format;
- [0090] b. are cross-reactive with *Macaca fascicularis* (Cynomolgus) IL4R, in particular bind to Cynomolgus IL-4R with a monovalent K_D of 1 pM to 5 nM, particularly of 1 pM to 3 nM, particularly of 1 pM to 2 nM, as measured by SPR, particularly wherein said IL4R-BDs are in an scFv format;
- [0091] c. are cross-reactive with *Callithrix jacchus* (Marmoset) IL-4R, in particular bind to Marmoset IL-4R with a monovalent K_D of 1 pM to 5 nM, particularly of 1 pM to 3 nM, particularly of 1 pM to 2 nM, as measured by SPR, particularly wherein said IL4R-BDs are in an scFv format;
- [0092] d. neutralize the human IL-4-induced signaling with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 3 ng/ml, particularly with an IC_{50} of 0.01 to 1.5 ng/ml, as measured in a stat6 reporter gene assay in HEK-Blue cells, wherein said IL4R-BDs are in an scFv format;
- [0093] e. neutralize the human IL-13-induced signaling with an IC_{50} of 0.01 to 10 ng/ml, particularly with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 2.5 ng/ml, as measured in a stat-6 reporter gene assay in HEK-Blue cells, wherein said IL4R-BDs are in an scFv format;
- [0094] f. inhibit the binding of human IL-4 to human IL-4R with an IC_{50} of 0.01 to 10 ng/ml, particularly with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 2 ng/ml, as measured in a competition ELISA, wherein said IL4R-BDs are in an scFv format;
- [0095] g. when being in scFv format, has a melting temperature (T_m), determined by differential scanning fluorimetry (DSF), of at least 63° C., particularly of at least 65° C., particularly of at least 67° C., particularly of at least 69° C., particularly of at least 70° C., particularly of at least 71° C., particularly of at least 72° C., in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
- [0096] h. when being in scFv format, has a loss in monomer content, after storage for four weeks at 4° C. of less than 5%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4; and/or
- [0097] i. when being in scFv format, has a loss in monomer content, after 5 freeze-thawing cycles, of less than 5%, e. g. less than 4%, less than 3%, preferably less than 2%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4.
- [0098] 22. The multispecific antibody of any one of the preceding items, wherein said IL4R-BDs comprise VH3 or VH4 domain framework sequences FR1 to FR4; particularly VH3 domain framework sequences FR1 to FR4.
- [0099] 23. The multispecific antibody of any one of item 23, wherein said IL4R-BDs, when being in scFv-format, comprise
- [0100] (i) a VL domain comprising a VL framework comprising framework regions FR1, FR2 and FR3, which are selected from VK subtypes, particularly from the VK1 and VK3 subtypes, particularly are of the VK1 subtype, and a framework FR4, which is selected from a VK FR4 and a VA FR4, particularly is a VA FR4 comprising an amino acid sequence having at least 70, 80, 90 percent identity to a VA FR4 sequence selected from SEQ ID NO: 26 to SEQ ID NO: 33, more particularly a VA FR4 selected from SEQ ID NO: 26 to SEQ ID NO: 33, particularly VA FR4 according to SEQ ID NO: 26 or 33.
- [0101] 24. The multispecific antibody of any one of the preceding items, wherein said IL4R-BDs comprise
- [0102] a) a VH sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to an amino acid sequence selected from SEQ ID NO: 4, 5 and 6; and
- [0103] b) a VL sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to an amino acid sequence selected from SEQ ID NO: 10 and 11.
- [0104] 25. The multispecific antibody of any one of the preceding items, wherein said IL4R-BDs comprise
- [0105] a) a VH sequence selected from SEQ ID NOs: 4, 5 and 6,
- [0106] b) a VL sequence selected from SEQ ID NOs: 10 and 11.
- [0107] 26. The multispecific antibody of any one of the preceding items, wherein said IL4R-BDs comprise
- [0108] a) a VH sequence of SEQ ID NO: 4 and a VL sequence of SEQ ID NO: 11; or
- [0109] b) a VH sequence of SEQ ID NO: 5 and a VL sequence of SEQ ID NO: 10; or
- [0110] c) a VH sequence of SEQ ID NO: 6 and a VL sequence of SEQ ID NO: 10.
- [0111] 27. The multispecific antibody of any of the preceding items, wherein said IL31-BDs:
- [0112] a. bind to human IL-31 with a monovalent dissociation constant (K_D) of 5 nM or less, particularly with a monovalent K_D of 5 pM to 5 nM, particularly of 5 pM to 2 nM, particularly of 5 to 1000 pM, as measured by surface plasmon resonance (SPR), particularly wherein said IL31-BDs are in an scFv;

- [0113] b. are cross-reactive with *Macaca fascicularis* (Cynomolgus) IL-31, in particular bind to Cynomolgus IL-31 with a monovalent K_D of 5 nM or less, particularly with a monovalent K_D of 5 pM to 5 nM, particularly of 5 pM to 2 nM, particularly of 5 to 1000 pM, as measured by SPR, particularly wherein said IL31-BDs are in an scFv;
- [0114] c. inhibit the human IL-31-induced signaling with an IC_{50} of 0.1 to 30 ng/ml, particularly with an IC_{50} of 0.1 to 20 ng/ml, particularly with an IC_{50} of 0.1 to 10 ng/ml, as measured in a Path Hunter IL-31RA/OSMRb dimerization assay, wherein said IL31-BDs are in an scFv format;
- [0115] d. block the binding of human IL-31 to human IL-31R with an IC_{50} of 0.1 to 20 ng/ml, particularly with an IC_{50} of 0.1 to 10 ng/ml, particularly with an IC_{50} of 0.2 to 6 ng/ml, as measured in a competition ELISA, wherein said IL31-BDs are in an scFv format;
- [0116] e. when being in scFv format, have a melting temperature (T_m), determined by differential scanning fluorimetry, of at least 65° C., preferably of at least 67° C., more preferably at least 69° C., in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
- [0117] f. when being in scFv format, have a loss in monomer content, after storage for at least four weeks at 4° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
- [0118] g. when being in scFv format, have a loss in monomer content, after storage for four weeks at 40° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
- [0119] h. when being in scFv format, have a loss in monomer content, after 3 freeze-thawing cycles, of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4; and/or
- [0120] i. when being in scFv format, have a loss in protein content, after storage for four weeks at 4° C. or 40° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFv is at a starting concentration of 10 mg/ml, and in particular wherein said scFv is formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4.
- [0121] 28. The multispecific antibody of any one of the preceding items, wherein said IL31-BDs comprise VH3 or VH4 domain framework sequences FR1 to FR4; particularly VH3 domain framework sequences FR1 to FR4.
- [0122] 29. The multispecific antibody of item 28, wherein said IL31-BDs, when being in scFv-format, comprise
- [0123] (i) a VL domain comprising a VL framework comprising framework regions FR1, FR2 and FR3, which are selected from VK subtypes, particularly from the VK1 and VK3 subtypes, particularly are of the VK1 subtype, and a framework FR4, which is selected from a VK FR4 and a VA FR4, particularly is a VA FR4 comprising an amino acid sequence having at least 70, 80, 90 percent identity to a VA FR4 sequence selected from SEQ ID NO: 26 to SEQ ID NO: 33, more particularly a VA FR4 selected from SEQ ID NO: 26 to SEQ ID NO: 33, particularly VA FR4 according to SEQ ID NO: 26 or 33.
- [0124] 30. The multispecific antibody of any one of the preceding items, wherein said IL31-BDs comprise
- [0125] a) a VH sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to an amino acid sequence selected from SEQ ID NOs: 15 and 16; and
- [0126] b) a VL sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to an amino acid sequence selected from SEQ ID NOs: 21 and 22.
- [0127] 31. The multispecific antibody of any one of the preceding items, wherein said IL31-BD comprises
- [0128] a) a VH sequence selected from SEQ ID NOs: 15 and 16, particularly the VH sequence of SEQ ID NO: 16; and
- [0129] b) a VL sequence selected from SEQ ID NOs: 21 and 22, particularly the VL sequence of SEQ ID NO: 21.
- [0130] 32. The multispecific antibody of any one of the preceding items, wherein at least one of said IL4R-BDs and/or at least one of said IL31-BDs comprises CDR regions derived from a parental rabbit antibody.
- [0131] 33. The multispecific antibody of any one of the preceding items, wherein at least one IL4R-BD and at least one IL31-BD are capable of binding simultaneously to their respective antigens.
- [0132] 34. The multispecific antibody of any one of the preceding items, wherein said multispecific antibody:
- [0133] a. binds to human IL-4R with a monovalent dissociation constant (K_D) of 1000 pM or less, particularly with a K_D of 1 to 1000 pM, particularly of 1 to 700 pM, particularly of 1 to 500 pM, as measured by surface plasmon resonance (SPR);
- [0134] b. is cross-reactive with *Macaca fascicularis* (Cynomolgus) IL4R, in particular bind to Cynomolgus IL-4R with a monovalent K_D of less than 50 nM, particularly less than 20 nM, particularly with a monovalent K_D of 1 pM to 5 nM, particularly of 1 pM to 2 nM, particularly of 1 to 1000 pM, particularly of 1 to 500 pM, as measured by SPR;
- [0135] c. binds to human IL-31 with a monovalent K_D of less than 5 nM, particularly with a monovalent K_D of 5 pM to 5 nM, particularly of 5 pM to 2 nM, particularly of 5 pM to 1000 pM, as measured by SPR;
- [0136] d. is cross-reactive with *Macaca fascicularis* (Cynomolgus) IL-31, in particular binds to Cynomolgus IL-31 with a monovalent K_D of 20 nM or less, particularly with a monovalent K_D of 5 pM to

- 20 nM, particularly of 5 pM to 10 nM, particularly of 5 pM to 5 nM, as measured by SPR;
- [0137] e. neutralizes the human IL-4-induced signaling with an IC_{50} of 0.2 pM to 200 pM, particularly with an IC_{50} of 0.2 pM to 100 pM, particularly with an IC_{50} of 0.2 pM to 50 pM, as measured in a stat-6 reporter gene assay in HEK-Blue cells;
- [0138] f. neutralizes the human IL-13-induced signaling with an IC_{50} of 0.2 pM to 200 pM, particularly with an IC_{50} of 0.2 pM to 100 pM, particularly with an IC_{50} of 0.2 pM to 50 pM, as measured in a stat-6 reporter gene assay in HEK-Blue cells;
- [0139] g. blocks the human IL-4- and the human IL-13-induced thymus activation regulated chemokine (TARC) secretion with an IC_{50} of 10 pM to 2 nM, particularly with an IC_{50} of 10 pM to 1 nM, as measured in a human whole blood assay;
- [0140] h. binds to HEK293T cells expressing human IL-4R with an EC_{50} of 0.01 to 10 nM, particularly with an EC_{50} of 0.01 to 7 nM, particularly with an EC_{50} of 0.01 to 5 nM;
- [0141] i. binds to HEK293T cells expressing Cynomolgus IL-4R with an EC_{50} of 0.01 to 10 nM, particularly with an EC_{50} of 0.01 to 7 nM, particularly with an EC_{50} of 0.01 to 5 nM;
- [0142] j. blocks the human IL-31-induced signaling with an IC_{50} of 5 pM to 2 nM, particularly with an IC_{50} of 5 pM to 1 nM, particularly with an IC_{50} of 5 to 700 pM, as measured in a Path Hunter IL-31RA/OSMRb dimerization assay;
- [0143] k. blocks the Cynomolgus IL-31-induced signaling with an IC_{50} of 0.05 to 20 nM, particularly with an IC_{50} of 0.05 to 10 nM, particularly with an IC_{50} of 0.05 to 5 nM, as measured in a Path Hunter IL-31RA/OSMRb dimerization assay;
- [0144] l. has a melting temperature (T_m), determined by differential scanning fluorimetry, of at least 60° C., preferably of at least 63° C., more preferably of at least 66° C., in particular wherein said multispecific antibody is formulated in 20 mM MES buffer at pH 7.0; and/or
- [0145] m. has a loss in monomer content, after storage for four weeks, of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said multispecific antibody is at a starting concentration of 100 mg/ml, and in particular wherein said multispecific antibody is formulated in 20 mM acetate buffer at pH 5.5.
- [0146] 35. The multispecific antibody of any one of the preceding items, which is selected from Morrison antibodies having the following heavy chain (HC) and light chain (LC) sequences:
- [0147] an HC sequence of SEQ ID NO: 38 and an LC sequence of SEQ ID NO: 37;
- [0148] an HC sequence of SEQ ID NO: 40 and an LC sequence of SEQ ID NO: 39;
- [0149] an HC sequence of SEQ ID NO: 42 and an LC sequence of SEQ ID NO: 41;
- [0150] an HC sequence of SEQ ID NO: 44 and an LC sequence of SEQ ID NO: 43;
- [0151] an HC sequence of SEQ ID NO: 46 and an LC sequence of SEQ ID NO: 45;
- [0152] an HC sequence of SEQ ID NO: 48 and an LC sequence of SEQ ID NO: 47; and
- [0153] an HC sequence of SEQ ID NO: 50 and an LC sequence of SEQ ID NO: 49.
- [0154] 36. A nucleic acid or two nucleic acids encoding the multispecific antibody of any one of items 1 to 35.
- [0155] 37. A vector or two vectors comprising the nucleic acid or two nucleic acids of item 36.
- [0156] 38. A host cell or host cells comprising the vector or the two vectors of item 37.
- [0157] 39. A method for producing the multispecific antibody of any one of items 1 to 35, comprising
- [0158] (i) providing the nucleic acid or two nucleic acids of item 36, or the vector or the two vectors of item 37, expressing said nucleic acid or nucleic acids, or said vector or vectors, and collecting said multispecific antibody from the expression system, or (ii) providing a host cell or host cells according to item 38, culturing said host cell or said host cells; and collecting said multispecific antibody from the cell culture.
- [0159] 40. A pharmaceutical composition comprising the multispecific antibody of any one of items 1 to 35 and a pharmaceutically acceptable carrier.
- [0160] 41. The multispecific antibody of any one of items 1 to 35 for use as a medicament.
- [0161] 42. The multispecific antibody of any one of items 1 to 35 for use in the treatment of a disease, particularly a human disease, more particularly a human disease selected from allergic, inflammatory and autoimmune diseases, particularly from inflammatory and autoimmune diseases.
- [0162] 43. The multispecific antibody of any one of items 1 to 35 for use in the treatment of a disease according to item 42, wherein said disease is selected from pruritus-causing allergic diseases, pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases, particularly from pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases.
- [0163] 44. The multispecific antibody of item 42 or 43, wherein said disease is selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma; particularly wherein said disease is atopic dermatitis.
- [0164] 45. A method for the treatment of a disease, particularly a human disease, more particularly a human disease selected from allergic, inflammatory and autoimmune diseases, particularly from inflammatory and autoimmune diseases, comprising the step of administering the multispecific antibody of any one of items 1 to 35 to a patient in need thereof.
- [0165] 46. The method of item 45, wherein said disease is selected from pruritus-causing allergic diseases, pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases, particularly from pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases.
- [0166] 47. The method of item 45 or 46, wherein said disease is selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma; particularly wherein said disease is atopic dermatitis.

[0167] 48. A method for the treatment of a disease, more particularly a human disease selected from allergic, inflammatory and autoimmune diseases; particularly from pruritus-causing allergic diseases, pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases;

[0168] said method comprising the steps of administering to a patient in need thereof

[0169] 1. a first antibody comprising

[0170] a) an IL4R-BD, as defined in any of the preceding items 1 to 26 and 32 to 35, and

[0171] b) an Fc region, in particular an IgG region, in particular an IgG4 region; and

[0172] 2. a second antibody comprising an IL31-BD, as defined in any of the preceding items 1 to 20 and 27 to 35.

[0173] 49. The method of item 48, wherein the administration is performed such that an effective dose level of said first antibody and said second antibody is simultaneously reached in the blood plasma of the patient in need thereof.

[0174] 50. The method of item 48 or 49, wherein said disease is selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma; particularly wherein said disease is atopic dermatitis.

[0175] 51. A kit comprising

[0176] 1. a first antibody comprising

[0177] a) an IL4R-BD, as defined in any of the preceding items 1 to 26 and 32 to 35, and

[0178] b) an Fc region, in particular an IgG region, in particular an IgG4 region; and

[0179] 2. a second antibody comprising an IL31-BD, as defined in any of the preceding items 1 to 20 and 27 to 35.

[0180] 52. The pharmaceutical composition of item 40, wherein said pharmaceutical composition is a therapeutic agent for a human disease selected from allergic, inflammatory and autoimmune diseases.

[0181] 53. The pharmaceutical composition of item 52, wherein said disease is selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma; particularly wherein said disease is atopic dermatitis.

BRIEF DESCRIPTION OF THE DRAWINGS

[0182] FIG. 1 shows the potency of the Morrison-H molecules PRO2198 and PRO2199 to inhibit IL-4R- and IL-31-induced signaling in the BEAS-2B assay compared to a combination of dupilumab and BMS-981164. PRO2198 and PRO2199 show similar potencies to inhibit IL-4R- and IL-31-induced signaling when compared to the combination of dupilumab and BMS-981164 (A). Individual blockage of IL-4R or IL-31 alone by the addition of dupilumab or BMS-981164 alone is less efficacious than the combined blockage of IL-4R- and IL-31-induced signaling (B).

[0183] FIG. 2 shows the schematic representation of the IgG4 Morrison-H format (left) and Morrison-L format (right) with labeled subdomain nomenclature. Linker sequences connecting the Fv domains (light grey) consist of

(G₄S) modules of different repetitions (two modules in between constant regions and scFv domains and 4 modules in between VL2 and VH2).

[0184] FIG. 3 shows the potency of the optimized anti-IL-4R scFvs PRO1898 and PRO1899 to block human IL-4-induced signaling (A) and human IL-13-induced signaling (B) of IL-4R in a Stat-6 reporter gene assay in HEK-Blue cells. The potency of the analyzed molecules is compared to dupilumab.

[0185] FIG. 4 shows the potency of the optimized anti-IL-4R scFvs PRO1898 (A) and PRO1899 (B) to inhibit the interaction between human IL-4 and human IL-4R assessed by competitive ELISA.

[0186] FIG. 5 shows the concomitant binding of Morrison antibody PRO2198 to human IL-4R/human IL-31 and to cynomolgus monkey IL-4R/cynomolgus monkey IL-31. Top: schematic representation of the experimental design; Bottom: sensorgram showing concomitant binding of PRO2198 to human IL-4R/human IL-31 (dark grey) and cynomolgus monkey IL-4R/cynomolgus monkey IL-31 (light grey).

[0187] FIG. 6 shows the binding of Morrison-H (MorrH) and Morrison-L (MorrL) molecules PRO2198, PRO2199, PRO2206 and PRO2207 and negative control molecule PRO2077 to human IL-4R (A) and to cynomolgus IL-4R (B) on transfected HEK293T cells. All molecules bound to human IL-4R (A). No binding to cynomolgus IL-4R was observed for the control molecule PRO2077 (B). An antibody targeting V5 Tag (V5 Tag), which is fused to the N-terminus of human and cynomolgus IL-4R expression constructs, was used as a positive control for plasma membranous binding.

[0188] FIG. 7 shows the potencies of Morrison-H (MorrH) and Morrison-L (MorrL) antibodies PRO2198, PRO2199, PRO2206 and PRO2207 to neutralize human IL-4- and human IL-13-induced signaling in the stat-6 reporter gene assay. The potency of IL-4R blockage by the Morrison-H and Morrison-L antibodies is comparable to dupilumab for both IL-4-induced signaling (A and C) and IL-13-induced signaling (B and D). The IC₅₀ values depicted in the FIGS. 5A to 5D represent single point measurements.

[0189] FIG. 8 shows the potencies of Morrison-H (MorrH) antibodies PRO2198 (A) and PRO2199 (B) to neutralize human IL-4-induced signaling in the stat-6 reporter gene assay with or without an excess of IL-31 in the assay medium. No potency loss is observed in presence of an excess of IL-31. The IC₅₀ values depicted in the FIGS. 6A and 6B represent single point measurements.

[0190] FIG. 9 shows the potencies of Morrison-H (MorrH) and Morrison-L (MorrL) antibodies PRO2198, PRO2199, PRO2206 and PRO2207 to neutralize human IL-31-induced signaling in the IL-31 RA/OSMR dimerization assay with or without an excess of IL-4R extracellular domain in the assay medium. The Morrison antibodies blocked the interaction between IL-31 and IL-31 RA with similar potency when compared to BMS-981164 (A and B). No potency loss was determined in presence of an excess of IL-4R (C and D).

[0191] FIG. 10 shows the potency of the Morrison-H antibodies PRO2198 and PRO2199 to neutralize cynomolgus IL-31-induced signaling in the IL-31RA/OSMR dimerization assay. The antibodies PRO2198 and PRO2199 were able to block the interaction between cynomolgus IL-31 and IL-31RA.

[0192] FIG. 11 shows the potency of Morrison-H molecules PRO2198 and PRO2199 and corresponding anti-IL-4R IgG4 molecules PRO2209 and PRO2210, which lack IL-31 scFv binding domains, to block IL-4- and IL-13-induced CD23 expression in human PBMCs. PRO2198 and PRO2199 inhibit IL-4- and IL-13-induced CD23 upregulation in human monocytes (A), naïve B cells (B) and memory B cells (C) with similar potency when compared to dupilumab. PRO2209 and PRO2210 show equal inhibition when compared to dupilumab.

[0193] FIG. 12 shows an SDS-Page analysis of PRO2198, PRO2199, PRO2206 and PRO2207 under reducing and non-reducing conditions.

[0194] FIG. 13 shows an analysis of clipping by SDS-PAGE. 1st lane: Marker—2nd: pH 3.5, t0—3rd: pH 3.5, t10d—4th: pH 5, t0—5th: pH 5, t10d—6th: pH 7, t0d—7th: pH 7, t10d—8th: pH 8.5, t0—9th: pH 8.5, t10d. *Presumably, incomplete reduction of HC dimer. #Presumably, incomplete reduction of LC.

DETAILED DESCRIPTION OF THE INVENTION

[0195] Known therapeutic anti-IL-4R antibodies often suffer from a lack of efficacy and low to moderate response rates in patients. They also do not directly address itching, which is a common symptom particularly in skin-associated inflammatory and autoimmune disorders. There is thus a need in the medical field for improved anti-IL-4R antibody-based therapeutics, which have higher efficacy for allergic, inflammatory and autoimmune diseases, in particular for pruritus-causing inflammatory and autoimmune diseases, such as moderate to severe atopic dermatitis, and at the same time are able to potentially reduce disease-related itching.

[0196] The present invention provides multispecific antibodies comprising one or two binding domains, which specifically bind to IL-4R (IL4R-BD), and one or two binding domains, which specifically bind to IL-31 (IL31-BD), wherein the multispecific antibody comprises an immunoglobulin Fc region. The multispecific antibodies of the present invention can simultaneously block IL-4/IL-13 as well as IL-31-induced signaling in a highly efficient manner.

[0197] It is believed by the inventors that the immunoglobulin Fc region provides additional FcγR interactions that enhance the binding of the multispecific antibodies of the invention to IL-4R on living cells, and thereby increasing their potency to block IL-4R-mediated signaling. The analysis of different antibody formats in a human PBMC assay further indicates that the interaction of the Fc region with Fc gamma receptor type II (FcγRII) seems to be of particular importance for the observed potency increase. Thus, in the context of the present invention, the term “functional Fc region” refers to an Fc region that is able to interact with FcγR, in particular with FcγRII, within the range of its natural functionality.

[0198] According to the best knowledge of the inventors, the simultaneous blockage of both the IL-4R signaling pathway and the IL-31 signaling pathway has not yet been reported in the prior art, let alone the highly efficient simultaneous blockage of said pathways by a single molecule.

[0199] Interleukin 31 (IL-31) is an inflammatory cytokine that helps trigger cell-mediated immunity against pathogens. IL-31 is preferentially produced by Th2-cells. IL-31 sends

signals through a heterodimeric receptor complex (IL-31R or IL31R) comprising the interleukin 31 receptor alpha (IL-31RA or IL31RA) and the oncostatin M receptor p (OSMR R), expressed in immune and epithelial cells. Binding of IL-31 to this receptor complex results in activation of the JAK/STAT and PI3K/AKT signal transduction pathways, and also activates different MAPK pathways (ERK, p38, and JNK).

[0200] IL-31 is implicated in various chronic inflammatory diseases. For example, it has been found that IL-31 over-expression in mice results in dermatitis-like symptoms, see, Dillon, et al, *Nature Immunol.* 5:752-760, (2004). Furthermore, in many chronic inflammatory diseases, such as for example in AD, IL-31 mediates activation of nerve fibers within the skin of the patients resulting in an aggressive itch-phenotype, which exacerbates the symptoms of these diseases upon patient scratching, see for example Oetjen et al., *Cell*, 171: 217-228 (2017). Scratching can lead to skin barrier disruption, access of microbial pathogens into the skin and further promotes inflammation at the site.

[0201] Also, it has been found that IL-31 is implicated in allergic asthma, allergic rhinitis, inflammatory bowel diseases, malignancies and osteoporosis, see Bagci et al., *J Allergy Clin Immunol*, 141(3):858-866 (2018).

[0202] Blockage of IL-31/IL-31RA signaling by anti-IL31RA antibodies, such as for example by the anti-IL-31RA antibody nemolizumab, is clinically proven to be effective at reducing itch in patients suffering from AD, see Ruzicka et al., *N Engl J Med*, 376:2092-2093 (2017).

[0203] Furthermore, the IL-31-neutralizing antibody BMS-981164 was developed to provide an effective targeted therapy for the treatment of chronic pruritic skin conditions, see Lewis et al, *J Eur Acad of Dermatol Venereol*, 31(1): 142-150 (2017).

[0204] Besides targeting IL-4R (first specificity), it was decided to target IL-31 (second specificity) and not the cell-bound IL-31RA, to avoid any potential cross linkage between cells, which might well happen when targeting two different types of receptors on a cell surface simultaneously (e. g. IL-4R and IL-31RA).

[0205] As stated above, the multispecific antibodies of the present invention can simultaneously block IL-4/IL-13- as well as IL-31-induced signaling in a very efficient manner. For example, at a concentration of 1 µg/ml, PRO2198 and PRO2199, two representative examples of the multispecific antibodies of the present invention, are able to reduce the combined IL-4R- and IL-31-mediated CCL2 secretion of BEAS-2B cells, that have been treated with 1 ng/ml IL-4 and 10 ng/ml IL-31, at least to a similar extent when compared to an equivalent amount of a 1:1 mixture of the anti-IL-4R antibody dupilumab and the anti-IL-31 antibody BMS-981164 (FIG. 1B).

[0206] The chemokine (C—C motif) ligand 2 (CCL2), also referred to as monocyte chemoattractant protein 1 (MCP1) and small inducible cytokine A2, is a small cytokine that belongs to the CC chemokine family. CCL2 recruits monocytes, memory T cells, and dendritic cells to the sites of inflammation produced by either tissue injury or infection. CCL2 is implicated in pathogenesis of several diseases characterized by monocytic infiltrates, such as psoriasis, rheumatoid arthritis and atherosclerosis (Xia et al, *Expert Opinion on Therapeutic Patents.*, 2009, 19 (3): 295-303).

[0207] Some of the multispecific antibodies of the present invention, e. g. PRO2198, are able to reduce the combined

IL-4R- and IL-31-mediated CCL2 secretion of said BEAS-2B cells stronger than an equivalent amount of such a 1:1 mixture of dupilumab and BMS-981164 and almost to the level of untreated BEAS-2B cells (FIG. 1B).

[0208] The term “BEAS-2B cells” or BEAS-2B”, as used herein, refers to cells of an immortalized non-tumorigenic epithelial cell line from human bronchial epithelium. This cell line has been established via transfection with an adenovirus 12-SV40 hybrid virus and subsequent immortalization via consecutive cell passaging, see Reddel et al., *Cancer Res.*, 48(7):1904-1909 (1988). BEAS-2B cells have been widely used to study cellular and molecular mechanisms in lung related diseases and as an in vitro cell model for lung-related toxicology studies.

[0209] In addition, the anti-IL-4R $\times$ IL-31 multispecific antibodies of the present invention exhibit very advantageous biophysical properties, in particular an outstanding formulation and storage stability at antibody concentrations well above 100 mg/ml. For example, the loss of the protein content as well as of the monomeric content for PRO2198, a representative multispecific antibody of the present invention, is less than 1% when stored for four months at 4° C. at a concentration above 100 mg/ml in 20 mM acetate buffer (pH 5.5).

[0210] The multispecific antibodies of the present invention thus provide distinct therapeutic advantages over conventional therapies.

[0211] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by those of ordinary skill in the art to which this invention pertains.

[0212] The terms “comprising” and “including” are used herein in their open-ended and non-limiting sense unless otherwise noted. With respect to such latter embodiments, the term “comprising” thus includes the narrower term “consisting of”.

[0213] The terms “a” and “an” and “the” and similar references in the context of describing the invention (especially in the context of the following claims) are to be construed to cover both the singular and the plural, unless otherwise indicated herein or clearly contradicted by context. For example, the term “a cell” includes a plurality of cells, including mixtures thereof. Where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

[0214] In one aspect, the present invention relates to a multispecific antibody comprising:

[0215] a) one or two binding domains, which specifically bind to IL-4R (IL4R-BD), and

[0216] b) one or two binding domains, which specifically bind to IL-31 (IL31-BD),

[0217] wherein the multispecific antibody comprises an immunoglobulin Fc region.

[0218] The term “antibody” and the like, as used herein, includes whole antibodies or single chains thereof; and any antigen-binding fragment (i. e., “antigen-binding portion”) or single chains thereof; and molecules comprising antibody CDRs, VH regions or VL regions (including without limitation multispecific antibodies). A naturally occurring “whole antibody” is a glycoprotein comprising at least two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as VH) and a heavy chain constant region. The heavy chain constant

region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein as VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDRs), flanked by regions that are more conserved, termed framework regions (FRs). Each VH and VL is composed of three CDRs and four FRs arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4. The variable regions of the heavy and light chains contain a binding domain that interacts with an antigen. The constant regions of the antibodies may mediate the binding of the immunoglobulin to host tissues or factors, including various cells of the immune system (e. g., effector cells) and the first component (C1q) of the classical complement system.

[0219] The term “immunoglobulin Fc region” or “Fc region”, as used herein, is used to define a C-terminal region of an immunoglobulin heavy chain, i. e. the CH2 and CH3 domains of the heavy chain constant regions. The term “Fc region” includes native-sequence Fc regions and variant Fc regions, i. e. Fc regions that are engineered to exhibit certain desired properties, such as for example altered Fc receptor binding function and/or reduced or suppressed Fab arm exchange. An example of such an engineered Fc region is the knob-into-hole (KiH) technology (see for example Ridgway et al., *Protein Eng.* 9:617-21 (1996) and Spiess et al., *J Biol Chem.* 288(37):26583-93 (2013)). Native-sequence Fc regions include human IgG1, IgG2 (IgG2A, IgG2B), IgG3 and IgG4. “Fc receptor” or “FcR” describes a receptor that binds to the Fc region of an antibody. Particularly, the FcR is a native sequence human FcR, which binds an IgG antibody (a gamma receptor) and includes receptors of the Fc γ RI, Fc γ RII, and Fc γ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors, Fc γ RII receptors including Fc γ RIIA (an “activating receptor”) and Fc γ RIIB (an “inhibiting receptor”), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating receptor Fc γ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc γ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain, (see M. Daeron, *Annu. Rev. Immunol.* 5:203-234 (1997). FcRs are reviewed in Ravetch and Kinet, *Annu. Rev. Immunol.* 9: 457-92 (1991); Capet et al, *Immunomethods* 4: 25-34 (1994); and de Haas et al, *J. Lab. Clin. Med.* 126: 330-41 (1995). Other FcRs, including those to be identified in the future, are encompassed by the term “FcR” herein. The term “Fc receptor” or “FcR” also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus. Guyer et al., *J. Immunol.* 117: 587 (1976) and Kim et al., *J. Immunol.* 24: 249 (1994). Methods of measuring binding to FcRn are known (see, e. g., Ghetie and Ward, *Immunol. Today* 18: (12): 592-8 (1997); Ghetie et al., *Nature Biotechnology* 15 (7): 637-40 (1997); Hinton et al., *J. Biol. Chem. TJI* (8): 6213-6 (2004); WO 2004/92219 (Hinton et al). Binding to FcRn in vivo and serum half-life of human FcRn high-affinity binding polypeptides can be assayed, e. g., in transgenic mice or transfected human cell lines expressing human FcRn, or in primates to which the polypeptides having a variant Fc region are administered. WO 2004/42072 (Presta) describes antibody variants which

improved or diminished binding to FcRs. See also, e. g., Shields et al., *J. Biol. Chem.* 9(2): 6591-6604 (2001).

[0220] In particular embodiments, the immunoglobulin Fc region comprised in the multispecific antibodies of the invention is selected from an Fc region of an IgG subclass, particularly from an Fc region of the IgG subclasses IgG1 and IgG4, particularly from an Fc region of IgG4.

[0221] In particular embodiments, the multispecific antibodies of the invention comprise an IgG region.

[0222] The term “IgG region”, as used herein, refers to the heavy and light chain of an immunoglobulin G, i. e. the Fc region, as defined above, and the Fab region, consisting of the VL, VH, CL and CH1 domains. The term “IgG region” includes native-sequence IgG regions, such as human IgG1, IgG2 (IgG2A, IgG2B), IgG3 and IgG4, as well as engineered IgG regions, which exhibit certain desired properties, as for example the properties defined above for the Fc region. The term “functional IgG region”, as used herein, refers to an IgG region comprising a functional Fc region.

[0223] In particular embodiments, the multispecific antibodies of the invention comprise an IgG region, wherein the IgG region is selected from the IgG subclasses IgG1 and IgG4, in particular from IgG4.

[0224] The terms “binding domain”, “antigen-binding fragment thereof”, “antigen-binding portion” of an antibody, and the like, as used herein, refer to one or more fragments of an intact antibody that retain the ability to specifically bind to a given antigen (e. g., IL4R, IL-31). Antigen-binding functions of an antibody can be performed by fragments of an intact antibody. Specifically, in case of the multispecific antibodies of the present invention, the terms “binding domain”, “antigen-binding fragment thereof”, “antigen-binding portion”, and the like, as used herein, refer to a Fab fragment, i. e. a monovalent fragment consisting of the VL, VH, CL and CH1 domains; a F(ab)₂ fragment, i. e. a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; an Fv fragment consisting of the VL and VH domains of a single arm of an antibody; a disulfide stabilized Fv fragment (dsFv); and a single chain Fv fragment (scFv). Preferably, the binding domains of the antibodies of the present invention are independently of each other selected from a Fab fragment, an Fv fragment, a dsFv fragment and a single-chain Fv fragment (scFv). In particular embodiments, the binding domains of the antibodies of the present invention are independently of each other selected from a Fab fragment and a single-chain Fv fragment (scFv). In other particular embodiments, the VL and VH domains of the scFv fragment are stabilized by an inter-domain disulfide bond, in particular said VH domain comprises a single cysteine residue in position 51 (AHO numbering) and said VL domain comprises a single cysteine residue in position 141 (AHO numbering).

[0225] The term “Complementarity Determining Regions” (“CDRs”) refers to amino acid sequences with boundaries determined using any of a number of well-known schemes, including those described by Kabat et al. (1991), “Sequences of Proteins of Immunological Interest,” 5th Ed. Public Health Service, National Institutes of Health, Bethesda, MD (“Kabat” numbering scheme); Al-Lazikani et al., (1997) *JMB* 273, 927-948 (“Chothia” numbering scheme); ImMunoGenTics (IMGT) numbering (Lefranc, M.-P., *The Immunologist*, 7, 132-136 (1999); Lefranc, M.-P. et al., *Dev. Comp. Immunol.*, 27, 55-77 (2003)) (“IMGT” numbering scheme); and the numbering scheme described in

Honegger & Plückthun, *J. Mol. Biol.* 309 (2001) 657-670 (“AHO” numbering). For example, for classic formats, under Kabat, the CDR amino acid residues in the heavy chain variable domain (VH) are numbered 31-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3); and the CDR amino acid residues in the light chain variable domain (VL) are numbered 24-34 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3). Under Chothia the CDR amino acids in the VH are numbered 26-32 (HCDR1), 52-56 (HCDR2), and 95-102 (HCDR3); and the amino acid residues in VL are numbered 24-34 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3). By combining the CDR definitions of both Kabat and Chothia, the CDRs consist of amino acid residues 26-35 (HCDR1), 50-65 (HCDR2), and 95-102 (HCDR3) in human VH and amino acid residues 24-34 (LCDR1), 50-56 (LCDR2), and 89-97 (LCDR3) in human VL. Under IMGT the CDR amino acid residues in the VH are numbered approximately 26-35 (HCDR1), 51-57 (HCDR2) and 93-102 (HCDR3), and the CDR amino acid residues in the VL are numbered approximately 27-32 (LCDR1), 50-52 (LCDR2), and 89-97 (LCDR3) (numbering according to “Kabat”). Under IMGT, the CDRs of an antibody can be determined using the program IMGT/DomainGap Align.

[0226] In the context of the present invention, the numbering system suggested by Honegger & Plückthun (“AHO”) is used (Honegger & Plückthun, *J. Mol. Biol.* 309 (2001) 657-670), unless specifically mentioned otherwise. In particular, the following residues are defined as CDRs according to AHO numbering scheme: LCDR1 (also referred to as CDR-L1): L24-L42; LCDR2 (also referred to as CDR-L2): L58-L72; LCDR3 (also referred to as CDR-L3): L107-L138; HCDR1 (also referred to as CDR-H1): H27-H42; HCDR2 (also referred to as CDR-H2): H57-H76; HCDR3 (also referred to as CDR-H3): H108-H138. For the sake of clarity, the numbering system according to Honegger & Plückthun takes the length diversity into account that is found in naturally occurring antibodies, both in the different VH and VL subfamilies and, in particular, in the CDRs, and provides for gaps in the sequences. Thus, in a given antibody variable domain usually not all positions 1 to 149 will be occupied by an amino acid residue.

[0227] The term “binding specificity” as used herein refers to the ability of an individual antibody to react with one antigenic determinant and not with a different antigenic determinant. As used herein, the term “specifically binds to” or is “specific for” refers to measurable and reproducible interactions such as binding between a target and an antibody, which is determinative of the presence of the target in the presence of a heterogeneous population of molecules including biological molecules. For example, an antibody that specifically binds to a target (which can be an epitope) is an antibody that binds this target with greater affinity, avidity, more readily, and/or with greater duration than it binds to other targets. In its most general form (and when no defined reference is mentioned), “specific binding” is referring to the ability of the antibody to discriminate between the target of interest and an unrelated molecule, as determined, for example, in accordance with specificity assay methods known in the art. Such methods comprise, but are not limited to Western blots, ELISA, RIA, ECL, IRMA, SPR (Surface plasmon resonance) tests and peptide scans. For example, a standard ELISA assay can be carried out. The scoring may be carried out by standard color development (e. g. secondary antibody with horseradish peroxidase and tetramethyl

benzidine with hydrogen peroxide). The reaction in certain wells is scored by the optical density, for example, at 450 nm. Typical background (=negative reaction) may be about 0.1 OD; typical positive reaction may be about 1 OD. This means the ratio between a positive and a negative score can be 10-fold or higher. In a further example, an SPR assay can be carried out, wherein at least 10-fold, particularly at least 100-fold difference between a background and signal indicates on specific binding. Typically, determination of binding specificity is performed by using not a single reference molecule, but a set of about three to five unrelated molecules, such as milk powder, transferrin or the like.

[0228] Suitably, the antibodies of the invention are isolated antibodies. The term “isolated antibody”, as used herein, refers to an antibody that is substantially free of other antibodies having different antigenic specificities (e. g., an isolated antibody that specifically binds IL-4R and IL-31 is substantially free of antibodies that specifically bind antigens other than IL-4R and IL-31. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

[0229] Suitably, the antibodies of the invention are monoclonal antibodies. The term “monoclonal antibody” as used herein refers to antibodies that have substantially identical amino acid sequences or are derived from the same genetic source. A monoclonal antibody displays a binding specificity and affinity for a particular epitope, or binding specificities and affinities for specific epitopes.

[0230] Antibodies of the invention include, but are not limited to, chimeric, human and humanized antibodies.

[0231] The term “chimeric antibody” (or antigen-binding fragment thereof), as used herein, refers to an antibody molecule (or antigen-binding fragment thereof) in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen-binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species; or (b) the variable region, or a portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity. For example, a mouse antibody can be modified by replacing its constant region with the constant region from a human immunoglobulin. Due to the replacement with a human constant region, the chimeric antibody can retain its specificity in recognizing the antigen while having reduced antigenicity in human as compared to the original mouse antibody.

[0232] The term “humanized” antibody (or antigen-binding fragment thereof), as used herein, refers to an antibody (or antigen-binding fragment thereof) that retains the reactivity of a non-human antibody while being less immunogenic in humans. This can be achieved, for instance, by retaining the non-human CDR regions and replacing the remaining parts of the antibody with their human counterparts (i. e., the constant region as well as the framework portions of the variable region). Additional framework region modifications may be made within the human framework sequences as well as within the CDR sequences derived from the germline of another mammalian species. The humanized antibodies of the invention may include amino acid residues not encoded by human sequences (e. g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo, or a conservative substitution to promote stability or manufacturing). See, e. g., Morrison et al., *Proc. Natl. Acad. Sci. USA*, 81:6851-

6855, 1984; Morrison and Oi, *Adv. Immunol.*, 44:65-92, 1988; Verhoeven et al., *Science*, 239: 1534-1536, 1988; Padlan, *Molec. Immun.*, 28:489-498, 1991; and Padlan, *Molec. Immun.*, 31: 169-217, 1994. Other examples of human engineering technology include but are not limited to the Xoma technology disclosed in U.S. Pat. No. 5,766,886.

[0233] The term “recombinant humanized antibody” as used herein, includes all human antibodies that are prepared, expressed, created or isolated by recombinant means, such as antibodies isolated from a host cell transformed to express the humanized antibody, e. g., from a transfectoma, and antibodies prepared, expressed, created or isolated by any other means that involve splicing of all or a portion of a human immunoglobulin gene, sequences to other DNA sequences.

[0234] Preferably, the multispecific antibodies of the invention are humanized. More preferably, the multispecific antibodies of the invention are humanized and comprises rabbit derived CDRs.

[0235] The term “multispecific antibody” as used herein, refers to an antibody that binds to two or more different epitopes on at least two or more different targets (e. g., IL-4R and IL-31). Preferably, the multispecific antibodies of the invention are bispecific. The term “bispecific antibody” as used herein, refers to an antibody that binds to at least two different epitopes on two different targets (e. g., IL-4R and IL-31).

[0236] The term “epitope” means a protein determinant capable of specific binding to an antibody. Epitopes usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three-dimensional structural characteristics, as well as specific charge characteristics. “Conformational” and “linear” epitopes are distinguished in that the binding to the former but not the latter is lost in the presence of denaturing solvents.

[0237] The term “conformational epitope” as used herein refers to amino acid residues of an antigen that come together on the surface when the polypeptide chain folds to form the native protein.

[0238] The term “linear epitope” refers to an epitope, wherein all points of interaction between the protein and the interacting molecule (such as an antibody) occurring linearly along the primary amino acid sequence of the protein (continuous).

[0239] The term “recognize” as used herein refers to an antibody antigen-binding fragment thereof that finds and interacts (e. g., binds) with its conformational epitope.

[0240] The term “avidity” as used herein refers to an informative measure of the overall stability or strength of the antibody-antigen complex. It is controlled by three major factors: antibody epitope affinity; the valency of both the antigen and antibody; and the structural arrangement of the interacting parts. Ultimately these factors define the specificity of the antibody, that is, the likelihood that the particular antibody is binding to a precise antigen epitope.

[0241] Suitably, the multispecific antibodies of the invention comprise one or two IL4R-BDs. Particularly, the multispecific antibodies of the invention comprise two IL4R-BDs.

[0242] The term “IL-4R” refers in particular to human IL-4R with UniProt ID number P24394. Suitably, the IL4R-BD of the present invention targets IL-4R, in particular human IL-4R. Particularly, the multispecific antibodies of

the invention comprises one or two IL4R-BDs that target human and Cynomolgus (*Macaca fascicularis*) IL-4R. More particularly, the multispecific antibody of the invention comprises one or two IL4R-BDs that targets human, Cynomolgus (*Macaca fascicularis*) and Marmoset (*Callithrix jacchus*) IL-4R.

[0243] Suitably, the IL4R-BDs are characterized by one or more of the following parameters:

[0244] a. bind to human IL-4R with a monovalent dissociation constant (K_D) of 0.1 to 200 pM, particularly with a K_D of 0.1 to 100 pM, particularly of 0.1 to 50 pM, as measured by surface plasmon resonance (SPR), particularly wherein said IL4R-BDs are in an scFv format;

[0245] b. are cross-reactive with *Macaca fascicularis* (Cynomolgus) IL4R, in particular bind to in particular bind to Cynomolgus IL-4R with a monovalent K_D of 1 pM to 5 nM, particularly of 1 pM to 3 nM, particularly of 1 pM to 2 nM, as measured by SPR, particularly wherein said IL4R-BDs are in an scFv format;

[0246] c. are cross-reactive with *Callithrix jacchus* (Marmoset) IL-4R, in particular bind to Marmoset IL-4R with a monovalent K_D of 1 pM to 5 nM, particularly of 1 pM to 3 nM, particularly of 1 pM to 2 nM, as measured by SPR, particularly wherein said IL4R-BDs are in an scFv format;

[0247] d. neutralize the human IL-4-induced signaling with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 3 ng/ml, particularly with an IC_{50} of 0.01 to 1.5 ng/ml, as measured in a stat6 reporter gene assay in HEK-Blue cells, wherein said IL4R-BDs are in an scFv format;

[0248] e. neutralize the human IL-13-induced signaling with an IC_{50} of 0.01 to 10 ng/ml, particularly with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 2.5 ng/ml, as measured in a stat-6 reporter gene assay in HEK-Blue cells, wherein said IL4R-BDs are in an scFv format; and/or

[0249] f. inhibit the binding of human IL-4 to human IL-4R with an IC_{50} of 0.01 to 10 ng/ml, particularly with an IC_{50} of 0.01 to 5 ng/ml, particularly with an IC_{50} of 0.01 to 2 ng/ml, as measured in a competition ELISA, wherein said IL4R-BDs are in an scFv format.

[0250] The term “HEK-Blue cells” or “HEK-Blue”, as used herein, refers to commercially available human embryonic kidney cells that are transfected with and stably express an optimized secreted embryonic alkaline phosphatase (SEAP) reporter gene under the control of a promoter inducible by NF- κ B transcription factor. The level of SEAP protein released into the culture media is typically used as a measure of NF- κ B activation.

[0251] As used herein, the term “affinity” refers to the strength of interaction between the antibody and the antigen at single antigenic sites. Within each antigenic site, the variable region of the antibody “arm” interacts through weak non-covalent forces with antigen at numerous sites; the more interactions, the stronger the affinity.

[0252] “Binding affinity” generally refers to the strength of the total sum of non-covalent interactions between a single binding site of a molecule (e. g., of an antibody) and its binding partner (e. g., an antigen or, more specifically, an epitope on an antigen). Unless indicated otherwise, as used herein, “binding affinity”, “bind to”, “binds to” or “binding to” refers to intrinsic binding affinity that reflects a 1:1

interaction between members of a binding pair (e. g., an antibody fragment and an antigen). The affinity of a molecule X for its partner Y can generally be represented by the dissociation constant (K_D). Affinity can be measured by common methods known in the art, including those described herein. Low-affinity antibodies generally bind antigens slowly and tend to dissociate readily, whereas high-affinity antibodies generally bind antigens faster and tend to remain bound longer. A variety of methods of measuring binding affinity are known in the art, any of which can be used for purposes of the present invention. Specific illustrative and exemplary embodiments for measuring binding affinity, i. e. binding strength are described in the following.

[0253] The term “ K_{assoc} ”, “ K_a ” or “ K_{on} ”, as used herein, are intended to refer to the association rate of a particular antibody-antigen interaction, whereas the term “ K_{dis} ”, “ K_d ” or “ K_{off} ”, as used herein, is intended to refer to the dissociation rate of a particular antibody-antigen interaction. In one embodiment, the term “ K_D ”, as used herein, is intended to refer to the dissociation constant, which is obtained from the ratio of K_d to K_a (i. e. K_d/K_a) and is expressed as a molar concentration (M). The “ K_D ” or “ K_D value” or “ K_D ” or “ K_D value” according to this invention is in one embodiment measured by using surface plasmon resonance assays.

[0254] Affinity to recombinant human IL-4R, recombinant Cynomolgus IL-4R and recombinant Marmoset IL-4R is determined by surface plasmon resonance (SPR) measurements, as described in the paragraphs [0187] and [0188] (scFvs) and [0265] (multispecific molecules). Affinity to recombinant human IL-31, recombinant Cynomolgus IL-31 and recombinant Marmoset IL-31 is determined by surface plasmon resonance (SPR) measurements, as described in the paragraphs [0209] (scFvs) and [0267] (multispecific molecules).

[0255] Suitably, the multispecific antibody of the invention acts as an antagonist of IL4R. In other words, the IL4R-BD of the multispecific antibody of the invention is an inhibitor of IL-4R signaling. The term “blocker” or “inhibitor” or “antagonist”, as used herein, refers to an antibody or binding domain thereof that inhibits or reduces a biological activity of the antigen it binds to. The IL4R-BDs of the multispecific antibody of the invention bind to the IL-4R, thereby blocking the binding of IL-4 to IL4R, in particular the binding of both IL-4 and IL-13 to IL-4R, which leads to reduced IL-4R function.

[0256] Suitably, the IL4R-BDs of the multispecific antibody of the present invention are further characterized by one or more of the following parameters:

[0257] g. when being in scFv format, has a melting temperature (T_m), determined by differential scanning fluorimetry (DSF), of at least 63° C., particularly of at least 65° C., particularly of at least 67° C., particularly of at least 69° C., particularly of at least 70° C., particularly of at least 71° C., particularly of at least 72° C., in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;

[0258] h. when being in scFv format, has a loss in monomer content, after storage for four weeks at 4° C. of less than 5%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4; and/or

- [0259] i. when being in scFv format, has a loss in monomer content, after 5 freeze-thawing cycles, of less than 5%, e. g. less than 4%, less than 3%, preferably less than 2%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4.
- [0260] DSF is described earlier (Egan, et al., MAbs, 9(1) (2017), 68-84; Niesen, et al., Nature Protocols, 2(9) (2007) 2212-2221). The midpoint of transition for the thermal unfolding of the scFv constructs is determined by Differential Scanning Fluorimetry using the fluorescence dye SYPRO® Orange (see Wong & Raleigh, Protein Science 25 (2016) 1834-1840). Samples in phosphate-citrate buffer at pH 6.4 are prepared at a final protein concentration of 50 µg/ml and containing a final concentration of 5xSYPRO® Orange in a total volume of 100 µl. Twenty-five microliters of prepared samples are added in triplicate to white-walled AB gene PCR plates. The assay is performed in a qPCR machine used as a thermal cycler, and the fluorescence emission is detected using the software's custom dye calibration routine. The PCR plate containing the test samples is subjected to a temperature ramp from 25° C. to 96° C. in increments of 1° C. with 30 s pauses after each temperature increment. The total assay time is about 2 h. The T_m is calculated by the software GraphPad Prism using a mathematical second derivative method to calculate the inflection point of the curve. The reported T_m is an average of three measurements.
- [0261] The loss in monomer content is as determined by area under the curve calculation of SE-HPLC chromatograms. SE-HPLC is a separation technique based on a solid stationary phase and a liquid mobile phase as outlined by the US Pharmacopeia (USP), chapter 621. This method separates molecules based on their size and shape utilizing a hydrophobic stationary phase and aqueous mobile phase. The separation of molecules is occurring between the void volume (V₀) and the total permeation volume (V_T) of a specific column. Measurements by SE-HPLC are performed on a Chromaster HPLC system (Hitachi High-Technologies Corporation) equipped with automated sample injection and a UV detector set to the detection wavelength of 280 nm. The equipment is controlled by the software EZChrom Elite (Agilent Technologies, Version 3.3.2 SP2) which also supports analysis of resulting chromatograms. Protein samples are cleared by centrifugation and kept at a temperature of 4-6° C. in the autosampler prior to injection. For the analysis of scFv samples the column Shodex KW403-4F (Showa Denko Inc., #F6989202) is employed with a standardized buffered saline mobile phase (50 mM sodium-phosphate pH 6.5, 300 mM sodium chloride) at the recommended flow rate of 0.35 ml/min. The target sample load per injection is 5 pg. Samples are detected by an UV detector at a wavelength of 280 nm and the data recorded by a suitable software suite. The resulting chromatograms are analyzed in the range of V₀ to V_T thereby excluding matrix associated peaks with >10 min elution time.
- [0262] Suitably, the IL4R-BDs of the multispecific antibodies of the invention are binding domains provided in the present disclosure. The IL4R-BDs of the multispecific antibodies of the invention include, but are not limited to, the humanized IL4R-BDs whose sequences are listed in Table 1.
- [0263] Suitably, the multispecific antibodies of the invention comprise one or two IL31-BDs. Particularly, the multispecific antibodies of the invention comprise two IL31-BDs.
- [0264] The term "IL-31" or "IL31" refers in particular to human IL-31 with UniProt ID number Q6EBC2. Suitably, the IL31-BD of the multispecific antibodies of the invention targets IL-31, in particular human IL31 as shown in UniProt ID number Q6EBC2. Particularly, the multispecific antibodies of the invention comprise one or two IL31-BDs that target human and cynomolgus (*Macaca fascicularis*) IL-31.
- [0265] Suitably, the IL31-BDs used in the present invention are characterized by one or more of the following parameters:
- [0266] a. bind to human IL-31 with a monovalent dissociation constant (K_D) of 5 nM or less, particularly with a monovalent K_D of 5 pM to 5 nM, particularly of 5 pM to 2 nM, particularly of 5 to 1000 pM, as measured by surface plasmon resonance (SPR), particularly wherein said IL31-BDs are in an scFv;
 - [0267] b. are cross-reactive with *Macaca fascicularis* (Cynomolgus) IL-31, in particular bind to Cynomolgus IL-31 with a monovalent K_D of 5 nM or less, particularly with a monovalent K_D of 5 pM to 5 nM, particularly of 5 pM to 2 nM, particularly of 5 to 1000 pM, as measured by SPR, particularly wherein said IL31-BDs are in an scFv;
 - [0268] c. inhibit the human IL-31-induced signaling with an IC₅₀ of 0.1 to 30 ng/ml, particularly with an IC₅₀ of 0.1 to 20 ng/ml, particularly with an IC₅₀ of 0.1 to 10 ng/ml, as measured in a Path Hunter IL-31RA/OSMRb dimerization assay, wherein said IL31-BDs are in an scFv format; and/or
 - [0269] d. block the binding of human IL-31 to human IL-31R with an IC₅₀ of 0.1 to 20 ng/ml, particularly with an IC₅₀ of 0.1 to 10 ng/ml, particularly with an IC₅₀ of 0.1 to 6 ng/ml, as measured in a competition ELISA, wherein said IL31-BDs are in an scFv format.
- [0270] Suitably, the IL31-BDs of the multispecific antibodies of the present invention are further characterized by one or more of the following parameters:
- [0271] e. when being in scFv format, have a melting temperature (T_m), determined by differential scanning fluorimetry, of at least 65° C., preferably of at least 67° C., more preferably at least 69° C., in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
 - [0272] f. when being in scFv format, have a loss in monomer content, after storage for four weeks at 4° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
 - [0273] g. when being in scFv format, have a loss in monomer content, after storage for four weeks at 40° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4;
 - [0274] h. when being in scFv format, have a loss in monomer content, after 3 freeze-thawing cycles, of less than 5%, e. g. less than 4%, less than 3%, less than 2%,

preferably less than 1%, when said scFvs are at a starting concentration of 10 mg/ml, and in particular wherein said scFvs are formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4; and/or

[0275] i. when being in scFv format, have a loss in protein content, after storage for four weeks at 4° C. or 40° C. of less than 5%, e. g. less than 4%, less than 3%, less than 2%, preferably less than 1%, when said scFv is at a starting concentration of 10 mg/ml, and in particular wherein said scFv is formulated in 50 mM phosphate citrate buffer with 150 mM NaCl at pH 6.4.

[0276] Suitably, the IL31-BDs of the multispecific antibodies of the invention are binding domains provided in the present disclosure. The IL31-BDs of the multispecific antibodies of the invention include, but are not limited to, the humanized IL31-BDs whose sequences are listed in Table 2.

[0277] The term “multivalent antibody” refers to a single binding molecule with more than one valency, where “valency” is described as the number of antigen-binding moieties that binds to epitopes on target molecules. As such, the single binding molecule can bind to more than one binding site on a target molecule and/or to more than one target molecule due to the presence of more than one copy of the corresponding antigen-binding moieties. Examples of multivalent antibodies include, but are not limited to bivalent antibodies, trivalent antibodies, tetravalent antibodies, pentavalent antibodies, hexavalent antibodies, and the like.

[0278] The term “monovalent antibody”, as used herein, refers to an antibody that binds to a single target molecule, and more specifically to a single epitope on a target molecule. Also, the term “binding domain” or “monovalent binding domain”, as used herein, refers to a binding domain that binds to a single epitope on a target molecule.

[0279] In particular embodiments, the multispecific antibodies of the invention comprise one IL4R-BD and one IL-31-BD, i. e. the multispecific antibodies of the invention are monovalent for both IL-4R and IL-31 specificity.

[0280] In further particular embodiments, the multispecific antibodies of the invention comprise one IL4R-BD and two IL-31-BDs, i. e. the multispecific antibodies of the invention are monovalent for IL-4R specificity and bivalent for IL-31 specificity.

[0281] In further particular embodiments, the multispecific antibodies of the invention comprise two IL4R-BDs and one IL-31-BD, i. e. the multispecific antibodies of the invention are bivalent for IL-4R specificity and monovalent for IL-31 specificity.

[0282] In preferred embodiments, the multispecific antibodies of the invention comprise two IL4R-BDs and two IL-31-BDs, i. e. the multispecific antibodies of the invention are bivalent for IL-4R specificity and bivalent for IL-31 specificity.

[0283] In case the multispecific antibodies of the invention comprise two IL4R-BDs, said two IL4R-BDs either bind the same epitope or different epitopes on the IL-4R target molecule. Preferably, the two IL4R-BDs bind the same epitope on the IL-4R target molecule. Suitably, the multispecific antibody of the present invention comprises two identical IL4R-BDs

[0284] In case the multispecific antibodies of the invention comprise two IL31-BDs, said two IL31-BDs either bind the same epitope or different epitopes on the IL-31 target molecule. Preferably, the two IL31-BDs bind the same

epitope on the IL-31 target molecule. Suitably, the multispecific antibody of the present invention comprises two identical IL31-BDs.

[0285] The term “same epitope”, as used herein, refers to an individual protein determinant on the protein capable of specific binding to more than one antibody, where that individual protein determinant is identical, i. e. consist of identical chemically active surface groupings of molecules such as amino acids or sugar side chains having identical three-dimensional structural characteristics, as well as identical charge characteristics for each of said antibodies. The term “different epitope”, as used herein in connection with a specific protein target, refers to individual protein determinants on the protein, each capable of specific binding to a different antibody, where these individual protein determinants are not identical for the different antibodies, i. e. consist of non-identical chemically active surface groupings of molecules such as amino acids or sugar side chains having different three-dimensional structural characteristics, as well as different charge characteristics. These different epitopes can be overlapping or non-overlapping.

[0286] In particular embodiments, the multispecific antibodies of the invention are bispecific and bivalent.

[0287] In further particular embodiments, the multispecific antibodies of the invention are bispecific and trivalent.

[0288] Preferably, the multispecific antibodies of the invention are bispecific and tetravalent, i. e. bivalent for IL-4R specificity and bivalent for IL-31 specificity.

[0289] In a particular aspect, the present invention relates to a multispecific antibody comprising:

[0290] a) two binding domains, which specifically bind to IL-4R (IL4R-BD), and

[0291] b) two binding domains, which specifically bind to IL-31 (IL31-BD),

[0292] wherein

[0293] said multispecific antibody comprises an IgG region;

[0294] each of said IL4R-BDs comprise

[0295] the HCDR1 sequence of SEQ ID NO: 1,

[0296] the HCDR2 sequences of SEQ ID NO: 2,

[0297] the HCDR3 sequence of SEQ ID NO: 3,

[0298] the LCDR1 sequence of SEQ ID NO: 7,

[0299] the LCDR2 sequence of SEQ ID NO: 8, and

[0300] the LCDR3 sequences of SEQ ID NO: 9; and

[0301] each of said IL31-BDs comprise

[0302] the HCDR1 sequence of SEQ ID NO: 12,

[0303] the HCDR2 of sequence of SEQ ID NO: 13,

[0304] the HCDR3 sequences of SEQ ID NO: 14,

[0305] the LCDR1 sequence of SEQ ID NO: 17 or 18,

[0306] the LCDR2 sequence of SEQ ID NO: 19, and

[0307] the LCDR3 sequences of SEQ ID NOs: 20.

[0308] Other variable domains used in the invention include amino acid sequences that have been mutated, yet have at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identity in the CDR regions with the CDR regions depicted in the sequences described in Tables 1 and 2. Other variable domains used the invention include mutant amino acid sequences wherein no more than 1, 2, 3, 4 or 5 amino acids have been mutated in the CDR regions when compared with the CDR regions depicted in the sequence described in Tables 1 and 2.

[0309] Suitably, the VH domains of the binding domains of the multispecific antibodies of the invention belong to the

VH3 or the VH4 family. In one embodiment, a binding domain used in the invention comprises a VH domain belonging to the VH3 family. In the context of the present invention, the term “belonging to the VHx family (or the VLx family)” means that the framework sequences FR1 to FR3 show the highest degree of homology to said VHx family (or VLx family, respectively). Examples of VH and VL families are given in Knappik et al., J. Mol. Biol. 296 (2000) 57-86, or in WO 2019/057787. A specific example of a VH domain belonging to the VH3 family is represented by SEQ ID NO: 23, and a specific example of a VH domain belonging to VH4 family is represented by SEQ ID NO: 24. In particular, framework regions FR1 to FR3 taken from SEQ ID NO: 23 belong to the VH3 family (Table 3, regions marked in non-bold). Suitably, a VH belonging to the VH3 family, as used herein, is a VH comprising FR1 to FR3 having at least 85%, particularly at least 90%, more particularly at least 95% sequence identity to FR1 to FR3 of SEQ ID NO: 23. Alternative examples of VH3 and VH4 sequences, and examples of other VHx sequences, may be found in Knappik et al., J. Mol. Biol. 296 (2000) 57-86 or in WO 2019/057787.

[0310] Suitably, the VL domains of the binding domains used in the invention comprise: VK frameworks FR1, FR2 and FR3, particularly VK1 or VK3 frameworks, particularly VK1 frameworks FR1 to FR3, and a framework FR4, which is selected from a VK FR4. In the case where said binding domains are in an scFv-format, said binding domains comprise: VK frameworks FR1, FR2 and FR3, particularly VK1 or VK3 frameworks, particularly VK1 frameworks FR1 to FR3, and a framework FR4, which is selected from a VK FR4 and a VA FR4, particularly a VA FR4.

[0311] Suitably VK1 frameworks FR1 to FR3 as well as an exemplary VA FR4 are set forth in SEQ ID NO: 25 (Table 3, FR regions are marked in non-bold). Alternative examples of VK1 sequences, and examples of VK2, VK3 or VK4 sequences, may be found in Knappik et al., J. Mol. Biol. 296 (2000) 57-86. Suitably VK1 frameworks FR1 to FR3 comprise the amino acid sequences having at least 70, 80, 90 percent identity to amino acid sequences corresponding to FR1 to FR3 and taken from SEQ ID NO: 25 (Table 3, FR regions are marked in non-bold). Suitably VA FR4 are as set forth in SEQ ID NO: 26 to SEQ ID NO: 32 and in SEQ ID NO: 33 comprising a single cysteine residue, particular in a case where a second single cysteine is present in the corresponding VH chain, particularly in position 51 (AHO numbering) of VH, for the formation of an inter-domain disulfide bond. In one embodiment, the VL domains of the binding domains of the multispecific antibodies of the invention, when being in scFv-format, comprises VA FR4 comprising the amino acid sequence having at least 70, 80, 90 percent identity to an amino acid sequence selected from any of SEQ ID NO: 26 to SEQ ID NO: 33, particularly to SEQ ID NO: 26 or 33.

[0312] The binding domains of the multispecific antibodies of the invention comprise a VH domain listed in Tables 1 and 2. Suitably, the binding domains used the invention comprise a VH amino acid sequence listed in one of Tables 1 and 2, wherein no more than 20 amino acids in the framework sequences (i. e., the sequences which are not CDR sequences) have been mutated (wherein a mutation is, as various non-limiting examples, an addition, substitution or deletion). Suitably, the binding domains of the multispecific antibodies of the invention comprise a VH amino acid

sequence listed in one of Tables 1 and 2, wherein no more than 15 amino acids, particularly no more than 10 amino acids, particularly no more than 5 amino acids in the framework sequences (i. e., the sequences which are not CDR sequences) have been mutated (wherein a mutation is, as various non-limiting examples, an addition, substitution or deletion). Other binding domains used in the invention include amino acids that have been mutated, yet have at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identity in the VH regions with the VH regions depicted in the corresponding sequences described in one of Tables 1 and 2, including VH domains comprising at least positions 5 to 140 (AHO numbering), particularly at least positions 3 to 145 of one of the sequences shown in Tables 1 and 2.

[0313] In particular, the binding domains of the multispecific antibodies of the invention comprise a VL domain listed in one of Tables 1 and 2. Suitably, the binding domains of the multispecific antibodies of the invention comprise a VL amino acid sequence listed in one of Tables 1 and 2, wherein no more than about 20 amino acids in the framework sequences (i. e., the sequences which are not CDR sequences) have been mutated (wherein a mutation is, as various non-limiting examples, an addition, substitution or deletion). Other binding domains used in the invention include amino acids that have been mutated, yet have at least 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identity in the VL regions with a VL region depicted in the sequences described in Tables 1 and 2, including VL domains comprising at least positions 5 to 140 (AHO numbering), particularly at least positions 3 to 145 of one of the sequences shown in Tables 1 and 2.

[0314] In the context of the present invention, the term “binding domain used in the present invention” relates to a binding domain as such, i. e. independent of a multispecific context, and, in particular, to a binding domain comprised in a multispecific construct, e. g. one of the binding domains comprised in a bispecific, trispecific or tetraspecific construct.

[0315] Suitably, the binding domains of the multispecific antibodies of the invention are selected from the group consisting of: a Fab, an Fv, a dsFv and an scFv.

[0316] Suitably, the binding domains of the multispecific antibodies of the invention are operably linked. The binding domains of the multispecific antibodies of the invention are capable of binding to their respective antigens or receptors simultaneously. The term “simultaneously”, as used in this connection refers to the simultaneous binding of at least one of the IL4R-BDs and at least one of the IL31-BDs.

[0317] The multispecific antibodies of the present invention comprising one or two IL4R-BDs and one or two IL31-BDs, wherein said one or two IL4R-BDs, and said one or two IL31-BDs are operably linked to each other.

[0318] The term “operably linked”, as used herein, indicates that two molecules (e. g., polypeptides, domains, binding domains) are attached in a way that each molecule retains functional activity. Two molecules can be “operably

linked” whether they are attached directly or indirectly (e. g., via a linker, via a moiety, via a linker to a moiety). The term “linker” refers to a peptide or other moiety that is optionally located between binding domains or antibody fragments used in the invention. A number of strategies may be used to covalently link molecules together. These include, but are not limited to, polypeptide linkages between N- and C-termini of proteins or protein domains, linkage via disulfide bonds, and linkage via chemical cross-linking reagents. In one aspect of this embodiment, the linker is a peptide bond, generated by recombinant techniques or peptide synthesis. Choosing a suitable linker for a specific case where two polypeptide chains are to be connected depends on various parameters, including but not limited to the nature of the two polypeptide chains (e. g., whether they naturally oligomerize), the distance between the N- and the C-termini to be connected if known, and/or the stability of the linker towards proteolysis and oxidation. Furthermore, the linker may contain amino acid residues that provide flexibility.

[0319] In the context of the present invention, the term “polypeptide linker” refers to a linker consisting of a chain of amino acid residues linked by peptide bonds that is connecting two domains, each being attached to one end of the linker. The polypeptide linker should have a length that is adequate to link two molecules in such a way that they assume the correct conformation relative to one another so that they retain the desired activity. In particular embodiments, the polypeptide linker has a continuous chain of between 2 and 30 amino acid residues (e. g., 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 amino acid residues). In addition, the amino acid residues selected for inclusion in the polypeptide linker should exhibit properties that do not interfere significantly with the activity of the polypeptide. Thus, the linker peptide on the whole should not exhibit a charge that would be inconsistent with the activity of the polypeptide, or interfere with internal folding, or form bonds or other interactions with amino acid residues in one or more of the monomers that would seriously impede the binding of receptor monomer domains. In particular embodiments, the polypeptide linker is non-structured polypeptide. Useful linkers include glycine-serine, or GS linkers. By “Gly-Ser” or “GS” linkers is meant a polymer of glycines and serines in series (including, for example, (Gly-Ser)_n, (GSGGS)_n, (GGGGS)_n, and (GGGS)_n, where n is an integer of at least one), glycine-alanine polymers, alanine-serine polymers, and other flexible linkers such as the tether for the shaker potassium channel, and a large variety of other flexible linkers, as will be appreciated by those in the art. Glycine-serine polymers are preferred since oligopeptides comprising these amino acids are relatively unstructured, and therefore may be able to serve as a neutral tether between components. Secondly, serine is hydrophilic and therefore able to solubilize what could be a globular glycine chain. Third, similar chains have been shown to be effective in joining subunits of recombinant proteins such as single-chain antibodies.

[0320] In one group of embodiments, the multispecific antibody of the invention is in a format selected from (scFv)₂-Fc-(scFv)₂ fusion (ADAPTIR); DVD-Ig; a DART™ and a TRIDENT™. The term “DART™” refers to an antibody format developed by MacroGenics that comprises an immunoglobulin Fc region polypeptide and one or two bispecific Fv binding domains fused to the N-terminus of

one heavy chain of the Fc region or to both N-termini of the heavy chains of the Fc region. The term “TRIDENT™” refers to an antibody format developed by MacroGenics that comprises an immunoglobulin Fc region polypeptide, one bispecific Fv binding domain and one Fab fragment. Both the bispecific Fv binding domain and the Fab fragment are fused to the respective N-termini of the two heavy chains of the Fc region polypeptide.

[0321] In another group of embodiments, the format of the multispecific antibodies of the present invention is selected from bivalent bispecific IgG formats, trivalent bispecific IgG formats and tetravalent bispecific IgG formats. In particular, the format of said multispecific antibodies is selected from KiH-based IgGs, such as DuoBodies (bispecific IgGs prepared by the Duobody technology) (MAbs. 2017 February/March; 9(2):182-212. doi: 10.1080/19420862.2016.1268307); DVD-Ig; IgG-scFv fusions, such as CODV-IgG, Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)), bsAb (scFv linked to C-terminus of light chain), Bs1Ab (scFv linked to N-terminus of light chain), Bs2Ab (scFv linked to N-terminus of heavy chain), Bs3Ab (scFv linked to C-terminus of heavy chain), Ts1Ab (scFv linked to N-terminus of both heavy chain and light chain) and Ts2Ab (dsScFv linked to C-terminus of heavy chain). More particularly, the format of said multispecific antibody is selected from KiH-based IgGs, such as DuoBodies; DVD-Ig; CODV-IgG and Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)), even more particularly from DVD-Ig and Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)).

[0322] In particular embodiments of the invention, the format of said multispecific antibodies is selected from Morrison formats, i. e. a Morrison-L and a Morrison-H format. The Morrison-L and Morrison-H formats used in the present invention are tetravalent and bispecific molecular formats bearing an IgG Fc region, in particular an IgG4 Fc region. Two highly stable scFv binding domains, wherein the light chain comprises VK FR1 to FR3 in combination with a VA FR4 (A-cap), herein also called A-cap scFv, are fused via a linker L1 to the heavy chain (Morrison-H) or light chain (Morrison-L) C-termini (see FIG. 2).

[0323] The linker L1 is a peptide of 2-30 amino acids, more particularly 5-25 amino acids, and most particularly 10-20 amino acids. In particular embodiments, said linker L1 comprises one or more units of four (4) glycine amino acid residues and one (1) serine amino acid residue (GGGGS)_n, wherein n=1, 2, 3, 4 or 5, particularly n=2.

[0324] In a special embodiment of the invention, the multispecific antibodies have a Morrison-L format, as defined above. In another special embodiment of the invention, the multispecific antibodies have a Morrison-H format, as defined above.

[0325] The scFv domains of the multispecific antibodies of the invention comprise a variable heavy chain domain (VH) and a variable light chain domain (VL) connected by a linker L2.

[0326] The linker L2 is a peptide of 10-40 amino acids, more particularly 15-30 amino acids, and most particularly 20-25 amino acids. In particular embodiments, said linker L2 comprises one or more units of four (4) glycine amino acid residues and one (1) serine amino acid residue (GGGGS)_n, wherein n=1, 2, 3, 4, 5, 6, 7 or 8, particularly n=4.

[0327] Specific but non-limiting examples the multispecific antibodies of the invention are the Morrison-L antibodies PRO2206, PRO2207, PRO2208 and the Morrison-H antibodies PRO2198, PRO2199, PRO2200, PRO2201, whose sequences are listed in table 4.

[0328] The multispecific antibodies of the invention can be produced using any convenient antibody-manufacturing method known in the art (see, e. g., Fischer, N. & Leger, O., *Pathobiology* 74 (2007) 3-14 with regard to the production of bispecific constructs; Hornig, N. & Farber-Schwarz, A., *Methods Mol. Biol.* 907 (2012) 713-727, and WO 99/57150 with regard to bispecific diabodies and tandem scFvs). Specific examples of suitable methods for the preparation of the bispecific construct further include, inter alia, the Genmab (see Labrijn et al., *Proc. Natl. Acad. Sci. USA* 110 (2013) 5145-5150) and Merus (see de Kruif et al., *Biotechnol. Bioeng.* 106 (2010) 741-750) technologies. Methods for production of bispecific antibodies comprising a functional antibody Fc part are also known in the art (see, e. g., Zhu et al., *Cancer Lett.* 86 (1994) 127-134; and Suresh et al., *Methods Enzymol.* 121 (1986) 210-228).

[0329] These methods typically involve the generation of monoclonal antibodies, for example by means of fusing myeloma cells with the spleen cells from a mouse that has been immunized with the desired antigen using the hybridoma technology (see, e. g., Yokoyama et al., *Curr. Protoc. Immunol.* Chapter 2, Unit 2.5, 2006) or by means of recombinant antibody engineering (repertoire cloning or phage display/yeast display) (see, e. g., Chames & Baty, *FEMS Microbiol. Letters* 189 (2000) 1-8), and the combination of the antigen-binding domains or fragments or parts thereof of two or more different monoclonal antibodies to give a bispecific or multispecific construct using known molecular cloning techniques.

[0330] The multispecific antibodies of the invention can be prepared by conjugating the constituent binding specificities, using methods known in the art. For example, each binding specificity of the bispecific molecule can be generated separately and then conjugated to one another. When the binding specificities are proteins or peptides, a variety of coupling or cross-linking agents can be used for covalent conjugation. Examples of cross-linking agents include protein A, carbodiimide, N-succinimidyl-5-acetyl-thioacetate (SATA), 5,5'-dithiobis (2-nitrobenzoic acid) (DTNB), o-phenylenedimaleimide (oPDM), N-succinimidyl-3-(2-pyridyldithio)propionate (SPDP), and sulfo-succinimidyl 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (sulfo-SMCC) (see e. g., Karpovsky et al., 1984 *J. Exp. Med.* 160: 1686; Liu, M A et al., 1985 *Proc. Natl. Acad. Sci. USA* 82:8648). Other methods include those described in Paulus, 1985 *Behring Ins. Mitt. No.* 78, 118-132; Brennan et al., 1985 *Science* 229:81-83, and Glennie et al., 1987 *J. Immunol.* 139: 2367-2375. Conjugating agents are SATA and sulfo-SMCC, both available from Pierce Chemical Co. (Rockford, Ill, USA).

[0331] When the binding specificities are antibodies, they can be conjugated by sulfhydryl bonding of the C-terminus hinge regions of the two heavy chains. In a particular embodiment, the hinge region is modified to contain an odd number of sulfhydryl residues, for example one, prior to conjugation.

[0332] Alternatively, two or more binding specificities can be encoded in the same vector and expressed and assembled in the same host cell. This method is particularly useful

where the bispecific molecule is a mAb $\times$ Fab, a mAb $\times$ scFv, a mAb $\times$ dsFv or a mAb $\times$ Fv fusion protein. Methods for preparing multispecific antibodies and molecules are described for example in U.S. Pat. Nos. 5,260,203; 5,455,030; 4,881,175; 5,132,405; 5,091,513; 5,476,786; 5,013,653; 5,258,498; and 5,482,858.

[0333] Binding of the multispecific antibodies to their specific targets can be confirmed by, for example, enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (REA), FACS analysis, bioassay (e. g., growth inhibition), or Western Blot assay. Each of these assays generally detects the presence of protein-antibody complexes of particular interest by employing a labeled reagent (e. g., an antibody) specific for the complex of interest.

[0334] In a further aspect, the invention provides a nucleic acid or two nucleic acids encoding the multispecific antibody of the invention. Such nucleic acids can be optimized for expression in mammalian cells.

[0335] The term "nucleic acid" is used herein interchangeably with the term "polynucleotide(s)" and refers to one or more deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. The term encompasses nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties when compared to the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphorates, 2-O-methyl ribonucleotides, peptide-nucleic acids (PNAs). Unless otherwise indicated, a particular nucleic acid also implicitly encompasses conservatively modified variants thereof (e. g., degenerate codon substitutions) and nucleic acids with complementary sequences, as well as the nucleic acid with the sequence explicitly indicated. Specifically, as detailed below, degenerate codon substitutions may be achieved by generating nucleic acids in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.* 19:5081, 1991; Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608, 1985; and Rossolini et al., *Mol. Cell. Probes* 8:91-98, 1994).

[0336] The invention provides substantially purified nucleic acid molecules which encode polypeptides comprising segments or domains of the multispecific antibody described above. When expressed from appropriate expression vectors, polypeptides encoded by these nucleic acid molecules are capable of exhibiting antigen-binding capacities of the multispecific antibody of the present invention.

[0337] The polynucleotide sequences can be produced by de novo solid-phase DNA synthesis or by PCR mutagenesis of an existing sequence (e. g., sequences as described in the Examples below) encoding the multispecific antibody of the invention or fragments thereof or binding domains thereof. Direct chemical synthesis of nucleic acids can be accomplished by methods known in the art, such as the phosphotriester method of Narang et al., 1979, *Meth. Enzymol.* 68:90; the phosphodiester method of Brown et al., *Meth. Enzymol.* 68: 109, 1979; the diethylphosphoramidite method of Beaucage et al., *Tetra. Lett.*, 22: 1859, 1981; and the solid support method of U.S. Pat. No. 4,458,066. Introducing mutations to a polynucleotide sequence by PCR can be performed as described in, e. g., PCR Technology:

Principles and Applications for DNA Amplification, H. A. Erlich (Ed.), Freeman Press, NY, N.Y., 1992; PCR Protocols: A Guide to Methods and Applications, Innis et al. (Ed.), Academic Press, San Diego, Calif, 1990; Mattila et al., Nucleic Acids Res. 19:967, 1991; and Eckert et al., PCR Methods and Applications 1:17, 1991.

[0338] Also provided in the invention are expression vectors and host cells for producing the multispecific antibody of the invention or fragments thereof or binding domains thereof.

[0339] The term “vector” is intended to refer to a polynucleotide molecule capable of transporting another polynucleotide to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e. g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e. g., non-episomal mammalian vectors) can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome.

[0340] Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e. g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions. In this particular context, the term “operably linked” refers to a functional relationship between two or more polynucleotide (e. g., DNA) segments. Typically, it refers to the functional relationship of a transcriptional regulatory sequence to a transcribed sequence. For example, a promoter or enhancer sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, i. e., they are cis-acting. However, some transcriptional regulatory sequences, such as enhancers, need not be physically contiguous or located in close proximity to the coding sequences whose transcription they enhance.

[0341] Various expression vectors can be employed to express the polynucleotides encoding the multispecific antibody chains. Both viral-based and non-viral expression vectors can be used to produce the antibodies in a mammalian host cell. Non-viral vectors and systems include plasmids, episomal vectors, typically with an expression cassette for expressing a protein or RNA, and human artificial chromosomes (see, e. g., Harrington et al., Nat Genet. 15:345, 1997). For example, non-viral vectors useful for expression of the polynucleotides encoding the multispecific antibody chains in mammalian (e. g., human) cells include pThioHis A, B and C, pcDNA3.1/His, pEBVHis A, B and C, (Invitrogen, San Diego, CA, USA), MPS V vectors, and

numerous other vectors known in the art for expressing other proteins. Useful viral vectors include vectors based on retroviruses, adenoviruses, adeno-associated viruses, herpes viruses, vectors based on SV40, papilloma virus, HBP Epstein Barr virus, vaccinia virus vectors and Semliki Forest virus (SFV). See, Brent et al., supra; Smith, Annu. Rev. Microbiol. 49:807, 1995; and Rosenfeld et al., Cell 68: 143, 1992.

[0342] The choice of expression vector depends on the intended host cells in which the vector is to be expressed. Typically, the expression vectors contain a promoter and other regulatory sequences (e. g., enhancers) that are operably linked to the polynucleotides encoding a multispecific antibody chain or a fragment. In one embodiment, an inducible promoter is employed to prevent expression of inserted sequences except under inducing conditions. Inducible promoters include, e. g., arabinose, lacZ, metallothionein promoter or a heat shock promoter. Cultures of transformed organisms can be expanded under non-inducing conditions without biasing the population for coding sequences whose expression products are better tolerated by the host cells. In addition to promoters, other regulatory elements may also be required or desired for efficient expression of a multispecific antibody chain or a fragment. These elements typically include an ATG initiation codon and adjacent ribosome binding site or other sequences. In addition, the efficiency of expression may be enhanced by the inclusion of enhancers appropriate to the cell system in use (see, e. g., Scharf et al., Results Probl. Cell Differ. 20: 125, 1994; and Bittner et al., Meth. Enzymol., 153:516, 1987). For example, the SV40 enhancer or CMV enhancer may be used to increase expression in mammalian host cells.

[0343] Vectors to be used typically encode the multispecific antibody light and heavy chain including constant regions or parts thereof. Such vectors allow expression of the variable regions as fusion proteins with the constant regions thereby leading to production of intact antibodies and antigen-binding fragments thereof. Typically, such constant regions are human.

[0344] The term “recombinant host cell” (or simply “host cell”) refers to a cell into which a recombinant expression vector has been introduced. It should be understood that such terms are intended to refer not only to the particular subject cell but to the progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term “host cell” as used herein.

[0345] The host cells for harboring and expressing the multispecific antibody of the invention can be either prokaryotic or eukaryotic. *E. coli* is one prokaryotic host useful for cloning and expressing the polynucleotides of the present invention. Other microbial hosts suitable for use include bacilli, such as *Bacillus subtilis*, and other enterobacteriaceae, such as *Salmonella*, *Serratia*, and various *Pseudomonas* species. In these prokaryotic hosts, one can also make expression vectors, which typically contain expression control sequences compatible with the host cell (e. g., an origin of replication). In addition, any number of a variety of well-known promoters will be present, such as the lactose promoter system, a tryptophan (trp) promoter system, a beta-lactamase promoter system, or a promoter system from phage lambda. The promoters typically control expression,

optionally with an operator sequence, and have ribosome binding site sequences and the like, for initiating and completing transcription and translation. Other microbes, such as yeast, can also be employed to express the multispecific antibodies of the invention. Insect cells in combination with baculovirus vectors can also be used.

[0346] In one embodiment, mammalian host cells are used to express and produce the multispecific antibody of the invention. For example, they can be either a hybridoma cell line expressing endogenous immunoglobulin genes or a mammalian cell line harboring an exogenous expression vector. These include any normal mortal or normal or abnormal immortal animal or human cell. For example, a number of suitable host cell lines capable of secreting intact immunoglobulins have been developed including the CHO cell lines, various COS cell lines, HeLa cells, myeloma cell lines, transformed B-cells and hybridomas. The use of mammalian tissue cell culture to express polypeptides is discussed generally in, e. g., Winnacker, *FROM GENES TO CLONES*, VCH Publishers, N.Y., N.Y., 1987. Expression vectors for mammalian host cells can include expression control sequences, such as an origin of replication, a promoter, and an enhancer (see, e. g., Queen, et al., *Immunol. Rev.* 89:49-68, 1986), and necessary processing information sites, such as ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. These expression vectors usually contain promoters derived from mammalian genes or from mammalian viruses. Suitable promoters may be constitutive, cell type-specific, stage-specific, and/or modulatable or regulatable. Useful promoters include, but are not limited to, the metallothionein promoter, the constitutive adenovirus major late promoter, the dexamethasone-inducible MMTV promoter, the SV40 promoter, the MRP polIII promoter, the constitutive MPS V promoter, the tetracycline-inducible CMV promoter (such as the human immediate-early CMV promoter), the constitutive CMV promoter, and promoter-enhancer combinations known in the art.

[0347] Methods for introducing expression vectors containing the polynucleotide sequences of interest vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts. (See generally Green, M. R., and Sambrook, J., *Molecular Cloning: A Laboratory Manual* (Fourth Edition), Cold Spring Harbor Laboratory Press (2012)). Other methods include, e. g., electroporation, calcium phosphate treatment, liposome-mediated transformation, injection and microinjection, ballistic methods, virosomes, immunoliposomes, polycation-nucleic acid conjugates, naked DNA, artificial virions, fusion to the herpes virus structural protein VP22 (Elliot and O'Hare, *Cell* 88:223, 1997), agent-enhanced uptake of DNA, and ex vivo transduction. For long-term, high-yield production of recombinant proteins, stable expression will often be desired. For example, cell lines which stably express the multispecific antibody of the invention can be prepared using expression vectors of the invention which contain viral origins of replication or endogenous expression elements and a selectable marker gene. Following the introduction of the vector, cells may be allowed to grow for 1 to 2 days in an enriched media before they are switched to selective media. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth of

cells which successfully express the introduced sequences in selective media. Resistant, stably transfected cells can be proliferated using tissue culture techniques appropriate to the cell type. The present invention thus provides a method of producing the antibody of the invention or antigen-binding fragment thereof, wherein said method comprises the step of culturing a host cell comprising a nucleic acid or a vector encoding the antibody of the invention or antigen-binding fragment thereof, whereby said antibody of the disclosure or a fragment thereof is expressed.

[0348] In one aspect, the present invention relates to a method of producing the multispecific antibody of the invention, the method comprising the step of culturing a host cell expressing a nucleic acid encoding the multispecific antibody of the invention. In particular, the present invention relates to a method of producing the multispecific antibody of the invention, the method comprising (i) providing a nucleic acid or two nucleic acids encoding the multispecific antibody of the invention or a vector or two vectors encoding the multispecific antibody of the invention, expressing said nucleic acid or nucleic acids, or said vector or vectors, and collecting said multispecific antibody or said binding domain from the expression system, or (ii) providing a host cell or host cells expressing a nucleic acid or two nucleic acids encoding the multispecific antibody of the invention, culturing said host cell or said host cells; and collecting said multispecific antibody or said binding domain from the cell culture.

[0349] In a further aspect, the present invention relates to a pharmaceutical composition comprising the multispecific antibody of the invention, and a pharmaceutically acceptable carrier. "Pharmaceutically acceptable carrier" means a medium or diluent that does not interfere with the structure of the antibodies. Pharmaceutically acceptable carriers enhance or stabilize the composition, or facilitate preparation of the composition. Pharmaceutically acceptable carriers include solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible.

[0350] Certain, of such carriers enable pharmaceutical compositions to be formulated as, for example, tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspension and lozenges for the oral ingestion by a subject. Certain of such carriers enable pharmaceutical compositions to be formulated for injection, infusion or topical administration. For example, a pharmaceutically acceptable carrier can be a sterile aqueous solution.

[0351] The pharmaceutical composition of the invention can be administered by a variety of methods known in the art. The route and/or mode of administration vary depending upon the desired results. Administration can be intravenous, intramuscular, intraperitoneal, or subcutaneous, or administered proximal to the site of the target. In particular embodiments, the administration is intramuscular, or subcutaneous, particularly subcutaneous. The pharmaceutically acceptable carrier should be suitable for intravenous, intramuscular, subcutaneous, parenteral, spinal or epidermal administration (e. g., by injection or infusion), particularly for intramuscular or subcutaneous administration. Depending on the route of administration, the active compound, i. e., the multispecific antibody of the invention, may be coated in a material to protect the compound from the action of acids and other natural conditions that may inactivate the compound.

[0352] The pharmaceutical compositions of the invention can be prepared in accordance with methods well known and routinely practiced in the art. See, e. g., Remington: The Science and Practice of Pharmacy, Mack Publishing Co., 20th ed., 2000; and Sustained and Controlled Release Drug Delivery Systems, J. R. Robinson, ed., Marcel Dekker, Inc., New York, 1978. Pharmaceutical compositions are preferably manufactured under GMP conditions. Typically, a therapeutically effective dose or efficacious dose of the multispecific antibody of the invention is employed in the pharmaceutical compositions of the invention. The multispecific antibodies of the invention are formulated into pharmaceutically acceptable dosage forms by conventional methods known to those of skill in the art. Dosage regimens are adjusted to provide the optimum desired response (e. g., a therapeutic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subjects to be treated; each unit contains a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier.

[0353] Actual dosage levels of the active ingredients in the pharmaceutical compositions of the invention can be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired therapeutic response for a particular patient, composition, and mode of administration, without being toxic to the patient. The selected dosage level depends upon a variety of pharmacokinetic factors including the activity of the particular compositions of the present invention employed, the route of administration, the time of administration, the rate of excretion of the particular compound being employed, the duration of the treatment, other drugs, compounds and/or materials used in combination with the particular compositions employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors.

[0354] The multispecific antibody of the invention is usually administered on multiple occasions. Intervals between single dosages can be weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of the multispecific antibody of the invention in the patient. Alternatively, the multispecific antibody of the invention can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, humanized antibodies show longer half-life than that of chimeric antibodies and non-human antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows

partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[0355] In one aspect, the present invention relates to the multispecific antibody of the invention or the pharmaceutical composition of the invention for use as a medicament. In a suitable embodiment, the present invention provides the multispecific antibody or the pharmaceutical composition for use in the treatment of a disease selected from allergic, inflammatory and autoimmune diseases, particularly from inflammatory and autoimmune diseases.

[0356] In another aspect, the present invention provides the pharmaceutical composition for use in the manufacture of a medicament for the treatment of an allergic, inflammatory or autoimmune disease, particularly of an inflammatory or autoimmune disease.

[0357] In another aspect, the present invention relates to the use of the multispecific antibody or the pharmaceutical composition for treating an allergic, inflammatory or autoimmune disease, particularly for treating an inflammatory or autoimmune disease, in a subject in need thereof.

[0358] In another aspect, the present invention relates to a method of treating a subject comprising administering to the subject a therapeutically effective amount of the multispecific antibody of the present invention. In a suitable embodiment, the present invention relates to a method for the treatment of an allergic, inflammatory or autoimmune disease, particularly for treating an inflammatory or autoimmune disease, in a subject comprising administering to the subject a therapeutically effective amount of the multispecific antibody of the present invention.

[0359] The term “subject” includes human and non-human animals.

[0360] The term “animals” include all vertebrates, e. g., non-human mammals and non-mammals, such as non-human primates, sheep, dog, cow, chickens, amphibians, and reptiles. Except when noted, the terms “patient” or “subject” are used herein interchangeably.

[0361] The terms “treatment”, “treating”, “treat”, “treated”, and the like, as used herein, refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease or delaying the disease progression. “Treatment”, as used herein, covers any treatment of a disease in a mammal, e. g., in a human, and includes: (a) inhibiting the disease, i. e., arresting its development; and (b) relieving the disease, i. e., causing regression of the disease.

[0362] The term “therapeutically effective amount” or “efficacious amount” refers to the amount of an agent that, when administered to a mammal or other subject for treating a disease, is sufficient to affect such treatment for the disease. The “therapeutically effective amount” will vary depending on the agent, the disease and its severity and the age, weight, etc., of the subject to be treated.

[0363] In one embodiment, the allergic, inflammatory and autoimmune diseases are selected from pruritus-causing allergic diseases, pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases, particularly from pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases. The terms “pruritus” and “itch”, which are herein used synonymously, refers to a sensation that causes the desire or reflex to scratch. Scratching can be problematic particularly in the case of chronic

skin diseases that induce itching, such as atopic dermatitis, dermatomycoses, psoriasis or urticaria, as the permanent itching stimulates patients to scratch the affected skin areas constantly and/or excessively, which often leads to skin injuries and further deterioration of the skin surface.

[0364] The term “allergic diseases” or “allergies” as used herein refers to a large number of conditions caused by hypersensitivity of the immune system to typically harmless substances in the environment.

[0365] The term “inflammatory diseases” as used herein refers to a vast number of inflammatory disorders, i. e. inflammatory abnormalities, which are often characterized by prolonged inflammation, known as chronic inflammation. The term “inflammation” refers to the complex biological response of body tissues to harmful stimuli, such as pathogens, damaged cells, or irritants. It is a protective response involving immune cells, blood vessels, and molecular mediators. While regular inflammation reactions are essential for the body to eliminate the initial cause of cell injury, clear out necrotic cells and tissues damaged from the original insult and the inflammatory process, and to initiate tissue repair, inflammatory disorders are generally characterized by persistent inflammation in the absence of harmful stimuli.

[0366] The term “autoimmune diseases” as used herein refers to a condition arising from an abnormal immune response to a functioning body part. It is the result of autoimmunity, i. e. the presence of self-reactive immune response (e. g., autoantibodies, self-reactive T cells), with or without damage or pathology resulting from it, which is typically restricted to certain organs or involve a particular tissue in different places.

[0367] In particular embodiments, the pruritus-causing inflammatory or autoimmune disease is selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma, in particular from atopic dermatitis.

[0368] In another aspect, the allergic, inflammatory and autoimmune diseases are selected from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urti-

caria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma, in particular from atopic dermatitis.

[0369] In another embodiment, the allergic, inflammatory and autoimmune diseases are selected from allergic asthma, allergic rhinitis, inflammatory airway disease, recurrent airway obstruction, airway hyperresponsiveness, chronic obstruction pulmonary disease, Crohn disease, chronic non-histamine related urticaria, antihistamine-unresponsive mastocytosis, lichen simplex chronicus, seborrheic dermatitis, xeroderma, Dermatitis herpetiformis, lichen planus and ulcerative colitis.

[0370] In another embodiment, the present invention provides the multispecific antibody or the pharmaceutical composition, as defined herein, for use in the treatment of a disease, which is a neuropathic pruritus disease selected from post-herpetic neuralgia, post-herpetic itch, notalgia paresthetica, multiple sclerosis and brachioradial pruritus.

[0371] In another embodiment, the present invention provides the multispecific antibody or the pharmaceutical composition for use in an antipruritic agent for the treatment of a systemic disease with itching, which disease is selected from Cholestasis, chronic kidney disease, Hodgkin’s disease, cutaneous T-cell lymphoma and other lymphomas or leukemias associated with chronic itch, polycythemia vera, hyperthyroidism, chronic post-arthropod itch (Id reaction), pregnancy-induced chronic itch (e. g. PUPPP), eosinophilic pustular folliculitis, drug hypersensitivity reactions, chronic pruritus of the elderly or dry skin itch (local, generalized), and pruritus at the scar portion after burn (post-burn itch), genetic or nevoid chronic itches (e. g. Netherton syndrome, Darier’s disease (Morbus Darier), Hailey-Hailey disease, inflammatory linear verrucous epidermal nevus (ILVEN), familial primary cutaneous amyloidosis, Olmsted syndrome), aquagenic pruritus, fiberglass dermatitis, mucous chronic itch, chemotherapy-induced itch and HIV.

Sequence Listing (Mutations Designated According to AHo Numbering Scheme; the CDRs Defined According to Numab CDR Definition, Unless Specified Otherwise)

TABLE 1

Examples of IL-4R binding domains as used in the present invention (CDR residues shown in bold and italic letters).		
SEQ ID NUMBER	Ab region	Sequence
44-34-C10		
SEQ ID NO: 1	HCDR1 (H27-H42; AHo numbering)	<i>GFSSSSAHYMC</i>
SEQ ID NO: 2	HCDR2 (H57-H76; AHo numbering)	<i>CIYAGSRGSTYYASWAKG</i>
SEQ ID NO: 3	HCDR3 (H108-H138; AHo numbering)	<i>RAAFVNRRGVSWIWPYYFSL</i>
SEQ ID NO: 4	VH (44-34-C10-sc07/ 44-34-C10-sc08)	EVQLVESGGGLVQPGGSLRLSCAAS <i>GFSSSSAHYMC</i> WVRQAPGKGLEWIGCIYAGSRGSTYYASWAKGRFTISRDN SKNTVYLQMNSLR AEDTAVYYCARAAAFVNRRGVSWIWPYYFSLWGQGTLVTVSS
SEQ ID NO: 5	VH (44-34-C10-sc09)	EVQLVESGGGRVQPGGSLRLSCAAS <i>GFSSSSAHYMC</i> WVRQAPGKGLEWIGCIYAGSRGSTYYASWAKGRFTISKASSTTVY LQMNSLR AEDTATYYCARAAAFVNRRGVSWIWPYYFSLWGQGTVTVSS

TABLE 1-continued

Examples of IL-4R binding domains as used in the present invention (CDR residues shown in bold and italic letters).		
SEQ ID NUMBER	Ab region	Sequence
SEQ ID NO: 6	VH (44-34-C10-sc09) mutations R12L, T103V, Q144L	EVQLVESGGGLVQPGGSLRLSCAAS <i>GFSSSSAHYMC</i> WVRQAPGKLEWI GCT <i>YAGSRG</i> <i>STYYASWAK</i> GRFTISKASSTTVYLQMNSLR <i>AEDTAVYYCA</i> <i>RAAFVNRGVSWIWPYYFSL</i> WGQGLVTVSS
SEQ ID NO: 7	LCDR1 (L24-L42; AHo numbering)	<i>QASEDIESYLA</i>
SEQ ID NO: 8	LCDR2 (L58-L72; AHo numbering) (present in 44-34-C10-sc06, 44-34-C10-sc08 and 44-34-C10-sc09)	<i>RASTLAY</i>
SEQ ID NO: 9	LCDR3 (L107-L138; AHo numbering)	<i>LGSSYSTGSNI</i>
SEQ ID NO: 10	VL (44-34-C10-sc06/ 44-34-C10-sc09)	DIQMTQSPSSLSASVGDRTITTC <i>QASEDIESYLA</i> WYQQKPKAPKLLI <i>YR</i> <i>ASTLAY</i> GVPSRFRSGSGSDTFTLTIS <i>SLQPEDFATYYCLGSSYSTGSNI</i>
SEQ ID NO: 11	VL (44-34-C10-sc08)	DIQMTQSPSSLSARVGDRTITTC <i>QASEDIESYLA</i> WYQQKPKAPKLLI <i>YR</i> <i>ASTLAY</i> GVPSRFRSGSGGRDFTLTIS <i>SLQPEDFATYYCLGSSYSTGSNI</i>

TABLE 2

Examples of IL-31 binding domains as used in the present invention (CDR residues shown in bold and italic letters).		
SEQ ID NUMBER	Ab region	Sequence
50-09-D07		
SEQ ID NO: 12	HCDR1 (H27-H42; AHo numbering)	<i>GFSSFSANYWIC</i>
SEQ ID NO: 13	HCDR2 (H57-H76; AHo numbering) (present in 50-09-D07-sc04, 50-09-D07-sc05 and 50-09-D07-sc07)	<i>CIIYIGSRADTTYAPWAKG</i>
SEQ ID NO: 14	HCDR3 (H108-H138; AHo numbering)	<i>RFSSGIYDLDRFFL</i>
SEQ ID NO: 15	VH (50-09-D07-sc03/ 50-09-D07-sc07)	QSQLVESGGGLVQPGGSLRLSCAAS <i>GFSSFSANYWIC</i> WVRQAPGKG LEWIV <i>CIIYIGSRADTTYAPWAK</i> GRFTISKDNSKNTVYLQMNSLR <i>A</i> <i>EDTAVYFCARFSSGIYDLDRFFL</i> WGQGLVTVSS
SEQ ID NO: 16	VH (50-09-D07-sc04)	QSQLVESGGGRVQPGGSLRLSCAAS <i>GFSSFSANYWIC</i> WVRQAPGKG LEWIV <i>CIIYIGSRADTTYAPWAK</i> GRFTISKDNSKNTVYLQMNSLR <i>A</i> <i>EDTATYFCARFSSGIYDLDRFFL</i> WGQGTQVTVSS
SEQ ID NO: 17	LCDR1 (L24-L42; AHo numbering)	<i>QASQSIDSWLA</i>
SEQ ID NO: 18	LCDR1 (L24-L42; AHo numbering)	<i>QASQSIESWLA</i>
SEQ ID NO: 19	LCDR2 (L58-L72; AHo numbering)	<i>QASKLAS</i>
SEQ ID NO: 20	LCDR3 (L107-L138; AHo numbering)	<i>QTYYGGGHIGWG</i>
SEQ ID NO: 21	VL (50-09-D07-sc03 to sc06)	AFQLTQSPSSLSASVGDRTITTC <i>QASQSIDSWLA</i> WYQQKPKPK <i>P</i> LLI <i>YQASKLAS</i> GVPSRFRSGSGSDTYTLTIS <i>SLQPEDFATYYCQT</i> <i>YGGGGHIGWG</i>

TABLE 2-continued

Examples of IL-31 binding domains as used in the present invention (CDR residues shown in bold and italic letters).		
SEQ ID NUMBER	Ab region	Sequence
SEQ ID NO: 22	VL (50-09-D07-sc07)	AFQLTQSPSSLSASVGDRTITC <i>QASQSI</i> <i>ES</i> <i>NL</i> AWYQQKPGKPPK LLIY <i>QASKLAS</i> GVPSRFSGSGSGTDYTLTISSLPEDFATYYC <i>QT</i> <i>YGGGHI</i> GWG

TABLE 3

Other sequences related to the present invention.		
SEQ ID NUMBER	Ab region	Sequence
SEQ ID NO: 23	VH3	EVQLVESGGGLVQPGGSLRLSCAAS <i>GFSFS</i> <i>SANY</i> YPCWVRQAPGKG LEWIG <i>CTFYGGSSDITYDANWTK</i> GRFTISRDNSKNTVYLMNSLRA EDTAVYYCARS <i>AWYSGWGGDL</i> WGQGLTVTVSS
SEQ ID NO: 24	VH4	QVQLQESGPGLVKPSETLSLTCKV <i>S</i> <i>GFSFS</i> <i>NSY</i> WICWIRQPPGKG LEWIG <i>CTFVGGSDSTYYANWAK</i> GRVTISVDSSKNQFSLKLSVTA ADTAVYYCAR <i>RHPSDAVYGYANNL</i> WGQGLTVTVSS
SEQ ID NO: 25	Vkappa1	DIQMTQSPSSLSASVGDRTITC <i>QASQS</i> <i>INN</i> VLAWYQQKPGKAPK LLIY <i>RASTLAS</i> GVPSRFSGSGSGTDFTLTISSELPEDFATYYC <i>QS</i> <i>SYGNYG</i> DPGTGKTVTLG
SEQ ID NO: 26	Vλ germline-based FR4 (Sk17)	FGTGTKVTVLG
SEQ ID NO: 27	Vλ germline-based FR4 (Sk12)	FGGGTKLTVLG
SEQ ID NO: 28	Vλ germline-based FR4	FGGGTQLIILG
SEQ ID NO: 29	Vλ germline-based FR4	FGEGTELTVLG
SEQ ID NO: 30	Vλ germline-based FR4	FGSGTKVTVLG
SEQ ID NO: 31	Vλ germline-based FR4	FGGGTQLTVLG
SEQ ID NO: 32	Vλ germline-based FR4	FGGGTQLTALG
SEQ ID NO: 33	Vλ germline-based FR4_G141C	FGCGTKVTVLG
SEQ ID NO: 34	Linker	GGGGSGGGSGGGSGGGGS
SEQ ID NO: 35	Linker	GGGGS
SEQ ID NO: 36	Linker	GGGGSGGGGS

TABLE 4

Examples of multispecific antibodies as used in the present invention (Linkers are shown in bold).		
SEQ ID NUMBER	Ab Format	Sequence
PRO2198		
SEQ ID NO: 37	Morrison-H, LC	DIQMTQSPSSLSARVGDRTITCQASEDIESYLAW YQQKPGKAPKLLIYRASTLAYGVPSRFSGSGSGRD FTLTISSELPEDFATYYCLGSSYSTGSNIFGGGTK VEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTY SLSSLTTLTKADYEEKHKVYACEVTHQGLSSPVTKS FNRGEC
SEQ ID NO: 38	Morrison-H, HC	EVQLVESGGGLVQPGGSLRLSCAASGFSSSSAHYM CWVRQAPGKGLEWIGCIYAGSRGSTYYASWAKGRF TISRDNSKNTVYLMNSLRAEDTAVYYCARAAFN RGVSWIWPYYFSLWGQGLTVTVSSASTKGPSVFPL

TABLE 4-continued

Examples of multispecific antibodies as used in the present invention (Linkers are shown in bold).		
SEQ ID NUMBER	Ab Format	Sequence
		APCSRSTSESTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKY TCNVDHKPSNTKVDKRVESEKYGPCCPAPPEFLG GPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDP EVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKG QPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTTPPVLDSDGSFPLYSRLT VDKSRWQEGNVFSCSVMEALHNHYTQKSLSLGLG GGGGSGGGGS AFQLTQSPSSLSASVGDRVTITCQA QSIDSWLAWYQQKPKPKLLIYQASKLASGVPS RFGSGSGTDYTLTISSLQPEDFATYYCQTYGGG HIGWFGTGTQVTVL GGGGSGGGSGGGSGGGG SQ SQLVESGGGRVQPGGSLRLSCAASGFSFSANYW ICWVRQAPGKGLEWIVCIYIGSRADTYAPWAKGR FTISKDNSKNTVYLMNSLRAEDTATYFCARFSSG IYDLDRFFLWGQGTQVTVSS
PRO2199		
SEQ ID NO: 39	Morrison-H, LC	DIQMTQSPSSLSASVGDRVTITCQASEDIESYLAW YQQKPGKAPKLLIYRASTLAYGVPSRFGSGSGTD FTLTISSLQPEDFATYYCLGSSYSTGSIQGGTK VEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTY SLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC
SEQ ID NO: 40	Morrison-H, HC	EVQLVESGGGRVQPGGSLRLSCAASGFSSSSAHYM CWVRQAPGKGLEWIGCIYAGSRGTYASWAKGRF TISKASSTTVYLMNSLRAEDTATYYCARAAFPVNR GVSWIWPIYFSLWGQGTQVTVSSASTKGPSVFPLA PCSRSTSESTAALGCLVKDYFPEPVTVSWNSGALT SGVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKYT CNVDHKPSNTKVDKRVESEKYGPCCPAPPEFLGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPE VQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQ PREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTTPPVLDSDGSFPLYSRLT DKSRLWQEGNVFSCSVMEALHNHYTQKSLSLGLG GGGGSGGGGS AFQLTQSPSSLSASVGDRVTITCQAS QSIDSWLAWYQQKPKPKLLIYQASKLASGVPSR FSGSGSGTDYTLTISSLQPEDFATYYCQTYGGGH IGWFGTGTQVTVL GGGGSGGGSGGGSGGGG SQ SQLVESGGGRVQPGGSLRLSCAASGFSFSANYWI CWVRQAPGKGLEWIVCIYIGSRADTYAPWAKGRF TISKDNSKNTVYLMNSLRAEDTATYFCARFSSGI YDLDRFFLWGQGTQVTVSS
PRO2200		
SEQ ID NO: 41	Morrison-H, LC	AFQLTQSPSSLSASVGDRVTITCQASQSIDSWLAW YQQKPGKPKLLIYQASKLASGVPSRFGSGSGTD YTLTISSLQPEDFATYYCQTYGGGHIGWFGQGT KVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLN NFYPREAKVQWKVDNALQSGNSQESVTEQDSKDST YSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC
SEQ ID NO: 42	Morrison-H, HC	QSQVESGGGRVQPGGSLRLSCAASGFSFSANYWI CWVRQAPGKGLEWIVCIYIGSRADTYAPWAKGRF TISKDNSKNTVYLMNSLRAEDTATYFCARFSSGI YDLDRFFLWGQGTQVTVSSASTKGPSVFPLAPCSR STSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPFAVLQSSGLYSLSSVTVPSSSLGKTKYTCNVD HKPSNTKVDKRVESEKYGPCCPAPPEFLGGPSVF LFPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFN WYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKGLPSSIEKTIKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEW ESNQGPENNYKTTPPVLDSDGSFPLYSRITVDKSR

Examples of multispecific antibodies as used in the present invention (Linkers are shown in bold).			
SEQ ID NUMBER	Ab Format	Sequence	
		WQEGNVFSCSVMHEALHNHYTQKSLSLSLGGGGG GGGSDIQMTQSPSSLSARVGDRVITITCQASEDIES YLAWYQKPGKAPKLLIYRASTLAYGVPSRFGSGSG SGRDFTLTIISSLQPEDFATYYCLGSSYSTGSGNIFG TGTKVTVLGGGGGSGGGGSGGGGSGGGGSEVQLVE SGGGLVQPGGSLRLSCAASGFSSSSAHYMCWVRQA PGKGLEWIGCIYAGSRGSTYYASWAKGRFTISRDN SKNTVYLMQNSLRAEDTAVYYCARAAFVNRGVSWI WPYYFSLWGQGTLTVTSS	
PRO2201			
SEQ ID NO: 43	Morrison-H, LC	AFQLTQSPSSLSASVGDRVITITCQASQSIDSWLAW YQKPGKPPKLLIYQASKLASGVPSRFGSGSGSDT YTLTIISSLQPEDFATYYCQTYGGGHIWGFGQGT KVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLN NFYPREAKVQWKVDNALQSGNSQESVTEQDSKDS TSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGEC	
SEQ ID NO: 44	Morrison-H, HC	QSQLVESGGGRVQPGGSLRLSCAASGFSFSANYWI CWVRQAPGKGLEWIVICIYIGSRADTYAPWAKGRF TISKDNSKNTVYLMQNSLRAEDTATYFCARFSSGI YDLDRFFLWGQGTQVTVSSASTKGPSVFPLAPCSR STSESTAALGCLVKDYFPEPVTVSWNSGALTSGVH TFPAVLQSSGLYSLSVVTVPPSSSLGKTITYTCND HKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVF LFPKPKDITLMI SRTPEVTCVVVDVSQEDPEVQFN WYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQ DWLNGDKYKCKVSNKGLPSSIEKTIKAKGQPREP QVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEW ESNGQPENNYKTPPVLDSDGSFFLYSRLTVDKSR WQEGNVFSCSVMHEALHNHYTQKSLSLSLGGGGG GGGSDIQMTQSPSSLSASVGDRVITITCQASEDIE SYLAWYQKPGKAPKLLIYRASTLAYGVPSRFGSG SGSGDTFTLTIISSLQPEDFATYYCLGSSYSTGSDN FTGTGKVTVLGGGGGSGGGGSGGGGSGGGGSEVQLV ESGGGRVQPGGSLRLSCAASGFSSSSAHYMCWVRQ APGKLEWIGCIYAGSRGSTYYASWAKGRFTISK SSTTVYLMQNSLRAEDTATYCARAAFVNRGVSWI WPYYFSLWGQGTQVTVSS	
PRO2206			
SEQ ID NO: 45	Morrison-L, LC	DIQMTQSPSSLSARVGDRVITITCQASEDIESYLA WYQKPGKAPKLLIYRASTLAYGVPSRFGSGSGGRD FTLTIISSLQPEDFATYYCLGSSYSTGSGNIFGQGT VEIKRTVAAPSVFIFPPSDEQLKSGTASVVCCLN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDS TSLSTLTLSKADYEKHKVYACEVTHQGLSSPVTK SFNRGECGGGGGGGGGSAFQLTQSPSSLSASVGDRV TITCQASQSIDSWLAWYQKPGKPPKLLIYQASKL ASGVPSRFGSGSGSDTYTLTIISSLQPEDFATYYC QTYGGGHIWGFGGTGKVTVLGGGGGSGGGGSGGG GSGGGGSQSLVESGGGRVQPGGSLRLSCAASGFS FSANYWICWVRQAPGKLEWIVICIYIGSRADTYA PWAKGRFTISKDNSKNTVYLMQNSLRAEDTATYFC ARFSSGIYDLDRFFLWGQGTQVTVSS	
SEQ ID NO: 46	Morrison-L, HC	EVQLVESGGGLVQPGGSLRLSCAASGFSSSSAHYM CWVRQAPGKLEWIGCIYAGSRGSTYYASWAKGRF TISRDNKNTVYLMQNSLRAEDTAVYYCARAAFPVN RGVSWIWPYYFSLWGQGTLTVTSSASTKGPSVFPL APCSRSTSESTAALGCLVKDYFPEPVTVSWNSGAL TSGVHTFPAVLQSSGLYSLSVVTVPPSSSLGKT ITYTCNDHKPSNTKVDKRVESKYGPPCPPCPAPEFLG GPSVFLFPKPKDITLMI SRTPEVTCVVVDVSQEDP EVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVL TVLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKG QPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSD IAVEWESNGQPENNYKTPPVLDSDGSFFLYSRLT VDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSG	

TABLE 4-continued

Examples of multispecific antibodies as used in the present invention (Linkers are shown in bold).		
SEQ ID NUMBER	Ab Format	Sequence
PRO2207		
SEQ ID NO: 47	Morrison-L, LC	DIQMTQSPSSLSASVGDRVITTCQASEDIESYLAW YQOKPGKAPKLLIYRASTLAYGVPSRFGSGSGTD FTLTISSLQPEDFATYYCLGSSYSTGSNIFGQGTK VEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTY SLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC GGGGSGGGGSA FLTQSPSSLSASVGDRV TITCQASQSIDSWLAWYQOKPGKPPKLLIYQASKL ASGVPSRFGSGSGTDYTLTISSLQPEDFATYYCQ TYYGGGHIWGFGGTGKVTVL GGGGSGGGSGGG SGGGGGSQ SQLVESGGGRVQPGGSLRLSCAASGFS FSANYWICWVRQAPGKLEWIVCIYIGSRADTYA PWAKGRFTISKDNSKNTVYLQMNSLRAEDTATYFC ARFSSGIYDLDRFFLWGQGTQTVSS
SEQ ID NO: 48	Morrison-L, HC	EVQLVESGGGRVQPGGSLRLSCAASGFSSTSAHYM CWVRQAPGKLEWIGCIYAGSRGSTYYASWAKGRF TISKASSTTVYLQMNSLRAEDTATYYCARAAAFVNR GVSWIWPPYFSLWGQGTQTVSSASTKGPSVFPLA PCSRSTSESTAALGCLVKDYFPEPVTWNSGALT SGVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYT CNVDHKPSNTKVDKRVESKYGPPCPCPAPEFLGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPE VQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQ PREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTV DKSRWQEGNVFSCSMHEALHNHYTQKSLSLSLG
PRO2208		
SEQ ID NO: 49	Morrison-L, LC	DIQMTQSPSSLSASVGDRVITTCQASEDIESYLAW YQOKPGKAPKLLIYRASTLAYGVPSRFGSGSGTD FTLTISSLQPEDFATYYCLGSSYSTGSNIFGQGTK VEIKRTVAAPSVFIFPPSDEQLKSGTASVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTY SLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKS FNRGEC GGGGSGGGGSA FLTQSPSSLSASVGDRV TITCQASQSIDSWLAWYQOKPGKPPKLLIYQASKL ASGVPSRFGSGSGTDYTLTISSLQPEDFATYYCQ TYYGGGHIWGFGGTGKVTVL GGGGSGGGSGGG SGGGGGSQ SQLVESGGGRVQPGGSLRLSCAASGFS FSANYWICWVRQAPGKLEWIVCIYIGSRADTYA PWAKGRFTISKDNSKNTVYLQMNSLRAEDTATYFC ARFSSGIYDLDRFFLWGQGTQTVSS
SEQ ID NO: 50	Morrison-L, HC	EVQLVESGGGLVQPGGSLRLSCAASGFSSTSAHYM CWVRQAPGKLEWIGCIYAGSRGSTYYASWAKGRF TISKASSTTVYLQMNSLRAEDTAVYYCARAAAFVNR GVSWIWPPYFSLWGQGTQTVSSASTKGPSVFPLA PCSRSTSESTAALGCLVKDYFPEPVTWNSGALT SGVHTFPAVLQSSGLYSLSVVTVPSSSLGTQTYT CNVDHKPSNTKVDKRVESKYGPPCPCPAPEFLGG PSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPE VQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLT VLHQDWLNGKEYKCKVSNKGLPSSIEKTIKAKGQ PREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDI AVEWESNGQPENNYKTPPVLDSDGSFFLYSRLTV DKSRWQEGNVFSCSMHEALHNHYTQKSLSLSLG

[0372] Throughout the text of this application, should there be a discrepancy between the text of the specification (e. g., Tables 1 to 4) and the sequence listing, the text of the specification shall prevail.

[0373] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination

in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination. All combinations of the embodiments pertaining to the invention are specifically embraced by the present invention and are disclosed herein just as if each and every combination was individually and

explicitly disclosed. In addition, all sub-combinations of the various embodiments and elements thereof are also specifically embraced by the present invention and are disclosed herein just as if each and every such sub-combination was individually and explicitly disclosed herein.

[0374] The present invention is not to be limited in scope by the specific embodiments described herein. Indeed, various modifications of the invention in addition to those described herein will become apparent to those skilled in the art from the foregoing description. Such modifications are intended to fall within the scope of the appended claims.

[0375] To the extent possible under the respective patent law, all patents, applications, publications, test methods, literature, and other materials cited herein are hereby incorporated by reference.

[0376] The following Examples illustrates the invention described above, but is not, however, intended to limit the scope of the invention in any way. Other test models known as such to the person skilled in the pertinent art can also determine the beneficial effects of the claimed invention.

EXAMPLES

Example 1: Generation and Testing of Anti-IL-4R Molecules

Aim of the Project

[0377] The goal of the project was to generate humanized monoclonal antibody fragments that specifically bind to and neutralize the biological effect of human IL-4R. A further goal was to identify anti-IL-4R antibody fragments that are potent and stable enough for incorporation into multispecific antibody formats.

1.1. Generation, Selection, Humanization and Production of the Anti-IL-4R Binding Domains Included in the Multispecific Antibodies of the Present Invention

[0378] The generation of anti-IL-4R rabbit antibodies, the screening and selection of suitable clones, as well as the

humanization and production of the humanized anti-IL-4R binding domains were performed as described in the parallel patent application EP20216928.0, which is herewith incorporated by reference. Anti-IL-4R binding domains that are potent and stable enough for incorporation into multispecific antibody formats were identified and selected based on their biophysical and functional properties.

1.2. Characterization of Suitable Anti-IL-4R Binding Domains

[0379] For further characterization of suitable anti-IL-4R binding domains, the following analyses were performed.

1.2.1. Manufacture of Suitable Anti-IL-4R Binding Domains for Characterization (scFv Format)

[0380] Expression of mammalian constructs was performed in CHO-S cells using CHOgro transient transfection kit (Mirus). Cultures were harvested after 5-7 days (when cell viability <70% was reached) expression at 37° C. by centrifugation followed by filtration. Proteins were purified from clarified culture supernatants by Protein L affinity chromatography. None of the molecules needed polishing by size exclusion chromatography as all molecules exhibited at least one affinity chromatography fraction with a monomeric content >95% post capture as assessed by SE-HPLC analysis. Samples were re-buffered to final buffer (50 mM phosphate-citrate buffer with 150 mM NaCl at pH 6.4) by dialysis. For quality control of the manufactured material standard analytical methods such as SE-HPLC, UV₂₈₀ and SDS-PAGE were applied. The manufacture characteristics of eight scFv molecules originating from four clones that were deemed to be suitable for incorporation into multispecific antibody formats are summarized in Table 6. Manufacture characteristics of the CDR graft (sc01) and the Full graft (sc04) of clone 44-18-C11 (PRO1510) and its variants (PRO1549-PRO1554) are summarized in Table 7.

TABLE 6

scFv manufacture of 8 selected clones. Sc01: CDR-graft, sc04: Full graft									
Protein ID	Description	Expression volume [ml]	Yield post affinity chromatography purification [mg]	Final yield [mg]	Titer [mg/l]	Purity SE-HPLC [% monomer]	Purity SDS-PAGE	T _m [° C.]	T _{onset} [° C.]
PRO1506	44-03-A03-sc01	200	14.6	11.7	58	99.0	identity confirmed	62.8	56.7
PRO1524	44-03-A03-sc04	100	6.5	5.0	50	99.0	identity confirmed	59.6	54.0
PRO1510	44-18-C11-sc01	200	12.5	10.0	50	98.0	identity confirmed	71.5	64.0
PRO1528	44-18-C11-sc04	100	5.7	4.7	47	99.0	identity confirmed	59.1	52.3
PRO1517	44-34-C10-sc01	50	1.5	0.3	7	97.6	identity confirmed	63.7	57.0
PRO1535	44-34-C10-sc04	100	7.3	4.6	46	99.1	identity confirmed	67.9	62.7
PRO1520	44-39-B07-sc01	100	9.5	3.1	31	98.2	identity confirmed	70.2	64.0
PRO1538	44-39-B07-sc04	100	8.4	2.2	22	95.0	identity confirmed	68.3	63.7

TABLE 7

scFv manufacture of all grafting variants of clone 44-18-C11. Sc01: CDR-graft, sc04: Full graft									
Protein ID	Description	Expression volume [ml]	Yield post capture [mg]	Final yield [mg]	Titer [mg/l]	Purity SE-HPLC [% monomer]	Purity SDS-PAGE	T _m [° C.]	T _{onset} [° C.]
PRO1510	44-18-C11-sc01	200	12.5	10.0	50	98.0	identity confirmed	71.5	64.0
PRO1528	44-18-C11-sc04	100	5.7	4.7	47	99.0	identity confirmed	59.1	52.3
PRO1549	44-18-C11-sc05	100	8.1	6.8	68	99.0	identity confirmed	66.9	61.0
PRO1550	44-18-C11-sc06	100	8.0	9.0	90	99.0	identity confirmed	69.8	62.0

TABLE 7-continued

scFv manufacture of all grafting variants of clone 44-18-C11. Sc01: CDR-graft, sc04: Full graft									
Protein ID	Description	Expression volume [ml]	Yield post capture [mg]	Final yield [mg]	Titer [mg/l]	Purity SE-HPLC [% monomer]	Purity SDS-PAGE	T _m [° C.]	T _{onset} [° C.]
PRO1551	44-18-C11-sc07	100	8.1	6.8	68	98.0	identity confirmed	68.1	62.7
PRO1552	44-18-C11-sc08	100	6.8	5.5	55	98.0	identity confirmed	55.6	47.0
PRO1553	44-18-C11-sc09	100	7.5	6.7	67	99.0	identity confirmed	62.8	56.0
PRO1554	44-18-C11-sc10	100	8.8	6.9	69	99.0	identity confirmed	69.5	61.3

1.2.2. Pharmacodynamic Characterization of Suitable Anti-IL-4R Binding Domains (scFvs Format)

[0381] The scFvs antibodies that were evaluated as suitable were characterized for primary pharmacodynamics properties.

Affinity to Human IL-4R

[0382] Affinity of the 14 humanized scFvs (derived from 4 rabbit monoclonal antibodies) to human IL-4R was determined by SPR analysis on a T200 device (Biacore, GE Healthcare). Human IL-4R-Fc was captured via an anti-human IgG-Fc coupled to a carboxymethylated dextran surface and the scFvs were injected as analyte. After each analyte injection cycle the sensor chip was regenerated and new antigen was captured. The scFvs were measured using a dose response multi-cycle kinetic assay with seven analyte concentrations ranging from 0.12 to 90 nM diluted in running buffer. Obtained sensorgrams were fitted using a 1:1 binding model. The results are shown in Table 8. As can be seen from Table 8, the scFv antibody variable domains based on clone 44-34-C10 exhibit the best overall binding affinities to human IL-4R.

TABLE 8

Affinity of scFvs to human IL-4R.				
Protein ID	Clone ID	scFv Affinity to human IL-4R		
		k _a (M ⁻¹ s ⁻¹)	k _d (s ⁻¹)	K _D (M)
PRO1506	44-03-A03 -sc01	2.17E+06	1.49E-04	6.89E-11
PRO1524	44-03-A03-sc04	1.13E+06	3.40E-05	3.00E-11
PRO1510	44-18-C11 -sc01	1.67E+06	9.58E-04	5.75E-10
PRO1528	44-18-C11-sc04	3.83E+06	2.07E-04	5.39E-11
PRO1549	44-18-C11-sc05	1.63E+06	1.11E-03	6.77E-10
PRO1550	44-18-C11-sc06	1.55E+06	1.32E-03	8.52E-10
PRO1551	44-18-C11-sc07	1.84E+06	1.05E-03	5.70E-10
PRO1552	44-18-C11-sc08	8.41E+06	3.54E-04	4.21E-11
PRO1553	44-18-C11-sc09	4.24E+06	2.24E-04	5.27E-11
PRO1554	44-18-C11-sc10	2.03E+06	1.09E-03	5.37E-10
PRO1517	44-34-C10 -sc01	3.55E+06	1.95E-05	5.50E-12
PRO1535	44-34-C10-sc04	2.72E+06	5.42E-05	1.99E-11
PRO1520	44-39-B07 -sc01	6.18E+05	1.18E-02	1.91E-08
PRO1538	44-39-B07-sc04	1.67E+06	2.29E-04	1.37E-10

sc01: CDR graft; sc04: Full graft.

Affinity to Cynomolgus Monkey, Marmoset Monkey and Mouse IL-4R

[0383] Cross-reactivity to cynomolgus IL-4R was measured using the same setup as for the analysis of human IL-4R affinities, with the difference that cynomolgus IL-4R-Fc was captured instead of human IL-4R-Fc. To determine affinities for marmoset IL-4R, a direct setup was used since

only extracellular domain (ECD) of marmoset IL-4R was available. Marmoset ECD was immobilized directly onto the sensor chip and a titration series of scFvs was injected as analyte. To allow for comparison between KD values obtained for cynomolgus and marmoset IL-4R, affinities to cynomolgus IL-4R ECD were also determined using a direct setup.

[0384] Only few scFvs bound to cynomolgus IL-4R ECD. Three scFvs showed cross-reactivity in the capture setup compared to one scFv in the direct setup, meaning that the binding of two scFvs was no longer measurable in the direct setup. This is most likely a consequence of lower KD values determined in the direct setup. The difference was due to the on-rates, which were slower, possibly due to lower accessibility. Off-rates were comparable for all scFvs.

[0385] Furthermore, cross-reactivity to marmoset IL-4R ECD was determined. The results are summarized in Table 9.

[0386] Binding to mouse IL-4R was also assessed for scFvs PRO1517, PRO1535 and PRO1538 but none of them showed cross-reactivity.

Stat-6 Reporter Gene Assay in HEK-Blue Cells (Blockage of Human IL-4- and IL-13-Induced Signaling)

[0387] The HEK-Blue assay tested the ability of the humanized scFvs to inhibit IL-4- and IL-13-induced signaling via the IL-4R. 50,000 cells per well were seeded in a 96-well plate. Serial dilution of the scFvs and of the control antibody dupilumab were added to the plates in presence of either 0.02 ng/ml IL-4 or 0.3 ng/ml IL-13. After 24 h incubation at 37° C. and 5% CO₂, supernatants were transferred to a new plate and secreted embryonic alkaline phosphatase was quantified using the Quanti-Blue substrate.

[0388] The scFvs were analyzed in this HEK-Blue assay for their ability to block IL-4R-mediated signaling. The potency of the analyzed molecules was compared to dupilumab. All potency data are summarized in Table 10. Relative IC₅₀ values were calculated in mass units (ng/ml) of dupilumab and the scFvs. Eight scFvs blocked IL-4- and IL-13-induced signaling with high potencies. The scFvs based on the clone 44-34-C10 as well as PRO1552 showed best overall blockage potencies. PRO1517, PRO1524, PRO1535, PRO1538 and PRO1552 could block IL-4- and IL-13-induced signaling with a potency similar to dupilumab.

Competitive ELISA (Inhibition of hIL-4 Binding to hIL-4R)

[0389] Potency of humanized scFvs was further determined using a competitive ELISA. The potency of each scFv to inhibit the interaction between human IL-4 and human IL-4R was assessed by ELISA using the same procedure as described in EP20216928.0. Similarly, to the stat-6 reporter

gene assay, individual IC_{50} values on each plate were calibrated against the IC_{50} of the reference molecule dupilumab.

[0390] Seven scFvs inhibited the interaction between human IL-4 and human IL-4R with good to high potency. The scFvs based on clone 44-34-C10 as well as PRO1552 showed the best overall blockage potencies. The blocking potencies of PRO1517 and PRO1552 were similar to dupilumab. On the other hand, PRO1538 showed only moderate inhibition. Five scFv molecules showed no inhibition.

Summary of Pharmacological Characterization of scFvs and Molecule Selection for Developability Assessment

[0391] The pharmacological characterization of 14 scFvs derived from four different B cell clones provided the basis for selection of molecules for extended biophysical characterization. Five molecules, namely PRO1517, PRO1535, PRO1524, PRO1506 and PRO1538 were selected for further extended characterization based on their overall better performance in terms of neutralization of IL-4- and IL-13-induced signaling in the stat-6 reporter gene assay and in the IL-4/IL-4R-blocking ELISA. All seven molecules showed affinities better than 500 pM. The affinity to cynomolgus monkey IL-4R was lower by a factor of between 4- and 20-fold when compared to human IL-4R, except for three molecules, which showed very low affinity to cynomolgus IL-4R or no binding. In particular, the scFvs based on clone 44-18-C11, i. e. PRO1510, PRO1528 and PRO1549 to PRO1554, showed no binding to cynomolgus IL-4R. Generally, also the affinities of these scFvs to marmoset monkey IL-4R were lower when compared to human IL-4R. As a result of these affinity studies, the scFvs based on clone 44-18-C11, i. e. PRO1510, PRO1528 and PRO1549 to PRO1554 were excluded from further evaluation due to their lack of cross-reactivity to cynomolgus IL-4R, even though PRO1552 showed very good inhibition potencies. For the same reasons, PRO1520 was excluded as well. PRO1538 however, which is based on the same clone 44-39-B07 was not excluded, despite its weak binding affinity towards cynomolgus IL-4R.

1.2.3. Biophysical Characterization of Suitable Anti-IL-4R Binding Domains (scFvs Format)

Manufacture of Material for Stability Measurements

[0392] PRO1517 and PRO1538 were produced again using the same manufacturing process as described above at slightly larger scale (0.25 l expression volume) to generate sufficient material for stability assessment. For other proteins which were selected for stability assessment, the amount of previously produced material was sufficient to perform stability assessment. Samples were formulated in 50 mM phosphate-citrate buffer with 150 mM NaCl at pH 6.4 (50 mM NaCl, pH 6.4) after purification. After purification and dialysis, protein samples were concentrated to ≥ 10 mg/ml using centrifugal concentration tubes with 5 MWCO as this is the desired concentration to perform storage stability assessment.

[0393] Monomer loss upon concentration to 10 mg/ml was between 0 and 12% depending on the molecule as shown in Table 11. Monomeric content of PRO1506 and PRO1524 dropped below 95% which is a critical value for molecules entering storage stability assessment and, therefore, a reason to exclude them from the study. Nevertheless, as the material was already prepared and everything was set up for the study, stability of those molecules was assessed alongside the remaining molecules which exhibited a monomeric content $>95\%$ post concentration. The scFvs based on clone 44-34-C10, i. e. PRO1517 and PRO1535, as well as PRO1538 showed virtually no loss of the monomeric content upon concentration.

Storage Stability Study

[0394] Humanized scFvs were subjected to stability studies such as a four-week stability study, in which the scFvs were formulated in an aqueous buffer (50 mM NaCl, 150 mM NaCl, pH 6.4) at 10 mg/ml and stored at $<-80^{\circ}\text{C}$., 4°C . and 40°C . for four weeks. At the minimum, the fractions of monomers and oligomers in the formulation were evaluated by integration of SE-HPLC peak areas after one week, two weeks and at the end of each study. Additional time points were recorded for most of the molecules. Table 12 compares day 14 (d14) and endpoint measurements obtained at day 28 (d28) of the study.

TABLE 9

Affinity of the scFvs to cynomolgus and marmoset monkey IL-4R.										
Protein	scFv	Affinity to cynomolgus IL-4R capture setup			Affinity to cynomolgus IL-4R direct setup			Affinity to marmoset IL-4R direct setup		
		k_a ($\text{M}^{-1} \text{s}^{-1}$)	k_d (s^{-1})	K_D (M)	k_a ($\text{M}^{-1} \text{s}^{-1}$)	k_d (s^{-1})	K_D (M)	k_a ($\text{M}^{-1} \text{s}^{-1}$)	k_d (s^{-1})	K_D (M)
PRO1506	44-03-A03 -sc01	NB	NB	NB	NB	NB	NB	3.78E+06	7.45E-04	1.97E-10
PRO1524	44-03-A03-sc04	1.37E+09	8.07E+01	5.89E-08	NB	NB	NB	2.27E+06	2.06E-04	9.09E-11
PRO1510	44-18-C11 -sc01	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1528	44-18-C11-sc04	NB	NB	NB	NB	NB	NB	6.43E+06	4.39E-03	6.83E-10
PRO1549	44-18-C11-sc05	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1550	44-18-C11-sc06	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1551	44-18-C11-sc07	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1552	44-18-C11-sc08	NB	NB	NB	NB	NB	NB	1.57E+07	1.28E-03	8.16E-11
PRO1553	44-18-C11-sc09	NB	NB	NB	NB	NB	NB	6.64E+06	5.38E-03	8.09E-10
PRO1554	44-18-C11-sc10	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1517	44-34-C10 -sc01	1.92E+06	1.82E-04	9.46E-11	5.94E+05	4.04E-04	6.81E-10	2.67E+06	1.05E-03	3.92E-10
PRO1535	44-34-C10-sc04	1.60E+06	2.64E-04	1.65E-10	6.34E+05	2.38E-04	3.76E-10	3.23E+06	2.54E-03	7.87E-10

TABLE 9-continued

Affinity of the scFvs to cynomolgus and marmoset monkey IL-4R.										
Protein	scFv	Affinity to cynomolgus IL-4R capture setup			Affinity to cynomolgus IL-4R direct setup			Affinity to marmoset IL-4R direct setup		
ID	Clone ID	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	K_D (M)	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	K_D (M)	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	K_D (M)
PRO1520	44-39-B07 -sc01	NB	NB	NB	NA	NA	NA	NA	NA	NA
PRO1538	44-39-B07-sc04	5.29E+04	8.06E-04	1.52E-08	NB	NB	NB	1.02E+06	1.83E-04	1.80E-10

NA: not analyzed;
 NB: no binding
 sc01: CDR graft;
 sc04: Full graft;
 sc05-sc10: Other grafting variant

TABLE 10

Potencies of the scFvs to neutralize IL-4- and IL-13- induced signaling in the stat-6 reporter gene assay.					
Protein	scFv	Neutralization of hIL-4-induced signaling in reporter gene assay		Neutralization of hIL-13-induced signaling in reporter gene assay	
ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
PRO1506	44-03-A03-sc01	0.93	0.78	2.06	0.98
PRO1524	44-03-A03-sc04	0.95	0.80	1.81	1.05
PRO1510	44-18-C11-sc01	no inhibition	n/a	no inhibition	n/a
PRO1528	44-18-C11-sc04	1.49	0.28	2.29	0.59
PRO1549	44-18-C11-sc05	no inhibition	n/a	no inhibition	n/a
PRO1550	44-18-C11-sc06	no inhibition	n/a	no inhibition	n/a
PRO1551	44-18-C11-sc07	no inhibition	n/a	no inhibition	n/a
PRO1552	44-18-C11-sc08	0.43	1.30	1.02	1.41
PRO1553	44-18-C11-sc09	1.19	0.47	6.05	0.24
PRO1554	44-18-C11-sc10	no inhibition	n/a	no inhibition	n/a
PRO1517	44-34-C10-sc01	1.44	0.39	1.83	0.57
PRO1535	44-34-C10-sc04	1.15	1.02	2.30	0.99
PRO1520	44-39-B07-sc01	NA	NA	NA	NA
PRO1538	44-39-B07-sc04	0.64	1.83	1.69	1.35

sc01: CDR graft; sc04: Full graft; sc05-sc10: Other grafting variant.

NA: not analyzed

*IC₅₀, dupilumab/IC₅₀, test sample

n/a: relative IC₅₀ not calculated (no inhibition or IC₅₀ not determined)

TABLE 11

% monomer loss (SE-HPLC) upon concentration to 10 mg/ml.				
Protein ID	Clone ID (scFv)	Initial monomer content [%]	Monomer content post concentration to 10 mg/ml [%]	% loss upon concentration to 10 mg/ml
PRO1506	44-03-A03 -sc01	99.0	92.5	6.6
PRO1517	44-34-C10 -sc01	97.6	97.4	0.2
PRO1524	44-03-A03-sc04	99.0	87.3	11.8
PRO1535	44-34-C10-sc04	99.1	99.0	0.1
PRO1538	44-39-B07-sc04	95.0	98.2	-3.4

Sc01: CDR graft, sc04: Full graft

[0395] As can be depicted from Table 12, the scFvs based on clone 44-34-C10, i. e. PRO1517 and PRO1535, showed a good storage stability at 4° C. The loss of monomeric content after storage at 4° C. for d28 was less than 5%. Also PRO1538 showed a good storage stability at least for 14d. However, none of the stable molecules showed considerable loss of protein content at any temperature.

Freeze-Thaw Stability

[0396] In addition to the storage stability study described above, the compatibility of the top performing scFv mol-

ecules with respect to freeze-thawing (F/T) cycles (colloidal stability) was assessed. As in the early development stages the drug substance is usually stored frozen, formulation and fill/finish processing will require some freezing and thawing operations.

[0397] For the F/T stability assessment the same analytical methods (SE-HPLC, SDS-PAGE) and parameters (% monomer content and % monomer loss) as for the storage stability study were applied to monitor the quality of the molecules over five F/T cycles. Table 13 illustrates the course of %

monomer content loss over five F/T cycles. As no dedicated freeze-thaw study was performed, freeze-thaw data obtained with the -80°C . samples of storage stability study, which was acquired over 28 days, is shown. As only four time points could be recorded for some of the proteins, also only four F/T-cycles were done with those proteins.

Scanning Fluorimetry (DSF). Resulting midpoint of thermal unfolding (T_m) and onset temperature of unfolding calculated at 10% of maximal signal (T_{onset} 10%) values were determined by fitting of data to a Boltzmann equation. Table 14 summarizes calculated melting temperatures measured by DSF. As can be deduced from Table 14, the scFvs based

TABLE 12

28 d storage stability study at 10 mg/ml and temperatures of -80°C ., 4°C . and 40°C .												
Protein	Clone ID	Temp.	monomer content [%]			% monomer change		protein concentration [mg/ml]			% protein content change	
ID	(scFv)	[$^{\circ}\text{C}$.]	d 0	d 14	d 28	d 14	d 28	d 0	d 14	d 28	d 14	d 28
PRO1506	44-03-A03-sc01	-80	92.5	91.9	94.1	-0.6	1.7	10.1	9.8	NA	-3.0	NA
		4	92.5	92.3	92.2	-0.3	-0.3	10.1	9.4	NA	-6.7	NA
		40	92.5	91.7	92.1	-0.9	-0.5	10.1	9.7	10.3	-3.7	2.0
PRO1517	44-34-C10-sc01	-80	97.4	97.1	96.8	-0.4	-0.7	10.0	9.9	10.9	-1.5	8.4
		4	97.4	96.4	95.2	-1.1	-2.3	10.0	7.9	11.9	-21.7	18.8
		40	97.4	87.2	86.6	-10.5	-11.2	10.0	10.2	10.7	2.0	6.9
PRO1524	44-03-A03-sc04	-80	87.3	88.7	88.9	1.6	1.8	10.5	10.3	9.6	-1.9	-8.1
		4	87.3	89.2	88.2	2.2	1.0	10.5	10.2	9.6	-2.4	-8.8
		40	87.3	88.4	88.4	1.2	1.3	10.5	11.4	13.5	8.4	28.5
PRO1535	44-34-C10-sc04	-80	99.0	98.5	98.1	-0.5	-0.9	10.5	10.9	9.6	3.3	-9.1
		4	99.0	96.9	95.0	-2.1	-4.1	10.5	10.4	7.8	-1.4	-25.6
		40	99.0	90.3	88.6	-8.8	-10.5	10.5	10.8	10.8	2.6	3.1
PRO1538	44-39-B07-sc04	-80	98.2	97.8	NA	-0.3	NA	10.4	10.5	NA	0.6	NA
		4	98.2	96.0	NA	-2.2	NA	10.4	10.2	NA	-2.0	NA
		40	98.2	88.3	NA	-10.0	NA	10.4	12.0	NA	15.2	NA

NA: not analyzed

Sc01: CDR graft,

sc04: Full graft

TABLE 13

F/T stability - monomer content in % and % monomeric change upon repetitive freeze thawing.												
Protein	monomer content [%]					% monomer change						
ID	Clone ID (scFv)	F/T 0	F/T 1	F/T 2	F/T 3	F/T 4	F/T 5	F/T 1	F/T 2	F/T 3	F/T 4	F/T 5
PRO1506	44-03-A03-sc01	92.51	92.4	92.6	91.9	94.1	NA*	-0.1	0.1	-0.6	1.7	NA*
PRO1517	44-34-C10-sc01	97.44	97.7	97.7	97.1	96.9	96.8	0.2	0.3	-0.4	-0.6	-0.7
PRO1524	44-03-A03-sc04	87.31	85.5	85.6	88.7	88.9	NA*	-2.1	-2.0	1.6	1.8	NA*
PRO1535	44-34-C10-sc04	98.98	99.0	98.8	98.5	98.1	NA*	0.0	-0.2	-0.5	-0.9	NA*
PRO1538	44-39-B07-sc04	98.17	98.2	98.2	97.8	97.8	NA*	0.0	0.0	-0.3	-0.4	NA*

*not applicable as only 4 time points could be measured

Sc01: CDR graft,

sc04: Full graft

TABLE 14

DSF of selected domains.				
Protein ID	Clone ID (scFv)	Purity SE-HPLC [% monomer]	T_m [$^{\circ}\text{C}$.]	T_{onset} [$^{\circ}\text{C}$.]
PRO1506	44-03-A03-sc01	99.0	62.8	56.7
PRO1517	44-34-C10-sc01	97.6	63.7	57.0
PRO1524	44-03-A03-sc04	99.0	59.6	54.0
PRO1535	44-34-C10-sc04	99.1	67.9	62.7
PRO1538	44-39-B07-sc04	95.0	68.3	63.7

Sc01: CDR graft, sc04: Full graft

Thermal Unfolding

[0398] Thermal unfolding measurements of the five selected scFv constructs were performed using Differential

on the clone 44-34-C10 exhibit high melting temperatures of above 63°C . Also, PRO1538 showed a high melting temperature.

Example 2: Generation and Testing of Anti-IL31 Molecules

Aim of the Project

[0399] The goal of the project was to generate humanized monoclonal antibody fragments that specifically bind to and neutralize the biological effect of human IL-31. A further goal was to identify anti-IL-31 antibody fragments that are potent and stable enough for incorporation into multispecific antibody formats.

2.1. Generation, Selection, Humanization and Production of the Anti-IL-31 Binding Domains Included in the Multispecific Antibodies of the Present Invention

[0400] The generation of anti-IL-31 rabbit antibodies, the screening and selection of suitable clones, as well as the

humanization and production of the humanized anti-IL-31 binding domains were performed as described in the parallel patent application EP20216938.9, which is herewith incorporated by reference. Anti-IL-31 binding domains that are potent and stable

[0401] enough for incorporation into multispecific antibody formats were identified and selected based on their biophysical and functional properties. Clone 50-09-D07 was identified to have the overall best biophysical and functional properties.

[0402] For further characterization of suitable anti-IL-31 binding domains, the following analyses, were performed. 2.2.1. Manufacture of Suitable Anti-IL-31 Binding Domains for Characterization (scFv Format)

[0403] Expression of mammalian constructs was performed in CHO-S cells, as described above in section 1.2.1. Manufacture of two scFv sc01 and sc03 constructs (PRO1632 and PRO1643, respectively) originating from one clone, i. e. clone 50-09-D07, is summarized in Table 15. 2.2.2. Pharmacodynamics Characterization of Suitable Anti-IL-31 Binding Domains (scFv Format)

2.2. Characterization of Suitable Anti-IL-31 Binding Domains

[0404] The scFv antibodies that were evaluated as suitable, i. e. PRO1632 and PRO1643, were characterized for primary pharmacodynamics properties.

Affinity to Human and Cynomolgus IL-31

[0405] Affinity of the two humanized scFvs to human and cynomolgus IL-31 was determined by SPR analysis on a T200 device (Biacore, GE Healthcare). Cynomolgus IL-31 was manufactured on demand by Sino Biological since it was not available commercially. Human and cynomolgus IL-31-His was captured via an anti-His tag antibody coupled to a carboxymethylated dextran surface and the scFvs were injected as analyte. After each analyte injection cycle the sensor chip was regenerated and new antigen was captured. The scFvs were measured using a dose response multi-cycle

kinetic assay in a high-throughput mode with two concentrations (30 and 10 nM) diluted in running buffer. Obtained sensorgrams were fitted using a 1:1 binding model.

[0406] As shown in Table 16, binding to human IL-31 was confirmed for both humanized scFvs.

IL-31RA/OSMR Dimerization Assay (Blockage of Human IL-31-Induced Signaling)

[0407] The IL-31RA/OSMR dimerization assay was used to test the ability of the humanized scFvs to inhibit IL-31-induced signaling via the IL-31RA/OSMR heterodimer. 10,000 cells per well were seeded in a 96-well plate. Serial dilutions of the scFvs and of the control antibody BMS-981164 were added to the plates on the next day in presence of 10 ng/ml IL-31. After 6 h incubation at 37° C. and 5% CO₂, detection solution was added, the plates were incubated for another hour and luminescence was measured.

[0408] The two scFvs were tested in the cellular assay. As described above the potency of the analyzed molecules was compared to BMS-981164.

[0409] Relative IC₅₀ values were calculated in mass units (ng/ml) of BMS-981164 and the scFvs. The potency data are summarized in Table 17.

TABLE 17

Potencies of scFvs to neutralize IL-31R- and IL-31-induced signaling in IL-31RA/OSMR dimerization assay.			
scFv		Neutralization of hIL-31-induced signaling in IL-31RA/OSMR dimerization assay	
Protein ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
PRO1632	50-09-D07-sc01	53.29	0.20
PRO1643	50-09-D07-sc03	11.14	1.16

sc01: CDR graft; sc03: Full graft.
*IC₅₀ BMS-981164/IC₅₀ test sample

TABLE 15

scFv manufacture of 2 selected clones.								
Protein ID	Clone ID (scFv)	Expression volume [ml]	Yield post affinity chromatography purification [mg]	Yield/l post affinity chromatography purification [mg/l]	Final yield [mg]	Titer [mg/l]	Purity SE-HPLC [% monomer]	Purity SDS-PAGE
PRO1632	50-09-D07-sc01	30	5.8	193	0.7	23	96.4	identity confirmed**
PRO1643	50-09-D07-sc03	30	0.7	23	0.2	7	90.4	identity confirmed**

**protein of interest detected only
Sc01: CDR graft,
sc03: Full graft

TABLE 16

Affinity of scFvs to human IL-31.							
Protein ID	scFv Clone ID	Affinity to human IL-31			Affinity to cyno IL-31		
		k _a (M ⁻¹ s ⁻¹)	k _d (s ⁻¹)	K _D (M)	k _a (M ⁻¹ s ⁻¹)	k _d (s ⁻¹)	K _D (M)
PRO1632	50-09-D07-sc01	1.02E+06	1.26E-03	1.24E-09	1.52E+06	3.06E-03	2.01E-09
PRO1643*	50-09-D07-sc03	8.84E+05	4.47E-04	5.05E-10	7.61E+05	4.44E-04	5.83E-10
PRO1643**	50-09-D07-sc03	8.20E+05	5.16E-04	6.30E-10	7.64E+05	4.26E-04	5.57E-10

*The K_D of PRO1643 has been determined twice. The first K_D (KD 1) has been determined in this first SPR analysis via screening mode.
**The second K_D (KD 2) has been determined in this second SPR analysissusing full titration mode.
sc01: CDR graft;
sc03: Full graft.

Competitive ELISA (Inhibition of hIL-31 Binding to hIL-31RA)

[0410] Potency of the two humanized scFvs was further determined using a competitive ELISA. The potency of each scFv to inhibit the interaction between human IL-31 and human IL-31RA was assessed by an ELISA using the same procedure as described in EP20216938.9. Comparably to the IL-31RA/OSMR dimerization assay, individual IC₅₀ values on each plate were calibrated against the IC₅₀ of the reference molecule BMS-981164. The Full graft scFv inhibited the interaction between human IL-31 and human IL-31RA with similar potency when compared to BMS-981164. The potency data are summarized in Table 18.

TABLE 18

Potencies of scFvs to inhibit the interaction between IL-31 and IL-31R.			
scFv		Potency in IL-31/IL-31RA ELISA	
Protein ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
PRO1632	50-09-D07-sc01	22.91	0.41
PRO1643	50-09-D07-sc03	4.58	2.04

sc01: CDR graft; sc03: Full graft.
*IC₅₀, BMS-981164/IC₅₀, test sample

Summary of Pharmacological Characterization of scFvs and Molecule Selection for Detailed Biophysical Evaluation

[0411] The pharmacological characterization of the two scFvs provided the basis for selection of molecules for detailed biophysical evaluation. Based on these results, PRO1643 was selected for detailed biophysical evaluation based on its overall better performance.

2.2.3 Biophysical Characterization of Suitable Anti-IL-31 Binding Domains (scFv Format)

Manufacture of Stability Material

[0412] PRO1643 was produced again using the same manufacturing process as described above at slightly larger scale (0.25 l expression volume) to generate sufficient material for stability assessment. Samples were formulated in 50 mM phosphate-citrate buffer with 150 mM NaCl at pH 6.4 (50 mM NaCiP, pH 6.4). The protein was concentrated to >10 mg/ml using centrifugal concentration tubes with 5MWCO after purification and dialysis.

[0413] Monomer loss upon concentration to 10 mg/ml was 0.1% as shown in Table 19.

TABLE 19

% monomer loss (SE-HPLC) upon concentration to 10 mg/ml.				
Protein ID	Clone ID (scFv)	Initial monomer content [%]	Monomer content post concentration to 10 mg/ml [%]	% loss upon concentration to 10 mg/ml
PRO1643	50-09-D07-sc03	98.3	98.2	0.1

Sc03: Full graft

Storage Stability Study

[0414] PRO1643 was subjected to stability studies such as a four-week stability study, in which the scFv was formulated in an aqueous buffer (final buffer, 50 mM NaCiP, 150 mM NaCl, pH 6.4) at 10 mg/ml and stored at <-80° C., 4°

C. and 40° C. for four weeks. The fractions of monomers and oligomers in the formulation were evaluated by integration of SE-HPLC peak areas at different time points of the study. Further, protein concentration was determined by UV₂₈₀ measurement at different time points. Table 20 compares d14 and endpoint measurements obtained at d28 of the study.

[0415] PRO1643 showed no considerable loss of monomeric content at 4° C. and d28 of the study. PRO1643 showed no considerable loss of protein content at any temperature.

Freeze-Thaw Stability

[0416] In addition to the storage stability study described above, the compatibility of PRO1643 with respect to freeze-thawing (F/T) cycles (colloidal stability) was assessed.

[0417] For the F/T stability assessment the same analytical methods (SE-HPLC, UV-Vis) and parameters (% monomer content and % monomer loss) as for the storage stability study were applied to monitor the stability of the molecules over three F/T cycles. Table 21 illustrates the course of monomer content in % and % monomer content loss over three F/T cycles. As no dedicated freeze-thaw study was performed, freeze-thaw data obtained with the -80° C. samples of storage stability study which was acquired over 28 days is shown in the table below. As only three time points could be recorded per sample, freeze-thaw stability data is available for three F/T-cycles only.

Thermal Unfolding

[0418] Thermal unfolding measurement of PRO1643 has been performed using Differential Scanning Fluorimetry (DSF). Resulting midpoint of thermal unfolding (T_m) and onset temperature of unfolding calculated at 10% of maximal signal (T_{onset} 10%) values has been determined by fitting of data to a Boltzmann equation. Table 22 summarizes calculated melting temperatures measured by DSF.

Example 3: Selection and Optimization of Anti-IL-4R Molecules for Multispecific Format

3.1. General Remarks on the Selection of Anti-IL-4R Domains

[0419] The anti-IL-4R domains PRO1517 (44-34-C10-sc01) and PRO1535 (44-34-C10-sc04) were selected in the first place, since they exhibit very good pharmacodynamic properties and an acceptable to good stability. PRO1538

(44-39-B07-sc04) was selected as well, albeit having improvable pharmacological properties, since it showed the best stability. PRO1517 (44-34-C10-sc01) was subjected to a stability optimization and PRO1538 (44-39-B07-sc04) was subjected to pI balance optimization, as shown below in 3.2.

[0420] The anti-IL-4R binding domains applied in the final multispecific antibodies showed exclusive cross-reactivity to cynomolgus and marmoset monkey.

3.2. Optimization of Anti-IL-4R Domains

[0421] The anti-IL-4R domains PRO1538 (44-39-B07-sc03) and PRO1517 (44-34-C10-sc01) were further optimized for assembly into multispecific formats.

TABLE 20

28 d storage stability study at 10 mg/ml and temperatures of -80° C., 4° C. and 40° C.												
Protein		Temp.	monomer content [%]			% monomer change		protein concentration [mg/ml]			% protein content change	
ID	Clone ID (scFv)	[° C.]	d 0	d 14	d 28	d 14	d 28	d 0	d 14	d 28	d 14	d 28
PRO1643	50-09-D07-sc03	-80	98.2	98.3	99.2	0.1	1.0	10.6	12.4	12.1	23.2	20.4
		4	98.2	98.2	99.0	-0.1	0.7	10.6	11.9	10.3	18.4	2.6
		40	98.2	97.2	97.8	-1.0	-0.4	10.6	12.1	13.0	20.6	29.1

Sc03: Full graft

TABLE 21

F/T stability - monomer content in % and % monomeric change upon repetitive freeze thawing.									
Protein	Clone ID	monomer content [%]				% monomer change			
ID	(scFv)	F/T 0	F/T 1	F/T 2	F/T 3	F/T 1	F/T 2	F/T 3	
PRO1643	50-09-D07-sc03	98.2	98.2	98.3	99.2	-0.1	0.1	1.0	

Sc03: Full graft

TABLE 22

DSF measurement of selected scFv domains.				
Protein ID	Clone ID (scFv)	Purity SE-HPLC [% monomer]	T _m [° C.]	T _{onset} 10% [° C.]
PRO1643	50-09-D07-sc03	90.4*	69.5	64.3

Sc03: Full graft

*initial material with 90.4% monomer content used for measurement

Optimization of Anti-IL-4R Domain 44-39-B07-Sc04 (PRO1538)

[0422] PRO1538 (44-39-B07-sc03) has a pI imbalance between VL and VH (pI_{VL}=8.2 and pI_{VH}=5.0). Imbalance in pI correlates with high viscosity and could negatively influence stability. Therefore, two new variants of this molecule were designed. These variants are described below. Briefly, all residues in VH have been analyzed in detail with the aim to design molecules with removed pI imbalance and without compromising binding affinity and without introducing critical sequence liabilities (chemical and post-translational modifications), T cell epitopes etc.

[0423] Two variants were designed, which are summarized in Table 23.

[0424] These variants of PRO1538 (44-39-B07-sc03) did, however, not show the desired improvement of stability. Since PRO1538 is significantly less potent in inhibiting IL-4- and IL-13-induced signaling, and has a more than 10-fold lower potency in blocking the interaction of IL-4 to IL-4R than PRO1517, PRO1538 and its variants were excluded from further consideration and efforts were focused on the improvement of RPO1517 (44-34-C10-sc01).

TABLE 23

Optimization variants for 44-39-B07-sc03				
Protein ID	Construct	Optimization Aim	Mutations VL	Mutations VH
PRO2097	44-39-B07-sc05	pI balance	I2V, A51P, P75S, G82R, F89Y	V2E, A25R, R82S, Y105F, T143R
PRO2098	44-39-B07-sc06	pI balance	I2V, A51P, P75S, G82R, F89Y	V2E, A24R, R82S, Y105F, T143R

Optimization of Anti-IL-4R Domain 44-34-C10-Sc01 (PRO1517)

[0425] PRO1517 (44-34-C10-sc01) is an anti-IL-4R scFv with favorable binding characteristics. During biophysical characterization the scFv PRO1517 exhibited good but yet still improvable solubility. Efforts were made to further improve PRO1517 with respect to solubility, long-term stability and concentration behavior. Therefore, four new variants of this molecule were designed. These variants are

described below. Briefly, hydrophobic patches in CDRs or in the former VH-CH interface have been analyzed in detail and pI imbalance has been reduced with the aim to design molecules with removed hydrophobic patches without compromising binding affinity and without introducing critical sequence liabilities (chemical and post-translational modifications), T cell epitopes etc.

[0426] Four variants were designed, which are summarized in Table 24.

TABLE 24

Optimization variants for 44-34-C10-sc01				
Protein ID	Construct	Optimization Aim	Mutations VL	Mutations VH
PRO1896	44-34-C10-sc06	Solubility, stability		L12R, V103T, L144Q
PRO1897	44-34-C10-sc07	Solubility, stability	Y72S	
PRO1898	44-34-C10-sc08	Solubility, stability	S14R, T87R	
PRO1899	44-34-C10-sc09	Solubility, stability, affinity		L12R, V103T, L144Q + RDNSKN changed to KASST (Aho 82-87)

3.3 Biophysical Characterization of Optimized Anti-IL-4R Binding Domains (scFv Format)

Storage Stability Study

[0427] PRO1897, PRO1898 and PRO1899 were subjected to storage stability studies as described above in section 1.2.3. The results are summarized in Table 25.

[0428] As can be depicted from Table 25, the scFvs PRO1897, PRO1898 and PRO1899 exhibit excellent storage stability. The loss of monomeric content after storage at 4° C. for 28 days was less than 5%. Also, none of these molecules showed considerable loss of protein content at any temperature.

Thermal Unfolding

[0429] Thermal stability of PRO1897, PRO1898 and PRO1899 was analyzed by nano dynamic scanning fluorimetry (nDSF) using a NanoTemper, allowing the determination of the onset of unfolding (T_{onset}), midpoint of unfolding (T_{m1}) and scattering onset temperature. T_{m1} and T_{onset} results of duplicate/triplicate measurements are summarized in Table 26. As can be deduced from Table 26, PRO1897, PRO1898 and PRO1899 exhibit high melting temperatures.

TABLE 25

28 d storage and F/T stability study at 10 mg/ml and temperatures of -80° C., 4° C. and 40° C.												
Protein		Temp.	monomer content [%]			% monomer change		protein concentration [mg/ml]			% protein content change	
ID	Clone ID (scFv)	[° C.]	d 0	d 1	d 28	d 1	d 28	d 0	d 1	d 28	d 1	d 28
PRO1897	44-34-C10-sc07	-80	97.7	97.7*	NA	-0.1*	NA	10.2	10.5*	NA	2.2*	NA
		4	97.7	98	97	-0.1	-0.3	10.2	10	11	0.3	3.6
		40	97.7	95	87	-2.6	-10.8	10.2	11	13	2.6	30.0
PRO1898	44-34-C10-sc08	-80	97.4	97.3*	NA	-0.1*	NA	10.2	10.4*	NA	1.8*	NA
		4	97.4	97	96	-0.1	-1.1	10.2	10	10	-0.3	-1.7
		40	97.4	93	86	-4.4	-11.8	10.2	10	11	-1.7	9.7
PRO1899	44-34-C10-sc09	-80	98.1	98.0*	NA	0.0*	NA	10.5	10.7*	NA	1.9*	NA
		4	98.1	98	98	-0.1	0.0	10.5	10	11	-0.2	1.8
		40	98.1	97	84	-0.6	-14.3	10.5	11	13	2.9	28.8

NA: not analyzed;

*Five repeated F/T cycles were performed prior to measurement

TABLE 26

nDSF of optimized domains.				
Protein ID	Clone ID (scFv)	Purity SE-HPLC [% monomer]	T _{m1} [° C.]	T _{onset} [° C.]
PRO1897	44-34-C10-sc07	98.0	70.4	65.7
PRO1898	44-34-C10-sc08	98.0	76.0	67.0
PRO1899	44-34-C10-sc09	98.7	71.3	64.9

3.4. Pharmacodynamic Characterization of Optimized Anti-IL-4R Binding Domains (scFv Format)

Affinity to Human IL-4R

[0430] Affinity of the optimized anti-IL-4R scFvs PRO1898 and PRO1899 to human IL-4R was determined by SPR analysis on a T200 device (Biacore, GE Healthcare), as described above in section 1.2.2. The scFvs were measured using a dose response multi-cycle kinetic assay with seven analyte concentrations ranging from 0.12 to 90 nM diluted in running buffer. Obtained sensorgrams were fitted using a 1:1 binding model.

TABLE 28

Affinity of the optimized anti-IL-4R scFvs to cynomolgus IL-4R.				
Protein ID	Clone ID	Affinity to cynomolgus IL-4R capture setup		
		k _a (M ⁻¹ s ⁻¹)	k _d (s ⁻¹)	K _D (M)
PRO1898	44-34-C10-sc08	1.20E+06	1.09E-04	9.08E-11
PRO1899	44-34-C10-sc09	1.1E+07	1.4E-04	1.2E-11

Stat-6 Reporter Gene Assay in HEK-Blue Cells (Blockage of Human IL-4- and IL-13-Induced Signaling)

[0433] The optimized anti-IL-4R scFvs PRO1898 and PRO1899 were analyzed in this HEK-Blue assay for their ability to block IL-4R mediated signaling, as described above in section 1.2.2. The potency of the analyzed molecules was compared to dupilumab. All potency data are summarized in Table 29. Relative IC₅₀ values were calculated in mass units (ng/ml) of dupilumab and the scFvs. The optimized scFvs blocked IL-4- and IL-13-induced signaling with high potencies similar to dupilumab (see FIG. 3).

TABLE 29

Potencies of the scFvs PRO1898 and PRO1899 to neutralize IL-4 and IL-13 induced signaling in the stat-6 reporter gene assay.					
Protein ID	Clone ID	Neutralization of hIL-4 induced signaling in reporter gene assay		Neutralization of hIL-13 induced signaling in reporter gene assay	
		IC ₅₀ [ng/ml]	rel. IC ₅₀ *	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
Dupilumab	NA	1.15	1.00	1.00	1.00
PRO1898	44-34-C10-sc08	1.00	1.15	1.15	0.87
PRO1899	44-34-C10-sc09	0.71	1.61	0.56	1.80

*IC₅₀, dupilumab/IC₅₀, test sample
NA: not applicable

[0431] As shown in Table 27, binding to human IL-4R and cynomolgus IL-4R was confirmed for all optimized scFvs.

TABLE 27

Affinity of optimized anti-IL-4R scFvs to human IL-4R.				
Protein ID	Clone ID	Affinity to human IL-4R		
		k _a (M ⁻¹ s ⁻¹)	k _d (s ⁻¹)	K _D (M)
PRO1898	44-34-C10-sc08	2.15E+06	1.70E-05	7.91E-12
PRO1899	44-34-C10-sc09	2.95E+06	2.04E-05	6.92E-12

Affinity to Cynomolgus IL-4R

[0432] Affinity of the optimized anti-IL-4R scFvs PRO1898 and PRO1899 to cynomolgus IL-4R was measured using the same setup as for the analysis of human IL-4R affinities, with the difference that cynomolgus IL-4R-Fc was captured instead of human IL-4R-Fc, as described above in section 1.2.2. The results of this analysis are summarized in Table 28.

Competitive ELISA (Inhibition of hIL-4 Binding to hIL-4R)

[0434] Potency of the optimized anti-IL-4R scFvs PRO1898 and PRO1899 scFvs was determined using a competitive ELISA. The potency of each scFv to inhibit the interaction between human IL-4 and human IL-4R was assessed by ELISA as described in section 1.2.2. Similarly, to the stat-6 reporter gene assay, individual IC₅₀ values on each plate were calibrated against the IC₅₀ of the reference molecule dupilumab.

[0435] The results of the competitive ELISA analysis are summarized in Table 30. The blocking potencies of PRO1898 and PRO1899 were similar to dupilumab (see FIG. 4).

TABLE 30

Potency of the scFvs PRO1898 and PRO1899 to inhibit the hIL-4 - IL-4R interaction in a competition ELISA assay.			
Protein ID	Clone ID	Inhibition in hIL-4 - IL-4R competition ELISA	
		IC ₅₀ [ng/ml]	rel. IC ₅₀ *
Dupilumab	NA	6.78	1.00
PRO1898	44-34-C10-sc08	3.41	1.95

TABLE 30-continued

Potency of the scFvs PRO1898 and PRO1899 to inhibit the hIL-4 - IL-4R interaction in a competition ELISA assay.			
scFv		Inhibition in hIL-4 - IL-4R competition ELISA	
Protein ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
Dupilumab	NA	6.18	1.00
PRO1899	44-34-C10-sc09	2.98	2.07

*IC₅₀, dupilumab/IC₅₀, test sample;
NA: not applicable

Example 4: Selection and Optimization of Anti-IL31 Molecules for Multispecific Format

4.1. General Remarks on the Selection of Anti-IL-31 Domains

[0436] The anti-IL-31 domain PRO1643 (50-09-D07-sc03) was selected in the first place for further development, since it exhibits the desired pharmacodynamic properties as well as an excellent stability. Nevertheless, the anti-IL-31 binding domain PRO1643 (50-09-D07-sc03) underwent further solubility improving as shown below in 4.2.

[0437] The anti-IL-31 binding domains applied in the final multispecific antibodies do cross-react with cynomolgus monkey IL-31.

4.2. Optimization of Anti-IL-31 Domains

[0438] The anti-IL-31 domain PRO1643 (50-09-D07-sc03) was further optimized for assembly into multispecific format.

Optimization of Anti-IL-31 Domain 50-09-D07-Sc03

[0439] PRO1643 (50-09-D07-sc03) is a very stable anti-IL-31 scFv, however, efforts were nevertheless made to improve it with respect to solubility, long-term stability and concentration behavior. Therefore, new variants of this molecule were designed. These variants are described below. Briefly, hydrophobic patches in CDRs or in the former VH-CH interface have been analyzed in detail with the aim to design molecules with removed hydrophobic patches without compromising binding affinity and without introducing critical sequence liabilities (chemical and post-translational modifications), T cell epitopes etc.

[0440] Four variants were designed, which are summarized in Table 31.

TABLE 31

Optimization variants for 50-09-D07-sc03				
Protein ID	Construct	Optimization Aim	Mutations VL	Mutations VH
PRO1900	50-09-D07-sc04	Solubility, stability	—	L12R, V103T, L144Q
PRO1901	50-09-D07-sc05	Solubility, stability	—	L12S, V103T, L144Q
PRO1902	50-09-D07-sc06	Solubility, stability	—	Y59S
PRO1903	50-09-D07-sc07	Solubility, stability, sequence liability (DS)	D32E	—

4.3. Biophysical Characterization of Optimized Anti-IL-31 Binding Domains (scFv Format) Storage Stability Study

[0441] PRO1900, PRO1901, PRO1902 and PRO1903 were subjected to storage stability studies as described above in section 1.2.3. except that the last readout was taken at day 14 and F/T as well as that stability at -80° C. was not assessed. The results are summarized in Table 32.

[0442] As summarized in Table 32, scFvs PRO1900, PRO1901, PRO1902 and PRO1903 exhibited excellent storage stability. The loss of monomeric content after storage at 4° C. for 14 days was less than 0.2%, and <3% after storage at 40° C. for 14 days. Also, none of these molecules showed considerable loss of protein content at any temperature and data point.

Thermal Unfolding

[0443] Thermal stability of PRO1900, PRO1901, PRO1902 and PRO1903 was analyzed by nano dynamic scanning fluorimetry (nDSF) using a NanoTemper, allowing the determination of the onset of unfolding (T_{onset}), mid-point of unfolding (T_{m1}) and scattering onset temperature. T_{m1} and T_{onset} results of duplicate/triplicate measurements are summarized in Table 33. As can be deduced from Table 33, PRO1900, PRO1901, PRO1902 and PRO1903 exhibited high melting temperatures.

4.4. Pharmacodynamics Characterization of Optimized Anti-IL-31 Binding Domains (scFv Format)

Affinity to Human IL-31

[0444] Affinity of PRO1643 and of the optimized anti-IL-31 scFvs to human IL-31 was determined by SPR analysis on a T200 device (Biacore, GE Healthcare), as described above in section 2.2.2. The scFvs were measured using a dose response multi-cycle kinetic assay in a high-throughput mode with two concentrations (30 and 10 nM) diluted in running buffer. Obtained sensorgrams were fitted using a 1:1 binding model.

[0445] As shown in Table 34, binding to human IL-31 and cynomolgus IL-31 was confirmed for all optimized scFvs.

Path Hunter IL-31RA/OSMR Dimerization Assay (Blockage of Human IL-31-Induced Signaling)

[0446] PRO1643 and the optimized scFvs were tested for their ability to inhibit IL-31-induced signaling via the IL-31RA/OSMR heterodimer by using an IL-31RA/OSMR dimerization assay. The assay was performed as described above. The potency of the analyzed molecules was compared to BMS-981164.

[0447] Relative IC₅₀ values were calculated in mass units (ng/ml) of BMS-981164 and the scFvs. The potency data are

summarized in Table 35. As can be depicted from Table 35, PRO1643 as well as the optimized variants PRO1900, PRO1901 and PRO1903 could potentially neutralize IL-31-induced signaling. The mutations introduced in the variant PRO1902 to optimize solubility and stability obviously resulted in the disruption of the ability to inhibit IL-31-induced signaling.

TABLE 32

14 d storage stability study at 10 mg/ml at temperatures of 4° C. and 40° C.												
Protein		Temp. [° C.]	monomer content [%]			% monomer change		protein concentration [mg/ml]			% protein content change	
ID	Clone ID (scFv)		d 0	d 1	d 14	d 1	d 14	d 0	d 1	d 14	d 1	d 14
PRO1900	50-09-D07-sc04	4	99.8	NA	99.8	NA	0.0	9.8	NA	10.0	NA	2.6
		40	99.8	99.3	98.9	-0.5	-0.9	9.8	10.1	10.1	3.4	3.5
PRO1901	50-09-D07-sc05	4	97.0	NA	97.0	NA	-0.0	10.5	NA	10.9	NA	3.6
		40	97.0	96.9	96.3*	-0.1	-0.7*	10.5	10.1	10.9*	-3.7	4.0*
PRO1902	50-09-D07-sc06	4	98.5	NA	98.4	NA	0.0	9.3	NA	9.4	NA	0.9
		40	98.5	97.2	95.8	-1.3	-2.7	9.3	9.3	9.4	0.7	1.4
PRO1903	50-09-D07-sc07	4	98.7	NA	98.7	NA	-0.1	9.6	NA	10.1	NA	4.5
		40	98.7	98.2	97.7	-0.5	-1.1	9.6	9.8	9.9	1.2	2.3

*sample incubated for first 7 days at 25° C. instead of 40° C.
NA: not analyzed

TABLE 34

Affinity of scFvs to human and cynomolgus IL-31.							
Protein		Affinity to human IL-31			Affinity to cynomolgus IL-31		
ID	Clone ID	k_a (M ⁻¹ s ⁻¹)	k_d (s ⁻¹)	K_D (M)	k_a (M ⁻¹ s ⁻¹)	k_d (s ⁻¹)	K_D (M)
PRO1643	50-09-D07-sc03	5.79E+05	6.53E-04	1.13E-09	5.01E+05	5.87E-04	1.17E-09
PRO1900	50-09-D07-sc04	6.61E+05	5.99E-04	9.07E-10	5.54E+05	5.94E-04	1.07E-09
PRO1901	50-09-D07-sc05	5.91E+05	5.65E-04	9.56E-10	5.01E+05	5.69E-04	1.14E-09
PRO1902	50-09-D07-sc06	8.52E+05	2.46E-03	2.89E-09	7.31E+05	2.47E-03	3.38E-09
PRO1903	50-09-D07-sc07	5.24E+05	6.38E-04	1.22E-09	4.34E+05	6.05E-04	1.39E-09

Sc03: Full graft,
sc04 to sc07: other grafting variants

TABLE 33

nDSF of optimized anti-IL-31 binding domains (scFv format).				
Protein ID	Clone ID (scFv)	Purity SE-HPLC [% monomer]	T_{m1} [° C.]	T_{onset} [° C.]
PRO1900	50-09-D07-sc04	99.8	70.7	60.5
PRO1901	50-09-D07-sc05	97.7	69.0	52.3
PRO1902	50-09-D07-sc06	98.3	73.8	62.5
PRO1903	50-09-D07-sc07	98.9	71.1	55.6

TABLE 35

Potencies of scFvs to neutralize IL-31R- and IL-31-induced signaling in IL-31RA/OSMR dimerization assay.			
scFv		Neutralization of hIL-31-induced signaling in IL-31RA/OSMR dimerization assay	
Protein ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
PRO1643	50-09-D07-sc03	8.078	0.85
PRO1900	50-09-D07-sc04	8.209	0.73
PRO1901	50-09-D07-sc05	8.071	0.78

TABLE 35-continued

Potencies of scFvs to neutralize IL-31R- and IL-31-induced signaling in IL-31RA/OSMR dimerization assay.			
scFv		Neutralization of hIL-31-induced signaling in IL-31RA/OSMR dimerization assay	
Protein ID	Clone ID	IC ₅₀ [ng/ml]	rel. IC ₅₀ *
PRO1902	50-09-D07-sc06	371.3	0.03
PRO1903	50-09-D07-sc07	7.982	1.49

Sc03: Full-graft,
sc04 to sc07: other grafting variants
*IC₅₀ BMS-981164/IC₅₀ test sample

Example 5: Anti-IL-4R×IL-31 bispecific antibodies based on Morrison-H IgG1-Fc-silenced

5.1. Morrison-H Design

[0448] The Morrison format (also IgG-(scFv)₂ format) is a bivalent and bispecific IgG-based molecular format comprising an Fc region and two scFv domains that are either fused via linkers to the C-termini of the heavy chains, commonly referred to as Morrison-H format (see FIG. 2A),

or that are fused via linkers to the C-termini of the light chains, commonly referred to as Morrison-L format (see FIG. 2B).

[0449] In a first series of anti-IL-4R $\times$ IL-31 bispecific antibodies, a Morrison-H format was chosen, wherein the Fc region is derived from the IgG subclass IgG1 that has been silenced by the double mutation Leu234Ala and Leu235Ala, also generally referred to as the “LALA mutation”. Two highly stable A-capped scFvs domains were fused to the heavy chain C-terminus.

[0450] In total, 19 bispecific antibodies having the above-mentioned Morrison-H format were designed, bearing the anti-IL-4R domains on the Fab-arms except for one molecule. 13 different anti-IL-4R domains derived from 5 clones and three anti-IL-31 domains derived from two clones were combined. Nine of these bispecific Morrison-H antibodies exhibited good pharmacological properties. These Morrison-H antibodies are summarized in Table 36.

TABLE 36

Overview of Morrison-H molecule architecture				
PRO ID	Fab-arm domains	Constant	scFv-domains	Linkers
PRO1929	44-34-C10-sc01	kappa/IgG1 (silent)	50-09-D07-sc03	(G ₄ S) ₄
PRO1930	44-18-C11-sc08	kappa/IgG1 (silent)	50-09-D07-sc03	(G ₄ S) ₄
PRO1982	44-18-C11-sc08	kappa/IgG1 (silent)	50-09-D07-sc03	(G ₄ S) ₂
PRO2077	44-18-C11-sc14	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂
PRO2078	44-18-C11-sc18	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂
PRO1973	44-39-B07-sc04	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂
PRO2079	44-39-B07-sc05	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂
PRO2080	44-39-B07-sc06	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂
PRO1977	44-03-A03-sc04	kappa/IgG1 (silent)	50-09-D07-sc04	(G ₄ S) ₂

5.2. Morrison-H Molecule Production

[0451] Expression of multispecific constructs was performed in FreeStyle CHO-S cells using the transient CHO-gro expression system (Mirus). The genes of interest were optimized for mammalian expression, synthesized and cloned into a standard pcDNA3.1 vector. Expression cultures were cultivated in batch using shaking flasks for 6 to 7 days (cell viability <70%) at 37° C. Expression temperature was lowered to 32° C. after 1 day of expression of all molecules listed in Table 37 except for PRO1982, PRO2078 and PRO2080 which were kept at 37° C. over the whole expression period. The culture supernatants were separated from cells by centrifugation followed by a 0.22 μ m sterile filtration. The target proteins were captured from the clarified culture supernatants by Protein L or A affinity chromatography followed by a size-exclusion chromatography polishing step (in case no fractions with a suitable monomeric content as assessed by SE-HPLC were available post capture already). For the quality control of the manufactured material standard analytical methods, such as SE-HPLC, SDS-PAGE, and UV₂₈₀ were used.

[0452] Table 37 shows a production overview of these molecules. All molecules were finalized with a monomeric content >98% after a SEC polishing step.

TABLE 37

Production overview of Morrison-H molecules						
Protein ID	Format	Fab-arm domain	scFv-domain	Expression titer post capture [mg/l]	Final titer [mg/l]	Purity SE-HPLC [% monomer]
PRO1929	Morrison-H	44-34-C10-sc01	50-09-D07-sc03	33.2	19.4	99.2
PRO1930	Morrison-H	44-18-C11-sc08	50-09-D07-sc03	16.9	7.4	98.4
PRO1982	Morrison-H	44-18-C11-sc08	50-09-D07-sc03	22.9	14.2	98.2
PRO2077	Morrison-H	44-18-C11-sc14	50-09-D07-sc04	19.1	11.6	99.4
PRO2078	Morrison-H	44-18-C11-sc18	50-09-D07-sc04	18.6	9.7	98.9
PRO1973	Morrison-H	44-39-B07-sc04	50-09-D07-sc04	37.2	18*	98.1
PRO2079	Morrison-H	44-39-B07-sc05	50-09-D07-sc04	43.1	19.6	99.4
PRO2080	Morrison-H	44-39-B07-sc06	50-09-D07-sc04	34.4	16.3	99.2
PRO1977	Morrison-H	44-03-A03-sc04	50-09-D07-sc04	21.8	11.8*	98.5

*calculated/extrapolated value as not all material was used for polishing step

5.3. Biophysical and Pharmacological Characterization

[0453] Nine multispecific molecules, i. e. PRO1929, PRO1930, PRO1982 and its improved variants PRO2077 and PRO2078; PRO1973 and its improved variants PRO2079 and PRO2080; and PRO1977, were selected for further characterization.

[0454] The nine molecules were subjected to an extended biophysical characterization to determine suitability for process and formulation development. These molecules were characterized by a range of methods to assess melting and aggregation temperatures, solubility at concentrations over 150 g/l, loss of monomer content after 2 weeks at 4° and 25° C., increase in hydrodynamic radius and charge variants and clipping of the proteins. Solubility was also assessed by PEG screen. For PRO1929, loss of monomer content and the increase of hydrodynamic radius were determined.

[0455] Based on these results PRO2077, PRO2080, PRO1977, PRO1929 are the best constructs. They harbor the anti-IL-4R domains 44-18-C11-sc14, 44-39-09-sc06, 44-03-A04-sc03 and 44-34-C10-sc01, respectively and the anti-IL-31 domain 50-09-D07-sc04. The intended target concentration of 150 g/l was reached for all molecules. In terms of monomeric stability during storage at 4° C. and 25° C., PRO1929 and PRO2080 had the best profile. With respect to chemical stability and process development PRO2077 and PRO2080 showed a good pH stability and low charge variant complexity.

[0456] From a pharmacological perspective, PRO2077 shows a 2-fold superior potency to block IL-4- and IL-13-induced signaling when compared to dupilumab. The anti-IL-4R domain in this molecule does not cross-react with cynomolgus IL-4R. PRO2080, PRO1977 and PRO1929 block IL-4R-signaling approximately 2- to 3-fold less potently than dupilumab. PRO2080 and PRO1977 appear to show cross-reactivity to cynomolgus IL-4R based on avidity binding. A variant of PRO1929 could function as surrogate. PRO1929 exhibits a strong cross-reactivity to cynomolgus IL-4R and similar potency to inhibit IL-4R-signaling when compared to PRO2080 and PRO1977. All molecules potentially block IL-31-induced signaling and are cross-reactive to cynomolgus IL-31. Furthermore, all molecules exhibit picomolar affinities for both human targets.

Example 6: Anti-IL-4R×IL-31 Bispecific Antibodies Based on Morrison-H and Morrison-L IgG4

6.1. Morrison Format Design

[0457] In a second series of anti-IL-4R×IL-31 bispecific antibodies, Morrison-H and Morrison-L format were chosen, wherein the Fc region is derived from the IgG subclass IgG4. Two highly stable A-capped scFvs domains were fused to the C-termini of the heavy chain (Morrison-H) or the light chain (Morrison-L).

[0458] Based upon cynomolgus monkey cross-reactivity, excellent potency and excellent biophysical properties, constructs including the IL-4R and IL-31 domains of the PRO1929 family were prioritized.

[0459] Two variants of the PRO1929 family IL-4R binder were included in the reformatting of the molecules mentioned above, these were 44-34-C10-sc08 and 44-34-C10-sc09. The same IL-31 binding domain was included in all constructs and this was 50-09-D07-sc04.

[0460] In total, seven IgG4-(scFv)₂ molecules were designed, four Morrison-H and three Morrison-L molecules. Two different anti-IL-4R domains derived from one clone and one anti-IL-31 domain were combined, as shown in Table 38 (Morrison-H) and Table 39 (Morrison-L).

TABLE 38

IgG4 Morrison-H constructs			
PRO ID	Fab-arm domain	Constant	scFv-domain
PRO2198	44-34-C10-sc08	IgG4 (S228P)	50-09-D07-sc04
PRO2199	44-34-C10-sc09	IgG4 (S228P)	50-09-D07-sc04
PRO2200	50-09-D07-sc04	IgG4 (S228P)	44-34-C10-sc08
PRO2201	50-09-D07-sc04	IgG4 (S228P)	44-34-C10-sc09

TABLE 39

IgG4 Morrison-L constructs			
PRO ID	Fab-arm domain	Constant	scFv-domain
PRO2206	44-34-C10-sc08	IgG4 (S228P)	50-09-D07_sc04
PRO2207	44-34-C10-sc09	IgG4 (S228P)	50-09-D07_sc04
PRO2208	44-34-C10-sc09	IgG4 (S228P)	50-09-D07_sc04

6.2. Assessment of Binding Affinities and Specificities by SPR

Affinity to Human and Cyno IL-4R

[0461] Binding kinetics (including affinity) of the bispecific Morrison-H (PRO2198, PRO2199) and Morrison-L (PRO2206, PRO2207) molecules to recombinant human IL-4R protein (ECD with His Tag, Sino Biological) were determined by SPR analysis on a T200 device (Biacore, Cytiva). In this SPR experiment, Morrison molecules were captured via an anti-human IgG-Fc antibody (Human Antibody Capture Kit; Biacore, Cytiva) covalently immobilized to a carboxymethylated dextran surface (CM5 sensorchip; Biacore, Cytiva) and a titration series of human IL-4R ECD was injected as analyte. After each analyte injection cycle, the sensor chip was regenerated (MgCl₂), and new Morrison antibody was subsequently re-captured. The binding kinetics to human IL-4R was measured using a multi-cycle kinetic assay, with nine analyte concentrations ranging from 0.175 to 45 nM diluted in running buffer (HEPES buffered saline, 0.05% Tween-20, pH 7.5). The apparent dissociation (k_d) and association (k_a) rate constants and the apparent dissociation equilibrium constant (K_D) were calculated with the Biacore analysis software (Biacore Evaluation software Version 3.2, Cytiva) using one-to-one Langmuir binding model and quality of the fits was monitored based on Chi2 and U-value, which are measures for the quality of the curve fitting. The binding level was calculated as the maximum stability binding achieved normalized to the theoretical R_{max}. Cross-reactivity to cynomolgus monkey IL-4R was measured using the same setup as for the analysis of human IL-4R kinetics, with the difference that recombinant cynomolgus monkey IL-4R (ECD with His-tag, Acro Biosystems) protein was used as analyte instead of human IL-4R.

[0462] As shown in Table 40 high affinity binding to human IL-4R was demonstrated for the Morrison-H and Morrison-L antibodies. The calculated K_D values are very

similar ranging from 291 pM to 48 pM, with Morrison-H antibody PRO2198 having 291 ± 136 pM affinity for binding to human IL-4R. Kinetic affinities to cynomolgus monkey IL-4R are also similar among the tested molecules (209 to 11 pM), with 144 ± 49 pM affinity measured for PRO2198 Morrison-H antibody, thus demonstrating cross-reactivity of all the molecules to IL-4R of cynomolgus monkey origin.

Affinity to Human and Cyno IL-31

[0463] Binding kinetics (including affinity) of the bispecific Morrison-H (PRO2198, PRO2199) and Morrison-L (PRO2206, PRO2207) molecules to recombinant human IL-31 protein (Peprotech) were determined by SPR analysis on a T200 device (Biacore, Cytiva). In this SPR experiment, Morrison antibodies were injected over a carboxymethylated dextran surface (CM5 sensorchip; Biacore, Cytiva) immobilized with human recombinant IL-4R protein (ECD with Fc Tag, R&D Systems) and a titration series of human IL-31 was injected as analyte. The affinity to human IL-31 was measured using a single-cycle kinetic assay with injections of five sequential analyte concentrations, ranging from 0.05 to 30 nM diluted in running buffer (HEPES buffered saline, 0.05% Tween-20, pH 7.5), without regeneration of the sensorchip surface after injection of the analyte. The calculation of the kinetics and the apparent dissociation equilibrium constant (K_D) as well as the measurement for the quality of curve fitting were done as described above for IL-4R. The binding level was calculated as the maximum stability binding achieved normalized to the theoretical Rmax. Cross-reactivity to cynomolgus monkey IL-31 was measured using the same setup as for the analysis of human IL-31 affinities, with the difference that recombinant cynomolgus monkey IL-31 (ECD with His Tag, Sino Biological) protein was used as analytes instead of human IL-31.

[0464] A prerequisite of the SPR setup employed to measure binding kinetics to recombinant IL-31 is the capture of Morrison-H and Morrison-L antibodies via immobilized human IL-4R. All previously tested Morrison antibodies showed stable binding to immobilized human IL-4R, and thus the setup was proven valid. The rationale to use this SPR setup is to guarantee a homogeneous orientation of the captured Morrison molecules, while minimizing steric hindrance for the IL-31 binding site.

[0465] After having shown a stable capture of Morrison antibodies via recombinant human IL-4R protein, IL-31 was

injected as analyte and the kinetics of the binding of IL-31 to captured Morrison molecules were calculated. As shown in Table 41, all tested Morrison-H and Morrison-L molecule bind to human IL-31 and cynomolgus monkey IL-31 with similar affinities (152 to 52 pM and 169 to 103 pM, respectively). PRO2198 Morrison-H antibody demonstrated binding to human IL-31 with a calculated K_D of 52 ± 18 pM and binding to recombinant cynomolgus monkey IL-31 protein with a binding affinity of 10 ± 1 pM, therefore confirming good cross-reactivity.

Concomitant Binding to Human IL-4R and IL-31 and to Cyno IL-4R and IL-31

[0466] Concomitant binding of Morrison-H and Morrison-L antibodies was assessed by SPR analysis on a T200 device (Biacore, Cytiva). As described above, binding of Morrison antibodies to IL-31 (both human and cynomolgus monkey) is best measured when the Morrison antibody is captured on the sensorchip surface via IL-4R interaction, due to homogenous molecular orientation of the antibody as well as minimal steric hindrance of IL-31 binding sites. Therefore, concomitant binding of the tested Morrisons to both human IL-4R and human IL-31 was assessed by injecting the Morrison antibodies over a carboxymethylated dextran surface (CM5 sensorchip; Biacore, Cytiva) immobilized with human recombinant IL-4R protein (Fc Tag, R&D Systems), followed by injecting a single high concentration (45 nM) of human IL-31 as analyte. PRO2199, PRO2206 and PRO2207 concomitant binding was assessed during the kinetic measurement of human IL-31 as reported above, by using the binding levels from the injection of human IL-31 at 45 nM. Concomitant binding of PRO2198 to both cynomolgus monkey IL-4R and cynomolgus monkey IL-31 was measured similarly to human IL-4R and human IL-31, except for using recombinant cynomolgus monkey IL-4R (ECD with His-tag, Acro Biosystems) immobilized on the sensorchip surface and recombinant cynomolgus monkey IL-31 (ECD with His Tag, Sino Biological) protein as analyte.

[0467] Concomitant binding to human IL-4R and human IL-31 was demonstrated by SPR for all tested Morrison antibodies, i. e. PRO2198, PRO2199, PRO2206 and PRO2207, as exemplarily shown for PRO2198 in FIG. 5. For PRO2198, also concomitant binding to cynomolgus monkey IL-4R and cynomolgus monkey IL-31 was shown (see FIG. 5).

TABLE 40

Binding kinetics of Morrison antibodies to human IL-4R and cynomolgus monkey IL-4R						
Protein	Fab arm	scFv domain	Affinity by SPR measurement			
ID	Clone ID	Clone ID	Antigen	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	K_D (M)
PRO2198	44-34-C10-sc08	50-09-D07-sc04	human IL-4R	3.52E+05	1.01E-04	2.91E-10
PRO2199	44-34-C10-sc09	50-09-D07-sc04	human IL-4R	2.93E+05	5.76E-05	1.97E-10
PRO2206	44-34-C10-sc08	50-09-D07-sc04	human IL-4R	4.09E+05	1.95E-05	4.77E-11
PRO2207	44-34-C10-sc09	50-09-D07-sc04	human IL-4R	2.44E+05	4.61E-05	1.89E-10
PRO2198	44-34-C10-sc08	50-09-D07-sc04	cynomolgus IL-4R	3.70E+05	4.90E-05	1.44E-10
PRO2199	44-34-C10-sc09	50-09-D07-sc04	cynomolgus IL-4R	9.80E+05	3.44E-05	3.51E-11
PRO2206	44-34-C10-sc08	50-09-D07-sc04	cynomolgus IL-4R	1.16E+05	2.42E-05	2.09E-10
PRO2207	44-34-C10-sc09	50-09-D07-sc04	cynomolgus IL-4R	2.53E+06	2.77E-05	1.10E-11

TABLE 41

Binding kinetics of Morrison antibodies to human IL-31 and cynomolgus monkey IL-31						
Protein	Fab arm	scFv domain	Affinity by SPR measurement			
ID	Clone ID	Clone ID	Antigen	k_a ($M^{-1} s^{-1}$)	k_d (s^{-1})	K_D (M)
PRO2198	44-34-C10-sc08	50-09-D07-sc04	human IL-31	2.75E+05	1.60E-05	5.20E-11
PRO2199	44-34-C10-sc09	50-09-D07-sc04	human IL-31	2.51E+05	1.75E-05	6.97E-11
PRO2206	44-34-C10-sc08	50-09-D07-sc04	human IL-31	2.20E+05	2.36E-05	1.07E-10
PRO2207	44-34-C10-sc09	50-09-D07-sc04	human IL-31	1.63E+05	2.47E-05	1.52E-10
PRO2198	44-34-C10-sc08	50-09-D07-sc04	cynomolgus IL-31	9.92E+04	<1.00E-05	1.03E-10
PRO2199	44-34-C10-sc09	50-09-D07-sc04	cynomolgus IL-31	8.10E+04	<1.00E-05	1.23E-10
PRO2206	44-34-C10-sc08	50-09-D07-sc04	cynomolgus IL-31	5.93E+04	<1.00E-05	1.69E-10
PRO2207	44-34-C10-sc09	50-09-D07-sc04	cynomolgus IL-31	6.09E+04	<1.00E-05	1.64E-10

6.3. Binding to Human or Cynomolgus IL-4R-Expressing Cells (FACS)

[0468] To test binding of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 to human or cynomolgus IL-4R-expressing HEK293T cells, HEK293T cells were transfected with human and cynomolgus IL-4R plasmids and mock plasmid using Lipofectamine3000. Binding of an antibody targeting V5 Tag, which is fused to the N-terminus of human and cynomolgus IL-4R expression constructs, would confirm successful expression of both targets and was thus used as a positive control for plasma membranous binding. The Morrison-H molecule PRO2077 was used as negative control for cynomolgus IL-4R binding since this molecule had not shown any cross-reactivity to cynomolgus IL-4R (see section 5.3.).

[0469] 50,000 transfected HEK293T cells per well were seeded in a 96-well plate. 5-fold serial dilutions of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 and of the control molecule PRO2077, which is not cross-reactive to cynomolgus IL-4R, were added at concentrations ranging from 5,000 to 0.32 ng/ml to the plates and incubated for 60 min at 4° C. on a shaker. Supernatants were removed and detection reagent was added for flow cytometry analysis. The geometric mean of the APC fluorescence intensity of single cells was calculated.

[0470] All potency data are summarized in Table 42. Mean relative IC_{50} values from repeated analysis are indicated in pM. Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 efficiently bound to human and cynomolgus IL-4R on transfected HEK293T cells (see FIG. 6).

6.4. Potency Assessment to Inhibit Human IL-4 and IL-13 Signaling (HEK-Blue™ IL-4™ L-13 Cells Stat-6 Reporter Gene Assay)

[0471] The HEK-Blue assay has been used to test the ability of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 to inhibit IL-4- and IL-13-induced signaling via IL-4R. 50,000 cells per well were seeded in a 96-well plate. Three-fold serial dilutions of the Morrison molecules and of the reference antibody dupilumab at concentrations ranging from 100 to 0.002 ng/ml were added to the plates in presence of either 0.05 ng/ml IL-4 or 0.3 ng/ml IL-13. After 24 h incubation at 37° C. and 5% CO₂, supernatants were transferred to a new plate and secreted embryonic alkaline phosphatase was quantified using the Quanti-Blue substrate. The Morrison-H

and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 were also tested for inhibition of IL-4-induced signaling with an excess of IL-31 (10 nM) in the assay medium.

[0472] All potency data are summarized in Table 43 (IL-4/IL-4R inhibition) and Table 44 (IL-13/IL-4R inhibition). Mean relative IC_{50} values from repeated analyses are also indicated in pM. Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 efficiently blocked the interaction between IL-4 or IL-13 and IL-4R, with comparable potency to dupilumab (see FIG. 7). Efficient IL-4R blockage was maintained when IL-31 was bound to the anti-IL-31 domains of the Morrison-H molecules (see FIG. 8).

6.5. Potency Assessment to Inhibit Human IL-31-Induced Receptor Dimerization (PathHunter® eXpress IL31RA/OSMRb Assay)

PathHunter IL-31RA/OSMRb Dimerization Assay

[0473] In the IL-31RA/OSMR dimerization assay the ability of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 to inhibit IL-31-induced signaling via the IL-31RA/OSMR heterodimer was assessed. 10,000 cells per well were seeded in a 96-well plate. 3-fold serial dilutions of the molecules and of the control antibody BMS-981164 at concentrations ranging from 1,000 to 0.2 ng/ml were added to the plates on the next day in presence of 10 ng/ml IL-31. After 6 h incubation at 37° C. and 5% CO₂, detection solution was added, the plates were incubated for another hour and luminescence was measured. All antibodies were also tested for inhibition of IL-31-induced signaling with an excess of IL-4R (at 50 nM) in the assay medium.

Inhibition of Human IL-31

[0474] In Table 45, a summary of all potency data is shown. Mean relative IC_{50} values from repeated analysis are indicated in pM. All Morrison-H and Morrison-L antibodies inhibited IL-31-induced signaling with similar potency when compared to BMS-981164. Efficient blockage of the interaction of IL-31 with IL-31R was maintained when the anti-IL-4R domains were bound to IL-4R (see FIG. 9).

Inhibition of Cynomolgus IL-31

[0475] To assess inhibition of signaling induced by cynomolgus IL-31, the IL-31RA/OSMR dimerization assay was used as described above with the difference that 10 ng/ml cynomolgus IL-31 were used instead of 10 ng/ml human

IL-31, and only the Morrison-H antibodies PRO2198 and PRO2199 were tested for inhibition. Both PRO2198 and PRO2199 efficiently blocked the interaction between cynomolgus IL-31 and human IL-31RA (see FIG. 10).

6.6. Potency Assessment for Simultaneous Inhibition of Human IL-4 and IL-31 Signaling (BEAS-2B Assay; CCL2)

[0476] To assess potency of the Morrison-H antibodies PRO2198 and PRO2199 to inhibit both IL-4- and IL-31-induced signaling simultaneously, a cell-based assay using BEAS-2B (Human bronchial epithelial) cells with a Sandwich ELISA for readout was implemented. In this assay, blockage of both signaling pathways results in decreased CCL2 secretion by BEAS-2B cells.

[0477] 25,000 cells per well were seeded in 96 well plate in 100 μ l growth medium and incubated overnight at 37° C. with 5% CO₂. The following day, cells were washed and 1 ng/ml IL-4 and 10 ng/ml IL-31 were added with serial dilutions of a combination of dupilumab and BMS-981164 or of the Morrison-H antibodies PRO2198 or PRO2199. The cells were then incubated for 24 h, supernatants were collected, and the concentration of secreted CCL-2 was measured by means of a Sandwich ELISA adapted from a commercial kit.

[0478] The BEAS-2B assay is the only available assay that allows for the assessment of the combined inhibition of IL-4 and IL-31. Both PRO2198 and PRO2199 were able to simultaneously block IL-4 and IL-31 signaling with similar potency when compared to the combination of dupilumab and BMS-981164. They reached a level of inhibition comparable to the combination of the reference molecules (see FIG. 1).

6.7. Potency Assessment to Inhibit Human IL-4-Induced CD23 Upregulation on Human PBMC Subsets

[0479] Human PBMCs were used to assess IL-4-induced upregulation of CD23 in monocytes and naïve and memory B cells. Human PBMC were isolated from fresh whole blood according to an adapted Leucosep protocol, using Leucosep tubes with an integrated porous barrier and gradient density centrifugation.

[0480] 400,000 PBMCs per well were seeded in a 96-well plate in 100 ml assay medium. Three-fold serial dilutions of the Morrison-H antibodies PRO2198 and PRO2199, of the reference anti-IL-4R IgG4 antibodies PRO2209 and PRO2210, which have no anti-IL-31 scFv domains attached, and of the reference antibody dupilumab at concentrations ranging from 10,000 to 0.169 ng/ml were added to the plates in presence of 2 ng/ml IL-4. After 48 h incubation at 37° C. and 5% CO₂, PBMCs were stained for flow cytometry analysis using mouse anti-human CD19-PE, mouse anti-human CD14-PerCP/Cy5.5, mouse anti-human CD14-PerCP/Cy5.5 and mouse anti-human CD23-APC dyes. Following FACS analysis, the geometric mean of gated CD23 monocytes, CD23 naïve B cells and CD23 memory B cells was used for calculations. A four-parameter logistic (4PL) curve fit was applied to calculate IC₅₀ values of inhibition-curve using GraphPad Prism software.

[0481] PBMC assay data are summarized in Table 46. IC₅₀ values obtained for different blood donors are shown in picomolar. The Morrison-H antibodies PRO2198 and PRO2199 as well as the reference IgG4 antibodies PRO2209

and PRO2210 showed similar potency when compared to dupilumab to inhibit IL-4-induced upregulation of CD23 in monocytes and naïve and memory B cells. When formatted as IgG4 molecules without the anti-IL-31 domains attached, the molecules blocked IL-4-induced signaling with the same potency when compared to dupilumab (see FIG. 11). Thus, the addition of the anti-IL-31 domains has little effect on IL-4 inhibition potency.

6.8. Potency Assessment to Inhibit Human IL-4-Induced TARC Production in Human Whole Blood

[0482] To determine potency of inhibiting IL-4-induced secretion of thymus activation regulated chemokine (TARC, also CCL17), a whole blood assay based on a commercial ELISA as readout was used. Blockage of both signaling pathways leads to decreased TARC levels in whole blood.

[0483] Whole human peripheral blood collected in sterile tubes containing the anticoagulant EDTA was used for the assay. 160 μ l per well of serial dilutions of dupilumab, PRO2198 or PRO2199 ranging from 64,800 to 1.10 pM were added to a 96 well plate with round bottom wells with either 1 ng/ml IL-4 or 1 ng/ml IL-13. Following addition of 40 μ l whole blood per well, the plates were incubated for 24 h at 37° C. with 5% CO₂. On the next day, the plates were centrifuged for 10 min at 1,000 g, supernatants were collected and frozen at -20° C. The supernatants were then analyzed using the human TARC/CCL17 DuoSet ELISA Kit from R&D systems, and TARC levels were determined.

[0484] The TARC assay was performed with blood from three different donors for IL-4-induced TARC secretion. PRO2198 and PRO2199 neutralized IL-4-induced TARC in human whole blood with a potency comparable to dupilumab, i. e. with IC₅₀ values of less than 500 pM.

Example 7: Biophysical Characterization

[0485] The Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 were subjected to several biophysical characterization procedures. The following biophysical properties were investigated:

- [0486]** 1. Confirmation of identity
- [0487]** 2. Multi-pH stress study
 - [0488]** a. Chemical stability and charge heterogeneity
 - [0489]** b. Thermal stability
- [0490]** 3. High concentration feasibility (HCF) study
- [0491]** 7.1. Confirmation of identity:

Mass Determination

[0492] Masses were analyzed by intact mass analysis using ESI-MS. All tested Morrison-H and Morrison-L antibodies, i. e. PRO2198, PRO2199, PRO2206 and PRO2207, showed the expected masses for the light chain and the heavy chain. In addition, the heavy chain shows the expected glycosylation. The determined mass for each Morrison antibody is shown in Table 47.

TABLE 42

Plasma membranous binding to human and cynomolgus IL-4R on transfected HEK293T-cells.							
PRO ID	Fab-arm domain	Constant	scFv-domain	Human IL-4R		Cynomolgus IL-4R	
				EC ₅₀ [nM]	rel. IC ₅₀ rel. EC ₅₀ (EC ₅₀ V5 antibody/ EC ₅₀ sample)	EC ₅₀ [nM]	rel. IC ₅₀ rel. EC ₅₀ (EC ₅₀ V5 antibody/ EC ₅₀ sample)
PRO2198	44-34-C10-sc08	IgG4 (S228P)	50-09-D07-sc04	1.11	3.31	0.97	21.56
PRO2199	44-34-C10-sc09	IgG4 (S228P)	50-09-D07-sc04	0.85	4.33	1.78	11.73
PRO2206	44-34-C10-sc08	IgG4 (S228P)	50-09-D07-sc04	0.83	4.46	0.90	23.18
PRO2207	44-34-C10-sc09	IgG4 (S228P)	50-09-D07-sc04	0.60	6.15	0.65	32.22
PRO2077	44-18-C11-sc14	kappa/IgG1 (silent)	50-09-D07-sc04	0.60	6.18	no binding	no binding

TABLE 43

Potencies of the Morrison-H and Morrison-L antibodies to neutralize human IL-4-induced signaling in the stat-6 reporter gene assay without and with IL-31 in the assay medium.					
PRO ID	Fab-arm domain	scFv-domain	Mean IC ₅₀ [pM]	Mean rel. IC ₅₀ [IC ₅₀ dupilumab/ IC ₅₀ sample]	Retained IC ₅₀ with and without IL-31 excess (n = 1) [IC ₅₀ , no IL-31/IC ₅₀ , with IL-31] ng/ml values
PRO2198	44-34-C10-sc08	50-09-D07-sc04	23.9**	0.42	0.71
PRO2199	44-34-C10-sc09	50-09-D07-sc04	11.5**	0.72	1.25
PRO2206*	44-34-C10-sc08	50-09-D07-sc04	11.8	0.45	not measured
PRO2207*	44-34-C10-sc09	50-09-D07-sc04	9.4	0.52	not measured

N = number of repeated analyses.

*IC₅₀ or relative IC₅₀ (single analysis)/NA: not applicable

**average value

TABLE 44

Potencies of the Morrison-H and Morrison-L molecules to neutralize human IL-13-induced signaling in the stat-6 reporter gene assay.				
PRO ID	Fab-arm domain	scFv-domain	Mean IC ₅₀ [pM]	Mean rel. IC ₅₀ [IC ₅₀ dupilumab/ IC ₅₀ sample]
PRO2198	44-34-C10-sc08	50-09-D07-sc04	22.1**	0.42
PRO2199*	44-34-C10-sc09	50-09-D07-sc04	11.3	0.69
PRO2206*	44-34-C10-sc08	50-09-D07-sc04	29.5	0.26
PRO2207*	44-34-C10-sc09	50-09-D07-sc04	12.6	0.62

N = number of repeated analyses.

*IC₅₀ or relative IC₅₀ (single analysis)/NA: not applicable

**average value.

TABLE 45

Potencies of the Morrison-H and Morrison-L molecules to neutralize human IL-31-induced signaling in IL-31RA/OSMR dimerization assay without and with IL-4R in the medium.					
PRO ID	Fab-arm domain	scFv-domain	Mean IC ₅₀ [pM]	Mean rel. IC ₅₀ [IC ₅₀ BMS-981164/ IC ₅₀ sample]	Retained IC ₅₀ with and without IL-4R excess (n = 1) [IC ₅₀ , no IL-4R/IC ₅₀ , with IL-4R] ng/ml values
PRO2198	44-34-C10-sc08	50-09-D07-sc04	198	0.70	1.02
PRO2199	44-34-C10-sc09	50-09-D07-sc04	142	0.61	0.90
PRO2206*	44-34-C10-sc08	50-09-D07-sc04	230	0.65	not measured
PRO2207*	44-34-C10-sc09	50-09-D07-sc04	215	0.55	not measured

N = number of repeated analyses.

*IC₅₀ or relative IC₅₀ (single analysis)/NA: not applicable

TABLE 46

Ability of the Morrison antibodies to neutralize IL-4-induced CD23 expression on human PBMCs										
PRO ID	Fab-arm domain	Constant	scFv-domain	blood donor	monocytes		naive B cells		memory B cells	
					IC ₅₀ [pM]	[IC ₅₀ dupilumab/ IC ₅₀ sample]	IC ₅₀ [pM]	[IC ₅₀ dupilumab/ IC ₅₀ sample]	IC ₅₀ [pM]	[IC ₅₀ Dupilumab/ IC ₅₀ sample]
PRO2198	44-34-C10-sc08	IgG4 (S228P)	50-09-D07-sc04	1	90.37	0.11	1366	0.31	4470	0.14
PRO2199	44-34-C10-sc09	IgG4 (S228P)	50-09-D07-sc04	1	69.74	0.15	1099	0.38	2531	0.25
PRO2206	44-34-C10-sc08	IgG4 (S228P)	50-09-D07-sc04	2	110.5	0.12	1480	0.45	4339	0.13
PRO2207	44-34-C10-sc09	IgG4 (S228P)	50-09-D07-sc04	3	*54.47	0.02	1036	0.12	1580	0.20

*approximate values, top constraint applied when fitting curves

TABLE 47

Masses for light and heavy chain of each Morrison antibody found under reducing condition				
Masses in [Da]	PRO2198	PRO2199	PRO2206	PRO2207
Calculated mass	23582	23458	51028	50904
Light chain (LC)				
Detected mass	23580	23453	51019	50895
Calculated mass	77368.8	77169	49923	49743
Heavy chain (HC)				
Detected mass	78798 (+1429)	78618 (+1429)	51358 (+1435)	51179 (+1436)
Main species				
Comments	HC: Not fully reduced and glycosylated	LC: Not fully reduced HC: Not fully reduced and glycosylated	LC: Not fully reduced HC: Not fully reduced and glycosylated	LC: Not fully reduced HC: Not fully reduced and glycosylated

Glycosylation Pattern

[0493] PRO2198 and PRO2199 were analyzed in more details for their glycosylation pattern. Both molecules show a similar glycosylation. An approximation of the relative glycosylation of the respective heavy chain is shown in Table 48. Both molecules are glycosylated mainly with G0F sugar. Lower amounts of G1F and G2F are also found. No non-glycosylated mass for the heavy chain was observed.

TABLE 48

Type and relative abundance of heavy chain glycosylation		
Glycosylation	PRO2198	PRO2199
G0F (+1445 Da)	65%	65%
G1F (+1608 Da)	30%	30%
G2F (+1770 Da)	5%	5%

SDS Page

[0494] The identity of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207 was additionally analyzed by SDS-PAGE under reducing and non-reducing condition (see FIG. 12).

[0495] For the Morrison-H molecules, the expected bands were observed both under reducing as well as non-reducing conditions. Some HC dimer is observed under reducing conditions, presumably due to incomplete reduction of the interchain disulfide.

[0496] In case of the Morrison-L molecules, all expected bands are visible under both reducing and non-reducing conditions and represent the most prominent species. An additional band between 37 kDa and 50 kDa is observed under reducing conditions suggesting the LC or HC not being fully denatured or reduced under the chosen condition. The band running between 75 kDa and 100 kDa presumably originates from a LC dimer. Under non-reducing conditions, LC and LC dimers are visible. For IgG molecules it is known that the LC and LC dimers are soluble and can be present using a generic purification process.

7.2. Multi-pH Stress Study:

[0497] Aim of the multi-pH stress study was to examine the chemical and thermal stability of the Morrison-H and Morrison-L antibodies PRO2198, PRO2199, PRO2206 and PRO2207. The molecules were dialyzed into a 20 mM MES/Acetate/Hepes buffer with pH 3.5, 5, 7, and 8.5. In the latter three conditions, samples were incubated for 10 days at 40° C. to force chemical degradation such as oxidation and clipping. pH 3.5 is a harsh stress condition leading to rapid degradation and hydrolysis, therefore these samples were incubated at 25° C. for 10 days. The formation of charge variants was analyzed by CEX-HPLC and clipping was investigated by SDS-PAGE. The molecules were incubated at a concentration of 1 mg/ml. Thermal stability was analyzed by nDSF (NanoTemper) determining the onset of unfolding (T_{onset}), midpoint of unfolding (T_{m1}) and onset of aggregation ($T_{Scattering}$).

Charge Variants Profile

[0498] During later DSP development, purification in high yields is an important measure. One of the main techniques used to purify IgG like molecules is ion exchange chromatography. Therefore, IgG antibodies showing a high target purity by CEX-HPLC are likely to be less challenging during the development of an ion exchange purification step.

[0499] All tested antibodies, i. e. PRO2198, PRO2199, PRO2206 and PRO2207, exhibited a comparable complexity (charge heterogeneity), whereby the Morrison-L molecules showed a slightly smaller complexity than the Morrison-H molecules. In addition, the molecules PRO2199 and PRO2207 that are based on the IL-4R clone sc09 had a slightly higher target purity.

[0500] In summary, all molecules showed a low charge heterogeneity after a generic purification process without substantial differences.

Formation of Charge Variants

[0501] The formation of charge variants was analyzed for PRO2198, PRO2199, PRO2206 and PRO2207 after incubation in 20 mM MES/Acetate/Hepes buffer at pH 3.5, pH 5, pH 7 and pH 8.5. A specific importance is given to the chemical stability at pH 5 and pH 7 as these pH values define approximate limits for the later formulation space. Table 49 shows the results obtained. At pH 5, all molecules showed a good stability towards chemical degradation. For higher pH values, chemical degradation towards the formation of acidic variants increased. All molecules showed similar stabilities except PRO2206, which had a higher target loss at pH 8.5. At pH 3.5, only minor amounts of chemical degradation were observed.

TABLE 49

Formation of charge variants at different pH				
Values in [rel. %]	PRO2198	PRO2199	PRO2206	PRO2207
t0	76.4	81.4	78.1	84.1
Δ Target after 10 days of storage at 40° C. ^{#1}				
pH 3.5	3.1	1.9	1.0	1.3
pH 5.0	1.1	0.2	-1.0	-0.7
pH 7.0	-5.5	-7.2	-6.8	-7.2
pH 8.5	-12.9	-11.8	-22.0	-15.5

^{#1}Storage temperature at pH 3.5 is 25° C.)

Analysis of Clipping by SDS-PAGE

[0502] All proteins were analyzed for clipping using SDS-PAGE analysis (see FIG. 13). No fragmentation was observed for any of the Morrison antibodies at pH 3.5, pH 5 and pH 7. Only little fragmentation of the HC of PRO2198 and PRO2199 was observed at pH 8.5. Although no HC clipping was observed for the Morrison-L molecules, it could be overlaid by the closely running LC.

[0503] In summary, all molecules showed a good resistance towards clipping.

Thermal Stability

[0504] Within the pH stress study, PRO2198, PRO2199, PRO2206 and PRO2207 were analyzed for their thermal stability between pH 3.5 and pH 8.5. Both thermal unfolding

as well as the onset of thermal aggregation were determined. During thermal unfolding, all molecules showed more than one transition due to the multi domain architecture of the molecules. For simplicity, only the first melting midpoint is shown. Table 50 summarizes the results.

TABLE 50

Thermal stability				
pH	PRO2198	PRO2199	PRO2206	PRO2207
Onset of unfolding (T_{onset}) by nDSF [° C.]				
3.5	45.6	43.4	42.4	40.9
5	55.4	55.1	55.8	56.3
7	60.5	59.3	58.7	57.5
8.5	59.7	58.3	57.0	55.1
Midpoint of the first unfolding transition (T_{m1}) by nDSF [° C.]				
3.5	50.7	50.6	49.7	50.0
5	62.9	62.8	62.1	62.0
7	66.9	66.7	66.2	66.1
8.5	66.5	66.2	65.1	65.1
Onset of aggregation ($T_{scattering}$) by nDSF [° C.]				
3.5	No aggregation observed			
5	No aggregation observed			
7	73.5	71.6	72.2	69.0
8.5	71.4	68.4	68.8	65.2

[0505] The onset of unfolding (T_{onset}) as well as the first melting point (T_{m1}) was highest at pH 7 for all molecules. A change to pH 8.5 decreased the thermal stability only slightly. Towards acidic pH, the thermal stability decreased. However, at pH 5 all onsets of unfolding were above 55° C. At pH 3.5, the thermal stability decreased to a T_{onset} below 50° C.

[0506] Thermal aggregation was only observed at pH 7 and pH 8.5. At acidic pH, both pH 3.5 and pH 5.0, all Morrison antibodies did not form larger aggregates, even in the unfolded state. At neutral and basic pH, the aggregation onset was above the first melting point, suggesting a non-native aggregation mechanism.

[0507] Although the differences between the molecules were not substantially, PRO2198 has overall the highest stability. This was most prominent at pH 3.5 with a 5° C. higher T_{onset} compared to PRO2207. This pH is relevant in the production process for a viral inactivation step.

7.3. High Concentration Study

[0508] For the concentration feasibility tests PRO2198 was formulated in two formulation buffers:

[0509] Formulation buffer F1: 20 mM acetate at pH 5.5;

[0510] Formulation buffer F2: 20 mM citrate with 50 mM NaCl at pH 5.5.

Solubility—Protein Concentration

[0511] PRO2198 could be concentrated above 100 mg/ml, without signs of precipitation or reaching the limit of solubility. Furthermore, PRO2198 showed no detectable decrease in protein concentration, i. e. PRO2198 was sufficiently soluble to reach and keep a target concentration of above 100 mg/ml for at least four weeks at 4° C. and 25° C.

Monomer Stability at Different Temperatures

[0512] PRO2198 was formulated in F1 and F2 and concentrated to >100 mg/ml. The concentrated samples were stored at 4° C., 25° C. and 40° C. for up to 4 weeks. At different time points, the monomeric content was analyzed by means of SE-HPLC. The results are summarized in Table 51.

TABLE 51

Monomer stability for PRO2198 at a concentration of >100 mg/ml					
Storage temperature [° C.]	time [weeks]	Formulation F1 Monomer [rel. %]		Formulation F2 Monomer [rel. %]	
		Absolut	Delta	Absolut	Delta
N/A	0	99.0	0.0	99.1	0.0
4.0	4	98.1	-0.9	98.3	-0.8
25.0	4	96.1	-2.9	96.2	-2.9
40.0	2	83.1	-15.9	82.0	-17.1
	4	72.4	-26.6	70.9	-28.2

7.4. General Methods Used in the Biophysical Characterization of the Morrison-H and Morrison-L Antibodies

Buffer Exchange

[0513] Buffer exchange was carried out by dialysis. The antibodies were dialyzed in Spectra Pro 3 dialysis membranes (Spectrum Laboratories) with at least a 200-fold excess of dialysis buffer.

Concentrating of the Antibodies

[0514] Antibodies were concentrated using centrifugal concentrators with a molecular weight cut-off (MWCO) of 10 kDa or 30 kDa. Samples were centrifuged in 5 min steps at 22° C. until target concentration was reached. In between steps, the samples are resuspended.

Determination of Protein Concentration

[0515] The concentration of the protein samples was determined using a Tecan plate reader and a NanoQuant Plate. The buffer was used as blank to be subtracted from the absorbance measured at 280 nm. Each measurement was corrected for scattering caused by visible particles determined at 310 nm. The corrected value was normalized to a path length of 1 cm and the protein concentration calculated using the theoretical extinction coefficient of the corresponding protein. For protein samples with a concentration higher than 10 mg/ml, the sample was diluted in the corresponding buffer at least 10 times or to 1 mg/ml nominal concentration.

SDS-PAGE

[0516] SDS-PAGE was done under reducing conditions. After denaturation and reduction 5 µg of the sample were applied on a gradient gel (4%-20%). After separation, protein bands were stained with Coomassie Blue and the gel destained with purified water.

Storage

[0517] To assess the protein stability at different temperatures, samples were incubated at 4° C., 25° C. and 40° C. Storage at 4° C. was carried out in a fridge at a nominal temperature of 4° C. For storage at 25° C. and 40° C. samples were placed in cabinets with controlled humidity of 65% rH and 75% rH, respectively.

Thermal Unfolding by nanoDSF

[0518] Thermal unfolding using a thermal shift assay was determined by the change in fluorescence intensity of Sypro Orange. The fluorescence of the dye is sensitive to hydrophobic interactions. As the protein unfolds, hydrophobic amino acids are exposed to the solvent leading to an increased fluorescence of Sypro Orange. The resulting midpoint of thermal unfolding (T_m) as well as the onset temperature calculated at 10% of the maximal signal (T_{onset} is the temperature at: min. signal+0.1*(max. signal-min. signal)) was determined by fitting the data to a Boltzmann equation.

Monomeric Content by SE-HPLC

[0519] The monomeric content was determined by analytical size-exclusion chromatography using a Shodex KW403-4F column running in 50 mM sodium phosphate, 300 mM NaCl, pH 6.5. For analysis, 5 µg of sample were injected and the absorption recorded at 280 nm. The quality of the sample is stated as relative percentages of monomer, HMWS, and LMWS.

Cation Exchange HPLC (CEX-HPLC)

[0520] For the quantification of charge variants, analytical cation exchange chromatography was applied. The samples were analyzed on a MAbPac™ SCX-10 column. The charge variants were separated using a pH gradient from pH 4 to pH 10. For analysis, 30 µg of sample were injected and the absorption recorded at 280 nm. The quality of the sample is stated as relative percentages of target, acidic charge variants, and basic charge variants.

Mass Spectrometry

[0521] For identification and the exclusion of severe post-translational modifications, the intact mass of the Morrison antibodies was determined. Samples were reduced with DTT prior to ESI-MS analyses, desalted and analyzed in MeOH:2-PrOH:0.2% FA (30:20:50). Nano ESI-MS analyses of the samples were performed on a Syn-apt G2-Si mass spectrometer. The recorded m/z data were then deconvoluted into mass spectra by applying the maximum entropy algorithm MaxEnt1 (MaxLynx) with a resolution of the output mass 0.5 Da/channel and Uniform Gaussian Damage Model at the half height of 0.7 Da.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 50

<210> SEQ ID NO 1

<211> LENGTH: 11

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 1

Gly Phe Ser Ser Ser Ser Ala His Tyr Met Cys
1 5 10

<210> SEQ ID NO 2

<211> LENGTH: 18

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 2

Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser Trp Ala
1 5 10 15

Lys Gly

<210> SEQ ID NO 3

<211> LENGTH: 19

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 3

Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp Pro Tyr Tyr
1 5 10 15

Phe Ser Leu

<210> SEQ ID NO 4

<211> LENGTH: 129

<212> TYPE: PRT

<213> ORGANISM: Artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 4

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Ser Ser Ser Ala
20 25 30

His Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
35 40 45

Ile Gly Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser
50 55 60

Trp Ala Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
65 70 75 80

Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
85 90 95

Tyr Cys Ala Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp
100 105 110

-continued

Pro	Tyr	Tyr	Phe	Ser	Leu	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser
							115				120				125

Ser

<210> SEQ ID NO 5
 <211> LENGTH: 128
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 5

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Ser	Ser	Ser	Ala
			20					25					30		

His	Tyr	Met	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp
		35					40					45			

Ile	Gly	Cys	Ile	Tyr	Ala	Gly	Ser	Arg	Gly	Ser	Thr	Tyr	Tyr	Ala	Ser
	50					55					60				

Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Ala	Ser	Ser	Thr	Thr	Val
65					70					75					80

Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Thr	Tyr	Tyr
				85					90					95	

Cys	Ala	Arg	Ala	Ala	Phe	Val	Asn	Arg	Gly	Val	Ser	Trp	Ile	Trp	Pro
			100					105					110		

Tyr	Tyr	Phe	Ser	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser
		115				120						125			

<210> SEQ ID NO 6
 <211> LENGTH: 128
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 6

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly
1				5					10					15	

Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Ser	Ser	Ser	Ala
			20					25					30		

His	Tyr	Met	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp
		35					40					45			

Ile	Gly	Cys	Ile	Tyr	Ala	Gly	Ser	Arg	Gly	Ser	Thr	Tyr	Tyr	Ala	Ser
	50					55					60				

Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Ala	Ser	Ser	Thr	Thr	Val
65					70					75					80

Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Val	Tyr	Tyr
				85					90					95	

Cys	Ala	Arg	Ala	Ala	Phe	Val	Asn	Arg	Gly	Val	Ser	Trp	Ile	Trp	Pro
			100					105					110		

Tyr	Tyr	Phe	Ser	Leu	Trp	Gly	Gln	Gly	Thr	Leu	Val	Thr	Val	Ser	Ser
		115				120						125			

<210> SEQ ID NO 7
 <211> LENGTH: 11

-continued

<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 7

Gln Ala Ser Glu Asp Ile Glu Ser Tyr Leu Ala
1 5 10

<210> SEQ ID NO 8
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 8

Arg Ala Ser Thr Leu Ala Tyr
1 5

<210> SEQ ID NO 9
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 9

Leu Gly Ser Ser Tyr Ser Thr Gly Ser Asn Ile
1 5 10

<210> SEQ ID NO 10
<211> LENGTH: 99
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 10

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
85 90 95

Ser Asn Ile

<210> SEQ ID NO 11
<211> LENGTH: 99
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 11

-continued

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Arg Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Arg Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
 85 90 95
 Ser Asn Ile

<210> SEQ ID NO 12
 <211> LENGTH: 11
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 12

Gly Phe Ser Phe Ser Ala Asn Tyr Trp Ile Cys
 1 5 10

<210> SEQ ID NO 13
 <211> LENGTH: 18
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 13

Cys Ile Tyr Ile Gly Ser Arg Ala Asp Thr Tyr Tyr Ala Pro Trp Ala
 1 5 10 15

Lys Gly

<210> SEQ ID NO 14
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 14

Arg Phe Ser Ser Gly Ile Tyr Asp Leu Asp Arg Phe Phe Leu
 1 5 10

<210> SEQ ID NO 15
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 15

Gln Ser Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Ser Ala Asn
 20 25 30

-continued

```

Tyr Trp Ile Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
   35                               40               45

Ile Val Cys Ile Tyr Ile Gly Ser Arg Ala Asp Thr Tyr Tyr Ala Pro
   50                               55               60

Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ser Lys Asn Thr
   65                               70               75               80

Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
                        85               90               95

Phe Cys Ala Arg Phe Ser Ser Gly Ile Tyr Asp Leu Asp Arg Phe Phe
   100                               105               110

Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
   115                               120

```

```

<210> SEQ ID NO 16
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 16

```

```

Gln Ser Gln Leu Val Glu Ser Gly Gly Gly Arg Val Gln Pro Gly Gly
 1         5               10               15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Ser Ala Asn
 20        25               30

Tyr Trp Ile Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
   35                               40               45

Ile Val Cys Ile Tyr Ile Gly Ser Arg Ala Asp Thr Tyr Tyr Ala Pro
   50                               55               60

Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ser Lys Asn Thr
   65                               70               75               80

Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Thr Tyr
                        85               90               95

Phe Cys Ala Arg Phe Ser Ser Gly Ile Tyr Asp Leu Asp Arg Phe Phe
   100                               105               110

Leu Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser
   115                               120

```

```

<210> SEQ ID NO 17
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 17

```

```

Gln Ala Ser Gln Ser Ile Asp Ser Trp Leu Ala
 1         5               10

```

```

<210> SEQ ID NO 18
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 18

```

-continued

Gln Ala Ser Gln Ser Ile Glu Ser Trp Leu Ala
1 5 10

<210> SEQ ID NO 19
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 19

Gln Ala Ser Lys Leu Ala Ser
1 5

<210> SEQ ID NO 20
<211> LENGTH: 12
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 20

Gln Thr Tyr Tyr Gly Gly Gly His Ile Gly Trp Gly
1 5 10

<210> SEQ ID NO 21
<211> LENGTH: 100
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 21

Ala Phe Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Ser Ile Asp Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Pro Pro Lys Leu Leu Ile
35 40 45

Tyr Gln Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Thr Tyr Tyr Gly Gly Gly His
85 90 95

Ile Gly Trp Gly
100

<210> SEQ ID NO 22
<211> LENGTH: 100
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 22

Ala Phe Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Ser Ile Glu Ser Trp
20 25 30

-continued

```

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Pro Pro Lys Leu Leu Ile
   35                               40               45

Tyr Gln Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly
   50                               55               60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
   65                               70               75               80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Thr Tyr Tyr Gly Gly Gly His
           85                               90               95

Ile Gly Trp Gly
           100

```

```

<210> SEQ ID NO 23
<211> LENGTH: 122
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 23

```

```

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1                               10               15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Ser Ala Asn
   20                               25               30

Tyr Tyr Pro Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
   35                               40               45

Ile Gly Cys Ile Tyr Gly Gly Ser Ser Asp Ile Thr Tyr Asp Ala Asn
   50                               55               60

Trp Thr Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr
   65                               70               75               80

Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr
           85                               90               95

Tyr Cys Ala Arg Ser Ala Trp Tyr Ser Gly Trp Gly Gly Asp Leu Trp
           100               105               110

Gly Gln Gly Thr Leu Val Thr Val Ser Ser
           115               120

```

```

<210> SEQ ID NO 24
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 24

```

```

Gln Val Gln Leu Gln Glu Ser Gly Pro Gly Leu Val Lys Pro Ser Glu
 1                               10               15

Thr Leu Ser Leu Thr Cys Lys Val Ser Gly Phe Ser Phe Ser Asn Ser
   20                               25               30

Tyr Trp Ile Cys Trp Ile Arg Gln Pro Pro Gly Lys Gly Leu Glu Trp
   35                               40               45

Ile Gly Cys Thr Phe Val Gly Ser Ser Asp Ser Thr Tyr Tyr Ala Asn
   50                               55               60

Trp Ala Lys Gly Arg Val Thr Ile Ser Val Asp Ser Ser Lys Asn Gln
   65                               70               75               80

Phe Ser Leu Lys Leu Ser Ser Val Thr Ala Ala Asp Thr Ala Val Tyr
           85                               90               95

```

-continued

Tyr Cys Ala Arg His Pro Ser Asp Ala Val Tyr Gly Tyr Ala Asn Asn
100 105 110

Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
115 120

<210> SEQ ID NO 25
<211> LENGTH: 108
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 25

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Ser Ile Asn Asn Val
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Ser Thr Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Ser Ser Tyr Gly Asn Tyr Gly
85 90 95

Asp Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
100 105

<210> SEQ ID NO 26
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 26

Phe Gly Thr Gly Thr Lys Val Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 27
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 27

Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 28
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 28

Phe Gly Gly Gly Thr Gln Leu Ile Ile Leu Gly
1 5 10

-continued

<210> SEQ ID NO 29
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 29

Phe Gly Glu Gly Thr Glu Leu Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 30
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 30

Phe Gly Ser Gly Thr Lys Val Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 31
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 31

Phe Gly Gly Gly Thr Gln Leu Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 32
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 32

Phe Gly Gly Gly Thr Gln Leu Thr Ala Leu Gly
1 5 10

<210> SEQ ID NO 33
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 33

Phe Gly Cys Gly Thr Lys Val Thr Val Leu Gly
1 5 10

<210> SEQ ID NO 34
<211> LENGTH: 20
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 34

-continued

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly
1 5 10 15

Gly Gly Gly Ser
20

<210> SEQ ID NO 35
<211> LENGTH: 5
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 35

Gly Gly Gly Gly Ser
1 5

<210> SEQ ID NO 36
<211> LENGTH: 10
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 36

Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser
1 5 10

<210> SEQ ID NO 37
<211> LENGTH: 216
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 37

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Arg Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Arg Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
85 90 95

Ser Asn Ile Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val
100 105 110

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
115 120 125

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
130 135 140

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
145 150 155 160

Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
165 170 175

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys

-continued

180	185	190
Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr		
195	200	205
Lys Ser Phe Asn Arg Gly Glu Cys		
210	215	

<210> SEQ ID NO 38
 <211> LENGTH: 720
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

 <400> SEQUENCE: 38

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly		
1	5	10
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Ser Ser Ser Ala		
20	25	30
His Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp		
35	40	45
Ile Gly Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser		
50	55	60
Trp Ala Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr		
65	70	75
Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr		
85	90	95
Tyr Cys Ala Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp		
100	105	110
Pro Tyr Tyr Phe Ser Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser		
115	120	125
Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser		
130	135	140
Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp		
145	150	155
Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr		
165	170	175
Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr		
180	185	190
Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys		
195	200	205
Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp		
210	215	220
Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala		
225	230	235
Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro		
245	250	255
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val		
260	265	270
Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val		
275	280	285
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln		
290	295	300
Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln		

-continued

305					310						315					320
Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	
				325					330						335	
Leu	Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	
			340					345						350		
Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	
	355						360					365				
Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	
	370					375					380					
Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	
385				390					395						400	
Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	
			405						410					415		
Ser	Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	
		420						425					430			
Ser	Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	
	435						440						445			
Ser	Leu	Ser	Leu	Ser	Leu	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	
	450					455					460					
Ser	Ala	Phe	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	
465				470					475						480	
Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Asp	Ser	
			485						490					495		
Trp	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Pro	Pro	Lys	Leu	Leu	
		500						505					510			
Ile	Tyr	Gln	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	
	515					520						525				
Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	
	530				535						540					
Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Thr	Tyr	Tyr	Gly	Gly	Gly	
545				550					555						560	
His	Ile	Gly	Trp	Gly	Phe	Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu	Gly	
			565						570					575		
Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	
		580						585					590			
Gly	Gly	Gly	Ser	Gln	Ser	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg	Val	
	595					600						605				
Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	
	610				615					620						
Phe	Ser	Ala	Asn	Tyr	Trp	Ile	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	
625				630					635						640	
Gly	Leu	Glu	Trp	Ile	Val	Cys	Ile	Tyr	Ile	Gly	Ser	Arg	Ala	Asp	Thr	
			645					650						655		
Tyr	Tyr	Ala	Pro	Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Asp	Asn	
		660						665					670			
Ser	Lys	Asn	Thr	Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	
		675					680					685				
Thr	Ala	Thr	Tyr	Phe	Cys	Ala	Arg	Phe	Ser	Ser	Gly	Ile	Tyr	Asp	Leu	
	690					695					700					
Asp	Arg	Phe	Phe	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser	
705				710					715						720	

-continued

<210> SEQ ID NO 39
 <211> LENGTH: 216
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 39

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
 1 5 10 15
 Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
 20 25 30
 Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
 35 40 45
 Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
 50 55 60
 Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
 65 70 75 80
 Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
 85 90 95
 Ser Asn Ile Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val
 100 105 110
 Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
 115 120 125
 Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
 130 135 140
 Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
 145 150 155 160
 Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
 165 170 175
 Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
 180 185 190
 Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
 195 200 205
 Lys Ser Phe Asn Arg Gly Glu Cys
 210 215

<210> SEQ ID NO 40
 <211> LENGTH: 719
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 40

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Arg Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Ser Ser Ser Ala
 20 25 30
 His Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
 35 40 45
 Ile Gly Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser
 50 55 60
 Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Ala Ser Ser Thr Thr Val

-continued

65	70					75					80				
Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Thr	Tyr	Tyr
				85					90					95	
Cys	Ala	Arg	Ala	Ala	Phe	Val	Asn	Arg	Gly	Val	Ser	Trp	Ile	Trp	Pro
			100					105					110		
Tyr	Tyr	Phe	Ser	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser
		115				120						125			
Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg
	130					135					140				
Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr
145					150					155					160
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser
				165					170					175	
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser
			180					185					190		
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr
	195						200					205			
Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
	210					215					220				
Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro
225					230					235					240
Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys
			245						250					255	
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val
		260						265					270		
Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp
	275						280					285			
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe
	290				295						300				
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp
305					310						315				320
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu
			325						330					335	
Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg
			340					345					350		
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys
	355						360					365			
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp
	370					375					380				
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys
385					390					395					400
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser
				405					410					415	
Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser
			420					425					430		
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
		435					440						445		
Leu	Ser	Leu	Ser	Leu	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
	450					455					460				
Ala	Phe	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
465					470					475					480

-continued

Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Asp	Ser	Trp	
				485					490				495			
Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Pro	Pro	Lys	Leu	Leu	Ile	
				500					505				510			
Tyr	Gln	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	
				515					520				525			
Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	
				530					535				540			
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Thr	Tyr	Tyr	Gly	Gly	Gly	His	
				545					550				555			
Ile	Gly	Trp	Gly	Phe	Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu	Gly	Gly	
				565					570				575			
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	
				580					585				590			
Gly	Gly	Ser	Gln	Ser	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg	Val	Gln	
				595					600				605			
Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Phe	
				610					615				620			
Ser	Ala	Asn	Tyr	Trp	Ile	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	
				625					630				635			
Leu	Glu	Trp	Ile	Val	Cys	Ile	Tyr	Ile	Gly	Ser	Arg	Ala	Asp	Thr	Tyr	
				645					650				655			
Tyr	Ala	Pro	Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Asp	Asn	Ser	
				660					665				670			
Lys	Asn	Thr	Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	
				675					680				685			
Ala	Thr	Tyr	Phe	Cys	Ala	Arg	Phe	Ser	Ser	Gly	Ile	Tyr	Asp	Leu	Asp	
				690					695				700			
Arg	Phe	Phe	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser		
				705					710				715			

```
<210> SEQ ID NO 41
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence
```

<400> SEQUENCE: 41

Ala	Phe	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser	Val	Gly
1				5				10						15	
Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Asp	Ser	Trp
			20				25						30		
Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Pro	Pro	Lys	Leu	Leu	Ile
		35				40						45			
Tyr	Gln	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
	50					55					60				
Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
65					70					75					80
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Thr	Tyr	Tyr	Gly	Gly	Gly	His
				85					90					95	
Ile	Gly	Trp	Gly	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr
			100					105					110		

-continued

Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu
115 120 125

Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro
130 135 140

Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
145 150 155 160

Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr
165 170 175

Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His
180 185 190

Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val
195 200 205

Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 42
<211> LENGTH: 719
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 42

Gln Ser Gln Leu Val Glu Ser Gly Gly Gly Arg Val Gln Pro Gly Gly
1 5 10 15

Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Phe Ser Ala Asn
20 25 30

Tyr Trp Ile Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp
35 40 45

Ile Val Cys Ile Tyr Ile Gly Ser Arg Ala Asp Thr Tyr Tyr Ala Pro
50 55 60

Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Asp Asn Ser Lys Asn Thr
65 70 75 80

Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Thr Tyr
85 90 95

Phe Cys Ala Arg Phe Ser Ser Gly Ile Tyr Asp Leu Asp Arg Phe Phe
100 105 110

Leu Trp Gly Gln Gly Thr Gln Val Thr Val Ser Ser Ala Ser Thr Lys
115 120 125

Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg Ser Thr Ser Glu
130 135 140

Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro
145 150 155 160

Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser Gly Val His Thr
165 170 175

Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val
180 185 190

Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr Tyr Thr Cys Asn
195 200 205

Val Asp His Lys Pro Ser Asn Thr Lys Val Asp Lys Arg Val Glu Ser
210 215 220

Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala Pro Glu Phe Leu Gly
225 230 235 240

-continued

Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	245	250	255
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	260	265	270
Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	275	280	285
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	290	295	300
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	305	310	315
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	325	330	335
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	340	345	350
Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	355	360	365
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	370	375	380
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	385	390	395
Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Arg	Leu	Thr	Val	405	410	415
Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	420	425	430
His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	435	440	445
Leu	Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Asp	Ile	Gln	Met	450	455	460
Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Arg	Val	Gly	Asp	Arg	Val	Thr	465	470	475
Ile	Thr	Cys	Gln	Ala	Ser	Glu	Asp	Ile	Glu	Ser	Tyr	Leu	Ala	Trp	Tyr	485	490	495
Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile	Tyr	Arg	Ala	Ser	500	505	510
Thr	Leu	Ala	Tyr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	515	520	525
Arg	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro	Glu	Asp	Phe	Ala	530	535	540
Thr	Tyr	Tyr	Cys	Leu	Gly	Ser	Ser	Tyr	Ser	Thr	Gly	Ser	Asn	Ile	Phe	545	550	555
Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu	Gly	Gly	Gly	Gly	Gly	Ser	Gly	565	570	575
Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Glu	Val	580	585	590
Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Leu	Val	Gln	Pro	Gly	Gly	Ser	Leu	595	600	605
Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Ser	Ser	Ser	Ala	His	Tyr	610	615	620
Met	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	Ile	Gly	625	630	635
																		640

-continued

Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser Trp Ala
645 650 655

Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Val Tyr
660 665 670

Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
675 680 685

Ala Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp Pro Tyr
690 695 700

Tyr Phe Ser Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
705 710 715

<210> SEQ ID NO 43
<211> LENGTH: 217
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 43

Ala Phe Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Ser Ile Asp Ser Trp
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Pro Pro Lys Leu Leu Ile
35 40 45

Tyr Gln Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Thr Tyr Tyr Gly Gly Gly His
85 90 95

Ile Gly Trp Gly Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr
100 105 110

Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu
115 120 125

Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro
130 135 140

Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly
145 150 155 160

Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr
165 170 175

Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His
180 185 190

Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val
195 200 205

Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 44
<211> LENGTH: 718
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 44

-continued

Gln	Ser	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg	Val	Gln	Pro	Gly	Gly	
1				5					10					15		
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Phe	Ser	Ala	Asn	
			20					25					30			
Tyr	Trp	Ile	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp	
		35					40					45				
Ile	Val	Cys	Ile	Tyr	Ile	Gly	Ser	Arg	Ala	Asp	Thr	Tyr	Tyr	Ala	Pro	
	50					55					60					
Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Asp	Asn	Ser	Lys	Asn	Thr	
65					70					75					80	
Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Thr	Tyr	
				85					90					95		
Phe	Cys	Ala	Arg	Phe	Ser	Ser	Gly	Ile	Tyr	Asp	Leu	Asp	Arg	Phe	Phe	
			100					105					110			
Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser	Ala	Ser	Thr	Lys	
		115					120					125				
Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	Ser	Thr	Ser	Glu	
	130					135					140					
Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	Phe	Pro	Glu	Pro	
145					150					155				160		
Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	Gly	Val	His	Thr	
			165					170						175		
Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	Leu	Ser	Ser	Val	
			180					185					190			
Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr	Tyr	Thr	Cys	Asn	
	195						200					205				
Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Ser	
	210					215				220						
Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Phe	Leu	Gly	
225					230					235				240		
Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	
			245					250						255		
Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	Gln	
		260						265					270			
Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	
	275					280					285					
His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	Asn	Ser	Thr	Tyr	
	290					295					300					
Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	
305					310					315				320		
Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	Pro	Ser	Ser	Ile	
			325						330					335		
Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	
		340						345					350			
Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	Asn	Gln	Val	Ser	
		355					360					365				
Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	
	370					375					380					
Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	
385					390					395				400		


```

<210> SEQ ID NO 45
<211> LENGTH: 481
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 45

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Arg Val Gly
1             5             10            15
Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
          20            25            30

```

Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Ala	Pro	Lys	Leu	Leu	Ile
35							40					45			
Tyr	Arg	Ala	Ser	Thr	Leu	Ala	Tyr	Gly	Val	Pro	Ser	Arg	Phe	Ser	Gly
50						55					60				
Ser	Gly	Ser	Gly	Arg	Asp	Phe	Thr	Leu	Thr	Ile	Ser	Ser	Leu	Gln	Pro
65					70					75					80
Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Leu	Gly	Ser	Ser	Tyr	Ser	Thr	Gly
				85					90					95	
Ser	Asn	Ile	Phe	Gly	Gln	Gly	Thr	Lys	Val	Glu	Ile	Lys	Arg	Thr	Val
				100				105					110		
Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	Leu	Lys
115							120					125			
Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	Pro	Arg
130						135					140				
Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	Gly	Asn
145					150					155					160
Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	Tyr	Ser
				165					170					175	
Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	His	Lys
				180				185					190		
Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr
195						200						205			
Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly
210						215					220				
Gly	Ser	Ala	Phe	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser
225					230					235					240
Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Asp
				245					250					255	
Ser	Trp	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Pro	Pro	Lys	Leu
260								265					270		
Leu	Ile	Tyr	Gln	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe
275						280						285			
Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Thr	Leu	Thr	Ile	Ser	Ser	Leu
290						295					300				
Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Thr	Tyr	Tyr	Gly	Gly
305					310					315					320
Gly	His	Ile	Gly	Trp	Gly	Phe	Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu
				325					330					335	
Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
				340					345				350		
Gly	Gly	Gly	Gly	Ser	Gln	Ser	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg
355							360					365			
Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe
370						375					380				
Ser	Phe	Ser	Ala	Asn	Tyr	Trp	Ile	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly
385					390										

-continued

435	440	445
Asp Thr Ala Thr Tyr Phe Cys Ala Arg Phe Ser Ser Gly Ile Tyr Asp		
450	455	460
Leu Asp Arg Phe Phe Leu Trp Gly Gln Gly Thr Gln Val Thr Val Ser		
465	470	475 480
Ser		
<210> SEQ ID NO 46		
<211> LENGTH: 455		
<212> TYPE: PRT		
<213> ORGANISM: Artificial sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: artificial antibody-related sequence		
<400> SEQUENCE: 46		
Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly		
1	5	10 15
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Ser Ser Ser Ala		
20	25	30
His Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp		
35	40	45
Ile Gly Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser		
50	55	60
Trp Ala Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr		
65	70	75 80
Val Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr		
85	90	95
Tyr Cys Ala Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp		
100	105	110
Pro Tyr Tyr Phe Ser Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser		
115	120	125
Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser		
130	135	140
Arg Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp		
145	150	155 160
Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr		
165	170	175
Ser Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr		
180	185	190
Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys		
195	200	205
Thr Tyr Thr Cys Asn Val Asp His Lys Pro Ser Asn Thr Lys Val Asp		
210	215	220
Lys Arg Val Glu Ser Lys Tyr Gly Pro Pro Cys Pro Pro Cys Pro Ala		
225	230	235 240
Pro Glu Phe Leu Gly Gly Pro Ser Val Phe Leu Phe Pro Pro Lys Pro		
245	250	255
Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val		
260	265	270
Val Asp Val Ser Gln Glu Asp Pro Glu Val Gln Phe Asn Trp Tyr Val		
275	280	285
Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln		
290	295	300

-continued

```

Phe Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln
305              310              315              320

Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Gly
              325              330              335

Leu Pro Ser Ser Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro
              340              345              350

Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Gln Glu Glu Met Thr
              355              360              365

Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser
              370              375              380

Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr
385              390              395              400

Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr
              405              410              415

Ser Arg Leu Thr Val Asp Lys Ser Arg Trp Gln Glu Gly Asn Val Phe
              420              425              430

Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys
              435              440              445

Ser Leu Ser Leu Ser Leu Gly
              450              455

```

```

<210> SEQ ID NO 47
<211> LENGTH: 481
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

```

```

<400> SEQUENCE: 47

```

```

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1           5           10           15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
20          25          30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35          40          45

Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
50          55          60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65          70          75          80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
85          90          95

Ser Asn Ile Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val
100         105         110

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
115         120         125

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
130         135         140

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
145         150         155         160

Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
165         170         175

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
180         185         190

```

-continued

Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	Val	Thr
	195						200					205			
Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly
	210					215					220				
Gly	Ser	Ala	Phe	Gln	Leu	Thr	Gln	Ser	Pro	Ser	Ser	Leu	Ser	Ala	Ser
225					230					235					240
Val	Gly	Asp	Arg	Val	Thr	Ile	Thr	Cys	Gln	Ala	Ser	Gln	Ser	Ile	Asp
			245						250					255	
Ser	Trp	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Lys	Pro	Pro	Lys	Leu
		260						265					270		
Leu	Ile	Tyr	Gln	Ala	Ser	Lys	Leu	Ala	Ser	Gly	Val	Pro	Ser	Arg	Phe
	275						280					285			
Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Tyr	Thr	Leu	Thr	Ile	Ser	Ser	Leu
	290					295					300				
Gln	Pro	Glu	Asp	Phe	Ala	Thr	Tyr	Tyr	Cys	Gln	Thr	Tyr	Tyr	Gly	Gly
305					310					315					320
Gly	His	Ile	Gly	Trp	Gly	Phe	Gly	Thr	Gly	Thr	Lys	Val	Thr	Val	Leu
			325						330					335	
Gly	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser	Gly	Gly	Gly	Gly	Ser
			340					345				350			
Gly	Gly	Gly	Gly	Ser	Gln	Ser	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg
			355				360					365			
Val	Gln	Pro	Gly	Gly	Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe
	370					375					380				
Ser	Phe	Ser	Ala	Asn	Tyr	Trp	Ile	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly
385				390						395					400
Lys	Gly	Leu	Glu	Trp	Ile	Val	Cys	Ile	Tyr	Ile	Gly	Ser	Arg	Ala	Asp
			405					410						415	
Thr	Tyr	Tyr	Ala	Pro	Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Asp
			420					425					430		
Asn	Ser	Lys	Asn	Thr	Val	Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu
		435					440					445			
Asp	Thr	Ala	Thr	Tyr	Phe	Cys	Ala	Arg	Phe	Ser	Ser	Gly	Ile	Tyr	Asp
	450				455					460					
Leu	Asp	Arg	Phe	Phe	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser
465					470					475					480

Ser

<210> SEQ ID NO 48
 <211> LENGTH: 454
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 48

Glu	Val	Gln	Leu	Val	Glu	Ser	Gly	Gly	Gly	Arg	Val	Gln	Pro	Gly	Gly
1				5					10					15	
Ser	Leu	Arg	Leu	Ser	Cys	Ala	Ala	Ser	Gly	Phe	Ser	Ser	Ser	Ser	Ala
			20				25						30		
His	Tyr	Met	Cys	Trp	Val	Arg	Gln	Ala	Pro	Gly	Lys	Gly	Leu	Glu	Trp
		35				40					45				

-continued

Ile	Gly	Cys	Ile	Tyr	Ala	Gly	Ser	Arg	Gly	Ser	Thr	Tyr	Tyr	Ala	Ser	50	55	60
Trp	Ala	Lys	Gly	Arg	Phe	Thr	Ile	Ser	Lys	Ala	Ser	Ser	Thr	Thr	Val	65	70	75
Tyr	Leu	Gln	Met	Asn	Ser	Leu	Arg	Ala	Glu	Asp	Thr	Ala	Thr	Tyr	Tyr	85	90	95
Cys	Ala	Arg	Ala	Ala	Phe	Val	Asn	Arg	Gly	Val	Ser	Trp	Ile	Trp	Pro	100	105	110
Tyr	Tyr	Phe	Ser	Leu	Trp	Gly	Gln	Gly	Thr	Gln	Val	Thr	Val	Ser	Ser	115	120	125
Ala	Ser	Thr	Lys	Gly	Pro	Ser	Val	Phe	Pro	Leu	Ala	Pro	Cys	Ser	Arg	130	135	140
Ser	Thr	Ser	Glu	Ser	Thr	Ala	Ala	Leu	Gly	Cys	Leu	Val	Lys	Asp	Tyr	145	150	155
Phe	Pro	Glu	Pro	Val	Thr	Val	Ser	Trp	Asn	Ser	Gly	Ala	Leu	Thr	Ser	165	170	175
Gly	Val	His	Thr	Phe	Pro	Ala	Val	Leu	Gln	Ser	Ser	Gly	Leu	Tyr	Ser	180	185	190
Leu	Ser	Ser	Val	Val	Thr	Val	Pro	Ser	Ser	Ser	Leu	Gly	Thr	Lys	Thr	195	200	205
Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys	210	215	220
Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro	225	230	235
Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	245	250	255
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	260	265	270
Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp	275	280	285
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe	290	295	300
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	305	310	315
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu	325	330	335
Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	340	345	350
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys	355	360	365
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	370	375	380
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	385	390	395
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	405	410	415
Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser	420	425	430
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	435	440	445
Leu	Ser	Leu	Ser	Leu	Gly													

-continued

450

<210> SEQ ID NO 49
<211> LENGTH: 481
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 49

Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser Val Gly
1 5 10 15

Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Glu Asp Ile Glu Ser Tyr
20 25 30

Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys Leu Leu Ile
35 40 45

Tyr Arg Ala Ser Thr Leu Ala Tyr Gly Val Pro Ser Arg Phe Ser Gly
50 55 60

Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Ser Leu Gln Pro
65 70 75 80

Glu Asp Phe Ala Thr Tyr Tyr Cys Leu Gly Ser Ser Tyr Ser Thr Gly
85 90 95

Ser Asn Ile Phe Gly Gln Gly Thr Lys Val Glu Ile Lys Arg Thr Val
100 105 110

Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln Leu Lys
115 120 125

Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr Pro Arg
130 135 140

Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser Gly Asn
145 150 155 160

Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr Tyr Ser
165 170 175

Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys His Lys
180 185 190

Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro Val Thr
195 200 205

Lys Ser Phe Asn Arg Gly Glu Cys Gly Gly Gly Gly Ser Gly Gly Gly
210 215 220

Gly Ser Ala Phe Gln Leu Thr Gln Ser Pro Ser Ser Leu Ser Ala Ser
225 230 235 240

Val Gly Asp Arg Val Thr Ile Thr Cys Gln Ala Ser Gln Ser Ile Asp
245 250 255

Ser Trp Leu Ala Trp Tyr Gln Gln Lys Pro Gly Lys Pro Pro Lys Leu
260 265 270

Leu Ile Tyr Gln Ala Ser Lys Leu Ala Ser Gly Val Pro Ser Arg Phe
275 280 285

Ser Gly Ser Gly Ser Gly Thr Asp Tyr Thr Leu Thr Ile Ser Ser Leu
290 295 300

Gln Pro Glu Asp Phe Ala Thr Tyr Tyr Cys Gln Thr Tyr Tyr Gly Gly
305 310 315 320

Gly His Ile Gly Trp Gly Phe Gly Thr Gly Thr Lys Val Thr Val Leu
325 330 335

Gly Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Gly Gly Gly Ser

-continued

340	345	350
Gly Gly Gly Gly Ser Gln Ser	Gln Leu Val Glu Ser	Gly Gly Gly Arg
355	360	365
Val Gln Pro Gly Gly Ser	Leu Arg Leu Ser Cys	Ala Ala Ser Gly Phe
370	375	380
Ser Phe Ser Ala Asn Tyr Trp	Ile Cys Trp Val Arg	Gln Ala Pro Gly
385	390	395
Lys Gly Leu Glu Trp Ile Val	Cys Ile Tyr Ile Gly	Ser Arg Ala Asp
405	410	415
Thr Tyr Tyr Ala Pro Trp	Ala Lys Gly Arg Phe	Thr Ile Ser Lys Asp
420	425	430
Asn Ser Lys Asn Thr Val Tyr	Leu Gln Met Asn Ser	Leu Arg Ala Glu
435	440	445
Asp Thr Ala Thr Tyr Phe	Cys Ala Arg Phe Ser	Ser Gly Ile Tyr Asp
450	455	460
Leu Asp Arg Phe Phe Leu	Trp Gly Gln Gly Thr	Gln Val Thr Val Ser
465	470	475
480		

Ser

<210> SEQ ID NO 50
 <211> LENGTH: 454
 <212> TYPE: PRT
 <213> ORGANISM: Artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: artificial antibody-related sequence

<400> SEQUENCE: 50

Glu Val Gln Leu Val Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly	
1 5 10 15	
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Ser Ser Ser Ala	
20 25 30	
His Tyr Met Cys Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp	
35 40 45	
Ile Gly Cys Ile Tyr Ala Gly Ser Arg Gly Ser Thr Tyr Tyr Ala Ser	
50 55 60	
Trp Ala Lys Gly Arg Phe Thr Ile Ser Lys Ala Ser Ser Thr Thr Val	
65 70 75 80	
Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr	
85 90 95	
Cys Ala Arg Ala Ala Phe Val Asn Arg Gly Val Ser Trp Ile Trp Pro	
100 105 110	
Tyr Tyr Phe Ser Leu Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser	
115 120 125	
Ala Ser Thr Lys Gly Pro Ser Val Phe Pro Leu Ala Pro Cys Ser Arg	
130 135 140	
Ser Thr Ser Glu Ser Thr Ala Ala Leu Gly Cys Leu Val Lys Asp Tyr	
145 150 155 160	
Phe Pro Glu Pro Val Thr Val Ser Trp Asn Ser Gly Ala Leu Thr Ser	
165 170 175	
Gly Val His Thr Phe Pro Ala Val Leu Gln Ser Ser Gly Leu Tyr Ser	
180 185 190	
Leu Ser Ser Val Val Thr Val Pro Ser Ser Ser Leu Gly Thr Lys Thr	
195 200 205	

-continued

Tyr	Thr	Cys	Asn	Val	Asp	His	Lys	Pro	Ser	Asn	Thr	Lys	Val	Asp	Lys
210						215					220				
Arg	Val	Glu	Ser	Lys	Tyr	Gly	Pro	Pro	Cys	Pro	Pro	Cys	Pro	Ala	Pro
225					230					235					240
Glu	Phe	Leu	Gly	Gly	Pro	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys
			245						250					255	
Asp	Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val
			260					265					270		
Asp	Val	Ser	Gln	Glu	Asp	Pro	Glu	Val	Gln	Phe	Asn	Trp	Tyr	Val	Asp
		275					280					285			
Gly	Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Phe
	290					295					300				
Asn	Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp
305					310						315				320
Trp	Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Gly	Leu
			325						330					335	
Pro	Ser	Ser	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg
			340					345					350		
Glu	Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Gln	Glu	Glu	Met	Thr	Lys
		355					360					365			
Asn	Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp
		370				375					380				
Ile	Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys
385					390					395					400
Thr	Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser
				405					410					415	
Arg	Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Glu	Gly	Asn	Val	Phe	Ser
			420					425					430		
Cys	Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser
		435					440					445			
Leu	Ser	Leu	Ser	Leu	Gly										
		450													

1. A multispecific antibody comprising:

- a) one or two binding domains, which specifically bind to IL-4R (IL4R-BD), and
- b) one or two binding domains, which specifically bind to IL-31 (IL31-BD),

wherein

the multispecific antibody comprises an immunoglobulin Fc region;

each of said IL4R-BDs comprise

the HCDR1 sequence of SEQ ID NO: 1,

the HCDR2 sequences of SEQ ID NO: 2,

the HCDR3 sequence of SEQ ID NO: 3,

the LCDR1 sequence of SEQ ID NO: 7,

the LCDR2 sequence of SEQ ID NO: 8, and

the LCDR3 sequences of SEQ ID NO: 9; and

each of said IL31-BDs comprises

the HCDR1 sequence of SEQ ID NO: 12,

the HCDR2 sequence of SEQ ID NO: 13,

the HCDR3 sequences of SEQ ID NO: 14,

the LCDR1 sequence of SEQ ID NO: 17 or 18,

the LCDR2 sequence of SEQ ID NO: 19, and

the LCDR3 sequences of SEQ ID NOs: 20.

2. The multispecific antibody of claim 1, wherein the immunoglobulin Fc region is selected from an IgG subclass, particularly from IgG subclasses IgG1 and IgG4.

3. The multispecific antibody of claim 2, wherein the format of said multispecific antibody is selected from bivalent bispecific IgG formats, trivalent bispecific IgG formats and tetravalent bispecific IgG formats.

4. The multispecific antibody of claim 3, wherein the format of said multispecific antibody is selected from Morrison formats.

5. The multispecific antibody of claim 1, wherein said multispecific antibody comprises two IL4R-BDs and/or two IL31-BDs.

6. The multispecific antibody of claim 5, wherein the IL4R-BDs are located in the Fab arms, and the IL31-BDs are located in the scFv part of the Morrison format.

7. The multispecific antibody of claim 4, wherein the format of the multispecific antibody is selected from a Morrison-L and a Morrison-H format.

8. (canceled)
9. The multispecific antibody of claim 8, wherein said IL4R-BDs and said IL31-BDs, when being in scFv-format, comprise
- (i) VL domains comprising a VL framework comprising framework regions FR1, FR2 and FR3, which are selected from VK subtypes, particularly from the VK1 and VK3 subtypes, particularly are of the VK1 subtype, and a framework FR4, which is selected from a VK FR4 and a VA FR4, particularly is a VA FR4 comprising an amino acid sequence having at least 70, 80, 90 percent identity to an amino acid sequence selected from SEQ ID NO: 26 to SEQ ID NO: 33, more particularly a VA FR4 selected from any of SEQ ID NO: 26 to SEQ ID NO: 33, particularly a VA FR4 according to SEQ ID NO: 26 or 33.
10. The multispecific antibody of claim 1, wherein said IL4R-BDs comprise
- a) a VH sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to any one of the amino acid sequences selected from SEQ ID NO: 4, 5 and 6; and
 - b) a VL sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to any one of the amino acid sequences selected from SEQ ID NO: 10 and 11,
- in particular wherein said IL4R-BDs comprise
- a) a VH sequence of SEQ ID NO: 4 and a VL sequence of SEQ ID NO: 11; or
 - b) a VH sequence of SEQ ID NO: 5 and a VL sequence of SEQ ID NO: 10; or
 - c) a VH sequence of SEQ ID NO: 6 and a VL sequence of SEQ ID NO: 10.
11. The multispecific antibody of claim 1, wherein said IL31-BDs comprise
- a) a VH sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to any one of the amino acid sequences selected from SEQ ID NOs: 15 and 16; and
 - b) a VL sequence at least 90, 91, 92, 93, 94, 95, 96, 97, 98 or 99 percent identical to any one of the amino acid sequences selected from SEQ ID NOs: 21 and 22,
- in particular wherein said IL31-BD comprises
- a) a VH sequence of SEQ ID NO: 15 and a VL sequence of SEQ ID NO: 22; or
 - b) a VH sequence of SEQ ID NO: 16 and a VL sequence of SEQ ID NO: 21.
12. (canceled)
13. A nucleic acid or two nucleic acids encoding the multispecific antibody of claim 1.
14. A vector or two vectors comprising the nucleic acid or the two nucleic acids of claim 13.
15. A host cell or host cells comprising the vector or the two vectors of claim 14.
16. A method for producing the multispecific antibody of claim 1, comprising (i) providing the nucleic acid or the two nucleic acids encoding the multispecific antibody of claim 1, expressing said nucleic acid or said two nucleic acids, and collecting said multispecific antibody from the expression system.
17. A pharmaceutical composition comprising the multispecific antibody of claim 1 and a pharmaceutically acceptable carrier.
18. The multispecific antibody of claim 1 for use in the treatment of a disease selected from allergic, inflammatory and autoimmune diseases, particularly from pruritus-causing allergic diseases, pruritus-causing inflammatory diseases and pruritus-causing autoimmune diseases, particularly from atopic dermatitis, acute allergic contact dermatitis, chronic spontaneous urticaria, bullous pemphigoid, alopecia areata, dermatomyositis, prurigo nodularis, psoriasis and atopic asthma; particularly wherein said disease is atopic dermatitis.
19. The multispecific antibody of claim 2, wherein the immunoglobulin Fc region is selected from IgG subclasses IgG1 and IgG4.
20. The multispecific antibody of claim 19, wherein the immunoglobulin Fc region is selected from IgG subclass IgG4.
21. The multispecific antibody of claim 3, wherein the format of said multispecific antibody is selected from KiH-based IgGs; DVD-Ig; CODV-IgG and Morrison (IgG CH₃-scFv fusion (Morrison-H) or IgG CL-scFv fusion (Morrison-L)).
22. The multispecific antibody of claim 7, wherein the format of the multispecific antibody is the Morrison-H format.
23. The multispecific antibody of claim 1, which is selected from
- (a) a Morrison antibody having the following heavy chain (HC) and light chain (LC) sequences: HC sequence of SEQ ID NO: 38 and LC sequence of SEQ ID NO: 37; and
 - (b) a Morrison antibody having the following heavy chain (HC) and light chain (LC) sequences: HC sequence of SEQ ID NO: 40 and an LC sequence of SEQ ID NO: 39.
24. A pharmaceutical composition comprising the multispecific antibody of claim 23 and a pharmaceutically acceptable carrier.
25. (canceled)

* * * * *