

(19) **United States**
(12) **Patent Application Publication**
IZHAR et al.
(10) **Pub. No.: US 2025/0283115 A1**
(43) **Pub. Date: Sep. 11, 2025**

(54) **ENGINEERED HIGH FIDELITY OMNI-50 NUCLEASE VARIANTS**
filed on Apr. 18, 2022, provisional application No. 63/333,037, filed on Apr. 20, 2022.

(71) Applicant: **EmendoBio Inc.**, Wilmington, DE (US)

(72) Inventors: **Lior IZHAR**, Tel Aviv (IL); **Rafi EMMANUEL**, Ramla (IL); **Liat ROCKAH**, Rishon Lezion (IL); **Asael HERMAN**, Ness-ziona (IL)

(73) Assignee: **EmendoBio Inc.**, Wilmington, DE (US)

(21) Appl. No.: **18/682,982**

(22) PCT Filed: **Aug. 12, 2022**

(86) PCT No.: **PCT/US2022/074930**

§ 371 (c)(1),
(2) Date: **Feb. 12, 2024**


Related U.S. Application Data

(60) Provisional application No. 63/232,571, filed on Aug. 12, 2021, provisional application No. 63/332,214,

Publication Classification
(51) **Int. Cl.**
C12N 15/90 (2006.01)
C12N 9/22 (2006.01)
C12N 15/11 (2006.01)
(52) **U.S. Cl.**
CPC **C12N 15/907** (2013.01); **C12N 9/226** (2025.05); **C12N 15/11** (2013.01); **C12N 2310/20** (2017.05)

(57) **ABSTRACT**
The present invention is directed to, inter alia, composition and methods for genome editing. Specifically, a non-naturally occurring OMNI-50 nuclease variant having a wild-type OMNI-50 protein sequence (SEQ ID NO: 1) comprising an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, V981, K965, and K1036.

Specification includes a Sequence Listing.






Fig. 1




Fig. 2




Fig. 3A





Fig. 3B




Fig. 4A




Fig. 4B




Fig. 4C




Fig. 5A




Fig. 5B




Fig. 5C




Fig. 6A




Fig. 6B




Fig. 6C




Fig. 7A




Fig. 7B




Fig. 8A




Fig. 8B




Fig. 8C




Fig. 8D




Fig. 8E




Fig. 8F




Fig. 8G




Fig. 8H




Fig. 9A




Fig. 9B




Fig. 9C




Fig. 9D




Fig. 9E




Fig. 9F

G




Fig. 9G

H




Fig. 9H




Fig. 9I




Fig. 9J




Fig. 9K

A

SNP ID	Heterozygosity frequency	
	Healthy population	Patient population
rs10414837	33%	43%
rs3761005	46%	45%
rs1683564	42%	42%

Fig. 10A

B




Fig. 10B




Fig. 11





Fig. 12






Fig. 13B




Fig. 13C




Fig. 14A




Fig. 14B




Fig. 15A




Fig. 15B




Fig. 15C




Fig. 16A

B




Fig. 16B




Fig. 16C





Fig. 16D






Fig. 17B



Fig. 18A



Fig. 18B




Fig. 19A




Fig. 19B




Fig. 20A




Fig. 20B




Fig. 20C

Chromosome	Position	Gene	Type	Sequence	Mismatches	Reads 5CN-P41	Reads 5CN-P55
sgRNA-Int4-const							
chr15	90957975	RCCD1	intron	aCAGTCCtGGCTGGGAGCaGGTGGG	3	0	0.03±0
chr17	74081352	LINC02074	intron	aCAGTCCGGGCTGGGAGCGgAGG	3	0	0.03±0
sgRNA-164133-alt							
chr22	28908576	ZNRF3	intron	GTG>CArTCCCCArAGsACACAGGG	3		0
chr2	26950233	DPYSL5	exon	GTGTCAAcCCCaAGAcGACACATGG	3		0.01±0.01
sgRNA-164133-ref							
chr12	58121885	NA	intergenic	GTGTgAAGgCCCAGAGGCCAaAGGG	3	0.07±0.02	
chr12	116915161	FBXW8	intron	GTtgCAAGCCCCAGAGcCCACAGGG	3	0.03±0.03	


Fig. 20D



Figs. 21A-J



Figs. 22A-I



Figs. 23A-I

ENGINEERED HIGH FIDELITY OMNI-50 NUCLEASE VARIANTS

[0001] This application is a U.S. National Stage of PCT International Application No. PCT/US2022/074930 filed Aug. 12, 2022, which claims priority to of U.S. Provisional Application Nos. 63/333,037 filed Apr. 20, 2022, 63/332,214 filed Apr. 18, 2022, and 63/232,571 filed Aug. 12, 2021, the contents of each of which are hereby incorporated by reference.

[0002] Throughout this application, various publications are referenced, including referenced in parenthesis. The disclosures of all publications mentioned in this application in their entireties are hereby incorporated by reference into this application in order to provide additional description of the art to which this invention pertains and of the features in the art which can be employed with this invention.

REFERENCE TO SEQUENCE LISTING

[0003] This application incorporates-by-reference nucleotide sequences which are present in the file named “ENGOMNI50_5798-0046US01.xml”, which is 290,097 bytes in size, and which was created on May 12, 2025 in the IBM-PC machine format, having an operating system compatibility with MS-Windows, which is contained in the XML file filed May 12, 2025 as part of this application.

BACKGROUND OF INVENTION

[0004] Targeted genome modification is a powerful tool that can be used to reverse the effect of pathogenic genetic variations and therefore has the potential to provide new therapies for human genetic diseases. Current genome engineering tools, including engineered zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs), and most recently, RNA-guided DNA endonucleases such as CRISPR/Cas, produce sequence-specific DNA breaks in a genome. The modification of the genomic sequence occurs at the next step and is the product of the activity of a cellular DNA repair mechanism triggered in response to the newly formed DNA break. These mechanisms may include, for example: (1) classical non-homologous end-joining (NHEJ) in which the two ends of the break are ligated together in a fast but also inaccurate manner (i.e. frequently resulting in mutation of the DNA at the cleavage site in the form of small insertion or deletions) or (2) homology-directed repair (HDR) in which an intact homologous DNA donor is used to replace the DNA surrounding the cleavage site in an accurate manner. Minimal off-target activity of the initial DNA damage inducer is required for efficient and safe genome editing.

SUMMARY OF THE INVENTION

[0005] Disclosed herein are engineered Clustered Regularly Interspaced Short Palindromic Repeats (CRISPRs)/CRISPR-associated OMNI-50 nucleases with altered and improved target specificity and their use in genomic engineering, epigenomic engineering, genome targeting, genome editing, and in vitro diagnostics.

[0006] In some embodiments, there is provided a variant of an OMNI-50 nuclease with increased specificity as compared to the wild-type OMNI-50 nuclease, as well as methods of using the improved variants. Advantageously, when the engineered variant OMNI-50 nucleases are active in a

CRISPR endonuclease system, the CRISPR endonuclease system displays reduced off-target editing activity and maintained on-target editing activity relative to a wild-type CRISPR endonuclease system in which a wild-type OMNI-50 nuclease is active. For example, an engineered variant OMNI-50 nuclease may display improved allele-specific discrimination, e.g. specific binding and activity at a target region which contains a heterozygous SNP present in only the targeted allele and not present in the non-targeted allele.

[0007] In some embodiments, there is provided a variant of an OMNI-50 nickase with increased specificity as compared to the wild-type OMNI-50 nickase. In some embodiments, there is provided a variant of an OMNI-50 dead nuclease with increased specificity as compared to the wild-type OMNI-50 dead nuclease. For example, the catalytic site of any one of the OMNI-50 nuclease variants provided herein may be modified such that the variant has nickase activity, such that it is capable of performing single-strand DNA cuts. Alternatively, the catalytic site of any one of the OMNI-50 nuclease variants provided herein may be modified such that the variant has no nuclease activity, i.e. a dead nuclease.

[0008] According to some embodiments of the present invention, there is provided a variant of OMNI-50 nuclease protein comprising a sequence that is at least 80% identical to the amino acid sequence of wild-type OMNI-50 nuclease protein (SEQ ID NO: 1).

[0009] According to some embodiments of the present invention, there is provided a non-naturally occurring OMNI-50 nuclease variant having a wild-type OMNI-50 protein sequence (SEQ ID NO: 1) comprising an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036.

[0010] According to some embodiments of the present invention, there is provided a CRISPR system comprising any one of the OMNI-50 nuclease variant disclosed herein complexed with a guide RNA molecule that targets a DNA target site, wherein the CRISPR system displays reduced off-target editing activity relative to a wild-type CRISPR system comprising a wild-type OMNI-50 nuclease protein and the guide RNA molecule.

[0011] In some embodiments, the OMNI-50 variant nuclease exhibits increased specificity to a target site when complexed with a guide RNA targeting the OMNI-50 variant to the target site compared to a wild-type OMNI-50 nuclease (SEQ ID NO: 1).

[0012] In some embodiments, the OMNI-50 nuclease variant is a nickase having an inactivated RuvC domain created by an amino acid substitution at a position provided for the CRISPR nuclease in column 1 of the table below. In some embodiments, the nickase further comprises an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036.

[0013] In some embodiments, the OMNI-50 nuclease variant is a nickase having an inactivated HNH domain created by an amino acid substitution at a position provided for the CRISPR nuclease in column 2 of the table below. In some embodiments, the nickase further comprises an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036.

[0014] In some embodiments, the OMNI-50 nuclease variant is a catalytically dead nuclease having an inactivated RuvC domain and an inactivated HNH domain created by substitutions at the positions provided for the CRISPR nuclease in column 3 of the table below. In some embodiments, the catalytically dead nuclease comprises an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036.

TABLE

Positions affecting OMNI-50 nuclease activity		
Nickase having inactivated RuvC domain	Nickase having inactivated HNH domain	Dead nuclease having inactivated RuvC and HNH domains
D12 or E785 or H1008 or D1011	D875* or H876 or N899	(D12 or E785 or H1008 or D1011) and (D875* or H876 or N899)

[0015] The above table lists alternative positions to be substituted to generate a nickase having an inactivated RUV domain, alternative positions to be substituted to generate a nickase having an inactivated HNH domain, and alternative positions to be substituted to generate a catalytically dead nuclease having inactivated RUV and HNH domains. Substitution to any other amino acid is permissible for each of the amino acid positions indicated in columns 1-3, except if followed by an asterisk, which indicates that any substitution other than aspartic acid (D) to glutamic acid (E) or glutamic acid (E) to aspartic acid (D) results in inactivation.

[0016] Throughout the text OMNI-50 nuclease variants are referred to, however, any of these variants may be modified to have nickase activity (i.e. nucleases which create a single-strand DNA break as opposed to a double-strand break) or to have no nuclease activity (i.e. a catalytically dead nuclease).

[0017] Accordingly, point mutations can be introduced into any one of the variants described herein to modify or abolish their nuclease activity while still retaining their ability to specifically bind DNA in a sgRNA-programmed manner. Any one of these variants can specifically target a desired DNA target sequence via a guide RNA molecule. The variant-guide complex will also carry any molecule attached to the complex to the target site. Thus, this disclosure also contemplates fusion proteins comprising such variants and a DNA modifying domain (e.g., a deaminase, a nuclease, a nickase, a recombinase, a methyltransferase, a methylase, an acetylase, an acetyltransferase, a transcriptional activator, or a transcriptional repressor domain), as well as the use of such fusion proteins in correcting mutations in a genome (e.g., the genome of a human subject) that are associated with disease, or generating mutations in a genome (e.g., the human genome) to decrease or prevent expression of a gene.

[0018] In some embodiments, any of the variants provided herein may be fused to a protein that has an enzymatic activity. In some embodiments, the enzymatic activity modifies a target DNA. In some embodiments, the enzymatic activity is nuclease activity, methyltransferase activity, demethylase activity, DNA repair activity, DNA damage activity, deamination activity, dismutase activity, alkylation activity, depurination activity, oxidation activity, pyrimidine dimer

forming activity, integrase activity, transposase activity, recombinase activity, polymerase activity, ligase activity, helicase activity, photolyase activity or glycosylase activity. In some cases, the enzymatic activity is nuclease activity. In some cases, the nuclease activity introduces a double strand break in the target DNA. In some cases, the enzymatic activity modifies a target polypeptide associated with the target DNA. In some cases, the enzymatic activity is methyltransferase activity, demethylase activity, acetyltransferase activity, deacetylase activity, kinase activity, phosphatase activity, ubiquitin ligase activity, deubiquitinating activity, adenylation activity, deadenylation activity, SUMOylating activity, deSUMOylating activity, ribosylation activity, deribosylation activity, myristoylation activity or demyristoylation activity. In some cases, the target polypeptide is a histone and the enzymatic activity is methyltransferase activity, demethylase activity, acetyltransferase activity, deacetylase activity, kinase activity, phosphatase activity, ubiquitin ligase activity or deubiquitinating activity.

[0019] Thus, any one of the OMNI-50 nuclease, nickase, or dead-nuclease variants may be fused (e.g. directly fused or fused via a linker) to another DNA modulating or DNA modifying enzyme, including, but not limited to, base editors such as a deaminase, a reverse transcriptase (e.g. for use in prime editing, see Anzaolone et al. (2019)), an enzyme that modifies the methylation state of DNA (e.g. a methyltransferase), or a modifier of histones (e.g. a histone acetyltransferase). Indeed, the OMNI-50 nuclease, nickase, inactive variants described herein may be fused to a DNA modifying enzyme or an effector domain thereof. Examples of DNA modifiers include but are not limited to: a deaminase, a nuclease, a nickase, a recombinase, a methyltransferase, a methylase, an acetylase, an acetyltransferase, a reverse transcriptase, an helicase, an integrase, a ligase, a transposase, a demethylase, a phosphatase, a transcriptional activator, or a transcriptional repressor. In some embodiments, any of the OMNI-50 variants provided herein are fused to a protein that has an enzymatic activity. In some embodiments, the enzymatic activity modifies a target DNA molecule. the OMNI-50 variants described herein or fusion proteins thereof, may be used to correct or generate one or more mutations in a gene associated with disease, or to increase, correct, decrease or prevent expression of a gene.

[0020] According to some embodiments of the present invention, there is provided a method for gene editing having reduced off-target editing activity, comprising contacting a DNA target site with an active CRISPR system comprising any one of the OMNI-50 nuclease variant proteins described herein.

[0021] According to some embodiments, there is provided a method for gene editing having reduced off-target editing activity and/or increased on-target editing activity, comprising: contacting a target site locus with an active CRISPR system comprising a variant OMNI-50 nuclease protein of any one of the variants described herein, wherein the active CRISPR system displays reduced off-target editing activity and maintained on-target editing activity relative to a wild-type CRISPR system having a wild-type OMNI-50 nuclease protein.

[0022] Further embodiments and the full scope of applicability of the present invention will become apparent from the detailed description given hereinafter. However, it should be understood that the detailed description and specific examples, while indicating preferred embodiments

of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE FIGURES

[0023] FIG. 1: A schematic showing the nuclease optimization platform is shown. Libraries of nuclease variants are created using a combination of rational design and random mutagenesis, followed by selection for activity, specificity, multiplexing (and any other desired trait) using specialized selection assays.

[0024] FIG. 2: A schematic of the ELANE gene and target sites used to test activity and fidelity of wild-type OMNI-50 nuclease and its variants. “Alt” and “ref” indicate guide RNA molecules that target a nuclease to a SNP position in an allele having either the “alt” version of the SNP or having the “ref” version of the SNP, respectively.

[0025] FIGS. 3A-3B: Allele specificity and off-target effects of wild-type OMNI-50 nuclease, demonstrating that the OMNI-50 nuclease could be improved by optimization. FIG. 3A: HSCs edited by wild-type OMNI-50 alt composition showed elimination of the alternative allele while the reference allele was kept intact. However, editing by wild-type OMNI-50 ref composition resulted in a reduction in the reference allele and only partial preservation of the alternative allele, providing a differential but not complete allele specific editing. FIG. 3B: rhAmpseq analysis in healthy (HD) and Patient (ESCN-2)-edited HSCs (RNP(ref) or RNP(alt)) on off-targets identified for constant guide (Sg(constant)), reference guide (Sg(ref)) and alternative guide (Sg(alt)). Results indicate that wild-type OMNI-50 has one off-target for each guide that should be eliminated to ensure target-specific editing. % Editing in RNPref treated HSCs is shown in the top panel; % Editing in RNPalt treated HSCs is shown in the bottom panel. Statistical significance is indicated as **P<0.01, ***P<0.0001.

[0026] FIGS. 4A-4C: Optimized variants show improved allele specificity and comparable activity compared to wild-type OMNI-50 nuclease. FIG. 4A: Allele specificity was determined in healthy HSCs edited with wild-type OMNI-50 or optimized variants. Variants show superior allele discrimination compared with wild-type OMNI-50. Specifically, a graph representing allele specific editing with sgRNA-564DS-ref or sgRNA-564DS-alt measured by ddPCR is shown. Editing specificity is determined by two competitive probes, FAM probe which binds the alternative allele and HEX probe which binds the reference allele, governing decrease signal from the edited allele. The ratio between the concentration of the reference allele to the alternative allele in heterozygote untreated cells is 1. The graph represents average concentration of reference allele (HEX), alternative allele (FAM), normalized to endogenous gene control RPP30 and STAT1 for each gDNA sample. FIG. 4B: % excision was measured in healthy HSCs edited with wild-type OMNI-50 (WT OMNI-50) or optimized variants. Variant 3795 shows comparable activity compared with WT OMNI-50. Variant 3795 showed the highest allele specificity without compromising on activity. FIG. 4C: Variant fidelity was measured by next-generation sequence (NGS) analysis specific to known OMNI-50 nuclease off-target. Note, all variants depicted high fidelity.

[0027] FIGS. 5A-5C: rhAmpSeq analysis of healthy HSCs edited using either a WT OMNI-50 or the Variant 3795

nuclease on off-targets identified for the constant guide (Sg(constant)), reference guide (Sg(ref)) (RNP(ref), FIG. 5A), and alternative guide (Sg(alt)) molecules (RNP(alt)), FIG. 5B). FIG. 5C shows additional results validating the fidelity of the Variant 3795 nuclease are demonstrated by examining the off-targets when using three different guide molecules (g35, g62Ref, or g62Alt) on HSC samples with either a WT OMNI-50 or Variant 3795 nuclease (FIG. 5C). These results show that there is a strong validated off-target effect for each guide when using the WT OMNI-50 nuclease, but no off-targets were validated with the Variant 3795 nuclease, indicating that the Variant 3795 nuclease eliminates off-targets compared to WT OMNI-50 nuclease.

[0028] FIGS. 6A-6C: Mono-allelic excision in ELANE restores neutrophil differentiation. FIG. 6A: A schematic of gene editing treatment to restore neutrophil differentiation. FIG. 6B: Flow cytometry analysis showing Healthy donor (HD-2) and Patient-derived (ESCN-2) HSCs that were edited ex-vivo only on the targeted mutated allele and differentiated into mature neutrophils. FIG. 6C: Quantification of data shown in FIG. 6B.

[0029] FIGS. 7A-7B: Linear representation of ELANE's five exons and four introns showing location of representative heterozygous mutations associated with SCN depicted as black inverted triangles (FIG. 7A). Based on Makaryan et al. FIG. 7B: Schematic of three identified SNPs (white inverted triangles), associated with the majority of ELANE mutations and a common cut site (gray inverted triangle), based on which three allele specific sgRNA guides and a constant guide were designed representing three mono-allelic excision strategies (I-III).

[0030] FIGS. 8A-8H: Allele specificity and excision efficiency of OMNI Variant 3795 nuclease compositions. FIG. 8A: Scheme depicting experimental workflow. HSCs from healthy donors and SCN patients were electroporated with RNPs or left non-treated followed by 3 days recovery in CD34⁺ expansion media. Cells were then subjected to differentiation by culturing for 7 days with IL-3, SCF, GM-CSF and G-CSF for proliferation and myeloid progenitor differentiation, followed by a 7-days culture in G-CSF for neutrophil differentiation. FIG. 8B: Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles at day 6 of differentiation in HSCs taken from either healthy donor (HD-V3) or SCN patient (SCN-P41) treated with RNP (ref) composition or left non-treated (NT), as measured by ddPCR. Average of each allele concentration was normalized to endogenous gene control RPP30 and STAT1 and presented relatively to non-treated cells. (n=3 groups of cells from HD-V3 healthy/SCN-P41 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. FIG. 8C: Bar graphs representing percentages of excision at days 6 and 14 of differentiation in HSCs taken from either healthy donor (HD-V3): non-treated (NT, black) or RNP (ref)-treated (gray) or SCN patient (SCN-P41): non-treated (NT, white) or RNP (ref)-treated (dark oblique lines), as measured by ddPCR. (n=3 groups of cells from HD-V3 healthy/SCN-P41 patient donors). Statistical significance is indicated as **P<0.01, ****P<0.0001. FIG. 8D: Bar graphs representing percentages of wild-type (black) and mutated (gray) alleles in cDNA taken from SCN-P41 patient HSCs that were either treated with RNP (ref) composition or left non-treated (NT), as measured by next-generation sequencing (NGS) targeting the mutation site. (n=3 groups of cells from SCN-P41

patient). Statistical significance is indicated as $***P<0.001$. FIG. 8E: Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles at day 6 of differentiation in HSCs taken from SCN patient (SCN-P55) treated with RNP (alt) composition or left non-treated (NT), as measured by ddPCR. Average of each allele concentration was normalized to endogenous gene control RPP30 and STAT1 and presented relatively to non-treated cells. (n=3 groups of cells from SCN-P55 patient donor). Statistical significance is indicated as $***P<0.0001$, ns=Not statistically significant. FIG. 8F: Bar graphs representing percentages of excision at days 6 and 14 of differentiation in HSCs taken from SCN patient (SCN-P55): non-treated (NT, white) or RNP (alt)-treated (dark oblique lines), as measured by ddPCR. (n=3 groups of cells from SCN-P55 patient donor). Statistical significance is indicated as $***P<0.001$, $****P<0.0001$. FIG. 8G: Bar graphs representing percentages of wild-type (black) and mutated (gray) alleles in cDNA taken from SCN-P55 patient HSCs that were either treated with RNP (alt) composition or left non-treated (NT), as measured by NGS targeting the mutation site. (n=3 groups of cells from SCN-P55 patient). Statistical significance is indicated as $***P<0.001$. FIG. 8H: Bar graphs representing ELANE mRNA levels in day 6 differentiated HSCs of SCN-P41 and SCN-P55 patients that were either not-treated (NT, Black) or RNP(ref)/RNP(alt)-treated, respectively (Excised, Gray). Data is presented relatively to the NT group. (n=3 groups of cells from SCN-P41 and SCN-P55 patients). Statistical significance is indicated as $*P<0.05$. Bars represent mean values with standard deviation. Bars represent mean values with standard deviation.

[0031] FIGS. 9A-9K: OMNI Variant 3795-facilitated editing boosts neutrophil differentiation and maturation in vitro. FIG. 9A: Representative FACS plots of non-treated (NT, left panel) and RNP(ref)-treated (right panel) healthy donor (HD-V3, upper panel) and SCN-P41 patient (lower panel) differentiated HSCs, analyzed for neutrophilic (CD66b+) and monocytic (CD14+/CD66b-) subsets. FIG. 9B: Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b+ cells) in healthy (HD-V3) and SCN patient (SCN-P41) differentiated HSCs that were non-treated (NT, black) or treated with RNP(ref) (gray). (n=3 groups of cells from HD-V3 healthy/SCN-P41 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9C: Quantitative analysis of respective FACS data for percentages of monocytes (CD14+/CD66b- cells) in healthy (HD-V3) and SCN patient (SCN-P41) differentiated HSCs that were non-treated (NT, black) or treated with RNP(ref) (gray). (n=3 groups of cells from HD-V3 healthy/SCN-P41 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9D: Quantification of percentages of Zymosan Green uptake by healthy (HD-V3) and SCN patient (SCN-P41) differentiated HSCs that were non-treated (NT, black) or treated with RNP(ref) (gray). ns=Not statistically significant. FIG. 9E: Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy non-treated (HD-V3, NT; square) or patient RNP(ref)-treated (SCN-P41, RNP(ref); triangle) HSCs compared to bacterial cells only control (e-coli, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as $*P<0.05$, $**P<0.01$. FIG. 9F: Representative FACS plots of non-treated healthy donor (HD-V4 NT, left panel), non-treated SCN patient (SCN-P55 NT, middle panel) and RNP(alt)-treated SCN patient (right panel) differentiated HSCs, analyzed for neutrophilic (CD66b+) and monocytic (CD14+/CD66b-) subsets. FIG. 9G: Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b+ cells) in differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9H: Quantitative analysis of respective FACS data for percentages of monocytes (CD14+/CD66b- cells) in differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9I: Diff-Quik staining of P55 SCN patient-derived differentiated HSCs treated with RNP(alt) or electroporated without a nuclease composition (SCN-P55 Mock). Microphotographs were taken on LEITZ LABORLUX S polarizing light microscope at 400x magnification using Nikon DSLR digital camera. FIG. 9J: Quantification of percentages of Zymosan Green uptake by differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $*P<0.05$. FIG. 9K: Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy non-treated (HD-V4, NT; square) HSCs and patient RNP(alt)-treated (SCN-P55, RNP(alt); triangle) HSCs compared to bacterial cells only control (*E. coli*, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as $*P<0.05$, $***P<0.001$. Bars represent mean values with standard deviation.

cated as $*P<0.05$, $**P<0.01$. FIG. 9F: Representative FACS plots of non-treated healthy donor (HD-V4 NT, left panel), non-treated SCN patient (SCN-P55 NT, middle panel) and RNP(alt)-treated SCN patient (right panel) differentiated HSCs, analyzed for neutrophilic (CD66b+) and monocytic (CD14+/CD66b-) subsets. FIG. 9G: Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b+ cells) in differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9H: Quantitative analysis of respective FACS data for percentages of monocytes (CD14+/CD66b- cells) in differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $****P<0.0001$, ns=Not statistically significant. FIG. 9I: Diff-Quik staining of P55 SCN patient-derived differentiated HSCs treated with RNP(alt) or electroporated without a nuclease composition (SCN-P55 Mock). Microphotographs were taken on LEITZ LABORLUX S polarizing light microscope at 400x magnification using Nikon DSLR digital camera. FIG. 9J: Quantification of percentages of Zymosan Green uptake by differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as $*P<0.05$. FIG. 9K: Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy non-treated (HD-V4, NT; square) HSCs and patient RNP(alt)-treated (SCN-P55, RNP(alt); triangle) HSCs compared to bacterial cells only control (*E. coli*, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as $*P<0.05$, $***P<0.001$. Bars represent mean values with standard deviation.

[0032] FIGS. 10A-10B: Heterozygosity frequency and coverage of patient and healthy populations by the three SNPs. FIG. 10A: Heterozygosity frequency of each of the three chosen SNPs in the healthy (left) or patient (right) population. Heterozygosity frequency was similar among the healthy and patient populations in each of the three SNPs. rs10414837, p value=0.126, Odds ratio=0.653; rs3761005, p value=0.9615, Odds ratio=1.014; rs1683564, p value=0.9475, Odds ratio=1.019, all analyzed by Chi-square. FIG. 10B: Pie chart presenting percentage of the population being heterozygous for at least one of the three chosen SNPs (gray) in the healthy (left) or patient (right) population. Similar coverage of the healthy and patient populations by the three SNPs (more than 75%, p value=0.1285, Odds ratio=0.56, Chi-square).

[0033] FIG. 11: Mutation-SNP linkage determination in SCN-P41 and SCN-P55 patients. Electropherograms of sequencing analyses of the mutation site and the rs1683564 SNP. SCN-P41 patient harbors a mutation on the same allele as the reference form of the SNP (C, cytosine), whereas SCN-P55 patient harbors a mutation on the same allele as the alternative form of the SNP (A, adenosine).

[0034] FIG. 12: Same editing outcomes with RNP(ref) and RNP(alt) compositions. The ELANE gene is cleaved in two locations: 1) intron 4, a biallelic site guided by sgRNA (constant) guide and 2) a heterozygous SNP site, rs1683564, a single allelic site guided by either sgRNA(ref) or sgRNA(alt) depending on the linkage to the mutation site. If the mutation is located at the allele harboring the reference form of the SNP (C, cytosine), RNP(ref) composition, including a nuclease, sgRNA(ref) and sgRNA(constant), is chosen and a section including the mutation is cleaved from the reference allele (Patient A, upper panel). If the mutation is located at the allele harboring the alternative form of the SNP (A, adenosine), RNP(alt) composition, including a nuclease, sgRNA(alt) and sgRNA(constant), is chosen and a section including the mutation is cleaved from the alternative allele (Patient B, lower panel). Illustration created with BioRender.com.

[0035] FIGS. 13A-13C: Inversion events following excision. FIG. 13A: Schematic of detection of inversion events: Specific primers were designed to amplify inverted variations of the excised fragment. EvaGreen dye, a fluorescent DNA-binding dye that binds dsDNA, was used in a ddPCR assay to measure all inversion events. Illustration created with BioRender.com. FIG. 13B: Quantification of total inversion events measured by EvaGreen-based ddPCR assay in unedited (black) and RNP(ref)-treated (gray) HD-V3 healthy donor and SCN-P41 patient derived differentiated HSCs. Statistical significance is indicated as ****P<0.0001. FIG. 13C: Quantification of total inversion events measured by EvaGreen-based ddPCR assay in unedited HD-V4 healthy donor and SCN-P55 patient-derived differentiated HSCs (black) and in RNP(alt)-treated (gray) SCN-P55 patient derived differentiated HSCs. Statistical significance is indicated as ****P<0.0001. Bars represent mean values with standard deviation. (n=3-4 groups of cells from HD-V3 or HD-V4 healthy/SCN-P41 or SCN-P55 patient donors).

[0036] FIGS. 14A-14B: Excision levels and allele specificity in additional healthy donors. FIG. 14A: Bar graphs representing percentages of excision in HSCs taken from healthy donors that were either non-treated (NT, black), RNP (ref)-treated (gray) or RNP(alt)-treated (white) as measured by ddPCR. (n=4 groups of cells from 2 healthy donors in each group). Statistical significance is indicated as ****P<0.0001. FIG. 14B: Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles in HSCs taken from healthy donors that were non-treated (NT), treated with RNP(ref) or RNP(alt), as measured by ddPCR. Average of each allele concentration was normalized to endogenous gene control RPP30 and STAT1 and presented relatively to non-treated cells. (n=4 groups of cells from two (2) healthy donors in each group). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. Bars represent mean values with standard deviation.

[0037] FIGS. 15A-15C: Excision levels in Long Term HSC population. FIG. 15A: Representative FACS plots of healthy donor derived CD34⁺ HSCs prior to sorting (Total, left panel) and following sorting to CD90⁻ (middle panel) and CD90⁺ (right panel) populations. (FIG. 15B and FIG. 15C) Bar graphs representing percentages of excision from HSCs taken from two healthy donors (MLP1; FIG. 15B—heterozygous to the alternative form of the SNP and MLP2, FIG. 15C—homozygous to the alternative form of the SNP) prior to sorting (Total, black) and following sorting to

CD90⁺ (light gray) and CD90⁻ (dark gray) populations as measured by ddPCR. Non-treated HSCs prior to sorting served as control (NT) (n=2 groups of cells from each healthy donor in each group). Bars represent mean values with standard deviation.

[0038] FIGS. 16A-16D: Differentiation into CD11b⁺/CD15⁺ neutrophils. FIG. 16A: Representative FACS plots of non-treated (NT, left panel) and RNP(ref)-treated (right panel) healthy donor (HD-V3, upper panel) and SCN-P41 patient (lower panel) differentiated HSCs, analyzed for neutrophilic (CD11b⁺/CD15⁺) subset. FIG. 16B: Quantitative analysis of respective FACS data for percentages of neutrophils (CD11b⁺/CD15⁺ cells) in healthy (HD-V3) and SCN patient (SCN-P41) differentiated HSCs that were non-treated (NT, black) or treated with RNP(ref) (gray). (n=3 groups of cells from HD-V3 healthy/SCN-P41 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. FIG. 16C: Representative FACS plots of non-treated healthy donor (HD-V4 NT, left panel), non-treated SCN patient (SCN-P55 NT, middle panel) and RNP(alt)-treated SCN patient (right panel) differentiated HSCs, analyzed for neutrophilic (CD11b⁺/CD15⁺) subset. FIG. 16D: Quantitative analysis of respective FACS data for percentages of neutrophils (CD11b⁺/CD15⁺ cells) in differentiated HSCs from non-treated healthy donor (HD-V4, NT; black) and SCN patient (SCN-P55) either non-treated (NT, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V4 healthy/SCN-P55 patient donors). Statistical significance is indicated as ****P<0.0001. Bars represent mean values with standard deviation.

[0039] FIGS. 17A-17B. OMNI-50 variants editing activity of ref and alt allele of g62 in LCL cells. Editing activity was determined by NGS analysis. Displayed average and standard deviation of three replicates.

[0040] FIGS. 18A-18B: OMNI-50 variants editing activity of g62 off-targets. Two different off-targets were tested: g62 OT1 and g62 OT2. Displayed average and standard deviation of three replicates.

[0041] FIGS. 19A-19B: Specificity of probes and guides. FIG. 19A: The binding of each probe (FAM, black; HEX, gray) to DNA extracted from healthy donor (HD) cells that are homozygous to either the reference or alternative forms of the SNP, was measured by ddPCR. Bar graphs representing concentration of positive events. (n=3 groups of cells). Statistical significance is indicated as ****P<0.0001. FIG. 19B: Bar graphs representing percentages of excision in a healthy donor HSCs that are homozygous to the reference form of the rs1683564 SNP and were either not-treated (NT), treated with RNP(alt) or RNP(ref) as measured by ddPCR. (n=3 groups of cells). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. Bars represent mean values with standard deviation.

[0042] FIGS. 20A-20D: No detected off targets following editing of OMNI Variant 3795 nuclease and each of the sgRNAs. (FIGS. 20A-C) An unbiased survey (GUIDE-seq) of whole-genome off-target cleavage using OMNI Variant 3795 nuclease and each of the constant guide (FIG. 20A, SgRNA(constant)), reference guide (FIG. 20B, SgRNA(ref)) and alternative guide (FIG. 20C, SgRNA(alt)), showing all reads are of the target sequence and no off-targets detected (4 mismatches). Note, analysis was done in U2OS cells that are homozygous to the reference form of rs1683564 SNP. Since OMNI Variant 3795 nuclease is

highly allele discriminatory, when using sgRNA(alt) there is only minor on-target editing of the reference allele (13 reads of the reference cytosine genotype) and no detectable off targets. (FIG. 20D) A table summarizing the results of an in-silico off-target analysis for constant, alternative and reference guides depicting a few potential off-targets. None of these off targets were validated by rhAmpSeq analysis performed on HSCs derived from SCN-P41 and SCN-P55 patients edited with RNP(ref) and RNP(alt), respectively, see two right columns. rhAmpSeq validation threshold was set to editing $\geq 0.2\%$.

[0043] FIGS. 21A-21J. Excision using RNP(ref) in SCN-P42 and HD-V5. (FIG. 21A) Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles at day 6 of differentiation in HSCs taken from either healthy donor (HD-V5) or SCN patient (SCN-P42) treated with RNP (ref) composition or electroporated without a nuclease composition (Mock), as measured by ddPCR. (n=3 groups of cells from HD-V5 healthy/SCN-P42 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 21B) Bar graphs representing percentages of excision at days 6 and 14 of differentiation in HSCs taken from either healthy donor (HD-V5): Mock-treated (Mock, black) or RNP (ref)-treated (gray), or SCN patient (SCN-P42): Mock-treated (Mock, white) or RNP (ref)-treated (dark oblique lines), as measured by ddPCR. (n=3 groups of cells from HD-V5 healthy/SCN-P42 patient donors). Statistical significance is indicated as ****P<0.0001. (FIG. 21C) Bar graphs representing ELANE mRNA levels in day 6 differentiated HSCs of SCN-P42 patient that were either Mock-treated (Black) or RNP(ref)-treated (Gray). Data is presented relatively to the mock group. (n=3 groups of cells from SCN-P42 patient). Statistical significance is indicated as **P<0.01. (FIG. 21D) Bar graphs representing percentages of wild-type (black) and mutated (gray) alleles in cDNA taken from SCN-P42 patient HSCs that were either RNP (ref)-treated or Mock-treated (Mock), as measured by NGS targeting the mutation site. (n=3 groups of cells from SCN-P42 patient). Statistical significance is indicated as *P<0.05. (FIG. 21E) Representative FACS plots of mock-treated (Mock, left panel) and RNP(ref)-treated (right panel) healthy donor (HD-V5, upper panel) and SCN-P42 patient (lower panel) differentiated HSCs, analyzed for neutrophilic (CD66b+) and monocytic (CD14+/CD66b-) subsets. (FIG. 21F) Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b+ cells) in healthy (HD-V5) and SCN patient (SCN-P42) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(ref) (gray). (n=3 groups of cells from HD-V5 healthy/SCN-P42 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 21G) Quantitative analysis of respective FACS data for percentages of monocytes (CD14+/CD66b- cells) in healthy (HD-V5) and SCN patient (SCN-P42) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(ref) (gray). (n=3 groups of cells from HD-V5 healthy/SCN-P42 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 21H) Diff-Quik staining of P42 SCN patient-derived differentiated HSCs treated with RNP(ref) or electroporated without a nuclease composition (SCN-P42 Mock). Microphotographs were taken on LEITZ LABORLUX S polarizing light microscope at 400x magnification using Nikon DSLR digital camera. (FIG. 21I)

Quantification of percentages of Zymosan Green uptake by healthy (HD-V5) and SCN patient (SCN-P42) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(ref) (gray). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 21J) Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy mock-treated (HD-V5, Mock; white square), patient mock-treated (SCN-P42 Mock; crossed circle) or patient RNP(ref)-treated (SCN-P42 RNP(ref); triangle) HSCs compared to bacterial cells only control (e-coli, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as ****P<0.0001. Bars represent mean values with standard deviation.

[0044] FIGS. 22A-22I. Excision using RNP(alt) in SCN-P12 and HD-V1. (FIG. 22A) Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles at day 6 of differentiation in HSCs taken from either healthy donor (HD-V1) or SCN patient (SCN-P12) treated with RNP (alt) composition or electroporated without a nuclease composition (Mock), as measured by ddPCR. (n=3 groups of cells from HD-V1 healthy/SCN-P12 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 22B) Bar graphs representing percentages of excision at days 6 and 14 of differentiation in HSCs taken from either healthy donor (HD-V1): Mock-treated (Mock, black) or RNP (alt)-treated (gray), or SCN patient (SCN-P12): Mock-treated (Mock, white) or RNP (alt)-treated (dark oblique lines), as measured by ddPCR. (n=3 groups of cells from HD-V1 healthy/SCN-P12 patient donors). Statistical significance is indicated as ***P<0.001, ****P<0.0001. (FIG. 22C) Bar graphs representing ELANE mRNA levels in day 6 differentiated HSCs of SCN-P12 patient that were either Mock-treated (Black) or RNP(alt)-treated (Gray). Data is presented relatively to the mock group. (n=3 groups of cells from SCN-P12 patient). Statistical significance is indicated as *P<0.05. (FIG. 22D) Bar graphs representing percentages of wild-type (black) and mutated (gray) alleles in cDNA taken from SCN-P12 patient HSCs that were either RNP (alt)-treated or Mock-treated (Mock), as measured by NGS targeting the mutation site. (n=3 groups of cells from SCN-P12 patient). Statistical significance is indicated as ***P<0.001. (FIG. 22E) Representative FACS plots of mock-treated (Mock, left panel) and RNP(alt)-treated (right panel) healthy donor (HD-V1, upper panel) and SCN-P12 patient (lower panel) differentiated HSCs, analyzed for neutrophilic (CD66b+) and monocytic (CD14+/CD66b-) subsets. (FIG. 22F) Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b+ cells) in healthy (HD-V1) and SCN patient (SCN-P12) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V1 healthy/SCN-P12 patient donors). Statistical significance is indicated as **P<0.01, ***P<0.001. (FIG. 22G) Quantitative analysis of respective FACS data for percentages of monocytes (CD14+/CD66b- cells) in healthy (HD-V1) and SCN patient (SCN-P12) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP (alt) (gray). (n=3 groups of cells from HD-V1 healthy/SCN-P12 patient donors). Statistical significance is indicated as ****P<0.0001, ns=Not statistically significant. (FIG. 22H) Quantification of percentages of Zymosan Green uptake by

healthy (HD-V1) and SCN patient (SCN-P12) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(alt) (gray). Statistical significance is indicated as *** $P < 0.001$, ns=Not statistically significant. (FIG. 22I) Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy mock-treated (HD-V1, Mock; white square), patient mock-treated (SCN-12 Mock; crossed circle) or patient RNP(alt)-treated (SCN-P12 RNP(alt); triangle) HSCs compared to bacterial cells only control (e-coli, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$. Bars represent mean values with standard deviation.

[0045] FIGS. 23A-23I: Excision using RNP(alt) in SCN-P56 and HD-V3. (FIG. 23A) Bar graphs representing percentages of un-edited reference (black) and alternative (gray) alleles at day 6 of differentiation in HSCs taken from either healthy donor (HD-V3) or SCN patient (SCN-P56) treated with RNP (alt) composition or electroporated without a nuclease composition (Mock), as measured by ddPCR. (n=3 groups of cells from HD-V3 healthy/SCN-P56 patient donors). Statistical significance is indicated as *** $P < 0.0001$, ns=Not statistically significant. (FIG. 23B) Bar graphs representing percentages of excision at day 6 of differentiation in HSCs taken from either healthy donor (HD-V3): Mock-treated (Mock, black) or RNP (alt)-treated (gray), or SCN patient (SCN-P56): Mock-treated (Mock, white) or RNP (alt)-treated (dark oblique lines), as measured by ddPCR. (n=3 groups of cells from HD-V3 healthy/SCN-P56 patient donors). Statistical significance is indicated as *** $P < 0.001$. (FIG. 23C) Bar graphs representing percentages of wild-type (black) and mutated (gray) alleles in cDNA taken from SCN-P56 patient HSCs that were either RNP (alt)-treated or Mock-treated (Mock), as measured by NGS targeting the mutation site. (n=3 groups of cells from SCN-P56 patient). Statistical significance is indicated as *** $P < 0.0001$. (FIG. 23D) Representative FACS plots of mock-treated (Mock, left panel) and RNP(alt)-treated (right panel) healthy donor (HD-V3, upper panel) and SCN-P56 patient (lower panel) differentiated HSCs, analyzed for neutrophilic (CD66b⁺) and monocytic (CD14⁺/CD66b⁻) subsets. (FIG. 23E) Quantitative analysis of respective FACS data for percentages of neutrophils (CD66b⁺ cells) in healthy (HD-V3) and SCN patient (SCN-P56) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP(alt) (gray). (n=3 groups of cells from HD-V3 healthy/SCN-P56 patient donors). Statistical significance is indicated as *** $P < 0.0001$, ns=Not statistically significant. (FIG. 23F) Quantitative analysis of respective FACS data for percentages of monocytes (CD14⁺/CD66b⁻ cells) in healthy (HD-V3) and SCN patient (SCN-P56) differentiated HSCs that were mock-treated (Mock, black) or treated with RNP (alt) (gray). (n=3 groups of cells from HD-V3 healthy/SCN-P56 patient donors). Statistical significance is indicated as $P = 0.04$, *** $P < 0.0001$. (FIG. 23G) Diff-Quik staining of P56 SCN patient-derived differentiated HSCs treated with RNP(alt) or electroporated without a nuclease composition (SCN-P56 Mock). Microphotographs were taken on LEITZ LABORLUX S polarizing light microscope at 400× magnification using Nikon DSLR digital camera. (FIG. 23H) Quantification of percentages of Zymosan Green uptake by healthy (HD-V3) and SCN patient (SCN-P56) differentiated

HSCs that were mock-treated (Mock, black) or treated with RNP(alt) (gray). Statistical significance is indicated as *** $P < 0.0001$, ns=Not statistically significant. (FIG. 23I) Graph depicts real time change in light emission, relative light units (RLUs), from 200,000 Luciferase expressing bacterial cells incubated with differentiated neutrophils from healthy mock-treated (HD-V3, Mock; white square), patient mock-treated (SCN-56 Mock; crossed circle) or patient RNP(alt)-treated (SCN-P56 RNP(alt); triangle) HSCs compared to bacterial cells only control (e-coli, circle). Statistical significance for each one of the groups versus e-coli control at the last time point presented, when RLU levels reached plateau, is indicated as ** $P < 0.01$. Bars represent mean values with standard deviation.

DETAILED DESCRIPTION

[0046] The present disclosure provides an engineered OMNI-50 nuclease exhibiting increased specificity to a target site compared to the wild-type OMNI-50 nuclease (SEQ ID NO: 1). The wild-type OMNI-50 nuclease is disclosed in PCT International Application Publication No. WO/2020-030782, incorporated herein by reference. When the engineered OMNI-50 nuclease variant is active in a CRISPR endonuclease system, the CRISPR endonuclease system displays reduced off-target editing activity and maintained on-target editing activity relative to a CRISPR endonuclease system comprising the wild-type OMNI-50 nuclease. In some embodiments, the engineered OMNI-50 nuclease is an OMNI-50 nuclease variant comprising at least one amino acid substitution relative to the wild-type OMNI-50 nuclease. In some embodiments, the engineered OMNI-50 nuclease comprises multiple amino acid substitutions compared to wild-type OMNI-50 nuclease.

[0047] In some embodiments, an OMNI-50 nuclease variant is at least 80%, e.g., at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the amino acid sequence of SEQ ID NO: 1. As a non-limiting example, an OMNI-50 nuclease variant may have amino acid sequence differences at up to 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 11%, 12%, 13%, 14%, 15%, or 20% of its residues relative to SEQ ID NO:1. Such sequence differences may be revealed by a sequence alignment. An OMNI-50 variant nuclease may be generated by replacing at least one amino acid residue of an OMNI-50 wild-type nuclease with another amino acid residue e.g. with a conservative or non-conservative amino acid substitution, and/or by inserting or deleting an amino acid residue of the OMNI-50 wild-type nuclease. Any such mutations, including but not limited to substitutions, insertions, or deletions, in addition to any other mutations described herein, or with mutations in addition to the mutations described herein, may be used to generate an OMNI-50 variant nuclease from an OMNI-50 wild-type nuclease. In some embodiments, the OMNI-50 variant nuclease retains a desired activity of the parent wild-type OMNI-50 nuclease, e.g., the ability to interact with a guide RNA and target DNA and/or the activity of the nuclease (e.g. ability to cause a double-strand DNA break, a single-strand DNA break, or lack of any nuclease or nickase activity). In some embodiments, the variant retains the desired activity of the parent, e.g. nuclease activity, at a level greater than or equal to the level of activity of the parent. In some embodiments, the variant retains the desired activity of the parent at a level of at least 100%, 95%, 90%, 80%, 70%, 60%, 50%, 40%, or 30% the

level of activity of the parent. In some embodiments, the OMNI-50 variant nuclease displays reduced off-target effects relative to OMNI-50 wild-type nuclease.

[0048] In some embodiments, there is provided a variant of OMNI-50 nuclease protein comprising a sequence that is at least 80% identical to the amino acid sequence of wild-type OMNI-50 (SEQ ID NO: 1) and having at least one amino acid substitution. In some embodiments, the amino acid substitution comprises an amino acid residue replacement to a positive, negative, uncharged, hydrophilic, hydrophobic, polar, or non-polar amino acid. In some embodiments, the amino acid substitution is selected from replacement of an amino acid to any one of a different amino acid selected from the group consisting of R, K, H, D, E, S, T, N, Q, C, U, G, P, A, I, L, M, F, W, Y and V.

[0049] Positive amino acids include any amino acid having a positively charged R-group, e.g. lysine (K), arginine (R), or Histidine (H). Negative amino acids include any amino acid having a negatively charged R-group, e.g. aspartic acid (D) or glutamic acid (E). Uncharged amino acids or neutral amino acids include amino acids whose R-group does not normally carry a charge. Polar amino acids include any amino acid having a polar R-group, e.g. serine (S), threonine (T), tyrosine (Y), asparagine (N), or glutamine (Q). Non-polar amino acids include any amino acid having a non-polar R-group, e.g. glycine (G), alanine (A), valine (V), cysteine (C), proline (P), leucine (L), isoleucine (I), methionine (M), tryptophan (W), or phenylalanine (F).

[0050] Properties of an original variant protein having an original amino acid substitution at a given position may be extended to a different variant having a different amino acid substitution at the same position if the different amino acid substitution has an R-group with similar properties to the original amino acid substitution. For example, if a variant protein is shown to have higher specificity compared to a wild-type protein by substituting a glutamic acid (E) residue for a lysine (K) residue, it is reasonable to consider that a similar variant substituting the glutamic acid (E) residue for an arginine (R) residue will also display higher specificity since both lysine (K) and arginine (R) share similar properties (e.g. they both contain positively charged R-groups). Conversely, a variant having a substitution of the glutamic acid (E) residue to an aspartic acid (D) is less likely to display the higher specificity property because both glutamic acid (E) and aspartic acid (D) share the similar property of both containing a negatively charged R-group.

[0051] In some embodiments, a variant OMNI-50 nuclease protein contains an amino acid substitution in at least one of the following positions in the wild-type OMNI-50 protein sequence (SEQ ID NO:1): R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036. Each possibility represents a separate embodiment of the present disclosure. In some embodiments, there is provided a variant of OMNI-50 nuclease protein comprising a sequence that is at least 80% identical to the amino acid sequence of the wild-type OMNI-50 nuclease (SEQ ID NO: 1) and having at least one amino acid substitution in at least one of the following positions in the wild-type OMNI-50 protein sequence: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, K965, V981, and K1036. In some embodiments, the variant OMNI-50 nuclease protein comprises at least one of the following amino acid substitutions in the following positions in the wild-type OMNI-50 protein

sequence: R478I, Y545H, Q803V, L805I, R61A, Y437W, R788K, L844N, V981M, G606P, L690V, E695Q, R478T, A493G, K688E, L718C, K965V, and K1036V. Each possibility represents a separate embodiment of the present disclosure. In some embodiments, the substitution corresponds to the mutations listed in Table 1.

[0052] In some embodiments, the variant OMNI-50 nuclease protein comprises at least one amino acid substitution in the following positions in the wild-type OMNI-50 protein sequence: R478, Y545, Q803, and L805. In some embodiments, the variant OMNI-50 nuclease protein comprises amino acid substitutions in the following positions in the wild-type OMNI-50 protein sequence: R478, Y545, Q803, and L805. In some embodiments, the variant OMNI-50 nuclease protein comprises the following amino acid substitutions from the wild-type OMNI-50 protein sequence: R478I, Y545H, Q803V, and L805I.

[0053] In some embodiments, the variant OMNI-50 nuclease protein comprises at least one amino acid substitution in the following positions in the wild-type OMNI-50 protein sequence: R61, Y437, R788, L844, and V981. In some embodiments, the variant OMNI-50 nuclease protein comprises amino acid substitutions in the following positions in the wild-type OMNI-50 protein sequence: R61, Y437, R788, L844, and V981. In some embodiments, the variant OMNI-50 nuclease protein comprises the following amino acid substitutions from the wild-type OMNI-50 protein sequence: R61A, Y437W, R788K, L844N, and V981M.

[0054] In some embodiments, the variant OMNI-50 nuclease protein comprises at least one amino acid substitution in the following positions in the wild-type OMNI-50 protein sequence: G606, L690, and E695. In some embodiments, the variant OMNI-50 nuclease protein comprises amino acid substitutions in the following positions in the wild-type OMNI-50 protein sequence: G606, L690, and E695. In some embodiments, the variant OMNI-50 nuclease protein comprises the following amino acid substitutions from the wild-type OMNI-50 protein sequence: G606P, L690V, and E695Q.

[0055] In some embodiments, the variant OMNI-50 nuclease protein comprises at least one amino acid substitution in the following positions in the wild-type OMNI-50 protein sequence: R478, A493, K688, L718, K965, and K1036. In some embodiments, the variant OMNI-50 nuclease protein comprises amino acid substitutions in the following positions in the wild-type OMNI-50 protein sequence: R478, A493, K688, L718, K965V, and K1036. In some embodiments, the variant OMNI-50 nuclease protein comprises the following amino acid substitutions from the wild-type OMNI-50 protein sequence: R478T, A493G, K688E, L718C, K965V, and K1036V.

[0056] In some embodiments, the OMNI-50 variant nuclease further comprises one or more of a nuclear localization sequence (NLS), cell penetrating peptide sequence, and/or affinity tag. In an embodiment, the OMNI-50 variant nuclease comprises one or more nuclear localization sequences of sufficient strength to drive accumulation of a CRISPR complex comprising the CRISPR nuclease in a detectable amount in the nucleus of a eukaryotic cell.

[0057] In some embodiments, the OMNI-50 variant nuclease comprises amino acid substitutions selected from amino acid substitutions corresponding to the substitutions displayed relative to wild-type OMNI-50 in Table 1.

[0058] According to some embodiments, there is provided an isolated OMNI-50 variant nuclease protein comprising one or more substitutions or mutations relative to the wild-type OMNI-50 nuclease sequence, wherein the isolated variant OMNI-50 variant nuclease is active in a CRISPR system, wherein the CRISPR system displays reduced off-target editing activity and maintained on-target editing activity relative to a wild-type CRISPR system.

[0059] According to some embodiments, additional mutations to the OMNI-50 variant nuclease described herein may be implemented. Examples include, but are not limited to, mutations which alter the PAM recognition sequence, alter the nuclease activity of the enzyme, and truncations or removal of portions of the nuclease. According to some embodiments, the variant OMNI-50 variant nuclease may be encoded by any nucleic acid sequence which produces the desired amino acid sequence of the variant. For example, the nucleic acid sequence may be codon-optimized for a cell, such as a bacterial cell, plant cell, or mammalian cell.

[0060] In embodiments of the present invention, a CRISPR nuclease and a targeting molecule form a CRISPR complex that binds to a target DNA sequence to effect cleavage of the target DNA sequence. A CRISPR nuclease may form a CRISPR complex comprising the CRISPR nuclease and a single-guide RNA (sgRNA) molecule. Alternatively, a CRISPR nucleases may form a CRISPR complex comprising the CRISPR nuclease, an crRNA molecule, and a tracrRNA molecule.

[0061] According to some embodiments of the present invention, there is provided a method of gene editing having reduced off-target editing activity and/or increased on-target editing activity, comprising: contacting a target site locus with an active CRISPR endonuclease system having a variant OMNI-50 protein complexed with a suitable guide RNA or guide RNA complex, wherein the active CRISPR endonuclease system displays reduced off-target editing activity and maintained on-target editing activity relative to a wild-type OMNI-50 CRISPR system.

[0062] According to some embodiments, there is provided a non-naturally occurring OMNI-50 nuclease variant having a wild-type OMNI-50 protein sequence (SEQ ID NO: 1) comprising an amino acid substitution in at least one of the following positions: R61, Y437, R478, A493, Y545, G606, K688, L690, E695, L718, R788, Q803, L805, L844, V981, K965, and K1036.

[0063] In some embodiments, the amino acid substitution at position R478 and/or position Y545, preferably at both position R478 and position Y545.

[0064] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478 and Y545.

[0065] In some embodiments, the amino acid substitution at position R478 is any one of the following substitutions: R478D, R478E, R478S, R478T, R478N, R478Q, R478G, R478P, R478C, R478A, R478V, R478I, R478L, R478M, R478F, R478Y, or R478W, preferably R478V, R478H, R478L, R478M, R478P, R478F, R478W, R478Y, R478S, R478C, R478T, R478N, or R478Q.

[0066] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is an amino acid having a negatively charged R-group or an R-group lacking a charge.

[0067] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is a polar amino acid or non-polar amino acid.

[0068] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is a non-polar amino acid.

[0069] In some embodiments, the amino acid substitution at position Y545 is any one of the following substitutions: Y545D, Y545E, Y545S, Y545T, Y545N, Y545Q, Y545G, Y545P, Y545C, Y545A, Y545V, Y545I, Y545L, Y545M, Y545F, Y545R, Y545K, Y545H, or Y545W, preferably Y545W, Y545F, Y545H, Y545V, or Y545G.

[0070] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is an amino acid having a negatively charged R-group or a positively charged R-group.

[0071] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is a non-polar amino acid.

[0072] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine lacks a phenyl ring or is phenylalanine.

[0073] In some embodiments, the amino acid substitutions are R478I and Y545H.

[0074] In some embodiments, the amino acid substitution is any one of the following substitutions: R478I, Y545H, Q803V, L805I, R61A, Y437W, R788K, L844N, V981M, G606P, L690V, E695Q, R478T, A493G, K688E, L718C, K965V, and K1036V.

[0075] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478, Y545, Q803, and L805.

[0076] In some embodiments, the amino acid substitutions are R478I, Y545H, Q803V, and L805I.

[0077] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R61, Y437, R788, L844, and V981.

[0078] In some embodiments, the amino acid substitutions are R61A, Y437W, R788K, L844N, and V981M.

[0079] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions G606, L690, and E695.

[0080] In some embodiments, the amino acid substitutions are G606P, L690V, and E695Q.

[0081] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478, A493, K688, L718, K965, and K1036.

[0082] In some embodiments, the amino acid substitutions are R478T, A493G, K688E, L718C, K965V, and K1036V.

[0083] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at position Q803.

[0084] In some embodiments, the amino acid substitution is Q803V.

[0085] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at position Y545.

[0086] In some embodiments, the amino acid substitution is Y545H.

[0087] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at position L805.

[0088] In some embodiments, the amino acid substitution is L805I.

[0089] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478, Y545, and Q803.

[0090] In some embodiments, the amino acid substitutions are R478I, Y545H, and Q803V.

[0091] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478, Y545, and L805.

[0092] In some embodiments, the amino acid substitutions are R478I, Y545H, and L805I.

[0093] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478, Q803, and L805.

[0094] In some embodiments, the amino acid substitutions are R478I, Q803V, and L805I.

[0095] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions Y545, Q803, and L805.

[0096] In some embodiments, the amino acid substitutions are R478I, Y545H, and L805I.

[0097] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at position R478 and/or position Y545.

[0098] In some embodiments, the amino acid substitution at position R478 is any one of the following substitutions: R478D, R478E, R478S, R478T, R478N, R478Q, R478G, R478P, R478C, R478A, R478V, R478I, R478L, R478M, R478F, R478Y, or R478W.

[0099] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is an amino acid having a negatively charged R-group or an R-group lacking a charge.

[0100] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is a polar amino acid or non-polar amino acid.

[0101] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is a non-polar amino acid.

[0102] In some embodiments, the amino acid substitution is at position R478 and the amino acid substituted for arginine is selected from a large hydrophobic amino acid (e.g., leucine, methionine, proline, valine), an aromatic amino acid (e.g., histidine, phenylalanine, tryptophan and tyrosine), and a polar uncharged amino acid (e.g., serine, cysteine, threonine, asparagine, and glutamine).

[0103] In some embodiments, the amino acid substitution at position Y545 is any one of the following substitutions: Y545D, Y545E, Y545S, Y545T, Y545N, Y545Q, Y545G, Y545P, Y545C, Y545A, Y545V, Y545I, Y545L, Y545M, Y545F, Y545R, Y545K, Y545H, or Y545W.

[0104] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is an amino acid having a negatively charged R-group or a positively charged R-group.

[0105] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is an amino acid having a positively charged R-group. Non-limiting examples of positively charged amino acids include histidine, lysine and arginine.

[0106] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is a non-polar amino acid. Non-limiting examples of non-polar amino acids include glycine, alanine, valine, proline, leucine, isoleucine, methionine, tryptophan and phenylalanine.

[0107] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine lacks a phenyl ring or is phenylalanine.

[0108] In some embodiments, the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine includes a phenyl ring (e.g., phenylalanine).

[0109] In some embodiments, the OMNI-50 nuclease variant comprises an amino acid substitution at each of positions R478 and Y545.

[0110] In some embodiments, the amino acid substitutions are R478I and Y545H. However, others amino acid substitutions at the R478 and Y545 positions are also contemplated. As a non-limiting example, amino acids having R-groups with similar non-polar to properties to isoleucine, such as alanine (A), valine (V), and leucine (L), are contemplated substitutions at the R478 position. In another non-limiting example, amino acids having R-groups with similar non-polar to properties to isoleucine, such as methionine (M), phenylalanine (F), tyrosine (Y), and tryptophan (W) are contemplated substitutions at the R478 position.

[0111] In some embodiments, the OMNI-50 nuclease variant has an amino acid sequence selected from the group consisting of SEQ ID NOs: 2-5, 63-70, and 89-97.

[0112] In some embodiments, the OMNI-50 nuclease variant has at least 80% sequence identity to the wild-type OMNI-50 protein sequence (SEQ ID NO:1).

[0113] In some embodiments, the OMNI-50 nuclease variant further comprises a nuclear localization sequence (NLS).

[0114] In some embodiments, the OMNI-50 nuclease variant exhibits increased specificity toward a DNA target site when complexed with a guide RNA molecule that targets variant to the said DNA target site relative to a wild-type OMNI-50 nuclease complexed with the guide RNA molecule.

[0115] According to some embodiments of the present invention, there is provided a CRISPR system comprising any one of the OMNI-50 nuclease variants described herein complexed with a guide RNA molecule that targets a DNA target site, wherein the CRISPR system displays reduced off-target editing activity relative to a wild-type CRISPR system comprising a wild-type OMNI-50 nuclease protein and the guide RNA molecule.

[0116] According to some embodiments of the present invention, there is provided a method for gene editing having reduced off-target editing activity, comprising contacting a DNA target site with an active CRISPR system comprising any one of the OMNI-50 nuclease variant proteins described herein.

[0117] In some embodiments, the active CRISPR system displays reduced off-target editing activity relative to a wild-type CRISPR system comprising a wild-type OMNI-50 nuclease protein.

[0118] In some embodiments, the gene editing occurs in a eukaryotic cell or prokaryotic cell.

[0119] In some embodiments, the eukaryotic cell is a plant cell or mammalian cell.

[0120] In some embodiments, the mammalian cell is a human cell.

[0121] In some embodiments, the DNA target site is located within or in proximity to a pathogenic allele of a gene.

[0122] In some embodiments, the DNA target site is located in a gene selected from the group consisting of ELANE, CXCR4, EMX, RyR2, KNCQ1, KCNH2, SCN5a,

GBA1, GBA2, Rhodopsin, GUCY2D, IMPDH1, FGA, BEST1, PRPH2, KRT5, KRT14, ApoA1, STAT3, STAT1, ADA2, RPS19, SBDS, GATA2, RPE65, LDLR, ANGPTL3, B2M, TRAC, TCF4, TGFB1, PAX6, C3, LRRK2, SARMI, SAMD9, SAMD9L, HAVCR2, CD3E, APLP2, CISH, TIGIT, TNNT2, TNN, MYH7, and HLA-E.

[0123] In some embodiments, the DNA target is repaired with an exogenous donor molecule.

[0124] In some embodiments, the off-target editing activity is reduced by at least 2-fold, 10-fold, 10^2 -fold, 10^3 -fold, 10^4 -fold, 10^5 -fold, or 10^6 -fold.

[0125] According to some embodiments of the present invention, there is provided a modified cell obtained by any one of the methods described herein.

[0126] In some embodiments, the cell is capable of engraftment.

[0127] In some embodiments, the cell is capable of giving rise to progeny cells after engraftment.

[0128] In some embodiments, the cell is capable of giving rise to progeny cells after an autologous engraftment.

[0129] In some embodiments, the cell is capable of giving rise to progeny cells for at least 12 months or at least 24 months after engraftment.

[0130] In some embodiments, the cell is selected from the group consisting of a hematopoietic stem cell, a progenitor cell, a CD34+ hematopoietic stem cell, a bone marrow cell, and a peripheral mononucleated cell.

[0131] According to some embodiments of the present invention, there is provided a composition comprising any one of the modified cells described herein and a pharmaceutically acceptable carrier. According to some embodiments of the present invention, there is provided an in vitro or ex vivo method of preparing the composition, comprising mixing the cells with the pharmaceutically acceptable carrier.

[0132] According to some embodiments of the present invention, there is provided a polynucleotide molecule encoding any one of the OMNI-50 variant proteins described herein.

Delivery

[0133] The OMNI-50 variant compositions described herein may be delivered as a protein, DNA molecules, RNA molecules, Ribonucleoproteins (RNP), nucleic acid vectors, or any combination thereof. In some embodiments, the RNA molecule comprises a chemical modification. Non-limiting examples of suitable chemical modifications include 2'-O-methyl (M), 2'-O-methyl, 3'-phosphorothioate (MS) or 2'-O-methyl, 3'-thioPACE (MSP), pseudouridine, and 1-methyl pseudo-uridine. Each possibility represents a separate embodiment of the present invention.

[0134] The OMNI-50 variants and/or polynucleotides encoding same described herein, and/or additional molecules, such as a single-guide RNA molecule, crRNA molecule, tracrRNA molecules or a nucleotide molecule that encodes any one of them, may be delivered to a target cell by any suitable means. The target cell may be any type of cell e.g., eukaryotic or prokaryotic, in any environment e.g., isolated or not, maintained in culture, in vitro, ex vivo, in vivo or in planta. A target site in a target cell may be within the nucleus of the cell.

[0135] The compositions described herein may be introduced into a cell as part of a vector molecule having additional sequences such as, for example, replication ori-

gins, promoters and genes encoding antibiotic resistance. Moreover, compositions may be introduced into a cell as naked nucleic acids or proteins, as nucleic acids or proteins complexed with or packaged within an agent such as a liposome, exosome, or poloxamer, or can be delivered by recombinant viruses (e.g., adenovirus, AAV, herpesvirus, retrovirus, lentivirus and integrase defective lentivirus (IDLV)) or virus-like particles. As non-limiting examples, the composition may be packaged into an adeno-associated virus (AAV), or into a lentivirus, such as a non-integrating lentivirus or a lentivirus lacking reverse transcription capability. Additional non-limiting examples include packaging the composition into liposomes, extracellular vesicles, or exosomes, which may be pseudotyped with vesicular stomatitis glycoprotein (VSVG) or conjugated to a cell-penetrating peptide, an antibody, a targeting moiety, or any combination thereof.

[0136] In some embodiments, the composition to be delivered includes mRNA of the nuclease and RNA of the guide. In some embodiments, the composition to be delivered includes mRNA of the nuclease, RNA of the guide and a donor template. In some embodiments, the composition to be delivered includes the CRISPR nuclease and guide RNA. In some embodiments, the composition to be delivered includes the CRISPR nuclease, guide RNA and a donor template for gene editing via, for example, homology directed repair. Optionally the lentivirus includes mRNA of the nuclease and a guide RNA molecule, e.g. a single-guide RNA molecule or crRNA molecule, which is used to target the nuclease to a target site. In some embodiments, the composition delivered to a cell includes mRNA of the nuclease, a guide RNA molecule and a donor template molecule. Optionally, the lentivirus includes the nuclease protein variant and a guide RNA molecule. Optionally, the composition delivered to a cell includes the nuclease protein variant, a guide RNA molecule and/or donor template for homology directed repair. Optionally, the composition delivered to a cell includes mRNA of the nuclease variant, a DNA-targeting crRNA molecule, and a tracrRNA molecule. the composition delivered to a cell includes mRNA of the nuclease variant, DNA-targeting crRNA molecule, and a tracrRNA molecule, and a donor template molecule. the composition delivered to a cell includes the nuclease protein variant, DNA-targeting crRNA molecule, and a tracrRNA molecule. Optionally, the composition delivered to a cell includes the nuclease protein variant, DNA-targeting crRNA molecule, and a tracrRNA molecule, and DNA donor template molecule for homology directed repair.

[0137] Any suitable viral vector system may be used to deliver such compositions. Conventional viral and non-viral based gene transfer methods can be used to introduce nucleic acids and/or OMNI-50 variant protein in cells (e.g., mammalian cells, plant cells, etc.) and target tissues. Such methods can also be used to administer nucleic acids encoding and/or OMNI-50 variant protein to cells in vitro. In certain embodiments, nucleic acids and/or a OMNI-50 variant protein are administered for in vivo or ex vivo gene therapy uses. Non-viral vector delivery systems include naked nucleic acid, and nucleic acid complexed with a delivery vehicle such as a liposome or poloxamer. For a review of gene therapy procedures, see Anderson, *Science* 256:808-813 (1992); Nabel & Felgner, *TIBTECH* 11:211-217 (1993); Mitani & Caskey, *TIBTECH* 11:162-166 (1993); Dillon, *TIBTECH* 11:167-175 (1993); Miller, *Nature* 357:455-460 (1992); Van Brunt, *Biotechnology*

6(10):1149-1154 (1988); Vigne, Restorative Neurology and Neuroscience 8:35-36 (1995); Kremer & Perricaudet, British Medical Bulletin 51(1):31-44 (1995); Haddada et al., in Current Topics in Microbiology and Immunology Doerfler and Bohm (eds.) (1995); and Yu et al., Gene Therapy 1:13-26 (1994).

[0138] Methods of non-viral delivery of nucleic acids and/or proteins include electroporation, lipofection, micro-injection, biolistics, particle gun acceleration, virosomes, virus-like particles, exosomes, liposomes, immunoliposomes, polycation or lipid:nucleic acid conjugates, artificial virions, and agent-enhanced uptake of nucleic acids or can be delivered to plant cells by bacteria or viruses (e.g., *Agrobacterium*, *Rhizobium* sp. NGR234, *Sinorhizobium meliloti*, *Mesorhizobium loti*, tobacco mosaic virus, potato virus X, cauliflower mosaic virus and cassava vein mosaic virus. See, e.g., Chung et al. Trends Plant Sci. (2006). Sonoporation using, e.g., the Sonitron 2000 system (Rich-Mar) can also be used for delivery of nucleic acids. Cationic-lipid mediated delivery of proteins and/or nucleic acids is also contemplated as an in vivo or in vitro delivery method. See Zuris et al., Nat. Biotechnol. (2015), Coelho et al., N. Engl. J. Med. (2013); Judge et al., Mol. Ther. (2006); and Basha et al., Mol. Ther. (2011).

[0139] Non-viral vectors, such as transposon-based systems e.g. recombinant Sleeping Beauty transposon systems or recombinant PiggyBac transposon systems, may also be delivered to a target cell and utilized for transposition of a polynucleotide sequence of a molecule of the composition or a polynucleotide sequence encoding a molecule of the composition in the target cell.

[0140] Additional exemplary nucleic acid delivery systems include those provided by Amaxa® Biosystems (Cologne, Germany), Maxcyte, Inc. (Rockville, Md.), BTX Molecular Delivery Systems (Holliston, Mass.) and Copernicus Therapeutics Inc., (see for example U.S. Pat. No. 6,008,336). Lipofection is described in e.g., U.S. Pat. Nos. 5,049,386, 4,946,787; and 4,897,355) and lipofection reagents are sold commercially (e.g., Transfectam™, Lipofectin™ and Lipofectamine™ RNAiMAX). Cationic and neutral lipids that are suitable for efficient receptor-recognition lipofection of polynucleotides include those disclosed in PCT International Publication Nos. WO/1991/017424 and WO/1991/016024. Delivery can be to cells (ex vivo administration) or target tissues (in vivo administration).

[0141] The preparation of lipid:nucleic acid complexes, including targeted liposomes such as immunolipid complexes, is well known to one of skill in the art (see, e.g., Crystal, Science 270:404-410 (1995); Blaese et al., Cancer Gene Ther. 2:291-297 (1995); Behr et al., Bioconjugate Chem. 5:382-389 (1994); Remy et al., Bioconjugate Chem. 5:647-654 (1994); Gao et al., Gene Therapy 2:710-722 (1995); Ahmad et al., Cancer Res. 52:4817-4820 (1992); U.S. Pat. Nos. 4,186,183, 4,217,344, 4,235,871, 4,261,975, 4,485,054, 4,501,728, 4,774,085, 4,837,028, and 4,946,787).

[0142] Additional methods of delivery include the use of packaging the nucleic acids to be delivered into EnGeneIC delivery vehicles (EDVs). These EDVs are specifically delivered to target tissues using bispecific antibodies where one arm of the antibody has specificity for the target tissue and the other has specificity for the EDV. The antibody brings the EDVs to the target cell surface and then the EDV

is brought into the cell by endocytosis. Once in the cell, the contents are released (see MacDiamid et al (2009) Nature Biotechnology 27(7) p. 643).

[0143] The use of RNA or DNA viral based systems for the delivery of nucleic acids take advantage of highly evolved processes for targeting a virus to specific cells in the body and trafficking the viral payload to the nucleus. Viral vectors can be administered directly to patients (in vivo) or they can be used to treat cells in vitro and the modified cells are administered to patients (ex vivo). Conventional viral based systems for the delivery of nucleic acids include, but are not limited to, retroviral, lentivirus, adenoviral, adeno-associated, vaccinia and herpes simplex virus vectors for gene transfer. Integration in the host genome is possible with the retrovirus, lentivirus, and adeno-associated virus gene transfer methods, often resulting in long term expression of the inserted transgene. Additionally, high transduction efficiencies have been observed in many different cell types and target tissues. A OMNI-50 variant or a nucleic acid expressing the variant, as well as any associated nucleic acids, may be delivered by a non-integrating lentivirus. Optionally, RNA delivery with lentivirus is utilized. Optionally, the lentivirus includes mRNA of the nuclease and a guide RNA molecule, e.g. a single-guide RNA molecule or crRNA molecule, which is used to target the nuclease to a target site. Optionally the lentivirus includes mRNA of the nuclease, guide RNA molecule and a donor template molecule. Optionally, the lentivirus includes the nuclease protein variant and a guide RNA molecule. Optionally, the lentivirus includes the nuclease protein variant, a guide RNA molecule and/or donor template molecule for homology directed repair. Optionally, the lentivirus includes mRNA of the nuclease variant, a DNA-targeting crRNA molecule, and a tracrRNA molecule. Optionally the lentivirus includes mRNA of the nuclease variant, DNA-targeting crRNA molecule, and a tracrRNA molecule, and a donor template molecule. Optionally, the lentivirus includes the nuclease protein variant, DNA-targeting crRNA molecule, and a tracrRNA molecule. Optionally, the lentivirus includes the nuclease protein variant, DNA-targeting crRNA molecule, and a tracrRNA molecule, and DNA donor template molecule for homology directed repair.

[0144] As mentioned above, the compositions described herein may be delivered to a target cell using a non-integrating lentiviral particle method, e.g. a LentiFlash® system. Such a method may be used to deliver mRNA or other types of RNAs into the target cell, such that delivery of the RNAs to the target cell results in assembly of the compositions described herein inside of the target cell. See also PCT International Publication Nos. WO2013/014537, WO2014/016690, WO2016185125, WO2017194902, and WO2017194903.

[0145] The tropism of a retrovirus can be altered by incorporating foreign envelope proteins, expanding the potential target population of target cells. Lentiviral vectors are retroviral vectors capable of transducing or infecting non-dividing cells and typically produce high viral titers. Selection of a retroviral gene transfer system depends on the target tissue. Retroviral vectors are comprised of cis-acting long terminal repeats with packaging capacity for up to 6-10 kb of foreign sequence. The minimum cis-acting LTRs are sufficient for replication and packaging of the vectors, which are then used to integrate the therapeutic gene into the target cell to provide permanent transgene expression. Widely used

retroviral vectors include those based upon murine leukemia virus (MuLV), gibbon ape leukemia virus (GaLV), Simian Immunodeficiency virus (SIV), human immunodeficiency virus (HIV), and combinations thereof (see, e.g., Buchscher Panganiban, J. Virol. (1992); Johann et al., J. Virol. (1992); Sommerfelt et al., Virol. (1990); Wilson et al., J. Virol. (1989); Miller et al., J. Virol. (1991); PCT International Publication No. WO/1994/026877A1).

[0146] At least six viral vector approaches are currently available for gene transfer in clinical trials, which utilize approaches that involve complementation of defective vectors by genes inserted into helper cell lines to generate the transducing agent.

[0147] pLASN and MFG-S are examples of retroviral vectors that have been used in clinical trials (Dunbar et al., Blood (1995); Kohn et al., Nat. Med. (1995); Malech et al., PNAS (1997)). PA317/pLASN was the first therapeutic vector used in a gene therapy trial. (Blaese et al., Science (1995)). Transduction efficiencies of 50% or greater have been observed for MFG-S packaged vectors. (Ellem et al., Immunol Immunother. (1997); Dranoff et al., Hum. Gene Ther. (1997)).

[0148] Packaging cells are used to form virus particles that are capable of infecting a host cell. Such cells include 293 cells, which package adenovirus, AAV, and .psi.2 cells or PA317 cells, which package retrovirus. Viral vectors used in gene therapy are usually generated by a producer cell line that packages a nucleic acid vector into a viral particle. The vectors typically contain the minimal viral sequences required for packaging and subsequent integration into a host (if applicable), other viral sequences being replaced by an expression cassette encoding the protein to be expressed. The missing viral functions are supplied in trans by the packaging cell line. For example, AAV vectors used in gene therapy typically only possess inverted terminal repeat (ITR) sequences from the AAV genome which are required for packaging and integration into the host genome. Viral DNA is packaged in a cell line, which contains a helper plasmid encoding the other AAV genes, namely rep and cap, but lacking ITR sequences. The cell line is also infected with adenovirus as a helper. The helper virus promotes replication of the AAV vector and expression of AAV genes from the helper plasmid. The helper plasmid is not packaged in significant amounts due to a lack of ITR sequences. Contamination with adenovirus can be reduced by, e.g., heat treatment to which adenovirus is more sensitive than AAV. Additionally, AAV can be produced at clinical scale using baculovirus systems (see U.S. Pat. No. 7,479,554).

[0149] In many gene therapy applications, it is desirable that the gene therapy vector be delivered with a high degree of specificity to a particular tissue type. Accordingly, a viral vector can be modified to have specificity for a given cell type by expressing a ligand as a fusion protein with a viral coat protein on the outer surface of the virus. The ligand is chosen to have affinity for a receptor known to be present on the cell type of interest. For example, Han et al., Proc. Natl. Acad. Sci. USA 92:9747-9751 (1995), reported that Moloney murine leukemia virus can be modified to express human heregulin fused to gp70, and the recombinant virus infects certain human breast cancer cells expressing human epidermal growth factor receptor. This principle can be extended to other virus-target cell pairs, in which the target cell expresses a receptor and the virus expresses a fusion protein comprising a ligand for the cell-surface receptor. For

example, filamentous phage can be engineered to display antibody fragments (e.g., Fab or Fv) having specific binding affinity for virtually any chosen cellular receptor. Although the above description applies primarily to viral vectors, the same principles can be applied to nonviral vectors. Such vectors can be engineered to contain specific uptake sequences which favor uptake by specific target cells.

[0150] Gene therapy vectors can be delivered in vivo by administration to an individual patient, typically by systemic administration (e.g., intravenous, intraperitoneal, intramuscular, subdermal, or intracranial infusion) or topical application, as described below. Alternatively, vectors can be delivered to cells ex vivo, such as cells explanted from an individual patient (e.g., lymphocytes, bone marrow aspirates, tissue biopsy) or universal donor hematopoietic stem cells, followed by reimplantation of the cells into a patient, usually after selection for cells which have incorporated the vector.

[0151] Ex vivo cell transfection for diagnostics, research, or for gene therapy (e.g., via re-infusion of the transfected cells into the host organism) is well known to those of skill in the art. In a preferred embodiment, cells are isolated from the subject organism, transfected with an RNA composition, and re-infused back into the subject organism (e.g., patient). Various cell types suitable for ex vivo transfection are well known to those of skill in the art (see, e.g., Freshney et al., Culture of Animal Cells, A Manual of Basic Technique (3rd ed. 1994)) and the references cited therein for a discussion of how to isolate and culture cells from patients).

[0152] Suitable cells include but not limited to eukaryotic and prokaryotic cells and/or cell lines. Non-limiting examples of such cells or cell lines generated from such cells include COS, CHO (e.g., CHO-S, CHO-K1, CHO-DG44, CHO-DUXB11, CHO-DUKX, CHO-K1SV), VERO, MDCK, WI38, V79, B14AF28-G3, BHK, HaK, NSO, SP2/0-Ag14, HeLa, HEK293 (e.g., HEK293-F, HEK293-H, HEK293-T), and perC6 cells, any plant cell (differentiated or undifferentiated) as well as insect cells such as *Spodoptera fugiperda* (Sf), or fungal cells such as *Saccharomyces*, *Pichia* and *Schizosaccharomyces*. In certain embodiments, the cell line is a CHO-K1, MDCK or HEK293 cell line. Additionally, primary cells may be isolated and used ex vivo for reintroduction into the subject to be treated following treatment with nuclease systems (e.g. CRISPR/Cas). Suitable primary cells include peripheral blood mononuclear cells (PBMC), and other blood cell subsets such as, but not limited to, CD4+ T cells or CD8+ T cells. Suitable cells also include stem cells such as, by way of example, embryonic stem cells, induced pluripotent stem cells, hematopoietic stem cells (CD34+), neuronal stem cells and mesenchymal stem cells.

[0153] In one embodiment, stem cells are used in ex vivo procedures for cell transfection and gene therapy. The advantage to using stem cells is that they can be differentiated into other cell types in vitro, or can be introduced into a mammal (such as the donor of the cells) where they will engraft in the bone marrow. Methods for differentiating CD34+ cells in vitro into clinically important immune cell types using cytokines such as GM-CSF, IFN- γ , and TNF- α are known (as a non-limiting example see, Inaba et al., J. Exp. Med. 176:1693-1702 (1992)).

[0154] Stem cells are isolated for transduction and differentiation using known methods. For example, stem cells are isolated from bone marrow cells by panning the bone

marrow cells with antibodies which bind unwanted cells, such as CD4+ and CD8+ (T cells), CD45+ (panB cells), GR-1 (granulocytes), and lad (differentiated antigen presenting cells) (as a non-limiting example see Inaba et al., J. Exp. Med. 176:1693-1702 (1992)). Stem cells that have been modified may also be used in some embodiments.

[0155] Notably, any one of the OMNI-50 variant described herein may be suitable for genome editing in post-mitotic cells or any cell which is not actively dividing, e.g., arrested cells. Examples of post-mitotic cells which may be edited using an OMNI-50 variant of the present invention include, but are not limited to, myocyte, a cardiomyocyte, a hepatocyte, an osteocyte and a neuron.

[0156] Vectors (e.g., retroviruses, liposomes, etc.) containing therapeutic RNA compositions can also be administered directly to an organism for transduction of cells in vivo. Alternatively, naked RNA or mRNA can be administered. Administration is by any of the routes normally used for introducing a molecule into ultimate contact with blood or tissue cells including, but not limited to, injection, infusion, topical application and electroporation. Suitable methods of administering such nucleic acids are available and well known to those of skill in the art, and, although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

[0157] Vectors suitable for introduction of transgenes into immune cells (e.g., T-cells) include non-integrating lentivirus vectors. See, for example, U.S. Patent Publication No. 2009/0117617.

[0158] Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions available, as described below (see, e.g., Remington's Pharmaceutical Sciences, 17th ed., 1989).

DNA Repair by Homologous Recombination

[0159] In some embodiments of the present invention, a variant OMNI-50 nuclease is utilized to affect a DNA break at a target site to induce cellular repair mechanisms, for example, but not limited to, non-homologous end-joining (NHEJ) or homology-directed repair (HDR).

[0160] The term "homology-directed repair" or "HDR" refers to a mechanism for repairing DNA damage in cells, for example, during repair of double-stranded and single-stranded breaks in DNA. HDR requires nucleotide sequence homology and uses a "nucleic acid template" (nucleic acid template or donor template used interchangeably herein) to repair the sequence where the double-stranded or single break occurred (e.g., DNA target sequence). This results in the transfer of genetic information from, for example, the nucleic acid template to the DNA target sequence. HDR may result in alteration of the DNA target sequence (e.g., insertion, deletion, mutation) if the nucleic acid template sequence differs from the DNA target sequence and part or all of the nucleic acid template polynucleotide or oligonucleotide is incorporated into the DNA target sequence. In some embodiments, an entire nucleic acid template polynucleotide, a portion of the nucleic acid template polynucleotide, or a copy of the nucleic acid template is integrated at the site of the DNA target sequence.

[0161] The terms "nucleic acid template" and "donor", refer to a nucleotide sequence that is inserted or copied into a genome. The nucleic acid template comprises a nucleotide sequence, e.g., of one or more nucleotides, that will be added to or will template a change in the target nucleic acid or may be used to modify the target sequence. A nucleic acid template sequence may be of any length, for example between 2 and 10,000 nucleotides in length (or any integer value there between or there above), preferably between about 100 and 1,000 nucleotides in length (or any integer there between), more preferably between about 200 and 500 nucleotides in length. A nucleic acid template may be a single stranded nucleic acid, a double stranded nucleic acid. In some embodiment, the nucleic acid template comprises a nucleotide sequence, e.g., of one or more nucleotides, that corresponds to wild type sequence of the target nucleic acid, e.g., of the target position. In some embodiment, the nucleic acid template comprises a ribonucleotide sequence, e.g., of one or more ribonucleotides, that corresponds to wild type sequence of the target nucleic acid, e.g., of the target position. In some embodiment, the nucleic acid template comprises modified ribonucleotides.

[0162] Insertion of an exogenous sequence (also called a "donor sequence," donor template" or "donor"), for example, for correction of a mutant gene or for increased expression of a wild-type gene can also be carried out. It will be readily apparent that the donor sequence is typically not identical to the genomic sequence where it is placed. A donor sequence can contain a non-homologous sequence flanked by two regions of homology to allow for efficient HDR at the location of interest. Additionally, donor sequences can comprise a vector molecule containing sequences that are not homologous to the region of interest in cellular chromatin. A donor molecule can contain several, discontinuous regions of homology to cellular chromatin. For example, for targeted insertion of sequences not normally present in a region of interest, said sequences can be present in a donor nucleic acid molecule and flanked by regions of homology to sequence in the region of interest.

[0163] The donor polynucleotide can be DNA or RNA, single-stranded and/or double-stranded and can be introduced into a cell in linear or circular form. See, e.g., U.S. Patent Publication Nos. 2010/0047805; 2011/0281361; 2011/0207221; and 2019/0330620. If introduced in linear form, the ends of the donor sequence can be protected (e.g., from exonucleolytic degradation) by methods known to those of skill in the art. For example, one or more dideoxynucleotide residues are added to the 3' terminus of a linear molecule and/or self-complementary oligonucleotides are ligated to one or both ends. See, for example, Chang and Wilson, Proc. Natl. Acad. Sci. USA (1987); Nehls et al., Science (1996). Additional methods for protecting exogenous polynucleotides from degradation include, but are not limited to, addition of terminal amino group(s) and the use of modified internucleotide linkages such as, for example, phosphorothioates, phosphoramidates, and O-methyl ribose or deoxyribose residues.

[0164] Accordingly, embodiments of the present invention using a donor template for repair may use a DNA or RNA, single-stranded and/or double-stranded donor template that can be introduced into a cell in linear or circular form. In embodiments of the present invention a gene-editing composition comprises: (1) an RNA molecule comprising a guide sequence to affect a double strand break in a gene prior

to repair and (2) a donor RNA template for repair, and the RNA molecule comprising the guide sequence is a first RNA molecule and the donor RNA template is a second RNA molecule. In some embodiments, the guide RNA molecule and template RNA molecule are connected as part of a single molecule.

[0165] A donor sequence may also be an oligonucleotide and be used for gene correction or targeted alteration of an endogenous sequence. The oligonucleotide may be introduced to the cell on a vector, may be electroporated into the cell, or may be introduced via other methods known in the art. The oligonucleotide can be used to ‘correct’ a mutated sequence in an endogenous gene (e.g., the sickle mutation in beta globin), or may be used to insert sequences with a desired purpose into an endogenous locus.

[0166] A polynucleotide can be introduced into a cell as part of a vector molecule having additional sequences such as, for example, replication origins, promoters and genes encoding antibiotic resistance. Moreover, donor polynucleotides can be introduced as naked nucleic acid, as nucleic acid complexed with or packaged within an agent such as a liposome, exosome, or poloxamer, or can be delivered by recombinant viruses (e.g., adenovirus, AAV, herpesvirus, retrovirus, lentivirus and integrase defective lentivirus (IDLV)) or virus-like particles. Non-viral vectors, such as transposon-based systems, e.g. recombinant Sleeping Beauty transposon systems or recombinant PiggyBac transposon systems, may also be utilized for transposition of a polynucleotide sequence in a target cell.

[0167] The donor is generally inserted so that its expression is driven by the endogenous promoter at the integration site, namely the promoter that drives expression of the endogenous gene into which the donor is inserted. However, it will be apparent that the donor may comprise a promoter and/or enhancer, for example a constitutive promoter or an inducible or tissue specific promoter.

[0168] The donor molecule may be inserted into an endogenous gene such that all, some or none of the endogenous gene is expressed. For example, a transgene as described herein may be inserted into an endogenous locus such that some (N-terminal and/or C-terminal to the transgene) or none of the endogenous sequences are expressed, for example as a fusion with the transgene. In other embodiments, the transgene (e.g., with or without additional coding sequences such as for the endogenous gene) is integrated into any endogenous locus, for example a safe-harbor locus, for example a CCR5 gene, a CXCR4 gene, a PPP1R12c (also known as AAVS1) gene, an albumin gene or a Rosa gene. See, e.g., U.S. Pat. Nos. 7,951,925 and 8,110,379; U.S. Publication Nos. 2008/0159996; 20100/0218264; 2010/0291048; 2012/0017290; 2011/0265198; 2013/0137104; 2013/0122591; 2013/0177983 and 2013/0177960 and U.S. Provisional Application No. 61/823,689).

[0169] When endogenous sequences (endogenous or part of the transgene) are expressed with the transgene, the endogenous sequences may be full-length sequences (wild-type or mutant) or partial sequences. Preferably the endogenous sequences are functional. Non-limiting examples of the function of these full length or partial sequences include increasing the serum half-life of the polypeptide expressed by the transgene (e.g., therapeutic gene) and/or acting as a carrier.

[0170] Furthermore, although not required for expression, exogenous sequences may also include transcriptional or

translational regulatory sequences, for example, promoters, enhancers, insulators, internal ribosome entry sites, sequences encoding 2A peptides and/or polyadenylation signals.

[0171] In certain embodiments, the donor molecule comprises a sequence selected from the group consisting of a gene encoding a protein (e.g., a coding sequence encoding a protein that is lacking in the cell or in the individual or an alternate version of a gene encoding a protein), a regulatory sequence and/or a sequence that encodes a structural nucleic acid such as a microRNA or siRNA.

Dna-Targeting RNA Molecules

[0172] In embodiments of the present invention, the DNA-targeting RNA sequence comprises a guide sequence portion. The “guide sequence portion” of an RNA molecule refers to a nucleotide sequence that is capable of hybridizing to a specific target DNA sequence, e.g., the guide sequence portion has a nucleotide sequence which is fully complementary to the DNA sequence being targeted along the length of the guide sequence portion. In some embodiments, the guide sequence portion is 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides in length, or approximately 17-30, 17-29, 17-28, 17-27, 17-26, 27-25, 17-24, 18-22, 19-22, 18-20, 17-20, or 21-22 nucleotides in length. The entire length of the guide sequence portion is fully complementary to the DNA sequence being targeted along the length of the guide sequence portion. The guide sequence portion may be part of an RNA molecule that can form a complex with a CRISPR nuclease with the guide sequence portion serving as the DNA targeting portion of the CRISPR complex. When the RNA molecule having the guide sequence portion is present contemporaneously with the CRISPR molecule, the RNA molecule is capable of targeting the CRISPR nuclease to the specific target DNA sequence. Each possibility represents a separate embodiment. An RNA molecule can be custom designed to target any desired sequence.

[0173] According to some aspects of the invention, the disclosed methods comprise a method of modifying a nucleotide sequence at a target site in a cell-free system or the genome of a cell comprising introducing into the cell the composition of any one of the embodiments described herein.

[0174] In some embodiments, the cell is a eukaryotic cell, preferably a mammalian cell or a plant cell. In some embodiments, genome modifying occurs within the nucleus of a cell.

[0175] According to some aspects of the invention, the disclosed methods comprise a use of any one of the compositions described herein for the treatment of a subject afflicted with a disease associated with a genomic mutation comprising modifying a nucleotide sequence at a target site in the genome of the subject.

[0176] According to some aspects of the invention, the disclosed methods comprise a method of treating subject having a mutation disorder comprising targeting any one of the compositions described herein to an allele associated with the mutation disorder.

[0177] In some embodiments, the mutation disorder is related to a disease or disorder selected from any of a neoplasia, age-related macular degeneration, schizophrenia, neurological, neurodegenerative, or movement disorder, Fragile X Syndrome, secretase-related disorders, prion-re-

lated disorders, ALS, addiction, autism, Alzheimer's Disease, neutropenia, inflammation-related disorders, Parkinson's Disease, blood and coagulation diseases and disorders, beta thalassemia, sickle cell anemia, cell dysregulation and oncology diseases and disorders, inflammation and immune-related diseases and disorders, metabolic, liver, hypercholesterolemia, kidney and protein diseases and disorders, muscular and skeletal diseases and disorders, dermatological diseases and disorders, neurological and neuronal diseases and disorders, pulmonary disease and disorders, corneal disease and disorders, retinal diseases and disorders, and ocular diseases and disorders.

Diseases and Therapies

[0178] Certain embodiments of the invention target a nuclease to a specific genetic locus associated with a disease or disorder as a form of gene editing, method of treatment, or therapy. For example, to induce editing or knockout of a gene, a novel nuclease disclosed herein may be specifically targeted to a pathogenic mutant allele of the gene using a custom designed guide RNA molecule. The guide RNA molecule is preferably designed by first considering the PAM requirement of the nuclease, which as shown herein is also dependent on the system in which the gene editing is being performed. For example, a guide RNA molecule designed to target an OMNI-50 nuclease to a target site is designed to contain a spacer sequence complementary to a DNA strand of a DNA double-stranded region that neighbors a OMNI-140 PAM sequence, e.g. "NGG." The guide RNA molecule is further preferably designed to contain a spacer region (i.e. the region of the guide RNA molecule having complementarity to the target allele) of sufficient and preferably optimal length in order to increase specific activity of the nuclease and reduce off-target effects.

[0179] As a non-limiting example, the guide RNA molecule may be designed to target the nuclease to a specific region of a mutant allele, e.g. near the start codon, such that upon DNA damage caused by the nuclease a non-homologous end joining (NHEJ) pathway is induced and leads to silencing of the mutant allele by introduction of frameshift mutations. This approach to guide RNA molecule design is particularly useful for altering the effects of dominant negative mutations and thereby treating a subject. As a separate non-limiting example, the guide RNA molecule may be designed to target a specific pathogenic mutation of a mutated allele, such that upon DNA damage caused by the nuclease a homology directed repair (HDR) pathway is induced and leads to template mediated correction of the mutant allele. This approach to guide RNA molecule design is particularly useful for altering haploinsufficiency effects of a mutated allele and thereby treating a subject.

[0180] Non-limiting examples of specific genes which may be targeted for alteration to treat a disease or disorder are presented herein below. Specific disease-associated genes and mutations that induce a mutation disorder are described in the literature. Such mutations can be used to design a DNA-targeting RNA molecule to target a CRISPR composition to an allele of the disease associated gene, where the CRISPR composition causes DNA damage and induces a DNA repair pathway to alter the allele and thereby treat the mutation disorder.

[0181] Mutations in the ELANE gene are associated with neutropenia. Accordingly, without limitation, embodiments of the invention that target ELANE may be used in methods of treating subjects afflicted with neutropenia. Guide RNA molecules which target the ELANE gene and are useful for treating neutropenia are disclosed in PCT International Application No. PCT/US2020/059186, incorporated herein by reference.

[0182] CXCR4 is a co-receptor for the human immunodeficiency virus type 1 (HIV-1) infection. Accordingly, without limitation, embodiments of the invention that target CXCR4 may be used in methods of treating subjects afflicted with HIV-1 or conferring resistance to HIV-1 infection in a subject.

[0183] Programmed cell death protein 1 (PD-1) disruption enhances CAR-T cell mediated killing of tumor cells and PD-1 may be a target in other cancer therapies. Accordingly, without limitation, embodiments of the invention that target PD-1 may be used in methods of treating subjects afflicted with cancer. In an embodiment, the treatment is CAR-T cell therapy with T cells that have been modified according to the invention to be PD-1 deficient.

[0184] In addition, BCL11A is a gene that plays a role in the suppression of hemoglobin production. Globin production may be increased to treat diseases such as thalassemia or sickle cell anemia by inhibiting BCL11A. See for example, PCT International Publication No. WO 2017/077394A2; U.S. Publication No. US2011/0182867A1; Humbert et al. Sci. Transl. Med. (2019); and Canver et al. Nature (2015). Accordingly, without limitation, embodiments of the invention that target an enhancer of BCL11A may be used in methods of treating subjects afflicted with beta thalassemia or sickle cell anemia.

[0185] Embodiments of the invention may also be used for targeting any disease-associated gene, for studying, altering, or treating any of the diseases or disorders listed in Table A or Table B below. Indeed, any disease-associated with a genetic locus may be studied, altered, or treated by using the nucleases disclosed herein to target the appropriate disease-associated gene, for example, those listed in U.S. Publication No. 2018/0282762A1 and European Patent No. EP3079726B1.

TABLE A

Diseases, Disorders and their associated genes	
DISEASE/ DISORDERS	GENE(S)
Neoplasia	PTEN; ATM; ATR; EGFR; ERBB2; ERBB3; ERBB4; Notch1; Notch2; Notch3; Notch4; AKT; AKT2; AKT3; HIF; HIF1a; HIF3a; Met; HRG; Bcl2; PPAR alpha; PPAR gamma; WT1 (Wilms Tumor); FGF Receptor Family members (5 members: 1, 2, 3, 4, 5); CDKN2a; APC; RB (retinoblastoma); MEN1; VHL; BRCA1; BRCA2; AR

TABLE A-continued

Diseases, Disorders and their associated genes	
DISEASE/ DISORDERS	GENE(S)
	(Androgen Receptor); TSG101; IGF; IGF Receptor; Igf1 (4 variants); gf2 (3 variants); Igf 1 Receptor; Igf 2 Receptor; Bax; Bcl2; caspases family (9 members: 1, 2, 3, 4, 6, 7, 8, 9, 12); Kras; Apc
Age-related Macular Degeneration	Abcr; Ccl2; Cc2; cp (ceruloplasmin); Timp3; cathepsinD; Vldlr; Cer2
Schizophrenia	Neuregulin1 (Nrg1); Erb4 (receptor for Neuregulin); Complexin1 (Cp1x1); Tph1 Tryptophan hydroxylase; Tph2 Tryptophan hydroxylase 2; Neurexin 1; GSK3; GSK3a; GSK3b
Neurological, Neuro-degenerative, and Movement Disorders	5-HTT (S1c6a4); COMT; DRD (Drd1a); SLC6A3; DAOA; DTNBP1; Dao (Dao1)
Trinucleotide Repeat Disorders	HTT (Huntington's Dx); SBMA/SMAX1/AR (Kennedy's Dx); FXN/X25 (Friedrich's Ataxia); ATX3 (Machado-Joseph's Dx); ATXN1 and ATXN2 (spinocerebellar ataxias); DMPK (myotonic dystrophy); Atrophin-1 and Atn1 (DRPLA Dx); CBP (Creb-BP - global instability); VLDLR (Alzheimer's); Atxn7; Atxn10
Fragile X Syndrome	FMR2; FXR1; FXR2; mGLUR5
Secretase Related Disorders	APH-1 (alpha and beta); Presenilin (Psen1); nicastrin (Ncstn); PEN-2
Others	Nos1; Parp1; Nat1; Nat2
Prion related disorders	Prp
ALS	SOD1; ALS2; STEX; FUS; TARDBP; VEGF (VEGF-a; VEGF-b; VEGF-c)
Addiction	Prkce (alcohol); Drd2; Drd4; ABAT (alcohol); GRIA2; Grm5; Grin1; Htr1b; Grin2a; Drd3; Pdyn; Gria1 (alcohol)
Autism	Mecp2; BZRAP1; MDGA2; Sema5A; Neurexin 1; Fragile X (FMR2 (AFF2); FXR1; FXR2; Mglur5)
Alzheimer's Disease	E1; CHIP; UCH; UBB; Tau; LRP; PICALM; Clusterin; PS1; SORL1; CR1; Vldlr; Uba1; Uba3; CHIP28 (Aqp1, Aquaporin 1); Uchl1; Uchl3; APP
Inflammation	IL-10; IL-1 (IL-1a; IL-1b); IL-13; IL-17 (IL-17a (CTLA8); IL-17b; IL-17c; IL-17d; IL-17f); IL-23; Cx3cr1; ptpn22; TNFa; NOD2/CARD15 for IBD; IL-6; IL-12 (IL-12a; IL-12b); CTLA4; Cx3cl1
Parkinson's Disease	x-Synuclein; DJ-1; LRRK2; Parkin; PINK1

TABLE B

Diseases, Disorders and their associated genes	
DISEASE CATEGORY	DISEASE AND ASSOCIATED GENES
Blood and coagulation diseases and disorders	Anemia (CDAN1, CDA1, RPS19, DBA, PKLR, PK1, NT5C3, UMPH1, PSN1, RHAG, RH50A, NRAMP2, SPTB, ALAS2, ANH1, ASB, ABCB7, ABC7, ASAT); Bare lymphocyte syndrome (TAPBP, TPSN, TAP2, ABCB3, PSF2, RING11, MHC2TA, C2TA, RFX5, RFXAP, RFX5); Bleeding disorders (TBXA2R, P2RX1, P2X1); Factor H and factor H-like 1 (HF1, CFH, HUS); Factor V and factor VIII (MCFD2); Factor VII deficiency (F7); Factor X deficiency (F10); Factor XI deficiency (F11); Factor XII deficiency (F12, HAF); Factor XIIIa deficiency (F13A1, F13A); Factor XIIIb deficiency (F13B); Fanconi anemia (FANCA, FACA, FA1, FA, FAA, FAAP95, FAAP90, FLJ34064, FANCB, FANCC, FACC, BRCA2, FANCD1, FANCD2, FANCD, FACD, FAD, FANCE, FACE, FANCE, XRCC9, FANCG, BRIP1, BACH1, FANCI, PHF9, FANCL, FANCM, KIAA1596); Hemophagocytic lymphohistiocytosis disorders (PRF1, HPLH2, UNC13D, MUNC13-4, HPLH3, HLH3, FHL3); Hemophilia A (F8, F8C, HEMA); Hemophilia B (F9, HEMB); Hemorrhagic disorders (PI, ATT, F5); Leukocyte deficiencies and disorders (ITGB2, CD18, LCAMB, LAD, EIF2B1, EIF2BA, EIF2B2, EIF2B3, EIF2B5, LVWM, CACH, CLE, EIF2B4); Sickle cell anemia (HBB); Thalassemia (HBA2, HBB, HBD, LCRB, HBA1)

TABLE B-continued

Diseases, Disorders and their associated genes	
DISEASE CATEGORY	DISEASE AND ASSOCIATED GENES
Cell dysregulation and oncology diseases and disorders	B-cell non-Hodgkin lymphoma (BCL7A, BCL7); Leukemia (TAL1, TCL5, SCL, TAL2, FLT3, NBS1, NBS, ZNFN1A1, IK1, LYF1, HOXD4, HOX4B, BCR, CML, PHL, ALL, ARNT, KRAS2, RASK2, GMPS, AF10, ARHGEF12, LARG, KIAA0382, CALM, CLTH, CEBPA, CEBP, CHIC2, BTL, FLT3, KIT, PBT, LPP, NPM1, NUP214, D9S46E, CAN, CAIN, RUNX1, CBFA2, AML1, WHSC1L1, NSD3, FLT3, AFIQ, NPM1, NUMA1, ZNF145, PLZF, PML, MYL, STAT5B, AF10, CALM, CLTH, ARL11, ARLTS1, P2RX7, P2X7, BCR, CML, PHL, ALL, GRAF, NF1, VRNF, WSS, NENS, PTPN11, PTP2C, SHP2, NS1, BCL2, CCND1, PRAD1, BCL1, TCRA, GATA1, GF1, ERYF1, NFE1, ABL1, NQO1, DIA4, NMOR1, NUP214, D9S46E, CAN, CAIN)
Inflammation and immune related diseases and disorders	AIDS (KIR3DL1, NKAT3, NKB1, AMB11, KIR3DS1, IFNG, CXCL12, SDF1); Autoimmune lymphoproliferative syndrome (TNFRSF6, APT1, FAS, CD95, ALPS1A); Combined immunodeficiency, (IL2RG, SCIDX1, SCIDX, IMD4); HIV-1 (CCL5, SCYA5, D17S136E, TCP228), HIV susceptibility or infection (IL10, CSIF, CMKBR2, CCR2, CMKBR5, CCCKR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFSF5, CD40LG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TAC1); Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-17f), IL-23, Cx3cr1, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b), CTLA4, Cx3cl1); Severe combined immunodeficiencies (SCIDs)(JAK3, JAKL, DCLRE1C, ARTEMIS, SCIDA, RAG1, RAG2, ADA, PTPRC, CD45, LCA, IL7R, CD3D, T3D, IL2RG, SCIDX1, SCIDX, IMD4)
Metabolic, liver, kidney and protein diseases and disorders	Amyloid neuropathy (TTR, PALB); Amyloidosis (APOA1, APP, AAA, CVAP, AD1, GSN, FGA, LYZ, TTR, PALB); Cirrhosis (KRT18, KRT8, CIRH1A, NAIC, TEX292, KIAA1988); Cystic fibrosis (CFTR, ABCC7, CF, MRP7); Glycogen storage diseases (SLC2A2, GLUT2, G6PC, G6PT, G6PT1, GAA, LAMP2, LAMPB, AGL, GDE, GBE1, GYS2, PYGL, PFKM); Hepatic adenoma, 142330 (TCF1, HNF1A, MODY3), Hepatic failure, early onset, and neurologic disorder (SCOD1, SCO1), Hepatic lipase deficiency (LIPC), Hepatoblastoma, cancer and carcinomas (CTNNB1, PDGFRL, PDGRL, PRLTS, AXIN1, AXIN, CTNNB1, TP53, P53, LFS1, IGF2R, MPRI, MET, CASP8, MCH5; Medullary cystic kidney disease (UMOD, HNFJ, FJHN, MCKD2, ADMCKD2); Phenylketonuria (PAH, PKU1, QDPR, DHPR, PTS); Polycystic kidney and hepatic disease (FCYT, PKHD1, ARPKD, PKD1, PKD2, PKD4, PKDTS, PRKCSH, G19P1, PCLD, SEC63)
Muscular/Skeletal diseases and disorders	Becker muscular dystrophy (DMD, BMD, MYF6), Duchenne Muscular Dystrophy (DMD, BMD); Emery-Dreifuss muscular dystrophy (LMNA, LMN1, EMD2, FPLD, CMD1A, HGPS, LGMD1B, LMNA, LMN1, EMD2, FPLD, CMD1A); Facioscapulohumeral muscular dystrophy (FSHMD1A, FSHD1A); Muscular dystrophy (FKRP, MDC1C, LGMD2I, LAMA2, LAMM, LARGE, KIAA0609, MDC1D, FCMD, TTID, MYOT, CAPN3, CANP3, DYSE, LGMD2B, SGCG, LGMD2C, DMDA1, SCG3, SGCA, ADL, DAG2, LGMD2D, DMDA2, SGCB, LGMD2E, SGCD, SGD, LGMD2F, CMD1L, TCAP, LGMD2G, CMD1N, TRIM32, HT2A, LGMD2H, FKRP, MDC1C, LGMD2I, TTN, CMD1G, TMD, LGMD2J, POMT1, CAV3, LGMD1C, SEP1, SELN, RSMD1, PLEC1, PLTN, EBS1); Osteopetrosis (LRP5, BMND1, LRP7, LR3, OPPG, VBCH2, CLCN7, CLC7, OPTA2, OSTM1, GL, TCIRG1, TIRC7, OC116, OPTB1); Muscular atrophy (VAPB, VAPC, ALS8, SMN1, SMA1, SMA2, SMA3, SMA4, BSCL2, SPG17, GARS, SMAD1, CMT2D, HEXB, IGHMBP2, SMUBP2, CATF1, SMARD1)
Dermatological diseases and disorders	Albinism (TYR, OCA2, TYRP1, SLC45A2, LYST), Ectodermal dysplasias (EDAR, EDARADD, WNT10A), Ehlers-Danlos syndrome (COL5A1, COL5A2, COL1A1, COL1A2, COL3A1, TNXB, ADAMTS2, PLOD1, FKBP14), Ichthyosis-associated disorders (FLG, STS,

TABLE B-continued

Diseases, Disorders and their associated genes	
DISEASE CATEGORY	DISEASE AND ASSOCIATED GENES
Neurological and Neuronal diseases and disorders	TGM1, ALOXE3/ALOX12B, KRT1, KRT10, ABCA12, KRT2, GJB2, TGM1, ABCA12, CYP4F22, ALOXE3, CERS3, NSHDL, EBP, MBTPS2, GJB2, SPINK5, AGHD5, PHYH, PEX7, ALDH3A2, ERCC2, ERCC3, GFT2H5, GBA), Incontinentia pigmenti (IKBKG, NEMO), Tuberous sclerosis (TSC1, TSC2), Premature aging syndromes (POLR3A, PYCR1, LMNA, POLD1, WRN, DMPK)
	ALS (SOD1, ALS2, STEX, FUS, TARDBP, VEGF (VEGF-a, VEGF-b, VEGF-c); Alzheimer disease (APP, AAA, CVAP, AD1, APOE, AD2, PSEN2, AD4, STM2, APBB2, FE65L1, NOS3, PLAUI, URK, ACE, DCP1, ACE1, MPO, PACIP1, PAXIP1L, PTIP, A2M, BLMH, BMH, PSEN1, AD3); Autism (Mecp2, BZRAP1, MDGA2, Sema5A, Neurexin 1, GLO1, MECP2, RTT, PPMX, MRX16, MRX79, NLGN3, NLGN4, KIAA1260, AUTSX2); Fragile X Syndrome (FMR2, FXR1, FXR2, mGLUR5); Huntington's disease and disease like disorders (HD, IT15, PRNP, PRIP, JPH3, JP3, HDL2, TBP, SCA17); Parkinson disease (NR4A2, NURR1, NOT, TINUR, SNCAIP, TBP, SCA17, SNCA, NACP, PARK1, PARK4, DJ1, PARK7, LRRK2, PARK8, PINK1, PARK6, UCHL1, PARK5, SNCA, NACP, PARK1, PARK4, PRKN, PARK2, PDJ, DBH, NDUFV2); Rett syndrome (MECP2, RTT, PPMX, MRX16, MRX79, CDKL5, STK9, MECP2, RTT, PPMX, MRX16, MRX79, x-Synuclein, DJ-1); Schizophrenia (Neuregulin1 (Nrg1), Erb4 (receptor for Neuregulin), Complexin1 (Cplx1), Tph1 Tryptophan hydroxylase, Tph2, Tryptophan hydroxylase 2, Neurexin 1, GSK3, GSK3a, GSK3b, 5-HTT (Slc6a4), COMT, DRD (Drd1a), SLC6A3, DAOA, DTNBP1, Dao (Dao1)); Secretase Related Disorders (APH-1 (alpha and beta), Presenilin (Psen1), nicastrin, (Ncstn), PEN-2, Nos1, Parp1, Natl, Nat2); Trinucleotide Repeat Disorders (HTT (Huntington's Dx), SBMA/SMAX1/AR (Kennedy's Dx), FXN/X25 (Friedrich's Ataxia), ATX3 (Machado-Joseph's Dx), ATXN1 and ATXN2 (spinocerebellar ataxias), DMPK (myotonic dystrophy), Atrophin-1 and Atn1 (DRPLA Dx), CBP (Creb-BP - global instability), VLDLR (Alzheimer's), Atxn7, Atxn10)
Ocular diseases and disorders	Age-related macular degeneration (Abcr, Ccl2, Cc2, cp (ceruloplasmin), Timp3, cathepsinD, Vldlr, Ccr2); Cataract (CRYAA, CRYA1, CRYBB2, CRYB2, PITX3, BFSP2, CP49, CP47, CRYAA, CRYA1, PAX6, AN2, MGDA, CRYBA1, CRYB1, CRYGC, CRYG3, CCL, LIM2, MP19, CRYGD, CRYG4, BFSP2, CP49, CP47, HSF4, CTM, HSF4, CTM, MIP, AQPO, CRYAB, CRYA2, CTPP2, CRYBB1, CRYGD, CRYG4, CRYBB2, CRYB2, CRYGC, CRYG3, CCL, CRYAA, CRYA1, GJA8, CX50, CAE1, GJA3, CX46, CZP3, CAE3, CCM1, CAM, KRIT1); Corneal clouding and dystrophy (APOA1, TGFBI, CSD2, CDGG1, CSD, BIGH3, CDG2, TACSTD2, TROP2, MIS1, VSX1, RINX, PPCD, PPD, KTCN, COL8A2, FECD, PPCD2, PIP5K3, CFD); Cornea plana congenital (KERA, CNA2); Glaucoma (MYOC, TIGR, GLC1A, JOAG, GPOA, OPTN, GLC1E, FIP2, HYPL, NRP, CYP1B1, GLC3A, OPA1, NTG, NPG, CYP1B1, GLC3A); Leber congenital amaurosis (CRB1, RP12, CRX, CORD2, CRD, RPGRIP1, LCA6, CORD9, RPE65, RP20, AIPL1, LCA4, GUCY2D, GUC2D, LCA1, CORD6, RDH12, LCA3); Macular dystrophy (ELOVL4, ADMD, STGD2, STGD3, RDS, RP7, PRPH2, PRPH, AVMD, AOFMD, VMD2)

[0186] Unless otherwise defined, all technical and/or scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the invention pertains. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of embodiments of the invention, exemplary methods and/or materials are described below. In

case of conflict, the patent specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and are not intended to be necessarily limiting.

[0187] In the discussion unless otherwise stated, adjectives such as “substantially” and “about” modifying a condition or relationship characteristic of a feature or features of

an embodiment of the invention, are understood to mean that the condition or characteristic is defined to within tolerances that are acceptable for operation of the embodiment for an application for which it is intended. Unless otherwise indicated, the word “or” in the specification and claims is considered to be the inclusive “or” rather than the exclusive or, and indicates at least one of, or any combination of items it conjoins.

[0188] It should be understood that the terms “a” and “an” as used above and elsewhere herein refer to “one or more” of the enumerated components. It will be clear to one of ordinary skill in the art that the use of the singular includes the plural unless specifically stated otherwise. Therefore, the terms “a,” “an” and “at least one” are used interchangeably in this application.

[0189] For purposes of better understanding the present teachings and in no way limiting the scope of the teachings, unless otherwise indicated, all numbers expressing quantities, percentages or proportions, and other numerical values used in the specification and claims, are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in the following specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained. At the very least, each numerical parameter should at least be construed in light of the number of reported significant digits and by applying ordinary rounding techniques.

[0190] In the description and claims of the present application, each of the verbs, “comprise,” “include” and “have” and conjugates thereof, are used to indicate that the object or objects of the verb are not necessarily a complete listing of components, elements or parts of the subject or subjects of the verb. Other terms as used herein are meant to be defined by their well-known meanings in the art.

[0191] As used herein, the term “targeting sequence” or “targeting molecule” refers a nucleotide sequence or molecule comprising a nucleotide sequence that is capable of hybridizing to a specific target sequence, e.g., the targeting sequence has a nucleotide sequence which is at least partially complementary to the sequence being targeted along the length of the targeting sequence. The targeting sequence or targeting molecule may be part of an RNA molecule that can form a complex with a CRISPR nuclease with the targeting sequence serving as the targeting portion of the CRISPR complex. When the molecule having the targeting sequence is present contemporaneously with the CRISPR molecule the RNA molecule is capable of targeting the CRISPR nuclease to the specific target sequence. Each possibility represents a separate embodiment. An RNA molecule can be custom designed to target any desired sequence.

[0192] The term “targets” as used herein, refers to a targeting sequence or targeting molecule’s preferential hybridization to a nucleic acid having a targeted nucleotide sequence. It is understood that the term “targets” encompasses variable hybridization efficiencies, such that there is preferential targeting of the nucleic acid having the targeted nucleotide sequence, but unintentional off-target hybridization in addition to on-target hybridization might also occur. It is understood that where an RNA molecule targets a sequence, a complex of the RNA molecule and a CRISPR nuclease molecule targets the sequence for nuclease activity.

[0193] As used herein the term “wild-type” is a term of the art understood by skilled persons and means the typical form of an organism, strain, gene or characteristic as it occurs in nature as distinguished from mutant or variant forms. Accordingly, as used herein, where a sequence of amino acids or nucleotides refers to a wild-type sequence, a variant refers to variant of that sequence, e.g., comprising substitutions, deletions, insertions. In embodiments of the present invention, an engineered CRISPR nuclease is a variant CRISPR nuclease comprising at least one amino acid modification (e.g., substitution, deletion, and/or insertion), also referred to as a “mutation,” compared to the wild-type OMNI-50 nuclease of SEQ ID NO:1.

[0194] The terms “non-naturally occurring” or “engineered” are used interchangeably and indicate human manipulation. The terms, when referring to nucleic acid molecules or polypeptides may mean that the nucleic acid molecule or the polypeptide is at least substantially free from at least one other component with which they are naturally associated in nature and as found in nature.

[0195] The terms “mutant” or “variant” are used interchangeably and indicate a molecule that is non-naturally occurring or engineered.

[0196] As used herein the term “amino acid” includes natural and/or unnatural or synthetic amino acids, including glycine and both the D- or L-, optical isomers, and amino acid analogs and peptidomimetics.

[0197] As used herein, “genomic DNA” refers to linear and/or chromosomal DNA and/or to plasmid or other extra-chromosomal DNA sequences present in the cell or cells of interest. In some embodiments, the cell of interest is a eukaryotic cell. In some embodiments, the cell of interest is a prokaryotic cell. In some embodiments, the methods produce double-stranded breaks (DSBs) at pre-determined target sites in a genomic DNA sequence, resulting in mutation, insertion, and/or deletion of DNA sequences at the target site(s) in a genome.

[0198] “Eukaryotic” cells include, but are not limited to, fungal cells (such as yeast), plant cells, animal cells, mammalian cells and human cells.

[0199] As used herein, the term “modified cells” refers to cells in which a double strand break is affected by a complex of an RNA molecule and the CRISPR nuclease variant as a result of hybridization with the target sequence, i.e. on-target hybridization. The term “modified cells” may further encompass cells in which a repair or correction of a mutation was affected following the double strand break induced by the variant. The modified cell may be any type of cell e.g., eukaryotic or prokaryotic, in any environment e.g., isolated or not, maintained in culture, in vitro, ex vivo, in vivo or in planta.

[0200] This invention provides a modified cell or cells obtained by use of any of the variants or methods described herein. In an embodiment these modified cell or cells are capable of giving rise to progeny cells. In an embodiment these modified cell or cells are capable of giving rise to progeny cells after engraftment. As a non-limiting example, the modified cells may be hematopoietic stem cell (HSC), or any cell suitable for an allogenic cell transplant or autologous cell transplant. The variants and methods described herein may also be utilized to generate chimeric antigen receptor T (CAR-T) cells.

[0201] This invention also provides a composition comprising these modified cells and a pharmaceutically accept-

able carrier. Also provided is an in vitro or ex vivo method of preparing this, comprising mixing the cells with the pharmaceutically acceptable carrier.

[0202] The term “nuclease” as used herein refers to an enzyme capable of cleaving the phosphodiester bonds between the nucleotide subunits of nucleic acid. A nuclease may be isolated or derived from a natural source. The natural source may be any living organism. Alternatively, a nuclease may be a modified or a synthetic protein which retains the phosphodiester bond cleaving activity.

[0203] The terms “protospacer adjacent motif” or “PAM” as used herein refers to a nucleotide sequence of a target DNA located in proximity to the targeted DNA sequence and recognized by the CRISPR nuclease. The PAM sequence may differ depending on the nuclease identity. For example, wild-type *Streptococcus pyogenes* Cas9 recognizes a “NGG” PAM sequence. A skilled artisan will appreciate that single-guide RNA molecules or crRNA:tracrRNA complexes capable of complexing with a CRISPR nuclease such as to associate with a target genomic DNA sequence of interest next to a protospacer adjacent motif (PAM). The nuclease then mediates cleavage of target DNA to create a double-stranded break within the protospacer.

[0204] As used herein, a sequence or molecule has an X % “sequence identity” to another sequence or molecule if X % of bases or amino acids between the sequences of molecules are the same and in the same relative position. For example, a first nucleotide sequence having at least a 95% sequence identity with a second nucleotide sequence will have at least 95% of bases, in the same relative position, identical with the other sequence.

[0205] The terms “nuclear localization sequence” and “NLS” are used interchangeably to indicate an amino acid sequence/peptide that directs the transport of a protein with which it is associated from the cytoplasm of a cell across the nuclear envelope barrier. The term “NLS” is intended to encompass not only the nuclear localization sequence of a particular peptide, but also derivatives thereof that are capable of directing translocation of a cytoplasmic polypeptide across the nuclear envelope barrier. NLSs are capable of directing nuclear translocation of a polypeptide when attached to the N-terminus, the C-terminus, or both the N- and C-termini of the polypeptide. In addition, a polypeptide having an NLS coupled by its N- or C-terminus to amino acid side chains located randomly along the amino acid sequence of the polypeptide will be translocated. Typically, an NLS consists of one or more short sequences of positively charged lysines or arginines exposed on the protein surface, but other types of NLS are known. Non-limiting examples of NLSs include an NLS sequence derived from: the SV40 virus large T-antigen, nucleoplasmin, c-myc, the hRNPA1 M9 NLS, the IBB domain from importin- α , myoma T protein, human p53, mouse c-abl IV, influenza vims NS1, Hepatitis virus delta antigen, mouse Mx1 protein, human poly(ADP-ribose) polymerase, and the steroid hormone receptors (human) glucocorticoid.

[0206] The term “CRISPR system” refers to a CRISPR endonuclease system that includes a CRISPR nuclease protein, such as the mutants or variants described herein, and a suitable guide RNA molecule or guide RNA complex, e.g. a single-guide RNA or a crRNA:tracrRNA complex, for targeting the CRISPR nuclease protein to a desired target DNA sequence based on complementarity between a portion of the guide RNA molecule or guide RNA complex and the

target DNA sequence. The term “wild-type CRISPR endonuclease system” refers to a CRISPR endonuclease system that includes wild-type CRISPR protein and a suitable guide RNA molecule or guide RNA complex, e.g. a single-guide RNA or a crRNA:tracrRNA complex, for targeting the wild-type CRISPR nuclease protein to a desired target DNA sequence based on complementarity between a portion of the guide RNA molecule or guide RNA complex and the target DNA sequence.

[0207] In the context of the invention, “maintained on-target editing activity” refers to the ability of an OMNI-50 variant to target a DNA target site that is targeted by a guide RNA molecule associated with, and thereby programming, the OMNI-50 variant. In some embodiments, the OMNI-50 variant maintains on-target editing activity of a DNA target at a percent editing level greater than or equal to the percent editing level of a wild-type OMNI-50 nuclease for the DNA target. In some embodiments, the OMNI-50 variant maintains on-target editing activity of a DNA target of at least 100%, 95%, 90%, 80%, 70%, 60%, 50%, 40%, or 30% the level of percent editing of a wild-type OMNI-50 nuclease for the DNA target.

[0208] For the foregoing embodiments, each embodiment disclosed herein is contemplated as being applicable to each of the other disclosed embodiment. For example, it is understood that any of the RNA molecules or compositions of the present invention may be utilized in any of the methods of the present invention.

[0209] As used herein, all headings are simply for organization and are not intended to limit the disclosure in any manner. The content of any individual section may be equally applicable to all sections.

[0210] Additional objects, advantages, and novel features of the present invention will become apparent to one ordinarily skilled in the art upon examination of the following examples, which are not intended to be limiting. Additionally, each of the various embodiments and aspects of the present invention as delineated hereinabove and as claimed in the claims section below finds experimental support in the following examples.

[0211] It is appreciated that certain features of the invention, which are, for clarity, described in the context of separate embodiments, may also be provided in combination in a single embodiment. Conversely, various features of the invention, which are, for brevity, described in the context of a single embodiment, may also be provided separately or in any suitable sub-combination or as suitable in any other described embodiment of the invention. Certain features described in the context of various embodiments are not to be considered essential features of those embodiments, unless the embodiment is inoperative without those elements.

[0212] Generally, the nomenclature used herein, and the laboratory procedures utilized in the present invention include molecular, biochemical, microbiological and recombinant DNA techniques. Such techniques are thoroughly explained in the literature. See, for example, Sambrook et al., “Molecular Cloning: A laboratory Manual” (1989); Ausubel, R. M. (Ed.), “Current Protocols in Molecular Biology” Volumes I-III (1994); Ausubel et al., “Current Protocols in Molecular Biology”, John Wiley and Sons, Baltimore, Maryland (1989); Perbal, “A Practical Guide to Molecular Cloning”, John Wiley & Sons, New York (1988); Watson et al., “Recombinant DNA”, Scientific American

Books, New York; Birren et al. (Eds.), “Genome Analysis: A Laboratory Manual Series”, Vols. 1-4, Cold Spring Harbor Laboratory Press, New York (1998); Methodologies as set forth in U.S. Pat. Nos. 4,666,828; 4,683,202; 4,801,531; 5,192,659 and 5,272,057; Cellis, J. E. (Ed.), “Cell Biology: A Laboratory Handbook”, Volumes I-III (1994); Freshney, “Culture of Animal Cells—A Manual of Basic Technique” Third Edition, Wiley-Liss, N. Y. (1994); Coligan J. E. (Ed.), “Current Protocols in Immunology” Volumes I-III (1994); Stites et al. (Eds.), “Basic and Clinical Immunology” (8th Edition), Appleton & Lange, Norwalk, CT (1994); Mishell and Shiigi (Eds.), “Strategies for Protein Purification and Characterization—A Laboratory Course Manual” CSHL Press (1996); Clokie and Kropinski (Eds.), “Bacteriophage Methods and Protocols”, Volume 1: Isolation, Characterization, and Interactions (2009), all of which are incorporated by reference. Other general references are provided throughout this document.

[0213] Examples are provided below to facilitate a more complete understanding of the invention. The following examples illustrate the exemplary modes of making and practicing the invention. However, the scope of the invention is not limited to specific embodiments disclosed in these Examples, which are for purposes of illustration only.

EXAMPLES

Example 1: Variant Selection

[0214] In order to select OMNI-50 nuclease variants with increased specificity to the target site (e.g. increased ratio between On-target cuts and Off-target cuts), amino acid substitutions were introduced into the open reading frame of the wild-type OMNI-50 sequence (SEQ ID NO: 1) as shown in Table 1.

TABLE 1

Summary of amino acid positions wild-type OMNI-50 nuclease and variants thereof. A blank box indicates the OMNI-50 variant has the same amino acid as the wild-type OMNI-50 nuclease in the indicated position.																		
Variant	Substitutions relative to	Amino acid identity by position																
name	WT OMNI-50	61	437	478	493	545	606	688	690	695	718	788	803	805	844	965	981	1036
WT OMNI-50	Not applicable	R	Y	R	A	Y	G	K	L	E	L	R	Q	L	L	K	V	K
Variant 3795	R478I; Y545H; Q803V; L805I			I		H							V	I				
Variant 3942	R61A; Y437W; R788K; L844N; V981M	A	W									K			N		M	
Variant 2902	G606P; L690V; E695V						P		V	V								
Variant 3954	R478T; A493G; L718C; K965V; K1036V			T	G						C					V		V
Variant 6918	Q803V												V					
Variant 6919	Y545H					H												
Variant 6920	L805I													I				
Variant 6864	R478I; Y545H; Q803V			I		H							V					
Variant 6865	R478I; Y545H			I		H												
Variant 6866	R478I; Q803V; L805I			I									V	I				
Variant 6867	Y545H; Q803V; L805I					H							V	I				
Variant 7085	R478H; Y545H; Q803V; L805I			H		H							V	I				
Variant 7086	R478V; Y545H; Q803V; L805I			V		H							V	I				

TABLE 1-continued

Summary of amino acid positions wild-type OMNI-50 nuclease and variants thereof. A blank box indicates the OMNI-50 variant has the same amino acid as the wild-type OMNI-50 nuclease in the indicated position.																	
Variant	Substitutions relative to	Amino acid identity by position															
		WT OMNI-50	61	437	478	493	545	606	688	690	695	718	788	803	805	844	965
Variant 7087	R478D; Y545H; Q803V; L805I				D		H							V	I		
Variant 7088	R478S; Y545H; Q803V; L805I				S		H							V	I		
Variant 7090	R478I; Y545C; Q803V; L805I				I		C							V	I		
Variant 7091	R478I; Y545D; Q803V; L805I				I		D							V	I		
Variant 7092	R478I; Y545T; Q803V; L805I				I		T							V	I		
Variant 7093	R478I; Y545F; Q803V; L805I				I		F							V	I		
Variant 7094	R478I; Y545A; Q803V; L805I				I		A							V	I		

TABLE 2

Guides Designed for discriminating SNPs and used for testing OMNI-50 nuclease variant activity and allele-specific editing.				
Target SNP	gRNA	Guide Sequence	Location	Mechanism
rs1683564	g62 REF 20 nt	GUCAAGCCCCA GAGGCCACA (SEQ ID NO: 29)	Downstream to 3'UTR	Excision, allele knock out
	g62 ALT 20 nt	GUCAAGCCCCA GAGGACACA (SEQ ID NO: 30)		
	sgRNA- 564DS-ref 22 nt	GUGUCAAGCCC CAGAGGCCACA (SEQ ID NO: 31)		
	sgRNA- 564DS-alt 22 nt	GUGUCAAGCCC CAGAGGACACA (SEQ ID NO: 32)	Intron 4	Excision, allele knock out
	g35 20 nt	AGUCCGGGCUG GGAGCGGGU (SEQ ID NO: 33)		
	sgRNA- Constant 22 nt	GCAGUCCGGGC UGGGAGCGGGU (SEQ ID NO: 34)		

Example 2: OMNI-50 Variants Depict Allele Specific Editing of ELANE Down-Stream Composition

[0215] Improved specificity of OMNI-50 variant nucleases relative to wild-type OMNI-50 nuclease was demonstrated by examining the ability of the nucleases to discriminate between the Alternative and Reference sequences of rs1683564 by utilizing g62alt and g62ref RNA guide sequences. See FIGS. 1-6C.

[0216] While editing with sgRNA-564DS-alt depicted high degree of specificity with wild-type OMNI-50 nuclease, sgRNA-564DS-ref showed lack of discrimination represented by the editing of both reference and alternative allele. Allele specificity was tested by excision with four OMNI-50 nuclease variants v3942, v2902, v3954 and v3795 in HSCs.

[0217] All variants of OMNI-50 revealed improved specificity for sgRNA-564DS-ref in comparison to OMNI-50 while maintaining sgRNA-564DS-alt allele discrimination determined by ddPCR (FIG. 4A). In addition to allele discrimination, one off-target to sgRNA-564DS-ref, sgRNA-564DS-alt and sg-constant were validated and showed high degree of variant fidelity in comparison to OMNI-50 characterized by <0.5% editing activity (FIG. 4B and FIG. 4C).

[0218] These results emphasize the relative high fidelity of OMNI-50 nuclease variants.

Example 3: Mutant Allele Knock-Out with a Novel CRISPR Nuclease Promotes Myelopoiesis in ELANE Neutropenia

[0219] Representative ELANE mutations and respective therapeutic strategies. Severe congenital neutropenia (SCN) is associated with numerous heterozygous mutations in ELANE gene (FIG. 7A). The current study presents a novel approach for removal of the mutated ELANE allele by targeting heterozygous sites of SNPs that are adjacent to the majority of ELANE-mediated SCN mutations, instead of independently editing each pathogenic mutation. Out of hundreds of potential SNPs in the ELANE gene, three SNPs retrieved from the healthy database 1000 Genomes Project Consortium were identified that are frequently heterozygous in the healthy population, termed herein: rs1683564, rs10414837 and rs3761005 (FIG. 7B upper panel and FIG. 10A). Analysis of the healthy and patient populations revealed similar heterozygosity frequency between the two populations in each of the SNPs. About 76-85% of each of the populations (healthy and patient, respectively) was heterozygous to at least one of the SNPs, indicating on the applicability of our strategy to more than about 75% of the SCN patient population. (FIG. 10B, Supplemental Table 1). For each of the three SNPs we developed an editing ribonucleoprotein (RNP) composition that includes two different guides and a novel optimized CRISPR/Cas9 nuclease termed OMNI Variant 3795. One guide (termed herein sgRNA constant) is common to all three compositions and cuts both ELANE alleles at intron 4. The second guide targets the heterozygous form of one of the three SNPs and therefore cuts only one allele. The composition which targets SNP rs1683564 excises exon 5 and the entire 3' UTR, causing the degradation of the destabilized mRNA transcript (FIG. 7B, I). The compositions which target either SNP rs10414837 or SNP rs3761005 lead to excision of most of

the coding region and the promotor, thus preventing the transcription of the mutated allele (FIG. 7B, II and III). The current study is focused on composition 1, targeting the rs1683564 SNP. The more prevalent form of rs1683564 SNP is cytosine and is referred to herein as reference (ref) allele. Adenosine is the less prevalence form of the SNP and is referred to herein as alternative (alt) allele. Prior to treatment, patient cells are genotyped to determine if the mutation and the SNP are on the same allele or different alleles in a process termed linkage determination (FIG. 11). If the pathogenic mutation is linked to the reference allele, a nuclease-guide RNA composition including a guide (sgRNA (ref)) that targets the cytosine form of the SNP is chosen (termed herein RNP(ref)). If the pathogenic mutation is linked to the alternative allele, a composition including a guide (sgRNA (alt)) that targets the adenosine form of the SNP is chosen (termed herein RNP(alt)). The guides differ by only one nucleotide and when used with the sgRNA constant result in the same editing outcome (FIG. 12).

[0220] Allele specificity and excision efficiency using OMNI Variant 3795 nuclease in a SNP-based knock-out strategy. Unlike most CRISPR-associated editing strategies that cut the target gene in both alleles, our approach is directed at removal of the mutated allele while keeping the wild-type functional allele intact. To demonstrate the feasibility of our mono-allelic editing strategy, HSCs heterozygous to SNP rs1683564, taken from healthy donors and SCN patients were excised using either RNP(ref) or RNP(alt) (according to their linkage) or left non-treated (NT). HSCs recovered for 3 days in CD34⁺ expansion media, then cultured for 7 days in the presence of IL-3, SCF, GM-CSF and G-CSF for proliferation and myeloid progenitor differentiation, and subsequently stimulated with G-CSF for further 7 days for neutrophil differentiation (FIG. 8A).

[0221] A fraction of cells was harvested at day 6 and 14 of differentiation for genomic DNA or RNA extraction. Allele specificity was determined by two competitive probes binding either the alternative allele or the reference allele. (For probes' specificity see Supplemental Methods and FIG. 19A and FIG. 19B). To test the RNP(ref) composition we used HSCs from P41 SCN patient (SCN-P41) harboring a mutation on the reference allele and HSCs from V3 healthy donor (HD-V3), both heterozygous to the reference form of the SNP. ddPCR revealed that editing using RNP(ref) was specific as only about 40% and about 20% of the reference allele remained intact in HSCs of HD-V3 and SCN-P41, respectively (FIG. 8B). The alternative allele, however, was not affected by the RNP(ref) composition as indicated by its high levels that were similar to those of non-treated cells (NT) (FIG. 8B). These results demonstrated that treatment with RNP(ref) composition leads to allele-specific editing.

[0222] Next, we evaluated excision efficiency at day 6 and 14 of neutrophil differentiation. Excision was determined by amplification of two regions in ELANE gene using two differently labeled probes, one for exon 1 that is not affected by the current excision strategy and a second probe for exon 5 that is degraded upon excision (FIG. 6B, I). The ratio between the signals of the two probes was translated to excision efficiency. Treatment with RNP(ref) resulted in about 13% excision in HSCs of HD-V3 and about 25% excision in HSCs of SCN-P41 (FIG. 8C). Given the high specificity of the nuclease (FIG. 8B), this excision correlated to about 50% of the cell population that has undergone excision at the reference allele in HSCs of SCN-P41. RNP

(ref) treatment also involved about 6% inversion events in HSCs of SCN-P41 as measured by EvaGreen staining (FIG. 13A and FIG. 13B). Of note, other experiments evaluating RNP(ref)-mediated excision levels in HSCs from healthy donors showed higher excision levels than reported above, that were comparable to those measured in HSCs of SCN-P41 (FIG. 14). In addition, NGS analysis of cDNA from SCN-P41 cells targeting exon 4-harboring mutation showed a 1:1 ratio between wild-type and mutated allele in non-treated cells. RNP(ref) treatment shifted the wild-type to mutated allele ratio to 3:1 by differentially reducing the mutated allele transcript, thereby enriching the wild-type allele (FIG. 8D).

[0223] To evaluate the specificity and efficiency of the RNP(alt) composition we used HSCs from P55 SCN patient (SCN-P55) harboring a mutation on the alternative allele and heterozygous to the alternative form of the SNP.

[0224] RNP(alt)-based treatment resulted in editing of about 90% of the alternative allele (only 10% remained intact), whereas the reference allele was kept intact at day 6 of differentiation (FIG. 8E). Excision levels in SCN-P55 HSCs were about 23% at day 6 and day 14 of neutrophil differentiation (FIG. 8F), indicating about 46% of the cell population has undergone excision at the alternative allele. RNP(alt) treatment resulted in 7.6% inversion events in SCN-P55 HSCs as measured by EvaGreen staining (FIG. 13C). Specificity and excision efficiency of RNP(alt)-edited HSCs of healthy donors was tested and found comparable to that obtained in patient-derived cells (FIG. 14). In addition, NGS analysis of cDNA from SCN-P55 cells targeting exon 5-harboring mutation showed enrichment of the wild-type allele (FIG. 8G). ELANE mRNA levels were decreased following excision in cells from SCN-P41 and SCN-P55 patients (FIG. 8H). In view of the increased WT:mutant allele ratio, obtained following excision (FIG. 8D and FIG. 8G), the reduced mRNA levels were mainly a result of the degradation of the mutated transcript.

[0225] Next, we confirmed RNP(alt)-based excision has occurred in a sub-population of HSCs (CD34⁺/CD90⁺ cells) that is considered essential for multilineage engraftment and hematopoietic reconstitution (FIGS. 15A-15C).

[0226] HSCs from both P41 and P55, edited using RNP(ref) and RNP(alt), respectively, were analyzed by Amplicon NGS to measure editing on off targets identified for constant guide (SgRNA(constant)), reference guide (SgRNA(ref)) and alternative guide (SgRNA(alt)).

[0227] Specifically, an unbiased survey (GUIDE-seq) of whole-genome off-target cleavage using the Variant 3795 nuclease and each of the constant guide (SgRNA(constant)), reference guide (SgRNA(ref)) and alternative guide (SgRNA(alt)), resulted in no identified off-targets (≤ 4 mismatches) (FIGS. 20A-20C).

[0228] In addition, in-silico off-target analysis was performed for each of the guides and identified a few potential off-targets. None of these off-targets were validated by a rhAmpSeq analysis done on edited HSCs from patients SCN-P41 and SCN-P55 (FIG. 20D), demonstrating the high fidelity of the novel nuclease compositions.

[0229] Taken together, the results provided above present an active, highly accurate nuclease that can target the mutant allele while preserving the wild-type functional allele intact.

[0230] OMNI Variant 3795 facilitated editing boosts neutrophil differentiation and maturation in vitro. To demonstrate the functional outcome of our SNP-based single-

allelic editing approach we evaluated neutrophil differentiation and maturation capacities of RNP(ref)-edited and non-treated healthy and patient-derived HSCs in vitro, at day 14 of differentiation. Flow cytometric analysis showed that about 74% of the HD-V3 HSCs, subjected to the differentiation protocol, differentiated into neutrophils (CD66b⁺; two right quarters (Q2+Q3) of the dot plot), compared to only 36% of the SCN-P41-derived HSCs. To the contrary, only about 15% of HD-V3 HSCs differentiated into monocytes (CD14⁺/CD66b⁻; upper left quarter (Q1) of the dot plot) compared to 38% of the SCN-P41-derived HSCs. These observations are consistent with the characteristic hematopoietic defect in SCN patients. (FIGS. 9A-9C; NT: HD-V3 versus SCN-P41). SCN-P41-derived HSCs treated with RNP(ref) depicted a 74% increase in neutrophils (CD66b⁺ cells) and a 2 fold reduction in the monocytic subset (CD14⁺/CD66b⁻) (FIGS. 9A-9C; SCN-P41: NT versus RNP(ref)). Similar increase in neutrophil count was observed in SCN-P41-derived edited HSCs by flow cytometric analysis of CD11b⁺/CD15⁺ cells (FIGS. 16A and 16B). RNP(ref)-mediated editing in HSCs of HD-V3 did not affect the monocytic and neutrophilic subsets, supporting the safety of this composition (FIGS. 9A-9C; HD-V3: NT versus RNP(ref)).

[0231] After demonstrating our allele-specific editing approach significantly improved the cellular abnormalities associated with SCN, we next assessed neutrophilic functions in RNP(ref)-treated and non-treated healthy (V3) and patient (P41) HSC-derived neutrophils. In vitro phagocytic capacity was tested by measurement of phagosomal uptake of zymosan green particles by neutrophils from the different groups. Flow cytometric analysis revealed equivalent levels of phagocytosis in RNP(ref)-treated and non-treated neutrophils from both HD-V3 and SCN-P41 (FIG. 9D). In addition, anti-bacterial killing capacity was examined by incubating neutrophils derived from non-treated healthy HSCs (HD-V3, NT) and RNP(ref)-treated patient HSCs (SCN-P41, RNP(ref)) with E-coli bacteria, expressing bacterial luciferase gene, and tracking real time changes in light emission, expressed as relative light units (RLUs). The two groups were compared with bacteria only (E-coli control without neutrophils). Healthy and RNP(ref)-treated patient derived neutrophils exhibited efficient bacterial killing as indicated by a 23-25% decrease in bacterial unit (RLU) in comparison to bacteria only control. Experiment was performed with 50,000 and 100,000 HSCs (FIG. 9E).

[0232] An experiment using the RNP(ref) composition was conducted on cells from another patient (SCN-P42) and depicted similar excision, differentiation and functional results, including histological staining demonstrating the restoration of neutrophil differentiation (FIGS. 21A-21J). Thus, specific excision of ELANE mutated allele by RNP(ref) composition ameliorated the aberrant phenotype of attenuated differentiation towards neutrophils and preserved neutrophil basic functions. A similar analysis was performed on the RNP(alt) composition. Flow cytometric analysis showed about 83% of HD-V4-derived HSCs differentiated into neutrophils (CD66b⁺; two right quarters (Q2+Q3) of the dot plot) and about 7% differentiated into monocytes (CD14⁺/CD66b⁻; upper left quarter (Q1) of the dot plot). SCN-P55-derived HSCs showed lower differentiation towards neutrophils (about 53%) and higher differentiation to monocytes (about 29%) (FIGS. 9F-9H; NT: HD-V4 versus SCN-P55), representing a typical SCN hematopoietic

defect. SCN-P55-derived HSCs treated with RNP(alt) presented a 1.5 fold increase in neutrophils (CD66b⁺ cells) and a 3 fold reduction in the monocytic subset (CD14⁺/CD66b⁻) (FIGS. 9F-9H; SCN-P41: NT versus RNP(alt)). A similar increase in neutrophil subset was observed in SCN-P55-derived edited HSCs by flow cytometric analysis of CD11b⁺/CD15⁺ cells (FIGS. 16C and 16D). Diff-Quik staining of SCN-P55-derived HSCs treated with RNP(alt) or electroporated without a nuclease composition (SCN-P55 Mock) revealed higher numbers of cells with classical polymorphonuclear neutrophilic morphology in the RNP(alt)-treated group compared with Mock group (FIG. 9I). Flow cytometric analysis of zymosan green particles revealed slightly higher levels of phagocytosis in RNP(alt)-treated neutrophils compared to non-treated neutrophils from SCN-P55 (FIG. 9J). In addition, healthy non-treated (HD-V4, NT) and patient RNP(alt)-treated (SCN-P55, RNP(alt)) neutrophils exhibited efficient bacterial killing as indicated by a significant 30% decrease in bacterial unit (RLU) in comparison to bacteria only control (FIG. 9K).

[0233] An experiment using the RNP(alt) composition was conducted on cells from another patient (SCN-P12) and depicted similar excision, differentiation and functional results (FIGS. 22A-22I). Moreover, another experiment performed on patient cells (SCN-P56) harboring a mutation on exon 2, located upstream to the mutations found in previous patients (exons 4 and 5), showed similar results (FIGS. 23A-23I). The results described herein indicate a single-allelic knock-out of ELANE mutated allele, using RNP(ref) and RNP(alt) novel nuclease compositions, is safe and effective both in boosting neutrophil differentiation and in maintaining essential neutrophilic core functions.

[0234] OMNI Variant 3795 showed high fidelity without traceable off targets. This is the first report demonstrating specific mono-allelic gene editing, which is not mediated by a gain of PAM sequence. This unique feature of our novel nuclease could be implemented in various indications that are dominant, dominant negative, and compound heterozygous, covering most genetic disorders that other technologies cannot address.

[0235] OMNI Variant 3795 composition showed excision efficiency of about 25%. Notably, given the high allele specificity of this nuclease composition, it is estimated that about 50% of the cell population had undergone excision at the mutant allele. With respect to the functional aspects, ELANE mono-allelic excision significantly enhanced neutrophil differentiation in-vitro. It also reduced the aberrant numbers of monocytes, consistent with the hematopoietic defect of SCN patients. Excised neutrophils showed normal phagocytic and bacterial killing capacities, indicating the editing was both effective and safe.

[0236] Thus, this Example presents a novel CRISPR/Cas9-based strategy of specific mono-allelic excision. Such a strategy was found to be efficient, functional, accurate and safe.

Example 3: Materials & Methods

[0237] Human HSC isolation: HSCs were isolated from SCN patients' bone marrow and healthy donors' mobilized peripheral blood.

[0238] CRISPR/Cas9 OMNI Variant 3795 ELANE gene editing: Ribonucleic protein (RNP) system at a molar ratio of 1:2.5 (OMNI Variant 3795 nuclease:sgRNA) was used.

Human CD34⁺ cells were electroporated using the CA-137 program (Lonza 4D, Nucleofector™).

[0239] Digital Droplet PCR (excision, allele specificity): Excision and allele specificity were measured using Digital Droplet PCR™ on genomic DNA. For excision reaction, amplification of two regions in ELANE gene, exon 1 and exon 5 was performed, using two different probes FAM(X1) and HEX(X1), respectively. The ratio between the probe signals was translated to excision efficiency. For allele specificity, a FAM probe (binding the alternative allele) and a HEX probe (binding the reference allele) were used (FAM+HEX). The ratio between the two probes was normalized to endogenous genes.

[0240] Assessment of mutated:wild-type allele ratio: cDNAs were mapped using next-generation sequencing (NGS) targeting exon 4 and 5 harboring S126L and R220Q mutations in patients 41 and 55, respectively. Relative ratio of mutated:wild-type alleles in treated cells was calculated and compared to that of non-treated cells.

[0241] Differentiation assay: Edited and non-treated HSCs were subjected to a differentiation protocol adopted from Nasri et al. On day 14, cells were analyzed by flow cytometry for monocytic (CD14⁺/CD66b⁻) and neutrophilic (CD66b⁺) subsets.

[0242] Bacterial killing assay: Day 13 differentiated HSCs were evaluated for their bacterial killing capacity as described in J. T. Atosuo. Relative light units (RLUs) were measured over 5 hrs. Last time point presented is when RLU levels reached plateau. Wells without differentiated HSCs (*E. coli* only) and with phagocytosis inhibitor, Cytochalasin D, (data not shown) served as controls.

[0243] Phagocytosis assay: Phagocytosis capacity was evaluated using the EZCell™ Phagocytosis Assay Kit (Green Zymosan), (BioVision, Cat no. K397). Cells were analyzed by flow cytometry for internalization of opsonized fluorescent Zymosan Green particles.

[0244] Further details on FACS antibodies, probes, sequences, assays, protocols, and additional methods are provided in Supplemental Methods.

Example 3: Supplemental Methods

[0245] Human SCN Patient HSC isolation: Three to 6 mls of freshly collected bone marrow was shipped overnight at ambient temperature. Hematopoietic stem and progenitor cells, HSPC's, were initially enriched using RosetteSep Human Bone Marrow Progenitor Cell Pre-Enrichment Cocktail, (Cat. No. 15027) and Lymphoprep (Cat. no. 07801) according to manufacturer's protocol. The HSC enriched cell population was expanded by culturing for 4 days in CD34⁺ expansion media (StemSpan SFEMII media (Cat. no. 09655) supplemented with 1% Penn Strep (Cat. no. 03-031-1B, Biological Industries), 1x StemSpan CD34 Expansion Supplement (10x) (Cat. no. 02691), and 1.0 μM UM729 (Cat. no. 72332), at 37° C. 5% CO₂. After expansion, CD34⁺ cells were further enriched using EasySep Human CD34 Positive Selection Kit II (Cat. no. 17856) according to manufacturer's protocol. Enriched CD34⁺ cells were cryopreserved at 1x10⁶ cells/ml in Cryostor CS10 (cat. no. 07931). Cells were stored in liquid nitrogen, vapor phase. All catalog numbers refer to materials from StemCell Technologies unless indicated otherwise.

[0246] Human healthy donor HSC isolation: Cryopreserved healthy human CD34⁺ progenitor cells from mobilized peripheral blood were obtained from Lonza (Cat no.

4Y-101C). Cells were suspended in CD34⁺ expansion media at 50,000 cells/ml and expanded for 4 days at 37° C., 5% CO₂ prior to electroporation.

[0247] CRISPR/Cas9 OMNI Variant 3795 ELANE gene editing: Editing of HSCs was carried out using a ribonucleic protein (RNP) system at a molar ratio of 1:2.5 (nuclease: sgRNA), including 17 µg nuclease and 262 pmol of each guide. Nuclease and sgRNA complex were incubate at 25° C. for 10 minutes. Human CD34⁺ cells were washed once with PBS. 2×10⁵ CD34⁺ cells were suspended into 20 µl of P3 electroporation buffer (Lonza P3 kit S) and were added to RNPs mix. After electroporation, using the CA-137 program (Lonza 4D, Nucleofector™), the cells were transferred to pre-warmed CD34⁺ expansion media at a concentration of 1.25.0×10⁵ cells/ml. Guides were manufactured by Agilent. Guide sequences are summarized in the table below:

sgRNA name	Guide sequence (scaffold + spacer)
sgRef	GUGUCAAGCCCCAGAGGCCACAgUUUGAGAGUUA UGAAAAUGACGAGUUCAAAUAAAAUUUAUUCAA ACCGCCUAAUUUAUAGGCCGAGAUUUCUGCUUU (SEQ ID NO: 35)
sgAlt	GUGUCAAGCCCCAGAGGACACAgUUUGAGAGUUA UGAAAAUGACGAGUUCAAAUAAAAUUUAUUCAA ACCGCCUAAUUUAUAGGCCGAGAUUUCUGCUUU (SEQ ID NO: 36)
sgConstant	GCAGUCCGGGCGGGAGCGGGUGUUUGAGAGUUA UGAAAAUGACGAGUUCAAAUAAAAUUUAUUCAA ACCGCCUAAUUUAUAGGCCGAGAUUUCUGCUUU (SEQ ID NO: 37)

[0248] Digital Droplet PCR for percentage excision and allele specificity: Percentage excision and allele specificity were measured using Digital Droplet PCR™ (ddPCR™, Bio-Rad, Hercules, CA, USA) on genomic DNA that was extracted using QIAamp DNA Micro Kit, Qiagen (Cat no. 56304). According to manufacturer’s protocol.

[0249] ddPCR reaction contained 1× ddPCR Supermix for probe without dUTP (#1863024), 25-100 ng of digested DNA using HindIII (diluted in X1 Cutsmart Buffer to 4 U/µL) and suitable primers/probes. For excision reaction, amplification of two regions in ELANE gene, exon 1 and exon 5 was performed, using two different probes labeled with FAM (X1) and HEX (X1), respectively. The ratio between the HEX and the FAM signals was translated to excision efficiency.

[0250] For allele specificity, two competitive probes: a FAM probe, which binds the alternative allele, and a HEX probe, which binds the reference allele were used (FAM+HEX). The ratio between the concentrations of the two in heterozygote non-treated cell is 1, which was normalized to the endogenous genes RPP30 and STAT1 for each gDNA sample. Reaction total volume was 22 µL. The binding of each probe to DNA extracted from healthy donor cells that were homozygous to either the reference or alternative forms of the SNP was measured by ddPCR, confirming the probes do not cross react, (FIG. 19A). Moreover, healthy donor cells homozygous to the reference form of the SNP depicted efficient excision when treated with RNP(ref) composition, compared to treatment with RNP(alt) composition

that resulted in excision levels comparable to non-treated cells. This further demonstrates the specific targeting of the sgRNAs (FIG. 19B).

[0251] Genomic DNA in the ddPCR mixture was partitioned into individual droplets using QX100 Droplet Generator, transferred to a 96-deep well PCR plate and amplified in a Bio-Rad PCR thermocycler. Bio-Rad Droplet Reader and QuantaSoft Software were used to read and analyzed the experiment following manufacturer’s guidelines (Bio-Rad).

[0252] The primers and probes were manufactured by Bio-Rad and are detailed in the table below:

Probe name	Catalog number
ELANE ddPCR assay Exon1 (FAM)	dHsaCNS328407057
ELANE ddPCR assay Exon5 (HEX)	qHsaCEP0055470
rs1683564 Editing (FAM + HEX) ddPCR assay	dHsaMDS873573221
RPP30	dHsaMDS117591774
STAT1	dHsaCNS850507320

[0253] Assessment of mutated:wild-type allele ratio: cDNAs from HSCs treated with either RNP(ref) or RNP(alt) were mapped using next-generation sequencing (NGS) targeting exon 4 and 5 harboring S126L and R220Q mutations in patients 41 and 55, respectively. The raw FASTQ files were analyzed, and BAM files (text-based format for storing biological sequences) were generated using FASTQ to BAM script. The relative ratio of the mutated allele to the wild-type allele was calculated and compared to the non-treated cells in both patients. Primers are detailed in the table below:

Mutation	Sequence
S126L (P41)	TCGTCGGCAGCGTCAGATGTGTATAAGA GACAGNNNTACGACCCGTAACCTTGCT GTCTCGTGGGCTCGGAGATGTGTATAAG AGACAGNNNCGGAGCGTGGATGATAGA G
R220Q (P55)	TCGTCGGCAGCGTCAGATGTGTATAAGA GACAGNNNTCGCAGTCCAGCTTCCCCAC GTCTCGTGGGCTCGGAGATGTGTATAAG AGACAGNNNACAGCCAAGGAGCATCAAA C

[0254] Differentiation assay: Edited and non-treated HSCs were allowed to recover for 3 days in CW34⁺ expansion media and were subjected to a differentiation protocol adopted from Nasri et al.²³ In brief, HSCs were cultured for 7 days in RPMI (Cat. no 11875093, Gibco™) supplemented with 1% Glutamax (Cat. no 35050061, Gibco™), 10% FBS (Cat. no 04-001-1A, Biological Industries), 5 ng/ml IL-3 (Cat. no 200-03), SCF (Cat. no 300-07), GM-CSF (Cat. no 300-03) & 10 ng/ml G-CSF (Cat. no 300-23), all from PeproTech, for proliferation and myeloid progenitor differentiation followed by a 7-day culture in RPMI, 1% Glutamax, 10% FBS, 1% Penn Strep (Cat. no 03-031-1B, Biological Industries), 10 ng/ml G-CSF for neutrophil differentiation and maturation. On day 14 cells were ana-

lyzed by flow cytometry for monocytic (CD14⁺/CD66b⁻) and neutrophilic (CD66b⁺) subsets. CD66b anti-human, Pacific Blue (Cat No. 305112, Biolegend) and CD14, anti-human, APC (Cat No. 130-110-520, Miltenyi Biotec) were used.

[0255] Cytospin staining: 8×10⁴ HSCs at day 15 of differentiation were spun onto Cytoslide microscope slides (ThermoFisher) using Cytospin 4 low speed cytocentrifuge (Thermo Scientific) and stained with Diff-Quick staining system (MilliporeSigma) according to manufacturer's recommendations. Microphotographs were taken on LEITZ LABORLUX S polarizing light microscope at 400× magnification using Nikon DSLR digital camera.

[0256] Bacterial killing assay: Day 13 differentiated HSCs (subjected to a differentiation protocol adopted from Nasri et al.) from healthy donors and SCN patients (edited with either RNP(ref), RNP(alt) or non-treated), were evaluated for their bacterial killing capacity as described in J. T. Atosuo. Briefly, 100,000 differentiated HSCs were incubated in the presence of 200,000 Luciferase expressing bacterial cells, pAKLUX2, per well (Addgene, Cat No. 14080). Cells were cultured in 200 ul HBSS++, 10% FBS at 37° C. At 30-minute intervals luminescence was measured by transferring the plate to a Luminometer (Berthold CentroXS3 LB960) and measuring luminescence for 0.5 sec per well). A real time change in light emission, relative light units (RLUs), was measured over 5 hrs. Last time point presented is when RLU levels reached plateau. Wells without differentiated HSCs (*E. coli* only) and with 10 ng/ml phagocytosis inhibitor (data not shown), Cytochalasin D (Santa Cruz Biotec, Cat No. sc-20144) served as controls.

[0257] Phagocytosis assay: Phagocytosis capacity was evaluated using the EZCell™ Phagocytosis Assay Kit (Green Zymosan), (BioVision, Cat no. K397 according to manufacturer's protocol). Day 14 differentiated HSCs (subjected to a differentiation protocol adopted from Nasri et al.) from healthy donors and SCN patients (edited with either RNP(ref), RNP(alt) or non-treated), were resuspended in HBSS++/10% FBS (0.5×10⁶ cells/ml) and incubated for 1.5 hours at 37° C. in the presence of 5 ml opsonized Alexa Fluor 488-conjugated zymosan particles per 200 ml suspended cells. As a negative control, cells were incubated with 10 ng/ml of the phagocytosis inhibitor, Cytochalasin D (Santa Cruz Biotec, Cat No. sc-201442) 1 hour prior and during incubation with Zymosan Green reagent. Cells were then washed and incubated in a quencher solution, based on kit's instructions, to remove fluorescence from particles that were not internalized. Cells were then analyzed by flow cytometry for internalization of opsonized fluorescent Zymosan Green particles.

[0258] Inversions: Inversion events were detected and quantified by a Droplet Digital PCR (ddPCR) mutation assay. First, a perfect inversion was mimicked using a SnapGene software and verified by NGS. Then, total inversion events were quantified by ddPCR using EvaGreen dye, a fluorescent DNA-binding dye that binds dsDNA (BIO-RAD, catalog number 186-4034, according to manufacturer's protocol). Specific primers were designed to amplify inverted variations of the excised fragment (See illustration in FIG. 11). Fluorescent signals were normalized to amplification of ELANE exon 1 region that was not affected by excision (performed by two different sets of primers. Averaged normalized data is presented). Primers are detailed in the table below:

Primer name	Sequence
Exon 4 primer	ACGTCCTGCACTCTCGTGAGG
Exon 5 primer (inverted)	TTCAGGCTCCACCCAGTTTGTC
Exon 1 primers	Mix3: GCACAGGGCTATAAGAGGAGC GCGGGAGGTTGGACTCAAAA Mix6: GGGAGAGGAAGTGGAGGGC GAGGGTCATGGTGGGGCT

[0259] Excision levels in Long Term HSC population: HSCs of two healthy donors (MLP1 (3055934); heterozygous to the alternative form of the SNP and MLP2 (3055940), homozygous to the alternative form of the SNP), isolated from leukopaks purchased from AllCells, were edited a day after thaw according to 'CRISPR/Cas9 OMNI Variant 3795 ELANE gene editing' section above, with minor changes. Cell number was 2M cells/electroporation. Upscale of guides and nuclease was performed accordingly. A molar ratio of 1:2.5 (nuclease:sgRNA) was used including 85 µg nuclease and 1310 pmol of each guide. Nucleofection was performed in P3 nucleofection solution (Lonza) and Lonza 4D-Nucleofector™ X Kit L (program CA-137). Three days after editing, cells were sorted in a FACS ARIA™ II SORP Flow Cytometer Cell (BD), using CD90-APC-Vio770, human (130-114-863 Milteny). Sorted cells were incubated for 7 days in a proliferation medium according to Nasri et al. Excision levels of CD90⁺, CD90⁻ and total population were evaluated using ddPCR as described in ddPCR section.

[0260] Mutation-SNP linkage: First, ELANE mutation and possible SNPs were identified in cells from SCN patients by targeted short read NGS. Then, a part of the gene encompassing both the mutation and the SNP was amplified by a PCR reaction with linkage primers and cloned into bacteria. Each clone bore an amplicon from one allele. A plasmid of multiple clones was Sanger sequenced (with T7 and SP6 primers) for the mutation and the SNP regions. If the mutation and the SNP were in cis, they were found at the same clone, if in trans, the mutation and SNP were found in different clones. Primers are detailed in the table below:

Primer name	Sequence
rs10414837	FP : TCGTCGGCAGCGTCAGATGTGTATAAGAGA CAGNNNGGTGGGCTCCTCAGTGACTCT RP : GTCTCGTGGGCTCGGAGATGTGTATAAGAG ACAGNNNGGAATTCCAGCCTGACCAA
rs3761005	FP : TCGTCGGCAGCGTCAGATGTGTATAAGAGA CAGNNNGAGTGAGGACCAAGCCTGAG RP : GTCTCGTGGGCTCGGAGATGTGTATAAGAG ACAGNNNAGGGCCATTGTCTCCCTAAC
rs1683564	FP : TCGTCGGCAGCGTCAGATGTGTATAA GAGACAGNNNTCTGTACCTCCCTCTCT

-continued

Primer name	Sequence
Linkage	RP: GTCTCGTGGGCTCGGAGATGTGTATA AGAGACAGNNNTTTAGGAGGGGCCACTG A
	FP: GAGGGTCATCATCACTGCCC
	RP: GCCAGACTCACACCAGAGTCGACAAG T
T7	TAATACGACTCACTATAGGG
SP6	ATTTAGGTGACACTATAG
R220Q	FP: TCGTCGGCAGCGTCAGATGTGTATAAGAGA CAGNNNTCGCAGTCCAGCTTCCCCAC
mutation	RP: GTCTCGTGGGCTCGGAGATGTGTATAAGAG ACAGNNNACAGCCAAGGAGCATCAAAC
S126L	FP: TCGTCGGCAGCGTCAGATGTGTATAAGAGA CAGNNNGAGGGTCATCATCACTGCCC
mutation	RP: GTCTCGTGGGCTCGGAGATGTGTATAAGAG ACAGNNNAGTCCGGGCTGGGAGCGGGT

[0261] Heterozygosity frequency of SNPs in the healthy and patient populations: Variant call files encompassing the ELANE gene region (± 3 kb of ELANE gene) were down-

loaded from the 1000 Genomes Project Consortium (phase 3) using the Data Slicer tool and analyzed in the R statistical computing environment. 3501 genotypes were available from 3501 individuals. Familial relationship was omitted from the analysis, which resulted in 2407 genotypes from unrelated individuals. The allele frequency for all common polymorphism ($>1\%$ MAF) was calculated. Three SNPs were chosen (rs3761005, rs1683564, and rs10414837 polymorphisms) to optimize the population coverage for allele-specific ELANE knock-out. The percentage of the population being heterozygous for at least one of the three chosen SNPs was calculated.

[0262] 53 patients' samples were sequenced for the pathogenic mutation and the three chosen SNPs. A total of 46 bone marrow samples and 7 iPSC lines were used. 44 of the samples were from SCN patients and nine of them were from patients with Cyclic Neutropenia. Heterozygosity frequency of each of the three chosen SNPs and the percentage of the population being heterozygous for at least one of them was calculated.

[0263] Statistical methods: The two-sample T-test for independent samples or the ANOVA model, as appropriate, was applied for testing the statistical significance of the difference in continuous variables between treatment groups. The two-ways Analysis of Variance with repeated measurements was used to analyze killing assays. Chi-square test or Fisher's Exact test, as appropriate, was applied to test the statistical significance of the difference in heterozygosity between healthy and patient populations. All tests were two-tailed, and a p-value of 5% or less was considered statistically significant. The data was analyzed using Prism software (GraphPad version 9.0.2).

SUPPLEMENTAL TABLE 1

Patient #	Diagnosis	Source	ELANE mutation	Location of mutation	rs10414837	rs3761005	rs1683564
1	SCN	Bone Marrow	S126L	4	CC	TA	CC
2	SCN	Bone Marrow	G210V	5	CT	TT	CC
3	SCN	Bone Marrow	C555	2	CC	TA	CC
4	SCN	Bone Marrow	G221ter	5	CC	TA	AA
5	SCN	Bone Marrow	D230fs	5	CC	AA	CC
6	SCN	Bone Marrow	A57T	2	CT	TA	AA
7	SCN	Bone Marrow	R103P	3	CC	TT	AA
8	SCN	Bone Marrow	T128del	4	CT	AA	CA
9	SCN	Bone Marrow	V83D	3	CT	TT	AA
10	SCN	Bone Marrow	L84P	3	CT	TA	CA
11	SCN	Bone Marrow	L8RP	3	CT	TT	CA
12	SCN	Bone Marrow	M154R	4	CC	TT	CA
13	SCN	Bone Marrow	G221ter	5	CT	TT	AA
14	SCN	Bone Marrow	G203R	5	CT	TA	CA
15	SCN	Bone Marrow	G214R	5	CC	AA	CA
16	SCN	Bone Marrow	P139L	4	CC	TA	CC
17	SCN	Bone Marrow	Y228tr	5	CT	TT	AA
18	SCN	Bone Marrow	P139L	4	CC	TA	CA
19	SCN	Bone Marrow	W156R	4	CT	TA	CC
20	SCN	Bone Marrow	P139L	4	CC	AA	CA
21	CyN	iPSC	P139L	4	CC	TT	CC
22	SCN	iPSC	I120N	3	CT	TT	AA
23	SCN	iPSC	G214R	5	CC	TA	CC
24	CyN	iPSC	S46F	2	CC	TA	CA
25	SCN	Bone Marrow	IVS4 + 5G > A	INT4	CC	TT	CA
26	SCN	Bone Marrow	S126L	4	CC	TA	CA
27	SCN	Bone Marrow	R103L	3	CC	TA	CA
28	SCN	Bone Marrow	M66R	2	TT	TT	CC
29	SCN	Bone Marrow	C20BX	5	CT	TA	CC
30	SCN	Bone Marrow	M1R	1	CC	TT	CA
31	SCN	Bone Marrow	P234fs	5	CC	AA	CC

-continued

SUPPLEMENTAL TABLE 1

Patient #	Diagnosis	Source	ELANE mutation	Location of mutation	rs10414837	rs3761005	rs1683564
32	SCN	Bone Marrow	160T	2	CC	TT	CA
33	CyN	iPSC	IVS4 + 1 sd	INT4	CT	TT	CC
34	SCN	iPSC	P139L	4	CT	TT	AA
35	SCN	iPSC	R191S	4	TT	TT	CC
41	SCN	Bone Marrow	S126L	4	CC	AA	CA
42	SCN	Bone Marrow	S126L	4	CC	AA	CA
43	CyN	Bone Marrow	IVS4 + 5G > A	INT4	CC	TT	CC
44	CyN	Bone Marrow	IVS4 + 1 G > A	INT4	CT	TA	CA
45	CyN	Bone Marrow	R220Q	5	CT	TT	CC
46	CyN	Bone Marrow	IVS4 + 3A > T	INT4	CC	TA	CC
47	CyN	Bone Marrow	W241ter	5	CT	AA	CC
48	SCN	Bone Marrow	S126L	4	CT	TA	CC
49	SCN	Bone Marrow	S126L	4	CT	TA	CC
50	SCN	Bone Marrow	P139L	4	CT	TA	CC
51	SCN	Bone Marrow	S46F	2	CC	TA	CC
52	SCN	Bone Marrow	IVS3 - 8	INT3	CT	TA	CC
53	SCN	Bone Marrow	P139L	4	CC	TA	CA
54	SCN	Bone Marrow	R220Q	5	CC	TA	CA
55	SCN	Bone Marrow	R220Q	5	CT	TA	CA
56	SCN	Bone Marrow	A57V	2	CT	AA	CA
57	SCN	Bone Marrow	R191S	4	TT	TT	CC
58	CyN	Bone Marrow	S126L	4	CC	AA	CA

[0264] Supplemental Table 1. Coverage of the three SNPs by the patient population. Sequencing results of the three SNPs in ELANE gene in samples obtained from SCN and Cyclic Neutropenia (CyN) patients. The pathogenic mutations in ELANE gene were also verified by sequencing. Green cells depict heterozygous SNPs.

Example 4: Contribution of Each of the Variant 3795 Mutations to the Specificity and Fidelity of the Nuclease

[0265] Variant 3795 is a specificity variant of CRISPR-based nuclease with extremely low off-target activity (<0.05%) and an ability to discriminate between two alleles of the same gene which have a single mismatch between them. The variant includes four mutations: R478I, Y545H, Q803V and L805I. To investigate the effect and contribution to the activity and specificity of each mutation found in OMNI-50 V3795, three out of the four single mutant variants were expressed and purified, as well as the four triple mutant variants, in which one of the four mutations is missing in each variant (Table A). RNP complexes were prepared using WT OMNI-50 nuclease, the V3795 variant, and the seven variants by assembly with sgRNA g62-Ref. These RNP complexes were electroporated into LCL cells that are heterozygote for the target of the guide sequence (Table B). The editing levels of the targeted allele (REF) and non-targeted allele (ALT) were measured by NGS. As can be seen in FIGS. 17A-17B, the discrimination towards the targeted allele (REF) of V3795 is contributed mostly by the R478I mutation (Table E). Only variants containing the R478I mutation (e.g. V6864, V6865, V6866, V3795) showed clear preference toward the targeted allele, while removing this mutation abolished the discrimination (e.g. V6867).

[0266] The specificity effect was validated by measuring the editing levels at two known off-target sites of g62 by NGS (g62OT1 and g62OT2, Table C). OMNI-50 V3795 has extremely low levels of editing at these sites in comparison

to WT OMNI-50 nuclease (FIGS. 18A-18B, Table F). In all the three single mutants lower than WT editing levels of the off-target sites were observed, indicating a contribution of the three mutations to the reduction in off-targets. However, the Y545H mutation had the most significant reduction in the off-target editing (OT1 0.5% and OT2 0.04%, Table F). All of the triple variants also exhibited extremely low off-target activity, similar to the V3795 variant nuclease. Remarkably, when the Y545H mutation was excluded from the set of mutations (e.g. V6866) the off-target editing slightly increased (OT1 1.90% OT2 0.2%, FIGS. 18A-18B, Table F). Altogether, these results led to the conclusion that the V3795 variant nuclease's high level of specificity is attributed to the R478I and Y545H mutations.

[0267] Once it was established that R478I and Y545H are responsible for the observed specificity of the V3795 variant nuclease, the effect of other substitutions was tested. For each of these positions, RNPs were generated with the g62-Ref sgRNA and several variant nucleases containing substitutions from different biochemical groups in the background of the other three mutations (Table A). The effect on discrimination and off-targets was measured as above (FIGS. 17A-17B and 18A-18B).

[0268] Regarding position 478: Charged amino acids, such as aspartic Acid (D) in variant V7087, abolishes the discrimination and increases the off-target editing (FIG. 18B). A similar effect on discrimination is seen for arginine (R, the reference amino acid) in the triple variant V6867 (FIGS. 17A-17B). However, large hydrophobic amino acids such as valine (V) in V7086 as well as isoleucine (I) in V3795 retain high discrimination and reduced off-targets. An aromatic amino acid such as histidine (H) in V7085 and a polar uncharged amino acid such as serine (S) in V7088 also supports a similar phenotype as OMNI-50 V3795.

[0269] Regarding position 545: Substitution to amino acids such as phenylalanine (F) in V7093, as well as histidine (H) in V3795, retain high discrimination and low off-targets similar to the variant V3795 (FIGS. 17A-17B and

18A-18B). Alanine (A) is a small hydrophobic residue and is used as a substitute in in V7094, which also showed a similar phenotype as V3795 (FIGS. 17B and 18B).

Example 4: Materials & Methods

HTP Protein Expression and Purification

[0270] Wild-type OMNI-50 nuclease and its variants were cultured in auto-induction TB media at 37° C., 350 rpm agitation for 3.5 hours, and then shifted to 18° C. for 17-20 hours. Cells were harvested by centrifugation at 4000×g and stored at −80° C. OMNI-50 variants and WT OMNI-50 cell pellets were thawed and incubated in lysis buffer for 30 minutes. Crude lysate was clarified by centrifugation at 4000×g for 1 hour, 4° C. Cleared protein lysates were incubated with Sepharose 6 Ni-NTA resin (Cytiva). Protein-bound Ni-NTA resin was loaded on a 96 Filter well plate and washed with buffer (HEPES 20 mM, NaCl 0.6 M, Imidazole

cuvette immediately after electroporation. Cells were incubated for 2-4 days (37° C., 5% CO₂).

NGS Analysis

[0272] At 72 hours, cells were harvested and their genomic DNA content was used in a PCR reaction which amplified the corresponding putative genomic targets. Amplicons were subjected to next-generation sequencing (NGS) and the resulting sequences were then used to calculate the percentage of editing events at each target site. Short insertions or deletions (indels) around the cut site are the typical outcome of DNA repair following nuclease induced DNA cleavage. The calculation of percent editing was therefore deduced from the fraction of indel-containing sequences within each amplicon.

TABLE A

CRISPR Nucleases			
OMNI-50 Variant	Mutations Relative to WT OMNI-50	Amino acid sequence (starting at second position)	DNA sequence (starting at second position)
Wild-type	None	SEQ ID NO: 62	SEQ ID NO: 71
V3795	R478I, Y545H, Q803V, L805I	SEQ ID NO: 63	SEQ ID NO: 72
V6918	Q803V	SEQ ID NO: 64	SEQ ID NO: 73
V6919	Y545H	SEQ ID NO: 65	SEQ ID NO: 74
V6920	L805I	SEQ ID NO: 66	SEQ ID NO: 75
V6864	R478I, Y545H, Q803V	SEQ ID NO: 67	SEQ ID NO: 76
V6865	R478I, Y545H, L805I	SEQ ID NO: 68	SEQ ID NO: 77
V6866	R478I, Q803V, L805I	SEQ ID NO: 69	SEQ ID NO: 78
V6867	Y545H, Q803V, L805I	SEQ ID NO: 70	SEQ ID NO: 79
V7085	R478H, Y545H, Q803V, L805I	SEQ ID NO: 89	SEQ ID NO: 98
V7086	R478V, Y545H, Q803V, L805I	SEQ ID NO: 90	SEQ ID NO: 99
V7087	R478D, Y545H, Q803V, L805I	SEQ ID NO: 91	SEQ ID NO: 100
V7088	R478S, Y545H, Q803V, L805I	SEQ ID NO: 92	SEQ ID NO: 101
V7090	R478I, Y545C, Q803V, L805I	SEQ ID NO: 93	SEQ ID NO: 102
V7091	R478I, Y545D, Q803V, L805I	SEQ ID NO: 94	SEQ ID NO: 103
V7092	R478I, Y545T, Q803V, L805I	SEQ ID NO: 95	SEQ ID NO: 104
V7093	R478I, Y545F, Q803V, L805I	SEQ ID NO: 96	SEQ ID NO: 105
V7094	R478I, Y545A, Q803V, L805I	SEQ ID NO: 97	SEQ ID NO: 106

60 mM) to remove contaminants. Protein was eluted from the resin with high concentrations of imidazole buffer (HEPES 20 mM, NaCl 0.6 M, Imidazole 0.4 M). Eluted proteins were desalted using 96 filter plates containing 1800 µl of G-25 Sephadex resin equilibrated with storage buffer. Desalted proteins were then concentrated (Amicon-ultra 0.5 ml 50 kDa, Millipore) and sterile filtered (Ultrafree-MC 0.22 m PVDF filters). Purified variants were then stored at −80° C. and analyzed for concentration, purity, and in vitro activity before transfection to cells.

RNP Electroporation of LCL Cells

[0271] An RNP mixture was prepared by mixing 124 pmol of sgRNA and 105 pmol of nuclease. LCL cells were centrifuged at room temperature, at 300×g for 5 minutes and washed with PBS. The pellet was re-suspended in an appropriate volume of Lonza Nucleofection SG electroporation solution and transferred to the RNP mixture. Electroporation was performed using 4D-Nucleofector device (Lonza Bioscience). Pre-warmed BLCL medium was added to the

TABLE B

sg62 Guide Molecule	
sgRNA Name	sgRNA Sequence (Spacer and Scaffold)
sg62	GUGUCAAGCCCCAGAGGCCACAGUUUGAGA GUUAUGAAAAUGACGAGUUCAAAUAAAAAU UUAUUCAAAACCGCCUAUUUAUAGGCCGCAG AUGUUCUGUUU (SEQ ID NO: 80)

TABLE C

Guide Molecule Target Sequences		
Target	DNA sequence	PAM
g62	GTGTCAAGCCCCAGAGGCCACA (SEQ ID NO: 81)	GGGACA
g620T1	GCCAAACCCCAAGGCCACA (SEQ ID NO: 82)	CGGTGT

TABLE C-continued

Guide Molecule Target Sequences		
Target	DNA sequence	PAM
g620T2	GCCAAGCCTCAGAGGCCACA	GGGCAG
	(SEQ ID NO: 83)	
g62Alt	GTGTCAAGCCCCAGAGGACACA	GGGACA
	(SEQ ID NO: 84)	

TABLE D

pET9a Plasmid Utilized		
Plasmid	Purpose	Elements
pET9a	Expressing OMNI polypeptide in the bacterial system	T7 promoter - SV40 NLS - Linker - OMNI ORF (Human optimized) - HA Tag - SV40 NLS - 8XHisTag - T7 terminator

TABLE D

Appendix: pET9a Elements		
Element	Protein Sequence	DNA sequence
HA Tag	YPYDVPDYA	TACCCATACGATGT
	(SEQ ID NO: 85)	TCCAGATTACGCT (SEQ ID NO: 86)
SV40 NLS	PKKKRKV	CCAAAAAGAAAAG
	(SEQ ID NO: 87)	AAAAGTT (SEQ ID NO: 88)

TABLE E

Sample Name	% Unedited ALT	SD	% Unedited REF	SD	Graphical Representation
Activity in LCL Cells as displayed in FIG. 17A					
NT	49.94	0.57	49.92	0.54	FIG. 17A
WT	31.48	0.87	5.87	2.98	FIG. 17A
V3795	55.31	0.27	5.52	0.81	FIG. 17A
V6918	24.21	0.67	7.86	3.64	FIG. 17A
V6919	24.08	6.60	4.98	1.33	FIG. 17A
V6920	27.60	5.28	5.43	1.39	FIG. 17A
V6864	57.41	0.30	10.34	1.08	FIG. 17A
V6865	58.18	1.95	8.71	1.61	FIG. 17A
V6866	60.44	0.76	7.30	2.38	FIG. 17A
V6867	30.83	1.97	4.68	0.28	FIG. 17A
Activity in LCL Cells as displayed in FIG. 17B					
NT	49.91	1.43	49.90	1.40	FIG. 17B
V3795	62.19	2.45	6.40	4.32	FIG. 17B
V7085	61.51	2.16	7.83	4.48	FIG. 17B
V7086	61.08	2.06	7.04	3.03	FIG. 17B
V7087	27.70	2.00	6.68	0.34	FIG. 17B
V7088	56.73	0.04	8.23	2.71	FIG. 17B
V7090	34.38	2.13	6.72	2.33	FIG. 17B
V7091	21.23	2.37	6.59	0.72	FIG. 17B
V7092	36.66	0.91	10.43	1.73	FIG. 17B
V7093	59.74	2.75	10.37	6.44	FIG. 17B
V7094	61.44	1.07	7.04	4.59	FIG. 17B

TABLE F

Off-target Editing as displayed in FIG. 18A					
Sample Name	% Editing OT1	SD	% Editing OT2	SD	Graphical Representation
NT	0.04	0.01	0.03	0.01	FIG. 18A
WT	38.5	11.10	6.14	0.31	FIG. 18A
V3795	0.05	0.01	0.02	0.00	FIG. 18A
V6918	20.4	1.30	2.52	0.49	FIG. 18A
V6919	0.5	0.05	0.04	0.02	FIG. 18A
V6920	31.0	7.39	4.78	1.56	FIG. 18A
V6864	0.04	0.01	0.0	0.01	FIG. 18A
V6865	0.07	0.03	0.0	0.01	FIG. 18A
V6866	1.90	0.11	0.2	0.06	FIG. 18A
V6867	0.07	0.03	0.1	0.05	FIG. 18A

TABLE F

Off-target Editing as displayed in FIG. 18B					
Sample Name	% Editing OT1	SD	% Editing OT2	SD	Graphical Representation
NT	0.03	0.01	0.02	0.00	FIG. 18B
V3795	0.04	0.01	0.03	0.01	FIG. 18B
V7085	0.18	0.21	0.04	0.02	FIG. 18B
V7086	0.13	0.00	0.03	0.00	FIG. 18B
V7087	15.92	0.00	2.56	0.10	FIG. 18B
V7088	0.12	0.08	0.04	0.03	FIG. 18B
V7090	7.49	1.27	0.88	0.14	FIG. 18B
V7091	68.40	3.49	0.30	0.18	FIG. 18B
V7092	6.87	1.53	1.05	0.33	FIG. 18B
V7093	0.05	0.01	0.02	0.00	FIG. 18B
V7094	0.10	0.00	0.03	0.01	FIG. 18B

REFERENCES

[0273] 1. Ahmad and Allen (1992) “Antibody-mediated Specific Binding and Cytotoxicity of Liposome-entrapped Doxorubicin to Lung Cancer Cells in Vitro”, Cancer Research 52:4817-20.

[0274] 2. Anderson (1992) “Human gene therapy”, Science 256:808-13.

[0275] 3. Atosuo J T, Lilius E-M. The Real-Time-Based Assessment of the Microbial Killing by the Antimicrobial Compounds of Neutrophils. Scientific World Journal. 2011; 11:2382-2390.

[0276] 4. Basha et al. (2011) “Influence of Cationic Lipid Composition on Gene Silencing Properties of Lipid Nanoparticle Formulations of siRNA in Antigen-Presenting Cells”, Mol. Ther. 19(12):2186-200.

[0277] 5. Behr (1994) “Gene transfer with synthetic cationic amphiphiles: Prospects for gene therapy”, Bioconjugate Chem 5:382-89.

[0278] 6. Blaese et al. (1995) “Vectors in cancer therapy: how will they deliver”, Cancer Gene Ther. 2:291-97.

[0279] 7. Blaese et al. (1995) “T lymphocyte-directed gene therapy for ADA-SCID: initial trial results after 4 years”, Science 270(5235):475-80.

[0280] 8. Briner et al. (2014) “Guide RNA functional modules direct Cas9 activity and orthogonality”, Molecular Cell 56:333-39.

[0281] 9. Buchschacher and Panganiban (1992) “Human immunodeficiency virus vectors for inducible expression of foreign genes”, J. Virol. 66:2731-39.

- [0282] 10. Burstein et al. (2017) "New CRISPR-Cas systems from uncultivated microbes", *Nature* 542:237-41.
- [0283] 11. Canver et al., (2015) "BCL11A enhancer dissection by Cas9-mediated in situ saturating mutagenesis", *Nature* Vol. 527, Pgs. 192-214.
- [0284] 12. Chang and Wilson (1987) "Modification of DNA ends can decrease end-joining relative to homologous recombination in mammalian cells", *Proc. Natl. Acad. Sci. USA* 84:4959-4963.
- [0285] 13. Charlesworth et al. (2019) "Identification of preexisting adaptive immunity to Cas9 proteins in humans", *Nature Medicine*, 25(2), 249.
- [0286] 14. Chung et al. (2006) "*Agrobacterium* is not alone: gene transfer to plants by viruses and other bacteria", *Trends Plant Sci.* 11(1):1-4.
- [0287] 15. Coelho et al. (2013) "Safety and efficacy of RNAi therapy for transthyretin amyloidosis" *N. Engl. J. Med.* 369, 819-829.
- [0288] 16. Crystal (1995) "Transfer of genes to humans: early lessons and obstacles to success", *Science* 270 (5235):404-10.
- [0289] 17. Dillon (1993) "Regulation gene expression in gene therapy" *Trends in Biotechnology* 11(5):167-173.
- [0290] 18. Dranoff et al. (1997) "A phase I study of vaccination with autologous, irradiated melanoma cells engineered to secrete human granulocyte macrophage colony stimulating factor", *Hum. Gene Ther.* 8(1):111-23.
- [0291] 19. Dunbar et al. (1995) "Retrovirally marked CD34-enriched peripheral blood and bone marrow cells contribute to long-term engraftment after autologous transplantation", *Blood* 85:3048-57.
- [0292] 20. Ellem et al. (1997) "A case report: immune responses and clinical course of the first human use of granulocyte/macrophage-colony-stimulating-factor-transduced autologous melanoma cells for immunotherapy", *Cancer Immunol Immunother* 44:10-20.
- [0293] 21. Gao and Huang (1995) "Cationic liposome-mediated gene transfer" *Gene Ther.* 2(10):710-22.
- [0294] 22. Haddada et al. (1995) "Gene Therapy Using Adenovirus Vectors", in: *The Molecular Repertoire of Adenoviruses III: Biology and Pathogenesis*, ed. Doerfler and Böhm, pp. 297-306.
- [0295] 23. Han et al. (1995) "Ligand-directed retro-viral targeting of human breast cancer cells", *Proc. Natl. Acad. Sci. USA* 92(21):9747-51.
- [0296] 24. Humbert et al., (2019) "Therapeutically relevant engraftment of a CRISPR-Cas9-edited HSC-enriched population with HbF reactivation in nonhuman primates", *Sci. Trans. Med.*, Vol. 11, Pgs. 1-13.
- [0297] 25. Inaba et al. (1992) "Generation of large numbers of dendritic cells from mouse bone marrow cultures supplemented with granulocyte/macrophage colony-stimulating factor", *J Exp Med.* 176(6):1693-702.
- [0298] 26. Jinek et al. (2012) "A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity", *Science* 337(6096):816-21.
- [0299] 27. Johan et al. (1992) "GLVR1, a receptor for gibbon ape leukemia virus, is homologous to a phosphate permease of *Neurospora crassa* and is expressed at high levels in the brain and thymus", *J Virol* 66(3):1635-40.
- [0300] 28. Judge et al. (2006) "Design of noninflammatory synthetic siRNA mediating potent gene silencing in vivo", *Mol Ther.* 13(3):494-505.
- [0301] 29. Kohn et al. (1995) "Engraftment of gene-modified umbilical cord blood cells in neonates with adenosine deaminase deficiency", *Nature Medicine* 1:1017-23.
- [0302] 30. Kremer and Perricaudet (1995) "Adenovirus and adeno-associated virus mediated gene transfer", *Br. Med. Bull.* 51(1):31-44.
- [0303] 31. Macdiarmid et al. (2009) "Sequential treatment of drug-resistant tumors with targeted minicells containing siRNA or a cytotoxic drug", *Nat Biotechnol.* 27(7): 643-51.
- [0304] 32. Makaryan V. et al. The diversity of mutations and clinical outcomes for ELANE-associated neutropenia. *Curr. Opin. Hematol.* 2015; 22(1):3-11.
- [0305] 33. Malech et al. (1997) "Prolonged production of NADPH oxidase-corrected granulocytes after gene therapy of chronic granulomatous disease", *PNAS* 94(22):12133-38.
- [0306] 34. Maxwell et al. (2018) "A detailed cell-free transcription-translation-based assay to decipher CRISPR protospacer adjacent motifs", *Methods* 14348-57
- [0307] 35. Miller et al. (1991) "Construction and properties of retrovirus packaging cells based on gibbon ape leukemia virus", *J Virol.* 65(5):2220-24.
- [0308] 36. Miller (1992) "Human gene therapy comes of age", *Nature* 357:455-60.
- [0309] 37. Mitani and Caskey (1993) "Delivering therapeutic genes—matching approach and application", *Trends in Biotechnology* 11(5):162-66.
- [0310] 38. Nabel and Felgner (1993) "Direct gene transfer for immunotherapy and immunization", *Trends in Biotechnology* 11(5):211-15.
- [0311] 39. Nasri M. et al. CRISPR/Cas9-mediated ELANE knockout enables neutrophilic maturation of primary hematopoietic stem and progenitor cells and induced pluripotent stem cells of severe congenital neutropenia patients. *Haematologica.* 2020; 105(3):598-609.
- [0312] 40. Nehls et al. (1996) "Two genetically separable steps in the differentiation of thymic epithelium" *Science* 272:886-889.
- [0313] 41. Remy et al. (1994) "Gene Transfer with a Series of Lipophilic DNA-Binding Molecules", *Bioconjugate Chem.* 5(6):647-54.
- [0314] 42. Sentmanat et al. (2018) "A Survey of Validation Strategies for CRISPR-Cas9 Editing", *Scientific Reports* 8:888, doi:10.1038/s41598-018-19441-8.
- [0315] 43. Sommerfelt et al. (1990) "Localization of the receptor gene for type D simian retroviruses on human chromosome 19", *J. Virol.* 64(12):6214-20.
- [0316] 44. Van Brunt (1988) "Molecular framing: transgenic animals as bioactors" *Biotechnology* 6:1149-54.
- [0317] 45. Vigne et al. (1995) "Third-generation adenovectors for gene therapy", *Restorative Neurology and Neuroscience* 8(1,2): 35-36.
- [0318] 46. Wagner et al. (2019) "High prevalence of *Streptococcus pyogenes* Cas9-reactive T cells within the adult human population" *Nature Medicine*, 25(2), 242

- [0319] 47. Wilson et al. (1989) "Formation of infectious hybrid virion with gibbon ape leukemia virus and human T-cell leukemia virus retroviral envelope glycoproteins and the gag and pol proteins of Moloney murine leukemia virus", *J. Virol.* 63:2374-78.
- [0320] 48. Yu et al. (1994) "Progress towards gene therapy for HIV infection", *Gene Ther.* 1(1):13-26.

- [0321] 49. Zetsche et al. (2015) "Cpf1 is a single RNA-guided endonuclease of a class 2 CRISPR-Cas system" *Cell* 163(3):759-71.
- [0322] 50. Zuris et al. (2015) "Cationic lipid-mediated delivery of proteins enables efficient protein based genome editing in vitro and in vivo" *Nat Biotechnol.* 33(1):73-80.

SEQUENCE LISTING

Sequence total quantity: 106
 SEQ ID NO: 1 moltype = AA length = 1370
 FEATURE Location/Qualifiers
 REGION 1..1370
 note = Wild-type OMNI-50 nuclease
 source 1..1370
 mol_type = protein
 organism = unidentified

SEQUENCE: 1
 MTKVKDYIIG LDIGTSSVGW AVTDEAYNVL KFNSKKMWGV RLFDDAKTAE ERRGQRGARR 60
 RLDRRKKERLS LLQDFFAAEEV AKVDPNFFLR LDNSDLYMED KDQKLKSKYT LFNDDKDFKDK 120
 NFHKKYPTIH HLLMDLIEDD SKKDIRLVYL ACHYLLKNRG HPIFEGQKFD TKSSFENSLN 180
 ELKVHLNDEY GLDLEFDNEN LINILTDPKL NKTAKKKELK SVIGDTKFLK AVSAIMIGSS 240
 QKLVDLFENP EDFDSSAISK VDFSTTSFDD KYSDYELALG DKIALVNILK EIYDSSILEN 300
 LLKEADKSKD GNKYISNAFV KKYNKHGQDL KEFKRLVRQY HKSAYFDIFR SEKVNDNYVS 360
 YTKSSISNNK RVKANKFTDQ EAFYKFAKKH LETIKYKINK VNGSKADLEL IDGMLRDMEF 420
 KNFMPKIKSS DNGVIPPYQLK LMELNKKILEN QSKHHEFLNV SDEYGSVCDK IASIMEFRIP 480
 YYVGPLNPNS KYAWIKKQKD SEITPWNFKD VVDLSSREE FIDSLIGRCT YLKDEKVLKP 540
 ASLLYNEYMV LNELNNLKLN DLPITEEMKK KIFDQLFKTR KKVTLKAVAN LLKKEFNING 600
 EILLSGTDGD FKQGLNSYND FKAIVGDKVD SDDYRDKIEE IIKLIVLYGD DKSYLQKKIK 660
 AGYGYFTDS EIKKMAGLNY KDWGRLSKKL LTGLEGANKI TGERGSIHF MREYNLNLME 720
 LMSASFTEE EIQKLNPDVDD RKLSYEMVDE LYLSPSVKRM LWQSLRIVDE IKNIMGTDSK 780
 KIFIEMARGK EEVKARKESR KNQLLKPYKD GKAFISEIG EERYSYLLSE IEGEENKFR 840
 WDNLYLYYTQ LGRCMYSLEP IDISELSSKN IYDQDHIYPK SKIYDDSIEN RVLVKKDLNS 900
 KKGNSYPIPD EILNKNCYAY WKILYDKGLI GQKKYTRLTR RTGFTDDELV QFISRQIVET 960
 RQATKETANL LKTICKNSEI VYKAENASR PRQEFDIVKC RAVNDLHHMH DAYINIIVGN 1020
 VYNTKFTKDP MNFVKQKEA RSYNLENMFK YDVKRGYTA WIADDEKGTV KNASIKRIRK 1080
 ELEGTYRFT RMNYIESGAL FNATLQKKNK GSRPLKDKGP KSSIEKYGGY TNINKACFAV 1140
 LDIKSKNKIE RKLMPVEREI YAKQKNDKKL SDEIFSKYLK DRFGIEDYRV VYPVVKMRTL 1200
 LKIDGSYTFI TGSQDTELEL RSALQLILPK KNEWAIKQID KSENNDYLT IRIQDLTEEL 1260
 VYNTFDIIVN FKFTSVFKKS FLNLFQDDKI ENIDFKFKSM DFKECKCTLL MLVKAIRASG 1320
 VRQDLKSIDL KSDYGRLLSSK TNNIGNYQEF KIINQSITGL FENEVDLLKL 1370

SEQ ID NO: 2 moltype = AA length = 1370
 FEATURE Location/Qualifiers
 REGION 1..1370
 note = Variant 3795
 source 1..1370
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 2
 MTKVKDYIIG LDIGTSSVGW AVTDEAYNVL KFNSKKMWGV RLFDDAKTAE ERRGQRGARR 60
 RLDRRKKERLS LLQDFFAAEEV AKVDPNFFLR LDNSDLYMED KDQKLKSKYT LFNDDKDFKDK 120
 NFHKKYPTIH HLLMDLIEDD SKKDIRLVYL ACHYLLKNRG HPIFEGQKFD TKSSFENSLN 180
 ELKVHLNDEY GLDLEFDNEN LINILTDPKL NKTAKKKELK SVIGDTKFLK AVSAIMIGSS 240
 QKLVDLFENP EDFDSSAISK VDFSTTSFDD KYSDYELALG DKIALVNILK EIYDSSILEN 300
 LLKEADKSKD GNKYISNAFV KKYNKHGQDL KEFKRLVRQY HKSAYFDIFR SEKVNDNYVS 360
 YTKSSISNNK RVKANKFTDQ EAFYKFAKKH LETIKYKINK VNGSKADLEL IDGMLRDMEF 420
 KNFMPKIKSS DNGVIPPYQLK LMELNKKILEN QSKHHEFLNV SDEYGSVCDK IASIMEFIIP 480
 YYVGPLNPNS KYAWIKKQKD SEITPWNFKD VVDLSSREE FIDSLIGRCT YLKDEKVLKP 540
 ASLLHNEYMV LNELNNLKLN DLPITEEMKK KIFDQLFKTR KKVTLKAVAN LLKKEFNING 600
 EILLSGTDGD FKQGLNSYND FKAIVGDKVD SDDYRDKIEE IIKLIVLYGD DKSYLQKKIK 660
 AGYGYFTDS EIKKMAGLNY KDWGRLSKKL LTGLEGANKI TGERGSIHF MREYNLNLME 720
 LMSASFTEE EIQKLNPDVDD RKLSYEMVDE LYLSPSVKRM LWQSLRIVDE IKNIMGTDSK 780
 KIFIEMARGK EEVKARKESR KNLVKKFYKD GKAFISEIG EERYSYLLSE IEGEENKFR 840
 WDNLYLYYTQ LGRCMYSLEP IDISELSSKN IYDQDHIYPK SKIYDDSIEN RVLVKKDLNS 900
 KKGNSYPIPD EILNKNCYAY WKILYDKGLI GQKKYTRLTR RTGFTDDELV QFISRQIVET 960
 RQATKETANL LKTICKNSEI VYKAENASR PRQEFDIVKC RAVNDLHHMH DAYINIIVGN 1020
 VYNTKFTKDP MNFVKQKEA RSYNLENMFK YDVKRGYTA WIADDEKGTV KNASIKRIRK 1080
 ELEGTYRFT RMNYIESGAL FNATLQKKNK GSRPLKDKGP KSSIEKYGGY TNINKACFAV 1140
 LDIKSKNKIE RKLMPVEREI YAKQKNDKKL SDEIFSKYLK DRFGIEDYRV VYPVVKMRTL 1200
 LKIDGSYTFI TGSQDTELEL RSALQLILPK KNEWAIKQID KSENNDYLT IRIQDLTEEL 1260
 VYNTFDIIVN FKFTSVFKKS FLNLFQDDKI ENIDFKFKSM DFKECKCTLL MLVKAIRASG 1320
 VRQDLKSIDL KSDYGRLLSSK TNNIGNYQEF KIINQSITGL FENEVDLLKL 1370

-continued

SEQ ID NO: 3 moltype = AA length = 1370
 FEATURE Location/Qualifiers
 REGION 1..1370
 note = Variant 3942
 source 1..1370
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 3

MTKVKDYYIG	LDIGTSSVGW	AVTDEAYNVL	KFNSKKMWGV	RLFDDAKTAE	ERRGQRGARR	60
ALDRKKERLS	LLQDFFAEV	AKVDPNFFLR	LDNSDLYMED	KDQKLKSKYT	LFNDKDFKDK	120
NFHKKYPTIH	HLLMDLIEDD	SKKDIRLVYL	ACHYLLKNRG	HFIFEGQKFD	TKSSFENSLN	180
ELKVHLNDEY	GLDLEFDNEN	LINILTDPKL	NKTAKKKELK	SVIGDTKFLK	AVSAIMIGSS	240
QKLVDLFENP	EDFDDSAIKS	VDFSTTSFDD	KYSDYELALG	DKIALVNILK	EIYDSSILEN	300
LLKEADKSKD	GNKYISNAFV	KKYNKHGQDL	KEFKRLVRQY	HKSAYFDIFR	SEKVNDNYVS	360
YTKSSISNNK	RVKANKFTDQ	EAFYKFAKKH	LETIKYKINK	VNGSKADLEL	IDGMLRDMEF	420
KNFMPIKISS	DNGVIPWQLK	LMELNKILEN	QSKHHEFLNV	SDEYGSVCDK	IASIMEFRIP	480
YYVGPLNPNS	KYAWIKKQKD	SEITPWNFKD	VVDLDSSREE	FIDSLIGRCT	YLKDEKVLPK	540
ASLLYNEYMV	LNELNNLKLN	DLPITEEMKK	KIFDQLFKTR	KKVTLKAVAN	LLKKEFNING	600
EILLSGTDGD	FKQGLNSYND	FKAIVGDKVD	SDDYRDKIEE	IIKLIVLYGD	DKSYLQKKIK	660
AGYGKYFTDS	EIKKMAGLNY	KDWGRLSKKL	LTGLEGANKI	TGERGSIHF	MREYNLNLME	720
LMSASFTFTE	EIQKLNVPDD	RKLSYEMVDE	LYLSPSVKRM	LWQSLRIVDE	IKNIMGTDSK	780
KFIEMAKGK	EEVKARKESR	KNQLLKIFYD	GKKAFISEIG	EERYSYLLSE	IEGEEENKFR	840
WDNNLYYYTQ	LGRCMYSLEP	IDISELSSKN	IYDQDHIYPK	SKIYDDSIEN	RVLVKKDLNS	900
KKGNSYPIPD	EILNKNYAY	WKILYDKGLI	GQKKYTRLTR	RTGFTDDELV	QFISRQIVET	960
ROATKETANL	LKTICKNSEI	MYSKAENASR	FRQEPDIVKC	RAVNDLHHMH	DAYINIIIVGN	1020
VYNTKFTKDP	MNFVKKQEKA	RSYNLENMPK	YDVKRGGYTA	WIADDEKGTV	KNASIKRIRK	1080
ELEGTYNRYFT	RMNYIESGAL	FNATLQQRKNK	GSRLPKDKGP	KSSIEKYGGY	TNINKACFAV	1140
LDIKSKNKIE	RKLMPVEREI	YAKQKNDKKL	SDEIFSKYLK	DRFGIEDYRV	VYPVVKMRTL	1200
LKIDGSYYFI	TGGSDDTLLEL	RSALQLILPK	KNEWAIKQID	KSSENDYLT	ERIQDLTEEL	1260
VYNTFDIIVN	KFKTSVFKKS	FLNLFQDDKI	ENIDFKFKSM	DFKECKTLL	MLVKAIRASG	1320
VRQDLKSIDL	KSDYGRLLSSK	TNNIGNYQEF	KIINQSITGL	FENEVDLLKL		1370

SEQ ID NO: 4 moltype = AA length = 1370
 FEATURE Location/Qualifiers
 REGION 1..1370
 note = Variant 2902
 source 1..1370
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 4

MTKVKDYYIG	LDIGTSSVGW	AVTDEAYNVL	KFNSKKMWGV	RLFDDAKTAE	ERRGQRGARR	60
RLDRKKERLS	LLQDFFAEV	AKVDPNFFLR	LDNSDLYMED	KDQKLKSKYT	LFNDKDFKDK	120
NFHKKYPTIH	HLLMDLIEDD	SKKDIRLVYL	ACHYLLKNRG	HFIFEGQKFD	TKSSFENSLN	180
ELKVHLNDEY	GLDLEFDNEN	LINILTDPKL	NKTAKKKELK	SVIGDTKFLK	AVSAIMIGSS	240
QKLVDLFENP	EDFDDSAIKS	VDFSTTSFDD	KYSDYELALG	DKIALVNILK	EIYDSSILEN	300
LLKEADKSKD	GNKYISNAFV	KKYNKHGQDL	KEFKRLVRQY	HKSAYFDIFR	SEKVNDNYVS	360
YTKSSISNNK	RVKANKFTDQ	EAFYKFAKKH	LETIKYKINK	VNGSKADLEL	IDGMLRDMEF	420
KNFMPIKISS	DNGVIPYQLK	LMELNKILEN	QSKHHEFLNV	SDEYGSVCDK	IASIMEFRIP	480
YYVGPLNPNS	KYAWIKKQKD	SEITPWNFKD	VVDLDSSREE	FIDSLIGRCT	YLKDEKVLPK	540
ASLLYNEYMV	LNELNNLKLN	DLPITEEMKK	KIFDQLFKTR	KKVTLKAVAN	LLKKEFNING	600
EILLSPDGD	FKQGLNSYND	FKAIVGDKVD	SDDYRDKIEE	IIKLIVLYGD	DKSYLQKKIK	660
AGYGKYFTDS	EIKKMAGLNY	KDWGRLSKKV	LTGLQGANIK	TGERGSIHF	MREYNLNLME	720
LMSASFTFTE	EIQKLNVPDD	RKLSYEMVDE	LYLSPSVKRM	LWQSLRIVDE	IKNIMGTDSK	780
KFIEMARGK	EEVKARKESR	KNQLLKIFYD	GKKAFISEIG	EERYSYLLSE	IEGEEENKFR	840
WDNLYYYTQ	LGRCMYSLEP	IDISELSSKN	IYDQDHIYPK	SKIYDDSIEN	RVLVKKDLNS	900
KKGNSYPIPD	EILNKNYAY	WKILYDKGLI	GQKKYTRLTR	RTGFTDDELV	QFISRQIVET	960
ROATKETANL	LKTICKNSEI	VYSKAENASR	FRQEPDIVKC	RAVNDLHHMH	DAYINIIIVGN	1020
VYNTKFTKDP	MNFVKKQEKA	RSYNLENMPK	YDVKRGGYTA	WIADDEKGTV	KNASIKRIRK	1080
ELEGTYNRYFT	RMNYIESGAL	FNATLQQRKNK	GSRLPKDKGP	KSSIEKYGGY	TNINKACFAV	1140
LDIKSKNKIE	RKLMPVEREI	YAKQKNDKKL	SDEIFSKYLK	DRFGIEDYRV	VYPVVKMRTL	1200
LKIDGSYYFI	TGGSDDTLLEL	RSALQLILPK	KNEWAIKQID	KSSENDYLT	ERIQDLTEEL	1260
VYNTFDIIVN	KFKTSVFKKS	FLNLFQDDKI	ENIDFKFKSM	DFKECKTLL	MLVKAIRASG	1320
VRQDLKSIDL	KSDYGRLLSSK	TNNIGNYQEF	KIINQSITGL	FENEVDLLKL		1370

SEQ ID NO: 5 moltype = AA length = 1370
 FEATURE Location/Qualifiers
 REGION 1..1370
 note = Variant 3954
 source 1..1370
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 5

MTKVKDYYIG	LDIGTSSVGW	AVTDEAYNVL	KFNSKKMWGV	RLFDDAKTAE	ERRGQRGARR	60
RLDRKKERLS	LLQDFFAEV	AKVDPNFFLR	LDNSDLYMED	KDQKLKSKYT	LFNDKDFKDK	120
NFHKKYPTIH	HLLMDLIEDD	SKKDIRLVYL	ACHYLLKNRG	HFIFEGQKFD	TKSSFENSLN	180
ELKVHLNDEY	GLDLEFDNEN	LINILTDPKL	NKTAKKKELK	SVIGDTKFLK	AVSAIMIGSS	240

-continued

QKLVDLFENP	EDFDDSAIKS	VDFSTTSFDD	KYSDYELALG	DKIALVNILK	EIYDSSILEN	300
LLKEADKSKD	GKNYISNAFV	KKNYKHGQDL	KEFKRLVRQY	HKSAYFDIFR	SEKVNNDNYVS	360
YTKSSISNNK	RVKANKFTDQ	EAFYKFAKKH	LETIKYKINK	VNGSKADLEL	IDGMLRDMEF	420
KNFMPKIKSS	DNGVIPYQLK	LMELNKILEN	QSKHHEFLNV	SDEYGSVCDK	IASIMEFTIP	480
YYVGPLNPNS	KYGWIKQKQD	SEITPWNFKD	VVDLDSSREE	FIDSLIGRCT	YLKDEKVLPK	540
ASLLYNEYMV	LNELNNLKLN	DLPIEEMKK	KIFDQLFKTR	KKVTLKAVAN	LLKKEFNING	600
EILLSGTDGD	FKQGLNSYND	FKAIIVGDKVD	SDDYRDKIEE	IKLIVLYGD	DKSYLQKKIK	660
AGYGKYFTDS	EIKKMAGLNY	KDWGRLESEK	LTGLEGANKI	TGERGSIHF	MREYNLNCME	720
LMSASFTFTE	EIQKLNVPDD	RKLSYEMVDE	LYLSPSVKRM	LWQSLRIVDE	IKNIMGTDSK	780
KIFIEMARGK	EEVKARKESR	KNQLLKFYKD	GKKAFISEIG	EERYSYLLSE	IEGEEENKFR	840
WDNLYLYYTQ	LGRCMYSLEP	IDISELSSKN	IYDQDHIYPK	SKIYDDSIEN	RVLVKKDLNS	900
KKGNSYPIPD	EILNKNCYAY	WKILYDKGLI	GQKQYTRLTR	RTGFTDDELV	QFISRQIVET	960
RQATVETANL	LKTICKNSEI	VYSKAENASR	FRQEPDIVKC	RAVNDLHHMH	DAYINIIVGN	1020
VYNTKFTKDP	MNFVKVQEKA	RSYNLNMFK	YDVKRGGYTA	WIADDEKGTV	KNASIKRIRK	1080
ELEGNTNYRT	RMNYESGAL	FNATLQQRNK	GSRLPKDKGP	KSSIEKYGGY	TNINKACFAV	1140
LDIKSKNKIE	RKLMPVEREI	YAKQKNDKKL	SDEIFSKYLK	DRFGIEDYRV	VYPVVKMRTL	1200
LKIDGSYYFI	TGGSDDKLEL	RSALQLILPK	KNEWAIKQID	KSENDYLTII	ERIQDLTEEL	1260
VYNTFDIIVN	KFKTSVFKKS	FLNLFQDDKI	ENIDFKFKSM	DFKEKCKTLL	MLVKAIRASG	1320
VRQDLKSIDL	KSDYGRLLSK	TNNIGNYQEF	KIINQISITGL	FENEVDLLKL		1370

SEQ ID NO: 6 moltype = DNA length = 4110
 FEATURE Location/Qualifiers
 misc_feature 1..4110
 note = Wild-type OMNI-50 nuclease DNA sequence
 source 1..4110
 mol_type = other DNA
 organism = unidentified

SEQUENCE: 6

atgaccaagg	tgaaggacta	ctacataggc	ttggacatcg	gcacctctag	cgtcggtgtg	60
gccgtcaccg	atgaagccta	taacgtgctt	aagtttaata	gcaagaaaaa	gtggggcgctg	120
cggctgtgtc	acgacgctaa	gacggcagag	gagcgtaggg	gccagcgagg	agcaagacga	180
cgtctggatc	ggaagaagga	gagactcagc	ctgctgcagg	actctctcgc	cgaagaggtta	240
gcaaaggctg	accccaactt	cttcctcagg	ctggacaatt	ccgatctgta	catgggaagat	300
aaggaccaga	aactgaaaag	caaatatata	ctgttcaacg	acaaggactt	caaggataag	360
aattttcata	agaagtaccc	cacaatacat	cacctgtcta	tggatctgat	cgaggacgac	420
agtaagaagg	acatccggct	cgtctacctg	gcctgtcact	atttgtctca	gaacagggggt	480
catttcactc	tcaggggcca	gaagttcgac	actaaatcaa	gcttcgagaa	cagtttgaa	540
gagctcaaa	ttcatttgaa	cgacgagtat	ggactggacc	tcgaatttga	caacgagaa	600
ctgattaaca	tcttgactga	cccaaaactc	aataaaacgg	ccaagaagaa	ggagctgaag	660
tcgtaaatcg	gcgacaccaa	gttctcctca	gccgtttccg	cgataatgat	cggctctagc	720
cagaaactcg	tcgacttggt	cgagaacccc	gaggatttcg	acgactctgc	gataaagtc	780
gttgactctc	caactacctc	tttcgacgac	aagtaactct	actatgaact	cgctctgggt	840
gacaagatcg	ctctggtcaa	catccttaag	gaaatttacg	atagctccat	cctcgagaa	900
ctgctcaaa	aggcagacaa	gtctaaggac	ggtaacaaat	atatcagtaa	tgcattcggt	960
aagaagtaca	ataaacacgg	acaagatctg	aaagagttca	aacgtctggt	acgacaatat	1020
cacaagagtg	cgtattttga	tattttcaga	tcgagagaag	tgaatgacaa	ttacgtcagc	1080
tacactaaaa	gctcaattag	caacaataaa	cgcgtcaaa	caaacaagtt	caactgatcaa	1140
gaggccttct	acaaattcgc	caagaacat	ctggagacaa	tcaagataaa	gatcaacaag	1200
gtaaacggct	ccaaggcaga	tctggagctg	attgacggga	tgctgcggga	catggagttc	1260
aagaacttta	tgcccaaaat	taagtccagt	gacaacgggg	tgattccata	ccagctcaag	1320
ctgatggaat	tgaacaaaat	actcgagaat	cagtcacaa	atcacgagtt	cctcaatgtc	1380
agcgacgagt	acggctccgt	gtgtgataaa	atcgcatcta	tcattggagt	ccgtatcccc	1440
tactacgtgg	gacctctgaa	ccccaatagc	aagtacgcct	ggatcaagaa	gcagaaagat	1500
agtgagatta	ctccctggaa	cttcaaggac	gtcgtggacc	tcgactccag	cagagaggag	1560
ttcattgact	cactgatcgg	acgctgtact	tacctaagg	acgagaaggt	ccttcccaaa	1620
gcttctttgc	tgtataacga	atacatggtg	ctgaacgagc	tgaataacct	gaagttgaac	1680
gaccttccca	tcaccgagga	gatgaagaag	aagatatttg	accagtttgt	caaaacaaga	1740
aagaaggtca	cccttaaaag	ggtggcaaac	ctgctgaaga	aggagttcaa	catcaacggc	1800
gagattctgc	tctctgggac	cgacggtgac	ttcaagcagg	gcttgaactc	atacaatgac	1860
ttcaagacta	tcgtgggcga	taaagtcgat	tccgatgatt	accgggacaa	gattgaggag	1920
atcattaaac	tgatagttct	ttacggtgac	gataaagagt	accttcagaa	gaagattaaa	1980
gctgggtatg	gaaaaactct	caccgacagt	gagattaa	aaatggcggg	gctgaactac	2040
aaggattggg	gaaggctctc	aaagaagctg	ctgacgggac	tcgaggggtg	aaacaagatc	2100
actggagagc	ggggctccat	tattcacttc	atgagggaat	ataaccttaa	tctgatggag	2160
cttatgtcag	cttcatttac	gttcacgcaa	gagatacaga	aacttaacct	cgtggatgac	2220
cgcaagctgt	catacgaaat	ggtggacgaa	ctgtaccttt	ctcccagtgt	gaaacggatg	2280
ctctggcagt	ccctgcgcgt	cgtcgacgag	ataaagaaca	tcatgggaac	cgacagtaag	2340
aagattttca	tcgagatggc	tcggggtaag	gaagaggtga	aagcccgcaa	ggagtcaagg	2400
aagaaccaac	tgctgaagtt	ctataaagac	ggaaagaagg	cattcatcag	cgagattggc	2460
gaggagaggt	actcttactt	gctttctgag	atagagggtg	aggaagagaa	taagtttcga	2520
ttgggataacc	tgtactttta	ttatactcaa	ctgggtcgct	gcatgtactc	tttggaaacct	2580
atcgacatat	ctgagctgtc	ttcaaagaat	atttacgata	aggatcatat	ctaccccaaa	2640
agcaagattt	acgacgacag	tatcgagaat	agggtgctgg	tgaagaagga	ccttaactcc	2700
aagaagggtta	acagctatcc	tatcccagac	gaaatcctga	acaagaactg	ttacgcctac	2760
tggaagatcc	tgatcgataa	aggctttatc	gggcagaaga	agtaactctg	gctgacccgg	2820
agaactggct	tcacggacga	cgagctcggt	cagttcatct	caagacagat	cgtggaaact	2880

-continued

```

agacaagcaa caaaggagac tgctaacctg ctcaagacaa tatgtaagaa ctccgagatc 2940
gtgtattcca aagcogagaa cgcaagtcgg tttaggcaag agttcgacat cgtgaagtgt 3000
agggcggtga acgatcttca tcatatgcac gatgcctaca tcaacatcat agtggggaac 3060
gtgtataaca ccaagttcac gaaggaccct atgaatttcg taaagaagca ggaaaaggcg 3120
cggagctaca atctcgagaa tatgttcaag tacgatgtga aacgtggcgg atacacgcgt 3180
tggatcgccg atgacgagaa gggcaccgtg aagaacgcga gtattaaacg tatccggaag 3240
gagctggaag gcacaaatta taggttcaca agaatgaact acattgagtc tggagcgctt 3300
ttcaacgcga ctctccagcg gaagaataag ggctccagac ccctgaagga caaaggcccg 3360
aaatcttcca tcgagaagta cggcggctac acaaacatca ataaagcctg tttcgctgtt 3420
cttgacatca agtctaagaa caagattgag aggaagctga tgcccgtcga gcgtgagatc 3480
tatgccaaac agaagaacga caagaagctg tccgacgaga tttctcaaa gtacctcaag 3540
gaccgatctg gcacgcagga ctacagggtt gctacccag tggtgaaaat gcgcacactg 3600
ctcaagatcg acggcagctc ctacttcac acaggcgggt ctgataagac cctggagttg 3660
cgatctgtct tgcagctgat tctccctaag aagaacgagt gggcgatcaa acagatcgac 3720
aagctcttcg aaaacgacta tctgacgatc gagcgatcc aggcctgac caggagctg 3780
gtgtataaca ctctcgacat catcgtaaac aagttcaaga ccagtgtctt caagaagtct 3840
tctcttaact tgtttcagga cgacaagatt gagaacattg acttcaagtt taagtccatg 3900
gacttcaagg agaattgcaa gacacttctc atgctggta aggcgatctg ggcatccggc 3960
gtgaggcagg atctcaagtc catcgacctc aagctgatt acggacggct cagttcaag 4020
accaacaaca tcggcaatta ccaggagttc aagattatta atcagtcctc cactggactg 4080
ttcgagaatg aggtcgatct cctgaagctg 4110

```

```

SEQ ID NO: 7      moltype = DNA length = 4110
FEATURE          Location/Qualifiers
misc_feature      1..4110
                  note = Variant 3795 DNA sequence
source            1..4110
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 7
atgaccaagg tgaaggacta ctacataggc ttggacatcg gcacctctag cgtcgggtgg 60
gccgtcaccc atgaagccta taacgtgctt aagtttaata gcaagaaaaa gtggggcggt 120
cggctgtttc acgacgctaa gacggcagag gagcgtaggg gccagcgagg agcaagacga 180
cgtctggatc ggaagaagga gagactcagc ctgctgcagg acttctctgc cgaagaggta 240
gcaaaggctc accccaactt ctctctcagg ctggacaatt ccgatctgta catggaagat 300
aaggaccaga aactgaaaag caaatatata ctgttcaacg acaaggactt caaggataag 360
aatcttcata agaagtcacc cacaatacat cacctgctga tggatctgat cgaggacgac 420
agtaagaagg acatccggct cgtctacctg gccgttcaat atttgcctaa gaacaggggt 480
catttcatct tcgagggcca gaagttcgac actaaatcaa gcttcgagaa cagtttgaac 540
gagctcaaa tctatttga gacgagatg ggactggacc tcgaatttga caacgagaac 600
ctgattaaca tcttgatgta cccaaaactc aataaaaacg ccaagaagaa ggagctgaag 660
tcctgaatcg gcgacaccaa gttcctcaaa gccgtttccg cgataatgat cggctctagc 720
cagaactctg tcgacttggt cgagaacccc gaggtattcg acgactctgc gataaagtcc 780
gttgacttct caactacctc ttctcgacgc aagtactctg actatgaact cgctctgggt 840
gacaagatcg ctctggctca catccttaag gaaatttacc atagctccat cctcgagaac 900
ctgctcaaa gtcataacga gctaaggac ggtaacaaat atatcagtaa tgcattctg 960
aagaagtaca ataaacacgg caagatctg aaagagtcca aacgtctggt acgacaatat 1020
cacaagatg cgtattttga tattttcaga tccgagaagg tgaatgacaa ttacgtcagc 1080
tacactaaaa gctcaattag caacaataaa ccgctcaaa ccaacaagtt cactgatcaa 1140
gaggccttct caaaattcgc caagaacat ctggagacaa tcaagtataa gatcaacaag 1200
gtaaacggct ccaaggcaga tctggagctg attgacggga tgctgcggga catggagttc 1260
aagaacttta tgcccaaaat taagtccagt gacaacgggg tgattccata ccagctcaag 1320
ctgatggaat tgaacaaaat actcgagaat cagtcaaaag atcacgagtt cctcaatgct 1380
agcgacgagt acggctccgt gtgtgataaa atcgcatcta tcatggaatt tattattccg 1440
tattacgtgg gcccgctgaa cccgaacagc aaatatgctg ggatcaagaa gcagaaagat 1500
agtgaatta tctcctggaa cttcaaggac gtcgtggacc tcgactccag cagagaggag 1560
ttcattgact cactgatcgg acgctgtact taccttaagg acgagaaggt ccttcccaaa 1620
gcttctttgc tcgataacga atacatggtg ctgaacgagc tgaataacct gaagttgaac 1680
gaccttccca tcaccgagga gatgaagaag aagatatttg accagttggt caaaacaaga 1740
aagaaggctc cccttaaaag ggtggcaaac ctgctgaaga aggagttcaa catcaacggc 1800
gagattctgc tctctgggac cgacggtgac ttcaagcagg gcttgaactc atacaatgac 1860
ttcaagcta tcgtgggcca taaagtcgat tccgatgatt accgggacaa gattgaggag 1920
atcattaaac tgatagttct ttacggtgac gataagagtt accttcagaa gaagattaaa 1980
gctgggtatg gaaaatactt acccgacagt gagattaaga aaatggcggg gctgaactac 2040
aaggattggg gaaggctctc aaagaagctg ctgacgggac tcgaggggtg aaacaagatc 2100
actggagagc ggggctccat tattcacttc atgagggaat ataaccttaa tctgatggag 2160
cttatgtcag cttcatttac gttcacggaa gagatcacga aacttaacct cgtggatgac 2220
cgcaagctgt catacgaaat ggtggacgaa ctgtaccttt ctcccagtg gaaacggatg 2280
ctctggcagt ccttcgcat cgtcgacgag ataaagaaca tcatgggaac cgacagtaag 2340
aagattttca tcgagatggc tggggtaag gaagaggtga aagcccgcaa ggagtcagg 2400
aagaacgtcc tactcaagtt ctataaagac ggaaagaagg catctacag cgagattggc 2460
gaggagaggt actcttactt gctttctgag atagagggtg aggaagagaa taagtttcta 2520
tgggataacc tgtaccttta ttatactcaa ctgggtcgct gcatgtactc tttggaacct 2580
atcgacatat ctgagctgtc ttcaagaat atttacgac aggatcatat ctaccccaaa 2640
agcaagattt acgacgacag tatcgagaat aggggtgctg tgaagaagga ccttaactcc 2700
aagaagggtt acagctatcc tatccagac gaaatcctga acaagaactg ttacgcctac 2760

```

-continued

tggaagatcc	tgtacgataa	aggtcttata	gggcagaaga	agtacactcg	gctgacccgg	2820
agaactggct	tcacggacga	cgagctcggt	cagttcatct	caagacagat	cgtggaaact	2880
agacaagcaa	caaaggagac	tgctaacctg	ctcaagacaa	tatgtaagaa	ctccgagatc	2940
gtgtattcca	aagccgagaa	cgcagtcggt	ttagggcaag	agttcgacat	cgtgaagtgt	3000
agggcggtga	acgatcttca	tcatatgcac	gatgcctaca	tcaacatcat	agtggggaac	3060
gtgtataaca	ccaagtccac	gaaggaccct	atgaatttcg	taaagaagca	ggaaaaggcg	3120
cggagctaca	atctcgagaa	tatgttcaag	tacgatgtga	aacgtggcgg	atacaccgct	3180
tggatcgccg	atgaocgagaa	gggcacccgt	aagaacgcga	gtattaaacg	tatccggaag	3240
gagctggaag	gcacaaatta	taggttcaca	agaatgaact	acattgagtc	tggagcgctt	3300
ttcaacgcca	ctctccagcg	gaagaataag	ggctccagac	ccctgaagga	caaaggcccc	3360
aaatcttcca	tcgagaagta	cggcggtcac	acaaacatca	ataaagcctg	tttcgctgtt	3420
cttgacatca	agtctaagaa	caagattgag	aggaagctga	tgcccgctga	gcgtgagatc	3480
tatgccaac	agaagaacga	caagaagctg	tccgacgaga	ttttctcaaa	gtacctcaag	3540
gaccgatttg	gcacgaggtt	ctacaggggt	gtctaccacg	tggtgaaaat	gcgcacactg	3600
ctcaagatcg	acggcagcta	ctacttcac	acaggcggtt	ctgataagac	cctggagttg	3660
cgatctgctc	tgcagctgat	tctccctaag	aagaacgagt	gggcatcaa	acagatcgac	3720
aagtcttccg	aaaacgacta	tctgacgatc	gagcgtatcc	aggacctgac	cgaggagctg	3780
gtgtataaca	ctttcgacat	catcgccaac	aagtccaaga	ccagtgtctt	caagaagtct	3840
ttctttaact	tgatttcagga	gcacaagatt	gagaacattg	acttcaagtt	taagtccatg	3900
gacttcaagg	agaaatgcga	gacactttct	atgctggcca	aggcgattcg	ggcatccggc	3960
gtgagggcag	atctcaagtc	catcgacctc	aagctctgatt	acggacggct	cagttcaaa	4020
accaacacaa	tcggcaatta	ccaggagttc	aagattatta	atcagtcctat	cactggactg	4080
ttcgagaatg	aggtcgatct	cctgaagctg				4110

SEQ ID NO: 8
 FEATURE
 misc_feature
 source

moltype = DNA length = 4110
 Location/Qualifiers
 1..4110
 note = Variant 3942 DNA sequence
 1..4110
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 8

atgaccaagg	tgaaggacta	ctacataggg	ttggacatcg	gcacctctag	cgtcggggtg	60
gocgtcaccc	atgaagccta	taacgtgctt	aagtttaata	gcaagaaaat	gtggggcggt	120
cggctgttcc	acgacgctaa	aaaccgagg	gagcgccgtg	gccagcgccg	cgcgcccggt	180
gcgctggatc	gcaaaaaggga	acgcctcagc	ctgctgcagg	acttcttcgc	cgaagaggtg	240
gcaaaggctg	accccaactt	cttctcagg	ctggacaatt	ccgatctgta	catgggaagt	300
aaggaccaga	aactgaaaag	caaatataca	ctgttcaacg	acaaggactt	caaggataag	360
aattttcata	agaagtaccc	cacaatacat	cacctgctga	tgatctctat	cgaggacgac	420
agtaagaagg	acatccggct	gctctacctg	gcctgtcact	atttgcctaa	gaacaggggt	480
catttcctct	tcgaggggcca	gaagttcgac	actaaatcaa	gcttcgagaa	cagtttgaaa	540
gagctcaaa	ttcatttgaa	cgacgagtat	ggactggacc	tcgaatttga	caacgagaa	600
ctgatttaaca	tcttgactga	cccaaaaact	aataaaaacg	ccaagaagaa	ggagctgaag	660
tccgtaactg	gcgacaccaa	gttctcctaa	gccgtttccg	cgataatgat	cggctctagc	720
cagaaactcg	tcgacttggt	cgagaacccc	gaggatttcg	acgactctgc	gataaagttc	780
gttgactttc	caactctcgc	tttcgacgac	aagtactctg	actatgaact	cgctctgggt	840
gacaagatcg	ctctgggtcaa	cactccttaag	gaaatttacg	atagctccat	cctcgagaa	900
ctgctcaaa	aggcagacaa	gtctaaggac	ggtaacaaat	atatacagta	tgcatctcgt	960
aagaagtaca	ataaacacgg	acaagatctg	aaagagtcca	aacgtctggt	acgacaatat	1020
cacaagagtg	cgtattttga	tattttcaga	tccgagaagg	tgaatgacaa	ttacgtcagc	1080
tacactaaaa	gctcaattag	caacaataaa	cgcgtcaaa	caaacaaagt	cactgatcaa	1140
gaaggccttc	acaaattcgc	caagaacacat	ctggagacaa	tcaagtataa	gatcaacaag	1200
gtaaaacggc	ccaaggcaga	tctggagctg	attgacggga	tgctgcgagg	catggagttc	1260
aagaacttta	tgcccaaaat	taaaagctcg	gataacggcg	tgattccgtg	gcagctgaaa	1320
ctgatggaat	tgaacaaaat	actcgagaat	cagtcaaaag	atcacgagtt	cctcaatgtc	1380
agcgacgagt	acggctccgt	gtgtgataaa	atcgcatcta	tcatggagtt	ccgtatcccc	1440
tactacgtgg	gacccctgaa	ccccaatagc	aagtacgcct	ggatcaagaa	gcagaaagat	1500
agtgagatta	ctccctggaa	cttcaaggac	gtcgtggacc	tcgactccag	cagagaggag	1560
ttcattgact	cactgatcgg	acgctgtact	taccttaagg	acgagaaggt	ccttcccaaa	1620
gcttcttttc	tgtataacga	atacatgggt	ctgaacgagc	tgaataacct	gaagtgtgac	1680
gaacttccca	tcaccgagga	gatgaagaag	aagatatttg	accagttgtt	caaaaacaga	1740
aagaagggtc	cccttaaaag	gggtggcaaac	ctgctgaaga	aggagttcaa	catcaacggc	1800
gagattctgc	tctctgggac	cgaacgtgac	ttcaagcagg	gcttgaactc	atacaatgac	1860
ttcaaaagta	tcgtggggcg	taaagtcgat	tccgatgatt	accgggacaa	gattgaggag	1920
atcattaaac	tgatagttct	ttacggtgac	gataaagagt	accttcagaa	gaagattaaa	1980
gctgggtatg	gaaaaatact	caccgacagt	gagattaaag	aaatggcggg	gctgaaactac	2040
aaggattggg	gaaggctctc	aaagaagctg	ctgacgggac	tcgaggggtg	aaacaagatc	2100
actggagagc	ggggctccat	tattcacttc	atgagggaa	ataaccttaa	tctgatggag	2160
cttatgtcag	cttcattttac	gttcaccgaa	gagatacaga	aacttaacct	cgtggatgac	2220
cgcaagctgt	catcacgaat	gggtggacgaa	ctgtaccttt	ctcccagttg	gaaacgggatg	2280
ctctggcagt	ccctgcgcac	cgtcgacgag	ataaagaaca	tcatgggaac	cgacagtaag	2340
aagattttta	tcgagatggc	taagggttaag	gaagaggtga	aagcccgcaa	ggagtcaagg	2400
aagaaccaac	tgctgaagtt	ctataaagac	ggaaagaagg	cattoatcag	cgagattggc	2460
gaggagaggt	actcttactt	gctttctgag	atagaggggt	aggaagagaa	taagtttcga	2520
tgggataaca	actaccttta	ttatactcaa	ctgggtcgct	gcattgactc	tttggaaact	2580
atcgacatat	ctgagctgtc	ttcaagaat	atttacgac	aggatcatat	ctaccccaaa	2640

-continued

```

agcaagattt acgacgacag tatcgagaat aggggtgctg tgaagaagga ccttaactcc 2700
aagaagggtt acagctatcc tatcccagac gaaatcctga acaagaactg ttacgcctac 2760
tggaagatcc tgtacgataa aggtcttata gggcagaaga agtacactcg gctgacccgg 2820
agaactggct tccaggacga cgagctcggt cagttcatct caagacagat cgtggaaact 2880
agacaagcaa caaaggagac tgctaacctg ctcaagacaa tatgtaagaa ctccgagatc 2940
atgtattcca aagccgagaa cgcaagtcgg tttaggcaag agttcgacat cgtgaagtgt 3000
agggcggtga acgatcttca tcatatgcac gatgcctaca tcaacatcat agtgggggaa 3060
gtgtataaca ccaagttcac gaaggacct atgaatttcg taaagaagca ggaaaaggcg 3120
cggagctaca atctcgagaa tatgttcaag tacgatgtga aacgtggcgg atacaccgct 3180
tggatcgccg atgacgagaa gggcaccgtg aagaacgcga gtattaaacg tatccggaag 3240
gagctggaag gcacaaatta taggttcaca agaatgaact acattgagtc tggagcgctt 3300
ttcaacgcca ctctccagcg gaagaataag ggctccagac ccctgaagga caaaggcccc 3360
aaatcttcca tcgagaagta cggcggtctac acaaacatca ataaagcctg ttctcgctgtt 3420
cttgacatac agctaagaa caagattgag aggaagctga tgcccgtcga gcgtgagatc 3480
tatgccaacaa agaagaacga caagaagctg tccgacgaga tttctcaaa gtacctcaag 3540
gaccgatttg gcacgagga ctacagggtt gtctaccag tggtgaaaa gcgcacactg 3600
ctcaagatcg acggcagcta ctacttcac acaggcggtt ctgataagac cctggatttg 3660
cgatctgctc tgcagctgat tctccctaag aagaacgagt gggcgatcaa acagatcgac 3720
aagtcttcgg aaaacgacta tctgacgatc gagcgtatcc aggaacctgac caggagctg 3780
gtgtataaca ctctcgacat catcgtaaac aagttcaaga ccagtgtctt caagaagtct 3840
ttccttaact tgtttcagga cgacaagatt gagaacattg acttcaagtt taagtccatg 3900
gacttcaagg agaaatgcaa gacacttctc atgctggtca aggcgattcg ggcctccggc 3960
gtgagggcagg atctcaagtc catcgacctc aagctcgatt acggacggct cagttcaaa 4020
accaacaaca tcggcaatta ccaggagttc aagattatta atcagtcctat cactggactg 4080
ttcgagaatg aggtcgatct cctgaagctg 4110

```

```

SEQ ID NO: 9      moltype = DNA length = 4110
FEATURE
misc_feature      1..4110
                  note = Variant 2902 DNA sequence
source            1..4110
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 9
atgaccaagg tgaaggacta ctacataggg ttggacatcg gcacctctag cgtcggtgtg 60
gccgtccacc atgaagccta taacgtgctt aagtttaata gcaagaaaaa gtggggcgctg 120
cgggtgttgc acgacgctaa gacggcagag gagcgtaggg gccagcgagg agcaagacga 180
cgtctggatc ggaagaagga gagactcagc ctgctgcagg actctctcgc cgaagaggtta 240
gcaaaggctc accccaactt ctctctcagg ctggacaatt ccgatctgta catggaagat 300
aaggaccaga aactgaaaag caaatatata ctgttcaacg acaaggactt caaggataag 360
aattttcata agaagtaccc cacaatacat cacctgtcta tggatctgat caggagacga 420
agtaagaagg acatccggct cgtctacctg gcctgtcact atttgtctca gaacagggg 480
catttcactc tcgagggccg gaagttcgac actaaatcaa gcttcgagaa cagtttgaac 540
gagctcaaa gttcattgaa cgacgagtat ggactggacc tcgaatttga caacgagaac 600
ctgattaaac tcttgactga cccaaaactc aataaaacgg ccaagaagaa ggagctgaag 660
tccgtaatcg gcgacacca gttctctcaa gccgtttccg cgataatgat cggctctagc 720
cagaaactcg tgcacttggt cgagaacccc gaggatctcg acgactctgc gataaagtc 780
gttgactctc caactacctt ttctgacgac aagtaactct actatgaact cgctctgggt 840
gacaagatcg ctctggtcaa catccttaag gaaatttacg atagctccat cctcgagaac 900
ctgctcaaa gggcagacaa gtctaaggac ggtaacaaat atatcagtaa tgcattcgtg 960
aagaagtaca ataaacacgg acaagatctg aaagagtcca aacgtctggt acgacaatat 1020
cacaagagtg cgtattttga tattttcaga tccgagaagg tgaatgacaa ttacgtcagc 1080
tacactaaaa gctcaattag caacaataaa ccgctcaaa caaacaagtt cactgatcaa 1140
gaggccttct acaaatctgc caagaacat ctggagacaa tcaagtataa gatcaacaag 1200
ttaaaccgct ccaaggcaga tctggagctg attgacggga tgcgtcgagg catggagtct 1260
aagaacttta tgcccaaaat taagtcaggt gacaacgggg tgattccata ccagctcaag 1320
ctgatggaat tgaacaaaat actcgagaat cagtcaaaag atcacgagtt cctcaatgtc 1380
agcgacgagt acggctccgt gtgtgataaa atcgcatcta tcatggagtt ccgtatcccc 1440
tactacgtgg gacccctgaa ccccaatagc aagtacgcct ggatcaagaa gcagaaagat 1500
agtgaagata ctccctggaa ctccaaggac gtctgtggac tcgactccag cagagaggag 1560
ttcattgact cactgtctcg acgtgtactt taccttaagg acgagaaggt ccttcccaaa 1620
gcttcttttc tgtataacga atacatgggt ctgaacgagc tgaataacct gaagttgaac 1680
gaccttccca tcaccgagga gatgaagaag aagatatttg accagttgtt caaaacaaga 1740
aagaaggcca cctttaaagc ggtggcaaac ctgctgaaga aggagttcaa catcaacggc 1800
gagattctcg tctctccgac cgacggtgac ttaagcagg gcttgaactc atacaatgac 1860
ttcaaaagcta tcgtggcgga taaagtcgat tccgatgatt accgggacaa gattgaggag 1920
atcattaaac tgatagttct ttacgggtgac gataaagatt accttcagaa gaagattaaa 1980
gctgggtatg gaaaataact caccgacagt gagattaaga aaatggcggg gctgaactac 2040
aaggattggg gaaggctctc aaagaagggt ctgacgggac tccagggtgc aaacaagatc 2100
actggagagc ggggctccat tattcacttc atgaggggat ataaccttaa tctgatggag 2160
cttatgtcag cttcatttac gttcaccgaa gagatacaga aacttaaccc cgtggatgac 2220
cgcaagctgt catacgaat ggtggacgaa ctgtaccttt ctcccagtg gaaacggatg 2280
ctctggcagc cctcgccgat cgtcgacgag ataaagaaca tcatgggaac cgacagtaag 2340
aagattttca tcgagatggc tccgggtaag gaagaggtga aagcccgcaa ggagtcagg 2400
aagaaccac tcgtgaagtc ctataaagac ggaaagaagg cattoatcag cgagattggc 2460
gaggagaggt actcttactt gctttctgag atagaggggt aggaagagaa taagtttcca 2520

```

-continued

tgggataaac	tgtaccttta	ttatactcaa	ctgggtcgct	gcatgtactc	tttggaaacct	2580
atcgacatat	ctgagctgtc	ttcaaagaat	atttacgata	aggatcatat	ctaccccaaa	2640
agcaagattt	acgacgacag	tatcgagaat	aggggtcgctg	tgaagaagga	ccttaactcc	2700
aagaagggtg	acagctatcc	tatcccagac	gaaatcctga	acaagaactg	ttacgcctac	2760
tggagagatc	tgtacgataa	aggtctttat	gggcagaaga	agtacactcg	gctgaccccg	2820
agaactggct	tcacggacga	cgagctcggt	cagttcatct	caagacagat	cgtggaaact	2880
agacaagcaa	caaaggagac	tgctaacctg	ctcaagacaa	tatgtaagaa	ctccgagatc	2940
gtgtattcca	aagcogagaa	cgcaagtcgg	tttaggcaag	agttcgacat	cgtgaagtgt	3000
agggcggtga	acgatcttca	tcatatgcac	gatgcctaca	tcaacatcat	agtgggggaa	3060
gtgtataaca	ccaagtccac	gaaggaccct	atgaatttcg	taaagaagca	ggaaaaggcg	3120
cggagctaca	atctcgagaa	tatgttcaag	tacgatgtga	aacgtggcgg	atacacgcgt	3180
tggatcgccg	atgacgagaa	gggcaccgtg	aagaacgcga	gtattaaacg	tatccggaag	3240
gagctggaag	gcacaaatta	taggttcaca	agaatgaact	acattgagtc	tggagcgctt	3300
ttcaacgcga	ctctccagcg	gaagaataag	ggctccagac	ccctgaagga	caaaggcccg	3360
aaatcttcca	tcgagaagta	cggcggctac	acaacatca	ataaagcctg	tttcgctgtt	3420
cttgacatca	agtctaagaa	caagattgag	aggaagctga	tgcccgtcga	gcgtgagatc	3480
tatgccaaac	agaagaacga	caagaagctg	tcgcacgaga	ttttctcaaa	gtacctcaag	3540
gaccgatttg	gcacgagaga	ctacaggggt	gtctaccacg	tggtgaaagt	gcgcacactg	3600
ctcaagatcg	acggcagcta	ctacttcac	acaggcgggt	ctgataagac	cctggagttg	3660
cgatctgtct	tgcagctgat	tctccctaag	aagaacgagt	gggcgatcaa	acagatcgac	3720
aagttctccg	aaaacgacta	tctgacgatc	gagcgtatcc	aggacctgac	caggagctg	3780
gtgtataaca	ctttcgacat	catcgtcaac	aagttcaaga	ccagtgtctt	caagaagtct	3840
ttccttaact	tgtttcagga	cgacaagatt	gagaacattg	acttcaagtt	taagtccatg	3900
gacttcaagg	agaaatgcaa	gacacttctc	atgctgggtc	aggcgattcg	ggcatccggc	3960
gtgaggcagg	atctcaagtc	catcgacctc	aagttcgatt	acggacggct	cagttcaaac	4020
accaacaaca	tcggcaatta	ccaggagttc	aagattatta	atcagttccat	cactggactg	4080
tcgagaatg	aggtcgatct	cctgaagctg				4110

SEQ ID NO: 10 moltype = DNA length = 4110
FEATURE Location/Qualifiers
misc_feature 1..4110
 note = Variant 3954 DNA sequence
source 1..4110
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 10

atgaccaagg	tgaaggacta	ctacataggc	ttggacatcg	gcacctctag	cgtcgggttg	60
gccgtcaccc	atgaagccta	taacgtgctt	aagtttaata	gcaagaaaaa	gtggggcgctg	120
cggctgttgc	acgacgctaa	gacggcagag	gagcgtatgg	gccagcgagg	agcaagacga	180
cgtctggatc	ggaagaagga	gagactcagc	ctgctgcagg	acttcttcgc	cgaagagggt	240
gcaaaggctc	accccaactt	ctctctcagg	ctggacaatt	ccgatctgta	catggaagat	300
aaggaccaga	aactgaaaaa	caaatataca	ctgttcaaac	acaaggactt	caaggataag	360
aattttcata	agaagtagcc	cacaatacat	cacctgctga	tggatctgat	cgaggacgac	420
agtaagaagg	acatccggct	cgtctacctg	gcctgtcact	atttgcctca	gaacaggggt	480
catttcatct	tcaggggcca	gaagttcgac	actaaatcaa	gcttcgagaa	cagtttgaac	540
gagctcaaa	ttcatttgaa	cgacgagtat	ggactggacc	tcgaatttga	caacgagaac	600
ctgattaaca	tcttgactga	ccccaaactc	aataaaaacg	ccaagaagaa	ggagctgaag	660
tcctgaatcg	gcgacaccaa	gttctccaaa	gccgtttccg	cgataatgat	cggctctagc	720
cagaaactcg	tgcacttggt	cgagaacccc	gaggatttgc	acgactctgc	gataaagtcc	780
gttgacttct	caactacctc	tttcgacgac	aagtactctg	actatgaact	cgtctcgggt	840
gacaagatcg	ctctgggtcaa	catccttaag	gaaattttac	atagctcaat	cctcgagaa	900
ctgctcaaa	atgcagacaa	gctaaggac	ggtaacaaat	atatcagtaa	tgcattctgt	960
aagaagtaca	atgaacacgg	acaagatctg	aaagagtcca	aacgtctcgt	acgacaatat	1020
cacaagagtg	cgtattttga	tattttcaga	tcgagaagg	tgaatgacaa	ttacgtcagc	1080
tacactaaaa	gctcaattag	caacaataaa	cgcgtcaaa	caaacaagtt	cactgatcaa	1140
gaggccttct	acaaattcgc	caagaaacat	ctggagacaa	tcaagtataa	gatcaacaag	1200
gtaaacggct	ccaaggcaga	tctggagctg	attgacggga	tgctgcggga	catggagttc	1260
aagaacttta	tgcccaaaat	taaaagctcg	gataacggcg	tgattccgta	tcagctgaaa	1320
ctgatggaat	tgaacaaaat	actcgagaat	cagtcaaaag	atcacgagtt	cctcaatgtc	1380
agcgacgagt	acggctccgt	gtgtgataaa	atcgcatcta	tcatggaatt	taccattccg	1440
tattacgtgg	gcccgttgaa	cccgaacagc	aaatatggct	ggatcaagaa	gcagaagat	1500
agtgagatta	ctccctggaa	cttcaaggac	gtcgtggacc	tcgactccag	cagagaggag	1560
ttcattgact	cactgatctg	acgctgtact	taccttaagg	acgagaaggt	ccttcccaaa	1620
gcttctttgc	tgtataacga	atacatgggt	ctgaacgagc	tgaataacct	gaagtgaac	1680
gaccttccca	tcaccgagga	gatgaagaag	aagatatttg	accagttgtt	caaaacaaga	1740
aagaaggctc	cccttaaaag	ggtggcaaac	ctgctgaaga	aggagttcaa	catcaacggc	1800
gagattctgc	tctctgggac	cgaacgtgac	ttcaagcagg	gcttgaactc	atacaatgac	1860
ttcaaaagcta	tcgtgggcga	taaagtcgat	tccgatgatt	accgggacaa	gattgaggag	1920
atcattaaac	tgatagttct	ttacgggtgac	gataaagagt	accttcagaa	gaagattaaa	1980
gctgggtatg	gaaaaacttt	caccgacagt	gagattaaag	aaatggcggg	gctgaactac	2040
aaggattggg	gaaggctctc	agaaaagctg	ctgacgggac	tcgagggtgc	aaacaagatc	2100
actggagagc	ggggtctccat	tattcacttc	atgaggggaat	ataaccttaa	ttgtatggag	2160
cttatgtcag	cttcattttac	gttcaccgaa	gagatacaga	aacttaaccc	cgtggatgac	2220
cgcaagctgt	catacgaat	ggtggacgaa	ctgtaccttt	ctccagatgt	gaaacggatg	2280
ctctggcagt	ccctgcgcac	cgtcgacgag	ataaagaaca	tcattgggaa	cgacagtaag	2340
aagattttca	tcgagatggc	tcggggtaag	gaagaggtga	aagcccgcaa	ggagtcaagg	2400

-continued

```

aagaaccaac tgctgaagtt ctataaagac ggaaagaagg cattcatcag cgagattggc 2460
gaggagaggt actcttactt gctttctgag atagaggggtg aggaagagaa taagtttcga 2520
tgggataacc tgtaccttta ttatactcaa ctgggtcgct gcatgtactc tttggaacct 2580
atcgacatat ctgagctgtc ttcaaagaat atttacgatac aggatcatat ctaccccaaa 2640
agcaagattt acgacgacag tatcgagaat aggggtgctgg tgaagaagga ccttaactcc 2700
aagaagggtt acagctatcc tatcccagac gaaatcctga acaagaactg ttacgcctac 2760
tggaagatcc tgtacgataa aggtctttatc gggcagaaga agtacactcg gctgaccggy 2820
agaactggct tcacggacga cgagctcggt cagttcatct caagacagat cgtggaaact 2880
agacaagcaa cagtcgagac tgcataacctg ctcaagacaa tatgtaagaa ctccgagatc 2940
gtgtattcca aagccgagaa cgcaagtcgg tttaggcaag agttcgacat cgtgaagtgt 3000
agggcggtga acgatcttca tcatatgcac gatgcctaca tcaacatcat agtggggaac 3060
gtgtataaca ccaagttcac gaaggacct atgaatttcg taaaggtcca ggaaaaggcg 3120
cggagctaca atctcgagaa tatgttcaag tacgatgtga aacgtggcgg ataccgcgt 3180
tggaatcgcg atgacgagaa gggcacgctg aagaacgca gtattaaacg tatccggaag 3240
gagctggaag gcaaaaatta taggttcaca agaatagaac acattgagtc tggagcgctt 3300
ttcaacgcca ctctccagcg gaagaataag ggctccagac ccctgaagga caaaggcccg 3360
aaatcttcca ctcgagaaga cggcggtctac acaaacatca ataaagcctg ttctcgctgt 3420
cttgacatca agtctaagaa caagattgag aggaagctga tgcccgctga gcgtgagatc 3480
tatgccaaac agaagaacga caagaagctg tccgacgaga tttctcaaa gtacctaaag 3540
gaccgatttg gcatcgagga ctacagggtt gtctaccagc ttgtgaaaaa gcgcacactg 3600
ctcaagatcg acggcagcta ctacttcac acaggcggtt ctgataagac cctggagttg 3660
cgatctgctc tgcagctgat tctccctaag aagaacgagt gggcgatcaa acagatcgac 3720
aagtcttccg aaaaacgacta tctgacgatac gagcgtatcc aggacctgac cgaggagctg 3780
gtgtataaca ctttcgacat catcgtaaac aagtccaaga ccagtgtctt caagaagtct 3840
ttctttaact tggttcagga gcacaagatt gagaacattg acttcaagtt taagtccatg 3900
gacttcaagg agaattgcaa gacacttctc atgctgggta aggcgattcg ggcactccgc 3960
gtgagggcagg atctcaagtc catcgacctc aagctgtatt acggacggct cagttcaaag 4020
accaacaaca tcggcaatta ccaggagttc aagattatta atcagtcctc cactggactg 4080
ttcgagaatg aggtcgactc cctgaagctg                                     4110

```

```

SEQ ID NO: 11      moltype = AA length = 1410
FEATURE           Location/Qualifiers
REGION            1..1410
                  note = Wild-type OMNI-50 nuclease with tags
source            1..1410
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 11
MPKKKKRVGT TKVKDYYIGL DIGTSSVGWA VTDEAYNVLK FNSKKMWGVR LFDDAKTAE 60
RRGQRGARRR LDRKKERLSL LQDFFAEVA KVDPNFFLRL DNSDLYMEDK DQKLKSKYTL 120
FNDKDFKDKN PHKKYPTIHH LLMDLIEDDS KKDRLVLYLA CHYLLKNRGR FIFEGQKFDT 180
KSSFENSLNE LKVHLNDEYG LDLEFDNENL INILTDPKLN KTAKKKELKS VIGDTKFLKA 240
VSAIMIGSSQ KLVDLFENPE DFDDSAIKSV DFSTTSFDDK YSDYELALGD KIALVNLKE 300
IYDSSILENL LKEADKSKDG NKYISNAFVK KYNKHGQDLK EFKRLVRQYH KSAYFDIFRS 360
EKVNDNYVSY TKSSISNNKR VKANKFTDQE AFYKFAKKHL ETIKYKINKV NGSKADLELI 420
DGMLRDMEFK NFMPKIKSSD NGVIPYQLKL MELNKILENQ SKHHEFLNVS DEYGSVCDKI 480
ASIMEFRIPY YVGPLNPNSK YAWIKKQKDS EITPWNFKDV VDLDSREEF IDSLIGRCTY 540
LKDEKVLPAK SLLYNEYMVL NELNNLKLND LPITEEMKKK IFDQLFKTRK KVTLKAVANL 600
LKKEFNINGE ILLSGTDGDF KQGLNSYNDF KAIVGDKVDS DDYRDKIEEI IKLIVLYGDD 660
KSYLQKKIKA GYGYFTDSE IKKMAGLNYK DWGRLSKKLL TGLEGANKIT GERGSIIHFM 720
REYNLNLML MSASFTFTEE IQKLNPVDDR KLSYEMVDEL YLSPSVKRM LQSLRIVDEI 780
KNIMGTDSKK IFIEMARGEK EVKARKESRK NQLLKFKYDG KKAIFSEIGE ERSYLLSEI 840
EGEENKFRW DNLYLYYTQL GRCMYSLEPI DISELSSKNI YDQDHIYPKS KIYDDSIENR 900
VLVKKDLNSK KGNYSYIPDE ILNKNCYAYW KILYDKGLIG QKKYTRLTRR TGFTDDELVQ 960
FISRQIVETR QATKETANLL KTKCNSEIV YSKAENASRF RQEPDIVKCR AVNDLHHMHD 1020
AYINIIVGNV YNTKFTKDPM NFVKKQEKAR SYNLENMFKY DVKRGGYTAW IADDEKGTVK 1080
NASIKRIRKE LEGTNYRFR MNYIESGALF NATLQRKNKG SRPLKDKGPK SSIEKYGGYT 1140
NINKACFAVL DIKSNKIER KLMPVEREIV AKQKNDKKLS DEIFSKYLLD RFGIEDYRVV 1200
YPVVKMRTL LKIDGSYYFIT GGSDKTLELR SALQLILPKK NEWAIKQIDK SSENDYLTIE 1260
RIQDLTEELV YNTFDIIVNK FKTSVPKKSF LNLQDDKIE NIDPKFKSMD FKEKCKTLLM 1320
LVKAIRASGV RQDLKSIDLK SDYGRLSSTK NNIGNYQEFK IINQSITGLF ENEVDLLKLK 1380
SYPYDVPDYA AAPKKKRKVA ASHHHHHHHH                                     1410

```

```

SEQ ID NO: 12      moltype = AA length = 1410
FEATURE           Location/Qualifiers
REGION            1..1410
                  note = Variant 3795 with tags
source            1..1410
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 12
MPKKKKRVGT TKVKDYYIGL DIGTSSVGWA VTDEAYNVLK FNSKKMWGVR LFDDAKTAE 60
RRGQRGARRR LDRKKERLSL LQDFFAEVA KVDPNFFLRL DNSDLYMEDK DQKLKSKYTL 120
FNDKDFKDKN PHKKYPTIHH LLMDLIEDDS KKDRLVLYLA CHYLLKNRGR FIFEGQKFDT 180
KSSFENSLNE LKVHLNDEYG LDLEFDNENL INILTDPKLN KTAKKKELKS VIGDTKFLKA 240
VSAIMIGSSQ KLVDLFENPE DFDDSAIKSV DFSTTSFDDK YSDYELALGD KIALVNLKE 300

```

-continued

IYDSSILENL	LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	360
EKVNDNYVSY	TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	420
DGMLRDMEFK	NFMPKIKSSD	NGVIPYQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	480
ASIMEFIIPY	YVGPLNPNK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	540
LKDEKVLPA	SLLNHEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	600
LKKEFNINGE	ILLSGTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	660
KSYLQKKIKA	GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	720
REYNLNLML	MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	780
KNIMGTDSKK	IFIEMARGKE	EVKARKESRK	NVLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	840
EGEENKFRW	DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	900
VLVKKDLNSK	KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	960
FISRQIVETR	QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	1020
AYINIIIVGNV	YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	1080
NASIKRIRKE	LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	1140
NINKACFAVL	DIKSKNKIER	KLMPVEREYI	AKQKNDKLS	DEIFSKYLKD	RFGIEDYRVV	1200
YPVVKMRTL	LKIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	1260
RIQDLTEELV	YNTFDIIVNK	FKTSVPKKS	LNLFDQDDKIE	NIDFKFKSMD	FKEKCKTLLM	1320
LVKAIRASGV	RQDLKSIDLK	SDYGRLSST	NNIGNYQEFK	IINQSITGLF	ENEVDLLKLG	1380
SYPYDVPDIA	AAPKKKRKVA	ASHHHHHHHH				1410

SEQ ID NO: 13 moltype = AA length = 1410
 FEATURE Location/Qualifiers
 REGION 1..1410
 note = Variant 3942 with tags
 source 1..1410
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 13

MPKKKKRVGT	TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAAE	60
RRGQRGARR	LDRKKERLSL	LQDFFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	120
FNDKDFKDN	PHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	180
KSSFENSLNE	LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	240
VSAIMIGSSQ	KLVDLFENPE	DFDSSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	300
IYDSSILENL	LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	360
EKVNDNYVSY	TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	420
DGMLRDMEFK	NFMPKIKSSD	NGVIPWQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	480
ASIMEFRIIPY	YVGPLNPNK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	540
LKDEKVLPA	SLLYNEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	600
LKKEFNINGE	ILLSGTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	660
KSYLQKKIKA	GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	720
REYNLNLML	MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	780
KNIMGTDSKK	IFIEMARGKE	EVKARKESRK	NQLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	840
EGEENKFRW	DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	900
VLVKKDLNSK	KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	960
FISRQIVETR	QATKETANLL	KTICKNSEIM	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	1020
AYINIIIVGNV	YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	1080
NASIKRIRKE	LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	1140
NINKACFAVL	DIKSKNKIER	KLMPVEREYI	AKQKNDKLS	DEIFSKYLKD	RFGIEDYRVV	1200
YPVVKMRTL	LKIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	1260
RIQDLTEELV	YNTFDIIVNK	FKTSVPKKS	LNLFDQDDKIE	NIDFKFKSMD	FKEKCKTLLM	1320
LVKAIRASGV	RQDLKSIDLK	SDYGRLSST	NNIGNYQEFK	IINQSITGLF	ENEVDLLKLG	1380
SYPYDVPDIA	AAPKKKRKVA	ASHHHHHHHH				1410

SEQ ID NO: 14 moltype = AA length = 1410
 FEATURE Location/Qualifiers
 REGION 1..1410
 note = Variant 2902 with tags
 source 1..1410
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 14

MPKKKKRVGT	TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAAE	60
RRGQRGARR	LDRKKERLSL	LQDFFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	120
FNDKDFKDN	PHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	180
KSSFENSLNE	LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	240
VSAIMIGSSQ	KLVDLFENPE	DFDSSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	300
IYDSSILENL	LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	360
EKVNDNYVSY	TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	420
DGMLRDMEFK	NFMPKIKSSD	NGVIPYQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	480
ASIMEFRIIPY	YVGPLNPNK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	540
LKDEKVLPA	SLLYNEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	600
LKKEFNINGE	ILLSPTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	660
KSYLQKKIKA	GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKVL	TGLQGANKIT	GERGSIHFH	720
REYNLNLML	MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	780
KNIMGTDSKK	IFIEMARGKE	EVKARKESRK	NQLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	840
EGEENKFRW	DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	900

-continued

VLVKKDLNSK	KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	960
FISRQIVETR	QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	1020
AYINIIVGNV	YNTKFTKDPM	NFVKKQEKAR	SYNLENMPKY	DVKRGGYTAW	IADDEKGTVK	1080
NASIKRIRKE	LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	1140
NINKACFAVL	DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	1200
YPVVKMRTL	L KIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	1260
RIQDLTEELV	YNTFDIIVNK	FKTSVFKKSF	LNLQDDKIE	NIDPKFKSMD	FKEKCKTLLM	1320
LVKAIRASGV	RQDLKSIDLK	SDYGRLLSSKT	NNIGNYQEFK	IINQSITGLF	ENEVDLLKLK	1380
SYPYDVDPYA	AAPKKKRKVA	ASHHHHHHHH				1410

SEQ ID NO: 15 moltype = AA length = 1410
 FEATURE Location/Qualifiers
 REGION 1..1410
 note = Variant 3954 with tags
 source 1..1410
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 15

MPKKKKRVGT	TKVKDYIYGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	60
RRGQRGARR	LDRKKERLSL	LQDFFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	120
FNDKDFKDN	PHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	180
KSSFENSLNE	LKVHLNDEYG	LDLEFDNENL	INILTDPLKN	KTAKKKELKS	VIGDTKFLKA	240
VSAIMIGSSQ	KLVDLFENPE	DFDSSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	300
IYDSSILENL	LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRYQH	KSAYFDIFRS	360
EKVNDNYVSY	TKSSISNNKR	VKANKPTDQE	AFYKPAKKHL	ETIKYKINKV	NGSKADLELI	420
DGMLRDMEPK	NFMPKIKSSD	NGVIPYQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	480
ASIMEFTIPY	YVGPLNPNSK	YGWIKKQKDS	EITPWNFKDV	VLDLSSREEF	IDSLIGRCTY	540
LKDEKVLPA	SLLYNEVMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	600
LKKEFNINGE	ILLSGTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	660
KSYLQKKIKA	GYGKYFTDSE	IKKMAGLNYK	DWGRLESEKL	TGLEGANKIT	GERGSIHF	720
REYNLNCMEL	MSASFTFTEE	IQKLNPVDDR	KLSEYEMVDEL	YLSPSVKRML	WQSLRIVDEI	780
KNIMGTDSKK	IFIEMARGKE	EVKARKESRK	NQLLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	840
EGEENKFRW	DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	900
VLVKKDLNSK	KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	960
FISRQIVETR	QATVETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	1020
AYINIIVGNV	YNTKFTKDPM	NFVKVQEKAR	SYNLENMPKY	DVKRGGYTAW	IADDEKGTVK	1080
NASIKRIRKE	LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	1140
NINKACFAVL	DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	1200
YPVVKMRTL	L KIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	1260
RIQDLTEELV	YNTFDIIVNK	FKTSVFKKSF	LNLQDDKIE	NIDPKFKSMD	FKEKCKTLLM	1320
LVKAIRASGV	RQDLKSIDLK	SDYGRLLSSKT	NNIGNYQEFK	IINQSITGLF	ENEVDLLKLK	1380
SYPYDVDPYA	AAPKKKRKVA	ASHHHHHHHH				1410

SEQ ID NO: 16 moltype = DNA length = 4233
 FEATURE Location/Qualifiers
 misc_feature 1..4233
 note = Wild-type OMNI-50 nuclease with tags DNA sequence
 source 1..4233
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 16

atgcctaaga	agaagagaaa	gggtgggtacc	accaaaggta	aggactacta	cataggcttg	60
gacatcgcca	cctctagcgt	cgggtggggcc	gtcaccgatg	aagcctataa	cgtgcttaag	120
ttaataagca	agaaaatgtg	gggcgtgctg	ctgttcgacg	acgctaagac	ggcagaggag	180
cgtaggggcc	agcaggagac	aagacgacgt	ctggatcgga	agaaggagag	actcagcctg	240
ctgcaggact	tcttcgccga	agaggttagca	aaggctcgacc	ccaacttctt	cctcaggctg	300
gacaattccg	atctgtacat	ggaagataag	gaccagaaac	tgaaaagcaa	atatacactg	360
ttcaacgaca	aggacttcaa	ggataagaat	tttcataaga	agtacccccc	aatacatcac	420
ctgctgatgg	atctgatcga	ggacgacagt	aagaaggaca	tccggctcgt	ctacctggcc	480
tgtcactatt	tgcctcaaga	cagggggtcat	ttcatcttcg	agggccagaa	gttcgacact	540
aaatacaagc	tcgagaacag	tttgaacgag	ctcaaaagtt	atttgaacga	cgagatggga	600
ctggacctcg	aatttgacaa	cgagaacctg	attaacatct	tgactgaccc	aaaactcaat	660
aaaacggcca	agaagaagga	gctgaagtcc	gtaatcgccg	acaccaagtt	cctcaaaagc	720
gtttccgcga	taatgatcgg	ctctagccag	aaactcgctg	acttggttcga	gaaccccgag	780
gatttcgacg	actctgcgat	aaagtccgtt	gactctctca	ctacctcttt	cgacgacaag	840
tactctgact	atgaactcgc	tctgggtgac	aagatcgctc	tggtcaacat	ccttaaggaa	900
atttacgata	gctccatcct	cgagaacctg	ctcaaaagag	cagacaagtc	taaggacggt	960
aacaaatata	tcagtaatgc	attcgtgaag	aagtacaata	aacacggaca	agatctgaaa	1020
gagttcaaac	gtctggtacg	acaatatcac	aagagtgctg	attttgatat	tttcagatcc	1080
gagaagggtg	atgacaatta	cgctcagctac	actaaaagct	caatttagcaa	caataaacgc	1140
gtcaaaagca	acaagttcac	tgatcaagag	gcctcttaca	aattcgccaa	gaaacatctg	1200
gagacaatca	agataagat	caacaaggta	aacggctcca	aggcagatct	ggagctgatt	1260
gacgggatgc	tgccggacat	ggagttcaag	aactttatgc	ccaaaataa	gtccagtgac	1320
aacgggggtg	ttccatacca	gctcaagctg	atggaattga	acaaaatact	cgagaatcag	1380
tcaagcatc	acgagttcct	caatgtcagc	gacgagtagc	gctccgtgtg	tgataaaatc	1440
gcatctatca	tgaggttccg	tatcccctac	tacgtgggac	ccctgaaccc	caatagcaag	1500

-continued

tacgcctgga	tcaagaagca	gaaagatagt	gagattactc	cctggaactt	caaggacgtc	1560
gtggacctcg	actccagcag	agaggagttc	attgactcac	tgatcggacg	ctgtacttac	1620
cttaaggacg	agaaggctct	tcccaagct	tctttgctgt	ataacgaata	catgggtgctg	1680
aacgagctga	ataacctgaa	gttgaacgac	cttcccatca	cagaggagat	gaagaagaag	1740
atatttgacc	agttgttcaa	aacaagaaa	aaggtcaccc	ttaaagcggg	ggcaaacctg	1800
ctgaagaagg	agttcaacat	caacggcgag	attctgctct	ctgggaccca	cgggtgacttc	1860
aagcagggct	tgaactcata	caatgacttc	aaagctatcg	tgggcgataa	agtcgattcc	1920
gatgattacc	gggacaagat	tgaggagatc	attaaactga	tagttcttta	cggtgacgat	1980
aagagttacc	ttcagaagaa	gattaaagct	gggtatggaa	aatacttcac	cgacagttag	2040
attaagaaaa	tggcggggct	gaactacaag	gattggggaa	ggctctcaaa	gaagctgctg	2100
acgggactcg	aggggtgcaa	caagatcact	ggagagcggg	gctccattat	tcacttcacg	2160
agggaaataa	accttaactc	gatggagctt	atgtcagctt	catttacgtt	caccgaagag	2220
atacagaaac	ttaaccccg	ggatgaccgc	aagctgtcat	acgaaatggt	ggacgaactg	2280
tacctttctc	ccagtgtgct	acggatgctc	tggcagtcct	tgcgcacgt	cgacgagata	2340
aagaacatca	tgggaaccca	cagtaagaag	attttcatcg	agatggctcg	gggttaaggaa	2400
gaggtgaaag	cccgaagga	gtcaaggga	aaccaactgc	tgaagttcta	taaagacgga	2460
aagaaggcat	tcatcagcga	gattggcgag	gagaggtaact	cttacttgct	ttctgagata	2520
gagggtaggg	aagagaataa	gtttcgatgg	gataacctgt	acctttatta	tactcaactg	2580
ggtcgctgca	tgtactcttt	ggaactctatc	gacatatctg	agctgtcttc	aaagaatatt	2640
tacgatcagg	atcatatcta	ccccaaaagc	aagatttacg	acgacagtat	cgagaatagg	2700
gtgctggtga	agaaggacct	taactccaag	aagggttaaca	gctatcctat	cccagacgaa	2760
atcctgaaca	agaactgtta	gcctactgg	aagatcctgt	acgataaagg	tcttatcggg	2820
cagaagaagt	acactcgctt	gacccggaga	actggcttca	cggacgacga	gctcgttcag	2880
ttcatctcaa	gacagatcgt	ggaactaga	caagcaacaa	aggagactgc	taacctgtct	2940
aagacaatat	gtaagaactc	cgagatcgtg	tattccaaag	ccgagaacgc	aagtcggttt	3000
aggcaagagt	tcgacatcgt	gaagtgtagg	gcggtgaacg	atcttcatca	tatgcacgat	3060
gctacatca	acatcatagt	ggggaacgtg	tataaacacca	agttcaccga	ggacctatg	3120
aatttcgtaa	agaagcagga	aaaggcgcgg	agctacaatc	tcgagaatat	gttcaagtac	3180
gatgtgaaac	gtggcgagata	caccgcttgg	atcgccgatg	acgagaaggg	caccgtgaag	3240
aacgcgagta	ttaaacgtat	ccggaaggag	ctggaaggca	caaatattag	gttcacaaga	3300
atgaactaca	ttagtctctg	agcgcctttc	aacgccaactc	tccagcggaa	gaataagggc	3360
tccagacccc	tgaaggacaa	aggcccga	tcttccatcg	agaagtacgg	cggctacaca	3420
aacatcaata	aagcctgttt	cgctgttctt	gacatcaagt	ctaagaacaa	gattgagagg	3480
aagctgatgc	ccgtcgagcg	tgagatctat	gccaaacaga	agaacgacaa	gaagctgtcc	3540
gacgagattt	tctcaaaagta	cctcaaggac	cgatttgcca	tcgaggacta	cagggttgct	3600
taccagtggt	tgaaaatgcg	cacactgctc	aagatcgacg	gcagctacta	cttcatcaca	3660
ggcggttctg	ataagaccct	ggagttgcga	tctgctctgc	agctgattct	ccctaagaag	3720
aacgagtggt	cgatcaaaaca	gatcgacaag	tcttccgaaa	acgactatct	gacgatcgag	3780
cgtatccagg	acctgaccca	ggagctgggt	tataaacactt	tcgacatcat	cgtcaacaag	3840
ttcaagacca	gtgtcttcaa	gaagtctttc	cttaacttgt	ttcaggacga	caagattgag	3900
aacattgact	tcaagtttaa	gtccatggac	ttcaaggaga	aatgcaagac	acttctcatg	3960
ctggtcaagg	cgattcgggc	atccggcgtg	aggcaggatc	tcaagtcac	cgacctcaag	4020
tctgattacg	gacggctcag	ttcaaaagacc	aacaacatcg	gcaattacca	ggagttcaag	4080
attattaatc	agtcctacac	tggactgttc	gagaatgagg	tcgatctcct	gaagctggga	4140
tctaccat	acgatgttcc	agattacgcg	gccgctccaa	aaaagaaaag	aaaagttgct	4200
gctagccatc	atcaccatca	ccatcatcat	taa			4233

SEQ ID NO: 17 moltype = DNA length = 4233
FEATURE Location/Qualifiers
misc_feature 1..4233
 note = Variant 3795 with tags DNA sequence
source 1..4233
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 17

atgcctaaga	agaagagaaa	gggtgggtacc	accaaggtga	aggactacta	cataggcttg	60
gacatcggca	cctctagcgt	cgggtggggcc	gtcaccgatg	aagcctataa	cgtgcttaag	120
tttaatatga	agaaaatggt	gggcgtgccc	ctgttcgacg	acgctaagac	ggcagaggag	180
cgtagggggc	agcagaggag	aagacgacgt	ctggatcgga	agaaggagag	actcagcctg	240
ctgcaggact	tcttcgccga	agaggttagca	aaggtcgacc	ccaacttctt	cctcaggctg	300
gacaattccg	atctgtacat	ggaagataag	gaccagaaac	tgaaaagcaa	atatacactg	360
ttcaacgaca	aggacttcaa	ggataagaat	tttcataaga	agtaccccat	aatacatcac	420
ctgctgatgg	atctgatcga	ggacgacagt	aagaaggaca	tccggctcgt	ctacctggcc	480
tgctactatt	tgctcaagaa	caggggtcat	ttcatcttcg	agggccagaa	gttcgacact	540
aaatcaagct	tcgagaacag	tttgaacgag	ctcaaaagttc	atttgaacga	cgagtatgga	600
ctggacctcg	aatttgacaa	cgagaacctg	attaacatct	tgactgaccc	aaaactcaat	660
aaaacggcca	agaagaagga	gctgaagtcc	gtaatcggcg	acaccaagtt	cctcaaaagc	720
gtttccgcga	taattgatcgg	ctctagccag	aaactcgtcg	acttggtcga	gaaccccgag	780
gatttcgacg	actctgcgat	aaagtccggt	gactttctcaa	ctacctcttt	cgacgacaag	840
tactctgact	atgaactcgc	tctgggtgac	aagatcgctc	tggtcaacat	ccttaaggaa	900
atttaacgata	tgtccatcct	cgagaacctg	ctcaaaagag	cagacaagtc	taaggcgggt	960
aacaaatata	tcagtaatgc	attcgtgaag	aagtacaata	aacacggaca	agatctgaaa	1020
gagttcaaac	gtctgggtacg	acaatatcac	aagagtgcgt	attttgatat	tttcagatcc	1080
gagaagggtg	atgacaatta	cgctcagctac	actaaaagct	caattagcaa	caataaacgc	1140
gtcaaaagcaa	acaagttcac	tgatcaagag	gccttctaca	aattcgccaa	gaaacatctg	1200
gagacaatca	agtataagat	caacaaggta	aacggctcca	aggcagatct	ggagctgatt	1260

-continued

```

gacgggatgc tgcgggacat ggagttcaag aactttatgc ccaaaattaa gtccagtgc 1320
aacgggggtga ttccataacca gctcaagctg atggaattga acaaaatact cgagaatcag 1380
tcaaagcatc acgagttcct caatgtcagc gacgagtagc gctccgtgtg tgataaaatc 1440
gcatctatca tggaaatttat tattccgtat tacgtgggcc cgctgaacc gaacagcaaa 1500
tatgcgtgga tcaagaagca gaaagatagt gagattactc cctggaactt caaggacgtc 1560
gtggacctcg actccagcag agaggagttc attgactcac tgatcggacg ctgtacttac 1620
cttaaggacg agaaggctct tcccaaagct tctttgtctg ataacgaata catggtgctg 1680
aacgagctga ataacctgaa gttgaacgac ctccccatca ccgaggagat gaagaagaag 1740
atatttgacc agttgttcaa aacaagaaag aaggtcacc ttaagcgtt ggcaaacctg 1800
ctgaagaagg agttcaacat caacggcgag attctgctct ctgggaccga cggtagcttc 1860
aagcagggtc tgaactcata caatgacttc aaagctatcg tgggcgataa agtcgattcc 1920
gatgattacc gggacaagat tgaggagatc attaaactga tagttcttta cggtagcat 1980
aagagttacc ttcagaagaa gattaaagct gggtaggaa aatacttcac cgacagtgc 2040
atgaagaaaa tgcgggggtc gaactacaag gattggggaa ggctctcaa gaagctgctg 2100
acgggactcg aggggtgcaaa caagatcact ggagagcggg gctccattat tcaactcatg 2160
agggaaatata acctaatct gatggagctt atgtcagctt cattacgtt caccgaagag 2220
atacagaaac ttaacctcgt ggatgaccgc aagctgtcat acgaaatggg ggacgaactg 2280
tacctttctc ccagtgtgaa acggatgctc tggcagtcct tgcgcactcg cgacgagata 2340
aagaacatca tgggaaccga cagtaagaag attttcatcg agatggctcg gggtaaggaa 2400
gaggtgaaag ccgcgaagga gtcaagggaag aacgtcctga tcaagttcta taaagacgga 2460
aagaaggcat tcatcagcga gattggcgag gagaggtagt ctactctgct tctgagata 2520
gagggtaggg aagagaataa gtttcgatgg gataacctgt acctttatta tactcaactg 2580
ggtcgctgca tgtactcttt ggaacctatc gacatatctg agctgtcttc aaagaatatt 2640
tacgatcagg atcatatcta ccccaaaagc aagatttacg acgacagtat cgagaatagg 2700
gtgctggtga agaaggacct taactccaag aagggttaaca gctatcctat cccagacgaa 2760
atcctgaaca agaactgtta cgctactctg aagatcctgt acgataaagg tcttatcggg 2820
cagaagaagt acactcggct gacccggaga actggttca cggacgacga gctcgttcag 2880
ttcatctcaa gacagatcgt ggaactaga caagcaacaa aggagactgc taacctgtc 2940
aagacaatat gtaagaactc cgagatcgtg tattccaaag ccgagaacgc aagtcgggtt 3000
aggcaagagt tcgacatcgt gaagtgtagg gcggtgaacg atcttcatca tatgcacgat 3060
gctacatca acatcatagt ggggaacgtg tataaacacca agttcacgaa ggaccctatg 3120
aatctcgtaa agaagcagga aaaggcgagg agctacaatc tcgagaatat gttcaagtac 3180
gatgtgaaac gtggcgagata caccgcttgg atcgccgatg acgagaaggg caccgtgaag 3240
aacgcgagta ttaaacgtat ccggaaggag ctggaaggca caaattatag gttcacaaga 3300
atgaactaca ttgagctcgg agcgcttttc aacgccactc tccagcggaa gaataagggc 3360
tccagacccc tgaaggacaa agggccgaaa tcttccatcg agaagtagcg cggtacaca 3420
aacatcaata aagcctgttt cgtgttctt gacatcaagt ctaagaacaa gattgagagg 3480
aagctgatgc ccgctgagcg tgagatctat gccaaacaga agaacgacaa gaagctgtcc 3540
gacgagattt tctcaaagta cctcaaggac cgatttggca tcgaggacta cagggttgtc 3600
taccagtggt tgaaaatgcg cacactgctc aagatcgacg gcagctacta cttcatcaca 3660
ggcggttctg ataagacctt ggagttcgca tctgctctgc agctgattct ccctaagaag 3720
aacgagtggt cgatcaacaa gatcgacaag tcttccgaaa acgactatct gacgatcgag 3780
cgtatccagg acctgaccga ggaagctggtg tataaacactt tcgacatcat cgtcaacaag 3840
ttcaagacca gtgtcttcaa gaagcttttc cttaacttgt ttcaggacga caagattgag 3900
aacattgact tcaagtttaa gtccatggac ttcaaggaga aatgcaagac acttctcatg 3960
ctggtcaagg gcattcggcg atccggcgtg aggcaggatc tcaagtccat cgacctcaag 4020
tctgattacg gacggctcag ttcaaagacc aacaacatcg gcaattacca ggagttcaag 4080
attattaatc agtccatcac tggactgttc gagaatgagg tcgatctcct gaagctggga 4140
tcctaccocat acgatgttcc agattacgag gccgctccaa aaaagaaaag aaaagttgag 4200
gctagccatc atcaccatca ccatcatcat taa 4233

```

```

SEQ ID NO: 18      moltype = DNA length = 4233
FEATURE
misc_feature       1..4233
                    note = Variant 3942 with tags DNA sequence
source              1..4233
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 18
atgctaaaga agaagagaaa ggtgggtacc accaagggtga aggactacta cataggcttg 60
gacatcgcca cctctagcgt cgggtgggcc gtcaccgatg aagcctataa cgtgcttaag 120
tttaaatgca agaaaatgtg ggggtgtggc ctgttcgacg acgctaaaaa cgcggaagag 180
cgccgtggcc agcgcggcgc gcgccgtgct ctggatcgca aaaaggaacg cctcagcctg 240
ctgcaggact tcttcgccga agaggtagca aaggtcgacc ccaacttctt cctcaggctg 300
gacaattccg atctgtacat ggaagataag gaccagaaac tgaaaagcaa atatacactg 360
ttcaacgaca aggacttcaa ggataagaat ttctataaga agtaccctac aatacatcac 420
ctgctgatgg atctgatcga ggacgacagt aagaaggaca tccggctcgt ctacctggcc 480
tgtcactatt tgcctaaaga cagggttcac ttcatcttcg agggccagaa gttcgacact 540
aaatcaagct tcgagaacag tttgaacgag ctcaaagtgc atttgaacga cgagtatgga 600
ctggacctcg aatttgacaa cgagaacctg attaacatct tgactgaccc aaaactcaat 660
aaaaaggcca agaagaagga gctgaagtcg gtaatcgagg acaccaagtt cctcaagacc 720
gtttccgcga taatgatcgg ctctagccag aaactcgtcg acttgttcga gaaccccgag 780
gatttcgacg actctcgat aaagtccgtt gacttctcaa ctacctttt cgacgacaag 840
tactctgact atgaactcgc tctgggtgac aagatcgctc tggtaacatc ccttaaggaa 900
atttacgata gctccatcct cgagaacctg ctcaaaggag cagacaagtc taaggacggt 960
aacaatatata tcagttaatgc attcgtgaag aagtacaata aacacggaca agatctgaaa 1020

```

-continued

gagttcaaac	gtctggtacg	acaatatcac	aagagtgcgt	atattgatat	tttcagatcc	1080
gagaagggtga	atgacaatta	cgtcagctac	actaaaagct	caattagcaa	caataaacgc	1140
gtcaaaagcaa	acaagttcac	tgatcaagag	gcctcttaca	aattcgccaa	gaaacatctg	1200
gagacaatca	agtataagat	caacaaggta	aacggctcca	aggcagatct	ggagctgatt	1260
gacgggagtc	tgccgggacat	ggagttcaag	aactttatgc	ccaaaattaa	aagctcggat	1320
aacggcggtga	ttccgtggca	gctgaaactg	atggaattga	acaaataact	cgagaatcag	1380
tcaaagcatc	acgagttcct	caatgtcagc	gacgagtagc	gctccgtgtg	tgataaaatc	1440
gcatactatca	tggaagtccg	tatcccctac	tacgtgggac	ccctgaaccc	caatagcaag	1500
tacgccttga	tcaagaagca	gaaagatagt	gagattactc	cctggaactt	caaggacgtc	1560
gtggacctcg	actccagcag	agaggagtgc	attgactcac	tgatcggacg	ctgtacttac	1620
cttaaggacg	agaaggctct	tcccaaagct	tctttgctgt	ataacgaata	catgggtgctg	1680
aacgagctga	ataacctgaa	gttgaaacgac	cttcccatca	ccgaggagat	gaagaagaag	1740
atatttgacc	agttgttcaa	aacaagaaaag	aaggtcacc	ttaaagcggg	ggcaaacctg	1800
ctgaagaagg	agttcaacat	caacggcag	attctgctct	ctgggacgca	cggtgacttc	1860
aagcagggct	tgaactcata	caatgacttc	aaagctatcg	tgggcgataa	agtcgatctc	1920
gatgattacc	gggacaagat	tgaggagatc	attaaactga	tagttcttta	cggtgacgat	1980
aagagttacc	ttcagaagaa	gattaaagct	gggtatggaa	aatacttcac	cgacagttag	2040
attaagaaaa	tggcggggct	gaactacaag	gattggggaa	ggctctcaaa	gaagctgctg	2100
acgggactcg	aggggtgcaa	caagatcact	ggagagcggg	gctccattat	tcacttcatg	2160
agggaaata	accttaactc	gatggagctt	atgtcagctt	catttacgtt	caccgaagag	2220
atacagaaac	ttaacccctg	ggaatgaccg	aagctgtcat	acgaaatggt	ggacgaactg	2280
tacctttctc	ccagtgtgaa	acggatgctc	tggcagtcct	tgcgcactgt	cgacgagata	2340
aagaacatca	tgggaaacga	cagtaagaag	attttcatcg	agatggctaa	gggtaaggaa	2400
gaggtgaaaag	cccgaagga	gtcaaggaag	aaccaactgc	tgaagttcta	taaagacgga	2460
aagaaggcat	tcatcagcga	gattggcgag	gagaggtact	cttacttgc	ttctgagata	2520
gagggtaggg	aagagaataa	gtttcgatgg	gataacaact	acctttatta	tactcaactg	2580
ggtcgctgca	tgtactcttt	ggaacctatc	gacatatctg	agctgtcttc	aaagaatat	2640
tacgatcagg	atcatatcta	ccccaaaagc	aagatttacg	acgacagtat	cgagaatagg	2700
gtgctgggtga	agaaggacct	taactccaag	aagggttaaca	gctatcctat	cccagacgaa	2760
atcctgaaca	agaactgtta	cgcctactgg	aagatcctgt	acgataaagg	tcttactcggg	2820
cagaagaagt	acactcggct	gaaccggaga	actggcttca	cggacgacga	gctcgttcag	2880
ttcatctcaa	gacagatcgt	ggaaaactaga	caagcaacaa	aggagactgc	taacctgctc	2940
aagacaatat	gtgaagaact	cgagatcatg	tattccaaag	ccgagaacgc	aagtcgggtt	3000
aggaagagt	tcatcaactg	gaagtgtagg	gcgggtgaacg	atcttcatca	tatgcacgat	3060
gacctacatca	acatcatagt	gggggaacgtg	tataaacacca	agttcacgaa	ggacctatg	3120
aatttcgtaa	agaagcagga	aaaggcgctg	agctacaatc	tcgagaatat	gttcaagtag	3180
gatgtgaaac	gtggcggata	caccgcttgg	atcgccgatg	acgagaaggg	caccgtgaag	3240
aacgcgagta	ttaaactgtat	ccggaaggag	ctggaaggca	caaattatag	gttcacaaga	3300
atgaactaca	ttgagtcctg	agcgcttttc	aacgccactc	tccagcggaa	gaataagggc	3360
tccagacccc	tgaaggacaa	agggccgaaa	tcttccatcg	agaagtagcg	cggctacaca	3420
aacatcaata	agaactgttt	cgctgttctt	gacatcaagt	ctaagaacaa	gattgagagg	3480
aagctgatgc	ccgtcgagcg	tgagatctat	gccaaacaga	agaacgacaa	gaagctgtcc	3540
gacgagattt	tctcaaatga	cctcaaggac	cgatttgcca	tcgaggacta	caggggtgtc	3600
taccagtggt	tgaaaatgct	cacactgctc	aagatcgacg	gcagctacta	cttcacacaa	3660
ggcggtctct	ataagaccct	ggagttgcga	tctgctctgc	agctgattct	ccctaagaag	3720
aacgagtggt	gcattcaaaa	gattcgacaa	tcttccgaaa	acgactatct	gacgatcgag	3780
cgtatccagg	acctgaccga	ggagctgggt	tataaacactt	tcgacatcat	cgtcaacaag	3840
ttcaagacca	gtgtcttcaa	gaagcttttc	cttaacttgt	ttcaggacga	caagatttga	3900
aaacattgact	tcaagtttaa	gtccatggac	ttcaaggaga	aatgcaagac	acttctcatg	3960
ctgggtcaagg	cgattcgggc	atccggcgtg	aggcaggatc	tcaagtcctat	cgacctcaag	4020
tctgattacg	gacgggtcag	ttcaaaagacc	aacaacatcg	gcaattacca	ggagtccaag	4080
attattaatc	agtcctatc	tgagactgttc	gagaatgagg	tcgatctctc	gaagctggga	4140
tcctaccatc	acgatgttcc	agattaccgc	gccgctccaa	aaaagaaaaa	aaaagttgctg	4200
gtagccatc	atcaccatca	ccatcatcat	taa			4233

SEQ ID NO: 19 moltype = DNA length = 4233
 FEATURE Location/Qualifiers
 misc_feature 1..4233
 note = Variant 2902 with tags DNA sequence
 source 1..4233
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 19

atgcctaaga	agaagagaaa	ggtgggtacc	accaagggtga	aggactacta	cataggcttg	60
gacatcgcca	cctctagcgt	cgggtgggccc	gtcaccgatg	aagcctataa	cgtgcttaag	120
tttaatatga	agaaaaatgtg	gggcgtgctg	ctgttcgacg	acgctaagac	ggcagaggag	180
cgtagggggc	agcgaggagc	aagacgacgt	ctggatcgga	agaaggagag	actcagcctg	240
ctgcaggact	tcttcgccga	agaggtagca	aaggtcgacc	ccaacttctt	cctcaggctg	300
gacaattccg	atctgtacat	ggaagataag	gaccagaaac	tgaaaagcaa	atatacactg	360
ttcaacgaca	aggacttcaa	ggataagaat	tttcataaga	agtacccacc	aatacatcac	420
ctgtctgagg	actctgcaga	ggacgacagt	aagaaggaca	tccggctcgt	ctacctggcc	480
tgtcaactatt	tgtccaagaa	caggggtcat	ttcatcttcg	agggccagaa	gttcgacact	540
aatcaagct	tcgagaacag	tttgaacgag	ctcaaagttc	atattgaacga	cgagtatgga	600
ctggacctcg	aatttgacaa	cgagaacctg	attaacatct	tgactgaccc	aaaactcaat	660
aaaacggcca	agaagaaggga	gctgaagtc	gtaactcggc	acaccaagtt	cctcaagacc	720
gtttccgcga	taatgatcgg	ctctagccag	aaactcgtcg	acttggtcga	gaaccccgag	780

-continued

```

gatttcgacg actctgcgat aaagtccggt gacttctcaa ctacctcttt cgacgacaag 840
tactctgact atgaactcgc tctgggtgac aagatcgctc tggtaacat ccttaaggaa 900
atttacgata gctccatcct cgagaacctg ctcaaagagg cagacaagtc taaggacggt 960
aacaataata tcagtaatgc attcgtgaag aagtacaata aacacggaca agatctgaaa 1020
gagttcaaac gtctggtagc acaatatcac aagagtgcgt attttgatat ttccagatcc 1080
gagaaggtga atgacaatta cgtcagctac actaaaagct caattagcaa caataaacgc 1140
gtcaaagcaa acaagttcac tgatcaagag gcctcttaca aattcgccaa gaaacatctg 1200
gagacaatca agtataagat caacaaggta aacggctcca aggcagatct ggagctgatt 1260
gacgggatgc tgcgggacat ggagtccaag aactttatgc ccaaaattaa gtccagtgc 1320
aacggggtga ttccatacca gctcaagctg atggaattga acaaaatact cgagaatcag 1380
tcaaagcatc acgagtctct caatgtcagc gacgagtacg gctccgtgtg tgataaaatc 1440
gcatctatca tggagtcccg tatccctac tacgtgggac ccctgaaccc caatagcaag 1500
tacgcctgga tcaagaagca gaaagatagt gagattactc cctggaaact caaggacgtc 1560
gtggacctcg actccagcag agaggagttc attgactcac tgatcggagc ctgtacttac 1620
cttaaggacg agaaggtcct tcccaaagct tctttgctgt ataacgaata catggtgctg 1680
aacgagctga ataactgaa gttgaacgac ctccccatca cggaggagat gaagaagaag 1740
atatattgacc atgtgttcaa aacaagaaaag aaggtcacc ttaagcgggt ggcaaacctg 1800
ctgaagaagg agttcaacat caacggcgag attctgctct ctccgaccca cggtgacttc 1860
aagcagggct tgaactcata caatgacttc aaagctatcg tgggcgataa agtcgattcc 1920
gatgattacc gggacaagat tgaggagatc attaaactga tagttcttta cgttgacgat 1980
aagagttacc ttcagaagaa gattaaagct gggataggaa aatacttcac cgacagttag 2040
attaagaaaa tggcggggct gaactacaag gattgggaa ggctctcaa gaaggtctc 2100
acgggactcc aggggtgcaaa caagatcact ggagagcggt gctccattat tcacttcag 2160
agggaaatata acctaatct gatggagctt atgtcagctt cattacggtt caccgaagag 2220
atacagaaac ttaaccccggt ggatgaccgc aagctgtcat acgaaatggt ggacgaactg 2280
tacctttctc ccagtgtgtc acggatgctc tggcagctcc tgcgcactcg cgacgagata 2340
aagaacatca tgggaaccca cagtaagaag attttcatcg agatggctcg gggtaaaggaa 2400
gaggtgaaag cccgcaagga gtcaaggaa aaccaactgc tgaagttcta taaagacgga 2460
aagaaggcat tctcagcgca gatggcgag gagaggtact cttaactgct ttctgagata 2520
gaggggtgag aagagaataa gtttcgatgg gataacctgt acctttatta tactcaactg 2580
ggtcgtgca tgtactcttt ggaactatc gacatctctg agctgtcttc aaagaatatt 2640
tacgatcagg atcatatcta ccccaaaagc aagatttacg acgacagtat cgagaatagg 2700
gtgctggtga agaaggacct taactccaag aagggtaaaca gctatcctat cccagacgaa 2760
atcctgaaca agaactgttg gcctactgg aagatcctgt acgataaagg tcttatcggt 2820
cagaagaagt acactcggtt gacccggaga actggcttca cggacgacga gctcgttcag 2880
ttcatctcaa gacagatcgt ggaactaga caagcaacaa aggagactgc taacctgctc 2940
aagacaatat gtaagaactc cgagatcgtg agcgtcttc cagagaacgc aagtcggtt 3000
aggcaagagt tcgacatcgt gaagtgtagg gcggtgaacg atcttcatca tatgcacgat 3060
gctcatatca acatcatagt ggggaacgtg tataacacca agttcacgaa ggacctatg 3120
aatttcgtaa agaagcagga aaaggcgcg agctacaatc tcgagaatat gttcaagtac 3180
gatgtgaaac gtggcgagata caccgcttgg atcgccgatg acgagaaggg caccgtgaag 3240
aacgcgagta ttaaacgtat ccggaaggag ctggaaggca caaattatag gttcacaaaga 3300
atgaactaca ttgagtctgg agcgtttttc aacgccactc tccagcgga gaataagggc 3360
tccagacccc tgaaggacaa agggccgaaa tcttccatcg agaagtagcg cggctacaca 3420
aacatcaata aagcctgttt cgctgttctt gacatcaagt ctaagaacaa gattgagagg 3480
aagctgatgc ccgtcgagcg tgagatctat gccaaacaga agaacgacaa gaagctgtcc 3540
gacgagattt tctcaaaagta cctcaaggac cgatttgcca tcgaggacta cagggttgtc 3600
taccagtggt tgaaaatgcg cacactgctc aagatcgacg gcagctacta ctccatcaca 3660
ggcgggtctg ataagaccct ggaagttgca tctgctctgc agctgattct ccataagaag 3720
aacgagtggt cgatcaaaac gatcgacaag tcttccgaaa acgactatct gacgatcgag 3780
cgtatccagg acctgaccca ggagctggtg tataacactt tcgacatcat cgtcaacaag 3840
ttcaagacca gtgtcttcaa gaagtctttc cttaacttgt ttcaggacga caagattgag 3900
aacattgact tcaagtttaa gtccatggac ttcaaggaga aatgcaagac acttctcatg 3960
ctggtcaagg cgattcgggc atccggcgtg aggcaggatc tcaagtccat cgacctcaag 4020
tctgattacg gacggctcag ttcaaaagacc aacaactacg gcaattacca ggagttcaag 4080
attattaatc agtccatcac tggactgttc gagaatgagg tcgatctcct gaagctggga 4140
tctacccat acgatgttcc agattacgcg gccgtccaa aaaagaaaaa aaaagttgcg 4200
gctagccatc atcaccatca ccatcatcat taa 4233

```

```

SEQ ID NO: 20      moltype = DNA length = 4233
FEATURE
misc_feature       1..4233
                    note = Variant 3954 with tags DNA sequence
source             1..4233
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 20
atgcctaaga agaagagaaa ggtgggtacc accaagggtga aggactacta cataggttgc 60
gacatcggca cctctagcgt cgggtggggc gtcaccgatg aagcctataa cgtgcttaag 120
ttaataagca agaaaatgtg gggcggtgagg ctgttcgacg acgctaagac ggacagggag 180
cgtaggggccc agcgaggagc aagacgacgt ctggatcgga agaaggagag actcagcctg 240
ctgcaggact tcttcgcca agaggtagca aaggtagacc ccaacttctt cctcaggctg 300
gacaattccg atctgtacat ggaagataag gaccagaaac tgaaaagcaa atatacactg 360
ttcaacgaca aggacttcaa ggataagaat ttctcataaga agtaccaccac aatacatcac 420
ctgctgatgg atctgatcga ggacgacagt aagaaggaca tccggctcgt ctacctggcc 480
tgtcactatt tgcctaagaa caggggtcat ttcattctcg agggccagaa gttcgacact 540

```

-continued

```

aatcaagct tgcgagaacag tttgaacgag ctcaaagttc atttgaacga cgagtatgga 600
ctggacctcg aatttgacaa cgagaacctg attaacatct tgactgacct aaaactcaat 660
aaaacggcca agaagaagga gctgaagtcg gtaatcggcg acaccaagtt cctcaaaagg 720
gtttccgcga taatgatcgg ctctagccag aaactcgtcg acttggtcga gaaccccgag 780
gatttcgacg actctgcatg aaagtcggtt gacttctcaa ctacctcttt cgacgacaag 840
tactctgact atgaactcgc tctgggtgac aagatcgctc tggccaacat ccttaaggaa 900
atttacgata gctcaatctc cgagaacctg ctcaaagagg cagacaagtc taaggacggg 960
aacaatatata tcagtaatgc attcgtgaag aagtacaata aacacggaca agatctgaaa 1020
gagttcaaac gtctggtagc acaatatcac aagagtgcgt attttgatat tttcagatcc 1080
gagaaggtag atgacaatta cgtcagctac actaaaagct caattagcaa caataaacgc 1140
gtcaaagcaa acaagttcac tgatcaagag gccttctaca aattcgccaa gaaacatctg 1200
gagacaatca agtataagat caacaaggta aacggctcca aggcagatct ggagctgatt 1260
gacgggtagc tgcgggacat ggagttcaag aactttatgc ccaaaattaa aagctcggat 1320
aacggcgtga ttccgtatca gctgaaactg atggaattga acaaaatact cgagaatcag 1380
tcaaagcatc acgagttcct caatgtcagc gacgagtagc gctcgtgtg tgataaaatc 1440
gcatctatca tgaatttacc cattccgtat tacgtggggc cgctgaacc gaacagcaaa 1500
tatggctgga tcaagaagca gaaagatagt gagattactc cctggaactc caaggacgtc 1560
gtggacctcg actccagcag agaggagttc attgactcac tgatcggagc ctgtacttac 1620
cttaaggacg agaaggtcct tcccgaagct tctttgctgt ataacgaata catggtgctg 1680
aacgagctga ataacctgaa gttgaacgac cttcccatca ccgaggagat gaagaagaag 1740
atatttgacc agttgttcaa aacaagaaag aaggtcacc ttaagcggg ggcaaacctg 1800
ctgaagaagg agttcaacat caacggcgcg attctgctct ctgggaccga cggtagcttc 1860
aagcagggtc tgaactcata caatgacttc aaagctatcg tgggcgataa agtcgattcc 1920
gatgattacc gggacaagat tgaggagatc attaaactga tagttcttta cggtagacgat 1980
aagagttacc ttcagaagaa gattaaagct gggtagtgaa aatacttcac cgacagttag 2040
attaagaaaa tggcggggct gaactacaag gattggggaa ggctctcaga aaagctgctg 2100
acgggactcg aggggtgaaa caagatcact ggagagcggg gctccattat tcaactcoatg 2160
agggaaatata accttaattg tatggagctt atgtcagctt cattacgtt caccgaagag 2220
atacagaaac ttaacctcgt ggaatgaccgc aagctgtcat acgaaatggg ggacgaactg 2280
tacctttctc ccagttgtga accgatgctc tggcagtcct tgcgcactcg cgacgagata 2340
aagaacatca tgggaacctga cagtaagaag attttcatcg agatggctcg gggtaaggaa 2400
gaggtgaaag cccgcaagga gtcaaggaaag aaccaactgc tgaagttcta taaagacgga 2460
aagaaggcat tcatcagcga gattggcgag gagaggtagc cttaactgct ttctgagata 2520
gagggtagag aagagaataa gtttcgatgg gataacctgt acctttatta tactcaactg 2580
ggtcgctgca tgactctttt ggaacctatc gacatatctg agctgtcttc aaagaatatt 2640
tacgatcagg atcatatcta ccccaaaagc aagatttacg acgacagtat cgagaatagg 2700
gtgctggtga agaaggacct taactccaag aagggttaaca gctatcctat cccagacgaa 2760
atcctgaaca agaactgtta cgctactcgg aagatcctgt acgataaagg tcttatcggg 2820
cagaagaagt acactcggct gacccggaga actggtctca cggacgacga gctcgttcag 2880
ttcatctcaa gacagatcgt ggaactaga caagcaacag tcgagactgc taacctgctc 2940
aagacaatat gtaagaactc cgagatcgtg tattccaaag ccgagaacgc aagtcgggtt 3000
aggcaagagt tcgacatcgt gaagtgtagg gcgggtgaac atcttcatca tatgcacgat 3060
gcctacatca acatcatagt ggggaacctg tataaacacca agttcacgaa ggacctatg 3120
aatctcgtaa aggtccagga aaaggcgcgg agctacaatc tcgagaatat gttcaagtag 3180
gatgtgaaac gtggcggata caccgcttgg atcgccgatg acgagaaggg caccgtgaag 3240
aacgcgagta ttaaacgtat ccggaaggag ctggaaggca caaattatag gttcacaaga 3300
atgaactaca ttgagtctgg agcgtctttc aacgccactc tccagcggaa gaataagggc 3360
tccagacccc tgaaggacaa agggccgaaa tcttccatcg agaagtacgg cggctacaca 3420
aacatcaata aagcctgttt cgctgttctt gacatcaagt ctaagaacaa gattgagagg 3480
aagctgatgc ccgtcgagcg tgagatctat gccaaacaga agaacgacaa gaagctgtcc 3540
gacgagattt tctcaaagta cctcaaggac cgatttgga tcgaggacta cagggttgtc 3600
taccagttcg tgaaaatgcg cacactgctc aagatcgacg gcagctacta cttcatcaca 3660
ggcgggttctg ataagacctt ggagttgcca tctgctctgc agctgattct ccctaagaag 3720
aacgagtggt cgatcaaaaca gatcgacaag tcttccgaaa acgactatct gacgatcgag 3780
cgtatccagg acctgaaccga ggaagctggtg tataaacact tcgacatcat cgtcaacaag 3840
ttcaagacca gtgtcttcaa gaagtctttc cttaacttgt ttcaggacga caagattgag 3900
aacattgact tcaagtttaa gtccatggac ttcaggaga aatgcaagac acttctcatg 3960
ctggtaaggc cgattcggcg atccggcgtg aggcaggatc tcaagtccat cgacctcaag 4020
tctgattacg gacggctcag ttcaaagacc aacaacatcg gcaattacca ggagttcaag 4080
attattaatc agtccatcac tggactgttc gagaatgagg tcgatctcct gaagctggga 4140
tcctacccat acgatgttcc agattacgcg gccgctccaa aaaagaaaag aaaagttgcg 4200
gctagccatc atcaccatca ccatcatcat taa 4233

```

```

SEQ ID NO: 21      moltype = AA length = 7
FEATURE           Location/Qualifiers
REGION           1..7
                  note = Nt-NLS
source           1..7
                  mol_type = protein
                  organism = synthetic construct

```

```

SEQUENCE: 21
PKKKRKV

```

7

```

SEQ ID NO: 22      moltype = AA length = 9
FEATURE           Location/Qualifiers
REGION           1..9

```

-continued

source	note = HA tag 1..9 mol_type = protein organism = synthetic construct	
SEQUENCE: 22 YPYDVPDYA		9
SEQ ID NO: 23 FEATURE REGION	moltype = AA length = 7 Location/Qualifiers 1..7 note = Ct-NLS	
source	1..7 mol_type = protein organism = synthetic construct	
SEQUENCE: 23 PKKKRKV		7
SEQ ID NO: 24 FEATURE REGION	moltype = AA length = 8 Location/Qualifiers 1..8 note = X8 His-tag	
source	1..8 mol_type = protein organism = synthetic construct	
SEQUENCE: 24 HHHHHHHH		8
SEQ ID NO: 25 FEATURE misc_feature	moltype = DNA length = 21 Location/Qualifiers 1..21 note = Nt-NLS DNA sequence	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 25 cctaagaaga agagaaaggt g		21
SEQ ID NO: 26 FEATURE misc_feature	moltype = DNA length = 27 Location/Qualifiers 1..27 note = HA tag DNA sequence	
source	1..27 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 26 taccatacag atgttcaga ttacgcg		27
SEQ ID NO: 27 FEATURE misc_feature	moltype = DNA length = 21 Location/Qualifiers 1..21 note = Ct-NLS DNA sequence	
source	1..21 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 27 ccaaaaaaga aaagaaaagt t		21
SEQ ID NO: 28 FEATURE misc_feature	moltype = DNA length = 24 Location/Qualifiers 1..24 note = X8 His-tag DNA sequence	
source	1..24 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 28 catcatcacc atcaccatca tcat		24
SEQ ID NO: 29 FEATURE misc_feature	moltype = RNA length = 20 Location/Qualifiers 1..20 note = g62 REF 20 nt	
source	1..20 mol_type = other RNA organism = synthetic construct	
SEQUENCE: 29 gtcaagcccc agaggccaca		20

-continued

```

SEQ ID NO: 30      moltype = RNA  length = 20
FEATURE           Location/Qualifiers
misc_feature       1..20
                   note = g62 ALT 20 nt
source            1..20
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 30
gtcaagcccc agaggacaca                               20

SEQ ID NO: 31      moltype = RNA  length = 22
FEATURE           Location/Qualifiers
misc_feature       1..22
                   note = sgRNA-564DS-ref 22 nt
source            1..22
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 31
gtgtcaagcc ccagaggcca ca                             22

SEQ ID NO: 32      moltype = RNA  length = 22
FEATURE           Location/Qualifiers
misc_feature       1..22
                   note = sgRNA-564DS-alt 22 nt
source            1..22
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 32
gtgtcaagcc ccagaggaca ca                             22

SEQ ID NO: 33      moltype = RNA  length = 20
FEATURE           Location/Qualifiers
misc_feature       1..20
                   note = g35 20 nt
source            1..20
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 33
agtcgggct gggagcgggt                                20

SEQ ID NO: 34      moltype = RNA  length = 22
FEATURE           Location/Qualifiers
misc_feature       1..22
                   note = sgRNA-Constant 22 nt
source            1..22
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 34
gcagtcggg ctgggagcgg gt                             22

SEQ ID NO: 35      moltype = RNA  length = 102
FEATURE           Location/Qualifiers
misc_feature       1..102
                   note = Synthetic sequence
source            1..102
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 35
gtgtcaagcc ccagaggcca cagtttgaga gttatgaaaa tgacgagttc aaataaaaat 60
ttattcaaac cgcctattta taggcgcag atgttctgct tt                    102

SEQ ID NO: 36      moltype = RNA  length = 102
FEATURE           Location/Qualifiers
misc_feature       1..102
                   note = Synthetic sequence
source            1..102
                   mol_type = other RNA
                   organism = synthetic construct

SEQUENCE: 36
gtgtcaagcc ccagaggaca cagtttgaga gttatgaaaa tgacgagttc aaataaaaat 60
ttattcaaac cgcctattta taggcgcag atgttctgct tt                    102

SEQ ID NO: 37      moltype = RNA  length = 102
FEATURE           Location/Qualifiers
misc_feature       1..102

```

-continued

```

source                note = Synthetic sequence
                      1..102
                      mol_type = other RNA
                      organism = synthetic construct

SEQUENCE: 37
gcagtcgcggg ctgggagcgg gtgtttgaga gttatgaaaa tgacgagttc aaataaaaat 60
ttattcaaac cgcctattta taggcgcgag atgttctgct tt 102

SEQ ID NO: 38         moltype = DNA length = 56
FEATURE              Location/Qualifiers
variation            1..56
                      note = Synthetic sequence
variation            34..36
                      note = n is a, c, g, or t
source               1..56
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 38
tcgtcggcag cgtcagatgt gtataagaga cagnnntacg accccgtaaa cttgct 56

SEQ ID NO: 39         moltype = DNA length = 57
FEATURE              Location/Qualifiers
variation            1..57
                      note = Synthetic sequence
variation            35..37
                      note = n is a, c, g, or t
source               1..57
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 39
gtctcgtggg ctcggagatg tgtataagag acagnnncgg agcgttggat gatagag 57

SEQ ID NO: 40         moltype = DNA length = 56
FEATURE              Location/Qualifiers
variation            1..56
                      note = Synthetic sequence
variation            34..36
                      note = n is a, c, g, or t
source               1..56
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 40
tcgtcggcag cgtcagatgt gtataagaga cagnnntcgc agtccagctt ccccac 56

SEQ ID NO: 41         moltype = DNA length = 57
FEATURE              Location/Qualifiers
variation            1..57
                      note = Synthetic sequence
variation            35..37
                      note = n is a, c, g, or t
source               1..57
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 41
gtctcgtggg ctcggagatg tgtataagag acagnnnaca gccaaggagc atcaaac 57

SEQ ID NO: 42         moltype = DNA length = 20
FEATURE              Location/Qualifiers
misc_feature         1..20
                      note = Synthetic sequence
source               1..20
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 42
acgtctgcac tctcgtgagg 20

SEQ ID NO: 43         moltype = DNA length = 22
FEATURE              Location/Qualifiers
misc_feature         1..22
                      note = Synthetic sequence
source               1..22
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 43
ttcaggctcc acccagtttg tc 22

```

-continued

SEQ ID NO: 44	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = Synthetic sequence	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 44		
gcacagggct ataagaggag c		21
SEQ ID NO: 45	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = Synthetic sequence	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 45		
gcgggagggtt ggactcaaaa		20
SEQ ID NO: 46	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
misc_feature	1..19	
	note = Synthetic sequence	
source	1..19	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 46		
gggagaggaa gtggagggt		19
SEQ ID NO: 47	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
misc_feature	1..18	
	note = Synthetic sequence	
source	1..18	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 47		
gagggtcatg gtggggct		18
SEQ ID NO: 48	moltype = DNA length = 56	
FEATURE	Location/Qualifiers	
variation	1..56	
	note = Synthetic sequence	
variation	34..36	
	note = n is a, c, g, or t	
source	1..56	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 48		
tcgtcggcag cgtcagatgt gtataagaga cagnnnngtg ggtcctcagt gactct		56
SEQ ID NO: 49	moltype = DNA length = 57	
FEATURE	Location/Qualifiers	
variation	1..57	
	note = Synthetic sequence	
variation	35..37	
	note = n is a, c, g, or t	
source	1..57	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 49		
gtctcgtggg ctcggagatg tgtataagag acagnnnggg aattccagcc tgaccaa		57
SEQ ID NO: 50	moltype = DNA length = 56	
FEATURE	Location/Qualifiers	
variation	1..56	
	note = Synthetic sequence	
variation	34..36	
	note = n is a, c, g, or t	
source	1..56	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 50		
tcgtcggcag cgtcagatgt gtataagaga cagnnnngagt gaggaccaag cctgag		56

-continued

```

SEQ ID NO: 51      moltype = DNA length = 57
FEATURE           Location/Qualifiers
variation         1..57
                  note = Synthetic sequence
variation         35..37
                  note = n is a, c, g, or t
source           1..57
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 51
gtctcgtggg ctccgagatg tgtataagag acagnnnagg gccattgtct ccctaac      57

SEQ ID NO: 52      moltype = DNA length = 56
FEATURE           Location/Qualifiers
variation         1..56
                  note = Synthetic sequence
variation         34..36
                  note = n is a, c, g, or t
source           1..56
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 52
tcgtcggcag cgtcagatgt gtataagaga cagnnntcct gctaccctcc cttctt      56

SEQ ID NO: 53      moltype = DNA length = 57
FEATURE           Location/Qualifiers
variation         1..57
                  note = Synthetic sequence
variation         35..37
                  note = n is a, c, g, or t
source           1..57
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 53
gtctcgtggg ctccgagatg tgtataagag acagnnnttt taggaggggc ccactga      57

SEQ ID NO: 54      moltype = DNA length = 20
FEATURE           Location/Qualifiers
misc_feature      1..20
                  note = Synthetic sequence
source           1..20
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 54
gagggtcatc atcactgccc      20

SEQ ID NO: 55      moltype = DNA length = 27
FEATURE           Location/Qualifiers
misc_feature      1..27
                  note = Synthetic sequence
source           1..27
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 55
gccagactca caccagagtc gacaagt      27

SEQ ID NO: 56      moltype = DNA length = 20
FEATURE           Location/Qualifiers
misc_feature      1..20
                  note = Synthetic sequence
source           1..20
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 56
taatacgact cactataggg      20

SEQ ID NO: 57      moltype = DNA length = 18
FEATURE           Location/Qualifiers
misc_feature      1..18
                  note = Synthetic sequence
source           1..18
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 57
atttaggtga cactatag      18

```

-continued

SEQ ID NO: 58 moltype = DNA length = 56
 FEATURE Location/Qualifiers
 variation 1..56
 note = Synthetic sequence
 variation 34..36
 note = n is a, c, g, or t
 source 1..56
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 58
 tcgtcggcgag cgctcagatgt gtataagaga cagnnntcgc agtccagctt ccccac 56

SEQ ID NO: 59 moltype = DNA length = 57
 FEATURE Location/Qualifiers
 variation 1..57
 note = Synthetic sequence
 variation 35..37
 note = n is a, c, g, or t
 source 1..57
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 59
 gtctcgtggg ctcggagatg tgtataagag acagnnnaca gccaaaggagc atcaaac 57

SEQ ID NO: 60 moltype = DNA length = 56
 FEATURE Location/Qualifiers
 variation 1..56
 note = Synthetic sequence
 variation 34..36
 note = n is a, c, g, or t
 source 1..56
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 60
 tcgtcggcgag cgctcagatgt gtataagaga cagnnngagg gtcacatca ctgccc 56

SEQ ID NO: 61 moltype = DNA length = 57
 FEATURE Location/Qualifiers
 variation 1..57
 note = Synthetic sequence
 variation 35..37
 note = n is a, c, g, or t
 source 1..57
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 61
 gtctcgtggg ctcggagatg tgtataagag acagnnnagt cggggctggg agcgggt 57

SEQ ID NO: 62 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = Wild-type OMNI-50
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 62
 TKVKDYIIGL DIGTSSVGWA VTDEAYNVLK FNSKKMWGVR LFDDAKTAE RRGQRGARRR 60
 LDRKKERLSL LQDFFAEVA KVDPNFFLR L DNSDLYMEDK DQKLKSKYTL FNDKDFKDKN 120
 FHKKYPTIIH LLMDLIEDDS KKDRLVYLA CHYLLKNRGH FIFEGQKFD T KSSFENSLNE 180
 LKVHLNDEYG LDLEFDNENL INILTDPKLN KTAKKKELKS VIGDTKFLKA VSAIMIGSSQ 240
 KLVDLFENPE DFDDSAIKSV DFSTTSFDDK YSDYELALGD KIALVNILKE IYDSSILENL 300
 LKEADKSKDG NKYISNAFVK KYNKHGQDLK EFKRLVRQYH KSAYFDIFRS EKVNDNYVSY 360
 TKSSISNNKR VKANKFTDQE AFYKFAKKHL ETIKYKINKV NGSKADLELI DGMLRDMFEK 420
 NFMPIKSSD NGVIPYQLKL MELNKKILEN SKHHEFLNVS DEYGSVCDKI ASIMEFRIPY 480
 YVGPLNPNSK YAWIKKQKDS EITPWNFKDV VLDLSSREEF IDSLIGRCTY LKDEKVLPA 540
 SLLYNEYMVL NELNNKLND LPITEEMKKK IFDQLFKTRK KVTLKAVANL LKKEFNINGE 600
 ILLSGTDGDF KQGLNSYNDF KAIVGDKVDS DDYRDKIEEI IKLIVLYGDD KSYLQKKIKA 660
 GYGKYFTDSE IKKMAGLNYK DWGRLSKKLL TGLEGANKIT GERGSIIHFM REYNLNLME 720
 MSASFTFTEE IQKLPVDDR KLSYEMVDEL YLSPSVKRM L WQSLRIVDEI KNIMGTD SKK 780
 IFIEMARGKE EVKARKESRK NQLLKPYKDG KKAFISEIGE ERYSYLLSEI EGEEENKFRW 840
 DNLYLYYTQL GRCMYSLEPI DISELSSKNI YDQDHIYPKS KIYDDSIENR VLVKKDLNSK 900
 KGNSYPIPDE ILNKNCYAYW KILYDKGLIG QKKYTRLTRR TGFTDDELVQ FISRQIVETR 960
 QATKETANLL KTICKNSEIV YSKAENASRF RQEFDIVKCR AVNDLHHMHD AYINIIVGNV 1020
 YNTKFTKDPM NFVKKQEKAR SYNLENMFKY DVKRGGYTAW IADDEKGTVK NASIKRIRKE 1080
 LBGTNYRFRF MNYIESGALF NATLQRKNKG SRPLKDKGPK SSIEKYGGYT NINKACFAVL 1140
 DIKSKNKIER KLMPVEREYI AKQKNDKKLS DEIFSKYLLD RFGIEDYRVV YPVVKMRTL 1200

-continued

KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLMM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEFK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 63 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V3795
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 63

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VDLDSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLNHEYMYL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNLYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHF	REYNLNLME	720
MSASPTFTTE	IQKLNPDVDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFR	MNYIESGALF	NATLQKKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLMM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEFK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 64 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6918
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 64

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFRIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VDLDSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLYNEYMYL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNLYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHF	REYNLNLME	720
MSASPTFTTE	IQKLNPDVDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLKIFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFR	MNYIESGALF	NATLQKKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLMM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEFK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 65 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6919
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 65

-continued

TKVKDYYIIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMED	DQKLKSKYT	FNDKDFDKN	120
FHKKYPTIHH	LLMDLIEDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMEPK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFRIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHNEVMV	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHF	REYNLNLME	720
MSASFTFTEE	IQKLNPFVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NQLLKFKYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPE	ILNKNYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LBGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREYI	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSITGLF	ENEVDLLKL		1369

SEQ ID NO: 66 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6920
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 66

TKVKDYYIIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMED	DQKLKSKYT	FNDKDFDKN	120
FHKKYPTIHH	LLMDLIEDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMEPK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFRIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHNEVMV	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHF	REYNLNLME	720
MSASFTFTEE	IQKLNPFVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NQLLKFKYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPE	ILNKNYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LBGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREYI	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSITGLF	ENEVDLLKL		1369

SEQ ID NO: 67 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6864
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 67

TKVKDYYIIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMED	DQKLKSKYT	FNDKDFDKN	120
FHKKYPTIHH	LLMDLIEDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMEPK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHNEVMV	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHF	REYNLNLME	720

-continued

MSASFTFTEE	IQKLNVPVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLLKFKYKD	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEPK	IINQSITEGLF	ENEVDLLKL		1369

SEQ ID NO: 68 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6865
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 68

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIIH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EPKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILEN	SKHHEFLNVS	DEYGSVCDKI	ASIMEPIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLNHEYVMV	NELNNLKLND	LPITEEMKKK	IPDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLME	720
MSASFTFTEE	IQKLNVPVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLLKFKYKD	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEPK	IINQSITEGLF	ENEVDLLKL		1369

SEQ ID NO: 69 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6866
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 69

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIIH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EPKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILEN	SKHHEFLNVS	DEYGSVCDKI	ASIMEPIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLNHEYVMV	NELNNLKLND	LPITEEMKKK	IPDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLME	720
MSASFTFTEE	IQKLNVPVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLLKFKYKD	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDDP	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFDQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSST	NNIGNYQEPK	IINQSITEGLF	ENEVDLLKL		1369

-continued

SEQ ID NO: 70 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V6867
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 70

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARRR	60
LDRKKERLSL	LQDFFAEVVA	KVDPNFRLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDSSAISKV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EK VNDNYVSY	360
TKSSISNNKR	VGANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMEPK	420
NFMPKIKSSD	NKVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFRIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLNHNEYML	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNLY	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLME	720
MSASPTFTTE	EQLKNPVDNR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	IVKIKFYSIV	NVLKFKYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLVLYYTQL	GRCMYSLEPI	DISELSSKNI	YQDQHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELQV	FISRQIVETR	960
QATKETANLL	KTICKNSIEIV	YSKAENASRF	RQFEDIVKCR	AVNDLHHMHD	AYINIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLDK	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRSSSKT	NNIGNVQEPK	IINQSITEGLF	ENEVDLLKL		1369

SEQ ID NO: 71 moltype = DNA length = 4107
 FEATURE Location/Qualifiers
 misc_feature 1..4107
 note = Wild-type OMNI-50
 source 1..4107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 71

accaaggtga	aggactacta	cataggcttg	gacatcgcca	cctctagcgt	cggggtgggcc	60
gtcaccgatg	aagcctataa	cgtgcttaag	tttaatagca	agaaaatgtg	gggcgtgcgg	120
ctgttcgacg	acgctaagac	ggcagaggag	cgtaggggcc	agcagaggag	aagacgacgt	180
ctggatcgga	agaaggagag	actcagcctg	ctgcaggact	tcttcgcgca	agaggtagca	240
aaggtcgacc	ccaactctct	cctcaggctg	gacaattccg	atctgtacat	ggaagataag	300
gaccagaaac	tgaagaagcaa	atatacactg	ttcaacgaca	aggacttcaa	ggataagaat	360
tttcataaga	agtacccac	aatacatcac	ctgctgatgg	atctgatcga	ggacgacagt	420
aagaaggaca	tccggctcgt	ctacctggcc	tgtcactatt	tgctcaagaa	cagggggtcat	480
ttcatcttcg	agggccagaa	gttcgacact	aaatcaagct	tcgagaacag	tttgacgag	540
ctcaaaagtc	atttgaacga	cgagatgga	ctggacctcg	aatttgacaa	cgagaacctg	600
attaacatct	tgaactgacc	aaaactcaat	aaaacggcca	agaagaagga	gctgaagtcc	660
gtaatcgccg	acaccaaagt	cctcaaaagc	gtttccgcga	taatgatcgg	ctctagccag	720
aaactcgtcg	acttggtcga	gaaccccgag	gatttcgacg	actctgcgat	aaagtccgtt	780
gacttctcaa	ctacctcttt	cgacgacaag	tactctgact	atgaactcgc	tctgggtgac	840
aagatcgctc	tggtaaacat	ccttaaggaa	atttacgata	gctccatcct	cgagaacctg	900
ctcaaaagag	cagacaagtc	taaggacggt	aacaaatata	tcagtaatgc	attcgtgaag	960
aagtacaata	aacacggaca	agatctgaaa	gagttcaaac	gtctggtacg	acaatatcac	1020
aagagtcgct	attttgatat	tttcagatcc	gagaagggtga	atgacaatta	cgtcagctac	1080
actaaaagct	caattagcaa	caataaacgc	gtcaaaagcaa	acaagttcac	tgatcaagag	1140
gccttctaca	aattcgccaa	gaacatctcg	gagacaatca	agtataagat	caacaaggta	1200
aacggctcca	aggcagatct	ggagctgatt	gacgggatgc	tgccggacat	ggagttcaag	1260
aactttatgc	ccaaaattaa	gtccagtgac	aacgggggtga	ttccatacca	gctcaagctg	1320
atggaattga	acaaaatact	cgagaatcag	tcaaaagcatc	acgagttcct	caatgtcagc	1380
gacgagtacg	gctccgtgtg	tgataaaatc	gcatctatca	tgaggttccg	tatcccttac	1440
tacgtgggac	caatagcaag	tacgcctgga	tcaagaagca	gaaagatagt		1500
gagattactc	cctggaactt	caaggacgtc	gtggacctcg	actccagcag	agaggagttc	1560
attgactcac	tgatcggacg	ctgtacttac	cttaaggacg	agaaggtcct	tcccaagact	1620
tctttgctgt	ataacgaata	catggtgctg	aacgagctga	ataacctgaa	gttgaaacgac	1680
cttcccacat	ccgaggagat	gaagaagaag	atatattgacc	agttgttcaa	aacaagaaaag	1740
aaggtcacc	ttaaagcggt	ggcaaacctg	ctgaagaagg	agttcaacat	caacggcgag	1800
attctgctct	ctgggaccca	cggtgacttc	aagcagggct	tgaactcata	caatgacttc	1860
aaagctatcg	tgggcgataa	agtcgattcc	gatgattacc	gggacaagat	tgaggagatc	1920
attaaactga	tagttcttta	cggtgacgat	aagagttacc	ttcagaagaa	gattaaagct	1980
gggtatggaa	aatacttcac	cgacagttag	attaagaaaa	tggcgggggt	gaactacaag	2040
gattggggaa	ggctctcaaa	gaagctgctg	acgggactcg	aggggtcaaa	caagatcact	2100
ggagagcggg	gctccattat	tcacttcacg	agggaatata	accttaactc	gatggagctt	2160

-continued

```

atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccg ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcoc tgcgcacgtg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
atthtcacgt agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggaa 2400
aaccaactgc tgaagttcta taaagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagagggtact ctactctgct ttctgagata gaggggtgag aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgctgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctgggtg agaaggacct taactccaag 2700
aagggttaaca gctatcctat ccagacgaa atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt acgataaagg tcttatcggt cagaagaagt acactcggtt gacccggaga 2820
actggcttca cggacgacga gctcgctcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggtt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca gtttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcaacaaga atgaactaca ttagtctgg agcgtctttc 3300
aacgccactc tccagcgga gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtagcg cggtacacaa aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt taaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga acaacgacaa gaagctgtcc gacgagattt tctcaaatga cctcaaggac 3540
cgatttggca tcgaggacta cagggttgtc taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctctatcaca ggcggttctg ataagacctt ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctgggtg 3780
tataacactt tcgactatct cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggctcaagg cgattcgggc atccggcggtg 3960
aggcaggatc tcaagtcacat gcacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 72      moltype = DNA length = 4107
FEATURE            Location/Qualifiers
misc_feature       1..4107
                    note = V3795
source             1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 72
accaaagggtga aggaactacta cataggettg gacatcgagg cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataa cgtgcttaag ttttaatagca agaaaatgtg gggcgctgcg 120
ctgttcgacg acgctaagac ggcagaggag cgtaggggcc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgccga agaggtagca 240
aaggctcgacc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaaacaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccaccac aatcacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa cagggggtcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag ttgaaacgag 540
ctcaaaagttc atttgaacga cagatattga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcgccg acaccaagtt cctcaagacc gtttcgcgca taatgatcgg ctctagccag 720
aaactcgtcg acttgttcga gaaccccgag gatttcgacg actctgcgat aaagtccggt 780
gacttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggccaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctgggtacg acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggagatct ggagctgatt gacgggatgc tgcgggacat ggagtccaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaaattga acaaaatact cgagaatcag tcaaaagcatc acgagttcct caatgtcagc 1380
gacgagtagc gctccgtgtg tgataaaatc gcatctatca tggaaattat tattccgtat 1440
tacgtggggc cgctgaaccc gaacagcaaa tatgcgtgga tcaagaagca gaaagatagt 1500
gagattactc cctggaaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcgagcg ctgtacttac cttaaggacg agaaggtcct tcccaaaagt 1620
tctttgtctg ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ctaggagatg gaagaagaag atatttgacc agttgttcaa aacaagaaga 1740
aaggtaaccc ttaaaagcgt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccca cggtgacttc aagcagggtc tgaactcata caatgacttc 1860
aaagctatcg tggcgcataa agtcgattcc gatgattacc gggacaagat ttaggagatc 1920
attaaactga tagttcttta cgggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagttag attaagaaaa tggcgggggt gaactacaag 2040

```

-continued

```

gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcacg aggggaatata accttaactc gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccggt ggatgaccgc 2220
aagctgtcat acgaaatggg ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcacgtg cgacgagata aagaacatca tgggaacgca cagtaagaag 2340
attttcatcg agatgggtcg gggtaaggaa gaggtgaaag cccgcaaggga gtcaaggaa 2400
aacgtcctga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcga 2460
gagaggtaact cttacttgct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgctgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctgggtga agaaggcact taactccaag 2700
aagggtaaca gctatcctat cccagacgaa atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt acgataaaag tcttatcggg cagaagaagt aactcgggt gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgctg 3120
agctacaatc tcgagaatat gttcaagtag gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gtccacaaga atgaactaca ttgagcttgg agcgtctttc 3300
aacgccactc tccagcggaa gaataagggt tccagacccc tgaaggacaa aggccgaaa 3360
tcttccatcg agaagtcagg gcgctacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gcgaacaga agaacgacaa gaagctgttc gacgagattt tctcaaagta cctcaaggag 3540
cgatttgcca tcgaggacta cagggttgct taccagtggt tgaaaaatgc cacactgctc 3600
aagatcgacg gcagctacta ctccatcaca ggcggttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct cctaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgacgga ggagctgggt 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagcttttc 3840
cttaacttgt ttaaggacga caagatttag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgaagac acttctcatg ctggctcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtcctat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggaagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tgcgtctcct ggaagctg 4107

```

```

SEQ ID NO: 73      moltype = DNA length = 4107
FEATURE            Location/Qualifiers
misc_feature       1..4107
                    note = V6918
source             1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 73
accaagggtga aggactacta cataggtctg gacatcgcca cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctgttcgacg acgctaagac ggcagaggag cgtagggggc agcagaggag aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgccga agaggtatga 240
aaggtcgacc ccaactcttt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggaactcaa ggataagaat 360
tttcataaga agtccccac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgcacaagaa caggggtcat 480
ttcatcttgc agggccagaa gttcgacact aaatcaagct tcgagaacag ttgaaacgag 540
ctcaaaagtc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgacc aaaaactcaat aaaacggcca agaagaaggga gctgaagctc 660
gtaatcggcg acaccaagtt cctcaaagcc gtttccgca taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgat aaagtccgtt 780
gacttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggteaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtaacaata aacacggaca agatctgaaa gagttcaaac gtcgtggtacg acaatatcac 1020
aagagtcgct attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaaagcac acgagttcct caatgtcagc 1380
gacgagtacg gctccgtgtg tgataaaatc gcatctatca tggagttccg tatccctac 1440
tacgtgggac cctgaaccc caatagcaag tacgcctgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac gctcggaag ctgtactttac cttaaaggacg agaaggtcct tccaaagct 1620
tctttgctgt ataacgaata catgggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaaaagcgt ggcaaacctg ctgaagaagg agttcaacat caacggcag 1800
attctgctct ctgggaccca cgggtacttc aagcagggtc tgaactcata caatgacttc 1860
aaagctatcg tggggcataa agtcgattcc gatgattacc gggacaagat tgaaggagatc 1920

```

-continued

```

attaaactga tagttcttta cgttgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagtggag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcacg agggaaatata acctaatct gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccg ggatgaccgc 2220
aagctgtcat acgaaatggg ggacgaactg tacctttctc ccagtggtga acggatgctc 2280
tggcagtcct tgcgcacgtg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag ccgcgaagga gtcaagggaag 2400
aacggtctgc tgaagtctta taagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact ctacttgct tcttgagata gagggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgctgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggttaaca gcctatcctat ccagacgaa atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gaccgcgaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gctacatca acatcatagt ggggaacgtg 3060
tataaaccca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgctg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcacaaaga atgaactaca ttgagtcgtg agcgttttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtcagg cggtacacaa aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgctgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttctc taccagtggt tgaaaatcg cactactgctc 3600
aagatcgacg ccagctacta ctctacaca ggcggttctg ataagacct ggagtgtcga 3660
tctgtctcgc agctgattct cctaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcagc cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagttcat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 74      moltype = DNA length = 4107
FEATURE
misc_feature       1..4107
                    note = V6919
source              1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 74
accaagggtga aggactacta cataggcttg gacatcgcca cctctagcgt cgggtgggccc 60
gtcaccgatg aagcctataa cgtgcttaag ttaatagca agaaatgtg gggcgtgctg 120
ctgttcgacg acgctaagag ggtaggggccc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcgca agaggtagca 240
aaggtcgacc ccaactctct cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggaacttcaa ggataagaat 360
tttcataaga agtaccaccac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa caggggctcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag tttgaacgag 540
ctcaaaagttc atttgaacga cgagatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaacgggcca agaagaagga gctgaagtcc 660
gtaatcgccg acaccaagtt cctcaagacc gtttcgcgca taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgat aaagtccgtt 780
gactctcaca ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtaaacat ctttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtacg acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtg atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagtccaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtg ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtacg gctccgtgtg tgataaaatc gcactatca tggagttccg tatcccttac 1440
tacgtgggac ccttgaaccc caatagcaag tacgcctgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcgagcg ctgtacttac cttaaggacg agaaggtcct tcccaaaagt 1620
cttttgctgc ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680
tctcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcacc c ttaaagcggg ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800

```

-continued

```

attctgctct ctgggaccga cgttgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggogataa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaacttga tagttcttta cgttgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggatgggaa aatacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggtctctcaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcctg agggaaata accttaactc gatggagctt 2160
atgtcagctt cattacgtt ccacgaagag atacagaaac ttaaccccggt ggatgaccgc 2220
aagctgtcat acgaaatggg ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcactcg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
atthtcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggag 2400
aaccaactgc tgaagttcta taagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggctact cttacttgct ttctgagata gaggggtgag aagagaataa gtttcgatgg 2520
gataacctgt accttatta tactcaactg ggtcgtcgca tgtactcttt ggaacctatc 2580
gacatctctg acgtgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttaccg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggttaaca gctatcctat ccacagcga atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt acgataaagg tcttatcggt cagaagaagt acactcggtt gacccggaga 2820
actggttcca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaactaga 2880
caagcaacaa cggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcgaagag tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcg 3120
agctacaatc tggagaatatt gttcaagtag gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcaacaaga atgaactaca ttgagtcgtg agcgtcttct 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtacgg cggctacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaaacaga atgaacgaca gaagctgtcc gacgagattt tctcaaatga cctcaaggac 3540
cgatttggca tcgaggacta cagggttgtc taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ttcatcaca ggcggttctg ataagacctt ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat gttcaacaag ttcaagacca gtgtcttcaa gaagtcttct 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggctcaagg cgattcgggc atccggcggt 3960
agcgaggatc tcaagtccat gcacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 75      moltype = DNA length = 4107
FEATURE            Location/Qualifiers
misc_feature       1..4107
                    note = V6920
source             1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 75
accaaaggta aggactacta cataggett gacatcgga cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcgtgctg 120
ctggttcgacg acgctaagac ggcagaggag cgtagggggc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgccga agaggtagca 240
aaggctgaccc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataaagaat 360
tttcataaga agtaccctac aatcacatc ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa caggggtcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag tttgaacgag 540
ctcaaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaagtt cctcaaagcc gtttccgca taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgat aaagtccgtt 780
gacttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc ttggtcaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaggag cagacaagtc taaggacggg aaacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtcgggtacg acaatatcac 1020
aagagtgctg attttgatat ttctcagatc gagaaaggta atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagca acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtcacgtgac aacggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaatcat cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtacg gctcgtgtg tgataaaatc gcatctatca tggagttccg tatccctac 1440
tacgtgggac cctcgaaccc caatagcaag tacgcctgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tctcggacg ctgtacttac cttaaggacg agaaggtcct tcccaagct 1620
tctttgctgt ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680

```

-continued

```

cttccccca cccaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaagcgggt ggcaaacctg ctgaagaagg agttcaacat caacggcggag 1800
attctgctct ctgggaccga cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tggggcgataa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatactctac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcacg agggaatata accttaactc gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccggt ggatgaccgc 2220
aagctgtcat acgaaatggg ggaagaaactg tacctttctc ccagtgtgaa acggatgctc 2280
tgccagctcc tgcccatcgt cgacgagata aagaacatca tgggaacgca cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaagggaag 2400
aaccaactga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggttat cttacttgct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgctgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctgggtga agaaggacct taactccaag 2700
aagggtaaca gctatcctat ccagacgcaa atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt aactcggct gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcgg 3120
agctacaatc tcgagaatat gttcaagta gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcacaaaga atgaactaca ttgagcttgg agcgtttttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg acaagtagcg cggctacaca aacatcaata aagcctgttt cgtgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaaacaga tgaacgacaa gaagctgttc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tggaggacta cagggttgtc taccagtggt tgaaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctccatcaca ggcggttctg ataagacct ggagtgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctgggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt tcaaggacga caagatttag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgaagac acttctcatg ctgggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtcctc cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca gaagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tgcgtctcct ggaagctg 4107

```

```

SEQ ID NO: 76      moltype = DNA length = 4107
FEATURE            Location/Qualifiers
misc_feature       1..4107
                    note = V6920
source             1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 76
accaagggtga aggactacta cataggtctg gacatcgcca cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa ctgctttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctgttcgacg acgctaagac ggcagaggag cgtaggggcc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcgga agaggttagca 240
aaggctcgacc ccaactctct cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccctac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt cctacgtgcc tgtcactatt tgcacaagaa cagggggtcat 480
ttcatcttct agggccagaa gttcgacact aaatcaagct tcgagaacag ttgaaacgag 540
ctcaaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
atttaacatct tgactgaccc aaaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaagtt cctcaaagcc gtttcgcgca taatgatcgg ctctagccag 720
aaactcgtcg acttgttcga gaaccccgag gatttcgacg actctcgat aaagtccgtt 780
gacttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtaaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagttaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctgggtacg acaatatcac 1020
aagagtgcgt attttgatc ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gcctctcaca aatcgcgcaa gaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca cggcagatct ggagctgatt gacgggatgc tgcgggacat ggagtccaag 1260
aactttatgc ccaaaattaa gtccagttag aacgggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaactact cgagaatcag tcaaaagcat acgagttcct caatgtcagc 1380
gacgagtagc gctccgtgtg tgataaaatc gcatctatca tggaaatttat tattccgtat 1440
tacgtgggccc cgtggaaccc gaacagcaaa tatcggtgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560

```

-continued

```

attgactcac tgateggagc ctgtacttac ctttaaggac agaaggctct tcccaaagct 1620
tctttgctgc ataacgaata catgggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaga 1740
aaggtcaccc ttaaagcggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct cttgggacga cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggcgataa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg agggtgcaaa caagatcact 2100
ggagagcggt gctccattat tcacttcacg agggaatata acctaatct gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccg ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcactcg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttactcg agatggctcg gggttaaggaa gaggtgaaag ccgcgaagga gtcaaggaa 2400
aacgctcctg tgaagttcta taagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact cttacttgct ttctgagata gagggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acccttatta taactcaact ggctcgtgca tgactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggttaaca gctatcctat ccagacgaa atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt acgataaagg tcttatcggt cagaagaagt acactcggtc gaccgggaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactcg taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgctg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcgagata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaaggca caaattatag gttcacaaaga atgaactaca ttgagctctg agcgtctttc 3300
aacgccactc tccagcgga gaataagggt tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtcagg cggtacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaaagta cctcaaggac 3540
cgatttgga ctaggagata cagggttgtc taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgagc gcagctacta ctctacaca ggcggttctg ataagacct ggagttgca 3660
tctgctctgc agctgattct cctcaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aacgaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtcact cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 77      moltype = DNA length = 4107
FEATURE
misc_feature       1..4107
                    note = V6865
source              1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 77
accagggtga aggactacta cataggcttg gacatcgga cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa cgtgcttaag ttaatagca agaaaatgtg gggcgtgctg 120
ctgttcgacg acgctaagac ggcagaggag cgtagggggc agcagaggag aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcca agaggtagca 240
aaggctgacc ccaacttctt cctcagggtg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccctac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa cagggtgcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag tttagacgag 540
ctcaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaacgggcca agaagaagga gctgaagtcc 660
gtaatcggtg acaccaagtt cctcaaagcc gtttcggcga taatgatcgg ctctagccag 720
aaactcgtcg acttgctcga gaaccccgag gatctcgacg actctcgcat aaagtccgtt 780
gacttctcaa ctacctctt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtaacatc ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaggag cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtacg acaatatcac 1020
aagagtcggt attttgatat ttccagatcc gagaaggtag atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggtctca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacggggtga ttccatacca gctcaagctg 1320
atggaattga aaaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtacg gtcctgtgtg tgataaaatc gcatctatca tgggaatttat tattccgtat 1440

```

-continued

tacgtggggc	cgctgaaccc	gaacagcaaa	tatgcgtgga	tcaagaagca	gaaagatagt	1500
gagattactc	cctggaactt	caaggacgtc	gtggacctcg	actccagcag	agaggagttc	1560
attgactcac	tgatcggacg	ctgtacttac	cttaaggacg	agaaggctct	tcccaagct	1620
tctttgtctg	ataacgaata	catggtgctg	aacgagctga	ataacctgaa	gttgaacgac	1680
cttcccatca	ccgaggagat	gaagaagaag	atatttgacc	agttgttcaa	aacaagaaaag	1740
aagggtcacc	ttaaacgggt	ggcaaacctg	ctgaagaagg	agttcaacat	caacggcgag	1800
attctgctct	ctgggaccga	cggtgacttc	aagcagggct	tgaactcata	caatgacttc	1860
aaagctatcg	tgggogataa	agtcgattcc	gatgattacc	gggacaagat	tgaggagatc	1920
attaacttga	tagttcttta	cggtgacgat	aagagttaac	ttcagaagaa	gattaaagct	1980
gggtatggaa	aatacttcac	cgacagttag	attaagaaaa	tggcggggct	gaactacaag	2040
gattggggaa	ggctctcaaa	gaagctgctg	acgggactcg	aggggtgcaa	caagatcact	2100
ggagagcggg	gctccattat	tcacttcatg	agggaatata	accttaactc	gatggagctt	2160
atgtcagctt	cattacgttt	accgaagag	atacagaaac	ttaacccctg	ggatgaccgc	2220
aagctgctat	acgaaatggg	ggacgaactg	tacctttctc	ccagtgtgaa	acggatgctc	2280
tggcagctcc	tgcccatctg	cgacgagata	aagaacatca	tgggaaccga	cagtaagaag	2340
atthtcatcg	agatggctcg	gggttaaggaa	gaggtgaaag	cccgaaggaa	gtcaagggaag	2400
aaccaactga	tcaagtctta	taagacgga	aagaaggcat	tcatacgcga	gattggcagag	2460
gagaggtact	cttacttgct	ttctgagata	gaggggtgag	aagagaataa	gtttcgatgg	2520
gataacctgt	acctttatta	tactcaactg	ggtcgtgca	tgtactcttt	ggaacctatc	2580
gacatatctg	agctgtcttc	aaagaatatt	tacgatcagg	atcatatcta	ccccaaaagc	2640
aagatttacg	acgacagtat	cgagatatag	gtgctgggtg	agaaggacct	taactccaag	2700
aaaggtaaca	gctatcttat	cccagacgaa	atcctgaaca	agaactgtta	cgcctactgg	2760
aagatcctgt	acgataaagg	tcttatcggt	cagaagaagt	acactcggct	gacccggaga	2820
actggcttca	cggaacgaga	gctcgttcag	ttcatctcaa	gacagatcgt	ggaactaga	2880
caagcaacaa	aggagactgc	taacctgctc	aagacaatat	gtaagaactc	cgagatcgtg	2940
tattccaaag	ccgagaacgc	aaagtcgggt	aggcaagagt	tcgacatcgt	gaagtgtagg	3000
gcggtgaacg	atcttcatca	tatgcacgat	gcctacatca	acatcatagt	ggggaacgtg	3060
tataacacca	agttcacgaa	ggaccctatg	aatttcgtaa	agaagcagga	aaaggcgcgg	3120
agctacaatc	tcgagaatat	gttcaagtac	gatgtgaaac	gtggcggata	caccgcttgg	3180
atcgccgatg	acgagaaggg	caccgtgaag	aacgcgagta	ttaaacgtat	ccggaaggag	3240
ctggaaaggca	caatttatag	gttcacaaga	atgaactaca	ttgagctctg	agcgtctttc	3300
aacgccactc	tccagcgga	gaataagggc	tccagacccc	tgaaggacaa	aggcccgaaa	3360
tcttccatcg	agaagtacgg	cggtctacaca	aacatcaata	aagcctgttt	cgctgttctt	3420
gacatcaagt	ctaagaacaa	gattgagagg	aagctgatgc	ccgtcgagcg	tgagatctat	3480
gccaacacaga	agaacgacaa	gaagctgtcc	gacgagattt	tctcaaagta	cctcaaggac	3540
cgatttggca	tcgaggacta	cagggttgtc	taccagtggt	tgaanaatgc	cacactgctc	3600
aaagtcgacg	gcagctacta	cttcatcaca	ggcgggtctg	ataagacctc	ggagttgcga	3660
tctgtctctg	agctgattct	ccctaagaag	aacgagtggt	cgatcaaaac	gatcgacaa	3720
tcttccgaaa	acgactatct	gacgatcgag	cgatccagg	acctgaccca	ggagctgggtg	3780
tataacactt	tcgacatcta	cgtaacaaga	ttcaagacca	gtgtcttcaa	gaagtctttc	3840
cttaacttgt	ttcaggacga	caagatttag	aacattgact	tcaagtttaa	gtccatggac	3900
ttcaaggaga	aatgcaagac	acttctcatg	ctggctcaag	cgattcgggc	atccggcgtg	3960
aggcaggatc	tcaagttccat	cgacctcaag	tctgattacg	gacgggtcag	ttcaagacc	4020
aacaacatcg	gcaattacca	ggagttcaag	attattaatc	agtcacatc	tggaactgtc	4080
gagaatgagg	tcgatctcct	gaagctg				4107

SEQ ID NO: 78 moltype = DNA length = 4107
 FEATURE Location/Qualifiers
 misc_feature 1..4107
 note = V6866
 source 1..4107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 78

accaaagtga	aggactacta	cataggett	gacatcgga	cctctagcgt	cggtggggcc	60
gtcaccgatg	aagcctataa	cgtgcttaag	tttaatagca	agaaaatgtg	gggcgtgctg	120
ctgttcgacg	acgctaagac	ggcagaggag	cgtagggggc	agcagaggagc	aagacgacgt	180
ctggatcgga	agaaggagag	actcagcctg	ctgcaggact	tcttcgccga	agaggtagca	240
aaggctcgacc	ccaacttctt	cctcaggctg	gacaattccg	atctgtacat	ggaagataag	300
gaccagaaac	tgaaaagcaa	atatacactg	ttcaacgaca	aggacttcaa	ggataagaat	360
tttcataaga	agtaacccac	aatacatcac	ctgctgatgg	atctgatcga	ggacgacagt	420
aagaaggaca	tccggctcgt	ctacctggcc	tgtcactatt	tgctcaagaa	cagggggtcat	480
ttcatcttcg	agggccagaa	gttcgacact	aaatcaagct	tcgagaacag	tttgaacgag	540
ctcaaagttc	atttgaacga	cgagtatgga	ctggacctcg	aatttgacaa	cgagaacctg	600
atttaacatc	tgactgacct	aaaactcaat	aaaacggcca	agaagaaggga	gctgaagtcc	660
gtaatcggcg	acaccaagtt	cctcaaagcc	gtttccgcga	taatgatcgg	ctctagccag	720
aaactcgtcg	acttgttcga	gaaccccgag	gatttcgacg	actctgcgat	aaagtccgtt	780
gacttctcaa	ctacctcttt	cgacgacaag	tactctgact	atgaactcgc	tctgggtgac	840
aagatcgctc	tggtaacat	ccttaaggaa	atttacgata	gctccatcct	cgagaacctg	900
ctcaaaggag	cagacaagtc	taaggacggt	aacaaatata	tcagttaatgc	attcgtgaag	960
aagtagaata	aacacggaca	agatctgaaa	gagttcaaac	gtctggtacg	acaatatcac	1020
aagagtgcgt	atthtgcgat	tttcagatcc	gagaagggtg	atgacaatta	cgtaagctac	1080
actaaaagct	caatttagcaa	caataaacgc	gtcaaagcaa	acaagttcac	tgatcaagag	1140
gccttctaca	aattcgccaa	gaaacatctg	gagacaatca	agtataagat	caacaaggta	1200
aacggctcca	aggcagatct	ggagctgatt	gacgggatgc	tgccgggacat	ggagttcaag	1260
aactttatgc	ccaaaattaa	gtccagtgac	aacgggggtg	ttccatacca	gctcaagctg	1320

-continued

atggaattga	acaaaatact	cgagaatcag	tcaaagcatc	acgagttcct	caatgtcagc	1380
gacgagtagc	gctcogtgtg	tgataaaaac	gcatactatca	tggaaatttat	tattccgtat	1440
tacgtggggcc	cgctgaaccc	gaacagcaaaa	tatgcgtgga	tcaagaagca	gaaagatagt	1500
gagattactc	cctggaactt	caaggacgtc	gtggacctcg	actccagcag	agaggagtgc	1560
attgactcac	tgtacggacg	ctgtacttac	cttaaggacg	agaaggctct	tcccaagct	1620
tctttgctgt	ataacgaata	catgggtgctg	aacgagctga	ataacctgaa	gttgaacgac	1680
cttcccatca	ccgaggagat	gaagaagaag	atatttgacc	agttgttcaa	aacaagaaag	1740
aaggtcaccc	ttaaagcggg	ggcaaacctg	ctgaagaagg	agttcaacat	caacggcag	1800
attctgctct	ctgggaccga	cggtgacttc	aagcagggct	tgaactcata	caatgacttc	1860
aaagctatcg	tggggcgata	agtcgattcc	gatgattacc	gggacaagat	tgaggagatc	1920
attaaactga	tagttcttta	cggtagcgat	aagagttacc	ttcagaagaa	gattaaagct	1980
gggtatggaa	aatactctac	cgacagtgag	attaagaaaa	tggcggggct	gaactacaag	2040
gattggggaa	ggctctcaaa	gaagctgctg	acgggactcg	aggggtgcaa	caagatcact	2100
ggagagcggg	tgcctcattat	tcacttcacg	agggaaatata	accttaactc	gatggagctt	2160
atgtcagctt	catttacgtt	caccgaagag	atacagaaac	ttaacccctg	ggatgaccgc	2220
aagctgtcat	acgaatagg	ggacgaactg	tacctttctc	ccagtgtgaa	acggatgctc	2280
tggcagtcct	tgcgcactcg	cgacgagata	aagaacatca	tgggaacgca	cagtaagaag	2340
attttcatcg	agatgggtcg	gggtaaggaa	gaggtgaaag	cccgaacgga	gtcaagggaag	2400
aacgtctcga	tcaagttcta	taaagacgga	aagaaggcat	tcatcagcga	gattggcag	2460
gagaggtact	cttacttgct	tctctgagata	gagggtgagg	aagagaataa	gtttcgatgg	2520
gataacctgt	acctttatta	tactcaactg	ggctcgctgca	tgtactcttt	ggaacctatc	2580
gacatatctg	agctgtcttc	aaagaatatt	tacgatcagg	atcatatcta	ccccaaaagc	2640
aagattttacg	acgacagtat	cgagaatagg	gtgctggtga	agaaggacct	taactccaag	2700
aagggtaaca	gctatcctat	cccagacgaa	atcctgaaca	agaactgtta	cgctactggt	2760
aaagtcctgt	acgataaagg	tcttatcggg	cagaagaagt	acactcggct	gaccgcgaga	2820
actggcttca	cggacgacga	gctcgttcag	tctatctcaa	gacagatcgt	ggaaactaga	2880
caagcaacaa	aggagactgc	taacctgctc	aagacaatat	gtaagaactc	cgagatcgtg	2940
tattccaaag	ccgagaacgc	aagtcggttt	aggcaagagt	tcgacatcgt	gaagtgtagg	3000
gcggtgaacg	atcttcatca	tatgcacgat	gcctacatca	acatcatagt	ggggaaacgtg	3060
tataaacacca	agttcacgaa	ggacctatg	aatttcgtaa	agaagcagga	aaaggcgcgg	3120
agctacaatc	tcgagaatat	gttcaagtag	gatgtgaaac	gtggcgggata	caccgcttgg	3180
atcgccgatg	acgagaaggg	caccgtgaag	aacgcgagta	ttaaacgtat	ccggaaggag	3240
ctggaaggca	caaattatag	gttcacaaga	atgaactaca	ttagtcttgg	agcgcttttc	3300
aaagccactc	tccagcggaa	gaataagggc	tccagacccc	tgaaggacaa	aggcccgaaa	3360
tcttccatcg	agaagtacgg	gggtacacac	aacatcaata	aagcctgttt	cgctgttctt	3420
gacatcaagt	ctaagaacaa	gattgagagg	aagctgatgc	ccgtcgagcg	tgagatctat	3480
gccaacacga	agaacgacaa	gaagctgtcc	gacgagattt	tctcaagta	cctcaaggac	3540
cgatttgcca	tgcaggacta	cagggttgct	taccagtggt	tgaaaaatgcg	cacactgctc	3600
aagatcgacg	gcagctacta	cttcatcaca	ggcgggtctg	ataagacctc	ggagtgtcga	3660
tctgctctgc	agctgattct	cctcaagaag	aacgagtggt	cgatcaaaaca	gatcgacaag	3720
tcttccgaaa	acgactatct	gacgatcgag	cgatccacgg	acctgaccga	ggagctgggtg	3780
tataacactt	tgcacatcat	cgtcaacaag	tccaagacca	gtgtcttcaa	gaagtctttc	3840
cttaacttgt	ttaaggacga	caagattgag	aacattgact	tcaagtttaa	gtccatggac	3900
ttcaaggaga	aatgaagac	acttctcatg	ctgggtcaagg	cgatttcgggc	atccggcgtg	3960
aggcaggatc	tcaagtcctat	cgaacctcaag	tctgattacg	gacggctcag	ttcaagagacc	4020
aacaacatcg	tcaattacca	ggaagttcaag	attattaatc	agtcacatcac	tggactgttc	4080
gagaatgagg	tgcattctct	ggaagctg				4107

SEQ ID NO: 79 moltype = DNA length = 4107
FEATURE Location/Qualifiers
misc_feature 1..4107
 note = V6867
source 1..4107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 79

accaagggtga	aggactacta	cataggcttg	gacatcgcca	cctctagcgt	cgggtggggcc	60
gtcaccgatg	aagcctataa	cgtgcttaag	tttaatagca	agaaaatgtg	gggcgtgcgg	120
ctgttcgacg	acgctaagac	ggcagaggag	cgtagggggc	agcagaggagc	aagacgacgt	180
ctggatcgga	agaaggagag	actcagcctg	ctgcaggact	tcttcgccga	agaggtagca	240
aaagtcgacc	ccaactctct	cctcaggctg	gacaattccg	atctgtacat	ggaagataag	300
gaccagaaac	tgaaaagcaa	atataactg	ttcaacgaca	aggacttcaa	ggataaagaat	360
tttcataaga	agtccccac	aatacatcac	ctgctgatgg	atctgatcga	ggacgacagt	420
aagaaggaca	tccggctcgt	ctacctggcc	tgtcactatt	tgtcgaagaa	cagggtgcat	480
tctatcttcg	agggccagaa	gttcgacact	aaatcaagct	tgcagaacag	tttgaacgag	540
ctcaaaagtc	atttgaacga	cgagtatgga	ctggacctcg	aatttgacaa	cgagaacctg	600
attaacatct	tgactgaccc	aaaactcaat	aaaacggcca	agaagaagga	gctgaagtcc	660
gtaatcggcg	acaccaagtt	cctcaaaagg	gtttccgcga	taatgatcgg	ctctagccag	720
aaactcgtcg	acttgctcga	gaaccccgag	gatttcgacg	actctgcgat	aaagtccgtt	780
gacttctcaa	ctacctcttt	cgcagacaag	tactctgact	atgaactcgc	tctgggtgac	840
aagatcgctc	tggtaaacat	ctttaaggaa	atttacgata	gctccatcct	cgagaacctg	900
ctcaaaaggg	gcagacaagtc	taaggacggg	aacaaatata	tcagtaatgc	attcgtgaag	960
aagtacaata	aacacggaca	agatctgaaa	gagttcaaac	gtctggtacg	acaatatcac	1020
aagagtgcgt	attttgatat	tttcagatcc	gagaagggtga	atgacaatta	cgtcagctac	1080
actaaaagct	caattagcaa	caataaacgc	gtcaaaagcaa	acaagttcac	tgatcaagag	1140
gccttctaca	aattcgccaa	gaaacatctg	gagacaatca	agtataagat	caacaaggta	1200

-continued

```

aacgggtcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtagc gctccgtgtg tgataaaatc gcatctatca tggaaattcg tattccgtat 1440
tacgtggggc cgctgaaccc gaacagcaaa tatgctgtga tcaagaagca gaaagatagt 1500
gagattactc cctggaaact caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcggagc ctgtacttac cttaaggacg agaaggtcct tcccaaagct 1620
tctttgctgc ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaaagcggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggacgga cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggcgataa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagtga attaagaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggtgtcaaa caagatcact 2100
ggagagcggg cctccattat tcacttcatt agggaaatata acctaatct gatggagctt 2160
atgtcagctt catctacgtt caccgaagag atacagaaac ttaacctcgt ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcoc tgcgcattct cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag ccgcgaagga gtcaaggaa 2400
aacgtcctga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact cttaactgtt tcttgagata gagggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acttttatta taactcaact ggtcgtgca tgtactctt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaca gctatcctat ccagacgaa atcctgaaca agaactgtta cgctactggt 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gaccocggaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcgaaggt tgcacatcgt gaagtgtagg 3000
gcggtgaacg atcttcacca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgctg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcacaaaga atgaactaca ttgagttctg agcgtctttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtacgg cggtcacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttgct taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta cttcatcaca ggcggttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtccat cgacctcaag tctgattacg gacggctcac ttcaaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 80      moltype = RNA  length = 102
FEATURE
misc_feature      1..102
                  note = sg62 spacer and scaffold sequence
source            1..102
                  mol_type = other RNA
                  organism = synthetic construct

```

```

SEQUENCE: 80
gtgtcaagcc ccagaggcca cagtttgaga gttatgaaaa tgacgagttc aaataaaaaa 60
ttattcaaac cgctatttta taggcgcgag atgttctgct tt 102

```

```

SEQ ID NO: 81      moltype = DNA  length = 22
FEATURE
misc_feature      1..22
                  note = g62 target
source            1..22
                  mol_type = other DNA
                  organism = synthetic construct

```

```

SEQUENCE: 81
gtgtcaagcc ccagaggcca ca 22

```

```

SEQ ID NO: 82      moltype = DNA  length = 20
FEATURE
misc_feature      1..20
                  note = g62OT1
source            1..20

```

-continued

	mol_type = other DNA organism = synthetic construct	
SEQUENCE: 82		
gccaaccccc aaaggccaca		20
SEQ ID NO: 83	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = g62OT2	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 83		
gccaagcctc agaggccaca		20
SEQ ID NO: 84	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
misc_feature	1..22	
	note = g62Alt target	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 84		
gtgtcaagcc ccagaggaca ca		22
SEQ ID NO: 85	moltype = AA length = 9	
FEATURE	Location/Qualifiers	
REGION	1..9	
	note = HA Tag	
source	1..9	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 85		
YPYDVFPDYA		9
SEQ ID NO: 86	moltype = DNA length = 27	
FEATURE	Location/Qualifiers	
misc_feature	1..27	
	note = HA Tag	
source	1..27	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 86		
taccatacag atgttccaga ttacgct		27
SEQ ID NO: 87	moltype = AA length = 7	
FEATURE	Location/Qualifiers	
REGION	1..7	
	note = SV40 NLS	
source	1..7	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 87		
PKKKRKV		7
SEQ ID NO: 88	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = SV40 NLS	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 88		
ccaaaaaaga aaagaaaagt t		21
SEQ ID NO: 89	moltype = AA length = 1369	
FEATURE	Location/Qualifiers	
REGION	1..1369	
	note = V7085 Variant	
source	1..1369	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 89		
TKVKDYYIGL DIGTSSVGWA VTDEAYNVLK FNSKKMWGVR LFDDAKTAAE RRGQRGARR		60
LDRKKERLSL LQDFFAEVA KVDPNFFLR L DNSDLYMEDK DQKLKSKYTL FNDKDFKDKN		120
PHKKYPTIHH LLMDLIEDDS KKDTRLVYLA CHYLLKNRGH FIFEGQKFD T KSSFENSLNE		180

-continued

LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFHIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VDLSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHEYMYL	NELNNKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDGF	KQGLNSYNDP	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFTR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 90 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V7086 Variant
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 90

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARRR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFVIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VDLSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHEYMYL	NELNNKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDGF	KQGLNSYNDP	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFTR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 91 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V7087 Variant
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 91

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARRR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFDIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VDLSSSREEF	IDSILIGRCTY	LKDEKVLPA	540
SLLNHEYMYL	NELNNKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDGF	KQGLNSYNDP	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900

-continued

KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMPKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 92 moltype = AA length = 1369
FEATURE Location/Qualifiers
REGION 1..1369
 note = V7088 Variant
source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 92

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLR	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILEN	SKHHEFLNVS	DEYGSVCDKI	ASIMEFSIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLCNEYMVL	NELNNLKLND	LPITEEMKKK	IPDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNPNVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMPKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 93 moltype = AA length = 1369
FEATURE Location/Qualifiers
REGION 1..1369
 note = V7090 Variant
source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 93

TKVKDYYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDPFAEVA	KVDPNFFLR	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFD	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILEN	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLCNEYMVL	NELNNLKLND	LPITEEMKKK	IPDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNPNVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEFDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMPKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	PKTSVFKKSF	LNLFQDDKIE	NIDFKFKSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEPK	IINQSIITGLF	ENEVDLLKL		1369

SEQ ID NO: 94 moltype = AA length = 1369
FEATURE Location/Qualifiers
REGION 1..1369

-continued

source note = V7091 Variant
1..1369
mol_type = protein
organism = synthetic construct

SEQUENCE: 94

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAAE	RRGQRGARRR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFDT	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDLSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLDNEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDFG	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNPNVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFTR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFQDDKIE	NIDPFKFSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEFK	IINQSITGLF	ENEVDLLKL		1369

SEQ ID NO: 95 moltype = AA length = 1369
FEATURE Location/Qualifiers
REGION 1..1369
note = V7092 Variant
source 1..1369
mol_type = protein
organism = synthetic construct

SEQUENCE: 95

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAAE	RRGQRGARRR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFDT	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDLSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLTNEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTGDFG	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIHFH	REYNLNLML	720
MSASFTFTEE	IQKLNPNVDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLIKFYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTYNRFTR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSNKIER	KLMPVEREIV	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFQDDKIE	NIDPFKFSMD	FKECKCTLML	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEFK	IINQSITGLF	ENEVDLLKL		1369

SEQ ID NO: 96 moltype = AA length = 1369
FEATURE Location/Qualifiers
REGION 1..1369
note = V7093 Variant
source 1..1369
mol_type = protein
organism = synthetic construct

SEQUENCE: 96

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAAE	RRGQRGARRR	60
LDRKKERLSL	LQDFFAEVA	KVDPNFFLR	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FHKKYPTIHH	LLMDLIEDDS	KKDIRLVYLA	CHYLLKNRGH	FIFEGQKFDT	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDLFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNLKE	IYDSSILENL	300
LKEADKSKDG	NKYISNAFVK	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420

-continued

NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLFNEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIIHFM	REYNLNLML	720
MSASFTFTTE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLKPYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREII	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFDQDDKIE	NIDPKFKSMD	FKEKCKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEFK	IINQSITEGLF	ENEVDLLKL		1369

SEQ ID NO: 97 moltype = AA length = 1369
 FEATURE Location/Qualifiers
 REGION 1..1369
 note = V7094 Variant
 source 1..1369
 mol_type = protein
 organism = synthetic construct

SEQUENCE: 97

TKVKDYIYIGL	DIGTSSVGWA	VTDEAYNVLK	FNSKKMWGVR	LFDDAKTAE	RRGQRGARR	60
LDRKKERLSL	LQDFFAEVA	KVDPNPFRL	DNSDLYMEDK	DQKLKSKYTL	FNDKDFKDKN	120
FKHKYPTIHH	LLMDLIEDDS	KKDRLVYLA	CHYLLKNRGH	FIFEGQKFDT	KSSFENSLNE	180
LKVHLNDEYG	LDLEFDNENL	INILTDPKLN	KTAKKKELKS	VIGDTKFLKA	VSAIMIGSSQ	240
KLVDFENPE	DFDDSAIKSV	DFSTTSFDDK	YSDYELALGD	KIALVNILKE	IYDSSILENL	300
LKEADKSDG	NKYISNAFPI	KYNKHGQDLK	EFKRLVRQYH	KSAYFDIFRS	EKVNDNYVSY	360
TKSSISNNKR	VKANKFTDQE	AFYKFAKHL	ETIKYKINKV	NGSKADLELI	DGMLRDMFEK	420
NFMPKIKSSD	NGVIPYQLKL	MELNKKILENQ	SKHHEFLNVS	DEYGSVCDKI	ASIMEFIIPY	480
YVGPLNPNSK	YAWIKKQKDS	EITPWNFKDV	VLDSSSREEF	IDSLIGRCTY	LKDEKVLPA	540
SLLANEYMVL	NELNNLKLND	LPITEEMKKK	IFDQLFKTRK	KVTLKAVANL	LKKEFNINGE	600
ILLSGTDGDF	KQGLNSYNDF	KAIVGDKVDS	DDYRDKIEEI	IKLIVLYGDD	KSYLQKKIKA	660
GYGKYFTDSE	IKKMAGLNYK	DWGRLSKKLL	TGLEGANKIT	GERGSIIHFM	REYNLNLML	720
MSASFTFTTE	IQKLNVPDDR	KLSYEMVDEL	YLSPSVKRML	WQSLRIVDEI	KNIMGTDSSK	780
IFIEMARGKE	EVKARKESRK	NVLKPYKDG	KKAFISEIGE	ERYSYLLSEI	EGEEENKFRW	840
DNLYLYYTQL	GRCMYSLEPI	DISELSSKNI	YDQDHIYPKS	KIYDDSIENR	VLVKKDLNSK	900
KGNSYPIPDE	ILNKNCYAYW	KILYDKGLIG	QKKYTRLTRR	TGFTDDELVQ	FISRQIVETR	960
QATKETANLL	KTICKNSEIV	YSKAENASRF	RQEPDIVKCR	AVNDLHHMHD	AYINIIVGNV	1020
YNTKFTKDPM	NFVKKQEKAR	SYNLENMFKY	DVKRGGYTAW	IADDEKGTVK	NASIKRIRKE	1080
LEGTNYRFR	MNYIESGALF	NATLQRKNKG	SRPLKDKGPK	SSIEKYGGYT	NINKACFAVL	1140
DIKSKNKIER	KLMPVEREII	AKQKNDKKLS	DEIFSKYLKD	RFGIEDYRVV	YPVVKMRTL	1200
KIDGSYYFIT	GGSDKTLELR	SALQLILPKK	NEWAIKQIDK	SSENDYLTIE	RIQDLTEELV	1260
YNTFDIIVNK	FKTSVFKKSF	LNLFDQDDKIE	NIDPKFKSMD	FKEKCKTLLM	LVKAIRASGV	1320
RQDLKSIDLK	SDYGRLLSKT	NNIGNYQEFK	IINQSITEGLF	ENEVDLLKL		1369

SEQ ID NO: 98 moltype = DNA length = 4107
 FEATURE Location/Qualifiers
 misc_feature 1..4107
 note = V7085 Variant
 source 1..4107
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 98

accagggtga	aggactacta	cataggcttg	gacatcgcca	cctctagcgt	cgggtgggcc	60
gtcaccgatg	aagcctataa	cgtgcttaag	tttaatagca	agaaaatgtg	gggcgtgcgg	120
ctgttcgacg	acgctaagac	ggcagaggag	cgtaggggcc	agcgaggagc	aagacgacgt	180
ctggatcgga	agaaggagag	actcagcctg	ctgcaggact	tcttcgccga	agaggtagca	240
aaggctcgacc	ccaacttctt	cctcaggctg	gacaattccg	atctgtacat	ggaagataag	300
gaccagaaac	tgaaaagcaa	atacactctg	ttcaacgaca	aggacttcaa	ggataagaat	360
tttcataaga	agtacccccc	aatacatcac	ctgctgatgg	atctgatcga	ggacgacagt	420
aagaaggaca	tccggctcgt	ctaccctggc	tgctactatt	tgctcaagaa	caggggtcat	480
ttcatcttcg	agggccagaa	gttcgacact	aaatcaagct	tcgagaacag	tttgaacgag	540
ctcaaagtct	atttgaacga	cgagtatgga	ctggacctcg	aatttgacaa	cgagaacctg	600
atatacatct	tgaactgacc	aaaactcaat	aaacggcca	agaagaagga	gctgaagtcc	660
gtaatcgccg	acaccaagtt	cctcaaagcc	gtttccgcga	taatgatcgg	ctctagccag	720
aaactcgtcg	acttgttcga	gaaccccgag	gatttcgacg	actctcgcat	aaagtccgtt	780
gacttctcaa	ctacactctt	cgacgacaag	tactctgact	atgaactcgc	tctgggtgac	840
aagatcgctc	tggtcaacat	ccttaaggaa	atttaacgata	gctccatcct	cgagaacctg	900
ctcaaagagg	cagacaagtc	taaggacggt	aacaaatata	tcagtaatgc	attcgtgaag	960
aagtacaata	aacacggaca	agatctgaaa	gagttcaaac	gtctggtacg	acaatatcac	1020
aagagtgcgt	attttgatat	tttcagatcc	gagaagggtga	atgacaatta	cgtcagctac	1080
actaaaagct	caattagcaa	caataaacgc	gtcaaaagcaa	acaagttcac	tgatcaagag	1140

-continued

```

gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggtgga ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtacg gctcogtggt tgataaaatc gcatctatca tggaaattca tattccgtat 1440
tacgtggggc cgctgaaccc gaacagcaaa tatgcgtgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac ttgacggacg ctgtacttac cttaaggacg agaaggtcct tcccaaagct 1620
tctttgctgc ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaagcgggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccga cggtgacttc aagcagggtc tgaactcata caatgacttc 1860
aaagctatcg tggggcgata agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcacg agggaaatata accttaactc gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccggt ggatgaccgc 2220
aagctgtcat cgaagaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcactcg cgacgagata aagaacatca tgggaaacga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggaa 2400
aacgtctcga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcga 2460
gagaggtact cttacttgct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtgcgtgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaca gctatcctat ccagacgcaa atcctgaaca agaactgtta cgctactggt 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaactaga 2880
caagcaacaa agagagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgagg 3120
agctacaatc tcgagaatat gttcaagtag gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gtccacaaga atgaactaca ttgagctggt agcgtttttc 3300
aacgccactc tccagcgaa gaataagggc tccagacccc tgaaggacaa aggccgaaa 3360
tcttccatcg agaagtacgg cggtacacaa aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaaacaga gaaacgacaa gaagctgttc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttgct taccagtggt tgaaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctccatcaca ggcggttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaaca gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctgggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtccat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca gaagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tgcattctct gaagctg 4107

```

```

SEQ ID NO: 99          moltype = DNA length = 4107
FEATURE               Location/Qualifiers
misc_feature          1..4107
                        note = V7086 Variant
source                1..4107
                        mol_type = other DNA
                        organism = synthetic construct

```

```

SEQUENCE: 99
accaagggtga aggactacta cataggcttg gacatcgcca cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataaa cgtgccttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctgttccgacg acgctaagac ggcagaggag cgtagggggc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgccga agaggtagca 240
aaggctcgacc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggaactcaa ggataaagaat 360
tttcataaga agtaccacac aatcacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa cagggggtcat 480
ttcatcttgc agggccagaa gttcgacact aaatcaagct tcgagaacag tttgaacgag 540
ctcaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaacgggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaagtt cctcaaaagg gtttccgcga taatgatcgg ctctagccag 720
aaactcgtcg acttgttcca gaaccccgag gatttcgacg actctcgcat aaagtcogtt 780
gactttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtaacaat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtacg acaatatcac 1020

```

-continued

```

aagagtgcgt attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaacatctcg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaactact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtacg gctccgtgtg tgataaaatc gcatctatca tggaaattgt gatcccgat 1440
tacgtgggccc cgctgaaccc gaacagcaaa tatgctgtga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcggacg ctgtacttac cttaaggacg agaaggtcct tcccaaagct 1620
tctttgtctg ataacgaata catggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcacc cttaaagcggg ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccca cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggcgataa agtcgattcc gatgattacc gggacaagat tgaaggagatc 1920
atataactga tagttcttta cggtgacgat aagagttacc ttcaagaaga gattaaagct 1980
gggtatggaa aatactctac cgacagtga attaagaaaa tggcggggct gaactacaa 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg agggtgcaaa caagatcact 2100
ggagagcggg cctccattat tcacttcag agggaaatata acctaatct gatggagctt 2160
atgtcagctt gatctacgtt caccgaagag atacagaaac ttaaccccg ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcacgtg gcacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag ccgcgaagga gtcaaggaa 2400
aacgtcctga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact cttactgtct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta taactcaactg ggtcgtctga tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaca gctatcctat ccagacgaa atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tgcacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgagg 3120
agctacaatc tcgagaatat gttcaagtag gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcacaaaga atgaactaca ttgagctctg agcgtctttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtcagg cggtacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcagcgc tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttgtc taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctctacaca ggcggttctg ataagacctt ggagttgcga 3660
ctctctctgc agctgattct cctcaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatcgcaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtcctat cgacctcaag tctgattacg gacgggtcac ttcaaaagac 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 100      moltype = DNA length = 4107
FEATURE
misc_feature        1..4107
                     note = V7087 Variant
source              1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 100
accaagggtga aggactacta cataggcttg gacatcggca cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa cgtgcttaag tttaatagca agaaatgtg gggcgtgcgg 120
ctggttcgacg agcctaagac ggcagaggag cgtaggggcc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcga agaggtagca 240
aaggtcgacc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
ttcataaga agtaccaccac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgcacaaaga cagggtgcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag ttgaaacgag 540
ctcaaaagttc attgaacga caggtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtc 660
gtaatcgcg acaccaagtt cctcaaagcc gtttccgca taatgatcgg ctctagccag 720
aaactcgtcg acttgttcga gaaccccgag gatttcgacg actctcgat aaagtccgtt 780
gactctcaa ctacctcttt cgacgacaag tactctgact atgaactgc tctgggtgac 840
aagatcgctc tggtaacat ccttaaggaa atttaacgata gtcctcatc cgagaacctg 900

```

-continued

```

ctcaaagagg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctgggtacg acaatatcac 1020
aagagtgctg attttgatat ttccagatcc gagaaaggta atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaagcaa acaagttcac tgatcaagag 1140
gcttcttaca aattcgccaa gaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aatcttatgc ccaaaattaa gtccagtgac aacgggggtg tccataacca gctcaagctg 1320
atggaaattg acaaaatctc cgagaatcag tcaaaagcat acgagttcct caatgtcagc 1380
gacgagtacg gctcgtgtg tgataaaatc gcatctatca tggaaattga tattccgtat 1440
tacgtggggc cgctgaaccg gaacagcaaa tatgctgtga tcaagaagca gaaagatagt 1500
gagattactc cctggaaact caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcggacg ctgtacttac cttaaggacg agaaggtcct tcccaagct 1620
tctttgtctg ataacgaata catgggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcacc cttaaagcgt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccca cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatct tgggocgata agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtagcagat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aataacttcac gcacagtga attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggtctctcaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcacg agggaatata accttaactc gatggagctt 2160
atgtcagctt catttacgtt ccacgaagag atacagaaac ttaacccctg ggatgaccgc 2220
aagctgtcat acgaaatggg ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcactcg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag ccgcgaagga gtcaaggaa 2400
aacgtcctga tcaagttcta taagacgga aagaaggcat tcatcagcga gattggcag 2460
gagaggtact ctacttctg ttctgagata gaggggtgag aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgtgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaaca gctatcttat ccacgacgaa atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt acgataaagg tcttatcggt cagaagaagt acactcggtg gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaactaga 2880
caagcaacaa aggagactcg taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcgaagat tgcacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcgg 3120
agctacaatc tcgagatatg gtccaagtac gatgtgaaac gtggcgagaa caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaaggca ccaactctat gttcacaaag atgaactaca ttgagtctgg agcgtttttc 3300
aacgccactc tccagcgga gataaagggc tccagacccc tgaaggacaa aggcgccgaa 3360
tcttccatcg agaagtacgg cggctacaca aacatcaata aagcctgttt cgtgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgttc gacgagattt tctcaaaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttgtc taccagtggt tgaanaatgc cactctgctc 3600
aagatcgacg gcagctacta ctctacaca ggcggttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct ccctaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctgggt 3780
tataacactt tcgacatcat gctcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttccaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggccaagg cgattccggc atccggcgtg 3960
aggaaggatc tcaagttccat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 101      moltype = DNA length = 4107
FEATURE             Location/Qualifiers
misc_feature         1..4107
                     note = V7088 Variant
source               1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 101
accaaggtag aggaactacta cataggcttg gacatcggca cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctggttcgacg acgctaagac ggacagaggag cgtagggggc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcgca agaggttagca 240
aaggtcgacc ccaactcttt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaggacaa atatacactg ttcaacgaca aggaactcaa ggataagaat 360
tttcataaga agtaccctac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgcacaagaa caggggtcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag ttggaacgag 540
ctcaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtc 660
gtaatcgccg acaccaagtt cctcaagacc gtttcgcgca taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctcgcat aaagtcggtt 780

```

-continued

```

gactctctcaa ctacctcttt cgaagacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtoaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggc cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctgggtacg acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaagcadc acgagttcct caatgtcagc 1380
gacgagtacg gctccgtgtg tgataaaatc gcatctatca tggaaattag catcccgat 1440
tacgtggggc cgctgaaccc gaacagcaaa tatgcgtgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac ttgctggacg ctgtactttac cttaaggacg agaaggctcct tcccaagct 1620
tctttgctgc ataacgaata catggtgtctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccaggagagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaaaacggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccga cggtgacttc aagcagggtc tgaactcata caatgacttc 1860
aaagctatcg tgggocgata agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cgggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagtgag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgtcg acgggactcg aggggtgcaaa caagatcact 2100
ggagagcggt gctccattat tcacttcacg agggaaatata acctaatct gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccccggt ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgtct 2280
tggcagctcc tgcgcacgt cgacgagata aagaacatca tgggaacgca cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggaa 2400
aacgtctcga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcag 2460
gagaggtact cttacttgct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgtctga tgtactctt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaaca gctatcctat ccacagacgaa atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt cggataaaag tcttatcggt cagaagaagt acactcggt gacccggaga 2820
actggcttca ccgacgacga gctcgttcag ttcatctcaa gacagatcgt ggaactaga 2880
caagcaacaa agagagactgc taacctgtctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcaagagt tcgacatcgt gaagtgtagg 3000
ggcgtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgagg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gtccacaaga atgaactaca ttgagctcgg agcgttttc 3300
aacgcccact tccagcgga ggaataaggc tccagacccc tgaaggacaa aggccgga 3360
tcttccatcg agaagtcagg gggtacacaa aacatcaata aagcctgtt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaaaagta cctcaaggac 3540
cgatttggca tcgaggacta cagggttgtc taccagtggt tgaataatg cactactgctc 3600
aagatcgacg gcagctacta ctccatcaca ggctgttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct ccctaagaag aacgagtggt cgatcaaaaca gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctgggtg 3780
tataacactt tcgacatcat cgtccacaag ttcaagacca gtgtotcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgaagac acttctcatg ctgggtcaagg cgattcgggc atccggcggtg 3960
aggcaggatc tcaagtcctat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 102      moltype = DNA length = 4107
FEATURE             Location/Qualifiers
misc_feature         1..4107
                     note = V7090 Variant
source               1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 102
accaagggtga aggactacta cataggcttg gacatcgcca cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataa cgtgcttaag ttttaatagca agaaaatgtg gggcggtggc 120
ctgttcgacg acgctaagag ggacagagg cgtaggggcc agcagaggag aagacgagc 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgccga agaggtagca 240
aaggtcgacc ccaactcttt cctcagggtg gacaattccg atctgtacat ggaagataag 300
gaccagaac tgaaaagcaa atatacactg ttcaacgaca aggaactcaa ggataagaat 360
tttcataaga agtaacccac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa cagggggtcat 480
ttcatcttgc agggccagaa gttcgacact aaatcaagct tcgagaacag tttgaacgag 540
ctcaaaagtc atttgaacga caggtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgacct aaaactcaat aaacggcca agaagaagga gctgaagtcc 660

```

-continued

```

gtaatcggcg acaccaagtt cctcaaagcc gtttcgcgca taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgat aaagtcogtt 780
gactttctcaa ctacctcttt cgacgcacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tgggtcaacat ccttaaggaa atttaacgata gctccatcct cgagaacctg 900
ctcaaagagg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtagc acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacgggggtga ttccatacca gctcaagctg 1320
atggaattga ccaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacgagtagc gctccgtgtg tgataaaatc gcatctatca tgggaatttat tatccgtat 1440
tacgtgggccc cgctgaaccc gaacagcaaaa tatgctgga tcaagaagca gaaagatagt 1500
gagattactc cctggaaactt caaggacgctc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcggagc ctgtacttac ctttaaggacg agaaggtcct tcccaaagct 1620
tctttgctgt gtaacgaata ctggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca cggaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcacc ctaaaagggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccca cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggagcata agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cgttgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aataacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggt cctccattat tcacttcag agggaaatata acctaatct gatggagctt 2160
atgtcagctt gattctacgt caccgaagag atacagaaac ttaacctcgt ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tacctttctc ccagtgtgaa acggtatgctc 2280
tggcagtcct tcgcctcatg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggaa 2400
aacgtcctga tcaagttcta taagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact tctactgtct ttctgagata gagggtagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgtcgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtag cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaaca gctatcttat ccagacgaa atcctgaaca agaactgtta cgctactagg 2760
aagatcctgt acgataaagg tcttatcggt cagaagaagt acactcggtc gacccgagga 2820
actggcttca cggagcagca gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgagg 3120
agctacaatc tcgagaatat gttcaagtag gatgtgaaac gtggcgagata caccgcttgg 3180
atcgccgatg aacgaagggt gttcacaaga atgaaactaca ttgagtctgg agcgcttttc 3240
ctggaaggca caaattatag gttcacaaga atgaaactaca ttgagtctgg agcgcttttc 3300
aacgccactc tccagcgga gaataagggt tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtagc gggtacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaagcaca gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggata cagggttgctc taccagtggt tgaaaatgcg cacactgctc 3600
aagatcgagc gcagctacta ctctacaca ggcggttctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagatttag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtccat cgacctcaag tctgattacg gacgggtcac ttcaaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 103      moltype = DNA length = 4107
FEATURE             Location/Qualifiers
misc_feature         1..4107
                     note = V7091 Variant
source               1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 103
accaaggtga aggactacta cataggtctg gacatcgcca cctctagcgt cgggtgggccc 60
gtcaccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcggtgcgg 120
ctggtcgcagc acgctaagac ggcagaggag cgtagggggcc agcgaggagc aagacgacgt 180
ctggatcgga aagaaggagag actcagcctg ctgcaggact tcttcgccga agaggttagca 240
aaggctgaccc ccaactctctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccaccac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccgggtcgt ctacctggcc tgtcactatt tgctcaagaa caggggtcat 480
ttcatcttcg agggccagaa gttcgacact aaatcaagct tcgagaacag ttgaaacgag 540

```

-continued

```

ctcaaagttc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaagtt cctcaaagcc gtttcgcgga taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgat aaagtccgtt 780
gactctctca ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggccaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaagagg cagacaagtc taaggacggg aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtagc acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtg atgacaatta cgtcagctac 1080
actaaaagct caattagcaa caataaacgc gtcaaagcaa acaagttcac tgatcaagag 1140
gcctctacaa aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagtccaag 1260
aactttatgc ccaaaattaa gtccagtgc aacgggggtg ttccatacca gctcaagctg 1320
atgggaattg acaaaatcag cgagaatcag tcaaagcatt acgagttcct caatgtcagc 1380
gacgagtacg gctccgtgtg tgataaaatc gcatctatca tggaaattat tattccgtat 1440
tacgtggggc cgtgaaccc gaacagcaaa tatgctgtga tcaagaagca gaaagatagt 1500
gagattactc cctgggaact caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgatcggaac ctgtacttac cttaaggacg agaaggtcct tcccaagct 1620
tctttgctcg ataacgaata cagtgtgtg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatac ccgagagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaaacgggt ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccga cgtgtacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggcgataa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta ccgtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aataactcac gcacagtgc attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggtctctcaa acggtgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcatg agggaatata accttaactc gatggagctt 2160
atgtcagctt catttactgt ccacgaagag atacagaaac ttaacccctg ggatgaccgc 2220
aagctgtcat acgaattggg ggcgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagtcct tgcgcactcg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agtagggctg gggtaaggaa gaggtgaaag ccgcgaagga gtcaaggaa 2400
aacgtctcga tcaagtctta taaagacgga aagaaggcat tcctcagcga gattggcgag 2460
gagaggtact ctactctgct ttctgagata gagggtagag aagagaataa gtttcgatgg 2520
gataacctgt caacttatta tactcaactg ggtcgtgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctgggtg agaaggacct taactccaag 2700
aagggttaaca gctatcctat ccacagcga atcctgaaca agaactgtta cgctactgg 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gacccggaga 2820
actggcttca cggacgacga gctcgttcag ttcactctca gacagatcgt ggaactaga 2880
caagcaacaa aggagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaag ccgagaacgc aagtcgggtt aggcgaagat tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataaaccca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcgg 3120
agctacaatc tcgagatat gtccaagtag gatgtgaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gtccaacaga atgaactaca ttgagtctgg agcgtttttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtacgg ccgtacacaa aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagct tgagctcat 3480
gccaacaga agaacgacaa gaagctgtcc gacgagattt tctcaaaagt cctcaaggac 3540
cgatttgca tcgaggacta cagggttgtc taccagtggt tgaaaatgag cacactgctc 3600
aagatcgaga gcagtcacta ctctcatcaca ggcggttctg ataagacct ggagttgcga 3660
tctgtctcgc agctgattct ccctaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccca ggagctgggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggtcaagg cgattcgggc atccggcgtg 3960
agggcaggat ccaagttccat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 104      moltype = DNA length = 4107
FEATURE             Location/Qualifiers
misc_feature         1..4107
                     note = V7092 Variant
source               1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 104
accaagggtga aggactacta cataggcttg gacatcgcca cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa cgtgccttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctgttcgacg acgctaagac ggcagaggag cgtagggggc agcgaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgcaggact tcttcgcca agaggtagca 240
aaggtcgacc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300
gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccacac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420

```

-continued

```

aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa caggggtcat 480
tcatctctcg agggccagaa gttagacact aaatcaagct tgcagaacag ttggaacgag 540
ctcaaaagttc atttgaacga ctagtatgga ctggacctcg aatttgacaa cgagaacctg 600
atatacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaaggt cctcaaaagc gtttcgcgga taatgatcgg ctctagccag 720
aaactcgtcg acttggtcga gaaccccgag gatttcgacg actctgcgag aaagtccggt 780
gactctctca ctacctcttt ctagcacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgcgc tggtaaacat ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtacg acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaagggtga atgacaatta cgtcagctac 1080
actaaaagct caatttagcaa caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gcctcttaca aattcgccaa gaaacatctg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggatgc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacggggtga ttccatacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaagcatc acgagttcct caatgtcagc 1380
gacagtgacg gctcgtgtg tgataaaatc gcatctatca tggaaattat tattccgtat 1440
tacgtggggc cgctgaaccc gaacagcaaa tatgctgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgtc gtggacctcg actccagcag agaggagttc 1560
attgactcac tgactggagc ctgtacttac ctttaaggacg agaaggtcct tcccaagct 1620
tctttgctga ctaacgaata catggtgtcg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccaggagatg gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcaccc ttaaagcggg ggcaaacctg ctgaagaagg agttcaacat caacggcgag 1800
attctgctct ctgggaccga cggtgacttc aagcagggtc tgaactcata caatgacttc 1860
aaagctatcg tggcgataaa agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cggtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcaactcatg agggaaatata acctaatct gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaacccctg gtagtaccgc 2220
aagctgtcat cgaaaatggt ggacgaactg tacctttctc ccagtgtgaa acggatgctc 2280
tggcagctcc tgccatcgt cgacgagata aagaacatca tgggaaacga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaa cccgcaagga gtcaaggag 2400
aacgtctcga tcaagttcta taagacgga aagaaggcat tcatcagcga gattggcgag 2460
gagaggtact cttacttgct ttctgagata gaggggtgagg aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgtgca tgtactctt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacagtat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaaca gctatcctat cccagacgaa atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt cggataaagg tcttatcggg cagaagaagt acactcggct gacccgga 2820
actggcttca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaaactaga 2880
caagcaacaa agagagactgc taacctgctc aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcgggtt aggcaagagt tcgacatcgt gaagtgtagg 3000
ggcgtgaacg atcttcatca tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcgg 3120
agctacaatc tcgagaatat gttcaagtac gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg acccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gtccacaaga atgaactaca ttgagctcgg agcgttttc 3300
aacgccactc tccagcggaa gaataagggc tccagacccc tgaaggacaa aggccgga 3360
tcttccatcg agaagtacgg ggctacaca aacatcaata aagcctgttt cgtgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcagcgg tgagatctat 3480
gccaaacaga gaaacgacaa gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta caggggtgtc tacccagtgg tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctccatcaca ggcgggtctg ataagacct ggagttgcga 3660
tctgctctgc agctgattct cctcaagaag aacgagtggt cgatcaaaaca gatcgacaag 3720
tcttccgaaa acgactatct gacgatcagc cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaaggaca caagattgag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctgggtcaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtccat cgaacctcaag tctgattacg gacgggtcag ttcaaaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 105      moltype = DNA length = 4107
FEATURE             Location/Qualifiers
misc_feature         1..4107
                     note = V7093 Variant
source               1..4107
                     mol_type = other DNA
                     organism = synthetic construct

```

```

SEQUENCE: 105
accaagggtga aggactacta cataggcttg gacatcgga cctctagcgt cgggtggggc 60
gtcaccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcgtgcgg 120
ctgttcgacg acgctaagac ggcagaggag cgtagggggc agcagaggagc aagacgacgt 180
ctggatcgga agaaggagag actcagcctg ctgacggact tcttcgccga agaggtagca 240
aaggtcgacc ccaacttctt cctcaggctg gacaattccg atctgtacat ggaagataag 300

```

-continued

```

gaccagaaac tgaaaagcaa atatacactg ttcaacgaca aggacttcaa ggataagaat 360
tttcataaga agtaccceac aatacatcac ctgctgatgg atctgatcga ggacgacagt 420
aagaaggaca tccggctcgt ctacctggcc tgtcactatt tgctcaagaa caggggtcat 480
ttcatcttcg agggccagaa gtctgacact aaatcaagct tcgagaacag tttgaacgag 540
ctcaaaagtc atttgaacga cgagtatgga ctggacctcg aatttgacaa cgagaacctg 600
attaacatct tgactgaccc aaaactcaat aaaacggcca agaagaagga gctgaagtcc 660
gtaatcggcg acaccaagtt cctcaaagcc gtttccgcga taatgatcgg ctctagccag 720
aaactcgtcg acttgttoga gaaccccgag gatttcgacg actctcgcat aaagtccgtt 780
gacttctcaa ctacctcttt cgacgacaag tactctgact atgaactcgc tctgggtgac 840
aagatcgctc tggtaacatc ccttaaggaa atttacgata gctccatcct cgagaacctg 900
ctcaaaaggg cagacaagtc taaggacggt aacaaatata tcagtaatgc attcgtgaag 960
aagtacaata aacacggaca agatctgaaa gagttcaaac gtctggtagc acaatatcac 1020
aagagtgcgt attttgatat ttccagatcc gagaaggtag atgacaatta cgtcagctac 1080
actaaaagct ccaaaattga caataaacgc gtcaaaagcaa acaagttcac tgatcaagag 1140
gccttctaca aattcgccaa gaacatctcg gagacaatca agtataagat caacaaggta 1200
aacggctcca aggcagatct ggagctgatt gacgggtagc tgcgggacat ggagttcaag 1260
aactttatgc ccaaaattaa gtccagtgac aacggggtag ttcataacca gctcaagctg 1320
atggaattga acaaaatact cgagaatcag tcaaaagcat acgagttcct caatgtcagc 1380
gacgagtacg gctcgtgtg tgataaaatc gcatctatca tggaaattat tattccgtat 1440
tacgtgggcc cgtcgaaccc gaacagcaaa tatgcgtgga tcaagaagca gaaagatagt 1500
gagattactc cctggaactt caaggacgct gtggacctcg actccagcag agaggagttc 1560
attgactcac tctactttac cttaaggacg agaaggtcct tcccaagct 1620
tctttgctgt ttaacgaata ctgggtgctg aacgagctga ataacctgaa gttgaacgac 1680
cttcccatca ccgaggagat gaagaagaag atatttgacc agttgttcaa aacaagaaag 1740
aaggtcacc ctaaaagcgt ggcaaacctg ctgaagaagg agttcaacat caacggcag 1800
attctgctct ctgggacgga cggtgacttc aagcagggct tgaactcata caatgacttc 1860
aaagctatcg tgggcatata agtcgattcc gatgattacc gggacaagat tgaggagatc 1920
attaaactga tagttcttta cgtgacgat aagagttacc ttcagaagaa gattaaagct 1980
gggtatggaa aatacttcac cgacagttag attaagaaaa tggcggggct gaactacaag 2040
gattggggaa ggctctcaaa gaagctgctg acgggactcg aggggtgcaa caagatcact 2100
ggagagcggg gctccattat tcacttcatg agggaatata acctaatct gatggagctt 2160
atgtcagctt catttacgtt caccgaagag atacagaaac ttaaccctgt ggatgaccgc 2220
aagctgtcat acgaaatggt ggacgaactg tactttcttc ccagtgtgaa acggatgctc 2280
tggcagtcoc ttaagcatcg cgacgagata aagaacatca tgggaaccga cagtaagaag 2340
attttcatcg agatggctcg gggtaaggaa gaggtgaaag cccgcaagga gtcaaggaaag 2400
aacgtcctga tcaagttcta taaagacgga aagaaggcat tcatcagcga gattggcag 2460
gagaggtact cttaactgtt ttctgagata gagggtagag aagagaataa gtttcgatgg 2520
gataacctgt acctttatta tactcaactg ggtcgtgca tgtactcttt ggaacctatc 2580
gacatatctg agctgtcttc aaagaatatt tacgatcagg atcatatcta ccccaaaagc 2640
aagatttacg acgacatgat cgagaatagg gtgctggtga agaaggacct taactccaag 2700
aagggtaaca gctatcttat ccacagcga atcctgaaca agaactgtta cgcctactgg 2760
aagatcctgt acgataaagg tcttatcggg cagaagaagt acactcggct gaccgcgaga 2820
actggcttca cggacgacga gctcgttcag ttcatctcaa gacagatcgt ggaaactaga 2880
caagcaacaa aggagactgc taacctgtct aagacaatat gtaagaactc cgagatcgtg 2940
tattccaaag ccgagaacgc aagtcggttt aggcaagagt tcgacatcgt gaagtgtagg 3000
gcggtgaacg atcttcata tatgcacgat gcctacatca acatcatagt ggggaacgtg 3060
tataacacca agttcacgaa ggacctatg aatttcgtaa agaagcagga aaaggcgcgg 3120
agctacaatc tcgagaatat gttcaagtag gatgtgaaac gtggcggata caccgcttgg 3180
atcgccgatg acgagaaggg caccgtgaag aacgcgagta ttaaacgtat ccggaaggag 3240
ctggaaggca caaattatag gttcacaaaga atgaactaca ttgagctctg agcgtctttc 3300
aacgccactc tccagcgga gaataagggc tccagacccc tgaaggacaa aggcccgaaa 3360
tcttccatcg agaagtcagg cgggtacaca aacatcaata aagcctgttt cgctgttctt 3420
gacatcaagt ctaagaacaa gattgagagg aagctgatgc ccgtcgagcg tgagatctat 3480
gccaacacga agaacgacaa gaagctgtcc gacgagattt tctcaagta cctcaaggac 3540
cgatttgcca tcgaggacta cagggttgct taccagtg tgaaaatgcg cacactgctc 3600
aagatcgacg gcagctacta ctctacaca ggcggttctg ataagacctt ggagttgcga 3660
tctgctctgc agctgattct ccttaagaag aacgagtggt cgatcaaaac gatcgacaag 3720
tcttccgaaa acgactatct gacgatcgag cgtatccagg acctgaccga ggagctggtg 3780
tataacactt tcgacatcat cgtcaacaag ttcaagacca gtgtcttcaa gaagtctttc 3840
cttaacttgt ttcaggacga caagatttag aacattgact tcaagtttaa gtccatggac 3900
ttcaaggaga aatgcaagac acttctcatg ctggtaagg cgattcgggc atccggcgtg 3960
aggcaggatc tcaagtccat cgacctcaag tctgattacg gacggctcag ttcaagacc 4020
aacaacatcg gcaattacca ggagttcaag attattaatc agtccatcac tggactgttc 4080
gagaatgagg tcgatctcct gaagctg 4107

```

```

SEQ ID NO: 106      moltype = DNA length = 4107
FEATURE            Location/Qualifiers
misc_feature        1..4107
                    note = V7094 Variant
source              1..4107
                    mol_type = other DNA
                    organism = synthetic construct

```

```

SEQUENCE: 106
accaaggtag aggaactacta cataggcttg gacatcgga cctctagcgt cgggtggggc 60
gtcacccgatg aagcctataa cgtgcttaag tttaatagca agaaaatgtg gggcgtggcg 120
ctgttcgacg acgctaagac ggcagaggag cgtaggggcc agcgaggagc aagacgacgt 180

```

-continued

ctggatcgga	agaaggagag	actcagcctg	ctgcaggact	tcttcgccga	agaggtagca	240
aaggctcgacc	ccaacttctt	cctcaggctg	gacaattccg	atctgtacat	ggaagataag	300
gaccagaaac	tgaaaagcaa	atatacactg	ttcaacgaca	aggacttcaa	ggataaagaat	360
tttcataaga	agtacccccc	aatacatcac	ctgctgatgg	atctgatcga	ggacgacagt	420
aagaaggaca	tccggctcgt	ctacctggcc	tgctcactatt	tgctcaagaa	caggggtcat	480
ttcatcttcg	aggcccgaaa	gttcgacact	aatcaagct	tcgagaacag	tttgaacgag	540
ctcaaagttc	atttgaacga	cgagtatgga	ctggacctcg	aatttgacaa	cgagaacctg	600
attaacatct	tgactgaccc	aaaactcaat	aaaacggcca	agaagaagga	gctgaagtcc	660
gtaatcggcg	acaccaagtt	cctcaaagcc	gtttccgcga	taatgatcgg	ctctagccag	720
aaactcgtcg	acttggtcga	gaaccccgag	gatttcgacg	actctcgcat	aaagtccgtt	780
gacttctcaa	ctacctcttt	cgacgacaag	tactctgact	atgaactcgc	tctgggtgac	840
aagatcgctc	tggccaacat	ccttaaggaa	atttacgata	gctccatcct	cgagaacctg	900
ctcaaagagg	cagacaagtc	taaggacggg	aacaaatata	tcagtaatgc	attcgtgaag	960
aagtacaata	aaccccgaaa	agatctgaaa	gagttcaaac	gtctggtagc	acaatatcac	1020
aagagtgcgt	attttgatat	tttcagatcc	gagaagggtga	atgacaatta	cgtagcgtac	1080
actaaaagct	caatttagcaa	caataaacgc	gtcaaagcaa	acaagttcac	tgatcaagag	1140
gccttctaca	aattcgccaa	gaaacatctg	gagacaatca	agtataagat	caacaaggta	1200
aacggctcca	aggcgatct	ggagctgatt	gacgggatgc	tgccggacat	ggagtccaag	1260
aactttatgc	ccaaaattaa	gtccagtgac	aacggggtga	ttccatacca	gctcaagctg	1320
atggaattga	acaaaatact	cgagaatcag	tcaaagcatc	acgagttcct	caatgtcagc	1380
gacgagtacg	gctccgtgtg	tgataaaatc	gcatctatca	tggaatttat	tattccgtat	1440
tacgtgggcc	cggaggagat	gaacagcaaa	tatgcgtgga	tcaagaagca	gaaagatagt	1500
gagattactc	cctggaactt	caaggacgtc	gtggacctcg	actccagcag	agaggagttc	1560
attgactcac	tgatcggacg	ctgtacttac	cttaaggacg	agaaggctct	tcccaagact	1620
tctttgtctg	ctaacgaata	catggtgtctg	aacgagctga	ataacctgaa	gttgaacgac	1680
cttcccatca	cggaggagat	gaagaagaag	atatttgacc	agttgttcaa	aacaagaaag	1740
aaggctaccc	ttaaagcggg	ggcaaacctg	ctgaagaagg	agttcaacat	caacggcgag	1800
attctgctct	ctgggaccga	cggtgacttc	aagcagggtc	tgaactcata	caatgacttc	1860
aaagctatcg	tggggcgtaa	agtcgattcc	gatgattacc	gggacaagat	tgaggagatc	1920
attaaactga	tagttcttta	cggtagcatg	aagagttacc	ttcagaagaa	gattaaagct	1980
gggtatggaa	aatacttcac	cgacagttag	attaaagaaa	tggcggggct	gaactacaag	2040
gattggggaa	ggctctcaaa	gaagctgctg	acgggactcg	aggggtgcaa	caagatcact	2100
ggagagcggg	gctccattat	tcacttcacg	agggaaatata	accttaactc	gatggagctt	2160
atgtcagctt	catttaccgt	gaacgaagag	atacagaaac	ttaaccccg	ggatgaccgc	2220
aagctgtcat	acgaaatggt	ggacgaactg	tacctttctc	ccagtgtgaa	acggatgctc	2280
tggcagtcct	tgccgcatcgt	cgacgagata	aagaacatca	tgggaaccga	cagtaagaag	2340
attttcatcg	agatggctcg	gggttaaggaa	gaggtgaaag	cccgaagga	gtcaagggaag	2400
aacgtcctga	tcaagtctta	taaaagcggg	aagaaggcat	tcacagcgga	gattggcgag	2460
gagagggtact	cttacttctg	ttctgagata	gaggggtgag	aagagaataa	gtttcgatgg	2520
gataacctgt	acctttatta	tactcaactg	ggtcgtcgca	tgtactcttt	ggaacctatc	2580
gacatatctg	agctgtcttc	aaagaatatt	tacgatcagg	atcatatcta	ccccaaaagc	2640
aagatttacg	acgacatagg	gtgctgggtga	agaaggacct	taactccaag	2700	
aagggttaaca	gctatcctat	cccagacgaa	atcctgaaca	agaactgtta	cgctactgg	2760
aagatcctgt	acgataaagg	tcttatcggg	cagaagaagt	acactcggct	gacccggaga	2820
actggcttca	cggacgacga	gctcgttcag	ttcatctcaa	gacagatcgt	ggaactaga	2880
caagcaacaa	aggagactcg	taacctgctc	aagacaatat	gtaagaactc	cgagatcggt	2940
tattccaaag	ccgagaacgc	aagtcgggtt	aggcaagagt	tcgacatcgt	gaagtgtagg	3000
gcggtgaacg	atcttcatca	tatgcacgat	gcctacatca	acatcatagt	ggggaacgtg	3060
tataacacca	agttcacgaa	ggacctatg	aatttcgtaa	agaagcagga	aaaggcgagg	3120
agctacaatc	tcgagaatat	gttcaagtac	gatgtgaaac	gtgcccagata	caccgcttgg	3180
atcgccgatg	acgagaaggg	caccgtgaag	aacgcgagta	ttaaacgtat	ccggaaggag	3240
ctggaaggca	caaattatag	gttcacaaga	atgaactaca	ttgagtctgg	agcgcttttc	3300
aacgcccactc	tccagcggaa	gaataagggc	tccagacccc	tgaaggacaa	aggcccgaag	3360
tcttccatcg	accaggtacg	cggtacacaa	aacatcaata	aagcctgttt	cgctgttctt	3420
gacatcaagt	ctaagaacaa	gattgagagg	aagctgatgc	ccgtcgagcg	tgagatctat	3480
gccaacacga	agaacgacaa	gaagctgtcc	gacgagattt	tctcaaagta	cctcaaggac	3540
cgatttggca	tcgaggacta	cagggttgtc	taccagtggt	tgaaaatgcg	cacactgtct	3600
aagatcgacg	gcagactact	cttcacacaa	ggcggttctg	ataagacctc	ggagttgcga	3660
tctgctctgc	agctgattct	ccctaagaag	aacgagtggt	cgatcaaaac	gatcgacaa	3720
tcttccgaaa	acgactatct	gacgatcgag	cgtatccagg	acctgaccga	ggagctgggt	3780
tataacacct	tcgacatcat	cgtcaacaag	tccaagacca	gtgtcttcaa	gaagtctttc	3840
cttaacttgt	ttcaggacga	caagatttag	aacattgact	tcaagtttaa	gtccatggac	3900
ttcaaggaga	aatgcgaagc	acttctcatg	ctgggtcaag	cgattcgggc	atcccgctg	3960
aggcaggatc	tcaagtccat	cgacctcaag	tctgattacg	gacggctcag	ttcaaagacc	4020
aacaacatcg	gcaattacca	ggagttcaag	attattaatc	agtcacatcac	tgagctgttc	4080
gagaatgagg	tcgatctcct	gaagctg				4107

1. A non-naturally occurring OMNI-50 nuclease variant having a wild-type OMNI-50 protein sequence (SEQ ID NO: 1) comprising an amino acid substitution in at least one of the following positions: R478, Y545, R61, Y437, A493, G606, K688, L690, E695, L718, R788, Q803, L805, L844, V981, K965, and K1036.

2. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at position R478 and/or position Y545.

3. The OMNI-50 nuclease variant of claim 2, comprising an amino acid substitution at each of positions R478 and Y545.

4. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution at position R478 is any one of the following substitutions: R478D, R478E, R478S, R478T, R478N, R478Q, R478G, R478P, R478C, R478A, R478V, R478I, R478L, R478M, R478F, R478Y, or R478W, preferably R478V, R478H, R478L, R478M, R478P, R478F, R478W, R478Y, R478S, R478C, R478T, R478N, or R478Q.

5. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position R478 and the amino acid substituted for arginine is an amino acid having a negatively charged R-group or an R-group lacking a charge.

6. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position R478 and the amino acid substituted for arginine is a polar amino acid or non-polar amino acid.

7. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position R478 and the amino acid substituted for arginine is a non-polar amino acid.

8. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution at position Y545 is any one of the following substitutions: Y545D, Y545E, Y545S, Y545T, Y545N, Y545Q, Y545G, Y545P, Y545C, Y545A, Y545V, Y545I, Y545L, Y545M, Y545F, Y545R, Y545K, Y545H, or Y545W.

9. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is an amino acid having a negatively charged R-group or a positively charged R-group.

10. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine is a non-polar amino acid.

11. The OMNI-50 nuclease variant of claim 2, wherein the amino acid substitution is at position Y545 and the amino acid substituted for tyrosine lacks a phenyl ring or is phenylalanine.

12. The OMNI-50 nuclease variant of claim 3, wherein the amino acid substitutions are R478I and Y545H.

13. The OMNI-50 nuclease variant of claim 1, wherein the amino acid substitution is any one of the following substitutions: R478I, Y545H, Q803V, L805I, R61A, Y437W, R788K, L844N, V981M, G606P, L690V, E695Q, R478T, A493G, K688E, L718C, K965V, and K1036V.

14. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions R478, Y545, Q803, and L805.

15. The OMNI-50 nuclease variant of claim 14, wherein the amino acid substitutions are R478I, Y545H, Q803V, and L805I.

16. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions R61, Y437, R788, L844, and V981.

17. The OMNI-50 nuclease variant claim 16, wherein the amino acid substitutions are R61A, Y437W, R788K, L844N, and V981M.

18. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions G606, L690, and E695.

19. The OMNI-50 nuclease variant of claim 18, wherein the amino acid substitutions are G606P, L690V, and E695Q.

20. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions R478, A493, K688, L718, K965, and K1036.

21. The OMNI-50 nuclease variant of claim 20, wherein the amino acid substitutions are R478T, A493G, K688E, L718C, K965V, and K1036V.

22. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at position Q803.

23. The OMNI-50 nuclease variant of claim 22, wherein the amino acid substitution is Q803V.

24. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at position Y545.

25. The OMNI-50 nuclease variant of claim 24, wherein the amino acid substitution is Y545H.

26. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at position L805.

27. The OMNI-50 nuclease variant of claim 26, wherein the amino acid substitution is L805I.

28. The OMNI-50 nuclease variant of claim 1 comprising an amino acid substitution at each of positions R478, Y545, and Q803.

29. The OMNI-50 nuclease variant of claim 28, comprising an amino acid substitutions are R478I, Y545H, AND Q803V.

30. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions R478, Q803, and L805.

31. The OMNI-50 nuclease variant of claim 30, wherein the amino acid substitutions are R478I, Q803V, and L805I.

32. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions R478, Q803, and L805.

33. The OMNI-50 nuclease variant of claim 32, wherein the amino acid substitutions are R478I, Q803V, and L805I.

34. The OMNI-50 nuclease variant of claim 1, comprising an amino acid substitution at each of positions Y545, Q803, and L805.

35. The OMNI-50 nuclease variant claim 34, wherein the amino acid substitutions are R478I, Y545H, and L805I.

36. The OMNI-50 nuclease variant of claim 1, having an amino acid sequence selected from the group consisting of SEQ ID NOs: 2-5, 63-70, and 89-97.

37. The OMNI-50 nuclease variant of claim 1, having at least 80% sequence identity to the wild-type OMNI-50 protein sequence (SEQ ID NO:1).

38. The OMNI-50 nuclease variant of claim 1, further comprising a nuclear localization sequence (NLS).

39. The OMNI-50 nuclease variant of claim 1, wherein the variant exhibits increased specificity toward a DNA target site when complexed with a guide RNA molecule that targets variant to the said DNA target site relative to a wild-type OMNI-50 nuclease complexed with the guide RNA molecule.

40. A CRISPR system comprising the OMNI-50 nuclease variant of claim 1, wherein the OMNI-50 nuclease variant is complexed with a guide RNA molecule that targets a DNA

target site, and wherein the CRISPR system displays reduced off-target editing activity relative to a wild-type CRISPR system comprising a wild-type OMNI-50 nuclease protein and the guide RNA molecule.

41. A method for gene editing having reduced off-target editing activity, comprising contacting a DNA target site with an active CRISPR system comprising an OMNI-50 nuclease variant protein of claim 1.

42. The method of claim 41, wherein the active CRISPR system displays reduced off-target editing activity relative to a wild-type CRISPR system comprising a wild-type OMNI-50 nuclease protein.

43. The method of claim 41, wherein the gene editing occurs in a eukaryotic cell or prokaryotic cell.

44. The method of claim 43, wherein the eukaryotic cell is a plant cell or mammalian cell.

45. The method of claim 44, wherein the mammalian cell is a human cell.

46. The method of claim 41, wherein the DNA target site is located within or in proximity to a pathogenic allele of a gene.

47. The method of claim 46, wherein the DNA target site is located in a gene selected from the group consisting of ELANE, CXCR4, EMX, RyR2, KNCQ1, KCNH2, SCN5a, GBA1, GBA2, Rhodopsin, GUCY2D, IMPDH1, FGA, BEST1, PRPH2, KRT5, KRT14, ApoA1, STAT3, STAT1, ADA2, RPS19, SBDS, GATA2, RPE65, LDLR, ANGPTL3, B2M, TRAC, TCF4, TGFB1, PAX6, C3, LRRK2, SARM1, SAMD9, SAMD9L, HAVCR2, CD3E, APLP2, CISH, TIGIT, TNNT2, TNN, MYH7, and HLA-E.

48. The method of claim 41 wherein the DNA target is repaired with an exogenous donor molecule.

49. The method of claim 41, wherein the off-target editing activity is reduced by at least 2-fold, 10-fold, 10^2 -fold, 10^3 -fold, 10^4 -fold, 10^5 -fold, or 10^6 -fold.

50. A modified cell obtained by the method of claim 43.

51. The modified cell of claim 50, wherein the cell is capable of engraftment.

52. The modified cell of claim 51, wherein the cell is capable of giving rise to progeny cells after engraftment.

53. The modified cell of claim 51, wherein the cell is capable of giving rise to progeny cells after an autologous engraftment.

54. The modified cell of claim 51, wherein the cell is capable of giving rise to progeny cells for at least 12 months or at least 24 months after engraftment.

55. The modified cell of claim 51, wherein the cell is selected from the group consisting of a hematopoietic stem cell, a progenitor cell, a CD34+ hematopoietic stem cell, a bone marrow cell, and a peripheral mononucleated cell.

56. A composition comprising a modified cell of claim 50 and a pharmaceutically acceptable carrier.

57. An in vitro or ex vivo method of preparing the composition of claim 56, comprising mixing the cells with the pharmaceutically acceptable carrier.

58. A polynucleotide molecule encoding the OMNI-50 variant protein claim 1.

* * * * *